



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**


(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/07/31
(87) Date publication PCT/PCT Publication Date: 2019/02/07
(85) Entrée phase nationale/National Entry: 2020/01/14
(86) N° demande PCT/PCT Application No.: US 2018/044508
(87) N° publication PCT/PCT Publication No.: 2019/027960
(30) Priorité/Priority: 2017/08/01 (US62/539,642)

(51) Cl.Int./Int.Cl. *C07D 403/12* (2006.01),
A61K 31/416 (2006.01), *A61K 31/437* (2006.01),
A61K 31/519 (2006.01), *A61P 29/00* (2006.01),
C07D 487/04 (2006.01)
(71) Demandeur/Applicant:
THERAVANCE BIOPHARMA R&D IP, LLC, US
(72) Inventeurs/Inventors:
FENSTER, ERIK, US;
LAM, TOM M., US;
LOO, MANDY, US;
MCKINNELL, ROBERT MURRAY, US;
PALERMO, ANTHONY FRANCESCO, US;
WANG, DIANA JIN, US;

(54) Titre : COMPOSES BICYCLIQUES PYRAZOLO ET TRIAZOLO UTILISES EN TANT QU'INHIBITEURS DE KINASE
JAK

(54) Title: PYRAZOLO AND TRIAZOLO BICYCLIC COMPOUNDS AS JAK KINASE INHIBITORS



(57) Abrégé/Abstract:

The invention provides compounds of formula (I) or a pharmaceutically-acceptable salt thereof, wherein the variables are defined in the specification, that are inhibitors of JAK kinases, particularly JAK3. The invention also provides crystalline forms, pharmaceutical

(72) **Inventeurs(suite)/Inventors(continued):** FRAGA, BREENA, US; NZEREM, JERRY, US; DABROS, MARTA, US; THALLADI, VENKAT R., US; RAPTA, MIROSLAV, US

(74) **Agent:** TORYS LLP

(57) **Abrégé(suite)/Abstract(continued):**

compositions comprising such compounds, methods of using such compounds to treat gastrointestinal and other inflammatory diseases, and processes and intermediates useful for preparing such compounds.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2019/027960 A1

(43) International Publication Date
07 February 2019 (07.02.2019)

(51) International Patent Classification:

C07D 403/12 (2006.01) *A61K 31/416* (2006.01)
C07D 487/04 (2006.01) *A61K 31/519* (2006.01)
A61P 29/00 (2006.01) *A61K 31/437* (2006.01)

(21) International Application Number:

PCT/US2018/044508

(22) International Filing Date:

31 July 2018 (31.07.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/539,642 01 August 2017 (01.08.2017) US

(71) Applicant: **THERAVANCE BIOPHARMA R&D IP, LLC** [US/US]; 901 Gateway Boulevard, South San Francisco, California 94080 (US).

(72) Inventors: **FENSTER, Erik**; 535 Redwood Avenue, San Bruno, California 94066 (US). **LAM, Tom M.**; 1280 15th Avenue, Apt. 201, San Francisco, California 94122 (US). **LOO, Mandy**; 2119 Imwalle Court, San Jose, California 95131 (US). **MCKINNELL, Robert Murray**; 321 Poplar Avenue, Millbrae, California 94030 (US). **PALERMO, Anthony Francesco**; 170 Scotia Avenue, San Francisco, California 94124 (US). **WANG, Diana Jin**; 3590 Cydonia Ct., Dublin, California 94568 (US). **FRAGA, Breena**; 9311 Road 29 1/2, Madera, California 93637 (US). **NZEREM, Jerry**; 29 Woods Circle, South San Francisco, California 94080 (US). **DABROS, Marta**; 888 Foster City Boulevard, Apt. Q2, Foster City, California 94404 (US). **THALLADI, Venkat R.**; 888 Foster City Boulevard, Apt. E3, Foster City, California 94404 (US). **RAPTA, Miroslav**; 1240 Highland Court, San Carlos, California 94070 (US).

(74) Agent: **HAGENAH, Jeffrey A.** et al.; Theravance Biopharma US, Inc., 901 Gateway Boulevard, South San Francisco, California 94080 (US).


(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PYRAZOLO AND TRIAZOLO BICYCLIC COMPOUNDS AS JAK KINASE INHIBITORS



(57) Abstract: The invention provides compounds of formula (I) or a pharmaceutically-acceptable salt thereof, wherein the variables are defined in the specification, that are inhibitors of JAK kinases, particularly JAK3. The invention also provides crystalline forms, pharmaceutical compositions comprising such compounds, methods of using such compounds to treat gastrointestinal and other inflammatory diseases, and processes and intermediates useful for preparing such compounds.

WO 2019/027960 A1

5

PYRAZOLO AND TRIAZOLO BICYCLIC COMPOUNDS AS JAK KINASE INHIBITORS

10

BACKGROUND OF THE INVENTION

Field of the Invention

15 The invention is directed to pyrazolo and triazolo bicyclic compounds useful as JAK kinase inhibitors and more particularly as JAK3 inhibitors that are selective for JAK3 over other members of the JAK kinase family such as JAK1, JAK2 and TYK2. The invention is also directed to crystalline forms, pharmaceutical compositions comprising such compounds, methods of using such compounds to treat inflammatory
20 diseases, and processes and intermediates useful for preparing such compounds.

State of the Art

 Ulcerative colitis is a chronic inflammatory disease of the colon. The disease is characterized by inflammation and ulceration of the mucosal layer of the rectum and the
25 large intestine. Common symptoms include diarrhea, bloody stools, and abdominal pain. The clinical course is intermittent, marked by alternating periods of exacerbation and remission. Incidence seems to be greater in developed than in developing countries. An estimated 1.2 million people in major industrialized countries suffer from ulcerative colitis and the numbers are expected to increase along with population growth. Patients
30 with ulcerative colitis are at an increased risk of developing colorectal cancer. (e.g. Danese et al. *N Engl J Med*, **2011**, 365, 1713-1725). Although there exists a variety of therapeutic options to promote and maintain remission of ulcerative colitis (UC) in patients, none is ideal. There remains an unmet medical need for an effective therapy to

promote and maintain remission of moderate to severe UC without the safety concerns resulting from chronic, systemic immunosuppression.

Although the precise pathogenesis of UC is unclear, it is apparent that proinflammatory cytokines play a pivotal role in the immunological response (Strober et al., *Gastroenterol*, **2011**, 140, 1756-1767). Many of the proinflammatory cytokines most commonly elevated in UC (e.g., IL-4, IL-6, IL-13, IL-15, IL-23, IL-24, IFN γ and leptin), rely on the JAK family of tyrosine kinases (i.e., JAK1, JAK2, JAK3 and Tyk2) for signal transduction.

Inhibition of the JAK3 enzyme blocks the signaling of many key pro-inflammatory cytokines. Thus JAK3 inhibitors are likely to be useful in the treatment of ulcerative colitis and other gastrointestinal inflammatory diseases such as Crohn's disease and immune checkpoint inhibitor induced colitis. JAK3 inhibitors are also likely to be useful for the treatment of inflammatory skin diseases such as atopic dermatitis and inflammatory respiratory disorders such as allergic rhinitis, asthma, and chronic obstructive pulmonary disease (COPD). In addition, JAK3 inhibitors may also be useful in the treatment of many ocular diseases for which inflammation plays a prominent role such as uveitis, diabetic retinopathy, diabetic macular edema, dry eye disease, age-related macular degeneration, retinal vein occlusion (RVO) and atopic keratoconjunctivitis.

Selectivity for JAK3 over JAK1 is anticipated to be beneficial as there is evidence that JAK3 selectivity allows sparing of potentially beneficial cytokines such as IL-10 which has been involved in mucosal healing, IL-22 which is involved in mucus barrier protection and epithelial regeneration, and IL-6 which is involved in the proliferation of intestinal epithelial cells. Selectivity for JAK3 over JAK2 also allows sparing of erythropoietin (EPO) and thrombopoietin (TPO) signaling. Therefore, it would be desirable to provide new compounds which are selective JAK3 inhibitors over other members of the JAK kinase family such as JAK1, JAK2 and TYK2.

Finally, due to the modulating effect of the JAK/STAT pathway on the immune system, systemic exposure to JAK inhibitors may have an adverse systemic immunosuppressive effect. It would be desirable, therefore, to provide new JAK3 inhibitors which have their effect at the site of action without significant systemic effects. In particular, for the treatment of gastrointestinal inflammatory diseases, such as ulcerative colitis, it would be desirable to provide new JAK3 inhibitors which can be administered orally and achieve therapeutically relevant exposure in the gastrointestinal tract with minimal systemic exposure. For skin diseases, it would be desirable to provide


new JAK3 inhibitors that could be administered topically to the skin with minimal systemic exposure.

Therefore, it would be desirable to provide new compounds which are selective JAK3 inhibitors over other members of the JAK kinase family such as JAK1, JAK2 and
5 TYK2, and have minimal systemic exposure.

SUMMARY OF THE INVENTION

In one aspect, the invention provides novel compounds having activity as JAK kinase inhibitors and more particularly as JAK3 inhibitors.

10 Accordingly, the invention provides a compound of formula (I):




(I)


or a pharmaceutically acceptable salt thereof, wherein


X¹ and X² are each independently selected from N and CH;

15 X³ is selected from the group consisting of N, CH, C-CH₃, C-CF₃, C-CHF₂, C-CH₂-O-CH₃, C-SMe, C-NMe₂, C-NH-CH₃, C-Cl, C-CN, and C-OMe;



is selected from the group consisting of





R^a , R^b , R^c , and R^f are each independently selected from the group consisting of H and C_{1-3} alkyl;

R^d , R^e , R^g , R^h , R^i , R^j , R^l , R^m , R^n and R^o are each independently selected from the group consisting of H and C_{1-3} alkyl wherein the C_{1-3} alkyl group may be optionally substituted with 1 to 3 halogens;

Optionally R^d and R^e may be joined to form a cyclopropyl ring;

A is selected from the group consisting of


(a) a 4 to 10 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, $S(O)_2$ and O, and

(b) a 6 to 10 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O,

wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 3 R^k groups;

each R^k is independently selected from the group consisting of F, CN, C_{1-3} alkoxy, cyclopropyl, and C_{1-3} alkyl, wherein the C_{1-3} alkyl group may be optionally substituted with OH, OMe or 1 to 3 halogens;

R^l is selected from the group consisting of



wherein R^p and R^q are each independently selected from the group consisting of H, C_{3-5} cycloalkyl and C_{1-6} alkyl, wherein the C_{1-6} alkyl group may be optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-3} alkoxy and $-S-C_{1-3}$ alkyl,

or R^p and R^q form a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N,

S, and O, wherein the 4 to 6 membered monocyclic heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₆ alkyl, C₁₋₃ alkoxy, -S-C₁₋₃ alkyl and -C₁₋₃ alkyl-C₁₋₃ alkoxy;

R² is selected from the group consisting of H, Cl, OMe, Me and F;

5 R³ is selected from the group consisting of H and F;

R⁴ is selected from the group consisting of H and F; and

R⁵ is selected from the group consisting of H, Me and F.

The disclosure also provides some crystalline forms of certain compounds, Form 1, Form 2, Form 2b, Form 3, and Form 4.

10 The invention also provides a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form of the disclosure, and a pharmaceutically-acceptable carrier.

The invention also provides a method of treating gastrointestinal inflammatory disease, in particular, ulcerative colitis, in a mammal, the method comprising
15 administering to the mammal a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form of the disclosure, or a pharmaceutical composition of the disclosure.

The invention also provides a method of treating inflammatory diseases or disorders of the skin in a mammal, the method comprising applying a compound of the
20 disclosure, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition of the disclosure to the skin of the mammal.

The invention also provides a method of treating cutaneous T-cell lymphoma in a mammal, the method comprising applying a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, to the skin of
25 the mammal.

In another aspect, the invention also provides a process described herein, which is useful for preparing the compounds of the disclosure.

The invention also provides a compound of the disclosure or a pharmaceutically acceptable salt thereof, as described herein for use in medical therapy, as well as the use
30 of a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, in the manufacture of a formulation or medicament for treating a gastrointestinal inflammatory disease, or an inflammatory disease of the skin in a mammal.

BRIEF DESCRIPTION OF THE DRAWINGS

Various aspects of the present invention are illustrated by reference to the accompanying drawings.

Figure 1 shows a powder x-ray diffraction (PXRD) pattern of crystalline Form 1 of compound 3 (hereinafter Form 1).

Figure 2 shows a differential scanning calorimetry (DSC) thermogram of crystalline Form 1.

Figure 3 shows a thermal gravimetric analysis (TGA) plot of crystalline Form 1.

Figure 4 shows a dynamic moisture sorption (DMS) isotherm of crystalline Form 1 observed at a temperature of about 25 °C.

Figure 5 shows a powder x-ray diffraction (PXRD) pattern of crystalline Form 2 of compound 3 (hereinafter Form 2).

Figure 6 shows a differential scanning calorimetry (DSC) thermogram of crystalline Form 2.

Figure 7 shows a thermal gravimetric analysis (TGA) plot of crystalline Form 2.

Figure 8 shows a dynamic moisture sorption (DMS) isotherm of crystalline Form 2 observed at a temperature of about 25 °C.

Figure 9 shows a powder x-ray diffraction (PXRD) pattern of crystalline Form 3 of compound 1 (hereinafter Form 3).

Figure 10 shows a differential scanning calorimetry (DSC) thermogram of crystalline Form 3.

Figure 11 shows a thermal gravimetric analysis (TGA) plot of crystalline Form 3.

Figure 12 shows a dynamic moisture sorption (DMS) isotherm of crystalline Form 3 observed at a temperature of about 25 °C.

Figure 13 shows a powder x-ray diffraction (PXRD) pattern of crystalline Form 4 of compound 1 (hereinafter Form 4).

Figure 14 shows a differential scanning calorimetry (DSC) thermogram of crystalline Form 4.

Figure 15 shows a thermal gravimetric analysis (TGA) plot of crystalline Form 4.

Figure 16 shows a dynamic moisture sorption (DMS) isotherm of crystalline Form 4 observed at a temperature of about 25 °C.


Figure 17 shows a powder x-ray diffraction (PXRD) pattern of a dehydrated crystalline form of compound 3 (hereinafter Form 2b).

DETAILED DESCRIPTION OF THE INVENTION

Among other aspects, the invention provides JAK kinase inhibitors of formula (I) which are selective for JAK3 over other members of the JAK kinase family such as JAK1, JAK2 and TYK2, pharmaceutically-acceptable salts thereof, and intermediates for the preparation thereof.

In one aspect, the invention provides novel compounds having activity as JAK kinase inhibitors, particularly as JAK3 kinase inhibitors.

Accordingly, the invention provides a compound of formula (I):




(I)


or a pharmaceutically acceptable salt thereof, wherein

X¹ and X² are each independently selected from N and CH;

X³ is selected from the group consisting of N, CH, C-CH₃, C-CF₃, C-CHF₂, C-CH₂-O-CH₃, C-SMe, C-NMe₂, C-NH-CH₃, C-Cl, C-CN, and C-OMe;



is selected from the group consisting of



R^a , R^b , R^c , and R^f are each independently selected from the group consisting of H and C₁₋₃ alkyl;

R^d , R^e , R^g , R^h , R^i , R^j , R^l , R^m , R^n and R^o are each independently selected from the group consisting of H and C₁₋₃ alkyl wherein the C₁₋₃ alkyl group may be optionally substituted with 1 to 3 halogens; optionally R^d and R^e may be joined to form a cyclopropyl ring;

A is selected from the group consisting of


(a) a 4 to 10 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, S(O)₂ and O, and

(b) a 6 to 10 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O,

wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 3 R^k groups;

each R^k is independently selected from the group consisting of F, CN, C₁₋₃ alkoxy, cyclopropyl, and C₁₋₃ alkyl, wherein the C₁₋₃ alkyl group may be optionally substituted with OH, OMe or 1 to 3 halogens;

R^1 is selected from the group consisting of



wherein R^p and R^q are each independently selected from the group consisting of H, C₃₋₅ cycloalkyl and C₁₋₆ alkyl, wherein the C₁₋₆ alkyl group may be optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₃ alkoxy and -S-C₁₋₃ alkyl,

or R^p and R^q form a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O, wherein the 4 to 6 membered monocyclic heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₆ alkyl, C₁₋₃ alkoxy, -S-C₁₋₃ alkyl and -C₁₋₃ alkyl-C₁₋₃ alkoxy;

R^2 is selected from the group consisting of H, Cl, OMe, Me and F;

R^3 is selected from the group consisting of H and F;

R^4 is selected from the group consisting of H and F; and

R^5 is selected from the group consisting of H, Me and F.

5 In some embodiments, X^3 is CH.


In some embodiments, R^p and R^q are each independently selected from the group consisting of H, cyclobutyl and C_{1-4} alkyl, wherein the C_{1-4} alkyl group may be optionally substituted with 1 or 2 substituents independently selected from the group consisting of C_{1-2} alkoxy and -S- C_{1-2} alkyl,


10 or R^p and R^q form a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from S, and O, wherein the 4 to 6 membered monocyclic heterocyclic group is optionally substituted with 1 or 2 substituents independently selected from the group consisting of C_{1-3} alkyl, C_{1-2} alkoxy, -S- C_{1-2} alkyl and - C_{1-3} alkyl- C_{1-2} alkoxy.

15 In some embodiments, R^p and R^q are each independently selected from the group consisting of H, cyclobutyl and C_{1-4} alkyl, wherein the C_{1-4} alkyl group may be optionally substituted with 1 or 2 substituents independently selected from the group consisting of OMe and -SEt,


20 or R^p and R^q form a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from S, and O, wherein the 4 to 6 membered monocyclic heterocyclic group is optionally substituted with 1 or 2 substituents independently selected from the group consisting of Me, OMe, CH_2OMe , and -SMe.

In some embodiments, R^1 is selected from the group consisting of








5

In some embodiments, R¹ is

or

In some embodiments, R¹ isIn some embodiments, R¹ is

The 6 to 10 membered multicyclic heterocyclic group may be spiro-cyclic, fused and/or bridged.

10

In some embodiments, the 6 to 10 membered multicyclic heterocyclic group is a spiro-cyclic heterocyclic group. In some embodiments, the 6 to 10 membered multicyclic heterocyclic group is a fused heterocyclic group. In some embodiments, the 6 to 10 membered multicyclic heterocyclic group is a bridged heterocyclic group.

15

In some embodiments, A is selected from the group consisting of

(a) a 4 to 8 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, S(O)₂ and O, and

20

(b) a 6 to 10 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O, wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 3 R^k groups.

In some embodiments, A is selected from the group consisting of

(a) a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, S(O)₂ and O, and

(b) a 6 to 10 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O, wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 3 R^k groups.


In some embodiments, A is selected from the group consisting of:


10 (a) a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, S(O)₂ and O, and


(b) a 7 or 8 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S and O,


15 wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to
3 R^k groups.


In some embodiments, A is selected from the group consisting of azetidine, pyrrolidine, piperidine, morpholine, 2-azaspiro[3.3]heptane, thiomorpholine and nortropane.

20 In some embodiments,  is selected from the group consisting of



In some embodiments,  is selected from the group consisting of:








10 In some embodiments, X^1 and X^2 are both CH. In some embodiments, X^1 is N and X^2 is CH. In some embodiments, X^1 and X^2 are both N. In some embodiments, X^1 is CH and X^2 is N.


In some embodiments, R^2 is H. In some embodiments, R^2 is F. In some embodiments, R^2 is Cl. In some embodiments, R^2 is OMe. In some embodiments, R^2 is

Me. In some embodiments, R^3 is H. In some embodiments, R^3 is F. In some embodiments, R^4 is H. In some embodiments, R^4 is F. In some embodiments, R^5 is H. In some embodiments, R^5 is F. In some embodiments, R^5 is Me.


In some embodiments,  is selected from the group consisting of:



In some embodiments,  is selected from the group consisting of:




The invention also provides a compound of formula (II):




(II)

15 or a pharmaceutically acceptable salt thereof, wherein

X^1 and X^2 are both CH, or X^1 and X^2 are both N, or X^1 is N and X^2 is CH;

 is selected from the group consisting of




R^c, R^d, and R^e are each independently selected from the group consisting of H and methyl;


5 A is selected from the group consisting of azetidine, pyrrolidine, piperidine, and morpholine;


wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 2 R^k groups;

each R^k is independently selected from the group consisting of F, CN, methyl, ethyl and C₁₋₂ haloalkyl;



R² is selected from the group consisting of H, Cl, and F.


In some embodiments,  is selected from the group consisting of




In some embodiments, R¹ is




In some embodiments, R¹ is





In some embodiments, the compound has the formula:




In some embodiments, the compound has the formula:




In some embodiments,  is selected from the group consisting of:




5

In some embodiments,  is selected from the group consisting of:




10

The disclosure also provides a compound, or a pharmaceutically acceptable salt thereof, of the formula (Ia) or (IIa):




(Ia)




(IIa)

wherein the variables are as defined in the embodiments above.

The disclosure also provides a compound, or a pharmaceutically acceptable salt thereof, of the formula (Ib) or (IIb):




(Ib)




(IIb)

wherein the variables are as defined in the embodiments above.

The disclosure also provides a compound, or a pharmaceutically acceptable salt thereof, of the formula (Ic) or (IIc):





(Ic)



(IIc)



wherein the variables are as defined in the embodiments above.

The disclosure also provides a compound, or a pharmaceutically acceptable salt thereof, of the formula (Id) or (IIId):






wherein the variables are as defined in the embodiments above.


The disclosure also provides a compound, or a pharmaceutically acceptable salt thereof, of the formula (Ie) or (IIe):




wherein the variables are as defined in the embodiments above.

The disclosure also provides a compound, or a pharmaceutically acceptable salt thereof, of the formula (If), (IIIf), (IIIIf), (IVf), (Vf), (VIIf), (VIIIf) or (VIIIf):







(IVf)




(Vf)



(VIf)






(VIIf)




(VIIf)

5 wherein the variables are as defined in the embodiments above.


The invention also provides a compound, or a pharmaceutically acceptable salt thereof, selected from the group consisting of



and



10 The invention also provides a compound, or a pharmaceutically acceptable salt thereof, having formula (B):



(B)


or a pharmaceutically acceptable salt thereof, wherein

X^1 and X^2 are each independently selected from N and CH;

5 R^d and R^e are each independently selected from the group consisting of H and C_{1-3} alkyl; optionally R^d and R^e may be joined to form a cyclopropyl ring;

R^{k1} is selected from the group consisting of H, F, CN, OMe, and C_{1-3} alkyl;


R^{k2} is selected from the group consisting of H and methyl;



R^2 is selected from the group consisting of H, Cl, and F.

In some embodiments, R^d and R^e are each independently selected from the group consisting of H and methyl; optionally R^d and R^e may be joined to form a cyclopropyl ring; and R^{k1} is selected from the group consisting of H, F, CN, OMe, methyl and ethyl.

15 The invention also provides a compound, or a pharmaceutically acceptable salt thereof, having formula (C):




(C)

or a pharmaceutically acceptable salt thereof, wherein

20 X^1 and X^2 are both CH, or X^1 and X^2 are both N, or X^1 is N and X^2 is CH;


R^d and R^e are each independently selected from the group consisting of H and methyl;

R^k is selected from the group consisting of H, CN, methyl and ethyl;




5 R^2 is selected from the group consisting of H, Cl, and F.

In some embodiments, the compound or a pharmaceutically acceptable salt thereof, is selected from the group consisting of:




10


The invention also provides a compound, or a pharmaceutically acceptable salt thereof, having the formula:



15 The invention also provides a compound, or a pharmaceutically acceptable salt thereof, having the formula:



The invention also provides a compound, or a pharmaceutically acceptable salt thereof, having the formula:



The invention also provides a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, and a pharmaceutically-acceptable carrier. In some embodiments, the pharmaceutical composition further comprises one or more other therapeutic agents. In some embodiments, the one or more other therapeutic agent is useful for treating a gastrointestinal inflammatory disease, an inflammatory disease of the skin, an inflammatory disease of the lungs or an inflammatory disease of the eye. In some embodiments, the one or more other therapeutic agent is useful for treating a gastrointestinal inflammatory disease. In some embodiments the gastrointestinal inflammatory disease is ulcerative colitis. In some embodiments the gastrointestinal inflammatory disease is Crohn's disease.

Furthermore, some compounds may sometimes exist in tautomeric forms. It will be understood that although structures are shown, or named, in a particular form, the invention also includes the tautomer thereof.

The compounds of the invention may contain one or more chiral centers and therefore, such compounds (and intermediates thereof) can exist as racemic mixtures; pure stereoisomers (i.e., enantiomers or diastereomers); stereoisomer-enriched mixtures and the like. Chiral compounds shown or named herein without a defined stereochemistry at a chiral center are intended to include any or all possible stereoisomer variations at the undefined stereocenter unless otherwise indicated. The depiction or naming of a particular stereoisomer means the indicated stereocenter has the designated stereochemistry with the understanding that minor amounts of other stereoisomers may also be present unless otherwise indicated, provided that the utility of the depicted or named compound is not eliminated by the presence of another stereoisomer.

This invention also includes isotopically-labeled compounds of the disclosure, for example isotopically-labeled compounds of formula (I), (II), (B), (C), compound 1, compound 3, compound 4, i.e., compounds of the disclosure and compounds of formula (I), (II), (B), (C), compound 1, compound 3, compound 4 where an atom has been replaced or enriched with an atom having the same atomic number but an atomic mass different from the atomic mass that predominates in nature. Examples of isotopes that

may be incorporated into a compounds of the disclosure and a compound of formula (I), (II), (B), (C), compound 1, compound 3, compound 4, include, but are not limited to, ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{35}S , and ^{18}F . Of particular interest are compounds of the disclosure and compounds of formula (I), (II), (B), (C), compound 1, compound 3, and compound 4, enriched in tritium or carbon-14, which compounds can be used, for example, in tissue distribution studies. Also of particular interest are compounds of the disclosure and compounds of formula (I), (II), (B), (C), compound 1, compound 3, compound 4 enriched in deuterium especially at a site of metabolism, which compounds are expected to have greater metabolic stability. Additionally, of particular interest are compounds of the disclosure and compounds of formula (I), (II), (B), (C), compound 1, compound 3, compound 4, enriched in a positron emitting isotope, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , which compounds can be used, for example, in Positron Emission Tomography (PET) studies.

Definitions

When describing this invention including its various aspects and embodiments, the following terms have the following meanings, unless otherwise indicated.

The term "alkyl" means a monovalent saturated hydrocarbon group which may be linear or branched or combinations thereof. Unless otherwise defined, such alkyl groups typically contain from 1 to 10 carbon atoms. Representative alkyl groups include, by way of example, methyl (Me), ethyl (Et), n-propyl (n-Pr) or (nPr), isopropyl (i-Pr) or (iPr), n-butyl (n-Bu) or (nBu), sec-butyl, isobutyl, tert-butyl (t-Bu) or (tBu), n-pentyl, n-hexyl, 2,2-dimethylpropyl, 2-methylbutyl, 3-methylbutyl, 2-ethylbutyl, 2,2-dimethylpentyl, 2-propylpentyl, and the like.

The term "haloalkyl" refers to an alkyl group, as defined above, that is substituted by one or more halogen, e.g., trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1,2 difluoroethyl, 3 bromo 2 fluoropropyl, 1,2 dibromoethyl, and the like.

When a specific number of carbon atoms are intended for a particular term, the number of carbon atoms is shown preceding the term. For example, the term "C₁₋₃alkyl" means an alkyl group having from 1 to 3 carbon atoms wherein the carbon atoms are in any chemically-acceptable configuration, including linear or branched configurations.

The term "alkoxy" means the monovalent group -O-alkyl, where alkyl is defined as above. Representative alkoxy groups include, by way of example, methoxy, ethoxy, propoxy, butoxy, and the like.

The term "cycloalkyl" means a monovalent saturated carbocyclic group which may be monocyclic or multicyclic. Unless otherwise defined, such cycloalkyl groups typically contain from 3 to 10 carbon atoms. Representative cycloalkyl groups include, by way of example, cyclopropyl (cPr), cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, adamantyl, and the like.

The term "heterocycle", "heterocyclic", or "heterocyclic ring" means a saturated or partially unsaturated cyclic non-aromatic group, having from 3 to 10 total ring atoms, wherein the ring contains from 2 to 9 carbon ring atoms and from 1 to 4 ring heteroatoms selected from nitrogen, oxygen, and sulfur. Heterocyclic groups may be monocyclic or multicyclic (i.e., fused, spiro or bridged). When the heterocyclic group is multicyclic, at least one but not necessarily all of the cyclic groups contains a heteroatom.

Representative heterocyclic groups include, by way of example, pyrrolidinyl, piperidinyl, piperazinyl, imidazolidinyl, morpholinyl, thiomorpholyl, indolin-3-yl, 2-imidazoliny, tetrahydropyranyl, 1,2,3,4-tetrahydroisoquinolin-2-yl, quinuclidinyl, 7-azanorbornanyl, nortropanyl, and the like, where the point of attachment is at any available carbon or nitrogen ring atom. Where the context makes the point of attachment of the heterocyclic group evident, such groups may alternatively be referred to as a non-valent species, i.e. pyrrolidine, piperidine, piperazine, imidazole, tetrahydropyran etc.

The term "therapeutically effective amount" means an amount sufficient to effect treatment when administered to a patient in need of treatment.

The term "treatment" as used herein means the treatment of a disease, disorder, or medical condition (such as a gastrointestinal inflammatory disease), in a patient, such as a mammal (particularly a human) which includes one or more of the following:

(a) preventing the disease, disorder, or medical condition from occurring, i.e., preventing the reoccurrence of the disease or medical condition or prophylactic treatment of a patient that is pre-disposed to the disease or medical condition;

(b) ameliorating the disease, disorder, or medical condition, i.e., eliminating or causing regression of the disease, disorder, or medical condition in a patient, including counteracting the effects of other therapeutic agents;

(c) suppressing the disease, disorder, or medical condition, i.e., slowing or arresting the development of the disease, disorder, or medical condition in a patient;

(d) alleviating the symptoms of the disease, disorder, or medical condition in a patient.

The term “pharmaceutically acceptable salt” means a salt that is acceptable for administration to a patient or a mammal, such as a human (e.g., salts having acceptable mammalian safety for a given dosage regime). Representative pharmaceutically acceptable salts include salts of acetic, ascorbic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, edisylic, fumaric, gentisic, gluconic, glucuronic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, lactobionic, maleic, malic, mandelic, methanesulfonic, mucic, naphthalenesulfonic, naphthalene-1,5-disulfonic, naphthalene-2,6-disulfonic, nicotinic, nitric, orotic, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic and xinafoic acid, and the like.

The term “salt thereof” means a compound formed when the hydrogen of an acid is replaced by a cation, such as a metal cation or an organic cation and the like. For example, the cation can be a protonated form of a compound of formula (I), i.e. a form where one or more amino groups have been protonated by an acid. Typically, the salt is a pharmaceutically acceptable salt, although this is not required for salts of intermediate compounds that are not intended for administration to a patient.

The term "amino-protecting group" means a protecting group suitable for preventing undesired reactions at an amino nitrogen. Representative amino-protecting groups include, but are not limited to, formyl; acyl groups, for example alkanoyl groups, such as acetyl and tri-fluoroacetyl; alkoxycarbonyl groups, such as *tert*-butoxycarbonyl (Boc); arylmethoxycarbonyl groups, such as benzyloxycarbonyl (Cbz) and 9-fluorenylmethoxycarbonyl (Fmoc); arylmethyl groups, such as benzyl (Bn), trityl (Tr), and 1,1-di-(4'-methoxyphenyl)methyl; silyl groups, such as trimethylsilyl (TMS), triisopropylsilyl (TIPS), *tert*-butyldimethylsilyl (TBS or TBDMS), [2-(trimethylsilyl)-ethoxy]methyl (SEM); and the like. Numerous protecting groups, and their introduction and removal, are described in T. W. Greene and P. G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York


General Synthetic Procedures

Compounds of this disclosure, and intermediates thereof, can be prepared according to the following general methods and procedures using commercially-available or routinely-prepared starting materials and reagents. The substituents and variables (e.g., A, X¹, X², R¹, R², R³, R^a, R^b etc.) used in the following schemes have the same meanings as those defined elsewhere herein unless otherwise indicated. Additionally, compounds having an acidic or basic atom or functional group may be used or may be produced as a salt unless otherwise indicated (in some cases, the use of a salt in a particular reaction will

require conversion of the salt to a non-salt form, e.g., a free base, using routine procedures before conducting the reaction).


Although a particular embodiment of the present invention may be shown or described in the following procedures, those skilled in the art will recognize that other
 5 embodiments or aspects of the present invention can also be prepared using such procedures or by using other methods, reagents, and starting materials known to those skilled in the art. In particular, it will be appreciated that compounds of the disclosure may be prepared by a variety of process routes in which reactants are combined in
 different orders to provide different intermediates en route to producing final products.

10 A general method of preparing final compounds of the disclosure wherein L is selected from:




is illustrated in Scheme 1.

Scheme 1




Starting material P1, where R^x and R^y are halogens which may be the same or different, is protected with a protected group PG such as tetrahydropyran to give P2. P2 is then reacted with P3 to give P4.

P3 may be:




where R^z is a second protecting group, for example Boc. In this case, P3 is deprotonated with a base such as NaH and reacted with P2 to give P4.

Alternatively, P3 may be:



where R^z is a second protecting group, for example Boc. In this case, P3 is reacted with P2 under Buchwald coupling conditions such as in the presence of Pd(0) and a base to give P4. Alternatively, P3 is reacted with P2 in presence of a base such as DIPEA to give P4.

Alternatively, P3 may be:



where R^Z is a second protecting group, for example Boc. In this case, P3 is reacted with P2 in presence of Pd(0), 9-BBN and a base to form P4.


5 P4 is coupled with boronic acid P5 (Suzuki coupling) in presence of Pd(0) and a base to give P6. P6 is deprotected to give P7 (when PG is tetrahydropyran and R^Z is Boc, simultaneous deprotection of the amines occur in presence of a strong acid such as TFA or HCl). Finally, P7 is derivatized into an amide by amide coupling (reaction with an acid in presence of a coupling agent such as HATU or hydroxybenzotriazole (HOBT)) or
10 reaction with an acyl chloride in presence of a base such as Hunig's base.

In this reaction scheme, the order of the reactions may be modified. For example the Suzuki coupling may be conducted before the introduction of the portion containing the A ring. This can for example be the case when the portion containing the A ring is introduced through a Buchwald coupling.

15 In this reaction scheme, protection of one or both of the amino groups is optional. The same synthetic scheme may be used without amino protection for one or both amino groups but may provide lower yields.


The sulfonyl linker can be obtained by oxidizing the corresponding sulfide, for example with oxone and basic alumina.

20 Accordingly, in a method aspect, the invention provides a method for preparing a compound of formula (I) or a pharmaceutically acceptable salt thereof,



the method comprising:

25 reacting a compound of formula (III):



(III)

with

- (i) Cl-R¹, or
 5 (ii) HO-R¹

wherein R¹, R², R³, R⁴, X¹, X², L and A are as defined above, and


optionally forming a pharmaceutically-acceptable salt

to provide a compound of formula (I), or a pharmaceutically acceptable salt thereof.

In separate and distinct aspects, the invention provides a compound of formula
 10 (III) wherein the variables take any of the values described above.

Crystalline Forms

In one aspect, the invention provides a crystalline form of the compound of formula:



15

Form 1

Crystalline Form 1 of the invention is a crystalline anhydrous free form of
 20 compound 3. In one aspect, Form 1 is characterized by a powder X-ray diffraction (PXRD) pattern having significant diffraction peaks, among other peaks, at 2θ values of 5.65±0.20, 14.22±0.20, 15.16±0.20, and 19.31±0.20. Form 1 may be further characterized by a PXRD pattern having additional diffraction peaks at 2θ values of 7.12±0.20, 10.02±0.20, 11.16±0.20, 17.06±0.20, and 24.43±0.20. Form 1 may be further
 25 characterized by a PXRD pattern having two or more additional diffraction peaks,

including three or more and four or more additional diffraction peaks at 2θ values selected from 13.10 ± 0.20 , 14.82 ± 0.20 , 16.55 ± 0.20 , 20.08 ± 0.20 , 21.08 ± 0.20 , 21.65 ± 0.20 , 22.51 ± 0.20 , 22.98 ± 0.20 , 25.02 ± 0.20 , 25.72 ± 0.20 , 26.80 ± 0.20 , 27.06 ± 0.20 , 28.31 ± 0.20 , 30.08 ± 0.20 , 30.31 ± 0.20 and 32.08 ± 0.20 . Form 1 is characterized by a PXRD pattern
5 having three, four, five, or six diffraction peaks at 2θ values selected from 5.65 ± 0.20 , 7.12 ± 0.20 , 10.02 ± 0.20 , 11.16 ± 0.20 , 14.22 ± 0.20 , 15.16 ± 0.20 , 17.06 ± 0.20 , 19.31 ± 0.20 and 24.43 ± 0.20 .

As is well known in the field of powder X-ray diffraction, peak positions of PXRD pattern are relatively less sensitive to experimental details, such as details of
10 sample preparation and instrument geometry, than are the relative peak heights. Thus, in one aspect, the crystalline Form 1 is characterized by a powder x-ray diffraction pattern in which the peak positions are substantially in accordance with those shown in Figure 1.

In another aspect, crystalline Form 1 is characterized by its behavior when exposed to high temperature. As demonstrated in Figure 2, the differential scanning
15 calorimetry (DSC) trace recorded at a heating rate of $10\text{ }^{\circ}\text{C}$ per minute exhibits a peak in endothermic heat flow, identified as a melt transition, with an onset at about $154.9\text{ }^{\circ}\text{C}$ and a peak at about $162.9\text{ }^{\circ}\text{C}$. Melting was followed immediately by decomposition.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of $10\text{ }^{\circ}\text{C}$ per minute which shows a maximum in endothermic
20 heat flow with a peak at about $162.9\text{ }^{\circ}\text{C}$. The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of $10\text{ }^{\circ}\text{C}$ per minute which shows a maximum in endothermic heat flow with a peak at $162.9 \pm 3\text{ }^{\circ}\text{C}$.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of $10\text{ }^{\circ}\text{C}$ per minute which shows a maximum in endothermic
25 heat flow at a temperature between about $154.9\text{ }^{\circ}\text{C}$ and about $171\text{ }^{\circ}\text{C}$, or between $158\text{ }^{\circ}\text{C}$ and $167\text{ }^{\circ}\text{C}$.

A representative TGA trace of the Form 1 crystalline free form is shown in Figure 3. The thermal gravimetric analysis (TGA) trace shows a small weight loss of about
0.14% at $100\text{ }^{\circ}\text{C}$. The compound decomposes at an onset temperature of about $175\text{ }^{\circ}\text{C}$.

30 A representative DMS trace for the Form 1 crystalline free form is shown in Figure 4. Form 1 demonstrated about 1.62 % weight gain in the humidity range of 5 % to 90 % relative humidity. Form 1 is considered to be slightly hygroscopic.

Form 1 may be prepared by dissolving compound 3 as an amorphous form in ethanol followed by stirring at a temperature of between about 20 °C and about 25 °C followed by filtration and drying to give Form 1. Optionally, the solid can be washed with ethanol before drying.

- 5 Form 1 may be prepared by adding acetone to the compound in an amorphous form and stirring at a temperature of between about 20 °C and about 25 °C followed by the addition of seeds. The resulting slurry is filtered and dried to give Form 1.

Form 2

- Crystalline Form 2 of the invention is a crystalline hydrate free form of compound
- 10 3. In one aspect, Form 2 is characterized by a powder X-ray diffraction (PXRD) pattern having significant diffraction peaks, among other peaks, at 2θ values of 6.90 ± 0.20 , 9.15 ± 0.20 , 10.00 ± 0.20 , and 18.31 ± 0.20 . Form 2 may be further characterized by a PXRD pattern having additional diffraction peaks at 2θ values of 11.18 ± 0.20 , 15.51 ± 0.20 , and 20.90 ± 0.20 . Form 2 may be further characterized by a PXRD pattern having two or more
- 15 additional diffraction peaks, including three or more and four or more additional diffraction peaks at 2θ values selected from 12.76 ± 0.20 , 13.33 ± 0.20 , 13.82 ± 0.20 , 14.43 ± 0.20 , 16.04 ± 0.20 , 17.00 ± 0.20 , 17.90 ± 0.20 , 22.06 ± 0.20 , 22.51 ± 0.20 , 25.00 ± 0.20 , 26.92 ± 0.20 , 27.26 ± 0.20 , 27.61 ± 0.20 , 29.37 ± 0.20 , 30.53 ± 0.20 , and 30.92 ± 0.20 . Form 2 is characterized by a PXRD pattern having three, four, five, or six diffraction peaks at
- 20 2θ values selected from 6.90 ± 0.20 , 9.15 ± 0.20 , 10.00 ± 0.20 , 11.18 ± 0.20 , 15.51 ± 0.20 , 18.31 ± 0.20 , and 20.90 ± 0.20 .

- As is well known in the field of powder X-ray diffraction, peak positions of PXRD pattern are relatively less sensitive to experimental details, such as details of sample preparation and instrument geometry, than are the relative peak heights. Thus, in
- 25 one aspect, the crystalline Form 2 is characterized by a powder x-ray diffraction pattern in which the peak positions are substantially in accordance with those shown in Figure 5.

- In another aspect, crystalline Form 2 is characterized by its behavior when exposed to high temperature. As demonstrated in Figure 6, the differential scanning calorimetry (DSC) trace recorded at a heating rate of 10 °C per minute exhibits a
- 30 desolvation endotherm with an onset at about 52.7 °C and a peak at about 84.4 °C, and a

melting endotherm with an onset at about 160.0 °C and a peak at about 167.6 °C. Melting was followed immediately by decomposition.

A representative TGA trace of the Form 2 crystalline free form of the invention is shown in Figure 7. The thermal gravimetric analysis (TGA) trace of Figure 7 shows a weight loss of about 6.73% at 75 °C. The compound desolvates at an onset temperature of about 25 °C. The compound decomposes at an onset temperature of about 185 °C.

A representative DMS trace for the Form 2 crystalline free form of the invention is shown in Figure 8. Form 2 converts to a hydrate (Form 2b) at RH above 65 %. The dehydration occurs at RH below 15 %. The total moisture uptake between 5%-90% RH is 7.99 %.

Form 2 may be prepared by dissolving compound 3 in amorphous form in methanol followed by the addition of an anti-solvent such as water, in a ratio of about 1:2 methanol:water. Optionally, the mixture is sonicated. The mixture is then stirred at a temperature of between about 20 °C and about 25 °C for about 12 to 24 hours. Form 2 is then isolated by filtration and drying. Optionally, the solid can be washed with methanol.

Form 2 can also be prepared by dissolution in ethanol and water or methanol and water by complete dissolution in about 10 volumes of alcohol followed by slow addition of about 8-10 volumes of water until cloud point. Seeds of Form 2 are added to form a slurry slowly over time. More water is added slowly (about 10 volumes) to provide a solid which can be filtered and dried to give Form 2.


Form 2b

Crystalline Form 2b of the invention is a crystalline dehydrated free form of compound 3. In one aspect, Form 2b is characterized by a powder X-ray diffraction (PXRD) pattern having significant diffraction peaks, among other peaks, at 2θ values of 7.61±0.20, 16.76±0.20, 17.90±0.20, and 20.67±0.20. Form 2b may be further characterized by a PXRD pattern having additional diffraction peaks at 2θ values of 10.33±0.20, 11.25±0.20, 12.71±0.20, 15.88±0.20. Form 2b may be further characterized by a PXRD pattern having two or more additional diffraction peaks, including three or more and four or more additional diffraction peaks at 2θ values selected from 13.23±0.20, 13.66±0.20, 13.90±0.20, 15.02±0.20, 15.27±0.20, 16.33±0.20, 18.26±0.20, 21.37±0.20, 21.92±0.20, 22.31±0.20, 22.90±0.20, 23.22±0.20, 23.61±0.20, 24.74±0.20, 25.78±0.20, 26.23±0.20, 26.73±0.20, 27.57±0.20, 29.10±0.20, 29.39±0.20, 30.72±0.20, 30.94±0.20, 31.69±0.20, 32.06±0.20, 33.76±0.20, and 34.35±0.20. Form 2b is characterized by a

PXRD pattern having three, four, five, or six diffraction peaks at 2θ values selected from 7.61 ± 0.20 , 10.33 ± 0.20 , 11.25 ± 0.20 , 12.71 ± 0.20 , 15.88 ± 0.20 , 16.76 ± 0.20 , 17.90 ± 0.20 , and 20.67 ± 0.20 .

As is well known in the field of powder X-ray diffraction, peak positions of
 5 PXRD pattern are relatively less sensitive to experimental details, such as details of sample preparation and instrument geometry, than are the relative peak heights. Thus, in one aspect, the crystalline Form 2b is characterized by a powder x-ray diffraction pattern in which the peak positions are substantially in accordance with those shown in Figure 17.

10 In another aspect, the invention provides a crystalline form of the compound of formula:



Form 3

Crystalline Form 3 of the invention is a crystalline anhydrous free form of
 15 compound 1. In one aspect, Form 3 is characterized by a powder X-ray diffraction (PXRD) pattern having significant diffraction peaks, among other peaks, at 2θ values of 9.67 ± 0.20 , 11.61 ± 0.20 , 17.61 ± 0.20 , 18.88 ± 0.20 , and 23.33 ± 0.20 . Form 3 may be further characterized by a PXRD pattern having additional diffraction peaks at 2θ values of 4.82 ± 0.20 , 15.69 ± 0.20 , and 16.19 ± 0.20 . Form 3 may be further characterized by a PXRD
 20 pattern having two or more additional diffraction peaks, including three or more and four or more additional diffraction peaks at 2θ values selected from 11.92 ± 0.20 , 12.98 ± 0.20 , 13.23 ± 0.20 , 16.45 ± 0.20 , 16.67 ± 0.20 , 19.39 ± 0.20 , 19.96 ± 0.20 , 20.14 ± 0.20 , 22.14 ± 0.20 , 23.84 ± 0.20 , 24.06 ± 0.20 , 24.29 ± 0.20 , 25.31 ± 0.20 , 25.63 ± 0.20 , 27.06 ± 0.20 , 27.31 ± 0.20 , 30.10 ± 0.20 , and 30.53 ± 0.20 . Form 3 is characterized by a PXRD pattern having three,
 25 four, five, or six diffraction peaks at 2θ values selected from 4.82 ± 0.20 , 9.67 ± 0.20 , 11.61 ± 0.20 , 15.69 ± 0.20 , 16.19 ± 0.20 , 17.61 ± 0.20 , 18.88 ± 0.20 , and 23.33 ± 0.20 .

As is well known in the field of powder X-ray diffraction, peak positions of PXRD pattern are relatively less sensitive to experimental details, such as details of sample preparation and instrument geometry, than are the relative peak heights. Thus, in

one aspect, the crystalline Form 3 is characterized by a powder x-ray diffraction pattern in which the peak positions are substantially in accordance with those shown in Figure 9.

In another aspect, crystalline Form 3 is characterized by its behavior when exposed to high temperature. As demonstrated in Figure 10, the differential scanning calorimetry (DSC) trace recorded at a heating rate of 10 °C per minute exhibits a peak in
5 endothermic heat flow, identified as a melt transition, with an onset at about 197.7 °C and a peak at about 201.3 °C. Melting was followed immediately by decomposition.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic
10 heat flow with a peak at 201.3 °C \pm 2 °C.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic heat flow at a temperature between 198 °C and 204 °C.

A representative TGA trace of the Form 3 crystalline free form of the invention is
15 shown in Figure 11. The thermal gravimetric analysis (TGA) trace of Figure 11 shows no significant weight loss at temperatures below the onset of decomposition at about 195 °C.

A representative DMS trace for the Form 3 crystalline free form of the invention is shown in Figure 12. Form 3 demonstrated about 0.33 % weight gain in the humidity range of 5 % to 90 % relative humidity. Form 3 is considered to be non-hygroscopic.

20 Form 3 may be prepared by suspending compound 1 in an amorphous form in a 1:1 mixture of acetonitrile and isopropanol. The resulting suspension is stirred for about 1 day at about 50 °C, filtered, optionally washed with a 1:1 mixture of acetonitrile and isopropanol and dried for a few hours to provide Form 3.

Form 3 may be prepared by dissolving compound 1 as an amorphous free base in
25 IPA at a temperature of between about 20 °C and about 25 °C. An equal amount of acetonitrile is added. More compound can be added until a saturated solution is formed. Seeds are added and the mixture is stirred overnight. The developing white slurry formed is filtered and dried to yield Form 3.

Form 4

30 Crystalline Form 4 of the invention is a crystalline hydrate free form of compound 1. In one aspect, Form 4 is characterized by a powder X-ray diffraction (PXRD) pattern having significant diffraction peaks, among other peaks, at 2θ values of 6.26 ± 0.20 , 16.55 ± 0.20 , 16.94 ± 0.20 , 18.33 ± 0.20 , 23.61 ± 0.20 , and 24.24 ± 0.20 . Form 4 may be

further characterized by a PXRD pattern having additional diffraction peaks at 2θ values of 11.86 ± 0.20 , 12.51 ± 0.20 , 13.16 ± 0.20 , and 14.98 ± 0.20 . Form 4 may be further characterized by a PXRD pattern having two or more additional diffraction peaks, including three or more and four or more additional diffraction peaks at 2θ values
5 selected from 17.61 ± 0.20 , 18.78 ± 0.20 , 19.39 ± 0.20 , 19.57 ± 0.20 , 19.84 ± 0.20 , 21.45 ± 0.20 , 21.82 ± 0.20 , 22.57 ± 0.20 , 24.67 ± 0.20 , 25.10 ± 0.20 , 25.39 ± 0.20 , 27.19 ± 0.20 , 27.39 ± 0.20 , 28.55 ± 0.20 , and 31.51 ± 0.20 . Form 4 is characterized by a PXRD pattern having three, four, five, or six diffraction peaks at 2θ values selected from 6.26 ± 0.20 , 11.86 ± 0.20 , 12.51 ± 0.20 , 13.16 ± 0.20 , 14.98 ± 0.20 , 16.55 ± 0.20 , 16.94 ± 0.20 , 18.33 ± 0.20 , 23.61 ± 0.20 ,
10 and 24.24 ± 0.20 .

As is well known in the field of powder X-ray diffraction, peak positions of PXRD pattern are relatively less sensitive to experimental details, such as details of sample preparation and instrument geometry, than are the relative peak heights. Thus, in one aspect, the crystalline Form 4 is characterized by a powder x-ray diffraction pattern in
15 which the peak positions are substantially in accordance with those shown in Figure 13.

In another aspect, crystalline Form 4 is characterized by its behavior when exposed to high temperature. As demonstrated in Figure 14, the differential scanning calorimetry (DSC) trace recorded at a heating rate of $10\text{ }^{\circ}\text{C}$ per minute exhibits a desolvation endotherm with an onset at about $60.9\text{ }^{\circ}\text{C}$ and a peak at about $103.6\text{ }^{\circ}\text{C}$, and a
20 melting endotherm characterized by an onset at about $167.3\text{ }^{\circ}\text{C}$. The compound decomposes at melting and the melting endotherm and the decomposition exotherm overlap.

A representative TGA trace of the Form 4 crystalline free form of the invention is shown in Figure 15. The thermal gravimetric analysis (TGA) trace of Figure 15 shows a
25 weight loss of about 3.54% at $100\text{ }^{\circ}\text{C}$. The compound desolvates at an onset temperature of about $50\text{ }^{\circ}\text{C}$. The compound decomposes at an onset temperature of about $165\text{ }^{\circ}\text{C}$.

A representative DMS trace for the Form 4 crystalline free form of the invention is shown in Figure 16. Form 4 demonstrated about 5.01 % weight gain in the humidity range of 5 % to 90 % relative humidity. Form 4 is considered to be moderately
30 hygroscopic.

Form 4 may be prepared by suspending compound 1 in water. The resulting suspension is stirred for about 1 to 2 days at about $50\text{ }^{\circ}\text{C}$, filtered, optionally washed with

water, and dried at a temperature of between about 20 °C and about 25 °C for about 2-6 hours to provide Form 4.

Alternatively, Form 4 may be prepared by dissolving compound 1 in ethanol and water or methanol and water by complete dissolution in about 10 volumes of alcohol followed by slow addition of about 8-10 volumes of water until cloud point. Seeds of Form 4 are added and a resulting slurry develops slowly over time. Then more water is added slowly (about 10 volumes) and the solid is filtered and dried to give Form 4.

Pharmaceutical Compositions

The compounds of the disclosure and pharmaceutically-acceptable salts thereof are typically used in the form of a pharmaceutical composition or formulation. Such pharmaceutical compositions may be administered to a patient by any acceptable route of administration including, but not limited to, oral, topical (including transdermal), rectal, nasal, inhaled, and parenteral modes of administration.

Accordingly, in one of its composition aspects, the invention is directed to a pharmaceutical composition comprising a pharmaceutically-acceptable carrier or excipient and a compound of formula (I), (II), (B), (C), compound 1, compound 3, or compound 4 or a pharmaceutically-acceptable salt thereof. Optionally, such pharmaceutical compositions may contain other therapeutic and/or formulating agents if desired. When discussing compositions and uses thereof, the "compound of the invention" or "compound of the disclosure" may also be referred to herein as the "active agent". As used herein, the term "compound(s) of the disclosure" is intended to include all compounds encompassed by formula (I), (II), (B), (C), (Ia), (IIa), (Ib), (IIb), (Ic), (IIc), (Id), (IId), (Ie), (IIe), (If), (IIIf), (IIIf), (IVf), (Vf), (VIf), (VIIIf), and (VIIIf), and pharmaceutically-acceptable salts thereof.

The pharmaceutical compositions of the disclosure typically contain a therapeutically effective amount of a compound of the disclosure. Those skilled in the art will recognize, however, that a pharmaceutical composition may contain more than a therapeutically effective amount, i.e., bulk compositions, or less than a therapeutically effective amount, i.e., individual unit doses designed for multiple administration to achieve a therapeutically effective amount.

Typically, such pharmaceutical compositions will contain from about 0.1 to about 95% by weight of the active agent; including from about 5 to about 70% by weight of the active agent.

Any conventional carrier or excipient may be used in the pharmaceutical compositions of the invention. The choice of a particular carrier or excipient, or combinations of carriers or excipients, will depend on the mode of administration being used to treat a particular patient or type of medical condition or disease state. In this
5 regard, the preparation of a suitable pharmaceutical composition for a particular mode of administration is well within the scope of those skilled in the pharmaceutical arts. Additionally, the carriers or excipients used in the pharmaceutical compositions of this invention are commercially-available. By way of further illustration, conventional formulation techniques are described in Remington: The Science and Practice of
10 Pharmacy, 20th Edition, Lippincott Williams & White, Baltimore, Maryland (2000); and H.C. Ansel et al., Pharmaceutical Dosage Forms and Drug Delivery Systems, 7th Edition, Lippincott Williams & White, Baltimore, Maryland (1999).

Representative examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, the following: sugars, such as lactose,
15 glucose and sucrose; starches, such as corn starch and potato starch; cellulose, such as microcrystalline cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene
20 glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances employed in pharmaceutical compositions.

25 Pharmaceutical compositions are typically prepared by thoroughly and intimately mixing or blending the active agent with a pharmaceutically-acceptable carrier and one or more optional ingredients. The resulting uniformly blended mixture can then be shaped or loaded into tablets, capsules, pills and the like using conventional procedures and equipment.

30 The pharmaceutical compositions of the disclosure are preferably packaged in a unit dosage form. The term "unit dosage form" refers to a physically discrete unit suitable for dosing a patient, i.e., each unit containing a predetermined quantity of active agent calculated to produce the desired therapeutic effect either alone or in combination

with one or more additional units. For example, such unit dosage forms may be capsules, tablets, pills, and the like, or unit packages suitable for parenteral administration.

In one embodiment, the pharmaceutical compositions of the disclosure are suitable for oral administration. Suitable pharmaceutical compositions for oral administration may be in the form of capsules, tablets, pills, lozenges, cachets, dragees, powders, granules; or as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil liquid emulsion; or as an elixir or syrup; and the like; each containing a predetermined amount of a compound of the present disclosure as an active ingredient.

When intended for oral administration in a solid dosage form (i.e., as capsules, tablets, pills and the like), the pharmaceutical compositions of the disclosure will typically comprise the active agent and one or more pharmaceutically-acceptable carriers. Optionally, such solid dosage forms may comprise: fillers or extenders, such as starches, microcrystalline cellulose, lactose, dicalcium phosphate, sucrose, glucose, mannitol, and/or silicic acid; binders, such as carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; humectants, such as glycerol; disintegrating agents, such as croscarmellose sodium, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and/or sodium carbonate; solution retarding agents, such as paraffin; absorption accelerators, such as quaternary ammonium compounds; wetting agents, such as cetyl alcohol and/or glycerol monostearate; absorbents, such as kaolin and/or bentonite clay; lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and/or mixtures thereof; coloring agents; and buffering agents.

Release agents, wetting agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the pharmaceutical compositions of the disclosure. Examples of pharmaceutically-acceptable antioxidants include: water-soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfate, sodium sulfite and the like; oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, lecithin, propyl gallate, alpha-tocopherol, and the like; and metal-chelating agents, such as citric acid, ethylenediamine tetraacetic acid, sorbitol, tartaric acid, phosphoric acid, and the like. Coating agents for tablets, capsules, pills and like, include those used for enteric coatings, such as cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropyl methylcellulose phthalate, methacrylic acid, methacrylic

acid ester copolymers, cellulose acetate trimellitate, carboxymethyl ethyl cellulose, hydroxypropyl methyl cellulose acetate succinate, and the like.

Pharmaceutical compositions of the disclosure may also be formulated to provide slow or controlled release of the active agent using, by way of example, hydroxypropyl methylcellulose in varying proportions; or other polymer matrices, liposomes and/or microspheres. In addition, the pharmaceutical compositions of the disclosure may optionally contain opacifying agents and may be formulated so that they release the active ingredient only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active agent can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Suitable liquid dosage forms for oral administration include, by way of illustration, pharmaceutically-acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. Liquid dosage forms typically comprise the active agent and an inert diluent, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (esp., cottonseed, groundnut, corn, germ, olive, castor and sesame oils), oleic acid, glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Alternatively, certain liquid formulations can be converted, for example, by spray drying, to a powder, which is used to prepare solid dosage forms by conventional procedures.

Suspensions, in addition to the active ingredient, may contain suspending agents such as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

The compounds of this disclosure, or a pharmaceutically acceptable salt thereof, can also be administered parenterally (e.g. by intravenous, subcutaneous, intramuscular or intraperitoneal injection). For parenteral administration, the active agent is typically admixed with a suitable vehicle for parenteral administration including, by way of example, sterile aqueous solutions, saline, low molecular weight alcohols such as propylene glycol, polyethylene glycol, vegetable oils, gelatin, fatty acid esters such as ethyl oleate, and the like. Parenteral formulations may also contain one or more anti-oxidants, solubilizers, stabilizers, preservatives, wetting agents, emulsifiers, buffering

agents, or dispersing agents. These formulations may be rendered sterile by use of a sterile injectable medium, a sterilizing agent, filtration, irradiation, or heat.

Alternatively, the pharmaceutical compositions of the disclosure are formulated for administration by inhalation. Suitable pharmaceutical compositions for administration
5 by inhalation will typically be in the form of an aerosol or a powder. Such compositions are generally administered using well-known delivery devices, such as a metered-dose inhaler, a dry powder inhaler, a nebulizer or a similar delivery device.

When administered by inhalation using a pressurized container, the pharmaceutical compositions of the disclosure will typically comprise the active
10 ingredient and a suitable propellant, such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. Additionally, the pharmaceutical composition may be in the form of a capsule or cartridge (made, for example, from gelatin) comprising a compound of the disclosure and a powder suitable for use in a powder inhaler. Suitable powder bases include, by way of example,
15 lactose or starch.

The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may also be formulated for topical administration to the skin as an ointment or cream. Ointment formulations are semisolid preparations having a base of an oily or greasy material that is typically clear. Suitable oily materials for use in ointment formulations
20 include petrolatum (petroleum jelly), beeswax, cocoa butter, shea butter, and cetyl alcohol. Ointments may optionally additionally include emollients and penetration enhancers, if desired.

Cream formulations may be prepared as emulsions comprising an oil phase and aqueous phase, typically including purified water. Components of cream formulations
25 may include: oil bases, such as petrolatum, mineral oils, vegetable and animal oils, and triglycerides; cream bases, such as lanolin alcohols, stearic acid, and cetostearyl alcohol; a gel base, such as polyvinyl alcohol; solvents, such as, propylene glycol and polyethylene glycol; emulsifiers, such as polysorbates, stearates, such as glyceryl stearate, octylhydroxystearate, polyoxyl stearate, PEG stearyl ethers, isopropyl palmitate, and
30 sorbitan monostearate; stabilizers, such as polysaccharides and sodium sulfite; emollients (i.e. moisturizers), such as medium chain triglycerides, isopropyl myristate, and dimethicone; stiffening agents, such as cetyl alcohol and stearyl alcohol; antimicrobial agents, such as methylparaben, propylparaben, phenoxyethanol, sorbic acid, diazolidinyl urea, and butylated hydroxyanisole; penetration enhancers, such as *N*-methylpyrrolidone,

propylene glycol, polyethylene glycol monolaurate, and the like; and chelating agents, such as edetate disodium.

The following non-limiting examples illustrate representative pharmaceutical compositions of the present invention.

5 *Tablet oral solid dosage form*

A compound of the disclosure or a pharmaceutically-acceptable salt thereof is dry blended with microcrystalline cellulose, polyvinyl pyrrolidone, and crosscarmellose sodium in a ratio of 4:5:1:1 and compressed into tablets to provide a unit dosage of, for example, 5 mg, 20 mg or 40 mg active agent per tablet.

10 *Capsule oral solid dosage form*

A compound of the disclosure or a pharmaceutically-acceptable salt thereof is combined with microcrystalline cellulose, polyvinyl pyrrolidone, and crosscarmellose sodium in a ratio of 4:5:1:1 by wet granulation and loaded into gelatin or hydroxypropyl methylcellulose capsules to provide a unit dosage of, for example, 5 mg, 20 mg or 40 mg
15 active agent per capsule.

Liquid formulation

A liquid formulation comprising a compound of the disclosure (0.1 %), water (98.9 %) and ascorbic acid (1.0 %) is formed by adding a compound of the disclosure to a mixture of water and ascorbic acid.

20 *Enteric coated oral dosage form*

A compound of the disclosure is dissolved in an aqueous solution containing polyvinyl pyrrolidone and spray coated onto microcrystalline cellulose or sugar beads in a ratio of 1:5 w/w active agent:beads and then an approximately 5 % weight gain of an enteric coating comprising an acrylic copolymer, for example a combination of acrylic
25 copolymers available under the trade names Eudragit-L® and Eudragit-S®, or hydroxypropyl methylcellulose acetate succinate is applied. The enteric coated beads are loaded into gelatin or hydroxypropyl methylcellulose capsules to provide a unit dosage of, for example, 30 mg active agent per capsule.

Enteric coated oral dosage form

30 An enteric coating comprising a combination of Eudragit-L® and Eudragit-S®, or hydroxypropyl methylcellulose acetate succinate is applied to a tablet oral dosage form or a capsule oral dosage form described above.

Ointment formulation for topical administration

A compound of the disclosure is combined with petrolatum, C₈-C₁₀ triglyceride, octylhydroxystearate, and *N*-methylpyrrolidone in a ratio to provide a composition containing 0.05 % to 5 % active agent by weight.

5 *Ointment formulation for topical administration*

A compound of the disclosure is combined with white petrolatum, propylene glycol, mono- and di-glycerides, paraffin, butylated hydroxytoluene, and edetate calcium disodium in a ratio to provide a composition containing 0.05 % to 5 % active agent by weight.

10 *Ointment formulation for topical administration*

A compound of the disclosure is combined with mineral oil, paraffin, propylene carbonate, white petrolatum and white wax to provide a composition containing 0.05 % to 5 % active agent by weight.

Cream formulation for topical administration

15 Mineral oil is combined with a compound of the disclosure, propylene glycol, isopropyl palmitate, polysorbate 60, cetyl alcohol, sorbitan monostearate, polyoxyl 40 stearate, sorbic acid, methylparaben and propylparaben to form an oil phase, which is combined with purified water by shear blending to provide a composition containing 0.05 % to 5 % active agent by weight.

20 *Cream formulation for topical administration*

A cream formulation comprising a compound of the disclosure, benzyl alcohol, cetyl alcohol, citric acid anhydrous, mono and di-glycerides, oleyl alcohol, propylene glycol, sodium cetostearyl sulphate, sodium hydroxide, stearyl alcohol, triglycerides, and water contains 0.05 % to 5 % active agent by weight.

25 *Cream formulation for topical administration*

A cream formulation comprising a compound of the disclosure, cetostearyl alcohol, isopropyl myristate, propylene glycol, cetomacrogol 1000, dimethicone 360, citric acid, sodium citrate, and purified water, with imidurea, methylparaben, and propylparaben, as preservatives, contains 0.05 % to 5 % active agent by weight.

30

Utility

Inhibition of JAK3 blocks the signaling of many key pro-inflammatory cytokines. Thus the compounds of the disclosure are expected to be useful in the treatment of inflammatory diseases.

The compounds of the disclosure have been designed to be selective for JAK3 over JAK1, JAK2 and TYK2. Selectivity for JAK3 over JAK1 is anticipated to be beneficial as there is some evidence that JAK3 selectivity allows sparing of potentially beneficial cytokines such as IL-10 which has been involved in mucosal healing, IL-22 which is involved in mucus barrier protection and epithelial regeneration, and IL-6 which is involved in the proliferation of intestinal epithelial cells. Selectivity for JAK3 over JAK2 allows sparing of erythropoietin (EPO) and thrombopoietin (TPO) signaling.

Without being limited by this theory, the compounds of the disclosure possess an electrophilic portion which may form a covalent bond with the cysteine (Cys909) present in JAK3, a residue replaced by a serine in the other three JAK isoforms (Goedken et al., *J Biol Chem.*, **2015**, 290, 8, 4573-89). Such covalent binding to JAK3 could be beneficial by providing an extended target engagement which may translate in better efficacy. As described in the experimental part, co-crystal structures of compounds 1, 3 and 4 covalently bound to human JAK3 have been obtained which confirms the irreversible binding nature for each of these ligands to JAK3.

Some compounds of the disclosure have been designed to have their effect at the site of action without significant systemic effects, thereby avoiding the potential adverse systemic immunosuppressive effects.

Gastrointestinal inflammatory disease

In addition to providing potent inhibition of JAK3, some compounds of the disclosure have been designed to be poorly absorbed to minimize systemic exposure. These compounds are designed to have their effect at the site of action, for example, in the colon. As described in Assay 6, certain compounds exhibit low permeabilities with K_p values less than about 5×10^{-6} cm/sec which is considered favorable to minimize systemic exposure and target the colon. Certain compounds have a K_p value less than about 10×10^{-6} cm/sec which may also be sufficient to minimize systemic exposure and target the colon. As described in Assay 7 below, compounds 1, 2, 3, 4, 6, 7, 8, 21 and 22 exhibited a ratio of exposure in the colon to exposure in plasma upon oral administration greater than about 1250. Compounds 9, 5, 19, and 20 exhibited a colon to plasma ratio in excess of about 200.

Oxazolone-induced colitis is an experimental model that has a histological resemblance to human ulcerative colitis. As described in Assay 8, the compounds 1, 2, 3, 4, 5, 6, 7, 8, 3-11, 5-10, 19, 15-1, 3-55, 3-34, 15-3, 21, 3-80, 3-81, 3-72, 3-57, 3-113 and 3-74 demonstrated activity in the oxazolone-induced colitis model in mice. Further, when

tested in Assay 9, an immunosuppression model in mice, which probes systemic functional activity, splenic NK cell counts were unaffected by compounds 1, 2, 4, 5, and 8 at the same or higher doses required to demonstrate efficacy in the oxazolone model.

Finally, compounds 1, 2, 3, 4, 5, 6, 7, and 8 were shown to demonstrate lack of systemic activity in a murine model of IL-2 induced pSTAT5 induction in the thymus.

Thus these compounds demonstrated anti-colitic activity without exhibiting systemic effects in preclinical models.

It is expected that a high colon to plasma ratio will provide robust, lumenally-driven anti-inflammatory activity without associated, systemically-driven, adverse effects. Such compounds may be useful for a variety of gastrointestinal inflammatory indications that include, but are not limited to, inflammatory bowel disease, ulcerative colitis (proctosigmoiditis, pancolitis, ulcerative proctitis and left-sided colitis), Crohn's disease, collagenous colitis, lymphocytic colitis, Behcet's disease, celiac disease, immune checkpoint inhibitor induced colitis, ileitis, eosinophilic esophagitis, graft versus host disease-related colitis, and infectious colitis. Ulcerative colitis (Reimund et al., *J Clin Immunology*, **1996**, 16, 144-150), Crohn's disease (Woywodt et al., *Eur J Gastroenterology Hepatology*, **1999**, 11, 267-276), collagenous colitis (Kumawat et al., *Mol Immunology*, **2013**, 55, 355-364), lymphocytic colitis (Kumawat et al., **2013**), eosinophilic esophagitis (Weinbrand-Goichberg et al., *Immunol Res*, **2013**, 56, 249-260), graft versus host disease-related colitis (Coghill et al., *Blood*, **2001**, 117, 3268-3276), infectious colitis (Stallmach et al., *Int J Colorectal Dis*, **2004**, 19, 308-315), Behcet's disease (Zhou et al., *Autoimmun Rev*, **2012**, 11, 699-704), celiac disease (de Nitto et al., *World J Gastroenterol*, **2009**, 15, 4609-4614), immune checkpoint inhibitor induced colitis (e.g., CTLA-4 inhibitor-induced colitis; (Yano et al., *J Translation Med*, **2014**, 12, 191), PD-1- or PD-L1-inhibitor-induced colitis), and ileitis (Yamamoto et al., *Dig Liver Dis*, **2008**, 40, 253-259) are characterized by elevation of certain pro-inflammatory cytokine levels. As many pro-inflammatory cytokines signal via JAK activation, compounds described in this application may be able to alleviate the inflammation and provide symptom relief.

In particular, the compounds of the disclosure may be useful for the induction and maintenance of remission of ulcerative colitis, and for the treatment of Crohn's disease, immune checkpoint inhibitor induced colitis, and the gastrointestinal adverse effects in graft versus host disease.

In one aspect, therefore, the invention provides a method of treating a gastrointestinal inflammatory disease in a mammal (e.g., a human), the method comprising administering to the mammal a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a compound of the disclosure, or a pharmaceutically acceptable salt thereof.

The invention further provides a method of treating ulcerative colitis in a mammal, the method comprising administering to the mammal a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a compound of the disclosure, or a pharmaceutically acceptable salt thereof.

When used to treat ulcerative colitis, the compounds of the disclosure, or a pharmaceutically acceptable salt thereof will typically be administered orally in a single daily dose or in multiple doses per day, although other forms of administration may be used. The amount of active agent administered per dose or the total amount administered per day will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

Suitable doses for treating ulcerative colitis and other gastrointestinal inflammatory disorders are expected to range from about 1 to about 400 mg/day of active agent, including from about 5 to about 300 mg/day and from about 20 to about 70 mg per day of active agent for an average 70 kg human.

Combination therapy

Compounds of the disclosure or a pharmaceutically acceptable salt thereof may also be used in combination with one or more agents which act by the same mechanism or by different mechanisms to effect treatment of gastrointestinal inflammatory disorders. The different agents may be administered sequentially or simultaneously (in separate compositions or in the same composition). Useful classes of agents for combination therapy include, but are not limited to, aminosalicylates, steroids, systemic immunosuppressants, anti-TNF α antibodies, TNF alpha ligand inhibitor, TNF binding agent, anti-VLA-4 antibodies, anti-integrin $\alpha_4\beta_7$ antibodies, anti-bacterial agents, Glucocorticoid agonists, Nuclear factor kappa B inhibitors, 5-Lipoxygenase inhibitors, integrin alpha-4/beta-7 antagonist, Cyclooxygenase inhibitors, IL-23 antagonists,

Leukotriene BLT receptor antagonist, IL-6 antagonists, IL-8 antagonists, integrin antagonists, nicotinic acetylcholine receptor agonists, PPAR gamma agonists, sphingosine-1-phosphate receptor-1 modulators, B-lymphocyte antigen CD20 inhibitors, calcineurin inhibitors, CD3 antagonist, cell adhesion molecule inhibitors, eosinophil
 5 peroxidase inhibitors, heparin agonists, ICAM1 gene inhibitors, IL-13 antagonists, IL-2 receptor alpha subunit inhibitors, insulin sensitizers, interferon beta ligands, interferon gamma receptor antagonists, interleukin-1 beta ligand modulators, MAdCAM inhibitors, PDE 4 inhibitors, sphingosine-1-phosphate receptor-1 agonists, TLR-9 agonists, acetylcholinesterase inhibitors, ACTH receptor agonists, activin receptor
 10 antagonists, CCR5 chemokine antagonists, CCR9 chemokine antagonists, and anti-diarrheal medicines.

Aminosalicylates that may be used in combination with the present JAK inhibitor compounds include, but are not limited to, mesalamine, osalazine and sulfasalazine. Examples of steroids include, but are not limited to, prednisone, prednisolone,
 15 hydrocortisone, budesonide, beclomethasone, and fluticasone. Systemic immunosuppressants useful for treatment of inflammatory disorders include, but are not limited to cyclosporine, azathioprine, methotrexate, 6-mercaptopurine, and tacrolimus. Further, anti-TNF α antibodies, which include, but are not limited to, infliximab, adalimumab, golimumab, and certolizumab, may be used in combination therapy. Useful
 20 compounds acting by other mechanisms include anti-VLA-4 antibodies, such as natalizumab, anti-integrin $\alpha_4\beta_7$ antibodies, such as vedolizumab, anti-bacterial agents, such as rifaximin, and anti-diarrheal medicines, such as loperamide. (Mozaffari et al., *Expert Opin. Biol. Ther.* **2014**, *14*, 583-600; Danese, *Gut*, **2012**, *61*, 918-932; Lam et al., *Immunotherapy*, **2014**, *6*, 963-971.)

25 Other compounds that may be used in combination with the present JAK inhibitor compounds include, but are not limited to opaganib, abatacept, mongersen, filgotinib, LYC-30937, BI-655130, mirikizumab, adalimumab, tacrolimus, rituximab, GSK-2982772, andecaliximab, naltrexone, risankizumab, QBECO, alicaforsen, etrolizumab, foralumab, ocrelizumab, vedolizumab, amiselimod, ozanimod, dolcanatide, catridecacog,
 30 budesonide, STNM-01, cannabidiol, telotristat etiprate, SHP-647, carotegrast methyl, peg-ilodecakin, TOP-1288, iberogast N, PF-06480605, peficitinib, beclomethasone, recombinant interferon beta-1a, infliximab, golimumab, tralokinumab, ustekinumab, certolizumab pegol, thalidomide, upadacitinib, apremilast, natalizumab, interferon beta-1a, rifaximin, RBX-2660, etrasimod, zileuton, fingolimod, cobitolimod, ropivacaine,

ABX-464, PF-06700841, prednisolone, GLPG-0974, valganciclovir, ciclosporin, VB-201, tulinercept, MDGN-002, PTG-100, dexamethasone, GED-0507-34-Levo, bertilimumab, brazikumab, KHK-4083, rosiglitazone, mocravimod, sotrastaurin, KAG-308, PUR-0110, E-6007, balsalazide, basiliximab, LP-02, ASP-3291, Trichuris suis ova, K(D)PT, midismase, DNVX-078, vatelizumab, alequel, low molecular weight heparin, metenkefalin, tridecactide, HMPL-004, SB-012, olsalazine, balsalazide, propionyl-L-carnitine, Clostridium butyricum, beclomethasone and acemannan.

In another aspect, therefore, the invention provides a therapeutic combination for use in the treatment of gastrointestinal inflammatory disorders, the combination comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, and one or more other therapeutic agents useful for treating gastrointestinal inflammatory disorders, such as the ones illustrated above. For example, the invention provides a combination comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, and one or more agents selected from aminosalicylates, steroids, systemic immunosuppressants, anti-TNF α antibodies, anti-VLA-4 antibodies, anti-integrin $\alpha_4\beta_7$ antibodies, anti-bacterial agents, and anti-diarrheal medicines. Secondary agent(s), when included, are present in a therapeutically effective amount, i.e. in any amount that produces a therapeutically beneficial effect when co-administered with a compound of the disclosure or a pharmaceutically acceptable salt thereof.

Also provided, therefore, is a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, and one or more other therapeutic agents useful for treating gastrointestinal inflammatory disorders.

Further, in a method aspect, the invention provides a method of treating gastrointestinal inflammatory disorders, the method comprising administering to the mammal a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, and one or more other therapeutic agents useful for treating gastrointestinal inflammatory disorders.

When used in combination therapy, the agents may be formulated in a single pharmaceutical composition, as disclosed above, or the agents may be provided in separate compositions that are administered simultaneously or at separate times, by the same or by different routes of administration. When administered separately, the agents are administered sufficiently close in time so as to provide a desired therapeutic effect. Such compositions can be packaged separately or may be packaged together as a kit. The

two or more therapeutic agents in the kit may be administered by the same route of administration or by different routes of administration.

Inflammatory skin disease

Atopic dermatitis and other inflammatory skin diseases have been associated with
 5 elevation of proinflammatory cytokines that rely on the JAK-STAT pathway. Therefore, the compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may be beneficial in a number of dermal inflammatory or pruritic conditions that include, but are not limited to atopic dermatitis, alopecia areata, vitiligo, psoriasis, dermatomyositis, cutaneous T cell lymphoma (Netchiporouk et al., *Cell*
 10 *Cycle*. **2014**; *13*, 3331-3335) and subtypes (Sezary syndrome, mycosis fungoides, pagetoid reticulosis, granulomatous slack skin, lymphomatoid papulosis, pityriasis lichenoides chronica, pityriasis lichenoides et varioliformis acuta, CD30+ cutaneous T-cell lymphoma, secondary cutaneous CD30+ large cell lymphoma, non-mycosis fungoides CD30- cutaneous large T-cell lymphoma, pleomorphic T-cell lymphoma,
 15 Lennert lymphoma, subcutaneous T-cell lymphoma, angiocentric lymphoma, blastic NK-cell lymphoma), prurigo nodularis, lichen planus, primary localized cutaneous amyloidosis, bullous pemphigoid, skin manifestations of graft versus host disease, pemphigoid, discoid lupus, granuloma annulare, lichen simplex chronicus, vulvar/scrotal/perianal pruritus, lichen sclerosus, post herpetic neuralgia itch, lichen
 20 planopilaris, and folliculitis decalvans. In particular, atopic dermatitis (Bao et al., *JAK-STAT*, **2013**, *2*, e24137), alopecia areata (Xing et al., *Nat Med*. **2014**, *20*, 1043-1049), vitiligo (Craiglow et al., *JAMA Dermatol*. **2015**, *151*, 1110-1112), prurigo nodularis (Sonkoly et al., *J Allergy Clin Immunol*. **2006**, *117*, 411-417), lichen planus (Welz-Kubiak et al., *J Immunol Res*. **2015**, ID:854747), primary localized cutaneous amyloidosis
 25 (Tanaka et al., *Br J Dermatol*. **2009**, *161*, 1217-1224), bullous pemphigoid (Feliciani et al., *Int J Immunopathol Pharmacol*. **1999**, *12*, 55-61), and dermal manifestations of graft versus host disease (Okiyama et al., *J Invest Dermatol*. **2014**, *134*, 992-1000) are characterized by elevation of certain cytokines that signal via JAK activation. Accordingly, compounds of the disclosure, or a pharmaceutically acceptable salt thereof,
 30 or a crystalline form thereof, may be able to alleviate associated dermal inflammation or pruritus driven by these cytokines. In particular, compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may be expected to be useful for the treatment of atopic dermatitis and other inflammatory skin diseases.

In one aspect, therefore, the invention provides a method of treating an inflammatory skin disease in a mammal (e.g., a human), the method comprising applying a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, and a pharmaceutical carrier to the skin of the mammal. In one aspect, the inflammatory skin disease is atopic dermatitis.

Compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may also be used in combination with one or more compound useful to treat inflammatory skin diseases. In some embodiments, the one or more compound is a steroid, Histamine H1 receptor antagonist, calcineurin inhibitor, IL-13 antagonist, PDE 4 inhibitor, G-protein coupled receptor-44 antagonist, IL-4 antagonist, 5-HT 1a receptor antagonist, 5-HT 2b receptor antagonist, Alpha 2 adrenoceptor agonist, cannabinoid CB1 receptor antagonist, CCR3 chemokine, antagonist, collagenase inhibitor, cytosolic phospholipase A2 inhibitor, eotaxin ligand inhibitor, GATA 3 transcription factor inhibitor, Histamine H4 receptor antagonist, IL-10 antagonist, IL-12 antagonist, IL-17 antagonist, IL-2 antagonist, IL-23 antagonist, IL-4 receptor modulator, IL-5 antagonist, immunoglobulin E antagonist, immunoglobulin E modulator, interferon gamma receptor antagonist, Interleukin 33 ligand inhibitor, Interleukin-31 receptor antagonist, Leukotriene antagonist, Liver X receptor agonist, Liver X receptor beta agonist, nuclear factor kappa B inhibitor, OX-40 receptor antagonist, PGD2 antagonist, phospholipase A2 inhibitor, SH2 domain inositol phosphatase 1 stimulator, thymic stromal lymphoprotein ligand inhibitor, TLR modulator, TNF alpha ligand modulator, or vanilloid VR1 antagonist. In some embodiments, the one or more compound is a gram positive antibiotic, such as mupirocin or fusidic acid. In some embodiments, the one or more compound is tranilast, tacrolimus, epinastine, SB-011, AM-1030, ZPL-521, MM-36, FB-825, PG-102, viomed, GBR-830, AVX-001, AMG-0101, E-6005, DMT-210, AX-1602, bertilimumab, rosiptor acetate, Q-301, ANB-020, VTP-38543, ZPL-389, lebrikizumab, tezepelumab, fexofenadine, pimecrolimus, bepotastine, crisaborole, tralokinumab, fevipiprant, doxycycline, desloratadine, ALX-101, nemolizumab, asivatrep, ciclosporin, mepolizumab, dupilumab, secukinumab, timapiprant, or ustekinumab.

In one aspect, therefore, the invention provides a method of treating an inflammatory skin disease in a mammal, the method comprising applying a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof,

and a gram positive antibiotic to the skin of the mammal. In another aspect, the invention provides a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, a gram positive antibiotic, and a pharmaceutically-acceptable carrier.

5 Respiratory Diseases

Cytokines which signal through the JAK-STAT pathway, in particular IL-2, IL-3, IL-4, IL-5, IL-6, IL-9, IL-11, IL-13, IL-23, IL-31, IL-27, thymic stromal lymphopoietin (TSLP), interferon- γ (IFN γ) and granulocyte-macrophage colony-stimulating factor (GM-CSF) have also been implicated in asthma inflammation and in other inflammatory
10 respiratory diseases. As described above, the compounds of the disclosure have been shown to be potent inhibitors of JAK3 and have also demonstrated potent inhibition of IL-2 pro-inflammatory cytokines in cellular assays.

The anti-inflammatory activity of JAK inhibitors has been robustly demonstrated in preclinical models of asthma (Malaviya et al., *Int Immunopharmacol*, **2010**, *10*, 829,-
15 836; Matsunaga et al., *Biochem and Biophys Res Commun*, **2011**, *404*, 261-267; Kudlacz et al., *Eur J Pharmacol*, **2008**, *582*, 154-161.) Accordingly, the compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may be useful for the treatment of inflammatory respiratory disorders such as asthma. Inflammation and fibrosis of the lung is characteristic of other respiratory diseases in
20 addition to asthma such as chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, and bronchiolitis obliterans. The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, therefore, may be useful for the treatment of chronic obstructive
25 pulmonary disease, cystic fibrosis, pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, bronchiolitis obliterans organizing pneumonia (also termed COS), primary Graft Dysfunction (PGD), organizing pneumonia (OP), acute rejection (AR), lymphocytic bronchiolitis (LB), chronic Lung Allograft
30 Dysfunction (CLAD), restrictive CLAD (rCLAD or RAS), neutrophilic allograft dysfunction, and sarcoidosis.

In one aspect, therefore, the disclosure provides a method of treating a respiratory disease in a mammal (e.g., a human), the method comprising administering to the

mammal a compound of the disclosure or a pharmaceutically-acceptable salt thereof, or a crystalline form thereof.

In one aspect, the respiratory disease is asthma, chronic obstructive pulmonary disease, cystic fibrosis, pneumonitis, chronic obstructive pulmonary disease (COPD),
 5 cystic fibrosis (CF), pneumonitis, interstitial lung diseases (including idiopathic pulmonary fibrosis), acute lung injury, acute respiratory distress syndrome, bronchitis, emphysema, bronchiolitis obliterans, bronchiolitis obliterans organizing pneumonia (also termed COS), primary Graft Dysfunction (PGD), organizing pneumonia (OP), acute rejection (AR), lymphocytic bronchiolitis (LB), chronic Lung Allograft Dysfunction
 10 (CLAD), restrictive CLAD (rCLAD or RAS), neutrophilic allograft dysfunction, allergic rhinitis or sarcoidosis. In another aspect, the respiratory disease is asthma or chronic obstructive pulmonary disease.

In a further aspect, the respiratory disease is a lung infection, a helminthic infection, pulmonary arterial hypertension, sarcoidosis, lymphangioleiomyomatosis,
 15 bronchiectasis, or an infiltrative pulmonary disease. In yet another aspect, the respiratory disease is drug-induced pneumonitis, fungal induced pneumonitis, allergic bronchopulmonary aspergillosis, hypersensitivity pneumonitis, eosinophilic granulomatosis with polyangiitis, idiopathic acute eosinophilic pneumonia, idiopathic chronic eosinophilic pneumonia, hypereosinophilic syndrome, Löffler syndrome,
 20 bronchiolitis obliterans organizing pneumonia, or immune-checkpoint-inhibitor induced pneumonitis.

The disclosure further provides a method of treating a respiratory disease, the method comprising administering to the mammal a pharmaceutical composition comprising a compound of the disclosure or a pharmaceutically-acceptable salt thereof, or
 25 a crystalline form thereof, and a pharmaceutically-acceptable carrier.

Compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may also be used in combination with one or more compound useful to respiratory diseases.

Ocular Diseases

30 Many ocular diseases have been shown to be associated with elevations of proinflammatory cytokines that rely on the JAK-STAT pathway.

The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, therefore, may be useful for the treatment of a number of ocular diseases that include, but are not limited to, uveitis, diabetic retinopathy, diabetic

macular edema, dry eye disease, age-related macular degeneration, retinal vein occlusion (RVO) and atopic keratoconjunctivitis.

In particular, uveitis (Horai and Caspi, *J Interferon Cytokine Res*, **2011**, 31, 733-744), diabetic retinopathy (Abcouwer, *J Clin Cell Immunol*, **2013**, Suppl 1, 1-12), diabetic macular edema (Sohn et al., *American Journal of Ophthalmology*, **2011**, 152, 686-694), dry eye disease (Stevenson et al, *Arch Ophthalmol*, **2012**, 130, 90-100), retinal vein occlusion (Shchuko et al, *Indian Journal of Ophthalmology*, **2015**, 63(12), 905-911) and age-related macular degeneration (Knickelbein et al, *Int Ophthalmol Clin*, **2015**, 55(3), 63-78) are characterized by elevation of certain pro-inflammatory cytokines that signal via the JAK-STAT pathway. Accordingly, compounds of the disclosure, or a pharmaceutically acceptable salt thereof, may be able to alleviate the associated ocular inflammation and reverse disease progression or provide symptom relief.

In one aspect, therefore, the disclosure provides a method of treating an ocular disease in a mammal, the method comprising administering a pharmaceutical composition comprising a compound of the disclosure, or a pharmaceutically-acceptable salt thereof, or a crystalline form thereof, and a pharmaceutical carrier to the eye of the mammal. In one aspect, the ocular disease is uveitis, diabetic retinopathy, diabetic macular edema, dry eye disease, age-related macular degeneration, retinal vein occlusion or atopic keratoconjunctivitis. In one aspect, the method comprises administering the compound of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, by intravitreal injection.

Compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may also be used in combination with one or more compound useful to ocular diseases.

Other diseases

The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may also be useful to treat other diseases such as other inflammatory diseases, autoimmune diseases or cancers.

The compounds of the disclosure, or a pharmaceutically acceptable salt thereof, or a crystalline form thereof, may be useful to treat one or more of arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, transplant rejection, xerophthalmia, psoriatic arthritis, diabetes, insulin dependent diabetes, motor neurone disease, myelodysplastic syndrome, pain, sarcopenia, cachexia, septic shock, systemic lupus erythematosus, leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, acute

lymphoblastic leukemia, acute myelogenous leukemia, ankylosing spondylitis, myelofibrosis, B-cell lymphoma, hepatocellular carcinoma, Hodgkins disease, breast cancer, Multiple myeloma, melanoma, non-Hodgkin lymphoma, non-small-cell lung cancer, ovarian clear cell carcinoma, ovary tumor, pancreas tumor, polycythemia vera, Sjoegrens syndrome, soft tissue sarcoma, sarcoma, splenomegaly, T-cell lymphoma, and thalassemia major.

Compounds of the disclosure have been demonstrated to be potent inhibitors of the JAK3 enzyme and to be selective for JAK3 over JAK1, JAK2 and TYK2 in enzyme binding assays and to have potent functional activity for JAK3 in a cellular assay as described in the following examples.

EXAMPLES

The following synthetic and biological examples are offered to illustrate the invention, and are not to be construed in any way as limiting the scope of the invention. In the examples below, the following abbreviations have the following meanings unless otherwise indicated. Abbreviations not defined below have their generally accepted meanings.

	ACN	=	acetonitrile
	Calcd	=	calculated
	Boc	=	tert-Butyloxycarbonyl
20	d	=	day(s)
	DIPEA	=	<i>N,N</i> -diisopropylethylamine
	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethyl sulfoxide
	EtOAc	=	ethyl acetate
25	EtOH	=	ethyl alcohol
	h	=	hour(s)
	HATU	=	<i>N,N,N',N'</i> -tetramethyl- <i>O</i> -(7-azabenzotriazol-1-yl)uronium hexafluorophosphate
	IPA	=	isopropyl alcohol
30	MeOH	=	methanol
	min	=	minute(s)
	RT or rt	=	room temperature
	SiG	=	Silica gel
	TEA	=	triethylamine

THF = tetrahydrofuran
THP = tetrahydropyran
TFA = trifluoroacetic acid

Reagents and solvents were purchased from commercial suppliers (Aldrich, Fluka,
5 Sigma, etc.), and used without further purification. Progress of reaction mixtures was
monitored by thin layer chromatography (TLC), analytical high performance liquid
chromatography (anal. HPLC), and mass spectrometry. Reaction mixtures were worked
up as described specifically in each reaction; commonly they were purified by extraction
and other purification methods such as temperature-, and solvent-dependent
10 crystallization, and precipitation. In addition, reaction mixtures were routinely purified
by column chromatography or by preparative HPLC, typically using C18 or BDS column
packings and conventional eluents. Typical preparative HPLC conditions are described
below.

Characterization of reaction products was routinely carried out by mass and
15 ¹H-NMR spectrometry. For NMR analysis, samples were dissolved in deuterated solvent
(such as CD₃OD, CDCl₃, or *d*₆-DMSO), and ¹H-NMR spectra were acquired with a
Varian Gemini 2000 instrument (400 MHz) under standard observation conditions. Mass
spectrometric identification of compounds was performed by an electrospray ionization
method (ESMS) with an Applied Biosystems (Foster City, CA) model API 150 EX
20 instrument or a Waters (Milford, MA) 3100 instrument, coupled to autopurification
systems.

Unless otherwise indicated the following conditions were used for preparative
HPLC purifications.

Column: C18, 5 μm 21.2 x 150 mm or C18, 5 μm 21 x 250 mm or
25 C14, 5 μm 21x150 mm
Column temperature: Room Temperature
Flow rate: 20.0 mL/min
Mobile Phases: A = Water + 0.05 % TFA
B = ACN + 0.05 % TFA,
30 Injection volume: (100-1500 μL)
Detector wavelength: 214 nm

Crude compounds were dissolved in 1:1 water:acetic acid at about 50 mg/mL. A
4 minute analytical scale test run was carried out using a 2.1 x 50 mm C18 column
followed by a 15 or 20 minute preparative scale run using 100 μL injection with the

gradient based on the % B retention of the analytical scale test run. Exact gradients were sample dependent. Samples with close running impurities were checked with a 21 x 250 mm C18 column and/or a 21 x 150 mm C14 column for best separation. Fractions containing desired product were identified by mass spectrometric analysis.

5 Analytic HPLC Conditions

Method A

Column: LUNA C18 (2), 150 x 4.60 mm, 3 μ m

Column temperature: 37 °C

Flow rate: 1.0 mL/min

10 Injection volume: 5 μ L

Sample preparation: Dissolve in 1:1 ACN:water

Mobile Phases: A = Water:ACN:TFA (98:2:0.05)

B = Water:ACN:TFA (2:98:0.05)

Detector wavelength: 250 nm

15 Gradient: 32 min total (time (min)/ % B): 0/2, 10/20, 24/90, 29/90, 30/2, 32/2

Method B

Column: LUNA C18 (2), 150 x 4.60 mm, 3 μ m

Column temperature: 37 °C

20 Flow rate: 1.0 mL/min

Injection volume: 10 μ L

Sample preparation: Dissolve in 1:1 ACN:water

Mobile Phases: A = Water:ACN:TFA (98:2:0.05)

B = Water:ACN:TFA (10:90:0.05)

25 Detector wavelength: 254 nm

Gradient: 35 min total (time (min)/ % B): 0/2, 20/25, 23/90, 26/90, 27/2, 35/2

Powder X-ray diffraction patterns were obtained with a Bruker D8-Advance X-ray diffractometer using Cu- $K\alpha$ radiation ($\lambda = 1.54051$ Å) with output voltage of 45 kV and current of 40 mA. The instrument was operated in Bragg-Brentano geometry with incident, divergence, and scattering slits set to maximize the intensity at the sample. For measurement, a small amount of powder (5-25 mg) was gently pressed onto a sample holder to form a smooth surface and subjected to X-ray exposure. The samples were scanned in 2 θ -2 θ mode from 2° to 35° in 2 θ with a step size of 0.02° and a scan speed of


0.30°seconds per step. The data acquisition was controlled by Bruker DiffracSuite measurement software and analyzed by Jade software (version 7.5.1). The instrument was calibrated with a corundum standard, within $\pm 0.02^\circ$ two-theta angle.

Differential scanning calorimetry (DSC) was performed using a TA Instruments Model Q-100 module with a Thermal Analyst controller. Data were collected and analyzed using TA Instruments Thermal Analysis software. A sample of each crystalline form was accurately weighed into a covered aluminum pan. After a 5 minute isothermal equilibration period at 5 °C, the sample was heated using a linear heating ramp of 10 °C/min from 0 °C to 300 °C.

Thermogravimetric analysis (TGA) measurements were performed using a TA Instruments Model Q-50 module equipped with high resolution capability. Data were collected using TA Instruments Thermal Analyst controller and analyzed using TA Instruments Universal Analysis software. A weighed sample was placed onto a platinum pan and scanned with a heating rate of 10 °C from ambient temperature to 300-350 °C. The balance and furnace chambers were purged with nitrogen flow during use.


Dynamic moisture sorption (DMS) measurements were performed using a VTI atmospheric microbalance, SGA-100 system (VTI Corp., Hialeah, FL 33016). A weighed sample was used and the humidity was lowest possible value (close to 0% RH) at the start of the analysis. The DMS analysis consisted of an initial drying step (0%RH) for 120 minutes, followed by two cycles of sorption and desorption with a scan rate of 5 % RH/step over the humidity range of 5 % RH to 90 % RH. The DMS run was performed isothermally at 25 °C.

Preparation 1: 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole



A mixture of 6-bromo-4-fluoro-1H-indazole (5 g, 23.25 mmol), 3,4-dihydro-2H-pyran (6.38 ml, 69.8 mmol) and p-toluenesulfonic acid monohydrate (0.442 g, 2.325 mmol) in DCM (76 ml) was stirred at rt overnight. The reaction was concentrated and the resulting residue was purified by flash column chromatography using a 0-60% EtOAc/Hexanes gradient to yield the desired product (6.08g, 87% yield).


Preparation 2: tert-butyl 3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate



Sodium hydride (1.003 g, 41.8 mmol) was slowly added to a stirring solution of 1-
 5 Boc-Azetidine-3-yl-methanol (6.89 g, 36.8 mmol) in DMF (60 mL) under an atmosphere
 of N₂ at 0 °C and the reaction was allowed to warm to rt. The frothy reaction mixture was
 stirred for 30 minutes at rt before being cooled once again to 0 °C. A solution of 6-
 bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (5.00 g, 16.71 mmol) in DMF
 (20 mL) was slowly cannulated into the reaction mixture, the reaction was warmed to rt
 10 and stirred for 2 hours at rt. The reaction was quenched with the slow addition of H₂O
 (150 mL) and EtOAc (100 mL) and allowed to stir for 5 minutes. Additional water (100
 mL) was added and the biphasic mixture was extracted with EtOAc (3 x 100 mL). The
 combined organic fractions were then washed with 1:1 H₂O:brine (3 x 100 mL) and dried
 over Na₂SO₄. The resulting clear slightly pink oil was purified by flash column
 15 chromatography using a 0-100% EtOAc/hexanes gradient to yield the desired product as a
 clear, colorless, viscous oil (7.34g, 15.74 mmol, 94% yield). (*m/z*): [M+H]⁺ calcd for
 C₂₁H₂₈BrN₃O₄ 466.13 found 466.1.

Preparation 3: tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate


20



Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane
 adduct (1.93 g, 2.36 mmol) was added to a solution of 4-hydroxybenzeneboronic acid


(3.26 g, 23.61 mmol), tert-butyl 3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (7.34 g, 15.74 mmol), and potassium phosphate (10.02 g, 47.2 mmol) in 1,4-dioxane (63.0 ml) and water (15.74 ml). The reaction mixture was degassed with N₂ for 10 minutes and then stirred at 110 °C for 2 hours. The reaction mixture was concentrated *in vacuo* to a volume of about 5mL. A saturated aqueous solution of ammonium chloride (20 mL) was added and the mixture was extracted with methylene chloride (3 x 20 mL). The methylene chloride extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via flash column chromatography using 50% ethyl acetate in hexanes to yield tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (7.55 g, 15.74 mmol, 100 % yield). (*m/z*): [M+H]⁺ calcd for C₂₇H₃₃N₃O₅ 480.24 found 480.1.

Preparation 4: 4-(4-(azetidin-3-ylmethoxy)-1H-indazol-6-yl)phenol




TFA (10.90 ml) was slowly added to a solution of tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (7.84 g, 16.35 mmol) in dichloromethane (21.80 ml). The clear solution was stirred at rt for 5 hours. The reaction was concentrated *in vacuo* to yield 4-(4-(azetidin-3-ylmethoxy)-1H-indazol-6-yl)phenol as a TFA salt (6.69 g, 16.35 mmol, 100% yield). (*m/z*): [M+H]⁺ calcd for C₁₇H₁₇N₃O₂ 296.13 found 296.1.

Example 1A: (E)-4-(dimethylamino)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidin-1-yl)but-2-en-1-one



HATU (2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium
 5 hexafluorophosphate, 817 mg, 2.150 mmol) was added to a solution of (E)-4-(
 (dimethylamino)but-2-enoic acid hydrochloride (405 mg, 2.44 mmol) in DMF (2.00 mL)
 and the reaction mixture was stirred at rt for 5 minutes. 4-(4-(azetidin-3-ylmethoxy)-1H-
 indazol-6-yl)phenol TFA salt (800 mg, 1.954 mmol) was added, followed by DIPEA
 (1.707 ml, 9.77 mmol) and the reaction mixture was stirred at rt for 15 minutes and then
 10 concentrated *in vacuo* to yield a yellow liquid. The crude liquid was purified via
 preparatory scale C18 column chromatography using a gradient of 20-80% acetonitrile in
 water with 0.05% trifluoroacetic acid to yield (E)-4-(dimethylamino)-1-(3-(((6-(4-
 hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidin-1-yl)but-2-en-1-one as a TFA salt
 (235.0 mg, 0.578 mmol, 29.6 % yield). (*m/z*): [M+H]⁺ calcd for C₂₃H₂₆N₄O₃ 407.20 found
 15 407.2. ¹H NMR (400 MHz, DMSO-d₆) δ 13.07 (s, 1H), 10.33 (s, 1H), 7.93 (s, 1H), 7.55
 (d, J = 7.9 Hz, 2H), 7.21 (s, 1H), 6.86 (d, J = 8.6 Hz, 2H), 6.79 (s, 1H), 6.66 – 6.58 (m,
 1H), 6.45 (d, J = 15.5 Hz, 1H), 4.45 – 4.36 (m, 2H), 4.17 – 4.07 (m, 2H), 3.91-3.83 (m,
 3H), 3.22-3.10 (m, 2H), 2.75 (s, 6H).

Example 1B: Crystalline (E)-4-(dimethylamino)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidin-1-yl)but-2-en-1-one Form 4



5 Step 1

To a suspension of 6-bromo-4-fluoro-1H-indazole (20.0g, 93 mmol) in 200ml of DCM, was added p-toluenesulfonic acid monohydrate (1.769 g, 9.30 mmol). The reaction mixture remained a suspension. 3,4-dihydro-2H-pyran (16.97 ml, 186 mmol) was then added. A complete dissolution of solids was observed after 5 minutes. The reaction mixture was stirred at RT overnight to form a dark solution. 200mL of aqueous bicarbonate were added, the phases were separated and the organic layer was washed by 200ml of brine and dried with sodium sulfate. The solution was filtered through silica plug to remove the dark color and the silica was washed with 300ml of DCM. The solvent was evaporated to give 25g of product as an off-white solid.

15 Step 2A

In a 500ml round-bottomed flask was added tert-butyl 3-(hydroxymethyl)azetidine-1-carboxylate (22.53 g, 120 mmol) and 100ml of dimethylacetamide (DMAc). The flask was purged with nitrogen. Cesium carbonate (39.2 g, 120 mmol) was added. 6-Bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (24g, 80 mmol) was added and the reaction mixture was stirred at 70°C for 2

days. An additional 0.5eq of tert-butyl 3-(hydroxymethyl)azetidine-1-carboxylate and cesium carbonate were added and the reaction was heated at 70°C overnight to provide full conversion. The reaction mixture was cooled to RT and then slowly poured into stirred ice-cold water (700ml). The resulting slurry was stirred for 20 minutes and then
5 filtered and dried to yield 32g of product. The product was crystallized from methanol-water by adding water slowly as antisolvent until cloudiness. A white slurry developed over time. The solid was filtered and dried to yield 28g of material over 98% pure.

Step 2B

Alternatively, potassium t-butoxide was added to a solution of tert-butyl 3-
10 (hydroxymethyl)azetidine-1-carboxylate in DMAc at 0°C. After 60 minutes, 6-Bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole was added at 0°C and the reaction mixture was warmed up to room temperature giving clean complete conversion in less than 6 hours.

Step 3

15 In a 250 mL Schlenk flask was added tert-butyl 3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (10.0g, 21.44 mmol) and 80ml of dioxane. (4-Hydroxyphenyl)boronic acid (4.44 g, 32.2 mmol) was added. Potassium phosphate tribasic anhydrous (13.65 g, 64.3 mmol) was added with 20ml of water and the reaction mixture was purged with nitrogen. PdCl₂(dppf)-CH₂Cl₂ adduct
20 (0.876 g, 1.072 mmol) was added and the flask with a condenser was re-filled with nitrogen three times. The reaction mixture was heated at 110°C for 2h and 30 minutes to show complete conversion by HPLC. The reaction mixture was cooled to RT followed by removal of most of the dioxane. 150ml of saturated ammonium chloride was added followed by 150ml of ethyl acetate. The pH was adjusted to neutral with 1M aq HCl. The
25 phases were separated and the organic layer was dried with sodium sulfate followed by solvent removal. The crude product was dissolved in 150ml of DCM and loaded on 300g SiG column, eluted with 20-50 % ethyl acetate in hexanes. The pure fractions were combined and the solvent was evaporated. 100ml of MeTHF was added, followed by addition of crystalline seeds. A slurry developed over time. 100ml of DIPE was added
30 and the slurry was stirred overnight. Filtration and drying yielded 7.9g of pure material (>99%).

Step 4

In a 50 mL round-bottomed flask was added tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (85g,

177 mmol) in 400ml of methanol. p-Toluenesulfonic acid monohydrate (101 g, 532 mmol) was added and the reaction mixture was stirred at RT to give about 90% conversion after 2 days. The reaction mixture was stirred for another 24 hours at room temperature to give about 96% conversion. 400ml of diisopropyl ether was added and the resulting slurry was stirred at room temperature overnight and formed a crystalline salt overnight. Filtration and drying under nitrogen yielded 100g of >99% pure material as a bis PTSA salt.

Step 5

In a 100 mL round-bottomed flask was added (E)-4-(dimethylamino)but-2-enoic acid, HCl (1.899 g, 11.46 mmol) and 20ml of DMF. HCTU o-(1h-6-chlorobenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (4.91 g, 11.46 mmol) was added and the reaction mixture was stirred at RT for 20 minutes. In a separate flask was added 4-(4-(azetidin-3-ylmethoxy)-1H-indazol-6-yl)phenol, 2TFA (5.0g, 9.55 mmol) and 20ml of DMF. The resulting solution was cooled to 0° C, followed by a slow addition of DIPEA (5 ml, 38.2 mmol, 3eq). The above solution was added (over 5 minutes) to the pre-activated acid reaction mixture and the reaction mixture was stirred at room temperature for 15 minutes. One additional equivalent of DIPEA was added dropwise. Complete conversion was observed by HPLC after 20 minutes at room temperature. The reaction mixture was poured into 200ml of stirred water. Sticky solids precipitated out at pH around 7. The pH was carefully adjusted to about 8 with aqueous ammonia. The solution was extracted three times with 100ml of MeTHF. The combined organic layers were dried with sodium sulfate followed by solvent removal. The crude free base product was dissolved in 25ml of 10% methanol in DCM, loaded on 125g silica gel column and eluted with isocratic 10% methanol in DCM containing 0.5% aqueous ammonia. The pure fractions were combined and the solvent was evaporated. Recrystallization from acetone yielded >98% pure material (50% yield).

25 mg of compound 1 was suspended in 1 mL of water. The resulting suspension was stirred for 2 days at 50 °C, filtered, washed with 2 mL of water and dried under ambient conditions for a few hours to provide Form 4.

Alternatively, compound 1 was dissolved in ethanol and water or methanol and water by complete dissolution in 10 volumes of alcohol followed by slow addition of approximately 8-10 volumes of water until cloud point. Seeds of Form 4 were added and a slurry developed slowly over time. Then more water was added slowly (about 10 volumes) and the solid was filtered and dried to give Form 4.

Example 1C: Crystalline (E)-4-(dimethylamino)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidin-1-yl)but-2-en-1-one Form 3

Form 3 is an anhydrous free base crystalline form of compound 1.

5 150 mg of compound 1 was suspended in 2 mL of a 1:1 mixture of acetonitrile and isopropanol. The resulting suspension was stirred for 1 day at 50 °C, filtered, washed with 2 mL of a 1:1 mixture of acetonitrile and isopropanol and dried under ambient conditions for a few hours to provide Form 3, which was determined to be a crystalline anhydrous free base.

10 Alternatively, 200mg of compound 1 as an amorphous free base was dissolved in 2ml of IPA at RT. An equal amount of acetonitrile was added. More compound (0.5g total) was added until a saturated solution formed. Seeds were added and the mixture was stirred overnight. A white slurry developed over time. Filtration and drying yielded 400mg of product as Form 3, which was determined to be an anhydrous free base.

15 Example 1D: Characterization of Form 3

Samples of Form 3 were analyzed by powder X-ray diffraction (PXRD), differential scanning calorimetry (DSC), thermogravimetric analysis (TGA), and dynamic moisture sorption (DMS).

Powder X-Ray Diffraction

20 The powder X-ray diffraction pattern of Form 3 is shown in Figure 9. Observed PXRD two-theta peak positions and *d*-spacings are shown below.

2-Theta	d(Å)	Area	A%
4.82	18.32	16367	2.7
9.67	9.14	610151	100.0
11.61	7.61	140214	23.0
11.92	7.42	49009	8.0
12.98	6.81	21804	3.6
13.23	6.69	21840	3.6
15.69	5.64	88944	14.6
16.19	5.47	89861	14.7
16.45	5.38	74091	12.1
16.67	5.31	57780	9.5

17.61	5.03	158810	26.0
18.88	4.70	261799	42.9
19.39	4.57	26844	4.4
19.96	4.45	164339	26.9
20.14	4.40	125226	20.5
22.14	4.01	52100	8.5
23.33	3.81	74095	12.1
23.84	3.73	166452	27.3
24.06	3.70	200590	32.9
24.29	3.66	79151	13.0
25.31	3.52	119197	19.5
25.63	3.47	88838	14.6
27.06	3.29	62513	10.2
27.31	3.26	47507	7.8
30.10	2.97	18475	3.0
30.53	2.93	16406	2.7

Thermal Analysis

A representative DSC thermogram of the Form 3 crystalline free form of the invention is shown in Figure 10. The differential scanning calorimetry (DSC) trace recorded at a heating rate of 10 °C per minute exhibits a peak in endothermic heat flow, identified as a melt transition, with an onset at about 197.7 °C and a peak at about 201.3 °C. Melting was followed immediately by decomposition.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic heat flow with a peak at about 201.3 °C.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic heat flow at a temperature between about 197.7 °C and about 204 °C.

A representative TGA trace of the Form 3 crystalline free form of the invention is shown in Figure 11. The thermal gravimetric analysis (TGA) trace of Figure 11 shows no significant weight loss at temperatures below the onset of decomposition at about 195 °C.

Dynamic Moisture Sorption Assessment

A representative DMS trace for the Form 3 crystalline free form of the invention is shown in Figure 12.

Form 3 demonstrated about 0.33 % weight gain in the humidity range of 5 % to 90 % relative humidity. Form 3 is considered to be non-hygroscopic.

5

Example 1E: Characterization of Form 4

Samples of Form 4 were analyzed by powder X-ray diffraction (PXRD), differential scanning calorimetry (DSC), thermogravimetric analysis (TGA), and dynamic moisture sorption (DMS).

10 Powder X-Ray Diffraction

The powder X-ray diffraction pattern of Form 4 is shown in Figure 13.

Observed PXRD two-theta peak positions and *d*-spacings are shown below.

2-Theta	d(Å)	Area	A%
6.26	14.10	161835	73.90
11.86	7.46	51703	23.60
12.51	7.07	82712	37.80
13.16	6.72	72445	33.10
14.98	5.91	53903	24.60
16.55	5.35	218849	100.00
16.94	5.23	133498	61.00
17.61	5.03	28496	13.00
18.33	4.84	170874	78.10
18.78	4.72	35644	16.30
19.39	4.57	92940	42.50
19.57	4.53	86983	39.70
19.84	4.47	19121	8.70
21.45	4.14	22435	10.30
21.82	4.07	22934	10.50
22.57	3.94	62931	28.80
23.61	3.76	162063	74.10
24.24	3.67	135952	62.10

24.67	3.61	38471	17.60
25.10	3.54	55236	25.20
25.39	3.51	32363	14.80
27.19	3.28	27010	12.30
27.39	3.25	32148	14.70
28.55	3.12	24879	11.40
31.51	2.84	18975	8.70

Thermal Analysis

A representative DSC thermogram of the Form 4 crystalline free form is shown in Figure 14. The differential scanning calorimetry (DSC) trace recorded at a heating rate of 10 °C per minute exhibits a desolvation endotherm with an onset at about 60.9 °C and a peak at about 103.6 °C, and a melting endotherm characterized by an onset at about 167.3 °C. The compound decomposes at melting and the melting endotherm and the decomposition exotherm overlap.


A representative TGA trace of the Form 4 crystalline free form is shown in Figure 15. The thermal gravimetric analysis (TGA) trace of Figure 15 shows a weight loss of about 3.54% at 100 °C. The compound desolvates at an onset temperature of about 50 °C. The compound decomposes at an onset temperature of about 165 °C.

Dynamic Moisture Sorption Assessment

A representative DMS trace for the Form 4 crystalline free form is shown in Figure 16.


Form 4 demonstrated about 5.01 % weight gain in the humidity range of 5 % to 90 % relative humidity. Form 4 is considered to be moderately hygroscopic.

Preparation 5: tert-butyl (3S)-3-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate



Sodium hydride (0.201 g, 8.36 mmol) was added to a solution of (S)-N-boc-3-pyridinol (1.25 g, 6.69 mmol) in DMF (12 ml) under an atmosphere of N₂ at 0 °C and the reaction mixture was stirred at rt for 20 minutes. 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (1.0 g, 3.34 mmol) was added and the reaction mixture was stirred at rt for 1 hour. Water (1 mL) was added and the reaction was concentrated *in vacuo*. The crude material was purified via flash column chromatography using 40% ethyl acetate in hexanes to yield tert-butyl (3S)-3-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate (1.40 g, 3.00 mmol, 90 % yield).
 10 (m/z): [M+H]⁺ calcd for C₂₁H₂₈BrN₃O₄ 466.13 found 466.1.

Preparation 6: tert-butyl (3S)-3-((6-(3-chloro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate



15


Palladium acetate (0.135 g, 0.600 mmol) was added to a solution of 3-chloro-4-hydroxyphenylboronic acid (0.776 g, 4.50 mmol), tert-butyl (3S)-3-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate (1.40 g, 3.00 mmol), 1,1'-bis(di-t-butylphosphino)ferrocene (0.285 g, 0.600 mmol) and potassium phosphate (1.912 g, 9.01 mmol) in 1,4-dioxane (12 ml) and water (3.00 ml). The reaction mixture was degassed with nitrogen for 10 minutes and then stirred at 110 °C for 2 hours.
 20

The reaction mixture was concentrated *in vacuo* to a volume of about 5mL. A saturated aqueous solution of ammonium chloride (20 mL) was added and the mixture was extracted with methylene chloride (3 x 20 mL). The methylene chloride extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid.

- 5 The crude liquid was purified via flash column chromatography using 50% ethyl acetate in hexanes to yield tert-butyl (3S)-3-((6-(3-chloro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate (1.35 g, 2.63 mmol, 87 % yield) as a clear yellow liquid. (*m/z*): [M+H]⁺ calcd for C₂₇H₃₂ClN₃O₅ 514.20 found 514.2.


10

Preparation 7: (S)-2-chloro-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol




- 4.0N HCl in dioxane (13.13 ml, 52.5 mmol) was added to a solution of tert-butyl (3S)-3-((6-(3-chloro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate (1.35 g, 2.63 mmol) in dioxane (6 ml), and the reaction mixture was stirred at 60 °C for 30 minutes. The reaction mixture was concentrated *in vacuo* to yield (S)-2-chloro-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol as an HCl salt (0.962 g, 2.63 mmol, 100 % yield). (*m/z*): [M+H]⁺ calcd for C₁₇H₁₆ClN₃O₂ 330.09 found 330.2.
- 15
- 20

Example 2: (S)-1-(3-((6-(3-chloro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one



5 N,N-Diisopropylethylamine (4.59 ml, 26.3 mmol) was added to a solution of (S)-2-chloro-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol HCl salt (0.962 g, 2.63 mmol) in DMF (13.0 ml) at 0 °C, followed by acryloyl chloride (0.277 ml, 3.41 mmol). The reaction mixture was stirred at rt for 15 minutes then concentrated to a volume of about 2 mL. The crude liquid was purified via preparatory scale C18 column chromatography
 10 using a gradient of 20-80% acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-1-(3-((6-(3-chloro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (344 mg, 0.691 mmol, 26.3 % yield). (*m/z*): [M+H]⁺ calcd for C₂₀H₁₈ClN₃O₃ 384.10 found 384.1. ¹H NMR (400 MHz, DMSO-d₆) δ 13.03 (s, 1H), 10.33 (s, 1H), 7.98 (s, 1H), 7.74-7.67 (m, 1H), 7.57-7.49 (m, 1H), 7.24 (s, 1H), 7.07 (d, J = 0.5 Hz, 1H), 6.80 (s, 1H), 6.69 – 6.56 (m, 1H), 6.18-6.11 (m, 1H), 5.71-5.62 (m, 1H),
 15 4.01-3.60 (m, 5H), 2.35-2.14 (m, 2H).

Preparation 8: tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate



20


To a dry scintillation vial purged with N₂, tert-butyl 3-(2-hydroxypropan-2-yl)azetidine-1-carboxylate (374 mg, 1.74 mmol) was added, dissolved in 2.4 mL DMF

and the solution was cooled to 0 °C. 60% by weight sodium hydride in mineral oil (134 mg, 3.34 mmol) was slowly added to the stirring solution and the reaction was allowed to warm to rt following the addition. The frothy reaction was stirred for 30 minutes before being cooled once again to 0 °C. A solution of 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (400 mg, 1.34 mmol) in 1 mL DMF was slowly added into the scintillation vial containing the organosodium solution. Following the addition, the reaction was warmed to rt and stirred for 2 hours upon which LCMS indicated full conversion of the starting material to the desired product. The reaction was quenched with the slow addition of 1 mL H₂O and 1 mL EtOAc, which was allowed to stir for 5 minutes.

The biphasic solution was then transferred to a separatory funnel and an additional 5 mL H₂O were added. The mixture was extracted with 3 times 10 mL of EtOAc and the aqueous layer was discarded. The combined organic fractions were then washed with 3x10 mL 1:1 H₂O:brine to remove residual DMF. The organic was then dried over Na₂SO₄, filtered, and concentrated to a lightly yellow oil. The oil was then purified by flash column chromatography using a 0-40% EtOAc:hexanes gradient. The product was isolated pure as a clear, colorless, viscous oil (288 mg, 44% yield).

Preparation 9: tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate

20




A 1:1 mixture of Palladium acetate (26.2 mg, 0.116 mmol) and 1,1'-Bis(di-*t*-butylphosphino)ferrocene (55.3 mg, 0.116 mmol) was added to a solution of 3-chloro-4-hydroxybenzeneboronic acid (151 mg, 0.874 mmol), tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (288 mg, 0.582 mmol), and potassium phosphate (371 mg, 1.747 mmol) in 1,4-dioxane (2.0 ml) and water (0.50 ml). The reaction mixture was degassed with nitrogen for 10 minutes and then stirred at 110 °C for 2 hours. The reaction mixture was concentrated *in vacuo* to a volume of about 5 mL. A saturated aqueous solution of ammonium chloride (5

mL) was added and the mixture was extracted with ethyl acetate (2 x 5 mL). The ethyl acetate extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via flash column chromatography using 40% ethyl acetate in hexanes to yield tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (230 mg, 0.424 mmol, 73 % yield). (m/z): [M+H]⁺ calcd for C₂₉H₃₆N₃O₅ 542.24 found 542.3.

Preparation 10: 4-(4-((2-azetidin-3-yl)propan-2-yl)oxy)-1-H-indazole-6-yl)-2-chlorophenol

10




15

Tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (230 mg, 0.424 mmol) was dissolved in dichloromethane (1 ml) and TFA (1 ml) was slowly added. The clear solution was stirred at rt for 5 hours upon which LCMS indicated good conversion to the desired product. The reaction was concentrated down to provide 4-(4-((2-azetidin-3-yl)propan-2-yl)oxy)-1-H-indazole-6-yl)-2-chlorophenol as a TFA salt (100% yield). (m/z): [M+H]⁺ calcd for C₁₉H₂₀N₃O₂ 358.13 found 358.1.

20

Example 3A: (E)-1-(3-(2-((6-(3-chloro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidin-1-yl)-4-(dimethylamino)but-2-en-1-one



HATU (2-(7-aza-1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate, 100 mg, 0.263 mmol) was added to a solution of (E)-4-(dimethylamino)but-2-enoic acid hydrochloride (34 mg, 0.263 mmol) in DMF (1 mL).

5 The reaction mixture was stirred at rt for 5 minutes then 4-(4-((2-azetidin-3-yl)propan-2-yl)oxy)-1-*H*-indazole-6-yl)-2-chlorophenol TFA salt (118 mg, 0.251 mmol) was added, followed by DIPEA (0.438 mL, 2.507 mmol). The reaction mixture was stirred at rt for 15 minutes and then concentrated *in vacuo* to yield a yellow liquid. The crude liquid was purified via preparatory scale SFC (supercritical fluid chromatography) using methanol

10 with liquid carbon dioxide to yield (E)-1-(3-(2-((6-(3-chloro-4-hydroxyphenyl)-1*H*-indazol-4-yl)oxy)propan-2-yl)azetidin-1-yl)-4-(dimethylamino)but-2-en-1-one (41.3 mg, 0.084 mmol, 33 % yield). (*m/z*): [*M*+*H*]⁺ calcd for C₂₅H₂₉ClN₄O₃ 469.20 found 469.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.07 (s, 1H), 10.33 (s, 1H), 7.95 (s, 1H), 7.62 (d, *J* = 2.3 Hz, 1H), 7.47 (dd, *J* = 8.5, 2.3 Hz, 1H), 7.34 (s, 1H), 7.04 (d, *J* = 8.5 Hz, 1H), 6.85 (d, *J*

15 = 1.1 Hz, 1H), 6.63 – 6.53 (m, 1H), 6.39 (d, *J* = 15.3 Hz, 1H), 4.37 – 4.23 (m, 2H), 4.11 – 3.91 (m, 2H), 3.69 (d, *J* = 6.4 Hz, 2H), 3.02 – 2.90 (m, 1H), 2.62 (s, 9H), 1.32 (s, 9H).

Example 3B: Crystalline (E)-1-(3-(2-((6-(3-chloro-4-hydroxyphenyl)-1*H*-indazol-4-yl)oxy)propan-2-yl)azetidin-1-yl)-4-(dimethylamino)but-2-en-1-one Form 1

20 Form 1 is an anhydrous freebase crystalline form of compound 3.

80 mg of compound 3 as an amorphous freebase was dissolved in 0.5 mL of ethanol. The resulting mixture was stirred for one day at room temperature and yielded a precipitate. The solid was isolated by filtration, washed with 1 mL of ethanol and dried under ambient conditions for a few hours to give Form 1.

25 Alternatively, in a 50 mL round-bottomed flask, was added compound 3 which had been purified by SiG (1.3g, 2.77 mmol). 10 mL of acetone was added and the mixture was stirred at RT. Seeds of crystalline free base Form 1 were added. A thick white slurry

developed over time. Filtration and drying yielded 1g of crystalline free base Form 1 with >98% purity.

Example 3B: Crystalline (E)-1-(3-(2-((6-(3-chloro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidin-1-yl)-4-(dimethylamino)but-2-en-1-one Form 2

5 Form 2 is a freebase hydrate crystalline form of compound 3.

55 mg of amorphous freebase of compound 3 was dissolved in 0.25 mL of methanol. At this stage, water was added as an antisolvent in a ratio of about 1:2 methanol:water. The resulting mixture was sonicated for several minutes at room temperature to yield a precipitate. The resulting suspension was stirred for 1 day at room
10 temperature, filtered, washed with 1 mL of methanol and dried to provide Form 2.

Alternatively, compound 3 was dissolved in ethanol and water or methanol and water by complete dissolution in 10 volumes of alcohol followed by slow addition of approximately 8-10 volumes of water until cloud point. Seeds of Form 2 were added and a slurry developed slowly over time. Then more water was added slowly (about 10
15 volumes) and the resulting solid was filtered and dried to give Form 2.

Example 3C: Characterization of Form 1

Samples of Form 1 were analyzed by powder X-ray diffraction (PXRD), differential scanning calorimetry (DSC), thermogravimetric analysis (TGA), and dynamic moisture sorption (DMS).

20 Powder X-Ray Diffraction

The powder X-ray diffraction pattern of Form 1 is shown in Figure 1.

Observed PXRD two-theta peak positions and *d*-spacings are shown below.

2-Theta	d(Å)	Area	A%
5.65	15.63	343682	100.0
7.12	12.40	17888	5.2
10.02	8.82	13064	3.8
11.16	7.92	60848	17.7
13.10	6.75	13327	3.9
14.22	6.22	343837	100.0
14.82	5.97	158470	46.1
15.16	5.84	338146	98.3
16.55	5.35	84525	24.6

17.06	5.19	188182	54.7
19.31	4.59	251362	73.1
20.08	4.42	57430	16.7
21.08	4.21	29843	8.7
21.65	4.10	45528	13.2
22.51	3.95	88807	25.8
22.98	3.87	75161	21.9
24.43	3.64	95427	27.8
25.02	3.56	74549	21.7
25.72	3.46	24251	7.1
26.80	3.32	23046	6.7
27.06	3.29	26069	7.6
28.31	3.15	10766	3.1
30.08	2.97	45168	13.1
30.31	2.95	38502	11.2
32.08	2.79	22050	6.4

Thermal Analysis

A representative DSC thermogram of the Form 1 crystalline free form of the invention is shown in Figure 2. The differential scanning calorimetry (DSC) trace
5 recorded at a heating rate of 10 °C per minute exhibits a peak in endothermic heat flow, identified as a melt transition, with an onset at about 154.9 °C and a peak at about 162.9 °C. Melting was followed immediately by decomposition.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic
10 heat flow with a peak at about 162.9 °C. The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic heat flow with a peak at 162.9 ± 3 °C.

The crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10 °C per minute which shows a maximum in endothermic
15 heat flow at a temperature between about 154.9 °C and about 171 °C.

A representative TGA trace of the Form 1 crystalline free form is shown in Figure 3. The thermal gravimetric analysis (TGA) trace of Figure 3 shows a small weight loss of

about 0.14% at 100 °C. The compound decomposes at an onset temperature of about 175 °C.

Dynamic Moisture Sorption Assessment

5 A representative DMS trace for the Form 1 crystalline free form is shown in Figure 4.

Form 1 demonstrated about 1.62 % weight gain in the humidity range of 5 % to 90 % relative humidity. Form 1 is considered to be slightly hygroscopic.

Example 3D: Characterization of Form 2

10

Samples of Form 2 were analyzed by powder X-ray diffraction (PXRD), differential scanning calorimetry (DSC), thermogravimetric analysis (TGA), and dynamic moisture sorption (DMS).

Powder X-Ray Diffraction

15 The powder X-ray diffraction pattern of Form 2 is shown in Figure 5. Observed PXRD two-theta peak positions and *d*-spacings are shown below.

2-Theta	d(Å)	Area	A%
6.90	12.80	248788	52.1
9.15	9.66	73363	15.4
10.00	8.84	92849	19.4
11.18	7.91	42354	8.9
12.76	6.93	34590	7.2
13.33	6.64	33198	6.9
13.82	6.40	75603	15.8
14.43	6.13	20656	4.3
15.51	5.71	93338	19.5
16.04	5.52	19890	4.2
17.00	5.21	58922	12.3
17.90	4.95	339924	71.1
18.31	4.84	477815	100.0

20.90	4.25	169039	35.4
22.06	4.03	68934	14.4
22.51	3.95	110745	23.2
25.00	3.56	23916	5.0
26.92	3.31	52661	11.0
27.26	3.27	63506	13.3
27.61	3.23	22650	4.7
29.37	3.04	15298	3.2
30.53	2.93	24930	5.2
30.92	2.89	22089	4.6

Thermal Analysis

A representative DSC thermogram of the Form 2 crystalline free form is shown in Figure 6. The differential scanning calorimetry (DSC) trace recorded at a heating rate of 10 °C per minute exhibits a desolvation endotherm with an onset at about 52.7 °C and a peak at about 84.4 °C, and a melting endotherm with an onset at about 160.0 °C and a peak at about 167.6 °C. Melting was followed immediately by decomposition.

A representative TGA trace of the Form 2 crystalline free form of the invention is shown in Figure 7. The thermal gravimetric analysis (TGA) trace of Figure 7 shows a weight loss of about 6.73% at 75 °C. The compound desolvates at an onset temperature of about 25 °C. The compound decomposes at an onset temperature of about 185 °C.

Dynamic Moisture Sorption Assessment

A representative DMS trace for the Form 2 crystalline free form of the invention is shown in Figure 8. Form 2 converts to a hydrate (Form 2b) at RH above 65 %. The dehydration occurs at RH below 15 %. The total moisture uptake between 5%-90% RH is 7.99 %.

Example 3E: Characterization of Form 2b

Samples of Form 2b were analyzed by powder X-ray diffraction (PXRD).


Powder X-Ray Diffraction

The powder X-ray diffraction pattern of Form 2b is shown in Figure 17. Observed PXRD two-theta peak positions and *d*-spacings are shown below.

2-Theta	d(Å)	Area	A%
7.61	11.60	182433	39.1
10.33	8.56	128795	27.6
11.25	7.86	13807	3
12.71	6.96	102263	21.9
13.23	6.69	34318	7.4
13.66	6.48	86861	18.6
13.90	6.37	60997	13.1
15.02	5.89	53064	11.4
15.27	5.80	166039	35.6
15.88	5.58	163970	35.2
16.33	5.42	87102	18.7
16.76	5.29	364312	78.2
17.90	4.95	346920	74.4
18.26	4.85	105152	22.6
20.67	4.29	465993	100
21.37	4.16	104882	22.5
21.92	4.05	69905	15
22.31	3.98	133265	28.6
22.90	3.88	42178	9.1
23.22	3.83	20810	4.5
23.61	3.76	39278	8.4
24.74	3.60	160353	34.4
25.78	3.45	67865	14.6
26.23	3.40	30059	6.5
26.73	3.33	120802	25.9
27.57	3.23	17553	3.8
29.10	3.07	75111	16.1
29.39	3.04	97911	21
30.72	2.91	66141	14.2
30.94	2.89	71606	15.4

31.69	2.82	14912	3.2
32.06	2.79	33582	7.2
33.76	2.65	16247	3.5
34.35	2.61	18479	4


Preparation 11: 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine



5

4,6-dichloro-1H-pyrazolo[3,4-d]pyrimidine (2.50 g, 13.23 mmol) was added to a flask and dissolved in 1,4-dioxane (52.9 ml). *p*-Toluenesulfonic acid monohydrate (pTsOH, 0.25 g, 1.32 mmol) was then added to the clear, pale yellow solution followed by 3,4-dihydro-2h-pyran (1.8 ml, 19.84 mmol) and the reaction was heated to 40 °C and stirred overnight. The resulting mixture was then cooled and concentrated *in vacuo*. The crude was purified by flash column chromatography using a 0-100% EtOAc:hexanes gradient to yield 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (2.6 g, 9.52 mmol, 72.0 % yield). (*m/z*): [M+H]⁺ calculated for C₁₀H₁₀Cl₂N₄O 274.0 found 188.9 (loss of THP fragment).

Preparation 12: tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidine-1-carboxylate




To a dry round bottom flask, 1-boc-azetidine-3-yl-methanol (1.96 g, 10.47 mmol) was added and dissolved in DMF (20.0 mL). The stirring solution was cooled to 0 °C and sodium hydride (0.762 g, 19.04 mmol) was then added. The reaction was warmed to room

20

temperature and stirred for 30 minutes, resulting in a pale pink frothy mixture. The mixture was cooled to 0 °C and a solution of 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (2.6 g, 9.52 mmol) in DMF (10 mL) was added via cannula. The reaction was warmed to room temperature and stirred for 3 hours. The reaction was then quenched with 120 mL H₂O and extracted 3 times with 30 mL EtOAc. The organic solution was then washed 3 times with 100 mL brine:H₂O solution (1:1). The organic was dried over Na₂SO₄, filtered, and concentrated to a pale yellow oil. The crude was purified by flash column chromatography using a 0-50% EtOAc:hexanes gradient to yield tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidine-1-carboxylate (1.95 g, 4.60 mmol, 48 % yield).

Preparation 13: tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidine-1-carboxylate




In a round bottom flask, tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidine-1-carboxylate (1.95 g, 4.60 mmol) was dissolved in 1,4-dioxane (23.0 mL). 4-hydroxyphenylboronic acid (0.95 g, 6.90 mmol) was added followed by water (7.7 mL). Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (0.56 g, 0.69 mmol) was added and the flask was equipped with a reflux condenser, set under nitrogen, heated to 110 °C, and stirred overnight. The reaction was then cooled to room temperature and the organic solvent was removed *in vacuo*. H₂O (20 mL) was added to the residue and was extracted three times with DCM (30 mL). The organic was dried over Na₂SO₄, filtered, and concentrated to a black oil. The crude was purified by flash column chromatography using a 0-100% EtOAc:hexanes gradient to yield tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidine-1-carboxylate (2.03

20

25


g, 4.22 mmol, 92 % yield) as a yellow/orange oil. (*m/z*): [M+H]⁺ calculated for C₂₅H₃₁N₅O₅ 482.2 found 482.2.

Preparation 14: 4-(4-(azetidin-3-ylmethoxy)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)phenol



In a round bottom flask, tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidine-1-carboxylate (2.03 g, 4.22 mmol) was dissolved in DCM (10.0 ml) and TFA (5.0 ml) was slowly added at room temperature. The reaction was stirred for 18 hours to yield an orange/purple solution. The reaction was concentrated *in vacuo* to yield a grey/orange solid. The title compound was isolated as the TFA salt in 100% yield (1.25 g, 4.22 mmol). (*m/z*): [M+H]⁺ calculated for C₁₅H₁₅N₅O₂ 298.1 found 298.3.

Example 4: 1-(3-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)azetidin-1-yl)prop-2-en-1-one




In a round bottom flask, 4-(4-(azetidin-3-ylmethoxy)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)phenol trifluoroacetic acid (1.25 g, 4.20 mmol) was dissolved in DMF (21.0 ml) and diisopropylethylamine (3.67 ml, 21.0 mmol) was added. The reaction was cooled to 0 °C and stirred for 5 minutes before the slow addition of acryloyl chloride (0.273 ml, 3.36 mmol), it was then stirred for 15 minutes at 0 °C. The reaction was

warmed to room temperature and concentrated *in vacuo*. The crude was dissolved in 1:1 MeCN:H₂O and purified by reverse phase preparatory HPLC using a 20-80% MeCN:H₂O gradient. The desired fractions were combined and lyophilized to give the title compound (434 mg, 1.235 mmol, 29 % yield). (*m/z*): [M+H]⁺ calculated for C₁₈H₁₇N₅O₃ 352.1

5 found 352.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.32 – 8.23 (d appt, 2H), 8.07 (s, 1H), 6.89 – 6.81 (d appt, 2H), 6.30 (dd, *J* = 17.0, 10.3 Hz, 1H), 6.07 (dd, *J* = 17.0, 2.3 Hz, 1H), 5.63 (dd, *J* = 10.3, 2.3 Hz, 1H), 4.81 (d, *J* = 6.5 Hz, 2H), 4.36 (t, *J* = 8.6 Hz, 1H), 4.12 (dd, *J* = 8.6, 5.4 Hz, 1H), 4.06 (t, 1H), 3.82 (dd, *J* = 10.2, 5.5 Hz, 1H), 3.24 – 3.10 (m, 1H).


10

Preparation 15: tert-butyl 3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate



1-Boc-azetidine-3-yl-methanol (3.44 g, 18.39 mmol) was added to an oven dried
 15 flask and dissolved in DMF (41.8 ml). The solution was cooled to 0 °C and stirred for 10 minutes before the addition of sodium hydride (0.50 g, 20.89 mmol). The frothy mixture was warmed to room temperature and stirred for 30 minutes. The reaction was then cooled to 0 °C and a solution of 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (2.5 g, 8.36 mmol) in DMF (10 mL) was added via cannula. The reaction was warmed to room
 20 temperature and stirred for 1.5 hours. The reaction was quenched by the addition of 120 mL H₂O and extracted 3 times with 50 mL EtOAc. The organic phase was combined and washed 3 times with 100 mL brine:H₂O solution (1:1). The organic phase was then collected and dried over Na₂SO₄, filtered, and concentrated to a clear, pale yellow oil (3.9 g, 8.36 mmol). The crude product was used without any further purification. (*m/z*): [M+H]⁺
 25 calculated for C₂₁H₂₈BrN₃O₄ 466.14 found 466.1.

Preparation 16: tert-butyl 3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate




5

Tert-butyl 3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (3.9 g, 8.36 mmol) was added to a flask and dissolved in 1,4-dioxane (44.6 ml). 3-fluoro-4-hydroxyphenylboronic acid (1.96 g, 12.54 mmol) was added followed by water (11.2 ml) and potassium phosphate, tribasic (5.33 g, 25.09 mmol). Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (1.02 g, 1.25 mmol) was added and the flask was equipped with a reflux condenser and placed under a nitrogen atmosphere. The reaction was heated to 110 °C and stirred for 18 hours. Once cooled to room temperature, the organic solvent was removed *in vacuo* to yield a black oil. To the oil, 10 mL H₂O was added and the solution was extracted 3 times with 40 mL DCM. The organic was collected and dried over Na₂SO₄, filtered, and concentrated to a black oil. The oil was purified via flash column chromatography using a 0-100% EtOAc:Hex gradient to yield tert-butyl 3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (2.58 g, 5.19 mmol, 62 % yield) as a clear yellow oil. (*m/z*): [M+H]⁺ calculated for C₂₇H₃₂FN₃O₅ 498.2 found 498.2.


15
20

Preparation 17: 4-(4-(azetidin-3-ylmethoxy)-1H-indazol-6-yl)-2-fluorophenol




In a 50 mL round bottom flask, tert-butyl 3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (2.58 g, 5.19 mmol) was dissolved in DCM (10.0 ml). TFA (5.00 ml) was slowly added to the clear yellow solution at room temperature. The reaction was stirred for 24 hours and the resulting dark yellow/purple solution was concentrated *in vacuo* to give the TFA salt of 4-(4-(azetidin-3-ylmethoxy)-1H-indazol-6-yl)-2-fluorophenol (3.46 g) as a grey/yellow solid (100 % yield). (*m/z*): [M+H]⁺ calculated for C₁₇H₁₆FN₃O₂ 314.1 found 314.2.

Example 5: (E)-4-(dimethylamino)-1-(3-(((6-(3-fluoro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidin-1-yl)but-2-en-1-one




(2E)-4-(dimethylamino)but-2-enoic acid (464 mg, 2.80 mmol) and HATU (976 mg, 2.57 mmol) were added to a round bottom flask, dissolved in DMF (5.5 mL) and stirred for 15 minutes. A solution of 4-(4-(azetidin-3-ylmethoxy)-1H-indazol-6-yl)-2-fluorophenol (1.0 g, 2.34 mmol) in DMF (5.5 mL) was then slowly added to the stirring reaction followed by diisopropylethylamine (2.0 ml, 11.70 mmol). The reaction was stirred for 24 hours and was then concentrated *in vacuo*. The crude was dissolved in 1:1 MeCN:H₂O and purified by reverse phase preparatory HPLC using a 20-80% MeCN:H₂O gradient. The desired fractions were combined and lyophilized to give the TFA salt of the title compound (271 mg, 0.50 mmol, 21 % yield). (*m/z*): [M+H]⁺ calculated for C₂₃H₂₅FN₄O₃ 425.2 found 425.2. ¹H NMR (400 MHz, MeOD-*d*₄) δ 7.94 (d, *J* = 0.8 Hz, 1H), 7.39 (dd, *J* = 12.5, 2.2 Hz, 1H), 7.35 – 7.28 (m, 1H), 7.24 (t, *J* = 1.0 Hz, 1H), 7.03 – 6.93 (m, 1H), 6.80 – 6.68 (m, 2H), 6.53 (dt, *J* = 15.2, 1.1 Hz, 1H), 4.53 (t, *J* = 8.7 Hz, 1H), 4.47 – 4.34 (m, 2H), 4.35 – 4.22 (m, 2H), 4.09 (dd, *J* = 10.8, 5.6 Hz, 1H), 3.93 (dd, *J* = 7.2, 1.1 Hz, 2H), 3.27 – 3.19 (m, 1H), 2.88 (s, 6H).

Preparation 18: 6-Bromo-4-nitro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole


A mixture of 6-bromo-4-nitro-1H-indazole (3.00 g, 12.4 mmol), 3,4-dihydro-2H-pyran, 97% (3.40 ml, 37.2 mmol) and p-toluenesulfonic acid monohydrate (pTsOH, 0.236 g, 1.24 mmol) in DCM (41.3 ml) was stirred at rt overnight. The reaction was filtered through a pad of celite and concentrated. The reaction was assumed to be 100% conversion (4.05 g) and was carried into the subsequent reaction without further purification.

Preparation 19: 6-(4-((tert-Butyldimethylsilyl)oxy)phenyl)-4-nitro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole



A mixture of 6-(4-((tert-butyldimethylsilyl)oxy)phenyl)-4-nitro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (1.00 g, 3.07mmol), 4-(tert-butyldimethylsilyloxy)phenylboronic acid (1.16 g, 4.60mmo), and potassium phosphate tribasic (1.30g, 6.13mmol) in dioxane (14.9 mL) and H₂O (4.98 mL) was purged with N₂ for 10mins. [1,1'-bis(diphenylphosphine)ferrocene]dichloropalladium(II) (0.224 g, 0.307mmol) was then added, after which the flask was sealed. The reaction was heated to 110 °C and stirred for 1hr. After confirmed full conversion to the desired product via LCMS, the reaction was quenched with 20mL H₂O and 20mL EtOAc. Both layers were filtered through a pad of celite and transferred to a separatory funnel. The mixture was extracted 3 times with 20mL EtOAc, and the aqueous layer was discarded. The combined organic fractions were concentrated and subsequently purified by flash column chromatography using a 0-10% EtOAc:hexanes gradient. The product was isolated pure as a pale-yellow oil (1.03g, 74.4% yield). (*m/z*): [M+H]⁺ calcd for C₂₄H₃₁N₃O₄Si 454.61 found 454.3.

Preparation 20: 6-(4-((tert-butyldimethylsilyl)oxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-amine




5

6-(4-((tert-butyldimethylsilyl)oxy)phenyl)-4-nitro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (0.980g, 2.16mmol) and palladium 10%wt on carbon (wet) (0.980 g) were dissolved in THF (2.8 mL) and isopropyl alcohol (11.2 mL). The reaction vessel was back-filled with N₂ for 5mins, afterwhich it was sealed and placed under a hydrogen atmosphere. The reaction mixture was left to stir at room temperature for 3hrs. After LCMS indicated full conversion to the desired product, the reaction mixture with filtered. The filter was then washed with additional THF. After concentrating and purifying by flash column chromatography (40g) using a 0-25% EtOAC:hexanes gradient, the product was isolated (0.562 g, 61.5% yield). (*m/z*): [M+H]⁺ calcd for C₂₄H₃₃N₃O₂Si 424.63 found 424.4.

15

Preparation 21: tert-butyl (2R)-2-((6-(4-((tert-butyldimethylsilyl)oxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)carbamoyl)morpholine-4-carboxylate



20


(R)-N-boc-2-morpholinecarboxylic acid (0.819 g, 3.54mmol) and 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxide hexafluorophosphate (HATU, 2.69 g, 7.08 mmol) were dissolved in DMF (11.8 mL) and left to stir at room temperature for 5 mins. 6-(4-((tert-Butyldimethylsilyl)oxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-amine (1.00 g, 2.36 mmol) was then added, and the reaction mixture was left to stir at room temperature for 2hrs. LCMS indicated full

25

conversion to the desired product. The reaction was quenched with 10 mL H₂O and 10 mL DCM. After transferring to a separatory funnel, the mixture was extracted three times with 10mL DCM, and the aqueous layer was discarded. Combined organic fractions were concentrated and subsequently purified by flash column chromatography (40g) using a 0-45% EtOAc:hexanes gradient. The products was isolated as a pale-yellow oil (1.01 g, 67 % yield). (*m/z*): [M+H]⁺ calcd for C₃₄H₄₈N₄O₆Si 637.87 found 637.6.


Preparation 22: (R)-N-(6-(4-hydroxyphenyl)-1H-indazol-4-yl)morpholine-2-carboxamide

10




tert-Buyl (2R)-2-((6-(4-((tert-butyldimethylsilyl)oxy)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)carbamoyl)morpholine-4-carboxylate (1.00 g, 1.58 mmol) was dissolved in MeOH (2.0 mL). While stirring, an HCl solution in 4.0M dioxane (12.1 mL, 48.5 mmol) was added slowly. The reaction mixture was heated to 60°C and stirred for 30mins, after which LCMS indicated full conversion to the desired product. The mixture was concentrated to give the product as an orange oil. The reaction was assumed to be 100% conversion (0.533 g) and was carried into the subsequent reaction without further purification.

Example 6: (R)-4-acryloyl-N-(6-(4-hydroxyphenyl)-1H-indazol-4-yl)morpholine-2-carboxamide



(R)-N-(6-(4-hydroxyphenyl)-1H-indazol-4-yl)morpholine-2-carboxamide (0.533g, 1.58mmol) was dissolved in DMF (7.9 mL). N,N-diisopropylethylamine (DIPEA, 2.75 mL, 15.8 mmol) was added, followed by slow addition of acryloyl chloride (0.0900 mL, 1.10 mmol). The reaction was stirred at room temperature for 10mins before conversion was monitored by LCMS. After full conversion was indicated via LCMS, the reaction mixture was concentrated and purified via reverse-phase HPLC prep using a 10-50% ACN:H₂O gradient. Fractions of the desired product were collected and concentrated via lyophilization. The product was isolated (130 mg, 21 % yield). (*m/z*): [M+H]⁺ calcd for C₂₁H₂₀N₄O₄ 393.42 found 393.4. ¹H NMR (400 MHz, Methanol-*d*₄) δ 8.13 (s, 1H), 7.74 (d, *J* = 16.3 Hz, 1H), 7.56 (d, *J* = 6.0 Hz, 2H), 7.51 (s, 1H), 6.91 (dd, *J* = 8.7, 2.9 Hz, 3H), 6.30 (dt, *J* = 16.8, 2.3 Hz, 1H), 5.84 (d, *J* = 10.6 Hz, 1H), 4.45 – 4.25 (m, 2H), 4.21 (d, *J* = 11.8 Hz, 1H), 4.12-4.02 (m, 1H), 3.78 (t, *J* = 11.5 Hz, 1H), 3.53 (m, 1H), 3.29 – 2.99 (m, 1H).


Preparation 23: tert-butyl (3S)-3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate




Sodium hydride (0.072 g, 3.01 mmol) was added to a solution of (Ss)-N-boc-pyrrolidine-3-methanol (0.444 g, 2.21 mmol) in DMF (6 ml) under an atmosphere of N₂ at 0 °C and the reaction mixture was stirred at rt for 20 minutes. 6-bromo-4-fluoro-1-

(tetrahydro-2H-pyran-2-yl)-1H-indazole (0.600 g, 2.01 mmol) was added and the reaction mixture was stirred at rt for 1 hour. Water (1 mL) was added and the reaction was concentrated *in vacuo*. The crude material was purified via flash column chromatography using 40% ethyl acetate in hexanes to yield tert-butyl (3S)-3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (0.964 g, 1.87 mmol, 93 % yield). (*m/z*): [M+H]⁺ calcd for C₂₂H₃₀BrN₃O₄ 480.15 found 480.1.


Preparation 24: tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate



Palladium acetate (0.084 g, 0.375 mmol) was added to a solution of 4-hydroxyphenylboronic acid (0.388 g, 2.81 mmol), tert-butyl (3S)-3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (0.900 g, 1.87 mmol), 1,1'-bis(di-*t*-butylphosphino)ferrocene (0.178 g, 0.375 mmol) and potassium phosphate (1.193 g, 5.62 mmol) in 1,4-dioxane (12 ml) and water (3.00 ml). The reaction mixture was degassed with nitrogen for 10 minutes and then stirred at 110 °C for 2 hours. The reaction mixture was concentrated *in vacuo* to a volume of about 5mL. A saturated aqueous solution of ammonium chloride (10 mL) was added and the mixture was extracted with methylene chloride (3 x 10 mL). The methylene chloride extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via flash column chromatography using 50% ethyl acetate in hexanes to yield tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (0.800 g, 1.62 mmol, 87 % yield) as a clear yellow liquid. (*m/z*): [M+H]⁺ calcd for C₂₈H₃₅N₃O₅ 494.27 found 494.2.

Preparation 25: (S)-4-(4-(pyrrolidin-3-ylmethoxy)-1H-indazol-6-yl)phenol


4.0N HCl in dioxane (8.10 ml, 32.4 mmol) was added to a solution of tert-butyl
 5 (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (0.800 g, 1.62 mmol) in dioxane (6 ml), and the reaction mixture was stirred at 60 °C for 30 minutes. The reaction mixture was concentrated *in vacuo* to yield ((S)-4-(4-(pyrrolidin-3-ylmethoxy)-1H-indazol-6-yl)phenol as an HCl salt (0.501 g, 1.62 mmol, 100 % yield). (*m/z*): [M+H]⁺ calcd for
 10 C₁₈H₁₉N₃O₂ 310.16 found 310.3.

Example 7: (S)-1-(3-(((6-(3-chloro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)pyrrolidin-1-yl)prop-2-en-1-one

15 N,N-Diisopropylethylamine (2.83 ml, 16.2 mmol) was added to a solution of (S)-4-(4-(pyrrolidin-3-ylmethoxy)-1H-indazol-6-yl)phenol HCl salt (0.560 g, 1.62 mmol) in DMF (5 ml) at 0 °C, followed by acryloyl chloride (0.171 ml, 2.11 mmol). The reaction mixture was stirred at rt for 15 minutes then concentrated to a volume of about 2 mL. The
 20 crude liquid was purified via preparatory scale C18 column chromatography using a gradient of 30-90% acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (226 mg, 0.473 mmol, 29.2 % yield). (*m/z*): [M+H]⁺ calcd for C₂₁H₂₁N₃O₃ 364.17 found 364.2. ¹H NMR (400 MHz, DMSO-d₆) δ 13.00 (s, 1H), 10.31 (s, 1H),


8.04-7.94 (m, 1H), 7.59-7.48 (m, 1H), 7.18 (s, 1H), 6.89-6.79 (m, 1H), 6.76-6.70 (m, 1H), 6.65-6.54 (m, 1H), 6.13 (dd, J = 16.6, 1.7 Hz, 1H), 5.69-5.62 (m, 1H), 4.25-3.78 (m, 4H), 3.76-3.28 (m, 2H), 2.88-2.63 (m, 1H), 2.22-1.72 (m, 2H).

5 **Preparation 26: tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate**



To a dry scintillation vial purged with N₂, tert-butyl 3-cyano-3-(hydroxymethyl)azetidine-1-carboxylate (851 mg, 4.01 mmol) was added, dissolved in 5 mL DMF and the solution was cooled to 0 °C. 60% by weight sodium hydride in mineral oil (267 mg, 6.69 mmol) was slowly added to the stirring solution and the reaction was allowed to warm to rt following the addition. The frothy reaction was stirred for 30 minutes before being cooled once again to 0 °C. A solution of 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (800 mg, 2.67 mmol) in 2 mL DMF was slowly added into the scintillation vial containing the organosodium solution. Following the addition, the reaction was warmed to rt and stirred for 2 hours upon which LCMS indicated full conversion of the starting material to the desired product. The reaction was quenched with the slow addition of 2 mL H₂O and 2 mL EtOAc, which was allowed to stir for 5 minutes. The biphasic solution was then transferred to a separatory funnel and an additional 10 mL H₂O were added. The mixture was extracted with 3 times 20 mL of EtOAc and the aqueous layer was discarded. The combined organic fractions were then washed with 3x10 mL 1:1 H₂O:brine to remove residual DMF. The organic was then dried over Na₂SO₄, filtered, and concentrated to a lightly yellow oil. The oil was then purified by flash column chromatography using a 0-40% EtOAc:hexanes gradient. The product was isolated pure as a clear, colorless, viscous oil (1.096 g, 83% yield). (m/z): [M+H]⁺ calcd for C₂₂H₂₇BrN₄O₄ 491.13 found 491.1.

Preparation 27: tert-butyl 3-cyano-3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate




5

A 1:1 mixture of Palladium acetate (25.1 mg, 0.112 mmol) and 1,1'-Bis(di-*t*-butylphosphino)ferrocene 98% (53.1 mg, 0.112 mmol) was added to a solution of 3-fluoro-4-hydroxybenzeneboronic acid (262 mg, 1.679 mmol), tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (550 mg, 1.119 mmol), and potassium phosphate (713 mg, 3.36 mmol) in 1,4-dioxane (4.5 ml) and water (1 ml). The reaction mixture was degassed with nitrogen for 10 minutes and then stirred at 110 °C for 2 hours. The reaction mixture was concentrated *in vacuo*. A saturated aqueous solution of ammonium chloride (5 mL) was added and the mixture was extracted with ethyl acetate (2 x 5 mL). The ethyl acetate extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via flash column chromatography using 40% ethyl acetate in hexanes to yield tert-butyl 3-cyano-3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (267 mg, 0.511 mmol, 46 % yield). (*m/z*): [M+H]⁺ calcd for C₂₈H₃₁FN₄O₅ 523.24 found 523.4.

20


Preparation 28: 3-(((6-(3-fluoro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidine-3-carbonitrile



25


Tert-butyl 3-cyano-3-((((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)methyl)azetidine-1-carboxylate (267 mg, 0.511 mmol) was dissolved in dichloromethane (1 ml) and TFA (1 ml) was slowly added. The clear solution was stirred at rt for 5 hours upon which LCMS indicated good conversion to the
 5 desired product. The reaction was concentrated down to provide 3-((((6-(3-fluoro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidine-3-carbonitrile as a TFA salt (100% yield). (*m/z*): [M+H]⁺ calcd for C₁₈H₁₅FN₄O₂ 339.13 found 339.2.

Example 8: (E)-1-(4-(dimethylamino)but-2-enoyl)-3-((((6-(3-fluoro-4-hydroxyphenyl)-1Hindazol-4-yl)oxy)methyl)azetidine-3-carbonitrile
 10



HATU (2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium, 121 mg, 0.318 mmol) was added to a solution of (E)-4-(dimethylamino)but-2-enoic acid hydrochloride (41.1 mg, 0.318 mmol) in DMF (2 mL). The reaction mixture was stirred at
 15 rt for 5 minutes then 3-((((6-(3-fluoro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)methyl)azetidine-3-carbonitrile TFA salt (160 mg, 0.354 mmol) was added, followed by DIPEA (0.618 ml, 3.54 mmol). The reaction mixture was stirred at rt for 15 minutes and then concentrated *in vacuo* to yield a yellow liquid. The crude liquid was purified via preparatory scale Zorbax Bonus-RP (2.1x30mm, 1.8 micron) column
 20 chromatography using 15-75% acetonitrile in water with 0.05% trifluoroacetic acid to yield (E)-1-(4-(dimethylamino)but-2-enoyl)-3-((((6-(3-fluoro-4-hydroxyphenyl)-1Hindazol-4-yl)oxy)methyl)azetidine-3-carbonitrile (56.2 mg, 0.121 mmol, 34 % yield). (*m/z*): [M+H]⁺ calcd for C₂₄H₂₄FN₅O₃ 450.19 found 450.0. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.94 (s, 1H), 9.77 (s, 1H), 7.85 (s, 1H), 7.47 (dd, *J* = 12.8, 2.2 Hz, 1H), 7.31 (dd, *J* =
 25 8.4, 1.7 Hz, 1H), 7.22 (s, 1H), 6.99 – 6.90 (m, 1H), 6.78 (s, 1H), 6.59 – 6.48 (m, 1H), 6.31 (d, *J* = 15.4 Hz, 1H), 4.67 – 4.60 (m, 3H), 4.40 (d, *J* = 9.1 Hz, 1H), 4.27 (d, *J* = 10.5 Hz, 1H), 4.12 (d, *J* = 10.4 Hz, 1H), 3.78 (d, *J* = 6.8 Hz, 2H), 2.66 (s, 6H).


Preparation 29: tert-butyl 4-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)-4-methylpiperidine-1-carboxylate



To a dry scintillation vial purged with N₂, 1-boc-4-methylpiperidin-4-ol (567 mg, 2.63 mmol) was added, dissolved in 3 mL DMF and the solution was cooled to 0 °C. 60% by weight sodium hydride in mineral oil (175 mg, 4.39 mmol) was slowly added to the stirring solution and the reaction was allowed to warm to rt following the addition. The frothy reaction was stirred for 30 minutes before being cooled once again to 0 °C. A solution of 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (525 mg, 1.755 mmol) in 1 mL DMF was slowly added into the scintillation vial containing the organosodium solution. Following the addition, the reaction was warmed to rt and stirred for 2 hours upon which LCMS indicated full conversion of the starting material to the desired product. The reaction was quenched with the slow addition of 2 mL H₂O and 2 mL EtOAc, which was allowed to stir for 5 minutes. The biphasic solution was then transferred to a separatory funnel and an additional 10 mL H₂O were added. The mixture was extracted with 3 times 20 mL of EtOAc and the aqueous layer was discarded. The combined organic fractions were then washed with 3x10 mL 1:1 H₂O:brine to remove residual DMF. The organic was then dried over Na₂SO₄, filtered, and concentrated to a lightly yellow oil. The oil was then purified by flash column chromatography using a 0-40% EtOAc:hexanes gradient. The product was isolated pure as a clear, colorless, viscous oil (389 mg, 45% yield). (*m/z*): [M+H]⁺ calcd for C₂₃H₃₂BrN₃O₄ 494.17 found 494.4.


Preparation 30: tert-butyl 4-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)-4-methylpiperidine-1-carboxylate

25



A 1:1 mixture of Palladium acetate (9.1 mg, 0.040 mmol) and 1,1'-Bis(di-
 5 butylphosphino)ferrocene 98% (19.2 mg, 0.040 mmol) was added to a solution of 4-
 hydroxybenzeneboronic acid (42 mg, 0.30 mmol), tert-butyl 4-((6-bromo-1-(tetrahydro-
 2H-pyran-2-yl)-1H-indazol-4-yl)oxy)-4-methylpiperidine-1-carboxylate (100 mg, 0.202
 mmol), and Potassium phosphate, tribasic, 97%, anhydrous (129 mg, 0.607 mmol) in 1,4-
 Dioxane (2 ml) and Water (0.5 ml). The reaction mixture was degassed with nitrogen for
 10 minutes and then stirred at 110 °C for 2 hours. The reaction mixture was concentrated
in vacuo. A saturated aqueous solution of ammonium chloride (5 mL) was added and the
 mixture was extracted with ethyl acetate (2 x 5 mL). The ethyl acetate extracts were
 combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid.
 The crude liquid was purified via combiflash column chromatography (12g) using 40%
 ethyl acetate in hexanes to yield tert-butyl 4-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-
 15 pyran-2-yl)-1H-indazol-4-yl)oxy)-4-methylpiperidine-1-carboxylate (96 mg, 0.189 mmol,
 94 % yield). (*m/z*): [M+H]⁺ calcd for C₂₉H₃₇N₃O₅ 5098.28 found 508.3.

Preparation 31: 4-(4-((4-methylpiperidin-4-yl)oxy)-1H-indazol-6-yl)phenol




Tert-butyl 4-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-
 yl)oxy)-4-methylpiperidine-1-carboxylate (96 mg, 0.189 mmol) was dissolved in 1,4-
 dioxane (1 ml) and 4N HCl in 1,4-dioxane (1.18 ml, 4.73 mmol) was slowly added. The
 25 clear solution was stirred at rt for 5 hours upon which LCMS indicated good conversion
 to the desired product. The reaction was concentrated down to provide 4-(4-((4-

methyloperidin-4-yl)oxy)-1H-indazol-6-yl)phenol as an HCl salt (100% yield).

(*m/z*): [M+H]⁺ calcd for C₁₉H₂₁N₃O₂ 324.17 found 324.3.


Example 9: (E)-4-(dimethylamino)-1-(4-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)-4-methylpiperidin-1-yl)but-2-en-1-one



5 HATU (2-(7-aza-1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate, 72 mg, 0.189 mmol) was added to a solution of (E)-4-(dimethylamino)but-2-enoic acid hydrochloride (24 mg, 0.186 mmol) in DMF (1 mL). The reaction mixture was stirred at rt for 5 minutes then 4-(4-((4-methylpiperidin-4-yl)oxy)-1H-indazol-6-yl)phenol HCl salt (68.0 mg, 0.189 mmol) was added, followed by
 10 DIPEA (0.330 ml, 1.890 mmol). The reaction mixture was stirred at rt for 15 minutes and then concentrated *in vacuo* to yield a yellow liquid. The crude liquid was purified via preparatory scale Zorbax Bonus-RP (2.1x30mm, 1.8 micron) column chromatography using 5-65% acetonitrile in water with 0.05% trifluoroacetic acid to yield (E)-4-(dimethylamino)-1-(4-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)-4-methylpiperidin-1-yl)but-2-en-1-one (39.0 mg, 0.063 mmol, 34 % yield). (*m/z*): [M+H]⁺ calcd for
 15 C₂₅H₃₀FN₄O₃ 435.24 found 435.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.99 (s, 1H), 9.66 (s, 1H), 8.02 (d, *J* = 0.9 Hz, 1H), 7.52 – 7.44 (m, 2H), 7.27 (s, 1H), 6.90 (d, *J* = 15.0 Hz, 1H), 6.86 – 6.78 (m, 3H), 6.58 – 6.47 (m, 1H), 3.99 – 3.86 (m, 1H), 3.83 (d, *J* = 7.2 Hz, 2H), 3.78 – 3.55 (m, 2H), 2.74 (s, 6H), 1.40 (s, 3H).


20

Preparation 32: 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridine




3,4-Dihydro-2H-pyran (2.0 g, 24.07 mmol) was added to a solution of 4,6-dichloro-1H-pyrazolo[3,4-b]pyridine (2.50 g, 13.23 mmol) in methylene chloride (30 mL), followed by p-toluenesulfonic acid monohydrate (0.275 g, 1.60 mmol), and the reaction mixture was stirred at rt for 1 h. Water (50 mL) was added and the mixture was extracted with methylene chloride (3 x 50 mL). The methylene chloride extracts were combined, washed with brine (1 x 50 mL), dried over sodium sulfate and concentrated to yield a clear yellow liquid. The crude liquid was purified via flash column chromatography using 10% EtOAc in hexanes to yield 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridine (3.0 g, 11.1 mmol, 67 % yield). (m/z): [M+H]⁺ calculated for C₁₁H₁₁Cl₂N₃O 272.04 found 272.21.

Preparation 33: tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate




NaH (60% dispersion in mineral oil, 160 mg, 4.0 mmol) was added to a solution of 1-boc-azetidine-3-yl-methanol (760 mg, 4.0 mmol) in DMF (5 mL) and diethyl ether (5 mL) at 0°C and the reaction mixture was stirred at 0 °C for 15 minutes. 4,6-Dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridine (1.0 g, 3.7 mmol) was added and the reaction mixture was stirred at rt for 1 h. Water (10 mL) was added and the reaction mixture was extracted with methylene chloride (3 x 10 mL). The methylene chloride extracts were combined, washed with brine (1 x 10 mL), dried over sodium sulfate and concentrated *in vacuo* to yield a clear yellow liquid. The crude liquid was purified via flash column chromatography using 30% EtOAc in hexanes to yield tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate (1.1 g, 2.60 mmol, 70 % yield). (m/z): [M+H]⁺ calculated for C₂₀H₂₇ClN₄O₄ 423.18 found 423.25.

Preparation 34: tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate



A solution of sodium carbonate (500 mg, 4.72 mmol) in water (4 mL) was added to 4-hydroxyphenylboronic acid (392 mg, 2.84 mmol) and tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate (1.0 g, 2.36 mmol) in dioxane (20 mL) and the reaction mixture was degassed with nitrogen for 10 minutes. PdCl₂(dppf).DCM (196 mg, 0.24 mmol) was added and the reaction mixture was degassed further for 5 minutes and then stirred at 110 °C for 4 hours. The reaction mixture was cooled to rt and then filtered through a bed of Celite. The filtered material was washed with methylene chloride and the filtrates were combined and washed with water (1 x 10 mL), brine (1 x 10 mL), dried over sodium sulfate and concentrated *in vacuo* to yield a clear brown liquid. The crude liquid was purified via flash column chromatography using 50% EtOAc in hexanes to yield tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate (1.02 g, 2.12 mmol, 76 % yield). (m/z): [M+H]⁺ calculated for C₂₆H₃₂N₄O₅ 481.25 found 481.65.


Preparation 35: 4-(4-(azetidin-3-ylmethoxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol



TFA (4.0 ml) was added slowly to a solution of tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate (500 mg, 1.04 mmol) in methylene chloride (10 mL) at 0 °C and the reaction mixture was stirred at rt for 3 hours. The reaction mixture was concentrated *in vacuo* and the resulting solid was triturated with 5% methanol:methylene chloride to yield the title

compound as the TFA salt (330 mg, 0.804 mmol, 77 % yield). (m/z): [M+H]⁺ calculated for C₁₆H₁₆N₄O₂ 297.14 found 297.15.

Example 10: 1-(3-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidin-1-yl)prop-2-en-1-one




To a solution of 4-(4-(azetidin-3-ylmethoxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol TFA salt (35.5 mg, 0.087 mmol) in DMF (1 mL) was added DIPEA (0.151 mL, 0.865 mmol), followed by the addition of acryloyl chloride (7.0 μ L, 0.087 mmol). The reaction mixture was stirred at rt for 15 minutes and then concentrated *in vacuo* to yield a yellow liquid.

The crude liquid was purified via preparatory scale Zorbax Bonus-RP (2.1x30mm, 1.8 micron) column chromatography using 5-75% acetonitrile in water with 0.05% trifluoroacetic acid to yield 1-(3-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidin-1-yl)prop-2-en-1-one (13.4 mg, 0.028 mmol, 33 % yield).

(m/z): [M+H]⁺ calcd for C₁₉H₂₈N₄O₃ 351.15 found 351.1. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.02 (s, 1H), 9.80 (s, 1H), 7.98 (d, *J* = 8.9 Hz, 3H), 7.13 (s, 1H), 6.85 (d, *J* = 8.7 Hz, 2H), 6.32 (dd, *J* = 17.0, 10.3 Hz, 1H), 6.09 (dd, *J* = 17.0, 2.3 Hz, 1H), 5.64 (dd, *J* = 10.3, 2.3 Hz, 1H), 4.55 (d, *J* = 6.7 Hz, 1H), 4.38 (t, *J* = 8.5 Hz, 1H), 4.15 – 4.02 (m, 2H), 3.80 (dd, *J* = 10.0, 5.4 Hz, 2H), 3.22 – 3.07 (m, 1H).


Preparation 36: 6-bromo-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole



3,4-Dihydro-2H-pyran (19.79 mL, 216.5 mmol) was added to a solution of 6-bromo-4-chloro-1H-indazole (10.0 g, 43.3 mmol) in ethyl acetate (200 mL) at 0 °C, followed by p-toluenesulfonic acid monohydrate (1.64 g, 8.66 mmol), and the reaction mixture was stirred at 50 °C for 4 h. Saturated aqueous sodium bicarbonate (100 mL) was added and


the mixture was extracted with ethyl acetate (3 x 100 mL). Ethyl acetate extracts were combined, washed with water (1 x 100 mL), brine (1 x 100 mL), dried over sodium sulfate and concentrated to yield a yellow solid. The crude liquid was purified via trituration in n-pentane to yield 6-bromo-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (13.0 g, 41.2 mmol, 95 % yield). (m/z): [M+H]⁺ calculated for C₁₂H₁₂BrClN₂O 314.99 found 314.93.

Preparation 37: 4-(4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-6-yl)phenol



A solution of sodium carbonate (8.77 g, 82.8 mmol), 4-hydroxyphenylboronic acid (6.85g, 49.7 mmol) and 6-bromo-4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (13.0 g, 41.4 mmol) in dioxane (200 mL) and water (50 mL) was degassed with nitrogen for 15 minutes. PdCl₂(dppf).DCM (3.37 g, 4.14 mmol) was added and the reaction mixture was degassed further for 5 minutes and then stirred at 125 °C for 2 hours. The reaction mixture was cooled to rt, ethyl acetate was added and the reaction mixture was filtered through a bed of Celite. Water was added (100 mL) and the mixture was extracted with ethyl acetate. The ethyl acetate extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a clear brown liquid. The crude liquid was purified via flash column chromatography using a 0-30% EtOAc in hexanes gradient to yield 4-(4-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-6-yl)phenol (7.5 g, 22.9 mmol, 55 % yield). (m/z): [M+H]⁺ calculated for C₁₈H₁₇ClN₂O₂ 329.11 found 329.08.


25 **Preparation 38: tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)(methyl)amino)methyl)azetidine-1-carboxylate**



A 1:2 mixture of palladium acetate (68.3 mg, 0.304 mmol) and 2-(di-
 butylphosphino)biphenyl (182.0 mg, 0.608 mmol) was added to a solution of 4-(4-chloro-
 1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-6-yl)phenol (500 mg, 1.521 mmol), tert-butyl
 3-((methylamino)methyl)azetidine-1-carboxylate (457 mg, 2.281 mmol), and sodium tert-
 5 butoxide, >99.9% (731 mg, 7.60 mmol) in toluene (15 ml). The reaction mixture was
 degassed with nitrogen for 10 minutes and then stirred at 110 °C for 16 hours. The
 reaction mixture was concentrated *in vacuo* to a volume of about 5 mL. A saturated
 aqueous solution of ammonium chloride (5 mL) was added and the mixture was extracted
 with ethyl acetate (2 x 5 mL). The ethyl acetate extracts were combined, dried over
 10 sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid was
 purified via combiflash column chromatography (24g) using 50% ethyl acetate in hexanes
 to yield tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-
 yl)(methylamino)methyl)azetidine-1-carboxylate (368 mg, 0.747 mmol, 49 % yield).
 (m/z): [M+H]⁺ calcd for C₂₈H₃₆N₄O₄ 493.28 found 493.4.


15

Preparation 39: 4-(4-((azetidin-3-ylmethyl)(methyl)amino)-1H-indazol-6-yl)phenol




20 Tert-butyl 3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-
 yl)(methylamino)methyl)azetidine-1-carboxylate (368 mg, 0.747 mmol) was dissolved in
 Dichloromethane (1.5 ml) and TFA (1.5 ml) was slowly added. The clear solution was
 stirred at rt for 5 hours upon which LCMS indicated good conversion to the desired
 product. The reaction was concentrated down to provide 4-(4-((azetidin-3-
 25 ylmethyl)(methyl)amino)-1H-indazol-6-yl)phenol as a TFA salt (100 % yield).
 (m/z): [M+H]⁺ calcd for C₁₈H₂₀N₄O 309.17 found 309.2.

Example 11: (E)-4-(dimethylamino)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)(methyl)amino)methyl)azetidin-1-yl)but-2-en-1-one



- 5 HATU (2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate, 59.5 mg, 0.157 mmol) was added to a solution of (E)-4-(dimethylamino)but-2-enoic acid hydrochloride (34 mg, 0.263 mmol) in DMF (1 mL). The reaction mixture was stirred at rt for 5 minutes then 4-(4-((azetidin-3-ylmethyl)(methyl)amino)-1H-indazol-6-yl)phenol TFA salt (63 mg, 0.149 mmol) was
- 10 added, followed by DIPEA (0.260 ml, 1.491 mmol). The reaction mixture was stirred at rt for 15 minutes and then concentrated *in vacuo* to yield a yellow liquid. The crude liquid was purified via preparatory scale Zorbax Bonus-RP (2.1x30mm, 1.8 micron) column chromatography using 5-65% acetonitrile in water with 0.05% trifluoroacetic acid to yield
- 15 (E)-4-(dimethylamino)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)(methyl)amino)methyl)azetidin-1-yl)but-2-en-1-one (47.6 mg, 0.086 mmol, 96 % yield). (m/z): [M+H]⁺ calcd for C₂₄H₂₉N₅O₂ 420.24 found 420.2.

Preparation 40: tert-butyl (3S)-3-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)thio)pyrrolidine-1-carboxylate




To a stirred solution of 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (600 mg, 2.0 mmol) and tert-butyl (S)-3-mercaptopyrrolidine-1-carboxylate (488 mg, 2.20 mmol) in DMSO (5 mL) was added Cs₂CO₃ (1.30 mg, 4.0 mmol). The reaction mixture was stirred at 120 °C for 16 hours. Water was added and the reaction mixture was
 5 extracted with EtOAc (3 × 50 ml). The EtOAc extracts were combined, washed with water followed by brine, dried over anhydrous sodium sulphate, filtered and concentrated *in vacuo* to get the crude intermediate product as clear viscous liquid.

To a stirred solution of the crude intermediate in dioxane (20 mL) and water (4 ml) was added 4-hydroxyphenylboronic acid (240 mg, 1.74 mmol) and Na₂CO₃ (461 mg, 4.35
 10 mmol) and the reaction mixture was degassed with nitrogen for 15 minutes.

Pd(dppf)Cl₂.DCM (118 mg, 0.0145 mmol) was added and the reaction mixture was degassed further for 5 minutes and then stirred at 110 °C for 4 hours. The reaction mixture was cooled to rt, water was added and the reaction mixture was extracted with ethyl acetate (3 x 50 mL). The ethyl acetate extracts were combined, dried over sodium
 15 sulfate and concentrated in vacuo to yield a clear brown liquid. The crude liquid was purified via flash column chromatography using a 30-50% EtOAc in hexanes gradient to yield tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)thio)pyrrolidine-1-carboxylate (600 mg, 1.21 mmol, 61%). (m/z): [M+H]⁺ calculated for C₂₇H₃₃ClN₃O₄S 496.23 found 496.42.


Preparation 41: (S)-4-(4-(pyrrolidin-3-ylthio)-1H-indazol-6-yl)phenol



4.0N HCl in dioxane (10.0 ml, 40.0 mmol) was added to a solution of tert-butyl (3S)-3-
 25 (((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)thio)pyrrolidine-1-carboxylate (400 mg, 0.808 mmol) in methanol (6 ml), and the reaction mixture was stirred at 40 °C for 16 h. The reaction mixture was concentrated *in vacuo* and the resulting residue was purified via preparatory scale C18 column chromatography using a


gradient of 20-80% acetonitrile in water with 0.05% trifluoroacetic acid to yield to yield (S)-4-(4-(pyrrolidin-3-ylthio)-1H-indazol-6-yl)phenol as an TFA salt (175 mg, 0.411 mmol, 51 % yield). (*m/z*): [M+H]⁺ calcd for C₁₇H₁₇N₃OS 312.12 found 312.22.

5 **Example 12: (S)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)thio)pyrrolidin-1-yl)prop-2-en-1-one**



N,N-Diisopropylethylamine (0.079 ml, 0.450 mmol) was added to a solution of (S)-4-(4-(pyrrolidin-3-ylthio)-1H-indazol-6-yl)phenol TFA salt (38.5 mg, 0.091 mmol) in DMF (0.2 ml) at 0 °C, followed by acryloyl chloride (8.0 μL, 0.099 mmol). The reaction mixture was stirred at rt for 15 minutes and the crude liquid was purified via preparatory scale C18 column chromatography using a gradient of acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-1-(3-(((6-(4-hydroxyphenyl)-1H-indazol-4-yl)thio)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (17.1 mg, 0.046 mmol, 51 % yield). (*m/z*): [M+H]⁺ calcd for C₂₀H₁₉N₃O₂S 366.13 found 366.1.


Preparation 42: tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)sulfonyl)pyrrolidine-1-carboxylate



Basic alumina (200 mg) and Oxone (744 mg, 2.42 mmol) were added to a solution of tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)thio)pyrrolidine-1-carboxylate (400 mg, 0.80 mmol) in chloroform (50 mL) and water


(1 mL) and the reaction mixture was stirred at 65 °C for 12 h. The reaction mixture was filtered through a bed of Celite, and the filtered material was washed with chloroform. The filtrates were combined, water was added and the mixture was extracted with chloroform (3 x 100 mL). The chloroform extracts were combined, washed with brine, dried over sodium sulfate and concentrated in vacuo. The crude liquid was purified via flash column chromatography using a 30% EtOAc in hexanes to yield tert-butyl (3S)-3-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)sulfonyl)pyrrolidine-1-carboxylate (200 mg, 0.379 mmol, 47 % yield). (m/z): [M+H]⁺ calculated for C₂₇H₃₃ClN₃O₆ 528.22 found 528.48.

Preparation 43: (S)-4-(4-(pyrrolidin-3-ylsulfonyl)-1H-indazol-6-yl)phenol




4.0N HCl in dioxane (10.0 ml, 40.0 mmol) was added to a solution of (3S)-3-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)sulfonyl)pyrrolidine-1-carboxylate (200 mg, 0.379 mmol) in methanol (2 ml), and the reaction mixture was stirred at rt for 8 h. The reaction mixture was concentrated *in vacuo* and the resulting residue was triturated with diethyl ether to yield ((S)-4-(4-(pyrrolidin-3-ylsulfonyl)-1H-indazol-6-yl)phenol as an HCl salt (140 mg, 0.369 mmol, 97 % yield). (m/z): [M+H]⁺ calcd for C₁₇H₁₇N₃O₃S 344.11 found 344.04.

Example 13: ((S)-1-(3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)sulfonyl)pyrrolidin-1-yl)prop-2-en-1-one



N,N-Diisopropylethylamine (0.079 ml, 0.450 mmol) was added to a solution of (((S)-4-(4-(pyrrolidin-3-ylsulfonyl)-1H-indazol-6-yl)phenol HCl salt (31.0 mg, 0.082 mmol) in DMF (0.2 ml) at 0 °C, followed by acryloyl chloride (8.0 µL, 0.099 mmol). The reaction mixture was stirred at rt for 10 minutes and the crude liquid was purified via preparatory scale C18 column chromatography using a gradient of acetonitrile in water with 0.05% trifluoroacetic acid to yield ((S)-1-(3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)sulfonyl)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (3.1 mg, 0.0061 mmol, 7 % yield). (*m/z*): [M+H]⁺ calcd for C₂₀H₁₉N₃O₄S 398.12 found 398.0.

Preparation 44: tert-butyl 4-((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)methyl)piperidine-1-carboxylate




15

A 0.5 M solution of 9-BBN in THF (7.92 mL, 3.96 mmol) was added to tert-butyl 4-methylenepiperidine-1-carboxylate (468 mg, 2.37 mmol) and the reaction mixture was stirred at 80 °C for 2 h. The reaction mixture was cooled to rt and then cannulated into a preformed solution of 4-bromo-6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (500 mg, 1.58 mmol), potassium carbonate (548 mg, 3.96 mmol) and Pd(dppf)Cl₂.DCM (129 mg, 0.15 mmol) in DMF (10.0 ml) & H₂O (2.0 ml). The reaction mixture was stirred at 80 °C for 16 h. The reaction mixture was filtered through a bed of Celite, and the filtered


20

material was washed with ethyl acetate. The filtrates were combined, water (100 mL) was added and the mixture was extracted with ethyl acetate (2 x 150 mL). The ethyl acetate extracts were combined, washed with brine, dried over sodium sulfate and concentrated in vacuo. The crude liquid was purified via flash column chromatography using a 30% EtOAc in hexanes to yield tert-butyl 4-((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)methyl)piperidine-1-carboxylate (460 mg, 1.06 mmol, 67 % yield). (m/z): [M+H]⁺ calculated for C₂₃H₃₂ClN₃O₃ 434.22 found 434.47.


Preparation 45: tert-butyl 4-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)methyl)piperidine-1-carboxylate



4-Hydroxyphenylboronic acid (195 mg, 1.41 mmol) was added to a suspension of tert-butyl 4-((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)methyl)piperidine-1-carboxylate (410 mg, 0.94 mmol) in dioxane (10.0 mL) and water (2.0 mL), followed by potassium phosphate (401 mg, 1.88 mmol) and the reaction mixture was degassed with nitrogen for 15 minutes. 2'-(Dimethylamino)-2-biphenyl-palladium(II) chloride dinorbornylphosphine complex (52 mg, 0.07 mmol) was added and the reaction mixture was heated under microwave irradiation at 100 °C for 1 hour. The reaction mixture was cooled to rt and then filtered through a bed of Celite. Water (100 mL) was added to the filtrate, which was then extracted with ethyl acetate (3 x 100 mL). The ethyl acetate extracts were combined, washed with water, brine, dried over sodium sulfate and concentrated *in vacuo*. The crude liquid was purified via flash column chromatography using 50% EtOAc in hexanes to yield tert-butyl 4-((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)methyl)piperidine-1-carboxylate (300 mg, 0.610 mmol, 65%). (m/z): [M+H]⁺ calculated for C₂₉H₃₇N₃O₄ 492.29 found 492.46.

Preparation 46: 4-(4-(piperidin-4-ylmethyl)-1H-indazol-6-yl)phenol

4.0N HCl in dioxane (10.0 ml, 40.0 mmol) was added to a solution of tert-butyl 4-((6-(4-
 5 hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)methyl)piperidine-1-
 carboxylate (300 mg, 0.610 mmol) in methanol (2 ml), and the reaction mixture was
 stirred at rt for 8 h. The reaction mixture was concentrated *in vacuo* and the resulting
 residue was triturated with diethyl ether to yield 4-(4-(piperidin-4-ylmethyl)-1H-indazol-
 6-yl)phenol as an HCl salt (195 mg, 0.567 mmol, 93 % yield). (*m/z*): [M+H]⁺ calcd for
 10 C₁₉H₂₁N₃O 308.18 found 308.10.


Example 14: 1-(4-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)methyl)piperidin-1-yl)prop-2-en-1-one

15

N,N-Diisopropylethylamine (0.079 ml, 0.450 mmol) was added to a solution of 4-(4-
 (piperidin-4-ylmethyl)-1H-indazol-6-yl)phenol HCl salt (27.5 mg, 0.080 mmol) in DMF
 (0.2 ml) at 0 °C, followed by acryloyl chloride (8.0 μL, 0.099 mmol). The reaction
 mixture was stirred at rt for 10 minutes and the crude liquid was purified via preparatory
 20 scale C18 column chromatography using a gradient of acetonitrile in water with 0.05%
 trifluoroacetic acid to yield 1-(4-((6-(4-hydroxyphenyl)-1H-indazol-4-

yl)methyl)piperidin-1-yl)prop-2-en-1-one as a TFA salt (7.2 mg, 0.015 mmol, 19 % yield). (m/z): $[M+H]^+$ calcd for $C_{22}H_{23}N_3O_2$ 362.19 found 362.1.


Preparation 47: tert-butyl (R,E)-2-(3-ethoxy-3-oxoprop-1-en-1-yl)pyrrolidine-1-carboxylate



Ethyl 3-(triphenylphosphoranylidene)propanoate (3.64 g, 10.0 mmol) was added to a solution of tert-butyl (R)-2-formylpyrrolidine-1-carboxylate (2.0 g, 10.0 mmol) in methylene chloride (40 mL) at 0 °C and the reaction mixture was stirred at rt for 4 h. Water (200 mL) was added and the reaction mixture was extracted with methylene chloride (2 x 100 mL). The methylene chloride extracts were combined, dried over sodium sulfate and concentrated *in vacuo*. The crude liquid was purified via flash column chromatography using 5% ethyl acetate in hexanes to yield tert-butyl (R,E)-2-(3-ethoxy-3-oxoprop-1-en-1-yl)pyrrolidine-1-carboxylate (1.70 g, 6.31 mmol, 63 % yield).

15

Preparation 48: ethyl (R,E)-3-(1-methylpyrrolidin-2-yl)acrylate




4M HCl in dioxane (7.20 mL, 28.8 mmol) was added to a solution of TFA (10.90 mL) was slowly added to a solution of tert-butyl (R,E)-2-(3-ethoxy-3-oxoprop-1-en-1-yl)pyrrolidine-1-carboxylate (1.70 g, 6.31 mmol) in dioxane (40 mL) and the reaction mixture was stirred at rt for 2 h. The reaction was concentrated *in vacuo* to yield the crude intermediate. A 37% solution of formaldehyde in water (4.19 mL, 41.4 mmol) was added to a solution of the crude intermediate in methylene chloride (50 mL) at 0 °C, followed by sodium triacetoxyborohydride (5.26 g, 24.8 mmol), and the reaction mixture was stirred at rt for 16 h. Ice cold water (20 mL) was added and the reaction mixture was extracted with methylene chloride (2 x 100 mL). The methylene chloride extracts were

25


combined, washed with brine, dried over sodium sulfate and concentrated *in vacuo* to yield ethyl (R,E)-3-(1-methylpyrrolidin-2-yl)acrylate (0.900 g, 4.915 mmol, 78 % yield). (*m/z*): [M+H]⁺ calcd for C₁₀H₁₇NO₂ 184.14 found 184.0.

5 Preparation 49: (R,E)-3-(1-methylpyrrolidin-2-yl)acrylic acid




Sodium hydroxide (294 mg, 7.36 mmol) was added to a solution of ethyl (R,E)-3-(1-methylpyrrolidin-2-yl)acrylate (0.900 g, 4.915 mmol, 78 % yield) in 2:1 THF/water (15 mL) and the reaction mixture was stirred at rt for 4 h. The reaction mixture was concentrated *in vacuo* to yield ethyl (R,E)-3-(1-methylpyrrolidin-2-yl)acrylic acid (668 mg, 4.30 mmol, 88% yield). (*m/z*): [M+H]⁺ calcd for C₈H₁₃NO₂ 156.10 found 156.26.

15 Example 15: (E)-1-((S)-3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)-3-((R)-1-methylpyrrolidin-2-yl)prop-2-en-1-one




A solution of HATU (0.031 g, 0.083 mmol) in DMF (0.1 mL) was added to ethyl (R,E)-3-(1-methylpyrrolidin-2-yl)acrylic acid (14.0 mg, 0.090 mmol), followed by a solution of (S)-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol hydrochloride (0.025 g, 0.075 mmol) and DIPEA (0.065 ml, 0.375 mmol), the reaction mixture was stirred at rt for 15 minutes and the crude liquid was purified via preparatory scale C18 column chromatography using a gradient of acetonitrile in water with 0.05% trifluoroacetic acid to yield ((E)-1-((S)-3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)-3-((R)-1-methylpyrrolidin-2-yl)prop-2-en-1-one as a TFA salt (7.8 mg, 0.014 mmol, 19 % yield). (*m/z*): [M+H]⁺ calcd for C₂₅H₂₈N₄O₃ 433.23 found 433.1.

Preparation 50: tert-butyl 2-((dimethylamino)methyl)acrylate


5

Paraformaldehyde (1.20 g, 43.2 mmol) was added to a solution of Malonic acid (2.0 g, 19.3 mmol) in 1,4-dioxane (20 mL), followed by a 2M solution of dimethylamine in THF (9.60 mL, 19.2 mmol) and the reaction mixture was stirred at 70 °C for 1 h. The reaction mixture was concentrated *in vacuo* and the crude product was recrystallized using diethyl ether and acetone to yield the intermediate as a white solid. The intermediate white solid was dissolved in t-BuOH (100 mL) and di-tert-butyl dicarbonate (4.60 mL, 20.7 mmol) was added to the reaction mixture followed by 4-dimethylaminopyridine (511 mg, 4.18 mmol) and the reaction mixture was stirred at rt for 4 h. Methylene chloride (500 mL) was added and the reaction mixture was washed with water (2 x 500 mL), dried over sodium sulfate and concentrated *in vacuo*. The crude residue was purified via flash column chromatography using 2% methanol in methylene chloride to yield tert-butyl 2-((dimethylamino)methyl)acrylate (220 mg, 1.19 mmol, 6% yield).

Preparation 51: 2-((dimethylamino)methyl)acrylic acid


A 1N aqueous solution of HCl (5.0 mL, 5 mmol) was added to tert-butyl 2-((dimethylamino)methyl)acrylate (220 mg, 1.19 mmol) and the reaction mixture was stirred at 100 °C for 15 minutes. The reaction mixture was concentrated *in vacuo* and the resulting residue was azeotroped with toluene and further triturated with diethyl ether to yield 2-((dimethylamino)methyl)acrylic acid as a HCl salt (119 mg, 0.719 mmol, 60% yield). (m/z): [M+H]⁺ calcd for C₆H₁₁NO₂ 130.09 found 130.22.

Example 16: (S)-2-((dimethylamino)methyl)-1-(3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one



- 5 A solution of HATU (0.031 g, 0.083 mmol) in DMF (0.1 mL) was added to 2-((dimethylamino)methyl)acrylic acid hydrochloride (14.9 mg, 0.090 mmol), followed by a solution of (S)-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol hydrochloride (0.025 g, 0.075 mmol) and DIPEA (0.065 ml, 0.375 mmol), the reaction mixture was stirred at rt for 15 minutes and the crude liquid was purified via preparatory scale C18 column chromatography using a gradient of acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-2-((dimethylamino)methyl)-1-(3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (2.6 mg, 0.005 mmol, 7 % yield). (m/z): [M+H]⁺ calcd for C₂₃H₂₆N₄O₃ 407.21 found 407.1.

Example 17: (S)-1-(3-((6-(4-hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-yn-1-one




- 20 A solution of HATU (0.031 g, 0.083 mmol) in DMF (0.1 mL) was added to propiolic acid (6.0 mg, 0.090 mmol), followed by a solution of (S)-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol hydrochloride (0.025 g, 0.075 mmol) and DIPEA (0.065 ml, 0.375 mmol), the reaction mixture was stirred at rt for 15 minutes and the crude liquid was purified via preparatory scale C18 column chromatography using a gradient of acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-1-(3-((6-(4-

hydroxyphenyl)-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-yn-1-one as a TFA salt (2.4 mg, 0.005 mmol, 7 % yield). (*m/z*): [M+H]⁺ calcd for C₂₀H₁₇N₃O₃ 348.14 found 348.0.

Example 18: (S)-4-(4-((1-(vinylsulfonyl)pyrrolidin-3-yl)oxy)-1H-indazol-6-yl)phenol


5



N,N-Diisopropylethylamine (0.052 ml, 0.301 mmol) was added to a solution of (S)-4-(4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol hydrochloride (0.020 g, 0.060 mmol) in DMF (0.2 ml) at 0 °C, followed by ethenesulfonyl chloride (8.4 mL, 0.066 mmol). The reaction mixture was stirred at rt for 10 minutes and the crude liquid was purified via preparatory scale C18 column chromatography using a gradient of acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-4-(4-((1-(vinylsulfonyl)pyrrolidin-3-yl)oxy)-1H-indazol-6-yl)phenol as a TFA salt (8.2 mg, 0.016 mmol, 27 % yield). (*m/z*): [M+H]⁺ calcd for C₁₉H₁₉N₃O₄ 386.12 found 386.3.

15

Preparation 52: tert-butyl 3-(2-((6-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate




Cesium carbonate (6.33 g, 19.42 mmol) was added to a solution of tert-butyl 3-(2-((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (3.20 g, 6.47 mmol) and 3-fluoro-4-hydroxybenzeneboronic acid (1.514 g, 9.71 mmol) in 1,4-dioxane (20.71 ml) and water (5.18 ml) and the reaction mixture was degassed with nitrogen for 10 minutes.

Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) (0.793 g, 0.971 mmol) was added and the reaction mixture was stirred 110 °C for 2 hours. The reaction


mixture was concentrated *in vacuo* to a volume of about 5 mL. A saturated aqueous solution of ammonium chloride (20 mL) was added and the mixture was extracted with methylene chloride (2 x 20 mL). The methylene chloride extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid
 5 was purified via flash column chromatography using 50% ethyl acetate in hexanes to yield tert-butyl 3-(2-((6-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (2.30 g, 4.38 mmol, 68 % yield). (m/z): [M+H]⁺ calcd for C₂₉H₃₇FN₃O₅ 526.27 found 526.3.

10 **Preparation 53: 4-(4-((2-azetidin-3-yl)propan-2-yl)oxy)-1-H-indazole-6-yl)-2-fluorophenol**




15 Tert-butyl 3-(2-((6-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (3.10g, 5.90 mmol) was dissolved in dichloromethane (10 ml) and TFA (10 ml) was slowly added. The clear solution was stirred at rt for 2 hours upon which LCMS indicated good conversion to the desired product. The reaction was concentrated down to provide 4-(4-((2-azetidin-3-yl)propan-2-yl)oxy)-1-H-indazole-6-yl)-2-fluorophenol as a TFA salt (100% yield). (m/z): [M+H]⁺
 20 calcd for C₁₉H₂₁FN₃O₂ 342.16 found 342.3.

Example 19: (E)-1-(3-(2-((6-(3-fluoro-4-hydroxyphenyl)-1H-indazol-4-yl)oxy)propan-2-yl)azetidin-1-yl)-4-(dimethylamino)but-2-en-1-one



HATU (2-(7-aza-1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium
 5 hexafluorophosphate, 2.92 g, 7.07 mmol) was added to a solution of (E)-4-(
 (dimethylamino)but-2-enoic acid hydrochloride (1.07 g, 6.48 mmol) in DMF (10 mL).
 The reaction mixture was stirred at rt for 10 minutes then 4-(4-((2-azetidin-3-yl)propan-2-
 yl)oxy)-1-*H*-indazole-6-yl)-2-fluorophenol TFA salt (2.01 g, 5.89 mmol) was added,
 followed by DIPEA (3.08 ml, 17.7 mmol). The reaction mixture was stirred at rt for 15
 10 minutes and then concentrated *in vacuo* to yield a yellow liquid. The crude liquid was
 purified via preparatory scale C18 column chromatography using a gradient of 20-80%
 acetonitrile in water with 0.05% trifluoroacetic acid to yield (E)-4-(dimethylamino)-1-(3-
 (2-((6-(3-fluoro-4-hydroxyphenyl)-1*H*-indazol-4-yl)oxy)propan-2-yl)cyclobutyl)but-2-en-
 1-one as a TFA salt (1.65 g, 3.65 mmol, 61.9 % yield). (*m/z*): [*M*+*H*]⁺ calcd for
 15 C₂₅H₃₀FN₄O₃ 453.23 found 453.3.


Preparation 54: tert-butyl 3-(2-((2,6-dichloropyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate



20 A solution of tert-butyl 3-(2-hydroxypropan-2-yl)azetidine-1-carboxylate (6.5 g,
 30.21 mmol) in DMSO (10 mL) was added dropwise to a solution of 60% by weight
 sodium hydride in mineral oil (1.65g, 41.20 mmol) in DMSO (15 mL) and the reaction
 mixture was stirred at rt for 10 minutes. A solution of 2,4,6-trichloropyridine (5.0 g ,
 27.62 mmol) in DMSO (25 mL) was added dropwise to the resulting suspension, and the
 25 reaction mixture was stirred at 60 °C for 16 h. The reaction mixture was quenched using

a saturated aqueous solution of ammonium chloride and extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with an aqueous saturated solution of sodium chloride, dried over sodium sulfate and concentrated in *vacuo* to yield the crude liquid. The crude liquid was purified via flash column chromatography using 5% ethyl acetate in hexanes to yield tert-butyl 3-(2-((2,6-dichloropyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (2.10 g, 5.80 mmol, 21.1 % yield). (*m/z*): [M+H]⁺ calculated for C₁₆H₂₃Cl₂N₂O₃ 361.11 found 361.11.

Preparation 55: tert-butyl 3-(2-((6-chloro-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate




A 2M solution of nBuLi in hexanes (3.05 mL, 6.11 mmol) was added dropwise to a solution of tert-butyl 3-(2-((2,6-dichloropyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (2.0 g, 5.55 mmol) in THF (40 mL) at -78 °C. DMF (0.64 mL, 8.33 mmol) was added in one portion and the reaction mixture was stirred at -78 °C for 10 min. 5% Acetic acid in diethyl ether was added and the mixture was further diluted with water. The aqueous layer was extracted with ethyl acetate (3 times). The ethyl acetate extracts were combined, washed with an aqueous saturated solution of sodium chloride, dried over sodium sulfate and concentrated in *vacuo* to yield the crude intermediate tert-butyl 3-(2-((2,6-dichloro-3-formylpyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate.

Hydrazine hydrate (1.0 mL, 20.56 mmol) was added dropwise to a solution of the crude tert-butyl 3-(2-((2,6-dichloro-3-formylpyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate in THF (40 mL) at 0 °C, and the reaction mixture was stirred at rt for 2 h. Triethylamine (1.45 mL, 10.28 mmol) was added, and the reaction mixture was stirred at 60 °C for 16h. The reaction mixture was concentrated in *vacuo* and the resulting residue was diluted with water and extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with an aqueous saturated solution of sodium chloride, dried over sodium sulfate and concentrated in *vacuo* to yield the crude liquid. The crude liquid was purified via flash column chromatography using 20% ethyl acetate in hexanes to yield tert-butyl 3-(2-((6-chloro-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidine-1-


carboxylate (1.20 g, 3.27 mmol, 58.9 % yield). (m/z): $[M+H]^+$ calculated for $C_{17}H_{23}ClN_4O_3$ 367.15 found 367.00.

Preparation 56: tert-butyl 3-(2-((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate



Sodium carbonate (636 mg, 6.00 mmol) was added to a solution of tert-butyl 3-(2-((6-chloro-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (1.1 g, 3.00 mmol) in 4:1 dioxane/water (15 mL), followed by phenylboronic acid (496 mg, 3.60 mmol) and the reaction mixture was degassed with argon for 10 minutes. Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (245 mg, 0.3 mmol) was added, the reaction mixture was degassed further with argon for 5 minutes and the reaction mixture was stirred under microwave irradiation at 130 °C for 1 hour. The reaction mixture was diluted with water and extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with a saturated aqueous solution of sodium chloride, dried over sodium sulfate and concentrated *in vacuo* to yield the crude liquid. The crude liquid was purified via flash column chromatography using 50% ethyl acetate in hexanes to yield tert-butyl 3-(2-((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (650 mg, 1.53 mmol, 51.0 % yield). (m/z): $[M+H]^+$ calcd for $C_{23}H_{29}N_4O_4$ 425.22 found 425.21.


Preparation 57: 4-(4-((2-(azetidin-3-yl)propan-2-yl)oxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol



Tert-butyl 3-(2-((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidine-1-carboxylate (630 mg, 1.48 mmol) was dissolved in dichloromethane (10 ml) and TFA (6.3 ml) was slowly added. The clear solution was


stirred at rt for 4 hours upon which LCMS indicated good conversion to the desired product. The reaction was concentrated in *vacuo* and the residue was triturated with acetonitrile and diethyl ether to yield 4-(4-((2-(azetidin-3-yl)propan-2-yl)oxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol as a TFA salt (100% yield). (*m/z*): [M+H]⁺ calcd for C₁₈H₂₁N₄O₂ 325.17 found 325.13.

Example 20: 1-(3-(2-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidin-1-yl)prop-2-en-1-one



Diisopropylethylamine (0.298 ml, 1.711 mmol) was added to a solution of 4-(4-((2-(azetidin-3-yl)propan-2-yl)oxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol TFA salt (150 mg, 0.342 mmol) in DMF (3.00 ml) at 0 °C, followed by acryloyl chloride (0.031 ml, 0.376 mmol) and the reaction mixture was stirred at rt for 15 min. The reaction mixture was concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via preparatory scale C18 column chromatography using a gradient of 10-70% acetonitrile in water with 0.05% trifluoroacetic acid to yield 1-(3-(2-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)propan-2-yl)azetidin-1-yl)prop-2-en-1-one as a TFA salt (80.5 mg, 0.163 mmol, 47.8 % yield). (*m/z*): [M+H]⁺ calcd for C₂₁H₂₃N₄O₃ 379.18 found 379.3.


Preparation 58: tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidine-1-carboxylate



Sodium hydride (211 mg, 8.79 mmol) was added to a solution of tert-butyl 3-((hydroxymethyl)-3-methylazetidine-1-carboxylate (973 mg, 4.83 mmol) in DMF (4.4

mL) at 0 °C and the reaction mixture was stirred at rt for 30 minutes. The reaction mixture was then added dropwise to a solution of 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidine (1.20 g, 4.39 mmol) in DMF (4.4 mL) at 0 °C and the reaction mixture was stirred at rt for 2 hours. The reaction was quenched with addition of water (1 mL) and the resulting mixture was concentrated *in vacuo* to yield a light brown liquid. Water (10 mL) was added and the mixture was extracted with ethyl acetate (3 x 20 mL). The ethyl acetate extracts were combined, washed with a saturated aqueous solution of sodium chloride (5 mL) and dried over sodium sulfate to yield a brown liquid. The crude liquid was purified via flash column chromatography using 40% ethyl acetate in hexanes to yield tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidine-1-carboxylate (1.06 g, 4.39 mmol, 54.9 % yield). (m/z): [M+H]⁺ calcd for C₂₀H₂₉ClN₅O₄ 438.19 found 438.6.


Preparation 59: tert-butyl 3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidine-1-carboxylate



Potassium phosphate tribasic (218 mg, 1.03 mmol) was added to a solution of tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidine-1-carboxylate (150 mg, 0.343 mmol) in 4:1 dioxane/water (1.71 mL), followed by 3-fluoro-4-hydroxyphenylboronic acid (80.0 mg, 0.514 mmol) and the reaction mixture was degassed with nitrogen for 10 minutes. Palladium acetate (15.4 mg, 0.069 mmol) and 1,1'-bis(di-t-butylphosphino)ferrocene (32.5 mg, 0.069 mmol) were added and the reaction mixture was stirred at 110 °C for overnight. The reaction mixture was concentrated *in vacuo* to a volume of about 5 mL. A saturated aqueous solution of ammonium chloride (20 mL) was added and the mixture was extracted with ethyl acetate (3 x 20 mL). The ethyl acetate extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via flash column chromatography using 40% ethyl acetate in hexanes to yield tert-butyl 3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-


pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidine-1-carboxylate (65 mg, 0.343 mmol, 37.0 % yield). (*m/z*): [*M*+*H*]⁺ calcd for C₂₆H₃₃FN₅O₅ 514.25 found 514.7.

Preparation 60: 2-fluoro-4-(4-((3-methylazetidin-3-yl)methoxy)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)phenol




Tert-butyl 3-(((6-(3-fluoro-4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidine-1-carboxylate (65 mg, 0.127 mmol) was dissolved in dichloromethane (0.56 ml) and TFA (0.56 ml) was added and the reaction mixture was stirred at rt for 4 hours. The reaction was concentrated *in vacuo* to yield 2-fluoro-4-(4-((3-methylazetidin-3-yl)methoxy)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)phenol as a TFA salt (100% yield). (*m/z*): [*M*+*H*]⁺ calcd for C₁₆H₁₇FN₅O₂ 330.14 found 330.3.

Example 21: 1-(3-(((6-(3-fluoro-4-hydroxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidin-1-yl)prop-2-en-1-one




Diisopropylethylamine (0.210 ml, 1.20 mmol) was added to a solution of 2-fluoro-4-(4-((3-methylazetidin-3-yl)methoxy)-1H-pyrazolo[3,4-d]pyrimidin-6-yl)phenol TFA salt (53.2 mg, 0.12 mmol) in DMF (0.60 ml) at 0 °C, followed by acryloyl chloride (9.75 µl, 0.120 mmol) and the reaction mixture was stirred at rt for 15 min. The reaction mixture was concentrated *in vacuo* to yield the crude liquid. The crude liquid was purified via preparatory scale C18 column chromatography using a gradient of 10-60% acetonitrile in water with 0.05% trifluoroacetic acid to yield 1-(3-(((6-(3-fluoro-4-hydroxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl)oxy)methyl)-3-methylazetidin-1-yl)prop-2-en-1-one as a TFA salt (10.5 mg, 0.120 mmol, 21.7 % yield). (*m/z*): [*M*+*H*]⁺ calcd for C₁₉H₁₉FN₅O₃ 384.15 found 384.3.

Preparation 61: tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)-3-ethylazetidine-1-carboxylate



5 60% By weight sodium hydride in mineral oil (353 mg, 8.82 mmol) was added to a solution of tert-butyl 3-ethyl-3-(hydroxymethyl)azetidine-1-carboxylate (1.1 g, 4.85 mmol) in DMF (10 mL) at 0 °C and the reaction mixture was stirred at rt for 40 minutes. A solution of 4,6-dichloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridine (1.20 g, 4.41 mmol) in DMF (5 mL) at 0 °C and the reaction mixture was stirred at rt for 10 1 hour. The reaction was quenched with addition of water and the mixture was extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with water and a saturated aqueous solution of sodium chloride and then dried over sodium sulfate to yield the crude residue. The crude residue was purified via flash column chromatography using 5% ethyl acetate in hexanes to yield tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)-3-ethylazetidine-1-carboxylate (900 mg, 15 4.39 mmol, 45.2 % yield). (m/z): [M+H]⁺ calcd for C₂₂H₃₂ClN₄O₄ 451.21 found 451.13.

Preparation 62: tert-butyl 3-ethyl-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate




20 A solution of phenolboronic acid (413 mg, 2.99 mmol) and sodium carbonate (421 mg, 3.98 mmol) in water (2 mL) was added to solution of tert-butyl 3-(((6-chloro-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)-3-

ethylazetidine-1-carboxylate (900 mg, 1.99 mmol) in dioxane (8 mL) and the reaction was degassed with argon for 5 minutes.


Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloromethane adduct (162 mg, 0.19 mmol) was added, the reaction mixture was degassed further with argon for 5 minutes and the reaction mixture was stirred at 110 °C for 1 hour. The reaction mixture was diluted with ethyl acetate and filtered through Celite. Water was added, the ethyl acetate layer was separated and the aqueous layer was extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with a saturated aqueous solution of sodium chloride, dried over sodium sulfate and concentrated *in vacuo* to yield the crude liquid. The crude liquid was purified via flash column chromatography using 30% ethyl acetate in hexanes to yield tert-butyl 3-ethyl-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate (800 mg, 1.57 mmol, 79.0 % yield). (m/z): [M+H]⁺ calcd for C₂₈H₃₇N₄O₅ 509.28 found 509.29.

Preparation 63: 4-(4-((3-ethylazetidin-3-yl)methoxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol




Tert-butyl 3-ethyl-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidine-1-carboxylate (800 mg, 1.57 mmol) was dissolved in dichloromethane (10 ml) and TFA (10 ml) was added at 0 °C and the reaction mixture was stirred at rt for 3 hours. The reaction was concentrated *in vacuo* and the resulting residue was triturated with diethyl ether. The crude residue was dissolved in a minimum of acetonitrile and precipitated with diethyl ether to yield 4-(4-((3-ethylazetidin-3-yl)methoxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol as a TFA salt (287 mg, 0.885 mmol, 56.3 % yield). (m/z): [M+H]⁺ calcd for C₁₈H₂₁N₄O₂ 325.17 found 325.07.

Example 22: 1-(3-ethyl-3-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidin-1-yl)prop-2-en-1-one




Diisopropylethylamine (0.246 ml, 1.41 mmol) was added to a solution of 4-(4-((3-ethylazetidin-3-yl)methoxy)-1H-pyrazolo[3,4-b]pyridin-6-yl)phenol TFA salt (124 mg, 0.283 mmol) in DMF (2.0 ml) at 0 °C, followed by acryloyl chloride (25 μ l, 0.311 mmol) and the reaction mixture was stirred at rt for 15 min. The reaction mixture was concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via preparatory scale C18 column chromatography using a gradient of 20-80% acetonitrile in water with 0.05% trifluoroacetic acid to yield 1-(3-ethyl-3-(((6-(4-hydroxyphenyl)-1H-pyrazolo[3,4-b]pyridin-4-yl)oxy)methyl)azetidin-1-yl)prop-2-en-1-one as a TFA salt (44.8 mg, 0.118 mmol, 41.9 % yield). (*m/z*): [M+H]⁺ calcd for C₂₁H₂₃N₄O₃ 379.18 found 379.3.

Preparation 64: 6-bromo-4-fluoro-1H-benzo[d][1,2,3]triazole




A solution of sodium nitrite (1.85 g, 26.8 mmol) in water (50 mL) was added to a solution of 5-bromo-3-fluorobenzene-1,2-diamine (5.00 g, 24.4 mmol) in water (50 mL) and acetic acid (18.0 mL) and the reaction mixture was stirred at rt for 15 minutes and then at 85 °C for 1 hour. The reaction mixture was concentrated *in vacuo* a water was added to the resulting residue. The mixture was extracted with ethyl acetate (2 x 300 mL), the ethyl acetate extracts were combined, dried over sodium sulfate and concentrated *in vacuo* to yield 6-bromo-4-fluoro-1H-benzo[d][1,2,3]triazole (5.0 g, 23.1 mmol, 94.0 % yield). (*m/z*): [M-H]⁻ calcd for C₆H₃BrFN₃ 213.94 found 213.94.

Preparation 65: 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazole




5 p-Toluenesulfonic acid (796 mg, 4.62 mmol) was added to a solution of 6-bromo-4-fluoro-1H-benzo[d][1,2,3]triazole (5.0 g, 23.1 mmol) in tetrahydrofuran (200 mL), followed by 3,4-dihydro-2H-pyran (7.7 g, 92.6 mmol) and the reaction mixture was stirred at rt for 4 hours. The reaction mixture was concentrated *in vacuo* and the resulting residue was purified via flash column chromatography using 10% ethyl acetate in hexanes
10 to yield 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazole (3.8 g, 12.7 mmol, 55.4 % yield). (m/z): [M+H]⁺ calcd for C₁₁H₁₂BrFN₃O 300.01 found 300.09.

Preparation 66: tert-butyl (3S)-3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate



Cesium carbonate (1.50 g, 4.64 mmol) was added to a solution of tert-butyl (S)-3-(hydroxymethyl)pyrrolidine-1-carboxylate (564 mg, 2.80 mmol) in DMSO (15 ml) at 0
20 °C, followed by 6-bromo-4-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazole (700 mg, 2.34 mmol) and the reaction mixture was stirred at 80 °C for 3 hours. Water was added and the mixture was extracted with ethyl acetate (3 x). The ethyl acetate extracts were combined, washed with water and an aqueous solution of sodium chloride, dried over sodium sulfate and concentrated *in vacuo* to yield the crude
25 residue. The crude residue was purified via flash column chromatography using 30% ethyl acetate in hexanes to yield tert-butyl (3S)-3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (500 mg, 1.04 mmol, 45.6 % yield). (m/z): [M+H]⁺ calcd for C₂₁H₃₀BrN₄O₄ 481.15 found 481.2.


Preparation 67: tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate



5 Phenolboronic acid (192 mg, 1.40 mmol) was added to solution of tert-butyl (3S)-3-(((6-bromo-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (450 mg, 0.93 mmol) in dioxane (10 mL) and water (2 mL), followed by potassium phosphate (394 mg, 1.86 mmol) and the reaction was degassed with argon for 15 minutes. 2'-(Dimethylamino)-2-biphenyl-palladium(II) chloride dinorbornylphosphine complex (52 mg, 0.093 mmol) was added and the reaction mixture was stirred at 110 °C for 40 minutes under microwave irradiation. The reaction mixture was filtered through Celite. Water was added and the mixture was extracted with ethyl acetate (3 x). The ethyl acetate extracts were combined, washed with a saturated aqueous solution of sodium chloride, dried over sodium sulfate and concentrated *in vacuo* to yield the crude residue. The crude residue was purified via flash column chromatography using 30% ethyl acetate in hexanes to yield tert-butyl (3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (300 mg, 0.607 mmol, 65.0 % yield). (m/z): [M+H]⁺ calcd for C₂₇H₃₅N₄O₅ 495.26 found 495.06.

20

Preparation 68: (S)-4-(4-(pyrrolidin-3-ylmethoxy)-1H-benzo[d][1,2,3]triazol-6-yl)phenol




25 4M hydrochloric acid in dioxane (10.0 mL, 40 mmol) was added to a solution of ((3S)-3-(((6-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidine-1-carboxylate (230 mg, 0.465 mmol) in methanol (2. mL) and the reaction mixture was stirred at rt for 5 hours. The reaction was concentrated *in*

vacuo and the resulting residue was triturated with diethyl ether to yield (S)-4-(4-(pyrrolidin-3-ylmethoxy)-1H-benzo[d][1,2,3]triazol-6-yl)phenol as a hydrochloride salt (225 mg, 0.649 mmol, 85.0 % yield). (*m/z*): [M+H]⁺ calcd for C₁₇H₁₉N₄O₂ 311.15 found 311.06.

5


Example 23: (S)-1-(3-(((6-(4-hydroxyphenyl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidin-1-yl)prop-2-en-1-one



Diisopropylethylamine (0.083 ml, 0.475 mmol) was added to a solution of (S)-4-(4-(pyrrolidin-3-ylmethoxy)-1H-benzo[d][1,2,3]triazol-6-yl)phenol hydrochloride salt (32.9 mg, 0.095 mmol) in DMF (0.475 ml) at 0 °C, followed by acryloyl chloride (7.7 µl, 0.095 mmol) and the reaction mixture was stirred at rt for 15 min. The reaction mixture was concentrated *in vacuo* to yield a brown liquid. The crude liquid was purified via preparatory scale C18 column chromatography using a gradient of 5-75% acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-1-(3-(((6-(4-hydroxyphenyl)-1H-benzo[d][1,2,3]triazol-4-yl)oxy)methyl)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (2.5 mg, 0.068 mmol, 7.2 % yield). (*m/z*): [M+H]⁺ calcd for C₂₀H₂₁N₄O₃ 365.16 found 365.1.

20


Preparation 69: 6-bromo-4-fluoro-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole



p-Toluenesulfonic acid (236 mg, 1.24 mmol) was added to a solution of 6-bromo-4-fluoro-3-methyl-1H-indazole (2.84 g, 12.4 mmol) in dichloromethane (41 mL), followed by 3,4-dihydro-2H-pyran (3.13 g, 37.2 mmol) and the reaction mixture was stirred at rt for 3 days. The reaction mixture filtered through a pad of silica gel and the

filtrated was concentrated *in vacuo* to yield 6-bromo-4-fluoro-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (3.3 g, 10.5 mmol, 85 % yield). (m/z): [M+Na]⁺ calcd for C₁₃H₁₄BrFN₂NaO 335.02 found 335.4.

5 **Example 24: (S)-1-(3-((6-(3-chloro-4-hydroxyphenyl)-3-methyl-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one**



60% By weight sodium hydride in mineral oil (319 mg, 7.98 mmol) was added to a solution of (S)-1-boc-3-hydroxypyrrolidine (1.3 g, 6.94 mmol) in DMF (16 mL) at 0 °C and the reaction mixture was stirred at rt for 20 minutes. 6-Bromo-4-fluoro-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazole (1.00 g, 3.19 mmol) was added and the reaction mixture was stirred at rt for overnight. The reaction was quenched with addition of water and the mixture was concentrated *in vacuo* to yield the crude intermediate tert-butyl (3S)-3-((6-bromo-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate.


3-chloro-4-hydroxyphenylboronic acid (0.484 g, 2.81 mmol) and potassium phosphate tribasic (1.19 g, 5.62 mmol) were added to a solution of the crude tert-butyl (3S)-3-((6-bromo-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate in 2:1 dioxane/water (9.5 mL), and the reaction mixture was degassed with nitrogen for 10 minutes. (2-Dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate, XPhos-G3-Palladacycle (95 mg, 0.112 mmol) was added and the reaction mixture was stirred at 110 °C for 3 hours under microwave irradiation. Water was added and the mixture was extracted with ethyl acetate. The ethyl acetate extracts were combined and concentrated *in vacuo* to yield the crude intermediate tert-butyl (3S)-3-((6-(3-chloro-4-hydroxyphenyl)-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate.

4.0N HCl in dioxane (3.75 ml, 15.0 mmol) was added to a solution of the crude tert-butyl (3S)-3-((6-(3-chloro-4-hydroxyphenyl)-3-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-4-yl)oxy)pyrrolidine-1-carboxylate in methanol (0.5 mL), and the reaction mixture was stirred at 60 °C for 30 minutes. The reaction mixture was concentrated *in vacuo* to yield the crude intermediate (S)-2-chloro-4-(3-methyl-4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol.




















N,N-Diisopropylethylamine (4.59 ml, 26.3 mmol) was added to a solution of the crude intermediate (S)-2-chloro-4-(3-methyl-4-(pyrrolidin-3-yloxy)-1H-indazol-6-yl)phenol in methylene chloride (3.75 ml), followed by acryloyl chloride (60.9 μ L, 0.750 mmol). The reaction mixture was stirred at rt for 30 minutes then concentrated *in vacuo* to yield the crude residue. The crude residue was purified via preparatory scale C18 column chromatography using a gradient of 0-40% acetonitrile in water with 0.05% trifluoroacetic acid to yield (S)-1-(3-((6-(3-chloro-4-hydroxyphenyl)-3-methyl-1H-indazol-4-yl)oxy)pyrrolidin-1-yl)prop-2-en-1-one as a TFA salt (7.6 mg, 0.015 mmol, 0.98 % yield). (*m/z*): [M+H]⁺ calcd for C₂₁H₂₁ClN₃O₃ 398.13 found 398.3.















The compounds of Tables 1-12 were prepared using similar synthetic methods.

Table 1



Ex No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	X ¹	X ²	R ^{k1}	R ^{k2}	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
1-1		H	H	H	H	H	CH	CH	H	H	C ₂₃ H ₂₆ N ₄ O ₃	407.20	407
1-2		H	H	H	H	H	CH	CH	H	H	C ₂₀ H ₁₉ N ₃ O ₃	350.13	350
1-3		Cl	H	H	H	H	CH	CH	H	H	C ₂₃ H ₂₅ ClN ₄ O ₃	441.15	441.1

Ex No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	X ¹	X ²	R ^{k1}	R ^{k2}	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
1-4		F	H	H	H	H	CH	CH	H	H	C ₂₃ H ₂₅ FN ₄ O ₃	425.18	425.1
1-5		F	H	H	H	H	CH	CH	H	H	C ₂₀ H ₁₈ FN ₃ O ₃	368.12	368
1-6		H	H	H	H	H	N	N	H	H	C ₁₈ H ₁₇ N ₅ O ₃	352.12	352.3
1-7		H	H	H	H	H	CH	N	H	H	C ₂₂ H ₂₅ N ₅ O ₃	408.18	408.2
1-8		H	H	H	H	H	CH	N	H	H	C ₁₉ H ₁₈ N ₄ O ₃	351.12	351.2
1-9		H	H	H	H	H	N	CH	H	H	C ₁₉ H ₁₈ N ₄ O ₃	351.12	351.1
1-10		H	H	H	H	H	CH	CH	Me	H	C ₂₄ H ₂₈ N ₄ O ₃	421.20	421.3
1-11		H	H	H	H	H	CH	CH	Me	H	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.2
1-12		H	H	H	H	H	CH	CH	H	Me	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364
1-13		F	F	H	H	H	CH	CH	H	H	C ₂₀ H ₁₇ F ₂ N ₃ O ₄	386.11	386
1-14		F	H	F	H	H	CH	CH	H	H	C ₂₀ H ₁₇ F ₂ N ₃ O ₃	386.11	386
1-15		F	Cl	H	H	H	CH	CH	H	H	C ₂₀ H ₁₇ ClFN ₇ O ₃	402.08	402
1-16		F	H	H	Me	H	CH	CH	H	H	C ₂₁ H ₂₀ FN ₃ O ₃	382.13	382.1
1-17		H	F	H	H	H	CH	CH	H	H	C ₂₀ H ₁₈ FN ₃ O ₃	368.12	368.0
1-18		H	F	H	F	H	CH	CH	H	H	C ₂₀ H ₁₇ F ₂ N ₃ O ₃	386.11	386.0
1-19		F	H	H	F	H	CH	CH	H	H	C ₂₀ H ₁₇ F ₂ N ₃ O ₃	386.11	386.0
1-20		H	H	H	H	Me	CH	CH	H	H	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.0
1-21		H	H	H	H	NHMe	CH	CH	H	H	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.0
1-22		H	H	H	H	CH ₂ -OMe	CH	CH	H	H	C ₂₂ H ₂₃ N ₃ O ₄	394.15	394.0

Ex No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	X ¹	X ²	R ^{k1}	R ^{k2}	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
1-23		F	H	H	H	Me	CH	CH	H	H	C ₂₁ H ₂₀ FN ₃ O ₃	382.13	382.0
1-24		H	H	H	H	H	N	N	Me	H	C ₁₉ H ₁₉ N ₅ O ₃	366.13	366.0
1-25		H	H	H	H	CHF ₂	N	N	H	H	C ₁₉ H ₁₇ F ₂ N ₅ O ₃	402.11	402.0
1-26		H	H	H	H	H	N	CH	Me	H	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0
1-27		F	H	H	H	H	CH	N	H	H	C ₁₉ H ₁₇ FN ₄ O ₃	369.11	369.2
1-28		F	H	H	H	H	CH	N	H	H	C ₂₂ H ₂₄ FN ₅ O ₃	426.17	426.1
1-29		F	H	H	Me	H	CH	CH	H	H	C ₂₄ H ₂₇ FN ₄ O ₃	439.19	439.1
1-30		F	H	H	F	Me	CH	CH	H	H	C ₂₁ H ₁₉ F ₂ N ₃ O ₃	400.12	400.4
1-31		H	H	H	H	Me	N	CH	H	H	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0
1-32		Cl	H	H	H	Me	N	CH	H	H	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.0
1-33		F	H	H	H	Me	N	CH	H	H	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.0
1-34		H	H	H	H	Me	CH	N	H	H	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0
1-35		Cl	H	H	H	Me	CH	N	H	H	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.0
1-36		F	H	H	H	Me	CH	N	H	H	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.0














Ex No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	X ¹	X ²	R ^{k1}	R ^{k2}	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
1-37		Cl	H	H	H	Me	N	N	H	H	C ₁₉ H ₁₈ ClN ₅ O ₃	400.09	400.0
1-38		F	H	H	H	Me	N	N	H	H	C ₁₉ H ₁₈ FN ₅ O ₃	384.12	384.0
1-39		H	H	H	H	SMe	CH	CH	H	H	C ₂₁ H ₂₁ N ₃ O ₃ S	396.11	396.4
1-40		F	H	H	H	SMe	CH	CH	H	H	C ₂₁ H ₂₀ FN ₃ O ₃ S	414.11	414.0
1-41		Cl	H	H	H	SMe	CH	CH	H	H	C ₂₁ H ₂₀ ClN ₃ O ₃ S	430.08	430.0
1-42		H	H	H	H	NMe ₂	CH	CH	H	H	C ₂₂ H ₂₄ N ₄ O ₃	393.17	393.0
1-43		H	H	H	H	CF ₃	CH	CH	H	H	C ₂₁ H ₁₈ F ₃ N ₃ O ₃	418.11	418.4
1-44		H	H	H	H	OMe	CH	CH	H	H	C ₂₁ H ₂₁ N ₃ O ₄	380.14	380.0
1-45		H	H	H	H	CN	CH	CH	H	H	C ₂₁ H ₁₈ N ₄ O ₃	375.12	375.0
1-46		Cl	H	H	H	OMe	CH	CH	H	H	C ₂₁ H ₁₇ ClN ₄ O ₃	409.08	409.0
1-47		H	H	H	H	Cl	CH	CH	H	H	C ₂₀ H ₁₈ ClN ₃ O ₃	384.09	384.1
1-48		H	H	H	H	Cl	CH	CH	H	H	C ₂₃ H ₂₅ ClN ₄ O ₃	441.15	441.8

Table 2



Ex No.	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
2-1		H	H	CH	CH	C ₂₄ H ₂₈ N ₄ O ₃	421.20	421.2
2-2		Cl	H	CH	CH	C ₂₄ H ₂₇ N ₄ O ₃ Cl	455.16	455.1
2-3		Cl	H	CH	CH	C ₂₁ H ₂₀ ClN ₃ O ₃	398.10	398.2
2-4		F	H	CH	CH	C ₂₄ H ₂₇ FN ₄ O ₃	439.19	439.3
2-5		F	H	CH	CH	C ₂₁ H ₂₀ FN ₃ O ₃	382.13	382.1
2-6		H	H	N	CH	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0
2-7		H	H	N	N	C ₁₉ H ₁₉ N ₅ O ₃	366.13	366.0
2-8		H	H	CH	N	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0
2-9		Cl	H	CH	N	C ₂₃ H ₂₆ ClN ₅ O ₃	456.16	456.2
2-10		F	H	CH	N	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.0
2-11		Cl	H	N	N	C ₁₉ H ₁₈ ClN ₅ O ₃	400.09	400.1
2-12		H	H	N	CH	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0
2-13		F	H	CH	N	C ₂₃ H ₂₆ FN ₅ O ₃	440.19	440.3
























































2-14		Cl	H	CH	N	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.0
2-15		F	H	N	N	C ₁₉ H ₁₈ FN ₅ O ₃	384.12	384.1
2-16		Cl	H	N	CH	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.1
2-17		F	H	N	CH	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.1





















Table 3











































Ex No.	R ¹	R ²	R ³	X ¹	X ²	R ^{k1}	R ^{k2}	R ^d	R ^e	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
3-1		H	H	CH	CH	H	H	H	H	C ₂₀ H ₁₉ N ₃ O ₃	350.13	350.2
3-2		Cl	H	CH	CH	H	H	H	H	C ₂₃ H ₂₅ ClN ₄ O ₃	441.15	441.3
3-3		Cl	H	CH	CH	H	H	H	H	C ₂₀ H ₁₈ ClN ₃ O ₃	384.09	384.1
3-4		F	H	CH	CH	H	H	H	H	C ₂₀ H ₁₈ FN ₃ O ₃	368.12	368.2
3-5		H	H	CH	CH	F	H	H	H	C ₂₃ H ₂₅ FN ₄ O ₃	425.18	425.1
3-6		H	H	CH	CH	F	H	H	H	C ₂₀ H ₁₈ FN ₃ O ₃	368.12	368.2
3-7		H	H	CH	CH	Me	H	H	H	C ₂₄ H ₂₈ N ₄ O ₃	421.20	421.1
3-8		H	H	CH	CH	Me	H	H	H	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.1
3-9		Cl	H	CH	CH	Me	H	H	H	C ₂₄ H ₂₇ ClN ₄ O ₃	455.16	456.1
3-10		Cl	H	CH	CH	Me	H	H	H	C ₂₁ H ₂₀ ClN ₃ O ₃	398.10	398.1
















3-11		F	H	CH	CH	Me	H	H	H	$C_{24}H_{27}FN_4O_3$	439.19	439.2
3-12		F	H	CH	CH	Me	H	H	H	$C_{21}H_{20}FN_3O_3$	382.13	382.1
3-13		H	H	CH	CH	Et	H	H	H	$C_{25}H_{30}N_4O_3$	435.22	435.1
3-14		H	H	CH	CH	Et	H	H	H	$C_{22}H_{23}N_3O_3$	378.16	378.1
3-15		Cl	H	CH	CH	Et	H	H	H	$C_{25}H_{29}ClN_4O_3$	469.18	469.1
3-16		Cl	H	CH	CH	Et	H	H	H	$C_{22}H_{22}ClN_3O_3$	412.12	412.1
3-17		F	H	CH	CH	Et	H	H	H	$C_{25}H_{29}FN_4O_3$	453.21	453.3
3-18		F	H	CH	CH	Et	H	H	H	$C_{22}H_{22}FN_3O_3$	396.15	396.1
3-19		H	H	CH	CH	CN	H	H	H	$C_{24}H_{25}N_5O_3$	432.18	432.1
3-20		H	H	CH	CH	CN	H	H	H	$C_{21}H_{18}N_4O_3$	375.12	375.1
3-21		Cl	H	CH	CH	CN	H	H	H	$C_{24}H_{24}ClN_5O_3$	466.14	466.2
3-22		Cl	H	CH	CH	CN	H	H	H	$C_{21}H_{17}ClN_4O_3$	409.08	409.1
3-23		F	H	CH	CH	CN	H	H	H	$C_{21}H_{17}FN_4O_3$	393.11	393.1
3-24		H	H	CH	CH	H	H	Me	Me	$C_{25}H_{30}N_4O_3$	435.22	435.0
3-25		H	H	CH	CH	H	H	Me	Me	$C_{22}H_{23}N_3O_3$	378.16	378.1
3-26		Cl	H	CH	CH	H	H	Me	Me	$C_{22}H_{22}ClN_3O_3$	412.12	412.2
3-27		F	H	CH	CH	H	H	Me	Me	$C_{25}H_{29}FN_4O_3$	453.21	453.2
3-28		F	H	CH	CH	H	H	Me	Me	$C_{22}H_{22}FN_3O_3$	393.15	396.3
3-29		H	H	CH	N	H	H	H	H	$C_{19}H_{18}N_4O_3$	351.12	351.0
3-30		H	H	CH	CH	H	Me	H	H	$C_{24}H_{28}N_4O_3$	421.20	421.2

3-31		H	H	N	N	H	Me	H	H	$C_{19}H_{19}N_5O_3$	366.13	366.1
3-32		H	H	CH	CH	OMe	H	H	H	$C_{24}H_{28}N_4O_4$	437.20	437.3
3-33		H	H	CH	CH	OMe	H	H	H	$C_{21}H_{21}N_3O_4$	380.14	380.2
3-34		Cl	H	CH	N	H	H	H	H	$C_{22}H_{24}ClN_5O_3$	442.14	442.1
3-35		Cl	H	CH	N	H	H	H	H	$C_{19}H_{17}ClN_4O_3$	385.08	385.0
3-36		F	H	CH	N	H	H	H	H	$C_{22}H_{24}FN_5O_3$	426.17	426.0
3-37		F	H	CH	N	H	H	H	H	$C_{19}H_{17}FN_4O_3$	369.11	369.1
3-38		Cl	H	N	N	H	H	H	H	$C_{18}H_{16}ClN_5O_3$	386.08	386.2
3-39		F	H	N	N	H	H	H	H	$C_{18}H_{16}FN_5O_3$	370.11	370.1
3-40		F	H	CH	CH	F	H	H	H	$C_{23}H_{24}F_2N_4O_3$	443.17	443.1
3-41		F	H	CH	CH	F	H	H	H	$C_{20}H_{17}F_2N_3O_3$	386.11	386.2
3-42		Cl	H	CH	CH	F	H	H	H	$C_{23}H_{24}ClFN_4O_3$	459.14	459.1
3-43		Cl	H	CH	CH	F	H	H	H	$C_{20}H_{17}ClFN_3O_3$	402.08	402.0
3-44		H	H	CH	N	F	H	H	H	$C_{22}H_{24}FN_5O_3$	426.17	426.2
3-45		H	H	N	N	F	H	H	H	$C_{18}H_{16}FN_5O_3$	370.11	370.1
3-46		H	H	N	CH	F	H	H	H	$C_{19}H_{17}FN_4O_3$	369.11	369.3
3-47		Cl	H	CH	N	F	H	H	H	$C_{19}H_{16}ClFN_4O_3$	403.07	403.0
3-48		F	H	CH	N	F	H	H	H	$C_{22}H_{23}F_2N_5O_3$	444.16	444.0
3-49		F	H	CH	N	F	H	H	H	$C_{19}H_{16}F_2N_4O_3$	387.10	387.0
3-50		Cl	H	N	N	F	H	H	H	$C_{21}H_{22}ClFN_6O_3$	461.13	461.1

3-51		F	H	N	N	F	H	H	H	C ₂₁ H ₂₂ F ₂ N ₆ O ₃	445.16	445.2
3-52		Cl	H	N	CH	F	H	H	H	C ₂₂ H ₂₃ ClFN ₅ O ₃	460.13	460.1
3-53		F	H	N	CH	F	H	H	H	C ₂₂ H ₂₃ F ₂ N ₅ O ₃	444.16	444.2
3-54		F	H	N	CH	F	H	H	H	C ₁₉ H ₁₆ F ₂ N ₄ O ₃	387.10	387.1
3-55		H	H	CH	N	Me	H	H	H	C ₂₃ H ₂₇ N ₅ O ₃	422.20	422.2
3-56		H	H	CH	N	Me	H	H	H	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.2
3-57		H	H	N	N	Me	H	H	H	C ₁₉ H ₁₉ N ₅ O ₃	366.13	366.1
3-58		H	H	N	CH	Me	H	H	H	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.2
3-59		F	H	CH	N	Me	H	H	H	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.3
3-60		Cl	H	N	N	H	H	H	H	C ₂₁ H ₂₃ ClN ₆ O ₃	443.14	443.1
3-61		H	H	CH	N	Et	H	H	H	C ₂₄ H ₂₉ N ₅ O ₃	436.21	436.1
3-62		H	H	CH	N	Et	H	H	H	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.0
3-63		H	H	N	N	Et	H	H	H	C ₂₀ H ₂₁ N ₅ O ₃	380.15	380.2
3-64		H	H	CH	N	Et	H	H	H	C ₂₄ H ₂₉ N ₅ O ₃	436.21	436.1
3-65		H	H	CH	N	Et	H	H	H	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.0
3-66		Cl	H	CH	N	Et	H	H	H	C ₂₁ H ₂₁ ClN ₄ O ₃	413.11	413.1
3-67		F	H	N	N	Et	H	H	H	C ₂₀ H ₂₀ FN ₅ O ₃	398.14	398.1
3-68		Cl	H	N	CH	Et	H	H	H	C ₂₄ H ₂₈ ClN ₅ O ₃	470.17	470.2
3-69		Cl	H	N	CH	Et	H	H	H	C ₂₁ H ₂₁ ClN ₄ O ₃	413.11	413.2
3-70		F	H	N	CH	Et	H	H	H	C ₂₁ H ₂₁ FN ₄ O ₃	397.14	397.2

3-71		Cl	H	CH	N	CN	H	H	H	$C_{23}H_{23}ClN_6O_3$	467.14	467.0
3-72		Cl	H	CH	CH	OMe	H	H	H	$C_{24}H_{27}ClN_4O_4$	471.16	471.2
3-73		Cl	H	CH	CH	OMe	H	H	H	$C_{21}H_{20}ClN_3O_4$	414.10	414.2
3-74		F	H	CH	CH	OMe	H	H	H	$C_{24}H_{27}FN_4O_4$	455.19	455.1
3-75		F	H	CH	CH	OMe	H	H	H	$C_{21}H_{20}FN_3O_4$	398.13	398.2
3-76		H	H	N	N	OMe	H	H	H	$C_{19}H_{19}N_5O_4$	382.13	382.1
3-77		H	H	N	N	H	H	H	Me	$C_{19}H_{19}N_5O_3$	366.13	366.1
3-78		H	H	N	N	H	H	H	Me (S)	$C_{19}H_{19}N_5O_3$	366.13	366.1
3-79		H	H	N	CH	H	H	H	H	$C_{22}H_{25}N_5O_3$	408.18	408.1
3-80		Cl	H	N	CH	H	H	Me	Me	$C_{21}H_{21}ClN_4O_3$	413.11	413.2
3-81		F	H	N	CH	H	H	Me	Me	$C_{21}H_{21}FN_4O_3$	397.14	397.1
3-82		F	F	CH	CH	H	H	H	H	$C_{23}H_{24}F_2N_4O_3$	443.17	443.7
3-83		OMe	H	CH	CH	H	H	H	H	$C_{21}H_{21}N_3O_4$	380.14	380.0
3-84		H	H	CH	CH	H	H	H	H	$C_{20}H_{17}N_3O_3$	348.11	348.1
3-85		H	H	CH	CH	H	H	H	H	$C_{25}H_{28}N_4O_3$	433.20	433.1
3-86		Cl	H	N	CH	H	H	H	H	$C_{19}H_{17}ClN_4O_3$	385.08	385.1
3-87		F	H	N	CH	H	H	H	H	$C_{19}H_{17}FN_4O_3$	369.11	369.1
3-88		H	H	N	N	F	H	H	H	$C_{21}H_{23}FN_6O_3$	427.17	427.2
3-89		H	H	N	CH	F	H	H	H	$C_{22}H_{24}FN_5O_3$	426.17	426.2
3-90		H	H	CH	N	F	H	H	H	$C_{19}H_{17}FN_4O_3$	369.11	369.1

3-91		Cl	H	CH	N	F	H	H	H	$C_{22}H_{23}ClFN_5O_3$	460.13	460.0
3-92		Cl	H	N	N	F	H	H	H	$C_{18}H_{15}ClFN_5O_3$	404.07	404.1
3-93		F	H	N	N	F	H	H	H	$C_{18}H_{15}F_2N_5O_3$	388.10	388.1
3-94		Cl	H	N	CH	F	H	H	H	$C_{19}H_{16}ClFN_4O_3$	403.07	403.1
3-95		Cl	H	CH	N	Me	H	H	H	$C_{23}H_{26}ClN_5O_3$	456.16	456.2
3-96		F	H	CH	N	Me	H	H	H	$C_{23}H_{26}FN_5O_3$	440.19	440.2
3-97		Cl	H	CH	N	Me	H	H	H	$C_{20}H_{19}ClN_4O_3$	399.10	399.2
3-98		F	H	N	CH	Me	H	H	H	$C_{20}H_{19}FN_4O_3$	383.13	383.2
3-99		Cl	H	CH	N	Et	H	H	H	$C_{24}H_{28}ClN_5O_3$	470.17	470.1
3-100		F	H	CH	N	Et	H	H	H	$C_{24}H_{28}FN_5O_3$	454.20	454.2
3-101		F	H	CH	N	Et	H	H	H	$C_{21}H_{21}FN_4O_3$	397.14	397.2
3-102		H	H	CH	N	CN	H	H	H	$C_{23}H_{24}N_6O_3$	433.18	433.2
3-103		H	H	CH	N	CN	H	H	H	$C_{20}H_{17}N_5O_3$	376.12	376.2
3-104		H	H	N	N	CN	H	H	H	$C_{19}H_{16}N_6O_3$	377.11	377.0
3-105		H	H	CH	N	CN	H	H	H	$C_{20}H_{17}N_5O_3$	376.12	376.2
3-106		F	H	CH	N	CN	H	H	H	$C_{23}H_{23}FN_6O_3$	451.17	451.1
3-107		Cl	H	CH	N	CN	H	H	H	$C_{20}H_{16}ClN_5O_3$	410.08	410.1
3-108		F	H	CH	N	CN	H	H	H	$C_{20}H_{16}FN_5O_3$	394.11	394.2
3-109		Cl	H	N	N	CN	H	H	H	$C_{19}H_{15}ClN_6O_3$	411.07	411.1
3-110		F	H	N	N	CN	H	H	H	$C_{19}H_{15}FN_6O_3$	395.10	395.1

3-111		F	H	N	CH	CN	H	H	H	$C_{20}H_{16}FN_5O_3$	394.11	394.1
3-112		H	H	N	N	H	H	H	H	$C_{21}H_{24}N_6O_3$	409.18	409.1
3-113		H	H	CH	CH	H	H	H	H	$C_{25}H_{30}N_4O_3$	435.22	435.1
3-114		H	H	CH	CH	H	Me	H	H	$C_{21}H_{21}N_3O_3$	364.14	364.1
3-115		H	H	CH	CH	H	H	H	H	$C_{25}H_{28}N_4O_3$	433.20	433.1
3-116		H	H	CH	CH	H	H	H	H	$C_{26}H_{32}N_4O_3$	449.23	449.2
3-117		H	H	CH	CH	H	H	H	H	$C_{26}H_{30}N_4O_3$	447.22	
3-118		H	H	CH	CH	H	H	H	H	$C_{26}H_{32}N_4O_3$	449.23	449.2
3-119		H	H	CH	CH	H	H	H	H	$C_{25}H_{30}N_4O_4$	451.21	451.1
3-120		H	H	CH	CH	H	H	H	H	$C_{27}H_{34}N_4O_5$	495.24	
3-121		H	H	CH	CH	H	H	H	H	$C_{26}H_{32}N_4O_5$	481.22	
3-122		Cl	H	CH	CH	H	H	H	H	$C_{23}H_{25}ClN_4O_3$	441.15	441.1
3-123		H	H	CH	CH	H	H	H	H	$C_{24}H_{26}N_4O_3$	419.18	419.1
3-124		H	H	CH	CH	H	H	H	H	$C_{25}H_{28}N_4O_3$	433.20	
3-125		H	H	CH	CH	H	H	H	H	$C_{26}H_{30}N_4O_3S$	479.19	479.2












3-126		H	H	CH	CH	H	H	H	H	C ₂₅ H ₃₀ N ₄ O ₃ S	467.18	
3-127		H	H	CH	CH	H	H	H	H	C ₂₇ H ₃₂ N ₄ O ₄	477.23	
3-128		H	H	CH	CH	H	H	H	H	C ₂₇ H ₃₂ N ₄ O ₄	477.23	477.2
3-129		H	H	CH	CH	H	H	H	H	C ₂₅ H ₂₈ N ₄ O ₄	449.20	449.1
3-130		H	H	CH	CH	H	H	H	H	C ₂₅ H ₂₈ N ₄ O ₃ S	465.17	465.1
3-131		H	H	CH	CH	H	H	H	H	C ₂₇ H ₃₂ N ₄ O ₃ S	493.20	493.2
3-132		H	H	CH	CH	H	H	H	H	C ₂₅ H ₂₈ N ₄ O ₄	449.20	
3-133		H	H	CH	CH	H	H	H	H	C ₂₆ H ₃₀ N ₄ O ₄	463.21	463.2
3-134		H	H	CH	CH	H	H	H	H	C ₂₇ H ₃₂ N ₄ O ₄	477.23	

Table 4



Ex No.	R ¹	R ²	R ³	R ^b	R ^k	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
4-1		H	H	Me	H	CH	CH	C ₂₁ H ₂₂ N ₄ O ₂	363.15	363.1



































4-2		H	H	H	Me	CH	CH	C ₂₁ H ₂₂ N ₄ O ₂	363.15	363
4-3		H	H	H	H	CH	CH	C ₂₀ H ₂₀ N ₄ O ₂	349.14	349.1
4-4		Cl	H	Me	H	CH	CH	C ₂₄ H ₂₈ ClN ₅ O ₂	454.18	454.1
4-5		Cl	H	Me	H	CH	CH	C ₂₁ H ₂₁ ClN ₄ O ₂	397.12	397.0
4-6		F	H	Me	H	CH	CH	C ₂₄ H ₂₈ FN ₅ O ₂	438.21	438.1
4-7		F	H	Me	H	CH	CH	C ₂₁ H ₂₁ FN ₄ O ₂	481.15	381.0
4-8		H	H	Me	H	CH	N	C ₂₃ H ₂₈ N ₆ O ₂	421.21	421.2

Table 5



Ex No.	R ¹	R ²	R ³	X ¹	X ²	R ^k	L	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
5-1		H	H	CH	CH	H	O	C ₂₂ H ₂₄ N ₄ O ₃	393.17	393.4
5-2		H	H	CH	CH	H	O	C ₁₉ H ₁₇ N ₃ O ₃	336.11	336.2
5-3		Cl	H	CH	CH	H	O	C ₂₂ H ₂₃ ClN ₄ O ₃	427.13	427.1
5-4		Cl	H	CH	CH	H	O	C ₁₉ H ₁₆ ClN ₃ O ₃	370.07	370.1
5-5		F	H	CH	CH	H	O	C ₂₂ H ₂₃ FN ₄ O ₃	411.16	411.3
5-6		F	H	CH	CH	H	O	C ₁₉ H ₁₆ FN ₃ O ₃	354.10	354.2

5-7		H	H	CH	CH	H	NH	C ₁₉ H ₁₈ N ₄ O ₂	335.12	335.2
5-8		H	H	CH	CH	H	NMe	C ₂₃ H ₂₇ N ₅ O ₂	406.20	406.1
5-9		H	H	CH	CH	H	NMe	C ₂₀ H ₂₀ N ₄ O ₂	349.14	349.1
5-10		H	H	CH	CH	CHF ₂	O	C ₂₃ H ₂₄ F ₂ N ₄ O ₃	443.17	443.1
5-11		H	H	CH	CH	Me	O	C ₂₃ H ₂₆ N ₄ O ₃	407.18	407.1
5-12		H	H	CH	CH	Me	O	C ₂₀ H ₁₉ N ₃ O ₃	350.13	350.1
5-13		H	H	CH	CH		O	C ₂₅ H ₂₈ N ₄ O ₃	433.20	433.2
5-14		H	H	CH	CH	CHF ₂	O	C ₂₀ H ₁₇ F ₂ N ₃ O ₃	386.11	386.0
5-15		H	H	N	CH	H	O	C ₁₈ H ₁₆ N ₄ O ₃	337.11	337.1
5-16		Cl	H	CH	CH	CHF ₂	O	C ₂₃ H ₂₃ ClF ₂ N ₄ O ₃	477.13	477.1
5-17		F	H	CH	CH	CHF ₂	O	C ₂₃ H ₂₃ F ₃ N ₄ O ₃	461.16	461.2
5-18		Cl	H	CH	CH	Me	O	C ₂₃ H ₂₅ ClN ₄ O ₃	441.15	441.2
5-19		F	H	CH	CH	Me	O	C ₂₃ H ₂₅ FN ₄ O ₃	425.18	425.3
5-20		H	H	N	N	H	O	C ₁₇ H ₁₅ N ₅ O ₃	338.10	338.2
5-21		Cl	H	CH	CH	CHF ₂	O	C ₂₀ H ₁₆ ClF ₂ N ₃ O ₃	420.07	420.0
5-22		F	H	CH	CH	CHF ₂	O	C ₂₀ H ₁₆ F ₃ N ₃ O ₃	404.10	404.1
5-23		Cl	H	CH	CH	Me	O	C ₂₀ H ₁₈ ClN ₃ O ₃	384.09	384.2
5-24		F	H	CH	CH	Me	O	C ₂₀ H ₁₈ FN ₃ O ₃	368.12	368.1
5-25		H	H	CH	CH	CF ₃	O	C ₂₃ H ₂₃ F ₃ N ₄ O ₃	461.16	461.2














5-26		H	H	CH	CH	CF ₃	O	C ₂₀ H ₁₆ F ₃ N ₃ O ₃	404.10	404.1
------	---	---	---	----	----	-----------------	---	--	--------	-------

Table 6



Ex No.	R ¹	R ²	R ³	X ¹	X ²	R ^{k1}	R ^{k2}	R ^{k3}	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
6-1		H	H	CH	CH	H	H	H	C ₂₄ H ₂₈ N ₄ O ₃	421.20	421.2
6-2		H	H	CH	CH	H	H	H	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.2
6-3		H	H	CH	CH	Me	H	H	C ₂₂ H ₂₃ N ₃ O ₃	378.16	378.1
6-4		Cl	H	CH	CH	Me	H	H	C ₂₅ H ₂₉ ClN ₄ O ₃	469.18	469.2
6-5		Cl	H	CH	CH	Me	H	H	C ₂₂ H ₂₂ ClN ₃ O ₃	412.12	412.2
6-6		F	H	CH	CH	Me	H	H	C ₂₅ H ₂₉ FN ₄ O ₃	453.21	453.2
6-7		F	H	CH	CH	Me	H	H	C ₂₂ H ₂₂ FN ₃ O ₃	396.15	396.1
6-8		H	H	CH	CH	H	F	F	C ₂₁ H ₁₉ F ₂ N ₃ O ₃	400.12	400.0
6-9		H	H	CH	CH	Me	H	H	C ₂₅ H ₃₀ N ₄ O ₃	435.22	435.2
6-10		H	H	CH	CH	Me	H	H	C ₂₂ H ₂₃ N ₃ O ₃	378.16	378.1
6-11		H	H	N	CH	H	H	H	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.0














6-12		H	H	N	N	H	H	H	C ₁₉ H ₁₉ N ₅ O ₃	366.13	366.1
6-13		Cl	H	CH	CH	H	H	H	C ₂₁ H ₂₀ ClN ₃ O ₃	398.10	398.1
6-14		F	H	CH	CH	H	H	H	C ₂₁ H ₂₀ FN ₃ O ₃	382.13	382.1
6-15		H	H	CH	N	H	H	H	C ₂₁ H ₂₀ N ₄ O ₃	365.14	365.1
6-16		F	H	CH	N	H	H	H	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.0
6-17		Cl	H	N	CH	H	H	H	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.3
6-18		F	H	N	CH	H	H	H	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.0
6-19		Cl	H	CH	CH	H	H	H	C ₂₄ H ₂₇ ClN ₄ O ₃	455.16	455.2
6-20		F	H	CH	CH	H	H	H	C ₂₄ H ₂₇ FN ₄ O ₃	439.19	439.1
6-21		Cl	H	CH	N	H	H	H	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.1
6-22		Cl	H	N	N	H	H	H	C ₁₉ H ₁₈ ClN ₅ O ₃	400.09	400.1
6-23		F	H	N	N	H	H	H	C ₁₉ H ₁₈ FN ₅ O ₃	384.12	384.0

Table 7



Ex No.	R ¹	R ²	R ³	X ¹	X ²	R ^{k1}	R ^{k2}	R ^{k3}	R ^b	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
7-1		H	H	CH	CH	Me	H	H	H	C ₂₅ H ₃₁ N ₅ O ₂	434.22	434.2
7-2		H	H	CH	CH	Me	H	H	H	C ₂₂ H ₂₄ N ₄ O ₂	377.17	377.3
7-3		H	H	CH	CH	H	H	H	Me	C ₂₂ H ₂₄ N ₄ O ₂	377.17	377
7-4		H	H	CH	CH	H	H	H	H	C ₂₁ H ₂₂ N ₄ O ₂	363.15	363
7-5		H	H	CH	CH	H	F	F	H	C ₂₄ H ₂₇ F ₂ N ₅ O ₂	456.19	456.2
7-6		H	H	CH	CH	H	F	F	H	C ₂₁ H ₂₀ F ₂ N ₄ O ₂	399.13	399.2
7-7		Cl	H	CH	CH	H	H	H	Me	C ₂₂ H ₂₃ ClN ₄ O ₂	411.14	411.2
7-8		Cl	H	CH	CH	Me	H	H	H	C ₂₅ H ₃₀ ClN ₅ O ₂	468.19	468.2
7-9		F	H	CH	CH	Me	H	H	H	C ₂₂ H ₂₃ FN ₄ O ₂	395.16	395.1
7-10		F	H	CH	CH	H	H	H	Me	C ₂₂ H ₂₃ FN ₄ O ₂	395.16	395.3
7-11		Cl	H	CH	CH	Me	H	H	H	C ₂₂ H ₂₃ ClN ₄ O ₂	411.13	411.2
7-12		F	H	CH	CH	Me	H	H	H	C ₂₅ H ₃₀ FN ₅ O ₂	452.21	452.2
7-13		H	H	CH	N	Me	H	H	H	C ₂₁ H ₂₃ N ₅ O ₂	378.16	378.2















7-14		Cl	H	CH	N	Me	H	H	H	C ₂₁ H ₂₂ ClN ₅ O ₂	412.12	412.2
------	---	----	---	----	---	----	---	---	---	---	--------	-------

Table 8



Ex No.	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
8-1		H	H	CH	CH	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364
8-2		Cl	H	CH	CH	C ₂₄ H ₂₇ ClN ₄ O ₃	455.16	455.1
8-3		Cl	H	CH	CH	C ₂₁ H ₂₀ ClN ₃ O ₃	398.10	398.0
8-4		F	H	CH	CH	C ₂₄ H ₂₇ FN ₄ O ₃	439.19	439.2
8-5		F	H	CH	CH	C ₂₁ H ₂₀ FN ₃ O ₃	382.13	382.0
8-6		H	H	N	CH	C ₂₀ H ₂₀ FN ₄ O ₃	365.14	365.1
8-7		H	H	CH	N	C ₂₀ H ₂₀ N ₄ O ₃	365.14	365.1
8-8		F	H	N	CH	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.0
8-9		F	H	CH	N	C ₂₃ H ₂₆ FN ₅ O ₃	440.19	440.2
8-10		Cl	H	CH	N	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.1
8-11		F	H	CH	N	C ₂₀ H ₁₉ FN ₄ O ₃	383.13	383.1
8-12		Cl	H	N	N	C ₁₉ H ₁₈ ClN ₅ O ₃	400.09	400.2




8-13		F	H	N	N	C ₁₉ H ₁₈ FN ₅ O ₃	384.12	384.0
8-14		Cl	H	N	CH	C ₂₀ H ₁₉ ClN ₄ O ₃	399.10	399.0

Table 9















Ex No.	R ¹	R ²	R ³	X ¹	X ²	L	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
9-1		H	H	CH	CH	O	C ₂₃ H ₂₃ N ₃ O ₃	390.16	390.1
9-2		Cl	H	CH	CH	O	C ₂₃ H ₂₂ ClN ₃ O ₃	424.12	424.0
9-3		Cl	H	CH	CH	O	C ₂₆ H ₂₉ ClN ₄ O ₃	481.18	481.1
9-4		F	H	CH	CH	O	C ₂₃ H ₂₂ FN ₃ O ₃	408.15	408.1
9-5		F	H	CH	CH	O	C ₂₆ H ₂₉ FN ₄ O ₃	465.21	465.1
9-6		H	H	CH	N	O	C ₂₂ H ₂₂ N ₄ O ₃	391.15	391.1
9-7		Cl	H	CH	N	O	C ₂₂ H ₂₁ ClN ₄ O ₃	425.11	425.0
9-8		Cl	H	CH	N	O	C ₂₅ H ₂₈ ClN ₅ O ₃	482.17	482.1
9-9		H	H	CH	CH	NH	C ₂₃ H ₂₄ N ₄ O ₂	389.17	389.2
9-10		F	H	CH	N	O	C ₂₂ H ₂₁ FN ₄ O ₃	409.14	409.0
9-11		H	H	N	N	O	C ₂₂ H ₂₁ N ₅ O ₃	392.15	392.2


Table 10



Ex No.	R ¹	R ²	R ³	X ¹	X ²	L	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
10-1		H	H	CH	CH	O	C ₂₆ H ₃₀ N ₄ O ₃	447.22	447.2
10-2		H	H	CH	CH	O	C ₂₃ H ₂₃ N ₃ O ₃	390.16	390.2
10-3		Cl	H	CH	CH	O	C ₂₆ H ₂₉ ClN ₄ O ₃	481.18	481.3
10-4		Cl	H	CH	CH	O	C ₂₃ H ₂₂ ClN ₃ O ₃	424.12	424.2
10-5		F	H	CH	CH	O	C ₂₆ H ₂₉ FN ₄ O ₃	465.21	465.2
10-6		F	H	CH	CH	O	C ₂₃ H ₂₂ FN ₃ O ₃	408.15	408.1
10-7		H	H	CH	CH	NH	C ₂₃ H ₂₄ N ₄ O ₂	389.17	389.3


5

Table 11

























































Ex No.	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
11-1		H	H	CH	CH	C ₂₁ H ₂₀ N ₄ O ₄	393.12	393.1
11-2		H	H	CH	CH	C ₂₄ H ₂₇ N ₅ O ₄	450.18	450.2























Table 12









































Ex No.	R	R ¹	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
12-1			C ₂₃ H ₂₆ N ₄ O ₂ S	423.16	423.1
12-2			C ₂₀ H ₁₉ N ₃ O ₂ S	366.10	366.2
12-3			C ₂₃ H ₂₆ N ₄ O ₂ S	423.16	423.2
12-4			C ₂₁ H ₂₁ N ₃ O ₂ S	380.12	380.3
12-5			C ₂₄ H ₂₈ N ₄ O ₂ S	437.18	437.2
12-6			C ₂₁ H ₂₁ N ₃ O ₂ S	380.12	380.2
12-7			C ₂₄ H ₂₈ N ₄ O ₂ S	437.18	437.2
12-8			C ₂₁ H ₂₁ N ₃ O ₄ S	412.11	412.1

12-9			C ₂₁ H ₂₁ N ₃ O ₄ S	412.11	412.1
12-10 (racemic)			C ₂₁ H ₂₁ N ₃ O ₂	348.15	348.2
12-11 (racemic)			C ₂₄ H ₂₈ N ₄ O ₂	405.21	405.2
12-12 (racemic)			C ₂₂ H ₂₃ N ₃ O ₂	362.16	362.3
12-13 (racemic)			C ₂₅ H ₃₀ N ₄ O ₂	419.22	419.2
12-14			C ₂₅ H ₃₀ N ₄ O ₂	149.22	419.2
12-15			C ₂₂ H ₂₄ N ₄ O ₂	377.17	377
12-16			C ₂₂ H ₂₄ N ₄ O ₂	377.17	377.2
12-17			C ₂₂ H ₂₄ N ₄ O ₂	377.17	377.3
12-18			C ₂₀ H ₂₀ N ₄ O ₂	349.14	349.2
12-19			C ₂₄ H ₂₈ N ₄ O ₃	421.20	421.3
12-20			C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.2
12-21			C ₂₂ H ₂₁ N ₃ O ₃	376.14	376.4
12-22			C ₂₃ H ₂₆ N ₄ O ₃	407.18	407.1
12-23			C ₂₀ H ₁₉ N ₃ O ₃	350.13	350.1

12-24			$C_{23}H_{26}N_4O_3$	407.18	407.2
12-25			$C_{20}H_{19}N_3O_3$	350.13	350.2
12-26			$C_{25}H_{30}N_4O_3$	435.22	435.1
12-27			$C_{22}H_{23}N_3O_3$	378.16	378.1
12-28			$C_{25}H_{30}N_4O_3$	435.22	435.2
12-29			$C_{22}H_{23}N_3O_3$	378.16	378.2
12-30			$C_{22}H_{23}N_3O_3$	378.16	378.2
12-31			$C_{22}H_{23}N_3O_3$	378.16	378.2
12-32			$C_{22}H_{23}N_3O_3$	378.16	378.1
12-33			$C_{22}H_{23}N_3O_3$	378.16	378.2
12-34			$C_{20}H_{19}N_3O_2S$	366.10	366.1
12-35			$C_{19}H_{17}N_3O_2S$	352.09	352.3

12-36			$C_{25}H_{27}FN_4O_3$	451.19	451.2
12-37			$C_{22}H_{20}FN_3O_3$	394.13	394.2
12-38			$C_{23}H_{26}N_4O_3$	407.18	407.3
12-39			$C_{20}H_{19}N_3O_3$	350.13	350.1
12-40			$C_{22}H_{22}N_4O_3$	391.15	391.1
12-41			$C_{23}H_{21}N_5O_3$	416.14	416.1
12-42			$C_{21}H_{20}N_4O_3S$	409.10	409.1
12-43			$C_{21}H_{20}N_4O_3S$	409.10	409.3
12-44			$C_{21}H_{20}N_4O_3S$	409.10	409.3
12-45			$C_{21}H_{20}N_4O_5S$	441.09	441.1
12-46			$C_{20}H_{18}N_4O_3$	363.11	363.1

12-47			$C_{21}H_{17}N_5O_3$	388.11	388.0
12-48			$C_{20}H_{17}FN_4O_3$	381.10	381.0
12-49			$C_{22}H_{24}N_4O_4S$	441.14	441.2
12-50			$C_{19}H_{17}N_3O_4S$	384.08	384.0
12-51			$C_{22}H_{24}N_4O_2S$	409.15	409.3
12-52			$C_{25}H_{28}N_4O_3$	433.20	433.2
12-53			$C_{25}H_{28}N_4O_3$	433.20	433.5
12-54			$C_{22}H_{21}N_3O_3$	376.14	376.5
12-55			$C_{22}H_{22}N_4O_3$	391.15	391.1
12-56			$C_{22}H_{22}N_4O_3$	391.15	391.0

12-57			$C_{23}H_{24}N_4O_3$	405.16	405.2
12-58			$C_{22}H_{21}FN_4O_3$	409.14	409.3
12-59			$C_{22}H_{22}N_4O_3$	391.15	391.1
12-60			$C_{22}H_{22}N_4O_4$	407.14	407.1
12-61			$C_{22}H_{22}N_4O_4$	407.14	407.1
12-62			$C_{23}H_{24}N_4O_4$	421.16	421.1
12-63			$C_{21}H_{20}N_4O_4$	393.12	393.1
12-64			$C_{21}H_{20}N_4O_3S$	409.10	409.1
12-65			$C_{24}H_{27}N_5O_3S$	466.16	466.1










































12-66			C ₂₁ H ₂₀ N ₄ O ₃ S	409.10	409.1
12-67			C ₂₃ H ₂₄ FN ₅ O ₃	438.16	438.1
12-68			C ₂₁ H ₂₀ N ₄ O ₃	377.13	377.1
12-69			C ₂₀ H ₁₈ N ₄ O ₃	363.11	363.0

Table 13



Ex No.	R	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
13-1			Cl	H	N	CH	C ₁₉ H ₁₇ ClN ₄ O ₂ S	401.06	401.1
13-2			Cl	H	CH	CH	C ₂₁ H ₁₉ ClN ₄ O ₄	427.09	427.0
13-3			F	H	CH	CH	C ₂₁ H ₁₉ FN ₄ O ₄	411.12	411.0
13-4			H	H	CH	CH	C ₂₃ H ₂₆ N ₄ O ₂ S	423.16	423.1
13-5			F	H	CH	CH	C ₂₃ H ₂₅ FN ₄ O ₂ S	441.15	441.2

13-6			Cl	H	CH	CH	C ₂₃ H ₂₅ ClN ₄ O ₂ S	457.12	457.1
13-7			F	H	CH	CH	C ₂₀ H ₁₈ FN ₃ O ₂ S	384.09	384.1
13-8			Cl	H	CH	CH	C ₂₀ H ₁₈ ClN ₃ O ₂ S	400.07	400.1
13-9			H	H	N	CH	C ₁₉ H ₁₈ N ₄ O ₂ S	367.10	367.0
13-10			H	H	CH	CH	C ₂₀ H ₂₁ N ₅ O ₃	380.15	380.2
13-11			H	H	N	CH	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.0
13-12			H	H	N	CH	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.0
13-13			H	H	N	CH	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.0
13-14	 (racemic)		H	Cl	CH	CH	C ₂₁ H ₂₀ ClN ₃ O ₃	398.10	398.3
13-15	 (racemic)		H	H	CH	CH	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.5
13-16			H	H	N	N	C ₂₀ H ₁₉ N ₅ O ₃	378.13	378.5
















13-17			Cl	H	N	N	C ₂₀ H ₁₈ ClN ₅ O ₃	412.09	412.4
13-18			F	H	N	N	C ₂₀ H ₁₈ FN ₅ O ₃	396.12	396.2
13-19			H	H	N	CH	C ₂₀ H ₁₉ N ₅ O ₄	394.12	394.1
13-20			Cl	H	CH	N	C ₂₀ H ₁₈ ClN ₅ O ₄	428.08	428.0
13-21			F	H	CH	N	C ₂₀ H ₁₈ FN ₅ O ₄	412.11	412.1
13-22			Cl	H	N	CH	C ₂₀ H ₁₈ ClN ₅ O ₄	428.08	428

Table 14



Ex No.	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
14-1		Cl	H	CH	CH	C ₂₂ H ₂₂ ClN ₃ O ₃	412.12	412.0
14-2		H	H	N	CH	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.1














14-3		Cl	H	CH	CH	C ₂₅ H ₂₉ ClN ₄ O ₃	469.18	469.1
14-4		F	H	CH	CH	C ₂₅ H ₂₉ FN ₄ O ₃	453.21	453.1
14-5		F	H	CH	CH	C ₂₂ H ₂₂ FN ₃ O ₃	396.15	396.1
14-6		H	H	CH	N	C ₂₁ H ₂₂ N ₄ O ₃	379.15	379.1
14-7		Cl	H	CH	N	C ₂₁ H ₂₁ ClN ₄ O ₃	413.11	413.0
14-8		F	H	CH	N	C ₂₁ H ₂₁ FN ₄ O ₃	397.14	397.1
14-9		F	H	N	CH	C ₂₁ H ₂₁ FN ₄ O ₃	397.14	397.1
14-10		Cl	H	CH	N	C ₂₄ H ₂₈ ClN ₅ O ₃	470.17	470.1
14-11		F	H	CH	N	C ₂₄ H ₂₈ FN ₅ O ₃	454.20	454.1
14-12		H	H	N	N	C ₂₀ H ₂₁ N ₅ O ₃	380.15	380.2

Table 15



Ex No.	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
15-1		Cl	H	CH	CH	C ₂₂ H ₂₀ ClN ₃ O ₃	410.10	409.9
15-2		F	H	CH	CH	C ₂₂ H ₂₀ FN ₃ O ₃	394.13	393.4















15-3		H	H	N	N	C ₂₀ H ₁₉ N ₅ O ₃	378.13	378.1
15-4		Cl	H	CH	CH	C ₂₅ H ₂₇ ClN ₄ O ₃	467.16	467.1
15-5		F	H	CH	CH	C ₂₅ H ₂₇ FN ₄ O ₃	451.19	451.1
15-6		H	H	CH	N	C ₂₄ H ₂₇ N ₅ O ₃	434.20	434.1
15-7		H	H	CH	N	C ₂₁ H ₂₀ N ₄ O ₃	377.14	377.2
15-8		H	H	N	CH	C ₂₁ H ₂₀ N ₄ O ₃	377.14	377.0
15-9		Cl	H	N	N	C ₂₀ H ₁₈ ClN ₅ O ₃	412.09	412.2
15-10		F	H	N	N	C ₂₀ H ₁₈ FN ₅ O ₃	396.12	396.1

Table 16

Ex No.	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
16-1		Cl	H	CH	CH	C ₂₃ H ₂₅ ClN ₄ O ₃	441.15	441.2
16-2		Cl	H	CH	CH	C ₂₀ H ₁₈ ClN ₃ O ₃	384.09	384.1
16-3		F	H	CH	CH	C ₂₀ H ₁₈ FN ₃ O ₃	368.12	368.2
16-4		H	H	N	CH	C ₁₉ H ₁₈ N ₄ O ₃	351.12	351.0
16-5		H	H	CH	N	C ₁₉ H ₁₈ N ₄ O ₃	351.12	351.0








16-6		H	H	CH	N	C ₂₂ H ₂₅ N ₅ O ₃	408.18	408.1
16-7		F	H	CH	CH	C ₂₃ H ₂₅ FN ₄ O ₃	425.18	425.2


Table 17

















Ex No.	R	R ¹	R ²	R ³	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
17-1			H	H	CH	CH	C ₁₉ H ₁₈ N ₄ O ₃	351.12	351.0
17-2			H	H	CH	CH	C ₁₉ H ₁₈ N ₄ O ₃	351.12	351.2

5

Table 18



Ex No.	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	X ¹	X ²	Formula	Calc [M+H] ⁺	Found [M+H] ⁺
18-1		Cl	H	H	H	Me	CH	CH	C ₂₄ H ₂₇ ClN ₄ O ₃	455.16	455.0
18-2		H	F	H	F	H	CH	CH	C ₂₃ H ₂₄ F ₂ N ₄ O ₃	443.17	443.2

18-3		F	H	H	F	H	CH	CH	C ₂₃ H ₂₄ F ₂ N ₄ O ₃	443.17	443.0
18-4		H	H	H	H	Me	CH	CH	C ₂₄ H ₂₈ N ₄ O ₃	421.2	421.0
18-5		H	H	H	H	Me	CH	CH	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364
18-6		Cl	H	H	H	Me	CH	CH	C ₂₁ H ₂₀ ClN ₃ O ₃	398.10	398.0
18-7		F	H	H	H	Me	CH	CH	C ₂₁ H ₂₀ FN ₃ O ₃	382.13	382.0
18-8		Me	H	H	H	H	CH	CH	C ₂₄ H ₂₈ N ₄ O ₃	421.2	421.0
18-9		Me	H	H	H	H	CH	CH	C ₂₁ H ₂₁ N ₃ O ₃	364.14	364.0
18-10		H	H	H	H	SMe	CH	CH	C ₂₁ H ₂₁ N ₃ O ₃ S	396.11	396.0
18-11		H	H	H	H	NMe ₂	CH	CH	C ₂₂ H ₂₄ N ₄ O ₃	393.17	393.0
18-12		H	H	H	H	CF ₃	CH	CH	C ₂₄ H ₂₅ F ₃ N ₄ O ₃	475.17	475.0
18-13		H	H	H	H	CF ₃	CH	CH	C ₂₁ H ₁₈ F ₃ N ₃ O ₃	418.11	418.0
18-14		H	H	H	H	Cl	CH	CH	C ₂₀ H ₁₈ ClN ₃ O ₃	384.09	384.2

Biological Assays

The compounds of the disclosure have been characterized in one or more of the following biological assays.

5 **Assay 1: Biochemical JAK and Tyk2 Kinase Assays**

A panel of four LanthaScreen JAK biochemical assays (JAK1, 2, 3 and Tyk2) were carried in a common kinase reaction buffer (50 mM HEPES, pH 7.5, 0.01% Brij-35, 10 mM MgCl₂, and 1 mM EGTA). Recombinant GST-tagged JAK enzymes and a GFP-tagged STAT1 peptide substrate were obtained from Life Technologies.

10 Serially or discretely diluted compounds were pre-incubated with each of the four JAK enzymes and the substrate in white 384-well microplates (Corning) at ambient temperature for 1h. ATP was subsequently added to initiate the kinase reactions in 10 μ L total volume, with 1% DMSO. The final enzyme concentrations for JAK1, 2, 3 and Tyk2

are 4.2 nM, 0.1 nM, 1 nM, and 0.25 nM respectively; the corresponding K_m ATP concentrations used are 25 μ M, 3 μ M, 1.6 μ M, and 10 μ M; while the substrate concentration is 200 nM for all four assays. Kinase reactions were allowed to proceed for 1 hour at ambient temperature before a 10 μ L preparation of EDTA (10mM final concentration) and Tb-anti-pSTAT1 (pTyr701) antibody (Life Technologies, 2nM final concentration) in TR-FRET dilution buffer (Life Technologies) was added. The plates were allowed to incubate at ambient temperature for 1h before being read on the EnVision reader (Perkin Elmer). Emission ratio signals (520 nm/495 nm) were recorded and utilized to calculate the percent inhibition values based on DMSO and background controls.

For dose-response analysis, percent inhibition data were plotted vs. compound concentrations, and IC_{50} values were determined from a 4-parameter robust fit model with the Prism software (GraphPad Software). Results were expressed as pIC_{50} (negative logarithm of IC_{50}) and subsequently converted to pK_i (negative logarithm of dissociation constant, K_i) using the Cheng-Prusoff equation.

Assay 2: Cellular JAK3 Potency Assay: Inhibition of IL-2 Stimulated pSTAT5 in Tall-1 T cells

The potency of test compounds for inhibition of interleukin-2 (IL-2) stimulated STAT5 phosphorylation was measured in the Tall-1 human T cell line (DSMZ) using AlphaLisa. Because IL-2 signals through JAK3, this assay provides a measure of JAK3 cellular potency.

Phosphorylated STAT5 was measured via the AlphaLISA SureFire Ultra pSTAT5 (Tyr694/699) kit (PerkinElmer).

Human T cells from the Tall-1 cell line were cultured in a 37°C, 5% CO₂ humidified incubator in RPMI (Life Technologies) supplemented with 15% Heat Inactivated Fetal Bovine Serum (FBS, Life Technologies), 2mM Glutamax (Life Technologies), 25mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). Compounds were serially diluted in DMSO and dispensed acoustically to empty wells. Assay media (phenol red-free DMEM (Life Technologies) supplemented with 10% FBS (ATCC)) was dispensed (4 μ L/well) and plates shaken at 900rpm for 10 mins. Cells were seeded at 45,000 cells/well in assay media (4 μ L/well), and incubated at 37°C, 5% CO₂ for 1 hour, followed by the addition of IL-2 (R&D Systems; final

concentration 300 ng/ml) in pre-warmed assay media (4 μ L) for 30 minutes. After cytokine stimulation, cells were lysed with 6 μ L of 3x AlphaLisa Lysis Buffer (PerkinElmer) containing 1x PhosStop and Complete tablets (Roche). The lysate was shaken at 900rpm for 10 minutes at room temperature (RT). Phosphorylated STAT5 was measured via the pSTAT5 AlphaLisa kit (PerkinElmer). Freshly prepared acceptor bead mixture was dispensed onto lysate (5 μ L) under green filtered <100 lux light. Plates were shaken at 900rpm for 2mins, briefly spun down, and incubated for 2hrs at RT in the dark. Donor beads were dispensed (5 μ L) under green filtered <100 lux light. Plates were shaken at 900rpm for 2 minutes, briefly spun down, and incubated overnight at RT in the dark. Luminescence was measured with excitation at 689 nm and emission at 570 nm using an EnVision plate reader (PerkinElmer) under green filtered <100 lux light.

To determine the inhibitory potency of test compounds in response to IL-2, the average emission intensity of beads bound to pSTAT5 was measured in a human T cell line. IC₅₀ values were determined from analysis of the inhibition curves of signal intensity versus compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values (mean \pm standard deviation).

Assay 3: Inhibition of IL-2 Stimulated pSTAT5 in CD4+ T cells isolated from murine splenocytes

The potency of test compounds for inhibition of interleukin-2 (IL-2) stimulated STAT5 phosphorylation was measured in the CD4+ T cells isolated from murine splenocytes using AlphaLisa. Because IL-2 signals through JAK3, this assay provides a measure of JAK3 cellular potency in mouse.

Phosphorylated STAT5 was measured via the AlphaLISA *SureFire Ultra* pSTAT5 (Tyr694/699) kit (PerkinElmer).

CD4+ T cells were isolated from murine splenocytes via negative selection on a magnetic column (Miltenyi Biotec) and re-suspended in assay media (phenol red-free DMEM (Life Technologies) supplemented with 10% FBS (ATCC)). Cells were seeded at 50,000 cells/well in assay media (2 μ L/well). Compounds were serially diluted in DMSO and diluted to 2x final concentration in assay media. Compound was added (4 μ L/well) and the cells incubated at 37°C, 5% CO₂ for 1 hour, followed by the addition of IL-2 (R&D Systems; final concentration 7 ng/ml) in pre-warmed assay media (2 μ L) for 30 minutes. After cytokine stimulation, cells were lysed with 2 μ L of 5x AlphaLisa Lysis Buffer (PerkinElmer). The lysate was shaken at 900rpm for 10 minutes at room temperature

(RT). Phosphorylated STAT5 was measured via the pSTAT5 AlphaLisa kit (PerkinElmer). Freshly prepared acceptor bead mixture was dispensed onto lysate (5ul) under green filtered <100 lux light. Plates were shaken at 900rpm for 2mins, briefly spun down, and incubated for 2hrs at RT in the dark. Donor beads were dispensed (5ul) under green filtered <100 lux light. Plates were shaken at 900rpm for 2mins, briefly spun down, and incubated overnight at RT in the dark. Luminescence was measured with excitation at 689 nm and emission at 570 nm using an EnVision plate reader (PerkinElmer) under green filtered <100 lux light.

To determine the inhibitory potency of test compounds in response to IL-2, the average emission intensity of beads bound to pSTAT5 was measured in primary CD4+ T cells isolated from murine splenocytes. IC₅₀ values were determined from analysis of the inhibition curves of signal intensity vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values (mean ± standard deviation).

Compounds 1-10, 19, 20, 21, and 22 all had pIC₅₀ values over 6.0 in this assay.

Assay 4: Inhibition of IL-2 Stimulated pSTAT5 in human PBMC CD4+ T cells

The potency of test compounds for inhibition of interleukin-2 (IL-2) stimulated STAT5 phosphorylation was measured in human peripheral blood mononuclear cell (PBMC) CD4+ T cells using flow cytometry. Because IL-2 signals through JAK3, this assay provides a measure of JAK3 cellular potency in human primary cells.

Phosphorylated STAT5 was measured via flow cytometry by measuring pSTAT5 response in CD4+ T cells (Becton Dickinson, AlexaFluor 647 Mouse anti-Human STAT5 (pY694) and PE Mouse anti-Human CD4).

Human PBMCs were isolated from donors via Ficoll gradient and frozen at -80°C. Frozen PBMCs were thawed and cultured for one hour in a 37°C, 5% CO₂ humidified incubator in RPMI (Life Technologies) supplemented with 10% Heat Inactivated Fetal Bovine Serum (FBS, Life Technologies), 2mM Glutamax (Life Technologies), 25mM HEPES (Life Technologies) and 1X Pen/Strep (Life Technologies). Cells were seeded into 2 ml deep well plates at 4e6 cells/ml (50 µl/well). Compounds were serially diluted in DMSO and diluted to 2x final concentration in assay media. Compound was added (100 µl/well) and the cells incubated at 37°C, 5% CO₂ for 1 hr, followed by the addition of IL-2 (R&D Systems; final assay concentration 100 ng/ml) in pre-warmed assay media (50 µL/well) for 30 minutes. After cytokine stimulation, cells were fixed for 10 mins at 37°C (Becton Dickinson). Cells were washed and re-suspended in Dulbecco's phosphate

buffered saline (DPBS, Life Technologies). Cells were permeabilized for 30 mins with ice cold Perm Buffer III (Becton Dickinson), and then washed and re-suspended with DPBS containing 2% fetal bovine serum (staining buffer, Life Technologies). Cells were stained for 1 hr with a 50x dilution of CD4 surface marker (PE Mouse anti-Human CD4) and 5x dilution of pSTAT5 antibody (Becton Dickinson, AlexaFluor 647 Mouse anti-Human STAT5 (pY694)) in staining buffer. Cells were washed and re-suspended in staining buffer before storage at 4°C.

To determine the inhibitory potency of test compounds in response to IL-2, the median fluorescence intensity of pSTAT5 in CD4+ gated cells was measured in human PBMCs with a BD LSRII with analysis in FCS Express 6. IC₅₀ values were determined from analysis of the inhibition curves of signal intensity vs compound concentration. Data are expressed as pIC₅₀ (negative decadic logarithm IC₅₀) values (mean ± standard deviation).

Compounds 1-6, 8 and 9 were tested in this assay and exhibited values very similar to the ones obtained in assay 2 (IL-2 Stimulated pSTAT5 in Tall-1 T cells assay).

Assay 5: JAK Cytotoxicity Assay

A CellTiter-Glo luminescent cell viability/cytotoxicity assay was carried out in BEAS-2B human lung epithelial cells (ATCC) under the normal growth condition.

Cells were grown at 37°C in a 5% CO₂ humidified incubator in 50% DMEM/50% F-12 medium (Life Technologies) supplemented with 10% FBS (Hyclone), 100 U/mL penicillin, 100 µg/mL streptomycin (Life Technologies), and 2 mM GlutaMAX (Life Technologies). On day 1 of the assay, cells were seeded at a 500 cells/well density in white 384-well tissue culture plates (Corning) with 25 µL medium, and were allowed to adhere overnight in the incubator. On day 2 of the assay, 5 µL of medium containing dose-responses of test compounds was added, and incubated at 37°C for 48 h. 30 µL of CellTiter-Glo detection solution (Promega) was subsequently added, mixed on an orbital shaker for 5 min, and incubated for additional 10 min before being read on the EnVision reader. Luminescence signals were recorded and percent DMSO control values were calculated.

For dose-response analysis, percent DMSO control data were plotted vs. compound concentrations to derive dose-response curves by line connecting each data point. The concentration at which each curve crosses the 15 % inhibition threshold is defined as CC₁₅. Results were expressed as the negative logarithm of the CC₁₅ value, pCC₁₅.

It is expected that test compounds exhibiting a lower pCC₁₅ value in this assay have less likelihood to cause cytotoxicity. Compounds of the disclosure tested in this assay typically exhibited pCC₁₅ values between 5 and about 6.

Assay 6: Caco-2 Permeation Assay

5 The Caco-2 permeation assay was performed to model the ability of test compounds to pass through the intestine and get into the blood stream after oral administration. The rate at which test compounds in solution permeate a cell monolayer designed to mimic the tight junction of human small intestinal monolayers was determined.

10 CacoReady 24-well transwell plates were obtained from ADMEcell (Alameda, CA). The compounds were evaluated at a concentration of 5 μ M from 10 mM DMSO stock solutions in duplicate (n=2). The passive permeability of the compounds tested was evaluated using Caco-2 cell monolayers along with Verapamil (25 μ M) to inhibit P-gp transport proteins in the apical to basolateral (A-B) direction. The experiment was
15 conducted in a 37°C, 5% CO₂ incubator. Caco-2 culture media consisted of standard filtered DMEM, FCS 10%, L-Glutamine 1% and PenStrep 1%. Basal assay plate was prepared by adding 750 μ L of transport buffer to A-B wells. A CacoReady™ plate was prepared by removing the Caco-2 media from the apical wells and replacing with fresh transport media (200 μ L repeated for a total of 3 washes). Blank media (200 μ L) was
20 then replaced with diluted compound for A-B wells. To begin the incubation, the basal plate was removed from the incubator and the apical section was added on top of it. Samples (40 μ L) were collected from the apical and basal compartments for time zero (t₀). Samples were collected again after 120 minutes (t₁₂₀) from the apical and basal compartments. All samples were diluted and prepared for bioanalysis by LC-MS/MS. The
25 permeation coefficient (K_p, mean A to B + Verapamil P_{apparent}) in cm/sec was calculated as dQ (flux)/(dt x Area x concentration).

In this assay, a K_p value less than about 5 x 10⁻⁶ cm/sec is considered favorable to minimize systemic exposure and target the colon. A K_p value less than about 10 x 10⁻⁶ cm/sec may also be sufficient to minimize systemic exposure and target the colon. By
30 comparison, PF-06651600, a JAK3 inhibitor available systemically (2-propen-1-one, 1-[(2*S*,5*R*)-2-methyl-5-(7*H*-pyrrolo[2,3-*d*]pyrimidin-4-ylamino)-1-piperidinyl]) exhibited a K_p value of 25.

In Vitro Assay Results

All of the compounds of Examples 1 to 18 and Tables 1 to 12 were tested in one or more of the assays described above.

In Table 13 below, for the JAK1, JAK 2, JAK3, and TYK2 enzyme assays, A represents a pK_i value ≥ 10 (K_i ≤ 0.1 nM), B represents a pK_i value between 9 and 10 (K_i between 1 nM and 0.1 nM), C represents a pK_i value between 8 and 9 (K_i between 10 nM and 1 nM), D represents a pK_i value between 7 and 8 (K_i between 100 nM and 10 nM), and E represents a pK_i value of 7 or below (K_i of 100 nM or above). For the Tall-1 Potency assay, A represents a pIC₅₀ value ≥ 7.5 (IC₅₀ ≤ 32 nM), B represents a pIC₅₀ value between 6.7 (included) and 7.5 (IC₅₀ between 200 nM and 32 nM), and C represents a pIC₅₀ value between 6 and 6.7 (IC₅₀ between 1 μ M and 200 nM). For the JAK3 (pK_i)-JAK1 (pK_i) values, A represents a value of 3 or above, B represents a value of 2.3 to 3 and C represents a value of 1.8 to 2.3. For the Caco assay, A represents a value below 5 x 10⁻⁶ cm/sec, B represents a value between 5 x 10⁻⁶ and 10 x 10⁻⁶ cm/sec, C represents a value between 10 x 10⁻⁶ and 22 x 10⁻⁶ cm/sec.

Table 13

Example Number	JAK 1 (pK_i)	JAK 2 (pK_i)	JAK 3 (pK_i)	Tyk 2 (pK_i)	Tall-1 (pIC₅₀)	JAK3 (pK_i)-JAK1 (pK_i)	Caco K_p 10⁻⁶ cm/sec
1	E	D	A	E	B	A	A
2	E	E	A	E	A	A	A
3	E	E	B	E	B	A	A
4	E	E	A	E	B	A	A
5	E	D	A	E	B	A	A
6	E	D	A	E	A	A	A
7	E	D	B	E	B	A	B
8	E	D	A	E	A	A	A
9	E		B		B	A	A
10	E	E	A	E	B	A	A
11	E	E	B	E	B	A	A
12	E	E	B	E	B	B	A
13	E	E	A	E	B	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
14	E	E	B	E	B	A	C
15	E		B		C	A	A
16	E		B		B	A	A
17	E		B		C	B	
18	E	D	B	E	A	A	C
19	E	E	A	E	B	A	A
20	E	E	A	E	A	A	A
21	E	E	A	E	B	A	B
22	E	E	A	E	B	A	A
23	E	D	A	E	B	A	A
24	E	E	B	E	A	A	A
1-1	E	D	B	E		A	
1-2	E	D	A	E	A	A	C
1-3	E		B		B	A	
1-4	E		B		C	A	
1-5	E	D	A	E	A	A	B
1-6	E		B		C	A	
1-7	E		B		C	B	
1-8	E	E	B	E	B	A	B
1-9	E		A		C	A	
1-10	E		B		C	A	
1-11	E	E	A	E	A	A	C
1-12	E		B		B	A	
1-13	E		B		B	B	A
1-14	E		A		B	A	A
1-15	E	E	A	E	B	A	A
1-16	E	D	B	E	B	B	A
1-17	E	D	B	E	A	A	A
1-18	D	D	B	E	A	B	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
1-19	E	D	B	E	B	A	A
1-20	E	D	B	E	A	B	A
1-21	E	E	B	E	B	A	A
1-22	E	D	A	E	B	A	A
1-23	D		B		A	B	A
1-24	E	E	B	E	C	A	C
1-25	E		B		B	A	C
1-26	E	E	A	E	B	A	B
1-27	E	D	A	E	B	A	
1-28	E		B		C	B	
1-29	D	D	B	E	B	B	
1-30	D		A		B	B	
1-31	E	E	A	E	B	A	
1-32	E		B		C	A	
1-33	E	E	B	E	B	A	
1-34	E	D	B	E	A	A	
1-35	D		A		A	A	
1-36	D	D	B	E	A	B	
1-37	E		B		C	A	
1-38	E	E	A	E	B	A	
1-39	E		B		B	A	B
1-40	E	E	B	E	B	B	
1-41	E	E	B	E	B	A	
1-42	E		B		C	A	B
1-43	E		A		A	A	
1-44	E	E	A	E	A	A	C
1-45	E	E	A	E	A	A	
1-46	E	E	B	E	C	A	
1-47	E	E	B	E	B	B	C

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K_p 10⁻⁶ cm/sec
1-48	E		C		B	B	
2-1	D		A		C	B	
2-2	E	D	B	E	B	A	A
2-3	E	D	B	E	B	A	B
2-4	E		B		C	B	
2-5	E	D	B	E	B	A	B
2-6	E		A		C	A	C
2-7	E		B		B	A	A
2-8	E	D	B	E	B	A	A
2-9	E		B		B	B	A
2-10	E	D	B	E	A	A	A
2-11	E	E	B	E	B	A	B
2-12	E		A		C	A	C
2-13	E		B		C	B	
2-14	E		A		A	A	
2-15	E	E	A	E	B	A	C
2-16	E		B		C	A	
2-17	E	E	A	E	B	A	
3-1	E	E	B	E	A	A	A
3-2	E	E	B	E	B	A	A
3-3	E	E	B	E	A	A	B
3-4	E	D	B	E	A	A	B
3-5	D	D	B	E	A	B	A
3-6	D		A		A	A	
3-7	E	D	A	E	B	A	A
3-8	E	E	B	E	A	A	A
3-9	E		B		B	B	A
3-10	E		B		A	A	A
3-11	E	D	A	E	B	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
3-12	E		B		B	B	A
3-13	E	D	B	E	A	A	A
3-14	E		B		B	A	
3-15	E	E	A	E	B	A	A
3-16	E		A		B	A	A
3-17	E	D	A	E	A	A	A
3-18	E		A		B	A	A
3-19	E	E	B	E	B	A	A
3-20	E	D	B	E	B	B	A
3-21	E	E	B	E	A	A	A
3-22	E	E	A	E	B	A	A
3-23	E	D	A	E	B	A	A
3-24	E	E	B	E	B	A	A
3-25	E		C		B	B	A
3-26	E		B		B	A	A
3-27	E	E	A	E	B	A	A
3-28	E		A		A	A	B
3-29	E		B		A	A	A
3-30	E	D	A	E	B	A	A
3-31	E	E	A	E	B	A	B
3-32	E	E	B	E	A	A	A
3-33	E	E	B	E	B	B	A
3-34	E	E	A	E	B	A	A
3-35	E	E	B	E	A	A	A
3-36	E	E	A	E	B	A	A
3-37	E	D	A	E	A	A	A
3-38	E	E	A	E	B	A	A
3-39	E	E	A	E	B	A	A
3-40	B	D	A	E	A	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K_p 10⁻⁶ cm/sec
3-41	B	C	B	E	A	B	A
3-42	B	D	B	E	A	B	A
3-43	E	D	B	E	A	A	A
3-44	D	D	B	E	A	B	A
3-45	E	E	B	E	A	A	A
3-46	E	E	A	E	A	A	A
3-47	E	D	B	E	A	B	A
3-48	D	D	A	E	A	A	A
3-49	E	D	B	E	A	B	A
3-50	E	E	B	E	B	A	A
3-51	E	E	B	E	C	A	A
3-52	E	E	B	E	B	A	A
3-53	E	E	B	E	B	A	A
3-54	E	E	B	E	A	A	A
3-55	E	E	A	E	B	A	A
3-56	E	D	A	E	A	A	A
3-57	E	E	A	E	B	A	A
3-58	E	E	A	E	B	A	A
3-59	E	D	B	E	A	B	A
3-60	E	E	A	E	B	A	
3-61	E	D	A	E	A	A	A
3-62	E	D	B	E	A	A	A
3-63	E	E	A	E	B	A	A
3-64	E	D	A	E	A	A	A
3-65	E	D	B	E	A	A	A
3-66	E	E	B	E	A	A	A
3-67	E	E	B	E	B	A	A
3-68	D	E	B	E	C	B	A
3-69	E	E	A	E	B	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
3-70	E	E	A	E	B	A	A
3-71	E	E	A	E	A	A	A
3-72	E	E	B	E	A	A	A
3-73	E	E	A	E	A	A	A
3-74	E	E	A	E	A	A	A
3-75	E	E	B	E	A	A	A
3-76	E		A		B	A	A
3-77	E	E	A	E	B	A	A
3-78	E	E	B	E	B	A	A
3-79	E	E	B	E	B	A	
3-80	E	E	A	E	B	A	A
3-81	E	E	A	E	A	A	A
3-82	D		A		A	B	A
3-83	E	E	A	E	A	A	B
3-84	E		A		B	A	
3-85	E	D	A	E	A	A	A
3-86	E	E	A	E	B	A	
3-87	E	E	B	E	B	A	
3-88	E	E	B	E	B	A	
3-89	E	E	A	E	B	A	
3-90	D		A		A	B	
3-91	D	D	A	E	A	A	
3-92	E	E	A	E	A	A	
3-93	E	E	A	E	A	A	
3-94	E	E	A	E	A	A	
3-95	E	E	A	E	A	A	
3-96	E	E	B	E	B	A	
3-97	E	E	B	E	A	B	
3-98	E	E	A	E	B	A	

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K_p 10⁻⁶ cm/sec
3-99	E	E	A	E	A	A	
3-100	E	E	B	E	A	A	
3-101	E	D	A	E	A	A	
3-102	E	E	B	E	B	A	
3-103	E	D	A	E	A	A	
3-104	E	E	A	E	A	A	
3-105	E	D	A	E	A	A	
3-106	E	E	B	E	B	A	
3-107	E	E	A	E	A	A	
3-108	E	D	A	E	A	A	
3-109	E	E	B	E	B	A	
3-110	E	E	B	E	B	A	
3-111	E	E	B	E	C	A	
3-112	E	E	B	E	B	A	A
3-113	E	D	A	E	A	A	A
3-114	E	E	B	E	A	A	
3-115	E	B	B	E	B	A	A
3-116	E	D	B	E	B	B	
3-117	E	D	B	E	B	B	
3-118	E	E	B	E	A	A	
3-119	E	D	A	E	B	B	A
3-120	E	D	A	E	B	A	A
3-121	E	D	B	E	B	A	A
3-122	E	D	A	E	A	A	
3-123	E		A		C	A	
3-124	E		B		B	A	
3-125	E	E	B	E	B	A	
3-126	E	D	A	E	C	A	
3-127	E	D	A	E	B	A	

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K_p 10⁻⁶ cm/sec
3-128	E	B	A	E	B	A	
3-129	E		B		C	A	A
3-130	E		B		B	A	A
3-131	E	B	B	E	A	B	
3-132	E	E	B	E	C	A	
3-133	E	D	A	E	C	A	
3-134	E		B		B	A	
4-1	E	E	B	E	B	A	A
4-2	D	D	B	E	B	B	A
4-3	E		B		B	B	A
4-4	E	E	A	E	B	A	A
4-5	E	E	B	E	A	A	A
4-6	E	E	B	E	B	A	A
4-7	E	E	C	E	B	B	A
4-8	E		B		B	B	
5-1	E		B		C	A	
5-2	E	E	B	E	B	A	A
5-3	E	E	A	E	A	A	A
5-4	E	E	B	E	A	A	A
5-5	E	D	A	E	B	A	A
5-6	E	E	B	E	B	A	A
5-7	E	D	A	E	B	A	A
5-8	E		B		C	A	
5-9	E		B		C	A	A
5-10	E		B		B	A	A
5-11	E		B		B	A	A
5-12	E	E	B	E	B	A	A
5-13	E		B		B	B	A
5-14	E	E	A	E	A	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
5-15	E		A		B	A	A
5-16	E	6	A	E	A	A	A
5-17	E	E	B	E	A	A	A
5-18	E	E	A	E	A	A	A
5-19	E	E	B	E	B	A	A
5-20	E		B		C	A	
5-21	E	E	B	E	A	A	
5-22	E	E	B	E	B	A	
5-23	E	E	B	E	B	A	
5-24	E		B		B	A	
5-25	E		B		A	A	
5-26	E		B		C	B	
6-1	D		A		B	B	
6-2	D	D	A	E	A	B	C
6-3	E	D	B	E	A	A	C
6-4	E	D	A	E	B	A	A
6-5	E		B		A	A	A
6-6	E	D	B	E	A	A	A
6-7	E		B		A	A	B
6-8	D		A		A	B	
6-9	E		B		B	A	A
6-10	E	D	B	E	A	A	C
6-11	E		A		B	A	B
6-12	E		B		B	A	A
6-13	E	D	B	E	A	B	A
6-14	D		A		A	B	B
6-15	E	D	B	E	B	A	B
6-16	E	D	B	E	B	A	A
6-17	E	E	A	E	B	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
6-18	E	E	A	E	B	A	B
6-19	D		B		B	B	
6-20	D	C	B	E	B	B	A
6-21	D	D	B	E	B	B	
6-22	E		B		C	A	
6-23	E		B		C	A	
7-1	E		B		C	A	
7-2	E	D	B	E	A	A	A
7-3	E	E	B	E	B	A	C
7-4	D	C	B	E	A	B	C
7-5	D		B		B	C	
7-6	D		A		A	B	
7-7	E	D	B	E	B	A	A
7-8	E	E	A	E	B	A	A
7-9	E	D	A	E	A	A	A
7-10	E	D	B	E	B	A	C
7-11	E	D	A	E	A	A	
7-12	E	D	B	E	B	A	
7-13	E	D	A	E	A	A	
7-14	E	D	B	E	A	B	
8-1	E	D	B	E	A	B	C
8-2	E	D	B	E	B	A	A
8-3	E	D	B	E	A	A	A
8-4	E	D	B	E	B	B	A
8-5	E	D	A	E	A	A	A
8-6	E		B		B	A	B
8-7	E	D	B	E	B	A	B
8-8	E		A		B	A	B
8-9	E		B		C	B	

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
8-10	D		A		A	A	
8-11	D	D	A	E	A	B	
8-12	E		B		C	A	
8-13	E		B		C	A	
8-14	E	E	A	E	B	A	C
9-1	E	D	A	E	A	A	C
9-2	E	E	B	E	A	A	B
9-3	E		B		C	C	
9-4	E	C	A	E	A	A	C
9-5	D		B		C	C	
9-6	E		B		A	A	C
9-7	E		A		B	A	
9-8	D		B		C	C	
9-9	D	C	A	D	A	B	
9-10	E		B		A	A	C
9-11	E		A		B	A	C
10-1	D		A		B	B	A
10-2	D	C	A	D	A	B	C
10-3	D		B		B	B	
10-4	E	D	B	E	B	A	A
10-5	D		B		B	C	
10-6	D		B		A	B	
10-7	E	D	B	E	B	A	B
11-1	E		B		A	A	A
11-2	E		B		C	A	
12-1	E	D	B	E	B	A	
12-2	E		B		B	B	
12-3	D		B		C	B	
12-4	E	E	B	E	B	A	B

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K_p 10⁻⁶ cm/sec
12-5	E		B		B	B	
12-6	D		B		B	B	
12-7	D		B		C	C	
12-8	E		B		C	A	A
12-9	E		B		C	A	A
12-10	E		C		B	B	
12-11	E		B		C	A	
12-12	E		B		B	A	
12-13	E		B		C	A	
12-14	E		B		C	C	
12-15	E		B		B	B	
12-16	D	C	B	E	A	B	
12-17	D	C	A	E	A	B	
12-18	D		A		B	B	
12-19	E		A		C	A	
12-20	E	D	B	E	A	A	B
12-21	E		B		A	B	A
12-22	E		B		C	A	A
12-23	E		B		B	A	B
12-24	E		B		C	A	A
12-25	E		B		C	B	A
12-26	E	D	B	E	C	B	
12-27	E	D	A	E	A	A	C
12-28	E	C	A	E	B	A	C
12-29	E	D	B	E	A	A	C
12-30	E	C	A	E	A	A	C
12-31	E	C	A	E	B	A	C
12-32	E	D	A	E	A	A	C
12-33	E	E	B	E	B	A	A

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
12-34	E	E	B	E	A	B	A
12-35	E	E	B	E	B	B	A
12-36	D	E	A	E	A	A	A
12-37	D		B		B	B	A
12-38	E	E	A	E	B	A	A
12-39	E	E	B	E	A	A	B
12-40	E		B		C	B	A
12-41	E	E	B	E	A	A	A
12-42	E	D	B	E	B	B	A
12-43	E	D	A	E	A	A	A
12-44	E	D	A	E	B	A	A
12-45	E	E	A	E	C	A	A
12-46	E	D	A	E	B	A	A
12-47	E	E	B	E	B	A	A
12-48	E	E	B	E	A	A	A
12-49	E	E	B	E	C	A	
12-50	E	E	A	E	B	A	
12-51	D	D	A	E	B	A	
12-52	D		B		C	B	
12-53	E	C	B	E	A	B	
12-54	E	D	B	E	A	B	
12-55	E		A		B	A	
12-56	E		A		B	A	
12-57	E	D	B	E	B	A	
12-58	E	D	A	E	B	A	
12-59	E		A		B	A	
12-60	E	E	B	E	B	A	
12-61	E	E	A	E	B	A	
12-62	E		A		B	A	

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
12-63	E	E	B	E	B	A	
12-64	E	E	A	E	A	A	
12-65	E	E	B	E	C	A	
12-66	E	E	B	E	B	A	
12-67	E	E	B	E	B	A	
12-68	E	E	B	E	C	A	
12-69	E		A		B	A	
13-1	E	E	B	E	B	A	A
13-2	E	E	B	E	A	A	A
13-3	E		A		A	A	A
13-4	E	D	A	E	B	A	
13-5	E	D	A	E	B	A	
13-6	E	E	A	E	A	A	
13-7	E	D	A	E	B	A	
13-8	E	E	B	E	A	A	
13-9	E	E	B	E	B	A	
13-10	E	E	B	E	B	A	
13-11	E	E	A	E	A	A	C
13-12	E	E	B	E	B	A	C
13-13	E	E	B	E	C	A	C
13-14	E	D	A	E	A	A	
13-15	E	D	B	E	B	A	
13-16	E	E	B	E	B	A	C
13-17	E		A		B	A	
13-18	E		B		B	A	
13-19	E	E	A	E	B	A	
13-20	E	E	B	E	B	B	
13-21	E	D	B	E	A	B	
13-22	E	E	B	E	B	A	

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K_p 10⁻⁶ cm/sec
14-1	E	E	B	E	A	A	A
14-2	E	E	A	E	B	A	A
14-3	E		B		B	B	
14-4	D		B		B	B	
14-5	D		B		A	B	
14-6	D		A		A	B	
14-7	D		A		A	B	
14-8	D		A		A	B	
14-9	E	E	B	E	B	A	
14-10	D		B		C	C	
14-11	D		B		C	B	
14-12	E	E	B	E	C	A	
15-1	E	E	B	E	A	A	A
15-2	D	D	A	E	A	B	A
15-3	E	E	A	E	A	A	A
15-4	E		A		B	A	
15-5	D	D	B	E	B	B	
15-6	D		A		C	B	
15-7	E	D	B	E	A	B	
15-8	D		B		B	B	
15-9	E	E	A	E	B	A	
15-10	E	E	A	E	A	A	
16-1	E	E	A	E	B	A	A
16-2	E	E	A	E	A	A	A
16-3	E	E	B	E	A	A	A
16-4	E		B		B	A	A
16-5	E	E	A	E	A	A	A
16-6	E	E	B	E	C	A	
16-7	E	E	A	E	B	A	

Example Number	JAK 1 (pKi)	JAK 2 (pKi)	JAK 3 (pKi)	Tyk 2 (pKi)	Tall-1 (pIC ₅₀)	JAK3 (pKi)-JAK1 (pKi)	Caco K _p 10 ⁻⁶ cm/sec
17-1	E	E	B	E	B	A	A
17-2	E		A		C	A	A
18-1	E	D	B	E	A	B	A
18-2	D		A		A	C	
18-3	D	D	A	E	A	B	
18-4	D		A		B	B	
18-5	D		B		A	B	
18-6	D		B		A	B	
18-7	D		B		A	C	
18-8	E	E	B	E	B	A	
18-9	E	E	A	E	A	A	
18-10	E	D	A	E	B	A	
18-11	E		B		C	A	
18-12	D	D	B	E	B	B	
18-13	E	D	A	E	A	A	
18-14	E		B		B	A	

Assay 7: Colon and Plasma Mouse Pharmacokinetics

6 male Balb/c mice were administered 10 mg/kg of compound in 1% HPMC + 0.1% Tween-80 by PO administration. At 0.5, 2 and 6 hours after dose administration, animals were anesthetized, and terminal blood samples were collected by cardiac puncture, followed by collection of colon contents and colon tissue.

Blood samples were collected into K₂EDTA and stored on wet ice until processed to plasma by centrifugation (12000 rpm at 4°C). Plasma samples were transferred to cluster tubes and placed on dry ice prior to freezer storage. The colon contents from each animal were collected at each terminal blood collection time point. The colon tissues were flushed with saline and patted dry. The colon and colon content tissues were homogenized using sterile water containing 0.1% formic acid 9:1 (water:tissue, v/w). The homogenized tissues and colon contents were transferred to cluster tubes and placed

on dry ice prior to freezer storage. All samples were analyzed using LC/MS/MS against analytical standards.

The composite pharmacokinetic parameters of the compounds were determined by non-compartmental analysis using Phoenix WinNonlin Version 6 (Certara, St. Louis, MO) and using mean values from 2 animals/time point. For plasma concentrations below the quantification limit (BQL), the lowest concentration measurable or the BLOQ (below limit of quantification) was used.

A colon to plasma ratio was determined as the ratio of the colon AUC to the plasma AUC. Compounds 1, 2, 3, 4, 6, 7, 8, 21 and 22 exhibited a colon to plasma ratio in excess of about 1250. Compounds 9, 5, 19, and 20 exhibited a colon to plasma ratio in excess of about 200.

In contrast, the reference compound (PF-06651600, a JAK3 inhibitor available systemically) 2-propen-1-one, 1-[(2*S*,5*R*)-2-methyl-5-(7*H*-pyrrolo[2,3-*d*]pyrimidin-4-ylamino)-1-piperidinyl], exhibited a colon to plasma ratio of 2.8

15

Assay 8: Mouse Model of Oxazolone-induced Colitis

Oxazolone-induced colitis is an experimental model that has a histological resemblance to human ulcerative colitis (Heller et al. *Immunology*, **2002**, 17, 629-638). Adult BALB/C mice from Harlan were used in the assay. On day 1, animals were lightly anesthetized with isoflurane and the hairs between the shoulder were carefully removed before oxazolone (4%, 150 μ L, 4:1 acetone: olive oil formulation) or vehicle solution was slowly applied for skin sensitization. Seven days after skin sensitization, the mice were fasted around 6 hours prior, anesthetized with ketamine/xylazine injection, and a 1 mL syringe equipped with a 3.5-F catheter, filled with oxazolone solution, was inserted carefully about 4 cm into the colon of the mouse. Following insertion, 50 μ L of the oxazolone solution (0.8% in 1:1 ethanol:water formulation) was injected very slowly (over 30 sec using an injection pump) into the colon. Drug treatment (PO, QD or BID or TID) or vehicle was initiated a day prior to the oxazolone intrarectal (IR) challenge. Two-day post oxazolone intrarectal challenge, the disease was assessed by treatment-blinded experimenters for each mouse according to the criteria score: stool consistency score (0, normal; 2, loose; 4, diarrhea), gross bleeding score (0, absence; 2, blood tinged; 4, presence); Combined stool score endpoint = stool consistency score + stool blood score.

30

Select compounds were tested in the assay. Efficacy in the model is evidenced by a statistically significant decrease in combined stool score endpoint as compared with the score from vehicle treated animals.

The compounds 1, 2, 3, 4, 5, 6, 7, 8, 3-11, 5-10, 19, 15-1, 3-55, 3-34, 15-3, 21, 3-80, 3-81, 3-72 and 3-57 exhibited a statistically significant decrease in combined stool score endpoint as compared with vehicle treated animals in the oxazolone model at a dose of 3 mg/kg (PO, BID). The compounds 3-113 and 3-74 exhibited a statistically significant decrease in the combined stool score endpoint as compared with vehicle treated animals in the oxazolone model at a dose of 1 and 10 mg/kg only (PO, BID).

10

Assay 9: Immunosuppression Effects in Mouse Splenic Natural Killer (NK) Cells

Depletion of mouse splenic cells is an experimental model of immunosuppression (Kudlacz et al., *Am. J. of Transplantation*, **2004**, 4, 51-57). Select compounds were assessed in the mouse splenic cell model following the same treatment paradigm used in the oxazolone-induced colitis model (assay 8).

Adult male Balb/C mice from Harlan were used for the study. The test compound and a Jak3 covalent inhibitor standard 2-propen-1-one, 1-[(2*S*,5*R*)-2-methyl-5-(7*H*-pyrrolo[2,3-*d*]pyrimidin-4-ylamino)-1-piperidinyl] (30 mg/kg, BID) as a positive control, were dosed orally for three days to naïve mice. Spleens were harvested up to 4 hours post last dose and crushed immediately for cell subtype staining. Prior to fixation, fluorophore-labelled antibodies for CD19 (FITC; B cells), CD3e (PE; pan T cells) and CD49e (APC; NK cells) were incubated with splenocyte samples from each animal to allow for simultaneous, multiple subtype % analysis on the flow cytometer. The number of total spleen cells for each animal was measured by Scepter™ 2.0 Handheld Automated Cell Counter.

The absolute number of lymphocyte subtype population (e.g., splenic B, T and NK cells) was calculated from the percentage of each subtype times the total spleen cells for each animal. A one-way ANOVA, with Fisher's LSD *post hoc* test, was used to compare the splenic lymphocytes number of the vehicle and test compound groups. The α level was set at $p < 0.05$. Data were presented as the mean \pm SEM for each group.

The positive control PF-06651600, a JAK3 inhibitor available systemically (2-propen-1-one, 1-[(2*S*,5*R*)-2-methyl-5-(7*H*-pyrrolo[2,3-*d*]pyrimidin-4-ylamino)-1-piperidinyl]) (30 mg/kg PO, BID), significantly decreased splenic NK cell counts. Splenic

NK cell counts were unaffected by compound 1 at (PO, BID) doses up to 300 mg/kg (the maximum dose tested). Splenic NK cell counts were unaffected by compounds 4, 6 and 8 at (PO, BID) doses up to 100 mg/kg (the maximum dose tested). Splenic NK cell counts were unaffected by compound 3 at (PO, BID) doses up to 85 mg/kg (the maximum dose tested). Splenic NK cell counts were unaffected by compound 2 at (PO, BID) doses up to 30 mg/kg (the maximum dose tested). Splenic NK cell counts were unaffected by compound 5 at (PO, BID) doses up to 80 mg/kg (the maximum dose tested). No treatment effects were observed for the B and T cell populations with any of the compounds tested.

10

Assay 10: Systemic Target Engagement Assay: Murine model of IL-2 induced pSTAT5 induction in thymus

IL-2 is an important cytokine underlying the pathophysiology of gastro-intestinal diseases such as irritable bowel disease (IBD; Guan and Zhang. Mediators Inflamm, 2017; 4810258). IL-2 binds to cell surface receptors activating members of the Janus family of kinases (JAK), specifically JAK3, which then phosphorylates STAT5 and subsequently activates further transcription pathways. In this model, a dose of IL-2 was delivered systemically to mice to induce the phosphorylation of STAT5 (pSTAT5) which was then measured as the endpoint. The compounds tested in this assay which do not exhibit significant inhibition of IL-2 induced pSTAT5 compared to the vehicle-treated; IL-2 challenged control animals, demonstrate lack of systemic activity.

Adult male Balb/c mice from Harlan were used in the assay. Animals were administered test compound via oral gavage (PO, 10 or 30 mg/kg for single dose studies; 10 – 60 mg/kg for dose response studies) on the afternoon of the day prior to study and then in the morning on the day of study. Two hours after the second PO dose administration, mice were injected intraperitoneally (i.p.) with 100µl of an appropriate IL-2 dose (for a total of 1 - 5 µg/mouse depending upon IL-2 batch; R&D Systems). 90 - 120 minutes after IL-2 insult, thymus samples were harvested. Phosphorylated STAT5 (pSTAT5) levels were determined in the thymus using AlphaLISA (AlphaLISA® SureFire® Ultra™ p-STAT5 (Tyr694/699) HV (high volume). Activity in the model is evidenced by a lack of significant inhibition in the level of pSTAT5 present in the thymus of treated animals at 1.5 -2 hours following IL-2 challenge.

30

Compounds 6 and 7 were tested at 10mg/kg and exhibited no significant inhibition of IL-2 induced pSTAT5 compared to the vehicle-treated; IL-2 challenged control

animals, therefore demonstrating lack of systemic activity. Compound 1 was tested at 10, 30, 60 and 100 mg/kg and exhibited no significant inhibition of IL-2 induced pSTAT5 compared to the vehicle. Compound 2 was tested at 10, 30 and 60 mg/kg and exhibited no significant inhibition of IL-2 induced pSTAT5 compared to the vehicle. Compound 3 was
5 tested at 30 and 100 mg/kg and exhibited no significant inhibition of IL-2 induced pSTAT5 compared to the vehicle. Compounds 8, 5 and 4 were tested at 30 mg/kg and exhibited no significant inhibition of IL-2 induced pSTAT5 compared to the vehicle.

In contrast, the reference compound (PF-06651600, a JAK3 inhibitor available systemically) 2-propen-1-one, 1-[(2*S*,5*R*)-2-methyl-5-(7*H*-pyrrolo[2,3-*d*]pyrimidin-4-ylamino)-1-piperidinyl], exhibited significant inhibition of pSTAT5 induction in the
10 thymus at 10, 30 and 50 mg/kg, demonstrating systemic activity.


Crystal Structures

Co-crystal structures were obtained for compounds 1, 3 and 4 each bound to
15 human JAK3. The resolutions for the complex structures were at 2.73 Å, 2.80 Å, and 2.64 Å respectively for compound 1, compound 3 and compound 4. In each case, the ligands were observed to bind in the ATP binding site. Based on a distance of less than 3.5 Å of the donor and acceptor atoms four specific hydrogen bonds were identified for each of the ligands, namely to the main chain atoms of Glu903, Leu905 and Phe968 as well as the side chain atoms of
20 Glu871. An additional hydrogen bond was identified for compound 1 to the side chain atoms of Asp 912. Of particular note, the ligands are each covalently bound to Cys909 of JAK3 located shortly after the hinge region. The S-H moiety of the cysteine residue performs an addition reaction to the Michael system. The observed results of the covalent binding interaction in the crystal structures confirms the irreversible binding nature for each of
25 these ligands to JAK3.

While the present invention has been described with reference to specific aspects or embodiments thereof, it will be understood by those of ordinary skilled in the art that various changes can be made or equivalents can be substituted without departing from the
30 true spirit and scope of the invention. Additionally, to the extent permitted by applicable patent statutes and regulations, all publications, patents and patent applications cited herein are hereby incorporated by reference in their entirety to the same extent as if each document had been individually incorporated by reference herein.

WHAT IS CLAIMED IS:


1. A compound of formula (I):



- 5 or a pharmaceutically acceptable salt thereof, wherein


X^1 and X^2 are each independently selected from N and CH;

X^3 is selected from the group consisting of N, CH, C-CH₃, C-CF₃, C-CHF₂, C-CH₂-O-CH₃, C-SMe, C-NMe₂, C-NH-CH₃, C-Cl, C-CN, and C-OMe;



is selected from the group consisting of

10



R^a , R^b , R^c , and R^f are each independently selected from the group consisting of H and C₁₋₃ alkyl;

- 15 R^d , R^e , R^g , R^h , R^i , R^j , R^l , R^m , R^n and R^o are each independently selected from the group consisting of H and C₁₋₃ alkyl wherein the C₁₋₃ alkyl group may be optionally substituted with 1 to 3 halogens;

optionally R^d and R^e may be joined to form a cyclopropyl ring;

A is selected from the group consisting of


(a) a 4 to 10 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, S(O)₂ and O, and

5 (b) a 6 to 10 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O,

wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 3 R^k groups;

each R^k is independently selected from the group consisting of F, CN, C₁₋₃ alkoxy, cyclopropyl, and C₁₋₃ alkyl, wherein the C₁₋₃ alkyl group may be optionally substituted with OH, OMe or 1 to 3 halogens;

R¹ is selected from the group consisting of



15 wherein R^p and R^q are each independently selected from the group consisting of H, C₃₋₅ cycloalkyl and C₁₋₆ alkyl, wherein the C₁₋₆ alkyl group may be optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₃ alkoxy and -S-C₁₋₃ alkyl,

or R^p and R^q form a 4 to 6 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, and O, wherein the 4 to 6 membered monocyclic heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₆ alkyl, C₁₋₃ alkoxy, -S-C₁₋₃ alkyl and -C₁₋₃ alkyl-C₁₋₃ alkoxy;

R² is selected from the group consisting of H, Cl, OMe, Me and F;


25 R³ is selected from the group consisting of H and F;

R⁴ is selected from the group consisting of H and F; and

R⁵ is selected from the group consisting of H, Me and F.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein X^3 is CH.

3. The compound of any one of claims 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^1 is selected from the group consisting of



4. The compound of any one of claims 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^1 is


15


or

5. The compound of any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, wherein A is selected from the group consisting of azetidine, pyrrolidine, piperidine, morpholine, 2-azaspiro[3.3]heptane, thiomorpholine, and nortropane.


20


6. The compound of any one of claims 1 to 5, or a pharmaceutically

acceptable salt thereof, wherein  is selected from the group consisting of




7. The compound of any one of claims 1 to 4, or a pharmaceutically


acceptable salt thereof, wherein  is selected from the group consisting of:



5

10





5


8. The compound of any one of claims 1 to 7, or a pharmaceutically acceptable salt thereof, wherein X^1 and X^2 are both CH.


9. The compound of any one of claims 1 to 7, or a pharmaceutically acceptable salt thereof, wherein X^1 is N and X^2 is CH.

10. The compound of any one of claims 1 to 7, or a pharmaceutically acceptable salt thereof, wherein X^1 and X^2 are both N.


11. The compound of any one of claims 1 to 7, or a pharmaceutically acceptable salt thereof, wherein X^1 is CH and X^2 is N.


12. The compound of any one of claims 1 to 11, or a pharmaceutically

acceptable salt thereof, wherein  is selected from the group consisting of:




5 13. The compound of any one of claims 1 to 11, or a pharmaceutically

acceptable salt thereof, wherein  is selected from the group consisting of:




14. The compound of claim 1 having the formula (II):




(II)

or a pharmaceutically acceptable salt thereof, wherein

X^1 and X^2 are both CH, or X^1 and X^2 are both N, or X^1 is N and X^2 is CH;

 is selected from the group consisting of




15

R^c , R^d , and R^e are each independently selected from the group consisting of H and methyl;

A is selected from the group consisting of azetidine, pyrrolidine, piperidine, and morpholine;


5 wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 2 R^k groups;


each R^k is independently selected from the group consisting of F, CN, methyl, ethyl and C_{1-2} haloalkyl;



10 R^2 is selected from the group consisting of H, Cl, and F.


15. The compound of claim 14, or a pharmaceutically acceptable salt thereof,

wherein  is selected from the group consisting of:




15

16. The compound of claim 14, or a pharmaceutically acceptable salt thereof, wherein the compound is selected from the group consisting of



5

17. A compound of formula (B):




10 or a pharmaceutically acceptable salt thereof, wherein

X^1 and X^2 are each independently selected from N and CH;

R^d and R^e are each independently selected from the group consisting of H and C_{1-3} alkyl; optionally R^d and R^e may be joined to form a cyclopropyl ring;

R^{k1} is selected from the group consisting of H, F, CN, OMe, and C_{1-3} alkyl;

15 R^{k2} is selected from the group consisting of H and methyl;




R^2 is selected from the group consisting of H, Cl, and F.

18. The compound of claim 17, or a pharmaceutically acceptable salt thereof, wherein

R^d and R^e are each independently selected from the group consisting of H and methyl; optionally R^d and R^e may be joined to form a cyclopropyl ring; and

5 R^{k1} is selected from the group consisting of H, F, CN, OMe, methyl and ethyl.

19. A compound of formula (C):




10 or a pharmaceutically acceptable salt thereof, wherein

X^1 and X^2 are both CH, or X^1 and X^2 are both N, or X^1 is N and X^2 is CH;


R^d and R^e are each independently selected from the group consisting of H and methyl;

R^k is selected from the group consisting of H, CN, methyl and ethyl;




R^2 is selected from the group consisting of H, Cl, and F.

20. The compound of claim 19, or a pharmaceutically acceptable salt thereof, wherein the compound is selected from the group consisting of




21. A compound of formula:




or a pharmaceutically acceptable salt thereof.

22. A compound of formula:




or a pharmaceutically acceptable salt thereof.

23. A compound of formula:



or a pharmaceutically acceptable salt thereof.

24. A crystalline form of the compound of formula:



5 wherein the crystalline form is characterized by a powder X-ray diffraction pattern comprising diffraction peaks at 2θ values of 5.65 ± 0.20 , 14.22 ± 0.20 , 15.16 ± 0.20 , and 19.31 ± 0.20 .

25. The crystalline form of claim 24 wherein the powder X-ray diffraction
10 pattern is further characterized by having additional diffraction peaks at 2θ values of 7.12 ± 0.20 , 10.02 ± 0.20 , 11.16 ± 0.20 , 17.06 ± 0.20 , and 24.43 ± 0.20 .


26. The crystalline form of claim 25 wherein the powder X-ray diffraction
15 pattern is further characterized by having two or more additional diffraction peaks at 2θ values selected from 13.10 ± 0.20 , 14.82 ± 0.20 , 16.55 ± 0.20 , 20.08 ± 0.20 , 21.08 ± 0.20 , 21.65 ± 0.20 , 22.51 ± 0.20 , 22.98 ± 0.20 , 25.02 ± 0.20 , 25.72 ± 0.20 , 26.80 ± 0.20 , 27.06 ± 0.20 , 28.31 ± 0.20 , 30.08 ± 0.20 , 30.31 ± 0.20 and 32.08 ± 0.20 .

27. The crystalline form of claim 24, wherein the crystalline form is
20 characterized by a powder X-ray diffraction pattern in which the peak positions are substantially in accordance with the peak positions of the pattern shown in Figure 1.

28. The crystalline form of claim 24 wherein the crystalline form is
characterized by a differential scanning calorimetry trace recorded at a heating rate of
25 $10\text{ }^{\circ}\text{C}$ per minute which shows a maximum in endothermic heat flow with a peak at $162.9\pm3\text{ }^{\circ}\text{C}$.

29. The crystalline form of claim 24, wherein the crystalline form is
characterized by a differential scanning calorimetry trace substantially in accordance with
30 that shown in Figure 2.

30. A crystalline form of the compound of formula:



wherein the crystalline form is characterized by a powder X-ray diffraction pattern comprising diffraction peaks at 2θ values of 9.67 ± 0.20 , 11.61 ± 0.20 , 17.61 ± 0.20 , 18.88 ± 0.20 , and 23.33 ± 0.20 .

31. The crystalline form of claim 30 wherein the powder X-ray diffraction pattern is further characterized by having additional diffraction peaks at 2θ values of 4.82 ± 0.20 , 15.69 ± 0.20 , and 16.19 ± 0.20 .

32. The crystalline form of claim 31 wherein the powder X-ray diffraction pattern is further characterized by having two or more additional diffraction peaks at 2θ values selected from 11.92 ± 0.20 , 12.98 ± 0.20 , 13.23 ± 0.20 , 16.45 ± 0.20 , 16.67 ± 0.20 , 19.39 ± 0.20 , 19.96 ± 0.20 , 20.14 ± 0.20 , 22.14 ± 0.20 , 23.84 ± 0.20 , 24.06 ± 0.20 , 24.29 ± 0.20 , 25.31 ± 0.20 , 25.63 ± 0.20 , 27.06 ± 0.20 , 27.31 ± 0.20 , 30.10 ± 0.20 , and 30.53 ± 0.20 .

33. The crystalline form of claim 30, wherein the crystalline form is characterized by a powder X-ray diffraction pattern in which the peak positions are substantially in accordance with the peak positions of the pattern shown in Figure 9.

34. The crystalline form of claim 30 wherein the crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10°C per minute which shows a maximum in endothermic heat flow with a peak at $201.3^\circ\text{C} \pm 2^\circ\text{C}$.

35. The crystalline form of claim 30 wherein the crystalline form is characterized by a differential scanning calorimetry trace recorded at a heating rate of 10°C per minute which shows a maximum in endothermic heat flow at a temperature between 198°C and 204°C .

36. The crystalline form of claim 30, wherein the crystalline form is characterized by a differential scanning calorimetry trace substantially in accordance with that shown in Figure 10.

5 37. A pharmaceutical composition comprising a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, or a crystalline form of anyone of claims 24 to 36, and a pharmaceutically-acceptable carrier.

10 38. The pharmaceutical composition of claim 37 further comprising one or more other therapeutic agents useful for treating a gastrointestinal inflammatory disease.

39. A compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, or a crystalline form of anyone of claims 24 to 36, for use in the treatment of a gastrointestinal inflammatory disease in a mammal.

15 40. The compound of claim 39, or a pharmaceutically acceptable salt thereof, or the crystalline form of claim 39, wherein the gastrointestinal inflammatory disease is selected from the group consisting of immune checkpoint inhibitor induced colitis, CTLA-4 inhibitor-induced colitis, graft versus host disease-related colitis, celiac disease, collagenous colitis, lymphocytic colitis, Behcet's disease, ileitis, eosinophilic esophagitis, and infectious colitis.

20 41. The compound of claim 39, or a pharmaceutically acceptable salt thereof, or the crystalline form of claim 39, wherein the gastrointestinal inflammatory disease is ulcerative colitis.

42. The compound of claim 39, or a pharmaceutically acceptable salt thereof, or the crystalline form of claim 39, wherein the gastrointestinal inflammatory disease is Crohn's disease.

30 43. The compound of claim 39, or a pharmaceutically acceptable salt thereof, or the crystalline form of claim 39, for use in combination with one or more other therapeutic agents useful for treating a gastrointestinal inflammatory disease.

44. Use of a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, or a crystalline form of anyone of claims 24 to 36, for the manufacture of a medicament for the treatment of a gastrointestinal inflammatory disease in a mammal.

5

45. The use of claim 44 wherein the gastrointestinal inflammatory disease is selected from the group consisting of immune checkpoint inhibitor induced colitis, CTLA-4 inhibitor-induced colitis, graft versus host disease-related colitis, celiac disease, collagenous colitis, lymphocytic colitis, Behcet's disease, ileitis, eosinophilic esophagitis, and infectious colitis.

10

46. The use of claim 44 wherein the gastrointestinal inflammatory disease is ulcerative colitis.

15

47. The use of claim 44 wherein the gastrointestinal inflammatory disease is Crohn's disease.

48. A compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, for use in the treatment of an inflammatory skin disease in a mammal.

20

49. A compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, for use in the treatment of cutaneous T-cell lymphoma.

25

50. Use of a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, or a crystalline form of anyone of claims 24 to 36, for the manufacture of a medicament for the treatment of an inflammatory skin disease in a mammal.

30

51. Use of a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, or a crystalline form of anyone of claims 24 to 36, for the manufacture of a medicament for the treatment of cutaneous T-cell lymphoma.

52. A method of treating a gastrointestinal inflammatory disease in a mammal, the method comprising administering to the mammal a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, or a crystalline form of any one of claims 24 to 36, and a pharmaceutically-acceptable carrier.

5

53. The method of claim 52, wherein the method further comprises administering one or more other therapeutic agents useful for treating a gastrointestinal inflammatory disease.

10

54. The method of claim 52, wherein the gastrointestinal inflammatory disease is selected from the group consisting of immune checkpoint inhibitor induced colitis, CTLA-4 inhibitor-induced colitis, graft versus host disease-related colitis, celiac disease, collagenous colitis, lymphocytic colitis, Behcet's disease, ileitis, eosinophilic esophagitis, and infectious colitis.

15

55. The method of claim 52, wherein the gastrointestinal inflammatory disease is ulcerative colitis.

56. The method of claim 52, wherein the gastrointestinal inflammatory disease is Crohn's disease.


57. A method of treating an inflammatory skin disease in a mammal, the method comprising applying a pharmaceutical composition comprising a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, to the skin of the mammal.

25

58. A method of treating cutaneous T-cell lymphoma in a mammal, the method comprising applying a pharmaceutical composition comprising a compound of any one of claims 1 to 23, or a pharmaceutically acceptable salt thereof, to the skin of the mammal.

30

59. A process for preparing a compound of formula (I) or a pharmaceutically acceptable salt thereof,




(I)

5 wherein


X^1 and X^2 are each independently selected from N and CH;

X^3 is selected from the group consisting of N, CH, C-CH₃, C-CF₃, C-CHF₂, C-CH₂-O-CH₃, C-SMe, C-NMe₂, C-NH-CH₃, C-Cl, C-CN, and C-OMe;



is selected from the group consisting of

10



and

;

R^a , R^b , R^c , and R^f are each independently selected from the group consisting of H and C₁₋₃ alkyl;

15

R^d , R^e , R^g , R^h , R^i , R^j , R^l , R^m , R^n and R^o are each independently selected from the group consisting of H and C₁₋₃ alkyl wherein the C₁₋₃ alkyl group may be optionally substituted with 1 to 3 halogens;

Optionally R^d and R^e may be joined to form a cyclopropyl ring;


A is selected from the group consisting of

- (a) a 4 to 10 membered monocyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S, S(O)₂ and O, and
- (b) a 6 to 10 membered multicyclic heterocyclic group containing one nitrogen atom and optionally containing one additional heteroatom selected from N, S and O,

wherein L is linked to a carbon atom in A and A is optionally substituted with 1 to 3 R^k groups;

each R^k is independently selected from the group consisting of F, CN, C₁₋₃ alkoxy, cyclopropyl, and C₁₋₃ alkyl, wherein the C₁₋₃ alkyl group may be optionally substituted with OH, OMe or 1 to 3 halogens;

R^1 is selected from the group consisting of



- 15 wherein R^p and R^q are each independently selected from the group consisting of H, C₃₋₅ cycloalkyl and C₁₋₆ alkyl, wherein the C₁₋₆ alkyl group may be optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₃ alkoxy and -S-C₁₋₃ alkyl.

or R^p and R^q form a 4 to 6 membered monocyclic heterocyclic group containing
20 one nitrogen atom and optionally containing one additional heteroatom selected from N,
S, and O, wherein the 4 to 6 membered monocyclic heterocyclic group is optionally
substituted with 1 to 3 substituents independently selected from the group consisting of
C₁₋₆ alkyl, C₁₋₃ alkoxy, -S-C₁₋₃ alkyl and -C₁₋₃ alkyl-C₁₋₃ alkoxy;


R² is selected from the group consisting of H, Cl, OMe, Me and F;

25 R^3 is selected from the group consisting of H and F;

R^4 is selected from the group consisting of H and F; and

R^5 is selected from the group consisting of H, Me and F.

the method comprising reacting a compound of formula (III):



with

- (i) Cl-R¹, or
 5 (ii) HO-R¹

and optionally forming a pharmaceutically-acceptable salt

to provide a compound of formula (I), or a pharmaceutically acceptable salt thereof.

1/9




FIG. 1




FIG. 2

2/9




FIG. 3




FIG. 4

3/9




FIG. 5




FIG. 6

4/9




FIG. 7




FIG. 8

5/9




FIG. 9




FIG. 10

6/9




FIG. 11




FIG. 12

7/9




FIG. 13




FIG. 14

8/9




FIG. 15




FIG. 16

9/9





FIG. 17



(I)