



(51) International Patent Classification:

C07D 209/34 (2006.01) A61K 31/352 (2006.01)
 C07D 231/56 (2006.01) A61K 31/404 (2006.01)
 C07D 249/18 (2006.01) A61K 31/416 (2006.01)
 C07D 263/54 (2006.01) A61K 31/4192 (2006.01)
 C07D 277/62 (2006.01) A61K 31/428 (2006.01)
 C07D 403/12 (2006.01) A61P 25/00 (2006.01)
 C07D 407/12 (2006.01)

(21) International Application Number:

PCT/GB2023/050493

(22) International Filing Date:

02 March 2023 (02.03.2023)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2202921.9 02 March 2022 (02.03.2022) GB

(71) Applicant: NRG THERAPEUTICS LTD [GB/GB];

Stevenage Bioscience Catalyst, Gunnels Wood Road,
 Stevenage, Herts SG1 2FX (GB).

(72) Inventors: MILLER, Neil; NRG Therapeutics Ltd, Steven-

nage Bioscience Catalyst, Gunnels Wood Road, Stevenage,
 Herts, SG1 2FX (GB). RUTTER, Richard; NRG Thera-
 peutics Ltd, Stevenage Bioscience Catalyst, Gunnels Wood
 Road, Stevenage, Herts, SG1 2FX (GB). KULAGOWSKI,
 Jan; Parkinson's UK, 215 Vauxhall Bridge Road, Lon-
 don SW1V 1EJ (GB). MORPHY, Richard; Parkinson's
 UK, 215 Vauxhall Bridge Road, London SW1V 1EJ (GB).
 LADDUWAHETTY, Tammy; Pharmaron UK Limited,
 The Old Glassworks, Nettlefold Road, Cardiff CF24 5JQ
 (GB). MACLEAN, John; Pharmaron UK Limited, The
 Old Glassworks, Nettlefold Road, Cardiff, CF24 5JQ (GB).
 MOROGLU, Mustafa; Pharmaron UK Limited, The Old
 Glassworks, Nettlefold Road, Cardiff CD24 5JQ (GB).
 TALBOT, Eric; Pharmaron UK Limited, The Old Glass-
 works, Nettlefold Road, Cardiff, CF24 5JQ (GB). ROW-
 LEY, Michael; Parkinson's UK, 215 Vauxhall Bridge
 Road, London SW1V 1EJ (GB). THOMSON, Christo-
 pher G.; Pharmaron UK Limited, The Old Glassworks, Net-
 tlefold Road, Cardiff, CF24 5JQ (GB).

(74) Agent: TEUTEN, Andrew et al.; Sagittarius IP, Mar-

low International, Parkway, Marlow Buckinghamshire SL7
 1YL (GB).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
 CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM,
 DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
 HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE,
 KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU,
 LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG,
 NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,
 RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH,
 TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS,
 ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every


kind of regional protection available): ARIPO (BW, CV,
 GH, GM, KE, LR, LS, MW, MZ, NA, RW, SC, SD, SL, ST,
 SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ,
 RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ,
 DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT,
 LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE,
 SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
 GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
 claims and to be republished in the event of receipt of
 amendments (Rule 48.2(h))

(54) Title: ACRYLAMIDE COMPOUNDS

(57) Abstract: The invention relates to compounds of formula (I): (I) and related aspects.



Field of the invention

The invention relates to novel compounds which are inhibitors of the mitochondrial permeability transition pore (mPTP). The invention also *inter alia* relates to such compounds for use as medicaments, in particular, for the treatment or prevention of degenerative, neurodegenerative or mitochondrial diseases or other diseases or disorders in which inhibition of mPTP provides a therapeutic or prophylactic effect.

Background to the invention

The mitochondria permeability transition pore (mPTP) is a high conductance channel residing on the inner mitochondrial membrane that is activated under certain conditions of cellular stress, in particular excessive Ca^{2+} loading and oxidative stress. It is permeable to solutes with molecular mass <1.5 kDa, is voltage and Ca^{2+} dependent and exhibits a characteristic large conductance. Once activated, oxidative phosphorylation is uncoupled resulting in the loss of the mitochondria membrane potential and disrupted mitochondria metabolism. In addition, solutes enter the mitochondrial matrix resulting in swelling, eventual rupture of the outer membrane with consequent release of apoptotic factors as well as sequestered Ca^{2+} , leading to eventual cell death via apoptosis or necrosis depending on the type and physiology of the cell. As such it has been implicated as a key pathological event in multiple degenerative and metabolic diseases.

Under normal physiological conditions mitochondria play a key role in regulating cellular Ca^{2+} homeostasis. Ca^{2+} entering the cell via cell surface channels, a common mechanism of cell signalling, is rapidly sequestered by mitochondria, preventing excessive and toxic Ca^{2+} accumulation in the cell cytoplasm. In cell types such as neurons, skeletal muscle myofibers and cardiomyocytes which undergo high levels of Ca^{2+} flux, this Ca^{2+} 'buffering' effect of mitochondria is critical to maintain cell health. However, there is a limit to the capacity of mitochondria to sequester Ca^{2+} and if intramitochondrial Ca^{2+} levels reach a certain threshold the Ca^{2+} sensitive mPTP is activated, resulting in collapse of the mitochondria and initiation of cell death. Activation of the mPTP in degenerative diseases may occur in a variety of ways depending on the disease, for example: 1) excessive Ca^{2+} entry into cells and overload of the mitochondria with Ca^{2+} 2) dysfunctional mitochondrial Ca^{2+} efflux mechanisms, in particular decreased activity of the Ca^{2+} efflux transporter NCLX resulting in Ca^{2+} overload 3) overactivity or upregulation of the Ca^{2+} uptake mechanisms in mitochondria 4) oxidative stress 5) sensitization of the mPTP due to compromised mitochondrial function i.e. mPTP activation at lower intramitochondrial concentrations of Ca^{2+} 6) excessive transfer of Ca^{2+} from the

endoplasmic reticulum into the mitochondria at contact points between the two organelles known as mitochondria-associated-membranes.

While the properties and function of the mPTP can be studied in simple *in vitro* assays in isolated mitochondria, the molecular identity of the mPTP is not known. Multiple proteins have been proposed to comprise the pore forming complex, including the ATP synthase and the adenine nucleotide transporter (ANT) family of proteins but no single protein is widely accepted as being responsible for formation of the pore. However, the peptidyl prolyl cis-trans isomerase F (Ppif), also known as cyclophilin D, is well accepted to be a key regulator of the pore, though not forming a transmembrane channel in its own right. Genetic or pharmacological inhibition of Ppif significantly decreases the sensitivity of pore opening in response to Ca^{2+} loading and other mPTP activators. Genetic ablation or pharmacological inhibition of Ppif has therefore been utilised to evaluate involvement of the mPTP in pathological pathways in cell and animal disease models. In this way, inhibition of the mPTP has been shown to be protective in numerous models of disease, in particular those where Ca^{2+} dysregulation and oxidative stress are known to contribute to cellular degeneration. Notably, genetic knockout of Ppif was shown to be protective in various preclinical *in vivo* transgenic models of neurodegenerative disease including Alzheimer's disease, Parkinson's disease and motor neuron disease, demonstrating the therapeutic potential of mPTP inhibition. In each of these diseases, genetic mutations in particular proteins that cause inherited forms of disease (i.e. amyloid precursor protein, alpha-synuclein and superoxide dismutase 1 respectively), and are expressed in the mouse models, have been shown to cause either Ca^{2+} overload of the mitochondria or sensitization of the mPTP. Recent evidence suggests this may occur through a common mechanism in Alzheimer's, Parkinson's and Friedreich's ataxia. In each case, it has been reported that in cells expressing the mutated disease associated proteins (amyloid precursor protein, PINK1 and frataxin respectively), the activity or expression of the mitochondrial Ca^{2+} efflux transporter, NCLX, is decreased, resulting in Ca^{2+} overload of the mitochondria. In the case of Parkinson's disease, the pathological aggregated form of the protein alpha-synuclein, a common misfolded protein in sporadic and inherited cases of Parkinson's disease, has also been shown to sensitise and activate the mPTP.

Genetic ablation of Ppif has been shown to be beneficial in numerous other preclinical models of degenerative disease, therefore demonstrating the potential of mPTP inhibitors in Duchenne and congenital forms of muscular dystrophy, ischemia-reperfusion injury, bone repair, pancreatitis and *inter alia* other associated disorders.

In addition to the demonstrated benefit of Ppif inhibition in preclinical models, mPTP function has been shown to be dysregulated in multiple other disease indications. In particular, in a number of diseases the threshold for mPTP activation in response to Ca^{2+} loading appears to be sensitised suggesting that mPTP activation may occur aberrantly under physiological conditions and drive tissue degeneration. For example, in muscle mitochondria from elderly human muscle biopsies, the threshold for mPTP activation is reduced compared to healthy control. In these diseases, this sensitization of mPTP activity underlies additional rationale for the therapeutic potential of mPTP inhibitors.

mPTP inhibitors may also have therapeutic potential in other diseases where mitochondrial dysfunction, oxidative stress, inflammatory stress or Ca^{2+} dysregulation occur during disease pathogenesis.

The discovery and development of inhibitors of the mPTP has largely been focussed on identification of Ppif inhibitors. Cyclosporin A (CsA), originally identified as an immunosuppressant by virtue of its inhibitory activity at calcineurin, was also found to inhibit Ppif as well as other members of the peptidyl prolyl cis-trans isomerase (Ppi) enzyme family. Several cyclosporin A derivatives e.g. Debio-25, NIM811 were subsequently developed that retained broad activity against the Ppi enzyme family without inhibiting calcineurin, however none of these progressed to market. To date, no potent brain penetrant selective Ppif inhibitors have been reported. Other more recent approaches to discover mPTP inhibitors, have utilised phenotypic screens in isolated mitochondria. These have successfully identified potent small molecule mPTP inhibitors with a Ppif-independent mode of action.

Yu et al., (2020, Cell, 183, 1-14) relates to the link between mPTP activation and the mechanism of TDP-43 proteinopathy such as TAR DNA-binding protein 43 (TDP-43) associated neurodegeneration. Cytoplasmic neuronal accumulation of the normally nuclear protein TDP-43 is a disease hallmark for almost all cases of ALS and 40-50% of Frontotemporal Lobar Degeneration (FTLD), with some familial cases caused by mutant forms of the protein. Both diseases are associated with a neuroinflammatory cytokine profile related to upregulation of NF- κ B and type I IFN pathways, directly suggesting a role for TDP-43 in neuroinflammation. Mutant or overexpressed WT TDP-43 in neurons mis-localises to the mitochondria and induces the release of mitochondrial DNA (mtDNA) into the cytoplasm. This mtDNA then activates the immune sensor cGAS-STING triggering the induction of innate immune genes such as IL-6, TNF α , and interferon β . Inhibition of the mPTP with cyclosporin A or via CypD knockout, prevents the TDP-43 induced release of mtDNA and subsequent induction of innate immune response genes. Furthermore, inhibition of cGAS-STING extends the survival of mutant mice

expressing a mutant TDP-43. The data implicates mPTP activation in mediating the toxic effects of TDP-43 in ALS and other disease where either mutations in the TDP-43 gene causes disease or where TDP-43 proteinopathy is observed.

- 5 Jang et al., (2021 American Journal of Physiology: Renal physiology, doi: 10.1152/ajprenal.00171.2021. Epub ahead of print. PMID: 34396791.) highlighted the potential therapeutic benefit of mPTP inhibition (via CypD knockout) in a mouse model a kidney fibrosis. Unilateral ureteral obstruction was used to induce kidney fibrosis in WT and CypD KO mice. Inflammation, proximal tubule atrophy and markers of fibrosis were reduced in the CypD KO
10 mice versus WT. Measures of fibrosis included collagen deposition, α -SMA and TGF β expression, and interstitial cell proliferation. This highlights the potential role of the mPTP in cell injury/death-mediated tissue remodelling and fibrogenesis. mPTP inhibitors may therefore be beneficial in diseases where fibrosis is a key pathological mechanism, e.g, chronic kidney disease, idiopathic pulmonary fibrosis, non-alcoholic steatohepatitis, primary biliary cholangitis
15 and systemic sclerosis.

CYP2D6 is one of the major members of the human drug metabolising cytochrome P450 enzyme system. It is involved in the hepatic metabolism of a significant proportion of clinically used drugs. Inhibition of CYP2D6 can drive drug-drug interactions with co-prescribed
20 medications metabolised by the same enzyme, which result in increased plasma concentrations potentially to levels which may cause adverse effects. CYP2D6 is predominately expressed in the liver but also, to a lesser degree, in the central nervous system (CNS). In the CNS it is involved in the synthesis of different neurotransmitters such as dopamine. Consequently, inhibition of CYP2D6 particularly at the level of the central nervous system can potentially have
25 detrimental effects via impairment of pathways such as the production of dopamine. In Parkinson's disease, which is characterized by the loss of dopaminergic neurons in the Substantia Nigra, further depletion of dopamine levels through inhibition of CYP2D6 may not be tolerated.


30 When dosed orally, the oral bioavailability and systemic exposure of a drug are largely determined by the degree of absorption from the gastrointestinal tract and the extent of first-pass metabolism in the liver. Properties such as high solubility (as measured in phosphate-buffered saline (PBS) or in the more biologically relevant Fasted State Simulated Intestinal Fluid (FaSSIF)) and high metabolic stability (as measured *in vitro* in either isolated liver microsomes
35 or in hepatocytes from rat and human) are therefore predictive of improved oral bioavailability and/or systemic exposure in patients.

WO2010/049768 relates to acrylamido derivatives and their use as therapeutic agents, particularly for the prevention and/or treatment of diseases associated with the activity of the mPTP (see also Plyte et al. J. Med Chem. 2014, 57, 5333-47). Chen et al. (Assay and Drug Development Technologies, 2018, 16, 445-455) relates to phenotypic screening for mPTP modulators using platelets, and discloses further acrylamido derivatives. CA2884607A1 relates to acrylamido and maleimide compounds which are said to be useful in the treatment of mitochondrial diseases.

There remains a need to find further compounds which are inhibitors of mPTP, and in particular those that combine inhibition of mPTP with low CYP2D6 inhibition. Such compounds may also display other desirable pharmacological properties, such as improved oral bioavailability and/or improved systemic exposure.

Summary of the Invention

In a first aspect the invention provides a compound of formula (I):




wherein:

R_{1a} is H or methyl;

R_{1b} is H or F;

A is group (Aa), (Ab), (Ac), (Ad) or (Ae):

wherein group (Aa) is:



wherein:

R₂ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, C₁₋₄alkoxy, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl), C₁₋₄alkylene(CO₂H); OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2ab}R_{2ac}), OC₁₋₄alkylene(NR_{2ab}R_{2ac}) or N(R_{2aa})C₁₋₄alkylene(NR_{2ab}R_{2ac});

wherein said aryl, heterocycloalkyl and cycloalkyl are optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{2a}R_{2b}, SO₂R_{2c} and NHSO₂R_{2c};

R_{2a} is selected from H and C₁₋₄alkyl;

5 R_{2b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{2c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{2aa} is independently selected from H and C₁₋₄alkyl;

10 each R_{2ab} is independently selected from H and C₁₋₄alkyl;

each R_{2ac} is independently selected from H and C₁₋₄alkyl; or

R_{2ab} and R_{2ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R_{2x} is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

15 each R₃ is independently halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

m is 0, 1, 2 or 3;

or;

20 R_{2x} is C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkylene(CO₂H), OC₁₋

4alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2xaa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2xab}R_{2xac}), OC₁₋₄alkylene(NR_{2xab}R_{2xac}) or N(R_{2xaa})C₁₋₄alkylene(NR_{2xab}R_{2xac});

wherein said heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each

25 independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{2xa}R_{2xb}, SO₂R_{2xc} and NHSO₂R_{2xc};

R_{2xa} is selected from H and C₁₋₄alkyl;

R_{2xb} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

30 R_{2xc} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{2xaa} is independently selected from H and C₁₋₄alkyl;

each R_{2xab} is independently selected from H and C₁₋₄alkyl;

each R_{2xac} is independently selected from H and C₁₋₄alkyl; or


35 R_{2xab} and R_{2xac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R_2 is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyleneOC $_{1-4}$ alkyl, C_{1-4} alkyleneOC $_{3-6}$ cycloalkyl or C_{1-4} alkyleneO(4-10 membered heterocycloalkyl);

each R_3 is independently halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy;

m is 0, 1, 2 or 3;

5 wherein group (Ab) is:



wherein:

R_4 is H, C_{1-4} alkyl or C_{1-4} alkylene(aryl); wherein said aryl is optionally substituted by 1, 2 or 3 substituents each independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{4a}R_{4b}$, SO_2R_{4c} and $NHSO_2R_{4c}$;

R_{4a} is selected from H and C_{1-4} alkyl;

R_{4b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;


R_{4c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_5 is H or C_{1-4} alkyl;

each R_6 is independently C_{1-4} alkyl or halo;

n is 0, 1, 2 or 3;

wherein group (Ac) is:




wherein:

R_7 is C_{1-4} alkyl, C_{1-4} alkylene(OH) or C_{1-4} alkyleneOC $_{1-4}$ alkyl;

o is 1 or 2;

wherein group (Ad) is:



wherein:

X is a bond, O or CH_2 ;

each R_8 is independently halo, C_{1-4} alkyl, C_{1-4} alkoxy, OC_{1-4} haloalkyl, OC_{1-4} alkylene(C_{3-6} cycloalkyl), OC_{1-4} alkylene(4-10 membered heterocycloalkyl) or OH; wherein said heterocycloalkyl and cycloalkyl are optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{8a}R_{8b}$, SO_2R_{8c} and $NHSO_2R_{8c}$;

R_{8a} is selected from H and C_{1-4} alkyl;

R_{8b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;


R_{8c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_9 is independently halo or C_{1-4} alkyl;

p is 0, 1 or 2;

q is 0, 1, 2, 3 or 4;

wherein group (Ae) is:



wherein:

R_{17} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkylene(aryl), C_{1-4} alkylene(OH), C_{1-4} alkylene(C_{3-6} cycloalkyl), C_{1-4} alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, C_{1-4} alkoxy, OC_{1-4} alkylene(aryl), C_{1-4} alkylene OC_{1-4} alkyl, C_{1-4} alkylene OC_{3-6} cycloalkyl, C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkyleneO(aryl), C_{3-6} alkynyl, C_{1-4} alkenylO(C_{3-6} alkynyl), C_{1-4} alkylene(CO_2H), OC_{1-4} alkylene(CO_2H), C_{1-4} alkylene OC_{1-4} alkylene(CO_2H), $N(R_{17aa})C_{1-4}$ alkylene(CO_2H), C_{1-4} alkylene($NR_{17ab}R_{17ac}$), OC_{1-4} alkylene($NR_{17ab}R_{17ac}$) or $N(R_{17aa})C_{1-4}$ alkylene($NR_{17ab}R_{17ac}$); wherein said aryl, cycloalkyl or heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{17a}R_{17b}$, SO_2R_{17c} , $NHSO_2R_{17c}$;

R_{17a} is selected from H and C_{1-4} alkyl;

R_{17b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{17c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{17aa} is independently selected from H and C_{1-4} alkyl;

each R_{17ab} is independently selected from H and C_{1-4} alkyl;

each R_{17ac} is independently selected from H and C₁₋₄alkyl; or

R_{17ab} and R_{17ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R₁₈ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₂₀ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

R₂₁ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

or;

R₁₇ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₁₈ is C₁₋₄haloalkyl, C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl), C₁₋₄alkylene(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{18aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{18ab}R_{18ac}), OC₁₋₄alkylene(NR_{18ab}R_{18ac}) or N(R_{18aa})C₁₋₄alkylene(NR_{18ab}R_{18ac}); wherein said aryl, cycloalkyl or heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{18a}R_{18b}, SO₂R_{18c}, NHSO₂R_{18c};

R_{18a} is selected from H and C₁₋₄alkyl;

R_{18b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{18c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{18aa} is independently selected from H and C₁₋₄alkyl;

each R_{18ab} is independently selected from H and C₁₋₄alkyl;

each R_{18ac} is independently selected from H and C₁₋₄alkyl; or

R_{18ab} and R_{18ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R₂₀ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

R₂₁ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

or;

R₁₇ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₁₈ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

5 R₂₀ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

R₂₁ is C₁₋₄haloalkyl, C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl),

10 C₁₋₄alkylene(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{21aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{21ab}R_{21ac}), OC₁₋₄alkylene(NR_{21ab}R_{21ac}) or N(R_{21aa})C₁₋₄alkylene(NR_{21ab}R_{21ac}); wherein said aryl, cycloalkyl or heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{21a}R_{21b}, SO₂R_{21c}, NHSO₂R_{21c};

15 R_{21a} is selected from H and C₁₋₄alkyl;

R_{21b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{21c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

20 each R_{21aa} is independently selected from H and C₁₋₄alkyl;

each R_{21ab} is independently selected from H and C₁₋₄alkyl;


each R_{21ac} is independently selected from H and C₁₋₄alkyl; or

R_{21ab} and R_{21ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

25

wherein B is group (Ba), (Bb) or (Bc):

wherein group (Ba) is:



wherein:

30 Y is C(R₁₁)(R₁₂), N(R₁₃), O or S;

each R₁₀ is independently halo or C₁₋₄alkyl;

r is 0, 1, 2 or 3;

R₁₁ is H or C₁₋₄alkyl;

R_{12} is H or C_{1-4} alkyl; or R_{11} and R_{12} together with the carbon atom to which they are attached form a C_{3-6} cycloalkyl;

R_{13} is H, C_{1-4} alkyl or C_{3-6} cycloalkyl;


wherein said cycloalkyl is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{13a}R_{13b}$, SO_2R_{13c} and $NHSO_2R_{13c}$;

R_{13a} is selected from H and C_{1-4} alkyl;

R_{13b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{13c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

wherein group (Bb) is:




wherein:

each R_{14} is independently halo or C_{1-4} alkyl;

s is 0, 1, 2 or 3;

wherein group (Bc) is:



wherein:

R_{15} is C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-4} haloalkyl, halo or CN; wherein said cycloalkyl is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{15a}R_{15b}$, SO_2R_{15c} and $NHSO_2R_{15c}$;

R_{15a} is selected from H and C_{1-4} alkyl;

R_{15b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{15c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{16} is H, halo or C_{1-4} alkyl; and

D, E and F are each independently C(R₁₆); or one of D, E and F is N, and the two remaining D, E and F groups are independently C(R₁₆);
or a pharmaceutically acceptable salt and/or solvate thereof.

5 In one embodiment, a compound of formula (I) is provided in the form of a pharmaceutically acceptable salt and solvate. In one embodiment, a compound of formula (I) is provided in the form of a pharmaceutically acceptable salt. In one embodiment, a compound of formula (I) is provided in the form of a pharmaceutically acceptable solvate. In one embodiment, a compound of formula (I) is provided.

10

The invention further provides pharmaceutical compositions comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, and a pharmaceutically acceptable carrier or excipient.

15 The invention also provides a compound of formula (I), or pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect.

20 The invention also provides a method of preventing or treating a disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect in a subject.

Suitably the disease or disorder is selected from degenerative or neurodegenerative diseases, disorders of the central nervous system, ischemia and re-perfusion injury, metabolic diseases, inflammatory or autoimmune diseases, diseases of aging and renal diseases.

30

The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a mitochondrial disease.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a mitochondrial disease.

35

The invention also provides a method of preventing or treating a mitochondrial disease in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof.

- 5 The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration. The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis
10 of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration.

- The invention also provides a method of treating or preventing a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, which comprises
15 administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof.

- The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a disease or disorder
20 associated with fibrosis.

- The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a disease or disorder associated with fibrosis.
25

- The invention also provides a method of treating or preventing a disease or disorder associated with fibrosis, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof.

- 30 Also provided are novel intermediate compounds of formula (II) to (XXXVII) and salts thereof such as compounds of formula (II), (IV) and (VI). Such compounds are of use in the preparation of compounds of formula (I).

Detailed description of the Invention

- 35 The term "alkyl" as used herein, such as in C₁₋₄alkyl, whether alone or forming part of a larger group, is a straight or branched fully saturated hydrocarbon chain containing the specified number of carbon atoms. Examples of C₁₋₄alkyl groups include methyl, ethyl, *n*-propyl, *iso*-

propyl, *n*-butyl, *iso*-butyl, *tert*-butyl and *sec*-butyl. Reference to “propyl” includes *n*-propyl and *iso*-propyl. Reference to “butyl” includes *n*-butyl, *iso*-butyl, *tert*-butyl and *sec*-butyl.

Me means methyl. Et means ethyl. Pr means propyl. Bu means butyl.

5

The term “alkylene” as used herein, such as C₁₋₄alkylene, whether alone or forming part of a larger group e.g. C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), OC₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), OC₁₋₄alkylene(4-10 membered heterocycloalkyl), C₁₋₄alkoxy, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl) or C₁₋₄alkyleneO(C₃₋₆alkynyl), e.g. C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), C₁₋₄alkoxy, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl) or C₁₋₄alkyleneO(C₃₋₆alkynyl), is a bifunctional straight or branched fully saturated hydrocarbon group containing the specified number of carbon atoms. Examples of C₁₋₄alkylene groups include methylene (i.e. -CH₂-), ethylene (i.e. -CH₂CH₂-) *n*-propylene (i.e. (-CH₂)₃-) and *n*-butylene (i.e. (-CH₂)₄-). A branched example of a C₁₋₄alkylene group is *i*-propylene (i.e. -CH(Me)CH₂-).

10

15

20

The term C₁₋₄alkylene(OH) means an C₁₋₄alkyl group substituted by OH, such as CH₂OH.

The term “alkoxy” as used herein, such as in C₁₋₄alkoxy, refers to an alkyl group (e.g. a C₁₋₄alkyl group) as defined above, singularly bonded to an oxygen atom. Examples of C₁₋₄alkoxy groups include methoxy, ethoxy, 1-propoxy, 2-propoxy, 1-butoxy, 2-butoxy and 3-butoxy, especially methoxy.

25

The term ‘halo’ or ‘halogen’ as used herein, refers to fluorine, chlorine, bromine or iodine. Particular examples of halo are bromine, fluorine and chlorine, especially fluorine.

30

The term “haloalkyl” as used herein, such as in C₁₋₄haloalkyl, whether alone or forming part of a larger group such as OC₁₋₄haloalkyl, is a straight or branched alkyl group containing the specified number of carbon atoms, substituted by one or more halo atoms, for example fluoromethyl (CH₂F), di-fluoromethyl (CHF₂), tri-fluoromethyl (CF₃), 1-fluoroethyl (CH₂FCH₂) and 2-fluoroethyl (CH₂CH₂F).

35

The term “cycloalkyl” as used herein, such as in C₃₋₆cycloalkyl, whether alone or forming part of a larger group such as C₁₋₄alkylene(C₃₋₆cycloalkyl) or C₁₋₄alkyleneOC₃₋₆cycloalkyl is a fully

saturated hydrocarbon ring containing the specified number of carbon atoms. Examples of C₃₋₆cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, in particular cyclopropyl. Optionally, the cycloalkyl may be substituted as defined herein.

5 The term “heterocycloalkyl” as used herein, such as in 4-10 membered heterocycloalkyl (e.g. 4-7 membered heterocycloalkyl), whether alone or forming part of a larger group such as C₁₋₄alkylene(4-10 membered heterocycloalkyl) and C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), is a fully saturated hydrocarbon ring containing the specified number of carbon atoms, wherein at least one of the carbon atoms is replaced by a heteroatom such as N,
10 S or O. Optionally, the heterocycloalkyl may be substituted as defined herein. A heterocycloalkyl group may be monocyclic. Alternatively, a heterocycloalkyl group may be polycyclic such as a fused bicyclic or a bridged bicyclic ring system. In some examples a heterocycloalkyl group may incorporate a fused ring. In some examples a heterocycloalkyl group may incorporate a bridged ring. In some examples a bicyclic heterocycloalkyl group is spirocyclic i.e. a bicyclic cycloalkyl
15 group wherein the two rings are connected through just one atom. The rings can be different or identical.

Examples of 4-10 membered heterocycloalkyl groups include those comprising one heteroatom such as containing one heteroatom (e.g. nitrogen) or containing two or more heteroatoms (such
20 as two heteroatoms e.g. two nitrogen atoms or one nitrogen atom and one oxygen atom).

Examples of 4-10 membered heterocycloalkyl groups containing one nitrogen atom include azetidiny, pyrrolidiny, piperidiny and azepanyl. Examples of 4-10 membered heterocycloalkyl groups containing two nitrogen atoms include diazetidiny, imidazolidiny, pyrazolidiny, diazinanyl, and diazepanyl.

25 Other examples of 4-10 membered heterocycloalkyl groups include oxetanyl, thietanyl, dioxetanyl, dithietanyl, tetrahydrofuranyl, tetrahydrothiophenyl, oxazolidiny, isoxazolidiny, thiazolidiny, isothiazolidiny, tetrahydropyranyl, thianyl, morpholiny, thiomorpholiny, dioxanyl, dithianyl, triazinanyl, trioxanyl, trithianyl, oxepanyl, and thiepanyl.

30 Examples of 4-10 membered heterocycloalkyl groups incorporating a bridged ring are 1,4-diazabicyclo[3.2.1]octane, or 1-azabicyclo[3.2.1]octane.

An example of bicyclic spirocyclic heterocycloalkyl groups is 2,6-diazaspiro[3.3]heptane.

35

The term "aryl" as used herein, whether alone or forming part of a larger group e.g. C₁₋₄alkylene(aryl), OC₁₋₄alkylene(aryl) or C₁₋₄alkyleneO(aryl), refers to a phenyl ring. Optionally, the aryl may be substituted as defined herein.

- 5 The term "alkynyl" as used herein, such as in C₃₋₆alkynyl, whether alone or forming part of a larger group such as C₁₋₄alkyleneO(C₃₋₆alkynyl), is a straight or branched divalent hydrocarbon chain with a least one carbon-carbon triple bond. Examples of C₃₋₆alkynyl include ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, pentynyl and hexynyl.
- 10 An example of C₁₋₄alkylene(aryl) is CH₂Ph where Ph means phenyl. Examples of C₁₋₄alkyleneOC₁₋₄alkyl include CH₂OMe, CH₂OEt and CH₂OPr. Examples of C₁₋₄alkyleneOC₃₋₆cycloalkyl include CH₂O-C₃cycloalkyl and CH₂O-C₄cycloalkyl, for example CH₂O-cyclopropyl or CH₂O-cyclobutyl. Examples of C₁₋₄alkylene(4-10 membered heterocycloalkyl) include CH₂(4-membered heterocycloalkyl), for example CH₂-azetidiny, CH₂CH₂-azetidiny. Examples of C₁₋₄alkyleneO(4-10 membered heterocycloalkyl) include C₁₋₄alkyIOC₄heterocycloalkyl, such as CH₂O-azetidiny. An example of C₁₋₄alkyleneO(C₃₋₆alkynyl) is CH₂OCH₂C≡CH.
- 15

Where substituents are indicated as *being optionally substituted in* formula (I) in the embodiments and preferences set out below, the optional substituent may be attached to an available carbon atom, which means a carbon atom which is attached to a hydrogen atom i.e. a C-H group or the optional substituent may be attached to an available nitrogen atom, which means a nitrogen atom which is attached to a hydrogen atom i.e. a N-H group. The optional substituent replaces the hydrogen atom attached to the carbon atom or the hydrogen atom attached to the nitrogen atom.

20


25 In one embodiment, R_{1a} is methyl. In one embodiment, R_{1a} is H.

In one embodiment, R_{1b} is H. In one embodiment, R_{1b} is F.

In one embodiment, R_{1a} is H and R_{1b} is H. In one embodiment, R_{1a} is H and R_{1b} is F. In one embodiment, R_{1a} is methyl and R_{1b} is H.

30

In one embodiment, A is group (Aa):



In one embodiment, R_2 is H. In one embodiment, R_2 is halo. In one embodiment R_2 is C_{1-4} alkyl, such as methyl, ethyl, propyl or butyl, especially methyl. In one embodiment, R_2 is C_{1-4} haloalkyl, such as CF_3 . In one embodiment R_2 is C_{1-4} alkylene(aryl), such as benzyl. In one embodiment R_2 is C_{1-4} alkylene(OH), such as CH_2OH . In one embodiment R_2 is C_{1-4} alkylene OC_{1-4} alkyl, such as CH_2OMe , CH_2OEt or CH_2OPr , especially CH_2OMe . In one embodiment R_2 is C_{1-4} alkylene OC_{3-6} cycloalkyl, such as CH_2O-C_3 cycloalkyl or CH_2O-C_4 cycloalkyl, for example CH_2O -cyclopropyl or CH_2O -cyclobutyl. In one embodiment R_2 is C_{1-4} alkyleneO(aryl), such as CH_2OPh . In one embodiment R_2 is C_{1-4} alkylene(4-10 membered heterocycloalkyl), such as CH_2 (4-membered heterocycloalkyl), for example CH_2 -azetidiny, or CH_2CH_2 (4-membered heterocycloalkyl) for example CH_2CH_2 -azetidiny. In one embodiment R_2 is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl) such as $C_{1-4}alkyLOC_4$ heterocycloalkyl, especially CH_2O -azetidiny. In one embodiment R_2 is C_{1-4} alkyleneO(C_{3-6} alkynyl), such as $CH_2OCH_2C\equiv CH$. In one embodiment R_2 is 4-10 membered heterocycloalkyl, such as 1-methyl piperaziny. In one embodiment R_2 is C_{1-4} alkylene(CO_2H), such as $CH_2CH_2(CO_2H)$. In one embodiment R_2 is OC_{1-4} alkylene(CO_2H). In one embodiment R_2 is C_{1-4} alkylene OC_{1-4} alkylene(CO_2H). In one embodiment R_2 is $N(R_{2aa})C_{1-4}$ alkylene(CO_2H). In one embodiment R_2 is C_{1-4} alkylene($NR_{2ab}R_{2ac}$). In one embodiment R_2 is OC_{1-4} alkylene($NR_{2ab}R_{2ac}$). In one embodiment R_2 is $N(R_{2aa})C_{1-4}$ alkylene($NR_{2ab}R_{2ac}$).

In one embodiment, R_2 is H, C_{1-4} alkyl, C_{1-4} alkylene(aryl), C_{1-4} alkylene(OH), C_{1-4} alkylene(C_{3-6} cycloalkyl), C_{1-4} alkylene(4-7 membered heterocycloalkyl), C_{1-4} alkoxy, OC_{1-4} alkylene(aryl), C_{1-4} alkylene OC_{1-4} alkyl, C_{1-4} alkylene OC_{3-6} cycloalkyl, C_{1-4} alkyleneO(4-7 membered heterocycloalkyl), C_{1-4} alkyleneO(aryl), C_{3-6} alkynyl or C_{1-4} alkenylo(C_{3-6} alkynyl).

In one embodiment, R_2 is halo, C_{1-4} haloalkyl, C_{1-4} alkylene(8-10 membered heterocycloalkyl), C_{1-4} alkyleneO(8-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl such as 1-methyl piperaziny, C_{1-4} alkylene(CO_2H), such as $CH_2CH_2(CO_2H)$, OC_{1-4} alkylene(CO_2H), C_{1-4} alkylene OC_{1-4} alkylene(CO_2H), $N(R_{2aa})C_{1-4}$ alkylene(CO_2H), C_{1-4} alkylene($NR_{2ab}R_{2ac}$), OC_{1-4} alkylene($NR_{2ab}R_{2ac}$) or $N(R_{2aa})C_{1-4}$ alkylene($NR_{2ab}R_{2ac}$).

The aryl, heterocycloalkyl and cycloalkyl groups present in R_2 may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe ; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; CN ; OH ; $NR_{2a}R_{2b}$; SO_2R_{2c} ; and $NHSO_2R_{2c}$. Suitably, the aryl, heterocycloalkyl and cycloalkyl groups present in R_2 may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe ; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro;

and CN. Suitably, the aryl, heterocycloalkyl and cycloalkyl groups present in R_2 may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from OH; $NR_{2a}R_{2b}$; SO_2R_{2c} ; and $NHSO_2R_{2c}$.

- 5 In one embodiment R_{2a} is H. In a second embodiment R_{2a} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{2a} is H or methyl.

In one embodiment R_{2b} is H. In a second embodiment R_{2b} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{2b} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{2b} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{2b} is C_{1-4} haloalkyl, such as CF_3 . In a sixth embodiment, R_{2b} is aryl, such as phenyl. In a seventh embodiment, R_{2b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment R_{2b} is H or methyl.

15 In one embodiment R_{2c} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{2c} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{2c} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{2c} is C_{1-4} haloalkyl, such as CF_3 . In a fifth embodiment, R_{2c} is aryl, such as phenyl. In a sixth embodiment, R_{2c} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment R_{2c} is methyl.

25 In one embodiment, R_{2aa} is H. In one embodiment, R_{2aa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl.

In one embodiment, R_{2ab} is H. In one embodiment, R_{2aa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{2ac} is H. In one embodiment, R_{2ac} is C_{1-4} alkyl. Alternatively, in one embodiment R_{2ab} and R_{2ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl.

30 In one embodiment R_{2x} is H. In one embodiment R_{2x} is halo, such as chloro or fluoro. In one embodiment R_{2x} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{2x} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{2x} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

35 Alternatively, R_{2x} is C_{1-4} alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, such as 1-methyl piperaziny, C_{1-4} alkyleneO(4-10 membered heterocycloalkyl),

C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2xaa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2xab}R_{2xac}), OC₁₋₄alkylene(NR_{2xab}R_{2xac}) or N(R_{2xaa})C₁₋₄alkylene(NR_{2xab}R_{2xac}). In one embodiment, R_{2x} is C₁₋₄alkylene(4-10 membered heterocycloalkyl). In one embodiment, R_{2x} is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl. In one embodiment, R_{2x} is C₁₋₄alkyleneO(4-10 membered heterocycloalkyl). In one embodiment, R_{2x} is C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H). In one embodiment, R_{2x} is OC₁₋₄alkylene(CO₂H). In one embodiment, R_{2x} is C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H). In one embodiment, R_{2x} is N(R_{2xaa})C₁₋₄alkylene(CO₂H). In one embodiment, R_{2x} is C₁₋₄alkylene(NR_{2xab}R_{2xac}). In one embodiment, R_{2x} is OC₁₋₄alkylene(NR_{2xab}R_{2xac}). In one embodiment, R_{2x} is N(R_{2xaa})C₁₋₄alkylene(NR_{2xab}R_{2xac}).

Within such embodiments, R₂ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl). In one embodiment R₂ is H. In one embodiment R₂ is halo, such as chloro or fluoro. In one embodiment R₂ is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R₂ is C₁₋₄haloalkyl, such as CF₃. In one embodiment, R₂ is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In one embodiment, R₂ is C₁₋₄alkyleneOC₁₋₄alkyl. In one embodiment, R₂ is C₁₋₄alkyleneOC₃₋₆cycloalkyl. In one embodiment, R₂ is C₁₋₄alkyleneO(4-10 membered heterocycloalkyl).

The heterocycloalkyl groups present in R_{2x} may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, aryl, heterocycloalkyl and cycloalkyl are optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{2xa}R_{2xb}, SO₂R_{2xc} and NHSO₂R_{2xc}. Suitably, the heterocycloalkyl groups present in R_{2x} may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C₁₋₄alkyl, such as methyl; C₃₋₆cycloalkyl, such as cyclopropyl; C₁₋₄alkoxy, such as OMe; C₁₋₄haloalkyl, such as CF₃; halo, such as chloro or fluoro; and CN. Suitably, the heterocycloalkyl groups present in R_{2x} may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from OH; NR_{2xa}R_{2xb}; SO₂R_{2xc}; and NHSO₂R_{2xc}.

In one embodiment R_{2xa} is H. In a second embodiment R_{2xa} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{2xa} is H or methyl.

In one embodiment R_{2xb} is H. In a second embodiment R_{2xb} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{2xb} is C₃₋₆cycloalkyl, such as cyclopropyl,

cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{2xb} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{2xb} is C_{1-4} haloalkyl, such as CF_3 . In a sixth embodiment, R_{2xb} is aryl, such as phenyl. In a seventh embodiment, R_{2xb} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment R_{2xb} is H or methyl.

In one embodiment R_{2xc} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{2xc} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{2xc} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{2xc} is C_{1-4} haloalkyl, such as CF_3 . In a fifth embodiment, R_{2xc} is aryl, such as phenyl. In a sixth embodiment, R_{2xc} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment R_{2xc} is methyl.


In one embodiment, R_{2xaa} is H. In one embodiment, R_{2xaa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl.

In one embodiment, R_{2xab} is H. In one embodiment, R_{2xab} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{2xac} is H. In one embodiment, R_{2xac} is C_{1-4} alkyl. Alternatively, in one embodiment R_{2xab} and R_{2xac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl.

Suitably, in any of the above embodiments the aryl is substituted by 1, 2 or 3 substituents, such as 1 or 2, e.g. 1 substituent, each independently selected from methyl, chloro and fluoro. In one embodiment, the aryl is not substituted.


Suitably, in any of the above embodiments the heterocycloalkyl is substituted by 1, 2 or 3 substituents, such as 1 or 2, e.g. 1 substituent, each independently selected from CH_2CH_2F (particularly as a substituent on a nitrogen atom) and fluoro (particularly as a substituent on a carbon atom). In one embodiment, the heterocycloalkyl is not substituted.

Suitably, when the heterocycloalkyl is azetidiny, the nitrogen atom is in the 1-position or the 3-position (i.e. 1-azetidiny or 3-azetidiny) relative to the point of attachment to the remainder of the R_2 group, for example:



1-azetidiny or 3-azetidiny

Suitably, when R_2 is C_{1-4} alkylene(4-10 membered heterocycloalkyl), such as CH_2 -azetidiny or CH_2CH_2 -azetidiny, the azetidiny is 1-azetidiny. Suitably, when R_2 is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), such as CH_2O -azetidiny, the azetidiny is 3-azetidiny. For example:



1-azetidiny, 1-azetidiny or 3-azetidiny

Suitably, in any of the above embodiments when the heterocycloalkyl contains one or more nitrogen atoms, as required by valency the nitrogen atom(s) may be connected to a hydrogen atom to form an NH group. Alternatively the nitrogen atom(s) may be substituted (such as one nitrogen atom is substituted), for example by C_{1-4} alkyl, such as methyl, C_{1-4} haloalkyl, such as CH_2CH_2F , $C(O)H$, $C(O)C_{1-4}$ alkyl, $C(O)OC_{1-4}$ alkyl, such as $C(O)OtBu$, $C(O)OC_{1-4}$ alkylene(aryl) such as $C(O)Obz$, $C(O)NHC_{1-4}$ alkyl, $C(O)NHC_{1-4}$ alkylene(aryl) such as $C(O)NHBz$, an Fmoc group, $C(O)C_{1-4}$ haloalkyl, $C(O)OC_{1-4}$ haloalkyl or $C(O)NHC_{1-4}$ haloalkyl. Suitably, when the heterocycloalkyl contains one or more S atoms, the S atom(s) is substituted (such as one S atom is substituted) by one or two oxygen atoms (i.e. $S(O)$ or $S(O)_2$). Alternatively, any sulphur atom(s) in the heterocycloalkyl ring is not substituted.

When the one or more (such as one) nitrogen atoms is substituted by C_{1-4} alkyl, such as methyl, C_{1-4} haloalkyl, such as CH_2CH_2F , $C(O)H$, $C(O)C_{1-4}$ alkyl, $C(O)OC_{1-4}$ alkyl, such as $C(O)OtBu$, $C(O)OC_{1-4}$ alkylene(aryl) such as $C(O)Obz$, $C(O)NHC_{1-4}$ alkyl, $C(O)NHC_{1-4}$ alkylene(aryl) such as $C(O)NHBz$, an Fmoc group, $C(O)C_{1-4}$ haloalkyl, $C(O)OC_{1-4}$ haloalkyl or $C(O)NHC_{1-4}$ haloalkyl, these substituents may be present in addition to the optional substituents described above for

heterocycloalkyl. The substituents on the nitrogen atom may be referred to or function as protecting groups, which can be added and removed by methods known to the person skilled in the art.

- 5 Suitably, in any of the above embodiments the cycloalkyl is substituted by 1, 2 or 3 C₁₋₄alkyl substituents, such as 1 or 2 e.g. 1 C₁₋₄alkyl substituent, such as methyl, ethyl and propyl, especially methyl. In one embodiment, the cycloalkyl is not substituted.

- 10 Suitably unless R₂ is H, C₁₋₄alkyl or C₁₋₄alkyleneOC₁₋₄alkyl, then m is 0. More suitably, unless R₂ is H, methyl or CH₂OMe, then m is 0.

- When present, in one embodiment each R₃ is independently halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy. In one embodiment at least one R₃ is halo, such as chloro or fluoro. In one embodiment at least one R₃ is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In one
15 embodiment, at least one R₃ is C₁₋₄haloalkyl, such as CF₃. In one embodiment, at least one R₃ is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In one embodiment, each R₃ is independently fluoro or methyl. In one embodiment each R₃ is independently halo, C₁₋₄alkyl (other than methyl, ethyl or n-propyl), C₁₋₄haloalkyl or C₁₋₄alkoxy.


- 20 In one embodiment, m is 0, 1 or 2. In one embodiment, m is 1 or 2. In one embodiment, m is 1. In one embodiment, m is 2. In one embodiment, m is 2, one R₃ is fluoro and one R₃ is methyl.

As used herein, R_{2x'} is as defined for R_{2x} save that it is not H.

- 25 In one preferred embodiment, m is 0 and R_{2x} is R_{2x'}. In another preferred embodiment, R_{2x} is H, m is 1 and R₃ is in the 6-position. In another preferred embodiment, m is 1, R_{2x} is R_{2x'}, and R₃ is in the 6-position. In one embodiment, m is 2, one R_{2x} is R_{2x'}, one R₃ is in the 4-position, and the other R₃ is in the 6-position. In one embodiment, m is 2, R_{2x} is R_{2x'}, one R₃ is in the 5-position, and the other R₃ is in the 6-position.

- 30 Suitably, when R₂ is methyl, m is 1, and R₃ is chloro, R_{2x} is H and R₃ is not in the 5-position. Suitably, when R₂ is methyl, m is 1 and R₃ is chloro, R₃ is in the 4-position or 6-position and R_{2x} is R_{2x'}. Suitably, when R₂ is H, m is 1 and R₃ is C₁₋₄alkyl, R_{2x} is H and R₃ is not in the 5-position. Suitably, when R₂ is H, m is 1, R_{2x} is H and R₃ is C₁₋₄alkyl, R₃ is in the 4-position or the 6-
35 position. Suitably, when m is 1, R_{2x} is R_{2x'} and the R₃ group is not in the 5-position. Suitably, when m is 2, R_{2x} is H and the two R₃ groups are in the 4-position and the 6-position.

References to substituent position are in respect of attachment to the amide moiety, for example:




- 5 Examples of suitable substituents include 3-fluoro-2-methyl; 3-fluoro-2-CH₂OCH₃; 3-chloro-2-methyl-; 4,5-difluoro-2-methyl; 5-chloro-2-isopropyl; 5-fluoro-2-methyl; 2-isopropyl-6-methyl; 2,6-dimethyl; 2-methyl; 2-isopropyl; 4-fluoro-3-methyl; 3-fluoro-4-methyl; 3,4-difluoro-2,6-dimethyl; 3,5-difluoro-2,6-dimethyl; and 3-fluoro-2,6-dimethyl. In one embodiment, each R₃ is the same. In one embodiment, each R₃ is different.

10

Suitably, one of R₂ and R_{2x} groups (i.e. the 2-position or 3-position substituent) is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl, C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2ab}R_{2ac}), OC₁₋₄alkylene(NR_{2ab}R_{2ac}) or N(R_{2aa})C₁₋

- 15 4alkylene(NR_{2ab}R_{2ac}), and R₃ is halo, such as chloro and fluoro or C₁₋₄alkyl such as methyl e.g. R₃ in the 5-position is halo such as fluoro and R₃ in the 6-position is C₁₋₄alkyl, such as methyl.

In one embodiment (Aa) is group (Aa'):



- 20 wherein:

R₂ is 4-10 membered heterocycloalkyl such as 1-methyl piperazinyl, C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2aa})C₁₋

4alkylene(CO₂H), C₁₋₄alkylene(NR_{2ab}R_{2ac}), OC₁₋₄alkylene(NR_{2ab}R_{2ac}) or N(R_{2aa})C₁₋₄alkylene(NR_{2ab}R_{2ac});

R_{2x} is H or C₁₋₄alkyl,

R_{2v} is H or C₁₋₄alkyl;

5 R_{2y} is H, halo or C₁₋₄alkyl;

R_{2z} is H, halo or C₁₋₄alkyl;

each R_{2aa} is independently selected from H and C₁₋₄alkyl;

each R_{2ab} is independently selected from H and C₁₋₄alkyl;

each R_{2ac} is independently selected from H and C₁₋₄alkyl; or

10 R_{2ab} and R_{2ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

or

R₂ is H or C₁₋₄alkyl;

15 R_{2x} is 4-10 membered heterocycloalkyl such as 1-methyl piperazinyl, C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2xaa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2xab}R_{2xac}), OC₁₋₄alkylene(NR_{2xab}R_{2xac}) or N(R_{2xaa})C₁₋₄alkylene(NR_{2xab}R_{2xac});

R_{2v} is H or C₁₋₄alkyl;

20 R_{2y} is H, halo or C₁₋₄alkyl;

R_{2z} is H, halo or C₁₋₄alkyl;

each R_{2xaa} is independently selected from H and C₁₋₄alkyl;

each R_{2xab} is independently selected from H and C₁₋₄alkyl;

each R_{2xac} is independently selected from H and C₁₋₄alkyl; or


25 R_{2xab} and R_{2xac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl.

In one embodiment, R₂ is CH₂CH₂(CO₂H), R_{2x} is H, R_{2v} is H, R_{2y} is fluoro and R_{2z} is methyl. In

one embodiment, R₂ is 1-methyl piperazinyl, R_{2x} is H, R_{2v} is H, R_{2y} is fluoro and R_{2z} is methyl.

30 In one embodiment, R₂ is H, R_{2x} is CH₂CH₂(CO₂H), R_{2v} is H, R_{2y} is fluoro and R_{2z} is methyl.

In one embodiment, A is group (Ab):



In one embodiment, R_4 is H. In a second embodiment, R_4 is C_{1-4} alkyl, such as methyl. In a third embodiment R_4 is C_{1-4} alkylene(aryl), such as benzyl. In one preferred embodiment, R_4 is H, methyl or benzyl, in particular methyl or benzyl. Suitably, the aryl present in R_4 is substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; CN; OH; $NR_{4a}R_{4b}$; SO_2R_{4c} ; and $NHSO_2R_{4c}$. Suitably, the aryl present in R_4 is substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; and CN. Suitably, the aryl present in R_4 is substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from OH; $NR_{4a}R_{4b}$; SO_2R_{4c} ; and $NHSO_2R_{4c}$. Suitably, the aryl is substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from methyl, chloro and fluoro. In one embodiment, the aryl is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 methyl groups. In one embodiment, the aryl is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 chloro groups. In one embodiment, the aryl is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 fluoro groups. Suitably, the aryl is not substituted.

In one embodiment R_{4a} is H. In a second embodiment R_{4a} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{4a} is H or methyl.

In one embodiment R_{4b} is H. In a second embodiment R_{4b} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{4b} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{4b} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{4b} is C_{1-4} haloalkyl, such as CF_3 . In a sixth embodiment, R_{4b} is aryl, such as phenyl. In a seventh embodiment, R_{4b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment, R_{4b} is H or methyl.

In one embodiment R_{4c} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{4c} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{4c} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{4c} is C_{1-4} haloalkyl, such as CF_3 . In a fifth embodiment, R_{4c} is aryl, such as phenyl. In a sixth embodiment, R_{4c} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment, R_{4c} is methyl.


In one embodiment, R^5 is H.

When present, in one embodiment each R_6 is independently fluoro or methyl. Suitably, n may be 1, 2 or 3, such as 1 or 2 e.g. 1.

5

In one embodiment, n is 0.

In one embodiment, group A is group (Ac):




10

In one embodiment, R_7 is C_{1-4} alkyl. In one embodiment, R_7 is methyl, CH_2OH or CH_2OMe , especially methyl.


In one embodiment, o is 2. Suitably, when o is 2, B is not group (Ba).

15

Suitably, the stereochemistry of R_7 is *trans* in respect of bond linking the (Ac) group to the amide moiety, for example having one of the two following stereochemical arrangements:



20 In one embodiment, A is group (Ad):



In one embodiment, X is a bond or O. Suitably, X is a bond. Alternatively, X is O. In a second embodiment, X is CH_2 .

25

In one embodiment, R_8 is halo. In a second embodiment, R_8 is C_{1-4} alkyl. In a third embodiment, R_8 is C_{1-4} alkoxy. In a fourth embodiment, R_8 is OH.

When present, in one embodiment each R_8 is independently methyl, OMe or fluoro. In one embodiment, each R_8 is independently OCH_2 -cyclopropyl, OCH_2 -oxetanyl, OCH_2CH_2F , methyl, OMe, OEt or fluoro, for example OCH_2CH_2F , OMe or OEt, especially OMe. In another embodiment, each R_8 is independently OCH_2 -cyclopropyl, OCH_2 -oxetanyl, OCH_2CH_2F or OEt, such as OCH_2 -cyclopropyl, OCH_2 -oxetanyl or OCH_2CH_2F .

Suitably, the cycloalkyl and heterocycloalkyl present in R_8 are each independently substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; CN; OH; $NR_{8a}R_{8b}$; SO_2R_{8c} ; and $NHSO_2R_{8c}$. Suitably, the cycloalkyl and heterocycloalkyl present in R_8 are each independently substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; and CN. Suitably, the cycloalkyl and heterocycloalkyl present in R_8 are each independently substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from OH; $NR_{8a}R_{8b}$; SO_2R_{8c} ; and $NHSO_2R_{8c}$.

In one embodiment R_{8a} is H. In a second embodiment R_{8a} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{8a} is H or methyl.

In one embodiment R_{8b} is H. In a second embodiment R_{8b} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{8b} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{8b} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{8b} is C_{1-4} haloalkyl, such as CF_3 . In a sixth embodiment, R_{8b} is aryl, such as phenyl. In a seventh embodiment, R_{8b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment R_{8b} is H or methyl.


In one embodiment R_{8c} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{8c} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{8c} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{8c} is C_{1-4} haloalkyl, such as CF_3 . In a fifth embodiment, R_{8c} is aryl, such as phenyl. In a sixth embodiment, R_{8c} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment, R_{8c} is methyl.

Suitably, the cycloalkyl present in R_8 (e.g. cyclopropyl) is substituted by 1, 2 or 3 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from fluoro and C_{1-4} alkyl, such as methyl, ethyl and propyl, especially methyl. In one embodiment, the cycloalkyl is not substituted.


Suitably, the heterocycloalkyl present in R_8 (e.g. oxetanyl) is substituted by 1, 2 or 3

- 5 substituents, such as 1 or 2 e.g. 1 substituent, each independently selected from fluoro and C_{1-4} alkyl, such as methyl, ethyl and propyl, especially methyl. In one embodiment, the heterocycloalkyl is not substituted.


Suitably, when the heterocycloalkyl is oxetanyl, the oxygen atom is in the 2- or 3-position (i.e. 2-oxetanyl or 3-oxetanyl) relative to the point of attachment to the remainder of the R_8 group, for example:



Suitably, when R_8 is OC_{1-4} alkylene(4-10 membered heterocycloalkyl), such as OCH_2 -oxetanyl, the oxetanyl is 3-oxetanyl, for example:



When p is 1 or 2, R_8 may be in the 2- and/or 3 position. In one embodiment, p is 1 and R_8 is in the 2-position. In a second embodiment, p is 1 and R_8 is in the 3-position. In a third embodiment, p is 2, one R_8 is in the 2-position and one R_8 is in the 3-position. For example:




Examples of suitable R_8 substituents include 2-methyl, 2-methoxy, 2-ethoxy, 2- OCH_2CH_2F , 2- OCH_2 -cyclopropyl and 2- OCH_2 -oxetanyl such as 2-methyl and 2-methoxy.

In one embodiment, p is 0 or 1.

Suitably, when X is O, p is 1 or 2. Suitably, when X is a bond and group B is (Ba), Y is C(R₁₁)(R₁₂), R₁₁ and R₁₂ are both H, and p is 1 or 2. Suitably, when X is CH₂, B is not (Ba).


Suitably, when X is a bond and p is 0, the compound has the *R* stereochemical configuration.

- 5 Suitably, when X is as defined above and p is 1, the stereochemistry of R₈ is *trans* in respect of the bond linking the (Ad) group to the amide substituent, for example having one of the following stereochemical arrangements:




- 10 When present, in one embodiment each R₉ is independently fluoro.

Suitably, when q is 1, 2, 3 or 4, R₉ may be in the 5-, 6-, 7- and/or 8-position(s). In one embodiment, q is 1 and R₉ is in the 5-position. In a second embodiment, q is 1 and R₉ is in the 6-position. For example:




- 15 In embodiments wherein X is a bond, it will be understood that the substituent position numbering will change accordingly with the total number of atoms in the bicyclic ring system, for example:



- 20 Suitable examples of R₉ substituents when X is a bond are 4-fluoro and 5-fluoro.

In one embodiment, q is 1 or 2.

In one embodiment, group A is (Ae):



In one embodiment, R₁₇ is H. In one embodiment, R₁₇ is halo. In one embodiment R₁₇ is C₁₋₄alkyl, such as methyl, ethyl, propyl or butyl, especially methyl. In one embodiment, R₁₇ is C₁₋₄haloalkyl, such as CF₃. In one embodiment R₁₇ is C₁₋₄alkylene(aryl), such as benzyl. In one embodiment R₁₇ is C₁₋₄alkylene(OH), such as CH₂OH. In one embodiment R₁₇ is C₁₋₄alkyleneOC₁₋₄alkyl, such as CH₂OMe, CH₂OEt or CH₂OPr, especially CH₂OMe. In one embodiment R₁₇ is C₁₋₄alkyleneOC₃₋₆cycloalkyl, such as CH₂O-C₃cycloalkyl or CH₂O-C₄cycloalkyl, for example CH₂O-cyclopropyl or CH₂O-cyclobutyl. In one embodiment R₁₇ is C₁₋₄alkyleneO(aryl), such as CH₂OPh. In one embodiment R₁₇ is C₁₋₄alkylene(4-10 membered heterocycloalkyl), such as CH₂(4-membered heterocycloalkyl), for example CH₂-azetidiny, or CH₂CH₂(4-membered heterocycloalkyl) for example CH₂CH₂-azetidiny. In one embodiment R₁₇ is C₁₋₄alkyleneO(4-10 membered heterocycloalkyl) such as C₁₋₄alkylOC₄heterocycloalkyl, especially CH₂O-azetidiny. In one embodiment R₁₇ is C₁₋₄alkyleneO(C₃₋₆alkynyl), such as CH₂OCH₂C≡CH. In one embodiment R₁₇ is 4-10 membered heterocycloalkyl, such as 1-methyl piperaziny. In one embodiment R₁₇ is C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H). In one embodiment R₁₇ is OC₁₋₄alkylene(CO₂H). In one embodiment R₁₇ is C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H). In one embodiment R₁₇ is N(R_{2aa})C₁₋₄alkylene(CO₂H). In one embodiment R₁₇ is C₁₋₄alkylene(NR_{17ab}R_{17ac}). In one embodiment R₁₇ is OC₁₋₄alkylene(NR_{17ab}R_{17ac}). In one embodiment R₁₇ is N(R_{17aa})C₁₋₄alkylene(NR_{17ab}R_{17ac}).

In one embodiment, R₁₇ is 4-10 membered heterocycloalkyl such as 1-methyl piperaziny, C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{17aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{17ab}R_{17ac}), OC₁₋₄alkylene(NR_{17ab}R_{17ac}) or N(R_{17aa})C₁₋₄alkylene(NR_{17ab}R_{17ac}).

The aryl, cycloalkyl or heterocycloalkyl groups present in R₁₇ may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C₁₋₄alkyl, such as methyl; C₃₋₆cycloalkyl, such as cyclopropyl; C₁₋₄alkoxy, such as OMe; C₁₋₄haloalkyl, such as CF₃; halo, such as chloro or fluoro; CN; OH; NR_{17a}R_{17b}; SO₂R_{17c}; and NHSO₂R_{17c}. Suitably, the aryl, heterocycloalkyl and cycloalkyl groups present in R₁₇ may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C₁₋₄alkyl, such as methyl; C₃₋₆cycloalkyl, such as

cyclopropyl; C₁₋₄alkoxy, such as OMe; C₁₋₄haloalkyl, such as CF₃; halo, such as chloro or fluoro; and CN. Suitably, the aryl, heterocycloalkyl and cycloalkyl groups present in R₂ may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from OH; NR_{17a}R_{17b}; SO₂R_{17c}; and NHSO₂R_{17c}.

5

In one embodiment R_{17a} is H. In a second embodiment R_{17a} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{17a} is H or methyl.

In one embodiment R_{17b} is H. In a second embodiment R_{17b} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{17b} is C₃₋₆cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{17b} is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{17b} is C₁₋₄haloalkyl, such as CF₃. In a sixth embodiment, R_{17b} is aryl, such as phenyl. In a seventh embodiment, R_{17b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment R_{17b} is H or methyl.

15

In one embodiment R_{17c} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{17c} is C₃₋₆cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{17c} is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{17c} is C₁₋₄haloalkyl, such as CF₃. In a fifth embodiment, R_{17c} is aryl, such as phenyl. In a sixth embodiment, R_{17c} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment R_{17c} is methyl.

20

In one embodiment, R_{17aa} is H. In one embodiment, R_{17aa} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl.

25

In one embodiment, R_{17ab} is H. In one embodiment, R_{17ab} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{17ac} is H. In one embodiment, R_{17ac} is C₁₋₄alkyl. Alternatively, in one embodiment R_{17ab} and R_{17ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl.

30

Within such embodiments, R₁₈ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl). In one embodiment R₁₈ is H. In one embodiment R₁₈ is halo, such as chloro or fluoro. In one embodiment R₁₈ is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R₁₈ is C₁₋₄haloalkyl, such as CF₃. In one embodiment, R₁₈ is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In one embodiment, R₁₈ is C₁₋₄alkyleneOC₁₋₄alkyl. In one

35

embodiment, R_{18} is C_{1-4} alkyleneOC₃₋₆cycloalkyl. In one embodiment, R_{18} is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl).

Within such embodiments, R_{20} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy. In one embodiment R_{20} is H. In one embodiment R_{20} is halo, such as chloro or fluoro. In one embodiment R_{20} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{20} is C_{1-4} haloalkyl, such as CF₃. In one embodiment, R_{20} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

Within such embodiments, R_{21} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy. In one embodiment R_{21} is H. In one embodiment R_{21} is halo, such as chloro or fluoro. In one embodiment R_{21} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{21} is C_{1-4} haloalkyl, such as CF₃. In one embodiment, R_{21} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

Alternatively, R_{18} is C_{1-4} haloalkyl, C_{1-4} alkylene(aryl), C_{1-4} alkylene(OH), C_{1-4} alkylene(C₃₋₆cycloalkyl), C_{1-4} alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl, OC₁₋₄alkylene(aryl), C_{1-4} alkyleneOC₁₋₄alkyl, C_{1-4} alkyleneOC₃₋₆cycloalkyl, C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkyleneO(aryl), C₃₋₆alkynyl, C_{1-4} alkenylO(C₃₋₆alkynyl), C_{1-4} alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C_{1-4} alkyleneOC₁₋₄alkylene(CO₂H), N(R_{18aa}) C_{1-4} alkylene(CO₂H), C_{1-4} alkylene(NR_{18ab}R_{18ac}), OC₁₋₄alkylene(NR_{18ab}R_{18ac}) or N(R_{18aa}) C_{1-4} alkylene(NR_{18ab}R_{18ac}).

In one embodiment, R_{18} is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl, C_{1-4} alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C_{1-4} alkyleneOC₁₋₄alkylene(CO₂H), N(R_{18aa}) C_{1-4} alkylene(CO₂H), C_{1-4} alkylene(NR_{18ab}R_{18ac}), OC₁₋₄alkylene(NR_{18ab}R_{18ac}) or N(R_{18aa}) C_{1-4} alkylene(NR_{18ab}R_{18ac}).

In one embodiment, R_{18} is C_{1-4} haloalkyl, such as CF₃. In one embodiment R_{18} is C_{1-4} alkylene(aryl), such as benzyl. In one embodiment R_2 is C_{1-4} alkylene(OH), such as CH₂OH. In one embodiment R_{18} is C_{1-4} alkyleneOC₁₋₄alkyl, such as CH₂OMe, CH₂OEt or CH₂OPr, especially CH₂OMe. In one embodiment R_{18} is C_{1-4} alkyleneOC₃₋₆cycloalkyl, such as CH₂O-C₃cycloalkyl or CH₂O-C₄cycloalkyl, for example CH₂O-cyclopropyl or CH₂O-cyclobutyl. In one embodiment R_{18} is C_{1-4} alkyleneO(aryl), such as CH₂OPh. In one embodiment R_{18} is C_{1-4} alkylene(4-10 membered heterocycloalkyl), such as CH₂(4-membered heterocycloalkyl), for example CH₂-azetidiny, or CH₂CH₂(4-membered heterocycloalkyl) for example CH₂CH₂-azetidiny. In one embodiment R_{18} is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl) such as C_{1-4} alkylOC₄heterocycloalkyl,

especially CH₂O-azetidinyI. In one embodiment R₁₈ is C₁₋₄alkyleneO(C₃₋₆alkynyl), such as CH₂OCH₂C≡CH. In one embodiment R₁₈ is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyI,. In one embodiment R₁₈ is C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H). In one embodiment R₁₈ is OC₁₋₄alkylene(CO₂H). In one embodiment R₁₈ is C₁₋₄alkyleneOC₁₋

5 4alkylene(CO₂H). In one embodiment R₁₈ is N(R_{18aa})C₁₋₄alkylene(CO₂H). In one embodiment R₁₈ is C₁₋₄alkylene(NR_{18ab}R_{18ac}). In one embodiment R₁₈ is OC₁₋₄alkylene(NR_{18ab}R_{18ac}). In one embodiment R₁₈ is N(R_{18aa})C₁₋₄alkylene(NR_{18ab}R_{18ac}).

The aryl, cycloalkyl or heterocycloalkyl groups present in R₁₈ may be optionally substituted by
 10 up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C₁₋₄alkyl, such as methyl; C₃₋₆cycloalkyl, such as cyclopropyl; C₁₋₄alkoxy, such as OMe; C₁₋₄haloalkyl, such as CF₃; halo, such as chloro or fluoro; CN; OH; NR_{18a}R_{18b}; SO₂R_{18c}; and NHSO₂R_{18c}. Suitably, the heterocycloalkyl groups present in R₁₈ may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each
 15 independently selected from C₁₋₄alkyl, such as methyl; C₃₋₆cycloalkyl, such as cyclopropyl; C₁₋₄alkoxy, such as OMe; C₁₋₄haloalkyl, such as CF₃; halo, such as chloro or fluoro; and CN. Suitably, the heterocycloalkyl groups present in R₁₈ may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from OH; NR_{18a}R_{18b}; SO₂R_{18c}; and NHSO₂R_{18c}.

20 In one embodiment R_{18a} is H. In a second embodiment R_{18a} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{18a} is H or methyl.

In one embodiment R_{18b} is H. In a second embodiment R_{18b} is C₁₋₄alkyl, such as methyl, ethyl or
 25 propyl, especially methyl. In a third embodiment, R_{18b} is C₃₋₆cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{18b} is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{18b} is C₁₋₄haloalkyl, such as CF₃. In a sixth embodiment, R_{18b} is aryl, such as phenyl. In a seventh embodiment, R_{18b} is 4-10 membered heterocycloalkyl, such as azetidinyI or oxetanyI. In an eighth embodiment R_{18b} is H
 30 or methyl.

In one embodiment R_{18c} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{18c} is C₃₋₆cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{18c} is C₁₋₄alkoxy, such as OMe or OEt, especially
 35 OMe. In a fourth embodiment, R_{18c} is C₁₋₄haloalkyl, such as CF₃. In a fifth embodiment, R_{18c} is aryl, such as phenyl. In a sixth embodiment, R_{18c} is 4-10 membered heterocycloalkyl, such as azetidinyI or oxetanyI. In a seventh embodiment R_{18c} is methyl.

In one embodiment, R_{18aa} is H. In one embodiment, R_{18aa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl.

- 5 In one embodiment, R_{18ab} is H. In one embodiment, R_{18aa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{18ac} is H. In one embodiment, R_{18ac} is C_{1-4} alkyl. Alternatively, in one embodiment R_{18ab} and R_{18ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl.
- 10 Within such embodiments, R_{17} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl C_{1-4} alkoxy, C_{1-4} alkyleneOC₁₋₄alkyl, C_{1-4} alkyleneOC₃₋₆cycloalkyl or C_{1-4} alkyleneO(4-10 membered heterocycloalkyl). In one embodiment R_{17} is H. In one embodiment R_{18} is halo, such as chloro or fluoro. In one embodiment R_{17} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{17} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{17} is C_{1-4} alkoxy, such as
- 15 OMe or OEt, especially OMe. In one embodiment, R_{17} is C_{1-4} alkyleneOC₁₋₄alkyl. In one embodiment, R_{17} is C_{1-4} alkyleneOC₃₋₆cycloalkyl. In one embodiment, R_{17} is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl).

- Within such embodiments, R_{20} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy. In one
- 20 embodiment R_{20} is H. In one embodiment R_{20} is halo, such as chloro or fluoro. In one embodiment R_{20} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{20} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{20} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

- 25 Within such embodiments, R_{21} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy. In one embodiment R_{21} is H. In one embodiment R_{21} is halo, such as chloro or fluoro. In one embodiment R_{21} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{21} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{21} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

- 30 Alternatively, R_{21} is C_{1-4} haloalkyl, C_{1-4} alkylene(aryl), C_{1-4} alkylene(OH), C_{1-4} alkylene(C_{3-6} cycloalkyl), C_{1-4} alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl, OC₁₋₄alkylene(aryl), C_{1-4} alkyleneOC₁₋₄alkyl, C_{1-4} alkyleneOC₃₋₆cycloalkyl, C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkyleneO(aryl), C_{3-6} alkynyl, C_{1-4} alkenylO(C_{3-6} alkynyl), C_{1-4} alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C_{1-4} alkyleneOC₁₋₄alkylene(CO₂H), N(R_{21aa}) C_{1-4} alkylene(CO₂H), C_{1-4} alkylene(NR_{21ab}R_{21ac}), OC₁₋₄alkylene(NR_{21ab}R_{21ac}) or N(R_{21aa}) C_{1-4} alkylene(NR_{21ab}R_{21ac}).
- 35

In one embodiment, R_{21} is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl, C_{1-4} alkylene(CO_2H), such as $CH_2CH_2(CO_2H)$, OC_{1-4} alkylene(CO_2H), C_{1-4} alkylene OC_{1-4} alkylene(CO_2H), $N(R_{21aa})C_{1-4}$ alkylene(CO_2H), C_{1-4} alkylene($NR_{21ab}R_{21ac}$), OC_{1-4} alkylene($NR_{21ab}R_{21ac}$) or $N(R_{21aa})C_{1-4}$ alkylene($NR_{21ab}R_{21ac}$).

In one embodiment, R_{21} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment R_{21} is C_{1-4} alkylene(aryl), such as benzyl. In one embodiment R_{21} is C_{1-4} alkylene(OH), such as CH_2OH . In one embodiment R_{21} is C_{1-4} alkylene OC_{1-4} alkyl, such as CH_2OMe , CH_2OEt or CH_2OPr , especially CH_2OMe . In one embodiment R_{21} is C_{1-4} alkylene OC_{3-6} cycloalkyl, such as CH_2O-C_3 cycloalkyl or CH_2O-C_4 cycloalkyl, for example CH_2O -cyclopropyl or CH_2O -cyclobutyl. In one embodiment R_{21} is C_{1-4} alkyleneO(aryl), such as CH_2OPh . In one embodiment R_{21} is C_{1-4} alkylene(4-10 membered heterocycloalkyl), such as CH_2 (4-membered heterocycloalkyl), for example CH_2 -azetidiny, or CH_2CH_2 (4-membered heterocycloalkyl) for example CH_2CH_2 -azetidiny. In one embodiment R_{21} is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl) such as C_{1-4} alkyl OC_4 heterocycloalkyl, especially CH_2O -azetidiny. In one embodiment R_{21} is C_{1-4} alkyleneO(C_{3-6} alkynyl), such as $CH_2OCH_2C\equiv CH$. In one embodiment R_{21} is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl. In one embodiment R_{21} is C_{1-4} alkylene(CO_2H), such as $CH_2CH_2(CO_2H)$. In one embodiment R_2 is OC_{1-4} alkylene(CO_2H). In one embodiment R_{21} is C_{1-4} alkylene OC_{1-4} alkylene(CO_2H). In one embodiment R_{21} is $N(R_{21aa})C_{1-4}$ alkylene(CO_2H). In one embodiment R_{21} is C_{1-4} alkylene($NR_{21ab}R_{21ac}$). In one embodiment R_{21} is OC_{1-4} alkylene($NR_{21ab}R_{21ac}$). In one embodiment R_{21} is $N(R_{21aa})C_{1-4}$ alkylene($NR_{21ab}R_{21ac}$).

The aryl, cycloalkyl or heterocycloalkyl groups present in R_{21} may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe ; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; CN ; OH ; $NR_{21a}R_{21b}$; SO_2R_{21c} ; and $NHSO_2R_{21c}$. Suitably, the heterocycloalkyl groups present in R_{21} may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from C_{1-4} alkyl, such as methyl; C_{3-6} cycloalkyl, such as cyclopropyl; C_{1-4} alkoxy, such as OMe ; C_{1-4} haloalkyl, such as CF_3 ; halo, such as chloro or fluoro; and CN . Suitably, the heterocycloalkyl groups present in R_{18} may be optionally substituted by up to 3 substituents, such as 1, 2 or 3, such as 1 or 2, e.g. 1 substituent, each independently selected from OH ; $NR_{21a}R_{21b}$; SO_2R_{21c} ; and $NHSO_2R_{21c}$.

In one embodiment R_{21a} is H . In a second embodiment R_{21a} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment R_{21a} is H or methyl.

In one embodiment R_{21b} is H. In a second embodiment R_{21b} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{21b} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{21b} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{21b} is C_{1-4} haloalkyl, such as CF_3 . In a sixth embodiment, R_{21b} is aryl, such as phenyl. In a seventh embodiment, R_{21b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment R_{21b} is H or methyl.

10 In one embodiment R_{21c} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{21c} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{21c} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{21c} is C_{1-4} haloalkyl, such as CF_3 . In a fifth embodiment, R_{21c} is aryl, such as phenyl. In a sixth embodiment, R_{21c} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment R_{21c} is methyl.

In one embodiment, R_{21aa} is H. In one embodiment, R_{21aa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl.

20 In one embodiment, R_{21ab} is H. In one embodiment, R_{21aa} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{21ac} is H. In one embodiment, R_{21ac} is C_{1-4} alkyl. Alternatively, in one embodiment R_{21ab} and R_{21ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl.

25 Within such embodiments, R_{17} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyleneOC $_{1-4}$ alkyl, C_{1-4} alkyleneOC $_{3-6}$ cycloalkyl or C_{1-4} alkyleneO(4-10 membered heterocycloalkyl). In one embodiment R_{17} is H. In one embodiment R_{17} is halo, such as chloro or fluoro. In one embodiment R_{17} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{17} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{17} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In one embodiment, R_{17} is C_{1-4} alkyleneOC $_{1-4}$ alkyl. In one embodiment, R_{17} is C_{1-4} alkyleneOC $_{3-6}$ cycloalkyl. In one embodiment, R_{17} is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl).


35 Within such embodiments, R_{18} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyleneOC $_{1-4}$ alkyl, C_{1-4} alkyleneOC $_{3-6}$ cycloalkyl or C_{1-4} alkyleneO(4-10 membered heterocycloalkyl). In one embodiment R_{18} is H. In one embodiment R_{18} is halo, such as chloro or fluoro. In one embodiment R_{18} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one

embodiment, R_{18} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{18} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In one embodiment, R_{18} is C_{1-4} alkyleneOC₃₋₆cycloalkyl. In one embodiment, R_{18} is C_{1-4} alkyleneO(4-10 membered heterocycloalkyl).

- 5 Within such embodiments, R_{20} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy. In one embodiment R_{20} is H. In one embodiment R_{21} is halo, such as chloro or fluoro. In one embodiment R_{20} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In one embodiment, R_{20} is C_{1-4} haloalkyl, such as CF_3 . In one embodiment, R_{20} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

- 10 Compounds of the invention which bear functional groups capable of carrying charge at or around neutral or physiological pH (i.e. at or around pH 7.0 or 7.4) are believed to be capable of increasing the solubility of the compounds. Therefore in one embodiment at least one of R_2 and R_{2x} is a heterocycloalkyl or carboxylic acid containing group (e.g. R_2 is selected from 4-10 membered heterocycloalkyl, C_{1-4} alkylene(4-10 membered heterocycloalkyl), C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkylene($NR_{2xab}R_{2xac}$), OC_{1-4} alkylene($NR_{2xab}R_{2xac}$), $N(R_{2xaa})C_{1-4}$ alkylene($NR_{2xab}R_{2xac}$), C_{1-4} alkylene(CO_2H), such as $CH_2CH_2(CO_2H)$, OC_{1-4} alkylene(CO_2H), C_{1-4} alkyleneOC C_{1-4} alkylene(CO_2H) or $N(R_{2aa})C_{1-4}$ alkylene(CO_2H); when R_{2x} is a heterocycloalkyl or carboxylic acid containing group it will be understood that the corresponding R_{2x} groups are selected). In one embodiment at least one of R_{17} , R_{18} or R_{21} is a heterocycloalkyl or carboxylic acid containing group (e.g. R_{17} is 4-10 membered heterocycloalkyl, C_{1-4} alkylene(4-10 membered heterocycloalkyl), C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkylene($NR_{17ab}R_{17ac}$), OC_{1-4} alkylene($NR_{17ab}R_{17ac}$), $N(R_{17aa})C_{1-4}$ alkylene($NR_{17ab}R_{17ac}$), C_{1-4} alkylene(CO_2H), such as $CH_2CH_2(CO_2H)$, OC_{1-4} alkylene(CO_2H), C_{1-4} alkyleneOC C_{1-4} alkylene(CO_2H) or $N(R_{17a})C_{1-4}$ alkylene(CO_2H); when R_{18} or R_{21} is a heterocycloalkyl or carboxylic acid containing group it will be understood that the corresponding R_{18} or R_{21} groups are selected).

- Suitably when any of R_2 , R_{2x} , R_{17} , R_{18} or R_{21} are 4-10 membered heterocycle (whether alone or forming part of a larger groups e.g. C_{1-4} alkylene(4-10 membered heterocycloalkyl)), the 4-10 membered heterocycloalkyl may be selected from:



wherein:

Za is CH or N;

each R_{c1} is independently selected from H and C_{1-4} alkyl, such as methyl;

R_{c2} is independently selected from H and C_{1-4} alkyl, such as methyl;

R_{c3} is independently selected from H or C_{1-4} alkyl, such as methyl;

x_1 is 0, 1 or 2;

5 x_2 is 0, 1 or 2;

x_3 is 0, 1 or 2; and

x_4 is 0, 1 or 2.


Suitably, the 4-10 membered heterocycle contains at least one N atom e.g. one or two N atoms.

10

In one embodiment, the 4-10 membered heterocycle is (Ca). In one embodiment, the 4-10 membered heterocycle is (Cb). In one embodiment, the 4-10 membered heterocycle is (Cc). In one embodiment, the 4-10 membered heterocycle is (Cd).

15 In one embodiment, R_{17} is C_{1-4} alkyl, such as methyl, R_{20} is C_{1-4} alkoxy, such as OMe, R_{21} is H and R_{18} is C_{1-4} alkyl, such as methyl.

In one embodiment, group B is (Ba):



20 When Y is $C(R_{11})(R_{12})$, in one embodiment R_{11} is H. In a second embodiment, R_{11} is C_{1-4} alkyl, such as methyl.

When Y is $C(R_{11})(R_{12})$, in one embodiment R_{12} is H. In a second embodiment, R_{12} is C_{1-4} alkyl, such as methyl.

25

In a preferred embodiment, when Y is $C(R_{11})(R_{12})$, R_{11} is H or methyl and R_{12} is H. In a second preferred embodiment, R_{11} and R_{12} are both H. In a third embodiment, R_{11} and R_{12} together to the carbon atom to which they are attached form a cyclopropyl ring. Suitably, the cyclopropyl ring is substituted by 1, 2 or 3 C_{1-4} alkyl substituents, such as 1 or 2, e.g. 1 C_{1-4} alkyl substituent,

30

Suitably, when Y is $C(R_{11})(R_{12})$, R_{11} is methyl and R_{12} is H, A is not group (Aa).

In one embodiment, Y is N(R₁₃). Suitably, R₁₃ is C₁₋₄alkyl, such as methyl, ethyl, propyl or butyl, especially methyl. In a second embodiment, R₁₃ is C₃₋₆cycloalkyl, such as cyclopropyl. Suitably, the C₃₋₆cycloalkyl present in R₁₃ e.g. cyclopropyl ring is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 substituents independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋

5 4haloalkyl, halo, CN, OH, NR_{13a}R_{13b}, SO₂R_{13c} and NHSO₂R_{13c}. Suitably, the C₃₋₆cycloalkyl present in R₁₃ e.g. cyclopropyl ring is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 substituents independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo and CN. Suitably, the C₃₋₆cycloalkyl present in R₁₃ e.g. cyclopropyl ring is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 substituents independently selected from OH, NR_{13a}R_{13b}, SO₂R_{13c} and
10 NHSO₂R_{13c}. Suitably, the C₃₋₆cycloalkyl e.g. cyclopropyl ring is substituted by 1, 2 or 3 C₁₋₄alkyl substituents, such as 1 or 2 e.g. 1 C₁₋₄alkyl substituent, such as methyl. In one embodiment, the C₃₋₆cycloalkyl group e.g. cyclopropyl ring is not substituted.

In one embodiment R_{13a} is H. In a second embodiment R_{13a} is C₁₋₄alkyl, such as methyl, ethyl or
15 propyl, especially methyl. In a third embodiment R_{13a} is H or methyl.

In one embodiment R_{13b} is H. In a second embodiment R_{13b} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{13b} is C₃₋₆cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{13b} is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{13b} is C₁₋₄haloalkyl, such as CF₃.
20 In a sixth embodiment, R_{13b} is aryl, such as phenyl. In a seventh embodiment, R_{13b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment, R_{13b} is H or methyl.

25 In one embodiment R_{13c} is C₁₋₄alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{13c} is C₃₋₆cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{13c} is C₁₋₄alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{13c} is C₁₋₄haloalkyl, such as CF₃. In a fifth embodiment, R_{13c} is aryl, such as phenyl. In a sixth embodiment, R_{13c} is 4-10 membered heterocycloalkyl, such as
30 azetidiny or oxetanyl. In a seventh embodiment R_{13c} is H or methyl.


In one embodiment, Y is O or S.

When present, in one embodiment each R₁₀ is independently fluoro, chloro or methyl.

35

In one embodiment, r is 0 or 1, especially 0.


When r is 1, 2 or 3, R_{10} may be in the 7-, 5- and/or 4-position(s). In one embodiment, r is 1 and R_{10} is in the 7-position. In a second embodiment, r is 1 and R_{10} is in the 5-position. In one embodiment, r is 1 and R_{10} is in the 4-position. For example:



5

Examples of suitable R_{10} substituents include 4-fluoro and 7-fluoro.

In one embodiment, group B is (Bb):




10

When present, in one embodiment each R_{14} is independently fluoro or methyl.

In one embodiment, s is 0 or 1, especially 0.

- 15 Suitably, when s is 1, 2 or 3, R_{14} may be in the 7-, 5- and/or 4-position(s). In one embodiment, s is 1 and R_{14} is in the 5-position. In a second embodiment, s is 2, one R_{14} is in the 7-position and one R_{14} is in the 4-position, for example:




Suitably, R_{14} is not in the 7-position. Suitably, R_{14} is not in the 4-position.

20

An example of a suitable R_{14} substituent is 5-fluoro.

In one embodiment, group B is (Bc):



R_{15} is C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-4} haloalkyl, halo or CN; wherein said cycloalkyl is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR^{15a}R^{15b}$, SO_2R^{15c} and $NHSO_2R^{15c}$.

In one embodiment, R_{15} is methyl, ethyl, cyclopropyl, CF_3 or CN, for example methyl or CN, especially methyl. In a second embodiment, R_{15} is methyl, ethyl, cyclopropyl, CF_3 , CN, OMe, chloro or fluoro, for example methyl, CN, chloro or fluoro, especially chloro or fluoro. In a third embodiment, R_{15} is OMe, chloro or fluoro. In a fourth embodiment, R_{15} is methyl, CN, chloro or fluoro.

Suitably, the C_{3-6} cycloalkyl e.g. cyclopropyl ring present in R_{15} is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR^{15a}R^{15b}$, SO_2R^{15c} and $NHSO_2R^{15c}$. Suitably, the C_{3-6} cycloalkyl present in R_{15} e.g. cyclopropyl ring is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo and CN. Suitably, the C_{3-6} cycloalkyl present in R_{15} e.g. cyclopropyl ring is substituted by 1, 2 or 3, such as 1 or 2 e.g. 1 substituents independently selected from OH, $NR^{15a}R^{15b}$, SO_2R^{15c} and $NHSO_2R^{15c}$. Suitably, the C_{3-6} cycloalkyl e.g. cyclopropyl ring is substituted by 1, 2 or 3 C_{1-4} alkyl substituents, such as 1 or 2 e.g. 1 C_{1-4} alkyl substituent, such as methyl. In one embodiment, the C_{3-6} cycloalkyl group e.g. cyclopropyl ring is not substituted.

In one embodiment R_{15a} is H. In a second embodiment R_{15a} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{15a} is H or methyl.

In one embodiment R_{15b} is H. In a second embodiment R_{15b} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a third embodiment, R_{15b} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a fourth embodiment, R_{15b} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fifth embodiment, R_{15b} is C_{1-4} haloalkyl, such as CF_3 . In a sixth embodiment, R_{15b} is aryl, such as phenyl. In a seventh embodiment, R_{15b} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In an eighth embodiment, R_{15b} is H or methyl.

In one embodiment R_{15c} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl. In a second embodiment, R_{15c} is C_{3-6} cycloalkyl, such as cyclopropyl, cyclobutyl and cyclopentyl, especially cyclopropyl. In a third embodiment, R_{15c} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe. In a fourth embodiment, R_{15c} is C_{1-4} haloalkyl, such as CF_3 . In a fifth embodiment, R_{15c} is aryl, such as phenyl. In a sixth embodiment, R_{15c} is 4-10 membered heterocycloalkyl, such as azetidiny or oxetanyl. In a seventh embodiment, R_{15c} is methyl.

In one embodiment, D, E and F are $C(R_{16})$. In a second embodiment, D is N, and E and F are $C(R_{16})$. In a third embodiment, E is N, and D and F are $C(R_{16})$. In a fourth embodiment, F is N, and D and E are $C(R_{16})$.

In one embodiment, R_{16} is H. In a second embodiment, R_{16} is halo, such as fluoro or chloro, especially chloro. In a third embodiment, R_{16} is C_{1-4} alkyl, such as methyl. Suitably each R_{16} is independently H, fluoro, chloro or methyl.

In one preferred embodiment, A is group (Aa) and B is group (Ba). In a second preferred embodiment, A is group (Aa) and B is group (Bb). In a third preferred embodiment, A is group (Ad) and B is group (Bc). In a fourth preferred embodiment, A is group (Ae) and B is group (Ba). In a fifth preferred embodiment, A is group (Ae) and B is group (Bb). In a sixth preferred embodiment, A is group (Ae) and B is group (Bc).

In one embodiment, the compound of formula (I) is selected from the group consisting of:

- (E)-3-(1H-benzo[d][1,2,3]triazol-6-yl)-N-(3-fluoro-2-methylphenyl) acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]thiazol-5-yl)acrylamide;
- (E)-3-(3,3-dimethyl-2-oxoindolin-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(2'-oxospiro[cyclopropane-1,3'-indolin]-6'-yl)acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(7-fluoro-2-oxoindolin-6-yl)acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl) acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(3-methyl-2-oxoindolin-6-yl)acrylamide;
- (E)-N-(3-chloro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
- (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
- (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
- (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)-N-(o-tolyl)acrylamide;
- (E)-N-(2-isopropylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
- (E)-N-(2-isopropyl-6-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;

- (E)-*N*-(5-chloro-2-isopropylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
(E)-*N*-(4,5-difluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
(E)-*N*-(5-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
(E)-*N*-(3-fluoro-2-methylphenyl)-3-(4-fluoro-2-oxoindolin-6-yl)acrylamide;
5 (E)-*N*-(2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
(E)-*N*-(3-fluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
(E)-*N*-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide;
(E)-3-(1-ethyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-*N*-(3-fluoro-2-methylphenyl)
acrylamide;
10 (E)-3-(1-cyclopropyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-*N*-(3-fluoro-2-methylphenyl)
acrylamide;
(E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)
acrylamide;
(E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
15 (E)-3-(3-cyano-1H-indazol-6-yl)-*N*-(2,3-dihydro-1H-inden-1-yl)acrylamide;
(E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(5-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide;
(E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(3-(trifluoromethyl)-1H-indazol-6-yl)acrylamide;
(E)-*N*-(2,6-dimethylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide;
(E)-*N*-(3-fluoro-2,6-dimethylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-
20 yl)acrylamide;
(E)-*N*-(2,6-dimethylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
(E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(3-ethyl-1H-indazol-6-yl)acrylamide;
(E)-3-(3-cyclopropyl-1H-indazol-6-yl)-*N*-(2,3-dihydro-1H-inden-1-yl)acrylamide;
(E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(4-fluoro-3-methyl-1H-indazol-6-yl)acrylamide;
25 (E)-*N*-(3,5-difluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
(E)-*N*-(3,4-difluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
(E)-*N*-(3-fluoro-2-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
(E)-3-(3-methyl-1H-indazol-6-yl)-*N*-(2-methyl-2,3-dihydro-1H-inden-1-yl)acrylamide;
(E)-3-(3-methyl-1H-indazol-6-yl)-*N*-(1-methyl-1H-indazol-7-yl)acrylamide;
30 (E)-*N*-(5-fluoro-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
(E)-*N*-(4-fluoro-3-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
(E)-*N*-(3-fluoro-4-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
Racemic-(E)-3-(3-methyl-1H-indazol-6-yl)-*N*-((1*R*,2*R*)-2-methylcyclohexyl)acrylamide;
(E)-3-(3-cyano-1H-indazol-6-yl)-*N*-(2-methyl-2,3-dihydro-1H-inden-1-yl)acrylamide;
35 (E)-*N*-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-
5-yl)acrylamide;
(E)-*N*-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;

- (Z)-2-fluoro-N-(3-fluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
 (E)-N-(3-chloro-2-methylphenyl)-N-methyl-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
 (E)-N-(2-methylcyclopentyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
 (E)-N-(3-fluoro-2-(methoxymethyl)phenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 5 (E)-3-(3-cyano-1H-indazol-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide;
 (E)-3-(3-methyl-1H-indazol-6-yl)-N-(3-methylchroman-4-yl)acrylamide;
 (E)-N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 (E)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 (R,E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 10 (E)-N-(chroman-4-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide;
 (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)acrylamide;
 (E)-N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-(2-oxoindolin-6-yl)acrylamide;
 (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide;
 (E)-N-(3,5-difluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide;
 15 (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(7-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide;
 (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(4-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide;
 (E)-N-(5-chloro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide; and
 (E)-N-(2-methylcyclohexyl)-3-(2-oxoindolin-6-yl)acrylamide;
 or a pharmaceutically acceptable salt and/or solvate of any one thereof.

- 20 In one embodiment, the compound of formula (I) is selected from the group consisting of:
 (E)-3-(3-methyl-1H-indazol-6-yl)-N-(3-methylchroman-4-yl)acrylamide;
 (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3R,4S)-3-methylchroman-4-yl)acrylamide;
 (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3S,4R)-3-methylchroman-4-yl)acrylamide;
 25 (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3R,4R)-3-methylchroman-4-yl)acrylamide;
 (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3S,4S)-3-methylchroman-4-yl)acrylamide;
 (E)-3-(3-methyl-1H-indazol-6-yl)-N-((1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-yl)acrylamide;
 (E)-N-((1S,2S)-2-(cyclopropylmethoxy)-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 30 (E)-N-((1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 (E)-N-((1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
 (E)-3-(3-cyclopropyl-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide;
 35 (E)-3-(3-methoxy-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide;
 (E)-3-(3-chloro-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide;

(E)-3-(3-fluoro-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide;
and
(E)-3-(3-cyano-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide;
or a pharmaceutically acceptable salt and/or solvate of any one thereof.

5

In one embodiment, the compound of formula (I) is selected from the group consisting of:

- (E)-3-(3-fluoro-4-methyl-5-(3-(3-methyl-1H-indazol-6-yl)acrylamido)phenyl)propanoic acid;
(E)-3-(3-(3-(3-chloro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(3-fluoro-5-(3-(3-fluoro-1H-indazol-6-yl)acrylamido)-4-methylphenyl)propanoic acid;
10 (E)-3-(3-(3-(3-cyano-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(3-fluoro-5-(3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamido)-4-methylphenyl)propanoic acid;
(E)-3-(3-(3-(3-chloro-5-fluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(3-(3-(3,5-difluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
15 (E)-3-(3-(3-(3-cyano-5-fluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(4-fluoro-3-methyl-2-(3-(3-methyl-1H-indazol-6-yl)acrylamido)phenyl)propanoic acid;
(E)-3-(2-(3-(3-chloro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
20 (E)-3-(4-fluoro-2-(3-(3-fluoro-1H-indazol-6-yl)acrylamido)-3-methylphenyl)propanoic acid;
(E)-3-(2-(3-(3-cyano-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-3-(4-fluoro-2-(3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamido)-3-methylphenyl)propanoic acid;
(E)-3-(2-(3-(3-chloro-5-fluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
25 acid;
(E)-3-(2-(3-(3,5-difluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-3-(2-(3-(3-cyano-5-fluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
30 (E)-3-(3-chloro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
35 (E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;

(E)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;

(E)-3-(3-chloro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

5 (E)-3-(3-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

(E)-3-(3-cyano-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

10 (E)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamide;

(E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

(E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide; and

15 (E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

or a pharmaceutically acceptable salt and/or solvate of any one thereof.

In one embodiment, the compound of formula (I) is selected from the group consisting of:

20 (2E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;
 (2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl) prop-2-enamide;
 (2E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;
 (2E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

25 (2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

30 (2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl) prop-2-enamide;

(2E)-N-(5-fluoro-4-methylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide;

35 (2E)-3-(3-chloro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3-methyl-1H-indazol-6-yl)-N-(2-methylpyridin-3-yl)prop-2-enamide;
(2E)-N-(2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;
(2E)-N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

5 or a pharmaceutically acceptable salt and/or solvate of any one thereof.

The definition of the compounds of formula (I) is intended to include all tautomers of said compounds.

10 The compounds of the invention may be provided in the form of a pharmaceutically acceptable salt and/or solvate thereof. In particular, the compound of formula (I) may be provided in the form of a pharmaceutically acceptable salt and/or solvate, such as a pharmaceutically acceptable salt.

15 It will be appreciated that for use in medicine the salts of the compounds of formula (I) should be pharmaceutically acceptable. Non-pharmaceutically acceptable salts of the compounds of formula (I) may be of use in other contexts such as during preparation of the compounds of formula (I). Suitable pharmaceutically acceptable salts will be apparent to those skilled in the art. Pharmaceutically acceptable salts include those described by Berge et al. (1977). Such
20 pharmaceutically acceptable salts include acid and base addition salts. Pharmaceutically acceptable acid additional salts may be formed with inorganic acids e.g. hydrochloric, hydrobromic, sulphuric, nitric or phosphoric acid and organic acids e.g. succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, p-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid. Other salts e.g. oxalates or formates, may be used, for example in the isolation of compounds
25 of formula (I) and are included within the scope of this invention. Pharmaceutically acceptable salts may also be formed with organic bases such as basic amines e.g. with ammonia, meglumine, tromethamine, piperazine, arginine, choline, diethylamine, benzathine or lysine. Pharmaceutically acceptable salts may also be formed with inorganic bases such as group 1 or 2 metal ions e.g. lithium, sodium, potassium, magnesium or calcium.

30 Certain compounds of formula (I) may form acid or base addition salts with one or more equivalents of the acid or base. The present invention includes within its scope all possible stoichiometric and non-stoichiometric forms. In one embodiment the compound of formula (I) is the free base form. Alternatively, there is provided a compound of formula (I) in the form of a free
35 acid. When the compound contains a basic group as well as the free acid it may be Zwitterionic.

The compounds of formula (I) may be prepared in crystalline or non-crystalline form and, if crystalline, may optionally be solvated, e.g. as the hydrate. This invention includes within its scope stoichiometric solvates (e.g. hydrates) as well as compounds containing variable amounts of solvent (e.g. water).

5

It is to be understood that the present invention encompasses all isomers of formula (I) and their pharmaceutically acceptable derivatives, including all geometric, tautomeric and optical forms, and mixtures thereof (e.g. racemic mixtures). Where additional chiral centres are present in compounds of formula (I), the present invention includes within its scope all possible

10 diastereoisomers, including mixtures thereof. The different isomeric forms may be separated or resolved one from the other by conventional methods, or any given isomer may be obtained by conventional synthetic methods or by stereospecific or asymmetric syntheses.

The present disclosure includes all isotopic forms of the compounds of the invention provided
15 herein, whether in a form (i) wherein all atoms of a given atomic number have a mass number (or mixture of mass numbers) which predominates in nature (referred to herein as the "natural isotopic form") or (ii) wherein one or more atoms are replaced by atoms having the same atomic number, but a mass number different from the mass number of atoms which predominates in nature (referred to herein as an "unnatural variant isotopic form"). It is understood that an atom
20 may naturally exist as a mixture of mass numbers. The term "unnatural variant isotopic form" also includes embodiments in which the proportion of an atom of given atomic number having a mass number found less commonly in nature (referred to herein as an "uncommon isotope") has been increased relative to that which is naturally occurring e.g. to the level of >20%, >50%, >75%, >90%, >95% or >99% by number of the atoms of that atomic number (the latter
25 embodiment referred to as an "isotopically enriched variant form"). The term "unnatural variant isotopic form" also includes embodiments in which the proportion of an uncommon isotope has been reduced relative to that which is naturally occurring. Isotopic forms may include radioactive forms (i.e. they incorporate radioisotopes) and non-radioactive forms. Radioactive forms will typically be isotopically enriched variant forms.

30

An unnatural variant isotopic form of a compound may thus contain one or more artificial or uncommon isotopes such as deuterium (^2H or D), carbon-11 (^{11}C), carbon-13 (^{13}C), carbon-14 (^{14}C), nitrogen-13 (^{13}N), nitrogen-15 (^{15}N), oxygen-15 (^{15}O), oxygen-17 (^{17}O), oxygen-18 (^{18}O), phosphorus-32 (^{32}P), sulphur-35 (^{35}S), chlorine-36 (^{36}Cl), chlorine-37 (^{37}Cl), fluorine-18 (^{18}F)
35 iodine-123 (^{123}I), iodine-125 (^{125}I) in one or more atoms or may contain an increased proportion of said isotopes as compared with the proportion that predominates in nature in one or more atoms.

Unnatural variant isotopic forms comprising radioisotopes may, for example, be used for drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, i.e. ^3H , and carbon-14, i.e. ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection. Unnatural variant isotopic forms which incorporate deuterium i.e. ^2H or D may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances. Further, unnatural variant isotopic forms may be prepared which incorporate positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , and would be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. In one embodiment, the compounds of the invention are provided in a natural isotopic form. In one embodiment, the compounds of the invention are provided in an unnatural variant isotopic form. In a specific embodiment, the unnatural variant isotopic form is a form in which deuterium (i.e. ^2H or D) is incorporated where hydrogen is specified in the chemical structure in one or more atoms of a compound of the invention. In one embodiment, the atoms of the compounds of the invention are in an isotopic form which is not radioactive. In one embodiment, one or more atoms of the compounds of the invention are in an isotopic form which is radioactive. Suitably radioactive isotopes are stable isotopes. Suitably the unnatural variant isotopic form is a pharmaceutically acceptable form.

In one embodiment, a compound of the invention is provided whereby a single atom of the compound exists in an unnatural variant isotopic form. In another embodiment, a compound of the invention is provided whereby two or more atoms exist in an unnatural variant isotopic form. Unnatural isotopic variant forms can generally be prepared by conventional techniques known to those skilled in the art or by processes described herein e.g. processes analogous to those described in the accompanying Examples for preparing natural isotopic forms. Thus, unnatural isotopic variant forms could be prepared by using appropriate isotopically variant (or labelled) reagents in place of the normal reagents employed in the Examples. Since the compounds of formula (I) are intended for use in pharmaceutical compositions it will readily be understood that they are each preferably provided in substantially pure form, for example at least 60% pure, more suitably at least 75% pure and preferably at least 85%, especially at least 98% pure (% are on a weight for weight basis). Impure preparations of the compounds may be used for preparing the more pure forms used in the pharmaceutical compositions.


In general, the compounds of formula (I) may be made according to the organic synthesis techniques known to those skilled in this field, as well as by the representative methods set forth below, those in the Examples, and modifications thereof. In the following schemes, reactive

groups can be protected with protecting groups and deprotected according to established techniques well known to the skilled person.

Generic Routes


- 5 Generic routes by which compound examples of the invention may be conveniently prepared are summarised below. In the following description, the groups R_{1a} , R_{1b} , R_2 , R_3 , R_{13} , A and B are as defined above for the compound of formula (I), unless otherwise stated.

Scheme 1




- 10 Compounds of formula (I) may be prepared by reacting a compound of formula (II) with a compound of formula (III) under palladium-catalyzed cross coupling conditions, using a palladium pre-catalyst, such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane ($\text{Pd}(\text{dppf})\text{Cl}_2\cdot\text{CH}_2\text{Cl}_2$), in the presence of a base, such as triethylamine, and a suitable solvent, such as dimethylformamide (DMF).

15 Scheme 2




- Alternatively, compounds of formula (I) may be prepared by reacting a compound of formula (IV) with a compound of formula (V) under amidation conditions, using an amide coupling reagent, such as 1-[bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate (HATU), in the presence of a base, such as *N,N*-diisopropylethylamine (DIPEA, also known as Hünig's base), in a suitable solvent, such as DMF.
- 20

Scheme 3




Compounds of formula (I) may be also prepared by reacting a compound of formula (VI) with a compound of (V) under basic conditions, using a base such as lithium bis(trimethylsilyl)amide (LiHMDS), in a suitable solvent, such as tetrahydrofuran (THF).

Scheme 4




Compounds of formula (II) are commercially available. Compounds of formula (II) may also be prepared by reacting a compound of formula (VII) with a compound of formula (VIII) in the presence of a base, such as *N,N*-diisopropylethylamine, in a suitable solvent, such as dichloromethane (DCM).

Scheme 5




Compounds of formula (VI) may be obtained by reacting a compound of formula (IX) with a compound of formula (III) under palladium-catalyzed cross coupling conditions, using a palladium-precatalyst, such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (Pd(dppf)Cl₂CH₂Cl₂), in the presence of a base, such as triethylamine, and a suitable solvent, such as dimethylformamide (DMF).

Scheme 6




Compounds of formula (IV) may be obtained by reacting a compound of formula (VI) under hydrolysis conditions, using a base, such as sodium hydroxide (NaOH), in a suitable solvent system, such as a mixture of methanol and water.

Scheme 7



Compounds of formula (VII), wherein R_{1a} is H, A is group (Aa), R₂ is C₁₋₄alkyl, m is 3 and R₃ is C₁₋₄alkyl or halo, may be prepared in two steps. Firstly, reacting a compound of formula (X) with a brominating agent, such as *N*-bromosuccinimide (NBS), in a solvent, such as acetonitrile, affords di-brominated compounds of formula (XI). The compound of formula (VII) may be further reacted under palladium-catalyzed cross coupling conditions, using a palladium-precatalyst, such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (Pd(dppf)Cl₂·CH₂Cl₂), an organoboron compound, such as trimethyl-1,3,5,2,4,6-trioxatriborinane, a base, such as caesium carbonate (Cs₂CO₃), and a suitable solvent, such as 1,4-dioxane, to give a compound of formula (VII).


Scheme 8



Compounds of formula (VII), wherein R_{1a} is H, A is group (Aa), R₂ is C₁₋₄alkyl, m is 3 and R₃ is C₁₋₄alkyl or halo, may be prepared in four steps. Firstly, reacting a compound of formula (XII) with a chlorinating agent, such as *N*-chlorosuccinimide (NCS), in a solvent, such as acetonitrile, affords a compound of formula (XIII). Further reacting a compound of formula (XIII) with a brominating agent, such as *N*-bromosuccinimide (NBS), in a solvent, such as acetonitrile, affords tri-halogenated compounds of formula (XIV). The compound of formula (XIV) may be further reacted under palladium-catalyzed cross coupling conditions, using a palladium-precatalyst, such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (Pd(dppf)Cl₂·CH₂Cl₂), an organoboron compound, such as trimethyl-


1,3,5,2,4,6-trioxatriborinane, a base, such as caesium carbonate (Cs_2CO_3), and a suitable solvent, such as 1,4-dioxane, to give a compound of formula (XV). The compound of formula may be reacted under reductive dehalogenation conditions, using a hydrogen atmosphere, such as hydrogen at 30 atm, a palladium catalyst, such as palladium on carbon (Pd/C), an acid, such as hydrochloric acid, and a solvent, such as ethanol (EtOH), to afford the compound of formula (VII).

Scheme 9



Compounds of formula (VII), wherein R_{1a} is H, A is group (Aa), R_2 is C_{1-4} alkyl, m is 2 and R_3 is C_{1-4} alkyl or halo, may be prepared in two steps. Nitro compounds of formula (XVI) may be reacted under palladium-catalyzed cross coupling conditions, using a palladium-precatalyst, such as [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane ($\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$), an organoboron compound, such as trimethyl-1,3,5,2,4,6-trioxatriborinane, a base, such as caesium carbonate (Cs_2CO_3), and a suitable solvent system, such as a mixture of water 1,4-dioxane, to give a compound of formula (XVII). The compound of formula (XVII) may be further reacted with a metal, such as iron, in the presence of an acid, such as acetic acid, to give the compound of formula (VII).


Scheme 10



Compounds of formula (III), wherein B is group (Ba), Y is $\text{N}(\text{R}_{13})$ and r is 0, may be prepared in three steps. Firstly, a compound of formula (XVIII) may be reacted with an amine in a suitable solvent, such as ethanol, to give a compound of formula (XIX). Subsequently, the compound of formula (XIX) may be subjected to reduction conditions using a metal, such as zinc, and an


inorganic salt, such as ammonium chloride (NH_4Cl), in a suitable solvent, such as acetone, to afford a compound of formula (XX). Finally, the compound of formula (XX) may be reacted with a carbonylating agent, such as triphosgene, in a suitable solvent, such as dichloromethane (DCM), to afford the compound of formula (III).

5

Scheme 11

Compounds of formula (V) wherein A is group (Ad), X is a bond, q is 0, p is 1 and R_8 is $\text{OC}_{1-4}\text{haloalkyl}$, $\text{OC}_{1-4}\text{alkylene(4-10 membered heterocycloalkyl)}$ or $\text{C}_{1-4}\text{alkoxy}$ may be prepared by reacting a compound of formula (XXI), wherein PG is a nitrogen protecting group, such as *tert*-butoxycarbonyl (BOC), with a compound of formula (XXII), wherein C is $\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-4}\text{haloalkyl}$ or $\text{C}_{1-4}\text{alkylene(4-10 membered heterocycloalkyl)}$, in the presence of a base e.g. NaH in a suitable solvent e.g. THF, to afford compounds of formula (XXIII). Deprotection of the compounds of formula (XXIII) e.g. with aqueous acid, such as 2 M HCl, in a suitable solvent, such as methanol affords compounds of formula (V).


15

Scheme 12

Compounds of formula (V) wherein A is group (Ad), X is a bond, q is 0, p is 1 and R_8 is $\text{OC}_{1-4}\text{alkylene(C}_{3-6}\text{cycloalkyl)}$ may be prepared by treating a compound of formula (XXIV), wherein PG is a nitrogen protecting group, such as *tert*-butoxycarbonyl (BOC), with a suitable reagent e.g. di-iodomethane, in the presence of an organometallic reagent e.g. ZnEt_2 in a suitable solvent, such as dichloromethane, to afford compounds of formula (XXV). Deprotection of the compounds of formula (XXV) e.g. with aqueous acid, such as 2 M HCl, in a suitable solvent, such as methanol affords compounds of formula (V).

25


Scheme 13



Compounds of formula (II) wherein one of R_2 or R_{2x} is C_{1-4} alkylene(CO_2H) (e.g. R_{2x} is C_{1-4} alkylene as shown in Scheme 13) may be prepared by reacting a compound of formula (XXX)

with a compound of formula (XXXI) under palladium catalysed cross-coupling conditions, using a palladium pre-catalyst such as $Pd(OAc)_2$, a phosphine ligand such as triphenylphosphine, an inorganic base such as $NaHCO_3$, an organic base such as triethylamine, in a solvent such as DMF to afford compounds of formula (XXXII). Treatment of nitro compounds (XXXII) under hydrogenation conditions, using a palladium catalyst such as palladium on carbon (Pd/C) and a solvent such as ethanol, affords compounds of formula (XXXIII). Compounds of formula (XXXIII) may be reacted with compounds of formula (XXXIV) in the presence of an organic base e.g. triethylamine to afford compounds of formula (II). Compounds of formula (I) may be prepared from the compounds of formula (II) using methods known to the skilled person, and/or the methods disclosed herein. The *tert*-butyl ester of compounds of formulae (XXX), (XXXII), (XXXIII) and (II) may be removed by methods known to the skilled person, such as treatment with trifluoroacetic acid in dichloromethane to afford the corresponding carboxylic acids.

Scheme 14




Compounds of formula (II) wherein one of R_2 or R_{2x} is 4-7 membered heterocycloalkyl (e.g. R_2 is 4-7 membered heterocycloalkyl as shown in Scheme 14) may be prepared by reacting a compound of formula (XXXV) with a compound of formula (XXXVI) under palladium catalysed

cross-coupling conditions, using a palladium precatalyst such as Pd(OAc)₂, a phosphine ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP), an inorganic base such as sodium butoxide and a solvent such as toluene. Subsequent reduction of the nitro group under


- 5 compounds of formula (XXXVII). Compounds of formula (XXXVII) may be reacted with compounds of formula (XXXIV) in the presence of an organic base e.g. triethylamine to afford compounds of formula (II). Compounds of formula (I) may be prepared from the compounds of formula (II) using methods known to the skilled person, and/or the methods disclosed herein.

- 10 Therefore, in one embodiment the invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt and/or solvate thereof, which comprises reacting a compound of formula (II):




wherein A, R_{1a} and R_{1b} are as defined for the compound of formula (I); or salt thereof;

- 15 with a compound of formula (III):




wherein X is halo, such as bromo or iodo, and B is as defined for the compound of formula (I); or a salt thereof.

- 20 The invention also provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt and/or solvate thereof, which comprises reacting a compound of formula (IV):




wherein R_{1b} and B are as defined for the compound of formula (I); or a salt thereof;

- 25 with a compound of formula (V):




wherein A and R_{1a} are as defined for the compound of formula (I); or a salt thereof.

The invention also provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt and/or solvate thereof, which comprises reacting a compound of formula (VI):




wherein R_{1b} and B are as defined for the compound of formula (I); or a salt thereof; with a compound of formula (V):




wherein A and R_{1a} are as defined for the compound of formula (I); or a salt thereof.

In one embodiment, there is provided a compound of formula (II):




or a salt thereof, wherein A, R_{1a} , and R_{1b} are as defined elsewhere herein.

In one embodiment, there is provided a compound of formula (IV):



or a salt thereof, wherein B and R_{1b} are as defined elsewhere herein.

In one embodiment, there is provided a compound of formula (VI):



or a salt thereof, wherein B and R_{1b} are as defined elsewhere herein.

Novel compounds of formulae (II) to (XXXVII) and salts thereof are also provided as an aspect
5 of invention

Therapeutic Methods

Compounds of formula (I) of the present invention have utility as inhibitors of mPTP.

- 10 Therefore, the invention provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use as a pharmaceutical, in particular in the treatment or prophylaxis of a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect, for example those diseases and disorders mentioned herein below.
- 15 The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use as a pharmaceutical, in particular in the treatment of a disease or disorder in which inhibition of mPTP provides a therapeutic effect, for example those diseases and disorders mentioned herein below.
- 20 The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use as a pharmaceutical, in particular in the prophylaxis of a disease or disorder in which inhibition of mPTP provides a prophylactic effect, for example those diseases and disorders mentioned herein below.
- 25 The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect, for example those diseases and disorders mentioned herein below.
- 30 The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment of a disease or disorder in which inhibition of mPTP provides a therapeutic effect, for example those diseases and disorders mentioned herein below.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the prophylaxis of a disease or disorder in which inhibition of mPTP provides a prophylactic effect, for example those diseases and disorders mentioned herein below.

5

The invention also provides a method of preventing or treating a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

10

The invention also provides a method of treating a disease or disorder in which inhibition of mPTP provides a therapeutic effect in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

15

The invention also provides a method of preventing a disease or disorder in which inhibition of mPTP provides a prophylactic effect in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

20

The term 'treatment' or 'treating' as used herein includes the control, mitigation, reduction, or modulation of the disease state or its symptoms.

25

The term 'prophylaxis' or 'preventing' is used herein to mean preventing symptoms of a disease or disorder in a subject or preventing recurrence of symptoms of a disease or disorder in an afflicted subject and is not limited to complete prevention of an affliction.

30

In one embodiment, the disease or disorder is selected from degenerative or neurodegenerative diseases, disorders of the central nervous system, ischemia or re-perfusion injury, metabolic diseases, inflammatory or autoimmune diseases, diseases of aging and renal diseases.

35

In one particular embodiment, the disease or disorder is a degenerative or neurodegenerative disease, such as Parkinson's disease, dementia with Lewy bodies, Alzheimer's disease, amyotrophic lateral sclerosis, multiple sclerosis, frontal temporal dementia, chemotherapy

induced neuropathy, Huntington's disease, spinocerebellar ataxias, progressive supranuclear palsy, hereditary spastic paraplegia, Duchenne muscular dystrophy, congenital muscular dystrophy, traumatic brain injury and Friedreich's ataxia. In one preferred embodiment, the disease or disorder is Parkinson's disease. In one preferred embodiment, the disease or disorder is Alzheimer's disease. In one preferred embodiment, the disease or disorder is amyotrophic lateral sclerosis.

In another particular embodiment, the disease or disorder is a disease of the central nervous system, such as AIDS dementia complex, depressive disorders, schizophrenia and epilepsy.

In another embodiment, the disease or disorder is ischemia or re-perfusion injury, such as acute myocardial infarction, stroke, kidney ischemia reperfusion injury, and organ damage during transplantation.

In another embodiment, the disease or disorder is a metabolic disease, such as hepatic steatosis, diabetes, diabetic retinopathy, cognitive decline and other diabetes associated conditions, obesity and feeding behaviours, and non-alcoholic fatty liver disease.

In another embodiment, the disease or disorder is an inflammatory or autoimmune disease, such as acute pancreatitis, systemic lupus, organ failure in sepsis and hepatitis. In another embodiment the disease or disorder is a disease associated with mtDNA release and innate immune activation e.g. macular degeneration.

In another embodiment, the disease or disorder is a disease of aging, such as bone repair, bone weakness in aging in osteoporosis and sarcopenia.

In another embodiment, the disease or disorder is a renal disease, such as chronic kidney disease associated with APOL1 genetic variants and chronic kidney disease.

In another embodiment, the disease or disorder is a pulmonary dysfunction or multi-organ failure associated with severe respiratory virus infections e.g. COVID19 infection. In another embodiment, the disease or disorder is adrenoleukodystrophy, in particular X-linked adrenoleukodystrophy.

The compounds of formula (I) are expected to be useful in the treatment or prophylaxis of a mitochondrial disease.

Therefore, the invention provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a mitochondrial disease, for example those diseases and disorders mentioned herein below.

- 5 The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment of a mitochondrial disease, for example those diseases and disorders mentioned herein below.

- 10 The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the prophylaxis of a mitochondrial disease, for example those diseases and disorders mentioned herein below.

- 15 The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a mitochondrial disease, for example those diseases and disorders mentioned herein below.

- 20 The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment of a mitochondrial disease, for example those diseases and disorders mentioned herein below.

- The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the prophylaxis of a mitochondrial disease, for example those diseases and disorders mentioned herein below.

- 25 The invention also provides a method of treating or preventing a mitochondrial disease in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

- 30 The invention also provides a method of treating a mitochondrial disease in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

- 35 The invention also provides a method of preventing a mitochondrial disease in a subject, which comprises administering to a subject in need thereof an effective amount of a compound of

formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

Suitably, the mitochondrial disease is selected from Reye syndrome, Leber's hereditary optic neuropathy and associated disorders and disorders, such as those diseases and disorders disclosed in CA2884607A1 (Stealth Peptides International Inc.)

The compounds of formula (I) are expected to be useful in the treatment or prophylaxis of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration.

Therefore, the invention provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, for example those diseases and disorders mentioned herein below.

The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, for example those diseases and disorders mentioned herein below.

The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the prophylaxis of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, for example those diseases and disorders mentioned herein below.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, for example those diseases and disorders mentioned herein below.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, for example those diseases and disorders mentioned herein below.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the prophylaxis of a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, for example those diseases and disorders mentioned herein below.

5

The invention also provides a method of treating or preventing a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

10

The invention also provides a method of treating a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

15

The invention also provides a method of preventing a disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for example those diseases and disorders mentioned herein below.

20

Suitably, the disease or disorder associated with TDP-43 proteinopathy such as TDP-43 associated neurodegeneration is selected from Amyotrophic Lateral Sclerosis, Frontotemporal dementia, Facial onset sensory and motor neuronopathy, Primary lateral sclerosis, Progressive muscular atrophy, Inclusion body myopathy associated with early-onset Paget disease of the bone and Frontotemporal lobar degeneration dementia, Perry disease, Chronic traumatic encephalopathy, Severe traumatic brain injury, Alzheimer's disease, Hippocampal sclerosis dementia, Limbic-predominant age-related TDP-43 encephalopathy, and Cerebral age-related TDP-43 with sclerosis.

25

30

The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment or prophylaxis of a disease or disorder associated with fibrosis.

35

The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the treatment of a disease or disorder associated with fibrosis.

The invention also provides a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, for use in the prophylaxis of a disease or disorder associated with fibrosis.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment or prophylaxis of a disease or disorder associated with fibrosis.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the treatment of a disease or disorder associated with fibrosis.

The invention also provides use of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof, in the manufacture of a medicament for the prophylaxis of a disease or disorder associated with fibrosis.

The invention also provides a method of treating or preventing a disease or disorder associated with fibrosis, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof.

The invention also provides a method of treating a disease or disorder associated with fibrosis, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof.

The invention also provides a method of preventing a disease or disorder associated with fibrosis, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof.

Suitably, the disease or disorder associated with fibrosis is selected from chronic kidney disease, idiopathic pulmonary fibrosis, non-alcoholic steatohepatitis, primary biliary cholangitis and systemic sclerosis.

Suitably the subject is a mammal, in particular the subject is a human.

Pharmaceutical Compositions

For use in therapy the compounds of the invention are usually administered as a pharmaceutical composition. The invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof, and a pharmaceutically acceptable carrier or excipient.

In one embodiment, there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof, for use in the treatment or prophylaxis of a disease or disorder as described herein. In one embodiment, there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof, for use in the treatment of a disease or disorder as described herein. In one embodiment, there is provided a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof, for use in the prophylaxis of a disease or disorder as described herein.

In a further embodiment, there is provided a method for the treatment or prophylaxis of a disease or disorder as described herein, which comprises administering to a subject in need thereof an effective amount of a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof. In a further embodiment, there is provided a method for the treatment of a disease or disorder as described herein, which comprises administering to a subject in need thereof an effective amount of a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof. In a further embodiment, there is provided a method for the prophylaxis of a disease or disorder as described herein, which comprises administering to a subject in need thereof an effective amount of a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt and/or solvate (e.g. salt) thereof. Pharmaceutical compositions of the invention may take the form of a pharmaceutical formulation as described below.

The invention also provides the use of a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof (e.g. salt) thereof, in the manufacture of a medicament for the treatment or prophylaxis of a disease or disorder as described herein. The invention also provides the use of a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof (e.g. salt) thereof, in the manufacture of a medicament for the treatment of a disease or disorder as described herein. The invention also provides the use of a pharmaceutical

composition comprising a compound of formula (I), or a pharmaceutically acceptable salt and/or solvate thereof (e.g. salt) thereof, in the manufacture of a medicament for the prophylaxis of a disease or disorder as described herein.

- 5 The amount of active ingredient which is required to achieve a therapeutic effect will, of course, vary with the particular compound, the route of administration, the subject under treatment or prophylaxis, including the type, species, age, weight, sex, and medical condition of the subject and the renal and hepatic function of the subject, and the particular disorder or disease being treated or prevented, as well as its severity. An ordinarily skilled physician, veterinarian or
10 clinician can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

Oral dosages of the present invention, when used for the indicated effects, will range between about 0.01 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day, suitably 0.01
15 mg per kg of body weight per day (mg/kg/day) to 10 mg/kg/day, and most suitably 0.1 to 5.0 mg/kg/day, for adult humans. For oral administration, the compositions are suitably provided in the form of tablets or other forms of presentation provided in discrete units containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A
20 medicament typically contains from about 0.01 mg to about 500 mg of the active ingredient, suitably from about 1 mg to about 100 mg of active ingredient. Intravenously, the most suitable doses will range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, compounds of the invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily.

25 Furthermore, suitably compounds of the invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

30 The pharmaceutical formulations according to the invention include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous [bolus or infusion], and intraarticular), intranasal (also known as nasal administration), inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose
35 pressurized aerosols, nebulizers or insufflators) insufflation, rectal, intraperitoneal, topical (including dermal, buccal, sublingual, and intraocular) and intrathecal administration, although

the most suitable route may depend upon, for example, the condition and disorder of the recipient.

Suitable pharmaceutical formulations according to the invention are those suitable for oral,
5 intrathecal and parenteral administration; and more suitably are those suitable for oral or intrathecal administration.

In one suitable embodiment a compound according to formula (I) is administered by intrathecal administration. Such a method of administration involves injection of the compound of the
10 invention into the spinal canal, or into the subarachnoid space so that it reaches the cerebrospinal fluid. This is advantageous for the administration of compounds which may not be able to pass the blood brain barrier *via* other routes of administration, such as oral administration.

Suitable pharmaceutical formulations may be administered intrathecally by continuous infusion such as with a catheter, or a pump, or intrathecally by a single bolus injection or by intermittent bolus injection. To be administered intrathecally, the pharmaceutical composition may be administered continuously or intermittently. The intermittent administration may be, for example,
15 every thirty minutes, every hour, every several hours, every 24 hours, every couple of days (for example every 48 or 72 hours) or any combination thereof.
20

When the pharmaceutical formulation of the invention is administered continuously, implantable delivery devices, such as an implantable pump may be employed. Examples of such delivery devices include devices which can be implanted subcutaneously in the body or in the cranium,
25 and provides an access port through which the pharmaceutical formulation may be delivered to the nerves or brain.

Intrathecal dosages of the present invention, when used for the indicated effects, will typically be less than 1 mg, such as less than 500 µg, for example less than 250 µg per kg of body
30 weight when administered in a single dose or intermittently for adult humans. When administered continuously, the intrathecal dosages of the present invention will typically be less than 250 µg per kg body weight per hour, such as less than 125 µg per kg body weight per hour for adult humans.

In another suitable embodiment a compound according to formula (I) is administered by intranasal, inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose pressurized aerosols, nebulizers or insufflators) or insufflation
35

administration. Such a method of administration allows for low doses of the compound of the invention to be administered, which can lead to a reduction in side-effects. For example, a daily dose of 10 to 0.01 μ g, suitably 1 to 0.01 μ g, and more suitably in the region of as low as 0.1 μ g (100ng) of compound of the invention may be used.

5

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into
10 association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets, pills or tablets each containing a predetermined
15 amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid, for example as elixirs, tinctures, suspensions or syrups; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

20 A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, lubricating, surface active or dispersing agent. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound
25 moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein. The compounds of formula (I) can, for example, be administered in a form suitable for immediate release or extended release. Immediate release or extended release can be achieved by the use of suitable pharmaceutical compositions comprising a compound of the present invention,
30 or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps. The compounds of the invention may also be administered liposomally.

Exemplary compositions for oral administration include suspensions which can contain, for
35 example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which can contain, for example,

microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate, calcium sulfate, sorbitol, glucose and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Disintegrators include without limitation starch, methylcellulose, agar, bentonite, xanthan gum and the like. The compounds of formula (I) can also be delivered through the oral cavity by sublingual and/or buccal administration. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms which may be used. Exemplary compositions include those formulating a compound of the present invention with fast dissolving diluents such as mannitol, lactose, sucrose and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (avicel) or polyethylene glycols (PEG). Such formulations can also include an excipient to aid mucosal adhesion such as hydroxy propyl cellulose (HPC), hydroxy propyl methyl cellulose (HPMC), sodium carboxy methyl cellulose (SCMC), maleic anhydride copolymer (e.g., Gantrez), and agents to control release such as polyacrylic copolymer (e.g. Carbopol 934). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. For oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like.

The compounds of formula (I) can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, 1,2-dipalmitoylphosphatidylcholine, phosphatidyl ethanolamine (cephaline), or phosphatidylcholine (lecithin).

Formulations for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of the sterile liquid carrier, for example saline or water-for-injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described. Exemplary compositions for

parenteral administration include injectable solutions or suspensions which can contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid, or Cremaphor.

Exemplary compositions for intranasal, aerosol or inhalation administration include solutions in saline, which can contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Formulations for rectal administration may be presented as a suppository with the usual carriers such as cocoa butter, synthetic glyceride esters or polyethylene glycol. Such carriers are typically solid at ordinary temperatures, but liquefy and/or dissolve in the rectal cavity to release the drug.

Formulations for topical administration in the mouth, for example buccally or sublingually, include lozenges comprising the active ingredient in a flavoured basis such as sucrose and acacia or tragacanth, and pastilles comprising the active ingredient in a basis such as gelatin and glycerine or sucrose and acacia. Exemplary compositions for topical administration include a topical carrier such as Plastibase (mineral oil gelled with polyethylene).

Suitable unit dosage formulations are those containing an effective dose, as hereinbefore recited, or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents.

The compounds of formula (I) are expected to display one or more of the following advantageous properties:

- inhibitory activity of mPTP as demonstrated in the assay of Biological Example 1 (preferably with a plC_{50} value of 6.0 and above); and
- low inhibition of CYP2D6 as demonstrated in the assays of Biological Example 2;

In addition to the properties described above, certain compounds of formula (I) may also display one or more of the following advantageous properties:

- improved (i.e. higher) solubility and/or improved (i.e. lower) intrinsic clearance (CL_{int}) resulting in e.g. improved (i.e. higher) oral bioavailability and/or improved (i.e. higher) systemic exposure as demonstrated in the assays of Biological Examples 3 and 4.

The invention is further exemplified by the following non-limiting examples.

EXAMPLES

The invention is illustrated by the compounds described below. The following examples describe the laboratory synthesis of specific compounds of the invention and are not meant to limit the scope of the invention in any way with respect to compounds or processes. It is understood that, although specific reagents, solvents, temperatures and time periods are used, there are many possible equivalent alternatives that can be used to produce similar results. The invention is meant to include such equivalents.

General Experimental Details

Starting materials, reagents and solvents were obtained from commercial suppliers and used without further purification unless otherwise stated. Unless otherwise stated, all compounds with chiral centres are racemic. Where reactions are described as having been carried out in a similar manner to earlier, more completely described reactions, the general reaction conditions used were essentially the same. Work up conditions used were of the types standard in the art, but may have been adapted from one reaction to another. The starting material may not necessarily have been prepared from the batch referred to. Compounds synthesised may have various purities, ranging from for example 85% to 99%. Calculations of number of moles and yield are in some cases adjusted for this.

Purity of final compounds was confirmed by HPLC/MS analysis and determined to be at least ≥ 90%, and in the significant majority of cases ≥ 95%. Analytical LCMS was conducted using the instrumentation shown in Table 1. ¹H NMR were recorded at 300K in a Bruker 300MHz instruments (ADVANCE III and ADVANCE III HD). Flash prep HPLC was conducted using the following columns: XBridge Prep C18 OBD Column, 5um, 19x150mm; XBridge Prep C18 OBD Column, 5um, 30x150mm; Welch Xtimate C18, 21.2x250mm, 5um; SunFire Prep C18 OBD

19x150mmx5um. SFC purification was conducted using the following columns; (a) CHIRALPAK AS-H, 3*25cm, 5um (b) SFC-YMC Cellulose-SB,, 4.6 x 100mm, 3um.

Table 1: Analytical LC-MS conditions

Instrument ID	Column	Mobile Phase	Flow Rate
LCMS01	Halo-C18, 30*3.0mm, 2.0µm	A:H ₂ O/0.05%TFA; B:ACN	1.5 mL/min
LCMS02	Cortecs C18+, 50*3.0mm, 2.7µm	A:H ₂ O/0.05%TFA; B:ACN/0.05%TFA	1.5 mL/min
LCMS03	Halo-C18, 30*3.0mm, 2.0µm	A:H ₂ O/0.1%TFA; B:ACN/0.05%FA	1.5 mL/min
LCMS04	Kinetex XB-C18, 50*3.0mm, 2.6µm	A:H ₂ O/0.01%TFA; B:ACN/0.01%FA	1.5 mL/min
LCMS05	Poroshell HPH-C18, 50*3.0mm, 2.7µm	A:H ₂ O/5mM NH ₄ HCO ₃ ; B:MeOH	1.0 mL/min
LCMS06	Xbridge C18, 50*3.0mm, 3.5µm	A:H ₂ O/5mM NH ₄ HCO ₃ +0.05% NH ₃ .H ₂ O; B:5%H ₂ O in ACN	1.2 mL/min
LCMS07	Poroshell HPH-C18, 50*3.0mm, 2.7µm	A:H ₂ O/0.05% NH ₃ .H ₂ O; B:ACN	1.2 mL/min
LCMS08	Halo C18, 50*3.0mm, 2.7µm	A:H ₂ O/0.05%TFA; B:ACN	1.5 mL/min
LCMS09	Poroshell HPH-C18, 50*3.0mm, 2.7µm	A:H ₂ O/0.05% NH ₃ .H ₂ O; B:ACN	1.2 mL/min


Abbreviations

	CH ₃ CN	Acetonitrile
	Cs ₂ CO ₃	Cesium Carbonate
5	DCM	Dichloromethane
	DIPEA	Diisopropylethylamine
	DMF	Dimethylformamide
	Et	Ethyl
	Et ₃ N	Triethylamine
10	EtOAc	Ethyl acetate
	EtOH	Ethyl alcohol
	HATU	1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxide hexafluorophosphate
	heps	hepatocytes
15	HCl	Hydrochloric acid
	K ₂ CO ₃	Potassium Carbonate
	K ₃ PO ₄	Potassium Phosphate
	LiHMDS	Lithium bis(trimethylsilyl)amide
	Me	Methyl
20	MeOH	Methanol
	NaHCO ₃	Sodium bicarbonate
	NaOAc	Sodium acetate
	NaOH	Sodium hydroxide
	NBS	N-Bromosuccinimide
25	NCS	N-Chlorosuccinimide
	NH ₄ Cl	Ammonium chloride
	PE	Petroleum ether
	Pd(OAc) ₂	Palladium(II) Acetate
30	Pd(dppf)Cl ₂ .CH ₂ Cl ₂	Bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane
	Pd(dppf)Cl ₂	Bis(diphenylphosphino)ferrocene]dichloropalladium(II)
	RT	room temperature
	o/n	overnight (16h)
	TFA	Trifluoroacetic acid
35	THF	Tetrahydrofuran
	THP	Tetrahydropyranyl
	T ₃ P	Propanephosphonic anhydride

uL microlitre
 uM micromolar

Preparation of Comparative Example 1


5 Comparative Example 1: (E)-N-(2-methyl-3-fluorophenyl)-3-(1H-indazol-6-yl)acrylamide



Comparative Example 1 was prepared according to methods described in Chen et al. (Assay and Drug Development Technologies, 2018, 16, 445-455). Comparative Example 1 may also be prepared using analogous synthetic methods to those described herein for Examples 1 to 64.


Preparation of Examples 1 to 117

Intermediate 1: N-(3-fluoro-2-methylphenyl)acrylamide



To a stirred solution of 3-fluoro-2-methyl aniline (1.0 g, 8.0 mmol, 1.0 eq.) and DIPEA (3.1 g, 23.9 mmol, 3.0 eq.) in DCM (40 mL) was added acryloyl chloride (0.71 mL, 8.8 mmol, 1.1 eq.) dropwise at 0 °C under nitrogen. The resulting mixture was stirred at 25 °C under an inert atmosphere of nitrogen for 3 h. The resulting mixture was washed with 2 x 30 mL of water and the organic layer was concentrated. The residue was purified by silica gel chromatography eluting with EtOAc/petroleum ether (1:5). This resulted in 1.04 g (73%) of N-(3-fluoro-2-methylphenyl)acrylamide as a white solid. LC-MS (ES, m/z): [M+H]⁺=180


Intermediate 2: N-(3-chloro-2-methylphenyl)acrylamide




Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3-chloro-2-methyl-aniline (300 mg, 2.12 mmol, 1.0 eq.), DCM (15 mL), Et₃N (1 mL, 7.19 mmol, 3.4 eq.). This was followed by the addition of acryloyl chloride (260 mg, 2.9 mmol, 1.4 eq.) dropwise with stirring at 25 °C. The resulting solution was stirred for 5 h at 25 °C. The resulting mixture was washed with 2 x 10 mL of water and the organic layer was concentrated. The residue was purified by silica gel chromatography eluting with EtOAc/PE

(1:5). This resulted in 300 mg (74%) of *N*-(3-chloro-2-methylphenyl)acrylamide as a light yellow solid. LC-MS (ES, *m/z*): $[M+H]^+=196$

Intermediate 3: (E)-3-(3-methyl-1H-indazol-6-yl)acrylic acid




Step 1: Into a 250-mL 3-necked round-bottom flask, was placed 6-bromo-3-methyl-1H-indazole (2.50 g, 11.84 mmol, 1.00 equiv), methyl acrylate (1.53 g, 17.76 mmol, 1.50 equiv), Et₃N (3.60 g, 35.53 mmol, 3.00 equiv), Pd(dppf)Cl₂ (0.87 g, 1.18 mmol, 0.10 equiv), DMF (100.00 mL). The resulting solution was stirred for 10 hr at 120 °C. The mixture was concentrated and the residue was applied onto a silica gel column with THF/PE (1/1). This resulted in 0.9 g (35% yield) of methyl (2E)-3-(3-methyl-1H-indazol-6-yl)prop-2-enoate as a light yellow solid.




Step 2: Into a 40-mL vial, was placed methyl (2E)-3-(3-methyl-1H-indazol-6-yl)prop-2-enoate (890.00 mg, 4.12 mmol, 1.00 equiv), NaOH (329.24 mg, 8.23 mmol, 2.00 equiv) in H₂O (10.00 mL) and MeOH (10.00 mL). The resulting solution was stirred for 2 hr at 20 °C. The pH value of the solution was adjusted to 3 with HCl (1 mol/L). The solids were collected by filtration. This resulted in 500 mg (60% yield) of (E)-3-(3-methyl-1H-indazol-6-yl)acrylic acid as a off-white solid.

Intermediate 4: N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide




Step 1: To a stirred solution of 2-hydroxy-4,6-dimethylpyridine-3-carbonitrile (70 g, 472 mmol) in Ac₂O (278 mL) were added c.HNO₃ (43 mL) dropwise in portions at 0°C. The ice bath was removed and the reaction stirred to room temperature for 10 h. The resulting mixture was cooled and diluted with H₂O (200 mL). The precipitated solids were collected by filtration and washed with H₂O (2x20 mL), then dried under reduced pressure to afford 2-hydroxy-4,6-dimethyl-5-nitronicotinonitrile (56g, 70% yield) as a yellow solid. LC MS (ES, *m/z*):

$[M+H]^+ = 194$




Step 2: A solution of 2-hydroxy-4,6-dimethyl-5-nitronicotinonitrile (55 g, 327 mmol) and H_2O (55 mL) in c. H_2SO_4 (275 mL) was stirred for 4 hours at 150°C . The mixture was cooled and diluted with H_2O (300 mL). The precipitated solids were collected by filtration and washed with H_2O (2x20 mL), then dried under reduced pressure to afford 4,6-dimethyl-5-nitropyridin-2-ol (24 g, 44% yield) as a light yellow solid. LCMS (ES, m/z):

$[\text{M}+\text{H}]^+ = 169$




Step 3: To a stirred solution of 4,6-dimethyl-5-nitropyridin-2-ol (10 g, 58 mmol) and dimethyl sulfate (15 g, 116 mmol, 2.0 equiv) in DMF (200 mL) was added 60% NaH in oil (2.86 g, 116 mmol, 2.0 equiv) in portions at 0°C . The reaction was stirred for 4 hours at 0°C , then quenched by the addition of water (200 mL). The resulting mixture was extracted with ethyl acetate (3x150 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , then concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/ AcOEt (3:1), to afford 6-methoxy-2,4-dimethyl-3-nitropyridine (2.4 g, 22% yield) as a light yellow solid. LC MS (ES, m/z): $[\text{M}+\text{H}]^+ = 183$




Step 4: 6-Methoxy-2,4-dimethyl-3-nitropyridine (2.4 g, 14 mmol) and 10% Pd/C (0.40 g) in MeOH (30 mL) were hydrogenated under an atmosphere of hydrogen for 2 h. The resulting mixture was filtered, and the filter cake washed with MeOH (3x2 mL). The combined filtrate was concentrated to afford 6-methoxy-2,4-dimethylpyridin-3-amine (1.61 g, 81% yield) as a light yellow oil. LC MS (ES, m/z):

$[\text{M}+\text{H}]^+ = 153$



Step 5: To a stirred solution of 6-methoxy-2,4-dimethylpyridin-3-amine (1.5 g, 9.8 mmol), TEA (2.99 g, 30 mmol, 3 equiv) and DMAP (0.24 g, 1.9 mmol, 0.2 equiv) in DCM (30 mL) was added acryloyl chloride (2.1 g, 24 mmol, 2.4 equiv) dropwise at 0°C over 30 mins. The resulting mixture was stirred for 2h and diluted with ice-water (20 mL). The aqueous layer was extracted with DCM (3x20 mL). The combined organics were dried over anhydrous Na₂SO₄ then concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:1), to afford N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (1.4 g, 69% yield) as an off-white solid.


10 Intermediate 5: 6-bromo-5-fluoro-3-methyl-1H-indazole



Step 1: To a solution of 1-(4-bromo-2,5-difluorophenyl)ethanone (4.0 g, 17 mmol) in ethylene glycol (100 mL) was added 98% NH₂NH₂·H₂O (2.2 g, 43 mmol, 2.5 eq) was added. The mixture was stirred for 16h at 160°C under N₂ atmosphere, then was cooled and quenched with water (200 mL). The precipitated solids were collected by filtration and washed with water (2x50 mL). Drying under reduced pressure gave 6-bromo-5-fluoro-3-methyl-1H-indazole (3.0 g, 77% yield) as an off-white solid.


LCMS (ES, m/z): 229, 231 [M+H]⁺

20 Intermediate 6: (2E)-3-[3-fluoro-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid



Step 1: A solution of 6-bromo-1H-indazole (4.4 g, 22 mmol), Selectfluor (15 g, 42 mmol, 1.9 equiv) and AcOH (3 mL, 53 mmol, 2.4 equiv) in MeCN (300 mL) was stirred for 4 hours at reflux. The mixture was cooled and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether: ethyl acetate (5:1) to afford 6-bromo-3-fluoro-1H-indazole (1.13 g, 23% yield) as a yellow solid.


LC-MS (ES, m/z): 215, 217 [M+H]



Step 2: A solution of 6-bromo-3-fluoro-1H-indazole (1.1 g, 5.1 mmol), DHP (6.0 g, 72 mmol, 14 equiv) and PPTS (642 mg, 2.5 mmol, 0.5 equiv) in THF (80 mL) was stirred for 10 hours at 70°C.


The mixture was cooled and diluted with saturated brine (100 mL) and saturated aqueous NaHCO₃ (20 mL). The aqueous layer was extracted with EA (2x100 mL). The combined were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (5:1) to afford 6-bromo-3-fluoro-1-(oxan-2-yl) indazole (1.0 g, 65% yield) as a yellow solid.

LC-MS (ES, m/z): 299, 301 [M+H]⁺



Step 3: A solution of 6-bromo-3-fluoro-1-(oxan-2-yl) indazole (700 mg, 2.3 mmol), methyl acrylate (604 mg, 7.0 mmol, 3 equiv), Pd(dppf)Cl₂ (191 mg, 0.3 mmol, 0.1 equiv) and TEA (1.06 g, 10.4 mmol, 4.5 equiv) in DMF (15 mL) was stirred for 4 hours at 110°C. The mixture was cooled and diluted with H₂O (20 mL), then extracted with ethyl acetate (4 x 50 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether: ethyl acetate (3:1) to afford methyl (2E)-3-[3-fluoro-1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (300 mg, 42% yield) as a light yellow solid.


LC-MS (ES, m/z): 305 [M+H]⁺



Step 4: A solution of (2E)-3-[3-fluoro-1-(oxan-2-yl) indazol-6-yl] prop-2-enoate (300 mg, 0.9 mmol), NaOH (107 mg, 2.7 mmol, 3.0 equiv) in H₂O (2 mL) and MeOH (2 mL) was stirred for 4 hours at room temperature. The resulting mixture was concentrated, then diluted with H₂O (3 mL). The mixture was neutralized to pH 6 with 1M HCl. The precipitated solids were collected by filtration and washed with H₂O (2x2 mL). The solid was dried under vacuum to give (2E)-3-[3-fluoro-1-(oxan-2-yl) indazol-6-yl] prop-2-enoic acid (234 mg, 82% yield) as a white solid.

LC-MS (ES, m/z): 291 [M+H]⁺


Intermediate 7: 6-bromo-3-chloro-5-fluoro-1H-indazole



A solution of 6-bromo-5-fluoro-1H-indazole (200 mg, 0.9 mmol) and NCS (248 mg, 1.8 mmol, 2 equiv) in DMF (4 mL) was stirred for 2 h at 60°C. The mixture was cooled and diluted with H₂O (8 mL). The precipitated solids were collected by filtration and washed with H₂O (2x5 mL). The precipitate was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:1) to afford 6-bromo-3-chloro-5-fluoro-1H-indazole (143 mg, 67% yield) as an off-white solid.


LCMS (ES, m/z): 249 [M+H]⁺

Intermediate 8: (2E)-3-[3,5-difluoro-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid




Step 1: A mixture of 6-bromo-5-fluoro-1H-indazole (2.8 g, 13 mmol), Selectfluor (9.5 g, 27 mmol, 2 equiv) and AcOH (2 mL, 35 mmol, 2.6 equiv) in MeCN (40 mL) was stirred for 2 hours at reflux. The mixture was cooled and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether: ethyl acetate (5:1) to afford 6-bromo-3,5-difluoro-1H-indazole (490 mg, 16% yield) as a brown solid.

LC-MS (ES, m/z): 233 [M+H]⁺



Step 2: A solution of 6-bromo-3,5-difluoro-1H-indazole (440 mg, 1.9 mmol), DHP (2.22 g, 26 mmol, 14 equiv) and PPTS (237 mg, 0.9 mmol, 0.5 equiv) in THF (10 mL) was stirred for 2 hours at 70°C. The reaction was cooled and diluted with saturated brine (20 mL) and saturated aqueous NaHCO₃ (5 mL). The resulting mixture was extracted with CH₂Cl₂ (3x20 mL). The combined organics were dried over anhydrous Na₂SO₄. The residue was purified by silica gel column chromatography, eluting with petroleum ether: ethyl acetate (3:1) to afford 6-bromo-3,5-difluoro-1-(oxan-2-yl)indazole (320 mg, 53% yield) as a brown solid.


LC-MS (ES, m/z): 317 [M+H]⁺



Step 3: A solution of 6-bromo-3,5-difluoro-1-(oxan-2-yl)indazole (320 mg, 1.0 mmol), methyl acrylate (344 mg, 4.0 mmol, 4 equiv), Pd(dppf)Cl₂ (163 mg, 0.2 mmol, 0.2 equiv) and TEA (505 mg, 5.0 mmol, 5 equiv) in DMF (10 mL) was stirred for 2 hours at 110°C under N₂ atmosphere.

The mixture was cooled and diluted with H₂O (20 mL), then extracted with ethyl acetate (3 x 25 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether: ethyl acetate (5:1) to afford methyl (2E)-3-[3,5-difluoro-1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (120 mg, 24% yield) as a light yellow solid.


LC-MS (ES, m/z): 323 [M+H]⁺



Step 4: A solution of methyl (2E)-3-[3,5-difluoro-1-(oxan-2-yl) indazol-6-yl]prop-2-enoate (120 mg, 0.3 mmol), NaOH (36 mg, 0.9 mmol, 3 equiv) in H₂O (1 mL) and MeOH (1 mL) was stirred for 4 hours at room temperature. The mixture was acidified to pH6 with 1M HCl, and the precipitated solids were collected by filtration and washed with THF (3x3 mL). Drying under vacuum gave (2E)-3-[3,5-difluoro-1-(oxan-2-yl) indazol-6-yl] prop-2-enoic acid (63 mg, 55% yield) as a white solid.

LC-MS (ES, m/z): 309 [M+H]⁺


Intermediate 9: 6-bromo-5-fluoro-1-(oxan-2-yl)indazole-3-carbonitrile




Step 1: A mixture of 6-bromo-5-fluoro-1H-indazole (3.0 g, 13 mmol), NaOH (0.56 g, 39 mmol, 3 equiv) and iodine (7.1 g, 26 mmol, 2 equiv) in dioxane (60 mL) and H₂O (10 mL) was stirred for 2 h then neutralized 1M aqueous HCl. The resulting mixture was extracted with EtOAc (3x50 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was

purified by silica gel column chromatography, with petroleum ether: EtOAc (2:1) to afford 6-bromo-5-fluoro-3-iodo-1H-indazole (2.90 g, 61% yield) as a white solid.

LCMS (ES, m/z): 340 [M+H]⁺




- 5 **Step 2:** A mixture of 6-bromo-5-fluoro-3-iodo-1H-indazole (2.9 g, 8.5 mmol), Zn(CN)₂ (1.2 g, 10 mmol, 1.2 equiv), Dppf (0.94 g, 1.7 mmol, 0.2 equiv), Pd₂(dba)₃ (0.78 g, 0.86 mmol, 0.1 equiv) and K₂CO₃ (3.5 g, 26 mmol, 3 equiv) in DMF (60 mL) was stirred for 2 h at 60°C under nitrogen atmosphere. The mixture was cooled and diluted with water (80 mL), then extracted with EtOAc (3 x 80 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with Petroleum ether: EtOAc (1:1) to afford 6-bromo-5-fluoro-1H-indazole-3-carbonitrile (1.3 g, 64% yield) as a white solid.
- 10 LCMS (ES, m/z): 240 [M+H]⁺




- 15 **Step 3:** A solution of 6-bromo-5-fluoro-1H-indazole-3-carbonitrile (1.3 g, 5.4 mmol), DHP (1.37 g, 16 mmol, 3 equiv) and TsOH (0.28 g, 1.6 mmol, 0.3 equiv) in THF (26 mL) was stirred for 13 h at 65°C. The reaction was cooled and diluted with saturated brine (20 mL) and saturated aqueous NaHCO₃ (5 mL). The mixture was extracted with EtOAc (2 x 10 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/EtOAc (8:1) to afford 6-bromo-5-fluoro-1-(oxan-2-yl)indazole-3-carbonitrile (1.6 g, 92% yield) as an off-white solid.
- 20

Intermediate 10: N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide




- 25 **Step 1:** To a solution of 2,4-dibromo-5-fluoro-3-nitropyridine (1.00 g, 3.3 mmol) in 1,4-dioxane (10 mL) and H₂O (1 mL) was added 50% trimethyl-1,3,5,2,4,6-trioxatriborinane in THF (1.7 g, 6.6 mmol, 2 eq), Pd(dppf)Cl₂ (0.24 g, 0.3 mmol, 0.1 eq) and Cs₂CO₃ (2.17 g, 6.6 mmol, 2 eq). The

mixture was stirred for 16 h at 100°C under N₂ atmosphere. The reaction was cooled and diluted with water (20 mL), and the resulting mixture extracted with EtOAc (3 x 10 mL). The combined organics were washed with brine (3 x 5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (9:1) to afford 5-fluoro-2,4-dimethyl-3-nitropyridine (0.46 g, 81% yield) as a light red oil.



Step 2: To a solution of 5-fluoro-2,4-dimethyl-3-nitropyridine (0.46 g, 1.8 mmol) in MeOH (10 mL) was added 10% Pd/C (120 mg). The mixture was stirred for 2h at room temperature under H₂ atmosphere. The resulting mixture was filtered, and the filter cake washed with MeOH (3 x 5 mL). The combined filtrate was concentrated under reduced pressure to afford 5-fluoro-2,4-dimethylpyridin-3-amine (0.32 g, 86% yield) as an off-white solid.


LCMS (ES, m/z): 141 [M+H]⁺



Step 3: To a solution of 5-fluoro-2,4-dimethylpyridin-3-amine (0.32 g, 2.2 mmol) in DCM (2 mL) was added TEA (0.69 g, 6.8 mmol, 3 eq). The reaction was cooled to 0°C, and acryloyl chloride (0.31 g, 3.4 mmol, 1.5 eq) in DCM (1 mL) was added dropwise over 2 min at 0°C. The reaction was stirred for 2h at 0°C, then concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:5) to afford N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (0.16 g, 36% yield) as an off-white solid.

LCMS (ES, m/z): 195 [M+H]⁺


Intermediate 11: N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]prop-2-enamide



Step 1: To 2-chloro-5-fluoropyridin-4-amine (4.0 g, 27 mmol) in H₂SO₄ (60 mL) at 0°C was added HNO₃ (3.44 g, 55 mmol, 2 eq) dropwise over 5 min. The mixture was stirred for 1h at 0°C, then 20h at room temperature. The reaction was quenched with water/ice and basified to pH9 with NH₃·H₂O. The resulting mixture was extracted with EtOAc (3 x 10 mL). The combined


organic layers were washed with brine (3x3 mL), dried over anhydrous Na_2SO_4 and concentrated to afford 2-chloro-5-fluoro-3-nitropyridin-4-amine (4.8 g, 92% yield) as a light yellow solid.

LCMS (ES, m/z): 192 $[\text{M}+\text{H}]^+$




Step 2: To a solution of 2-chloro-5-fluoro-3-nitropyridin-4-amine (2.0 g, 10 mmol) in 1,4-dioxane (11 mL) and H_2O (1 mL) was added 50% trimethyl-1,3,5,2,4,6-trioxatriborinane in THF (7.9 g, 31 mmol, 3eq), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.76 g, 1 mmol, 0.1 eq) and Cs_2CO_3 (6.8 g, 21 mmol, 2 eq). The reaction was stirred for 16h at 100°C under N_2 atmosphere, then cooled and diluted with water (20 mL). The resulting mixture was extracted with EtOAc (3 x 10mL). The combined organic layers were washed with brine (3x5 mL), dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:1) to afford 5-fluoro-2-methyl-3-nitropyridin-4-amine (1.2 g, 67% yield) as a light yellow solid.

LCMS (ES, m/z): 172 $[\text{M}+\text{H}]^+$




Step 3: To 5-fluoro-2-methyl-3-nitropyridin-4-amine (0.90, 5.2 mmol) in CH_3CN (9 mL) was added CuBr (3.0 g, 21 mmol, 4 eq) and $t\text{-BuONO}$ (1.1 g, 11 mmol, 2 eq) at 0°C . The mixture was stirred for 2h at 0°C , then concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (3:1) to afford 4-bromo-5-fluoro-2-methyl-3-nitropyridine (1.0 g, 81% yield) as an off-white solid.



Step 4: To 4-bromo-5-fluoro-2-methyl-3-nitropyridine (0.90 g, 3.9 mmol) in toluene (9 mL) was added tributyl(methoxymethyl)stannane (2.6 g, 7.8 mmol, 2 eq) and $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (273 mg, 0.39 mmol, 0.1 eq). The reaction was stirred for 2h at 120°C under N_2 atmosphere. The resulting mixture was diluted with water (100mL), extracted with EtOAc (3 x 5 mL), and the combined


organic layers were washed with brine (2x5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (2:1) to afford 5-fluoro-4-(methoxymethyl)-2-methyl-3-nitropyridine (600 mg, 78% yield) as a colorless oil.

- 5 ¹H-NMR (400 MHz, DMSO-*d*₆, ppm): δ 8.80 (s, 1H), 4.60 (d, *J* = 1.4 Hz, 2H), 3.24 (s, 3H), 2.49 (s, 3H).



- 10 **Step 5:** To a solution of 5-fluoro-4-(methoxymethyl)-2-methyl-3-nitropyridine (0.90 g, 4.5 mmol) in MeOH (10 mL) was added 10% Pd/C (0.48 g, 0.4 mmol, 0.1 eq). The reaction was stirred for 2h at room temperature under H₂ atmosphere. The resulting mixture was filtered, and the filter cake was washed with MeOH (3x3 mL). The filtrate was concentrated to afford 5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-amine (600 mg, 78% yield) as an off-white solid.


LCMS-6 (ES, *m/z*): [M+H]⁺=171



- 15 **Step 6:** To 5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-amine (0.30 g, 1.7 mmol) in DCM (2.0 mL) was added TEA (0.53 g, 5.3 mmol, 3 eq). The mixture was cooled to 0°C and acryloyl chloride (0.16 g, 1.8 mmol) in DCM (1.0 mL) was added dropwise over 2 min at 0°C. The resulting mixture was stirred for 2h at 0°C, then concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:1) to afford N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]prop-2-enamide (120 mg, 30% yield) as an off-white solid.

20 LCMS (ES, *m/z*): 225 [M+H]⁺


Intermediate 12: N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]prop-2-enamide



- 25 **Step 1:** To tert-butyl N-(5-fluoropyridin-3-yl)carbamate (0.50 g, 2.3 mmol) in THF (5 mL) was added TMEDA (0.68 g, 5.9 mmol, 2.5 eq). The reaction mixture was cooled to -30°C, and 2.5M *n*-BuLi in hexanes (2.4 mL, 5.9 mmol, 2.5 eq) was added, keeping the internal temperature below -20°C. On complete addition, the reaction mixture was stirred at -30°C for a further 50 mins. Iodine


(0.96 g, 3.7 mmol, 1.6 eq) was slowly added, then the mixture was stirred at -30° C for a further 45 mins. The reaction was quenched by the slow addition of a saturated ammonium chloride aqueous solution (5mL) and water (5mL). The resulting mixture was extracted with EtOAc (3 x 7 mL). The combined organics were washed with brine (3x3 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether/ethyl acetate (5:1) to afford tert-butyl N-(5-fluoro-4-iodopyridin-3-yl)carbamate (0.50 g, 63% yield) as an off-white solid.

LCMS (ES, m/z): 339 [M+H]⁺




Step 2: To a solution of tert-butyl N-(5-fluoro-4-iodopyridin-3-yl)carbamate (0.50 g, 1.5 mmol) in toluene (5 mL) was added tributyl(methoxymethyl)stannane (0.99 g, 2.9 mmol, 2 eq) and Pd(PPh₃)₂Cl₂ (0.10 g, 0.1 mmol, 0.1 eq). The reaction was stirred for 8 h at 120°C, then cooled and quenched by water (20 mL). The resulting mixture was extracted with EtOAc (3 x 10mL). The combined organic layers were washed with brine (3 x 5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether/ethyl acetate (1:1) to afford tert-butyl N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]carbamate (0.28g, 74% yield) as a light yellow oil.

LCMS (ES, m/z): 257 [M+H]⁺



Step 3: A solution of tert-butyl N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]carbamate (280 mg, 1.1 mmol) in 4M HCl in 1,4-dioxane (2 mL) was stirred for 10 min at 60°C, then concentrated to give 5-fluoro-4-(methoxymethyl)pyridin-3-amine (160 mg, crude) as an off-white solid. The crude product was used in the next step directly without further purification.


LCMS (ES, m/z): 157 [M+H]⁺



Step 4: To a solution of 5-fluoro-4-(methoxymethyl)pyridin-3-amine (150 mg, 0.9 mmol) in DCM (1 mL) was added TEA (97 mg, 2.7 mmol, 3 eq). The solution was cooled in an ice bath and acryloyl chloride (44 mg, 1.3 mmol, 1.5 eq) in DCM (0.5 mL) was added dropwise. The reaction was stirred to room temperature over 1 h, then concentrated. The residue was dissolved in DCM (5 mL) and purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:1) to afford N-[5-fluoro-4-(methoxymethyl)-pyridin-3-yl]prop-2-enamide (120 mg, 30% yield) as an off-white solid.


LCMS (ES, m/z): 211 [M+H]⁺

Intermediate 13: (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid




Step 1: A solution of 6-bromo-3-methyl-1H-indazole (4.0 g, 19 mmol) in THF (80 mL) was treated with DHP (4.8 g, 57 mmol, 3.0 equiv) and PPTS (480 mg, 1.9 mmol, 0.1 equiv). The solution was stirred for 2 hours at 70°C, then cooled and quenched with water. The resulting mixture was extracted with CH₂Cl₂ (4 x 50 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether: THF (5:1) to afford 6-bromo-3-methyl-1-(oxan-2-yl)indazole (4.9 g, 88% yield) as an off-white solid.

LC-MS (ES, m/z): 295 [M+H]⁺



Step 2: A solution of 6-bromo-3-methyl-1-(oxan-2-yl)indazole (4.9 g, 17 mmol) and methyl acrylate (5.7 g, 66 mmol, 4.0 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (2.7 g, 3.3 mmol, 0.2 equiv) and TEA (8.4 g, 83 mmol, 5.0 equiv) in DMF (100 mL) was stirred for 2 hours at 110°C under N₂ atmosphere. The mixture was allowed to cool and was quenched with water. The resulting mixture was extracted with ethyl acetate (4x50 mL). The combined organics were washed with brine (3x5 mL) and dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether/ethyl acetate (5:1) to afford methyl (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (4.0 g, 80% yield) as a yellow solid.


LC-MS (ES, m/z): 301 [M+H]⁺



Step 3: A solution of methyl (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (4.0 g, 13 mmol) in H₂O (27 mL), MeOH (27 mL) and THF (27 mL) was treated with NaOH (1.6 g, 40 mmol, 3 equiv). The mixture was stirred for 4 hours at room temperature, then acidified to pH6 with 1M HCl. The precipitated solids were collected by filtration and washed with THF (3x10 mL). Drying under vacuum gave (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (3.2 g, 84% yield) as a white solid.


LC-MS (ES, m/z): 287 [M+H]⁺

10 Intermediate 14: (2E)-3-[1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid




Step 1: To a solution of 6-bromo-1H-indazole (35 g, 178 mmol) and PPTS (4.5 g, 18 mmol, 0.1 eq) in THF (400 mL) was added dihydropyran (37.4 g, 444 mmol, 2.5 eq). The reaction was stirred for 3h at 70°C, then cooled, concentrated and diluted with water (300 mL). The resulting mixture was extracted with EtOAc (3 x 500 mL). The combined organics were washed with brine (3 x 200 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/EtOAc (3:1) to afford 6-bromo-1-(oxan-2-yl)indazole (30 g, 60% yield) as an off-white solid.

LCMS (ES, m/z): 281 [M+H]⁺



Step 2: To a stirred solution of 6-bromo-1-(oxan-2-yl)indazole (30 g, 107 mmol) in DMF (300 mL) was added methyl acrylate (18.4 g, 213 mmol, 2.0 eq), TEA (32.4 g, 320 mmol, 3.0 eq) and Pd(dppf)Cl₂ (7.8 g, 11 mmol, 0.1 eq) at room temperature under nitrogen atmosphere. The reaction was stirred for 3h at 110°C, then cooled, quenched with water (500 mL) and extracted with EtOAc (3 x 300 mL). The combined organics were washed with brine (2 x 200 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/EtOAc (1:1) to afford methyl (2E)-3-[1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (25 g, 82% yield) as a light yellow oil.


LCMS (ES, m/z): 287 [M+H]⁺



Step 3: To a stirred solution of methyl (2E)-3-[1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (25 g, 87 mmol) in MeOH (250 mL) and H₂O (100 mL) was added NaOH (7.0 g, 175 mmol, 2.0 eq). The resulting mixture was stirred for 3h at room temperature, then concentrated. The remaining aqueous was washed with EtOAc (2 x 300 mL), then acidified to pH4 with 0.1M HCl (aq.). The resulting mixture was extracted with EtOAc (3 x 300 mL). The combined organics were washed with brine (2 x 200 mL), dried over anhydrous Na₂SO₄ and concentrated to afford (2E)-3-[1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (10 g, 42% yield) as an off-white solid.

LCMS (ES, m/z): 273 [M+H]⁺

Example 1: (E)-3-(1H-benzo[d][1,2,3]triazol-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide




To a microwave vial with N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 100 mg, 0.56 mmol, 1.0 eq.), 5-bromo-3H-1,2,3-benzotriazole (110 mg, 0.56 mmol, 1.0 equiv), Pd(OAc)₂ (19 mg, 0.084 mmol, 0.15 equiv), tris(2-methylphenyl)phosphane (34 mg, 0.112 mmol, 0.20 equiv) and tetrabutylammonium chloride (155 mg, 0.558 mmol, 1.0 equiv) was added DMF (1.5 mL). The resulting solution was stirred for 12h at 115 °C. The resulting mixture was diluted with 20 mL EtOAc washed with 2 x 10 mL of 1 M aq. K₂CO₃ and the organic layer was concentrated and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column. The resulting crude product was purified by Flash-Prep-HPLC. This resulted in 9 mg (5%) of (E)-3-(1H-benzo[d][1,2,3]triazol-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide as a pale grey solid. LC-

MS(ES,m/z):[M+H]⁺=297;

¹H NMR (400 MHz, DMSO-*d*₆, ppm): δ 9.65 (s, 1H), 8.15 (s, 1H), 7.96-7.92 (m, 1H), 7.79 (d, *J* = 16.0 Hz, 1H), 7.71-7.66 (m, 1H), 7.49 (s, 1H) 7.27-7.20 (m, 1H), 7.10-6.96 (m, 2H), 2.18 (s, 3H).

Example 2: (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]thiazol-5-yl)acrylamide




To a microwave vial with N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 180 mg, 1.0 mmol, 1.0 equiv), 5-bromo-3H-1,3-benzothiazol-2-one (231 mg, 1.0 mmol, 1.0 equiv), Pd(OAc)₂ (34 mg, 0.151 mmol, 0.15 equiv), tris(2-methylphenyl)phosphane (62 mg, 0.201 mmol, 0.20 equiv) and tetrabutylammonium chloride (279 mg, 1.0 mmol, 1.0 equiv) was added DMF (2.8 mL). The resulting solution was stirred for 12 h at 115 °C. The resulting mixture was diluted with 30 mL EtOAc washed with 2 x 15 mL of 1 M aq. K₂CO₃ and the organic layer was concentrated and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column. The resulting crude product was purified by Flash-Prep-HPLC.

This resulted in 6 mg (2%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]thiazol-5-yl)acrylamide as a white solid. LC-MS (ES, m/z):

[M+H]⁺=329

¹H NMR (400 MHz, DMSO-*d*₆, ppm): δ 12.12 (s, 1H), 9.66 (s, 1H), 7.66-7.56 (m, 2H), 7.48 (d, *J* = 8.0 Hz, 1H), 7.39-7.30 (m, 2H), 7.29-7.21 (m, 1H) 7.06-6.92 (m, 2H), 2.17 (s, 3H).

Example 3: (E)-3-(3,3-dimethyl-2-oxoindolin-6-yl)-N-(3-fluoro-2-methylphenyl) acrylamide




To a microwave vial with N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 185 mg, 1.03 mmol, 1.0 equiv), 6-bromo-3,3-dimethyl-1H-indol-2-one (240 mg, 1.03 mmol, 1.0 equiv), Pd(OAc)₂ (35 mg, 0.155 mmol, 0.15 equiv), tris(2-methylphenyl)phosphane (63 mg, 0.206 mmol, 0.20 equiv) and tetrabutylammonium chloride (286 mg, 1.03 mmol, 1.0 equiv) was added DMF (2.9 mL). The resulting solution was stirred for 12 h at 115 °C. The resulting mixture was diluted with 30 mL EtOAc washed with 2 x 15 mL of 1 M aq. K₂CO₃ and the organic layer was concentrated and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column. The resulting crude product was purified by Flash-Prep-HPLC. This resulted in 76 mg (22%) of (E)-3-(3,3-dimethyl-2-oxoindolin-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide as a white solid. LC-MS (ES, m/z):

[M+H]⁺=339

¹H NMR (400 MHz, DMSO-*d*₆, ppm): δ 10.54 (s, 1H), 9.63 (s, 1H), 7.75 (d, *J* = 15.7 Hz, 1H), 7.48 (d, *J* = 8.1 Hz, 1H), 7.37 (d, *J* = 7.6 Hz, 1H) 7.28-7.17 (m, 2H), 7.09 (d, *J* = 1.5 Hz, 1H), 7.05-6.97 (m, 1H), 6.93 (d, *J* = 15.7 Hz, 1H), 2.16 (s, 3H), 1.27 (s, 6H).

5 **Example 4: (E)-N-(3-fluoro-2-methylphenyl)-3-(2'-oxospiro[cyclopropane-1,3'-indolin]-6'-yl)acrylamide**




To a microwave vial with N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 130 mg, 0.725 mmol, 1.0 equiv), 6'-bromo-1'H-spiro[cyclopropane-1,3'-indol]-2'-one (172 mg, 0.725 mmol, 1.0 equiv), Pd(OAc)₂ (24 mg, 0.109 mmol, 0.15 equiv) tris(2-methylphenyl)phosphane (44 mg, 0.145 mmol, 0.20 equiv) and tetrabutylammonium chloride (278 mg, 0.725 mmol, 1.0 equiv) was added DMF (2 mL). The resulting solution was stirred for 12 h at 115 °C. The resulting mixture was diluted with 30 mL EtOAc washed with 2 x 15 mL of 1 M aq. K₂CO₃ and the organic layer was concentrated and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column. The resulting crude product was purified by Flash-Prep-HPLC. This resulted in 10 mg (4%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(2'-oxospiro[cyclopropane-1,3'-indolin]-6'-yl)acrylamide as a white solid.

LC-MS (ES, *m/z*):

[M+H]⁺=337

20 ¹H NMR (400 MHz, DMSO-*d*₆, ppm): δ 10.77 (s, 1H), 9.62 (s, 1H), 7.57 (d, *J* = 15.7 Hz, 1H), 7.49 (d, *J* = 8.1 Hz, 1H), 7.29-7.18 (m, 2H), 7.16 (s, 1H), 7.06 (d, *J* = 7.7 Hz, 1H), 7.01 (t, *J* = 9.0 Hz, 1H), 6.95 (d, *J* = 15.7 Hz, 1H), 2.17 (d, *J* = 2.0 Hz, 3H), 1.66-1.60 (m, 2H), 1.55-1.50 (m, 2H).


25 **Example 5: (E)-N-(3-fluoro-2-methylphenyl)-3-(7-fluoro-2-oxoindolin-6-yl)acrylamide**



Prepared according to the methodology described above using **Intermediate 1** and 6-bromo-7-fluoro-1,3-dihydro-2*H*-Indol-2-one. LC-MS (ES, *m/z*): 327 [M-H]⁺

30 ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.59 (s, 1H), 9.72 (s, 1H), 7.64 (d, *J* = 15.9 Hz, 1H), 7.45-7.19 (m, 3H), 7.04-6.98 (m, 3H), 3.56 (s, 2H), 2.16 (s, 3H).

Example 6: (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide



5


Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, (50 mg, 0.28 mmol, 1.0 equiv), DMF (4 mL), 5-bromo-2-benzoxazolinone, (66 mg, 0.31 mmol, 1.10 equiv), Et₃N (0.12 mL, 0.84 mmol, 3.0 equiv), and Pd(dppf)Cl₂.CH₂Cl₂ (11 mg, 0.014 mmol, 0.05 equiv). The resulting solution was stirred for 12 h at 120 °C. The reaction was then quenched by the addition of 3 mL of water. The resulting solution was extracted with 5 mL of DCM and the organic layer was dried over anhydrous sodium sulfate and concentrated. The crude product was purified by Flash-Prep-HPLC. This resulted in 21 mg (24%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a white solid. LC MS (ES, m/z):

15 [M+H]⁺=313

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 11.89 (brs, 1H), 9.59 (s, 1H), 7.62 (d, *J* = 15.6 Hz, 1H), 7.48 (d, *J* = 8.1 Hz, 1H), 7.44-7.31 (m, 3H), 7.26-7.19 (m, 1H), 7.07-6.95 (m, 2H), 2.16 (s, 3H).

Example 7: (E)-N-(3-fluoro-2-methylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide

20



Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 90 mg, 0.50 mmol, 1.0 equiv), DMF (5 mL), 5-bromo-1-methyl-3H-1,3-benzodiazol-2-one (137 mg, 0.60 mmol, 1.20 equiv), Et₃N (0.21 mL, 1.51 mmol, 3.0 equiv), and Pd(dppf)Cl₂.CH₂Cl₂ (20 mg, 0.025 mmol, 0.05 equiv). The resulting solution was stirred for 12 h at 120 °C and cooled to 25 °C. The mixture was purified by Prep-HPLC. This resulted in 7 mg (4%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(1-methyl-2-oxo-2,3-


25

dihydro-1H-benzo[d]imidazol-5-yl)acrylamide as an off-white solid. LC MS (ES, m/z):


$[M+H]^+=326$

^1H NMR (300 MHz, DMSO- d_6 , ppm): δ 11.03 (s, 1H), 9.51 (s, 1H), 7.60 (d, J = 15.6 Hz, 1H), 7.49 (d, J = 8.1 Hz, 1H), 7.33 (dd, J = 8.1, 1.5 Hz, 1H), 7.29-7.16 (m, 3H), 7.0-6.98 (m, 1H), 6.88 (d, J = 15.6 Hz, 1H), 2.16 (s, 3H).


Example 8: (E)-N-(3-fluoro-2-methylphenyl)-3-(3-methyl-2-oxoindolin-6-yl)acrylamide



Step 1: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 6-bromo-1H-indole-2,3-dione (3.5 g, 15.48 mmol, 1.0 eq), THF (70 mL). The 3M Methylmagnesium bromide (5.2 mL, 15.48 mmol, 1.0 eq) was added dropwise at -78°C over 0.5 h. The resulting solution was allowed to react, with stirring, for an additional 5h at -78°C and then warmed to 25°C . The reaction was then quenched by the addition of 10 mL of 0.2 M HCl. The resulting solution was extracted with 2x50 mL of EtOAc, and the organic was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE (35/65). This resulted in 3 g (80%) of 6-bromo-3-hydroxy-3-methyl-1H-indol-2-one as an off-white solid. LC-MS (ES, m/z): $[M-H]^+=240$



Step 2: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 6-bromo-3-hydroxy-3-methyl-1H-indol-2-one (2.0 g, 8.26 mmol, 1.0 eq) and THF (30 mL). This was followed by the addition of diethylaminosulfur trifluoride (DAST) (2.0 g, 12.39 mmol, 1.50 eq) at -78°C . The resulting solution was stirred for 1 h from -78°C and warmed to 25°C . The reaction was then quenched by the addition of 10 mL of Sat. NaHCO_3 . The resulting solution was extracted with 2x50 mL of EtOAc, and the organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE (25/75). This resulted in 1.65 g (82%) of 6-bromo-3-fluoro-3-methyl-1H-indol-2-one as a white solid. LC-MS (ES, m/z): $[M-H]^+=242$



Step 3: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed 6-bromo-3-fluoro-3-methyl-1H-indol-2-one (200 mg, 0.82 mmol, 1.0 eq), N-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 147 mg, 0.82 mmol, 1.0 eq), Pd(dppf)Cl₂·CH₂Cl₂ (13 mg, 0.02 mmol, 0.02 eq), DMF (4 mL), Et₃N (0.23 mL, 1.64 mmol, 2.0 eq). The resulting solution was stirred for 2 h at 110 °C. The solids were filtered out. The filtrate was concentrated. The crude product was purified by Flash-Prep-HPLC. This resulted in 13 mg (4.5%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(3-fluoro-3-methyl-2-oxoindolin-6-yl)acrylamide and 12 mg (4%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(3-methyl-2-oxoindolin-6-yl)acrylamide.

(E)-N-(3-fluoro-2-methylphenyl)-3-(3-fluoro-3-methyl-2-oxoindolin-6-yl)acrylamide

LC-MS (ES, m/z):

[M+H]⁺=343

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.87 (s, 1H), 9.68 (s, 1H), 7.60-7.46 (m, 3H), 7.34-7.19 (m, 2H), 7.12 (s, 1H), 7.04-6.97 (m, 2H), 2.16 (s, 3H), 1.70 (d, *J* = 22.5 Hz, 3H).


(E)-N-(3-fluoro-2-methylphenyl)-3-(3-methyl-2-oxoindolin-6-yl)acrylamide

LC-MS (ES, m/z):

[M+H]⁺=325

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.53 (s, 1H), 9.62 (s, 1H), 7.58-7.46 (m, 2H), 7.34-7.21 (m, 3H), 7.06-6.90 (m, 3H), 3.48-3.45 (m, 1H), 2.16 (s, 3H), 1.34 (d, *J* = 7.8 Hz, 3H).

Example 9: (E)-N-(3-chloro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-chloro-2-methylphenyl)prop-2-enamide (**Intermediate 2**, 55 mg, 0.28 mmol, 1.0 equiv), DMF (4 mL), 2-benzoxazolinone, 5-bromo- (66 mg, 0.31 mmol, 1.10 equiv), Et₃N (0.12 mL, 0.84 mmol, 3.0 equiv) and Pd(dppf)Cl₂.CH₂Cl₂ (11 mg, 0.014 mmol, 0.05 equiv). The resulting solution was stirred for 12 h at 120 °C. The reaction was then quenched by the addition of 3 mL of water. The resulting solution was extracted with 5 mL of DCM and the organic layer was dried over anhydrous sodium sulfate and concentrated. The crude product was purified by Flash-Prep-HPLC. This resulted in 21 mg (24%) of (E)-N-(3-chloro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a white solid. LC MS (ES, m/z):

[M+H]⁺=329

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 11.83 (s, 1H), 9.68 (s, 1H), 7.61 (d, *J* = 15.6 Hz, 1H), 7.53-7.51 (m, 1H), 7.43-7.30 (m, 5H), 6.91 (d, *J* = 15.6 Hz, 1H), 2.28 (s, 3H).


Example 10: (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide




Into a 40-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-3-fluoro-2-methylphenyl)prop-2-enamide (**Intermediate 1**, (500 mg, 2.79 mmol, 1.0 eq), 6-bromo-1,3-dihydroindol-2-one (592 mg, 2.79 mmol, 1.0 eq), DMF (20 mL), Et₃N (1.2 mL, 8.37 mmol, 3.0 eq) and Pd(dppf)Cl₂ (41 mg, 0.05 mmol, 0.02 eq). The reaction mixture was stirred for 2 h at 110 °C. The resulting mixture was concentrated. The crude product was purified by Flash-Prep-HPLC. This resulted in 102 mg (11%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide as a white solid. LC-MS (ES, m/z): [M-
H]⁺=309

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.57 (s, 1H), 9.62 (s, 1H), 7.55 (d, *J*=15.9 Hz, 1H), 7.48-7.45 (m, 1H), 7.29-7.18 (m, 3H), 7.05-6.89 (m, 3H), 3.52 (s, 2H), 2.26 (s, 3H).


Example 11: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide




Step 1: Into a 500-mL sealed tube, was placed 5-bromobenzo[d]oxazol-2(3H)-one (10.0 g, 46.96 mmol, 1.0 equiv), methyl acrylate (12.1 g, 140.8 mmol, 3.0 equiv), Et₃N (19.6 mL, 140.8 mmol, 3.0 equiv), Pd(dppf)Cl₂ (350 mg, 4.7 mmol, 0.01 equiv) in DMF (200 mL). The resulting solution was stirred for 2 h at 120 °C. The resulting solution was concentrated. The residue was applied onto a silica gel column eluting with THF/hexane (20/80). This resulted in 11g (82%) of methyl (E)-3-(3-(3-methoxy-3-oxopropyl)-2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate as a yellow solid.



Step 2: Into a 500 mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed THF (300 mL), methyl (E)-3-(3-(3-methoxy-3-oxopropyl)-2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate (11.0 g, 36.1 mmol, 1.0 equiv), t-BuOK (13.2 g, 108.2 mmol, 3.0 equiv). The resulting solution was stirred for 2 h at 60 °C. The reaction was then quenched by the addition of 600 mL of Satd. NH₄Cl. The resulting solution was extracted with 2x300 mL of EtOAc. The organic layer was washed with 2x400 ml of Water. The organic layer was dried over anhydrous sodium sulfate and concentrated. The resulting solution was concentrated. The residue was applied onto a silica gel column with EtOAc/hexane (40/60). This resulted in 5.5 g (69%) of methyl (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate as a light-red solid.




Step 3: Into a 100 mL 3-necked round-bottom flask purged, was placed THF (20 mL), methyl (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate (2.5 g, 11.4 mmol, 1.0 equiv), 2 M NaOH (17.1 mL, 34.2 mmol, 3.0 equiv). The resulting solution was stirred for 2 h at 25 °C. The reaction mixture was concentrated under low temperature (<30 °C). The residue was dissolved into water (30 mL), and the pH was adjusted to 2-3 with 2 M HCl. The solid was collected. This resulted in 2.5 g (65%) of (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylic acid as an off-white solid.



Step 4: In each vial was added (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylic acid (30 mg, 0.146 mmol, 1.0 equiv) in DMF (2 mL). Then T₃P (70 mg, 0.22 mmol, 1.50 equiv) and DIPEA (28 mg, 0.22 mmol, 1.50 equiv), 1-indanamine (0.161 mmol, 1.1 equiv) were added. The reaction mixture was stirred for 2 h at room temperature. The reaction mixture was quenched by water and extracted with EtOAc, the organic layer was concentrated under vacuum to afford crude product. The crude product was then purified by Prep-HPLC directly. The collected fraction was lyophilized to get the final compounds. LC-MS (ES, *m/z*): 321 [M+H]⁺


¹H NMR: (300 MHz, DMSO-*d*₆, *ppm*): δ 11.77 (s, 1H), 8.44 (d, *J* = 8.4 Hz, 1H), 7.51 (d, *J* = 15.6 Hz, 1H), 7.34-7.18 (m, 7H), 6.65 (d, *J* = 15.9 Hz, 1H), 5.42-5.33 (m, 1H), 2.96-2.81 (m, 2H), 2.48-2.42 (m, 1H), 1.85-1.81 (m, 1H).

General Procedure A:



To each vial was added methyl (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate (**Example 11, Step 2**, 30 mg, 0.14 mmol, 1.0 equiv) in THF (2 mL). The amine (0.18 mmol, 1.30 equiv) was added and the reaction mixture was cooled to 0°C. Then 1M LiHMDS (0.68 mL, 0.68 mmol, 5.0 equiv) was added. The reaction mixture was stirred for 1 h at 25 °C. The reaction mixture was quenched by water and extracted with EtOAc, the organic was concentrated under vacuum to afford crude product which was then purified by Prep-HPLC. The collected fraction was lyophilized to get the final compounds.


Example 12: (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)-*N*-(*o*-tolyl)acrylamide



Synthesised using general procedure A using 2-methyl aniline and (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate to give (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)-N-(o-tolyl)acrylamide as a solid. LC-MS (ES, m/z): 295 $[M+H]^+$

^1H NMR: (300 MHz, DMSO- d_6 , ppm): δ 9.40 (brs, 1H), 7.63-7.57 (m, 2H), 7.38-7.32 (m, 3H),
 5 7.25-6.91 (m, 4H), 2.26(s, 3H).


Example 13: (E)-N-(2-isopropylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide



Synthesised using general procedure A using 2-isopropyl aniline and (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate to give (E)-N-(2-isopropylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a solid. LC-MS (ES, m/z): 323 $[M+H]^+$

15 ^1H NMR (300 MHz, DMSO- d_6 , ppm): δ 11.83 (brs, 1H), 9.48 (brs, 1H), 7.59 (d, $J=15.6$ Hz, 1H), 7.40-7.34 (m, 5H), 7.24-7.18 (m, 2H), 6.93 (d, $J=15.3$ Hz, 1H), 3.23-.319 (m, 1H), 1.17 (d, $J=6.6$ Hz, 6H)


Example 14: (E)-N-(2-isopropyl-6-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide



Synthesised using general procedure A using 2-isopropyl-6-methyl aniline and (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate to give (E)-N-(2-isopropyl-6-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a solid. LC-MS (ES, m/z): 337 $[M+H]^+$

25 ^1H NMR (300 MHz, DMSO- d_6 , ppm): δ 11.82 (brs, 1H), 9.43 (brs, 1H), 7.57 (d, $J=15.6$ Hz, 1H), 7.40-7.33 (m, 3H), 7.27-7.09 (m, 3H), 6.85 (d, $J=15.6$ Hz, 1H), 3.15-.3.06 (m, 1H), 2.18 (s, 3H), 1.173 (d, $J=6.9$ Hz, 6H)


Example 15: (E)-N-(5-chloro-2-isopropylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide



- 5 Synthesised using general procedure A using 5-chloro-2-isopropyl aniline and (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate to give (E)-N-(5-chloro-2-isopropylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a solid. LC-MS (ES, m/z): 357[M+H]⁺
¹H NMR (300 MHz, DMSO-*d*₆, *ppm*): δ 11.79 (brs, 1H), 9.56 (brs, 1H), 7.64-7.58 (m, 2H), 7.37-7.23 (m, 5H), 6.95 (d, $J=15.6$ Hz, 1H), 3.25-3.20 (m, 1H), 1.16 (d, $J=6.9$ Hz, 6H).

10


Example 16: (E)-N-(4,5-difluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide



- 15 Synthesised using general procedure A using 4,5-difluoro-2-methyl aniline and (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate to give (E)-N-(4,5-difluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a solid. LC-MS (ES, m/z): 331[M+H]⁺, 373 [M+CH₃CN]⁺.


Example 17: (E)-N-(5-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide

20



- 25 Synthesised using general procedure A using 5-fluoro-2-methyl aniline and (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate to give (E)-N-(5-fluoro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a solid. LC-MS (ES, m/z): 311[M+H]⁺

25


Example 18: (E)-N-(3-fluoro-2-methylphenyl)-3-(4-fluoro-2-oxoindolin-6-yl) acrylamide

Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-2-methylphenyl)acrylamide (60.0 mg, 0.34 mmol, 1.0 eq), 6-bromo-4-fluoroindolin-2-one (78 mg, 0.34 mmol, 1.0 eq), Et₃N (0.14 mL, 1.0 mmol, 3.0 eq), DMF (4.0 mL), Pd(dppf)Cl₂ (25.2 mg, 0.034 mmol, 0.10 eq). The resulting solution was stirred for 2 h at 120 °C. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 11.8 mg (11%) of (E)-N-(3-fluoro-2-methylphenyl)-3-(4-fluoro-2-oxoindolin-6-yl)acrylamide as an off-white solid.


LC-MS (ES, m/z):

[M+H]⁺=329

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.80 (brs, 1H), 9.66 (s, 1H), 7.55 (d, *J*=15.9 Hz, 1H), 7.47 (d, *J*=8.4 Hz, 1H), 7.24-7.22 (m, 1H), 7.10-6.93 (m, 4H), 3.60 (s, 2H), 2.15 (s, 3H)


Example 19: (E)-N-(2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide

Step 1: Into a 500-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 6-bromo-1,3-dihydroindol-2-one (20.0 g, 94.32 mmol, 1.0 equiv), DMF (350 mL), methyl acrylate (6.50 g, 75.50 mmol, 0.80 equiv), Et₃N (26.3 mL, 188.64 mmol, 2.0 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (0.77 g, 0.94 mmol, 0.01 equiv). The resulting solution was stirred for 2 hr at 120 °C. The reaction mixture was cooled to RT. The mixture was applied onto a silica gel column with THF/PE (2/1). This resulted in 7.4 g (36%) of methyl (2E)-3-(2-oxo-1,3-dihydroindol-6-yl)prop-2-enoate as a light yellow solid.



Step 2: Into a 250-mL round-bottom flask, was placed methyl (2E)-3-(2-oxo-1,3-dihydroindol-6-yl)prop-2-enoate (3.0 g, 13.81 mmol, 1.0 equiv), MeOH/H₂O (80/40 mL), NaOH (1.7 g, 41.43 mmol, 3.0 equiv). The resulting solution was stirred for 12 h at 25 °C. The resulting mixture was


concentrated. The resulting solution was diluted with 40 mL of water. The resulting solution was extracted with 2x50 mL of DCM. The pH value of the aqueous phase was adjusted to 5 with 2M HCl. The solids were collected by filtration. This resulted in 1.8 g (64%) of (2E)-3-(2-oxo-1,3-dihydroindol-6-yl)prop-2-enoic acid as a light yellow solid.



Step 3: Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen was added (E)-3-(2-oxoindolin-6-yl)acrylic acid (60.0 mg, 0.30 mmol, 1.0 equiv) in DMF (4.0 mL). Then HATU (167.6 mg, 0.44 mmol, 1.50 equiv) and DIPEA (56.8 mg, 0.44 mmol, 1.50 equiv), 2,6-dimethylaniline (39.5 mg, 0.33 mmol, 1.10 equiv) were added. The reaction mixture was stirred for 2 h at room temperature. The crude mixture was then purified by Prep-HPLC. This resulted in 12 mg (13%) of (E)-N-(2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide as a white solid. LC-MS (ES, m/z): [M+H]⁺=307

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 10.53 (brs, 1H), 9.47 (brs, 1H), 7.51 (d, *J*=15.9 Hz, 1H), 7.29-7.18 (m, 2H), 7.10-7.05 (m, 4H), 6.83 (d, *J*=15.9 Hz, 1H), 3.52 (s, 2H), 2.18 (s, 6H).

Example 20: (E)-N-(3-fluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide




Into a 8-mL round-bottom flask, was placed 6-bromoindolin-2-one (100 mg, 0.47 mmol, 1.0 equiv), N-(3-fluoro-2,6-dimethylphenyl)prop-2-enamide (Prepared according to the procedure described for **Intermediate 1** from 2,6-dimethyl-3-fluoroaniline and acryloyl chloride; 90.7 mg, 0.47 mmol, 1.0 equiv), Pd(dppf)Cl₂ (34.4 mg, 0.047 mmol, 0.10 equiv), Et₃N (0.2 mL, 1.41 mmol, 3.0 equiv), DMF (4.0 mL). The resulting solution was stirred for 2 hr at 120 °C in an oil bath. The mixture was applied onto a silica gel column with PE/THF (1/1). This resulted in 11 mg (7%) of (E)-N-(3-fluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide as a solid.

LC-MS (ES, m/z):


[M+H]⁺=325

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.55 (s, 1H), 9.63 (s, 1H), 7.55 (d, *J* = 15.6 Hz, 1H), 7.29-7.03 (m, 5H), 6.84 (d, *J* = 15.9 Hz, 1H), 3.53 (s, 2H), 2.14 (s, 3H), 2.07 (s, 3H).


Example 21: (E)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide

Into a 8-mL vial, was placed (E)-3-(2-oxoindolin-6-yl)acrylic acid (50.0 mg, 0.25 mmol, 1.0 equiv), 2-methyl-2,3-dihydro-1H-inden-1-amine hydrochloride (45.20 mg, 0.25 mmol, 1.0 equiv), HATU (141.38 mg, 0.37 mmol, 1.50 equiv) and DIPEA (95.41 mg, 0.74 mmol, 3.0 equiv) in DMF (2.0 mL). The resulting solution was stirred for 2 h at 20 °C. The mixture was purified by Flash-Prep-HPLC. This resulted in 27 mg (33%) of ((E)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=333

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 10.51 (brs, 1H), 8.45-8.20 (m, 1H), 7.50-7.45 (m, 1H), 7.26-7.13 (m, 6H), 7.05-6.96 (m, 1H), 6.68-6.63 (m, 1H), 5.45-5.0 (m, 1H), 3.51 (s, 2H), 3.10-3.0 (m, 1H), 2.70-2.20 (m, 2H), 1.20-0.90 (m, 3H).


Example 22: (E)-3-(1-ethyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-N-(3-fluoro-2-methylphenyl)acrylamide

Step 1: Into a 100-mL sealed tube, was placed 4-bromo-1-fluoro-2-nitrobenzene (2.0 g, 9.09 mmol, 1.0 eq), EtOH (8.0 mL, 1.0 eq), ethylamine (22.73 mL, 45.45 mmol, 5.0 eq, 2 M in ethanol). The resulting solution was stirred for 4 h at 50 °C. The resulting mixture was concentrated. The residue was diluted with 15 mL of H₂O and stirred 15 min. The solids were collected by filtration. This resulted in 2.2 g (98%) of 4-bromo-N-ethyl-2-nitroaniline as a red solid. LC-MS (ES, m/z): [M+H]⁺=245



Step 2: Into a 50-mL round-bottom flask, was placed 4-bromo-N-ethyl-2-nitroaniline (2.0 g, 8.16 mmol, 1.0 eq), acetone (16.0 mL), H₂O (2.0 mL), NH₄Cl (4.37 g, 81.61 mmol, 10.0 eq), Zn (2.67 g, 40.80 mmol, 5.0 eq). The resulting solution was stirred for 3 h at room temperature. The


solids were filtered out. The filtrate was concentrated. The residue was diluted with 20 mL of H₂O and stirred 15 min. The solids were collected by filtration. This resulted in 1.1 g (63%) of 4-bromo-N¹-ethylbenzene-1,2-diamine as a light yellow solid. LC-MS (ES, m/z): [M+H]⁺=215



5

Step 3: Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 4-bromo-N¹-ethylbenzene-1,2-diamine (400.0 mg, 1.86 mmol, 1.0 eq), DCM (8.0 mL), Triphosgene (441.5 mg, 1.48 mmol, 0.80 eq). This was followed by the addition of Et₃N (0.8 mL, 5.58 mmol, 3.0 eq) at 0 °C. The resulting solution was stirred for 3 h at room temperature. The reaction was then quenched by the addition of 5 mL of water. The organic phase was separated and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with (THF/PE=18/82). This resulted in 210 mg (47%) of 5-bromo-1-ethyl-3H-1,3-benzodiazol-2-one as a light brown solid.

LC-MS (ES, m/z): [M+H]⁺=240



15


Step 4: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-2-methylphenyl)acrylamide (120.0 mg, 0.67 mmol, 1.0 eq), 5-bromo-1-ethyl-3H-1,3-benzodiazol-2-one (161.5 mg, 0.67 mmol, 1.0 eq), Et₃N (0.28 mL, 2.01 mmol, 3.0 eq), DMF (5.0 mL), Pd(dppf)Cl₂ (9.8 mg, 0.01 mmol, 0.02 eq). The resulting solution was stirred for 2 h at 120 °C. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 25 mg (11%) of (E)-3-(1-ethyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-N-(3-fluoro-2-methylphenyl)acrylamide as an off-white solid. LC-MS (ES, m/z):

[M+H]⁺=340


¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 11.05 (brs, 1H), 9.52 (s, 1H), 7.62-7.48 (m, 2H), 7.33-7.19 (m, 4H), 7.11-6.85 (m, 2H), 3.88-3.81 (m, 2H), 2.27 (s, 3H), 1.31-1.19 (m, 3H).

25


Example 23: (E)-3-(1-cyclopropyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-N-(3-fluoro-2-methylphenyl)acrylamide



- 5 **Step 1:** Into a 40-mL sealed tube, was placed 4-bromo-1-fluoro-2-nitrobenzene (2.0 g, 9.09 mmol, 1.0 eq), aminocyclopropane (2.60 g, 45.45 mmol, 5.0 eq), EtOH (20.0 mL). The resulting solution was stirred for 4 h at 50 °C. The resulting mixture was concentrated. The residue compound was diluted with 15 mL of H₂O and stirred 15 min. The solids were collected by filtration. The solid was dried in an oven under reduced pressure. This resulted in 2.4 g (96%) of
10 4-bromo-N-cyclopropyl-2-nitroaniline as a red solid. LC-MS (ES, m/z): [M+H]⁺=257




- Step 2:** Into a 40-mL sealed tube, was placed 4-bromo-N-cyclopropyl-2-nitroaniline (2.0 g, 7.8
15 mmol, 1.0 eq), acetone (16.0 mL), H₂O (2.0 mL), Zn (2.54 g, 38.89 mmol, 5.0 eq), NH₄Cl (4.16 g, 77.8 mmol, 10.0 eq). The resulting solution was stirred for 3 h at 25 °C. The solids were filtered out. The filtrate was concentrated. The residue was diluted with 20 mL of H₂O and stirred 15 min. The solids were collected by filtration. This resulted in 1 g (56%) of 4-bromo-N¹-cyclopropylbenzene-1,2-diamine as a light brown solid.
20 LC-MS (ES, m/z): [M+H]⁺=227



- Step 3:** Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert
25 atmosphere of nitrogen, was placed 4-bromo-N¹-cyclopropylbenzene-1,2-diamine (300.0 mg, 1.32 mmol, 1.0 eq), Triphosgene (313.60 mg, 1.06 mmol, 0.80 eq), DCM (6.0 mL). This was followed by the addition of Et₃N (0.6 mL, 3.96 mmol, 3.0 eq) at 0 °C. The resulting solution was stirred for 3 h at room temperature. The reaction was then quenched by the addition of 5 mL of water. The organic phase was separated and dried over anhydrous sodium sulfate and

concentrated. This resulted in 180 mg (53%) of 5-bromo-1-cyclopropyl-3H-1,3-benzodiazol-2-one as a light brown solid. LC-MS (ES, m/z): $[M+H]^+=253$




- 5 **Step 4:** Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-2-methylphenyl)acrylamide (120.0 mg, 0.67 mmol, 1.0 eq), 5-bromo-1-cyclopropyl-3H-1,3-benzodiazol-2-one (169.5 mg, 0.67 mmol, 1.0 eq), Et₃N (0.2 mL, 1.34 mmol, 2.0 eq), DMF (5.0 mL), Pd(dppf)Cl₂ (9.80 mg, 0.013 mmol, 0.02 equiv). The resulting solution was stirred for 2 h at 120 °C. The solids were filtered out. The crude mixture was
- 10 purified by Flash-Prep-HPLC. This resulted in 21 mg (9%) of (E)-3-(1-cyclopropyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-N-(3-fluoro-2-methylphenyl)acrylamide as an off-white solid. LC-MS (ES, m/z):


$[M+H]^+=352$

- ¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.96 (brs, 1H), 9.54 (brs, 1H), 7.61-7.47 (m, 2H), 7.33-7.18 (m, 4H), 7.02-6.85 (m, 2H), 2.95-2.87 (m, 1H), 2.28 (s, 3H), 1.04-0.88 (m, 4H).
- 15

Example 24: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide



- 20 **Step 1:** Into a 500-mL 3-necked round-bottom flask, acryloyl chloride (6.80 g, 75.08 mmol, 1.0 equiv), was added to indanamine (10.0 g, 75.08 mmol, 1.0 equiv) and Et₃N (20.9 mL, 150.16 mmol, 2.0 equiv) in DCM (200.0 mL) at 0 °C. The resulting solution was stirred for 15 h at 25 °C. The reaction was then quenched by the addition of 100 mL of water/ice. The resulting solution was extracted with 2x100 mL of dichloromethane and the organic layer was dried over
- 25 anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE(10/1, resulting in 7.3 g (52%) of N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide as a white solid. LC-MS (ES, m/z): $[M+H]^+=188$




Into a 8-mL vial, was placed N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (100.0 mg, 0.53 mmol, 1.0 equiv), 5-bromo-1-methyl-3H-1,3-benzodiazol-2-one (121.27 mg, 0.53 mmol, 1.0 equiv), Pd(dppf)Cl₂ (39.08 mg, 0.05 mmol, 0.10 equiv) and Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv) in DMF (4.0 mL, 25.844 mmol). The resulting solution was stirred for 15 h at 120 °C in an oil bath. The reaction mixture was cooled. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 7.8 mg (4%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide as a white solid. LC-MS (ES, m/z):

[M+H]⁺=334

¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.98 (brs, 1H), 8.37 (d, *J* = 8.4 Hz, 1H), 7.50 (d, *J* = 15.9 Hz, 1H), 7.34-7.03 (m, 7H), 6.56 (d, *J* = 15.6 Hz, 1H), 5.42-5.39 (m, 1H), 3.29 (s, 3H), 3.02-2.77 (m, 2H), 2.47-2.37 (m, 1H), 1.93-1.72 (m, 1H).

Example 25: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Example 24, Step 1**, 100.0 mg, 0.53 mmol, 1.0 equiv), 6-bromo-3-methyl-1H-indazole (135.27 mg, 0.64 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (87.23 mg, 0.11 mmol, 0.20 equiv), DMF (4.0 mL), Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 64 mg (38%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a off-white solid. LC-MS (ES, m/z):

[M+H]⁺=318

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.77 (brs, 1H), 8.46 (d, *J* = 8.4 Hz, 1H), 7.73-7.60 (m, 3H), 7.31-7.17 (m, 5H), 6.74 (d, *J* = 15.9 Hz, 1H), 5.46-5.38 (m, 1H), 3.02-2.93 (m, 1H), 2.89-2.79 (m, 1H), 2.51 (s, 3H), 2.48-2.41 (m, 1H), 1.90-1.77 (m, 1H).

5 **Example 26: (E)-3-(3-cyano-1H-indazol-6-yl)-N-(2,3-dihydro-1H-inden-1-yl)acrylamide**




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Example 24, Step 1**, 100.0 mg, 0.53 mmol, 1.0 equiv), 6-bromo-1H-indazole-3-carbonitrile (142.30 mg, 0.64 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (43.61 mg, 0.05 mmol, 0.10 equiv), DMF (4 mL), Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 24 mg (14%) of (E)-3-(3-cyano-1H-indazol-6-yl)-N-(2,3-dihydro-1H-inden-1-yl)acrylamide as an off-white solid. LC-MS (ES, *m/z*):

[M+H]⁺=329

¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 14.33 (brs, 1H), 8.55 (d, *J* = 8.1 Hz, 1H), 7.93-7.91 (m, 2H), 7.70 (d, *J* = 15.9 Hz, 1H), 7.62-7.59 (m, 1H), 7.30-7.17 (m, 4H), 6.84 (d, *J* = 15.6 Hz, 1H), 5.47-5.39 (m, 1H), 3.03-2.94 (m, 1H), 2.90-2.80 (m, 1H), 2.47-2.42 (m, 1H), 1.91-1.76 (m, 1H).

20 **Example 27: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide**




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Example 24, Step 1**, 100.0 mg, 0.53 mmol, 1.0 equiv), 5-bromo-6-fluoro-3H-1,2,3-benzotriazole (138.44 mg, 0.64 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (43.61 mg, 0.05 mmol, 0.10 equiv), DMF (4.0 mL), Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv). The resulting solution was stirred overnight at 140 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted

in 6.3 mg (4%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(5-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide as a off-white solid. LC-MS (ES, m/z):

[M+H]⁺=323

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.63 (d, *J* = 8.1 Hz, 1H), 8.27 (d, *J* = 6.6 Hz, 1H), 7.82 (d, *J* = 10.8 Hz, 1H), 7.68 (d, *J* = 16.2 Hz, 1H), 7.29-7.18 (m, 4H), 6.88 (d, *J* = 15.9 Hz, 1H), 5.47-5.40 (m, 1H), 3.01-2.94 (m, 1H), 2.90-2.79 (m, 1H), 2.49-2.44 (m, 1H), 1.92-1.79 (m, 1H).

Example 28: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-(trifluoromethyl)-1H-indazol-6-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Example 24, Step 1**, 100.0 mg, 0.53 mmol, 1.0 equiv), 6-bromo-3-(trifluoromethyl)-1H-indazole (169.9 mg, 0.64 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (43.61 mg, 0.05 mmol, 0.10 equiv), DMF (4.0 mL), Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 80 mg (40%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-(trifluoromethyl)-1H-indazol-6-yl)acrylamide as an off-white solid. LC-MS (ES, m/z):

[M+H]⁺=372

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 14.16 (brs, 1H), 8.55 (d, *J* = 8.1 Hz, 1H), 7.89-7.83 (m, 2H), 7.70 (d, *J* = 15.9 Hz, 1H), 7.58-7.54 (m, 1H), 7.30-7.17 (m, 4H), 6.83 (d, *J* = 15.6 Hz, 1H), 5.47-5.39 (m, 1H), 3.03-2.94 (m, 1H), 2.90-2.80 (m, 1H), 2.47-2.42 (m, 1H), 1.91-1.78 (m, 1H).

Example 29: (E)-N-(2,6-dimethylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide




Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed N-(2,6-dimethylphenyl)prop-2-enamide ((Prepared as for Intermediate 1 from acryloyl chloride and 2,6-dimethylaniline) 50.0 mg, 0.29 mmol, 1.0 equiv), DMF (4.0 mL), 5-bromo-1-methyl-3H-1,3-benzodiazol-2-one (71.3 mg, 0.31 mmol, 1.10 equiv), Et₃N (0.12 mL, 0.86 mmol, 3.0 equiv),
 5 Pd(dppf)Cl₂.CH₂Cl₂ (11.65 mg, 0.014 mmol, 0.05 equiv). The resulting solution was stirred for 5 h at 120 °C. The reaction mixture was cooled to RT. The crude mixture was applied onto a silica gel column with EtOAc/PE (1/1). This resulted in 20 mg (22%) of (E)-N-(2,6-dimethylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide as an off-white solid.

LC-MS (ES, m/z):


10 [M+H]⁺=322

¹H NMR (300 MHz, DMSO-*d*₆, ppm): 11.02 (s, 1H), 9.38 (s, 1H), 7.56 (d, *J* = 15.6 Hz, 1H), 7.33-7.31 (m, 1H), 7.24 (s, 1H), 7.16-7.09 (m, 4H), 6.78 (d, *J* = 15.6 Hz, 1H), 3.32 (s, 1H), 2.17 (s, 6H).

15 **Example 30: (E)-N-(3-fluoro-2,6-dimethylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide**




Step 1: Into a 40-mL vial, was placed 1-bromo-4-fluoro-3-methyl-2-nitrobenzene (500.0 mg, 2.14 mmol, 1.0 equiv), trimethyl-1,3,5,2,4,6-trioxatriborinane (268.20 mg, 2.13 mmol, 1.0 equiv),
 20 K₂CO₃ (590.6 mg, 4.27 mmol, 2.0 equiv) and Pd(dppf)Cl₂ (156.3 mg, 0.21 mmol, 0.10 equiv) in dioxane (10.0 mL) and H₂O (2.0 mL). The resulting solution was stirred for 1 h at 110 °C in an oil bath. The crude mixture was applied onto a silica gel column with EtOAc/PE (1/10). This resulted in 250 mg (69%) of 1-fluoro-2,4-dimethyl-3-nitrobenzene as a solid.




25 **Step 2:** Into a 40-mL vial, was placed 1-bromo-4-fluoro-3-methyl-2-nitrobenzene (0.50 g, 2.14 mmol, 1.0 equiv) in AcOH (10.0 mL) was added Fe (596.6 mg, 10.68 mmol, 5.0 equiv), The resulting solution was stirred for 1 h at RT. The solids were filtered out. The resulting solution was diluted with 20 mL water and extracted with 40 mL of EtOAc. The organic layer was dried and concentrated. This resulted in 250 mg (84%) of 3-fluoro-2,6-dimethylaniline as a solid.

30 LC-MS (ES, m/z): [M+H]⁺=140



Step 3: Into a 8-mL vial, was placed 6-bromo-3-fluoro-2-methylaniline (100.0 mg, 0.49 mmol, 1.0 equiv) and Et₃N (0.2 mL, 1.47 mmol, 3.0 equiv) in DCM (4 mL) was added acryloyl chloride (53.2 mg, 0.59 mmol, 1.20 equiv), The resulting solution was stirred for 1 h at 0 °C. The resulting mixture was concentrated under vacuum. The residue was purified with flash chromatography (PE/EA=10/1). This resulted in 120 mg (126.72%) of *N*-(3-fluoro-2,6-dimethylphenyl)prop-2-enamide as a white solid. LC-MS (ES, *m/z*): [M+H]⁺=194




Step 4: Into a 8-mL round-bottom flask, was placed 5-bromo-1-methyl-3H-1,3-benzodiazol-2-one (100 mg, 0.44 mmol, 1.0 equiv), *N*-(3-fluoro-2,6-dimethylphenyl)prop-2-enamide (85.1 mg, 0.44 mmol, 1.0 equiv), Pd(dppf)Cl₂ (32.2 mg, 0.044 mmol, 0.10 equiv), Et₃N (0.18 mL, 1.32 mmol, 3.0 equiv), DMF (4.0 mL). The resulting solution was stirred for 2 h at 120 °C in an oil bath. The mixture was applied onto a silica gel column with PE/THF (1/1). This resulted in 45 mg (30%) of (E)-*N*-(3-fluoro-2,6-dimethylphenyl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide as a solid. LC-MS (ES, *m/z*):

[M+H]⁺=340


¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 11.04 (s, 1H), 9.53 (s, 1H), 7.58 (d, *J* = 15.3 Hz, 1H), 7.33 (d, *J* = 8.4 Hz, 1H), 7.24 (s, 1H), 7.17-7.01 (m, 3H), 6.79 (d, *J* = 15.6 Hz, 1H), 3.31 (s, 3H), 2.15 (s, 3H), 2.07 (s, 3H).

Example 31: (E)-*N*-(2,6-dimethylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide



Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 2,6-dimethylaniline (160.0 mg, 1.32 mmol, 1.0 equiv), DCM (10.0 mL), Et₃N (0.55 mL, 3.96 mmol, 3.0 equiv). This was followed by the addition of acryloyl chloride (143.40 mg, 1.58 mmol, 1.20 equiv) dropwise with stirring at 0 °C. The resulting solution was

stirred for 3 h at 25 °C. The reaction was then quenched by the addition of 15 mL of water. The resulting solution was extracted with 2x10 mL of dichloromethane and the organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE (1/3). This resulted in 150 mg (65%) of *N*-(2,6-dimethylphenyl)prop-2-enamide as a white solid. LC-MS (ES, *m/z*): [M+H]⁺=176




Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed *N*-(2,6-dimethylphenyl)prop-2-enamide (50.0 mg, 0.29 mmol, 1.0 equiv), DMF (5.0 mL), 6-bromo-3-methyl-1H-indazole (66.3 mg, 0.31 mmol, 1.10 equiv), Et₃N (0.12 mL, 0.86 mmol, 3.0 equiv) and Pd(dppf)Cl₂.CH₂Cl₂ (11.7 mg, 0.014 mmol, 0.05 equiv). The resulting solution was stirred for 5 h at 120 °C. The reaction mixture was cooled to 25 °C. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 30 mg (34%) of (E)-*N*-(2,6-dimethylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as an off-white solid. LC-MS (ES, *m/z*):

[M+H]⁺=306

¹H NMR (300 MHz, DMSO-*d*₆, ppm): 12.81 (s, 1H), 9.49 (s, 1H), 7.77-7.67 (m, 3H), 7.40 (d, *J*=8.4 Hz, 1H), 7.11 (s, 3H), 6.97 (d, *J* = 15.6 Hz, 1H), 2.51 (s, 3H), 2.19 (s, 6H).

Example 32: (E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(3-ethyl-1H-indazol-6-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed *N*-(**Example 24, Step 1**, 2,3-dihydro-1H-inden-1-yl)prop-2-enamide (100.0 mg, 0.53 mmol, 1.0 equiv), 6-bromo-3-ethyl-1H-indazole (144.3 mg, 0.64 mmol, 1.20 equiv), Pd(dppf)Cl₂.CH₂Cl₂ (43.6 mg, 0.05 mmol, 0.10 equiv), DMF (4.0 mL) and Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude product was purified by Prep-HPLC. This resulted in 86 mg (49%) of (E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(3-ethyl-1H-indazol-6-yl)acrylamide as a

off-white solid. LC MS (ES, m/z):

$[M+H]^+=332$

^1H NMR (300 MHz, DMSO- d_6 , ppm): δ 12.77 (brs, 1H), 8.47 (d, J = 8.1 Hz, 1H), 7.75 (d, J = 8.4 Hz, 1H), 7.63-7.61 (m, 2H), 7.31-7.17 (m, 5H), 6.75 (d, J = 15.9 Hz, 1H), 5.45-5.39 (m, 1H),
 5 3.03-2.79 (m, 4H), 2.47-2.41 (m, 1H), 1.90-1.78 (m, 1H), 1.34-1.29 (m, 3H).

Example 33: (E)-3-(3-cyclopropyl-1H-indazol-6-yl)-N-(2,3-dihydro-1H-inden-1-yl)acrylamide




10 Into a 8-mL vial, was placed N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Example 24, Step 1**, 100.0 mg, 0.53 mmol, 1.0 equiv), 6-bromo-3-cyclopropyl-1H-indazole (126.6 mg, 0.53 mmol, 1.0 equiv), Pd(dppf)Cl₂ (39.1 mg, 0.05 mmol, 0.10 equiv), Et₃N (0.22 mL, 1.60 mmol, 3.0 equiv), DMF (4 mL). The resulting solution was stirred for 2 h at 120°C in an oil bath. The reaction mixture was cooled. The mixture was applied onto a silica gel column with PE/THF (1/1). This
 15 resulted in 15 mg (8%) of (E)-3-(3-cyclopropyl-1H-indazol-6-yl)-N-(2,3-dihydro-1H-inden-1-yl)acrylamide as a solid. LC MS (ES, m/z):

$[M+H]^+=344$

^1H NMR (300 MHz, DMSO- d_6 , ppm): δ 12.70 (s, 1H), 8.48 (d, J = 8.4 Hz, 1H), 7.78 (d, J = 8.4 Hz, 1H), 7.63 (m, 2H), 7.37-7.16 (m, 5H), 6.74 (d, J = 15.6 Hz, 1H), 5.49-5.37 (m, 1H), 3.10 -
 20 2.75 (m, 2H), 2.48-2.39 (m, 1H), 2.33-2.20 (m, 1H), 1.91-1.77 (m, 1H), 1.06-0.89 (m, 4H).

Example 34: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(4-fluoro-3-methyl-1H-indazol-6-yl)acrylamide




25 Into a 8-mL vial, was placed 6-bromo-4-fluoro-3-methyl-1H-indazole (**Example 24, Step 1**, 100.0 mg, 0.44 mmol, 1.0 equiv), N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (81.8 mg, 0.44 mmol, 1.0 equiv), Pd(dppf)Cl₂ (31.9 mg, 0.04 mmol, 0.10 equiv), Et₃N (0.18 mL, 1.31 mmol, 3.0 equiv), DMF (4.0 mL). The resulting solution was stirred for 2 h at 120 °C in an oil bath. The reaction mixture was cooled. The residue was applied onto a silica gel column with PE/THF
 30 (1/1). This resulted in 17 mg (12%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(4-fluoro-3-methyl-

1H-indazol-6-yl)acrylamide as a solid. LC MS (ES, m/z):


[M+H]⁺=336

¹H-NMR (300 MHz, DMSO-*d*₆, ppm) δ 13.09 (s, 1H), 8.47 (d, J = 8.1 Hz, 1H), 7.61 (d, J = 15.6 Hz, 1H), 7.48 (s, 1H), 7.35-7.15 (m, 4H), 7.01 (d, J = 12.0 Hz, 1H), 6.73 (d, J = 15.6 Hz, 1H),
 5.49-5.36 (m, 1H), 3.06-2.76 (m, 2H), 2.56 (s, 3H), 2.47-2.38 (m, 1H), 1.92-1.76 (m, 1H).


Example 35: (E)-N-(3,5-difluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide



Step 1: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3,5-difluoroaniline (1.0 g, 7.8 mmol, 1.0 equiv), CH₃CN (30 mL), NCS (1.1g, 8.16 mmol, 1.05 equiv). The resulting solution was stirred for 5 h at 80 °C. The reaction mixture was cooled to room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:4-1:1). This resulted in 500 mg (39%) of 4-chloro-3,5-difluoroaniline as a grey solid.




Step 2: Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 4-bromo-3,5-difluoroaniline (500.0 mg, 2.40 mmol, 1.0 equiv), CH₃CN (20.0 mL), NBS (1.3 g, 7.21 mmol, 3.0 equiv). The resulting solution was stirred for 1 h at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 700 mg (91%) of 2,6-dibromo-4-chloro-3,5-difluoroaniline as a yellow solid.




Step 3: Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 2,6-dibromo-4-chloro-3,5-difluoroaniline (3.0 g, 9.36 mmol, 1.0 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (0.76 g, 0.94 mmol, 0.10 equiv), dioxane (60.0 mL), Cs₂CO₃ (12.20 g, 37.46 mmol, 4.0 equiv), trimethyl-1,3,5,2,4,6-trioxatriborinane (8.23 g, 32.77 mmol, 3.50 equiv, 50%). The resulting solution was stirred overnight at 100 °C. The reaction mixture was cooled to room


temperature. The resulting mixture was concentrated. The crude mixture was purified by Prep-HPLC. This resulted in 600 mg (33%) of 4-chloro-3,5-difluoro-2,6-dimethylaniline as an off-white solid.



- 5 **Step 4:** Into a 100-mL pressure tank reactor, was placed 4-chloro-3,5-difluoro-2,6-dimethylaniline (280.0 mg, 1.46 mmol, 1.0 equiv), EtOH (20.0 mL), HCl (1.0 mL), Pd/C (77.76 mg). The flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The resulting solution was stirred overnight at 70 °C under an atmosphere of hydrogen (30atm). The reaction mixture was cooled to room temperature. The solids were
- 10 filtered out and the resulting mixture was concentrated. This resulted in 220 mg (crude) of 3,5-difluoro-2,6-dimethylaniline hydrochloride as an off-white solid.



- Step 5:** Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3,5-difluoro-2,6-dimethylaniline hydrochloride (220.0 mg, 1.14 mmol, 1.0 equiv), DCM (20.0 mL), Et₃N (0.48 mL, 3.41 mmol, 3.0 equiv). This was followed
- 15 by the addition of acryloyl chloride (123.4 mg, 1.36 mmol, 1.20 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred overnight at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 160 mg (67%) of N-(3,5-difluoro-2,6-dimethylphenyl)prop-2-enamide as a off-white
- 20 solid.



- Step 6:** Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3,5-difluoro-2,6-dimethylphenyl)prop-2-enamide (160.0 mg, 0.76 mmol, 1.0 equiv), 6-bromo-1,3-dihydroindol-2-one (192.8 mg, 0.91 mmol, 1.20 equiv), Pd(dppf)Cl₂.CH₂Cl₂
- 25 (61.7 mg, 0.076 mmol, 0.10 equiv), DMF (5.0 mL), Et₃N (0.32 mL, 2.27 mmol, 3.0 equiv). The resulting solution was stirred for 2 h at 100 °C The reaction mixture was cooled to room


temperature. The crude mixture was purified by Prep-HPLC. This resulted in 14.8 mg (6%) of (E)-N-(3,5-difluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide as a solid.

LC MS (ES, m/z):


[M+H]⁺=343

- 5 ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 10.56 (s, 1H), 9.77 (s, 1H), 7.56 (d, *J* = 15.6 Hz, 1H), 7.29-7.06 (m, 4H), 6.85 (d, *J* = 15.9 Hz, 1H), 3.53 (s, 2H), 2.05 (s, 6H).


Example 36: (E)-N-(3,4-difluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide



- 10 **Step 1:** Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed benzenamine, 3,4-difluoro- (1.0 g, 7.75 mmol, 1.0 equiv), CH₃CN (30.0 mL), NBS (2.9 g, 16.27 mmol, 2.10 equiv). The resulting solution was stirred overnight at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 1.4 g (63%) of 2,6-
15 dibromo-3,4-difluoroaniline as a dark brown solid.




- Step 2:** Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 2,6-dibromo-3,4-difluoroaniline (1.20 g, 4.18 mmol, 1.0 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (340.7 mg, 0.42 mmol, 0.10 equiv), Cs₂CO₃ (4.8 g, 14.64 mmol, 3.50 equiv),
20 dioxane (50.0 mL), trimethyl-1,3,5,2,4,6-trioxatriborinane (3.2 g, 12.55 mmol, 3.0 equiv, 50%). The resulting solution was stirred overnight at 100 °C. The reaction mixture was cooled to room temperature. The resulting mixture was concentrated. The crude mixture was purified by Prep-HPLC. This resulted in 350 mg (53%) of 3,4-difluoro-2,6-dimethylaniline as a off-white solid.



- 25 **Step 3:** Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3,4-difluoro-2,6-dimethylaniline (210.0 mg, 1.34 mmol, 1.0 equiv), DCM (20 mL), Et₃N (0.28 mL, 2.0 mmol, 1.50 equiv). This was followed by the addition

of acryloyl chloride (145.13 mg, 1.60 mmol, 1.20 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 2 hr at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 180 mg (64%) of *N*-(3,4-difluoro-2,6-dimethylphenyl)prop-2-enamide as a off-white solid.




Step 4: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed *N*-(3,4-difluoro-2,6-dimethylphenyl)prop-2-enamide (100.0 mg, 0.47 mmol, 1.0 equiv), 6-bromo-1,3-dihydroindol-2-one (120.47 mg, 0.57 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (38.57 mg, 0.05 mmol, 0.10 equiv), DMF (4.0 mL), Et₃N (0.2 mL, 1.42 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 11.3 mg (7%) of (E)-*N*-(3,4-difluoro-2,6-dimethylphenyl)-3-(2-oxoindolin-6-yl)acrylamide as an off-white solid.

LCMS (ES, *m/z*):

[*M*+*H*]⁺=343

¹H NMR (300 MHz, DMSO-*d*₆, *ppm*): δ 10.55 (s, 1H), 9.63 (s, 1H), 7.53 (d, *J* = 15.9 Hz, 1H), 7.29-7.18 (m, 3H), 7.06 (s, 1H), 6.82 (d, *J* = 15.9 Hz, 1H), 3.53 (s, 2H), 2.05 (s, 3H), 2.10 (s, 3H).

Example 37: (E)-*N*-(3-fluoro-2-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed *N*-(3-fluoro-2-methylphenyl)acrylamide (**Intermediate 1**, 110.0 mg, 0.61 mmol, 1.0 equiv), 6-bromo-3-methyl-1H-indazole (155.47 mg, 0.74 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (100.01 mg, 0.12 mmol, 0.20 equiv), DMF (4.0 mL), Et₃N (0.26 mL, 1.84 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 26 mg (14%) of (E)-*N*-(3-fluoro-2-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as an off-white solid.

LC MS (ES, m/z):

$[M+H]^+=310$

^1H NMR 300 MHz, DMSO- d_6 , ppm): δ 12.83 (brs, 1H), 9.62 (brs, 1H), 7.78-7.69 (m, 3H), 7.49 (d, $J = 7.8$ Hz, 1H), 7.39 (d, $J = 8.4$ Hz, 1H), 7.27-7.20 (m, 1H), 7.09-6.98 (m, 2H), 2.51 (s, 3H),
5 2.17 (s, 3H).

Example 38: (E)-3-(3-methyl-1H-indazol-6-yl)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)acrylamide




- 10 Into a 8-mL vial, was placed 6-bromo-3-methyl-1H-indazole (40.0 mg, 0.19 mmol, 1.0 equiv), N-(2-methyl-2,3-dihydro-1H-inden-1-yl)prop-2-enamide (Prepared according to Example 24, Step 1 using 2-methylindanamine, 38.14 mg, 0.19 mmol, 1.0 equiv), Pd(dppf)Cl₂ (13.87 mg, 0.02 mmol, 0.10 equiv) and Et₃N (0.8 mL, 0.57 mmol, 3.0 equiv) in DMF (2.0 mL). The resulting solution was stirred for 2 hr at 120 °C in an oil bath. The reaction solution was cooled to RT. The
15 crude mixture was purified by Flash-Prep-HPLC . This resulted in 29 mg (44%) of (E)-3-(3-methyl-1H-indazol-6-yl)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)acrylamide as a off- white solid.

LC MS (ES, m/z):

$[M+H]^+=332$


- ^1H -NMR (300 MHz, DMSO- d_6 , ppm) δ 12.78 (brs, 1H), 8.45-8.21 (m , 1H), 7.82-7.57 (m, 3H),
20 7.42-7.11 (m, 5H), 6.88-6.74 (m, 1H), 5.40-5.0 (m, 1H), 3.12-2.95 (m, 1H), 2.75-2.58 (m, 2H), 2.49 (s, 3H), 2.38-2.24 (m, 1H), 1.24-0.88 (m, 3H).

Example 39: (E)-3-(3-methyl-1H-indazol-6-yl)-N-(1-methyl-1H-indazol-7-yl)acrylamide



- 25 **Step 1:** Into a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 1-methylindazol-7-amine (400.0 mg, 2.71 mmol, 1.0 eq), DCM (10.0 mL), Et₃N (0.6 mL, 4.07 mmol, 1.50 eq). This was followed by the addition of acryloyl chloride (246 mg, 2.71 mmol, 1.0 eq) at -30 °C. The resulting solution was stirred for 10 min at -30 °C. The reaction was then quenched by the addition of 8 mL of water. The organic

layer was separated and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE(15:85). This resulted in 120 mg (21%) of *N*-(1-methylindazol-7-yl)prop-2-enamide as an off-white solid. LC MS (ES, *m/z*): $[M+H]^+=202$




5 **Step 2:** Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed *N*-(1-methylindazol-7-yl)prop-2-enamide (100.0 mg, 0.49 mmol, 1.0 eq), 6-bromo-3-methyl-1H-indazole (104.8 mg, 0.49 mmol, 1.0 eq), DMF (4.0 mL), Et₃N (0.14 mL, 0.99 mmol, 2.0 eq), Pd(dppf)Cl₂·CH₂Cl₂ (41 mg, 0.05 mmol, 0.10 eq). The resulting solution was stirred for 2 h at 120 °C. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 33 mg (20%)
10 of (E)-3-(3-methyl-1H-indazol-6-yl)-*N*-(1-methyl-1H-indazol-7-yl)acrylamide as an off-white solid. LC MS (ES, *m/z*):


$[M+H]^+=332$

¹H NMR (300 MHz, DMSO-*d*₆, ppm): 12.83 (brs, 1H), 10.14 (brs, 1H), 8.07 (s, 1H), 7.81-7.67 (m, 4H), 7.44-7.41 (m, 1H), 7.25-7.11 (m, 2H), 7.01 (d, *J* = 15.9 Hz, 1H), 4.12 (s, 3H), 2.50 (s,
15 3H).


Example 40: (E)-*N*-(5-fluoro-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide




20 **Step 1:** Into a 40-mL sealed tube, was placed 5-fluoro-2,3-dihydroinden-1-one (800.0 mg, 5.33 mmol, 1.0 eq), NaOAc (874.14 mg, 10.66mmol, 2.0 eq), MeOH (15.0 mL), Hydroxylamine hydrochloride (1.10 g, 15.98 mmol, 3.0 eq). The resulting solution was stirred for 16 h at 60 °C. The resulting mixture was concentrated. The crude product was diluted with of EtOAc (30.0 mL)
25 and H₂O (15.0 mL). The organic phase was washed with 20 ml of H₂O. The organic layer was concentrated. This resulted in 810 mg (92%) of *N*-[5-fluoro-2,3-dihydroinden- 1-ylidene]hydroxylamine as an off-white solid. LC MS (ES, *m/z*): $[M+H]^+=166$



Step 2: Into a 50-mL round-bottom flask, was placed N-[5-fluoro-2,3-dihydroinden-1-ylidene]hydroxylamine (810.0 mg, 4.90 mmol, 1.0 eq), MeOH (20.0 mL). This was followed by the addition of Pd/C (104.38 mg) under H₂. The resulting solution was stirred for 16 h at room temperature. The solid was filtered out and washed with 10 mL MeOH. The combined solution was concentrated. This resulted in 530 mg (71%) of 5-fluoro-2,3-dihydro-1H-inden-1-amine as an off-white solid. LC MS (ES, m/z): [M+H]⁺=152



Step 3: Into a 25-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 5-fluoro-2,3-dihydro-1H-inden-1-amine (210.0 mg, 1.38 mmol, 1.0 eq), DCM (5.0 mL), Et₃N (0.4 mL, 2.77 mmol, 2.0 eq). This was followed by the addition of acryloyl chloride (188.58 mg, 2.08 mmol, 1.50 eq) at -30°C. The resulting solution was stirred for 10 min at -30 °C. The reaction was then quenched by the addition of 5 mL of water. The organic layer was separated and concentrated. The residue was applied onto a silica gel column with EtOAc/PE (80:20). This resulted in 200 mg (70%) of N-(5-fluoro-2,3-dihydro-1H-inden-1-yl)prop-2-enamide as an off-white solid. LC MS (ES, m/z): [M+H]⁺=206




Step 4: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(5-fluoro-2,3-dihydro-1H-inden-1-yl)acrylamide (100.0 mg, 0.48 mmol, 1.0 eq), 6-bromo-3-methyl-1H-indazole (102.8 mg, 0.48 mmol, 1.0 eq), Et₃N (0.14 mL, 0.97 mmol, 2.0 eq), DMF (5.0 mL), Pd(dppf)Cl₂·CH₂Cl₂ (39.69 mg, 0.05 mmol, 0.10 eq). The resulting solution was stirred for 2 h at 120 °C. The reaction solution was cooled to RT. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 34 mg (20%) of (E)-N-(5-fluoro-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a white solid. LC MS (ES, m/z):


[M+H]⁺=336

¹H NMR (300 MHz, DMSO-*d*₆, ppm): 12.78 (s, 1H), 8.48 (d, *J* = 8.1 Hz, 1H), 7.74-7.61 (m, 3H), 7.32-7.24 (m, 2H), 7.13-7.09 (m, 2H), 6.73 (d, *J* = 15.9 Hz, 1H), 5.40-5.37 (m, 1H), 3.16-2.73 (m, 2H), 2.52 (s, 3H), 2.50-1.83 (m, 2H).

5 **Example 41: (E)-N-(4-fluoro-3-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide**



Step 1: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 4-fluoro-3-methylaniline (600.0 mg, 4.79 mmol, 1.0 equiv), DCM (20.0 mL), Et₃N (2.0 mL, 14.38 mmol, 3.0 equiv). This was followed by the addition of acryloyl chloride (520.73 mg, 5.75 mmol, 1.20 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 1 h at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 400 mg (56%) of N-(4-fluoro-3-methylphenyl)prop-2-enamide as a light yellow solid.




Step 2: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(4-fluoro-3-methylphenyl)prop-2-enamide (110.0 mg, 0.61 mmol, 1.0 equiv), 6-bromo-3-methyl-1H-indazole (155.47 mg, 0.74 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (100.01 mg, 0.12 mmol, 0.20 equiv), DMF (4.0 mL), Et₃N (0.26 mL, 1.84 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120 °C. The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 20.4 mg (11%) of (E)-N-(4-fluoro-3-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a off-white solid.

LC MS (ES, m/z):


[M+H]⁺=310

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.82 (brs, 1H), 10.16 (brs, 1H), 7.77-7.67 (m, 3H), 7.62-7.59 (m, 1H), 7.56-7.51 (m, 1H), 7.36 (d, *J* = 8.4 Hz, 1H), 7.14-7.08 (m, 1H), 6.88 (d, *J* = 15.6 Hz, 1H), 2.51 (s, 3H), 2.24 (s, 3H).

Example 42: (E)-N-(3-fluoro-4-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide



Step 1: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3-fluoro-4-methylaniline (550.0 mg, 4.40 mmol, 1.0 equiv), DCM (20.0 mL), Et_3N (1.2 mL, 8.79 mmol, 2.0 equiv). This was followed by the addition of acryloyl chloride (477.34 mg, 5.27 mmol, 1.20 equiv) dropwise with stirring at 0°C . The resulting solution was stirred for 1 h at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 450 mg (57%) of N-(3-fluoro-4-methylphenyl)prop-2-enamide as a off-white solid.




Step 2: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-4-methylphenyl)prop-2-enamide (110.0 mg, 0.61 mmol, 1.0 equiv), 6-bromo-3-methyl-1H-indazole (155.5 mg, 0.74 mmol, 1.20 equiv), $\text{Pd(dppf)Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (100.0 mg, 0.12 mmol, 0.20 equiv), DMF (4.0 mL), Et_3N (0.26 mL, 1.84 mmol, 3.0 equiv). The resulting solution was stirred overnight at 120°C . The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 35.8 mg (19%) of (E)-N-(3-fluoro-4-methylphenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a off-white solid.

LC MS (ES, m/z):

$[\text{M}+\text{H}]^+=310$

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): δ 12.83 (brs, 1H), 10.30 (brs, 1H), 7.77-7.67 (m, 4H), 7.39-7.35 (m, 1H), 7.30-7.20 (m, 2H), 6.87 (d, $J = 15.6$ Hz, 1H), 2.51 (s, 3H), 2.20 (s, 3H).


Example 43: Racemic- (E)-3-(3-methyl-1H-indazol-6-yl)-N-((1R,2R)-2-methylcyclohexyl)acrylamide



Step 1: Into a 25-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed *Racemic-trans*-(1R,2R)-2-methylcyclohexan-1-amine hydrochloride (300.0 mg, 2.0 mmol, 1.0 eq), DCM (6.0 mL), Et_3N (0.84 mL, 6.01 mmol, 3.0 eq). This was followed by the addition of acryloyl chloride (181.43 mg, 2.0 mmol, 1.0 eq) at 0°C .

The resulting solution was stirred for 40 min at room temperature. The reaction was then quenched by the addition of 5 mL of water. The resulting solution was diluted with 10 mL of DCM. The resulting mixture was washed with 10 mL of brine. The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (22:78). This resulted in 200 mg (60%) of *N*-[(1*R*,2*R*)-2-methylcyclohexyl]prop-2-enamide as an off-white solid.

LC-MS (ES, *m/z*): $[M+H]^+ = 168$




Step 2: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed DMF (4.0 mL), *N*-[(1*R*,2*R*)-2-methylcyclohexyl]prop-2-enamide (100.0 mg, 0.60 mmol, 1.0 eq), 6-bromo-3-methyl-1*H*-indazole (126. mg, 0.60 mmol, 1.0 eq), Et₃N (0.25 mL, 1.79 mmol, 3.0 eq), Pd(dppf)Cl₂·CH₂Cl₂ (48.71 mg, 0.06 mmol, 0.10 eq). The resulting solution was stirred for 2 h at 120 °C. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 41 mg (23%) of Racemic (*E*)-3-(3-methyl-1*H*-indazol-6-yl)-*N*-((1*R*,2*R*)-2-methylcyclohexyl)acrylamide as an off-white solid. LC MS (ES, *m/z*):

$[M+H]^+ = 298$

¹H NMR (300 MHz, CD₃OD-*d*₄, ppm): 7.75-7.60 (m, 3H), 7.43-7.40 (m, 1H), 6.70 (d, *J* = 15.6 Hz, 1H), 3.54-3.50 (m, 1H), 2.57 (s, 3H), 1.96-1.71 (m, 4H), 1.48-1.37 (m, 5H), 1.30 (d, *J* = 9.3 Hz, 3H).

Example 44: (*E*)-3-(3-cyano-1*H*-indazol-6-yl)-*N*-(2-methyl-2,3-dihydro-1*H*-inden-1-yl)acrylamide




Into a 8-mL vial, was placed 6-bromo-1*H*-indazole-3-carbonitrile (40.0 mg, 0.18 mmol, 1.0 equiv), *N*-(2-methyl-2,3-dihydro-1*H*-inden-1-yl)prop-2-enamide (**Prepared according to Example 24, Step 1 using 2-methylindanamine**, 36.26 mg, 0.18 mmol, 1.0 equiv), Pd(dppf)Cl₂ (13.18 mg, 0.018 mmol, 0.10 equiv) and Et₃N (0.075 mL, 0.54 mmol, 3.0 equiv) in DMF (2.0 mL). The resulting solution was stirred for 2 h at 120 °C in an oil bath. The mixture was

cooled to RT. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 16 mg (26%) of (E)-3-(3-cyano-1H-indazol-6-yl)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)acrylamide as a off-white solid. LC MS (ES, m/z):

[M+H]⁺=343

- 5 ¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 8.59-8.30 (m, 1H), 8.05-7.87 (m, 2H), 7.77-7.55 (m, 2H), 7.34-7.08 (m, 4H), 6.97-6.79 (m, 1H), 5.50-4.98 (m, 1H), 3.06-2.95 (m, 1H), 2.84-2.62 (m, 1H), 2.39-2.19 (m, 1H), 1.25-0.88 (m, 3H).

10 **Example 45: (E)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide**




Into a 8-mL vial, was placed 5-bromo-1-methyl-3H-1,3-benzodiazol-2-one (50.0 mg, 0.22 mmol, 1.0 equiv), N-(2-methyl-2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Prepared according to Example 24, Step 1 using 2-methylindanamine** 44.3 mg, 0.22 mmol, 1.0 equiv), Pd(dppf)Cl₂ (16.1 mg, 0.022 mmol, 0.10 equiv) and Et₃N (0.092 mL, 0.66 mmol, 3.0 equiv) in DMF (2.0 mL). The resulting solution was stirred for 2 hat 120 °C in an oil bath. The reaction solution was cooled to RT. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 29 mg (38%) of (E)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)acrylamide as a off-white solid. LC MS (ES, m/z):

20 [M+H]⁺=348

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 10.99 (s, 1H), 8.32-8.12 (m, 1H), 7.54-7.46 (m, 1H), 7.28-7.10 (m, 7H), 6.69-6.57 (m, 1H), 5.40-5.02 (m, 1H), 3.30 (s, 3H), 3.07-3.0 (m, 1H), 2.69-2.50 (m, 1H), 2.27-2.20 (m, 1H), 1.21- 0.90 (m, 3H).

25 **Example 46: (E)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide**




Into a 8-mL vial, was placed 2-benzoxazolinone, 5-bromo- (50.0 mg, 0.23 mmol, 1.0 equiv), N-(2-methyl-2,3-dihydro-1H-inden-1-yl)prop-2-enamide (**Prepared according to Example 24,**

Step 1 using 2-methylindanamine 47.0 mg, 0.23 mmol, 1.0 equiv), Pd(dppf)Cl₂ (17.1 mg, 0.023 mmol, 0.10 equiv) and Et₃N (0.098 mL, 0.70 mmol, 3.0 equiv) in DMF (2.0 mL). The resulting solution was stirred for 2 hr at 120 °C in an oil bath. The reaction solution was cooled to RT. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 29 mg (37%) of (E)-N-(2-methyl-2,3-dihydro-1H-inden-1-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a off-white solid. LCMS (ES, m/z):


[M+H]⁺=335

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 11.76 (brs, 1H), 8.40-8.19 (m, 1H), 7.56-7.48 (m, 1H), 7.33-7.12 (m, 6H), 6.74-6.63 (m, 1H), 5.39-4.99 (m, 1H), 3.10-2.94 (m, 1H), 2.69-2.20 (m, 2H), 1.21-0.91(m, 3H).

Example 47: (Z)-2-fluoro-N-(3-fluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide



Step 1: Into a 40-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed DMF (20.0 mL), 6-bromo-1,3-dihydroindol-2-one (1.0 g, 4.72 mmol, 1.0 eq), methyl 2-fluoroacrylate (0.59 g, 5.66 mmol, 1.20 eq), Pd(dppf)Cl₂·CH₂Cl₂ (77.0 mg, 0.094 mmol, 0.02 eq), Et₃N (1.3 mL, 9.43 mmol, 2.0 eq). The resulting solution was stirred for 2 h at 110 °C. The resulting mixture was concentrated. The residue was applied onto a silica gel column with (25/75). This resulted in 420 mg (37%) of methyl 2-fluoro-3-(2-oxo-1,3-dihydroindol-6-yl)prop-2-enoate as a yellow solid. LC-MS (ES, m/z): [M-H]⁺=234




Step 2: Into a 8-mL sealed tube, was placed methyl 2-fluoro-3-(2-oxo-1,3-dihydroindol-6-yl)prop-2-enoate (150.0 mg, 0.64 mmol, 1.0 eq), 3-fluoro-2-methyl-aniline (239.42 mg, 1.91 mmol, 3.0 eq), THF (3.0 mL). This was followed by the addition of LiHMDS (1.92 mL, 1.92 mmol, 3.0 eq) at 0 °C. The resulting solution was stirred for 30 min at room temperature. The reaction was then quenched by the addition of 15 mL of H₂O. The resulting solution was extracted with 2x10 mL of EtOAc. The combined organic phase was dried over anhydrous sodium sulfate and concentrated. The crude product was purified by Flash-Prep-HPLC. This resulted in 28 mg (13%) of (Z)-2-fluoro-N-(3-fluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide as a white solid. LC MS (ES, m/z):

[M+H]⁺=329


^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): δ 10.50 (s, 1H), 10.16 (s, 1H), 7.31-6.93 (m, 7H), 3.53 (s, 2H), 2.12 (s, 3H).

Example 48: (E)-N-(3-chloro-2-methylphenyl)-N-methyl-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed (E)-methyl 3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate (**Prepared according to Example 11, Step 2**, 30.0 mg, 0.14 mmol, 1.0 equiv) in THF (2.0 mL). Then 3-chloro-N,2-dimethylbenzenamine (27.60 mg, 0.18 mmol, 1.30 equiv) was added. The reaction mixture was cooled to 0°C. Then 1M LiHMDS (0.68 mL, 0.68 mmol, 5.0 equiv) was added. The reaction mixture was stirred 1 h at RT. The reaction mixture was quenched by 1 mL water and extracted with 2x5 mL EtOAc, the organic was concentrated under vacuum to afford crude product. The crude product was then purified by Prep-HPLC. This resulted in 9.6 mg (21%) of (E)-N-(3-chloro-2-methylphenyl)-N-methyl-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a white solid. LC-MS (ES, m/z): $[\text{M}+\text{H}]^+=343$

Example 49: (E)-N-(2-methylcyclopentyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed (E)-methyl 3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate (**Prepared according to Example 11, Step 2**, 30.0 mg, 0.14 mmol, 1.0 equiv) in THF (2.0 mL). Then 2-methylcyclopentanamine (17.6 mg, 0.18 mmol, 1.30 equiv) was added. The reaction mixture was cooled to 0 °C. Then 1M LiHMDS (0.68 mL, 0.68 mmol, 5.0 equiv) was added. The reaction mixture was stirred 1 h at RT. The reaction mixture was quenched by 1 mL water and extracted with 2x5 mL EtOAc, the organic was concentrated under vacuum to afford crude product. The crude product was then purified by Prep-HPLC. This resulted in 19 mg (48%) of (E)-N-(2-methylcyclopentyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a white solid. LC-MS (ES, m/z): $[\text{M}+\text{H}]^+=287$


Example 50: (E)-N-(3-fluoro-2-(methoxymethyl)phenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide



- 5 **Step 1:** Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed THF (50.0 mL), 2-fluoro-6-nitrobenzoic acid (2.0 g, 10.80 mmol, 1.0 eq). This was followed by the addition of borane (32.41 mL, 32.41 mmol, 3.0 eq, 1 M/THF) at 0°C. The resulting solution was stirred for 10 h at 50 °C. The reaction was then quenched by the addition of 20 mL of 3 M HCl. The resulting solution was diluted with 50 mL of EtOAc. The organic layer was washed with 2x30 ml of brine. The solution was dried over sodium sulfate and evaporated under reduced pressure. This resulted in 1.3 g (70%) of (2-fluoro-6-nitrophenyl)methanol as a light brown solid.




- 15 **Step 2:** Into a 40-mL sealed tube, was placed (2-fluoro-6-nitrophenyl)methanol (1.30 g, 7.59 mmol, 1.0 eq), DMF (20.0 mL), methyl iodide (5.39 g, 37.98 mmol, 5.0 eq). This was followed by the addition of Cs₂CO₃ (3.71 g, 11.39 mmol, 1.50 eq) at 0 °C. The resulting solution was stirred for 5 h at 50 °C. The solids were filtered out. The filtrate solution was diluted with 100 mL of EtOAc. The resulting mixture was washed with 3x50 ml of H₂O and 50 mL of brine. The organic layer was dried over sodium sulfate and evaporated. The crude product was purified by Flash-Prep-HPLC. This resulted in 0.91 g (64%) of 1-fluoro-2-(methoxymethyl)-3-nitrobenzene as yellow oil.




- 25 **Step 3:** Into a 40-mL sealed tube, was placed 1-fluoro-2-(methoxymethyl)-3-nitrobenzene (0.70 g, 3.78 mmol, 1.0 eq), MeOH (20.0 mL), H₂O (3.0 mL), NH₄Cl (1.21 g, 22.68 mmol, 6.0 eq). This was followed by the addition of Zn (1.24 g, 18.90 mmol, 5.0 eq) at 10 °C. The resulting solution was stirred for 1 h at 25 °C. The solids were filtered out. The resulting solution was diluted with

20 mL of DCM. The mixture was dried over anhydrous sodium sulfate and concentrated (low temp). This resulted in 0.21 g (35%) of 3-fluoro-2-(methoxymethyl)aniline as light brown oil (the amine was unstable and was used in next step directly and quickly). LC-MS (ES, m/z):

$[M+H]^+=156$



Step 4: Into a 8-mL sealed tube, was placed 3-fluoro-2-(methoxymethyl)aniline (200.0 mg, 1.28 mmol, 1.0 eq), DCM (4.0 mL), Et_3N (0.36 mL, 2.57 mmol, 2.0 eq). This was followed by the addition of acryloyl chloride (116.6 mg, 1.28 mmol, 1.0 eq) at 0°C . The resulting solution was stirred for 0.5 h at 0°C . The reaction was then quenched by the addition of 2 mL of water. The resulting mixture was washed with 2 mL of brine. The organic layer was dried over sodium sulfate and concentrated. This resulted in 160 mg (59%) of N-[3-fluoro-2-(methoxymethyl)phenyl]prop-2-enamide as an off-white solid. LC-MS (ES, m/z): $[M+H]^+=210$




Step 5: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-[3-fluoro-2-(methoxymethyl)phenyl]prop-2-enamide (100.0 mg, 0.47 mmol, 1.0 eq), DMF (4.0 mL), 6-bromo-3-methyl-1H-indazole (100.88 mg, 0.47 mmol, 1.0 eq), Et_3N (0.2 mL, 1.43 mmol, 3.0 eq), $\text{Pd(dppf)Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (38.94 mg, 0.048 mmol, 0.10 eq). The resulting solution was stirred for 2 h at 120°C . The crude mixture was purified by Flash-Prep-HPLC. This resulted in 71 mg (43%) of (E)-N-(3-fluoro-2-(methoxymethyl)phenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a white solid. LC-MS (ES, m/z):


$[M+H]^+=340$

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): δ 12.84 (s, 1H), 9.55 (s, 1H), 7.78-7.69 (m, 4H), 7.44-7.38 (m, 2H), 7.11-7.02 (m, 2H), 4.56 (s, 2H), 3.32 (s, 3H), 2.51 (s, 3H).


Example 51: (E)-3-(3-cyano-1H-indazol-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide



Step 1: Into a 40-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed 6-bromo-1H-indazole-3-carbonitrile (220.0 mg, 0.99 mmol, 1.0 equiv), dihydropyran (416.7 mg, 4.95 mmol, 5.0 equiv), DCM (10.0 mL), TsOH (34.1 mg, 0.20 mmol, 0.20 equiv). The resulting solution was stirred overnight at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:20-1:8). This resulted in 300 mg (99%) of 6-bromo-1-(oxan-2-yl)indazole-3-carbonitrile as a off-white solid.




Step 2: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-fluoro-2-methylphenyl)acrylamide (150.0 mg, 0.84 mmol, 1.0 equiv), 6-bromo-1-(oxan-2-yl)indazole-3-carbonitrile (307.5 mg, 1.0 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (68.2 mg, 0.08 mmol, 0.10 equiv), DMF (4.0 mL), Et₃N (0.35 mL, 2.51 mmol, 3.0 equiv). The resulting solution was stirred for 3 h at 120 °C. The reaction mixture was cooled to room temperature. The mixture was applied onto a silica gel column with THF:PE (1:4-1:1). This resulted in 150 mg (44%) of (2E)-3-[3-cyano-1-(oxan-2-yl)indazol-6-yl]-N-(3-fluoro-2-methylphenyl)acrylamide as a light brown solid.



Step 3: Into a 40-mL sealed tube, was placed (2E)-3-[3-cyano-1-(oxan-2-yl)indazol-6-yl]-N-(3-fluoro-2-methylphenyl)acrylamide (150.0 mg, 0.37 mmol, 1.0 equiv), 4M HCl/dioxane (10 mL). The resulting solution was stirred overnight at room temperature. The resulting mixture was concentrated. The crude product was purified by Prep-HPLC. This resulted in 12.5 mg (11%) of (E)-3-(3-cyano-1H-indazol-6-yl)-N-(3-fluoro-2-methylphenyl)acrylamide as a off-white solid. LC-MS (ES, m/z):


[M+H]⁺=321

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 14.55 (brs, 1H), 9.68 (s, 1H), 8.0-7.95 (m, 2H), 7.80 (d, *J* = 15.9 Hz, 1H), 7.68 (d, *J* = 9.0 Hz, 1H), 7.49 (d, *J* = 9.6 Hz, 1H), 7.28-7.20 (m, 1H), 7.15 (d, *J* = 15.9 Hz, 1H), 7.05-6.99 (m, 1H), 2.17 (s, 3H).

Example 52: (E)-3-(3-methyl-1H-indazol-6-yl)-N-(3-methylchroman-4-yl)acrylamide

Step 1: Into a 25-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3-methyl-3,4-dihydro-2H-1-benzopyran-4-amine (110.0 mg, 0.67 mmol, 1.0 equiv), DCM (10.0 mL), Et₃N (0.19 mL, 1.35 mmol, 2.0 equiv). This was followed by the addition of acryloyl chloride (67.10 mg, 0.74 mmol, 1.10 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred for 2 h at room temperature. The reaction was then quenched by the addition of 10 mL of water. The resulting solution was extracted with 2x15 mL of dichloromethane dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE (1/5). This resulted in 110 mg (75%) of N-(3-methyl-3,4-dihydro-2H-1-benzopyran-4-yl)prop-2-enamide as a light yellow solid.

LC-MS (ES, m/z): [M+H]⁺=218




Step 2: Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-methyl-3,4-dihydro-2H-1-benzopyran-4-yl)prop-2-enamide (120.0 mg, 0.55 mmol, 1.0 equiv), DMF (5.0 mL), 6-bromo-3-methyl-1H-indazole (116.6 mg, 0.55 mmol, 1.0 equiv), Et₃N (0.23 mL, 1.66 mmol, 3.0 equiv), Pd(dppf)Cl₂.CH₂Cl₂ (22.50 mg, 0.028 mmol, 0.05 equiv). The resulting solution was stirred for 5 h at 120 °C. The reaction mixture was cooled to room temperature. The residue was applied onto a silica gel column with EtOAc (1/3). This resulted in 55 mg (29%) of (E)-3-(3-methyl-1H-indazol-6-yl)-N-(3-methylchroman-4-yl)acrylamide as a off-white solid. LC-MS (ES, m/z):


[M+H]⁺=348

¹H NMR (300 MHz, DMSO-*d*₆, ppm): 12.80 (brs, 1H), 8.52-8.33 (m, 1H), 7.74-7.62 (m, 3H),

7.33-7.28 (m, 1H), 7.18-7.13 (m, 2H), 6.92-6.73 (m, 3H), 5.24-4.81 (m, 1H), 4.24-4.10 (m, 1H), 3.98-3.88 (m, 1H), 2.50-2.49 (m, 3H), 2.28-2.04 (m, 1H), 0.98-0.90 (m, 3H).

Example 53: (E)-N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide


Step 1: Into a 25-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 2-methyl-1,2,3,4-tetrahydronaphthalen-1-amine (160.0 mg, 0.99 mmol, 1.0 equiv), DCM (10.0 mL), Et_3N (0.28 mL, 1.98 mmol, 2.0 equiv). This was followed by the addition of acryloyl chloride (98.79 mg, 1.09 mmol, 1.10 equiv) dropwise with stirring at 0°C . The resulting solution was stirred for 2 h at room temperature. The reaction was then quenched by the addition of 5 mL of water. The resulting solution was extracted with 2x10 mL of DCM and the organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with EtOAc/PE (1/5). This resulted in 140 mg (66%) of N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)prop-2-enamide as a light yellow solid. LC-MS (ES, m/z): $[\text{M}+\text{H}]^+=216$




Step 2: Into a 8-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)prop-2-enamide (140.0 mg, 0.65 mmol, 1.0 equiv), DMF (5.0 mL), 6-bromo-3-methyl-1H-indazole (137.3 mg, 0.65 mmol, 1.0 equiv), Et_3N (0.27 mL, 1.95 mmol, 3.0 equiv), $\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (26.5 mg, 0.033 mmol, 0.05 equiv). The resulting solution was stirred for 5 h at 120°C . The reaction mixture was cooled to room temperature. The crude mixture was purified by Prep-HPLC. This resulted in 30 mg (13%) of (E)-N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a off-white solid. LC-MS (ES, m/z):

$[\text{M}+\text{H}]^+=346$

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): 12.77 (brs, 1H), 8.39-8.15 (m, 1H), 7.74-7.59 (m, 3H), 7.34-7.11 (m, 5H), 6.84-6.76 (m, 1H), 5.23-4.76 (m, 1H), 2.84-2.77 (m, 2H), 2.51-2.48 (m, 3H), 2.08-1.98 (m, 1H), 1.97-1.52 (m, 2H), 1.10-0.90 (m, 3H).

Example 54: (E)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide


Step 1: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-amine hydrochloride (300.0 mg, 1.50 mmol, 1.0 equiv), DCM (20.0 mL), Et_3N (0.31 mL, 2.25 mmol, 1.50 equiv). This was followed by the addition of acryloyl chloride (163.2 mg, 1.80 mmol, 1.20 equiv) dropwise with stirring at 0°C . The resulting solution was stirred for 2 h at room temperature. The resulting mixture was concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 200 mg (61%) of N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide as a off-white solid.



Step 2: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (140.0 mg, 0.64 mmol, 1.0 equiv), 6-bromo-3-methyl-1H-indazole (163.2 mg, 0.77 mmol, 1.20 equiv), $\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (52.5 mg, 0.064 mmol, 0.10 equiv), DMF (4 mL), Et_3N (0.27 mL, 1.93 mmol, 3.0 equiv). The resulting solution was stirred for 1 h at 120°C . The reaction mixture was cooled to room temperature. The crude mixture was purified by Flash-Prep-HPLC. This resulted in 28.6 mg (13%) of (E)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as an off-white solid. LC-MS (ES, m/z):


$[\text{M}+\text{H}]^+=348$

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): δ 12.78 (brs, 1H), 8.58 (d, $J = 8.7$ Hz, 1H), 7.74-7.62 (m, 3H), 7.33-7.18 (m, 5H), 6.75 (d, $J = 15.6$ Hz, 1H), 5.34-5.30 (m, 1H), 4.07-4.01 (m, 1H), 3.38 (s, 3H), 3.35-3.27 (m, 1H), 2.84-2.76 (m, 1H), 2.49 (s, 3H).

Example 55: (R,E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide

Step 1: Into a 8-mL sealed tube, was placed (1R)-2,3-dihydro-1H-inden-1-amine (100.0 mg, 0.75 mmol, 1.0 equiv), DCM (5.0 mL), Et_3N (0.21 mL, 1.50 mmol, 2.0 equiv). This was followed by the addition of acryloyl chloride (81.54 mg, 0.90 mmol, 1.20 equiv) at -30°C . The resulting solution was stirred for 10 min at -30°C . The reaction was then quenched by the addition of 5 mL of water. The organic phase was dried over Na_2SO_4 and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (15:75). This resulted in 93 mg (66%) of N-[(1R)-2,3-dihydro-1H-inden-1-yl]prop-2-enamide as a white solid.


LC-MS (ES, m/z): $[\text{M}+\text{H}]^+=188$



Step 2: Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed N-[(1R)-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (60.0 mg, 0.32 mmol, 1.0 equiv), 6-bromo-3-methyl-1H-indazole (67.63 mg, 0.32 mmol, 1.0 equiv), DMF (2.0 mL), Et_3N (0.09 mL, 0.64 mmol, 2.0 equiv), $\text{Pd(dppf)Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (5.2 mg, 0.06 mmol, 0.02 equiv). The resulting solution was stirred for 2 h at 120°C . The crude mixture was purified by Flash-Prep-HPLC. This resulted in 41 mg (40%) of (R,E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a white solid. LC-MS (ES, m/z):


$[\text{M}+\text{H}]^+=318$

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): 12.78 (s, 1H), 8.47 (d, $J = 8.4$ Hz, 1H), 7.73-7.62 (m, 3H), 7.32-7.17 (m, 5H), 6.74 (d, $J = 15.9$ Hz, 1H), 5.47-5.39 (m, 1H), 3.03-2.79 (m, 2H), 2.51-2.41 (m, 4H), 1.90-1.81 (m, 1H).

Example 56: (E)-N-(chroman-4-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide


Prepared according to Example 11, Step 4 using (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylic acid and chroman-4-amine to give (E)-N-(chroman-4-yl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a white solid. LC-MS (ES, m/z): [M+H]⁺=337

5 **Example 57: (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)acrylamide**



10 Prepared according to Example 11, Step 4 using (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylic acid and 1,2,3,4-tetrahydronaphthalen-1-amine to give (E)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)acrylamide as a white solid. LC-MS (ES, m/z): [M+H]⁺=335


Example 58: (E)-N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-(2-oxoindolin-6-yl)acrylamide



15 Into a 8-mL vial, was placed (E)-3-(2-oxoindolin-6-yl)acrylic acid (50.0 mg, 0.25 mmol, 1.00 equiv), 2-methyl-1,2,3,4-tetrahydronaphthalen-1-amine (40.3 mg, 0.25 mmol, 1.00 equiv), HATU (141.4 mg, 0.37 mmol, 1.50 equiv) and DIPEA (95.4 mg, 0.74 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 2 h at 20 °C. The mixture was purified by Flash-Prep-
20 HPLC. This resulted in 19 mg (22%) of (E)-N-(2-methyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-(2-oxoindolin-6-yl)acrylamide as a off-white solid. LC-MS (ES, m/z): [M+H]⁺=347


¹H NMR (300 MHz, DMSO-d₆, ppm) δ 10.51 (brs, 1H), 8.40-8.20 (m, 1H), 7.50-7.45 (m, 1H), 7.27-7.14 (m, 6H), 7.05-6.96 (m, 1H), 6.68-6.63 (m, 1H), 5.45-5.00 (m, 1H), 3.51 (s, 2H), 3.10-3.00 (m, 1H), 2.70-2.20 (m, 4H), 1.20-0.91 (m, 3H).

25


Example 59: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide

Into a 8-mL vial, was placed (E)-3-(2-oxoindolin-6-yl)acrylic acid (50.0 mg, 0.25 mmol, 1.00 equiv), 2,3-dihydro-1H-inden-1-amine (33.3 mg, 0.25 mmol, 1.00 equiv), HATU (141.4 mg, 0.37 mmol, 1.50 equiv) and DIPEA (95.4 mg, 0.74 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 2 h at 20 °C. The mixture was purified by Flash-Prep-HPLC. This resulted in 15 mg (19%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide as a off-white solid. LC-MS (ES, m/z): [M+H]⁺=319

¹H NMR (300 MHz, DMSO-d₆, ppm) δ 10.50 (brs, 1H), 8.46 (d, J=8.1 Hz, 1H), 7.45 (d, J=15.6 Hz, 1H), 7.29-7.12 (m, 6H), 6.98 (s, 1H), 6.62 (d, J=15.9 Hz, 1H), 5.45-5.37 (m, 1H), 3.51 (s, 2H), 3.02-2.79 (m, 2H), 2.50-2.40 (m, 1H), 1.89-1.82 (m, 1H).


Example 60: (E)-N-(3,5-difluoro-2-methylphenyl)-3-(2-oxoindolin-6-yl)acrylamide

Step 1: Into a 100-mL pressure tank reactor, was placed 4-chloro-3,5-difluoro-2-methylaniline (160.00 mg, 0.90 mmol, 1.00 equiv), EtOH (30.00 mL), 4M HCl/EtOH (1.00 mL), Pd/C (47.94 mg). The flask was evacuated and flushed three times with nitrogen, followed by flushing with hydrogen. The resulting solution was stirred overnight at 70 °C under an atmosphere of hydrogen (30 atm). The reaction mixture was cooled to room temperature. The solids were filtered out. The resulting mixture was concentrated. This resulted in 120 mg (crude) of 3,5-difluoro-2-methylaniline hydrochloride.



Step 2: Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 3,5-difluoro-2-methylaniline hydrochloride (120.00 mg, 0.67 mmol, 1.00 equiv), DCM (20 mL), Et₃N (0.28 mL, 2.01 mmol, 3.00 equiv). This was followed by the addition of acryloyl chloride (90.7 mg, 1.00 mmol, 1.50 equiv) dropwise with stirring at 0 °C. The resulting solution was stirred overnight at room temperature. The resulting mixture was

concentrated. The residue was applied onto a silica gel column with THF:PE (1:5-1:3). This resulted in 120 mg (91%) of *N*-(3,5-difluoro-2-methylphenyl)prop-2-enamide as a off-white solid.




- 5 **Step 3:** Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed *N*-(3,5-difluoro-2-methylphenyl)prop-2-enamide (120.00 mg, 0.61 mmol, 1.00 equiv), 6-bromo-1,3-dihydroindol-2-one (154.9 mg, 0.73 mmol, 1.20 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (49.6 mg, 0.06 mmol, 0.10 equiv), DMF (4.00 mL), Et₃N (0.25 mL, 1.83 mmol, 3.00 equiv). The resulting solution was stirred for 2 h at 100 °C. The reaction mixture was cooled to room
- 10 temperature. The crude mixture was purified by Prep-HPLC. This resulted in 11.3 mg (6%) of (E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(2-oxoindolin-6-yl)acrylamide as a off-white solid.

LC-MS (ES, *m/z*):

[*M*+*H*]⁺=329

- ¹H NMR (300 MHz, DMSO-*d*₆, *ppm*): δ 10.57 (s, 1H), 9.88 (s, 1H), 7.95-7.90 (m, 1H), 7.54 (d, *J* = 15.6 Hz, 1H), 7.31-7.17 (m, 3H), 7.04 (s, 1H), 6.96 (d, *J* = 15.3 Hz, 1H), 3.52 (s, 2H), 2.22 (s, 3H).
- 15

Example 61: (E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(7-fluoro-1H-benzo[*d*][1,2,3]triazol-6-yl)acrylamide




- 20 Into a 8-mL round-bottom flask, was placed 5-bromo-4-fluoro-3H-1,2,3-benzotriazole (50.0 mg, 0.23 mmol, 1.00 equiv), *N*-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (43.3 mg, 0.23 mmol, 1.00 equiv), Pd(dppf)Cl₂ (16.9 mg, 0.023 mmol, 0.10 equiv), Et₃N (0.096 mL, 0.70 mmol, 3.00 equiv), DMF (3.00 mL). The resulting solution was stirred for 2 h at 120 °C in an oil bath. The reaction mixture was cooled. The residue was applied onto a silica gel column with PE/THF (1/1). This resulted in 19 mg (25%) of (E)-*N*-(2,3-dihydro-1H-inden-1-yl)-3-(7-fluoro-1H-benzo[*d*][1,2,3]triazol-6-yl)acrylamide as a solid. LC-MS (ES, *m/z*):
- 25

[*M*+*H*]⁺=323

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): δ 8.63 (d, J = 8.1 Hz, 1H), 7.93-7.63 (m, 3H), 7.32-7.15 (m, 4H), 6.85 (d, J = 15.9 Hz, 1H), 5.47-5.35 (m, 1H), 3.06 -2.81 (m, 2H), 2.47-2.39 (m, 1H), 1.90-1.78 (m, 1H).

5 **Example 62: (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(4-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide**




Into a 8-mL round-bottom flask, was placed 6-bromo-4-fluoro-1H-1,2,3-benzotriazole (40.00 mg, 0.18 mmol, 1.00 equiv), N-(2,3-dihydro-1H-inden-1-yl)prop-2-enamide (34.7 mg, 0.18 mmol, 1.00 equiv), Pd(dppf)Cl₂ (13.6 mg, 0.019 mmol, 0.10 equiv) and Et₃N (0.077 mL, 0.55 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 1 h at 120 °C in an oil bath. The reaction mixture was cooled. The mixture was applied onto a silica gel column with PE/THF (1/1). This resulted in 17.8 mg (30%) of (E)-N-(2,3-dihydro-1H-inden-1-yl)-3-(4-fluoro-1H-benzo[d][1,2,3]triazol-6-yl)acrylamide as a white solid. LC MS (ES, m/z):

10 [M+H]⁺=323

^1H NMR (300 MHz, $\text{DMSO}-d_6$, ppm): δ 8.51 (d, J = 8.1 Hz, 1H), 7.90 (s, 1H), 7.68 (d, J = 15.6 Hz, 1H), 7.41 (d, J = 12.0 Hz, 1H), 7.28-7.15 (m, 4H), 6.77 (d, J = 15.6 Hz, 1H), 5.47-5.34 (m, 1H), 3.03-2.81 (m, 2H), 2.44-2.35 (m, 1H), 1.93-1.78 (m, 1H).

20 **Example 63: (E)-N-(5-chloro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide**




Into a 8-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed (E)-methyl 3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylate (30.0 mg, 0.14 mmol, 1.00 equiv) in THF (2.00 mL). Then 5-chloro-2-methylbenzenamine (25.4 mg, 0.18 mmol, 1.30 equiv) was added. The reaction mixture was cooled to 0 °C. Then 1M LiHMDS (0.68 mL, 0.68 mmol, 5.00 equiv) was added. The reaction mixture was stirred 1 h at RT. The reaction mixture was

25

quenched by 1 mL water and extracted with 2x5 mL ETOAc, the organic was concentrated under vacuum to afford crude product. The crude product was then purified by Prep-HPLC. This resulted in 5 mg (11%) of (E)-N-(5-chloro-2-methylphenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-5-yl)acrylamide as a white solid. LC-MS (ES, m/z): [M+H]⁺=329


- 5 ¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 9.42 (s, 1H), 7.85 (s, 1H), 7.60 (d, *J*=15.9 Hz, 1H), 7.36-7.11 (m, 5H), 6.97 (d, *J*=15.6 Hz, 1H), 2.26 (s, 3H).

Example 64: (E)-N-(2-methylcyclohexyl)-3-(2-oxoindolin-6-yl)acrylamide




- 10 Into a 8-mL vial, was placed (E)-3-(2-oxoindolin-6-yl)acrylic acid (50.00 mg, 0.25 mmol, 1.00 equiv), 2-methylcyclohexanamine (28.25 mg, 0.25 mmol, 1.00 equiv), HATU (141.38 mg, 0.37 mmol, 1.50 equiv) and DIPEA (95.41 mg, 0.74 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 2 h at 20°C. The mixture was purified by Flash-Prep-HPLC. This resulted in 8.3 mg (11%) of (E)-N-(2-methylcyclohexyl)-3-(2-oxoindolin-6-yl)acrylamide as a
15 off-white solid. LC-MS (ES, m/z): [M+H]⁺=299


Example 65: (E)-3-(3-methyl-1H-indazol-6-yl)-N-(3-methylchroman-4-yl)acrylamide




- Step 1:** Into a 500-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 2,3-dihydro-1-benzopyran-4-one (15.00 g, 101.24 mmol, 1.00 equiv) and THF (200 mL). This was followed by the addition of LiHMDS (1 M in THF, 1.20 equiv) dropwise with stirring at -78°C. The resulting solution was stirred for 40 min at -78°C. To this was added a solution of MeI (17.24 g, 121.49 mmol, 1.20 equiv) in THF (10 mL) dropwise with stirring at -78°C. The resulting solution was allowed to react, with stirring, for an additional
25 40 min at -78°C. Then stirred for an additional 1 h at 25°C. The reaction was then quenched by the addition of 150 mL of NH₄Cl. The resulting solution was extracted with 2x150 mL of ethyl acetate and the organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/10). This resulted in 5 g (30% yield) of 3-methyl-2,3-dihydro-1-benzopyran-4-one as a light yellow oil.



Step 2: Into a 40-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed 3-methyl-2,3-dihydro-1-benzopyran-4-one (1.10 g, 6.78 mmol, 1.00 equiv), MeOH (20.00 mL), $\text{NH}_2\text{OH}\cdot\text{HCl}$ (1.41 g, 20.35 mmol, 3.00 equiv), Et_3N (2.06 g, 20.35 mmol, 3.00 equiv). The resulting solution was stirred for 15 h at 70°C . The resulting mixture was concentrated. The resulting solution was diluted with 10 mL of water. The resulting solution was extracted with 2x10 mL of ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The resulting mixture was concentrated. This resulted in 1 g (83% yield) of N-[(4E)-3-methyl-2,3-dihydro-1-benzopyran-4-ylidene]hydroxylamine as a white solid.




Step 3: Into a 100-mL 1-necked round-bottom flask purged and maintained with an inert atmosphere of H_2 , was placed N-[(4E)-3-methyl-2,3-dihydro-1-benzopyran-4-ylidene]hydroxylamine (0.70 g, 3.95 mmol, 1.00 equiv), MeOH (20.00 mL), Pd/C (0.06 g). The resulting solution was stirred for 12 h at 40°C . The solids were filtered out. The resulting mixture was concentrated. This resulted in 530 mg (82% yield) of 3-methyl-3,4-dihydro-2H-1-benzopyran-4-amine as light yellow oil.



Step 4: Into a 40-mL vial, was placed 3-methyl-3,4-dihydro-2H-1-benzopyran-4-amine (300.00 mg, 1.838 mmol, 1.00 equiv), DCM (10.00 mL), Et_3N (371.98 mg, 3.676 mmol, 2.00 equiv). This was followed by the addition of a solution of acryloyl chloride (182.99 mg, 2.022 mmol, 1.10 equiv) in DCM (2 mL) dropwise with stirring at 0°C . The resulting solution was stirred for 5 h at 0°C . The reaction was then quenched by the addition of 8 mL of water. The resulting solution was extracted with 2x15 mL of dichloromethane and the organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum

ether (1/2). This resulted in 280 mg (70% yield) of N-(3-methyl-3,4-dihydro-2H-1-benzopyran-4-yl)prop-2-enamide as an off-white solid.



5

Step 5: Into a 20-mL vial purged and maintained with an inert atmosphere of nitrogen, was placed N-(3-methyl-3,4-dihydro-2H-1-benzopyran-4-yl)prop-2-enamide (150.00 mg, 0.690 mmol, 1.00 equiv), DMF (6.00 mL), 6-bromo-3-methyl-1H-indazole (145.72 mg, 0.690 mmol, 1.00 equiv), Et₃N (174.65 mg, 1.725 mmol, 2.50 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (33.74 mg, 0.041 mmol, 0.06 equiv). The resulting solution was stirred overnight at 120°C. The reaction mixture was cooled to room temperature. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 110 mg racemic mixture of cis/trans isomers as a light yellow solid. The mixture was purified by Chiral-Prep-HPLC to give the 4 separated stereoisomers, including (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3R,4R)-3-methylchroman-4-yl)acrylamide (Example 65c).

The cis-trans isomers were assigned using NOESY NMR. The stereochemistry of each enantiomeric pair has been assigned arbitrarily.

10

15

Example 65a: (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3R,4S)-3-methylchroman-4-yl)acrylamide

LC-MS (ES, m/z): [M+H]⁺=348

¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.77 (s, 1H), 8.33 (d, *J* = 6.6 Hz, 1H), 7.73-7.61 (m, 3H), 7.30-7.16 (m, 3H), 6.91-6.77 (m, 3H), 5.23-5.20 (m, 1H), 4.13 (d, *J* = 8.1 Hz, 1H),

20

Example 65b: (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3S,4R)-3-methylchroman-4-yl)acrylamide

25

LC-MS1 (ES, m/z): [M+H]⁺=348

¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.77 (s, 1H), 8.33 (d, *J* = 6.9 Hz, 1H), 7.73-7.61 (m, 3H), 7.30-7.16 (m, 3H), 6.91-6.77 (m, 3H), 5.23-5.20 (m, 1H), 4.13 (d, *J* = 8.1 Hz, 1H), 3.94-3.89 (m, 1H), 2.55-2.50 (m, 3H), 2.29-2.33 (m, 1H), 0.91 (d, *J* = 4.5 Hz, 3H).

5

Example 65c: (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3R,4R)-3-methylchroman-4-yl)acrylamide

LC-MS (ES, m/z): [M+H]⁺=348

¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.77 (s, 1H), 8.49 (d, *J* = 6.3 Hz, 1H), 7.73-7.62 (m, 3H), 7.32-7.30 (m, 1H), 7.18-7.13 (m, 2H), 6.91-6.87 (m, 1H), 6.81-6.74 (m, 2H), 4.86-4.82 (m, 1H), 4.23-4.21 (m, 1H), 3.97-3.95 (m, 1H), 2.55-2.50 (m, 3H), 2.08-2.05 (m, 1H), 0.97 (d, *J* = 5.1 Hz, 3H).


Example 65d: (E)-3-(3-methyl-1H-indazol-6-yl)-N-((3S,4S)-3-methylchroman-4-yl)acrylamide

LC-MS (ES, m/z): [M+H]⁺=348

¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.77 (s, 1H), 8.49 (d, *J* = 6.3 Hz, 1H), 7.73-7.62 (m, 3H), 7.32-7.30 (m, 1H), 7.18-7.13 (m, 2H), 6.91-6.87 (m, 1H), 6.81-6.74 (m, 2H), 4.86-4.82 (m, 1H), 4.23-4.21 (m, 1H), 3.97-3.95 (m, 1H), 2.55-2.50 (m, 3H), 2.08-2.05 (m, 1H), 0.97 (d, *J* = 5.1 Hz, 3H).

20


Example 66: (E)-3-(3-methyl-1H-indazol-6-yl)-N-((1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-yl)acrylamide




25

Step 1: Into a 100-mL 3-necked round-bottom flask, was placed oxetan-3-ylmethanol (2.00 g, 22.70 mmol, 1.00 equiv), triflic anhydride (9.61 g, 34.050 mmol, 1.50 equiv), Et₃N (4.59 g, 45.40 mmol, 2.00 equiv), DCM (50.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The resulting mixture was washed with 2x50 ml of NaCO₃ aq. and 1x50 mL of NaCl aq. The organic layer was dried over anhydrous sodium sulfate and concentrated and used for next step without purification.


30



Step 2: Into a 100-mL 3-necked round-bottom flask, was placed tert-butyl N-[(1S,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]carbamate (1.00g, 4.01mmol, 1.00equiv), oxetan-3-ylmethyl trifluoromethanesulfonate (0.88 g, 4.01 mmol, 1.00 equiv), NaH (0.14 g, 6.01 mmol, 1.50 equiv) in THF (20.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The reaction was then quenched by the addition of 5 mL of water. The resulting solution was extracted with 2x20 mL of ethyl acetate and the organic layers combined. The organic layer was concentrated and the residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 150 mg of tert-butyl N-[(1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-yl]carbamate as a solid.



Step 3: Into a 8-mL vial, was placed tert-butyl N-[(1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-yl]carbamate (140.00 mg, 0.438 mmol, 1.00 equiv) in 2M HCl(g) in MeOH (2.00 mL) and MeOH (2.00 mL). The resulting solution was stirred for 2 hr at 10 °C. The resulting mixture was concentrated. This resulted in 35 mg of (1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-amine as a solid.




Step 4: Into a 8-mL vial, was placed (1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-amine (25.00 mg, 0.114 mmol, 1.00 equiv), (E)-3-(3-methyl-1H-indazol-6-yl)acrylic acid (**Intermediate 3**, 22.83 mg, 0.114 mmol, 1.00 equiv), HATU (65.02 mg, 0.171 mmol, 1.50 equiv) and Et₃N (34.61 mg, 0.342 mmol, 3.00 equiv) in DMF (3.00 mL). The resulting solution was stirred for 2 hr at 20 °C. The crude mixture was purified by Prep-HPLC. This resulted in 15 mg of (E)-3-(3-methyl-1H-indazol-6-yl)-N-((1S,2S)-2-(oxetan-3-ylmethoxy)-2,3-dihydro-1H-inden-1-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=404

¹H-NMR (300 MHz, DMSO-*d*₆, *ppm*) δ 12.78 (s, 1H), 8.58 (d, *J* = 8.4 Hz, 1H), 7.74-7.63 (m, 3H), 7.33-7.22 (m, 5H), 6.75 (d, *J* = 15.9 Hz, 1H), 5.34-5.29 (m, 1H), 4.65-4.60 (m, 2H), 4.32-4.28 (m, 2H), 4.17-4.14 (m, 1H), 3.89-3.86 (m, 1H), 3.84-3.74 (m, 1H), 3.36-3.34 (m, 1H), 3.32-3.31 (m, 1H), 2.86-2.78 (m, 1H), 2.50 (s, 3H).

5


Example 67: (E)-N-((1S,2S)-2-(cyclopropylmethoxy)-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide



10

Step 1: Into a 8-mL vial, was placed tert-butyl N-(2-hydroxy-octahydro-1H-inden-1-yl)carbamate (200.00 mg, 0.783 mmol, 1.00 equiv), allyl bromide (94.75 mg, 0.783 mmol, 1.00 equiv) and NaH (5.00 mg, 0.783 mmol, 60%) in THF (3.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The reaction was then quenched by the addition of 0.5 mL of water. The mixture was concentrated and the residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 150 mg (66% yield) of tert-butyl N-[(1S,2S)-2-(prop-2-en-1-yloxy)-2,3-dihydro-1H-inden-1-yl]carbamate as a light yellow solid.


15




20

Step 2: Into a 40-mL vial, was placed tert-butyl N-[(1S,2S)-2-(prop-2-en-1-yloxy)-2,3-dihydro-1H-inden-1-yl]carbamate (140.00 mg, 0.474 mmol, 1.00 equiv), 1M ZnEt₂ (4.74 mL, 4.739 mmol, 10.00 equiv), CH₂I₂ (1015.41 mg, 3.791 mmol, 8.00 equiv) and TFA (432.28 mg, 3.791 mmol, 8.00 equiv) in DCM (5.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The mixture was concentrated and the residue was applied onto a silica gel column with ethyl acetate/petroleum ether (10/1). This resulted in 70 mg (48% yield) of tert-butyl N-[(1S,2S)-2-(cyclopropylmethoxy)-2,3-dihydro-1H-inden-1-yl]carbamate as a solid.

25




Step 3: Into a 8-mL vial, was placed tert-butyl N-[2-(cyclopropylmethoxy)-octahydro-1H-inden-1-yl]carbamate (50.00 mg, 1.00 equiv). in 2M HCl(g) in MeOH (3.00 mL) and MeOH (3.00 mL). The resulting solution was stirred for 2 hr at 20 °C. The resulting mixture was concentrated. This resulted in 30 mg of (1S,2S)-2-(cyclopropylmethoxy)-2,3-dihydro-1H-inden-1-amine as an off-white solid.




Step 4: Into a 8-mL vial, was placed (1S,2S)-2-(cyclopropylmethoxy)-2,3-dihydro-1H-inden-1-amine (20.00 mg, 0.098 mmol, 1.00 equiv), (E)-3-(3-methyl-1H-indazol-6-yl)acrylic acid (19.70 mg, 0.098 mmol, 1.00 equiv), HATU (56.11 mg, 0.148 mmol, 1.50 equiv) and Et₃N (29.87 mg, 0.295 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 2 hr at 20 °C. The mixture was purified by Flash-Prep-HPLC. This resulted in 20 mg (52% yield) of (E)-N-((1S,2S)-2-(cyclopropylmethoxy)-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=388

¹H-NMR (300 MHz, DMSO-*d*₆, ppm) δ 8.57 (d, *J* = 8.7 Hz, 1H), 7.74-7.62 (m, 3H), 7.31 (d, *J* = 7.8 Hz, 1H), 7.25-7.19 (m, 4H), 6.74 (d, *J* = 15.6 Hz, 1H), 5.31-5.27 (m, 1H), 4.19-4.12 (m, 1H), 3.48-3.44 (m, 1H), 3.38-3.32 (m, 2H), 2.84-2.73 (m, 1H), 1.04-1.00 (m, 1H), 0.53-0.42 (m, 2H), 0.24-0.12 (m, 2H)


Example 68: (E)-N-((1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide




Step 1: Into a 100-mL 3-necked round-bottom flask, was placed 2-fluoroethanol (3.20 g, 49.95 mmol, 1.00 equiv), triflic anhydride (14.09 g, 49.95 mmol, 1.00 equiv) and Et₃N (10.11 g, 99.91 mmol, 2.00 equiv) in DCM (60.00 mL). The resulting solution was stirred for 10 hr at -78 °C in a liquid nitrogen bath. The resulting mixture was washed with 2x50 mL of NaCO₃ aq. and 1x50 mL of NaCl aq. The organic layer was dried over anhydrous sodium sulfate and concentrated, the resulted in 9.7 g was used for the next step without purification.



Step 2: Into a 20-mL vial, was placed tert-butyl N-[(1S,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]carbamate (400.00 mg, 1.60 mmol, 1.00 equiv), 2-fluoroethyl trifluoromethanesulfonate (629.32 mg, 3.21 mmol, 2.00 equiv), NaH (77.01 mg, 3.21 mmol, 2.00 equiv) in THF (10.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The reaction was then quenched by the addition of 5 mL of water. The resulting solution was extracted with 2x20 mL of ethyl acetate and the organic layers combined. The organic layer was concentrated and the residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 300 mg (63% yield) of tert-butyl N-[(1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-yl]carbamate as a light yellow solid.




Step 3: Into a 20-mL vial, was placed tert-butyl N-[(1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-yl]carbamate (300.00 mg) in 2M HCl(g) in MeOH (5.00 mL) and MeOH (5.00 mL) was stirred for 10 hr at 20 °C. The resulting mixture was concentrated. This resulted in 120 mg of (1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-amine as a light yellow solid.



Step 4: Into a 8-mL vial, was placed (1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-amine (40.00 mg, 0.205 mmol, 1.00 equiv), ((E)-3-(3-methyl-1H-indazol-6-yl)acrylic acid (**Intermediate 3**, 41.03 mg, 0.205 mmol, 1.00 equiv), HATU (116.85 mg, 0.307 mmol, 1.50 equiv), DIPEA (79.44 mg, 0.615 mmol, 3.00 equiv), DMF (2.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The mixture was purified by Flash-Prep-HPLC. This resulted in 21 mg of (E)-N-((1S,2S)-2-(2-fluoroethoxy)-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=380

¹H-NMR(300 MHz, DMSO-*d*₆, ppm) δ12.78(s, 1H), 8.58 (d, *J* = 8.1 Hz, 1H), 7.74-7.62 (m, 3H), 7.33-7.22(m, 5H), 6.75 (d, *J* = 15.6 Hz, 1H), 5.34-5.29 (m, 1H), 4.65-4.62 (m, 1H), 4.49-4.46 (m, 1H), 4.20-4.16 (m, 1H), 3.97-3.71 (m, 2H), 3.37-3.35 (m, 1H), 2.88-2.80(m, 1H), 2.50 (s, 3H)


Example 69: (E)-N-((1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide



Step 1: Into a 20-mL vial, was placed tert-butyl N-[(1S,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl]carbamate (400.00 mg, 1.604 mmol, 1.00 equiv), ethyl iodide (500.47 mg, 3.209 mmol, 2.00 equiv) and NaH (128.36 mg, 3.209 mmol, 2.00 equiv, 60%) in THF (10.00 mL). The resulting solution was stirred for 10 hr at 20 °C. The mixture was quenched with 5 ml H₂O and extracted with 30 mL of ethyl acetate. The organic layers combined and dried over anhydrous sodium sulfate and concentrated. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 200 mg (45% yield) of tert-butyl N-[(1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-yl]carbamate as a light yellow solid.



Step 2: Into a 20-mL vial, was placed tert-butyl N-[(1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-yl]carbamate (170.00 mg) in 2M HCl(g) in MeOH (5.00 mL) and MeOH (5.00 mL) was stirred for 10 hr at 20 °C. The resulting mixture was concentrated. This resulted in 80 mg of (1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-amine as a light yellow solid.




Step 3: Into a 8-mL vial, was placed (1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-amine (40.00 mg, 0.226 mmol, 1.00 equiv), (E)-3-(3-methyl-1H-indazol-6-yl)acrylic acid (**Intermediate 3**, 45.19 mg, 0.226 mmol, 1.00 equiv), HATU (128.71 mg, 0.339 mmol, 1.50 equiv) and DIPEA (87.50 mg, 0.677 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 2 hr at 20 °C. The mixture was purified by Prep-HPLC. This resulted in 25 mg of (E)-N-((1S,2S)-2-ethoxy-2,3-dihydro-1H-inden-1-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide as a off-white solid.

LC-MS (ES, m/z): [M+H]⁺=362

¹H-NMR (300 MHz, DMSO-*d*₆, ppm) δ 8.57 (d, *J* = 8.4 Hz, 1H), 7.74 -7.63 (m, 3H), 7.33-7.17 (m, 5H), 6.75 (d, *J* = 15.6 Hz, 1H), 5.32-5.28 (m, 1H), 4.14-4.10 (m, 1H), 3.69-3.55 (m, 2H), 3.34-3.26 (m, 1H), 2.83-2.73 (m, 1H), 2.78-2.76 (m, 3H), 1.16-1.12 (m, 3H)


Example 70: (E)-3-(3-cyclopropyl-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide



Into a 8-mL vial, was placed N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (see Example 54, Step 1, 60.00 mg, 0.276 mmol, 1.00 equiv), 6-bromo-3-cyclopropyl-1H-indazole (65.48 mg, 0.276 mmol, 1.00 equiv), Pd(dppf)Cl₂ (20.21 mg, 0.028 mmol, 0.10 equiv), Et₃N (83.83 mg, 0.828 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 3 hr at 120 °C. The mixture was purified by Prep-HPLC. This resulted in 31 mg of (E)-3-(3-cyclopropyl-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=374

¹H-NMR- (300 MHz, DMSO-*d*₆, ppm) δ 8.61-8.58(d, J = 8.7 Hz, 1H), 7.87-7.78 (d, J = 8.7 Hz, 1H), 7.68 -7.62(m, 2H), 7.32-7.20 (m, 5H), 6.77- 6.72 (d, J = 15.9 Hz, 1H), 5.34-5.32 (m, 1H), 4.08-4.07 (m, 1H), 3.38 (s, 3H). 3.35-3.27 (m, 1H), 2.84-2.77 (m, 1H). 2.29-2.24 (m, 1H), 1.01-0.94 (m, 4H).


Example 71: (E)-3-(3-methoxy-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide



Into a 8-mL vial, was placed N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (see Example 54, Step 1, 60.00 mg, 0.276 mmol, 1.00 equiv), 6-bromo-3-methoxy-1H-indazole (62.70 mg, 0.276 mmol, 1.00 equiv), Pd(dppf)Cl₂ (20.21 mg, 0.028 mmol, 0.10 equiv), Et₃N (83.83 mg, 0.828 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 3 hr at 120 °C. The mixture was purified by Prep-HPLC. This resulted in 33 mg of (E)-3-(3-methoxy-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide as an off-white solid. The compounds was converted to the HCl. LC-MS (ES, m/z): [M+H]⁺=364

¹H-NMR (300 MHz, DMSO-*d*₆, ppm) δ 12.09 (s, 1H), 8.59 (d, J = 8.4 Hz, 1H), 7.67-7.54 (m, 3H), 7.26-7.17 (m, 5H), 6.74 (d, J = 15.9 Hz, 1H), 5.34-5.30 (m, 1H), 4.07-4.01 (m, 4H), 3.38 (s, 3H), 3.37-3.29 (m, 1H), 2.84-2.78 (m, 1H)


Example 72: (E)-3-(3-chloro-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide




Into a 8-mL vial, was placed N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (60.00 mg, 0.276 mmol, 1.00 equiv), 6-bromo-3-chloro-1H-indazole (63.92 mg, 0.276 mmol, 1.00 equiv), Pd(dppf)Cl₂ (20.21 mg, 0.028 mmol, 0.10 equiv), Et₃N (83.83 mg, 0.828 mmol, 3.00 equiv) in DMF (4.00 mL). The resulting solution was stirred for 3 hr at 120 °C. The mixture was purified by Prep-HPLC. This resulted in 32 mg of (E)-3-(3-chloro-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=368

¹H-NMR (300 MHz, DMSO-*d*₆, ppm) δ 13.45 (s, 1H), 8.62 (d, *J* = 8.7 Hz, 1H), 7.75-7.67 (m, 3H), 7.46 (d, *J* = 9.6 Hz, 1H), 7.28-7.17 (m, 4H), 6.80 (d, *J* = 15.9 Hz, 1H), 5.35-5.30 (m, 1H), 4.08-4.01 (m, 1H), 3.38 (s, 3H), 3.35-3.27 (m, 1H), 2.84-2.77 (m, 1H)

Example 73: (E)-3-(3-fluoro-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide



Step 1: Into a 40-mL vial, was placed 6-bromo-1H-indazole (500.00 mg, 2.54 mmol, 1.00 equiv), Selectfluor (1797.95 mg, 5.075 mmol, 2.00 equiv) in CH₃CN (10.00 mL) and AcOH (1.00 mL). The resulting solution was stirred for 15 hr at 95 °C. The mixture was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 110 mg (20% yield) of 6-bromo-3-fluoro-1H-indazole as a light yellow solid.




Step 2: Into a 8-mL vial, was placed N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (see Example 54, Step 1, 60.00 mg, 0.276 mmol, 1.00 equiv), 6-bromo-3-fluoro-1H-indazole (59.38 mg, 0.276 mmol, 1.00 equiv), Pd(dppf)Cl₂ (20.21 mg, 0.028 mmol, 0.10 equiv),


Et₃N (83.83 mg, 0.828 mmol, 3.00 equiv) in DMF (2.00 mL). The resulting solution was stirred for 14 hr at 120 °C. The mixture was purified by Prep-HPLC. This resulted in 34 mg of (E)-3-(3-fluoro-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide as a off-white solid. LC-MS1 (ES, m/z): [M+H]⁺=352

- 5 ¹H-NMR(300 MHz, DMSO-*d*₆, ppm) δ 12.72 (s, 1H), 8.62 (d, *J* = 8.7 Hz, 1H), 7.75-7.66 (m, 3H), 7.41 (d, *J* = 8.7 Hz, 1H), 7.27-7.18 (m, 4H), 6.79 (d, *J* = 15.9 Hz, 1H), 5.35-5.30 (m, 1H), 4.08-4.01 (m, 1H), 3.39 (s, 3H), 3.35-3.27 (m, 1H), 2.84-2.78 (m, 1H)

Example 74: (E)-3-(3-cyano-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide



- Step 1:** Into a 8-mL vial, was placed N-[(1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl]prop-2-enamide (see Example 54, Step 1, 70.0 mg, 0.322 mmol, 1.00 equiv), 6-bromo-1-(oxan-2-yl)indazole-3-carbonitrile (98.64 mg, 0.322 mmol, 1.00 equiv), Pd(dppf)Cl₂ (23.57 mg, 0.032 mmol, 0.10 equiv), Et₃N (97.81 mg, 0.967 mmol, 3.00 equiv) in DMF (4.00 mL). The resulting solution was stirred for 3 hr at 120 °C. The mixture was applied onto a silica gel column and eluted with ethyl acetate/petroleum ether (1/1). This resulted in 60 mg (42% yield) of (E)-3-(3-cyano-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide as a light yellow solid.



- Step 2:** Into a 8-mL vial, was placed (E)-3-(3-cyano-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide (60.00 mg, 0.136 mmol, 1.00 equiv) in 2M HCl(g) in MeOH (2.00 mL) and MeOH (2.00 mL). The resulting solution was stirred for 2 hr at 10 °C. The resulting mixture was concentrated. The residue was purified by Prep-HPLC. This resulted in 31 mg of (E)-3-(3-cyano-1H-indazol-6-yl)-N-((1S,2S)-2-methoxy-2,3-dihydro-1H-inden-1-yl)acrylamide as an off-white solid. LC-MS (ES, m/z): [M+H]⁺=359

¹H-NMR (300 MHz, DMSO-*d*₆, ppm) δ 14.52 (s, 1H), 8.64 (d, *J* = 8.4 Hz, 1H), 7.94-7.91 (m, 2H), 7.73 (d, *J* = 15.9 Hz, 1H), 7.62 (d, *J* = 8.7 Hz, 1H), 7.26-7.18 (m, 4H), 6.83 (d, *J* = 15.6 Hz, 1H), 5.34-5.30 (m, 1H), 4.07-4.01 (m, 1H), 3.38 (s, 3H), 3.35-3.27 (m, 1H), 2.84-2.73 (m, 1H).

Example compounds 75 to 106 shown in Table 1a below may be prepared according to methods known to the skilled person, or using the methods disclosed herein:

5 Table 1a: Example compounds 75 to 106

Example No	Structure	Name	Representative scheme(s)
75		(E)-3-(3-(3-fluoro-4-methyl-5-(3-(3-methyl-1H-indazol-6-yl)acrylamido)phenyl)propanoic acid	1 and 13
76		(E)-3-(3-(3-(3-chloro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid	1 and 13
77		(E)-3-(3-(3-fluoro-5-(3-(3-fluoro-1H-indazol-6-yl)acrylamido)-4-methylphenyl)propanoic acid	1 and 13
78		(E)-3-(3-(3-(3-cyano-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid	1 and 13
79		(E)-3-(3-(3-fluoro-5-(3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamido)-4-methylphenyl)propanoic acid	1 and 13
80		(E)-3-(3-(3-(3-chloro-5-fluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid	1 and 13


Exempl e No	Structure	Name	Representati ve scheme(s)
81		(E)-3-(3-(3-(3,5-difluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid	1 and 13
82		(E)-3-(3-(3-(3-cyano-5-fluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid	1 and 13
83		(E)-3-(4-fluoro-3-methyl-2-(3-(3-methyl-1H-indazol-6-yl)acrylamido)phenyl)propanoic acid	1 and 13
84		(E)-3-(2-(3-(3-chloro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid	1 and 13
85		(E)-3-(4-fluoro-2-(3-(3-fluoro-1H-indazol-6-yl)acrylamido)-3-methylphenyl)propanoic acid	1 and 13
86		(E)-3-(2-(3-(3-cyano-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid	1 and 13

Exempl e No	Structure	Name	Representati ve scheme(s)
87		(E)-3-(4-fluoro-2-(3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamido)-3-methylphenyl)propanoic acid	1 and 13
88		(E)-3-(2-(3-(3-chloro-5-fluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid	1 and 13
89		(E)-3-(2-(3-(3,5-difluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid	1 and 13
90		(E)-3-(2-(3-(3-cyano-5-fluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid	1 and 13
91		(E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide	1
92		(E)-3-(3-chloro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1
93		(E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1

Exempl e No	Structure	Name	Representati ve scheme(s)
94		(E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1
95		(E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1
96		(E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1
97		(E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1
98		(E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide	1
99		(E)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)-3-(3-methyl-1H-indazol-6-yl)acrylamide	1 and 14
100		(E)-3-(3-chloro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide	1 and 14


Exempl e No	Structure	Name	Representati ve scheme(s)
101		(E)-3-(3-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide	1 and 14
102		(E)-3-(3-cyano-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide	1 and 14
103		(E)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamide	1 and 14
104		(E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide	1 and 14
105		(E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide	1 and 14
106		(E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide	1 and 14

Example 91: (2E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide



Step 1: A solution of (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (**Intermediate 13**, 0.22 g, 0.7 mmol), 6-methoxy-2,4-dimethylpyridin-3-amine (0.12 g, 0.9 mmol, 1.2 equiv), HATU (0.58 g, 1.5 mmol, 2 equiv) and DIEA (0.10g, 1.5 mmol, 2 equiv) in DMF (5 mL) was stirred for 2 h at 60°C. The mixture was cooled and diluted with water (8 mL), then extracted with EtOAc (3 x 5 mL). The combined organics were washed with water (3x5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with hexane/AcOEt (1:1) to afford (2E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enamide (0.15 g, 46% yield) as a white solid.

LCMS (ES, m/z): 421 [M+H]⁺




Step 2: A solution of (2E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enamide (0.15 g, 0.4 mmol) in 4M HCl in 1,4-dioxane (2 mL) was stirred for 1 h at 60 °C. The mixture was cooled and concentrated, then basified with NaHCO₃ solution. The crude product was purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5µm; Mobile Phase: 24-58% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 10 mins; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide (60 mg, 37% yield) as a white solid.


LCMS (ES, m/z): 337 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 12.82 (s, 1H), 9.51 (s, 1H), 7.87-7.57 (m, 3H), 7.39 (d, *J* = 8.5 Hz, 1H), 6.93 (d, *J* = 15.7 Hz, 1H), 6.59 (s, 1H), 3.82 (s, 3H), 2.51 (s, 3H), 2.28 (s, 3H), 2.14 (s, 3H).

Example 93: (2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl) prop-2-enamide



Step 1: A solution of (2E)-3-[3-fluoro-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (**Intermediate 6**, 150 mg, 0.5 mmol), 6-methoxy-2,4-dimethylpyridin-3-amine (102 mg, 0.6 mmol, 1.3 equiv), HATU (392 mg, 1.0 mmol, 2.0 equiv) and DIEA (133 mg, 1.0 mmol, 2.0 equiv) in DMF (3 mL) was stirred for 1 h at 60°C. The reaction was cooled and diluted with H₂O (10 mL). The resulting mixture was extracted with ethyl acetate (3 x 10 mL), and the combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether: ethyl acetate (3:1) to afford (2E)-3-[3-fluoro-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (164 mg, 75% yield).




Step 2: A solution of (2E)-3-[3-fluoro-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (162 mg, 0.4 mmol) and TFA (3.5 mL) in DCM (11 mL) was stirred for 2 hours at room temperature. The resulting mixture was diluted with H₂O (5 mL) and basified with NaHCO₃. The resulting mixture was extracted with CH₂Cl₂ (3 x 5 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH (3 mL) and purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5µm; Mobile Phase: 25-57% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 9 mins; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (32 mg, 25% yield) as an off-white solid.

LC-MS(ES, m/z): 341 [M+H]⁺


¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 12.75 (s, 1H), 9.56 (s, 1H), 7.79-7.69 (m, 3H), 7.50-7.47 (m, 1H), 6.97 (d, *J* = 15.8 Hz, 1H), 6.60 (s, 1H), 3.82 (s, 3H), 2.29 (s, 3H), 2.14 (s, 3H).

Example 94: (2E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide




Step 1: A solution of 6-bromo-1H-indazole-3-carbonitrile (0.30 g, 1.3 mmol), DHP (0.68 g, 8.1 mmol, 6 equiv) and PPTS (0.10 g, 0.3 mmol, 0.3 equiv) in THF (6 mL) was stirred for 10 h at 70°C. The mixture was cooled and diluted with saturated brine (10 mL) and saturated aqueous NaHCO₃ (2 mL). The aqueous was extracted with EtOAc (2x8 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (5:1) to afford 6-bromo-1-(oxan-2-yl)indazole-3-carbonitrile (0.32 g, 77% yield) as an light yellow solid.

LCMS (ES, m/z): 306 [M+H]⁺



Step 2: A mixture of N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 4**, 0.06 g, 0.3 mmol), 6-bromo-1-(oxan-2-yl)indazole-3-carbonitrile (0.11 g, 0.4 mmol, 1.2 equiv), Pd(dppf)Cl₂ (0.07 g, 0.1 mmol, 0.3 equiv) and TEA (0.09 g, 0.9 mmol, 3 equiv) in DMF (1.1 mL) was stirred for 2 h at 120°C under N₂ atmosphere. The mixture was cooled and diluted with water (5 mL), then extracted with EtOAc (3 x8 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/ EtOAc (1:1) to afford (2E)-3-[3-cyano-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (0.06 g, 49% yield) as a white solid.

LCMS (ES, m/z): 432 [M+H]⁺




Step 3: To (2E)-3-[3-cyano-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (0.05 g, 0.1 mmol) was added 4M HCl in 1,4-dioxane (1.5 mL) and the reaction stirred for 30 mins, before being concentrated. The residue was dissolved in MeOH (3 mL) and basified with NaHCO₃ solution. The aqueous was extracted with EtOAc (3x10 mL), and the combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was

dissolved in MeOH (1 mL) and purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5 μ m; Mobile Phase: 21-63% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 12 mins; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (8.0 mg, 18% yield) as a white solid.

LCMS (ES, m/z): 348 [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 9.56 (s, 1H), 7.99 (s, 1H), 7.93 (d, *J* = 8.6 Hz, 1H), 7.76 (d, *J* = 15.8 Hz, 1H), 7.67 (d, *J* = 8.5 Hz, 1H), 7.01 (d, *J* = 15.5 Hz, 1H), 6.60 (s, 1H), 3.83 (s, 3H), 2.29 (s, 3H), 2.15 (s, 3H).

Example 95: (2E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide




A solution of N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 4**, 150 mg, 0.7 mmol), 6-bromo-5-fluoro-3-methyl-1H-indazole (**Intermediate 5**, 200 mg, 0.8 mmol, 1.2 equiv), TEA (220 mg, 2.1 mmol, 3.0 equiv) and Pd(dppf)Cl₂ (178 mg, 0.2 mmol, 0.3 equiv) in DMF (3 mL) was stirred for 4 hours at 120°C under N₂ atmosphere. The mixture was cooled, filtered, and the filter cake washed with THF (3 mL). The combined filtrate was concentrated, and the residue purified by Prep-HPLC (Column: XBridge Prep OBD C18 Column, 30*150 mm, 5 μ m; Mobile Phase: 18-55% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 45 mins; Flow rate: 60 mL/min). Concentration of the appropriate fractions gave (2E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (77 mg, 29% yield) as an off-white solid.

LCMS (ES, m/z): 355 [M+H]⁺.

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 12.93 (s, 1H), 9.61 (s, 1H), 7.79 (d, *J* = 5.9 Hz, 1H), 7.72 (d, *J* = 15.9 Hz, 1H), 7.61 (s, 1H), 7.05 (d, *J* = 15.9 Hz, 1H), 6.59 (s, 1H), 3.83 (s, 3H), 2.49 (s, 3H), 2.29 (s, 3H), 2.14 (s, 3H).

Example 96: (2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide




A solution of 6-bromo-3-chloro-5-fluoro-1H-indazole (**Intermediate 7**, 109 mg, 0.4 mmol, 1.2 equiv), N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 4**, 75 mg, 0.3 mmol, 1 equiv), Pd(dppf)Cl₂•CH₂Cl₂ (89 mg, 0.1 mmol, 0.3 equiv) and TEA (110 mg, 1.0 mmol, 3 equiv) in DMF (1.5 mL) was stirred for 6 h at 110°C under N₂ atmosphere. The mixture was cooled and diluted with water (10 mL). The resulting mixture was extracted with EtOAc (3 x 10 mL). The combined organics were washed with brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5µm; Mobile Phase: 20-56% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 45 mins; Flow rate: 60 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (31 mg, 23% yield) as an off-white solid.

LCMS (ES, m/z): 375 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 13.61 (s, 1H), 9.66 (s, 1H), 7.95 (d, *J* = 5.8 Hz, 1H), 7.73 (d, *J* = 15.9 Hz, 1H), 7.60 (d, *J* = 10.7 Hz, 1H), 7.10 (d, *J* = 15.9 Hz, 1H), 6.60 (s, 1H), 3.83 (s, 3H), 2.29 (s, 3H), 2.14 (s, 3H).


Example 97: (2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide



Step 1: A solution of (2E)-3-[3,5-difluoro-1-(oxan-2-yl) indazol-6-yl] prop-2-enoic acid (**Intermediate 8**, 63 mg, 0.2 mmol), 6-methoxy-2,4-dimethylpyridin-3-amine (30 mg, 0.2 mmol, 1.0 equiv), HATU (152 mg, 0.4 mmol, 2 equiv) and DIEA (52 mg, 0.4 mmol, 2 equiv) in DMF (1

mL) was stirred for 1 hours at 60°C. The mixture was cooled and diluted with water, then extracted with ethyl acetate (4x5 mL). The combined organics were washed with water (3x5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether/THF (2:1) to afford (2E)-3-[3,5-difluoro-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (73 mg, 78% yield) as a light yellow solid.

LC-MS (ES, m/z): 443 [M+H]⁺




Step 2: A solution of (2E)-3-[3,5-difluoro-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl) prop-2-enamide (70 mg, 0.2 mmol) and TFA (0.3 mL) in DCM (1.2 mL) was stirred for 2 hours at room temperature. The mixture was basified with saturated aqueous NaHCO₃ and extracted with CH₂Cl₂ (3 x 5 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH (3 mL) and purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5μm; Mobile Phase: 25-58% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 13 min; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl) prop-2-enamide (9.8 mg, 19% yield) as an off-white solid.

LC-MS (ES, m/z): 359 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 12.88 (s, 1H), 9.66 (s, 1H), 7.86 (dd, *J* = 5.9, 2.4 Hz, 1H), 7.75-7.64 (m, 2H), 7.09 (d, *J* = 16.0 Hz, 1H), 6.60 (s, 1H), 3.83 (s, 3H), 2.29 (s, 3H), 2.14 (s, 3H).

Example 98: (2E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide



Step 1: A mixture of N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 4**, 100 mg, 0.5 mmol), 6-bromo-5-fluoro-1-(oxan-2-yl)indazole-3-carbonitrile (**Intermediate 9**, 190 mg,

0.6 mmol, 1.2 equiv), Pd(dppf)Cl₂·CH₂Cl₂ (110 mg, 0.1 mmol, 0.2 equiv) and TEA (150 mg, 1.5 mmol, 3 equiv) in DMF (2 mL) was stirred for 4 h at 110 °C under nitrogen atmosphere. The mixture was cooled and diluted with H₂O (10 mL), then extracted with EtOAc (3x 6 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/EtOAc (2:1) to afford (E)-3-(3-cyano-5-fluoro-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide. (123 mg, 56% yield) as an off-white solid.




Step 2: A solution of (2E)-3-[3-cyano-5-fluoro-1-(oxan-2-yl)indazol-6-yl]-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (123 mg, 0.3 mmol) in DCM (1.5 mL) and TFA (0.5 mL) was stirred for 3 h then diluted with H₂O (3 mL). The mixture was basified with NaHCO₃ solution, and extracted with EtOAc (3 x 10 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH (3 mL) and purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5µm; Mobile Phase: 21-52% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃·H₂O) over 10 min; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide (7.8 mg, 7% yield) as an off-white solid.

LCMS (ES, m/z): 366 [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 14.69 (br, s, 1H), 9.69 (s, 1H), 8.14 (d, *J* = 5.9 Hz, 1H), 7.88 (d, *J* = 10.7 Hz, 1H), 7.74 (d, *J* = 15.9 Hz, 1H), 7.13 (d, *J* = 16.0 Hz, 1H), 6.60 (s, 1H), 3.83 (s, 3H), 2.29 (s, 3H), 2.14 (s, 3H).

Example 107: (2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)prop-2-enamide



To a solution of 6-bromo-5-fluoro-3-methyl-1H-indazole (**Intermediate 5**, 0.10 g, 0.4 mmol) in DMF (1 mL) was added N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 10**,


0.10 g, 0.5 mmol, 1.2 eq), TEA (0.13 g, 1.3 mmol, 3 eq) and Pd(dppf)Cl₂ (0.09 g, 0.1 mmol, 0.3 eq). The reaction was stirred for 2h at 110°C under N₂ atmosphere, then cooled and quenched with water (10 mL). The resulting mixture was extracted with EtOAc (3 x 10mL). The combined organics were washed with brine (3 x 10 mL), dried over anhydrous Na₂SO₄ and concentrated.

The residue was dissolved in MeOH (2mL) and purified by Prep-HPLC: Column: YMC-Actus Triart C18, 30*150 mm, 5µm; Mobile Phase: 24-49% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 7 min; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)prop-2-enamide (35 mg, 23% yield) as a white solid.

LCMS (ES, m/z): 343 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 12.97 (s, 1H), 9.99 (s, 1H), 8.34 (s, 1H), 7.84-7.78 (m, 2H), 7.73 (d, *J* = 11.4 Hz, 1H), 7.64 (d, *J* = 11.3 Hz, 1H), 2.50 (s, 3H), 2.37 (s, 3H), 2.14 (s, 3H).

Example 108: (2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide




To a solution of N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 10**, 0.10 g, 0.5 mmol) in DMF (1 mL) was added 6-bromo-3-fluoro-1H-indazole (0.13 g, 0.6 mmol, 1.2 equiv), Pd(dppf)Cl₂•CH₂Cl₂ (0.11 g, 0.1 mmol, 0.2 eq) and TEA (0.16 g, 1.5 mmol, 3.0 eq). The reaction was stirred for 2h at 110°C under N₂ atmosphere. The resulting mixture was quenched with water (10 mL) and extracted with EtOAc (3 x 10mL). The combined organic layers were washed with brine (3 x 5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH (2 mL) and purified by Prep-HPLC (Column: YMC-Actus Triart C18, 30*150 mm, 5µm; Mobile Phase: 24-49% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 7 min; Flow rate: 30 mL/min). Concentration of the appropriate fractions gave (2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (24 mg, 13% yield) as a white solid.

LCMS (ES, m/z): 329 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆, ppm) δ 12.77 (s, 1H), 9.93 (s, 1H), 8.34 (s, 1H), 7.93-7.65 (m, 3H), 7.50 (d, *J* = 8.5 Hz, 1H), 7.01 (d, *J* = 15.8 Hz, 1H), 2.37 (s, 3H), 2.14 (s, 3H).

Example 109: (2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide




A solution of N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 10**, 100 mg, 0.5 mmol), 6-bromo-3-chloro-5-fluoro-1H-indazole (**Intermediate 7**, 154 mg, 0.6 mmol, 1.2 equiv), TEA (156 mg, 1.5 mmol, 3 equiv) and Pd(dppf)Cl₂·CH₂Cl₂ (126 mg, 0.1 mmol, 0.2 equiv) in DMF (2 mL) was stirred for 4 h at 110°C under N₂ atmosphere. The mixture cooled, filtered, and the filter cake was washed with THF (5 mL). The combined filtrate was concentrated and purified by Prep-HPLC: Column: XBridge Prep OBD C18 Column, 30*150 mm, 5μm; Mobile Phase: 20-56% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃·H₂O) over 15 min; Flow rate: 60 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl) prop-2-enamide (30 mg, 16% yield) as an off-white solid.

LCMS (ES, m/z): 363 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 13.95 (s, 1H), 10.46 (s, 1H), 8.54 (s, 1H), 7.97 (s, 1H), 7.82-7.76 (m, 1H), 7.60 (d, *J* = 10.6 Hz, 1H), 7.23 (d, *J* = 16.1 Hz, 1H), 2.46 (s, 3H), 2.21 (s, 3H).


Example 110: (2E)-N-(5-fluoro-4-methylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide



Step 1: To a solution of (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (**Intermediate 13**, 0.15 g, 0.5 mmol) in DMF (1.0 mL) was added 5-fluoro-4-methylpyridin-3-amine (**Intermediate 14**, 0.13 g, 1.0 mmol, 2.0 eq) and DIEA (0.20 g, 1.6 mmol, 3.0 eq), followed by T₃P (0.67 g, 1.0 mmol, 2.0 eq, 50% in EtOAc). The mixture was stirred for 1h, then quenched with water. The resulting mixture was extracted with EtOAc (3 x 10 mL), and the combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated. The crude product was triturated with petroleum ether/ethyl acetate (1:5) to afford ((2E)-N-(5-fluoro-4-

methylpyridin-3-yl)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enamide (150 mg, 72% yield) as an off-white solid.

LCMS (ES, m/z): 395 $[M+H]^+$




- 5 **Step 2:** To (2E)-N-(5-fluoro-4-methylpyridin-3-yl)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enamide (90 mg, 0.2 mmol) was added 4M HCl in 1,4-dioxane (2.0 mL). The mixture was stirred for 10 min at 60°C. The resulting mixture was concentrated and the residue was dissolved in MeOH (2 mL) and was basified to pH8 with $NH_3 \cdot H_2O$. The solids were filtered and washed with MeOH, then dried to afford (2E)-N-(5-fluoro-4-methylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide (30 mg, 43% yield) as an off-white solid.

LCMS (ES, m/z): 311 $[M+H]^+$

1H NMR (400 MHz, $DMSO-d_6$, ppm) δ 12.84 (s, 1H), 9.86 (s, 1H), 8.69 (s, 1H), 8.35 (s, 1H), 7.82-7.74 (m, 2H), 7.71 (s, 1H), 7.41 (d, $J = 8.4$ Hz, 1H), 7.06 (d, $J = 15.7$ Hz, 1H), 2.50 (s, 3H), 2.23 (s, 3H).

15

Example 111: (2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide




- To a solution of 6-bromo-3,5-difluoro-1H-indazole (0.1 g, 0.4 mmol) in DMF (1 mL) was added N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 10**, 0.1 g, 0.5 mmol, 1.2 eq), TEA (130 mg, 1.3 mmol, 3 eq) and $Pd(dppf)Cl_2$ (10 mg, 0.13 mmol, 0.3 eq). The reaction was stirred for 2h at 110°C under N_2 atmosphere, then was cooled and quenched with water (10 mL). The resulting mixture was extracted with EtOAc (3 x 10mL). The combined organics were washed with brine (3 x 5 mL), dried over anhydrous Na_2SO_4 and concentrated. The residue was dissolved in MeOH (2 mL) and purified by Prep-HPLC: Column: YMC-Actus Triart C18, 30*150 mm, 5 μ m; Mobile Phase: 24-49% MeCN / Water (10 mmol/L NH_4HCO_3 + 0.1% $NH_3 \cdot H_2O$) over 9 min; Flow

rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (45 mg, 31% yield) as a white solid.

LCMS (ES, m/z): 347 $[M+H]^+$

1H NMR (400 MHz, $DMSO-d_6$, ppm) δ 12.84 (br, s, 1H), 10.04 (s, 1H), 8.34 (s, 1H), 7.88 (d, J = 6.1 Hz, 1H), 7.81-7.65 (m, 2H), 7.13 (d, J = 15.9 Hz, 1H), 2.37 (s, 3H), 2.14 (s, 3H).

Example 112: (2E)-3-(3-chloro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide




To a solution of N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 10**, 0.10 g, 0.5 mmol) in DMF (1 mL) was added 6-bromo-3-chloro-1H-indazole (0.14 g, 0.6 mmol, 1.2 equiv), TEA (0.16 g, 1.5 mmol, 3 eq) and Pd(dppf)Cl₂ (0.11 g, 0.15 mmol, 0.3 eq). The reaction was stirred for 2h at 110°C under N₂ atmosphere, then quenched with water (10 mL). The resulting mixture was extracted with EtOAc (3 x 10mL). The combined organics were washed with brine (3 x 5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH (2 mL) and purified by Prep-HPLC: Column: YMC-Actus Triart C18, 30*150 mm, 5 μ m; Mobile Phase: 24-49% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 9 min; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-chloro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (64 mg, 36% yield) as a white solid.

LCMS (ES, m/z): 345 $[M+H]^+$

1H NMR (300 MHz, $DMSO-d_6$, ppm) δ 13.50 (s, 1H), 9.93 (s, 1H), 8.34 (s, 1H), 7.86-7.69 (m, 3H), 7.55 (d, J = 8.5 Hz, 1H), 7.02 (d, J = 15.8 Hz, 1H), 2.37 (s, 3H), 2.14 (s, 3H).

Example 113: (2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide




To a solution of N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide (**Intermediate 10**, 0.16 g, 0.8 mmol) in DMF (1.5 mL) was added 6-bromo-3-methyl-1H-indazole (0.21 g, 0.9 mmol, 1.2 eq),

TEA (0.25 g, 2.4 mmol, 3 eq) and Pd(dppf)Cl₂ (0.06 g, 0.1 mmol, 0.1 eq). The mixture was stirred for 2h at 110°C under N₂ atmosphere. The resulting mixture was cooled and quenched with water (10 mL), then extracted with EtOAc (3 x 10mL). The combined organics were washed with brine (3x5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH (2 mL) and purified by Prep-HPLC: Column: YMC-Actus Triart C18, 30*150 mm, 5µm; Mobile Phase: 24-49% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 7 mins; Flow rate: 30 mL/min. Concentration of the appropriate fractions gave (2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide (38.5 mg, 14% yield) as a white solid.

LCMS (ES, m/z): 325 [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 12.83 (s, 1H), 9.88 (s, 1H), 8.33 (s, 1H), 7.85-7.67 (m, 3H), 7.41 (d, *J* = 8.5 Hz, 1H), 6.98 (d, *J* = 15.7 Hz, 1H), 2.51 (s, 3H), 2.38 (s, 3H), 2.14 (s, 3H).

Example 114: (2E)-N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide




To a solution of N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]prop-2-enamide (**Intermediate 12**, 0.10 g, 0.5 mmol) in DMF (2 mL) was added 6-bromo-3-methyl-1H-indazole (0.12 g, 0.6 mmol, 1.2 eq), TEA (0.14 g, 1.4 mmol, 3 eq) and Pd(dppf)Cl₂ (69.6 mg, 0.1 mmol, 0.2 eq). The reaction was stirred for 2h at 110°C under N₂ atmosphere, then cooled and quenched by water. The resulting mixture was extracted with EtOAc (3 x 5 mL). The combined organics were washed with brine (3 x 5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with CH₂Cl₂/MeOH (10:1) to afford (2E)-N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide (56 mg, 34% yield) as an off-white solid.

LCMS (ES, m/z): 341 [M+H]⁺


¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 12.85 (s, 1H), 9.83 (s, 1H), 8.88 (s, 1H), 8.43 (s, 1H), 7.80 (d, *J* = 7.5 Hz, 1H), 7.77 (s, 1H), 7.72 (s, 1H), 7.43 (d, *J* = 8.5 Hz, 1H), 7.09 (d, *J* = 15.7 Hz, 1H), 4.59 (s, 2H), 3.30 (s, 3H), 2.51 (s, 3H).

Example 115: (2E)-3-(3-methyl-1H-indazol-6-yl)-N-(2-methylpyridin-3-yl)prop-2-enamide



Step 1: To a solution of (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (**Intermediate 13**, 0.12 g, 0.4 mmol) in DMF (2 mL) was added HATU (0.23 g, 0.6 mmol, 1.5 eq) and DIEA (0.16 g, 1.3 mmol, 3.0 eq). The mixture was stirred for 5 min then 2-methylpyridin-3-amine (68 mg, 0.6 mmol, 1.5 eq) was added. The reaction was stirred for 2h before being quenched by water (10 mL). The resulting mixture was extracted with EtOAc (3 x 10 mL). The combined organics were washed with brine (3x5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with n-hexane/AcOEt (1:1) to afford (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]-N-(2-methylpyridin-3-yl)prop-2-enamide (100 mg, 63% yield) as a white solid.

LCMS (ES, m/z): 377 [M+H]⁺




Step 2: A solution of (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]-N-(2-methylpyridin-3-yl)prop-2-enamide (95 mg, 0.3 mmol) in 4M HCl in 1,4-dioxane (1 mL) was stirred for 4h at 60°C. The resulting mixture was concentrated and the residue was dissolved in MeOH (2 mL). The solution was basified with aq. NH₃•H₂O, then purified by prep-HPLC: Column: XBridge C18, 19*150 mm, 5 um; Mobile Phase: 20-65% MeCN / 20 mM aqueous NH₄HCO₃ + 0.05% NH₃•H₂O over 8 mins; Flow rate: 40mL/min. Concentration of the appropriate fractions gave (2E)-3-(3-methyl-1H-indazol-6-yl)-N-(2-methylpyridin-3-yl)prop-2-enamide (31 mg, 41% yield) as a white solid.

LCMS (ES, m/z): 293 [M+H]⁺


¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 12.88 (br, s, 1H), 10.28 (s, 1H), 8.77 (d, *J* = 8.4 Hz, 1H), 8.53 (d, *J* = 4.8 Hz, 1H), 7.87-7.76 (m, 3H), 7.74 (s, 1H), 7.42 (d, *J* = 8.4 Hz, 1H), 7.22 (d, *J* = 16.0 Hz, 1H), 2.74 (s, 3H), 2.51 (s, 3H).

Example 116: (2E)-N-(2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide



Step 1: A solution of (2E)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enoic acid (**Intermediate 13**, 100 mg, 0.4 mmol), 2,4-dimethylpyridin-3-amine (51 mg, 0.4 mmol, 1.0 equiv) and DIEA (90 mg, 0.7 mmol, 2.0 equiv) in DMF (2 mL) were treated with 50% T₃P in EtOAc (445 mg, 1.4 mmol, 4.0 equiv), then stirred for 2 hours at 60°C. The mixture was allowed to cool, then quenched with water. The resulting mixture was extracted with ethyl acetate (4x50 mL). The combined organics were washed with water (3x5 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether:ethyl acetate (5:1) to afford methyl (2E)-3-[3,5-difluoro-1-(oxan-2-yl)indazol-6-yl]prop-2-enoate (43 mg, 42% yield) as a brown solid.

LC-MS (ES, m/z): 391 [M+H]⁺




Step 2: A solution of (2E)-N-(2,4-dimethylpyridin-3-yl)-3-[3-methyl-1-(oxan-2-yl)indazol-6-yl]prop-2-enamide (65 mg, 0.2 mmol) and TFA (0.4 mL) in DCM (4 mL) was stirred for 2 hours at room temperature. The mixture was basified with saturated aqueous NaHCO₃, and the resulting mixture extracted with CH₂Cl₂ (2 x 10 mL). The combined organics were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography, eluting with petroleum ether:ethyl acetate (1:1) to afford (2E)-N-(2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide (17 mg, 34% yield) as a white solid.

LC-MS (ES, m/z): 307 [M+H]⁺

¹H NMR (300 MHz, DMSO-*d*₆) δ 12.81 (br, s, 1H), 9.71 (s, 1H), 8.23 (d, *J* = 4.9 Hz, 1H), 7.77-7.65 (m, 3H), 7.40 (dd, *J* = 8.4, 1.3 Hz, 1H), 7.16 (d, *J* = 5.0 Hz, 1H), 6.96 (d, *J* = 15.8 Hz, 1H), 2.50 (s, 3H), 2.38 (s, 3H), 2.20 (s, 3H).

Example 117: (2E)-N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide



To N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]prop-2-enamide (**Intermediate 11**, 0.10 g, 0.4 mmol) in DMF (3 mL) was added 6-bromo-3-methyl-1H-indazole (0.11g, 0.5 mmol, 1.2 eq), Pd(dppf)Cl₂ (98 mg, 0.1 mmol, 0.3 eq) and TEA (135 mg, 1.3 mmol, 3 eq). The mixture was stirred for 2h at 110°C under N₂ atmosphere, then cooled and quenched with water (10 mL). The resulting mixture was extracted with EtOAc (3 x 10mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated. The residue was dissolved in MeOH and purified by Prep-HPLC (Column: YMC-Actus Triart C18, 30*150 mm, 5µm; Mobile phase: 24-49% MeCN / Water (10 mmol/L NH₄HCO₃ + 0.1% NH₃•H₂O) over 7 mins; Flow rate: 30 mL/min; Wave Length: 254 nm). Concentration of the appropriate fractions gave (2E)-N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide (38 mg, 24% yield) as an off-white solid.

LCMS (ES, m/z): 355 [M+H]⁺

¹H NMR (400 MHz, DMSO-*d*₆, ppm) δ 12.84 (s, 1H), 9.96 (s, 1H), 8.44 (s, 1H), 7.81-7.68 (m, 3H), 7.41 (d, *J* = 8.4 Hz, 1H), 7.00 (d, *J* = 15.8 Hz, 1H), 4.44 (s, 2H), 3.23 (s, 3H), 2.50 (s, 3H), 2.39 (s, 3H).

Biological Examples

Biological Example 1 – mPTP activity assay in isolated rat liver mitochondria, human platelet mitochondria and isolated rat brain mitochondria

Rat liver mitochondria assay

Pharmacological inhibition or modulation of the mPTP can be measured in well characterised 'Ca²⁺ retention' assays performed in isolated mitochondria. *In vitro*, isolated mitochondria rapidly sequester exogenous Ca²⁺ until the intramitochondrial Ca²⁺ concentration reaches the threshold for mPTP activation. Once the pore is activated, mitochondrial integrity is compromised and the stored Ca²⁺ is released. The distribution of Ca²⁺ between extra- and intra-mitochondrial compartments can be measured in real time with the use of membrane-impermeant Ca²⁺ sensitive fluorescent dyes. Depending on the configuration of the assay, inhibition or modulation of the mPTP either delays the opening of the pore or increases the concentration of Ca²⁺ required to induce mPTP opening.

MPTP activity was measured in mitochondria freshly isolated from female Sprague Dawley (250 to 300 gram) rat livers using the following method. Cervical dislocation was performed on the rat. The liver was then perfused *in-situ* with ~40 ml cold Dulbecco's Phosphate Buffered Saline (DPBS) prior to dissection and transferred into 30 ml Isolation Buffer (250mM Sucrose, 10mM KCl, 1mM EGTA, 1mM EDTA, 25mM HEPES, adjusted to pH 7.5 with 1M NaOH). Each lobe of the liver was then removed from the buffer, minced using tweezers and a scalpel into ~5mm pieces then transferred into a 50 ml Potterton dounce homogenization tube on ice containing 30 ml ice-cold centrifugation buffer (300mM Trehalose, 25mM HEPES, 1mM EGTA, 1mM EDTA, 10mM KCl, adjusted to pH 7.5 with 1M NaOH and supplemented with 0.1% bovine serum albumin (BSA) and complete protease inhibitor cocktail (one tablet of inhibitor per 50mls of buffer). Homogenisation was carried out using a teflon pestle at 1800 rpm. The slurry was centrifuged at 800 g for 10 min at 4 °C, then the supernatant centrifuged at 10,000 g for 10 min. The pellet was washed once with FLIPR assay buffer (75mM Mannitol, 25mM Sucrose, 5mM Potassium Phosphate Monobasic, 20mM Tris base, 100mM KCl, 0.1 % BSA adjusted to pH 7.4 with 5M HCl) centrifuged again, then resuspended in FLIPR assay buffer to a concentration of 8.8 mg/ml protein.

Tested compounds (10 mM stock in DMSO) were serially diluted in DMSO in half log steps to generate 10 test concentrations (final concentrations in assay 30 μ M to 1 nM). An intermediate dilution of 5 μ l DMSO samples into 247 μ l FLIPR assay buffer was carried out prior to transfer of 5 μ l into duplicate wells of a 384 well polypropylene assay plate. Control wells were 0.5 % (v/v) DMSO and 5 μ M cyclosporin A.

A stock mitochondria/Fluo5N assay solution was prepared in 5.6 ml FLIPR assay buffer (at RT) supplemented with succinate disodium salt (10mM), rotenone (1 μ M), Fluo5N pentapotassium salt (2 μ M) and 1 ml mitochondria suspension, then transferred (15 μ l) into the assay plate containing test compounds and incubated for 10 min at RT. Assay plates were then transferred to a FLIPR Tetra plate reader (Molecular Devices). Dye fluorescence was then measured every 3 sec for a total of 10 min. After 12 sec, a 2.5 μ l bolus of CaCl_2 (75 μ M) was added from a source plate containing 675 μ M CaCl_2 in FLIPR assay buffer. IC₅₀ values for tested compounds were calculated using the fluorescence value collected at the 10 min timepoint with % inhibition calculated using the DMSO control and cyclosporin A values as 100 and 0 % respectively.

Human platelet mitochondria assay

The mPTP cell based assay was performed using a mitochondria membrane potential flow cytometry assay in stimulated human platelets. Simulation of platelets results in the rapid influx of Ca^{2+} across the platelet membrane. The Ca^{2+} is then sequestered by mitochondria until the

threshold for mPTP opening is reached, at which point the pore opens and the mitochondrial membrane potential is dissipated. Changes in mitochondria membrane potential due to mPTP opening can be quantified in live platelets using standard mitochondrial membrane potential dyes e.g. 3,3'-dihexyloxacarbocyanine Iodide; DiOC6(3), enabling pharmacological characterisation of mPTP inhibitors.

Fresh human blood (20 ml) was collected from consenting donors into 3.2 % sodium citrate. Platelets were isolated by centrifugation at 200 g for 20 min at room temperature, then the platelet rich plasma layer transferred to a fresh tube. Prostaglandin 12 is added to the platelets at a final concentration of 20 ng/ml. After centrifugation at 640 g for 10 min, the platelet rich pellet was resuspended in 4 ml HEPES assay buffer (137 mM NaCl, 2.7 mM KCl, 11.9 mM NaHCO₃, 0.42 mM NaH₂PO₄, 1 mM MgCl₂, 5.5 mM glucose, 0.1 % bovine serum albumin, 10 mM HEPES adjusted to pH 7.4) and stored on ice.

Test compounds were prepared from 10 mM stocks in DMSO and serially diluted in assay buffer containing 0.4 % DMSO and 25 µl transferred into a 96 well plate. Platelets were loaded with the mitochondrial membrane potential dye DiOC6(3) (3,3'-dihexyloxacarbocyanine Iodide; Invitrogen) at 200 nM for 30 min, then 50 µl plated into each well of the 96 well plate containing diluted test compound and incubated for 15 min. Control wells included DMSO (0.1 % final concentration) only or 5 µM cyclosporin A. Platelets were then stimulation with the addition of assay buffer (25µl) containing CaCl₂, alpha thrombin and Convulxin to achieve a final concentrations of 2mM, 0.017 U/ml and 0.167 µg/ml respectively and incubated for 14 min. The reaction was stopped by the addition of 25 µl 15 mM EDTA in assay buffer. The mitochondrial membrane potential across the population of platelets in each well was then quantified by flow cytometry using a Guava easyCyte 5 Benchtop Flow Cytometer with 3000 events per well. The percentage of platelets with a depolarised mitochondrial membrane potential was calculated for each well. A pIC50 for each compound was then calculated using a standard four parameter curve fit model (GraphPad Prism).

Rat brain mitochondria assay

mPTP activity was measured in brain mitochondria freshly isolated from female Sprague Dawley (250 to 300 gram) rats. Anaesthetised rats were perfused in-situ with ~40 ml cold Dulbecco's Phosphate Buffered Saline (DPBS), then brains dissected and transferred into 30 ml Isolation Buffer (225mM mannitol, 75 mM sucrose, 1mM EGTA, adjusted to pH 7.4 with 1M NaOH). The brain was minced using tweezers and a scalpel into ~5mm pieces then transferred into a 50 ml Potterton Dounce homogenization tube on ice containing 10 ml ice-cold isolation buffer (as above with addition of Complete Protease inhibitor; 1 tablet per 50 ml buffer).

Homogenisation was carried out using a 170eflon pestle at 1800 rpm. The slurry was centrifuged at 2000 g for 10 min at 4 °C, then the supernatant centrifuged at 12,000 g for 9 min. The pellet was resuspended with a dounce homogeniser in isolation buffer as above but with the addition of 0.02 % digitonin, centrifuged at 12,000g for 11 min and finally resuspended in 5 ml modified isolation buffer (as above but with EGTA reduced to 0.1 mM).

Test compounds were prepared in 384 well polypropylene assay plates as described above for the liver mitochondria assay. A stock mitochondria/Fluo5N assay solution was prepared in 5.6 ml assay buffer (120 mM mannitol, 40 mM MOPS, 5 mM KH_2PO_4 , 60 mM KCl, 10 mM pyruvate, 2 mM malate, 2 mM MgCl_2 , 20 μM ADP, 1.26 μM oligomycin A, adjusted to pH 7.4) supplemented with Fluo5N pentapotassium salt (2 μM) and 1 ml mitochondria suspension, then transferred (15 μl) into the assay plate containing test compounds and incubated for 10 min at RT. Assay plates were then transferred to a FLIPR Tetra plate reader (Molecular Devices). Dye fluorescence was then measured every 3 sec for a total of 10 min. After 12 sec, a 2.5 μl bolus of Ca^{2+} (75 μM) was added from a source plate containing 675 μM CaCl_2 in FLIPR assay buffer. IC₅₀ values for test compounds were calculated using the fluorescence value collected at the 10 min timepoint with % inhibition calculated using the DMSO control and cyclosporin A values at 100 % and 0 % respectively.

General cytotoxicity was assessed using standard cell viability methods (Cell Titre Glo; Promega) in HEK293 and SHSY5Y cells, following incubation of test compound for between 24 and 96 hours.

Results: mPTP pIC₅₀ values for certain Example compounds of the invention in a range of mPTP assays are provided in Table 3 below. Table 3 also provides the pIC₅₀ values for Comparative Example 1. The results indicate that the tested compounds of the invention display inhibition of mPTP, with many Example compounds displaying pIC₅₀ values of 6.0 or greater. Examples 37 and 51 showed the highest activity in the rat liver mitochondria assay and Example 51 also showed the highest activity in the rat brain mitochondria assay. Table 3 also presents mPTP human platelet pIC₅₀ values for certain Example compounds and Comparative Example 1. Table 3 also presents mPTP rat brain mitochondria pIC₅₀ values for certain Example compounds and Comparative Example 1. These results indicate that the tested Example compounds are active against isolated rat liver mitochondria, isolated rat brain mitochondria and human platelet mitochondria.

Biological Example 2 – Cytochrome P450 Assays

Studies to assess tested compound mediated inhibition of cytochrome P450 enzyme isoform CYP2D6 were performed using human liver microsomes (BD Gentest) using either a single concentration (1 μ M) of test compound or concentration response (0.1, 0.3, 1, 3, 10 and 30 μ M) to derive an IC₅₀. Tested compound solutions were prepared from 10 mM stocks in DMSO and diluted to 200 μ M in DMSO. Reactions were prepared in a 96 deep well plate by combining 1 μ l test compound with 179 μ l reaction mixture (100 mM phosphate buffered saline (PBS), 0.2 mg/mL microsomes and 2 μ M Dextromethorphan prepared from stocks as detailed below).

Table 2: Summary of incubation mixtures

Buffer	Stock Concentration	Volume	Final Concentration
Microsomes	20 mg/mL	2 μ L	0.2 mg/mL
Phosphate buffer	100 mM	176 μ L	100 mM
Substrate	-	1 μ L	-

The positive control inhibitor, quinidine, was used at a final concentration of 0.5 μ M when used at a single concentration. The final concentrations of quinidine used to derive an IC₅₀ were 0, 0.1, 0.3, 1, 3, 10 and 30 μ M. Plates were warmed at 37 °C for 15 min before starting reactions with 20 μ l 10 mM NADPH solution in PBS and incubated for 20 min at 37°C. The assay is performed in duplicate. Reactions were quenched with 200 μ l cold acetonitrile containing internal standards (200 nM labetalol, 200 nM alprazolam and 100 nM tolbutamide). The plate was centrifuged at 4000 rpm for 30 minutes, placed on ice for 20 minutes and then centrifuged at 4000 rpm for 30 minutes again to precipitate protein. 100 μ L of the supernatant was transferred to a new plate and diluted with 100 μ L pure water before being analysed using UPLC/MS/MS. The products of the transformation for dextromethorphan to dextrophan was monitored by UPLC-MS/MS. The inhibition of CYP2D6 in human liver microsomes was measured as the percentage decrease in the activity of dextrophan formation compared to non-inhibited controls (= 100% activity). The IC₅₀ value was calculated (test compound concentration which produces 50% inhibition) by using Excel Xlfit.

Results: CYP2D6 % inhibition values for certain compounds of the invention are presented in Table 3. Table 3 also presents the CYP2D6 % inhibition value for Comparative Example 1. The results indicate that the tested compounds displayed a significantly reduced inhibition of CYP2D6 than Comparative Example 1. Negative values indicate that there was no effective inhibition of CYP2D6 at 1uM concentration of the tested compound. Table 3 also presents CYP2D6 IC₅₀ values for certain Example compounds and for Comparative Example 1. The results show that Comparative Example 1 is a highly potent inhibitor of CYP2D6, and significantly more potent than the tested Example compounds. This is consistent with the potent

CYP2D6 % inhibition value displayed for Comparative Example 1. Therefore, the tested compounds of the invention are expected to display improved *in vivo* properties, such as the reduction of deleterious drug-drug interactions and reduced inhibition in the production of neurotransmitters in the central nervous system, in particular dopamine.

5

Table 3: Summary of results from Biological Examples 1 and 2

Example No.	mPTP Rat liver pIC50*	mPTP Human Platelets pIC50	mPTP Rat brain pIC50	CYP2D6 % inhibition @ 1uM**	CYP2D6 IC50 uM**
Comparative Example 1	8.0	6.9	7.6	86.5	0.074
	7.8***		7.5***		
1	7.3			-2.11	
2	6.0				
3	6.4			-1.87	
4	6.5			2.89	
5	6.7			-2.57	
6	6.8	7.0		5.08	19.02
7	6.8	7.2	6.9	2.27	>30
8	5.9			3.13	
9	7.4			-2.59	
10	6.7	7.0	6.5	3.66	>30
11	6.1			-0.63	
12	6.5			3.05	
13	6.0			3.84	
14	6.5			-0.52	
15	6.9			-0.28	
16	6.8			1.72	
17	6.4			1.91	
18	6.0			1.20	
19	6.2			-3.04	
20	6.8			-1.76	
21	6.1			0.91	
22	6.6			2.05	
23	6.6			0.7	
24	6.1			0.77	

Example No.	mPTP Rat liver pIC50*	mPTP Human Platelets pIC50	mPTP Rat brain pIC50	CYP2D6 % inhibition @ 1uM**	CYP2D6 IC50 uM**
25	6.9			2.05	13.80
26	7.2			3.49	8.80
27	6.6			14.28	
28	6.7			8.27	
29	6.9			-3.03	
30	7.4		7.6	-3.33	
31	7.7			19.89	
32	6.6			4.01	
33	6.3			-0.55	
34	6.7			4.37	
35	7.2			2.83	
36	7.2			-2.57	
37	8.1			5.04	
38	7.2			7.30	
39	7.5			3.83	
40	6.6			6.65	
41	7.1			1.20	
42	6.1			-1.97	
43	6.3				
44	7.1			8.94	
45	6.3				
46	6.3				
47	5.9			1.91	
48	5.9			6.34	
49	5.8			4.52	
50	7.3			1.30	
51	7.9		8.3		
52	6.9				
53	6.7				
54	7.6		7.7	8.2	
55	7.1				
56	5.8				

Example No.	mPTP Rat liver pIC50*	mPTP Human Platelets pIC50	mPTP Rat brain pIC50	CYP2D6 % inhibition @ 1uM**	CYP2D6 IC50 uM**
57	5.8				
58	5.5				
59	5.5				
60	5.7				
61	5.6				
62	5.6				
63	4.9				
64	4.9				
65c	7.7		7.8		
66	7.2		7.3		
67	7.2		7.3		
68	7.8		7.7		
69	7.7		7.6		
70	7.5			7.5	
71	7.0			7.4	
72	8.2		8.2	9.98	
73	8.1		7.7	34.57	
74	8.0		8.0	7.5	
91	6.2		7.2		
93	7.6		6.8		
94	7.5		7.4		
95	7.6		7.9		
96	7.6		7.8		
97	7.5		7.5		
98	6.9		7.0		
107	8.1		8.0		
108	7.8		7.6		
109	7.9		7.9		
110	7.1		6.7		
111	8.1		8.1		
112	8.2		8.2		
113	7.7		7.4		

Example No.	mPTP Rat liver pIC50*	mPTP Human Platelets pIC50	mPTP Rat brain pIC50	CYP2D6 % inhibition @ 1uM**	CYP2D6 IC50 uM**
114	6.9		6.4		
115	6.2		5.7		
116	6.5		6.2		
117	6.6		5.9		

*average value from multiple experiments (n≥2)

**average value from two experiments.

***average value from a larger number of experiments

5 Biological Example 3 - PBS and FaSSIF Solubility

Test compounds were prepared as 10 mM stocks in DMSO and 15 µl samples transferred in duplicate into 1.5 mL glass flat bottom vials (BioTech Solutions). Fasted state simulated intestinal fluid (FaSSIF) or PBS (pH 7.4) was added to each vial to final volume of 500 µl. One PTFE encapsulated stir stick (V&P Scientific) was placed in each vial before sealing with PTFE/SIL plugs (BioTech Solutions). Vials were shaken at 1100 rpm for 2 hr at 25°C. Samples were then filtered through MultiScreen Solvinert filter plates (Millipore) via vacuum filtration. Aliquots of filtrate (5 µl) plus 5 µl DMSO were diluted in 490 µl 50 % acetonitrile in water containing internal standard. The filtrate was analysed and quantified against a standard of known concentration using LC-MS/MS. Solubility values of the test compound and control compound were calculated as follows:

$$[\text{Sample}] = (\text{Area ratio}_{\text{sample}} * \text{INJ VOL STD} * \text{DF}_{\text{Sample}} * [\text{STD}]) / (\text{Area ratio STD} * \text{INJ VOL}_{\text{Sample}}).$$

Results: The solubility values for certain compounds of the invention are provided in Table 4 below. Table 4 also presents results for Comparative Example 1. The results indicate that certain compounds of the invention display higher solubility in PBS and/or FaSSIF than Comparative Example 1. Certain compounds of the invention display solubility in either PBS or FaSSIF, whilst certain compounds display higher solubility values in both PBS and FaSSIF. Therefore, certain compounds of the invention may be expected to display improved bioavailability and/or improved systemic exposure than Comparative Example 1, particularly when said compounds are dosed orally.

Biological Example 4 – Liver microsome and hepatocyte intrinsic clearance assaysHepatocyte clearance assay

In vitro clearance studies were performed in primary rat and human hepatocytes (BioIVT). Vials of cryopreserved rat or human hepatocytes were thawed in a 37°C water bath for 2 min. Cells were transferred into thawing medium (Williams' Medium E containing 30% Percoll, 1 x GlutaMAX-1, 15 mM HEPES, 5 % fetal bovine serum (FBS), 4 µg/ml insulin, 1 µM dexamethasone), centrifuged at 100 g for 10 min then resuspended in culture medium (Leibovitz's L-15 Medium) at a concentration of 0.5×10^6 viable cells/mL (number of viable cells assessed using AO/PI staining). Hepatocytes (198 µL) were transferred into wells of a 96-well non-coated plate and placed in a 37°C incubator for 10 min. Test compound solutions were prepared from 10 mM stocks in DMSO, diluted to 100 µM in 50 % (v/v in water) acetonitrile. Test compound samples (2 µl) were added to each well of hepatocytes and incubated at 37°C. Samples (25 µl) were collected at t=0, 15, 30, 60, 60 and 120 min, mixed with 6 volumes (150 µl) of acetonitrile containing internal standard (100 nM alprazolam, 200 nM caffeine and 100 nM tolbutamide), vortexed for 5 min and centrifuged for 45 min at 3220 g. An aliquot of supernatant (100 µL) was diluted with 100 µL ultra-pure water, and the mixture was used for LC/MS/MS analysis. All incubations were performed in duplicate. Peak areas were determined from extracted ion chromatograms. The slope value, k, was determined by linear regression of the natural logarithm of the remaining percentage of the parent drug vs. incubation time curve. The *in vitro* half-life (*in vitro* t_{1/2}) was determined from the slope value: *in vitro* t_{1/2} = 0.693 / k. Conversion of the *in vitro* t_{1/2} (in min) into the *in vitro* intrinsic clearance (*in vitro* CL_{int}, in µL/min/1×10⁶ cells) was done using the following equation (mean of duplicate determinations):

$$\text{in vitro CL}_{\text{int}} = kV/N$$

V = incubation volume (0.2 mL)

N = number of hepatocytes per well (0.1×10^6 cells).

Microsomal clearance assay

The microsomal stability of test compounds was evaluated using rat liver microsomes (BioIVT) with and without the cofactors nicotinamide adenine dinucleotide phosphate (NADPH) and uridine-diphosphate-glucuronic acid (UDPGA). Reactions were performed in a final volume of 250 µl pre-warmed (37°C) 100 mM phosphate buffer containing 5 mM MgCl₂, 0.025 mg/ml alamethicin, and 0.5 mg/ml rat liver microsomes. NADPH and UDPGA were included at 1 mM and 2 mM respectively, where appropriate. Reactions were started with the addition of 1 µM (final concentration) test compound. Verapamil was used as the positive control. Solutions were incubated in a water bath at 37°C and aliquots collected at 0.5, 5, 15, 30 and 60 min. Reactions

were stopped by the addition of 5 volumes of cold acetonitrile with internal standards (200 nM caffeine and 100 nM tolbutamide). Samples were centrifuged at 3220 g for 40 min. An aliquot of supernatant was diluted 1:1 In ultra-pure H₂O, then used for LC-MS/MS analysis. Peak areas were determined from extracted ion chromatograms. The slope value, k, was determined by linear regression of the natural logarithm of the remaining percentage of the parent drug vs. incubation time curve. The *in vitro* half-life (*in vitro* t_{1/2}) was determined from the slope value: *in vitro* t_{1/2} = 0.693 / k. Conversion of the *in vitro* t_{1/2} (min) into the *in vitro* intrinsic clearance (*in vitro* CL_{int}, in µL/min/mg protein) was done using the following equation (mean of duplicate determinations):

$$\text{in vitro CL}_{\text{int}} = (0.693/t_{1/2}) * (\text{Vol of incubation } (\mu\text{L}) / \text{amount of protein (mg)})$$

Results: Intrinsic clearance values for certain compounds of the invention are presented in Table 4. Table 4 also presents intrinsic clearance values for Comparative Example 1. These results indicate that certain compounds of the invention may be expected to have improved oral bioavailability and/or improved systemic exposure when compared to Comparative Example 1 i.e. they exhibited lower intrinsic clearance (CL_{int}) values in at least human or rat species. Certain compounds exhibited lower intrinsic clearance (CL_{int}) values than Comparative Example 1 in both human and rat species.

Table 4: Summary of results from Biological Examples 3 and 4

Example	Solubility PBS uM (pH 7.4)	Solubility FaSSIF uM	Rat heps CL _{int} uL/min/ 10 ⁶ cells*	Rat liver microsomes CL _{int} uL/min/mg*	Human heps CL _{int} uL/min/10 ⁶ cells*
Comparative Example 1	0.9	4	89	43	13
1	22.2			106	
3	1.75				
4	0.31				
5	6.78	66	27	17	
6	3.71	16	53.3	17	<3
7	1.4	22.5	27	6.5	7.63
8	33.64	219.97	46	13	
9	0.6				
10	4.7	74	21.1	13	3.3
13	5.3				

Example	Solubility PBS uM (pH 7.4)	Solubility FaSSIF uM	Rat heps Cl _{int} uL/min/ 10 ⁶ cells*	Rat liver microsomes Cl _{int} uL/min/mg*	Human heps Cl _{int} uL/min/10 ⁶ cells*
14	16				
15	0.03				
17	0.15				
18	1.11			23	
19	134.23	196.48	9.1	20.04	4.02
20		210.47	11.9		6.08
21	25.99			38.04	
22	0.4		26.12	9.85	16.2
23	6.13		35.74	9.8	17.4
25	3.69		36.3	52.88	21.8
26			87.55		
27		43.13			9.60
28			73.64		
29	33.48	69.83	16.1		
30	5.73	135.43	12.08		18.1
35		258.2			8.33
36	65.93		9.41		7.26
37		2.95	87.63		18.9
47	0.5			36.8	4.78
50		23.7	47.81		15.59
51			48.08		11.02
54		21.97	39.38		10.79
65c		9.2	37.9		9.4
66			36.1		25.4
67			73.3		42.1
68		61.7	34.9		12
69		180.7	73		17.9
70		18.4	70.1		24.6
71		19.94	72		8.9
72		24.4	77		11.3
73		37.5	73.5		7.7
74	1.2				

Example	Solubility PBS uM (pH 7.4)	Solubility FaSSIF uM	Rat heps Cl _{int} uL/min/ 10 ⁶ cells*	Rat liver microsomes Cl _{int} uL/min/mg*	Human heps Cl _{int} uL/min/10 ⁶ cells*
91		86			
93		151			
94		43			
95		76	5		3
96		14	13		6
97		35			
98			7		9
107					
108					
109					
110			307		45
111					
112					
113		8	3		3
114		19	76		3
115					
116					
117		86			3

* average value from two experiments.

Conclusion: The results of Biological Examples 1 and 2 demonstrate that the tested compounds of the invention are inhibitors of mPTP in a range of mPTP assays. The tested compounds of the invention also showed a reduced inhibition of CYP2D6 compared to Comparative Example 1. The results of Biological Examples 3 and 4 demonstrate that certain compounds of the invention show improved solubility and/or lower intrinsic clearance compared to Comparative Example 1 and as such are expected to display improved oral bioavailability and/or improved systemic exposure compared to Comparative Example 1.

Therefore, the compounds of the invention are believed to be useful pharmaceuticals, particularly for the treatment or prophylaxis of diseases and disorders in which inhibition of mPTP provides a therapeutic or prophylactic effect.

Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer, step, group of integers or group of steps but not to the exclusion of any other integer, step, group of integers or group of steps.

5

The application of which this description and claims forms part may be used as a basis for priority in respect of any subsequent application. The claims of such subsequent application may be directed to any feature or combination of features described herein. They may take the form of product, composition, process, or use claims and may include, by way of example and without limitation, the claims which follow.

10

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

15

REFERENCES

Yu et al., TDP-43 Triggers Mitochondrial DNA Release via mPTP to Activate cGAS/STING in ALS, Cell, Volume 183, Issue 3, 2020, pages 636-649.e18.

20

Jang et al., Proximal tubule cyclophilin D mediates kidney fibrogenesis in obstructive nephropathy, 2021, American Journal of Physiology: Renal physiology, doi: 10.1152/ajprenal.00171.2021. Epub ahead of print. PMID: 34396791.


Plyte et al., Cinnamic Anilides as New Mitochondrial Permeability Transition Pore Inhibitors Endowed with Ischemia-Reperfusion Injury Protective Effect in Vivo, J. Med Chem. 2014, 57, 5333-47

25

Chen et al., Probing Mitochondrial Permeability Transition Pore Activity in Nucleated Cells and Platelets by High-Throughput Screening Assays Suggests Involvement of Protein Phosphatase 2B in Mitochondrial Dynamics, Assay and Drug Development Technologies, 2018, 16, 445-45.

CLAIMS

1. A compound of formula (I):




5 wherein:

R_{1a} is H or methyl;

R_{1b} is H or F;

A is group (Aa), (Ab), (Ac), (Ad) or (Ae):

wherein group (Aa) is:



10

wherein:

R_2 is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkylene(aryl), C_{1-4} alkylene(OH), C_{1-4} alkylene(C_{3-6} cycloalkyl), C_{1-4} alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, C_{1-4} alkoxy, OC_{1-4} alkylene(aryl), C_{1-4} alkylene OC_{1-4} alkyl, C_{1-4} alkylene OC_{3-6} cycloalkyl, C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkyleneO(aryl), C_{3-6} alkynyl, C_{1-4} alkenylO(C_{3-6} alkynyl), C_{1-4} alkylene(CO_2H); OC_{1-4} alkylene(CO_2H), C_{1-4} alkylene OC_{1-4} alkylene(CO_2H), $N(R_{2aa})C_{1-4}$ alkylene(CO_2H), C_{1-4} alkylene($NR_{2ab}R_{2ac}$), OC_{1-4} alkylene($NR_{2ab}R_{2ac}$) or $N(R_{2aa})C_{1-4}$ alkylene($NR_{2ab}R_{2ac}$); wherein said aryl, heterocycloalkyl and cycloalkyl are optionally substituted by 1, 2 or 3 substituents each independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{2a}R_{2b}$, SO_2R_{2c} and $NHSO_2R_{2c}$;

15

20

R_{2a} is selected from H and C_{1-4} alkyl;

R_{2b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

25

R_{2c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{2aa} is independently selected from H and C_{1-4} alkyl;

each R_{2ab} is independently selected from H and C_{1-4} alkyl;

each R_{2ac} is independently selected from H and C_{1-4} alkyl; or

30

R_{2ab} and R_{2ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R_{2x} is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy;

each R_3 is independently halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy;

m is 0, 1, 2 or 3;

5 or;

R_{2x} is C_{1-4} alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, C_{1-4} alkyleneO(4-10 membered heterocycloalkyl), C_{1-4} alkylene(CO_2H), OC_{1-4} alkylene(CO_2H), C_{1-4} alkyleneOC C_{1-4} alkylene(CO_2H), $N(R_{2xaa})C_{1-4}$ alkylene(CO_2H), C_{1-4} alkylene($NR_{2xab}R_{2xac}$), OC_{1-4} alkylene($NR_{2xab}R_{2xac}$) or $N(R_{2xaa})C_{1-4}$ alkylene($NR_{2xab}R_{2xac}$);

10 wherein said heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{2xa}R_{2xb}$, SO_2R_{2xc} and $NHSO_2R_{2xc}$;

R_{2xa} is selected from H and C_{1-4} alkyl;

15 R_{2xb} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{2xc} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{2xaa} is independently selected from H and C_{1-4} alkyl;

20 each R_{2xab} is independently selected from H and C_{1-4} alkyl;

each R_{2xac} is independently selected from H and C_{1-4} alkyl; or


R_{2xab} and R_{2xac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R_2 is H, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} alkyleneOC C_{1-4} alkyl, C_{1-4} alkyleneOC C_{3-6} cycloalkyl or C_{1-4} alkyleneO(4-10 membered heterocycloalkyl);

25 each R_3 is independently halo, C_{1-4} alkyl, C_{1-4} haloalkyl or C_{1-4} alkoxy;

m is 0, 1, 2 or 3;

wherein group (Ab) is:



30 wherein:

R_4 is H, C_{1-4} alkyl or C_{1-4} alkylene(aryl); wherein said aryl is optionally substituted by 1, 2 or 3 substituents each independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{4a}R_{4b}$, SO_2R_{4c} and $NHSO_2R_{4c}$;

R_{4a} is selected from H and C₁₋₄alkyl;

R_{4b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;


R_{4c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R₅ is H or C₁₋₄alkyl;

each R₆ is independently C₁₋₄alkyl or halo;

n is 0, 1, 2 or 3;

wherein group (Ac) is:




wherein:

R₇ is C₁₋₄alkyl, C₁₋₄alkylene(OH) or C₁₋₄alkyleneOC₁₋₄alkyl;

o is 1 or 2;

wherein group (Ad) is:



wherein:

X is a bond, O or CH₂;

each R₈ is independently halo, C₁₋₄alkyl, C₁₋₄alkoxy, OC₁₋₄haloalkyl, OC₁₋₄alkylene(C₃₋₆cycloalkyl), OC₁₋₄alkylene(4-10 membered heterocycloalkyl) or OH; wherein said heterocycloalkyl and cycloalkyl are optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{8a}R_{8b}, SO₂R_{8c} and NHSO₂R_{8c};

R_{8a} is selected from H and C₁₋₄alkyl;

R_{8b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;


R_{8c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R₉ is independently halo or C₁₋₄alkyl;

p is 0, 1 or 2;

q is 0, 1, 2, 3 or 4;

wherein group (Ae) is:



wherein:

R₁₇ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, C₁₋₄alkoxy, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl), C₁₋₄alkylene(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{17aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{17ab}R_{17ac}), OC₁₋₄alkylene(NR_{17ab}R_{17ac}) or N(R_{17aa})C₁₋₄alkylene(NR_{17ab}R_{17ac}); wherein said aryl, cycloalkyl or heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{17a}R_{17b}, SO₂R_{17c}, NHSO₂R_{17c};

R_{17a} is selected from H and C₁₋₄alkyl;

R_{17b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{17c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{17aa} is independently selected from H and C₁₋₄alkyl;

each R_{17ab} is independently selected from H and C₁₋₄alkyl;

each R_{17ac} is independently selected from H and C₁₋₄alkyl; or

R_{17ab} and R_{17ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R₁₈ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₂₀ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

R₂₁ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

or;

R₁₇ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₁₈ is C₁₋₄haloalkyl, C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl), C₁₋₄alkylene(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{17aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{17ab}R_{17ac}), OC₁₋₄alkylene(NR_{17ab}R_{17ac}) or N(R_{17aa})C₁₋₄alkylene(NR_{17ab}R_{17ac});

4alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl), C₁₋₄alkylene(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{18aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{18ab}R_{18ac}), OC₁₋₄alkylene(NR_{18ab}R_{18ac}) or N(R_{18aa})C₁₋₄alkylene(NR_{18ab}R_{18ac}); wherein said aryl, cycloalkyl or heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{18a}R_{18b}, SO₂R_{18c}, NHSO₂R_{18c};

R_{18a} is selected from H and C₁₋₄alkyl;

R_{18b} is selected from H, C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{18c} is selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, aryl and 4-10 membered heterocycloalkyl;

each R_{18aa} is independently selected from H and C₁₋₄alkyl;

each R_{18ab} is independently selected from H and C₁₋₄alkyl;

each R_{18ac} is independently selected from H and C₁₋₄alkyl; or

R_{18ab} and R_{18ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

R₂₀ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

R₂₁ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

or;

R₁₇ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₁₈ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₁₋₄alkoxy, C₁₋₄alkyleneOC₁₋₄alkyl C₁₋₄alkyleneOC₃₋₆cycloalkyl or C₁₋₄alkyleneO(4-10 membered heterocycloalkyl);

R₂₀ is H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl or C₁₋₄alkoxy;

R₂₁ is C₁₋₄haloalkyl, C₁₋₄alkylene(aryl), C₁₋₄alkylene(OH), C₁₋₄alkylene(C₃₋₆cycloalkyl), C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, OC₁₋₄alkylene(aryl), C₁₋₄alkyleneOC₁₋₄alkyl, C₁₋₄alkyleneOC₃₋₆cycloalkyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkyleneO(aryl), C₃₋₆alkynyl, C₁₋₄alkenylO(C₃₋₆alkynyl), C₁₋₄alkylene(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{21aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{21ab}R_{21ac}), OC₁₋₄alkylene(NR_{21ab}R_{21ac}) or N(R_{21aa})C₁₋₄alkylene(NR_{21ab}R_{21ac}); wherein said aryl, cycloalkyl or heterocycloalkyl is optionally substituted by 1, 2 or 3 substituents each independently selected from C₁₋₄alkyl, C₃₋₆cycloalkyl, C₁₋₄alkoxy, C₁₋₄haloalkyl, halo, CN, OH, NR_{21a}R_{21b}, SO₂R_{21c}, NHSO₂R_{21c};

R_{21a} is selected from H and C₁₋₄alkyl;

R_{21b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{21c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

5 each R_{21aa} is independently selected from H and C_{1-4} alkyl;

each R_{21ab} is independently selected from H and C_{1-4} alkyl;


each R_{21ac} is independently selected from H and C_{1-4} alkyl; or

R_{21ab} and R_{21ac} together with the nitrogen atom to which they are attached form a 4-10 membered heterocycloalkyl;

10

wherein B is group (Ba), (Bb) or (Bc):

wherein group (Ba) is:



wherein:

15 Y is $C(R_{11})(R_{12})$, $N(R_{13})$, O or S;

each R_{10} is independently halo or C_{1-4} alkyl;

r is 0, 1, 2 or 3;

R_{11} is H or C_{1-4} alkyl;

R_{12} is H or C_{1-4} alkyl; or R_{11} and R_{12} together with the carbon atom to which they are attached form a C_{3-6} cycloalkyl;

20

R_{13} is H, C_{1-4} alkyl or C_{3-6} cycloalkyl;


wherein said cycloalkyl is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{13a}R_{13b}$, SO_2R_{13c} and $NHSO_2R_{13c}$;

25 R_{13a} is selected from H and C_{1-4} alkyl;

R_{13b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{13c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

30 wherein group (Bb) is:




wherein:

each R_{14} is independently halo or C_{1-4} alkyl;

s is 0, 1, 2 or 3;

wherein group (Bc) is:



wherein:

R_{15} is C_{1-4} alkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyl, C_{1-4} haloalkyl, halo or CN; wherein said cycloalkyl is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, halo, CN, OH, $NR_{15a}R_{15b}$, SO_2R_{15c} and $NHSO_2R_{15c}$;

R_{15a} is selected from H and C_{1-4} alkyl;

R_{15b} is selected from H, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{15c} is selected from C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, aryl and 4-10 membered heterocycloalkyl;

R_{16} is H, halo or C_{1-4} alkyl; and


D, E and F are each independently $C(R_{16})$; or one of D, E and F is N, and the two remaining D, E and F groups are independently $C(R_{16})$;

or a pharmaceutically acceptable salt and/or solvate thereof.

2. The pharmaceutically acceptable salt according to claim 1.

3. The compound according to claim 1.

4. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 3, wherein A is group (Aa)



5. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 4, wherein R_2 is halo, C_{1-4} haloalkyl, C_{1-4} alkylene(8-10 membered

heterocycloalkyl), C₁₋₄alkyleneO(8-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl such as 1-methyl piperazinyl, C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2aa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2ab}R_{2ac}), OC₁₋₄alkylene(NR_{2ab}R_{2ac}) or N(R_{2aa})C₁₋₄alkylene(NR_{2ab}R_{2ac}).

5

6. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 5, wherein R₂ is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl.

7. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 5, wherein R₂ is C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H).

8. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claim 5 to 7, wherein R_{2x} is H.

9. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 4, wherein R_{2x} is C₁₋₄alkylene(4-10 membered heterocycloalkyl), 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl, C₁₋₄alkyleneO(4-10 membered heterocycloalkyl), C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H), OC₁₋₄alkylene(CO₂H), C₁₋₄alkyleneOC₁₋₄alkylene(CO₂H), N(R_{2xaa})C₁₋₄alkylene(CO₂H), C₁₋₄alkylene(NR_{2xab}R_{2xac}), OC₁₋₄alkylene(NR_{2xab}R_{2xac}) or N(R_{2xaa})C₁₋₄alkylene(NR_{2xab}R_{2xac}).

10. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 9, wherein R_{2x} is 4-10 membered heterocycloalkyl, such as 1-methyl piperazinyl.

11. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 9, wherein R_{2x} is C₁₋₄alkylene(CO₂H), such as CH₂CH₂(CO₂H).


12. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 9 to 11, wherein R₂ is H.

30

13. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 4 to 12, wherein m is 2 and each R₃ is independently fluoro and methyl.

14. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 3, wherein A is group (Ae)

35



15. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 14, wherein R_{17} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl.

5

16. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 14 or 15, wherein R^{18} is C_{1-4} alkyl, such as methyl, ethyl or propyl, especially methyl.


17. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 14 to 16, wherein R_{20} is C_{1-4} alkoxy, such as OMe or OEt, especially OMe.

10

18. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 14 to 17, wherein R_{21} is H.

19. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 18, wherein B is group (Bc):

15



20. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 19, wherein R_{15} is methyl, ethyl, cyclopropyl, CF_3 , CN, OMe, chloro or fluoro e.g. methyl, ethyl, cyclopropyl, CF_3 or CN.

20

21. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 19 or 20, wherein D, E and F are $C(R_{16})$ and each R_{16} is independently H, fluoro, chloro or methyl.

25

22. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 1, which is selected from the group consisting of:

(E)-3-(3-fluoro-4-methyl-5-(3-(3-methyl-1H-indazol-6-yl)acrylamido)phenyl)propanoic acid;

(E)-3-(3-(3-(3-chloro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;

30

- (E)-3-(3-fluoro-5-(3-(3-fluoro-1H-indazol-6-yl)acrylamido)-4-methylphenyl)propanoic acid;
(E)-3-(3-(3-(3-cyano-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(3-fluoro-5-(3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamido)-4-methylphenyl)propanoic acid;
5 (E)-3-(3-(3-(3-chloro-5-fluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(3-(3-(3,5-difluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
(E)-3-(3-(3-(3-cyano-5-fluoro-1H-indazol-6-yl)acrylamido)-5-fluoro-4-methylphenyl)propanoic acid;
10 (E)-3-(4-fluoro-3-methyl-2-(3-(3-methyl-1H-indazol-6-yl)acrylamido)phenyl)propanoic acid;
(E)-3-(2-(3-(3-chloro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-3-(4-fluoro-2-(3-(3-fluoro-1H-indazol-6-yl)acrylamido)-3-methylphenyl)propanoic acid;
(E)-3-(2-(3-(3-cyano-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-3-(4-fluoro-2-(3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamido)-3-methylphenyl)propanoic
15 acid;
(E)-3-(2-(3-(3-chloro-5-fluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-3-(2-(3-(3,5-difluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic acid;
(E)-3-(2-(3-(3-cyano-5-fluoro-1H-indazol-6-yl)acrylamido)-4-fluoro-3-methylphenyl)propanoic
20 acid;
(E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)acrylamide;
(E)-3-(3-chloro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
25 (E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)acrylamide;
(E)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)-3-(3-methyl-1H-indazol-6-
30 yl)acrylamide;
(E)-3-(3-chloro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;
(E)-3-(3-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;
35 (E)-3-(3-cyano-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

(E)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)acrylamide;

(E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide;

5 (E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide; and

(E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(3-fluoro-2-methyl-6-(4-methylpiperazin-1-yl)phenyl)acrylamide,

or a pharmaceutically acceptable salt and/or solvate of any one thereof.

10

23. The compound, pharmaceutically acceptable salt and/or solvate thereof according to claim 1, which is selected from the group consisting of:

(2E)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl) prop-2-enamide ;

15 (2E)-3-(3-cyano-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

20 (2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(3-cyano-5-fluoro-1H-indazol-6-yl)-N-(6-methoxy-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(5-fluoro-3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide;

25 (2E)-3-(3-chloro-5-fluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl) prop-2-enamide;

(2E)-N-(5-fluoro-4-methylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3,5-difluoro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-3-(3-chloro-1H-indazol-6-yl)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)prop-2-enamide;

(2E)-N-(5-fluoro-2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

30 (2E)-N-[5-fluoro-4-(methoxymethyl)pyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-3-(3-methyl-1H-indazol-6-yl)-N-(2-methylpyridin-3-yl)prop-2-enamide;

(2E)-N-(2,4-dimethylpyridin-3-yl)-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

(2E)-N-[5-fluoro-4-(methoxymethyl)-2-methylpyridin-3-yl]-3-(3-methyl-1H-indazol-6-yl)prop-2-enamide;

35 or a pharmaceutically acceptable salt and/or solvate of any one thereof.

24. A pharmaceutical composition comprising a compound according to any one of claims 1 to 23 and a pharmaceutically acceptable carrier or excipient.

25. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 23, or the pharmaceutical composition according to claim 24 for use as a pharmaceutical.

26. The compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 23, or the pharmaceutical composition according to claim 24 for use in the treatment or prophylaxis of a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect.

27. Use of a compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 23 or a pharmaceutical composition according to claim 24 in the manufacture of a medicament for treating or preventing a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect.

28. A method of treating or preventing a disease or disorder in which inhibition of mPTP provides a therapeutic or prophylactic effect, which comprises administering to a subject in need thereof an effective amount of a compound, pharmaceutically acceptable salt and/or solvate thereof according to any one of claims 1 to 23 or a pharmaceutical composition according to claim 24.


29. The compound, pharmaceutically acceptable salt and/or solvate thereof for use, pharmaceutical composition for use, use or method, according to any one of claims 26 to 28, wherein the disease or disorder is selected from degenerative or neurodegenerative diseases (such as Parkinson's disease, dementia with Lewy bodies, Alzheimer's disease, amyotrophic lateral sclerosis, multiple sclerosis, frontal temporal dementia, chemotherapy induced neuropathy, Huntington's disease, spinocerebellar ataxias, progressive supranuclear palsy, hereditary spastic paraplegia, Duchenne muscular dystrophy, congenital muscular dystrophy, traumatic brain injury and Friedreich's ataxia, in particular Parkinson's disease, Alzheimer's disease and amyotrophic lateral sclerosis), disorders of the central nervous system (such as AIDS dementia complex, depressive disorders, schizophrenia and epilepsy), ischemia and reperfusion injury (such as acute myocardial infarction, stroke, kidney ischemia reperfusion injury, and organ damage during transplantation), metabolic diseases (such as hepatic steatosis, diabetes, diabetic retinopathy, cognitive decline and other diabetes associated conditions, obesity and feeding behaviours, and non-alcoholic fatty liver disease), inflammatory or

autoimmune diseases (such as acute pancreatitis, systemic lupus, organ failure in sepsis and hepatitis), diseases of aging (such as bone repair, bone weakness in aging in osteoporosis and sarcopenia), renal diseases (such as chronic kidney disease associated with APOL1 genetic variants and chronic kidney disease), mitochondrial diseases (such as Reye syndrome, Leber's hereditary optic neuropathy and associated disorders and disorders), and TDP-43 diseases or disorders, such as TDP-43 associated neurodegeneration (e.g. Amyotrophic Lateral Sclerosis, Frontotemporal dementia, Facial onset sensory and motor neuronopathy, Primary lateral sclerosis, Progressive muscular atrophy, Inclusion body myopathy associated with early-onset Paget disease of the bone and Frontotemporal lobar degeneration dementia, Perry disease, Chronic traumatic encephalopathy, Severe traumatic brain injury, Alzheimer's disease, Hippocampal sclerosis dementia, Limbic-predominant age-related TDP-43 encephalopathy, and Cerebral age-related TDP-43 with sclerosis).

30. The compound, pharmaceutically acceptable salt and/or solvate thereof for use, pharmaceutical composition for use, use or method according to any one of claims 26 to 29, wherein the compound is for administration to a human subject.

31. The compound, pharmaceutically acceptable salt and/or solvate thereof for use, pharmaceutical composition for use, use or method according to any one of claims 26 to 30, for use in combination with a further therapeutic agent.


32. A compound of formula (II):



wherein:

A and R_{1a} are as defined in any one of claims 1 to 24; or a salt thereof.


33. A compound of formula (IV):



wherein:

B and R_{1b} are as defined in any one of claims 1 to 24; or a salt thereof.

34. A compound of formula (VI):



wherein:

5 B and R_{1b} are as defined in any one of claims 1 to 24;
or a salt thereof.

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2023/050493

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D209/34 C07D231/56 C07D249/18 C07D263/54 C07D277/62
C07D403/12 C07D407/12 A61K31/352 A61K31/404 A61K31/416
A61K31/4192 A61K31/428 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP 2006 160916 A (FUJI PHOTO FILM CO LTD) 22 June 2006 (2006-06-22) paragraph 66, compound 8 ----- -/--	1-31



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

2 June 2023

Date of mailing of the international search report

07/08/2023

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Beyss-Kahana, Ellen

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2023/050493

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEN MAO XIANG ET AL: "Probing Mitochondrial Permeability Transition Pore Activity in Nucleated Cells and Platelets by High-Throughput Screening Assays Suggests Involvement of Protein Phosphatase 2B in Mitochondrial Dynamics", ASSAY AND DRUG DEVELOPMENT TECHNOLOGIES, vol. 16, no. 8, 1 December 2018 (2018-12-01), pages 445-455, XP055848095, US ISSN: 1540-658X, DOI: 10.1089/adt.2018.872 Retrieved from the Internet: URL:https://www.liebertpub.com/doi/pdf/10.1089/adt.2018.872> cited in the application figure 3 -----	1-31
A	WO 2010/049768 A1 (CONGENIA SRL [IT]; FANCELLI DANIELE [IT] ET AL.) 6 May 2010 (2010-05-06) cited in the application claims; examples -----	1-31
A	Song Yonghong ET AL: "Design and synthesis of factor Xa inhibitors and their prodrugs", Bioorganic & medicinal chemistry letters, 1 January 2003 (2003-01-01), pages 297-300, XP055848774, Oxford DOI: 10.1016/S0960-894X(02)00921-6 Retrieved from the Internet: URL:https://www.sciencedirect.com/science/article/pii/S0960894X02009216/pdf?md5=f6377183b7df5153080a68f535e170fb&pid=1-s2.0-S0960894X02009216-main.pdf [retrieved on 2021-10-07] table 3, compound 18 -----	1-31
A	WO 2019/196714 A1 (UNIV EAST CHINA SCIENCE & TECH [CN]) 17 October 2019 (2019-10-17) examples -----	1-31

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2023/050493

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims;; it is covered by claims Nos.:

1-31

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-31

A compound according to formula (I)

2. claim: 32

A compound according to formula (II)

3. claims: 33, 34

Compound according to formulae (IV), (VI)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2023/050493

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 2006160916 A	22-06-2006	NONE	
WO 2010049768 A1	06-05-2010	AU 2009309378 A1	06-05-2010
		BR PI0920506 A2	27-09-2016
		CA 2741433 A1	06-05-2010
		CL 2011000903 A1	11-11-2011
		CN 102264693 A	30-11-2011
		DK 2349982 T3	02-11-2015
		EA 201170607 A1	30-12-2011
		EP 2179984 A1	28-04-2010
		EP 2349982 A1	03-08-2011
		ES 2552051 T3	25-11-2015
		HK 1164274 A1	21-09-2012
		HU E026249 T2	28-06-2016
		IL 212250 A	30-11-2015
		JP 5670338 B2	18-02-2015
		JP 2012506852 A	22-03-2012
		JP 2015003921 A	08-01-2015
		KR 20110089852 A	09-08-2011
		NZ 592224 A	28-09-2012
		PL 2349982 T3	29-02-2016
		PT 2349982 E	23-11-2015
		SI 2349982 T1	31-12-2015
		UA 105018 C2	10-04-2014
		US 2011195977 A1	11-08-2011
		US 2013245019 A1	19-09-2013
		WO 2010049768 A1	06-05-2010
		ZA 201103047 B	28-12-2011
WO 2019196714 A1	17-10-2019	CN 110357789 A	22-10-2019
		WO 2019196714 A1	17-10-2019