



US 20030022848A1

(19) **United States**

(12) **Patent Application Publication**

Baker et al.

(10) **Pub. No.: US 2003/0022848 A1**

(43) **Pub. Date: Jan. 30, 2003**

(54) **ANTISENSE OLIGONUCLEOTIDE
MODULATION OF TUMOR NECROSIS
FACTOR-ALPHA (TNF-ALPHA)
EXPRESSION**

continuation-in-part of application No. 09/166,186,
filed on Oct. 5, 1998, now patented.

Publication Classification

(76) Inventors: **Brenda F. Baker**, Carlsbad, CA (US);
C. Frank Bennett, Carlsbad, CA (US);
Madeline M. Butler, Rancho Sante Fe,
CA (US); **William R. Shanahan JR.**,
Sugar land, TX (US)

(51) **Int. Cl.⁷** **A61K 48/00**; C07H 21/04

(52) **U.S. Cl.** **514/44**; 536/23.5

Correspondence Address:
LICATLA & TYRRELL P.C.
66 E. MAIN STREET
MARLTON, NJ 08053 (US)

(57) **ABSTRACT**

(21) Appl. No.: **09/824,322**

(22) Filed: **Apr. 2, 2001**

Related U.S. Application Data

(63) Continuation-in-part of application No. 09/313,932,
filed on May 18, 1999, now patented, which is a

Compositions and methods are provided for inhibiting the expression of human tumor necrosis factor- α (TNF- α). Antisense oligonucleotides targeted to nucleic acids encoding TNF- α are preferred. Methods of using these oligonucleotides for inhibition of TNF- α expression and for treatment of diseases, particularly inflammatory and autoimmune diseases, associated with overexpression of TNF- α are provided.

FIGURE 1A





FIGURE 1B






FIGURE 2

FIGURE 3A




FIGURE 3B




FIGURE 4A




FIGURE 4B




FIGURE 5A




FIGURE 5B




FIGURE 6A





FIGURE 6B






FIGURE 7A




FIGURE 7B

FIGURE 8A





FIGURE 8B






FIGURE 9

ANTISENSE OLIGONUCLEOTIDE MODULATION OF TUMOR NECROSIS FACTOR-ALPHA (TNF-ALPHA) EXPRESSION

INTRODUCTION

[0001] This application is a continuation-in part of allowed U.S. application Ser. No. 09/313,932, filed May 18, 1999, which is a continuation-in-part of U.S. application Ser. No. 09/166,186 filed Oct. 5, 1998 (U.S. Pat. No. 6,080,580)

FIELD OF THE INVENTION

[0002] This invention relates to compositions and methods for modulating expression of the human tumor necrosis factor- α (TNF- α) gene, which encodes a naturally present cytokine involved in regulation of immune function and implicated in infectious and inflammatory disease. This invention is also directed to methods for inhibiting TNF- α mediated immune responses; these methods can be used diagnostically or therapeutically. Furthermore, this invention is directed to treatment of conditions associated with expression of the human TNF- α gene.

BACKGROUND OF THE INVENTION

[0003] Tumor necrosis factor α (TNF- α also cachectin) is an important cytokine that plays a role in host defense. The cytokine is produced primarily in macrophages and monocytes in response to infection, invasion, injury, or inflammation. Some examples of inducers of TNF- α include bacterial endotoxins, bacteria, viruses, lipopolysaccharide (LPS) and cytokines including GM-CSF, IL-1, IL-2 and IFN- γ .

[0004] TNF- α interacts with two different receptors, TNF receptor I (TNFRI, p55) and TNFRII (p75), in order to transduce its effects, the net result of which is altered gene expression. Cellular factors induced by TNF- α include interleukin-1 (IL-1), interleukin-6 (IL-6), interleukin-8 (IL-8), interferon- γ (IFN- γ), platelet derived growth factor (PDGF) and epidermal growth factor (EGF), and endothelial cell adhesion molecules including endothelial leukocyte adhesion molecule 1 (ELAM-1), intercellular adhesion molecule-1 (ICAM-1) and vascular cell adhesion molecule-1 (VCAM-1) (Tracey, K. J., et al., *Annu. Rev. Cell Biol.*, 1993, 9, 317-343; Arvin, B., et al., *Ann. NY Acad. Sci.*, 1995, 765, 62-71).

[0005] Despite the protective effects of the cytokine, over-expression of TNF- α often results in disease states, particularly in infectious, inflammatory and autoimmune diseases. This process may involve the apoptotic pathways (Ksontini, R., et al., *J. Immunol.*, 1998, 160, 4082-4089). High levels of plasma TNF- α have been found in infectious diseases such as sepsis syndrome, bacterial meningitis, cerebral malaria, and AIDS; autoimmune diseases such as rheumatoid arthritis, inflammatory bowel disease (including Crohn's disease), sarcoidosis, multiple sclerosis, Kawasaki syndrome, graft-versus-host disease and transplant (allograft) rejection; and organ failure conditions such as adult respiratory distress syndrome, congestive heart failure, acute liver failure and myocardial infarction (Eigler, A., et al., *Immunol. Today*, 1997, 18, 487-492). Other diseases in which TNF- α is involved include asthma (Shah, A., et al., *Clinical and Experimental Allergy*, 1995, 25, 1038-1044), brain injury following ischemia (Arvin, B., et al., *Ann. NY*

Acad. Sci., 1995, 765, 62-71), non-insulin-dependent diabetes mellitus (Hotamisligil et al., *Science*, 1993, 259, 87-90), insulin-dependent diabetes mellitus (Yang et al., *J. Exp. Med.*, 1994, 180, 995-1004), hepatitis (Ksontini et al., *J. Immunol.*, 1998, 160, 4082-4089), atopic dermatitis (Sumimoto et al., *Arch. Dis. Child.*, 1992, 67, 277-279), and pancreatitis (Norman et al., *Surgery*, 1996, 120, 515-521). Further, inhibitors of TNF- α have been suggested to be useful for cancer prevention (Suganuma et al. (*Cancer Res.*, 1996, 56, 3711-3715). Elevated TNF- α expression may also play a role in obesity (Kern, *J. Nutr.*, 1997, 127, 1917S-1922S). TNF- α was found to be expressed in human adipocytes and increased expression, in general, correlated with obesity.

[0006] There are currently several approaches to inhibiting TNF- α expression. Approaches used to treat rheumatoid arthritis include a chimeric anti-TNF- α antibody, a humanized monoclonal anti-TNF- α antibody, and recombinant human soluble TNF- α receptor (Camussi, *Drugs*, 1998, 55, 613-620). Other examples are indirect TNF- α inhibitors including phosphodiesterase inhibitors (e.g., pentoxifylline) and metalloprotease inhibitors (Eigler et al., *Immunol. Today*, 1997, 18, 487-492). An additional class of direct TNF- α inhibitors is oligonucleotides, including triplex-forming oligonucleotides, ribozymes, and antisense oligonucleotides. Several publications describe the use of oligonucleotides targeting TNF- α by non-antisense mechanisms. U.S. Pat. No. 5,650,316, WO 95/33493 and Aggarwal et al. (*Cancer Research*, 1996, 56, 5156-5164) disclose triplex-forming oligonucleotides targeting TNF- α . WO 95/32628 discloses triplex-forming oligonucleotides especially those possessing one or more stretches of guanosine residues capable of forming secondary structure. WO 94/10301 discloses ribozyme compounds active against TNF- α mRNA. WO 95/23225 discloses enzymatic nucleic acid molecules active against TNF- α mRNA.

[0007] A number of publications have described the use of antisense oligonucleotides targeting nucleic acids encoding TNF- α . The TNF- α gene has four exons and three introns. WO 93/09813 discloses TNF- α antisense oligonucleotides conjugated to a radioactive moiety, including sequences targeted to the 5'-UTR, AUG start site, exon 1, and exon 4 including the stop codon of human TNF- α . EP 0 414 607 B1 discloses antisense oligonucleotides targeting the AUG start codon of human TNF- α . WO 95/00103 claims antisense oligonucleotides to human TNF- α including sequences targeted to exon 1 including the AUG start site. Hartmann et al. (*Mol. Med.*, 1996, 2, 429-438) disclose uniform phosphorothioates and mixed backbone phosphorothioate/phosphodiester oligonucleotides targeted to the AUG start site of human TNF- α . Hartmann et al. (*Antisense Nucleic Acid Drug Devel.*, 1996, 6, 291-299) disclose antisense phosphorothioate oligonucleotides targeted to the AUG start site, the exon 1/intron 1 junction, and exon 4 of human TNF- α . d'Hellencourt et al. (*Biochim. Biophys. Acta*, 1996, 1317, 168-174) designed and tested a series of unmodified oligonucleotides targeted to the 5'-UTR, and exon 1, including the AUG start site, of human TNF- α . Additionally, one oligonucleotide each was targeted to exon 4 and the 3'-UTR of human TNF- α and one oligonucleotide was targeted to the AUG start site of mouse TNF- α . Rojanasakul et al. (*J. Biol. Chem.*, 1997, 272, 3910-3914) disclose an antisense phosphorothioate oligonucleotide targeted to the AUG start site of mouse TNF- α . Taylor et al. (*J. Biol. Chem.*, 1996, 271,

17445-17452 and *Antisense Nucleic Acid Drug Devel.*, 1998, 8, 199-205) disclose morpholino, methyl-morpholino, phosphodiester and phosphorothioate oligonucleotides targeted to the 5'-UTR and AUG start codon of mouse TNF- α . Tu et al. (*J. Biol. Chem.*, 1998, 273, 25125-25131) designed and tested 42 phosphorothioate oligonucleotides targeting sequences throughout the rat TNF- α gene.

[0008] Interestingly, some phosphorothioate oligodeoxynucleotides have been found to enhance lipopolysaccharide-stimulated TNF- α synthesis up to four fold due to nonspecific immunostimulatory effects (Hartmann et al. *Mol. Med.*, 1996, 2, 429-438).

[0009] Accordingly, there remains an unmet need for therapeutic compositions and methods for inhibiting expression of TNF- α , and disease processes associated therewith.

SUMMARY OF THE INVENTION

[0010] The present invention provides oligonucleotides which are targeted to nucleic acids encoding TNF- α and are capable of modulating TNF- α expression. The present invention also provides chimeric oligonucleotides targeted to nucleic acids encoding human TNF- α . The oligonucleotides of the invention are believed to be useful both diagnostically and therapeutically, and are believed to be particularly useful in the methods of the present invention.

[0011] The present invention also comprises methods of modulating the expression of human TNF- α in cells and tissues using the oligonucleotides of the invention. Methods of inhibiting TNF- α expression are provided; these methods are believed to be useful both therapeutically and diagnostically. These methods are also useful as tools, for example, for detecting and determining the role of TNF- α in various cell functions and physiological processes and conditions and for diagnosing conditions associated with expression of TNF- α .

[0012] The present invention also comprises methods for diagnosing and treating infectious and inflammatory diseases, particularly diabetes, rheumatoid arthritis, Crohn's disease, pancreatitis, multiple sclerosis, atopic dermatitis and hepatitis using the oligonucleotides of the present invention. These methods are believed to be useful, for example, in diagnosing TNF- α -associated disease progression. These methods are believed to be useful both therapeutically, including prophylactically, and as clinical research and diagnostic tools.

[0013] One embodiment of the present invention is a method of treating an inflammatory disorder in an individual comprising administering to said individual an effective amount of an oligonucleotide up to 30 nucleotides in length complementary to a nucleic acid molecule encoding human tumor necrosis factor- α , wherein the oligonucleotide inhibits the expression of said human tumor necrosis factor- α and comprises at least an 8 nucleobase portion of SEQ ID NO: 24, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 34, SEQ ID NO: 39, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 149, SEQ ID NO: 157, SEQ ID NO: 264, SEQ ID NO: 271, SEQ ID NO: 272, SEQ ID NO: 290, SEQ ID NO: 297, SEQ ID NO: 299, SEQ ID NO: 315, SEQ ID NO: 334, SEQ ID NO: 418, SEQ ID NO: 423, SEQ ID NO: 425, SEQ ID NO:

427, SEQ ID NO: 431, SEQ ID NO: 432, SEQ ID NO: 435, SEQ ID NO: 437, SEQ ID NO: 438, SEQ ID NO: 439, SEQ ID NO: 441, SEQ ID NO: 455, SEQ ID NO: 457, SEQ ID NO: 458, SEQ ID NO: 460, SEQ ID NO: 463, SEQ ID NO: 465, SEQ ID NO: 466, SEQ ID NO: 468, SEQ ID NO: 472, SEQ ID NO: 474, SEQ ID NO: 475, SEQ ID NO: 483, SEQ ID NO: 485, SEQ ID NO: 494 or SEQ ID NO: 496. Preferably, the antisense oligonucleotide is administered orally. In one aspect of this preferred embodiment, the inflammatory disorder is inflammatory bowel disease, Crohn's disease, colitis or rheumatoid arthritis. Preferably, the oligonucleotide comprises at least one modified intersugar linkage. Preferably, the modified intersugar linkage is a phosphorothioate or methylene(methylimino) intersugar linkage. In another aspect of this preferred embodiment, the oligonucleotide comprises at least one 2'-O-methoxyethyl modification. Preferably, the oligonucleotide comprises at least one 5-methyl cytidine. In one aspect of this preferred embodiment, every cytidine residue is a 5-methyl cytidine.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] FIGS. 1A-B are graphs showing collagen-induced arthritis (CIA) onset as determined by percent incidence in mice. Incidence=number of mice with at least one affected paw/total number of mice per group. FIG. 1A shows the effect of low dose range of ISIS 25302 anti-TNF- α antisense oligonucleotide in comparison to treatment by an anti-TNF- α mAb. FIG. 1B shows the effect of high dose range treatment by ISIS 25302 in comparison to treatment by an 8 mismatch control oligonucleotide (ISIS 30782).

[0015] FIG. 2 is a graph showing "total" histological scores for colon tissue from IL-10^{-/-} mice treated with saline (vehicle), ISIS 25302 or 8 MM Con. As recorded in Table 27. Results are expressed as mean $\pm$ standard deviation (n=6). The asterisk indicates a significant difference (p<0.05) in comparison to the vehicle group.

[0016] FIGS. 3A-B show the basal (FIG. 3A) and LPS-induced (FIG. 3B) levels of TNF- α secretion from colon tissue of IL-10^{-/-} mice post-treatment with ISIS 25302 and the 8 base mismatch control oligonucleotide 30782 (8 MM). Doses of oligonucleotide are shown in parentheses (mg/kg). Secretion levels (pg/gm-tissue) are shown in the y-axis. The mean values $\pm$ standard deviation (n=7 to 9) are shown.

[0017] FIGS. 4A-B show the basal (FIG. 4A) and LPS-induced (FIG. 4B) levels of IFN- γ secretion from colon tissue of IL-10^{-/-} mice post-treatment with ISIS 25302 and the 8 base mismatch control oligonucleotide 30782 (8 MM). Doses of oligonucleotide are shown in parentheses (mg/kg). Secretion levels (pg/gm-tissue) are shown in the y-axis. The mean values $\pm$ standard deviation (n=6 to 9) are shown.

[0018] FIGS. 5A-B show the efficacy of ISIS 25302 versus anti-mouse TNF- α mAb in the acute model of DSS-induced colitis. FIG. 5A shows the disease activity index (DAI). FIG. 5B shows the effect of different treatments on colon length. Results are expressed as the mean $\pm$ S.E.M., where n=7. Asterisks show a significant difference from saline treated (*) or normal (*) group (p<0.05).

[0019] FIGS. 6A-B show that the prevention of acute colitis by ISIS 25302 in the DSS-induced colitis molecule is sequence-dependent. FIG. 5A shows DAI versus treatment. FIG. 5B shows the effect of different treatments on colon

length. Asterisks indicate significant differences from saline (*) or 1.0 mg/kg 8 MM Con (*) treated group ($p < 0.05$).

[0020] FIGS. 6A-B are graphs showing the efficacy of ISIS 25302 in the DSS-induced mouse model of chronic colitis based on DAI. FIG. 6A shows the mean DAI of each group over the course of the two cycle DSS-induced chronic colitis study. FIG. 6B shows the mean DAI at representative cycle times. The doses are indicated in parentheses (mg/kg). Results are expressed as the mean S.E.M., where $n = 8$ to 10. Asterisks indicate statistical significance in comparison to the Vehicle group ($P < 0.05$).

[0021] FIGS. 8A-B show histopathology of colon tissue from mice administered DSS in the two cycle chronic colitis model. Results are expressed as mean S.E.M. FIG. 8A shows the total inflammation and crypt scores. Acute inflammatory infiltrates consist of granulocytes, lymphocytes and plasma cells. Chronic inflammatory infiltrates consist of granulocytes, lymphocytes, plasma cells, monocytes and macrophages. FIG. 8B shows histological scores of different regions of the colon. PA=proximal acute inflammation score, DA=distal acute inflammation score, PC=proximal chronic inflammation score, DC=distal chronic inflammation score, PCS=proximal crypt score and DCS=distal crypt score. Asterisks indicate statistical significance in comparison to the Vehicle group ($p < 0.05$).

[0022] FIG. 9 shows TNF- α mRNA levels from longitudinal sections of colon tissue derived from each mouse at time of sacrifice in the chronic colitis model (mean S.E.M.). Group A=0.25 mg/kg ISIS 25302, group B=Vehicle, group C=anti-TNF mAb, group D=no treatment, group E=2.5 mg/kg ISIS 25302, group F=12.5 mg/kg ISIS 25302.

DETAILED DESCRIPTION OF THE INVENTION

[0023] TNF- α plays an important regulatory role in the immune response to various foreign agents. Overexpression of TNF- α results in a number of infectious and inflammatory diseases. As such, this cytokine represents an attractive target for treatment of such diseases. In particular, modulation of the expression of TNF- α may be useful for the treatment of diseases such as Crohn's disease, diabetes mellitus, multiple sclerosis, rheumatoid arthritis, hepatitis, pancreatitis and asthma.

[0024] The present invention employs antisense compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding TNF- α , ultimately modulating the amount of TNF- α produced. This is accomplished by providing oligonucleotides which specifically hybridize with nucleic acids, preferably mRNA, encoding TNF- α .

[0025] This relationship between an antisense compound such as an oligonucleotide and its complementary nucleic acid target, to which it hybridizes, is commonly referred to as "antisense". "Targeting" an oligonucleotide to a chosen nucleic acid target, in the context of this invention, is a multistep process. The process usually begins with identifying a nucleic acid sequence whose function is to be modulated. This may be, as examples, a cellular gene (or mRNA made from the gene) whose expression is associated with a particular disease state, or a foreign nucleic acid from an infectious agent. In the present invention, the targets are

nucleic acids encoding TNF- α ; in other words, a gene encoding TNF- α , or mRNA expressed from the TNF- α gene. mRNA which encodes TNF- α is presently the preferred target. The targeting process also includes determination of a site or sites within the nucleic acid sequence for the antisense interaction to occur such that modulation of gene expression will result.

[0026] In accordance with this invention, persons of ordinary skill in the art will understand that messenger RNA includes not only the information to encode a protein using the three letter genetic code, but also associated ribonucleotides which form a region known to such persons as the 5'-untranslated region, the 3'-untranslated region, the 5' cap region and intron/exon junction ribonucleotides. Thus, oligonucleotides may be formulated in accordance with this invention which are targeted wholly or in part to these associated ribonucleotides as well as to the informational ribonucleotides. The oligonucleotide may therefore be specifically hybridizable with a transcription initiation site region, a translation initiation codon region, a 5' cap region, an intron/exon junction, coding sequences, a translation termination codon region or sequences in the 5'- or 3'-untranslated region. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon." A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formyl-methionine (prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding TNF- α , regardless of the sequence(s) of such codons. It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region," "AUG region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. This region is a preferred target region. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon. This region is a preferred target region. The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other preferred target regions include the 5'untranslated region (5'UTR), known in the art to refer to

the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

[0027] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a pre-mRNA transcript to yield one or more mature mRNAs. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., exon-exon or intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. Targeting particular exons in alternatively spliced mRNAs may also be preferred. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

[0028] Once the target site or sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired modulation.

[0029] "Hybridization", in the context of this invention, means hydrogen bonding, also known as Watson-Crick base pairing, between complementary bases, usually on opposite nucleic acid strands or two regions of a nucleic acid strand. Guanine and cytosine are examples of complementary bases which are known to form three hydrogen bonds between them. Adenine and thymine are examples of complementary bases which form two hydrogen bonds between them.

[0030] "Specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity such that stable and specific binding occurs between the DNA or RNA target and the oligonucleotide.

[0031] It is understood that an oligonucleotide need not be 100% complementary to its target nucleic acid sequence to be specifically hybridizable. An oligonucleotide is specifically hybridizable when binding of the oligonucleotide to the target interferes with the normal function of the target molecule to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the oligonucleotide to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment or, in the case of in vitro assays, under conditions in which the assays are conducted.

[0032] Hybridization of antisense oligonucleotides with mRNA interferes with one or more of the normal functions

of mRNA. The functions of mRNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in by the RNA. Binding of specific protein(s) to the RNA may also be interfered with by antisense oligonucleotide hybridization to the RNA.

[0033] The overall effect of interference with mRNA function is modulation of expression of TNF- α . In the context of this invention "modulation" means either inhibition or stimulation; i.e., either a decrease or increase in expression. This modulation can be measured in ways which are routine in the art, for example by Northern blot assay of mRNA expression, or reverse transcriptase PER, as taught in the examples of the instant application or by Western blot or ELISA assay of protein expression, or by an immunoprecipitation assay of protein expression. Effects of antisense oligonucleotides of the present invention on TNF- α expression can also be determined as taught in the examples of the instant application. Inhibition is presently a preferred form of modulation.

[0034] The oligonucleotides of this invention can be used in diagnostics, therapeutics, prophylaxis, and as research reagents and in kits. Since the oligonucleotides of this invention hybridize to nucleic acids encoding TNF- α , sandwich, calorimetric and other assays can easily be constructed to exploit this fact. Provision of means for detecting hybridization of oligonucleotides with the TNF- α gene or mRNA can routinely be accomplished. Such provision may include enzyme conjugation, radiolabelling or any other suitable detection systems. Kits for detecting the presence or absence of TNF- α may also be prepared.

[0035] The present invention is also suitable for diagnosing abnormal inflammatory states in tissue or other samples from patients suspected of having an inflammatory disease such as rheumatoid arthritis. The ability of the oligonucleotides of the present invention to inhibit inflammatory processes may be employed to diagnose such states. A number of assays may be formulated employing the present invention, which assays will commonly comprise contacting a tissue sample with an oligonucleotide of the invention under conditions selected to permit detection and, usually, quantitation of such inhibition. In the context of this invention, to "contact" tissues or cells with an oligonucleotide or oligonucleotides means to add the oligonucleotide(s), usually in a liquid carrier, to a cell suspension or tissue sample, either in vitro or ex vivo, or to administer the oligonucleotide(s) to cells or tissues within an animal.

[0036] The oligonucleotides of this invention may also be used for research purposes. Thus, the specific hybridization exhibited by the oligonucleotides may be used for assays, purifications, cellular product preparations and in other methodologies which may be appreciated by persons of ordinary skill in the art.

[0037] In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid or deoxyribonucleic acid. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent intersugar (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oli-

gonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced binding to target and increased stability in the presence of nucleases.

[0038] The antisense compounds in accordance with this invention preferably comprise from about 5 to about 50 nucleobases. Particularly preferred are antisense oligonucleotides comprising from about 8 to about 30 nucleobases (i.e., from about 8 to about 30 linked nucleosides). As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidine. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

[0039] Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

[0040] Preferred modified oligonucleotide backbones include, for example, phosphorothioates, choral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and choral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and borano-phosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

[0041] Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050.

[0042] Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones

that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

[0043] Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439.

[0044] In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262. Further teaching of PNA compounds can be found in Nielsen et al. (*Science*, 1991, 254, 1497-1500).

[0045] Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular —CH₂—NH—O—CH₂—, —CH₂—N(CH₃)—O—CH₂— [known as a methylene (methylimino) or MMI backbone], —CH₂—O—N(CH₃)—CH₂—, —CH₂—N(CH₃)—N(CH₃)—CH₂— and —O—N(CH₃)—CH₂—CH₂— [wherein the native phosphodiester backbone is represented as —O—P—O—CH₂—] of the above referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above referenced U.S. Pat. No. 5,602,240. Also referred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

[0046] Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O—, S—, or N-alkyl, O-alkyl-O-alkyl, O—, S—, or N-alkenyl, or O—, S— or N-alkynyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)₂ON(CH₃)₂, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to

about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, poly-alkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O—CH₂CH₂OCH₃, also known as 2'-O—(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta* 1995, 78, 486-504) i.e., an alkoxyalkoxy group.

[0047] Other preferred modifications include 2'-methoxy (2'-O—CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugars structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920.

[0048] Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C or m5c), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in the *Concise Encyclopedia Of Polymer Science And Engineering* 1990, pages 858-859, Kroschwitz, J. I., ed. John Wiley & Sons, those disclosed by Englisch et al. (*Angewandte Chemie, International Edition* 1991, 30, 613-722), and those disclosed by Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., eds., *Antisense Research and Applications* 1993, CRC Press, Boca Raton, pages 289-302. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidine, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-Methylcytosine substitutions have been shown to increase nucleic

acid duplex stability by 0.6-1.2° C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., eds., *Antisense Research and Applications* 1993, CRC Press, Boca Raton, pages 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0049] Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. No. 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; and 5,681,941.

[0050] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA* 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.* 1994, 4, 1053-1059), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.* 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.* 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.* 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.* 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.* 1990, 259, 327-330; Svinarchuk et al., *Biochimie* 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.* 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.* 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides* 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.* 1995, 36, 3651-3654), a palmitoyl moiety (Mishra et al., *Biochim. Biophys. Acta* 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxcholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.* 1996, 277, 923-937).

[0051] Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S. Pat. No. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

[0052] The present invention also includes oligonucleotides which are chimeric oligonucleotides. "Chimeric" oligonucleotides or "chimeras," in the context of this invention, are oligonucleotides which contain two or more chemically distinct regions, each made up of at least one nucleotide. These oligonucleotides typically contain at least one

region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of antisense inhibition of gene expression. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art. This RNase H-mediated cleavage of the RNA target is distinct from the use of ribozymes to cleave nucleic acids. Ribozymes are not comprehended by the present invention.

[0053] Examples of chimeric oligonucleotides include but are not limited to "gapmers," in which three distinct regions are present, normally with a central region flanked by two regions which are chemically equivalent to each other but distinct from the gap. A preferred example of a gapmer is an oligonucleotide in which a central portion (the "gap") of the oligonucleotide serves as a substrate for RNase H and is preferably composed of 2'-deoxynucleotides, while the flanking portions (the 5' and 3' "wings") are modified to have greater affinity for the target RNA molecule but are unable to support nuclease activity (e.g., fluoro- or 2'-O-methoxyethyl-substituted). Chimeric oligonucleotides are not limited to those with modifications on the sugar, but may also include oligonucleosides or oligonucleotides with modified backbones, e.g., with regions of phosphorothioate (P=S) and phosphodiester (P=O) backbone linkages or with regions of MMI and P=S backbone linkages. Other chimeras include "wingmers," also known in the art as "hemimers," that is, oligonucleotides with two distinct regions. In a preferred example of a wingmer, the 5' portion of the oligonucleotide serves as a substrate for RNase H and is preferably composed of 2'-deoxynucleotides, whereas the 3' portion is modified in such a fashion so as to have greater affinity for the target RNA molecule but is unable to support nuclease activity (e.g., 2'-fluoro- or 2'-O-methoxyethyl-substituted), or vice-versa. In one embodiment, the oligonucleotides of the present invention contain a 2'-O-methoxyethyl (2'-O-CH₂CH₂O₂H)₃ modification on the sugar moiety of at least one nucleotide. This modification has been shown to increase both affinity of the oligonucleotide for its target and nuclease resistance of the oligonucleotide. According to the invention, one, a plurality, or all of the nucleotide subunits of the oligonucleotides of the invention may bear a 2'-O-methoxyethyl (-O-CH₂CH₂OCH₃) modification. Oligonucleotides comprising a plurality of nucleotide subunits having a 2'-O-methoxyethyl modification can have such a modification on any of the nucleotide subunits within the oligonucleotide, and may be chimeric oligonucleotides. Aside from or in addition to 2'-O-methoxyethyl modifications, oligonucleotides containing other modifications which enhance antisense efficacy, potency or target affinity are also preferred. Chimeric oligonucleotides comprising one or more such modifications are presently preferred.

[0054] The oligonucleotides used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including

Applied Biosystems. Any other means for such synthesis may also be employed; the actual synthesis of the oligonucleotides is well within the talents of the routineer. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and 2'-alkoxy or 2'-alkoxyalkoxy derivatives, including 2'-O-methoxyethyl oligonucleotides (Martin, P., *Helv. Chim. Acta* 1995, 78, 486-504). It is also well known to use similar techniques and commercially available modified amidites and controlled-pore glass (CPG) products such as biotin, fluorescein, acridine or psoralen-modified amidites and/or CPG (available from Glen Research, Sterling, Va.) to synthesize fluorescently labeled, biotinylated or other conjugated oligonucleotides.

[0055] The antisense compounds of the present invention include bioequivalent compounds, including pharmaceutically acceptable salts and prodrugs. This is intended to encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to pharmaceutically acceptable salts of the nucleic acids of the invention and prodrugs of such nucleic acids. A pharmaceutically acceptable salts@ are physiologically and pharmaceutically acceptable salts of the nucleic acids of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.* 1977, 66, 1-19).

[0056] For oligonucleotides, examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; 8 salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

[0057] The oligonucleotides of the invention may additionally or alternatively be prepared to be delivered in a Aprodug@ form. The term Aprodug@ indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510.

[0058] For therapeutic or prophylactic treatment, oligonucleotides are administered in accordance with this invention. Oligonucleotide compounds of the invention may be formulated in a pharmaceutical composition, which may include pharmaceutically acceptable carriers, thickeners,

diluents, buffers, preservatives, surface active agents, neutral or cationic lipids, lipid complexes, liposomes, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients and the like in addition to the oligonucleotide. Such compositions and formulations are comprehended by the present invention.

[0059] Pharmaceutical compositions comprising the oligonucleotides of the present invention may include penetration enhancers in order to enhance the alimentary delivery of the oligonucleotides. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., fatty acids, bile salts, chelating agents, surfactants and non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems* 1991, 8, 91-192; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems* 1990, 7, 1-33). One or more penetration enhancers from one or more of these broad categories may be included. Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprinate, recinleate, monoolein (a.k.a. 1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, mono- and di-glycerides and physiologically acceptable salts thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems* 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems* 1990, 7, 1; El-Hariri et al., *J. Pharm. Pharmacol.* 1992 44, 651-654).

[0060] The physiological roles of bile include the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 In: *Goodman & Gilman's The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al., eds., McGraw-Hill, New York, N.Y., 1996, pages 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus, the term "bile salt" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives.

[0061] Complex formulations comprising one or more penetration enhancers may be used. For example, bile salts may be used in combination with fatty acids to make complex formulations.

[0062] Chelating agents include, but are not limited to, disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, lauroth-9 and N-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems* 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems* 1990, 7, 1-33; Buur et al., *J. Control Rel.* 1990, 14, 43-51). Chelating agents have the added advantage of also serving as DNase inhibitors.

[0063] Surfactants include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems* 1991, page 92); and perfluorochemical emulsions, such as FC-43 (Takahashi et al., *J. Pharm. Pharmacol.* 1988, 40, 252-257).

[0064] Non-surfactants include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems* 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.* 1987, 39, 621-626).

[0065] As used herein, "carrier compound" refers to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity per se) but is recognized as a nucleic acid by in vivo processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. In contrast to a carrier compound, a "pharmaceutically acceptable carrier" (excipient) is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The pharmaceutically acceptable carrier may be liquid or solid and is selected with the planned manner of administration in mind so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutically acceptable carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinyl-pyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrates (e.g., starch, sodium starch glycolate, etc.); or wetting agents (e.g., sodium lauryl sulphate, etc.). Sustained release oral delivery systems and/or enteric coatings for orally administered dosage forms are described in U.S. Pat. Nos. 4,704,295; 4,556,552; 4,309,406; and 4,309,404.

[0066] The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional compatible pharmaceutically-active materials such as, e.g., antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the invention.

[0067] Regardless of the method by which the oligonucleotides of the invention are introduced into a patient, colloidal dispersion systems may be used as delivery vehicles to enhance the in vivo stability of the oligonucleotides and/or to target the oligonucleotides to a particular organ, tissue or cell type. Colloidal dispersion systems include, but are not limited to, macromolecule complexes, nanocapsules, microspheres, beads and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, liposomes and lipid:oligonucleotide complexes of uncharacterized structure. A preferred colloidal dispersion system is a plurality of liposomes. Liposomes are microscopic spheres having an aqueous core surrounded by one or more outer layers made up of lipids arranged in a bilayer configuration (see, generally, Chonn et al., *Current Op. Biotech.* 1995, 6, 698-708).

[0068] The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic, vaginal, rectal, intranasal, epidermal, and transdermal), oral or parenteral. Parenteral administration includes intravenous drip, subcutaneous, intraperitoneal or intramuscular injection, pulmonary administration, e.g., by inhalation or insufflation, or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

[0069] Formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

[0070] Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

[0071] Compositions for parenteral administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives. In some cases it may be more effective to treat a patient with an oligonucleotide of the invention in conjunction with other traditional therapeutic modalities in order to increase the efficacy of a treatment regimen. In the context of the invention, the term "treatment regimen" is meant to encompass therapeutic, palliative and prophylactic modalities. For example, a patient may be treated with conventional chemotherapeutic agents such as those used for tumor and cancer treatment. When used with the compounds of the invention, such chemotherapeutic agents may be used individually, sequentially, or in combination with one or more other such chemotherapeutic agents.

[0072] The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC_{50} s found to be effective in vitro and in in vivo animal models. In general, dosage is from 0.01 μ g to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 μ g to 100 g per kg of body weight, once or more daily, to once every 20 years.

[0073] Thus, in the context of this invention, by "therapeutically effective amount" is meant the amount of the

compound which is required to have a therapeutic effect on the treated individual. This amount, which will be apparent to the skilled artisan, will depend upon the age and weight of the individual, the type of disease to be treated, perhaps even the gender of the individual, and other factors which are routinely taken into consideration when designing a drug treatment. A therapeutic effect is assessed in the individual by measuring the effect of the compound on the disease state in the animal.

[0074] The following examples illustrate the present invention and are not intended to limit the same.

EXAMPLES

Example 1

[0075] Synthesis of Oligonucleotides

[0076] Unmodified oligodeoxynucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine. β -cyanoethyl-diisopropyl-phosphoramidites are purchased from Applied Biosystems (Foster City, Calif.). For phosphorothioate oligonucleotides, the standard oxidation bottle was replaced by a 0.2 M solution of 3 H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise titration of the phosphite linkages. The thiation cycle wait step was increased to 68 seconds and was followed by the capping step. Cytosines may be 5-methyl cytosines. (5-methyl deoxycytidine phosphoramidites available from Glen Research, Sterling, Va. or Amersham Pharmacia Biotech, Piscataway, N.J.)

[0077] 2'-methoxy oligonucleotides are synthesized using 2'-methoxy β -cyanoethyl-diisopropyl-phosphoramidites (Chemgenes, Needham, Mass.) and the standard cycle for unmodified oligonucleotides, except the wait step after pulse delivery of tetrazole and base is increased to 360 seconds. Other 2'-alkoxy oligonucleotides are synthesized by a modification of this method, using appropriate 2'-modified amidites such as those available from Glen Research, Inc., Sterling, Va.

[0078] 2'-fluoro oligonucleotides are synthesized as described in Kawasaki et al. (*J. Med. Chem.* 1993, 36, 831-841). Briefly, the protected nucleoside N⁶-benzoyl-2'-deoxy-2'-fluoroadenosine is synthesized utilizing commercially available 9- β -D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'- α -fluoro atom is introduced by a S_N2 -displacement of a 2'- β -O-triflyl group. Thus N⁶-benzoyl-9- β -D-arabinofuranosyladenine is selectively protected in moderate yield as the 3', 5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N⁶-benzoyl groups is accomplished using standard methodologies. Standard methods are also used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

[0079] The synthesis of 2'-deoxy-2'-fluoroguanosine is accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9- β -D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutryl-arabinofuranosylguanine. Deprotection of the TPDS group is followed by protection of the hydroxyl group with THP to give diisobutryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation is followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies are used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

[0080] Synthesis of 2'-deoxy-2'-fluorouridine is accomplished by the modification of a known procedure in which 2, 2'-anhydro-1- β -D-arabinofuranosyluracil is treated with 70% hydrogen fluoride-pyridine. Standard procedures are used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

[0081] 2'-deoxy-2'-fluorocytidine is synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N⁴-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures are used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

[0082] 2'-(2-methoxyethyl)-modified amidites were synthesized according to Martin, P. (*Helv. Chim. Acta* 1995, 78, 486-506). For ease of synthesis, the last nucleotide may be a deoxynucleotide. 2'-O—CH₂CH₂OCH₃cytosines may be 5-methyl cytosines.

[0083] Synthesis of 5-Methyl Cytosine Monomers:

2,2'-Anhydro [1-(β -D-arabinofuranosyl)-5-methyluridine]

[0084] 5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenyl-carbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60EC at 1 mm Hg for 24 hours) to give a solid which was crushed to a light tan powder (57 g, 85% crude yield). The material was used as is for further reactions.

2'-O-Methoxyethyl-5-methyluridine

[0085] 2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160EC. After heating for 48 hours at 155-160EC, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH₂Cl₂/acetone/MeOH (20:5:3) containing 0.5% Et₃NH. The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

[0086] 2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxy-trityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product.

The solvent was evaporated and triturated with CH₃CN (200 mL). The residue was dissolved in CHCl₃ (1.5 L) and extracted with 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl. The organic phase was dried over Na₂SO₄, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/Hexane/Acetone (5:5:1) containing 0.5% Et₃NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

31'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-uridine

[0087] 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by tlc by first quenching the tlc sample with the addition of MeOH. Upon completion of the reaction, as judged by tlc, MeOH (50 mL) was added and the mixture evaporated at 35EC. The residue was dissolved in CHCl₃ (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl₃. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/Hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%).

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

[0088] A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH₃CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH₃CN (1 L), cooled to -5EC and stirred for 0.5 hours using an overhead stirrer. POCl₃ was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10EC, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the later solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO₃ and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

[0089] A solution of 31'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH₄OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH₃ gas was added and the vessel heated to 100EC for 2 hours (tlc showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried

over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

N⁴-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-cytidine

[0090] 2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, tlc showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl₃ (700 mL) and extracted with saturated NaHCO₃ (2×300 mL) and saturated NaCl (2×300 mL), dried over MgSO₄ and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/Hexane (1:1) containing 0.5% Et₃NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

N⁴ Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

[0091] N⁴-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH₂Cl₂ (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (tlc showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO₃ (1×300 mL) and saturated NaCl (3×300 mL). The aqueous washes were back-extracted with CH₂Cl₂ (300 mL), and the extracts were combined, dried over MgSO₄ and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/Hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

[0092] 5-methyl-2'-deoxycytidine (5-me-C) containing oligonucleotides were synthesized according to published methods (Sanghvi et al., *Nucl. Acids Res.* 1993, 21, 3197-3203) using commercially available phosphoramidites (Glen Research, Sterling Va. or ChemGenes, Needham Mass.).

[0093] Oligonucleotides having methylene (methylimino) (MMI) backbones were synthesized according to U.S. Pat. No. 5,378,825, which is coassigned to the assignee of the present invention and is incorporated herein in its entirety. For ease of synthesis, various nucleoside dimers containing MMI linkages were synthesized and incorporated into oligonucleotides. Other nitrogen-containing backbones are synthesized according to WO 92/20823 which is also coassigned to the assignee of the present invention and incorporated herein in its entirety.

[0094] Oligonucleotides having amide backbones are synthesized according to De Mesmaeker et al. (*Acc. Chem. Res.* 1995, 28, 366-374). The amide moiety is readily accessible by simple and well-known synthetic methods and is compatible with the conditions required for solid phase synthesis of oligonucleotides.

[0095] Oligonucleotides with morpholino backbones are synthesized according to U.S. Pat. No. 5,034,506 (Summerton and Weller).

[0096] Peptide-nucleic acid (PNA) oligomers are synthesized according to P. E. Nielsen et al. (*Science* 1991, 254, 1497-1500). After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concen-

trated ammonium hydroxide at 55EC for 18 hours, the oligonucleotides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically checked by ³¹P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al. (*J. Biol. Chem.* 1991, 266, 18162). Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

Example 2

[0097] Human TNF-α Oligodeoxynucleotide Sequences

[0098] Antisense oligonucleotides were designed to target human TNF-α. Target sequence data are from the TNF-α cDNA sequence published by Nedwin, G. E. et al. (*Nucleic Acids Res.* 1985, 13, 6361-6373); Genbank accession number X02910, provided herein as SEQ ID NO: 1. Oligodeoxynucleotides were synthesized primarily with phosphorothioate linkages. Oligonucleotide sequences are shown in Table 1. Oligonucleotide 14640 (SEQ ID NO: 2) is a published TNF-α antisense oligodeoxynucleotide targeted to the start site of the TNF-α gene (Hartmann, G., et al., *Antisense Nucleic Acid Drug Dev.* 1996, 6, 291-299). Oligonucleotide 2302 (SEQ ID NO: 41) is an antisense oligodeoxynucleotide targeted to the human intracellular adhesion molecule-1 (ICAM-1) and was used as an unrelated (negative) target control. Oligonucleotide 13664 (SEQ ID NO: 42) is an antisense oligodeoxynucleotide targeted to the Herpes Simplex Virus type 1 and was used as an unrelated target control.

[0099] NeoHK cells, human neonatal foreskin keratinocytes (obtained from Cascade Biologicals, Inc., Portland, Ore.) were cultured in Keratinocyte medium containing the supplied growth factors (Life Technologies, Rockville, Md.).

[0100] At assay time, the cells were between 70% and 90% confluent. The cells were incubated in the presence of Keratinocyte medium, without the supplied growth factors added, and the oligonucleotide formulated in LIPOFECTIN7 (Life Technologies), a 1:1 (w/w) liposome formulation of the cationic lipid N-[1-(2,3-dioleoyloxy)propyl]-n,n-trimethylammonium chloride (DOTMA), and dioleoyl phosphatidylethanolamine (DOPE) in membrane filtered water. For an initial screen, the oligonucleotide concentration was 300 nM in 9 μg/mL LIPOFECTIN7. Treatment was for four hours. After treatment, the medium was removed and the cells were further incubated in Keratinocyte medium containing the supplied growth factors and 100 nM phorbol 12-myristate 13-acetate (PMA, Sigma, St. Louis, Mo.). mRNA was analyzed 2 hours post-induction with PMA. Protein levels were analyzed 12 to 20 hours post-induction.

[0101] Total MRNA was isolated using the RNEASY7 Mini Kit (Qiagen, Valencia, Calif.; similar kits from other manufacturers may also be used), separated on a 1% agarose gel, transferred to HYBOND™-N+membrane (Amersham Pharmacia Biotech, Piscataway, N.J.), a positively charged nylon membrane, and probed. A TNF-α probe consisted of the 505 bp EcoRI-HindIII fragment from BBG 18 (R&D Systems, Minneapolis, Minn.), a plasmid containing human TNF-α cDNA. A glyceraldehyde 3-phosphate dehydrogenase (G3PDH) probe consisted of the 1.06 kb HindIII

fragment from pHcGAP (American Type Culture Collection, Manassas, Va.), a plasmid containing human G3PDH cDNA. The restriction fragments were purified from low-melting temperature agarose, as described in Maniatis, T., et al., *Molecular Cloning: A Laboratory Manual*, 1989 and labeled with REDIVUE™³²P-dCTP (Amersham Pharmacia

Biotech, Piscataway, N.J.) and PRIME-A-GENE7 labeling kit (Promega, Madison, Wis.). mRNA was quantitated by a PhosphorImager (Molecular Dynamics, Sunnyvale, Calif.). Secreted TNF-α protein levels were measured using a human TNF-α ELIZA kit (R&D Systems, Minneapolis, Minn. or Genzyme, Cambridge, Mass.).

TABLE 1

Nucleotide Sequences of Human TNF-α Phosphorothioate Oligodeoxynucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO: CO-ORDINATES ²	TARGET GENE NUCLEOTIDE	GENE TARGET
14640	CATGCTTTCTAGTGCAT	2	0796-0813	AUG
14641	TGAGGAGCGTCTGCTGGCT	3	0615-0634	5'-UTR
14642	GTGCTCATGGTGTCTTTCC	4	0784-0803	AUG
14643	TAATCACAAGTGCAAAACATA	5	3038-3057	3'-UTR
14644	TACCCCGGTCTCCCAAATAA	6	3101-3120	3'-UTR
14810	GTGCTCATGGTGTCTTTCC	4	0784-0803	AUG
14811	AGCACCGCCTGGAGCCCT	7	0869-0886	coding
14812	GCTGAGGAACAAGCACCGCC	8	0878-0897	coding
14813	AGGCAGAAGAGCGTGGTGGC	9	0925-0944	coding
14814	AAAGTCAGCAGGCAGAAGA	10	0935-0954	coding
14815	TTAGAGAGAGGTCCCTGG	11	1593-1610	coding
14816	TGACTGCCTGGGCCAGAG	12	1617-1634	junction
14817	GGGTCGAGAAGATGATC	13	1822-1839	junction
14818	GGGCTACAGGCTTGTCACCTC	14	1841-1860	coding
14820	CCCCTCAGCTTGAGGGTTTG	15	2171-2190	junction
14821	CCATTGGCCAGGAGGCATT	16	2218-2237	coding
14822	ACCACCAGCTGGTTATCTCT	17	2248-2267	coding
14823	CTGGGAGTAGATGAGGTACA	18	2282-2301	coding
14824	CCCTTGAAGAGGACCTGGGA	19	2296-2315	coding
14825	GGTGTGGGTGAGGAGCACAT	20	2336-2355	coding
14826	GTCTGGTAGGAGACGGCGAT	21	2365-2384	coding
14827	GCAGAGAGGAGGTTGACCTT	22	2386-2405	coding
14828	GCTTGGCCTCAGCCCCCTCT	23	2436-2455	coding
14829	CCTCCCAGATAGATGGGCTC	24	2464-2483	coding
14830	CCCTTCTCCAGCTGGAAGAC	25	2485-2504	coding
14831	ATCTCAGCGCTGAGTCGGTC	26	2506-2525	coding
14832	TCGAGATAGTCGGGCCGATT	27	2527-2546	coding
14833	AAGTAGACCTGCCCAGACTC	28	2554-2573	coding
14834	GGATGTTTCGTCCTCCTCACA	29	2588-2607	STOP
14835	ACCCTAAGCCCCCAATTCTC	30	2689-2708	3'-UTR

TABLE 1-continued

Nucleotide Sequences of Human TNF-α Phosphorothioate Oligodeoxynucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO: CO-ORDINATES ²	TARGET GENE NUCLEOTIDE	GENE TARGET
			REGION	
14836	CCACACATTCTGAATCCCA	31	2758-2777	3'-UTR
14837	AGGCCCCAGTGAGTTCTGGA	32	2825-2844	3'-UTR
14838	GTCTCCAGATTCCAGATGTC	33	2860-2879	3'-UTR
14839	CTCAAGTCCTGCAGCATTCT	34	2902-2921	3'-UTR
14840	TGGGTCCCCCAGGATACCCC	35	3115-3134	3'-UTR
14841	ACGGAAAACATGTCTGAGCC	36	3151-3170	3'-UTR
14842	CTCCGTTTTCACGGAAAACA	37	3161-3180	3'-UTR
14843	GCCTATTGTTTCAGCTCCGTT	38	3174-3193	3'-UTR
14844	GGTCACCAATCAGCATTTGT	39	3272-3292	3'-UTR
14845	GAGGCTCAGCAATGAGTGAC	40	3297-3316	3'-UTR
2302	GCCCAAGCTGGCATCCGTCA	41	target	con- trol
13664	GCCGAGGTCCATGTCGTACGC	42	target control	

¹ "C" residues are 5-methyl-cytosines except "c" residues are unmodified cytidines; all linkages are phosphorothioate linkages.
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.

[0102] Results are shown in Table 2. Oligonucleotides 14828 (SEQ ID NO. 23), 14829 (SEQ ID NO. 24), 14832 (SEQ ID NO. 27), 14833 (SEQ ID NO. 28), 14834 (SEQ ID NO. 29), 14835 (SEQ ID NO. 30), 14836 (SEQ ID NO. 31), 14839 (SEQ ID NO. 34), 14840 (SEQ ID NO. 35), and 14844 (SEQ ID NO. 39) inhibited TNF-α expression by approximately 50% or more. Oligonucleotides 14828 (SEQ ID NO. 23), 14834 (SEQ ID NO. 29), and 14840 (SEQ ID NO. 35) gave better than 70% inhibition.

TABLE 2

Inhibition of Human TNF-α mRNA Expression by Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
basal	—	—	16%	—
induced	—	—	100%	0%
13664	42	control	140%	—
14640	2	AUG	61%	39%
14641	3	5'-UTR	95%	5%
14642	4	AUG	131%	—
14810	4	AUG	111%	—
14815	11	coding	85%	15%
14816	12	junction	106%	—
14817	13	junction	97%	3%
14818	14	coding	64%	36%
14820	15	junction	111%	—
14821	16	coding	91%	9%

TABLE 2-continued

Inhibition of Human TNF-α mRNA Expression by Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
14822	17	coding	57%	43%
14827	22	coding	67%	33%
14828	23	coding	27%	73%
14829	24	coding	33%	67%
14830	25	coding	71%	29%
14831	26	coding	62%	38%
14832	27	coding	40%	60%
14833	28	coding	43%	57%
14834	29	STOP	26%	74%
14835	30	3'-UTR	32%	68%
14836	31	3'-UTR	40%	60%
14837	32	3'-UTR	106%	—
14838	33	3'-UTR	70%	30%
14839	34	5'-UTR	49%	51%
14840	35	3'-UTR	28%	72%
14841	36	3'-UTR	60%	40%
14842	37	3'-UTR	164%	—
14843	38	3'-UTR	67%	33%
14844	39	3'-UTR	46%	54%
14845	40	3'-UTR	65%	35%

Example 3

[0103] Dose Response of Antisense Phosphorothioate Oligodeoxynucleotide Effects on Human TNF- α mRNA Levels in NeoHK Cells

[0104] Four of the more active oligonucleotides from the initial screen were chosen for dose response assays. These include oligonucleotides 14828 (SEQ ID NO. 23), 14833 (SEQ ID NO. 28), 14834 (SEQ ID NO. 29) and 14839 (SEQ ID NO. 34). NeoHK cells were grown, treated and processed as described in Example 2. LIPOFECTIN7 was added at a ratio of 3 μ g/mL per 100 nM of oligonucleotide. The control included LIPOFECTIN7 at a concentration of 9 μ g/mL. The effect of the TNF- α antisense oligonucleotides was normalized to the non-specific target control. Results are shown in Table 3. Each oligonucleotide showed a dose response effect with maximal inhibition greater than 70%. Oligonucleotides 14828 (SEQ ID NO. 23) had an IC₅₀ of approximately 185 nM. Oligonucleotides 14833 (SEQ ID NO. 28) had an IC₅₀ of approximately 150 nM. Oligonucleotides 14834 (SEQ ID NO. 29) and 14839 (SEQ ID NO. 34) had an IC₅₀ of approximately 140 nM.

TABLE 3

Dose Response of NeoHK Cells to TNF- α Antisense Phosphorothioate Oligodeoxynucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
2302	41	control	25 nM	100%	—
"	"	"	50 nM	100%	—
"	"	"	100 nM	100%	—
"	"	"	200 nM	100%	—
"	"	"	300 nM	100%	—
14828	23	coding	25 nM	122%	—
"	"	"	50 nM	97%	3%
"	"	"	100 nM	96%	4%
"	"	"	200 nM	40%	60%
"	"	"	300 nM	22%	78%
14833	28	coding	25 nM	89%	11%
"	"	"	50 nM	8%	22%

TABLE 3-continued

Dose Response of NeoHK Cells to TNF- α Antisense Phosphorothioate Oligodeoxynucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
"	"	"	100 nM	64%	36%
"	"	"	200 nM	36%	64%
"	"	"	300 nM	25%	75%
14834	29	STOP	25 nM	94%	6%
"	"	"	50 nM	69%	31%
"	"	"	100 nM	65%	35%
"	"	"	200 nM	26%	74%
"	"	"	300 nM	11%	89%
14839	34	3'-UTR	25 nM	140%	—
"	"	"	50 nM	112%	—
"	"	"	100 nM	65%	35%
"	"	"	200 nM	29%	71%
"	"	"	300 nM	22%	78%

Example 4

[0105] Design and Testing of Chimeric (Deoxy Gapped) 2'-O-methoxyethyl TNF- α Antisense Oligonucleotides on TNF- α Levels in NeoHK Cells

[0106] Oligonucleotides having SEQ ID NO: 28 and SEQ ID NO: 29 were synthesized as uniformly phosphorothioate or mixed phosphorothioate/phosphodiester chimeric oligonucleotides having variable regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides. The sequences and the oligonucleotide chemistries are shown in Table 4. All 2'-MOE cytosines were 5-methyl-cytosines.

[0107] Dose response experiments, as discussed in Example 3, were performed using these chimeric oligonucleotides. The effect of the TNF- α antisense oligonucleotides was normalized to the non-specific target control. Results are shown in Table 5. The activities of the chimeric oligonucleotides tested were comparable to the parent phosphorothioate oligonucleotide.

TABLE 4

Nucleotide Sequences of TNF- α Chimeric (deoxy gapped) 2'-O-methoxyethyl Oligonucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE (5' -> 3') ¹	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION
14833	AsAsGsTsAsGsAsCsTsGsCsCsAsGsAsCsTsC	28	2554-2573	coding
16467	AoAoGoToAsGsAsCsTsGsCsCsAsGoAoCoToC	28	2554-2573	coding
16468	AsAsGsTsAsGsAsCsTsGsCsCsAsGsAsCsTsC	28	2554-2573	coding
16469	AsAsGsTsAsGsAsCsTsGsCsCsAsGsAsCsTsC	28	2554-2573	coding
16470	AsAsGsTsAsGsAsCsTsGsCsCsAsGsAsCsTsC	28	2554-2573	coding
16471	AsAsGsTsAsGsAsCsTsGsCsCsAsGsAsCsTsC	28	2554-2573	coding
14834	GsGsAsTsGsTsTsCsGsTsCsCsTsCsAsCsA	29	2588-2607	STOP
16472	GoGoAoToGsTsTsCsGsTsCsCsTsCsToCoAoCoA	29	2588-2607	STOP

TABLE 4-continued

Nucleotide Sequences of TNF-α Chimeric (deoxy gapped) 2'-O-methoxy-ethyl Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE (5' -> 3') ¹	SEQ ID NO:	TARGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET	GENE REGION
16473	GsGsAsTsGsTsTsCsGsTsCsCsTsCsCsTsCsAsCsA	29	2588-2607	STOP	
16474	GsGsAsTsGsTsTsCsGsTsCsCsTsCsCsTsCsAsCsA	29	2588-2607	STOP	
16475	GsGsAsTsGsTsTsCsGsTsCsCsTsCsCsTsCsAsCsA	29	2588-2607	STOP	
16476	GsGsAsTsGsTsTsCsGsTsCsCsTsCsCsTsCsAsCsA	29	2588-2607	STOP	

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-methoxy-ethoxy cytidines are 5-methyl-cytidines; "s" linkages are phosphorothioate linkages, "o" linkages are phosphodiester linkages.
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.

[0108]

TABLE 5

Dose Response of NeoHK Cells to TNF-α Chimeric (deoxy gapped) 2'-O-methoxyethyl Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
13664	42	Control	50 nM	100%	—
"	"	"	100 nM	100%	—
"	"	"	200 nM	100%	—
"	"	"	300 nM	100%	—
14833	28	Coding	50 nM	69%	31%
"	"	"	100 nM	64%	36%
"	"	"	200 nM	56%	44%
"	"	"	300 nM	36%	64%
16468	28	Coding	50 nM	66%	34%
"	"	"	100 nM	53%	47%
"	"	"	200 nM	34%	66%
"	"	"	300 nM	25%	75%
16471	28	Coding	50 nM	77%	23%
"	"	"	100 nM	56%	44%
"	"	"	200 nM	53%	47%
"	"	"	300 nM	31%	69%
14834	29	STOP	50 nM	74%	26%
"	"	"	100 nM	53%	47%
"	"	"	200 nM	24%	76%
"	"	"	300 nM	11%	89%
16473	29	STOP	50 nM	71%	29%
"	"	"	100 nM	51%	49%
"	"	"	200 nM	28%	72%
"	"	"	300 nM	23%	77%
16476	29	STOP	50 nM	74%	26%
"	"	"	100 nM	58%	42%
"	"	"	200 nM	32%	68%
"	"	"	300 nM	31%	69%

Example 3

[0109] Design and Testing of Chimeric Phosphorothioate/MMI TNF-α Antisense Oligodeoxynucleotides on TNF-α Levels in NeoHK Cells

[0110] Oligonucleotides having SEQ ID NO. 29 were synthesized as mixed phosphorothioate/methylene(methylimino) (MMI) chimeric oligodeoxynucleotides. The sequences and the oligonucleotide chemistries are shown in Table 6. Oligonucleotide 13393 (SEQ ID NO. 49) is an

antisense oligonucleotide targeted to the human intracellular adhesion molecule-1 (ICAM-1) and was used as an unrelated target control. All cytosines were 5-methyl-cytosines.

[0111] Dose response experiments were performed using these chimeric oligonucleotides, as discussed in Example 3 except quantitation of TNF-α mRNA levels was determined by real-time PER (RT-PER) using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, Calif.) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PER) products in real-time. As opposed to standard PER, in which amplification products are quantitated after the PER is completed, products in RT-PER are quantitated as they accumulate. This is accomplished by including in the PER reaction an oligonucleotide probe that anneals specifically between the forward and reverse PER primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE or FAM, PE-Applied Biosystems, Foster City, Calif.) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, PE-Applied Biosystems, Foster City, Calif.) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PER amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular (six-second) intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

[0112] RT-PER reagents were obtained from PE-Applied Biosystems, Foster City, Calif. RT-PER reactions were carried out by adding 25 μl PER cocktail (1×TAQMAN7 buffer A, 5.5 mM MgCl2, 300 μM each of dATP, dCTP and dGTP,

600 μ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 U RNase inhibitor, 1.25 units AMPLITAQ GOLD7, and 12.5 U MuLV reverse tran-

The IC₅₀ value was reduced from approximately 90 nM, for oligonucleotide 14834 (SEQ ID NO: 29), to approximately 45 nM for oligonucleotide 16922 (SEQ ID NO: 29).

TABLE 6

Nucleotide Sequences of Human TNF- α Chimeric Phosphorothioate/MMI Oligodeoxynucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATED ²	GENE TARGET REGION
14834	GsGsAsTsGsTsTsCsGsTsCsCsTsCsCsTsCsAsCsA	29	2588-2607	STOP
16922	GmGmAmTmGsTsTsCsGsTsCsCsTsCsCsTmCmAmCmA	29	2588-2607	STOP
16923	GmGmAmTmGmTmTsCsGsTsCsCsTsCmCmTmCmAmCmA	29	2588-2607	STOP
13393	TsCsTsGsAsGsTsAsGsCsAsGsAsGsGsAsGsCsTsC	49	target control	

¹ All cytosine residues are 5-methyl-cytosines; "s" linkages, "m" linkages are methylene(methylimino) (MMI).
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.

scriptase) to 96 well plates containing 25 μ l poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48° C. following a 10 minute incubation at 95° C. to activate the AMPLITAQ GOLD7, 40 cycles of a two-step PER protocol were carried out: 95° C. for 15 seconds (denaturation) followed by 60° C. for 1.5 minutes (annealing/extension).

[0113] For TNF- α the PER primers were:

[0114] Forward: 5'-CAGGCGGTGCTTGTTCCT-3' SEQ ID NO. 43 Reverse: 5'-GCCAGAGGGCT-GATTAGAGAGA-3' SEQ ID NO. 44 and the PER probe was: FAM-CTTCTCCTTCCTGATCGTG-GCAGGC-TAMRA (SEQ ID NO. 45) where FAM or JOE (PE-Applied Biosystems, Foster City, Calif.) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, Calif.) is the quencher dye.

[0115] For GAPDH the PER primers were:

[0116] Forward primer: 5'-GAAGGTGAAGGTGCG-GAGTC-3' SEQ ID NO. 46 Reverse primer: 5'-GAA-GATGGTGATGGGATTTTC-3' SEQ ID NO. 47 and the PER probe was: 5'JOE-CAAGCTTCCCCTTCT-CAGCC -TAMRA 3'(SEQ ID NO. 48) where FAM or JOE (PE-Applied Biosystems, Foster City, Calif.) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, Calif.) is the quencher dye.

[0117] Results are shown in Table 7. The oligonucleotide containing MMI linkages was more effective in reducing TNF- α mRNA levels than the uniformly phosphorothioate oligonucleotide. The IC₅₀ value was reduced from approximately 75 nM, for oligonucleotide 14834 (SEQ ID NO: 29), to approximately 30 nM for oligonucleotide 16922 (SEQ ID NO: 29).

[0118] Dose response experiments were also performed measuring the effect on TNF- α protein levels. Protein levels were measured as described in Example 2. Results are shown in Table 8. The oligonucleotide containing four MMI linkages on each end was more effective in reducing protein levels than the uniformly phosphorothioate oligonucleotide.

[0119]

TABLE 7

Dose Response of Chimeric Phosphorothioate/MMI TNF- α Antisense Oligodeoxynucleotides on TNF- α mRNA Levels in PMA-Induced NeoHK Cells					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
induced	—	—	—	100%	—
13393	49	control	25 nM	87.3%	12.7%
"	"	"	50 nM	98.5%	1.5%
"	"	"	100 nM	133.1%	—
"	"	"	200 nM	139.6%	—
14834	29	STOP	25 nM	98.7%	1.3%
"	"	"	50 nM	70.8%	29.2%
"	"	"	100 nM	36.0%	64.0%
"	"	"	200 nM	38.2%	61.8%
16922	29	STOP	25 nM	58.9%	41.1%
"	"	"	50 nM	28.2%	71.8%
"	"	"	100 nM	22.2%	77.8%
"	"	"	200 nM	18.9%	81.1%

[0120]

TABLE 8

Dose Response of Chimeric Phosphorothioate/MMI TNF- α Antisense Oligodeoxynucleotides on TNF- α Protein Levels in PMA-Induced NeoHK Cells					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
induced	—	—	—	100.0%	—
13393	49	control	25 nM	117.0%	—
"	"	"	50 nM	86.6%	13.4%
"	"	"	100 nM	98.7%	1.3%
"	"	"	200 nM	78.0%	22.0%
14834	29	STOP	25 nM	84.8%	15.2%
"	"	"	50 nM	76.9%	23.1%
"	"	"	100 nM	44.5%	55.5%
"	"	"	200 nM	18.7%	81.3%

TABLE 8-continued

Dose Response of Chimeric Phosphorothioate/MMI TNF- α Antisense Oligodeoxynucleotides on TNF- α Protein Levels in PMA-Induced NeoHK Cells					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
16922	29	STOP	25 nM	67.1%	32.9%
"	"	"	50 nM	48.6%	51.4%
"	"	"	100 nM	20.0%	80.0%
"	"	"	200 nM	7.9%	92.1%
16923	29	STOP	25 nM	79.9%	20.1%
"	"	"	50 nM	69.9%	30.1%
"	"	"	100 nM	56.0%	44.0%
"	"	"	200 nM	44.5%	55.5%

Example 6

[0121] Additional Human TNF- α Antisense Oligonucleotide Sequences

[0122] A second screening of human TNF- α antisense oligonucleotides was performed. Oligonucleotides were designed specifically against specific regions of the TNF- α gene. A series of oligonucleotides was designed to target introns 1 and 3, and exon 4. Sequences targeting introns 1 or 3 were synthesized as uniformly phosphorothioate oligodeoxynucleotides or mixed phosphorothioate/phosphodiester chimeric backbone oligonucleotides having variable regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides. Sequences targeting exon 4 were synthesized as mixed phosphorothioate/phosphodiester chimeric backbone oligonucleotides having variable regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides. The sequences of the chimeric oligonucleotides are shown in Table 9. Sequences of the uniformly phosphorothioate oligodeoxynucleotides are shown in Table 11. These oligonucleotides were screened at 50 nM and 200 nM for their ability to inhibit TNF- α protein secretion, essentially as

described in Example 2. Results for the chimeric backbone oligonucleotides are shown in Table 10; results for the uniformly phosphorothioate oligodeoxynucleotides are shown in Table 12.

[0123] For the chimeric backbone oligonucleotides targeting introns 1 or 3, oligonucleotide 21688 (SED ID NO. 69) gave 60% inhibition or greater. For chimeric backbone oligonucleotides targeting exon 4, two-thirds of the oligonucleotides gave nearly 60% inhibition or greater (SEQ ID NOs. 88, 90, 91, 92, 93, 94, 97, and 98). See Table 10. For the uniformly phosphorothioate oligodeoxynucleotides, five of nine oligonucleotides targeting intron 3 were effective in reducing TNF- α expression by nearly 60% or greater (SEQ ID NOs. 79, 80, 81, 82, and 84). See Table 12.

[0124] Oligonucleotides having SEQ ID NO. 91 and SEQ ID NO. 98 were synthesized as a uniformly phosphorothioate oligodeoxynucleotides or mixed phosphorothioate/phosphodiester chimeric backbone oligonucleotides having variable regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides. The sequences and the oligonucleotide chemistries are shown in Table 13. All 2'-MOE cytosines and 2'-deoxy cytosines were 5-methyl-cytosines.

[0125] Dose response experiments, as discussed in Example 3, were performed using these oligonucleotides. Included in this experiment were two oligonucleotides targeting intron 1 and two oligonucleotides targeting intron 3. Results are shown in Tables 14 and 15. The oligonucleotides targeting exon 4 with variable regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides and/or uniformly phosphorothioate or mixed phosphorothioate/phosphodiester were, in general, comparable to the parent compound.

[0126] Oligonucleotides targeting introns 1 or 3 having SEQ ID NOs 66, 69 and 80 were effective in reducing TNF- α mRNA levels by greater than 80% and showed a dose response effect with an IC₅₀, approximately 110 nM. See Tables 14 and 15.

TABLE 9

Nucleotide Sequences of TNF- α Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl Oligonucleotides			
ISIS NO.	NEUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE GENE NUCLEOTIDE TARGET CO-ORDINATES ² REGION
21669	ToGoCoGoTsCsTsCsTsCsAsTsTsTsCsCoCoCoToT	50	1019-1038 intron 1
21670	ToCoCoCoAsTsCsTsCsTsCsTsCsCsCsToCoToCoT	51	1039-1058 intron 1
21671	CoAoGoCoGsCsAsCsAsTsCsTsTsTsCsAoCoCoCoA	52	1059-1078 intron 1
21672	ToCoToCoTsCsTsCsAsTsCsCsCsTsCsCoCoToAoT	53	1079-1098 intron 1
21673	CoGoToCoTsTsTsCsTsCsCsAsTsGsTsToToToToT	54	1099-1118 intron 1
21674	CoAoCoAoTsCsTsCsTsTsTsCsTsGsCsAoToCoCoC	55	1119-1138 intron 1
21675	CoToCoToCsTsTsCsCsCsCsAsTsCsTsCoToToGoC	56	1139-1158 intron 1
21676	GoToCoToCsTsCsCsAsTsCsTsTsTsCsCoToToCoT	57	1159-1178 intron 1
21677	ToToCoCoAsTsGsTsGsCsCsAsGsAsCsAoToCoCoT	58	1179-1198 intron 1

TABLE 9-continued

Nucleotide Sequences of TNF- α Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl Oligonucleotides				
ISIS NO.	NEUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION
21678	AoToAoCoAsCsAsCsTsTsAsGsTsGsAsGoCoAoCoC	59	1199-1218	intron 1
21679	ToToCoAoTsTsCsAsTsTsCsAsTsTsCsAoCoToCoC	60	1219-1238	intron 1
21680	ToAoToAoTsCsTsGsCsTsTsGsTsTsCsAoToToCoA	61	1239-1258	intron 1
21681	CoToGoToCsTsCsCsAsTsAsTsCsTsTsAoToToToA	62	1259-1278	intron 1
21682	ToCoToCoTsTsCsTsCsAsCsAsCsCsCsCoAoCoAoT	63	1279-1298	intron 1
21683	CoAoCoToTsGsTsTsTsCsTsTsCsCsCsCoCoAoToC	64	1299-1318	intron 1
21684	CoToCoAoCsCsAsTsCsTsTsTsAsTsTsCoAoToAoT	65	1319-1338	intron 1
21685	AoToAoToTsTsCsCsGsCsTsTsCsTsTsToCoToGoT	66	1339-1358	intron 1
21686	CoAoToCoTsCsTsCsTsCsTsTsAsGsCoToGoToC	67	1359-1378	intron 1
21687	ToCoToToCsTsCsTsCsTsTsAsTsCsToCoCoCoC	68	1379-1398	intron 1
21688	GoToGoToGsCsCsAsGsAsCsAsCsCsCsToAoToCoT	69	1399-1418	intron 1
21689	ToCoToToTsCsCsCsTsGsAsGsTsGsTsCoToToCoT	70	1419-1438	intron 1
21690	AoCoCoToTsCsCsAsGsCsAsTsTsCsAsAoCoAoGoC	71	1439-1458	intron 1
21691	CoToCoCoAsTsTsCsAsTsCsTsGsTsGsToAoToToC	72	1459-1478	intron 1
21692	ToGoAoGoGsTsGsTsCsTsGsGsTsTsTsToCoToCoT	73	1479-1498	intron 1
21693	AoCoAoCoAsTsCsCsTsCsAsGsAsGsCsToCoToToA	74	1871-1890	intron 3
21694	CoToAoGoCsCsCsTsCsCsAsAsGsTsTsCoCoAoAoG	75	1891-1910	intron 3
21695	CoGoGoGoCsTsTsCsAsAsTsCsCsCsCsAoAoAoToC	76	1911-1930	intron 3
21696	AoAoGoToTsCsTsGsCsCsTsAsCsCsAsToCoAoGoC	77	1931-1950	intron 3
21697	GoToCoCoTsTsCsTsCsAsCsAsTsTsGsToCoToCoC	78	1951-1970	intron 3
21698	CoCoToToCsCsCsTsTsGsAsGsCsTsCsAoGoCoGoA	79	1971-1990	intron 3
21699	GoGoCoCoTsGsTsGsCsTsGsTsTsCsCsToCoCoAoC	80	1991-2010	intron 3
21700	CoGoToToCsTsGsAsGsTsAsTsCsCsCsAoCoToAoA	81	2011-2030	intron 3
21701	CoAoCoAoTsCsCsCsAsCsCsTsGsGsCsCoAoToGoA	82	2031-2050	intron 3
21702	GoToCoCoTsCsTsCsTsGsTsCsTsGsTsCoAoToCoC	83	2051-2070	intron 3
21703	CoCoAoCoCsCsCsAsCsAsTsCsCsGsGsToToCoCoT	84	2071-2090	intron 3
21704	ToCoCoToGsGsCsCsTsCsGsAsGsCsToCoToGoC	85	2091-2110	intron 3
21705	AoToGoToCsGsGsTsTsCsAsCsTsCsTsCoCoAoCoA	86	2111-2130	intron 3
21706	AoGoAoGoGsAsGsAsGsTsCsAsGsTsGsToGoGoCoC	87	2131-2150	intron 3
21722	GoAoToCoCsCsAsAsAsGsTsAsGsAsCsCoToGoCoC	88	2561-2580	exon 4
21723	CoAoGoAoCsTsCsGsGsCsAsAsAsGsTsCoGoAoGoA	89	2541-2560	exon 4
21724	ToAoGoToCsGsGsGsCsCsGsAsTsTsGsAoToCoToC	90	2521-2540	exon 4
21725	AoGoCoGoCsTsGsAsGsTsCsGsGsTsCsAoCoCoCoT	91	2501-2520	exon 4
21726	ToCoToCoCsAsGsCsTsGsGsAsAsGsAsCoCoCoCoT	92	2481-2500	exon 4
21727	CoCoCoAoGsAsTsAsGsAsTsGsGsGsCsToCoAoToA	93	2461-248	exon 4

TABLE 9-continued

Nucleotide Sequences of TNF-α Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl Oligonucleotides					
ISIS NO.	NEUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	
21728	C o C o A o G o G sGsCsTsTsGsGsCsCsTsCs A o G o C o C	94	2441-2460	exon 4	
21729	C o C o T o C o T sGsGsGsGsTsCsTsCsCsCs T o C o T o G o G	95	2421-2440	exon 4	
21730	C o A o G o G o G sGsCsTsCsTsTsGsAsTsGs G o C o A o G o A	96	2401-2420	exon 4	
21731	G o A o G o G o A sGsGsTsTsTsGsAsCsCsTsTs G o G o T o C o T	97	2381-2400	exon 4	
21732	G o G o T o A o G sGsAsGsAsCsGsGsCsGsAs T o G o C o G o G	98	2361-2380	exon 4	
21733	C o T o G o A o T sGsGsTsGsTsGsGsGsTsGs A o G o G o A o G	99	2341-2360	exon 4	

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; "s" linkages are phosphorothioate linkages, "o" linkages are phosphodiester linkages.
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.

[0127]

TABLE 10

Dose Response of PMA-Induced neoHK Cells to Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl TNF-α Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Express- ion	% protein Inhibit- ion
induced	—	—	—	100%	—
14834	29	STOP	50 nM	76%	24%
"	"	"	200 nM	16%	84%
21669	50	intron 1	50 nM	134%	—
"	"	"	200 nM	114%	—
21670	51	intron 1	50 nM	122%	—
"	"	"	200 nM	101%	—
21671	52	iritron 1	50 nM	90%	10%
"	"	"	200 nM	58%	42%
21672	53	intron 1	50 nM	122%	—
"	"	"	200 nM	131%	—
21673	54	intron 1	50 nM	102%	—
"	"	"	200 nM	110%	—
21674	55	intron 1	50 nM	111%	—
"	"	"	200 nM	96%	4%
21675	56	intron 1	50 nM	114%	—
"	"	"	200 nM	99%	1%
21676	57	intron 1	50 nM	107%	—
"	"	"	200 nM	96%	4%
21677	58	intron 1	50 nM	86%	14%
"	"	"	200 nM	95%	5%
21678	59	intron 1	50 nM	106%	—
"	"	"	200 nM	107%	—
21679	60	intron 1	50 nM	75%	25%
"	"	"	200 nM	73%	27%
21680	61	intron 1	50 nM	76%	24%
"	"	"	200 nM	80%	20%
21681	62	intron 1	50 nM	79%	21%
"	"	"	200 nM	82%	18%
21682	63	intron 1	50 nM	102%	—
"	"	"	200 nM	88%	12%
21683	64	intron 1	50 nM	80%	20%
"	"	"	200 nM	66%	34%
21684	65	intron 1	50 nM	91%	9%
"	"	"	200 nM	69%	31%

TABLE 10-continued

Dose Response of PMA-Induced neoHK Cells to Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl TNF-α Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Express- ion	% protein Inhibit- ion
21685	66	intron 1	50 nM	98%	2%
"	"	"	200 nM	90%	10%
21686	67	intron 1	50 nM	97%	3%
"	"	"	200 nM	72%	28%
21687	68	intron 1	50 nM	103%	—
"	"	"	200 nM	64%	36%
21688	69	intron 1	50 nM	87%	13%
"	"	"	200 nM	40%	60%
21689	70	intron 1	50 nM	78%	22%
"	"	"	200 nM	74%	26%
21690	71	intron 1	50 nM	84%	16%
"	"	"	200 nM	80%	20%
21691	72	intron 1	50 nM	86%	14%
"	"	"	200 nM	75%	25%
21692	73	intron 1	50 nM	85%	15%
"	"	"	200 nM	61%	39%
21693	74	intron 3	50 nM	81%	19%
"	"	"	200 nM	83%	17%
21694	75	intron 3	50 nM	99%	1%
"	"	"	200 nM	56%	44%
21695	76	intron 3	50 nM	87%	13%
"	"	"	200 nM	84%	16%
21696	77	intron 3	50 nM	103%	—
"	"	"	200 nM	86%	14%
21697	78	intron 3	50 nM	99%	1%
"	"	"	200 nM	52%	48%
21698	79	intron 3	50 nM	96%	4%
"	"	"	200 nM	47%	53%
21699	80	intron 3	50 nM	73%	27%
"	"	"	200 nM	84%	16%
21700	81	intron 3	50 nM	80%	20%
"	"	"	200 nM	53%	47%
21701	82	intron 3	50 nM	94%	6%
"	"	"	200 nM	56%	44%
21702	83	intron 3	50 nM	86%	14%
"	"	"	200 nM	97%	3%

TABLE 10-continued

Dose Response of PMA-Induced neoHK Cells to Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl TNF- α Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Express-ion	% protein Inhibit-ion
21703	84	intron 3	50 nM	88%	12%
"	"	"	200 nM	74%	26%
21704	85	intron 3	50 nM	69%	31%
"	"	"	200 nM	65%	35%
21705	86	intron 3	50 nM	92%	8%
"	"	"	200 nM	77%	23%
21706	87	intron 3	50 nM	95%	5%
"	"	"	200 nM	82%	18%
21722	88	exon 4	50 nM	81%	19%
"	"	"	200 nM	41%	59%
21723	89	exon 4	50 nM	87%	13%
"	"	"	200 nM	74%	26%
21724	90	exon 4	50 nM	68%	32%
"	"	"	200 nM	33%	67%
21725	91	exon 4	50 nM	55%	45%
"	"	"	200 nM	30%	70%
21726	92	exon 4	50 nM	72%	28%
"	"	"	200 nM	40%	60%
21727	93	exon 4	50 nM	67%	33%
"	"	"	200 nM	40%	60%

TABLE 10-continued

Dose Response of PMA-Induced neoHK Cells to Chimeric Backbone (deoxy gapped) 2'-O-methoxyethyl TNF- α Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Express-ion	% protein Inhibit-ion
21728	94	exon 4	50 nM	62%	38%
"	"	"	200 nM	41%	59%
21729	95	exon 4	50 nM	78%	22%
"	"	"	200 nM	53%	47%
21730	96	exon 4	50 nM	68%	32%
"	"	"	200 nM	48%	52%
21731	97	exon 4	50 nM	77%	23%
"	"	"	200 nM	41%	59%
21732	98	exon 4	50 nM	62%	38%
"	"	"	200 nM	28%	72%
21733	99	exon 4	50 nM	92%	8%
"	"	"	200 nM	74%	26%

[0128]

TABLE 11

Nucleotide Sequences of Additional Human TNF- α Phosphorothioate Oligodeoxynucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION
21804	TGCGTCTCTCATTTCCCTT	50	1019-1038	intron 1
21805	TCCCATCTCTCTCCCTCTCT	51	1039-1058	intron 1
21806	CAGCGCACATCTTTCACCCA	52	1059-1078	intron 1
21807	TCTCTCTCATCCCTCCCTAT	53	1079-1098	intron 1
21808	CGTCTTTCTCCATGTTTTTT	54	1099-1118	intron 1
21809	CACATCTCTTTCTGCATCCC	55	1119-1138	intron 1
21810	CTCTCTTCCCATCTCTTGC	56	1139-1158	intron 1
21811	GTCTCTCCATCTTTCCTTCT	57	1159-1178	intron 1
21812	TTCCATGTGCCAGACATCCT	58	1179-1198	intron 1
21813	ATACACACTTAGTGAGCACC	59	1199-1218	intron 1
21814	TTCATTCAATTCATCACTCC	60	1219-1238	intron 1
21815	TATATCTGCTTGTTCAATCA	61	1239-1258	intron 1
21816	CTGTCTCCATATCTTATTTA	62	1259-1278	intron 1
21817	TCTCTTCTCACACCCACAT	63	1279-1298	intron 1
21818	CACTTGTTTCTTCCCCCATC	64	1299-1318	intron 1
21819	CTCACCATCTTTATTCATAT	65	1319-1338	intron 1
21820	ATATTTCCGCTCTTTCTGT	66	1339-1358	intron 1

TABLE 11-continued

Nucleotide Sequences of Additional Human TNF- α Phosphorothioate Oligodeoxynucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	
21821	CATCTCTCTCCTTAGCTGTC	67	1359-1378	intron 1	
21822	TCTTCTCTCCTTATCTCCCC	68	1379-1398	intron 1	
21823	GTGTGCCAGACACCCTATCT	69	1399-1418	intron 1	
21824	TCTTTCCTGAGTGTCTTCT	70	1419-1438	intron 1	
21825	ACCTTCCAGCATTC AACAGC	71	1439-1458	intron 1	
21826	CTCCATTCACTGTGTATTC	72	1459-1478	intron 1	
21827	TGAGGTGTCTGGTTTTCTCT	73	1479-1498	intron 1	
21828	ACACATCCTCAGAGCTCTTA	74	1871-1890	intron 3	
21829	CTAGCCCTCCAAGTTCCAAG	75	1891-1910	intron 3	
21830	CGGGCTTCAATCCCCAAATC	76	1911-1930	intron 3	
21831	AAGTTCTGCCTACCATCAGC	77	1931-1950	intron 3	
21832	GTCCTTCTCACATTGTCTCC	78	1951-1970	intron 3	
21833	CCTTCCCTTGAGCTCAGCGA	79	1971-1990	intron 3	
21834	GGCCTGTGCTGTTCCCTCCAC	80	1991-2010	intron 3	
21835	CGTTCTGAGTATCCCACTAA	81	2011-2030	intron 3	
21836	CACATCCCACCTGGCCATGA	82	2031-2050	intron 3	
21837	GTCCTCTCTGTCTGTCAATCC	83	2051-2070	intron 3	
21838	CCACCCACATCCGGTTCCT	84	2071-2090	intron 3	
21839	TCCTGGCCCTCGAGCTCTGC	85	2091-2110	intron 3	
21840	ATGTCGGTTCACCTCCACA	86	2111-2130	intron 3	
21841	AGAGGAGAGTCAGTGTGGCC	87	2131-2150	intron 3	

¹ All "C" residues are 5-methyl-cytosines; all linkages are phosphorothioate linkages.
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.

[0129]

TABLE 12

Dose Response of PMA-Induced neoHK Cells to TNF- α Antisense Phosphorothioate Oligodeoxynucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
induced	—	—	—	100%	—
14834	29	STOP	50 nM	80%	20%
"	"	"	200 nM	13%	87%
21812	58	intron 1	50 nM	110%	—
"	"	"	200 nM	193%	—
21833	79	intron 3	50 nM	88%	12%
"	"	"	200 nM	8%	92%

TABLE 12-continued

Dose Response of PMA-Induced neoHK Cells to TNF- α Antisense Phosphorothioate Oligodeoxynucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
21834	80	intron 3	50 nM	70%	30%
"	"	"	200 nM	18%	82%
21835	81	intron 3	50 nM	106%	—
"	"	"	200 nM	42%	58%
21836	82	intron 3	50 nM	71%	29%
"	"	"	200 nM	12%	88%
21837	83	intron 3	50 nM	129%	—
"	"	"	200 nM	74%	26%

TABLE 12-continued

Dose Response of PMA-Induced neoHK Cells to TNF- α Antisense Phosphorothioate Oligodeoxynucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
21838	84	intron 3	50 nM	85%	15%
"	"	"	200 nM	41%	59%
21839	85	intron 3	50 nM	118%	—
"	"	"	200 nM	58%	42%
21840	86	intron 3	50 nM	120%	—
"	"	"	200 nM	96%	4%
21841	87	intron 3	50 nM	117%	—
"	"	"	200 nM	78%	22%

TABLE 14-continued

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF- α Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
21823	69	intron 1	75 nM	79.5%	20.5%
"	"	"	150 nM	31.8%	68.2%
"	"	"	300 nM	16.2%	83.8%
21725	91	exon 4	75 nM	74.8%	25.2%
"	"	"	150 nM	58.4%	41.6%
"	"	"	300 nM	45.2%	54.8%
25655	91	exon 4	75 nM	112.0%	—
"	"	"	150 nM	55.0%	45.0%
"	"	"	300 nM	39.3%	60.7%

[0130]

TABLE 13

Nucleotide Sequences of TNF- α Chimeric (deoxy gapped) 2'-O-Methoxyethyl Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TRAGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	GENE TARGET REGION
21725	AoGoCoGoCsTsGsAsGsTsCsGsGsTsCsAoCoCoCoT	91	2501-2520	exon 4	
25655	AsGsCsGsCsTsGsAsGsTsCsGsGsTsCsAsCsCsCsT	"	"	"	
25656	AsGsCsGsCsTsGsAsGsTsCsGsGsTsCsAsCsCsCsT	"	"	"	
25660	AoGoCoGsCsTsGsAsGsTsCsGsGsTsCsAsCoCoCoT	"	"	"	
21732	GoGoToAoGsGsAsGsAsCsGsGsCsGsAsToGoCoGoG	98	2361-2380	exon 4	
25657	GsGsTsAsGsGsAsGsAsCsGsGsCsGsAsTsGsCsGsG	"	"	"	
25658	GsGsTsAsGsGsAsGsAsCsGsGsCsGsAsTsGsCsGsG	"	"	"	
25661	GoGoToAsGsGsAsGsAsCsGsGsCsGsAsTsGoCoGoG	"	"	"	

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; "s" linkages are phosphorothioate linkages, "o" linkages are phosphodiester linkages.
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.

[0131]

TABLE 14

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF- α Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
induced	—	—	—	100%	—
14834	29	STOP	75 nM	91.2%	8.8%
"	"	"	150 nM	42.0%	58.0%
"	"	"	300 nM	16.9%	83.1%
21820	66	intron 1	75 nM	79.0%	21.0%
"	"	"	150 nM	34.5%	65.5%
"	"	"	300 nM	15.6%	84.4%

TABLE 14-continued

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF- α Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
25656	91	exon 4	75 nM	108.3%	—
"	"	"	150 nM	60.7%	39.3%
"	"	"	300 nM	42.8%	57.2%
25660	91	exon 4	75 nM	93.2%	6.8%
"	"	"	150 nM	72.8%	27.2%
"	"	"	300 nM	50.3%	49.7%

[0132]

TABLE 15

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF- α Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
induced	—	—	—	100%	—
14834	29	STOP	75 nM	44.9%	55.1%
"	"	"	150 nM	16.3%	83.7%
"	"	"	300 nM	2.2%	97.8%
21834	80	intron 3	75 nM	102.9%	—
"	"	"	150 nM	24.5%	75.5%
"	"	"	300 nM	19.1%	80.9%
21836	82	intron 3	75 nM	70.8%	29.2%
"	"	"	150 nM	55.9%	44.1%
"	"	"	300 nM	32.7%	67.3%
21732	98	exon 4	75 nM	42.4%	57.6%
"	"	"	150 nM	34.9%	65.1%
"	"	"	300 nM	15.4%	84.6%
25657	98	exon 4	75 nM	46.7%	53.3%
"	"	"	150 nM	72.0%	28.0%
"	"	"	300 nM	50.6%	49.4%
25658	98	exon 4	75 nM	83.7%	16.3%
"	"	"	150 nM	56.6%	43.4%
"	"	"	300 nM	36.9%	63.1%

TABLE 15-continued

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF- α Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
25661	98	exon 4	75 nM	54.9%	45.1%
"	"	"	150 nM	34.4%	65.6%
"	"	"	300 nM	8.6%	91.4%

Example 7

[0133] Activity of Fully 2'-MOE Modified TNF- α Antisense Oligonucleotides

[0134] A series of antisense oligonucleotides were synthesized targeting the terminal twenty nucleotides of each exon at every exon-intron junction of the TNF- α gene. These oligonucleotides were synthesized as fully 2'-methoxyethoxy modified oligonucleotides. The oligonucleotide sequences are shown in Table 16. Oligonucleotide 12345 (SEQ ID NO. 106) is an antisense oligonucleotide targeted to the human intracellular adhesion molecule-1 (ICAM-1) and was used as an unrelated target control.

[0135] The oligonucleotides were screened at 50 nM and 200 nM for their ability to inhibit TNF- α mRNA levels, as described in Example 3. Results are shown in Table 17. Oligonucleotide 21794 (SEQ ID NO. 102) showed an effect at both doses, with greater than 75% inhibition at 200 nM.

TABLE 16

Nucleotide Sequences of Human TNF- α Uniform 2'-MOE Oligonucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ 5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION ³
21792	AGGCACTCACCTCTTCCCTC	100	0972-0991	E1/I1
21793	CCCTGGGGAACGTGTGGGGA	101	1579-1598	I1/E2
21794	AGACACTTACTGACTGCCTG	102	1625-1644	E2/I2
21795	GAAGATGATCCTGAAGAGGA	103	1812-1831	I2/E3
21796	GAGCTCTTACCTACAACATG	104	1860-1879	E3/I3
21797	TGAGGGTTTGCTGGAGGGAG	105	2161-2180	I3/E4
12345	GATCGCGTCGGACTATGAAG	106	target control	

¹ Emboldened residues are 2'-methoxyethoxy residues, 2'-methoxyethoxy cytosine residues are 5-methyl-cytosines; all linkages are phosphorothioate linkages.
² Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO.1.
³ Each target region is an exon-intron junction and is represented in the form, for example, I1/E2, where I, followed by a number, refers to the intron number and E, followed by a number, refers to the exon number.

[0136]

TABLE 17

Dose Response of neoHK Cells to TNF- α Antisense 2'-MOE Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
induced	—	—	—	100%	—
12345	106	control	50 nM	121%	—
"	"	"	200 nM	134%	—
13393	49	control	50 nM	110%	—
"	"	"	200 nM	112%	—
14834	29	STOP	50 nM	92%	8%
"	"	"	200 nM	17%	83%
21792	100	E1/I1	50 nM	105%	—
"	"	"	200 nM	148%	—
21793	101	I1/E2	50 nM	106%	—
"	"	"	200 nM	172%	—
21794	102	E2/I2	50 nM	75%	25%
"	"	"	200 nM	23%	77%
21795	103	I2/E3	50 nM	79%	21%
"	"	"	200 nM	125%	—
21796	104	E3/I3	50 nM	56%	44%
"	"	"	200 nM	150%	—
21797	105	I3/E4	50 nM	90%	10%
"	"	"	200 nM	128%	—

Example 8

[0137] Mouse TNF- α Oligonucleotide Sequences

[0138] Antisense oligonucleotides were designed to target mouse TNF- α . Target sequence data are from the TNF- α CDNA sequence published by Semon et al. (*Nucleic Acids Res.* 1987, 15, 9083-9084); Genbank accession number Y00467, provided herein as SEQ ID NO: 107. Oligonucleotides were synthesized primarily as phosphorothioate oligodeoxynucleotides. Oligonucleotide sequences are shown in Table 18. Oligonucleotide 3082 (SEQ ID NO. 141) is an antisense oligodeoxynucleotide targeted to the human intracellular adhesion molecule-1 (ICAM-1) and was used as an unrelated target control. Oligonucleotide 13108 (SEQ ID NO. 142) is an antisense oligodeoxynucleotide targeted to the herpes simplex virus type 1 and was used as an unrelated target control.

[0139] P388D1, mouse macrophage cells (obtained from American Type Culture Collection, Manassas, Va.) were cultured in RPMI 1640 medium with 15% fetal bovine serum (FBS) (Life Technologies, Rockville, Md.).

[0140] At assay time, cell were at approximately 90% confluency. The cells were incubated in the presence of OPTI-MEM7 medium (Life Technologies, Rockville, Md.), and the oligonucleotide formulated in LIPOFECTIN7 (Life Technologies), a 1:1 (w/w) liposome formulation of the cationic lipid N-[1-(2,3-dioleyloxy)propyl]-n,n,n-trimethylammonium chloride (DOTMA), and dioleoyl phosphatidylethanolamine (DOPE) in membrane filtered water. For an initial screen, the oligonucleotide concentration was 100 nM in 3 μ g/ml LIPOFECTIN7. Treatment was for four hours. After treatment, the medium was removed and the cells were further incubated in RPMI medium with 15% FBS and induced with 10 ng/ml LPS. mRNA was analyzed 2 hours post-induction with PMA.

[0141] Total mRNA was isolated using the TOTALLY RNAT kit (Ambion, Austin, Tex.), separated on a 1% agarose gel, transferred to HYBONDTM-N+ membrane (Amersham, Arlington Heights, Ill.), a positively charged nylon membrane, and probed. A TNF- α probe consisted of the 502 bp EcoRI-HindIII fragment from BBG 56 (R&D Systems, Minneapolis, Minn.), a plasmid containing mouse TNF- α cDNA. A glyceraldehyde 3-phosphate dehydrogenase (G3PDH) probe consisted of the 1.06 kb HindIII fragment from pHcGAP (American Type Culture Collection, Manassas, Va.), a plasmid containing human G3PDH cDNA. The fragments were purified from low-melting temperature agarose, as described in Maniatis, T., et al., *Molecular Cloning: A Laboratory Manual*, 1989 and labeled with REDIVUETM32P-dCTP (Amersham Pharmacia Biotech, Piscataway, N.J.) and PRIME-A-GENE7 labeling kit (Promega, Madison, Wis.). mRNA was quantitated by a PhosphorImager (Molecular Dynamics, Sunnyvale, Calif.).

[0142] Secreted TNF- α protein levels were measured using a mouse TNF- α ELISA kit (R&D Systems, Minneapolis, Minn. or Cambridge, Ma.).

TABLE 18

Nucleotide Sequences of Mouse TNF- α Phosphorothioate Oligodeoxynucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION
14846	GAGCTTCTGCTGGCTGGGCTG	108	4351-4370	5'-UTR
14847	CCTTGCTGTCCTCGCTGAGG	109	4371-4390	5'-UTR
14848	TCATGGTGTCTTTTCTGGAG	110	4511-4530	AUG
14849	CTTTCTGTGCTCATGGTGTC	111	4521-4540	AUG
14850	GCGGATCATGCTTTCTGTGC	112	4531-4550	coding
14851	GGGAGGCCATTTGGGAATT	113	5225-5244	junction
14852	CGAATTTTGAGAAGATGATC	114	5457-5476	junction
14853	CTCCTCCACTTGGTGGTTTG	115	5799-5818	junction

TABLE 18-continued

Nucleotide Sequences of Mouse TNF- α Phosphorothioate Oligodeoxynucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION
14854	CCTGAGATCTTATCCAGCCT	116	6540-6559	3'-UTR
14855	CAATTACAGTCACGGCTCCC	117	6927-6946	3'-UTR
15921	CCCTTCATTCTCAAGGCACA	118	5521-5540	junction
15922	CACCCCTCAACCCGCCCCCC	119	5551-5570	intron
15923	AGAGCTCTGTCTTTCTCAG	120	5581-5600	intron
15924	CACTGCTCTGACTCTCACGT	121	5611-5630	intron
15925	ATGAGGTCCCGGGTGGCCCC	122	5651-5670	intron
15926	CACCCCTGTCTTTCCACAT	123	5681-5700	intron
15927	CTCCACATCCTGAGCCTCAG	124	5731-5750	intron
15928	ATTGAGTCAGTGTACCCCTC	125	5761-5780	intron
15929	GCTGGCTCAGCCACTCCAGC	126	5821-5840	coding
15930	TCTTTGAGATCCATGCCGTT	127	5861-5880	coding
15931	AACCCATCGGCTGGCACCAC	128	5891-5910	coding
15932	GTTTGAGCTCAGCCCCCTCA	129	6061-6080	coding
15933	CTCCTCCCAGGTATATGGGC	130	6091-6110	coding
15934	TGAGTTGGTCCCCTTCTCC	131	6121-6140	coding
15935	CAAAGTAGACCTGCCCGGAC	132	6181-6200	coding
15936	ACACCCATTCCCTTCACAGA	133	6211-6230	STOP
15937	CATAATCCCCTTCTAAGTT	134	6321-6340	3'-UTR
15938	CACAGAGTTGGA CTCTGAGC	135	6341-6360	3'-UTR
15939	CAGCATCTTGTTTCTGAG	136	6381-6400	3'-UTR
15940	CACAGTCCAGGTCACTGTCC	137	6401-6420	3'-UTR
15941	TGATGGTGGTGCATGAGAGG	138	6423-6442	3'-UTR
15942	GTGAATTCGGAAGCCCAT	139	6451-6470	3'-UTR
15943	CCTGACCACCTCCCTTTGC	140	6501-6520	3'-UTR
3082	TGCATCCCCCAGGCCACCAT	141	target control	
13108	GCCGAGGTCCATGTCGTACGC	142	target control	

¹ All "C" residues are 5-methyl-cytosines except underlined "C" residues are unmodified cytosines; all linkages are phosphorothioate linkages.
²Co-ordinates from Genbank Accession No. Y00467, locus name "MMTNFAB", SEQ ID NO.107.

[0143] Results are shown in Table 19. Oligonucleotides 14853 (SEQ ID NO. 115), 14854 (SEQ ID NO. 116), 14855 (SEQ ID NO. 117), 15921 (SEQ ID NO. 118), 15923 (SEQ ID NO. 120), 15924 (SEQ ID NO. 121), 15925 (SEQ ID NO. 122), 15926 (SEQ ID NO. 123), 15929 (SEQ ID NO. 126), 15930 (SEQ ID NO. 127), 15931 (SEQ ID NO. 128), 15932 (SEQ ID NO. 129), 15934 (SEQ ID NO. 131), 15935 (SEQ ID NO. 132), 15936 (SEQ ID NO. 133), 15937 (SEQ ID NO. 134), 15939 (SEQ ID NO. 136), 15940 (SEQ ID NO. 137), 15942 (SEQ ID NO. 139), and 15943 (SEQ ID NO. 140) gave better than 50% inhibition. Oligonucleotides 15931 (SEQ ID NO. 128), 15932 (SEQ ID NO. 129), 15934 (SEQ ID NO. 131), and 15943 (SEQ ID NO. 140) gave 75% inhibition or better.

TABLE 19

Inhibition of Mouse TNF- α mRNA expression in P388D1 Cells by Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
Induced	—	—	100%	0%
3082	141	control	129%	—
13664	42	control	85%	15%
14846	108	5'-UTR	84%	16%
14847	109	5'-UTR	88%	12%
14848	110	AUG	60%	40%
14849	111	AUG	75%	25%
14850	112	coding	67%	33%
14851	113	junction	62%	38%
14852	114	junction	69%	31%
14853	115	junction	49%	51%
14854	116	3'-UTR	31%	69%
14855	117	3'-UTR	39%	61%
15921	118	junction	42%	58%
15922	119	intron	64%	36%
15923	120	intron	31%	69%
15924	121	intron	29%	71%
15925	122	intron	30%	70%
15926	123	intron	29%	71%
15928	125	intron	59%	41%
15929	126	coding	38%	62%
15930	127	coding	43%	57%
15931	128	coding	23%	77%
15932	129	coding	25%	75%
15933	130	coding	52%	48%
15934	131	coding	21%	79%
15935	132	coding	39%	61%
15936	133	STOP	35%	65%
15937	134	3'-UTR	45%	55%
15938	135	3'-UTR	76%	24%
15939	136	3'-UTR	33%	67%
15940	137	3'-UTR	38%	62%
15941	138	3'-UTR	54%	46%
15942	139	3'-UTR	42%	58%
15943	140	3'-UTR	25%	75%

Example 9

Dose Response of Antisense Phosphorothioate
Oligodeoxynucleotide Effects on Mouse TNF- α
mRNA Levels in P388D1 Cells

[0144] Four of the more active oligonucleotides from the initial screen were chosen for dose response assays. These include oligonucleotides 15924 (SEQ ID NO. 121), 15931 (SEQ ID NO. 128), 15934 (SEQ ID NO. 131) and 15943 (SEQ ID NO. 140). P388D1 cells were grown, treated and processed as described in Example 8. LIPOFECTIN7 was added at a ratio of 3 μ g/ml per 100 nM of oligonucleotide. The control included LIPOFECTIN7 at a concentration of 6 μ g/ml. Results are shown in Table 20. Each oligonucleotide tested showed a dose response effect with maximal inhibition about 70% or greater and IC₅₀ values less than 50 nM.

TABLE 20

Dose Response of LPS-Induced P388D1 Cells to TNF- α Antisense Phosphorothioate Oligodeoxynucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Express- ion	% mRNA Inhibit- ion
induced	—	—	—	100%	—
13108	142	control	25 nM	68%	32%
"	"	"	50 nM	71%	29%
"	"	"	100 nM	64%	36%
"	"	"	200 nM	75%	25%
15924	121	intron	25 nM	63%	37%
"	"	"	50 nM	49%	51%
"	"	"	100 nM	36%	64%
"	"	"	200 nM	31%	69%
15931	128	coding	25 nM	42%	58%
"	"	"	50 nM	30%	70%
"	"	"	100 nM	17%	83%
"	"	"	200 nM	16%	84%
15934	131	coding	25 nM	37%	63%
"	"	"	50 nM	26%	74%
"	"	"	100 nM	13%	87%
"	"	"	200 nM	13%	87%
15943	140	3'-UTR	25 nM	38%	62%
"	"	"	50 nM	38%	62%
"	"	"	100 nM	16%	84%
"	"	"	200 nM	16%	84%

Example 10

[0145] Design and Testing of 2'-O-methoxyethyl (Deoxy Gapped) TNF- α Antisense Oligonucleotides on TNF- α Levels in P388D1 Cells

[0146] Oligonucleotides having SEQ ID NO: 128, SEQ ID NO: 131, and SEQ ID NO: 140 were synthesized as uniformly phosphorothioate oligodeoxynucleotides or mixed phosphorothioate/phosphodiester chimeric oligonucleotides having variable regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides. The sequences and the oligonucleotide chemistries are shown in Table 21. All 2'-MOE cytosines were 5-methyl-cytosines. Oligonucleotides were screened as described in Example 8. Results are shown in Table 22. All the oligonucleotides tested, except oligonucleotide 16817 (SEQ ID NO. 140) showed 44% or greater inhibition of TNF- α mRNA expression. Oligonucleotides 16805 (SEQ ID NO: 131), 16813 (SEQ ID NO: 140), and 16814 (SEQ ID NO: 140) showed greater than 70% inhibition.

TABLE 21

Nucleotide Sequences of Mouse 2'-O-inethoxyethyl (deoxy gapped) TNF-α Oligonucleotides				
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION
15931	AsAsCsCsCsAsTsCsGsGsCsTsGsGsCsAsCsCsAsC	128	5891-5910	coding
16797	AoAoCoCsCsAsTsCsGsGsCsTsGsGsCsAsCoCoAoC	"	5891-5910	coding
16798	AsAsCsCsCsAsTsCsGsGsCsTsGsGsCsAsCsCsAsC	"	5891-5910	coding
16799	AoAoCoCoCsAsTsCsGsGsCsTsGsGsCsAoCoCoAoC	"	5891-5910	coding
16800	AsAsCsCsCsAsTsCsGsGsCsTsGsGsCsAsCsCsAsC	"	5891-5910	coding
16801	AoAoCoCoCoAoToCoGsGsCsTsGsGsCsAsCsCsAsC	"	5891-5910	coding
16802	AsAsCsCsCsAsTsCsGsGsCsTsGsGsCsAsCsCsAsC	"	5891-5910	coding
16803	AsAsCsCsCsAsTsCsGsGsCsToGoGoCoAoCoCoAoC	"	5891-5910	coding
16804	AsAsCsCsCsAsTsCsGsGsCsTsGsGsCsAsCsCsAsC	"	5891-5910	coding
15934	TsGsAsGsTsTsGsGsTsCsCsCsCsCsTsTsCsTsCsC	131	6121-6140	coding
16805	ToGoAoGsTsTsGsGsTsCsCsCsCsCsTsTsCoToCoC	"	6121-6140	coding
16806	TsGsAsGsTsTsGsGsTsCsCsCsCsCsTsTsCsTsCsC	"	6121-6140	coding
16807	ToGoAoGoTsTsGsGsTsCsCsCsCsCsTsTsCoToCoC	"	6121-6140	coding
16808	TsGsAsGsTsTsGsGsTsCsCsCsCsCsTsTsCsTsCsC	"	6121-6140	coding
16809	ToGoAoGoToToGoGoTsCsCsCsCsCsTsTsCsTsCsC	"	6121-6140	coding
16810	TsGsAsGsTsTsGsGsTsCsCsCsCsCsTsTsCsTsCsC	"	6121-6140	coding
16811	TsGsAsGsTsTsGsGsTsCsCsCoCoCoToToCoToCoC	"	6121-6140	coding
16812	TsGsAsGsTsTsGsGsTsCsCsCsCsCsTsTsCsTsCsC	"	6121-6140	coding
15943	CsCsTsGsAsCsCsAsCsTsCsTsCsCsCsTsTsTsGsC	140	6501-6520	3'-UTR
16813	CoCoToGsAsCsCsAsCsTsCsTsCsCsCsTsToToGoC	"	6501-6520	3'-UTR
16814	CsCsTsGsAsCsCsAsCsTsCsTsCsCsCsTsTsTsGsC	"	6501-6520	3'-UTR
16815	CoCoToGoAsCsCsAsCsTsCsTsCsCsCsToToToGoC	"	6501-6520	3'-UTR
16816	CsCsTsGsAsCsCsAsCsTsCsTsCsCsCsTsTsTsGsC	"	6501-6520	3'-UTR
16817	CoCoToGoAoCoCoAoCsTsCsTsCsCsCsTsTsTsGsC	"	6501-6520	3'-UTR
16818	CsCsTsGsAsCsCsAsCsTsCsTsCsCsCsTsTsTsGsC	"	6501-6520	3'-UTR
16819	CsCsTsGsAsCsCsAsCsTsCsToCoCoCoToToToGoC	"	6501-6520	3'-UTR
16820	CsCsTsGsAsCsCsAsCsTsCsTsCsCsCsTsTsTsGsC	"	6501-6520	3'-UTR

¹ Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines are 5-methyl-cytidines; "s" linkages are phosphorothioate linkages, "o" linkages are phosphodiester linkages, "o" linkages are phosphodiester linkages.

² Co-ordinates from Genbank Accession No. Y00467, locus name "MMTNFAB", SEQ ID NO.107.

[0147]

TABLE 22

Inhibition of mouse TNF- α mRNA expression in P388D1 Cells by 2'-O-methoxyethyl (deoxy gapped) Oligonucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
induced	—	—	100%	0%
13108	142	control	87%	13%
15934	131	coding	28%	72%
16797	128	coding	33%	67%
16798	"	coding	34%	66%
16799	"	coding	56%	44%
16800	"	coding	35%	65%
16801	"	coding	34%	66%
16802	"	coding	38%	62%
16803	"	coding	35%	65%
16804	"	coding	39%	61%
16805	131	coding	29%	71%
16806	"	coding	31%	69%
16807	"	coding	46%	54%
16808	"	coding	43%	57%
16809	"	coding	33%	67%
16810	"	coding	37%	63%
16811	"	coding	40%	60%
16812	"	coding	31%	69%
16813	140	3'-UTR	28%	72%
16814	"	3'-UTR	28%	72%
16815	"	3'-UTR	46%	54%
16816	"	3'-UTR	49%	51%
16817	"	3'-UTR	172%	—
16818	"	3'-UTR	34%	66%
16819	"	3'-UTR	51%	49%
16820	"	3'-UTR	44%	56%

Example 11

[0148] Effect of TNF- α Antisense Oligonucleotides in a Murine Model for Non-Insulin-dependent Diabetes Mellitus

[0149] The db/db mouse model, a standard model for non-insulin-dependent diabetes mellitus (NIDDM; Hottel, G. S., et al., Science, 1993, 259, 87-90), was used to assess the activity of TNF- α antisense oligonucleotides on blood glucose levels and TNF- α mRNA levels in whole mice. These mice have elevated blood glucose levels and TNF- α mRNA levels compared to wild type mice. Female db/db mice and wild-type littermates were purchased from Jackson Laboratories (Bar Harbor, Me.). The effect on oligonucleotide 15931 (SEQ ID NO. 128) on blood glucose levels was determined. For determination of TNF- α mRNA levels, oligonucleotide 15931 (SEQ ID NO. 128), a uniformly modified phosphorothioate oligodeoxynucleotide, was compared to oligonucleotide 25302 (SEQ ID NO. 128), a mixed phosphorothioate/phosphodiester chimeric oligonucleotide having regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides. The sequences and chemistries are shown in Table 23. Oligonucleotide 18154 (SEQ ID NO. 143) is an antisense mixed phosphorothioate/phosphodiester chimeric oligonucleotide, having regions of 2'-O-methoxyethyl (2'-MOE) nucleotides and deoxynucleotides, targeted to the human vascular cell adhesion molecule-1 (VCAM-1) and was used as an unrelated target control.

TABLE 23

Nucleotide Sequence of TNF- α Antisense Oligonucleotide					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	
15931	AACCCATCGGCTGGCACCAC	128	5891-5910	coding	
25302	AACCCATCGGCTGGCACCAC	128	5891-5910	coding	
18154	TCAAGCAGTGCCACCGATCC	143	target control		

¹ All 2'-methoxyethyl cytosines and 2'-deoxy cytosines residues are 5-methyl-cytosines; all linkages are phosphorothioate linkages.
² Co-ordinates from Genbank Accession No. Y00467, locus name "MMTNFAB", SEQ ID NO.107.

[0150] db/db mice, six to ten weeks old, were dosed intraperitoneally with oligonucleotide every other day for 2 weeks at 10 mg/kg. The mice were fasted for seven hours prior to administration of the oligonucleotide. The mice were bled via retro orbital sinus every other day, and glucose measurements were performed on the blood. Results are shown in Table 24. Oligonucleotide 15931 (SEQ ID NO. 128) was able to reduce blood glucose levels in db/db mice to levels comparable with wild type mice. Food intake between wild type mice, treated and untreated, did not differ. Food intake between db/db mice, treated and untreated, although higher than wild type mice, did not differ significantly.

[0151] Samples of the fat (adipose) tissue from the inguinal fat pads were taken for RNA extraction. RNA was extracted according to *Current Protocols in Molecular Biology*, 1997, Ausubel, F., et al. ed., John Wiley & Sons. RNA was purified using the RNA clean up procedure of the RNEASY7 Mini kit (Qiagen, Valencia, Calif.). TNF- α mRNA levels were measured using the RIBOQUANT7 kit (PharMingen, San Diego, Calif.) with 15 μ g of RNA per lane. The probe used was from the mCK-3b Multi-Probe Template set (PharMingen, San Diego, Calif.) labeled with [α^{32} p] UTP (Amersham Pharmacia Biotech, Piscataway, N.J.). Results are shown in Table 25. Both oligonucleotide 15931 (SEQ ID NO. 128) and 25302 (SEQ ID NO. 128) were able to reduce TNF- α levels in fat, with 25302 (SEQ ID NO. 128) reducing TNF- α to nearly wild-type levels.

TABLE 24

Level of Blood Glucose in Normal and db/db Mice After Treatment with TNF- α Antisense Oligonucleotides					
Mouse Strain	ISIS #	SEQ ID NO:	ASO Gene Target	Time (days)	blood glucose (mg/dL)
wild type	—	—	—	1	140
"	15931	128	coding	"	138
db/db	—	—	—	1	260
"	15931	128	coding	"	254
wild type	—	—	—	9	175
"	15931	128	coding	"	163
db/db	—	—	—	9	252
"	15931	128	coding	"	128

[0152]

TABLE 25

Level of TNF- α mRNA in Fat of db/db Mice After Treatment with TNF- α Antisense Oligonucleotides			
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION
wt saline	—	—	100%
db/db saline	—	—	362%
18154	142	control	130%
15931	128	coding	210%
25302	128	coding	417%

Example 12

[0153] Effect of TNF- α Antisense Oligonucleotides in a Murine Model for Rheumatoid Arthritis

[0154] Collagen-induced arthritis (CIA) was used as a murine model for arthritis (Mussener, A., et al., Clin. Exp. Immunol., 1997, 107, 485-493). Female DBA/1LacJ mice (Jackson Laboratories, Bar Harbor, Me.) between the ages of 6 and 9 weeks were used to assess the activity of TNF- α antisense oligonucleotides. In all studies, 10 mice were used per treatment group.

[0155] On day 0, the mice were immunized at the base of the tail with 100 μ g of bovine type II collagen which was emulsified in Complete Freund's Adjuvant (CFA). On day 7, a second booster dose of collagen was administered by the same route. On day 14, the mice were injected subcutaneously with 100 μ g of LPS. Oligonucleotide was administered intraperitoneally (bolus) three times per week, starting on day 0, for the duration of the 7 week study at the indicated doses. The anti-TNF- α mAb (MM350D, Endogen, Woburn, Mass.) was administered intraperitoneally at 2 mg/kg once per week, starting on day 0. This antibody was formulated free of preservatives and carrier, and had an endotoxin level of 9.06 EU/mg.

[0156] Weights were recorded weekly. Mice were inspected daily for the onset of CIA, characterized by erythema and edema. Upon the onset of the disease, an assessment chart for each animal was started. Paw widths are rear ankle widths of affected and unaffected joints were measured three times a week using a constant tension caliper. Limbs were clinically evaluated and graded on a scale from 0-4, where 0=normal, 1=one digit swollen, 2=inflammation present in more than one digit, 3=joint distortion with or without inflammation, and 4=ankylosis as detected by joint manipulation. The progression of all measurements recorded to day 50. On day 50, animals were euthanized by cervical dislocation. All paws were removed and fixed in 10% neutral buffered formalin, from which histopathology slides were prepared.

[0157] Arthritis was classified into four stages based on histological evaluation of the degrees of inflammation, cartilage damage, pannus formation, bone erosion, osteolysis, fibrosis and ankylosis. Stage I is described by inflammatory cell infiltration in the tissues surrounding the joint and/or superficial layers of the synovium. Stage II is described by pannus formation with damage to the superficial layers of the cartilage. Stage III is described by subchondral bone

erosion with some degree of osteolysis. Stage IV is described by severe destruction of cartilage and bone with areas of fibrosis and/or bony ankylosis. The clinical data was analyzed for differences in the incidence of disease, the onset of disease and the severity of the disease. Descriptive statistics and an analysis of variance (ANOVA) were performed. If a statistically significant difference was detected, a Dunnett's test was performed.

[0158] Two independent studies, which differed in dose range, showed that mice treated with ISIS 25302 had a reduced incidence of arthritis (FIGS. 1A-1B). The two dose ranges were 0.03 to 3.0 mg/kg (low range, FIG. 1A), and 2.5 to 20 mg/kg (high range, FIG. 1B). The lowest incidence of disease was observed in mice treated at doses of 3.0 (22%) and 2.5 mg/kg (38%) of ISIS 25302 respectively, as compared to the vehicle control incidence of 88% in both studies. No further reduction in the incidence of disease occurred in mice treated at higher doses. The onset of disease was delayed in groups treated with ISIS 25302, but varied between experiments (Table 1). The severity of the disease and the percent affected paws were also reduced by treatment with ISIS 25302. Best effects on these clinical outcomes were observed at 3.0 mg/kg in the low dose range study, and 2.5 and 20 mg/kg in the high dose range study.

[0159] Treatment of mice with the eight mismatch control, ISIS 30782 (5' CACCAAGCTGCGGTCCCCAA 3'; SEQ ID NO: 502), yielded variable results between the low dose (Table 26A) and high dose (Table 26B) range studies. In the low dose range study, the one group treated with the control oligonucleotide, at a dose of 3.0 mg/kg, showed comparable improvements in the clinical outcome in comparison to the group treated with the anti-TNF- α oligonucleotide of equivalent dose. In contrast, the eight mismatch control oligonucleotide had minimal effects on the clinical outcome in the high dose range study, at doses of 2.5, 5.0, and 10 mg/kg; but did show effects in the clinic at the highest dose of 20 mg/kg.

TABLE 26A

Treatment	Schedule	Dose (mg/kg)	% incidence	Day of onset	Severity ("SEM)	% affected paws
Vehicle	3x/wk	—	88	18.1"0.7	7.1"2.1	59
ISIS 25302	3x/wk	0.03	70	18.6"1.1	3.1"1.2	28
ISIS 25302	3x/wk	0.1	70	17.6"0.2	3.5"1.5	30
ISIS 25302	3x/wk	0.3	44	21.5"4.5	2.9"1.4	25
ISIS 25302	3x/wk	1.0	67	21.0"3.6	3.4"1.0	36
ISIS 25302	3x/wk	3.0	22	21.5"3.5	1.2"0.8	14
TNF mAb	1x/wk	2.0	30	28.0"1.5	1.3"0.7	8.3
8MM ctrl	3x/wk	3.0	22	17.5"0.5	1.0"0.7	8.3

[0160]

TABLE 26B

Treatment	Schedule	Dose (mg/kg)	% incidence	Day of Onset	Severity ("SEM)	% affected paws
Vehicle	3x/wk	—	88	17.6"0.4	6.0"1.6	53
ISIS 25302	3x/wk	2.5	38	28.3"10.8	2.1"1.5	19
ISIS 25302	3x/wk	5.0	50	23.2"5.7	4.5"1.7	40
ISIS 25302	3x/wk	10	44	17.0"0.4	4.0"1.7	33
ISIS 25302	3x/wk	20	56	23.8"5.1	2.2"1.4	19

TABLE 26B-continued

Treatment	Schedule	Dose (mg/kg)	% inci- dence	Day of Onset	Severity ("SEM)	% affected paws
8MM ctrl	3x/wk	2.5	71	17.4*0.7	6.3*2.2	57
8MM ctrl	3x/wk	5.0	86	20.7*3.1	6.6*2.1	57
8MM ctrl	3x/wk	10	80	18.0*0.6	6.4*1.5	55
8MM ctrl	3x/wk	20	44	19.5*1.6	1.7*1.3	17

[0161] In both tables, the incidence is the number of mice with at least one affected paw/total number of mice per group. Severity is the total clinical score/total number of mice in the group. Percent affected paws=(number of affected paws at termination/total number of paws in group)×100. 8 MM ctrl=eight mismatch control (ISIS 30782).

[0162] Efficacy of ISIS 25302 (3 mg/kg, three times per week) was found to be comparable to that of an anti-TNF-α mAb (2 mg/kg, once per week) as described in Table 26A. The disease incidence in mice treated with ISIS 25302 was 22% versus 30% for the group treated with the anti-TNF-α mb. Disease severity and percent affected paws were also reduced to a similar degree in the 3 mg/kg ISIS 25302 and anti-TNF-α mAb treated groups.

[0163] Mice treated with the anti-mTNF-α oligonucleotide, ISIS 25302, showed an improvement in the disease outcome when treated three times per week starting on the initial day of collagen-induction. Reduction of symptoms by the ISIS 25302 was dose dependent, and showed equivalent effects when compared to mice treated with an anti-TNF-α monoclonal antibody once per week from the time of collagen-induction. Histological evaluation of the joints showed a reduction in the incidence and severity of arthritic lesions in mice treated with ISIS 25302, but to a lesser extent than those mice treated with the anti-TNF-α mAb.

[0164] The efficacy of ISIS 25302 compares favorably to other anti-TNF biological agents which have been evaluated in the “classical” CIA model. For instance, treatment of mice with the recombinant human TNF receptor FC fusion protein prior to onset of disease resulted in a 28% incidence of disease as compared to 86% incidence in the saline control treated animals (Wooley, *J. Immunol.* 151:6602-6607, 1993). In addition, preventative treatment by an anti-TNF-α antibody in the “classical” model showed 40% reduction in paw swelling in the clinic, as well as reduction in histopathological severity (Williams, *Proc. Natl. Acad. Sci. U.S.A.* 89:9784-9788, 1992).

[0165] A marked difference was observed between the two independent studies of ISIS 25302 in this model of CIA, with respect to responsiveness of the animals to oligonucleotide treatment. Mice were more responsive to oligonucleotide treatment in the low dose range study. This responsiveness was reflected in the histological results, where all oligonucleotide treated groups showed a notable reduction in paw incidence in comparison to the vehicle group. In comparison to the high dose study, mice in the low dose study overall displayed a lower percentage of paws with arthritic changes at the histological level.

[0166] In conclusion, evaluation of ISIS 25302 in the accelerated CIA model has shown that an anti-TNF-α oli-

gonucleotide provides an alternative approach to treatment of related human disease indications. Potential advantages of the antisense oligonucleotide therapeutic approach, over the current anti-arthritic (biological) agents, include ease of administration and a lower potential for adverse effects from long term usage.

Example 13

[0167] Effect of TNF-α Antisense Oligonucleotides in a Murine Model for Contact Sensitivity

[0168] Contact sensitivity is a type of immune response resulting from contact of the surface of the skin with a sensitizing chemical. A murine model for contact sensitivity is widely used to develop therapies for chronic inflammation, autoimmune disorder, and organ transplant rejection (Goebeler, M., et al., *Int Arch. Allergy Appl. Immunol.*, 1990, 93, 294-299). One example of such a disease is atopic dermatitis. Female Balb/c mice between the ages of 8 and 12 weeks are used to assess the, activity of TNF-α antisense oligonucleotides in a contact sensitivity model.

[0169] Balb/c mice receive injections of oligonucleotide drug in saline via i.v. injection into the tail vein. The abdomen of the mice is shaved using an Oster hair clipper. The animals are anesthetized using isoflurane, and 25 μl of 0.2% 2,4-dinitrofluorobenzene (DNFB) in 4:1 acetone:olive oil is applied to the shaved abdomen two days in a row. After five days, 10 ml of 0.2% DNFB in the same vehicle is applied to the right ear. After each exposure, the mouse is suspended in air for two minutes to allow the DNFB to absorb into the skin. 24 and 48 hours after application of DNFB to the ear, the ear thickness is measured using a micrometer. Inflammation (dermatitis) is indicated by a ranked thickening of the ear. Thickness of the treated ear is compared to untreated (contralateral) ear thickness.

Example 14

[0170] Effect of TNF-α Antisense Oligonucleotides in an IL10(-/-) Murine Model for Colitis

[0171] The effects of antisense oligonucleotide-inhibition of TNF-α was studied in the IL-10^{-/-} mouse model of colitis. IL10 deficient mice IL-10^{-/-} display some of the features that are observed in Crohn’s disease such as discontinuous lesions throughout the gastrointestinal tract and have a cytokine profile that is characteristic of a Th1 immune response. Unlike Crohn’s disease, however, IL-10^{-/-} mice show a marked crypt hyperplasia and absence of fissures and fistulas. In addition, IL-10^{-/-} mice have elevated levels of TNF-α expression.

[0172] Animals were treated in a prophylactic manner with one of four doses of ISIS 25302 or ISIS. Dosing extended from two weeks of age, before the development of colitis, to eight weeks of age, a time at which IL-10^{-/-} mice typically exhibit advanced stages of colitis. Colitis was assessed by histological evaluation at the conclusion of the study, and the basal and induced secretion of IFN-γ and TNF-α from colon organ culture supernatants was also measured at that time.

[0173] Homozygous Interleukin-10 gene-deficient mice, generated on a 129 Sv/Ev background, and 129 Sv/Ev controls were housed under specific pathogen-free conditions. Mice were housed in micro-isolator cages with tight-

fitting lids containing spun-polyester fiber filters. Mice were injected every other day with either ISIS 25302 or ISIS 30782 (the 8 mismatch control) at 0.01, 0.1, 1.0, and 10 mg/kg from 2-8 weeks of age via subcutaneous injection.

[0174] Animals were sacrificed using sodium pentobarbital (160 mg/kg). Whole colons were harvested, cut lengthwise, and fixed in 10% phosphate-buffered formalin, paraffin-embedded, sectioned at 4 μ m, and stained with haematoxylin and eosin for light microscopic examination. The slides were reviewed independently by a pathologist in a blinded fashion and assigned a histological score for intestinal inflammation (Table 27). The total histological score represents the numerical sum of the four scoring criteria: mucosal ulceration, epithelial hyperplasia, lamina propria mononuclear cell infiltration, and lamina propria neutrophilic infiltration.

TABLE 27

Score	Mucosal ulceration	Epithelial hyperplasia	LP mononuclear infiltration	LP neutrophil infiltrate
0	Normal	Normal	Normal	Normal
1	Surface inflammation	Mild	Slight increase	Slight increase
2	Erosions	Moderate	Marked increase	Marked increase
3	Ulcerations	Pseudopolyps		

[0175] Colonic organ cultures were prepared from IL-10 gene-deficient mice treated for six weeks. Due to the patchy nature of colitis in IL-10 gene-deficient mice, whole colons were removed, cut lengthwise, flushed with PBS, and resuspended in tissue culture plates (Falcon 3046; Becton Dickinson Labware, Lincoln Park, N.J.) in RPMI-1640 medium supplemented with 10% fetal calf serum, 50 mM 2-mercaptoethanol, penicillin (100 U/mL), and streptomycin (100 U/mL). Cultures were incubated at 37° C. in 5% CO₂. After 24 hours in the absence (basal) or presence of 10 μ g/ mL LPS (*E. coli*, 0111:B4; Sigma), supernatants were harvested and stored at -70° C. for analysis of cytokine levels. TNF- α and IFN- γ levels in cell supernatants were measured using ELISA kits purchased from Biosource Cytoscreen (Montreal, Quebec).

[0176] Differences between treatment groups were evaluated by analysis of variance (ANOVA). Single arm analysis was performed by the Dunnett's test (SAS Institute Inc., Cary N.C.).

[0177] Over the 6-week treatment period, all treatment groups of IL-10 deficient mice gained weight at a similar rate (data not shown). At 8 weeks of age, IL-10^{-/-} mice displayed a patchy distribution of transmural acute and chronic inflammation, extensive mucosal ulceration, and epithelial hyperplasia. Table 28 shows the histological scores for colon tissue from IL-10^{-/-} mice treated with saline (vehicle), ISIS 25302 or ISIS 30782 (8 MM ctrl) from 2 to 8 weeks of age at the indicated doses (n=6). The "total" histological score is the summation of the scores determined for each of the four histological parameters: mucosal ulceration, epithelial hyperplasia, lamina propria (LP) mononuclear cell infiltration, and lamina propria neutrophilic infiltration. Mice receiving the 0.1 mg/kg dose of the anti-TNF- α oligonucleotide, ISIS 25302, demonstrated a marked improvement in their mucosal architecture, which was sta-

tistically significant (p<0.05) in comparison to the Vehicle (saline) group (FIG. 2). No other group showed a significant histological difference in comparison to Vehicle.

TABLE 28

Treatment	Score	Mucosal ulceration	Mucosal hyperplasia	Mononuclear infiltrate	Neutrophil infiltrate	Total
Saline	Mean	1.00	1.83	2.00	1.83	6.67
	Std.	0.89	0.41	0.00	0.41	1.21
	Dev.					
0.01 mg/kg	Mean	0.50	1.50	1.50	1.50	5.00
ISIS 25302	Std.	0.55	0.55	0.55	0.55	0.63
	Dev.					
0.1 mg/kg	Mean	0.50	0.83	1.33	1.00	3.67
ISIS 25302	Std.	0.55	0.41	0.52	0.63	0.52
	Dev.					
1 mg/kg	Mean	0.67	2.00	1.67	1.67	6.00
ISIS 25302	Std.	1.21	0.89	0.52	0.52	2.61
	Dev.					
10 mg/kg	Mean	1.17	1.83	1.83	1.17	6.00
ISIS 25302	Std.	1.47	0.98	0.41	0.75	2.83
	Dev.					
0.01 mg/kg	Mean	0.83	1.83	1.33	1.67	5.67
8MM ctrl	Std.	1.17	0.75	0.52	0.52	2.58
	Dev.					
0.1 mg/kg	Mean	1.00	1.67	1.33	1.17	5.17
8MM ctrl	Std.	0.63	0.52	0.52	0.52	0.63
	Dev.					
1 mg/kg	Mean	0.67	1.67	1.33	1.33	5.00
8MM ctrl	Std.	0.52	0.52	0.52	0.52	0.63
	Dev.					
10 mg/kg	Mean	0.83	2.00	1.33	1.50	5.67
8 MM ctrl	Std.	1.17	0.63	0.52	0.55	2.25
	Dev.					

[0178] Reduction of secreted TNF- α protein levels was observed in colon tissue isolated from mice treated every other day with 0.1 mg/kg of ISIS 25302 under both basal (FIG. 3A) and LPS-induced (FIG. 3B) conditions. IFN- γ protein secretion from the isolated colon tissue was also reduced in the 0.1 mg/kg ISIS 25302 treated group relative to the saline treated group under both culture conditions (basal, FIG. 4A; LPS-induced, FIG. 4B). These effects were sequence specific, as the eight base mismatch oligonucleotide at the same dose of 0.1 mg/kg had no effect on basal or LPS-induced TNF- α protein secretion, or LPS-induced IFN- γ secretion.

[0179] Although treatment of IL-10^{-/-} mice with an antisense oligonucleotide targeted to TNF- α had no effect on the rate at which these animals gained weight, anti-TNF- α oligonucleotide treatment did have effects on several key disease parameters. Most importantly, antisense treatment at a relatively low dose (0.1 mg/kg) significantly reduced histological signs of colitis in the mice. This included reductions in mucosal ulceration, mucosal hyperplasia, and infiltrations of mononuclear cells and neutrophils into the lamina propria of the colon. These effects were not seen with the eight-base mismatch control oligonucleotide, ISIS 30782, which indicated that the effect was sequence specific.

[0180] The histological improvement is most likely due to specific reduction in TNF- α protein levels with antisense treatment. Both the basal and LPS-induced secretion of TNF- α from colons of mice treated with 0.1 mg/kg of ISIS 25302 were reduced, while the control oligonucleotide had

no effect. A decrease in basal and induced IFN- γ levels also occurred in the mice treated with 0.1 mg/kg ISIS 25302. Interruption of the proinflammatory cytokine cascade by inhibition of TNF- α expression may have abrogated the recruitment and activation of CD4+ T cells that produce IFN- γ . TNF- α is known to activate expression of key inflammatory intermediates which promote this process, including expression of cell adhesion molecules, chemokines, and other proinflammatory cytokines (Zhang et al. "Tumor necrosis factor" in *The Cytokine Handbook*, 3rd ed., Academic Press Ltd., pp. 517-547; van Deventer, *Gut* 40:443-448, 1997).

[0181] A biphasic response to the anti-TNF- α oligonucleotide was observed in this genetically engineered mouse model of colitis, where optimal efficacy of the anti-TNF- α oligonucleotide occurred at the mid range dose of 0.1 mg/kg. Treatment at the higher doses of 1.0 and 10 mg/kg resulted in complete loss of efficacy, as observed histologically and by cytokine expression levels. The basis of this response may lie in the undefined roles of the pro- and anti-inflammatory cytokines in the absence of IL-10; and/or the pharmacokinetics and mechanism of action of the oligonucleotide.

[0182] In conclusion, ISIS 25302 reduced TNF- α expression levels in a dose and sequence-dependent manner in the IL-10 deficient mice. Specific reduction of this proinflammatory molecule diminished the pathological features associated with the intestinal injury and inflammation which occurs in the absence of IL-10 in these mice. The results from this mouse model of colitis indicate that antisense oligonucleotides to TNF- α represent a new treatment of Crohn's disease in man.

Example 15

[0183] Effect of TNF- α Antisense Oligonucleotides in a DSS-induced Murine Model for Colitis

[0184] The pathological features of DSS-induced colitis in mice are similar in many respects to human ulcerative colitis (UC) (Table 29). This model is characterized by ulceration, epithelial damage, mucosal or transmural inflammatory infiltrate, and lymphoid hyperplasia of the colon. These effects are attributed to inappropriate macrophage function, alterations of the lumina bacteria, and the direct toxic effects of DSS on the colonic epithelium (Okayasu, *Gastroenterol.* 98:694-702, 1990). Both acute and chronic colitis may be studied in this model, by alteration of the DSS administration schedule (Okayasu, 1990, supra.; Cooper et al., *Lab. Invest.* 69:238-249, 1993). The efficacy of an anti-TNF- α mAb has been shown in both the acute and chronic model of DSS-induced colitis (Murthy et al., *Aliment. Pharmacol. Ther.* 13:251-260, 1999; Kojouharoff et al., *Clin. Exp. Immunol.* 107:353-358, 1997), as well as efficacy of an antisense oligonucleotide to ICAM-1 in the acute model of DSS-induced colitis (Bennett et al., *J. Pharmacol. Exp. Ther.* 280:988-1000, 1997).

TABLE 29

Feature	Crohn's	Ulcerative colitis	DSS-induced colitis
Location	GI tract	Colon	Colon
Depth	Transmural	Mucosal	Mucosal
Extent	Discontinuous	Continuous	Continuous

TABLE 29-continued

Feature	Crohn's	Ulcerative colitis	DSS-induced colitis
Symptoms	Non-bloody diarrhea, fistula	Bloody diarrhea, no fistula	BD, no fistula
Granuloma	Yes	No	No
Genetic	Yes	Yes	Yes
Microbial	Yes	Yes	Yes
Immunological	Yes	Yes	Yes
Inflammation	Transmural	Epithelium	Epithelium
TNF- α	Elevated	Elevated	Elevated

[0185] ISIS 25302 was evaluated for efficacy in both the acute and chronic models of DSS-induced colitis. ISIS 25302 is similar in design to the human anti-TNF- α oligonucleotide, ISIS 104838, with respect to the phosphorothioate modified backbone, methylated cytosine residues, and modification of each of the five 5' and 3' sugar residues with 2'-O-(2-methoxyethyl).

[0186] Female Swiss-Webster mice, 7 to 8 weeks of age weighing 25 to 30 grams, were obtained from Taconic or Jackson Laboratory. The animals were housed at 22° C. and 12 hours of dark and light cycles. Mouse chow and water were made available ab libitum.

[0187] Female Swiss-webster mice (n=2) were intravenously injected with 20 mg/kg of ISIS 13920 in saline or with saline alone on day 1, 3, and 5 of the acute DSS-induced colitis protocol as described below. ISIS 13920 is a fully modified phosphorothioate oligodeoxynucleotide, 5'TCCGTCATCGCTCCTCAGGG 3' (SEQ ID NO: 503), with 2'-O-(2-methoxyethyl) modified indicated by underline. This oligonucleotide is directed to the human ras-Ha gene. Two additional groups (n=2) of normal mice (no DSS) were subjected to the same oligonucleotide administration protocol. Mice were sacrificed on day 7. Colons were removed, trimmed longitudinally, fixed in 10% neutral buffered formaldehyde, and processed through paraffin. Four micron sections were cut from paraffin-embedded tissues, and deparaffinized by standard histological procedures. Endogenous tissue peroxidase activity was quenched with Peroxidase Blocking Reagent (DAKO; Carpinteria, Calif.) for 10 min at room temperature (r.t.). Tissue was treated with proteinase K (DAKO) for 10 min at r.t. to make it permeable for staining. After blocking with normal donkey serum (Jackson Laboratory; Bar Harbor, Me.), the sections were incubated for 45 min at r.t. with the 2E1-B5 anti-oligonucleotide mAb (Butler et al., *Lab. Invest.*, 77:379-388, 1997). Sections were rinsed with PBS and then incubated with peroxidase conjugated rabbit anti-mouse IgG1 (Zymed Laboratories; San Francisco, Calif.) diluted 1:200 for 30 min at r.t. Slides were washed thoroughly with PBS and then stained for peroxidase activity by addition of 3,3'-diaminobenzidine (DAKO) for 5 min at r.t.

[0188] Mice received 4% dextran sodium sulfate (MW 40,000, ICN Biomedicals, Inc., Aurora Ohio) in double distilled water ad libitum from day 0 until day 5 to induce colitis. On day 5, the 4% DSS was replaced with plain drinking water.

[0189] Mice were first weighed and randomized into groups of seven or eight animals. Mice were administered

oligonucleotide every other day (q2d) by i.v. or s.c. injection at the indicated doses from day-2 to day 6. The vehicle group was administered 1 mL/kg 0.9% saline (Baxter Healthcare Corporation, Deerfield, Ill.) under a similar treatment protocol.

[0190] Disease activity index was calculated on day 7 based on the summation of the weight, hemocult, and stool consistency scores (Table 30). Mice were weighed initially on day 0, and then every day beginning on day 3 until time of sacrifice. The stool consistency from each mouse was evaluated daily by visible appearance, beginning on day 3. On the day of sacrifice, day 7, stool from each mouse was evaluated for occult blood using the Hemocult test (Smith-Kline Diagnostics, Inc., San Jose Calif.). After sacrifice, the colon was removed from the ileocecal junction to the anal verge. The entire colon was then measured and observed for gross changes in the appearance of the mucosa, the total length of the colon was noted, and sections of the colon were dissected for histopathological evaluation.

TABLE 30

Score	Weight loss	Stool consistency	Hemocult
0	None	Normal	Negative
1	1-5%		
2	6-10%	Loose stool	Positive
3	11-15%		
4	>15%	Diarrhea	Gross bleeding

[0191] Mice were first weighed and randomized into groups of eight to ten animals. Chronic colitis was induced by giving the mice 4% DSS in their drinking water for two cycles. For each cycle, DSS was administered until the disease activity index (DAI) reached a score of 2.0 to 2.5 (see scoring criteria below) in at least one group, at which time the 4% DSS was replaced with plain drinking water. The first cycle of DSS administration was followed by 14 days of plain drinking water. The second cycle of DSS was followed by 8 to 9 days of plain drinking water, at which time the mice were sacrificed.

[0192] Oligonucleotide was administered subcutaneously (s.c.) for four consecutive days starting on the second day of the first cycle, and then every other day thereafter at doses of 0.25 mg/kg, 2.5 mg/kg, and 12.5 mg/kg; or 0.5 and 2.5 mg/kg. TNF- α mAb was administered s.c. one time at the beginning of each cycle for a total of two treatments at 30 μ g/mouse.

[0193] Chronic colitis progression was determined by daily measurement of the Disease Activity Index (DAI), consisting of weight loss, stool consistency and hemocult scores (Cooper et al., 1993, supra.). Each parameter was given a score (Table 30) and the combined score was divided by three to obtain the disease activity index (DAI). This method has been shown to correlate with the histological measures of inflammation and crypt damage.

[0194] The damage to the crypts and extent of recovery were determined by histological analysis of the proximal and distal sections of the colon based on the crypt grade and percent involvement in each section. Crypt grades were scored as Grade 0=intact crypt; Grade 1=loss of 1/3 crypt; Grade 2=loss of 2/3 of crypt; Grade 3=loss of entire crypt w/intact epithelium; and Grade 4=loss of entire crypt w/loss

of epithelium (ulceration). Percent involvement was scored as 1=1-25%; 2=26-50%; 3=51-75%; and 4=76-100%. Total crypt score is the combined scores of the distal and proximal colon sections. The inflammation score is the product of the grade of inflammation and the extent of involvement, where Grade 0=normal; Grade 1=mild; Grade 2=moderate; Grade 3=Severe; and Percent Involvement 1=1-25%; 2=26-50%; 3=51-75%; 4=76-100%.

[0195] Total RNA was isolated from a 1 mm full length colon strip from each animal using the RNeasy Mini Kit (Qiagen, Valencia Calif.). Mouse TNF- α and G3PDH mRNA levels were determined by standard northern blot procedures. TNF- α probe signals were normalized to G3PDH probe signal.

[0196] Differences between treatment groups were evaluated by analysis of variance (ANOVA). If a statistically significant difference was detected by ANOVA then the Dunnett's test was applied (SAS Institute Inc., Cary N.C.).

[0197] Previous studies have examined the distribution of the "first-generation" phosphorothioate oligodeoxynucleotides in colon tissue of normal and DSS-treated mice, and demonstrated localization of oligonucleotide in both the lamina propia and the epithelial cells of the mucosal layer (Bennett, 1997, supra.). In this case, differences were observed between the two groups of mice with respect to degree of tissue accumulation as well as relative distribution between the lamina propia and epithelial cells. Disruption of the epithelial mucosa layer and influx of immune cells into the lamina propia in the DSS-treated mice coincided with increased accumulation of the oligonucleotide in the tissue, particularly in the epithelial layer.

[0198] To obtain information on the localization of a 2'-O-(2-methoxyethyl) modified (2'-MOE) phosphorothioate oligodeoxynucleotide a similar experiment was performed using immunohistochemical staining techniques, instead of autoradiographic or fluorescent techniques, to detect the oligonucleotide (Butler et al., 1997, supra.) in the colon tissue. Immunohistochemical staining allows for direct detection of the oligonucleotide without further labeling steps during oligonucleotide synthesis. The previously identified anti-oligonucleotide monoclonal antibody, 2E1, was utilized for this purpose (Butler, 1997, supra.). Cumulative studies have shown that the strength of the signal obtained from histological staining of an oligonucleotide with the 2E1 antibody is dependent on the oligonucleotide sequence. In this respect, the staining signal for ISIS 25302 proved to be modest. For this reason we utilized ISIS 13920, a 2'-MOE modified phosphorothioate oligodeoxynucleotide with enhanced histological staining properties, to evaluate the distribution of this type of oligonucleotide in colon tissue of normal and DSS-treated mice. A similar distribution and accumulation profile was observed with the "second-generation" 2'-MOE modified phosphorothioate oligodeoxynucleotide, as had previously been observed for a rhodamine labeled "first-generation" phosphorothioate oligodeoxynucleotide (Bennett, 1997, supra.). Enhanced staining by the anti-oligonucleotide antibody, 2E1, was observed in the colon tissue of DSS-treated mice, in comparison to the normal mice.

[0199] Mice treated with ISIS 25302 every other day at a dose of 1 mg/kg in the acute model of DSS-induced colitis showed a 44% reduction in the disease activity index (DAI)

relative to the saline treated control group (1.4 ± 0.2 vs 2.6 ± 0.2 ; **FIG. 5A**). In comparison, mice treated one time with 25 micrograms of the anti-TNF- α mAb, at the commencement of DSS-induction, showed a 57% reduction in the DAI. In both cases, the reduction in DAI was significant ($p < 0.05$) in comparison to the saline treated group. In contrast to the other two treatments, mice treated with 50 micrograms of antibody showed no improvement in the DAI. Improvement in the DAI correlated with an increase in colon length (**FIG. 5B**). The mean colon length of the saline treated DSS-induced mice was 57% the length of normal mice (see also Okayasu, 1990, *supra.*), whereas those of the ISIS 25302 and anti-TNF- α antibody (25 μ g) treated mice were 76% and 79% respectively. The mean colon lengths of each of the two anti-TNF- α treated groups were significantly different from both the saline treated DSS-induced mice and normal mice ($p < 0.05$).

[0200] The effect of ISIS 25302 on the development of acute colitis was dose and sequence dependent (**FIG. 6A-6B**). A reduction of the clinical symptoms of DSS-induced colitis, as measured by the DAI, was observed in mice treated with 0.04 (60%), 0.2 (60%), and 1 mg/kg (80%) of ISIS 25302 relative to saline treated control mice. Mice treated with the eight base mismatch control oligonucleotide, ISIS 30782, showed no reduction in the DAI in comparison to the saline treated group. The reduction in DAI in mice treated with ISIS 25302 at 0.04, 0.2, and 1.0 mg/kg was statistically significant in comparison to mice treated with the eight base mismatch control oligonucleotide at 1.0 mg/kg ($p < 0.05$). A statistically significant difference was also observed between the 1.0 mg/kg ISIS 25302 group and the saline treated group. Treatment of the mice with ISIS 25302 at the higher dose of 5 mg/kg, yielded no improvement in the DAI; as previously observed in mice treated with 50 micrograms of the anti-TNF- α mAb (described below). A partial loss of efficacy was also observed in the acute DSS-induced colitis model with the anti-ICAM-1 oligonucleotide, ISIS 3082, at a dose of 5 mg/kg (Bennett, 1997, *supra.*). In the ICAM-1 study mice were administered oligonucleotide once a day for five consecutive days, instead of every other day for a total of five injections. Loss of efficacy, in all applications, may have resulted from an excessive accumulation of the oligonucleotide (or antibody) in the inflamed tissue, which in turn had an adverse effect on the animals (immune) response to intestinal injury by DSS.

[0201] ISIS 25302 was also tested for efficacy in the chronic model of DSS-induced mouse colitis. In this model, DSS was administered a second time, fourteen days after the first period of DSS administration. Animals were treated with ISIS 25302 prior to establishment of disease, starting on Day 2 of the first cycle of DSS administration. A dose-dependent reduction in the clinical signs of chronic colitis was observed in the mice treated with ISIS 25302 (**FIG. 7A**). For example, a 49% reduction (0.88 ± 0.17) in the disease activity index (DAI) was observed in mice treated at the lowest dose of 0.25 mg/kg of ISIS 25302, in comparison to the saline treated control group (1.7 ± 0.3) at the end of the second cycle (Day 10, **FIG. 7B**). A greater reduction in the DAI, 86 to 87%, was observed in mice treated at the higher doses of 2.5 and 12.5 mg/kg of ISIS 25302 (0.22 ± 0.11 and 0.27 ± 0.11 , respectively). In comparison, animals treated with the anti-TNF- α mAb showed a 61% reduction in DAI (0.67 ± 0.14). At this time the reductions in DAI scores were statistically significant ($p < 0.05$) in mice treated with either

the anti-TNF- α mAb or ISIS 25302, at all three doses, in comparison to the vehicle group.

[0202] Mice that showed an improvement in DAI also showed a reduction in inflammatory infiltrates and crypt damage at the histological level as compared to the untreated and vehicle groups (**FIG. 8A-B**). For example, mice treated with ISIS 25302 at 2.5 and 12.5 mg/kg demonstrated a 43% and 52% reduction in total inflammatory infiltrates (respectively), and a 43% and 48% reduction in total crypt damage relative to vehicle (**FIG. 8A**). The proximal region of the colon was more responsive to treatment by ISIS 25302, than the distal region (**FIG. 8B**). However, the severity of the disease was greater in the distal region of the colon.

[0203] Although not statistically significant, a thirty percent reduction in target TNF- α mRNA levels was observed in the colon tissue of mice treated at the higher doses of 2.5 and 12.5 mg/kg ISIS 25302 (**FIG. 9**). The TNF- α mRNA levels in colons from mice treated at the lower dose of 0.25 mg/kg of ISIS 25302 were not reduced in comparison to the vehicle group. The reduced levels of TNF- α mRNA observed for mice treated with the two higher doses of ISIS 25302 supports the dose-dependent response observed in the clinic, as measured by the DAI.

[0204] The anti-mTNF- α oligonucleotide, ISIS 25302, showed dose and sequence-specific efficacy in both the acute and chronic indications of DSS-induced colitis. ISIS 25302 treatment was also comparable in effect to treatment with an anti-TNF mAb in both indications. The reduction in the clinical symptoms observed in DSS-induced mice treated with ISIS 25302 correlated with a reduction of inflammatory infiltrates and crypt damage. Target TNF- α mRNA levels were also reduced in colon tissue derived from DSS-induced animals treated with ISIS 25302, relative to vehicle controls. The efficacy of ISIS 25302 in both the acute and chronic models of DSS-induced mouse colitis indicates that an antisense oligonucleotide which targets TNF- α mRNA represents a novel approach for treatment of human inflammatory bowel disease.

Example 16

[0205] Effect of TNF- α Antisense Oligonucleotides in a Murine Model for Crohn's Disease

[0206] C3H/HeJ, SJL/JK and IL10^{-/-} mice are used in a TNBS (2,4,5-trinitrobenzene sulfonic acid) induced colitis model for Crohn's disease (Neurath, M. F., et al., J. Exp. Med., 1995, 182, 1281-1290). Mice between the ages of 6 weeks and 3 months are used to assess the activity of TNF- α antisense oligonucleotides.

[0207] C3H/HeJ, SJL/JK and IL10^{-/-} mice are fasted overnight prior to administration of TNBS. A thin, flexible polyethylene tube is slowly inserted into the colon of the mice so that the tip rests approximately 4 cm proximal to the anus. 0.5 mg of the TNBS in 50% ethanol is slowly injected from the catheter fitted onto a 1 ml syringe. Animals are held inverted in a vertical position for approximately 30 seconds. TNF- α antisense oligonucleotides are administered either at the first sign of symptoms or simultaneously with induction of disease. Animals, in most cases, are dosed every day. Administration is by i.v., i.p., s.q., minipumps or intracolonic injection. Experimental tissues are collected at the end of the treatment regimen for histochemical evaluation.

Example 17

[0208] Effect of TNF-α Antisense Oligonucleotides in a Murine Model for Multiple Sclerosis

[0209] Experimental autoimmune encephalomyelitis (EAE) is a commonly accepted murine model for multiple sclerosis (Myers, K. J., et al., J. Neuroimmunol., 1992, 41, 1-8). SJL/H, PL/J, (SJLxPL/J) F1, (SJLxBalb/c) F1 and Balb/c female mice between the ages of 6 and 12 weeks are used to test the activity of TNF-α antisense oligonucleotides.

[0210] The mice are immunized in the two rear foot pads and base of the tail with an emulsion consisting of encephalitogenic protein or peptide (according to Myers, K. J., et al., J. of Immunol., 1993, 151, 2252-2260) in Complete Freund's Adjuvant supplemented with heat killed Mycobacterium tuberculosis. Two days later, the mice receive an intravenous injection of 500 ng Bordatella pertussis toxin and additional adjuvant.

[0211] Alternatively, the disease may also be induced by the adoptive transfer of T-cells. T-cells are obtained from the draining of the lymph nodes of mice immunized with encephalitogenic protein or peptide in CFA. The T cells are grown in tissue culture for several days and then injected intravenously into naive syngeneic recipients.

[0212] Mice are monitored and scored daily on a 0-5 scale for signals of the disease, including loss of tail muscle tone, wobbly gait, and various degrees of paralysis.

Example 18

[0213] Effect of TNF-α Antisense Oligonucleotides in a Murine Model for Pancreatitis

[0214] Swiss Webster, C57BL/56, C57BL/6 lpr and gld male mice are used in an experimental pancreatitis model (Niederau, C., et al., Gastroenterology, 1985, 88, 1192-1204). Mice between the ages of 4 and 10 weeks are used to assess the activity of TNF-α antisense oligonucleotides.

[0215] Caerulein (5-200 μg/kg) is administered i.p. every hour for one to six hours. At varying time intervals, the mice are given i.p. injection of avertin and bled by cardiac puncture. The pancreas and spleen are evaluated for histopathology and increased levels of IL-1β, IL-6, and TNF-α. The blood is analyzed for increased levels of serum amylase and lipase. TNF-α antisense oligonucleotides are administered by intraperitoneal injection at 4 hours pre-caerulein injections.

Example 19

[0216] Effect of TNF-α Antisense Oligonucleotides in a Murine Model for Hepatitis

[0217] Concanavalin A-induced hepatitis is used as a murine model for hepatitis (Mizuhara, H., et al., J. Exp. Med., 1994, 179, 1529-1537). It has been shown that this type of liver injury is mediated by Fas (Seino, K., et al., Gastroenterology 1997, 113, 1315-1322). Certain types of viral hepatitis, including Hepatitis C, are also mediated by Fas (J. Gastroenterology and Hepatology, 1997, 12, S223-S226). Female Balb/c and C57BL/6 mice between the ages of 6 weeks and 3 months are used to assess the activity of TNF-α antisense oligonucleotides.

[0218] Mice are intravenously injected with oligonucleotide. The pretreated mice are then intravenously injected with 0.3 mg concanavalin A (Con A) to induce liver injury. Within 24 hours following Con A injection, the livers are removed from the animals and analyzed for cell death (apoptosis) by in vitro methods. In some experiments, blood is collected from the retro-orbital vein.

Example 20

[0219] Effect of Antisense Oligonucleotide Targeted to TNF-α on Survival in Murine Heterotopic Heart Transplant Model

[0220] To determine the therapeutic effects of TNF-α antisense oligonucleotides in preventing allograft rejection, murine TNF-α-specific oligonucleotides are tested for activity in a murine vascularized heterotopic heart transplant model. Hearts from Balb/c mice are transplanted into the abdominal cavity of C3H mice as primary vascularized grafts essentially as described by Isobe et al., Circulation 1991, 84, 1246-1255. Oligonucleotide is administered by continuous intravenous administration via a 7-day Alzet pump. The mean survival time for untreated mice is usually approximately 9-10 days. Treatment of the mice for 7 days with TNF-α antisense oligonucleotides is expected to increase the mean survival time.

Example 21

[0221] Optimization of Human TNF-α Antisense Oligonucleotide

[0222] Additional antisense oligonucleotides targeted to intron 1 of human TNF-α were designed. These are shown in Table 31. Oligonucleotides are screened by RT-PCR as described in Example 5 hereinabove.

TABLE 31

Nucleotide Sequences of Human TNF-α Intron 1 Antisense Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5'→ 3')	TARGET GENE			
		SEQ ID NO:	NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	
100181	AGTGTCTTCTGTGTGCCAGA	144	1409-1428	intron 1	
100201	AGTGTCTTCTGTGTGCCAGA	"	"	intron 1	
100230	AGTGTCTTCTGTGTGCCAGA	"	"	intron 1	
100250	AGTGTCTTCTGTGTGCCAGA	"	"	intron 1	
100182	GTGTCTTCTGTGTGCCAGAC	145	1408-1427	intron 1	
100202	GTGTCTTCTGTGTGCCAGAC	"	"	intron 1	
100231	GTGTCTTCTGTGTGCCAGAC	"	"	intron 1	
100251	GTGTCTTCTGTGTGCCAGAC	"	"	intron 1	
100183	TGTCTTCTGTGTGCCAGACA	146	1407-1426	intron 1	
100203	TGTCTTCTGTGTGCCAGACA	"	"	intron 1	
100232	TGTCTTCTGTGTGCCAGACA	"	"	intron 1	
100252	TGTCTTCTGTGTGCCAGACA	"	"	intron 1	
100184	GTCTTCTGTGTGCCAGACAC	147	1406-1425	intron 1	
100204	GTCTTCTGTGTGCCAGACAC	"	"	intron 1	
100233	GTCTTCTGTGTGCCAGACAC	"	"	intron 1	
100253	GTCTTCTGTGTGCCAGACAC	"	"	intron 1	
100185	TCTTCTGTGTGCCAGACACC	148	1405-1424	intron 1	
100205	TCTTCTGTGTGCCAGACACC	"	"	intron 1	
100234	TCTTCTGTGTGCCAGACACC	"	"	intron 1	
100254	TCTTCTGTGTGCCAGACACC	"	"	intron 1	
100186	CTTCTGTGTGCCAGACACCC	149	1404-1423	intron 1	
100206	CTTCTGTGTGCCAGACACCC	"	"	intron 1	
100235	CTTCTGTGTGCCAGACACCC	"	"	intron 1	
100255	CTTCTGTGTGCCAGACACCC	"	"	intron 1	

TABLE 31-continued

Nucleotide Sequences of Human TNF- α Intron 1 Antisense Oligonucleotides					TARGET GENE		
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	CO-ORDINATES ²	GENE TARGET REGION			
100187	TTCTGTGTGCCAGACACCCCT	150	1403-1422	intron 1			
100207	TTCTGTGTGCCAGACACCCCT	"	"	intron 1			
100236	TTCTGTGTGCCAGACACCCCT	"	"	intron 1			
100256	TTCTGTGTGCCAGACACCCCT	"	"	intron 1			
100188	TCTGTGTGCCAGACACCCCTA	151	1402-1421	intron 1			
100208	TCTGTGTGCCAGACACCCCTA	"	"	intron 1			
100237	TCTGTGTGCCAGACACCCCTA	"	"	intron 1			
100257	TCTGTGTGCCAGACACCCCTA	"	"	intron 1			
100189	CTGTGTGCCAGACACCCCTAT	152	1401-1420	intron 1			
100209	CTGTGTGCCAGACACCCCTAT	"	"	intron 1			
100238	CTGTGTGCCAGACACCCCTAT	"	"	intron 1			
100258	CTGTGTGCCAGACACCCCTAT	"	"	intron 1			
100190	TGTGTGCCAGACACCCCTATC	153	1400-1419	intron 1			
100210	TGTGTGCCAGACACCCCTATC	"	"	intron 1			
100239	TGTGTGCCAGACACCCCTATC	"	"	intron 1			
100259	TGTGTGCCAGACACCCCTATC	"	"	intron 1			
100191	TGTGCCAGACACCCCTATCTT	154	1398-1417	intron 1			
100211	TGTGCCAGACACCCCTATCTT	"	"	intron 1			
100240	TGTGCCAGACACCCCTATCTT	"	"	intron 1			
100260	TGTGCCAGACACCCCTATCTT	"	"	intron 1			
100192	GTGCCAGACACCCCTATCTTC	155	1397-1416	intron 1			
100212	GTGCCAGACACCCCTATCTTC	"	"	intron 1			
100241	GTGCCAGACACCCCTATCTTC	"	"	intron 1			
100261	GTGCCAGACACCCCTATCTTC	"	"	intron 1			
100193	TGCCAGACACCCCTATCTTCT	156	1396-1415	intron 1			
100213	TGCCAGACACCCCTATCTTCT	"	"	intron 1			
100242	TGCCAGACACCCCTATCTTCT	"	"	intron 1			
100262	TGCCAGACACCCCTATCTTCT	"	"	intron 1			
100194	GCCAGACACCCCTATCTTCTT	157	1395-1414	intron 1			
100214	GCCAGACACCCCTATCTTCTT	"	"	intron 1			
100243	GCCAGACACCCCTATCTTCTT	"	"	intron 1			
100263	GCCAGACACCCCTATCTTCTT	"	"	intron 1			
100195	CCAGACACCCCTATCTTCTTC	158	1394-1413	intron 1			
100215	CCAGACACCCCTATCTTCTTC	"	"	intron 1			
100244	CCAGACACCCCTATCTTCTTC	"	"	intron 1			
100264	CCAGACACCCCTATCTTCTTC	"	"	intron 1			
100196	CAGACACCCCTATCTTCTTCT	159	1393-1412	intron 1			
100216	CAGACACCCCTATCTTCTTCT	"	"	intron 1			
100245	CAGACACCCCTATCTTCTTCT	"	"	intron 1			
100265	CAGACACCCCTATCTTCTTCT	"	"	intron 1			
100197	AGACACCCCTATCTTCTTCTC	160	1392-1411	intron 1			
100217	AGACACCCCTATCTTCTTCTC	"	"	intron 1			
100246	AGACACCCCTATCTTCTTCTC	"	"	intron 1			
100266	AGACACCCCTATCTTCTTCTC	"	"	intron 1			
100198	GACACCCCTATCTTCTTCTCT	161	1391-1410	intron 1			
100218	GACACCCCTATCTTCTTCTCT	"	"	intron 1			
100247	GACACCCCTATCTTCTTCTCT	"	"	intron 1			
100267	GACACCCCTATCTTCTTCTCT	"	"	intron 1			
100199	ACACCCCTATCTTCTTCTCTC	162	1390-1409	intron 1			
100219	ACACCCCTATCTTCTTCTCTC	"	"	intron 1			
100248	ACACCCCTATCTTCTTCTCTC	"	"	intron 1			
100268	ACACCCCTATCTTCTTCTCTC	"	"	intron 1			
100200	CACCCCTATCTTCTTCTCTCC	163	1389-1408	intron 1			
100220	CACCCCTATCTTCTTCTCTCC	"	"	intron 1			
100249	CACCCCTATCTTCTTCTCTCC	"	"	intron 1			
100269	CACCCCTATCTTCTTCTCTCC	"	"	intron 1			
100270	GTCCTTCTGTGTGCCAGAC	164	1408-1425	intron 1			
100271	TCTTCTGTGTGCCAGACA	165	1407-1424	intron 1			
100272	CTTCTGTGTGCCAGACAC	166	1406-1423	intron 1			
100273	TTCTGTGTGCCAGACACC	167	1405-1422	intron 1			
100274	TCTGTGTGCCAGACACCC	168	1404-1421	intron 1			
100275	CTGTGTGCCAGACACCCCT	169	1403-1420	intron 1			
100276	TGTGTGCCAGACACCCCTA	170	1402-1419	intron 1			
100277	GTGTGCCAGACACCCCTAT	171	1401-1418	intron 1			
100278	TGTGCCAGACACCCCTATC	172	1400-1417	intron 1			
100279	TGCCAGACACCCCTATCTT	173	1398-1415	intron 1			
100280	GCCAGACACCCCTATCTTC	174	1397-1414	intron 1			
100281	CCAGACACCCCTATCTTCT	175	1396-1413	intron 1			

TABLE 31-continued

Nucleotide Sequences of Human TNF- α Intron 1 Antisense Oligonucleotides					TARGET GENE		
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	CO-ORDINATES ²	GENE TARGET REGION			
100282	CAGACACCCCTATCTTCTT	176	1395-1412	intron 1			
100283	AGACACCCCTATCTTCTTC	177	1394-1411	intron 1			
100284	GACACCCCTATCTTCTTCT	178	1393-1410	intron 1			
100285	ACACCCCTATCTTCTTCTC	179	1392-1409	intron 1			

¹Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethyl cytosines and 2'-deoxy cytosines residues are 5-methyl-cytosines; all linkages are phosphorothioate linkages.
²Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO. 1.

Example 22

[0223] Design of Antisense Oligonucleotides Targeting Human TNF- α Intron 2

[0224] Additional antisense oligonucleotides targeted to intron 2 and coding regions of human TNF- α were designed. These are shown in Table 32. Oligonucleotides are screened by RT-PCR as described in Example 5 hereinabove.

TABLE 32

Nucleotide Sequences of Human TNF- α Intron 2 Antisense Oligonucleotides					TARGET GENE		
ISIS No.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	CO-ORDINATES ²	GENE TARGET REGION			
100549	AGAGGTTTGGAGACACTTAC	180	1635-1654	intron 2			
100566	AGAGGTTTGGAGACACTTAC	"	"	intron 2			
100550	GAATTAGGAAAGAGGTTTGG	181	1645-1664	intron 2			
100567	GAATTAGGAAAGAGGTTTGG	"	"	intron 2			
100551	CCCAAACCCAGAAATTAGGAA	182	1655-1674	intron 2			
100568	CCCAAACCCAGAAATTAGGAA	"	"	intron 2			
100552	TACCCCCAAACCCAAACCCA	183	1665-1684	intron 2			
100569	TACCCCCAAACCCAAACCCA	"	"	intron 2			
100553	GTACTAACCCCTACCCCCAAA	184	1675-1694	intron 2			
100570	GTACTAACCCCTACCCCCAAA	"	"	intron 2			
100554	TTCCATACCGGTACTAACCC	185	1685-1704	intron 2			
100571	TTCCATACCGGTACTAACCC	"	"	intron 2			
100555	CCCCCACTGCTTCCATACCG	186	1695-1714	intron 2			
100572	CCCCCACTGCTTCCATACCG	"	"	intron 2			
100556	CTTTAAATTCCCCCACTGC	187	1705-1724	intron 2			
100573	CTTTAAATTCCCCCACTGC	"	"	intron 2			
100557	AAGACCAAACTTTAAATTT	188	1715-1734	intron 2			
100571	AAGACCAAACTTTAAATTT	"	"	intron 2			
100558	ATCCTCCCCCAAGACCAAAA	189	1725-1744	intron 2			
100640	ATCCTCCCCCAAGACCAAAA	"	"	intron 2			
100559	ACCTCCATCCATCTCCCCC	190	1735-1754	intron 2			
100641	ACCTCCATCCATCTCCCCC	"	"	intron 2			
100560	CCCTACTTTCACCTCCATCC	191	1745-1764	intron 2			
100642	CCCTACTTTCACCTCCATCC	"	"	intron 2			
100561	GAAATATACCCCTACTTTTC	192	1755-1774	intron 2			
100643	GAAATATACCCCTACTTTTC	"	"	intron 2			
100562	AAACTTCCTAGAAAATACCC	193	1765-1784	intron 2			
100644	AAACTTCCTAGAAAATACCC	"	"	intron 2			
100563	TGAGACCCCTTAACTTCCTA	194	1775-1794	intron 2			
100645	TGAGACCCCTTAACTTCCTA	"	"	intron 2			
100564	AAGAAAAAGCTGAGACCCCT	195	1785-1804	intron 2			
100646	AAGAAAAAGCTGAGACCCCT	"	"	intron 2			

TABLE 32-continued

Nucleotide Sequences of Human TNF-α Intron 2 Antisense Oligonucleotides					
ISIS No.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	TARGET GENE			
		SEQ ID NO:	CO-ORDINATES ²	GENE	TARGET REGION
100565	GGAGAGAGAAAAGAAAAGC	196	1795-1814	intron 2	
100647	GGAGAGAGAAAAGAAAAGC	"	"	intron 2	
100575	TGAGCCAGAAGAGGTTGAGG	197	2665-2684	coding	
100576	ATTCTCTTTTGGAGCCAGAA	198	2675-2694	coding	
100577	TAAGCCCCCAATTCTCTTTT	199	2685-2704	coding	
100578	GTTCCGACCCTAAGCCCCCA	200	2695-2714	coding	
100579	CTAAGCTTGGGTCCGACCC	201	2705-2724	coding	
100580	GCTTAAAGTTCTAAGCTTGG	202	2715-2734	coding	
100581	TGGTCTTGTGTCTTAAAGTT	203	2725-2744	coding	
100582	TTCGAAGTGGTGTCTTGT	204	2735-2754	coding	
100583	AATCCACGGTTTCGAAGTGG	205	2745-2764	coding	
100584	CACATTCTGAATCCAGGT	206	2755-2774	coding	
100585	GTGCAGGCCACACATTCTG	207	2765-2784	coding	
100586	GCACCTTCACTGTGCAGGCCA	208	2775-2794	coding	
100587	GTGGTTGCCAGCACTTCACT	209	2785-2804	coding	
100588	TGAATTCCTTAGTGGTTGCCA	210	2795-2814	coding	
100589	GGCCCCAGTTTGAATCTTA	211	2805-2824	coding	
100590	GAGTTCTGGAGGCCCAAGTT	212	2815-2834	coding	
100591	AGGCCCCAGTGAGTTCTGGA	32	2825-2844	coding	
100592	TCAAGCTGTAGGCCCCAGT	214	2835-2854	coding	
100593	ATGTCAGGGATCAAGCTGT	215	2845-2864	coding	
100594	CAGATCCAGATGTCAGGGA	216	2855-2874	coding	
100595	CCCTGGTCTCCAGATTCCAG	217	2865-2884	coding	
100596	ACCAAAGGCTCCCTGGTCTC	218	2875-2894	coding	
100597	TCGGCCAGAACCAAGGCT	219	2885-2904	coding	
100598	CCTGCAGCATTCTGGCCAGA	220	2895-2914	coding	
100599	CTTCTCAAGTCCTGCAGCAT	221	2905-2924	coding	
100600	TAGGTGAGGTCTTCTCAAGT	222	2915-2934	coding	
100601	TGTCAATTCTAGGTGAGGT	223	2925-2944	coding	
100602	GGTCCACTTGTGTCAATTTT	224	2935-2954	coding	
100603	GAAGGCTAAGGTCCACTTG	225	2945-2964	coding	
100604	CTGGAGAGAGGAGGCCATA	226	2955-2974	coding	
100605	CTGGAACATCTGGAGAGAG	227	2965-2984	coding	
100606	TCAGGGAAGTCTGGAACAT	228	2975-2994	coding	
100607	GCTCCGTGTCTCAGGAAGT	229	2985-3004	coding	
100608	ATAATACATTATCTGTAA	230	3085-3104	coding	
100609	GGTCTCCCAATAAATACAT	231	3095-3114	coding	
100610	AGGATACCCCGTCTCCCAA	232	3105-3124	coding	
100611	TGGGTCCCCCAGGATACCCC	35	3115-3134	coding	
100612	GCTCCTACATTGGGTCCCCC	234	3125-3144	coding	
100613	AGCCAAGGCAGCTCCTACAT	235	3135-3154	coding	
100614	AACATGTCTGAGCCAAGGCA	236	3145-3164	coding	
100615	TTTCACGGAAGAACATGTCTG	237	3155-3174	coding	
100616	TCAGCTCCGTTTTCACGGAA	238	3165-3184	coding	
100617	AGCCTATTGTTCAGCTCCGT	239	3175-3194	coding	
100618	ACATGGGAACAGCCTATTGT	240	3185-3204	coding	
100619	ATCAAAAGAAGGCACAGAGG	241	3215-3234	coding	
100620	GTTTAGACAACCTTAATCAGA	242	3255-3274	coding	
100621	AATCAGCATTGTTTAGACAA	243	3265-3284	coding	
100622	TTGGTCACCAAATCAGCATT	244	3275-3294	coding	
100623	TGAGTGACAGTTGGTCACCA	245	3285-3304	coding	
100624	GGCTCAGCAATGAGTGACAG	246	3295-3314	coding	
100625	ATTACAGACACAACCTCCCT	247	3325-3344	coding	
100626	TAGTAGGGCGATTACAGACA	248	3335-3354	coding	
100627	CQCCACTGAATAGTACCCCG	249	3345-3364	coding	
100628	CTTTATTTCTCGCCACTGAA	250	3355-3374	coding	

¹Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethyl cytosines and 2'-deoxy cytosines residues are 5-methyl-cytosines; all linkages are phosphorothioate linkages.
²Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO. 1.

[0225] Several of these oligonucleotides were chosen for dose response studies. Cells were grown and treated as described in Example 3. Results are shown in Table 33. Each

oligonucleotide tested showed a dose response curve with maximum inhibition greater than 75%.

TABLE 33

Dose Response of PMA-Induced neoHK Cells to TNF-α Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
induced	—	—	—	100%	—
100235	149	intron 1	75 nM	77%	23%
"	"	"	150 nM	25%	75%
"	"	"	300 nM	6%	94%
100243	157	intron 1	75 nM	68%	32%
"	"	"	150 nM	15%	85%
"	"	"	300 nM	6%	94%
100263	157	intron 1	75 nM	79%	21%
"	"	"	150 nM	30%	70%
"	"	"	300 nM	23%	77%

Example 23

[0226] Optimization of Human TNF-α Antisense Oligonucleotide Chemistry

[0227] Analogs of oligonucleotides 21820 (SEQ ID NO. 66) and 21823 (SEQ ID NO. 69) were designed and synthesized to find an optimum gap size. The sequences and chemistries are shown in Table 34.

[0228] Dose response experiments were performed as described in Example 3. Results are shown in Table 35.

TABLE 34

Nucleotide Sequences of TNF-α Chimeric Backbone (deoxy gapped) Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET GENE	GENE	TARGET REGION
21820	ATATTTCCTCCGCTCTTTCTGT	66	1339-1358	intron 1	
28086	ATATTTCCTCCGCTCTTTCTGT	"	"	"	"
28087	ATATTTCCTCCGCTCTTTCTGT	"	"	"	"
21823	GTGTGCCAGACACCCATATCT	69	1399-1418	intron 1	
28088	GTGTGCCAGACACCCATATCT	"	"	"	"
28089	GTGTGCCAGACACCCATATCT	"	"	"	"

¹Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.
²Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO. 1.

[0229]

TABLE 35

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF-α Chimeric (deoxy gapped) Antisense Oligonucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition
induced	—	—	—	100%	—
13393	49	control	75 nM	150.0%	—
"	"	"	150 nM	135.0%	—
"	"	"	300 nM	90.0%	10.0%

TABLE 35-continued

Dose Response of 20 Hour PMA-Induced neoHK Cells to TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides (ASOs)						
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% protein Expression	% protein Inhibition	
21820	66	intron 1	75 nM	65.0%	35.0%	
"	"	"	150 nM	28.0%	72.0%	
"	"	"	300 nM	9.7%	90.3%	
28086	66	intron 1	75 nM	110.0%	—	
"	"	"	150 nM	83.0%	17.0%	
"	"	"	300 nM	61.0%	39.0%	
28087	66	intron 1	75 nM	127.0%	—	
"	"	"	150 nM	143.0%	—	
"	"	"	300 nM	147.0%	—	
21823	69	intron 1	75 nM	35.0%	65.0%	
"	"	"	150 nM	30.0%	70.0%	
"	"	"	300 nM	6.4%	93.6%	
28088	69	intron 1	75 nM	56.0%	44.0%	
"	"	"	150 nM	26.0%	74.0%	
"	"	"	300 nM	11.0%	89.0%	
28089	69	intron 1	75 nM	76.0%	24.0%	
"	"	"	150 nM	53.0%	47.0%	
"	"	"	300 nM	23.0%	77.0%	

Example 24

[0230] Screening of Additional TNF- α Chimeric (Deoxy Gapped) Antisense Oligonucleotides

[0231] Additional oligonucleotides targeting the major regions of TNF- α were synthesized. Oligonucleotides were synthesized as uniformly phosphorothioate chimeric oligonucleotides having regions of five 2'-O-methoxyethyl (2'-MOE) nucleotides at the wings and a central region of ten deoxynucleotides. Oligonucleotide sequences are shown in Table 36.

[0232] Oligonucleotides were screened as described in Example 5. Results are shown in Table 37.

TABLE 36

Nucleotide Sequence of Additional Human TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides						
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET CO-ORDINATES ²	GENE	GENE	REGION
104649	CTGAGGGAGCGTCTGCTGGC	251	0616-0635	5'-UTR		
104650	CCTTGCTGAGGGAGCGTCTG	252	0621-0640	5'-UTR		
104651	CTGGTCCCTCTGCTGTCCCTTG	253	0636-0655	5'-UTR		
104652	CCTCTGCTGTCTCTGCTGAG	254	0631-0650	5'-UTR		
104653	TTCTCTCCCTCTTAGCTGGT	255	0651-0670	5'-UTR		
104654	TCCCTCTTAGCTGGTCCCTCT	256	0646-0665	5'-UTR		
104655	TCGTAGGGTTGTTTTCAGGG	257	0686-0705	5'-UTR		
104656	CTGTAGTTGCTTCTCTCCCT	258	0661-0680	5'-UTR		
104657	ACCTGCCCTGGCAGCTTGTC	259	0718-0737	5'-UTR		
104658	GGATGTGGCTCTGAGGGTT	260	0696-0715	5'-UTR		
104659	TGTGAGAGGAAGAGAACCTG	261	0733-0752	5'-UTR		
104660	GAGGAAGAGAACCTGCTGG	262	0728-0747	5'-UTR		
104661	AGCCGTGGGTCTGATGTGA	263	0748-0767	5'-UTR		
104662	TGGCTCAGTATGTGAGAGGA	264	0743-0762	5'-UTR		
104663	GAGAGGGTGAAGCCGTGGGT	265	0758-0777	5'-UTR		
104664	TCATGGTGTCTTTCAGGG	266	0780-0799	AUG		
104665	CTTTCAGTGTCTCATGTGTC	267	0790-0809	AUG		
104666	TCATGCTTTTCAGTGTCTCATG	268	0795-0814	AUG		
104667	ACGTCCCGGATCATGCTTTC	269	0805-0824	coding		
104668	GCTCCACGTCCCGGATCATG	270	0810-0829	coding		
104669	TCCTCGGCCAGCTCCACGTC	271	0820-0839	coding		
104670	CGCGCTGCTCGGCCAGCTCC	272	0825-0844	coding		

TABLE 36-continued

Nucleotide Sequence of Additional Human TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides						
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET CO-ORDINATES ²	GENE	GENE	REGION
104671	AGGAACAAGCACCGCCTGGA	273	0874-0893	coding		
104672	CAAGCACCGCCTGGAGCCCT	274	0869-0888	coding		
104673	AAGGAGAAGAGGCTGAGGAA	275	0889-0908	coding		
104674	GAAGAGGCTGAGGAACAAGC	276	0884-0903	coding		
104675	CCTGCCACGATCAGGAAGGA	277	0904-0923	coding		
104676	CACGATCAGGAAGGAGAAGA	278	0899-0918	coding		
104677	AAGAGCGTGGTGGCCCTGTC	279	0919-0938	coding		
104678	CGTGGTGGCGCCTGCCACGA	280	0914-0933	coding		
104679	AAGTGCAGCAGGCAGAAGAG	281	0934-0953	coding		
104680	CAGCAGGCAGAAAGAGCGTGG	282	0929-0948	coding		
104681	GATCACTCCAAAGTGCAGCA	283	0944-0963	coding		
104682	GGCCCGATCACTCCAAAGTG	284	0949-0968	coding		
104683	GGGCCAGAGGGCTGATTAGA	285	1606-1625	coding		
104684	AGAGGGCTGATTAGAGAGAG	286	1601-1620	coding		
104685	GCTACAGGCTTGTCACTCGG	287	1839-1858	coding		
104686	CTGACTGCCTGGGCCAGAGG	288	1616-1635	E2/I2 ³		
104687	TACAACTATGGGTACAGGCT	289	1849-1868	coding		
104688	AGCCACTGGAGCTGCCCTCTC	290	2185-2204	coding		
104689	CTGGAGCTGCCCTCAGCTT	291	2180-2199	coding		
104690	TTGGCCCGCGGTTCAAGCA	292	2200-2219	coding		
104691	TTGGCCAGGAGGCATTGGGC	293	2215-2234	coding		
104692	CCGGCGGTTTCAAGCCACTGGA	294	2195-2214	coding		
104693	CTCAGCTCCACGCCATTGGC	295	2230-2249	coding		
104694	CAGGAGGGCATTTGGCCCGGC	296	2210-2229	coding		
104695	CTCCACGCCATTGGCCAGGA	297	2225-2244	coding		
104696	ACCAGCTGGTTATCTCTCAG	298	2245-2264	coding		
104697	CTGGTTATCTCTCAGCTCCA	299	2240-2259	coding		
104698	CCCTCTGATGGCACCACGAG	300	2260-2279	coding		
104699	TGATGGCACCACGAGCTGGT	301	2255-2274	coding		
104700	TAGATGAGGTACAGGCCCTC	302	2275-2294	coding		
104701	AAGAGGACCTGGGAGTAGAT	303	2290-2309	coding		
104702	GAGGTACAGGCCCTCTGATG	304	2270-2289	coding		
104703	CAGCCTTGGCCCTTGAAGAG	305	2305-2324	coding		
104704	GACCTGGGAGTAGATGAGGT	306	2285-2304	coding		
104705	TTGGCCCTTGAAGAGGACCT	307	2300-2319	coding		
104706	TGGTGTGGGTGAGGAGCACA	308	2337-2356	coding		
104707	CGGCGATGCGGCTGATGGTG	309	2352-2371	coding		
104708	TGGGTGAGGAGACATGGGT	310	2332-2351	coding		
104709	TGGTCTGGTAGGAGACGGCG	311	2367-2386	coding		
104710	ATGCGGCTGATGGTTGGGT	312	2347-2366	coding		
104711	AGAGGAGGTGTGACCTTGTC	313	2382-2401	coding		
104712	TGGTAGGAGACGGCGATCGG	314	2362-2381	coding		
104713	AGGTTGACCTTGGTCTGGTA	315	2377-2396	coding		
104714	GGCTCTTATGATGGCAGAGAG	316	2397-2416	coding		
104715	TCATACCGAGCTTGGCCCTC	317	2446-2465	coding		
104716	TTGATGGCAGAGAGGAGGTT	318	2392-2411	coding		
104717	CCCAGATAGATGGGCTCATA	93	2461-2480	coding		
104718	CCAGGGCTTGGCCTCAGCCC	94	2441-2460	coding		
104719	AGCTGGAAGACCCCTCCCAG	319	2476-2495	coding		
104720	ATAGATGGGCTCATACGAGG	320	2456-2475	coding		
104721	CGGTCACCCCTTCTCCAGCTG	321	2491-2510	coding		
104722	GAAGACCCCTCCCAGATAGA	322	2471-2490	coding		
104723	ATCTCAGCGCTGAGTCGGTC	26	2506-2525	coding		
104724	ACCCCTCTCCAGCTGGAAGA	323	2486-2505	coding		
104725	TAGTCGGGCGGATTCATCTC	90	2521-2540	coding		
104726	AGCGCTGAGTCGGTCAACCT	91	2501-2520	coding		
104727	TCGGCAAAGTCGAGATAGTC	324	2536-2554	coding		
104728	GGGCGGATTGATCTCAGCGC	325	2516-2535	coding		
104729	TAGACCTGCCAGAGCTCGGC	326	2551-2570	coding		
104730	AAAGTCGAGATAGTCGGGCC	327	2531-2550	coding		
104731	GCAATGATCCCAAAGTAGAC	328	2566-2585	coding		
104732	CTGCCCAGACTCGGCAAAAGT	329	2546-2565	coding		
104733	CGTCTCTCTCACAGGGCAAT	330	2581-2600	stop		
104734	GATCCCAAAGTAGACCTGCC	88	2561-2580	coding		
104735	GGAAGGTTGGATGTTTCGTCC	331	2596-2615	3'-UTR		
104736	TCCTCACAGGGCAATGATCC	332	2576-2595	stop		
104737	GTTGAGGGTGTCTGAAGGAG	333	2652-2671	3'-UTR		

TABLE 36-continued

Nucleotide Sequence of Additional Human TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	GENE
104738	GTTCGATGTTTCGTCCTCCTC	334	2591-2610	stop	
104739	TTTGAGCCAGAAGAGGTGTA	335	2667-2686	3'-UTR	
104740	GAGGCGTTTGGGAAGGTTGG	336	2606-2625	3'-UTR	
104741	GCCCCCAATTCCTTTTGA	337	2682-2701	3'-UTR	
104742	GCCAGAAGAGGTTGAGGGTG	338	2662-2681	3'-UTR	
104743	GGGTTCCGACCCTAACCCCC	339	2697-2716	3'-UTR	
104744	CAATTCCTCTTTTGGAGCCAG	340	2677-2696	3'-UTR	
104745	TAAAGTTCTAAGCTTGGGTT	341	2712-2731	3'-UTR	
104746	CCGACCCTAAGCCCCCAATT	342	2692-2711	3'-UTR	
104747	GGTGGTCTTGTGCTTAAAG	343	2727-2746	3'-UTR	
104748	TTCTAAGCTTGGGTTCCGAC	344	2707-2726	3'-UTR	
104749	CCCAGGTTTCGAAGTGGTGG	345	2742-2761	3'-UTR	
104750	TCTTGTGCTTAAAGTTCTA	346	2722-2741	3'-UTR	
104751	CACACATTCCTGAATCCCGA	347	2757-2776	3'-UTR	
104752	GTTTCGAAGTGGTGGTCTTG	348	2737-2756	3'-UTR	
104753	CTTCACGTGTCAGGCCACAC	349	2772-2791	3'-UTR	
104754	ATTCTCTGAATCCAGGTTTC	350	2752-2771	3'-UTR	
104755	TAGTGGTTCGCAGCAGCTTCA	351	2787-2806	3'-UTR	
104756	CCCAGTTTGAATTCCTTAGTG	352	2802-2821	3'-UTR	
104757	CTGTGCAAGGCCACACATTC	353	2767-2786	3'-UTR	
104758	GTGAGTTCTGGAGGCCCCAG	354	2817-2836	3'-UTR	
104759	GTTGCCAGCAGCTTCACTGTG	355	2782-2801	3'-UTR	
104760	TTTGAATTCCTAGTGGTTG	356	2797-2816	3'-UTR	
104761	AAGCTGTAGGCCCCAGTGAG	357	2832-2851	3'-UTR	
104762	TTCTGGAGGCCCCAGTTTGA	358	2812-2831	3'-UTR	
104763	AGATGTTCTAGGATCAAGCT	359	2847-2866	3'-UTR	
104764	TGGTCTCCAGATTCAGATG	360	2862-2881	3'-UTR	
104765	GTAGGCCCCAGTGAGTTCTG	361	2827-2846	3'-UTR	
104766	GAAACCAAGGCTCCCTGGTC	362	2877-2896	3'-UTR	
104767	TCAGGGATCAAGCTGTAGG	363	2842-2861	3'-UTR	
104768	TCCAGATTCAGGATGTCAGG	364	2857-2876	3'-UTR	
104769	GCAGCATTCCTGGCCAGAAC	365	2892-2911	3'-UTR	
104770	GTCTTCTCAAGTCTTGAGC	366	2907-2926	3'-UTR	
104771	AAAGGCTCCCTGGTCTCCAG	367	2872-2891	3'-UTR	
104772	CAATTTCTAGGTAGGTCATT	368	2922-2941	3'-UTR	
104773	ATTCTGGCCAGAACCAAGG	369	2887-2906	3'-UTR	
104774	CTCAAGTCTGCAGCATTTCT	370	2902-2921	3'-UTR	
104775	AAGGTCCACTTGTGTCAATT	371	2937-2956	3'-UTR	
104776	GAGAGAGGAAGGCTTAAGGT	372	2952-2971	3'-UTR	
104777	TCTAGGTGAGGTTCTCTCAA	373	2917-2936	3'-UTR	
104778	CCACTTGTGTCAATTTCTAG	374	2932-2951	3'-UTR	
104779	GTCGTGGAACATCTGGAGAG	375	2967-2986	3'-UTR	
104780	CCGTGTCTCAAGGAAGTCTG	376	2982-3001	3'-UTR	
104781	AGGAAGGCCCTAAGGTCAC	377	2947-2966	3'-UTR	
104782	GAGGGAGCTGGCTCCATGGG	378	3014-3033	3'-UTR	
104783	GAAACATCTGGAGAGAGGAA	379	2962-2981	3'-UTR	
104784	GTGCAACATATAAGTAGAGG	380	3029-3048	3'-UTR	
104785	TCTCAAGGAAGTCTGGAAC	381	2977-2996	3'-UTR	
104786	AATAAATAATCACAAGTGCA	382	3044-3063	3'-UTR	
104787	GGGCTGGGCTCCGTGCTCTA	383	2992-3011	3'-UTR	
104788	TACCCCGGTCTCCCAATAAA	384	3101-3120	3'-UTR	
104789	AACATAAATAGAGGGAGCTG	385	3024-3043	3'-UTR	
104790	TTGGGTCCCCCAGGATACCC	386	3116-3135	3'-UTR	
104791	ATAATCACAAGTGCAACAT	387	3039-3058	3'-UTR	
104792	AAGGCACTCCTACATTGGG	388	3131-3150	3'-UTR	
104793	CGGTCTCCCAAAATAATACA	389	3096-3115	3'-UTR	
104794	AAACATGTCGTAGCCAAAGG	390	3146-3165	3'-UTR	
104795	TCCCCAGGATACCCCGGTC	391	3111-3130	3'-UTR	
104796	AGCTCCTACATTGGGTCCTCC	392	3126-3145	3'-UTR	
104797	CTCCGTTTTCACGGAATAACA	393	3161-3180	3'-UTR	
104798	TGCTGAGCCCAAGCAGCTC	394	3141-3160	3'-UTR	
104799	CAGCCTATTGTTCTAGTCCG	395	3176-3195	3'-UTR	
104800	AGAGGACACAGAGGCCAGGG	396	3209-3228	3'-UTR	
104801	TTTTCACGGAAGCTGCTCT	397	3156-3175	3'-UTR	
104802	TATTGTTCAAGCTCCGTTTTC	398	3171-3190	3'-UTR	
104803	AAAAACATAATCAAAAGAA	399	3224-3243	3'-UTR	
104804	CAGATAAATTTTAAAAAA	400	3239-3258	3'-UTR	
104805	TACATGGGAACAGCCTATTG		3186-3205	3'-UTR	
104806	TTTAGACAACCTTAATCAGAT		3254-3273	3'-UTR	

TABLE 36-continued

Nucleotide Sequence of Additional Human TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	GENE
104807	CATAATCAAAAGAAGGCACA	401	3219-3238	3'-UTR	
104808	ACCAATCAGCATTGTTTAG	402	3269-3288	3'-UTR	
104809	AAATATTTTAAAAACATAA	403	3234-3253	3'-UTR	
104810	GAGTGACAGTTGGTCACCAA	404	3284-3303	3'-UTR	
104811	ACAACCTAATCAGATAAATA	405	3249-3268	3'-UTR	
104812	CAGAGGCTCAGCAATGAGTG	406	3299-3318	3'-UTR	
104813	ATCAGCATTGTTTAGACAAC	407	3264-3283	3'-UTR	
104814	AGGCGGATTACAGACACAC	408	3331-3350	3'-UTR	
104815	ACAGTTGGTCACCAAATCAG	409	3279-3298	3'-UTR	
104816	TCGCCACTGAATAGTAGGGC	410	3346-3365	3'-UTR	
104817	GCTCAGCAATGAGTGACAGT	411	3294-3313	3'-UTR	
104818	AGCAAACCTTATTCTCGCC	412	3361-3380	3'-UTR	
104819	GATTACAGACACAACCTCCC	413	3326-3345	3'-UTR	
104820	ACTGAATAGTAGGGCGATTA	414	3341-3360	3'-UTR	
104821	ACTTTATTCTCGCCACTGA	415	3356-3375	3'-UTR	
104822	GCTGTCCTTGCTGAGGGAGC	416	0626-0645	5'-UTR	
104823	CTTAGCTGGTCTCTGCTGT	417	0641-0660	5'-UTR	
104824	GTTGCTTCTCTCCCTCTTAG	418	0656-0675	5'-UTR	
104825	TGGCGTCTGAGGGTTGTTTT	419	0691-0710	5'-UTR	
104826	AGAGAACCTGCCTGGCAGCT	420	0723-0742	5'-UTR	
104827	CAGTATGTGAGAGGAAGAGA	421	0738-0757	5'-UTR	
104828	GGTGAAGCCGTGGGTCAGTA	422	0753-0772	5'-UTR	
104829	AGTGCTCATGGGTGCTCTTC	423	0785-0804	AUG	
104830	CCGGATCATGCTTTCACTGC	424	0800-0819	coding	
104831	GGCCAGCTCCACGTCCTCCGA	425	0815-0834	coding	
104832	GGCCCCCTGTCTTCTTGGG	426	0847-0866	coding	
104833	GGCTGAGGAACAGCACCAGC	427	0879-0898	coding	
104834	TCAGGAAGGAGAAGAGGCTG	428	0894-0913	coding	
104835	TGGCGCCTGCCACGCTGAGT	429	0909-0918	coding	
104836	GGCAGAGAGCGGTGGTGGCG	430	0924-0943	coding	
104837	CTCCAAAGTGCAGCAGGCGC	431	0939-0958	coding	
104838	GCTGATTAGAGAGAGGCTCCC	432	1596-1615	coding	
104839	TGCCTGGGCCAGAGGGCTGA	433	1611-1630	coding	
104840	GCTGCCCTCAGCTTGAGGG	434	2175-2194	coding	
104841	GGTTAGCCACTGGAGCTGC	435	2190-2209	coding	
104842	GGGCATTGGCCCGCGGCTC	436	2205-2224	coding	
104843	CGCCATTGGCCAGGAGGGCA	437	2220-2239	coding	
104844	TATCTCTCAGCTCCACGCCA	438	2235-2254	coding	
104845	GCACCACAGCTGGTTATCT	439	2250-2269	coding	
104846	ACAGGCCCTCTGATGGCACC	440	2265-2284	coding	
104847	GGGAGTAGATGAGGTACAGG	441	2280-2299	coding	
104848	CCTTGAAGAGGACCTGGGAG	442	2295-2314	coding	
104849	GAGGAGCACATGGGTGGAGG	443	2327-2346	coding	
104850	GCTGATGGTGTGGGTGAGGA	444	2342-2361	coding	
104851	GGAGACGGCGATGCGGCTGA	445	2357-2376	coding	
104852	GACCTTGGTCTGGTAGGAGA	446	2372-2391	coding	
104853	GGCAGAGAGGAGGTTGACCT	447	2387-2406	coding	
104854	GCTTGGCCTCAGCCCCCTCT	23	2436-2455	coding	
104855	TGGGCTCATACAGGGGCTTG	448	2451-2470	coding	
104856	CCCCCTCCAGATAGATGGGC	449	2466-2485	coding	
104857	TCTCCAGCTGGAAGACCCCT	92	2481-2500	coding	
104858	TGAGTCGGTCAACCTTCTCC	450	2496-2515	coding	
104859	GATTGATCTCAGCGCTGAGT	451	2511-2530	coding	
104860	CGAGATAGTCGGGCCGATTG	452	2526-2545	coding	
104861	CAGACTCGGCAAGTCCAGA	89	2541-2560	coding	
104862	CAAAAGTAGACCTGCCACAG	453	2556-2575	coding	
104863	ACAGGGCAATGATCCCAAG	454	2571-2590	stop	
104864	ATGTTCTGCTCTCTCACAGG	455	2586-2605	stop	
104865	GTTTGGGAAGGTTGGATGTT	456	2601-2620	3'-UTR	
104866	AAGAGGTTGAGGGTGTCTGA	457	2657-2676	3'-UTR	
104867	CTCTTTTTCAGGCCAAGAG	458	2672-2691	3'-UTR	
104868	CCTAAGCCCCCAATTCTCTT	459	2687-2706	3'-UTR	
104869	AGCTTGGGTTCCGACCTTAA	460	2702-2721	3'-UTR	
104870	TTGCTTAAAGTTCTAAGCTT	461	2717-2736	3'-UTR	
104871	GAAGTGGTGGTCTTGTGCT	462	2732-2751	3'-UTR	
104872	TGAATCCAGGTTTTCGAAGT	463	2747-2766	3'-UTR	
104873	CAGCCACACATTCTGTAAT	464	2762-2781	3'-UTR	
104874	CAGCACTTCACTGTGCAGGC	465	2777-2796	3'-UTR	
104875	ATTCTTAGTGGTGGCAGCA	466	2792-2811	3'-UTR	

TABLE 36-continued

Nucleotide Sequence of Additional Human TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET NUCLEOTIDE CO-ORDINATES ²	GENE TARGET REGION	GENE
104876	GAGGCC CCAGTTTGAATTCT	467	2807-2826	3'-UTR	
104877	CCCCAG TGAGTTCTGGAGGC	468	2822-2841	3'-UTR	
104878	GATCAA AGCTGTAGGCCCA	469	2837-2856	3'-UTR	
104879	ATTCC AGATGTCAGGGATCA	470	2852-2871	3'-UTR	
104880	CTCCCT GGTCTCCAGATTCC	471	2867-2886	3'-UTR	
104881	GGCCAGA ACCAAGGCTCCC	472	2882-2901	3'-UTR	
104882	GTCTCT GAGCATTTCTGGCCA	473	2897-2916	3'-UTR	
104883	GTGAGG TCTTCTCAAGTCCT	474	2912-2931	3'-UTR	
104884	TGTGT CAATTTCTAGGTGAG	475	2927-2946	3'-UTR	
104885	GGCCTA AGGTCCACTTGTGT	476	2942-2961	3'-UTR	
104886	ATCTGG AAGGAGGAAGGCCT	477	2957-2976	3'-UTR	
104887	AGGAAG TCTGGAAACATCTG	478	2972-2991	3'-UTR	
104888	GGGCTC CGTGTCTCAAGGAA	479	2987-3006	3'-UTR	
104889	AAATAG AGGGAGCTGGCTCC	480	3019-3038	3'-UTR	
104890	CACAAG TGCAAAACATAATA	481	3034-3053	3'-UTR	
104891	TCCCAA ATAAATACATTTCAT	482	3091-3110	3'-UTR	
104892	CAGGAT ACCCCGTCTCCCA	483	3106-3125	3'-UTR	
104893	CTACAT TGGTCTCCCCAGGA	484	3121-3140	3'-UTR	
104894	GAGCCA AGGCGAGCTCTACA	485	3136-3155	3'-UTR	
104895	ACGGAAA ACATGTCTGAGCC	486	3151-3170	3'-UTR	
104896	TTCAGC TCCGTTTTCACGGA	487	3166-3185	3'-UTR	
104897	GGGAAC AGCCTATTGTTTCAG	488	3181-3200	3'-UTR	
104898	TCAAAA GGAAGGCACAGAGGC	489	3214-3233	3'-UTR	
104899	TTTTAA AAAAACATAATCAAA	490	3229-3248	3'-UTR	
104900	TTAATC AGATAAATATTTTA	491	3244-3263	3'-UTR	
104901	CATTGT TTTAGACAACTTAAT	492	3259-3278	3'-UTR	
104902	TGGTC ACCACAAATCAGCATTG	493	3274-3293	3'-UTR	
104903	GCAAT GAGTGACAGTTGGTC	494	3289-3308	3'-UTR	
104904	GGGAGC AGAGGCTCAGCAAT	495	3304-3323	3'-UTR	
104905	ATAGTA GGGGGATTACAGAC	496	3336-3355	3'-UTR	
104906	ATTTCT CGCCACTGAATAGT	497	3351-3370	3'-UTR	

¹Emboldened residues are 2'-O-methoxyethyl residues (others are 2'-deoxy-). All 2'-O-methoxyethyl cytosines and 2'-deoxy cytosines residues are 5-methyl-cytosines; all linkages are phosphorothioate linkages.
²Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO. 1.
³This target region is an exon-intron junction and is represented in the form, for example, I1/E2, where I, followed by a number, refers to the intron number and E, followed by a number, refers to the exon number.

[0233]

TABLE 37

Inhibition of Human TNF- α mRNA Expression by Chimeric (deoxy gapped) Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
basal	—	—	0.0%	—
induced	—	—	100.0%	0.0%
28089	69	intron 1	42.3%	57.7%
104649	251	5'-UTR	165.6%	—
104650	252	5'-UTR	75.8%	24.2%
104651	253	5'-UTR	58.2%	41.8%
104652	254	5'-UTR	114.5%	—
104653	255	5'-UTR	84.9%	15.1%
104654	256	5'-UTR	80.8%	19.2%
104655	257	5'-UTR	94.3%	5.7%
104656	258	5'-UTR	78.4%	21.6%
104657	259	5'-UTR	87.4%	12.6%

TABLE 37-continued

Inhibition of Human TNF- α mRNA Expression by Chimeric (deoxy gapped) Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
104658	260	5'-UTR	213.4%	—
104659	261	5'-UTR	96.3%	3.7%
104660	262	5'-UTR	153.1%	—
104661	263	5'-UTR	90.0%	10.0%
104662	264	5'-UTR	33.3%	66.7%
104663	265	5'-UTR	144.2%	—
104664	266	AUG	76.3%	23.7%
104665	267	AUG	185.3%	—
104666	268	AUG	67.4%	32.6%
104667	269	Coding	94.3%	5.7%
104668	270	Coding	63.1%	36.9%
104669	271	Coding	50.8%	49.2%
104670	272	Coding	43.7%	56.3%
104671	273	Coding	52.2%	47.8%
104672	274	Coding	51.8%	48.2%
104673	275	Coding	102.3%	—
104674	276	Coding	135.4%	—
104675	277	Coding	83.1%	16.9%
104676	278	Coding	87.5%	12.5%
104677	279	Coding	53.6%	46.4%
104678	280	Coding	75.2%	24.8%
104679	281	Coding	114.0%	—
104680	282	Coding	142.5%	—
104681	283	Coding	58.5%	41.5%
104682	284	Coding	101.9%	—
104683	285	Coding	77.1%	22.9%
104684	286	Coding	61.0%	39.0%
104685	287	Coding	65.9%	34.1%
104686	288	E2/I2	59.2%	40.8%
104687	289	Coding	77.0%	23.0%
104688	290	Coding	40.1%	59.9%
104689	291	Coding	78.6%	21.4%
104690	292	Coding	90.9%	9.1%
104691	293	Coding	107.6%	—
104692	294	Coding	63.4%	36.6%
104693	295	Coding	74.1%	25.9%
104694	296	Coding	108.3%	—
104695	297	Coding	48.2%	51.8%
104696	298	Coding	120.3%	—
104697	299	Coding	45.0%	55.0%
104698	300	Coding	77.1%	22.9%
104699	301	Coding	143.7%	—
104700	302	Coding	96.1%	3.9%
104701	303	Coding	106.8%	—
104702	304	Coding	157.4%	—
104703	305	Coding	84.3%	15.7%
104704	306	Coding	182.8%	—
104705	307	Coding	125.1%	—
104706	308	Coding	81.8%	18.2%
104707	309	Coding	104.8%	—
104708	310	Coding	163.0%	—
104709	311	Coding	95.0%	5.0%
104710	312	Coding	182.1%	—
104711	313	Coding	82.1%	17.9%
104712	314	Coding	118.1%	—
104713	315	Coding	31.1%	68.9%
104714	316	Coding	90.5%	9.5%
104715	317	Coding	96.7%	3.3%
104716	318	Coding	180.7%	—
104717	93	Coding	71.6%	28.4%
104718	94	Coding	187.0%	—
104719	319	Coding	88.8%	11.2%
104720	320	Coding	166.5%	—
104721	321	Coding	65.0%	35.0%
104722	322	Coding	59.6%	40.4%
104723	26	Coding	90.1%	9.9%
104724	323	Coding	88.7%	11.3%
104725	90	Coding	94.7%	5.3%
104726	91	Coding	84.1%	15.9%

TABLE 37-continued

Inhibition of Human TNF- α mRNA Expression by Chimeric (deoxy gapped) Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
104727	324	Coding	125.3%	—
104728	325	Coding	221.7%	—
104729	326	Coding	102.4%	—
104730	327	Coding	151.6%	—
104731	328	Coding	102.2%	—
104732	329	Coding	53.2%	46.8%
104733	330	Stop	57.0%	43.0%
104734	88	Coding	119.2%	—
104735	331	3'-UTR	71.2%	28.8%
104736	332	Stop	79.0%	21.0%
104737	333	3'-UTR	87.4%	12.6%
104738	334	Stop	36.8%	63.2%
104739	335	3'-UTR	106.0%	—
104740	336	3'-UTR	130.9%	—
104741	337	3'-UTR	79.2%	20.8%
104742	338	3'-UTR	159.0%	—
104743	339	3'-UTR	96.1%	3.9%
104744	340	3'-UTR	129.9%	—
104745	341	3'-UTR	80.2%	19.8%
104746	342	3'-UTR	168.8%	—
104747	343	3'-UTR	89.2%	10.8%
104748	344	3'-UTR	103.4%	—
104749	345	3'-UTR	89.0%	11.0%
104750	346	3'-UTR	160.0%	—
104751	347	3'-UTR	60.1%	39.9%
104752	348	3'-UTR	72.4%	27.6%
104753	349	3'-UTR	70.0%	30.0%
104754	350	3'-UTR	115.6%	—
104755	351	3'-UTR	71.7%	28.3%
104756	352	3'-UTR	91.5%	8.5%
104757	353	3'-UTR	85.6%	14.4%
104758	354	3'-UTR	97.6%	2.4%
104759	355	3'-UTR	68.6%	31.4%
104760	356	3'-UTR	182.4%	—
104761	357	3'-UTR	110.9%	—
104762	358	3'-UTR	161.4%	—
104763	359	3'-UTR	102.0%	—
104764	360	3'-UTR	113.5%	—
104765	361	3'-UTR	154.8%	—
104766	362	3'-UTR	126.4%	—
104767	363	3'-UTR	116.1%	—
104768	364	3'-UTR	177.7%	—
104769	365	3'-UTR	89.8%	10.2%
104770	366	3'-UTR	94.3%	5.7%
104771	367	3'-UTR	191.2%	—
104772	368	3'-UTR	80.3%	19.7%
104773	369	3'-UTR	133.9%	—
104774	34	3'-UTR	94.8%	5.2%
104775	370	3'-UTR	80.6%	19.4%
104776	371	3'-UTR	90.1%	9.9%
104777	372	3'-UTR	84.7%	15.3%
104778	373	3'-UTR	121.3%	—
104779	374	3'-UTR	97.8%	2.2%
104780	375	3'-UTR	67.6%	32.4%
104781	376	3'-UTR	141.5%	—
104782	377	3'-UTR	96.5%	3.5%
104783	378	3'-UTR	153.2%	—
104784	379	3'-UTR	85.4%	14.6%
104785	380	3'-UTR	163.9%	—
104786	381	3'-UTR	82.9%	17.1%
104787	382	3'-UTR	89.7%	10.3%
104788	383	3'-UTR	103.9%	—
104789	384	3'-UTR	75.8%	24.2%
104790	385	3'-UTR	106.3%	—
104791	386	3'-UTR	165.3%	—
104792	387	3'-UTR	71.8%	28.2%
104793	388	3'-UTR	101.9%	—
104794	389	3'-UTR	70.7%	29.3%
104795	390	3'-UTR	68.8%	31.2%

TABLE 37-continued

Inhibition of Human TNF- α mRNA Expression by Chimeric (deoxy gapped) Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
104796	391	3'-UTR	93.4%	6.6%
104797	37	3'-UTR	131.7%	—
104798	392	3'-UTR	89.4%	10.6%
104799	393	3'-UTR	89.6%	10.4%
104800	394	3'-UTR	89.0%	11.0%
104801	395	3'-UTR	196.8%	—
104802	396	3'-UTR	189.3%	—
104803	397	3'-UTR	119.7%	—
104804	398	3'-UTR	102.4%	—
104805	399	3'-UTR	90.6%	9.4%
104806	400	3'-UTR	89.1%	10.9%
104807	401	3'-UTR	152.6%	—
104808	402	3'-UTR	96.8%	3.2%
104809	403	3'-UTR	178.8%	—
104810	404	3'-UTR	94.9%	5.1%
104811	405	3'-UTR	234.4%	—
104812	406	3'-UTR	114.3%	—
104813	407	3'-UTR	153.7%	—
104814	408	3'-UTR	86.3%	13.7%
104815	409	3'-UTR	153.9%	—
104816	410	3'-UTR	79.9%	20.1%
104817	411	3'-UTR	196.5%	—
104818	412	3'-UTR	94.3%	5.7%
104819	413	3'-UTR	143.3%	—
104820	414	3'-UTR	123.8%	—
104821	415	3'-UTR	129.2%	—
104822	416	5'-UTR	76.6%	23.4%
104823	417	5'-UTR	63.9%	36.1%
104824	418	5'-UTR	22.0%	78.0%
104825	419	5'-UTR	109.4%	—
104826	420	5'-UTR	45.2%	54.8%
104827	421	5'-UTR	68.9%	31.1%
104828	422	5'-UTR	70.9%	29.1%
104829	423	AUG	46.6%	53.4%
104830	424	Coding	55.0%	45.0%
104831	425	Coding	49.5%	50.5%
104832	426	Coding	106.0%	—
104833	427	Coding	23.7%	76.3%
104834	428	Coding	91.8%	8.2%
104835	429	Coding	72.3%	27.7%
104836	430	Coding	63.4%	36.6%
104837	431	Coding	31.0%	69.0%
104838	432	Coding	18.0%	82.0%
104839	433	Coding	67.9%	32.1%
104840	434	Coding	93.8%	6.2%
104841	435	Coding	43.0%	57.0%
104842	436	Coding	73.2%	26.8%
104843	437	Coding	48.1%	51.9%
104844	438	Coding	39.2%	60.8%
104845	439	Coding	37.6%	62.4%
104846	440	Coding	81.7%	18.3%
104847	441	Coding	50.8%	49.2%
104848	442	Coding	56.7%	43.3%
104849	443	Coding	51.8%	48.2%
104850	444	Coding	91.8%	8.2%
104851	445	Coding	93.9%	6.1%
104852	446	Coding	100.9%	—
104853	447	Coding	67.7%	32.3%
104854	23	Coding	11.0%	89.0%
104855	448	Coding	62.5%	37.5%
104856	449	Coding	67.8%	32.2%
104857	92	Coding	28.1%	71.9%
104858	450	Coding	76.2%	23.8%
104859	451	Coding	52.3%	47.7%
104860	452	Coding	93.6%	6.4%
104861	89	Coding	79.3%	20.7%
104862	453	Coding	63.1%	36.9%
104863	454	Stop	64.5%	35.5%
104864	455	Stop	43.2%	56.8%

TABLE 37-continued

Inhibition of Human TNF- α mRNA Expression by Chimeric (deoxy gapped) Phosphorothioate Oligodeoxynucleotides				
ISIS No:	SEQ ID NO:	GENE TARGET REGION	% mRNA EXPRESSION	% mRNA INHIBITION
104865	456	3'-UTR	83.1%	16.9%
104866	457	3'-UTR	49.4%	50.6%
104867	458	3'-UTR	49.5%	50.5%
104868	459	3'-UTR	89.6%	10.4%
104869	460	3'-UTR	21.4%	78.6%
104870	461	3'-UTR	118.0%	—
104871	462	3'-UTR	55.8%	44.2%
104872	463	3'-UTR	49.0%	51.0%
104873	464	3'-UTR	92.6%	7.4%
104874	465	3'-UTR	33.4%	66.6%
104875	466	3'-UTR	36.2%	63.8%
104876	467	3'-UTR	73.4%	26.6%
104877	468	3'-UTR	40.9%	59.1%
104878	469	3'-UTR	78.7%	21.3%
104879	470	3'-UTR	75.4%	24.6%
104880	471	3'-UTR	50.2%	49.8%
104881	472	3'-UTR	47.0%	53.0%
104882	473	3'-UTR	82.7%	17.3%
104883	474	3'-UTR	46.4%	53.6%
104884	475	3'-UTR	46.1%	53.9%
104885	476	3'-UTR	156.9%	—
104886	477	3'-UTR	102.4%	—
104887	478	3'-UTR	59.1%	40.9%
104888	479	3'-UTR	64.7%	35.3%
104889	480	3'-UTR	83.7%	16.3%
104890	481	3'-UTR	52.9%	47.1%
104891	482	3'-UTR	87.9%	12.1%
104892	483	3'-UTR	39.8%	60.2%
104893	484	3'-UTR	71.1%	28.9%
104894	485	3'-UTR	34.0%	66.0%
104895	486	3'-UTR	129.8%	—
104896	487	3'-UTR	57.6%	42.4%
104897	488	3'-UTR	49.6%	50.4%
104898	489	3'-UTR	71.7%	28.3%
104899	490	3'-UTR	101.5%	—
104900	491	3'-UTR	142.1%	—
104901	492	3'-UTR	55.9%	44.1%
104902	493	3'-UTR	85.3%	14.7%
104903	494	3'-UTR	46.0%	54.0%
104904	495	3'-UTR	59.9%	40.1%
104905	496	3'-UTR	47.2%	52.8%
104906	497	3'-UTR	56.3%	43.7%

[0234] Oligonucleotides 104662 (SEQ ID No: 264), 104669 (SEQ ID NO: 271), 104670 (SEQ ID NO: 272), 104688 (SEQ ID NO: 290), 104695 (SEQ ID NO: 297), 104697 (SEQ ID NO: 299), 104713 (SEQ ID NO: 315), 104738 (SEQ ID NO:334), 104824 (SEQ ID NO: 418), 104826 (SEQ ID NO: 420), 104829 (SEQ ID NO: 423), 104831 (SEQ ID NO: 425), 104833 (SEQ ID NO: 427), 104837 (SEQ ID NO: 431), 104838 (SEQ ID NO: 431), 104841 (SEQ ID NO: 435), 104843 (SEQ ID NO: 437), 104844 (SEQ ID NO: 438), 104845 (SEQ ID NO: 439), 104847 (SEQ ID NO: 441), 104854 (SEQ ID NO: 23), 104857 (SEQ ID NO: 92), 104864 (SEQ ID NO: 455), 104866 (SEQ ID NO: 457), 104867 (SEQ ID NO: 458), 104869 (SEQ ID NO: 460), 104872 (SEQ ID NO: 463), 104874 (SEQ ID NO: 465), 104875 (SEQ ID NO: 466), 104877 (SEQ ID NO: 468), 104880 (SEQ ID NO: 471), 104881 (SEQ ID NO: 472), 104883 (SEQ ID NO: 474), 104884 (SEQ ID NO: 475), 104892 (SEQ ID NO: 483), 104894 (SEQ ID NO: 485), 104897 (SEQ ID NO: 488), 104903 (SEQ ID NO: 494) and 104905 (SEQ ID NO: 496)

gave approximately 50% or greater reduction in TNF- α mRNA expression in this assay. Oligonucleotides 104713 (SEQ ID NO: 315), 104824 (SEQ ID NO: 418), 104833 (SEQ ID NO: 427), 104837 (SEQ ID NO: 431), 104838 (SEQ ID NO: 432), 104854 (SEQ ID NO: 23), 104857 (SEQ ID NO: 92), and 104869 (SEQ ID NO: 460) gave approximately 70% or greater reduction in TNF- α mRNA expression in this assay.

Example 25

[0235] Dose Response of Chimeric (Deoxy Gapped) Antisense Phosphorothioate Oligodeoxynucleotide Effects on TNF- α mRNA and Protein Levels

[0236] Several oligonucleotides from the initial screen were chosen for dose response assays. NeoHk cells were grown, treated and processed as described in Example 3. LIPOFECTIN7 was added at a ratio of 3 μ g/ml per 100 nM of oligonucleotide. The control included LIPOFECTIN7 at a concentration of 9 μ g/ml.

[0237] The human promonocytic leukaemia cell line, THP-1 (American Type Culture Collection, Manassas, Va.) was maintained in RPMI 1640 growth media supplemented with 10% fetal serum (FCS; Life Technologies, Rockville, Md.).

[0238] A total of 8 $\times$ 10⁵ cell were employed for each treatment by combining 50 μ l of cell suspension in OPTI-MENT™, 1% FBS with oligonucleotide at the indicated concentrations to reach a final volume of 100 μ l with OPTIMENT™, 1% FBS. Cells were then transferred to a 1 mm electroporation cuvette and electroporated using an Electroporation Manipulator 600 instrument (Biotechnologies and Experimental Research, INC.) employing 90 V, 1000 μ F, at 13 Ω . Electroporated cells were then transferred to 24 well plates. 400 μ l of RPMI 1640, 10% FCS was added to the cells and the cells were allowed to recover for 6 hrs. Cells were then induced with LPS at a final concentration of 100 ng/ml for 2 hours. RNA was isolated and processed as described in Example 3. Results with NeoHK cells are shown in Table 38 for mRNA, and Table 39 for protein. Results with THP-1 cells are shown in Table 40.

[0239] Most of the oligonucleotides tested showed dose response effects with a maximum inhibition of mRNA greater than 70% and a maximum inhibition of protein greater than 85%.

TABLE 38

Dose Response of NeoHK Cells to TNF- α chimeric (deoxy gapped) Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
induced	—	—	—	100%	—
16798	128	coding	30 nM	87%	13%
"	"	"	100 nM	129%	—
"	"	"	300 nM	156%	—
21823	69	intron 1	30 nM	82%	18%
"	"	"	100 nM	90%	10%
"	"	"	300 nM	59%	41%
28088	68	intron 1	30 nM	68%	32%
"	"	"	100 nM	43%	57%
"	"	"	300 nM	42%	58%
28089	69	intron 1	30 nM	59%	41%
"	"	"	100 nM	44%	56%
"	"	"	300 nM	38%	62%

TABLE 38-continued

Dose Response of NeoHK Cells to TNF- α chimeric (deoxy gapped) Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
104697	299	coding	30 nM	60%	40%
"	"	"	100 nM	45%	55%
"	"	"	300 nM	27%	73%
104777	372	3' -UTR	30 nM	66%	34%
"	"	"	100 nM	55%	45%
"	"	"	300 nM	43%	57%

[0240]

TABLE 39

Dose Response of NeoHK Cells to TNF- α Chimeric (deoxy gapped) Antisense Oligonucleotides					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% Protein Expression	% Protein Inhibition
induced	—	—	—	100.0%	—
16798	128	coding	30 nM	115.0%	—
"	"	"	100 nM	136.0%	—
"	"	"	300 nM	183.0%	—
28089	69	intron 1	30 nM	87.3%	12.7%
"	"	"	100 nM	47.4%	52.6%
"	"	"	300 nM	22.8%	77.0%
104681	283	coding	30 nM	91.3%	8.7%
"	"	"	100 nM	62.0%	38.0%
"	"	"	300 nM	28.5%	71.5%
104697	299	coding	30 nM	87.1%	12.9%
"	"	"	100 nM	59.6%	40.4%
"	"	"	300 nM	29.1%	70.9%
104838	432	coding	30 nM	91.9%	8.1%
"	"	"	100 nM	56.9%	43.1%
"	"	"	300 nM	14.8%	85.2%
104854	23	coding	30 nM	64.4%	35.6%
"	"	"	100 nM	42.3%	57.7%
"	"	"	300 nM	96.1%	3.9%
104869	460	3' -UTR	30 nM	88.9%	11.1%
"	"	"	100 nM	56.8%	43.2%
"	"	"	300 nM	42.3%	57.7%

[0241]

TABLE 40

Dose Response of LPS-Induced THP-1 Cells to Chimeric (deoxy gapped) TNF- α Antisense Phosphorothioate Oligodeoxynucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
induced	—	—	—	100%	—
16798	128	coding	1 μ M	102%	—
"	"	"	3 μ M	87%	13%
"	"	"	10 μ M	113%	—
"	"	"	30 μ M	134%	—
28089	69	intron 1	1 μ M	39%	61%
"	"	"	3 μ M	79%	21%
"	"	"	10 μ M	91%	9%
"	"	"	30 μ M	63%	37%
104697	299	coding	1 μ M	99%	1%
"	"	"	3 μ M	96%	4%
"	"	"	10 μ M	92%	8%
"	"	"	30 μ M	52%	48%

TABLE 40-continued

Dose Response of LPS-Induced THP-1 Cells to Chimeric (deoxy gapped) TNF- α Antisense Phosphorothioate Oligodeoxynucleotides (ASOs)					
ISIS #	SEQ ID NO:	ASO Gene Target	Dose	% mRNA Expression	% mRNA Inhibition
104838	432	coding	1 μ M	31%	69%
"	"	"	3 μ M	20%	80%
"	"	"	10 μ M	15%	85%
"	"	"	30 μ M	7%	93%
104854	23	coding	1 μ M	110%	—
"	"	"	3 μ M	90%	10%
"	"	"	10 μ M	95%	5%
"	"	"	30 μ M	61%	39%

Example 26

[0242] Further Optimization of Human TNF- α Antisense Oligonucleotide Chemistry

[0243] Additional analogs of TNF- α oligonucleotides were designed and synthesized to find an optimum gap size. The sequences and chemistries are shown in Table 36.

[0244] Dose response experiments are performed as described in Example 3.

TABLE 41

Nucleotide Sequences of TNF- α Chimeric Backbone (deoxy gapped) Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID	TARGET NUCLEOTIDE NO: CO-ORDINATES ²	GENE TARGET REGION	
110554	GCTGATTAGAGAGAGGTCCC	432	104838	analog	
110555	GCTGATTAGAGAGAGGTCCC	"	"	"	
110556	GCTGATTAGAGAGAGGTCCC	"	"	"	
110557	GCTGATTAGAGAGAGGTCCC	"	"	"	
110583	GCTGATTAGAGAGAGGTCCC	"	"	"	
110558	CTGATTAGAGAGAGGTCCC	498	1596-1614	coding	
110559	CTGATTAGAGAGAGGTCCC	"	"	"	
110560	CTGATTAGAGAGAGGTCCC	"	"	"	
110561	CTGATTAGAGAGAGGTCCC	"	"	"	
110562	CTGATTAGAGAGAGGTCCC	"	"	"	
110563	CTGATTAGAGAGAGGTCCC	"	"	"	
110564	CTGATTAGAGAGAGGTCCC	"	"	"	
110565	CTGATTAGAGAGAGGTCCC	"	"	"	
110566	CTGATTAGAGAGAGGTCCC	"	"	"	
110567	CTGATTAGAGAGAGGTCCC	"	"	"	
110584	CTGATTAGAGAGAGGTCCC	"	"	"	
108371	CTGATTAGAGAGAGGTCC	499	1597-1614	coding	
110568	CTGATTAGAGAGAGGTCC	"	"	"	
110569	CTGATTAGAGAGAGGTCC	"	"	"	
110570	CTGATTAGAGAGAGGTCC	"	"	"	
110585	CTGATTAGAGAGAGGTCC	"	"	"	
110571	CTGGTTATCTCTCAGCTCCA	299	104697	analog	
110572	CTGGTTATCTCTCAGCTCCA	"	"	"	
110573	CTGGTTATCTCTCAGCTCCA	"	"	"	
110586	CTGGTTATCTCTCAGCTCCA	"	"	"	
110574	GATCACTCCAAAGTGCAGCA	283	104681	analog	
110575	GATCACTCCAAAGTGCAGCA	"	"	"	
110576	GATCACTCCAAAGTGCAGCA	"	"	"	
110587	GATCACTCCAAAGTGCAGCA	"	"	"	
110577	AGCTTGGGTTCCGACCCCTAA	460	104689	analog	
110578	AGCTTGGGTTCCGACCCCTAA	"	"	"	
110579	AGCTTGGGTTCCGACCCCTAA	"	"	"	
110588	AGCTTGGGTTCCGACCCCTAA	"	"	"	
110580	AGGTTGACCTTGGTCTGGTA	315	104713	analog	
110581	AGGTTGACCTTGGTCTGGTA	"	"	"	

TABLE 41-continued

Nucleotide Sequences of TNF-α Chimeric Backbone (deoxy gapped) Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET CO-ORDINATES ²	GENE TARGET REGION	GENE
110582	AGGTTGACCTTGGTCTGGTA	"	"	"	"
110589	AGGTTGACCTTGGTCTGGTA	"	"	"	"
110637	GTGTGCCAGACACCCCTATCT	69	21823	analog	"
110651	GTGTGCCAGACACCCCTATCT	"	"	"	"
110665	GTGTGCCAGACACCCCTATCT	"	"	"	"
110679	GTGTGCCAGACACCCCTATCT	"	"	"	"
110693	GTGTGCCAGACACCCCTATCT	"	"	"	"
110707	GTGTGCCAGACACCCCTATCT	"	"	"	"
110590	TGAGTGCTCTCTGTGTGCCA	500	1411-1430	intron	"
110597	TGAGTGCTCTCTGTGTGCCA	"	"	1	"
110604	TGAGTGCTCTCTGTGTGCCA	"	"	"	"
110611	TGAGTGCTCTCTGTGTGCCA	"	"	"	"
110618	TGAGTGCTCTCTGTGTGCCA	"	"	"	"
110625	TGAGTGCTCTCTGTGTGCCA	"	"	"	"
110591	GAGTGCTCTCTGTGTGCCAG	501	1410-1429	intron	"
110598	GAGTGCTCTCTGTGTGCCAG	"	"	1	"
110605	GAGTGCTCTCTGTGTGCCAG	"	"	"	"
110612	GAGTGCTCTCTGTGTGCCAG	"	"	"	"
110619	GAGTGCTCTCTGTGTGCCAG	"	"	"	"
110626	GAGTGCTCTCTGTGTGCCAG	"	"	"	"
110592	AGTGTCTTCTGTGTGCCAGA	144	100181	analog	"
110599	AGTGTCTTCTGTGTGCCAGA	"	"	"	"
110606	AGTGTCTTCTGTGTGCCAGA	"	"	"	"
110613	AGTGTCTTCTGTGTGCCAGA	"	"	"	"
110620	AGTGTCTTCTGTGTGCCAGA	"	"	"	"
110627	AGTGTCTTCTGTGTGCCAGA	"	"	"	"
110593	GTGTCTTCTGTGTGCCAGAC	145	100182	analog	"
110600	GTGTCTTCTGTGTGCCAGAC	"	"	"	"
110607	GTGTCTTCTGTGTGCCAGAC	"	"	"	"
110614	GTGTCTTCTGTGTGCCAGAC	"	"	"	"
110621	GTGTCTTCTGTGTGCCAGAC	"	"	"	"
110628	GTGTCTTCTGTGTGCCAGAC	"	"	"	"
110594	TGTCTTCTGTGTGCCAGACA	146	100183	analog	"
110601	TGTCTTCTGTGTGCCAGACA	"	"	"	"
110608	TGTCTTCTGTGTGCCAGACA	"	"	"	"
110615	TGTCTTCTGTGTGCCAGACA	"	"	"	"
110622	TGTCTTCTGTGTGCCAGACA	"	"	"	"
110629	TGTCTTCTGTGTGCCAGACA	"	"	"	"
110595	GTCTTCTGTGTGCCAGACAC	147	100184	analog	"
110602	GTCTTCTGTGTGCCAGACAC	"	"	"	"
110609	GTCTTCTGTGTGCCAGACAC	"	"	"	"
110616	GTCTTCTGTGTGCCAGACAC	"	"	"	"
110623	GTCTTCTGTGTGCCAGACAC	"	"	"	"
110630	GTCTTCTGTGTGCCAGACAC	"	"	"	"
110596	TCTTCTGTGTGCCAGACACC	148	100185	analog	"
110603	TCTTCTGTGTGCCAGACACC	"	"	"	"
110610	TCTTCTGTGTGCCAGACACC	"	"	"	"
110617	TCTTCTGTGTGCCAGACACC	"	"	"	"
110624	TCTTCTGTGTGCCAGACACC	"	"	"	"
110631	TCTTCTGTGTGCCAGACACC	"	"	"	"
110632	CTTCTGTGTGCCAGACACC	149	100186	analog	"
110646	CTTCTGTGTGCCAGACACC	"	"	"	"
110660	CTTCTGTGTGCCAGACACC	"	"	"	"
110674	CTTCTGTGTGCCAGACACC	"	"	"	"
110688	CTTCTGTGTGCCAGACACC	"	"	"	"
110702	CTTCTGTGTGCCAGACACC	"	"	"	"
110633	TTCTGTGTGCCAGACACCCCT	150	100187	analog	"
110647	TTCTGTGTGCCAGACACCCCT	"	"	"	"
110661	TTCTGTGTGCCAGACACCCCT	"	"	"	"
110675	TTCTGTGTGCCAGACACCCCT	"	"	"	"
110689	TTCTGTGTGCCAGACACCCCT	"	"	"	"
110703	TTCTGTGTGCCAGACACCCCT	"	"	"	"
110634	TCTGTGTGCCAGACACCCCTA	151	100188	analog	"
110648	TCTGTGTGCCAGACACCCCTA	"	"	"	"
110662	TCTGTGTGCCAGACACCCCTA	"	"	"	"
110676	TCTGTGTGCCAGACACCCCTA	"	"	"	"
110690	TCTGTGTGCCAGACACCCCTA	"	"	"	"

TABLE 41-continued

Nucleotide Sequences of TNF-α Chimeric Backbone (deoxy gapped) Oligonucleotides					
ISIS NO.	NUCLEOTIDE SEQUENCE ¹ (5' -> 3')	SEQ ID NO:	TARGET CO-ORDINATES ²	GENE TARGET REGION	GENE
110704	TCTGTGTGCCAGACACCCCTA	"	"	"	"
110635	CTGTGTGCCAGACACCCCTAT	152	100189	analog	"
110649	CTGTGTGCCAGACACCCCTAT	"	"	"	"
110663	CTGTGTGCCAGACACCCCTAT	"	"	"	"
110677	CTGTGTGCCAGACACCCCTAT	"	"	"	"
110691	CTGTGTGCCAGACACCCCTAT	"	"	"	"
110705	CTGTGTGCCAGACACCCCTAT	"	"	"	"
110636	TGTGTGCCAGACACCCCTATC	153	100190	analog	"
110650	TGTGTGCCAGACACCCCTATC	"	"	"	"
110664	TGTGTGCCAGACACCCCTATC	"	"	"	"
110678	TGTGTGCCAGACACCCCTATC	"	"	"	"
110692	TGTGTGCCAGACACCCCTATC	"	"	"	"
110706	TGTGTGCCAGACACCCCTATC	"	"	"	"
110638	TGTGTGCCAGACACCCCTATCT	154	100191	analog	"
110652	TGTGTGCCAGACACCCCTATCT	"	"	"	"
110666	TGTGTGCCAGACACCCCTATCT	"	"	"	"
110680	TGTGTGCCAGACACCCCTATCT	"	"	"	"
110694	TGTGTGCCAGACACCCCTATCT	"	"	"	"
110708	TGTGTGCCAGACACCCCTATCT	"	"	"	"
110639	GTGCCAGACACCCCTATCTTC	155	100192	analog	"
110653	GTGCCAGACACCCCTATCTTC	"	"	"	"
110667	GTGCCAGACACCCCTATCTTC	"	"	"	"
110681	GTGCCAGACACCCCTATCTTC	"	"	"	"
110695	GTGCCAGACACCCCTATCTTC	"	"	"	"
110709	GTGCCAGACACCCCTATCTTC	"	"	"	"
110640	TGCCAGACACCCCTATCTTCT	156	100193	analog	"
110654	TGCCAGACACCCCTATCTTCT	"	"	"	"
110668	TGCCAGACACCCCTATCTTCT	"	"	"	"
110682	TGCCAGACACCCCTATCTTCT	"	"	"	"
110696	TGCCAGACACCCCTATCTTCT	"	"	"	"
110710	TGCCAGACACCCCTATCTTCT	"	"	"	"
110641	GCCAGACACCCCTATCTTCTT	157	100194	analog	"
110655	GCCAGACACCCCTATCTTCTT	"	"	"	"
110669	GCCAGACACCCCTATCTTCTT	"	"	"	"
110683	GCCAGACACCCCTATCTTCTT	"	"	"	"
110697	GCCAGACACCCCTATCTTCTT	"	"	"	"
110711	GCCAGACACCCCTATCTTCTT	"	"	"	"
110642	CCAGACACCCCTATCTTCTTC	158	100195	analog	"
110656	CCAGACACCCCTATCTTCTTC	"	"	"	"
110670	CCAGACACCCCTATCTTCTTC	"	"	"	"
110684	CCAGACACCCCTATCTTCTTC	"	"	"	"
110698	CCAGACACCCCTATCTTCTTC	"	"	"	"
110712	CCAGACACCCCTATCTTCTTC	"	"	"	"
110643	CAGACACCCCTATCTTCTTCT	159	100196	analog	"
110657	CAGACACCCCTATCTTCTTCT	"	"	"	"
110671	CAGACACCCCTATCTTCTTCT	"	"	"	"
110685	CAGACACCCCTATCTTCTTCT	"	"	"	"
110699	CAGACACCCCTATCTTCTTCT	"	"	"	"
110713	CAGACACCCCTATCTTCTTCT	"	"	"	"
110644	AGACACCCCTATCTTCTTCTC	160	100197	analog	"
110658	AGACACCCCTATCTTCTTCTC	"	"	"	"
110672	AGACACCCCTATCTTCTTCTC	"	"	"	"
110686	AGACACCCCTATCTTCTTCTC	"	"	"	"
110700	AGACACCCCTATCTTCTTCTC	"	"	"	"
110714	AGACACCCCTATCTTCTTCTC	"	"	"	"
110645	GACACCCCTATCTTCTTCTCT	161	100198	analog	"
110659	GACACCCCTATCTTCTTCTCT	"	"	"	"
110673	GACACCCCTATCTTCTTCTCT	"	"	"	"
110687	GACACCCCTATCTTCTTCTCT	"	"	"	"
110701	GACACCCCTATCTTCTTCTCT	"	"	"	"
110715	GACACCCCTATCTTCTTCTCT	"	"	"	"

¹Emboldened residues are 2'-methoxyethoxy residues (others are 2'-deoxy-). All 2'-methoxyethoxy cytidines and 2'-deoxycytidines are 5-methyl-cytidines; all linkages are phosphorothioate linkages.
²Co-ordinates from Genbank Accession No. X02910, locus name "HSTNFA", SEQ ID NO. 1.

Example 26

[0245] Effect of TNF- α Antisense Oligonucleotides in TNF- α Transgenic Mouse Models

[0246] The effect of TNF- α antisense oligonucleotides is studied in transgenic mouse models of human diseases. Such experiments can be performed through contract laboratories (e.g., The Laboratory of Molecular Genetics at The Hellenic Pasteur Institute, Athens, Greece) where such transgenic mouse models are available. Such models are available for testing human oligonucleotides in arthritis (Keffler, J., et al., *EMBO J.*, 1991, 10, 4025-4031) and multiple sclerosis (Akassoglou et al., *J. Immunol.*, 1997, 158, 438-445) mod-

els. A model for inflammatory bowel disease is available for testing mouse oligonucleotides (Kontoyiannis et al., *Immunity*, 1999, 10, 387-398).

[0247] Briefly, litters of the appropriate transgenic mouse strain are collected and weighed individually. Twice weekly from birth, oligonucleotide in saline is administered intraperitoneally or intravenously. Injections continue for 7 weeks. Each week the animals are scored for manifestations of the appropriate disease. After the final treatment, the mice are sacrificed and histopathology is performed for indicators of disease as indicated in the references cited for each model.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 503

<210> SEQ ID NO 1
<211> LENGTH: 3634
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (796..981,1589..1634,1822..1869,2171..2592)
<221> NAME/KEY: exon
<222> LOCATION: (615)..(981)
<221> NAME/KEY: intron
<222> LOCATION: (982)..(1588)
<221> NAME/KEY: exon
<222> LOCATION: (1589)..(1634)
<221> NAME/KEY: intron
<222> LOCATION: (1635)..(1821)
<221> NAME/KEY: exon
<222> LOCATION: (1822)..(1869)
<221> NAME/KEY: intron
<222> LOCATION: (1870)..(2070)
<221> NAME/KEY: exon
<222> LOCATION: (2171)..(3381)
<300> PUBLICATION INFORMATION:
<301> AUTHORS: Nedwin, G.E.
Naylor, S.L.
Sakaguchi, A.Y.
Smith, D.
Jarrett-Nedwin, J.
Pennica, D.
Goeddel, D.V.
Gray, P.W.
<302> TITLE: Human lymphotoxin and tumor necrosis factor genes:
structure, homology and chromosomal localization
<303> JOURNAL: Nucleic Acids Res.
<304> VOLUME: 13
<305> ISSUE: 17
<306> PAGES: 6361-6373
<307> DATE: 1985-09-11
<308> DATABASE ACCESSION NUMBER: X02910 Genbank
<309> DATABASE ENTRY DATE: 1997-02-17

<400> SEQUENCE: 1

gaattccggg tgatttcact cccggctgtc caggcttgct ctgctacccc acccagcctt	60
tcctgaggcc tcaagcctgc caccaagccc ccagctcctt ctccccgcag gacccaaaca	120
caggcctcag gactcaacac agcttttccc tccaaccctg ttctctcccc tcaacggact	180
cagotttctg aagccctcc cagttctagt tctatctttt tctgcaccc tgtctggaag	240
ttagaaggaa acagaccaca gacctggtcc ccaaaagaaa tggagggaat aggttttgag	300
gggcatgggg acgggggttca gcctccaggg tcctacacac aaatcagtcg gtggcccgag	360

-continued

agacccccct cggaatcgga gcagggagga tggggagtgt gaggggtatc ctgtatgctt	420
gtgtgtcccc aactttccaa atccccgccc ccgcgatgga gaagaaccg agacagaagg	480
tgcagggccc actaccgctt cctccagatg agtcctatggg tttctccacc aaggaagttt	540
tccgctgtgtt gaatgattct ttccccgccc tcctctcgcc ccagggacat ataaaggcag	600
ttgttggcac acccagccag cagacgctcc ctacagcaagg acagcagagg accagctaag	660
agggagagaa gcaactacag accccccctg aaaacaaccc tcagacgcca catccccctga	720
caagctgcca ggcaggttct cttcctctca catactgacc cacggttcca ccctctctcc	780
cctggaaagg acacc atg agc act gaa agc atg atc cgg gac gtg gag ctg	831
Met Ser Thr Glu Ser Met Ile Arg Asp Val Glu Leu	
1 5 10	
gcc gag gag gcg ctc ccc aag aag aca ggg ggg ccc cag ggc tcc agg	879
Ala Glu Glu Ala Leu Pro Lys Lys Thr Gly Gly Pro Gln Gly Ser Arg	
15 20 25	
cgg tgc ttg ttc ctc agc ctc ttc tcc ttc ctg atc gtg gca ggc gcc	927
Arg Cys Leu Phe Leu Ser Leu Phe Ser Phe Leu Ile Val Ala Gly Ala	
30 35 40	
acc acg ctc ttc tgc ctg ctg cac ttt gga gtg atc ggc ccc cag agg	975
Thr Thr Leu Phe Cys Leu Leu His Phe Gly Val Ile Gly Pro Gln Arg	
45 50 55 60	
gaa gag gtgagtgcct ggccagcctt catccactct cccacccaag gggaaatgag	1031
Glu Glu	
agacgcaaga gaggggagaga gatgggatgg gtgaaagatg tgcgctgata gggagggatg	1091
agagagaaaa aaacatggag aaagacgggg atgcagaaag agatgtggca agagatgggg	1151
aagagagaga gagaaagatg gagagacagg atgtctggca catggaaggt gctcactaag	1211
tgtgtatgga gtgaatgaat gaatgaatga atgaacaagc agatatataa ataagatatg	1271
gagacagatg tggggtgtga gaagagagat gggggaagaa acaagtgata tgaataaaga	1331
tggtgagaca gaaagacggg gaaatatgac agctaaggag agagatgggg gagataagga	1391
gagaagaaga taggggtgtct ggcacacaga agacactcag gaaaagagct gttgaatgct	1451
ggaagggtgaa tacacagatg aatggagaga gaaaaccaga cacctcaggg ctaagagcgc	1511
aggccagaca ggcagccagc tgctcctcct ttaagggtga ctccctcgat gttaaccatt	1571
ctccttctcc ccaacag ttc ccc agg gac ctc tct cta atc agc cct ctg	1621
Phe Pro Arg Asp Leu Ser Leu Ile Ser Pro Leu	
65 70	
gcc cag gca gtc agtaagtgtc tccaaacctc tttcctaatt ctgggtttgg	1673
Ala Gln Ala Val	
75	
gtttgggggt agggttagta ccggtatgga agcagtgggg gaaatttaaa gttttggtct	1733
tgggggagga tggatggagg tgaaagtagg ggggtatttt ctagggaagtt taagggtctc	1793
agctttttct tttctctctc ctcttca gga tca tct tct cga acc ccg agt gac	1847
Arg Ser Ser Ser Arg Thr Pro Ser Asp	
80 85	
aag cct gta gcc cat gtt gta ggtaagagct ctgaggatgt gtcttggaac	1898
Lys Pro Val Ala His Val Val	
90	
ttggagggct aggatttggg gattgaagcc cggctgatgg taggcagaac ttggagacaa	1958
tgtgagaagg actcgtgag ctcaagggaa gggtaggagga acagcacagg ccttagtggg	2018
atactcagaa cgctcatggcc aggtgggatg tgggatgaca gacagagagg acaggaaccg	2078

-continued

gatgtgggggt gggcagagct cgagggccag gatgtggaga gtgaaccgac atggccacac	2138
tgactctcct ctccctctct ccctccctcc a gca aac cct caa gct gag ggg	2190
Ala Asn Pro Gln Ala Glu Gly	
95 100	
cag ctc cag tgg ctg aac cgc cgg gcc aat gcc ctc ctg gcc aat ggc	2238
Gln Leu Gln Trp Leu Asn Arg Arg Ala Asn Ala Leu Leu Ala Asn Gly	
105 110 115	
gtg gag ctg aga gat aac cag ctg gtg gtg cca tca gag ggc ctg tac	2286
Val Glu Leu Arg Asp Asn Gln Leu Val Val Pro Ser Glu Gly Leu Tyr	
120 125 130	
ctc atc tac tcc cag gtc ctc ttc aag ggc caa ggc tgc ccc tcc acc	2334
Leu Ile Tyr Ser Gln Val Leu Phe Lys Gly Gln Gly Cys Pro Ser Thr	
135 140 145	
cat gtg ctc ctc acc cac acc atc agc cgc atc gcc gtc tcc tac cag	2382
His Val Leu Leu Thr His Thr Ile Ser Arg Ile Ala Val Ser Tyr Gln	
150 155 160	
acc aag gtc aac ctc ctc tct gcc atc aag agc ccc tgc cag agg gag	2430
Thr Lys Val Asn Leu Leu Ser Ala Ile Lys Ser Pro Cys Gln Arg Glu	
165 170 175 180	
acc cca gag ggg gct gag gcc aag ccc tgg tat gag ccc atc tat ctg	2478
Thr Pro Glu Gly Ala Glu Ala Lys Pro Trp Tyr Glu Pro Ile Tyr Leu	
185 190 195	
gga ggg gtc ttc cag ctg gag aag ggt gac cga ctc agc gct gag atc	2526
Gly Gly Val Phe Gln Leu Glu Lys Gly Asp Arg Leu Ser Ala Glu Ile	
200 205 210	
aat cgg ccc gac tat ctc gac ttt gcc gag tct ggg cag gtc tac ttt	2574
Asn Arg Pro Asp Tyr Leu Asp Phe Ala Glu Ser Gly Gln Val Tyr Phe	
215 220 225	
ggg atc att gcc ctg tga ggaggacgaa catccaacct tcccaaacgc	2622
Gly Ile Ile Ala Leu	
230	
ctcccttgcc ccaatccctt tattaccccc tccttcagac acctcaacc tcttctggct	2682
caaaaagaga attgggggct taggggtcga acccaagctt agaacttta gcaacaagac	2742
caccacttgg aaacctggga ttcaggaatg tgtggcctgc acagtgaagt gctggcaacc	2802
actaagaatt caaactgggg cctccagaac tcaactggggc ctacagcttt gatccctgac	2862
atctggaatc tggagaccag ggagcctttg gttctggcca gaatgctgca ggacttgaga	2922
agacctcacc tagaaattga cacaagtgga ccttaggcct tcctctctcc agatgtttcc	2982
agacttcctt gagacacgga gcccgaccct ccccatggag ccagctccct ctatttatgt	3042
ttgcacttgt gattattttat tattttatta ttatttttt atttacagat gaatgtattt	3102
atttgggaga cgggggtatc ctgggggacc caatgtagga gctgccttgg ctacagacatg	3162
ttttccgtga aaacggagct gaacaatagc ctgttcccat gtagccccct ggccctctgtg	3222
ccttcttttt attatgtttt taaaatatatt tatctgatta agttgtctaa acaatgctga	3282
tttggtgacc aactgtcact cattgctgag cctctgctcc ccaggggagt tgtgtctgta	3342
atcgccctac tattcagtgg cgagaaataa agtttgctta gaaaagaaac atggtctcct	3402
tcttggaatt aattctgcat ctgcctcttc ttgtgggtgg gaagaagctc cctaagtcct	3462
ctctccacag gctttaagat ccctcggacc cagtccatc cttagactcc tagggccctg	3522
gagaccctac ataaacaaag cccaacagaa tattcccat cccccaggaa acaagagcct	3582
gaacctaatt acctctccct cagggcctgg gaatttccaa ctctgggaat tc	3634

-continued

<210> SEQ ID NO 2
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 2

catgctttca gtgctcat 18

<210> SEQ ID NO 3
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 3

tgagggagcg tctgctggct 20

<210> SEQ ID NO 4
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 4

gtgctcatgg tgtcctttcc 20

<210> SEQ ID NO 5
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 5

taatcacaag tgcaaacata 20

<210> SEQ ID NO 6
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 6

taccccggtc tcccaaataa 20

<210> SEQ ID NO 7
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 7

agcaccgcct ggagccct 18

<210> SEQ ID NO 8
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 8

gctgaggaac aagcaccgcc

20

<210> SEQ ID NO 9
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 9

aggcagaaga gcgtggtggc

20

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 10

aaagtgcagc aggcagaaga

20

<210> SEQ ID NO 11
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 11

ttagagagag gtccctgg

18

<210> SEQ ID NO 12
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 12

tgactgcctg ggccagag

18

<210> SEQ ID NO 13
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 13

gggttcgaga agatgatc

18

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 14

gggctacagg cttgtcactc

20

<210> SEQ ID NO 15

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 15

cccctcagct tgagggtttg

20

<210> SEQ ID NO 16

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 16

ccattggcca ggagggcatt

20

<210> SEQ ID NO 17

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 17

accaccagct ggttatctct

20

<210> SEQ ID NO 18

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 18

ctgggagtag atgaggtaca

20

<210> SEQ ID NO 19

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 19

cccttgaaga ggacctggga

20

<210> SEQ ID NO 20

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 20

ggtgtgggtg aggagcaca

20

-continued

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 21

gtctggtagg agacggcgat 20

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 22

gcagagagga ggttgacctt 20

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 23

gcttggcctc agccccctct 20

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 24

cctcccagat agatgggctc 20

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 25

cccttctcca gctggaagac 20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 26

atctcagcgc tgagtcggtc 20

<210> SEQ ID NO 27
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 27

tcgagatagt cgggccgatt 20

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 28

aagtagacct gccagactc 20

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 29

ggatgttcgt cctcctcaca 20

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 30

accctaagcc cccaattctc 20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 31

ccacacattc ctgaatccca 20

<210> SEQ ID NO 32
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 32

aggccccagt gagttctgga 20

<210> SEQ ID NO 33
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 33
gtctccagat tccagatgtc 20

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 34
ctcaagtcct gcagcattct 20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 35
tgggtccccc aggatacccc 20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 36
acggaaaaca tgtctgagcc 20

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 37
ctccgttttc acggaaaaca 20

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 38
gcctattgtt cagctccgtt 20

<210> SEQ ID NO 39
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 39
ggtcacaaaa tcagcattgt t 21

-continued

<210> SEQ ID NO 40
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 40

gaggctcagc aatgagtgac 20

<210> SEQ ID NO 41
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: control sequence

<400> SEQUENCE: 41

gccaagctg gcatccgtca 20

<210> SEQ ID NO 42
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: control sequence

<400> SEQUENCE: 42

gccgaggtcc atgtcgtacg c 21

<210> SEQ ID NO 43
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 43

caggcgggtgc ttgttcct 18

<210> SEQ ID NO 44
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 44

gccagagggc tgattagaga ga 22

<210> SEQ ID NO 45
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR probe

<400> SEQUENCE: 45

cttctccttc ctgacgtgg caggc 25

<210> SEQ ID NO 46
<211> LENGTH: 19

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 46

gaaggtgaag gtcggagtc

19

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer

<400> SEQUENCE: 47

gaagatggtg atgggatttc

20

<210> SEQ ID NO 48
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR probe

<400> SEQUENCE: 48

caagcttccc gttctcagcc

20

<210> SEQ ID NO 49
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: control sequence

<400> SEQUENCE: 49

tctgagtagc agaggagctc

20

<210> SEQ ID NO 50
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 50

tgcgtctctc atttcccctt

20

<210> SEQ ID NO 51
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 51

tcccatctct ctccctctct

20

<210> SEQ ID NO 52
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 52

cagcgcacat ctttcaccca

20

<210> SEQ ID NO 53

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 53

tctctctcat ccctccctat

20

<210> SEQ ID NO 54

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 54

cgtctttctc catgtttttt

20

<210> SEQ ID NO 55

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 55

cacatctctt tctgcatccc

20

<210> SEQ ID NO 56

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 56

ctctcttccc catctcttgc

20

<210> SEQ ID NO 57

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 57

gtctctccat ctttccttct

20

<210> SEQ ID NO 58

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 58

ttccatgtgc cagacatcct

20

-continued

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 59

atacacactt agtgagcacc 20

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 60

ttcattcatt cattcactcc 20

<210> SEQ ID NO 61
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 61

tatatctgct tgttcattca 20

<210> SEQ ID NO 62
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 62

ctgtctccat atcttattta 20

<210> SEQ ID NO 63
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 63

tctcttctca caccacat 20

<210> SEQ ID NO 64
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 64

cacttgtttc ttccccatc 20

<210> SEQ ID NO 65
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 65

ctcaccatct ttattcatat

20

<210> SEQ ID NO 66
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 66

atatttcccg ctctttctgt

20

<210> SEQ ID NO 67
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 67

catctctctc cttagctgtc

20

<210> SEQ ID NO 68
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 68

tcttctctcc ttatctcccc

20

<210> SEQ ID NO 69
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 69

gtgtgccaga caccctatct

20

<210> SEQ ID NO 70
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 70

tccttccctg agtgtcttct

20

<210> SEQ ID NO 71
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 71
accttcacgc attcaacagc 20

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 72
ctccattcat ctgtgtattc 20

<210> SEQ ID NO 73
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 73
tgaggtgtct ggttttctct 20

<210> SEQ ID NO 74
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 74
acacatcctc agagctctta 20

<210> SEQ ID NO 75
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 75
ctagccctcc aagttccaag 20

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 76
cgggcttcaa tcccaaatac 20

<210> SEQ ID NO 77
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 77
aagttctgcc taccatcagc 20

-continued

<210> SEQ ID NO 78
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 78

gtccttctca cattgtctcc 20

<210> SEQ ID NO 79
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 79

ccttcccttg agctcagcga 20

<210> SEQ ID NO 80
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 80

ggcctgtgct gttcctccac 20

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 81

cgttctgagt atcccactaa 20

<210> SEQ ID NO 82
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 82

cacatcccac ctggccatga 20

<210> SEQ ID NO 83
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 83

gtcctctctg tctgtcatcc 20

<210> SEQ ID NO 84
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 84

ccaccccaca tccggttcct

20

<210> SEQ ID NO 85
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 85

tcctggccct cgagctctgc

20

<210> SEQ ID NO 86
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 86

atgtcggttc actctccaca

20

<210> SEQ ID NO 87
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 87

agaggagagt cagtggtggc

20

<210> SEQ ID NO 88
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 88

gatcccaaag tagacctgcc

20

<210> SEQ ID NO 89
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 89

cagactcggc aaagtcgaga

20

<210> SEQ ID NO 90
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 90
tagtcgggcc gattgatctc 20

<210> SEQ ID NO 91
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 91
agcgctgagt cggtcaccct 20

<210> SEQ ID NO 92
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 92
tctccagctg gaagaccctc 20

<210> SEQ ID NO 93
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 93
cccagataga tgggctcata 20

<210> SEQ ID NO 94
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 94
ccagggcttg gcctcagccc 20

<210> SEQ ID NO 95
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 95
cctctggggt ctccctctgg 20

<210> SEQ ID NO 96
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 96
caggggctct tgatggcaga 20

-continued

<210> SEQ ID NO 97
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 97

gaggaggttg accttggtct 20

<210> SEQ ID NO 98
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 98

ggtaggagac ggcgatgcgg 20

<210> SEQ ID NO 99
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 99

ctgatgggtg gggtaggag 20

<210> SEQ ID NO 100
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 100

aggcactcac ctcttcctc 20

<210> SEQ ID NO 101
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 101

ccctggggaa ctgtgggga 20

<210> SEQ ID NO 102
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 102

agacacttac tgactgcctg 20

<210> SEQ ID NO 103
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 103

gaagatgatac ctgaagagga

20

<210> SEQ ID NO 104
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 104

gagctcttac ctacaacatg

20

<210> SEQ ID NO 105
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 105

tgagggtttg ctggaggag

20

<210> SEQ ID NO 106
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: control sequence

<400> SEQUENCE: 106

gatacgctcg gactatgaag

20

<210> SEQ ID NO 107
<211> LENGTH: 7208
<212> TYPE: DNA
<213> ORGANISM: Mus musculus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (4527..4712,5225..5279,5457..5504,5799..6217)
<221> NAME/KEY: exon
<222> LOCATION: (4371)..(4712)
<221> NAME/KEY: intron
<222> LOCATION: (4713)..(5224)
<221> NAME/KEY: exon
<222> LOCATION: (5225)..(5279)
<221> NAME/KEY: intron
<222> LOCATION: (5280)..(5456)
<221> NAME/KEY: exon
<222> LOCATION: (5457)..(5504)
<221> NAME/KEY: intron
<222> LOCATION: (5505)..(5798)
<221> NAME/KEY: exon
<222> LOCATION: (5799)..(>6972)
<300> PUBLICATION INFORMATION:
<301> AUTHORS: Semon, D.
Kawashima, E.
Jongeneel, C.V.
Shakhov, A.N.
Nedospasov, S.A.
<302> TITLE: Nucleotide sequence of the murine TNF locus, including the
TNF-alpha (tumor necrosis factor) and TNF-beta (lymphotoxin) genes
<303> JOURNAL: Nucleic Acids Res.

-continued

<304> VOLUME: 15

<305> ISSUE: 21

<306> PAGES: 9083-9084

<307> DATE: 1987-11-11

<308> DATABASE ACCESSION NUMBER: Y00467 Genbank

<309> DATABASE ENTRY DATE: 1993-05-11

<400> SEQUENCE: 107

gaattctgaa gctccctctg tacagagcat tgggaagcctg ggggtgtacat ttgggggttac	60
atgatcttgg ggttctaaga gaataccccc aaatcatctt ccagacctgg aacattctag	120
gacagggttc tcaaccttcc taactccatg accctttaat acagttcctc atgttggtgt	180
gaccccaacc atacaattat ttctgttgct atttcataac tgtaatttcg ctgctattat	240
gaatcataat gtaaatattt gttttaaata gaggtttgcc aaaggacact tgcccacagg	300
ttgagaactg ccgtccaga gagtaagggg acacagttaa gattgttaca caccaggatg	360
ccccagattt ggggagaggg cactgtaatg gaacttcttg acatgaaact ggcagatgaa	420
actggcagaa aaaaaaaaaa aagctgggca gtggtggcac acaccttaa tccagcact	480
tgggaggcag aggcaggcgg atttctgagt tctaggccag cctggtctac agagtgagtt	540
tcaggacagc cagggtcaca cagagaaacc ctgtctcgaa aaaagcaaaa aaaaaaaaaa	600
aaaaaaaaa aaactggcag atgaccagaa aatacagata tattggaata actgtgactt	660
gaacccccaa agacaagaga ggaaataggc ctgaaggggc ggcaggcatg tcaagcatcc	720
agagccctgg gttcgaacct gaaaaacaa aggtgccgct aaccacatgt ggcttcggag	780
ccctccagac atgaccatga tcgacagaga gggaaatgtg cagagaagcc tgtgagcagt	840
caagggtgca gaagtgatat aaaccatcac tcttcaggga accaggcttc cagtccacagc	900
ccagctgcac cctctccacg aattgctcgg ccgttcaactg gaactcctgg gcctgaccca	960
gtccctctgt agtccctcgg gccacagatt ccccggaacc gactcccttt ccagaaacgc	1020
agtagtctaa gcccttagcc tgcggttctc tcttagggcc cagcctttcc tgccttcgac	1080
tgaaacagca gcatcttcta agccctgggg gcttccccaa gcccagccc cgacctagaa	1140
ccgcccgcgt gcctgccaca ctgccgcttc ctctataaag ggacccgagc gccagcggcc	1200
aggaccccgc acagcagggtg agcctctcct accctgtctc cttgggctta ccctgggtatc	1260
aggcatccct caggatccta cctcctttct tgagccacag cctttcttat acaacctgcc	1320
tggtatccca gccttaatgg gtctggtcct cctgtcgtgg ctttgatatt ttggtctgttc	1380
ctgtggcggc cttatcagtc tctctctctc tctctctctc tctctctctc tctctctctc	1440
tctctctctc tctccctctc tctctctctc tctctctctc ttctctctct ctgcctctgt	1500
tagccattgt ctgattctat ggtggagctt tcctcttccc ctctgtctct ccttatccct	1560
gtcacttca gggttccctt gcctgtcccc ttttctgtct gtgcgcctgt ctctcagggt	1620
ggctgtctca gctgggagggt aaggctctgc ttccgctgtg tgccccgcct ccgtacaca	1680
cacacactct ctctctctct ctccagaggt tctccacatg aactgctcg gccgtctcca	1740
cctcttgagg gtgcttgga cccctcctgt cttcctcctg gggtgctgc tggccctgcc	1800
tctaggggccc caggtaggac agcaagagat tgggggtgct ggggtggcct agctaactca	1860
gagtcctaga gtctcttcca ctctcttctg tcccaggac tctctggtgt ccgttctctc	1920
gctgccagga cagcccatcc actccctcag aagcacttga cccatggcat cctgaaacct	1980
gctgctcacc ttgttggtaa acttctgcct ccagaggaga ggtccagtcc ctgccttttg	2040

-continued

tcctacttgc ccaggggctc aggcgatctt cccatctccc cacaccaact tttcttacct	2100
ctaagggcag gcacccccact cccatctccc taccaacct cccacttgct cagtgcctgc	2160
tcctcagggg tggggacctc tgatcttgat agcccccaa tgtcttgctc ctcttcccag	2220
ggtacccccag caagcagaac tcaactgctt ggagagcaag cacggatcgt gcctttctcc	2280
gacatggctt ctctttgagc aacaactccc tcctgatccc caccagtggc ctctactttg	2340
tctactccca ggtgggtttt tctggagaaa gctgctccc cagggccatt cccactcca	2400
tctacttgcc acacgaggtc cagctctttt cctcccaata ccccttccat gtgcctctcc	2460
tcagtgcgca gaagtctgtg tatccgggac ttcaaggacc gtgggtgcgc tcaatgtacc	2520
agggggctgt gttcctgctc agtaaggag accagctgtc caccacacc gacggcatct	2580
cccatctaca cttcagcccc agcagtgtat tctttggagc ctttgactg tagattctaa	2640
agaaacccaa gaattggatt ccaggcctcc atcctgaccg ttgtttcaag ggtcacatcc	2700
ccacagtctc cagccttccc cactaaaata acctggagct ctcacgggag tctgagacac	2760
ttcaggggag tacatcttcc ccagggccac tcagatgct caggggacga ctcaagccta	2820
cctagaagtt cctgcacaga gcagggtttt tgtgggtcta ggtcgacag agacctggac	2880
atgaaggagg gacagacatg ggagagggtg ctgggaacag gggaagggtg actatttatg	2940
gagagaaaaa ttaagttatt tatttataga gaatagaaag aggggaaaaa tagaaagccg	3000
tcagatgaca actaggctcc agacacaaa gtgtctcacc tcagacagga cccatctaag	3060
agagagatgg cgagagaatt agatgtgggt gaccaagggg ttctagaaga aagcacgaag	3120
ctctaaaagc cagccactgc ttggctagac atccacaggg accccctgca ccatctgtga	3180
aaccaataa acctcttttc tctgagattc tgtctgcttg tgtctgtctt gcgttggggg	3240
agaaacttcc tggctctctt aaggagtggg gcaggggaca gaggcctcag ttggtccatg	3300
ggatccgggc agagcaaaga gacatgagga gcaggcagct cccagagaca tgggtggattc	3360
acgggagtga ggcagcttaa ctgccgagag acccaaagga tgagctaggg agatccatcc	3420
aagggtggag agagatgagg gttctgggga gaagtgactc cactggaggg tgggagagtg	3480
tttaggagtg ggagggtggg ggaggggaat ccttgaaga ccggggagtc atacggattg	3540
ggagaaatcc tggaagcagg gctgtgggac ctaaatgtct gagttgatgt accgcagtca	3600
agatatggca gaggctccgt ggaaaactca cttgggagca gggacccaaa gcagcagcct	3660
gagctcatga tcagagtgaaggaggagggc ttgtgagggt cgtgaattcc cagggtgag	3720
ttcattccct ctgggctgcc ccatactcat cccattaccc cccccaccag cctcccaaa	3780
gcccagtcac acttccaac tctcaagctg ctctgccttc agccacttcc tccaagaact	3840
caaacagggg gctttccctc ctcaatatca tgtctcccc cttatgcacc cagctttcag	3900
aagcaccccc ccatgctaag ttctccccc tggatgtccc atttagaaat caaaggaaa	3960
tagacacagg catggctctt ctacaaagaa acagacaatg attagctctg gaggacagag	4020
aagaaatggg tttcagttct cagggtccta tacaacacac acacacacac acacacacac	4080
acacacacac acacacccct ctgattggcc ccagattgcc acagaatcct ggtggggacg	4140
acgggggaga gattccttga tgctgggtg tccccaaact tccaaaccct ctgccccgc	4200
gatggagaag aaaccgagac agaggtgtag ggccactacc gcttctcca catgagatca	4260
tgggttttct caccaaggaa gttttccgag ggttgaatga gagcttttcc ccgccctctt	4320

-continued

ccccaagggc tataaaggcg gccgtctgca cagccagcca gcagaagctc cctcagcgag	4380
gacagcaagg gactagccag gagggagaac agaaactcca gaacatcttg gaaatagctc	4440
ccagaaaagc aagcagccaa ccaggcaggt tctgtccctt tcactcactg gccaaggcg	4500
ccacatctcc ctccagaaaa gacacc atg agc aca gaa agc atg atc cgc gac	4553
Met Ser Thr Glu Ser Met Ile Arg Asp	
1 5	
gtg gaa ctg gca gaa gag gca ctc ccc caa aag atg ggg ggc ttc cag	4601
Val Glu Leu Ala Glu Glu Ala Leu Pro Gln Lys Met Gly Gly Phe Gln	
10 15 20 25	
aac tcc agg cgg tgc cta tgt ctc agc ctc ttc tca ttc ctg ctt gtg	4649
Asn Ser Arg Arg Cys Leu Cys Leu Ser Leu Phe Ser Phe Leu Leu Val	
30 35 40	
gca ggg gcc acc acg ctc ttc tgt cta ctg aac ttc ggg gtg atc ggt	4697
Ala Gly Ala Thr Thr Leu Phe Cys Leu Leu Asn Phe Gly Val Ile Gly	
45 50 55	
ccc caa agg gat gag gtgagtgtct gggcaaccct tattctcgtc cacaagcaaa	4752
Pro Gln Arg Asp Glu	
60	
acgggttagg agggcaagaa ggacagtgtg agggaaagaa gtgggctaata gggcagggca	4812
aggtggagga gagtgtggag gggacagagt caggacctcg gacctatgcg tccagctgac	4872
taaacatcct tcgtcggatg cacagagaga tgaatgaacg aacaagtgtg ttcacacgtg	4932
gagagatctg gaaagatgtg gccaggggaa gaggggataa gcaagagata aaactcagag	4992
acagaaatga gagaggcatg agagataagg aggaagatga aggggagata acgggagatc	5052
aagcacagag ggcaccgcag aaagaagccg tgggttgac agatgaatga atgaagaaga	5112
aaacacaaa tgggggttggt gtggggcaaa gaggaactgt aagcggggca atcagccggg	5172
agcttctcct ttgggttgag tctgtcttaa ctaacctcct ttctctacac ag aag ttc	5230
Lys Phe	
cca aat ggc ctc cct ctc atc agt tct atg gcc cag acc ctc aca ctc	5278
Pro Asn Gly Leu Pro Leu Ile Ser Ser Met Ala Gln Thr Leu Thr Leu	
65 70 75 80	
agtaagtgtt cccacacctc tctcttaatt taagatggag aagggcagtt aggcagggga	5338
Arg	
tgagatgggg tggggggaaa acttaaagct ttggtttggg aggaaagggg tctaagtgca	5398
tagatgcttg ctgggaagcc taaaaggctc atccttgctt ttgtctcttc ccttcca	5455
gga tca tct tct caa aat tcg agt gac aag cct gta gcc cac gtc gta	5503
Ser Ser Ser Gln Asn Ser Ser Asp Lys Pro Val Ala His Val Val	
85 90 95	
ggtaaagattt ctttacatgt gccttgagaa tgaaggggca tgattttggg gggcgggttg	5563
aggggtgtcg agccaggctg agaaaagaca gagctcttag agacagcacg tgagagtcag	5623
agcagtgact caaaagcaag gcatcagggg gccacccggg acctcatagc caatgggatg	5683
tggaagaca gaggggtcag gaaccggaag tgaagtgtgg gtagctgctg aggctcagga	5743
tgtggagtgt gaactaagag ggtgacactg actcaatcct ccccccccc ctca gca	5800
Ala	
aac cac caa gtg gag gag cag ctg gag tgg ctg agc cag cgc gcc aac	5848
Asn His Gln Val Glu Glu Gln Leu Glu Trp Leu Ser Gln Arg Ala Asn	
100 105 110	
gcc ctc ctg gcc aac ggc atg gat ctc aaa gac aac caa cta gtg gtg	5896
Ala Leu Leu Ala Asn Gly Met Asp Leu Lys Asp Asn Gln Leu Val Val	

-continued

115	120	125	
cca gcc gat ggg ttg tac ctt gtc tac tcc cag gtt ctc ttc aag gga			5944
Pro Ala Asp Gly Leu Tyr Leu Val Tyr Ser Gln Val Leu Phe Lys Gly			
130	135	140	145
caa ggc tgc ccc gac tac gtg ctc ctc acc cac acc gtc agc cga ttt			5992
Gln Gly Cys Pro Asp Tyr Val Leu Leu Thr His Thr Val Ser Arg Phe			
150	155	160	
gct atc tca tac cag gag aaa gtc aac ctc ctc tct gcc gtc aag agc			6040
Ala Ile Ser Tyr Gln Glu Lys Val Asn Leu Leu Ser Ala Val Lys Ser			
165	170	175	
ccc tgc ccc aag gac acc cct gag ggg gct gag ctc aaa ccc tgg tat			6088
Pro Cys Pro Lys Asp Thr Pro Glu Gly Ala Glu Leu Lys Pro Trp Tyr			
180	185	190	
gag ccc ata tac ctg gga gga gtc ttc cag ctg gag aag ggg gac caa			6136
Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys Gly Asp Gln			
195	200	205	
ctc agc gct gag gtc aat ctg ccc aag tac tta gac ttt gcg gag tcc			6184
Leu Ser Ala Glu Val Asn Leu Pro Lys Tyr Leu Asp Phe Ala Glu Ser			
210	215	220	225
ggg cag gtc tac ttt gga gtc att gct ctg tga agggaaatggg tgttcaccca			6237
Gly Gln Val Tyr Phe Gly Val Ile Ala Leu			
230	235		
ttctctaccc agccccact ctgaccctt tactctgacc cttttattgt ctactcctca			6297
gagccccag tctgtgtcct tctaacttag aaaggggatt atggctcaga gtccaactct			6357
gtgtcagag ctttcaacaa ctactcagaa acacaagatg ctgggacagt gacctggact			6417
gtgggcctct catgcaccac catcaaggac tcaaatgggc tttccgaatt cactggagcc			6477
tcgaatgtcc attcctgagt tctgcaaagg gagagtggtc aggttgccctc tgtctcagaa			6537
tgaggctgga taagatctca ggccctccta ccttcagacc tttccagact cttccctgag			6597
gtgcaatgoa cagccttctct cacagagcca gccccctct atttatattt gcacttatta			6657
tttattattt atttattatt tatttatttg cttatgaatg tatttatttg gaagccggg			6717
gtgtcctgga ggaccagtg tgggaagctg tcttcagaca gacatgtttt ctgtgaaaac			6777
ggagctgagc tgtccccacc tggcctctct accttgttgc ctctctttt gcttatgttt			6837
aaaacaaaat atttatctaa cccaattgtc ttaataacgc tgatttggtg accaggctgt			6897
cgctacatca ctgaacctct gctccccacg ggagccgtga ctgtaattgc cctacagtca			6957
attgagagaa ataaagatcg cttggaanaa aaatgtgatt tctgtcttgg gatgaagtct			7017
gcatccatct ctttgccgag gcctaaagtc tctgggtcca gatctcagtc tttatacccc			7077
tggggcatta agacccccaa gacccccgtg gaacaaaagg cagccaacat ccctacctct			7137
cccccgaaa caggagccta accctaatta cctttgcctt ggggcatggg aatttccac			7197
tctggaatt c			7208
<210> SEQ ID NO 108			
<211> LENGTH: 20			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic			
<400> SEQUENCE: 108			
gagcttctgc tggctggctg			20

-continued

<210> SEQ ID NO 109
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 109

ccttgctgtc ctcgctgagg 20

<210> SEQ ID NO 110
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 110

tcatggtgtc ttttctggag 20

<210> SEQ ID NO 111
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 111

ctttctgtgc tcatggtgtc 20

<210> SEQ ID NO 112
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 112

gcggatcatg ctttctgtgc 20

<210> SEQ ID NO 113
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 113

gggaggccat ttgggaactt 20

<210> SEQ ID NO 114
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 114

cgaattttga gaagatgatc 20

<210> SEQ ID NO 115
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 115

ctcctccact tgggtggttg 20

<210> SEQ ID NO 116
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 116

cctgagatct tatccagcct 20

<210> SEQ ID NO 117
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 117

caattacagt cacggctccc 20

<210> SEQ ID NO 118
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 118

cccttcattc tcaaggcaca 20

<210> SEQ ID NO 119
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 119

caccctcaa cccgcccccc 20

<210> SEQ ID NO 120
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 120

agagctctgt cttttctcag 20

<210> SEQ ID NO 121
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 121

cactgctctg actctcacgt 20

<210> SEQ ID NO 122

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 122

atgagggtccc ggggtggcccc 20

<210> SEQ ID NO 123

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 123

caccctctgt ctttccacat 20

<210> SEQ ID NO 124

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 124

ctccacatcc tgagcctcag 20

<210> SEQ ID NO 125

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 125

attgagtcag tgtcacccctc 20

<210> SEQ ID NO 126

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 126

gctggctcag ccactccagc 20

<210> SEQ ID NO 127

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 127

tccttgagat ccatgccgtt 20

-continued

<210> SEQ ID NO 128
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 128

aacccatcgg ctggcaccac

20

<210> SEQ ID NO 129
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 129

gtttgagetc agccccctca

20

<210> SEQ ID NO 130
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 130

ctcctcccag gtatatgggc

20

<210> SEQ ID NO 131
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 131

tgagttggtc ccccttctcc

20

<210> SEQ ID NO 132
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 132

caaagtagac ctgcccggac

20

<210> SEQ ID NO 133
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 133

acacccattc ccttcacaga

20

<210> SEQ ID NO 134
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 134

cataatcccc tttctaagtt 20

<210> SEQ ID NO 135
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 135

cacagagttg gactctgagc 20

<210> SEQ ID NO 136
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 136

cagcatcttg tgtttctgag 20

<210> SEQ ID NO 137
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 137

cacagtccag gtcactgtcc 20

<210> SEQ ID NO 138
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 138

tgatggtggt gcatgagagg 20

<210> SEQ ID NO 139
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 139

gtgaattcgg aaagcccatt 20

<210> SEQ ID NO 140
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 140

cctgaccact ctcctttgc 20

<210> SEQ ID NO 141

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 141

tgcattcccc aggccacat 20

<210> SEQ ID NO 142

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 142

gccgaggtcc atgtcgtacg c 21

<210> SEQ ID NO 143

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 143

tcaagcagtg ccaccgatcc 20

<210> SEQ ID NO 144

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 144

agtgtcttct gtgtccaga 20

<210> SEQ ID NO 145

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 145

gtgtcttctg tgtgccagac 20

<210> SEQ ID NO 146

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 146

tgtcttctgt gtgccagaca 20

-continued

<210> SEQ ID NO 147
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 147

gtcttctgtg tgccagacac

20

<210> SEQ ID NO 148
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 148

tcttctgtgt gccagacacc

20

<210> SEQ ID NO 149
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 149

cttctgtgtg ccagacaccc

20

<210> SEQ ID NO 150
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 150

ttctgtgtgc cagacaccct

20

<210> SEQ ID NO 151
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 151

tctgtgtgcc agacacccta

20

<210> SEQ ID NO 152
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 152

ctgtgtgcc gacaccctat

20

<210> SEQ ID NO 153
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 153

tgtgtgccag acaccctatc 20

<210> SEQ ID NO 154
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 154

tgtgccagac accctatctt 20

<210> SEQ ID NO 155
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 155

gtgccagaca ccctatcttc 20

<210> SEQ ID NO 156
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 156

tgccagacac cctatcttct 20

<210> SEQ ID NO 157
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 157

gccagacacc ctatcttctt 20

<210> SEQ ID NO 158
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 158

ccagacaccc tatcttcttc 20

<210> SEQ ID NO 159
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 159

cagacaccct atcttcttct 20

<210> SEQ ID NO 160

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 160

agacacccta tcttcttctc 20

<210> SEQ ID NO 161

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 161

gacaccctat cttcttctct 20

<210> SEQ ID NO 162

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 162

acaccctatc ttcttctctc 20

<210> SEQ ID NO 163

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 163

caccctatct tcttctctcc 20

<210> SEQ ID NO 164

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 164

gtcttctgtg tgccagac 18

<210> SEQ ID NO 165

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 165

tcttctgtgt gccagaca 18

-continued

<210> SEQ ID NO 166
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 166

cttctgtgtg ccagacac

18

<210> SEQ ID NO 167
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 167

ttctgtgtgc cagacacc

18

<210> SEQ ID NO 168
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 168

ttctgtgtgcc agacaccc

18

<210> SEQ ID NO 169
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 169

ctgtgtgccca gacaccct

18

<210> SEQ ID NO 170
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 170

tgtgtgccag acacccta

18

<210> SEQ ID NO 171
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 171

gtgtgccaga caccctat

18

<210> SEQ ID NO 172
<211> LENGTH: 18
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 172

tgtgccagac accctatc 18

<210> SEQ ID NO 173
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 173

tgccagacac cctatctt 18

<210> SEQ ID NO 174
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 174

gccagacacc ctatcttc 18

<210> SEQ ID NO 175
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 175

ccagacacc tctcttct 18

<210> SEQ ID NO 176
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 176

cagacaccct atcttctt 18

<210> SEQ ID NO 177
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 177

agacacccta tcttcttc 18

<210> SEQ ID NO 178
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 178

gacaccctat cttcttct 18

<210> SEQ ID NO 179

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 179

acaccctatc ttcttctc 18

<210> SEQ ID NO 180

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 180

agaggtttgg agacacttac 20

<210> SEQ ID NO 181

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 181

gaattaggaa agaggtttgg 20

<210> SEQ ID NO 182

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 182

cccaaaccga gaattaggaa 20

<210> SEQ ID NO 183

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 183

tacccccaga cccaaaccga 20

<210> SEQ ID NO 184

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 184

gtactaaccc tacccccaga 20

-continued

<210> SEQ ID NO 185
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 185

ttccataccg gtactaaccc 20

<210> SEQ ID NO 186
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 186

cccccaactgc ttccataccg 20

<210> SEQ ID NO 187
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 187

ctttaaatTT cccccactgc 20

<210> SEQ ID NO 188
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 188

aagaccacaaa ctttaaatTT 20

<210> SEQ ID NO 189
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 189

atcctccccc aagaccacaaa 20

<210> SEQ ID NO 190
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 190

acctccatcc atcctccccc 20

<210> SEQ ID NO 191
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 191

ccctactttc acctccatcc 20

<210> SEQ ID NO 192
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 192

gaaaataccc ccctactttc 20

<210> SEQ ID NO 193
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 193

aaacttccta gaaaataccc 20

<210> SEQ ID NO 194
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 194

tgagaccctt aaacttccta 20

<210> SEQ ID NO 195
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 195

aagaaaaagc tgagaccctt 20

<210> SEQ ID NO 196
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 196

ggagagagaa aagaaaaagc 20

<210> SEQ ID NO 197
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 197

tgagccagaa gaggttgagg 20

<210> SEQ ID NO 198

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 198

attctctttt tgagccagaa 20

<210> SEQ ID NO 199

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 199

taagcccca attctctttt 20

<210> SEQ ID NO 200

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 200

gttccgaccc taagcccca 20

<210> SEQ ID NO 201

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 201

ctaagcttgg gttccgaccc 20

<210> SEQ ID NO 202

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 202

gcttaaagtt ctaagcttgg 20

<210> SEQ ID NO 203

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 203

tggtcttggt gcttaaagtt 20

-continued

<210> SEQ ID NO 204
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 204

ttcgaagtgg tggctctgtt 20

<210> SEQ ID NO 205
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 205

aatcccaggt ttcgaagtgg 20

<210> SEQ ID NO 206
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 206

cacattcctg aatcccaggt 20

<210> SEQ ID NO 207
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 207

gtgcaggcca cacattcctg 20

<210> SEQ ID NO 208
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 208

gcacttcact gtgcaggcca 20

<210> SEQ ID NO 209
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 209

gtggttgcca gcacttcact 20

<210> SEQ ID NO 210
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 210

tgaattctta gtggttgcca 20

<210> SEQ ID NO 211
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 211

ggccccagtt tgaattctta 20

<210> SEQ ID NO 212
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 212

gagttctgga ggccccagtt 20

<210> SEQ ID NO 213
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 213

aggccccagt gaggttctgga 20

<210> SEQ ID NO 214
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 214

tcaaagctgt aggccccagt 20

<210> SEQ ID NO 215
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 215

atgtcaggga tcaaagctgt 20

<210> SEQ ID NO 216
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 216

cagattccag atgtcagggg 20

<210> SEQ ID NO 217

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 217

ccctggtctc cagattccag 20

<210> SEQ ID NO 218

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 218

accaaaggct ccctggtctc 20

<210> SEQ ID NO 219

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 219

tctggccaga accaaaggct 20

<210> SEQ ID NO 220

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 220

cctgcagcat tctggccaga 20

<210> SEQ ID NO 221

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 221

cttctcaagt cctgcagcat 20

<210> SEQ ID NO 222

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 222

taggtgaggt cttctcaagt 20

-continued

<210> SEQ ID NO 223
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 223

tgtcaatttc tagtgaggt 20

<210> SEQ ID NO 224
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 224

ggtccacttg tgtcaatttc 20

<210> SEQ ID NO 225
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 225

gaaggcctaa ggtccacttg 20

<210> SEQ ID NO 226
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 226

ctggagagag gaaggcctaa 20

<210> SEQ ID NO 227
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 227

ctggaaacat ctggagagag 20

<210> SEQ ID NO 228
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 228

tcaaggaggt ctggaaacat 20

<210> SEQ ID NO 229
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 229

gctccgtgtc tcaaggaagt 20

<210> SEQ ID NO 230
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 230

ataaatacat tcattctgtaa 20

<210> SEQ ID NO 231
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 231

ggtctcccaa ataaatacat 20

<210> SEQ ID NO 232
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 232

aggatacccc ggtctcccaa 20

<210> SEQ ID NO 233
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 233

tgggtcccc aggatacccc 20

<210> SEQ ID NO 234
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 234

gctcctacat tgggtcccc 20

<210> SEQ ID NO 235
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 235

agccaaggca gctcctacat 20

<210> SEQ ID NO 236

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 236

aacatgtctg agccaaggca 20

<210> SEQ ID NO 237

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 237

tttcacggaa aacatgtctg 20

<210> SEQ ID NO 238

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 238

tcagctccgt tttcacggaa 20

<210> SEQ ID NO 239

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 239

agcctattgt tcagctccgt 20

<210> SEQ ID NO 240

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 240

acatgggaac agcctattgt 20

<210> SEQ ID NO 241

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 241

atcaaaagaa ggcacagagg 20

-continued

<210> SEQ ID NO 242
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 242

gtttagacaa cttaatcaga 20

<210> SEQ ID NO 243
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 243

aatcagcatt gtttagacaa 20

<210> SEQ ID NO 244
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 244

ttggtcacca aatcagcatt 20

<210> SEQ ID NO 245
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 245

tgagtgcagc ttggtcacca 20

<210> SEQ ID NO 246
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 246

ggctcagcaa tgagtgcagc 20

<210> SEQ ID NO 247
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 247

attacagaca caactcccct 20

<210> SEQ ID NO 248
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 248

tagtagggcg attacagaca

20

<210> SEQ ID NO 249
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 249

cgccactgaa tagtagggcg

20

<210> SEQ ID NO 250
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 250

ctttatttct cgccactgaa

20

<210> SEQ ID NO 251
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 251

ctgaggggagc gtctgctggc

20

<210> SEQ ID NO 252
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 252

ccttgctgag ggagcgtctg

20

<210> SEQ ID NO 253
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 253

ctggtcctct gctgtccttg

20

<210> SEQ ID NO 254
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 254

cctctgctgt ccttgctgag 20

<210> SEQ ID NO 255

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 255

ttctctccct cttagctggt 20

<210> SEQ ID NO 256

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 256

tccctcttag ctggtcctct 20

<210> SEQ ID NO 257

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 257

tctgaggggt gttttcaggg 20

<210> SEQ ID NO 258

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 258

ctgtagttgc ttctctccct 20

<210> SEQ ID NO 259

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 259

acctgcctgg cagcttgta 20

<210> SEQ ID NO 260

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 260

ggatgtggcg tctgaggggt 20

-continued

<210> SEQ ID NO 261
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 261

tgtgagagga agagaacctg 20

<210> SEQ ID NO 262
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 262

gaggaagaga acctgcctgg 20

<210> SEQ ID NO 263
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 263

agccgtgggt cagtatgtga 20

<210> SEQ ID NO 264
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 264

tgggtcagta tgtgagagga 20

<210> SEQ ID NO 265
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 265

gagagggtga agccgtgggt 20

<210> SEQ ID NO 266
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 266

tcatggtgtc cttccaggg 20

<210> SEQ ID NO 267
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 267

ctttcagtg ctcagtggtgc 20

<210> SEQ ID NO 268
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 268

tcatgctttc agtgctcatg 20

<210> SEQ ID NO 269
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 269

acgtcccga tcatgctttc 20

<210> SEQ ID NO 270
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 270

gctccacgtc ccggatcatg 20

<210> SEQ ID NO 271
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 271

tcctcggcca gctccacgtc 20

<210> SEQ ID NO 272
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 272

gcgcctcctc ggccagctcc 20

<210> SEQ ID NO 273
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 273

aggaacaagc accgcctgga 20

<210> SEQ ID NO 274

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 274

caagcaccgc ctggagccct 20

<210> SEQ ID NO 275

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 275

aaggagaaga ggctgaggaa 20

<210> SEQ ID NO 276

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 276

gaagaggctg aggaacaagc 20

<210> SEQ ID NO 277

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 277

cctgccacga tcaggaagga 20

<210> SEQ ID NO 278

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 278

cacgatcagg aaggagaaga 20

<210> SEQ ID NO 279

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 279

aagagcgtgg tggcgcctgc 20

-continued

<210> SEQ ID NO 280
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 280

cgtggtggcg cctgccacga

20

<210> SEQ ID NO 281
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 281

aagtgcagca ggcagaagag

20

<210> SEQ ID NO 282
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 282

cagcaggcag aagagcgtgg

20

<210> SEQ ID NO 283
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 283

gatcactcca aagtcagca

20

<210> SEQ ID NO 284
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 284

gggccgatca ctccaaagtg

20

<210> SEQ ID NO 285
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 285

gggccagagg gctgattaga

20

<210> SEQ ID NO 286
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 286

agagggctga ttagagagag 20

<210> SEQ ID NO 287
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 287

gctacaggct tgtcactcgg 20

<210> SEQ ID NO 288
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 288

ctgactgcct gggccagagg 20

<210> SEQ ID NO 289
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 289

tacaacatgg gctacaggct 20

<210> SEQ ID NO 290
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 290

agccactgga gctgccctc 20

<210> SEQ ID NO 291
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 291

ctggagctgc ccctcagctt 20

<210> SEQ ID NO 292
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 292

ttggcccggc ggttcagcca 20

<210> SEQ ID NO 293

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 293

ttggccagga gggcattggc 20

<210> SEQ ID NO 294

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 294

ccggcggttc agccactgga 20

<210> SEQ ID NO 295

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 295

ctcagctcca cgccattggc 20

<210> SEQ ID NO 296

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 296

caggagggca ttggcccggc 20

<210> SEQ ID NO 297

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 297

ctccacgcca ttggccagga 20

<210> SEQ ID NO 298

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 298

accagctggt tatctctcag 20

-continued

<210> SEQ ID NO 299
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 299

ctggttatct ctcagctcca 20

<210> SEQ ID NO 300
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 300

ccctctgatg gcaccaccag 20

<210> SEQ ID NO 301
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 301

tgatggcacc accagctggt 20

<210> SEQ ID NO 302
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 302

tagatgaggt acagggccctc 20

<210> SEQ ID NO 303
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 303

aagaggacct gggagtagat 20

<210> SEQ ID NO 304
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 304

gaggtacagg ccctctgatg 20

<210> SEQ ID NO 305
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 305

cagccttggc ccttgaagag 20

<210> SEQ ID NO 306
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 306

gacctgggag tagatgaggt 20

<210> SEQ ID NO 307
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 307

ttggcccttg aagaggacct 20

<210> SEQ ID NO 308
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 308

tggtgtgggt gaggagcaca 20

<210> SEQ ID NO 309
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 309

cggcgatgcg gctgatggtg 20

<210> SEQ ID NO 310
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 310

tgggtgagga gcacatgggt 20

<210> SEQ ID NO 311
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 311

tggtctggta ggagacggcg 20

<210> SEQ ID NO 312

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 312

atgcggctga tgggtgggt 20

<210> SEQ ID NO 313

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 313

agaggagggtt gaccttggtc 20

<210> SEQ ID NO 314

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 314

tggtaggagac ggcgatgcg 20

<210> SEQ ID NO 315

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 315

aggttgacct tggctggta 20

<210> SEQ ID NO 316

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 316

ggctcttgat ggcagagagg 20

<210> SEQ ID NO 317

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 317

tcataccaggg cttgcctc 20

-continued

<210> SEQ ID NO 318
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 318

ttgatggcag agaggaggtt

20

<210> SEQ ID NO 319
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 319

agctggaaga cccctcccag

20

<210> SEQ ID NO 320
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 320

atagatgggc tcataccagg

20

<210> SEQ ID NO 321
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 321

cggtcacct tctccagctg

20

<210> SEQ ID NO 322
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 322

gaagaccct cccagataga

20

<210> SEQ ID NO 323
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 323

acccttctcc agctggaaga

20

<210> SEQ ID NO 324
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 324

tcggcaaagt cgagatagtc 20

<210> SEQ ID NO 325
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 325

gggccgattg atctcagcgc 20

<210> SEQ ID NO 326
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 326

tagacctgcc cagactcggc 20

<210> SEQ ID NO 327
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 327

aaagtcgaga tagtcgggcc 20

<210> SEQ ID NO 328
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 328

gcaatgatcc caaagtagac 20

<210> SEQ ID NO 329
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 329

ctgcccagac tcggcaaagt 20

<210> SEQ ID NO 330
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 330

cgctcctcctc acagggcaat 20

<210> SEQ ID NO 331

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 331

ggaagggttg atgttcgtcc 20

<210> SEQ ID NO 332

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 332

tcctcacagg gcaatgatcc 20

<210> SEQ ID NO 333

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 333

gttgagggtg tctgaaggag 20

<210> SEQ ID NO 334

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 334

gttggatgtt cgctcctcctc 20

<210> SEQ ID NO 335

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 335

tttgagccag aagaggttga 20

<210> SEQ ID NO 336

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 336

gaggcgtttg ggaaggttgg 20

-continued

<210> SEQ ID NO 337
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 337

gcccccaatt ctctttttga

20

<210> SEQ ID NO 338
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 338

gccagaagag gttgagggtg

20

<210> SEQ ID NO 339
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 339

gggttccgac cctaagcccc

20

<210> SEQ ID NO 340
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 340

caattctctt tttgagccag

20

<210> SEQ ID NO 341
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 341

taaagttcta agcttggggtt

20

<210> SEQ ID NO 342
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 342

ccgaccctaa gcccccaatt

20

<210> SEQ ID NO 343
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 343

ggtggtcttg ttgcttaaag 20

<210> SEQ ID NO 344
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 344

ttctaagctt gggttccgac 20

<210> SEQ ID NO 345
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 345

cccaggtttc gaagtggtag 20

<210> SEQ ID NO 346
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 346

tcttggtgct taaagttcta 20

<210> SEQ ID NO 347
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 347

cacacattcc tgaatcccag 20

<210> SEQ ID NO 348
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 348

gtttcgaagt ggtggtcttg 20

<210> SEQ ID NO 349
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 349

cttcactgtg caggccacac 20

<210> SEQ ID NO 350

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 350

attcctgaat cccaggtttc 20

<210> SEQ ID NO 351

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 351

tagtggttgc cagcacttca 20

<210> SEQ ID NO 352

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 352

cccagtttga attcttagtg 20

<210> SEQ ID NO 353

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 353

ctgtgcaggc cacacattcc 20

<210> SEQ ID NO 354

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 354

gtgagttctg gaggccccag 20

<210> SEQ ID NO 355

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 355

gttgccagca cttcactgtg 20

-continued

<210> SEQ ID NO 356
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 356

tttgaattct tagtggttgc

20

<210> SEQ ID NO 357
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 357

aagctgtagg cccagtgag

20

<210> SEQ ID NO 358
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 358

ttctggaggc cccagtttga

20

<210> SEQ ID NO 359
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 359

agatgtcagg gatcaaagct

20

<210> SEQ ID NO 360
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 360

tggtctccag attccagatg

20

<210> SEQ ID NO 361
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 361

gtaggcccc a gtgagttctg

20

<210> SEQ ID NO 362
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 362

gaaccaaagg ctccctggtc 20

<210> SEQ ID NO 363
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 363

tcagggatca aagctgtagg 20

<210> SEQ ID NO 364
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 364

tccagattcc agatgtcagg 20

<210> SEQ ID NO 365
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 365

gcagcattct gccagaacc 20

<210> SEQ ID NO 366
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 366

gtcttctcaa gtctgcagc 20

<210> SEQ ID NO 367
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 367

aaaggctccc tggctccag 20

<210> SEQ ID NO 368
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 368

caatttctag gtgaggtcctt 20

<210> SEQ ID NO 369

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 369

attctggcca gaaccaaagg 20

<210> SEQ ID NO 370

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 370

aaggtccact tgtgtcaatt 20

<210> SEQ ID NO 371

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 371

gagagaggaa ggcctaaggt 20

<210> SEQ ID NO 372

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 372

tctaggtgag gtcttctcaa 20

<210> SEQ ID NO 373

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 373

ccacttgtgt caatttctag 20

<210> SEQ ID NO 374

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 374

gtctggaaac atctggagag 20

-continued

<210> SEQ ID NO 375
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 375

ccgtgtctca aggaagtctg

20

<210> SEQ ID NO 376
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 376

aggaaggcct aaggtccact

20

<210> SEQ ID NO 377
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 377

gaggagctg gctccatggg

20

<210> SEQ ID NO 378
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 378

gaaacatctg gagagaggaa

20

<210> SEQ ID NO 379
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 379

gtgcaaacat aaatagaggg

20

<210> SEQ ID NO 380
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 380

tctcaaggaa gtctggaaac

20

<210> SEQ ID NO 381
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 381

aataaataat cacaagtgc 20

<210> SEQ ID NO 382
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 382

gggctgggct ccgtgtctca 20

<210> SEQ ID NO 383
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 383

taccccggtc tcccaaataa 20

<210> SEQ ID NO 384
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 384

aacataaata gagggagctg 20

<210> SEQ ID NO 385
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 385

ttgggtcccc caggataccc 20

<210> SEQ ID NO 386
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 386

ataatcacia gtgcaaakat 20

<210> SEQ ID NO 387
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 387

aaagcagctc ctacattggg 20

<210> SEQ ID NO 388

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 388

cgggtctccca aataaataca 20

<210> SEQ ID NO 389

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 389

aaacatgtct gagccaaggc 20

<210> SEQ ID NO 390

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 390

tccccccagga taccgccggtc 20

<210> SEQ ID NO 391

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 391

agctcctaca ttgggtcccc 20

<210> SEQ ID NO 392

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 392

tgtctgagcc aagcagctc 20

<210> SEQ ID NO 393

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 393

cagcctattg ttcagctccg 20

-continued

<210> SEQ ID NO 394
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 394

agaaggcaca gaggccaggg

20

<210> SEQ ID NO 395
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 395

ttttcacgga aaacatgtct

20

<210> SEQ ID NO 396
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 396

tattgttcag ctccgttttc

20

<210> SEQ ID NO 397
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 397

aaaaacataa tcaaaagaag

20

<210> SEQ ID NO 398
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 398

cagataaata ttttaaaaaa

20

<210> SEQ ID NO 399
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 399

tacatgggaa cagcctattg

20

<210> SEQ ID NO 400
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 400

tttagacaac ttaatcagat 20

<210> SEQ ID NO 401
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 401

cataatcaaa agaaggcaca 20

<210> SEQ ID NO 402
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 402

accaaatacag cattgtttag 20

<210> SEQ ID NO 403
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 403

aaatatattta aaaaacataa 20

<210> SEQ ID NO 404
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 404

gagtgacagt tggtcaccaa 20

<210> SEQ ID NO 405
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 405

acaacttaatc agataaata 20

<210> SEQ ID NO 406
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 406

cagaggctca gcaatgagtg 20

<210> SEQ ID NO 407

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 407

atcagcattg tttagacaac 20

<210> SEQ ID NO 408

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 408

agggcgatta cagacacaac 20

<210> SEQ ID NO 409

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 409

acagttggtc accaaatcag 20

<210> SEQ ID NO 410

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 410

tcgccactga atagtagggc 20

<210> SEQ ID NO 411

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 411

gctcagcaat gagtgcagt 20

<210> SEQ ID NO 412

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 412

agcaaaacttt atttctcgcc 20

-continued

<210> SEQ ID NO 413
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 413

gattacagac acaactcccc

20

<210> SEQ ID NO 414
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 414

actgaatagt agggcgatta

20

<210> SEQ ID NO 415
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 415

actttatttc tcgccactga

20

<210> SEQ ID NO 416
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 416

gctgtccttg ctgagggagc

20

<210> SEQ ID NO 417
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 417

cttagctggt cctctgctgt

20

<210> SEQ ID NO 418
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 418

gttgcttctc tccctcttag

20

<210> SEQ ID NO 419
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 419

tggcgtctga gggttgtttt 20

<210> SEQ ID NO 420
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 420

agagaacctg cctggcagct 20

<210> SEQ ID NO 421
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 421

cagtatgtga gaggaagaga 20

<210> SEQ ID NO 422
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 422

ggtgaagcgc tgggtcagta 20

<210> SEQ ID NO 423
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 423

agtgtcatg gtgtcctttc 20

<210> SEQ ID NO 424
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 424

ccggatcatg ctttcagtgc 20

<210> SEQ ID NO 425
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 425

ggccagctcc acgtcccga 20

<210> SEQ ID NO 426

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 426

ggccccctg tcttcttggg 20

<210> SEQ ID NO 427

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 427

ggctgaggaa caagcaccgc 20

<210> SEQ ID NO 428

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 428

tcaggaagga gaagaggctg 20

<210> SEQ ID NO 429

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 429

tggcgctgc cagatcagg 20

<210> SEQ ID NO 430

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 430

ggcagaagag cgtggtggcg 20

<210> SEQ ID NO 431

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 431

ctccaaagtg cagcaggcag 20

-continued

<210> SEQ ID NO 432
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 432

gctgattaga gagaggtccc

20

<210> SEQ ID NO 433
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 433

tgccctgggcc agagggtga

20

<210> SEQ ID NO 434
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 434

gctgcccctc agcttgaggg

20

<210> SEQ ID NO 435
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 435

ggttcagcca ctggagctgc

20

<210> SEQ ID NO 436
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 436

gggcattggc ccggcggttc

20

<210> SEQ ID NO 437
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 437

cgccattggc caggagggca

20

<210> SEQ ID NO 438
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 438

tatctctcag ctccacgccca 20

<210> SEQ ID NO 439
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 439

gcaccaccag ctggttatct 20

<210> SEQ ID NO 440
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 440

acaggccctc tgatggcacc 20

<210> SEQ ID NO 441
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 441

gggagtagat gaggtacagg 20

<210> SEQ ID NO 442
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 442

ccttgaagag gacctgggag 20

<210> SEQ ID NO 443
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 443

gaggagcaca tgggtggagg 20

<210> SEQ ID NO 444
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 444

gctgatggtg tgggtgagga

20

<210> SEQ ID NO 445

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 445

ggagacggcg atgcggctga

20

<210> SEQ ID NO 446

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 446

gaccttggtc tggtaggaga

20

<210> SEQ ID NO 447

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 447

ggcagagagg aggttgacct

20

<210> SEQ ID NO 448

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 448

tgggctcata ccagggcttg

20

<210> SEQ ID NO 449

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 449

cccctcccag atagatgggc

20

<210> SEQ ID NO 450

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 450

tgagtcggtc acccttctcc

20

-continued

<210> SEQ ID NO 451
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 451

gattgatctc agcgctgagt

20

<210> SEQ ID NO 452
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 452

cgagatagtc gggccgattg

20

<210> SEQ ID NO 453
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 453

caaagtagac ctgccagac

20

<210> SEQ ID NO 454
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 454

acagggcaat gatcccaaag

20

<210> SEQ ID NO 455
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 455

atgttcgtcc tcctcacagg

20

<210> SEQ ID NO 456
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 456

gtttgggaag gttgatgtt

20

<210> SEQ ID NO 457
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 457

aagagggtga ggggtgtctga 20

<210> SEQ ID NO 458
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 458

ctctttttga gccagaagag 20

<210> SEQ ID NO 459
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 459

cctaagcccc caattctctt 20

<210> SEQ ID NO 460
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 460

agcttggggt cgcaccctaa 20

<210> SEQ ID NO 461
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 461

ttgcttaaag ttctaagctt 20

<210> SEQ ID NO 462
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 462

gaagtgggtg tcttggtgct 20

<210> SEQ ID NO 463
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 463

tgaatcccag gtttcgaagt 20

<210> SEQ ID NO 464

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 464

caggccacac attcctgaat 20

<210> SEQ ID NO 465

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 465

cagcacttca ctgtgcaggc 20

<210> SEQ ID NO 466

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 466

attcttagtg gttgccagca 20

<210> SEQ ID NO 467

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 467

gaggccccag tttgaattct 20

<210> SEQ ID NO 468

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 468

ccccagtgag ttctggaggc 20

<210> SEQ ID NO 469

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 469

gatcaaagct gtaggccccca 20

-continued

<210> SEQ ID NO 470
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 470

attccagatg tcagggatca

20

<210> SEQ ID NO 471
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 471

ctccctgggc tccagattcc

20

<210> SEQ ID NO 472
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 472

ggccagaacc aaaggctccc

20

<210> SEQ ID NO 473
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 473

gtcctgcagc attctggcca

20

<210> SEQ ID NO 474
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 474

gtgaggtctt ctcaagtcct

20

<210> SEQ ID NO 475
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 475

tgtgtcaatt tctaggtgag

20

<210> SEQ ID NO 476
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 476

ggcctaaggt ccacttgtgt 20

<210> SEQ ID NO 477
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 477

atctggagag aggaagcct 20

<210> SEQ ID NO 478
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 478

aggaagtctg gaaacatctg 20

<210> SEQ ID NO 479
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 479

gggctccgtg tctcaaggaa 20

<210> SEQ ID NO 480
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 480

aaatagaggg agctggctcc 20

<210> SEQ ID NO 481
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 481

cacaagtgca aacataaata 20

<210> SEQ ID NO 482
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 482

tcccaaataa atacattcat 20

<210> SEQ ID NO 483

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 483

caggataccc cggctctcca 20

<210> SEQ ID NO 484

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 484

ctacattggg tccccagga 20

<210> SEQ ID NO 485

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 485

gagccaaggc agctcctaca 20

<210> SEQ ID NO 486

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 486

acggaaaaca tgtctgagcc 20

<210> SEQ ID NO 487

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 487

ttcagctccg ttttcacgga 20

<210> SEQ ID NO 488

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 488

gggaacagcc tattgttcag 20

-continued

<210> SEQ ID NO 489
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 489

tcaaaagaag gcacagaggc

20

<210> SEQ ID NO 490
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 490

ttttaaaaa cataatcaaa

20

<210> SEQ ID NO 491
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 491

ttaatcagat aaatatttta

20

<210> SEQ ID NO 492
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 492

cattgttttag acaacttaat

20

<210> SEQ ID NO 493
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 493

tggtcaccaa atcagcattg

20

<210> SEQ ID NO 494
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 494

gcaatgagtg acagttggtc

20

<210> SEQ ID NO 495
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 495

gggagcagag gctcagcaat

20

<210> SEQ ID NO 496
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 496

atagtagggc gattacagac

20

<210> SEQ ID NO 497
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 497

atttctcgcc actgaatagt

20

<210> SEQ ID NO 498
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 498

ctgattagag agaggtccc

19

<210> SEQ ID NO 499
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 499

ctgattagag agaggtcc

18

<210> SEQ ID NO 500
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 500

tgagtgtctt ctgtgtgcca

20

<210> SEQ ID NO 501
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic

-continued

<400> SEQUENCE: 501

gagtgtcttc tgtgtgccag

20

<210> SEQ ID NO 502

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 502

caccaagctg cggccccaa

20

<210> SEQ ID NO 503

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 503

tccgtcatcg ctctcaggg

20

What is claimed is:

1. A method of treating an inflammatory disorder in an individual comprising administering to said individual an effective amount of an oligonucleotide up to 30 nucleotides in length complementary to a nucleic acid molecule encoding human tumor necrosis factor- α .

2. The method of claim 1, wherein said oligonucleotide inhibits the expression of said human tumor necrosis factor- α and comprises at least an 8 nucleobase portion of SEQ ID NO: 24, SEQ ID NO: 27, SEQ ID NO: 287 SEQ ID NO: 29, SEQ ID NO: 30, SEQ ID NO: 34, SEQ ID NO: 39, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 149, SEQ ID NO: 157, SEQ ID NO: 264, SEQ ID NO: 271, SEQ ID NO: 272, SEQ ID NO: 290, SEQ ID NO: 297, SEQ ID NO: 299, SEQ ID NO: 315, SEQ ID NO: 334, SEQ ID NO: 418, SEQ ID NO: 423, SEQ ID NO: 425, SEQ ID NO: 427, SEQ ID NO: 431, SEQ ID NO: 432, SEQ ID NO: 435, SEQ ID NO: 437, SEQ ID NO: 438, SEQ ID NO: 439, SEQ ID NO: 441, SEQ ID NO: 455, SEQ ID NO: 457, SEQ ID NO: 458, SEQ ID NO: 460, SEQ ID NO: 463, SEQ ID NO: 465, SEQ ID NO: 466, SEQ ID NO: 468, SEQ ID NO: 472, SEQ ID NO: 474, SEQ ID NO: 475, SEQ ID NO: 483, SEQ ID NO: 485, SEQ ID NO: 494 or SEQ ID NO: 496.

3. The method of claim 1, wherein said antisense oligonucleotide is administered orally, topically or parenterally.

4. The method of claim 1, wherein said inflammatory disorder is inflammatory bowel disease, Crohn's disease, colitis or rheumatoid arthritis.

5. The method of claim 1, wherein said oligonucleotide comprises at least one modified intersugar linkage.

6. The method of claim 4, wherein said intersugar linkage is a phosphorothioate linkage.

7. The method of claim 1, wherein said oligonucleotide comprises at least one 2'-O-methoxyethyl modification.

8. The method of claim 1, wherein said oligonucleotide comprises at least one 5-methyl cytidine.

9. The method of claim 7, wherein every 2'-O-methoxyethyl modified cytidine residue is a 5-methyl cytidine.

10. The method of claim 7, wherein every cytidine residue is a 5-methyl cytidine.

11. The method of claim 1, wherein said modified intersugar linkage is a methylene(methylimino) intersugar linkage.

* * * * *