




(86) Date de dépôt PCT/PCT Filing Date: 2003/06/26
 (87) Date publication PCT/PCT Publication Date: 2004/01/15
 (45) Date de délivrance/Issue Date: 2009/04/07
 (85) Entrée phase nationale/National Entry: 2004/12/20
 (86) N° demande PCT/PCT Application No.: EP 2003/006726
 (87) N° publication PCT/PCT Publication No.: 2004/005266
 (30) Priorité/Priority: 2002/07/03 (EP02014828.4)

(51) Cl.Int./Int.Cl. *C07D 263/32* (2006.01),
A61K 31/421 (2006.01), *A61P 3/10* (2006.01)
 (72) Inventeurs/Inventors:
BRODBECK, BERND, DE;
HILPERT, HANS, CH;
HUMM, ROLAND, DE
 (73) Propriétaire/Owner:
F. HOFFMANN-LA ROCHE AG, CH
 (74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : DERIVES OXAZOLE ET UTILISATION DE CEUX-CI EN TANT QUE SENSIBILISATEURS A L'INSULINE
 (54) Title: OXAZOLE DERIVATIVES AND THEIR USE AS INSULIN SENSITIZERS



(57) Abrégé/Abstract:

Compound of formula (I): and pharmaceutically acceptable salts and esters thereof, wherein R¹ is aryl; R² is hydrogen, alkyl or cycloalkyl; R³ is hydrogen, alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl-S(O)₂- or aryl-S(O)₂-; R⁴ is aralkyl; R⁵, R⁶, R⁷ and R⁸ are independently selected from hydrogen, alkyl or cycloalkyl; and n is 1,2,3,4 or 5, are useful as insulin sensitizers, particularly PPAR activators.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/005266 A1


- (51) International Patent Classification⁷: C07D 263/32, A61K 31/421, A61P 3/10
- (21) International Application Number: PCT/EP2003/006726
- (22) International Filing Date: 26 June 2003 (26.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 02014828.4 3 July 2002 (03.07.2002) EP
- (71) Applicant: F. HOFFMANN-LA ROCHE AG [CH/CH]; 124, Grenzacherstrasse, CH-4070 Basle (CH).
- (72) Inventors: BRODBECK, Bernd; Friedrichstrasse 62, 79585 Steinen-Hoellstein (DE). HILPERT, Hans; Gustav Bay-Strasse 34, CH-4142 Muenchenstein (CH). HUMM, Roland; In den Sigrismatten 2a, 79115 Freiburg (DE).
- (74) Agent: WASCHBUESCH, Klaus; 124, Grenzacherstrasse, CH-4070 Basle (CH).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: OXAZOLE DERIVATIVES AND THEIR USE AS INSULIN SENSITIZERS




(57) Abstract: Compound of formula (I): and pharmaceutically acceptable salts and esters thereof, wherein R¹ is aryl; R² is hydrogen, alkyl or cycloalkyl; R³ is hydrogen, alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl-S(O)₂- or aryl-S(O)₂-; R⁴ is aralkyl; R⁵, R⁶, R⁷ and R⁸ are independently selected from hydrogen, alkyl or cycloalkyl; and n is 1,2,3,4 or 5, are useful as insulin sensitizers, particularly PPAR activators.

OXAZOLE DERIVATIVES AND THEIR USE AS INSULIN SENSITIZERS

The present invention is concerned with novel oxazole derivatives useful as insulin sensitizers, particularly PPAR activators.

The invention is concerned especially with compounds of formula I

5



(I)

and pharmaceutically acceptable salts and esters thereof, wherein

R¹ is aryl;

10 R² is hydrogen, alkyl or cycloalkyl;

R³ is hydrogen, alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl-S(O)₂- or aryl-S(O)₂- ;

R⁴ is aralkyl;

R⁵, R⁶, R⁷ and R⁸ are independently selected from hydrogen, alkyl or cycloalkyl; and

n is 1, 2, 3, 4 or 5.

- 2 -

The compounds of formula I and their pharmaceutically acceptable salts and esters are novel and have valuable pharmacological properties. They are insulin sensitizers, particularly PPAR activators.

Peroxisome Proliferator Activated Receptors (PPAR's) are members of the nuclear hormone receptor super family, which are ligand-activated transcription factors regulating gene expression. Various subtypes thereof have been identified and cloned. These include PPAR α , PPAR β (also known as PPAR δ), and PPAR γ . There exist at least two major isoforms of PPAR γ . While PPAR γ 1 is ubiquitously expressed in most tissues, the longer isoform PPAR γ 2 is almost exclusively found in adipocytes. In contrast, PPAR α is predominantly expressed in the liver, kidney and heart. PPAR's modulate a variety of body responses including glucose- and lipid- homeostasis, cell differentiation, inflammatory responses and cardiovascular events.

Diabetes is a disease in which a patient's ability to control glucose levels in blood is impaired, because he has partially lost the ability to respond properly to the action of insulin. In type II diabetes (T2D), often referred to as non-insulin dependent diabetes mellitus (NIDDM), which afflicts 80-90 % of all diabetic patients in developed countries, the Isles of Langerhans in the pancreas still produce insulin. However, the target organs, mainly muscle, liver and adipose tissue, exhibit a profound resistance to insulin stimulation, and the body compensates by producing unphysiologically high levels of insulin. In later stage of disease, however, insulin secretion decreases due to exhaustion of the pancreas. In addition to that T2D is a metabolic-cardiovascular disease syndrome. Among the comorbidities associated with T2D are for example insulin resistance, dyslipidemia, hypertension, endothelial dysfunction and inflammatory atherosclerosis.

Current first line treatment for diabetes generally involves low fat - and glucose - diet and exercise. However, compliance can be moderate and as the disease progresses, treatment with hypoglycemic drugs, e.g. sulfonylureas or metformin, becomes necessary. A promising new class of drugs has recently been introduced that resensitizes patients to their own insulin (insulin sensitizers), thereby reverting blood glucose and triglyceride levels to normal, and thus abolishing, or at least reducing, the requirement for exogenous insulin. Pioglitazone (ActosTM) and rosiglitazone (AvandiaTM) belong to the thiazolidinediones (TZD) class of PPAR γ -agonists and were the first representatives who had been approved for NIDDM in several countries. These compounds, however, suffer from side effects including rare but severe liver toxicity (as seen with troglitazone), and they increase body weight in humans. Therefore, new, better and more efficacious drugs for the treatment of NIDDM are urgently needed. Recent studies provide evidence that a coagonism on PPAR α and PPAR γ would result in compounds with enhanced therapeutic

potential, i. e. with an improved lipid profile effect on top of the normalization of glucose- and insulin-levels (Keller and Wahli: Trends Endocrin. Metab. 1993; 4:291-296, Macdonald and Lane: Current Biology Vol.5 pp.618-621 (1995)).

The novel compounds of the present invention exceed the compounds known in the art, inasmuch as they bind to and activate both, PPAR α and PPAR γ , simultaneously and very efficiently. Therefore, these compounds combine the anti-glycemic effect of PPAR γ activation with the anti-dyslipidemic effect of PPAR α activation. Consequently, plasma glucose and insulin are reduced (=insulin sensitization), triglycerides lowered and HDL cholesterol increased (=improved lipid profile). In addition, such compounds may also lower LDL cholesterol, decrease blood pressure and counteract inflammatory atherosclerosis. Since multiple facets of the T2D disease syndrome are addressed by PPAR α and γ coagonists, they are expected to have an enhanced therapeutic potential compared to the compounds already known in the art.

Accordingly, the compounds of formula I can be used in the prophylaxis and/or treatment of diabetes, particularly non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome.

Objects of the present invention are the compounds of formula I and their aforementioned pharmaceutically acceptable salts and esters per se and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments comprising the said compounds, their pharmaceutically acceptable salts and esters, the use of the said compounds, esters and salts for the prophylaxis and/or therapy of illnesses, especially in the treatment and/or prophylaxis of diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome and particularly for the prophylaxis and/or therapy of non-insulin dependent diabetes mellitus, and the use of the said compounds, salts and esters for the production of medicaments for the treatment and/or prophylaxis of illnesses, especially in the treatment and/or prophylaxis of diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome.

In the present description the term "alkyl", alone or in combination, signifies a straight-chain or branched-chain alkyl group with 1 to 8 carbon atoms, preferably a straight or branched-chain alkyl group with 1 to 6 carbon atoms and particularly preferred a straight or branched-chain alkyl group with 1 to 4 carbon atoms Examples of straight-

chain and branched C₁-C₈ alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert.-butyl, the isomeric pentyls, the isomeric hexyls, the isomeric heptyls and the isomeric octyls, preferably methyl and ethyl and most preferred methyl.

The term "cycloalkyl", alone or in combination, signifies a cycloalkyl ring with 3 to 8
5 carbon atoms and preferably a cycloalkyl ring with 3 to 6 carbon atoms. Examples of C₃-C₈ cycloalkyl are cyclopropyl, methyl-cyclopropyl, dimethylcyclopropyl, cyclobutyl, methyl-cyclobutyl, cyclopentyl, methyl-cyclopentyl, cyclohexyl, methyl-cyclohexyl, dimethyl-cyclohexyl, cycloheptyl and cyclooctyl, preferably cyclopropyl.

The term "alkoxy", alone or in combination, signifies a group of the formula alkyl-
10 O- in which the term "alkyl" has the previously given significance, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec. butoxy and tert.butoxy, 2-hydroxyethoxy, 2-methoxyethoxy preferably methoxy and ethoxy and most preferred methoxy.

The term "aryloxy", alone or in combination, signifies a group of the formula aryl-O-
15 in which the term "aryl" has the previously given significance, such as phenyloxy.

The term "aryl", alone or in combination, signifies a phenyl or naphthyl group, preferably a phenyl group which optionally carries one or more substituents each independently selected from halogen, trifluoromethyl, amino, alkyl, alkoxy, alkylcarbonyl, cyano, carbamoyl, alkoxycarbamoyl, methylenedioxy, carboxy, alkoxycarbonyl,
20 aminocarbonyl, alkyaminocarbonyl, dialkylaminocarbonyl, hydroxy, nitro and the like, such as phenyl, chlorophenyl, trifluoromethylphenyl, chlorofluorophenyl, aminophenyl, methylcarbonylphenyl, methoxyphenyl, methylenedioxyphenyl, 1-naphthyl and 2-naphthyl. Preferred is phenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, 3-aminophenyl, 4-methylcarbonylphenyl, 4-methoxyphenyl and particularly phenyl.

25 The term "aralkyl", alone or in combination, signifies an alkyl or cycloalkyl group as previously defined in which one or more, preferably one hydrogen atom has been replaced by an aryl group as previously defined. Preferred are benzyl, benzyl substituted with hydroxy, alkoxy or halogen, preferably fluorine. Particularly preferred is benzyl.

The term "amino", alone or in combination, signifies a primary, secondary or
30 tertiary amino group bonded via the nitrogen atom, with the secondary amino group carrying an alkyl or cycloalkyl substituent and the tertiary amino group carrying two similar or different alkyl or cycloalkyl substituents or the two nitrogen substituents together forming a ring, such as, for example, -NH₂, methylamino, ethylamino,

dimethylamino, diethylamino, methyl-ethylamino, pyrrolidin-1-yl or piperidino etc., preferably amino, dimethylamino and diethylamino and particularly primary amino.

The term "halogen" signifies fluorine, chlorine, bromine or iodine and preferably fluorine, chlorine or bromine.

5 The term "carbonyl", alone or in combination signifies the -C(O)- group.

The term "cyano", alone or in combination signifies the group -CN.

The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as
10 hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, preferably hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the
15 like. In addition these salts may be prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring
20 substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polymine resins and the like. The compound of formula I can also be present in the form of zwitterions. Particularly preferred pharmaceutically acceptable salts of compounds of formula I are the hydrochloride salts.

25 The compounds of formula I can also be solvated, e.g. hydrated. The solvation can be effected in the course of the manufacturing process or can take place e.g. as a consequence of hygroscopic properties of an initially anhydrous compound of formula I (hydration). The term pharmaceutically acceptable salts also includes physiologically acceptable solvates.

30 "Pharmaceutically acceptable esters" means that compounds of general formula (I) may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compounds in vivo. Examples of such compounds include physiologically acceptable and metabolically labile ester derivatives, such as

methoxymethyl esters, methylthiomethyl esters and pivaloyloxymethyl esters. Additionally, any physiologically acceptable equivalents of the compounds of general formula (I), similar to the metabolically labile esters, which are capable of producing the parent compounds of general formula (I) in vivo, are within the scope of this invention.

5

The term "lipase inhibitor" refers to compounds which are capable of inhibiting the action of lipases, for example gastric and pancreatic lipases. For example orlistat and lipstatin as described in U.S. Patent No. 4,598,089 are potent inhibitor of lipases. Lipstatin is a natural product of microbial origin, and orlistat is the result of a hydrogenation of lipstatin. Other lipase inhibitors include a class of compound commonly referred to as panclicins. Panclicins are analogues of orlistat (Mutoh et al, 1994). The term "lipase inhibitor" refers also to polymer bound lipase inhibitors for example described in International Patent Application WO99/34786 (Geltex Pharmaceuticals Inc.). These polymers are characterized in that they have been substituted with one or more groups that inhibit lipases. The term "lipase inhibitor" also comprises pharmaceutically acceptable salts of these compounds. The term "lipase inhibitor" preferably refers to orlistat.

Orlistat is a known compound useful for the control or prevention of obesity and hyperlipidemia. See, U.S. Patent No. 4,598,089, issued July 1, 1986, which also discloses processes for making orlistat and U.S. Patent No. 6,004,996, which discloses appropriate pharmaceutical compositions. Further suitable pharmaceutical compositions are described for example in International Patent Applications WO 00/09122 and WO 00/09123. Additional processes for the preparation of orlistat are disclosed in European Patent Applications Publication Nos. 185,359, 189,577, 443,449, and 524,495.

Orlistat is preferably orally administered from 60 to 720 mg per day in divided doses two to three times per day. Preferred is wherein from 180 to 360 mg, most preferably 360 mg per day of a lipase inhibitor is administered to a subject, preferably in divided doses two or, particularly, three times per day. The subject is preferably an obese or overweight human, i.e. a human with a body mass index of 25 or greater. Generally, it is preferred that the lipase inhibitor be administered within about one or two hours of ingestion of a meal containing fat. Generally, for administering a lipase inhibitor as defined above it is preferred that treatment be administered to a human who has a strong family history of obesity and has obtained a body mass index of 25 or greater.

Orlistat can be administered to humans in conventional oral compositions, such as, tablets, coated tablets, hard and soft gelatin capsules, emulsions or suspensions.

Examples of carriers which can be used for tablets, coated tablets, dragées and hard gelatin capsules are lactose, other sugars and sugar alcohols like sorbitol, mannitol, maltodextrin, or other fillers; surfactants like sodium lauryl sulfate, Brij™ 96, or Tween™ 80; disintegrants like sodium starch glycolate, maize starch or derivatives thereof; polymers like povidone, 5 crospovidone; talc; stearic acid or its salts and the like. Suitable carriers for soft gelatin capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like. Moreover, the pharmaceutical preparations can contain preserving agents, solubilizers, stabilizing agents, wetting agents, emulsifying agents, sweetening agents, coloring agents, flavoring agents, salts for varying the osmotic pressure, buffers, coating 10 agents and antioxidants. They can also contain still other therapeutically valuable substances. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods known in the pharmaceutical art. Preferably, orlistat is administered according to the formulation shown in the Examples and in U.S. Patent No. 6,004,996, respectively.

15

The compounds of formula I can contain several asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates.

20 Preferred are the compounds of formula I and pharmaceutically acceptable salts thereof, particularly the compounds of formula I.

Further preferred are compounds of formula I, wherein R^1 is phenyl.

A preferred embodiment of the present invention are compounds of formula I, wherein R^2 is alkyl, preferably methyl.

25 Also preferred are compounds of formula I, wherein R^3 is alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl-S(O)₂- or aryl-S(O)₂- . Further preferred are those compounds, wherein R^3 is methyl, propyl, benzyl, methylcarbonyl, phenylcarbonyl, methyl-S(O)₂- , phenyl-S(O)₂- , phenyl or phenyl substituted with one or more, preferably one or two substituents independently selected from alkyl, halogen, trifluoromethyl and 30 alkoxy.

A further preferred aspect of the present invention are compounds of formula I, wherein R^3 is alkyl or phenyl, wherein phenyl is optionally mono- or disubstituted with

- 8 -

halogen. Particularly preferred are those compounds of formula I, wherein R³ is propyl, phenyl, fluorophenyl or difluorophenyl.

Further preferred compounds of formula I are those, wherein R⁴ is arylmethyl. Particularly preferred are the compounds of formula I, wherein R⁴ is benzyl substituted
5 with carboxy and optionally further substituted with fluoro, chloro, trifluoromethyl or alkoxy.

Also preferred are compounds according to formula I, wherein R⁵ is hydrogen.

Further preferred are compounds according to formula I, wherein R⁶ is hydrogen.

Another preferred aspect of the present invention are compounds of formula I,
10 wherein R⁷ is hydrogen.

Also preferred are compounds according to formula I, wherein R⁸ is hydrogen.

Particularly preferred compounds according to formula I are those, wherein R⁵, R⁶, R⁷ and R⁸ are hydrogen.

Another preferred embodiment of the present invention are the compounds of
15 formula I, wherein n is 1, 2 or 3. Particularly preferred compounds of formula I are those, wherein n is 2.

Examples of preferred compounds of formula (I) are

1. 2-[(Acetyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 20 2. 2-[(Benzoyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
3. 2-[(Methanesulfonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
4. 2-[(Benzenesulfonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-
25 amino)-methyl]-benzoic acid;
5. 2-[(Methyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;

- 9 -

6. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-propyl-amino)-methyl]-benzoic acid;
7. 2-[(Benzyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 5 8. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-o-tolyl-amino)-methyl]-benzoic acid;
9. 2-[(3-Fluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 10 10. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-m-tolyl-amino)-methyl]-benzoic acid;
11. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-(3-trifluoromethyl-phenyl)-amino)-methyl]-benzoic acid;
12. 2-[(4-Fluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 15 13. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-p-tolyl-amino)-methyl]-benzoic acid;
14. 2-[(4-Methoxy-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 20 15. 2-[(3,4-Dimethyl-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
16. 2-[(3,4-Difluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
17. 2-[(4-Fluoro-3-methyl-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 25 18. 3-Fluoro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;
19. 3-Chloro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;

20. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-3-trifluoromethyl-benzoic acid;
21. 3-Methoxy-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;
- 5 22. 4-Chloro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;
23. 4-Methoxy-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;
24. 5-Fluoro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;
- 10 25. 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid; and
26. 2-Methoxy-6-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid.

15

Examples of particularly preferred compounds of formula (I) are


- 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid;
- 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-propyl-amino)-methyl]-benzoic acid;
- 20 2-[(3-Fluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 2-[(3,4-Difluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;
- 25 3-Fluoro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid; and
- 5-Fluoro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid.

Processes for the manufacture of compounds of formula I are an object of the invention.

The substituents and indices used in the following description of the processes have the significance given above unless indicated to the contrary.

Compounds of general formula I, wherein R^1 , R^2 , R^4 to R^8 and n are as previously defined and, wherein R^3 is hydrogen, alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl-S(O)₂- or aryl-S(O)₂- can be prepared according to Scheme I and II:

Scheme I




Boc means tert-butoxycarbonyl

- 10 The oxazole ether (3) can be prepared by coupling of the alcohol (1) and the monoprotected aniline (2) under Mitsunobu conditions in THF (O. Mitsunobu, *Synthesis*, 1, 1981).

Alkylation of (3) with (4) can be accomplished with KOH in DMSO followed by deprotection of the Boc group using CF_3COOH to give compound (5).

Scheme II




R' is alkyl or aryl,
 R'' is alkyl or aryl
 R''' is alkyl or aralkyl

5 Acylations and sulfonylations of the amine (5) can be carried out with the corresponding chlorides. Alkylations of (5) require preferably the corresponding alkyl iodids or benzyl bromide. All reactions can be performed in the presence of a base, e.g. NEt_3 in THF to give the precursor of (I) as the esters. In a second step the esters can be hydrolyzed with LiOH in a solvent mixture of THF, MeOH and water affording the target compounds (I). Arylations of the amine (5) can be performed with the corresponding boronic acids
 10 according to the method of Lam (P.Y.S. Lam, G. Vincent, C.G. Clark, S. Deudon, P.K. Jadhav, Tetrahedron Lett. 42, 3415, 2001) to give the precursor of (I) as the esters which can be hydrolyzed as described above.

In case a free carboxy group is present e.g. in substituent R^4 this carboxy group can be protected by methods known in the art e.g. as ethyl ester. Present OH residues can be
 15 protected using appropriate protecting groups such as e.g. as ethyl ester.

Alternatively, compounds of general formula I, wherein R^1 , R^2 , R^4 to R^8 and n are as previously defined and, wherein R^3 is hydrogen, alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl- $\text{S}(\text{O})_2$ - or aryl- $\text{S}(\text{O})_2$ - can be prepared according to Scheme III:

Scheme III




The alternative preparation of (I) according to Scheme III preferentially be used when R³ is fixed and R⁴ will be varied follows the same type of reactions as described in Schemes I and II.

- 5 In case a free carboxy group is present e.g. in substituent R⁴ this carboxy group can be protected by methods known in the art e.g. as ethyl ester

Starting compound (1) can be obtained e.g. according to Scheme IV or V:

Scheme IV




Aldehydes (1a) are commercially available or known. They are condensed with diketo-
 monoximes (2a) according to literature precedence (Goto, Y.; Yamazaki, M.; Hamana, M.;
 Chem Pharm Bull (1971), 19, 2050) in the presence of a strong acid, typically HCl, in a
 5 polar solvent like AcOH to yield the oxazole-N-oxides (3a) (step a). Subsequent treatment
 with POCl₃ in dichloromethane under reflux provides the corresponding primary
 chlorides (4a) (Goto, Y.; Yamazaki, M.; Hamana, M.; Chem Pharm Bull (1971), 19, 2050,
 step b). These intermediates are either used as such, transformed according to well
 established methods into the corresponding alcohols or activated alcohols like mesylates or
 10 tosylates or into the bromides or iodides, or finally further elaborated via S_N2-reaction
 with NaCN to give, via nitrils 5 (step c), exhaustive hydrolysis (step d) and reduction (step
 e), e. g. with borane in tetrahydrofuran, the building blocks (7a).

4-Chloromethyl-2-aryl or 2-heteroaryl-oxazoles (4a) with R² equal hydrogen are
 preferably prepared from the corresponding aryl or heteroaryl carboxamides and 1,3-
 15 dichloroacetone as described e. g. in Bioorg. Med. Chem. Lett. (2000), 10(17), 2041-2044.

Compounds of formula (I), wherein n is 1 can be obtained by reacting a compound of formula 4a with a compound of formula 6 analogous to the reactions shown in Scheme III.

Scheme V



5

N-Acyl-glycine esters (1b) are either commercially available, known, or can be prepared by standard operations of N-acylation. Mono-allylated esters (2b) can easily be obtained by double deprotonation of (1b) with a strong, non-nucleophilic base like LiHMDS in an aprotic solvent like THF, typically at -78°C , followed by treatment with allyl bromide to produce selectively the C-alkylated products (2b) (step a). Standard hydrolysis generates intermediate acids (3b) (step b), which are then transformed, following well established literature precedence (J. Med. Chem. (1996), 39, 3897), into compounds (4b) (step c). Ring-closure to the oxazole using trifluoro-acetic acid and trifluoro-acetic anhydride as

10

reagents generates key intermediates (5b) (step d), which, finally, are elaborated via hydroboration to the target alcohols (6b), e. g. with 9-BBN in THF and ensuing oxidative work-up with H₂O₂ and NaOH (step e).

Starting compounds of formula (1), wherein n is 4 can be obtained as follows:

- 5 1) mesylation of a compound according to formula (6b)
- 2) SN₂-reaction with NaCN in order to obtain the corresponding nitril
- 3) hydrolysis of the nitril
- 4) reduction e.g. with borane

10 Starting compounds of formula (1), wherein n is 5 can be obtained as follows:

- 1) mesylation of a compound according to formula 1, wherein n is 4
- 2) SN₂-reaction with NaCN in order to obtain the corresponding nitril
- 3) hydrolysis of the nitril
- 4) reduction e.g. with borane

15 Starting compounds (2) and (6) are known or can be synthesized by methods known in the art, e.g. by nitration of the phenole using sulfuric- and salpetric-acid followed by reduction of the corresponding nitro phenole using catalytic hydrogenation or iron in hydrochloric acid. Boc protection of the primary amine can be accomplished with Boc anhydride in pyridine to give (2) or, alternatively, the unprotected aniline can be
20 converted to compound (6) using the arylation procedure developed by Lam (P.Y.S. Lam, G. Vincent, C.G. Clark, S. Deudon, P.K. Jadhav, Tetrahedron Lett. 42, 3415, 2001).

Compounds according to formula (4) and the corresponding protected analogs can be prepared as follows: The substituted toluol carbonic acid is protected as the ester using well known procedures, e.g. esterification with an alcohol and hydrochloric acid.


25 Bromination of the methyl group follows also well established procedures, e.g. using N-bromo succinimide and a catalytic amount of dibenzoyl peroxide in an halogenated solvent such as CCl₄.

The conversion of a compound of formula I into a pharmaceutically acceptable salt can be carried out by treatment of such a compound with an inorganic acid, for example a hydrohalic acid, such as, for example, hydrochloric acid or hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid etc., or with an organic acid, such as, for example, acetic acid, citric acid, maleic acid, fumaric acid, tartaric acid, methanesulfonic acid or p-toluenesulfonic acid. The corresponding carboxylate salts can also be prepared from the compounds of formula I by treatment with physiologically compatible bases.

The conversion of compounds of formula I into pharmaceutically acceptable esters or amides can be carried out e.g. by treatment of suited amino or hydroxyl groups present in the molecules with an carboxylic acid such as acetic acid, with a condensating reagent such as benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP) or N,N-dicyclohexylcarbodiimide (DCCI) to produce the carboxylic ester or carboxylic amide.

Further preferred is a process for the preparation of a compound according to formula I comprising one of the following reactions:

a) reaction of a compound according to formula (5)




in the presence of R^3 -Hal and following reaction in the presence of a hydroxide, preferably LiOH, in order to obtain a compound of formula I, wherein Hal means chlorine, bromine or iodine and R^1 to R^8 and n are defined as before. Particularly preferred is the above reaction, wherein R^3 means alkylcarbonyl or arylcarbonyl and Hal means chlorine. Further preferred is the above reaction, wherein R^3 means alkyl or aralkyl and Hal means bromine or iodine.

b) reaction of a compound according to formula (5) in the presence of $(HO)_2B$ -aryl and following reaction in the presence of a hydroxide, preferably LiOH, in order to obtain a compound of formula I, wherein R^1 , R^2 , R^4 to R^8 and n are defined as before;

c) reaction of a compound according to formula (7)

- 18 -



(7)

in the presence of R⁴-Hal and following reaction in the presence of a hydroxide, preferably LiOH, in order to obtain a compound of formula I, wherein Hal means chlorine, bromine or iodine and R¹ to R⁸ and n are defined as before. Preferred is the above reaction, wherein

5 Hal means bromine.

Preferred intermediates are:

{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-carbamic acid tert-butyl ester

10 2-[(tert-Butoxycarbonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid ethyl ester

2-({4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenylamino}-methyl)-benzoic acid ethyl ester and

{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amine

15

The compounds of formula I described above for use as therapeutically active substances are a further object of the invention. Preferred is the use as therapeutically active substances for the prophylaxis and/or therapy of diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels,

20 atherosclerotic diseases or metabolic syndrome and particularly preferred non-insulin dependent diabetes mellitus.

Also an object of the invention are compounds described above for the preparation of medicaments for the prophylaxis and/or therapy of diseases which are modulated by PPAR α and/or PPAR γ agonists, preferably for the production of medicaments for the

25 prophylaxis and/or therapy of diabetes, non-insulin dependent diabetes mellitus, elevated

blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome and particularly preferred non-insulin dependent diabetes mellitus.

Likewise an object of the invention are pharmaceutical compositions comprising a compound of formula I described above and a therapeutically inert carrier. Another object
5 of the present invention is the above pharmaceutical composition further comprising a therapeutically effective amount of a lipase inhibitor particularly, wherein the lipase inhibitor is orlistat.

An object of the invention is also the use of the compounds described above for the production of medicaments, particularly for the treatment and/or prophylaxis of diseases
10 which are modulated by PPAR α and/or PPAR γ agonists, preferably diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome and particularly preferred non-insulin dependent diabetes mellitus.

A further object of the present invention is the use of a compound of formula I in
15 the manufacture of a medicament for the treatment and/or prophylaxis of diseases which are modulated by PPAR α and/or PPAR γ agonists in a patient who is also receiving treatment with a lipase inhibitor. Preferred is the above use, wherein the lipase inhibitor is orlistat. Particularly preferred is the above use for the treatment and/or prophylaxis of diseases, wherein the diseases are diabetes, non-insulin dependent diabetes mellitus,
20 elevated blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome and particularly preferred non-insulin dependent diabetes mellitus.

A further object of the invention comprises compounds which are manufactured according to one of the described processes.

A further object of the invention is a method for the treatment and/or prophylaxis of
25 diseases which are modulated by PPAR α and/or PPAR γ agonists, preferably diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, atherosclerotic diseases or metabolic syndrome and particularly preferred non-insulin dependent diabetes mellitus, whereby an effective amount of a compound of formula I is administered. Another object of the present invention is the
30 above method which further comprises administration to the human a therapeutically effective amount of a lipase inhibitor, particularly, wherein the lipase inhibitor is orlistat. The above method for simultaneous, separate or sequential administration is also an object of the present invention.

Assay Procedures

The following tests can be used in order to determine the activity of the compounds of formula I.

- 5 Background information on the performed assays can be found in: Nichols JS et al. "Development of a scintillation proximity assay for peroxisome proliferator-activated receptor gamma ligand binding domain", (1998) *Anal. Biochem.* 257: 112-119.

Full-length cDNA clones for human PPAR α and mouse PPAR γ were obtained by RT-PCR from human adipose and mouse liver cRNA, respectively, cloned into plasmid vectors and
10 verified by DNA sequencing. Bacterial and mammalian expression vectors were constructed to produce glutathione-s-transferase (GST) and Gal4 DNA binding domain proteins fused to the ligand binding domains (LBD) of PPAR γ (aa 174 to 476) and PPAR α (aa 167 to 469). To accomplish this, the portions of the cloned sequences encoding the LBDs were amplified from the full-length clones by PCR and then subcloned into the
15 plasmid vectors. Final clones were verified by DNA sequence analysis.

Induction, expression, and purification of GST-LBD fusion proteins were performed in *E. coli* strain BL21(pLysS) cells by standard methods (Ref: *Current Protocols in Molecular Biology*, Wiley Press, edited by Ausubel et al.).

Radioligand Binding Assay

- 20 PPAR α receptor binding was assayed in TKE10 (10 mM Tris-HCl, pH 8, 50 mM KCl, 2mM EDTA, 0.1mg/ml fatty acid free BSA and 10 mM DTT). For each 96 well 2.4 ug equivalent of GST-PPAR α -LBD fusion protein and radioligand, e.g. 40000 dpm 2(S)-(2-benzoyl-phenylamino)-3-{4-[1,1-ditritio-2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid, were incubated in 100 ul volume at RT for 2 hrs. Bound ligand
25 was removed from unbound ligand by solid phase separation using MultiScreen plates (Millipore™) filled with 80 ul of SG25 according to the manufacturer's recommendations.

- PPAR γ receptor binding was assayed in TKE50 (50mM Tris-HCl, pH 8, 50 mM KCl, 2mM EDTA, 0.1 mg/ml fatty acid-free BSA and 10 mM DTT). For each 96 well reaction an 140 ng equivalent of GST-PPAR γ -LBD fusion protein was bound to 10 ug SPA beads
30 (PharmaciaAmersham) in a final volume of 50 ul by shaking. The resulting slurry was

- 21 -

incubated for 1h at RT and centrifuged for 2 min at 1300g. The supernatant containing unbound protein was removed and the semidry pellet containing the receptor-coated beads was resolved in 50 ul of TKE. For radioligand binding e.g. 10000 dpm 2(S)-(2-benzoyl-phenylamino)-3-{4-[1,1-ditritio-2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid in 50 ul were added, the reaction incubated at RT for 1h and scintillation proximity counting performed. All binding assays were performed in 96 well plates and the amount of bound ligand measured on a Packard TopCount™ using OptiPlates™ (Packard). Nonspecific binding was determined in the presence of 10⁻⁴ M unlabelled compound. Dose response curves were done in triplicates within a range of concentration from 10⁻¹⁰ M to 10⁻⁴ M.

Luciferase Transcriptional Reporter Gene Assays

Baby hamster kidney cells (BHK21 ATCC CCL10) were grown in DMEM medium containing 10% FBS at 37°C in a 95%O₂:5%CO₂ atmosphere. Cells were seeded in 6 well plates at a density of 10⁵ Cells/well and then batch-transfected with either the pFA-PPAR γ -LBD or pFA-PPAR α -LBD expression plasmids plus the pFR-luc reporter plasmid and an expression plasmid encoding the secretable form of alkaline phosphatase (SEAP) as a normalization control. Transfection was accomplished with the Fugene 6 reagent (Roche Molecular Biochemicals) according to the suggested protocol. Six hours following transfection, the cells were harvested by trypsinization and seeded in 96 well plates at a density of 10⁴ cells/well. After 24 hours to allow attachment of cells, the medium was removed and replaced with 100 ul of phenol red-free medium containing the test substances or control ligands (final. 0.1% DMSO). Following incubation of the cells for 24 hours with substances, 50 ul of the supernatant was recovered and analyzed for SEAP activity (Roche Molecular Biochemicals). The remainder of the supernatant was discarded, 50 ul PBS was added per well followed by one volume of Luciferase Constant-Light Reagent (Roche Molecular Biochemicals) to lyse the cells and initiate the luciferase reaction. Luminescence for both SEAP and luciferase was measured in a Packard TopCount. Luciferase activity was normalized to the SEAP control and transcriptional activation in the presence of a test substance was expressed as fold-activation over cells incubated in the absence of the substance. EC₅₀ values were calculated using the XLfit program (ID Business Solutions Ltd. UK).

The compounds of the present invention exhibit IC₅₀ values of 0.1 nM to 50 μ M, preferably 1 nM to 10 μ M, particularly 1 - 3500 nM, more preferred 1 to 500 nM, for PPAR α and PPAR γ . The compounds further exhibit EC₅₀ values of 0.1 nM to 50 μ M,

preferably 1 nM to 10 μ M, more preferably 1 to 3500 nM, particularly 1 to 500 nM, for PPAR α and PPAR γ .

| | PPAR γ IC ₅₀ (μ M) | PPAR α EC ₅₀ (μ M) | PPAR γ EC ₅₀ (μ M) |
|---------------|--|--|--|
| Example 6 | 0.93 | 2.70 | 1.32 |
| Example 12 | 1.08 | 2.59 | 2.32 |
| Example 18 | 0.24 | 10 | 1.98 |
| Example 26 | 0.98 | 2.05 | 1.41 |
| Rosiglitazone | 0.465 | inactive | 0.025 |

5 The compounds of formula I and their pharmaceutically acceptable salts and esters can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories, parenterally, e.g. in
10 the form of injection solutions or infusion solutions, or topically, e.g. in the form of ointments, creams or oils.

 The production of the pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described compounds of formula I and their pharmaceutically acceptable, into a galenical
15 administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

 Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard
20 gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers are, however, required in the case of soft gelatine

- 23 -

capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

The dosage of the compounds of formula I can vary within wide limits depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 mg to about 1000 mg, especially about 1 mg to about 100 mg, comes into consideration. Depending on the dosage it is convenient to administer the daily dosage in several dosage units.

The pharmaceutical preparations conveniently contain about 0.5-500 mg, preferably 0.5-100 mg, of a compound of formula I.

The following Examples serve to illustrate the present invention in more detail. They are, however, not intended to limit its scope in any manner.

Examples

Preparation of the starting material of examples 1 – 17

a) {4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-carbamic acid tert-butyl ester

To a solution of 5.00 g of (4-hydroxy-phenyl)-carbamic acid tert-butyl ester, 7.28 g of 2-
5 (5-methyl-2-phenyl-1,3-oxazol-4-yl)ethan-1-ol and 9.40 g of triphenylphosphine in 100
ml of THF was added at 0°C a solution of 7.25 g of diisopropyl azodicarboxylate in 50 ml
of THF over 30 min and stirring was continued at 22°C for 16 h. A further portion of 1.88
g of triphenylphosphine and 1.45 g of diisopropyl azodicarboxylate in 10 ml of THF was
added at 0°C and stirring was continued at 22°C for 2 h after which time conversion was
10 complete. The mixture was evaporated and the residue purified by chromatography (SiO₂,
n-hexane/AcOEt 4:1) to give 7.5 g of the title compound as a colorless solid. MS: (M+H)⁺
395.4.

b) 2-[(tert-Butoxycarbonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-
amino)-methyl]-benzoic acid ethyl ester

15 To a suspension of 3.41 g of powdered KOH in 110 ml of DMSO was added at 22°C 6.00 g
of {4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-carbamic acid tert-butyl ester
and the suspension was stirred for 25 min. A solution of 7.40 g of 2-bromomethyl-benzoic
acid ethyl ester in 10 ml of DMSO was added slowly keeping the temperature at 15-20°C
and stirring was continued at 22°C for 2.5 h. The dark mixture was partitioned between
20 500 ml of a saturated aqueous NH₄Cl and 200 ml of AcOEt, the organic layer was washed
with saturated aqueous NH₄Cl and water, dried and evaporated. The residue was purified
by chromatography (SiO₂, n-hexane/AcOEt 4:1) to give 7.96 g of the title compound as a
pale yellow oil. MS: (M+H)⁺ 557.3.

25 c) 2-({4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenylamino}-methyl)-benzoic acid
ethyl ester

To a solution of 7.00 g of 2-[(tert-butoxycarbonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-
ethoxy]-phenyl}-amino)-methyl]-benzoic acid ethyl ester in 40 ml of dichloromethane was
added 9.6 ml of trifluoroacetic acid and stirring was continued at 22°C for 2.5 h. The
30 mixture was partitioned between aqueous saturated Na₂CO₃ and dichloromethane, the
organic layer was dried and evaporated. The residue was purified by chromatography

(SiO₂, n-hexane/AcOEt 4:1) to give 3.14 g of the title compound as a pale yellow oil. MS: (M+H)⁺ 457.5.

General description for the preparation of examples 1-7

To a solution of 0.2 mmol of 2-({4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenylamino}-methyl)-benzoic acid ethyl ester prepared as described above and 0.4 mmol of triethylamine in 2 ml of THF was added 0.22 mmol of the corresponding acyl chloride or sulfo chloride or 0.22-2 mmol of the alkyl iodide or benzyl bromide followed by addition of a catalytic amount of dimethylamino pyridine in cases of slow conversions. The reaction mixtures were stirred at 22-55°C until completion of the conversion. The suspension was filtered, the filtrate evaporated and the residue purified by preparative HPLC chromatography (RP-18, CH₃CN/H₂O gradient) to give the product as the ester.

The esters (0.1 mmol) were hydrolyzed with 0.3 mmol of LiOH·H₂O in 1 ml of THF, 0.5 ml of MeOH and 0.5 ml of water followed by purification using preparative HPLC chromatography (RP-18, CH₃CN/H₂O gradient).

15

Example 1

2-[(Acetyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid ethyl ester was obtained in 66% yield as a pale yellow gum. MS: (M+H)⁺ 499.3. After hydrolysis, 2-[(acetyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid was obtained in 61% yield as a white solid. MS: (M-
20 H) 469.2.

Example 2

2-[(Benzoyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid ethyl ester was obtained in 65% yield as a colorless gum. MS: (M+H)⁺ 561.4. After hydrolysis, 2-[(benzoyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid was obtained in 71% yield as a white solid. MS: (M-H)
25 531.1.

Example 3

2-[(Methanesulfonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid ethyl ester was obtained in 57% yield as a yellow gum. MS: (M+H)⁺
30 535.3. After hydrolysis, 2-[(methanesulfonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-

- 26 -

ethoxy]-phenyl}-amino)-methyl]-benzoic acid was obtained in 75% yield as a white solid.
MS: (M-H) 505.2.

Example 4

2-[(Benzenesulfonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-
5 methyl]-benzoic acid ethyl ester was obtained in 89% yield as a pale yellow gum. MS:
(M+H)⁺ 597.1. After hydrolysis, 2-[(benzenesulfonyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-
yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid was obtained in 66% yield as a white
solid. MS: (M-H) 567.1.

Example 5

10 2-[(Methyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-
benzoic acid ethyl ester was obtained in 29% yield as a pale brown gum. MS: (M+H)⁺
471.1. After hydrolysis, 2-[(methyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-
phenyl}-amino)-methyl]-benzoic acid was obtained in 64% yield as a pale yellow gum.
MS: (M+H)⁺ 443.4.

Example 6

15 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-propyl-amino)-methyl]-
benzoic acid ethyl ester was obtained in 59% yield as a yellow gum. MS: (M+H)⁺ 499.3.
After hydrolysis, 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-propyl-
amino)-methyl]-benzoic acid was obtained in 83% yield as a pale yellow gum. MS:
20 (M+H)⁺ 471.3.

Example 7

2-[(Benzyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-
benzoic acid ethyl ester was obtained in 55% yield as a pale yellow gum. MS: (M+H)⁺
547.2. After hydrolysis, 2-[(benzyl-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-
25 phenyl}-amino)-methyl]-benzoic acid was obtained in 59% yield as a pale yellow gum.
MS: (M+H)⁺ 519.3.

General description for the preparation of examples 8-17

According to the method described by P.Y.S. Lam et al., Tetrahedron Letters 42, 3415,
30 2001, a suspension of 0.44 mmol of the corresponding boronic acid and 0.25 g of

- 27 -

molecular sieves in 3 ml of dichloromethane was treated subsequently with 0.22 mmol of 2-({4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenylamino}-methyl)-benzoic acid ethyl ester prepared as described above, 0.22 mmol of copper(II)acetate, 0.24 mmol of 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) and 0.44 mmol of NEt₃ and the mixture
5 was stirred at 22°C without sealing the flask (oxygen required) for 16 h. The mixture was filtered over a short pad of silica and the filtrate was purified by preparative HPLC chromatography (RP-18, CH₃CN/H₂O gradient) to give the product as the ester. The ester was hydrolyzed as described above.

Example 8

10 2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-o-tolyl-amino]-methyl]-benzoic acid ethyl ester was obtained in 22% yield as a brown gum. MS: (M+H)⁺ 547.3. After hydrolysis, 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-o-tolyl-amino]-methyl]-benzoic acid was obtained in 78% yield as a pale yellow gum. MS: (M+H)⁺ 519.3.

Example 9

15 2-[(3-Fluoro-phenyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl]-amino)-methyl]-benzoic acid ethyl ester was obtained in 22% yield as a pale yellow gum. MS: (M+H)⁺ 551.1. After hydrolysis, 2-[(3-fluoro-phenyl)-4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl]-amino)-methyl]-benzoic acid was obtained in 29% yield as a pale
20 yellow gum. MS: (M+H)⁺ 523.2.

Example 10

2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-m-tolyl-amino)-methyl]-benzoic acid ethyl ester was obtained in 33% yield as a pale yellow gum. MS: (M+H)⁺ 547.2. After hydrolysis, 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-m-tolyl-amino)-methyl]-benzoic acid was obtained in 43% yield as a pale yellow gum. MS:
25 (M+H)⁺ 519.3.

Example 11

2-[(4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-(3-trifluoromethyl-phenyl)-amino)-methyl]-benzoic acid ethyl ester was obtained in 19% yield as a pale yellow gum.
30 MS: (M+H)⁺ 601.1. After hydrolysis, 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-(3-trifluoromethyl-phenyl)-amino)-methyl]-benzoic acid was obtained in 31% yield as a pale yellow gum. MS: (M+H)⁺ 573.1.

- 28 -

Example 12

2-(((4-Fluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid ethyl ester was obtained in 17% yield as a pale yellow gum. MS: (M+H)⁺ 551.3. After hydrolysis, 2-(((4-fluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid was obtained in 70% yield as a pale yellow gum. MS: (M+H)⁺ 523.2.

Example 13

2-(((4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-p-tolyl-amino)-methyl)-benzoic acid ethyl ester was obtained in 31% yield as a pale yellow gum. MS: (M+H)⁺ 547.2. After hydrolysis, 2-(((4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-p-tolyl-amino)-methyl)-benzoic acid was obtained in 81% yield as a pale yellow gum. MS: (M+H)⁺ 519.2.

Example 14

2-(((4-Methoxy-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid ethyl ester was obtained in 37% yield as a pale yellow gum. MS: (M+H)⁺ 563.4. After hydrolysis, 2-(((4-methoxy-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid was obtained in 43% yield as a pale yellow gum. MS: (M+H)⁺ 535.3.

Example 15

2-(((3,4-Dimethyl-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid ethyl ester was obtained in 31% yield as a pale yellow gum. MS: (M+H)⁺ 561.4. After hydrolysis, 2-(((3,4-dimethyl-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid was obtained in 12% yield as a pale yellow gum. MS: (M+H)⁺ 533.3.

Example 16

2-(((3,4-Difluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid ethyl ester was obtained in 9% yield as a pale yellow gum. MS: (M+H)⁺ 569.2. After hydrolysis, 2-(((3,4-difluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid was obtained in 57% yield as a pale yellow gum. MS: (M+H)⁺ 541.2.

- 29 -

Example 17

2-(((4-Fluoro-3-methyl-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid ethyl ester was obtained in 10% yield as a yellow gum. MS: (M+H)⁺ 565.4. After hydrolysis, 2-(((4-fluoro-3-methyl-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl)-benzoic acid was obtained in 39%
5 yield as a white solid. MS: (M+H)⁺ 537.5.

Preparation of the starting material of examples 18 -26{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amine

To a solution of 0.50 g 4-hydrox-phenylaniline, 0.83 g of 2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethan-1-ol and 1.06 g of triphenylphosphine in 20 ml of THF was added at 0°C
10 a solution of 0.82 g of diisopropyl azodicarboxylate in 10 ml of THF over 30 min and stirring was continued at 22°C for 5 h. The mixture was evaporated and the residue purified by chromatography (SiO₂, n-hexane/AcOEt 6:1) to give 0.62 g of the title compound as a colorless solid. MS: (M+H)⁺ 371.4.

15

General description for the preparation of examples 18 – 26

To a suspension of 0.76 mmol of powdered KOH in 2.5 ml of DMSO was added at 22°C 0.19 mmol of the amine and the suspension was stirred for 5 min. A solution of 0.38 mmol of the corresponding benzylbromide in 0.5 ml of DMSO was added slowly keeping the
20 temperature at 15-20°C and stirring was continued at 22°C until the conversion was complete. The pH of the reaction mixture was adjusted to 2-3 using formic acid and the product was purified by preparative HPLC chromatography (RP-18, CH₃CN/H₂O gradient) to give the product as the ester.

The esters were hydrolyzed as described for examples 1 – 7.

25

Example 18

Starting from the amine and 2-bromomethyl-3-fluoro-benzoic acid methyl ester, 3-fluoro-2-(((4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amino)-methyl)-benzoic acid methyl ester was obtained in 55% yield as a pale yellow oil. MS: (M+H)⁺ 537.3. After hydrolysis, 3-fluoro-2-(((4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amino)-methyl)-benzoic acid was obtained in 90% yield as a pale brown
30 solid. MS: (M+H)⁺ 523.1.

- 30 -

Example 19

Starting from the amine and 2-bromomethyl-3-chloro-benzoic acid methyl ester, 3-chloro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid methyl ester was obtained in 43% yield as a pale yellow oil. MS: (M+H)⁺ 553.1 and 555.3 (Cl isotopes). After hydrolysis, 3-chloro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid was obtained in 74% yield as a colorless solid. MS: (M-H) 537.1 and 539.3 (Cl isotopes).

Example 20

Starting from the amine and 2-bromomethyl-3-trifluoromethyl-benzoic acid methyl ester, 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-3-trifluoromethyl-benzoic acid methyl ester was obtained in 59% yield as a colorless oil. MS: (M+H)⁺ 587.2. After hydrolysis, 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-3-trifluoromethyl-benzoic acid was obtained in 87% yield as colorless solid. MS: (M-H) 571.0.

Example 21

Starting from the amine and 2-bromomethyl-3-methoxy benzoic acid methyl ester, 3-methoxy-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid methyl ester was obtained in 36% yield as a colorless oil. MS: (M+H)⁺ 549.2. After hydrolysis, 3-methoxy-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid was obtained in 77% yield as a pale yellow solid. MS: (M-H) 533.2.

Example 22

Starting from the amine and 2-bromomethyl-4-chloro-benzoic acid methyl ester, 4-chloro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid methyl ester was obtained in 15% yield as a pale yellow oil. MS: (M+H)⁺ 553.2 and 555.1 (Cl isotopes). After hydrolysis, 4-chloro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid was obtained in 72% yield as a yellow solid. MS: (M-H) 537.1 and 539.2 (Cl isotopes).

Example 23

Starting from the amine and 2-bromomethyl-4-methoxy-benzoic acid methyl ester, 4-methoxy-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid methyl ester was obtained in 14% yield as a colorless oil. MS: (M+H)⁺ 549.2. After hydrolysis, 4-methoxy-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-

- 31 -

ethoxy]-phenyl}-phenyl-amino)-methyl]-benzoic acid was obtained in 93% yield as a pale brown solid. MS: (M-H) 533.2.

Example 24

Starting from the amine and 2-bromomethyl-5-fluorobenzoic acid methyl ester, 5-fluoro-
 5 2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-
 benzoic acid methyl ester was obtained in 44% yield as a colorless oil. MS: (M+H)⁺ 537.3.
 After hydrolysis, 5-fluoro-2-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-
 phenyl-amino)-methyl]-benzoic acid was obtained in 96% yield as a pale yellow foam. MS:
 (M-H) 521.1.

10

Example 25

Starting from the amine and 2-bromomethyl-benzoic acid ethyl ester, 2-[(4-[2-(5-
 methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-benzoic acid ethyl
 ester was obtained in 48% yield as a pale yellow oil. (M+H)⁺ 533.4. After hydrolysis, 2-
 [(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-methyl]-
 15 benzoic acid was obtained in 29% yield as a colorless solid. (M-H) 503.2.

Example 26

Starting from the amine and 2-bromomethyl-6-methoxy-benzoic acid ethyl ester, 2-
 methoxy-6-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl)-phenyl-amino)-
 methyl]-benzoic acid ethyl ester was obtained in 67% yield as a colorless oil. (M+H)⁺
 20 563.3. After hydrolysis, 2-methoxy-6-[(4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-
 phenyl)-phenyl-amino)-methyl]-benzoic acid was obtained in 29% yield as a colorless oil.
 (M+H)⁺ 535.3.

Example A

Tablets comprising the following ingredients can be manufactured in a conventional
 25 manner:

| <u>Ingredients</u> | <u>Per tablet</u> |
|-----------------------|-------------------|
| Compound of formula I | 10.0 - 100.0 mg |
| Lactose | 125.0 mg |
| Maize starch | 75.0 mg |
| Talc | 4.0 mg |
| Magnesium stearate | 1.0 mg |

- 32 -

Example B

Capsules comprising the following ingredients can be manufactured in a conventional manner:

| <u>Ingredients</u> | <u>Per capsule</u> |
|-----------------------|--------------------|
| Compound of formula I | 25.0 mg |
| Lactose | 150.0 mg |
| Maize starch | 20.0 mg |
| Talc | 5.0 mg |

5


Example C

Injection solutions can have the following composition:

| | |
|-------------------------------|-----------|
| Compound of formula I | 3.0 mg |
| Gelatine | 150.0 mg |
| Phenol | 4.7 mg |
| Water for injection solutions | ad 1.0 ml |

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OF PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A compound of formula



wherein

R¹ is aryl;

R² is hydrogen, alkyl or cycloalkyl;

R³ is hydrogen, alkyl, aralkyl, aryl, alkylcarbonyl, arylcarbonyl, alkyl-S(O)₂- or aryl-S(O)₂-;

R⁴ is aralkyl;

R⁵, R⁶, R⁷ and R⁸ are independently selected from hydrogen, alkyl and cycloalkyl;

n is 1, 2, 3, 4 or 5;

or a pharmaceutically acceptable salt or ester thereof,

wherein each aryl, alone or in combination is optionally substituted with one or more substituents independently selected from the group consisting of halogen, trifluoromethyl, amino, alkyl, alkoxy, alkylcarbonyl, cyano, carbamoyl, alkoxy-carbamoyl, methylenedioxy, carboxy, alkoxy-carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, hydroxyl and nitro.

2. The compound according to claim 1, wherein R¹ is phenyl.
3. The compound according to claim 1 or 2, wherein R² is alkyl.
4. The compound according to claim 3, wherein R² is methyl.
5. The compound according to any one of claims 1 to 4, wherein R³ is alkyl or phenyl, wherein phenyl is optionally mono- or disubstituted with halogen.
6. The compound according to claim 5, wherein R³ is propyl, phenyl, fluorophenyl or difluorophenyl.
7. The compound according to any one of claims 1 to 6, wherein R⁴ is arylmethyl.
8. The compound according to claim 7, wherein R⁴ is benzyl substituted with carboxy and optionally further substituted with fluoro, chloro, trifluoromethyl or alkoxy.
9. The compound according to any one of claims 1 to 8, wherein R⁵, R⁶, R⁷ and R⁸ are hydrogen.
10. The compound according to any one of claims 1 to 9, wherein n is 2.
11. The compound according to any one of claims 1 to 10 selected from

5

2-[[{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amino)-methyl]-benzoic acid;

10

2-[[{4-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propyl-amino)-methyl]-benzoic acid;

2-[[{(3-Fluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;

2-[[{(3,4-Difluoro-phenyl)-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-amino)-methyl]-benzoic acid;

15


3-Fluoro-2-[[{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amino)-methyl]-benzoic acid; and

5-Fluoro-2-[[{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-phenyl-amino)-methyl]-benzoic acid.

20

12. A process for the preparation of the compound defined in any one of claims 1 to 11, comprising one of the following reactions

a) reaction of a compound of formula (5)




- 36 -

in the presence of R^3 -Hal and following reaction in the presence of a hydroxide in order to obtain the compound defined in any one of claims 1 to 11, wherein Hal means chlorine, bromine or iodine and R^1 to R^8 and n are defined as in claim 1;

- 5 b) reaction of the compound of formula (5) in the presence of $(HO)_2B$ -aryl and following reaction in the presence of a hydroxide in order to obtain the compound defined in any one of claims 1 to 11, wherein R^1 , R^2 , R^4 to R^8 and n are defined as in claim 1;

c) reaction of a compound of formula (7)



(7)

- 10 in the presence of R^4 -Hal and following reaction in the presence of a hydroxide in order to obtain a compound according to any one of claims 1 to 11, wherein Hal means chlorine, bromine or iodine and R^1 to R^8 and n are defined as in claim 1.
13. The compound according to any one of claims 1 to 11 for use in the prophylaxis and/or therapy of a disease modulated by a $PPAR\alpha$ agonist and/or a $PPAR\gamma$ agonist.
- 15 14. The compound according to any one of claims 1 to 11 for the preparation of a medicament for the prophylaxis and/or therapy of a disease which is modulated by a $PPAR\alpha$ agonist and/or a $PPAR\gamma$ agonist.
15. A pharmaceutical composition comprising the compound defined in any one of claims 1 to 11 and a therapeutically inert carrier.
- 20 16. The pharmaceutical composition of claim 15 further comprising a lipase inhibitor.
17. The pharmaceutical composition according to claim 16, wherein the lipase inhibitor is orlistat.

18. A use of the compound defined in any one of claims 1 to 11 for the preparation of a medicament for the treatment and/or prophylaxis of a disease which is modulated by a PPAR α agonist and/or a PPAR γ agonist.
19. The compound according to any one of claims 1 to 11, when manufactured according to the process of claim 12.
20. A use of the compound defined in any one of claims 1 to 11 for the treatment and/or prophylaxis of a disease which is modulated by a PPAR α agonist and/or a PPAR γ agonist.
21. A use of the compound defined in any one of claims 1 to 11 with a lipase inhibitor, for simultaneous, separate or sequential administration, in the treatment and/or prophylaxis of a disease modulated by a PPAR α agonist and/or a PPAR γ agonist.
22. The use according to claim 21, wherein the lipase inhibitor is orlistat.
23. A use of the compound defined in any one of claims 1 to 11 and a lipase inhibitor, separately or in combination, in the manufacture of one or more medicaments for administration simultaneously, separately or sequentially in the treatment and/or prophylaxis of a disease which is modulated by a PPAR α agonist and/or a PPAR γ agonist in a patient.
24. The use according to claim 23, wherein the lipase inhibitor is orlistat.
25. The use according to any one of claims 18 and 20-24, wherein the disease is diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, an atherosclerotic disease or metabolic syndrome.
26. The use of claim 25, wherein the disease is non-insulin dependent diabetes mellitus.

27. The compound according to claim 13 or 14, wherein the disease is diabetes, non-insulin dependent diabetes mellitus, elevated blood pressure, increased lipid and cholesterol levels, an atherosclerotic disease or metabolic syndrome.
28. The compound according to claim 27, wherein the disease is non-insulin dependent diabetes mellitus.

