

(19) **DANMARK**

(10) **DK/EP 2220107 T3**



(12)

Oversættelse af
europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 K 1/22 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2017-02-13**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-11-02**
- (86) Europæisk ansøgning nr.: **08850855.1**
- (86) Europæisk indleveringsdag: **2008-11-12**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-08-25**
- (86) International ansøgning nr.: **EP2008065346**
- (87) Internationalt publikationsnr.: **WO2009062942**
- (30) Prioritet: **2007-11-12 EP 07120454**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **CHRETO ApS, Lejrvej 17, 1., 3500 Værløse, Danmark**
- (72) Opfinder: **KYHSE-ANDERSEN, Jan, Dalsøe Park 8, 3500 Værløse, Danmark**
- (74) Fuldmægtig i Danmark: **Inspicos P/S, Kogle Alle 2, 2970 Hørsholm, Danmark**
- (54) Benævnelse: **Polypeptid med dobbelt affinitet til oprensning**
- (56) Fremdragne publikationer:
EP-A1- 0 962 771
WO-A2-01/95857
US-A- 5 328 985
SHINAGAWA K ET AL: "Purification of staphylococcal enterotoxins A and E by immunoaffinity chromatography using a murine monoclonal antibody with dual specificity for both of these toxins" JOURNAL OF IMMUNOLOGICAL METHODS, ELSEVIER SCIENCE PUBLISHERS B.V.,AMSTERDAM, NL, vol. 139, no. 1, 17 May 1991 (1991-05-17), pages 49-53, XP023974529 ISSN: 0022-1759 [retrieved on 1991-05-17]

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates to a process for purification of a target biomolecule on a solid support comprising the steps: (a) contacting (i) a target biomolecule, (ii) a dual affinity polypeptide, and (iii) a solid support comprising a catching ligand, as defined in the claims.

BACKGROUND OF THE INVENTION

[0002] Recovery and purification of therapeutic proteins accounts for approximately 50% of the manufacturing cost of biological drugs. The general industrial purification process often includes a number of unit operation steps, like extraction, precipitation, as well as anion- and cation-exchange chromatography. Affinity chromatography is the preferred downstream process step due to its high recovery, yield and specificity, but the current cost and limitations of affinity chromatography is very substantial and in many cases prohibitive for a more general use of this unit operation. For a general description of conventional purification procedures including affinity chromatography see e.g. Jason and Rydén 1998 (Jason, J-C and Rydén, L., Protein Purification: Principles, high-Resolution, Methods and Applications, 2nd edition, Wiley & sons Inc. New York, 1998).

[0003] Conventional affinity chromatography is in general characterized by having a capturing ligand immobilised to a solid phase matrix. The ligand reversibly binds a target molecule present in a fluid such as liquid culture medium or serum. Target molecules are recovered by dissociating the complex at eluting conditions. Commercially available affinity matrices are in a ready to use format including capturing ligands covalently attached to the matrices. In conventional affinity chromatography the dissociation constant, K_D , between the ligand and the target protein is in the range of about 10^{-5} - 10^{-7} M. Interactions with dissociation constants exceeding 10^{-10} - 10^{-11} M are often impossible to use, as the conditions required to dissociate the complex are then the same as those that will result in denaturation of the target proteins. The prior art include alternative variations of affinity chromatography purification methods described in the literature (Wilchek, M. and Gorecki, M. (1973), A New Approach for isolation of Biologically Active Compounds by Affinity Chromatography: Isolation of Trypsin).

[0004] FEBS Letters. 31, 1, 149-152, describes antibodies immobilized on an insoluble material. The antibodies have affinity for a certain ligand attached to a complex of two or more proteins, and are independent of the chemical, physical and biological properties of the complex itself. The immobilized antibody matrix serves as means for concentrating the complex. The adsorbed complex can then be recovered from the column by elution. The authors use the trypsin enzyme reacted with dinitrophenylated soybean trypsin inhibitor (DNP-STI) to form the complex. The complex is adsorbed to anti DNP-column and eluted under conditions that dissociate the antigen-antibody binding. The affinity column is then ready for the next purification cycle. The target trypsin is obtained by separation of the trypsin enzyme - dinitrophenylated soybean trypsin inhibitor complex into its components in a later step.

[0005] This procedure is different from the present invention in that the affinity column is reusable and it is the binding between the immobilized agent and the linker that is dissociated during elution and not the bond between linker and target biomolecule.

[0006] Another concept described by Hammarbergh, B. et al., (Proc.Natl.Acad.Sci USA, 86, 4367-4371 (1989)), is a fusion protein affinity approach and its use to express recombinant human insulin-like growth factor II. The procedure relates to a recombinant target protein of interest (X) fused between two different affinity protein tails (A and B). The protein (X) has a protease-sensitive site. A cell lysate containing the recombinant tripartite fusion protein is first passed through an affinity column containing a tail B-specific ligand. A mixture of full-length protein and proteolytic fragments containing the C-terminal fusion protein region can thus be obtained. In a second passage through a tail A-specific affinity column, the degraded proteins flow through while full-length fusion protein is retained. After site-specific cleavage of the tails, the protein of interest (X) is obtained by passing the cleavage mixture through a mixed affinity column for tails A and B and collecting the flow-through. The authors describe a procedure to obtain the target protein by expressing the target protein as an integrated part in between a dual affinity protein construct.

[0007] This is different from the present invention as the described affinity procedure requires two different affinity columns and that the immobilized ligand on the column and the dual affinity fusion protein is dissociated to recover the target biomolecule. Following the elution step and a regeneration procedure, the affinity columns are ready for the next affinity purification cycle. The

target protein is only part of the fusion protein and is obtained following enzymatic degrading steps.

[0008] In a review article by, Ford, C.F., Suominen, I, Glatz, C.E. (1991) Fusion Tails for the Recovery and Purification of Recombinant Proteins. Protein Expression and Purification, 2, 95-107, the authors discuss the applications and advantages of using fusion tail systems to promote efficient recovery and purification of recombinant proteins from crude cell extracts or culture media. In these systems, a target protein is genetically engineered to contain a C- or N-terminal polypeptide tail, which provides the biochemical basis for specificity in recovery and purification. Fusion tails are useful for enhancing recovery methods for industrial downstream processing. Nevertheless, for the purification of target proteins a site for specific enzymatic cleavage is included, allowing removal of the tail after recovery. The article describes the application of fusion proteins with one binding partner having affinity for the ligand immobilized on a matrix. The procedures include an enzymatic cleavage step to recover the target protein from the fusion tail as required.

[0009] This is different from the present invention as the described affinity procedure requires that the fusion protein is dissociated from the ligand immobilized on the column matrix to recover the protein. Following the elution step and a regeneration procedure, the affinity column is ready for the next affinity purification cycle. Also, different from the present invention is that the target protein is part of the fusion protein and is only obtained following an enzymatic processing step.

[0010] In Rigaut, G. et al. (1991)(A Generic Protein Purification Method for Protein Complex Characterization and Proteom Exploration. Nature Biotechnology, 17, 1030-1032), is described a generic procedure for purification of protein complexes using tandem affinity purification (TAP) tag. The purification requires one affinity step followed by an enzymatic step cleaving the first affinity tag from the complex and a second affinity purification step to recover the target protein complex from the protease. Overall, the method involves two binding partners in combination both for binding to a ligand immobilized to a column matrix and a protease cleavage step to expose the second binding partner.

[0011] This is different from the present invention as the described affinity procedure requires that the fusion protein is dissociated from the ligand immobilized on the column matrix to recover protein. Following the elution step and a regeneration procedure, the affinity column is ready for the next affinity purification cycle. Also, different from the present invention is that the target protein is part of the fusion protein and is obtained following an enzymatic processing step.

[0012] EP1529844 describes a method for altering the properties of a recombinant target protein involving co-expression of target protein and the binding partner. The target protein and the binding partner form a complex in the cell. The complex formation result in altered properties such as accumulation, stability and/or integrity, sub-cellular localization, post-translational modifications, purification, and phase partitioning behavior of natural or recombinant target proteins expressed in a host organism. The binding partner may provide an affinity tag that enables co-purification of the complex and the target protein contained therein.

This description is different from the present invention as it describes a co-expression of the binder and the target in order to form a complex in the cell. The disclosed method is for alteration of the target protein properties in general, whereas the present invention describes a dual affinity polypeptide specifically designed to facilitate a dedicated purification process, wherein the dual affinity polypeptides needs to possess specific binding properties.

[0013] Linder et al., (Linder, M., Nevanen, T., Söderholm, L., Bengs, O. and Teeri, T., 1998, Biotechnology and Bioengineering, 60(5): 642-647) describes the use of CBD in fusionproteins for use as an affinity tag for purification. Some leakage from the column was observed.

[0014] Shpigel, E. et al. (Biotechnol. Appl. Biochem. (2000) 31, 197-203, "Expression, purification and application of Staphylococcal Protein A fused to cellulose-binding domain"), describes an example of purifying IgG using Protein A-CBD dual affinity polypeptide.

[0015] They claim that they save expensive coupling procedures by choosing immobilization of the Protein A functionality to a solid phase through the cellulose-binding domain (CBD) of a fusion protein. The fusion protein is immobilized on the column before adding the target.

[0016] Due to leakage problems this choice of dual affinity molecule is unsuitable for biopharmaceutical applications.

[0017] Sano et al. (US patent 5,328,985) describes a fusion protein consisting of streptavidin and one or two immunoglobulin G (IgG) binding domains of protein A expressed in Escherichia coli. The streptavidin-protein A (ST-PA) fusion protein has functional biotin and IgG binding sites. Sano further describes complexes of the streptavidin-protein A fusion protein, a monoclonal antibody to bovine serum albumin (BSA) and biotinylated horseradish peroxidase. Sano also describes a method of labelling cell using the

ST-PA fusion protein. Cells are incubated with an antibody to the cell surface antigen, Thy-1. The chimeric protein biotinylated marker complex is subsequently added to the cell suspension. This technique was used to deliver biotinylated FITC to the surface of the cells having Thy-1 antigens on their surface.

[0018] However, Sano does not describe or suggest using the ST-PA fusion protein as a tool for purification purposes nor does he describe a procedure of single use affinity chromatography column materials, nor recovery of a target protein.

[0019] WO 97/19957 describes an invention related to delivering toxins or nucleic acids into specific cell types using ST-PA fusion proteins for the purpose. Similar to Sano et al. (vide supra), an antibody recognise a surface antigen on the cell surface. The ST-PA binds to the antibody and facilitates a linkage to a biotinylated toxin bound to the biotin-binding site. However, it is not described or suggested to use the ST-PA fusion protein as a tool for purification purposes.

[0020] WO 01/95857 discloses a method and components for extracting toxic substances from mammalian blood. The method includes preparing an affinity column (~extracorporeal device) and a procedure for extracorporeal extraction of toxic material from mammalian body fluids in connection with diagnosis or treatment of a mammalian condition or disease.

The extracorporeal affinity column exemplified in the patent is made by coupling biotinylated entities to a matrix containing immobilized avidin. The biotinylated entity includes a part that binds strongly to the toxin in the mammalian blood. The toxic material is removed (i.e. immobilized but not recovered by elution from the column) from the blood following a conventional affinity chromatography procedure. The product from the flow through chromatography procedure is purified blood as the target (toxic materials) stays immobilized on the column after the process.

This is different from the present invention as it describes a procedure that bind the target tightly with high affinity in order to remove target from the product. The purification procedure is also different from the present invention as the product does not bind to the affinity column, but flows through and is collected as depleted from the toxic material (the target). The toxic material is not released or recovered.

[0021] WO 97/09068 discloses a method and chemical components that alter the equilibrium dissociation constant between two pairs of bio-molecules. The chemical component is a polymer that can be stimulated to change conformation and thus binding efficiency. The polymer is coupled e.g. to a specific site of the binding partner (the ligand) immobilised to the matrix of the affinity chromatography column. WO 97/09068 does not describe or suggest the use of a dual affinity component for affinity purification, nor recovering of target molecules.

[0022] In general, methods that will improve the capturing efficiency and simplify the purification process as well as reduce costs are desirable.

SUMMARY OF THE INVENTION

[0023] The present invention significantly improves and simplifies the downstream processing and lowers the cost of affinity chromatography processes in general. The present invention includes a generic capturing ligand immobilised to a matrix, a target biomolecule and a semi generic dual affinity polypeptide with different binding affinity toward the target and the capturing ligand respectively. The dual affinity polypeptide reacts with the target biomolecule to form a complex of medium binding affinity, and the complex binds non-covalently to a generic affinity matrix with a strong binding affinity. The target biomolecule is recovered by specific elution from the generic matrix leaving the dual affinity polypeptide attached to the capturing ligand on the matrix, due to the tight binding to the ligand preventing leakage from the solid phase matrix.

[0024] In a first aspect the present invention provides a process for purification of a target biomolecule, comprising the steps: (a) contacting (i) a target biomolecule, (ii) a dual affinity polypeptide, and (iii) a solid support comprising a catching ligand or dual affinity polypeptide binding site, wherein the ratio between the equilibrium dissociation constants of the dual affinity polypeptide, $[K_{D,T} / K_{D,S}]$, is at least 10^0 at standard conditions; and (b) recovering the target biomolecule by elution.

[0025] In a second aspect the present invention provides a process for purification of a target biomolecule, comprising the steps: (a) contacting (i) a target polypeptide, (ii) a dual affinity polypeptide, and (iii) a solid support comprising a catching ligand, wherein the dual affinity polypeptide has an equilibrium dissociation constant, $K_{D,T}$ towards the target biomolecule in the range from 10^{-2} to 10^{-13} M, more particularly from 10^{-4} to 10^{-13} M at standard conditions, and wherein binding of the dual affinity polypeptide to the catching ligand on the solid support is provided by cleavage of a para-substituted benzyl guanine resulting in a thioether bond; and (b) recovering the target biomolecule by elution.

DETAILED DESCRIPTION OF THE INVENTION

[0026] In conventional affinity chromatography the capturing ligand is attached directly to the support. The main technical challenges are to optimize the entire system with regard to e.g. ligand coupling, nature of the support material, flow, backpressure and physical dimensions of the column. It should be understood that several of the technical limitations in high performance affinity columns are closely linked, making performance and cost optimization as well as scale up difficult. The ligand, in traditional affinity chromatography, preferably possesses the following characteristics:

1. a) The ligand should have chemical properties that allow easy covalent attachment to the matrix.
2. b) The ligand must be able to form a reversible complex with the target molecule.
3. c) The specificity of the ligand's affinity for the target molecule must be appropriate for the planned application.
4. d) The dissociation constant for the ligand-target molecule complex under "loading conditions" should be strong enough to enable formation of stable complexes or to give sufficient retardation in the elution of the target molecule.
5. e) It should be easy to dissociate the ligand from the target molecule by changing the conditions, e.g. pH or salt concentration, without irreversibly damaging either.

[0027] Furthermore, in traditional affinity chromatography, the ligand is normally covalently attached to the matrix and is also the component binding to the target molecule.

[0028] The capacity and quality of purification is greatly influenced by the contact time between target and ligand in the affinity column, the so-called residence time.

[0029] In addition to the association rate of the target protein to an immobilized ligand, diffusion into the pores within the chromatography beads in the column and mass transfer of the protein from the solute will impact the dynamic binding capacity of a chromatography matrix.

[0030] The mass transfer of the target protein from the solute depends on a variety of factors, including type and degree of cross-linking, compressibility of the support material, the size of the pores and the physical size of the target protein.

[0031] Flow rates, protein concentrations, column length, temperature, buffer, conductivity, and pH can also influence on pore diffusion and the dynamic binding capacity of the adsorbent.

[0032] Due to the requirement for rapid development of downstream processes and regulatory constraints, the residence time for a particular type of biological product such as for example a therapeutic antibody is typically fixed in the early development. Therefore, often the flow rate in the large scale column with e.g. larger bed height is tried adjusted to maintain the desired residence time used during the small scale development.

[0033] Due to technical constraints and the major investment required to purchase process-scale chromatography equipment, the scale up of traditional high performance affinity chromatography is a major challenge.

[0034] The present invention suggests a more simple and flexible scale up process with less technical constraints.

[0035] Furthermore, conventional affinity chromatography is characterised by regeneration procedures to provide repeated uses of the column materials. These cleaning procedures require extensive validation to allow multiple uses of the column.

[0036] The present invention differs in several aspects from the conventional affinity chromatography e.g.

- the immobilized ligand binds tightly to the dual affinity polypeptide (DAP) in order to prevent dissociation at elution conditions
- it is intended for single use applications

[0037] It is clear from the above that the role of the ligand in the present invention is to bind the DAP molecule and not the target molecule.

[0038] The attractive benefit of affinity chromatography is that it provides a large increase in purity with a minimal loss of target molecule material in a single unit operation. However, affinity chromatography is also characterised by the high cost prohibiting the use of large columns and thus favouring repetitive use of smaller columns. This leads to extended production processes and capacity loss proportional to the number of column reuses, increased loss and/or modification of the target molecule. In principle a typical affinity chromatography matrix can be used for up to 100 or more runs, but the average number of runs in manufacturing scale appears to be several fold lower. One of the reasons that the matrix is discarded long before the end of its theoretical lifespan is that the affinity columns used in manufacturing are dimensioned to process the entire fermentation batch in far less than 100 runs - in order to save cost, but also reduce the risk of contamination and handling failures. Rather than using the same matrix for several fermentation batches, the matrix may be discarded after processing of one fermentation batch, which leads to the relatively low number of average runs on an affinity matrix.

[0039] Controlling the flow rate through an affinity chromatography support is important in achieving binding. Flow rate through the column support is inextricably related to the efficiency of the separation; too fast a flow will cause the mobile phase to move past the beads faster than the diffusion time necessary to reach the internal bead volume. For each application a flow rate can be selected to achieve an optimal balance between efficient binding and elution of the target protein and a fast separation. Gravity driven flow chromatography is very slow and resolution of the protein separation can be adversely affected by secondary diffusion effects. Therefore modern systems have active pumping to control flow rates and continuous monitoring of back pressure to ensure that the maximum operating back pressure is not being exceeded.

[0040] In conventional columns fouling is of major concern. Debris, proteins and salts can slowly build up fouling layers in the channels of high performance affinity chromatography supports resulting in changed flow rates, reduced mass transfer rates, increased back pressure and hidden and deactivated affinity ligands. Especially lipids and lipoproteins material can rapidly clog chromatography columns and it is often necessary to remove them before affinity purification. This is especially important for samples derived from ascites fluid.

[0041] This pre purification step can be done by precipitation steps with for example dextran sulphate and polyvinylpyrrolidone followed by centrifugation and dialysis or desalting. The step can result in the loss of 5-10% of the target protein.

[0042] Omitting the delipidation step may be possible for the affinity purification system of the present invention as the column is single use. This will result in a higher overall target recovery and a more efficient downstream processing workflow.

[0043] Because of the intended single use of the columns according to the invention the elution step is also simplified. As the support material of the invention is not to be reused, one can more freely select elution conditions. For example, it is possible to select any elution buffers with an unconventional high concentration of salt, strong chaotropes, organic solvents etc. which will allow the recovery of the intact target and leave the DAP molecule attached to the support. It is of no importance if the properties of the support material are irreversibly changed with respect to e.g. structure and flow characteristics and cannot be reused.

[0044] This flexibility in selecting elution conditions is often not possible when using traditional high-performance affinity purification systems as the internal structures and surfaces are highly optimized and sensitive to polymer swelling or precipitation. Additionally, due to the cost of traditional affinity columns, the operators can be reluctant to test new elution conditions further reducing the flexibility in elution optimization.

[0045] The traditional operation of affinity purification includes cycles of equilibration, sample loading, elution and cleaning-in-place (CIP).

[0046] The cleaning steps or sanitization protocols have to be designed for each specific target purification. As described above, a major concern during operation is the build up of fouling layers or cross contamination between runs.

[0047] The cleaning step often includes using chemically harsh buffers like 0.1 M NaOH/1 M NaCl or 0.1 M phosphoric acid in a combination with sodium chloride or ethanol, followed by regeneration. It is well established that in general the dynamic binding capacity decreases as the number of CIP cycles increases. Therefore, one needs to find an optimum between the quality of the purified target, the number of runs, CIP's and the size and cost of the particular column. Further, the change of purification quality needs to be monitored for most pharmaceutical purifications.

[0048] As suggested by the present invention, the cost of this quality validation and the optimization of the CIP and runs can be greatly reduced.

[0049] In the conventional affinity chromatography outlined above for purification of e.g. monoclonal antibodies, the capturing ligand (Protein A) is attached to a solid phase matrix and has the affinity towards the target biomolecule (monoclonal antibodies). The present invention provides advantages compared to conventional affinity purification technologies for the downstream processing industry due to lower costs, high specificity and ease of use without compromising the quality of the downstream process. An essential feature of the present invention is the use of a dual affinity polypeptide as a linker between the target molecule and the solid support comprising a ligand. These dual affinity polypeptides are particularly useful for the downstream processing of biopharmaceutical and diagnostic proteins and peptides.

[0050] The invention suggests the improvement of the entire method of affinity purification by eliminating several of the constraints in current systems.

[0051] By using the dual affinity polypeptide (DAP) and the generic supports of the invention, the majority of the above problems and limitations can be completely eliminated or reduced.

[0052] According to the present invention the dual affinity purification technology is characterized by a generic solid support, which in one embodiment is a solid phase matrix, plus ready-to-use specific dual affinity polypeptides serving as linker molecules. A dual affinity polypeptide reacts with the target biomolecule. The dual affinity polypeptide - target biomolecule complex subsequently connects non-covalently to a capturing ligand immobilized on a solid support by contacting the complex and the solid support. The target biomolecule is recovered by specific elution. The dual affinity polypeptide remains attached to the ligand on the solid support during elution.

[0053] In one aspect the present invention therefore relates to a process for purification of a target biomolecule, comprising the steps: (a) contacting (i) a target biomolecule, (ii) a dual affinity polypeptide, and (iii) a solid support comprising a catching ligand or dual affinity polypeptide binding site, wherein the ratio between the equilibrium dissociation constants of the dual affinity polypeptide, $[K_{D,t} / K_{D,s}]$, is at least 10^0 at standard conditions; and (b) recovering the target biomolecule by elution.

[0054] The dual affinity polypeptide acts as the linking partner between the solid support and the target molecule. In one particular embodiment the affinity of the dual affinity polypeptide towards the immobilized ligand is stronger than the affinity towards the target molecule. Furthermore this difference in binding affinity, can be expressed as the ratio between the equilibrium dissociation constants. In one embodiment this ratio is at least 1.

[0055] The dual affinity polypeptide according to the invention comprises at least two binding sites, of which one binding site has affinity for the ligand and another binding site has affinity for the target molecule. These binding sites are polypeptide based meaning that they comprise either complete proteins or fragments of proteins. Such fragments should at least comprise the part of the protein containing the binding site for the specific target. The dual affinity polypeptide could be a fusion polypeptide or could be two or more polypeptides chemically linked in any suitable way e.g. by a linker segment.


[0056] Therefore the present invention in further embodiments relates to a dual affinity polypeptide having an equilibrium dissociation constant towards a target biomolecule, $K_{D,t}$ in the range from 10^{-2} to 10^{-13} M, e.g. 10^{-8} M, and an equilibrium dissociation constant towards a catching ligand, $K_{D,s}$ in the range from 10^{-9} to 10^{-16} M, e.g. 10^{-10} M, and at the same time the ratio, $K_{D,t}/K_{D,s}$, should be matched such that the ratio is at least 10^0 , more particularly at least 10^1 , more particularly 10^2 , more particularly 10^3 and even more particularly 10^4 .

[0057] The above in other words means that binding of DAP to the target is in preferred embodiments weaker than binding of DAP to the ligand.

[0058] Particularly the said dual affinity polypeptide has an equilibrium dissociation constant, $K_{D,t}$ towards the target polypeptide in the range from 10^{-4} to 10^{-13} M, more particularly in the range from 10^{-6} to 10^{-13} M, and an equilibrium dissociation constant, $K_{D,s}$ towards the catching ligand in the range from 10^{-9} to 10^{-16} M, more particularly in the range from 10^{-11} to 10^{-16} M.

[0059] In general the binding towards the ligand or the column cannot be too strong. Therefore the value at the upper end of the range is not important in respect of $K_{D,s}$.

[0060] In the context of the present invention the equilibrium dissociation constant are measured according to the reaction:



[0061] A and B represents the binding partners: the target biomolecule and the dual affinity polypeptide or the dual affinity polypeptide and the catching ligand immobilized on the solid phase matrix.

[0062] The rate constants for the reaction above represent the rate at which the two molecules A and B associates and dissociates

$$\text{Dissociation rate} \quad -\frac{d[AB]}{dt} = k_d [AB]$$

$$\text{Association rate:} \quad \frac{d[AB]}{dt} = k_a [A][B]$$

[0063] When the rates are equal at equilibrium $k_a[A][B] = k_d [AB]$, which gives

$$\frac{k_d}{k_a} = \frac{[A][B]}{[AB]} = K_D$$

$$\frac{k_a}{k_d} = \frac{[AB]}{[A][B]} = K_A$$

[0064] The candidate binding domains to be employed in the dual affinity polypeptide should be evaluated according to the apparent equilibrium dissociation constants based on the total binding affinity of each of the dual affinities in a given DAP molecule irrespective of whether it contains one or several binding domains for each specificity (target/capturing ligand). If e.g. A and B represent protein A (has four to five binding domains) and avidin (having four binding sites) respectively the above ranges should apply for one protein A molecule fused to one avidin molecule. However, this does not exclude the possibility that e.g. the DAP molecule could be composed of several binding candidates for the target and several candidates for the ligand on the matrix. The DAP could e.g. in another embodiment consist of 3 protein A molecules linked to one or more avidin molecules. Therefore the specified ranges as defined above should in the context of the present invention be evaluated based on the apparent binding constants for the binding domains in common.

[0065] In the context of the present invention the specified equilibrium dissociation constants are determined by surface plasmon resonance (SPR) technology using a Biacore Instrument as illustrated in detail in the examples. The conditions described herein represent the standard conditions. As a suitable starting point for selecting different binding domains to be combined in the DAP molecule published K_D 's may be used.

[0066] The two binding pairs should be selected based on the K_D 's during specific binding conditions, but also considering the planned elution conditions, when the target is recovered and the DAP molecule remains on the support.

[0067] As described above determination of dissociation affinities of various binding domains in the context of a DAP molecule was accomplished by using surface plasmon resonance (SPR). Such evaluation can be done with the Biacore system. Biacore has commercial instrumentation where measurements based on SPR make determinations on protein-protein interactions. The evaluation was conducted having the complete DAP immobilized on the sensor chip used in the Biacore instrument. The Biacore system defines the characteristics of proteins in terms of their specificity of interaction with other molecules, the rates at which they interact (association and dissociation), and their affinity (how tightly they bind to another molecule). This technique has been described e.g. for determining the binding interactions between specific antibodies and their target (see e.g. Rönmark, 2002, Eur. J. Biochem., 269: 2647-2655).

[0068] In the examples below several DAP candidates have been evaluated and their binding affinities under standard conditions (as described in the examples) have been measured for the complete DAP. Other methods may also be used, however, results may then differ. A list of alternative methods has been described below.

[0069] Quantitative measurement of non-covalent protein-ligand interactions is well known. The methods suited for quantitative measurement of binding constants of particular relevance for the present invention include various versions of surface plasmon resonance (SPR) and circular dichroism (CD).

[0070] Other methods include mass spectrometry methods for dynamic titrations like ESI-MS titration, HPLC-ESI-MS titration or MALDI-SUPREX titration.

[0071] Other methods are based on determining the dissociation constant of a ligand at a binding site indirectly by competitive displacement of a radioactive ligand or by measurement of NMR chemical shift as function of concentration, fluorescence spectroscopy analysis of e.g. signal quenching, X-ray crystallographic measurement of the ligand occupancy, isothermal calorimetry (ITC) or enzyme inhibition.

[0072] Yet other methods use labeled ligands, for example capillary electrophoresis with laser-induced fluorescence detection of enzyme labeled ligands.

[0073] Alternatively, binding constants can be found from computational techniques by using de novo design, data mining and sophisticated algorithms.

[0074] In the context of the present invention the appropriate ranges for the equilibrium dissociation constants as specified in the claims should apply to the complete dual affinity polypeptide and not to the individual binding parts measured separately.

Moreover, if a single candidate binding domain has a weaker binding affinity towards the target or ligand than required according to the present invention, it still could be applicable by combining several such candidate binding domains into one DAP.

This is due to the valence effect. It is possible to obtain an increased binding strength due to an avidity gain. Single domains with a low intrinsic affinity combined into multimers often generates avidity effects which lead to slower dissociation rates and increased functional affinities by more than 100 fold (MacKenzie, C.R. et al (1996), Analysis by surface plasmon resonance of the influence of valence on the ligand binding affinities and kinetics of and anti carbohydrate antibody. Journal of Biological Chemistry, 271, 1527-1533). It is possible to measure effects from monovalent and bivalent bindings, but at higher binding valences the situation becomes so complex that it is impossible to distinguish between different binding valences. Nevertheless relative data can be obtained and are used in the context of the present invention

[0075] The invention provides a purification procedure wherein the first reaction between the target molecule and the dual affinity polypeptide is completed in free solution. Reaction binding kinetics is about 1000 times faster in free solution compared to interface reactions (Nygren, H. and Stenberg, M. (1989) Immunochemistry at interfaces. Immunology, 66, 321-327).

[0076] The target molecule-dual affinity polypeptide complexes are subsequently presented to and bind efficiently to the ligand on the solid support. The strong binding (fast association rate and slow dissociation rate of the ligand towards the dual affinity polypeptide) depletes the mobile phase of target-DAP complexes. The target molecules are recovered from solution through this sequential procedure facilitated by the second binding functionality of the dual affinity polypeptide.

[0077] Due to the described differences in equilibrium dissociation constants the target polypeptide can be efficiently eluted without eluting the dual affinity polypeptide. Elution can in one embodiment be performed by changing either pH, ionic strength or chaotropic ions in solution, or any combination thereof.

[0078] The K_D value can be influenced by changing conditions like pH, ionic strength, temperature and polar properties. Unfortunately, the literature values for K_D are not always listed at relevant elution conditions. Though, the skilled in the art will be able to find elution conditions which will only break the weakest binding without disturbing the stronger binding in cases where the binding to the solid matrix is sufficiently strong (i.e. $K_{D,s} < 10^{-9}$ M and the ratio between K_D values is at least 1 when measured at standard binding conditions).

[0079] The criteria for selecting the target specific binding pairs of the invention resemble those for the traditional affinity chromatography with regard to dissociation constant, specificity, binding and possible elution conditions. However, since elution conditions are usually different from the conditions applied when measuring K_D 's on the Biacore instrument in the present invention the limits set for the applicable ranges of the two distinct binding affinities of the DAP has been determined under standard conditions, which equals the conditions used in the examples.

[0080] The criteria for selecting the specific ligand binding domains of the invention are somewhat different from the criteria used

in the traditional affinity chromatography, as the DAP molecule is not to be eluted from the support.

[0081] Binding domains which are specific and strong, but cannot be broken under normal elution conditions are not suited for traditional affinity chromatography. Such binding domains can be used in the present invention. Examples include the very specific biotin-Streptavidin binding, which for most practical applications cannot be reversed under elution conditions and consequently is well suited as one of the binding pairs of the invention.

[0082] In general, the binding between DAP and the ligand should be stronger than the binding between DAP and the target and strong enough to prevent leakage of the DAP molecule from the support during elution of the target.

[0083] Preferred ligand-DAP binding pairs are strong and exhibit no or little reduction in binding strength due to changing pH, ionic strength, solvents, chaotropic agents, temperature etc.

[0084] It should be clear that when changing the scale of purification, using the system of the invention, the amount of DAP added is adjusted to the amount and concentration of target protein. As the DAP molecule can be supplied as a concentrate, the binding conditions can be adjusted with respect to e.g. pH and salts. Also, the temperature and time can be selected to give the best binding and subsequently purification.

[0085] The size and capacity of the generic column is selected to be large enough to capture the DAP molecules. Potentially several columns are used in parallel or in a bundle.

[0086] If purifying another target, another appropriate DAP molecule is selected. The same or another column can be used.

[0087] In one embodiment the dual affinity polypeptide is a fusion polypeptide. Such fusion polypeptides can either be prepared by chemically linking two appropriate proteins or alternatively in another embodiment the fusion protein can be synthesized as a recombinant polypeptide. The fusion polypeptide can be linked in any suitable way e.g. by a linker segment. and the fusion polypeptide should at least comprise the binding domains of the selected proteins. The linker peptide should be selected in such a way that it is not unstable resulting in degradation. The linker could e.g. be a highly O-glycosylated linker as linkers between catalytic domains and carbohydrate binding domains known from fungal carbohydrases, or it could be proline rich linkers.

[0088] The dual affinity polypeptide comprises at least one binding domain capable of binding to the target biomolecule with the desired binding specificity as described. The binding domain can be comprised in the complete protein or it can be a fragment of the protein which has retained its binding specificity. Many proteins have been described in the literature displaying affinity towards biomolecules, e.g. peptides, proteins, DNA, RNA, carbohydrates, and all such proteins or fragments thereof are potentially useful in the context of the present invention as candidates for the dual affinity polypeptide.

[0089] The said binding domain directed towards the target biomolecule can in one embodiment therefore be selected from but not limited to the group consisting of protein A, protein A fragments, protein A derived domains (e.g. domains known as an affibody®), antibodies, antibody fragments, lipocalins, and lectins.

[0090] Combinatorial protein engineering has been applied to develop artificial proteins that can bind to selected targets with high affinity and be used as alternatives to antibodies (Nygren, P.-Å. & Skerra, A. (2004). Binding proteins from alternative scaffolds. *J. Immunol. Methods*, 290, p. 3-28; Binz, H. K. & Plückthun, A. (2005). Engineered proteins as specific binding reagents. *Curr. Opin. Biotech.* 16, p. 459-469). In the context of the present invention the term "affibody" defines a class of engineered proteins selected for their specific binding activity towards a desired target and based on the Z domain, which is a 58 residue three-helical bundle derived by a single amino acid substitution in the B domain of staphylococcal protein A (SPA) (Nilsson, B., Moks, T., Jansson, B., Abrahmsén, L., Elmlad, A., Holmgren, E. et al. (1987) *Protein Eng.* 1, p. 107-113). The Z domain binds to the Fc region of immunoglobulins as do the five homologous SPA domains, but unlike the parental domain it does not bind to the Fab region. Such affibodies are examples of a protein A derived binding domain.

[0091] The dual affinity polypeptide also comprises at least one binding domain capable of binding to the catching ligand immobilized on the solid support. This second binding domain can be comprised in the complete protein or it can be a fragment of the protein which has retained its binding specificity. In one embodiment the second binding domain is selected from but not limited to the group consisting of avidin, streptavidin, neutravidin, steroid receptor, antibody, antibody fragment, amyloglucosidase (AMG), enzyme domain (e.g. cellulose binding domain, CBD), lipocalins, and lectins. As stated above these candidates, for the second binding domain, are meant as examples illustrating the invention, however, these examples should not be seen as the only usable combinations.

[0092] In one embodiment the antibody is selected from the group consisting of lama and camel antibodies.

[0093] In a particular embodiment the dual affinity polypeptide according to the invention comprises at least one binding domain of protein A fused to at least one biotin binding domain of avidin, streptavidin or neutravidin.

[0094] In a particular embodiment the dual affinity polypeptide according to the invention comprises at least one binding domain of a protein A derived binding domain fused to at least one biotin binding domain of avidin, streptavidin or neutravidin.

[0095] In another particular embodiment the dual affinity polypeptide comprises at least one binding domain of an affibody fused to at least one biotin binding domain of avidin, streptavidin or neutravidin.

[0096] In another particular embodiment the dual affinity polypeptide comprises at least one binding domain of an antibody fused to at least one biotin binding domain of avidin, streptavidin or neutravidin.

[0097] In another particular embodiment the dual affinity polypeptide comprises at least one binding domain of protein A fused to AMG, CBD or (VhhRR6(R2)).

[0098] In another particular embodiment the dual affinity polypeptide comprises at least one binding domain of a protein A derived binding domain fused to AMG, CBD or (VhhRR6(R2)).

[0099] In another particular embodiment the dual affinity polypeptide comprises at least one binding domain of an affibody fused to AMG, CBD or (VhhRR6(R2)).

[0100] In another particular embodiment the dual affinity polypeptide comprises at least one binding domain of an antibody fused to AMG, CBD or (VhhRR6(R2)).

[0101] The dual affinity polypeptide can as illustrated in the examples be linked chemically; however, a more cost efficient way to produce the dual affinity polypeptide would be to express it as a recombinant fusion protein.

[0102] In one embodiment of the invention, the fusion polypeptide is produced as a recombinant polypeptide.

[0103] Another possibility also envisioned would be to co-express the fusion protein and the target biomolecule in the host cell making it possible to load the crude cell culture extract directly on the solid support.

[0104] In a further embodiment the target biomolecule and the DAP is expressed separately but in the same type of host cell.

[0105] In a particular embodiment the fusion protein is expressed as a recombinant protein, particularly the fusion protein is in one embodiment recombinant Streptavidin linked to protein A. Such fusion protein can be produced intracellular in *E. coli* as described in Sano (T. Sano and C. R. Cantor (1991) BioTechnology 9 p 1378 - 1381), preferentially using the construct pTSAPA-2 carrying two IgG binding domains. However this construct is not industrially feasible as intracellular production with recovery of inclusion bodies in *E. coli* do not give industrially relevant yields and the production process is highly complex. A process based on a secreted fusion produced in e.g. *Bacillus* or *Aspergillus* is of much higher industrial relevance.

[0106] The nucleotide sequence encoding the fusion protein according to the invention may preferably be expressed by inserting the nucleotide sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0107] The recombinant expression vector may be any vector (e.g., a plasmid or virus) which can be conveniently subjected to recombinant DNA procedures and can bring about expression of the nucleotide sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids.

[0108] The vector may be an autonomously replicating vector, i.e., a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the

chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the host cell, or a transposon may be used.

[0109] The vectors of the present invention preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

[0110] A conditionally essential gene may function as a non-antibiotic selectable marker. Example of bacterial conditionally essential non-antibiotic selectable markers are the *dal* genes from *Bacillus subtilis*, *Bacillus licheniformis*, or other *Bacilli*, that are only essential when the bacterium is cultivated in the absence of D-alanine. Also the genes encoding enzymes involved in the turnover of UDP-galactose can function as conditionally essential markers in a cell when the cell is grown in the presence of galactose or grown in a medium which gives rise to the presence of galactose. Non-limiting examples of such genes are those from *B. subtilis* or *B. licheniformis* encoding UTP-dependent phosphorylase (EC 2.7.7.10), UDP-glucose-dependent uridylyltransferase (EC 2.7.7.12), or UDP-galactose epimerase (EC 5.1.3.2). Also a xylose isomerase gene such as *xylA*, of *Bacilli* can be used as selectable markers in cells grown in minimal medium with xylose as sole carbon source. The genes necessary for utilizing gluconate, *gntK*, and *gntP* can also be used as selectable markers in cells grown in minimal medium with gluconate as sole carbon source. Other examples of conditionally essential genes are known in the art. Antibiotic selectable markers confer antibiotic resistance to such antibiotics as ampicillin, kanamycin, chloramphenicol, erythromycin, tetracycline, neomycin, hygromycin or methotrexate.

[0111] Suitable markers for yeast host cells are ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*.

[0112] The vectors of the present invention preferably contain an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

[0113] For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleotide sequences for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, preferably 400 to 10,000 base pairs, and most preferably 800 to 10,000 base pairs, which have a high degree of identity with the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleotide sequences. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

[0114] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication which functions in a cell. The term "origin of replication" or "plasmid replicator" is defined herein as a nucleotide sequence that enables a plasmid or vector to replicate *in vivo*.

[0115] Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM β 1 permitting replication in *Bacillus*.

[0116] Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

[0117] Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems et al., 1991, Gene 98:61-67; Cullen et al., 1987, Nucleic Acids Research 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

[0118] More than one copy of a polynucleotide of the present invention may be inserted into the host cell to increase production of the gene product. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional

copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

[0119] The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, e.g., Sambrook *et al.*, 1989, *supra*).

[0120] The present invention also relates to recombinant host cells, comprising a polynucleotide of the present invention, which are advantageously used in the recombinant production of the polypeptides. A vector comprising a polynucleotide of the present invention is introduced into a host cell so that the vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier. The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

[0121] The host cell may be a unicellular microorganism, e.g., a prokaryote, or a non-unicellular microorganism, e.g., a eukaryote.

[0122] Useful unicellular microorganisms are bacterial cells such as gram positive bacteria including, but not limited to, a *Bacillus* cell, e.g., *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*; or a *Streptomyces* cell, e.g., *Streptomyces lividans* and *Streptomyces murinus*, or gram negative bacteria such as *E. coli* and *Pseudomonas* sp. In a preferred aspect, the bacterial host cell is a *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus stearothermophilus*, or *Bacillus subtilis* cell. In another preferred aspect, the *Bacillus* cell is an alkalophilic *Bacillus*.

[0123] The introduction of a vector into a bacterial host cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, Molecular General Genetics 168: 111-115), using competent cells (see, e.g., Young and Spizzin, 1961, Journal of Bacteriology 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, Journal of Molecular Biology 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, Biotechniques 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, Journal of Bacteriology 169: 5771-5278).

[0124] The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

[0125] In a preferred aspect, the host cell is a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota (as defined by Hawksworth *et al.*, In, Ainsworth and Bisby's Dictionary of The Fungi, 8th edition, 1995, CAB International, University Press, Cambridge, UK) as well as the Oomycota (as cited in Hawksworth *et al.*, 1995, *supra*, page 171) and all mitosporic fungi (Hawksworth *et al.*, 1995, *supra*).

[0126] In a more preferred aspect, the fungal host cell is a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in Biology and Activities of Yeast (Skinner, F.A., Passmore, S.M., and Davenport, R.R., eds, Soc. App. Bacteriol. Symposium Series No. 9, 1980).

[0127] In an even more preferred aspect, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

[0128] In a most preferred aspect, the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis* or *Saccharomyces oviformis* cell. In another most preferred aspect, the yeast host cell is a *Kluyveromyces lactis* cell. In another most preferred aspect, the yeast host cell is a *Yarrowia lipolytica* cell.

[0129] In another more preferred aspect, the fungal host cell is a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

[0130] In an even more preferred aspect, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filobasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolyopocladium*, *Trametes*, or *Trichoderma* cell.

[0131] In a most preferred aspect, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus fumigatus*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger* or *Aspergillus oryzae* cell. In another most preferred aspect, the filamentous fungal host cell is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cell. In another most preferred aspect, the filamentous fungal host cell is a *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*, *Ceriporiopsis subrufa*, or *Ceriporiopsis subvermispora*, *Coprinus cinereus*, *Coriolus hirsutus*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

[0132] Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238 023 and Yelton et al., 1984, Proceedings of the National Academy of Sciences USA 81: 1470-1474. Suitable methods for transforming *Fusarium* species are described by Malardier et al., 1989, Gene 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al., 1983, Journal of Bacteriology 153: 163; and Hinnen et al., 1978, Proceedings of the National Academy of Sciences USA 75: 1920.

[0133] The contact between the target biomolecule, the dual affinity polypeptide (DAP), and the solid support can be performed in several optional ways. In one embodiment all components could be brought into contact in one step, eg. by loading the target polypeptide and the fusion protein on the solid support without pre-incubation in solution. The dual affinity polypeptide can however, be contacted with the target before loading this complex on the solid support. In this embodiment the target biomolecule and the dual affinity polypeptide are contacted first, e.g. in solution, and subsequently the formed complex is contacted with the solid support. Depending on the nature of the solid support preferred embodiments of this principle could differ.

[0134] In one preferred embodiment the solid support is a solid phase matrix. This includes conventional solid phase matrixes. In the case of solid phase matrixes in the form of columns, the target and the dual affinity polypeptide are contacted first in solution and subsequently contacted with the solid phase matrix by loading the complex onto the column.

[0135] In another embodiment the solid support is in the form of particles.

[0136] The catching ligand according to the invention is covalently attached to the solid support. As explained above the ligand according to the present invention is different from the ligand used in traditional affinity chromatography where the purpose of the ligand is to bind the target. In the present invention the ligand should bind to the DAP. Ligands are well known in the art and below are given examples that can be applied according to the invention. In the context of the present invention in one particular embodiment instead of a ligand attached to the solid phase the solid phase could alternatively comprise a binding affinity or binding site towards the DAP. An example could be cellulose as the solid phase and CBD (cellulose binding domain) as part of the DAP.

[0137] In one embodiment the ligand is chosen from but not limited to the group consisting of biotin, acarbose, steroids, haptens, epitope-peptides, dyes and enzyme inhibitors. In a particular embodiment the ligand is biotin. The ligand can be chemically attached to the solid support as described in the examples where the chemical attachment of acarbose and reactive red 6 is illustrated.

[0138] The coupling of affinity ligands to supports strongly influences the specificity, capacity and cost of traditional affinity chromatography columns.

[0139] The current state of the art in covalent coupling technology allows for chemo and regio selective coupling of the binding

ligands to the support, often using spacers or linkers to anchor the ligand to the surface.

[0140] Great care is taken to avoid using functional groups that are close to a binding site or that play a role in the interaction between the ligand and target molecule.

[0141] If a suitable functional group does not exist on the ligand, further derivatizing of the ligand can be done to add an appropriate functional group. Numerous references describe appropriate chemistries, including "Bioconjugate Techniques", by Greg T. Hermanson, Academic Press, 2008 and "Chemistry of Protein Conjugation and Cross-linking", by Shan S. Wong, CRC Press, 1991.

[0142] It is commonly accepted that a high concentration of coupled ligand often has adverse effects on affinity chromatography, also the binding efficiency of the medium may be reduced due to steric hindrance between the active sites. This is particularly pronounced when large molecules such as antibodies, antigens and enzymes interact with small ligands.

[0143] In addition, the target substances may become more strongly bound to closely packed ligands making elution difficult and also the extent of nonspecific binding increases at very high ligand concentrations, thus reducing the selectivity of the affinity column.

[0144] Ligand-surface interface interaction is known to be important for the affinity ligand performance. The length of spacer arms between the ligand and the surface is critical. If it is too short, the arm is ineffective and the ligand fails to bind the target in the sample. If it is too long, proteins may bind non-specifically to the spacer arm and reduce the selectivity of the separation. Often 4-12 atom long hydrophilic arms are used.

[0145] MabSelect™ Media and HiTrap MabSelect™ (GE Healthcare) are examples of affinity columns using oriented coupling of recombinant Protein A to the matrix via an engineered C-terminal cysteine and a hydrophilic spacer arm.

[0146] The present invention suggests the use of soluble dual affinity polypeptide which can be characterized and used in any concentration appropriate for the specific target concentration. The technical challenge of coupling delicate target specific binding ligands to a solid support is substituted with more simple preparation of soluble molecules making it possible to utilize the entire arsenal of analytical methods.

[0147] There are numerous types of support material for affinity chromatography.

[0148] The size and uniformity of beads, the distribution of internal channels and the nature of the surfaces has all been optimized to produce numerous types of supports.

[0149] In general, smaller particle size and greater porosity, ensures increased dynamic binding capacity. On the other hand, resistance to mechanical collapses is reduced.

[0150] Both compressible and the incompressible support material needs to be robust enough to survive multiple cycles without change of flow rates which will influence the residence time.

[0151] The solid support are in the form of beads, gels or granulates. The quality of packing of the solid support material in columns for traditional affinity purification and the flow rates during operation greatly influence the performance.

[0152] Specialized equipment is used to successfully pack large columns above 5-10 cm in diameter. High performance columns are normally purchased pre-packed and in standard sizes. Consequently, the practical dimensions during scale up depend on available column systems for the purification of the particular target molecule.

[0153] The present invention suggests a general method using target DAP molecules and a generic column.

[0154] One of the most important factors in determining the cost and quality of the large scale purification is the chemical and mechanical stability of the adsorbent.

[0155] Traditional affinity columns with immobilized protein ligands are susceptible to further degradation due to for example oxidation or microbial growth.

[0156] Therefore, due to the cost of large affinity columns, great care has to be taken to control the storage condition between uses. Often the column is washed and stored with a special buffer solution containing anti microbial agents, alcohols or similar. These storage solutions must be washed away before use.

[0157] Some affinity ligands are also sensitive to proteases and the column lifetime will be reduced unless special cleaning and regeneration procedures are followed rigorously. The freedom to design efficient affinity purification procedures is therefore somewhat restricted.

[0158] A single use column system according to the invention or a column system using synthetic ligands will not have the above technical limitations.

In one embodiment of the invention the solid support is in the form of a solid phase matrix. The solid phase matrix may comprise, as the matrix backbone, any natural or synthetic and organic or inorganic material known per se to be applicable in solid phase separation of proteins and other biomolecules, e.g. natural or synthetic polysaccharides such as agar-agar and agaroses; celluloses, cellulose ethers such as hydroxypropyl cellulose, carboxymethyl cellulose; starches; gums such as guar gum, and gum arabic, gum ghatti, gum tragacanth, locust bean gum, xanthan gum; pectins; mucins; dextrans; chitins; chitosans; alginates; carrageenans; heparins; gelatins; synthetic polymers such as polyamides such as polyacrylamides and polymethacrylamides; polyimides; polyesters; polyethers; polymeric vinyl compounds such as polyvinylalcohols and polystyrenes; polyalkenes; inorganic materials such as silicious materials such as silicon dioxide including amorphous silica and quartz; silicas; metal silicates, controlled pore glasses and ceramics; metal oxides and sulfides, or combinations of these natural or synthetic and organic or inorganic materials.

[0159] The matrix backbone is preferably selected from agar-agar, agaroses, celluloses, cellulose ethers such as hydroxypropyl cellulose, carboxymethyl cellulose, polyamides such as poly(meth)acrylamides, polyvinylalcohols, silicas, and controlled pore glasses.

[0160] Especially interesting solid phase materials as matrix backbones are e.g. agar or agarose beads such as Sepharose and Superose beads from GE Healthcare, USA, and Biogel A from Biorad, USA; dextran based beads such as Sephadex, GE Healthcare; cellulose based beads and membranes such as Perloza cellulose from Iontosorb, Czech Republic; composite beads such as Sephacryl and Superdex, GE Healthcare, USA; beads of synthetic organic polymers such as Fractogel from Tosoh Lifesciences LLC, USA; POROS media from Applied Biosystems, USA, Bio-Rex, Bio-Gel P and Macro Prep from Biorad, HEMA and Separon from TESSEK and Hyper D and Trisacryl media from Pall Corporation, USA, Enzacryl and Azlactone, 3M, USA; beads of siliceous materials such as controlled pore glass, PROSEP, from Millipore, USA, and Spherozil, Pall Corporation, USA; and coated silica composites in the form of beads or membranes such as ACTI-DISK, ACTI-MOD and CycloSep from Arbor Technologies, USA.

[0161] The ligand (e.g. biotin or similar specific molecules of low molecular weight (LMW)) is then covalently attached to this material. Several coupling chemistries of ligand molecules to the solid support can be selected from text books on the subject (Protein Purification, 1998, 2ed, eds. Janson, J.-C., Rydén, L, Wiley & sons inc. New York). Based on the particular purification task the best candidate of ligand derivatives is coupled to the best choice of solid support, e.g. solid phase matrix or particles. Production process properties of the affinity solid matrix are analyzed through practical laboratory and pilot testing.

[0162] The ligands may be attached to the solid phase material by any type of covalent bond known per se to be applicable for this purpose, either by a direct chemical reaction between the ligand and the solid phase material or by a preceding activation of the solid phase material or of the ligand with a suitable reagent known per se making it possible to link the matrix backbone and the ligand. Examples of such suitable activating reagents are epichlorohydrin, epibromohydrin, allyl glycidylether; bis-epoxides such as butanedioldiglycidylether; halogen-substituted aliphatic compounds such as di-chloro-propanol, divinyl sulfone; carbonyldiimidazole; aldehydes such as glutaric dialdehyde; quinones; cyanogen bromide; periodates such as sodium-meta-periodate; carbodiimides; chloro-triazines such as cyanuric chloride; sulfonyl chlorides such as tosyl chlorides and tresyl chlorides; N-hydroxy succinimides; 2-fluoro-1-methylpyridinium toluene-4-sulfonates; oxazolones; maleimides; pyridyl disulfides; and hydrazides. Among these, the activating reagents leaving a spacer group SP1 different from a single bond, e.g. epichlorohydrin, epibromohydrin, allylglycidylether; bis-epoxides; halogen-substituted aliphatic compounds; divinyl sulfone; aldehydes; quinones; cyanogen bromide; chloro-triazines; oxazolones; maleimides; pyridyl disulfides; and hydrazides, are preferred.

[0163] In one embodiment the solid support is in the form of particles. Particles can be selected from the group comprising microspheres, latex particles or beads. The particles can be made from but not limited to the group consisting of e.g. polystyrene, silica, naphthalene, polybutylmethacrylate.

[0164] The generic solid support can be produced at costs comparable to ion exchange matrices and the recombinant dual affinity protein can also be produced as a recombinant fusion protein by fermentation at low cost. Due to the lowered cost of the novel downstream procedure materials, the purification technology may be provided as disposables, which eliminate the need for expensive cleaning in place (CIP) and certain validations. Another consequence of the reduced cost is optional large column-volume applications, which saves manufacturing labour expenses, prevent repeated purification re-runs and limit time occupations of the downstream process plant.

[0165] The use of a generic solid support including the capturing ligand and the potentially improved binding efficiency and capacity due to complex formation in solution poses several advantages over the conventional affinity chromatography. These advantages are listed below.

- No time consuming and expensive chemical conjugation reactions, purifications and QC procedures of protein ligands to prepare the affinity column material prior to affinity purification of the target molecule.
- The generic matrix is more cost efficient to manufacture compared to the present commercial affinity matrices (e.g. Protein A). Only one type of capturing column material is required for all affinity purifications using the dual affinity polypeptide principle. The low molecular weight ligand, e.g. biotin, dye molecules or similar specific low molecular weight (LMW) molecules are covalently attached by simple low cost conjugation procedures to make the generic solid phase matrix.
- The preferred fermentation of a dual affinity polypeptide fusion protein is "simple" , based on known technology and provide the conjugation needed between binding domains in DAP
- The manufacturing cost of DAP molecules is comparable or cheaper than recombinant Protein A molecules.
- The DAP fusion protein and the generic matrix required for purification purposes costs a fraction of the ready-to-use Protein A affinity matrix for similar purposes
- The DAP transport and immobilize the target molecules to the generic matrix during the purification process. There is no need for an expensive and time consuming immobilization of a dedicated ligand to make a specific purification matrix as known in conventional affinity chromatography.
- The low cost of the components in the presented invention facilitate a disposable affinity purification process featuring
 - o elimination of CIP procedures
 - o elimination of validation procedures
 - o save time on regulatory issues
 - o exclude repetitions of down stream process cycles
 - o limit operational failures
 - o lower labor expenses during processing
 - o shorter manufacturing run time
 - o limit risk of contamination
 - o easy to use
 - o lower capacity cost investments due to flexible plant designs
 - o better down stream process economy
- Substitution of conventional multi-cycle protein separation procedures to a single step using the disposable affinity purification technology.

[0166] In a particular modified form of the invention it could be envisioned that the DAP molecule could bind covalently to the solid support. This would still allow the possibility of having the DAP and the target reacting in solution. Such a covalent bond could in one embodiment be formed by cleavage of a para-substituted benzyl guanine resulting in a thioether bond.

[0167] One embodiment of this modified form of the invention therefore relates to a process for purification of a target biomolecule, comprising the steps: (a) contacting (i) a target polypeptide, (ii) a dual affinity polypeptide, and (iii) a solid support comprising a catching ligand, wherein the dual affinity polypeptide has an equilibrium dissociation constant, K_{D1} towards the

target biomolecule in the range from 10^{-2} to 10^{-13} M, more particularly from 10^{-4} to 10^{-13} M at standard conditions, and wherein binding of the dual affinity polypeptide to the catching ligand on the solid support is provided by cleavage of a para-substituted benzyl guanine resulting in a thioether bond; and (b) recovering the target biomolecule by elution.

[0168] The basic principle of the affinity purification technology, for purifying a target molecule (polyclonal antibody) is illustrated below.

EXAMPLES

Example 1. Preparation of dual affinity linker by chemical conjugation

[0169] Based on published values for binding affinities, dual linker binding functionalities were selected that fit both to the binding to the ligand matrix ($K_{D,S} \sim 10^{-9}$ to 10^{-16} M) and to the target biomolecules (products, $K_{D,T} \sim 10^{-2}$ to 10^{-13} M). To investigate the influence of the $K_{D,S}$, some components with $K_{D,S}$ -values outside the above interval were also tested.

[0170] In order to prepare a conjugate made from Protein A and a biotin binding protein e.g. Avidin, Streptavidin or Neutravidin the two proteins were chemically activated separately as a first step and joined together by cross linking in a second step afterwards.

Protein A do not have accessible sulphhydryl (-SH) on the surface, so these were introduced by reaction with SATA (N-succinimidyl S-acetylthioacetate) to primary amine (-NH₂) functional groups on Protein A. SATA (or SATP) is a reagent for introducing protected sulphhydryls into proteins, peptides and other molecules. They are the N-hydroxysuccinimide (NHS) esters of S-acetylthioacetic and propionic acid. A stable, covalent amide bond was formed from the reaction of the NHS ester with primary amines. The amine was reacted with the NHS ester by nucleophilic attack, with N-hydroxysuccinimide being released as a by-product. Deprotection (deacylation) to generate a sulphhydryl for use in cross-linking and other applications was accomplished using hydroxylamine•HCl.

[0171] The maleimide groups were introduced to the Avidin using Sulfo-SMCC. Sulfo-SMCC is a heterobifunctional cross-linker that contains a N-hydroxysuccinimide (NHS) ester and a maleimide group. NHS esters react with primary amines at pH 7-9 to form covalent amide bonds. SMCC and Sulfo-SMCC are often used to prepare protein-protein conjugates in a two-step reaction scheme. First, the amine-containing protein was reacted with a several-fold molar excess of the cross linker, followed by removal of excess (nonreacted) reagent by desalting or dialysis; finally, the sulphhydryl-containing molecule is added to react with the maleimide groups already attached to the first protein.

[0172] The conjugates prepared by cross linking were obtained by reacting maleimides with sulphhydryl groups at pH 6.5-7.5 to form stable thioether bonds.

[0173] An alternative to the procedure above is to use commercially available Maleimide activated Neutravidin instead of the activated Avidin. Maleimide Activated NeutrAvidin™ Protein is for directly preparing NeutrAvidin™ Protein (NAP) conjugates with proteins, peptides, and other molecules that contain a free sulphhydryl (-SH) group.

Preparation of Dual Affinity Polypeptide (DAP) by chemical cross-linking

[0174] Procedure for chemically cross-linking Protein A, and either Avidin or Neutravidin into a conjugate with the required properties of a DAP linker.

Materials

[0175] SATA (N-Succinimidyl S-Acetylthioacetate), (Pierce, no.26102)

D-Salt™ Excellulose™ Desalting Column, 5 ml (Pierce No. 20449)

Hydroxylamine•HCl (Pierce, No. 26103), DMSO (Dimethylsulfoxide, Pierce, No. 20688), Sulfo-SMCC: (sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexan-1-carboxylate) (Pierce, 22322), Protein A (GE Health Care, 17-0872-50), Avidin (Kem-En-Tec, 4020H), Maleimide activated neutravidin (Pierce, no. 31007), PD-10 Sephadex G-25M (GE; 17-0851-01),

HiPrep 26/60 Sephacryl S-100 HR (MW range 1.000-100.000) (GE, 17-1194-01)
 Anti-IgG Affibody (Affibody, 10.0623.01.0050)
 Dithiothreitol ([3483-12-3], Sigma-Aldrich D0632).

[0176] Buffers:

PBS Reaction Buffer: 200-500 ml of PBS: 0.1 M phosphate, 0.15 M NaCl, pH 7.2

Deacetylation Solution: 0.5 M Hydroxylamine, 25 mM EDTA in PBS, pH 7.2

PBS-EDTA solution: 200-500 ml of PBS: 0.1 M phosphate, 0.15 M NaCl, 5 mM EDTA, pH 7.2.

Phosphate-buffered saline (PBS, pH 7.2; or other amine- and sulfhydrylfree buffer at pH 6.5-7.5 adding EDTA to 1-5 mM helps to chelate divalent metals, thereby reducing disulfide formation in the sulfhydryl-containing protein.

Procedure for Sulfhydryl Modification of Protein A

A. Reaction of Protein A with SATA (or SATP)

[0177] Immediately before reaction, 6.4 mg of SATA was dissolved in 0.5 ml of DMSO (resulting in ~55 mM solution).

[0178] 1.0 ml of Protein A solution (2.6 mg/mL) was then combined with 10 µl of the SATA solution. The contents were mixed and incubated at room temperature for 30 minutes.

[0179] The level of sulfhydryl incorporation may be altered by using different molar ratios of SATA to protein. The default reaction uses 60 nmol Protein A and 550 nmol SATA, a 9:1 molar ratio of SATA to protein. The amount of SATA in the reaction may be increased or decreased by adding more or less than 10 µl of the SATA solution per ml of Protein Solution.

B. Desalt to purify acylated Protein A from excess reagent and by-products

[0180] A desalting column was equilibrated with two column volumes of Reaction Buffer. Use at least a 5 ml desalting column for each 1 ml of reaction volume to be processed.

[0181] 1.01 ml reaction mixture was applied to the column. Collection of 1 ml fractions was started immediately. After the reaction mixture had completely entered the column bed and the first fraction was collected, at least 10 mL Reaction Buffer was added to the column and collection continued as separate 1 ml fractions as they emerged from the column.

[0182] Fraction(s) that contain Protein A were identified as those having peak absorbance at 280 nm. With a 5 ml desalting column, fractions 2 and 3 contained most of the protein, while excess SATA came out in the following fractions. The fractions that contain the modified Protein A were pooled.

[0183] The modified Protein A may be stored indefinitely for later deacetylation and generation of sulfhydryl groups (Section C).

C. Deacetylate SATA-modified Protein A to generate sulfhydryl groups

[0184] 1.0 ml of SATA-modified (acylated) Protein A was combined with 100 µl of the Deacetylation Solution. The contents were mixed and incubated 2 hours at room temperature.

[0185] A desalting column was used to purify the sulfhydryl-modified protein from the Hydroxylamine in the Deacetylation Solution.

[0186] Desalting was done into Reaction Buffer containing 10 mM EDTA to minimize disulfide bond formation using the same procedure as in Section B. Fractions that contained the modified Protein A were pooled. The protein concentration should be ~1.3

mg/ml. In order to minimize disulfide formation Section D was performed immediately.

[0187] Before or after desalting, the protein may be assayed for sulfhydryl content using Ellman's Reagent (Pierce, no 23460 (kit for sulfhydryl group detection)).

D. Conjugation of SATA modified Protein A to maleimide activated Avidin or Neutravidin.

[0188] This method uses approximately equimolar amounts of activated Protein A to Avidin or Neutravidin.

Example 1a. Procedure for Maleimide Modification of Avidin and preparation of the DAP molecule [Protein A - Avidin]

[0189] Generally, a 10- to 50-fold molar excess of cross-linker over the amount of amine-containing protein results in sufficient maleimide activation to enable several sulfhydryl-containing proteins to be conjugated to each amine-containing protein.

[0190] More dilute protein solutions require greater fold molar excess of reagent to achieve the same activation level. Empirical testing is necessary to determine optimal activation levels and final conjugation ratios for the intended application.

Protocol

[0191] For best results, ensure that Protein A-SH is prepared as described above and ready to combine with maleimide activated Avidin in step 5.

[0192] 32 mg Avidin was prepared in 5 mL PBS Buffer, and 4.36 mg sulfo-SMCC was prepared in 1 mL PBS/EDTA buffer. Then 500 μ L of the activation solution was transferred to the Avidin solution. The mixture was incubated 30 minutes at room temperature. Excess cross-linker was removed using a desalting column equilibrated with PBS-EDTA Buffer.

[0193] Protein A-SH and desalted maleimide activated Avidin were combined and mixed in a molar ratio corresponding to approximately 1:1. The reaction mixture was incubated at room temperature overnight.

[0194] Generally, there is no harm in allowing the reaction to proceed for several hours or overnight, although usually the reaction will be complete in about 30 min. To stop the conjugation reaction before completion, add buffer containing reduced cysteine at a concentration several times greater than the sulfhydryls of Protein A-SH.

Example 1 b. Preparation of the DAP molecule [Protein A - Neutravidin]

[0195] For best results, ensure that Protein A-SH is prepared as described above and ready to combine with maleimide activated Avidin.

[0196] Maleimide activated Neutravidin (Pierce, no 31007) is commercially available for directly preparing NeutrAvidin™ Protein (NAP) conjugates with proteins, peptides, and other molecules that contain a free sulfhydryl (-SH) group. NeutrAvidin™ Protein is a modified avidin derivative with several key features that provide a biotin-binding protein with exceptionally low nonspecific binding properties. NeutrAvidin™ Protein does not contain carbohydrates, rendering lectin-binding activity to undetectable levels. Additionally, the isoelectric point of NAP is 6.3 ± 0.3 , which is much lower than native Avidin and not as acidic as streptavidin.

Protocol

[0197] 1.0 mL of ultra pure water was added to suspend 5 mg lyophilized Neutravidin.

Protein A-SH and maleimide activated Neutravidin were combined and mixed in a molar ratio corresponding to 1:1. The reaction mixture was incubated at room temperature overnight.

[0198] Generally, there is no harm in allowing the reaction to proceed for several hours or overnight, although usually the

reaction will be complete in the specified time. To stop the conjugation reaction before completion, add buffer containing reduced cysteine at a concentration several times greater than the sulfhydryls of Protein A-SH.

Example 1c. Preparation of the DAP molecule [Affibody (IgG) - Avidin]

Protocol

[0199] Avidin (10 mg) was activated with sulfo-SMCC as described in Example 1 a.

Anti-IgG Affibody disulfide dimers were reduced to monomers:

Anti-IgG Affibody (5 mg) is dissolved in PBS-buffer (5 mL), and 3.8 mL of this solution is transferred to a vial containing 12.3 mg dithiothreitol (DTT) giving a final DTT concentration of 20 mM solution. This mixture is turned at RT for 2 h.

Upon this, excess DTT is removed by splitting the reaction mixture in two portions, passing each portion through a PD-10 column (bedvolume 8 mL). The columns had been equilibrated with 25 mL PBS buffer before use. The monomeric Anti-IgG Affibody is eluted from the columns in 2 x 9-10 fractions, each containing 1 mL.

By measuring A₂₈₀ of the fractions the protein was located in 2 fractions from each column. These fractions were pooled and mixed with the desalted maleimid activated avidin solution (20 mL) in a molar ratio corresponding to approximately 1:1 (avidin calculated as monomer; MW=17.000), and the coupling was allowed to proceed overnight at room temperature with gently turning of the coupling mixture.

The following day, 1500 µL of the conjugation mixture was concentrated in an Amicon Ultra (cut off 3 kDa) to a total volume of 400 µL, which was used for analysis by SDS PAGE. This showed that all avidin had reacted, and that there was still some unreacted anti-IgG Affibody present.

The conjugation mixture was freeze-dried until purified by SEC.

[0200] The above protocol can be used for the preparation of all derivatives of Affibody-Avidin dual affinity polypeptides.

Example 1 d. Preparation of the DAP molecule [Affibody (Insulin) - Avidin]

[0201] Avidin (9 mg) was activated with sulfo-SMCC as described in Example 1a.

Anti-Insulin Affibody (His₆-Z000810-Cys; *PBS00014*) disulfide dimers were reduced to monomers as described in Example 1 c.

The pooled fractions from the PD-10 columns were mixed with the desalted maleimid activated avidin solution (16.1 mL) in a molar ratio corresponding to approximately 1:1, and the coupling was allowed to proceed overnight at room temperature with gently turning of the coupling mixture.

The following day, the conjugation mixture was analyzed by SDS PAGE. This showed that all avidin had reacted, and that there was still some unreacted anti-Insulin Affibody present.

[0202] The conjugation mixture was freeze-dried until purified by SEC.

Example 1e. Preparation of the DAP molecule [Affibody (Insulin) - Avidin]

[0203] Avidin (9 mg) was activated with sulfo-SMCC as described in Example 1a.

Anti-Insulin Affibody (Insulin, His₆-Z01139-Cys; *PB00022*) disulfide dimers were reduced to monomers as described in Example 3c.

The pooled fractions from the PD-10 columns were mixed with the desalted maleimid activated avidin solution (16.1 mL) in a molar ratio corresponding to approximately 1:1, and the coupling was allowed to proceed overnight at room temperature with gently turning of the coupling mixture.

The following day, the conjugation mixture was analyzed by SDS PAGE. This showed that all avidin had reacted, and that there was still some unreacted anti-Insulin Affibody present. The conjugation mixture was freeze-dried until purified by SEC.

Example 2. Recombinant dual affinity constructs for expression in *Aspergillus*

oryzae.**Strains**

[0204] *Aspergillus oryzae* BECh2 is described in WO 00/39322, example 1, which is further referring to JaL228 described in WO 98/12300, example 1.

JaL1168 is described in example 2g.

JaL1171 is described in example 2g.

JaL1174 is described in example 2g.

JaL1176 is described in example 2g.

JaL1181 is described in example 2g.

JaL1210 is described in example 2g.

MT173 is a derivative of MC1000 (Casadaban & Cohen J. Mol. Biol. 138 (1980) 179-207) which are *ara*⁺ and *leuB6*.

Genes

[0205] AMG: indicate the *Aspergillus niger* glucoamylase gene (Boel et al. EMBO Journal 3 (1984) 1581-1585)

Z: indicated the Z domain from *Staphylococcus aureus* preotein A (Nilsson et al. Protein Engineering 1 (1987) 107-113).

Pre-CBD(C315): indicate the *Meripilus giganteus* endoglucanase II (DSM971) signal (pre) + cellulose binding domain (CBD) + linker region.

CBD(egv): indicated the *Humicola insolens* endoglucanase V (DSM1800) linker region + cellulose binding domain.

VhhRR6(2): indicated the variable region from a Llama single chain antibody reacting against the hapten azo-dye Reactive Red (RR6) (Frenken et al. J. Biotechnology 78 (2000) 11-21).

Plasmids

[0206] p775 is described in EP 238023.

pA2C315 is deposited at DSM under the accession no. DSM971. The plasmid contains a cDNA clone from *Meripilus giganteus* encoding an endoglucanase II gene.

pCAMG91 is described in Boel et al. EMBO Journal 3 (1984), 1581-1585.

pJaL790 is described in WO2005070962, example 1.

pJaL1153 is described in example 2c.

pJaL1154 is described in example 2a.

pJaL1158 is described in example 2d.

pJaL1159 is described in example 2a.

pJaL1164 is described in example 2c.

pJaL1165 is described in example 2d.

pJaL1168 is described in example 2e.

pJaL1169 is described in example 2b.

pJaL1170 is described in example 2f.

pJaL1171 is described in example 2b.

pMT2786 is described in WO2006050737 example 2.

pSX320 is described in EP 0 531 372.

Primer and DNA sequences**[0207]**

Synthetic DNA 1 (SEQ ID NO 1)

Synthetic DNA 2 (SEQ ID NO 4)

Adaptor sequence 1 (SEQ ID NO 5)

Adaptor sequence 2 (SEQ ID NO 6)

primer 8683 (SEQ ID NO 10)

primer CBD:Z-NA (SEQ ID NO 11)

primer Z-NA (SEQ ID NO 12)

primer Z-CA (SEQ ID NO 13)

primer Z-CA:CBD (SEQ ID NO 14) primer 8654 (SEQ ID NO 15)

Primer CBD:Z-NB (SEQ ID NO 19)

Primer Z-NB (SEQ ID NO 20)

Primer Z-CB (SEQ ID NO 21)

Primer Z-NB:CBD (SEQ ID NO 22)

Methods

[0208] General methods of PCR, cloning, ligation nucleotides etc. are well-known to a person skilled in the art and may for example be found in "Molecular cloning: A laboratory manual", Sambrook et al. (1989), Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.); "Current protocols in Molecular Biology", John Wiley and Sons, (1995); Harwood, C. R., and Cutting, S. M. (eds.); "DNA Cloning: A Practical Approach, Volumes I and II", D.N. Glover ed. (1985); "Oligonucleotide Synthesis", M.J. Gait ed. (1984); "Nucleic Acid Hybridization", B.D. Hames & S.J. Higgins eds (1985); "A Practical Guide To Molecular Cloning", B. Perbal, (1984).

PCR amplification

[0209] All PCR amplifications were performed in a volume of 100 microL containing 2.5 units Taq po-lymerase, 100 ng of pSO₂, 250 nM of each dNTP, and 10 pmol of each of the two primers described above in a reaction buffer of 50 mM KCl, 10 mM Tris-HCl pH 8.0, 1.5 mM MgCl₂. Amplification was carried out in a Perkin-Elmer Cetus DNA Thermal 480, and consisted of one cycle of 3 minutes at 94°C, followed by 25 cycles of 1 minute at 94°C, 30 seconds at 55°C, and 1 minute at 72°C.

Aspergillus transformation

[0210] *Aspergillus* transformation was done as described by Christensen et al.; Biotechnology 1988 6 1419-1422. In short, *A. oryzae* mycelia were grown in a rich nutrient broth. The mycelia were separated from the broth by filtration. The enzyme preparation Novozyme[®] (Novo Nordisk) was added to the mycelia in osmotically stabilizing buffer such as 1.2 M MgSO₄ buffered to pH 5.0 with sodium phosphate. The suspension was incubated for 60 minutes at 37degrees C with agitation. The protoplast was filtered through mira-cloth to remove mycelial debris. The protoplast was harvested and washed twice with STC (1.2 M sorbitol, 10 mM CaCl₂, 10 mM Tris-HCl pH 7.5). The protoplasts were finally re-suspended in 200-1000 microl STC.

[0211] For transformation, 5 microg DNA was added to 100 microl protoplast suspension and then 200 microl PEG solution (60% PEG 4000, 10 mM CaCl₂, 10 mM Tris-HCl pH 7.5) was added and the mixture is incubated for 20 minutes at room temperature. The protoplast were harvested and washed twice with 1.2 M sorbitol. The protoplast was finally re-suspended 200 microl 1.2 M sorbitol. Transformants containing an intact niaD gene were selected for its ability to used nitrate as the sole source for nitrogen on minimal plates (Cove D.J. 1966. Biochem. Biophys. Acta. 113:51-56) containing 1.0 M sucrose as carbon source, 10 mM Sodium nitrate as nitrogen source. After 4-5 days of growth at 37 degrees C, stable transformants appeared as vigorously growing and sporulating colonies. Transformants were purified twice through conidiospores.

Media and reagents

[0212] YPM medium (2 g/l yeast extract, 2 g/l peptone, and 2% maltose)

Growth of *Aspergillus* transformants

[0213] Shake flask containing 10 ml YPM medium was inoculated with spores from the respective transformants and incubated at 30 degrees C, at 200 rpm for 4 days.

SDS-page

[0214] SDS gel used was Criterion™ XT precast gels, 10% Bis-Tris, from BIO-RAD and was run and stained with Coomassie blue as recommend by the manufactory.

Example 2a**Construction of *Aspergillus* expression cassette pJaL1159 (pre-CBD_(C315)-KR::VhhRR6(R2)::Z::Z)**

[0215] Plasmid pJaL1154 contains a synthetic DNA SEQ ID NO 1 in pUC19 encoding a fusion protein composed of: signal + cellulose binding domain + linker from C315, the amino acids KR, the variable region of a llama single chain antibody raised against the reactive dye RR6, and a repeat of the Z domain from protein A (pre-CBD_(C315)-KR::VhhRR6(R2)::Z::Z).

[0216] Expression vector pJaL1159 was constructed for transcription of the fusion protein pre-CBD_(C315)-KR::VhhRR6(R2)::Z::Z (SEQ ID NO 2). The plasmid pJaL1154 harboring the fusion protein was digested with *Bam*HI-*Xho*I. The 966 bp fragment was gel-purified and ligated into the *Aspergillus* expression cassette pMT2786 digested with *Bam*HI I-*Xho*I (a 6936 bp fragment). The ligation mixture was transformed into *E. coli* MT173 using the *Saccharomyces cerevisiae* Leu2 gene as selective marker to create the expression plasmid pJaL1159. The amplified plasmid was recovered using a QIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

[0217] Plasmid pMT2786 comprise an expression cassette based on the *Aspergillus niger* neutral amylase II promoter fused to the *Aspergillus nidulans* triose phosphate isomerase non translated leader sequence (Na2/tpi promoter) and the *Aspergillus niger* amyloglycosidase terminator (AMG terminator), the selective marker *amdS* from *Aspergillus nidulans* enabling growth on acetamide as nitrogen source and having the *Saccharomyces cerevisiae* Leu2 gene for selection in *E. coli*.

Example 2b**Construction of *Aspergillus* expression cassette pJaL1171 pre-CBD_(C315)-KR::VhhRR6(R2)::Z**

[0218] For construction of the fusion protein pre-CBD_(C315)-KR::VhhRR6(R2)::Z (SEQ ID NO 3) plasmid pJaL1154 was digested with *Bgl*II and the 3472 bp fragment was gel-purified and ligated with itself resulting in pJaL1169. The 795 bp *Bam*HI-*Xho*I fragment from pJaL1169 was purified and ligated into the *Aspergillus* expression cassette pMT2780 digested with *Bam*HI I and *Xho*I (a 6936 bp fragment). The ligation mixture was transformed into *E. coli* MT173 using the *Saccharomyces cerevisiae* Leu2 gene as selective marker to create the expression plasmid pJaL1171. The amplified plasmid was recovered using a QIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

Example 2c

Construction of *Aspergillus* expression cassette pJaL1164 AMG_(1-526aa)::Z::Z

[0219] Plasmid pJaL1153 contains a synthetic DNA SEQ ID NO 4 in pUC19 encoding a fusion protein composed of: *Aspergillus niger* AMG DNA encoding amino acids 488 - 526 and a repeat of the Z domain from protein A (AMG_(488-526aa)::Z::Z).

[0220] Plasmid pToC100 contains the *Aspergillus niger* AMG (Boel et al. EMBO Journal 3 (1984), 1581-1585) regulated by the TAKA promoter from p775 and at the same time a BamHI site was introduced upfront of the AMG start codon. pToC100 was constructed by ligating the following fragments together: 4306 bp *BamHI-NcoI* fragment from p775, an adapter SEQ ID NO: 5 and SEQ ID NO: 6, 860 bp *BssHII-PstI* from pCAMG91 and 1410 bp *PstI-NcoI* fragment from pCAMG91.

[0221] Expression vector pJaL1164 was constructed for transcription of the fusion protein AMG_(1-526aa)::Z::Z (SEQ ID NO 7). A 1723 bp *BamHI-DraIII* fragment and a 458 bp *DraIII-XhoI* fragment was gel-purified from plasmid pToC100 and pJaL1153, respectively, and ligated into the *Aspergillus* expression cassette pMT2786 digested with *BamHI-XhoI* (a 6936 bp fragment). The ligation mixture was transformed into *E. coli* MT173 using the *Saccharomyces cerevisiae* *Leu2* gene as selective marker to create the expression plasmid pJaL1164. The amplified plasmid was recovered using a QIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

Example 2d**Construction of *Aspergillus* expression cassette pJaL1165 AMG_(1-526aa)::Z**

[0222] For construction of an expression plasmid encoding for the fusion protein AMG_(1-526aa)::Z (SEQ ID NO 8) plasmid pJaL1153 was digested with *BglII* and the 2969 bp fragment was gel-purified and ligated with itself resulting in pJaL1158. A 1723 bp *BamHI-DraIII* fragment from pToC100 and a 287 bp fragment from pJaL1158 was purified and ligated into the *Aspergillus* expression cassette pMT2786 digested with *BamHI* and *XhoI* (a 6936 bp fragment). The ligation mixture was transformed into *E. coli* MT173 using the *Saccharomyces cerevisiae* *Leu2* gene as selective marker to create the expression plasmid pJaL1165. The amplified plasmid was recovered using a QIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

Example 2e**Construction of *Aspergillus* expression cassette pJaL1168 pre-CBD_(C315)::Z::Z::CBD_(EGV)**

[0223] Construction of the expression plasmid pJaL1168 encoding for the fusion protein pre-CBD_(C315)::Z::Z::CBD_(EGV) (SEQ ID NO 9) was done by amplification by PCR: 1) of the pre-CBD_(C315) region using pA2C315 as template and the primer pair 8683/CBD:Z-NA (SEQ ID NO 10 and 11), 2) of the Z::Z region using pJaL1153 as template and the primer pair Z-NA/Z-CA (SEQ ID NO 12 and 13) and 3) of the CBD_(EGV) region using pSX320 as template and the primer pair Z-CA: CBD/8654 (SEQ ID NO 14 and 15), resulting in 3 DNA fragments of 337 bp, 382 bp and 344 bp, respectively. The 3 fragments were mixed and used as template for amplification by PCR with primer pair 8653/8654 of a 983 bp fragment. The PCR fragment was digested with *BamHI-HindIII* and the 798 bp fragment was purified and clone ligated into the *Aspergillus* expression cassette pJaL790 digested with *BamHI-HindIII* (a 7386 bp fragment). The ligation mixture was transformed into *E. coli* DB6507 using the *Saccharomyces cerevisiae* *Ura3* gene as selective marker to create the expression plasmid pJaL1168. The amplified plasmid was recovered using a QIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

[0224] Plasmid pJaL790 comprised an expression cassette based on the *Aspergillus niger* neutral amylase II promoter fused to the *Aspergillus nidulans* triose phosphate isomerase non translated leader sequence (Na2/tpi promoter) and the *Aspergillus niger* amyloglycosidase terminator (AMG terminator), the selective marker *amdS* from *Aspergillus oryzae* enabling growth on acetamide as nitrogen source.

Example 2f

Construction of *Aspergillus* expression cassette pJaL1170 pre-CBD_(C315)-KR::Z::Z::CBD_(C315)::CBD_(EGV)

[0225] Plasmid pJaL802 is an *Aspergillus* expression plasmid builds on pJaL790 which contains a DNA (SEQ ID NO 16) encoding for the fusion protein pre-CBD_(C315)::CBD_(EGV) (SEQ ID NO 17).

[0226] Construction of the expression plasmid pJaL1170 encoding for the fusion protein pre-CBD_(C315)-KR::Z::Z::CBD_(C315)::CBD_(EGV) (SEQ ID NO 18) was done by amplification by PCR: 1) of the pre-CBD_(C315)-KR region using pA2C315 as template and the primer pair 8683/CBD:Z-NB (SEQ ID NO 10 and 19), 2) of the Z::Z region using pJaL1153 as template and the primer pair Z-NB/Z-CB (SEQ ID NO 20 and 21) and 3) of the CBD_(C315)-CBD_(EGV) region using pJaL802 as template and the primer pair Z-CB:CBD/8654 (SEQ ID NO 22 and 15), resulting in 3 DNA fragments of 343 bp, 382 bp and 443 bp, respectively. The 3 fragments were mixed and used as template for amplification by PCR with primer pair 8653/8654 of a 1088 bp fragment. The PCR fragment was digested with BamHI-HindIII and the 894 bp fragment was purified and clone ligated into the *Aspergillus* expression cassette pJaL790 digested with *Bam*H I-*Hind*III (a 7386 bp fragment). The ligation mixture was transformed into *E. coli* DB6507 using the *Saccharomyces cerevisiae* *Ura3* gene as selective marker to create the expression plasmid pJaL1170. The amplified plasmid was recovered using a QIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

Example 2g**Expression of DAP in *Aspergillus oryzae* strains**

[0227] The *Aspergillus oryzae* strains BECh2 was transformed with the expression plasmid pJaL1159, pJaL1164, pJaL1165, pJaL1168, pJaL1170 and pJaL1171 as described under methods.

A shake flask containing 10 ml YPM medium (2 g/l yeast extract, 2 g/l peptone, and 2% maltose) was inoculated with spores from the generated transformants and the host BECh2 and incubated at 30°C, with shaking (200 rpm) for 4 days. Supernatants (10 µl) were analysed on SDS-page. A transformant producing the desired protein from each plasmid pJaL1159, pJaL1164, pJaL1165, pJaL1168, pJaL1170 and pJaL1171 was named JaL1210, JaL1168, JaL1171, JaL1174, JaL1176 and JaL1181, respectively. Products of the expected size from each transformant were confirmed by SDS-page. The amino acids sequence of each construct produced in JaL1210 (VhhRR6(R2)::Z::Z), JaL1168 (AMG_(1-526aa)::Z::Z), JaL1171 (AMG_(1-526aa)::Z), JaL1174 (CBD_(C315)::Z::Z::CBD_(egv)), JaL1176 (Z::Z::CBD_(C315)::CBD_(egv)) and JaL1181 (VhhRR6(R2)::Z) are shown in SEQ ID NO 23, SEQ ID NO 24, SEQ ID NO 25, SEQ ID NO 26, SEQ ID NO 27 and SEQ ID NO 28, respectively.

Example 3. Recombinant dual affinity constructs for expression in *Bacillus licheniformis*.**Media**

[0228] LB agar, TY buillon medium and BPX shake flask medium have all been described in Patent Publication WO 94/14968.

[0229] CAL 18-2 media (11): Yeast extract (#0127-17-9 Difco Laboratories, MI, USA) 40g; Magnesium Sulfate (#5886 Merck, Darmstadt, Germany) 1.3g; Glucidex 12 (Roquette Feres, France) 50g; Sodium Di-hydrogenphosphate (#6346 Merck, Darmstadt, Germany) 20g; EDF-Tracemetals (recipe see below) 6.7ml; Na₂MoO₄-Tracemetal (recipe see below) 6.7ml; Pluronic PE6100 (BASF, Germany) 0.1ml; Ionexchanged water adjust to 1000ml. All is mixed, volume is adjusted, pH is measured and adjusted to pH 6.0 using NaOH. The media is sterilised by autoclaving at 121 °C for 20 min. EDF-Tracemetals (11): Mangan (II) sulphate (#5963 Merck, Darmstadt, Germany) 4.48g; Iron (III) chloride (#3943 Merck, Darmstadt, Germany) 3.33g; Copper (II) sulphate (#2790 Merck, Darmstadt, Germany) 0.625g; Zinksulphate (#8883 Merck, Darmstadt, Germany) 7.12 g; Ionexchanged water adjust to 1000ml. All is mixed, volume is adjusted. Solution is filter-sterilized and kept at 4°C. Na₂MoO₄-Tracemetals (11): SodiumMolybdat (#6521 Merck, Darmstadt, Germany) 2.0g; Ionexchanged water adjust to 1000ml. All is mixed, volume is

adjusted. Solution is filtersterilized and kept at 4°C.

Strains and Donor Organisms

[0230] *Bacillus subtilis* PL1801. This strain is the *B. subtilis* DN1885 with disrupted *apr* and *npr* genes (Diderichsen, B. , Wedsted, U., Hedegaard, L., Jensen, B. R., Sjöholm, C. (1990) Cloning of *aldB*, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*. J. Bacteriol., 172, 4315-4321).

Bacillus subtilis PP289-5. This strain is a donor strain for conjugation of plasmids to *Bacillus licheniformis* described in US5843720 *Bacillus licheniformis* MDT223 described in patent WO2005/123915

Genes

[0231] Z: indicate the Z domain from *Staphylococcus aureus* preotein A (Nilsson et al. Protein Engineering 1 (1987) 107-113). The Z gene is a synthetic gene assembled by primers (SEQ ID NO 29)

Streptavidin gene: indicate the gene encoding Streptavidin from *Streptomyces avidinii* as described by Argarana (Argarana et. Al. (1986) Nucleic Acids Res. 14, 1871-1882). The gene encoding streptavidin is a synthetic gene (SEQ ID NO 30)

[0232] A synthetic gene can be constructed by PCR assembly of overlapping oligonucleotides in various methods described eg. by Stemmer et al, Gene 164, pp-49-53, 1995; Dillon and Rossen, BioTechniques 9, 298-300, 1990; Prodromou and Pearl, Protein Engineering 5, 827-829, 1992; Chn et al., Journal of American Chemical Society 11, 8799-8800, 1994 and others. Such genes may also simply be purchased through one of many commercial companies.

Plasmids

[0233] pSJ6208 is an *E. coli* pUC derivative described in SEQ ID NO 31.

pSJ6321 is a pE194 derivative with an erythromycin marker gene. The plasmid also holds *cryIIIA* stabilizer sequence, DNA encoding the signal peptide of *amyl* fused to a protease followed by a downstream sequence of *amyl* (SEQ ID NO 32).

pMOL2743 is described in this example (SEQ ID NO 33)

pMOL2744 is described in this example (SEQ ID NO 34)

pMOL2746b is described in this example (SEQ ID NO 35)

Construction of integration vector Z::Z::streptavidin for *B. licheniformis* expression

[0234] The synthetic gene (SEQ ID NO 29) encoding the Z::Z domaine was amplified by the primers SEQ ID NO 36 and SEQ ID NO 37:

SEQ ID NO 36: TCATTCTGCAGCAGCGGCGGATAACAAATTTAACAAGAAGACAGCA-GAACGCGTTTTATGAAA

SEQ ID NO 37: AACTAAGCGGCCGCTAGCGACTACACTTTAGGAGCTTGCGCGTCAT-TAAGCT

[0235] The PCR fragment was digested with *PstI*- *EagI* and the 368 bp fragment was purified and ligated into the *E. coli* pUC derivative plasmid pSJ6208 (SEQ ID NO 31) digested with *PstI* - *EagI* giving a 3389 bp fragment. The ligation mixture was transformed into *E. coli* SJ2 (Diderichsen, B. , Wedsted, U. , Hedegaard, L. , Jensen, B. R. , Sjöholm, C. (1990). Cloning of *aldB*, which encodes acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*). The plasmid holding the sub cloned Z::Z gene pMOL2743 (SEQ ID NO 33), was recovered using a OIAprep® Spin Miniprep kit (QIAGEN, Chatsworth, CA) according to the manufacturer's instructions.

[0236] The synthetic gene (SEQ ID NO 30) encoding streptavidin was digested with *HindIII* giving a fragment of 620 bp. The plasmid pMOL2743 was restricted with *HindIII* and treated with alkaline phosphatase to avoid relegation. The two fragments were ligated and transformed to the *E. coli* SJ2. The colonies were screened for presence of the streptavidin gene and clones were picked where the streptavidin gene is inserted in the right orientation giving rise to the plasmid pMOL2744 (SEQ ID NO 34). In this

plasmid the genes encoding the Z::Z domaine and Streptavidin if translationally fused.

[0237] The hybrid gene encoding Z::Z::Streptavidin was transferred to an integration vector designed to allow integration of the amylase expression cassette into the chromosome of a *B. licheniformis* strain, that already contains an artificial tandem promoter integrated at the *amyL* locus, as described in example 6 of WO2005/123915. This was achieved by a ligation of three fragments . The first fragment is a PstI - BglII restriction digest of pMOL2744 giving rise to a 931 bp fragment. The second fragment is a BglII - BamHI restriction digest of the plasmid pSJ6321 (SEQ ID NO 32) isolating a 4288 bp fragment. The third fragment is a BamHI - PstI digest of pSJ 6321 isolating a 1234 bp fragment. The three fragments were ligated and introduced by transformation into PL1801 giving rise to an integration vector pMOL2746b (SEQ ID NO 35).

[0238] This pMOL2746b plasmid is then re-transformed by either competence, electroporation or conjugation into a protease deficient *Bacillus licheniformis* and inserted by double homologous integration at the *amyL* locus using an already inserted cryIIIA sequence and the *amyL* downstream sequence. The resulting *Bacillus licheniformis* strain has the artificial tandem promoter and cryIIIA sequence driving the Z::Z::Streptavidin expression from the *amyL* locus. The *Bacillus licheniformis* host is preferred to be protease deficient to allow expression of the Z::Z::Streptavidin hybrid protein. The following proteases can be deleted by standard techniques using double homologous recombination: *mpr*, *aprE*, *nprE*, *vpr*, *bpr*, *epr*, *wprA* and *ispA*.

[0239] The protease deficient *Bacillus licheniformis* host with the expression cassette encoding the Z::Z::Streptavidin hybrid DAP protein is fermented in 100 ml shake flasks with CAL18-2 medium described above at 30°C, 300 rpm for 2 days. Samples are taken out day one and day two to evaluate the DAP expression on an SDS gel. The data show a protein band at the right size of 30 KDa.

[0240] The PCR fragment is digested with the restriction enzymes Rsa I and Hind III and the resulting 489 bp fragment is cloned into the 5327 pStrExp1 digested with the restriction enzymes Nru I and Hind III by DNA ligation and cloning into *B. subtilis* PL1801.

[0241] The DNA sequence of the open reading frame encoding the fusion protein is shown in SEQ ID NO: 38, and the protein sequence in SEQ ID NO: 39.

Example 4: Expression of the DAP construct in *Bacillus subtilis*

[0242] Transformants of *B. Subtilis* PL1801 is grown in shake flasks as described in patent WO 2000/075344 and the fusion protein is recovered from the supernatant. The antibody binding and biotin binding properties of the fusion protein is confirmed as shown elsewhere herein.

Example 5. Purification of DAP

5a. Chemically synthesized DAP purified by size exclusion chromatography

[0243] The conjugation mixtures from example 1 were loaded on a size exclusion column in order to purify the DAP-molecules by removing reactants. The size exclusion chromatography was performed on a prepacked Superdex 200 16/60 column. 1 mL of DAP reaction solution was loaded on the column. The pump flow was 1.00 mL/min, the eluent was 150 mM NaCl, 50 mM Hepes pH 7.0 and fractions of 1 mL were collected. Fractions were pooled based on absorbance measurements at 280 nm in order to remove reactants.

[0244] The collected and pooled fractions were concentrated 10x using Amicon Ultracentrifuge tubes with a NWCO of 3000.

5b. Recombinant DAP by IgG-sepharose affinity chromatography

[0245] Recombinant DAP molecules were purified from the fermentation broth by sterile filtration and subsequent column purification by IgG affinity chromatography.

[0246] The IgG-sepharose column was prepared by following the procedure supplied by the vendor:

15 g CNBr-activated sepharose 4B from GE Healthcare was washed for 15 min with 3 L 1 mM HCl. The washed medium was added to 25 mL 20 mg/mL IgG solution from DAKO A/SA (X0903) and 50 mL 0.75 M NaCl, 0.15 M NaHCO₃ pH 8.3. The mixture was gently rotated for 95 min. at room temperature. Excess IgG was washed away with 75 mL 0.1 M NaHCO₃ pH 8.3 containing 0.5 M NaCl before the medium was incubated in 0.1 M Tris/HCl pH 8.0 for 2 hours. The medium was stored in 20% ethanol until use.

Generic procedure for purification of recombinant DAP

[0247] The recombinant DAP molecules from examples 2 and 3 were purified by a generic affinity chromatographic method that takes advantage of the IgG binding domain shared by all the recombinant DAP constructs. The DAP molecules were purified from the sterile filtered fermentation broth.

[0248] The chromatography was performed on a XK26/20 column packed with approx. 30 mL IgG-sepharose. The fermentation broths were sterile filtered and between 65 mL and 80 mL was loaded depending on the volume of the filtered fermentation broth. The pump flow was 1.50 mL/min during sample load and 2 mL/min during wash and elution. Buffer A was 0.1 M NaH₂PO₄ pH 7.2, 0.15 M NaCl and buffer B: 0.1 M Citric acid pH 3.5. The column was washed with 15 column volumes buffer A following the sample load. The bound material was eluted with 5 column volumes buffer B before the column was regenerated with 10 column volumes buffer A. Fractions of 10 mL were collected. Eluate having an increase of absorbance at 280 nm was pooled and the pH was adjusted to 7.2 using 1 M Tris. The concentration of the DAP protein was calculated from the absorption at 280 nm and the theoretical absorption coefficients calculated from the primary sequence using GPMaw 8.0 (Trends in Biochemical Science, Vol 26, No. 11, November 2001, pp 687-689, "GPMaw - a software tool for analyzing proteins and peptides"; see also <http://www.gpmaw.com/>). The Mw of the purified proteins were determined by SDS-PAGE. The samples were stored at -18°C before further analysis.

Example 6. Characterization of DAP molecules with regard to binding strength measured on the Biacore instrument.

[0249] The commercialised surface plasmon resonance (SPR) technology for real time monitoring biomolecular binding event is used to measure the binding affinities for the prepared DAP candidates. The general principle of this technology is that a SPR sensor chip measures changes in refractive index, and the changes in refractive index correlate to changes in mass in the aqueous layer close to the sensor surface. When target molecules in solution bind to ligands, immobilised on the sensor surface, the mass increases, and when they dissociate from the ligands the mass decreases. This principle facilitates a continuous, real time monitoring of the association and dissociation of interacting molecules. The graphical presentation of the relationship provides quantitative information in real time on the binding specificity, active concentration of molecule in a sample, reaction kinetics and affinity.

[0250] In order to evaluate the binding affinities for both of the binding affinities present in the DAP either IgG or the capturing ligand had to be immobilized onto the sensor chip. For IgG rabbit anti-Mannanase was used and for the ligand the chip was coated with streptavidin and the ligand was then immobilized via biotin linked to the appropriate ligand.

[0251] In order to measure the binding to RR6 and acarbose the following compounds were prepared; biotin-linked acarbose and biotin-linked RR6.

Preparation of biotin linked acarbose

[0252] Biotin (5 mg, 26 µmol) was dissolved in DMF (250 µL) in a 2 mL eppendorf tube, and to this EDC-HCl (5 mg, 26 µmol) was added. The mixture was stirred at RT for 30 min. Acarbose (16.1 mg, 25 µmol) was dissolved in DMF (250 µL) in a 2 mL eppendorf tube, and to this the activated biotin solution was added dropwise, while the reaction mixture was gently stirred. When all activated biotin solution was added, 100 µL DMF was used to wash the reaction container, and these 100 µL were also added to the acarbose solution. The reaction mixture was left stirring for 2h at RT. Upon this, the DMF was removed by freeze-drying overnight at -5°C. The crude product was stored at -18°C, until used in the Biacore experiments.

Preparation of biotin linked Reactive Red 6

[0253] Biotin (5 mg, 26 μ mol) was dissolved in DMF (250 μ L) in a 2 mL eppendorf tube, and to this EDC-HCl (5 mg, 26 μ mol) was added. The mixture was stirred at RT for 30 min. 1,4-Diaminobutane (25.1 μ L, 25 μ mol) was dissolved in DMF (250 μ L) in a 2 mL eppendorf tube, and to this the activated biotin solution was added dropwise, while the reaction mixture was gently stirred. When all activated biotin solution was added, 100 μ L DMF was used to wash the reaction container, and these 100 μ L were also added to the diaminobutane solution. The reaction mixture was left stirring for 2h at RT.

[0254] Reactive Red 6 (24.4 mg, 25 μ mol) was dissolved in DMF (250 μ L) in a 2 mL eppendorf tube, and to this the biotin-amide solution was added dropwise, while the reaction mixture was gently stirred. When all biotin-amide solution was added, 100 μ L DMF was used to wash the reaction container and these 100 μ L were also added to the RR6 solution.

[0255] The reaction mixture was left stirring overnight at RT. The DMF was removed by freeze-drying overnight at -5°C. The crude product was stored at -18°C, until used in the Biacore experiments.

Biacore evaluation:

[0256] The DAP candidates were analysed for binding to the capturing ligand sensor chip and the target biomolecule sensor chip respectively. A Biacore 3000 instrument was used.

[0257] To study the interactions between the IgG-binding end of the DAP molecule and IgG on the one hand, and the interaction between the ligand-binding end and the ligand (biotin, acarbose, reactive red) on the other hand, IgG and ligand were immobilized onto the sensor surface of a sensor chip as described below.

[0258] Immobilisation occurred by direct covalent coupling to the surface (using the Amine Coupling Kit, Biacore, GE Health Care) or via a capturing molecule as prescribed by the manufacturer (Biacore, GE Health Care). The amount of coupled target was quantified and expressed in Refractive Units (RU).

[0259] Interactions were monitored by injecting samples (20 μ L/min) over the prepared sensor surface of the chip. Unless stated otherwise, the binding was assessed in 10 mM sodium acetatbuffer pH 5.0 at room temperature.

Experiment 1:

[0260] Chip: CM5

Immobilized target: rabbit anti-Mannanase (10 μ g/ml) in 10 mM acetatbuffer pH 5.0. Target RU: 1250 RU

Actual RU:

FC1: 1349

FC2: 1492

FC3: 1338

FC4: 1331

Samples:

[0261] FC1: Protein A, 1 μ g/ml injected

FC2: Protein A - Avidin, 0.5 μ g/ml injected

FC3: Protein A - Avidin, 0.5 μ g/ml injected

FC4: Affibody (IgG) - Avidin, 0.6 μ g/ml injected

Results:

[0262]

Protein A	ka	kd	KA	KD	Final RU
Chi ₂ = 0.06	4.35e5	3.4e-6	1.28e11	7.81 e-12	140
Protein A-Avidin	ka	kd	KA	KD	Final RU
Chi ₂ = 0.737	6.99e5	1.75e-7	3.99e12	2.5e-13	130
Protein A-Avidin	ka	kd	KA	KD	Final RU
Chi ₂ = 1.82	1.21 e6	5.03e-5	2.41e10	4.15e-11	130
Affibody(IgG)-Avidin	ka	kd	KA	KD	Final RU
Chi ₂ = 0.505	9.56e4	6.41 e-8	1.49e12	6.7e-13	105

Experiment 2:**[0263]** Chip: CM5

Immobilized target: rabbit anti-Mannanase (10 µg/ml) in 10 mM acetatbuffer pH 5.0.

Target RU: 1250 RU

Actual RU:

FC2: 1697

FC3: 1665

Samples:**[0264]** FC2: CBD-Z-Z-CBD, 10 µg/ml injected

FC3: CBD-Z-Z-CBD, 10 µg/ml injected

Results:**[0265]**

CBD-Z-Z-CBD	ka	kd	KA	KD	Final RU
Chi ₂ = 2.3	1.46e5	2e-4	7.3e8	1.37e-9	200
CBD-Z-Z-CBD	ka	kd	KA	KD	Final RU
Chi ₂ = 3.78	1.23e5	2.08e-4	5.93e8	1.69e-9	200

Experiment 3:**[0266]** Chip: CM5

Immobilized target: rabbit anti-Mannanase (10 µg/ml) in 10 mM acetatbuffer pH 5.0.

Target RU: 1250 RU

Actual RU:

FC3: 1485

FC4: 1760

Samples:**[0267]** FC3: AMG-Z, 10 µg/ml injected

FC4: AMG-ZZ, 10 µg/ml injected

Results:**[0268]**

AMG-Z	ka	kd	KA	KD	Final RU
Chi ₂ = 6.76	2.69e6	1.08e-2	2.49e8	4.02e-9	5
AMG-ZZ	ka	kd	KA	KD	Final RU
Chi ₂ = 2.97	3.46e5	4.1 e-4	8.42e8	1.19e-9	150

Experiment 4:**[0269]** Chip: CM5

Immobilized target: rabbit anti-Mannanase (1 µg/ml) in 10 mM acetatbuffer pH 5.0.

Target RU: 625 RU

Actual RU:

FC1: 683

FC2: 731

FC3: 881

FC4: 716

Samples:**[0270]** FC1: Protein A, 1 µg/ml injected

FC2: ZZ-CBD-CBD, 0.1 µg/ml injected

FC3: VhhRR6(R2)-Z, 1 µg/ml injected

FC4: CBD-Z-Z-CBD, 1 µg/ml injected

Results:**[0271]**

Protein A	ka	kd	KA	KD	Final RU
Chi ₂ = 1.22	4.27e5	1.26e-4	3.38e9	2.95e-10	55
ZZ-CBD-CBD	ka	kd	KA	KD	Final RU
Chi ₂ = 0.116	4.6e6	3.3e-4	1.39e10	7.18e-11	70
VhhRR6(R2)-Z	ka	kd	KA	KD	Final RU
Chi ₂ = 2.35	3.2e5	1e-3	3.2e8	3.13e-9	70
CBD-Z-Z-CBD	ka	kd	KA	KD	Final RU
Chi ₂ = 1.96	2.65e5	2.97e-4	8.94e8	1.12e-9	50

Experiment 5:**[0272]** Chip: SA

Immobilization: None. The Chip is pre-coated with Streptavidin.

[0273] Immobilisation of the ligand through Biotin-Streptavidin binding

[0274] FC1: Biotin- Acarbose, 10 µg/ml

FC2: Biotin- Acarbose, 10 µg/ml

FC3: Biotin- Reactive Red 6, 10 µg/ml

[0275] Binding of the DAP via the ligand-binding end of the molecule to the ligand.

[0276] FC1: AMG-Z-Z, 10 µg/ml

FC2: AMG-Z, 10 µg/ml

FC3: VhhRR6(R2)-Z, 10 µg/ml

Results:

[0277]

AMG-Z-Z	ka	kd	KA	KD	Final RU
Chi ₂ = 1.38	6.29e3	2.69e-3	2.34e6	4.27e-7	15
AMG-Z	Ka	Kd	KA	KD	Final RU
Chi ₂ = 2.9	1.02e4	1.8e-3	5.64e6	1.77e-7	20
VhhRR6(R2)-Z	Ka	Kd	KA	KD	Final RU
Chi ₂ = 3.24	1.72e4	3.99e-3	4.31 e6	2.32e-7	15

Experiment 6:

[0278] Chip: SA

Immobilization: None. The Chip is pre-coated with Streptavidin.

Immobilisation of the ligand through Biotin-Streptavidin binding

[0279] FC4: Biotin-Reactive Red 6, 10 µg/ml

[0280] Binding of the DAP via the ligand-binding end of the molecule to the ligand in the presence of IgG binding to the Z-domain of the DAP molecule.

[0281] FC4: VhhRR6(R2)-Z + IgG 9.3 µg/ml

Result:

[0282]

VhhRR6(R2)-Z + IgG	ka	kd	KA	KD	Final RU
Chi ₂ = 5.3	225	6.18e-3	3.65e4	2.74e-5	5

Table 1

DAP	IgG binding			Ligand	Ligand binding		
	k _a (1/Ms)	k _d (1/s)	K _D (M)		k _a (1/Ms)	k _d (1/s)	K _D (M)
Avidin-Protein A	1×10 ⁶	3×10 ⁻⁵	2×10 ⁻¹¹	Biotina	-	-	-
Avidin-Affibody (IgG)	1×10 ³	6×10 ⁻⁸	7×10 ⁻¹³	Biotin	-	-	-
AMG-ZZ	4×10 ⁵	4×10 ⁻⁴	1×10 ⁻⁹	Acarbose	6×10 ³	3×10 ⁻³	4×10 ⁻⁷
AMG-Z	3×10 ⁶	1×10 ⁻²	4×10 ⁻⁹	Acarbose	1×10 ⁴	2×10 ⁻³	2×10 ⁻⁷

DAP	IgG binding				Ligand	Ligand binding	
	k_a (1/Ms)	k_d (1/s)	K_D (M)		k_a (1/Ms)	k_d (1/s)	K_D (M)
CBD-ZZ-CBD	2×10^5	2×10^{-4}	1×10^{-9}	Cellulose			
ZZ-CBD-CBD	5×10^6	3×10^{-4}	7×10^{-11}	Cellulose			
VhhRR6(R2)-Z	3×10^5	1×10^{-3}	3×10^{-9}	RR6b	2×10^4	4×10^{-3}	2×10^{-7}
Protein A control	5×10^5	7×10^{-5}	2×10^{-10}	-	-	-	-

^aBinding between biotin and avidin/streptavidin is not measured, since it is known to be very tight, and no dissociation can therefore be measured. ^bIf VhhRR6(R2)-Z is mixed with IgG before loading to the Biacore, the K_D is 10^{-5} M, however, the binding is broken completely as soon as the injection is stopped.

Example 7: Purification of antibodies using dual affinity polypeptide purification technology

[0283] Below is a short description of a generally applicable procedure for immunoglobulin purification:

1. A disposable generic solid phase with a low molecular weight capturing ligand molecule.
2. The dual affinity polypeptide DAP (e.g. Avidin-Protein A) reacts with the IgG (target biomolecule) in the solution. The DAP molecules are immobilized on the solid phase (e.g. biotin-agarose) in a complex together with the target protein (IgG).
3. The column is washed to eliminate interfering non-product components.
4. The immunoglobulin is eluted from the column using an appropriate buffer at low pH.
5. The antibody containing fractions are collected and pH is neutralized.
6. The generic matrix containing immobilized dual affinity polypeptide may be discarded.

Materials and Methods:

[0284] All buffer ingredients were pro-analysis. Rabbit serum proteins and Rabbit IgG fraction, code X0903 were from DAKO A/S, Denmark. Biotin Agarose was from Sigma-Aldrich; B6885-5ML.

[0285] The purification experiments were performed using a chromatography system comprising a fraction collector (Frac-100), a recorder (Rec-1), an optical unit and a control UV-1, all from GE Healthcare. For all experiments we used a BioRad Econo-column ID 1.0 cm with a flow adaptor. The equilibration and dilution buffer was 0.1 M phosphate, 0.15 M NaCl, pH=7.2 (PBS). The elution buffer was 0.1 M citrate, pH=3.5, or alternatively 0.1 M glycine, pH=2.8.

[0286] The data processing was based on Abs 280 nm measurement using a Pharmacia Gene Quant II and the extinction coefficient for Rabbit IgG (1 g/L) of 1.35.

IgG purification analysis

[0287] All experiments were performed at room temperature.

Example 7a - Recovery of Rabbit IgG with free DAP protein A - Neutravidin]

[0288] The sorbent (Biotin Agarose, 5 mL) was allowed to settle in the column for 10 min. The column was packed at a flow of 1.46 mL/min. The column was equilibrated with 7.5 column volumes (CV) PBS. DAP (Protein A-Neutravidin) solution (4.5 mL from example 1 b) was mixed with 100 μ L Rabbit IgG stock solution (20 g/L) and incubated for 5 minutes on a magnetic stirrer. The reaction solution was loaded on the column and washed with 7.5 CV of PBS to remove excess of target protein. The IgG was

recovered by elution with 3 CV of 0.1 M glycine buffer (pH 2.8). 5 mL fractions were collected and analyzed for IgG content by Abs 280. See Table 2 for results.

Example 7b - Recovery of Rabbit IgG with Immobilized DAP [Protein A - Neutravidin]

[0289] The gel material from example 7a with immobilized DAP [Protein A - Neutravidin] was regenerated with 7.5 CV PBS before analyzing the conventional affinity purification capabilities.

[0290] We loaded 2 mg IgG in 4.6 mL PBS solution on the column. After sample loading, the column was washed with 7.5 CV of PBS to remove excess of protein. Then the column was eluted with 3 CV of 0.1 M Glycine buffer (pH 2.8) to recover the IgG. 5 mL fractions were collected and analyzed for IgG content by Abs 280 and SDS-PAGE. See Table 2 for results.

Table 2.

	Flow through	Recovered
Example 1b - Free DAP	74%	26%
Example 1b - Immobilized DAP	86%	14%

[0291] As seen in Table 2, we obtained approximately twice the binding capability (26% versus 14%) when DAP and IgG are reacted in solution prior to loading on the column compared to the conventional affinity chromatography applying immobilized DAP.

Example 7c Purification of Rabbit IgG from serum with free DAP [Protein A - Neutravidin]

[0292] We studied purification of IgG from rabbit serum to show the specificity of the DAP purification technology.

[0293] Approximately 1 mL of sorbent (Biotin Agarose) was allowed to settle in the column for 10 min. The column was packed at a flow of 1.46 mL/min. The column was equilibrated with 7.5 column volumes (CV) PBS. A 4.5 mL DAP (Protein A - Neutravidin) solution (from example 1 b) were mixed with 115 μ L rabbit serum and incubated for 5 minutes on a magnetic stirrer. The reaction solution was loaded on the column following a wash with 7.5 CV of PBS to remove excess of target protein. The IgG was recovered by elution with 3 CV of glycine buffer. 2.5 mL fractions were collected and analyzed for IgG content by Abs 280 and SDS-PAGE. See Table 3 for results.

Example 7d Purification of Rabbit IgG from serum with immobilized DAP protein A - Neutravidin]

[0294] The gel from example 1 c with immobilized DAP [Protein A - Neutravidin] was regenerated with 7.5 CV PBS before analyzing the conventional affinity purification capabilities

[0295] We loaded rabbit serum (115 μ L rabbit serum in 4.5 mL PBS) solution on the column. After sample loading, the column was washed with 7.5 CV of PBS to remove excess of protein. Then the column was eluted with 3 CV of 0.1 M Glycine buffer (pH 2.8) to recover the target IgG. The gel was regenerated with 7.5 CV PBS before the next affinity purification cycle. 2.5 mL fractions were collected and analyzed for IgG content by Abs 280 and SDS-PAGE. See Table 3 for results.

Table 3

	Recovered IgG (mg) from serum
Example 1b - Free DAP	0,31
Example 1b - Immobilised DAP	0,18

[0296] As seen in Table 3, we obtained approximately twice the binding capability (0.31 mg versus 0.18 mg IgG), when DAP was reacted with rabbit serum in solution prior to contacting the biotin-agarose column compared to the conventional affinity chromatography applying immobilized DAP.

[0297] SDS-PAGE showed that only IgG molecules were obtained from serum, showing that the DAP purification technology is

specific.

[0298] We repeated the analysis using the other DAP conjugate (Protein A-Avidin) and performed similar tests as above but included an analysis of the leakage of DAP from the column by repeated binding analysis to the immobilized DAP.

Example 7e - Recovery of Rabbit IgG with free DAP Protein A - Avidin]

[0299] The sorbent (Biotin Agarose, 1 mL) was allowed to settle in the column for 10 min. The column was packed at a flow of 1.46 mL/min. The column was equilibrated with 7.5 column volumes (CV) PBS. 2 mL of DAP (Protein A - Avidin) solution (from example 3a) was mixed with 160 µL Rabbit IgG stock solution (20 g/L) and incubated for 5 minutes on a magnetic stirrer. The reaction solution (~2 CV) was loaded on the column following a wash with 7.5 CV of PBS to remove excess of target protein. The IgG was recovered by elution with 3 CV of 0.1 M Citrate, pH=3.5. 2.5 mL fractions were collected and analyzed for IgG content by Abs 280. The results are shown in Table 4.

Table 4. Free DAP versus reuses of immobilised DAP technology (Protein A - Avidin conjugate) Rabbit IgG recovery

	Free DAP	1. reuse of immobilised DAP	2. reuse of immobilised DAP	3. reuse of immobilised DAP	4. reuse of immobilised DAP
IgG Recovered (mg)	1,20	0,30	0,40	0,40	0,30

Example 7f - Recovery of Rabbit IgG with immobilized DAP [Protein A - Avidin]

[0300] The gel from example 7e with immobilized DAP [Protein A - Avidin] was regenerated with 7.5 CV PBS before analyzing the conventional affinity purification capabilities. The rabbit IgG stock solution (20 g/L) was diluted to a concentration of 1.5 mg/mL with PBS. In each runs 3.2 mg of IgG was loaded (in 2.16 mL) on the column. After sample loading (~2 CV), the column was washed with 7.5 CV of PBS to remove excess of protein. Then the column was eluted with 3 CV of elution buffer (0.1 M Citrate, pH=3.5) to recover the target IgG. The gel was regenerated with 7.5 CV PBS before the next affinity purification cycle. 2.5 mL fractions were collected and analyzed for IgG content by Abs 280.

[0301] When DAP is reacted with 3.2 mg of IgG in solution prior to contacting with the Biotin-Agarose we recovered 1.2 mg of IgG compared to the 0.35 mg IgG which was recovered on average in four repeated cycles with immobilized DAP.

[0302] The above results thus illustrate the advantageous effect of using a DAP according to the invention compared to conventional chromatography.

Example 8. Recovery of IgG using dual affinity chromatography

[0303] The purified DAP molecules from examples 5a and 5b were evaluated in a generic purification assay. The experiments were conducted at room temperature using an Äkta explorer system. 0.6 mL solid phase material was packed at a flow rate of 1.2 mL/min in an empty glass column (6.6 x 100 mm) equipped with an adjustable flow adaptor (Omnifit). The column was packed in 0.1 M sodium phosphate, 0.15 M NaCl, pH=7.2 (PBS) and allowed to equilibrate with 10 column volumes PBS followed by 3 column volumes 0.1 M citrate, pH=3.5 and finally 10 column volumes PBS before use.

[0304] ZZ-CBD-CDB and CBD-ZZ-CBD were analyzed using a column packed with cellulose. 1.2 g Avicel (Merck product no. 1.02331) was suspended in 8 ml PBS in a test tube and the suspension was allowed to settle for 30 min. Subsequently the fine particles were decanted before the column was packed.

[0305] AMG-Z and AMG-ZZ were evaluated using a column packed with Acarbose-agarose. Approx. 0.6 mL of the Acarbose-agarose from Example 9 was transferred to the column and allowed to settle for 10 min before the column was packed.

[0306] VhhRR6(R2)-Z was analysed using a column packed with RR6-agarose. Approx. 0.6 mL of the Reactive Red-agarose from Example 9 was transferred to the column and allowed to settle for 10 min before the column was packed.

[0307] Avidin-Protein A, Avidin-Affibody and ZZ-Streptavidin were evaluated using a column packed with Biotin-agarose. Approx. 0.6 mL of Biotin-agarose (Sigma B6885) was transferred to the column and allowed to settle for 10 min before the column was packed.

[0308] The packed column was operated at a flow rate of 0.6 mL/min. Buffer A was 0.1 M sodium phosphate pH 7.2, 0.15 M NaCl and buffer B was 0.1 M Citric acid pH 3.5. The column was initially washed with 10 column volumes buffer A before 0.6 mL sample was injected. The column was washed with 7.5 column volumes buffer A and the bound target protein was eluted with 5 column volumes buffer B. The column was finally regenerated with 10 column volumes buffer A. Detection was at 280 nm. The data were evaluated by determining the height of the peak observed during elution.

[0309] Purified DAP (8 nmole) was mixed with IgG (code X0903, DAKO A/S, 8 nmole) and water was added ad 660 μ L. The reaction mixture was incubated on a magnetic stirrer for 10 minutes before it was injected onto the column. The solution of target protein was prepared as a 2 mg/mL IgG solution in water. The following sequence of injections was carried out in all experiments: Water; target protein (7.1 nmole); target protein and DAP reaction mixture; and finally 10 times subsequent injections of target protein (7.1 nmole).

[0310] The column employed for evaluating the Protein A - Avidin DAP molecule was subsequently used for assessing the effect of changing the load of target protein by varying the injection volume. Four injections were made: (0.6; 0.45; 0.3; 0.15)mL of the same IgG solution (12 μ M). The results showed that the height of the peak observed during elution was almost constant whereas the height of peak observed in the flow through decreased markedly as the column load was lowered (Table 5). These results are in accordance with the nature of affinity chromatography and demonstrate that the applied approach of evaluating the data by using the height of the peak observed during elution is valid.

Table 5 Peak heights determined from injections of different volumes of IgG

Injection volume	Peak height of flow through	Peak height eluate
mL	mAU	mAU
0.6	154	126
0.45	116	124
0.3	61	120
0.15	14	109

[0311] The non-specific binding of target protein to the column was evaluated by injecting water and subsequently the target protein before the DAP molecule was introduced to the column material. The peak heights observed from injections of water and IgG were comparable in all the experiments performed. This demonstrates that the peak observed during elution is unaffected of potential non-specific binding of target protein to the column. Thus the peak observed during elution is a measure of the recovered amount of target protein from the non-covalently immobilized DAP.

[0312] The ability of the DAP molecules to recover the target protein was analyzed by comparing the chromatograms obtained from injection of water, target protein and the target protein/DAP reaction mixture. The results are shown in Table 6. Only two DAP molecules were unable to recover the target protein. 1) VhhRR6(R2)-Z did not recover IgG, which may be explained by results from the Biacore analysis showing that the binding between DAP and ligand is broken completely as soon as the injection is stopped (Table 1 footnote). This indicates that the DAP molecule is quickly released from the solid phase and thus not suitable for affinity chromatography. 2) The ZZ-streptavidin DAP did not recover IgG which is likely explained by a blocking of the biotin binding sites due to the reaction with endogenous biotin present in the fermentation broth. Thus this ZZ-streptavidin preparation is likely not to bind to the solid phase.

Table 6 Ability of DAP molecules to recover the target protein

DAP	Column materiale	Binding and elution of target protein
AMG-ZZ	Acarbose-agarose	+
AMG-Z	Acarbose-agarose	+
ZZ-CBD-CBD	Cellulose	+
CBD-ZZ-CBD	Cellulose	+
VhhRR6(R2)-Z	Reactive red-agarose	-
ZZ-streptavidin	Biotin-agarose	-

DAP	Column materiale	Binding and elution of target protein
Affibody(IgG)-avidin	Biotin-agarose	+
ProteinA-avidin	Biotin-agarose	+

[0313] The leakage of DAP from the column was assessed by 10 consecutive injections of the target protein following the initial injection of the DAP/target protein reaction mixture. The heights of the peaks observed during elution were determined and the relative responses were calculated relative to the first injection of IgG. The relative responses were plotted as a function of the injection number and the relative decrease in peak height was calculated by linear regression. The results from the different DAP - ligand combinations are shown in Table 7 together with the dissociation constants.

Table 7 Dissociation constants and relative leakage

DAP	$K_{D,t}$ M	$K_{D,s}$ M	$K_{D,t}/K_{D,s}$	Rel decrease %/inj	r^2
AMG-ZZ	1×10^{-9}	4×10^{-7}	3×10^{-3}	-1.1	0.8
AMG-Z	4×10^{-9}	2×10^{-7}	2×10^{-2}	-1.7	1.0
ZZ-CBD-CBDa	7×10^{-11}	10^{-6}	7×10^{-5}	-1.7	1.0
CBD-ZZ-CBDa	1×10^{-9}	10^{-6}	1×10^{-3}	-1.7	0.9
Affibody(IgG)-avidin ^b	7×10^{-13}	10^{-15} ^a	7×10^2	-0.2	0.3 ^c
Protein A-avidin ^b	2×10^{-11}	10^{-15} ^a	2×10^4	-0.7	0.9

^aThe dissociation constant (K_D) of CBDs binding to cellulose is generally considered to be $\approx 10^{-6}$ M (Linder et al, Biotechnology and Bioengineering, Vol. 60, No. 5, Dec. 5, 1998). ^bThe dissociation constant (K_D) of avidin binding to biotin is well known to be 10^{-15} M (Green, N. (1963). Biochem J, 89, 585-591). ^cThe correlation coefficient reflects that the calculated leakage is mainly determined by a single point. The relative leakage is -0.05% with $r^2 = 0.0$ if this point is removed.

[0314] The results in Table 7 show purification schemes of the same target molecule (IgG) using various compositions of DAP molecules. It is concluded that the most efficient DAP molecules in affinity chromatography are those with tighter binding to the ligand on the matrix, i.e. those having a relative $K_{D,t}/K_{D,s} > 10^0 = 1$.

[0315] Specifically, the strong binding towards the column provided by the avidin-biotin bond prevents leakage of the bound DAP molecules.

Example 9. Preparation of functionalized resins

Materials

[0316]

Resin:	Mini-Leak-Low (loading 2-5 mM, Kem-En-Tec).
Ligands:	1,4-Diaminobutane ([110-60-1], Sigma-Aldrich, D13208), Reactive Red 6 (Cherry red #14, Grateful Dyes inc.), Acarbose ([56180-94-0] Sigma-Aldrich, A8980).
Coupling buffer:	0.5M K_2HPO_4 - pH 8.5
Washing buffer:	0.5M K_2HPO_4 - pH 7.0
Blocking buffer	0.1 M Ethanolamine in Milli-Q water

Preparation of an RR6-agarose resin

[0317] Resin (10 ml, suspended) was washed 2 X with distilled water and the water removed by filtration. 1,4-Diaminobutane (2.0 mL) was dissolved in coupling buffer (20 mL) and the resin was added slowly with gentle shaking. The resin was left shaking overnight at RT, whereupon it was washed with coupling buffer and sucked dry.

[0318] Reactive Red 6 (15.9 g) was dissolved in coupling buffer (50 mL) and to this solution, the amino-functionalized resin was added slowly with shaking. Again the resin was left shaking overnight at RT. After washing with water and washing buffer, the resin was transferred to blocking buffer (20 mL) and shaken for 2 h. Finally the resin was washed in water until the filtrate was colorless, and the red resin was suspended in 30% ethanol in Milli-Q water.

Preparation of an acarbose-agarose resin

[0319] Resin (10 ml, suspended) was washed 2 X with distilled water and the water removed by filtration. Acarbose (500 mg) was dissolved in coupling buffer (20 mL) and the resin was added slowly with gentle shaking. The resin was left shaking overnight at RT, whereupon it was washed with water and washing buffer. The resin was transferred to blocking buffer (20 mL) and shaken for 2 h. Finally the resin was washed in water, and the resulting resin was suspended in 30% ethanol in Milli-Q water.

SEQUENCE LISTING

[0320]

<110> Novozymes A/S Kyhse Andersen, Jan

<120> Dual affinity polypeptides for purification

<130> 11297.204-WO

<160> 39

<170> PatentIn version 3.5

<210> 1

<211> 972

<212> DNA

<213> Artificial

<220>

<223> fusion protein coding sequence

<400> 1

```

ggatccacca tgaaggcgat cctctctctc gctgccgcgc tcttgagcgc cgcgcccgcg      60
ttctcgaccg cagtctgggg ccagtcggtt ggcatcggct tctccggcga caccacgtgc      120
accgcttcca cctgcgtgaa ggtcaacgac tactactcgc agtgccagcc tggcgcgtcc      180
gcacccacgt caaccgcctc cgcgcccggc ccgtcccgct gccctttggt gaagcgtcag      240
gtgcagtttg aggaatcggg tggaggactc gtgcaggcag gtggctcgct ccggttgcg      300
tgtgcagcct cgggcagggc gacatcggga catggccatt acggtatggg atggttcgc      360
cagggtgcctg gaaaggaacg agaattcgtc gcagccatcc ggtggtccgg caaagagacc      420
tgggtacaaag attcggtgaa aggacgggtc accatctcca gggataacgc gaaaactact      480
gtctacctcc agatgaactc gctcaagccc gaggatacgg cagtgtatta ctgtgccgca      540
aggcctgtcc gagtggacga catctcgctc ccggtgggtt tcgattactg gggacagggt      600
acgcagggtg cggctctcctc cgataacaaa ttcaacaagg aacagcagaa cgcgttctac      660
gagatcttgc acctcccaa cctcaacgag gaacagagga acggtttcat tcagtcgctc      720
aaggacgacc cctcccgatc cgccaacctc ttggcagaag ccaagaagtt gaacgatgca      780
caggcaccga aagacaacaa gttcaacaaa gagcagcaga acgccttcta tgagatcttg      840
catctcccgga acttgaacga ggagcagcgg aacggcttca tccagtcctt gaaagatgat      900
ccctcgagtc cggccaacct cctcgagaaa gcgaagaagc tcaacgacgc gcaggcacct      960
aagtgactcg ag                                         972

```

<210> 2

<211> 318

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 2

```

Met Lys Ala Ile Leu Ser Leu Ala Ala Ala Leu Leu Ser Ala Ala Pro
1           5           10          15

```

Ala Phe Ser Thr Ala Val Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser
20 25 30

Gly Asp Thr Thr Cys Thr Ala Ser Thr Cys Val Lys Val Asn Asp Tyr
35 40 45

Tyr Ser Gln Cys Gln Pro Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser
50 55 60

Ala Pro Gly Pro Ser Ala Cys Pro Leu Val Lys Arg Gln Val Gln Leu
65 70 75 80

Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly Ser Leu Arg Leu
85 90 95

Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly His Gly His Tyr Gly
100 105 110

Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu Arg Glu Phe Val Ala
115 120 125

Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr Lys Asp Ser Val Lys
130 135 140

Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Thr Thr Val Tyr Leu
145 150 155 160

Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
165 170 175

Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu Pro Val Gly Phe Asp
180 185 190

Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser Asp Asn Lys Phe
195 200 205

Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn
210 215 220

Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp
225 230 235 240

Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp
245 250 255

Ala Gln Ala Pro Lys Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala
260 265 270

Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn
275 280 285

Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu
290 295 300

Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys
305 310 315

<210> 3

<211> 261

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 3

Met Lys Ala Ile Leu Ser Leu Ala Ala Ala Leu Leu Ser Ala Ala Pro
1 5 10 15
Ala Phe Ser Thr Ala Val Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser
20 25 30
Gly Asp Thr Thr Cys Thr Ala Ser Thr Cys Val Lys Val Asn Asp Tyr
35 40 45
Tyr Ser Gln Cys Gln Pro Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser
50 55 60
Ala Pro Gly Pro Ser Ala Cys Pro Leu Val Lys Arg Gln Val Gln Leu
65 70 75 80
Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly Ser Leu Arg Leu
85 90 95
Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly His Gly His Tyr Gly
100 105 110
Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu Arg Glu Phe Val Ala
115 120 125
Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr Lys Asp Ser Val Lys
130 135 140
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Thr Thr Val Tyr Leu
145 150 155 160
Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys Ala
165 170 175
Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu Pro Val Gly Phe Asp
180 185 190
Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser Asp Asn Lys Phe
195 200 205
Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn
210 215 220
Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp
225 230 235 240
Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp
245 250 255
Ala Gln Ala Pro Lys
260

<210> 4

<211> 469

<212> DNA

<213> Artificial

<220>

<223> fusion protein coding sequence

<400> 4

cacctcgtgg ccgagtatcg tggctactgg cggcaccact acgacggcta cccccactgg	60
atcaggcagc gtgacctcga ccagcaagac caccgcgact gctagcaaga ccagcaccga	120
taacaaattc aacaaggaac agcagaacgc gttctacgag atcttgacc tccccaacct	180
caacgaggaa cagaggaacg gtttcattca gtcgctcaag gacgaccct cccagtccgc	240
caacctcttg gcagaagcca agaagttgaa cgaatgcacag gcaccgaaag acaacaagtt	300
caacaagag cagcagaacg ctttctatga gatcttgcac ctcccgaact tgaacgagga	360
gcagcggaaac ggcttcatcc agtccctgaa agatgatccc tcgcagtcgg ccaacctcct	420
cgcagaagcg aagaagctca acgacgcgca ggcacctaag tgactcgag	469

<210> 5

<211> 77

<212> DNA

<213> Artificial

<220>

<223> adaptor sequence

<400> 5

cgcgcttgga aatcacattt gccaaccttg tgcagacgag gccgctcagg gcgagtagag 60
atcggaacga catggtg 77

<210> 6

<211> 77

<212> DNA

<213> Artificial

<220>

<223> adaptor sequence

<400> 6

gatccaccat gtcgttccga tctctactcg ccctgagcgg cctcgtctgc acagggttgg 60
caaattgat ttccaag 77

<210> 7

<211> 639

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 7

Met Ser Phe Arg Ser Leu Leu Ala Leu Ser Gly Leu Val Cys Thr Gly
1 5 10 15
Leu Ala Asn Val Ile Ser Lys Arg Ala Thr Leu Asp Ser Trp Leu Ser
20 25 30
Asn Glu Ala Thr Val Ala Arg Thr Ala Ile Leu Asn Asn Ile Gly Ala
35 40 45
Asp Gly Ala Trp Val Ser Gly Ala Asp Ser Gly Ile Val Val Ala Ser
50 55 60
Pro Ser Thr Asp Asn Pro Asp Phe Tyr Thr Trp Thr Arg Asp Ser Gly
65 70 75 80
Leu Val Leu Lys Thr Leu Val Asp Leu Phe Arg Asn Gly Asp Thr Ser
85 90 95
Leu Leu Ser Thr Ile Glu Asn Tyr Ile Ser Ala Gln Ala Ile Val Gln
100 105 110
Gly Ile Ser Asn Pro Ser Gly Asp Leu Ser Ser Gly Ala Gly Leu Gly
115 120 125
Glu Pro Lys Phe Asn Val Asp Glu Thr Ala Tyr Thr Gly Ser Trp Gly
130 135 140
Arg Pro Gln Arg Asp Gly Pro Ala Leu Arg Ala Thr Ala Met Ile Gly
145 150 155 160
Phe Gly Gln Trp Leu Leu Asp Asn Gly Tyr Thr Ser Thr Ala Thr Asp
165 170 175
Ile Val Trp Pro Leu Val Arg Asn Asp Leu Ser Tyr Val Ala Gln Tyr
180 185 190
Trp Asn Gln Thr Gly Tyr Asp Leu Trp Glu Glu Val Asn Gly Ser Ser
195 200 205
Phe Phe Thr Ile Ala Val Gln His Arg Ala Leu Val Glu Gly Ser Ala
210 215 220

Phe Ala Thr Ala Val Gly Ser Ser Cys Ser Trp Cys Asp Ser Gln Ala
 225 230 235 240
 Pro Glu Ile Leu Cys Tyr Leu Gln Ser Phe Trp Thr Gly Ser Phe Ile
 245 250 255
 Leu Ala Asn Phe Asp Ser Ser Arg Ser Gly Lys Asp Ala Asn Thr Leu
 260 265 270
 Leu Gly Ser Ile His Thr Phe Asp Pro Glu Ala Ala Cys Asp Asp Ser
 275 280 285
 Thr Phe Gln Pro Cys Ser Pro Arg Ala Leu Ala Asn His Lys Glu Val
 290 295 300
 Val Asp Ser Phe Arg Ser Ile Tyr Thr Leu Asn Asp Gly Leu Ser Asp
 305 310 315 320
 Ser Glu Ala Val Ala Val Gly Arg Tyr Pro Glu Asp Thr Tyr Tyr Asn
 325 330 335
 Gly Asn Pro Trp Phe Leu Cys Thr Leu Ala Ala Ala Glu Gln Leu Tyr
 340 345 350
 Asp Ala Leu Tyr Gln Trp Asp Lys Gln Gly Ser Leu Glu Val Thr Asp
 355 360 365
 Val Ser Leu Asp Phe Phe Lys Ala Leu Tyr Ser Asp Ala Ala Thr Gly
 370 375 380
 Thr Tyr Ser Ser Ser Ser Ser Thr Tyr Ser Ser Ile Val Asp Ala Val
 385 390 395 400
 Lys Thr Phe Ala Asp Gly Phe Val Ser Ile Val Glu Thr His Ala Ala
 405 410 415
 Ser Asn Gly Ser Met Ser Glu Gln Tyr Asp Lys Ser Asp Gly Glu Gln
 420 425 430
 Leu Ser Ala Arg Asp Leu Thr Trp Ser Tyr Ala Ala Leu Leu Thr Ala
 435 440 445
 Asn Asn Arg Arg Asn Ser Val Val Pro Ala Ser Trp Gly Glu Thr Ser
 450 455 460
 Ala Ser Ser Val Pro Gly Thr Cys Ala Ala Thr Ser Ala Ile Gly Thr
 465 470 475 480
 Tyr Ser Ser Val Thr Val Thr Ser Trp Pro Ser Ile Val Ala Thr Gly
 485 490 495
 Gly Thr Thr Thr Thr Ala Thr Pro Thr Gly Ser Gly Ser Val Thr Ser
 500 505 510
 Thr Ser Lys Thr Thr Ala Thr Ala Ser Lys Thr Ser Thr Asp Asn Lys
 515 520 525
 Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro
 530 535 540
 Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp
 545 550 555 560
 Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu Asn
 565 570 575
 Asp Ala Gln Ala Pro Lys Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn
 580 585 590
 Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg
 595 600 605
 Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn
 610 615 620
 Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys
 625 630 635

<210> 8

<211> 583

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 8

```

Met Ser Phe Arg Ser Leu Leu Ala Leu Ser Gly Leu Val Cys Thr Gly
1      5      10      15

Leu Ala Asn Val Ile Ser Lys Arg Ala Thr Leu Asp Ser Trp Leu Ser
20     25     30

Asn Glu Ala Thr Val Ala Arg Thr Ala Ile Leu Asn Asn Ile Gly Ala
35     40     45

Asp Gly Ala Trp Val Ser Gly Ala Asp Ser Gly Ile Val Val Ala Ser
50     55     60

Pro Ser Thr Asp Asn Pro Asp Tyr Phe Tyr Thr Trp Thr Arg Asp Ser
65     70     75     80

Gly Leu Val Leu Lys Thr Leu Val Asp Leu Phe Arg Asn Gly Asp Thr
85     90     95

Ser Leu Leu Ser Thr Ile Glu Asn Tyr Ile Ser Ala Gln Ala Ile Val
100    105    110

Gln Gly Ile Ser Asn Pro Ser Gly Asp Leu Ser Ser Gly Ala Gly Leu
115    120    125

Gly Glu Pro Lys Phe Asn Val Asp Glu Thr Ala Tyr Thr Gly Ser Trp
130    135    140

Gly Arg Pro Gln Arg Asp Gly Pro Ala Leu Arg Ala Thr Ala Met Ile
145    150    155    160

Gly Phe Gly Gln Trp Leu Leu Asp Asn Gly Tyr Thr Ser Thr Ala Thr
165    170    175

Asp Ile Val Trp Pro Leu Val Arg Asn Asp Leu Ser Tyr Val Ala Gln
180    185    190

Tyr Trp Asn Gln Thr Gly Tyr Asp Leu Trp Glu Glu Val Asn Gly Ser
195    200    205

Ser Phe Phe Thr Ile Ala Val Gln His Arg Ala Leu Val Glu Gly Ser
210    215    220

Ala Phe Ala Thr Ala Val Gly Ser Ser Cys Ser Trp Cys Asp Ser Gln
225    230    235    240

Ala Pro Glu Ile Leu Cys Tyr Leu Gln Ser Phe Trp Thr Gly Ser Phe
245    250    255

Ile Leu Ala Asn Phe Asp Ser Ser Arg Ser Gly Lys Asp Ala Asn Thr
260    265    270

Leu Leu Gly Ser Ile His Thr Phe Asp Pro Glu Ala Ala Cys Asp Asp
275    280    285

Ser Thr Phe Gln Pro Cys Ser Pro Arg Ala Leu Ala Asn His Lys Glu
290    295    300

Val Val Asp Ser Phe Arg Ser Ile Tyr Thr Leu Asn Asp Gly Leu Ser
305    310    315    320

Asp Ser Glu Ala Val Ala Val Gly Arg Tyr Pro Glu Asp Thr Tyr Tyr
325    330    335

Asn Gly Asn Pro Trp Phe Leu Cys Thr Leu Ala Ala Ala Glu Gln Leu
340    345    350

Tyr Asp Ala Leu Tyr Gln Trp Asp Lys Gln Gly Ser Leu Glu Val Thr
355    360    365

Asp Val Ser Leu Asp Phe Phe Lys Ala Leu Tyr Ser Asp Ala Ala Thr

```

```

370          375          380
Gly Thr Tyr Ser Ser Ser Ser Thr Tyr Ser Ser Ile Val Asp Ala
385          390          395
Val Lys Thr Phe Ala Asp Gly Phe Val Ser Ile Val Glu Thr His Ala
405          410          415
Ala Ser Asn Gly Ser Met Ser Glu Gln Tyr Asp Lys Ser Asp Gly Glu
420          425          430
Gln Leu Ser Ala Arg Asp Leu Thr Trp Ser Tyr Ala Ala Leu Leu Thr
435          440          445
Ala Asn Asn Arg Arg Asn Ser Val Val Pro Ala Ser Trp Gly Glu Thr
450          455          460
Ser Ala Ser Ser Val Pro Gly Thr Cys Ala Ala Thr Ser Ala Ile Gly
465          470          475
Thr Tyr Ser Ser Val Thr Val Thr Ser Trp Pro Ser Ile Val Ala Thr
485          490          495
Gly Gly Thr Thr Thr Thr Ala Thr Pro Thr Gly Ser Gly Ser Val Thr
500          505          510
Ser Thr Ser Lys Thr Thr Ala Thr Ala Ser Lys Thr Ser Thr Asp Asn
515          520          525
Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu
530          535          540
Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys
545          550          555
Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu
565          570          575
Asn Asp Ala Gln Ala Pro Lys
580

<210> 9
<211> 259
<212> PRT
<213> Artificial

<220>
<223> fusion protein

<400> 9
Met Lys Ala Ile Leu Ser Leu Ala Ala Ala Leu Leu Ser Ala Ala Pro
1          5          10          15

```

Ala Phe Ser Thr 20 Ala Val Trp Gly 25 Gln Cys Gly Gly Ile 30 Gly Phe Ser
 Gly Asp Thr 35 Thr Cys Thr Ala 40 Ser Thr Cys Val Lys 45 Val Asn Asp Tyr
 Tyr Ser 50 Gln Cys Gln Pro 55 Gly Ala Ser Ala Pro 60 Thr Ser Thr Ala Ser
 Ala Pro Gly Pro Ser 70 Ala Cys Pro Gly Ser 75 Asp Asn Lys Phe Asn 80 Lys
 Glu Gln Gln Asn 85 Ala Phe Tyr Glu Ile 90 Leu His Leu Pro Asn Leu Asn 95
 Glu Glu Gln Arg 100 Asn Gly Phe Ile 105 Gln Ser Leu Lys Asp 110 Asp Pro Ser
 Gln Ser 115 Ala Asn Leu Leu Ala 120 Glu Ala Lys Lys Leu 125 Asn Asp Ala Gln
 Ala Pro 130 Lys Asp Asn Lys 135 Phe Asn Lys Glu Gln 140 Gln Asn Ala Phe Tyr
 Glu Ile Leu His Leu 150 Pro Asn Leu Asn Glu 155 Glu Gln Arg Asn Gly Phe 160
 Ile Gln Ser Leu 165 Lys Asp Asp Pro Ser 170 Gln Ser Ala Asn Leu Leu Ala 175
 Glu Ala Lys 180 Lys Leu Asn Asp Ala 185 Gln Ala Pro Lys Ser 190 Ser Ser Thr
 Ser Ser 195 Pro Val Asn Gln Pro 200 Thr Ser Thr Ser Thr 205 Thr Ser Thr Ser
 Thr Thr 210 Ser Ser Pro Pro 215 Val Gln Pro Thr Thr 220 Pro Ser Gly Cys Thr
 Ala Glu Arg Trp 225 Ala Gln 230 Cys Gly Gly Asn 235 Gly Trp Ser Gly Cys Thr 240
 Thr Cys Val Ala 245 Gly Ser Thr Cys Thr 250 Lys Ile Asn Asp Trp Tyr His 255
 Gln Cys Leu

<210> 10

<211> 20

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 10

gcaagggatg ccatgcttgg 20

<210> 11

<211> 38

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 11

ctgttgaat ttgtatccg agccagggca cgcggacg 38

<210> 12

<211> 38

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 12
 cgtccgctg cctggctcg gataacaaat tcaacaag 38

<210> 13
 <211> 42
 <212> DNA
 <213> Artificial

<220>

<223> Primer

<400> 13
 ggagagctgg tgctgctga ctaggtgcc tgcgcgtgt tg 42

<210> 14
 <211> 42
 <212> DNA
 <213> Artificial

<220>

<223> Primer

<400> 14
 caacgacgog caggcaccta agtcacgag caccagctct cc 42

<210> 15
 <211> 19
 <212> DNA
 <213> Artificial

<220>

<223> Primer

<400> 15
 catataacca attgccctc 19

<210> 16
 <211> 372
 <212> DNA
 <213> Artificial

<220>

<223> fusion protein coding sequence

<400> 16
 atgaaggcga tcctctctct cgctgccgog ctcttgagcg ccgcgcccgc gttctcgacc 60
 gcagtctggg gccagtgcgg tggcatcggc ttctccggcg acaccacgtg caccgcttcc 120
 acctgcgtga aggtcaacga ctactactcg cagtgccagc ctggcgcgtc cgcacccacg 180
 tcaaccgcct ccgcgcccgg cccgtccgog tgccctggct cgcagtcgag cccgccagtc 240
 cagcctacga ctcccagcgg ctgcactgct gagaggtggg ctgagtcggg cggcaatggc 300
 tggagcggct gcaccacctg cgctcgctggc agcacttgca cgaagattaa tgactggtac 360
 catcagtgcc tg 372

<210> 17
 <211> 124
 <212> PRT
 <213> Artificial

<220>

<223> fusion protein

<400> 17

```

Met Lys Ala Ile Leu Ser Leu Ala Ala Ala Leu Leu Ser Ala Ala Pro
1      5      10      15

Ala Phe Ser Thr Ala Val Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser
20      25      30

Gly Asp Thr Thr Cys Thr Ala Ser Thr Cys Val Lys Val Asn Asp Tyr
35      40      45

Tyr Ser Gln Cys Gln Pro Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser
50      55      60

Ala Pro Gly Pro Ser Ala Cys Pro Gly Ser Gln Ser Ser Pro Pro Val
65      70      75      80

Gln Pro Thr Thr Pro Ser Gly Cys Thr Ala Glu Arg Trp Ala Gln Cys
85      90      95

Gly Gly Asn Gly Trp Ser Gly Cys Thr Thr Cys Val Ala Gly Ser Thr
100     105     110

Cys Thr Lys Ile Asn Asp Trp Tyr His Gln Cys Leu
115     120

```

<210> 18

<211> 294

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 18

Met Lys Ala Ile Leu Ser Leu Ala Ala Ala Leu Leu Ser Ala Ala Pro
1 5 10 15
Ala Phe Ser Thr Ala Val Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser
20 25 30
Gly Asp Thr Thr Cys Thr Ala Ser Thr Cys Val Lys Val Asn Asp Tyr
35 40 45
Tyr Ser Gln Cys Gln Pro Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser
50 55 60
Ala Pro Gly Pro Ser Ala Cys Pro Leu Val Lys Arg Asp Asn Lys Phe
65 70 75 80
Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn
85 90 95
Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp
100 105 110
Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp
115 120 125
Ala Gln Ala Pro Lys Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala
130 135 140
Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn
145 150 155 160
Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu
165 170 175
Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Val
180 185 190
Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser Gly Asp Thr Thr Cys Thr
195 200 205
Ala Ser Thr Cys Val Lys Val Asn Asp Tyr Tyr Ser Gln Cys Gln Pro
210 215 220
Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser Ala Pro Gly Pro Ser Ala
225 230 235 240
Cys Pro Gly Ser Gln Ser Ser Pro Pro Val Gln Pro Thr Thr Pro Ser
245 250 255
Gly Cys Thr Ala Glu Arg Trp Ala Gln Cys Gly Gly Asn Gly Trp Ser
260 265 270
Gly Cys Thr Thr Cys Val Ala Gly Ser Thr Cys Thr Lys Ile Asn Asp
275 280 285
Trp Tyr His Gln Cys Leu
290

<210> 19

<211> 38

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 19

ctgttgaat ttgtatcac gcttcacaa agggcacg 38

<210> 20

<211> 38

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 20
cgtgcccttt ggtgaagcgt gataacaaat tcaacaag 38

<210> 21
<211> 42
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 21
ccgcactggc cccagactgc cttagggtcc tgcgcgtcgt tg 42

<210> 22
<211> 42
<212> DNA
<213> Artificial

<220>
<223> Primer

<400> 22
caacgacggc caggcaccta aggcagtcgt gggccagtcg gg 42

<210> 23
<211> 242
<212> PRT
<213> Artificial

<220>
<223> fusion protein

<400> 23

Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly His
20 25 30
Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu Arg
35 40 45
Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr Lys
50 55 60
Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Thr
65 70 75 80
Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val
85 90 95
Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu Pro
100 105 110
Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120 125
Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu
130 135 140
His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser
145 150 155 160
Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys
165 170 175
Lys Leu Asn Asp Ala Gln Ala Pro Lys Asp Asn Lys Phe Asn Lys Glu
180 185 190
Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu
195 200 205
Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln
210 215 220
Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala
225 230 235 240

Pro Lys

<210> 24

<211> 615

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 24

Ala Thr Leu Asp Ser Trp Leu Ser Asn Glu Ala Thr Val Ala Arg Thr
1 5 10 15

Ala Ile Leu Asn Asn Ile Gly Ala Asp Gly Ala Trp Val Ser Gly Ala
20 25 30

Asp Ser Gly Ile Val Val Ala Ser Pro Ser Thr Asp Asn Pro Asp Phe
35 40 45

Tyr Thr Trp Thr Arg Asp Ser Gly Leu Val Leu Lys Thr Leu Val Asp
50 55 60

Leu Phe Arg Asn Gly Asp Thr Ser Leu Leu Ser Thr Ile Glu Asn Tyr
65 70 75 80

Ile Ser Ala Gln Ala Ile Val Gln Gly Ile Ser Asn Pro Ser Gly Asp
85 90 95

Leu Ser Ser Gly Ala Gly Leu Gly Glu Pro Lys Phe Asn Val Asp Glu
100 105 110

Thr Ala Tyr Thr Gly Ser Trp Gly Arg Pro Gln Arg Asp Gly Pro Ala
115 120 125

Leu Arg Ala Thr Ala Met Ile Gly Phe Gly Gln Trp Leu Leu Asp Asn
130 135 140

Gly Tyr Thr Ser Thr Ala Thr Asp Ile Val Trp Pro Leu Val Arg Asn
145 150 155 160

Asp Leu Ser Tyr Val Ala Gln Tyr Trp Asn Gln Thr Gly Tyr Asp Leu
165 170 175

Trp Glu Glu Val Asn Gly Ser Ser Phe Phe Thr Ile Ala Val Gln His
180 185 190

Arg Ala Leu Val Glu Gly Ser Ala Phe Ala Thr Ala Val Gly Ser Ser
195 200 205

Cys Ser Trp Cys Asp Ser Gln Ala Pro Glu Ile Leu Cys Tyr Leu Gln
210 215 220

Ser Phe Trp Thr Gly Ser Phe Ile Leu Ala Asn Phe Asp Ser Ser Arg
225 230 235 240

Ser Gly Lys Asp Ala Asn Thr Leu Leu Gly Ser Ile His Thr Phe Asp
245 250 255

Pro Glu Ala Ala Cys Asp Asp Ser Thr Phe Gln Pro Cys Ser Pro Arg
260 270

Ala Leu Ala Asn His Lys Glu Val Val Asp Ser Phe Arg Ser Ile Tyr
275 280 285

Thr Leu Asn Asp Gly Leu Ser Asp Ser Glu Ala Val Ala Val Gly Arg
290 295 300

Tyr Pro Glu Asp Thr Tyr Tyr Asn Gly Asn Pro Trp Phe Leu Cys Thr
305 310 315 320

Leu Ala Ala Ala Glu Gln Leu Tyr Asp Ala Leu Tyr Gln Trp Asp Lys
325 330 335

Gln Gly Ser Leu Glu Val Thr Asp Val Ser Leu Asp Phe Phe Lys Ala
340 345 350

Leu Tyr Ser Asp Ala Ala Thr Gly Thr Tyr Ser Ser Ser Ser Thr
355 360 365

Tyr Ser Ser Ile Val Asp Ala Val Lys Thr Phe Ala Asp Gly Phe Val
370 375 380

Ser Ile Val Glu Thr His Ala Ala Ser Asn Gly Ser Met Ser Glu Gln
385 390 395 400

Tyr Asp Lys Ser Asp Gly Glu Gln Leu Ser Ala Arg Asp Leu Thr Trp
405 410 415

Ser Tyr Ala Ala Leu Leu Thr Ala Asn Asn Arg Arg Asn Ser Val Val
420 425 430

Pro Ala Ser Trp Gly Glu Thr Ser Ala Ser Ser Val Pro Gly Thr Cys
435 440 445

Ala Ala Thr Ser Ala Ile Gly Thr Tyr Ser Ser Val Thr Val Thr Ser
450 455 460

Trp Pro Ser Ile Val Ala Thr Gly Gly Thr Thr Thr Thr Ala Thr Pro
465 470 475 480

Thr Gly Ser Gly Ser Val Thr Ser Thr Ser Lys Thr Thr Ala Thr Ala
485 490 495

Ser Lys Thr Ser Thr Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala
500 505 510

Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn
515 520 525

Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu
530 535 540

Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Asp Asn
545 550 555 560

Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu
565 570 575

Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys
580 585 590

Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu
595 600 605

Asn Asp Ala Gln Ala Pro Lys
610 615

<210> 25

<211> 559

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 25

Ala Thr Leu Asp Ser Trp Leu Ser Asn Glu Ala Thr Val Ala Arg Thr
 1 5 10 15
 Ala Ile Leu Asn Asn Ile Gly Ala Asp Gly Ala Trp Val Ser Gly Ala
 20 25 30
 Asp Ser Gly Ile Val Val Ala Ser Pro Ser Thr Asp Asn Pro Asp Tyr
 35 40 45
 Phe Tyr Thr Trp Thr Arg Asp Ser Gly Leu Val Leu Lys Thr Leu Val
 50 55 60
 Asp Leu Phe Arg Asn Gly Asp Thr Ser Leu Leu Ser Thr Ile Glu Asn
 65 70 75 80
 Tyr Ile Ser Ala Gln Ala Ile Val Gln Gly Ile Ser Asn Pro Ser Gly
 85 90 95
 Asp Leu Ser Ser Gly Ala Gly Leu Gly Glu Pro Lys Phe Asn Val Asp
 100 105 110
 Glu Thr Ala Tyr Thr Gly Ser Trp Gly Arg Pro Gln Arg Asp Gly Pro
 115 120 125
 Ala Leu Arg Ala Thr Ala Met Ile Gly Phe Gly Gln Trp Leu Leu Asp
 130 135 140
 Asn Gly Tyr Thr Ser Thr Ala Thr Asp Ile Val Trp Pro Leu Val Arg
 145 150 155 160
 Asn Asp Leu Ser Tyr Val Ala Gln Tyr Trp Asn Gln Thr Gly Tyr Asp
 165 170 175
 Leu Trp Glu Glu Val Asn Gly Ser Ser Phe Phe Thr Ile Ala Val Gln
 180 185 190
 His Arg Ala Leu Val Glu Gly Ser Ala Phe Ala Thr Ala Val Gly Ser
 195 200 205
 Ser Cys Ser Trp Cys Asp Ser Gln Ala Pro Glu Ile Leu Cys Tyr Leu
 210 215 220
 Gln Ser Phe Trp Thr Gly Ser Phe Ile Leu Ala Asn Phe Asp Ser Ser
 225 230 235 240
 Arg Ser Gly Lys Asp Ala Asn Thr Leu Leu Gly Ser Ile His Thr Phe
 245 250 255
 Asp Pro Glu Ala Ala Cys Asp Asp Ser Thr Phe Gln Pro Cys Ser Pro
 260 265 270
 Arg Ala Leu Ala Asn His Lys Glu Val Val Asp Ser Phe Arg Ser Ile
 275 280 285
 Tyr Thr Leu Asn Asp Gly Leu Ser Asp Ser Glu Ala Val Ala Val Gly
 290 295 300
 Arg Tyr Pro Glu Asp Thr Tyr Tyr Asn Gly Asn Pro Trp Phe Leu Cys
 305 310 315 320
 Thr Leu Ala Ala Ala Glu Gln Leu Tyr Asp Ala Leu Tyr Gln Trp Asp
 325 330 335
 Lys Gln Gly Ser Leu Glu Val Thr Asp Val Ser Leu Asp Phe Lys
 340 345 350
 Ala Leu Tyr Ser Asp Ala Ala Thr Gly Thr Tyr Ser Ser Ser Ser
 355 360 365
 Thr Tyr Ser Ser Ile Val Asp Ala Val Lys Thr Phe Ala Asp Gly Phe
 370 375 380
 Val Ser Ile Val Glu Thr His Ala Ala Ser Asn Gly Ser Met Ser Glu
 385 390 395 400
 Gln Tyr Asp Lys Ser Asp Gly Glu Gln Leu Ser Ala Arg Asp Leu Thr
 405 410 415

Trp Ser Tyr Ala Ala Leu Leu Thr Ala Asn Asn Arg Arg Asn Ser Val
420 425 430
Val Pro Ala Ser Trp Gly Glu Thr Ser Ala Ser Ser Val Pro Gly Thr
435 440 445
Cys Ala Ala Thr Ser Ala Ile Gly Thr Tyr Ser Ser Val Thr Val Thr
450 455 460
Ser Trp Pro Ser Ile Val Ala Thr Gly Gly Thr Thr Thr Thr Ala Thr
465 470 475 480
Pro Thr Gly Ser Gly Ser Val Thr Ser Thr Ser Lys Thr Thr Ala Thr
485 490 495
Ala Ser Lys Thr Ser Thr Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn
500 505 510
Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg
515 520 525
Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn
530 535 540
Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys
545 550 555

<210> 26

<211> 239

<212> PRT

<213> Artificial

<220>

<223> fusion protein

<400> 26

Ala Val Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser Gly Asp Thr Thr
1 5 10 15
Cys Thr Ala Ser Thr Cys Val Lys Val Asn Asp Tyr Tyr Ser Gln Cys
20 25 30
Gln Pro Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser Ala Pro Gly Pro
35 40 45
Ser Ala Cys Pro Gly Ser Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn
50 55 60
Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg
65 70 75 80
Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn
85 90 95

Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Asp
 100 105 110
 Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His
 115 120 125
 Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
 130 135 140
 Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys
 145 150 155 160
 Leu Asn Asp Ala Gln Ala Pro Lys Ser Ser Ser Thr Ser Ser Pro Val
 165 170 175
 Asn Gln Pro Thr Ser Thr Ser Thr Thr Ser Thr Ser Thr Thr Ser Ser
 180 185 190
 Pro Pro Val Gln Pro Thr Thr Pro Ser Gly Cys Thr Ala Glu Arg Trp
 195 200 205
 Ala Gln Cys Gly Gly Asn Gly Trp Ser Gly Cys Thr Thr Cys Val Ala
 210 215 220
 Gly Ser Thr Cys Thr Lys Ile Asn Asp Trp Tyr His Gln Cys Leu
 225 230 235
 <210> 27
 <211> 218
 <212> PRT
 <213> Artificial
 <220>
 <223> fusion protein
 <400> 27
 Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu
 1 5 10 15
 His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser
 20 25 30
 Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys
 35 40 45
 Lys Leu Asn Asp Ala Gln Ala Pro Lys Asp Asn Lys Phe Asn Lys Glu
 50 55 60
 Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu
 65 70 75 80
 Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln
 85 90 95
 Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala
 100 105 110
 Pro Lys Ala Val Trp Gly Gln Cys Gly Gly Ile Gly Phe Ser Gly Asp
 115 120 125
 Thr Thr Cys Thr Ala Ser Thr Cys Val Lys Val Asn Asp Tyr Tyr Ser
 130 135 140
 Gln Cys Gln Pro Gly Ala Ser Ala Pro Thr Ser Thr Ala Ser Ala Pro
 145 150 155 160
 Gly Pro Ser Ala Cys Pro Gly Ser Gln Ser Ser Pro Pro Val Gln Pro
 165 170 175
 Thr Thr Pro Ser Gly Cys Thr Ala Glu Arg Trp Ala Gln Cys Gly Gly
 180 185 190
 Asn Gly Trp Ser Gly Cys Thr Thr Cys Val Ala Gly Ser Thr Cys Thr
 195 200 205
 Lys Ile Asn Asp Trp Tyr His Gln Cys Leu
 210 215

<210> 28
 <211> 185
 <212> PRT
 <213> Artificial

<220>
 <223> fusion protein

<400> 28
 Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Ala Thr Ser Gly His
 20 25 30
 Gly His Tyr Gly Met Gly Trp Phe Arg Gln Val Pro Gly Lys Glu Arg
 35 40 45
 Glu Phe Val Ala Ala Ile Arg Trp Ser Gly Lys Glu Thr Trp Tyr Lys
 50 55 60
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Thr
 65 70 75 80
 Thr Val Tyr Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val
 85 90 95
 Tyr Tyr Cys Ala Ala Arg Pro Val Arg Val Asp Asp Ile Ser Leu Pro
 100 105 110
 Val Gly Phe Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
 115 120 125
 Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu
 130 135 140
 His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser
 145 150 155 160
 Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys
 165 170 175
 Lys Leu Asn Asp Ala Gln Ala Pro Lys
 180 185

<210> 29
 <211> 390
 <212> DNA
 <213> Artificial

<220>
 <223> z:z domain coding sequence

<400> 29
 tcattctgca gcagcggcgg ataacaaatt taacaaagaa cagcagaacg cgttttatga 60
 aatcctgcat ctgccgaacc tgaacgaaga acagcgcaac gcgtttatcc agagcctgaa 120
 agatgatccg agccagagcg cgaacctgct ggcggaagcg aaaaaactga acgatgcgca 180
 ggcgccgaaa gtcgacaaca aatttaataa ggaacaacaa aacgccttct acgaaattct 240
 ccatttacc aatcttaatg aagagcagcg caacgctttt attcaatcgc ttaaggacga 300
 tcccagccaa tcggcaaatc ttctggctga agctaaaaag cttaatgacg cgcaagctcc 360
 taaagtgtag tcgctagcgg ccgcttagtt 390

<210> 30
 <211> 626
 <212> DNA
 <213> Artificial

<220>

<223> streptavidin coding sequence

<400> 30

aagcttaatg acgcgcaagc tcctaaagtg gacccgtcaa aagattccaa agcccaggta	60
agtcagctg aagcaggcat aacgggaaca tggtaacaatc aactgggaag cacgtttatc	120
gtaacagcag gagcggacgg agcgttgaca ggcacatacg aatctgccgt tggaaacgca	180
gaatcgctgt acacacttac aggtagatac gattcagcac cggctaccga tggaaagcga	240
actgcattgg gatggagagt ggcttggaag aacaactata gaaatgcgca tagcgccacg	300
acatggtctg gccaatacgt cggaggagca gaagctagaa tcaatacaca atggacatta	360
acgagcggga ctacagaagc aaatgcctgg aaatcgacgt taagaggta tgataccttt	420
accaaagtaa agccgtccgc agcatcaatt gatgcagcga agaaagcagg agtgaataac	480
ggtaatcctt tggatgcagt gcaacaatag tcgctagcgg ccgcgtcgac tagaagagca	540
gagaggacgg atttctgaa ggaaatccgt ttttttattt tgcccgtctt ataaatttcg	600
ttgagatctt gatcatcgat aagctt	626

<210> 31

<211> 4084

<212> DNA

<213> Artificial

<220>

<223> plasmid pSJ6208

<400> 31

tcgcgcgttt cgggtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggtca	60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg	120
ttggcgggtg tcggggctgg cttactatg cggcatcaga gcagattgta ctgagagtgc	180
accatattcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc	240
attcgccatt caggctgcgc aactgttggg aaggcgatc ggtcggggc tcttcgctat	300
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagtgggta acgccagggt	360
tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt cagatctaaa gataatatct	420
ttgaattgta acsccttca aaagtaagaa ctacaaaaa agaatacgtt atatagaaat	480
atgtttgaac cttcttcaga ttacaaatat attcggacgg actctacctc aaatgcttat	540
ctaactatag aatgacatac aagcacaaac ttgaaaattt gaaaataata ctaccaatga	600
acttgttcat gtgaattatc gctgtattta attttctca ttcaatatat aatatgccaa	660
tacattgtta caagtagaaa ttaagacacc cttgatagcc ttactatacc taacatgatg	720
tagtattaaa tgaatatgta aatatattta tgataagaag cgacttattt ataataatta	780
catatttttc tattggaatg attaagattc caatagaata gtgtataaat tatttatctt	840
gaaaggaggg atgcctaaaa acgaagaaca ttaaaaacat atatttgcac cgtctaattg	900
atztatgaaa aatcatttta tcagtttgaa aattatgtat tatggccaca ttgaaagggg	960
aggagaatca tgaaacaaca aaaacggctt tacgcccgat tgctgacgct gttatttgcg	1020
ctcatcttct tgctgcctca ttctgcagca gcggcgatcg tcggcggtag gaaggccagt	1080
acctcgacct acccgcttct ggtcttcctg actgacagca ccggtttcca gttctgcggt	1140
ggcacgctgg tcaagccgaa caaggtggtc acggcggcgc actgcaccgt cggtgagtcc	1200
gcggccaaca tccgcgttgt cgccggtcgc gacgacaagc agagcaccgc cggcactgtc	1260
tcgaaggta gcaagatctg gatccaccgg agttaccagg acgccaccaa gggcagcgac	1320
gtgtcgggtc tgaccctgtc gaccagcctg acccagttca cgccgttgcc gctgggtgcc	1380
accactgaca ccgcgctgta caaggagggc accgccgca ccacccctcg ctgggggtgac	1440
accaccgagg gcgggtcggc ctctcggtag ctgctcaagg cgacagtgcc gctgaccagc	1500

```

gacgccacct gcaagaaggc gtacggcgag tacagttcca ccgcatgggt ctgtgccgga 1560
taccgcgagg gtggcacgga cacctgccag ggcgactccg gcggtccgct cgtcgccggc 1620
aacaagctga tcggcatcac ctctgtgggc cagggtctgc ccgaggccgg ttatccaggc 1680
gtctacaccc gggctgccac ctacagttcg ctgatcacc agcagctcgg ctagtgccta 1740
gcggccgcgt cgactagaag agcagagagg acggatttcc tgaaggaaat ccgttttttt 1800
attttgccg tcttataaat ttcgttgaga tcttgatcat cgataagctt ggcgtaatca 1860
tggtcatagc tgtttctctgt gtgaaattgt tatccgctca caattccaca caacatacga 1920
gccggaagca taaagtgtaa agcctggggt gcctaagag tgagctaact cacattaatt 1980
gcgttgcgct cactgccgcg tttccagtcg ggaaacctgt cgtgccagct gcattaatga 2040
atcgccaac gcgcggggag aggcggtttg cgtattgggc gctcttccgc ttcctcgctc 2100
actgactcgc tgcgctcgggt cgttcggctg cggcgagcgg tatcagctca ctcaaggcgg 2160
gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacaatgt agcaaaaggc 2220
cagcaaaaagg ccaggaacgg taaaaaggcc gcgttgctgg cgtttttcca taggtccgc 2280
ccccctgacg agcatcacaa aaatcgacgc tcaagtcaga ggtggcgaaa cccgacagga 2340
ctataaagat accaggcgtt tccccctgga agctccctcg tgcgctctcc tgttccgacc 2400
ctgccgtta ccggatacct gtccgccttt ctcccttcgg gaagcgtggc gctttctcaa 2460
tgctcacgct gtaggtatct cagttcgggt taggtcgttc gctccaagct gggctgtgtg 2520
cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg tcttgagtcc 2580
aaccgcgtaa gacacgactt atcgccactg gcagcagcca ctggtaacag gattagcaga 2640
gcgaggtatg taggcggtgc tacagagttc ttgaagtggg gccctaacta cggctacact 2700
agaaggacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg aaaaagagtt 2760
ggtagctctt gatccggcaa acaaacaccg gctggtagcg gtgggttttt tgtttgcaag 2820
cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt ttctacgggg 2880
tctgacgctc agtggaacga aaactcacgt taagggattt tggatcatgag attatcaaaa 2940
aggatcttca cctagatcct tttaaattaa aaatgaagtt ttaaatcaat ctaaagtata 3000
tatgagtaaa cttggtctga cagttaccaa tgcttaataca gtgaggcacc tatctcagcg 3060
atctgtctat ttcgttcatc catagttgcc tgactccccg tctgttagat aactacgata 3120
cgggagggct taccatctgg cccagtgct gcaatgatac cgcgagaccc acgctaccg 3180
gctccagatt tatcagcaat aaaccagcca gccggaaggg ccgagcgagc aagtggtcct 3240
gcaactttat ccgcctccat ccagtctatt aattgttgcc gggagctag agtaagtagt 3300
tcgccagtta atagtttgcg caacgttgtt gccattgcta caggcatcgt ggtgtcacgc 3360
tcgtcgtttg gtatggcttc attcagctcc ggttcccaac gatcaaggcg agttacatga 3420
tccccatgt tgtgcaaaaa agcgggttagc tccttcggtc ctccgatcgt tgtcagaagt 3480
aagttggccg cagtggtatc actcatggtt atggcagcac tgcataattc tcttactgtc 3540
atgccatccg taagatgctt tctgtgact ggtgagtact caaccaagtc attctgagaa 3600

tagtgtatgc ggcgaccgag ttgctcttgc ccggcgtcaa tacgggataa taccgcgcca 3660
catagcagaa ctttaaaagt gctcatcatt ggaaaacgtt cttcggggcg aaaactctca 3720
aggatcttac cgctgttgag atccagttcg atgtaacca ctctgcacc caactgatct 3780
tcagcatctt ttactttcac cagcgtttct gggtagagca aaacaggaag gcaaaatgcc 3840
gcaaaaaagg gaataaggc gacacggaaa tgttgaatac tcatactctt cctttttcaa 3900
tattattgaa gcatttatca gggttattgt ctcatgagcg gatacatatt tgaatgtatt 3960
tagaaaaata aacaaatagg ggttcgcgc acatttcccc gaaaagtgcc acctgacgtc 4020
taagaaacca ttattatcat gacattaacc tataaaaata ggcgtatcac gagcccttt 4080
cgtc 4084

```

<210> 32

<211> 6691

<212> DNA

<213> Artificial

<220>

<223> Plasmid sequence pSJ6321

<400> 32

gaattcagat ctaaagataa tatctttgaa ttgtaacsc cctcaaaagt aagaactaca	60
aaaaaagaat acgttatata gaaatatgtt tgaaccttct tcagattaca aatatattcg	120
gacggactct acctcaaatg cttatctaac tatagaatga catacaagca caaccttgaa	180
aatttgaaaa tataactacc aatgaacttg ttcattgtgaa ttatcgctgt atttaatttt	240
ctcaattcaa tatataatat gccaatacat tgttacaagt agaaattaag acacccttga	300
tagccttact atacctaaaca tgatgtagta ttaaatgaat atgtaaatat atttatgata	360
agaagcgact tatttataat cattacatat ttttctattg gaatgattaa gattccaata	420
gaatagtgtg taaattatgt atcttgaaag gagggatgcc taaaaacgaa gaacattaaa	480
aacatatatt tgcaccgtct aatggattta tgaaaaatca ttttatcagt ttgaaaatta	540
tgtattatgg ccacattgaa aggggaggag aatcatgaaa caacaaaaac ggctttacgc	600
ccgattgctg acgctgttat ttgcgctcat cttcttgctg cctcattctg cagcagcggc	660
ggccaccgga gcgctcccc agtcacccac cccggaggcc gacgcgtct ccatgcagga	720
ggcgctccag cgcgacctg acctgacctc cgcggaggcc gaggagctgc tggccgccca	780
ggacaccgcc ttcgaggctg acgaggccgc ggccgaggcc gccggggacg cctacggcgg	840
ctccgtcttc gacaccgaga gcctggaact gaccgtcctg gtcaccgatg ccgccgcggt	900
cgaggccgtg gaggccaccg gcgccgggac cgagctggtc tcctacggca tcgacggtct	960
cgacgagatc gtccaggagc tcaacgccgc cgacgccgtt cccggtgtgg tcggctggtgta	1020
cccgacgtg gcgggtgaca ccgtcgtcct ggaggtcctg gaggttccg gagccgacgt	1080
cagcggcctg ctcgaggacg ccggcgtgga cgcctcggcc gtcgaggatga ccacgagcga	1140
ccagcccag ctctacgccg acatcatcgg tggcttgccc tacaccatgg gcggccgctg	1200

ttcggtcggc ttcgcgcca ccaacgccgc cggtcagccc gggttcgtca cgcgcggtca	1260
ctgcgccgc gtgggacccc aggtgacct cggcaacggc aggggcgtct tcgagcagtc	1320
cgtcttcccc ggcaacgacg cggccttcgt ccgcggtacg tccaacttca cgtgaccaa	1380
cctggtcagc cgtacaaca cggcgggta cgcacggtc gccggtcaca accaggcccc	1440
catcggtccc tccgtctgcc gctccggctc caccaccggt tggcactcgc gcaccatcca	1500
ggcccgggc cagtcggtga gctacccga gggcaccgtc accaacatga cccggaccac	1560
cgtgtgcgcc gagcccggcg actccggcgg ctctacatc tccggcacc aggcccagg	1620
cgtgacctcc ggcggtccg gcaactgccg caccggcggg accaccttct accaggaggt	1680
cacccccatg gtgaactcct gggcggtccg tctccggacc tgaacgcgtg ctacggccg	1740
cgtcgactag aagagcagag aggacggatt tcctgaagga aatccgtttt tttattttgc	1800
ccgtcttata aatttcgttg agatctacgc gtccatgggc tagcgcggcc gcgtcgacag	1860
gcctctttga ttacatttta taattaattt taacaaagtg tcatcagccc tcaggaagga	1920
cttgcgaca gtttgaatcg cataggtgaag cgggggatga aatggcaacg ttatctgatg	1980
tagcaaagaa agcaaatgtg tcgaaaatga cggtatcgcg ggtgatcaat catctgaga	2040
ctgtgacgga tgaattgaaa aagcttggtc attccgcaat gaaggagctc aattatatac	2100
cgaactatgc agcaagagcg ctctttcaaa acagaacaca ggtcgtcaag ctgctcatac	2160
tggaagaagt ggatacaaca gaaccttatt atatgaatct gttaacggga atcagccgcg	2220
agctggaccg tcactattat gctttgcagc ttgtcacaag gaaatctctc aatatcggcc	2280
agtgcgacgg cattattgcy acggggttga gaaaagccga ttttgaaggg ctcatcaagg	2340
tttttgaaaa gcgtgtcgtt gtattcggga cgtcgattca caaaaatagg cacacgaaaa	2400
acaagtaagg gatgcagttt atgcacctt taacttactt attaaataat ttatagctat	2460
tgaaaagaga taagaattgt tcaaagctaa tattgtttaa atcgcaatt cctgcatgtt	2520
ttaaggaatt gttaaatga tttttgttaa atattttctt gtattctttg ttaacccatt	2580
tcataacgaa ataattatc tttgtttat ctttgtgtga tattcttgat tttttctac	2640
ttaatctgat aagtgagcta ttcactttag gtttaggatg aaaatatctt ctggaacca	2700
tacttaatat agaaatatca acttctgcca ttaaaagtaa tgccaatgag cgttttgtat	2760
ttaataatct ttagcaaac ccgtattcca cgattaaata aatctcatta gctatactat	2820
caaaaacaat tttgcgtatt atatccgtac ttatgttata aggtatatta ccatatattt	2880
tataggattg gtttttagga aatttaaaact gcaatatatc cttgtttaaa acttggaagt	2940
tatcgtgatc aacaagttta tttctgtag ttttgcataa tttatggtct atttcaatgg	3000
cagttacgaa attacacctc ttactaatt caagggtaaa atggcccttt cctgagccga	3060
ttcacaagat attatcatgt tcatttaatc ttatatattg cattatttta tctatattat	3120
gttttgaggt aataaagttt tgactgtgtt ttatatattt ctggttcatt ataacctct	3180
ttaatttggg tatatgaatt ttgcttatta acgattcatt ataaccactt atttttgtt	3240
tggttgataa tgaactgtgc tgattacaaa aatactaaaa atgcccatat tttttctcc	3300

ttataaaatt agtataatta tagcacgagc tctgataaat atgaacatga tgagtgatcg	3360
ttaaatttat actgcaatcg gatgcgatta ttgaataaaa gatatgagag atttatctaa	3420
ttctttttt cttgtaaaaa aagaaagttc ttaaaggttt tatagttttg gtcgtagagc	3480
acacggttta acgacttaat tacgaagtaa ataagtctag tgtgttagac ttatgaaat	3540
ctatatacgt ttatatatat ttattatccg gaggtgtagc atgtctcatt caattttgag	3600
ggttgccaga gttaaaggat caagtaatac aaacgggata caaagacata atcaaagaga	3660
gaataaaaac tataataata aagacataaa tcatgaggaa acatataaaa attatgattt	3720
gattaacgca caaaatataa agtataaaga taaaattgat gaaacgattg atgagaatta	3780
ttcagggaaa cgtaaaattc ggtcagatgc aattcgacat gtggacggac tggttacaag	3840
tgataaagat ttcttttgat atttaagcgg agaagaata gaacgatttt ttaaagatag	3900
cttgagttt ctgaaaaatg aatacggtaa ggaatatatg ctgtatgcga ctgtccatct	3960
ggatgaaaga gtccccacata tgcactttgg ttttgtccct ttaacagagg acgggagatt	4020
gtctgcaaaa gaacagttag gcaacaagaa agactttact caattacaag atagatttaa	4080
tgagtatgtg aatgagaag gttatgaact tgaagaggc acgtccaaag aggttacaga	4140
acgagaacat aaagcgatg atcagtacaa gaaagatact gtatttcata aacaggaaact	4200
gcaagaagtt aaggatgagt tacagaaggc aaataagcag ttacagagtg gaatagagca	4260
tatgaggctc acgaaaccct ttgattatga aaatgagcgt acaggtttgt tctctggagc	4320
tgaaagagact ggtagaaga tattaactgc tgatgaattt gaacgcctcg aagaaacaat	4380
ctctctgca gaacggattg ttgatgatta cgaaaatatt aagagcacag actattacac	4440
agaaaatcaa gaattaaaa aacgtagaga gagtttgaaa gaagtagtga atacatggaa	4500
agaggggtat cacgaaaaa gtaagagggt taataaatta aagcgagaga atgatatgtt	4560
gaatgagcag ttgaatgtat cagagaaatt tcaagctagt acagtgactt tatactgtgc	4620
tcgagggcg aatttccctg ggtttgagaa agggtttaat aggcctaaag agaaattcct	4680
taatgattcc aaatttgagc gtgtgggaca gtttatggat gttgtacagg ataatgtcca	4740
gaaggtcgat agaaagcgtg agaaacagcg tacagacgat ttagagatgt agaggtactt	4800
ttatgccgag aaaacttttt gcgtgtgaca gtccctaaaa tatacttaga gcgtaagcga	4860
aagtagtagc gacagctatt aactttcggg ttcaaagctc taggattttt aatggacgca	4920
gcgcacaca cgcaaaaagg aaattggaat aaatgcgaaa ttgagatgt taattaaaga	4980
cctttttgag gtcttttttt cttagatttt tggggttatt taggggagaa aacatagggg	5040
ggtactacga cctccccctt aggtgtccat tgtccattgt ccaaacaaat aaataaatat	5100
tgggttttta atgttaaaag gttgtttttt atgttaaagt gaaaaaaca gatgttggga	5160
ggtacagtga tgggtgtaga tagaaaagaa gagaaaaaag ttgctgttac ttttaagactt	5220
acaacagaag aaatgagat attaaataga atcaagaaa aatataatat tagcaaatca	5280
gatgcaaccg gtatttctaataaaaaatat gcaaaggagg aatacgggtgc attttaaaca	5340

```

aaaaagata gacagcactg gcatgctgcc tatctatgac taaattttgt taagtgtatt 5400
agcaccgtta ttatatcatg agcgaaaatg taataaaaga aactgaaaac aagaaaaatt 5460
caagaggacg taattggaca tttgttttat atccagaatc agcaaaagcc gagtggttag 5520
agtattttaa agagttacac attcaatttg tagtgtctcc attacatgat agggatactg 5580
atacagaagg taggatgaaa aaagagcatt atcatattct agtgatgtat gagggtaata 5640
aatcttatga acagataaaa ataattacag aagaattgaa tgcgactatt ccgcagattg 5700
caggaagtgt gaaaggctct gtgagatata tgcttcacat ggacgatcct aataaattta 5760
aatatcaaaa agaagatatg atagtttatg gcggtgtaga tgttgatgaa ttattaaaga 5820
aaacaacaac agatagatat aaattaatta aagaaatgat tgagtttatt gatgaacaag 5880
gaatcgtaga atttaagagt ttaatggatt atgcaatgaa gtttaaatgt gatgattggt 5940
tcccgccttt atgtgataac tcggcgtatg ttattcaaga atataataaa tcaaatcggg 6000
ataaatctga ccgatagatt ttgaatttag gtgtcacaag acactctttt ttcgcaccag 6060
cgaaaactgg tttaagccga ctgcgcaaaa gacataatcg actctagagg atccccgggt 6120
accgagctct gccttttagt ccagctgatt tcactttttg cattctacaa actgcataac 6180
tcatatgtaa atcgctcctt tttaggtggc acaaatgtga ggcattttcg ctctttccgg 6240
caaccacttc caagtaaagt ataacacact atactttata ttcataaagt gtgtgctctg 6300
cgaggctgtc ggcagtgccc accaaaacca taaaaccttt aagacctttc ttttttttac 6360
gagaaaaaag aaacaaaaaa acctgccctc tgccacctca gcaaaggggg gttttgctct 6420
cgtgtcgttt taaaaatcag caagggacag gtagtatttt ttgagaagat cactcaaaaa 6480
atctccacct ttaaaccttt gccaattttt atttgtccg ttttgtctag cttaaccgaa 6540
gccagactca gcaagaataa aatttttatt gtctttcggg ttctagtgt aacggacaaa 6600
accactcaaa ataaaaaaga tacaagagag gtctctcgta tcttttattc agcaatcgcg 6660
cccgattgct gaacagatta ataatgagct c 6691

```

<210> 33

<211> 3757

<212> DNA

<213> Artificial

<220>

<223> Plasmid sequence pMOL2743

<400> 33

```

tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc 240
attcgccatt caggctgcgc aactgttggg aagggcgatc ggtgcgggcc tcttcgctat 300
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggta acgccagggt 360
tttccagatc acgacgttgt aaaacgacgy ccagtgaatt cagatctaaa gataatatct 420

```

ttgaattgta	acscctctca	aaagtaagaa	ctacaaaaa	agaatacgtt	atatagaaat	480
atgtttgaac	cttcttcaga	ttacaaatat	attcggacgg	actctacctc	aaatgcttat	540
ctaactatag	aatgacatac	aagcaccaacc	ttgaaaattt	gaaaatataa	ctaccaatga	600
acttggtcat	gtgaattatc	gctgtattta	atcttctcaa	ttcaatatat	aatatgcaa	660
tacattgtta	caagttagaa	ttaagacacc	cttgatagcc	ttactatacc	taacatgatg	720
tagtattaaa	tgaatatgta	aatatattta	tgataagaag	cgacttattt	ataatcatta	780
catatttttc	tattggaatg	attaagattc	caatagaata	gtgtataaat	tatttatctt	840
gaaaggagg	atgcctaaaa	acgaagaaca	ttaaaaacat	atatttgac	cgtcctaatg	900
atztatgaaa	aatcatttta	tcagtttgaa	aattatgtat	tatggccaca	ttgaaaggg	960
aggagaatca	tgaacaaca	aaaacggctt	tacgccgat	tgctgacgct	gttatttgcg	1020
ctcatcttct	tgctgcctca	ttctgcagca	gcggcggata	acaaatttaa	caaagaacag	1080
cagaacgcgt	tttatgaaat	cctgcattcg	ccgaacctga	acgaagaaca	gcgcaacgcg	1140
tttatccaga	gcctgaaaga	tgatccgagc	cagagcgcga	acctgctggc	ggaagcgaaa	1200
aaactgaacg	atgcgcaggc	gccgaaagtc	gacaacaaat	ttaataagga	acaacaaaac	1260
gccttctacg	aaattctcca	tctacccaat	cttaatgaag	agcagcgcaa	cgctttttatt	1320
caatcgctta	aggacgatcc	cagccaatcg	gcaaatcttc	tggtgaagc	taaaaagctt	1380
aatgacgcgc	aagctcctaa	agtgtagtcg	ctagcggccg	cgctgactag	aagagcagag	1440
aggacggatt	tcctgaagga	aatccgtttt	tttattttgc	ccgtcttata	aatctcgttg	1500
agatcttgat	catcgataag	cttgccgtaa	tcattggtcat	agctgtttcc	tgtgtgaaat	1560
tgttatccgc	tcacaattcc	acacaacata	cgagccggaa	gcataaagt	taaagcctgg	1620
ggtgccta	gagtgagcta	actcacatta	attgcgttgc	gctcactgcc	cgctttccag	1680
tcgggaaacc	tgctgtgcca	gctgcattaa	tgaatcgccc	aacgcgcggg	gagagcggt	1740
ttgcgtattg	ggcgtctctc	cgcttcctcg	ctcactgact	cgctgcgctc	ggtcgttcgg	1800
ctgcggcgag	cggtatcagc	tactcaaaag	gcggtaatac	ggttatccac	agaatcaggg	1860
gataacgcag	gaaagaacat	gtgagcaaaa	ggccagcaaa	aggccaggaa	ccgtaaaaag	1920
gccgcgttgc	tggcgttttt	ccataggctc	cgccccctcg	acgagcatca	caaaaatcga	1980
cgctcaagtc	agaggtggcg	aaacccgaca	ggactataaa	gataccaggc	gtttccccct	2040
ggaagctccc	tcgtgcgctc	tctgttccg	accctgccgc	ttaccggata	cctgtccgcc	2100
tttctccctt	cgggaagcgt	ggcgctttct	caatgctcac	gctgtaggta	tctcagttcg	2160
gtgtaggctg	ttcgtccaa	gctgggctgt	gtgcacgaac	cccccgttca	gcccgaaccgc	2220
tgcgcttat	ccggtaaacta	tcgtcttgag	tccaacccgg	taagacacga	cttatcgcca	2280
ctggcagcag	ccactggtaa	caggattagc	agagcgaggt	atgtaggcgg	tgctacagag	2340
ttcttgaagt	ggtggcctaa	ctacggctac	actagaagga	cagtatttgg	tatctgcgct	2400
ctgctgaagc	cagttacctt	cggaaaaaga	gttggtagct	cttgatccgg	caaacaaacc	2460

```

accgctggta gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag aaaaaaagga 2520
tctcaagaag atcctttgat cttttctacg gggctctgacg ctacgtggaa cgaaaactca 2580
cgtaaaggga ttttggtcat gagattatca aaaaggatct tcacctagat ctttttaaat 2640
taaaaatgaa gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac 2700
caatgcttaa tcagtgaggc acctatctca gcgatctgtc tatttcgttc atccatagtt 2760
gcctgactcc ccgtcgtgta gataactacg atacgggagg gcttaccatc tggccccagt 2820
gctgcaatga taccgcgaga cccacgctca ccggctccag atttatcagc aataaaccag 2880
ccagccggaa gggccgagcg cagaagtggc cctgcaactt tatccgcctc catccagtct 2940
attaattggt gccgggaagc tagagtaagt agttcccgag ttaatagttt gcgcaacggt 3000
gttgccattg ctacaggcat cgtggtgtca cgctcgtcgt ttggtatggc ttcattcagc 3060
tcgggttccc aacgatcaag gcgagttaca tgatcccca tgttgtgcaa aaaagcgggt 3120
agtccttcg gtccctcgat cgttgtcaga agtaagtgg ccgcagtggt atcactcatg 3180
gttatggcag cactgcataa ttctcttact gtcattgccat ccgtaagatg cttttctgtg 3240
actggtgagt actcaaccaa gtcattctga gaatagtgtg tgcggcgacc gagttgctct 3300
tgcccgcggt caatacggga taataccgcg ccacatagca gaactttaaa agtgctcatc 3360
attggaaaac gttcttcggg gcgaaaactc tcaaggatct taccgctggt gagatccagt 3420
tcgatgtaac ccactcgtgc acccaactga tcttcagcat cttttacttt caccagcggt 3480
tctgggtgag caaaaacagg aaggcaaaat gccgcaaaa agggataaag ggcgacacgg 3540
aaatgttgaa tactcatact cttccttttt caatattatt gaagcattta tcagggttat 3600
tgtctcatga gcggatacat atttgaatgt atttagaaaa ataaacaaat aggggttccg 3660
cgcacatttc cccgaaaagt gccacctgac gtctaagaaa ccattattat catgacatta 3720
acctataaaa ataggcgtat cagaggccc tttcgtc 3757

```

<210> 34

<211> 4234

<212> DNA

<213> Artificial

<220>

<223> plasmid sequence pMOL2744

<400> 34

```

tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctccc gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagccc tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgct gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc 240
attcgccatt caggctgcgc aactgttggg aaggcgatc ggtgcgggcc tcttcgctat 300
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggtt acgccagggt 360
tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt cagatctaaa gataatatct 420
ttgaattgta acscccctca aaagtaagaa ctacaaaaa agaatacgtt atatagaaat 480

```

atgtttgaac	cttcttcaga	ttacaaatat	attcggacgg	actctacctc	aaatgcttat	540
ctaactatag	aatgacatac	agcacacaacc	ttgaaaattt	gaaaatataa	ctaccaatga	600
acttgttcat	gtgaattatc	gctgtattta	atcttctcaa	ttcaatatat	aatatgcaa	660
tacattgtta	caagtagaaa	ttaagacacc	cttgatagcc	ttactatacc	taacatgatg	720
tagtattaaa	tgaatatgta	aatatattta	tgataagaag	cgacttattt	ataatcatta	780
catatttttc	tattggaatg	attaagattc	caatagaata	gtgtataaat	tatttatctt	840
gaaaggaggg	atgcctaaaa	acgaagaaca	ttaaaaacat	atatttgcac	cgctcaatgg	900
atztatgaaa	aatcatttta	tcagtttgaa	aattatgtat	tatggccaca	ttgaaagggg	960
aggagaatca	tgaacaaca	aaaacggctt	tacgccgat	tgctgacgct	gttatttgcg	1020
ctcatcttct	tgctgcctca	ttctgcagca	gcggcggata	acaaatttaa	caaagaacag	1080
cagaacgcgt	tttatgaaat	cctgcattctg	ccgaacctga	acgaagaaca	gcgcaacgcg	1140
tttatccaga	gcctgaaaga	tgatccgagc	cagagcgcga	acctgctggc	ggaagcgaaa	1200
aaactgaacg	atgcgcaggg	gccgaaagtc	gacaacaaat	ttaataagga	acaacaaaac	1260
gccttctacg	aaattctcca	tctaccaaat	cttaatgaag	agcagcgcaa	cgcttttatt	1320
caatcgctta	aggacgatcc	cagccaatcg	gcaaatcttc	tggtgaagc	taaaaagctt	1380
aatgacgcgc	aagctcctaa	agtggacccg	tcaaaagatt	ccaaagccca	ggtaagtgc	1440
gctgaagcag	gcataacggg	aacatggtag	aatcaactgg	gaagcacgtt	tatcgtaaca	1500
gcaggagcgg	acggagcgtt	gacaggcaca	tacgaatctg	ccgttggaag	cgcagaatcg	1560
cgttacacac	ttacaggtag	atacgattca	gcaccggcta	ccgatggaag	cggaaactgc	1620
ttgggatgga	gagtggtctg	gaaaaacaac	tatagaaatg	cgcatagcgc	cacgacatgg	1680
tctggccaat	acgtcggagg	agcagaagct	agaatcaata	cacaatggac	attaacgagc	1740
gggactacag	aagcaaatgc	ctggaaatcg	acgttaagag	gtcatgatac	ctttaccaa	1800
gtaaagccgt	ccgcagcatc	aattgatgca	gcgaagaaag	caggagtga	taacggtaat	1860
cctttggatg	cagtgaaca	atagtcgcta	gcggccgcgt	cgactagaag	agcagagagg	1920
acggatttcc	tgaaggaaat	ccgttttttt	atcttgcccg	tcttataaat	ttcgttgaga	1980
tcttgatcat	cgataagctt	ggcgtaatac	tggtcatagc	tgtttcctgt	gtgaaattgt	2040
tatccgcctca	caattccaca	caacatacga	gccggaagca	taaagtgtaa	agcctggggg	2100
gcctaattgag	tgagctaaat	cacattaatt	gcgttcgcgt	cactgcccgc	ttccagtcg	2160
ggaaacctgt	cgtagccagct	gcattaatga	atcggccaac	gcgcggggag	aggcggtttg	2220
cgtattgggc	gctcttccgc	ttcctcgctc	actgactcgc	tgcgctcggt	cgttcggctg	2280
cgcgagcgg	tatcagctca	ctcaaaggcg	gtaatacggg	tatccacaga	atcaggggat	2340
aacgcaggaa	agaacatgtg	agcaaaaggc	cagcaaaagg	ccagggaaccg	taaaaaggcc	2400
gcgttgctgg	cgtttttcca	taggctccgc	ccccctgacg	agcatcaca	aaatcgacgc	2460
tcaagtcaga	ggtggcgaaa	cccgcagga	ctataaagat	accaggcggt	tccccctgga	2520

```

agctccctcg tgcgtctcc tgttccgacc ctgccgctta ccggatacct gtccgccttt 2580
ctcccttcgg gaagcgtggc gctttctcaa tgctcacgct gtaggtatct cagttcggtg 2640
taggtcggtc gctccaagct gggctgtgtg cacgaacccc ccgttcagcc cgaccgctgc 2700
gccttatccg gtaactatcg tcttgagtcc aacccggtaa gacacgactt atcgccactg 2760
gcagcagcca ctggtaacag gattagcaga gcgaggatg taggcggtgc tacagagttc 2820
ttgaagtggg ggcctaacta cggctacact agaaggacag tatttggat ctgcgctctg 2880
ctgaagccag ttaccttcgg aaaaagagtt ggtagctctt gatccggcaa acaaccacc 2940
gctggtagcg gtggtttttt tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct 3000
caagaagatc ctttgatctt ttctacgggg tctgacgctc agtggaaacga aaactcacgt 3060
taagggtatt tgggtcatgag attatcaaaa aggatcttca cctagatcct tttaaattaa 3120
aaatgaagtt ttaaatacat ctaaagtata tatgagtaaa cttggtctga cagttacca 3180
tgcttaatca gtgaggcacc tatctcagcg atctgtctat ttctgttcac catagttgcc 3240
tgactccccg tctgttagat aactacgata cgggagggct taccatctgg cccagtgct 3300
gcaatgatac cgcgagaccc acgctcaccg gtccagatt tatcagcaat aaaccagcca 3360
gccggaaggg ccgagcgtag aagtggctct gcaactttat ccgcctcat ccagtctatt 3420
aattgttgcg ggaagctag agtaagtagt tcgccagtta atagtttgcg caacgttgtt 3480
gccattgcta caggcatcgt ggtgtcacgc tctgtcttgc gtatggcttc attcagctcc 3540
ggttcccaac gatcaaggcg agttacatga tccccatgt tgtcagaaa agcggttagc 3600
tccttcggtc ctccgacgt tgtcagaagt aagtggccg cagtgttatc actcatggtt 3660
atggcagcac tgcataattc tcttactgtc atgccatccg taagatgctt ttctgtgact 3720
ggtgagtact caaccaagtc attctgagaa tagtgtatgc ggcgacggag ttgctcttgc 3780
ccggcgtaaa tacgggataa taccgcgcca catagcagaa ctttaaaagt gctcatcatt 3840
ggaaaacggt cttcggggcg aaaaactctc aggatcttac cgctgttgag atccagttcg 3900
atgtaacca ctcgtgcacc caactgatct tcagcatctt ttactttcac cagcgtttct 3960
gggtgagcaa aaacaggaag gcaaatgcc gcaaaaaagg gaataaggcg gacacggaaa 4020
tgttgaaac tcatactctt ctttttcaa tattattgaa gcatttatca gggttattgt 4080
ctcatgagcg gatacatatt tgaatgtatt tagaaaaata aacaaatagg ggttccgcgc 4140
acatttcccc gaaaagtgcc acctgacgtc taagaaacca ttattatcat gacattaacc 4200
tataaaaata ggcgtatcac gaggcccttt cgtc 4234

```

<210> 35

<211> 6453

<212> DNA

<213> Artificial

<220>

<223> plasmid sequence pMOL2746b

<400> 35

gaattcagat ctaaagataa tatctttgaa ttgtaacsc cctcaaaagt aagaactaca

60

aaaaaagaat	acgttatata	gaaatatgtt	tgaaccttct	tcagattaca	aatatatctg	120
gacggactct	acctcaaatg	cttatctaac	tatagaatga	catacaagca	caaccttgaa	180
aatttgaaaa	tataactacc	aatgaacttg	tccatgtgaa	ttatcgctgt	atttaatttt	240
ctcaattcaa	tatataatat	gccaatacat	tgttacaagt	agaaattaag	acacccttga	300
tagccttact	atacctaaca	tgatgtagta	ttaaatgaat	atgtaaatat	atttatgata	360
agaagcgact	tattttataat	cattacatat	ttttctattg	gaatgattaa	gattccaata	420
gaatagtgtg	taaattattt	atcttgaaa	gagggatgcc	taaaaacgaa	gaacattaaa	480
aacatatatt	tgcaaccgtct	aatggattta	tgaaaaatca	ttttatcagt	ttgaaaatta	540
tgtattatgg	ccacattgaa	aggggaggag	aatcatgaaa	caacaaaaac	ggctttacgc	600
ccgattgctg	acgctgttat	ttgcgctcat	cttcttgctg	cctcattctg	cagcagcggc	660
ggataacaaa	tttaacaaag	aacagcagaa	cgcgttttat	gaaatcctgc	atctgccgaa	720
cctgaacgaa	gaacagcgca	acgcgtttat	ccagagcctg	aaagatgata	cagaccagag	780
cgcgaacctg	ctggcggaag	cgaaaaaact	gaacgatgcg	caggcgccga	aagtcgacaa	840
caaatttaat	aaggaacaac	aaaacgcctt	ctacgaaatt	ctccatctac	ccaatcttaa	900
tgaagagcag	cgcaacgctt	ttattcaatc	gcttaaggac	gatcccgacc	aatcgcaaaa	960
tcttctggct	gaagctaaaa	agcttaatga	cgcgcaagct	cctaaagtgg	accgctcaaa	1020
agattccaaa	gccaggttaa	gtgcagctga	agcaggcata	acgggaacat	ggtacaatca	1080
actgggaagc	acgtttatcg	taacagcagg	agcggacgga	gcgttgacag	gcacatacga	1140
atctgccgtt	ggaagcgcag	aatcgcggtt	cacacttaca	ggtagatacg	attcagcacc	1200
ggctaccgat	ggaagcggaa	ctgcattggg	atggagagtg	gcttggaata	acaactatag	1260
aaatgcgcac	agcgccacga	catggtcttg	ccaatacgtc	ggaggagcag	aagctagaat	1320
caatacacia	tggacattaa	cgagcgggac	tacagaagca	aatgcctgga	aatcgacgtt	1380
aagaggtcat	gataacctta	ccaaagtaaa	gccgtccgca	gcatcaattg	atgcagcgaa	1440
gaaagcagga	gtgaataacg	gtaatccttt	ggatgcagtg	caacaatagt	cgctagcggc	1500
cgcgtcgact	agaagagcag	agaggacgga	tttcctgaag	gaaatccgtt	tttttatatt	1560
gcccgtctta	taaatttcgt	tgagatctac	gcgtccatgg	gctagcgcgg	ccgcgtcgac	1620
aggcctcttt	gattacattt	tataattaat	tttaacaaag	tgctatcagc	cctcaggaag	1680
gacttgctga	cagtttgaat	cgcataggta	aggcggggat	gaaatggcaa	cgttatctga	1740
tgtagcaaa	aaagcaaatg	tgtcgaaaa	gacggtatcg	cgggtgatca	atcatcctga	1800
gactgtgacg	gatgaattga	aaaagcttgt	tcattccgca	atgaaggagc	tcaattatat	1860
accgaactat	gcagcaagag	cgctcgttca	aaacagaaca	caggctgtca	agctgctcat	1920
actggaagaa	atggatataa	cagaacctta	ttatatgaat	ctgttaacgg	gaatcagccg	1980
cgagctggac	cgctatcatt	atgctttgca	gcttgtcaca	aggaaatctc	tcaatatcgg	2040
ccagtgcgac	ggcattattg	cgacgggggt	gagaaaagcc	gattttgaag	ggctcatcaa	2100

```

ggtttttgaa aagcgtgtcg ttgtattcgg gacgtcgatt cacaaaaata ggcacacgaa 2160
aaacaagtaa gggatgcagt ttatgcatcc ctttaacttac ttattaaata atttatagct 2220
attgaaaaga gataagaatt gttaaagct aatattgttt aaatcgtcaa ttcctgcatg 2280
ttttaaggaa ttgttaaatt gattttttgt aaatatattc ttgtattctt tgtaaaccca 2340
tttcataacg aaataattat acttttgttt atctttgtgt gatattcttg atttttttct 2400
acttaactcg ataagtgagc ttttactttt aggttttaga tgaaaaatatt ctcttggaac 2460
catacttaat atagaaatat caacttctgc cattaaaagt aatgccaatg agcgttttgt 2520
atttaataat ctttttagcaa acccgatttc cagattaaa taaatctcat tagctatact 2580
atcaaaaaca attttgcgta ttatatccgt acttatgtta taaggatatat taccatatat 2640
tttataggat tgggtttttag gaaatttaaa ctgcaatata tccttgitta aaacttgga 2700
attatcgtga tcaacaagtt tattttctgt agttttgcat aatttatggt ctatttcaat 2760
ggcagttacg aaattacacc tctttactaa ttcaagggtt aaatggcctt ttcctgagcc 2820
gatttcaaag atattatcat gtccatttaa tcttatattt gtcattattt tatctatatt 2880
atgttttgaa gtaataaagt ttgtactgtg ttttatattt ttctcgttca ttataaccct 2940
ctttaatttg gttatatgaa ttttgcttat taacgattca ttataaccac ttattttttg 3000
tttggttgat aatgaactgt gctgattaca aaaatactaa aaatgccat attttttcct 3060
ccttataaaa ttagtataat tatagcacga gctctgataa atatgaacat gatgagtgat 3120
cgtaaaattt atactgcaat cggatgcgat tattgaataa aagatatgag agatttatct 3180
aatttctttt ttcttgtaaa aaaagaaagt tcttaagggt ttatagttt tggtcgtaga 3240
gcacacggtt taacgactta attacgaagt aaataagtct agtggttag actttatgaa 3300
atctatatac gtttatatat atttattatc cggagggtga gcatgtctca ttcaattttg 3360
agggttgcca gagttaaagg atcaagtaat acaaacggga tacaagaca taatcaaaga 3420
gagaataaaa actataataa taaagacata aatcatgagg aaacatataa aaattatgat 3480
ttgattaacg cacaaaaat aaagtataaa gataaaattg atgaaacgat tgatgagaat 3540
tattcagga aacgtaaaat tcggtcagat gcaattcgac atgtggacgg actggttaca 3600
agtgataaag atttctttga tgatttaagc ggagaagaaa tagaacgatt ttttaagat 3660
agcttgaggt ttctagaaaa tgaatacgtt aaggaaaata tgctgtatgc gactgtccat 3720
ctggatgaaa gagtcccaca tatgcacttt ggttttgtcc ctttaacaga ggcgggaga 3780
ttgtctgcaa aagaacagtt aggaacaag aaagacttta ctcaattaca agatagattt 3840
aatgagtatg tgaatgagaa aggttatgaa cttgaaagag gcacgtccaa agaggttaca 3900
gaacgagaac ataaagcgtt ggatcagtac aagaagata ctgtatttca taaacaggaa 3960
ctgcaagaag ttaaggatga gttacagaag gcaaataagc agttacagag tggaaatagag 4020
catatgaggt ctacgaaacc ctttgattat gaaaatgagc gtacagggtt gttctctgga 4080
cgtgaagaga ctggtagaaa gatattaact gctgatgaat ttgaacgcct gcaagaaca 4140
atctcttctg cagaacggat tgttgatgat tacgaaaata ttaagagcac agactattac 4200

```

```

acagaaaatc aagaattaaa aaacgtaga gagagtttga aagaagtagt gaatacatgg 4260
aaagaggggt atcacgaaaa agtaaaagag gttaataaat taaagcgaga gaatgatagt 4320
ttgaatgagc agttgaatgt atcagagaaa tttcaagcta gtacagtgac tttatatcgt 4380
gctgcgaggg cgaatttccc tgggtttgag aaagggttta ataggcttaa agagaaattc 4440
tttaatgatt ccaaatttga gcgtgtggga cagtttatgg atgttgtaca ggataatgtc 4500
cagaaggtcg atagaaagcg tgagaaacag cgtacagacg atttagagat gtagaggtag 4560
ttttatgccg agaaaacttt ttgctgtgta cagtccttaa aatatactta gagcgttaagc 4620
gaaagttagt gcgacagcta ttaactttcg gtttcaaagc tctaggattt ttaatggacg 4680
cagcgcatca cgcgaaaaa ggaaattgga ataatgcga aatttgagat gtttaattaaa 4740
gaccttttgg aggtcttttt ttcttagatt tttgggggta tttaggggag aaaacatagg 4800
gggggtactac gacctcccc ctaggtgtcc attgtccatt gtccaaacaa ataaataaat 4860
attgggtttt taatgtttaa aggtgttttt ttatgtttaa gtgaaaaaaa cagatgttgg 4920
gaggtacagt gatggttgtg gatagaaaag aagagaaaaa agttgctgtt actttaagac 4980
ttacaacaga agaaaatgag atattaaata gaatcaaaga aaaatataat attagcaaat 5040
cagatgcaac cggtatttcta ataaaaaaat atgcaaagga ggaatacggg gcatttttaa 5100
caaaaaaaga tagacagcac tggcatgctg cctatctatg actaaatttt gttaagtgtg 5160
ttagcaccgt tattatatca tgagcgaaaa tgtaataaaa gaaactgaaa acaagaaaaa 5220
ttcaagagga cgtaatttga cttttgtttt atatccagaa tcagcaaaaag ccgagtgggt 5280
agagtattta aaagagttac acattcaatt tgtagtgtct ccattacatg atagggatac 5340
tgatacagaa ggtagtagta aaaaagagca ttatcatatt ctagtgtagt atgagggtaa 5400
taaatcttat gaacagataa aaataattac agaagaattg aatgcgacta ttccgcagat 5460
tgcaggaagt gtgaaaggtc ttgtgagata tatgtctcac atggacgac ctaataaatt 5520
taaatatcaa aaagaagata tgatagttaa tggcgggtga gatgttgatg aattattaaa 5580
gaaaacaaca acagatagat ataaattaat taaagaaatg attgagttaa ttgatgaaca 5640
aggaatcgta gaatttaaga gtttaatgga ttatgcaatg aagtttaaat ttgatgattg 5700
gttcccgcct ttatgtgata actcggcgta tgttattcaa gaatatataa aatcaaatcg 5760
gtataaatct gaccgataga ttttgaattt aggtgtcaca agacactctt ttttcgcacc 5820
agcgaaaact ggtttaagcc gactgcgcaa aagacataat cgactctaga ggatccccgg 5880
gtaccgagct ctgcctttta gtccagctga tttcactttt tgcatctcac aaactgcata 5940
actcatatgt aaatcgctcc tttttagggt gcacaaatgt gaggcatttt cgctctttcc 6000
ggcaaccact tccaagtaaa gtataacaca ctatacttta tattcataaa gtgtgtgctc 6060
tgcgagggctg tcggcagtg cgcacaaaac cataaaacct ttaagacctt tctttttttt 6120
acgagaaaaa agaaacaaaa aaacctgccc tctgccacct cagcaaaagg gggttttgct 6180
ctcgtgctcg tttaaaaatc agcaagggac aggtagtatt ttttgagaag atcactcaaa 6240

aaatctccac ctttaaacc ttgccaattt ttattttgtc cgttttgtct agcttaccga 6300
aagccagact cagcaagaat aaaattttta ttgtctttcg gttttctagt gtaacggaca 6360
aaaccactca aaataaaaaa gatacaagag aggtctctcg tatcttttat tcagcaatcg 6420
cgccccgattg ctgaacagat taataatgag ctc 6453

```

<210> 36

<211> 62

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 36

```

tcattctgca gcagcgcgga ataacaatt taacaaagaa cagcagaacg cgttttatga 60
aa 62

```

<210> 37

<211> 52

<212> DNA

<213> Artificial

<220>

<223> Primer

<400> 37

aactaagcgg cgcctagcga ctacacttta ggagcttgcg cgtcattaag ct 52

<210> 38

<211> 1062

<212> DNA

<213> Artificial

<220>

<223> ORF of protein A fused to streptavidin

<400> 38

atgaaacaac aaaaacggct ttacgcccgga ttgctgacgc tgttatttgc gctcatcttc	60
ttgctgcctc attctgcagc cgcggcagac cctccaagg actcgaaggc ccaggctctg	120
gccgccgagg ccggcatcac cggcacctgg tacaaccagc tcggctcgac cttcatcgtg	180
accgcggggc ccgacggcgc cctgaccgga acctacgagt cggccgtcgg caacgccgag	240
agccgctacg tcctgaccgg tcgttacgac agcgcgccgg ccaccgacgg cagcggcacc	300
gccctcggtt ggacgggtggc ctggaagaat aactaccgca acgcccactc cgcgaccacg	360
tggagcggcc agtacgtcgg cggcgccgag gcgaggatca acaccagtg gctgctgacc	420
tccggcacca ccgaggccaa cgcctggaag tccacgctgg tcggccacga caccttcacc	480
aagggtgaagc cgtccgccgc ctccatcgac gcggcgaaga aggcgggctg caacaacggc	540
aaccgcctcg acgccgttca gcagtcgaca ttacttatat ctggtggcgt aacacctgct	600
gcaaatgctg cgcaacacga tgaagctcaa caaatgctt tttatcaagt cttaaataatg	660
cctaacttaa atgctgatca acgcaatggt tttatccaaa gccttaaga tgatccaagc	720
caaagtgcta acgttttagg tgaagctcaa aaacttaatg actctcaagc tccaaaagct	780

gatgcgcaac aaaataactt caacaaagat caacaaagcg ctttctatga aatcttgaac	840
atgcctaact taaacgaagc gcaacgtaac ggcttcattc aaagtcttaa agacgacca	900
agccaaagca ctaacgtttt aggtgaagct aaaaaattaa acgaatctca agcaccgaaa	960
gctgataaca atttcaacaa agaacaacaa aatgctttct atgaaatctt gaatatgcct	1020
aacttaaacg aagaacaacg caatggtttc atccaagct ta	1062

<210> 39

<211> 354

<212> PRT

<213> Artificial

<220>

<223> ProteinA streptavidin fusion protein

<400> 39

Met Lys Gln Gln Lys Arg Leu Tyr Ala Arg Leu Leu Thr Leu Leu Phe
 1 5 10 15
 Ala Leu Ile Phe Leu Leu Pro His Ser Ala Ala Ala Ala Asp Pro Ser
 20 25 30
 Lys Asp Ser Lys Ala Gln Val Ser Ala Ala Glu Ala Gly Ile Thr Gly
 35 40 45
 Thr Trp Tyr Asn Gln Leu Gly Ser Thr Phe Ile Val Thr Ala Gly Ala
 50 55 60
 Asp Gly Ala Leu Thr Gly Thr Tyr Glu Ser Ala Val Gly Asn Ala Glu
 65 70 75 80
 Ser Arg Tyr Val Leu Thr Gly Arg Tyr Asp Ser Ala Pro Ala Thr Asp
 85 90 95
 Gly Ser Gly Thr Ala Leu Gly Trp Thr Val Ala Trp Lys Asn Asn Tyr
 100 105 110
 Arg Asn Ala His Ser Ala Thr Thr Trp Ser Gly Gln Tyr Val Gly Gly
 115 120 125
 Ala Glu Ala Arg Ile Asn Thr Gln Trp Leu Leu Thr Ser Gly Thr Thr
 130 135 140
 Glu Ala Asn Ala Trp Lys Ser Thr Leu Val Gly His Asp Thr Phe Thr
 145 150 155 160
 Lys Val Lys Pro Ser Ala Ala Ser Ile Asp Ala Ala Lys Lys Ala Gly
 165 170 175
 Val Asn Asn Gly Asn Pro Leu Asp Ala Val Gln Gln Ser Thr Leu Leu
 180 185 190
 Ile Ser Gly Gly Val Thr Pro Ala Ala Asn Ala Ala Gln His Asp Glu
 195 200 205
 Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn Met Pro Asn Leu Asn
 210 215 220
 Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser
 225 230 235 240
 Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys Leu Asn Asp Ser Gln
 245 250 255
 Ala Pro Lys Ala Asp Ala Gln Gln Asn Asn Phe Asn Lys Asp Gln Gln
 260 265 270
 Ser Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Ala Gln
 275 280 285
 Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Thr
 290 295 300
 Asn Val Leu Gly Glu Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys
 305 310 315 320
 Ala Asp Asn Asn Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile
 325 330 335
 Leu Asn Met Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln
 340 345 350
 Ser Leu

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [EP1529844A \[0012\]](#)
- [US5328985A \[0017\]](#)
- [WO9719957A \[0019\]](#)
- [WO0195857A \[0020\]](#)
- [WO9709668A \[0021\] \[0021\]](#)
- [WO0024883A \[0117\] \[0117\]](#)
- [EP238023A \[0132\] \[0206\]](#)
- [WO9600787A \[0132\]](#)
- [WO0039322A \[0204\]](#)
- [WO9812300A \[0204\]](#)
- [WO2005070962A \[0206\]](#)
- [WO2006050737A \[0206\]](#)
- [EP0531372A \[0206\]](#)
- [WO9414968A \[0228\]](#)
- [US5643720A \[0230\]](#)
- [WO2005123915A \[0230\] \[0237\]](#)
- [WO2000075344A \[0242\]](#)

Non-patent literature cited in the description

- **JASON, J-CRYDÉN, L.** Protein Purification: Principles, high-Resolution, Methods and Applications Wiley & sons Inc. 19980000 [\[0002\]](#)
- **WILCHEK, M. GORECKI, M.** A New Approach for isolation of Biologically Active Compounds by Affinity Chromatography: Isolation of Trypsin, 1973, [\[0003\]](#)
- FEBS Letters, vol. 31, 1149-152 [\[0004\]](#)
- **HAMMARBERGH, B. et al.** Proc. Natl. Acad. Sci USA, 1989, vol. 86, 4367-4371 [\[0006\]](#)
- **FORD, C.F. SUOMINEN, IGLATZ, C.E.** Fusion Tails for the Recovery and Purification of Recombinant Proteins Protein Expression and Purification, 1991, vol. 2, 95-107 [\[0008\]](#)
- **RIGAUT, G. et al.** A Generic Protein Purification Method for Protein Complex Characterization and Proteome Exploration Nature Biotechnology, 1991, vol. 17, 1030-1032 [\[0010\]](#)
- **LINDER, M. NEVANEN, T. SÖDERHOLM, L. BENGS, O. TEERI, T.** Biotechnology and Bioengineering, 1998, vol. 60, 5642-647 [\[0012\]](#)
- **SHPIGEL, E. et al.** Expression, purification and application of Staphylococcal Protein A fused to cellulose-binding domain Biotechnol. Appl. Biochem., 2000, vol. 31, 197-203 [\[0014\]](#)
- **RÖNNMARKE** Eur. J. Biochem., 2002, vol. 269, 2647-2655 [\[0067\]](#)
- **MACKENZIE, C.R. et al.** Analysis by surface plasmon resonance of the influence of valence on the ligand binding affinities and kinetics of and anti carbohydrate antibody Journal of Biological Chemistry, 1996, vol. 271, 1527-1533 [\[0074\]](#)
- **NYGREN, H. STENBERG, M.** Immunochimistry at interfaces Immunology, 1989, vol. 66, 321-327 [\[0075\]](#)
- **P.-Å. & SKERRA, A.** Binding proteins from alternative scaffolds J. Immunol. Methods, 2004, vol. 290, 3-28 [\[0090\]](#)
- **BINZ, H. K. PLÜCKTHUN, A.** Engineered proteins as specific binding reagents Curr. Opin. Biotech., 2005, vol. 16, 459-469 [\[0090\]](#)
- **NILSSON, B. MOKS, T. JANSSEN, B. ABRAHMSÉN, L. ELMBLAD, A. HOLMGREN, E. et al.** Protein Eng., 1987, vol. 1, 107-113 [\[0090\]](#)
- **T. SANOC. R. CANTOR** BioTechnology, 1991, vol. 9, 1378-1381 [\[0105\]](#)
- **GEMS et al.** Gene, 1991, vol. 98, 61-67 [\[0117\]](#)
- **CULLEN et al.** Nucleic Acids Research, 1987, vol. 15, 9163-9175 [\[0117\]](#)
- **CHANGCOHEN** Molecular General Genetics, 1979, vol. 168, 111-115 [\[0123\]](#)
- **YOUNGSPIZIN** Journal of Bacteriology, 1961, vol. 81, 823-829 [\[0123\]](#)
- **DUBNAU DAVIDOFF-ABELSON** Journal of Molecular Biology, 1971, vol. 56, 209-221 [\[0123\]](#)
- **SHIGEKAWA DOWER** Biotechniques, 1988, vol. 6, 742-751 [\[0123\]](#)

- **KOEHLER** **THORNE** *Journal of Bacteriology*, 1987, vol. 169, 5771-5278 [0123]
- In, Ainsworth and Bisby's Dictionary of The Fungi **HAWKSWORTH et al.** CAB International University Press 1995 0000 [0125]
- *Biology and Activities of Yeast* Soc. App. Bacteriol. Symposium Series 1980 0000 [0126]
- **YELTON et al.** *Proceedings of the National Academy of Sciences USA*, 1984, vol. 81, 1470-1474 [0132]
- **MALARDIER et al.** *Gene*, 1989, vol. 78, 147-156 [0132]
- *Guide to Yeast Genetics and Molecular Biology* **BECKER** **GUARENTE** *Methods in Enzymology* Academic Press, Inc. vol. 194, 182-187 [0132]
- **ITO et al.** *Journal of Bacteriology*, 1983, vol. 153, 163- [0132]
- **HINNEN et al.** *Proceedings of the National Academy of Sciences USA*, 1978, vol. 75, 1920- [0132]
- **GREG T. HERMANSON** *Bioconjugate Techniques* Academic Press 2008 0000 [0141]
- **SHAN S. WONG** *Chemistry of Protein Conjugation and Cross-linking* CRC Press 1991 0000 [0141]
- **JANSON, J.-C. RYDÉN, L.** *Protein Purification* Wiley & sons inc. 1998 0000 [0161]
- **CASADABAN** **COHEN J.** *Mol. Biol.*, 1980, vol. 138, 179-207 [0204]
- **BOEL et al.** *EMBO Journal*, 1984, vol. 3, 1581-1585 [0205] [0206] [0220]
- **NILSSON et al.** *Protein Engineering*, 1987, vol. 1, 107-113 [0205] [0231]
- **FRENKEN et al.** *J. Biotechnology*, 2000, vol. 78, 11-21 [0205]
- **SAMBROOK et al.** *Molecular cloning: A laboratory manual* Cold Spring Harbor lab. 1989 0000 [0208]
- *Current protocols in Molecular Biology* John Wiley and Sons 1995 0000 [0208]
- *Land and DNA Cloning: A Practical Approach* 1985 0000 [0208]
- *Oligonucleotide Synthesis* 1984 0000 [0208]
- *Nucleic Acid Hybridization* 1985 0000 [0208]
- **B. PERBALA** *Practical Guide To Molecular Cloning* 1984 0000 [0208]
- **CHRISTENSEN et al.** *Biotechnology*, 1988, vol. 6, 1419-1422 [0210]
- **COVE D.J.** *Biochem. Biophys. Acta.*, 1966, vol. 113, 51-56 [0211]
- **DIDERICHSEN, B. WEDSTED, U. HEDEGAARD, L. JENSEN, B. R. SJOHOLM, C.** *Cloning of aldB, which encodes alpha-acetolactate decarboxylase, an exoenzyme from i Bacillus brevis* *J. Bacteriol.*, 1990, vol. 172, 4315-4321 [0230]
- **ARGARANA** *Nucleic Acids Res.*, 1986, vol. 14, 1871-1882 [0231]
- **STEMMER et al.** *Gene*, 1995, vol. 164, 49-53 [0232]
- **DILLON** **ROSSEN** *BioTechniques*, 1990, vol. 9, 298-300 [0232]
- **PRODROMOU** **PEARL** *Protein Engineering*, 1992, vol. 5, 827-829 [0232]
- **CHN et al.** *Journal of American Chemical Society*, 1994, vol. 11, 8799-8800 [0232]
- **GPMAW** - a software tool for analyzing proteins and peptides *Trends in Biochemical Science*, 2001, vol. 26, 11687-689 [0240]
- **LINDER et al.** *Biotechnology and Bioengineering*, 1998, vol. 60, 5 [0313]
- **GREEN, N.** *Biochem J*, 1963, vol. 89, 585-591 [0313]

Patentkrav

1. Fremgangsmåde til oprensning af et målbiomolekyle, omfattende trinnene: (a) kontakte (i) et målbiomolekyle, (ii) et polypeptid med dobbelt affinitet, og (iii) en fast bærer omfattende en indfangende ligand eller et dobbeltaffinitetpolypeptidbindingssted, hvori forholdet mellem ligevægtsdissociationskonstanterne for polypeptidet med dobbelt affinitet, $[K_{D,t} / K_{D,s}]$, er mindst 10^0 ved standardbetingelser; og (b) udvinde målbiomolekylet ved eluering, hvori målaaffinitetspolypeptidet og polypeptidet med dobbelt affinitet bringes i kontakt i opløsning, før blandingen bringes i kontakt med den faste bærer omfattende en indfangende ligand eller et dobbeltaffinitetpolypeptidbindingssted.
2. Fremgangsmåde ifølge krav 1, hvori den faste bærer er valgt fra gruppen omfattende fastfasematrixer og partikler.
3. Fremgangsmåde ifølge krav 1, hvori polypeptidet med dobbelt affinitet har en ligevægtsdissociationskonstant, $K_{D,t}$ mod målbiomolekylet i intervallet fra 10^{-2} til 10^{-13} M, især fra 10^{-4} til 10^{-13} M, fortrinsvis i intervallet fra 10^{-6} til 10^{-13} M, og en ligevægtsdissociationskonstant, $K_{D,s}$ mod den indfangende ligand i intervallet fra 10^{-9} til 10^{-16} M, fortrinsvis i intervallet fra 10^{-11} til 10^{-16} M.
4. Fremgangsmåde ifølge ethvert af de foregående krav, hvori forholdet mellem ligevægtsdissociationskonstanterne for polypeptidet med dobbelt affinitet, $[K_{D,t} / K_{D,s}]$, er mindst 10^1 , mere specielt mindst 10^2 , mere specielt 10^3 , og endnu mere specielt mindst 10^4 .
5. Fremgangsmåde ifølge ethvert af de foregående krav, hvori eluering af målet opnås ved ændring af enten pH, ionstyrke eller indhold af chaotrope ioner i opløsning, eller enhver kombination deraf.
6. Fremgangsmåde ifølge ethvert af de foregående krav, hvori polypeptidet med dobbelt affinitet er et fusionspolypeptid, fortrinsvis valgt fra gruppen bestående af protein A, antistoffer, antistoffragmenter, protein A-fragmenter, protein A-afledte IgG-bindingsdomæner, lipocaliner, lectiner.
7. Fremgangsmåde ifølge ethvert af de foregående krav, hvori den ligandbindende del af polypeptidet med dobbelt affinitet er valgt fra gruppen bestående af avidin, streptavidin, neutravidin, steroid-receptor, antistof, antistoffragment, lipocaliner, lectiner, amyloglucosidase, cellulose-bindingsdomæner.
8. Fremgangsmåde ifølge ethvert af kravene 6-7, hvori antistoffet er valgt fra gruppen bestående af lama- og kamel-antistoffer.
9. Fremgangsmåde ifølge krav 6, hvori fusionspolypeptidet fremstilles ved fusion af mindst et IgG-bindingsdomæne af protein A eller protein A-afledt IgG-bindingsdomæne og mindst et biotinbindingsdomæne af avidin, streptavidin eller neutravidin.

10. Fremgangsmåde ifølge ethvert af de foregående krav, hvori liganden er valgt fra gruppen bestående af biotin, acarbose, steroider, haptener, epitop-peptider, farvestoffer og enzyminhibitorer.

11. Fremgangsmåde ifølge krav 9, hvori den indfangende ligand, der er bundet til den faste bærer, er biotin, og målbiomolekylet er IgG.

12. Fremgangsmåde ifølge ethvert af de foregående krav, hvori den faste bærer er en fastfasematrix, fortrinsvis valgt fra gruppen bestående af agar-agar, agaroser, celluloser, cellulose-ethere, carboxymethylcellulose, polyamider, polyvinylalkoholer, silicaer og glas med kontrollerede porer.

13. Fremgangsmåde ifølge krav 6, hvori fusionspolypeptidet fremstilles som et rekombinant polypeptid i en rekombinant værtselle.

14. Fremgangsmåde ifølge krav 13, hvori fusionspolypeptidet og målbiomolekylet udtrykkes i samme type værtselle.

15. Fremgangsmåde ifølge ethvert af kravene 13-14, hvori værtsellen er valgt fra gruppen bestående af bakterieceller, svampeceller, pattedyrsceller, planteceller og insektceller.

16. Fremgangsmåde ifølge krav 1, hvori polypeptidet med dobbelt affinitet er kemisk fusioneret.

17. Fremgangsmåde til oprensning af et målbiomolekyle, omfattende trinnene: (a) kontakte (i) et målpolypeptid, (ii) et polypeptid med dobbelt affinitet, og (iii) en fast bærer omfattende en indfangende ligand, hvori polypeptidet med dobbelt affinitet har en ligevægtsdissociationskonstant, $K_{D,t}$ mod målbiomolekylet i intervallet fra 10^{-2} til 10^{-13} M, fortrinsvis fra 10^{-4} til 10^{-13} M, mere foretrukket fra 10^{-6} til 10^{-13} M ved standardbetingelser, og hvori bindingen af polypeptidet med dobbelt affinitet til den indfangende ligand på den faste bærer tilvejebringes ved spaltning af en para-substitueret benzylguanin, der resulterer i en thioetherbinding; og (b) udvinde målbiomolekylet ved eluering, hvor målpolypeptidet og polypeptidet med dobbelt affinitet bringes i kontakt i opløsning, før blandingen bringes i kontakt med den faste bærer omfattende en indfangende ligand.