

(19)



(11)

EP 2 891 662 A1

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:

08.07.2015 Bulletin 2015/28

(21) Application number: 15152164.8

(22) Date of filing: 05.02.2010

(51) Int Cl.:

C07K 14/415 (2006.01)

C07K 7/06 (2006.01)

C07K 7/08 (2006.01)

A61K 38/04 (2006.01)

A61K 38/16 (2006.01)

A61K 39/36 (2006.01)

(84) Designated Contracting States:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL  
PT RO SE SI SK SM TR

(30) Priority: 05.02.2009 GB 0901928

05.02.2009 GB 0901927

20.07.2009 GB 0912578

14.08.2009 PCT/GB2009/001995

12.10.2009 GB 0917871

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

10704399.4 / 2 393 830

(71) Applicant: Circassia Limited

Oxford OX4 4GA (GB)

(72) Inventors:

• Hafner, Rod

Oxford OX4 4GA (GB)

- Laidler, Paul  
Oxford OX4 4GA (GB)
- Layton, Guy  
Oxford OX4 4GA (GB)
- Larche, Mark  
Hamilton  
Ontario L8N 3Z5 (CA)

(74) Representative: Woods, Geoffrey Corlett  
J A Kemp  
14 South Square  
Gray's Inn  
London WC1R 5JJ (GB)

### Remarks:

This application was filed on 22-01-2015 as a divisional application to the application mentioned under INID code 62.

(54) Grass peptides for vaccine

(57) The present invention relates to compositions comprising peptides for preventing or treating allergy to grass pollen, and in particular to optimal combinations of peptides for preventing or treating said allergy.

**Description****Incorporating by Reference**

5 [0001] The contents of the following priority applications are incorporated herein by reference: United Kingdom Patent Application No. 0901928.2 filed 5 February 2009, United Kingdom Patent Application No. 0901927.4 filed 5 February 2009, United Kingdom Patent Application No. 0912578.2 filed 20 July 2009, International Patent Application No. PCT/GB09/01995 filed 14 August 2009 and United Kingdom Patent Application No. 0917871.6 filed 12 October 2009,

**10 Field of the Invention**

[0002] The present invention relates to compositions for preventing or treating allergy to grass.

**Background of the Invention**

15 [0003] T-cell antigen recognition requires antigen presenting cells (APCs) to present antigen fragments (peptides) on their cell surface in association with molecules of the major histocompatibility complex (MHC). T cells use their antigen specific T-cell receptors (TCRs) to recognise the antigen fragments presented by the APC. Such recognition acts as a trigger to the immune system to generate a range of responses to eradicate the antigen which has been recognised.

20 [0004] Recognition of external antigens by the immune system of an organism, such as man, can in some cases result in diseases, known as atopic conditions. Examples of the latter are the allergic diseases including asthma, atopic dermatitis and allergic rhinitis. In this group of diseases, B lymphocytes generate antibodies of the IgE class (in humans) which bind externally derived antigens, which are referred to in this context as allergens since these molecules elicit an allergic response. Production of allergen-specific IgE is dependent upon T lymphocytes which are also activated by (are 25 specific for) the allergen. Allergen-specific IgE antibodies bind to the surface of cells such as basophils and mast cells by virtue of the expression by these cells of surface receptors for IgE.

30 [0005] Crosslinking of surface bound IgE molecules by allergen results in degranulation of these effector cells causing release of inflammatory mediators such as histamine, 5-hydroxytryptamine and lipid mediators such as the sulphidoleukotrienes. In addition to IgE-dependent events, certain allergic diseases such as asthma are characterised by IgE-independent events.

35 [0006] Allergic IgE-mediated diseases are currently treated with agents which provide symptomatic relief or prevention. Examples of such agents are antihistamines,  $\beta$ 2 agonists, and glucocorticosteroids. In addition, some IgE-mediated diseases are treated by desensitisation procedures that involve the periodic injection of allergen components or extracts. Desensitisation treatments may induce an IgG response that competes with IgE for allergen, or they may induce specific 40 suppressor T cells that block the synthesis of IgE directed against allergen. This form of treatment is not always effective and poses the risk of provoking serious side effects, particularly general anaphylactic shock. This can be fatal unless recognised immediately and treated with adrenaline. A therapeutic treatment that would decrease or eliminate the unwanted allergic-immune response to a particular allergen, without altering the immune reactivity to other foreign antigens or triggering an allergic response itself would be of great benefit to allergic individuals.

45 [0007] Grass allergens are universally recognised as a major cause of allergic diseases in humans and animals, including asthma, allergic rhinitis and allergic dermatitis. Proteins present in grass pollen are particularly important. For example, approximately 90% of hayfever sufferers are allergic to grass pollen. Hayfever is the common term for a form of seasonal allergy characterised by sneezing, runny nose and itching eyes. The term "hayfever" arose because this form of allergic disease is most prevalent during "haying season", which corresponds to the flowering season of many grasses, when grass plants release the highest quantities of pollen. It is particularly prevalent in summer, typically from the end of May to the end of August (in the Northern Hemisphere).

50 [0008] It has been calculated that for adults in the United States, hayfever is the 5th leading chronic disease and a major cause of work absenteeism, resulting in nearly 4 million missed or lost workdays each year, resulting in a total cost of more than \$700 million in total lost productivity. Allergies are also the most frequently reported chronic condition in children, limiting activities for more than 40% of them. Each year, allergies account for more than 17 million outpatient office visits in the United States; seasonal allergies such as hayfever account for more than half of these allergy visits.

[0009] A therapeutic or preventative treatment would therefore be of great benefit to humans that suffer or are at risk of suffering from grass allergy.

**55 Summary of the Invention**

[0010] Grass pollen allergens are typically classified into groups, with most species expressing at least one pollen allergen in each group. Examples of groups include:

- the Group 1 grass pollen allergens, which include, for example, the proteins Lol p 1 from Rye grass (*Lolium perenne*) and Phl p 1 from Timothy grass (*Phleum pratense*); and
- the Group 5 grass pollen allergens, which include, for example, the proteins Lol p 5a and Lol p 5b from Rye, Phl p 5 from Timothy, and Cyn d 5 from Bermuda grass (*Cynodon dactylon*).

5

**[0011]** The present inventors have discovered that certain peptide fragments derived from the major allergens in the pollens of grass species are particularly useful in desensitising individuals to these allergens. The polypeptide combinations of the invention have been selected for their ability to bind to many MHC Class II molecules, and cause T cell proliferation with minimal histamine release. The compositions, products, vectors and formulations of the invention may therefore be provided to individuals for preventing or treating allergy to grass by tolerisation.

10

**[0012]** The peptides of the invention were selected as MHC class II-binding T cell epitopes through use of *in silico* analysis to predict peptide-MHC interactions and MHC class II binding assays. Additional epitopes were identified by homology. Peptides and peptide combinations of the invention were further selected on the basis of *in vitro* T cell response assays.

15

**[0013]** The peptide combinations of the invention, however, provide a broad coverage of efficacy over the human population by targeting multiple different MHC molecules. A vaccine formulated with peptides of the invention would therefore have broad utility.

**[0014]** The inventors' work has produced peptide combinations with the following characteristics:

20

- the combination binds to many different MHC Class II molecules
- the combination gives significant stimulation of cytokine release in grass allergic individuals
- the peptides of the combinations do not give significant histamine release.

25

The peptide combinations are selected from an extensive analysis of grass epitope sequences to identify core peptides having particularly good MHC binding properties and cytokine response profiles in grass allergic individuals. The combinations may comprise a core of three such peptides, one of each being selected from the most prevalent grasses, Timothy, Perennial Rye and Bermuda. The combinations may comprise a core of four such peptides, each of these peptides being selected from peptides shown to be particularly effective in combination with each other.

30

**[0015]** Providing a core group of peptides in this manner, including a core group of polypeptides representing the three most prevalent grasses, preferably a core group of four or more different individual polypeptides in the same composition provides a variety of MHC-binding epitopes, and so builds in redundancy to allow for the polymorphic nature of the MHC. Providing a core group of the invention, preferably four or more different individual polypeptides having particularly good MHC binding properties and cytokine response profiles also provides multiple T cell epitopes, allowing recruitment of a broad range of T cell specificities for induction of tolerance. Thus, a composition of the invention has beneficial additive effects over use of single polypeptides. The Inventors have also shown that compositions of the invention were able to provide extremely high coverage of a polymorphic study population as measured by cytokine responses.

35

**[0016]** Accordingly, the present invention provides a composition suitable for use in preventing or treating allergy to grass pollen by tolerisation comprising:

40

- (a) at least one of the polypeptides Tim07B (KIPAGELQIIDKIDA), Tim 10B (KYTVFETALKKAITAMSE), Tim 04A (WGAIWRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof;
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQQED), Ber02 (FIPMKSSWGA), Ber02C (KSSWGAI-WRIDPKKPLK) and Ber 02B KDSDEFIPMKSSWGAIR or a variant of any thereof; and
- (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEALTKAIT), Bio02A (KY-DAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIIPSLEAAVKQAY), Rye 05C (NAGFKAAVAAAANAPPK), or a variant of any thereof,

wherein said variant is:

50

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a), (b) or (c), or
- II) a polypeptide of 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a), (b) or (c), which sequence is capable of tolerising to said corresponding polypeptide; or
- III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a), (b) or (c), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

[0017] The invention further provides a composition suitable for use in preventing or treating allergy to grass pollen by tolerisation comprising at least four different polypeptides selected from:

- (a) Tim07B (KIPAGELQIIDKIDA) or a variant thereof;
- (b) Ber01 (SGKAFGAMAKKGQED) or a variant thereof;
- (c) Bio04A (LKKAVTAMSEAEK) or a variant thereof;
- (d) Rye09B (PEVKYAVFEAALTAKAIT) or a variant thereof;
- (e) Ber02 (FIPMKSSWGA) or a variant thereof;
- (f) Ber02C (KSSWGAIWRIDPKKPLK) or a variant thereof;
- (g) Bio03A (KFIPTLVAAVKQAYAAKQ) or a variant thereof; and
- (h) Bio02A (KYDAYVATLTEALR) or a variant thereof;

wherein said variant is:

- 15 I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a) to (h), or
- II) a polypeptide 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a) to (h), which sequence is capable of tolerising to said corresponding polypeptide; or
- 20 III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a) to (h), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

25 [0018] The compositions of the invention are typically capable of tolerising at least 50% or at least 60% of a panel of grass pollen allergic individuals in the population and/or comprises at least one further polypeptide up to a total of fourteen polypeptides, wherein the further polypeptides:(a) comprise a sequence having at least 65% sequence identity to at least 9 or more contiguous amino acids in any of SEQ ID NO: 1 to 73 described below; and (b) are 9 to 30 amino acids in length.

#### Description of the sequences mentioned herein

30 [0019] SEQ ID NOS: 1 to 73 provide the polypeptide sequences of the invention as set out in Tables 2 to 4. SEQ ID NOS: 1 to 27 correspond to peptides derived from Group 1 grass allergens. SEQ ID NOS: 28 to 73 correspond to peptides derived from Group 5 allergens.

#### Detailed description of the invention

35 [0020] The invention concerns peptides which can be used in tolerisation. Such peptides may be derived from any grass allergen described herein or be variants thereof as described below. A difficulty associated with approaches to desensitisation based on peptide immunisation lies in how to select an appropriate size and region of the allergen as the basis for the peptide to be used for immunisation. The size of the peptide of choice is crucial. If the peptide is too small, the vaccine would not be effective in inducing an immunological response. If the peptides are too large, or if the whole antigen is introduced into an individual, there is the risk of inducing adverse reactions, such as anaphylaxis, which may be fatal.

40 [0021] The polypeptides of the invention have been selected to retain T cell specificity whilst being small enough in size to not possess significant tertiary structure that would enable them to retain the conformation of an IgE-binding epitope of the whole molecule. The polypeptides of the invention therefore do not induce significant crosslinking of adjacent specific IgE molecules on cells such as mast cells and basophils and consequently do not cause significant histamine release.

45 [0022] An advantage of the invention is the ability of the peptides to broadly target Major Histocompatibility Complex (MHC) molecules. T cell receptors (TCRs) are highly variable in their specificity. Variability is generated, as with antibody molecules, through gene recombination events within the cell. TCRs recognise antigen in the form of short peptides bound to molecules encoded by the genes of the Major Histocompatibility Complex (MHC). These gene products are 50 the same molecules that give rise to "tissue types" used in transplantation and are also referred to as Human Leukocyte Antigen molecules (HLAs) which terms may be used interchangeably. Individual MHC molecules possess peptide binding grooves which, due to their shape and charge are only capable of binding a limited group of peptides. The peptides bound by one MHC molecule may not necessarily be bound by other MHC molecules.

[0023] When a protein molecule such as an antigen or allergen is taken up by antigen presenting cells such as B lymphocytes, dendritic cells, monocytes and macrophages, the molecule is enzymatically degraded within the cell. The process of degradation gives rise to peptide fragments of the molecule which, if they are of the appropriate size, charge and shape, may then bind within the peptide binding groove of certain MHC molecules and be subsequently displayed upon the surface of antigen presenting cells. If the peptide/MHC complexes are present upon the antigen presenting cell surface in sufficient numbers they may then activate T cells which bear the appropriate peptide/MHC-specific T cell receptors.

Due to the polymorphic nature of the MHC, individuals in an outbred population such as man will express different combinations of MHC molecules on their cell surfaces. Since different MHC molecules can bind different peptides from the same molecule based on the size, charge and shape of the peptide, different individuals will display a different repertoire of peptides bound to their MHC molecules. Identification of universal MHC-binding peptide epitopes in an outbred population such as man is more difficult than in inbred animals (such as certain strains of laboratory mice). On the basis of differential MHC expression between individuals and the inherent differences in peptide binding and presentation which this brings, it is unlikely that a single peptide can be identified which will be of use for desensitisation therapy in man.

[0024] As discussed below, the invention addresses this problem by providing compositions based on more than one peptide i.e. peptide combinations. These combinations are based on core groups of peptides identified experimentally as showing surprisingly strong responses in grass allergic patients when tested individually, and particularly when provided in combination. These core subsets of peptides have advantageous properties when included in grass peptide vaccines.

[0025] The core groups may comprise at least one peptide derived from Timothy grass, at least one peptide derived from Perennial Rye and at least one peptide derived from Bermuda grass, each peptide showing particularly good individual response characteristics. The inclusion of such peptides in combination allows for coverage of the three most prevalent grasses, and additionally extends to coverage of other grasses by homology, as discussed further below.

[0026] Additionally a core group may comprise at least four peptides selected from peptides having the most highly ranked individual response characteristics from the group tested, and further shown to provide optimal response characteristics for a grass vaccine when provided together combination.

[0027] Such peptide combinations preferably comprise at least one peptide derived from a Group I grass allergen or a variant thereof and at least one peptide derived from a Group V grass allergen or a variant thereof. The classification of grass allergens as Group I and Group V grass allergens is well known to the skilled person, and is further discussed herein. A preferred Group I grass allergen is the Bermuda Grass allergen, Cyn d 1. A preferred Group V grass allergen is the Rye Grass allergen, Lol p 5. In some embodiments, the invention relates to a composition comprising at least one peptide derived from Cyn d 1 or a variant thereof and at least one peptide derived from Lol p 5 or a variant thereof. Compositions based on combinations of Cyn d 1 and Lol p 5 peptides or variants thereof are further discussed below.

[0028] Preferred peptides of the invention may comprise, consist of, or consist essentially of the sequences shown in any of SEQ ID NOS: 1 to 73. Variants of these specific peptides may also be used. The variants may comprise, consist of, or consist essentially of sequences which are fragments of either any of SEQ ID NOS: 1 to 73 or homologues of any of SEQ ID NOS: 1 to 73.

[0029] The invention thus provides a composition suitable for use in preventing or treating allergy to grass pollen by tolerisation comprising:

- (a) at least one of the polypeptides Tim07B (KIPAGELQIIDKIDA), , Tim 10B (KYTVFETALKKAITAMSE), Tim 04A (WGAIWRIDTPDKL), Tim 07G (FKVAATAANAAPANDK), or a variant of any thereof;
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), Ber02C (KSSWGAI-WRIDPKKPLK), and Ber 02B KDSDEFIPMKSSWGAIR, or a variant of any thereof; and
- (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTKAIT), Bio02A (KY-DAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIIPSLEAAVKQAY), Rye 05C (NAGFKAAVAAAANAPPK), or a variant of any thereof,

50 wherein said variant is:

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a), (b) or (c), or
- II) a polypeptide of 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a), (b) or (c), which sequence is capable of tolerising to said corresponding polypeptide; or
- III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a), (b) or (c), or a sequence that has at least 65%

homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

**[0030]** Other preferred variants of a) to c) are:

5 i) polypeptides having a length of 9 to 30 amino acids and comprising a region consisting of:

- any of the corresponding sequences of a) to c), or
- a sequence which has at least 65% homology to any of the corresponding sequences of a) to c) which sequence is capable of tolerising an individual to any of the sequences of a) to c), or

10 ii) polypeptides having a length of 9 to 30 amino acids and comprising a region consisting of a sequence that represents either:

15 - a fragment of any of the sequences of a) to c), or

- a homologue of a fragment of any of the sequences of a) to c),

20 which sequence is capable of tolerising an individual to any of the sequences of a) to c) and has a length of at least 9 amino acids, and wherein said homologue has at least 65% homology to any 9 contiguous amino acids in any of the sequences of a) to c); wherein optionally none of the four polypeptides are variants of the same original sequence defined by any one of a) to c).

**[0031]** The composition thus comprises a minimum of three polypeptides, a first polypeptide being a polypeptide of a) or a variant thereof, a second polypeptide being a polypeptide of b) or a variant thereof, and a third polypeptide being a polypeptide of c) or a variant thereof. Thus, for instance the composition may comprise a variant of Tim07B, Ber01 and a variant of Rye09B. Preferably, the above composition comprises in addition to the core three polypeptides a fourth polypeptide of a), b), or c) or a variant of any thereof. The composition may comprise four, five, six, seven, eight or more polypeptides of a), b), c) or variants thereof.

**[0032]** In some aspects, the composition may comprise two, three, four or more polypeptides of a) or variants thereof.

30 In additional aspects, the composition may comprise two, three, four or more polypeptides of b) or variants thereof. In further aspects the composition may comprise two, three, four, five, six or more polypeptides of c) or variants thereof. These selections are subject to the composition comprising at least one polypeptide of a) or a variant thereof, at least one polypeptide of b) or a variant thereof. A preferred combination of three polypeptides selected from groups a), b) and c) is the polypeptide Tim07B or a variant thereof, the polypeptide Ber01 or a variant thereof, and the polypeptide Bio04A or a variant thereof.

**[0033]** The composition may comprise at least two polypeptides selected from group a) or variants thereof, at least two polypeptides selected from group b) or variants thereof and at least two polypeptides selected from group c) or variants thereof.

**[0034]** A preferred composition may comprise the polypeptide Tim07B or a variant thereof and at least two, three or four polypeptides selected from group b) or variants thereof. This preferred composition may comprise two or three polypeptides selected from group c) or variants thereof, preferably including Bio04A or a variant thereof. Another preferred composition may comprise the polypeptide Tim07B or a variant thereof and at least two, three or four polypeptides selected from group c) or variants thereof. This preferred composition may comprise two or three polypeptides selected from group b) or variants thereof, preferably including Ber01 or a variant thereof.

**[0035]** All the above compositions based on a) to c) may comprise further polypeptides selected from any of SEQ ID NOS 1 to 74 or variants thereof not previously selected according to a) to c).

**[0036]** The invention also provides a composition suitable for use in preventing or treating allergy to grass pollen by tolerisation comprising at least four different polypeptides selected from:

50 (a) Tim07B (KIPAGELQIIDKIDA) or a variant thereof;

(b) Ber01 (SGKAFGAMAKKGQED) or a variant thereof;

(c) Bio04A (LKKAVTAMSEAK) or a variant thereof;

(d) Rye09B (PEVKYAVFEAALTKAIT) or a variant thereof;

(e) Ber02 (FIPMKSSWGA) or a variant thereof;

55 (f) Ber02C (KSSWGAIWRIDPKKPLK) or a variant thereof;

(g) Bio03A (KFIPTLVAAVKQAYAAKQ) or a variant thereof; and

(h) Bio02A (KYDAYVATLTEALR) or a variant thereof;

wherein said variant is:

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a) to (h), or
- 5 II) a polypeptide 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a) to (h), which sequence is capable of tolerising to said corresponding polypeptide; or
- 10 III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a) to (h), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

**[0037]** Other preferred variants of a) to h) are:

- 15 i) polypeptides having a length of 9 to 30 amino acids and comprising a region consisting of:
  - any of the corresponding sequences of a) to h), or
  - a sequence which has at least 65% homology to any of the corresponding sequences of a) to h) which sequence is capable of tolerising an individual to any of the sequences of a) to h), or
- 20 ii) polypeptides having a length of 9 to 30 amino acids and comprising a region consisting of a sequence that represents either:
  - a fragment of any of the sequences of a) to h), or
  - a homologue of a fragment of any of the sequences of a) to h),

which sequence is capable of tolerising an individual to any of the sequences of a) to h) and has a length of at least 9 amino acids, and wherein said homologue has at least 65% homology to any 9 contiguous amino acids in any of the sequences of a) to h); wherein optionally none of the four polypeptides are variants of the same original sequence defined by any one of a) to h).

**[0038]** The composition thus comprises a minimum of four polypeptides, each of said four polypeptides being selected from polypeptides of a) to h) or variants thereof. The above composition may comprise additional polypeptides of a) to h) or a variant of any thereof, such as five, six, seven, eight or more polypeptides of a) to h) or variants thereof.

**[0039]** In some aspects, the composition may comprise a polypeptide of a) or a variant thereof, at least one polypeptide selected from b), e) or f) or a variant of any thereof, and at least one polypeptide selected from c), d), g) or h) or a variant thereof. Alternatively, the composition may preferably comprise a polypeptide of a) or a variant thereof and at least two or three polypeptides selected from b), e) or f) or a variant of any thereof. This composition may preferably include the polypeptide Ber01 or a variant thereof. In another embodiment, the composition may comprise a polypeptide of a) or a variant thereof and at least two, three or four polypeptides selected from c), d), g) or h) or a variant of any thereof. This composition may preferably include the polypeptide Bio04A or a variant thereof.

**[0040]** All such compositions may preferably comprise the polypeptide Tim07B or a variant thereof, the polypeptide Ber01 or a variant thereof, and the polypeptide Bio04A or a variant thereof.

**[0041]** The above compositions selected from a) to h) preferably comprise at least four different polypeptides of a) to h) or comprise at least four variants each corresponding to different original or baseline sequences of a) to h). Thus, the compositions are typically selected on the basis of four different epitope sequences of origin. Preferably, the composition comprises at least one epitope sequence of Bermuda origin, at least one epitope sequence of Rye Grass origin and at least one epitope sequence of Timothy origin. The relevant epitope sequences may be selected from any of SEQ ID NOS 1 to 74 or variants thereof.

**[0042]** All of the above compositions based on a) to h) may thus comprise further polypeptides selected from any of SEQ ID NOS 1 to 74 or variants thereof not previously selected according to a) to h). All selections are subject to the composition comprising at least four polypeptides selected from a) to h) or variants thereof.

**[0043]** Optionally, any composition of specific polypeptides described herein may comprise further polypeptides up to a total of fourteen unique polypeptides. The further polypeptides will typically relate to (i.e. are typically homologues and/or fragments of) the other sequences, i.e. SEQ ID NOS: 1 to 73, that are not amongst the polypeptides already selected. The further peptides are typically functional variants of one of the peptides of SEQ ID NOS: 1 to 73. The further polypeptides may be identical to any of SEQ ID NOS: 1 to 73. The composition may therefore comprise up to fourteen different polypeptides as provided in any of SEQ ID NOS: 1 to 73, subject to at least one polypeptide being selected

from each of (a) and (b) as defined above. However, the optional further polypeptides do not need to be 100% identical to any of SEQ ID NOS: 1 to 73. They are preferably at least 65% identical to at least 9 (for example at least 10, 11, 12 or 13) or more contiguous amino acids in any of SEQ ID NOS: 1 to 73, not already selected amongst the at least one polypeptide selected from each of (a) and (b). These contiguous amino acids may comprise a MHC class II epitope, for example which binds to any of the MHC molecules mentioned herein.

5 [0044] The invention also provides products and formulations comprising the polypeptides of the invention and compositions, products and vectors comprising polynucleotides capable of expressing the polypeptides of the invention for use in preventing or treating grass allergy by tolerisation. Such tolerisation will typically be to an epitope (for example a MHC class II-binding T cell epitope) present in any of SEQ ID NOS: 1 to 74.

10 *Grass species*

15 [0045] The grass species Rye (*Lolium perenne*), Timothy (*Phleum pratense*) and Bermuda (*Cynodon dactylon*) are responsible for a high proportion of grass allergy worldwide, particularly allergies associated with grass pollen, such as hayfever. Other important grass species include Velvet grass (*Holcus lanatus*), Orchard grass (*Dactylis glomerata*), Canary grass (*Phalaris canariensis*) and Meadow grass / Kentucky bluegrass (*Poa pratensis*).

20 [0046] Rye is one of the most common grasses in the world, and is widely used as a source of animal fodder. It is native to Europe, but has been introduced on all continents worldwide and is common in all temperate zones. It can flower throughout the summer, but typically between May and July in the northern hemisphere. Rye is well adapted to mild, humid temperate climates, and grows best on rather heavy, rich, moist soils. It also grows well on well-manured lighter soils with sufficient moisture. It dislikes shade and requires a well-drained soil. Rye typically grows in locations with a soil pH in the range of about 6 to about 7, though it can tolerate a range from about 4.5 to about 8.5, preferably with an annual precipitation in the range of about 21 to about 180cm.

25 [0047] Timothy is another of the most common grasses in the world and is a primary source of animal fodder. It is native and widespread in most of Europe, North Africa and northern Asia. It has been introduced and is widespread in North and South America, South Africa and Australia. Timothy grass typically grows in locations with an annual precipitation in the range of about 35 to about 180cm, an annual temperature range of about 4 to about 22°C, and a soil pH of about 4.5 to about 8. It is best adapted to a cool, humid, temperate climate, growing best on heavy, deep and moist or even wet soils. The optimum temperature for growth is 18° - 21°C varying with day/night temperatures of 15°/10°C and 21°/15°C. The Timothy flowering season is typically confined to the early summer and Timothy pollen is a major cause of grass allergy during this season. In studies, up to 21% of patients with allergic rhinitis or other symptoms have been found to be responsive to Timothy pollen.

30 [0048] Bermuda grass typically grows in locations with an annual precipitation of about 9 to about 429cm, an annual temperature range of about 5 to about 28°C, and a soil pH in the range of about 4 to about 8.5, preferably about 6 to about 7. The Bermuda flowering season is typically in late summer (August to October in the northern hemisphere). Bermuda plants produce high quantities of pollen and are therefore a major cause of hayfever. Bermuda plants have also been reported to cause contact dermatitis. Bermuda is widely grown throughout the world, predominantly in warm climates, and is typically found between the latitudes of about 30° north and about 30° south.

40 *Peptide fragments of Group 1 and Group 5 grass pollen allergens*

45 [0049] The major allergens of grass include the Group 1 grass pollen allergens and Group 5 grass pollen allergens. Proteins from different species are assigned to groups based on amino acid sequence homology. For example, Group 1 major grass pollen allergens include the Timothy protein Phl p 1 and the Rye protein Lol p 1, and the Group 5 major grass pollen allergens include the Timothy protein Phl p 5 and the Rye protein Lol p 5.

50 [0050] The present inventors have identified the regions in certain grass pollen allergen proteins which comprise MHC Class II-binding T cell epitopes. The present inventors have also shown that regions corresponding to MHC Class II-binding T cell epitopes within the major grass pollen allergens are highly conserved between representatives in a given group, see for example, Example 2. Based on this information, peptides derived from the relevant regions of a protein in a given Group are suitable for preventing or treating grass allergy by tolerisation to the grass allergens in that Group. For example, the relevant regions from, for example, Phl p 1 of Timothy or Lol p 1 of Rye are suitable for use in preventing or treating grass allergy by tolerisation to the Group 1 grass allergens.

55 [0051] The peptides of the invention are derived from the Group 1 (SEQ ID NOS: 1 to 27) and Group 5 (SEQ ID NOS: 28 to 73) grass allergens. The terms "peptide" and "polypeptide" are used interchangeably herein. The above proteins are also referred to herein as "the allergens".

Tables 2 to 4 set out the sequences of the peptides of the invention, indicating the parent protein from which each peptide derives.

[0052] Where a composition of the invention comprises four or more polypeptides selected from polypeptides of a) to

h) or variants thereof, the composition typically comprises at least one polypeptide or variant thereof (for example a functional variant) selected from each of a) SEQ ID NOS: 1 to 27; and b) SEQ ID NOS: 28 to 73. The composition may comprise at least two, at least three, at least four, at least five, at least six, or at least seven polypeptides or variants thereof (for example functional variants) selected from each of a) SEQ ID NOS: 1 to 27; and b) SEQ ID NOS: 28 to 73.

5 In some embodiments, the composition may comprise at least one polypeptide or variant thereof (for example a functional variant) selected from each of a) SEQ ID NOS: 1 to 10; and b) SEQ ID NOS: 28 to 52. The composition may comprise at least two, at least three, at least four, at least five, at least six, or at least seven polypeptides or variants thereof (for example functional variants) selected from each of a) SEQ ID NOS: 1 to 10; and b) SEQ ID NOS: 28 to 52.

10 [0053] As outlined above, preferred compositions based on the polypeptides of a) to h) or variants thereof therefore include those which comprise at least one peptide derived from Cyn d 1 or a variant thereof and at least one peptide derived from Lol p5 or a variant thereof. Particularly preferred compositions are those which comprise at least one peptide derived from Cyn d 1 or a variant thereof, at least one peptide derived from Lol p5 or a variant thereof and at least one peptide derived from Phl p 5. In some embodiments, the compositions of a) to h) may comprise two, three, four or more peptides derived from Cyn d 1 or variants thereof and/or two, three, four or more peptides derived from Lol p 5 or variants thereof. In some embodiments, the compositions may consist or consist essentially of peptides derived from Cyn d 1 and Lol p 5 or variants thereof. In other embodiments, the composition optionally does not comprise a peptide derived from Lol p1 or a variant thereof.

15 [0054] As discussed below, compositions based on Cyn d 1 and Lol p 5 peptides are of particular utility in the treatment of grass allergy.

20 [0055] Particularly preferred compositions of the invention include those which comprise, consist or consist essentially of:

- a) at least one of the polypeptides of SEQ ID NOS: 1, 2 and 5, or variants thereof as defined herein;
- b) at least one of the polypeptides of SEQ ID NO:s 28, 29, 31 and 46, or variants thereof as defined herein; and
- 25 optionally
- c) the polypeptide of SEQ ID NO: 69 or a variant thereof as defined in herein.

20 [0056] These compositions provide a particularly preferred Cyn d 1 peptide, a particularly preferred Lol p5 peptide, and optionally a particularly preferred Phl p5 peptide. More preferably, the compositions comprise at least two or all three polypeptides as defined in a) and at least two or at least three polypeptides as defined in b). Still preferably the composition includes a polypeptide as defined in c). It is particularly preferred that the composition comprises i) SEQ ID NO: 1, ii) one of SEQ ID NO: 2 and SEQ ID NO: 5; and iii) SEQ ID NO: 46.

30 [0057] Specific preferred compositions include Combinations 1 to 10 as set out in Table 7. In such compositions, one or more of the individual polypeptides of SEQ ID NO:s 1, 2, 5, 28, 29, 31, 46 and 69 can optionally be substituted for a variant thereof. In some embodiments two, three, four, five or more of the polypeptides of SEQ ID NO:s 1, 2, 5, 28, 29, 31, 46 and 69 present in Combinations 1 to 10 can be substituted for variants thereof. The number of polypeptides substituted for variants thereof typically depends on the total number of polypeptides included in the composition. For example, in combinations based on four polypeptides (i.e. Combinations 6, 8, 10), it is preferred that only one or two polypeptides are substituted for variants thereof. Alternatively, any composition comprising four or more polypeptides selected from any of SEQ ID NOS: 1, 69, 31, 46, 2, 5, 29, 28, 43, 53, 35, 27, 4, and 70 or variants thereof is also preferred.

35 [0058] A even more preferred composition comprises, consists or consists essentially of the polypeptides of SEQ ID NO:s 1, 2, 5, 28, 31, 91 and 93 or variants thereof. Other preferred compositions comprise SEQ ID NO:s 1, 2, 5, 28, 31, 91 and 93 or in place of one, two or three, four or more of these polypeptides, variants thereof. In some embodiments, the above seven peptide compositions do not comprise any further polypeptides or do not comprise any further polypeptides derived from grass allergens. In preferred embodiments, the compositions comprise SEQ ID NO:s 1, 2, 5, 28, 31, 91 and 93 or variants thereof and do not contain any further proteins with more than 20% homology to a protein of any species of grass mentioned herein. The above compositions may comprise further non-peptide constituents such as carriers or adjuvants.

40 [0059] The invention also provides a product comprising,

- 50 (a) the polypeptide Tim07B (KIPAGELQIIDKIDA), Tim10B (KYTVFETALKAITAMSE), Tim 04A (WGAIWRIDTP-DKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof as described herein;
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED) Ber02 (FIPMKSSWGA), Ber02C (KSSWGAI-WRIDPKKPLK) and Ber02B KDSDEFIPMKSSWGAIR or a variant of any thereof as described herein; and
- 55 (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTKAIT), and Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIIPSLEAAVKQAY), Rye 05C (NAGFKAAVAAAANAPPK) or a variant of any thereof as described herein, wherein each different polypeptide is for simultaneous, separate or sequential use in preventing or treating allergy to grass pollen by tolerisation.

**[0060]** The four or more polypeptides in the product of the invention may be selected according to the same criteria outlined above in relation to the composition for use in the prevention or treatment of grass allergy by tolerisation of the invention.

5 *Variants*

**[0061]** The composition or products of the invention may therefore comprise variants of any of SEQ ID NOS: 1 to 73. The variant typically comprises 1, 2, 3 or more of the MHC class II epitopes present in the corresponding peptide of SEQ ID NOS: 1 to 73.

10 **[0062]** Functional variants are mentioned herein. Such a variant may be able to tolerise an individual to a class II MHC epitope present in the corresponding peptide of SEQ ID NOS: 1 to 59, and thus it will typically comprise sequence that binds to the same MHC class II molecule and/or is recognised by a T cell which recognises the corresponding epitope in the polypeptide of SEQ ID NOS: 1 to 73. The sequence is thus capable of tolerising against the corresponding sequence in the native polypeptide. The variant polypeptide may thus tolerise against the corresponding native polypeptide.

15 **[0063]** Variants of SEQ ID NOS: 1 to 73 may be fragments derived by truncation, e.g. by removal of one or more amino acids from the N and/or C-terminal ends of a polypeptide. Fragments may also be generated by one or more internal deletions, provided that the core 9 amino acids that makes up the T cell epitope is not substantially disrupted.

20 **[0064]** For example, a variant of SEQ ID NO: 1 may comprise a fragment of SEQ ID NO: 1, i.e. a shorter sequence. This may include a deletion of one, two, three or four amino acids from the N-terminal end of SEQ ID NO: 1 or from the 25 C-terminal end of SEQ ID NO: 1. Such deletions may be made from both ends of SEQ ID NO: 1. A variant of SEQ ID NO: 1 may include additional amino acids (for example from the sequence of the parent protein from which the peptide derives) extending beyond the end(s) of SEQ ID NO: 1. A variant may include a combination of the deletions and additions discussed above. For example, amino acids may be deleted from one end of SEQ ID NO: 1, but additional amino acids from the full length parent protein sequence may be added at the other end of SEQ ID NO: 1. The same discussion of variants above also applies to SEQ ID NOS: 2 to 73.

30 **[0065]** A variant peptide may include one or more amino acid substitutions from the amino acid sequence of any of SEQ ID NOS: 1 to 73 or a fragment thereof. A variant peptide may comprise sequence having at least 65% sequence identity to at least 9 or more contiguous amino acids in any of SEQ ID NOS: 1 to 73. More preferably a suitable variant may comprise at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% amino acid identity to at least 9 contiguous amino acids of any of SEQ ID NOS: 1 to 73. This level of amino acid identity may be seen at any section of the peptide, although it is preferably the core region. The level of amino acid identity is over at least 9 contiguous amino acids but it may be at least 10, 11, 12, 13, 14, 15 or at least 16 or 17 amino acids, depending on the size of the peptides of comparison. Accordingly, any of the above-specified levels of identity may be across the entire length of sequence.

35 **[0066]** Examples of preferred variants selected on the basis of sequence identity to particularly preferred peptides of the invention are:

Variants of Ber01 (SEQ ID NO: 1): SGKAFGAMAKKGEED (SEQ ID NO: 74), Holcus lanatus, Hol 11; Poa pratensis Poa p 1; SGHAFGSMAKKGEED (SEQ ID NO: 75), Dactylis glomerata, Dac g 1; Lolium perenne Lol p 1; SGIAFGS-40 MAKKGDED (SEQ ID NO: 76), Phleum pratense, Phl p 1, SGTAFGAMAKKGEEE (SEQ ID NO: 77), Zea mays, Zea m 1.

Variants of Bio02A (SEQ ID NO: 28): KYDAYVATLTESLR (SEQ ID NO: 78), Hordeum vulgare, Hor v 5; KYDAY-VATLSEALR (SEQ ID NO: 79), Phleum pratense, Phl p 5; Poa pratensis, Poa p 9; KYDAFVAALTEALR (SEQ ID NO: 80), Dactylis glomerata, Dac g 5; KYDAFVTTLTEALR (SEQ ID NO: 81), Holcus lanatus, Hol 15.

45 Variants of Bio03A (SEQ ID NO: 29): KFIPITLEAAVKQAYAA (SEQ ID NO: 82), Dactylis glomerata, Dac g 5; Phalaris aquatica, Pha a 5; KFIPALEAAVKQAYAA (SEQ ID NO: 83), Phleum pratense, Phl p 5.

Variants of Tim07B (SEQ ID NO: 69): KIPAGELQIVDKIDA (SEQ ID NO: 84), Dactylis glomerata, Dac g 5; Phalaris aquatica, Pha a 5; KIPTGELQIVDKIDA (SEQ ID NO: 85), Lolium perenne, Lol p 5; KIPAGEQQIIDKIDA (SEQ ID NO: 86), Poa pratensis, Poa p 5. Tim07B is also conserved in Holcus lanatus, Hol 15b.

50 Variants of Bio04A (SEQ ID NO: 31): LKKAITAMSEAQK (SEQ ID NO: 87), Phleum pratense, Phl p 5; Holcus lanatus, Hol 15b; LKKAITAMSQAQK (SEQ ID NO: 88), Poa pratensis, Poa p 5.

Variants of Ber02C (SEQ ID NO: 5): KESWGAIWRIDTPDK (SEQ ID NO: 89), Phalaris aquatica, Pha a 1; Phleum pratense, Phl p 1.

Variants of Rye09B (SEQ ID NO: 46): PQVKYAVFEAALTKAIT (SEQ ID NO: 90), Phleum pratense, Phl p 5.

55

**[0067]** Thus, SEQ ID NO:s 74 to 90 are preferred variants which may be used as substitutes for the above native peptides in a composition of the invention. Other referred variants for native peptides of the invention are derived from maize allergens. Maize allergens have close homology to grass pollen allergens and so sequences derived from maize

may contain cross-reactive epitopes of use in tolerisation against grass pollen allergy.

[0068] In connection with amino acid sequences, "sequence identity" refers to sequences which have the stated value when assessed using ClustalW (Thompson et al., 1994, *supra*) with the following parameters:

5 Pairwise alignment parameters -Method: accurate, Matrix: PAM, Gap open penalty: 10.00, Gap extension penalty: 0.10; Multiple alignment parameters -Matrix: PAM, Gap open penalty: 10.00, % identity for delay: 30, Penalize end gaps: on, Gap separation distance: 0, Negative matrix: no, Gap extension penalty: 0.20, Residue-specific gap penalties: on, Hydrophilic gap penalties: on, Hydrophilic residues: GPSNDQEKR. Sequence identity at a particular residue is intended to include identical residues which have simply been derivatized.

10 [0069] A variant peptide may comprise 1, 2, 3, 4, 5 or more, or up to 10 amino acid substitutions from any of SEQ ID NOS: 1 to 73. Substitution variants preferably involve the replacement of one or more amino acids with the same number of amino acids and making conservative amino acid substitutions. For example, an amino acid may be substituted with an alternative amino acid having similar properties, for example, another basic amino acid, another acidic amino acid, another neutral amino acid, another charged amino acid, another hydrophilic amino acid, another hydrophobic amino acid, another polar amino acid, another aromatic amino acid or another aliphatic amino acid. Some properties of the 20 main amino acids which can be used to select suitable substituents are as follows:

20	Ala	aliphatic, hydrophobic, neutral	Met	hydrophobic, neutral
	Cys	polar, hydrophobic, neutral	Asn	polar, hydrophilic, neutral
	Asp	polar, hydrophilic, charged (-)	Pro	hydrophobic, neutral
	Glu	polar, hydrophilic, charged (-)	Gln	polar, hydrophilic, neutral
25	Phe	aromatic, hydrophobic, neutral	Arg	polar, hydrophilic, charged (+)
	Gly	aliphatic, neutral	Ser	polar, hydrophilic, neutral
	His	aromatic, polar, hydrophilic, charged (+)	Thr	polar, hydrophilic, neutral
30	Ile	aliphatic, hydrophobic, neutral	Val	aliphatic, hydrophobic, neutral
	Lys	polar, hydrophilic, charged(+)	Trp	aromatic, hydrophobic, neutral
	Leu	aliphatic, hydrophobic, neutral	Tyr	aromatic, polar, hydrophobic

35 [0070] Further variants include those in which instead of the naturally occurring amino acid the amino acid which appears in the sequence is a structural analog thereof. Amino acids used in the sequences may also be modified, e.g. labelled, providing the function of the peptide is not significantly adversely affected. Where the peptide has a sequence that varies from the sequence of any of SEQ ID NOS: 1 to 73 or a fragment thereof, the substitutions may occur across the full length of the sequence, within the sequence of any of SEQ ID NOS: 1 to 73 or outside the sequence of any of SEQ ID NOS: 1 to 73. For example, the variations described herein, such as additions, deletions, substitutions and modifications, may occur within the sequence of any of SEQ ID NOS: 1 to 73. A variant peptide may comprise or consist essentially of the amino acid sequence of any of SEQ ID NOS: 1 to 73 in which one, two, three, four or more amino acid substitutions have been made. A variant peptide may comprise a fragment of the parent protein that is larger than any of SEQ ID NOS: 1 to 73. In this embodiment, the variations described herein, such as substitutions and modifications, may occur within and/or outside the sequence of any of SEQ ID NOS: 1 to 73.

40 [0071] The variant peptides of the invention are 9 to 30 amino acids in length inclusive. Preferably, they may be from 9 to 20 or more preferably 13 to 17 amino acids in length. The peptides may be the same length as the peptide sequences in any one of SEQ ID NOS: 1 to 73.

45 [0072] The peptides may be chemically derived from the polypeptide allergen, for example by proteolytic cleavage or can be derived in an intellectual sense from the polypeptide allergen, for example by making use of the amino acid sequence of the polypeptide allergen and synthesising peptides based on the sequence. Peptides may be synthesised using methods well known in the art.

50 [0073] The term "peptide" includes not only molecules in which amino acid residues are joined by peptide (-CO-NH-) linkages but also molecules in which the peptide bond is reversed. Such retro-inverso peptidomimetics may be made using methods known in the art, for example such as those described in Meziere et al (1997) *J. Immunol.* 159, 3230-3237. This approach involves making pseudopeptides containing changes involving the backbone, and not the orientation of side chains. Meziere et al (1997) show that, at least for MHC class II and T helper cell responses, these pseudopeptides are useful. Retro-inverse peptides, which contain NH-CO bonds instead of CO-NH peptide bonds, are much more

resistant to proteolysis.

**[0074]** Similarly, the peptide bond may be dispensed with altogether provided that an appropriate linker moiety which retains the spacing between the carbon atoms of the amino acid residues is used; it is particularly preferred if the linker moiety has substantially the same charge distribution and substantially the same planarity as a peptide bond. It will also be appreciated that the peptide may conveniently be blocked at its N- or C-terminus so as to help reduce susceptibility to exoproteolytic digestion. For example, the N-terminal amino group of the peptides may be protected by reacting with a carboxylic acid and the C-terminal carboxyl group of the peptide may be protected by reacting with an amine. Other examples of modifications include glycosylation and phosphorylation. Another potential modification is that hydrogens on the side chain amines of R or K may be replaced with methylene groups (-NH<sub>2</sub> → -NH(Me) or -N(Me)<sub>2</sub>).

**[0075]** Analogues of peptides according to the invention may also include peptide variants that increase or decrease the peptide's half-life *in vivo*. Examples of analogues capable of increasing the half-life of peptides used according to the invention include peptoid analogues of the peptides, D-amino acid derivatives of the peptides, and peptide-peptoid hybrids. A further embodiment of the variant polypeptides used according to the invention comprises D-amino acid forms of the polypeptide. The preparation of polypeptides using D-amino acids rather than L-amino acids greatly decreases any unwanted breakdown of such an agent by normal metabolic processes, decreasing the amounts of agent which needs to be administered, along with the frequency of its administration.

**[0076]** The peptides provided by the present invention may be derived from splice variants of the parent proteins encoded by mRNA generated by alternative splicing of the primary transcripts encoding the parent protein chains. The peptides may also be derived from amino acid mutants, glycosylation variants and other covalent derivatives of the parent proteins which retain at least an MHC-binding property of the allergens. Exemplary derivatives include molecules wherein the peptides of the invention are covalently modified by substitution, chemical, enzymatic, or other appropriate means with a moiety other than a naturally occurring amino acid. Further included are naturally occurring variants of the parent proteins found in different mites. Such a variant may be encoded by an allelic variant or represent an alternative splicing variant.

**[0077]** Variants as described above may be prepared during synthesis of the peptide or by post-production modification, or when the peptide is in recombinant form using the known techniques of site- directed mutagenesis, random mutagenesis, or enzymatic cleavage and/or ligation of nucleic acids.

**[0078]** In accordance with the invention, the further one or more peptides that the composition may comprise are preferably functional variants of any of SEQ ID NOS: 1 to 73. That is, the peptides are preferably capable of inducing an immune response. In particular, they may be capable of inducing a late phase response in an individual with grass allergy. This may be tested by the ability of the peptide to induce T cell proliferation in a sample of T cells. Methods of testing the induction of T cell proliferation are well known in the art and one such method is exemplified in Example 4. Preferably the one or more further peptides are capable of causing T cell proliferation in at least 20 % of samples of T cells, wherein each sample is obtained from different grass allergic individuals in the population. The compositions of the invention are preferably capable of inducing T cell proliferation in 30 % or more samples of T cells obtained from a panel of grass allergic individuals. More preferably, the compositions are capable of inducing T cell proliferation in 35% or more, 40 % or more, 45 %, 50 %, 55 %, 60 %, 65 %, 70 %, 75 %, 80 %, 85 %, or 90 % or more of samples obtained from sensitized individuals in a panel.

**[0079]** The compositions may also be capable of inducing a significant release of one or more cytokines, preferably including one or more of Interferon gamma, Interleukin-10 and Interleukin-13 in samples of T cells obtained from sensitized individuals in a panel. A "significant" release may be determined by criteria similar to that described below in Example 5. Thus, the compositions may be capable of inducing a release of one or more cytokines in 35% or more, 40 % or more, 45 %, 50 %, 55 %, 60 %, 65 %, 70 %, 75 %, 80 %, 85 %, or 90 % or more of samples obtained from sensitized individuals in a panel. The number of individuals in a panel of grass allergic individuals may be any number greater than one, for example at least 2, 3, 5, 10, 15, 20, 30, 50, 80, or at least 100 individuals.

**[0080]** It is preferred if the peptides cause T cell proliferation, but do not lead to the release of histamine from basophils or mast cell preparations from a sensitised individual. There may be some histamine release, but preferably the composition does not cause significant amounts of histamine to be released. Significant histamine release may be considered to be the release of 20% or more of the total available leukocyte histamine when a sample of leukocytes from an individual is stimulated with a composition *in vitro*. A normal individual typically has an approximate leukocyte histamine content of 150ng/10<sup>7</sup> cells.

**[0081]** Suitable variants capable of binding to TCRs may be derived empirically or selected according to known criteria. Within a single peptide there are certain residues which contribute to binding within the MHC antigen binding groove and other residues which interact with hypervariable regions of the T cell receptor (Allen et al (1987) *Nature* 327: 713-5).

**[0082]** Within the residues contributing to T cell receptor interaction, a hierarchy has been demonstrated which pertains to dependency of T cell activation upon substitution of a given peptide residue. Using peptides which have had one or more T cell receptor contact residues substituted with a different amino acid, several groups have demonstrated profound effects upon the process of T cell activation. Evavold & Allen (1991) *Nature* 252: 1308-10) demonstrated the dissociation

of T cell proliferation and cytokine production. In this *in vitro* model, a T cell clone specific for residues 64-76 of haemoglobin (in the context of I-E<sup>k</sup>), was challenged with a peptide analogue in which a conservative substitution of aspartic acid for glutamic acid had been made. This substitution did not significantly interfere with the capacity of the analogue to bind to I-E<sup>k</sup>.

5 [0083] Following *in vitro* challenge of a T cell clone with this analogue, no proliferation was detected although IL-4 secretion was maintained, as was the capacity of the clone to help B cell responses. In a subsequent study the same group demonstrated the separation of T cell-mediated cytotoxicity from cytokine production. In this instance, the former remained unaltered while the latter was impaired. The efficacy of altered peptide ligands *in vivo* was initially demonstrated in a murine model of EAE (experimental allergic encephalomyelitis) by McDevitt and colleagues (Smilek et al (1991) 10 Proc Natl Acad Sci USA 88 : 9633-9637). In this model EAE is induced by immunisation with the encephalitogenic peptide Ac1-11 of MBP (myelin basic protein). Substitution at position four (lysine) with an alanine residue generated a peptide which bound well to its restricting element (A $\alpha^u$ A $\beta^u$ ), but which was non-immunogenic in the susceptible PL/JxSJLF1 strain and which, furthermore prevented the onset of EAE when administered either before or after immunisation with the encephalitogenic peptide. Thus, residues can be identified in peptides which affect the ability of the peptides to induce various functions of T-cells.

15 [0084] Advantageously, peptides may be designed to favour T-cell proliferation and induction of desensitisation. Metzler and Wraith have demonstrated improved tolerogenic capacity of peptides in which substitutions increasing peptide-MHC affinity have been made (Metzler & Wraith(1993) Int Immunol ~ : 1159-65). That an altered peptide ligand can cause long-term and profound anergy in cloned T cells was demonstrated by Sloan-Lancaster et al (1993) Nature 363: 156-9.

20 [0085] The compositions of the invention may be capable of inducing a late phase response in an individual that is sensitised to the allergens. The term "late phase response" includes the meaning as set forth in Allergy and Allergic Diseases (1997) A. B. Kay (Ed.), Blackwell Science, pp 1113-1130. The late phase response may be any late phase response (LPR). Preferably, the peptides may be capable of inducing a late asthmatic response (LAR) or a late rhinitic response, or a late phase skin response or a late phase ocular response. Whether or not a particular peptide can give 25 rise to a LPR can be determined using methods well known in the art; a particularly preferred method is that described in Cromwell O, Durham SR, Shaw RJ, Mackay J and Kay AB. Provocation tests and measurements of mediators from mast cells and basophils in asthma and allergic rhinitis. In: Handbook of Experimental Immunology (4) Chapter 127, Editor: Weir DM, Blackwell Scientific Publications, 1986.

30 [0086] Thus, preferably, the individual peptides of the invention are able to induce a LPR in an individual who has been sensitised to the allergens. Whether or not an individual has been sensitised to the allergens may be determined by well known procedures such as skin prick testing with solutions of allergen extracts, induction of cutaneous LPRs, clinical history, allergen challenge and radioallergosorbent test (RAST) for measurement of allergen specific IgE. Whether or not a particular individual is expected to benefit from treatment may be determined by the physician based, for example, on such tests.

35 [0087] Desensitising or tolerising an individual to the allergens means inhibition or dampening of allergic tissue reactions induced by the allergens in appropriately sensitised individuals. It has been shown that T cells can be selectively activated, and then rendered unresponsive. Moreover the anergising or elimination of these T-cells leads to desensitisation of the patient for a particular allergen. The desensitisation manifests itself as a reduction in response to an allergen or allergen-derived peptide, or preferably an elimination of such a response, on second and further administrations of the allergen 40 or allergen-derived peptide. The second administration may be made after a suitable period of time has elapsed to allow desensitisation to occur; this is preferably any period between one day and several weeks. An interval of around two weeks is preferred.

45 [0088] Although the compositions of the invention may be able to induce a LPR in a grass allergic individual, it should be appreciated that when a composition is used to treat a patient it is preferable that a sufficiently low concentration of the composition is used such that no observable LPR will occur but the response will be sufficient to partially desensitise the T cells such that the next (preferably higher) dose may be given, and so on. In this way the dose is built up to give full desensitisation but often without ever inducing a LPR in the patient. Although, the composition or peptide is able to do so at a higher concentration than is administered.

50 [0089] The compositions of the invention may be capable of inducing a late phase response in 50 % or more of a panel of grass allergic individuals from the population. More preferably, the compositions are capable of inducing a LPR in 55% or more, 60 % or more, 65 % or more, 70% or more, 75% or more, 80% or more, 85% or more, or 90 % or more of sensitized individuals in a panel. Whether or not the compositions are able to induce a LPR in a certain percentage of a panel of subjects can be determined by methods which are well known in the art.

55 [0090] It will be understood that the peptides of the invention comprise a T cell epitope that consists of a core 9 amino acids which are the minimal essential sequence required for MHC class II binding. However, the peptides may also comprise additional residues flanking the core 9 amino acids. The peptides may therefore comprise a region containing a T cell epitope, in which some residues may be modified without affecting the function of the epitope. Accordingly, functional variants of the peptides as defined above include peptides which are altered to improve their solubility relative

to the native sequence of the peptides. Improved solubility is advantageous for the tolerisation of subjects to allergens from which the peptides of the invention derive, since administration of poorly soluble agents to subjects causes undesirable, non-tolerising inflammatory responses. The solubility of the peptides may be improved by altering the residues which flank the region containing a T cell epitope. A peptide of the invention may be engineered to be more soluble such that it comprises:

- 5 i) N terminal to the residues of the peptide which flank a T cell epitope: one to six contiguous amino acids corresponding to the two to six contiguous amino acids immediately N terminal to said residues in the sequence of the protein from which the peptide derives; and/or
- 10 ii) C terminal to the residues of the peptide which flank a T cell epitope: one to six contiguous amino acids corresponding to the one to six contiguous amino acids immediately C terminal to the said residues in the sequence of the protein from which the peptide derives; or
- 15 iii) both N and C terminal to the residues of the peptide which flank a T cell epitope, at least one amino acid selected from arginine, lysine, histidine, glutamate and aspartate.

[0091] Optionally, the peptides may additionally be engineered to be more soluble such that:

- 20 i) any cysteine residues in the native sequence of the peptide are replaced with serine or 2-aminobutyric acid; and /or
- ii) any hydrophobic residues in the upto three amino acids at the N or C terminus of the native sequence of the peptide, which are not comprised in a T cell epitope, are deleted; and/or
- iii) any two consecutive amino acids comprising the sequence Asp-Gly in the upto four amino acids at the N or C terminus of the native sequence of the peptide, which are not comprised in a T cell epitope, are deleted; and/or
- iv) one or more positively charged residues are added at the N- and/or C-terminus.

[0092] Examples of peptides engineered to improve solubility are Rye 09B and Tim07B as described in Example 8. The variants with improved solubility are shown in Table 8. Thus, peptides Rye 09B1 (SEQ ID NO: 91), KPE-VKYAVFEAALTKAIT; Rye 09B2 (SEQ ID NO: 92), KKPEVKYAVFEAALTKAIT, Tim 07B1, (SEQ ID NO: 93), KKIPAGEL-QIIDKIDA, Tim 07B2 (SEQ ID NO: 94), KKIPAGELQIIDKIDAK are preferred examples of variants with improved solubility. SEQ ID NO:s 91 to 94 may therefore preferentially be used as substitutes for the above native peptides in a composition of the invention.

#### *Nucleic acids and vectors*

[0093] The individual peptides that make up the compositions and products of the invention may be administered directly, or may be administered indirectly by expression from an encoding sequence. For example, a polynucleotide may be provided that encodes a peptide combination of the invention, such as any combination of peptides described above. A peptide combination of the invention may thus be produced from or delivered in the form of a polynucleotide which encodes, and is capable of expressing, it. Any reference herein to the use, delivery or administration of a peptide combination of the invention is intended to include the indirect use, delivery or administration of such a peptide combination via expression from a polynucleotide that encodes it.

[0094] Accordingly, the invention provides a composition for use in preventing or treating allergy to grass by tolerisation comprising at least one polynucleotide sequence which when expressed causes the production of a composition suitable for use in preventing or treating allergy to grass by tolerisation as described above.

[0095] The invention also provides a product comprising four or more different polynucleotides encoding polypeptides of a) to h) or variants thereof as described above; and wherein each different encoded polypeptide is for simultaneous, separate or sequential use in the prevention or treatment of allergy to grass in a human.

[0096] The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably herein and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Non-limiting examples of polynucleotides include a gene, a gene fragment, messenger RNA (mRNA), cDNA, recombinant polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers. A polynucleotide of the invention may be provided in isolated or purified form.

[0097] A nucleic acid sequence which "encodes" a selected polypeptide is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. For the purposes of the invention, such nucleic acid sequences can include, but are not limited to, cDNA from viral, prokaryotic or eukaryotic mRNA, genomic sequences from viral or prokaryotic DNA or RNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence.

[0098] Polynucleotides of the invention can be synthesised according to methods well known in the art, as described by way of example in Sambrook et al (1989, Molecular Cloning - a laboratory manual; Cold Spring Harbor Press).

[0099] The polynucleotide molecules of the present invention may be provided in the form of an expression cassette which includes control sequences operably linked to the inserted sequence, thus allowing for expression of the peptide of the invention *in vivo* in a targeted subject. These expression cassettes, in turn, are typically provided within vectors (e.g., plasmids or recombinant viral vectors) which are suitable for use as reagents for nucleic acid immunization. Such an expression cassette may be administered directly to a host subject. Alternatively, a vector comprising a polynucleotide of the invention may be administered to a host subject. Preferably the polynucleotide is prepared and/or administered using a genetic vector. A suitable vector may be any vector which is capable of carrying a sufficient amount of genetic information, and allowing expression of a peptide of the invention.

[0100] The present invention thus includes expression vectors that comprise such polynucleotide sequences. Thus, the present invention provides a vector for use in preventing or treating allergy to grass by tolerisation comprising four or more polynucleotide sequences which encode different polypeptides of the invention and optionally one or more further polynucleotide sequences which encode different polypeptides as defined herein. The vector may comprise 4, 5, 6 or 7 polynucleotide sequences which encode different polypeptides of the invention. The vector preferably comprises a first polynucleotide sequence that encodes a polypeptide selected from SEQ ID NO:s 1 to 27 or a variant thereof as described above, and a second polynucleotide sequence that encodes a polypeptide selected from SEQ ID NO:s 28 to 73 or a variant thereof as described above. This (first) vector may be used in combination with one or more other vectors providing coding sequences for different polypeptides of the invention not encoded by the first vector.

[0101] Furthermore, it will be appreciated that the compositions and products of the invention may comprise a mixture of polypeptides and polynucleotides. Accordingly, the invention provides a composition or product as defined herein, wherein in place of any one of the polypeptide is a polynucleotide capable of expressing said polypeptide.

[0102] Expression vectors are routinely constructed in the art of molecular biology and may for example involve the use of plasmid DNA and appropriate initiators, promoters, enhancers and other elements, such as for example polyadenylation signals which may be necessary, and which are positioned in the correct orientation, in order to allow for expression of a peptide of the invention. Other suitable vectors would be apparent to persons skilled in the art. By way of further example in this regard we refer to Sambrook et al.

[0103] Thus, a polypeptide of the invention may be provided by delivering such a vector to a cell and allowing transcription from the vector to occur. Preferably, a polynucleotide of the invention or for use in the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector.

[0104] "Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given regulatory sequence, such as a promoter, operably linked to a nucleic acid sequence is capable of effecting the expression of that sequence when the proper enzymes are present. The promoter need not be contiguous with the sequence, so long as it functions to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the nucleic acid sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

[0105] A number of expression systems have been described in the art, each of which typically consists of a vector containing a gene or nucleotide sequence of interest operably linked to expression control sequences. These control sequences include transcriptional promoter sequences and transcriptional start and termination sequences. The vectors of the invention may be for example, plasmid, virus or phage vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. A "plasmid" is a vector in the form of an extrachromosomal genetic element. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a resistance gene for a fungal vector. Vectors may be used *in vitro*, for example for the production of DNA or RNA or used to transfect or transform a host cell, for example, a mammalian host cell. The vectors may also be adapted to be used *in vivo*, for example to allow *in vivo* expression of the polypeptide.

[0106] A "promoter" is a nucleotide sequence which initiates and regulates transcription of a polypeptide-encoding polynucleotide. Promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is repressed by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters. It is intended that the term "promoter" or "control element" includes full-length promoter regions and functional (e.g., controls transcription or translation) segments of these regions.

[0107] A polynucleotide, expression cassette or vector according to the present invention may additionally comprise a signal peptide sequence. The signal peptide sequence is generally inserted in operable linkage with the promoter such that the signal peptide is expressed and facilitates secretion of a polypeptide encoded by coding sequence also in operable linkage with the promoter.

[0108] Typically a signal peptide sequence encodes a peptide of 10 to 30 amino acids for example 15 to 20 amino

acids. Often the amino acids are predominantly hydrophobic. In a typical situation, a signal peptide targets a growing polypeptide chain bearing the signal peptide to the endoplasmic reticulum of the expressing cell. The signal peptide is cleaved off in the endoplasmic reticulum, allowing for secretion of the polypeptide via the Golgi apparatus. Thus, a peptide of the invention may be provided to an individual by expression from cells within the individual, and secretion from those cells.

**[0109]** Alternatively, polynucleotides of the invention may be expressed in a suitable manner to allow presentation of a peptide of the invention by an MHC class II molecule at the surface of an antigen presenting cell. For example, a polynucleotide, expression cassette or vector of the invention may be targeted to antigen presenting cells, or the expression of encoded peptide may be preferentially stimulated or induced in such cells.

**[0110]** In some embodiments, the polynucleotide, expression cassette or vector will encode an adjuvant, or an adjuvant will otherwise be provided. As used herein, the term "adjuvant" refers to any material or composition capable of specifically or non-specifically altering, enhancing, directing, redirecting, potentiating or initiating an antigen-specific immune response.

**[0111]** Polynucleotides of interest may be used *in vitro*, *ex vivo* or *in vivo* in the production of a peptide of the invention. Such polynucleotides may be administered or used in the prevention or treatment of allergy by tolerisation.

**[0112]** Methods for gene delivery are known in the art. See, e.g., U.S. Patent Nos. 5,399,346, 5,580,859 and 5,589,466. The nucleic acid molecule can be introduced directly into the recipient subject, such as by standard intramuscular or intradermal injection; transdermal particle delivery; inhalation; topically, or by oral, intranasal or mucosal modes of administration. The molecule alternatively can be introduced *ex vivo* into cells that have been removed from a subject.

For example, a polynucleotide, expression cassette or vector of the invention may be introduced into APCs of an individual *ex vivo*. Cells containing the nucleic acid molecule of interest are re-introduced into the subject such that an immune response can be mounted against the peptide encoded by the nucleic acid molecule. The nucleic acid molecules used in such immunization are generally referred to herein as "nucleic acid vaccines."

**[0113]** The polypeptides, polynucleotides, vectors or cells of the invention may be present in a substantially isolated form. They may be mixed with carriers or diluents which will not interfere with their intended use and still be regarded as substantially isolated. They may also be in a substantially purified form, in which case they will generally comprise at least 90%, e.g. at least 95%, 98% or 99%, of the proteins, polynucleotides, cells or dry mass of the preparation.

#### *Antigen presenting cells (APCs)*

**[0114]** The invention encompasses the use *in vitro* of a method of producing a population of APCs that present the peptides of the invention on their surface, that may be subsequently used in therapy. Such a method may be carried out *ex vivo* on a sample of cells that have been obtained from a patient. The APCs produced in this way therefore form a pharmaceutical agent that can be used in the treatment or prevention of grass allergy by tolerisation. The cells should be accepted by the immune system of the individual because they derive from that individual. Delivery of cells that have been produced in this way to the individual from whom they were originally obtained, thus forms a therapeutic embodiment of the invention.

#### *Formulations and compositions*

**[0115]** The peptides, polynucleotides, vectors and cells of the invention may be provided to an individual either singly or in combination. Each molecule or cell of the invention may be provided to an individual in an isolated, substantially isolated, purified or substantially purified form. For example, a peptide of the invention may be provided to an individual substantially free from the other peptides. Alternatively, four or more peptides in the composition may be coupled chemically together, using standard peptide coupling reagents, to provide a single peptide containing the preferred epitopes. Such peptides would be screened for basophil histamine release to confirm lack of histamine release as per the individual peptides. In a further embodiment, four or more peptides in the composition may be provided as part of a single polypeptide chain i.e by recombinant means from an encoding polynucleotide. The four or more peptides may be fused contiguously, or may alternatively be separated by appropriate linkers.

**[0116]** Whilst it may be possible for the peptides, polynucleotides or compositions according to the invention to be presented in raw form, it is preferable to present them as a pharmaceutical formulation. Thus, according to a further aspect of the invention, the present invention provides a pharmaceutical formulation for use in preventing or treating allergy to grass by tolerisation comprising a composition, vector or product according to the invention together with one or more pharmaceutically acceptable carriers or diluents and optionally one or more other therapeutic ingredients. The carrier (s) must be 'acceptable' in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Typically, carriers for injection, and the final formulation, are sterile and pyrogen free. Preferably, the carrier or diluent is thioglycerol.

**[0117]** Formulation of a composition comprising the peptide, polynucleotides or cells of the invention can be carried

out using standard pharmaceutical formulation chemistries and methodologies all of which are readily available to the reasonably skilled artisan.

[0118] For example, compositions containing one or more molecules or cells of the invention can be combined with one or more pharmaceutically acceptable excipients or vehicles. Auxiliary substances, such as wetting or emulsifying agents, pH buffering substances and the like, may be present in the excipient or vehicle. These excipients, vehicles and auxiliary substances are generally pharmaceutical agents that do not induce an immune response in the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, polyethyleneglycol, hyaluronic acid, glycerol, thioglycerol and ethanol. Pharmaceutically acceptable salts can also be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable excipients, vehicles and auxiliary substances is available in Remington's Pharmaceutical Sciences (Mack Pub. Co., N.J. 1991).

[0119] Such compositions may be prepared, packaged, or sold in a form suitable for bolus administration or for continuous administration. Injectable compositions may be prepared, packaged, or sold in unit dosage form, such as in ampoules or in multidose containers containing a preservative. Compositions include, but are not limited to, suspensions, solutions, emulsions in oily or aqueous vehicles, pastes, and implantable sustained-release or biodegradable formulations. Such compositions may further comprise one or more additional ingredients including, but not limited to, suspending, stabilizing, or dispersing agents. In one embodiment of a composition for parenteral administration, the active ingredient is provided in dry (for e.g., a powder or granules) form for reconstitution with a suitable vehicle (e. g., sterile pyrogen-free water) prior to parenteral administration of the reconstituted composition. The pharmaceutical compositions may be prepared, packaged, or sold in the form of a sterile injectable aqueous or oily suspension or solution. This suspension or solution may be formulated according to the known art, and may comprise, in addition to the active ingredient, additional ingredients such as the dispersing agents, wetting agents, or suspending agents described herein. Such sterile injectable formulations may be prepared using a non-toxic parenterally-acceptable diluent or solvent, such as water or 1,3-butane diol, for example. Other acceptable diluents and solvents include, but are not limited to, Ringer's solution, isotonic sodium chloride solution, and fixed oils such as synthetic mono- or di-glycerides. Other parentally-administrable compositions which are useful include those which comprise the active ingredient in microcrystalline form, in a liposomal preparation, or as a component of a biodegradable polymer systems. Compositions for sustained release or implantation may comprise pharmaceutically acceptable polymeric or hydrophobic materials such as an emulsion, an ion exchange resin, a sparingly soluble polymer, or a sparingly soluble salt.

[0120] Alternatively, the peptides or polynucleotides of the present invention may be encapsulated, adsorbed to, or associated with, particulate carriers. Suitable particulate carriers include those derived from polymethyl methacrylate polymers, as well as PLG microparticles derived from poly(lactides) and poly(lactide-co-glycolides). See, e.g., Jeffery et al. (1993) *Pharm. Res.* 10:362-368. Other particulate systems and polymers can also be used, for example, polymers such as polylysine, polyarginine, polyornithine, spermine, spermidine, as well as conjugates of these molecules.

[0121] The formulation of any of the peptides, polynucleotides or cells mentioned herein will depend upon factors such as the nature of the substance and the method of delivery. Any such substance may be administered in a variety of dosage forms. It may be administered orally (e.g. as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules), topically, parenterally, subcutaneously, by inhalation, intravenously, intramuscularly, intraternally, transdermally, intradermally, sublingually, intranasally, buccally or by infusion techniques. The substance may also be administered as suppositories. A physician will be able to determine the required route of administration for each particular individual.

[0122] The compositions or formulations of the invention will comprise a suitable concentration of each peptide/polynucleotide/cell to be effective without causing adverse reaction. Typically, the concentration of each peptide in the composition will be in the range of 0.03 to 200 nmol/ml. More preferably in the range of 0.3 to 200 nmol/ml, 3 to 180 nmol/ml, 10 to 150 nmol/ml, 5 to 200nmol/ml or 30 to 120 nmol/ml. The composition or formulations should have a purity of greater than 95% or 98% or a purity of at least 99%.

[0123] In one aspect of the invention an adjuvant may be used in combination with the polypeptide/polynucleotides/cells of the invention. The adjuvant is preferably administered in an amount which is sufficient to augment the effect of the polypeptide/polynucleotides/cells of the invention or vice versa. The adjuvant or other therapeutic agent may be an agent that potentiates the effects of the molecule of the invention. For example, the other agent may be an immunomodulatory molecule or an adjuvant which enhances the response to the peptide or cell of the invention.

[0124] In one embodiment, therefore, the peptides, polynucleotides, cells or compositions of the invention are used for therapy in combination with one or more other therapeutic agents. The agents may be administered separately, simultaneously or sequentially. They may be administered in the same or different compositions. Accordingly, in a method of the invention, the subject may also be treated with a further therapeutic agent.

[0125] A composition may therefore be formulated which comprises a molecule and/or cell of the invention and also one or more other therapeutic molecules. A composition of the invention may alternatively be used simultaneously,

sequentially or separately with one or more other therapeutic compositions as part of a combined treatment.

**[0126]** Non-limiting examples of adjuvants include vitamin D, rapamycin and glucocorticoid steroids such as dexamethasone, fluticasone, budesonide, mometasone, beclomethasone, hydrocortisone, cortisone acetate, prednisone, prednisolone, methylprednisolone, betamethasone and triamcinolone. A preferred glucocorticoid is dexamethasone.

5

*Therapeutic methods and individual to be treated*

**[0127]** The present invention relates to peptides, polynucleotides, vectors and cells that are capable of desensitising or tolerising human individuals to the allergens described above and are therefore useful in the prevention or treatment of grass allergy. The invention provides compositions, products, vectors and formulations for use in preventing or treating allergy to grass by tolerisation. The compositions of the invention may be used to reduce allergic symptoms or improve the condition of an allergic individual. The invention also provides a method of tolerising or desensitizing a grass allergic individual comprising administering, either singly or in combination the polypeptides/polynucleotides/cells of the invention as described above.

10

**[0128]** The individual to be treated or provided with the composition or formulation of the invention is preferably human. It will be appreciated that the individual to be treated may be known to be sensitised to the allergens, at risk of being sensitised or suspected of being sensitised. The individual can be tested for sensitisation using techniques well known in the art and as described herein. Alternatively, the individual may have a family history of allergy to grass. It may not be necessary to test an individual for sensitisation to grass because the individual may display symptoms of allergy when exposed to grass. By exposure is meant proximity to, for example, a grass plant, or a substance or product derived from a grass plant, or a substance or product containing or comprising either of the above. The substance or product derived from a grass plant is typically grass pollen. By proximity is meant 10 metres or less, 5 metres or less, 2 metres or less, 1 metre or less, or 0 metres from the items described above. Symptoms of allergy can include itchy eyes, runny nose, breathing difficulties, red itchy skin or rash.

15

**[0129]** The individual to be treated may be of any age. However, preferably, the individual may be in the age group of 1 to 90, 5 to 60, 10 to 40, or more preferably 18 to 35.

**[0130]** Preferably, the individual to be treated is from a population that has MHC allele frequencies within the range of frequencies that are representative of the Caucasian population. Reference population allele frequencies for 11 common DRB1 allele families are shown in Table A (Data from HLA Facts Book, Parham and Barber).

20

Table A

DRB1	1	3	4	7	8	11	12	13	14	15	16
%	6.4	14.7	15.7	8.8	3.4	8.3	3.9	14.7	2.9	17.6	2.5
Reference population %	9.4	11.1	12.8	13.2	3.7	13.4	2.3	10.2	3.2	10.7	3.6

Reference frequencies were obtained by analysis of multiple studies reporting frequencies and the figures shown are mean values. Preferably therefore, the individual to be treated is from a population that has equivalent MHC allele frequencies as the reference population for the alleles referred to Table A (such as for at least 1, 2, 3, 4, 5 or all of the alleles), for example within the ranges of those figures plus or minus 1, 2, 3, 5, 10, 15 or 20%.

**[0131]** Preferably the individual is from a population where the allele frequencies of the following DRB1 alleles is:

- 4 - at least 9%
- 7 - at least 10%
- 11 - at least 8%.

45

**[0132]** The individual may have had allergy to grass for at least 2 weeks, 1 month, 6 months, 1 year or 5 years. The individual may suffer from a rash, nasal congestion, nasal discharge and/or coughing caused by the allergy. The individual may or may not have been administered with other compositions/compounds which treat grass allergy. The individual may live in a geographical region which has:

50

- a temperate climate, and/or
- a typical soil pH in the range of about 3.5, 4 or 4.5 to about 5.5, 6, 7 or 8; and/or
- a mean annual precipitation no less than about 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20cm per year and no greater than about 180cm, 250cm, 300cm, 400cm or 500cm per year; and/or
- an annual minimum temperature of no less than about -5°C, -4°C, -3°C, -2°C, -1°C, 0°C, 1°C, 2°C, 3°C, 4°C or 5°C and/or

- an annual maximum temperature of no greater than about 35°C, about 30°C, about 29°C, about 28°C, about 27°C, about 26°C about 25°C, about 24°C, about 23°C or about 22°C; and/or
- lies between the latitudes of about 30° north and about 30° south.

5 [0133] The individual typically suffers from allergy to grass in a particular season. The season typically corresponds to the flowering season of grass, which is typically summer, preferably early summer (for example from May to June in the Northern hemisphere) or late summer (for example August to October in the Northern hemisphere). The grass allergic individual is typically allergic to grass pollen.

10 *Combination immunotherapy*

[0134] Since many individuals are allergic, or may require desensitizing to several polypeptide antigens, the current invention also provides means of desensitizing individuals that are allergic to multiple antigens. "Tolerance" induced in an individual to a first polypeptide antigen or allergen can create in the individual a "tolerogenic environment" wherein 15 inappropriate immune responses to other antigens can be downregulated in order to provide tolerance to other antigens.

[0135] This finding means that individuals allergic to multiple allergens can be treated in a greatly reduced time period, and that individuals seriously allergic to some allergens (e.g., peanuts) but more mildly allergic to other allergens (e.g., 20 cat dander) can benefit from a therapy wherein tolerance to the milder allergen is established and then this tolerogenic environment is used to provide tolerance to the other, more extreme allergen. In addition, individuals suffering from an autoimmune disorder who are additionally sensitised (or otherwise immune) to an unrelated antigen or allergen can benefit from a treatment regime wherein tolerance to the unrelated antigen or allergen is first established and then this tolerogenic environment is used to provide tolerance to the autoantigen associated with the autoimmune disorder.

[0136] A method is therefore provided for desensitising a grass allergic individual to grass allergen as described above and one or more further different polypeptide antigens. The method entails, in a first step, administering to the individual 25 a composition/product/formulation (primary composition) according to the invention as described herein and wherein the administration is carried out in a manner sufficient to generate a hyporesponsive state against grass allergen. Once a hyporesponsive state has been established toward grass allergen, or at least a shift toward desensitisation has occurred, the method entails administration of a secondary composition comprising a second, different polypeptide antigen to which the individual is to be sensitised. Administration of the secondary composition is carried out in such a way as to 30 take advantage of the tolerogenic environment established by use of the primary composition, where it is now possible to establish tolerance to the second, different polypeptide antigen. The secondary composition is coadministered with either the first primary composition or a larger fragment of the grass allergen(s). By "coadministered" it is meant either the simultaneous or concurrent administration, e.g., when the two are present in the same composition or administered in separate compositions at nearly the same time but at different sites, as well as the delivery of polypeptide antigens 35 in separate compositions at different times. For example, the secondary composition may be delivered prior to or subsequent to delivery of the first composition at the same or a different site. The timing between deliveries can range from about several seconds apart to about several minutes apart, several hours apart, or even several days apart. Furthermore, different delivery methods can be employed.

[0137] Classes of suitable allergens include, but are not limited to dust mite allergens, pollens, animal dander (especially 40 cat dander), grasses, molds, dusts, antibiotics, stinging insect venoms, and a variety of environmental (including chemicals and metals), drug and food allergens. Common tree allergens include pollens from cottonwood, popular, ash, birch, maple, oak, elm, hickory, and pecan trees; common plant allergens include those from mugwort, ragweed, English plantain, sorrel-dock and pigweed; plant contact allergens include those from poison oak, poison ivy and nettles; common 45 grass allergens include rye grass, Timothy, Johnson, Bermuda, fescue and bluegrass allergens; common allergens can also be obtained from molds or fungi such as Alternaria, Fusarium, Hormodendrum, Aspergillus, Micropolyspora, Mucor and thermophilic actinomycetes; epidermal allergens can be obtained from house or organic dusts (typically fungal in origin), or from animal sources such as feathers, and dog dander; common food allergens include milk and cheese (dairy), egg, wheat, nut (e.g., peanut), seafood (e.g., shellfish), pea, bean and gluten allergens; common environmental 50 allergens include metals (nickel and gold), chemicals (formaldehyde, trinitrophenol and turpentine), Latex, rubber, fiber (cotton or wool), burlap, hair dye, cosmetic, detergent and perfume allergens; common drug allergens include local anesthetic and salicylate allergens; antibiotic allergens include penicillin, tetracycline and sulfonamide allergens; and common insect allergens include bee, wasp and ant venom, and cockroach calyx allergens. Particularly well characterized 55 allergens include, but are not limited to, the major cat allergen Fel d 1, bee venom phospholipase A2 (PLA) (Akdis et al. (1996) J. Clin. Invest. 98:1676-1683), birch pollen allergen Bet v 1 (Bauer et al. (1997) Clin. Exp. Immunol. 107:536-541), and the multi-epitopic recombinant grass allergen rKBG8.3 (Cao et al. (1997) Immunology 90:46-51).

[0138] Particularly preferred T cell epitopes are derived from the allergens: cat dander protein Fel d 1; House dust mite proteins Der p 1, Der p 2 and Der p 7; Ragweed protein amb a 1.1, a 1.2, a 1.3 or a 1.4; Rye grass proteins Lol p 1 and Lol p 5; Timothy grass proteins Phl p 1 and Phl p 5; Bermuda grass protein Cyn d 1; Alternaria alternate proteins

Alt a 1, Alt a 2 and Enolase (Alt a 6); Birch protein Bet v 1 and P14; German Cockroach proteins Bla g 1, Bla g 2, Bla g 3, Bla g 4, Bla g 5 and Bla g 6; Mugwort protein Art v 1; Russian thistle protein Sal k 1 and Sal k 2; peanut Ara h 1, Ara h 2, Ara h 3, Ara h 4, Ara h 5, Ara h 6, plant profilins or lipid transfer proteins or a human leukocyte antigen.

**[0139]** These and other suitable allergens are commercially available and/or can be readily prepared as extracts following known techniques.

**[0140]** Preferably, the second polypeptide allergen is a grass allergen or a grass allergen fragment, more particularly a grass allergen or grass allergen fragment selected from the list of grass allergen sequences indexed by database accession numbers (NCBI Entrez accession numbers) below. NCBI is the National Center for Biotechnology information and is a division of the US National Institutes of Health. The NCBI web site, from which access to the database may be sought, is [www.ncbi.nlm.nih.gov/](http://www.ncbi.nlm.nih.gov/). Allergen sequences indexed by database accession numbers.

75232277 Dac g 1, 75139988 Dac g 1, 75232276 Dac g 1, 33149335 Dac g 1, 33149333 Dac g 1, 75163303 Dac g 5, 75163399 Dac g 5, 75163400 Dac g 5, 14423124 Dac g 5, 14423122 Dac g 5, 14423120 Dac g 5, 1171005 Hol l 1, 3860384 Hol l 1, 414703 Hol l 1, 1167836 Hol l 1, 1085628 Hol l 1, 2266625 Hol l 5, 2266623 Hol l 5, 75172041 Hol 1 5, 75098038 Hol l 5, 75098037 Hol l 5, 11991229 Hol l 5, 2266623 Hol l 5, 2506771 Hor v 1, 282991 Hor v 1, 75219009 Hor v 5, 126385 Lol p 1, 168314 Lol p 1, 6599300 Lol p 1, 75274600 Lol p 1, 168316 Lol p 1, 3210053 Lol p 1, 3210049 Lol p 1, 3210047 Lol p 1, 3210050 Lol p 1, 3210044 Lol p 1, 3210043 Lol p 1, 3210041 Lol p 1, 3210039 Lol p 1, 3210037 Lol p 1, 3210036 Lol p 1, 3210035 Lol p 1, 3210034 Lol p 1, 3210033 Lol p 1, 3210032 Lol p 1, 3210030 Lol p 1, 100636 Lol p 1320616 Lol p 1, 320614 Lol p 1, 100638 Lol p 1, 100636 Lol p 1, 320614 Lol p 1, 100637 Lol p 1, 126386 Lol p 2a, 126387 Lol p 3, 2498581 Lol p 5 a, 4416516 Lol p 5 a, 485371 Lol p 5 a, 100639 Lol p 5 a, 3409495 Lol p 5 a, 3409494 Lol p 5 a, 3409493 Lol p 5 a, 3409489 Lol p 5 a, 3409488 Lol p 5 a, 3409487 Lol p 5 a, 3409486 Lol p 5 a, 3409484 Lol p 5 a, 3409483 Lol p 5 a, 3409481 Lol p 5 a, 3409479 Lol p 5 a, 3409478 Lol p 5 a, 3409477 Lol p 5 a, 3409476 Lol p 5 a, 3409475 Lol p 5 a, 3409474 Lol p 5 a, 3409473 Lol p 5 a, 3409472 Lol p 5 a, 3409471 Lol p 5 a, 3409470 Lol p 5 a, 3409469 Lol p 5 a, 3409468 Lol p 5 a, 3409467 Lol p 5 a, 3409466 Lol p 5 a, 3409456 Lol p 5 a, 3209999 Lol p 5 a, 3210002 Lol p 5 a, 3210003 Lol p 5 a, 3210004 Lol p 5 a, 3210005 Lol p 5 a, 3210006 Lol p 5 a, 3210007 Lol p 5 a, 3210008 Lol p 5 a, 3210009 Lol p 5 a, 3210010 Lol p 5 a, 3210011 Lol p 5 a, 3210012 Lol p 5 a, 3210013 Lol p 5 a, 3210014 Lol p 5 a, 3210015 Lol p 5 a, 3210017 Lol p 5 a, 3210018 Lol p 5 a, 3210019 Lol p 5 a, 3210020 Lol p 5 a, 3210021 Lol p 5 a, 3210022 Lol p 5 a, 3210023 Lol p 5 a, 3210024 Lol p 5 a, 3210025 Lol p 5 a, 3210026 Lol p 5 a, 542129 Lol p 5 a, 2498582 Lol p 5 b, 3409457 Lol p 5 b, 626028 Lol p 5 b, 542131 Lol p 5 b, 455288 Lol p 5 b, 6634467 Lol p 5c, 455288 Lol p isoform 9, 1582249 Lol p 11, Additional Lolium sequences : 135480; 417103; 687261; 687259; 1771355; 2388662; 631955; 542131; 542130; 542129; 100636; 626029; 542132; 320616; 320615; 320614; 100638; 100634; 82450; 626028; 100639; 283345; 542133; 1771353; 1763163; 1040877; 1040875; 250525; 551047; 515377; 510911; 939932; 439950; 2718; 168316; 168314; 485371; 2388664; 2832717; 2828273; 548867; 3409458 Pha a 1, 3210038 Pha a 1, 2498576 Pha a 1, 1246116 Pha a 1, 3210031 Pha a 1, 3210027 Pha a 1, 3210028 Pha a 1, 3210029 Pha a 1, 82450 Pha a 1, 409328 Pha a 1, 2498580 Pha a 5, 2498579 Pha a 5, 2498578 Pha a 5, 2498577 Pha a 5, 1246120 Pha p 5, 1246119 Pha p 5, 1246118 Pha p 5, 1246117 Pha p 5, 3409480 Pha p 5, 3409482 Pha p 5, 3409490 Pha p 5, 1171008 Phl p 1, 28373838 Phl p 1, 28373839 Phl p 1, 473360 Phl p 1, 3901094 Phl p 1, 1582250 Phl p 1, 75221090 Phl p 1, 3210052 Phl p 1, 3210046 Phl p 1, 3210040 Phl p 1, 629812 Phl p 1, 481432 Phl p 1, 1684718 Phl p 5, 13430402 Phl p 5, 3135501 Phl p 5, 3135499 Phl p 5, 3135497 Phl p 5, 1684720 Phl p 5, 40644796 Phl p 5, 3309039 Phl p 5, 3309041 Phl p 5, 739542 Phl p 5, 3309047 Phl p 5, 3309045 Phl p 5, 3309043 Phl p 5, 3135503 Phl p 5, 626037 Phl p 5, 2851456 Phl p 5a, 2398757 Phl p 5a, 1092249 Phl p 5a, 29500897 Phl p 5a, 422005 Phl p 5a, 3409492 Phl p 5a, 2851457 Phl p 5b, 481397 Phl p 5b, 1096197 Phl p 5b, 2398759 Phl p 5b, 3409491 Phl p 5b; Additional Phleum sequences: 458878; 548863; 2529314; 2529308; 2415702; 2415700; 2415698; 542168; 542167; 626037; 542169; 541814; 542171; 253337; 253336; 453976; 439960; 75267691 Poa p 1, 4090265 Poa p 1, 280414 Poa p 1, 320620 Poa p 1, 250525 Poa p 5, 75172042 Poa p 5, 113562 Poa p 9, 113561 Poa p 9, 113560 Poa p 9, 729944 Zea m 1, 115502168 Zea m 1b, 11550238 Zea m 1d, 115502167 Zea m 1c, 122238295 Zea m 1, 75272187 Zea m 1, 115502389 Zea m 1, 162459584 Zea m 1, 89892723 Zea m 1, .293902 Zea m 1, 89892721 Zea m 1, 114794319 Zea m 1, 478272 Zea m 1.

#### Delivery methods

**[0141]** Once formulated the compositions of the invention can be delivered to a subject *in vivo* using a variety of known routes and techniques. For example, a composition can be provided as an injectable solution, suspension or emulsion and administered via parenteral, subcutaneous, epidermal, intradermal, intramuscular, intraarterial, intraperitoneal, intravenous injection using a conventional needle and syringe, or using a liquid jet injection system. Compositions can also be administered topically to skin or mucosal tissue, such as nasally, intratracheally, intestinal, rectally or vaginally, or provided as a finely divided spray suitable for respiratory or pulmonary administration. Other modes of administration include oral administration, suppositories, sublingual administration, and active or passive transdermal delivery techniques.

[0142] Where a peptide of the invention is to be administered, it is preferred to administer the peptide to a site in the body where it will have the ability to contact suitable antigen presenting cells, and where it, or they, will have the opportunity to contact T cells of the individual. Where an APC is to be administered, it is preferred to administer the APC to a site in the body where it will have the ability to contact, and activate, suitable T cells of the individual.

5

#### *Delivery regimes*

[0143] Administration of the peptides/polynucleotides/cells (such as the composition containing a plurality of peptides) may be by any suitable method as described above. Suitable amounts of the peptide may be determined empirically, but typically are in the range given below. A single administration of each peptide may be sufficient to have a beneficial effect for the patient, but it will be appreciated that it may be beneficial if the peptide is administered more than once, in which case typical administration regimes may be, for example, once or twice a week for 2-4 weeks every 6 months, or once a day for a week every four to six months. As will be appreciated, each peptide or polynucleotide, or combination of peptides and/or polynucleotides may be administered to a patient singly or in combination.

10

[0144] Dosages for administration will depend upon a number of factors including the nature of the composition, the route of administration and the schedule and timing of the administration regime. Suitable doses of a molecule of the invention may be in the order of up to 15 $\mu$ g, up to 20 $\mu$ g, up to 25 $\mu$ g, up to 30 $\mu$ g, up to 50 $\mu$ g, up to 100 $\mu$ g, up to 500 $\mu$ g or more per administration. Suitable doses may be less than 15 $\mu$ g, but at least 1ng, or at least 2ng, or at least 5ng, or at least 50ng, or least 100ng, or at least 500ng, or at least 1 $\mu$ g, or at least 10 $\mu$ g. For some molecules of the invention, the dose used may be higher, for example, up to 1 mg, up to 2 mg, up to 3 mg, up to 4 mg, up to 5 mg or higher. Such doses may be provided in a liquid formulation, at a concentration suitable to allow an appropriate volume for administration by the selected route.

15

#### *Kits*

20

[0145] The invention also relates to a combination of components described herein suitable for use in a treatment of the invention which are packaged in the form of a kit in a container. Such kits may comprise a series of components to allow for a treatment of the invention. For example, a kit may comprise one or more different peptides, polynucleotides and/or cells of the invention, or one or more peptides, polynucleotides or cells of the invention and one or more additional therapeutic agents suitable for simultaneous administration, or for sequential or separate administration. The kit may optionally contain other suitable reagent(s) or instructions and the like.

25

[0146] The invention is illustrated by the following Examples:

#### Example 1

30

##### *MHC Class II binding search*

35

[0147] The aim of this study is to identify a distinct panel of peptides with strong affinities for the eight most common human MHC Class II HLA-DRB1 \* allotypes. In order to identify binding peptides in the major grass allergens Rye Lol p 1, Rye Lol p Va, p Vb, p 5a and p 5b, Bermuda Cyn d 1 and Timothy Phl p 5, an *in silico* approach known as "peptide threading" was employed using the commercially available EpiMatrix algorithm (EpiVax Inc.) This is a bioinformatic analysis of peptides from a sequence for the potential to be accommodated within the binding groove of MHC class II HLA-DR molecules. EpiMatrix is a matrix-based algorithm that ranks 9 amino acid long segments, overlapping by 8 amino acids, from any polypeptide sequence by estimated probability of binding to each of the selected MHC molecules. (De Groot et al., AIDS Research and Human Retroviruses 13:539-41 (1997)). The procedure for developing matrix motifs was published by Schafer et al, 16 Vaccine 1998 (1998). In this Example, binding potential for HLA DR1, DR3, DR4, DR7, DR8, DR11, DR13 and DR15 is assessed. Putative MHC ligands are selected by scoring each 9-mer frame in a protein sequence. This score is derived by comparing the sequence of the 9-mer to the matrix of amino acid sequences known to bind to each MHC allele. Retrospective studies have demonstrated that EpiMatrix accurately predicts published MHC ligands (Jesdale et al., in Vaccines '97 (Cold Spring Harbor Press, Cold Spring Harbor, N.Y., 1997)). Successful prediction of peptides which bind to multiple MHC molecules has also been confirmed.

40

[0148] Estimated probability of binding to a selected MHC molecule is calculated by EpiMatrix as follows. The peptides are scored by estimating the relative promotion or inhibition of binding for each amino acid, compared to known MHC binders for a given MHC allele. This information is summed across the peptide and a summary score (EMX score) is assigned to the entire peptide. After comparing the EMX score to the scores of known MHC ligands, EpiMatrix arrives at an "estimated binding probability" (abbreviated as EBP, but not strictly a probability). The EBP describes the proportion of peptides with EpiMatrix scores as high or higher that will bind to a given MHC molecule. EBPs range from 100% (highly likely to bind) to less than 1% (very unlikely to bind).

The polypeptide sequences analysed by EpiMatrix are shown in Table 1:

Table 1

5	Ber01 ori	SGKAFGAMAKKGQEDKLRKA
	Ber02 ori	PKDSDEFIPMKSSWGAIRIDPKPLKGP
	Ber03 ori	RLTSEGGGAHLVQDDVIPANWKPDTVYTSK
	Bio01 ori	QKLIEKINAGFKAAVAA
10	Bio02 ori	AYVATLTEALRVIAGTL
	Bio03 ori	KFIPTLVAAVKQAYAAKQAT
	Bio04 ori	TALKKAVTAMSEAEKEA
15	Bio05 ori	NDKFTVFESAFNKALNE
	Rye01 ori	LDAKSTWYGKPTGAGPKDNG
	Rye02 ori	GHAFGSMAKKGEEQNVRSG
	Rye03 ori	GSNPNYLAILVKYVDGDGDV
20	Rye04 ori	KESWGAVWRIDTPDKLTGPF
	Rye05 ori	DVNAGFKAAVAAAANAPPAD
	Rye06 ori	GATPEAKYDAFVTALTEALR
25	Rye07 ori	GELQIVDKIDAAFKIAATAANAAAPTNDKF
	Rye08 ori	GAYETYKFIPSLEAAVKQAY
	Rye09 ori	PEVKYAVFEAALTAKAITAMTQAQKAGKPA
30	Tim10 ori	PEVKYTVFETALKKAITAMSEAQ

**[0149]** Based on the results of the EpiMatrix analysis of these sequences, core peptides were identified which were predicted to have good MHC binding properties. The selected peptides are shown in Table 2. Peptides highlighted in grey and marked \* are not selected peptides. These correspond to the original sequences analysed in EpiMatrix, from which the subsequent selected peptides derive. For example, Ber01 derives from Ber01 ori.

35

40

45

50

55

Table 2

## Group 1 peptides

SEQ ID NO:	Peptide name	Peptide Sequence	Allergen Group	Protein of origin
74	Ber01 ori	SGKAFGAMAKKGQEDKLRKA	1	Ber Cyn d1
1	Ber01	SGKAFGAMAKKGQED	1	Ber Cyn d1
75	Ber02 ori	PKDSDEFIPMKSSWGAIWRI DPKKPLKGP	1	Ber Cyn d1
2	Ber02	FIPMKSSWGA	1	Ber Cyn d1
3	Ber02A	WGAIWRIDPKKPL	1	Ber Cyn d1
4	Ber02B	KDSDEFIPMKSSWGAIWR	1	Ber Cyn d1
5	Ber02C	KSSWGAIWRIDPKKPLK	1	Ber Cyn d1
6	Ber02D	MKSSWGAIWRIDPKKPLK	1	Ber Cyn d1
7	Ber02E	MKSSWGAIWRIDPPKPLK	1	Ber Cyn d1
76	Ber03 ori	RLTSEGGAAHLVQDDVIPANW KPDTVYTSK	1	Ber Cyn d1
8	Ber03A	IPANWKPDIVYTSK	1	Ber Cyn d1
9	Ber04	KATFYGSDPRGAAP	1	Ber Cyn d1
10	Ber05	AYHFDSLGSKAFG	1	Ber Cyn d1
77	Rye01 ori	LDAKSTWYGKPTGAGPKDNG	1	Rye Lol p1
11	Rye01	LDAKSTWYGKPTGAG	1	Rye Lol p1

45

50

55

12	Rye01A	KWLDAKSTWYKGKPTGAG	1	Rye Lol p1	*
78	Rye02 ori	GHAFGSMAKKGEEQNVRSAG	1	Rye Lol p1	*
13	Rye02	FGSMAKKGEEQNVRSAG	1	Rye Lol p1	
14	Rye02A	HAFGMAKKGEEQNVRSAG	1	Rye Lol p1	
79	Rye03 ori	GSNPNYLAILVKYVDGDGDV	1	Rye Lol p1	*
15	Rye03A	SNPNYLAILVKYVD	1	Rye Lol p1	*
80	Rye04 ori	KESWGAVWRIDTPDKL TGPF	1	Rye Lol p1	*
16	Rye04	WGAVWRIDTPDKL	1	Rye Lol p1	
17	Rye04A	KESWGAVWRIDTPDKL	1	Rye Lol p1	
18	Rye04B	KESWGAVWRIDTPDKLGP	1	Rye Lol p1	
19	Rye12	APYHFDSLGHAFGS	1	Rye Lol p1	

30 *Group 5 peptides*

81	Bio02 ori	AYVATLTEALRVIAGTL	5	Rye Lol p 5b	*
28	Bio02A	KYDAYVATLTEALR	5	Rye Lol p 5b	*
82	Bio03 ori	KFIPTLVAAVKQAYAAKQAT	5	Rye Lol p 5b	*
29	Bio03A	KFIPTLVAAVKQAYAAKQ	5	Rye Lol p 5b	
30	Bio03B	YKFIPTLVAAVKQAYAAKQ	5	Rye Lol p 5b	
83	Bio04 ori	TALKKAVTAMSEAEKEA	5	Rye Lol p 5b	*
31	Bio04A	LKKAVTAMSEAEK	5	Rye Lol p 5b	
32	Bio04B	PETALKKAVTAMSEAEK	5	Rye Lol p 5b	
84	Bio05 ori	NDKFTVFESAFNKALNE	5	Rye Lol p Va	*
33	Bio05B	KFTVFESAFNKALNE	5	Rye Lol p Va	

85	Rye05 ori	DVNAGFKAAVAAAANAPPAD	5	Rye Lol p Va	*
34	Rye05A	FKAAVAAAANAPPADKFK	5	Rye Lol p Va	
35	Rye05C	NAGFKAAVAAAANAPPK	5	Rye Lol p Va	
86	Rye06 ori	GATPEAKYDAFVTALTEALR	5	Rye Lol p Va	*
36	Rye06A	KYDAFVTALTEALR	5	Rye Lol p Va	
37	Rye06B	PEAKYDAFVTALTEALR	5	Rye Lol p Va	
87	Rye07 ori	GELOIVDKIDAAFKIAATAA NAAPTNDF	5	Rye Lol p Va	*
38	Rye07A	GELQIVDKIDAAFK	5	Rye Lol p Va	
39	Rye07B	KIPTGELQIVDKIDA	5	Rye Lol p Va	
40	Rye07G	FKIAATAANAAPTN DK	5	Rye Lol p Va	
41	Rye07H	AFKIAATAANAAPTN DK	5	Rye Lol p Va	
88	Rye08 ori	GAYETYKFIPSLEAAVKQAY	5	Rye Lol p Va	*
42	Rye08	YKFIPSLEAAVKQAY	5	Rye Lol p Va	
43	Rye08A **	ETYKFIPSLEAAVKQAY	5	Rye Lol p Va	
44	Rye08B	DSYKFIPTLVAVK	5	Rye Lol p Va	
89	Rye09 ori	PEVKYAVFEAALT KAITAMT QAQKAGKPA	5	Rye Lol p Va	*
45	Rye09A	LTKAITAMTQAQKAGK	5	Rye Lol p Va	
46	Rye09B	PEVKYAVFEAALT KAIT	5	Rye Lol p Va	
47	Rye09D	KYAVFEAALT KAITAMT	5	Rye Lol p Va	
48	Rye11	DKFKIFEAAFSESSK	5	Rye Lol p Va	
49	Rye13	TPLRRRTSSRSSRP	5	Rye Lol p Va	
50	Rye14	DVAYKAAEAHPRGQ	5	Rye Lol p Va	
51	Rye15	ALRVIAGTLEVHA	5	Rye Lol p Vb	
52	Rye16	FENTFNNAIKVSLG	5	Rye Lol p Vb	

90	Tim10 ori	PEVKYTVFETALKKAITAMS EAQ	5	Tim Ph1 p5	*
53	Tim10B	KYTVFETALKKAITAMSE	5	Tim Ph1 p 5	
54	Tim10C	LKKAITAMS	5	Tim Ph1 p 5	
55	Tim10D	PETALKKAITAMS	5	Tim Ph1 p 5	

Tim = Timothy grass; Rye = Rye grass; Cyn = Bermuda grass

10 [0150] Any peptides indicated above which have an N terminal glutamate (E) or glutamine (Q) residue, for example Rye 08A, may have this residue replaced with pyroglutamate to improve stability during manufacture, without affecting function of the peptide. The data from further testing of these peptides is typically obtained using peptides where such replacement has taken place.

15 [0151] A further EpiMatrix analysis was performed on the entire sequences of three known Timothy grass sequences: Timothy Grass Ph1 p 1 (NCBI Accession no. 1N10A), Timothy Grass Ph1 p Va (NCBI Accession no. Q40962), and Timothy Grass Ph1 p Vb (Accession no. Q40963). This analysis identified further core peptides (and their flanking sequences) which are predicted to have good MHC class-II binding. These sequences are shown below in Tables 3A-C. In each table:

20 "Residues in main sequence" gives the location of the peptide within the sequences that were analysed. The core peptide (underscored middle amino acids in bold) defines the actual binding sequence that was identified during the analysis. The stabilizing flanks (N-terminal and C-terminal, not bold) were included for use with the core sequence and are typically required to aid manufacture of the peptides. "Number of hits" refers to the number of high predicted binding affinities for all MHC types tested within the sequence. The "EpiMatrix Cluster Score" is derived from the number of hits normalized for the length of the cluster. Cluster Score is thus the excess or shortfall in predicted 25 aggregate MHC binding properties relative to a random peptide standard. A score above 10 is considered to indicate broad MHC binding properties.

Table 3

A) 1N10A – Ph1 p 1

35

40

45

50

55

5	RESIDUES IN MAIN SEQUENC E (INCLUDI NG FLANKS)	SEQUENCE	Hydro- phobicit y	EpiMatrix HITS (EXCLUDI NG FLANKS)	EpiMatrix CLUSTER SCORE (EXCLUDIN G FLANKS)	SEQ ID NO:
10	<u>120</u> - <u>142</u>	GDE <b>QKLRSAGELELQFRRVK</b> CKY	-1.15	10	13.23	20
15	<u>157</u> - <u>175</u>	NPN <b>YLLALLVKYVNGDG</b> DVV	0.19	8	14.11	21
20	<u>191</u> - <u>209</u>	KES <b>WGAIRIDTPDKLTG</b> PP	-0.85	8	13.79	22
25	<u>229</u> - <u>241</u>	PEG <b>WKADTSYESK</b>	-1.77	6	12.78	23

## B) Q40962 – Phl p Va

25	RESIDUES IN MAIN SEQUENCE (INCLUDING FLANKS)	SEQUENCE	Hydro- phobicity	EpiMatrix HITS (EXCLUDING FLANKS)	EpiMatrix CLUSTER SCORE (EXCLUDING FLANKS)	SEQ ID NO:
30	<u>42</u> - <u>60</u>	<b>TEKINAGFKAALAGAGVQP</b>	0.41	8	13.04	56
35	<u>55</u> - <u>78</u>	GAG <b>VQPADKYRTFVATFGPAS</b> NKA	-0.32	12	17.23	57
40	<u>98</u> - <u>122</u>	AET <b>SKLDAAYKLAYKTAEGATP</b> PEAK	0.38	11	14.26	58
45	<u>164</u> - <u>184</u>	<b>IEKVDAAFKVAATAANAA</b> PAN	0.37	9	12.44	59
50	<u>206</u> - <u>226</u>	<b>YESYKFTPALEAAVKQAYAAT</b>	0.04	8	14.18	60
55	<u>232</u> - <u>253</u>	EVK <b>YTVFETALKKAITAMS</b> EAQ	-0.06	12	17.18	61

## C) Q40963 – Phl p Vb

40	RESIDUES IN MAIN SEQUENC E (INCLUDI NG FLANKS)	SEQUENCE	Hydro- phobicit y	EpiMatrix HITS (EXCLUDI NG FLANKS)	EpiMatrix CLUSTER SCORE (EXCLUDIN G FLANKS)	SE Q ID NO
45	<u>49</u> - <u>69</u>	<b>DINVGKAAVAAAASVPAADK</b>	0.63	13	21.17	62
50	<u>115</u> - <u>138</u>	<b>FDSFVASLTEALRVIAGALEV</b> HAV	1.10	11	15.59	63
55	<u>158</u> - <u>176</u>	<b>IDKIDAAFKVAATAAA</b> TAP	0.66	8	13.93	64

206 - 232	IPS <b>LEAAVKQAYAATVAAAPQ</b> <b>VKYAVF</b>	0.69	11	12.1	65
5 229 - 247	<b>YAVFEAALTAKITAMS</b> <sup>EVO</sup>	0.66	9	14.1	66

**Example 2**10 *Homology search*

[0152] The sequences of each of the peptides identified above as MHC Class II-binding were used to probe the sequence of alternative proteins in the grass allergen group from which the parent sequence derived. For example, peptide Rye01 is from Lol p 1, therefore the sequence of Rye01 was used to probe for conserved sequences in Group 1 from other grass species, in particular Timothy. The results of this analysis are shown below for the residues of Rye 1 from other grass species, in particular Timothy. The results of this analysis are shown below for the residues of Rye Lol p 1 as indicated, compared with corresponding sequences from Timothy Phl p 1 and Bermuda Cyn d 1:

Rye 19-38:	LDAKSTWYGKPTGAGPKDNG (RYE 01 ori)
Timothy	LDAKSTWYGKPTGAGPKDNG
Rye 109-128:	GHAFGSMAKKGEEQNVRSAG (Rye 02 ori)
Timothy	GHAFGAMAKKGDEQKLRSAG
Cyn d 1 109-128:	SGKAFGAMAKKGQEDKLRLKA
Rye 154-173:	GSNPYNYLAILVKYVGDGKV (Rye03 ori)
Timothy	GSNPYNYLALLVKYVNGDGV
Rye 190-209:	KESWGAVWRIDTPDKLTGPF (Rye04 ori)
Timothy	GKDKWIELKESWGAIWRIDTPDKLTGPF
Cyn d 1 181-209	PKDSDEFIPMKSSWGAIWRIDPKKPLK
Cyn d 1 217-241	
30 Rye 19-38: EGGAHLVQDDVIPANWPDTVYTSK	
Timothy	EGGKTEAEDVIPEGWKADTSYESK

[0153] Similarly, shown below are results for the residues of Rye Lol p 5 as indicated, compared with corresponding sequences from Timothy Phl p5 variants as indicated:

Rye 37-56:	DVNAGFKAAVAAAANAPPAD (Rye 05 ori)
Phl p 5a	KINAGFKAALAGAGVQPAD
Phl p Va	KINAGFKAALAGAGVQPAD
40 Phl p 5	KINDGFKAALAAAAGVPPAD
Phl p Vb	DINVGFKAAVAAAASVPAAD
Phl p Vb	DINVGFKAAVAAAASVPAAD
Rye 100-119:	GATPEAKYDAFVTALTEALR (Rye 06 ori)
Phl p 5a	GATPEAKYDAYVATLSEALR
Phl p Va	GATPEAKYDAYVATLSEALR
45 Phl p 5	GATPEAKYDAYVATLSEALR
Phl p Vb	GATPEAKFDSFVASLTEALR
Phl p Vb	GATPEAKFDSFVASLTEALR
50 Rye 145-164:	GELQIVDKIDAQFVAAATAA (from Rye 07 ori)
Pha p 5a	GELQVIEKDAQFVAAATAA
Phl p Va	GELQVIEKVDAAQFVAAATAA
Phl p 5	GELQFIEKVDSALKVAAATAA
55 Phl p Vb	GELQIIDKIDAQFVAAATAA
Phl p Vb	GELQIIDKIDAQFVAAATAA
Rye 154-173:	DAAFKVAAATAANAAAPTNDKF (from Rye 07 ori)
Phl p 5a	DAAFKVAAATAANAAAPNDKF

(continued)

5	Phl p Va	DAAFKVAATAANAAPANDKF
	Phl p 5	DSALKVAATAANAAAANDKF
	Phl p Vb	DAAFKVAATAATAPADDKF
	Phl p Vb	DAAFKVAATAATAPADDKF
	Rye 190-209:	GAYETYKFIPSLEAAVKQAY (Rye 08 ori)
	Phl p 5a	GAYESYKFIPALEAAVKQAY
10	Phl p Va	GAYESYKFIPALEAAVKQAY
	Phl p 5	GAYESYKFIPALEAAVKQAY
	Phl p Vb	GAYDTYKCIPSLEAAVKQAY
	Phl p Vb	GAYDTYKCIPSLEAAVKQAY
	Rye 217-236:	PEVKYAVFEAALTKAITAMT (from Rye 09 ori)
15	Phl p 5a	PEVKYTVFETALKKAITAMS
	Phl p Va	PEVKYTVFETALKKAITAMS
	Phl p 5	PEVKYTVFETALKKAITAMS
	Phl p Vb	PEVKYTVFETALKKAITAMS
	Phl p Vb	AEVKYAVFEAALTKAITAMS
20		
	Rye 226-245:	AALTKAITAMTQAQKAGKPA (from Rye 09 ori)
	Phl p 5a	TALKKAITAMSEAQKAAKPA
	Phl p Va	TALKKAITAMSEAQKAAKPA
25	Phl p 5	TALKKAITAMSEAQKAAKPA
	Phl p Vb	AALTKAITAMSEVQKVSQPA
	Phl p Vb	AALTKAITAMSEVQKVSQPA

30 [0154] Based on the sequences from Timothy grass proteins which were highly conserved with the Rye sequences selected in Example 1, additional peptides derived from these sequences were predicted as having good MHC binding properties. These additional peptides are shown in Table 4.

Table 4

<u>Group 1 peptides</u>				
SEQID NO:	Peptide name	Peptide Sequence	Group	Protein
35	24	FGAMAKKGDEQKLRSG	1	Tim Phl p 1
	25	HAFGAMAKKGDEQKLRSG	1	Tim Phl p 1
40	26	SNPNYLALLVKYVNGD	1	Tim Phl p 1
	27	WGAIWRIDTPDKL	1	Tim Phl p 1
<u>Group 5 peptides</u>				
45	67	FKAAVAAAASVPAADKFK	5	Tim Phl p 5
	168	KFDSFVASLTEALR	5	Tim Phl p 5
	169	KIPAGELQIIDKIDA	5	Tim Phl p 5
50	70	FKVAATAANAAPANDK	5	Tim Phl p 5
	71	YKFIPALEAAVKQAY	5	Tim Phl p 5
	72	PEESYKFIPALEAAVKQAY	5	Tim Phl p 5
55	73	LTKAITAMSEVQKVSQ	5	Tim Phl p 5

**Example 3***In vitro binding analysis*

5 [0155] The peptides identified as being potential MHC Class II-binding are pre-screened for solubility in an aqueous, acidic milieu and the peptides are tested in an in vitro MHC Class II binding assay.

*Methods*

10 [0156] The assay employed is a competitive MHC class II binding assay, wherein each peptide is analysed for its ability to displace a known control binder from each of the human MHC class II allotypes investigated. The allotypes and control peptides used in this study are typically those shown below:

Allotype	Control Peptide	Sequence
DRB1 *0301	Myco. tuberculosis/leprae hsp 65 2-16	AKTIAYDEEARRGLE
DRB1*1101	Influenza haemagglutinin 307-319	PKYVKQNTLKLAT
DRB1*1501	Human myelin basic protein 85-99	ENPVVHFFKNIVTPR
Control peptides used in the in vitro binding assays		

25 [0157] Each of the peptides from Tables 2 to 4 (excluding those marked \*) are analysed in the competition assay and screened for relative binding compared to the control peptides. Due to the nature of the competitive assay the data for each peptide is determined as a ratio of its own IC50 to that of the control peptide. Thus, a peptide that has an IC50 value that is parity to the control peptide has an identical binding affinity, while peptides with a ratio less than one have a higher affinity and those with a ratio greater than one have a lower affinity.

30 [0158] Solubility in aqueous solution is an essential criterion for a peptide to be an effective therapeutic agent. Therefore, as a consequence of the solubility screen very hydrophobic peptides with a high frequency of large hydrophobic amino acid residues in multiple binding registers will be eliminated. This is a characteristic of promiscuous HLA-DRB1\* binders. Peptides which bind to one or more of the MHC Class II allotypes are identified. It would be expected that such peptides would have the ability to bind similar allotypes that have not been tested through the homology of MHC structures.

**Example 4**

35 [0159] The following method is applied to the same peptides as in Example 3.

*Cell proliferation assay*

40 [0160] The cell proliferation assay is performed on PBMC's ( $140 \times 10^6$  cells required for all parameters to be tested). Proliferation is measured by the incorporation of the radiolabelled compound 3H-thymidine. In more detail, 100 $\mu$ l of the appropriate antigen or peptide concentration is distributed into the appropriate wells of 96 well plates. The plates are then placed into a humidified 5% CO<sub>2</sub> incubator set at 37°C for a maximum of 4 hours. PBMC's isolated as described above are prepared to a concentration of  $2 \times 10^6$  cells/ml in complete medium at room temperature. 100 $\mu$ l of cell solution is then distributed into each of the wells of the 96 well plates containing antigen/peptide. The plates are then incubated for 6 to 8 days. The cultures are pulsed with tritiated thymidine solution by adding 10 $\mu$ l of tritiated thymidine stock solution (1.85MBq/ml in serum-free RPMI medium) to each well. The plates are then returned to the incubator for between 8 and 16 hours. Cultures are then harvested using a Canberra Packard FilterMate 196 cell harvester. Dried filter mats are counted using an appropriate beta scintillation counter.

45 [0161] Counts from wells containing peptide are compared statistically to wells containing media alone (12 wells per group). The non-parametric Mann-Whitney test is used. The same statistical test is used for all subjects. A statistically significant difference between media only wells and peptide-stimulated wells is considered a positive stimulation of PBMC's by the peptide.

**Example 5**

55 [0162] 59 peptides identified by the EpiMatrix analysis of Example 1 that encompassed one or more epitopes predicted to bind to at least 5/8 MHC Class II alleles (HLA DR01, 03, 04, 07, 08, 11, 13, 15) were selected for further studies. In

many cases the epitopes were predicted to bind all 8 of these alleles. Some sequences had two or more overlapping epitopes or non-overlapping but close epitopes binding to the same or different MHC Class II alleles. The selected peptides are shown in Tables 5A and 5B below.

**[0163]** 44 peptides were tested for activity in the *in vitro* T cell cytokine release assay described using peripheral blood mononuclear cells (PBMCs) from 48 grass allergic individuals. Peripheral blood mononuclear cells (PBMCs) were isolated from heparinised blood by Ficoll density gradient centrifugation. Results are shown in Table 5A. Another group of 15 peptides were tested in 28 grass allergic subjects. Results are shown in Table 5B. The subjects were from Hamilton and surrounding area in Ontario, Canada. The peptides were tested for stimulation of production of Interferon gamma (IFN-gamma), Interleukin-10 (IL-10) and Interleukin-13 (IL-13) in the supernatants of the PBMC cultures.

10 *Cytokine release assay*

**[0164]** Cytokine secretion profiles from PBMC's were analysed in response to the peptide stimulation. Supernatants from the cytokine release assay were tested for the presence of 3 cytokines, IFN- $\gamma$ , IL-10 and IL-13, using ELISA assays. The presence of all 3 cytokines was assayed using a multiplex bead array(Luminex Corporation). The cytokine release assay typically required  $40 \times 10^6$  PBMC's per subject. In more detail,  $250 \mu\text{l}$  of a  $200 \mu\text{g/ml}$  solution of the appropriate antigen or peptide concentration was distributed into the appropriate wells of 48 well plates. Plates were then incubated in a humidified 5%  $\text{CO}_2$  incubator at  $37^\circ\text{C}$  for a maximum of 4 hours.  $250 \mu\text{l}$  of a  $5 \times 10^6$  cell/ml PBMC suspension is then added to each well and the plates returned to the incubator for 5 days. Following stimulation, samples of culture supernatant were harvested for testing by multiplex bead assay according to standard protocols. Typically, the samples were harvested into 3 aliquots and frozen until the ELISA assays were performed..

**[0165]** A positive result for stimulation of cytokine secretion was taken where the reading was greater than four times the control well, where no peptide was added. Peptides giving a positive result for one or more cytokines in more than 18 out of the 48 subjects in the first group or 9 out of the 28 subjects in the second group were deemed to be particularly useful in the treatment of grass allergy. 20 of the 44 peptides tested in the first group of 48 subjects, and 8 of the 15 peptides tested in the second group of 28 subjects fulfilled the above criteria.

**[0166]** Table 5A shows the number of positives for the three cytokines out of the group of 48, and Table 5B shows the number of positives for the three cytokines out of the group of 28. Preferred peptides fulfilling the criteria defined above are highlighted in bold.

30 Table 5A

Peptide	IFN- $\gamma$ +ve/48	IL-13 +ve/48	IL-10 +ve/48	>18/48 +ve for one or more cytokine?
<b>Ber01</b>	35	22	34	Yes
<b>Ber02</b>	26	21	15	Yes
Ber02A	17	15	7	No
<b>Ber02B</b>	22	17	3	Yes
<b>Ber02C</b>	29	18	9	Yes
<b>Ber03A</b>	8	7	23	Yes
<b>Bio02A</b>	15	12	31	Yes
<b>Bio03A</b>	26	19	9	Yes
<b>Bio04A</b>	28	16	37	Yes
<b>Bio04B</b>	22	16	27	Yes
<b>Bio05B</b>	8	8	34	Yes
Rye01	12	5	2	No
Rye01A	2	4	4	No
Rye02	4	5	1	No
Rye02A	12	6	1	No
Rye03A	3	5	0	No
Rye04	6	3	1	No

(continued)

	Peptide	IFN- $\gamma$ +ve/48	IL-13 +ve/48	IL-10 +ve/48	$>18/48$ +ve for one or more cytokine?
5	Rye04A	8	4	2	No
	Rye05A	13	10	3	No
	<b>Rye05C</b>	20	19	12	Yes
	Rye06A	11	15	5	No
10	Rye06B	11	7	5	No
	<b>Rye07A</b>	8	3	25	Yes
	Rye07B	10	3	1	No
	Rye07G	7	3	2	No
15	Rye07H	10	8	4	No
	Rye08	9	6	2	No
	<b>Rye08A</b>	18	9	32	Yes
	Rye09A	6	5	11	No
20	<b>Rye09B</b>	28	24	16	Yes
	Tim02	16	6	3	No
	Tim02A	6	9	2	No
	Tim03A	9	9	4	No
25	<b>Tim04A</b>	16	4	31	Yes
	Tim05A	15	12	9	No
	Tim06A	17	11	11	No
	<b>Tim07B</b>	23	18	41	Yes
30	<b>Tim07G</b>	19	20	10	Yes
	Tim08	16	17	14	No
	<b>Tim08A</b>	21	10	8	Yes
	<b>Tim10B</b>	16	24	22	Yes
35	Tim10C	6	4	3	No
	<b>Tim10D</b>	28	8	3	Yes

Table 5B

	Peptide	IFN- $\gamma$ +ve/28	IL-13 +ve/28	IL-10 +ve/28	$\geq 9/28$ +ve for one or more cytokine?
45	<b>Bio02D</b>	21	8	21	Yes
	<b>Bio02E</b>	13	2	24	Yes
	<b>Bio04</b>	12	5	8	Yes
	<b>Bio05</b>	11	5	1	Yes
50	<b>Bio03B</b>	13	8	4	Yes
	Rye04B	3	6	0	No
	Rye07A	1	4	1	No
	Rye08B	2	3	1	No

(continued)

	Peptide	IFN- $\gamma$ +ve/28	IL-13 +ve/28	IL-10 +ve/28	$\geq 9/28$ +ve for one or more cytokine?
5	<b>Rye09D</b>	6	4	13	Yes
	<b>Rye11</b>	4	10	3	Yes
	<b>Rye12</b>	10	8	5	Yes
	Rye13	6	1	0	No
	Rye14	5	3	0	No
	Rye15	1	0	1	No
	Rye16	4	3	4	No

15 Example 6Ranking criteria for individual grass peptides

20 [0167] The 28 peptides which were selected as being particularly useful in the treatment of grass allergy on the basis of the criteria used in Example 5 were further evaluated in order to compare their characteristics. This involved calculation of the cumulative response observed to all 3 cytokines for each peptide i.e the total score (sum of the number of responders for the three cytokines). Pharmaceutical development aspects for each peptide were also considered on the basis of physical and chemical properties, in particular solubility, pI and hydrophobicity index (GRAVY). Epimatrix software was also used to predict binding strength to MHC class II HLA DR alleles, with values shown as high (1%) medium (5%) and low (10%) affinity for each specific MHC class II HLA DR allele shown.

25 [0168] The results of this analysis for 14 particularly preferred peptides are shown in Table 6. These particularly preferred peptides are allocated an internal ranking order of 1 to 14 which corresponds to the Inventors evaluation of their relative utility for the treatment of grass allergy. The 14 peptides were ranked according principally to the total score (sum of the number of responders for the three cytokines) with some weighting placed on the IFN-gamma response. Account was also taken of pharmaceutical development aspects and MHC coverage, as outlined above.

Table 6

	Peptide	Total score	Pi	GRAVY	Epimatrix analysis of MHC binding affinity			Solubility mg/ml	RANK
					1%	5%	10%		
35	Ber01	91	8.22	-0.94	08, 11	01, 08, 15	03, 04, 08	>20	1
40	Ber02	62	8.75	-0.26		01, 04, 08, 11, 13	11, 15	>20	5
45	Ber02B	42	6.12	-0.75		01, 04, 08, 11, 13	11, 15	>20	13
50	Ber02C	56	10.46	-0.941	03, 07, 08	01, 11	04, 13, 15	>20	6
55	Bio02A	58	6.07	-0.157	01, 04, 07	08, 11, 13	03, 15	>20	8
	Bio03A	54	10.00	0.344	03	01, 03, 04, 08, 11, 13, 15		>20	7
	Bio04A	81	8.50	-0.377	04	01, 03, 04, 08, 11, 13		>20	3
	Rye05C	51	10.00	0.176	01, 04	03, 08, 11, 13	04, 15	10	11
	Rye08A	58	6.24	-0.194	01, 03, 04, 11	13, 15	07	>20	9
	Rye09B	68	6.56	0.447	01, 07	01, 04, 11, 15	03, 08, 13	5	4
	Tim04A	51	5.96	-0.408	03, 04, 07	01, 11	08, 15	2	12
	Tim07B	82	4.56	0.107		03, 11, 15	01, 13	0.56	2
	Tim07G	49	8.59	-0.062		01, 04, 08, 15	03, 07, 11	>20	14

(continued)

5	Peptide	Total score	Pi	GRAV Y	Epimatrix analysis of MHC binding affinity			Solubility mg/ml	RANK
					1%	5%	10%		
10	Tim10B	62	8.43	-0.017	04	01, 03, 04, 08, 11, 13		1.65	10

[0169] The order of ranking for particularly preferred grass peptides is thus: 1, Ber01, (SEQ ID NO:1), Cyn d 1; 2, Tim07B (SEQ ID NO: 69), Phl p 5; 3, Bio04A (SEQ ID NO: 31), Lol p 5; 4, Rye09B (SEQ ID NO: 46), Lol p 5; 5, Ber02 (SEQ ID NO: 2), Cyn d 1; 6, Ber02C (SEQ ID NO: 5), Cyn d 1; 7, Bio03A (SEQ ID NO: 29), Lol p 5; 8, Bio02A (SEQ ID NO: 28), Lol p 5; 9, Rye08A, (SEQ ID NO: 43), Lol p 5; 10, Tim10B, (SEQ ID NO: 53), Phl p 5; 11, Rye05C (SEQ ID NO: 35), Lol p 5; 12, Tim04A (SEQ ID NO: 27), Phl p 1; 13, Ber02B (SEQ ID NO:4), Cyn d 1; 14, Tim07G (SEQ ID NO: 70), Phl p 5.

[0170] Surprisingly, the top ranked peptide, Ber01 and also 3 other peptides in the top 14 (Ber02, Ber02C and Ber02B) were derived from the Bermuda grass allergen Cyn d 1. Bermuda grass is a warm season perennial species adapted to tropical and subtropical climates. It grows best under extended periods of high temperatures, mild winters and moderate to high rainfall. Temperature is the main environmental factor that limits its adaptability to tropical and subtropical areas of the world. The northern limits of Bermuda grass extend into the transitional zone of the United States where low temperatures seldom drop below 10°F. For this reason it would not be expected that Bermuda grass allergens would be the major allergens recognised by subjects in Canada. Furthermore, Perennial ryegrass, although present, is not the most common grass in Canada and 6 of the top 14 peptides (Bio02A, Bio03A, Bio04A, Rye05C, Rye08A, Rye09B) were derived from Perennial rye allergens.

[0171] The following is a summary of grass types prevalent in Canada. Almost all forage grasses in Canada are improved cultivars of European species. Different grasses are adapted to grow in different areas of Canada, depending on soil and climate conditions. Timothy (*Phleum pratense*) is the most widely grown grass outside dry parts of the region, and is a dominant forage grass in eastern Canada. Crested wheatgrass (*Agropyron cristatum*) is a dominant forage grass in western Canada. Orchard grass (*Dactylis glomerata*) and Russian wild ryegrass (*Elymus junceus*) are dominant forage grasses in British Columbia. Bromegrass (*Bromus inermis*) is grown in eastern Canada and on the Prairies. Kentucky bluegrass (*Poa pratensis*) is commonly grown in many areas. In terms of prairie grasses, 3-awn grass (*Aristida longiseta*) is found in arid regions of British Columbia, and wild rice (*Zizania aquatica*) in eastern Canadian lakes. Certain genera (eg, *Arctagrostis* and *Arctophila*) are native to the Canadian Arctic. Turf grasses are developed from species that show desirable characteristics, eg, density of growth, fast growth after seeding, ability to remain green, etc. In Canada cold-hardiness and frequently drought resistance are also important. Popular Canadian lawn grass mixtures often include species of *Poa* (eg, Kentucky bluegrass, roughstalk bluegrass) and *Festuca* (especially creeping red fescue, chewing fescue), although other useful species have been developed.

[0172] Based on the prevalence of grass types in Canada, the observation that grass allergic individuals of Canadian origin are highly responsive to peptides from Bermuda grass and Perennial rye is unexpected. Such peptides therefore have the potential to have broad utility in treatment of grass allergic individuals worldwide. Also, the peptides which were found to induce high levels of response (ie a high total score) are commonly derived from allergen proteins conventionally described as "minor allergens", e.g Lol p 5 and Phl p 5. The major Timothy and Perennial Rye allergens recognized by IgE antibodies in grass allergic subjects are, respectively, Phl p 1 and Lol p 1. The induction of antibodies to these major allergens in allergic individuals is a T cell dependent process and so it would be expected that the peptides inducing high levels of T cell response would be mainly from Phl p 1 and Lol p 1.

[0173] The determination of the top 14 T cell stimulatory peptides from the group of 59 peptides tested in Example 5, and particularly identifying the top 8 peptides which can be used most optimally in combination involved a narrow and specific selection. A narrow subset of combinations were identified from a massive number of initially possible combinations of peptides. The total number of possible combinations for selecting 8 peptides from the top 14 ranked peptides is 3003. This number of combinations (3003) represents a very small proportion of the possible combinations of 8 peptides out of the original group of 73 peptides (13,442,126,049). The possibility of identifying the top eight ranked peptides by chance is therefore minuscule.

#### Example 7

##### 55 Combinations of grass peptides

[0174] Peptide combinations based on selections from the top 8 ranked peptides in Table 6 were investigated with a

view to identifying optimal vaccines for the treatment of grass allergy. Cytokine assays were performed as in Example 5 for each mix. The results for ten optimal combinations are listed in Table 7. Combination 1 is the most optimal combination and comprises all 8 top ranked peptides. It should be noted that these specific combinations represent a minuscule proportion of the number of possible combinations of peptides originally screened in Examples 1 and 2.

[0175] Optimal grass vaccine mixes were selected on the basis of showing a significant release of IFN- $\gamma$ , IL-10 and IL-13 in a large proportion of the study group of 48 allergic individuals. As such, providing preferred individual grass peptides in combination increases MHC coverage and provides an optimised product of general utility as a grass vaccine. Optimal grass vaccine mixes were also selected on the basis of manufacturing considerations, including physical and chemical characteristics of each peptide in the combination.

10

Table 7

Peptide combination	IL-10 +ve/48	IL-13 +ve/48	IFN- $\gamma$ 10 +ve/48
1	47	41	48
2	47	41	48
3	47	39	48
4	46	39	48
5	44	39	47
6	41	39	46
7	47	40	48
8	46	40	48
9	44	40	48
10	40	40	48

Combinations:

1: Ber01; Ber02; Ber02C; Bio02A; Bio03A; Rye09B; Tim07B; Bio04A  
 2: Ber01; Ber02; Ber02C; Bio02A; Bio03A; Rye09B; Tim07B  
 3: Ber01; Ber02; Ber02C; Rye09B; Tim07B; Bio04A  
 4: Ber01; Ber02; Ber02C; Rye09B; Tim07B  
 5: Ber01; Ber02; Ber02C; Rye09B; Bio04A  
 6: Ber01; Ber02; Ber02C; Rye09B  
 7: Ber01; Ber02C; Bio03A; Rye09B; Tim07B; Bio04A  
 8: Ber01; Ber02C; Rye 09B; Tim07B  
 9: Ber01; Ber02C; Bio03A; Rye09B; Bio04A  
 10: Ber01; Ber02C; Bio03A; Rye09B

40

[0176] Combination 1 is the most optimal combination based on cytokine release characteristics and comprises all 8 top ranked peptides. This combination provides peptides derived from three grass allergens, Cyn d1 (Ber01; Ber02; Ber02C), Lol p 5 (Bio03A; Rye09B, Bio04A) and Phl p 5 (Tim07B). The combination therefore accounts for regional variations in exposure to specific allergens, even though the data from the Canadian study group suggests this may not be an important factor.

45

[0177] It should be noted that each specific optimal combination represents a minuscule proportion of the number of possible combinations of peptides originally screened in the Examples. As outlined above, for example, the eight peptide mix of combination 1 represents one out of 13,442,126,049 potential combinations of the 73 peptides screened in Examples 1 and 2. The optimal combinations each provide a significant release of IFN- $\gamma$ , IL-10 and IL-13 in  $\geq 39/48$  individuals.

50

[0178] It is noteworthy that high MHC coverage is maintained in Combinations 2 to 10 despite significant variation both in the number of peptides and in which specific peptides are included from the top 8 identified in Example 6. It can be seen that even selecting a mix of 4 peptides (Combination 10) still gives 100% coverage for IFN- $\gamma$  response and 83% coverage for IL-10/IL-13 response. Combination 10 is based on 2 peptides from Bermuda grass and 2 peptides from Perennial Rye grass. The coverage obtained with such a mix is surprising, given both the lack of prevalence of these grasses in Canada, and the fact that peptides from Lol p 1, a major allergen recognised by IgE antibodies in grass allergic individuals are not included.

Example 8*Improving solubility*

5 [0179] Two of the top 8 peptides, Rye09B and Tim07B, were identified as having solubility characteristics which could be improved (see Table 6). To improve solubility, various analogues were designed utilising one or more lysine residues at the N- or C-terminus of the peptide. The peptides were assessed in the Epimatrix software to ensure that the modifications had not affected the T cell epitope within the peptide and had not created a neoepitope. Two variant peptides were selected for Rye09B (Rye09B1 & Rye09B2) and Tim07B & Tim07B2). The sequences are shown in Table 8 together  
 10 with the solubility values. The variants of Rye09B are twice as soluble as the native peptide. The variants of Tim07B are more than twenty times as soluble as the native peptide. In addition to improved solubility, both variants also retained their ability to induce T cell cytokine release when tested in a group of 10 subjects. These variants of Rye09B and Tim07B therefore are preferred alternatives to the native peptides, for example in any mix of grass peptides for therapeutic vaccine use.

15

Table 8

Peptide	Sequence	Solubility(mg/ml)	SEQ ID NO:
Rye09B1	KPEVKYAVFEAALTKAIT	10	91
Rye09B2	KKPEVKYAVFEAALTKAIT	10	92
Tim07B 1	KKIPAGELQIIDKIDA	20	93
Tim07B2	KKIPAGELQIIDKIDAK	20	94

25

Example 9Selection of a preferred grass vaccine

30 [0180] During trial manufacturing, one peptide from the top ranked peptides present in Combination 1 above (Bio03A) was found to be difficult to manufacture due to having a string of 5 contiguous hydrophobic amino acids. When assessing the impact of not including Bio03A in a grass vaccine, the overall coverage of the remaining seven top ranked peptides present in Combination 1 was analysed for the 48 subjects. In this seven peptide mix (Ber01, Ber02, Ber02C, Bio02A, Rye09B, Tim07B and Bio04A) Tim07B and Rye07B are optionally substituted for the variants Tim07B 1 and Rye07B1 having improved solubility.

35 [0181] With the 8 peptides present, including Bio03A, 47/48 subjects (IL-10), 41/48 subjects (IL-13) and 48/48 subjects (IFN-gamma) subjects showed positive cytokine responses. In comparison, with seven peptides present, excluding Bio03A, 47/48 subjects (IL-10), 40/48 subjects (IL-13) and 48/48 subjects (IFN-gamma) showed positive cytokine responses. Therefore it was concluded that there would be little impact of not including Bio03A in a grass vaccine for manufacture.

40 [0182] The response properties of the above peptide combinations was compared with whole grass pollen extract containing Timothy, Perennial Rye, and Bermuda grass pollens (Greer Laboratories) and a further positive control, the mitogen SEB. The whole grass pollen extract induced cytokine responses in 11/48 (IL-10), 42/48 (IL-13) and 43/48 (IFN-gamma) whilst the other positive control, the mitogen SEB, induced responses in 47/48 (IL-10), 48/48 (IL-13) and 48/48 (IFN-gamma). The seven peptide mix thus provides strong cytokine responses in similar percentages of the population as for whole grass allergen.

45 [0183] The selection of the peptides Ber01, Ber02, Ber02C, Bio02A, Rye09B, Tim07B and Bio04A for a preferred seven peptide grass vaccine was also based on homology considerations. When these seven peptide sequences are compared between different grasses, there is considerable homology in many cases increasing the probability that grass-allergic individuals will respond to the peptides. This means that the peptides have utility even in those subjects whose dominant allergic response is to a grass other than Bermuda, Timothy or Perennial rye. Homologous sequences from other grasses for the above seven peptides are shown as SEQ ID NOs 74 to 90 above. For example, the Ber01 peptide contains the 9 mer epitope FGAMAKKGQ which has close homologues in many other common grasses.

50 [0184] Furthermore, the seven vaccine peptides are derived from both group 1 and group 5 allergens from the three most prevalent grasses (Timothy, Perennial Rye and Bermuda grasses) and include epitopes with complete or significant homology to the other common grasses (Orchard, Velvet, Kentucky blue and Canary). The seven peptide sequences therefore maximise the breadth of coverage of grass pollen allergic individuals making the vaccine suitable for treating all grass pollen allergic individuals.

Example 10

## Histamine release assay

5 [0185] The purpose of this assay was to identify whether the preferred seven peptide combination of Example 9 was capable of activating blood basophils (as a surrogate for tissue mast cells) resulting in histamine release that may result in allergic reactions during therapy. A combination of peptides that induces histamine release frequently may be considered unsuitable for use as a peptide vaccine.

10 [0186] Histamine release requires the crosslinking of adjacent specific IgE molecules on the surface of the basophil. The peptides being evaluated were small (10 to 18 amino acids in length) and should not, therefore, possess significant tertiary structure that would enable them to retain the conformation of an IgE-binding epitope of the whole molecule. Furthermore, peptide monomers in solution, even if they are bound by IgE, should not be able to crosslink adjacent IgE molecules. Histamine release from peripheral blood basophils isolated from peripheral whole blood obtained from grass allergic subjects was evaluated. Peripheral blood basophils were used as a surrogate for tissue mast cells which were not practical to assay. The assay required  $3 \times 10^6$  peripheral blood mononuclear cells (PBMC) per subject. PBMCs were incubated *in vitro* with the seven grass peptides from Example 9 in combination. Histamine release in response to whole Grass pollen allergen extract containing Timothy, Perennial Rye, and Bermuda grass pollens (Greer Laboratories) was included as a control. A positive control, representing total histamine release, generated by freeze/thawing the cells twice was also included in each assay.

15 [0187] Histamine concentrations were measured by ELISA and results expressed as a percentage of the positive control (% positive control). The assay was performed using the Immunotech Histamine Release Immunoassay kit according to the manufacturer's instructions. Following the *in vitro* challenge of PBMCs with peptides, peptide mixes, whole allergen or buffer in microtitre plate wells, supernatants were removed and the histamine in the samples converted to acyl histamine. Acylated samples were tested by a competitive acyl histamine ELISA.

20 [0188] Peptides were assayed for their ability to induce histamine release over a 5 log10 range (1 to 10,000 ng/mL). The concentration range assayed was selected based on theoretical *in vivo* doses of peptide that may be achieved during therapy. Based on delivery of peptides by intradermal injection, high local peptide concentrations of up to 10  $\mu\text{g}/\text{mL}$  per peptide could be present. Although unlikely, there is a risk that the full dose may be injected into the bloodstream. In this unlikely event, the maximum clinical dose of 20  $\mu\text{g}$  (12 nmole) of each peptide entering a blood volume of 5 litres, 25 would result in a theoretical maximum blood concentration of 4.0 ng/mL. This is at the lower end of the histamine release assay dose range and 2000 times lower than the top concentration used in the assay.

30 [0189] A whole grass allergen preparation containing Timothy, Perennial Rye and Bermuda grass pollens (Greer Laboratories) was used as a control for histamine release over a 5 log10 range from 10 to 100,000 ng/mL. A negative control for spontaneous histamine release was generated by incubating cells in buffer only.

35 [0190] Single measurements were performed for each dilution. After completion of the ELISA, individual histamine levels were determined by interpolation from the standard curve generated in the ELISA assay. Results from samples were adjusted to allow for dilution. Where two or more consecutive dilutions of a peptide/allergen preparation elicited >15% of the total histamine release seen in the freeze thawed positive control (>15% of positive control), or where a single value of >15% of positive control was achieved at the highest concentration tested (10  $\mu\text{g}/\text{mL}$  for peptides), this 40 was considered a "positive histamine release".

45 [0191] A total of 45 histamine release assays were completed during the study. Of these, 3 assays were rejected, due to unacceptably high levels (>15%) of histamine release in the medium plus buffer negative control wells or no response at all in the positive control wells. Therefore a total of 42 subjects were included in the analysis. The study findings are summarised in Table 9.

Protein Concentration ( $\mu\text{g}/\text{mL}$ )	Subjects with positive histamine release (>=15%)	Mean histamine release (% pos control) n=42	Range (% pos control)
Seven peptide combination * (10)	0/42	0	0-4
Whole grass (100)	32/42	41	2-152
Whole grass (0.01)	21/42	24	0-84

55 \* Seven peptide combination: Ber01, Ber02, Ber02C, Bio02A, Rye09B1, Tim07B 1 and Bio04A.

[0192] The whole grass allergen preparation induced 15% or higher histamine release in 32/42 (76%) of subjects at 100  $\mu\text{g}/\text{mL}$ . Even at the lowest concentration of 10  $\mu\text{g}/\text{mL}$ , the whole allergen induced high levels of histamine release

in 21/42 individuals (50%).

**[0193]** In contrast, the seven peptide combination failed to elicit significant histamine release in any of the 42 subjects tested, even at the highest concentration for which data is shown, where each peptide was present at 10 µg/mL. This is a concentration 1000 fold higher than the concentration of whole allergen still giving very high levels of histamine release in over half of the subjects (10 ng/mL). The seven peptide combination therefore has negligible potential for causing IgE mediated reactions even in highly sensitive individuals.

**[0194]** Given the large excess of peptide dose tested in this assay compared to likely concentrations of the peptides following clinical dosing of a grass vaccine, it is not anticipated that administration of the seven peptide combination would cause significant histamine release either by IgE-mediated or direct peptide-mediated basophil or mast cell activation and degranulation. The histamine release data indicate that the seven peptide combination does not induce the activation of basophils and therefore has a very low potential for inducing IgE-mediated acute allergic reactions in grass allergic individuals, either systemically or locally in the skin.

#### ASPECTS OF THE INVENTION

15

**[0195]**

1. A composition suitable for use in preventing or treating allergy to grass pollen by tolerisation comprising:

20 (a) at least one of the polypeptides Tim07B (KIPAGELQIIDKIDA), Tim 10B (KYTVFETALKAITAMSE), Tim 04A (WGAIWRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof;  
 (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), Ber02C (KSS-WGAIWRIDPKKPLK) and Ber 02B KDSDEFIPMKSSWGAiWR or a variant of any thereof; and  
 25 (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTAKIT), Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIPSLEAAVKQAY), Rye 05C (NAGFKAAVAAAANAPPK), or a variant of any thereof,

wherein said variant is:

30 I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a), (b) or (c), or  
 II) a polypeptide of 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a), (b) or (c), and that is capable of tolerising to said corresponding polypeptide; or  
 35 III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a), (b) or (c), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

40 2. A composition according to aspect 1 comprising:

a) the polypeptide Tim07B (KIPAGELQIIDKIDA) or a variant thereof,  
 b) the polypeptide Ber01 (SGKAFGAMAKKGQED) or a variant thereof, and  
 c) the polypeptide Bio04A (LKKAVTAMSEAEK) or a variant thereof.

45 3. A composition suitable for use in preventing or treating allergy to grass pollen by tolerisation comprising at least four different polypeptides selected from:

50 (a) Tim07B (KIPAGELQIIDKIDA) or a variant thereof;  
 (b) Ber01 (SGKAFGAMAKKGQED) or a variant thereof;  
 (c) Bio04A (LKKAVTAMSEAEK) or a variant thereof;  
 (d) Rye09B (PEVKYAVFEAALTAKIT) or a variant thereof;  
 (e) Ber02 (FIPMKSSWGA) or a variant thereof;  
 (f) Ber02C (KSSWGAiWRIDPKKPLK) or a variant thereof;  
 55 (g) Bio03A (KFIPTLVAAVKQAYAAKQ) or a variant thereof; and  
 (h) Bio02A (KYDAYVATLTEALR) or a variant thereof;

wherein said variant is:

5 I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a) to (h), or

II) a polypeptide 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a) to (h), and that is capable of tolerising to said corresponding polypeptide; or

10 III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a) to (h), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

15 4. A composition according to aspect 3 comprising

(a) the polypeptide Tim07B (KIPAGELQIIDKIDA) or a variant thereof;

15 (b) at least one of the polypeptides of Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), and Ber02C (KSSWGAIWRIDPKKPLK) or a variant of any thereof; and

(c) at least one of the polypeptides of Bio04A (LKKAVTAMSEAEK), ye09B (PEVKYAVFEAALTAKIT), Bio02A (KYDAYVATLTEALR), and Bio03A (KFIPTLVAVKQAYAAKQ) or a variant of any thereof; and

(a) at least one additional polypeptide of b) or c) not selected above.

20 5. A composition according to aspect 4, comprising the polypeptide Tim07B (KIPAGELQIIDKIDA) or a variant thereof, the polypeptide Ber01 (SGKAFGAMAKKGQED) or a variant thereof, the polypeptide Bio04A (LKKAVTAMSEAEK) or a variant thereof, and at least one additional polypeptide of b) or c) not selected above.

25 6. The composition according to any one of the preceding aspects further comprising one or more polypeptides of any of SEQ ID NOs 1 to 73 as shown in Tables 2 to 4 or variants thereof, not previously selected above.

7. The composition according to any one of the preceding aspects, wherein the composition:

30 - is capable of tolerising at least 50% or at least 60% of a panel of grass pollen allergic individuals in the population, and/or

- comprises at least one further polypeptide up to a total of fourteen different polypeptides, wherein the further polypeptides:

35 (a) comprise a sequence having at least 65% sequence identity to at least 9 or more contiguous amino acids in any of SEQ ID NOs: 1 to 73 not selected above; and

(b) are 9 to 30 amino acids in length.

40 8. The composition according to any one of the preceding aspects, comprising at least one said polypeptide which is 9 to 20 or 13 to 17 amino acids in length.

45 9. A composition according to any one of the preceding aspects which comprises the polypeptides of any one of Combinations 1 to 10 as set forth in Table 7, wherein optionally one or more of said polypeptides can be substituted for a variant thereof as defined in 1 (I) or (II).

10. A composition according to any one of the preceding aspects which comprises the polypeptides:

(a) Tim07B1 (KKIPAGELQIIDKIDA);  
 (b) Ber01 (SGKAFGAMAKKGQED);  
 (c) Bio04A (LKKAVTAMSEAEK);  
 50 (d) Rye09B1 (KPEVKYAVFEAALTAKIT);  
 (e) Ber02 (FIPMKSSWGA);  
 (f) Ber02C (KSSWGAIWRIDPKKPLK); and  
 (g) Bio02A (KYDAYVATLTEALR)

55 11. The composition according to any one of the preceding aspects, wherein one or more of the polypeptides have one or more modifications selected from the following:

(i) N terminal acetylation;

5 (ii) C terminal amidation;  
 (iii) one or more hydrogens on the side chain amines of Arginine and/or Lysine replaced with a methylene group;  
 (iv) glycosylation; and  
 (v) phosphorylation.

10 12. The composition according to any one of the preceding aspects wherein at least one of the peptides has been engineered to be soluble such that it comprises:

15 (i) N terminal to the residues of the peptide which flank a T cell epitope: one to six contiguous amino acids corresponding to the two to six contiguous amino acids immediately N terminal to said residues in the sequence of the protein from which the peptide derives; and/or  
 (ii) C terminal to the residues of the peptide which flank a T cell epitope: one to six contiguous amino acids corresponding to the one to six contiguous amino acids immediately C terminal to the said residues in the sequence of the protein from which the peptide derives; or  
 (iii) both N and C terminal to the residues of the peptide which flank a T cell epitope, at least one amino acid selected from arginine, lysine, histidine, glutamate and aspartate,

20 wherein the polypeptide has a solubility of at least 3.5mg/ml and the T cell epitope has a solubility of less than 3.5mg/ml.

25 13. The composition according to any one of the preceding aspects wherein at least one of the peptides has been engineered to be soluble such that additionally:

30 (i) any cysteine residues in the native sequence of the peptide are replaced with serine or 2-aminobutyric acid; and /or  
 (ii) hydrophobic residues in the up to three amino acids at the N or C terminus of the native sequence of the peptide, which are not comprised in a T cell epitope, are deleted; and/or  
 (iii) any two consecutive amino acids comprising the sequence Asp-Gly in the up to four amino acids at the N or C terminus of the native sequence of the peptide, which are not comprised in a T cell epitope, are deleted; and/or  
 (iv) one or more positively charged residues are added at the N-and/or C-terminus.

35 14. The composition according to any one of the preceding aspects wherein each polypeptide has a concentration in the range of 0.03 to 200 nmol/ml, 0.3 to 200 nmol/ml, 50 to 200 nmol/ml or 30 to 120 nmol/ml.

40 15. A composition for use in preventing or treating allergy to grass pollen by tolerisation comprising at least one polynucleotide sequence which when expressed causes the production of a composition as defined in any one of aspects 1 to 13.

16. A vector for use in preventing or treating allergy to grass pollen by tolerisation comprising four or more polynucleotide sequences which each encode a different polypeptide as defined in any one of aspects 1 to 13.

45 17. A product comprising:

(a) the polypeptide Tim07B (KIPAGELQIIDKIDA), Tim 10B (KYTVFETALKAITAMSE), Tim 04A (WGAI-WRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof as defined in aspect 1 (I) or (II);  
 (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED) Ber02 (FIPMKSSWGA), Ber02C (KSS-WGAIWRIDPKKPLK) and Ber02B KDSDEFIPMKSSWGAIWR or a variant of any thereof as defined in aspect 1 (I) or (II); and  
 (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTAKIT), and Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIPSLEAAVKQAY), Rye 05C (NAGFKAAVAAAANAPPK) or a variant of any thereof as defined in aspect 1 (I) or (II),

55 wherein each different polypeptide is for simultaneous, separate or sequential use in preventing or treating allergy to grass pollen by tolerisation.

18. A pharmaceutical formulation for use in preventing or treating allergy to grass pollen by tolerisation comprising a composition according to any one of aspects 1 to 15; a vector according to aspect 16; or a product according to aspect 17; and a pharmaceutically acceptable carrier or diluent.

**EP 2 891 662 A1**

19. The composition, vector or product according to aspect 18, formulated for oral administration, topical administration, nasal administration, subcutaneous administration, sublingual administration, intradermal administration, buccal administration, epidermal or patch administration or for administration by inhalation or by injection.

5 20. The composition as defined in any one of aspects 1 to 15 or product as defined in aspect 17, additionally comprising a further polypeptide allergen for use in tolerising an individual to the further polypeptide allergen.

10 21. An in vitro method of determining whether T cells recognize a composition as defined in any one of aspects 1 to 14 comprising contacting said T cells with said polypeptide and detecting whether said T cells are stimulated by said polypeptide.

22. A method according to aspect 21 which is carried out to determine whether an individual has, or is at risk of having, an allergy to grass pollen.

15

20

25

30

35

40

45

50

55

## SEQUENCE LISTING

5 <110> CIRCASSIA LIMITED

10 <120> GRASS PEPTIDES FOR VACCINE

15 <130> N102658F-EP

20 <150> GB 0901928.2

25 <151> 2009-02-05

30 <150> GB 0901927.4

35 <151> 2009-02-05

40 <150> GB 0912578.2

45 <151> 2009-07-20

50 <150> PCT/GB09/01995

55 <151> 2009-08-14

60 <150> GB 0917871.6

65 <151> 2009-10-12

70 <160> 144

75 <170> PatentIn version 3.5

80 <210> 1

85 <211> 15

90 <212> PRT

95 <213> Cynodon dactylon

100 <400> 1

105 Ser Gly Lys Ala Phe Gly Ala Met Ala Lys Lys Gly Gln Glu Asp  
1 5 10 15

110 <210> 2

115 <211> 10

120 <212> PRT

125 <213> Cynodon dactylon

130 <400> 2

135 Phe Ile Pro Met Lys Ser Ser Trp Gly Ala  
1 5 10

140 <210> 3

145 <211> 13

150 <212> PRT

155 <213> Cynodon dactylon

160 <400> 3

165 Trp Gly Ala Ile Trp Arg Ile Asp Pro Lys Lys Pro Leu  
1 5 10

170 <210> 4

175 <211> 18

180 <212> PRT

&lt;213&gt; Cynodon dactylon

&lt;400&gt; 4

5	Lys	Asp	Ser	Asp	Glu	Phe	Ile	Pro	Met	Lys	Ser	Ser	Trp	Gly	Ala	Ile
	1				5					10					15	

Trp Arg

10

&lt;210&gt; 5

&lt;211&gt; 17

&lt;212&gt; PRT

15

&lt;213&gt; Cynodon dactylon

&lt;400&gt; 5

20

Lys	Ser	Ser	Trp	Gly	Ala	Ile	Trp	Arg	Ile	Asp	Pro	Lys	Lys	Pro	Leu
1					5				10					15	

Lys

25

&lt;210&gt; 6

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Cynodon dactylon

30

&lt;400&gt; 6

Met	Lys	Ser	Ser	Trp	Gly	Ala	Ile	Trp	Arg	Ile	Asp	Pro	Lys	Lys	Pro
1						5				10				15	

35

Leu Lys

40

&lt;210&gt; 7

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Cynodon dactylon

&lt;400&gt; 7

45

Met	Lys	Ser	Ser	Trp	Gly	Ala	Ile	Trp	Arg	Ile	Asp	Pro	Pro	Lys	Pro
1						5				10				15	

Leu Lys

50

&lt;210&gt; 8

&lt;211&gt; 14

&lt;212&gt; PRT

55

&lt;213&gt; Cynodon dactylon

&lt;400&gt; 8

Ile Pro Ala Asn Trp Lys Pro Asp Thr Val Tyr Thr Ser Lys  
 1 5 10

5 <210> 9  
 <211> 14  
 <212> PRT  
 <213> *Cynodon dactylon*

10 <400> 9  
 Lys Ala Thr Phe Tyr Gly Ser Asp Pro Arg Gly Ala Ala Pro  
 1 5 10

15 <210> 10  
 <211> 12  
 <212> PRT  
 <213> *Cynodon dactylon*

20 <400> 10  
 Ala Tyr His Phe Asp Leu Ser Gly Lys Ala Phe Gly  
 1 5 10

25 <210> 11  
 <211> 15  
 <212> PRT  
 <213> *Lolium perenne*  
 30 <400> 11  
 Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly  
 1 5 10 15

35 <210> 12  
 <211> 17  
 <212> PRT  
 <213> *Lolium perenne*  
 40 <400> 12  
 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala  
 1 5 10 15

45 Gly  
 50 <210> 13  
 <211> 17  
 <212> PRT  
 <213> *Lolium perenne*  
 55 <400> 13  
 Phe Gly Ser Met Ala Lys Lys Gly Glu Glu Gln Asn Val Arg Ser Ala  
 1 5 10 15

## Gly

5                   <210> 14  
                   <211> 18  
                   <212> PRT  
                   <213> *Lolium perenne*

10                  <400> 14  
                   His Ala Phe Gly Met Ala Lys Lys Gly Glu Glu Gln Asn Val Arg Ser  
                   1                                   5                                   10                           15

15                  Ala Gly

20                  <210> 15  
                   <211> 14  
                   <212> PRT  
                   <213> *Lolium perenne*

25                  <400> 15  
                   Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys Tyr Val Asp  
                   1                                   5                                   10

30                  <210> 16  
                   <211> 13  
                   <212> PRT  
                   <213> *Lolium perenne*

35                  <400> 16  
                   Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu  
                   1                                   5                                   10

40                  <210> 17  
                   <211> 16  
                   <212> PRT  
                   <213> *Lolium perenne*

45                  <400> 17  
                   Lys Glu Ser Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu  
                   1                                   5                                   10                           15

50                  <210> 18  
                   <211> 18  
                   <212> PRT  
                   <213> *Lolium perenne*

55                  <400> 18  
                   Lys Glu Ser Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu  
                   1                                   5                                   10                           15

## Gly Pro

5 &lt;210&gt; 19

&lt;211&gt; 14

&lt;212&gt; PRT

<213> *Lolium perenne*

10 &lt;400&gt; 19

Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ser  
1 5 10

15 &lt;210&gt; 20

&lt;211&gt; 23

&lt;212&gt; PRT

<213> *Phleum pratense*

20 &lt;400&gt; 20

Gly Asp Glu Gln Lys Leu Arg Ser Ala Gly Glu Leu Glu Leu Gln Phe  
1 5 10 1525 Arg Arg Val Lys Cys Lys Tyr  
20

30 &lt;210&gt; 21

&lt;211&gt; 19

&lt;212&gt; PRT

<213> *Phleum pratense*

&lt;400&gt; 21

35 Asn Pro Asn Tyr Leu Ala Leu Leu Val Lys Tyr Val Asn Gly Asp Gly  
1 5 10 15

40 Asp Val Val

45 &lt;210&gt; 22

&lt;211&gt; 19

&lt;212&gt; PRT

<213> *Phleum pratense*

45 &lt;400&gt; 22

50 Lys Glu Ser Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys Leu  
1 5 10 15

55 Thr Gly Pro

55 &lt;210&gt; 23

&lt;211&gt; 13

<212> PRT  
 <213> *Phleum pratense*

<400> 23

5

Pro Glu Gly Trp Lys Ala Asp Thr Ser Tyr Glu Ser Lys  
 1 5 10

10

<210> 24  
 <211> 17  
 <212> PRT  
 <213> *Phleum pratense*

<400> 24

15

Phe Gly Ala Met Ala Lys Lys Gly Asp Glu Gln Lys Leu Arg Ser Ala  
 1 5 10 15

20

Gly

25

<210> 25  
 <211> 19  
 <212> PRT  
 <213> *Phleum pratense*

<400> 25

30

His Ala Phe Gly Ala Met Ala Lys Lys Gly Asp Glu Gln Lys Leu Arg  
 1 5 10 15

Ser Ala Gly

35

<210> 26  
 <211> 16  
 <212> PRT  
 <213> *Phleum pratense*

40

<400> 26

Ser Asn Pro Asn Tyr Leu Ala Leu Leu Val Lys Tyr Val Asn Gly Asp  
 1 5 10 15

45

<210> 27  
 <211> 13  
 <212> PRT  
 <213> *Phleum pratense*

50

<400> 27

Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys Leu  
 1 5 10

55

<210> 28  
 <211> 14

<212> PRT  
 <213> **Lolium perenne**

<400> 28

5

Lys Tyr Asp Ala Tyr Val Ala Thr Leu Thr Glu Ala Leu Arg  
 1 5 10

10

<210> 29  
 <211> 18  
 <212> PRT  
 <213> **Lolium perenne**

<400> 29

15

Lys Phe Ile Pro Thr Leu Val Ala Ala Val Lys Gln Ala Tyr Ala Ala  
 1 5 10 15

20

**Lys Gln**

25

<210> 30  
 <211> 19  
 <212> PRT  
 <213> **Lolium perenne**

<400> 30

30

Tyr Lys Phe Ile Pro Thr Leu Val Ala Ala Val Lys Gln Ala Tyr Ala  
 1 5 10 15

**Ala Lys Gln**

35

<210> 31  
 <211> 13  
 <212> PRT  
 <213> **Lolium perenne**

40

<400> 31

Leu Lys Lys Ala Val Thr Ala Met Ser Glu Ala Glu Lys  
 1 5 10

45

<210> 32  
 <211> 17  
 <212> PRT  
 <213> **Lolium perenne**

50

<400> 32

Pro Glu Thr Ala Leu Lys Lys Ala Val Thr Ala Met Ser Glu Ala Glu  
 1 5 10 15

55

**Lys**

<210> 33  
 <211> 15  
 <212> PRT  
 <213> **Lolium perenne**

5

<400> 33

Lys Phe Thr Val Phe Glu Ser Ala Phe Asn Lys Ala Leu Asn Glu  
 1 5 10 15

10

<210> 34  
 <211> 18  
 <212> PRT  
 <213> **Lolium perenne**

15

<400> 34

Phe Lys Ala Ala Val Ala Ala Ala Asn Ala Pro Pro Ala Asp Lys  
 1 5 10 15

20

**Phe Lys**

25

<210> 35  
 <211> 17  
 <212> PRT  
 <213> **Lolium perenne**

30

<400> 35

Asn Ala Gly Phe Lys Ala Ala Val Ala Ala Ala Asn Ala Pro Pro  
 1 5 10 15

35

**Lys**

40

<210> 36  
 <211> 14  
 <212> PRT  
 <213> **Lolium perenne**

45

Lys Tyr Asp Ala Phe Val Thr Ala Leu Thr Glu Ala Leu Arg  
 1 5 10

50

<210> 37  
 <211> 17  
 <212> PRT  
 <213> **Lolium perenne**

55

Pro Glu Ala Lys Tyr Asp Ala Phe Val Thr Ala Leu Thr Glu Ala Leu  
 1 5 10 15

## Arg

5           <210> 38  
       <211> 14  
       <212> PRT  
       <213> *Lolium perenne*

10          <400> 38  
           Gly Glu Leu Gln Ile Val Asp Lys Ile Asp Ala Ala Phe Lys  
           1                           5                           10

15          <210> 39  
       <211> 15  
       <212> PRT  
       <213> *Lolium perenne*

20          <400> 39  
           Lys Ile Pro Thr Gly Glu Leu Gln Ile Val Asp Lys Ile Asp Ala  
           1                           5                           10                   15

25          <210> 40  
       <211> 16  
       <212> PRT  
       <213> *Lolium perenne*

30          <400> 40  
           Phe Lys Ile Ala Ala Thr Ala Ala Asn Ala Ala Pro Thr Asn Asp Lys  
           1                           5                           10                   15

35          <210> 41  
       <211> 17  
       <212> PRT  
       <213> *Lolium perenne*

40          <400> 41  
           Ala Phe Lys Ile Ala Ala Thr Ala Ala Asn Ala Ala Pro Thr Asn Asp  
           1                           5                           10                   15

45          Lys

50          <210> 42  
       <211> 15  
       <212> PRT  
       <213> *Lolium perenne*

55          <400> 42  
           Tyr Lys Phe Ile Pro Ser Leu Glu Ala Ala Val Lys Gln Ala Tyr  
           1                           5                           10                   15

<210> 43  
 <211> 17  
 <212> PRT  
 <213> **Lolium perenne**

5

<400> 43

Glu Thr Tyr Lys Phe Ile Pro Ser Leu Glu Ala Ala Val Lys Gln Ala  
 1 5 10 15

10

**Tyr**

15

<210> 44  
 <211> 13  
 <212> PRT  
 <213> **Lolium perenne**

20

<400> 44

Asp Ser Tyr Lys Phe Ile Pro Thr Leu Val Ala Val Lys  
 1 5 10

25

<210> 45  
 <211> 16  
 <212> PRT  
 <213> **Lolium perenne**

30

<400> 45

Leu Thr Lys Ala Ile Thr Ala Met Thr Gln Ala Gln Lys Ala Gly Lys  
 1 5 10 15

35

<210> 46  
 <211> 17  
 <212> PRT  
 <213> **Lolium perenne**

40

<400> 46

Pro Glu Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala Ile  
 1 5 10 15

45

**Thr**

50

<210> 47  
 <211> 17  
 <212> PRT  
 <213> **Lolium perenne**

<400> 47

55

Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala Ile Thr Ala Met  
 1 5 10 15

**Thr**

5           <210> 48  
       <211> 15  
       <212> PRT  
       <213> *Lolium perenne*

10          <400> 48  
           Asp Lys Phe Lys Ile Phe Glu Ala Ala Phe Ser Glu Ser Ser Lys  
           1                           5                                   10                           15

15          <210> 49  
       <211> 13  
       <212> PRT  
       <213> *Lolium perenne*

20          <400> 49  
           Thr Pro Leu Arg Arg Thr Ser Ser Arg Ser Ser Arg Pro  
           1                           5                                   10

25          <210> 50  
       <211> 14  
       <212> PRT  
       <213> *Lolium perenne*

30          <400> 50  
           Asp Val Ala Tyr Lys Ala Ala Glu Ala His Pro Arg Gly Gln  
           1                           5                                   10

35          <210> 51  
       <211> 13  
       <212> PRT  
       <213> *Lolium perenne*

40          <400> 51  
           Ala Leu Arg Val Ile Ala Gly Thr Leu Glu Val His Ala  
           1                           5                                   10

45          <210> 52  
       <211> 14  
       <212> PRT  
       <213> *Lolium perenne*

50          <400> 52  
           Phe Glu Asn Thr Phe Asn Asn Ala Ile Lys Val Ser Leu Gly  
           1                           5                                   10

55          <210> 53  
       <211> 18

<212> PRT  
 <213> *Phleum pratense*

<400> 53

5

Lys Tyr Thr Val Phe Glu Thr Ala Leu Lys Lys Ala Ile Thr Ala Met  
 1 5 10 15

10 Ser Glu

<210> 54

<211> 9

15 <212> PRT

<213> *Phleum pratense*

<400> 54

20 Leu Lys Lys Ala Ile Thr Ala Met Ser  
 1 5

25 <210> 55

<211> 13

<212> PRT

<213> *Phleum pratense*

<400> 55

30 Pro Glu Thr Ala Leu Lys Lys Ala Ile Thr Ala Met Ser  
 1 5 10

35 <210> 56

<211> 19

<212> PRT

<213> *Phleum pratense*

<400> 56

40 Ile Glu Lys Ile Asn Ala Gly Phe Lys Ala Ala Leu Ala Gly Ala Gly  
 1 5 10 15

Val Gln Pro

45

<210> 57

<211> 24

<212> PRT

<213> *Phleum pratense*

50

<400> 57

Gly Ala Gly Val Gln Pro Ala Asp Lys Tyr Arg Thr Phe Val Ala Thr  
 1 5 10 15

55

Phe Gly Pro Ala Ser Asn Lys Ala  
 20

<210> 58  
 <211> 25  
 <212> PRT  
 <213> *Phleum pratense*

5

<400> 58

Ala Leu Thr Ser Lys Leu Asp Ala Ala Tyr Lys Leu Ala Tyr Lys Thr  
 1 5 10 15

10

Ala Glu Gly Ala Thr Pro Glu Ala Lys  
 20 25

15

<210> 59  
 <211> 21  
 <212> PRT  
 <213> *Phleum pratense*

20

Ile Glu Lys Val Asp Ala Ala Phe Lys Val Ala Ala Thr Ala Ala Asn  
 1 5 10 15

25

Ala Ala Pro Ala Asn  
 20

30

<210> 60  
 <211> 21  
 <212> PRT  
 <213> *Phleum pratense*

<400> 60

35

Tyr Glu Ser Tyr Lys Phe Ile Pro Ala Leu Glu Ala Ala Val Lys Gln  
 1 5 10 15

40

Ala Tyr Ala Ala Thr  
 20

45

<210> 61  
 <211> 22  
 <212> PRT  
 <213> *Phleum pratense*

<400> 61

50

Glu Val Lys Tyr Thr Val Phe Glu Thr Ala Leu Lys Lys Ala Ile Thr  
 1 5 10 15

Ala Met Ser Glu Ala Gln  
 20

55

<210> 62

<211> 21  
 <212> PRT  
 <213> Phleum pratense

5 <400> 62

Asp Ile Asn Val Gly Phe Lys Ala Ala Val Ala Ala Ala Ala Ser Val  
 1 5 10 15

10 Pro Ala Ala Asp Lys  
20

<210> 63  
 <211> 24  
 <212> PRT  
 <213> Phleum pratense

15 <400> 63

Phe Asp Ser Phe Val Ala Ser Leu Thr Glu Ala Leu Arg Val Ile Ala  
 1 5 10 15

Gly Ala Leu Glu Val His Ala Val  
 20

25

<210> 64  
 <211> 19  
 <212> PRT  
 <213> Phleum pratense

30 <400> 64

Ile Asp Lys Ile Asp Ala Ala Phe Lys Val Ala Ala Thr Ala Ala Ala  
 1 5 10 15

35

Thr Ala Pro

40 <210> 65  
<211> 27  
<212> PRT  
<213> Phleum pratense

45 <400> 65

Ile Pro Ser Leu Glu Ala Ala Val Lys Gln Ala Tyr Ala Ala Thr Val  
 1 5 10 15

50 Ala Ala Ala Pro Gln Val Lys Tyr Ala Val Phe  
20 25

55 <210> 66  
 <211> 19  
 <212> PRT  
 <213> Phleum pratense

&lt;400&gt; 66

Tyr	Ala	Val	Phe	Glu	Ala	Ala	Leu	Thr	Lys	Ala	Ile	Thr	Ala	Met	Ser
1				5					10				15		

5

Glu Val Gln

10

&lt;210&gt; 67

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

15

&lt;400&gt; 67

Phe	Lys	Ala	Ala	Val	Ala	Ala	Ala	Ala	Ser	Val	Pro	Ala	Ala	Asp	Lys
1				5					10			15			

20

Phe Lys

25

&lt;210&gt; 68

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

&lt;400&gt; 68

30

Lys	Phe	Asp	Ser	Phe	Val	Ala	Ser	Leu	Thr	Glu	Ala	Leu	Arg
1				5					10				

35

&lt;210&gt; 69

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

&lt;400&gt; 69

40

Lys	Ile	Pro	Ala	Gly	Glu	Leu	Gln	Ile	Ile	Asp	Lys	Ile	Asp	Ala
1				5					10		15			

45

&lt;210&gt; 70

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

&lt;400&gt; 70

50

Phe	Lys	Val	Ala	Ala	Thr	Ala	Ala	Asn	Ala	Ala	Pro	Ala	Asn	Asp	Lys
1				5					10			15			

55

&lt;210&gt; 71

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

&lt;400&gt; 71

Tyr	Lys	Phe	Ile	Pro	Ala	Leu	Glu	Ala	Ala	Val	Lys	Gln	Ala	Tyr
1						5				10			15	

5

&lt;210&gt; 72

&lt;211&gt; 19

&lt;212&gt; PRT

10 <213> **Phleum pratense**

&lt;400&gt; 72

Pro	Glu	Glu	Ser	Tyr	Lys	Phe	Ile	Pro	Ala	Leu	Glu	Ala	Ala	Val	Lys
1										5			10		15

15

**Gln Ala Tyr**

20

&lt;210&gt; 73

&lt;211&gt; 16

&lt;212&gt; PRT

<213> **Phleum pratense**

25 &lt;400&gt; 73

Leu	Thr	Lys	Ala	Ile	Thr	Ala	Met	Ser	Glu	Val	Gln	Lys	Val	Ser	Gln
1										5			10		15

30

&lt;210&gt; 74

&lt;211&gt; 15

&lt;212&gt; PRT

<213> **Holcus lanatus**

35 &lt;400&gt; 74

Ser	Gly	Lys	Ala	Phe	Gly	Ala	Met	Ala	Lys	Lys	Gly	Glu	Glu	Asp
1									5			10		15

40

&lt;210&gt; 75

&lt;211&gt; 15

&lt;212&gt; PRT

<213> **Dactylis glomerata**

45 &lt;400&gt; 75

Ser	Gly	His	Ala	Phe	Gly	Ser	Met	Ala	Lys	Lys	Gly	Glu	Glu	Asp
1									5			10		15

50

&lt;210&gt; 76

&lt;211&gt; 15

&lt;212&gt; PRT

<213> **Phleum pratense**

55 &lt;400&gt; 76

Ser	Gly	Ile	Ala	Phe	Gly	Ser	Met	Ala	Lys	Lys	Gly	Asp	Glu	Asp
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

	1	5	10	15
5	<210> 77			
	<211> 15			
	<212> PRT			
	<213> Zea mays			
	<400> 77			
10	Ser Gly Thr Ala Phe Gly Ala Met Ala Lys Lys Gly Glu Glu Glu			
	1	5	10	15
15	<210> 78			
	<211> 14			
	<212> PRT			
	<213> Hordeum vulgare			
	<400> 78			
20	Lys Tyr Asp Ala Tyr Val Ala Thr Leu Thr Glu Ser Leu Arg			
	1	5	10	
25	<210> 79			
	<211> 14			
	<212> PRT			
	<213> Phleum pratense			
	<400> 79			
30	Lys Tyr Asp Ala Tyr Val Ala Thr Leu Ser Glu Ala Leu Arg			
	1	5	10	
35	<210> 80			
	<211> 14			
	<212> PRT			
	<213> Dactylis glomerata			
	<400> 80			
40	Lys Tyr Asp Ala Phe Val Ala Ala Leu Thr Glu Ala Leu Arg			
	1	5	10	
45	<210> 81			
	<211> 14			
	<212> PRT			
	<213> Holcus lanatus			
	<400> 81			
50	Lys Tyr Asp Ala Phe Val Thr Thr Leu Thr Glu Ala Leu Arg			
	1	5	10	
55	<210> 82			
	<211> 16			
	<212> PRT			
	<213> Dactylis glomerata			

&lt;400&gt; 82

Lys	Phe	Ile	Pro	Thr	Leu	Glu	Ala	Ala	Val	Lys	Gln	Ala	Tyr	Ala	Ala
1					5					10				15	

5

&lt;210&gt; 83

&lt;211&gt; 16

&lt;212&gt; PRT

10 &lt;213&gt; Phleum pratense

&lt;400&gt; 83

Lys	Phe	Ile	Pro	Ala	Leu	Glu	Ala	Ala	Val	Lys	Gln	Ala	Tyr	Ala	Ala
1				5						10				15	

15

&lt;210&gt; 84

&lt;211&gt; 15

&lt;212&gt; PRT

20 &lt;213&gt; Dactylis glomerata

&lt;400&gt; 84

Lys	Ile	Pro	Ala	Gly	Glu	Leu	Gln	Ile	Val	Asp	Lys	Ile	Asp	Ala	
1				5						10				15	

25

&lt;210&gt; 85

&lt;400&gt; 85

000

30

&lt;210&gt; 86

&lt;211&gt; 15

&lt;212&gt; PRT

35 &lt;213&gt; Poa pratensis

&lt;400&gt; 86

Lys	Ile	Pro	Ala	Gly	Glu	Gln	Gln	Ile	Ile	Asp	Lys	Ile	Asp	Ala	
1				5						10				15	

40

&lt;210&gt; 87

&lt;211&gt; 13

&lt;212&gt; PRT

45 &lt;213&gt; Phleum pratense

&lt;400&gt; 87

Leu	Lys	Lys	Ala	Ile	Thr	Ala	Met	Ser	Glu	Ala	Gln	Lys	
1				5						10			

50

&lt;210&gt; 88

&lt;211&gt; 13

&lt;212&gt; PRT

55 &lt;213&gt; Poa pratensis

&lt;400&gt; 88

EP 2 891 662 A1

Leu Lys Lys Ala Ile Thr Ala Met Ser Gln Ala Gln Lys  
1 5 10

5 <210> 89  
<211> 15  
<212> PRT  
<213> Phalaris aquatica

10 <400> 89  
Lys Glu Ser Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys  
1 5 10 15

15 <210> 90  
<211> 17  
<212> PRT  
<213> Phleum pratense

20 <400> 90  
Pro Gln Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala Ile  
1 5 10 15

25 Thr

30 <210> 91  
<211> 18  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Rye09B1 peptide

35 <400> 91

Lys Pro Glu Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala  
1 5 10 15

40 Ile Thr

45 <210> 92  
<211> 19  
<212> PRT  
<213> Artificial Sequence

50 <220>  
<223> Rye09B2 peptide

<400> 92

Lys Lys Pro Glu Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys  
1 5 10 15

55 Ala Ile Thr

<210> 93  
 <211> 16  
 <212> PRT  
 <213> Artificial Sequence

5

<220>  
 <223> Tim07B1 peptide

<400> 93

10

Lys Lys Ile Pro Ala Gly Glu Leu Gln Ile Ile Asp Lys Ile Asp Ala  
 1 5 10 15

15

<210> 94  
 <211> 17  
 <212> PRT  
 <213> Artificial Sequence

20

<220>  
 <223> Tim07B2 peptide

<400> 94

25

Lys Lys Ile Pro Ala Gly Glu Leu Gln Ile Ile Asp Lys Ile Asp Ala  
 1 5 10 15

Lys

30

<210> 95  
 <211> 20  
 <212> PRT  
 <213> Cynodon dactylon

35

<400> 95

Ser Gly Lys Ala Phe Gly Ala Met Ala Lys Lys Gly Gln Glu Asp Lys  
 1 5 10 15

40

Leu Arg Lys Ala  
 20

45

<210> 96  
 <211> 29  
 <212> PRT  
 <213> Cynodon dactylon

50

<400> 96

Pro Lys Asp Ser Asp Glu Phe Ile Pro Met Lys Ser Ser Trp Gly Ala  
 1 5 10 15

55

Ile Trp Arg Ile Asp Pro Lys Lys Pro Leu Lys Gly Pro  
 20 25

<210> 97  
 <211> 29  
 <212> PRT  
 <213> *Cynodon dactylon*

5

<400> 97

Arg Leu Thr Ser Glu Gly Gly Ala His Leu Val Gln Asp Asp Val Ile  
 1 5 10 15

10

Pro Ala Asn Trp Lys Pro Asp Thr Val Tyr Thr Ser Lys  
 20 25

15

<210> 98  
 <211> 20  
 <212> PRT  
 <213> *Lolium perenne*

20

Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly Pro  
 1 5 10 15

25

Lys Asp Asn Gly  
 20

30

<210> 99  
 <211> 20  
 <212> PRT  
 <213> *Lolium perenne*

<400> 99

35

Gly His Ala Phe Gly Ser Met Ala Lys Lys Gly Glu Glu Gln Asn Val  
 1 5 10 15

40

Arg Ser Ala Gly  
 20

45

<210> 100  
 <211> 20  
 <212> PRT  
 <213> *Lolium perenne*

<400> 100

50

Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys Tyr Val Asp Gly  
 1 5 10 15

55

Asp Gly Asp Val  
 20

<210> 101

<211> 20  
 <212> PRT  
 <213> Lolium perenne

5 <400> 101

Lys Glu Ser Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu  
 1 5 10 15

10 Thr Gly Pro Phe  
20

<210> 102

15 <211> 17  
 <212> PRT  
 <213> Lolium perenne

<400> 102

Ala Tyr Val Ala Thr Leu Thr Glu Ala Leu Arg Val Ile Ala Gly Thr  
 1 5 10 15

Leu

25

<210> 103

<211> 20  
 <212> PRT  
 30 <213> Lolium perenne

<400> 103

Lys Phe Ile Pro Thr Leu Val Ala Ala Val Lys Gln Ala Tyr Ala Ala  
 1 5 10 15

35

Lys Gln Ala Thr  
 20

40

<210> 104  
 <211> 17  
 <212> PRT  
 <213> Lolium perenne

45 <400> 104

Thr Ala Leu Lys Lys Ala Val Thr Ala Met Ser Glu Ala Glu Lys Glu  
 1 5 10 15

50 Ala

<210> 105

55 <211> 17  
 <212> PRT  
 <213> Lolium perenne

&lt;400&gt; 105

Asn	Asp	Lys	Phe	Thr	Val	Phe	Glu	Ser	Ala	Phe	Asn	Lys	Ala	Leu	Asn
1					5				10					15	

5

Glu

10

&lt;210&gt; 106

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Lolium perenne

15

&lt;400&gt; 106

Asp	Val	Asn	Ala	Gly	Phe	Lys	Ala	Ala	Val	Ala	Ala	Ala	Ala	Asn	Ala
1					5				10					15	

20

Pro Pro Ala Asp

20

25

&lt;210&gt; 107

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Lolium perenne

&lt;400&gt; 107

30

Gly	Ala	Thr	Pro	Glu	Ala	Lys	Tyr	Asp	Ala	Phe	Val	Thr	Ala	Leu	Thr
1					5				10					15	

35

Glu Ala Leu Arg

20

40

&lt;210&gt; 108

&lt;211&gt; 29

&lt;212&gt; PRT

&lt;213&gt; Lolium perenne

&lt;400&gt; 108

45

Gly	Glu	Leu	Gln	Ile	Val	Asp	Lys	Ile	Asp	Ala	Ala	Phe	Lys	Ile	Ala
1					5				10				15		

50

Ala Thr Ala Ala Asn Ala Ala Pro Thr Asn Asp Lys Phe

20

25

55

&lt;210&gt; 109

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Lolium perenne

&lt;400&gt; 109

Gly Ala Tyr Glu Thr Tyr Lys Phe Ile Pro Ser Leu Glu Ala Ala Val  
 1 5 10 15

5 Lys Gln Ala Tyr  
 20

10 <210> 110  
 <211> 29  
 <212> PRT  
 <213> *Lolium perenne*

<400> 110

15 Pro Glu Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala Ile  
 1 5 10 15

20 Thr Ala Met Thr Gln Ala Gln Lys Ala Gly Lys Pro Ala  
 20 25

25 <210> 111  
 <211> 23  
 <212> PRT  
 <213> *Phleum pratense*

<400> 111

30 Pro Glu Val Lys Tyr Thr Val Phe Glu Thr Ala Leu Lys Lys Ala Ile  
 1 5 10 15

35 Thr Ala Met Ser Glu Ala Gln  
 20

40 <210> 112  
 <211> 17  
 <212> PRT  
 <213> *Lolium perenne*

<400> 112

45 Gln Lys Leu Ile Glu Lys Ile Asn Ala Gly Phe Lys Ala Ala Val Ala  
 1 5 10 15

50 Ala  
 <210> 113  
 <211> 20  
 <212> PRT  
 <213> *Phleum pratense*

<400> 113

55 Gly His Ala Phe Gly Ala Met Ala Lys Lys Gly Asp Glu Gln Lys Leu  
 1 5 10 15

Arg Ser Ala Gly  
20

5           <210> 114  
       <211> 20  
       <212> PRT  
       <213> *Phleum pratense*

10           <400> 114

Gly Ser Asn Pro Asn Tyr Leu Ala Leu Leu Val Lys Tyr Val Asn Gly  
1                       5                                   10                           15

15           Asp Gly Asp Val  
20                       20

20           <210> 115  
       <211> 28  
       <212> PRT  
       <213> *Phleum pratense*

<400> 115

Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser Trp Gly Ala Ile Trp  
1                       5                                   10                           15

30           Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro Phe  
35                       20                               25

35           <210> 116  
       <211> 27  
       <212> PRT  
       <213> *Cynodon dactylon*

<400> 116

40           Pro Lys Asp Ser Asp Glu Phe Ile Pro Met Lys Ser Ser Trp Gly Ala  
1                       5                                   10                           15

Ile Trp Arg Ile Asp Pro Lys Lys Pro Leu Lys  
45                       20                               25

45           <210> 117  
       <211> 25  
       <212> PRT  
       <213> *Cynodon dactylon*

50           <400> 117

Glu Gly Gly Ala His Leu Val Gln Asp Asp Val Ile Pro Ala Asn Trp  
1                       5                                   10                           15

55           Lys Pro Asp Thr Val Tyr Thr Ser Lys  
60                       20                               25

<210> 118  
 <211> 25  
 <212> PRT  
 <213> *Phleum pratense*

5

<400> 118

Glu Gly Gly Thr Lys Thr Glu Ala Glu Asp Val Ile Pro Glu Gly Trp  
 1 5 10 15

10

Lys Ala Asp Thr Ser Tyr Glu Ser Lys  
 20 25

15

<210> 119  
 <211> 19  
 <212> PRT  
 <213> *Phleum pratense*

20

<400> 119

Lys Ile Asn Ala Gly Phe Lys Ala Ala Leu Ala Gly Ala Gly Val Gln  
 1 5 10 15

25

**Pro Ala Asp**

30

<210> 120  
 <211> 20  
 <212> PRT  
 <213> *Phleum pratense*

<400> 120

35

Lys Ile Asn Asp Gly Phe Lys Ala Ala Leu Ala Ala Ala Ala Gly Val  
 1 5 10 15

40

**Pro Pro Ala Asp**

20

45

<210> 121  
 <211> 20  
 <212> PRT  
 <213> *Phleum pratense*

<400> 121

50

Asp Ile Asn Val Gly Phe Lys Ala Ala Val Ala Ala Ala Ala Ser Val  
 1 5 10 15

55

**Pro Ala Ala Asp**

20

<210> 122

<211> 20  
 <212> PRT  
 <213> *Phleum pratense*

5 <400> 122

Gly Ala Thr Pro Glu Ala Lys Tyr Asp Ala Tyr Val Ala Thr Leu Ser  
 1 5 10 15

10 Glu Ala Leu Arg  
 20

<210> 123  
 <211> 20  
 <212> PRT  
 <213> *Phleum pratense*

<400> 123

20 Gly Ala Thr Pro Glu Ala Lys Phe Asp Ser Phe Val Ala Ser Leu Thr  
 1 5 10 15

Glu Ala Leu Arg  
 20

25

<210> 124  
 <211> 20  
 <212> PRT  
 <213> *Lolium perenne*

<400> 124

Gly Glu Leu Gln Ile Val Asp Lys Ile Asp Ala Ala Phe Lys Ile Ala  
 1 5 10 15

35

Ala Thr Ala Ala  
 20

40

<210> 125  
 <211> 19  
 <212> PRT  
 <213> *Phleum pratense*

45

<400> 125

Gly Glu Leu Gln Val Ile Glu Lys Asp Ala Ala Phe Lys Val Ala Ala  
 1 5 10 15

50

Thr Ala Ala

55

<210> 126  
 <211> 20  
 <212> PRT  
 <213> *Phleum pratense*

&lt;400&gt; 126

Gly	Glu	Leu	Gln	Val	Ile	Glu	Lys	Val	Asp	Ala	Ala	Phe	Lys	Val	Ala
1					5				10						15

5

Ala	Thr	Ala	Ala
		20	

10

&lt;210&gt; 127

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

15

&lt;400&gt; 127

Gly	Glu	Leu	Gln	Phe	Ile	Glu	Lys	Val	Asp	Ser	Ala	Leu	Lys	Val	Ala
1					5				10						15

20

Ala	Thr	Ala	Ala
		20	

25

&lt;210&gt; 128

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

&lt;400&gt; 128

30

Gly	Glu	Leu	Gln	Ile	Ile	Asp	Lys	Ile	Asp	Ala	Ala	Phe	Lys	Val	Ala
1					5				10						15

35

Ala	Thr	Ala	Ala
		20	

&lt;210&gt; 129

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Lolium perenne

&lt;400&gt; 129

45

Asp	Ala	Ala	Phe	Lys	Ile	Ala	Ala	Thr	Ala	Ala	Asn	Ala	Ala	Pro	Thr
1					5				10						15

Asn	Asp	Lys	Phe
		20	

50

&lt;210&gt; 130

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Phleum pratense

&lt;400&gt; 130

Asp Ala Ala Phe Lys Val Ala Ala Thr Ala Ala Asn Ala Ala Pro Ala  
 1 5 10 15

5 Asn Asp Lys Phe  
 20

10 <210> 131  
 <211> 20  
 <212> PRT  
 <213> Phleum pratense  
 <400> 131

15 Asp Ser Ala Leu Lys Val Ala Ala Thr Ala Ala Asn Ala Ala Ala Ala  
 1 5 10 15

20 Asn Asp Lys Phe  
 20

25 <210> 132  
 <211> 20  
 <212> PRT  
 <213> Phleum pratense  
 <400> 132

30 Asp Ala Ala Phe Lys Val Ala Ala Thr Ala Ala Ala Thr Ala Pro Ala  
 1 5 10 15

Asp Asp Lys Phe  
 20

35 <210> 133  
 <211> 20  
 <212> PRT  
 <213> Phleum pratense  
 <400> 133

40 Gly Ala Tyr Glu Ser Tyr Lys Phe Ile Pro Ala Leu Glu Ala Ala Val  
 1 5 10 15

45 Lys Gln Ala Tyr  
 20

50 <210> 134  
 <211> 20  
 <212> PRT  
 <213> Phleum pratense  
 <400> 134

55 Gly Ala Tyr Asp Thr Tyr Lys Cys Ile Pro Ser Leu Glu Ala Ala Val  
 1 5 10 15

Lys Gln Ala Tyr  
20

5           <210> 135  
       <211> 20  
       <212> PRT  
       <213> *Lolium perenne*

10           <400> 135  
          Pro Glu Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala Ile  
           1                   5                                   10                           15

15           Thr Ala Met Thr  
20           20

20           <210> 136  
       <211> 20  
       <212> PRT  
       <213> *Phleum pratense*

25           <400> 136  
          Pro Glu Val Lys Tyr Thr Val Phe Glu Thr Ala Leu Lys Lys Ala Ile  
           1                   5                                   10                           15

30           Thr Ala Met Ser  
35           20

35           <210> 137  
       <211> 20  
       <212> PRT  
       <213> *Phleum pratense*

40           <400> 137  
          Ala Glu Val Lys Tyr Ala Val Phe Glu Ala Ala Leu Thr Lys Ala Ile  
           1                   5                                   10                           15

45           Thr Ala Met Ser  
50           20

45           <210> 138  
       <211> 20  
       <212> PRT  
       <213> *Lolium perenne*

50           <400> 138  
          Ala Ala Leu Thr Lys Ala Ile Thr Ala Met Thr Gln Ala Gln Lys Ala  
           1                   5                                   10                           15

55           Gly Lys Pro Ala  
60           20

<210> 139  
 <211> 20  
 <212> PRT  
 <213> **Phleum pratense**

5

&lt;400&gt; 139

Thr Ala Leu Lys Lys Ala Ile Thr Ala Met Ser Glu Ala Gln Lys Ala  
 1 5 10 15

10

Ala Lys Pro Ala  
 20

15

<210> 140  
 <211> 20  
 <212> PRT  
 <213> **Phleum pratense**

20

&lt;400&gt; 140

Ala Ala Leu Thr Lys Ala Ile Thr Ala Met Ser Glu Val Gln Lys Val  
 1 5 10 15

25

Ser Gln Pro Ala  
 20

30

<210> 141  
 <211> 15  
 <212> PRT  
 <213> **Mycobacterium tuberculosis**

&lt;400&gt; 141

35

Ala Lys Thr Ile Ala Tyr Asp Glu Glu Ala Arg Arg Gly Leu Glu  
 1 5 10 15

40

<210> 142  
 <211> 13  
 <212> PRT  
 <213> **Influenza virus**

&lt;400&gt; 142

45

Pro Lys Tyr Val Lys Gln Asn Thr Leu Lys Leu Ala Thr  
 1 5 10

50

<210> 143  
 <211> 15  
 <212> PRT  
 <213> **Homo sapiens**

&lt;400&gt; 143

55

Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg  
 1 5 10 15

<210> 144  
<211> 9  
<212> PRT  
<213> *Cynodon dactylon*  
<200> 144

Phe Gly Ala Met Ala Lys Lys Gly Gln  
1 5

## Claims

1. A composition suitable for use in preventing or treating allergy to grass pollen, comprising:

- (a) at least one of the polypeptides Tim07B (KIPAGELQIIDKID A), Tim 10B (KYTVFETALKKAITAMSE), Tim 04A (WGAIRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof;
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), Ber02C (KSS-WGAIWRIDPKKPLK), Ber 02B (KDSDEFIPMKSSWGAIR) or a variant of any thereof; and
- (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTKAIT), Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIPSLEAAVKQAY), Rye 05C (NAGFKAVALAAANAPPK) or a variant of any thereof.

wherein said variant is:

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a), (b) or (c), or
- II) a polypeptide of 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a), (b) or (c), and that is capable of tolerising to said corresponding polypeptide; or
- III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a), (b) or (c), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

2. A composition according to claim 1, comprising at least four different polypeptides selected from:

- (a) Tim07B (KIPAGELQIIDKIDA) or a variant thereof;
- (b) Ber01 (SGKAFGAMAKKGQED) or a variant thereof;
- (c) Bio04A (LKKAVTAMSEAEK) or a variant thereof;
- (d) Rye09B (PEVKYAVFEAALTAKIT) or a variant thereof;
- (e) Ber02 (FIPMKSSWGA) or a variant thereof;
- (f) Ber02C (KSSWGAIWRIDPKKPLK) or a variant thereof;
- (g) Bio03A (KFIPTLVAAVKQAYAAKQ) or a variant thereof; and
- (h) Bio02A (KYDAYVATI TEAL R) or a variant thereof;

wherein said variant is:

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a) to (h), or
- II) a polypeptide 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a) to (h), and that is capable of tolerising to said corresponding polypeptide; or
- III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a) to (h), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide.

3. A composition according to claim 2, comprising

- (a) the polypeptide Tim07B (KIPAGELQIIDKIDA) or a variant thereof;
- (b) at least one of the polypeptides of Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), Ber02C (KSS-WGAIWRIDPKKPLK) or a variant of any thereof; and
- (c) at least one of the polypeptides of Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTCAIT), Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ) or a variant of any thereof; and
- (d) at least one additional polypeptide of b) or c) not selected above.

10 4. A composition according to claim 3, comprising the polypeptide Tim07B (KIPAGELQIIDKIDA) or a variant thereof, the polypeptide Ber01 (SGKAFGAMAKKGQED) or a variant thereof, the polypeptide Bio04A (LKKAVTAMSEAEK) or a variant thereof, and at least one additional polypeptide of b) or c) not selected above.

15 5. A composition according to any one of the preceding claims, wherein said variants are 9 to 20 or 13 to 17 amino acids in length.

20 6. A composition according to any one of the preceding claims, comprising a said variant which has 1, 2, 3 or 4 amino acids deleted from the N-terminal and/or C-terminal end of the sequence of the corresponding said polypeptide and/or has 1, 2, 3, 4 or 5 conservative amino acid substitutions in the sequence of the corresponding said polypeptide.

25 7. A composition according to any one of the preceding claims, which comprises the polypeptides:

- (a) Tim07B1 (KKIPAGELQIIDKIDA);
- (b) Ber01 (SGKAFGAMAKKGQED);
- (c) Bio04A (LKKAVTAMSEAEK);
- (d) Rye09B1 (KPEVKYAVFEAALTCAIT);
- (e) Ber02 (FIPMKSSWGA);
- (f) Ber02C (KSSWGAIWRIDPKKPLK); and
- (g) Bio02A (KYDAYVATLTEALR)

30 8. A composition according to any one of the preceding claims, which is

- a solution wherein each polypeptide has a concentration in the range of 0.03 to 200 nmol/ml, 0.3 to 200 nmol/ml, 5 to 200 nmol/ml or 30 to 120 nmol/ml, or
- a dry powder or dry granules.

35 9. A composition according to any one of the preceding claims, which is a pharmaceutical formulation comprising a pharmaceutically acceptable carrier or diluent.

40 10. A composition according to claim 9, for use in a method of preventing or treating allergy to grass pollen.

11. A composition according to claim 10, wherein said method is for preventing or treating allergy to grass pollen in a human, optionally wherein said human is grass allergic and optionally wherein said grass allergic human is:

- sensitised to one or more grass pollen allergens;
- has a family history of allergy to grass; or
- displays symptoms of allergy when exposed to grass.

45 12. A kit comprising:

- (a) the polypeptide Tim07B (KIPAGELQIIDKIDA), Tim 10B (KYTVFETALKAITAMSE), Tim 04A (WGAI-WRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof as defined in claim 1 (I) or (II);
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED) Ber02 (FIPMKSSWGA), Ber02C (KSS-WGAIWRIDPKKPLK), Ber02B (KDSDEFIPMKSSWGAIWR) or a variant of any thereof as defined in claim 1 (I) or (II); and
- (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTCAIT), Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFPISLEAAVKQAY), Rye 05C (NAGFKAVALAAAANAPPK) or a variant of any thereof as defined in claim 1 (I) or (II),

wherein each different polypeptide is for simultaneous, separate or sequential use in preventing or treating allergy to grass pollen.

13. A process for preparing a pharmaceutical formulation suitable for use in preventing or treating allergy to grass pollen,  
5 which process comprises formulating:

- (a) at least one of the polypeptides Tim07B (KIPAGELQIIDKID A), Tim 10B (KYTVFETALKKAITAMSE), Tim 04A (WGAIWRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof;
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), Ber02C (KSS-  
10 WGAIWRIDPKKPLK), Ber 02B (KDSDEFIPMKSSWGAIWR) or a variant of any thereof; and
- (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTAKIT), Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIPSLEAAVKQAY), Rye 05C (NAGFKAVALAAANAPPK) or a variant of any thereof,

15 wherein said variant is:

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a), (b) or (c), or
- II) a polypeptide of 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a), (b) or (c), and that is capable of tolerising to said corresponding polypeptide; or
- III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a), (b) or (c), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide;

together with a pharmaceutically acceptable carrier or diluent.

14. Use of:

30

- (a) at least one of the polypeptides Tim07B (KIPAGELQIIDKID A), Tim 10B (KYTVFETALKKAITAMSE), Tim 04A (WGAIWRIDTPDKL), Tim 07G (FKVAATAANAAPANDK) or a variant of any thereof;
- (b) at least one of the polypeptides Ber01 (SGKAFGAMAKKGQED), Ber02 (FIPMKSSWGA), Ber02C (KSS-  
35 WGAIWRIDPKKPLK), Ber 02B (KDSDEFIPMKSSWGAIWR) or a variant of any thereof; and
- (c) at least one of the polypeptides Bio04A (LKKAVTAMSEAEK), Rye09B (PEVKYAVFEAALTAKIT), Bio02A (KYDAYVATLTEALR), Bio03A (KFIPTLVAAVKQAYAAKQ), Rye 08A (ETYKFIPSLEAAVKQAY), Rye 05C (NAGFKAVALAAANAPPK) or a variant of any thereof,

40 wherein said variant is:

45

- I) a longer polypeptide of up to 30 amino acids in length which comprises the sequence of the corresponding polypeptide specified in (a), (b) or (c), or
- II) a polypeptide of 9 to 30 amino acids in length which comprises a sequence that has at least 65% homology to the sequence of the corresponding polypeptide specified in (a), (b) or (c), and that is capable of tolerising to said corresponding polypeptide; or
- III) a polypeptide of length 9 to 30 amino acids which comprises a sequence of at least 9 contiguous amino acids of the sequence of the corresponding polypeptide specified in (a), (b) or (c), or a sequence that has at least 65% homology to said at least 9 contiguous amino acids, which sequence of at least 9 contiguous amino acids or homologous sequence is capable of tolerising to said corresponding polypeptide,

50 for the manufacture of a medicament for use in preventing or treating allergy to grass pollen.

55 15. A polypeptide selected from Bio04A (LKKAVTAMSEAEK), Tim07B1 (KKIPAGELQIIDKIDA) and Ber01 (SGKAF-  
GAMAKKGQED).



## EUROPEAN SEARCH REPORT

Application Number

EP 15 15 2164

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	Y WO 03/024998 A1 (UNIV MONASH [AU]; O'HEHIR ROBYN [AU]; ROLLAND JENNIFER [AU]) 27 March 2003 (2003-03-27) * page 3, last paragraph; claim 9; figure 2; sequences 14,22,23 * * page 39, last paragraph * -----	1-3,5, 8-10, 12-14	INV. C07K14/415 C07K7/06 C07K7/08 A61K38/04 A61K38/16 A61K39/36
15	Y WO 2007/140505 A2 (BIOMAY AG [AT]; VALENTE RUDOLF [AT]; FOCKE-TEJKL MARGARETE [AT]; LINHA) 13 December 2007 (2007-12-13) * examples 14-18; table 3; sequences 11,13,14 *	1-3,5, 8-10, 12-14	
20	X, P WO 2009/022154 A2 (CIRCASSIA LTD [GB]; HAFNER RODERICK PETER [GB]; LARCHE MARK [CA]) 19 February 2009 (2009-02-19) * example 2 * * page 27, line 29 - page 29 * * page 31 - page 33 * * page 48 * * page 54, paragraph 2 - last paragraph * * page 73, * * page 76 * -----	15	
25	Y, P WO 2009/022157 A2 (CIRCASSIA LTD [GB]; HAFNER RODERICK PETER [GB]; LAIDLER PAUL [GB]) 19 February 2009 (2009-02-19) * sequences 16,19,25,29,31,33,35 *	1-3,5, 8-10, 12-14	TECHNICAL FIELDS SEARCHED (IPC)
30	X, P WO 2009/022157 A2 (CIRCASSIA LTD [GB]; HAFNER RODERICK PETER [GB]; LAIDLER PAUL [GB]) 19 February 2009 (2009-02-19) * sequences 16,19,25,29,31,33,35 *	15	C07K A61K
35	Y, P WO 96/03106 A2 (UNIV MANITOBA [CA]; MOHAPATRA SHYAM S [CA]) 8 February 1996 (1996-02-08) * page 29; claim 7; sequences 19,24 * ----- -/-	1-3,5, 8-10, 12-14	
40	Y WO 96/03106 A2 (UNIV MANITOBA [CA]; MOHAPATRA SHYAM S [CA]) 8 February 1996 (1996-02-08) * page 29; claim 7; sequences 19,24 * ----- -/-	1-3,5, 8-10, 12-14	
45	The present search report has been drawn up for all claims		
1	Place of search Munich	Date of completion of the search 24 April 2015	Examiner Roscoe, Richard
50	CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document
55	EPO FORM 1503/03.82 (P04C01)		



## EUROPEAN SEARCH REPORT

Application Number

EP 15 15 2164

5

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
10	Y WO 2006/008018 A1 (MERCK PATENT GMBH [DE]; FIEBIG HELMUT [DE]; WALD MARTIN [DE]; NANDY AN) 26 January 2006 (2006-01-26) * page 8, paragraph 2; claim 2; figure 1 * -----	1-3,5, 8-10, 12-14	
15	Y JUTEL ET AL: "Allergen-specific immunotherapy with recombinant grass pollen allergens", JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, MOSBY, INC, US LNKD-DOI:10.1016/J.JACI.2005.06.004, vol. 116, no. 3, 1 September 2005 (2005-09-01), pages 608-613, XP005068191, ISSN: 0091-6749 * abstract *	1-3,5, 8-10, 12-14	
20			
25	A SWOBODA INES ET AL: "Mutants of the major ryegrass pollen allergen, Lol p 5, with reduced IgE-binding capacity: candidates for grass pollen-specific immunotherapy.", EUROPEAN JOURNAL OF IMMUNOLOGY JAN 2002 LNKD- PUBMED:11782018, vol. 32, no. 1, January 2002 (2002-01), pages 270-280, XP002581444, ISSN: 0014-2980 * abstract *	1-15	TECHNICAL FIELDS SEARCHED (IPC)
30			
35	X WO 95/06728 A2 (IMMULOGIC PHARMA CORP [US]; GRIFFITH IRWIN J [US]; KUO MEI CHANG [US];) 9 March 1995 (1995-03-09) * Whole doc., in particular Fig.2, 3; Claims *	1-15	
40		----- -/-	
45			
1	The present search report has been drawn up for all claims		
50	Place of search Munich	Date of completion of the search 24 April 2015	Examiner Roscoe, Richard
55	CATEGORIE OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		
	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document		



## EUROPEAN SEARCH REPORT

Application Number

EP 15 15 2164

5

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
10	X WO 03/082924 A1 (UNIV MONASH [AU]; O'HEHIR ROBYN [AU]; ROLLAND JENNIFER [AU]) 9 October 2003 (2003-10-09) * Whole doc., in particular Seq. on p.10, 11, claims. *	1-15	
15	X US 7 112 333 B1 (GRIFFITH IRWIN J [CA] ET AL) 26 September 2006 (2006-09-26) * Similar to D9. Whole doc., in particular peptide allergens. *	1-15	
20		-----	
25		-----	
30		-----	TECHNICAL FIELDS SEARCHED (IPC)
35		-----	
40		-----	
45		-----	
1	The present search report has been drawn up for all claims		
50	Place of search Munich	Date of completion of the search 24 April 2015	Examiner Roscoe, Richard
55	EPO FORM 1503/03.82 (P04C01) CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		
	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document		

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 15 15 2164

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24-04-2015

10

	Patent document cited in search report		Publication date	Patent family member(s)	Publication date
15	WO 03024998	A1	27-03-2003	AT 502046 T	15-04-2011
				DK 1436320 T3	27-06-2011
				EP 1436320 A1	14-07-2004
				ES 2362040 T3	27-06-2011
				US 2005075481 A1	07-04-2005
				US 2011217325 A1	08-09-2011
				WO 03024998 A1	27-03-2003
20	WO 2007140505	A2	13-12-2007	AT 503690 A1	15-12-2007
				AU 2007257308 A1	13-12-2007
				BR PI0712421 A2	19-06-2012
				CA 2657971 A1	13-12-2007
				CN 101466738 A	24-06-2009
				CN 103319601 A	25-09-2013
				CN 103319602 A	25-09-2013
				EP 2035457 A2	18-03-2009
				JP 5399895 B2	29-01-2014
				JP 2009539354 A	19-11-2009
				JP 2014064573 A	17-04-2014
				JP 2014087341 A	15-05-2014
25				RU 2008152045 A	20-07-2010
				US 2009324501 A1	31-12-2009
				WO 2007140505 A2	13-12-2007
				-----	-----
				-----	-----
30	WO 2009022154	A2	19-02-2009	AT 482721 T	15-10-2010
				AU 2008288283 A1	19-02-2009
				BR PI0815395 A2	10-02-2015
				CA 2696563 A1	19-02-2009
				CN 101820907 A	01-09-2010
				CN 103272227 A	04-09-2013
				DK 2083856 T3	24-01-2011
				DK 2190473 T3	25-03-2013
				EA 201070270 A1	30-08-2010
				EA 201070271 A1	30-12-2010
				EP 2083856 A2	05-08-2009
				EP 2190473 A2	02-06-2010
35				EP 2286833 A2	23-02-2011
				EP 2397154 A2	21-12-2011
				ES 2402956 T3	10-05-2013
				HK 1132194 A1	15-07-2011
				HR P20100662 T1	31-01-2011
				HR P20130270 T1	30-04-2013
				IL 203943 A	30-05-2013
40				JP 5697447 B2	08-04-2015
				JP 2010536333 A	02-12-2010
				JP 2015057393 A	26-03-2015
				-----	-----
45				-----	-----
				-----	-----
				-----	-----
50				-----	-----
				-----	-----
				-----	-----
				-----	-----
				-----	-----
55				-----	-----
				-----	-----

EPO FORM P0459  
For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 15 15 2164

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24-04-2015

10

	Patent document cited in search report	Publication date	Patent family member(s)	Publication date
			KR 20100068252 A	22-06-2010
			NZ 583361 A	26-10-2012
			NZ 583362 A	29-06-2012
15			PT 2083856 E	23-12-2010
			SI 2083856 T1	28-02-2011
			SI 2190473 T1	31-05-2013
			US 2010260805 A1	14-10-2010
20			US 2010298239 A1	25-11-2010
			US 2011123558 A1	26-05-2011
			US 2013336999 A1	19-12-2013
			US 2014186403 A1	03-07-2014
25			WO 2009022154 A2	19-02-2009
			WO 2009022155 A2	19-02-2009
			WO 2009022156 A2	19-02-2009
			WO 2009022157 A2	19-02-2009
	<hr/>			
	WO 2009022157 A2 19-02-2009	AT 482721 T		15-10-2010
		AU 2008288283 A1		19-02-2009
30		BR PI0815395 A2		10-02-2015
		CA 2696563 A1		19-02-2009
		CN 101820907 A		01-09-2010
		CN 103272227 A		04-09-2013
		DK 2083856 T3		24-01-2011
		DK 2190473 T3		25-03-2013
35		EA 201070270 A1		30-08-2010
		EA 201070271 A1		30-12-2010
		EP 2083856 A2		05-08-2009
		EP 2190473 A2		02-06-2010
		EP 2286833 A2		23-02-2011
		EP 2397154 A2		21-12-2011
40		ES 2402956 T3		10-05-2013
		HK 1132194 A1		15-07-2011
		HR P20100662 T1		31-01-2011
		HR P20130270 T1		30-04-2013
		IL 203943 A		30-05-2013
45		JP 5697447 B2		08-04-2015
		JP 2010536333 A		02-12-2010
		JP 2015057393 A		26-03-2015
		KR 20100068252 A		22-06-2010
		NZ 583361 A		26-10-2012
		NZ 583362 A		29-06-2012
50		PT 2083856 E		23-12-2010
		SI 2083856 T1		28-02-2011
		SI 2190473 T1		31-05-2013
		US 2010260805 A1		14-10-2010
		US 2010298239 A1		25-11-2010

55

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 15 15 2164

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24-04-2015

10

	Patent document cited in search report		Publication date	Patent family member(s)	Publication date
15				US 2011123558 A1	26-05-2011
				US 2013336999 A1	19-12-2013
				US 2014186403 A1	03-07-2014
				WO 2009022154 A2	19-02-2009
				WO 2009022155 A2	19-02-2009
				WO 2009022156 A2	19-02-2009
				WO 2009022157 A2	19-02-2009
20	WO 9603106	A2	08-02-1996	AU 3338195 A	22-02-1996
				WO 9603106 A2	08-02-1996
25				WO 2006008018 A1 26-01-2006	AT 432944 T 15-06-2009
				AU 2005263477 A1	26-01-2006
				BR P10513504 A	06-05-2008
				CA 2574449 A1	26-01-2006
				CN 1989151 A	27-06-2007
				CN 103467583 A	25-12-2013
				DE 102004035337 A1	16-03-2006
				EP 1768996 A1	04-04-2007
				EP 2028189 A2	25-02-2009
				ES 2328050 T3	06-11-2009
30				JP 5149002 B2	20-02-2013
				JP 2008507262 A	13-03-2008
				PT 1768996 E	07-09-2009
				RU 2007106170 A	27-08-2008
				US 2008267985 A1	30-10-2008
				WO 2006008018 A1	26-01-2006
				WO 9506728 A2 09-03-1995	AU 7559794 A 22-03-1995
				CA 2168565 A1	09-03-1995
				EP 0714440 A1	05-06-1996
				FI 960629 A	12-04-1996
40				JP H09504167 A	28-04-1997
				NO 960553 A	12-04-1996
				NZ 271818 A	24-11-1997
				WO 9506728 A2	09-03-1995
				ZA 9406068 A	28-03-1995
				WO 03082924 A1 09-10-2003	EP 1499337 A1 26-01-2005
				EP 2092937 A2	26-08-2009
50				US 2006084789 A1	20-04-2006
				WO 03082924 A1	09-10-2003
				US 7112333 B1 26-09-2006	NONE

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- WO 0901928 A [0001]
- WO 09019274FILED A [0001]
- WO 0912578 A [0001]
- GB 0901995 W [0001]
- WO 0917871 A [0001]
- US 5399346 A [0112]
- US 5580859 A [0112]
- US 5589466 A [0112]

## Non-patent literature cited in the description

- MEZIERE et al. *J. Immunol.*, 1997, vol. 159, 3230-3237 [0073]
- ALLEN et al. *Nature*, 1987, vol. 327, 713-5 [0081]
- EVAVOLD ; ALLEN. *Nature*, 1991, vol. 252, 1308-10 [0082]
- SMILEK et al. *Proc Natl Acad Sci USA*, 1991, vol. 88, 9633-9637 [0083]
- METZLER ; WRAITH. *Int Immunol*, 1993, 1159-65 [0084]
- SLOAN-LANCASTER et al. *Nature*, 1993, vol. 363, 156-9 [0084]
- Allergy and Allergic Diseases. Blackwell Science, 1997, 1113-1130 [0085]
- Provocation tests and measurements of mediators from mast cells and basophils in asthma and allergic rhinitis. CROMWELL O ; DURHAM SR ; SHAW RJ ; MACKAY J ; KAY AB. *Handbook of Experimental Immunology*. Blackwell Scientific Publications, 1986 [0085]
- SAMBROOK et al. Molecular Cloning - a laboratory manual. Cold Spring Harbor Press, 1989 [0098]
- Remington's Pharmaceutical Sciences. Mack Pub. Co, 1991 [0118]
- JEFFERY et al. *Pharm. Res.*, 1993, vol. 10, 362-368 [0120]
- AKDIS et al. *J. Clin. Invest.*, 1996, vol. 98, 1676-1683 [0137]
- BAUER et al. *Clin. Exp. Immunol.*, 1997, vol. 107, 536-541 [0137]
- CAO et al. *Immunology*, 1997, vol. 90, 46-51 [0137]
- DE GROOT et al. *AIDS Research and Human Retroviruses*, 1997, vol. 13, 539-41 [0147]
- SCHAFER et al. *Vaccine* 1998, 1998, vol. 16 [0147]
- JESDALE et al. *Vaccines '97*. Cold Spring Harbor Press, 1997 [0147]

## 摘要

本發明涉及包括用於預防或治療草花粉過敏症的肽的組合物，並且特別涉及用於預防和治療所述過敏症的肽的最佳組合。