



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/00	A2	(11) International Publication Number: WO 00/23091 (43) International Publication Date: 27 April 2000 (27.04.00)
(21) International Application Number: PCT/DK99/00567 (22) International Filing Date: 15 October 1999 (15.10.99) (30) Priority Data: PA 1998 01321 15 October 1998 (15.10.98) DK PA 1998 01322 15 October 1998 (15.10.98) DK PA 1998 01323 15 October 1998 (15.10.98) DK (71) Applicant (for all designated States except US): BIOIMAGE A/S [DK/DK]; Mørkhøj Bygade 28, DK-2860 Søborg (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): ARKHAMMAR, Per, O., G. [SE/SE]; Helmfeltsgatan 13, S-254 40 Helsingborg (SE). TERRY, Bernard, Robert [GB/DK]; Frederiksberg Allé 15,1., DK-1820 Frederiksberg C (DK). SCUDDER, Kurt, Marshall [US/DK]; Lavendelhaven 70, DK-2830 Virum (DK). BJØRN, Sara, Petersen [DK/DK]; Klampenborgvej 102, DK-2800 Lyngby (DK). THASTRUP, Ole [DK/DK]; Birkevej 37, DK-3460 Birkerød (DK). (74) Agent: PLOUGMANN, VINGTOFT & PARTNERS A/S; Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: SPECIFIC THERAPEUTIC INTERVENTIONS OBTAINED BY INTERFERENCE WITH REDISTRIBUTION AND/OR TARGETTING (57) Abstract The application describes a novel mechanism of action, that is modulation of the specific effectiveness of I-kappa-kinases or cyclic nucleotide phosphodiesterases (PDEs) which have the ability to cleave cGMP or cAMP. The preferred mode of action is dislocation, disruption of targeting or interference with redistribution of specific isoforms or splice variants of PDE4, PDE5, or I-kappa-kinases from their anchoring sites within cells, thereby modulating their specific effectiveness, not their enzymatic capacity. The chemical entities may be useful in preventing or treating in an animal, preferably a human, in need thereof an adverse condition which may be reduced or abolished by modulating the specific effectiveness of PDE4, PDE5, or I-kappa-kinases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

SPECIFIC THERAPEUTIC INTERVENTIONS OBTAINED BY INTERFERENCE WITH REDISTRIBUTION AND/OR TARGETTING.

SUMMARY OF THE INVENTION

This application describes a novel mechanism of action of chemical entities in order to
5 prevent or treat adverse conditions which may be reduced or abolished by modulating
the effectiveness of I-kappaB kinase (IKK) or cyclic nucleotide phosphodiesterases
(PDE:s) by modulation of their targeting or localisation in the cell. The preferred mode of
action being sought is dislocation or interference with the targeting of specific isoforms of
IKK or PDE:s and interference with their anchoring sites within cells, thereby reducing
10 their specific effectiveness, not directly their enzymatic capacity.

In its broadest aspect, the present application relates to a novel method for preventing or
treating, in an animal in need thereof, an adverse condition which may be reduced or
abolished by modulating the activity of one or more IKKs or PDE:s having the ability to
15 cleave cAMP or cGMP. The method comprises modulation of the specific effectiveness
of IKKs or PDE:s by modulating their spatial distribution within cells of the animal.
The IKK is chosen from the group consisting of IKK α , IKK β , IKK γ and NIK. In one
embodiment IKK β is the preferred isoform. The PDE:s are chosen from the group
consisting of PDE1, PDE2, PDE3, PDE4, PDE 5, PDE6, PDE7, PDE8, PDE9 and
20 PDE10. More specifically, the method relates to PDE4 and isoforms thereof, such as
PDE4D, and splice variants of PDE4D, such as PDE4D1, PDE4D2, PDE4D3, PDE4D4
and PDE4D5. The animal with the adverse condition may be a mammal and preferably a
human.

In one embodiment of the invention modulation of the specific effectiveness of the PDE
25 is a dislocation of the PDE from a native location within the cell.

In another embodiment of the invention modulation of the specific effectiveness of the
PDE involves a disruption of its targeting to a native location within the cell.

In another embodiment of the invention modulation of the specific effectiveness of the
PDE involves interference with the redistribution of the PDE, the redistribution being
30 associated with an increase or a decrease of the specific effectiveness of the PDE.

The modulation of the specific effectiveness of the PDE may involve both an up-
regulation or a down-regulation of the effectiveness of the PDE to perform its function
within the cell.

The present invention provides compositions and methods for modifying the activation of NF-kappaB by mis-targeting and/or modulation of the redistribution of specific IKKs.

In one embodiment we specifically modulate the targeting of IKK β . We have developed
5 two molecular probes PS473 and PS474 that upon expression in a relevant cell system
will dislocate endogenous IKK β from its anchoring site. The mis-targeting has, as shown
in example 1, significant functional consequences that can be related to a diminished
ability of cytokines and other stimuli to activate NFkappaB. We thus show that IL-1
induced translocation of NFkappaB from cytoplasm to the nucleus is effectively inhibited,
10 and furthermore as a consequence thereof we found that NFkappaB-induced
transcriptional activation was inhibited.

NFkappaB has been shown to rescue transformed cells from undergoing apoptosis
when exposed to pro-inflammatory cytokines like TNF α (Baichwal, V.R. & Baeuerle, P.A.
15 (1997) Curr Biol 7, R94-6). To substantiate that mis-targeting of IKK β is an effective way
of blocking the functional effect of IKK β , we analysed whether PS473 was able to
influence TNF α -induced apoptosis. As seen in example 1 the probe (PS473) was found
to hypersensitise cells to apoptotic stimuli.

20 In another embodiment the present invention provides agents that modulate the
targeting and/or redistribution of IKKs. Such agents include polypeptides that comprise a
putative leucine zipper region of IKK β . Included are DNA molecules and expression
vectors that encode for the described peptides, furthermore host cells are provided that
express said peptides in a stable or transient expression system.

25

In another embodiment the invention provides a method for finding compounds that
modulate targeting and redistribution of IKK β and of derivatives thereof. The method
renders itself to screening for compounds that modulate the functional activity of I-
kappaB kinase β through modulation of one or more of multiple targeting sites of IKK β
30 (or other IKKs) and which thereby cause either a partial or a complete inhibition of the
NF-kappaB activation. The method will allow for identification of compounds that
modulate said targeting or redistribution in specific cell types.

The presented novel mechanism of action will be useful in the treatment of the following
35 diseases/conditions: chronic inflammation, asthma and chronic bronchial hyperreactivity

of non-asthma etiology, rheumatoid arthritis and pelvospondylitis, ulcerative colitis and Crohn's disease, diabetes mellitus type I, systemic lupus erythematosus, myasthenia gravis, Hashimoto's thyroiditis, Graves' disease and immune thrombocytopenic purpura, acute respiratory distress syndrome (ARDS) and septic shock as well as
5 depression.

Background

Chronic inflammation is the result of unbalanced and continued production of
10 inflammatory cytokines. Cytokines are produced in cascades, the pro-inflammatory $\text{TNF}\alpha$ and $\text{IL-1}\beta$ often responsible for initiating a process, which leads to a more general production of further cytokines. This cascade of gene expression is largely under the control of NF-kappaB, a ubiquitous transcription factor that, by regulating the expression of multiple inflammatory and immune genes, plays a critical role in host defence and in
15 chronic inflammatory diseases (Sen and Baltimore, 1986; Mukaida *et al.*, 1990; Beg *et al.*, 1993; Cogswell *et al.*, 1993). NF-kappaB is activated not only by cytokines, but also by reactive oxygen species (ROS), viruses, and a range of other generally noxious and pathogenic stimuli (Blackwell *et al.*, 1997; Schulzwe-Osthoff *et al.*, 1997). Activation of NF-kappaB via ROS has been implicated in neurodegenerative disorders such as
20 Parkinson's and Alzheimer's (Lesoualc'h *et al.*, 1998; O'Neill *et al.*, 1997) and also in inflammatory bowel disease (Jourd'heuil *et al.*, 1997). Tissue inflammatory response to x-rays is mediated directly by NF-kappaB (Hallahan *et al.*, 1995). Activation of NF-kappaB has been implicated in the production of atherosclerotic lesions of smooth muscle cells (Bourcier *et al.*, 1997) and in cardiac inflammatory disorders (Hattori *et al.*, 1997). NF-
25 kappaB/Rel transcription factors are also known to play a role in the pathogenesis of certain tumours, especially those of haematopoietic origin (Neumann *et al.*, 1997), and constitutive (autocrine) activation of NF-kappaB is known to promote a resistance to apoptotic stimuli (Giri *et al.*, 1998). Inhibitors of NF-kappaB should increase the cytotoxic efficacy of anticancer chemotherapies (Bours *et al.*, 1998).
30 The inflammatory pathways are notoriously complex, yet the feasibility of reducing or eliminating inflammatory responses through modulation of NF-kappaB activity has already been demonstrated in a number of different cells (Makarov *et al.*, 1997).

The NF-kappaB/Rel group of transcription activators and their co-evolved regulatory
35 proteins, the inhibitors of kappa B (I-kappaBs), play important roles in many cellular

- signalling processes in vertebrates, which include controlling communication between cells, embryo development, maintenance of cell type specific expression of genes as well as co-ordinating the inflammatory response to stressors and viral infection (Wulczyn *et al.*, 1996). The key proteins involved in this control system divide into distinct groups:
- 5 a) Those that bind DNA. These belong to the Rel family of transcription factors (Ghosh *et al.*, 1990) and include p50, p65, p52/49, p75/Rel and RelB. Only dimers bind DNA, but these can be homodimers or heterodimers. p65/p50 heterodimer is the most abundant, and plays a more elaborate role than other factors in regulating gene expression (Baldwin, 1996).
 - 10 b) Those that interact with the DNA-binding subunits in cytoplasm, which include the inhibitory I-kappaB α and I-kappaB β molecules (Bauerle and Baltimore, 1988), and the precursor molecule p105 (Naumann *et al.*, 1993).
 - c) Those transcriptional coactivators which interact with the DNA-binding subunits in the nucleus, such as Bcl3 (Nolan *et al.*, 1993; Watanabe *et al.*, 1997) and Cbp/p300 (Zhong *et al.*, 1998).
 - d) Kinases which activate proteasomal destruction of I-kappaB α and β subunits - the I-
 - 15 kappaB kinases (Beg *et al.*, 1993).
 - e) Kinases which directly phosphorylate the DNA-binding subunits in cytoplasm and nucleus to modulate their activity, such as PKA (Zhong *et al.*, 1998), casein kinase II (Bird *et al.*, 1997) and others (Hayashi *et al.*, 1993; Schulze-Osthoff *et al.*, 1997).
- 20 Inactive p65/p50 NF-kappaB dimers are held in the cytoplasm coupled to inhibitory I-kappaB molecules (α and β isoforms) via the p65 subunits. Activated I-kappaB kinases (IKK) phosphorylate the inhibitors, targeting them for ubiquitination and subsequent proteasomal digestion (Beg *et al.*, 1993). The released subunits translocate to the nucleus and there activate transcription.
- 25 The I-kappa kinases (IKK- α , IKK- β and IKK- γ) have been shown to be part of a large multi-component complex (Chen *et al.* 1996; Rothwarf *et al.*, 1998). It is likely to assume that the assembly and disassembly of the IKK complex is controlled by a scaffold protein termed IKK-complex-associated protein, IKAP (Cohen *et al.* 1998). It is expected that a tight assembly of the complex is necessary for the IKKs to be activated by the NF-kappa-
- 30 B-inducing kinase (NIK) and thereby induce phosphorylation of the I-kappaB subunits. Interestingly the affinity of IKK- β for IKAP diminishes upon phosphorylation of IKK- β by NIK.

Glucocorticoids (GC) are powerfully efficient modulators of inflammation, but suffer from
35 the potential hazards of suppressing necessary protective responses to infection and

decreasing some essential healing processes. They modulate cytokine expression by a combination of genomic mechanisms. The activated GC-receptor complex can (i) bind to and inactivate AP-1 or NF-kappaB, (ii) upregulate I-kappaB production via GC response elements (iii) reduce the half-life of cytokine mRNAs (Brattsand & Linden 1996). But
5 steroid treatment broadly attenuates all cytokine production from all lymphocytes, so not only do levels of the inflammatory cytokines fall, but also that of the anti-inflammatory IL-10. Specific modulation of Th1-type pathways would be an initial goal of this project. It is also known that some fibroblast cell NF-kappaB-mediated responses are likely governors of inflammatory progression, so inhibition of such responses could have
10 detrimental effects (Smith et al., 1997). Therapies, which maintain appropriate feedback systems, but modulate inappropriate cytokine production represent an unmet medical need.

An attractive therapeutic intervention to be used in the treatment of chronic inflammatory
15 conditions is inhibition of the I-kappaB degradation. Blocking the ubiquitin proteasome pathway (PharmaProjects, Accession no. 023654 and 027675), can directly inhibit this degradation. Another mechanism that is being pursued is inhibition of the enzymatic activity of either of the IKKs or NIK (public statement from Signal Pharmaceuticals).

20 Very many extracellular signals are transduced via intracellular systems employing the cyclic nucleotides cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP) as intermediaries, or second messengers. The processes mediated by cAMP and cGMP include control of smooth muscle tone, learning, vision, cellular differentiation, control of pro-inflammatory mediator production and action,
25 apoptosis, lipogenesis, glycogenolysis and gluconeogenesis, circadian rhythms, cardiac function, and mood control through noradrenergic potentiation.

Cyclic nucleotides are generated by adenylate and guanylate cyclases (ACs and GCs, respectively) from ATP and GTP, signal to cAMP- and cGMP-dependent effector proteins such as protein kinases (cAKs and cGKs, respectively) and certain ion
30 channels. cAMP and cGMP are removed by phosphodiesterases (PDE:s). The required specificity of signals generated by these systems arises from diversity of type, tissue-specific expression and intracellular placement of the enzymes involved. For instance there are nine isoforms of ACs known plus additional splice variants, soluble and membrane located forms of GCs, multiple isoforms of the cAK and cGK kinases, and
35 very many isoforms of PDE:s of which over 30 have been identified (Perry and Higgs, 1998; Houslay and Milligan, 1997; Beavo, 1995). Additional specificity arises from

targeting particular signalling enzymes to restricted locations within cells; this is the function of scaffold and anchoring proteins, such as the AKAP family, and not only may they place enzymes close to their substrates, but they may also serve to recruit multiple enzymes into functional signalling units (Pawson and Scott, 1997).

- 5 Inactivation of cAMP/cGMP occurs by hydrolysis of the 3'-ester bond, catalysed by the PDEs. The PDE:s are key components of the cyclic nucleotide signalling systems, allowing local concentration differences of the cyclic nucleotide messengers to be established, between adjacent tissues, between adjacent cells, even within a single cell between different volumes of cytoplasm. The ability to generate such heterogeneity in
- 10 the distribution of concentrations of a commonly shared signalling molecule, such as cAMP, is at the heart of directed signalling processes. To be of therapeutic value, cyclic nucleotide control has to be achieved with defined cellular selectivity (Perry and Higgs, 1998). It is the therapeutic opportunities offered by certain of the PDE:s that concerns this application.
- 15 Ten families of PDE:s have been identified, designated simply PDE1 to PDE10. Within each family there are two or more related but distinct gene products (A, B, C, etc.) and for each of these alternative mRNA processing gives rise to multiple splice variants, identified by an additional arabic numeral in accordance with the most recent nomenclature recommendation (Molecular Pharmacology 46:399-405, 1994). All PDE
- 20 gene products identified so far have two functional domains per molecule, one catalytic, and one regulatory. The catalytic domain lies towards the carboxylic acid terminus of each PDE protein and has the greatest homology between the PDE families, being >75% homologous at the amino acid level (Perry and Higgs, 1998). Nevertheless, each of the more than 30 PDE:s known have individually distinct substrate specificities, kinetic
- 25 characteristics, regulatory properties and cellular and subcellular distributions (Houslay and Milligan, 1997).
- PDE:s 4, 7 and 8 are highly specific for cAMP. PDE:s 5, 6, 9 and 10 are selective for cGMP. PDE3s bind cAMP and cGMP with similar affinity, but hydrolyse cAMP most efficiently, cGMP rather poorly. PDE3s are therefore negatively regulated in their cAMP
- 30 hydrolysing ability by cGMP. PDE:s 1 and 2 hydrolyse both cAMP and cGMP, but with PDE1 the relative efficiencies vary with isoenzyme subtype (Perry and Higgs, 1998). The amino terminal ends of PDE:s consist of the regulatory domains, which are very different both between families and between variants within families. This region contains variously: a binding domain for Ca^{2+} -calmodulin (CaM) in PDE1; non-catalytic cGMP-
- 35 binding sites in PDE:s 2, 5 and 6; a binding domain for the signalling G-protein

transducin in PDE6. The amino terminal region also contains protein- and membrane-targeting sequences in several PDE3:s and PDE4:s, as well as protein kinase phosphorylation sites in PDE:s 1, 3, 4 and 5. These phosphorylation sites are likely to be important in regulation of catalytic activity and/or subcellular location (Perry and Higgs, 5 1998).

Amongst the cAMP degrading phosphodiesterases, we focus here on the largest and most diverse family known, the PDE4:s. PDE4 enzymes share a common structure, as deduced from their amino acid sequences (Beavo and Reifsnnyder, 1990; Bolger *et al.*, 10 1993, Houslay, Sullivan and Bolger, 1998). Members of each gene family (PDE4A, PDE4B, PDE4C, PDE4D) share common C-terminal regions, different for each family, and catalytic domains that for all PDE4 isoforms are very similar (84% homology over about 360 amino acids across all PDE4:s; Houslay, Sullivan and Bolger, 1998). From N-terminus to catalytic region, the sequence in "long form" PDE4s can be divided into 5 15 regions, three of which are isoform-specific (N-terminal region, linker regions 1 and 2, or LR1 and LR2) and two, more conserved regions, that are broadly similar between all isoforms, the upstream conserved regions 1 and 2 (UCR1 and UCR2). "Short form" PDE4:s, e.g. PDE4A1, PDE4B2, PDE4D1, PDE4D2, lack UCR1 and LR1 plus differing amounts of the N-terminal region of UCR2. Throughout all regions are potential 20 phosphorylation sites for a variety of kinases, including PKA (e.g. Ser 54 in human PDE4D3), mitogen activated protein kinases (e.g. Ser 487 of human PDE4B2), casein kinase II (e.g. Ser 489 of PDE4B2) and calcium-diacylglycerol dependent protein kinases (Houslay, Sullivan and Bolger, 1998). Phosphorylations at some of these sites have been shown to activate the PDEs (e.g. Ser 54), others serve to inhibit. There is also 25 evidence that some phosphorylations serve to prime the enzymes ready for subsequent activation by further phosphorylation at a different site or sites (Houslay, Sullivan and Bolger, 1998). Other auto-regulatory sites may be found in the N-terminal sequence of certain PDE4:s (Bolger *et al.*, 1996, McPhee *et al.*, 1995).

The identification of rolipram (Schering AG, Berlin, Germany) as an effective inhibitor of 30 PDE4:s (Wachtel, 1982, Nemoz *et al.*, 1985) gave an important tool by which to determine the role of PDE4:s in different cell types. Originally developed as a neurotropic agent, rolipram indicated the therapeutic potential of PDE4 inhibition in control of depressive disorders. Analysis of the pharmacological properties of rolipram, and over 800 publications covering these properties have appeared over the period 1993 to 1998 35 alone, now indicates that specific PDE4 inhibition may be useful over a very wide range of disease areas. These include: asthma, atopic dermatitis, depression, reperfusion

injury, septic shock, toxic shock, autoimmune diabetes, AIDS, Crohn's disease, multiple sclerosis, cerebral ischemia, psoriasis, allograft rejection, restenosis, ulcerative colitis, cachexia, cerebral malaria, allergic rhinoconjunctivitis, osteoarthritis, rheumatoid arthritis, autoimmune encephalomyelitis (Houslay, Sullivan and Bolger, 1998).

- 5 In the area of asthma, PDE4 inhibition helps to increase cAMP in bronchial smooth muscle, thereby producing a modest bronchodilatory effect, of use in the alleviation of asthmatic symptoms. But perhaps most importantly, inhibition of PDE4:s is now a recognised method by which to suppress immune and inflammatory cell responses (Hughes *et al.*, 1997; Torphy, 1998; Teixeira *et al.*, 1997).
- 10 PDE4:s play major roles in modulating the activity of virtually every cell type involved in the inflammatory process. Immune and inflammatory conditions occur when recruitment of leukocytes from the blood compartment into tissues is either uncontrolled, inappropriate, prolonged or directed against self. In asthma, rheumatoid arthritis and multiple sclerosis, infiltration of tissues with inflammatory cells is prolonged and intense,
- 15 leading ultimately to severe (and self-perpetuating) damage and loss of function. Acute dysregulation of the immune system occurs in such conditions as acute respiratory distress syndrome (ARDS) where an overwhelming and generalised inflammatory response can frequently lead to death. There is also substantial evidence which suggests that inflammation may play a part in defining the extent of injury resulting from
- 20 reperfusion following ischaemia, at least in brain and lung (Entman and Smith, 1994). Chronic inflammatory conditions such as asthma are currently treatable with steroids, but long term treatment brings unavoidable side-effects including immunosuppression, metabolic disturbance and hypertension (Teixeira *et al.*, 1997). Symptoms of rheumatoid arthritis can be alleviated by non-steroidal anti-inflammatories (NSAIDS), but again their
- 25 side effects are of great concern. Acute conditions such as ARDS have no current treatment as such, only supportive care. Effective anti-inflammatories able to control dysregulated responses, but without the side effects associated with NSAIDS and steroids, have not yet been found.

Within the context of asthma, elevation of intracellular cAMP by PDE inhibition has been

30 associated with inhibition of the function of various types of cells involved in the inflammatory response, including lymphocytes, monocytes, macrophages, neutrophils, eosinophils, mast cells, basophils, endothelial cells and lung epithelial cells (Nicholson and Shahid, 1994); PDE4:s appear to play the dominant role in neutrophils, basophils, eosinophils and mast cells, PDE3s being dominant in monocytes/macrophages and

35 lymphocytes. Inhibitors of PDE3s and PDE4:s often interact synergistically in control of

inflammatory response in asthma models (Teixeira *et al.*, 1997). Other PDE:s may be important in inflammatory cells, but their involvement has yet to be clarified or demonstrated.

Increased cAMP modulates myosin light chain kinase (MLCK) activity causing relaxation, and this is the primary effect in bronchial smooth muscle. Useful compounds will relax bronchial smooth muscle slowly and maintain relaxation for sustained periods, but also help reduce inflammatory immune responses to allergens. Although a combined inhibition of PDE3 and PDE4 isozymes seems to relax bronchial smooth muscle most effectively (Raeburn & Advenier, 1995) in humans, the possibility of cardiovascular complications is increased by the use of PDE3 inhibitors, and in fact PDE4 inhibitors such as rolipram, alone or in combination with agonists of the β_2 adrenoceptors such as salbutamol, are effective bronchorelaxants.

Possible mechanisms (Teixeira *et al.*, 1997) involved in the anti-inflammatory benefits of PDE4 inhibition *in vivo* include:

- 15 - Inhibition of the production and release of inflammatory mediators/cytokines.
- Inhibition of leukocyte migration.
- Induction of cytokines with suppressive activity.
- Inhibition of leukocyte activation (degranulation, respiratory burst).
- Inhibition of the expression/upregulation of cell adhesion molecules.
- 20 - Induction of apoptosis amongst inflammatory cells.
- Also, stimulation of endogenous steroid and catecholamine release (Pettipher *et al.*, 1996).

Perhaps the most important consequence *in vivo* of selective PDE4 inhibition may be to inhibit chemokine production, especially those that are chemoattractants of leukocytes (Teixeira *et al.*, 1997). Inhibitors of PDE4 are effective suppressers of cytokine production *in vitro* and reduce serum levels of tumor necrosis factor alpha (TNF- α) in animal models of septic shock (Sekut *et al.*, 1995; Pettipher *et al.*, 1996; Prabhakar *et al.*, 1994). Inhibition of TNF- α production may be central to the beneficial effects of PDE4 inhibition in treatment of inflammatory conditions, but inhibition of the release of chemoattractants such as the α -chemokine interleukin-8 and the lipid leukotriene (LT) B_4 may also be important for reducing leukocyte recruitment to sites of inflammation (Turner *et al.*, 1994; Griswold *et al.*, 1993).

It is also known however that there are protective effects of PDE4 inhibition which are quite separate from inhibition of release and action of TNF- α and other pro-inflammatory mediators. At higher concentrations than are necessary to inhibit TNF- α release,

rolipram appears to have a direct effect on eosinophils (Teixeira *et al.*, 1994) and eosinophilia. PDE4 inhibition also stimulates macrophages to produce and release the antiinflammatory cytokine interleukin 10 (IL-10) when challenged with lipopolysaccharide (LPS) *in vitro* (Kambayashi *et al.*, 1995; Jilg *et al.*, 1996), and this same effect may be
5 involved in the protective action of methylxanthines, which are general PDE inhibitors, in a murine model of septic shock (Jilg *et al.*, 1996).

Inhibition of neutrophil activation *in vivo* may also be how PDE4 inhibition protects against acute lung injury induced by LPS followed by zymosan in a murine model (Miotla *et al.*, 1995), and in animal models of asthma, it is likely that PDE4 inhibition suppresses
10 allergic inflammation by inhibition of eosinophil activation together with inhibition of mast cell de-granulation (Hughes *et al.*, 1996).

PDE4 inhibition has also been shown to affect the *in vitro* expression and presentation of cell adhesion molecules such as E-selectin by endothelial cells of the microvasculature (Blease *et al.*, 1998; Morandini *et al.*, 1996) and increased cAMP also prevents mediator-
15 induced upregulation of $\beta 2$ integrins on the surface of eosinophils and neutrophils (Teixeira *et al.*, 1996). Inhibition of the cell adhesion components responsible for recruitment of leukocytes and for initiation of tissue infiltration by the inflammatory cells is an important aspect of therapeutic control for inflammatory conditions.

cAMP-elevating agents also enhance apoptotic clearance of various leukocytes *in vitro* (Hallsworth *et al.*, 1996), and this too may be useful effect in the control of inflammation
20 through PDE4 inhibition.

The major cGMP-degrading PDEs are types 1,2,5, 6, 9 and 10 but here we focus on PDE5, since this is the principal cGMP-specific PDE found in airway and vascular
25 smooth muscle, and it is one of the better documented families of cGMP-specific PDEs. Little is known yet concerning the role of the newly discovered PDE9 and PDE10 isoforms (Soderling *et al.*, 1998; Fisher *et al.*, 1998; Soderling *et al.*, 1999; Fujishige *et al.*, 1999), and the situation is similar for PDE2s, since good inhibitors are as yet unknown for these (Perry and Higgs, 1998). PDE5 is activated by cAK and (10-fold
30 faster) by cGK (Thomas *et al.*, 1990). Phosphorylation of PDE5 is enhanced in the presence of cGMP, and apparently increases the enzyme's V_{max} by 10-fold (Burns *et al.*, 1992). Coupled with PDE3, these interactions form a feedback system to limit cGMP signaling: increased cGMP will increase cAMP through inhibition of PDE3, high cAMP will activate cAK which, in the presence of elevated cGMP will activate PDE5 and
35 therefore stimulate cGMP breakdown. cAMP levels return to baseline as cGMP falls, by re-activation of PDE3. Recent evidence (Pyne *et al.*, 1996; Lochhead *et al.*, 1997)

suggests that PDE5 may have additional protein components associated with it analogous to the gamma subunits of PDE6. The PDE6 γ subunits serve to link activation of the G-protein transducin to activation of the PDE. They are subsequently involved in turning off the signal by helping to activate the transducin GTPase. In the case of PDE5, these associated proteins (14 to 18 kDa) may serve to block activation of the enzyme by cGK and cAK, and the blocking ability of these polypeptides appears to be controlled by a G-protein regulated kinase (Pyne *et al.*, 1996).

cGMP-degrading PDEs work in concert with the action of guanylate cyclases, just as cAMP PDEs and adenylate cyclases together control cAMP levels in cells. Two groups of GCs are known in mammals, the soluble ones and those that are membrane located. GCs from both groups are central to systemic control of blood pressure. Soluble GCs are expressed in almost all cell types of the cardiovascular system including cardiomyocytes, vascular smooth muscle cells (VSMCs), endothelial cells and platelets (Drewett and Garbers, 1994). Soluble GCs contain a prosthetic heme group which binds NO (and CO) and leads to activation of the enzyme: the vasoactive properties of NO are mediated through the cGMP pathway in this way. The membrane located GCs act as receptors for various ligands (among them, natriuretic peptides and guanylin). cGMP-mediated functions of the natriuretic hormone receptors include vascular smooth muscle relaxation as well as regulation of blood volume (Benner *et al.*, 1990).

cGMP interacts with a number of different effector proteins:

- a) with certain ion channels e.g. in photoreceptors and olfactory cells, also in heart and kidney (Lincoln & Cornwell, 1993; Biel *et al.*, 1994; Light *et al.*, 1990);
- b) with cGMP-dependent protein kinases (cGKI and cGKII), of which "cytosolic" cGKI predominates in the cardiovascular system and has at least 2 splice variants, α and β . cGKI α has 10-fold higher affinity for cGMP than the β variant. Both cGKI variants are found in vascular smooth muscle (Keilbach *et al.*, 1992, Hofmann *et al.*, 1992);
- c) at high concentrations, with cAMP-dependent protein kinases (cAK), which being similar to the cGKs have a certain affinity for cGMP, just as the reverse is also true (Vaandrager & de Jonge, 1996). The functional significance of this potential cross-talk between pathways is not yet fully known, but may be connected with the anti-proliferative effects of cGMP (Lincoln *et al.*, 1994);
- d) with cGMP-modulated PDEs: cGMP binds to a non-catalytic site of PDE2 and lowers its K_m for cAMP, lowering the baseline level of cAMP achievable by the enzyme. PDE3 catalysis of cAMP is effectively inhibited by cGMP (Pyne *et al.*, 1987), thus in cells where PDE3 predominates, increased cGMP leads to increased cAMP.

Smooth muscle contracts following Ca^{2+} -calmodulin activation of myosin light chain kinase (MLCK). cGKI relaxes smooth muscle by lowering free cytoplasmic Ca^{2+} levels, but the principal means by which this is accomplished varies considerably between types of smooth muscle, animal species, and the nature of the contractile stimulus being

5 antagonised (Vaandrager & de Jonge, 1996). cGKI has been implicated in: inhibition of G-protein activation of phospholipase C β ; activation of Ca^{2+} -ATPase activity at plasma membrane and sarcoplasmic reticulum (SR); hyperpolarisation of membrane potential through activation of Ca^{2+} -activated K^+ channels; inhibition of voltage operated Ca^{2+} channels; stimulation of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger; inhibition of SR IP_3 receptors. All of

10 these actions require that the normally cytoplasmic cGKs must find membrane located targets, and specific anchor proteins may be involved. cGKI is already known to be targeted to specific anchor proteins of the cytoskeleton (MacMillan-Crow & Lincoln, 1994), and the discovery of further interactions is likely.

Blood pressure elevation to a degree that requires medical treatment is often

15 encountered in up to 15% of an adult population. In only 10-15% of these, a definite cause for the hypertension can be found and in the rest, the "essential hypertension" has to be treated without a hope for cure of the underlying disease. Long-standing elevation of blood pressure, even quite moderate, damages vessels in the heart, kidneys and brain and dramatically increases the risk for coronary heart disease, renal failure and

20 stroke. It has been shown that effective pharmacologic treatment of hypertension substantially reduces morbidity and mortality from these conditions. The finding that endothelial cells produce a local vascular relaxation factor, identified as nitric oxide (NO), that activates guanylyl cyclase and increases cGMP that in turn leads to reduction in vascular smooth muscle cell tone, has opened new possibilities for blood pressure

25 regulation / vasorelaxation based on modulation of the cellular levels of cGMP. A number of the components in the cGMP system displays tissue specific distribution (Vaandrager & de Jonge, 1996; Pyne *et al.*, 1996). This increases the likelihood for improved pharmacological specificity and fewer side-effects when using these as targets for antihypertensive treatment instead of the traditional ones. It is the cGMP-dependent

30 protein kinase (PKG) (Vaandrager & de Jonge, 1996) that is thought to mediate the intracellular effects of cGMP. The cGMP -dependent and -specific phosphodiesterases can serve as connectors to the cAMP system and terminators of cGMP effects (Pyne *et al.*, 1996).

PDE5 has attracted attention since it is selective for degradation of cGMP versus cAMP.

35 Isoform-specific inhibitors for PDE5 are being developed by several companies and one

compound from Pfizer, Sildenafil, has proven selectivity for PDE5 and is currently being marketed as treatment against impotence (Viagra), originally a side-effect resulting from vasorelaxation in the corpus cavernosum. However the screening procedures currently used search only for direct enzymatic inhibitors of PDE and the compounds found are
5 often not selective, inhibiting for instance both PDE 1 and 5 (e.g. Zaprinast (M&B 22948 RPR), Sch 59498 and Sch 51866). By the methods described herein and within appendix A, new chemical entities can be found which primarily will be specific modulators of PDE action, not inhibitors of the enzymatic action *per se*. Preferred compounds will inhibit the site-specific anchoring of PDEs which hydrolyse cGMP, and
10 thereby reduce their effectiveness in controlling local concentrations cGMP within living cells.

The therapeutic potential of selective modulators of cGMP-related PDE action is not restricted to relaxation of smooth muscle cells but also encompasses other effects ascribed to PKG, such as inhibition of platelet activation (Chiu *et al.*, 1997; Vemulapalli
15 *et al.*, 1996), inhibition of endothelial permeability increases in response to vasoactive substances (Raeburn & Karlsson, 1993), inhibition of the differentiation of osteoclasts (Holliday *et al.*, 1997) and light-induced resetting of circadian rhythms (Mathur *et al.*, 1996; Liu *et al.*, 1997).

20 The search for chemical inhibitors of the catalytic activity of specific PDE:s is currently one of the most intensive areas of pharmaceutical research, particularly so for PDE:s 4 and 5. Much progress has been made in this area, with several compounds known to have selective activity for particular families of PDE:s (reviewed in Perry and Higgs,
25 1998; Hughes *et al.*, 1997; Teixeira *et al.*, 1997). However, there has not yet been found a class of compounds able to select between isoenzymes within the same family, which is where the greatest opportunities lie. Without isoform specificity, certain difficulties can be expected with the use of enzymic inhibitors of PDE:s. Some of these difficulties are outlined below.

30 In general, the effects a known inhibitor of the catalytic activity of a particular class of PDE:s may have on cyclic nucleotide levels often varies between different cell types. The reasons for this are several, but include: differences in the basal level of cyclase activity in distinct cell types, crosstalk between cAMP and cGMP systems, and differences in
35 local concentrations of substrate within a cell which influences the degree of inhibition that can be attained by a simple competitive enzyme inhibitor (Perry and Higgs, 1998).

First, PDE inhibition is only useful if it produces the appropriate change in the activity of the dependent effectors, for instance activation of cAK when the concentration of cAMP can be increased above a threshold level. The rate of change in concentration depends in part on the activity of the cyclases which generate the cyclic nucleotides, and that
5 basal level of activity differs from isoform to isoform, and therefore from cell type to cell type. In adipocytes, for example, AC activity is high and cAMP levels are kept at baseline only by a correspondingly high PDE activity. Hepatocytes on the other hand have a rather low AC activity. If both cell types share PDE:s of the same family, and are treated with a chemical inhibitor targeting that family, there will be a rapid increase in cAMP
10 within adipocytes and activation of their cAKs, but no activation in hepatocytes, unless the AC is also stimulated.

Second, general inhibition of a particular isoform of PDE can have certain unavoidable consequences on other cyclic nucleotide pathways since cAMP and cGMP systems are often closely interlinked. Much of this crosstalk arises from PDE regulation by cyclic
15 nucleotides. When cGMP increases in platelets (e.g. following nitric oxide stimulation of soluble GC, or PDE5 inhibition) it inhibits PDE3 and causes a concomitant rise in cAMP (Ashida and Sakuma, 1992). In adrenal glomerulosa cells, atrial natriuretic factor elevates cGMP but inhibits cAMP-stimulated aldosterone synthesis via cGMP-stimulation of PDE2 (MacFarland *et al.*, 1991).

20 Third, the expected effects of PDE inhibition may be modified by differences in local concentrations of substrates, the reason being that most chemical inhibitors of PDE action are competitive with substrate, so their therapeutic profile is dependent on both the Michaelis-Menton equilibrium constant (K_M) and the substrate concentration in which they are operating (Perry and Higgs, 1998). Most effective inhibition will always occur at
25 lowest substrate levels, but as a corollary, a locally increased substrate level will reduce the inhibition attained. In combination with subtle differences in isoform K_M values for an inhibitor, the desired spatial modulation of cyclic nucleotide levels within a cell could be difficult to obtain by simple competitive inhibition of catalytic activity.

Fourth, there is increasing evidence that cells respond to the prolonged use of agents
30 that increase cyclic nucleotide concentrations by increasing the activity of endogenous levels of appropriate phosphodiesterases (Torphy *et al.* 1995), and that one class of mechanism whereby this occurs is by increasing expression levels of PDE proteins (Swinnen *et al.*, 1989, 1991). There is even evidence to suggest that the use of selective inhibitors of different PDE families (eg rolipram for PDE4:s, cilostimide for PDE3,
35 zaprinast for PDE5 etc.), encourages cells and tissues to respond to catalytic inhibition

by upregulating PDE:s specifically of the family type that is under inhibition. Full catalytic inhibition of PDE:s may therefore have self-defeating results, as cells attempt to compensate for lack of specific PDE activity. Careful modulation of local cyclic nucleotide levels within a cell through dislocation or inhibition of redistribution, which may not
5 greatly affect global levels of cyclic nucleotide, may therefore prove to be a better and more effective means to achieve long term therapy.

The radically different methods of interference with PDE action as proposed below in this application should avoid many of the problems outlined above, principally because
10 interference will be family and isoform specific and targeted not against catalytic activity of the PDE:s, but their spatial organisation within the cell.

Targeting of signalling enzymes is a recognised mechanism by which sensitivity, specificity, precision and control may be introduced into intracellular signalling pathways
15 (Pawson and Scott, 1997; Faux and Scott, 1996). The importance and occurrence of targeting as a phenomenon are described and discussed in appendix A. Of central importance to this application is the modulation of the effectiveness of signalling PDE:s through interference with their intracellular targeting. As already described, the many PDE:s known share much structural homology, and this is especially true within the
20 catalytic regions found towards the carboxylic acid terminals of the proteins. At the amino terminals much more heterogeneity is found, between families of PDE:s, between isoforms within families, and between splice variants derived from individual gene isoforms (Houslay and Milligan, 1997). Much of this heterogeneity appears to be associated with differences in targeting behaviour, at least in PDE4 isoforms and
25 variants (Scotland *et al.*, 1998, Bolger *et al.*, 1997), and by extension should apply to other PDEs as well since they are in overall character similar protein molecules with similar roles in cellular signalling.

Evidence suggests that the amino terminal regions of PDE:s can serve to target isoforms to specific intracellular sites (Shakur *et al.*, 1995; McPhee *et al.*, 1995; Bolger *et al.*,
30 1996; Pooley *et al.*, 1997) and that they can regulate the functioning of the catalytic unit either through interaction with binding proteins (Shakur *et al.*, 1995; O'Connell *et al.*, 1996; Pyne *et al.*, 1996) or through phosphorylation (Sette and Conti, 1996). Targeting appears to occur through protein-protein interactions with membrane- or cytoskeletally-located proteins (Houslay, Sullivan and Bolger, 1998), and of these the membrane
35 associated proteins include both integral and peripherally adherent species. Such

interactions have been probed at a gross level through the use of nonionic detergents and elevated ionic strength (Scotland *et al.*, 1998).

Four separate genes are known to produce PDE4:s in human and rat (PDE4A-D), and each of these produces multiple splice variants (more than 20 described to June 98),
5 many with unique amino terminal regions (Huston *et al.*, 1997; Bolger *et al.*, 1997; Obernolte *et al.*, 1997). Some variants have extensive deletions, even to the point of removing catalytic activity (Obernolte *et al.*, 1997). Differences in the amino terminal regions are presently contemplated to be important for determining differences in the subcellular localisation, activity and sensitivity to inhibitors amongst PDE4 isozymes
10 (Bolger, 1997; Scotland *et al.*, 1998). As an example, PDE4D1 and PDE4D2 are found only in cytosolic fractions, PDE4D3, D4 & D5 are all represented in both cytosolic and particulate fractions. PDE4D3 and D5 are both more sensitive to rolipram inhibition in the cytosolic phase than they are in the particulate fraction (Bolger *et al.*, 1997). Of the 3 "B" isozymes, PDE4B2 is approximately 10 fold more sensitive to rolipram in the particulate
15 fraction than in the cytosolic (Huston *et al.*, 1997). Certain PDE4 isozymes are known to have restricted tissue distributions, e.g. PDE4A8 and PDE4C-delta54 are found only in testis, PDE4C-791 in lung and a melanoma cell line G361 (Bolger *et al.*, 1996; Obernolte *et al.*, 1997). In other cells the expression of isozymes changes with cellular differentiation (Verghese *et al.*, 1995; Giorgi *et al.*, 1997; Bolger *et al.*, 1994; Essayan *et al.*, 1997).
20

Certain PDE4 isozymes are known to associate with membranes, some with proteins bearing SH3 domains, and some to be purely cytosolic (Scotland *et al.*, 1998; Bolger *et al.*, 1997). A variant of PDE4A ("RD1") transfected into human thyroid carcinoma lines accumulates specifically in Golgi, and at the same time inhibits all expression of "native"
25 PDE1 in those cells (Pooley *et al.*, 1997). These distinct locations are believed to reflect very different functions of the specific phosphodiesterases. A very clear demonstration of functional separation of PDE:s has been seen in renal mesangial cells. Immuno-inflammatory stimulation of these cells increases their production of reactive oxygen metabolites (ROM) and simultaneously increases proliferation. Specific inhibition of
30 PDE4 suppresses ROM production, but not proliferation. Specific inhibition of PDE3 inhibits proliferation but not ROM production (Chini *et al.*, 1997). Both responses are mediated by PKA but control of the cAMP pool is effectively separated.

Location of PDE:s to membranes brings them into contact with phospholipids. Certain PDE4 isozymes are activated by anionic phospholipids such as phosphatidyl serine and

phosphatidic acid (Disanto *et al.*, 1995; Nemoz *et al.*, 1997). Dislocation from the membrane will inhibit such activation, and crosstalk with phospholipid signalling systems. Targeting or anchoring of PDE4:s is likely to have its greatest effect through compartmentalisation of cAMP signalling within cells (Houslay and Milligan, 1997).

5 Associated with the PDE4:s will be specific ACs together with specific isoforms of the effector cAK, or cAMP-operated ion channels. cAKs will likely be attached to specific AKAPs (A-kinase anchoring proteins). Specific subcellular distributions of these components have been mapped in cells (Houslay and Milligan, 1997; Scott and Pawson, 1997; Coghlan *et al.*, 1995) and allow for spatial and temporal gradients of cAMP to be

10 established within cellular compartments. Targeted PDE4 species might serve to control threshold levels of cAMP in the environs of specific cAK molecules, perhaps protecting certain protein complexes from cAK-mediated phosphorylation or manipulating the activity levels of ACs that are necessary before cAK activation may occur.

15 Competitive chemical inhibitors are known which can selectively inhibit members of the PDE4 family. There are none known which can effectively select between the different gene products or splice variants of the PDE4 family (Perry and Higgs, 1998). This may be due to the particularly high degree of sequence homology within the proteins of this family around the catalytic site. Without splice-variant selectivity, there are likely to be

20 problems with long-term administration of PDE4 inhibitors, such as immunosuppression and metabolic disturbances, possibly with significant CNS effect as well (Teixeira *et al.*, 1997) since PDE4:s are clearly involved in such a wide range of systems at the organismal level. For the family of PDE4 enzymes, the pyrrolidone compound rolipram remains the "gold standard" reference inhibitor. However, its profile of serious side

25 effects prevented rolipram from becoming a compound of clinical utility. Principal side effects of rolipram are headaches, nausea, emesis and an unacceptable increase in gastric acid secretion (Barnes, 1995). The PDE4 family is likely to consist of more than the 20 or so isoforms already known in humans (Houslay, Sullivan and Milligan, 1998). Although a potent inhibitor of all known isoforms of PDE4s, the kinetics of inhibition are

30 complex and sensitivity varies significantly from isoform to isoform, and even for individual isoforms in different cell backgrounds or cellular compartments (Bolger *et al.*, 1996; Huston *et al.*, 1996; Jacobitz *et al.*, 1996; McPhee *et al.*, 1995; Owens *et al.*, 1997; Wilson *et al.*, 1994). The side effects of rolipram clearly indicate the potential problems associated with general PDE4 inhibition, while different isoform sensitivities, and

35 changing sensitivities in different cellular contexts, highlights the potential functional

diversity of the many PDE4 isoforms known, and therefore the therapeutic potential that lies in selective inhibition of individual isoforms.

So far only two PDE5 genes are known and two enzyme variants have been reported. In parallel with other PDE isoforms more splicing variants are to be expected from each gene. The enzyme is a homodimer, each subunit being 93 kDa. The structural organisation of the dimer is very similar to that of the cGKs.

PDE5s exist in two distinct forms: one membrane-bound (mPDE5) and one cytosolic (cPDE5) (Pyne *et al.*, 1996). The mPDE5 is activated by PKA and is inhibited by a G-protein dependent mechanism. It is assumed that cPDE5 is part of a "signalling cassette" with NO-regulated guanylate cyclase and PDE3. The latter construction will lead to very short-lived messages whereas the former allows for generation of prolonged cGMP signals

Targeting or anchoring of PDE5s is likely to have its greatest effect through compartmentalisation of cGMP signalling within cells. Associated with the PDE5s will be specific GCs together with specific isoforms of the effector cGK, or cGMP-operated ion channels. cGKs may be attached to specific G-kinase anchoring proteins. Specific subcellular distributions of these components will allow for spatial and temporal gradients of cGMP to be established within cellular compartments. Targeted PDE5 species might serve to control threshold levels of cGMP in the environs of specific cGK molecules, perhaps protecting certain protein complexes from cGK-mediated phosphorylation or manipulating the activity levels of GCs that are necessary before cGK activation may occur.

Competitive chemical inhibitors are known which can selectively inhibit PDE5s. Relatively few isoforms of PDE5 are known to date. PDE5 is found rather specifically in vascular and airway smooth muscle. That sildenafil, with its 5 nM IC₅₀ for PDE5, affects only a subset of vascular smooth muscle is puzzling, but strongly suggests that either multiple PDE5 isoforms or states exist in different vascular smooth muscle, presumably with different sensitivities to sildenafil, or more likely, other cGMP-hydrolysing PDEs are important in different vascular smooth muscles.

As to other potentially important cGMP-hydrolysing PDE targets, many are doubtless yet to be discovered. PDE9:s have only been known since the end of 1997, PDE10:s since late 1998. PDE9:s have a rather general distribution (kidney, brain, lung), have a very high affinity for cGMP (about 70 nM) and are inhibitable by the PDE1/5 inhibitor SCH51866 (1.55 μ M), but "not by sildenafil" (7 μ M, Soderling *et al.*, 1998). Their

physiological roles and regulation have not been defined (Soderling *et al.*, 1998; Fisher *et al.*, 1998), but the best suggestions are that they may be involved in keeping cGMP at very low levels when activated, and may, in kidney, be involved in termination of ANP signalling, and therefore inhibition may help potentiate natriuresis without causing
5 deleterious drops in blood pressure (Soderling *et al.*, 1998).

It is clear that PDEs possess heterogeneity, particularly in their amino terminal, or "regulatory" regions, and the approach outlined in this application exploits those differences between isoforms and splice variants to produce what should be confined
10 and defined therapeutic effects. Furthermore, in many cases it may be expected that dislocation of an active enzyme from a targeted site of action will have little effect on average cellular concentrations of their preferred cyclic nucleotide substrate, although significant increases may occur at the now PDE-free site of action. This may have significance where an acute short-term process is the therapeutic target, but an
15 integrative gene-regulation effect may occur upon general, non-specific PDE inhibition and overall cyclic nucleotide increase in the cell.

Detailed disclosure

In the present specification and claims, the term "influence" covers any influence to
20 which the cellular response comprises a redistribution. Thus, e.g., heating, cooling, high pressure, low pressure, humidifying, or drying are influences on the cellular response on which the resulting redistribution can be quantified, but perhaps the most important influence is the influence of contacting or incubating the cell or cells with a substance which is known or suspected to cause a redistribution or modify a change of
25 redistribution. In another embodiment of the invention the influence could be substances from a compound drug library.

In the present context, the term "green fluorescent protein" (GFP) is intended to indicate a protein which, when expressed by a cell, emits fluorescence upon exposure to light of
30 the correct excitation wavelength (cf. Chalfie, M. *et al.* (1994) *Science* 263, 802-805). In the following, GFP in which one or more amino acids have been substituted, inserted or deleted is also termed "modified GFP". "GFP" as used herein includes wild-type GFP derived from the jelly fish *Aequorea victoria* and modifications of GFP, such as the blue fluorescent variant of GFP disclosed by Heim *et al.* (Heim, R. *et al.* (1994)).

- Proc.Natl.Acad.Sci. 91:26, pp 12501-12504), and other modifications that change the spectral properties of the GFP fluorescence, or modifications that exhibit increased fluorescence when expressed in cells at a temperature above about 30°C described in PCT/DK96/00051, published as WO 97/11094 on 27 March 1997 and hereby
- 5 incorporated by reference, and which comprises a fluorescent protein derived from *Aequorea* Green Fluorescent Protein or any functional analogue thereof, wherein the amino acid in position 1 upstream from the chromophore has been mutated to provide an increase of fluorescence intensity when the fluorescent protein of the invention is expressed in cells. Preferred GFP variants are F64L-GFP, F64L-Y66H-GFP and F64L-S65T-GFP. An
- 10 especially preferred variant of GFP for use in all the aspects of this invention is EGFP (DNA encoding EGFP which is a F64L-S65T variant with codons optimized for expression in mammalian cells is available from Clontech, Palo Alto, plasmids containing the EGFP DNA sequence, cf. GenBank Acc. Nos. U55762, U55763).
- 15 The terms "intracellular signalling pathway" and "signal transduction pathway" are intended to indicate the coordinated intracellular processes whereby a living cell transduces an external or internal signal into cellular responses. Said signal transduction will involve an enzymatic reaction said enzymes include but are not limited to protein kinases, GTPases, ATPases, protein phosphatases, phospholipases and cyclic
- 20 nucleotide phosphodiesterases. The cellular responses include but are not limited to gene transcription, secretion, proliferation, mechanical activity, metabolic activity, cell death.

The term "second messenger" is used to indicate a low molecular weight component

25 involved in the early events of intracellular signal transduction pathways.

The term "luminophore" is used to indicate a chemical substance which has the property of emitting light either inherently or upon stimulation with chemical or physical means. This includes but is not limited to fluorescence, bioluminescence, phosphorescence,

30 chemiluminescence.

The term "mechanically intact living cell" is used to indicate a cell which is considered living according to standard criteria for that particular type of cell such as maintenance of normal membrane potential, energy metabolism, proliferative capability, and has not

experienced any physically invasive treatment designed to introduce external substances into the cell such as microinjection.

In the present context, the term "permeabilised living cell" is used to indicate cells where
5 a pore forming agent such as Streptolysin O or *Staphylococcus Aureus* α -toxin has been applied and thereby incorporated into the plasma membrane in the cells. This creates proteinaceous pores with a defined pore size in the plasma membranes of the exposed cells. Pores could also be made by electroporation, i.e. exposing the cells to high voltage discharges, a procedure that creates small holes in the plasma membrane by
10 coagulating integral membrane proteins. Treatment with a mild detergent such as saponin may accomplish the same thing. Common to all these treatments is that pores are formed only in the plasma membrane without affecting the integrity of cytoplasmic structural elements and organelles. The term living in this context means that the permeabilised cell or cells bathed in a solution mimicking the intracellular milieu still have
15 functional organelles, such as actively respiring mitochondria and endoplasmatic reticulum that can take up and release calcium ions, and functional structural elements. In one embodiment this method is applied so that substances that normally can not traverse the plasma membrane, but most likely exert their influence intracellularly, can be introduced and their influence studied. In another embodiment this method is used to
20 record the response to an influence from many cells simultaneously.

In the present context, the term "permeabilisation" is intended to indicate the selective disruption of the plasma membrane barrier so that soluble substances freely mobile in the cytosol may be lost from the interior of the cells. The permeabilisation can be
25 achieved as described above under "permeabilised living cells" or by using other chemical detergents such as Triton X-100 or digitonin in carefully titrated amounts.

The term "physiologically relevant", when applied to an experimentally determined redistribution of an intracellular component, as measured by a change in the
30 luminescence properties or distribution, is used to indicate that said redistribution can be explained in terms of the underlying biological phenomenon which gives rise to the redistribution.

The terms "image processing" and "image analysis" are used to describe a large family
35 of digital data analysis techniques or combination of such techniques which reduce

ordered arrays of numbers (images) to quantitative information describing those ordered arrays of numbers. When said ordered arrays of numbers represent measured values from a physical process, the quantitative information derived is therefore a measure of the physical process.

5

The term "mammalian cell" is intended to indicate any living cell of mammalian origin. The cell may be an established cell line, many of which are available from The American Type Culture Collection (ATCC, Virginia, USA) or a primary cell with a limited life span derived from a mammalian tissue, including tissues derived from a transgenic animal, or
10 a newly established immortal cell line derived from a mammalian tissue including transgenic tissues, or a hybrid cell or cell line derived by fusing different celltypes of mammalian origin e.g. hybridoma cell lines. The cells may optionally express one or more non-native gene products, e.g. receptors, enzymes, enzyme substrates, prior to or in addition to the fluorescent probe. Preferred cell lines include but are not limited to
15 those of fibroblast origin, e.g. BHK, CHO, BALB, or of endothelial origin, e.g. HUVEC, BAE (bovine artery endothelial), CPAE (cow pulmonary artery endothelial), HLMVEC (human lung microvascular endothelial cells), or of airway epithelial origin, e.g. BEAS-2B, or of pancreatic origin, e.g. RIN, INS-1, MIN6, bTC3, aTC6, bTC6, HIT, or of hematopoietic origin, e.g. primary isolated human monocytes, macrophages, neutrophils,
20 basophils, eosinophils and lymphocyte populations, AML-14, AML-193, HL-60, RBL-1, U937, RAW, JAWS, or of adipocyte origin, e.g. 3T3-L1, human pre-adipocytes, or of neuroendocrine origin, e.g. AtT20, PC12, GH3, muscle origin, e.g. SKMC, A10, C2C12, renal origin, e.g. HEK 293, LLC-PK1, or of neuronal origin, e.g. SK-N-DZ, SK-N-BE(2), HCN-1A, NT2/D1.

25

The term "hybrid polypeptide" is intended to indicate a polypeptide which is a fusion of at least a portion of each of two proteins, in this case at least a portion of the green fluorescent protein, and at least a portion of a catalytic and/or regulatory domain of a protein kinase. Furthermore a hybrid polypeptide is intended to indicate a fusion
30 polypeptide comprising a GFP or at least a portion of the green fluorescent protein that contains a functional fluorophore, and at least a portion of a biologically active polypeptide as defined herein provided that said fusion is not the Glucocorticoid Receptor-GFP disclosed by Carey, KL et al. and Guiliano, KA et al., respectively. Thus, GFP may be N- or C-terminally tagged to a biologically active polypeptide, optionally via
35 a linker portion or linker peptide consisting of a sequence of one or more amino acids.

The hybrid polypeptide or fusion polypeptide may act as a fluorescent probe in mechanically intact or permeabilised living cells carrying a DNA sequence encoding the hybrid polypeptide under conditions permitting expression of said hybrid polypeptide.

The term hybrid polypeptide or fusion polypeptide is intended also to include the term
5 "fluorescent probe", where the latter is used to indicate a fluorescent fusion polypeptide comprising a GFP or any functional part thereof which is N- or C-terminally fused to a biologically active polypeptide as defined herein, optionally via a peptide linker consisting of one or more amino acid residues, where the size of the linker peptide in itself is not critical as long as the desired functionality of the fluorescent probe is maintained. A
10 fluorescent probe according to the invention is expressed in a cell and basically mimics the physiological behaviour of the biologically active polypeptide moiety of the fusion polypeptide.

The term "kinase" is intended to indicate an enzyme that is capable of phosphorylating a
15 cellular component.

The term "protein kinase" is intended to indicate an enzyme that is capable of phosphorylating serine and/or threonine and/or tyrosine in peptides and/or proteins.

20 The term "phosphatase" is intended to indicate an enzyme that is capable of dephosphorylating phosphoserine and/or phosphothreonine and/or phosphotyrosine in peptides and/or proteins.

The term "cyclic nucleotide phosphodiesterase" is intended to indicate an enzyme that is
25 capable of inactivating the second messengers cAMP and cGMP by hydrolysis of their 3'-ester bond.

In the present context, the term "biologically active polypeptide" is intended to indicate a polypeptide affecting intracellular processes upon activation, such as an enzyme which
30 is active in intracellular processes or a portion thereof comprising a desired amino acid sequence which has a biological function or exerts a biological effect in a cellular system. In the polypeptide one or several amino acids may have been deleted, inserted and/or replaced to alter its biological function, e.g. by rendering a catalytic site inactive or by disrupting the targeting sequence. In another embodiment, one or several amino acids
35 may have been deleted, inserted and/or replaced without altering the biological function

of the polypeptide, that is, it remains biologically equivalent. Preferably, the biologically active polypeptide is selected from the group consisting of proteins taking part in an intracellular signalling pathway, such as enzymes involved in the intracellular phosphorylation and dephosphorylation processes including kinases, protein kinases
5 and phosphorylases as defined herein, but also proteins making up the cytoskeleton play important roles in intracellular signal transduction and are therefore included in the meaning of "biologically active polypeptide" herein. More preferably, the biologically active polypeptide is a protein which according to its state as activated or non-activated changes localisation within the cell, preferably as an intermediary component in a signal
10 transduction pathway. Included in this preferred group of biologically active polypeptides are cAMP dependent protein kinases, 'inhibitor of NF-kappaB' kinases, and cyclic nucleotide phosphodiesterases.

The term "a substance" is intended to indicate any sample which has a biological
15 function or exerts a biological effect in a cellular system. The sample may be a sample of a biological material such as a sample of a body fluid including blood, plasma, saliva, milk, urine, or a microbial or plant extract, an environmental sample containing pollutants including heavy metals or toxins, or it may be a sample containing a compound or mixture of compounds prepared by organic synthesis or genetic techniques.

20

The phrase "any change in fluorescence" means any change in absorption properties, such as wavelength and intensity, or any change in spectral properties of the emitted light, such as a change of wavelength, fluorescence lifetime, intensity or polarisation, or any change in the intracellular localisation of the fluorophore. It may thus be localised to
25 a specific cellular component (e.g. organelle, membrane, cytoskeleton, molecular structure) or it may be evenly distributed throughout the cell or parts of the cell.

The term "organism" as used herein indicates any unicellular or multicellular organism preferably originating from the animal kingdom including protozoans, but also organisms
30 that are members of the plant kingdoms, such as algae, fungi, bryophytes, and vascular plants are included in this definition.

The term "nucleic acid" is intended to indicate any type of poly- or oligonucleic acid sequence, such as a DNA sequence, a cDNA sequence, or an RNA sequence.

The term "biologically equivalent" as it relates to proteins is intended to mean that a first protein is equivalent to a second protein if the cellular functions of the two proteins may substitute for each other, e.g. if the two proteins are closely related isoforms encoded by different genes, if they are splicing variants, or allelic variants derived from the same
5 gene, if they perform identical cellular functions in different cell types, or in different species. The term "biologically equivalent" as it relates to DNA is intended to mean that a first DNA sequence encoding a polypeptide is equivalent to a second DNA sequence encoding a polypeptide if the functional proteins encoded by the two genes are biologically equivalent.

10

The term "fixed cells" is used to mean cells treated with a cytological fixative such as glutaraldehyde or formaldehyde, treatments which serve to chemically cross-link and stabilize soluble and insoluble proteins within the structure of the cell. Once in this state, such proteins cannot be lost from the structure of the now-dead cell.

15

In the present context a "quantitative fluorescence redistribution assay" is intended to indicate an assay whereby it is possible to observe and quantify the subcellular localisation and possible redistribution of an biologically active polypeptide, or part thereof, genetically or chemically tagged with a luminophore inside an intact living cell or
20 cells or permeabilised living cells. The subcellular location and redistribution may be monitored using fluorescence microscopy or fluorescence imaging microscopy but is preferably monitored using a fluorescence imaging plate reader or a fluorescence plate reader for improved throughput. A more thorough description is given in Appendix A.

25 In the present context a "mortal cell line" is used to indicate animal cells that may grow in vitro, given the right conditions, but that have a definite life span of a number of cell divisions or days, week or months beyond which it is not at present possible to keep them alive.

30 In the present context an "immortalised cell line" is used to indicate cells of animal origin where the normal limitations for cell life and number of cell divisions do not apply. Essentially, such cells can live, grow and divide for an unlimited or very long (years to decades) time.

The term "targeting sequence" is used to indicate the amino-acid sequence of a biologically active polypeptide that contains the actual structure or structures necessary for association of the biologically active polypeptide with its native intracellular binding sites. The term "targeting sequence" is also used to indicate the amino-acid sequence of
5 a protein that contains the actual structure or structures necessary for association of a biologically active polypeptide with the protein.

The term "targeting" is used to indicate the process whereby a spatially distributed protein is directed to the intracellular sites and maintained at the intracellular sites to
10 which it is normally anchored or associated. These anchoring sites are normally assumed to be the intracellular sites where the protein has its optimal function for the cell.

The term "dislocate" and derivatives thereof is used to indicate the process whereby an
15 intracellularly spatially distributed protein is forced to detach from its normal anchoring or association structures in the cells due to intercalation of another, preferably smaller, compound at the site of anchoring or association. This usually means that the optimal function of the protein within the cell is lost or reduced and that a larger portion of the protein molecules are freely mobile within the cytoplasm.

20

In the present context a "screening assay" is intended to mean any measurement protocol, including materials, cells, instruments, chemicals, reagents, detection units, calibration and quantification procedures used to measure a response from mechanically intact or permeabilised living cells relevant to influences on an intracellular pathway.

25

In the present context a "primary screening assay" is used to indicate the first screening assay in a discovery project that is used to select and sort all compounds available to the project according to the quantified effect of the compounds in the assay.

30 In the present context a "counterscreen" is intended to mean a screening assay that is relevant to a phenomenon that is undesirable seen from the point of view of the discovery project.

In the present context a "discovery project" is intended to mean the process whereby
35 general or specific ideas about ways of how to modulate an intracellular signalling

pathway are exploited in order to find new chemical compounds that can be used to modulate the intracellular signalling pathway and thereby treat, reduce or abolish symptoms associated with a condition or a disease that is lethal, degenerative, performance-reducing or just uncomfortable to an animal, preferably a human being. The
5 aim of the discovery project is to produce drug candidates that can be tested as potential drugs in an animal, preferably in human beings. The term "discovery project" also encompasses the actual group of individuals, screening assays, tests, machinery, cells, animals and compounds involved in different aspects of the project.

10 The term "tagging" is used to indicate the process whereby a luminophore is genetically or chemically attached to the protein, or part of the protein, of interest to the discovery project.

The term "primary hit" is used to indicate compounds identified in the primary screening
15 assay as having at least the minimum level of desired effect that has been specified in the discovery project.

The term "primary lead compound" is used to indicate a primary hit that has at least the minimal level of desired potency and specificity predetermined by the discovery project.
20

The term "dose-response relationship" is in the present context intended to mean a clear correlation between the quantified response of cells in a screening assay to application of an influence, such as a compound, and the concentration of the applied influence. The response to the influence may be both an up-regulation and a down-regulation of the
25 quantitated parameter used in the screening assay.

In the present context, the term "potency" is intended to mean the ability of an influence to affect the process under study. The process under study may be, for example a screening assay or a specific physiological or pathophysiological response in an animal.
30

In the present context, the term "selectivity" is intended to mean the difference in potency on the desired process, such as a screening assay, and an undesired process, such as a counterscreen, with the view of the discovery project. An influence or a compound is said to display selectivity if the potency for the desired process is higher than for the
35 undesired process.

In the present context, the term "structure-activity relationship" or "SAR" is intended to mean the situation where a direct relationship exists between a compound and modifications made to the compound and the activity of the compound and the
5 modifications made to the compound in one or more screening assays. The process of building a SAR may be used to direct the chemical construction of new compounds with higher potency and selectivity than the original compound.

The term "drug candidate lead" is used to indicate compounds that may be pursued by a
10 discovery project as potential candidates for the final outcome of the project.

In the present context, the term "efficacy" is intended to mean the ability of a compound to affect the process or condition under study. It is closely related to the term "potency" but is in the present context used when relating to effects of a compound on more
15 complex screening assays than the primary screening assay or counterscreens and when relating to effects of a compound in animals.

In the present context, the term "toxicity" is intended to mean that a compound in some way is toxic to cells, tissues or animals. The toxicity means that the cells, tissues or
20 animals will in some way be harmed if the compound is applied at a sufficient concentration. The effects may ultimately lead to cell, tissue or animal death or a limited life compared to the normal condition.

In the present context, the term "physiology" is intended to mean the normal function of
25 biological and biochemical processes inside cells, between cells and in the whole organism or animal.

In the present context, the term "pathophysiology" is intended to mean deviations from the normal function of biological and biochemical processes inside cells, between cells
30 and in the whole organism or animal that may be part of a condition or disease.

In the present context, the term "pathogenesis" is intended to mean the process, be it genetical, biological, biochemical, chemical or environmental, that ultimately may explain, at least in part, the apparent pathophysiology associated with a condition or
35 disease in an animal.

In the present context, the term "fractionated cells" is intended to mean the outcome of a simple division of initially mechanically intact living cells into two fractions, particulate (the components that can be sedimented by centrifugation at more than 10 000xg and
5 not more than 100 000xg for 10 minutes) and soluble fraction (the soluble components and small membrane fragments that do not sediment), after subjecting the cells to plasma membrane disruption either mechanically with some form of homogeniser or sonicator or osmotically (hypoosmotic shock) or through some kind of permeabilisation of the plasma membrane with detergents, toxins or electroporation.

10

The term "parenteral route of administration" is used to indicate the administration of a drug or compound in solution to an animal, such as a mammal or a human, by injection or infusion of the drug or compound into the bloodstream of the animal via an injection needle inserted into one of the animal's blood vessels, preferably a vein.

15

The term "oral route of administration" is used to indicate the administration of a drug or compound in solution or as a solid to an animal, such as a mammal or a human, by placing the drug or compound in the mouth of the animal so that the animal itself can swallow the drug or compound or have it delivered to the stomach or intestine by
20 intubation. When the drug or compound enters the stomach and intestine it will be taken up over the mucosa into the bloodstream and administered via the blood stream to the tissues and organs where it is to exert its effect, or it will be acting locally in the stomach and intestine.

25 The term "pulmonary route of administration" is used to indicate the administration of a drug or compound as an aerosol with either solid or liquid particles to an animal, such as a mammal or a human, by placing the drug or compound container close to or in contact with the mouth and/or nose of the animal so that the animal itself can inhale the drug or compound aerosol. When the drug or compound enters the peripheral bronchioles and
30 alveoli it will be taken up over the alveolar membrane, either into the bloodstream and administered via the blood stream to the tissues and organs where it is to exert its effect or it will act locally in the lungs on lung, vessel and muscle cells as well as any other cell type present there.

The term "cutaneous route of administration" is used to indicate the administration of a drug or compound in solution or as a solid to an animal, such as a mammal or a human, by placing the drug or compound on the skin of the animal. The drug can then enter the blood vessels under the skin as it is permeating the skin and thereby be taken up into the bloodstream and administered via the blood stream to the tissues and organs where it is to exert its effect. It may also exert an effect locally on the site of application on the skin.

The term "rectal route of administration" is used to indicate the administration of a drug or compound in solution or as a solid to an animal, such as a mammal or a human, by placing the drug or compound in the rectal cavity of the animal. When the drug or compound enters the rectum and parts of the large intestine it will be taken up over the mucosa into the bloodstream and administered via the blood stream to the tissues and organs where it is to exert its effect, or it will act locally in the rectum and parts of the large intestine.

Several IKKs and very many phosphodiesterases (PDE:s) are known. They are grouped in families according to functional criteria. Within each family there may be several members - isoforms- encoded by different genes. Each isoform may give rise to several splice variants. This hierarchy is evidenced at the sequence level: isoforms are more similar to each other than to members of other families; splice variants are more similar to each other than to other PDE:s. Each specific PDE thus contains sequences that are unique to itself, as well as sequences that are shared between isoforms and/or families. When setting up a program to identify pharmacological agents that affect the intracellular distribution of a target IKK or PDE, it is first necessary to choose the target from the IKKs and PDE:s known. This may be done according to various criteria. A first criterion is that it is imperative that the target IKK or PDE be present in the tissue or cell type(s) where the pharmacological agent is to exert its effect. A second criterion is that it is desirable that either the target or a specific anchoring/targeting site not be present in tissues or cell types where no pharmacological effects are desired.

Establishing the expression patterns of IKKs and PDE:s in relation to tissues and cell types is best done using the methods of detection of mRNA, e.g. Northern analysis, which is a well established procedure. Briefly, mRNA isolated from a given source is probed with a labelled nucleotide, whose sequence is complementary to the mRNA or a region in a mRNA of interest. The assay allows the investigator to determine the

stringency of the probing, i.e. to correlate the resulting signal(s) with sequence similarities.

As a first step, the nucleotide sequences of IKKs or PDE:s are compiled and inspected to identify regions that are unique to specific IKKs or PDE:s as well as regions that are
5 shared among several, many, or all IKKs or PDE:s. Nucleotide sequences may be found in a depository of genetic information, e.g. GenBank, which is a well known resource.

The inspection of the sequences may be aided by using computer programs that were developed to align several or many sequences, and in so doing highlighting regions of similarity or lack of the same. Many of these are presented and explained in great detail
10 in e.g. Sequence Data Analysis Guidebook /edited by S.R.Swindell, Methods in Molecular Biology vol. 70 (1997), from Humana Press Inc. Totowa, New Jersey.

When sequences have been identified that are unique to an IKK, or a PDE, or respectively shared by several or many IKKs or PDE:s, oligonucleotide probes based on these sequences may be designed and synthesized. The use of such probes to detect
15 mRNA is well established in the research community, see e.g. Basic DNA and RNA Protocols/edited by A.J.Harwood, Methods in Molecular Biology vol. 58 (1996), from Humana Press Inc. Totowa, New Jersey. E.g. Life Technologies offer to synthesize specified oligonucleotides.

20 In addition to oligonucleotide probes, mRNA extracted from the tissues and cell types of interest is required, preferably in a form ready to use in Northern analysis. Several companies offer such material, e.g. Invitrogen and Clontech. Briefly, they provide RNA extracted from a great many human and non-human tissues or cell types immobilized on membranes, as an array or size-fractionated.

25 In a next step, a detectable label needs to be attached to the oligonucleotide probe(s). The label is traditionally in the form of a radioactive isotope, but may to advantage be a chemiluminescent reagent or a fluorescent agent. See e.g. DNA Probes by Keller and Manak (1993), from Macmillan Publishers. Several companies offer reagents to label nucleotide probes, e.g. Ambion (Austin, Texas) and Molecular Probes (Eugene, Oregon).

30 The actual probing procedure involves contacting the immobilized mRNA (s) with the probe(s), washing away unbound probe(s) and detecting the signal(s) from the probe(s) that bound under the conditions tested, a positive signal indicating that the target(s) of the probe(s) was present in the sample(s) subjected to the test. In its simplest form, the test is "one-to-one", i.e. each sample of mRNA is exposed to each probe. However, it
35 may be advantageous to exploit the sequence hierarchy of the IKKs or PDE:s, by first

probing arrays of mRNA from multiple sources with family-specific probes, then examining first positives with isotype-specific probes, and then examining the secondary positives in detail with very specific probes. One could also multiplex the probing by adding different distinguishable fluorescent labels to the probes, thus obtaining

5 information from several probes in one experiment.

The outcome of the analysis is information regarding the expression pattern(s) of IKKs and PDE:s.

Based on their expression pattern(s) specific IKKs and/or PDE:s are then selected for further study, and genetic probes are constructed.

10

In general, a genetic probe, i.e. a "GeneX"-GFP fusion or a GFP-"GeneX" fusion, is constructed using PCR with "GeneX"-specific primers followed by a cloning step to fuse "GeneX" in frame with GFP. The fusion may contain a short vector derived sequence between "GeneX" and GFP (e.g. part of a multiple cloning site region in the plasmid)

15 resulting in a peptide linker between "GeneX" and GFP in the resulting fusion protein.

The fusion may be made using ploymerase chain reaction techniques, which are common laboratory procedures, see e.g. PCR Protocols/edited by B.A.White, Methods in Molecular Biology vol. 15 (1993), from Humana Press Inc. Totowa, New Jersey.

20 In more detail, the steps involved include:

- Design of gene-specific primers. Inspection of the sequence of the gene allows design of gene-specific primers to be used in a PCR reaction. Typically, the top-strand primer encompasses the ATG start codon of the gene and the following ca. 20 nucleotides, while the bottom-strand primer encompasses the stop codon and the ca. 20
- 25 preceding nucleotides, if the gene is to be fused behind GFP, i.e. a GFP-"GeneX" fusion. If the gene is to be fused in front of GFP, i.e. a "GeneX"-GFP fusion, a stop codon must be avoided. Optionally, the full length sequence of GeneX may not be used in the fusion, but merely the part which localizes and redistributes like GeneX in response to a signal.

30 In addition to gene-specific sequences, the primers contain at least one recognition sequence for a restriction enzyme, to allow subsequent cloning of the PCR product. The sites are chosen so that they are unique in the PCR product and compatible with sites in the cloning vector. Furthermore, it may be necessary to include an exact number of nucleotides between the restriction enzyme site and the gene-specific

35 sequence in order to establish the correct reading frame of the fusion gene and/or a

translation initiation consensus sequence. Lastly, the primers always contain a few nucleotides in front of the restriction enzyme site to allow efficient digestion with the enzyme.

- Identifying a source of the gene to be amplified. In order for a PCR reaction to produce a product with gene-specific primers, the gene-sequence must initially be present in the reaction, e.g. in the form of cDNA. The results of the extensive expression analysis performed previously will provide clear information regarding what tissue(s) are useful as source material. cDNA libraries from a great variety of tissues or cell types from various species are commercially available, e.g. from Clontech (Palo Alto), Stratagene (La Jolla) and Invitrogen (San Diego). Many genes are also available in cloned form from The American Type Tissue Collection (Virginia).
- Optimizing the PCR reaction. Several factors are known to influence the efficiency and specificity of a PCR reaction, including the annealing temperature of the primers, the concentration of ions, notably Mg^{2+} and K^{+} , present in the reaction, as well as pH of the reaction. If the result of a PCR reaction is deemed unsatisfactory, it might be because the parameters mentioned above are not optimal. Various annealing temperatures should be tested, e.g. in a PCR machine with a built-in temperature gradient, available from e.g. Stratagene (La Jolla), and/or various buffer compositions should be tried, e.g. the OptiPrime buffer system from Stratagene (La Jolla).
- Cloning the PCR product. The vector into which the amplified gene product will be cloned and fused with GFP will already have been taken into consideration when the primers were designed. When choosing a vector, one should at least consider in which cell types the probe subsequently will be expressed, so that the promoter controlling expression of the probe is compatible with the cells. Most expression vectors also contain one or more selective markers, e.g. conferring resistance to a drug, which is a useful feature when one wants to make stable transfectants. The selective marker should also be compatible with the cells to be used.

The actual cloning of the PCR product should present no difficulty for the person skilled in the art as it typically will be a one-step cloning of a fragment digested with two different restriction enzymes into a vector digested with the same two enzymes. If the cloning proves to be problematic, it may be because the restriction enzymes did not work well with the PCR fragment. In this case one could add longer extensions to the end of the primers to overcome a possible difficulty of digestion close to a fragment end, or one could introduce an intermediate cloning step not based on restriction enzyme digestion.

Several companies offer systems for this approach, e.g. Invitrogen (San Diego) and Clontech (Palo Alto).

Once the gene has been cloned and, in the process, fused with the GFP gene, the resulting product, usually a plasmid, should be carefully checked to make sure it is as
5 expected. The most exact test would be to obtain the nucleotide sequence of the fusion-gene.

Once a DNA construct for a probe has been generated, its functionality and usefulness may be tested by subjecting it to the following tests:

- 10 - Transfecting it into cells capable of expressing the probe. The fluorescence of the cell is inspected soon after, typically the next day. At this point, two features of cellular fluorescence are noted:
 - The intensity should usually be at least as strong as that of unfused GFP in the cells. If it is not, the sequence or quality of the probe-DNA might be faulty, and should be
15 carefully checked.
 - The sub-cellular localization is an indication of whether the probe is likely to perform well.

If it localizes as expected for the gene in question, e.g. is excluded from the nucleus, it can immediately go on to a functional test. If the probe is not localized soon after the
20 transfection procedure, it may be because of overexpression at this point in time, as the cell typically will have taken of very many copies of the plasmid, and localization will occur in time, e.g. within a few weeks, as plasmid copy number and expression level decreases. If localization does not occur after prolonged time, it may be because the fusion to GFP has destroyed a localization function, e.g. masked a protein sequence
25 essential for interaction with its normal cellular anchor-protein. In this case the opposite fusion might work, e.g. if GeneX-GFP does not work, GFP-GeneX might, as two different parts of GeneX will be affected by the proximity to GFP. If this does not work, the proximity of GFP at either end might be a problem, and it could be attempted to increase the distance by incorporating a longer linker between GeneX and GFP in the DNA
30 construct.

If there is no prior knowledge of localization, and no localization is observed, it may be because the probe should not be localized at this point, because such is the nature of the protein fused to GFP. It should then be subjected to a functional test.

In a functional test, the cells expressing the probe are treated with at least one compound known to perturb, usually by activating, the signalling pathway on which the probe is expected to report by redistributing itself within the cell.

If the redistribution is as expected, e.g. if prior knowledge tell that it should translocate
5 from location X to location Y, it has passed the first critical test. In this case it can go on to further characterization and quantification of the response.

If it does not perform as expected, it may be because the cell lacks at least one component of the signalling pathway, e.g. a cell surface receptor, or there is species incompatibility, e.g. if the probe is modelled on sequence information of a human
10 geneproduct, and the cell is of hamster origin. In both instances one should identify other cell types for the testing process where these potential problems would not apply.

If there is no prior knowledge about the pattern of redistribution, the analysis of the redistribution will have to be done in greater depth to identify what the essential and indicative features are, and when this is clear, it can go on to further characterization and
15 quantification of the response.

If no feature of redistribution can be identified, the problem might be as mentioned above, and the probe should be retested under more optimal cellular conditions.

Libraries for cloning of cDNA libraries in the present discovery plan are naturally related
20 to the target tissues of the projects. For ultimately finding lead compounds useful in the treatment of asthma the cloning libraries should preferably be obtained from one or more of the following tissue or cells types: Bronchial smooth muscle, Lung microvascular endothelial cells, eosinophil granulocytes, Th1 or 2 lymphocytes and alveolar macrophages.

25 For ultimately finding lead compounds useful in the treatment of chronic inflammatory diseases the cloning libraries should preferably be obtained from one or more of the following tissue or cell types: Th1 or 2 lymphocytes, T-lymphocytes, B-lymphocytes, Monocytes, Eosinophil granulocytes, Neutrophil granulocytes, Basophil granulocytes, Tissue specific macrophages (such as the liver Kupffer cells and skin Langerhans cells),
30 microvascular endothelial cells, vascular endothelial cells, antigen presenting cells, joint connective and synovial cells. For ultimately finding lead compounds useful in the treatment of depression the cloning libraries should preferably be obtained from one or more of the various tissue regions of the brain containing noradrenergic neurons. For ultimately finding lead compounds useful in the treatment of jet lag or circadian clock

resetting the cloning libraries should preferably be obtained from one or more of the various tissues of the brain such as the pineal gland, hypothalamus and substantia nigra. For ultimately finding lead compounds useful in the treatment of hyper- and hypotension and erectile dysfunction the cloning libraries should preferably be obtained from one or
5 more of the following tissue or cell types: vascular smooth muscle, vascular smooth muscle from resistance vessels on the arterial side of the vascular system, vascular smooth muscle from capacitance vessels on the venous side of the vascular system, vascular smooth muscle cells from small arteries, arterioles, venules or veins, smooth vascular cells lines such as T/G HA-VSMCA10 and A7r5.

10

The cells should always be of animal origin, most likely of mammalian origin and preferably of human origin. The cells could be derived from normal tissue or from tissue of an individual animal having a disease or condition of interest for the project. The cells may also be a mortal or immortalised cell line where the initial cell clone has been
15 derived from a tissue or cell type as described above. Depending on the discovery project the cells of interest for screening assays will vary but may be chosen from the above mentioned categories.

Once a genetic construct containing the protein of interest and the luminophore, from
20 here on referred to as "the original fluorescent probe", has been transfected into a relevant cell type, as described above under 'preferred cell types for cloning libraries' the cells are monitored for the appearance of spatially distributed or randomly distributed intracellular fluorescence. Based on prior knowledge regarding the distribution of the actual protein different patterns can be expected. If for example previous studies have
25 found the protein associated only with the particulate fraction of fractionated cells, it can be expected to find a spatial distribution of the original fluorescent probe to the plasma membrane, internal membrane/organelle structures or structural cytoplasmic elements such as microtubules and microfilaments. If on the other hand previous studies report that the protein has been found mostly in the soluble fraction of fractionated cells one
30 can expect to find a homogenous or nonhomogenous distribution of the original fluorescent probe throughout the cytoplasm and perhaps also in the nucleus. For proteins where previous studies have found a mixed localisation to both the particulate and soluble fraction of fractionated cells any mixture in the two distribution patterns mentioned above for the original fluorescent probe can be expected. For proteins where
35 no prior knowledge is at hand a simple cell fractionation and Western Blotting can be

made, one can use immunohistochemistry of fixed cells of relevance or one can decide to rely on the distribution observed for the original fluorescent probe. At this stage of the project, a normal distribution pattern of the original fluorescent probe may be established after such studies as outlined above. The effects of physiologically important and
5 relevant cellular activation on the distributed pattern of the original fluorescent probe is also established. It will also become evident if the pattern of distribution changes, i.e. if a redistribution of the original fluorescent probe occurs as a consequence of applying a physiologically important and relevant influence.

10

The strategy described herein is used to search for chemical entities which can interfere with the protein-protein interactions that occur amongst biologically active polypeptides and their anchoring/regulating partners, and thereby interfere with the effectiveness of a biologically active polypeptide's action within its cellular environment. The strategy will
15 have different effects, and require slightly different discovery methods depending on the nature of the interaction. The possibilities are as follows:

- 1) A biologically active polypeptide is permanently located at its targeting point, and either remains permanently active there, or its activity is modulated in some way by post-
20 translational modification such as phosphorylation or by binding of modulators to non-catalytic regulatory sites. Dislocation from the targeting site will remove the biologically active polypeptide from a localised site of action, and may also lead to inactivation of its inherent catalytic activity.
- 2) A biologically active polypeptide is permanently located at its targeting point, and
25 remains inactive there until its activity is modulated in some way by post-translational modification, such as phosphorylation or by binding of modulators to non-catalytic regulatory sites. Dislocation from the targeting site will remove the biologically active polypeptide from a localised site of action, and may also lead to activation of its inherent catalytic activity, albeit away from its original anchoring site.
- 30 3) A biologically active polypeptide is inactive in its unattached or untargeted form, and when activated (as described in "1" above), or partially activated, it redistributes within the cell and becomes attached to its targeting site, its activity being restricted to the anchoring site and possibly enhanced by interaction with the anchoring protein or some associated factor, or at some later time inhibited by the anchoring protein or an
35 associated regulatory factor. Any agent which prevents association of the biologically active polypeptide with its anchoring or targeting site will prevent it from locating to the

preferred site of action, and may also prevent the biologically active polypeptide from becoming fully activated by the appropriate stimulus whilst in the untargeted state.

4) A biologically active polypeptide is active in its unattached or untargeted form, and when inactivated (as described in "1" above), or partially inactivated, it redistributes
5 within the cell and becomes attached to its targeting site, whereby its activity is inhibited by interaction with the anchoring protein or an associated regulatory factor. Subsequent stimuli may then activate and release the biologically active polypeptide. Any agent which prevents association of the biologically active polypeptide with its anchoring or targeting site will prevent it from relocating to the anchoring position, and may also
10 prevent the biologically active polypeptide from ever being inactivated. In addition, if the biologically active polypeptide cannot target to its anchoring site, it may not be possible subsequently to activate the biologically active polypeptide in the appropriate way in the untargeted state.

15 When a specific subcellular distribution of a GFP-based IKK or PDE probe has been identified, it may be advantageous to narrow down which part of the IKK or PDE is responsible for this effect. The advantage is twofold: It may suggest the design of peptide leads, and it may eventually aid in defining the binding partner. Knowledge of both partners involved in specific binding may aid in the selection of compound libraries
20 to screen for inhibition of the specific binding.

To identify the region of the IKK or PDE involved in specific binding, one may make GFP-based fusions with progressively shorter parts of the IKK or PDE, and examine the cellular distribution of these constructs. If there is prior knowledge of functional domains,
25 one may start with the domain believed to confer specific binding to a subcellular structure. The generation of constructs to test may consist of selecting a particular part of the IKK or PDE to fuse to GFP, or it may involve the generation of in-frame deletions in the IKK or PDE part of the fusion. Both approaches have been widely used in molecular genetic studies.

30 When a region has been identified that appears responsible for conferring a specific subcellular distribution upon an IKK or a PDE, the amino acid residues most important for this trait may be identified by a more detailed analysis, e.g. substituting them one by one with e.g. an alanine residue, a so called Ala-scan, which also has been used extensively in molecular genetic studies.

35 To identify the identity of the cellular protein partaking in the specific distribution of the IKK or PDE, one may exploit the knowledge about the region of the IKK or PDE

responsible for the subcellular distribution; for example, one may use the region of the IKK or PDE as bait in a genetic two hybrid screen to pull out its binding partner. Several companies offer two hybrid systems, e.g. Life Technologies.

- 5 The knowledge about the normal distribution of the original fluorescent probe is used to establish which part or which parts of the terminal (or entire) amino-acid sequence that is important for the attachment of this fluorescent probe to subcellular structures, giving it its specific spatially distributed pattern in the cell or cells, when such a pattern has been established as the normal distribution of this fluorescent probe. This may be
- 10 accomplished by creating new fluorescent probes where a systematic deletion of short N- or C-terminal or internal sequences (number of DNA bases) of the original fluorescent probe are made. These new shorter variants of the of the original fluorescent probe construct are transfected into the cells of interest and then the cells are examined for spatial distribution of the new fluorescent probes as described above for the original
- 15 fluorescent probe. In those cells where the new fluorescent probe distribution pattern is different from the original fluorescent probe distribution pattern it is evident that part of the, or the entire, targeting sequence has been deleted. The DNA- or amino-acid sequence of the missing part therefore contains the structural information necessary for association of the original fluorescent probe with its intracellular binding sites.

20

- Peptides for inhibition of the established normal distribution of the original fluorescent probe are designed according to the hypothesis, that the deduced targeting sequence, or sequences, in the original fluorescent probe amino-acid sequence are the important sequences for the actual spatial distribution of the original fluorescent probe in intact
- 25 living cells, is tested. This is done by producing peptides of identical amino-acid sequence as the deduced targeting sequence or parts thereof and introducing them into the cytoplasm, either by microinjection or transient or permanent permeabilisation, of cells containing the original fluorescent probe and thereafter monitoring the spatial distribution of the original fluorescent probe in the cells. If the deduced targeting
- 30 sequence or sequences are of importance for the actual spatial distribution of the original fluorescent probe in intact living cells, the introduced peptides will self-associate with the anchoring sites for the original fluorescent probe and thereby disrupt the normal distribution of the original fluorescent probe. In order to have this effect, the introduction of the peptides should change the original distribution pattern so that a decrease in
- 35 fluorescence of 10% or more, compared to the pattern before their introduction, can be

detected. This is done by observing the same cells before and after administration of the peptides. When peptides that fulfil this criterion have been found they are called 'peptide leads' and will hereafter be referred to using this expression. These peptide leads can now be used as a basis for the design of organic molecules that can be used eventually
5 to disrupt the spatial distribution of the original fluorescent probe but also as control compounds in screening assays.

PS473 and derivatives thereof show a discrete intracellular localisation that allow establishment of assay systems valuable in the screening for compounds that modulate
10 targeting of said probes. IKK β interacts with multiple components of the I κ B complex. Construction of the described assay systems has allowed us to screen for compounds that interact with specific or multiple targeting sites. This approach allow for development of compounds that through modulation of one (or several) of multiple targeting sites of IKK β (or other IKKs) will provoke either a partial or a complete inhibition
15 of the NF-kappaB activation. In addition cell specific anchoring will allow design of compounds that only affect defined cell types.

In parallel to the above mentioned step wherein peptide leads are defined, the distribution pattern found for the original fluorescent probe is compared to the naturally
20 occurring spatial distribution of the protein on which the original fluorescent probe is based. This may be accomplished by observing fixed primary cells separated from or still within the tissue of interest and fixed cells that contain the original fluorescent probe. Thereafter the protein is stained using ordinary immunocytochemical or immunohistochemical methods and the spatial distribution revealed by this staining
25 procedure is compared to the spatial distribution of the original fluorescent probe. It is desirable, but not required, that a high degree of correlation between the two patterns obtained in this step can be observed.

Establishment of a primary screening assay is normally done by making use of the cells
30 of interest containing the original fluorescent probe as the basis for a screening assay. Depending on the knowledge acquired about the behaviour of the original fluorescent probe when subjecting the cells to physiologically relevant influences the assay procedure can be chosen: 1. If the fluorescent probe normally is targeted to specific sites and stays associated with these sites during stimulation of the intracellular pathway, the
35 assay should preferably be designed to detect dislocation of the original fluorescent

probe from the targeting sites in mechanically intact or permeabilised living cells. This is an assay where the dislocation can be detected within minutes after application of an influence and the time frame for the detection and time for exposing the cells to an influence should be chosen to match this. 2. If the desire is to disrupt the actual targeting event rather than dislocate already targeted fluorescent probe the influence may need hours to produce a detectable response. The actual measurement, still of a change in the fluorescence or luminescence distribution pattern compared to the normal distribution pattern for the original fluorescent probe, may be made at two time points; before and after the influence has exerted any effect it may have. This is an assay where the effect of an influence may require several hours to produce a detectable response and the time frame for the detection and time for exposing the cells to an influence should be chosen to match this. 3. If the fluorescent probe normally redistributes between two intracellular sites upon activation of the intracellular pathway one may either want to disrupt the initial targeting or dislocate the original fluorescent probe from its initial or resting anchoring site. In this case procedure no. 1 above may be used. If the desire instead is to inhibit the association of the original fluorescent probe with the site it redistributes to during activation of the intracellular pathway the targeting sequence of this site should be in focus for the lead peptide generation. This is an assay where the redistribution may be detected within minutes after application of an influence and the time frame for the detection and time for exposing the cells to an influence should be chosen to match this. Furthermore, any influence applied to inhibit the targeting of the original fluorescent probe upon its redistribution may need to be added to the cells before activation of the intracellular pathway.

While the original fluorescent probe and peptide leads will be used in the actual primary screening assay, it is also desirable to have a counterscreen or counterscreens directed at protein isoforms that one does not wish to affect. In order to accomplish this, constructs are made for new fluorescent probes encoding the protein isoforms tagged with GFP. These constructs are subsequently transfected into the cells of interest. When the new fluorescent probes are expressed in the cells, some of the cells are chosen as the basis for new cell lines that can be used in the counterscreen or counterscreens.

Suitable probes for this purpose comprise DNA constructs encoding fusion polypeptides comprising forms of IKK α , IKK β , IKK γ or NIK and GFP; PDE1, PDE2, PDE3, PDE4, PDE5, PDE6, PDE7, PDE8, PDE9 or PDE10 and GFP; PKA catalytic subunit and GFP.

In a preferred embodiment the DNA constructs will encode fusion polypeptides comprising isoforms of IKK β , PDE 4, mPDE5, PKA catalytic subunit and GFP.

- 5 In a much preferred embodiment the DNA construct is selected from table 1.

Table 1 list of the fusion constructs of the invention by the names used herein as well as by reference to relevant SEQ ID NOs of sequences of DNA encoding the construct and full amino acid sequences

Fusion construct	DNA sequence SEQ ID NO:	Protein Sequence SEQ ID NO:
PDE 4D3 - EGFP	1	2
PDE 4D4 - EGFP	3	4
PDE 4D5 - EGFP	5	6
PDE 5 - EGFP	7	8
IKK β - EGFP	9	10
NF-KappaB - EGFP	11	12
EGFP - IKK β	13	14
EGFP - IKK β L2	15	16

10

The cell lines established for the primary screen and the counterscreen, or counterscreens, are used to establish peptide leads that more specifically dislocate the desired isoform of the protein of interest compared to other isoforms of the same protein. The peptide leads are introduced into the cells as described above and the changes in

- 15 spatial distribution of the original and counterscreen fluorescent probes are quantified and dose-response relationships are established for each lead peptide. Thereafter the dose-response relationships are compared. A peptide lead is considered specific for the original fluorescent probe if the dose of the peptide required to dislocate at least 10% of the fluorescent probes in the counterscreen or counterscreens are at least two times
- 20 higher than the dose required to dislocate 10% of the original fluorescent probe. The lead peptides with the biggest dose difference when comparing the primary and the counterscreen dose-response relationships are chosen as the basis for the next step in the discovery project.

- 25 In one embodiment the primary screening assay and counterscreen or counterscreens are used to define specificity of the peptide leads by using a procedure that compares their ability to cause a dislocation, disruption of targeting or inhibition of redistribution of the original fluorescent probe in the primary screening assay to their ability to cause a

dislocation, disruption of targeting or inhibition of redistribution of the new fluorescent probes in the counterscreen or counterscreens.

In a preferred embodiment the dose of a peptide lead required to cause a quantified
5 dislocation, disruption of targeting or inhibition of redistribution of the original fluorescent probe of at least 10% in the primary screening assay is 50% or less of the dose required to cause a quantified dislocation, disruption of targeting or inhibition of redistribution of the new fluorescent probes of at least 10% in the counterscreen or counterscreens.

The invention provides for a specificity index which may be constructed describing a
10 numerical relationship, with the primary screening assay result first, of the dose required to produce half-maximal effect in the primary assay compared to the dose required to produce half-maximal effect in the counterscreen or counterscreens.

In one embodiment the peptide leads chosen for further use in the discovery project have a specificity index of 1 to 2.

15 In another embodiment the peptide leads chosen for further use in the discovery project have a specificity index between 1 to 2 and 1 to 10.

In a further embodiment the peptide leads chosen for further use in the discovery project have a specificity index between 1 to 11 and 1 to 100.

In yet a further preferred embodiment the peptide leads chosen for further use in the
20 discovery project have a specificity index better than 1 to 100.

Lead peptides are used to create and select libraries of small organic molecules that can be useful in screening assays to find bioactive substances useful as drugs to treat the condition or disease of interest for the project. In this step the amino-acid sequence
25 information and other structural information about the lead peptide or peptides is used to extract information useful for finding and/or defining and synthesising bioactive organic molecules that can mimic the effect of the lead peptides on the normal spatial distribution pattern of the original fluorescent probe. Such compounds may be useful as drugs to treat the condition or disease of interest for the project. Peptide leads selected by the
30 discovery project are used to design and assemble compound libraries based on the structural and chemical information inherent in the lead peptides using prior chemical knowledge and computational chemistry approaches so that the compounds have a structure that give them the ability to interact with or bind to the targeting sequence of IKK β , PDE 4D X or mPDE5 thereafter testing the compound libraries at a concentration
35 of 10 or 100 micromolar of each compound in the primary screening assay.

When the libraries of compounds have been defined and are at hand it is time to initiate primary screening. In this procedure, cells containing the original fluorescent probe are contacted with the compounds. The compounds are all tested at just one or a few concentrations, typically 10 and 100 micromolar, in a highly parallel fashion using a quantitative fluorescence redistribution assay. Compounds that cause a change in the quantitated response (the response scale defined by the range 0 (no change in redistribution) – 100%) of the assay by more than a predetermined value, typically between 10 and 100%, are considered to be “primary hits”. The primary hits are then further characterised: 1. for potency by establishing a dose-response relationship compared to the lead peptide(s) using the primary screening assay 2. for selectivity by establishing a dose-response relationship in the counterscreen or counterscreens. Primary hits that have low potency, typically when the half-maximal effect of the compound in the primary assay is achieved at a concentration of the compound between 10 and 100 micromolar, may not need testing in the counterscreen or counterscreens since the likelihood that they will be used beyond this step in the discovery project is small. Primary hits that have equal or lower potency in the primary screening assay compared to the counterscreen or counterscreens are regarded as non-selective and the likelihood that they will be used beyond this step in the discovery project is small.

Primary hits that display some degree of selectivity, typically half maximal effect in the primary screening assay at a concentration 50% or less of the concentration that gives half maximal effect in the counterscreen or counterscreens are considered interesting as the basis for further chemical synthesis or construction of new libraries of compounds and will hereafter be referred to as “primary lead compounds”.

Compounds that cause a change in the quantitated response, with a response scale from 0 to 100% based on the absence of a response and the maximal response observed with the peptide leads in the primary screening assay, of the assay by more than a predetermined value are selected and called “primary hits”.

In one embodiment the predetermined value is 10%.

In another embodiment the predetermined value is 50%.

In yet another embodiment the predetermined value is 70%.

In one embodiment the primary hits are further characterised for potency and maximal effect by establishing a dose-response relationship and comparing that to the effects of the lead peptides using the primary screening assay and for selectivity by establishing a dose-response relationship in the counterscreen or counterscreens.

Primary hits may be deselected by the discovery project when they display a half-maximal potency at a dose corresponding to a concentration of more than 10 micromolar or because they display a selectivity index less than 1 to 2.

Primary hits may be selected by the discovery project when they display a half-maximal
5 potency at a dose corresponding to a concentration of 10 micromolar or less or because they display a selectivity index higher than 1 to 2, the compounds hereafter also referred to as "primary lead compounds".

A Structure-Activity Relationship (SAR) is built by iterations of compound library
10 composition and screening to define drug candidate leads. This step is included to further improve the possibilities of finding bioactive compounds with desirable properties for treatment of the diseases or conditions of interest to the project. The primary lead compounds are here used to provide chemical structural information that can be used as the basis for composition or chemical synthesis of new, directed, compound libraries. By
15 systematic chemical modification of part of the structure of one or more primary lead compounds new libraries are assembled. These new libraries of compounds are also investigated using the primary screening assay and counterscreen or counterscreens. Preferably, dose-response relationships are recorded for each chemical modification of the primary lead compound and compared to the primary lead compound itself. Thereby
20 SAR is established. Among the new compounds, the ones that in this step has the best combination of potency and specificity are chosen either as the basis for a new round of compound library synthesis or composition or, as the final step of the SAR building process, as compounds that will be further for actual pharmacological effects in assay systems and animals that are relevant to the underlying physiological and
25 pathophysiological processes of interest to the project. The latter compounds will hereafter be referred to as "drug candidate leads".

In one embodiment drug candidate leads have a half-maximal potency at a dose corresponding to a concentration of less than 1 micromolar and a selectivity index higher than 1 to 2.

30 In one embodiment the drug candidate leads have a half-maximal potency at a dose corresponding to a concentration of less than 1 micromolar and a selectivity index higher than 1 to 10.

In one embodiment the drug candidate leads have a half-maximal potency at a dose corresponding to a concentration of less than 1 micromolar and a selectivity index higher
35 than 1 to 100.

In one embodiment the drug candidate leads have a half-maximal potency at a dose corresponding to a concentration of less than 0,1 micromolar and a selectivity index higher than 1 to 2.

In a preferred embodiment the drug candidate leads have a half-maximal potency at a
5 dose corresponding to a concentration of less than 0,1 micromolar and a selectivity index higher than 1 to 10.

In another preferred embodiment the drug candidate leads have a half-maximal potency at a dose corresponding to a concentration of less than 0,1 micromolar and a selectivity index higher than 1 to 100.

10

Drug candidate leads may be further characterised in tissue based, cell based and biochemical assays to validate *in vitro* their efficacy and toxicity. There are many ways to test efficacy of a drug candidate lead. Preferably, the drug candidate lead is tested in assay systems with high relevance to the underlying physiological and
15 pathophysiological processes involved in the pathogenesis and pathophysiology of the disease or condition of interest to the project. Likewise, the drug candidate leads are tested for toxic effects, preferably testing for genetic effects (influence on the integrity and arrangement of DNA), metabolic effects (influence on cellular metabolic processes) and cytotoxic effects (influence on cell integrity and organelle integrity). There is a high
20 likelihood that drug candidate leads, that do not show appropriate efficacy or that display toxicity will not be used beyond this step in the discovery project because it is expected that such compounds are less suitable as actual drugs to be used in an animal.

In one embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying
25 physiological and patophysiological processes involved in hypotension, inflammatory diseases, and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In another embodiment drug candidate leads chosen by the discovery project are tested
30 *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying physiological and patophysiological processes involved in inflammatory airway diseases, and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In another embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying physiological and patophysiological processes involved in inflammatory joint diseases, and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter
5 the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In another embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying physiological and patophysiological processes involved in inflammatory bowel diseases,
10 and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In another embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying
15 physiological and patophysiological processes involved in autoimmune diseases, and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

20 In a preferred embodiment of the present invention I-kappaB degradation is inhibited by a novel mechanism namely by mis-targeting and/or modulation of the redistribution of specific IKKs. In contrast to previous interventions involving IKK the presented invention does not involve direct inhibition of the IKK enzymatic activity.

25 This completely novel mechanism for inhibition of the overall effect of the IKK complex provides clear advantages as it opens for a higher IKK isoform selectivity and a higher cell specificity of the therapy. In addition cell specific anchoring will allow design of compounds that only affect defined cell types.

30 In one aspect of the invention the substance is an organic compound, the organic compound being a weak acid in that it is a neutral molecule that can reversibly dissociate into an anion (a negatively charged molecule) and a proton (a hydrogen ion). In another aspect, the organic compound is a weak base in that it is a neutral molecule that can form a cation (a positively charged molecule) by combining with a proton. The functional
35 groups of the targeting sequences include functional groups selected from the group

consisting of: methyl-, isopropyl-, isobutyl-, hydroxyl-, thiol-, benzyl-, benzyloyl-, methylindolyl-, methylimidazolyl-, amine-, imine-, carboxyl- and acetamide-groups as parts of amino acids in the targeting sequences.

- 5 In another aspect of the invention the organic compound is a compound having one or more chemical domains capable of interacting with one or more functional groups of the targeting sequence of the native anchoring site of the cyclic nucleotide phosphodiesterase or I-kappaB kinase. In yet another aspect the organic compound is a compound having at least two chemical domains capable of interacting with at least two
- 10 functional groups of the targeting sequence of the native anchoring site for the cyclic nucleotide phosphodiesterase or I-kappaB kinase. In a further aspect the organic compound is a compound having at least three chemical domains capable of interacting with at least three functional groups of the targeting sequence of the native anchoring site for the cyclic nucleotide phosphodiesterase or I-kappaB kinase.

15

- The organic compound is, in one aspect of the invention, a compound having at least two chemical domains capable of interacting with at least two functional groups of the targeting sequence of the cyclic nucleotide phosphodiesterase. In a specific embodiment, the organic compound is a compound having at least three chemical
- 20 domains capable of interacting with at least three functional groups of the targeting sequence of the cyclic nucleotide phosphodiesterase.

- In the next part of the discovery process the drug candidate leads are tested *in vivo* for toxic and unwanted effects in animals such as mice and rats. The drug candidate leads
- 25 are also tested for efficacy in animals that have a disease or condition with high degree of relevance to the disease or condition of interest to the project. The drug candidate leads may also be tested for efficacy in animals which have been treated in a way that make them experience a disease or condition with high degree of relevance to the disease or condition of interest to the project. Drug candidate leads that display efficacy
- 30 in one or more of such animal tests and that does not display any apparent toxicity at a dosage level, preferably 2 –10 times higher than the level that gives satisfactory efficacy are chosen to be the final drug candidates that should be considered for further animal testing and initial testing in humans. These compounds are hereafter referred to as “discovery project leads”.

35

In one embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying physiological and pathophysiological processes involved in depression, and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug

5 candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In another embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying physiological and pathophysiological processes involved in jet-lag, and for toxicity,

10 preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In another embodiment drug candidate leads chosen by the discovery project are tested *in vitro* for efficacy, in assay systems with high degree of relevance to the underlying

15 physiological and patophysiological processes involved in erectile dysfunction, and for toxicity, preferably testing for genetic, metabolic and cytotoxic effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity are chosen to be the candidates that will enter testing in animals.

In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in hypotension, and for toxicity and unwanted side effects, after which the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter

25 further testing in animals and testing in humans.

In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in inflammatory diseases, and for toxicity and unwanted side effects, after which the drug candidate

30 leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to

35 the underlying physiological and pathophysiological processes involved in hypertension,

and for toxicity and unwanted side effects, after which the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

- 5 In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in jet-lag and circadian rhythm resetting, and for toxicity and unwanted side effects, after which the drug candidate leads that display the best efficacy and the least, or no, indications of
- 10 toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

- In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in erectile
- 15 dysfunction, and for toxicity and unwanted side effects, after which the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

- 20 In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in inflammatory airway diseases, and for toxicity and unwanted side effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity
- 25 or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

- In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in inflammatory
- 30 joint diseases, and for toxicity and unwanted side effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

- In one embodiment drug candidate leads chosen by the discovery project are tested for
- 35 efficacy, in healthy animals and animals with a condition with high degree of relevance to

the underlying physiological and pathophysiological processes involved in inflammatory bowel diseases, and for toxicity and unwanted side effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, 5 that will enter further testing in animals and testing in humans.

In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to the underlying physiological and pathophysiological processes involved in autoimmune diseases, and for toxicity and unwanted side effects, whereafter the drug candidate 10 leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

In one embodiment drug candidate leads chosen by the discovery project are tested for efficacy, in healthy animals and animals with a condition with high degree of relevance to 15 the underlying physiological and pathophysiological processes involved in depression, and for toxicity and unwanted side effects, whereafter the drug candidate leads that display the best efficacy and the least, or no, indications of toxicity or unwanted side effects are chosen to be the candidates, called discovery project leads, that will enter further testing in animals and testing in humans.

20

The administration route of any of the compounds of the invention may be of any suitable route which leads to a concentration in the blood corresponding to a therapeutic concentration by the oral route, the parenteral route, the cutaneous route, the nasal route, the rectal route, the vaginal route and the ocular route. It should be clear to a 25 person skilled in the art that the administration route is dependant on the compound in question, particularly, the choice of administration route depends on the physico-chemical properties of the compound together with the age and weight of the patient and on the particular disease and the severity of the same.

The compounds of the invention may be contained in any appropriate amount in a 30 pharmaceutical composition, and are generally contained in an amount of about 1-95% by weight of the total weight of the composition. The composition may be in form of, e.g., tablets, capsules, pills, powders, granulates, suspensions, emulsions, solutions, gels including hydrogels, pastes, ointments, creams, plasters, drenches, delivery devices, suppositories, enemas, injectables, implants, sprays, aerosols and in other suitable form. 35 The pharmaceutical compositions may be formulated according to conventional

pharmaceutical practice, see, e.g., "Remington's Pharmaceutical Sciences" and "Encyclopedia of Pharmaceutical Technology".

- Pharmaceutical compositions according to the present invention may be formulated to release the active compound substantially immediately upon administration or at any
- 5 substantially predetermined time or time period after administration. The latter type of compositions are generally known as controlled release formulations. Controlled release formulations may also be denoted "sustained release", "prolonged release", "programmed release", "time release", "rate-controlled" and/or "targeted release" formulations.
- 10 In the present context every pharmaceutical composition is an actual drug delivery system, since upon administration it presents the active drug substance to the body of the organism.

- The compounds of the invention are preferably administered in an amount of about 0.1-30 mg per kg body weight per day, such as about 0.5-15 mg per kg body weight per day.
- 15 The compound in question may be administered orally in the form of tablets, capsules, elixirs or syrups, or rectally in the form of suppositories. Parenteral administration of the compounds of the invention, is suitably performed in the form of saline solutions of the compounds or with the compound incorporated into liposomes. In cases where the
- 20 compound in itself is not sufficiently soluble to be dissolved, an acid addition salt of a basic compound can be used, or a solubilizer such as ethanol can be applied.

Oral administration. For compositions adapted for oral administration for systemic use, the dosage is normally 1 mg to 1 g per dose administered 1-4 times daily for 1 week, 12 months or even lifelong depending on the disease to be treated.

- 25 Rectal administration. For compositions adapted for rectal a somewhat higher amount of compound is usually preferred, i.e. from approximately 1 mg to 100 mg per kg body weight per day.

- Parenteral administration. For parenteral administration a dose of about 0.1 mg to about 50 mg per kg body weight per day is convenient. For intravenous administration a dose
- 30 of about 0.1 mg to about 20 mg per kg body weight per day. For intraarticular administration a dose of about 0.1 mg to about 20 mg per kg body weight per day is usually preferable. For parenteral administration in general, a solution in an aqueous medium of 0.5-2% or more of the active ingredients may be employed.

- Cutaneous administration. For topical administration on the skin a dose of about 1 mg to
- 35 about 5 g administered 1-10 times daily is usually preferable.

EXAMPLES

Example 1: Probes for detection of PDE4D dislocation.

These are specific PDE4D variants fused to a GFP. Currently 5 PDE4D splice variants are known: PDE4D1, PDE4D2, PDE4D3, PDE4D4 and PDE4D5. These all share C-

5 terminal sequences but differ in their N-termini.

Inspection of the scientific literature indicates that the PDE4D1 and PDE4D2 subtypes are found only in the cytosolic fraction, whereas PDE4D3, PDE4D4 and PDE4D5 subtypes appear to associate with some form of cellular structure(s). Targetting sequences of PDE4Ds are presently believed to be located in their N-terminal domain(s).

10 In accordance with this, PDE4D1 and PDE4D2 have much shorter N-terminal domains than PDE4D3, PDE4D4 and PDE4D5. To best preserve the normal distribution of PDE4Ds, the fusions are made between the C-terminus of the PDE4D species and the N-terminal of the GFP.

To construct PDE4D-GFP fusions, PDE4D sequences are amplified using PCR
15 according to standard protocols with specific top-primers as listed below, and the common bottom-primer listed below. The PCR products are digested with restriction enzymes Hind3 and EcoR1, and ligated into pEGFP-N1 (Clontech, Palo Alto; GenBank Accession number U55762) digested with Hind3 and EcoR1. This produces PDE4D-EGFP fusions under the control of a CMV promoter (SEQ ID NOs: 5 and 6 (PDE4D5-EGFP); SEQ ID NOs: 3 and 4 (PDE4D4-EGFP); SEQ ID NOs: 1 and 2 (PDE4D3-EGFP)).
20

Top primers all include specific sequences following the ATG, a Kozak sequence, and a cloning site (Hind3). The bottom primer includes the common C-terminal sequence minus the stop codon, an EcoR1 cloning site, and an extra nucleotide to preserve the
25 reading frame in EGFP-N1.

Sequences of top-primers:

5'-GTAAGCTTCGAACATGATGCACGTGAATAATTTTCCC-3' ; specific for PDE4D3A and PDE4D3B (GenBank Acc. nos. L20970 & U50159).

30

5'-GTAAGCTTCGAACATGGAGGCAGAGGGCAGCAGC-3'; specific for PDE4D4A (GenBank Acc. no. L20969).

5'-GTAAGCTTCGAACATGGCTCAGCAGACAAGCCCG-3'; specific for PDE4D5A (GenBank Acc. no. AF012073).

Sequence of common bottom-primer:

5 5'-GTGAATTCCCGTCGTGTCAGGAGAAGCATCATCTATG-3'.

The resulting plasmids are transfected into a suitable cell line, e.g. MVLEC. The subcellular distribution of the probes is examined carefully by fluorescence microscopy, both under resting conditions, and upon elevation of cAMP, e.g. by activation of
10 adenylate cyclase with forskolin, which may or may not have an effect on the normal distribution.

Example 2: Probes for detection of PDE5 dislocation:

These are specific PDE5 variants fused to a GFP. Currently only one main human variant is known (GenBank Acc.nos. AJ004865 and D89094).

15 Inspection of the scientific literature indicates that the catalytic domain is contained in the C-terminal part of the protein, so potential targeting sequences of PDE5 may be located in the N-terminal part. To best preserve the normal distribution of PDE5, the first fusion is made between the C-terminus of the PDE5 species and the N-terminal of the GFP.

20 To construct the PDE5-GFP fusions, PDE5 sequences are amplified using PCR according to standard protocols with the specific primers listed below. The PCR product is digested with restriction enzymes EcoR1 and Acc65I, and ligated into pEGFP-N1 (Clontech, Palo Alto; GenBank Accession number U55762) digested with EcoR1 and Acc65I. This produces a PDE5-EGFP fusion under the control of a CMV promoter (SEQ
25 ID NOs: 7 and 8).

The top primer includes specific sequences following the ATG, a Kozak sequence, and a cloning site (EcoR1). The bottom primer includes specific C-terminal sequences minus the stop codon, an Acc65I cloning site, and two extra nucleotides to preserve the reading frame in EGFP-N1.

30

PDE5-top :

5'-GTGAATTCAACCATGGAGCGGGCC-3'

PDE5-bottom:

35 5'-GTGGTACCCAGTTCGCTTGGCC

The resulting plasmids are transfected into a suitable cell line, e.g. MVLEC. The subcellular distribution of the probes is examined carefully by fluorescence microscopy, both under resting conditions, and upon elevation of cGMP, e.g. by activation of cyclase
5 with NO or nitroprusside, which may or may not have an effect on the normal distribution.

EXAMPLE 3: Probes for detection of IKK redistribution.

Modulation of IKK β redistribution by mis-targeting provoke an inhibition of cytokine-induced NF-kappaB activation. In the present example it is shown that specific mis-targeting of IKK β inhibits cytokine-induced NF-kappaB activation. Dislocation of
10 endogenous IKK β from its anchoring sites is achieved by expression of a C-terminal part of IKK β (PS473). The PS473 probe, which is a GFP fusion, allows a simultaneous monitoring of its localisation and redistribution.

Expression of the PS473 probe has a clear inhibitory activity on cytokine-induced
15 activation of NF-kappaB. For the first time we hereby show that dislocating IKK β , without directly affecting its kinase activity, effectively hampers the functional activity of NF-kappaB. This causal relationship between mis-targeting of IKK β and a lacking NF-kappaB activity is studied in two different systems: a) Real-time measurement of NF-kappaB translocation from the cytoplasm to the nucleus, and b) measurement of NF-
20 kappaB induced transcriptional activity.

These are specific IKK subunit variants fused to a GFP. As examples, the following three subunits have been chosen: IKK α (GenBank Acc.no. AF009225) , IKK β (GenBank Acc. No. AF031416), IKK γ (GenBank Acc. No. AF074382) and NIK (GenBank Acc. No.
25 NM003954).

Inspection of the scientific literature indicates that IKK β dissociates transiently from the IKAP complex during activation, and so becomes the first choice for a probe to detect redistribution.

To construct the IKK β -GFP fusion, IKK β sequences are amplified using PCR according
30 to standard protocols with the specific primers listed below. The PCR product is digested with restriction enzymes Hind3 and Acc65I, and ligated into pEGFP-N1 (Clontech, Palo Alto; GenBank Accession number U55762) digested with Hind3 and Acc65I. This produces an IKK β -EGFP fusion under the control of a CMV promoter (SEQ ID NOs: 9 and 10).

The top primer includes specific sequences following the ATG and a cloning site (Hind3). The bottom primer includes specific C-terminal sequences minus the stop codon, an Acc65I cloning site, and two extra nucleotides to preserve the reading frame in EGFP-N1.

5

IKK β -top:

5'-GTAAGCTTACATGAGCTGGTCACCTTCCCTG-3'

IKK β -bottom:

10 5'-GTGGTACCCATGAGGCCTGCTCCAG-3'

The resulting plasmids are transfected into a suitable cell line. The subcellular distribution of the probes is examined carefully by fluorescence microscopy, both under resting conditions, and upon activation, e.g. with TNF α .

15

Probes for detection of activation of the NFkappaB signal transduction pathway.

Plasmid PS377 contains an NFkappaBp65-EGFP fusion. The GenBank accession number of the p65 subunit of NFkappaB is M62399. It is constructed by performing PCR on human cDNA (from Clontech) with specific primers p65-top and p65-bottom. The resulting ca. 1.7 kb PCR product is cut with restriction enzymes Xho1 and Hind3 and cloned into pEGFP-N1 (Clontech) cut with Xho1 and Hind3. This produces an NFkappaB-EGFP fusion (SEQ ID NOs: 11 and 12) under the control of the CMV promoter.

25

p65-top: 5'-TTTACTCGAGATGGACGAACTGTTCCCCCTCA-3'

p65-bottom: 5'-TTTGAAGCTTGGAGCTGATCTGACTCAGCAGG-3'

Construction of a reporter gene assay for monitoring NFkappaB-induced transcriptional activation:

30

Plasmid PS397 contains a selectable NFkappaB reporter construct. It is constructed through ligation of two BamH1-Not1 fragments: A 2.4 kb fragment from pNFkappaB-Luc (from Clontech,), which contains a luciferase gene and NFkappaB response elements, and a 2.8 kb BamH1-Not1 fragment from pZeoSV (from Invitrogen), which contains

essential plasmid elements and a zeocin selective marker for use in E.coli and mammalian cells.

Construction of probes for monitoring IKK β localisation, mis-targeting and redistribution
5 in live cells:

Plasmid PS410 contains an EGFP-IKK β fusion. The GenBank accession number of the beta subunit of IkappaB kinase is AF031416. It is constructed by performing PCR on human cDNA (from Clontech) with specific primers IKK β -top and IKK β -stop. The resulting 2.2 kb PCR product is cut with restriction enzymes Hind3 and Acc65I and
10 cloned into pEGFP-C1 (Clontech) cut with Hind3 and Acc65I. This produces an EGFP-IKK β fusion (SEQ ID NOs: 13 and 14) under the control of the CMV promoter.

IKK β -top: 5'-GTAAGCTTACATGAGCTGGTCACCTTCCCTG-3'

IKK β -stop: 5'-GTGGTACCTCATGAGGCCTGCTCCAG-3'

15

Plasmid PS472 contains a full length IKK β under the control of the CMV promoter. It is constructed by cutting PS410 with restriction enzymes Nhe1 and Hind3, which flank EGFP. This excises EGFP sequences from the plasmid, while placing IKK β immediately downstream of the CMV promoter. The protruding ends generated by the enzymes are
20 then made blunt using Klenow polymerase according to standard protocol, and the plasmid is recircularized with DNA ligase.

PS473 contains EGFP fused to the C-terminal part of IKK β . This part of IKK β contains a putative leucine zipper region, but is without catalytic activity as this function resides in
25 the N-terminal part of IKK β . It is constructed by performing PCR on PS410 with primers IKK β -LZ-top and IKK β -stop. IKK β -LZ-top contains a Hind3 site and specific IKK β sequence from amino acid position 455 in the predicted amino acid sequence. This is almost immediately upstream of the first leucine of the predicted leucine zipper, which is at position 458. The resulting 0.9 kb PCR product is cut with restriction enzymes Hind3
30 and Acc65I and cloned into pEGFP-C1 (Clontech) cut with Hind3 and Acc65I. This produces an EGFP-IKK β -LZdomain fusion (SEQ ID NOs: 15 and 16) under the control of the CMV promoter.

IKK β -LZ-top: 5'-GTAAGCTTCCACCATGATGAATCTCCTCCGAAAC-3'

35

Plasmid PS474 contains the IKK β C-terminal part under the control of the CMV promoter. It is constructed by cutting PS473 with restriction enzymes Age1 and BspE1, which flank EGFP. This excises EGFP sequences from the plasmid, while placing IKK β sequences immediately downstream of the CMV promoter. As Age1 and BspE1 produce
5 compatible ends, the plasmid is simply recircularized with DNA ligase. The ATG methionine codon at position 455 in the predicted amino acid sequence of IKK β , may serve as initiation codon in this construct.

Transfections and cell culture conditions.

- 10 Chinese hamster ovary cells (CHO), Human epithelial kidney cells (HEK293) and Human epithelial adenocarcinoma cells (HeLa), were transfected with above mentioned plasmids using FuGENE transfection reagent (Boehringer Mannheim). Stable transfectants were selected using 1000 μ g Zeocin/ml (Invitrogen) or 500 μ g G418/ml (*Neo* marker) in the growth medium [DMEM (HEK293 and HeLa) or HAM F12 (CHO)
15 with 1000 mg glucose/l, 10 % fetal bovine serum (FBS), 100 μ g penicillin-streptomycin mixture ml⁻¹, 2 mM L-glutamine purchased from Life Technologies Inc., Gaithersburg, MD, USA).

For fluorescence microscopy, cells were allowed to adhere to Lab-Tek chambered coverglasses (Nalge Nunc Int., Naperville, IL, USA) for at least 24 hours and cultured to
20 about 80% confluence. Prior to experiments, the cells were cultured over night without selection pressure in DMEM or HAM F-12 medium with glutamax (Life Technologies), 100 μ g penicillin-streptomycin mixture ml⁻¹ and 0.3 % FBS. This medium has low autofluorescence enabling fluorescence microscopy of cells straight from the incubator.

- 25 Microscope imaging of localisation and redistribution in live cells:

Image aquisition of live cells were gathered using a Zeiss Axiovert 135M fluorescence microscope fitted with a Fluar 40X, NA: 1.3 oil immersion objective and coupled to a Photometrics CH250 charged coupled device (CCD) camera. The cells were illuminated with a 100 W HBO arc lamp. For imaging of GFP-based probes we
30 inserted in the light path was a 470 $\pm$ 20 nm excitation filter, a 510 nm dichroic mirror and a 515 $\pm$ 15 nm emission filter. For imaging of the Hoechst 33342 (H1399, Molecular Probes) nuclear stain we used a 380 $\pm$ 20 nm excitation filter, a 410 nm dichroic mirror and a 555 $\pm$ 15 nm emission filter

The cells were kept and monitored to be at 37°C with a custom built stage heater.

Quantification of NF-kappaB redistribution:

Cells are stained with the vital nuclear stain, Hoechst.

A sequence of images with a time separation of 10 sec is acquired. At each time point the sequence consists of one NF-kappaB-GFP image and one image of the Hoechst
5 stained nucleus.

The image sequence is corrected for dark current by performing a pixel-by-pixel subtraction of a dark image (an image taken under the same conditions as the actual image, except the camera shutter is not allowed to open).

The image sequence is corrected for non-uniformity of the illumination by performing a
10 pixel-by-pixel ratio with a flat field correction image (an image taken under the same conditions as the actual image of a uniformly fluorescent specimen).

At each time point the accumulated intensity of the NFkappaB probe in the nucleus is ratioed over the total cytoplasmic intensity. The Hoechst image is used to mask the
nucleus.

15

Results:

The full length IKK β probe (PS410) show an even distribution throughout the cytoplasm when expressed in CHO (Fig. 2) and HEK293 cells. PS473 show a similar localisation after its expression (Fig. 3A). Interestingly however the probe has sensitised the cells to
20 stimuli that induce apoptosis. It is thus observed that the PS473 expressing cells upon 2 hrs of serum starvation undergo apoptosis, in comparison non-transfected cells or PS410 expressing cells did show no sign on apoptosis after similar treatment. The induction of apoptosis could be visualised as a change in the localisation of the PS473 probe from an even distribution throughout the cytoplasm to a discrete punctate localisation (Fig. 3B).

25

The PS473 provoked mis-tageting of IKK β had pronounced functional consequences. We thus observed a prominent inhibition of IL-1 induced NFkappaB redistribution (Fig. 4). Furthermore we observed an inhibition of IL-1 and TNF α induced activation of the NFkappaB regulated transcription as monitored with the above described luciferase
30 reporter construct (PS397) (Fig. 5).

Figure legends

Figure 1

CHO cells expressing PS377 for monitoring NFkappaB redistribution in live cells. A) Before stimulation and B) 10 minutes after stimulation with IL-1 (10 ng/ml).

5

Figure 2

The full length IKK β probe (PS410) show an even distribution throughout the cytoplasm when expressed in CHO cells.

10 Figure 3

PS473 expressed in CHO cells. (A) show an even distribution throughout the cytoplasm. (B) The distributaion change when cells undergo appoptosis as observed after two hours of serum starvation.

15 Figure 4

Expression of PS473 inhibits IL-1 (0.5 ng/ml) induced redistribution of NF-kappaB in CHO cells.

Figure 5

20 Expression of PS473 inhibits IL-1 (0.5 ng/ml) and TNF- α (0.5 ng/ml) induced NF-kappaB regulated transcription in HEK293 cells.

References

- Ashida, S., Sakuma, K., (1992) Adv. Second Messenger Phosphoprotein Res. 25:229-239.
- Baeuerle, P.A., Baltimore, D. (1988) I-kappaB: a specific inhibitor of the NF-kappaB transcription factor. Science 242:540-546.
- 5 Baeuerle, P.A., Baltimore, D. (1989) A 65-kappaD subunit of NF-kappaB is required for inhibition of NF-kappaB by I-kappaB. Genes and Dev. 3:1689-1698.
- Baldwin, A.S. Jr (1996) The NF-kappaB and I-kappaB proteins: new discoveries and insights. Annu. Rev. Immunol. 14:649-683.
- 10 Barnes, P.J. (1995) Ann, Med. 27:531-535.
- Beavo, J.A., (1995) Physiol. Rev 75:725-748.
- Beavo, J.A., Reifsnnyder, D.H. (1990) Trends Pharmacol. Sci. 11:150-155
- Beg, A.A., Finco, T.S., Nantermet, P.V., Baldwin, A.S. (1993) Tumor necrosis factor and interleukin-1 lead to phosphorylation and loss of I-kappa-B α - a mechanism of NF-
- 15 kappaB activation. Mol. Cell. Biol. 13:3301-3310.
- Benner, B.M., Ballermann, B.J., Gunning, M.E., Zeidel, M.L. (1990) Physiol. Rev. 70:665-699
- Biel, M., Zong, X., Distler, M., Bosse, E., Klugbauer, N., Murakami, M., Flockerzi, V. and Hofmann, F. (1994) Comment in: Proc Natl Acad Sci USA. 91: 3505-3509.
- 20 Bird, T.A., Schooley, K., Dower, S.K., Hagen, H., Virca, G.D., (1997) Activation of nuclear transcription factor NF-kappaB by interleukin-1 is accompanied by casein kinase II-mediated phosphorylation of the p65 subunit. 272:32606-32612.
- Blackwell, T.S., Christman, J.W. (1997) The role of nuclear factor-kappa B in cytokine gene regulation. Am. J. Resp. Cell and Mol. Biol. 17:3-9.
- 25 Blease, K., Burkegaffney, A., Hellewell, P.G. (1998) Br. J. Pharm. 124:229-237.
- Bolger *et al.*, (1997) Biochem. J. 328:539-548.
- Bolger, G., Michaeli, T., Martins, T., St John, T., Steiner, B., Rodgers, L., Riggs, M., Wigler, M., Ferguson, K. (1993) Mol. Cell Biol. 13:6558-6571.
- Bolger, G.B., McPhee, I., Houslay, M.D. (1996) J. Biol. Chem. 271:1065-1071
- 30 Bolger, G.B., Rodgers, L., Riggs, M. (1994) Gene 149:237-244.
- Bourcier, T., Sukhove, G., Libby P. (1997) The nuclear factor kappa-B signalling pathway participates in dysregulation of vascular smooth muscle cells in vitro and in human atherosclerosis. J.Biol. Chem. 272:15817-15824

- Bours, V., Dejardin, E., Bonizzi, G., Merville, M.P., Piette, J. (1998). The NF-kappaB transcription factor - role in oncogenesis and in response to anticancer therapeutics. *Medecine Sciences* 14:566-571
- Brattsand, R., Linden, M. (1996) Cytokine modulation by glucocorticoids: mechanisms
5 and actions in cellular studies. *Alimentary Pharmacol. Therapeutics* 10:81-90
- Burns, F., Rodger, I.W. and Pyne, N.J. (1992) *Biochem. J.* 283 (Pt 2): 487-491.
- Chen, ZJ ; Parent, L; Maniatis, T. (1996) Site-specific phosphorylation of i-kappa-b-alpha by a novel ubiquitination-dependent protein-kinase activity. *Cell* 84: 853-862.
- Chini, C.C.S., Grande, J.P., Chini, E.N., Dousa, T.P. (1997) *J. Biol. Chem.* 272:9854-
10 9859.
- Chiu, P.J., Vemulapalli, S., Chintala, M., Kurowski, S., Tetzloff, G.G., Brown, A.D. and Sybertz, E.J. (1997) *Naunyn-schmiedeberg's Archives of Pharmacology*, 355: 463-469.
- Coghlan, V.M, Hausken, Z.E., Scott, J.D. (1995) *Biochem. Soc. Trans.* 23:592-596.
- 15 Cogswell, P.C., Schienman, R.I., Baldwin, A.S. (1993) Promoter of the human NF-kappaB p50/p105 gene - regulation by NF-kappaB subunits and by c-rel. *J. Immunol.* 150:2794-2804.
- Cohen, L.; Henzel, W.J.; Baeuerle, P.A. (1998) IKAP is a scaffold protein of the Ikb kinase complex . *Nature* 395: 292-296.
- 20 Disanto, M.E., Glaser, K.B., Heaslip, R.J. (1995) *Cell. Sig.* 7:827-835.
- Drewett, J.G. and Garbers, D.L. (1994) *Endocrine Reviews*, 15: 135-162.
- Entman, M.L., Smith, C.W. (1994) *Cardiovasc. Res.* 28:1301-1311
- Essayan, D.M., Kageysobotka, A., Lichenstein, L.M., Huang, S.K.(1997) *J. Pharm. Exp. Ther.* 282:505-512.
- 25 Faux, M.C., Scott, J.D., (1996) *Cell* 85:9-12.
- Ferrari, D., Wesselborg, S., Bauer, M.K.A., Schulze-Osthoff, K., (1997) Extracellular ATP activates transcription factor NF-kappaB through the P2Z purinoreceptor by selectively targeting NF-kappaB p65 (RelA). *J. Cell Biol.* 139:1635-1643
- Fisher, D.A., Smith, J.F., Pillar, J.S., Stdenis, S.H. and Cheng, J.B. (1998) *J. Biol. Chem.*
30 273: 15559-15564.
- Fujishige, K., Kotera, J., Michibata, H., Yuasa, K., Takebayashi, S., Okumura, K., Omori, K. (1999) *J. Biol. Chem.* 274:18438-18445
- Ghosh, S., *et al.*, (1990) Cloning of the p50 DNA-binding subunit of NF-kappaB: homology to rel and dorsal. *Cell*:62:1007-1018.

- Giorgi, M., Giordano, D., Caniglia, C., Biagioni, S., Augusticocco, G. (1997) *Int. J. Dev. Neurosci.* 15:309-319.
- Giri, D.K., Aggarwal, B.B. (1998) Constitutive activation of NF-kappaB causes resistance to apoptosis in human cutaneous t-cell lymphoma HuT-78 cells - autocrine role of tumor-necrosis factor and reactive oxygen species. 273:14008-14014
- 5 Griswold, D.E. *et al.*, (1993) *Inflammation* 17:333-343.
- Grumont, R.J., Gerondakis, S. (1994) The subunit composition of NF-kappaB complexes changes during B-cell development. *Cell Growth and Diffn.* 5:1321-1331.
- Hallahan, D., Clark, E.T., Kuchibholta, J., Gewertz, B.L., Collins, T. (1995) E-selectin
- 10 gene induction by ionizing-radiation is independent of cytokine induction. *Biochem. Biophys. Res. Comm.* 217:784-795.
- Hallsworth, M.P., Giembycz, M.A., Barnes, P.J., Lee, T.H. (1996) *Br. J. Pharmacol.* 117:79-86.
- Hattori, Y., Akimoto, K., Murakami, Y., Kasai, K. (1997) Pyrrolidine dithiocarbamate
- 15 inhibits cytokine-induced VCAM-1 gene expression in rat cardiac myocytes. *Mol. Cell. Biochem.* 177:177-181.
- Hayashi, T., Sekine, T., Okamoto, T. (1993) Identification of a new serine kinase that activates NF-kappaB by direct phosphorylation. *J. Biol. Chem.* 268:26790-26795.
- Hiramoto, M., *et al.*, (1998) Nuclear targeted suppression of NF-kappaB activity by the
- 20 quinone derivative E3330. *J. Immunol.* 160:810-819.
- Hofmann, F., Dostmann, W., Keilbach, A., Landgraf, W. and Ruth, P. (1992) *Biochimica Et Biophysica Acta*, 1135: 51-60.
- Holliday, L.S., Dean, A.D., Lin, R.H., Greenwald, J.E. and Gluck, S.L. (1997) *Am. J. Physiol.* 272: F283-F291
- 25 Houslay, M.D., Milligan, G. (1997) *Trends in Biochem. Sci.* 22:217-224.
- Houslay, M.D., Sullivan, M., Bolger, G.B. (1998) *Adv. Pharmacol.* 44:225-242
- Hughes B. *et al.*, (1997) *Drug Discov. Today* 2:89-101.
- Hughes, B. *et al.*, (1996) *Br. J. Pharmacol.* 118:1183-1191.
- Huston *et al.*, (1997) *Biochem. J.* 328:549-558.
- 30 Huston, E., Pooley, L., Julien, J., Scotland, G., McPhee, I., Sullivan, M., Bolger, G., Houslay, M.D. (1996) *J. Biol. Chem.* 271:31334-31344.
- Jacobitz, S., McLaughlin, M.M., Livi, G.P., Burman, M., Torphy, T.J. (1996) *Mol. Pharmacol.* 50:891-899.
- Jilg, S. *et al.*, (1996) *J. Pharmacol. Exp. Ther.* 278:421-431.

- Jourd'heuil, D., Morise, Z., Conner, E.M., Grisham, M.B. (1997) Oxidants, transcription factors and intestinal inflammation. *J. Clin. Gastroenterol.* 25:S61-S72.
- Kambayashi, T. *et al.*, (1995) *J. Immunol.* 155:4909-4916.
- Keilbach, A., Ruth, P. and Hofmann, F. (1992) *Eur. J. Biochem.* 208: 467-473.
- 5 Lezoual'ch, F., Behl, C. (1998) Transcription factor NF-kappaB: Friend or foe of neurons? *Mol. Psychiatry* 3:15-20
- Light, D.B., Corbin, J.D. and Stanton, B.A. (1990) *Nature*, 344: 336-339.
- Lincoln, T.M. and Cornwell, T.L. (1993) *Faseb Journal*, 7: 328-338.
- Lincoln, T.M., Komalavilas, P. and Cornwell, T.L. (1994) *Hypertension*, 23: 1141-1147.
- 10 Liu, C., Ding, J.M., Faiman, L.E. and Gillette, M.U. (1997) *J. Neuroscience*, 17: 659-666.
- Lochhead, A., Nekrasova, E., Arshavsky, V.Y. and Pyne, N.J. (1997) *J. Biol. Chem.* 272: 18397-18403.
- MacFarland, R.T., Zelus, B.D., Beavo, J.A., (1991) *J. Biol. Chem.* 266:136-142.
- MacMillan-Crow, L.A. and Lincoln, T.M. (1994) *Biochemistry*, 33: 8035-8043.
- 15 Makarov, S.S., Johnston, W.N., Olsen, J.C., Watson, J.M., Mondal, K., Rinehart, C., Haskill, J.S. (1997) NF-kappaB as a target for anti-inflammatory gene therapy - suppression of inflammatory responses in monocytic and stromal cells by stable gene-transfer of I-kappaB α cDNA. *Gene Therapy* 4:846-852
- Mathur, A., Golombek, D.A. and Ralph, M.R. (1996) *Am. J. Physiol.* 270: R1031-R1036
- 20 McPhee, I., *et al.*, (1995) *Biochem. J.* 310:965-974.
- Miotla, J.M., Teixeira, M.M., Jeffery, P.K., Hellewell, P.G. (1995) *Br. J. Pharmacol.* 116:5P.
- Morandini, R., *et al.*, (1996) *Am. J. Physiol.* 270:H807-H816.
- Mukaida, N. Mahe, Y., Matsushima, K., (1990) Co-operative interaction of NF-kappaB and cis-regulatory enhancer binding protein-like factor binding elements in activation
- 25 the interleukin-8 gene by pro-inflammatory cytokines. *J. Biol. Chem.* 265:21128-21133.
- Naumann, M., Wulczyn, F.G., Scheidereit, C., (1993) The NF-kappaB precursor p105 and the proto-oncogene product bcl-3 are I-kappaB molecules and control nuclear
- 30 translocation of NF-kappaB. *EMBO J.* 12:213-222
- Nemoz, G., Prigent, A.F., Moueqqit, M., Fougles, S., Machovschi, O., Pacheco, H. (1985) *Biochem. Pharmacol.* 34:2997-3000.
- Nemoz, G., Sette, C., Conti, M. (1997) *Mol. Pharm.* 51:242-249.

- Neumann, M., Marienfeld, R., Serfling, E. (1997) Rel/NF-kappaB transcription factors and cancer - oncogenesis by dysregulated transcription. *Int. J. Oncology* 11:1335-1347
- Nicholson, C.D., Shahid, M. (1994) *Pulm. Pharmacol.* 7:1-17.
- 5 Nolan, G.P., Fujita, T., Bhatia, K., Huppi, C., Liou, H.C., Scott, M.L., Baltimore, D., (1993) The bcl-3 proto-oncogene encodes a nuclear I-kappaB-like molecule that preferentially interacts with NF-kappaB p50 and p52 in a phosphorylation-dependent manner. *Mol. Cell. Biol.*, 13:3557-3566
- O'Connell, J.C., *et al.*, (1996) *Biochem. J.* 318:255-262.
- 10 O'Neill, L.A., Kaltschmidt, C. (1997) NF-kappaB: a crucial transcription factor for glial and neuronal cell function. *Trends Neurosci.* 20:252-258.
- Oberholte, R., Ratzliff, J., Baecker, P.A., Daniels, D.V., Zuppan, P., Jarnagin, K., Shelton, E.R. (1997) *Biochim. Biophys. Acta* 1353:287-297.
- Owens, R.J., Catterall, C., Batty, D., Jappy, J., Russell, A., Smith, B., O'Connell, J.,
- 15 Perry, M. (1997) *Biochem. J.* 326:53-60.
- Pawson, T., Scott, J.D. (1997) *Science* 278:2075-2080.
- Perry, M.J., Higgs, G.A. (1998) *Curr. Opin. Chem. Biol.* 2:472-481.
- Pettipher, E.R., *et al.*, (1996) *Br. J. Pharmacol.* 117:1530-1534.
- Pooley, L., Shakur, Y., Rena, G., Houslay, M.D. (1997) *Bioch. J.* 321:177-185.
- 20 Prabhakar, U., *et al.*, (1994) *Int. J. Immunopharmacol.* 16:805-816.
- Pyne, N.J., Arshavsky, V. and Lochhead, A. (1996) *Biochem. Soc. Trans.* 24: 1019-1022.
- Pyne, N.J., Cooper, M.E. and Houslay, M.D. (1987) *Biochemical Journal*, 242: 33-42.
- Raeburn, D. and Karlsson, J.A. (1993) *J. Pharm. Exp. Ther.* 267: 1147-1152.
- 25 Raeburn, D., Advenier, C. (1995) *Int. J. Biochem. Cell Biol.* 27:29-37.
- Rothwarf, D.M., Zandi, E., Natoli, G., Karin, M. (1998) IKK-gamma is an essential regulatory subunit of the I-kappaB kinase complex. *Nature* 395: 297-300.
- Schulze-Osthoff, K., Ferrari, D., Riehemann, K., Wesselborg, S. (1997) Regulation of NF-kappaB activation by MAP kinase cascades. *Immunobiol.* 198:35-49.
- 30 Scotland *et al.*, (1998) *Methods - A companion to Meth. Enzymology* 14:65-79.
- Sekut, L. *et al.*, (1995) *Clin. Exp. Immunol.* 100:126-132.
- Sen, R., Baltimore, D., 1986. Inducibility of kappa immunoglobulin enhancer-binding protein NF-kappaB by a post-translational mechanism. *Cell* 47:921-928.
- Sette, C., Conti, M., (1996) *J. Biol. Chem.* 271:16526-16534.
- 35 Sette, C., Conti, M., (1996) *J. Biol. Chem.* 271:16526-16534.

- Shakur, Y. *et al.*, (1995) *Biochem. J.* 310:965-974.
- Smith, R.S., Smith, T.J., Blieden, T.M., Phipps, R.P. (1997) Fibroblasts as sentinel cells. Synthesis of chemokines and regulation of inflammation. *Am. J. Path.* 151:317-322.
- Soderling, S.H., Bayuga, S.J. and Beavo, J.A. (1998) *J. Biol. Chem.* 273: 15553-15558.
- 5 Soderling, S.H., Bayuga, S.J., Beavo, J.A. (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:7071-7076
- Swinnen, J.V., Joseph, D.R., Conti, M. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:8197-8201.
- Swinnen, J.V., Tsikalas, K.E., Conti, M. (1991) *J. Biol. Chem.* 266:18370-18377.
- 10 Teixeira, M.M., Gristwood, R.W., Cooper, N., Hellewell, P.G. (1997) *Trends in Pharm. Science* 18:164-171.
- Teixeira, M.M., Rossi, A.G., Giembycz, M.A., Hellewell, P.G. (1996) *Br. J. Pharmacol.* 118:2099-2106.
- Teixeira, M.M., Rossi, A.G., Williams, T.J., Hellewell, P.G. (1994) *Br. J. Pharmacol.* 15 112:332-340.
- Thomas, M.K., Francis, S.H. and Corbin, J.D. (1990) *J. Biol. Chem.* 265: 14971-14978.
- Torphy, T.J. (1998) *Am. J. Respir. Crit. Care Med.* 157:351-370.
- Turner, C.R., Andresen, C.J., Smith, W.B., Watson, J.W. (1994) *Am. J. Respir. Crit. Care Med.* 149:1153-1159.
- 20 Vaandrager, A.B. and de Jonge, H.R. (1996) *Mol. Cell. Biochem.* 157: 23-30.
- Vemulapalli, S., Watkins, R.W., Chintala, M., Davis, H., Ahn, H.S., Fawzi, A., Tulshian, D., Chiu, P., Chatterjee, M., Lin, C.C. and Sybertz, E.J. (1996) *J. Cardiovasc. Pharm.* 28: 862-869.
- Verghese, M.W., McConnell, R.T., Lenhard, J.M., Hamacher, L., Jin, S.L.C. (1995) *Mol. Pharm.* 47:1164-1171.
- 25 Wachtel, H. (1982) *Psychopharmacology* 77:309-314.
- Watanabe, N., Iwamura, T., Shinoda, T., Fujita, T. (1997) Regulation of NF-kappaB 1 proteins by the candidate oncoprotein bcl3 - generation of NF-kappaB homodimers from the cytoplasmic pool of p50-p105 and nuclear translocation. *EMBO J.* 16:3609-3620.
- 30 Wilson, M., Sullivan, M, Brown, N., Houslay, M.D. (1994) *Biochem. J.* 304:407-415.
- Wulczyn, F.G., Krappmann, D., Scheidereit, C., (1996) the NF-kappaB B/Rel and I-kappaB gene families: mediators of immune response and inflammation. *J. Mol. Medicine* 74:749-769.

Zhong, H.H., Voll, R.E., Ghosh, S., (1998) Phosphorylation of NF-kappaB p65 by PKA stimulates transcriptional activity by promoting a novel bivalent interaction with the coactivator CBP/p300. *Molecular Cell* 5:661-671.

Claims

1. Use of a substance, capable of modulating the specific effectiveness of a cyclic nucleotide phosphodiesterase or I-kappaB kinases through modulating the spatial
5 distribution or change in spatial distribution of the cyclic nucleotide phosphodiesterases or I-kappaB kinases within cells of an animal, for the preparation of a medicament for the prevention or treatment in an animal of an adverse condition which may be reduced or abolished by modulating the activity of one or more cyclic nucleotide phosphodiesterases having the ability to cleave cyclic AMP or cyclic GMP or by
10 modulating the activity of one or more I-kappaB.
2. Use according to claim 1, wherein the I-kappaB kinase is selected from the group consisting of I-kappaB kinase α , I-kappaB kinase β , I-kappaB kinase γ and NIK.
- 15 3. Use according to claim 2, wherein the I-kappaB kinase is I-kappaB kinase β .
4. Use according to claim 1, wherein the cyclic nucleotide phosphodiesterase is selected from the group consisting of PDE3, PDE4, PDE7 and PDE8.
- 20 5. Use according to claim 4, wherein the cyclic nucleotide phosphodiesterase is PDE4.
6. Use according to claim 5, wherein the cyclic nucleotide phosphodiesterase is a splice variant of PDE4, selected from the group consisting of PDE4A, PDE4B, PDE4C and PDE4D.
25
7. Use according to claim 6, wherein the PDE4 species is a splice variant of PDE4D.
8. Use according to claim 7, wherein the splice variant is PDE4D1, PDE4D2, PDE4D3, PDE4D4, PDE4D5 and PDE4A1.
30
9. Use according to claim 8, wherein the splice variant is PDE4D3, PDE4D4 or PDE4D5.
10. Use according to claim 6, wherein the PDE4 splice variant is PDE4A1.

11. Use according to any of the preceding claims, wherein the adverse condition is an inflammatory diseases such as chronic inflammation.
12. Use according to any of claims 1-10, wherein the adverse condition is chronic
5 inflammatory airway diseases such as asthma and chronic bronchial hyperreactivity of non-asthma etiology.
13. Use according to any of claims 1-10, wherein the adverse condition is chronic inflammatory joint diseases such as rheumatoid arthritis and pelvospondylitis.
- 10 14. Use according to any of claims 1-10, wherein the adverse condition is chronic inflammatory bowel diseases such as ulcerative colitis and Crohn's disease.
- 15 15. Use according to any of claims 1-10, wherein the adverse condition is autoimmune diseases with chronic inflammation such as rheumatoid arthritis, diabetes mellitus type I, systemic lupus erythematosus, myasthenia gravis, Hashimoto's thyroiditis, Graves' disease and immune thrombocytopenic purpura.
- 20 16. Use according to any of claims 1-10, wherein the adverse condition involves a disregulation of the immune system such as acute respiratory distress syndrome (ARDS) and septic shock.
17. Use according to claim 10, wherein the adverse condition is depression.
- 25 18. Use according to claim 1, wherein the cyclic nucleotide phosphodiesterase is selected from the group consisting of PDE1, PDE2, PDE5, PDE6, PDE9 and PDE10.
19. Use according to claim 18, wherein the nucleotide phosphodiesterase is a splice variant of PDE5.
- 30 20. Use according to claim 18 or 19, wherein the adverse condition is hypo- or hypertension, erectile dysfunction, circadian rhythm resetting or jet-lag.
21. Use according to any of the preceding claims wherein the animal is a mammal.

22. Use according to claim 21, wherein the mammal is a human being.
23. Use according to any of the preceding claims, wherein the substance is an organic compound having a molecular weight of around 3000 Da
- 5 24. Use according to any of claims 1-22, wherein the substance is an organic compound having a molecular weight of at the most 1200 Da.
- 10 25. Use according to claim 24, wherein the substance is an organic compound having a molecular weight of at the most 900 Da.
26. Use according to claim 25, wherein the substance is an organic compound having a molecular weight of at the most 600 Da.
- 15 27. Use according to claim 26, wherein the substance is an organic compound having a molecular weight of at the most 300 Da.
28. Use according to any of the preceding claims, wherein the substance is a peptide.
- 20 29. Use according to any of claim 1-27, wherein the substance is a carbon-containing non-peptide.
30. Use according to any of the preceding claims, wherein the organic compound is a compound having one or more chemical domains capable of interacting with one or
- 25 more functional groups of the targeting sequence of the native anchoring site of the cyclic nucleotide phosphodiesterase or I-kappaB kinase.
31. Use according to any of the preceding claims, wherein the substance interacts with the targeting sequence or part thereof in a manner that dislocates, disrupts targeting, or
- 30 interferes with redistribution of the fluorescent probe as measured in quantitative fluorescence redistribution assay.
32. A method for extracting quantitative information relating to an influence on a cellular response, the method comprising recording variation, caused by the influence on a
- 35 mechanically intact living cell or mechanically intact living cells, in spatially distributed

light emitted from a luminophore, the luminophore being part of a fluorescent probe further comprising at least a part of a cyclic nucleotide phosphodiesterase or I-kappaB kinase, the fluorescent probe being present in the cell or cells and being capable of being redistributed in a manner which is related with the degree of the influence, and/or
5 of being modulated by a component which is capable of being redistributed in a manner which is related to the degree of the influence, the association resulting in a modulation of the luminescence characteristics of the luminophore, and processing the recorded variation in the spatially distributed light to provide quantitative information correlating the spatial distribution to the degree of the influence on the cellular response.

10

33. A screening assay for carrying out the method of claim 32.

34. A screening assay according to claim 32 or 33 wherein the fluorescent probe is modified in a systematic way, still keeping the GFP coding sequence intact, so that the
15 new fluorescent probes are fusion polypeptides where parts of the suspected targeting sequences are altered.

35. A screening assay according to claim 34, wherein the modification of the suspected targeting sequence is a deletion.

20

36. A screening assay according to any of claims 33-35, wherein the spatial distribution of the fluorescent probe is compared to the spatial distribution of the unmodified fluorescent probe to deduce the targeting sequence.

25 37. A screening assay according to any of claims 33-36, wherein the quantitative fluorescence redistribution assay is a primary screening assay used in a discovery project

38. A nucleotide sequence encoding the protein corresponding to amino acids 331-552
30 of SEQ ID NO: 16 or any sub-sequence thereof of more than 25 contiguous amino acids, able to dislocate IKK β when expressed in CHO cells under the control of the CMV promoter.

39. A nucleotide sequence according to claim 38, wherein the sub-sequence is the
35 predicted leucine zipper contained in amino acids 331-360 of SEQ ID NO: 16.

40. A screening assay according to any of claims 33-37, wherein the fluorescent probe comprises a nucleotide sequence according to claim 38 or 39.

5 41. A method according to claim 32 wherein the fluorescent probe is able to dislocate IKK β when expressed in CHO cells under the control of the CMV promoter.

42. A method for preventing or treating, in an animal in need thereof, an adverse condition which may be reduced or abolished by modulating the activity of one or more cyclic nucleotide phosphodiesterases having the ability to cleave cyclic AMP, or cyclic
10 AMP, or by modulating the activity of one or more I-kappaB kinases, the method comprising modulating the specific effectiveness of the cyclic nucleotide phosphodiesterase or I-kappaB kinase by modulating the spatial distribution within cells of the animal.

Figures

Fig. 1A




Fig. 1B




Fig. 2






Fig. 3A**Fig. 3B****Fig. 4**

Fig. 5

SEQUENCE LISTING

<110> BioImage A/S

<120> Specific therapeutic interventions
obtained by interference with redistribution and/or
targetting

<130> 22130PC1

<160> 16

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2793

<212> DNA

<213> Aequorea victoria and human

<220>

<221> CDS

<222> (1)...(2793)

<400> 1

atg atg cac gtg aat aat ttt ccc ttt aga agg cat tcc tgg ata tgt	48
Met Met His Val Asn Asn Phe Pro Phe Arg Arg His Ser Trp Ile Cys	
1 5 10 15	
ttt gat gtg gac aat ggc aca tct gcg gga cgg agt ccc ttg gat ccc	96
Phe Asp Val Asp Asn Gly Thr Ser Ala Gly Arg Ser Pro Leu Asp Pro	
20 25 30	
atg acc agc cca gga tcc ggg cta att ctc caa gca aat ttt gtc cac	144
Met Thr Ser Pro Gly Ser Gly Leu Ile Leu Gln Ala Asn Phe Val His	
35 40 45	
agt caa cga cgg gag tcc ttc ctg tat cga tcc gac agc gat tat gac	192
Ser Gln Arg Arg Glu Ser Phe Leu Tyr Arg Ser Asp Ser Asp Tyr Asp	
50 55 60	
ctc tct cca aag tct atg tcc cgg aac tcc tcc att gcc agt gat ata	240
Leu Ser Pro Lys Ser Met Ser Arg Asn Ser Ser Ile Ala Ser Asp Ile	
65 70 75 80	
cac gga gat gac ttg att gtg act cca ttt gct cag gtc ttg gcc agt	288
His Gly Asp Asp Leu Ile Val Thr Pro Phe Ala Gln Val Leu Ala Ser	
85 90 95	
ctg cga act gta cga aac aac ttt gct gca tta act aat ttg caa gat	336
Leu Arg Thr Val Arg Asn Asn Phe Ala Ala Leu Thr Asn Leu Gln Asp	
100 105 110	
cga gca cct agc aaa aga tca ccc atg tgc aac caa cca tcc atc aac	384
Arg Ala Pro Ser Lys Arg Ser Pro Met Cys Asn Gln Pro Ser Ile Asn	
115 120 125	
aaa gcc acc ata aca gag gag gcc tac cag aaa ctg gcc agc gag acc	432
Lys Ala Thr Ile Thr Glu Glu Ala Tyr Gln Lys Leu Ala Ser Glu Thr	

130	135	140	
ctg gag gag ctg gac tgg tgt ctg gac cag cta gag acc cta cag acc			480
Leu Glu Glu Leu Asp Trp Cys Leu Asp Gln Leu Glu Thr Leu Gln Thr			
145	150	155	160
agg cac tcc gtc agt gag atg gcc tcc aac aag ttt aaa agg atg ctt			528
Arg His Ser Val Ser Glu Met Ala Ser Asn Lys Phe Lys Arg Met Leu			
165	170		175
aat cgg gag ctc acc cat ctc tct gaa atg agt cgg tct gga aat caa			576
Asn Arg Glu Leu Thr His Leu Ser Glu Met Ser Arg Ser Gly Asn Gln			
180	185		190
gtg tca gag ttt ata tca aac aca ttc tta gat aag caa cat gaa gtg			624
Val Ser Glu Phe Ile Ser Asn Thr Phe Leu Asp Lys Gln His Glu Val			
195	200		205
gaa att cct tct cca act cag aag gaa aag gag aaa aag aaa aga cca			672
Glu Ile Pro Ser Pro Thr Gln Lys Glu Lys Glu Lys Lys Lys Arg Pro			
210	215		220
atg tct cag atc agt gga gtc aag aaa ttg atg cac agc tct agt ctg			720
Met Ser Gln Ile Ser Gly Val Lys Lys Leu Met His Ser Ser Ser Leu			
225	230		240
act aat tca agt atc cca agg ttt gga gtt aaa act gaa caa gaa gat			768
Thr Asn Ser Ser Ile Pro Arg Phe Gly Val Lys Thr Glu Gln Glu Asp			
245	250		255
gtc ctt gcc aag gaa cta gaa gat gtg aac aaa tgg ggt ctt cat gtt			816
Val Leu Ala Lys Glu Leu Glu Asp Val Asn Lys Trp Gly Leu His Val			
260	265		270
ttc aga ata gca gag ttg tct ggt aac cgg ccc ttg act gtt atc atg			864
Phe Arg Ile Ala Glu Leu Ser Gly Asn Arg Pro Leu Thr Val Ile Met			
275	280		285
cac acc att ttt cag gaa cgg gat tta tta aaa aca ttt aaa att cca			912
His Thr Ile Phe Gln Glu Arg Asp Leu Leu Lys Thr Phe Lys Ile Pro			
290	295		300
gta gat act tta att aca tat ctt atg act ctc gaa gac cat tac cat			960
Val Asp Thr Leu Ile Thr Tyr Leu Met Thr Leu Glu Asp His Tyr His			
305	310		320
gct gat gtg gcc tat cac aac aat atc cat gct gca gat gtt gtc cag			1008
Ala Asp Val Ala Tyr His Asn Asn Ile His Ala Ala Asp Val Val Gln			
325	330		335
tct act cat gtg cta tta tct aca cct gct ttg gag gct gtg ttt aca			1056
Ser Thr His Val Leu Leu Ser Thr Pro Ala Leu Glu Ala Val Phe Thr			
340	345		350
gat ttg gag att ctt gca gca att ttt gcc agt gca ata cat gat gta			1104
Asp Leu Glu Ile Leu Ala Ala Ile Phe Ala Ser Ala Ile His Asp Val			
355	360		365
gat cat cct ggt gtg tcc aat caa ttt ctg atc aat aca aac tct gaa			1152

Asp His Pro Gly Val Ser Asn Gln Phe Leu Ile Asn Thr Asn Ser Glu	
370 375 380	
ctt gcc ttg atg tac aat gat tcc tca gtc tta gag aac cat cat ttg	1200
Leu Ala Leu Met Tyr Asn Asp Ser Ser Val Leu Glu Asn His His Leu	
385 390 395 400	
gct gtg ggc ttt aaa ttg ctt cag gaa gaa aac tgt gac att ttc cag	1248
Ala Val Gly Phe Lys Leu Leu Gln Glu Glu Asn Cys Asp Ile Phe Gln	
405 410 415	
aat ttg acc aaa aaa caa aga caa tct tta agg aaa atg gtc att gac	1296
Asn Leu Thr Lys Lys Gln Arg Gln Ser Leu Arg Lys Met Val Ile Asp	
420 425 430	
atc gta ctt gca aca gat atg tca aaa cac atg aat cta ctg gct gat	1344
Ile Val Leu Ala Thr Asp Met Ser Lys His Met Asn Leu Leu Ala Asp	
435 440 445	
ttg aag act atg gtt gaa act aag aaa gtg aca agc tct gga gtt ctt	1392
Leu Lys Thr Met Val Glu Thr Lys Lys Val Thr Ser Ser Gly Val Leu	
450 455 460	
ctt ctt gat aat tat tcc gat agg att cag gtt ctt cag aat atg gtg	1440
Leu Leu Asp Asn Tyr Ser Asp Arg Ile Gln Val Leu Gln Asn Met Val	
465 470 475 480	
cac tgt gca gat ctg agc aac cca aca aag cct ctc cag ctg tac cgc	1488
His Cys Ala Asp Leu Ser Asn Pro Thr Lys Pro Leu Gln Leu Tyr Arg	
485 490 495	
cag tgg acg gac cgg ata atg gag gag ttc ttc cgc caa gga gac cga	1536
Gln Trp Thr Asp Arg Ile Met Glu Glu Phe Phe Arg Gln Gly Asp Arg	
500 505 510	
gag agg gaa cgt ggc atg gag ata agc ccc atg tgt gac aag cac aat	1584
Glu Arg Glu Arg Gly Met Glu Ile Ser Pro Met Cys Asp Lys His Asn	
515 520 525	
gct tcc gtg gaa aaa tca cag gtg ggc ttc ata gac tat att gtt cat	1632
Ala Ser Val Glu Lys Ser Gln Val Gly Phe Ile Asp Tyr Ile Val His	
530 535 540	
ccc ctc tgg gag aca tgg gca gac ctc gtc cac cct gac gcc cag gat	1680
Pro Leu Trp Glu Thr Trp Ala Asp Leu Val His Pro Asp Ala Gln Asp	
545 550 555 560	
att ttg gac act ttg gag gac aat cgt gaa tgg tac cag agc aca atc	1728
Ile Leu Asp Thr Leu Glu Asp Asn Arg Glu Trp Tyr Gln Ser Thr Ile	
565 570 575	
cct cag agc ccc tct cct gca cct gat gac cca gag gag ggc cgg cag	1776
Pro Gln Ser Pro Ser Pro Ala Pro Asp Asp Pro Glu Glu Gly Arg Gln	
580 585 590	
ggt caa act gag aaa ttc cag ttt gaa cta act tta gag gaa gat ggt	1824
Gly Gln Thr Glu Lys Phe Gln Phe Glu Leu Thr Leu Glu Glu Asp Gly	
595 600 605	

gag tca gac acg gaa aag gac agt ggc agt caa gtg gaa gaa gac act Glu Ser Asp Thr Glu Lys Asp Ser Gly Ser Gln Val Glu Glu Asp Thr 610 615 620	1872
agc tgc agt gac tcc aag act ctt tgt act caa gac tca gag tct act Ser Cys Ser Asp Ser Lys Thr Leu Cys Thr Gln Asp Ser Glu Ser Thr 625 630 635 640	1920
gaa att ccc ctt gat gaa cag gtt gaa gag gag gca gta ggg gaa gaa Glu Ile Pro Leu Asp Glu Gln Val Glu Glu Glu Ala Val Gly Glu Glu 645 650 655	1968
gag gaa agc cag cct gaa gcc tgt gtc ata gat gat cgt tct cct gac Glu Glu Ser Gln Pro Glu Ala Cys Val Ile Asp Asp Arg Ser Pro Asp 660 665 670	2016
acg acg gga att ctg cag tcg acg gta ccg cgg gcc cgg gat cca ccg Thr Thr Gly Ile Leu Gln Ser Thr Val Pro Arg Ala Arg Asp Pro Pro 675 680 685	2064
gtc gcc acc atg gtg agc aag ggc gag gag ctg ttc acc ggg gtg gtg Val Ala Thr Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val 690 695 700	2112
ccc atc ctg gtc gag ctg gac ggc gac gta aac ggc cac aag ttc agc Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser 705 710 715 720	2160
gtg tcc ggc gag ggc gag ggc gat gcc acc tac ggc aag ctg acc ctg Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu 725 730 735	2208
aag ttc atc tgc acc acc ggc aag ctg ccc gtg ccc tgg ccc acc ctc Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu 740 745 750	2256
gtg acc acc ctg acc tac ggc gtg cag tgc ttc agc cgc tac ccc gac Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp 755 760 765	2304
cac atg aag cag cac gac ttc ttc aag tcc gcc atg ccc gaa ggc tac His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr 770 775 780	2352
gtc cag gag cgc acc atc ttc ttc aag gac gac ggc aac tac aag acc Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr 785 790 795 800	2400
cgc gcc gag gtg aag ttc gag ggc gac acc ctg gtg aac cgc atc gag Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu 805 810 815	2448
ctg aag ggc atc gac ttc aag gag gac ggc aac atc ctg ggg cac aag Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys 820 825 830	2496
ctg gag tac aac tac aac agc cac aac gtc tat atc atg gcc gac aag Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys 835 840 845	2544

cag aag aac ggc atc aag gtg aac ttc aag atc cgc cac aac atc gag 2592
 Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu
 850 855 860
 gac ggc agc gtg cag ctc gcc gac cac tac cag cag aac acc ccc atc 2640
 Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile
 865 870 875 880
 ggc gac ggc ccc gtg ctg ctg ccc gac aac cac tac ctg agc acc cag 2688
 Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln
 885 890 895
 tcc gcc ctg agc aaa gac ccc aac gag aag cgc gat cac atg gtc ctg 2736
 Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu
 900 905 910
 ctg gag ttc gtg acc gcc gcc ggg atc act ctc ggc atg gac gag ctg 2784
 Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu
 915 920 925
 tac aag taa 2793
 Tyr Lys *
 930

<210> 2
 <211> 930
 <212> PRT
 <213> Aequorea victoria and human

<400> 2
 Met Met His Val Asn Asn Phe Pro Phe Arg Arg His Ser Trp Ile Cys
 1 5 10 15
 Phe Asp Val Asp Asn Gly Thr Ser Ala Gly Arg Ser Pro Leu Asp Pro
 20 25 30
 Met Thr Ser Pro Gly Ser Gly Leu Ile Leu Gln Ala Asn Phe Val His
 35 40 45
 Ser Gln Arg Arg Glu Ser Phe Leu Tyr Arg Ser Asp Ser Asp Tyr Asp
 50 55 60
 Leu Ser Pro Lys Ser Met Ser Arg Asn Ser Ser Ile Ala Ser Asp Ile
 65 70 75 80
 His Gly Asp Asp Leu Ile Val Thr Pro Phe Ala Gln Val Leu Ala Ser
 85 90 95
 Leu Arg Thr Val Arg Asn Asn Phe Ala Ala Leu Thr Asn Leu Gln Asp
 100 105 110
 Arg Ala Pro Ser Lys Arg Ser Pro Met Cys Asn Gln Pro Ser Ile Asn
 115 120 125
 Lys Ala Thr Ile Thr Glu Glu Ala Tyr Gln Lys Leu Ala Ser Glu Thr
 130 135 140
 Leu Glu Glu Leu Asp Trp Cys Leu Asp Gln Leu Glu Thr Leu Gln Thr
 145 150 155 160
 Arg His Ser Val Ser Glu Met Ala Ser Asn Lys Phe Lys Arg Met Leu
 165 170 175
 Asn Arg Glu Leu Thr His Leu Ser Glu Met Ser Arg Ser Gly Asn Gln
 180 185 190
 Val Ser Glu Phe Ile Ser Asn Thr Phe Leu Asp Lys Gln His Glu Val
 195 200 205
 Glu Ile Pro Ser Pro Thr Gln Lys Glu Lys Glu Lys Lys Lys Arg Pro

210	215	220
Met Ser Gln Ile Ser Gly Val Lys Lys Leu Met His Ser Ser Ser Leu		
225	230	235
Thr Asn Ser Ser Ile Pro Arg Phe Gly Val Lys Thr Glu Gln Glu Asp		240
	245	250
Val Leu Ala Lys Glu Leu Glu Asp Val Asn Lys Trp Gly Leu His Val		255
	260	265
Phe Arg Ile Ala Glu Leu Ser Gly Asn Arg Pro Leu Thr Val Ile Met		270
	275	280
His Thr Ile Phe Gln Glu Arg Asp Leu Leu Lys Thr Phe Lys Ile Pro		285
	290	295
Val Asp Thr Leu Ile Thr Tyr Leu Met Thr Leu Glu Asp His Tyr His		300
305	310	315
Ala Asp Val Ala Tyr His Asn Asn Ile His Ala Ala Asp Val Val Gln		320
	325	330
Ser Thr His Val Leu Leu Ser Thr Pro Ala Leu Glu Ala Val Phe Thr		335
	340	345
Asp Leu Glu Ile Leu Ala Ala Ile Phe Ala Ser Ala Ile His Asp Val		350
	355	360
Asp His Pro Gly Val Ser Asn Gln Phe Leu Ile Asn Thr Asn Ser Glu		365
	370	375
Leu Ala Leu Met Tyr Asn Asp Ser Ser Val Leu Glu Asn His His Leu		380
385	390	395
Ala Val Gly Phe Lys Leu Leu Gln Glu Glu Asn Cys Asp Ile Phe Gln		400
	405	410
Asn Leu Thr Lys Lys Gln Arg Gln Ser Leu Arg Lys Met Val Ile Asp		415
	420	425
Ile Val Leu Ala Thr Asp Met Ser Lys His Met Asn Leu Leu Ala Asp		430
	435	440
Leu Lys Thr Met Val Glu Thr Lys Lys Val Thr Ser Ser Gly Val Leu		445
	450	455
Leu Leu Asp Asn Tyr Ser Asp Arg Ile Gln Val Leu Gln Asn Met Val		460
465	470	475
His Cys Ala Asp Leu Ser Asn Pro Thr Lys Pro Leu Gln Leu Tyr Arg		480
	485	490
Gln Trp Thr Asp Arg Ile Met Glu Glu Phe Phe Arg Gln Gly Asp Arg		495
	500	505
Glu Arg Glu Arg Gly Met Glu Ile Ser Pro Met Cys Asp Lys His Asn		510
	515	520
Ala Ser Val Glu Lys Ser Gln Val Gly Phe Ile Asp Tyr Ile Val His		525
	530	535
Pro Leu Trp Glu Thr Trp Ala Asp Leu Val His Pro Asp Ala Gln Asp		540
545	550	555
Ile Leu Asp Thr Leu Glu Asp Asn Arg Glu Trp Tyr Gln Ser Thr Ile		560
	565	570
Pro Gln Ser Pro Ser Pro Ala Pro Asp Asp Pro Glu Glu Gly Arg Gln		575
	580	585
Gly Gln Thr Glu Lys Phe Gln Phe Glu Leu Thr Leu Glu Glu Asp Gly		590
	595	600
Glu Ser Asp Thr Glu Lys Asp Ser Gly Ser Gln Val Glu Glu Asp Thr		605
	610	615
Ser Cys Ser Asp Ser Lys Thr Leu Cys Thr Gln Asp Ser Glu Ser Thr		620
625	630	635
Glu Ile Pro Leu Asp Glu Gln Val Glu Glu Ala Val Gly Glu Glu		640
	645	650
Glu Glu Ser Gln Pro Glu Ala Cys Val Ile Asp Asp Arg Ser Pro Asp		655
	660	665
Thr Thr Gly Ile Leu Gln Ser Thr Val Pro Arg Ala Arg Asp Pro Pro		670
	675	680
		685

Val Ala Thr Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val
 690 695 700
 Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser
 705 710 715 720
 Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu
 725 730 735
 Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu
 740 745 750
 Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp
 755 760 765
 His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr
 770 775 780
 Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr
 785 790 795 800
 Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu
 805 810 815
 Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys
 820 825 830
 Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys
 835 840 845
 Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu
 850 855 860
 Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile
 865 870 875 880
 Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln
 885 890 895
 Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu
 900 905 910
 Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu
 915 920 925
 Tyr Lys
 930

<210> 3
 <211> 3201
 <212> DNA
 <213> Aequorea victoria and human

<220>
 <221> CDS
 <222> (1)...(3201)

<400> 3
 atg gag gca gag ggc agc agc gcg ccg gcc cgg gcg ggc agc gga gag 48
 Met Glu Ala Glu Gly Ser Ser Ala Pro Ala Arg Ala Gly Ser Gly Glu
 1 5 10 15
 ggc agc gac agc gcc ggc ggg gcc acg ctc aaa gcc ccc aag cat ctc 96
 Gly Ser Asp Ser Ala Gly Gly Ala Thr Leu Lys Ala Pro Lys His Leu
 20 25 30
 tgg agg cac gag cag cac cac cag tac ccg ctc cgg cag ccc cag ttc 144
 Trp Arg His Glu Gln His His Gln Tyr Pro Leu Arg Gln Pro Gln Phe
 35 40 45
 cgc ctc ctg cat ccc cat cac cac ctg ccc ccg ccg ccg cca ccc tcg 192
 Arg Leu Leu His Pro His His His Leu Pro Pro Pro Pro Pro Pro Ser
 50 55 60

ccc cag ccc cag ccc cag tgt ccg cta cag ccg ccg ccg ccg ccc ccc	240
Pro Gln Pro Gln Pro Gln Cys Pro Leu Gln Pro Pro Pro Pro Pro Pro	
65 70 75 80	
ctg ccg ccg ccc ccg ccg ccg ccc ggg gct gcc cgc ggc cgc tac gcc	288
Leu Pro Pro Pro Pro Pro Pro Pro Gly Ala Ala Arg Gly Arg Tyr Ala	
85 90 95	
tcg agc ggg gcc acc ggc cgc gtc cgg cat cgc ggc tac tcg gac acc	336
Ser Ser Gly Ala Thr Gly Arg Val Arg His Arg Gly Tyr Ser Asp Thr	
100 105 110	
gag cgc tac ctg tac tgt cgc gcc atg gac cgc acc tcc tac gcg gtg	384
Glu Arg Tyr Leu Tyr Cys Arg Ala Met Asp Arg Thr Ser Tyr Ala Val	
115 120 125	
gag acc ggc cac cgg ccc ggc ctg aag aaa tcc agg atg tcc tgg ccc	432
Glu Thr Gly His Arg Pro Gly Leu Lys Lys Ser Arg Met Ser Trp Pro	
130 135 140	
tcc tcg ttc cag gga ctc agg cgt ttt gat gtg gac aat ggc aca tct	480
Ser Ser Phe Gln Gly Leu Arg Arg Phe Asp Val Asp Asn Gly Thr Ser	
145 150 155 160	
gcg gga cgg agt ccc ttg gat ccc atg acc agc cca gga tcc ggg cta	528
Ala Gly Arg Ser Pro Leu Asp Pro Met Thr Ser Pro Gly Ser Gly Leu	
165 170 175	
att ctc caa gca aat ttt gtc cac agt caa cga cgg gag tcc ttc ctg	576
Ile Leu Gln Ala Asn Phe Val His Ser Gln Arg Arg Glu Ser Phe Leu	
180 185 190	
tat cga tcc gac agc gat tat gac ctc tct cca aag tct atg tcc cgg	624
Tyr Arg Ser Asp Ser Asp Tyr Asp Leu Ser Pro Lys Ser Met Ser Arg	
195 200 205	
aac tcc tcc att gcc agt gat ata cac gga gat gac ttg att gtg act	672
Asn Ser Ser Ile Ala Ser Asp Ile His Gly Asp Asp Leu Ile Val Thr	
210 215 220	
cca ttt gct cag gtc ttg gcc agt ctg cga act gta cga aac aac ttt	720
Pro Phe Ala Gln Val Leu Ala Ser Leu Arg Thr Val Arg Asn Asn Phe	
225 230 235 240	
gct gca tta act aat ttg caa gat cga gca cct agc aaa aga tca ccc	768
Ala Ala Leu Thr Asn Leu Gln Asp Arg Ala Pro Ser Lys Arg Ser Pro	
245 250 255	
atg tgc aac caa cca tcc atc aac aaa gcc acc ata aca gag gag gcc	816
Met Cys Asn Gln Pro Ser Ile Asn Lys Ala Thr Ile Thr Glu Glu Ala	
260 265 270	
tac cag aaa ctg gcc agc gag acc ctg gag gag ctg gac tgg tgt ctg	864
Tyr Gln Lys Leu Ala Ser Glu Thr Leu Glu Glu Leu Asp Trp Cys Leu	
275 280 285	
gac cag cta gag acc cta cag acc agg cac tcc gtc agt gag atg gcc	912
Asp Gln Leu Glu Thr Leu Gln Thr Arg His Ser Val Ser Glu Met Ala	
290 295 300	

tcc aac aag ttt aaa agg atg ctt aat cgg gag ctc acc cat ctc tct Ser Asn Lys Phe Lys Arg Met Leu Asn Arg Glu Leu Thr His Leu Ser 305 310 315 320	960
gaa atg agt cgg tct gga aat caa gtg tca gag ttt ata tca aac aca Glu Met Ser Arg Ser Gly Asn Gln Val Ser Glu Phe Ile Ser Asn Thr 325 330 335	1008
ttc tta gat aag caa cat gaa gtg gaa att cct tct cca act cag aag Phe Leu Asp Lys Gln His Glu Val Glu Ile Pro Ser Pro Thr Gln Lys 340 345 350	1056
gaa aag gag aaa aag aaa aga cca atg tct cag atc agt gga gtc aag Glu Lys Glu Lys Lys Lys Arg Pro Met Ser Gln Ile Ser Gly Val Lys 355 360 365	1104
aaa ttg atg cac agc tct agt ctg act aat tca agt atc cca agg ttt Lys Leu Met His Ser Ser Ser Leu Thr Asn Ser Ser Ile Pro Arg Phe 370 375 380	1152
gga gtt aaa act gaa caa gaa gat gtc ctt gcc aag gaa cta gaa gat Gly Val Lys Thr Glu Gln Glu Asp Val Leu Ala Lys Glu Leu Glu Asp 385 390 395 400	1200
gtg aac aaa tgg ggt ctt cat gtt ttc aga ata gca gag ttg tct ggt Val Asn Lys Trp Gly Leu His Val Phe Arg Ile Ala Glu Leu Ser Gly 405 410 415	1248
aac cgg ccc ttg act gtt atc atg cac acc att ttt cag gaa cgg gat Asn Arg Pro Leu Thr Val Ile Met His Thr Ile Phe Gln Glu Arg Asp 420 425 430	1296
tta tta aaa aca ttt aaa att cca gta gat act tta att aca tat ctt Leu Leu Lys Thr Phe Lys Ile Pro Val Asp Thr Leu Ile Thr Tyr Leu 435 440 445	1344
atg act ctc gaa gac cat tac cat gct gat gtg gcc tat cac aac aat Met Thr Leu Glu Asp His Tyr His Ala Asp Val Ala Tyr His Asn Asn 450 455 460	1392
atc cat gct gca gat gtt gtc cag tct act cat gtg cta tta tct aca Ile His Ala Ala Asp Val Val Gln Ser Thr His Val Leu Leu Ser Thr 465 470 475 480	1440
cct gct ttg gag gct gtg ttt aca gat ttg gag att ctt gca gca att Pro Ala Leu Glu Ala Val Phe Thr Asp Leu Glu Ile Leu Ala Ala Ile 485 490 495	1488
ttt gcc agt gca ata cat gat gta gat cat cct ggt gtg tcc aat caa Phe Ala Ser Ala Ile His Asp Val Asp His Pro Gly Val Ser Asn Gln 500 505 510	1536
ttt ctg atc aat aca aac tct gaa ctt gcc ttg atg tac aat gat tcc Phe Leu Ile Asn Thr Asn Ser Glu Leu Ala Leu Met Tyr Asn Asp Ser 515 520 525	1584
tca gtc tta gag aac cat cat ttg gct gtg ggc ttt aaa ttg ctt cag Ser Val Leu Glu Asn His His Leu Ala Val Gly Phe Lys Leu Leu Gln 530 535 540 545 550	1632

530	535	540	
gaa gaa aac tgt gac att ttc cag aat ttg acc aaa aaa caa aga caa			1680
Glu Glu Asn Cys Asp Ile Phe Gln Asn Leu Thr Lys Lys Gln Arg Gln			
545	550	555	560
tct tta agg aaa atg gtc att gac atc gta ctt gca aca gat atg tca			1728
Ser Leu Arg Lys Met Val Ile Asp Ile Val Leu Ala Thr Asp Met Ser			
	565	570	575
aaa cac atg aat cta ctg gct gat ttg aag act atg gtt gaa act aag			1776
Lys His Met Asn Leu Leu Ala Asp Leu Lys Thr Met Val Glu Thr Lys			
	580	585	590
aaa gtg aca agc tct gga gtt ctt ctt ctt gat aat tat tcc gat agg			1824
Lys Val Thr Ser Ser Gly Val Leu Leu Leu Asp Asn Tyr Ser Asp Arg			
	595	600	605
att cag gtt ctt cag aat atg gtg cac tgt gca gat ctg agc aac cca			1872
Ile Gln Val Leu Gln Asn Met Val His Cys Ala Asp Leu Ser Asn Pro			
	610	615	620
aca aag cct ctc cag ctg tac cgc cag tgg acg gac cgg ata atg gag			1920
Thr Lys Pro Leu Gln Leu Tyr Arg Gln Trp Thr Asp Arg Ile Met Glu			
	625	630	640
gag ttc ttc cgc caa gga gac cga gag agg gaa cgt ggc atg gag ata			1968
Glu Phe Phe Arg Gln Gly Asp Arg Glu Arg Glu Arg Gly Met Glu Ile			
	645	650	655
agc ccc atg tgt gac aag cac aat gct tcc gtg gaa aaa tca cag gtg			2016
Ser Pro Met Cys Asp Lys His Asn Ala Ser Val Glu Lys Ser Gln Val			
	660	665	670
ggc ttc ata gac tat att gtt cat ccc ctc tgg gag aca tgg gca gac			2064
Gly Phe Ile Asp Tyr Ile Val His Pro Leu Trp Glu Thr Trp Ala Asp			
	675	680	685
ctc gtc cac cct gac gcc cag gat att ttg gac act ttg gag gac aat			2112
Leu Val His Pro Asp Ala Gln Asp Ile Leu Asp Thr Leu Glu Asp Asn			
	690	695	700
cgt gaa tgg tac cag agc aca atc cct cag agc ccc tct cct gca cct			2160
Arg Glu Trp Tyr Gln Ser Thr Ile Pro Gln Ser Pro Ser Pro Ala Pro			
	705	710	720
gat gac cca gag gag ggc cgg cag ggt caa act gag aaa ttc cag ttt			2208
Asp Asp Pro Glu Glu Gly Arg Gln Gly Gln Thr Glu Lys Phe Gln Phe			
	725	730	735
gaa cta act tta gag gaa gat ggt gag tca gac acg gaa aag gac agt			2256
Glu Leu Thr Leu Glu Glu Asp Gly Glu Ser Asp Thr Glu Lys Asp Ser			
	740	745	750
ggc agt caa gtg gaa gaa gac act agc tgc agt gac tcc aag act ctt			2304
Gly Ser Gln Val Glu Glu Asp Thr Ser Cys Ser Asp Ser Lys Thr Leu			
	755	760	765
tgt act caa gac tca gag tct act gaa att ccc ctt gat gaa cag gtt			2352

Cys	Thr	Gln	Asp	Ser	Glu	Ser	Thr	Glu	Ile	Pro	Leu	Asp	Glu	Gln	Val	
770						775					780					
gaa	gag	gag	gca	gta	ggg	gaa	gaa	gag	gaa	agc	cag	cct	gaa	gcc	tgt	2400
Glu	Glu	Glu	Ala	Val	Gly	Glu	Glu	Glu	Glu	Ser	Gln	Pro	Glu	Ala	Cys	
785					790					795					800	
gtc	ata	gat	gat	cgt	tct	cct	gac	acg	acg	gga	att	ctg	cag	tcg	acg	2448
Val	Ile	Asp	Asp	Arg	Ser	Pro	Asp	Thr	Thr	Gly	Ile	Leu	Gln	Ser	Thr	
				805					810					815		
gta	ccg	cgg	gcc	cgg	gat	cca	ccg	gtc	gcc	acc	atg	gtg	agc	aag	ggc	2496
Val	Pro	Arg	Ala	Arg	Asp	Pro	Pro	Val	Ala	Thr	Met	Val	Ser	Lys	Gly	
			820					825					830			
gag	gag	ctg	ttc	acc	ggg	gtg	gtg	ccc	atc	ctg	gtc	gag	ctg	gac	ggc	2544
Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	Val	Glu	Leu	Asp	Gly	
		835					840					845				
gac	gta	aac	ggc	cac	aag	ttc	agc	gtg	tcc	ggc	gag	ggc	gag	ggc	gat	2592
Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	Glu	Gly	Glu	Gly	Asp	
	850					855					860					
gcc	acc	tac	ggc	aag	ctg	acc	ctg	aag	ttc	atc	tgc	acc	acc	ggc	aag	2640
Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	Cys	Thr	Thr	Gly	Lys	
865					870					875					880	
ctg	ccc	gtg	ccc	tgg	ccc	acc	ctc	gtg	acc	acc	ctg	acc	tac	ggc	gtg	2688
Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	Leu	Thr	Tyr	Gly	Val	
				885					890					895		
cag	tgc	ttc	agc	cgc	tac	ccc	gac	cac	atg	aag	cag	cac	gac	ttc	ttc	2736
Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	His	Met	Lys	Gln	His	Asp	Phe	Phe	
			900					905					910			
aag	tcc	gcc	atg	ccc	gaa	ggc	tac	gtc	cag	gag	cgc	acc	atc	ttc	ttc	2784
Lys	Ser	Ala	Met	Pro	Glu	Gly	Tyr	Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	
		915					920					925				
aag	gac	gac	ggc	aac	tac	aag	acc	cgc	gcc	gag	gtg	aag	ttc	gag	ggc	2832
Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	
	930					935					940					
gac	acc	ctg	gtg	aac	cgc	atc	gag	ctg	aag	ggc	atc	gac	ttc	aag	gag	2880
Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	Leu	Lys	Gly	Ile	Asp	Phe	Lys	Glu	
945					950					955					960	
gac	ggc	aac	atc	ctg	ggg	cac	aag	ctg	gag	tac	aac	tac	aac	agc	cac	2928
Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	
				965					970					975		
aac	gtc	tat	atc	atg	gcc	gac	aag	cag	aag	aac	ggc	atc	aag	gtg	aac	2976
Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	
			980					985					990			
ttc	aag	atc	cgc	cac	aac	atc	gag	gac	ggc	agc	gtg	cag	ctc	gcc	gac	3024
Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	Gln	Leu	Ala	Asp	
		995					1000					1005				

cac tac cag cag aac acc ccc atc ggc gac ggc ccc gtg ctg ctg ccc 3072
 His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro
 1010 1015 1020

 gac aac cac tac ctg agc acc cag tcc gcc ctg agc aaa gac ccc aac 3120
 Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn
 1025 1030 1035 1040

 gag aag cgc gat cac atg gtc ctg ctg gag ttc gtg acc gcc gcc ggg 3168
 Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly
 1045 1050 1055

 atc act ctc ggc atg gac gag ctg tac aag taa 3201
 Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys *
 1060 1065

<210> 4
 <211> 1066
 <212> PRT
 <213> Aequorea victoria and human

<400> 4
 Met Glu Ala Glu Gly Ser Ser Ala Pro Ala Arg Ala Gly Ser Gly Glu
 1 5 10 15
 Gly Ser Asp Ser Ala Gly Gly Ala Thr Leu Lys Ala Pro Lys His Leu
 20 25 30
 Trp Arg His Glu Gln His His Gln Tyr Pro Leu Arg Gln Pro Gln Phe
 35 40 45
 Arg Leu Leu His Pro His His His Leu Pro Pro Pro Pro Pro Pro Ser
 50 55 60
 Pro Gln Pro Gln Pro Gln Cys Pro Leu Gln Pro Pro Pro Pro Pro Pro
 65 70 75 80
 Leu Pro Pro Pro Pro Pro Pro Pro Gly Ala Ala Arg Gly Arg Tyr Ala
 85 90 95
 Ser Ser Gly Ala Thr Gly Arg Val Arg His Arg Gly Tyr Ser Asp Thr
 100 105 110
 Glu Arg Tyr Leu Tyr Cys Arg Ala Met Asp Arg Thr Ser Tyr Ala Val
 115 120 125
 Glu Thr Gly His Arg Pro Gly Leu Lys Lys Ser Arg Met Ser Trp Pro
 130 135 140
 Ser Ser Phe Gln Gly Leu Arg Arg Phe Asp Val Asp Asn Gly Thr Ser
 145 150 155 160
 Ala Gly Arg Ser Pro Leu Asp Pro Met Thr Ser Pro Gly Ser Gly Leu
 165 170 175
 Ile Leu Gln Ala Asn Phe Val His Ser Gln Arg Arg Glu Ser Phe Leu
 180 185 190
 Tyr Arg Ser Asp Ser Asp Tyr Asp Leu Ser Pro Lys Ser Met Ser Arg
 195 200 205
 Asn Ser Ser Ile Ala Ser Asp Ile His Gly Asp Asp Leu Ile Val Thr
 210 215 220
 Pro Phe Ala Gln Val Leu Ala Ser Leu Arg Thr Val Arg Asn Asn Phe
 225 230 235 240
 Ala Ala Leu Thr Asn Leu Gln Asp Arg Ala Pro Ser Lys Arg Ser Pro
 245 250 255
 Met Cys Asn Gln Pro Ser Ile Asn Lys Ala Thr Ile Thr Glu Glu Ala
 260 265 270
 Tyr Gln Lys Leu Ala Ser Glu Thr Leu Glu Glu Leu Asp Trp Cys Leu
 275 280 285

Asp Gln Leu Glu Thr Leu Gln Thr Arg His Ser Val Ser Glu Met Ala
 290 295 300
 Ser Asn Lys Phe Lys Arg Met Leu Asn Arg Glu Leu Thr His Leu Ser
 305 310 315 320
 Glu Met Ser Arg Ser Gly Asn Gln Val Ser Glu Phe Ile Ser Asn Thr
 325 330 335
 Phe Leu Asp Lys Gln His Glu Val Glu Ile Pro Ser Pro Thr Gln Lys
 340 345 350
 Glu Lys Glu Lys Lys Lys Arg Pro Met Ser Gln Ile Ser Gly Val Lys
 355 360 365
 Lys Leu Met His Ser Ser Ser Leu Thr Asn Ser Ser Ile Pro Arg Phe
 370 375 380
 Gly Val Lys Thr Glu Gln Glu Asp Val Leu Ala Lys Glu Leu Glu Asp
 385 390 395 400
 Val Asn Lys Trp Gly Leu His Val Phe Arg Ile Ala Glu Leu Ser Gly
 405 410 415
 Asn Arg Pro Leu Thr Val Ile Met His Thr Ile Phe Gln Glu Arg Asp
 420 425 430
 Leu Leu Lys Thr Phe Lys Ile Pro Val Asp Thr Leu Ile Thr Tyr Leu
 435 440 445
 Met Thr Leu Glu Asp His Tyr His Ala Asp Val Ala Tyr His Asn Asn
 450 455 460
 Ile His Ala Ala Asp Val Val Gln Ser Thr His Val Leu Leu Ser Thr
 465 470 475 480
 Pro Ala Leu Glu Ala Val Phe Thr Asp Leu Glu Ile Leu Ala Ala Ile
 485 490 495
 Phe Ala Ser Ala Ile His Asp Val Asp His Pro Gly Val Ser Asn Gln
 500 505 510
 Phe Leu Ile Asn Thr Asn Ser Glu Leu Ala Leu Met Tyr Asn Asp Ser
 515 520 525
 Ser Val Leu Glu Asn His His Leu Ala Val Gly Phe Lys Leu Leu Gln
 530 535 540
 Glu Glu Asn Cys Asp Ile Phe Gln Asn Leu Thr Lys Lys Gln Arg Gln
 545 550 555 560
 Ser Leu Arg Lys Met Val Ile Asp Ile Val Leu Ala Thr Asp Met Ser
 565 570 575
 Lys His Met Asn Leu Leu Ala Asp Leu Lys Thr Met Val Glu Thr Lys
 580 585 590
 Lys Val Thr Ser Ser Gly Val Leu Leu Leu Asp Asn Tyr Ser Asp Arg
 595 600 605
 Ile Gln Val Leu Gln Asn Met Val His Cys Ala Asp Leu Ser Asn Pro
 610 615 620
 Thr Lys Pro Leu Gln Leu Tyr Arg Gln Trp Thr Asp Arg Ile Met Glu
 625 630 635 640
 Glu Phe Phe Arg Gln Gly Asp Arg Glu Arg Glu Arg Gly Met Glu Ile
 645 650 655
 Ser Pro Met Cys Asp Lys His Asn Ala Ser Val Glu Lys Ser Gln Val
 660 665 670
 Gly Phe Ile Asp Tyr Ile Val His Pro Leu Trp Glu Thr Trp Ala Asp
 675 680 685
 Leu Val His Pro Asp Ala Gln Asp Ile Leu Asp Thr Leu Glu Asp Asn
 690 695 700
 Arg Glu Trp Tyr Gln Ser Thr Ile Pro Gln Ser Pro Ser Pro Ala Pro
 705 710 715 720
 Asp Asp Pro Glu Glu Gly Arg Gln Gly Gln Thr Glu Lys Phe Gln Phe
 725 730 735
 Glu Leu Thr Leu Glu Glu Asp Gly Glu Ser Asp Thr Glu Lys Asp Ser
 740 745 750
 Gly Ser Gln Val Glu Glu Asp Thr Ser Cys Ser Asp Ser Lys Thr Leu

755 760 765
 Cys Thr Gln Asp Ser Glu Ser Thr Glu Ile Pro Leu Asp Glu Gln Val
 770 775 780
 Glu Glu Glu Ala Val Gly Glu Glu Glu Glu Ser Gln Pro Glu Ala Cys
 785 790 795 800
 Val Ile Asp Asp Arg Ser Pro Asp Thr Thr Gly Ile Leu Gln Ser Thr
 805 810 815
 Val Pro Arg Ala Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly
 820 825 830
 Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly
 835 840 845
 Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp
 850 855 860
 Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys
 865 870 875 880
 Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val
 885 890 895
 Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe
 900 905 910
 Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe
 915 920 925
 Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly
 930 935 940
 Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu
 945 950 955 960
 Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His
 965 970 975
 Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn
 980 985 990
 Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp
 995 1000 1005
 His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro
 1010 1015 1020
 Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn
 1025 1030 1035 1040
 Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly
 1045 1050 1055
 Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
 1060 1065

<210> 5
 <211> 3009
 <212> DNA
 <213> Aequorea victoria and human

<220>
 <221> CDS
 <222> (1)...(3009)

<400> 5
 atg gct cag cag aca agc ccg gac act tta aca gta cct gaa gtg gat 48
 Met Ala Gln Gln Thr Ser Pro Asp Thr Leu Thr Val Pro Glu Val Asp
 1 5 10 15
 aat ccg cat tgt cca aac ccg tgg ctg aac gaa gac ctt gtg aaa tcc 96
 Asn Pro His Cys Pro Asn Pro Trp Leu Asn Glu Asp Leu Val Lys Ser
 20 25 30
 ttg cga gaa aac ctg ttg cag cat gag aag tcc aag aca gcg agg aaa 144

Leu	Arg	Glu	Asn	Leu	Leu	Gln	His	Glu	Lys	Ser	Lys	Thr	Ala	Arg	Lys	
		35					40					45				
tcg	gtt	tct	ccc	aag	ctc	tct	cca	gtg	atc	tct	ccg	aga	aat	tcc	ccc	192
Ser	Val	Ser	Pro	Lys	Leu	Ser	Pro	Val	Ile	Ser	Pro	Arg	Asn	Ser	Pro	
	50					55					60					
agg	ctt	ctg	cgc	aga	atg	ctt	ctc	agc	agc	aac	atc	ccc	aaa	cag	cgg	240
Arg	Leu	Leu	Arg	Arg	Met	Leu	Leu	Ser	Ser	Asn	Ile	Pro	Lys	Gln	Arg	
65					70					75					80	
cgt	ttc	acg	gtg	gca	cat	aca	tgt	ttt	gat	gtg	gac	aat	ggc	aca	tct	288
Arg	Phe	Thr	Val	Ala	His	Thr	Cys	Phe	Asp	Val	Asp	Asn	Gly	Thr	Ser	
				85					90					95		
gcg	gga	cgg	agt	ccc	ttg	gat	ccc	atg	acc	agc	cca	gga	tcc	ggg	cta	336
Ala	Gly	Arg	Ser	Pro	Leu	Asp	Pro	Met	Thr	Ser	Pro	Gly	Ser	Gly	Leu	
			100					105					110			
att	ctc	caa	gca	aat	ttt	gtc	cac	agt	caa	cga	cgg	gag	tcc	ttc	ctg	384
Ile	Leu	Gln	Ala	Asn	Phe	Val	His	Ser	Gln	Arg	Arg	Glu	Ser	Phe	Leu	
		115					120					125				
tat	cga	tcc	gac	agc	gat	tat	gac	ctc	tct	cca	aag	tct	atg	tcc	cgg	432
Tyr	Arg	Ser	Asp	Ser	Asp	Tyr	Asp	Leu	Ser	Pro	Lys	Ser	Met	Ser	Arg	
	130					135					140					
aac	tcc	tcc	att	gcc	agt	gat	ata	cac	gga	gat	gac	ttg	att	gtg	act	480
Asn	Ser	Ser	Ile	Ala	Ser	Asp	Ile	His	Gly	Asp	Asp	Leu	Ile	Val	Thr	
145					150					155					160	
cca	ttt	gct	cag	gtc	ttg	gcc	agt	ctg	cga	act	gta	cga	aac	aac	ttt	528
Pro	Phe	Ala	Gln	Val	Leu	Ala	Ser	Leu	Arg	Thr	Val	Arg	Asn	Asn	Phe	
				165					170					175		
gct	gca	tta	act	aat	ttg	caa	gat	cga	gca	cct	agc	aaa	aga	tca	ccc	576
Ala	Ala	Leu	Thr	Asn	Leu	Gln	Asp	Arg	Ala	Pro	Ser	Lys	Arg	Ser	Pro	
			180					185					190			
atg	tgc	aac	caa	cca	tcc	atc	aac	aaa	gcc	acc	ata	aca	gag	gag	gcc	624
Met	Cys	Asn	Gln	Pro	Ser	Ile	Asn	Lys	Ala	Thr	Ile	Thr	Glu	Glu	Ala	
		195					200					205				
tac	cag	aaa	ctg	gcc	agc	gag	acc	ctg	gag	gag	ctg	gac	tgg	tgt	ctg	672
Tyr	Gln	Lys	Leu	Ala	Ser	Glu	Thr	Leu	Glu	Glu	Leu	Asp	Trp	Cys	Leu	
	210					215					220					
gac	cag	cta	gag	acc	cta	cag	acc	agg	cac	tcc	gtc	agt	gag	atg	gcc	720
Asp	Gln	Leu	Glu	Thr	Leu	Gln	Thr	Arg	His	Ser	Val	Ser	Glu	Met	Ala	
225					230					235					240	
tcc	aac	aag	ttt	aaa	agg	atg	ctt	aat	cgg	gag	ctc	acc	cat	ctc	tct	768
Ser	Asn	Lys	Phe	Lys	Arg	Met	Leu	Asn	Arg	Glu	Leu	Thr	His	Leu	Ser	
				245					250					255		
gaa	atg	agt	cgg	tct	gga	aat	caa	gtg	tca	gag	ttt	ata	tca	aac	aca	816
Glu	Met	Ser	Arg	Ser	Gly	Asn	Gln	Val	Ser	Glu	Phe	Ile	Ser	Asn	Thr	
			260					265					270			

ttc tta gat aag caa cat gaa gtg gaa att cct tct cca act cag aag	864
Phe Leu Asp Lys Gln His Glu Val Glu Ile Pro Ser Pro Thr Gln Lys	
275 280 285	
gaa aag gag aaa aag aaa aga cca atg tct cag atc agt gga gtc aag	912
Glu Lys Glu Lys Lys Lys Arg Pro Met Ser Gln Ile Ser Gly Val Lys	
290 295 300	
aaa ttg atg cac agc tct agt ctg act aat tca agt atc cca agg ttt	960
Lys Leu Met His Ser Ser Ser Leu Thr Asn Ser Ser Ile Pro Arg Phe	
305 310 315 320	
gga gtt aaa act gaa caa gaa gat gtc ctt gcc aag gaa cta gaa gat	1008
Gly Val Lys Thr Gln Glu Asp Val Leu Ala Lys Glu Leu Glu Asp	
325 330 335	
gtg aac aaa tgg ggt ctt cat gtt ttc aga ata gca gag ttg tct ggt	1056
Val Asn Lys Trp Gly Leu His Val Phe Arg Ile Ala Glu Leu Ser Gly	
340 345 350	
aac cgg ccc ttg act gtt atc atg cac acc att ttt cag gaa cgg gat	1104
Asn Arg Pro Leu Thr Val Ile Met His Thr Ile Phe Gln Glu Arg Asp	
355 360 365	
tta tta aaa aca ttt aaa att cca gta gat act tta att aca tat ctt	1152
Leu Leu Lys Thr Phe Lys Ile Pro Val Asp Thr Leu Ile Thr Tyr Leu	
370 375 380	
atg act ctc gaa gac cat tac cat gct gat gtg gcc tat cac aac aat	1200
Met Thr Leu Glu Asp His Tyr His Ala Asp Val Ala Tyr His Asn Asn	
385 390 395 400	
atc cat gct gca gat gtt gtc cag tct act cat gtg cta tta tct aca	1248
Ile His Ala Ala Asp Val Val Gln Ser Thr His Val Leu Leu Ser Thr	
405 410 415	
cct gct ttg gag gct gtg ttt aca gat ttg gag att ctt gca gca att	1296
Pro Ala Leu Glu Ala Val Phe Thr Asp Leu Glu Ile Leu Ala Ala Ile	
420 425 430	
ttt gcc agt gca ata cat gat gta gat cat cct ggt gtg tcc aat caa	1344
Phe Ala Ser Ala Ile His Asp Val Asp His Pro Gly Val Ser Asn Gln	
435 440 445	
ttt ctg atc aat aca aac tct gaa ctt gcc ttg atg tac aat gat tcc	1392
Phe Leu Ile Asn Thr Asn Ser Glu Leu Ala Leu Met Tyr Asn Asp Ser	
450 455 460	
tca gtc tta gag aac cat cat ttg gct gtg ggc ttt aaa ttg ctt cag	1440
Ser Val Leu Glu Asn His His Leu Ala Val Gly Phe Lys Leu Leu Gln	
465 470 475 480	
gaa gaa aac tgt gac att ttc cag aat ttg acc aaa aaa caa aga caa	1488
Glu Glu Asn Cys Asp Ile Phe Gln Asn Leu Thr Lys Lys Gln Arg Gln	
485 490 495	
tct tta agg aaa atg gtc att gac atc gta ctt gca aca gat atg tca	1536
Ser Leu Arg Lys Met Val Ile Asp Ile Val Leu Ala Thr Asp Met Ser	
500 505 510	

aaa cac atg aat cta ctg gct gat ttg aag act atg gtt gaa act aag	1584
Lys His Met Asn Leu Leu Ala Asp Leu Lys Thr Met Val Glu Thr Lys	
515 520 525	
aaa gtg aca agc tct gga gtt ctt ctt ctt gat aat tat tcc gat agg	1632
Lys Val Thr Ser Ser Gly Val Leu Leu Leu Asp Asn Tyr Ser Asp Arg	
530 535 540	
att cag gtt ctt cag aat atg gtg cac tgt gca gat ctg agc aac cca	1680
Ile Gln Val Leu Gln Asn Met Val His Cys Ala Asp Leu Ser Asn Pro	
545 550 555 560	
aca aag cct ctc cag ctg tac cgc cag tgg acg gac cgg ata atg gag	1728
Thr Lys Pro Leu Gln Leu Tyr Arg Gln Trp Thr Asp Arg Ile Met Glu	
565 570 575	
gag ttc ttc cgc caa gga gac cga gag agg gaa cgt ggc atg gag ata	1776
Glu Phe Phe Arg Gln Gly Asp Arg Glu Arg Glu Arg Gly Met Glu Ile	
580 585 590	
agc ccc atg tgt gac aag cac aat gct tcc gtg gaa aaa tca cag gtg	1824
Ser Pro Met Cys Asp Lys His Asn Ala Ser Val Glu Lys Ser Gln Val	
595 600 605	
ggc ttc ata gac tat att gtt cat ccc ctc tgg gag aca tgg gca gac	1872
Gly Phe Ile Asp Tyr Ile Val His Pro Leu Trp Glu Thr Trp Ala Asp	
610 615 620	
ctc gtc cac cct gac gcc cag gat att ttg gac act ttg gag gac aat	1920
Leu Val His Pro Asp Ala Gln Asp Ile Leu Asp Thr Leu Glu Asp Asn	
625 630 635 640	
cgt gaa tgg tac cag agc aca atc cct cag agc ccc tct cct gca cct	1968
Arg Glu Trp Tyr Gln Ser Thr Ile Pro Gln Ser Pro Ser Pro Ala Pro	
645 650 655	
gat gac cca gag gag ggc cgg cag ggt caa act gag aaa ttc cag ttt	2016
Asp Asp Pro Glu Glu Gly Arg Gln Gly Gln Thr Glu Lys Phe Gln Phe	
660 665 670	
gaa cta act tta gag gaa gat ggt gag tca gac acg gaa aag gac agt	2064
Glu Leu Thr Leu Glu Glu Asp Gly Glu Ser Asp Thr Glu Lys Asp Ser	
675 680 685	
ggc agt caa gtg gaa gaa gac act agc tgc agt gac tcc aag act ctt	2112
Gly Ser Gln Val Glu Glu Asp Thr Ser Cys Ser Asp Ser Lys Thr Leu	
690 695 700	
tgt act caa gac tca gag tct act gaa att ccc ctt gat gaa cag gtt	2160
Cys Thr Gln Asp Ser Glu Ser Thr Glu Ile Pro Leu Asp Glu Gln Val	
705 710 715 720	
gaa gag gag gca gta ggg gaa gaa gag gaa agc cag cct gaa gcc tgt	2208
Glu Glu Glu Ala Val Gly Glu Glu Glu Glu Ser Gln Pro Glu Ala Cys	
725 730 735	
gtc ata gat gat cgt tct cct gac acg acg gga att ctg cag tcg acg	2256
Val Ile Asp Asp Arg Ser Pro Asp Thr Thr Gly Ile Leu Gln Ser Thr	

740	745	750	
gta ccg cgg gcc cgg gat cca ccg gtc gcc acc atg gtg agc aag ggc Val Pro Arg Ala Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly 755 760 765			2304
gag gag ctg ttc acc ggg gtg gtg ccc atc ctg gtc gag ctg gac ggc Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly 770 775 780			2352
gac gta aac ggc cac aag ttc agc gtg tcc ggc gag ggc gag ggc gat Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp 785 790 795 800			2400
gcc acc tac ggc aag ctg acc ctg aag ttc atc tgc acc acc ggc aag Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys 805 810 815			2448
ctg ccc gtg ccc tgg ccc acc ctg gtg acc acc ctg acc tac ggc gtg Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val 820 825 830			2496
cag tgc ttc agc cgc tac ccc gac cac atg aag cag cac gac ttc ttc Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe 835 840 845			2544
aag tcc gcc atg ccc gaa ggc tac gtc cag gag cgc acc atc ttc ttc Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe 850 855 860			2592
aag gac gac ggc aac tac aag acc cgc gcc gag gtg aag ttc gag ggc Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly 865 870 875 880			2640
gac acc ctg gtg aac cgc atc gag ctg aag ggc atc gac ttc aag gag Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu 885 890 895			2688
gac ggc aac atc ctg ggg cac aag ctg gag tac aac tac aac agc cac Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His 900 905 910			2736
aac gtc tat atc atg gcc gac aag cag aag aac ggc atc aag gtg aac Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn 915 920 925			2784
ttc aag atc cgc cac aac atc gag gac ggc agc gtg cag ctg gcc gac Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp 930 935 940			2832
cac tac cag cag aac acc ccc atc ggc gac ggc ccc gtg ctg ctg ccc His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro 945 950 955 960			2880
gac aac cac tac ctg agc acc cag tcc gcc ctg agc aaa gac ccc aac Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn 965 970 975			2928
gag aag cgc gat cac atg gtc ctg ctg gag ttc gtg acc gcc gcc ggc			2976

Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly
 980 985 990

atc act ctc ggc atg gac gag ctg tac aag taa
 Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys *
 995 1000

3009

<210> 6
 <211> 1002
 <212> PRT
 <213> Aequorea victoria and human

<400> 6
 Met Ala Gln Gln Thr Ser Pro Asp Thr Leu Thr Val Pro Glu Val Asp
 1 5 10 15
 Asn Pro His Cys Pro Asn Pro Trp Leu Asn Glu Asp Leu Val Lys Ser
 20 25 30
 Leu Arg Glu Asn Leu Leu Gln His Glu Lys Ser Lys Thr Ala Arg Lys
 35 40 45
 Ser Val Ser Pro Lys Leu Ser Pro Val Ile Ser Pro Arg Asn Ser Pro
 50 55 60
 Arg Leu Leu Arg Arg Met Leu Leu Ser Ser Asn Ile Pro Lys Gln Arg
 65 70 75 80
 Arg Phe Thr Val Ala His Thr Cys Phe Asp Val Asp Asn Gly Thr Ser
 85 90 95
 Ala Gly Arg Ser Pro Leu Asp Pro Met Thr Ser Pro Gly Ser Gly Leu
 100 105 110
 Ile Leu Gln Ala Asn Phe Val His Ser Gln Arg Arg Glu Ser Phe Leu
 115 120 125
 Tyr Arg Ser Asp Ser Asp Tyr Asp Leu Ser Pro Lys Ser Met Ser Arg
 130 135 140
 Asn Ser Ser Ile Ala Ser Asp Ile His Gly Asp Asp Leu Ile Val Thr
 145 150 155 160
 Pro Phe Ala Gln Val Leu Ala Ser Leu Arg Thr Val Arg Asn Asn Phe
 165 170 175
 Ala Ala Leu Thr Asn Leu Gln Asp Arg Ala Pro Ser Lys Arg Ser Pro
 180 185 190
 Met Cys Asn Gln Pro Ser Ile Asn Lys Ala Thr Ile Thr Glu Glu Ala
 195 200 205
 Tyr Gln Lys Leu Ala Ser Glu Thr Leu Glu Glu Leu Asp Trp Cys Leu
 210 215 220
 Asp Gln Leu Glu Thr Leu Gln Thr Arg His Ser Val Ser Glu Met Ala
 225 230 235 240
 Ser Asn Lys Phe Lys Arg Met Leu Asn Arg Glu Leu Thr His Leu Ser
 245 250 255
 Glu Met Ser Arg Ser Gly Asn Gln Val Ser Glu Phe Ile Ser Asn Thr
 260 265 270
 Phe Leu Asp Lys Gln His Glu Val Glu Ile Pro Ser Pro Thr Gln Lys
 275 280 285
 Glu Lys Glu Lys Lys Lys Arg Pro Met Ser Gln Ile Ser Gly Val Lys
 290 295 300
 Lys Leu Met His Ser Ser Ser Leu Thr Asn Ser Ser Ile Pro Arg Phe
 305 310 315 320
 Gly Val Lys Thr Glu Gln Glu Asp Val Leu Ala Lys Glu Leu Glu Asp
 325 330 335
 Val Asn Lys Trp Gly Leu His Val Phe Arg Ile Ala Glu Leu Ser Gly
 340 345 350
 Asn Arg Pro Leu Thr Val Ile Met His Thr Ile Phe Gln Glu Arg Asp

355	360	365
Leu Leu Lys Thr Phe Lys Ile Pro Val Asp Thr Leu Ile Thr Tyr Leu		
370	375	380
Met Thr Leu Glu Asp His Tyr His Ala Asp Val Ala Tyr His Asn Asn		
385	390	395
Ile His Ala Ala Asp Val Val Gln Ser Thr His Val Leu Leu Ser Thr		400
405	410	415
Pro Ala Leu Glu Ala Val Phe Thr Asp Leu Glu Ile Leu Ala Ala Ile		
420	425	430
Phe Ala Ser Ala Ile His Asp Val Asp His Pro Gly Val Ser Asn Gln		
435	440	445
Phe Leu Ile Asn Thr Asn Ser Glu Leu Ala Leu Met Tyr Asn Asp Ser		
450	455	460
Ser Val Leu Glu Asn His Leu Ala Val Gly Phe Lys Leu Leu Gln		
465	470	475
Glu Glu Asn Cys Asp Ile Phe Gln Asn Leu Thr Lys Lys Gln Arg Gln		480
485	490	495
Ser Leu Arg Lys Met Val Ile Asp Ile Val Leu Ala Thr Asp Met Ser		
500	505	510
Lys His Met Asn Leu Leu Ala Asp Leu Lys Thr Met Val Glu Thr Lys		
515	520	525
Lys Val Thr Ser Ser Gly Val Leu Leu Leu Asp Asn Tyr Ser Asp Arg		
530	535	540
Ile Gln Val Leu Gln Asn Met Val His Cys Ala Asp Leu Ser Asn Pro		
545	550	555
Thr Lys Pro Leu Gln Leu Tyr Arg Gln Trp Thr Asp Arg Ile Met Glu		560
565	570	575
Glu Phe Phe Arg Gln Gly Asp Arg Glu Arg Glu Arg Gly Met Glu Ile		
580	585	590
Ser Pro Met Cys Asp Lys His Asn Ala Ser Val Glu Lys Ser Gln Val		
595	600	605
Gly Phe Ile Asp Tyr Ile Val His Pro Leu Trp Glu Thr Trp Ala Asp		
610	615	620
Leu Val His Pro Asp Ala Gln Asp Ile Leu Asp Thr Leu Glu Asp Asn		
625	630	635
Arg Glu Trp Tyr Gln Ser Thr Ile Pro Gln Ser Pro Ser Pro Ala Pro		640
645	650	655
Asp Asp Pro Glu Glu Gly Arg Gln Gly Gln Thr Glu Lys Phe Gln Phe		
660	665	670
Glu Leu Thr Leu Glu Glu Asp Gly Glu Ser Asp Thr Glu Lys Asp Ser		
675	680	685
Gly Ser Gln Val Glu Glu Asp Thr Ser Cys Ser Asp Ser Lys Thr Leu		
690	695	700
Cys Thr Gln Asp Ser Glu Ser Thr Glu Ile Pro Leu Asp Glu Gln Val		
705	710	715
Glu Glu Glu Ala Val Gly Glu Glu Glu Glu Ser Gln Pro Glu Ala Cys		720
725	730	735
Val Ile Asp Asp Arg Ser Pro Asp Thr Thr Gly Ile Leu Gln Ser Thr		
740	745	750
Val Pro Arg Ala Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly		
755	760	765
Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly		
770	775	780
Asp Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp		
785	790	795
Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys		800
805	810	815
Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val		
820	825	830

Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe
835 840 845
Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe
850 855 860
Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly
865 870 875 880
Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu
885 890 895
Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His
900 905 910
Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn
915 920 925
Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp
930 935 940
His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro
945 950 955 960
Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn
965 970 975
Glu Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly
980 985 990
Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
995 1000

<210> 7
<211> 3381
<212> DNA
<213> *Aequorea victoria*

<220>
<221> CDS
<222> (1)...(3381)

<400> 7

atg gag cgg gcc ggc ccc agc ttc ggg cag cag cga cag cag cag cag	48
Met Glu Arg Ala Gly Pro Ser Phe Gly Gln Gln Arg Gln Gln Gln Gln	
1 5 10 15	
ccc cag cag cag aag cag cag cag agg gat cag gac tcg gtc gaa gca	96
Pro Gln Gln Gln Lys Gln Gln Gln Arg Asp Gln Asp Ser Val Glu Ala	
20 25 30	
tgg ctg gac gat cac tgg gac ttt acc ttc tca tac ttt gtt aga aaa	144
Trp Leu Asp Asp His Trp Asp Phe Thr Phe Ser Tyr Phe Val Arg Lys	
35 40 45	
gcc acc aga gaa atg gtc aat gca tgg ttt gct gag aga gtt cac acc	192
Ala Thr Arg Glu Met Val Asn Ala Trp Phe Ala Glu Arg Val His Thr	
50 55 60	
atc cct gtg tgc aag gaa ggt atc aga ggc cac acc gaa tct tgc tct	240
Ile Pro Val Cys Lys Glu Gly Ile Arg Gly His Thr Glu Ser Cys Ser	
65 70 75 80	
tgt ccc ttg cag cag agt cct cgt gca gat aac agt gtc cct gga aca	288
Cys Pro Leu Gln Gln Ser Pro Arg Ala Asp Asn Ser Val Pro Gly Thr	
85 90 95	
cca acc agg aaa atc tct gcc tct gaa ttt gac cgg cct ctt aga ccc	336
Pro Thr Arg Lys Ile Ser Ala Ser Glu Phe Asp Arg Pro Leu Arg Pro	

100	105	110	
att gtt gtc aag gat tct gag gga act gtg agc ttc ctc tct gac tca			384
Ile Val Val Lys Asp Ser Glu Gly Thr Val Ser Phe Leu Ser Asp Ser			
115	120	125	
gaa aag aag gaa cag atg cct cta acc cct cca agg ttt gat cat gat			432
Glu Lys Lys Glu Gln Met Pro Leu Thr Pro Pro Arg Phe Asp His Asp			
130	135	140	
gaa ggg gac cag tgc tca aga ctc ttg gaa tta gtg aag gat att tct			480
Glu Gly Asp Gln Cys Ser Arg Leu Leu Glu Leu Val Lys Asp Ile Ser			
145	150	155	160
agt cat ttg gat gtc aca gcc tta tgt cac aaa att ttc ttg cat atc			528
Ser His Leu Asp Val Thr Ala Leu Cys His Lys Ile Phe Leu His Ile			
165	170	175	
cat gga ctg ata tct gct gac cgc tat tcc ctg ttc ctt gtc tgt gaa			576
His Gly Leu Ile Ser Ala Asp Arg Tyr Ser Leu Phe Leu Val Cys Glu			
180	185	190	
gac agc tcc aat gac aag ttt ctt atc agc cgc ctc ttt gat gtt gct			624
Asp Ser Ser Asn Asp Lys Phe Leu Ile Ser Arg Leu Phe Asp Val Ala			
195	200	205	
gaa ggt tca aca ctg gaa gaa gtt tca aat aac tgt atc cgc tta gaa			672
Glu Gly Ser Thr Leu Glu Glu Val Ser Asn Asn Cys Ile Arg Leu Glu			
210	215	220	
tgg aac aaa ggc att gtg gga cat gtg gca gcg ctt ggt gag ccc ttg			720
Trp Asn Lys Gly Ile Val Gly His Val Ala Ala Leu Gly Glu Pro Leu			
225	230	235	240
aac atc aaa gat gca tat gag gat cct cgg ttc aat gca gaa gtt gac			768
Asn Ile Lys Asp Ala Tyr Glu Asp Pro Arg Phe Asn Ala Glu Val Asp			
245	250	255	
caa att aca ggc tac aag aca caa agc att ctt tgt atg cca att aag			816
Gln Ile Thr Gly Tyr Lys Thr Gln Ser Ile Leu Cys Met Pro Ile Lys			
260	265	270	
aat cat agg gaa gag gtt gtt ggt gta gcc cag gcc atc aac aag aaa			864
Asn His Arg Glu Glu Val Val Gly Val Ala Gln Ala Ile Asn Lys Lys			
275	280	285	
tca gga aac ggt ggg aca ttt act gaa aaa gat gaa aag gac ttt gct			912
Ser Gly Asn Gly Gly Thr Phe Thr Glu Lys Asp Glu Lys Asp Phe Ala			
290	295	300	
gct tat ttg gca ttt tgt ggt att gtt ctt cat aat gct cag ctc tat			960
Ala Tyr Leu Ala Phe Cys Gly Ile Val Leu His Asn Ala Gln Leu Tyr			
305	310	315	320
gag act tca ctg ctg gag aac aag aga aat cag gtg ctg ctt gac ctt			1008
Glu Thr Ser Leu Glu Glu Asn Lys Arg Asn Gln Val Leu Leu Asp Leu			
325	330	335	
gct agt tta att ttt gaa gaa caa caa tca tta gaa gta att ttg aag			1056

Ala Ser Leu Ile Phe Glu Glu Gln Gln Ser Leu Glu Val Ile Leu Lys	
340	345 350
aaa ata gct gcc act att atc tct ttc atg caa gtg cag aaa tgc acc	1104
Lys Ile Ala Ala Thr Ile Ile Ser Phe Met Gln Val Gln Lys Cys Thr	
355	360 365
att ttc ata gtg gat gaa gat tgc tcc gat tct ttt tct agt gtg ttt	1152
Ile Phe Ile Val Asp Glu Asp Cys Ser Asp Ser Phe Ser Ser Val Phe	
370	375 380
cac atg gag tgt gag gaa tta gaa aaa tca tct gat aca tta aca agg	1200
His Met Glu Cys Glu Glu Leu Glu Lys Ser Ser Asp Thr Leu Thr Arg	
385	390 395 400
gaa cat gat gca aac aaa atc aat tac atg tat gct cag tat gtc aaa	1248
Glu His Asp Ala Asn Lys Ile Asn Tyr Met Tyr Ala Gln Tyr Val Lys	
405	410 415
aat act atg gaa cca ctt aat atc cca gat gtc agt aag gat aaa aga	1296
Asn Thr Met Glu Pro Leu Asn Ile Pro Asp Val Ser Lys Asp Lys Arg	
420	425 430
ttt ccc tgg aca act gaa aat aca gga aat gta aac cag cag tgc att	1344
Phe Pro Trp Thr Thr Glu Asn Thr Gly Asn Val Asn Gln Gln Cys Ile	
435	440 445
aga agt ttg ctt tgt aca cct ata aaa aat gga aag aag aat aaa gtt	1392
Arg Ser Leu Leu Cys Thr Pro Ile Lys Asn Gly Lys Lys Asn Lys Val	
450	455 460
ata ggg gtt tgc caa ctt gtt aat aag atg gag gag aat act ggc aag	1440
Ile Gly Val Cys Gln Leu Val Asn Lys Met Glu Glu Asn Thr Gly Lys	
465	470 475 480
gtt aag cct ttc aac cga aat gac gaa cag ttt ctg gaa gct ttt gtc	1488
Val Lys Pro Phe Asn Arg Asn Asp Glu Gln Phe Leu Glu Ala Phe Val	
485	490 495
atc ttt tgt ggc ttg ggg atc cag aac acg cag atg tat gaa gca gtg	1536
Ile Phe Cys Gly Leu Gly Ile Gln Asn Thr Gln Met Tyr Glu Ala Val	
500	505 510
gag aga gcc atg gcc aag caa atg gtc aca ttg gag gtt ctg tcg tat	1584
Glu Arg Ala Met Ala Lys Gln Met Val Thr Leu Glu Val Leu Ser Tyr	
515	520 525
cat gct tca gca gca gag gaa gaa aca aga gag cta cag tcg tta gcg	1632
His Ala Ser Ala Ala Glu Glu Glu Thr Arg Glu Leu Gln Ser Leu Ala	
530	535 540
gct gct gtg gtg cca tct gcc cag acc ctt aaa att act gac ttt agc	1680
Ala Ala Val Val Pro Ser Ala Gln Thr Leu Lys Ile Thr Asp Phe Ser	
545	550 555 560
ttc agt gac ttt gag ctg tct gat ctg gaa aca gca ctg tgc aca att	1728
Phe Ser Asp Phe Glu Leu Ser Asp Leu Glu Thr Ala Leu Cys Thr Ile	
565	570 575

24

cgg atg ttt act gac ctc aac ctt gtg cag aac ttc cag atg aaa cat	1776
Arg Met Phe Thr Asp Leu Asn Leu Val Gln Asn Phe Gln Met Lys His	
580 585 590	
gag gtt ctt tgc aga tgg att tta agt gtt aag aag aat tat cgg aag	1824
Glu Val Leu Cys Arg Trp Ile Leu Ser Val Lys Lys Asn Tyr Arg Lys	
595 600 605	
aat gtt gcc tat cat aat tgg aga cat gcc ttt aat aca gct cag tgc	1872
Asn Val Ala Tyr His Asn Trp Arg His Ala Phe Asn Thr Ala Gln Cys	
610 615 620	
atg ttt gct gct cta aaa gca ggc aaa att cag aac aag ctg act gac	1920
Met Phe Ala Ala Leu Lys Ala Gly Lys Ile Gln Asn Lys Leu Thr Asp	
625 630 635 640	
ctg gag ata ctt gca ttg ctg att gct gca cta agc cac gat ttg gat	1968
Leu Glu Ile Leu Ala Leu Leu Ile Ala Ala Leu Ser His Asp Leu Asp	
645 650 655	
cac cgt ggt gtg aat aac tct tac ata cag cga agt gaa cat cca ctt	2016
His Arg Gly Val Asn Asn Ser Tyr Ile Gln Arg Ser Glu His Pro Leu	
660 665 670	
gcc cag ctt tac tgc cat tca atc atg gaa cac cat cat ttt gac cag	2064
Ala Gln Leu Tyr Cys His Ser Ile Met Glu His His His Phe Asp Gln	
675 680 685	
tgc ctg atg att ctt aat agt cca ggc aat cag att ctc agt ggc ctc	2112
Cys Leu Met Ile Leu Asn Ser Pro Gly Asn Gln Ile Leu Ser Gly Leu	
690 695 700	
tcc att gaa gaa tat aag acc acg ttg aaa ata atc aag caa gct att	2160
Ser Ile Glu Glu Tyr Lys Thr Thr Leu Lys Ile Ile Lys Gln Ala Ile	
705 710 715 720	
tta gct aca gac cta gca ctg tac att aag agg cga gga gaa ttt ttt	2208
Leu Ala Thr Asp Leu Ala Leu Tyr Ile Lys Arg Arg Gly Glu Phe Phe	
725 730 735	
gaa ctt ata aga aaa aat caa ttc aat ttg gaa gat cct cat caa aag	2256
Glu Leu Ile Arg Lys Asn Gln Phe Asn Leu Glu Asp Pro His Gln Lys	
740 745 750	
gag ttg ttt ttg gca atg ctg atg aca gct tgt gat ctt tct gca att	2304
Glu Leu Phe Leu Ala Met Leu Met Thr Ala Cys Asp Leu Ser Ala Ile	
755 760 765	
aca aaa ccc tgg cct att caa caa cgg ata gca gaa ctt gta gca act	2352
Thr Lys Pro Trp Pro Ile Gln Gln Arg Ile Ala Glu Leu Val Ala Thr	
770 775 780	
gaa ttt ttt gat caa gga gac aga gag aga aaa gaa ctc aac ata gaa	2400
Glu Phe Phe Asp Gln Gly Asp Arg Glu Arg Lys Glu Leu Asn Ile Glu	
785 790 795 800	
ccc act gat cta atg aac agg gag aag aaa aac aaa atc cca agt atg	2448
Pro Thr Asp Leu Met Asn Arg Glu Lys Lys Asn Lys Ile Pro Ser Met	
805 810 815	

caa gtt ggg ttc ata gat gcc atc tgc ttg caa ctg tat gag gcc ctg	2496
Gln Val Gly Phe Ile Asp Ala Ile Cys Leu Gln Leu Tyr Glu Ala Leu	
820 825 830	
acc cac gtg tca gag gac tgt ttc cct ttg cta gat ggc tgc aga aag	2544
Thr His Val Ser Glu Asp Cys Phe Pro Leu Leu Asp Gly Cys Arg Lys	
835 840 845	
aac agg cag aaa tgg cag gcc ctt gca gaa cag cag gag aag atg ctg	2592
Asn Arg Gln Lys Trp Gln Ala Leu Ala Glu Gln Gln Glu Lys Met Leu	
850 855 860	
att aat ggg gaa agc ggc cag gcc aag cgg aac tgg gta ccg cgg gcc	2640
Ile Asn Gly Glu Ser Gly Gln Ala Lys Arg Asn Trp Val Pro Arg Ala	
865 870 875 880	
cgg gat cca ccg gtc gcc acc atg gtg agc aag ggc gag gag ctg ttc	2688
Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly Glu Glu Leu Phe	
885 890 895	
acc ggg gtg gtg ccc atc ctg gtc gag ctg gac ggc gac gta aac ggc	2736
Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly	
900 905 910	
cac aag ttc agc gtg tcc ggc gag ggc gag ggc gat gcc acc tac ggc	2784
His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly	
915 920 925	
aag ctg acc ctg aag ttc atc tgc acc acc ggc aag ctg ccc gtg ccc	2832
Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro	
930 935 940	
tgg ccc acc ctc gtg acc acc ctg acc tac ggc gtg cag tgc ttc agc	2880
Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser	
945 950 955 960	
cgc tac ccc gac cac atg aag cag cac gac ttc ttc aag tcc gcc atg	2928
Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met	
965 970 975	
ccc gaa ggc tac gtc cag gag cgc acc atc ttc ttc aag gac gac ggc	2976
Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly	
980 985 990	
aac tac aag acc cgc gcc gag gtg aag ttc gag ggc gac acc ctg gtg	3024
Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val	
995 1000 1005	
aac cgc atc gag ctg aag ggc atc gac ttc aag gag gac ggc aac atc	3072
Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile	
1010 1015 1020	
ctg ggg cac aag ctg gag tac aac tac aac agc cac aac gtc tat atc	3120
Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile	
1025 1030 1035 1040	
atg gcc gac aag cag aag aac ggc atc aag gtg aac ttc aag atc cgc	3168
Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg	

1045	1050	1055	
cac aac atc gag gac ggc agc gtg cag ctc gcc gac cac tac cag cag			3216
His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln			
1060	1065	1070	
aac acc ccc atc ggc gac ggc ccc gtg ctg ctg ccc gac aac cac tac			3264
Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr			
1075	1080	1085	
ctg agc acc cag tcc gcc ctg agc aaa gac ccc aac gag aag cgc gat			3312
Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp			
1090	1095	1100	
cac atg gtc ctg ctg gag ttc gtg acc gcc gcc ggg atc act ctc ggc			3360
His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly			
1105	1110	1115	1120
atg gac gag ctg tac aag taa			3381
Met Asp Glu Leu Tyr Lys *			
1125			

<210> 8
 <211> 1126
 <212> PRT
 <213> Aequorea victoria and human

<400> 8

Met	Glu	Arg	Ala	Gly	Pro	Ser	Phe	Gly	Gln	Gln	Arg	Gln	Gln	Gln	Gln
1				5				10					15		
Pro	Gln	Gln	Gln	Lys	Gln	Gln	Gln	Arg	Asp	Gln	Asp	Ser	Val	Glu	Ala
			20					25					30		
Trp	Leu	Asp	Asp	His	Trp	Asp	Phe	Thr	Phe	Ser	Tyr	Phe	Val	Arg	Lys
		35					40					45			
Ala	Thr	Arg	Glu	Met	Val	Asn	Ala	Trp	Phe	Ala	Glu	Arg	Val	His	Thr
	50					55				60					
Ile	Pro	Val	Cys	Lys	Glu	Gly	Ile	Arg	Gly	His	Thr	Glu	Ser	Cys	Ser
65				70					75					80	
Cys	Pro	Leu	Gln	Gln	Ser	Pro	Arg	Ala	Asp	Asn	Ser	Val	Pro	Gly	Thr
			85						90					95	
Pro	Thr	Arg	Lys	Ile	Ser	Ala	Ser	Glu	Phe	Asp	Arg	Pro	Leu	Arg	Pro
			100					105					110		
Ile	Val	Val	Lys	Asp	Ser	Glu	Gly	Thr	Val	Ser	Phe	Leu	Ser	Asp	Ser
		115					120					125			
Glu	Lys	Lys	Glu	Gln	Met	Pro	Leu	Thr	Pro	Pro	Arg	Phe	Asp	His	Asp
	130					135					140				
Glu	Gly	Asp	Gln	Cys	Ser	Arg	Leu	Leu	Glu	Leu	Val	Lys	Asp	Ile	Ser
145				150					155					160	
Ser	His	Leu	Asp	Val	Thr	Ala	Leu	Cys	His	Lys	Ile	Phe	Leu	His	Ile
			165						170					175	
His	Gly	Leu	Ile	Ser	Ala	Asp	Arg	Tyr	Ser	Leu	Phe	Leu	Val	Cys	Glu
		180						185					190		
Asp	Ser	Ser	Asn	Asp	Lys	Phe	Leu	Ile	Ser	Arg	Leu	Phe	Asp	Val	Ala
	195					200						205			
Glu	Gly	Ser	Thr	Leu	Glu	Glu	Val	Ser	Asn	Asn	Cys	Ile	Arg	Leu	Glu
	210					215					220				
Trp	Asn	Lys	Gly	Ile	Val	Gly	His	Val	Ala	Ala	Leu	Gly	Glu	Pro	Leu
225				230					235					240	

Asn Ile Lys Asp Ala Tyr Glu Asp Pro Arg Phe Asn Ala Glu Val Asp
 245 250 255
 Gln Ile Thr Gly Tyr Lys Thr Gln Ser Ile Leu Cys Met Pro Ile Lys
 260 265 270
 Asn His Arg Glu Glu Val Val Gly Val Ala Gln Ala Ile Asn Lys Lys
 275 280 285
 Ser Gly Asn Gly Gly Thr Phe Thr Glu Lys Asp Glu Lys Asp Phe Ala
 290 295 300
 Ala Tyr Leu Ala Phe Cys Gly Ile Val Leu His Asn Ala Gln Leu Tyr
 305 310 315 320
 Glu Thr Ser Leu Leu Glu Asn Lys Arg Asn Gln Val Leu Leu Asp Leu
 325 330 335
 Ala Ser Leu Ile Phe Glu Glu Gln Gln Ser Leu Glu Val Ile Leu Lys
 340 345 350
 Lys Ile Ala Ala Thr Ile Ile Ser Phe Met Gln Val Gln Lys Cys Thr
 355 360 365
 Ile Phe Ile Val Asp Glu Asp Cys Ser Asp Ser Phe Ser Ser Val Phe
 370 375 380
 His Met Glu Cys Glu Glu Leu Glu Lys Ser Ser Asp Thr Leu Thr Arg
 385 390 395 400
 Glu His Asp Ala Asn Lys Ile Asn Tyr Met Tyr Ala Gln Tyr Val Lys
 405 410 415
 Asn Thr Met Glu Pro Leu Asn Ile Pro Asp Val Ser Lys Asp Lys Arg
 420 425 430
 Phe Pro Trp Thr Thr Glu Asn Thr Gly Asn Val Asn Gln Gln Cys Ile
 435 440 445
 Arg Ser Leu Leu Cys Thr Pro Ile Lys Asn Gly Lys Lys Asn Lys Val
 450 455 460
 Ile Gly Val Cys Gln Leu Val Asn Lys Met Glu Glu Asn Thr Gly Lys
 465 470 475 480
 Val Lys Pro Phe Asn Arg Asn Asp Glu Gln Phe Leu Glu Ala Phe Val
 485 490 495
 Ile Phe Cys Gly Leu Gly Ile Gln Asn Thr Gln Met Tyr Glu Ala Val
 500 505 510
 Glu Arg Ala Met Ala Lys Gln Met Val Thr Leu Glu Val Leu Ser Tyr
 515 520 525
 His Ala Ser Ala Ala Glu Glu Glu Thr Arg Glu Leu Gln Ser Leu Ala
 530 535 540
 Ala Ala Val Val Pro Ser Ala Gln Thr Leu Lys Ile Thr Asp Phe Ser
 545 550 555 560
 Phe Ser Asp Phe Glu Leu Ser Asp Leu Glu Thr Ala Leu Cys Thr Ile
 565 570 575
 Arg Met Phe Thr Asp Leu Asn Leu Val Gln Asn Phe Gln Met Lys His
 580 585 590
 Glu Val Leu Cys Arg Trp Ile Leu Ser Val Lys Lys Asn Tyr Arg Lys
 595 600 605
 Asn Val Ala Tyr His Asn Trp Arg His Ala Phe Asn Thr Ala Gln Cys
 610 615 620
 Met Phe Ala Ala Leu Lys Ala Gly Lys Ile Gln Asn Lys Leu Thr Asp
 625 630 635 640
 Leu Glu Ile Leu Ala Leu Leu Ile Ala Ala Leu Ser His Asp Leu Asp
 645 650 655
 His Arg Gly Val Asn Asn Ser Tyr Ile Gln Arg Ser Glu His Pro Leu
 660 665 670
 Ala Gln Leu Tyr Cys His Ser Ile Met Glu His His His Phe Asp Gln
 675 680 685
 Cys Leu Met Ile Leu Asn Ser Pro Gly Asn Gln Ile Leu Ser Gly Leu
 690 695 700
 Ser Ile Glu Glu Tyr Lys Thr Thr Leu Lys Ile Ile Lys Gln Ala Ile

```

705          710          715          720
Leu Ala Thr Asp Leu Ala Leu Tyr Ile Lys Arg Arg Gly Glu Phe Phe
          725          730          735
Glu Leu Ile Arg Lys Asn Gln Phe Asn Leu Glu Asp Pro His Gln Lys
          740          745          750
Glu Leu Phe Leu Ala Met Leu Met Thr Ala Cys Asp Leu Ser Ala Ile
          755          760          765
Thr Lys Pro Trp Pro Ile Gln Gln Arg Ile Ala Glu Leu Val Ala Thr
          770          775          780
Glu Phe Phe Asp Gln Gly Asp Arg Glu Arg Lys Glu Leu Asn Ile Glu
785          790          795          800
Pro Thr Asp Leu Met Asn Arg Glu Lys Lys Asn Lys Ile Pro Ser Met
          805          810          815
Gln Val Gly Phe Ile Asp Ala Ile Cys Leu Gln Leu Tyr Glu Ala Leu
          820          825          830
Thr His Val Ser Glu Asp Cys Phe Pro Leu Leu Asp Gly Cys Arg Lys
          835          840          845
Asn Arg Gln Lys Trp Gln Ala Leu Ala Glu Gln Gln Glu Lys Met Leu
          850          855          860
Ile Asn Gly Glu Ser Gly Gln Ala Lys Arg Asn Trp Val Pro Arg Ala
865          870          875          880
Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly Glu Glu Leu Phe
          885          890          895
Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly
          900          905          910
His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly
          915          920          925
Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro
          930          935          940
Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser
945          950          955          960
Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met
          965          970          975
Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly
          980          985          990
Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val
          995          1000          1005
Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile
          1010          1015          1020
Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile
1025          1030          1035          1040
Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg
          1045          1050          1055
His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln
          1060          1065          1070
Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr
          1075          1080          1085
Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp
          1090          1095          1100
His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly
1105          1110          1115          1120
Met Asp Glu Leu Tyr Lys
          1125

```

<210> 9

<211> 3024

<212> DNA

<213> Aequorea victoria and human

<220>

<221> CDS

<222> (1)...(3024)

<400> 9

atg agc tgg tca cct tcc ctg aca acg cag aca tgt ggg gcc tgg gaa	48
Met Ser Trp Ser Pro Ser Leu Thr Thr Gln Thr Cys Gly Ala Trp Glu	
1 5 10 15	
atg aaa gag cgc ctt ggg aca ggg gga ttt gga aat gtc atc cga tgg	96
Met Lys Glu Arg Leu Gly Thr Gly Gly Phe Gly Asn Val Ile Arg Trp	
20 25 30	
cac aat cag gaa aca ggt gag cag att gcc atc aag cag tgc cgg cag	144
His Asn Gln Glu Thr Gly Glu Gln Ile Ala Ile Lys Gln Cys Arg Gln	
35 40 45	
gag ctc agc ccc cgg aac cga gag cgg tgg tgc ctg gag atc cag atc	192
Glu Leu Ser Pro Arg Asn Arg Glu Arg Trp Cys Leu Glu Ile Gln Ile	
50 55 60	
atg aga agg ctg acc cac ccc aat gtg gtg gct gcc cga gat gtc cct	240
Met Arg Arg Leu Thr His Pro Asn Val Val Ala Ala Arg Asp Val Pro	
65 70 75 80	
gag ggg atg cag aac ttg gcg ccc aat gac ctg ccc ctg ctg gcc atg	288
Glu Gly Met Gln Asn Leu Ala Pro Asn Asp Leu Pro Leu Leu Ala Met	
85 90 95	
gag tac tgc caa gga gga gat ctc cgg aag tac ctg aac cag ttt gag	336
Glu Tyr Cys Gln Gly Gly Asp Leu Arg Lys Tyr Leu Asn Gln Phe Glu	
100 105 110	
aac tgc tgt ggt ctg cgg gaa ggt gcc atc ctc acc ttg ctg agt gac	384
Asn Cys Cys Gly Leu Arg Glu Gly Ala Ile Leu Thr Leu Leu Ser Asp	
115 120 125	
att gcc tct gcg ctt aga tac ctt cat gaa aac aga atc atc cat cgg	432
Ile Ala Ser Ala Leu Arg Tyr Leu His Glu Asn Arg Ile Ile His Arg	
130 135 140	
gat cta aag cca gaa aac atc gtc ctg cag caa gga gaa cag agg tta	480
Asp Leu Lys Pro Glu Asn Ile Val Leu Gln Gln Gly Glu Gln Arg Leu	
145 150 155 160	
ata cac aaa att att gac cta gga tat gcc aag gag ctg gat cag ggc	528
Ile His Lys Ile Ile Asp Leu Gly Tyr Ala Lys Glu Leu Asp Gln Gly	
165 170 175	
agt ctt tgc aca tca ttc gtg ggg acc ctg cag tac ctg gcc cca gag	576
Ser Leu Cys Thr Ser Phe Val Gly Thr Leu Gln Tyr Leu Ala Pro Glu	
180 185 190	
cta ctg gag cag cag aag tac aca gtg acc gtc gac tac tgg agc ttc	624
Leu Leu Glu Gln Gln Lys Tyr Thr Val Thr Val Asp Tyr Trp Ser Phe	
195 200 205	
ggc acc ctg gcc ttt gag tgc atc acg ggc ttc cgg ccc ttc ctc ccc	672
Gly Thr Leu Ala Phe Glu Cys Ile Thr Gly Phe Arg Pro Phe Leu Pro	

210	215	220	
aac tgg cag ccc gtg cag tgg cat tca aaa gtg cgg cag aag agt gag			720
Asn Trp Gln Pro Val Gln Trp His Ser Lys Val Arg Gln Lys Ser Glu			
225	230	235	240
gtg gac att gtt gtt agc gaa gac ttg aat gga acg gtg aag ttt tca			768
Val Asp Ile Val Val Ser Glu Asp Leu Asn Gly Thr Val Lys Phe Ser			
	245	250	255
agc tct tta ccc tac ccc aat aat ctt aac agt gtc ctg gct gag cga			816
Ser Ser Leu Pro Tyr Pro Asn Asn Leu Asn Ser Val Leu Ala Glu Arg			
	260	265	270
ctg gag aag tgg ctg caa ctg atg ctg atg tgg cac ccc cga cag agg			864
Leu Glu Lys Trp Leu Gln Leu Met Leu Met Trp His Pro Arg Gln Arg			
	275	280	285
ggc acg gat ccc acg tat ggg ccc aat ggc tgc ttc aag gcc ctg gat			912
Gly Thr Asp Pro Thr Tyr Gly Pro Asn Gly Cys Phe Lys Ala Leu Asp			
	290	295	300
gac atc tta aac tta aag ctg gtt cat atc ttg aac atg gtc acg ggc			960
Asp Ile Leu Asn Leu Lys Leu Val His Ile Leu Asn Met Val Thr Gly			
305	310	315	320
acc atc cac acc tac cct gtg aca gag gat gag agt ctg cag agc ttg			1008
Thr Ile His Thr Tyr Pro Val Thr Glu Asp Glu Ser Leu Gln Ser Leu			
	325	330	335
aag gcc aga atc caa cag gac acg ggc atc cca gag gag gac cag gag			1056
Lys Ala Arg Ile Gln Gln Asp Thr Gly Ile Pro Glu Glu Asp Gln Glu			
	340	345	350
ctg ctg cag gaa gcg ggc ctg gcg ttg atc ccc gat aag cct gcc act			1104
Leu Leu Gln Glu Ala Gly Leu Ala Leu Ile Pro Asp Lys Pro Ala Thr			
	355	360	365
cag tgt att tca gac ggc aag tta aat gag ggc cac aca ttg gac atg			1152
Gln Cys Ile Ser Asp Gly Lys Leu Asn Glu Gly His Thr Leu Asp Met			
	370	375	380
gat ctt gtt ttt ctc ttt gac aac agt aaa atc acc tat gag act cag			1200
Asp Leu Val Phe Leu Phe Asp Asn Ser Lys Ile Thr Tyr Glu Thr Gln			
385	390	395	400
atc tcc cca cgg ccc caa cct gaa agt gtc agc tgt atc ctt caa gag			1248
Ile Ser Pro Arg Pro Gln Pro Glu Ser Val Ser Cys Ile Leu Gln Glu			
	405	410	415
ccc aag agg aat ctc gcc ttc ttc cag ctg agg aag gtg tgg ggc cag			1296
Pro Lys Arg Asn Leu Ala Phe Phe Gln Leu Arg Lys Val Trp Gly Gln			
	420	425	430
gtc tgg cac agc atc cag acc ctg aag gaa gat tgc aac cgg ctg cag			1344
Val Trp His Ser Ile Gln Thr Leu Lys Glu Asp Cys Asn Arg Leu Gln			
	435	440	445
cag gga cag cga gcc gcc atg atg aat ctc ctc cga aac aac agc tgc			1392

Gln	Gly	Gln	Arg	Ala	Ala	Met	Met	Asn	Leu	Leu	Arg	Asn	Asn	Ser	Cys	
450						455					460					
ctc	tcc	aaa	atg	aag	aat	tcc	atg	gct	tcc	atg	tct	cag	cag	ctc	aag	1440
Leu	Ser	Lys	Met	Lys	Asn	Ser	Met	Ala	Ser	Met	Ser	Gln	Gln	Leu	Lys	
465					470				475						480	
gcc	aag	ttg	gat	ttc	ttc	aaa	acc	agc	atc	cag	att	gac	ctg	gag	aag	1488
Ala	Lys	Leu	Asp	Phe	Phe	Lys	Thr	Ser	Ile	Gln	Ile	Asp	Leu	Glu	Lys	
				485					490					495		
tac	agc	gag	caa	acc	gag	ttt	ggg	atc	aca	tca	gat	aaa	ctg	ctg	ctg	1536
Tyr	Ser	Glu	Gln	Thr	Glu	Phe	Gly	Ile	Thr	Ser	Asp	Lys	Leu	Leu	Leu	
			500					505					510			
gcc	tgg	agg	gaa	atg	gag	cag	gct	gtg	gag	ctc	tgt	ggg	cgg	gag	aac	1584
Ala	Trp	Arg	Glu	Met	Glu	Gln	Ala	Val	Glu	Leu	Cys	Gly	Arg	Glu	Asn	
	515						520					525				
gaa	gtg	aaa	ctc	ctg	gta	gaa	cgg	atg	atg	gct	ctg	cag	acc	gac	att	1632
Glu	Val	Lys	Leu	Leu	Val	Glu	Arg	Met	Met	Ala	Leu	Gln	Thr	Asp	Ile	
	530					535					540					
gtg	gac	tta	cag	agg	agc	ccc	atg	ggc	cgg	aag	cag	ggg	gga	acg	ctg	1680
Val	Asp	Leu	Gln	Arg	Ser	Pro	Met	Gly	Arg	Lys	Gln	Gly	Gly	Thr	Leu	
545					550					555					560	
gac	gac	cta	gag	gag	caa	gca	agg	gag	ctg	tac	agg	aga	cta	agg	gaa	1728
Asp	Asp	Leu	Glu	Glu	Gln	Ala	Arg	Glu	Leu	Tyr	Arg	Arg	Leu	Arg	Glu	
			565						570					575		
aaa	cct	cga	gac	cag	cga	act	gag	ggg	gac	agt	cag	gaa	atg	gta	cgg	1776
Lys	Pro	Arg	Asp	Gln	Arg	Thr	Glu	Gly	Asp	Ser	Gln	Glu	Met	Val	Arg	
			580					585					590			
ctg	ctg	ctt	cag	gca	att	cag	agc	ttc	gag	aag	aaa	gtg	cga	gtg	atc	1824
Leu	Leu	Leu	Gln	Ala	Ile	Gln	Ser	Phe	Glu	Lys	Lys	Val	Arg	Val	Ile	
		595					600					605				
tat	acg	cag	ctc	agt	aaa	act	gtg	gtt	tgc	aag	cag	aag	gcg	ctg	gaa	1872
Tyr	Thr	Gln	Leu	Ser	Lys	Thr	Val	Val	Cys	Lys	Gln	Lys	Ala	Leu	Glu	
	610					615					620					
ctg	ttg	ccc	aag	gtg	gaa	gag	gtg	gtg	agc	tta	atg	aat	gag	gat	gag	1920
Leu	Leu	Pro	Lys	Val	Glu	Glu	Val	Val	Ser	Leu	Met	Asn	Glu	Asp	Glu	
625					630					635					640	
aag	act	gtt	gtc	cgg	ctg	cag	gag	aag	cgg	cag	aag	gag	ctc	tgg	aat	1968
Lys	Thr	Val	Val	Arg	Leu	Gln	Glu	Lys	Arg	Gln	Lys	Glu	Leu	Trp	Asn	
				645					650					655		
ctc	ctg	aag	att	gct	tgt	agc	aag	gtc	cgt	ggg	cct	gtc	agt	gga	agc	2016
Leu	Leu	Lys	Ile	Ala	Cys	Ser	Lys	Val	Arg	Gly	Pro	Val	Ser	Gly	Ser	
			660					665					670			
cgg	gat	agc	atg	aat	gcc	tct	cga	ctt	agc	cag	cct	ggg	cag	ctg	atg	2064
Pro	Asp	Ser	Met	Asn	Ala	Ser	Arg	Leu	Ser	Gln	Pro	Gly	Gln	Leu	Met	
		675					680					685				

tct cag ccc tcc acg gcc tcc aac agc tta cct gag cca gcc aag aag	2112
Ser Gln Pro Ser Thr Ala Ser Asn Ser Leu Pro Glu Pro Ala Lys Lys	
690 695 700	
agt gaa gaa ctg gtg gct gaa gca cat aac ctc tgc acc ctg cta gaa	2160
Ser Glu Glu Leu Val Ala Glu Ala His Asn Leu Cys Thr Leu Leu Glu	
705 710 715 720	
aat gcc ata cag gac act gtg agg gaa caa gac cag agt ttc acg gcc	2208
Asn Ala Ile Gln Asp Thr Val Arg Glu Gln Asp Gln Ser Phe Thr Ala	
725 730 735	
cta gac tgg agc tgg tta cag acg gaa gaa gaa gag cac agc tgc ctg	2256
Leu Asp Trp Ser Trp Leu Gln Thr Glu Glu Glu Glu His Ser Cys Leu	
740 745 750	
gag cag gcc tca tgg gta ccg cgg gcc cgg gat cca ccg gtc gcc acc	2304
Glu Gln Ala Ser Trp Val Pro Arg Ala Arg Asp Pro Pro Val Ala Thr	
755 760 765	
atg gtg agc aag ggc gag gag ctg ttc acc ggg gtg gtg ccc atc ctg	2352
Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu	
770 775 780	
gtc gag ctg gac ggc gac gta aac ggc cac aag ttc agc gtg tcc ggc	2400
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly	
785 790 795 800	
gag ggc gag ggc gat gcc acc tac ggc aag ctg acc ctg aag ttc atc	2448
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile	
805 810 815	
tgc acc acc ggc aag ctg ccc gtg ccc tgg ccc acc ctc gtg acc acc	2496
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr	
820 825 830	
ctg acc tac ggc gtg cag tgc ttc agc cgc tac ccc gac cac atg aag	2544
Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys	
835 840 845	
cag cac gac ttc ttc aag tcc gcc atg ccc gaa ggc tac gtc cag gag	2592
Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu	
850 855 860	
cgc acc atc ttc ttc aag gac gac ggc aac tac aag acc cgc gcc gag	2640
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu	
865 870 875 880	
gtg aag ttc gag ggc gac acc ctg gtg aac cgc atc gag ctg aag ggc	2688
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly	
885 890 895	
atc gac ttc aag gag gac ggc aac atc ctg ggg cac aag ctg gag tac	2736
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr	
900 905 910	
aac tac aac agc cac aac gtc tat atc atg gcc gac aag cag aag aac	2784
Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn	
915 920 925	

ggc atc aag gtg aac ttc aag atc cgc cac aac atc gag gac ggc agc 2832
 Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
 930 935 940

 gtg cag ctc gcc gac cac tac cag cag aac acc ccc atc ggc gac ggc 2880
 Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
 945 950 955 960

 ccc gtg ctg ctg ccc gac aac cac tac ctg agc acc cag tcc gcc ctg 2928
 Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
 965 970 975

 agc aaa gac ccc aac gag aag cgc gat cac atg gtc ctg ctg gag ttc 2976
 Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
 980 985 990

 gtg acc gcc gcc ggg atc act ctc ggc atg gac gag ctg tac aag taa 3024
 Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys *
 995 1000 1005

<210> 10
 <211> 1007
 <212> PRT
 <213> Aequorea victoria and human

<400> 10
 Met Ser Trp Ser Pro Ser Leu Thr Thr Gln Thr Cys Gly Ala Trp Glu
 1 5 10 15
 Met Lys Glu Arg Leu Gly Thr Gly Gly Phe Gly Asn Val Ile Arg Trp
 20 25 30
 His Asn Gln Glu Thr Gly Glu Gln Ile Ala Ile Lys Gln Cys Arg Gln
 35 40 45
 Glu Leu Ser Pro Arg Asn Arg Glu Arg Trp Cys Leu Glu Ile Gln Ile
 50 55 60
 Met Arg Arg Leu Thr His Pro Asn Val Val Ala Arg Asp Val Pro
 65 70 75 80
 Glu Gly Met Gln Asn Leu Ala Pro Asn Asp Leu Pro Leu Leu Ala Met
 85 90 95
 Glu Tyr Cys Gln Gly Gly Asp Leu Arg Lys Tyr Leu Asn Gln Phe Glu
 100 105 110
 Asn Cys Cys Gly Leu Arg Glu Gly Ala Ile Leu Thr Leu Leu Ser Asp
 115 120 125
 Ile Ala Ser Ala Leu Arg Tyr Leu His Glu Asn Arg Ile Ile His Arg
 130 135 140
 Asp Leu Lys Pro Glu Asn Ile Val Leu Gln Gln Gly Glu Gln Arg Leu
 145 150 155 160
 Ile His Lys Ile Ile Asp Leu Gly Tyr Ala Lys Glu Leu Asp Gln Gly
 165 170 175
 Ser Leu Cys Thr Ser Phe Val Gly Thr Leu Gln Tyr Leu Ala Pro Glu
 180 185 190
 Leu Leu Glu Gln Gln Lys Tyr Thr Val Thr Val Asp Tyr Trp Ser Phe
 195 200 205
 Gly Thr Leu Ala Phe Glu Cys Ile Thr Gly Phe Arg Pro Phe Leu Pro
 210 215 220
 Asn Trp Gln Pro Val Gln Trp His Ser Lys Val Arg Gln Lys Ser Glu
 225 230 235 240
 Val Asp Ile Val Val Ser Glu Asp Leu Asn Gly Thr Val Lys Phe Ser

Asn Ala Ile Gln Asp Thr Val Arg Glu Gln Asp Gln Ser Phe Thr Ala
 725 730 735
 Leu Asp Trp Ser Trp Leu Gln Thr Glu Glu Glu Glu His Ser Cys Leu
 740 745 750
 Glu Gln Ala Ser Trp Val Pro Arg Ala Arg Asp Pro Pro Val Ala Thr
 755 760 765
 Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 770 775 780
 Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
 785 790 795 800
 Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
 805 810 815
 Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
 820 825 830
 Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
 835 840 845
 Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
 850 855 860
 Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
 865 870 875 880
 Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
 885 890 895
 Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
 900 905 910
 Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
 915 920 925
 Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
 930 935 940
 Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
 945 950 955 960
 Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
 965 970 975
 Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
 980 985 990
 Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
 995 1000 1005

<210> 11
 <211> 2430
 <212> DNA
 <213> Aequorea victoria and human

<220>
 <221> CDS
 <222> (1)...(2430)

<400> 11

atg gac gaa ctg ttc ccc ctc atc ttc ccg gca gag cca gcc cag gcc	48
Met Asp Glu Leu Phe Pro Leu Ile Phe Pro Ala Glu Pro Ala Gln Ala	
1 5 10 15	
tct ggc ccc tat gtg gag atc att gag cag ccc aag cag cgg ggc atg	96
Ser Gly Pro Tyr Val Glu Ile Ile Glu Gln Pro Lys Gln Arg Gly Met	
20 25 30	
cgc ttc cgc tac aag tgc gag ggg cgc tcc gcg ggc agc atc cca ggc	144
Arg Phe Arg Tyr Lys Cys Glu Gly Arg Ser Ala Gly Ser Ile Pro Gly	
35 40 45	

gag agg agc aca gat acc acc aag acc cac ccc acc atc aag atc aat	192
Glu Arg Ser Thr Asp Thr Thr Lys Thr His Pro Thr Ile Lys Ile Asn	
50 55 60	
ggc tac aca gga cca ggg aca gtg cgc atc tcc ctg gtc acc aag gac	240
Gly Tyr Thr Gly Pro Gly Thr Val Arg Ile Ser Leu Val Thr Lys Asp	
65 70 75 80	
cct cct cac cgg cct cac ccc cac gag ctt gta gga aag gac tgc cgg	288
Pro Pro His Arg Pro His Pro His Glu Leu Val Gly Lys Asp Cys Arg	
85 90 95	
gat ggc ttc tat gag gct gag ctc tgc ccg gac cgc tgc atc cac agt	336
Asp Gly Phe Tyr Glu Ala Glu Leu Cys Pro Asp Arg Cys Ile His Ser	
100 105 110	
ttc cag aac ctg gga atc cag tgt gtg aag aag cgg gac ctg gag cag	384
Phe Gln Asn Leu Gly Ile Gln Cys Val Lys Lys Arg Asp Leu Glu Gln	
115 120 125	
gct atc agt cag cgc atc cag acc aac aac aac ccc ttc caa gtt cct	432
Ala Ile Ser Gln Arg Ile Gln Thr Asn Asn Asn Pro Phe Gln Val Pro	
130 135 140	
ata gaa gag cag cgt ggg gac tac gac ctg aat gct gtg cgg ctc tgc	480
Ile Glu Glu Gln Arg Gly Asp Tyr Asp Leu Asn Ala Val Arg Leu Cys	
145 150 155 160	
ttc cag gtg aca gtg cgg gac cca tca ggc agg ccc ctc cgc ctg ccg	528
Phe Gln Val Thr Val Arg Asp Pro Ser Gly Arg Pro Leu Arg Leu Pro	
165 170 175	
cct gtc ctt cct cat ccc atc ttt gac aat cgt gcc ccc aac act gcc	576
Pro Val Leu Pro His Pro Ile Phe Asp Asn Arg Ala Pro Asn Thr Ala	
180 185 190	
gag ctc aag atc tgc cga gtg aac cga aac tct ggc agc tgc ctc ggt	624
Glu Leu Lys Ile Cys Arg Val Asn Arg Asn Ser Gly Ser Cys Leu Gly	
195 200 205	
ggg gat gag atc ttc cta ctg tgt gac aag gtg cag aaa gag gac att	672
Gly Asp Glu Ile Phe Leu Leu Cys Asp Lys Val Gln Lys Glu Asp Ile	
210 215 220	
gag gtg tat ttc acg gga cca ggc tgg gag gcc cga ggc tcc ttt tcg	720
Glu Val Tyr Phe Thr Gly Pro Gly Trp Glu Ala Arg Gly Ser Phe Ser	
225 230 235 240	
caa gct gat gtg cac cga caa gtg gcc att gtg ttc cgg acc cct ccc	768
Gln Ala Asp Val His Arg Gln Val Ala Ile Val Phe Arg Thr Pro Pro	
245 250 255	
tac gca gac ccc agc ctg cag gct cct gtg cgt gtc tcc atg cag ctg	816
Tyr Ala Asp Pro Ser Leu Gln Ala Pro Val Arg Val Ser Met Gln Leu	
260 265 270	
cgg cgg cct tcc gac cgg gag ctc agt gag ccc atg gaa ttc cag tac	864
Arg Arg Pro Ser Asp Arg Glu Leu Ser Glu Pro Met Glu Phe Gln Tyr	
275 280 285	

ctg cca gat aca gac gat cgt cac cgg att gag gag aaa cgt aaa agg	912
Leu Pro Asp Thr Asp Asp Arg His Arg Ile Glu Glu Lys Arg Lys Arg	
290 295 300	
aca tat gag acc ttc aag agc atc atg aag aag agt cct ttc agc gga	960
Thr Tyr Glu Thr Phe Lys Ser Ile Met Lys Lys Ser Pro Phe Ser Gly	
305 310 315 320	
ccc acc gac ccc cgg cct cca cct cga cgc att gct gtg cct tcc cgc	1008
Pro Thr Asp Pro Arg Pro Pro Pro Arg Arg Ile Ala Val Pro Ser Arg	
325 330 335	
agc tca gct tct gtc ccc aag cca gca ccc cag ccc tat ccc ttt acg	1056
Ser Ser Ala Ser Val Pro Lys Pro Ala Pro Gln Pro Tyr Pro Phe Thr	
340 345 350	
tca tcc ctg agc acc atc aac tat gat gag ttt ccc acc atg gtg ttt	1104
Ser Ser Leu Ser Thr Ile Asn Tyr Asp Glu Phe Pro Thr Met Val Phe	
355 360 365	
cct tct ggg cag atc agc cag gcc tcg gcc ttg gcc ccg gcc cct ccc	1152
Pro Ser Gly Gln Ile Ser Gln Ala Ser Ala Leu Ala Pro Ala Pro Pro	
370 375 380	
caa gtc ctg ccc cag gct cca gcc cct gcc cct gct cca gcc atg gta	1200
Gln Val Leu Pro Gln Ala Pro Ala Pro Ala Pro Ala Pro Ala Met Val	
385 390 395 400	
tca gct ctg gcc cag gcc cca gcc cct gtc cca gtc cta gcc cca ggc	1248
Ser Ala Leu Ala Gln Ala Pro Ala Pro Val Pro Val Leu Ala Pro Gly	
405 410 415	
cct cct cag gct gtg gcc cca cct gcc ccc aag ccc acc cag gct ggg	1296
Pro Pro Gln Ala Val Ala Pro Pro Ala Pro Lys Pro Thr Gln Ala Gly	
420 425 430	
gaa gga acg ctg tca gag gcc ctg ctg cag ctg cag ttt gat gat gaa	1344
Glu Gly Thr Leu Ser Glu Ala Leu Leu Gln Leu Gln Phe Asp Asp Glu	
435 440 445	
gac ctg ggg gcc ttg ctt ggc aac agc aca gac cca gct gtg ttc aca	1392
Asp Leu Gly Ala Leu Leu Gly Asn Ser Thr Asp Pro Ala Val Phe Thr	
450 455 460	
gac ctg gca tcc gtc gac aac tcc gag ttt cag cag ctg ctg aac cag	1440
Asp Leu Ala Ser Val Asp Asn Ser Glu Phe Gln Gln Leu Leu Asn Gln	
465 470 475 480	
ggc ata cct gtg gcc ccc cac aca act gag ccc atg ctg atg gag tac	1488
Gly Ile Pro Val Ala Pro His Thr Thr Glu Pro Met Leu Met Glu Tyr	
485 490 495	
cct gag gct ata act cgc cta gtg aca ggg gcc cag agg ccc ccc gac	1536
Pro Glu Ala Ile Thr Arg Leu Val Thr Gly Ala Gln Arg Pro Pro Asp	
500 505 510	
cca gct cct gct cca ctg ggg gcc ccg ggg ctc ccc aat ggc ctc ctt	1584
Pro Ala Pro Ala Pro Leu Gly Ala Pro Gly Leu Pro Asn Gly Leu Leu	

515	520	525	
tca gga gat gaa gac ttc tcc tcc att gcg gac atg gac ttc tca gcc			1632
Ser Gly Asp Glu Asp Phe Ser Ser Ile Ala Asp Met Asp Phe Ser Ala			
530	535	540	
ctg ctg agt cag atc agc tcc aag ctt cga att ctg cag tcg acg gta			1680
Leu Leu Ser Gln Ile Ser Ser Lys Leu Arg Ile Leu Gln Ser Thr Val			
545	550	555	560
ccg cgg gcc cgg gat cca ccg gtc gcc acc atg gtg agc aag ggc gag			1728
Pro Arg Ala Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly Glu			
565	570	575	
gag ctg ttc acc ggg gtg gtg ccc atc ctg gtc gag ctg gac ggc gac			1776
Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp			
580	585	590	
gta aac ggc cac aag ttc agc gtg tcc ggc gag ggc gag ggc gat gcc			1824
Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala			
595	600	605	
acc tac ggc aag ctg acc ctg aag ttc atc tgc acc acc ggc aag ctg			1872
Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu			
610	615	620	
ccc gtg ccc tgg ccc acc ctc gtg acc acc ctg acc tac ggc gtg cag			1920
Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln			
625	630	635	640
tgc ttc agc cgc tac ccc gac cac atg aag cag cac gac ttc ttc aag			1968
Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys			
645	650	655	
tcc gcc atg ccc gaa ggc tac gtc cag gag cgc acc atc ttc ttc aag			2016
Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys			
660	665	670	
gac gac ggc aac tac aag acc cgc gcc gag gtg aag ttc gag ggc gac			2064
Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp			
675	680	685	
acc ctg gtg aac cgc atc gag ctg aag ggc atc gac ttc aag gag gac			2112
Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp			
690	695	700	
ggc aac atc ctg ggg cac aag ctg gag tac aac tac aac agc cac aac			2160
Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn			
705	710	715	720
gtc tat atc atg gcc gac aag cag aag aac ggc atc aag gtg aac ttc			2208
Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe			
725	730	735	
aag atc cgc cac aac atc gag gac ggc agc gtg cag ctc gcc gac cac			2256
Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His			
740	745	750	
tac cag cag aac acc ccc atc ggc gac ggc ccc gtg ctg ctg ccc gac			2304

Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp
 755 760 765
 aac cac tac ctg agc acc cag tcc gcc ctg agc aaa gac ccc aac gag 2352
 Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu
 770 775 780
 aag cgc gat cac atg gtc ctg ctg gag ttc gtg acc gcc gcc ggg atc 2400
 Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile
 785 790 795 800
 act ctc ggc atg gac gag ctg tac aag taa 2430
 Thr Leu Gly Met Asp Glu Leu Tyr Lys *
 805

<210> 12
 <211> 809
 <212> PRT
 <213> Aequorea victoria and human

<400> 12
 Met Asp Glu Leu Phe Pro Leu Ile Phe Pro Ala Glu Pro Ala Gln Ala
 1 5 10 15
 Ser Gly Pro Tyr Val Glu Ile Ile Glu Gln Pro Lys Gln Arg Gly Met
 20 25 30
 Arg Phe Arg Tyr Lys Cys Glu Gly Arg Ser Ala Gly Ser Ile Pro Gly
 35 40 45
 Glu Arg Ser Thr Asp Thr Thr Lys Thr His Pro Thr Ile Lys Ile Asn
 50 55 60
 Gly Tyr Thr Gly Pro Gly Thr Val Arg Ile Ser Leu Val Thr Lys Asp
 65 70 75 80
 Pro Pro His Arg Pro His Pro His Glu Leu Val Gly Lys Asp Cys Arg
 85 90 95
 Asp Gly Phe Tyr Glu Ala Glu Leu Cys Pro Asp Arg Cys Ile His Ser
 100 105 110
 Phe Gln Asn Leu Gly Ile Gln Cys Val Lys Lys Arg Asp Leu Glu Gln
 115 120 125
 Ala Ile Ser Gln Arg Ile Gln Thr Asn Asn Asn Pro Phe Gln Val Pro
 130 135 140
 Ile Glu Glu Gln Arg Gly Asp Tyr Asp Leu Asn Ala Val Arg Leu Cys
 145 150 155 160
 Phe Gln Val Thr Val Arg Asp Pro Ser Gly Arg Pro Leu Arg Leu Pro
 165 170 175
 Pro Val Leu Pro His Pro Ile Phe Asp Asn Arg Ala Pro Asn Thr Ala
 180 185 190
 Glu Leu Lys Ile Cys Arg Val Asn Arg Asn Ser Gly Ser Cys Leu Gly
 195 200 205
 Gly Asp Glu Ile Phe Leu Leu Cys Asp Lys Val Gln Lys Glu Asp Ile
 210 215 220
 Glu Val Tyr Phe Thr Gly Pro Gly Trp Glu Ala Arg Gly Ser Phe Ser
 225 230 235 240
 Gln Ala Asp Val His Arg Gln Val Ala Ile Val Phe Arg Thr Pro Pro
 245 250 255
 Tyr Ala Asp Pro Ser Leu Gln Ala Pro Val Arg Val Ser Met Gln Leu
 260 265 270
 Arg Arg Pro Ser Asp Arg Glu Leu Ser Glu Pro Met Glu Phe Gln Tyr
 275 280 285
 Leu Pro Asp Thr Asp Asp Arg His Arg Ile Glu Glu Lys Arg Lys Arg

290		295		300
Thr Tyr Glu Thr Phe Lys Ser Ile Met Lys Lys Ser Pro Phe Ser Gly				
305		310		315
Pro Thr Asp Pro Arg Pro Pro Pro Arg Arg Ile Ala Val Pro Ser Arg				
		325		330
Ser Ser Ala Ser Val Pro Lys Pro Ala Pro Gln Pro Tyr Pro Phe Thr				
		340		345
Ser Ser Leu Ser Thr Ile Asn Tyr Asp Glu Phe Pro Thr Met Val Phe				
		355		360
Pro Ser Gly Gln Ile Ser Gln Ala Ser Ala Leu Ala Pro Ala Pro Pro				
		370		375
Gln Val Leu Pro Gln Ala Pro Ala Pro Ala Pro Ala Pro Ala Met Val				
385		390		395
Ser Ala Leu Ala Gln Ala Pro Ala Pro Val Pro Val Leu Ala Pro Gly				
		405		410
Pro Pro Gln Ala Val Ala Pro Pro Ala Pro Lys Pro Thr Gln Ala Gly				
		420		425
Glu Gly Thr Leu Ser Glu Ala Leu Leu Gln Leu Gln Phe Asp Asp Glu				
		435		440
Asp Leu Gly Ala Leu Leu Gly Asn Ser Thr Asp Pro Ala Val Phe Thr				
		450		455
Asp Leu Ala Ser Val Asp Asn Ser Glu Phe Gln Gln Leu Leu Asn Gln				
465		470		475
Gly Ile Pro Val Ala Pro His Thr Thr Glu Pro Met Leu Met Glu Tyr				
		485		490
Pro Glu Ala Ile Thr Arg Leu Val Thr Gly Ala Gln Arg Pro Pro Asp				
		500		505
Pro Ala Pro Ala Pro Leu Gly Ala Pro Gly Leu Pro Asn Gly Leu Leu				
		515		520
Ser Gly Asp Glu Asp Phe Ser Ser Ile Ala Asp Met Asp Phe Ser Ala				
		530		535
Leu Leu Ser Gln Ile Ser Ser Lys Leu Arg Ile Leu Gln Ser Thr Val				
545		550		555
Pro Arg Ala Arg Asp Pro Pro Val Ala Thr Met Val Ser Lys Gly Glu				
		565		570
Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp				
		580		585
Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala				
		595		600
Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu				
		610		615
Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln				
625		630		635
Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys				
		645		650
Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys				
		660		665
Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp				
		675		680
Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp				
		690		695
Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn				
705		710		715
Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe				
		725		730
Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His				
		740		745
Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp				
		755		760
				765

Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu
 770 775 780
 Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile
 785 790 795 800
 Thr Leu Gly Met Asp Glu Leu Tyr Lys
 805

<210> 13
 <211> 3018
 <212> DNA
 <213> Aequorea victoria and human

<220>
 <221> CDS
 <222> (1)...(3018)

<400> 13
 atg gtg agc aag ggc gag gag ctg ttc acc ggg gtg gtg ccc atc ctg 48
 Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 1 5 10 15
 gtc gag ctg gac ggc gac gta aac ggc cac aag ttc agc gtg tcc ggc 96
 Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
 20 25 30
 gag ggc gag ggc gat gcc acc tac ggc aag ctg acc ctg aag ttc atc 144
 Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
 35 40 45
 tgc acc acc ggc aag ctg ccc gtg ccc tgg ccc acc ctc gtg acc acc 192
 Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
 50 55 60
 ctg acc tac ggc gtg cag tgc ttc agc cgc tac ccc gac cac atg aag 240
 Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
 65 70 75 80
 cag cac gac ttc ttc aag tcc gcc atg ccc gaa ggc tac gtc cag gag 288
 Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
 85 90 95
 cgc acc atc ttc ttc aag gac gac ggc aac tac aag acc cgc gcc gag 336
 Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
 100 105 110
 gtg aag ttc gag ggc gac acc ctg gtg aac cgc atc gag ctg aag ggc 384
 Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
 115 120 125
 atc gac ttc aag gag gac ggc aac atc ctg ggg cac aag ctg gag tac 432
 Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
 130 135 140
 aac tac aac agc cac aac gtc tat atc atg gcc gac aag cag aag aac 480
 Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
 145 150 155 160
 ggc atc aag gtg aac ttc aag atc cgc cac aac atc gag gac ggc agc 528
 Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser

165									170					175					
gtg	cag	ctc	gcc	gac	cac	tac	cag	cag	aac	acc	ccc	atc	ggc	gac	ggc	576			
Val	Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly				
			180						185			190							
ccc	gtg	ctg	ctg	ccc	gac	aac	cac	tac	ctg	agc	acc	cag	tcc	gcc	ctg	624			
Pro	Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln	Ser	Ala	Leu				
			195						200			205							
agc	aaa	gac	ccc	aac	gag	aag	cgc	gat	cac	atg	gtc	ctg	ctg	gag	ttc	672			
Ser	Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe				
			210			215						220							
gtg	acc	gcc	gcc	ggg	atc	act	ctc	ggc	atg	gac	gag	ctg	tac	aag	tcc	720			
Val	Thr	Ala	Ala	Gly	Ile	Thr	Leu	Gly	Met	Asp	Glu	Leu	Tyr	Lys	Ser				
225						230			235						240				
gga	ctc	aga	tct	cga	gct	caa	gct	tac	atg	agc	tgg	tca	cct	tcc	ctg	768			
Gly	Leu	Arg	Ser	Arg	Ala	Gln	Ala	Tyr	Met	Ser	Trp	Ser	Pro	Ser	Leu				
			245						250						255				
aca	acg	cag	aca	tgt	ggg	gcc	tgg	gaa	atg	aaa	gag	cgc	ctt	ggg	aca	816			
Thr	Thr	Gln	Thr	Cys	Gly	Ala	Trp	Glu	Met	Lys	Glu	Arg	Leu	Gly	Thr				
			260						265						270				
ggg	gga	ttt	gga	aat	gtc	atc	cga	tgg	cac	aat	cag	gaa	aca	ggt	gag	864			
Gly	Gly	Phe	Gly	Asn	Val	Ile	Arg	Trp	His	Asn	Gln	Glu	Thr	Gly	Glu				
			275						280			285							
cag	att	gcc	atc	aag	cag	tgc	cgg	cag	gag	ctc	agc	ccc	cgg	aac	cga	912			
Gln	Ile	Ala	Ile	Lys	Gln	Cys	Arg	Gln	Glu	Leu	Ser	Pro	Arg	Asn	Arg				
			290			295						300							
gag	cgg	tgg	tgc	ctg	gag	atc	cag	atc	atg	aga	agg	ctg	acc	cac	ccc	960			
Glu	Arg	Trp	Cys	Leu	Glu	Ile	Gln	Ile	Met	Arg	Arg	Leu	Thr	His	Pro				
305						310			315						320				
aat	gtg	gtg	gct	gcc	cga	gat	gtc	cct	gag	ggg	atg	cag	aac	ttg	gcg	1008			
Asn	Val	Val	Ala	Ala	Arg	Asp	Val	Pro	Glu	Gly	Met	Gln	Asn	Leu	Ala				
			325						330						335				
ccc	aat	gac	ctg	ccc	ctg	ctg	gcc	atg	gag	tac	tgc	caa	gga	gga	gat	1056			
Pro	Asn	Asp	Leu	Pro	Leu	Leu	Ala	Met	Glu	Tyr	Cys	Gln	Gly	Gly	Asp				
			340						345			350							
ctc	cgg	aag	tac	ctg	aac	cag	ttt	gag	aac	tgc	tgt	ggt	ctg	cgg	gaa	1104			
Leu	Arg	Lys	Tyr	Leu	Asn	Gln	Phe	Glu	Asn	Cys	Cys	Gly	Leu	Arg	Glu				
			355			360						365							
ggt	gcc	atc	ctc	acc	ttg	ctg	agt	gac	att	gcc	tct	gcg	ctt	aga	tac	1152			
Gly	Ala	Ile	Leu	Thr	Leu	Leu	Ser	Asp	Ile	Ala	Ser	Ala	Leu	Arg	Tyr				
			370			375						380							
ctt	cat	gaa	aac	aga	atc	atc	cat	cgg	gat	cta	aag	cca	gaa	aac	atc	1200			
Leu	His	Glu	Asn	Arg	Ile	Ile	His	Arg	Asp	Leu	Lys	Pro	Glu	Asn	Ile				
385						390			395						400				
gtc	ctg	cag	caa	gga	gaa	cag	agg	tta	ata	cac	aaa	att	att	gac	cta	1248			

Val	Leu	Gln	Gln	Gly	Glu	Gln	Arg	Leu	Ile	His	Lys	Ile	Ile	Asp	Leu	
				405					410					415		
gga	tat	gcc	aag	gag	ctg	gat	cag	ggc	agt	ctt	tgc	aca	tca	ttc	gtg	1296
Gly	Tyr	Ala	Lys	Glu	Leu	Asp	Gln	Gly	Ser	Leu	Cys	Thr	Ser	Phe	Val	
			420					425					430			
ggg	acc	ctg	cag	tac	ctg	gcc	cca	gag	cta	ctg	gag	cag	cag	aag	tac	1344
Gly	Thr	Leu	Gln	Tyr	Leu	Ala	Pro	Glu	Leu	Leu	Glu	Gln	Gln	Lys	Tyr	
		435					440					445				
aca	gtg	acc	gtc	gac	tac	tgg	agc	ttc	ggc	acc	ctg	gcc	ttt	gag	tgc	1392
Thr	Val	Thr	Val	Asp	Tyr	Trp	Ser	Phe	Gly	Thr	Leu	Ala	Phe	Glu	Cys	
	450					455					460					
atc	acg	ggc	ttc	cgg	ccc	ttc	ctc	ccc	aac	tgg	cag	ccc	gtg	cag	tgg	1440
Ile	Thr	Gly	Phe	Arg	Pro	Phe	Leu	Pro	Asn	Trp	Gln	Pro	Val	Gln	Trp	
465					470				475					480		
cat	tca	aaa	gtg	cgg	cag	aag	agt	gag	gtg	gac	att	ggt	ggt	agc	gaa	1488
His	Ser	Lys	Val	Arg	Gln	Lys	Ser	Glu	Val	Asp	Ile	Val	Val	Ser	Glu	
			485					490						495		
gac	ttg	aat	gga	acg	gtg	aag	ttt	tca	agc	tct	tta	ccc	tac	ccc	aat	1536
Asp	Leu	Asn	Gly	Thr	Val	Lys	Phe	Ser	Ser	Ser	Leu	Pro	Tyr	Pro	Asn	
			500					505					510			
aat	ctt	aac	agt	gtc	ctg	gct	gag	cga	ctg	gag	aag	tgg	ctg	caa	ctg	1584
Asn	Leu	Asn	Ser	Val	Leu	Ala	Glu	Arg	Leu	Glu	Lys	Trp	Leu	Gln	Leu	
		515					520					525				
atg	ctg	atg	tgg	cac	ccc	cga	cag	agg	ggc	acg	gat	ccc	acg	tat	ggg	1632
Met	Leu	Met	Trp	His	Pro	Arg	Gln	Arg	Gly	Thr	Asp	Pro	Thr	Tyr	Gly	
	530					535					540					
ccc	aat	ggc	tgc	ttc	aag	gcc	ctg	gat	gac	atc	tta	aac	tta	aag	ctg	1680
Pro	Asn	Gly	Cys	Phe	Lys	Ala	Leu	Asp	Asp	Ile	Leu	Asn	Leu	Lys	Leu	
545					550					555				560		
gtt	cat	atc	ttg	aac	atg	gtc	acg	ggc	acc	atc	cac	acc	tac	cct	gtg	1728
Val	His	Ile	Leu	Asn	Met	Val	Thr	Gly	Thr	Ile	His	Thr	Tyr	Pro	Val	
				565				570						575		
aca	gag	gat	gag	agt	ctg	cag	agc	ttg	aag	gcc	aga	atc	caa	cag	gac	1776
Thr	Glu	Asp	Glu	Ser	Leu	Gln	Ser	Leu	Lys	Ala	Arg	Ile	Gln	Gln	Asp	
			580					585					590			
acg	ggc	atc	cca	gag	gag	gac	cag	gag	ctg	ctg	cag	gaa	gcg	ggc	ctg	1824
Thr	Gly	Ile	Pro	Glu	Glu	Asp	Gln	Glu	Leu	Leu	Gln	Glu	Ala	Gly	Leu	
		595					600					605				
gcg	ttg	atc	ccc	gat	aag	cct	gcc	act	cag	tgt	att	tca	gac	ggc	aag	1872
Ala	Leu	Ile	Pro	Asp	Lys	Pro	Ala	Thr	Gln	Cys	Ile	Ser	Asp	Gly	Lys	
	610					615					620					
tta	aat	gag	ggc	cac	aca	ttg	gac	atg	gat	ctt	gtt	ttt	ctc	ttt	gac	1920
Leu	Asn	Glu	Gly	His	Thr	Leu	Asp	Met	Asp	Leu	Val	Phe	Leu	Phe	Asp	
625					630					635					640	

aac agt aaa atc acc tat gag act cag atc tcc cca cgg ccc caa cct	1968
Asn Ser Lys Ile Thr Tyr Glu Thr Gln Ile Ser Pro Arg Pro Gln Pro	
645 650 655	
gaa agt gtc agc tgt atc ctt caa gag ccc aag agg aat ctc gcc ttc	2016
Glu Ser Val Ser Cys Ile Leu Gln Glu Pro Lys Arg Asn Leu Ala Phe	
660 665 670	
ttc cag ctg agg aag gtg tgg ggc cag gtc tgg cac agc atc cag acc	2064
Phe Gln Leu Arg Lys Val Trp Gly Gln Val Trp His Ser Ile Gln Thr	
675 680 685	
ctg aag gaa gat tgc aac cgg ctg cag cag gga cag cga gcc gcc atg	2112
Leu Lys Glu Asp Cys Asn Arg Leu Gln Gln Gly Gln Arg Ala Ala Met	
690 695 700	
atg aat ctc ctc cga aac aac agc tgc ctc tcc aaa atg aag aat tcc	2160
Met Asn Leu Leu Arg Asn Asn Ser Cys Leu Ser Lys Met Lys Asn Ser	
705 710 715 720	
atg gct tcc atg tct cag cag ctc aag gcc aag ttg gat ttc ttc aaa	2208
Met Ala Ser Met Ser Gln Gln Leu Lys Ala Lys Leu Asp Phe Phe Lys	
725 730 735	
acc agc atc cag att gac ctg gag aag tac agc gag caa acc gag ttt	2256
Thr Ser Ile Gln Ile Asp Leu Glu Lys Tyr Ser Glu Gln Thr Glu Phe	
740 745 750	
ggg atc aca tca gat aaa ctg ctg ctg gcc tgg agg gaa atg gag cag	2304
Gly Ile Thr Ser Asp Lys Leu Leu Leu Ala Trp Arg Glu Met Glu Gln	
755 760 765	
gct gtg gag ctc tgt ggg cgg gag aac gaa gtg aaa ctc ctg gta gaa	2352
Ala Val Glu Leu Cys Gly Arg Glu Asn Glu Val Lys Leu Leu Val Glu	
770 775 780	
cgg atg atg gct ctg cag acc gac att gtg gac tta cag agg agc ccc	2400
Arg Met Met Ala Leu Gln Thr Asp Ile Val Asp Leu Gln Arg Ser Pro	
785 790 795 800	
atg ggc cgg aag cag ggg gga acg ctg gac gac cta gag gag caa gca	2448
Met Gly Arg Lys Gln Gly Gly Thr Leu Asp Asp Leu Glu Glu Gln Ala	
805 810 815	
agg gag ctg tac agg aga cta agg gaa aaa cct cga gac cag cga act	2496
Arg Glu Leu Tyr Arg Arg Leu Arg Glu Lys Pro Arg Asp Gln Arg Thr	
820 825 830	
gag ggt gac agt cag gaa atg gta cgg ctg ctg ctt cag gca att cag	2544
Glu Gly Asp Ser Gln Glu Met Val Arg Leu Leu Leu Gln Ala Ile Gln	
835 840 845	
agc ttc gag aag aaa gtg cga gtg atc tat acg cag ctc agt aaa act	2592
Ser Phe Glu Lys Lys Val Arg Val Ile Tyr Thr Gln Leu Ser Lys Thr	
850 855 860	
gtg gtt tgc aag cag aag gcg ctg gaa ctg ttg ccc aag gtg gaa gag	2640
Val Val Cys Lys Gln Lys Ala Leu Glu Leu Leu Pro Lys Val Glu Glu	
865 870 875 880	

gtg gtg agc tta atg aat gag gat gag aag act gtt gtc cgg ctg cag 2688
Val Val Ser Leu Met Asn Glu Asp Glu Lys Thr Val Val Arg Leu Gln
885 890 895

gag aag cgg cag aag gag ctc tgg aat ctc ctg aag att gct tgt agc 2736
Glu Lys Arg Gln Lys Glu Leu Trp Asn Leu Leu Lys Ile Ala Cys Ser
900 905 910

aag gtc cgt ggt cct gtc agt gga agc ccg gat agc atg aat gcc tct 2784
Lys Val Arg Gly Pro Val Ser Gly Ser Pro Asp Ser Met Asn Ala Ser
915 920 925

cga ctt agc cag cct ggg cag ctg atg tct cag ccc tcc acg gcc tcc 2832
Arg Leu Ser Gln Pro Gly Gln Leu Met Ser Gln Pro Ser Thr Ala Ser
930 935 940

aac agc tta cct gag cca gcc aag aag agt gaa gaa ctg gtg gct gaa 2880
Asn Ser Leu Pro Glu Pro Ala Lys Lys Ser Glu Glu Leu Val Ala Glu
945 950 955 960

gca cat aac ctc tgc acc ctg cta gaa aat gcc ata cag gac act gtg 2928
Ala His Asn Leu Cys Thr Leu Leu Glu Asn Ala Ile Gln Asp Thr Val
965 970 975

agg gaa caa gac cag agt ttc acg gcc cta gac tgg agc tgg tta cag 2976
Arg Glu Gln Asp Gln Ser Phe Thr Ala Leu Asp Trp Ser Trp Leu Gln
980 985 990

acg gaa gaa gaa gag cac agc tgc ctg gag cag gcc tca tga 3018
Thr Glu Glu Glu Glu His Ser Cys Leu Glu Gln Ala Ser *
995 1000 1005

<210> 14
<211> 1005
<212> PRT
<213> Aequorea victoria and human

<400> 14
Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
1 5 10 15
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
20 25 30
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
35 40 45
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
50 55 60
Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
65 70 75 80
Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85 90 95
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100 105 110
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115 120 125
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130 135 140
Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn

145		150		155		160
Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser						
	165		170			175
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly						
	180		185			190
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu						
	195		200			205
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe						
	210		215			220
Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys Ser						
225		230		235		240
Gly Leu Arg Ser Arg Ala Gln Ala Tyr Met Ser Trp Ser Pro Ser Leu						
	245		250			255
Thr Thr Gln Thr Cys Gly Ala Trp Glu Met Lys Glu Arg Leu Gly Thr						
	260		265			270
Gly Gly Phe Gly Asn Val Ile Arg Trp His Asn Gln Glu Thr Gly Glu						
	275		280			285
Gln Ile Ala Ile Lys Gln Cys Arg Gln Glu Leu Ser Pro Arg Asn Arg						
	290		295			300
Glu Arg Trp Cys Leu Glu Ile Gln Ile Met Arg Arg Leu Thr His Pro						
305		310		315		320
Asn Val Val Ala Ala Arg Asp Val Pro Glu Gly Met Gln Asn Leu Ala						
	325		330			335
Pro Asn Asp Leu Pro Leu Leu Ala Met Glu Tyr Cys Gln Gly Gly Asp						
	340		345			350
Leu Arg Lys Tyr Leu Asn Gln Phe Glu Asn Cys Cys Gly Leu Arg Glu						
	355		360			365
Gly Ala Ile Leu Thr Leu Leu Ser Asp Ile Ala Ser Ala Leu Arg Tyr						
	370		375			380
Leu His Glu Asn Arg Ile Ile His Arg Asp Leu Lys Pro Glu Asn Ile						
385		390		395		400
Val Leu Gln Gln Gly Glu Gln Arg Leu Ile His Lys Ile Ile Asp Leu						
	405		410			415
Gly Tyr Ala Lys Glu Leu Asp Gln Gly Ser Leu Cys Thr Ser Phe Val						
	420		425			430
Gly Thr Leu Gln Tyr Leu Ala Pro Glu Leu Leu Glu Gln Gln Lys Tyr						
	435		440			445
Thr Val Thr Val Asp Tyr Trp Ser Phe Gly Thr Leu Ala Phe Glu Cys						
	450		455			460
Ile Thr Gly Phe Arg Pro Phe Leu Pro Asn Trp Gln Pro Val Gln Trp						
465		470		475		480
His Ser Lys Val Arg Gln Lys Ser Glu Val Asp Ile Val Val Ser Glu						
	485		490			495
Asp Leu Asn Gly Thr Val Lys Phe Ser Ser Ser Leu Pro Tyr Pro Asn						
	500		505			510
Asn Leu Asn Ser Val Leu Ala Glu Arg Leu Glu Lys Trp Leu Gln Leu						
	515		520			525
Met Leu Met Trp His Pro Arg Gln Arg Gly Thr Asp Pro Thr Tyr Gly						
	530		535			540
Pro Asn Gly Cys Phe Lys Ala Leu Asp Asp Ile Leu Asn Leu Lys Leu						
545		550		555		560
Val His Ile Leu Asn Met Val Thr Gly Thr Ile His Thr Tyr Pro Val						
	565		570			575
Thr Glu Asp Glu Ser Leu Gln Ser Leu Lys Ala Arg Ile Gln Gln Asp						
	580		585			590
Thr Gly Ile Pro Glu Glu Asp Gln Glu Leu Leu Gln Glu Ala Gly Leu						
	595		600			605
Ala Leu Ile Pro Asp Lys Pro Ala Thr Gln Cys Ile Ser Asp Gly Lys						
	610		615			620

```

Leu Asn Glu Gly His Thr Leu Asp Met Asp Leu Val Phe Leu Phe Asp
625                               630                               635                               640
Asn Ser Lys Ile Thr Tyr Glu Thr Gln Ile Ser Pro Arg Pro Gln Pro
                               645                               650                               655
Glu Ser Val Ser Cys Ile Leu Gln Glu Pro Lys Arg Asn Leu Ala Phe
                               660                               665                               670
Phe Gln Leu Arg Lys Val Trp Gly Gln Val Trp His Ser Ile Gln Thr
                               675                               680                               685
Leu Lys Glu Asp Cys Asn Arg Leu Gln Gln Gly Gln Arg Ala Ala Met
690                               695                               700
Met Asn Leu Leu Arg Asn Asn Ser Cys Leu Ser Lys Met Lys Asn Ser
705                               710                               715                               720
Met Ala Ser Met Ser Gln Gln Leu Lys Ala Lys Leu Asp Phe Phe Lys
                               725                               730                               735
Thr Ser Ile Gln Ile Asp Leu Glu Lys Tyr Ser Glu Gln Thr Glu Phe
                               740                               745                               750
Gly Ile Thr Ser Asp Lys Leu Leu Leu Ala Trp Arg Glu Met Glu Gln
                               755                               760                               765
Ala Val Glu Leu Cys Gly Arg Glu Asn Glu Val Lys Leu Leu Val Glu
770                               775                               780
Arg Met Met Ala Leu Gln Thr Asp Ile Val Asp Leu Gln Arg Ser Pro
785                               790                               795                               800
Met Gly Arg Lys Gln Gly Gly Thr Leu Asp Asp Leu Glu Glu Gln Ala
                               805                               810                               815
Arg Glu Leu Tyr Arg Arg Leu Arg Glu Lys Pro Arg Asp Gln Arg Thr
820                               825                               830
Glu Gly Asp Ser Gln Glu Met Val Arg Leu Leu Leu Gln Ala Ile Gln
835                               840                               845
Ser Phe Glu Lys Lys Val Arg Val Ile Tyr Thr Gln Leu Ser Lys Thr
850                               855                               860
Val Val Cys Lys Gln Lys Ala Leu Glu Leu Leu Pro Lys Val Glu Glu
865                               870                               875                               880
Val Val Ser Leu Met Asn Glu Asp Glu Lys Thr Val Val Arg Leu Gln
885                               890                               895
Glu Lys Arg Gln Lys Glu Leu Trp Asn Leu Leu Lys Ile Ala Cys Ser
900                               905                               910
Lys Val Arg Gly Pro Val Ser Gly Ser Pro Asp Ser Met Asn Ala Ser
915                               920                               925
Arg Leu Ser Gln Pro Gly Gln Leu Met Ser Gln Pro Ser Thr Ala Ser
930                               935                               940
Asn Ser Leu Pro Glu Pro Ala Lys Lys Ser Glu Glu Leu Val Ala Glu
945                               950                               955                               960
Ala His Asn Leu Cys Thr Leu Leu Glu Asn Ala Ile Gln Asp Thr Val
                               965                               970                               975
Arg Glu Gln Asp Gln Ser Phe Thr Ala Leu Asp Trp Ser Trp Leu Gln
980                               985                               990
Thr Glu Glu Glu Glu His Ser Cys Leu Glu Gln Ala Ser
995                               1000                               1005

```

<210> 15
 <211> 1659
 <212> DNA
 <213> *Aequorea victoria* and human

<220>
 <221> CDS
 <222> (1)...(1659)

<400> 15

atg gtg agc aag ggc gag gag ctg ttc acc ggg gtg gtg ccc atc ctg Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu 1 5 10 15	48
gtc gag ctg gac ggc gac gta aac ggc cac aag ttc agc gtg tcc ggc Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly 20 25 30	96
gag ggc gag ggc gat gcc acc tac ggc aag ctg acc ctg aag ttc atc Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile 35 40 45	144
tgc acc acc ggc aag ctg ccc gtg ccc tgg ccc acc ctc gtg acc acc Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr 50 55 60	192
ctg acc tac ggc gtg cag tgc ttc agc cgc tac ccc gac cac atg aag Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys 65 70 75 80	240
cag cac gac ttc ttc aag tcc gcc atg ccc gaa ggc tac gtc cag gag Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu 85 90 95	288
cgc acc atc ttc ttc aag gac gac ggc aac tac aag acc cgc gcc gag Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu 100 105 110	336
gtg aag ttc gag ggc gac acc ctg gtg aac cgc atc gag ctg aag ggc Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly 115 120 125	384
atc gac ttc aag gag gac ggc aac atc ctg ggg cac aag ctg gag tac Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr 130 135 140	432
aac tac aac agc cac aac gtc tat atc atg gcc gac aag cag aag aac Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn 145 150 155 160	480
ggc atc aag gtg aac ttc aag atc cgc cac aac atc gag gac ggc agc Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser 165 170 175	528
gtg cag ctc gcc gac cac tac cag cag aac acc ccc atc ggc gac ggc Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly 180 185 190	576
ccc gtg ctg ctg ccc gac aac cac tac ctg agc acc cag tcc gcc ctg Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu 195 200 205	624
agc aaa gac ccc aac gag aag cgc gat cac atg gtc ctg ctg gag ttc Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe 210 215 220	672
gtg acc gcc gcc ggg atc act ctc ggc atg gac gag ctg tac aag tcc Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys Ser 225 230 235 240	720

gga ctc aga tct cga gct caa gct tcc acc atg atg aat ctc ctc cga	768
Gly Leu Arg Ser Arg Ala Gln Ala Ser Thr Met Met Asn Leu Leu Arg	
245 250 255	
aac aac agc tgc ctc tcc aaa atg aag aat tcc atg gct tcc atg tct	816
Asn Asn Ser Cys Leu Ser Lys Met Lys Asn Ser Met Ala Ser Met Ser	
260 265 270	
cag cag ctc aag gcc aag ttg gat ttc ttc aaa acc agc atc cag att	864
Gln Gln Leu Lys Ala Lys Leu Asp Phe Phe Lys Thr Ser Ile Gln Ile	
275 280 285	
gac ctg gag aag tac agc gag caa acc gag ttt ggg atc aca tca gat	912
Asp Leu Glu Lys Tyr Ser Glu Gln Thr Glu Phe Gly Ile Thr Ser Asp	
290 295 300	
aaa ctg ctg ctg gcc tgg agg gaa atg gag cag gct gtg gag ctc tgt	960
Lys Leu Leu Leu Ala Trp Arg Glu Met Glu Gln Ala Val Glu Leu Cys	
305 310 315 320	
ggg cgg gag aac gaa gtg aaa ctc ctg gta gaa cgg atg atg gct ctg	1008
Gly Arg Glu Asn Glu Val Lys Leu Leu Val Glu Arg Met Met Ala Leu	
325 330 335	
cag acc gac att gtg gac tta cag agg agc ccc atg ggc cgg aag cag	1056
Gln Thr Asp Ile Val Asp Leu Gln Arg Ser Pro Met Gly Arg Lys Gln	
340 345 350	
ggg gga acg ctg gac gac cta gag gag caa gca agg gag ctg tac agg	1104
Gly Gly Thr Leu Asp Asp Leu Glu Glu Gln Ala Arg Glu Leu Tyr Arg	
355 360 365	
aga cta agg gaa aaa cct cga gac cag cga act gag ggt gac agt cag	1152
Arg Leu Arg Glu Lys Pro Arg Asp Gln Arg Thr Glu Gly Asp Ser Gln	
370 375 380	
gaa atg gta cgg ctg ctg ctt cag gca att cag agc ttc gag aag aaa	1200
Glu Met Val Arg Leu Leu Leu Gln Ala Ile Gln Ser Phe Glu Lys Lys	
385 390 395 400	
gtg cga gtg atc tat acg cag ctc agt aaa act gtg gtt tgc aag cag	1248
Val Arg Val Ile Tyr Thr Gln Leu Ser Lys Thr Val Val Cys Lys Gln	
405 410 415	
aag gcg ctg gaa ctg ttg ccc aag gtg gaa gag gtg gtg agc tta atg	1296
Lys Ala Leu Glu Leu Leu Pro Lys Val Glu Glu Val Val Ser Leu Met	
420 425 430	
aat gag gat gag aag act gtt gtc cgg ctg cag gag aag cgg cag aag	1344
Asn Glu Asp Glu Lys Thr Val Val Arg Leu Gln Glu Lys Arg Gln Lys	
435 440 445	
gag ctc tgg aat ctc ctg aag att gct tgt agc aag gtc cgt ggt cct	1392
Glu Leu Trp Asn Leu Leu Lys Ile Ala Cys Ser Lys Val Arg Gly Pro	
450 455 460	
gtc agt gga agc ccg gat agc atg aat gcc tct cga ctt agc cag cct	1440
Val Ser Gly Ser Pro Asp Ser Met Asn Ala Ser Arg Leu Ser Gln Pro	

465	470	475	480	
ggg cag ctg atg tct cag ccc tcc acg gcc tcc aac agc tta cct gag				1488
Gly Gln Leu Met Ser Gln Pro Ser Thr Ala Ser Asn Ser Leu Pro Glu				
	485	490	495	
cca gcc aag aag agt gaa gaa ctg gtg gct gaa gca cat aac ctc tgc				1536
Pro Ala Lys Lys Ser Glu Glu Leu Val Ala Glu Ala His Asn Leu Cys				
	500	505	510	
acc ctg cta gaa aat gcc ata cag gac act gtg agg gaa caa gac cag				1584
Thr Leu Leu Glu Asn Ala Ile Gln Asp Thr Val Arg Glu Gln Asp Gln				
	515	520	525	
agt ttc acg gcc cta gac tgg agc tgg tta cag acg gaa gaa gaa gag				1632
Ser Phe Thr Ala Leu Asp Trp Ser Trp Leu Gln Thr Glu Glu Glu Glu				
	530	535	540	
cac agc tgc ctg gag cag gcc tca tga				1659
His Ser Cys Leu Glu Gln Ala Ser *				
545	550			

<210> 16
 <211> 552
 <212> PRT
 <213> Aequorea victoria and human

<400> 16

Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu				
1	5	10	15	
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly				
	20	25	30	
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile				
	35	40	45	
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr				
	50	55	60	
Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys				
65	70	75	80	
Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu				
	85	90	95	
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu				
	100	105	110	
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly				
	115	120	125	
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr				
	130	135	140	
Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn				
145	150	155	160	
Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser				
	165	170	175	
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly				
	180	185	190	
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu				
	195	200	205	
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe				
	210	215	220	
Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys Ser				
225	230	235	240	

Gly Leu Arg Ser Arg Ala Gln Ala Ser Thr Met Met Asn Leu Leu Arg
 245 250 255
 Asn Asn Ser Cys Leu Ser Lys Met Lys Asn Ser Met Ala Ser Met Ser
 260 265 270
 Gln Gln Leu Lys Ala Lys Leu Asp Phe Phe Lys Thr Ser Ile Gln Ile
 275 280 285
 Asp Leu Glu Lys Tyr Ser Glu Gln Thr Glu Phe Gly Ile Thr Ser Asp
 290 295 300
 Lys Leu Leu Leu Ala Trp Arg Glu Met Glu Gln Ala Val Glu Leu Cys
 305 310 315 320
 Gly Arg Glu Asn Glu Val Lys Leu Leu Val Glu Arg Met Met Ala Leu
 325 330 335
 Gln Thr Asp Ile Val Asp Leu Gln Arg Ser Pro Met Gly Arg Lys Gln
 340 345 350
 Gly Gly Thr Leu Asp Asp Leu Glu Glu Gln Ala Arg Glu Leu Tyr Arg
 355 360 365
 Arg Leu Arg Glu Lys Pro Arg Asp Gln Arg Thr Glu Gly Asp Ser Gln
 370 375 380
 Glu Met Val Arg Leu Leu Leu Gln Ala Ile Gln Ser Phe Glu Lys Lys
 385 390 395 400
 Val Arg Val Ile Tyr Thr Gln Leu Ser Lys Thr Val Val Cys Lys Gln
 405 410 415
 Lys Ala Leu Glu Leu Leu Pro Lys Val Glu Glu Val Val Ser Leu Met
 420 425 430
 Asn Glu Asp Glu Lys Thr Val Val Arg Leu Gln Glu Lys Arg Gln Lys
 435 440 445
 Glu Leu Trp Asn Leu Leu Lys Ile Ala Cys Ser Lys Val Arg Gly Pro
 450 455 460
 Val Ser Gly Ser Pro Asp Ser Met Asn Ala Ser Arg Leu Ser Gln Pro
 465 470 475 480
 Gly Gln Leu Met Ser Gln Pro Ser Thr Ala Ser Asn Ser Leu Pro Glu
 485 490 495
 Pro Ala Lys Lys Ser Glu Glu Leu Val Ala Glu Ala His Asn Leu Cys
 500 505 510
 Thr Leu Leu Glu Asn Ala Ile Gln Asp Thr Val Arg Glu Gln Asp Gln
 515 520 525
 Ser Phe Thr Ala Leu Asp Trp Ser Trp Leu Gln Thr Glu Glu Glu Glu
 530 535 540
 His Ser Cys Leu Glu Gln Ala Ser
 545 550