

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 January 2012 (05.01.2012)

(10) International Publication Number
WO 2012/000957 A1

(51) International Patent Classification:

A 61K 8/36 (2006.01) A61Q 19/00 (2006.01)
A61K 8/37 (2006.01) A61Q 19/02 (2006.01)
A61K 8/66 (2006.01) A61Q 19/08 (2006.01)
A61K 8/97 (2006.01) A23L 1/30 (2006.01)
A61K 8/99 (2006.01)

(21) International Application Number:

PCT/EP201 1/060762

(22) International Filing Date:

28 June 2011 (28.06.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10167858.9 30 June 2010 (30.06.2010) EP

(71) Applicant (for all designated States except US):
NESTEC S.A. [CH/CH]; Av. Nestle 55, CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GUITARD, Marjorie [FR/CH]; Chemin de la Daumuse 6, CH-1073 Savigny (CH). BEL RHLID, Rachid [CA/CH]; Route de l'Ancienne Poste IB, CH-1073 Savigny (CH). MOODY-CLIFFE, Angus [GB/CH]; Chemin du Grammont 7, CH-1073 Savigny (CH). DIONISI, Fabiola [IT/CH]; Chemin des Roches 32B, CH-1066 Epalinges (CH).

(74) Agent: CHAUTARD, Cecile; Av. Nestle 55, CH-1800 Vevey (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.1 7(H))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(Hi))
- of inventorship (Rule 4.1 7(iv))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))

(54) Title: USE OF CHICORIC ACID AND LACTIC BACTERIUM FOR REGULATING SKIN PIGMENTATION

(57) Abstract: The present invention relates generally to the field of cosmetic and/or food products. More specifically, the present invention aims to provide the use of at least an ingredient containing chioric acid and/or derivatives and a micro-organism and/or an enzyme capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid, for improving skin tone and preventing and/or treating hyper-pigmentation of skin and/or skin color imperfections such as age-spots and other skin disorders characterized by abnormal pigments. Also, the present invention aims at providing a skin lightening agent.



WO 2012/000957 A1

USE OF CHICORIC ACID AND LACTIC BACTERIUM FOR REGULATING SKIN PIGMENTATION

The present invention relates generally to the field of food or beverage products for cosmetic purpose. More specifically, the present invention aims to provide the use of an ingredient containing chicoric acid and/or derivatives and a micro-organism and/or an enzyme capable of hydrolysing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid, for preventing and/or treating hyper-pigmentation of skin, skin color imperfections such as age-spots and other skin disorders characterized by abnormal pigments. The present invention also aims at improving skin tone as well as providing a skin lightening or whitening agent.

Background of the invention

Skin color is primarily determined by the amount and type of melanin, a brown pigment present in the skin. Lower amounts of melanin result in lighter skin color while higher amounts result in darker skin color. Also, hyper-pigmentation in the skin is caused by the over expression or accumulation of melanin in the skin. As a result, the pathway involved in melanin production has been the target for many inhibitors so as to reduce the levels produced. One of the principal enzymes involved in the melanin pathway is tyrosinase.

The synthesis of melanin is a process under hormonal control, including the melanocyte stimulating hormone (MSH) and adrenocorticotrophic hormone (ACTH) peptides that are produced from the precursor proopiomelanocortin. It is stimulated by the DNA damages that are caused by UVB-radiations as well.

Then, exposure to the sun over time can induce many biochemical reactions in the skin, leading to sunburn and tanning, for example. Other consequences of exposure to the sun accumulate over time. These changes can result in the development of age spots and create an uneven, mottled skin tone. Unfortunately, many of the commercially available products in today's market are either only marginally effective, or contain active agents that are unstable and lose their potency when incorporated into a final formula.

The ability to modify the expression of pigment content in the skin, to promote an evenness skin tone or lightening skin tone, is highly desired in today's society. Many people desire to modify their skin tone, to reduce aging spots, etc., or for purely cosmetic reasons.

As a result, efforts to develop effective compositions have focused on agents that inhibit the activity of tyrosinase. For example, a variety of tyrosinase inhibitors, such as hydroquinone, vitamin C, cystein, kojic acid, arbutin and glutathione among others have been proposed in topical compositions. Also, a variety of dermatological compositions have been suggested for improving the appearance of pigment disorders such as that observed in melasma, freckles, vitiligo, piebaldism, phenylketonuria, and the like, and/or for cosmetic purposes.

Also, the use of skin bleaching compositions is widely expanded. However, they either destroy melanin or inhibit its formation. Many of these contain harsh chemicals such as peroxides, acids or formaldehyde, or thiolated materials. Less stringent therapies have other disadvantages.

Topical retinoid and topical corticosteroids have been suggested as hypo-pigmenting agents, as have laser treatment and chemical peels, but these fall short of desirable responses.

Other compositions suggested the use on the skin of natural materials, which have in some cases been used for centuries in Asia or Europe to bleach skin and skin areas, or enhance the appearance of fair skin. These include the use of lemon, orange, cucumber, ginkgo, carob, rose fruit, geranium herb, cinnamon, sweet marjoram, rosemary, etc...

In order to combat disorders related to abnormal pigment or to lighten skin tone, various compounds which, when applied topically to the skin, are capable of reducing tyrosinase activity and consequently limiting melanin production, have thus been proposed.

Unfortunately, the treatments currently available are not entirely satisfactory, in particular in terms of the side effects which are frequently associated therewith, such as irritant side effects with certain topical agents.

It would thus be highly desirable to have alternative preparations that do not have the drawbacks of those described in the prior art. In particular, it would be highly desirable to develop nutritional cosmetic compositions to be administered via oral route that have improved stability and efficacy to promote an evenness skin tone or to lighten skin tone.

There also remains a need to active agents that are effective for treating and/or preventing skin pigmentation disorders, in particular those due to environmental factors or aging.

The object of the present invention is to meet these needs.

Summary of the invention

- 5 The present inventors could achieve this object by providing a food or beverage composition that comprises at least one ingredient containing chicoric acid and/or derivatives and a lactic bacterium and/or an enzyme capable of hydrolysing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid.
- 10 Thus, according to a first subject, the invention relates to the cosmetic use of least one ingredient containing chicoric acid and/or derivatives and a lactic bacterium and/or an enzyme capable of hydrolysing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid, for treating and/or preventing skin pigmentation disorders. Such skin disorder are in particular those due to age or to environmental factors (e.g. UV), such as age-
- 15 spots. It may also be skin disorders that are observed in melasma, freckles, vitiligo, piebaldism, phenylketonuria, and the like.

The present inventors have discovered that ingredient containing chicoric acid treated with a lactic bacterium effectively suppress the formation of melanin, melanogenesis, despite the

20 fact that the extracts show little to no inhibition of tyrosinase activity. This result is surprising and unexpected considering the pivotal role of tyrosinase in melanogenesis and the focus of development efforts in the art to inhibit this enzyme.

For the purpose of the present invention, the term "skin" is intended to mean the skin of the

25 face or of the body.

For the purpose of the present invention the term "effective amount", is intended to mean an amount sufficient to obtain the expected effect.

30 For the purpose of the present invention the term "prevent" is intended to mean the fact of reducing the risk of occurrence of the manifestation of the disorder under consideration.

The present invention is also directed towards the cosmetic use of the abovementioned ingredient, as an active agent for treating and/or composition for preventing the skin

35 imperfections, in particular to improve skin tone or skin complexion.

The present invention is also directed towards the cosmetic use of the abovementioned ingredient, as an active agent for treating and/or preventing the skin pigment imperfections. As a result, the complexion becomes brighter and more homogeneous, without areas of dyschromia or of dryness.

5

The present invention is also directed towards the cosmetic use of an effective amount of at least one ingredient containing chicoric acid and/or derivatives and a lactic bacterium and/or an enzyme capable of hydrolysing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid, as an active agent for whitening or lightening skin tone.

10

The present inventors have also discovered that the ingredient according to the invention further improves hydration and/or skin barrier function.

A use in accordance with the present invention may also comprises the use of at least one ingredient containing chicoric acid and/or derivatives and a lactic bacterium and/or an enzyme capable of hydrolysing chicoric acid to generate tartaric and/or caffeic acid, in combination with an effective amount of at least one active agent for further improving skin hydration or skin ageing, in particular as described hereinafter.

According to another of its aspects, the subject of the invention is a method, in particular a cosmetic method, for treating and/or preventing skin tone imperfections and the disorders associated with hyper-pigmentation, in particular aesthetic disorders, in an individual, comprising at least one step of administering, to said individual, at least one ingredient containing chicoric acid and/or derivatives and a lactic bacterium and/or an enzyme capable of hydrolysing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid.


Compositions according to the present invention are orally administrable. This has the advantage of acting globally on the entire skin, in its deep layers (dermis, hypodermis), by means of a rapid and relatively non-restrictive mode of administration. Specifically, the metabolites and other active nutrients are in particular distributed within the dermal matrix by means of the bloodstream. Oral administration also has the advantage of a rapid and relatively non-restrictive mode of administration.

35

Detailed description of the invention


Ingredient containing chicoric acid and or derivatives

Chicoric acid is



5

and derivatives of chicoric acid include



10


R_1 , R_2 , R_3 and/or R_4 may be identical or may differ from one another.

In one embodiment, R_1 , R_2 , R_3 and/or R_4 may be selected from the group consisting of H; CH_3 ; CrC_3 -alkyl; aryl, such as phenyl, benzyl, tolyl, o-xylylalkyl; CrC_3 -acyl; amino acids; mono-, di- or oligosaccharides. Oligosaccharides contain between two and nine monosaccharide units. R_1 , R_2 , R_3 and R_4 may be identical and/or may differ from one another.

15

One typical derivative of chicoric acid is for example the following compound :

20



The ingredient containing chicoric acid and/or derivatives thereof may be any ingredient that contains chicoric acid and/or derivatives thereof, either naturally or in added form, but is preferably a natural foodstuff such as lettuce, chicory, dandelion, grape, grape pomace; or combinations or extracts thereof.

In an embodiment, the plant material is added to the oral cosmetic composition in the form of an extract, such as a chicory extract. The chicory extract can be made from any suitable part of the plant material includes, for example, the root, the pulp, the like or combinations thereof.

Suitable extracts of chicory for the purpose of the present invention are also extracts that are commercially available, such as for example Leroux MS55 (commercially available from Leroux SAS, France)

In a particular preferred embodiment of the present invention, suitable extracts of chicory may be prepared by any means that are known in the art, e.g., by steam extraction, solvent extraction, distillation, pressing or grinding. In particular, the extract is obtainable by extraction with a solvent from Chicory plant material, by a water extraction or an alcohol/water extraction, for example by a ethanol/water extraction or methanol/water extraction. The extract can be used in liquid form (e.g. Leroux MS55, Leroux MS70) or powder form (e.g. Leroux Sol B).

For ease of handling, the plant material is preferably in a dried and comminuted or powder form. As described below, the processes utilize dried, comminuted chicory and/or extracts thereof. However, it is to be understood that any suitable plant material may be used in any suitable form and added to the product according to the present invention.

The extract is processed such that its flavor can be enhanced. For example, bitter flavors

which are typically associated with plant materials, such as chicory, can be removed by processing the plant into an extract. The extract can also be prepared such that the amount of bioactive agent in the final extract product can be desirably controlled.

5 It should be appreciated that the plant material can be processed to form an extract in a variety of different and suitable ways. In general, the plant material, such as the chicory root, is ground, powdered or provided in any suitable form. The plant material can then be further processed in a number of different stages to produce the product extract. In an embodiment, a defatting procedure is performed on the plant material to produce an extract that results
10 from fats removed from the plant material. The defatting procedure can be conducted under any suitable defatting process conditions with any suitable types and amounts of solvents including, for example, hexane.

In an embodiment, the resultant extract of the defatting procedure can be further processed
15 via acid hydrolysis to produce another type of plant extract that can be added to the nutritional composition of the present invention. The acid hydrolysis procedure can be conducted under any suitable process condition with any suitable types and amounts of solvents, including, for example, ethyl acetate.

20 In an embodiment, the extract from the defatting procedure can be further processed via a solvent extraction procedure. The solvent extraction can be carried out under any suitable process conditions and in the presence of any suitable amount and type of solvent. In an embodiment, the solvent includes a solution of methanol ("MeOH") and water mixed in a 1:1 volume ratio. The resultant solution of the solvent extraction procedure can be further
25 processed by evaporation of the solvent under suitable conditions to produce another extract. Alternatively, the resultant solution can be treated with an adsorbant agent, such as polyvinylpyrrolidone or the like, to trap polyphenols. The adsorbant agent treatment can be carried out under any suitable process conditions.

30 The amount of Chicoric acid or a plant extract containing it, in the product will depend on several factors, such as the nature of the extract, the condition of the plant, the age, condition and size of the person or animal to be treated, the frequency, the product will be administered and/or the specific kind of skin disorder or damage to be treated or prevented or desired cosmetic effect.

35

The present inventors have found that the effectiveness of chicoric acid or an extract

containing it according to the present invention is generally dose dependant and follows a dose response curve. If generally mild skin disorders or damages are to be prevented and the product will be used frequently, very small amounts of chicoric acid or an extract thereof will be sufficient to achieve the desired effect. If a severe skin pigment disorder is to be treated, larger amounts of the ingredient will be more appropriate, although also small amounts will produce an effect.

In a preferred embodiment the ingredient is enriched in chicoric acid and/or derivatives thereof. For example, the ingredient and/or the composition may comprise chicoric acid and/or derivatives thereof in an amount in the range of 0,001-99,99 weight-% of dry weight, preferably 0,1-50 weight-% of dry weight, most preferred 0,1-10 weight-% of dry weight. The ingredient and/or the composition may comprise the micro-organism and/or the enzyme capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid in an amount in the range of 0,001-99,99 weight-% of dry weight, preferably 0,1-50 weight-% of dry weight, most preferred 0,1-10 weight-% of dry weight.

Generally, it is preferred if the product contains chicory or an extract thereof in an amount in the range of about 0,1 g/l to 10 g/l, preferably in the range of 0,5 g/l to 3 g/l product. If the total amount of product cannot be measured in litres it is preferred if the product contains chicory or an extract thereof in an amount in the range of about 0,1 g/kg to 10 g/kg, preferably in the range of 0,5 g/kg to 3 g/kg product. Preferably the product contains chicory or an extract thereof in a daily dose of 0,01 g-100g, preferably 0,25 g-10g.


The ingredient to be mixed with the ingredient containing chicoric acid and/or derivatives thereof comprises a lactic acid bacterium and/or an enzyme capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid. These two ingredients may be mixed briefly prior to consumption or may be provided as a ready-to-consume composition.

Microorganisms or enzyme capable of hydrolysing chicoric acid

Chicoric acid and/or its derivatives can then be hydrolyzed by a lactic acid bacterium capable of hydrolyzing chicoric acid and/or derivatives thereof. This hydrolysis step will generate tartaric and/or caffeic acid.

Without wishing to be bound by theory, the inventors believe that this hydrolysis occurs as

outlined in the following scheme. Note, that all reaction steps catalyzed by "enzyme" may also be catalyzed by the lactic acid bacterium described in the present application and/or by combinations of lactic acid bacterium and enzymes.



The inventors have surprisingly found that treating an ingredient comprising chicoric acid and/or derivatives thereof with lactic acid bacterium or an enzyme capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid results for example in improved activity of the ingredient.

Furthermore, it has been found that this treatment can take place *in vivo* when a human or an animal ingests a composition comprising chicoric acid and/or derivatives thereof in combination with a lactic acid bacterium capable of hydrolyzing chlorogenic acids to generate phenolic acids.

Preferred lactic acid bacterium capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid are probiotic lactic acid bacterium having an esterase

activity, such as chlorogenate esterase and/or feruloyl esterase, preferably *Lactobacillus* or *Bifidobacterium*, for example *L. johnsonii*, *B. longum*, and *B. lactis* (CNCM 1-3446), even more preferred *Lactobacillus johnsonii* La1 (CNCM 1-1225), *B. longum* BB 536, and *B. lactis* BB12.

5

B. longum BB 536 is commercially available from Morinaga Nutritional Foods, Inc.

B. lactis BB12 is commercially available, e.g., from Chr. Hansen, DK-2970 Horsholm.

10 In one embodiment of the present invention, the lactic acid bacterium may be used in a non-replicating form.

The ability of a lactic acid bacterium or of a fraction thereof to hydrolyze chicoric acid may be tested as described in detail for *L. johnsonii* in examples.

15

In another embodiment, an enzyme capable of hydrolyzing chicoric acid to generate tartaric and/or caffeic acid is further added to the lactic acid bacterium. Preferably, such enzyme is selected from the group consisting of esterases, such as chlorogenate esterase, tannase and/or feruloyl esterase. It may be added in an amount such as preferably at least 5%, such
20 as at least 30%, at least 50%, or at least 75% of chicoric acid present in the composition is hydrolyzed prior to and/or during consumption.

Suitable enzymes that can be used in the framework of the present invention include e.g. esterases, e.g. chlorogenate esterase derived from *Aspergillus japonicus*. (Commercially
25 available from Kikkoman, Japan), tannase from *Aspergillus oryzae* (EC 3.1.1.20) (commercially available from Kikkoman, Japan). The enzyme may be present as a purified enzyme (immobilized or not) or e.g. in the form of a cell lysate of a microorganism. Suitable cells may e.g. be cells of the microorganisms mentioned above. Suitable methods for producing cell lysate are known in the art.

30

The composition and/or ingredients of the invention should be formulated such that the lactic acid bacterium strain will not ferment or react with the composition during storage. This may be achieved e.g. by formulating the composition as a dry powder, and/or by encapsulating the lactic acid bacterium so that it will only be released when the composition is mixed with at
35 least one other ingredient or during digestion.

The lactic acid bacterium should be present in an amount sufficient for hydrolyzing a substantial amount of chicoric acid to generate tartaric and/or caffeic acid during digestion. The amount of lactic acid bacterium and/or enzyme needed may e.g. be determined by those skilled in the art, for example dependent on the subject to be treated or on the speed by which the tartaric and/or caffeic acid should be liberated. Preferably at least 5%, such as at least 30%, at least 50%, or at least 75% of chicoric acid present in the composition is hydrolysed prior to and/or during consumption.

The compositions according to the invention may be in any of the galenical forms usually available for the method of administration selected.

Galenical forms

The compositions according to the invention may be in any of the galenical forms normally available for the method of administration selected. The carrier may be of diverse nature depending on the type of composition under consideration.

Nutritional complete formula, dairy product such as milk, yogurt, cheese, fermented milks, milk-based fermented products, ice-creams, cereal-based products or fermented cereal-based products, milk-based powders, infant and baby formulas, chilled or shelf stable beverages, food products of confectionary, chocolate or cereal type, animal feed, in particular for domestic animals. In particular, the chicory or extract thereof may be incorporated into any other form of enriched food, for example food bars or compacted or non-compacted powders. The powders may be diluted in water, soda, milk products or soya bean derivatives, or be incorporated into food bars.

The food product may also further comprise a protein source, a carbohydrate source, a lipid source, a mineral source and/or a vitamin source. The presence of proteins, carbohydrates, lipids, minerals and/or vitamins may have several advantages. These compounds generally contribute to the taste and mouthfeel of the final product. They also provide the body with nutrients that it may need urgently when it is affected by skin disorders. They also allow formulating the product of the present invention as a complete nutritional formula, so that no additional nutrition is needed.

The individual ingredients may comprise any kind of edible compound. Typical ingredients for food compositions, in particular drinks, are well known in the art, e.g. milk, cream, coffee

whiteners, and coffee creamers. Alternatively, ingredients could also be a salad dressing or parts thereof, for example. Such compounds are used by consumers to modify e.g. the aroma, appearance and texture of a composition. The ingredients may be in liquid or dry form, e.g. as powders, that are dissolved and/or suspended in a drink.

5

The composition of the invention may further comprise further ingredient suitable for inclusion in a food composition. Usual ingredients may e.g. be sugars, artificial sweeteners, emulsifiers, stabilisers, thickeners, flowing agents, colours, flavours, aromas, and the like. Suitable artificial sweeteners include saccharin, cyclamates, acetosulfame, L-aspartyl based
10 sweeteners such as aspartame, and mixtures of these. Suitable emulsifiers include monoglycerides, diglycerides, lecithin, diacetyl tartaric acid esters of mono-diglycerides, emulsifying starches, and mixtures thereof. Suitable stabilisers include dipotassium phosphate and sodium citrate. A suitable flowing agent is sodium silica aluminate. In one embodiment the composition comprises milk protein and/or vegetable protein. In a further
15 embodiment the composition comprises milk fat and/or vegetable fat.

In one embodiment the invention relates to a kit for preparing a food product, for example a beverage, comprising at least two parts: a) a first part comprising an ingredient containing
20 chicoric acid ; and b) a second part comprising a lactic acid bacterium capable of hydrolyzing chicoric acid to generate tartaric and/or caffeic acid. The two parts may be sold together for the preparation of a food product, for example a beverage, but may be physically separated in the packing of the product. The final food product to be consumed may be prepared by mixing the two parts shortly before consumption. If one or both parts are in a liquid form they
25 may be mixed directly, optionally further liquid, e.g. water or milk, may be added. The two parts may also be mixed by dissolving or suspending them in a liquid, e.g. water or milk. When liquid is used this may be hot or cold. If hot liquid is used, it may preferably have a temperature which is not so high as to inactivate the lactic acid bacterium and enzyme if any, before ingestion of the foodstuff or beverage.

30

In a preferred embodiment the first part is in a dry form, e.g. in the form of a powder. The first part may also be in a liquid form. The first part may additionally comprise any other suitable ingredient, e.g., aroma additives, stabilisers, salts, and/or sweeteners. The first part may be packed in any suitable way, e.g. in a sachet, bottle or can.

35

The second part comprises a lactic acid bacterium capable of hydrolysing chicoric acid into

tartaric and/or caffeic acid. This part may preferably be in the form of a composition to be mixed with the first part. It may be in dry form, e.g. as a powder, or in liquid form, and may be packed in any suitable way, e.g. in a sachet, bottle or can. It should be formulated such that the lactic acid bacterium will not ferment or react with other ingredients during storage. This may be achieved e.g. by formulating the composition as a dry powder, and/or by encapsulating the lactic acid bacterium.

The at least two parts may be packed together in any suitable way. They may e.g. be packed in a combined container wherein the parts are kept physically separated during storage and mixed when the container is opened, or they may be packed in separate containers which are sold together for the preparation of a foodstuff or beverage.

Those skilled in the art will understand that they can freely combine all features of the present invention described herein, without departing from the scope of the invention as disclosed. In particular, features described for the uses of the present invention may be applied to the composition and/or the kit of the present invention and vice versa.

Use

The products of the invention may be efficiently used for treating or preventing skin pigmentation disorders or cosmetically lightening skin tone e.g. by decreasing the production of melanin. Indeed ingredient containing chicoric acid treated with probiotic bacterium La-1 was shown to decrease in vitro the synthesis of melanin (Example 1, Figure 1). The production of tyrosinase was also decreased but to a limited extent (Figure 2), suggesting that the decrease in melanin was not due to tyrosinase inhibition but rather to mechanisms acting upstream or downstream of this enzyme.

The compositions according to the present invention have further a positive effect on strengthening skin barrier and maintaining skin hydration.

As a result, the pigment imperfections are reduced, the complexion becomes brighter and more homogeneous, without areas of dyschromia, or of dryness.

Thus, according to one subject, the invention relates to the cosmetic use of an effective amount of at least one ingredient containing chicoric acid and/or derivatives and a lactic bacterium and/or an enzyme capable of hydrolyzing chicoric acid and/or derivatives thereof

to generate tartaric and/or caffeic acid, as an active agent for treating and/or preventing skin pigmentation disorders, in particular those due to age or environmental factors such as UV.

The present compositions may also be used as active agent for whitening or lightening skin tone, which is particularly desirable for asian population.

A use in accordance with the present invention may also comprises the use of the compositions, in combination with an effective amount of at least one active agent for improving skin hydration or skin ageing, in particular as described hereinafter.

According to another of its aspects, the subject of the invention is a method, in particular a cosmetic method, for treating and/or preventing skin tone imperfections and the disorders associated with hyper-pigmentation, in particular aesthetic disorders, in an individual, comprising at least one step of administering, to said individual, compositions as described above.

The cosmetic treatment method of the invention may be carried out in particular by orally administering the composition as described above. Oral administration comprises ingesting, in one or more intakes, an oral composition as defined above.

It may comprise a single application. According to another embodiment, the application is repeated, for example, 2 to 3 times a day, for one day or more, and generally for a sustained period of at least 4, or even 1 to 15, weeks.

In addition, combinations of treatment with, optionally, oral or topical forms may be envisaged in order to supplement or reinforce the activity of the compositions as defined by the invention.

Thus, a topical or oral treatment with a composition containing chicoric acid in accordance with the invention, combined with an oral or topical composition optionally containing another active ingredient, in particular a probiotic microorganism, or other probiotics in dead, live or semi-active form or a hydrating or anti-ageing agent could be imagined as a kit. The ingredients are mixed, before they are formulated, in the order and under conditions readily determined by those skilled in the art.

The ingredients are mixed, before they are formulated, in the order and under conditions

readily determined by those skilled in the art.

Further advantages and features of the present invention are apparent from the following Examples and Figures. The examples hereinafter are thus presented by way of non-limiting illustration of the field of the invention. In these examples, unless otherwise indicated, the percentages are percentages by weight and the ranges of values written in the form "between ... and ..." include the upper and lower limits specified. The term "cfu" denotes "colony forming unit". This is the unit of measurement used to quantify live bacteria.

10 Figures

Figure 1 : Melanin production by murine melanocytes pre-treated with chicoric acid + La1 vs. controls (positive / negative).

Figure 2 : tyrosinase production by murine melanocytes pre-treated with chicoric acid + La1 vs. controls (positive / negative).

Figure 3 : Filaggrin synthesis by human primary epidermal keratinocytes pre-treated with chicoric acid + La1 vs. controls (positive / negative).

EXAMPLES

20

Example 1: Hydrolysis of chicoric acid

1- Hydrolysis of chicoric acid with chlorogenate esterase

25 A solution of chlorogenate esterase (0.8 mg, 24 U/g, from Kikkoman Japan) in 100 μ l phosphate buffer (50 mM, pH 7.0) was added to a solution of chicoric acid (0.57 mg) in 100 μ l phosphate buffer (50 mM, pH 7.0). The mixture was then incubated at 37°C for 4 h. After the reaction time, the enzymatic activity was stopped by heat treatment (5 min, 90°C) and the mixture was centrifuged (microcon YM10, 30 min, 14000g). The supernatant was then
30 analysed by HPLC. A control reaction was run in parallel under the same reaction conditions but without enzyme.

2- Hydrolysis of chicoric acid with *L. johnsonii* (La1) fresh cells

35 Cells of *L. johnsonii* (CNCM 1-1225) were grown (7.0×10^8 cfu/ml) and centrifuged (5000 g, 10 min), the pellets were resuspended in phosphate buffer (50 mM, pH 7.0) at a concentration

of 0.61 g/ml. To 100 μ l of this cells solution, 100 μ l of a solution of chicoric acid (12 mM) was added and the mixture was incubated at 37°C. Samples were withdrawn at different reaction times, centrifuged (3000 g, 5 min), filtered through 0.45 μ m pore size syringe filters (Millipore SLHA 025 BS) and analysed by HPLC.

- 5 A control reaction was run in parallel under the same reaction conditions but without bacterium.

3- Hydrolysis of chicoric acid with *L. johnsonii* extract (lysed cells)

- 10 Cells of *L. johnsonii* (CNCM 1-1225) were grown (7.0 E08 cfu/ml) and centrifuged (5000 g, 10 min), the pellets were resuspended in phosphate buffer (50 mM, pH 7.0) at a concentration of 0.61 g/ml. The cells were then lysed using the glass-beads method. 600 μ l of cells preparation were put into a Mini-Beadbeater for 1 min of intense shaking, cooled in ice, and put another 1 min in the Mini-Beadbeater. The crude cell extract (100 μ l) was then added to
15 100 μ l of a solution of chicoric acid (12 mM, phosphate buffer 50 mM, pH 7.0) and the mixture was incubated at 37°C. Samples were withdrawn at different reaction times, centrifuged (3000g, 5 min), filtered through 0.45 μ m pore size syringe filters (Millipore SLHA 025 BS) and analysed by HPLC.

- 20 4- Hydrolysis of chicoric acid with a spray-dried preparation of La1

- 10 mg of a spray-dried preparation of La1 (3.3 E09 cfu/g) were dissolved in 100 μ l of phosphate buffer (50 mM, pH 7.0). To this solution, 100 μ l of a chicoric acid solution (12 mM, phosphate buffer 50 mM, pH 7.0) were added. The mixture was then incubated at 37°C and
25 samples were withdrawn at different reaction times. After centrifugation (3000 g, 5 min) and filtration (0.45 μ m pore size syringe filters, Millipore SLHA 025 BS) the samples were analysed by HPLC.

HPLC Analysis

- 30 HPLC-DAD analysis of chicoric acid and hydrolysis products was performed on Agilent 1100 system equipped with Atlantis C18 reverse-phase column (4.6 x 100 mm, particle size 3 μ m) and a diode array detector. The column was equilibrated with water containing 0.1 % formic acid. After injection, a linear gradient to a final solvent composition of 55 % water and 45 % acetonitrile (containing 0.1 % formic acid) was run within 12 min at a flow rate of 1 ml/min.
35 Chicoric acid and caffeic acid were monitored by UV at 320 nm and were quantified using standard calibration curves.

RESULTS

Tested bacteria

Bacteria	Culture Media
<i>Lactobacillus rhamnosus</i> GG (NCC 4007)	MRS
<i>Lactobacillus johnsonii</i> La1 (CNCM I-1225)	MRS
<i>Lactobacillus paracasei</i> ST11 (NCC 2461)	MRS + Cysteine
<i>Bifidobacterium longum</i> BB 536 (ATCC BAA-999)	MRS + Cysteine
<i>Bifidobacterium lactis</i> BB12 (CNCM I-3446)	MRS
<i>Streptococcus thermophilus</i> TH4 (NCC 2496)	HJL

5 Table 1

In particular *L. johnsonii* (La1), *B. longum* BB 536, and *B. lactis* BB12 were able to hydrolyse chicoric acid. The best results in term of reaction rate and reaction yield were obtained with *L. johnsonii* (La1)

10

Tested enzymes

Enzyme	Supplier
Chlorogenate esterase	Kikkoman, Japan
Feruloyl esterase	Novozymes
Porcine liver esterase	Sigma E-3019
Hog liver esterase immobilised on Eupergit C	Fluka 46064
Esterase from <i>Saccharomyces cerevisiae</i>	Fluka 46071
Esterase from <i>L. johnsonii</i> (La1) CNCM I-1225	Internal production

Table 2

15 Chlorogenate esterase, feruloyl esterase and La1 esterase were able to hydrolyse chicoric acid into caffeic and tartaric acids.

1- Hydrolysis of chicoric acid with chlorogenate esterase

20 Table 3: Hydrolysis of chicoric acid into caffeic and caftaric acids by chlorogenate esterase. Concentration in % relative to untreated reference at t=0

Time (min)	0	15	30	60	120	240
chicoric Acid	98	85	83	77	65	50
Caftaric Acid	0	6	7	10	16	24
Caffeic Acid	2	9	10	13	19	26

Table 3

2- Hydrolysis of chicoric acid with La1 fresh cells

After 4h reaction time all chicoric acid was transformed into caffeic acid and tartaric acid as analysed by HPLC

3- Hydrolysis of chicoric acid with La1 extract (lysed cells)

As compared to whole cells, the hydrolysis of chicoric acid with lysed cells resulted in an increase of the reaction rate. Indeed, after only 2h all chicoric acid was transformed into caffeic and tartaric acids

4- Hydrolysis of chicoric acid with a spray-dried preparation of La1

Table 4: Hydrolysis of chicoric acid into caffeic and caftaric acids by a spray-dried preparation of *L. johnsonii* (CNCM 1-1225). Concentration in % relative to untreated reference at t=0

	0 min	15 min	30 min	1h	2h	3h
Chicoric Acid (mmole)	365	172	137	91	49	23
Caftaric Acid (mmole)	0	62	63	50	31	15
Caffeic Acid (mmole)	0	324	393	498	601	669

Table 4

Exemple 2 : Effect on Skin pigmentation

In order to evaluate the potential beneficial effect of ingredients towards skin de- or pro-pigmentation we used 2D culture of murine melanocytes (B16) and we performed 2 tests: 1- assessment of melanin production and 2- assessment of tyrosinase production.

1. The cell culture conditions.

B16 cells were cultured in DMEM 1g/L glucose without phenol red supplemented with

10 % foetal calf serum, in a humidified chamber at 37 °C and containing 5% CO₂.

2. The production of melanin by B16 murine melanocyte cell line.

Cells were incubated with the selected ingredients or the test references (Kojic acid at 400 µg/mL) for 72 hours, in the presence or absence of NDP-MSH an analog of MSH. The total quantity of melanin (extracellular and intracellular) was evaluated by measurement of the optical density at 405 nm of each sample against melanin standards in presence or in absence of NDP-MSH.

3. The production of tyrosinase by B16 murine melanocyte cell line.

Cells were incubated with the selected ingredients or the test references (Kojic acid at 400 µg/mL) for 48 hours. The production of tyrosinase was evaluated by immunolabeling.

Ingredients:

The selected ingredients are listed in the Table 1 below. Chicoric acid has been pre-treated with La-1 [(spray dried culture Lactobacillus Johnsonii CNCM 1-1225, 1.19E10 cfu/g) for 24 hours in a thermomixer at 40°C, under shaking conditions. After incubation, samples have been centrifuged 5 minutes at 3000g. After this treatment the probiotic as such is not present anymore in the sample however the presence of its metabolites can't be excluded. The tested concentrations of chicoric acid are also indicated in Table 5.

Ingredient	Highest non cytotoxic conc. on HDF (mg/mL)	Highest non cytotoxic conc. on HPEK (mg/mL)	Tested conc. on HDF (mg/mL)	Tested conc. on HPEK (mg/mL)
MS55 10mg/ml in NaPO ₄ 10mM pH 7.0 + 10mg/ml La1	2	10	0.4	10
Chicoric acid 10mM ie 4.74mg/ml + 60mg La1	0.2 mM	0.04 mM	0.2 mM	0.04 mM
La-1 (10EE11 cfu/g)	0.01	0.01	0.01	0.01

Table 5

Results

Results are expressed in percentage relative to the control. Test reference (Kojic acid)

induced, as expected a decrease in melanin content. Figure 1 shows the melanin production by B16 melanocytes treated for 72 hours with the selected ingredients.

Chicoric acid treated with the probiotic CNCM 1-1225 appeared to be efficient for skin depigmentation. Indeed chicoric acid treated with the probiotic decreased melanin production by 90% (Figure 1). The production of tyrosinase was also decreased by this ingredient but to a limited extent (less than 20%, Figure 2), suggesting that the decrease in melanin was not due to tyrosinase inhibition but rather to mechanisms acting upstream or downstream of this enzyme.

Example 3: Effect on skin barrier function and hydration

The potential beneficial effect of the Extracts of Example 2 towards skin barrier function and skin hydration was evaluated by using 2D culture of human primary epidermal keratinocytes and we assessed their synthesis of filaggrin after treatment with selected ingredients.

The cell culture conditions .

Human epidermal keratinocytes were cultured in control keratinocytes-SFM medium, in a humidified chamber at 37 °C and containing 5% CO₂.

The synthesis of filaggrin by human epidermal keratinocytes.

Cells were incubated with the selected ingredients or the test references (CaCl₂ at 1.5mM) for 144 hours. The production of filaggrin by was evaluated by immunolabeling.

Results

The results are shown in Figure 3. Pre-treatment of the cells with chicoric acid + La1 showed a significant increase of filaggrin (500 % of the control, Figure 3), suggesting that these extracts could strengthen skin barrier. A stronger skin barrier ensures a better protection of the body from the environment and pathogens' attack. It also limits the loss of water through the epidermis, thus ensuring an appropriate skin hydration.

PCT

Print Out (Original in Electronic Form)

(This sheet is not part of and does not count as a sheet of the international application)

0-1	Form PCT/RO/134 (SAFE) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared Using	PCT Online Filing Version 3.5.000.225 MT/FOP 20020701/0.20.5.20
0-2	International Application No.	
0-3	Applicant's or agent's file reference	10886-WO-PCT

1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	Paragraph number	
1-3	Identification of deposit	
1-3-1	Name of depositary institution	CNCM Collection nationale de cultures de micro-organismes
1-3-2	Address of depositary institution	Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cedex 15, France
1-3-3	Date of deposit	02 July 1992 (02.07.1992)
1-3-4	Accession Number	CNCM 1-1225
1-5	Designated States for Which Indications are Made	All designations
2	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
2-1	Paragraph number	
2-3	Identification of deposit	
2-3-1	Name of depositary institution	CNCM Collection nationale de cultures de micro-organismes
2-3-2	Address of depositary institution	Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cedex 15, France
2-3-3	Date of deposit	09 September 2005 (09.09.2005)
2-3-4	Accession Number	CNCM 1-3446
2-5	Designated States for Which Indications are Made	All designations

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	YES
0-4-1	Authorized officer	Dobbelaere, Julie

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

Claims

- 5 1. Cosmetic use of an effective amount of at least an ingredient containing chicoric acid and/or derivatives and a lactic acid bacterium and/or an enzyme capable of hydrolysing chicoric acid to generate tartaric and/or caffeic acid, for the preparation of a food product, a beverage or a pharmaceutical oral composition for treating and/or preventing skin pigment disorders and/or skin imperfections.
- 10 2. Use in accordance with one of the preceding claims, characterized in that the skin disorders are that observed in melasma, freckles, vitiligo, piebaldism, phenylketonuria, and the like, and/or age spots.
- 15 3. Use in accordance with one of the preceding claims for improving skin tone and/or skin complexion.
- 20 4. Cosmetic use of an effective amount of at least an ingredient containing chicoric acid and/or derivatives and a lactic acid bacterium and/or an enzyme capable of hydrolysing chicoric acid to generate tartaric and/or caffeic acid, for the preparation of a food product, a beverage or a pharmaceutical oral composition for whitening or lightening skin tone.
- 25 5. Use in accordance with one of the preceding claims, wherein the composition further improves hydration of the skin and/or skin barrier function..
- 30 6. Use in accordance with one of the preceding claims, wherein the chicoric acid containing ingredient is a natural foodstuff such as lettuce, chicory, dandelion, grape, grape pomace; or combinations or extracts thereof.
- 35 7. Use in accordance with one of the preceding claims, which further contains an enzyme capable of hydrolyzing chicoric acid to generate tartaric and/or caffeic acid.
8. Use in accordance with one of the preceding claims, wherein the lactic acid bacterium capable of hydrolyzing chicoric acid to generate tartaric and/or caffeic acid is a probiotic lactic acid bacterium having an esterase activity, such as chlorogenate esterase and/or feruloyl esterase activity.

9. Use in accordance with one of the preceding claims, wherein the lactic acid bacterium is selected from the group consisting of *L. johnsonii*, *B. longum*, and *B. lactis* in particular *Lactobacillus johnsonii* (CNCM 1-1225), *B. longum* BB 536, *B. lactis* (CNCM I-3446) and *B. lactis* BB12.
10. Use in accordance with one of the preceding claims, in combination with an effective amount of at least one active agent for improving skin hydration or skin ageing.
11. A kit for treating and/or preventing skin pigment disorders and/ or skin imperfections, comprising an oral treatment with a food, beverage or pharmaceutical composition containing at least an ingredient containing chicoric acid and/or derivatives and a lactic acid bacterium and/or an enzyme capable of hydrolyzing chicoric acid to generate tartaric and/or caffeic acid, combined with an oral or topical composition optionally containing a probiotic microorganism in dead, live or semi-active form, or an hydrating or anti-ageing agent.
12. A kit for preparing a food product or beverage, comprising at least two parts:
- a) a first part comprising a chicoric acid and/or derivatives thereof containing food ingredient; and
- b) a second part comprising a lactic acid bacterium and/or an enzyme capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid.
13. The kit of claim 11 wherein the lactic acid bacterium capable of hydrolyzing chicoric acid and/or derivatives thereof to generate tartaric and/or caffeic acid is a probiotic lactic acid bacterium having an esterase activity, such as chlorogenate esterase and/or feruloyl esterase activity.

1 / 2




FIG. 1




FIG. 2




FIG. 3

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/060762

A. CLASSIFICATION OF SUBJECT MATTER				
INV.	A61K8/36	A61K8/37	A61K8/66	A61K8/97
	A61Q19/00	A61Q19/02	A61Q19/08	A23L1/30
ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED				
Minimum documentation searched (classification system followed by classification symbols)				
A61K A61Q A23L				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)				
EPO-Internal				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages			Relevant to claim No.
X,P	EP 2 210 504 AI (NESTEC SA [CH]) 28 July 2010 (2010-07-28) paragraphs [0043] , [0049] , [0050] - [0052] ; claims 12-14 -----			1-13
X	EP 0 227 490 AI (LEROUX CHICOREE SARL [FR]) 1 July 1987 (1987-07-01) page 3, lines 19-23 ; claims 1-3 -----			1-13
A	EP 0 450 040 BI (OREAL [FR]) 9 May 2001 (2001-05-09) claims ; examples -----			1-13
A	US 2007/183996 AI (OKOMBI SABRINA [FR] ET AL) 9 August 2007 (2007-08-09) claims 1, 30-32 -----			1-13
-/- .				
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.				
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family				
Date of the actual completion of the international search			Date of mailing of the international search report	
14 November 2011			23/11/2011	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016			Authorized officer Mi tchel 1, Gemma	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/060762

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 1 570 839 A1 (KOSE CORP [JP]) 7 September 2005 (2005-09-07) c l a i m s 1, 11-15 -----	1-13
A	EP 1 261 315 B1 (UNI LEVER PLC [GB] ; UNI LEVER NV [NL]) 19 July 2006 (2006-07-19) c l a i m s -----	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/060762

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 2210504	AI	28-07-2010	AR 075188 AI 16-03-2011 CA 2750685 AI 05-08-2010 EP 2210504 AI 28-07-2010 EP 2391228 AI 07-12-2011 Wo 2010086322 AI 05-08-2010
EP 0227490	AI	01-07-1987	DE 227490 TI 05-11-1987 DE 3664785 DI 07-09-1989 EP 0227490 AI 01-07-1987 FR 2591073 AI 12-06-1987 JP 3038254 B 10-06-1991 JP 62138435 A 22-06-1987 US 4671962 A 09-06-1987
EP 0450040	BI	09-05-2001	AR 244545 AI 30-11-1993 AT 200977 T 15-05-2001 AU 638768 B2 08-07-1993 AU 6628890 A 16-05-1991 BE 1003833 A3 23-06-1992 BR 9006960 A 12-11-1991 CA 2044269 AI 21-04-1991 CH 682129 A5 30-07-1993 DE 69033731 DI 13-06-2001 DE 69033731 T2 07-02-2002 EP 0450040 AI 09-10-1991 ES 2085192 AI 16-05-1996 FR 2653336 AI 26-04-1991 GB 2244005 A 20-11-1991 IT 1251741 B 23-05-1995 JP 2659618 B2 30-09-1997 JP H04502476 A 07-05-1992 MX 173203 B 08-02-1994 US 5164185 A 17-11-1992 Wo 9105543 AI 02-05-1991 ZA 9008398 A 28-08-1991
US 2007183996	AI	09-08-2007	DE 102006052963 AI 16-05-2007 ES 2292362 AI 01-03-2008 FR 2892923 AI 11-05-2007 GB 2431876 A 09-05-2007 GB 2465703 A 02-06-2010 JP 2007131622 A 31-05-2007 KR 20070049582 A 11-05-2007 US 2007183996 AI 09-08-2007 US 2011237551 AI 29-09-2011
EP 1570839	AI	07-09-2005	AU 2003252305 AI 07-06-2004 EP 1570839 AI 07-09-2005 HK 1082916 AI 15-05-2009 KR 20050084971 A 29-08-2005 TW 1280138 B 01-05-2007 US 2006147397 AI 06-07-2006 wo 2004041236 AI 21-05-2004
EP 1261315	BI	19-07-2006	AT 333263 T 15-08-2006 AU 768908 B2 08-01-2004 AU 4832501 A 17-09-2001 CA 2397198 AI 13-09-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/060762

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		CN 1424904 A	18-06 -2003
		DE 60121549 T2	26-07 -2007
		EP 1261315 A2	04-12 -2002
		JP 4709459 B2	22-06 -2011
		JP 2003525890 A	02-09 -2003
		MX PA02008518 A	13-12 -2002
		WO 0166076 A2	13-09 -2001
		ZA 200204157 A	30-07 -2003
