



US00PP33710P2

(12) **United States Plant Patent**  
**Van Dijk**

(10) **Patent No.:** **US PP33,710 P2**

(45) **Date of Patent:** **Nov. 30, 2021**

- (54) **ANTHURIUM PLANT NAMED ‘ANTHGOUWHI’**
- (50) Latin Name: *Anthurium andraeanum* L.  
Varietal Denomination: **ANTHGOUWHI**
- (71) Applicant: **ANTHURA B.V.**, Bleiswijk (NL)
- (72) Inventor: **Jan Van Dijk**, Bleiswijk (NL)
- (73) Assignee: **Anthura B.V.**, Bleiswijk (NL)
- (\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **17/300,449**
- (22) Filed: **Jul. 1, 2021**
- (51) **Int. Cl.**  
*A01H 6/00* (2018.01)  
*A01H 5/02* (2018.01)

- (52) **U.S. Cl.**  
USPC ..... **Plt./365**
- (58) **Field of Classification Search**  
USPC ..... **Plt./365**  
CPC ..... **A01H 6/00; A01H 5/02**  
See application file for complete search history.

*Primary Examiner* — Keith O. Robinson  
(74) *Attorney, Agent, or Firm* — Jondle & Associates, P.C.

(57) **ABSTRACT**  
A new *Anthurium* plant named ‘ANTHGOUWHI’ particularly distinguished by having weakly blistered and weakly glossy, white, cordate, and durable spathes that retain the original color for a very long period of time, dark green and ovate-cordate, durable leaves, white spadices with yellow tips, early and rich flowering continuously throughout the year, and a plant height of 30.0 cm to 35.0 cm is disclosed.

**3 Drawing Sheets**

**1**

Genus and species: *Anthurium andraeanum* L.  
Variety denomination: ‘ANTHGOUWHI’.

**BACKGROUND OF THE NEW PLANT**

The present invention comprises a new and distinct variety of *Anthurium*, botanically known as *Anthurium andraeanum* L., and hereinafter referred to by the variety name ‘ANTHGOUWHI’. The new *Anthurium* plant is a product of a planned breeding program conducted by the inventor in Bleiswijk, the Netherlands. The objective of this breeding program was to create a new plant with a height of 30.0 cm to 35.0 cm having weakly blistered and weakly glossy, white, cordate, and very durable spathes.

The new variety originated from a cross-pollination made in August 2011 in Bleiswijk, the Netherlands. The female parent was a white *Anthurium* pot plant designated ‘15108-02’ (unpatented), and the male parent was a red *Anthurium* pot plant designated ‘12902-01’ (unpatented).

A single plant was selected from the progeny of the stated cross in June 2013. Asexual reproduction of the new variety by tissue culture over an 8-year period in Bleiswijk, the Netherlands, has demonstrated that the new variety reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations.

Community Plant Variety Rights for this variety have been applied for in the European Union on Nov. 12, 2019 (Application no. 2019/2947), by Applicant who obtained the subject matter disclosed directly from the inventor. ‘ANTHGOUWHI’ has not been made publicly available or sold anywhere in the world prior to the effective filing date of this application with the exception of sales or disclosures made one year or less before the effective filing date of this claimed invention by Applicant who obtained ‘ANTHGOUWHI’ directly from the inventor.

**2**

**SUMMARY OF THE INVENTION**

The following are the most outstanding and distinguishing characteristics of this new variety when grown under normal horticultural practices in Bleiswijk, the Netherlands:

- 1) Weakly blistered, weakly glossy, white, and cordate spathes;
- 2) White spadices with yellow tips;
- 3) Dark green and ovate-cordate leaves; and
- 4) Plant is rich in shooting.

**DESCRIPTION OF THE PHOTOGRAPHS**

This new *Anthurium* plant is illustrated by the accompanying photographs which show the overall plant habit including blooms and foliage of the plant; the colors shown are as true as can be reasonably obtained by conventional photographic procedures. The photographs are of a 34-week-old plant grown in a greenhouse in Bleiswijk, the Netherlands, in April 2021. Colors in the photographs may differ from the color values cited in the detailed botanical description, which accurately describe the actual colors of the new variety.

FIG. 1 shows the overall plant habit, including blooms and foliage.

FIG. 2 shows a close-up of the mature spathe.

FIG. 3 shows a close-up of the upper leaf blade surface.

**DESCRIPTION OF THE NEW VARIETY**

The following detailed description sets forth the distinctive characteristics of ‘ANTHGOUWHI’. The data which define these characteristics were collected from asexual reproductions carried out in Bleiswijk, the Netherlands. The plant history was taken on 34-week-old plants which were planted from tissue culture in 12-centimeter (diameter) pots and grown in a glass greenhouse between 19° C. and 24° C. Observations were made in April 2021. Color readings were

taken under 5000 lux natural light in the greenhouse. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.) (2015).

#### DETAILED BOTANICAL DESCRIPTION

##### Classification:

*Family*.—Araceae.

*Botanical*.—*Anthurium andraeanum* L.

*Common name*.—*Anthurium*.

*Denomination*.—‘ANTHGOUWHI’.

##### Parentage:

*Female parent*.—*Anthurium* plant ‘15108-02’ (unpatented).

*Male parent*.—*Anthurium* plant ‘12902-01’ (unpatented).

##### Plant:

*Propagation*.—Tissue culture.

*Root description*.—Fleshy-yellowish-white (RHS NN155A) colored roots with small hairy lateral roots having small light greenish-yellow (RHS 3C) colored root tips.

*Time to produce a finished flowering plant*.—32 to 36 weeks after planting in a 12-cm (diameter) pot.

*Growth habit*.—Upright.

*Height (measured from soil, including inflorescence)*.—30.0 cm to 35.0 cm.

*Width (measured from leaf tips)*.—28.0 cm to 33.0 cm.

##### Leaves:

*Immature leaves*.—Length: 10.0 cm to 12.0 cm. Width: 6.0 cm to 7.0 cm. Color: Upper surface: RHS 146A. Lower surface: RHS 146B. Texture (both upper and lower surfaces): Leathery, thin, smooth, and glossy.

*Mature leaves*.—Length (fully expanded): 12.0 cm to 14.0 cm. Width: 7.5 cm to 8.5 cm. Shape: Ovate cordate. Apex: Acuminate. Base: Cordate. Leaf blade angle with the petiole: Between 110 degrees and 130 degrees. Leaf margin: Entire. Color: Upper surface: RHS 147A. Lower surface: RHS 146B. Texture: Upper surface: Leathery, thick, and smooth. Lower surface: Leathery, thick, smooth, and glossy. Venation: Pinnate veining; the mid-vein and primary veins (the veins that radiate out from the junction of petiole and leaf) protrude at the underside of the leaf blade. Venation color: Upper surface: RHS 144A. Lower surface: RHS 144B.

*Lobes*.—Present. Arrangement: Leaf blade has two lobes extending past the petiole. The lobes are non-touching. Length of lobes of mature leaf blades: 2.0 cm to 2.5 cm. Width of lobes of mature leaf blades: 3.0 cm to 3.5 cm. Distance from petiole/leaf junction to highest point on lobes of mature leaf blades: 2.5 cm to 3.0 cm.

*Petiole*.—Cross-section: Round. Diameter: 0.3 cm to 0.4 cm. Length: 12.0 cm to 15.0 cm for a mature leaf size. Color: Mature leaf: RHS 144A. Immature leaf: RHS 144B. Cataphyll color surrounding the petiole: Outside: RHS 144B. Inside: RHS 145C.

*Geniculum*.—Length: 1.5 cm to 2.0 cm. Width: 0.3 cm to 0.4 cm. Color: RHS 144C.

##### Inflorescence:

*Arrangement*.—Single.

*Flowering habit (length of flowering season)*.—Continuous.

*Number of inflorescences per plant*.—6 to 8.

*Fragrance*.—Absent.

*Longevity of inflorescence on plant*.—Over a year.

##### Spathe:

*Buds*.—The spathe is tightly rolled around the spadix and extrudes from the peduncle sheath. After the spathe is fully opened the peduncle elongates some extra centimeters.

*Arrangement*.—Spathe angle with the peduncle is between 100 degrees and 110 degrees; the spathe stands on a wiry peduncle about 4.0 cm to 7.0 cm above the foliage.

*Shape*.—Cordate.

*Apex*.—Acuminate.

*Base*.—Cordate.

*Texture*.—Weakly blistered and weakly glossy.

*Margin*.—Undulated.

*Size*.—Length: 8.0 cm to 8.5 cm. Width: 7.0 cm to 7.5 cm.

*Lobes*.—Present. Arrangement: The spathe has two lobes extending past the peduncle. The lobes are non-touching. Length: 0.5 cm to 1.5 cm. Width: 3.0 cm to 3.5 cm.

*Color*.—Just fully open: Upper surface: RHS 155C. Lower surface: RHS 155C. This white color remains for a very long period, at least more than 30 weeks after opening. The spathe turns green after some weeks.

##### Peduncle:

*Shape*.—Erect.

*Cross-section*.—Round.

*Length*.—21.0 cm to 24.0 cm.

*Diameter*.—0.3 cm to 0.4 cm.

*Color*.—RHS 144B.

##### Flowering time:

*General*.—One small, rooted, untreated tissue culture plant of 8.0 cm tall will flower, depending on the season, after 32 to 36 weeks and 6 to 7 blossoms appear. More blossoms appear after some additional weeks so that a full flowering and commercial plant will have 7 to 8 white spathes. Smaller blossoms may occur on immature plants.

##### Spadix:

*Size*.—Length: 2.0 cm to 2.5 cm (depending on flower size). Width (at apex): 0.4 cm to 0.5 cm. Width (at base): 0.5 cm to 0.6 cm.

*Shape*.—Columnar.

*Angle of spadix tip with peduncle*.—150 degrees to 160 degrees.

*Texture*.—When the spathe is unfurling the spadix is smooth. When the spadix matures, small stigmata protrude. The stigmata are evenly distributed around the spadix. The spadix matures from base to top, slowly giving the spadix a somewhat rough appearance.

*Color*.—Immature: RHS 14A. Mature: RHS 155B. Ages to: RHS 144A.

##### Flowers:

*Quantity per spadix*.—50 to 80.

*Spadix flower arrangement*.—Bisexual, rounded in cross-section.

*Shape*.—Rounded.

*Size*.—Length: 0.05 cm to 0.10 cm. Diameter (maximum): 0.10 cm.

*Color*.—RHS 156D.

Reproductive organs:

*Stamens*.—Not visible.

*Pollen amount*.—Absent.

*Pistil*.—Quantity: Many. Length: Less than 0.01 cm.

Color: RHS 156D.

*Style*.—Not observed to date.

*Stigma*.—Shape: Ovoid. Diameter: Less than 0.01 cm.

Color: RHS 156D.

*Ovary*.—Rarely visible.

*Ovary color*.—Not measured.

Fruit and seed set: None observed to date.

Disease and pest resistance: No specific resistance or susceptibility observed to pathogens or pests common to *Anthurium* under commercial conditions to date.

COMPARISON WITH PARENTAL AND SIMILAR COMMERCIAL VARIETIES

‘ANTHGOUWHI’ differs from the female parent plant ‘15108-02’ (unpatented) in that ‘ANTHGOUWHI’ has short spadices and an angle of the spadix tip with the peduncle of

150 degrees to 160 degrees, whereas ‘15108-02’ has long spadices and an angle of the spadix tip with the peduncle of 170 degrees to 180 degrees.

‘ANTHGOUWHI’ differs from the male parent plant ‘12902-01’ (unpatented) in that ‘ANTHGOUWHI’ has white spathes with weak blistering, whereas ‘12902-01’ has red spathes with medium blistering.

‘ANTHGOUWHI’ differs from similar commercial variety ‘ANTHGROXOL’ (U.S. Plant Pat. No. 32,777) in that ‘ANTHGOUWHI’ has short spadices and acuminate spathe apices, whereas ‘ANTHGROXOL’ has medium spadices and mucronate spathe apices.

‘ANTHGOUWHI’ differs from similar variety ‘ANTH-CAPBUK’ (unpatented) in that ‘ANTHGOUWHI’ has short spadices and acuminate spathe apices, whereas ‘ANTH-CAPBUK’ has medium spadices and mucronate spathe apices.

I claim:

1. A new and distinct variety of *Anthurium* plant named ‘ANTHGOUWHI’, substantially as illustrated and described herein.

\* \* \* \* \*




FIG. 1



FIG. 2




FIG. 3