



(86) Date de dépôt PCT/PCT Filing Date: 2005/04/20
(87) Date publication PCT/PCT Publication Date: 2005/11/10
(45) Date de délivrance/Issue Date: 2012/12/11
(85) Entrée phase nationale/National Entry: 2006/10/17
(86) N° demande PCT/PCT Application No.: EP 2005/004199
(87) N° publication PCT/PCT Publication No.: 2005/105754
(30) Priorité/Priority: 2004/04/28 (EP04101792.2)

(51) Cl.Int./Int.Cl. *C07D 231/12* (2006.01),
A61K 31/41 (2006.01), *A61P 3/00* (2006.01)


(72) Inventeurs/Inventors:
ACKERMANN, JEAN, CH;
AEBI, JOHANNES, CH;
BINGGELI, ALFRED, CH;
GRETHER, UWE, DE;
HIRTH, GEORGES, FR;
KUHN, BERND, CH;
...

(73) Propriétaire/Owner:
F. HOFFMANN-LA ROCHE AG, CH


(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : DERIVES DE PHENYL PYRAZOLE UTILISES EN TANT QU'ACTIVATEURS DES PPAR

(54) Title: PYRAZOLE PHENYL DERIVATIVES AS PPAR ACTIVATORS



I



II

(57) Abrégé/Abstract:

This invention is concerned with compounds of the formula (I) wherein one of R⁵, R⁶ and R⁷ is (a) and X¹, X², R¹ to R¹⁵ and n are as defined in the description, and pharmaceutically acceptable salts and/or esters thereof. The invention further relates to pharmaceutical compositions containing such compounds, to a process for their preparation and to their use for the treatment and/or prevention of diseases which are modulated by PPAR δ and/or PPAR α agonists.



(72) **Inventeurs(suite)/Inventors(continued)**: MAERKI, HANS-PETER, CH; MEYER, MARKUS, DE; MOHR, PETER, CH;
WRIGHT, MATTHEW BLAKE, CH

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 November 2005 (10.11.2005)

PCT

(10) International Publication Number
WO 2005/105754 A1

(51) International Patent Classification⁷: **C07D 231/12**,
A61K 31/41, A61P 3/00

(21) International Application Number:
PCT/EP2005/004199

(22) International Filing Date: 20 April 2005 (20.04.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04101792.2 28 April 2004 (28.04.2004) EP

(71) Applicant (for all designated States except US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ACKERMANN, Jean** [CH/CH]; Im Gloeglihof 16, CH-4125 Riehen (CH). **AEBI, Johannes** [CH/CH]; Wallstrasse 13, CH-4051 Basel (CH). **BINGGELI, Alfred** [CH/CH]; Im Kugelfang 50, CH-4102 Binningen (CH). **GRETHER, Uwe** [DE/DE]; Kirchgasse 4, 79588 Efringen-Kirchen (DE). **HIRTH, Georges** [FR/FR]; 17, rue de la Fecht, F-68000 Colmar (FR). **KUHN, Bernd** [DE/CH]; Sonnmattweg 42, CH-4410 Liestal (CH). **MAERKI, Hans-Peter** [CH/CH]; Seltisbergerstrasse 75, CH-4059 Basel (CH). **MEYER,**

Markus [DE/DE]; Erasmusstrasse 23, 79395 Neuenburg (DE). **MOHR, Peter** [CH/CH]; Bekenstrasse 26, CH-4054 Basel (CH). **WRIGHT, Matthew, Blake** [US/CH]; Peter Rot-Strasse 76, CH-4058 Basel (CH).

(74) Agent: **KLOSTERMEYER, Dörte**; Grenzacherstrasse 124, CH-4070 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.



(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

[Continued on next page]

(54) Title: PYRAZOLE PHENYL DERIVATIVES AS PPAR ACTIVATORS



(57) Abstract: This invention is concerned with compounds of the formula (I) wherein one of R⁵, R⁶ and R⁷ is (a) and X¹, X², R¹ to R¹⁵ and n are as defined in the description, and pharmaceutically acceptable salts and/or esters thereof. The invention further relates to pharmaceutical compositions containing such compounds, to a process for their preparation and to their use for the treatment and/or prevention of diseases which are modulated by PPAR δ and/or PPAR α agonists.

WO 2005/105754 A1


WO 2005/105754 A1



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Pyrazole phenyl derivatives as PPAR activators

The present invention is concerned with novel phenyl derivatives of the formula



and pharmaceutically acceptable salts and/or esters thereof, wherein

X^1 is selected from the group consisting of O, S, and CH_2 ;

5 R^1 is hydrogen or C_{1-7} -alkyl;

R^2 is hydrogen or C_{1-7} -alkyl,


or, if X^1 is CH_2 , R^2 is selected from the group consisting of hydrogen, C_{1-7} -alkyl and C_{1-7} -alkoxy;

R^3 is hydrogen or C_{1-7} -alkyl;

10 R^4 and R^8 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

15 R^5 , R^6 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

and one of R^5 , R^6 and R^7 is



wherein

X^2 is selected from the group consisting of S, O, and NR^9 ,

5 R^9 is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, fluoro- C_{1-7} -alkyl, hydroxy- C_{2-7} -alkyl and C_{1-7} -alkoxy- C_{2-7} -alkyl;

R^{10} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl and fluoro- C_{1-7} -alkyl;

10 R^{11} is selected from the group consisting of hydrogen, C_{1-7} -alkyl and C_{1-7} -alkoxy- C_{1-7} -alkyl;

one of R^{12} or R^{13} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, C_{2-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl and fluoro- C_{1-7} -alkyl; and the other one is a lone pair;

15 R^{14} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl and fluoro- C_{1-7} -alkyl;

R^{15} is 4-trifluoromethoxyphenyl; and

n is 1, 2 or 3.

20 It has been found that compounds of formula I are useful as lipid modulators and insulin sensitizers. In particular, compounds of formula I are PPAR activators.

Peroxisome Proliferator Activated Receptors (PPARs) are members of the nuclear hormone receptor superfamily. The PPARs are ligand-activated transcription factors that regulate gene expression and control multiple metabolic pathways. Three subtypes have been described which are $PPAR\alpha$, $PPAR\delta$ (also known as $PPAR\beta$), and $PPAR\gamma$. $PPAR\delta$ is 25 ubiquitously expressed. $PPAR\alpha$ is predominantly expressed in the liver, kidney and heart. There are at least two major isoforms of $PPAR\gamma$. $PPAR\gamma 1$ is expressed in most tissues, and the longer isoform, $PPAR\gamma 2$ is almost exclusively expressed in adipose tissue.

The PPARs modulate a variety of physiological responses including regulation of glucose- and lipid- homeostasis and metabolism, energy balance, cell differentiation, inflammation and cardiovascular events.

Approximately half of all patients with coronary artery disease have low
5 concentrations of plasma HDL cholesterol. The atheroprotective function of HDL was first highlighted almost 25 years ago and stimulated exploration of the genetic and environmental factors that influence HDL levels. The protective function of HDL comes from its role in a process termed reverse cholesterol transport. HDL mediates the removal of cholesterol from cells in peripheral tissues including those in the
10 atherosclerotic lesions of the arterial wall. HDL then delivers its cholesterol to the liver and sterol-metabolizing organs for conversion to bile and elimination. Data from the Framingham study showed that HDL-C levels are predictive of coronary artery disease risk independently of LDL-C levels. The estimated age-adjusted prevalence among Americans age 20 and older who have HDL-C of less than 35 mg/dl is 16% (males) and
15 5.7% (females). A substantial increase of HDL-C is currently achieved by treatment with niacin in various formulations. However, the substantial side-effects limit the therapeutic potential of this approach.

As many as 90% of the 14 million diagnosed type 2 diabetic patients in the US are overweight or obese, and a high proportion of type 2 diabetic patients have abnormal
20 concentrations of lipoproteins. The prevalence of total cholesterol > 240 mg/dl is 37% in diabetic men and 44% in women. The respective rates for LDL-C > 160 mg/dl are 31% and 44%, respectively, and for HDL-C < 35 mg/dl 28% and 11%, respectively. Diabetes is a disease in which a patient's ability to control glucose levels in blood is decreased because of partial impairment in response to the action of insulin. Type II diabetes
25 (T2D) is also called non-insulin dependent diabetes mellitus (NIDDM) and afflicts 80-90 % of all diabetic patients in developed countries. In T2D, the pancreatic Islets of Langerhans continue to produce insulin. However, the target organs for insulin action, mainly muscle, liver and adipose tissue, exhibit a profound resistance to insulin stimulation. The body continues to compensate by producing unphysiologically high
30 levels of insulin, which ultimately decreases in later stage of disease, due to exhaustion and failure of pancreatic insulin-producing capacity. Thus T2D is a cardiovascular-metabolic syndrome associated with multiple comorbidities including insulin resistance, dyslipidemia, hypertension, endothelial dysfunction and inflammatory atherosclerosis.

First line treatment for dyslipidemia and diabetes generally involves a low-fat and
35 low-glucose diet, exercise and weight loss. However, compliance can be moderate, and as the disease progresses, treatment of the various metabolic deficiencies becomes necessary with e.g. lipid-modulating agents such as statins and fibrates for dyslipidemia and

hypoglycemic drugs, e.g. sulfonylureas or metformin for insulin resistance. A promising new class of drugs has recently been introduced that resensitizes patients to their own insulin (insulin sensitizers), thereby restoring blood glucose and triglyceride levels to normal, and in many cases, obviating or reducing the requirement for exogenous insulin.

5 Pioglitazone (ActosTM) and rosiglitazone (AvandiaTM) belong to the thiazolidinedione (TZD) class of PPAR γ -agonists and were the first in their class to be approved for NIDDM in several countries. These compounds, however, suffer from side effects, including rare but severe liver toxicity (as seen with troglitazone). They also increase body weight in patients. Therefore, new, more efficacious drugs with greater safety and

10 lower side effects are urgently needed. Recent studies provide evidence that agonism of PPAR δ would result in compounds with enhanced therapeutic potential, i. e. such compounds should improve the lipid profile, with a superior effect on HDL-C raising compared to current treatments and with additional positive effects on normalization of insulin-levels (Oliver et al; Proc Nat Acad Sci USA 2001; 98: 5306-11). Recent

15 observations also suggest that there is a independent PPAR α mediated effect on insulin-sensitization in addition to its well known role in reducing triglycerides (Guerre-Millo et al; J Biol Chem 2000; 275: 16638-16642). Thus selective PPAR δ agonists or PPAR δ agonists with additional PPAR α activity may show superior therapeutic efficacy without the side-effects such as the weight gain seen with PPAR γ agonists.

20 The novel compounds of the present invention exceed the compounds known in the art, inasmuch as they bind to and selectively activate PPAR δ or coactivate PPAR δ and PPAR α simultaneously and very efficiently, and with much improved pharmacokinetic properties. Therefore, these compounds combine the anti-dyslipidemic and anti-glycemic effects of PPAR δ and PPAR α activation with no effect on PPAR γ .

25 Consequently, HDL cholesterol is increased, triglycerides lowered (=improved lipid profile) and plasma glucose and insulin are reduced (=insulin sensitization). In addition, such compounds may also lower LDL cholesterol, decrease blood pressure and counteract inflammatory atherosclerosis. Furthermore, such compounds may also be useful for treating inflammatory diseases such as rheumatoid arthritis, osteoarthritis, and

30 psoriasis. Since multiple facets of combined dyslipidemia and the T2D disease syndrome are addressed by PPAR δ -selective agonists and PPAR δ and α coagonists, they are expected to have an enhanced therapeutic potential compared to the compounds already known in the art.

The compounds of the present invention further exhibit improved

35 pharmacological properties compared to known compounds.

Unless otherwise indicated the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

The term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon atoms.

5 The term "lower alkyl" or "C₁₋₇-alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent alkyl radical of one to seven carbon atoms, preferably one to four carbon atoms. This term is further exemplified by such radicals as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, t-butyl and the groups specifically exemplified herein.

10 The term "halogen" refers to fluorine, chlorine, bromine and iodine.

The term "fluoro-lower alkyl" or "fluoro-C₁₋₇-alkyl" refers to lower alkyl groups which are mono- or multiply substituted with fluorine. Examples of fluoro-lower alkyl groups are e.g. -CF₃, -CH₂CF₃, -CH(CF₃)₂ and the groups specifically exemplified herein.

15 The term "alkoxy" refers to the group R'-O-, wherein R' is alkyl. The term "lower-alkoxy" or "C₁₋₇-alkoxy" refers to the group R'-O-, wherein R' is lower-alkyl. Examples of lower-alkoxy groups are e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy and hexyloxy. Preferred are the lower-alkoxy groups specifically exemplified herein.

The term "lower fluoroalkoxy" or "fluoro-C₁₋₇-alkoxy" refers to lower alkoxy groups as defined above which are mono- or multiply substituted with fluorine.
20 Examples of lower fluoroalkoxy groups are e.g. -OCF₃, and -OCH₂CF₃.

The term "lower alkenyl" or "C₂₋₇-alkenyl", alone or in combination, signifies a straight-chain or branched hydrocarbon residue comprising an olefinic bond and up to 7, preferably up to 6, particularly preferred up to 4 carbon atoms. Examples of alkenyl groups are ethenyl, 1-propenyl, 2-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl
25 and isobutenyl. A preferred example is 2-propenyl.

The term "lower alkynyl" or "C₂₋₇-alkynyl", alone or in combination, signifies a straight-chain or branched hydrocarbon residue comprising a triple bond and up to 7, preferably up to 6, particularly preferred up to 4 carbon atoms. Examples of alkynyl groups are ethynyl, 1-propynyl, or 2-propynyl.

30 The term "cycloalkyl" or "C₃₋₇-cycloalkyl" denotes a saturated carbocyclic group containing from 3 to 7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

The term "aryl" relates to the phenyl or naphthyl group, preferably the phenyl

group, which can optionally be mono- or multiply-substituted, particularly mono- or di-substituted by halogen, hydroxy, CN, CF₃, NO₂, NH₂, N(H, lower-alkyl), N(lower-alkyl)₂, carboxy, aminocarbonyl, lower-alkyl, lower fluoro-alkyl, lower-alkoxy, lower fluoro-alkoxy, aryl and/or aryloxy. Preferred substituents are halogen, -CF₃, -OCF₃,
5 lower-alkyl and/or lower-alkoxy. Preferred are the specifically exemplified aryl groups.

The term "heteroaryl" refers to an aromatic 5- or 6-membered ring which can comprise 1, 2 or 3 atoms selected from nitrogen, oxygen and/or sulphur such as furyl, pyridyl, 1,2-, 1,3- and 1,4-diazinyl, thienyl, isoxazolyl, oxazolyl, imidazolyl, or pyrrolyl. The term "heteroaryl" further refers to bicyclic aromatic groups comprising two 5- or 6-
10 membered rings, in which one or both rings can contain 1, 2 or 3 atoms selected from nitrogen, oxygen or sulphur such as e.g. indole or quinoline, or partially hydrogenated bicyclic aromatic groups such as e.g. indolinyl. A heteroaryl group may have a substitution pattern as described earlier in connection with the term "aryl". Preferred heteroaryl groups are e.g. thienyl and furyl which can optionally be substituted as
15 described above, preferably with halogen, lower fluoro-alkyl such as -CF₃, lower fluoro-alkoxy such as -OCF₃, lower-alkyl and/or lower-alkoxy.

A "lone pair" is a pair of electrons in the outermost shell of an atom, in particular a nitrogen atom, that are not used in bonding.

The term "protecting group" refers to groups such as e.g. acyl, alkoxycarbonyl, aryloxy carbonyl, silyl, or imine-derivatives, which are used to temporarily block the
20 reactivity of functional groups. Well known protecting groups are e.g. t-butyloxycarbonyl, benzyloxycarbonyl, fluorenylmethyloxycarbonyl or diphenylmethylene which can be used for the protection of amino groups, or lower-alkyl-, β -trimethylsilylethyl- and β -trichloroethyl-esters, which can be used for the
25 protection of carboxy groups.

"Isomers" are compounds that have identical molecular formulae but that differ in the nature or the sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed
"stereoisomers". Stereoisomers that are not mirror images of one another are termed
30 "diastereoisomers", and stereoisomers that are non-superimposable mirror images are termed "enantiomers", or sometimes optical isomers. A carbon atom bonded to four nonidentical substituents is termed a "chiral center".


The term "pharmaceutically acceptable salts" embraces salts of the compounds of formula (I) with pharmaceutically acceptable bases such as alkali salts, e.g. Na- and K-
35 salts, alkaline earth salts, e.g. Ca- and Mg-salts, and ammonium or substituted

ammonium salts, such as e.g. trimethylammonium salts. The term "pharmaceutically acceptable salts" also relates to such salts.

The compounds of formula (I) can also be solvated, e.g. hydrated. The solvation can be effected in the course of the manufacturing process or can take place e.g. as a consequence of hygroscopic properties of an initially anhydrous compound of formula (I) (hydration). The term pharmaceutically acceptable salts also includes pharmaceutically acceptable solvates.

The term "pharmaceutically acceptable esters" embraces derivatives of the compounds of formula (I), in which a carboxy group has been converted to an ester. Lower-alkyl, hydroxy-lower-alkyl, lower-alkoxy-lower-alkyl, amino-lower-alkyl, mono- or di-lower-alkyl-amino-lower-alkyl, morpholino-lower-alkyl, pyrrolidino-lower-alkyl, piperidino-lower-alkyl, piperazino-lower-alkyl, lower-alkyl-piperazino-lower-alkyl and aralkyl esters are examples of suitable esters. The methyl, ethyl, propyl, butyl and benzyl esters are preferred esters. The methyl and ethyl esters are especially preferred. The term "pharmaceutically acceptable esters" furthermore embraces compounds of formula (I) in which hydroxy groups have been converted to the corresponding esters with inorganic or organic acids such as, nitric acid, sulphuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic acid, succinic acid, tartaric acid, methanesulphonic acid, p-toluenesulphonic acid and the like, which are non toxic to living organisms.

In detail, the present invention relates to compounds of formula (I)



wherein

X^1 is selected from the group consisting of O, S and CH_2 ;

R^1 is hydrogen or C_{1-7} -alkyl;


R^2 is hydrogen or C_{1-7} -alkyl,
or, if X^1 is CH_2 , R^2 is selected from the group consisting of hydrogen, C_{1-7} -alkyl and C_{1-7} -alkoxy;

R^3 is hydrogen or C_{1-7} -alkyl;

R^4 and R^8 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

R^5 , R^6 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

and one of R^5 , R^6 and R^7 is



wherein

X^2 is selected from the group consisting of S, O and NR^9 ;

R^9 is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, fluoro- C_{1-7} -alkyl, hydroxy- C_{2-7} -alkyl and C_{1-7} -alkoxy- C_{2-7} -alkyl;

R^{10} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl and fluoro- C_{1-7} -alkyl;

R^{11} is selected from the group consisting of hydrogen, C_{1-7} -alkyl and C_{1-7} -alkoxy- C_{1-7} -alkyl;

one of R^{12} or R^{13} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, C_{2-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl and fluoro- C_{1-7} -alkyl; and the other one is a lone pair;


R^{14} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl and fluoro- C_{1-7} -alkyl;

R^{15} is 4-trifluoromethoxyphenyl;

n is 1, 2 or 3; and

all isomers and pharmaceutically acceptable salts and/or esters thereof.

Preferred compounds of formula I of the present invention are compounds of formula



5 wherein


X^1 , X^2 , R^1 to R^4 , R^8 , R^{10} to R^{15} and n are as defined herein before;

R^5 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl
 10 and cyano; and

all isomers and pharmaceutically acceptable salts and/or esters thereof.

More preferred are those compounds of formula I-A in accordance with the present invention, wherein at least one of R^4 , R^5 , R^7 and R^8 is C_{1-7} -alkyl or C_{1-7} -alkoxy, with those compounds of formula I-A wherein R^4 is C_{1-7} -alkyl or C_{1-7} -alkoxy being
 15 especially preferred. Even more preferred are those compounds of formula I-A, wherein one of R^4 and R^8 is methyl and the other one hydrogen.

Also preferred are compounds of formula I having the formula



wherein


X^1, X^2, R^1 to R^4, R^8, R^{10} to R^{15} and n are as defined herein before;

R^5 and R^6 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano; and

all isomers and pharmaceutically acceptable salts and/or esters thereof.

More preferred are those compounds of formula I-B in accordance with the present invention, wherein at least one of R^4, R^5, R^6 and R^8 is C_{1-7} -alkyl or C_{1-7} -alkoxy, with those compounds of formula I-B wherein R^4 is C_{1-7} -alkyl or C_{1-7} -alkoxy being especially preferred. Even more preferred are those compounds of formula I-B, wherein one of R^4 and R^8 is methyl and the other one hydrogen.

Further preferred compounds of formula I have the formula



wherein

X^1, X^2, R^1 to R^4, R^8, R^{10} to R^{15} and n are as defined herein before;

R^6 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano; and

all isomers and pharmaceutically acceptable salts and/or esters thereof.

More preferred are those compounds of formula I-C in accordance with the present invention, wherein at least one of R^4, R^6, R^7 and R^8 is C_{1-7} -alkyl or C_{1-7} -alkoxy, with those compounds of formula I-C wherein R^4 is C_{1-7} -alkyl or C_{1-7} -alkoxy being

especially preferred. Even more preferred are those compounds of formula I-C, wherein one of R^4 and R^8 is methyl and the other one hydrogen.

Preferred compounds of formula I are those, wherein R^1 is hydrogen.

X^1 is selected from the group consisting of O, S and CH_2 . Compounds of formula I, wherein X^1 is O are preferred. More preferred are those compounds of formula I, wherein X^1 is O and at least one of R^2 and R^3 is C_{1-7} -alkyl with those compounds of formula I wherein X^1 is O and R^2 and R^3 are C_{1-7} -alkyl being especially preferred.

Also preferred are compounds of formula I, wherein X^1 is CH_2 .


X^2 is selected from the group consisting of S, O and NR^9 . Preferred are compounds of formula I, wherein X^2 is O.

R^{10} is selected from hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl or fluoro- C_{1-7} -alkyl and R^{11} is selected from hydrogen, C_{1-7} -alkyl or C_{1-7} -alkoxy- C_{1-7} -alkyl. Preferred are compounds of formula I, wherein R^{10} and R^{11} are hydrogen.

The integer n is 1, 2 or 3. Preferred are compounds of formula I, wherein n is 1. Further preferred are compounds of formula I, wherein n is 2.

Furthermore, compounds of formula I are also preferred, wherein n is 3.


Further preferred compounds are those compounds of formula I, wherein one of R^5 , R^6 and R^7 is



and X^2 , R^{10} to R^{12} , R^{14} , R^{15} and n are as defined herein before.


Especially preferred are those compounds, wherein R^{12} is C_{1-7} -alkyl or fluoro- C_{1-7} -alkyl.

Also especially preferred are compounds of formula I, wherein one of R^5 , R^6 and R^7 is



and R¹⁰ to R¹², R¹⁴, R¹⁵ and n are as defined herein above.

Also preferred are compounds of formula I, wherein one of R⁵, R⁶ and R⁷ is



5 and X², R¹⁰, R¹¹, R¹³ to R¹⁵ and n are as defined herein before.

Examples of preferred compounds of formula I are the following:

- 2-methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid,
- 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid,
- 10 2-methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid,
- 3-{2-methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid,
- 15 2-{2,3-dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid,
- 2-methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid,
- 2-methyl-2-{2-methyl-4-[1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid,
- 20 2-{2,5-dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid,
- 2-{4-[1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid,
- 25 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid,
- 2-(2,5-dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-

ethoxy}-phenoxy)-2-methyl-propionic acid,
 2-methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-
 2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid,
 2-{5-methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-
 5 ylmethoxy]-phenoxy}-2-methyl-propionic acid,
 2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-
 yl]-propoxy}-phenoxy)-2-methyl-propionic acid, and
 2-{4-[2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-
 methyl-phenoxy}-2-methyl-propionic acid.

10 Particularly preferred compounds of formula I of the present invention are the following:

2-methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-
 ylmethoxy]-phenoxy}-propionic acid,
 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-
 15 ethoxy}-phenoxy)-propionic acid,
 2-methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-
 phenoxy)-propionic acid,
 2-{2,3-dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-
 phenoxy}-2-methyl-propionic acid,
 20 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-
 propoxy}-phenoxy)-propionic acid,
 2-(2,5-dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-
 ethoxy}-phenoxy)-2-methyl-propionic acid,
 2-methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-
 25 2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid,
 2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-
 yl]-propoxy}-phenoxy)-2-methyl-propionic acid, and
 2-{4-[2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-
 methyl-phenoxy}-2-methyl-propionic acid.

30 Especially preferred are also the following compounds of formula I of the present invention:

2-methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-
 ylmethoxy]-phenoxy}-propionic acid,
 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-
 35 ethoxy}-phenoxy)-propionic acid,
 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-
 propoxy}-phenoxy)-propionic acid, and

2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid.

Furthermore, the pharmaceutically acceptable salts of the compounds of formula I and the pharmaceutically acceptable esters of the compounds of formula I individually
5 constitute preferred embodiments of the present invention.


Compounds of formula I can have one or more asymmetric carbon atoms and can exist in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates. The optically
10 active forms can be obtained for example by resolution of the racemates, by asymmetric synthesis or asymmetric chromatography (chromatography with a chiral adsorbent or eluant). The invention embraces all of these forms.

Compounds of formula I may also contain C₁₋₇-alkenyl groups. All forms of cis- and trans-isomers are embraced by the present invention.

It will be appreciated, that the compounds of general formula I in this invention may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compound in vivo. Physiologically acceptable and metabolically labile derivatives, which are capable of producing the parent compounds of general formula I in vivo are also within the scope of this invention.
15


A further aspect of the present invention is the process for the manufacture of compounds of formula (I) as defined above, which process comprises
20

reacting a compound of formula




wherein R¹ is C₁₋₇-alkyl, R², R³, R⁴ and R⁸ are as defined as above and R⁵, R⁶ and R⁷ are
25 selected from hydrogen, C₁₋₇-alkyl, C₁₋₇-alkoxy, C₃₋₇-cycloalkyl, halogen, C₁₋₇-alkoxy-C₁₋₇-alkyl, C₂₋₇-alkenyl, C₂₋₇-alkynyl, fluoro-C₁₋₇-alkyl, fluoro-C₁₋₇-alkoxy, cyano-C₁₋₇-alkyl, and cyano with the proviso that one of R⁵, R⁶ or R⁷ is -OH, -SH or -NHR⁹, wherein R⁹ is as defined above,

with a compound of formula



wherein R^{10} to R^{15} and n are as defined as above and R^{16} is $-OH$, $-Cl$, $-Br$, $-I$ or another leaving group, to obtain a compound of formula



5

wherein R^1 is C_{1-7} -alkyl, X^1 and R^2 to R^8 are as defined as above, and optionally hydrolysing the ester group to obtain a compound of formula I, wherein R^1 is hydrogen.

As described above, the compounds of formula (I) of the present invention can be used as medicaments for the treatment and/or prevention of diseases which are modulated by $PPAR\delta$ and/or $PPAR\alpha$ agonists. Examples of such diseases are diabetes, particularly non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, particularly low HDL-cholesterol, high LDL-cholesterol, or high triglyceride levels, atherosclerotic diseases, metabolic syndrome, syndrome X, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases (such as e.g. Crohn's disease, inflammatory bowel disease, colitis, pancreatitis, cholestasis/fibrosis of the liver, rheumatoid arthritis, osteoarthritis, psoriasis and other skin disorders, and diseases that have an inflammatory component such as e.g. Alzheimer's disease or impaired/improvable cognitive function) and proliferative diseases (cancers such as e.g. liposarcoma, colon cancer, prostate cancer, pancreatic cancer and breast cancer). The use as medicament for the treatment of low HDL cholesterol levels, high LDL cholesterol levels, high triglyceride levels, metabolic syndrome and syndrome X is preferred.

The invention therefore also relates to pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

25

Further, the invention relates to compounds as defined above for use as therapeutically active substances, particularly as therapeutic active substances for the treatment and/or prevention of diseases which are modulated by PPAR δ and/or PPAR α agonists. Examples of such diseases are diabetes, particularly non-insulin dependent
5 diabetes mellitus, increased lipid and cholesterol levels, particularly low HDL-cholesterol, high LDL-cholesterol, or high triglyceride levels, atherosclerotic diseases, metabolic syndrome, syndrome X, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases such as rheumatoid arthritis, osteoarthritis, psoriasis and other skin disorder, and proliferative
10 diseases.

In another embodiment, the invention relates to a method for the treatment and/or prevention of diseases which are modulated by PPAR δ and/or PPAR α agonists, which method comprises administering a compound of formula (I) to a human or animal. Preferred examples of such diseases are diabetes, particularly non-insulin
15 dependent diabetes mellitus, increased lipid and cholesterol levels, particularly low HDL-cholesterol, high LDL-cholesterol, or high triglyceride levels, atherosclerotic diseases, metabolic syndrome, syndrome X, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases such as rheumatoid arthritis, osteoarthritis, psoriasis and other skin disorder, and proliferative
20 diseases.


The invention further relates to the use of compounds as defined above for the treatment and/or prevention of diseases which are modulated by PPAR δ and/or PPAR α agonists. Preferred examples of such diseases are diabetes, particularly non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, particularly low HDL-
25 cholesterol, high LDL-cholesterol, or high triglyceride levels, atherosclerotic diseases, metabolic syndrome, syndrome X, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases such as rheumatoid arthritis, osteoarthritis, psoriasis and other skin disorder, and proliferative diseases.

30 In addition, the invention relates to the use of compounds as defined above for the preparation of medicaments for the treatment and/or prevention of diseases which are modulated by PPAR δ and/or PPAR α agonists. Preferred examples of such diseases are diabetes, particularly non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, particularly low HDL-cholesterol, high LDL-cholesterol, or high
35 triglyceride levels, atherosclerotic diseases, metabolic syndrome, syndrome X, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases such as rheumatoid arthritis, osteoarthritis,

psoriasis and other skin disorder, and proliferative diseases. Such medicaments comprise a compound as defined above.

The compounds of formula (I) can be manufactured by the methods given below, by the methods given in the examples or by analogous methods. Appropriate reaction conditions for the individual reaction steps are known to a person skilled in the art. Starting materials are either commercially available or can be prepared by methods analogous to the methods given below, by methods described in references cited in the text or in the examples, or by methods known in the art.

Scheme 1




10

Hydroxy aldehydes or hydroxy aryl alkyl ketones 1 are known or can be prepared by methods known in the art. Reaction of phenols 1 with α halo esters of formula 2 in the presence of a base like potassium or cesium carbonate in solvents like acetone, methyl-ethyl ketone, acetonitrile or N,N-dimethylformamide in a temperature range between room temperature and 140 °C leads to the corresponding ether compounds 3 (steps a). *Baeyer Villiger* oxidation e. g. with meta chloro perbenzoic acid in a solvent like dichloromethane, leads to compounds 4 (step b). Pyrazoles 5 (prepared as outlined in schemes 6 to 9) are condensed with phenols 4 according to well known procedures (step c): if R^{16} represents a hydroxy group e. g. via *Mitsunobu*-reaction, with triphenylphosphine and di-tert-butyl-, diisopropyl- or diethyl-azodicarboxylate as

20


reagents; this transformation is preferably carried out in a solvent like toluene, dichloromethane or tetrahydrofuran at ambient temperature. Alternatively, if R¹⁶ represents a halide, mesylate, tosylate or triflate moiety, pyrazoles 5 can be reacted with phenols 4 in solvents like N,N-dimethylformamide, dimethylsulfoxide, acetonitrile, acetone or methyl-ethyl ketone in the presence of a weak base like cesium or potassium carbonate at a temperature ranging from room temperature to 140 °C, preferably around 50 °C to yield ether compounds Ia (step c). Those can optionally be hydrolyzed according to standard procedures, e. g. by treatment with an alkali hydroxide like LiOH or NaOH in a polar solvent mixture like tetrahydrofuran/ethanol/water leading to carboxylic acids Ia.

An analogous reaction scheme with the same reaction sequences applies for the isomeric compound series leading to compounds of general formula I, particularly compounds according to formula Ib:



The synthesis of compounds with the general structure I, particularly compounds according to formula Ic, with X¹ equal to O and X² equal to nitrogen can be accomplished according to schemes 2 and 3.


Scheme 2



Nitro-phenols 2 of scheme 2 are commercially available, known or can be synthesized from anisols 3 by demethylation with aqueous 62% HBr in acetic acid between RT and 120 °C (step b). Alternatively, phenols 1 can be nitrated in para-position according to well established methods, e. g. with a solution of $NaNO_3$ in water/concentrated hydrochloric acid in a solvent like Et_2O , followed by the addition of acetic acid anhydride at RT [following a procedure of P. Keller, *Bull. Soc. Fr.* 1994, 131, 27-29] leading to phenols 2 (step a). Nitro-phenols 2 are then reduced in an alcohol like EtOH or MeOH with hydrogen in the presence of Pd/C and optionally an acid like HCl or AcOH at RT to give anilines 4 (step c). Intermediates 4 are then O-alkylated with electrophile 5, e.g. a bromo-acetate 5, in the presence of K_2CO_3 or Cs_2CO_3 in a solvent like acetonitrile or acetone between 10 °C and RT to give intermediates 6 (step d). Electrophiles 5 are commercially available or can be synthesized by methods known in the art. Triflates 5 can be prepared from the corresponding alcohols. Anilines 6 can alternatively be synthesized from compounds 5 and nitrophenols 2 in a two step procedure: first by O-alkylation as described above, followed by hydrogenation with Pd/C in an alcohol like MeOH or EtOH optionally in the presence of AcOH or HCl (step e). BOC-protection with di-tert-butyl dicarbonate in tetrahydrofuran at RT to reflux yields compounds 7 (step f). Compounds 7 can also be synthesized directly from

electrophiles 5 and BOC-protected anilines 8 with K_2CO_3 or Cs_2CO_3 as described for the synthesis of compounds 6 (step g).


Scheme 3



- 5 Intermediates 7 of scheme 3 can optionally be alkylated at nitrogen using sodium hydride and a reactive alkyl halogenide/mesylate or triflate to give compounds 9 (step h). Standard BOC-deprotection (TFA/ CH_2Cl_2 , or HCl in dioxane) at 0 °C to RT affords anilines 10 (step i). Reaction with activated pyrazoles 11 (R^{16} being a halide or a methanesulfonate) using sodium hydride or sodium, potassium or cesium carbonate in
- 10 N,N-dimethylformamide, dimethylsulfoxide, dimethylacetamide or tetrahydrofuran, at 0 °C to RT, leads to compounds Ic (step k). Alternatively, pyrazoles 11 with $R^{16}=OH$ can be transformed in situ to the corresponding triflates by treatment with trifluoromethanesulfonic anhydride/2,6-di-tert-butylpyridine in CH_2Cl_2 at 0 °C. These

triflates are then reacted with anilines 10 in the presence of 2,6-di-tert-butylpyridine as base in nitromethane between RT and 60 °C to yield compounds Ic [following a procedure of Belostotskii, Anatoly M., Hassner, A., *Tetrahedron Lett.* 1994, 35(28), 5075-6] (step k). Secondary aniline compounds Ic ($R^9=H$) can be reductively methylated with an aqueous solution of NaH_2PO_3 and formaldehyde between RT and 65 °C [Loibner, H., Pruckner, A., Stuetz, A., *Tetrahedron Lett.* 1984, 25, 2535-2536] to give compounds Ic with $R^9=Me$. Ensuing hydrolysis with aqueous LiOH, NaOH or KOH in tetrahydrofuran/EtOH or another suitable solvent produces compounds Ic of scheme 3 in the form of the free acid.


10 An analogous reaction scheme with the same reaction sequences applies for the isomeric compound series leading to compounds of general formula I, particularly compounds according to formula Id:



As alternative to the sequences described in scheme 2, the nitrogen containing intermediates can be prepared from suitable intermediates carrying a phenolic hydroxyl moiety. In such intermediates, optionally carrying one or more protective functions, the phenolic OH group can be replaced by the corresponding aromatic NH_2 function by methods known in the art. For example by a three step sequence as described in Tetrahedron Letters 43(42), 7617-7619 (2002): i) transformation of the phenol moiety into its trifluoromethanesulfonate (triflic anhydride, 2,6-lutidine, 4-dimethylaminopyridine, dichloromethane, 0 °C to room temperature; ii) treatment of the triflate with benzophenone imine, di-palladium-tris(dibenzylideneacetone) complex, S-(-)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, cesium carbonate, toluene, in a Schlenk tube at temperatures around 120 °C; iii) treatment with catalytic amounts of hydrochloric acid in wet tetrahydrofuran preferably at room temperature to liberate the aromatic NH_2 moiety.

The synthesis of compounds with the general structure I, particularly compounds according to formula Ie, with X^1 equal to CH_2 and X^2 equal to oxygen can be accomplished according to scheme 4.

Scheme 4




Aldehydes 1 are known, commercially available or can be prepared by methods known in the art. Aldehydes 1 can be reacted with a *Wittig* salt 2 such as (1,2-diethoxy-2-oxoethyl)triphenyl phosphonium chloride or (1,2-dimethoxy-2-oxoethyl)triphenyl phosphonium bromide in solvents like isopropanol, dichloromethane or tetrahydrofuran or mixtures thereof in the presence of a base like potassium carbonate, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 1,1,3,3-tetramethyl-guanidine or sodium tert butylate, preferably between 0 °C and the reflux temperature of the solvents, giving
 10 acrylic esters 3 as E and/or Z isomers (step a). Alternatively, a *Horner-Wadsworth-Emmons* reaction can be used for the transformation of compounds 1 into unsaturated esters 3, e. g. using dimethyl(methoxycarbonyl)methyl phosphonate, optionally substituted at the methylene group, and a base like sodium hydride in a solvent like tetrahydrofuran. Hydrogenation of acrylic esters 3 using palladium on charcoal as
 15 catalyst, preferably at room temperature and 1 atm. pressure of hydrogen, in solvents like methanol, ethanol, tetrahydrofuran, acetic acid, dichloromethane and mixtures thereof, affords esters 7, provided that the protecting group can be cleaved reductively (step e).

Alternatively, aldehydes 1 are reacted with the enolate of an acetic acid ester 4 (preferably the lithium-enolate, prepared at -78 °C by treatment of 4 with a strong, non-
 20 nucleophilic base like lithium diisopropylamide in an inert solvent like tetrahydrofuran), preferably at temperatures around -78 °C, in solvents like tetrahydrofuran giving the


aldol product 5 as a mixture of diastereomers (step b). Removal of the benzylic hydroxy group in compounds 5 can be performed with a reducing agent like e. g. triethylsilane in the presence of a Lewis acid, like boron-trifluoride, or a protic acid, like trifluoroacetic acid, in a suitable solvent like trifluoroacetic acid itself or dichloromethane between 0 °C and 60 °C to yield protected phenol compounds 6 (step d). Subsequent removal of the protecting group, e. g. a benzyl group, by standard technology, e. g. catalytic hydrogenation using hydrogen and a catalyst like palladium or by using dimethyl sulfide and boron trifluoride diethyl etherate in a solvent like dichloromethane between room temperature and the reflux temperature of the solvent gives phenolic compounds 7 (step g). Catalytic hydrogenation can be used to transform unsaturated esters 3 into compounds 6 (step f). In case the protective group in compounds 3 is a benzyl group, then a one step hydrogenation procedure directly gives phenolic compounds 7. Catalytic hydrogenation can also be used for the simultaneous removal of the benzylic hydroxy function and a benzyl protecting group, preferably using palladium on charcoal as catalyst in the presence of an acid like oxalic acid in solvents like alcohols at temperatures around room temperature and a hydrogen pressure up to 100 bar, thus giving the transformation of compounds 5 into compounds 7 in one step (step d and g). As an alternative method, compounds 5 can be treated with catalytic amounts of an acid like *p*-toluene sulfonic acid in a solvent like benzene or toluene, preferably under conditions allowing the removal of the water formed (e. g. with a Dean Stark trap or in the presence of molecular sieves) at temperatures between room temperature and the reflux temperature of the solvents to yield acrylic esters 3 (step c). The condensation of phenols 7 with pyrazoles 8 to form compounds Ie can be performed as outlined in scheme 1.

An analogous reaction scheme with the same reaction sequences applies for the isomeric compound series leading to compounds of general formula I, particularly compounds according to formula If:




The synthesis of compounds with the general structure I, particularly compounds according to formula Ig, with X¹ equal to CH₂ and X² equal to nitrogen can be accomplished according to scheme 5.

Scheme 5



Nitro-phenyl compounds 3 and 5 are prepared from nitro aldehydes 1, which are known, commercially available or can be prepared by methods known in the art, e.g. by Wittig/Horner-Wadsworth-Emmons or aldol reactions analogous to the reactions described for the synthesis of compounds 3 and 5 in scheme 4 (steps a and b). Catalytic hydrogenation can be used for the simultaneous removal of the benzylic hydroxy function (compounds 5) or the reduction of the double bond (compounds 3) and the reduction of the nitro group, preferably using palladium on charcoal as catalyst optionally in the presence of an acid like oxalic acid in solvents like alcohols at temperatures around room temperature and a hydrogen pressure up to 100 bar (step c). Compounds 7 with R⁹ substituents different from hydrogen are obtained by first introduction of a BOC group, alkylation and removal of the BOC protective function as described in schemes 2 and 3. The condensation of anilines 7 with pyrazoles 8 to form compounds Ig can be performed as outlined in scheme 3.

An analogous reaction scheme with the same reaction sequences applies for the isomeric compound series leading to compounds of general formula I, particularly compounds according to formula Ih:

I_h


As alternative to the sequences described in scheme 5, the nitrogen containing intermediates can be prepared from suitable intermediates carrying a phenolic hydroxy function. In such intermediates, optionally carrying one or more protective functions, the phenolic OH group can be replaced by the corresponding aromatic NH₂ function by methods known in the art. For example by a three step sequence as described in Tetrahedron Letters 43(42), 7617-7619 (2002) and discussed in the context of schemes 2 and 3.

The synthesis of compounds with the general structure I, particularly compounds with X¹ and / or X² equal to S can be accomplished in close analogy to the synthesis of the corresponding analogues with X¹ and / or X² equal to oxygen. Suitable sulfur containing intermediates are known, can be prepared by methods known in the art or are prepared from phenolic intermediates as described by W Zhi-Liang and AP Kozikowski (*J. Org. Chem.* 2003, 68, 9116-9118): treatment of a phenolic intermediate with sodium thiocyanate, sodium bromide and bromine in a solvent like methanol preferably between 0 °C and room temperature gives the corresponding 4-thiocyanato-phenols; subsequent reduction with lithium aluminium hydride in a solvent like tetrahydrofuran at temperatures around 0 °C then liberates the corresponding 4-mercapto-phenol. Alternatively, intermediates carrying an aromatic SH moiety can be prepared from suitable intermediates carrying a phenolic hydroxy function. In such intermediates, optionally carrying one or more protective functions, the phenolic OH group can be replaced by the corresponding aromatic SH function by methods known in the art. For example by a three step sequence as described in J. Labelled Compounds & Radiopharmaceuticals 43(7), 683-691, (2000): i) transformation of the phenol moiety into its trifluoromethanesulfonate (triflic anhydride, triethylamine, dichloromethane, at low temperature, preferably around -30 °C); ii) treatment of the triflate with triisopropylsilanethiolate, tetrakis(triphenylphosphine)-palladium(0) in solvent mixtures like toluene and tetrahydrofuran in a temperature range between 60 °C and 150 °C; iii) treatment of the silyl sulfide with hydrogen chloride in methanol preferably around 0 °C to liberate the phenolic SH moiety.

Compounds of the general formula I may be obtained in the form of racemates. Racemic compounds can be separated into their antipodes by methods known in the art, such as separation of the antipodes via diastereomeric salts by crystallization with optically pure amines such as e. g. (R) or (S)-1-phenyl-ethylamine, (R) or (S)-1-naphthalen-1-yl-ethylamine, brucine, quinine or quinidine or by separation of the antipodes by specific chromatographic methods using either a chiral adsorbent or a chiral eluent.

Pyrazoles 5 (scheme 1), identical to 11 (scheme 3), 8 (scheme 4) and 8 (scheme 5) are commercially available, known or can be synthesized by methods known in the art. Representative examples of possible syntheses of these key intermediates are given in schemes 6-9.


Scheme 6



Substituted acetophenones and heteroaryl ketones 1 are commercially available, known or can be prepared by methods known in the art. Acylation of compounds 1 with oxalate derivatives can be performed under standard conditions, e. g. with diethyl oxalate in the presence of a base like sodium ethoxide at temperatures between -78 °C and 50 °C in solvents like ethanol, or with lithium hexamethyldisilazide at temperatures between -78 °C and ambient temperature in solvents like ether, to form after subsequent acidification free ethyl pyruvates 2 (step a). Alternatively, pyruvates 2 can be synthesized via i) transforming ketones 1 into the corresponding silyl enol ethers 3, e. g. through treatment with trimethylsilyl chloride in the presence of a base like triethylamine at temperatures between 0 °C and 40 °C in a solvent like acetonitrile (step b); ii) in situ formation of a metal enol ether, e. g. with zinc chloride and subsequent acylation with an acylation reagent like ethyl oxalyl chloride at temperatures between 0 °C and 50 °C in a solvent like toluene or dichloromethane (step c). Pyruvates 2 can be converted to regioisomeric pyrazoles 4 and 5 through condensation with monosubstituted hydrazines $\text{H}_2\text{NNHR}^{12/13}$ which are commercially available, known or can be prepared by methods known in the art, e. g. at temperatures between ambient temperature and the reflux temperature of the solvent in solvents like ethanol (step d). Alternatively, pyrazoles 4 and 5 can be synthesized via i) reacting pyruvates 2 with hydrazine, preferably at reflux temperature in ethanol (step e); ii) conversion of the obtained pyrazole 6 into regioisomeres 4 and 5 under standard conditions, e. g. through alkylation with an alkyl halogenide in the presence of a base like potassium hydroxide at temperatures between -20 °C and the reflux temperature of the solvent in solvents like ethanol (step f). Regioisomeric pyrazoles 4 and 5 can easily be separated by standard techniques, e. g. through column chromatography on silica. Reduction of esters 4 and 5 can be performed by methods well known in the art, e. g. with lithium aluminium hydride at temperatures between 0 °C and the reflux temperature of the solvents in solvents like tetrahydrofuran or diethyl ether (step g).

The alcohol compounds 7 and 8 correspond to or can be converted into compounds of general formula 5 (scheme 1), identical to 11 (scheme 3), 8 (scheme 4) and 8 (scheme 5), e. g. by treatment with methanesulfonyl chloride in dichloromethane in the presence of a base like triethylamine preferably in a temperature range between -20 °C and room temperature, or e. g. by reaction with carbon tetrachloride or carbon tetrabromide and triphenylphosphine in solvents like tetrahydrofuran, preferably in a temperature range between room temperature and the reflux temperature of the solvents.

Scheme 7




Reduction of pyrazole esters **1** (compounds **4**, **5** and **6** in scheme 6), preferably using lithium aluminum hydride in a solvent like ether or tetrahydrofuran, preferably between 0 °C and room temperature, gives primary alcohols **2** (step a), which can be used as such or can be converted into the corresponding halides **3**, e. g. by treatment with methanesulfonyl chloride in dichloromethane in the presence of 2,6-lutidine, preferably

between -20°C and the reflux temperature of dichloromethane, by treatment with thionyl chloride in a solvent like dichloromethane or chloroform, preferably at temperatures between -20°C and $+50^{\circ}\text{C}$, or by treatment with tetrabromomethane and triphenylphosphine in solvents like tetrahydrofuran at temperatures between 0°C and the reflux temperature of tetrahydrofuran (step b). Esters 1 can further be converted into tertiary alcohols 4 with $\text{R}^{10} = \text{R}^{11}$ through reaction with alkyl organometallic reagents, preferably using alkyl Grignard compounds in a solvent like tetrahydrofuran or ether, preferably between -15°C and the reflux temperature of the solvent (step c). Alcohols 4 with R^{10} not equal to R^{11} can be prepared by a sequential procedure: i) saponification to the acid; ii) treatment with R^{10}Li , optionally in the presence of a Cu(I) salt, in ether or tetrahydrofuran to yield the alkyl ketones $-\text{COR}^{10}$; iii) subsequent reaction with R^{11}Li or lithium aluminium hydride in ether or tetrahydrofuran (step c). Primary alcohols 2 can be oxidized to aldehydes 5 by methods known in the art, e. g. by treatment with pyridinium chlorochromate in dichloromethane, preferably at temperatures between room temperature and the reflux temperature of dichloromethane, or by treatment with manganese dioxide in solvents like dichloromethane, preferably at room temperature (step d). These aldehydes 5 can be converted to the corresponding secondary alcohols 6 through reaction with alkyl organometallic compounds, preferably under the conditions given for the transformation of esters 1 to tertiary alcohols 4 (step e). Ketones 7 can be obtained from secondary alcohols 6 by methods known in the art, e. g. by treatment with Cr(VI) reagents like the Jones reagent (Jones et al., *J. Chem. Soc.* 1953, 2548.) (step f). These ketones 7 can be reduced back to the corresponding secondary alcohols 6 in an enantioselective fashion leading to the (R)- or (S)-alcohols 6, e. g. by treatment with borane-dimethylsulfide complex and (S)- or (R)-2-methyl-CBS-oxazaborolidine as chiral catalyst in tetrahydrofuran, preferably at temperatures between -78°C and ambient temperature, according to Corey et al. (E. J. Corey, R. K. Bakshi, S. Shibata, *J. Am. Chem. Soc.* 1987, 109, 5551-5553), or by treatment with (+)- or (-)-B-chlorodiisopinocampheylborane (DIP-Cl), according to Brown et al. (P. V. Ramachandran, B. Gong, A. V. Teodorovic, H. C. Brown, *Tetrahedron: Asymmetry* 1994, 5, 1061-1074) (step g). Ketones 7 can in addition be converted to the corresponding tertiary alcohols 4 through reaction with alkyl organometallic compounds, preferably under the conditions given for the transformation of esters 1 to tertiary alcohols 4 in step c (step h). If the alcohol compounds 2, 4, or 6 contain one or more chiral centers and are not optically pure, they can optionally be separated into optically pure antipodes by methods well known in the art, e. g. chromatography on a chiral HPLC column, or by derivatization with an optically pure acid to form esters, which can then be separated by conventional HPLC chromatography and converted back to the original alcohol.

The alcohol compounds 2, 4, and 6, and the halide compound 3, correspond to or can be converted into compounds of general formula 5 (scheme 1), identical to 11 (scheme 3), 8 (scheme 4) and 8 (scheme 5), e. g. by treatment with methanesulfonyl chloride in dichloromethane in the presence of a base like triethylamine preferably in a temperature range between -20°C and room temperature, or e. g. by reaction with carbon tetrachloride or carbon tetrabromide and triphenylphosphine in solvents like tetrahydrofuran, preferably in a temperature range between room temperature and the reflux temperature of the solvents.

Scheme 8



10

Pyrazole alkanols 1 with a chain length of n carbon atoms can be converted into analogues with a chain length of $n+1$ carbon atoms by methods well known in the art, e. g. by conversion of the primary alcohol function into a suitable leaving group, e. g. a halide (step a), reaction with cyanide ion (step b), saponification (step c) followed by reduction of the acid formed (compounds 4) to the primary alcohols 5, e. g. by using diborane in tetrahydrofuran (step d). In order to introduce substituents R^{10} and/or R^{11} different from hydrogen, cyano intermediates 3 of this elongation process can be reacted with alkyl Grignard reagents R^{10}MgX in solvents like ether or tetrahydrofuran between 0°C and the reflux temperature of the solvent to form the corresponding R^{10}CO -alkyl ketones, which upon treatment with an alkyllithium reagent R^{11}Li or lithium aluminum

20

hydride in solvents like ether or tetrahydrofuran give alcohols 5. $R^{10}CO$ -alkyl ketones can also be reduced, e. g. by treatment with sodium borohydride in alcohol, preferably at temperatures between $-15^{\circ}C$ and $40^{\circ}C$. This reaction can also be carried out in an enantioselective fashion leading to the (R)- or (S)-alcohols 5, e. g. by treatment with


5 borane-dimethylsulfide complex and (S)- or (R)-2-methyl-CBS-oxazaborolidine as chiral catalyst in tetrahydrofuran, preferably at temperatures between $-78^{\circ}C$ and ambient temperature according to Corey et al. (E. J. Corey, R. K. Bakshi, S. Shibata, *J. Am. Chem. Soc.* 1987, 109, 5551-5553), or by treatment with (+)- or (-)-B-chlorodiisopinocampheylborane (DIP-Cl), according to Brown et al. (P. V.

10 Ramachandran, B. Gong, A. V. Teodorovic, H. C. Brown, *Tetrahedron: Asymmetry* 1994, 5, 1061-1074). Alternatively, alcohol compounds 5 which contain one or more chiral centers can optionally be separated into optically pure antipodes by methods well known in the art, e. g. chromatography on a chiral HPLC column, or by derivatization with an optically pure acid to form esters, which can then be separated by conventional HPLC

15 and converted back to the original alcohol. The alcohol compounds 5 correspond to or can be transformed into compounds of general formula 5 (scheme 1), identical to 11 (scheme 3), 8 (scheme 4) and 8 (scheme 5), e. g. by treatment with methanesulfonyl chloride in dichloromethane in the presence of a base like triethylamine, preferably in a temperature range between $-20^{\circ}C$ and room temperature, or e. g. by reaction with

20 carbon tetrachloride or carbon tetrabromide and triphenylphosphine in solvents like tetrahydrofuran, preferably in a temperature range between room temperature and the reflux temperature of the solvents.

Scheme 9



Alcohols 1 (compounds 5 with $R^{14} = H$ and $R^{16} = OH$ in scheme 1, compounds 11 with $R^{14} = H$ and $R^{16} = OH$ in scheme 3, compounds 8 with $R^{14} = H$ and $R^{16} = OH$ in schemes 4 and 5, compounds 7 and 8 with $R^{14} = H$ in scheme 6, compounds 2, 4 and 6 with $R^{14} = H$ in scheme 7, compounds 1 and 5 with $R^{14} = H$ in scheme 8), can be protected by methods known in the literature, e. g. by treating them with tert-butyldimethylsilyl chloride in the presence of imidazole, preferably at room temperature in solvents like N,N-dimethylformamide, to obtain the corresponding tert-butyldimethylsilyl ethers 2 (step a). Halogenation of protected pyrazoles 2, e. g. through reaction with bromine preferably at temperatures between 0 °C and ambient temperature in solvents like dichloromethane delivers 4-halo pyrazoles 3 (step b). Compounds 3 can – following halogen metal exchange, preferably with tert-butyllithium at -78 °C in solvents like tetrahydrofuran - be reacted with alkylating reagents 4 with X e. g. being a chlorine, bromine or iodine atom, preferably with alkyl iodides, at temperatures between -78 °C and ambient temperature in solvents like tetrahydrofuran, to form pyrazoles 5 bearing a substituent in position 4 (step c). Alternatively, transition metal catalyzed reactions can be used to transform 4-halo pyrazoles 3 into compounds 5, e. g. by treatment with a stannane (X being trialkyl stannyl) in the presence of a Pd(0) catalyst like $[Pd_2(dba)_3]$ and triphenyl arsine at temperatures between 0 °C and the reflux temperature of the solvent in solvents like dioxane. Residues R^{14} can further be introduced by i) formylation of pyrazoles 2 through methods well known in the art, e. g. with phosphorus oxychloride and N,N-dimethylformamide preferably at temperatures between 0 °C and 100 °C; ii) subsequent transformation of the intermediate formyl pyrazole to 4-substituted pyrazoles 5, e. g. through reduction with sodium cyano borohydride in the presence of zinc iodide at temperatures between -78 °C and the reflux temperature of the solvent in solvents like diethyl ether (step d). O-Deprotection of compounds 5 leading to building blocks 6 can be performed by methods described in the literature, e. g. by treatment with tetrabutyl ammonium fluoride at temperatures between -15 °C and ambient temperature in a solvent like tetrahydrofuran, if the protecting groups are silyl ethers (step e). The alcohol compounds 6 correspond to or can be transformed into compounds of general formula 5 (scheme 1), identical to 11 (scheme 3), 8 (scheme 4) and 8 (scheme 5), e. g. by treatment with methanesulfonyl chloride in dichloromethane in the presence of a base like triethylamine, preferably in a temperature range between -20 °C and room temperature, or e. g. by reaction with carbon tetrachloride or carbon tetrabromide and triphenylphosphine in solvents like tetrahydrofuran, preferably in a temperature range between room temperature and the reflux temperature of the solvents.

The following tests were carried out in order to determine the activity of the compounds of formula (I).

Background information on the performed assays can be found in: Nichols JS et al. "Development of a scintillation proximity assay for peroxisome proliferator-activated receptor gamma ligand binding domain", (1998) Anal. Biochem. 257: 112-119.

Full-length cDNA clones for humans PPAR δ and PPAR α and mouse PPAR γ were
5 obtained by RT-PCR from human adipose and mouse liver cRNA, respectively, cloned into plasmid vectors and verified by DNA sequencing. Bacterial and mammalian expression vectors were constructed to produce glutathione-s-transferase (GST) and Gal4 DNA binding domain proteins fused to the ligand binding domains (LBD) of PPAR δ (aa 139 to 442), PPAR γ (aa 174 to 476) and PPAR α (aa 167 to 469). To
10 accomplish this, the portions of the cloned sequences encoding the LBDs were amplified from the full-length clones by PCR and then subcloned into the plasmid vectors. Final clones were verified by DNA sequence analysis.

Induction, expression, and purification of GST-LBD fusion proteins were performed in *E. coli* strain BL21(pLysS) cells by standard methods (Ref: Current
15 Protocols in Molecular Biology, Wiley Press, edited by Ausubel et al.).

Radioligand Binding Assay

PPAR δ receptor binding was assayed in HNM10 (50mM Hepes, pH 7.4, 10 mM NaCl, 5mM MgCl₂, 0.15 mg/ml fatty acid-free BSA and 15 mM DTT). For each 96 well reaction a 500 ng equivalent of GST-PPAR δ -LBD fusion protein and radioligand, e.g.
20 20000 dpm {2-methyl-4-[4-methyl-2-(4-trifluoromethyl-phenyl)-thiazol-5-yl-ditritiomethylsulfanyl]-phenoxy}-acetic acid, was bound to 10 μ g SPA beads (PharmaciaAmersham) in a final volume of 50 μ l by shaking. The resulting slurry was incubated for 1h at RT and centrifuged for 2 min at 1300g. The supernatant containing unbound protein was removed and the semidry pellet containing the receptor-coated
25 beads was resuspended in 50 μ l of HNM. Radioligand was added and the reaction incubated at RT for 1h and scintillation proximity counting performed in the presence of test compounds was determined. All binding assays were performed in 96 well plates and the amount of bound ligand was measured on a Packard TopCount using OptiPlates (Packard). Dose response curves were done in triplicates within a range of concentration
30 from 10⁻¹⁰ M to 10⁻⁴ M.

PPAR α receptor binding was assayed in TKE50 (50mM Tris-HCl, pH 8, 50 mM KCl, 2mM EDTA, 0.1 mg/ml fatty acid-free BSA and 10 mM DTT). For each 96 well reaction an 140 ng equivalent of GST-PPAR α -LBD fusion protein was bound to 10 μ g SPA beads (PharmaciaAmersham) in a final volume of 50 μ l by shaking. The resulting
35 slurry was incubated for 1h at RT and centrifuged for 2 min at 1300g. The supernatant

containing unbound protein was removed and the semidry pellet containing the receptor-coated beads was resolved in 50 μ l of TKE. For radioligand binding e.g. 10000 dpm of 2(S)-(2-benzoyl-phenylamino)-3-{4-[1,1-ditritio-2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid or 2,3-ditritio-2(S)-methoxy-3-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-benzo[b]thiophen-7-yl}-propionic acid in 50 μ l were added, the reaction incubated at RT for 1h and scintillation proximity counting performed. All binding assays were performed in 96 well plates and the amount of bound ligand measured on a Packard TopCount using OptiPlates (Packard). Nonspecific binding was determined in the presence of 10^{-4} M unlabelled compound. Dose response curves were done in triplicates within a range of concentration from 10^{-10} M to 10^{-4} M.

PPAR γ receptor binding was assayed in TKE50 (50mM Tris-HCl, pH 8, 50 mM KCl, 2mM EDTA, 0.1 mg/ml fatty acid-free BSA and 10 mM DTT). For each 96 well reaction an 140 ng equivalent of GST-PPAR γ -LBD fusion protein was bound to 10 μ g SPA beads (PharmaciaAmersham) in a final volume of 50 μ l by shaking. The resulting slurry was incubated for 1h at RT and centrifuged for 2 min at 1300g. The supernatant containing unbound protein was removed and the semidry pellet containing the receptor-coated beads was resolved in 50 μ l of TKE. For radioligand binding e.g. 10000 dpm 2(S)-(2-benzoyl-phenylamino)-3-{4-[1,1-ditritio-2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid in 50 μ l were added, the reaction incubated at RT for 1h and scintillation proximity counting performed. All binding assays were performed in 96 well plates and the amount of bound ligand measured on a Packard TopCount using OptiPlates (Packard). Nonspecific binding was determined in the presence of 10^{-4} M unlabelled compound. Dose response curves were done in triplicates within a range of concentration from 10^{-10} M to 10^{-4} M.

25 Luciferase Transcriptional Reporter Gene Assays (Transactivation Assay)

Baby hamster kidney cells (BHK21 ATCC CCL10) were grown in DMEM medium containing 10% FBS at 37 $^{\circ}$ C in a 95%O₂:5%CO₂ atmosphere. Cells were seeded in 6 well plates at a density of 10^5 cells/well and then batch-transfected with either the pFA-PPAR δ -LBD, pFA-PPAR γ -LBD or pFA-PPAR α -LBD expression plasmids plus a reporter plasmid (to monitor transfection efficiency). Transfection was accomplished with the Fugene 6 reagent (Roche Molecular Biochemicals) according to the suggested protocol. Six hours following transfection, the cells were harvested by trypsinization and seeded in 96 well plates at a density of 10^4 cells/well. After 24 hours the medium was removed and replaced with 100 μ l of phenol red-free medium containing the corresponding test compounds or control ligands (at a final DMSO concentration of 0.1%). Following incubation of the cells for 24 hours with compounds, 50 μ l of the supernatant was

discarded and then 50 μ l of Luciferase Constant-Light Reagent (Roche Molecular Biochemicals) was added to lyse the cells and initiate the luciferase reaction. Luminescence for luciferase was measured in a Packard TopCount. Transcriptional activation in the presence of a test substance was expressed as fold-activation over cells
 5 incubated in the absence of the substance. The signals were normalized against plate-specific controls (DMSO alone) and the fold-stimulation of luciferase activity observed with specific and selective PPAR α ((2(S)-2-(2-benzoyl-phenylamino)-3-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid), PPAR γ (Rosiglitazone) and PPAR δ ({2-methyl-4-[4-methyl-2-(4-trifluoromethyl-phenyl)-thiazol-5-ylmethylsulfanyl]-phenoxy}-acetic acid) reference compounds was set to 100 %. EC₅₀
 10 values were calculated using the XLfit program (ID Business Solutions Ltd. UK).

The free acids of the compounds of the present invention (R¹ is hydrogen) exhibit IC₅₀ values of 0.1 nM to 10 μ M, preferably 1 nM to 500 nM for PPAR δ and/or IC₅₀ values of 1 nM to 10 μ M, preferably 10 nM to 500 nM for PPAR α . Further the free acids of the
 15 compounds of the present invention (R¹ is hydrogen) exhibit EC₅₀ values of 1 nM to 10 μ M, preferably 10 nM to 1 μ M for PPAR δ and/or EC₅₀ values of 1 nM to 10 μ M, preferably 10 nM to 1 μ M for PPAR α . Compounds, in which R¹ is not hydrogen are converted in vivo to compounds in which R¹ is hydrogen. The following table shows measured values for some selected compounds of the present invention.

	PPAR α IC ₅₀ (μ mol/l)	PPAR γ IC ₅₀ (μ mol/l)	PPAR δ IC ₅₀ (μ mol/l)	PPAR α EC ₅₀ (μ mol/l) (fold activation [%])	PPAR δ EC ₅₀ (μ mol/l) (fold activation [%])
Example 1	0.166	10	0.144	0.38 (100)	0.2 (59)
Example 6	0.212	10	0.222	0.16 (120)	0.096 (140)

20

The compounds of formula (I) and their pharmaceutically acceptable salts and esters can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine
 25 capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories, parenterally, e.g. in the form of injection solutions or infusion solutions, or topically, e.g. in the form of ointments, creams or oils.

The production of the pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described compounds of formula (I) and their pharmaceutically acceptable, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible
5 solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for
10 example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers are, however, required in the case of soft gelatine capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injection solutions are, for example, water, alcohols, polyols, glycerol and
15 vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

20 Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

The dosage of the compounds of formula (I) can vary within wide limits
25 depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 mg to about 1000 mg, especially about 1 mg to about 100 mg, comes into consideration. Depending on the dosage it is convenient to administer the daily dosage in several dosage
30 units.

The pharmaceutical preparations conveniently contain about 0.1-500 mg, preferably 0.5-100 mg, of a compound of formula (I).

The following examples serve to illustrate the present invention in more detail. They are, however, not intended to limit its scope in any manner.

Examples

Abbreviations:

AcOEt = ethyl acetate, DMF = N,N-dimethylformamide, DMPU = 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone, MeOH = methanol, quant. = quantitative, RT = room
5 temperature.

Example 1a] (Z)-2-Hydroxy-4-oxo-4-(4-trifluoromethoxy-phenyl)-but-2-enoic acid ethyl ester

A solution of 1-(4-trifluoromethoxy-phenyl)-ethanone (5 g, 24 mmol) and diethyl
oxalate (3.25 ml, 24 mmol) in ethanol (5 ml) was added within 20 min to an ice cooled
10 solution of metallic sodium (552 mg, 24 mmol) in ethanol (15 ml) under an argon
atmosphere. The cooling bath was removed and the reaction stirred 30 min after
reaching ambient temperature. After standing 14 h, the precipitated yellow solid was
filtered. The solid was partitioned between 1 M HCl/ice water 1/1 and tert butyl methyl
ether. The aqueous layer was extracted two times with tert butyl methyl ether, the
15 combined extracts were washed with brine/ice water 1/1 and dried over sodium sulfate.
Evaporation of the solvent under reduced pressure gave 7.2 g (23.8 mmol, 99 %) of the
title compound as orange crystals.

MS: 305.0 (M+H)⁺.

b] 5-(4-Trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester

20 Hydrazine monohydrate (0.78 ml, 24 mmol) was added at ambient temperature to a
solution of (Z)-2-hydroxy-4-oxo-4-(4-trifluoromethoxy-phenyl)-but-2-enoic acid ethyl
ester (7.2 g, 24 mmol) in ethanol (37 ml) under an argon atmosphere. The suspension
was stirred for 4 h at reflux temperature, the solvent was removed under reduced
pressure and the residue partitioned between 1 M HCl/ice water and ethyl acetate. The
25 aqueous layer was extracted two times with ethyl acetate, the combined extracts were
washed with brine (3 times) and dried over sodium sulfate. Removal of the solvent under
reduced pressure left yellow crystals which were recrystallized from
dichloromethane/heptane to give 4.3 g (14.5 mmol, 61 %) of the title compound as
yellow crystals.

30 MS: 301.0 (M+H)⁺.

c) 2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester and 1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester

5-(4-Trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester (3 g, 10 mmol) was added to a solution of KOH (701 mg, 12 mmol) in absolute ethanol (91 ml). The solution was stirred at ambient temperature for 15 min. Methyl iodide (1.25 ml, 20 mmol) was added and the reaction solution was heated under reflux for 3 h. The solvent was removed under reduced pressure and the residue dissolved in brine/ice water 1/1 and ethyl acetate. The layers were separated and the aqueous layer was extracted two times with ethyl acetate. The combined organic layers were washed with brine and dried over sodium sulfate. The solvent was removed under reduced pressure and the residue purified by column chromatography (silica gel, heptane/AcOEt) to give 2.2 g (7 mmol, 70 %) 2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester as white crystals and 170 mg (0.54 mmol, 5 %) 1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester as yellow oil.

2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester: MS: 315.0 (M+H)⁺.

d) [2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol

A solution of 2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester (2.2 g, 7 mmol) in diethyl ether (68 ml) was added within 5 min to a suspension of lithium aluminium hydride (584 mg, 15 mmol) in diethyl ether (68 ml) under an argon atmosphere at ambient temperature. The mixture was heated to reflux for 12 h, cooled to 0 °C and treated cautiously with water (20 ml) and 10 % aqueous NaOH (10 ml). The reaction mixture was filtered over celite, ice water/tert butyl methyl ether 1/1 was added and the layers were separated. The aqueous layer was extracted one more time with tert butyl methyl ether, the combined organic layers were washed with ice water/brine 1/1 and dried over sodium sulfate. Removal of the solvent under reduced pressure gave 1.84 g (6.8 mmol, 97 %) of the title compound as white crystals.

MS: 273.1 (M+H)⁺.

e) 2-Methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]methoxy]-phenoxy}-propionic acid ethyl ester

To an ice cold solution of 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (50 mg, 210 μmol; PCT Int. Appl. (2002), WO 2002092590 A1), [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (57 mg, 210 μmol) and tributylphosphine (70 μl, 250 μmol) in tetrahydrofuran (5 ml) was added N,N,N',N'-

tetramethyl azodicarboxamide (43 mg, 250 μ mol). The cooling bath was removed and stirring continued for 14 h. The mixture was filtered over celite and the solvent removed under reduced pressure to give a yellow oil which was purified by column chromatography (silica gel, heptane/AcOEt) to obtain 77 mg (160 μ mol, 75 %) of the
5 title compound as colorless oil.

MS: 493.5 (M+H)⁺.

f] 2-Methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid

To a solution of 2-methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester (10 mg, 20 μ mol) in
10 THF/methanol 2/1 (1.5 ml) was added 1 N aqueous LiOH solution (120 μ l). The reaction mixture was stirred for 14 h at ambient temperature and concentrated under reduced pressure. The residue was dissolved in 1 N HCl/ice water 1/1 and ethyl acetate, the layers were separated and the aqueous layer was extracted with ethyl acetate. The combined
15 extracts were washed with ice water/brine 1/1, dried over sodium sulfate and the solvent was evaporated in vacuo to give the title compound (9 mg, 20 μ mol, 95 %) as off-white solid.

MS: 463.1 (M-H)⁻.

Example 2

20 a] 5-Chloromethyl-1-methyl-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole

To a solution of [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (1 g, 3.7 mmol; example 1 d) in chloroform (100 ml) was added thionyl chloride (0.53 ml, 7.3 mmol) at 0 °C under an argon atmosphere. The solution was stirred at ambient temperature for 4 h. Additional thionyl chloride (1.6 ml, 22 mmol) was added at 0 °C
25 and the solution was stirred for 12 h at ambient temperature. The mixture was poured onto ice water/aqueous NaHCO₃ 1/1, extracted two times with dichloromethane and the combined extracts were dried over sodium sulfate. Evaporation of the solvent under reduced pressure gave 2.15 g (quant.) of the title compound as colorless oil which was used in the next step without further purification.

30 MS: 291.0 (M+H)⁺.

b) [2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetonitrile

Tetrabutylammonium cyanide (1.27 g, 4.7 mmol) was added to a solution of 5-chloromethyl-1-methyl-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole (1.06 g, 3.7 mmol) in acetonitrile (24 ml). The solution was stirred at ambient temperature for 16 h,
5 saturated aqueous sodium bicarbonate solution/ice water 1/1 and ethyl acetate were added and the layers were separated. The aqueous layer was extracted with ethyl acetate, the combined organic layers were washed with ice water/brine 1/1, dried over sodium sulfate and the solvent was evaporated in vacuo to give a red oil which was purified by column chromatography (silica gel, n-heptane/AcOEt) to yield 627 mg (2.2 mmol, 61 %)
10 of the title compound as yellow crystals.

MS: 300.4 (M+NH₄)⁺.

c) [2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetic acid

A mixture of [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetonitrile (620 mg, 2.2 mmol), sodium hydroxide (882 mg, 22 mmol), water (9 ml) and ethanol
15 (9 ml) was stirred vigorously at 85 °C for 7 h. The reaction mixture was poured onto crushed ice and aqueous HCl and extracted three times with ethyl acetate. The combined extracts were washed with water and brine, and dried over anhydrous sodium sulfate. Evaporation of the solvent under reduced pressure gave 680 mg (2.26 mmol, quant.) of the title compound as off-white crystals.

20 MS: 301.0 (M+H)⁺.

d) 2-[2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol

A solution of [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetic acid (680 mg, 2.7 mmol) in tetrahydrofuran (9.5 ml) was treated at 0 °C with a 1 M solution of BH₃*THF in tetrahydrofuran (5.7 ml, 5.7 mmol). The cooling bath was removed and
25 the reaction mixture stirred at ambient temperature for 16 h. Careful quenching with MeOH and ice water, twofold extraction with AcOEt, washing with ice water/brine 1/1, drying over magnesium sulfate, and evaporation of the solvent left a crude product which was refluxed for 30 min in MeOH to liberate quantitatively the free alcohol. The solvent was evaporated in vacuo to yield a colorless oil which was purified by column
30 chromatography (silica gel, n-heptane/AcOEt) to give 346 mg (1.2 mmol, 53 %) of the title compound as colorless crystals.

MS: 287.0 (M+H)⁺.

e] 2-Methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590 A1) was reacted with 2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol in the presence of di-tert-butyl azodicarboxylate and triphenylphosphine to give 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester as colorless oil.

MS: 507.5 (M+H)⁺.

10 f] 2-Methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid

In analogy to the procedure described in example 1 f], 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester was treated with LiOH to obtain 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid as yellow crystals.

MS: 479.4 (M+H)⁺.

Example 3

a] 2-Methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 2-(3-hydroxy-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2001), WO 20010161120 A1) was reacted with 2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol (example 2 d]) in the presence of di-tert-butyl azodicarboxylate and triphenylphosphine to give 2-methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester as colorless oil.

MS: 493.5 (M+H)⁺.

b] 2-Methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid

30 In analogy to the procedure described in example 1 f], 2-methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl

ester was treated with LiOH to obtain 2-methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid as colorless foam.

MS: 465.3 (M+H)⁺.

5

Example 4

a) 3-{2-Methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 3-(4-hydroxy-2-methoxy-phenyl)-propionic acid ethyl ester (PCT Int. Appl. (2003), WO 2004000315 A1) was
10 reacted with [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (example 1 d]) in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 3-{2-methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid ethyl ester as colorless liquid.

MS: 479.4 (M+H)⁺.

15 b) 3-{2-Methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid

In analogy to the procedure described for example 1 f], 3-{2-methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid ethyl ester
was treated with LiOH to obtain 3-{2-methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid as colorless foam.
20

MS: 451.1 (M+H)⁺.

Example 5

a) 2-(4-Hydroxy-2,3-dimethyl-phenoxy)-2-methyl-propionic acid ethyl ester

A suspension of 8.0 g (57.9 mmol) 2,3-dimethylhydroquinone and 39.6 g (121.6 mmol)
25 cesium carbonate in 100 ml DMF was treated with 9.45 ml (63.7 mmol) ethyl 2-bromo-2-methylpropionate and stirred for 2 days at RT. The reaction was poured on a mixture of saturated NH₄Cl-solution and ice and extracted with AcOEt (3x). The organic phase was washed with aqueous 10 % NaCl, dried (Na₂SO₄) and evaporated. The crude product was purified by flash chromatography over silica gel with heptane/AcOEt 9:1, to
30 give 5.4 g of the title compound as dark brown oil.

MS: 252.1 (M⁺).

b] 2-{2,3-Dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2,3-dimethyl-phenoxy)-2-methyl-propionic acid ethyl ester was reacted with [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (example 1 d]) in the presence of
5 N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-{2,3-dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester as colorless liquid.

MS: 507.5 (M+H)⁺.

10 c] 2-{2,3-Dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid

In analogy to the procedure described for example 1 f], 2-{2,3-dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid ethyl ester was treated with LiOH to obtain 2-{2,3-dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid
15 as colorless foam.

MS: 479.4 (M+H)⁺.

Example 6

a] 2-(2,2,2-Trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester
20

Sodium hydride (55 % dispersion in mineral oil, 96 mg, 2 mmol) was added to an ice cooled solution of 5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester (500 mg, 2 mmol; example 1 b]) in DMF (30 ml) under an argon atmosphere. The solution was stirred for 10 min at 0 °C and for 40 min at ambient temperature.
25 Trifluoroethyltriflate (506 mg, 2 mmol) was added and the mixture was stirred for 3 h at ambient temperature. The solution was cooled to 0 °C, 1 N HCl/ice water 1/2 and dichloromethane were added. The layers were separated, the aqueous layer was extracted two times with dichloromethane, the combined extracts were washed with brine/ice water 1/1 and dried over sodium sulfate. Evaporation of the solvent under reduced
30 pressure gave a yellow oil which was purified by column chromatography (silica gel, n-heptane/AcOEt) to yield 553 mg (1.5 mmol, 87 %) of the title compound as colorless crystals.

MS: 382.1 (M)⁺.

b] [2-(2,2,2-Trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol

In analogy to the procedure described for example 1 d], 2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester was reduced with lithium aluminium hydride to give [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol as colorless crystals.

MS: 341.0 (M+H)⁺.

c] 2-Methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methylphenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590 A1) was reacted with [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester as yellow liquid.

MS: 561.3 (M+H)⁺.

d] 2-Methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid

In analogy to the procedure described for example 1 f], 2-methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester was treated with LiOH to obtain 2-methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid as yellow solid.

MS: 533.5 (M+H)⁺.

Example 7

a] [1-Methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-yl]-methanol

In analogy to the procedure described for example 1 d], 1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester (example 1 c]) was reduced with lithium aluminium hydride to give [1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-yl]-methanol as brown solid.

MS: 273.0 (M+H)⁺.

b] 2-Methyl-2-{2-methyl-4-[1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590
 5 A1) was reacted with [1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-yl]-methanol in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-methyl-2-{2-methyl-4-[1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester as colorless oil.

MS: 493.5 (M+H)⁺.

10 c] 2-Methyl-2-{2-methyl-4-[1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid

In analogy to the procedure described for example 1 f], 2-methyl-2-{2-methyl-4-[1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid ethyl ester was treated with LiOH to obtain 2-methyl-2-{2-methyl-4-[1-methyl-5-
 15 (4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid as yellow oil.

MS: 465.3 (M+H)⁺.

Example 8

a] Benzoic acid 2,5-dichloro-4-(1-ethoxycarbonyl-1-methyl-ethoxy)-phenyl ester

20 To a suspension of benzoic acid 2,5-dichloro-4-hydroxy-phenyl ester (1.4 g, 5 mmol; D. Koike, *Gunma Daigaku Kyoyobu Kiyo* 1968, 2, 13-28), cesium carbonate (2.6 g, 7.9 mmol) and a trace of potassium iodide in acetonitrile (80 ml) under an argon atmosphere was added bromo-acetic acid ethyl ester (1.1 ml, 7.4 mmol). The mixture was stirred for 14 h at ambient temperature, poured onto 1 N HCl/ice water 1/1 and
 25 extracted two times with ethyl acetate. The combined organic layers were washed with brine/water 1/1 and dried over sodium sulfate. The solvent was removed under reduced pressure and the residue purified by column chromatography (silica gel, heptane/AcOEt) to give 0.5 g (1.3 mmol, 25 %) of the title compound as colorless oil.

MS: 396.1 (M)⁺.

b) 2-(2,5-Dichloro-4-hydroxy-phenoxy)-2-methyl-propionic acid methyl ester

To an ice cold solution of benzoic acid 2,5-dichloro-4-(1-ethoxycarbonyl-1-methyl-ethoxy)-phenyl ester (500 mg, 1.3 mmol) in methanol (11.5 ml) was added a solution of sodium (145 mg, 6.3 mmol) in methanol (11.5 ml) within 5 min under an argon atmosphere. The solution was stirred for 4 h at ambient temperature, cooled to 0 °C and carefully neutralized with 1 N HCl. The solvent was removed under reduced pressure and the residue was dissolved in ethyl acetate and brine/ice water 1/1. The layers were separated and the aqueous layer was extracted two times with ethyl acetate. The combined organic layers were washed with brine/ice water 1/1 and dried over sodium sulfate. The solvent was removed under reduced pressure and the residue purified by column chromatography (silica gel, heptane/AcOEt) to give 219 mg (0.8 mmol, 62 %) of the title compound as colorless oil.

MS: 279.1 (M+H)⁺.

c) 2-{2,5-Dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid methyl ester

In analogy to the procedure described for example 1 e), 2-(2,5-dichloro-4-hydroxy-phenoxy)-2-methyl-propionic acid methyl ester was reacted with [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (example 1 d)) in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-{2,5-dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid methyl ester as colorless oil.

MS: 533.3 (M+H)⁺.

d) 2-{2,5-Dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid

In analogy to the procedure described for example 1 f), 2-{2,5-dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid methyl ester was treated with LiOH to obtain 2-{2,5-dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid as colorless solid.

MS: 517.1 (M-H)⁻.

Example 9

a] 2-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester and 1-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester

- 5 Chlorodifluoromethane (28.6 g, 331 mmol) was introduced to a suspension of 5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester (2 g, 7 mmol; example 1 b]) and anhydrous potassium carbonate (2.76 g, 20 mmol) in dry N,N-dimethylformamide (120 ml). The reaction mixture was stirred at 90 °C for 2 h. After cooling, the mixture was poured into ice water (400 ml) and extracted four times with
 10 dichloromethane. The combined extracts were washed two times with ice water/brine and dried over sodium sulfate. The solvent was removed under reduced pressure to give a yellow solid which was purified by column chromatography (silica gel, heptane/AcOEt) to give 281 mg (0.8 mmol, 12 %) 2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester as white solid and 1.29 g (3.7 mmol, 55 %) 1-
 15 difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester as white solid.

2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester: MS: 351.3 (M+H)⁺.

- 1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl
 20 ester: MS: 351.3 (M+H)⁺.

b] [1-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-yl]-methanol

- In analogy to the procedure described for example 1 d], 1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazole-3-carboxylic acid ethyl ester was reduced with lithium aluminium hydride to give [1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-
 25 1H-pyrazol-3-yl]-methanol as yellow oil.

c] 2-{4-[1-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid ethyl ester

- In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590
 30 A1) was reacted with [1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-yl]-methanol in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-{4-[1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-

pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid ethyl ester as colorless oil.

MS: 529.2 (M+H)⁺.

5 d] 2-{4-[1-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid

In analogy to the procedure described for example 1 f], 2-{4-[1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid ethyl ester was treated with LiOH to obtain 2-{4-[1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid as orange oil.

MS: 501.1 (M+H)⁺.

Example 10

a] 5-Iodomethyl-1-methyl-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole

15 A suspension of 5-chloromethyl-1-methyl-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole (3.2 g, 11 mmol; example 2 a]) and sodium iodide (8.25 g, 55 mmol) in acetone (56 ml) was heated under reflux conditions for 30 min. Tert butyl methyl ether was added, the solid was filtered off and the filtrate was brought to dryness under reduced pressure. The residue was dissolved in tert butyl methyl ether, washed with ice water/brine 1/1 and the aqueous layer was extracted two times with tert butyl methyl ether. The combined
20 extracts were washed with aqueous sodium thiosulfate solution and brine and dried over sodium sulfate. The solvent was removed under reduced pressure to give the title compound as yellow oil which was used in the next step without further purification.

b] 3-[2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propionic acid ethyl ester

25 A solution of lithium diisopropylamide (16.5 ml of a 2 M solution in tetrahydrofuran/heptane/ethylbenzol, 33 mmol) in tetrahydrofuran (25 ml) was cooled to -78 °C. Within 30 min a solution of ethyl acetate (3.77 ml, 38 mmol) in tetrahydrofuran (10 ml) was added. The solution was stirred for 45 min at -78 °C, DMPU (6.63 ml, 55 mmol) was added within 20 min and the mixture was kept for
30 additional 30 min at -78 °C. Within 20 min a solution of 5-iodomethyl-1-methyl-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole (4.2 g, 11 mmol) in tetrahydrofuran (25 ml) was added. The solution was stirred for 40 min at -78 °C, the cooling bath was removed and stirring was continued for 1 h. The reaction mixture was poored onto aqueous NH₄Cl

solution/ice water and extracted two times with ethyl acetate. The combined extracts were washed three times with ice water/brine and dried over sodium sulfate. The solvent was removed under reduced pressure to give an orange oil which was purified by column chromatography (silica gel, heptane/AcOEt) to give 1.5 g (4.4 mmol, 40 %) of the title
 5 compound as yellow oil.

MS: 343.1 (M+H)⁺.

c] 3-[2-Methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propan-1-ol

In analogy to the procedure described for example 1 d], 3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propionic acid ethyl ester was reduced with
 10 lithium aluminium hydride in diethyl ether to give 3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propan-1-ol as yellow oil.

MS: 300.2 (M)⁺.

d] 2-Methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid ethyl ester

15 In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590 A1) was reacted with 3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propan-1-ol in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid ethyl ester as colorless oil.
 20

MS: 521.5 (M+H)⁺.

e] 2-Methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid

In analogy to the procedure described for example 1 f], 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid ethyl ester was treated with LiOH to obtain 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid as colorless solid.
 25

MS: 493.5 (M+H)⁺.

Example 11

a] 2-(2,5-Dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid methyl ester

In analogy to the procedure described for example 1 e], 2-(2,5-dichloro-4-hydroxy-
5 phenoxy)-2-methyl-propionic acid methyl ester (example 8 b]) was reacted with 2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol in the presence of di-tert-butyl azodicarboxylate and triphenylphosphine to give 2-(2,5-dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid methyl ester as colorless oil.

10 MS: 547.3 (M+H)⁺.

b] 2-(2,5-Dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid

In analogy to the procedure described in example 1 f], 2-(2,5-dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-2-methyl-propionic
15 acid methyl ester was treated with LiOH to obtain 2-(2,5-dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid as colorless solid.

MS: 533.3 (M+H)⁺.

Example 12

20 a] 5-Chloromethyl-1-(2,2,2-trifluoro-ethyl)-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole

In analogy to the procedure described for example 2 a], [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (example 6 b]) was reacted with thionyl chloride in chloroform to yield 5-chloromethyl-1-(2,2,2-trifluoro-ethyl)-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole as colorless oil.
25

MS: 359.0 (M+H)⁺.

b] [2-(2,2,2-Trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetonitrile

In analogy to the procedure described for example 2 b], 5-chloromethyl-1-(2,2,2-trifluoro-ethyl)-3-(4-trifluoromethoxy-phenyl)-1H-pyrazole was reacted with
30

tetrabutylammonium cyanide in acetonitrile to give [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetonitrile as yellow oil.

MS: 350.3 (M+NH₄)⁺.

c] [2-(2,2,2-Trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetic acid

In analogy to the procedure described for example 2 c], [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetonitrile was treated with sodium hydroxide in water/ethanol 1/1 at 85 °C to give [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetic acid as brown crystals.

10 MS: 369.1 (M+H)⁺.

d] 2-[2-(2,2,2-Trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol

In analogy to the procedure described for example 2 d], [2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-acetic acid was reduced with a 1 M solution of BH₃*THF in tetrahydrofuran to give 2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol as colorless oil.

MS: 355.3 (M+H)⁺.

e] 2-Methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester

20 In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590 A1) was reacted with 2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethanol in the presence of di-tert-butyl azodicarboxylate and triphenylphosphine to give 2-methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid ethyl ester as colorless oil.

MS: 575.5 (M+NH₄)⁺.

f] 2-Methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid

30 In analogy to the procedure described in example 1 f], 2-methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-

phenoxy)-propionic acid ethyl ester was treated with LiOH to obtain 2-methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid as colorless oil.

MS: 547.3 (M+H)⁺.

5

Example 13

a) 1-(4-Hydroxy-2-methoxy-5-methyl-phenyl)-ethanone

Acetylchloride (1.16 ml, 16 mmol) was added within 5 min to a ice cooled suspension of AlCl₃ (2.4 g, 16.4 mmol) in 1,2-dichloroethane (5 ml) under an argon atmosphere. A solution of 5-methoxy-2-methyl-phenol (1.13 g, 8.2 mmol; PCT Int. Appl. (2003), WO
10 2003084916 A2) in 1,2-dichloroethane (2.4 ml) was added within 5 min. The mixture was naturally warmed to ambient temperature, poured after 4 h onto ice water and extracted twice with dichloromethane. The combined extracts were washed with ice water/0.5 M NaOH solution 1/1 and brine and dried over sodium sulfate. Removal of the solvent under reduced pressure gave a yellow oil which was dissolved in a mixture of 3.5
15 ml methanol, 7 ml THF and 7 ml 1 M LiOH solution. The solution was stirred for 30 min at ambient temperature and the solvent was partially removed under reduced pressure. Ice water/1 M HCl solution 1/1 was added and the solution was extracted twice with ethyl acetate. The combined extracts were washed with brine and dried over sodium sulfate. Evaporation of the solvent left a yellow solid which was crystallized from
20 dichloromethane/methanol/heptane to yield 441 mg (2.45 mmol, 30 %) of the title compound as colorless crystals.

MS: 179.4 (M-H)⁻.

b) 2-(4-Acetyl-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester

A suspension of 1-(4-hydroxy-2-methoxy-5-methyl-phenyl)-ethanone (416 mg, 2.3 mmol), 2-bromo-2-methyl-propionic acid ethyl ester (0.69 ml, 4.6 mmol), cesium carbonate (1.58 g, 4.9 mmol) and a trace of potassium iodide in acetonitrile (25 ml) was
25 heated under reflux conditions for 14 h. The mixture was poured onto 1 M HCl solution/ice water 1/1 and extracted two times with ethyl acetate. The combined extracts were washed with brine/ice water and dried over sodium sulfate. Removal of the solvent under reduced pressure gave a yellow oil which was purified by column chromatography
30 (silica gel, heptane/AcOEt) to give 450 mg (1.5 mmol, 66 %) of the title compound as colorless oil.

MS: 295.5 (M+H)⁺.

c] 2-(4-Acetoxy-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester

A solution of 2-(4-acetyl-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (531 mg, 1.8 mmol), 3-chloroperbenzoic acid (545 mg, 3.2 mmol) and 4-toluenesulfonic acid (34 mg, 0.2 mmol) in dichloromethane (24 ml) was heated under
5 reflux conditions for 72 h. The mixture was cooled to room temperature and washed two times with ice water/sodium iodide solution and two times with ice water/aqueous NaHSO₃ solution. The organic layer was dried over sodium sulfate, the solvent was removed under reduced pressure and the resulting brown solid was purified by column chromatography (silica gel, heptane/AcOEt) to give 235 mg (0.8 mmol, 42 %) of the title
10 compound as yellow oil.

MS: 311.3 (M+H)⁺.

d] 2-(4-Hydroxy-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid methyl ester

A freshly prepared solution of sodium (91 mg, 4 mmol) in methanol (5.4 ml) was added within 5 min to an ice cooled solution of 2-(4-acetoxy-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (235 mg, 0.8 mmol) in methanol (5.4 ml). The solution
15 was naturally warmed to ambient temperature and after 5 h the solvent was removed under reduced pressure. Ice water/1 M HCl 1/1 was added and the mixture was extracted two times with dichloromethane. The combined extracts were dried over sodium sulfate and the solvent was removed under reduced pressure to yield 153 mg (0.6 mmol, 76 %) of the title compound as brown oil which was used in the next step without further
20 purification.

MS: 254.2 (M)⁺.

e] 2-{5-Methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid methyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid methyl ester was reacted with [2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol (example 1 d]) in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-{5-methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid methyl ester as colorless oil.
30

MS: 509.5 (M+H)⁺.

e] 2-{5-Methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid

In analogy to the procedure described for example 1 f], 2-{5-methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid methyl ester was treated with LiOH to obtain 2-{5-methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid as colorless oil.

MS: 495.5 (M+H)⁺.

Example 14

10 a] 2-(5-Methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid methyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-5-methoxy-2-methyl-phenoxy)-2-methyl-propionic acid methyl ester (example 13 d]) was reacted with 3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propan-1-ol (example 10 c]) in the presence of N,N,N',N'-tetramethyl azodicarboxamide and tributylphosphine to give 2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid methyl ester as colorless oil.

MS: 537.2 (M+H)⁺.

20 b] 2-(5-Methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid

In analogy to the procedure described for example 1 f], 2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid methyl ester was treated with LiOH to obtain 2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid as colorless oil.

MS: 523.5 (M+H)⁺.

Example 15

a] [2-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol

30 In analogy to the procedure described for example 1 d], 2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazole-3-carboxylic acid ethyl ester (example 9 a]) was

reduced with lithium aluminium hydride to give [2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol as white solid.

MS: 309.4 (M+H)⁺.

5 b) 2-{4-[2-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid ethyl ester

In analogy to the procedure described for example 1 e], 2-(4-hydroxy-2-methyl-phenoxy)-2-methyl-propionic acid ethyl ester (PCT Int. Appl. (2002), WO 2002092590 A1) was reacted with [2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-methanol in the presence of N,N,N',N'-tetramethyl azodicarboxamide and
10 tributylphosphine to give 2-{4-[2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid ethyl ester as yellow oil.

MS: 529.3 (M+H)⁺.

15 c) 2-{4-[2-Difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid

In analogy to the procedure described for example 1 f], 2-{4-[2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid ethyl ester was treated with LiOH to obtain 2-{4-[2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-
20 propionic acid as colorless oil.

MS: 501.4 (M+H)⁺.

Example A

Film coated tablets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per tablet</u>	
Kernel:		
Compound of formula (I)	10.0 mg	200.0 mg
Microcrystalline cellulose	23.5 mg	43.5 mg
Lactose hydrous	60.0 mg	70.0 mg
Povidone K30	12.5 mg	15.0 mg
Sodium starch glycolate	12.5 mg	17.0 mg
Magnesium stearate	1.5 mg	4.5 mg
(Kernel Weight)	120.0 mg	350.0 mg
Film Coat:		
Hydroxypropyl methyl cellulose	3.5 mg	7.0 mg
Polyethylene glycol 6000	0.8 mg	1.6 mg
Talc	1.3 mg	2.6 mg
Iron oxyde (yellow)	0.8 mg	1.6 mg
Titan dioxide	0.8 mg	1.6 mg

- 5 The active ingredient is sieved and mixed with microcrystalline cellulose and the mixture is granulated with a solution of polyvinylpyrrolidone in water. The granulate is mixed with sodium starch glycolate and magnesium stearate and compressed to yield kernels of 120 or 350 mg respectively. The kernels are lacquered with an aqueous solution / suspension of the above mentioned film coat.

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula (I)	25.0 mg
Lactose	150.0 mg
Maize starch	20.0 mg
Talc	5.0 mg

- 5 The components are sieved and mixed and filled into capsules of size 2.

Example C

Injection solutions can have the following composition:

Compound of formula (I)	3.0 mg
Gelatine	150.0 mg
Phenol	4.7 mg
Sodium carbonate	to obtain a final pH of 7
Water for injection solutions	ad 1.0 ml

Example D

Soft gelatin capsules containing the following ingredients can be manufactured in a conventional manner:

Capsule contents

Compound of formula (I)	5.0 mg
Yellow wax	8.0 mg
Hydrogenated Soya bean oil	8.0 mg
Partially hydrogenated plant oils	34.0 mg
Soya bean oil	110.0 mg
Weight of capsule contents	165.0 mg

Gelatin capsule

Gelatin	75.0 mg
Glycerol 85 %	32.0 mg
Karion 83	8.0 mg (dry matter)
Titan dioxide	0.4 mg
Iron oxide yellow	1.1 mg

- 5 The active ingredient is dissolved in a warm melting of the other ingredients and the mixture is filled into soft gelatin capsules of appropriate size. The filled soft gelatin capsules are treated according to the usual procedures.

Example E


Sachets containing the following ingredients can be manufactured in a conventional manner:

Compound of formula (I)	50.0 mg
Lactose, fine powder	1015.0 mg
Microcrystalline cellulose (AVICEL PH 102)	1400.0 mg
Sodium carboxymethyl cellulose	14.0 mg
Polyvinylpyrrolidon K 30	10.0 mg
Magnesiumstearate	10.0 mg
Flavoring additives	1.0 mg

- 5 The active ingredient is mixed with lactose, microcrystalline cellulose and sodium carboxymethyl cellulose and granulated with a mixture of polyvinylpyrrolidon in water. The granulate is mixed with magnesiumstearate and the flavouring additives and filled into sachets.

Claims

1. A compound of the formula



wherein

X^1 is selected from the group consisting of O, S and CH_2 ;

R^1 is hydrogen or C_{1-7} -alkyl;

R^2 is hydrogen or C_{1-7} -alkyl,

or, if X^1 is CH_2 , R^2 is selected from the group consisting of hydrogen,


C_{1-7} -alkyl and C_{1-7} -alkoxy;

R^3 is hydrogen or C_{1-7} -alkyl;

R^4 and R^8 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

R^5 , R^6 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

and one of R^5 , R^6 and R^7 is



wherein

X^2 is S, O and NR^9 ,

R^9 is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, fluoro- C_{1-7} -alkyl, hydroxy- C_{2-7} -alkyl and C_{1-7} -alkoxy- C_{2-7} -alkyl;

R^{10} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl and fluoro- C_{1-7} -alkyl;

R^{11} is selected from the group consisting of hydrogen, C_{1-7} -alkyl and C_{1-7} -alkoxy- C_{1-7} -alkyl;

one of R^{12} or R^{13} is selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, C_{2-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl and fluoro- C_{1-7} -alkyl; and the other one is a lone pair;


R^{14} is hydrogen, C_{1-7} -alkyl, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl or fluoro- C_{1-7} -alkyl;

R^{15} is 4-trifluoromethoxyphenyl;

n is 1, 2 or 3; or

a pharmaceutically acceptable salt or ester thereof.

2. The compound of formula I according to claim 1 having the formula



wherein

X^1 , X^2 , R^1 to R^4 , R^8 , R^{10} to R^{15} and n are as defined in claim 1;


R^5 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano;

or a pharmaceutically acceptable salt or ester thereof.

3. The compound of formula I-A according to claim 2, wherein at least one of R^4 , R^5 , R^7 and R^8 is C_{1-7} -alkyl or C_{1-7} -alkoxy.

4. The compound of formula I-A according to claim 3, wherein R^4 is C_{1-7} -alkyl or C_{1-7} -alkoxy.

5. The compound of formula I according to claim 1 having the formula




wherein

X^1 , X^2 , R^1 to R^4 , R^8 , R^{10} to R^{15} and n are as defined in claim 1;

R^5 and R^6 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano; or a pharmaceutically acceptable salt or ester thereof.

6. The compound of formula I-B according to claim 5, wherein at least one of R^4 , R^5 , R^6 and R^8 is C_{1-7} -alkyl or C_{1-7} -alkoxy.

7. The compound of formula I according to claim 1 having the formula



wherein

X^1 , X^2 , R^1 to R^4 , R^8 , R^{10} to R^{15} and n are as defined in claim 1;

R^6 and R^7 independently from each other are selected from the group consisting of hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl and cyano; or a pharmaceutically acceptable salt or ester thereof.

8. The compound of formula I-C according to claim 6, wherein at least one of R^4 , R^6 , R^7 and R^8 is C_{1-7} -alkyl or C_{1-7} -alkoxy.

9. The compound of formula I according to any one of claims 1 to 8, wherein R^1 is hydrogen.

10. The compound of formula I according to any one of claims 1 to 9, wherein X^1 is O.

11. The compound of formula I according to claim 10, wherein R^2 and R^3 are C_{1-7} -alkyl.

12. The compound of formula I according to any one of claims 1 to 9, wherein X^1 is CH_2 .


13. The compound of formula I according to any one of claims 1 to 12, wherein X^2 is O.

14. The compound of formula I according to any one of claims 1 to 13, wherein n is 1 or 2.

15. The compound of formula I according to any one of claims 1 to 14, wherein n is 2.

16. The compound of formula I according to any one of claims 1 to 13, wherein n is 3.

17. The compound of formula I according to any one of claims 1 to 14, wherein one of R^5 , R^6 and R^7 is



and R^{10} to R^{12} , R^{14} , R^{15} and n are as defined in claim 1.

18. The compound of formula I according to claim 15, wherein R^{12} is C_{1-7} -alkyl or fluoro- C_{1-7} -alkyl.

19. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-{2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid or a pharmaceutically acceptable salt or ester thereof.

20. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-(2-methyl-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid or a pharmaceutically acceptable salt or ester thereof.

21. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-(3-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid or a pharmaceutically acceptable salt or ester thereof.

22. The compound of formula I according to claim 1, wherein the compound is 3-{2-methoxy-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenyl}-propionic acid or a pharmaceutically acceptable salt or ester thereof.

23. The compound of formula I according to claim 1, wherein the compound is 2-{2,3-dimethyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.

24. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-{2-methyl-4-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid or a pharmaceutically acceptable salt or ester thereof.

25. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-{2-methyl-4-[1-methyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-phenoxy}-propionic acid or a pharmaceutically acceptable salt or ester thereof.

26. The compound of formula I according to claim 1, wherein the compound is 2-{2,5-dichloro-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.

27. The compound of formula I according to claim 1, wherein the compound is 2-{4-[1-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-1H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.

28. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-(2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-propionic acid or a pharmaceutically acceptable salt or ester thereof.

29. The compound of formula I according to claim 1, wherein the compound is 2-(2,5-dichloro-4-{2-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.

30. The compound of formula I according to claim 1, wherein the compound is 2-methyl-2-(2-methyl-4-{2-[2-(2,2,2-trifluoro-ethyl)-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-ethoxy}-phenoxy)-propionic acid or a pharmaceutically acceptable salt or ester thereof.


31. The compound of formula I according to claim 1, wherein the compound is 2-{5-methoxy-2-methyl-4-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-phenoxy}-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.

32. The compound of formula I according to claim 1, wherein the compound is 2-(5-methoxy-2-methyl-4-{3-[2-methyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-yl]-propoxy}-phenoxy)-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.


33. The compound of formula I according to claim 1, wherein the compound is 2-{4-[2-difluoromethyl-5-(4-trifluoromethoxy-phenyl)-2H-pyrazol-3-ylmethoxy]-2-methyl-phenoxy}-2-methyl-propionic acid or a pharmaceutically acceptable salt or ester thereof.

34. A process for the manufacture of the compound defined in any one of claims 1 to 33, which process comprises


reacting a compound of formula



wherein R^1 is C_{1-7} -alkyl, R^2 , R^3 , R^4 and R^8 are as defined as in claim 1 and R^5 , R^6 and R^7 are selected from hydrogen, C_{1-7} -alkyl, C_{1-7} -alkoxy, C_{3-7} -cycloalkyl, halogen, C_{1-7} -alkoxy- C_{1-7} -alkyl, C_{2-7} -alkenyl, C_{2-7} -alkynyl, fluoro- C_{1-7} -alkyl, fluoro- C_{1-7} -alkoxy, cyano- C_{1-7} -alkyl, and cyano with the proviso that one of R^5 , R^6 or R^7 is -OH, -SH or -NHR⁹, wherein R^9 is as defined in claim 1, with a compound of formula



wherein R¹⁰ to R¹⁵ and n are as defined in claim 1 and R¹⁶ is -OH, -Cl, -Br, -I or another leaving group, to obtain a compound of formula



wherein R¹ is C₁₋₇-alkyl and X¹, R² to R⁸ are as defined in claim 1, and optionally hydrolysing the ester group to obtain a compound of formula I, wherein R¹ is hydrogen.

35. The compound according to any one of claims 1 to 33 when manufactured by a process according to claim 34.

36. A pharmaceutical composition comprising the compound defined in any one of claims 1 to 33 and a pharmaceutically acceptable carrier and/or adjuvant.

37. The pharmaceutical composition according to claim 36 for the treatment and/or prevention of diabetes, non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, atherosclerotic diseases, metabolic syndrome, syndrome X, obesity, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases, or a proliferative disease.

38. The pharmaceutical composition according to claim 37, for the treatment and/or prevention of low HDL cholesterol levels, high LDL cholesterol levels, high triglyceride levels, metabolic syndrome or syndrome X.

39. The compound according to any one of claims 1 to 33 for use as a therapeutically active substance for the treatment and/or prevention of diabetes, non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, atherosclerotic diseases, metabolic syndrome, syndrome X,

obesity, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases, or a proliferative disease.


40. The compound according to any one of claims 1 to 33 for use as a therapeutically active substance for treatment and/or prevention of low HDL cholesterol levels, high LDL cholesterol levels, high triglyceride levels, metabolic syndrome or syndrome X.

41. A use of the compound defined in any one of claims 1 to 33 for the preparation of a medicament for the treatment and/or prevention of diabetes, non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, atherosclerotic diseases, metabolic syndrome, syndrome X, obesity, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases, or a proliferative disease.


42. A use of the compound defined in any one of claims 1 to 33 for the preparation of a medicament for the treatment and/or prevention of low HDL cholesterol levels, high LDL cholesterol levels, high triglyceride levels, metabolic syndrome or syndrome X.

43. A use of the compound defined in any one of claims 1 to 33 for the treatment and/or prevention of diabetes, non-insulin dependent diabetes mellitus, increased lipid and cholesterol levels, atherosclerotic diseases, metabolic syndrome, syndrome X, obesity, elevated blood pressure, endothelial dysfunction, procoagulant state, dyslipidemia, polycystic ovary syndrome, inflammatory diseases, or a proliferative disease.

44. A use of the compound defined in any one of claims 1 to 33 for the treatment and/or prevention of low HDL cholesterol levels, high LDL cholesterol levels, high triglyceride levels, metabolic syndrome or syndrome X.



I



II