



(51) International Patent Classification:

A61K 38/05 (2006.01) A61P 35/02 (2006.01)
A61P 19/00 (2006.01)

(21) International Application Number:

PCT/GB2009/000687

(22) International Filing Date:

13 March 2009 (13.03.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0805035.3 18 March 2008 (18.03.2008) GB
0805036.1 18 March 2008 (18.03.2008) GB

(71) Applicant (for all designated States except US): **BTG INTERNATIONAL LIMITED** [GB/GB]; 5 Fleet Place, London EC4M 7RD (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JANSEN, Gerrit** [NL/—]; Dept. of Rheumatology, VU University Medical Center, De Boelelaan 1117, NL-1081 HV Amsterdam (NL). **JACKMAN, Ann** [GB/GB]; CRC Centre for Cancer Therapeutics, The Institute of Cancer Research, 15 Cotswold Road, SUTTON SM2 5NG (GB).

(74) Agent: **ENGLAND, C.D.**; BTG INTERNATIONAL LIMITED, 5 Fleet Place, London EC4M 7RD (GB).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: CYCLOPENTA[G]QUINAZOLINE DERIVATIVES FOR THE TREATMENT OF RHEUMATOID ARTHRITIS OR ACUTE MYELOID LEUKAEMIA



(57) Abstract: A cyclopenta[g]quinazoline derivative, containing an L-Glu-γ-D-Glu dipeptide group, of formula (I): wherein: R¹ is amino, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ fluoroalkyl or methoxy-C₁₋₄-alkyl; R² is hydrogen, C₁₋₄ alkyl, C₃₋₄ alkenyl, C₃₋₄ alkynyl, C₂₋₄ hydroxyalkyl C₂₋₄ halogenoalkyl or C₁₋₄ cyanoalkyl; and Ar is phenylene, thiophenediyl, thiazolediyl, pyridinediyl or pyrimidinediyl which may optionally bear one or two substituents selected from halogeno, hydroxy, amino, nitro, cyano, trifluoromethyl, C₁₋₄ alkyl and C₁₋₄ alkoxy. the compound (I) optionally being in the form of a pharmaceutically acceptable salt or ester; may be used for the treatment of rheumatoid arthritis or acute myeloid leukaemia.


CYCLOPENTA[G]QUINAZOLINE DERIVATIVES FOR THE TREATMENT OF RHEUMATOID ARTHRITIS OR ACUTE MYELOID LEUKAEMIA

BACKGROUND OF THE INVENTION

This invention relates to the use of cyclopenta[g]quinazoline derivatives. More particularly it relates to cyclopenta[g]quinazoline derivatives for the treatment of rheumatoid arthritis (RA) and acute myeloid leukaemia (AML).

Cyclopenta[g]quinazoline derivatives showing a good level of activity both as regards their ability to inhibit thymidylate synthase (TS) and also as regards their anticancer activity against various cell lines have been developed.

WO 94/11354 A1 (British Technology Group Limited) discloses tricyclic compounds of formula:




wherein R^1 is hydrogen, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl or C_{1-4} fluoroalkyl;

R^2 is hydrogen, C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, C_{2-4} hydroxyalkyl, C_{2-4} halogenoalkyl or C_{1-4} cyanoalkyl;

Ar is phenylene, thiophenediyl, thiazolediyl, pyridinediyl or pyrimidinediyl which may optionally bear one or two substituents selected from halogeno, hydroxy, amino, nitro, cyano, trifluoromethyl, C_{1-4} alkyl and C_{1-4} alkoxy; and


R^3 is a group of one of the following formulae:



or R^3 is a *N*-linked naturally-occurring amino acid selected from the group consisting of L-alanine, L-leucine, L-isoleucine, L-valine and L-phenylalanine. Among the compounds disclosed is the L-Glu- γ -D-Glu compound CB300638, also mentioned in *Clinical Cancer Research*, **5**, November 1999 (Supplement) at #566 (Theti *et al.*) and *Proceedings of the American Association for Cancer Research*, **41**, March 2000 at #33 (Jackman *et al.*), as well as in *J. Med. Chem.*, 2000, **43**, 1910–1926, where it is

disclosed on page 1923 as compound **7b**.

WO 95/30673 A1 (British Technology Group Limited) discloses cyclopenta-[g]quinazolines of formula:




5

wherein R^1 is hydrogen, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl or C_{1-4} fluoroalkyl;

R^2 is hydrogen, C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, C_{2-4} hydroxyalkyl, C_{2-4} halogenoalkyl or C_{1-4} cyanoalkyl;

10 Ar^1 is phenylene, thiophenediyl, thiazolediyl, pyridinediyl or pyrimidinediyl which may optionally bear one or two substituents selected from halogeno, hydroxy, amino, nitro, cyano, trifluoromethyl, C_{1-4} alkyl and C_{1-4} alkoxy; and

R^3 is a group of one of the following formulae:



The α -isoform of the folate receptor (α -FR; membrane-associated folate-binding protein) is a glycosylphosphatidylinositol anchored cell membrane protein that has very high affinity for folic acid and the more biologically relevant reduced-
20 -folates ($K_d \sim 0.1$ nM). The mechanism of folate internalization is receptor-mediated endocytosis. The α -FR is overexpressed in many carcinomas, particularly those of ovarian origin where it is overexpressed highly and homogeneously in 90% of cases; see *Cancer Res.* **51**, 5329–5338, 1991 (Campbell *et al.*, 1991). Furthermore, high α -FR expression has been linked to aggressive, platinum resistant disease and poor
25 prognosis—see *Int. J. Cancer* **74**, 193–198, 1997 and *Int. J. Cancer* **79**, 121–126, 1998 (both Toffoli *et al.*). The β -isoform is widely expressed in tumours of epithelial and non-epithelial origin with expression levels being generally low / moderate and high, respectively, reviewed in *Critical Rev. Therap. in Drug Carrier Systems* **15**, 587–627, 1998 (Reddy and Low).

30 Folate receptors (α and β) are expressed in some adult normal tissues (low to

moderate expression). Certain compounds within the general class of cyclopenta-
[g]quinazolines have been reported to have a high level of selectivity for α -folate
receptor expressing human tumour cell lines versus the affinity for the RFC (reduced-
folate carrier). Such compounds are disclosed in WO 03/020300 A1, WO 03/020706
5 A1 and WO 03/020748 A1 (BTG International Limited). Among the compounds
disclosed is the L-Glu- γ -D-Glu compound CB300945, also mentioned in *Tetrahedron*,
63 (7), 12 February 2007, 1537–1543 (Bavetsias *et al.*) and *Cancer Research* 65, 15
December 2005, 11721–11728 (Gibbs *et al.*).

FR- β is normally found in placenta tissues and in hematopoietic cells, where it
10 is expressed in the myelomonocytic lineage and is particularly elevated during
neutrophil maturation or during monocyte or macrophage activation. However, the
FR- β expressed on normal hematopoietic cells, unlike that on activated macrophages
for example, is nonfunctional in that it cannot bind and internalize folate. FR- β is
expressed on malignant cells from patients with chronic myelogenous leukaemia
15 (CML), and on malignant cells from approximately 70% of patients with AML.


WO 03/072091-A1 (The Ohio State University Research Foundation) uses the
discovery that expression of FR- β is increased in malignant cells from myeloid leu-
kaemia patients by FR- β inducers. The FR- β expressed in myeloid leukaemia cells,
preferably AML cells, is functional in that it binds and internalizes folate, unlike the
20 FR- β expressed in the majority of normal hematopoietic cells which is non-
functional. Such functional FR- β is a target for folate-conjugated therapeutics.

Jansen, in an abstract from the 13th International Symposium on Chemistry &
Biology of Pteridines & Folates entitled “Antifolates in chronic inflammatory
diseases / rheumatoid arthritis: what can we learn from cancer and vice versa,” *Pteri-*
25 *dines*, 16: 46, 2005, indicates that methotrexate (MTX) is the anchor drug in
treatment regimens for patients with rheumatoid arthritis. He furthermore speculates
that cyclopenta[g]quinazoline-based TS inhibitors exhibited binding affinities close to
folic acid and could thus be interesting FR-targeted drugs in the treatment of cancer
as well as inflammatory diseases.

SUMMARY OF THE INVENTION

We have now discovered that certain compounds within the general class of
cyclopenta[g]quinazolines have an unexpectedly high level of selectivity for β -folate
receptor expressing cell lines. Such a compound has two structural features, that is to

say a three-ring structure and modification to the glutamate side chain when compared with folic acid. Accordingly the present invention comprises a cyclopenta[g]-quinazoline derivative, containing an L-Glu-γ-D-Glu dipeptide group, of formula (I):



wherein:

R¹ is amino, C₁-₄ alkyl, C₁-₄ hydroxyalkyl, C₁-₄ fluoroalkyl or methoxy-C₁-₄-alkyl;

R² is C₁-₄ alkyl, C₃-₄ alkenyl, C₃-₄ alkynyl, C₂-₄ hydroxyalkyl C₂-₄ halogeno-alkyl or C₁-₄ cyanoalkyl; and

Ar is phenylene, thiophenediyl, thiazolediyl, pyridinediyl or pyrimidinediyl which may optionally bear one or two substituents selected from halogeno, hydroxy, amino, nitro, cyano, trifluoromethyl, C₁-₄ alkyl and C₁-₄ alkoxy.


the compound (I) optionally being in the form of a pharmaceutically acceptable salt or ester;

for the treatment of rheumatoid arthritis or acute myeloid leukaemia.

The compounds of the invention display one or more of the following advantages:

1. high selectivity for cells over-expressing the β-FR, when grown in physiological concentrations of folate and possessing normal expression of the RFC;
2. a potent TS inhibition, a low affinity for the RFC and a moderate to high affinity for the β-FR;
3. TS-specific activity and are resistant to *in vivo* hydrolases; and
4. selective activity in primary Chinese Hamster Ovarian cell line screen with moderate β-FR expression.

In this specification the terms alkyl, alkenyl and alkynyl include both straight and branched chain groups but references to individual alkyl groups, such as propyl, are specific for the straight chain group only. An analogous convention applies to other generic terms. Moreover, the numbering system used for the cyclopenta[g]-quinazoline nucleus is the conventional one as shown below:



Amino-acid residues are designated herein in the standard manner (*Pure and Applied Chemistry*, 1974, **40**, 317 and *European Journal of Biochemistry*, 1984, **138**, 9). Thus, for example, γ -glutamyl denotes the radical $\text{H}_2\text{NCH}(\text{CO}_2\text{H})\text{CH}_2\text{CH}_2\text{CO}-$ or
 5 $-\text{NHCH}(\text{CO}_2\text{H})\text{CH}_2\text{CH}_2\text{CO}-$ according to the context, the carbon atoms in these radicals being numbered from the carbon atom of the α -carboxyl group as position 1.

It will be observed that a cyclopenta[g]quinazoline of the invention contains at least three asymmetric carbon atoms [present at the point of attachment of the group $-\text{N}(\text{R}^2)-$ to the tricyclic ring system and at the α -carbon atoms of the group L-Glu- γ -
 10 D-Glu] and can therefore exist in racemic and optically active forms. It is to be understood that this invention encompasses both racemic and optically active forms, it being a matter of common general knowledge how such optically active forms may be obtained by stereospecific synthesis or by separation of a mixture of isomeric compounds. It will be appreciated that one isomer may be of more interest than
 15 another owing to the nature of the activity which it exhibits or owing to superior physical properties, for example aqueous solubility.

It is also to be understood that a cyclopenta[g]quinazoline of the formula (I) may exhibit the phenomenon of tautomerism and that the formulae shown in this specification represent only one of the possible tautomeric forms.

20 It is also to be understood that certain cyclopenta[g]quinazolines of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms.

A suitable value for R^1 or R^2 when it is C_{1-4} alkyl, or for a C_{1-4} alkyl substituent which may be present on Ar, is, for example, methyl, ethyl, propyl or
 25 isopropyl.

A suitable value for a C_{1-4} alkoxy substituent which may be present on Ar is, for example, methoxy, ethoxy, propoxy, isopropoxy or butoxy.

A suitable value for a halogeno substituent which may be present on Ar is, for example, fluoro, chloro or bromo.

30 A suitable value for R^2 when it is C_{3-4} alkenyl is, for example, prop-2-enyl, but-2-enyl, but-3-enyl or 2-methylprop-2-enyl; and when it is C_{3-4} alkynyl is, for

example, prop-2-ynyl or but-3-ynyl.

A suitable value for R^2 when it is C_{2-4} hydroxyalkyl is, for example, 2-hydroxyethyl or 3-hydroxypropyl; when it is C_{2-4} halogenoalkyl is, for example, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 3-fluoropropyl, 3-chloropropyl or
5 3-bromopropyl; and when it is C_{1-4} cyanoalkyl is, for example, cyanomethyl, 2-cyanoethyl or 3-cyanopropyl.

A suitable value for Ar when it is phenylene is, for example, 1,3- or 1,4-phenylene, especially 1,4-phenylene.

A suitable value for Ar when it is thiophenediyl is, for example, thio-
10 phene-2,4-diyl or thiophene-2,5-diyl; when it is thiazolediyl is, for example thiazole-2,4-diyl or thiazole-2,5-diyl; when it is pyridinediyl is, for example, pyridine-2,4-diyl, pyridine-2,5-diyl, pyridine-2,6-diyl or pyridine-3,5-diyl; and when it is pyrimidinediyl is, for example, pyrimidine-2,4-diyl, pyrimidine-2,5-diyl or pyrimidine-4,6-diyl.

15 As indicated, Ar may carry one or two substituents. A preferred level of substitution in Ar, where substitution is present, is either two substituents or especially one substituent; and the one or two substituents may conveniently be at positions adjacent to the atom bonded to the group -CO-L-Glu- γ -D-Glu, halogeno substituents such as fluoro being preferred.

20 A suitable pharmaceutically acceptable salt form of a cyclopenta[g]-quinazoline of the invention is, for example, an acid addition salt with an inorganic or organic acid, for example hydrochloric, hydrobromic, trifluoroacetic or maleic acid; or an alkali metal, for example sodium, an alkaline earth metal, for example calcium, or ammonium, for example tetra(2-hydroxyethyl)ammonium, salt.

25 A suitable pharmaceutically acceptable ester form of a cyclopenta[g]quinazoline of the invention is, for example, an ester with an aliphatic alcohol of up to 6 carbon atoms, for example a methyl, ethyl or *tert*-butyl ester.

The compound contains three carboxyl groups. A salt or ester may be mono-acid-di-salt or -ester, di-acid-mono-salt or -ester or even tri-salt or -ester.

30 Preferably R^1 is C_{1-4} alkyl or C_{1-4} hydroxyalkyl. More preferably R^1 is methyl or, especially, hydroxymethyl.

Preferably R^2 is methyl, ethyl, propyl, prop-2-enyl, prop-2-ynyl, 2-hydroxyethyl, 2-fluoroethyl, 2-bromoethyl or 2-cyanoethyl. More preferably R^2 is methyl or, especially, prop-2-ynyl.

Preferably Ar is 1,4-phenylene which may optionally bear one or two substituents selected from the group consisting of chloro and especially fluoro, thiophene-2,5-diyl, thiazole-2,5-diyl or pyridine-2,5-diyl. More preferably Ar is 1,4-phenylene or 1,4-phenylene having a 2-fluoro substituent as in 2,6-difluoro-1,4-phenylene or especially 2-fluoro-1,4-phenylene or is pyridine 2,5-diyl. Most preferably Ar is 1,4-phenylene or 2-fluoro-1,4-phenylene.

A preferred cyclopenta[g]quinazoline of the invention has the formula (I) wherein R¹ is C₁₋₄ alkyl or C₁₋₄ hydroxyalkyl, especially hydroxymethyl;

R² is methyl, ethyl, propyl, prop-2-enyl, prop-2-ynyl, 2-hydroxyethyl, 2-fluoroethyl, 2-bromoethyl or 2-cyanoethyl; and

Ar is 1,4-phenylene which may optionally bear one or two substituents selected from the group consisting of chloro and especially fluoro, thiophene-2,5-diyl, thiazole-2,5-diyl or pyridine-2,5-diyl.

A further preferred cyclopenta[g]quinazoline of the invention has the formula (I) wherein R¹ is methyl or hydroxymethyl;

R² is methyl or prop-2-ynyl; and

Ar is 1,4-phenylene or 1,4-phenylene having a 2-fluoro substituent as in 2,6-difluoro-1,4-phenylene or especially 2-fluoro-1,4-phenylene or is pyridine 2,5-diyl.

An especially preferred cyclopenta[g]quinazoline of the invention has the formula (I) wherein R¹ is methyl or hydroxymethyl;

R² is methyl or preferably prop-2-ynyl; and

Ar is 1,4-phenylene or 2-fluoro-1,4-phenylene.

Specific particularly preferred cyclopenta[g]quinazolines of the invention are:

N-{*N*-{4-[*N*-(2-methyl-4-oxo-3,4,7,8-tetrahydro-6*H*-cyclopenta[g]quinazolin-6-yl)-*N*-(prop-2-ynyl)amino]benzoyl}-L-γ-glutamyl}-D-glutamic acid; or

N-{*N*-{4-[*N*-(2-hydroxymethyl-4-oxo-3,4,7,8-tetrahydro-6*H*-cyclopenta[g]quinazolin-6-yl)-*N*-(prop-2-ynyl)amino]benzoyl}-L-γ-glutamyl}-D-glutamic acid; or a pharmaceutically acceptable salt or ester thereof;

as well as the 6*S* isomers of these two compounds.

Although the compounds of the present invention can exist as a mixture of stereoisomers it is preferred that they are resolved into one optically active isomeric form. Such a requirement complicates the synthesis of the compounds and it is preferred therefore that they contain as few asymmetric carbon atoms as possible con-

sistent with achieving the desired activity.

As indicated previously, however, the cyclopenta[g]quinazolines of the present invention contain at least three asymmetric carbon atoms. Of these, that at the 6 position of the ring system preferably has the 6*S* orientation rather than the 6*R* orientation.. The preferred compounds (I) described hereinbefore thus preferably have
5 such a configuration at this asymmetric carbon atoms or less preferably are a mixture in which one or both of these asymmetric carbon atoms is unresolved.

A cyclopenta[g]quinazoline of the invention may be prepared by any process known to be applicable to the preparation of chemically-related compounds.

10 A cyclopenta[g]quinazoline of the present invention may itself be active or may be a prodrug converted *in vivo* to an active compound. A cyclopenta[g]quinazoline of the invention may be administered to a warm-blooded animal, including a human, in the form of a pharmaceutical composition which comprises the cyclopenta-
[g]quinazoline in association with a pharmaceutically acceptable diluent or carrier.

15 The composition may be in a form suitable for oral use, for example a tablet, capsule, aqueous or oily solution, suspension or emulsion; for topical use, for example a cream, ointment, gel or aqueous or oily solution or suspension; for nasal use, for example a snuff, nasal spray or nasal drops; for vaginal or rectal use, for example a suppository; for administration by inhalation, for example as a finely divided powder
20 such as a dry powder, a microcrystalline form or a liquid aerosol; for sub-lingual or buccal use, for example a tablet or capsule; or for parenteral use (including intravenous, subcutaneous, intramuscular, intravascular or infusion use), for example a sterile aqueous or oily solution, emulsion or suspension. In general the above compositions may be prepared in a conventional manner using conventional excipients.

25 The cyclopenta[g]quinazoline will normally be administered to a warm-blooded animal at a dose within a range of 50–25000, particularly 50–5000, mg per square metre body area of the animal, i.e. approximately 1500, particularly 1–100, mg/kg. Where desired, however, dosages outside this range may be employed and, in particular, where the preferred mode of administration involving subcutaneous
30 infusion is used then the dose range may be increased to 1–1000 mg/kg. Preferably a daily dose in the range 10–250 mg/kg is employed, particularly 30–150 mg/kg. However, the daily dose will necessarily be varied depending upon the host treated, the particular route of administration and the severity of the illness being treated. Accordingly, the optimum dosage may be determined by the practitioner who is

treating any particular patient.

Accordingly the present invention also includes a method for treating rheumatoid arthritis or acute myeloid leukaemia in a patient in need of such treatment which comprises administering to said patient an effective amount of a cyclopenta[g]quinazoline derivative as defined hereinbefore.

The compound will normally be administered at a dose within the range 5–25000, particularly 5–500, mg per square metre body area of the animal, i.e. approximately 0.1–500, particularly 0.1–10, mg/kg. Where desired, however, dosages outside this range may be employed. Topical administration of a cyclopenta[g]quinazoline of the invention may be used. Thus, for example, for topical administration a daily dose in the range, for example, of 0.1 to 10 mg/kg may be used.

Compositions containing the quinazolines may be formulated in unit dosage form, i.e. in the form of discrete portions each comprising a unit dose, or a multiple or sub-multiple of a unit dose, for example as a tablet or capsule. Such a unit dosage form may, for example, contain an amount of the cyclopenta[g]quinazoline in the range of 1–250 or 1–500 mg.

The invention is illustrated by the following Examples.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Immunohistochemistry of RA-synovial tissue: Light microscopy of (A) rabbit isotype staining, (B) FR- β staining and (C) macrophage staining (3A5). Immunofluorescence (double) staining of (D) FR- β , (E) CD-68 (macrophages) and (F) merge of D and E. Of note: nuclei are stained blue (Dapi-staining).

Figure 2: (A) Correlation between 3A5 (macrophages) and FR- β positive cells/mm² in synovial tissue of RA patients (n = 15) before MTX treatment. Median positive cell counts in the synovial sublining layer/mm² were 126 (range: 9–630) for FR- β and 219 (range: 11–622) for 3A5. Linear regression: R = 0.64; p = 0.04. (B) Correlation between DAS28 improvement (Δ DAS28) and FR- β expression on macrophages (positive cells/mm²) after 4 months treatment with MTX (R = 0.31; p = 0.11).

Figure 3: FR- β mRNA expression in synovial tissue and *ex vivo* cultured RA Peripheral Blood Lymphocytes (PBLs). Median FR- β mRNA expression was determined in peripheral blood lymphocytes (PBLs; n = 9), monocytes (n = 9), *ex vivo* cultured macrophages (n = 25), *ex vivo* activated T-cells (n = 22) and synovial tissue of RA patients (n = 7). FR- β expression in these samples is shown as a percentage of

the expression in CHO- β cells (Chinese Hamster Ovarian cells transfected with FR- β ; expression in these cells was set to 1.00).

Figure 4: Relative binding affinities of FR- α (white bars) and FR- β (grey bars) for novel generation folate antagonists. FR- α and FR- β binding affinities of the drugs were determined by [3 H]-folic acid displacement of FR α expressing KB-cells and FR β transfected CHO- β cells as described in the Materials and Methods section. Binding affinities are presented as percentage relative of folic acid binding affinity. Results are the mean $\pm$ SD of 3–5 separate experiments (SD < 30%). For comparison, affinities of RFC for folate antagonists are presented as: ++++ (high affinity), +++ (moderate affinity), ++ (low affinity), + (poor affinity), +/- (very poor affinity) based on previously published data.

Figure 5: Growth inhibition of CHO/WT and CHO/FR- β cells by BGC 945 with or without 1 μ M folic acid (A), CB300635 (B) and AG2034 (C/D) with or without supplementation of folic acid (1 μ M) or leucovorin (20 nM). Results are the mean of 3 experiments (SD < 20%).

DETAILED DESCRIPTION

Example 1: Exploitation of folate receptor- β as a potential delivery route for novel generation folate antagonists to activated macrophages in synovial tissue of rheumatoid arthritis patients

INTRODUCTION

The folate antagonist methotrexate (MTX) is the anchor-drug most widely applied disease modifying antirheumatic drug (DMARD) in the treatment of patients with rheumatoid arthritis (RA). It is used either as single agent or in combination with other DMARDs (e.g. sulfasalazine and hydroxychloroquine) and MTX use is obligate in most treatment strategies involving biological agents (anti-TNF α or -CD20 monoclonal antibodies).

The first pivotal step in the cellular pharmacology of MTX is its cell entry which can be mediated by at least 3 different routes; the reduced folate carrier (RFC), membrane-associated folate receptors (MFR) or a proton-coupled (low pH) folate transporter (PCFT). The latter transporter is mainly involved in intestinal folate uptake. The other transport routes harbour physiological and pharmacological

relevance for immune-competent cells by facilitating the uptake of natural reduced folate cofactors and folate antagonists like MTX. The RFC and MFR differ considerably in mechanism of (anti)folate uptake (transmembrane carrier vs. endocytosis / potocytosis), substrate specificity (low affinity folic acid / high affinity MTX vs. high affinity folic acid / low affinity MTX) and tissue specificity (constitutive vs. restricted expression).

For MFR, three isoforms (α , β and γ) have been identified. The α -isoform of MFR is overexpressed in specific types of cancer (ovarian cancer) while the γ -isoform is a secreted protein from haematopoietic cells. Selective expression of FR- β isoform has been described on activated macrophages in inflamed synovial fluid of RA patients and animal models of arthritis. Subsequently, FR- β was recognized as an attractive target for imaging of arthritis and therapeutically for selective antibody-guided or folate-conjugate guided delivery of toxins and other small / macromolecules. Thus far, targeting of FR with folate antagonists has only been explored on cancer cells / tissues overexpressing FR α .

Over the past decades, a second generation of folate antagonists has been designed and clinically evaluated from a perspective to circumvent common mechanisms of resistance to MTX, including impaired transport via the RFC, defective polyglutamylation, increased activity of the target enzyme DHFR and/or enhanced drug efflux. Based on this background, second generation antifolates included compounds that were more efficiently transported via RFC, were more efficiently polyglutamylated or independent of polyglutamylation, or target other key enzymes in folate metabolism other than DHFR, e.g. thymidylate synthase (TS) or glycinamide ribonucleotide transformylase. In the present study we set out to investigate whether distinct second generation antifolates may serve as selective targeting drugs for FR- β expressing cells in synovial tissue and/or immune-competent cells of RA patients. We identified the TS inhibitor BGC 945 as a prototypical antifolate drug that fulfilled the criteria of a high FR- β binding affinity and a low RFC affinity, thereby enabling selective drug uptake in FR- β expressing cells.

MATERIALS AND METHODS

Drugs

Folic acid was obtained from Sigma Chem. Co, St. Louis, MO, L-leucovorin from Merck Eprova, Schaffhausen, Switzerland, methotrexate from Pharmachemie, Haarlem, Netherlands and Pemetrexed / ALIMTA[®] (Eli Lilly) via the VUmc pharmacy department. The following folate antagonist drugs were obtained from the indicated companies / institutions; *Raltitrexed* / Tomudex[®] / ZD1694 (AstraZeneca, UK), *PT523* and *PT644* (Dr. A. Rosowsky, Harvard Medical School, Boston, MA), *GW1843* (Glaxo Wellcome, USA), *CB300635* (Institute of Cancer Research, Sutton, UK), *BGC 9331* and *BGC 945* (6*RS* and 6*S*) (BTG International Limited, London, UK), *5,10-dideazatetrahydrofolate* (*DDATHF*) (Eli Lilly, Indianapolis, IN), and *AG2034* (Agouron / Pfizer Pharmaceuticals, San Diego, CA). The chemical structure of these folate antagonists are depicted in Tables 2 and 3. [³,⁵,⁷,⁹-³H]Folic acid (20–40 Ci/mmol, MT783) was purchased from Moravek, Brea, CA.

Cell lines

Wild type Chinese Hamster Ovary (CHO-WT) cells, CHO cells transfected with FR- β (CHO-FR- β) and human nasopharyngeal epidermoid KB cells, expressing FR- α (American Type Culture Collection, Manassas, VA) were grown in folic-acid-free RPMI 1640 medium (Gibco, Grand Island, NY), supplemented with 10% foetal calf serum, 2 mM L-glutamine, 0.15 mg/ml proline and 100 units/ml penicillin and streptomycin. [³H]folic acid binding capacities of CHO-FR- β and KB cells were 0.5–1 pmol/10⁶ cells and 20–40 pmol/10⁶ cells, respectively. Human monocytic-macrophage THP1 cells (American Type Culture Collection, Manassas, VA) were grown in RPMI 1640 medium (Gibco, Grand Island, NY) containing 2.2 μ M folic acid, supplemented with 10% foetal calf serum, 2 mM L-glutamine, 0.15 mg/ml proline and 100 units/ml penicillin and streptomycin. All cell lines were kept at 37 °C in a humidified atmosphere containing 5% CO₂.

Synovial tissue samples

In this study we analysed synovial tissue biopsies derived from the knee joints of 15 RA-patients with active disease before treatment and after 4 months of treatment with MTX (starting dose of MTX: 7,5 mg/week; increasing stepwise to 15 mg/week over 12 weeks). Active disease was defined as ≥ 6 swollen or tender joints and levels of moderate or worse on the physician's and patient's assessments of

disease activity (Disease Activity Score-28; DAS-28). All patients had at least 1 clinically involved knee joint. Low dose prednisone (< 10 mg/day) and concomitant stable doses of nonsteroidal anti-inflammatory drug (NSAID) treatment were allowed. None of the patients ever used MTX before enrolling the study. In patients
5 taking other DMARDs, the treatment was terminated following a washout phase of 28 days. The arthroscopy procedure was performed as described previously as part of a joint study approved by the Medical Ethics committees of Leiden University Medical Center (Netherlands) and Leeds University Medical Centre (UK). See *Arthritis Rheum.* 2002; 46(8):2034–2038 and 2000; 43(8):1820–1830 (both Kraan *et al.*).

10 As non-inflammatory control synovial tissue we included 7 samples from patients with mechanical joint injury, provided by Dr. B. J. van Royen, department of Orthopedic Surgery, VU University Medical Center, Amsterdam, Netherlands. For peripheral blood sample collection, all patients signed an informed consent form and the study on ‘DMARD resistance’ was approved by the Medical Ethics committee of
15 the VU University Medical Center, Amsterdam, Netherlands.

FR- β immunohistochemistry

Immunohistochemical staining of cryostat sections (4 μ m) of synovial tissue biopsies from RA patients and controls was performed using a 3-step immunoperoxidase method as described previously. See *Cancer Res.* 2001; 61(8):3458–3464
20 (Maliepaard *et al.*) and 2000; 60(18): 5269–5277 (Scheffer *et al.*). Sections were stained with a specific antibody for FR- β (dilution 1: 3000), isotype control: normal rabbit-serum). Macrophages and T-cells were stained with 3A5 (dilution 1: 100) and anti-CD3-PE (dilution 1: 25; Dako, Glostrup, Denmark) monoclonal antibodies (isotype control: mouse immunoglobulin), respectively. Biotinylated swine anti-rabbit
25 IgG (Dako; dilution 1:200) and rabbit anti-mouse IgG (Dako; dilution 1:300) were used as secondary antibodies. Colour development was performed using 0.4 mg/ml AEC (aminoethyl carbazole). After counterstaining with haematoxylin, slides were mounted. Stained sections were analysed for FR- β , 3A5 and CD3 expression by digital image analysis, as described previously. See *Arthritis Res. Ther.* 2005; 7(4):
30 R862–R867 (Haringman *et al.*). In short, for each marker representative regions were used for image acquisition, using 400 $\times$ magnification. These regions were divided into 6 high-power fields (hpfs) with a 3-pixel overlap. Positive cells were evaluated by analysing 18 consecutive hpfs, scoring numbers of positive cells in the intimal

lining layer and the synovial sublining per mm².

Double-labelling immunofluorescence

FR- β was detected by swine-anti-rabbit HRP-labelled antibodies (dilution 1:200; Dako, Glostrup, Denmark) and development was with rhodamine / thyramine (red fluorescence) according to the instructions of the manufacturer (dilution 1:1000).
CD68 was detected by goat-anti-mouse biotinylated antibodies (dilution 1:100; Dako, Glostrup, Denmark) utilizing streptavidin Alexa-488 as a substrate (dilution 1:750; green fluorescence; Molecular Probes, Eugene, OR). Slides were mounted with Vectashield, containing 1 μ g/ml DAPI (for staining of nuclei) (Vector Laboratories Inc., Burlingame, CA). Cells were examined using a fluorescence microscope (Leica DMRB, Rijswijk, Netherlands).

Isolation of peripheral blood cells of RA-patients and culture conditions

Peripheral blood mononuclear cells were isolated from freshly obtained blood samples by gradient centrifugation (35 minutes at 400 $\times$ g) on Ficoll-Paque Plus (Amersham Pharmacia Biotechnologies, UK). After centrifugation the interphase was carefully collected and washed 3 times using Phosphate Buffered Salt solution (PBS) supplemented with 1% BSA. The lymphocyte fraction was counted and resuspended in IMDM culture medium (Invitrogen, Breda, Netherlands) which contained 10% FCS, 2 mM L-glutamine and 100 μ g/ml penicillin and streptomycin. Monocytes were isolated by adherence after 2 hours incubation at 37° C in culture flasks followed by RNA extraction or macrophage differentiation by culturing the monocytes for 7 days in the presence of 10 ng/ml Macrophage Colony Stimulating Factor (M-CSF) (Strathmann Biotech, Hamburg, Germany).

Peripheral blood lymphocytes (PBLs) remaining in suspension after monocyte adherence were collected for RNA isolation or used for T-cell activation by incubating them at a density of 1 $\times$ 10⁶ cells/ml with monoclonal anti-CD28 (5 μ g/ml, CLB-CD28/1, Sanquin, Amsterdam, Netherlands) and anti-CD3 (1 μ g/ml, CLB-T3/4.E, Sanquin, Netherlands) in goat anti-mouse (Dako, Glostrup, Denmark) coated 24 well plates. After 48 hours stimulation, activated T-cells were harvested for RNA isolation and the activation status was determined by measuring CD25 expression using flow-cytometry (FACScalibur, Becton & Dickinson).

FR- β mRNA expression in synovial tissue and peripheral blood cells of RA-patients

RNA from synovial tissue (n = 7), PBLs (n = 9), monocytes (n = 9), macrophages (n = 25), and activated T-cells (n = 22) from RA patients was isolated using the Qiagen RNeasy Plus isolation kit (Qiagen, Venlo, Netherlands) following the instructions provided by the manufacturer. Prior to RNA isolation, the frozen synovial tissue samples were powdered by grinding them in a liquid nitrogen prechilled mortar where after RPE buffer was added. Total RNA concentrations were determined using a Nanodrop ND-1000 spectrophotometer (Nanodrop Technologies, Wilmington, USA). Real-time reverse transcription-PCR (RT-PCR) methodology described previously by was used to measure simultaneously mRNA levels for FR- β and glyceraldehyde-3-phosphate (reference gene). See *Cancer Res.* 2006; 66(11): 5875–5882 (Qi *et al.*).

The reverse transcription step was carried out using Taqman Reverse Transcript Reagents from Applied Biosystems (Foster City, CA), following the protocol of the manufacturer. Briefly, 400 ng of total RNA was mixed with random hexamer primers (50 μ mol/L), RNase inhibitor (1 unit/ μ L), MultiScribe reverse transcriptase (5 units/ μ L), and deoxynucleoside triphosphates mix (2.5 mmol/L each) in reverse transcriptase buffer. The 10- μ L reaction mixture was first incubated at 25 °C for 10 minutes, then at 48 °C for 30 minutes and finally at 95 °C for 5 minutes.

The subsequent real-time PCR step for FR- β was carried out in the presence of 12.5 μ L of PCR Mastermix (Applied Biosystems), 0.5 μ L each of forward and reverse primer (CTGGCTCCTTGGCTG-AGTTC, 'GCCCAGCCTGGTTATCCA), and 0.5 μ L of Taqman probe (6FAM-TCCTCCCAGACTACCTGCCCTCAGC-TAMERA). The primers and the Taqman probe for control *GAPDH* gene were purchased from Applied Biosystems. The PCR conditions were 2 minutes at 50 °C, then 10 minutes at 95 °C, followed by 40 cycles of 15 seconds each at 95 °C, and finally 1 minute at 60 °C. Fluorescence data generated were monitored and recorded by the Gene Amp 5700 sequence detection system (Applied Biosystems). All samples were set up in triplicate and normalized to GAPDH values.

Analysis of FR- β / FR- α and RFC binding affinity for novel generation folate antagonists

An intact cell binding assay for competitive binding of [3 H]-folic acid and

novel anti-folate drugs to FR- β and FR- α was performed essentially as described previously. See *Mol. Pharmacol.* 1995; 48: 459–47 and *Cancer Res.* 1995; 55(17): 3795–3802 (both Westerhof *et al.*). Briefly, CHO- β cells (FR- β transfected Chinese Hamster Ovary cells) and KB cells (FR- α expressing cells) were detached by
5 incubation in PBS + 1 mM EDTA. Detached cells were suspended in ice-cold HEPES-buffered saline (140 mM NaCl, 20 mM HEPES, 6 mM KCl, 2 mM MgCl₂, 6 mM D-glucose, pH 7.4 with NaOH) to a cell concentration of 3×10^6 and 1×10^6 cells/ml, respectively. One ml of cell suspension was added to a series of Eppendorf tubes containing 100 pmol [³H]-folic acid (specific radioactivity: 2,000 dpm/pmol) in
10 the absence or presence of increasing concentrations of natural folate or folate antagonists. Following 10 minutes at 4 °C, cells were centrifuged in an Eppendorf centrifuge (30 s, 10,000 rpm), the supernatant was aspirated and cell pellets were resuspended in 200 μ l water and analysed for radioactivity (Optima Gold scintillation fluid, United Technologies, Packard, Brussels, Belgium). Non-specific binding of
15 [³H] (usually < 2% of specific binding) was determined by measuring cell-associated radioactivity in the presence of a 1000-fold molar excess of unlabeled folic acid. Concentrations of natural folates and selected folate antagonists required to displace 50% of [³H]folic acid from FR- β or FR- α were determined and presented as binding affinities relative to folic acid. For comparison, data for affinities of the RFC for
20 natural folates and folate antagonists were presented from previous studies. See. *Mol Pharmacol* 1995; 48: 459–471 Westerhof *et al.*) and *Cancer Res.* 2005; 65(24): 11721–11728 (Gibbs *et al.*).

Cell proliferation inhibition assay CHO-WT and CHO-FR β cells were seeded in individual wells of a 24-well tissue culture plate at a density of 1×10^4 /cm².
25 After 24 hours, 8 concentrations (with 2.5-fold increments) of folate antagonist drugs were added in the absence or presence (to block FR) of 1 μ M folic acid. After 72 hours incubation, cells were harvested by trypsinization and counted for cell viability as described before. See *Mol. Pharmacol.* 1999; 55(4): 761–769 (Jansen *et al.*).

In other experiments human monocytic-macrophage THP1 cells were tested
30 for antiproliferative effects of folate antagonists. To this end, 1 ml of cell suspension containing 1.25×10^5 cells were plated in individual wells of a 24-well tissue culture plate and incubated with 8 concentrations (with 2.5-fold increments) of folate antagonist drugs. After 72 hours drug exposure cell counts were performed with a haemocytometer and viability was checked by trypan blue exclusion.

RESULTS

FR- β synovial tissue immunohistochemistry

Immunohistochemical staining of all RA synovial tissue showed high expression of FR- β in the intimal lining layer as well as in the synovial sublining. The staining pattern for FR- β was consistent with 3A5 (macrophage) staining, whereas T-cell areas showed no staining (Figure 1 A-C). In fact, more detailed fluorescence microscopic analysis demonstrated co-localization of FR- β and CD68 (macrophage marker) on the cellular membranes of synovial tissue infiltrating macrophages and intimal macrophages (Figure 1 D-F). In non-inflammatory synovial tissue of orthopaedic controls, no staining for FR- β was observed, consistent with low numbers of macrophages (not shown).

Staining results were analysed by computer-assisted digital image analysis. A significant correlation was found between 3A5 and FR- β expression in the synovial sublining layer (cell counts/mm²) ($p = 0.04$) (Figure 2A). Median positive cell counts in the synovial sublining layer/mm² were 126 (range 9–630) for FR- β and 219 (range 11–622) for 3A5. Median numbers of macrophages decreased upon 4 months of treatment with MTX (from 219 to 119 positive cell counts in the synovial sublining layer/mm², $p = 0.14$). FR- β expression after 4 months of treatment with MTX was positively correlated ($r = 0.31$) with DAS28 improvement (Δ DAS28), but did not reach statistical significance ($p = 0.11$) (Figure 2B).

FR- β mRNA expression in RA-synovial tissue and peripheral blood cells of RA patients

To further confirm the differential expression of FR- β in synovial tissue, FR- β mRNA levels were determined by PCR analysis in synovial tissue biopsies as compared to those in peripheral blood cells of RA-patients, including lymphocytes, *ex vivo* activated T-cells, and peripheral blood monocytes and *ex vivo* monocyte-derived macrophages. FR- β mRNA levels were presented relative to CHO-FR- β cells (set at 100%). Ranking of FR- β mRNA expression was highest for RA-synovial tissue (median 17% compared to CHO- β cells) > *ex vivo* monocyte-derived macrophages (3% compared to CHO- β cells) > peripheral blood lymphocytes (PBLs) (0.7%) >> monocytes (0.02%) and *ex vivo* activated T-cells (< 0.001%) (Figure 3).

Binding affinities of FR- β vs. FR- α for folate antagonists

Binding affinities of FR α for selected folate antagonists were previously

reported by Westerhof *et al.* but to what extent they overlap or differ for the FR- β isoform has not been established. To this end, binding affinities by FR- β versus FR α for a series of folate-based inhibitors of DHFR, TS and GARTFase were determined and shown in Figure 4. With respect to folic acid, both FR- β and FR α displayed a rather low affinity for the group of folate antagonist inhibitors of DHFR. Binding affinity of FR- β for MTX is approximately 50-fold lower than for folic acid. The binding affinity for PT523 is markedly lower than for MTX (0.3% of folic acid) while PT644, the 5-methyl analogue of PT523, showed an affinity comparable to MTX. Of note, FR- α exhibits a good binding affinity for all tested folate-based TS inhibitors, but binding affinities of FR- β for pemetrexed, raltitrexed and BGC 9331 were markedly lower (16–30 fold) than for FR α . Retention of a high binding affinity of FR- β was observed for the TS inhibitors CB300635 (161% of folic acid) and (6*RS*)-BGC 945 (89% of folic acid) and (6*S*)-BGC 945 (46% of folic acid). FR- β also exhibited a proficient binding affinity for the GARTFase inhibitors DDATHF (27% of folic acid) and AG2034 (54% of folic acid) even though FR α binding affinities for these compounds were 2.5-fold higher. Together, these results demonstrate a broad differential in binding affinities of FR- β for folate antagonists, among which several folate antagonists revealed a markedly higher binding affinity than for MTX.

Folate antagonist induced growth inhibition of FR- β expressing cells

To investigate whether the folate antagonists used in the current study would all convey a potential growth inhibitory effects against macrophage-like type of cells, this parameter was investigated for human monocytic-macrophage THP1 cells (Table 1). Consistent with RFC as the dominant transport route in THP1 cells, potent growth inhibitory effects were observed for all folate antagonists, except CB300635 and BGC 945, two compounds that have a poor affinity for RFC. Since THP1 cell line model is FR-negative (in contrast to activated synovial macrophage), three folate antagonists for which FR- β displayed the highest binding affinity (CB300635, AG2034 and BGC 945) were evaluated for their potency to target FR- β by provoking cell growth inhibition of CHO-FR- β cells. Against CHO/WT cells, BGC 945 only induced growth inhibition at extracellular concentrations > 1000 nM (Figure 5A). Remarkably, growth inhibition of CHO-FR- β cells was induced at markedly lower concentrations (10–50 nM) of BGC 945. The addition of folic acid to these cell cultures completely abrogated the activity of BGC 945, consistent with a blockade of

FR. Despite displaying the highest binding affinity for FR- β , CB300635 was not markedly potent in inducing growth inhibition in CHO/FR- β cells (Figure 5B). The notion that co-administration of folic acid abrogated activity of CB300635 suggest that FR- β is involved in the cellular uptake of this compound. Finally, AG2034 may
5 utilize both the constitutively expressed RFC and FR as route for cell entry (Fig. 5). As such, AG2034 displayed a growth inhibitory potential against CHO/WT cells and to a greater extent to CHO/FR- β cells (Figure 5C/D). Consistently, abrogation of AG2034 growth inhibitory effects by FR- β blocking (with folic acid) and RFC blocking (with LV) are only partial (Figure 5C/D).

10

DISCUSSION

Since MTX is the anchor-drug in many therapeutic regimens for RA treatment, delineation of genetic, biochemical and metabolic parameters that could assist in predicting and/or improving the therapeutic response to MTX have received considerable recent interest. This study focused specifically on the role of cell membrane
15 transport of MTX, which, in activated synovial macrophages, is mediated predominantly by the folate receptor β isoform. Given the notion that the molecular and functional properties of FRs and the constitutively expressed RFC differ considerably, a better understanding of the properties of FR- β may facilitate a better
20 therapeutic window by selective targeting of FR- β over RFC.

Here we showed that FR- β expression primarily co-localized with macrophages in the intimal lining layer and the synovial sublining of RA patients and may therefore be an attractive target for folate antagonists. Screening for binding affinities of a series of second generation folate antagonists, some of which with proven
25 anticancer activity, revealed that the group of DHFR inhibitors all had a rather low FR- β affinity. This is consistent with previously reported structure activity relationships demonstrating that the α isoform of FR had low affinities for folate antagonists with a 2,4-NH₂-based structure (see Table 2). Interestingly, while FR α demonstrated a relatively high binding affinity for all tested folate-based inhibitors of
30 thymidylate synthase, for FR- β this was only retained for 3 compounds (CB300635, GW1843 and BGC 945) that share a common chemical property of 3-ring structures and/or glutamate side chain modifications (see Tables 2 and 3). The latter modification also markedly suppresses its ability to be transported via the RFC and thus

contributes to a greater FR-selectivity. In fact, selective targeting by BGC 945 for FR- α and not RFC was demonstrated in FR α over expressing cell lines. In addition to folate-based TS inhibitors, FR- β also exhibited moderate to high binding affinities for folate-based GARTFase inhibitors (AG2034 and DDATHF) which classifies them as folate antagonist drugs that can be transported both via RFC and FR.

FR- β -transfected CHO cells were used as a model system to evaluate the efficiency of FR- β -mediated cellular uptake of folate antagonists by conveying anti-proliferative effects. This cell line model may be clinically representative, based on [^3H]-folic acid binding levels and FR- β mRNA levels that are compatible with FR- β mRNA levels in synovial tissue of RA patients (Fig. 4). The largest differential in activity between control (RFC-expressing) CHO cells and FR- β transfected cells was observed for BGC 945, consistent with a poor affinity for transport via RFC and a high FR- β binding affinity. FR- β mediated uptake of BGC 945 could be inhibited by blocking of the receptor with excess folic acid, implying that circulating natural folates in synovial tissue / plasma could attenuate the potential activity of BGC 945 *in vivo* either by receptor occupancy / competition or by receptor down-regulation.

Example 2: Compounds of the invention

Table 3 shows the structures of the following compounds of the invention:

- | | |
|----------------------|----------------------|
| • CB300638 (BGC 638) | • CB300944 |
| • (6S)-CB300638 | • CB300945 (BGC 945) |
| • CB300935 | • (6S)-CB300945 |
| • CB300936 | • CB300947 |
| • CB300940 | • CB300951 |

These were prepared according to the methods given in WO 94/11354 A1, WO 03/020300 A1, WO 03/020706 A1, WO 03/020748 A1, *J. Med. Chem.*, 2000, **43**, 1910–1926, *Tetrahedron*, **63** (7), 2007, 1537–1543 (Bavetsias *et al.*) and *Cancer Research* **65**, 2005, 11721–11728 (Gibbs *et al.*).

Example 3: Formulation

The following illustrate representative pharmaceutical dosage forms containing a cyclopenta[g]quinazoline of formula (I), particularly in pharmaceutically ac-

ceptable salt form, for therapeutic or prophylactic use in humans:

5	(a)	<u>Tablet I</u>	<u>mg/tablet</u>
		Cyclopenta[g]quinazoline salt	100
		Lactose Ph.Eur.	182.75
		Croscarmellose sodium	12.0
		Maize starch paste (5% w/v paste)	2.25
		Magnesium stearate	3.0
10	(b)	<u>Tablet II</u>	<u>mg/tablet</u>
		Cyclopenta[g]quinazoline salt	50
		Lactose Ph.Eur.	223.75
		Croscarmellose sodium	6.0
		Maize starch	15.0
15		Polyvinylpyrrolidone (5% w/v paste)	2.25
		Magnesium stearate	3.0
20	(c)	<u>Tablet III</u>	<u>mg/tablet</u>
		Cyclopenta[g]quinazoline salt	1.0
		Lactose Ph.Eur.	93.25
		Croscarmellose sodium	4.0
		Maize starch paste (5% w/v paste)	0.75
		Magnesium stearate	1.0
25	(d)	<u>Capsule</u>	<u>mg/capsule</u>
		Cyclopenta[g]quinazoline salt	10.0
		Lactose Ph.Eur.	488.5
		Magnesium stearate	1.5
30	(e)	<u>Injection I</u>	<u>(50 mg/ml)</u>
		Cyclopenta[g]quinazoline salt	5.0% w/v
		1M Sodium hydroxide solution	15.0% v/v
		0.1M Hydrochloric acid (to adjust pH to 7.6)	
		Polyethylene glycol 400	4.5% w/v

Water for injection to 100%

(f) Injection II (10 mg/ml)

	Cyclopenta[g]quinazoline salt	1.0% w/v
5	Sodium phosphate BP	3.6% w/v
	0.1M Sodium hydroxide solution	15.0% v/v
	Water for injection to 100%	

(g) Injection III (1 mg/ml, buffered to pH 6)

10	Cyclopenta[g]quinazoline salt	0.1% w/v
	Sodium phosphate BP	2.26% w/v
	Citric acid	0.38% w/v
	Polyethylene glycol 400	3.5% w/v
	Water for injection to 100%	

15

The above formulations may be prepared by conventional procedures well known in the pharmaceutical art. The tablets (a) to (c) may be enteric coated by conventional means, for example with a coating of cellulose acetate phthalate.







20 **Table 1. Growth inhibitory effects of folate antagonists against human monocytic-macrophage THP1 cells[#]**

Folate antagonist	IC ₅₀ (nM)
MTX	7.1 ± 0.5
PT523	2.2 ± 0.3
PT644	1.0 ± 0.4
Raltitrexed	2.4 ± 0.5
Pemetrexed	10.7 ± 2.2
GW1843	1.6 ± 0.1
BGC 9331	8.7 ± 0.8
CB300635	4850 ± 285
BGC 945	3630 ± 350
DDATHF	9.8 ± 0.8
AG2034	3.2 ± 0.2

[#]THP-1 cells were grown in RPMI-1640 medium containing 2.2 µM folic acid (which would block any FR activity).



Drug exposure time; 72 hours. Results are the mean ± S.D. of 3 separate experiments.

Table 2: Structures of comparative compounds



Designation	Structure
Folic acid	
Leucovorin	
MTX	
PT523	
PT644	
GW1843	

Designation	Structure
Pemetrexed	
Raltitrexed	
BGC 9331	
CB300635	
DDATHF	
AG2034	


Table 3: Structures of the compounds of the invention




CB300638**(6S)-CB300638**

5



CB300935**CB300936**

10

CB300940

CB300944**CB300945 (BGC 945)****(6S)-CB300945**


5

CB300947**CB300951**

10

CLAIMS

1. A cyclopenta[g]quinazoline derivative, containing an L-Glu-γ-D-Glu dipeptide group, of formula (I):



wherein:

R¹ is amino, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ fluoroalkyl or methoxy-C₁₋₄-alkyl;

R² is hydrogen, C₁₋₄ alkyl, C₃₋₄ alkenyl, C₃₋₄ alkynyl, C₂₋₄ hydroxyalkyl C₂₋₄ halogenoalkyl or C₁₋₄ cyanoalkyl; and

Ar is phenylene, thiophenediyl, thiazolediyl, pyridinediyl or pyrimidinediyl which may optionally bear one or two substituents selected from halogeno, hydroxy, amino, nitro, cyano, trifluoromethyl, C₁₋₄ alkyl and C₁₋₄ alkoxy.

the compound (I) optionally being in the form of a pharmaceutically acceptable salt or ester;

for the treatment of rheumatoid arthritis or acute myeloid leukaemia.

2. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 1 wherein R¹ is C₁₋₄ alkyl or C₁₋₄ hydroxyalkyl.

3. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 2 wherein R¹ is methyl or hydroxymethyl.

4. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in any preceding claim wherein R² is methyl, ethyl, propyl, prop-2-enyl, prop-2-ynyl, 2-hydroxyethyl, 2-fluoroethyl, 2-bromoethyl or 2-cyanoethyl.

5. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 4 wherein R² is methyl or prop-2-ynyl.

6. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in any preceding claim wherein Ar is 1,4-phenylene which may optionally bear one or two substituents selected from the group consisting of chloro and especially fluoro, thiophene-2,5-diyl, thiazole-2,5-diyl or pyridine-2,5-diyl.
7. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 6 wherein Ar is 1,4-phenylene, 1,4-phenylene having a 2-fluoro substituent or pyridine 2,5-diyl.
8. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 7 wherein Ar is 1,4-phenylene or 2-fluoro-1,4-phenylene.
9. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 1 wherein the cyclopenta[g]-quinazoline derivative comprises:
- N*-{*N*-{4-[*N*-(2-methyl-4-oxo-3,4,7,8-tetrahydro-6*H*-cyclopenta[g]quinazolin-6-yl)-*N*-(prop-2-ynyl)amino]benzoyl}-L- γ -glutamyl}-D-glutamic acid; or
- N*-{*N*-{4-[*N*-(2-hydroxymethyl-4-oxo-3,4,7,8-tetrahydro-6*H*-cyclopenta[g]quinazolin-6-yl)-*N*-(prop-2-ynyl)amino]benzoyl}-L- γ -glutamyl}-D-glutamic acid;
- or a pharmaceutically acceptable salt or ester thereof.
10. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in claim 9 wherein the cyclopenta[g]-quinazoline derivative is present as its 6*S* isomer.
11. A cyclopenta[g]quinazoline derivative for the treatment of rheumatoid arthritis or acute myeloid leukaemia as claimed in any preceding claim wherein the cyclopenta[g]quinazoline derivative is administered together with a pharmaceutically acceptable diluent or carrier.
12. A method for treating rheumatoid arthritis or acute myeloid leukaemia in a patient in need of such treatment which comprises administering to said patient an effective amount of a cyclopenta[g]quinazoline derivative claimed in any preceding claim.

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5



INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2009/000687

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K38/05 A61P19/00 A61P35/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HENDERSON E A ET AL: "Targeting the alpha-folate receptor with cyclopenta[g]quinazoline-based inhibitors of thymidylate synthase" BIOORGANIC & MEDICINAL CHEMISTRY, ELSEVIER SCIENCE LTD, GB, vol. 14, no. 14, 15 July 2006 (2006-07-15), pages 5020-5042, XP025133399 ISSN: 0968-0896 [retrieved on 2006-07-15]. ----- -/--	1-12



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

13 May 2009

Date of mailing of the international search report

26/05/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Schleifenbaum, A

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2009/000687

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JACKMAN ANN L ET AL: "Antifolates targeted specifically to the folate receptor." ADVANCED DRUG DELIVERY REVIEWS 29 APR 2004, vol. 56, no. 8, 29 April 2004 (2004-04-29), pages 1111-1125, XP002527440 ISSN: 0169-409X table 3	1-12
A	GIBBS DAVID D ET AL: "BGC 945, a novel tumor-selective thymidylate synthase inhibitor targeted to alpha-folate receptor-overexpressing tumors." CANCER RESEARCH 15 DEC 2005, vol. 65, no. 24, 15 December 2005 (2005-12-15), pages 11721-11728, XP002527441 ISSN: 0008-5472 cited in the application table 1	1-12
A	THETI DAVINDER S ET AL: "Selective delivery of CB300638, a cyclopenta[g]quinazoline-based thymidylate synthase inhibitor into human tumor cell lines overexpressing the alpha-isoform of the folate receptor." CANCER RESEARCH 1 JUL 2003, vol. 63, no. 13, 1 July 2003 (2003-07-01), pages 3612-3618, XP002527442 ISSN: 0008-5472 abstract	1-12
A	WO 03/020748 A (BTG INT LTD [GB]; BAVETSIAS VASSILIOS [GB]; JACKMAN ANN LESLEY [GB]) 13 March 2003 (2003-03-13) cited in the application claim 10; example 4	1-12
A	WO 03/020300 A (BTG INT LTD [GB]; BAVETSIAS VASSILIOS [GB]; JACKMAN ANN LESLEY [GB]) 13 March 2003 (2003-03-13) cited in the application claims 1,8,9; example 4	1-12
	----- -/-- -----	

INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2009/000687

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	NAKASHIMA-MATSUSHITA N ET AL: "Selective expression of folate receptor beta and its possible role in methotrexate transport in synovial macrophages from patients with rheumatoid arthritis" ARTHRITIS AND RHEUMATISM, LIPPINCOTT, PHILADELPHIA, US, vol. 42, no. 8, 1 August 1999 (1999-08-01), pages 1609-1616, XP002284073 ISSN: 0004-3591 the whole document -----	1-12
A	WO 03/072091 A (UNIV OHIO STATE RES FOUND [US]; LEE ROBERT J [US]; RATNAM MANOHAR [US]) 4 September 2003 (2003-09-04) cited in the application the whole document -----	1-12
A	XIN W ET AL: "Differential stereospecificities and affinities of folate receptor isoforms for folate compounds and antifolates" BIOCHEMICAL PHARMACOLOGY, PERGAMON, OXFORD, GB, vol. 44, no. 9, 3 November 1992 (1992-11-03), pages 1898-1901, XP025567617 ISSN: 0006-2952 [retrieved on 1992-11-03] page 1901, left-hand column, paragraph 2 -----	1-12
A	BRIGLE K E ET AL: "Increased expression and characterization of two distinct folate binding proteins in murine erythroleukemia cells" BIOCHEMICAL PHARMACOLOGY, PERGAMON, OXFORD, GB, vol. 47, no. 2, 20 January 1994 (1994-01-20), pages 337-345, XP023764830 ISSN: 0006-2952 [retrieved on 1994-01-20] page 342, right-hand column, paragraph 2 - page 343, left-hand column, paragraph 1 -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2009/000687

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03020748	A	13-03-2003	AU 2002321576 B2	11-09-2008
			BR 0212215 A	21-09-2004
			CA 2458813 A1	13-03-2003
			EP 1421105 A1	26-05-2004
			JP 2005509600 T	14-04-2005
			MX PA04001905 A	19-04-2005
			US 2004242606 A1	02-12-2004
			US 2008015206 A1	17-01-2008
			US 2009023760 A1	22-01-2009
			ZA 200401587 A	10-03-2005
WO 03020300	A	13-03-2003	EP 1420809 A1	26-05-2004
			US 2005009851 A1	13-01-2005
WO 03072091	A	04-09-2003	AU 2003216445 A1	09-09-2003
			CA 2503094 A1	04-09-2003
			EP 1485076 A1	15-12-2004
			JP 2005519078 T	30-06-2005