



HU000026041T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 026 041**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 12 717685**(22) A bejelentés napja: **2012. 04. 26.**

(96) Az európai bejelentés bejelentési száma:

EP 20120717685

(97) Az európai bejelentés közzétételi adatai:

EP 2702061 A1 **2012. 11. 01.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2702061 B1 **2015. 07. 29.**(51) Int. Cl.: **C07D 473/34** (2006.01)**A61K 314/37** (2006.01)**A61K 3149/85** (2006.01)**C07D 487/04** (2006.01)**A61P 35/00** (2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/EP 12/057671

(87) A nemzetközi közzétételi szám:

WO 12146666

(30) Elsőbbségi adatok:

11382124	2011. 04. 29.	EP
201161502550 P	2011. 06. 29.	US

(72) Feltaláló(k):

BERNAL ANCHUELA, Francisco Javier, Sant Feliu de Llobregat, Barcelona (ES)
CARRASCAL RIERA, Marta, Sant Feliu de Llobregat, Barcelona (ES)
CATURLA JAVALOYES, Juan Francisco, Sant Feliu de Llobregat, Barcelona (ES)
GRACIA FERRER, Jordi, Sant Feliu de Llobregat, Barcelona (ES)
MATASSA, Victor Giulio, Sant Feliu de Llobregat, Barcelona (ES)
TERRICABRAS BELART, Emma, Sant Feliu de Llobregat, Barcelona (ES)
TALTAVULL MOLL, Joan, Sant Feliu de Llobregat, Barcelona (ES)
ERRA SOLA, Montserrat, Sant Feliu de Llobregat, Barcelona (ES)

(73) Jogosult(ak):

Almirall S.A., 08022 Barcelona (ES)

(74) Képviselő:

PINTZ ÉS TÁRSAI Szabadalmi, Védjegy és Jogi Iroda Kft., Budapest

(54)

Pirrolo-triazinon származékok P13K inhibitoroként

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 702 061 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
29.07.2015 Bulletin 2015/31

(51) Int Cl.:
C07D 473/34 (2006.01) **C07D 487/04** (2006.01)
A61K 31/4985 (2006.01) **A61K 31/437** (2006.01)
A61P 35/00 (2006.01)

(21) Application number: **12717685.7**

(86) International application number:
PCT/EP2012/057671

(22) Date of filing: **26.04.2012**

(87) International publication number:
WO 2012/146666 (01.11.2012 Gazette 2012/44)

(54) **PYRROLOTRIAZINONE DERIVATIVES AS PI3K INHIBITORS**

PYRROLOTRIAZINONDERIVATIVE ALS PI3K-INHIBITOREN

DÉRIVÉS DE PYRROLOTRIAZINON EN TANT QU'INHIBITEURS PI3K

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

- **TALTAVULL MOLL, Joan**
Sant Feliu de Llobregat, Barcelona (ES)
- **ERRA SOLA, Montserrat**
Sant Feliu de Llobregat, Barcelona (ES)

(30) Priority: **29.04.2011 EP 11382124**
29.06.2011 US 201161502550 P

(74) Representative: **Srinivasan, Ravi Chandran**
J A Kemp
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(43) Date of publication of application:
05.03.2014 Bulletin 2014/10

(56) References cited:
WO-A1-03/035075 WO-A1-2007/023186
WO-A1-2010/111432

(73) Proprietor: **Almirall S.A.**
08022 Barcelona (ES)

(72) Inventors:
• **BERNAL ANCHUELA, Francisco Javier**
Sant Feliu de Llobregat, Barcelona (ES)
• **CARRASCAL RIERA, Marta**
Sant Feliu de Llobregat, Barcelona (ES)
• **CATURLA JAVALOYES, Juan Francisco**
Sant Feliu de Llobregat, Barcelona (ES)
• **GRACIA FERRER, Jordi**
Sant Feliu de Llobregat, Barcelona (ES)
• **MATASSA, Victor Giulio**
Sant Feliu de Llobregat, Barcelona (ES)
• **TERRICABRAS BELART, Emma**
Sant Feliu de Llobregat, Barcelona (ES)

- **KIM K S ET AL: "Synthesis and SAR of pyrrolotriazine-4-one based Eg5 inhibitors", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, ELSEVIER SCIENCE, GB, vol. 16, no. 15, 1 August 2006 (2006-08-01), pages 3937-3942, XP025107103, ISSN: 0960-894X, DOI: 10.1016/J.BMCL.2006.05.037 [retrieved on 2006-08-01]**
- **MCNEILL HELEN ET AL: "When pathways collide: collaboration and connivance among signalling proteins in development", NATURE REVIEWS MOLECULAR CELL BIOLOGY, NATURE PUBLISHING, GB, vol. 11, no. 6, 1 June 2010 (2010-06-01), pages 404-413, XP009140410, ISSN: 1471-0072, DOI: 10.1038/NRM2902**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] When cells are activated by extracellular stimuli, intracellular signalling cascades involving the regulation of second messengers are initiated that eventually produce a response of the cell to the stimuli. Phosphoinositide 3-Kinases (PI3Ks) are among the enzymes involved in early signalling events to a plethora of different types of stimuli. PI3Ks phosphorylate the 3-hydroxyl group of the inositol ring of phosphatidylinositol (PtdIns), PtdIns-4-phosphate (PtdIns4P), and PtdIns-4,5-bisphosphate (PtdIns(4,5)P₂). The resulting 3-phosphoinositides mediate correct localization and subsequent activation of a number of downstream effector proteins that bind to the lipids via specific lipid binding sequences such as the pleckstrin homology (PH) domain (Vanhaesebroeck B, 2010, Nat Rev Mol Cell Biol 5:11381-6).

[0002] The PI3K family is divided into 3 different classes (PI3K class I, class II, and class III), depending on substrate preference and structural features.

[0003] The best characterized is the PI3K class I with the preferential substrate PtdIns-(4,5)P₂. It englobes 4 different isoforms which originally were further subdivided into class IA (p110a, p110b, p110d), binding to a p85 type of regulatory subunit, and class IB (p110g) which is regulated by p101 and p87 subunits. Whereas p110a (PI3K α or PI3K α) and p110b (PI3K β or PI3K β) isoforms are expressed ubiquitously, p110g (PI3K γ or PI3K γ) and especially p110d (PI3K δ or PI3K δ) have a more restricted expression pattern and seem to play a major role in leukocytes (Kok K, Trends Biochem Science 34:115-127, 2009).

[0004] Both, PI3K δ and PI3K γ are involved in activation of immune cells by a large variety of different stimuli. Pharmacological inhibition or genetic deficiency in active p110d has been shown to inhibit T cell proliferation and cytokine production in response to different stimuli such as anti-CD3, anti-CD3/CD28, superantigen or antigen in vitro (Ji H, Blood 2007; Okkenhaug K, Science 2002; Garcon F, 2009; Soond DR, Blood 2010; Herman SEM, Blood June 3, 2010; William O, Chemistry & Biology 17, 2010) and to suppress concanavalin A and anti-CD3 induced cytokine production as well as antigen-dependent tissue retention in vivo (Soond DR, Blood 2010; Jarmin SJ, JCI 2008). In addition, B cell function is critically dependent on functional PI3K δ activity as demonstrated by suppressed B cell proliferation and cytokine release in vitro in response to anti-IgM (Bilancio A, Blood 107, 2006), toll like receptor agonists such as LPS and oligodeoxynucleotides (Dil N, Mol Immunol 46, 2009) or impaired ability to stimulate antigen-specific T cells (Al-Alwan M, JI 2007) in the absence of functional p110d or pharmacological inhibition. In vivo, PI3K γ deficient mice display partially suppressed antibody production upon immunization (Garcon F, 2009; Durand CA, JI 2009). Further studies have demonstrated an important role of PI3K δ in inhibition of T cell apoptosis and in TH17 differentiation (Haylock-Jacobs S, J. Autoimmun 2010).

[0005] In addition, mast cell degranulation was reduced in cells from mice with inactivated PI3K δ or by pharmacological inhibition of PI3K δ (Ali K, Nature 431:1007-1011, 2004; Ali K, Journal of Immunology 180:2538-2544, 2008) and basophil activation via the Fc ϵ receptor is suppressed by pharmacological inhibition of PI3K δ (Lannutti BJ, Blood Oct. 2010).

[0006] In terms of neutrophil function, PI3K δ inhibition inhibits migration of mouse neutrophils to fMLP in an under-agarose migration assay by inhibiting cell polarization and directional movement (Sadhu C, JI 170, 2003) and mouse PI3K δ deficient or inhibitor treated neutrophils show slightly (25%) reduced in vitro chemotaxis to LTB₄, whereas in vivo accumulation in the lung in response to LPS was reduced by more than 80%, indicating an important role of PI3K δ in endothelial cells for mediating PMN transendothelial migration (Puri KD, Blood 103, 2004). Furthermore, TNF induced neutrophil infiltration to an air pouch in mice and elastase release is partially inhibited by a PI3K δ selective inhibitor (Sadhu C, Biochem Biophys Res Comm 308, 2003). In addition, TNF mediated priming of oxidative burst by human neutrophils depends on PI3K δ activity (Condliffe AM, Blood 106, 2005).

[0007] In contrast to the dominant role of PI3K δ in lymphocyte activation, PI3K γ seems to affect primarily chemotaxis of different immune cells induced by various mediators and chemokines (Martin AL, JI 180, 2008; Thomas MS, J Leukoc Biol 84, 2008; Jarmin SJ, JCI 2008; Matthew T, Immunology 126, 2008), as well as degranulation and oxidative burst of innate immune cells induced by GPCR mediated stimuli such as fMLP, IL-8 or C5a (Condliffe AM, Blood 106, 2005; Yum HK, JI 167, 2001; Pinho V, JI 179, 2007).

[0008] The above mentioned findings suggest that selective PI3K δ or dual PI3K δ /PI3K γ pharmacological inhibition represents a promising approach for treating a variety of diseases such as respiratory diseases (asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis), allergic diseases (allergic rhinitis), inflammatory or autoimmune diseases (rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, myasthenia gravis, acute disseminated encephalomyelitis, idiopathic thrombocytopenic purpura, Sjogren's syndrome, autoimmune hemolytic anemia, type I diabetes, psoriasis, acrodermatitis, angiodermatitis, atopic dermatitis, contact dermatitis, eczema, acne, chronic urticaria, blistering diseases including but not limited to bullous pemphigoid, scleroderma, dermatomyositis, etc.), cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain (such as pain associated with rheumatoid arthritis or osteoarthritis, back pain, general inflammatory pain, inflammatory neuropathic pain, trigeminal neuralgia or central pain) as well as in bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors (such

as pancreatic cancer; bladder cancer; colorectal cancer; breast cancer; prostate cancer; renal cancer; hepatocellular cancer; lung cancer; ovarian cancer; cervical cancer; gastric cancer; esophageal cancer; head and neck cancer; non-small cell lung cancer and small-cell lung cancer; melanoma; neuroendocrine cancers; central nervous system cancers; brain tumors; bone cancer; soft tissue sarcoma; chronic lymphocytic leukemia, B-cell acute lymphoblastic leukemia, T-cell acute lymphoblastic leukaemia, non-hodgkins lymphoma, B-cell lymphoma, acute myeloid leukaemia; cutaneous T cell lymphoma, premalignant and malignant skin conditions including but not limited to basal cell carcinoma (BCC), squamous cell carcinoma (SCC) or actinic keratosis (AK)).

[0009] There is substantial experimental evidence supporting this view. In rodent models of allergic lung inflammation, genetic or pharmacological inactivation of PI3Kd or dual PI3Kd/g dual inhibition reduces cell influx, mucus production, cytokine production and airway hyperreactivity (Nashed et al. 2007, Eur J Immunol 37:416; Lee et al. 2006, FASEB J 20:455 & Lee KS et al. 2006, J Allergy Clin Immunol 118:403; Doukas J, JPET 2009;328:758; Par SJ, ERJ 2010). Moreover, LPS induced lung neutrophil infiltration is blocked by PI3Kd inhibition (Puri KD, Blood 2004;103:3448) and inflammation in response to LPS or tobacco smoke exposure is suppressed by a dual PI3Kd/g inhibitor (Doukas J, JPET 2009;328:758). Moreover, PI3Kd seems to be involved in the reduction of responsiveness to corticosteroid treatment associated with oxidative stress and chronic obstructive pulmonary disease (COPD). This notion is based on the findings that tobacco smoke induced inflammation remains responsive to treatment with budesonide, whereas wild type or PI3Kg deficient mice develop resistance to corticosteroid treatment (Marwick JA, JRCCM 179:542-548, 2009). Similar results were obtained with a PI3Kd selective inhibitor (To Y, AJRCCM 182:897-904, 2010). In addition, in vitro induction of corticosteroid resistance by oxidative stress is prevented by PI3Kd inhibition (To Y, AJRCCM 2010). In COPD patients, lung macrophages display increased expression of PI3Kd and phosphorylation of its downstream effector Akt and non-selective PI3K or PI3Kd- selective inhibition restored the impaired inhibitory efficacy of dexamethasone in PBMC from COPD patients (To Y, AJRCCM 182:897-904, 2010; Marwick JA, JACI 125:1146-53, 2010).

[0010] Furthermore, PI3Kd inhibition was effective in a model of contact hypersensitivity (Soond DR, Blood Jan 2010). In a model of experimental autoimmune encephalomyelitis, PI3Kd deficiency or pharmacological inhibition of PI3Kd attenuated T cell activation and function and reduced T cell numbers in the CNS, suggesting a therapeutic benefit of PI3Kd inhibitor in multiple sclerosis and other Th17-mediated autoimmune diseases (Haylock-Jacobs S, J. Autoimmun 2010). In line with that, genetic deficiency or pharmacological inhibition of PI3Kd diminished joint erosion in a mouse model of inflammatory arthritis (Randis TM, Eur J Immunol 38, 2008). Concerning metabolic diseases, PI3Kd overexpression seems to contribute to excessive vascular contraction and PI3Kd inhibition normalized vascular contractile responses in a mouse model of type I diabetes, suggesting a therapeutic potential of PI3Kd blockade to treat vascular dysfunction in diabetic patients (Pinho JF, Br. J. Pharmacol 161, 2010).

[0011] There is also substantial experimental evidence supporting that genetic or pharmacological inactivation of PI3Kd or dual PI3Kd/g dual inhibition is effective in the treatment of cancers including but not restricted to leukemias, such as chronic lymphocytic leukemia, B-cell acute lymphoblastic leukemia, T-cell acute lymphoblastic leukaemia, non-hodgkins lymphoma, B-cell lymphoma, acute myeloid leukaemia, myelo-dysplastic syndrome or myelo-proliferative diseases. In this aspect, the selective PI3Kd inhibitor CAL-101 demonstrated anti-proliferative properties on different tumor cells in vitro and efficacy in cancer patients with a dysregulated PI3Kd activity, such as chronic lymphocytic leukemia (Hermann SE, Blood 116:2078-88, 2010; Lannutti BJ, Blood Oct. 2010).

[0012] Conditions in which targeting of the PI3K pathway or modulation of the PI3 Kinases, particularly PI3Kd or PI3Kd/g, are contemplated to be therapeutically useful for the treatment or prevention of diseases including: respiratory diseases (asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis), allergic diseases (allergic rhinitis), inflammatory or autoimmune-mediated diseases (rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, myasthenia gravis, acute disseminated encephalomyelitis, idiopathic thrombocytopenic purpura, Sjogren's syndrome, autoimmune hemolytic anemia, type I diabetes, psoriasis, acrodermatitis, angiodermatitis, atopic dermatitis, contact dermatitis, eczema, acne, chronic urticaria, scleroderma, dermatomyositis and blistering diseases including but not limited to bullous pemphigoid), cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain (such as pain associated with rheumatoid arthritis or osteoarthritis, back pain, general inflammatory pain, inflammatory neuropathic pain, trigeminal neuralgia or central pain) as well as in bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors (such as pancreatic cancer; bladder cancer; colorectal cancer; breast cancer; prostate cancer; renal cancer; hepatocellular cancer; lung cancer; ovarian cancer; cervical cancer; gastric cancer; esophageal cancer; head and neck cancer; non-small cell lung cancer and small-cell lung cancer; melanoma; neuroendocrine cancers; central nervous system cancers; brain tumors; bone cancer; soft tissue sarcoma; chronic lymphocytic leukemia, B-cell acute lymphoblastic leukemia, T-cell acute lymphoblastic leukaemia, non-hodgkins lymphoma, B-cell lymphoma, acute myeloid leukaemia; cutaneous T cell lymphoma, premalignant and malignant skin conditions including but not limited to basal cell carcinoma (BCC), squamous cell carcinoma (SCC) or actinic keratosis (AK)).

[0013] In view of the numerous conditions that are contemplated to benefit by treatment involving modulation of the

PI3K pathway or modulation of the PI3 Kinases it is immediately apparent that new compounds that modulate PI3K pathways and use of these compounds should provide substantial therapeutic benefits to a wide variety of patients.

[0014] Provided herein are novel pyrrolotriazinone derivatives for use in the treatment of conditions in which targeting of the PI3K pathway or inhibition of PI3 Kinases can be therapeutically useful.

[0015] The compounds described in the present invention are potent PI3K inhibitors, particularly PI3Kd or dual PI3Kd/g inhibitors. This property makes them useful for the treatment or prevention of pathological conditions or diseases such as respiratory diseases (asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis), allergic diseases (allergic rhinitis), inflammatory or autoimmune diseases (rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, myasthenia gravis, acute disseminated encephalomyelitis, idiopathic thrombocytopenic purpura, Sjogren's syndrome, autoimmune hemolytic anemia, type I diabetes, psoriasis, acrodermatitis, angiodermatitis, atopic dermatitis, contact dermatitis, eczema, acne, chronic urticaria, scleroderma, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis and blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa), cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain (such as pain associated with rheumatoid arthritis or osteoarthritis, back pain, general inflammatory pain, inflammatory neuropathic pain, trigeminal neuralgia or central pain) as well as in bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (such as polycythemia vera, essential thrombocythemia or myelofibrosis); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors (such as pancreatic cancer; bladder cancer; colorectal cancer; breast cancer; prostate cancer; renal cancer; hepatocellular cancer; lung cancer; ovarian cancer; cervical cancer; gastric cancer; esophageal cancer; head and neck cancer; non-small cell lung cancer and small-cell lung cancer; melanoma; neuroendocrine cancers; central nervous system cancers; brain tumors; bone cancer; soft tissue sarcoma; chronic lymphocytic leukemia, B-cell acute lymphoblastic leukemia, T-cell acute lymphoblastic leukaemia, non-hodgkins lymphoma, B-cell lymphoma, acute myeloid leukaemia; cutaneous T cell lymphoma, premalignant and malignant skin conditions including but not limited to basal cell carcinoma (BCC), squamous cell carcinoma (SCC) or actinic keratosis (AK)).


[0016] The compounds described in the present invention are particularly useful for the treatment or prevention of pathological conditions or diseases such as neoplastic diseases (e.g. leukemia, lymphomas, solid tumors); transplant rejection, bone marrow transplant applications (e.g., graft-versus-host disease); autoimmune diseases (e.g. rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis and blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa); respiratory inflammation diseases (e.g. asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis); skin inflammatory diseases (e.g., atopic dermatitis, contact dermatitis, eczema or psoriasis); premalignant and malignant skin conditions (e.g. basal cell carcinoma (BCC), squamous cell carcinoma (SCC) or actinic keratosis (AK)); neurological disorders and pain (such as pain associated with rheumatoid arthritis or osteoarthritis, back pain, general inflammatory pain, inflammatory neuropathic pain, trigeminal neuralgia or central pain)

[0017] The compounds described in the present invention are particularly useful for the treatment or prevention of pathological conditions or diseases selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0018] In an embodiment, the compounds described in the present invention are particularly useful for the treatment or prevention of pathological conditions or diseases selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0019] It has now been found that certain pyrrolotriazinone derivatives are novel and potent PI3K inhibitors and can therefore be used in the treatment or prevention of these diseases.

[0020] Thus the present invention is directed to compounds of formula (I), or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof:



Formula (I)

X represents a nitrogen atom or a -CR₆ group;

n represents 0, 1, 2 or 3;

R_a and R_b each independently represent a hydrogen atom, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group or a linear or branched C₁-C₄ alkyl group;

R₁ represents a hydrogen atom, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a monocyclic or bicyclic C₆-C₁₄aryl group, a 5- to 14- membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N, or a 5- to 14- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N,

wherein the cycloalkyl, cycloalkenyl, aryl, heteroaryl, and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxy group, a cyano group, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₁₋₃CN group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group,

a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R₈ group, a -C(O)-(CH₂)₀₋₃-NR₇R₈ group, a -S(O)₂(CH₂)₀₋₃R₈ group, a -S(O)₂(CH₂)₀₋₃NR₇R₈ group or a -(CH₂)₀₋₃(phenyl)-OR₈ group;

R₂ and R₃ each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₇ cycloalkyl group, a -NR'R" group, or a linear or branched C₁-C₆ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₇ cycloalkyl group;

R₄ represents a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₇ cycloalkyl group, a -(CH₂)₁₋₄NR'R" group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group, a C₃-C₄ cycloalkyl group, a -C(O)-(CH₂)₀₋₃-R group or a -C(O)-(CH₂)₀₋₃-NR'R" group;

R₆ represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a linear or branched C₁-C₄ hydroxyalkyl group; a C₃-C₇ cycloalkyl group; a -(CH₂)₀₋₃NR'R" group; a -(CH₂)₁₋₃O(C₁-C₄ alkyl group); a -(CH₂)₀₋₃OC(O)-(C₁-C₄ alkyl group); a -(CH₂)₀₋₃C(O)O-(C₁-C₄ alkyl group); a -C(O)-(CH₂)₀₋₃-NR'R" group; a -(CH₂)₀₋₃C(O)OH group; a -(CH₂)₀₋₃-(5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N); a -(CH₂)₀₋₃-(5- to 14- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N); or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group;


wherein the heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxy group, a cyano group, a linear or branched C₁-C₄ alkyl group or a C₁-C₄ haloalkyl group,

R₇ and R₈ each independently represent a hydrogen atom, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group;

R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group.


R₅ represents a group selected from:

i) a group of formula (IIa)



formula (IIa)

ii) a group of formula (IIb)



formula (IIb)

and

iii) a group of formula (IIc)



formula (IIc)

wherein

Y represents a linker selected from a -NR'- group, -O- or -S-; wherein R' is as defined above;

(*) represents where R₅ is bonded to the carbon atom attached to R₄ and to the pyrrolo[2,1-b]triazine group;

W₁ represents a -CR₁₁ group and W₂ represents a nitrogen atom, or W₁ represents a nitrogen atom and W₂ represents a -CR₁₂ group;


G₁ represents a -CR₁₄ group and G₂ represents a nitrogen atom, or G₁ represents a nitrogen atom and G₂ represents a -CR₁₅ group, or G₁ represents a -CR₁₄ group and G₂ represents a -CR₁₅ group;

G₃ represents a nitrogen atom or a -CR₁₆ group;


R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom; a halogen atom; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a C₁-C₄ hydroxyalkyl group; a C₃-C₄ cycloalkyl group; a -(CH₂)₀₋₃CN group; a -C(O)-(CH₂)₁₋₃-CN group; a -C(O)-(CH₂)₀₋₃-R' group; a -C(O)-(CH₂)₀₋₃-NR'R''; a -(CH₂)₀₋₃NR'R'' group; or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' are as defined above; R₁₇ represents a group selected from

a) a group of formula (IIIa), b) a group of formula (IIIb),

(continued)




formula (IIIa)




formula (IIIb)

c) a group of formula (IIIc), and d) a group of formula (IIId),



formula (IIIc)



formula (IIId)


wherein

G₄ represents a nitrogen atom or a -CR₂₂ group;G₅ and G₆ each independently represents a nitrogen atom or a carbon atom, wherein when one of G₅ and G₆ represents a nitrogen atom the remaining represents a carbon atom;G₇ represents a -NH group or a -CH group;G₈ represents a nitrogen atom or a -CR₂₃ group;G₉ represents a nitrogen atom or a -CR₂₄ group;G₁₀ represents a nitrogen atom or a -CR₂₅ group;G₁₁ represents a nitrogen atom or a -CR₂₆ group;G₁₂ represents a nitrogen atom or a -CR₂₇ group;G₁₃ represents a nitrogen atom or a -CR₂₈ group;G₁₄ and G₁₅ each independently represents a nitrogen atom or a -carbon atom, wherein when one of G₁₄ and G₁₅ represents a nitrogen atom the remaining represents a carbon atom;G₁₆ represents a -NH group or a -CH group;G₁₇ represents a nitrogen atom or a -CR₂₉ group;G₁₈ represents a nitrogen atom or a -CR₃₀ group;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉, and R₃₀ each independently represent a hydrogen atom, a halogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' are as defined above; and wherein Y is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0021] The dotted line in the group of formula (IIIa)



formula (IIIa)

denotes that there are two double bounds in the C_5 heteroaryl ring, whose position may vary depending on which G_5 , G_6 , G_7 , G_8 or G_9 represents a nitrogen atom or a carbon atom.

[0022] The invention further provides synthetic processes and intermediates described herein, which are useful for preparing said compounds.

[0023] The invention is also directed to a compound of the invention as described herein for use in the treatment of the human or animal body by therapy.

[0024] The invention also provides a pharmaceutical composition comprising the compounds of the invention and a pharmaceutically-acceptable diluent or carrier.

[0025] The invention is also directed to the compounds of the invention as described herein, for use in the treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinases (PI3Ks), in particular wherein the pathological condition or disease is selected from respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis (RA), multiple sclerosis (MS), amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), idiopathic pulmonary fibrosis, sarcoidosis, atopic dermatitis, allergic rhinitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma (BCC), squamous cell carcinoma (SCC) and actinic keratosis (AK); preferably wherein the pathological condition or disease is selected from respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis (RA), multiple sclerosis (MS), amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, dermatomyositis, blistering diseases including but not limited to asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), idiopathic pulmonary fibrosis, sarcoidosis, atopic dermatitis, allergic rhinitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma (BCC), squamous cell carcinoma (SCC) and actinic keratosis (AK); even more in particular wherein the pathological condition or disease is leukemia, lymphomas and solid tumors, rheumatoid arthritis (RA), multiple sclerosis (MS), amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), idiopathic pulmonary fibrosis, sarcoidosis, atopic dermatitis, allergic rhinitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma (BCC), squamous cell carcinoma (SCC) and actinic keratosis (AK).

[0026] The invention is also directed to use of the compounds of the invention as described herein, in the manufacture of a medicament for treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinases (PI3Ks), in particular wherein the pathological condition or disease is as defined above.

[0027] The invention also provides a method of treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinases (PI3Ks), in particular wherein the pathological condition or disease is as defined above.

[0028] The invention also provides a combination product comprising (i) the compounds of the invention as described herein; and (ii) one or more additional active substances which are known to be useful in the treatment of respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs);

cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis (RA), multiple sclerosis (MS), amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), idiopathic pulmonary fibrosis, sarcoidosis, atopic dermatitis, allergic rhinitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma (BCC), squamous cell carcinoma (SCC) and actinic keratosis (AK); even more in particular wherein the pathological condition or disease is leukemia, lymphomas and solid tumors, rheumatoid arthritis (RA), multiple sclerosis (MS), amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF), idiopathic pulmonary fibrosis, sarcoidosis, atopic dermatitis, allergic rhinitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma (BCC), squamous cell carcinoma (SCC) and actinic keratosis (AK).

[0029] As used herein the term C₁-C₆ alkyl embraces linear or branched radicals having 1 to 6 carbon atoms, preferably 1 to 4 carbon atoms. Examples include methyl, ethyl, n-propyl, i-propyl, n-butyl, sec-butyl, t-butyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, isopentyl, 1-ethylpropyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, n-hexyl, 1-ethylbutyl, 2-ethylbutyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 2-methylpentyl, 3-methylpentyl and iso-hexyl radicals.

[0030] When it is mentioned that the alkyl radical may be optionally substituted it is meant to include linear or branched alkyl radical as defined above, which may be unsubstituted or substituted in any position by one or more substituents, for example by 1, 2 or 3 substituents. When two or more substituents are present, each substituent may be the same or different.

[0031] As used herein, the term C₁-C₄ haloalkyl group is an alkyl group, for example a C₁-C₄ or C₁-C₂ alkyl group, which is bonded to one or more, preferably 1, 2 or 3 halogen atoms. Preferably, said haloalkyl group is chosen from -CCl₃, -CHF₂ and -CF₃.

[0032] As used herein, the term C₁-C₄ hydroxyalkyl embraces linear or branched alkyl radicals having 1 to 4 carbon atoms, any one of which may be substituted by one or more, preferably 1 or 2, more preferably 1 hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, and hydroxybutyl.

[0033] As used herein, the term C₁-C₄ alkoxy (or alkyloxy) embraces linear or branched oxy-containing radicals each having alkyl portions of 1 to 4 carbon atoms.

[0034] As used herein, the term C₃-C₁₀ cycloalkyl embraces saturated monocyclic or polycyclic carbocyclic radicals having from 3 to 10 carbon atoms, preferably from 3 to 7 carbon atoms. An optionally substituted C₃-C₁₀ cycloalkyl radical is typically unsubstituted or substituted by 1, 2 or 3 substituents which may be the same or different. When a C₃-C₁₀ cycloalkyl radical carries 2 or more substituents, the substituents may be the same or different. Typically the substituents on a C₃-C₁₀ cycloalkyl group are themselves unsubstituted. Polycyclic cycloalkyl radicals contains two or more fused cycloalkyl groups, preferably two cycloalkyl groups. Typically, polycyclic cycloalkyl radicals are selected from decahydronaphthyl (decalyl), bicyclo[2.2.2]octyl, adamantly, camphyl or bornyl groups.

[0035] Examples of monocyclic cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl and cyclodecyl.

[0036] As used herein, the term C₃-C₁₀ cycloalkenyl embraces partially unsaturated carbocyclic radicals having from 3 to 10 carbon atoms, preferably from 3 to 7 carbon atoms. A C₃-C₁₀ cycloalkenyl radical is typically unsubstituted or substituted by 1, 2 or 3 substituents which may be the same or different. When a C₃-C₁₀ cycloalkenyl radical carries 2 or more substituents, the substituents may be the same or different. Typically, the substituents on a cycloalkenyl group are themselves unsubstituted.

[0037] Examples include cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclononenyl and cyclodecenyl.

[0038] As used herein, the term C₆-C₁₄ aryl radical embraces typically a C₆-C₁₄, more preferably C₆-C₁₀ monocyclic or bicyclic aryl radical such as phenyl, naphthyl, anthranyl and phenanthryl. Phenyl is preferred. A said optionally substituted C₆-C₁₄ aryl radical is typically unsubstituted or substituted by 1, 2 or 3 substituents which may be the same or different. When a C₆-C₁₄ aryl radical carries 2 or more substituents, the substituents may be the same or different. Unless otherwise specified, the substituents on a C₆-C₁₄ aryl group are typically themselves unsubstituted.

[0039] As used herein, the term 5- to 14- membered heteroaryl radical embraces typically a 5- to 14- membered ring system, preferably a 5- to 10- membered ring system, more preferably a 5- to 6- membered ring system, comprising at least one heteroatom in the ring and containing at least one heteroatom selected from O, S and N. A 5- to 14-membered heteroaryl radical may be a single ring or two fused rings wherein at least one ring contains a heteroatom.

[0040] A said optionally substituted 5- to 14- membered heteroaryl radical is typically unsubstituted or substituted by 1, 2 or 3 substituents which may be the same or different. When a 5- to 14- membered heteroaryl radical carries 2 or more substituents, the substituents may be the same or different. Unless otherwise specified, the substituents on a 5- to 14- membered heteroaryl radical are typically themselves unsubstituted.

[0041] Examples include pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, furyl, benzofuranyl, oxadiazolyl, oxazolyl, isoxazolyl, benzoxazolyl, imidazolyl, benzimidazolyl, thiazolyl, thiadiazolyl, thienyl, pyrrolyl, benzothiazolyl, indolyl, indazolyl, purinyl, quinolyl, isoquinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, quinoliziny, cinnolyl, triazolyl, indoliziny, indolyl, isoindolyl, isoindolyl, imidazolidinyl, pteridinyl, thianthrenyl, pyrazolyl, 2H-pyrazolo[3,4-d]pyrimidinyl, 1 H-pyrazolo[3,4-d]pyrimidinyl, thieno[2,3-d] pyrimidinyl and the various pyrrolopyridyl radicals.

[0042] As used herein, the term 5- to 14-membered heterocyclyl radical embraces typically a non-aromatic, saturated or unsaturated C₅-C₁₄ carbocyclic ring system, preferably C₅-C₁₀ carbocyclic ring system, more preferably C₅-C₆ carbocyclic ring system, in which one or more, for example 1, 2, 3 or 4 of the carbon atoms preferably 1 or 2 of the carbon atoms are replaced by a heteroatom selected from N, O and S. A heterocyclyl radical may be a single ring or two fused rings wherein at least one ring contains a heteroatom. When a 5 to 14-membered heterocyclyl radical carries 2 or more substituents, the substituents may be the same or different.

[0043] A said optionally substituted 5- to 14-membered heterocyclyl radical is typically unsubstituted or substituted by 1, 2 or 3 substituents which may be the same or different. Typically, the substituents on a 5 to 14-membered heterocyclyl radical are themselves unsubstituted.

[0044] Examples of 5- to 14-membered heterocyclyl radicals include piperidyl, pyrrolidyl, pyrrolinyl, piperazinyl, morpholinyl, thiomorpholinyl, pyrrolyl, pyrazolinyl, pirazolidinyl, quinuclidinyl, triazolyl, pyrazolyl, tetrazolyl, imidazolidinyl, imidazolyl, oxiranyl, thiaranyl, aziridinyl, oxetanyl, thiatanyl, azetidyl, 4,5-dihydro-oxazolyl, 2-benzofuran-1(3H)-one, 1,3-dioxol-2-one, tetrahydrofuran, 3-aza-tetrahydrofuran, tetrahydrothiophenyl, tetrahydropyran, tetrahydrothiopyran, 1,4-azathianyl, oxepanyl, thiephanyl, azepanyl, 1,4-dioxepanyl, 1,4-oxathiepanyl, 1,4-oxaazepanyl, 1,4-dithiepanyl, 1,4-thiezepanyl, 1,4-diazepanyl, tropanyl, (1S,5R)-3-aza-bicyclo[3.1.0]hexyl, 3,4-dihydro-2H-pyran, 5,6-dihydro-2H-pyran, 2H-pyran, 2,3-hydrobenzofuran, 1,2,3,4-tetrahydropyridyl, 1,2,5,6-tetrahydropyridyl, isoindolyl and indolyl.

[0045] Where a 5- to 14-membered heterocyclyl radical carries 2 or more substituents, the substituents may be the same or different.

[0046] As used herein, the bicyclic N-containing heteroaryl group is a C₈-C₁₀ membered ring system where two rings have been fused and wherein at least in one ring one of the carbon atoms is replaced by N and optionally in which 1, 2, 3, or 4, preferably 1, 2, or 3 further carbon atoms of any ring which form the group are replaced by N.

[0047] Examples include indolyl, benzimidazolyl, indazolyl, benzotriazolyl, pyrrolo[2,3-b]pyridinyl, pyrrolo[2,3-c]pyridinyl, pyrrolo[3,2-c]pyridinyl, pyrrolo[3,2-b]pyridinyl, imidazo[4,5-b]pyridinyl, imidazo[4,5-c]pyridinyl, pyrazolo[4,3-d]pyridinyl, pyrazolo[4,3-d]pyridinyl, pyrazolo[3,4-c]pyridinyl, pyrazolo[3,4-b]pyridinyl, isoindolyl, indazolyl, purinyl, indolyl, imidazo[1,2-a]pyridinyl, imidazo[1,5-a]pyridinyl, pyrazolo[1,5-a]pyridinyl, pyrrolo[1,2-b]pyridazinyl, imidazo[1,2-c]pyrimidinyl, quinolyl, isoquinolyl, cinnolyl, azaquinazolinyl, quinoxalinyl, phthalazinyl, naphthyridinyl, pyrido[3,2-d]pyrimidinyl, pyrido[4,3-d]pyrimidinyl, pyrido[3,4-d]pyrimidinyl, pyrido[2,3-d]pyrimidinyl, pyrazolo[1,5-a]pyrimidinyl, pyrido[2,3-b]pyrazinyl, pyrido[3,4-b]pyrazinyl, pyrimido[5,4-d]pyrimidinyl, pyrazino[2,3-b]pyrazinyl and pyrimido[4,5-d]pyrimidinyl.

[0048] As used herein, some of the atoms, radicals, moieties, chains and cycles present in the general structures of the invention are "optionally substituted". This means that these atoms, radicals, moieties, chains and cycles can be either unsubstituted or substituted in any position by one or more, for example 1, 2, 3 or 4, substituents, whereby the hydrogen atoms bound to the unsubstituted atoms, radicals, moieties, chains and cycles are replaced by chemically acceptable atoms, radicals, moieties, chains and cycles. When two or more substituents are present, each substituent may be the same or different. The substituents are typically themselves unsubstituted.

[0049] As used herein, the term halogen atom embraces chlorine, fluorine, bromine and iodine atoms. A halogen atom is typically a fluorine, chlorine or bromine atom, most preferably chlorine or fluorine. The term halo when used as a prefix has the same meaning. Compounds containing one or more chiral centre may be used in enantiomerically or diastereoisomerically pure form, in the form of racemic mixtures and in the form of mixtures enriched in one or more stereoisomer. The scope of the invention as described and claimed encompasses the racemic forms of the compounds as well as the individual enantiomers, diastereomers, and stereoisomer-enriched mixtures.

[0050] Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate using, for example, chiral high pressure liquid chromatography (HPLC). Alternatively, the racemate (or a racemic precursor) may be reacted with a suitable optically active compound, for example, an alcohol, or, in the case where the compound contains an acidic or basic moiety, an acid or base such as tartaric acid or 1-phenylethylamine. The resulting diastereomeric mixture may be separated by chromatography and/or fractional crystallization and one or both of the diastereoisomers converted to the corresponding pure enantiomer(s) by means well known to one skilled in the art. Chiral compounds of the invention (and chiral precursors thereof) may be obtained in enantiomerically-enriched form using chromatography, typically HPLC, on an asymmetric resin with a mobile phase consisting of a hydrocarbon, typically heptane or hexane, containing from 0 to 50% isopropanol, typically from 2 to 20%, and from 0 to 5% of an alkylamine, typically 0.1 % diethylamine. Concentration of the eluate affords the enriched mixture. Stereoisomer conglomerates may be separated by conventional techniques known to those skilled in the art. See, e.g. "Stereochemistry of Organic Compounds" by Ernest L. Eliel (Wiley, New York, 1994).

[0051] Atropisomers are stereoisomers resulting from hindered rotation about single bonds where the steric strain barrier to rotation is high enough to allow for the isolation of the conformers. Oki (Oki, M; Topics in Stereochemistry 1983, 1) defined atropisomers as conformers that interconvert with a half-life of more than 1000 seconds at a given temperature. The scope of the invention as described and claimed encompasses the racemic forms of the compounds as well as the individual atropisomers (an atropisomer "substantially free" of its corresponding enantiomer) and stereoisomer-enriched mixtures, i.e. mixtures of atropisomers.

[0052] Separation of atropisomers is possibly by chiral resolution methods such as selective crystallization. In an atropo-enantioselective or atroposelective synthesis one atropisomer is formed at the expense of the other. Atroposelective synthesis may be carried out by use of chiral auxiliaries like a Corey-Bakshi-Shibata (CBS) catalyst (asymmetric catalyst derived from proline) in the total synthesis of knipholone or by approaches based on thermodynamic equilibration when an isomerization reaction favors one atropisomer over the other.

[0053] As used herein, the term pharmaceutically acceptable salt refers to a salt prepared from a base or acid which is acceptable for administration to a patient, such as a mammal. Such salts can be derived from pharmaceutically-acceptable inorganic or organic bases and from pharmaceutically-acceptable inorganic or organic acids.

[0054] Pharmaceutically acceptable acids include both inorganic acids, for example hydrochloric, sulphuric, phosphoric, diphosphoric, hydrobromic, hydroiodic and nitric acid; and organic acids, for example citric, fumaric, gluconic, glutamic, lactic, maleic, malic, mandelic, mucic, ascorbic, oxalic, pantothenic, succinic, tartaric, benzoic, acetic, methanesulphonic, ethanesulphonic, benzenesulphonic, p-toluenesulphonic acid, xinafoic (1-hydroxy-2-naphthoic acid), napadisilic (1,5-naphthalenedisulfonic acid) and the like. Particularly preferred are salts derived from fumaric, hydrobromic, hydrochloric, acetic, sulfuric, methanesulfonic, xinafoic, and tartaric acids.

[0055] Salts derived from pharmaceutically-acceptable inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, zinc and the like. Particularly preferred are ammonium, calcium, magnesium, potassium and sodium salts.

[0056] Salts derived from pharmaceutically-acceptable organic bases include salts of primary, secondary and tertiary amines, including alkyl amines, arylalkyl amines, heterocyclic amines, cyclic amines, naturally-occurring amines and the like, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

[0057] Other preferred salts according to the invention are quaternary ammonium compounds wherein an equivalent of an anion (X^-) is associated with the positive charge of the N atom. X^- may be an anion of various mineral acids such as, for example, chloride, bromide, iodide, sulphate, nitrate, phosphate, or an anion of an organic acid such as, for example, acetate, maleate, fumarate, citrate, oxalate, succinate, tartrate, malate, mandelate, trifluoroacetate, methanesulphonate and p-toluenesulphonate. X^- is preferably an anion selected from chloride, bromide, iodide, sulphate, nitrate, acetate, maleate, oxalate, succinate or trifluoroacetate. More preferably X^- is chloride, bromide, trifluoroacetate or methanesulphonate.

[0058] As used herein, an N-oxide is formed from the tertiary basic amines or imines present in the molecule, using a convenient oxidising agent.

[0059] The present invention also embraces tautomeric forms of the compounds of formula (I), or pharmaceutically acceptable salts, solvates, N-oxides, stereoisomers or deuterated derivatives thereof.

[0060] The compounds of the invention may exist in both unsolvated and solvated forms. The term solvate is used herein to describe a molecular complex comprising a compound of the invention and an amount of one or more pharmaceutically acceptable solvent molecules. The term hydrate is employed when said solvent is water. Examples of solvate forms include, but are not limited to, compounds of the invention in association with water, acetone, dichloromethane, 2-propanol, ethanol, methanol, dimethylsulfoxide (DMSO), ethyl acetate, acetic acid, ethanolamine, or mixtures thereof. It is specifically contemplated that in the present invention one solvent molecule can be associated with one molecule of the compounds of the present invention, such as a hydrate.

[0061] Furthermore, it is specifically contemplated that in the present invention, more than one solvent molecule may be associated with one molecule of the compounds of the present invention, such as a dihydrate. Additionally, it is specifically contemplated that in the present invention less than one solvent molecule may be associated with one molecule of the compounds of the present invention, such as a hemihydrate. Furthermore, solvates of the present invention are contemplated as solvates of compounds of the present invention that retain the biological effectiveness of the non-solvate form of the compounds.

[0062] The invention also includes isotopically-labeled compounds of the invention, wherein one or more atoms is replaced by an atom having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes suitable for inclusion in the compounds of the invention include isotopes of hydrogen, such as ^2H and ^3H , carbon, such as ^{11}C , ^{13}C and ^{14}C , chlorine, such as ^{36}Cl ,

fluorine, such as ^{18}F , iodine, such as ^{123}I and ^{125}I , nitrogen, such as ^{13}N and ^{15}N , oxygen, such as ^{15}O , ^{17}O and ^{18}O , phosphorus, such as ^{32}P , and sulfur, such as ^{35}S . Certain isotopically-labeled compounds of the invention, for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, ^3H , and carbon-14, ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection. Substitution with heavier isotopes such as deuterium, ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life or reduced dosage requirements, and hence may be preferred in some circumstances. Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy.

[0063] Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed.

[0064] Preferred isotopically-labeled compounds include deuterated derivatives of the compounds of the invention. As used herein, the term deuterated derivative embraces compounds of the invention where in a particular position at least one hydrogen atom is replaced by deuterium. Deuterium (D or ^2H) is a stable isotope of hydrogen which is present at a natural abundance of 0.015 molar %.

[0065] Hydrogen deuterium exchange (deuterium incorporation) is a chemical reaction in which a covalently bonded hydrogen atom is replaced by a deuterium atom. Said exchange (incorporation) reaction can be total or partial.

[0066] Typically, a deuterated derivative of a compound of the invention has an isotopic enrichment factor (ratio between the isotopic abundance and the natural abundance of that isotope, i.e. the percentage of incorporation of deuterium at a given position in a molecule in the place of hydrogen) for each deuterium present at a site designated as a potential site of deuteration on the compound of at least 3500 (52.5% deuterium incorporation).

[0067] In a preferred embodiment, the isotopic enrichment factor is at least 5000 (75% deuterium). In a more preferred embodiment, the isotopic enrichment factor is at least 6333.3 (95% deuterium incorporation). In a most preferred embodiment, the isotopic enrichment factor is at least 6633.3 (99.5% deuterium incorporation). It is understood that the isotopic enrichment factor of each deuterium present at a site designated as a site of deuteration is independent from the other deuteration sites.

[0068] The isotopic enrichment factor can be determined using conventional analytical methods known to an ordinary skilled in the art, including mass spectrometry (MS) and nuclear magnetic resonance (NMR).

[0069] Prodrugs of the compounds described herein are also within the scope of the invention. Thus certain derivatives of the compounds of the present invention, which derivatives may have little or no pharmacological activity themselves, when administered into or onto the body may be converted into compounds of the present invention having the desired activity, for example, by hydrolytic cleavage. Such derivatives are referred to as 'prodrugs'. Further information on the use of prodrugs may be found in Pro-drugs as Novel Delivery Systems, Vol. 14, ACS Symposium Series (T. Higuchi and W. Stella) and Bioreversible Carriers in Drug Design, Pergamon Press, 1987 (ed. E. B. Roche, American Pharmaceutical Association).

[0070] Prodrugs in accordance with the invention can, for example, be produced by replacing appropriate functionalities present in the compounds of the present invention with certain moieties known to those skilled in the art as 'pro-moieties' as described, for example, in Design of Prodrugs by H. Bundgaard (Elsevier, 1985).

[0071] In the case of compounds that are solids, it is understood by those skilled in the art that the inventive compounds and salts may exist in different crystalline or polymorphic forms, or in an amorphous form, all of which are intended to be within the scope of the present invention.

[0072] As used herein, the term PI3Kd inhibitor generally refers to a compound that inhibits the activity of the PI3Kd isoform more effectively than other isoforms of the PI3K family.

[0073] As used herein, the term PI3Kd/g inhibitor generally refers to a compound that inhibits the activity of both the PI3Kd isoform and the PI3Kg isoform more effectively than other isoforms of the PI3K family.

[0074] The relative efficacies of compounds as inhibitors of an enzyme activity (or other biological activity) can be established by determining the concentrations at which each compound inhibits the activity to a predefined extent and then comparing the results. Typically, the preferred determination is the concentration that inhibits 50% of the activity in a biochemical assay, i.e., the 50% inhibitory concentration or " IC_{50} ." IC_{50} determinations can be accomplished using conventional techniques known in the art. In general, an IC_{50} can be determined by measuring the activity of a given enzyme in the presence of a range of concentrations of the inhibitor under study. The experimentally obtained values of enzyme activity then are plotted against the inhibitor concentrations used. The concentration of the inhibitor that shows 50% enzyme activity (as compared to the activity in the absence of any inhibitor) is taken as the IC_{50} value.

[0075] Accordingly, a PI3Kd inhibitor alternatively can be understood to refer to a compound that exhibits a 50% inhibitory concentration (IC_{50}) with respect to PI3Kd that is at least of less than about 100 μM , preferably of less than about 50 μM , more preferably of less than about 20 μM , even more preferably of less than about 10 μM PI3K HTRF assay (as described in Gray et al. Anal Biochem, 2003; 313: 234-45)

[0076] Typically, in the compound of formula (I), X represents a nitrogen atom or a -CR₆ group.

[0077] Typically, in the compound of formula (I), R_a and R_b each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group.

[0078] Preferably, R_a and R_b each independently represent a hydrogen atom, a methyl group or an ethyl group.

[0079] Typically, n represents 0, 1 or 2, preferably 0 or 1, more preferably 0.

[0080] Typically, in the compound of formula (I) R₁ represents a hydrogen atom, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a monocyclic or bicyclic C₆-C₁₀ aryl group, a 5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N, or a 5- to 14-membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N; wherein the cycloalkyl, cycloalkenyl, aryl, heteroaryl, and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxy group, a cyano group, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₁₋₃CN group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃R₈ group, a -C(O)-(CH₂)₀₋₃NR₇R₈ group, a -S(O)₂(CH₂)₀₋₃R₈ group, a -S(O)₂(CH₂)₀₋₃NR₇R₈ group or a -(CH₂)₀₋₃(phenyl)-OR₈ group; wherein R₇ and R₈ are as defined above.

[0081] Preferably, R₁ represents a hydrogen atom, C₁-C₃ alkyl group, a C₁-C₃ haloalkyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a naphthyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, or a 5- to 10-membered monocyclic or bicyclic heterocyclyl group containing one, two or three heteroatoms selected from O, S and N, wherein the cycloalkyl, phenyl, naphthyl, heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxy group, a cyano group, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group, a -C(O)-(CH₂)₀₋₃R₈ group, a -C(O)-(CH₂)₀₋₃NR₇R₈ group or a -(CH₂)₀₋₃(phenyl)-OR₈ group; wherein R₇ and R₈ are as defined above.

[0082] More preferably R₁ represents a phenyl group or a pyridinyl group, which phenyl or pyridinyl is unsubstituted or substituted by one, two or three substituents selected from a halogen atom, a hydroxyl group, a linear or branched C₁-C₃ alkyl group, a C₁-C₃ haloalkyl group, or a -(CH₂)₀₋₃OCH₃ group; for example from a halogen atom, a linear or branched C₁-C₃ alkyl group or a -(CH₂)₀₋₃OCH₃ group.

[0083] Preferably, when R₁ represents a phenyl group or a pyridinyl group, said phenyl and pyridinyl groups are directly bonded to the pyrrolotriazinone group. In other words, the linker -(R_a-C-R_b)_n- is not present. More preferably, R₁ represents a phenyl group.

[0084] Preferably, when R₁ is a phenyl group, it is unsubstituted or substituted by one, two or three substituents selected from a halogen atom (preferably a fluorine atom or a chlorine atom), a hydroxyl group, a linear or branched C₁-C₃ alkyl group (preferably a methyl group), a C₁-C₃ haloalkyl group, or a -OCH₃ group; for example when R₁ is a phenyl group, it is unsubstituted or substituted by one, two or three substituents selected from a halogen atom (preferably a fluorine atom or a chlorine atom), a linear or branched C₁-C₃ alkyl group (preferably a methyl group) or a -OCH₃ group. Preferably, when R₁ is a pyridinyl or piperidinyl group, said groups are linked to the rest of the molecule via a ring carbon atom, in other words they are linked to the pyrrolotriazinone group via a ring carbon atom. Substituents on a pyridinyl group may be present on any ring atom but are preferably present on a carbon atom. Substituents on a piperidinyl group may be present on any ring atom but are preferably present on the nitrogen atom.

[0085] In one embodiment, in the compound of formula (I) R₁ represents a C₁-C₃ alkyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a naphthyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, or a 5- to 10-membered monocyclic or bicyclic heterocyclyl group containing one, two or three heteroatoms selected from O, S and N; wherein the cycloalkyl, phenyl, naphthyl, heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group, a -C(O)-(CH₂)₀₋₃R₈ group or a -C(O)-(CH₂)₀₋₃NR₇R₈ group; wherein R₇ and R₈ are as defined above.

[0086] In this embodiment R₁ preferably represents a C₃-C₇ cycloalkyl group, a phenyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, a tetrahydropyranyl group or a morpholinyl group; wherein the cycloalkyl, phenyl, heteroaryl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydropyranyl or morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group, a -C(O)-(CH₂)₀₋₃R₈ group or a -C(O)-(CH₂)₀₋₃NR₇R₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a C₁-C₄ alkyl group.

[0087] Typically, in the compound of formula (I) R₂ represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group,

a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_3 alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1-C_3 alkyl group.

[0088] Preferably R_2 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_1-C_3 alkoxy group, a C_1-C_3 haloalkyl group, a C_3-C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_2 alkoxy group.

[0089] More preferably R_2 represents a hydrogen atom, a halogen atom, $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_3 alkyl group. Most preferably R_2 represents a hydrogen atom or a methyl group.

[0090] Typically, in the compound of formula (I) R_3 represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1-C_4 alkoxy group, a C_1-C_4 haloalkyl group, a C_1-C_4 hydroxyalkyl group, a C_3-C_4 cycloalkyl group, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_3 alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1-C_3 alkyl group.

[0091] Preferably, in the compound of formula (I) R_3 represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1-C_3 alkoxy group, a C_1-C_3 haloalkyl group, a C_3-C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_2 alkoxy group.

[0092] More preferably R_3 represents a hydrogen atom, a halogen atom, a cyano group, a C_1-C_3 alkoxy group, a C_1-C_3 haloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_3 alkyl group;

[0093] In one embodiment, in the compound of formula (I) R_3 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_1-C_3 alkoxy group, a C_1-C_3 haloalkyl group, a C_3-C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_2 alkoxy group. In this embodiment, more preferably R_3 represents a hydrogen atom, a halogen atom, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_3 alkyl group. Most preferably R_3 represents a hydrogen atom or a methyl group.

[0094] Typically, in the compound of formula (I), R_4 represents a hydrogen atom, a C_1-C_4 haloalkyl group, a C_1-C_4 hydroxyalkyl group, a C_3-C_4 cycloalkyl group, a $-(CH_2)_{1-4}NR'R''$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_3 alkoxy group, a $-C(O)-(CH_2)_{0-3}-R'$ group or a $-C(O)-(CH_2)_{0-3}-NR'R''$ group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1-C_3 alkyl group. More preferably, R_4 represents a hydrogen atom, a C_1-C_3 haloalkyl group, a C_1-C_3 hydroxyalkyl group, a C_3-C_4 cycloalkyl group, or a linear or branched C_1-C_3 alkyl group.

[0095] In one embodiment, in the compound of formula (I), R_4 represents a hydrogen atom, a C_1-C_4 alkoxy group, a C_1-C_4 haloalkyl group, a C_1-C_4 hydroxyalkyl group, a C_3-C_4 cycloalkyl group, a $-(CH_2)_{1-4}NR'R''$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_3 alkoxy group, a $-C(O)-(CH_2)_{0-3}-R'$ group or a $-C(O)-(CH_2)_{0-3}-NR'R''$ group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1-C_3 alkyl group. In this embodiment, more preferably, R_4 represents a hydrogen atom, a C_1-C_3 alkoxy group, a C_1-C_3 haloalkyl group, a C_3-C_4 cycloalkyl group, or a linear or branched C_1-C_3 alkyl group. In this embodiment, most preferably R_4 represents a hydrogen atom, a C_1-C_3 haloalkyl group or a linear or branched C_1-C_3 alkyl group.

[0096] Typically, in the compound of formula (I), R_6 represents a hydrogen atom; a halogen atom; a hydroxy group; a cyano group; a C_1-C_4 alkoxy group; a C_1-C_4 haloalkyl group; a linear or branched C_1-C_4 hydroxyalkyl group; a C_3-C_7 cycloalkyl group; a $-(CH_2)_{0-3}NR'R''$ group; a $-(CH_2)_{1-3}O(C_1-C_4 \text{ alkyl group})$; a $-(CH_2)_{0-3}OC(O)-(C_1-C_4 \text{ alkyl group})$; a $-(CH_2)_{0-3}C(O)O-(C_1-C_4 \text{ alkyl group})$; a $-C(O)-(CH_2)_{0-3}-NR'R''$ group; a $-(CH_2)_{0-3}C(O)OH$ group; a $-(CH_2)_{0-3}-(5- \text{ to } 10\text{-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N})$; a $-(CH_2)_{0-3}-(5- \text{ to } 10\text{-membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N})$;

or a linear or branched C_1-C_3 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1-C_4 alkoxy group, a cyano group or a C_3-C_4 cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1-C_3 alkyl group; and

wherein the heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1-C_3 alkyl group or a C_1-C_3 haloalkyl group. Preferably, in the compound of formula (I), R_6 represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1-C_4 alkoxy group, a C_1-C_4 haloalkyl group, a C_1-C_4 hydroxyalkyl group, a C_3-C_4 cycloalkyl group, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1-C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1-C_3 alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1-C_3 alkyl group.


[0097] More preferably, R_6 represents a hydrogen atom, a halogen atom, a C_1-C_3 alkoxy group, a C_1-C_3 haloalkyl group, a C_3-C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1-C_4 alkyl

group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group. Even more preferably, R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group (preferably a -CHF₂ group or a -CF₃ group), or a linear or branched C₁-C₃ alkyl group.


[0098] In one embodiment, in the compound of formula (I), R₆ represents a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₃ alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group. In this embodiment, preferably, R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group. In this embodiment, more preferably, R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group (preferably a -CHF₂ group or a -CF₃ group), or a linear or branched C₁-C₃ alkyl group.

[0099] In a particular embodiment, R₅ represents a group selected from

i) a group of formula (IIa-1), and ii) a group of formula (IIa-2)



formula (IIa-1)



formula (IIa-2)

wherein


R₉, R₁₀, R₁₁ and R₁₂ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group.

[0100] In this particular embodiment, preferably R₅ represents a group of formula (IIa-2) wherein R₉, R₁₀, and R₁₂ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group,

a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group. More preferably, R₉ and R₁₂ each independently represent a hydrogen atom and R₁₀ represents a -(CH₂)₀₋₃NR'R'' group wherein R' and R'' each independently represent a hydrogen atom or methyl group. Even more preferably, R₉ and R₁₂ each independently represent a hydrogen atom and R₁₀ represents a -NH₂ group.


[0101] In another particular embodiment, R₅ represents a group selected from

i) a group of formula (IIb-1),




formula (IIb-1)

ii) a group of formula (IIb-2),



formula (IIb-2)


iii) a group of formula (IIb-3),




formula (IIb-3)

iv) a group of formula (IIb-4), and v) a group of formula (IIb-5),

(continued)



formula (IIb-4)



formula (IIb-5)

wherein

R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a halogen atom; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a C₁-C₄ hydroxyalkyl group; a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group; a -C(O)-(CH₂)₀₋₃R' group; a -C(O)-(CH₂)₀₋₃NR'R''; a -(CH₂)₀₋₃NR'R'' group; or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group;

wherein R' and R'' each independently represent a hydrogen atom, a hydroxy group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a linker selected from a -NR'- group, -O- or -S-; wherein R' is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0102] In this particular embodiment, preferably R₅ represents a group selected from a group of formula (IIb-1), group of formula (IIb-2) and a group of formula (IIb-3) wherein R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a halogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃NR'R'' or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a -NR'- group, wherein R' is as defined before.

[0103] Preferably, when R₅ is a group of formula (IIb-1) R₁₄ and R₁₆ each independently represent a hydrogen atom, and R₁₃ and R₁₅ each independently represent a hydrogen atom, a -C(O)-(CH₂)₀₋₃NR'R'' or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group, wherein R' is as defined before. Even more preferably, R₁₄ and R₁₆ each independently represent a hydrogen atom, and R₁₃ and R₁₅ each independently represent a hydrogen atom, a -C(O)NR'R'' group or a -NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom or a methyl group; wherein Y represents a -NH- group.

[0104] Preferably, when R₅ is a group of formula (IIb-2) R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a halogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃NR'R'' or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxy group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group, wherein R' is as defined before. Even more preferably, R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a halogen atom, a -CN group or a -NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom or a methyl group; and wherein Y represents a -NH- group. Still more preferably, R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a -CN group or a -NH₂ group.

[0105] In a particular embodiment, when R₅ is a group of formula (IIb-2) R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a halogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃NR'R'' or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group, wherein R' is as defined before.

[0106] Even more preferably, when R₅ is a group of formula (IIb-2) R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a halogen atom, a -CN group or a -NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom or a methyl group; and wherein Y represents a -NH- group. Still more preferably, when R₅ is a group of formula (IIb-2) R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a -CN group or a -NH₂ group.

[0107] Preferably, when R₅ is a group of formula (IIb-3) R₁₃ and R₁₅ each independently represent a hydrogen atom, a -C(O)-(CH₂)₀₋₃NR'R'' or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group, wherein R' is as defined before. Even more preferably R₁₃ and R₁₅ each independently represent a hydrogen atom, a -C(O)NR'R'' group or a

-NR'R" group; wherein R' and R" each independently represent a hydrogen atom or a methyl group; wherein Y represents a -NH- group. Still more preferably, R₁₃ and R₁₅ each independently represent a hydrogen atom, a -CN group or a -NH₂ group.

[0108] In a particular embodiment, when R₅ represents a group selected from a group of formula (IIb-1), a group of formula (IIb-2), a group of formula (IIb-3), a group of formula (IIb-4), and a group of formula (IIb-3) as described above, R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃R' group, a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxy group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a linker selected from a -NR'-group, -O- or -S-; wherein R' is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0109] In this particular embodiment, preferably R₅ represents a group selected from a group of formula (IIb-1) and a group of formula (IIb-2) wherein


R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom,

a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R" or a -(CH₂)₀₋₃NR'R" group; wherein R' and R" each independently represent a hydrogen atom or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a -NR'- group, wherein R' is as defined before.

[0110] In this particular embodiment, preferably, when R₅ is a group of formula (IIb-1) R₁₄ and R₁₆ each independently represent a hydrogen atom, and R₁₃ and R₁₅ each independently represent a hydrogen atom, a -C(O)-(CH₂)₀₋₃-NR'R" or a -(CH₂)₀₋₃NR'R" group; wherein R' and R" each independently represent a hydrogen atom, a hydroxy group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group, wherein R' is as defined before. Even more preferably, R₁₄ and R₁₆ each independently represent a hydrogen atom, and R₁₃ and R₁₅ each independently represent a hydrogen atom, a -C(O)-NR'R' group or a -NR'R" group; wherein R' and R" each independently represent a hydrogen atom or a methyl group; wherein Y represents a -NH- group. In this particular embodiment, preferably, when R₅ is a group of formula (IIb-2) R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R" or a -(CH₂)₀₋₃NR'R" group; wherein R' and R" each independently represent a hydrogen atom, a hydroxy group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group, wherein R' is as defined before. Even more preferably, R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a -CN group or a -NR'R" group; wherein R' and R" each independently represent a hydrogen atom or a methyl group; and wherein Y represents a -NH- group. Still more preferably, R₁₃ represents a hydrogen atom, and R₁₄ and R₁₅ each independently represent a hydrogen atom, a -CN group or a -NH₂ group.


[0111] In a further particular embodiment, R₅ represents a group selected from

i) a group of formula (IIIa-1),




formula (IIIa-1)

ii) a group of formula (IIIa-2),




formula (IIIa-2)

iii) a group of formula (IIIa-3),




formula (IIIa-3)

iv) a group of formula (IIIa-4),




formula (IIIa-4)

v) a group of formula (IIIa-5),



formula (IIIa-5)

vi) a group of formula (IIIa-6),




formula (IIIa-6)

vii) a group of formula (IIIa-7),


viii) a group of formula (IIIa-8),

ix) a group of formula (IIIa-9),


(continued)



formula (IIIa-7)




formula (IIIa-8)




formula (IIIa-9)

X) a group of formula (IIIa-10),




formula (IIIa-10)

xi) a group of formula (IIIa-11),




formula (IIIa-11)

xii) a group of formula (IIIa-12),




formula (IIIa-12)

xiii) a group of formula (IIIa-13),




formula (IIIa-13)

xiv) a group of formula (IIIa-14),




formula (IIIa-14)

xv) a group of formula (IIIa-15),




formula (IIIa-15)

xvi) a group of formula (IIIa-16),




formula (IIIa-16)

xvii) a group of formula (IIIa-17),




formula (IIIa-17)

xviii) a group of formula (IIIa-18),




formula (IIIa-18)

xix) a group of formula (IIIa-19),




formula (IIIa-19)

xx) a group of formula (IIIa-20),




formula (IIIa-20)

xxi) a group of formula (IIIa-21), and



formula (IIIa-21)

xxii) a group of formula (IIIa-22)



formula (IIIa-22)

wherein

R₁₈, R₂₂, R₂₃, and R₂₄ each independently represent a hydrogen atom, a halogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a linker selected from a -NR'- group, -O- or -S-; wherein R' is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0112] In this particular embodiment, preferably R₅ represents a group selected from a group of formula (IIIa-1), a group of formula (IIIa-3), a group of formula (IIIa-5) and a group of formula (IIIa-14) wherein

R₁₈, R₂₂, R₂₃ and R₂₄ each independently represent a hydrogen atom, a halogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined above;

and wherein more preferably, R₁₈, R₂₂, R₂₃ and R₂₄ each independently represent a hydrogen atom, a halogen atom, a -CN group or a -NH₂ group;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0113] In a particular embodiment, preferably R₅ represents a group of formula (IIIa-1), wherein

R₁₈ and R₂₃ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group,

a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0114] In a particular embodiment, when R₅ is a group of formula (IIIa-1) R₁₈ and R₂₃ each independently represent a hydrogen atom, a -C(O)-(CH₂)₀₋₃-NR'R" or a -(CH₂)₀₋₃NR'R" group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; wherein Y represents a -NR'- group or -S-, wherein R' is as defined before. Even more preferably, R₁₈ and R₂₃ each independently represent a hydrogen atom or a -NR'R" group; wherein R' and R" each independently represent a hydrogen atom or a methyl group; wherein Y represents a -NR'- group or -S-, wherein R' is as defined before.

[0115] Preferably, when R₅ is a group of formula (IIIa-1) wherein Y represents a -NR'- group, wherein R' is a linear or branched C₁-C₄ alkyl group; R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group. More preferably, R₄ together with the -NR'- group of R₅ and the carbon atom to which both R₄ and the -NR'- group are bonded form an azetidiny group, a pyrrolidiny group, a piperidiny group or a piperaziny group; even more preferably a pyrrolidiny group or a piperidiny group.

[0116] In a particular embodiment, R₅ represents a group of formula (IIIa-1), wherein

R₁₈ and R₂₃ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group,

a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0117] In a particular embodiment, when R₅ represents a group selected from a group of formula (IIIa-1), a group of

formula (IIIa-2), a group of formula (IIIa-3), a group of formula (IIIa-4), a group of formula (IIIa-5), a group of formula (IIIa-6), a group of formula (IIIa-7), a group of formula (IIIa-8), a group of formula (IIIa-9), a group of formula (IIIa-10), a group of formula (IIIa-11), a group of formula (IIIa-12), a group of formula (IIIa-13), a group of formula (IIIa-14), a group of formula (IIIa-15), a group of formula (IIIa-16), a group of formula (IIIa-17), a group of formula (IIIa-18), a group of formula (IIIa-19), a group of formula (IIIa-20), a group of formula (IIIa-21), and a group of formula (IIIa-22), as described above,

R_{18} , R_{22} , R_{23} , and R_{24} each independently represent a hydrogen atom, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{0-3}CN$ group, a $-C(O)-(CH_2)_{1-3}CN$ group, a $-C(O)-(CH_2)_{0-3}R'$ group, a $-C(O)-(CH_2)_{0-3}NR'R''$, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxy group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group; and wherein Y represents a linker selected from a $-NR'$ -group, $-O$ - or $-S$ -; wherein R' is as defined above;

or in the case that Y represents a $-NR'$ - group, R_4 together with the $-NR'$ - group and the carbon atom to which both R_4 and the $-NR'$ - group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a $-CHF_2$ group or a $-CF_3$ group.

[0118] In this particular embodiment, preferably R_5 represents a group of formula (IIIa-1) wherein

R_{18} and R_{23} each independently represent a hydrogen atom, a $-(CH_2)_{0-3}CN$ group, a $-C(O)-(CH_2)_{1-3}CN$ group, a $-C(O)-(CH_2)_{0-3}R'$ group,

a $-C(O)-(CH_2)_{0-3}NR'R''$, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxy group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group; and wherein Y is as defined above;

or in the case that Y represents a $-NR'$ - group, R_4 together with the $-NR'$ - group and the carbon atom to which both R_4 and the $-NR'$ - group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a $-CHF_2$ group or a $-CF_3$ group.

[0119] In this embodiment, preferably, when R_5 is a group of formula (IIIa-1) R_{18} and R_{23} each independently represent a hydrogen atom, a $-C(O)-(CH_2)_{0-3}NR'R''$ or a $-(CH_2)_{0-3}NR'R''$ group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxy group, or a linear or branched C_1 - C_3 alkyl group; wherein Y represents a $-NR'$ - group or $-S$ -, wherein R' is as defined before. Even more preferably, R_{18} and R_{23} each independently represent a hydrogen atom or a $-NR'R''$ group; wherein R' and R'' each independently represent a hydrogen atom or a methyl group; wherein Y represents a $-NR'$ - group or $-S$ -, wherein R' is as defined before.


[0120] In this embodiment, preferably, when R_5 is a group of formula (IIIa-1) wherein Y represents a $-NR'$ - group, wherein R' is a linear or branched C_1 - C_4 alkyl group; R_4 together with the $-NR'$ - group and the carbon atom to which both R_4 and the $-NR'$ -group are bonded form a 4- to 7- membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a $-CHF_2$ group or a $-CF_3$ group. More preferably, R_4 together with the $-NR'$ - group of R_5 and the carbon atom to which both R_4 and the $-NR'$ - group are bonded form an azetidiny group, a pyrrolidiny group, a piperidiny group or a piperazinyl group; even more preferably a pyrrolidiny group or a piperidiny group.

[0121] In another particular embodiment, R_5 represents a group selected from


i) a group of formula (IIIb-1),

ii) a group of formula (IIIb-2),


iii) a group of formula (IIIb-3),



formula (IIIb-1)



formula (IIIb-2)




formula (IIIb-3)

iv) a group of formula (IIIb-4),


v) a group of formula (IIIb-5),

vi) a group of formula (IIIb-6),


(continued)



formula (IIIb-4)




formula (IIIb-5)




formula (IIIb-6)

vii) a group of formula (IIIb-7), and viii) a group of formula (IIIb-8),



formula (IIIb-7)



formula (IIIb-8)


wherein

R₁₈, R₂₂, R₂₅, R₂₆, and R₂₇ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a linker selected from a -NR'-group, -O- or -S-; wherein R' is as defined above;


or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0122] In a further particular embodiment, R₅ represents a group selected from

i) a group of formula (IIIc-1), and ii) a group of formula (IIIc-2),



formula (IIIc-1)



formula (IIIc-2)

wherein




R₁₉, R₂₀, and R₂₈ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y represents a linker selected from a -NR'-group, -O- or -S-; wherein R' is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a

cyano group, a $-\text{CHF}_2$ group or a $-\text{CF}_3$ group.

[0123] In another particular embodiment, R_5 represents a group selected from

i) a group of formula (IIId-1), ii) a group of formula (IIId-2), and iii) a group of formula (IIId-3),



wherein

R_{21} , R_{28} , R_{29} , and R_{30} each independently represent a hydrogen atom, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(\text{CH}_2)_{0-3}\text{CN}$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{1-3}\text{CN}$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}'$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$, a $-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group; and wherein Y represents a linker selected from a $-\text{NR}'$ -group, $-\text{O}-$ or $-\text{S}-$; wherein R' is as defined above;

or in the case that Y represents a $-\text{NR}'$ - group, R_4 together with the $-\text{NR}'$ - group and the carbon atom to which both R_4 and the $-\text{NR}'$ - group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a $-\text{CHF}_2$ group or a $-\text{CF}_3$ group.

[0124] Typically, in the compound of formula (I), Y represents a linker selected from a $-\text{NR}'$ -group, $-\text{O}-$ or $-\text{S}-$; wherein R' represents a hydrogen atom, a hydroxyl group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group. Preferably Y represents a linker selected from a $-\text{NR}'$ - group or $-\text{S}-$; wherein R' represents a hydrogen atom, a hydroxyl group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group. More preferably Y represents a linker selected from a $-\text{NR}'$ - group or $-\text{S}-$; wherein R' represents a hydrogen atom or a linear or branched C_1 - C_3 alkyl group. Most preferably Y represents a $-\text{NR}'$ - group; wherein R' represents a hydrogen atom or a linear or branched C_1 - C_3 alkyl group.

[0125] When R' and/or R'' are attached to a nitrogen atom, preferably R' and/or R'' do not represent a hydroxyl group or alkoxy group.

[0126] Where any of the above moieties represent $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}_8$ or $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}'$, it is preferable that R_8 and R' do not represent a hydrogen atom if the alkylene spacer moiety is absent.

[0127] Preferably in the compound of formula (I):

R_a and R_b each independently represent a hydrogen atom or a linear or branched C_1 - C_3 alkyl group;

n represents 0, 1 or 2;

R_1 represents a hydrogen atom, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_3 - C_7 cycloalkyl group, a phenyl group, a 5- to 10- membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, a tetrahydropyranyl group or a morpholinyl group;

wherein the cycloalkyl, phenyl, heteroaryl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydropyranyl or morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(\text{CH}_2)_{0-3}\text{OR}_8$ group, a $-(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}_8$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group or a $-(\text{CH}_2)_{0-3}(\text{phenyl})-\text{OR}_8$ group; wherein R_7 and R_8 each independently represent a hydrogen atom or a C_1 - C_4 alkyl group;


R_2 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, a $-\text{NH}_2$ group, a $-\text{N}(\text{CH}_3)\text{H}$ group, a $-\text{N}(\text{CH}_3)_2$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_2 alkoxy group;

R₃ represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group;

R₄ represents a hydrogen atom, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C₁-C₃ alkoxy group; a C₁-C₃ haloalkyl group; a linear or branched C₁-C₄ hydroxyalkyl group; a C₃-C₄ cycloalkyl group; a -(CH₂)₀₋₃NR'R'' group; a -(CH₂)₁₋₃O(C₁-C₄ alkyl group); a -(CH₂)₀₋₃OC(O)-(C₁-C₄ alkyl group); a -(CH₂)₀₋₃C(O)O-(C₁-C₄ alkyl group); a -C(O)-(CH₂)₀₋₃NR'R'' group; a -(CH₂)₀₋₃C(O)OH group; a -(CH₂)₀₋₃-(5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N); a -(CH₂)₀₋₃-(5- to 14- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N); or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group; wherein the heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C₁-C₄ alkyl group or a C₁-C₄ haloalkyl group,

R₅ represents a moiety of formula (IIa-2), (IIb-1), (IIb-2), (IIb-3), (IIIa-1), (IIIa-3), (IIIa-5) or (IIIa-14):




formula (IIa-2)

formula (IIb-1)

formula (IIb-2)

formula (IIb-3)



formula (IIIa-1)

formula (IIIa-3)

formula (IIIa-5)

formula (IIIa-14)

wherein:

- R₉, R₁₀, and R₁₂ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;
- R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a halogen atom, a C₁-C₄ haloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group; a phenyl group, which phenyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom or a hydroxyl group; a 5- to 7- membered monocyclic heteroaryl group containing at least one heteroatom selected from O, S and N, which heteroaryl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom or a linear or branched C₁-C₄ alkyl group;
- R₁₈, R₂₂, R₂₃ and R₂₄ each independently represent a hydrogen atom, a halogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group,

- a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;
- Y represents a -NR'-group, -O- or -S-; wherein R' represents hydrogen or a linear or branched C₁-C₄ alkyl group; or in the case that Y represents a -NR'-group, R₄ together with the -NR'-group and the carbon atom to which both R₄ and the -NR'-group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0128] In a particular preferred embodiment, in the compound of formula (I)

X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom or a methyl group;

R₁ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a methyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a pyridinyl group, a pyrazolyl group, an isoxazolyl group, a piperidinyl group or a tetrahydropyranyl group; wherein the cycloalkyl, phenyl, pyridinyl, pyrazolyl, isoxazolyl, piperidinyl or tetrahydropyranyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ alkyl group, a -(CH₂)₀₋₃-(phenyl)-O-(C₁-C₃ alkyl group), a -NR₇R₈ group or a -OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₂ and R₃ each independently represent a hydrogen atom, a halogen atom, a cyano group, a C₁-C₃ haloalkyl group or a linear or branched C₁-C₃ alkyl group;

R₄ represents a hydrogen atom, a C₁-C₃ haloalkyl group, a C₁-C₃ hydroxyalkyl group or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ hydroxyalkyl group, a linear or branched C₁-C₃ alkyl group or a cyclopropyl group;

R₆ represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a linear or branched C₁-C₄ hydroxyalkyl group; a C₃-C₇ cycloalkyl group; a linear or branched C₁-C₃ alkyl group;

a -(CH₂)₀₋₃NR'R" group; a -(CH₂)₁₋₃O(C₁-C₃ alkyl group); a -(CH₂)₀₋₃OC(O)-(C₁-C₃ alkyl group); a -(CH₂)₀₋₃C(O)O-(C₁-C₃ alkyl group); a -C(O)-NR'R" group; a -(CH₂)₀₋₃C(O)OH group; a -(CH₂)₀₋₃-(imidazolyl group); a -(CH₂)₀₋₃-(oxazolyl group);

a -(CH₂)₀₋₃-(oxadiazolyl group); a -(CH₂)₀₋₃-(pyrazolyl group) or a -(CH₂)₀₋₃-(morpholinyl group); wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; and wherein the imidazolyl, oxazolyl, oxadiazolyl, pyrazolyl and morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₃ alkyl group or a C₁-C₃ haloalkyl group;

R₅ represents a group selected from:

i) a group of formula (IIa), which group is a purinyl group unsubstituted or substituted by a -NR'R" group;

ii) a group of formula (IIb), which group is selected from a -NR'-pyridinyl group, a -S-pyridinyl group, a -NR'-pyrimidinyl group, a -S-pyrimidinyl group or a -NR'-triazinyl group; wherein the pyridinyl, pyrimidinyl and triazinyl groups are unsubstituted or substituted by one, two or three substituents selected from a halogen atom, a C₁-C₃ haloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R", a -(CH₂)₀₋₃NR'R" group; and

iii) a group of formula (IIc), which group is selected from a -NR'-purinyl group, a -S-purinyl group, a -NR'-7H-pyrrolo[2,3-d]pyrimidinyl group, a -NR'-1H-pyrazolo[3,4-d]pyrimidinyl group or a -NR'-pyrazolo[1,5-a]pyrimidinyl group;

wherein the purinyl, 7H-pyrrolo[2,3-d]pyrimidinyl, 1 H-pyrazolo[3,4-d]pyrimidinyl pyrazolo[1,5-a]pyrimidinyl and groups are unsubstituted or substituted by a halogen atom or a -(CH₂)₀₋₃NR'R" group; or

R₄ and R₅ together with the carbon atom to which they are attached form a pyrrolidinylpurinyl group or a pyrrolidinylpyrimidinyl; wherein the pyrrolidinyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom or a hydroxyl group; and wherein the purinyl group is unsubstituted or substituted by a -(CH₂)₀₋₃NR'R" group; and wherein the pyrimidinyl group is unsubstituted or substituted by one, two or three substituents selected from a -(CH₂)₀₋₃CN group or a -(CH₂)₀₋₃NR'R" group; and

[0129] R' and R" each independently represent a hydrogen atom, a C₁-C₃ alkoxy group or a linear or branched C₁-C₃ alkyl group.

[0130] In another particularly preferred embodiment, in the compound of formula (I)


X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom or a methyl group;

n represents 0 or 1;

R₁ represents a methyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a pyridinyl group, a piperidinyl group or a tetrahydropyranyl group;

wherein the cycloalkyl, phenyl, pyridinyl, piperidinyl or tetrahydropyranyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₃ alkyl group, a -NR₇R₈ group or a -OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group; R₂ and R₃ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group; R₄ represents a hydrogen atom, a C₁-C₃ haloalkyl group, or a linear or branched C₁-C₃ alkyl group; R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ alkyl group or a cyclopropyl group; R₅ represents a moiety of formula (II-a2), (IIb-1), (IIb-2), (IIb-3), (IIIa-1), (IIIa-3), (IIIa-5) or (IIIa-14):



wherein:

- R₉, R₁₀ and R₁₂ independently represent a hydrogen atom or a -NR'R" group;
- R₁₃ to R₁₆ independently represent a hydrogen atom, a halogen atom, a -CN group, a -C(O)-NR'R" group or a -NR'R" group;
- R₁₈, R₂₂, R₂₃ and R₂₄ represent a hydrogen atom, a halogen atom or a -NR'R" group;
- R' and R" each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group; and
- Y represents -NH- or -S-; or
- Y represents a nitrogen atom and Y, R₄ and the carbon atom to which both R₄ and Y are bonded form a pyrrolidinyl ring, which pyrrolidinyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom or a hydroxyl group.

[0131] In one embodiment, in the compound of formula (I)

X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group or a linear or branched C₁-C₄ alkyl group;

n represents 0, 1, 2 or 3;

R₁ represents a linear or branched C₁-C₄ alkyl group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a monocyclic or bicyclic C₆-C₁₄ aryl group, a 5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N, or a 5- to 14-membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N,

wherein the cycloalkyl, cycloalkenyl, aryl, heteroaryl, and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₁₋₃CN group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group,

a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R₈ group, a -C(O)-(CH₂)₀₋₃-NR₇R₈ group, a -S(O)₂(CH₂)₀₋₃R₈ group or a -S(O)₂(CH₂)₀₋₃NR₇R₈ group;

R₂ and R₃ each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄

alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₇ cycloalkyl group, a -NR'R'' group, or a linear or branched C₁-C₆ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₇ cycloalkyl group;

R₄ represents a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₇ cycloalkyl group, a -(CH₂)₁₋₄NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group, a C₃-C₄ cycloalkyl group, a -C(O)-(CH₂)₀₋₃-R' group or a -C(O)-(CH₂)₀₋₃-NR'R'' group;


R₆ represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₇ cycloalkyl group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group;

R₇ and R₈ each independently represent a hydrogen atom, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group;

R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;


R₅ represents a group selected from:

i) a group of formula (IIa)



formula (IIa)

ii) a group of formula (IIb)



formula (IIb)

and

iii) a group of formula (IIc)



formula (IIc)

wherein

Y represents a linker selected from a -NR'- group, -O- or -S-; wherein R' is as defined above;

(*) represents where R₅ is bonded to the carbon atom attached to R₄ and to the pyrrolo[1,2-a]triazinone group;

W₁ represents a -CR₁₁ group and W₂ represents a nitrogen atom, or W₁ represents a nitrogen atom and W₂ represents a -CR₁₂ group;

G₁ represents a -CR₁₄ group and G₂ represents a nitrogen atom, or G₁ represents a nitrogen atom and G₂ represents

a -CR₁₅ group, or G₁ represents a -CR₁₄ group and G₂ represents a -CR₁₅ group;


G₃ represents a nitrogen atom or a -CR₁₆ group;

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group,

a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' are as defined above;


R₁₇ represents a group selected from

a) a group of formula (IIIa),




formula (IIIa)

b) a group of formula (IIIb),




formula (IIIb)

c) a group of formula (IIIc), and



formula (IIIc)

d) a group of formula (IIId),



formula (IIId)

wherein

G₄ represents a nitrogen atom or a -CR₂₂ group;

G₅ and G₆ each independently represents a nitrogen atom or a -carbon atom, wherein when one of G₅ and G₆ represents a nitrogen atom the remaining represents a carbon atom;

G₇ represents a -NH group or a -CH group;

G₈ represents a nitrogen atom or a -CR₂₃ group;

G₉ represents a nitrogen atom or a -CR₂₄ group;

G₁₀ represents a nitrogen atom or a -CR₂₅ group;

G₁₁ represents a nitrogen atom or a -CR₂₆ group;

G₁₂ represents a nitrogen atom or a -CR₂₇ group;

G₁₃ represents a nitrogen atom or a -CR₂₈ group;

G₁₄ and G₁₅ each independently represents a nitrogen atom or a carbon atom, wherein when one of G₁₄ and G₁₅ represents a nitrogen atom the remaining represents a carbon atom;

G₁₆ represents a -NH group or a -CH group;

G₁₇ represents a nitrogen atom or a -CR₂₉ group;

G₁₈ represents a nitrogen atom or a -CR₃₀ group;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉, and R₃₀ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' are as defined above; and wherein Y is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

[0132] In this embodiment, it is preferred that in the compound of formula (I):

R_a and R_b each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

n represents 0, 1 or 2;

R₁ represents a linear or branched C₁-C₄ alkyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a 5- to 10- membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, a tetrahydropyranyl group or a morpholinyl group; wherein the cycloalkyl, phenyl, heteroaryl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydropyranyl or morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group, a -C(O)-(CH₂)₀₋₃-R₈ group or a -C(O)-(CH₂)₀₋₃-NR₇R₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a C₁-C₄ alkyl group;


R₂ represents a hydrogen atom, a halogen atom, a hydroxyl group, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group;

R₃ represents a hydrogen atom, a halogen atom, a hydroxyl group, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group;

R₄ represents a hydrogen atom, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group;

R₅ represents a moiety of formula (II-a2), (IIb-1), (IIb-2) or (IIIa-1):



wherein:

- R₉, R₁₀, and R₁₂ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;
- R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R'' group or a -(CH₂)₀₋₃NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom or a linear or branched C₁-C₄ alkyl group;
- R₁₈ and R₂₃ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group;

wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;

- Y represents a -NR'-group, -O- or -S-; wherein R' represents hydrogen or a linear or branched C₁-C₄ alkyl group; or in the case that Y represents a -NR'-group, R₄ together with the -NR'-group and the carbon atom to which both R₄ and the -NR'-group are bonded form a 4- to 7-membered, saturated N-containing heterocycl group.

[0133] In a particularly preferred embodiment, in the compound of formula (I)

X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom or a methyl group;

n represents 0 or 1;

R₁ represents a methyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a pyridinyl group, a piperidinyl group or a tetrahydropyranyl group;

wherein the cycloalkyl, phenyl, pyridinyl, piperidinyl or tetrahydropyranyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₃ alkyl group, a -NR₇R₈ group or a -OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₂ and R₃ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₄ represents a hydrogen atom, a C₁-C₃ haloalkyl group, or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ alkyl group or a cyclopropyl group;

R₅ represents a group selected from:

i) a group of formula (IIa), which group is a purinyl group unsubstituted or substituted by a -NR'R" group;

ii) a group of formula (IIb), which group is selected from a -NH-pyridinyl group, a -S-pyridinyl group, a -NH-pyrimidinyl group or a -S-pyrimidinyl group and preferably from a -NH-pyridinyl group and a -NH-pyrimidinyl group; wherein the pyridinyl or pyrimidinyl groups are unsubstituted or substituted by one, two or three substituents selected from a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R" or a -(CH₂)₀₋₃NR'R" group and preferably from a -CN group, a -C(O)NH₂ or a -NH₂ group; and

iii) a group of formula (IIc), which group is selected from a -NH-purinyl group or a -S-purinyl group; wherein the purinyl group is unsubstituted or substituted by a -(CH₂)₀₋₃NR'R" group; or

R₄ and R₅ together with the carbon atom to which they are attached form a pyrrolidinylpurinyl group, wherein the purinyl group is unsubstituted or substituted by a -(CH₂)₀₋₃NR'R" group;

R' and R" each independently represent a hydrogen atom, a C₁-C₃ alkoxy group or a linear or branched C₁-C₃ alkyl group, preferably a hydrogen atom or a linear or branched C₁-C₃ alkyl group.

[0134] In another particularly preferred embodiment, in the compound of formula (I)

X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom or a methyl group;

n represents 0 or 1;

R₁ represents a methyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a pyridinyl group, a piperidinyl group or a tetrahydropyranyl group;


wherein the cycloalkyl, phenyl, pyridinyl, piperidinyl or tetrahydropyranyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₃ alkyl group, a -NR₇R₈ group or a -OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₂ and R₃ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₄ represents a hydrogen atom, a C₁-C₃ haloalkyl group, or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ alkyl group or a cyclopropyl group;


R₅ represents a moiety of formula (II-a2), (IIb-1), (IIb-2) or (IIIa-1):



wherein:

- R₉, R₁₀ and R₁₂ independently represent a hydrogen atom or a -NR'R" group;
- R₁₃ to R₁₆ independently represent a hydrogen atom, a -CN group, a -C(O)-NR'R" or a -NR'R" group;
- R₁₈ and R₂₃ represent hydrogen or a -NR'R" group;
- R' and R" each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group; and
- Y represents -NH- or -S-; or
- Y represents a nitrogen atom and Y, R₄ and the carbon atom to which both R₄ and Y are bonded form a pyrrolidinyl ring.


[0135] In a particularly preferred embodiment, the compound of the invention is of formula (Ia)



Formula (Ia)

wherein R₁, R₂, R₃, R₄, R₅, R₆, R_a, R_b and n are as defined above.

[0136] In an alternative particularly preferred embodiment, the compound is of formula (Ib):



Formula (Ib)

wherein R₁, R₂, R₃, R₄, R₅, R_a, R_b and n are as defined above.

[0137] Particular individual compounds of the invention include:

- 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Aminopyrimidin-4-ylamino)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-amino-9H-purin-9-yl)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one; 2-((6-aminopyrimidin-4-ylami-
 5 no)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 4-((4-Oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)picolinamide;
 2-((2-aminopyridin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((9H-purin-6-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-cyclohexylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 10 2-((6-amino-9H-purin-9-yl)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((9H-purin-6-ylthio)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-amino-9H-purin-9-yl)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((9H-purin-6-ylthio)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-(1-(6-amino-9H-purin-9-yl)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 15 (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(6-aminopyrimidin-4-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
 (R)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 20 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(6-aminopyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 2-(1-(6-amino-9H-purin-9-yl)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 25 2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(3-methoxyphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(2,4-difluorophenyl)pyrrolo-[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-benzyl-5-chloropyrrolo[1,2-f][1,2,4]-triazin-4(3H)-one;
 30 2-((6-amino-9H-purin-9-yl)methyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-amino-9H-purin-9-yl)methyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(pyridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(tetrahydro-2H-pyran-4-yl)pyrrolo-[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-methylpiperidin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 35 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-
 carbonitrile;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-
 40 carbonitrile;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-methylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-((1 r,4r)-4-aminocyclohexyl)-5-chloropyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (R)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 45 (S)-4-amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-car-
 bonitrile;
 (S)-2-(1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimi-
 dine-5-carbonitrile;
 50 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimi-
 dine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 55 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 2-(1-(9H-purin-6-ylamino)-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 4-amino-6-(3,3,3-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-
 carbonitrile;

- (S)-4-amino-6-(2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile;
- (S)-3-phenyl-2-(1-(pyrazolo[1,5-a]pyrimidin-7-ylamino)ethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 2-((6-amino-9H-purin-9-yl)methyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 5 (S)-2-(1-(2-amino-9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(4,6-diamino-1,3,5-triazin-2-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-((6-amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-(5-fluoropyridin-2-yl)ethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 10 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (R)-2-(1-(9H-purin-6-ylamino)-2-hydroxyethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (R)-4-amino-6-(2-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 15 (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(methyl(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethyl)amino)pyrimidine-5-carbonitrile;
- 20 (S)-2-(1-(methyl(9H-purin-6-yl)amino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-7-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 25 (S)-4-amino-6-(1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(4,4-difluoro-1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(4,4-difluoro-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile;
- 30 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-6-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 2-((S)-1-(9H-purin-6-ylamino)ethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 4-amino-6-((S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 35 (S)-4-amino-6-(1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-((9H-purin-6-ylamino)methyl)-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-((4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile;
- 40 (S)-2-(1-(5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 45 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 50 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- 4-amino-6-((1S)-1-(5-(1,2-dihydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 55 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-Amino-6-(1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

- (S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)imidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(5-(difluoromethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 2-(1-(9H-Purin-6-ylamino)-2,2,2-trifluoroethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)propyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (R)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-carbamoylpyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(2-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(2-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 ((S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(2H-tetrazol-5-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(3-((5-methylisoxazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-7-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-7-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(2,6-Diamino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-4-Amino-6-(1-(5-(morpholinomethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

- (S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 5 4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(3-(5-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylic acid;
- 10 2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (S)-4-Amino-6-(1-(3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 15 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 20 (S)-2,4-Diamino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-(1-(3-((1H-Pyrazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)-6-aminopyrimidine-5-carbonitrile;
- 25 (S)-4-Amino-6-(1-(4-oxo-3-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(4-oxo-3-(2,2,2-trifluoroethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 30 (S)-2-Amino-4-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 4-Amino-6-(1-(5-(1-methyl-1H-pyrazol-4-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 35 (S)-4-Amino-6-(1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 40 2-((3-Iodo-1H-pyrazolo[3,4-d]pyrimidin-4-ylamino)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- 4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 45 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-5-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-Amino-6-(1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-N,N-dimethyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;
- 50 (S)-4-Amino-6-(1-(3-(1-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile
- (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-N-propyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;
- 55 2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- 2-((S)-1-(9H-purin-6-ylamino)ethyl)-3-((S)-tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-

carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile;

4-Amino-6-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)-3-hydroxypropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-(2-methyloxazol-5-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-Propyl 2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylate;

(S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-3-hydroxypropylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(4-oxo-3-(6-(trifluoromethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-yl)ethyl acetate;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((2S,4R)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-hydroxypyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

4-Amino-6-((2S,4R)-2-(5-(aminomethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-4-hydroxypyrrolidin-1-yl)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-(4-methyl-1H-imidazol-1-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)cyclopropylamino)pyrimidine-5-carbonitrile;

2-(1-(9H-Purin-6-ylamino)cyclopropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one; (S)-4-Amino-6-(1-(4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;

or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof.

[0138] Examples of the preferred compounds are:

- (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-Amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile;
- (S)-2-(1-(2-amino-9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (R)-4-amino-6-(2-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-7-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-amino-6-(4,4-difluoro-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-6-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- 4-amino-6-((S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-amino-6-(1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-4-amino-6-((4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-4-amino-6-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
- (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
- (S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethyl-

amino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-Amino-6-(1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-Amino-6-(1-(5-(difluoromethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)propyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(R)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(2-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-7-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(2,6-Diamino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2,4-Diamino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(4-oxo-3-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(4-oxo-3-(2,2,2-trifluoroethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-Amino-4-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

trile;

(S)-4-Amino-6-(1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylami-

no)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylami-

no)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(1-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

4-Amino-6-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(4-oxo-3-(6-(trifluoromethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-yl)ethyl acetate;

(S)-4-Amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof.

[0139] In one embodiment, particular compounds of the invention include:

2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-Aminopyrimidin-4-ylamino)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-Amino-9H-purin-9-yl)methyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-aminopyrimidin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

4-((4-Oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)picolinamide;

2-((2-aminopyridin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((9H-purin-6-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-Amino-9H-purin-9-yl)methyl)-3-cyclohexylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-amino-9H-purin-9-yl)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((9H-purin-6-ylthio)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-amino-9H-purin-9-yl)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((9H-purin-6-ylthio)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 5 2-(1-(6-amino-9H-purin-9-yl)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(6-aminopyrimidin-4-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
 10 (R)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(6-aminopyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 15 2-(1-(6-amino-9H-purin-9-yl)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(3-methoxyphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(2,4-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 20 2-((6-Amino-9H-purin-9-yl)methyl)-3-benzyl-5-chloropyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-amino-9H-purin-9-yl)methyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-amino-9H-purin-9-yl)methyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(pyridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(tetrahydro-2H-pyran-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 25 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-methylpiperidin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 30 (S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-methylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-((1r,4r)-4-aminocyclohexyl)-5-chloropyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (R)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 35 (S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 40 (S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 45 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
 50 2-(1-(9H-purin-6-ylamino)-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 4-amino-6-(3,3,3-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;

or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof.

[0140] Examples of the preferred compounds in this embodiment are:

(S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 5 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 10 (S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-((6-amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 15 (S)-4-amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof.

[0141] The compounds of the invention can be prepared using the methods and procedures described herein, or using similar methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given; other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

[0142] Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group, as well as suitable conditions for protection and deprotection, are well known in the art. For example, numerous protecting groups, and their introduction and removal are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York, 1999, and references cited therein.

[0143] The term amino-protecting group refers to a protecting group suitable for preventing undesired reactions at amino nitrogen. Representative amino-protecting groups include, but are not limited to, formyl; acyl groups, for example alkanoyl groups such as acetyl; alkoxycarbonyl groups such as tert-butoxycarbonyl (Boc); arylmethoxycarbonyl groups such as benzyloxycarbonyl (Cbz) and 9-fluorenylmethoxycarbonyl (Fmoc); arylmethyl groups such as benzyl (Bn), trityl (Tr), and 1,1-di-(4'-methoxyphenyl)methyl; silyl groups such as trimethylsilyl (TMS) and tert-butyldimethylsilyl (TBS); and the like. The term hydroxy-protecting group refers to a protecting group suitable for preventing undesired reactions at a hydroxy group. Representative hydroxy-protecting groups include, but are not limited to, alkyl groups, such as methyl, ethyl, and tert-butyl; acyl groups, for example alkanoyl groups, such as acetyl; arylmethyl groups, such as benzyl (Bn), p-methoxybenzyl (PMB), 9-fluorenylmethyl (Fm), and diphenylmethyl (benzhydryl, DPM); silyl groups, such as trimethylsilyl (TMS) and tert-butyldimethylsilyl (TBS); and the like.

[0144] According to one embodiment of the present invention, compounds of general Formula (I) may be prepared by the synthetic route illustrated in Scheme 1, from compounds of Formula (Va), where the group Z₁ represents a halogen atom such as chlorine, bromine and iodine or another suitable leaving group such as methanesulfonate or trifluoromethanesulfonate or other groups such as hydroxyl, that can be converted to suitable leaving groups by standard methods described in the literature, such as the Mitsunobu reaction and others.

[0145] Compounds of Formula (I) can be obtained directly from compounds of Formula (Va) by treatment of (Va) with compounds of Formula (IVa), (IVb) or (IVc) in the presence of a suitable base such as potassium carbonate, diisopropylethylamine or sodium hydride in an appropriate solvent such as *tert*-butanol, *N,N*-dimethylformamide or tetrahydrofuran at temperatures ranging from room temperature to 160 °C, with or without the use of microwaves irradiation.


[0146] When Z₁ is a halogen atom such as chlorine, it can be converted to another more reactive halogen atom such as iodine by treating the compound with the chlorine atom with sodium iodide in acetone at a temperature from room temperature to reflux.

5

10

15

20



Scheme 1


25 Alternatively, compounds of general Formula (I) can be obtained directly from compounds of Formula (Vb), where the group Y represents a -NR'- group, wherein R' is a hydrogen atom, as illustrated in Scheme 2.

30 [0147] Thus, compounds of Formula (Vb) can be treated with electrophiles of Formula (IVd) or (IVe), where the group Z₁ represents a leaving group such as a halogen atom, methanesulphonate or trifluoromethanesulphonate, in the presence of a suitable base such as potassium carbonate, diisopropylethylamine or sodium hydride in an appropriate solvent such as *tert*-butanol, *N,N*-dimethylformamide or tetrahydrofuran at temperatures ranging from room temperature to 220 °C, with or without the use of microwaves irradiation.

35

40

45



Scheme 2

50 [0148] Alternatively, compounds of Formula (Va) where Z₁ is for instance a halogen atom can be converted to compounds of Formula (Vb) where Y is a -NR'- group, wherein R' is as defined above, by treating compounds (Va) with a solution of ammonia in a solvent such as methanol at a temperature between 60 to 120 °C.

55 [0149] Compounds of general Formula (V), which comprises compounds of subFormula (Va) and subFormula (Vb), can be prepared directly from compounds of Formula (VII) as illustrated in Scheme 3 by treatment of compounds with Formula (VII) with the appropriate acid chlorides of Formula (VIII) in a solvent such as acetic acid at a temperature ranging from room temperature to 150°C with or without the use of microwaves irradiation.

[0150] In the particular case where Z₂ is a chlorine atom, the compounds of Formula (V) can also be prepared by treating the compounds of Formula (VII) with 2-chloro-1,1,1-trimethoxyethane in the presence of pyridinium *p*-tolue-

nesulfonate at a temperature between 50 °C and 150 °C.

[0151] Alternatively, compounds of Formula (V) can be obtained in two steps from compounds of Formula (VII), isolating the intermediate amides of Formula (VI).


[0152] Compounds of Formula (VII) can be transformed in amides of Formula (VI) by treatment with carboxylic acids of Formula (IX), where Z_2 represents a leaving group of Formula Z_1 or a nucleophile of Formula Y unprotected or protected by known protecting groups described in the literature, in the presence of an activating agent by methods and conditions well described in the literature, for example using EDC or HATU as an activating agent in a solvent such as tetrahydrofuran or dichloromethane or mixtures of these solvents at temperatures ranging from room temperature to 80 °C.

[0153] Alternatively, amides of Formula (VI) can be obtained from compounds of Formula (VII) by treatment with acid chlorides of Formula (VIII) at room temperature in a suitable solvent such as acetic acid or 1,4-dioxane or alternatively in the presence of a base such as triethylamine in a suitable solvent such as dichloromethane.

[0154] In a second step, compounds of Formula (VI) can yield compounds of Formula (V) by treatment with phosphorous oxychloride at temperatures ranging from room temperature to 100 °C, with or without a subsequent treatment with a solution of a base such as ammonia, pyrrolidine, piperidine, potassium carbonate or sodium methanethiolate in a solvent such as methanol, ethyl acetate or *N,N*-dimethylformamide at a temperature between room temperature and 100 °C.

[0155] Alternatively, amides of Formula (VI) can yield compounds of Formula (V) by heating these amides in a solvent such as xylene or toluene with the presence of pyridinium *p*-toluenesulfonate or *p*-toluenesulfonic acid at a temperature between 80 °C and 160 °C.

[0156] Alternatively, compounds of Formula (VI) can yield compounds of Formula (V) by treatment of compounds of Formula (VI) with the complex resulting from the treatment of triphenylphosphine with bromine in a solvent such as dichloromethane in the presence of a base such as triethylamine at a temperature from room temperature to reflux, with or without a subsequent treatment with a base such as ammonia, pyrrolidine, piperidine, potassium carbonate or sodium methanethiolate in a solvent such as methanol, ethyl acetate or *N,N*-dimethylformamide at a temperature between room temperature and 100 °C.




Scheme 3

[0157] The acid chlorides of Formula (VIII) and the carboxylic acids of Formula (IX) can be used in a protected form to prevent certain functional groups from undergoing undesired reactions. In these cases, standard methods for the removal of these protecting groups can be used at the suitable step of the synthesis. Numerous protecting groups, their introduction and their removal are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York, 1999, and references cited therein.

[0158] Compounds of Formula (VII) can be prepared from carboxylic acids of Formula (XII) following the Scheme described in Scheme 4.

[0159] Carboxylic acids (XII) can be activated with any activating reagent described in the literature such as thionyl chloride, oxalyl chloride, phosphorous oxychloride, 2-(3*H*-[1,2,3]triazolo[4,5-*b*]pyridin-3-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate(V), 3-((ethylimino)methyleneamino)-*N,N*-dimethylpropan-1-aminium chloride, 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphorinane-2,4,6-trioxide and treated with amines of Formula (XI) in the presence of a base such as diisopropylethylamine when needed in a suitable solvent such as dioxane, dichloromethane, *N,N*-dimethylformamide, ethyl acetate or tetrahydrofuran at temperatures ranging from 0 °C to reflux to give amides of Formula (X).

[0160] Subsequently, amides of Formula (X) can be aminated on the nitrogen atom in position 1 by any of the aminating reagents described in the literature, such as *O*-(mesitylenesulfonyl)hydroxylamine, *O*-(*p*-nitrobenzoyl)-hydroxylamine, *O*-(diphenyl-phosphinyl)-hydroxylamine, *O*-(2,4-dinitrophenyl)-hydroxylamine, hydroxylamine-*O*-sulfonic acid using a suitable base such as triethylamine, potassium carbonate, sodium hydride or butyl lithium in an appropriate solvent such as *N,N*-dimethylformamide, tetrahydrofuran, 1,4-dioxane at temperatures ranging from -78 to 100 °C. Alternatively, the amination reaction can be carried out in a biphasic system using an aqueous solution of ammonia, sodium hydroxide, ammonium chloride and sodium hypochlorite and a suitable organic solvent such as dialkyl ethers and adding a phase transfer catalyst such as Aliquat 336® at temperatures ranging from 0 °C to room temperature.



Scheme 4


[0161] In another embodiment of the present invention, compounds of general Formula (I) may be prepared by the synthetic route illustrated in Scheme 5, from compounds of Formula (XV), where the group Z₁ represents a halogen atom such as chlorine, bromine and iodine or another suitable leaving group such as methanesulfonate or trifluoromethanesulfonate or other groups such as hydroxyl, that can be converted to suitable leaving groups by standard methods described in the literature, such as the Mitsunobu reaction and others.

[0162] Compounds of Formula (I) can be obtained from compounds of Formula (XIV) by treatment of (XIV) with the corresponding amines of Formula (XI) in the presence or not of a suitable base such as sodium hexamethyldisilazide or a Lewis acid such as trimethyl aluminium at a temperature ranging from room temperature to 150 °C in an appropriate solvent such as 1,4-dioxane, tetrahydrofuran or dichloromethane. The intermediate diamides of Formula (XIII) obtained were subsequently cyclized to afford compounds of Formula (I) by treatment with phosphorous oxychloride at temperatures ranging from room temperature to 100 °C, with or without a subsequent treatment with a solution of a base such as ammonia, pyrrolidine, piperidine or sodium methanethiolate in a solvent such as methanol, tetrahydrofuran or ethyl acetate at a temperature between room temperature and 100 °C.

[0163] Alternatively, compounds of Formula (XIII) can yield compounds of Formula (I) by treatment of compounds of Formula (XIII) with the complex resulting from the treatment of triphenylphosphine with bromine in a solvent such as dichloromethane in the presence of a base such as triethylamine at a temperature from room temperature to reflux, with or without a subsequent treatment with a base such as ammonia, pyrrolidine, piperidine or sodium methanethiolate in a solvent such as methanol or ethyl acetate at a temperature between room temperature and 100 °C.

[0164] Compounds of Formula (XIV) can be obtained directly from compounds of Formula (XV) by treatment of (XV) with compounds of Formula (IVa), (IVb) or (IVc) in the presence of a suitable base such as potassium carbonate, diisopropylethylamine or sodium hydride in an appropriate solvent such as *tert*-butanol, *N,N*-dimethylformamide or tetrahydrofuran at temperatures ranging from room temperature to 160 °C, with or without the use of microwaves irradiation.

[0165] When Z₁ is a halogen atom such as chlorine, it can be converted to another more reactive halogen atom such as iodine by treating the compound with the chlorine atom with sodium iodide in acetone at a temperature from room temperature to reflux.




Scheme 5

Compounds of general Formula (XV) can be prepared directly from compounds of Formula (XVI) as illustrated in Scheme 6 by treatment of compounds with Formula (XVI) with the appropriate acid chlorides of Formula (VIII) in a solvent such as acetic acid at a temperature ranging from room temperature to 150°C with or without the use of microwaves irradiation.

[0166] Alternatively, compounds of Formula (XV) can be obtained in two steps from compounds of Formula (XVI), isolating the intermediate amides of Formula (XVII).

[0167] Compounds of Formula (XVI) can be transformed in amides of Formula (XVII) by treatment with acid chlorides of Formula (VIII) at a temperature ranging from 0°C to room temperature in a suitable solvent such as acetic acid or 1,4-dioxane.

[0168] In a second step, compounds of Formula (XVII) can yield compounds of Formula (XV) by treatment with phosphorous oxychloride at temperatures ranging from room temperature to 100 °C in a suitable solvent such as 1,4-dioxane. Alternatively, compounds of Formula (XVII) can yield compounds of Formula (XV) by treatment of compounds of Formula (XVII) with the complex resulting from the treatment of triphenylphosphine with bromine in a solvent such as dichloromethane in the presence of a base such as triethylamine at a temperature from room temperature to reflux, with or without a subsequent treatment with a base such as ammonia, pyrrolidine, piperidine or sodium methanethiolate or potassium carbonate in a solvent such as methanol, ethyl acetate or *N,N*-dimethylformamide at a temperature between room temperature and 100 °C.




Scheme 6

[0169] Compounds of Formula (XVI) may be obtained from compounds of Formula (XII) as illustrated in Scheme 7 by treatment with benzyl bromide and a suitable base such as triethylamine or cesium carbonate in an appropriate solvent such as *N,N*-dimethyl-formamide or acetonitrile at temperatures ranging from 0 to 60 °C following the protocol described elsewhere in the literature. Alternatively, compounds of Formula (XII) may be coupled with benzyl alcohol activating the carboxylic group of (XII) through the formation of the corresponding acid chloride or any other activated ester.

[0170] Subsequently, esters of Formula (XIX) can be aminated on the nitrogen atom in position 1 by any of the aminating reagents described in the literature, such as *O*-(mesitylenesulfonyl)-hydroxylamine, *O*-(*p*-nitrobenzoyl)-hydroxylamine, *O*-(diphenyl-phosphinyl)-hydroxylamine, *O*-(2,4-dinitrophenyl)-hydroxylamine, hydroxylamine-*O*-sulfonic acid using a suitable base such as triethylamine, potassium carbonate, sodium hydride or butyl lithium in an appropriate solvent such as *N,N*-dimethylformamide, tetrahydrofuran, 1,4-dioxane at temperatures ranging from -78 to 100 °C. Alternatively, the amination reaction can be carried out in a biphasic system using an aqueous solution of ammonia, sodium hydroxide, ammonium chloride and sodium hypochlorite and a suitable organic solvent such as dialkyl ethers and adding a phase transfer catalyst such as Aliquat 336® at temperatures ranging from 0 °C to room temperature.

[0171] Preparation of compounds of Formula (XVI) can be done by hydrogenolysis using an appropriate catalyst such as 10% palladium on charcoal in a suitable solvent such as an alkyl alcohol under a hydrogen atmosphere at pressures ranging from atmospheric pressure to 60 psi and at temperatures ranging from room temperature to 60°C. Alternatively, it is also possible to add an acid to the reaction media such as hydrochloric acid to favour the hydrogenolysis process. Also, compounds of Formula (XVI) can be obtained by saponification of esters (XVIII) using an acid such as hydrochloric acid or sulphuric acid or a base such as sodium hydroxide in an appropriate solvent such as water or alkyl alcohols at temperatures ranging from room temperature to 100°C.




Scheme 7

[0172] Compounds (XII) can either be commercially available compounds or can be prepared by the synthetic Schemes illustrated in Schemes 8 and 9. In the particular case when X represents CR₆ being R₆ a C₃-C₇ cycloalkyl group, or a linear or branched C₁-C₄ alkyl group, saturated or unsaturated C₁-C₄ alkyl group compounds of Formula XIIa can be prepared, as illustrated in Scheme 8, from bromopyrrole of Formula (XXIa)² by Suzuki or Stille coupling with the corresponding boronic acids or organotin compound in the presence of a palladium catalyst such as tetrakis(triphenylphosphane) palladium(0) or palladium acetate with or without an appropriate base such as potassium carbonate or cesium carbonate and in a suitable solvent such as toluene or dioxane or *N,N*-dimethylformamide at temperatures ranging from 60°C to 150°C.

[0173] In the particular case where X represents CR₆, being R₆ a trifluoromethyl group, the bromine atom of compound of Formula (XXIa) can be converted into a iodine atom by treatment of (XXIa) with sodium iodide in the presence of a catalysts such as copper (I) iodide and a chelating amine such as *trans*-1,2-bis(methylamino)cyclohexane in an appropriate solvent such as 1,4-dioxane at a temperature ranging from 60°C to reflux. Next, treatment of the iodine intermediate with methyl 2,2-difluoro-2-(fluorosulfonyl)acetate or any other trifluoro methylating agent using a suitable catalyst such as copper (I) iodide in the presence or not of a chelating agent such as hexamethylphosphoramide and in an appropriate solvent such as *N,N*-dimethylformamide afforded intermediate compound of Formula (XXa).


[0174] In the particular case where R₆ is a cyano group, the bromine atom of compound of Formula (XXIa) can be converted first into a iodine with the methods previously described or treated directly with dicyanozinc in the presence of a palladium catalyst such as tetrakis(triphenylphosphane) palladium (0) in an appropriate solvent such as *N,N*-dimethylformamide at a temperature ranging from 60°C to 150°C or by using copper cyanide in a solvent such pyridine at temperatures ranging from 60°C to 150°C. In the particular case where R₆ is a difluoromethyl group, vinyl intermediates synthesized by Stille reaction previously described can be treated with diethylaminosulfur trifluoride (DAST) in an appropriate solvent such as dichloromethane at a temperature between -78°C and room temperature to yield the compounds of Formula (XXa).

[0175] Finally, compounds of Formula (XIIa) can be obtained by simultaneous cleavage of the sulphone and ester groups of compounds of Formula (XXa) by means of a base such as lithium hydroxide in a suitable solvent or mixture of solvents such as water or tetrahydrofurane at temperatures ranging from room temperature to 220 °C, with or without the use of microwaves irradiation. Alternatively, the cleavage of the sulphone and ester groups of compounds of Formula (XXa) can be done sequentially by treatment of compounds (XXa) with tetrabutylammonium fluoride in an appropriate solvent such as tetrahydrofurane at a temperature from room temperature to reflux and subsequent hydrolysis of the ester group by any of the methods well known in the literature.



Scheme 8


[0176] In the particular case when X represents CR₆ being R₆ hydrogen or C₃-C₇ cycloalkyl group, or a linear or branched C₁-C₄ alkyl group, and R₂ independently represents hydrogen or C₃-C₇ cycloalkyl group, or a linear or branched C₁-C₄ alkyl group, compounds Xa can be prepared, as illustrated in Scheme 9, from pyrroles of Formula (XXIIIa). Pyrroles of Formula (XXIIIa) can be reacted with 2,2,2-trichloroacetyl chloride in a suitable solvent such as diethyl ether at a temperature ranging from room temperature to reflux affording ketones of Formula (XXIIa). These intermediate compounds of Formula (XXIIa) can be reacted with the corresponding amines of Formula (XI) with or without solvent in the presence of a base such as triethylamine at a temperature ranging from room temperature to 150°C to afford compounds of Formula (Xa).



Scheme 9


[0177] In another embodiment of the present invention compounds of Formula (X) may be prepared by the synthetic route illustrated in Scheme 10 from treating compounds of Formula (XXV) with amines of Formula (XI) in the presence or not of a Lewis acid such as trimethyl aluminium in an appropriate solvent such as toluene and a temperature ranging from room temperatures to 120°C.

[0178] Compounds of general Formula (XXV) can be obtained from commercial sources or prepared as described in the literature². Additionally, conventionally protecting groups may be necessary to prevent NH group from pyrrole from undergoing undesired reactions such as phenylsulphonyl group.




Scheme 10

[0179] Alternatively, compounds of general Formula (VI) may be prepared by the synthetic route illustrated in Scheme 11. Thus, compounds of Formula (VI) can be prepared from compounds of Formula (XXVII) by known amide formation methods such as those described above. Compounds of Formula (XXVII) can be prepared by the known coupling methods previously described. Compounds of Formula (XXVI) can be obtained by amination of compounds of Formula (XXV) by the methods already described.



Scheme 11

[0180] Compounds of general Formula (V) where X represents CR_6 being R_6 as defined in claim 1 can also be synthesized from compounds of Formula (Vc) as shown in Scheme 12 by the general methods described as it follows.



Scheme 12

[0181] In the particular case where R_6 is a trifluoromethyl group, the bromine atom of compound of Formula (Vc) can be converted first into a iodine and subsequently transformed to a trifluoromethyl group following the general methods previously described.

[0182] In the particular case where R_6 is a cyano group, the bromine atom of compound of Formula (Vc) can be converted first into a iodine with the methods previously described or treated directly with dicyanozinc in the presence of a palladium catalyst such as tetrakis(triphenylphosphane) palladium (0) in an appropriate solvent such as *N,N'*-dimethylformamide at a temperature ranging from 60°C to 150°C or by using copper cyanide in a solvent such pyridine at temperatures ranging from 60°C to 150°C.


[0183] In the particular case where R_6 is an alkyl or cycloalkyl group, or an aromatic or heteroaromatic ring compounds of Formula (V) can be obtained by standard Suzuki or Stille couplings with the corresponding boronic acid or organotin compound in the presence of a palladium catalyst such as tetrakis(triphenylphosphane) palladium(0) or palladium acetate with or without an appropriate base such as potassium carbonate or cesium carbonate and in a suitable solvent such as toluene or dioxane or *N,N*-dimethylformamide at temperatures ranging from 60°C to 150°C.

[0184] Alternatively, non commercial heteroaromatic rings can be prepared from compounds of Formula (V) where X is CR_6 being R_6 a cyano groups or a carboxylic acid with the standard methods described in the literature.

[0185] In the particular case where R_6 is a fluorine, compound of Formula (Vc) can be treated with a lithiated agent such as *n*-BuLi, in a non protic solvent such as hexanes and at a temperature between -78°C and 0°C and subsequently

treated with a suitable fluorine source such as N-fluoro-N-(phenylsulfonyl)-benzenesulfonamide at a temperature between -78°C and room temperature.

[0186] Compounds of general Formula (Vf) to (Vk) can be synthesized by the general methods illustrated in Scheme 13.



Scheme 13

[0187] Compounds of Formula (Vi) can be prepared by treating compounds of general Formula (Vh) with a reducing reagent such as NaBH₄ in a protic solvent such as methanol at room temperature. Compound of Formula (Vi) can then be further derivatized by treating those compounds with haloalkanes or carboxylic acids to obtain the corresponding ethers or esters.

[0188] Compounds of Formula (Vj) can be synthesized by treating compounds of Formula (Vh) with diethylaminosulfur trifluoride (DAST) in an appropriate solvent such as dichloromethane at a temperature between -78°C and room temperature. Compounds of Formula (Vk) can be prepared by treating compounds of Formula (Vh) with an amine of Formula NHR_cR_d in an appropriate solvent such as acetic acid at room temperature and subsequently adding a reducing agent such as sodium cyanoborohydride.

[0189] Compounds of Formula (Vh) can be prepared by treating compounds of Formula (Ve) with osmium tetroxide and 4-methylmorpholine-4-oxide in an appropriate solvent such as tetrahydrofuran at room temperature obtaining the 1,2-dihydroxyethyl derivative and then by treatment with sodium periodate in an appropriate solvent such as tetrahydrofuran at room temperature.

[0190] Alternatively, compounds of general Formula (Vh) may be prepared by ozonolysis of compounds of Formula (Ve) in a mixture of acetone and water at temperatures ranging from -25°C to 0°C.

[0191] Compounds of Formula (Ve) can be prepared by treating compounds of Formula (Vc) with ethenyl(tributyl)tin in the presence of a palladium catalyst such as tetrakis(triphenylphosphane) palladium(0) in a suitable solvent such as *N,N*-dimethylformamide at temperatures ranging from 60°C to 150°C.

[0192] Compounds of Formula (Vf) can be prepared by ozonolysis of compounds of Formula (Ve) in a mixture of solvents such as ethyl acetate and pyridine at temperatures ranging from -25°C to 0°C. In the particular case where R₆ is an ester, compounds of Formula (Vf) can be treated with the corresponding haloalkane in the presence of a base such as potassium carbonate in a solvent such as *N,N*-dimethylformamide at temperatures ranging from 0°C to 150°C.

[0193] In the particular case of compounds of (Vd) where R₆ is an amide, compounds can be obtained by treating compounds of Formula (Vf) with the corresponding amine with a coupling reagent such as *N*-(3-dimethylaminopropyl)-*N*'-

ethylcarbodiimide hydrochloride in the presence of 1-hydroxybenzotriazole in a solvent such as *N,N'*-dimethylformamide at temperatures ranging from 0°C to 150°C.


[0194] In the particular case of the primary amide it can be obtained from compounds where R₆ is a cyanogroup by treating with concentrated sulphuric acid at room temperature.

[0195] Compounds of Formula (Vg) can be prepared by treating compounds of Formula (Ve) with an hydroborating agent such as 9-borabicyclo[3.3.1]nonane in an appropriate solvent such as tetrahydrofurane at temperatures between -5°C and room temperature and subsequently treating with an oxidative reagent such as hydrogen peroxide in the presence of a base such as sodium hydroxide at temperatures ranging from -5°C to room temperature.

[0196] Compounds of formula (Vg) can be treated with haloalkanes in the presence of a base such as sodium hydride in an aprotic solvent such as tetrahydrofurane to obtain the corresponding ethers. In another embodiment compounds of formula (Vg) can also be treated with carboxylic acids at temperatures ranging from 25°C to 150°C to obtain the corresponding esters.

[0197] In the particular case of compounds of Formula (Vd) where R₆ is hydrogen, compounds can alternatively be obtained by hydrogenolysis of compounds of Formula (Vc) using an appropriate catalyst such as 10% palladium on charcoal in a suitable solvent such as an alkyl alcohol under a hydrogen atmosphere at pressures ranging from atmospheric pressure to 60 psi and at temperatures ranging from room temperature to 60°C.

[0198] In another embodiment of the present invention, compounds of Formula (V) can alternatively be prepared from compounds of Formula (Vm) as shown in Scheme 14 using the derivatization methods described above. In some particular cases the bromine atom of compound of Formula (Vm) can be converted first into a iodine of Formula (Vn) and subsequently derivatized following the general methods previously described.



Scheme 14

EXAMPLES

General

[0199] The syntheses of the compounds of the invention and of the intermediates for use therein are illustrated by the following Examples (1-187) (including Preparation Examples (Preparations 1-146)) are given in order to provide a person skilled in the art with a sufficiently clear and complete explanation of the present invention, but should not be considered as limiting of the essential aspects of its subject, as set out in the preceding portions of this description.

[0200] Reagents, starting materials, and solvents were purchased from commercial suppliers and used as received. Concentration or evaporation refers to evaporation under vacuum using a Büchi rotatory evaporator.

[0201] Reaction products were purified, when necessary, by flash chromatography on silica gel (40-63 µm) with the solvent system indicated. Purifications in reverse phase were made in a Biotage SP1® automated purification system equipped with a C₁₈ column and using a gradient of water-acetonitrile/MeOH (1:1) (0.1% v/v ammonium formate both phases) from 0% to 100% acetonitrile/MeOH (1:1) in 40 column volumes. The appropriate fractions were collected and the solvents evaporated under reduced pressure and/or lyophilized.

[0202] Preparative HPLC-MS were performed on a Waters instrument equipped with a 2767 injector/collector, a 2525

binary gradient pump, a 2996 PDA detector, a 515 pump as a make-up pump and a ZQ4000 Mass spectrometer detector or on a Agilent 1200 Series coupled to an Agilent 6120 Mass spectrometer detector. Both systems were equipped with a Symmetry Prep C₁₈ (19 x 300 mm, 7 µm) column or a XBridge Prep C₁₈ (19 x 100 mm, 5 µm) column. The mobile phase was formic acid (0.4 mL), ammonia (0.1 mL), methanol (500 mL) and acetonitrile (500 mL) (B) and formic acid (0.5 mL), ammonia (0.125 mL) and water (1000 mL) (A), the specific gradients used are specified in each particular case. The flow rate was 20 mL/min.

[0203] Purity and MS identification was performed in a Waters 2795 system coupled to a 2996 Diode array detector and to a Waters ZQ mass spectrometer detector or in a Waters Acquity UPLC system coupled to a SQD mass spectrometer detector. The injection volume was 5 microliter on the HPLC and 0.5 microliter on the UPLC. Chromatograms were processed at 210 nM or 254 nM. Mass spectra of the chromatograms were acquired using positive and negative electrospray ionization. The mobile phase was formic acid (0.4 mL), ammonia (0.1 mL), methanol (500 mL) and acetonitrile (500 mL) (B) and formic acid (0.5 mL), ammonia (0.125 mL) and water (1000 mL) (A) and a gradient between 0 to 95% of B was used. Columns: HPLC: Waters Symmetry (2.1x50mm, 3.5 µm); UPLC: ACQUITY UPLC BEH C-18 (2.1x50mm, 1.7 µm)

[0204] ¹H Nuclear Magnetic Resonance Spectra were recorded on a Varian Gemini-2000 spectrometer operating at a frequency of 300 MHz for the ¹H spectra or in a Varian Mercury plus operating at a frequency of 400 MHz for the ¹H spectra. Samples were dissolved in the specified deuterated solvent. Tetramethylsilane was used as reference.

Abbreviations:

[0205]

DMF	Dimethylformamide
DMSO	Dimethylsulfoxide
CDCl ₃	Deuterated chloroform
NMR	Nuclear magnetic resonance
s	Singlet
d	Doublet
dd	Doublet doublet
td	Triple doublet
br	Broad
q	Quartet
t	Triplet
m	Multiplet
LRMS	Low resolution mass spectrometry
h	hour
min	minutes
NMM	N-methylmorpholine
DMF	N,N-dimethylformamide
DCM	dichloromethane, methylene chloride
AcOEt	ethyl acetate
DMSO	dimethylsulfoxide
EDC·HCl	3-((ethylimino)methyleneamino)-N,N-dimethylpropan-1-aminium chloride
THF	tetrahydrofuran
DIEA	diisopropylethylamine
HOBt	1-Hydroxybenzotriazole hydrate
MeOH	methanol
DPPONH ₂	P,P-diphenylphosphinic amide
PPTS	pyridinium p-toluenesulphonate
Pd(PPh ₃) ₄	Tetrakis(triphenylphosphane) palladium(0)
HMPA	hexamethylphosphoramide
Cellite®	diatomaceous earth
BINAP	2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl
T3P®	2,4,6-Tripropyl-1,3,5,2,4,6-trioxatriphosphorinane-2,4,6-trioxide
HATU	2-(1 H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate

PREPARATION 1

1-Amino-3-chloro-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide**a) 3-Chloro-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide**

[0206] A suspension of the 3-chloro-1*H*-pyrrole-2-carboxylic acid¹ (1.2 g, 8.3 mmol) in thionyl chloride (6 ml) was heated to reflux during 30 minutes. At the end of this period, volatiles were removed by distillation under reduced pressure coevaporating with tetrahydrofuran a couple of times. The residue obtained after evaporation was dissolved in a small amount of dry dioxane and was added to a solution of *o*-toluidine (1.33 g, 12.4 mmol) and DIEA (4.32 mL, 25 mmol) in 70 mL of dry 1,4-dioxane at 0°C. Once the addition was finished, the reaction crude was heated to 60°C during 2 hours. Afterwards, this crude was evaporated under vacuum and the residue was taken up with ethyl acetate and washed successively with water, saturated solution of sodium carbonate, water, hydrochloric acid 2*N*, water and brine. The organic phase was separated, dried (sodium sulphate, Na₂SO₄) and concentrated under vacuum to give a residue that was triturated with hexane affording 950mg (88% yield) of a solid after filtration. LRMS (m/z): 235 (M+1)⁺.

b) 1-Amino-3-chloro-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide

[0207] In a 100 mL three-necked flask it was placed 11 mL of a 28% aqueous solution of sodium hydroxide, 4.1 mL of a 28% ammonium hydroxide solution, 1.23 g of ammonium chloride and 0.12 mL of Aliquat 336. Afterwards, a solution of 3-chloro-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide (0.9 g, 3.84 mmol) in 30 mL of diethyl ether and 30 mL of methyl *tert*-butyl ether was added and placed at 0°C affording a suspension. Over this suspension, a 10% aqueous solution of sodium hypochlorite (26 mL) was added drop wise with vigorous stirring maintaining the temperature during 20 min. more. Subsequently, the reaction mixture was stirred at room temperature during a further 1.5 h producing the consumption of the starting material. Next, the reaction crude was diluted with ethyl acetate until no suspended material was observed. The layers were separated and the organic phase was washed with a 25% aqueous solution of sodium thiosulphate, water and brine, dried (Na₂SO₄) and concentrated under vacuum to give a residue that was triturated with hexane to produce a solid (870 mg, 86% yield) after filtration. LRMS (m/z): 250 (M+1)⁺.
¹H NMR (400 MHz, DMSO) δ 11.05 (s, 1H), 8.04 (d, J = 7.5 Hz, 1H), 7.28 - 7.13 (m, 2H), 7.07 - 6.96 (m, 2H), 6.82 (s, 2H), 6.16 (d, J = 3.0 Hz, 1H), 2.29 (s, 3H).

PREPARATION 2

5-Chloro-2-(chloromethyl)-3-*o*-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0208] To a suspension of 1-amino-3-chloro-*N*-phenyl-1*H*-pyrrole-2-carboxamide (0.56 g, 2.24 mmol) in glacial acetic acid (19 mL) was added 0.93 mL (11.7 mmol) of chloroacetyl chloride getting a solution. The reaction mixture was stirred during 2 hours to produce the desired compound. Then, the reaction mixture was poured into ice-water precipitating a solid that was filtered and washed with more water. This solid was taken up with ethyl acetate and the organic phase was washed with 4% aqueous solution of sodium bicarbonate and water, dried and evaporated under vacuum to yield 650 mg (89% yield) of 3-chloro-1-(2-chloroacetamido)-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide as a solid.

[0209] This intermediate compound was dissolved in toluene (40 mL) and 50 mg of pyridinium *p*-toluenesulphonate (PPTS) were added heating the reaction to reflux with a Dean-Stark to remove the water from the reaction media. After 40 hours, the starting material has disappeared and then, the reaction was elaborated by cooling down to room temperature and adding ethyl acetate. This organic phase was washed with water, 4% aqueous solution of sodium bicarbonate and brine and dried (Na₂SO₄). Removal of the volatiles under vacuum produced a residue of 580 mg of the desired product (85% yield).

LRMS (m/z): 308 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 7.48 - 7.32 (m, 4H), 7.23 (d, J = 7.8 Hz, 1 H), 6.56 (d, J = 3.0 Hz, 1H), 4.21 (d, J = 12.1 Hz, 1H), 4.04 (d, J = 12.1 Hz, 1H), 2.22 (s, 3H).

PREPARATION 3

2-(Aminomethyl)-5-chloro-3-*o*-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0210] A solution of the starting 5-chloro-2-(chloromethyl)-3-*o*-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (200 mg, 0.65

mmol) in a 7M ammonia solution in methanol was heated at 85°C with stirring in a sealed tub during 4 h. At the end of this period, the volatiles were removed under vacuum and the residue was taken-up with ethyl acetate and washed with a 4% aqueous solution of sodium bicarbonate, water and brine. The organic layer was dried (Na₂SO₄) and concentrated to dryness giving 140 mg of a residue corresponding to the title compound of the preparation (76% yield).

LRMS (m/z): 289 (M+1)⁺.

PREPARATION 4

3-Cyclopropyl-1*H*-pyrrole-2-carboxylic acid

a) Methyl 3-cyclopropyl-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate

[0211] In a schlenk flask, a mixture of methyl 3-bromo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate² (1.45 g, 4.2 mmol), cyclopropylboronic acid (1.09 g, 12.6 mmol) and potassium carbonate (1.75 g, 12.7 mmol) was suspended in toluene and the system was degassed doing 3 cycles of vacuum-Ar. Next, tetrakis(triphenylphosphane) palladium(0) (Pd(PPh₃)₄) was added as a solid and 3 new cycles of vacuum-Ar were done stirring the reaction at 100°C overnight. At the end of this period, the starting material was consumed and the reaction was worked-up pouring the crude over water and extracting the resulting mixture with ethyl acetate (3x50 mL). The organic solution was washed with water and brine, dried over Na₂SO₄ and concentrated under vacuum to give a residue that was purified by flash chromatography silica (hexane/ethyl acetate). After the purification were obtained 976 mg of the title compound (76% yield).

LRMS (m/z): 306 (M+1)⁺.

b) 3-Cyclopropyl-1*H*-pyrrole-2-carboxylic acid

[0212] To a suspension of the starting methyl 3-cyclopropyl-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate (0.9 g, 3 mmol) in tetrahydrofuran (4.6 mL) and water (2.3 mL) was added lithium hydroxide (0.29 g, 12 mmol) and this mixture was heated to 100°C in a microwave vial during 4 hours. Next, the reaction mixture was poured into water (50 mL) and this aqueous phase was washed with diethyl ether (2x). Afterwards, the aqueous layer was acidified until pH=3 by adding solid phosphoric acid and extracted with ethyl acetate (3x). The organic solution was washed with water and brine, dried (Na₂SO₄) and concentrated in vacuum to give 370 mg of the title compound.

LRMS (m/z): 150 (M-1)⁻.

PREPARATION 5

1-Amino-3-cyclopropyl-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide

a) 3-Cyclopropyl-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide

[0213] To a suspension of the 3-cyclopropyl-1*H*-pyrrole-2-carboxylic acid (50 mg, 0,33 mmol) in dry dichloromethane (0.4 mL) was added a solution of oxalyl chloride (0.04 mL, 0,37 mmol) in 0.3 mL of dichloromethane at room temperature and 3 drops of *N,N*-dimethylformamide. After 1.5 h at this temperature, the mixture was concentrated under vacuum and the residue was redissolved in dichloromethane (1 mL). To this solution was added a solution of *o*-toluidine in dichloromethane (0,2 mL) and the mixture was stirred at room temperature overnight. The reaction was worked-up diluting with dichloromethane and washing with sodium bicarbonate (2x) and brine. The organic phase was dried and concentrated to give 74 mg of the title compound (88% yield).

LRMS (m/z): 241 (M+1)⁺.

b) 1-Amino-3-cyclopropyl-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide.

[0214] This compound was prepared starting from 3-cyclopropyl-*N*-*o*-tolyl-1*H*-pyrrole-2-carboxamide (106 mg, 0.44 mmol) and following the experimental procedure described in Preparation 1 b to afford 113 mg (93% yield) of the title compound.

LRMS (m/z): 256 (M+1)⁺.

PREPARATION 6**2-(Chloromethyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0215] This compound was prepared starting from 1-amino-3-cyclopropyl-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide (130 mg, 0.4 mmol) and following the experimental procedure described in Preparation 2 to afford 107 mg (73% yield) of the title compound.

LRMS (m/z): 314 (M+1)⁺.

PREPARATION 7**1-Amino-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide****a) *N*-o-Tolyl-1*H*-pyrrole-2-carboxamide**

[0216] 15.0 g (135 mmol) of 1*H*-pyrrol-2-carboxylic acid (purchased from Aldrich®, cat. no. P7,360-9) were suspended in a mixture of DMF (1.2 mL) and dichloromethane (150 mL). To this solution, 18 mL (207 mmol) of oxalyl chloride in dichloromethane (105 mL) were added dropwise over 30 minutes. The reaction was stirred two hours and then concentrated under reduced pressure to dryness.

[0217] The residual black oil was redissolved in dichloromethane (150 mL) and a solution of 15.9 g (148 mmol) of *o*-toluidine in dichloromethane (16 mL) was added dropwise. The reaction was stirred overnight then the solution was washed with a saturated aqueous solution of sodium bicarbonate. The organic phase was concentrated in vacuum. The product was purified by flash chromatography (30% AcOEt in hexane) to give 15.45 g (100% yield) of the title compound.

LRMS (m/z): 201 (M+1)⁺.

b) 1-Amino-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide

[0218] Prepared from 14.25 g (71.2 mmol) of *N*-o-tolyl-1*H*-pyrrole-2-carboxamide following the experimental procedure described in preparation 1 b. The crude product was suspended in diisopropyl ether and sonicated and the solid was filtered and washed with diethylether to give 10.04 g (66% yield) of the title compound.

LRMS (m/z): 216 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 11.15 (s, 1H), 8.06 (d, *J* = 7.7 Hz, 1H), 7.26 - 7.15 (m, 2H), 7.02 (t, *J* = 6.8 Hz, 1H), 6.96 (t, *J* = 2.3 Hz, 1H), 6.79 - 6.72 (m, 3H), 6.05 (dd, *J* = 4.2, 2.7 Hz, 1H), 2.29 (s, 3H).

PREPARATION 8**2-(Chloromethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0219] 9.5 mL (119 mmol) of 2-chloroacetyl chloride were added to a suspension of 5.14 g (23.9 mmol) of 1-amino-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide in 188 mL of glacial acetic acid, and the mixture was stirred at 120 °C for 3.5 hours. Then, the reaction mixture was cooled down to room temperature and it was concentrated in vacuum. The residue obtained was dissolved in ethyl acetate, washed with a saturated aqueous solution of sodium bicarbonate, water and brine. It was dried over magnesium sulphate, filtered and concentrated in vacuum. 4.99 g (65% yield) of the title compound were obtained.

LRMS (m/z): 274 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.66-7.83 (m, 1H), 7.29 - 7.60 (m, 4H), 7.05 (d, *J* = 4.30 Hz, 1H), 6.59 - 6.78 (m, 1H), 4.31 (dd, *J* = 10.55 Hz, 2H), 2.10 (s, 3H).

PREPARATION 9**2-(Aminomethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0220] A solution of 450 mg (1.64 mmol) of 2-(chloromethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one in 20 mL of a 7M solution of ammonia in methanol was heated at 85 °C in a sealed tube overnight, then cooled to room temperature and concentrated in vacuum to afford the title 491 mg (100% yield) of the title compound.

LRMS (m/z): 255 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.65 - 7.75 (m, 1H), 7.32 - 7.52 (m, 4H), 7.03 (d, *J* = 3.91 Hz, 1H), 6.42 - 6.97 (m, 3H), 2.09 (s, 3H).

PREPARATION 10**1-Amino-*N*-cyclohexyl-1*H*-pyrrole-2-carboxamide****a) *N*-Cyclohexyl-1*H*-pyrrole-2-carboxamide**

[0221] The mixture of methyl 1*H*-pyrrole-2-carboxylate (2.5 g, 20 mmol) and cyclohexylamine (13.8 mL, 120 mmol) was heated at 160°C in a sealed tub overnight with stirring. Next day, volatiles were removed under reduced pressure and the residue was taken-up with ethyl acetate and washed water, acidic water (pH=1) and brine. The organic layer was dried (Na₂SO₄) and concentrated in vacuum to give 1.88 g of a residue that was purified by flash chromatography silica (hexane/ethyl acetate) to obtain 0.88 g of the title compound (23% yield).
LRMS (m/z): 193(M+1)⁺.

b) 1-Amino-*N*-cyclohexyl-1*H*-pyrrole-2-carboxamide

[0222] This compound was prepared starting from *N*-cyclohexyl-1*H*-pyrrole-2-carboxamide (880 mg, 4.6 mmol) and following the experimental procedure described in Preparation 1 b to afford 700 mg (74% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 208 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.53 (d, *J* = 7.8 Hz, 1 H), 6.84 - 6.77 (m, 1 H), 6.61 (dd, *J* = 4.2, 2.0 Hz, 1 H), 6.55 (s, 2H), 5.91 (dd, *J* = 4.2, 2.6 Hz, 1 H), 3.80 - 3.64 (m, 1 H), 1.87 - 1.48 (m, 5H), 1.39 - 1.20 (m, 5H).

PREPARATION 11**2-(Chloromethyl)-3-cyclohexylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0223] To a solution of 1-amino-*N*-cyclohexyl-1*H*-pyrrole-2-carboxamide (160 mg, 0.77 mmol) in 1,4-dioxane (10 mL) was added 77 μL (0.97 mmol) of chloroacetyl chloride and the reaction mixture was stirred at 100°C during 1 h. Afterwards, the mixture was concentrated to dryness and the residue was dissolved in phosphorous oxychloride (3 mL) and heated to 50°C with stirring overnight. Next day, the cooled reaction mixture was poured slowly into an aqueous solution of potassium carbonate. The resulting mixture was extracted with ethyl acetate and the organic layer was washed with water (2x), dried (Na₂SO₄) and concentrated to afford 120 mg of a residue that was used in the next step without further purification.

LRMS (m/z): 266 (M+1)⁺.

PREPARATION 12**3-Methyl-1*H*-pyrrole-2-carboxylic acid**

[0224] 2.0 g (14.37 mmol) of methyl 3-methyl-1*H*-pyrrole-2-carboxylate (purchased from Otava Chemicals®, cat. no.1056278) were dissolved in 50 mL of methanol and 21.5 mL of a 2N aqueous solution of sodium hydroxide were added. The mixture was stirred at room temperature overnight and then at 60 °C for 20 hours. Then the methanol was evaporated in vacuum and the remaining aqueous solution was neutralized with 21.5 mL of a 2N solution of hydrochloric acid. The product was extracted with a 95:5 mixture of chloroform/methanol and the organic phase was washed with brine, dried over magnesium sulphate, filtered and evaporated under vacuum. 1.38 g (77% yield) of the title compound was obtained as a brown solid.

LRMS (m/z): 126 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.06 (s, 1H), 11.30 (s, 1H), 6.84 - 6.75 (m, 1H), 6.02 - 5.93 (m, 1 H), 2.24 (s, 3H).

PREPARATION 13**1-Amino-3-methyl-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide****a) 3-Methyl-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide**

[0225] 2.0 g (15.98 mmol) of 3-methyl-1*H*-pyrrole-2-carboxylic acid were dissolved in 50 mL of dichloromethane. 5.60 mL (63.9 mmol) of oxalyl chloride were added, followed by 5 drops of DMF. The reaction mixture was stirred at room temperature for 2 hours and then the solvents were evaporated. The black oil residue was dissolved in 50 mL of

dichloromethane and 6.85 g (63.9 mmol) of o-toluidine were added dropwise. The reaction was stirred at room temperature for 1 hour and then the solvent was evaporated. The crude product was purified by flash chromatography (dichloromethane to dichloromethane/methanol 95:5) and then triturated in diisopropylether to give 2.97 g (87% yield) of the title compound as a brown solid.

LRMS (m/z): 215 (M+1)⁺.

b) 1-Amino-3-methyl-N-o-tolyl-1H-pyrrole-2-carboxamide

[0226] 3-Methyl-N-o-tolyl-1H-pyrrole-2-carboxamide (1.5 g, 7.00 mmol) was dissolved in 60 mL of DMF. 294 mg (7.35 mmol) of sodium hydride (60 wt% dispersion in mineral oil) were added and the reaction mixture was stirred at room temperature for 1 hour. Then O-(mesitylsulfonyl)hydroxylamine (1.658 g, 7.70 mmol) was added and the reaction mixture was stirred for 30 minutes. The solvent was evaporated to dryness and the crude product was purified by flash chromatography (0 to 50% heptane/AcOEt) to give 1.17 g (73% yield) of the title compound as a yellow solid.

LRMS (m/z): 230 (M+1)⁺.

PREPARATION 14

2-(Chloromethyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0227] 140 mg (0.61 mmol) of 1-amino-3-methyl-N-o-tolyl-1H-pyrrole-2-carboxamide were dissolved in 3 mL of acetic acid. Chloroacetyl chloride (0.245 mL, 3.05 mmol) was added under vigorous stirring and the reaction mixture was then heated at 120 °C for 30 minutes. Then the mixture was allowed to cool to room temperature, poured into a mixture of water/ice and extracted twice with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium bicarbonate, dried over magnesium sulphate, filtered and evaporated to dryness. The crude product was purified by flash chromatography (0 to 40% heptane/ethyl acetate) to give 125 mg (71% yield) of the title compound as a white solid.

LRMS (m/z): 288 (M+1)⁺.

PREPARATION 15

2,2,2-Trichloro-1-(4-methyl-1H-pyrrol-2-yl)ethanone

[0228] To a solution of 2,2,2-trichloroacetyl chloride (5.05 mL, 45.3 mmol) in dry diethyl ether (12 mL) was slowly added a solution of 3-methyl-1H-pyrrole (3.15 g, 39.37 mmol) in 30 mL of dry diethyl ether during 1 h 15 min. Once the addition was finished, the reaction mixture was stirred at 45°C during 1 hour 30 minutes more. Next, more diethyl ether was added and the organic phase was washed with an aqueous solution of potassium carbonate to neutralize de media, water and brine. The organic phase was dried (Na₂SO₄) and concentrated to dryness to give a residue that was purified by flash chromatography silica (hexane/dichloromethane) to afford 4.7 g of the title compound (55% yield).

LRMS (m/z): 224 (M-1)⁻.

PREPARATION 16

1-Amino-4-methyl-N-o-tolyl-1H-pyrrole-2-carboxamide

a) 4-Methyl-N-o-tolyl-1H-pyrrole-2-carboxamide

[0229] A solution of 2,2,2-trichloro-1-(4-methyl-1H-pyrrol-2-yl)ethanone (2 g, 8.8 mmol) in a mixture of o-toluidine (1.76 mL, 16.5 mmol) and triethylamine (2.1 mL, 15.07 mmol) under argon was heated to 80°C with stirring during 75 h. Afterwards, the reaction mixture was concentrated to dryness giving a residue that was treated with hexane and the resulting solid filtered to afford 848 mg of the title compound (45% yield).

LRMS (m/z): 215 (M+1)⁺.

b) 1-Amino-4-methyl-N-o-tolyl-1H-pyrrole-2-carboxamide

[0230] This compound was prepared starting from 4-methyl-N-o-tolyl-1H-pyrrole-2-carboxamide (845 mg, 3.9 mmol) and following the experimental procedure described in Preparation 1 b to afford 516 mg (54% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 230 (M+1)⁺.

PREPARATION 17**2-(Chloromethyl)-6-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0231] 465 mg (2.03 mmol) of 1-amino-4-methyl-*N*-o-tolyl-1*H*-pyrrole-2-carboxamide were dissolved in 10 mL of acetic acid and 759 μ L (9.53 mmol) of chloroacetyl chloride were added. The mixture was stirred at 120 °C for 4 hours and the solvent was removed under vacuum. The residue was dissolved in AcOEt and washed with a saturated aqueous solution of sodium bicarbonate and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. 660 mg (69% yield) of the title compound were obtained as a beige solid.

LRMS (m/z): 288 (M+1)⁺.

PREPARATION 18**1-Amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide****a) *N*-Phenyl-1*H*-pyrrole-2-carboxamide**

[0232] Prepared following the experimental method described in preparation 1a starting from 10.0 g (90.0 mmol) of 1*H*-pyrrole-2-carboxylic acid (purchased from Aldrich®, cat. no. P7,360-9) and 9.22 g (99.0 mmol) of aniline. 13.0 g (78% yield) of the title compound were obtained as a brownish solid.

LRMS (m/z): 187 (M+1)⁺.

b) 1-Amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide

[0233] The title compound was prepared from 12.9 g (69.8 mmol) of *N*-phenyl-1*H*-pyrrole-2-carboxamide following the experimental procedure described in preparation 1 b. 10.3 g (73% yield) of the title compound were obtained as a solid.

LRMS (m/z): 202 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 10.74 (s, 1 H), 7.67 (d, *J* = 7.7 Hz, 2H), 7.33 (t, *J* = 7.9 Hz, 2H), 7.06 (t, *J* = 7.4 Hz, 1 H), 6.95 (t, *J* = 2.2 Hz, 1 H), 6.83 (dd, *J* = 4.2, 1.9 Hz, 1 H), 6.64 (s, 2H), 6.02 (dd, *J* = 4.2, 2.7 Hz, 1 H).

PREPARATION 19**2-(1-Chloroethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0234] 2.50 mL (25.75 mmol) of 2-chloropropanoyl chloride was added to a solution of 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide in glacial acetic acid and it was stirred at room temperature for 1 hour. Then it was concentrated in vacuum and the remaining acetic acid was co-evaporated with cyclohexane. 20 mL (218 mmol) of phosphorous oxychloride were added to the residue obtained and the resulting solution was heated at 125 °C. After 10 hours it was allowed to cool down to room temperature and the reaction mixture was carefully poured into a cold over-saturated aqueous solution of sodium bicarbonate. The aqueous solution was extracted with ethyl acetate, and the organic phases were collected together and washed with brine and then concentrated in vacuum. The residue obtained was dissolved in a mixture of hexane and ethyl acetate (10:1), filtered through silica and concentrated in vacuum. This residue was dissolved in 4 mL of a 7M solution of ammonia in methanol and then was heated to 60 °C overnight. The reaction mixture was concentrated in vacuum to obtain 75 mg (6% yield) of the title compound.

LRMS (m/z): 274 (M+1)⁺.

PREPARATION 20**(*S*)-2-(1-Aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (*S*)-*tert*-Butyl 1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate**

[0235] 2.0 g (9.94 mmol) of 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide, 2.45 g (12.05 mmol) of (*S*)-2-(*tert*-butoxycarbonylamino)butanoic acid (purchased from Aldrich®, cat. no. 15533) and 1.90 g (12.24 mmol) of EDC·HCl were dissolved in a mixture of 90 mL of THF and 30 mL of dichloromethane. The resulting solution was heated at 55 °C overnight. Then the solvents were evaporated and the crude residue was taken up in dichloromethane and washed with an aqueous solution of sodium bicarbonate and brine. The organic layer was dried over magnesium sulphate, filtered and the solvent was evaporated. The solid obtained was triturated in diethyl ether to furnish 2.47 g (64% yield) of the

title compound.

LRMS (m/z): 387 (M+1)⁺.

b) (S)-2-(1-Aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0236] 1.0 g (2.59 mmol) of (S)-*tert*-butyl 1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate were suspended in 13 mL of phosphorous oxychloride and heated to 80 °C for 6 hours. Then the excess reagent was removed under vacuum and the residue was taken up in AcOEt and treated with an aqueous solution of sodium bicarbonate. The two layers were separated and the aqueous phase was extracted with more AcOEt. The combined organic extracts were washed with brine, dried and the solvent was evaporated to give 1.2 g of a black oil. This intermediate was purified by flash chromatography (0-100% AcOEt in hexane) to give 421 mg of a yellow syrup that was treated with 50 mL of a 7M methanolic solution of ammonia at 80 °C in a sealed vessel. The solvent was then evaporated and the final product was purified by flash chromatography (0-100% AcOEt in hexane) to give 125 mg (11% yield) of the title compound.

LRMS (m/z): 269 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.23 (s, 1H), 7.71 - 7.37 (m, 5H), 6.94 (dd, J = 4.3, 1.7 Hz, 1 H), 6.60 (dd, J = 4.3, 2.7 Hz, 1 H), 3.14 (dd, J = 7.5, 5.6 Hz, 1 H), 1.81 - 1.64 (m, 1 H), 1.48 - 1.32 (m, 1 H), 0.72 (t, J = 7.4 Hz, 3H).

PREPARATION 21

2-(1-Aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) *tert*-Butyl 1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate

[0237] The title compound was prepared following the experimental procedure described in preparation 20a from 1.50 g (7.45 mmol) of 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide and 1.82 g (8.95 mmol) of racemic 2-(*tert*-butoxycarbonylamino)butanoic acid (purchased from ABCR, cat. no. AB154485). After recrystallisation of the crude product in diethylether, 2.43 g (84% yield) of the title compound were obtained.

LRMS (m/z): 387 (M+1)⁺.

b) 2-(1-Aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0238] 1.0 g (2.59 mmol) of *tert*-butyl 1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate were suspended in 16 mL of phosphorous oxychloride and heated to 80 °C for 6 hours. Then the mixture was evaporated to dryness under vacuum the residue was taken up in chloroform and treated with an aqueous solution of sodium bicarbonate. The two layers were separated and the aqueous phase was extracted with more chloroform. The combined organic extracts were washed with brine, dried and the solvent was evaporated. This intermediate was treated with a solution of 0.72 g (5.18 mmol) of potassium carbonate in DMF at 60 °C for 2.5 hours. Then the mixture was evaporated to dryness and the crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound (70 mg, 10% yield) as a solid.

LRMS (m/z): 269 (M+1)⁺.

PREPARATION 22

(S)-2-(1-Aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (S)-*tert*-Butyl 1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate

[0239] 2.00 g (9.94 mmol) of 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide were dissolved in 50 mL of DMF. To this solution, 2.07 g (10.94 mmol) of (S)-2-(*tert*-butoxycarbonylamino)propanoic acid (purchased from Aldrich®, cat. no. 13,451-1) and 2.10 g (10.95 mmol) of EDC-HCl were added and the resulting reaction mixture was stirred at room temperature overnight. The solvent was then evaporated under vacuum, the residue was taken up in ethyl acetate and the organic solution was washed with an aqueous solution of sodium bicarbonate and brine, it was dried over magnesium sulphate, filtered and the solvent was evaporated. The product was purified by flash chromatography (0-5% methanol in dichloromethane). 2.21 g (60% yield) of the final product were obtained as a white solid.

LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 10.11 (s, 1 H), 7.74 (s, 1 H), 7.55 - 7.48 (m, 2H), 7.36 - 7.28 (m, 2H), 7.15 - 7.07 (m, 1H),

7.03 (dd, $J = 2.9, 1.7$ Hz, 1H), 6.67 (dd, $J = 4.3, 1.7$ Hz, 1H), 6.15 (dd, $J = 4.2, 2.9$ Hz, 1H), 5.08 (d, $J = 7.5$ Hz, 1H), 4.40 (s, 1 H), 1.46 (s, 9H), 1.44 (d, $J = 7.1$ Hz, 3H).

b) (S)-2-(1-Aminoethyl)-3-phenylpyrrolo[1,2- η][1,2,4]triazin-4(3H)-one

[0240] 2.21 g (5.93 mmol) of (S)-*tert*-butyl 1-oxo-1-(2-(phenylcarbamoyl)-1H-pyrrol-1-ylamino)propan-2-ylcarbamate were treated with 27 mL of phosphorous oxychloride at 80 °C for 6 hours and then it was evaporated under vacuum until a dark solid formed. This residue was dissolved with chloroform and then treated with an aqueous solution of sodium bicarbonate. After stirring the mixture for 1 hour, the two layers were separated and the organic phase was washed with water and brine, dried over magnesium sulphate and the solvent was evaporated under vacuum. The residue was then treated in a sealed vessel with 30 mL of a 7M methanolic solution of ammonia at 80 °C overnight. The solvent was then evaporated and the product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound (350 mg, 23%) as a white solid.

LRMS (m/z): 255 ($M+1$)⁺.

¹H NMR (400 MHz, CDCl₃) δ 7.58 - 7.49 (m, 3H), 7.41 (dd, $J = 2.6, 1.7$ Hz, 1 H), 7.32 - 7.26 (m, 2H), 7.07 (dd, $J = 4.3, 1.7$ Hz, 1 H), 6.56 (dd, $J = 4.3, 2.7$ Hz, 1 H), 3.67 (q, $J = 6.6$ Hz, 1 H), 1.30 (d, $J = 6.6$ Hz, 3H).

PREPARATION 23

1-Amino-3-methyl-N-phenyl-1H-pyrrole-2-carboxamide

a) 3-Methyl-N-phenyl-1H-pyrrole-2-carboxamide

[0241] Prepared following the experimental method described in preparation 1a starting from 1.38 g (11.3 mmol) of 3-methyl-1H-pyrrole-2-carboxylic acid and 1.13 g (12.13 mmol) of aniline. After purification by flash chromatography (0-40% AcOEt in hexane), 1.08 g (49% yield) of the title compound were obtained as a white solid.

LRMS (m/z): 202 ($M+1$)⁺.

b) 1-Amino-3-methyl-N-phenyl-1H-pyrrole-2-carboxamide

[0242] The title compound was prepared from 1.08 g (5.39 mmol) of 3-methyl-N-phenyl-1H-pyrrole-2-carboxamide following the experimental procedure described in preparation 1 b. 1.16 g (47% yield) of the title compound was obtained as a solid.

LRMS (m/z): 216($M+1$)⁺.

¹H NMR (400 MHz, CDCl₃) δ 9.24 (s, 1 H), 7.63 - 7.56 (m, 2H), 7.37 - 7.30 (m, 2H), 7.14 - 7.06 (m, 1 H), 6.79 (d, $J = 2.7$ Hz, 1 H), 5.94 (d, $J = 2.3$ Hz, 1 H), 5.49 (s, 2H), 2.44 (s, 3H).

PREPARATION 24

2-(1-Chloroethyl)-5-methyl-3-phenylpyrrolo[1,2- η][1,2,4]triazin-4(3H)-one

[0243] To a solution of 1.13 g (5.25 mmol) of 1-amino-3-methyl-N-phenyl-1H-pyrrole-2-carboxamide in 45 mL of acetic acid were added 2.54 mL (26.21 mmol) of 2-chloropropanoyl chloride. The reaction was stirred at room temperature for 1 hour and then the solvent was removed in vacuum. The dark oil residue was treated with 20 mL of phosphorous oxychloride, and the mixture was heated to reflux for 16 hours and then it was evaporated under vacuum to dryness. The resulting residue was redissolved in dichloromethane and the organic solution was treated with a saturated aqueous solution of sodium bicarbonate. This mixture was vigorously stirred until the gas release stopped and the two layers were separated. The organic layer was washed with water and brine, dried over magnesium sulphate and the solvent was evaporated. The dark oil that resulted was treated with 40 mL of a 7M ammonia solution in methanol in a sealed vessel at 60 °C overnight. The solution was then evaporated to dryness and the product was purified by flash chromatography (0-5% methanol in dichloromethane) to furnish 0.28 g (18% yield) of the title compound as a white solid.

LRMS (m/z): 288 ($M+1$)⁺.

PREPARATION 25**2-(1-Iodoethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0244] 276 mg (0.96 mmol) of 2-(1-chloroethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one were dissolved in 10 mL of acetone and 287 mg (1.91 mmol) of sodium iodide were added. The reaction was stirred at 65 °C for 8 hours and at room temperature overnight. The solvent was evaporated to dryness under vacuum and the residue was dissolved in ethyl acetate. This solution was washed twice with water and brine, dried over magnesium sulphate and the solvent was evaporated. 340 mg (94% yield) of the title product were obtained as a brownish solid.

LRMS (m/z): 380 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 7.68 - 7.62 (m, 1H), 7.59 - 7.45 (m, 3H), 7.33 - 7.28 (m, 1H), 7.15 - 7.10 (m, 1H), 6.41 - 6.32 (m, 1H), 4.49 (q, J = 6.9 Hz, 1H), 2.50 (s, 3H), 2.16 (d, J = 7.0 Hz, 3H).

PREPARATION 26**Methyl 1-(phenylsulfonyl)-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxylate****a) Methyl 3-iodo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate**

[0245] A solution of methyl 3-bromo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate² (2.0 g, 5.8 mmol), sodium iodide (3.5 g, 23 mmol), *trans*-1,2-bis(methylamino)cyclohexane (0.83 g, 5.84 mmol) and copper iodide (0.55 g, 2.9 mmol) in 1,4-dioxane (23 mL) was stirred under reflux during 3 days. At the end of this period, the crude was allowed to reach room temperature and filtered through Celite® washing with ethyl acetate. The filtrate was concentrated to dryness, suspended in 80 mL of HCl 1 N and extracted with ethyl acetate (3x). The organic mixture was washed with water and brine, dried (Mg₂SO₄) and concentrated to give an oily residue that was purified by flash chromatography silica (hexane/ethyl acetate). Concentration of the fractions containing the compound afforded 1.72 g (50%) of the title compound. LRMS (m/z): 392 (M+1)⁺.

b) Methyl 1-(phenylsulfonyl)-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxylate

[0246] In a schlenk flask were placed methyl 3-iodo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate (2.44 g, 6.24 mmol) and copper iodide (1.46 g, 7.7 mmol) and it was established an inert atmosphere doing 3 cycles of vacuum-Ar. Subsequently, were added dimethyl formamide (44 mL) as solvent, hexamethylphosphoramide (HMPA) (5.4 mL, 31 mmol) and methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (3.9 mL, 30.7 mmol) and the reaction mixture was heated to 80 °C during 24 hours. Having consumed the starting material, the crude was poured into abundant ice-water and extracted with ethyl acetate (3x). The organic mixture was washed with water (2x) and brine, dried (Na₂SO₄) and concentrated in vacuum to give 2.58 g of a residue that was purified by flash chromatography silica (hexane/ethyl ether). Concentration of the fractions containing the compound afforded 370 mg (18%) of the title compound.

LRMS (m/z): 334 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.07 - 7.99 (m, 2H), 7.69 (m, 1H), 7.62 - 7.55 (m, 2H), 7.48 (d, J = 3.4 Hz, 1H), 6.51 (d, J = 3.4 Hz, 1H), 3.91 (s, 3H).

PREPARATION 27**1-Amino-*N*-o-tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide****a) 1-(Phenylsulfonyl)-*N*-o-tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide**

[0247] In a three-necked round-bottom flask *o*-toluidine (0.53 g, 5 mmol) was dissolved in 15 mL of toluene under inert atmosphere. To this solution was added trimethyl aluminium (2.5 mL, 5 mmol) and the mixture was stirred at room temperature during 10 minutes. Afterwards, a solution of methyl 1-(phenylsulfonyl)-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxylate (207 mg, 0.62 mmol) in 15 mL of toluene were added and the reaction mixture was heated at 80 °C over night. Next, the mixture was allowed to cool to room temperature and 2-3 mL of water were added to hydrolyze unreacted trimethyl aluminium and a 0.5M aqueous solution of disodium tartrate dihydrate were added stirring for a while. Afterwards, the two layers were separated and the aqueous phase was extracted with ethyl acetate. The organic mixture was washed with the same 0.5M aqueous solution of disodium tartrate dihydrate (25 mL), water and brine, dried and concentrated in vacuum to afford 560 mg of a residue that was used in the following step without further purification.

LRMS (m/z): 409 (M+1)⁺.

b) *N*-o-Tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide

[0248] To a solution of 1-(phenylsulfonyl)-*N*-o-tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide (560 mg of crude material) in 12 mL of methanol was added 2 mL of an aqueous 1N solution of sodium hydroxide and the mixture was stirred at room temperature during 1 h. At the end of this period, no starting material was detected and the reaction was elaborated in the following way: methanol was evaporated and water was added basifying the mixture with saturated aqueous solution of potassium carbonate. This mixture was extracted with ethyl acetate and this organic phase was washed with water and brine, dried (Na₂SO₄) and concentrated to dryness affording a reddish residue. This residue was purified by flash chromatography silica (hexane/ethyl ether). Concentration of the fractions containing the compound afforded 120 mg (50%) of the title compound.

LRMS (m/z): 269 (M+1)⁺.

c) 1-Amino-*N*-o-tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide

[0249] This compound was prepared starting from *N*-o-tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide (120 mg, 0.35 mmol) and following the experimental procedure described in Preparation 1b to afford 44 mg (31% purity, 14% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 284 (M+1)⁺.

PREPARATION 28**2-(Chloromethyl)-3-o-tolyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0250] This compound was prepared starting from 1-amino-*N*-o-tolyl-3-(trifluoromethyl)-1*H*-pyrrole-2-carboxamide (44 mg, 31% purity, 0.05 mmol) and following the experimental procedure described in Preparation 2 to afford 2 mg (67% purity, 8% yield) of the title compound that was used in the following step without further purification.

LRMS (m/z): 342 (M+1)⁺.

PREPARATION 29**1-Amino-3-chloro-1*H*-pyrrole-2-carboxylic acid****a) Benzyl 3-chloro-1*H*-pyrrole-2-carboxylate**

[0251] To a solution of 3-chloro-1*H*-pyrrole-2-carboxylic acid (15 g, 0.1 mol) in *N,N*-dimethylformamide (300 mL) and triethylamine (72 mL, 0.52 mmol) under argon atmosphere was added benzyl bromide (61 mL, 0.52 mmol) at 0-5°C and the reaction was stirred mechanically overnight at room temperature. Next day, the reaction mixture was concentrated in vacuum and the residue was suspended in 4% aqueous solution of sodium bicarbonate (300 mL) and extracted with ethyl acetate (2x250 mL). The organic layers were mixed and washed with more 4% aqueous solution of sodium bicarbonate, water and brine, and were dried (magnesium sulphate, MgSO₄) and concentrated under reduced pressure to give 21.4 g of a residue corresponding to the title compound (88% yield).

LRMS (m/z): 236 (M+1)⁺.

b) Benzyl 1-amino-3-chloro-1*H*-pyrrole-2-carboxylate

[0252] This compound was prepared starting from benzyl 3-chloro-1*H*-pyrrole-2-carboxylate (21.3 g, 0.09 mol) and following the experimental procedure described in Preparation 1 b to afford 22.19 g (92% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 251 (M+1)⁺.

c) 1-Amino-3-chloro-1*H*-pyrrole-2-carboxylic acid

[0253] To a solution of benzyl 1-amino-3-chloro-1*H*-pyrrole-2-carboxylate (20.56 g, 0.08 mol) in methanol (205 mL) was added 32.1 mL of a 3M methanolic solution of HCl (0.1 mol) and 7.7 g (0.07 mol) of 10% palladium on charcoal and this mixture was submitted to hydrogenation in a Parr apparatus working at 25 psi and room temperature during 21 hours. At the end of this period, the reaction was stopped filtering palladium catalyst through a pad of Celite® and the filtrate was concentrated in vacuum to give a light brown solid that was macerated with 40 mL of diethyl ether. This solid was filtrated and washed with petroleum ether to yield 13.32 g of the title compound with a purity of 55% (59% yield).

being the dehalogenated compound the main impurity. No further purifications were done proceeding to the next step.

LRMS (m/z): 159(M-1)⁻.

¹H NMR (400 MHz, DMSO) δ 7.51 (s, 2H), 7.05 (d, J = 2.9 Hz, 1 H), 6.10 (d, J = 2.9 Hz, 1 H).

PREPARATION 30

5-Chloro-2-(iodomethyl)-4*H*-pyrrolo[1,2-d][1,3,4]oxadiazin-4-one

a) 3-Chloro-1-(2-chloroacetamido)-1*H*-pyrrole-2-carboxylic acid

[0254] To a solution of 1-amino-3-chloro-1*H*-pyrrole-2-carboxylic acid (12.51 g, 55% purity, 0.04 mol) in 500 mL of glacial acetic acid were added under argon 17 mL of chloroacetyl chloride (0.21 mol) and the mixture was stirred at room temperature during 1.5 hours. Afterwards, the reaction mixture was poured into ice-water and extracted with ethyl acetate (2x400 mL). The organic layers were washed with water and brine, dried (MgSO₄) and concentrated in vacuum giving a dark oil that was crystallized from a 1/1 diethyl ether/petroleum ether mixture. Filtration of the solid obtained afforded 7.28 g of a white solid corresponding to the title compound (61% yield, 85% purity). This solid was used in the next step without further purifications.

LRMS (m/z): 237 (M+1)⁺.

b) 5-Chloro-2-(chloromethyl)-4*H*-pyrrolo[1,2-d][1,3,4]oxadiazin-4-one

[0255] To a solution of 3-chloro-1-(2-chloroacetamido)-1*H*-pyrrole-2-carboxylic acid (7.22 g, 25.9 mmol) in 110 mL of 1,4-dioxane was added phosphorous oxychloride (23.6 mL, 259mmol) solved in 35 mL of 1,4-dioxane at room temperature. Then, the reaction mixture was stirred to reflux during 2 hours. At the end of this period, the reaction mixture was poured into 500 mL of a 4% aqueous solution of sodium bicarbonate and extracted with ethyl acetate (2x400 mL). The organic layers were mixed and washed with more 4% aqueous solution of sodium bicarbonate, water and brine, and were dried (MgSO₄) and concentrated under reduced pressure to give 6.97 g of a dark oil that was purified by flash chromatography silica (hexane/ethyl acetate). After purification were obtained 2.85 g of the title compound (57% yield). LRMS (m/z): 219 (M+1)⁺.

c) 5-Chloro-2-(iodomethyl)-4*H*-pyrrolo[1,2-d][1,3,4]oxadiazin-4-one

[0256] To a solution of 5-chloro-2-(chloromethyl)-4*H*-pyrrolo[1,2-d][1,3,4]oxadiazin-4-one (2.87 g, 13.1 mmol) in 57 mL of dry acetone under inert atmosphere was added 3.93 g (26.2 mmol) of sodium iodide and this mixture was stirred at room temperature overnight. Next day, the reaction mixture was poured into a 1/1 mixture of water/brine (100 mL) and extracted with diethyl ether (2x 75 mL). The organic layers were washed with water and brine, dried (MgSO₄) and concentrated in vacuum to afford 3.95 g of the title compound (95% yield).

LRMS (m/z): 311 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 7.26 (d, J = 3.0 Hz, 1 H), 6.52 (d, J = 3.0 Hz, 1 H), 4.10 (s, 2H).

PREPARATION 31

Di-*tert*-butyl 9-((5-chloro-4-oxo-4*H*-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9*H*-purin-6-ylimidodicarbonate

[0257] To a solution of 5-chloro-2-(iodomethyl)-4*H*-pyrrolo[1,2-d][1,3,4]oxadiazin-4-one (0.93 g, 3 mmol) in 18.6 mL of dry *N,N*-dimethylformamide under argon atmosphere were added di-*tert*-butyl 9*H*-purin-6-ylimidodicarbonate (1.41 g, 4.2 mmol) and sodium bicarbonate (0.35 g, 4.2 mmol) and the reaction mixture was stirred overnight at room temperature. At the end of this period, the reaction mixture was poured into 50 mL of a 4% aqueous solution of sodium bicarbonate and extracted with ethyl acetate (2x40 mL). The organic layers were mixed and washed with more 4% aqueous solution of sodium bicarbonate, water and brine, and were dried (MgSO₄) and concentrated under reduced pressure to give 1.73 g of a solid that was purified by flash chromatography silica (hexane/ethyl acetate). After purification were obtained 0.81 g of the title compound (51% yield).

LRMS (m/z): 517 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.89 (s, 1H), 8.23 (s, 1H), 7.16 (d, J = 3.0 Hz, 1 H), 6.49 (d, J = 3.0 Hz, 1 H), 5.37 (s, 2H), 1.48 (s, 18H).

PREPARATION 32**1*H*-Imidazole-2-carboxylic acid**

[0258] 21.75 mL (43.5 mmol) of a 2M solution of lithium hydroxide in water were added to a solution of 1.22 g (8.71 mmol) of ethyl 1*H*-imidazole-2-carboxylate (ref) in a mixture of tetrahydrofuran (20 mL) and water (20 mL). The reaction mixture was warmed up to reflux, and stirred for 1.5 hours. The reaction mixture was cooled to room temperature and the solvent was evaporated under vacuum. The crude residue (3.0 g) was used in next step without further purification. LRMS (m/z): 113(M+1)⁺.

PREPARATION 33**1-Amino-*N*-phenyl-1*H*-imidazole-2-carboxamide****a) *N*-Phenyl-1*H*-imidazole-2-carboxamide**

[0259] To a solution of 1*H*-imidazole-2-carboxylic acid (0.975 g, 8.7 mmol) in DMF (30 mL) were added aniline (0.67 mL, 8.7 mmol), EDC·HCl (2.54 g, 13.05 mmol) and HOBt (1.76 g, 13.05 mmol). The reaction mixture was stirred at room temperature for 21 hours. Then, it was poured into water and extracted with ethyl acetate. The combined organic layer was dried over sodium sulphate, filtered and concentrated. The crude residue was purified by flash chromatography (2% to 3% methanol in dichloromethane) to yield 1.55 g (96% yield) of the title compound.

[0260] LRMS (m/z): 188(M+1)⁺. **b) 1-Amino-*N*-phenyl-1*H*-imidazole-2-carboxamide**

[0261] 0.430 g (10.75 mmol) of sodium hydride (60 % dispersion in mineral oil) was added to a 0 °C cooled solution of *N*-phenyl-1*H*-imidazole-2-carboxamide (1.55 g, 8.27 mmol) in DMF (50 mL). The mixture was stirred at 0 °C for 30 minutes and 2.70 g (11.57 mmol) of DPPONH₂ (*P,P*-diphenylphosphinic amide, available from Sigma Aldrich®, cat. no. 5994-87-6) were added portionwise. A thick suspension formed and additional 100 mL of DMF were added. The mixture was stirred at room temperature for 3 hours and then it was poured into 150 mL of a saturated aqueous solution of sodium thiosulphate and extracted with ethyl acetate. The combined organic layer was dried over sodium sulphate, filtered and concentrated. The crude product was purified by flash chromatography (20% to 40% AcOEt/Hexanes) to yield 1.27 g (76% yield) of the title compound as a pale yellow solid. LRMS (m/z): 203 (M+1)⁺.

PREPARATION 34**2-(Chloromethyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0262] 62 mg (0.25 mmol) of pyridinium *p*-toluenesulfonate were added to a suspension of 500 mg (2.47 mmol) of 1-amino-*N*-phenyl-1*H*-imidazole-2-carboxamide in 3.3 mL of 2-chloro-1,1,1-trimethoxyethane. The mixture was stirred at 100 °C for 5 hours and the solvent was evaporated. The crude product was purified by flash chromatography (1% to 3% MeOH/DCM) to yield 0.227 g (35%) of the title compound as a beige solid. LRMS (m/z): 261 (M+1)⁺.

PREPARATION 35**1-Amino-*N*-o-tolyl-1*H*-imidazole-2-carboxamide****a) *N*-o-Tolyl-1*H*-imidazole-2-carboxamide**

[0263] To a solution of 1*H*-imidazole-2-carboxylic acid (0.52 g, 4.64 mmol) in DMF (20 mL) was added o-toluidine (0.50 mL, 4.64 mmol), EDC·HCl (1.35 g, 6.96 mmol) and HOBt (0.94 g, 6.96 mmol). The reaction mixture was stirred at room temperature for 16 hours. Then, it was poured into water and extracted with ethyl acetate. The combined organic layer was dried over sodium sulphate, filtered and concentrated to dryness. The crude product was purified by flash chromatography (2% to 3% MeOH/DCM) to yield 0.93 g (99%) of the title compound as a beige solid. LRMS (m/z): 202 (M+1)⁺.

b) 1-Amino-*N*-o-tolyl-1*H*-imidazole-2-carboxamide

[0264] 240 mg (6.02 mmol) of sodium hydride (60% dispersion in mineral oil) were added to a 0° C cooled solution of

N-*o*-tolyl-1*H*-imidazole-2-carboxamide (0.93 g, 4.63 mmol) in DMF (120 mL). The mixture was stirred at 0 °C for 30 minutes and DPPONH₂ (1.51 g, 6.49 mmol, available from Sigma Aldrich®, cat. no. 5994-87-6) was added portionwise. The mixture was stirred at room temperature for 2 hours and then it was poured into 200 mL of a saturated aqueous solution of sodium thiosulphate and extracted with ethyl acetate. The combined organic layer was dried over sodium sulphate, filtered and concentrated. The crude product was purified by flash chromatography (20% to 50% AcOEt/Hexanes) to yield 0.55 g (55%) of the title compound as a beige solid.
LRMS (m/z): 217 (M+1)⁺.

PREPARATION 36

2-(Chloromethyl)-3-*o*-tolylimidazo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0265] Pyridinium *p*-toluenesulfonate (0.067 g, 0.27 mmol) was added to a suspension of 1-amino-*N*-*o*-tolyl-1*H*-imidazole-2-carboxamide (0.58 g, 2.68 mmol) in 2-chloro-1,1,1-trimethoxyethane (3.62 mL). The mixture was stirred at 100 °C for 5 hours. The solvent was evaporated to dryness and the residue was purified by flash chromatography (10% to 50% AcOEt/hexanes) to yield 0.360 g (49% yield) of the title compound.
LRMS (m/z): 275 (M+1)⁺.

PREPARATION 37

1-Amino-*N*-(3-fluorophenyl)-1*H*-pyrrole-2-carboxamide

[0266] This compound was prepared starting from 1*H*-pyrrole-2-carboxylic acid (0.67 g, 6 mmol) and following the experimental procedure described in Preparation 1 to afford 0.525 g (39% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 220 (M+1)⁺.

PREPARATION 38

1-Amino-*N*-(3,5-difluorophenyl)-1*H*-pyrrole-2-carboxamide

[0267] This compound was prepared starting from 1*H*-pyrrole-2-carboxylic acid (0.67 g, 6 mmol) and following the experimental procedure described in Preparation 1 to afford 0.393 g (26% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 238 (M+1)⁺.

PREPARATION 39

(*S*)-2-(1-Aminoethyl)-3-(3-fluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0268] This compound was prepared starting from 1-amino-*N*-(3-fluorophenyl)-1*H*-pyrrole-2-carboxamide (525 mg, 2.39 mmol) and following the experimental procedure described in Preparation 22 to afford 42 mg (7% yield) of the title compound.
LRMS (m/z): 273 (M+1)⁺.

PREPARATION 40

(*S*)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0269] This compound was prepared starting from 1-Amino-*N*-(3,5-difluorophenyl)-1*H*-pyrrole-2-carboxamide (393 mg, 1.66 mmol) and following the experimental procedure described in Preparation 22 to afford 36 mg (7% yield) of the title compound.
LRMS (m/z): 291 (M+1)⁺.

PREPARATION 41**1-Amino-*N*-(pyridin-2-yl)-1*H*-pyrrole-2-carboxamide**

[0270] This compound was prepared starting from 1*H*-pyrrole-2-carboxylic acid (2 g, 18 mmol) and 2-aminopyridine (3.40 g, 36 mmol), following the experimental procedure described in Preparation 1 to afford 0.34 g (9% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 203 (M+1)⁺.

PREPARATION 42**(*S*)-2-(1-Aminoethyl)-3-(pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-n-one dihydrochloride****a) (*S*)-*tert*-Butyl 1-oxo-1-(2-(pyridin-2-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate**

[0271] This compound was prepared starting from 1-amino-*N*-(pyridin-2-yl)-1*H*-pyrrole-2-carboxamide (340 mg, 1.68 mmol) and following the experimental procedure described in Preparation 22a to afford 440 mg (56% yield) of the title compound of preparation 42a.

b) (*S*)-2-(1-Aminoethyl)-3-(pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one dihydrochloride

[0272] A solution of bromine (113 μ L, 2.21 mmol) in dichloromethane (2 mL) was added dropwise to a solution of triphenylphosphine (558 mg, 2.13 mmol) in dichloromethane (5 mL) under nitrogen. The solution was stirred for 30 min, and triethylamine (494 μ L, 3.51 mmol) and a solution of (*S*)-*tert*-butyl 1-oxo-1-(2-(pyridin-2-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate (440 mg, 0.94 mmol) in 3 mL of dichloromethane were added. The reaction mixture was stirred at room temperature for 3.5 h, and then, volatiles were removed under reduced pressure and the residue was triturated with toluene affording a solid that was removed by filtration. The filtrate was concentrated to dryness under reduced pressure and the residue was redissolved in 20 mL of a 7M methanolic solution of ammonia and stirred overnight at 80 °C in a sealed vessel. The solvent was then evaporated and the residue was purified by flash chromatography (0% to 50% AcOEt/hexanes) to yield 0.19 g (57% yield) of (*S*)-*tert*-butyl 1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate. This compound (190 mg, 0.43 mmol) was dissolved in 2 mL of dioxane and 2 mL of a 4M hydrogen chloride solution in dioxane were added. The mixture was stirred at room temperature overnight. Then, the solid present in the reaction media was filtered-off and washed with hexane to afford 127 mg of the title compound of preparation 42 (90% yield).

LRMS (m/z): 256 (M+1)⁺.

PREPARATION 43**(*S*)-3-Phenyl-2-(pyrrolidin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (*S*)-*tert*-Butyl 2-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylcarbamoyl)pyrrolidine-1-carboxylate**

[0273] (*S*)-1-(*tert*-Butoxycarbonyl)pyrrolidine-2-carboxylic acid (1.80 g, 8.36 mmol) were dissolved in 20 mL of *N,N*-dimethylformamide and HATU (3.40 g, 8.95 mmol), DIEA (2.60 mL, 14.89 mmol) and 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide were added. The resulting solution was stirred at room temperature overnight and the solvent was removed in vacuum. The residue was taken up in ethyl acetate and the organic solution was washed with water and brine, dried over magnesium sulphate, filtered and evaporated to dryness. The crude product was purified by flash chromatography (0% to 30% hexane/AcOEt) to yield 1.90 g (62%) of the title compound as a white solid.
LRMS (m/z): 399 (M+1)⁺.

b) (*S*)-*tert*-Butyl 2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

[0274] To a solution of 2.50 g (9.54 mmol) of triphenylphosphine in 20 mL of methylene chloride was added a solution of 1.52 g (9.54 mmol) of bromine in 10 mL of dichloromethane dropwise under nitrogen atmosphere. At the end of the addition the colourless solution was stirred for 5 minutes and then 3.32 mL (23.84 mmol) of triethylamine and 1.90 g (4.77 mmol) of (*S*)-*tert*-butyl 2-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylcarbamoyl)pyrrolidine-1-carboxylate were added. The reaction mixture was then stirred at reflux for 3 hours and then cooled and the solvent was evaporated. The residue was taken up in cold toluene and the insoluble salts were filtered. The filtrate was evaporated and the residue was then

redissolved in a mixture of tetrahydrofuran (30 ml) and methanol (10 ml) and 1.00 g (14.31 mmol) of sodium methanethiolate was added. The solution was stirred at 60 °C for 3 hours. Then the solvents were evaporated and the residue was partitioned between water and ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. The product was purified by flash chromatography (20% to 40% hexane/AcOEt) to yield 1.20 g (66%) of the title compound.
LRMS (m/z): 381 (M+1)⁺.

c) (S)-3-Phenyl-2-(pyrrolidin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0275] 1.20 g (3.16 mmol) of (S)-*tert*-butyl 2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate were dissolved in 2 ml of methylene chloride and 2 ml of trifluoroacetic acid were added. The resulting solution was stirred at room temperature for 1 hour and the reaction mixture was evaporated to dryness. The residue was then redissolved in dichloromethane and the solution was washed with an aqueous solution of sodium bicarbonate and brine, dried over magnesium sulphate, filtered and the solvent was removed under vacuum to yield 0.80 g (91%) of the title compound.
LRMS (m/z): 281 (M+1)⁺.
¹H NMR (400 MHz, DMSO-d₆) δ 7.56 (dd, 1H), 7.55 - 7.42 (m, 4H), 7.41 - 7.37 (m, 1H), 6.93 (dd, 1H), 6.57 (dd, 1H), 3.14 - 2.97 (m, 1H), 2.39 (dd, 2H), 2.07 - 1.93 (m, 1H), 2.00 (s, 1H), 1.91 - 1.78 (m, 1H), 1.69 (tt, 1H), 1.60 - 1.49 (m, 1H).

PREPARATION 44

1-Amino-3-bromo-N-phenyl-1H-pyrrole-2-carboxamide

a) 3-Bromo-N-phenyl-1-(phenylsulfonyl)-1H-pyrrole-2-carboxamide

[0276] In a three-necked round-bottom flask aniline (1.57 mL, 17.20 mmol) was dissolved in 80 mL of toluene under inert atmosphere. To this solution was added trimethyl aluminium (7.82 mL, 15.64 mmol) and the mixture was stirred at room temperature during 10 minutes. Afterwards, a solution methyl 3-bromo-1-(phenylsulfonyl)-1H-pyrrole-2-carboxylate (2.0 g, 5.81 mmol) in 20 mL of toluene were added and the reaction mixture was heated at 80 °C for 3h. Next, the mixture was allowed to cool to room temperature and 20-30 mL of water was added to hydrolyze unreacted trimethyl aluminium and a 0.5M aqueous solution of disodium tartrate dihydrate was also added. After stirring for a while, the two layers were separated and the aqueous phase was extracted with ethyl acetate. The organic mixture was washed with the same 0.5M aqueous solution of disodium tartrate dihydrate (200 mL), water and brine, dried and concentrated in vacuum to afford 2.7 g of a residue that was used in the following step without further purification.
LRMS (m/z): 405, 407 (M+1)⁺.

b) 3-Bromo-N-phenyl-1H-pyrrole-2-carboxamide

[0277] To a solution of 3-bromo-N-phenyl-1-(phenylsulfonyl)-1H-pyrrole-2-carboxamide (2.70 g of crude material) in 50 mL of methanol was added 15 mL of an aqueous 1 N solution of sodium hydroxide and the mixture was stirred at room temperature during 1.5 h. At the end of this period, no starting material was detected and the reaction was elaborated in the following way: methanol was evaporated and a precipitate was formed which was filtered off and washed several times with water. The solid was dried in the vacuum oven to afford 1.14g and used in the following step without further purification.
LRMS (m/z): 265, 267 (M+1)⁺.

c) 1-Amino-3-bromo-N-phenyl-1H-pyrrole-2-carboxamide

[0278] This compound was prepared starting from 3-bromo-N-phenyl-1H-pyrrole-2-carboxamide (1.11 g, 4.19 mmol) and following the experimental procedure described in Preparation 1 b to afford 0.78 g (67% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 280, 282 (M+1)⁺.

PREPARATION 45

(S)-tert-Butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**a) (S)-tert-Butyl 1-(3-bromo-2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0279] The title compound was prepared following the experimental procedure described in preparation 20a from 810 mg (2.89 mmol) of 1-amino-3-bromo-N-phenyl-1*H*-pyrrole-2-carboxamide and 656 mg (3.47 mmol) of (S)-2-(*tert*-butoxycarbonylamino)propanoic acid (purchased from Aldrich). The crude product was purified by flash chromatography in hexane/ethyl acetate to afford 670 mg (49% yield) of the title compound.

LRMS (m/z): 306, 308 (M+1)⁺.

b) (S)-tert-Butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0280] This compound was prepared starting from (S)-*tert*-butyl 1-(3-bromo-2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (670 mg, 1.48 mmol) and following the experimental procedure described in Preparation 42b. The residue was purified by flash chromatography in hexane/ethyl acetate to afford 500 mg (78% yield) of the title compound were obtained.

LRMS (m/z): 433, 435 (M+1)⁺.

PREPARATION 46

(S)-2-(1-Aminoethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (S)-tert-Butyl 1-(5-iodo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0281] A solution of (S)-*tert*-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (254 mg, 0.59 mmol), *trans*-1,2-bis(methylamino)cyclohexane (53 mg, 0.37 mmol) and copper iodide (50 mg, 0.26 mmol) in 1,4-dioxane (5 mL) was stirred at 120°C overnight. The crude was allowed to reach room temperature and filtered through Celite® washing with ethyl acetate. The filtrate was concentrated to dryness, suspended in 10 mL of HCl 1 N and extracted with ethyl acetate (3x). The organic mixture was washed with water and brine, dried (Mg₂SO₄) and concentrated to give 259mg (92% yield) of the title compound that was used without further purification.

LRMS (m/z): 481 (M+1)⁺.

b) (S)-tert-Butyl 1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0282] The title compound was prepared from (S)-*tert*-butyl 1-(5-iodo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (259 mg, 0.47 mmol) following the experimental procedure described in Preparation 26b. 185mg (98% yield) of the desired compound were obtained.

LRMS (m/z): 423 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0283] (S)-*tert*-Butyl 1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (180mg, 0.43mmol) was dissolved in 2ml of dioxane and 2 ml of a 4M hydrogen chloride solution were added. The mixture was stirred at room temperature overnight. Reaction mixture was then concentrated and the residue dissolved in ethyl acetate and a 2N solution of NaOH. The organic layer was separated and the aqueous layer extracted with more ethyl acetate. The combined organic layer was dried over magnesium sulphate and concentrated to dryness. The title compound was obtained (146mg, 97% yield) as colourless oil.

LRMS (m/z): 323 (M+1)⁺.

PREPARATION 47**(S)-2-(1-Aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile****a) (S)-*tert*-butyl 1-(5-cyano-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0284] A mixture of methyl (S)-*tert*-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (238 mg, 0.55 mmol), dicyanozinc (129 mg, 1.10 mmol) and Tetrakis(triphenylphosphine)palladium(0) (64 mg, 0.06 mmol) in DMF, was heated at 120°C in a sealed tub overnight with stirring. Next day, ethyl acetate was added and filtered through Celite®. Phases were separated and the aqueous layer extracted with more ethyl acetate. The combined organic layer was dried (Na₂SO₄) and concentrated in vacuum to give 580 mg of a residue that was purified by flash chromatography silica (hexane/ethyl acetate) to obtain 119 mg (57% yield) of the title compound.
LRMS (m/z): 380 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0285] The title compound was prepared from (S)-*tert*-butyl 1-(5-cyano-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (119 mg, 0.31 mmol) following the experimental procedure described in Preparation 46c. 67 mg (77% yield) of the desired compound were obtained.
LRMS (m/z): 280 (M+1)⁺.

PREPARATION 48**1-(4-Methoxybenzyl)-4-nitro-1H-pyrazole**

[0286] 4-Nitro-1H-pyrazole (1 g, 8.84 mmol) was dissolved in dimethylformamide (40 ml). Potassium carbonate (1.2 g, 8.68 mmol) and 1-(chloromethyl)-4-methoxybenzene (1.2 g, 8.96 mmol) were added and the mixture was heated with stirring at 80 °C for 3h. Once at room temperature, the reaction mixture was poured onto water and extracted with ethyl acetate (x3). The organic phase was washed with water and brine, dried over magnesium sulphate, filtered and the solvent evaporated under reduced pressure to yield 2.27g of the final compound as an oil. The product was used in the next synthetic step without further purification.
LRMS (m/z): 234 (M+1)⁺.

PREPARATION 49**1-(4-Methoxybenzyl)-1H-pyrazol-4-amine**

[0287] 1-(4-Methoxybenzyl)-4-nitro-1H-pyrazole (2.17 g, 9.30 mmol) was dissolved in ethanol (20 ml). Ammonium chloride (50 mg, 0.93 mmol) and iron (powder, 2.60g, 46.56 mmol) were added and the mixture was refluxed for 1 h. The reaction mixture was filtered through Celite® and the solvent was evaporated to dryness under reduced pressure. Usual work-up with water and ethyl acetate afforded 1.77g (94% yield) of the title compound as an oil, which was used in the next synthetic step without further purification.
LRMS (m/z): 204 (M+1)⁺.

PREPARATION 50**(S)-Methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate**

[0288] The title compound was prepared following the experimental procedure described in preparation 20a from 6.1 g (18.38 mmol) of methyl 1-amino-3-bromo-1H-pyrrole-2-carboxylate and 3.5 g (28.50 mmol) of (S)-2-(*tert*-butoxycarbonylamino)propanoic acid (purchased from Aldrich). The crude product was purified by flash chromatography in hexane/ethyl acetate (0 to 25%) to afford 4.34 g (61% yield) of the title compound.
LRMS (m/z): 389, 391 (M+1)⁺.

PREPARATION 51

(S)-tert-Butyl 1-(3-bromo-2-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)carbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0289] This compound was prepared starting from (S)-methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (300 mg, 0.77 mmol) and 1-(4-methoxybenzyl)-1H-pyrazol-4-amine (469 mg, 2.31 mmol) following the experimental procedure described in Preparation 27a to afford 271 mg (96% purity, 60% yield) of the title compound after purification by flash chromatography (0% to 70% AcOEt/hexanes).

LRMS (m/z): 562 (M+1)⁺.

PREPARATION 52

(S)-tert-Butyl 1-(5-bromo-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0290] Bromine (35 μ L, 0.68 mmol) was added dropwise to a solution of triphenylphosphine (177 mg, 0.67 mmol) in dichloromethane (3 mL) under nitrogen. The solution was stirred for 30 min, and triethylamine (269 μ L, 1.93 mmol) and a solution of (S)-*tert*-butyl 1-(3-bromo-2-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)carbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (271 mg, 0.48 mmol) in 7 mL of dichloromethane were added. The reaction mixture was stirred at 60 °C for 1.5 h, and then, volatiles were removed under reduced pressure and the residue was redissolved in 10 mL of a 7M methanolic solution of ammonia and stirred overnight at 80 °C in a sealed vessel. The solvent was then evaporated and the residue was purified by flash chromatography (0% to 50% AcOEt/hexanes) to yield 154 mg (59% yield) of the title compound.

LRMS (m/z): 544 (M+1)⁺.

PREPARATION 53

(S)-2-(1-Aminoethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**a) (S)-tert-Butyl 1-(5-iodo-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0291] A solution of (S)-*tert*-butyl 1-(5-bromo-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (154 mg, 0.28 mmol), *trans*-1,2-bis(methylamino)cyclohexane (24 mg, 0.17 mmol) and copper iodide (27 mg, 0.14 mmol) in 1,4-dioxane (15 mL) was stirred in a sealed tube at 120 °C overnight. The crude was allowed to reach room temperature and filtered through Celite® washing with ethyl acetate. The filtrate was concentrated to dryness, suspended in 10 mL of HCl 1N and extracted with ethyl acetate (3x). The organic mixture was washed with water and brine, dried (Mg₂SO₄) and concentrated to give 131 mg (69% yield) of the title compound that was used without further purification.

LRMS (m/z): 591 (M+1)⁺.

b) (S)-tert-Butyl 1-(5-cyano-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0292] In a sealed tube provided with magnetic stirring were placed (S)-*tert*-butyl 1-(5-iodo-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (131 mg, 0.20 mmol) and copper cyanide (210 mg, 2.34 mmol). Subsequently, pyridine was added (10 mL) as solvent, and the reaction mixture was heated to 120 °C during 18 hours in a microwave apparatus. Having consumed the starting material, the crude was filtered through Celite® and washed with ethyl acetate. The organic mixture was washed with water (2x) and brine, dried (Na₂SO₄) and concentrated in vacuum to give 132 mg of a residue that was used in the next synthetic step without further purification.

LRMS (m/z): 490 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0293] The title compound was prepared from (S)-*tert*-butyl 1-(5-cyano-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-

oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (132 mg, 0.27 mmol) following the experimental procedure described in Preparation 46c. 90mg (81% yield) of the desired compound were obtained.

LRMS (m/z): 390 (M+1)⁺.

PREPARATION 54

tert-Butyl (S)-1-(5-bromo-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

a) tert-Butyl (S)-1-(3-bromo-2-((R)-1-phenylethylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0294] The title compound was prepared from (S)-methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (1.27 g, 3.25 mmol) following the experimental procedure described in Preparation 27a. 600 mg (39% yield) of the desired compound were obtained.

LRMS (m/z): 480 (M+1)⁺.

b) tert-Butyl (S)-1-(5-bromo-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0295] Bromine (90 μ L, 1.76 mmol) was added dropwise to a solution of triphenylphosphine (460 mg, 1.75 mmol) in dichloromethane (4 mL) under nitrogen. The solution was stirred for 30 min, and triethylamine (700 μ L, 7.02 mmol) and a solution of *tert*-butyl (S)-1-(3-bromo-2-((R)-1-phenylethylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (0.60 g, 1.25 mmol) in 10 ml of dichloromethane were added. The reaction mixture was stirred at 60 °C for 2 h, poured onto 4% sodium bicarbonate solution and extracted with dichloromethane. The organic phase is passed through a phase separator and the solvent removed under reduced pressure. The residue was redissolved in DMF (10 ml) and sodium thiomethoxide (263 mg, 3.75 mmol) was added. The reaction mixture was stirred for 2h, poured onto 4% sodium bicarbonate solution and extracted with ethyl acetate (x3). The organic mixture was washed with water and brine, dried (Mg₂SO₄) and concentrated to give 1.18g of a crude which was purified by flash chromatography (0% to 20% AcOEt/hexanes) to yield 470 mg (81% yield) of the title compound.

LRMS (m/z): 462 (M+1)⁺.

PREPARATION 55

2-((S)-1-Aminoethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

a) tert-Butyl (S)-1-(5-iodo-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0296] The title compound was prepared from *tert*-butyl (S)-1-(5-bromo-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (470 mg, 1 mmol) following the experimental procedure described in Preparation 53a. 507 mg (90% yield) of the desired compound were obtained.

LRMS (m/z): 509 (M+1)⁺.

b) tert-Butyl (S)-1-(5-cyano-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0297] The title compound was prepared from *tert*-butyl (S)-1-(5-iodo-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (0.51 g, 1 mmol) following the experimental procedure described in Preparation 53b. 410 mg (100% yield) of the desired compound were obtained.

LRMS (m/z): 408 (M+1)⁺.

c) 2-((S)-1-Aminoethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0298] The title compound was prepared from *tert*-butyl (S)-1-(5-cyano-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (498 mg, 1.22 mmol) following the experimental procedure described in Preparation 46c. 241 mg (53% yield) of the desired compound were obtained.

LRMS (m/z): 308 (M+1)⁺.

PREPARATION 56**(S)-2-(1-Aminoethyl)-3-(5-methyl-1H-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one****a) (S)-*tert*-Butyl 1-(2-(5-methyl-1H-pyrazol-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0299] The title compound was prepared from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (900 mg, 2.89 mmol) following the experimental procedure described in Preparation 27a. 188 mg (15% yield) of the desired compound were obtained.

LRMS (m/z): 377 (M+1)⁺.

b) (S)-*tert*-Butyl 1-(3-(5-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0300] The title compound was prepared from (S)-*tert*-butyl 1-(2-(5-methyl-1H-pyrazol-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (180 mg, 0.41 mmol) following the experimental procedure described in Preparation 54b. 64 mg (44% yield) of the desired compound were obtained.

LRMS (m/z): 359 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(5-methyl-1H-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0301] The title compound was prepared from (S)-*tert*-butyl 1-(3-(5-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (64 mg, 0.18 mmol) following the experimental procedure described in Preparation 46c. 46 mg (100% yield) of the desired compound were obtained.

LRMS (m/z): 259 (M+1)⁺.

PREPARATION 57**(S)-2-(1-Aminoethyl)-3-(1H-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one****a) (S)-*tert*-Butyl 1-(2-(1H-pyrazol-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0302] The title compound was prepared from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (900 mg, 2.89 mmol) following the experimental procedure described in Preparation 27a. 247 mg (17% yield) of the desired compound were obtained.

LRMS (m/z): 363 (M+1)⁺.

b) (S)-*tert*-Butyl 1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0303] The title compound was prepared from (S)-*tert*-butyl 1-(2-(1H-pyrazol-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (247 mg, 0.49 mmol) following the experimental procedure described in Preparation 54b. 220 mg (46% yield) of the desired compound were obtained.

LRMS (m/z): 345 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(1H-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0304] The title compound was prepared from (S)-*tert*-butyl 1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (220 mg, 0.22 mmol) following the experimental procedure described in Preparation 46c. 54 mg (100% yield) of the desired compound were obtained.

LRMS (m/z): 245 (M+1)⁺.

PREPARATION 58**(S)-3-((1H-Pyrazol-3-yl)methyl)-2-(1-aminoethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one****a) (S)-*tert*-Butyl 1-(2-((1H-pyrazol-3-yl)methylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0305] The title compound was prepared from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-

2-carboxylate (900 mg, 2.89 mmol) following the experimental procedure described in Preparation 27a. 370 mg (14% yield) of the desired compound were obtained.

LRMS (m/z): 377 (M+1)⁺.

b) (S)-tert-Butyl 1-(3-((1H-pyrazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0306] The title compound was prepared from (S)-tert-butyl 1-(2-((1H-pyrazol-3-yl)methylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (370 mg, 0.41 mmol) following the experimental procedure described in Preparation 54b. 784 mg (37% yield) of the desired compound were obtained.

LRMS (m/z): 359 (M+1)⁺.

c) (S)-3-((1H-Pyrazol-3-yl)methyl)-2-(1-aminoethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0307] The title compound was prepared from (S)-tert-butyl 1-(3-((1H-pyrazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (794 mg, 0.16 mmol) following the experimental procedure described in Preparation 46c. 41 mg (100% yield) of the desired compound were obtained.

LRMS (m/z): 259 (M+1)⁺.

PREPARATION 59

(S)-2-(1-Aminoethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one

a) (S)-tert-Butyl 1-oxo-1-(2-(phenylcarbamoyl)-1H-imidazol-1-ylamino)propan-2-ylcarbamate

[0308] The title compound was prepared following the experimental procedure described in preparation 20a, from 1.12 g (5.54 mmol) of 1-amino-N-phenyl-1H-imidazole-2-carboxamide, 1.26 g (6.65 mmol) of (S)-2-(tert-butoxycarbonylamino)propanoic acid and 1.03 g (6.65 mmol) of EDC·HCl. The solid obtained was triturated in diisopropylether to give 1.51 g (67% yield) of the title compound.

LRMS (m/z): 374 (M+1)⁺.

b) (S)-tert-Butyl 1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0309] A solution of bromine (370 µl, 7.39 mmol) in 10 mL of dichloromethane was added dropwise to a solution of triphenylphosphine (1.94 g, 7.39 mmol) in 10 mL of dichloromethane. The resulting solution was stirred for 10 min., and triethylamine (2.58 mL, 18.48 mmol) and (S)-tert-butyl 1-oxo-1-(2-(phenylcarbamoyl)-1H-imidazol-1-ylamino)propan-2-ylcarbamate (1.50 g, 3.70 mmol) in 30 mL of dichloromethane were added. The reaction mixture was refluxed under nitrogen for 2 hours, cooled and concentrated to dryness. The solid thus obtained was dissolved in a mixture of tetrahydrofuran/dimethylformamide (90:10) and 0.78 g (11.09 mmol) of sodium methanethiolate was added. The mixture was heated at 60 °C for 2 hours and then partitioned between a saturated aqueous solution of sodium bicarbonate and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum. The product was purified by flash chromatography (20% to 100% hexane/AcOEt) to yield 0.72 g (55% yield) of the title compound.

LRMS (m/z): 356 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one

[0310] 0.72 g (2.21 mmol) of (S)-tert-butyl 1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate were dissolved in 3 mL of dichloromethane and 3 mL of trifluoroacetic acid were added. The mixture was stirred at room temperature for 1 hour. The solvents were evaporated to dryness and the residue was partitioned between ethyl acetate and an aqueous solution of sodium bicarbonate. The organic layer was washed with brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum. 372 mg (66% yield) of the title compound were obtained.

LRMS (m/z): 256 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.06 (d, 1H), 7.57 (d, 6H), 3.46 - 3.37 (m, 1H), 1.87 (s, 2H), 1.18 (d, 3H).

PREPARATION 60

2-(*tert*-Butoxycarbonylamino)-4,4,4-trifluorobutanoic acid

[0311] 350 mg (2.23 mmol) of 2-amino-4,4,4-trifluorobutanoic acid (purchased from Alfa Aesar® cat. no. L13131) were suspended in 10 mL of dichloromethane and cooled in an ice bath. 534 mg (2.45 mmol) of *di-tert*-butyl dicarbonate and 621 µL (4.46 mmol) of triethylamine were added and the resulting solution was stirred at room temperature overnight. The solution was extracted with water and the aqueous layer was acidified to pH = 3 with 2N hydrochloric acid and extracted with dichloromethane (x3). The combined organic layers were washed with brine, dried over magnesium sulphate, filtered and evaporated to yield 425 mg (74% yield) of a white solid.

LRMS (m/z): 258 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 13.05 (s, 1 H), 7.36 (d, 1 H), 4.20 (td, 1 H), 2.84 - 2.53 (m, 2H), 1.38 (s, 9H).

PREPARATION 61

2-(1-Amino-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) *tert*-Butyl 4,4,4-trifluoro-1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate**

[0312] 421 mg (1.64 mmol) of 2-(*tert*-butoxycarbonylamino)-4,4,4-trifluorobutanoic acid were suspended in a mixture of *N,N*-dimethylformamide (5 mL) and dichloromethane (5 mL) and HATU (624 mg, 1.64 mmol) and DIEA (286 µL, 1.64 mmol) were added. The resulting suspension was stirred at room temperature for 2 hours and then 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide (300 mg, 1.49 mmol) was added. The resulting solution was stirred at room temperature overnight and the dichloromethane was removed in vacuum. 25 mL of water were added to the remaining solution and, after vigorous stirring, the solid that formed was filtered and washed with water. 582 mg (89% yield) of the title compound were obtained as a white solid.

LRMS (m/z): 441 (M+1)⁺.

b) *tert*-Butyl 3,3,3-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate

[0313] A solution of bromine (81 µL, 1.58 mmol) in dichloromethane (5 mL) was added dropwise to a solution of triphenylphosphine (415 mg, 1.58 mmol) in dichloromethane (10 mL) under nitrogen. The solution was stirred for 15 min, and triethylamine (552 µL, 3.97 mmol) and a solution of *tert*-butyl 4,4,4-trifluoro-1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate (582 mg, 1.32 mmol) were added. The reaction mixture was stirred at room temperature for 4 hours, and then the volatiles were removed under reduced pressure. The residue was redissolved in 80 mL of a 7M methanolic solution of ammonia and stirred overnight at 80 °C in a sealed vessel. The solvent was then evaporated and the residue was purified by flash chromatography (0% to 50% AcOEt/hexanes) to yield 228 mg (41% yield) of the title compound as a white solid.

LRMS (m/z): 423 (M+1)⁺.

c) 2-(1-Amino-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0314] 228 mg (0.54 mmol) of *tert*-butyl 3,3,3-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate were dissolved in 5 mL of dichloromethane and 125 µL (1.62 mmol) of trifluoroacetic acid were added. The solvent was evaporated and the residue was partitioned between water and ethyl acetate. The organic layer was washed three times with a diluted aqueous solution of potassium carbonate and brine, dried over magnesium sulphate, filtered and the solvent was removed under vacuum to yield 139 mg (80%) of the title product as a white solid.

LRMS (m/z): 323 (M+1)⁺.

PREPARATION 62

(*S*)-2-(1-Aminoethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (*S*)-*tert*-Butyl 1-(3-methyl-2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0315] Prepared from 2.00 g (9.29 mmol) of 1-amino-3-methyl-*N*-phenyl-1*H*-pyrrole-2-carboxamide and 2.11 g (11.15 mmol) of (*S*)-2-(*tert*-butoxycarbonylamino)propanoic acid following the experimental procedure described in Preparation 22a. After purification by flash chromatography (0 to 10% methanol in dichloromethane), 825 mg (23% yield) of the title

product were obtained.
LRMS (m/z): 387 (M+1)⁺.

b) (S)-tert-Butyl 1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0316] The title compound was obtained from 825 mg (2.13 mmol) of (S)-tert-butyl 1-(3-methyl-2-(phenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate following the experimental procedure described in Preparation 61b. After purification by flash chromatography (0 to 50% AcOEt/hexanes), 330 mg (42% yield) of the title compound were obtained.

LRMS (m/z): 368 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0317] Prepared from 330 mg (0.90 mmol) of (S)-tert-butyl 1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate following the experimental procedure described in Preparation 61c. 201 mg (84% yield) of the title compound were obtained as a white solid.

LRMS (m/z): 268 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.59 - 7.38 (m, 6H), 6.41 (d, 1 H), 3.41 - 3.35 (m, 1 H), 2.39 (s, 3H), 1.15 (d, 3H).

PREPARATION 63

(S)-1-(tert-Butoxycarbonyl)-4,4-difluoropyrrolidine-2-carboxylic acid

[0318] 1.08 g (4.08 mmol) of (S)-1-tert-butyl 2-methyl 4,4-difluoropyrrolidine-1,2-dicarboxylate (purchased from Aldrich®; cat. no. 702463) were dissolved in a mixture of 10 mL of methanol and 10 mL of tetrahydrofurane and 6.12 mL of a 2M aqueous solution of sodium hydroxide were added. The solution was stirred at room temperature for 1 hour and then the organic solvents were removed. The remaining solution was diluted with water and 6.12 mL of 2M hydrochloric acid were added. The product that precipitated was extracted with dichloromethane (x3), the combined organic layers were washed with brine, dried over magnesium sulphate, filtered and the solvent was removed. 1.01 g (99% yield) of the title compound were obtained as a white solid.

LRMS (m/z): 250 (M-1)⁺.

PREPARATION 64

(S)-2-(4,4-Difluoropyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

a) (S)-tert-Butyl 4,4-difluoro-2-(2-(phenylcarbamoyl)-1H-pyrrol-1-ylcarbamoyl)pyrrolidine-1-carboxylate

[0319] 1-Amino-N-phenyl-1H-pyrrole-2-carboxamide (500 mg, 2.48 mmol), (S)-1-(tert-butoxycarbonyl)-4,4-difluoropyrrolidine-2-carboxylic acid (687 mg, 2.73 mmol), HATU (1.04 g, 2.73 mmol) and diisopropylethylamine (476 µl, 2.73 mmol) were dissolved in a mixture of 6 mL of dichloromethane and 6 mL of DMF and stirred at room temperature overnight. The dichloromethane was removed under vacuum and 25 mL of water were poured over the remaining solution. A precipitated appeared and the suspension was stirred overnight. The solid was filtered and washed with water to furnish 1.05 g (97% yield) of the title compound.

LRMS (m/z): 435 (M+1)⁺.

b) (S)-tert-Butyl 4,4-difluoro-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

[0320] To a solution of 760 mg (1.20 mmol) of triphenylphosphine in 20 mL of dichloromethane was added a solution of 150 µl (1.18 mmol) of bromine in 10 mL of dichloromethane dropwise under nitrogen atmosphere. At the end of the addition the colourless solution was stirred for 10 minutes and then 1.34 mL (9.61 mmol) of triethylamine and 1.05 g (2.42 mmol) of (S)-tert-butyl 4,4-difluoro-2-(2-(phenylcarbamoyl)-1H-pyrrol-1-ylcarbamoyl)pyrrolidine-1-carboxylate were added. The reaction mixture was then stirred at 40 °C for 3 hours. In a separate vessel, an additional 760 mg (1.20 mmol) of triphenylphosphine in 20 mL of dichloromethane was added a solution of 150 µl (1.18 mmol) of bromine in 10 mL of dichloromethane dropwise under nitrogen atmosphere and stirred for 15 minutes. Then this solution was added to the reaction mixture and 1.34 mL of triethylamine were also added. After 3 hours at 40 °C, the solvent was evaporated and the residue was redissolved in a mixture of N,N-dimethylformamide (20 mL) and methanol (20 mL) and 500 mg

(7.13 mmol) of sodium methanethiolate was added. The solution was stirred at room temperature overnight. The solvents were evaporated and the residue was partitioned between water and ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. The product was purified by flash chromatography (0% to 30% hexane/AcOEt) to yield 823 mg (82%) of the title compound.

LRMS (m/z): 417 (M+1)⁺.

c) (S)-2-(4,4-Difluoropyrrolidin-2-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0321] 580 mg (1.39 mmol) of (S)-*tert*-butyl 4,4-difluoro-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate were dissolved in dichloromethane (5 mL) and 536 μ l (6.96 mmol) of trifluoroacetic acid were added. The solution was stirred at room temperature overnight and then the volatiles were removed in vacuum. The residue was partitioned between ethyl acetate and a diluted aqueous solution of potassium carbonate, and the organic layer was washed twice with water and brine, dried over magnesium sulphate, filtered and the solvent was removed under vacuum. 418 mg (95% yield) of the title compound were obtained as a yellowish solid.

LRMS (m/z): 317 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.78 - 7.46 (m, 5H), 7.46 - 7.20 (m, 1H), 7.06 - 6.85 (m, 1H), 6.71 - 6.56 (m, 1H), 4.47 (ddd, 1H), 3.93 - 3.67 (m, 2H), 2.87 (m, 1H), 2.47 - 2.27 (m, 1H).

PREPARATION 65

1-Amino-N-(3,5-difluorophenyl)-1*H*-imidazole-2-carboxamide

a) N-(3,5-Difluorophenyl)-1*H*-imidazole-2-carboxamide

[0322] To a solution of 1*H*-imidazole-2-carboxylic acid (2.50 g, 22.3 mmol) in DMF (30 mL) were added 3,5-difluoroaniline (2.23 mL, 22.3 mmol), EDC·HCl (6.41 g, 33.46 mmol) and HOBT (4.52 g, 33.46 mmol). The reaction mixture was stirred at room temperature overnight. The solvent was removed in vacuum and the crude was dissolved in dichloromethane. The solution was washed with a diluted aqueous solution of potassium carbonate, dried over sodium sulphate, filtered and concentrated. 2.75 g (55% yield) of the title compound were obtained as a solid.

LRMS (m/z): 224 (M+1)⁺.

b) 1-Amino-N-(3,5-difluorophenyl)-1*H*-imidazole-2-carboxamide

[0323] Prepared following the experimental procedure described in Preparation 33b from 2.68 g (12.01 mmol) of N-(3,5-difluorophenyl)-1*H*-imidazole-2-carboxamide. The product was purified by flash chromatography (dichloromethane/methanol/ammonium hydroxide, 100/8/1) to yield 1.50 g (52% yield) of the title compound as a pale yellow solid. LRMS (m/z): 239(M+1)⁺.

PREPARATION 66

(S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)imidazo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (S)-*tert*-Butyl 1-(2-(3,5-difluorophenylcarbamoyl)-1*H*-imidazol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0324] 1.60 g (6.18 mmol) of 1-amino-N-(3,5-difluorophenyl)-1*H*-imidazole-2-carboxamide, 1.64 g (8.65 mmol) of (S)-2-(*tert*-butoxycarbonylamino)butanoic acid and 1.34 g (8.65 mmol) of EDC·HCl were dissolved in 45 mL of THF and stirred at 55 °C overnight and at room temperature for 2 days. Then the solvent was evaporated and the crude residue was taken up in ethyl acetate and washed with brine. The organic layer was dried over magnesium sulphate, filtered and the solvent was evaporated. The solid obtained was redissolved in the minimum amount of ethyl acetate and diisopropylether was added until a solid precipitated, which was filtered to furnish 1.51 g (57% yield) of the title compound. LRMS (m/z): 410 (M+1)⁺.

b) (S)-*tert*-Butyl 1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroimidazo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0325] Prepared following the experimental procedure described in Preparation 64b from 1.51 g (3.28 mmol) of (S)-*tert*-butyl 1-(2-(3,5-difluorophenylcarbamoyl)-1*H*-imidazol-1-ylamino)-1-oxopropan-2-ylcarbamate, using 1.05 g (6.57 mmol) of bromine, 1.72 g (6.57 mmol) of triphenylphosphine, 2.29 mL (16.41 mmol) of triethylamine. After 3 hours, the dichloromethane was removed and 0.69 g (9.85 mmol) of sodium methanethiolate was used in a mixture of

N,N-dimethylformamide and methanol at 55 °C for 5 hours. The product was purified by flash chromatography (20% to 60% hexane/AcOEt) to yield 2.46 g (40% purity) of a mixture of the title compound and triphenylphosphine oxide, which was used without further purification in the next step.

LRMS (m/z): 392 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)imidazo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0326] The mixture of (*S*)-*tert*-butyl 1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroimidazo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate and triphenylphosphine oxide obtained in Preparation 66b was dissolved in 20 mL of dichloromethane and 2 mL of trifluoroacetic acid were added. The solution was stirred at room temperature for 1.2 hours and the solvent was removed. The residue was partitioned between water and diethylether, the two layers were separated and the aqueous layer was washed twice more with diethylether. Then the aqueous layer was basified with a diluted aqueous solution of sodium hydroxide and the product was extracted 3 times with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulphate, filtered and the solvent removed under vacuum. 0.26 g of the title compound

were obtained as a solid. Yield = 24% in two steps.

LRMS (m/z): 292 (M+1)⁺.

PREPARATION 67

2-(*tert*-Butoxycarbonylamino)-3,3,3-trifluoropropanoic acid

[0327] 650 mg (4.54 mmol) of 2-amino-3,3,3-trifluoropropanoic acid (purchased from Aldrich®; cat. no. 307556) and 866 mg (9.50 mmol) of tetramethylammonium were suspended in 30 mL acetonitrile. The mixture was stirred for 30 minutes until a clear solution was observed. Then 1.98 g (9.09 mmol) of di-*tert*-butyl dicarbonate were added and the resulting solution was stirred for 2 hours at room temperature. The solvent was removed in vacuum and the residue was dissolved in water and this solution was washed twice with diethylether. The aqueous layer was acidified with 2M hydrochloric acid and the product was extracted with ethyl acetate (x3). The combined organic layers were washed with brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum to furnish 830 mg (75% yield) of a white solid.

LRMS (m/z): 242 (M-1)⁺.

¹H NMR (400 MHz, DMSO) δ 13.83 (s, 1H), 8.07 (d, 1H), 5.10 - 4.76 (m, 1H), 1.41 (s, 9H).

PREPARATION 68

2-(1-Amino-2,2,2-trifluoroethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) *tert*-Butyl 1,1,1-trifluoro-3-oxo-3-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate

[0328] 1-Amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide (570 mg, 2.83 mmol), 2-(*tert*-butoxycarbonylamino)-3,3,3-trifluoropropanoic acid (826 mg, 3.40 mmol), HATU (1.29 g, 3.40 mmol) and *N*-methylmorpholine (685 μl, 6.23 mmol) were suspended in 20 mL of dichloromethane and stirred overnight. The reaction mixture was diluted with more dichloromethane and the solution was washed with water and brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound in a 47% yield (562 mg) as a white solid.

LRMS (m/z): 427 (M+1)⁺.

b) *tert*-Butyl 2,2,2-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0329] To a solution of 414 mg (1.58 mmol) of triphenylphosphine in 10 mL of dichloromethane was added a solution of 81 μl (1.58 mmol) of bromine in 5 mL of dichloromethane dropwise under nitrogen atmosphere. At the end of the addition the colourless solution was stirred for 15 minutes and then 735 μl (5.27 mmol) of triethylamine and 562 mg (1.32 mmol) of *tert*-butyl 1,1,1-trifluoro-3-oxo-3-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate were added. The reaction mixture was then stirred at 40 °C for 3 days. In a separate vessel, an additional 414 mg (1.58 mmol) of triphenylphosphine in 10 mL of dichloromethane was added a solution of 81 μl (1.58 mmol) of bromine in 5 mL of dichloromethane dropwise under nitrogen atmosphere and stirred for 15 minutes. Then this solution was added to the reaction mixture and 734 μl of triethylamine were also added. After 3 hours at 40 °C, the solvent was evaporated and the residue was redissolved in a mixture of *N,N*-dimethylformamide (10 mL) and methanol (10 mL) and 277 mg (3.95

mmol) of sodium methanethiolate was added. The solution was stirred at 60 °C for 4 hours and at room temperature overnight. Then the solvents were evaporated and the residue was partitioned between water and ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. The product was purified by flash chromatography (0% to 40% hexane/AcOEt) to yield 218 mg (41 %) of the title compound. LRMS (m/z): 409 (M+1)⁺.

c) 2-(1-Amino-2,2,2-trifluoroethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0330] 218 mg (0.53 mmol) of *tert*-butyl 2,2,2-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate were dissolved in dichloromethane (2 mL) and 205 µl of trifluoroacetic acid were added. The solution was stirred at room temperature for 3 hours and then the solvent was removed to dryness. The residue was partitioned between ethyl acetate and a diluted aqueous solution of potassium carbonate, and the organic layer was washed twice with water and brine, dried over magnesium sulphate, filtered and the solvent was removed under vacuum. 155 mg (94% yield) of the title compound were obtained as a brownish solid.

LRMS (m/z): 309 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.74 (s, 1 H), 7.65 - 7.50 (m, 4H), 7.40 (d, 1 H), 7.05 - 6.99 (m, 1 H), 6.70 - 6.64 (m, 1 H), 4.00 - 3.86 (m, 1 H), 2.63 (d, 2H).

PREPARATION 69

6-Bromo-9-(4-methoxybenzyl)-9*H*-purine

[0331] 1.00 g (5.02 mmol) of 6-bromo-9*H*-purine (purchased from Aldrich®; cat. no. 104981) was suspended in 10 mL of DMF and potassium carbonate (2.08 g, 15.05 mmol) was added. The mixture was stirred at room temperature for 20 minutes and then 1-(chloromethyl)-4-methoxybenzene (1.40 mL, 10.06 mmol) were added. The reaction mixture was stirred at 45 °C overnight. Then the mixture was evaporated to dryness and the residue was partitioned between water and dichloromethane. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum. The product was purified by flash chromatography (0% to 50% hexane/AcOEt) to yield 447 mg (28%) of the title compound.

LRMS (m/z): 320 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.83 (s, 1 H), 8.75 (s, 1 H), 7.35 (d, 2H), 6.90 (d, 2H), 5.44 (s, 2H), 3.71 (s, 3H).

PREPARATION 70

3-Phenyl-2-(2,2,2-trifluoro-1-(9-(4-methoxybenzyl)-9*H*-purin-6-ylamino)ethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0332] 2-(1-Amino-2,2,2-trifluoroethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (120 mg, 0.39 mmol), 6-bromo-9-(4-methoxybenzyl)-9*H*-purine (150 mg, 0.47 mmol), BINAP (73 mg, 0.12 mmol), caesium carbonate (190 mg, 0.58 mmol) and palladium (II) acetate (9 mg, 0.04 mmol) were suspended in toluene (2 mL) and stirred at 120 °C under nitrogen atmosphere overnight. The solvent was removed in vacuum and the residue was partitioned between water and dichloromethane. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and the solvent was evaporated to dryness. The product was purified by flash chromatography (0% to 30% hexane/AcOEt) to give 138 mg (65%) of the title compound as a white solid.

LRMS (m/z): 547 (M+1)⁺.

PREPARATION 71

(*S*)-2-(1-Aminoethyl)-3-benzylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (*S*)-*tert*-Butyl 1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0333] (*S*)-*tert*-Butyl 1-(4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (150 mg, 0.43 mmol) was suspended in hexane (2 mL) and benzyl bromide (110 µl, 0.92 mmol) and silver (I) carbonate (150 mg, 0.54 mmol) were added. The mixture was heated at 150 °C for 20 minutes using microwave irradiation. The reaction mixture was filtered through Celite® and the solvent was evaporated. The product was isolated from the mixture of by-products by flash chromatography (0% to 40% hexane/AcOEt) to give 20 mg (13%) of the title compound as a white solid.

LRMS (m/z): 369 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-3-benzylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0334] 20 mg (0.05 mmol) of (S)-*tert*-butyl 1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate were dissolved in 1 mL of dichloromethane and 1 mL of trifluoroacetic acid were added. The reaction mixture was stirred at room temperature for 1.5 hours and the volatiles were removed in vacuum to furnish 21 mg (100% yield) of (S)-2-(1-Aminoethyl)-3-benzylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one as its trifluoroacetate salt. LRMS (m/z): 269 (M+1)⁺.

PREPARATION 72**1-Amino-3-bromo-*N*-(3,5-dichlorophenyl)-1*H*-pyrrole-2-carboxamide****a) 3-Bromo-*N*-(3,5-dichlorophenyl)-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxamide**

[0335] Prepared from 2.00 g (5.81 mmol) of methyl 3-bromo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate and 2.82 g (17.41 mmol) of 3,5-dichloroaniline following the experimental procedure described in Preparation 44a. The product was purified by flash chromatography (0% to 40% hexane/AcOEt) to give 2.64 g (96% yield) of the title compound as a beige solid. LRMS (m/z): 474 (M+1)⁺. BrCl₂ isotopic pattern.

b) 3-Bromo-*N*-(3,5-dichlorophenyl)-1*H*-pyrrole-2-carboxamide

[0336] Prepared from 2.64 g (5.57 mmol) of 3-bromo-*N*-(3,5-dichlorophenyl)-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxamide following the experimental procedure described in Preparation 44b. 1.51 g (81% yield) of the title compound were obtained as a white solid. LRMS (m/z): 334 (M+1)⁺. BrCl₂ isotopic pattern.

c) 1-Amino-3-bromo-*N*-(3,5-dichlorophenyl)-1*H*-pyrrole-2-carboxamide

[0337] This compound was prepared starting from 3-Bromo-*N*-(3,5-dichlorophenyl)-1*H*-pyrrole-2-carboxamide (1.51 g, 4.52 mmol) and following the experimental procedure described in Preparation 1 b to afford 1.58 g (100% yield) of the title compound that was used in the next step without any further purification. LRMS (m/z): 350 (M+1)⁺. BrCl₂ isotopic pattern. ¹H NMR (400 MHz, DMSO) δ 11.23 (s, 1 H), 7.78 (d, 2H), 7.32 (t, 1 H), 7.02 (d, 1 H), 6.54 (s, 2H), 6.21 (d, 1 H).

PREPARATION 73**(S)-*tert*-Butyl 1-(5-bromo-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate****a) (S)-*tert*-Butyl 1-(3-bromo-2-(3,5-dichlorophenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0338] To a solution of 1.58 g (4.53 mmol) of 1-amino-3-bromo-*N*-(3,5-dichlorophenyl)-1*H*-pyrrole-2-carboxamide in 30 mL DMF were added 2.60 mL (14.93 mmol) of DIEA and 0.94 g (4.97 mmol) of (S)-2-(*tert*-butoxycarbonylamino)propanoic acid. The solution was cooled to 0 °C and 4 mL of T3P® (50% solution in DMF, 6.78 mmol) were added dropwise. The mixture was stirred at room temperature overnight and an additional 2.60 mL (14.93 mmol) of DIEA, 0.94 g (4.97 mmol) of (S)-2-(*tert*-butoxycarbonylamino)propanoic acid and 4 mL of T3P® (50% solution in DMF, 6.78 mmol) were added. The stirring was kept for 24 hours more and water was added to the reaction mixture. The product was extracted three times with ethyl acetate and the combined organic extracts were washed with water and brine, dried over magnesium sulphate, filtered and the solvents evaporated in vacuum. The product was purified by flash chromatography (0% to 50% hexane/AcOEt) to give 1.11 g (47% yield) of the title compound as a beige solid. LRMS (m/z): 521 (M+1)⁺. BrCl₂ isotopic pattern.

b) (S)-*tert*-Butyl 1-(5-bromo-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0339] A solution of bromine (130 µL, 2.56 mmol) in dichloromethane (3 mL) was added dropwise to a solution of triphenylphosphine (670 mg, 2.55 mmol) in dichloromethane (20 mL) under nitrogen. The solution was stirred for 10

min, and triethylamine (1.2 mL, 2.56 mmol) and a solution of (*S*)-*tert*-butyl 1-(3-bromo-2-(3,5-dichlorophenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (1.11 g, 2.13 mmol) were added. The reaction mixture was stirred at room temperature for 3 hours. In a separate vessel, an additional 670 mg (2.55 mmol) of triphenylphosphine in 20 mL of dichloromethane was added a solution of 130 μ L (2.56 mmol) of bromine in 5 mL of dichloromethane dropwise under nitrogen atmosphere and stirred for 15 minutes. Then this solution was added to the reaction mixture and an additional 1.2 mL of triethylamine was also added. The reaction mixture was stirred overnight at 40 °C and then the volatiles were removed under reduced pressure. The residue was redissolved in 40 mL of a 7M methanolic solution of ammonia and stirred overnight at 100 °C in a sealed vessel. The solvent was then evaporated and the residue was purified by flash chromatography (0% to 30% AcOEt/hexanes) to yield 576 mg (54% yield) of the title compound as a white solid.

LRMS (m/z): 503 (M+1)⁺. BrCl₂ isotopic pattern.

PREPARATION 74

(*S*)-2-(1-Aminoethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

a) (*S*)-*tert*-Butyl 1-(5-cyano-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0340] Prepared according to the experimental procedure described in Preparation 47a from 570 mg (1.14 mmol) of (*S*)-*tert*-butyl 1-(5-bromo-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate. The product was purified by flash chromatography (0% to 30% AcOEt/hexanes) to give 418 mg (82% yield) of the title compound as a white solid.

LRMS (m/z): 509 (M+1)⁺. Cl₂ isotopic pattern.

b) (*S*)-2-(1-Aminoethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0341] 418 mg (0.93 mmol) of (*S*)-*tert*-butyl 1-(5-cyano-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate were dissolved in 5 mL of dichloromethane and 431 μ L of trifluoroacetic acid were added. The reaction mixture was stirred at room temperature for 5 hours and the volatiles were removed in vacuum. The residue was partitioned between water and dichloromethane and 2M aqueous solution of sodium hydroxide was added until pH= 9 was reached. Then the organic layer was washed with brine, dried over magnesium sulphate, filtered and the solvent was evaporated in vacuum. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound in a 58% yield (188 mg) as a white solid.

LRMS (m/z): 349 (M+1)⁺. Cl₂ isotopic pattern.

¹H NMR (400 MHz, DMSO) δ 7.91 - 7.82 (m, 2H), 7.78 (s, 2H), 7.21 (d, 1H), 3.53 - 3.41 (m, 1H), 1.87 (s, 2H), 1.22 (d, 3H).

PREPARATION 75

(*R*)-*tert*-Butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)-2-*tert*-butoxyethylcarbamate

a) (*S*)-*tert*-Butyl 1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-3-*tert*-butoxy-1-oxopropan-2-ylcarbamate

[0342] Prepared according to the experimental procedure described in Preparation 61a from 1.07 g (3.39 mmol) of 1-amino-3-bromo-*N*-(3,5-difluorophenyl)-1*H*-pyrrole-2-carboxamide and 1.65 g (3.73 mmol) of (*S*)-3-*tert*-butoxy-2-(*tert*-butoxycarbonylamino)propanoic acid. After stirring the reaction mixture overnight, an excess of an additional 1.65 g (3.73 mmol) of the acid was added and the reaction stirred for yet 24 hours more. After precipitation and filtration of the product, two purification steps were needed. First by flash chromatography (70% to 100% DCM/hexanes) and a second purification by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%). 494 mg (26% yield) of the title compound were isolated as a white solid.

LRMS (m/z): 559, 561 (M+1)⁺. Br isotopic pattern.

b) (*R*)-*tert*-Butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)-2-*tert*-butoxyethylcarbamate

[0343] Prepared according to the experimental procedure described in Preparation 68b from 494 mg (0.88 mmol) of

(*S*)-*tert*-butyl 1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-3-*tert*-butoxy-1-oxopropan-2-ylcarbamate. The reaction was stirred overnight and then after a second addition of reagents for 2 hours more. The final product was purified by flash chromatography (0% to 40% AcOEt/hexanes) to give 313 mg (66% yield) of the title compound as a white solid.

LRMS (*m/z*): 541, 543 (*M*+1)⁺. Br isotopic pattern.

¹H NMR (400 MHz, CDCl₃) δ 7.32 (d, 1 H), 7.05 - 6.87 (m, 3H), 6.62 (d, 1 H), 5.01 (s, 1 H), 4.58 - 4.45 (m, 1 H), 3.57 - 3.38 (m, 2H), 1.41 (s, 9H), 1.10 (s, 9H).

PREPARATION 76

(*R*)-2-(1-Amino-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

a) (*R*)-*tert*-Butyl 2-*tert*-butoxy-1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0344] Prepared from 150 mg (0.28 mmol) of (*R*)-*tert*-butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)-2-*tert*-butoxyethylcarbamate following the experimental procedure described in Preparation 47a. The product was purified by flash chromatography (0% to 40% AcOEt/hexanes) to give 105 mg (78% yield) of the title compound as a white solid.

LRMS (*m/z*): 488 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 7.38 (d, 1H), 7.08 - 6.92 (m, 3H), 6.90 (d, 1H), 5.03 (d, 1H), 4.63 - 4.52 (m, 1H), 3.57 - 3.41 (m, 2H), 1.40 (s, 9H), 1.10 (s, 9H).

b) (*R*)-2-(1-Amino-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0345] 105 mg (0.22 mmol) of (*R*)-*tert*-butyl 2-*tert*-butoxy-1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate were stirred in 5.5 mL of a 4M solution of hydrogen chloride in dioxane overnight at room temperature and then 2 hours at 50 °C. The volatiles were evaporated under reduced pressure and the residue was partitioned between water and dichloromethane and 2M aqueous solution of sodium hydroxide was added until pH= 8 was reached. Then the organic layer was washed with brine, dried over magnesium sulphate, filtered and the solvent was evaporated. 71 mg (100% yield) of the title product were obtained.

LRMS (*m/z*): 332 (*M*+1)⁺.

PREPARATION 77

(*R*)-*tert*-Butyl 2-*tert*-butoxy-1-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0346] A solution of (*R*)-*tert*-butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)-2-*tert*-butoxyethylcarbamate (150 mg, 0.28 mmol), *trans*-1,2-bis(methylamino)cyclohexane (24 mg, 0.17 mmol), sodium iodide (165, 1.11 mmol) and copper (I) iodide (16 mg, 0.08 mmol) in 1,4-dioxane (2 mL) was stirred under argon atmosphere at 120 °C overnight. The crude was allowed to reach room temperature and filtered through Celite® washing with ethyl acetate. The organic solution was washed with water (x3) and brine, dried over magnesium sulphate and concentrated to give 148 mg (91% yield) of the title compound, which was used without further purification.

LRMS (*m/z*): 589 (*M*+1)⁺.

PREPARATION 78

(*R*)-2-(1-Amino-2-hydroxyethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (*R*)-*tert*-Butyl 2-*tert*-butoxy-1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0347] Prepared following the experimental procedure described in Preparation 26b from 148 mg (0.25 mmol) of (*R*)-*tert*-butyl 2-*tert*-butoxy-1-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate. 133 mg (100% yield) of the title compound were obtained.

LRMS (m/z): 531 (M+1)⁺.

b) (R)-2-(1-Amino-2-hydroxyethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)-pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0348] Prepared from (R)-*tert*-butyl 2-*tert*-butoxy-1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (134 mg, 0.25 mmol) following the method described in Preparation 76b. 61 mg (65% yield) of the title compound were obtained and used directly in the next step.
LRMS (m/z): 375 (M+1)⁺.

PREPARATION 79

(S)-*tert*-Butyl 1-(7-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0349] To a solution of (S)-*tert*-butyl 1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (500 mg, 1.41 mmol) in a mixture of methanol (5 mL) and tetrahydrofuran (5 mL), *N*-bromosuccinimide (252 mg, 1.42 mmol) were added. The reaction mixture was stirred at room temperature for 40 hours and then an excess of *N*-bromosuccinimide (252 mg, 1.42 mmol) was added. After 2 hours at room temperature the solvents were removed in vacuum and the residue was partitioned between water and dichloromethane. The organic layer was washed with brine, dried over magnesium sulphate, filtered and the solvent was evaporated. Purification by flash chromatography (0% to 15% AcOEt/hexanes) yielded 233 mg (38% yield) of the title compound as a white solid.
LRMS (m/z): 433, 435 (M+1)⁺. Br isotopic pattern.

PREPARATION 80

(S)-*tert*-Butyl 1-(7-iodo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0350] Prepared from 233 mg (0.54 mmol) of (S)-*tert*-butyl 1-(7-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate following the experimental procedure described in Preparation 77. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 130 mg (50% yield) of the title compound.
LRMS (m/z): 481 (M+1)⁺.
¹H NMR (400 MHz, CDCl₃) δ 7.61 - 7.47 (m, 4H), 7.39 (d, 1 H), 7.28 (t, 1 H), 7.12 (d, 1H), 6.75 (d, 1H), 5.19 (d, 1H), 4.58 - 4.45 (m, 1H), 1.43 (s, 9H), 1.29 (d, 3H).

PREPARATION 81

(S)-2-(1-Aminoethyl)-3-phenyl-7-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

a) (S)-*tert*-Butyl 1-(4-oxo-3-phenyl-7-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0351] Prepared following the experimental procedure described in Preparation 26b from 22 mg (0.05 mmol) of (S)-*tert*-butyl 1-(7-iodo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate. After purification by flash chromatography (0% to 50% AcOEt/hexanes), 17 mg (87% yield) of the title compound were obtained.
LRMS (m/z): 423 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-3-phenyl-7-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0352] (S)-*tert*-Butyl 1-(4-oxo-3-phenyl-7-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (17 mg, 0.04 mmol) was stirred in 5 mL of a 4M solution of hydrogen chloride in dioxane at room temperature for 4 hours. The volatiles were evaporated under reduced pressure and 14 mg (100% yield) of the title compound, isolated as the hydrochloric salt form, were obtained and used directly in the next step.
LRMS (m/z): 323 (M+1)⁺.

PREPARATION 82

(S)-2-(1-Aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-7-carbonitrile**a) (S)-*tert*-Butyl 1-(7-cyano-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0353] A mixture of (*S*)-*tert*-butyl 1-(7-iodo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (130 mg, 0.27 mmol), dicyanozinc (64 mg, 0.55 mmol) and tetrakis(triphenylphosphine)palladium (0) (31 mg, 0.03 mmol) in DMF, was heated at 120 °C in a sealed tub with stirring. Additional amounts of dicyanozinc (64 mg, 0.55 mmol) and tetrakis(triphenylphosphine)palladium (0) (31 mg, 0.03 mmol) were added after 16 hours and 40 hours. After 64 hours of reaction, the mixture was allowed to cool to room temperature and ethyl acetate was added. The resulting suspension was filtered through Celite®, the two phases were separated and the aqueous layer extracted with more ethyl acetate. The combined organic layer were dried over magnesium sulphate and concentrated in vacuum. The product was purified by flash chromatography (0% to 50% AcOEt/hexanes) to obtain 37 mg (36% yield) of the title compound.

LRMS (m/z): 380 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-7-carbonitrile

[0354] (*S*)-*tert*-Butyl 1-(7-cyano-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (37 mg, 0.20 mmol) was stirred in 5 mL of a 4M solution of hydrogen chloride in dioxane at room temperature for 4 hours. The volatiles were evaporated under reduced pressure and the residue was partitioned between dichloromethane and diluted aqueous solution of potassium carbonate. The organic layer was washed with brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum to give 53 mg (100% yield) of the title compound which was used directly in the next step.

LRMS (m/z): 280 (M+1)⁺.

PREPARATION 83

(S)-2-(1-Aminoethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**a) (S)-*tert*-Butyl 1-(3-bromo-2-(pyridin-2-ylmethylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0355] Same procedure as described in preparation 133a was used from (*S*)-methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (374mg, 0.96 mmols) and pyridine-2-ylmethanamine (0.30ml, 2.87 mmols). After reverse phase chromatography the title compound was obtained (74mg, 17%).

LRMS (m/z): 467 (M+1)⁺

b) (S)-*tert*-Butyl 1-(5-bromo-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0356] Same procedure as described in preparation 133b was used from (*S*)-*tert*-butyl 1-(3-bromo-2-(pyridin-2-ylmethylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (271 mg, 0.41 mmols). After reverse phase chromatography the title compound was obtained (50mg, 27%)

LRMS (m/z): 449 (M+1)⁺

c) (S)-*tert*-Butyl 1-(5-iodo-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0357] Sodium iodide (67mg, 0.45 mmols), Copper (I) iodide (11mg, 0.06 mmols) and (1*S*,2*S*)-N1,N2-dimethylcyclohexane-1,2-diamine (10mg, 0.07 mmols) were added to a solution of (*S*)-*tert*-butyl 1-(5-bromo-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (50mg, 0.11 mmols) in dioxane (5.4ml). It was stirred at 120°C in a sealed tube for 5 days. It was concentrated in vacuum. Ethyl acetate was added and it was washed with water and brine. The title compound was obtained (59mg, 88%).

LRMS (m/z): 496 (M+1)⁺

d) (S)-tert-Butyl 1-(5-cyano-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0358] Copper (I) cyanide (98mg, 1.09mmols) was added to a solution of (S)-tert-butyl 1-(5-iodo-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (59mg, 0.01 mmols) in pyridine (5ml). It was stirred at 115°C in a sealed tube overnight. It was concentrated in vacuum. Ethyl acetate and water were added and it was filtered through celite. The organic phase was washed, dried, filtered and concentrated in vacuum. The title compound was obtained (59mg, 87% purity, 100%).

LRMS (m/z): 395 (M+1)+

e) (S)-2-(1-Aminoethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile

[0359] Same procedure as described in preparation 133c was used from (S)-tert-butyl 1-(5-cyano-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (49mg, 0.11 mmols). The title compound was obtained (39mg, 45% purity, 4%) pure enough to be used in the next synthetic step without further purification.

LRMS (m/z): 295 (M+1)+

PREPARATION 84**2-((S)-1-Aminoethyl)-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile****a) tert-Butyl (2S)-1-(3-bromo-2-(tetrahydro-2H-pyran-3-yl)carbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0360] Same procedure as described in preparation 137a was used from (S)-methyl 3-bromo-1-(2-(tert-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (4.00g, 10.25 mmols) and tetrahydro-2H-pyran-3-amine.HCl (2.12g, 15.41 mmols). The title compound was obtained (2.24g, 48%).

LRMS (m/z): 460 (M+1)+

b) tert-Butyl (1S)-1-(5-bromo-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0361] Same procedure as described in preparation 137b was used from tert-butyl (2S)-1-(3-bromo-2-(tetrahydro-2H-pyran-3-yl)carbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (2.14g, 4.66 mmols). The title compound was obtained (3.91g, 50% purity, 95%) pure enough to be used in the next reaction step without further purification.

LRMS (m/z): 442 (M+1)+

c) tert-Butyl (1S)-1-(5-iodo-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0362] Same procedure as described in Preparation 83c was used from tert-butyl (1S)-1-(5-bromo-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (3.91g, 4.43 mmols). After reverse phase chromatography the title compound was obtained (0.60g, 28%).

LRMS (m/z): 489 (M+1)+

d) tert-Butyl (1S)-1-(5-cyano-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0363] Same procedure as described in Preparation 83d was used from tert-butyl (1S)-1-(5-iodo-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (280mg, 0.57 mmols). The title compound was obtained (0.33g, 60% purity, 88%) pure enough to be used in the next synthetic step without further purification.

LRMS (m/z): 388 (M+1)+

e) 2-((S)-1-Aminoethyl)-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile

[0364] Same procedure as described in preparation 133c was used from tert-butyl (1S)-1-(5-cyano-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (329mg, 0.50 mmols). The title compound

was obtained (0.12g, 57% purity, 47%) pure enough to be used in the next synthetic step without further purification.
LRMS (m/z): 288 (M+1)+

PREPARATION 85

2-((S)-1-Aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

a) *tert*-Butyl (1S)-1-(4-oxo-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0365] Copper (I) iodide (131mg, 0.69 mmols), Hexamethylphosphoramide (HMPA) (0.5ml, 2.87mmols) and methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (0.37ml, 2.87mmols) were added to a solution of *tert*-butyl (1S)-1-(5-iodo-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (280mg, 0.57mmols) in dimethylformamide (5.5ml). It was stirred at 80°C overnight in a sealed tube. It was concentrated in vacuum, ethyl acetate was added and it was filtered through celite. It was washed with water and brine. The title compound was obtained (303mg, 51% purity, 63%) pure enough to be used in the next synthetic step without further purification.
LRMS (m/z): 431 (M+1)+

b) 2-((S)-1-Aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0366] Same procedure as described in preparation 133c was used from *tert*-butyl (1S)-1-(4-oxo-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (303mg, 0.36mmols). The title compound was obtained (0.16g, 41% purity, 55%) pure enough to be used in the next synthetic step without further purification.
LRMS (m/z): 331 (M+1)+

PREPARATION 86

2-(1-Aminoethyl)-3-((5-methylisoxazol-3-yl)methyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

a) (S)-*tert*-Butyl 1-(2-((5-methylisoxazol-3-yl)methylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0367] The title compound was prepared from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (750 mg, 2.41 mmol) and (5-methylisoxazol-3-yl)methanamine bromhydrate (698 mg, 3.62 mmol) following the experimental procedure described in Preparation 27a. 613 mg (63% yield) of the desired compound were obtained.
LRMS (m/z): 392 (M+1)+.

b) (S)-*tert*-Butyl 1-(3-((5-methylisoxazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0368] The title compound was prepared from (S)-*tert*-butyl 1-(2-((5-methylisoxazol-3-yl)methylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (613 mg, 1.57 mmol) following the experimental procedure described in Preparation 54b. 1.07 g (62% yield) of the desired compound were obtained.
LRMS (m/z): 374 (M+1)+.

c) 2-(1-Aminoethyl)-3-((5-methylisoxazol-3-yl)methyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0369] The title compound was prepared from (S)-*tert*-butyl 1-(3-((5-methylisoxazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (1.07, 0.97 mmol) following the experimental procedure described in Preparation 46c. 300 mg (93% yield) of the desired compound were obtained.
LRMS (m/z): 274 (M+1)+.

PREPARATION 87**2-(1-Aminoethyl)-3-(1-methyl-1*H*-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) *tert*-Butyl 1-(2-(1-methyl-1*H*-pyrazol-3-yl)carbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0370] The title compound was prepared from (*S*)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (900 mg, 2.89 mmol) and 1-methyl-1*H*-pyrazol-3-amine (421 mg, 4.33 mmol) following the experimental procedure described in Preparation 27a. 680 mg (36% yield) of the desired compound were obtained.

LRMS (*m/z*): 377 (*M*+1)⁺.

b) (*S*)-*tert*-Butyl 1-(3-(1-methyl-1*H*-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0371] The title compound was prepared from (*S*)-*tert*-butyl 1-(2-(1-methyl-1*H*-pyrazol-3-yl)carbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (680 mg, 1.05 mmol) following the experimental procedure described in Preparation 54b. 750 mg (50% yield) of the desired compound were obtained.

LRMS (*m/z*): 359 (*M*+1)⁺.

c) 2-(1-Aminoethyl)-3-(1-methyl-1*H*-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0372] The title compound was prepared from (*S*)-*tert*-butyl 1-(3-(1-methyl-1*H*-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (750mg, 0.52 mmol) following the experimental procedure described in Preparation 46c. 100 mg (72% yield) of the desired compound were obtained.

LRMS (*m/z*): 259 (*M*+1)⁺.

PREPARATION 88**(*S*)-2-(1-Aminoethyl)-7-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) 5-Methyl-N-phenyl-1*H*-pyrrole-2-carboxamide**

[0373] The title compound was prepared from ethyl 5-methyl-1*H*-pyrrole-2-carboxylate (1g, 6.53 mmol, purchased from Matrix) following the experimental procedure described in Preparation 44a. Trituration with diisopropyl ether gave the title compound as a beige solid (0.65g, 98% yield).

LRMS (*m/z*): 201 (*M*+1)⁺.

b) 1-Amino-5-methyl-N-phenyl-1*H*-pyrrole-2-carboxamide

[0374] This compound was prepared starting from 5-methyl-N-phenyl-1*H*-pyrrole-2-carboxamide (0.64 g, 3.20 mmol) and following the experimental procedure described in Preparation 1b to afford 0.68 g (50% yield) of the title compound that was used in the next step without any further purification.

LRMS (*m/z*): 216 (*M*+1)⁺.

c) (*S*)-*tert*-Butyl 1-(2-methyl-5-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0375] The title compound was prepared following the experimental procedure described in preparation 20a from 380 mg (1.77 mmol) of 1-amino-5-methyl-N-phenyl-1*H*-pyrrole-2-carboxamide and 334mg (1.77 mmol) of (*S*)-2-(*tert*-butoxycarbonylamino)propanoic acid (purchased from Aldrich). The crude product was purified by flash chromatography in hexane/ethyl acetate to afford 500 mg (59% yield) of the title compound.

LRMS (*m/z*): 387 (*M*+1)⁺.

d) (*S*)-*tert*-Butyl 1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0376] This compound was prepared starting from (*S*)-*tert*-butyl 1-(2-methyl-5-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (500 mg, 1.04 mmol) and following the experimental procedure described in Preparation 42b. The residue was purified by flash chromatography in hexane/ethyl acetate to afford 95 mg (24% yield) of the title compound were obtained.

LRMS (m/z): 369 (M+1)⁺.

e) (S)-2-(1-Aminoethyl)-7-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

5 **[0377]** The title compound was prepared from (S)-*tert*-butyl 1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (95 mg, 0.26 mmol) following the experimental procedure described in Preparation 46c. 70 mg (97% yield) of the desired compound were obtained.

LRMS (m/z): 269 (M+1)⁺.

10 PREPARATION 89

7-Chloropyrazolo[1,5-*a*]pyrimidine

15 **[0378]** Pyrazolo[1,5-*a*]pyrimidin-7(4*H*)-one (0.50g, 3.70 mmols), phosphorus oxychloride (0.88ml, 9.62 mmols) and diisopropylethylamine (DIEA, 0.13ml, 0.74 mmols) were mixed and stirred at 90°C overnight. It was poured onto water/ice, extracted with dichloromethane and washed with brine. It was dried, filtered and concentrated in vacuum. It was purified by chromatography (Silica gel, Hexane/Ethyl acetate 9:1) to afford the expected compound (83mg, 71%).

LRMS (m/z): 154 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 6.93 (d, 1 H) 7.43 (d, 1 H) 8.37 (d, 1 H) 8.52 (d, 1 H)

20

PREPARATION 90

(S)-2-(1-(methylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

25 **a) (S)-*tert*-butyl 1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0379] To a solution of (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (300mg, 0.87 mmol) and triethylamine (302 μl, 2.17 mmol) in DCM (15 ml) was added di-*tert*-butyl dicarbonate (227 mg, 1.04 mmol) and the reaction mixture stirred overnight at room temperature. Ethyl acetate was added and the organic phase washed with

30

water, then dried, filtered and concentrated in vacuum to yield the title compound as an oil (99% yield).

LRMS (m/z): 355 (M+1)⁺

b) (S)-*tert*-Butyl methyl(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethyl)carbamate

35 **[0380]** To a solution of (S)-*tert*-butyl 1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (415 mg, 1.17 mmol) in tetrahydrofuran (20 mL), sodium *tert*-butoxide (169 mg, 1.75mmol) was added. The reaction mixture was stirred at room temperature for 1 hour and then methyl iodide (109 μl, 1.75 mmol) was added. After overnight stirring at room temperature ethyl acetate was added and the organic layer washed with water and brine. The organic layer was then dried over magnesium sulphate, filtered and concentrated. Purification by reverse phase flash chromatography

40

(0% to 50% ACN/water) yielded 225 mg (52% yield) of the title compound as a white solid.

LRMS (m/z): 369 (M+1)⁺

c) (S)-2-(1-(Methylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

45 **[0381]** The title compound was prepared from (S)-*tert*-butyl methyl(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethyl)carbamate (225 mg, 0.61 mmol) following the experimental procedure described in Preparation 46c. 170 mg (87% yield) of the desired compound were obtained.

LRMS (m/z): 305 (M+1)⁺.

50 PREPARATION 91

1-Amino-3-bromo-N-(3,5-difluorophenyl)-1*H*-pyrrole-2-carboxamide

a) 3-Bromo-N-(3,5-difluorophenyl)-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxamide

55

[0382] The title compound was prepared from methyl 3-bromo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate² (18 g, 52.2 mmol) following the experimental procedure described in Preparation 44a. 23 g (100% yield) of the desired compound were obtained.

LRMS (m/z): 441, 443 (M+1)⁺.

b) 3-Bromo-N-(3,5-difluorophenyl)-1H-pyrrole-2-carboxamide

5 **[0383]** The title compound was prepared from 3-bromo-N-(3,5-difluorophenyl)-1-(phenylsulfonyl)-1H-pyrrole-2-carboxamide (23 g, 52.2 mmol) following the experimental procedure described in Preparation 44b. 14.6 g (93% yield) of the desired compound were obtained.

LRMS (m/z): 301, 303 (M+1)⁺.

10 **c) 1-Amino-3-bromo-N-(3,5-difluorophenyl)-1H-pyrrole-2-carboxamide**

[0384] The title compound was prepared from 3-bromo-N-(3,5-difluorophenyl)-1H-pyrrole-2-carboxamide (14.6 g, 48.6 mmol) following the experimental procedure described in Preparation 44c. 8.9 g (58% yield) of the desired compound were obtained.

15 LRMS (m/z): 316, 318 (M+1)⁺.

PREPARATION 92

20 **(S)-tert-Butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate**

a) (S)-tert-Butyl 1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

25 **[0385]** The title compound was prepared from 1-amino-3-bromo-N-(3,5-difluorophenyl)-1H-pyrrole-2-carboxamide (7.87 g, 24.9 mmol) and (S)-2-(tert-butoxycarbonylamino)propanoic acid (purchased from Aldrich, 5.65 g, 29.9 mmol) following the experimental procedure described in Preparation 45a. 6.6 g (51% yield) of the desired compound were obtained.

LRMS (m/z): 487, 489 (M+1)⁺.

30 **b) (S)-tert-Butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate**

[0386] The title compound was prepared from (S)-tert-butyl 1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (6.12 g, 12.6 mmol) following the experimental procedure described in Preparation 45b. 2.6 g (44% yield) of the desired compound were obtained.

35 LRMS (m/z): 469, 471 (M+1)⁺.

PREPARATION 93

40 **(S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one**

a) (S)-tert-Butyl 1-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

45 **[0387]** The title compound was prepared from (S)-tert-butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (500 mg, 1.07 mmol) following the experimental procedure described in Preparation 46a. 578 mg (77.5% yield) of the desired compound were obtained.

LRMS (m/z): 517 (M+1)⁺.

50 **b) (S)-tert-Butyl 1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate**

[0388] The title compound was prepared from (S)-tert-butyl 1-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (578 mg, 0.83 mmol) following the experimental procedure described in Preparation 46b. 342 mg (87.5% yield) of the desired compound were obtained.

55 LRMS (m/z): 459 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0389] The title compound was prepared from (*S*)-*tert*-butyl 1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (342 mg, 0.75 mmol) following the experimental procedure described in Preparation 46c. 132 mg (47% yield) of the desired compound were obtained.
LRMS (*m/z*): 359 (*M*+1)⁺.

PREPARATION 94**(S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile****a) (S)-*tert*-Butyl 1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0390] The title compound was prepared from (*S*)-*tert*-butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (500 mg, 1.07 mmol) following the experimental procedure described in Preparation 47a. 326 mg (73.7% yield) of the desired compound were obtained.
LRMS (*m/z*): 416 (*M*+1)⁺.

b) (S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0391] The title compound was prepared from (*S*)-*tert*-butyl 1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (326 mg, 0.78 mmol) following the experimental procedure described in Preparation 47b. 250 mg (90% yield) of the desired compound were obtained.
LRMS (*m/z*): 352 (*M*+1)⁺.

PREPARATION 95**(S)-2-(1-Aminopropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile****a) (S)-*tert*-Butyl 1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxobutan-2-ylcarbamate**

[0392] The title compound was prepared from 1-amino-3-bromo-N-(3,5-difluorophenyl)-1*H*-pyrrole-2-carboxamide (1 g, 3.16 mmol) and (*S*)-2-(*tert*-butoxycarbonylamino)butanoic acid following the experimental procedure described in Preparation 45a. 0.26 g (14% yield) of the desired compound were obtained.
LRMS (*m/z*): 501, 503 (*M*+1)⁺.

b) (S)-*tert*-Butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate

[0393] The title compound was prepared from (*S*)-*tert*-butyl 1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxobutan-2-ylcarbamate (263 mg, 0.52 mmol) following the experimental procedure described in Preparation 45b. 113 mg (44% yield) of the desired compound were obtained.
LRMS (*m/z*): 483, 485 (*M*+1)⁺.

c) (S)-*tert*-Butyl 1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate

[0394] The title compound was prepared from (*S*)-*tert*-butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate (113 mg, 0.23 mmol) following the experimental procedure described in Preparation 47a. 73 mg (71% yield) of the desired compound were obtained.
LRMS (*m/z*): 430 (*M*+1)⁺.

d) (S)-2-(1-Aminopropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0395] The title compound was prepared from (*S*)-*tert*-butyl 1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate (73 mg, 0.17 mmol) following the experimental procedure described in Preparation 47b. 62 mg (100% yield) of the desired compound were obtained.

LRMS (m/z): 330 (M+1)⁺.

PREPARATION 96

(S)-2-(1-Aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxamide

[0396] A suspension of 60 mg (0.17 mmol) of (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile in sulphuric acid (2 ml) was stirred at room temperature overnight. The reaction mixture was slowly poured into a mixture of ice/water, neutralized with a 2N solution of NaHCO₃ and extracted with ethyl acetate. The organic layer was then washed with brine, dried over magnesium sulphate and concentrated. The title compound was obtained as a white solid (38mg, 64% yield).
LRMS (m/z): 334 (M+1)⁺.

PREPARATION 97

2-Amino-4-chloropyrimidine-5-carbonitrile

[0397] To a solution of 2,4-dichloropyrimidine-5-carbonitrile (600mg, 3.45mmol) in dioxane (20ml) was added a 0.5M solution of NH₃ in dioxane (20ml, 10 mmol) and the mixture stirred at room temperature for 4h. A mixture of two isomers were obtained and separated by column chromatography using a mixture of hexane/ethyl acetate (from 0% to 45% of ethyl acetate). The title compound (304mg, 56% yield) was found to be the less polar isomer.
LRMS (m/z): 153 (M-1)⁻.

PREPARATION 98

2,4-Diamino-6-chloropyrimidine-5-carbonitrile

[0398] A mixture of 2,4,6-trichloropyrimidine-5-carbonitrile (200mg, 0.96 mmol) and a 0.5M solution of NH₃ in dioxane (12ml, 6 mmol) and the mixture stirred at room temperature for 2h. The monosubstituted intermediate was obtained and 10 ml more of NH₃ in dioxane were added and the mixture stirred at 80°C over the weekend. A suspension was obtained, the solid filtered off and the filtrate concentrated to give a solid which was triturated in diethyl ether. The title compound was obtained as a beige solid (156mg, 78% yield).
LRMS (m/z): 170 (M+1)⁺.

PREPARATION 99

(S)-2-(1-Aminoethyl)-3-phenyl-5-(thiazol-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (S)-*tert*-Butyl 1-(4-oxo-3-phenyl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0399] 100 mg (0.23 mmol) of (S)-*tert*-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate, 95μl (0.3 mmol) of 2-(tributylstannyl)thiazole and 8 mg (0.01 mmol) of tetrakis(triphenylphosphine)palladium (0) under argon were stirred in dimethylformamide (2ml) at 100°C overnight. Then 95μl (0.3 mmol) of 2-(tributylstannyl)thiazole and 8 mg (0.01 mmol) of tetrakis(triphenylphosphine) palladium (0) were added and the mixture was stirred at 100°C for 2 more days. The crude was filtered over celite washing with ethyl acetate. Then the organic phase was washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by normal phase chromatography (hexane-diethyl ether, 0-60% in 30CV) to obtain the title compound (48 mg, 45% yield) as a white solid.
LRMS (m/z): 438(M+1)⁺.

b) (S)-2-(1-Aminoethyl)-3-phenyl-5-(thiazol-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0400] This compound was prepared starting from (S)-*tert*-Butyl 1-(4-oxo-3-phenyl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (48 mg, 0.11 mmol) and following the experimental procedure described in Preparation 46c to afford 26 mg (70% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 338 (M+1)⁺.

PREPARATION 100**(S)-tert-Butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0401] The title compound was prepared from (S)-tert-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (245 mg, 0.57 mmol) and ethenyl(tributyl)tin (214 μ l, 0.74 mmol) following the experimental procedure described in Preparation 99a. 144 mg (65% yield) of the desired compound were obtained.
LRMS (m/z): 381 (M+1)⁺.

PREPARATION 101**(S)-tert-Butyl 1-(5-formyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0402] (S)-tert-butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (120mg, 0.32 mmol) in a mixture of acetone/water (25ml, 95/5) was ozonolysed in a Sander Labor-Ozonisator (300.5) at -20°C with an air flow of 20 l/h and 40 mA for 20 min. Reaction mixture was concentrated and the crude was purified by reverse phase chromatography to yield the final compound as a white solid (93mg, 77% yield).
LRMS (m/z): 383 (M+1)⁺.

PREPARATION 102**(S)-2-(1-Aminoethyl)-5-(morpholinomethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (S)-tert-Butyl 1-(5-(morpholinomethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0403] To a solution of (S)-tert-butyl 1-(5-formyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (93 mg, 0.24 mmol) in methanol (9 ml), was added morpholine (27 μ l, 0.31 mmol) and acetic acid (58 μ l, 1.01mmol) and the reaction mixture was stirred 3h at room temperature. Then sodium cyanoborohydride (10mg, 0.16 mmol) was added and the reaction mixture stirred overnight at room temperature. The solvents were evaporated and the residue was partitioned between water and ethyl acetate. The organic layer was washed with 4% NaHCO₃ and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. The product was obtained as an oil (125mg, 99% yield)
LRMS (m/z): 454 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-5-(morpholinomethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0404] This compound was prepared starting from (S)-tert-butyl 1-(5-(morpholinomethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (125 mg, 0.24 mmol) and following the experimental procedure described in Preparation 46c to afford 95 mg (86% yield) of the title compound as a dihydrochloride salt that was used in the next step without any further purification.
LRMS (m/z): 354 (M+1)⁺.

PREPARATION 103**(S)-2-(1-(tert-Butoxycarbonylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxylic acid**

[0405] (S)-tert-Butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (108mg, 0.28 mmol) in a mixture of ethyl acetate/pyridine (15 mL, 80/20) was ozonolysed in a Sander Labor-Ozonisator (300.5) at -25°C with an air flow of 15 l/h and 30 mA for 10h. Reaction mixture was concentrated and the crude was purified by normal phase chromatography (hexane/ethyl acetate) to yield the final compound (26mg, 23% yield).
LRMS (m/z): 399 (M+1)⁺.

PREPARATION 104

(S)-2-(1-Aminoethyl)-5-(1-methyl-1H-pyrazol-4-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**a) (S)-tert-Butyl 1-(5-(1-methyl-1H-pyrazol-4-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0406] 50 mg (0.12 mmol) of (S)-tert-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate, 48 mg (0.23 mmol) of 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole, 8 mg (0.01 mmol) of tetrakis(triphenylphosphine)palladium(0) and 184 μ l of sodium carbonate 2M in water under argon were stirred in dimethylformamide (1 ml) at 120°C for 2 h. The crude was filtered over celite washing with ethyl acetate. Then the organic phase was washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%) to obtain the title compound (50 mg, 87% yield) as a white solid.

LRMS (m/z): 435 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-5-(1-methyl-1H-pyrazol-4-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0407] This compound was prepared starting from (S)-tert-butyl 1-(5-(1-methyl-1H-pyrazol-4-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (50 mg, 0.1 mmol) and following the experimental procedure described in Preparation 46c to afford 50 mg (100% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 335 (M+1)⁺.

PREPARATION 105

(S)-2-(1-Aminoethyl)-5-(2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**a) (S)-tert-Butyl 1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0408] 100 mg (0.26 mmol) of (S)-tert-butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate in tetrahydrofuran (8 ml) were cooled to 0°C in an ice bath. Then 3.15 ml (1.58 mmol) of 9-BBN (0.5M in THF) were slowly added. The reaction mixture was led at 0°C for an additional hour and then 4 h at room temperature. Then it was cooled at 0°C and 1.7 ml (3.4 mmol) of sodium hydroxide 2M and 3.8 ml (0.03 mmol) of hydrogen peroxide (35% in water) were added. The reaction mixture was stirred at room temperature for 2 h.

LRMS (m/z): 399 (M+1)⁺.

b) (S)-2-(1-aminoethyl)-5-(2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0409] This compound was prepared starting from (S)-tert-butyl 1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (160 mg, 0.2 mmol) and following the experimental procedure described in Preparation 46c to afford 160 mg (100% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 299 (M+1)⁺.

PREPARATION 106

(S)-2-(1-Aminoethyl)-5-bromo-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0410] This compound was prepared starting from (S)-tert-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (50 mg, 0.12 mmol) and following the experimental procedure described in Preparation 46c to afford 42 mg (100% yield) of the title compound that was used in the next step without any further purification.

LRMS (m/z): 333, 335 (M+1)⁺.

PREPARATION 107

(S)-2-(1-Aminoethyl)-5-(2-methoxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

a) (S)-*tert*-Butyl 1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0411] To a solution of (S)-*tert*-butyl 1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (47mg, 0.12mmol) in tetrahydrofuran (1 mL), sodium hydride (6 mg, 0.15mmol) was added. The reaction mixture was stirred at room temperature for 10 min and then methyl iodide (11 μ l, 0.18 mmol) was added. After overnight stirring at room temperature ethyl acetate was added and the organic layer washed with water and brine. The organic layer was then dried over magnesium sulphate, filtered and concentrated. The crude was used without further purification

LRMS (m/z): 413 (M+1)⁺

b) (S)-2-(1-Aminoethyl)-5-(2-methoxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0412] This compound was prepared starting from (S)-*tert*-butyl 1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (50mg, 0.08 mmol) and following the experimental procedure described in Preparation 46c to afford 50 mg (100% yield) of the hydrochloride salt of the title compound, that was used in the next step without any further purification.

LRMS (m/z): 313 (M+1)⁺.

PREPARATION 108

(S)-*tert*-Butyl 1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

a) (S)-*tert*-Butyl 1-(3-bromo-2-(3-(trifluoromethyl)phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0413] The title compound was prepared from (S)-methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (2 g, 5.13 mmol) following the experimental procedure described in Preparation 27a. 2.35 g (81% yield) of the desired compound were obtained.

LRMS (m/z): 520 (M+1)⁺.

b) (S)-*tert*-Butyl 1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0414] The title compound was prepared from (S)-*tert*-butyl 1-(3-bromo-2-(3-(trifluoromethyl)phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (2.30 g, 4.43 mmol) following the experimental procedure described in Preparation 54b. 1.40 g (63% yield) of the desired compound were obtained.

LRMS (m/z): 502 (M+1)⁺.

PREPARATION 109

Methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate

a) Methyl 1-amino-1*H*-pyrrole-2-carboxylate

[0415] Sodium hydride (4.40 g, 0.11 mol, 60% in hexanes) was suspended in DMF (550 ml) under nitrogen atmosphere. Once cooled at -5°C, methyl 1*H*-pyrrole-2-carboxylate (11.0 g, 0.09 mol) dissolved in DMF (182 ml) was dropwise added and vigorously stirred for 30. 277ml more of DMF was added and then O-(diphenylphosphoryl)hydroxylamine (32.8g, 0.14 mol) was introduced into the reaction mixture. The reaction mixture was stirred at room temperature for 4h. Once the reaction is over, 1l of saturated sodium thiosulfate solution (x5H₂O) was added and the mixture was warmed to 80°C for 1 h. Once at room temperature, 1l of ethyl ether was added and the phases separated. The aqueous phase was twice extracted with ethyl ether. The organic phase was washed with water and brine, dried over magnesium sulphate, filtered and the solvent evaporated under reduced pressure. 10.41 g (81.1% yield) of the final compound were obtained.

LRMS (m/z): 141 (M+1)⁺

b) Methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate

- 5 **[0416]** Methyl 1-amino-1*H*-pyrrole-2-carboxylate (6g, 48.1 mmol) and (*S*)-2-(*tert*-butoxycarbonylamino)propanoic acid (8.10g, 48.1 mmol) were dissolved in ethyl acetate (40 ml) and cooled in an ice-bath. Under an argon atmosphere, diisopropylethylamine (24.6 ml, 141.2 mmol) was added and, after stirring for 15 min, T3P solution (35.7 ml, 60 mmol, 50% in ethyl acetate) was dropwise added. After stirring for 20 min at 0°C, the reaction was left overnight at room temperature. The reaction mixture was poured onto water and extracted with ethyl ether. The organic phase was washed
10 with water and brine, dried over magnesium sulphate and the solvent evaporated under reduced pressure. 8.6g (83% yield) of the final product was obtained and use in the next synthetic step without further purification.
LRMS (m/z): 312 (M+1)⁺

PREPARATION 110

- 15 **(*S*)-*tert*-Butyl 1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

a) (*S*)-*tert*-Butyl 1-(3-bromo-2-(3-methoxyphenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

- 20 **[0417]** The title compound was prepared from (*S*)-methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (2 g, 5.13 mmol) following the experimental procedure described in Preparation 27a. 2.10 g (83% yield) of the desired compound were obtained.
LRMS (m/z): 481, 483 (M+1)⁺.

- 25 **b) (*S*)-*tert*-Butyl 1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

- [0418]** The title compound was prepared from (*S*)-*tert*-butyl 1-(3-bromo-2-(3-methoxyphenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (2.10 g, 4.36 mmol) following the experimental procedure described in Preparation
30 54b. 1.00 g (50% yield) of the desired compound were obtained.
LRMS (m/z): 463, 465 (M+1)⁺.

PREPARATION 111

- 35 **(*S*)-2-(1-Aminoethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

a) (*S*)-*tert*-Butyl 1-(5-cyano-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

- 40 **[0419]** The title compound was prepared from (*S*)-*tert*-butyl 1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (400 mg, 0.80 mmol) following the experimental procedure described in Preparation 47a. 228 mg (64% yield) of the desired compound were obtained.
LRMS (m/z): 448 (M+1)⁺.

- 45 **b) (*S*)-2-(1-Aminoethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

- [0420]** The dihydrochloride salt of the title compound was prepared from (*S*)-*tert*-butyl 1-(5-cyano-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (228 mg, 0.51 mmol) following the experimental procedure described in Preparation 47b. 229 mg (100% yield) of the desired compound were obtained.
50 LRMS (m/z): 348 (M+1)⁺.

55

PREPARATION 112

(S)-2-(1-Aminoethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**a) (S)-*tert*-Butyl 1-(5-cyano-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0421] The title compound was prepared from (S)-*tert*-butyl 1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (453 mg, 0.98 mmol) following the experimental procedure described in Preparation 47a. 197 mg (49% yield) of the desired compound were obtained. LRMS (m/z): 410 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0422] The dihydrochloride salt of the title compound was prepared from (S)-*tert*-butyl 1-(5-cyano-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (195 mg, 0.48 mmol) following the experimental procedure described in Preparation 47b. 250 mg (100% yield) of the desired compound were obtained. LRMS (m/z): 310 (M+1)⁺.

PREPARATION 113

2-(Chloromethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) Methyl 1-(phenylsulfonyl)-3-vinyl-1*H*-pyrrole-2-carboxylate.**

[0423] Tetrakis(triphenylphosphane) palladium(0) (170mg, 0.15 mmol) was added to a solution of 3-bromo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate (2 g, 5.8 mmol) and ethenyl(tributyl)tin (2.3 ml, 7.6 mmol) in *N,N*-dimethylformamide (60 mL). The resulting mixture was stirred for 23 h under Ar at 100 °C, cooled and evaporated in vacuum. The residue was dissolved in a saturated solution of potassium fluoride in methanol and stirred for 2 hours. The mixture was evaporated in vacuum and the product was purified by flash chromatography silica (hexane/ethyl acetate) to give a 1640 mg (97% yield) of the title compound.

b) Methyl 3-formyl-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate

[0424] To a solution of methyl 1-(phenylsulfonyl)-3-vinyl-1*H*-pyrrole-2-carboxylate (1640 mg, 5.63 mmol) in 45 ml of tetrahydrofuran were added 4-methylmorpholine 4-oxide (1.36 g, 11.3 mmol) and 2.4 ml (0.39 mmol) of a 4% aqueous solution of osmium tetroxide and the reaction was leaved with stirring at room temperature overnight. Afterwards, the starting material was completely consumed and the reaction mixture was filtered through a pad of Celite® using tetrahydrofuran. The filtrate was evaporated to dryness, taken up with ethyl acetate and washed with water and brine. The organic layer was dried (Na₂SO₄) and concentrated to give a brown residue that was immediately submitted to the next step. In this sense, this residue was dissolved in tetrahydrofuran (28 ml) and water (3.4 ml) and solid sodium periodate (1.8 g, 8.4 mmol) was added stirring the reaction vigorously at room temperature overnight. Next day, a suspension was formed and reaction was finished. The work-up was done by adding 4% aqueous solution of sodium bicarbonate (200 ml) and extracting with ethyl acetate (3x). The organic mixture was washed with water and brine, dried (Na₂SO₄) and concentrated to dryness to give a residue that was purified by flash chromatography silica (hexane/ethyl acetate) to give a 1.46 g (88% yield) of the title compound.

c) Methyl 3-(difluoromethyl)-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate

[0425] To a solution of methyl 3-formyl-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate (1.46 g, mmol) in dry dichloromethane (25 ml) in a schlenck flask at -75°C under Ar was added diethylaminosulfur trifluoride (DAST) (1.64 ml, 12.5 mmol) and the mixture was allowed to warm-up to room temperature during 3 h and then stirred at this temperature overnight. Next day, the reaction was finished and 200 ml of 4% aqueous solution of sodium bicarbonate were added carefully maintaining a vigorous stirring during 20 minutes. Afterwards, the mixture was extracted with ethyl acetate (3x) and the organic layers were washed with brine, dried (Na₂SO₄) and concentrated in vacuum to give a residue that was purified by flash chromatography silica (hexane/ethyl acetate) to give a 1.4 g (85% yield) of the title compound.

d) 3-(Difluoromethyl)-N-phenyl-1-(phenylsulfonyl)-1H-pyrrole-2-carboxamide

[0426] The title compound was prepared from methyl 3-(difluoromethyl)-1-(phenylsulfonyl)-1H-pyrrole-2-carboxylate (1.16 g, 3.7 mmol) following the experimental procedure described in Preparation 44a. 0.96 g (70% yield) of the desired compound were obtained.
LRMS (m/z): 377 (M+1)⁺.

e) 3-(Difluoromethyl)-N-phenyl-1H-pyrrole-2-carboxamide

[0427] The title compound was prepared from 3-(difluoromethyl)-N-phenyl-1-(phenylsulfonyl)-1H-pyrrole-2-carboxamide (0.96 g, 2.6 mmol) following the experimental procedure described in Preparation 44b. 0.6 g (98% yield) of the desired compound were obtained.
LRMS (m/z): 237 (M+1)⁺.

f) 1-Amino-3-(difluoromethyl)-N-phenyl-1H-pyrrole-2-carboxamide

[0428] 1.1 ml (1.1 mmol) of a 1M solution of lithium bis(trimethylsilyl)amide was added to a solution of 3-(difluoromethyl)-N-phenyl-1H-pyrrole-2-carboxamide (100 mg, 0.42 mmol) and DPPONH₂ (*P,P*-diphenylphosphinic amide, available from Sigma Aldrich®, cat. no. 5994-87-6) (250 mg, 1.1 mmol) in DMF (4 mL) at room temperature. A thick suspension formed and additional 4 mL of DMF were added. The mixture was stirred at room temperature for 1 hour and then it was poured into 50 mL of water and extracted with ethyl acetate. The aqueous layer was further extracted with ethyl acetate (2x). The combined organic layer was washed with water and brine and was dried over sodium sulphate, filtered and concentrated. The crude product was purified by flash chromatography (0% to 33% AcOEt/Hexanes) to yield 40 mg (38% yield) of the title compound.
LRMS (m/z): 252 (M+1)⁺.

g) 2-(Chloromethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0429] The title compound was prepared from 1-amino-3-(difluoromethyl)-N-phenyl-1H-pyrrole-2-carboxamide (0.167 g, 0.66 mmol) following the experimental procedure described in Preparation 2. 57 mg (28% yield) of the desired compound were obtained.
LRMS (m/z): 310 (M+1)⁺.

PREPARATION 114**(S)-2-(1-Aminoethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (S)-*tert*-Butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0430] Tetrakis(triphenylphosphane) palladium(0) (90mg, 0.01 mmol) was added to a solution of (S)-*tert*-butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (100 mg, 0.23 mmol) and ethenyl(tributyl)tin (90 µl, 0.31 mmol) in *N,N*-dimethylformamide (2.4 mL). The resulting mixture was stirred for 24 h under Ar at 100 °C, cooled and evaporated in vacuum. The residue was dissolved in a saturated solution of potassium fluoride in methanol and stirred for 1 hour. The mixture was evaporated in vacuum and the product was purified by flash chromatography silica (hexane/ethyl acetate) to give a 65 mg (74% yield) of the title compound.

b) (S)-*tert*-Butyl 1-(5-formyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0431] To a solution of *tert*-butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (37 mg, 0.1 mmol) in 0.8 ml of tetrahydrofuran were added 4-methylmorpholine 4-oxide (23 mg, 0.2 mmol) and 42 µl (0.01 mmol) of a 4% aqueous solution of osmium tetroxide and the reaction was leaved with stirring at room temperature overnight. Afterwards, the starting material was completely consumed and the reaction mixture was filtered through a pad of Celite® using tetrahydrofuran. The filtrate was evaporated to dryness, taken up with ethyl acetate and washed with water and brine. The organic layer was dried (Na₂SO₄) and concentrated to give a brown residue that was immediately submitted to the next step. In this sense, this residue was dissolved in tetrahydrofuran (0.5 ml) and water (60 µl) and solid sodium periodate (31 mg, 0.14 mmol) was added stirring the reaction vigorously at room temperature overnight. Next day, a suspension was formed and reaction was finished. The work-up was done by adding 4% aqueous solution of sodium bicarbonate and extracting with ethyl acetate (3x). The organic mixture was washed with water and brine,

dried (Na_2SO_4) and concentrated to dryness to give a residue that was purified by flash chromatography silica (hexane/ethyl acetate) to give a 26 g (70% yield) of the title compound.

c) (S)-tert-Butyl 1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0432] To a solution of *tert*-butyl 1-(5-formyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (400 mg, 1.05 mmol) in dry dichloromethane (10 ml) in a schlenk flask at -75°C under Ar was added diethylaminosulfur trifluoride (DAST) (1 ml, 7.63 mmol) and the mixture was allowed to warm-up till room temperature during 3 h and then stirred at this temperature overnight. Next day, the reaction was finished and 4% aqueous solution of sodium bicarbonate were added carefully maintaining a vigorous stirring during 20 minutes. Afterwards, the mixture was extracted with ethyl acetate (3x) and the organic layers were washed with brine, dried (Na_2SO_4) and concentrated in vacuum to give a residue that was purified by flash chromatography silica (hexane/ethyl acetate) to give a 316 mg (75% yield) of the title compound.

d) (S)-2-(1-Aminoethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0433] The title compound was prepared from *tert*-butyl 1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (316 mg, 0.78 mmol) following the experimental procedure described in Preparation 46c. 219 mg (92% yield) of the desired compound were obtained.

LRMS (*m/z*): 305 ($\text{M}+1$)⁺.

PREPARATION 115

(R)-2-(1-Amino-2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (R)-tert-Butyl 3-tert-butoxy-1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate

[0434] The title compound was prepared from 1-amino-*N*-phenyl-1*H*-pyrrole-2-carboxamide (1 g, 5 mmol) following the experimental procedure described in Preparation 42a. 0.72 g (32% yield) of the desired compound was obtained.

LRMS (*m/z*): 445 ($\text{M}+1$)⁺.

b) (R)-tert-Butyl 2-tert-butoxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0435] The title compound was prepared from (*R*)-*tert*-butyl 3-tert-butoxy-1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate (0.62 g, 1.4 mmol) following the experimental procedure described in Preparation 42b. 0.12 g (20% yield) of the desired compound was obtained.

LRMS (*m/z*): 427 ($\text{M}+1$)⁺.

c) (R)-2-(1-Amino-2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0436] 110 mg (0.26 mmol) of (*R*)-*tert*-butyl 2-tert-butoxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate were dissolved in 40 μl of methylene chloride and 40 μl of trifluoroacetic acid were added. The resulting solution was stirred at room temperature overnight and the reaction mixture was evaporated to dryness. The residue was then redissolved in dioxane (2ml) and the solution was treated with 4M HCl solution in dioxane (0.5 ml) stirring 10 minutes at this temperature and evaporated to dryness to obtain the corresponding hydrochloride 75 mg (91%) of the title compound.

LRMS (*m/z*): 271 ($\text{M}+1$)⁺.

PREPARATION 116

(S)-2-(1-Aminoethyl)-6-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) 4-Fluoro-*N*-phenyl-1*H*-pyrrole-2-carboxamide

[0437] The title compound was prepared from methyl 4-fluoro-1*H*-pyrrole-2-carboxylate³ (1.14 g, 8.0 mmol) and aniline (2.2 ml, 24.0 mmol) following the experimental procedure described in Preparation 44a. 1.09 g (64% yield) of the desired compound were obtained.

LRMS (*m/z*): 205 ($\text{M}+1$)⁺.

b) 1-Amino-4-fluoro-N-phenyl-1H-pyrrole-2-carboxamide

[0438] This compound was prepared starting from 4-fluoro-N-phenyl-1H-pyrrole-2-carboxamide (1.05 g, 5.1 mmol) and following the experimental procedure described in Preparation 1b to afford 1.14 g (84% yield) of the title compound.
LRMS (m/z): 220 (M+1)⁺.

c) (S)-tert-Butyl 1-(4-fluoro-2-(phenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0439] The title compound was prepared from 1-amino-4-fluoro-N-phenyl-1H-pyrrole-2-carboxamide (0.95 g, 3.6 mmol) and (S)-2-(tert-butoxycarbonylamino)propanoic acid following the experimental procedure described in Preparation 47b. 1.28 g (91% yield) of the desired compound were obtained.
LRMS (m/z): 391 (M+1)⁺.

d) (S)-tert-Butyl 1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0440] This compound was prepared starting from (S)-tert-butyl 1-(4-fluoro-2-(phenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (1.15 g, 2.95 mmol) and following the experimental procedure described in Preparation 73b to afford 0.68 g (62% yield) of the title compound.
LRMS (m/z): 373 (M+1)⁺.

e) (S)-2-(1-Aminoethyl)-6-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0441] This compound was prepared starting from (S)-tert-butyl 1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (0.64 g, 1.64 mmol) and following the experimental procedure described in Preparation 71b but heating the reaction mixture at 35°C during 5 hours to afford 0.59 g (96% yield) of the title compound, isolated as the trifluoroacetate salt form.
LRMS (m/z): 273 (M+1)⁺.

PREPARATION 117**2-((S)-1-Aminoethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one****a) (S)-N-(1-Phenylethyl)-1H-pyrrole-2-carboxamide**

[0442] The title compound was prepared from methyl 1H-pyrrole-2-carboxylate (8.0 g, 0.06 mol) and (S)-1-phenylethylamine (24.7 ml, 0.19 mol) following the experimental procedure described in Preparation 44a. 10.4 g (75% yield) of the desired compound were obtained.
LRMS (m/z): 215 (M+1)⁺.

b) (S)-1-Amino-N-(1-phenylethyl)-1H-pyrrole-2-carboxamide

[0443] This compound was prepared starting from (S)-N-(1-phenylethyl)-1H-pyrrole-2-carboxamide (11.5 g, 0.05 mol) and following the experimental procedure described in Preparation 1b to afford 9.8 g (79% yield) of the title compound.
LRMS (m/z): 230 (M+1)⁺.

c) tert-Butyl (S)-1-oxo-1-(2-((S)-1-phenylethylcarbamoyl)-1H-pyrrol-1-ylamino)propan-2-ylcarbamate

[0444] The title compound was prepared from (S)-1-amino-N-(1-phenylethyl)-1H-pyrrole-2-carboxamide (3.50 g, 15.3 mmol) and (S)-2-(tert-butoxycarbonylamino)propanoic acid (2.89 g, 15.3 mmol) following the experimental procedure described in Preparation 73a. 5.78 g (92% yield) of the desired compound were obtained.

d) tert-Butyl (S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0445] This compound was prepared starting from tert-butyl (S)-1-oxo-1-(2-((S)-1-phenylethylcarbamoyl)-1H-pyrrol-1-ylamino)propan-2-ylcarbamate (5.23 g, 13.1 mmol) and following the experimental procedure described in Preparation 68b to afford 4.40 g (85% yield) of the title compound.
LRMS (m/z): 383 (M+1)⁺.

e) 2-((S)-1-Aminoethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0446] This compound was prepared starting from *tert*-butyl (S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (4.34 g, 11.4 mmol) and following the experimental procedure described in Preparation 71 b but heating the reaction mixture at 35°C during 5 hours to afford 4.27 g (95% yield) of the title compound, isolated as the trifluoroacetate salt form.
LRMS (m/z): 283 (M+1)⁺.

PREPARATION 118**(S)-2-(1-Aminoethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) *N*-(2,6-Dimethylphenyl)-1*H*-pyrrole-2-carboxamide**

[0447] The title compound was prepared from 1*H*-pyrrole-2-carboxylic acid (2 g, 18 mmol) and 2,6-dimethylaniline (2.75 g, 22.7 mmol) following the experimental procedure described in Preparation 44a. 900 mg (23% yield) of the desired compound were obtained.
LRMS (m/z): 215 (M+1)⁺.

b) 1-Amino-*N*-(2,6-dimethylphenyl)-1*H*-pyrrole-2-carboxamide

[0448] This compound was prepared starting from *N*-(2,6-dimethylphenyl)-1*H*-pyrrole-2-carboxamide (1080 mg, 5.04 mmol) and following the experimental procedure described in Preparation 1 b to afford 324 mg (25% yield) of the title compound.
LRMS (m/z): 230 (M+1)⁺.

c) (S)-Benzyl 1-(2-(2,6-dimethylphenylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0449] The title compound was prepared from 1-amino-*N*-(2,6-dimethylphenyl)-1*H*-pyrrole-2-carboxamide (324 mg, 1.41 mmol) and (S)-2-(benzyloxycarbonylamino)propanoic acid (347 g, 1.55 mmol) following the experimental procedure described in Preparation 73a. 458 mg (75% yield) of the desired compound were obtained.
LRMS (m/z): 435 (M+1)⁺.

d) (S)-Benzyl 1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0450] This compound was prepared starting from (S)-benzyl 1-(2-(2,6-dimethylphenyl-carbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (458 mg, 1.05 mmol) and following the experimental procedure described in Preparation 68b to afford 252 mg (58% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 417 (M+1)⁺.

e) (S)-2-(1-Aminoethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0451] This compound was prepared starting from (S)-benzyl 1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (126 mg, 0.3 mmol) and following the experimental procedure described in Preparation 128c to afford 85 mg (100% yield) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 283 (M+1)⁺.

PREPARATION 119**(S)-2-(Aminomethyl)-3-(1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (S)-*tert*-Butyl 2-oxo-2-(2-(1-phenylethylcarbamoyl)-1*H*-pyrrol-1-ylamino)ethylcarbamate**

[0452] The title compound was prepared from (S)-1-amino-*N*-(1-phenylethyl)-1*H*-pyrrole-2-carboxamide (3.50 g, 15.3 mmol) and 2-(*tert*-butoxycarbonylamino)acetic acid (2.67 g, 15.3 mmol) following the experimental procedure described in Preparation 73a. 5.79 g (89% yield) of the desired compound were obtained.
LRMS (m/z): 387 (M+1)⁺.

b) (S)-tert-Butyl (4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)methylcarbamate

[0453] This compound was prepared starting from (S)-tert-butyl 2-oxo-2-(2-(1-phenylethylcarbamoyl)-1*H*-pyrrol-1-ylamino)ethylcarbamate (5.75 g, 14.9 mmol) and following the experimental procedure described in Preparation 68b to afford 2.34 g (35% yield, 81% purity) of the title compound that was used in the next step without any further purification. LRMS (m/z): 369 (M+1)⁺.

c) (S)-2-(Aminomethyl)-3-(1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0454] This compound was prepared starting from (S)-tert-butyl (4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)methylcarbamate (2.30 g, 81% purity, 5.1 mmol) and following the experimental procedure described in Preparation 71b but heating the reaction mixture at 35°C during 4 hours to afford 2.22 g (75% yield, 65% purity) of the title compound, isolated as the trifluoroacetate salt form. LRMS (m/z): 269 (M+1)⁺.

PREPARATION 120**(S)-2-(1-Aminoethyl)-5-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (S)-tert-Butyl 1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0455] To a solution of (S)-tert-Butyl 1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (500 mg, 1.15 mmol) in anhydrous THF (5 ml) placed in a schlenk tube under Ar was added dropwise a 1.6 M solution of n-BuLi in hexanes (1.8 ml, 2.88 mmol) at -78°C. This mixture was stirred during 30 min. at -78°C in order to accomplish the halogen-metal exchange. After this period, a solution of *N*-fluoro-*N*-(phenylsulfonyl)-benzenesulfonamide (475 mg, 1.5 mmol) in THF (4 ml) was added dropwise and the reaction mixture was allowed to warm up overnight till room temperature and then it was quenched by addition of a saturated solution of ammonium chloride (15 ml). Some additional water was added and the mixture was extracted with ethyl acetate (3x). The total organic phase was washed with brine, dried (sodium sulphate) and concentrated in vacuum to get 606 mg of a residue. This crude material was purified by flash chromatography (0% to 20% dichloromethane/AcOEt) to yield 78 mg (18% yield) of the title compound. LRMS (m/z): 373 (M+1)⁺.

b) 2-(1-Aminoethyl)-5-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0456] This compound was prepared starting from (S)-tert-butyl 1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (78 mg, 0.21 mmol) and following the experimental procedure described in Preparation 81b to afford 65 mg (quantitative yield) of the title compound, isolated as the hydrochloric salt form. LRMS (m/z): 273 (M+1)⁺.

PREPARATION 121**2-((S)-1-Aminoethyl)-5-(1,2-dihydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) tert-Butyl (1*S*)-1-(5-(1,2-dihydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0457] To a solution of (S)-tert-butyl 1-(4-oxo-3-phenyl-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (1.85 g, 4.9 mmol) in 40 ml of tetrahydrofuran were added 4-methylmorpholine 4-oxide (1.15 g, 9.8 mmol) and 4% aqueous solution of osmium tetroxide (2.1 ml, 0.3 mmol) and the reaction was leaved with stirring at room temperature overnight. Afterwards, the starting material was completely consumed and the reaction mixture was filtered through a pad of Celite® using tetrahydrofuran. The filtrate was evaporated to dryness, taken up with ethyl acetate and washed with water and brine. The organic layer was dried (Na₂SO₄) and concentrated to afford 1.98 g (98% yield) of the title compound. LRMS (m/z): 415 (M+1)⁺.

b) 2-((S)-1-Aminoethyl)-5-(1,2-dihydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0458] This compound was prepared starting from tert-butyl (1*S*)-1-(5-(1,2-dihydroxyethyl)-4-oxo-3-phenyl-3,4-dihy-

dropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (200 mg, 0.5 mmol) and following the experimental procedure described in Preparation 81b to afford 112 mg (66% yield) of the title compound, isolated as the hydrochloric salt form. LRMS (m/z): 315 (M+1)⁺.

PREPARATION 122

(S)-2-(1-Aminoethyl)-5-(hydroxymethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (S)-*tert*-Butyl 1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0459] To a solution of (S)-*tert*-butyl 1-(5-formyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (400 mg, 1.1 mmol) in 20 mL of methanol was added NaBH₄ (30 mg, 0.8 mmol) and the mixture was stirred at room temperature during 3.5 h. The solvent was evaporated and the residue was partitioned between ammonium chloride saturated aqueous solution and ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. The product was purified by flash chromatography (0% to 30% hexane/AcOEt) to yield 301 mg (75%) of the title compound. LRMS (m/z): 385 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-5-(hydroxymethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0460] This compound was prepared starting from (S)-*tert*-butyl 1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (300 mg, 0.8 mmol) and following the experimental procedure described in Preparation 81 b to afford 233 mg (93% yield) of the title compound, isolated as the hydrochloric salt form. LRMS (m/z): 285 (M+1)⁺.

PREPARATION 123

(S)-2-(1-Aminoethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (S)-*tert*-Butyl 1-(3-(3,5-difluorophenyl)-4-oxo-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0461] This compound was prepared starting from (S)-*tert*-butyl 1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (1.00 g, 2.1 mmol) and following the experimental procedure described in Preparation 114a to afford 0.71 g (76% yield) of the title compound. LRMS (m/z): 417 (M+1)⁺.

b) (S)-*tert*-Butyl 1-(3-(3,5-difluorophenyl)-5-formyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0462] This compound was prepared starting from (S)-*tert*-butyl 1-(3-(3,5-difluorophenyl)-4-oxo-5-vinyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (0.70 g, 1.7 mmol) and following the experimental procedure described in Preparation 114b to afford 0.62 g (88% yield) of the title compound. LRMS (m/z): 419 (M+1)⁺.

c) (S)-*tert*-Butyl 1-(5-(difluoromethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0463] This compound was prepared starting from (S)-*tert*-butyl 1-(3-(3,5-difluorophenyl)-5-formyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (0.62 g, 1.5 mmol) and following the experimental procedure described in Preparation 114c to afford 0.45 g (68% yield) of the title compound. LRMS (m/z): 441 (M+1)⁺.

d) (S)-2-(1-Aminoethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0464] This compound was prepared starting from (S)-*tert*-butyl 1-(5-(difluoromethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (0.45 g, 1.0 mmol) and following the experimental procedure de-

scribed in Preparation 81b to afford 0.36 g (94% yield) of the title compound, isolated as the hydrochloric salt form.
LRMS (m/z): 341 (M+1)⁺.

PREPARATION 124

(S)-2-(1-Aminoethyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

a) (S)-*tert*-Butyl 1-(3-bromo-2-(2-chlorobenzylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0465] The title compound was prepared from (S)-Methyl 3-bromo-1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (310 mg, 0.8 mmol) and (2-chlorophenyl)methanamine (385 μ L, 3.2 mmol) following the experimental procedure described in Preparation 44a. 259 mg (65% yield) of the desired compound were obtained.
LRMS (m/z): 499, 501 (M+1)⁺.

b) (S)-*tert*-Butyl 1-(5-bromo-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0466] This compound was prepared starting from (S)-*tert*-butyl 1-(3-bromo-2-(2-chlorobenzylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (296 mg, 0.6 mmol) and following the experimental procedure described in Preparation 73b to afford 38 mg (13% yield) of the title compound.

LRMS (m/z): 481, 483 (M+1)⁺.

c) (S)-*tert*-Butyl 1-(3-(2-chlorobenzyl)-5-cyano-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0467] This compound was prepared starting from (S)-*tert*-butyl 1-(5-bromo-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (38 mg, 0.08 mmol) and following the experimental procedure described in Preparation 47a to afford 20 mg (59% yield) of the title compound.

LRMS (m/z): 428 (M+1)⁺.

d) (S)-2-(1-Aminoethyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0468] This compound was prepared starting from (S)-*tert*-butyl 1-(5-bromo-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (16 mg, 0.04 mmol) and following the experimental procedure described in Preparation 81b to afford 13 mg (95% yield) of the title compound, isolated as the hydrochloric salt form.

LRMS (m/z): 328 (M+1)⁺.

PREPARATION 125

(S)-Methyl 1-(2-(benzyloxycarbonylamino)propanamido)-3-bromo-1*H*-pyrrole-2-carboxylate

a) Methyl 3-bromo-1*H*-pyrrole-2-carboxylate

[0469] To a solution of methyl 3-bromo-1-(phenylsulfonyl)-1*H*-pyrrole-2-carboxylate² (6.59 g, 19.2 mmol) in 132 mL of methanol was added MeONa (1.55 g, 28.7 mmol) and the mixture was stirred at room temperature during 4 h. The solvent was evaporated and the residue was partitioned between ammonium chloride saturated aqueous solution and ethyl acetate. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and evaporated under vacuum. The product was purified by flash chromatography (0% to 30% hexane/AcOEt) to yield 3.32 g (85%) of the title compound.

LRMS (m/z): 203, 205 (M+1)⁺.

b) Methyl 1-amino-3-bromo-1*H*-pyrrole-2-carboxylate

[0470] This compound was prepared starting from methyl 3-bromo-1*H*-pyrrole-2-carboxylate (3.30 g, 16.2 mmol) and following the experimental procedure described in Preparation 35b to afford 3.42 g (48% yield, 50% purity) of the title compound.

LRMS (m/z): 218, 220 (M+1)⁺.

c) (S)-Methyl 1-(2-(benzyloxycarbonylamino)propanamido)-3-bromo-1H-pyrrole-2-carboxylate

[0471] The title compound was prepared from methyl 1-amino-3-bromo-1H-pyrrole-2-carboxylate (9.05 g, 25.6 mmol) and (S)-2-(benzyloxycarbonylamino)propanoic acid (5.72 g, 25.6 mmol) following the experimental procedure described in Preparation 73a. 9.73 g (84% yield) of the desired compound were obtained.

LRMS (m/z): 424, 426 (M+1)⁺.

PREPARATION 126**(S)-2-(1-Aminoethyl)-3-(5-fluoropyridin-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one****a) (S)-Benzyl 1-(3-bromo-2-(5-fluoropyridin-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0472] The title compound was prepared from (S)-methyl 1-(2-(benzyloxycarbonylamino)propanamido)-3-bromo-1H-pyrrole-2-carboxylate (2.00 g, 4.7 mmol) and 5-fluoropyridin-3-amine (4.23 g, 37.6 mmol) following the experimental procedure described in Preparation 44a. 1.65 g (69% yield) of the desired compound were obtained.

LRMS (m/z): 504, 506 (M+1)⁺.

b) (S)-Benzyl 1-(5-bromo-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0473] This compound was prepared starting from (S)-benzyl 1-(3-bromo-2-(5-fluoropyridin-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (1.50 g, 3.0 mmol) and following the experimental procedure described in Preparation 73b to afford 0.40 g (28% yield) of the title compound.

LRMS (m/z): 486, 488 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(5-fluoropyridin-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0474] This compound was prepared starting from (S)-benzyl 1-(5-bromo-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (100 mg, 0.21 mmol) and following the experimental procedure described in Preparation 127c to afford 53 mg (95% yield) of the title compound.

LRMS (m/z): 274 (M+1)⁺.

PREPARATION 127**(S)-2-(1-Aminoethyl)-3-(pyrimidin-5-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one****a) (S)-Benzyl 1-(3-bromo-2-(pyrimidin-5-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0475] The title compound was prepared from (S)-methyl 1-(2-(benzyloxycarbonylamino)propanamido)-3-bromo-1H-pyrrole-2-carboxylate (0.81 g, 1.91 mmol) and pyrimidin-5-amine (1.45 g, 15.3 mmol) following the experimental procedure described in Preparation 44a. 0.58 g (62% yield) of the desired compound were obtained.

LRMS (m/z): 487, 489 (M+1)⁺.

b) (S)-Benzyl 1-(5-bromo-4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0476] This compound was prepared starting from (S)-benzyl 1-(3-bromo-2-(pyrimidin-5-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (0.57 g, 1.17 mmol) and following the experimental procedure described in Preparation 73b to afford 0.33 g (61% yield) of the title compound.

LRMS (m/z): 469, 471 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(pyrimidin-5-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0477] To a solution of (S)-benzyl 1-(5-bromo-4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (40 mg, 0.09 mmol) in 4 mL of methanol were added Et₃N (24 μ L, 0.18 mmol) and Pd/C 10% (20 mg). The reaction was stirred at room temperature under hydrogen (30 psi) for 15 hours. The reaction mixture was filtered through Celite® and the filtrate was evaporated to dryness to give 20 mg (89%) of the title compound.

LRMS (m/z): 257 (M+1)⁺.

PREPARATION 128

(S)-2-(1-Amino-3-hydroxypropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (S)-*tert*-Butyl 4-(benzyloxy)-1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate**

[0478] The title compound was prepared from 1-amino-N-phenyl-1*H*-pyrrole-2-carboxamide (0.65 g, 3.2 mmol) and (S)-4-(benzyloxy)-2-(*tert*-butoxycarbonylamino)butanoic acid (1.0 g, 3.2 mmol) following the experimental procedure described in Preparation 73a. 1.57 g (96% yield) of the desired compound were obtained.

LRMS (m/z): 493 (M+1)⁺.

b) (S)-*tert*-Butyl 3-(benzyloxy)-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate

[0479] This compound was prepared starting from (S)-*tert*-butyl 4-(benzyloxy)-1-oxo-1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylamino)butan-2-ylcarbamate (1.57 g, 3.2 mmol) and following the experimental procedure described in Preparation 73b to afford 0.81 g (53% yield) of the title compound.

LRMS (m/z): 475 (M+1)⁺.

c) (S)-*tert*-Butyl 3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate

[0480] To a solution of (S)-*tert*-butyl 3-(benzyloxy)-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate (0.65 g, 1.37 mmol) in 33 mL of methanol was added Pd/C 10% (0.65 g). The reaction was stirred at room temperature under hydrogen (30 psi) for 15 hours. The reaction mixture was filtered through Celite® and the filtrate was evaporated to dryness to give 0.52 g (99%) of the title compound.

LRMS (m/z): 385 (M+1)⁺.

d) (S)-2-(1-Amino-3-hydroxypropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0481] This compound was prepared starting from (S)-*tert*-butyl 3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate (0.30 g, 0.8 mmol) and following the experimental procedure described in Preparation 81b to afford 0.24 g (85% yield) of the title compound, isolated as the hydrochloric salt form.

LRMS (m/z): 285 (M+1)⁺.

PREPARATION 129

(S)-2-(1-Amino-3-hydroxypropyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (S)-*tert*-Butyl 3-(benzyloxy)-1-(4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate**

[0482] This compound was obtained as subproduct in preparation 128b.

LRMS (m/z): 399 (M+1)⁺.

b) (S)-*tert*-Butyl 3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate

[0483] This compound was prepared starting from (S)-*tert*-butyl 3-(benzyloxy)-1-(4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate (200 mg, 0.5 mmol) and following the experimental procedure described in Preparation 128c to afford 152 mg (93% yield) of the title compound.

LRMS (m/z): 309 (M+1)⁺.

c) (S)-2-(1-Amino-3-hydroxypropyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0484] This compound was prepared starting from (S)-*tert*-butyl 3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylcarbamate (150 mg, 0.5 mmol) and following the experimental procedure described in Preparation 81b to afford 96 mg (73% yield, 91% purity) of the title compound, isolated as the hydrochloric salt form.

LRMS (m/z): 245 (M+1)⁺.

PREPARATION 130

(S)-2-(1-Aminoethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (S)-Benzyl 1-(3-bromo-2-(6-(trifluoromethyl)pyridin-2-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0485] The title compound was prepared from (S)-methyl 1-(2-(benzyloxycarbonylamino)propanamido)-3-bromo-1*H*-pyrrole-2-carboxylate (2.00 g, 4.7 mmol) and 6-(trifluoromethyl)pyridin-2-amine (3.00 g, 18.5 mmol) following the experimental procedure described in Preparation 44a. 1.32 g (51% yield) of the desired compound were obtained.

LRMS (m/z): 554, 556 (M+1)⁺.

b) (S)-Benzyl 1-(5-bromo-4-oxo-3-(6-(trifluoromethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0486] This compound was prepared starting from (S)-benzyl 1-(3-bromo-2-(6-(trifluoromethyl)pyridin-2-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (1.32 g, 2.4 mmol) and following the experimental procedure described in Preparation 68b to afford 0.71 g (55% yield) of the title compound.

LRMS (m/z): 536, 538 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0487] This compound was prepared starting from (S)-benzyl 1-(5-bromo-4-oxo-3-(6-(trifluoromethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (1.00 mg, 1.9 mmol) and following the experimental procedure described in Preparation 127c to afford 0.55 g (92% yield) of the title compound.

LRMS (m/z): 324 (M+1)⁺.

PREPARATION 131

(S)-2-(1-Aminoethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**a) (S)-Benzyl 1-(5-cyano-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate**

[0488] This compound was prepared starting from (S)-benzyl 1-(5-bromo-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (240 mg, 0.5 mmol) and following the experimental procedure described in Preparation 47a to afford 170 mg (71 % yield, 90% purity) of the title compound.

LRMS (m/z): 433 (M+1)⁺.

b) (S)-2-(1-Aminoethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0489] This compound was prepared starting from (S)-benzyl 1-(5-cyano-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (170 g, 0.4 mmol) and following the experimental procedure described in Preparation 128c but the mixture was hydrogenated at 14 psi to afford 114 mg (97% yield) of the title compound.

LRMS (m/z): 299 (M+1)⁺.

PREPARATION 132

2-(1-Aminocyclopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) *tert*-Butyl 1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylcarbamoyl)cyclopropylcarbamate**

[0490] The title compound was prepared from 1-amino-N-phenyl-1*H*-pyrrole-2-carboxamide (0.30 g, 1.5 mmol) and 1-(*tert*-butoxycarbonylamino)cyclopropanecarboxylic acid (0.30 g, 1.5 mmol) following the experimental procedure described in Preparation 73a. 0.55 g (91% yield) of the desired compound were obtained.

LRMS (m/z): 385 (M+1)⁺.

b) *tert*-Butyl 1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)cyclopropylcarbamate

[0491] This compound was prepared starting from *tert*-butyl 1-(2-(phenylcarbamoyl)-1*H*-pyrrol-1-ylcarbamoyl)cyclopropylcarbamate (0.54 g, 1.4 mmol) and following the experimental procedure described in Preparation 68b to afford 0.42 g (74% yield) of the title compound.
LRMS (m/z): 367 (M+1)⁺.

c) 2-(1-Aminocyclopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0492] This compound was prepared starting from *tert*-butyl 1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)cyclopropylcarbamate (0.41 g, 1.1 mmol) and following the experimental procedure described in Preparation 81b to afford 0.33 g (98% yield) of the title compound, isolated as the hydrochloric salt form.
LRMS (m/z): 267 (M+1)⁺.

PREPARATION 133**(*S*)-2-(1-Aminoethyl)-3-(tetrahydro-2*H*-pyran-4-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one****a) (*S*)-*tert*-Butyl 1-oxo-1-(2-(tetrahydro-2*H*-pyran-4-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate**

[0493] Tetrahydro-2*H*-pyran-4-amine (900 μl, 8.69 mmols) was added to a solution of (*S*)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (900 mg, 2.89 mmols, preparation 109) in toluene (36 ml). A 2M solution of Trimethyl aluminium in toluene (7 ml, 14.00 mmols) was added and it was stirred at 80°C overnight. Water (50 ml) and a 0.5M solution of sodium tartrate (25 ml) were added. It was extracted with ethylacetate. The combined organic layers were washed with water and brine. It was dried, filtered and concentrated in vacuum. The title compound was obtained (617 mg, 56%).
LRMS (m/z): 381 (M+1)⁺

b) (*S*)-*tert*-Butyl 1-(4-oxo-3-(tetrahydro-2*H*-pyran-4-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0494] Bromine (200 μl, 3.905 mmols) was added to a solution of triphenylphosphine (500 mg, 1.91 mmols) in dry dichloromethane (10 ml) and it was stirred under inert atmosphere at room temperature for 30 min. Triethylamine (1.14 ml, 8.179 mmols) was added and it was stirred under inert atmosphere at room temperature for 5 min. A solution of (*S*)-*tert*-butyl 1-oxo-1-(2-(tetrahydro-2*H*-pyran-4-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate (517 mg, 1.36 mmols) in dry dichloromethane (5 ml) was added to the previous solution and it was stirred under inert atmosphere at 60°C for 1 h.
[0495] A 4% aqueous solution of NaHCO₃ (100 ml) and dichloromethane (75 ml) were added to the reaction crude. The organic layer was passed through a phase separator cartridge and it was concentrated in vacuum. The residue was dissolved in 10 ml of THF/DMF 9:1 and sodium methanethiolate (172 mg, 2.45 mmols) was added. It was stirred under inert atmosphere at room temperature overnight. A 4% aqueous solution of NaHCO₃ and ethylacetate were added to the reaction crude. The organic phase was washed with water and brine. It was dried, filtered and concentrated in vacuum. The title compound was obtained (1180 mg, 60% approx purity, 100%) pure enough to be used in the next synthetic step without further purification.
LRMS (m/z): 363 (M+1)⁺

c) (*S*)-2-(1-Aminoethyl)-3-(tetrahydro-2*H*-pyran-4-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0496] HCl 4M dioxane (24 ml, 96 mmols) was added to (*S*)-*tert*-butyl 1-(4-oxo-3-(tetrahydro-2*H*-pyran-4-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (1180 mg, 60% purity, 1.95 mmols) and it was stirred at room temperature for 1 h. It was concentrated in vacuum. Water was added and it was washed with dichloromethane. A saturated aqueous solution of potassium carbonate was added to the aqueous phase and it was extracted with dichloromethane. The organic phase was dried, filtered and concentrated in vacuum. The title compound (287 mg, 85% purity, 48%) was obtained.

LRMS (m/z): 263 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.41 (d, 3 H) 1.55 - 1.82 (m, 2 H) 2.13 (s, 2 H) 2.64 - 2.89 (m, 2 H) 3.34 - 3.45 (m, 2 H) 3.80 - 4.01 (m, 2 H) 4.10 - 4.27 (m, 1 H) 4.58 - 4.75 (m, 1 H) 6.52 (dd, 1 H) 6.82 (dd, 1 H) 7.41 - 7.55 (m, 1 H)

PREPARATION 134

(S)-2-(1-Aminoethyl)-3-(2,2,2-trifluoroethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (S)-*tert*-Butyl 1-oxo-1-(2-(2,2,2-trifluoroethylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate**

[0497] 2,2,2-Trifluoroethanamine (680 µmol, 8.66 mmols) was added to a solution of (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (900 mg, 2.89 mmols) in toluene (36 ml). A 2M solution of trimethyl aluminium in toluene (7 ml, 14.00 mmols) was added and it was stirred at 80°C overnight. Water (50 ml) and a 0.5M solution of sodium tartrate (25 ml) were added. It was extracted with ethylacetate. The combined organic layers were washed with water and brine. It was dried, filtered and concentrated in vacuum. The title compound was obtained (1.021 g, 93%).

LRMS (m/z): 379 (M+1)+

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.20 - 1.30 (d, 3 H) 1.34 - 1.43 (s, 9 H) 3.80 - 4.03 (m, 2 H) 4.03 - 4.17 (m, 1 H) 6.00 - 6.20 (m, 1 H) 6.80 - 6.93 (m, 2 H) 6.96 - 7.17 (m, 1 H) 8.39 - 8.56 (m, 1 H) 11.09 - 11.27 (s, 1 H)

b) (S)-*tert*-Butyl 1-(4-oxo-3-(2,2,2-trifluoroethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0498] Bromine (350 µl, 6.83 mmols) was added to a solution of triphenylphosphine (895 mg, 3.41 mmols) in dry dichloromethane (18 ml) and it was stirred under inert atmosphere at room temperature for 30 min. Triethylamine (2.00 ml, 14.35 mmols) was added and it was stirred under inert atmosphere at room temperature for 5 min. A solution of (S)-*tert*-butyl 1-oxo-1-(2-(2,2,2-trifluoroethylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate (921 mg, 2.43 mmols) in dry dichloromethane (9 ml) was added to the previous solution and it was stirred under inert atmosphere at 60°C for 1 h. A 4% aqueous solution of NaHCO₃ (180 ml) and dichloromethane (135 ml) were added to the reaction crude. The organic layer was passed through a phase separator cartridge and it was concentrated in vacuum. The residue was dissolved in 18 ml of THF/DMF 9:1 and sodium methanethiolate (307 mg, 4.38 mmols) was added. It was stirred under inert atmosphere at room temperature overnight. A 4% aqueous solution of NaHCO₃ and ethylacetate were added to the reaction crude. The organic phase was washed with water and brine. It was dried, filtered and concentrated in vacuum. The title compound was obtained (2300 mg, 50% approx purity, 100%) pure enough to be used in the next synthetic step without further purification.

LRMS (m/z): 361 (M+1)+

c) (S)-2-(1-Aminoethyl)-3-(2,2,2-trifluoroethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0499] HCl 4M dioxane (37 ml, 148 mmols) was added to (S)-*tert*-butyl 1-(4-oxo-3-(2,2,2-trifluoroethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (2.30 g, 50% purity, 3.19 mmols) and it was stirred at room temperature for 1 h. It was concentrated in vacuum. Water was added and it was washed with dichloromethane. A saturated aqueous solution of potassium carbonate (50 ml) was added to the aqueous phase and it was extracted with dichloromethane. The organic phase was dried, filtered and concentrated in vacuum. The title compound (526 mg, 58% yield) was obtained.

LRMS (m/z): 261 (M+1)+

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.42 (d, 3 H) 2.20 (s, 2 H) 3.86 - 4.16 (q, 1 H) 4.88 - 5.40 (m, 2 H) 6.60 (dd, 1 H) 6.99 (dd, 1 H) 7.47 - 7.81 (m, 1 H)

PREPARATION 135

(S)-2-(1-Aminoethyl)-3-cyclobutylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**a) (S)-*tert*-Butyl 1-(2-(cyclobutylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0500] Same procedure as described in preparation 133a was used from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (900 mg, 2.66 mmols) and cyclobutanamine (0.68 ml, 7.98 mmols). After reverse phase chromatography the title compound was obtained (229 mg, 25%).

LRMS (m/z): 351 (M+1)+

b) (S)-*tert*-Butyl 1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0501] Same procedure as described in preparation 133b was used from (S)-*tert*-butyl 1-(2-(cyclobutylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (204 mg, 0.58 mmols). The title compound was obtained (424 mg, 25%)

purity, 50%) pure enough to be used in the next synthetic step without further purification.
LRMS (m/z): 333 (M+1)+

c) (S)-2-(1-Aminoethyl)-3-cyclobutylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0502] Same procedure as described in preparation 133c was used from (S)-*tert*-butyl 1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (424 mg, 25% purity, 0.32 mmols). The title compound was obtained (107 mg, 83% purity, 100%)
LRMS (m/z): 233 (M+1)+

PREPARATION 136

(S)-2-(1-Aminoethyl)-3-cyclopropylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) (S)-*tert*-Butyl 1-(2-(cyclopropylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0503] Same procedure as described in preparation 133a was used from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (900 mg, 80% purity, 2.28 mmols) and cyclopropanamine (0.48 ml, 6.86 mmols). After 3.5h stirring at 80°C the title compound was obtained (589 mg, 77%).
LRMS (m/z): 337 (M+1)+

b) (S)-*tert*-Butyl 1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0504] Same procedure as described in preparation 133b was used from (S)-*tert*-butyl 1-(2-(cyclopropylcarbamoyl)-1*H*-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (589 mg, 1.75 mmols). The title compound was obtained (1.18 g, 100%)
LRMS (m/z): 319 (M+1)+

c) (S)-2-(1-Aminoethyl)-3-cyclopropylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0505] Same procedure as described in preparation 133c was used from (S)-*tert*-butyl 1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (1.18g, 3.37mmols). The title compound was obtained (311 mg, 69% purity, 29%)
LRMS (m/z): 219 (M+1)+

PREPARATION 137

2-((S)-1-Aminoethyl)-3-(tetrahydro-2*H*-pyran-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

a) *tert*-Butyl (2*S*)-1-oxo-1-(2-(tetrahydro-2*H*-pyran-3-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate

[0506] Anhydrous triethylamine (600 μ l, 4.31 mmols) was added to a solution of tetrahydro-2*H*-pyran-3-amine.HCl (600 mg, 4.36 mmols) in toluene (10 ml) and it was stirred at room temperature for 30min. A solution of (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1*H*-pyrrole-2-carboxylate (900 mg, 2.90 mmols) in toluene (36 ml) and a 2M solution of trimethylaluminium in toluene (7 ml, 14 mmols) were added. The solution obtained was stirred at 80°C for 2h.
[0507] Water (70 ml) and a 0.5M sodium tartrate solution (35ml) were added. It was extracted with ethylacetate. The combined organic layers were washed with water and brine. It was dried, filtered and concentrated in vacuum. The title compound was obtained (750 mg, 68%).
LRMS (m/z): 381 (M+1)+

b) *tert*-Butyl (1*S*)-1-(4-oxo-3-(tetrahydro-2*H*-pyran-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate

[0508] Bromine (283 μ l, 5.52 mmols) was added to a solution of triphenylphosphine (727 mg, 2.77 mmols) in dry dichloromethane (15 ml) and it was stirred under inert atmosphere at room temperature for 30min. Triethylamine (1.65 ml, 11.83 mmols) was added and it was stirred under inert atmosphere at room temperature for 5 min. A solution of *tert*-butyl (2*S*)-1-oxo-1-(2-(tetrahydro-2*H*-pyran-3-ylcarbamoyl)-1*H*-pyrrol-1-ylamino)propan-2-ylcarbamate (750 mg, 1.97 mmols) in dry dichloromethane (7 ml) was added to the previous solution and it was stirred under inert atmosphere at

60°C for 1 h. A 4% aqueous solution of NaHCO₃ (145 ml) and dichloromethane (110 ml) were added to the reaction crude. The organic layer was passed through a phase separator cartridge and it was concentrated in vacuum. The residue was dissolved in 15ml of THF/DMF 9:1 and sodium methanethiolate (250 mg, 3.57 mmols) was added. It was stirred under inert atmosphere at room temperature overnight. A 4% aqueous solution of NaHCO₃ and ethylacetate were added to the reaction crude. The organic phase was washed with water and brine. It was dried, filtered and concentrated in vacuum. The title compound was obtained (1163 mg, 60% approx purity, 100%) pure enough to be used in the next synthetic step without further purification.

LRMS (m/z): 363 (M+1)⁺

c) 2-((S)-1-Aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0509] HCl 4M dioxane (24ml, 96mmols) was added to *tert*-butyl (1S)-1-(4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (1163 mg, 1.93 mmols, 60% purity) and it was stirred at room temperature for 1 h. It was concentrated in vacuum. Water was added and it was washed with dichloromethane. A saturated aqueous solution of potassium carbonate (50ml) was added to the aqueous phase and it was extracted with dichloromethane. The organic phase was dried, filtered and concentrated in vacuum. The title compound (278mg, 55%) was obtained.

LRMS (m/z): 263 (M+1)⁺

PREPARATION 138

(S)-2-(1-Aminoethyl)-3-(isoxazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

a) (S)-*tert*-Butyl 1-(2-(isoxazol-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate

[0510] Same procedure as described in preparation 133a was used from (S)-methyl 1-(2-(*tert*-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (854 mg, 2.17 mmols) and isoxazol-3-amine (0.48ml, 6.50 mmols). The title compound was obtained (919mg, 55% purity, 64%).

LRMS (m/z): 364 (M+1)⁺

b) (S)-*tert*-Butyl 1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0511] Same procedure as described in preparation 133b was used from (S)-*tert*-butyl 1-(2-(isoxazol-3-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (919 mg, 55% purity, 1.39 mmols). The title compound was obtained (1.34g, 17% purity, 47%).

LRMS (m/z): 346 (M+1)⁺

c) (S)-2-(1-Aminoethyl)-3-(isoxazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0512] Same procedure as described in preparation 133c was used from (S)-*tert*-butyl 1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (1.34 g, 17% purity, 0.66 mmols). The title compound was obtained (188 mg, 100%) LRMS (m/z): 246 (M+1)⁺

PREPARATION 139

(S)-*tert*-Butyl 3-(benzyloxy)-1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate

a) (S)-Methyl 1-(4-(benzyloxy)-2-(*tert*-butoxycarbonylamino)butanamido)-3-bromo-1H-pyrrole-2-carboxylate

[0513] The title compound was prepared from methyl 1-amino-3-bromo-1H-pyrrole-2-carboxylate (2.83 g, 12.9 mmol) and (S)-4-(benzyloxy)-2-(*tert*-butoxycarbonylamino)butanoic acid (4.0 g, 12.9 mmol) following the experimental procedure described in Preparation 73a. 6.12 g (93% yield) of the desired compound were obtained.

LRMS (m/z): 510, 512 (M+1)⁺.

b) (S)-tert-Butyl 4-(benzyloxy)-1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxobutan-2-ylcarbamate

[0514] The title compound was prepared from (S)-methyl 1-(4-(benzyloxy)-2-(tert-butoxycarbonylamino)butanamido)-3-bromo-1H-pyrrole-2-carboxylate (4.0 g, 7.8 mmol) and 3,5-difluoroaniline (5.1 g, 39.0 mmol) following the experimental procedure described in Preparation 44a. 3.2 g (66% yield) of the desired compound were obtained.
LRMS (m/z): 607, 609 (M+1)⁺.

c) (S)-tert-Butyl 3-(benzyloxy)-1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate

[0515] This compound was prepared starting from (S)-tert-butyl 4-(benzyloxy)-1-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxobutan-2-ylcarbamate (2.14 g, 3.52 mmol) and following the experimental procedure described in Preparation 68b to afford 1.66 g (80% yield) of the title compound.
LRMS (m/z): 589, 591 (M+1)⁺.

PREPARATION 140**(S)-2-(1-Amino-3-hydroxypropyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one****a) (S)-tert-Butyl 3-(benzyloxy)-1-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate**

[0516] This compound was prepared starting from (S)-tert-butyl 3-(benzyloxy)-1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate (0.81 g, 1.37 mmol) and following the experimental procedure described in Preparation 26a to afford 0.80 g (54% yield, 59 % purity) of the title compound.
LRMS (m/z): 637 (M+1)⁺.

b) (S)-tert-Butyl 3-(benzyloxy)-1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate

[0517] This compound was prepared starting from (S)-tert-butyl 3-(benzyloxy)-1-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate (0.40 g, 0.63 mmol) and following the experimental procedure described in Preparation 26b to afford 0.32 g (87% yield) of the title compound.
LRMS (m/z): 579 (M+1)⁺.

c) (S)-tert-Butyl 1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-3-hydroxypropylcarbamate

[0518] This compound was prepared starting from (S)-tert-butyl 3-(benzyloxy)-1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate (0.32 g, 0.55 mmol) and following the experimental procedure described in Preparation 128c to afford 0.26 g (98% yield) of the title compound.
LRMS (m/z): 489 (M+1)⁺.

d) (S)-2-(1-Amino-3-hydroxypropyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0519] This compound was prepared starting from (S)-tert-butyl 1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-3-hydroxypropylcarbamate (0.25 g, 0.51 mmol) and following the experimental procedure described in Preparation 81 b to afford 0.21 g (95% yield) of the title compound, isolated as the hydrochloric salt form.
LRMS (m/z): 389 (M+1)⁺.

PREPARATION 141

a) (S)-tert-Butyl 3-(benzyloxy)-1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate

[0520] This compound was prepared starting from (S)-tert-butyl 3-(benzyloxy)-1-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate (0.81 g, 1.37 mmol) and following the experimental procedure described in Preparation 47a to afford 0.57 g (77% yield) of the title compound.
LRMS (m/z): 536 (M+1)⁺.

b) (S)-2-(1-Amino-3-hydroxypropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile

[0521] Under a nitrogen atmosphere, to a solution of (S)-tert-butyl 3-(benzyloxy)-1-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylcarbamate (0.25 g, 0.47 mmol) in 1.9 mL of dichloromethane, boron tribromide 1.0 M methylene chloride solution (1.41 mL, 1.41 mmol) was added at -78°C, and the mixture was stirred at the same temperature for 1 hour. After saturated aqueous NaHCO₃ was added and the mixture was extracted three times with ethyl acetate. The combined organic extracts were washed with water and brine, dried over magnesium sulphate, filtered and the solvent evaporated in vacuum to afford 0.17 g (77% yield, 73 % purity) of the title compound that was used in the next step without any further purification.
LRMS (m/z): 346 (M+1)⁺.

PREPARATION 142

(S)-4-Amino-6-(1-(3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**a) (S)-tert-Butyl 1-(2-(1-(4-methoxybenzyl)-1H-pyrazol-4-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate**

[0522] This compound was prepared starting from (S)-methyl 1-(2-(tert-butoxycarbonylamino)propanamido)-1H-pyrrole-2-carboxylate (900 mg, 2.89 mmol) and 1-(4-methoxybenzyl)-1H-pyrazol-4-amine (1.24 g, 6.10 mmol) following the experimental procedure described in Preparation 27a to afford 1.02 g (100% purity, 73% yield) of the title compound after purification by flash chromatography (0% to 100% AcOEt/hexanes).
LRMS (m/z): 483 (M+1)⁺.

b) (S)-tert-Butyl 1-(3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate

[0523] Bromine (151 µL, 2.95 mmol) was added dropwise to a solution of triphenylphosphine (780 mg, 2.97 mmol) in dichloromethane (8 mL) under nitrogen. The solution was stirred for 30 min, and triethylamine (1.18 mL, 8.47 mmol) and a solution of (S)-tert-butyl 1-(2-(1-(4-methoxybenzyl)-1H-pyrazol-4-ylcarbamoyl)-1H-pyrrol-1-ylamino)-1-oxopropan-2-ylcarbamate (1.02 g, 2.11 mmol) in 16 mL of dichloromethane was added. The reaction mixture was stirred at 60 °C for 2 h, and then poured onto 4% NaHCO₃. After extraction with dichloromethane, the organic phase was dried over magnesium sulphate and the volatiles were removed under reduced pressure. The residue was redissolved in a mixture of 40 mL of tetrahydrofuran and 4 mL of dimethylformamide, sodium thiomethoxide (0.44 g, 6.28 mmol) was added and the mixture was stirred at room temperature for 2h. After pouring onto 4% NaHCO₃, it was extracted with dichloromethane, the organic phase was dried over magnesium sulphate and the volatiles were removed under reduced pressure. The residue was purified by flash chromatography (0% to 70% AcOEt/hexanes) to yield 0.86 g (88% yield) of the title compound.
LRMS (m/z): 465 (M+1)⁺.

c) (S)-2-(1-Aminoethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0524] The title compound was prepared from (S)-tert-butyl 1-(3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylcarbamate (860 mg, 1.85 mmol) following the experimental procedure described in Preparation 46c. 540 mg (75% yield) of the desired compound were obtained.
LRMS (m/z): 365 (M+1)⁺.

d) (S)-4-Amino-6-(1-(3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile


[0525] The title compound (290 mg, 39% yield) was obtained from (S)-2-(1-Aminoethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 483 (M+1)⁺.

PREPARATION 143

(2S,4R)-4-(Benzyloxy)-1-(tert-butoxycarbonyl)pyrrolidine-2-carboxylic acid

[0526]



a) (2S,4R)-1-tert-Butyl 2-methyl 4-(benzyloxy)pyrrolidine-1,2-dicarboxylate

[0527] (2S,4R)-1-tert-Butyl 2-methyl 4-hydroxypyrrolidine-1,2-dicarboxylate (2.65 g, 10.80 mmol) was dissolved in dimethylformamide (75 ml) and cooled in an ice bath. Sodium hydride (60% in hexanes, 0.57 g, 23.75 mmol) was added and stirred for 10 min. To this solution, benzyl bromide (1.39 ml, 11.69 mmol) dissolved in dichloromethane (9 ml) was dropwise added and the reaction mixture was overnight stirred at room temperature. The solvent was evaporated under reduced pressure and the residue redissolved in ethyl acetate. This organic phase was washed with water, dried over magnesium sulphate, filtered and the solvent evaporated under reduced pressure. 3.93g (77% yield) of the final product were obtained, pure enough to perform the next synthetic step.

LRMS (m/z): 336 (M+1)⁺

b) (2S,4R)-4-(Benzyloxy)-1-(tert-butoxycarbonyl)pyrrolidine-2-carboxylic acid

[0528] (2S,4R)-1-tert-Butyl 2-methyl 4-(benzyloxy)pyrrolidine-1,2-dicarboxylate (2.78 g, 8.29 mmol) was dissolved in a 1:1 mixture of MeOH and THF (30 ml) and 2N NaOH (12.5 ml, 26 mmol) was added. After stirring at room temperature for 2h, the organic solvents were evaporated under reduced pressure and the aqueous phase was extracted twice with dichloromethane. The aqueous phase was then cooled at 0 °C and then acidified with concentrated chlorhydric acid. This phase was extracted twice with dichloromethane. The organic phase was dried over magnesium sulphate, filtered and the solvent evaporated under reduced pressure. 2.01 g (76% yield) of the final compound were obtained, pure enough to perform the next synthetic step.

LRMS (m/z): 322 (M+1)⁺

PREPARATION 144

(2S,4R)-tert-Butyl 4-(benzyloxy)-2-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

a) (2S,4R)-tert-Butyl 4-(benzyloxy)-2-(3-bromo-2-(3,5-difluorophenylcarbamoyl)-1H-pyrrol-1-ylcarbamoyl)pyrrolidine-1-carboxylate

[0529] 1-Amino-3-bromo-N-(3,5-difluorophenyl)-1H-pyrrole-2-carboxamide (1.5 g, 4.75 mmol), (2S,4R)-4-(benzyloxy)-1-(tert-butoxycarbonyl)pyrrolidine-2-carboxylic acid (0.99 g, 3.08 mmol) and diisopropylethylamine (2.8 ml, 16.08 mmol) were dissolved in dimethylformamide (30 ml) and the mixture cooled at 0 °C. To this mixture, T3P (50% solution in ethyl acetate, 2.1 ml, 7.19 mmol) dissolved in DMF (5 ml) was dropwise added. Once the addition is over, the mixture was stirred at room temperature for 48h. To this mixture, water was added and extracted twice with ethyl acetate. The

organic phase was washed with water and brine, dried over sodium sulphate, filtered and the solvents evaporated under reduced pressure. The residue was purified by flash chromatography (0% to 25% hexanes/AcOEt) to yield 1.42 g (74% yield) of the title compound.

LRMS (m/z): 620 (M+1)⁺.

b) (2*S*,4*R*)-*tert*-Butyl 4-(benzyloxy)-2-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

[0530] The title compound was prepared from (2*S*,4*R*)-*tert*-butyl 4-(benzyloxy)-2-(3-bromo-2-(3,5-difluorophenyl)carbamoyl)-1*H*-pyrrol-1-ylcarbamoylpyrrolidine-1-carboxylate (1.42 g, 2.29 mmol) following the experimental procedure described in Preparation 142b. 648 mg (47% yield) of the desired compound were obtained.

LRMS (m/z): 602 (M+1)⁺.

PREPARATION 145

(2*S*,4*R*)-*tert*-Butyl 4-(benzyloxy)-2-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

a) (2*S*,4*R*)-*tert*-Butyl 4-(benzyloxy)-2-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

[0531] This compound was prepared starting from (2*S*,4*R*)-*tert*-butyl 4-(benzyloxy)-2-(5-bromo-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate (648 mg, 1.08 mmol) and following the experimental procedure described in Preparation 26a to afford 326 mg (47% yield) of the title compound.

LRMS (m/z): 649 (M+1)⁺.

b) (2*S*,4*R*)-*tert*-Butyl 4-(benzyloxy)-2-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate

[0532] (2*S*,4*R*)-*tert*-Butyl 4-(benzyloxy)-2-(3-(3,5-difluorophenyl)-5-iodo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate (125 mg, 0.19 mmol) was dissolved in pyridine (10 ml) and copper (I) cyanide (210 mg, 2.32 mmol) was added. The reaction vessel was closed, purged with nitrogen and heated at 120 °C for 5h in a microwave apparatus. The reaction mixture was filtered through Celite®, and the solvents evaporated under reduced pressure. The residue was redissolved in water and extracted with ethyl acetate. The organic phase was washed with water and brine, dried over magnesium sulphate, filtered and the solvent evaporated under reduced pressure. 105 mg (100% yield) of the final compound were obtained, pure enough to perform the next synthetic step.

LRMS (m/z): 548 (M+1)⁺

PREPARATION 146

2-((2*S*,4*R*)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-(benzyloxy)pyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

.a) 2-((2*S*,4*R*)-4-(Benzyloxy)pyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0533] This compound was prepared starting from (2*S*,4*R*)-*tert*-butyl 4-(benzyloxy)-2-(5-cyano-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidine-1-carboxylate (105 mg, 0.19 mmol) and following the experimental procedure described in Preparation 46c to afford 37 mg (43% yield) of the title compound.

LRMS (m/z): 448 (M+1)⁺.

b) 2-((2*S*,4*R*)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-(benzyloxy)pyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0534] 2-((2*S*,4*R*)-4-(Benzyloxy)pyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (37 mg, 0.08 mmol), 23 mg (0.12 mmol) of 4-amino-6-chloropyrimidine-5-carbonitrile (prepared according to the procedure described in WO2010151735A2) and 60 µL (0.34 mmol) of diisopropylamine in 5 mL of *tert*-butanol was heated with stirring at 100 °C overnight. Then the solvent was removed under vacuum and the product was purified

by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1 % v/v formic acid buffered] 0% to 100%) to obtain the title compound (20 mg, 43% yield) as a white solid.

LRMS (m/z): 566 (M+1)⁺.

EXAMPLE 1

2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0535] A mixture of 5-chloro-2-(chloromethyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (90 mg, 0.25 mmol) and adenine (43 mg, 0.32 mmol) was suspended in *N,N*-dimethylformamide (2 mL) and potassium carbonate (44 mg, 0.32 mmol) was added stirring the reaction at room temperature overnight. At the end of this period, dichloromethane was added and the insolubles were filtered out. The filtrate was concentrated to dryness and macerated with dimethylsulphoxide affording 57 mg (51% yield) of a solid corresponding to the title compound.

LRMS (m/z): 407 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.06 (s, 1H), 7.92 (s, 1H), 7.61 (d, J = 3.0 Hz, 1 H), 7.54 - 7.33 (m, 4H), 7.25 (s, 2H), 6.67 (d, J = 3.0 Hz, 1H), 5.03 (d, J = 17.1 Hz, 1 H), 4.79 (d, J = 17.1 Hz, 1H), 2.11 (s, 3H).

EXAMPLE 2

2-((6-Aminopyrimidin-4-ylamino)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0536] In a microwave tub, a mixture of 2-(aminomethyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (85 mg, 0.29 mmol), 6-bromopyrimidin-4-amine (102 mg, 0.59 mmol), and DIEA (205 μL, 1.2 mmol) was dissolved in *tert*-butanol (3 mL) and was heated at 140°C with stirring during 20 hours. Next day, ethyl acetate was added and the organic phase was washed with water and brine, dried (Na₂SO₄) and concentrated to give 180 mg of a residue that was purified using a Bond Elut 5g silica cartridge eluting with dichloromethane/methanol mixtures obtain 5 mg of the title compound (4% yield).

LRMS (m/z): 382 (M+1)⁺.

EXAMPLE 3

2-((6-Amino-9H-purin-9-yl)methyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0537] This compound was prepared starting from 2-(chloromethyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (107 mg, 0.29 mmol) and following the experimental procedure described in Example 1. Isolation of the compound was done by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to afford 12 mg (10% yield) of the title compound.

LRMS (m/z): 413 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.05 (s, 1 H), 7.47-7.38 (m, 5H), 7.22 (s, 2H), 6.14 (d, J = 2.8 Hz, 1H), 5.01 (d, J = 16.9 Hz, 1H), 4.78 (d, J = 16.9 Hz, 1H), 2.50 (m, 1 H), 2.10 (s, 3H), 0.94 (d, J = 8.6 Hz, 2H), 0.70 - 0.56 (m, 2H).

EXAMPLE 4

2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0538] To a solution of 2-(chloromethyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (90 mg, 0.23 mmol) in dry *N,N*-dimethylformamide (4 mL), 9H-purin-6-amine (38 mg, 0.28 mmol) and potassium carbonate (38 mg, 0.27 mmol) were added. It was stirred at room temperature overnight. It was filtered through Celite® and it was concentrated in vacuum. The residue obtained was purified by flash chromatography silica (dichloromethane/methanol). The expected product was obtained (25 mg, 29% yield).

LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.06 (s, 1 H), 7.92 (s, 1 H), 7.53 - 7.62 (m, 1 H), 7.50 (d, J=7.42 Hz, 1 H), 7.34 - 7.48 (m, 3 H), 7.25 (s, 2 H), 7.00 (dd, J=4.30, 1.56 Hz, 1 H), 6.58 (dd, J=4.30, 2.74 Hz, 1 H), 5.07 (d, J=16.80 Hz, 1 H), 4.82 (d, J=16.80 Hz, 1 H), 2.09 (s, 3 H).

EXAMPLE 5**2-((6-Aminopyrimidin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0539] 100 mg (0.39 mmol) of 2-(aminomethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 76 mg (0.44 mmol) of 6-bromopyrimidin-4-amine and 140 μ L (0.80 mmol) of DIEA were suspended in 2 mL of *tert*-butanol and the resulting mixture was stirred at 80 °C overnight. After an extra addition of 76 mg (0.44 mmol) of 6-bromopyrimidin-4-amine and 140 μ L (0.80 mmol) of DIEA the reaction was heated at 80 °C for 70 hours. Then the solvents were evaporated under vacuum and the crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) and then by preparative HPLC (Symmetry Prep C₁₈ column, mixture of eluents A/B from 20% B to 20% B, in a 10 min. gradient) to give 18 mg (13% yield) of the title compound.
LRMS (m/z): 348 (M+1)⁺.

EXAMPLE 6**4-((4-Oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)methylamino)picolinamide**

[0540] A mixture of 2-(aminomethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (102 mg, 0.4 mmol), 4-bromopicolinamide (105 mg, 0.52 mmol) and DIEA (200 μ L, 1.13 mmol) in *n*-butanol (2.2 mL) was reacted under microwave irradiation at 190 °C during 22 h. After cooled to room temperature, the mixture was concentrated in vacuum and was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain 7 mg of the title compound (4,6%).

[0541] LRMS (m/z): 375 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.38 (s, 2 H), 8.01 (d, J=5.86 Hz, 1 H), 7.90 (m, 1 H), 7.61 - 7.73 (m, 1 H), 7.48 - 7.58 (m, 1 H), 7.39 - 7.48 (m, 1 H), 7.23 - 7.40 (m, 2 H), 7.03 - 7.14 (m, 1 H), 6.93 - 7.04 (m, 1 H), 6.62 (dd, J=4.30, 2.74 Hz, 1 H), 6.38 - 6.53 (m, 1 H), 3.86 - 3.99 (m, 2 H), 2.08 (s, 3 H)

EXAMPLE 7**2-((2-Aminopyridin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0542] 150 mg (0.59 mmol) of 2-(aminomethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 104 mg (0.60 mmol) of 4-bromopyridin-2-amine and 105 μ L (0.60 mmol) DIEA were dissolved in 2 mL of *tert*-butanol and stirred at 180 °C under microwave irradiation for 5.5 hours. Then the solvent was evaporated in vacuum and the crude product was purified by flash chromatography (dichloromethane to dichloromethane/MeOH/NH₄OH, 100:8:1) to give 57 mg (28% yield) of the title compound.

LRMS (m/z): 347 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.69 (s, 1 H), 7.55 - 7.29 (m, 5H), 6.99 (s, 1 H), 6.62 (s, 1H), 6.37 (s, 1H), 5.76 (s, 1H), 5.54 (s, 2H), 5.40 (s, 1H), 3.85 - 3.66 (m, 2H), 2.08 (s, 3H).

EXAMPLE 8**2-((9*H*-Purin-6-ylamino)methyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0543] 100 mg (0.39 mmol) of 2-(aminomethyl)-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 86 mg (0.43 mmol) of 6-bromo-9*H*-purine and 151 μ L (0.87 mmol) DIEA were dissolved in 5 mL of *tert*-butanol and stirred at 80 °C overnight. Then the solvent was evaporated in vacuum and the residue was dissolved in ethyl acetate and washed with a saturated aqueous solution of sodium bicarbonate and brine. It was dried over magnesium sulphate, filtered and the solvent was evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to give 42 mg (29% yield) of the title compound.

LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.95 (s, 1H), 8.14 (s, 1H), 8.08 (s, 1H), 7.87 (s, 1H), 7.62 (s, 1H), 7.46 (d, J = 7.1 Hz, 1H), 7.42 - 7.22 (m, 3H), 6.97 (dd, J = 4.3, 1.6 Hz, 1 H), 6.57 (dd, J = 4.3, 2.7 Hz, 1 H), 4.20 (br s, 2H), 2.19 (s, 3H).

EXAMPLE 9**2-((6-Amino-9H-purin-9-yl)methyl)-3-cyclohexylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0544] 2-(Chloromethyl)-3-cyclohexylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (140 mg, 0.53 mmol), 9H-purin-6-amine (91 mg, 0.67 mmol) and potassium carbonate (93 mg, 0.67 mmol) were suspended in *N,N*-dimethylformamide under argon atmosphere and the reaction was stirred overnight at room temperature. Next day, dichloromethane was added and the resulting solid was filtered off. The filtrate was concentrated to dryness giving a residue of 176 mg that was purified by flash chromatography silica (dichloromethane/methanol) to obtain 7 mg of the title compound (3,4% yield).

LRMS (*m/z*): 365 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.19 (s, 1H), 8.21 (s, 1H), 7.46 (br s, 1H), 7.32 (brs, 2H), 6.90 - 6.77 (m, 1 H), 6.57 - 6.47 (m, 1 H), 5.58 (s, 2H), 3.86 (s, 1 H), 1.63 - 0.96 (m, 10H).

EXAMPLE 10**2-((6-Amino-9H-purin-9-yl)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0545] 50 mg (0.17 mmol) of 2-(chloromethyl)-5-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 25.8 mg (0.19 mmol) of 9H-purin-6-amine and 26.4 mg of potassium carbonate were dissolved in 2.5 mL of DMF and stirred at room temperature for 3 hours. Then the reaction mixture was diluted with dichloromethane, filtered and evaporated to dryness. The oil that resulted was purified by flash chromatography (DCM to 5% MeOH/DCM) to give 52 mg (77% yield) of the title compound.

LRMS (*m/z*): 387 (*M*+1)⁺.

EXAMPLE 11**2-((9H-Purin-6-ylthio)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0546] To a solution of 2-(chloromethyl)-5-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (108 mg, 0.38 mmol) in 7.5 mL of *N,N*-dimethylformamide was added 9H-purine-6-thiol (57 mg, 0.37 mmol) and potassium carbonate (52 mg, 0.38 mmol) and the stirring was continued overnight at room temperature. Next day, the reaction was concentrated to dryness and the residue was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain 35 mg of the title compound (23%).

LRMS (*m/z*): 404 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.46 (s, 1H), 8.39 (s, 1H), 7.53 (d, *J* = 2.7 Hz, 1H), 7.44 (m, 1H), 7.24 (m, 3H), 6.42 (d, *J* = 2.7 Hz, 1H), 4.38 (d, *J* = 15.2 Hz, 1H), 4.26 (d, *J* = 15.2 Hz, 1H), 2.39 (s, 3H), 2.15 (s, 3H).

EXAMPLE 12**2-((6-Amino-9H-purin-9-yl)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0547] 289 mg (1.00 mmol) of 2-(chloromethyl)-6-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 163 mg (1.21 mmol) of 9H-purin-6-amine and 278 mg (2.01 mmol) of potassium carbonate were dissolved in 8 mL of DMF and stirred at room temperature overnight. The solvent was removed in vacuum and the residue was taken up in ethyl acetate, washed with brine, filtered and evaporated to dryness. The product was purified by preparative HPLC (Symmetry Prep C₁₈ column, mixture of eluents A/B from 40% B to 52% B, in a 12 min. gradient) to give 116 mg (29% yield) of the title compound.

LRMS (*m/z*): 387 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.04 (s, 1H), 7.88 (s, 1H), 7.52 - 7.29 (m, 5H), 7.24 (s, 2H), 6.80 (s, 1H), 5.04 (d, *J* = 17.0 Hz, 1H), 4.80 (d, *J* = 16.9 Hz, 1H), 2.13 (s, 3H), 2.04 (s, 3H).

EXAMPLE 13**2-((9H-Purin-6-ylthio)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0548] 289 mg (1.00 mmol) of 2-(chloromethyl)-6-methyl-3-o-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 183 mg (1.21 mmol) of 7H-purine-6-thiol and 277 mg (2.01 mmol) of potassium carbonate were dissolved in 8 mL of DMF and stirred at room temperature overnight. The solvent was removed in vacuum and the product was purified by preparative HPLC

(Symmetry Prep C₁₈ column, mixture of eluents A/B from 50% B to 63% B, in a 13 min. gradient) to give 133 mg (50% yield) of the title compound.

LRMS (m/z): 404 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.47 (s, 1H), 8.43 (s, 1H), 7.51 - 7.16 (m, 5H), 6.79 (s, 1H), 4.41 (d, J = 15.2 Hz, 1H), 4.29 (d, J = 15.2 Hz, 1H), 2.16 (s, 3H), 2.11 (s, 3H).

EXAMPLE 14

2-(1-(6-Amino-9H-purin-9-yl)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0549] To a solution of 100 mg (0.37 mmol) of 2-(1-chloroethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one in 5 mL of DMF, 55 mg (0.41 mmol) of 9H-purin-6-amine and 55 mg (0.40 mmol) of potassium carbonate were added. It was stirred at 60 °C overnight. It was then filtered through Celite® and it was concentrated in vacuum. The residue that was obtained was purified by flash chromatography (5% MeOH in dichloromethane). 20 mg (15% yield) of the title product were obtained.

LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ ppm 8.27 - 7.80 (m, 2H), 7.82 - 6.88 (d, J = 7.03 Hz, 1 H), 6.64 (m, 1 H), 5.49 (q, J = 6.25 Hz, 1 H), 1.71 (d, J = 5.86 Hz, 3H).

EXAMPLE 15

(S)-2-(1-(9H-Purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0550] 90 mg (0.34 mmol) of (S)-2-(1-aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 134 mg (0.67 mmol) of 6-bromo-9H-purine and 174 mg (1.35 mmol) of DIEA were suspended in 3 mL of *tert*-butanol and the mixture was heated to 100 °C for 40 hours. Then the solvent was removed in vacuum and the residue was taken up in AcOEt, washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound (59 mg, 45% yield) as a white solid.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.92 (s, 1H), 8.10 (m, 2H), 7.94 (s, 1H), 7.60 (s, 1 H), 7.53-7.34 (m, 4H), 6.93 (dd, 1 H), 6.64 - 6.54 (m, 1 H), 4.65 (s, 1 H), 1.98 (m, 2H), 0.77 (t, 3H).

EXAMPLE 16

(S)-2-(1-(6-Aminopyrimidin-4-ylamino)propyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0551] 100 mg (0.37 mmol) of (S)-2-(1-aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 130 mg (0.75 mmol) of 6-bromopyrimidine-4-amine and 130 μL (0.75 mmol) of diisopropylethylamine were suspended in 2 mL *tert*-butanol and the mixture was heated to 190 °C for 3 hours under microwave irradiation. Then the solvent was evaporated and the product was purified by preparative HPLC (Symmetry Prep C₁₈ column, mixture of eluents A/B from 5% B to 45% B, in a 30 min. gradient). 10 mg (7% yield) were obtained as a white solid.

LRMS (m/z): 362 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.52 (s, 1H), 7.80 (s, 1H), 7.67 - 7.38 (m, 5H), 7.04 (d, J = 6.9 Hz, 1 H), 6.93 (d, J = 2.7 Hz, 1 H), 6.64 - 6.49 (m, 1 H), 6.05 (s, 2H), 5.43 (s, 1 H), 4.28 (s, 1 H), 1.90 - 1.71 (m, 2H), 0.70 (t, J = 7.2 Hz, 3H).

EXAMPLE 17

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0552] 60 mg (0.34 mmol) of (S)-2-(1-aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 76 mg (0.45 mmol) of 6-chloro-9H-purin-2-amine and 78 μL (0.45 mmol) of diisopropylethylamine were suspended in 2 mL *tert*-butanol and the mixture was heated to 150 °C for 1.5 hours under microwave irradiation. Then the solvent was evaporated and the product was purified by preparative HPLC (Symmetry Prep C₁₈ column, mixture of eluents A/B from 10% B to 40% B, in a 25 min. gradient). 12 mg (13% yield) were obtained as a white solid.

LRMS (m/z): 402 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.11 (s, 1H), 8.27 (s, 1H), 7.69 (s, 2H), 7.62 - 7.42 (m, 4H), 7.34 (s, 1 H), 6.92 (dd, J = 4.2, 1.6 Hz, 1 H), 6.56 (dd, J = 4.3, 2.7 Hz, 1 H), 5.59 (s, 2H), 4.55 (s, 1 H), 1.83 (m, 2H), 0.65 (t, J = 7.2 Hz, 3H).

EXAMPLE 18**(S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile**

[0553] A suspension of 125 mg (0.33 mmol) of (S)-2-(1-aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 56 mg (0.36 mmol) of 4-amino-6-chloropyrimidine-5-carbonitrile (prepared according to the procedure described in WO2010151735A2) and 170 μ L (0.98 mmol) of diisopropylamine in 4 mL of *tert*-butanol was heated with stirring at 120 °C overnight. Then the solvent was removed under vacuum and the product was purified by flash chromatography (dichloromethane to dichloromethane/MeOH/NH₄OH, 100:8:1) followed by a second purification by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound (58 mg, 45% yield) as a white solid.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.79 (s, 1H), 7.67 (s, 1H), 7.60 (d, J = 7.2 Hz, 1H), 7.48 (d, J = 3.5 Hz, 2H), 7.40 - 7.29 (m, 3H), 7.23 (s, 2H), 6.95 (d, J = 2.8 Hz, 1H), 6.64 - 6.57 (m, 1H), 4.68 (dd, J = 13.3, 7.3 Hz, 1H), 1.86 (ddt, J = 28.8, 13.9, 7.1 Hz, 2H), 0.75 (t, J = 7.2 Hz, 3H).

EXAMPLE 19**(R)-2-(1-(9*H*-Purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0554] 70 mg (0.26 mmol) of racemic 2-(1-aminopropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 106 mg (0.53 mmol) of 6-bromo-9*H*-purine and 134 mg (1.04 mmol) of DIEA were suspended in 4 mL of *tert*-butanol and the mixture was heated to 80 °C for 40 hours. Then the solvent was removed in vacuum and the crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain 50 mg (50% yield) of racemic 2-(1-(9*H*-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one.

[0555] The two enantiomers were separated by chiral HPLC with a Chiralpack IA column (5 μ m, 20x250 mm), eluting with a mixture of heptane/isopropanol/diethylamine (85/15/0.1) and 15 mg (15% yield) of the title enantiomer were obtained, corresponding to the second peak to elute (e.e. > 99.5%).

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.88 (s, 1H), 8.11 (m, 2H), 7.97 (s, 1H), 7.62 - 7.56 (m, 1H), 7.56 - 7.41 (m, 3H), 7.40 (s, 1H), 7.29 (s, 1H), 6.92 (dd, J = 4.2, 1.4 Hz, 1H), 6.56 (dd, J = 4.2, 2.7 Hz, 1H), 4.65 (s, 1H), 1.95 (m, 2H), 0.75 (t, J = 7.0 Hz, 3H).

EXAMPLE 20**(S)-2-(1-(9*H*-Purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0556] 90 mg (0.34 mmol) of (S)-2-(1-Aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 134 mg (0.67 mmol) of 6-bromo-9*H*-purine and 174 μ L (1.35 mmol) of diisopropylethylamine were suspended in 3 mL *tert*-butanol and the mixture was heated to 100 °C for 40 hours. Then the solvent was removed in vacuum and the residue was taken up in AcOEt, washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound (59 mg, 45% yield) as a white solid.

LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.92 (s, 1H), 8.19 - 7.95 (m, 3H), 7.68 - 7.42 (m, 4H), 7.29 (d, J = 17.2 Hz, 1H), 7.17 (s, 1H), 6.93 (dd, J = 4.3, 1.6 Hz, 1H), 6.58 (dd, J = 4.2, 2.7 Hz, 1H), 4.95 - 4.69 (m, 1H), 1.45 (d, J = 6.7 Hz, 3H).

EXAMPLE 21**(S)-2-(1-(2-Amino-9*H*-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0557] 75 mg (0.29 mmol) of (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 100 mg (0.59 mmol) of 6-chloro-9*H*-purin-2-amine and 154 μ L (0.88 mmol) of diisopropylethylamine were suspended in 2 mL of 2-propanol and the mixture was heated at 170 °C for 1 hour under microwave irradiation. Then the solvent was removed in vacuum and the residue was taken up in AcOEt, washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1

acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to obtain the title compound (26 mg, 23% yield) as a white solid.

LRMS (m/z): 388 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.07 (s, 1H), 8.17 (s, 1H), 7.67 (s, 1H), 7.63 - 7.56 (m, 2H), 7.51 (s, 2H), 7.46 - 7.28 (m, 2H), 6.93 (dd, J = 4.3, 1.6 Hz, 1H), 6.57 (dd, J = 4.3, 2.7 Hz, 1H), 5.57 (s, 2H), 4.77 (s, 1H), 1.37 (d, J = 6.8 Hz, 3H).

EXAMPLE 22

(S)-2-(1-(6-Aminopyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0558] 100 mg (0.39 mmol) of (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 136 mg (0.78 mmol) of 6-bromopyrimidin-4-amine and 274 μL (1.57 mmol) of diisopropylethylamine were suspended in 2 mL of N-methylpyrrolidone and the mixture was heated at 170 °C for 1 hour under microwave irradiation, then at 180 °C for 2 hours and then at 200 °C for 4 hours. Then water was added to the reaction mixture and the product was extracted with dichloromethane. The organic layer was washed with water and brine, dried over MgSO₄, filtered and the solvent was evaporated under vacuum. The product was purified by preparative HPLC (Symmetry Prep C₁₈ column, mixture of eluents A/B from 5% B to 45% B, in a 30 min. gradient). 23 mg (17% yield) were obtained as a solid.

LRMS (m/z): 348 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.80 (s, 1H), 7.64 - 7.57 (m, 1H), 7.55 - 7.47 (m, 3H), 7.47 - 7.38 (m, 2H), 7.19 (s, 1H), 6.93 (dd, J = 4.2, 1.5 Hz, 1H), 6.58 (dd, J = 4.3, 2.7 Hz, 1H), 6.12 (s, 2H), 5.39 (s, 1H), 4.48 (s, 1H), 1.30 (d, J = 6.8 Hz, 3H).

EXAMPLE 23

(S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0559] 35 mg (0.14 mmol) of (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 23 mg (0.15 mmol) of 4-amino-6-chloropyrimidine-5-carbonitrile (prepared according to the procedure described in WO2010151735A2) and 72 μL (0.41 mmol) of diisopropylethylamine were heated in *tert*-butanol (2 mL) for 21 hours. Then the solvent was removed under vacuum and the crude product was purified by flash chromatography (0-10% methanol in dichloromethane) to give 26 mg (51% yield) of the title compound as a white solid.

LRMS (m/z): 348 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.76 (s, 1H), 7.73 - 7.64 (m, 2H), 7.52 - 7.46 (m, 1H), 7.43 (ddd, J = 8.0, 4.6, 2.3 Hz, 1H), 7.38 - 7.26 (m, 3H), 7.20 (s, 2H), 6.95 (dd, J = 4.3, 1.7 Hz, 1H), 6.61 (dd, J = 4.3, 2.7 Hz, 1H), 4.99 - 4.77 (m, 1H), 1.37 (d, J = 6.7 Hz, 3H).

EXAMPLE 24

2-(1-(6-Amino-9*H*-purin-9-yl)ethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0560] 340 mg (0.90 mmol) of 2-(1-iodoethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 194 mg (1.44 mmol) of 9*H*-purin-6-amine and 310 mg (2.24 mmol) of potassium carbonate were heated in DMF (10 mL) at 50 °C overnight. Then the solvent was removed under vacuum and water was added to the residue. A white solid precipitated and was stirred in water overnight. Then the solid was filtered and washed with water and the product was purified by flash chromatography (dichloromethane to dichloromethane/MeOH/NH₄OH, 100:8:1) to give 116 mg (34% yield) of the title compound as a white solid.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.03 (s, 1H), 7.94 (s, 1H), 7.61 (dd, J = 6.7, 1.8 Hz, 1H), 7.55 (d, J = 2.7 Hz, 1H), 7.49 (td, J = 7.6, 1.3 Hz, 1H), 7.30 (tt, J = 7.5, 1.2 Hz, 1H), 7.19 (s, 2H), 7.09 (td, J = 7.7, 1.4 Hz, 1H), 6.76 (d, J = 8.0 Hz, 1H), 6.45 (dd, J = 2.7, 0.7 Hz, 1H), 5.44 (q, J = 6.8 Hz, 1H), 2.39 (s, 3H), 1.68 (d, J = 6.8 Hz, 3H).

EXAMPLE 25

2-((6-Amino-9*H*-purin-9-yl)methyl)-3-*o*-tolyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0561] This compound was prepared starting from 2-(chloromethyl)-3-*o*-tolyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (2 mg, 0.006 mmol) and following the experimental procedure described in Example 1. Isolation of the compound was done by purification on a 5 g Bond Elut silica cartridge eluting with a mixture of dichloromethane/methanol

to afford 1 mg (50% yield) of the pure title compound.
LRMS (m/z): 441 (M+1)⁺.

EXAMPLE 26**2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one****[0562]**

a) To a solution of di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (125 mg, 0.2 mmol) in 2.5 mL of dry 1,4-dioxane under argon atmosphere 132 μ L of aniline (1.45 mmol) were added and the mixture was heated to reflux with stirring overnight. At the end of this period, the reaction mixture was poured into 50 mL of a 4% aqueous solution of sodium bicarbonate and extracted with ethyl acetate (2x40 mL). The organic layers were mixed and washed with more 4% aqueous solution of sodium bicarbonate, water and brine, and were dried (MgSO₄) and concentrated under reduced pressure to give 187 mg of a solid that was purified by flash chromatography silica (hexane/ethyl acetate). After purification were obtained 136 mg of di-*tert*-butyl [9-(2-([2-(anilino-carbonyl)-3-chloro-1H-pyrrol-1-yl]amino)-2-oxoethyl)-9H-purin-6-yl]imidodicarbonate (91% yield).

b) To a solution of di-*tert*-butyl [9-(2-([2-(anilino-carbonyl)-3-chloro-1H-pyrrol-1-yl]amino)-2-oxoethyl)-9H-purin-6-yl]imidodicarbonate (120 mg, 0.2 mmol) in 1.6 mL of dry 1,4-dioxane under argon atmosphere 179 μ L of phosphorous oxychloride (2 mmol) in 0.8 mL of dry 1,4-dioxane were added and the mixture was heated to reflux with stirring during 2 hours. Afterwards, the reaction mixture was concentrated to dryness co-evaporating with toluene to remove traces of phosphorous oxychloride and the residue was suspended in 1.2 mL of 7N ammonia in methanol heating the reaction at 60°C overnight. At the end of this period, the cooled reaction mixture was poured into 25 mL of a 1/1 mixture of water/brine and extracted with ethyl acetate (2x20 mL). The organic layers were mixed and washed brine, dried (MgSO₄) and concentrated under reduced pressure to give 45 mg of a solid that was purified by flash chromatography silica (dichloromethane/methanol). After purification were obtained 14 mg of the title compound of this example (18% yield).

LRMS (m/z): 393 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 8.04 (s, 1H), 7.95 (s, 1H), 7.59 (d, J = 3.1 Hz, 1 H), 7.53 - 7.46 (m, 5H), 7.23 (s, 2H), 6.66 (d, J = 3.1 Hz, 1 H), 4.97 (s, 2H).

EXAMPLE 27**2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(3-methoxyphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one****[0563]**

a) To a solution of di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (110 mg, 0.21 mmol) in 2 mL of dry 1,4-dioxane under argon atmosphere 160 μ L of 3-methoxyaniline (1.25 mmol) were added and the mixture was heated to reflux with stirring overnight. At the end of this period, the reaction mixture was poured into 50 mL of a 4% aqueous solution of sodium bicarbonate and extracted with ethyl acetate (2x40 mL). The organic layers were mixed and washed with more 4% aqueous solution of sodium bicarbonate, water and brine, and were dried (MgSO₄) and concentrated under reduced pressure to give a residue of 280 mg that was submitted to purification using a 10g Bond Elut silica cartridge eluting with a mixture of hexane/ethyl acetate. After purification were obtained 65 mg of di-*tert*-butyl [9-(2-([2-(3-methoxyphenylcarbamoyl)-3-chloro-1H-pyrrol-1-yl]amino)-2-oxoethyl)-9H-purin-6-yl]imidodicarbonate (42% yield).

b) To a solution of di-*tert*-butyl [9-(2-([2-(3-methoxyphenylcarbamoyl)-3-chloro-1H-pyrrol-1-yl]amino)-2-oxoethyl)-9H-purin-6-yl]imidodicarbonate (65 mg, 0.09 mmol) in 1 mL of dry 1,4-dioxane under argon atmosphere 100 μ L of phosphorous oxychloride (1.1 mmol) in 0.5 mL of dry 1,4-dioxane were added and the mixture was heated to reflux with stirring during 2 hours. Afterwards, the reaction mixture was concentrated to dryness co-evaporating with toluene to remove traces of phosphorous oxychloride and the residue was suspended in 1 mL of 7N ammonia in methanol heating the reaction at 60°C overnight. At the end of this period, the cooled reaction mixture was poured into 25 mL of a 4% aqueous solution of sodium bicarbonate and extracted with ethyl acetate (3x25 mL). The organic layers were mixed and washed brine, dried (MgSO₄) and concentrated under reduced pressure to give a residue that was purified by flash chromatography silica (dichloromethane/methanol). After purification were obtained 12 mg of the

title compound of this example (30% yield).

LRMS (m/z): 423 (M+1)⁺.

¹H NMR (400 MHz, DMSO) 512.02(s, 1H), 8.03 (s, 1H), 7.95 (s, 1H), 7.60 (d, J = 3.0 Hz, 1 H), 7.39 (dd, J = 9.1, 7.7 Hz, 1 H), 7.22 (s, 2H), 7.06 - 6.96 (m, 3H), 6.66 (d, J = 3.0 Hz, 1H), 5.07 (d, J = 16.8 Hz, 1H), 4.99 (d, J = 16.8 Hz, 1H), 3.73 (s, 3H).

EXAMPLE 28

2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(2,4-difluorophenyl)pyrrolo-[1,2-f][1,2,4]triazin-4(3H)-one

[0564]

a) Sodium hexamethyldisilazide (1 M solution in tetrahydrofuran, 965 μ L, 0.97 mmol) was added to a solution of 2,4-difluoroaniline (118 μ L, 1.16 mmol) in 0.5 mL of dry tetrahydrofuran under argon and the mixture was stirred at room temperature during 15 minutes. At the end of this period, the mixture was put in an ice-water bath and a solution of di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (100 mg, 0.19 mmol) in 2 mL of tetrahydrofuran was added maintaining the stirring during 30 minutes at room temperature. Next, the reaction mixture was poured into 25 mL of a saturated aqueous solution of ammonium chloride and extracted with ethyl acetate (2x20 mL). The organic layers were mixed and washed with water and brine, dried (MgSO₄) and concentrated under reduced pressure to give 145 mg of an oil that was purified by flash chromatography silica (dichloromethane/methanol). After purification were obtained 68 mg of *tert*-butyl 9-(2-(3-chloro-2-(2,4-difluorophenylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (65% yield).

b) To a solution of *tert*-butyl 9-(2-(3-chloro-2-(2,4-difluorophenylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (60 mg, 0.09 mmol) in 0.8 mL of dry 1,4-dioxane under argon atmosphere 85 μ L of phosphorous oxychloride (0.93 mmol) in 0.4 mL of dry 1,4-dioxane were added and the mixture was heated to reflux with stirring during 2 hours. Afterwards, the reaction mixture was concentrated to dryness co-evaporating with toluene to remove traces of phosphorous oxychloride and the residue was suspended in 2.4 mL of 7N ammonia in methanol heating the reaction at 60°C overnight. At the end of this period, the cooled reaction mixture was poured into 25 mL of a 1/1 mixture of water/brine and extracted with chloroform (3x25 mL). The organic layers were mixed and washed with brine, dried (MgSO₄) and concentrated under reduced pressure to give 41 mg of a solid that was purified by flash chromatography silica (dichloromethane/methanol). After purification were obtained 9 mg of the title compound of this example (23% yield).

LRMS (m/z): 429 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 7.95 (s, 1 H), 7.89 (s, 1 H), 7.70 (m, 1 H), 7.64 (d, J = 3.0 Hz, 1 H), 7.34 (m, 1 H), 7.22 (m, 1 H), 7.20 (s, 2H), 6.67 (d, J = 3.0 Hz, 1 H), 5.04 (dd, J = 16.7 Hz, 2H).

EXAMPLE 29

2-((6-Amino-9H-purin-9-yl)methyl)-3-benzyl-5-chloropyrrolo[1,2-f][1,2,4]-triazin-4(3H)-one

[0565]

a) To a solution di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (100 mg, 0.2 mmol) in 2 mL of dry 1,4-dioxane under argon atmosphere 127 μ L of benzylamine (1.16 mmol) were added and the mixture was stirred at 30°C during 2 hours. At the end of this period, the reaction mixture was concentrated to dryness under reduced pressure and the residue was purified by flash chromatography silica (dichloromethane/methanol). After purification were obtained 95 mg of *tert*-butyl 9-(2-(2-(benzylcarbamoyl)-3-chloro-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (92% yield).

b) To a solution of *tert*-butyl 9-(2-(2-(benzylcarbamoyl)-3-chloro-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (95 mg, 0.18 mmol) in 1.32 mL of dry 1,4-dioxane under argon atmosphere 165 μ L of phosphorous oxychloride (1.8 mmol) in 0.6 mL of dry 1,4-dioxane were added and the mixture was heated to reflux with stirring during 2 h. Afterwards, the reaction mixture was concentrated to dryness co-evaporating with toluene to remove traces of phosphorous oxychloride and the residue was suspended in 3.8 mL of 7N ammonia in methanol heating the reaction at 60°C overnight. At the end of this period, the cooled reaction mixture was poured into 25 mL of a 1/1 mixture of water/brine and extracted with chloroform (3x25 mL). The organic layers were mixed and washed with brine, dried (MgSO₄) and concentrated under reduced pressure to give 57 mg of a solid that was purified by reverse phase

chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%). After purification were obtained 11 mg of the title compound of this example (15% yield).

LRMS (m/z): 407 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 8.13 (s, 1H), 8.08 (s, 1H), 7.51 (d, J = 3.0 Hz, 1 H), 7.49 - 7.28 (m, 5H), 7.25 (s, 2H), 6.64 (d, J = 3.0 Hz, 1 H), 5.46 (s, 2H), 5.33 (s, 2H).

EXAMPLE 30

2-((6-Amino-9H-purin-9-yl)methyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one

[0566] 9H-purin-6-amine (85 mg, 0.63 mmol) and potassium carbonate (87 mg, 0.63 mmol) were added to a solution of 2-(chloromethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one (137 mg, 0.53 mmol) in 10 mL of DMF. The mixture was stirred at room temperature for 21 hours. The solvent was evaporated to dryness and the crude product was purified by flash chromatography (10% to 20% MeOH/DCM) to yield 100 mg (53% yield) of the title compound as a white solid.

LRMS (m/z): 360 (M+1)⁺.

¹H NMR (250 MHz, DMSO) δ 8.05 (s, 1 H), 7.98 (m, 2H), 7.53 (m, 6H), 7.25 (br s, 2H), 5.04 (s, 2H).

EXAMPLE 31

2-((6-amino-9H-purin-9-yl)methyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-one

[0567] 9H-purin-6-amine (207 mg, 1.53 mmol) and potassium carbonate (211 mg, 1.53 mmol) were added to a solution of 2-(chloromethyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-one (350 mg, 1.27 mmol) in 20 mL of DMF. The mixture was stirred at room temperature for 5 hours and poured into water. A 10% aqueous solution of sodium hydroxide was added until the solution reached pH=11 and then the product was extracted with dichloromethane. The combined organic layers were dried over sodium sulphate, filtered and the solvent was evaporated. The crude product was triturated with dichloromethane (5 mL) and methanol (5 mL) and 130 mg (27% yield) of the title compound were obtained as a white solid.

LRMS (m/z): 374 (M+1)⁺.

¹H NMR (250 MHz, DMSO) δ 8.02 (m, 3H), 7.51 (m, 5H), 7.24 (br s, 2H), 5.12 (s, J = 16.0 Hz, 1H), 4.86 (s, J = 16.0 Hz, 1H), 2.13 (s, 3H)

EXAMPLE 32

2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(pyridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0568]

a) To a suspension of 4-aminopyridine (164 mg, 1.74 mmol) in 3.3 mL of dichloromethane under inert atmosphere was added a 2M solution of trimethyl aluminium in toluene (0.87 mL, 1.74 mmol) and the mixture was stirred during 20 minutes at room temperature. Next, this mixture was cooled in an ice-water bath and a solution of di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (150 mg, 0.29 mmol) in 2.2 mL of dry dichloromethane was added stirring the mixture during 3 days at room temperature. At the end of this period, the reaction mixture was cooled-down with an ice-water bath and 5 mL of water were added followed by a 5% aqueous solution disodium tartrate dihydrate. The resulting mixture was purified directly by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%). After purification were obtained 139 mg of *tert*-butyl 9-(2-(3-chloro-2-(pyridin-4-ylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (93% yield).

b) Starting from *tert*-butyl 9-(2-(3-chloro-2-(pyridin-4-ylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (131 mg, 0.26 mmol) and following the experimental procedure described in Example 26b were obtained 4.2 mg (3.5% yield) of the title compound of this example.

LRMS (m/z): 394 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 8.61 (dd, J = 4.5, 1.6 Hz, 2H), 7.95 (s, 1 H), 7.90 (s, 1H), 7.59 (d, J = 3.0 Hz, 1H), 7.44 (dd, J = 4.5, 1.6 Hz, 2H), 7.18 (s, 2H), 6.64 (d, J = 3.1 Hz, 1 H), 4.97 (s, 2H).

EXAMPLE 33**2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(tetrahydro-2H-pyran-4-yl)pyrrolo-[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0569]

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (131 mg, 0.26 mmol) and following the experimental procedure described in Example 26a but heating the reaction mixture at 80°C during 4 hours were obtained 170 mg (70% yield) of di-*tert*-butyl {9-[2-oxo-2-((2-((tetrahydro-2H-pyran-4-ylamino)carbonyl)-1H-pyrrol-1-yl)amino)ethyl]-9H-purin-6-yl}imidodicarbonate.

b) Starting from di-*tert*-butyl {9-[2-oxo-2-((2-((tetrahydro-2H-pyran-4-ylamino)carbonyl)-1H-pyrrol-1-yl)amino)ethyl]-9H-purin-6-yl}imidodicarbonate (131 mg, 0.26 mmol) and using the conditions described in Example 26b but isolating the product in the following way: the crude was concentrated to dryness and the residue obtained was precipitated from a 1/1 mixture of dimethylsulfoxide/4% aqueous solution of sodium bicarbonate to afford 43 mg after filtration (36% yield) of the title compound of this example.

LRMS (m/z): 401 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 8.22 (s, 1H), 8.21 (s, 1H), 7.51 (d, J = 3.0 Hz, 1H), 7.38 (s, 2H), 6.62 (d, J = 3.0 Hz, 1H), 5.62 (s, 2H), 4.18 - 4.09 (m, 1H), 3.84 - 3.76 (m, 2H), 3.08 - 2.99 (m, 2H), 2.67 - 2.57 (m, 2H), 1.35 - 1.26 (m, 2H).

EXAMPLE 34**2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-methylpiperidin-4-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0570]

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (150 mg, 0.29 mmol) and following the experimental procedure described in Example 26a but heating the reaction mixture at 100°C during 4 hours were obtained 136 mg (86% yield) of *tert*-butyl 9-(2-(3-chloro-2-(1-methylpiperidin-4-ylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate.

b) Starting from *tert*-butyl 9-(2-(3-chloro-2-(1-methylpiperidin-4-ylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (128 mg, 0.24 mmol) and following the experimental procedure described in Example 33b were obtained 21 mg (21% yield) of the title compound of this example.

LRMS (m/z): 414 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 8.21 (s, 1H), 8.21 (s, 1H), 7.52 (d, J = 2.4 Hz, 1H), 7.36 (s, 2H), 6.62 (d, J = 2.5 Hz, 1H), 5.55 (s, 2H), 3.90 - 3.67 (m, 1H), 2.74 - 2.54 (m, 4H), 2.05 (s, 3H), 1.63 - 1.44 (m, 2H), 1.31 - 1.13 (m, 2H).

EXAMPLE 35**(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0571] Starting from (S)-2-(1-aminoethyl)-3-(3-fluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (23 mg, 0.08 mmol) and following the experimental procedure described in Example 20 were obtained 18 mg (50% yield) of the title compound of this example.

LRMS (m/z): 391 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.92 (s, 1H), 8.11-8.00 (2s, 1H), 7.68 (m, 1H), 7.58 - 7.07 (m, 2H), 6.95 (m, 1H), 6.60 (m, 1H), 4.95 (m, 1H), 4.84 (m, 1H), 1.47(d,J=6.6Hz,3H).

EXAMPLE 36**(S)-4-Amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0572] Starting from (S)-2-(1-aminoethyl)-3-(3-fluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (23 mg, 0.08 mmol) and following the experimental procedure described in Example 23 were obtained 13 mg (36% yield) of the title compound of this example.

LRMS (m/z): 391 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.78, 7.74 (2s, 1H), 7.73-7.67 (m, 1H), 7.55-7.27 (m, 2H), 7.22 (br s, 2H), 7.17-7.11 (m, 2H), 6.97 (m, 1 H), 6.63 (m, 1 H), 5.07 - 4.92 (m, 1 H), 4.84 (m, 1 H), 1.38 (d, J = 6.5 Hz, 3H).

EXAMPLE 37

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0573] Starting from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (17 mg, 0.06 mmol) and following the experimental procedure described in Example 20 were obtained 18 mg (71% yield) of the title compound of this example.

LRMS (m/z): 409 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 12.92 (s, 1H), 8.15 - 7.95 (m, 3H), 7.71 (s, 1H), 7.48 (m, 1 H), 7.10 - 7.01 (m, 1 H), 6.99 - 6.94 (m, 1 H), 6.66 - 6.58 (m, 1 H), 5.15 - 5.04 (m, 1H), 4.64 (m, 1 H), 1.48 (d, J = 6.7 Hz, 2H).

EXAMPLE 38

(S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0574] Starting from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (17 mg, 0.06 mmol) and following the experimental procedure described in Example 23 were obtained 13 mg (41% yield) of the title compound of this example.

LRMS (m/z): 409 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 7.78 (s, 1H), 7.75 (dd, J = 2.7, 1.7 Hz, 1H), 7.66 (d, J = 7.4 Hz, 1H), 7.47 (m, 1H), 7.25 (br s, 2H), 7.18 (m, 1H), 6.99 (dd, J = 4.2, 1.7 Hz, 1H), 6.64 (dd, J = 4.3, 2.7 Hz, 1H), 5.15 - 5.04 (m, 1H), 4.68 - 4.61 (m, 1 H), 1.39 (d, J = 6.6 Hz, 2H).

EXAMPLE 39

2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-methylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0575]

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-*d*][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodibutyl carbonate (150 mg, 0.29 mmol) and following the experimental procedure described in Example 26a but using tetrahydrofuran as solvent and stirring the reaction mixture at room temperature during 2 hours were obtained 126 mg (97% yield) of *tert*-butyl 9-(2-(3-chloro-2-(methylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate.

b) Starting from *tert*-butyl 9-(2-(3-chloro-2-(methylcarbamoyl)-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (125 mg, 0.28 mmol) and following the experimental procedure described in Example 26b were obtained 46 mg (48% yield) of the title compound of this example.

LRMS (m/z): 331 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.15 (s, 1H), 8.15 (s, 1H), 7.42 (d, J = 3 Hz, 1H), 7.32 (s, 2H), 6.57 (d, J = 3 Hz, 1 H), 5.58 (s, 2H), 3.46 (s, 3H).

EXAMPLE 40

2-((6-Amino-9H-purin-9-yl)methyl)-3-((1*r*,4*r*)-4-aminocyclohexyl)-5-chloropyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0576]

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-*d*][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodibutyl carbonate (150 mg, 0.29 mmol) and following the experimental procedure described in Example 26a but heating the reaction mixture at 50°C during 2 hours were obtained 138 mg (82% yield) of *tert*-butyl 9-(2-(2-((1*r*,4*r*)-4-aminocyclohexylcarbamoyl)-3-chloro-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate.

b) Starting from *tert*-butyl 9-(2-(2-((1*r*,4*r*)-4-aminocyclohexylcarbamoyl)-3-chloro-1H-pyrrol-1-ylamino)-2-oxoethyl)-9H-purin-6-ylcarbamate (130 mg, 0.24 mmol) and using the conditions described in Example 26b but isolating the

product in the following way: the crude was concentrated to dryness and the residue obtained was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%). After purification were obtained 14 mg (13% yield) of the title compound of this example.

LRMS (m/z): 414 (M+1)⁺.

¹H NMR (600 MHz, DMSO) δ 8.20 (s, 1H), 8.20 (s, 1H), 7.50 (d, J = 3.0 Hz, 1 H), 7.35 (s, 2H), 6.61 (d, J = 3.0 Hz, 1 H), 5.55 (s, 2H), 3.88 - 3.79 (m, 1 H), 2.40 - 2.37 (m, 1 H), 1.67 - 1.58 (m, 4H), 1.35 - 1.27 (m, 2H), 0.83 - 0.74 (m, 4H).

EXAMPLE 41

(R)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0577]

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodibutyl carbonate (150 mg, 0.29 mmol) and following the experimental procedure described in Example 26a but heating the reaction mixture at 40°C during 1.5 hours were obtained 138 mg (78% yield) of *tert*-butyl [9-(2-([3-chloro-2-((1R)-1-phenylethyl)amino)carbonyl]-1H-pyrrol-1-yl)amino]-2-oxoethyl)-9H-purin-6-yl]carbamate.

b) A solution of bromine (37 mg, 0.24 mmol) in dichloromethane (100 µL) was added dropwise to a solution of triphenylphosphine (63 mg, 0.24 mmol) in dichloromethane (1 ml) under nitrogen. The solution was stirred for 30 min, and triethylamine (78 µL, 0.56 mmol) and *tert*-butyl [9-(2-([3-chloro-2-((1R)-1-phenylethyl)amino)carbonyl]-1H-pyrrol-1-yl)amino]-2-oxoethyl)-9H-purin-6-yl]carbamate (100 mg, 0.19 mmol) were added. The reaction mixture was refluxed for 1.5 h, and quenched with 10 % aqueous sodium bicarbonate solution. The organic phase was separated, dried (sodium sulphate) and concentrated to obtain an oil that was treated with 6 ml of a 7M methanolic solution of ammonia at 60 °C in a sealed vessel. The solvent was then evaporated to obtain 66 mg (34% yield, 48% purity) of *tert*-butyl [9-((5-chloro-4-oxo-3-((1R)-1-phenylethyl)-3,4-dihydropyrrolo[2,1-f][1,2,4]triazin-2-yl)methyl)-9H-purin-6-yl]carbamate.

c) *tert*-Butyl [9-((5-chloro-4-oxo-3-((1R)-1-phenylethyl)-3,4-dihydropyrrolo[2,1-f][1,2,4]triazin-2-yl)methyl)-9H-purin-6-yl]carbamate (66 mg, 48% purity, 0.061 mmol) were dissolved in 4 ml of a 4M hydrogen chloride solution in dioxane and heated to 45°C for 1 hour. Then, the crude was concentrated to dryness and the residue obtained was purified by flash chromatography silica (dichloromethane/methanol). After purification were obtained 22 mg of the title compound of this example (85% yield).

LRMS (m/z): 421 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.21 (s, 1H), 8.11 (s, 1H), 7.47 (d, J = 3.0 Hz, 1 H), 7.40 - 7.21 (m, 7H), 6.58 (d, J = 3.0 Hz, 1 H), 5.80 (m, 1 H), 5.58 (m, 2H), 1.81 (d, J = 6.6 Hz, 3H).

EXAMPLE 42

(S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0578]

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodibutyl carbonate (150 mg, 0.29 mmol) and following the experimental procedure described in Example 26a but heating the reaction mixture at 40°C during 1.5 hours were obtained 132 mg (75% yield) of *tert*-butyl [9-(2-([3-chloro-2-((1S)-1-phenylethyl)amino)carbonyl]-1H-pyrrol-1-yl)amino]-2-oxoethyl)-9H-purin-6-yl]carbamate.

b) Starting from *tert*-butyl [9-(2-([3-chloro-2-((1S)-1-phenylethyl)amino)carbonyl]-1H-pyrrol-1-yl)amino]-2-oxoethyl)-9H-purin-6-yl]carbamate (132 mg, 0.24 mmol) and following the experimental procedure described in Example 41 b were obtained 132 mg (55% yield, 52% purity) of *tert*-butyl [9-((5-chloro-4-oxo-3-((1S)-1-phenylethyl)-3,4-dihydropyrrolo[2,1-f][1,2,4]triazin-2-yl)methyl)-9H-purin-6-yl]carbamate.

c) Starting from *tert*-butyl *tert*-butyl [9-((5-chloro-4-oxo-3-((1S)-1-phenylethyl)-3,4-dihydropyrrolo[2,1-f][1,2,4]triazin-2-yl)methyl)-9H-purin-6-yl]carbamate (132 mg, 52 % purity, 0.13 mmol) and following the experimental procedure described in Example 41c were obtained 48 mg of the title compound of this example (87% yield).

LRMS (m/z): 421 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.21 (s, 1H), 8.11 (s, 1H), 7.47 (d, J = 3.0 Hz, 1 H), 7.39 - 7.20 (m, 7H), 6.59 (d, J =

3.0 Hz, 1 H), 5.80 (s, 1 H), 5.53 (m, 2H), 1.81 (d, J = 6.7 Hz, 3H).

EXAMPLE 43

5 (S)-4-Amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

10 [0579] Starting from (S)-2-(1-aminoethyl)-3-(pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one dihydrochloride (60 mg, 0.13 mmol) and following the experimental procedure described in Example 23 were obtained 8 mg (9% yield) of the title compound of this example.

LRMS (m/z): 374 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.47 (d, J = 4.0 Hz, 1 H), 7.83 (t, J = 7.8 Hz, 1 H), 7.75 (m, 2H), 7.69 (d, J = 7.4 Hz, 1 H), 7.44 (d, J = 6.6 Hz, 1 H), 7.37 - 7.29 (m, 1H), 7.21 (bs, 2H), 7.01 (dd, J = 4.3, 1.6 Hz, 1H), 6.65 (dd, J = 4.3, 2.7 Hz, 1 H), 5.10 (m, 1 H), 1.40 (d, J = 6.7 Hz, 4H).

15 EXAMPLE 44

(S)-2-(1-(9*H*-Purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

20 [0580] 45 mg (0.16 mmol) of (S)-3-phenyl-2-(pyrrolidin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 35 mg (0.17 mmol) of 6-bromo-9*H*-purine and 25 μl (0.17 mmol) of diisopropylethylamine were stirred in *tert*-butanol (5 ml) at 80 °C overnight. Then the solvent was removed in vacuum and the residue was taken up in AcOEt, washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to

25 obtain the title compound (22 mg, 80% yield) as a white solid.

LRMS (m/z): 399 (M+1)⁺.

EXAMPLE 45

30 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0581] The title compound (53 mg, 58% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

35 LRMS (m/z): 441 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.83 (d, J = 2.8 Hz, 1H), 7.75 (s, 1H), 7.62 (d, J = 6.8 Hz, 1 H), 7.53 (d, J = 7.9 Hz, 1H), 7.44 (m, 1 H), 7.37 - 7.17 (m, 5H), 6.99 (d, J = 2.8 Hz, 1 H), 4.89 (p, J = 6.5 Hz, 1 H), 1.36 (d, J = 6.6 Hz, 3H).

40 EXAMPLE 46

(S)-2-(1-(9*H*-Purin-6-ylamino)ethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

45 [0582] The title compound (53 mg, 58% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 6-bromo-9*H*-purine following the experimental procedure described in example 20.

LRMS (m/z): 441 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.95 (s, 1H), 8.19 - 8.02 (m, 3H), 7.78 (d, J = 2.7 Hz, 1H), 7.59 (m, 1 H), 7.49 (d, J = 7.5 Hz, 2H), 7.34 (m, 1 H), 7.15 (t, J = 7.8 Hz, 1 H), 6.98 (d, J = 2.8 Hz, 1 H), 4.85 (m, 1 H), 1.47 (d, J = 6.7 Hz, 3H).

50 EXAMPLE 47

(S)-4-Amino-6-(1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

55 [0583] The title compound (67 mg, 59% yield) was obtained from (S)-2-(1-Aminoethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 423 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.81 (d, 1 H), 7.77 (s, 1 H), 7.66 (d, 1 H), 7.54 (d, 1 H), 7.50 - 7.41 (m, 1 H), 7.40 - 7.12 (m, 6H), 6.91 (s, 1 H), 4.97 - 4.85 (m, 1 H), 1.38 (d, 3H).

EXAMPLE 48

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0584] The title compound (42 mg, 60% yield) was obtained from (S)-2-(1-Aminoethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20.
LRMS (m/z): 423 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.94 (s, 1H), 8.20 - 7.99 (m, 2H), 7.86 - 6.80 (m, 9H), 4.86 (s, 1 H), 1.46 (d, 3H).

EXAMPLE 49

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0585] (S)-2-(1-Aminoethyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3H)-one (80 mg, 0.31 mmol), 6-bromo-9H-purine (125 mg, 0.63 mmol) and DIPEA (162 μl, 1.25 mmol) were stirred in *tert*-butanol at 100 °C for 24 hours. The solvent was removed in vacuum and the residue was dissolved in dichloromethane. The organic solution was washed with water and brine, dried and the solvent was removed under reduced pressure. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 47 mg (40% yield) of the title compound.

LRMS (m/z): 374 (M+1)⁺.

¹H RMN (400 MHz, DMSO) δ 8.14 (s, 1H), 8.13 - 8.00 (m, 1H), 7.70 - 7.42 (m, 4H), 7.27 (m, 2H), 4.83 (q, 1 H), 1.47 (d, J = 6.7 Hz, 3H).

EXAMPLE 50

(S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0586] Prepared according to the experimental procedure described in Example 48 from (S)-2-(1-Aminoethyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3H)-one (70 mg, 0.27 mmol), and 4-amino-6-chloropyrimidine-5-carbonitrile (51 mg, 0.33 mmol). The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 43 mg (42% yield) of the title compound.

LRMS (m/z): 374 (M+1)⁺.

¹H RMN (400 MHz, DMSO) δ 8.14 (s, 1 H), 7.78 (s, 1 H), 7.64 (d, 1 H), 7.60 (s, 1 H), 7.55 (d, 1 H), 7.47 (dd, 1 H), 7.40 - 7.28 (m, 2H), 7.25 (s, 2H), 4.90 (m, 1 H), 1.39 (d, 3H).

EXAMPLE 51

2-(1-(9H-Purin-6-ylamino)-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0587] 2-(1-Amino-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (70 mg, 0.22 mmol), 6-bromo-9H-purine (86 mg, 0.43 mmol) and DIEA (151 μl, 0.87 mmol) were stirred in *tert*-butanol at 100 °C for 16 hours. An excess of 150 μl of DIEA was added and the mixture was stirred at 100 °C for 24 hours more. The solvent was removed in vacuum and the residue was dissolved in ethyl acetate. The organic solution was washed with water and brine, dried and the solvent was removed under reduced pressure. The product was purified by flash chromatography (dichloromethane to dichloromethane/MeOH/NH₄OH, 100:8:1) to give 24 mg (25% yield) of the title compound as a white solid.

LRMS (m/z): 441 (M+1)⁺.

¹H RMN (400 MHz, DMSO) δ 13.02 (s, 1H), 8.39 (d, 1H), 8.17 (s, 1H), 8.14 (s, 1 H), 7.60 (dd, 2H), 7.54 (t, 2H), 7.42 (t, 1 H), 7.32 (t, 1 H), 6.96 (dd, 1 H), 6.59 (dd, 1H), 5.22 (s, 1 H), 3.17 - 2.97 (m, 2H).

EXAMPLE 52**4-Amino-6-(3,3,3-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile**

[0588] 2-(1-Amino-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (70 mg, 0.22 mmol), 4-amino-6-chloropyrimidine-5-carbonitrile (67 mg, 0.43 mmol) and DIEA (151 μ l, 0.87 mmol) were stirred in *tert*-butanol at 100 °C for 16 hours. An excess of 150 μ l of DIEA was added and the mixture was stirred at 100 °C for 24 hours more. The solvent was removed in vacuum and the residue was dissolved in ethyl acetate. The organic solution was washed with water and brine, dried and the solvent was removed under reduced pressure. The product was purified by flash chromatography (0% to 5% MeOH/DCM) to give 40 mg (42% yield) of the title compound as a white solid.

LRMS (m/z): 441 (M+1)⁺.

¹H RMN (400 MHz, DMSO) δ 7.94 (d, 1 H), 7.84 (s, 1 H), 7.69 (dd, 1 H), 7.57 - 7.46 (m, 2H), 7.38 (ddd, 4H), 6.97 (dd, 1 H), 6.63 (dd, 1 H), 5.22 (dd, 1 H), 3.09 - 2.91 (m, 2H).

EXAMPLE 53**(S)-4-Amino-6-(2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile**

[0589] (S)-3-Phenyl-2-(pyrrolidin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (50 mg, 0.18 mmol), 4-amino-6-chloropyrimidine-5-carbonitrile (61 mg, 0.40 mmol) and DIEA (150 μ l, 0.86 mmol) were stirred in *tert*-butanol at 120 °C for 16 hours. An excess of 50 mg of 4-amino-6-chloropyrimidine-5-carbonitrile was added and the mixture was stirred at 100 °C for 24 hours more. The solvent was removed in vacuum and the residue was dissolved in dichloromethane. The organic solution was washed with water and brine, dried and the solvent was removed under reduced pressure. The product was purified by flash chromatography (0% to 3% MeOH/DCM) to give 30 mg (41% yield) of the title compound as a white solid.

LRMS (m/z): 399 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.03 (s, 1H), 7.69 - 7.46 (m, 6H), 7.28 (s, 2H), 6.91 (dd, 1 H), 6.53 (m, 1 H), 2.26 - 2.08 (m, 3H), 2.00 - 1.90 (m, 2H), 1.86 - 1.74 (m, 2H).

EXAMPLE 54**(S)-3-Phenyl-2-(1-(pyrazolo[1,5-a]pyrimidin-7-ylamino)ethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0590] The title compound (38 mg, 35% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (preparation 22) and 7-Chloropyrazolo[1,5-a]pyrimidine (preparation 89) following the experimental procedure described in example 26.

LRMS (m/z): 372 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.22 (d, J = 7.9 Hz, 1 H), 8.09 (d, J = 2.1 Hz, 1 H), 7.93 (d, J = 5.2 Hz, 1 H), 7.76 - 7.70 (m, 1 H), 7.55 (d, J = 7.9 Hz, 1 H), 7.45 (t, J = 7.5 Hz, 1H), 7.22 (dd, J = 12.8, 7.2 Hz, 2H), 6.97 (dd, J = 4.2, 1.5 Hz, 1H), 6.87 (t, J = 7.5 Hz, 1 H), 6.63 (dd, J = 4.1, 2.8 Hz, 1 H), 6.41 (d, J = 2.1 Hz, 1 H), 5.59 (d, J = 5.3 Hz, 1 H), 4.62 - 4.47 (m, 1 H), 1.53 (d, J = 6.5 Hz, 3H).

EXAMPLE 55**2-((6-Amino-9*H*-purin-9-yl)methyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one**

[0591] The title compound (36 mg, 52% yield) was obtained from 2-(chloromethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and adenine following the experimental procedure described in example 1.

LRMS (m/z): 409 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.04 (s, 1H), 7.96 (s, 1H), 7.66 (d, 1H), 7.58 - 7.42 (m, 5H), 7.35 - 7.11 (m, 3H), 6.86 (d, 1H), 5.03 (s, 2H).

EXAMPLE 56**(S)-2-(1-(2-Amino-9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0592] (S)-3-Phenyl-2-(pyrrolidin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (70 mg, 0.25 mmol), 6-bromo-9H-purin-2-amine (169 mg, 1.00 mmol) and DIEA (174 μ l, 1.00 mmol) were stirred in *tert*-butanol at 150 °C for 1.5 hours under microwave irradiation. The solvent was removed in vacuum and the residue was dissolved in dichloromethane. The organic solution was washed with brine, dried and the solvent was removed under reduced pressure. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 104 mg (100% yield) of the title compound.

LRMS (m/z): 414 (M+1)⁺.

EXAMPLE 57**(S)-2-(1-(4,6-diamino-1,3,5-triazin-2-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0593] The title compound (26 mg, 24% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (preparation 22) and 6-chloro-1,3,5-triazine-2,4-diamine (purchased from Aldrich) following the experimental procedure described in example 26.

LRMS (m/z): 364 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.65 - 7.57 (m, 2H), 7.45 (m, 4H), 6.91 (dd, J = 4.3, 1.7 Hz, 1 H), 6.83 (d, J = 7.4 Hz, 1 H), 6.56 (dd, J = 4.3, 2.7 Hz, 1 H), 5.99 (br s, 4H), 4.48 (p, J = 6.6 Hz, 1 H), 1.20 (d, J = 6.8 Hz, 3H).

EXAMPLE 58**(S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-(5-fluoropyridin-2-yl)ethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one****[0594]**

a) Starting from di-*tert*-butyl 9-((5-chloro-4-oxo-4H-pyrrolo[1,2-d][1,3,4]oxadiazin-2-yl)methyl)-9H-purin-6-ylimidodicarbonate (150 mg, 0.29 mmol) and following the experimental procedure described in Example 26a but heating the reaction mixture at 30°C during 2 hours were obtained 188 mg (81% yield) of di-*tert*-butyl 9-(2-([3-chloro-2-((1S)-1-(5-fluoropyridin-2-yl)ethyl)amino]carbonyl)-1H-pyrrol-1-yl]amino)-2-oxoethyl)-9H-purin-6-yl]imidodicarbonate.

b) Starting from di-*tert*-butyl 9-(2-([3-chloro-2-((1S)-1-(5-fluoropyridin-2-yl)ethyl)amino]carbonyl)-1H-pyrrol-1-yl]amino)-2-oxoethyl)-9H-purin-6-yl]imidodicarbonate (185 mg, 0.28 mmol) and following the experimental procedure described in Example 41b were obtained 232 mg (47% yield, 31% purity) of (S)-*tert*-butyl 9-((5-chloro-3-(1-(5-fluoropyridin-2-yl)ethyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)methyl)-9H-purin-6-yl)carbamate.

c) Starting from (S)-*tert*-butyl 9-((5-chloro-3-(1-(5-fluoropyridin-2-yl)ethyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)methyl)-9H-purin-6-yl)carbamate (232 mg, 31 % purity, 0.13 mmol) and following the experimental procedure described in Example 41c were obtained 44 mg of the title compound of this example (75% yield).

LRMS (m/z): 440 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.41 (s, 1H), 8.19 (s, 1H), 8.11 (s, 1H), 7.80 - 7.63 (m, 1 H), 7.61 - 7.45 (m, 2H), 7.30 (s, 2H), 6.59 (d, 1 H), 5.91 - 5.44 (m, 3H), 1.85 (d, 3H).

EXAMPLE 59**(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0595] The title compound (11 mg, 20% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-chloro-9H-purin-2-amine following the experimental procedure described in example 21.

LRMS (m/z): 424 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.04 (d, 1H), 7.69 (dd, 1H), 7.67 - 7.65 (m, 1H), 7.53 - 7.38 (m, 2H), 7.20 - 7.09 (m, 2H), 6.99 - 6.93 (dd, 1 H), 6.61 (dd, 1 H), 5.57 (s, 2H), 4.95 - 4.85 (m, 1 H), 1.43 (d, 3H).

EXAMPLE 60**(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0596] The title compound (35 mg, 74% yield) was obtained from (S)-2-(1-aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 398 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.89 (d, J = 2.9 Hz, 1 H), 7.78 (s, 1 H), 7.62 (d, J = 6.8 Hz, 1 H), 7.54 (d, J = 7.9 Hz, 1 H), 7.50 - 7.44 (m, 1 H), 7.39 - 7.30 (m, 3H), 7.24 (br s, 1 H), 7.21 (d, J = 2.9 Hz, 1 H), 4.91 (p, J = 6.6 Hz, 1 H), 1.38 (d, J = 6.7 Hz, 3H).

EXAMPLE 61**(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0597] The title compound (35 mg, 74% yield) was obtained from (S)-2-(1-aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 6-bromo-9H-purine following the experimental procedure described in example 20.

LRMS (m/z): 398 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.96 (s, 1H), 8.10 (m, 3H), 7.83 (d, J = 2.8 Hz, 1H), 7.64 - 7.45 (m, 3H), 7.42 - 7.29 (m, 1 H), 7.18 (d, J = 3.0 Hz, 2H), 4.92 - 4.74 (m, 1 H), 1.46 (d, J = 6.7 Hz, 3H).

EXAMPLE 62**(R)-2-(1-(9H-Purin-6-ylamino)-2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0598] The title compound (25 mg, 50% yield) was obtained from (R)-2-(1-amino-2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20.

LRMS (m/z): 389 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.98 (s, 1 H), 8.17 (br s, 1 H), 8.07 (s, 1 H), 7.62 (dd, 1 H), 7.52 - 7.21 (m, 6H), 6.92 (dd, 1 H), 6.56 (dd, 1 H), 5.04 (t, 1 H), 4.84 - 4.74 (m, 1 H), 3.86 (td, 1 H), 3.69 (td, 1 H).

EXAMPLE 63**(R)-4-Amino-6-(2-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0599] The title compound (20 mg, 39% yield) was obtained from (R)-2-(1-amino-2-hydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 389 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.82 (s, 1H), 7.67 (dd, 1 H), 7.50 - 7.24 (m, 8H), 6.93 (dd, 1 H), 6.59 (dd, 1 H), 4.95 (t, 1 H), 4.80 - 4.74 (m, 1 H), 3.79 (dt, 1 H), 3.55 (dt, 1 H).

EXAMPLE 64**(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0600] (S)-2-(1-Aminoethyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3H)-one (80 mg, 0.31 mmol), 6-chloro-9H-purin-2-amine (106 mg, 0.63 mmol) and DIEA (109 μl, 0.63 mmol) were stirred in 1-butanol (2 mL) at 150 °C for 2 hours under microwave irradiation. The solvent was removed in vacuum and the residue was dissolved in ethyl acetate. The organic solution was washed with water and brine, dried and the solvent was removed under reduced pressure. The product was purified by flash chromatography (dichloromethane to dichloromethane/MeOH/NH₄OH, 100:8:1) to give 25 mg (20% yield) of the title compound as a white solid.

LRMS (m/z): 389 (M+1)⁺.

¹H RMN (400 MHz, DMSO) δ 12.10 (s, 1H), 8.02 (d, 1H), 7.75 - 7.27 (m, 7H), 5.59 (s, 2H), 4.71 (s, 1 H), 2.48 (m, 1 H),

1.36 (d, 3H).

EXAMPLE 65

5 (S)-2-(1-(7H-Pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0601] The title compound (17 mg, 13% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (preparation 22) and 4-Chloropyrrolo[2,3-D]pyrimidine (purchased from Alfa Aesar) following the experimental procedure described in example 26.

10 LRMS (m/z): 372 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 11.53 (s, 1 H), 7.99 (s, 1 H), 7.88 (br s, 1 H), 7.62 (m, 1 H), 7.57 - 7.46 (m, 3H), 7.34 (t, J = 7.2 Hz, 1 H), 7.20 (t, J = 7.4 Hz, 1 H), 7.10 (s, 1 H), 6.95 (dd, J = 4.2, 1.6 Hz, 1 H), 6.59 (dd, J = 4.9, 2.1 Hz, 2H), 4.93-4.69 (m, 1 H), 1.45 (d, J = 6.7 Hz, 3H).

15 EXAMPLE 66

(S)-4-Amino-6-(methyl(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethyl)amino)pyrimidine-5-carbonitrile

20 **[0602]** The title compound (24 mg, 22% yield) was obtained from (S)-2-(1-(methylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 387 (M+1)⁺.

25 ¹H NMR (400 MHz, DMSO) δ 7.74 (dd, J = 2.6, 1.7 Hz, 1H), 7.64 (s, 1H), 7.50 (dd, J = 6.9, 1.1 Hz, 1H), 7.35 (td, J = 7.7, 1.7 Hz, 1H), 7.24 - 7.11 (m, 4H), 6.97 (dd, J = 4.3, 1.6 Hz, 1 H), 6.64 (dd, J = 4.3, 2.7 Hz, 1 H), 5.79 (q, J = 6.6 Hz, 1 H), 3.02 (s, 3H), 1.38 (d, J = 6.6 Hz, 3H).

EXAMPLE 67

30 (S)-2-(1-(Methyl(9H-purin-6-yl)amino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0603] The title compound (35 mg, 32% yield) was obtained from (S)-2-(1-(methylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20.

LRMS (m/z): 387 (M+1)⁺.

35

EXAMPLE 68

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

40 **[0604]** 100 mg (0.37 mmol) of (S)-2-(1-aminoethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one, 148 mg (0.74 mmol) of 6-bromo-9H-purine and 260 μL (1.49 mmol) of diisopropylethylamine were heated at 100 °C in *tert*-butanol (2 mL) for 16 hours. Then the solvent was removed under vacuum and the crude product was purified by flash chromatography (dichloromethane to dichloromethane/MeOH/NH₄OH, 100:8:1) to give 94 mg (65% yield) of the title compound as a white solid.

45 LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.92 (s, 1H), 8.13 (s, 1H), 8.08 (s, 1H), 8.01 (s, 1H), 7.48 (t, 4H), 7.33 (s, 1H), 7.20 (s, 1H), 6.39 (d, 1H), 4.88 - 4.73 (m, 1H), 2.38 (s, 3H), 1.43 (d, 3H).

EXAMPLE 69

50

(S)-4-Amino-6-(1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0605] The title compound was prepared following the experimental procedure described in Example 68 from (S)-2-(1-aminoethyl)-5-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (100 mg, 0.37 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (115 mg, 0.74 mmol). The product was purified by flash chromatography (0% to 5% MeOH/DCM) to give 94 mg (65% yield) of the title compound as a white solid.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 13.02 (s, 1H), 8.39 (d, 1H), 8.17 (s, 1H), 8.14 (s, 1H), 7.60 (dd, 2H), 7.54 (t, 2H), 7.42 (t, 1H), 7.32 (t, 1H), 6.96 (dd, 1H), 6.59 (dd, 1H), 5.22 (s, 1H), 3.17 - 2.97 (m, 2H).

EXAMPLE 70

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-7-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0606] The title compound (35 mg, 68% yield) was obtained from (S)-2-(1-aminoethyl)-7-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.92 (s, 1H), 8.12 (s, 1H), 8.07 - 7.98 (m, 2H), 7.49 (m, 2H), 7.41 (d, J = 7.8 Hz, 1H), 7.33 - 7.24 (m, 1H), 7.15 - 7.05 (m, 1H), 6.87 (d, J = 4.2 Hz, 1H), 6.41 (d, J = 4.1 Hz, 1H), 4.94 (m, 1H), 2.41 (s, 3H), 1.47 (d, J = 6.6 Hz, 3H).

EXAMPLE 71

(S)-4-Amino-6-(1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0607] The title compound (35 mg, 68% yield) was obtained from (S)-2-(1-aminoethyl)-7-methyl-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.75 (s, 1H), 7.63 (d, J = 7.1 Hz, 1H), 7.49 - 7.38 (m, 2H), 7.27 (m, 5H), 6.88 (d, J = 4.1 Hz, 1H), 6.44 (d, J = 3.9 Hz, 1H), 5.02 - 4.90 (m, 1H), 2.47 (s, 3H).

EXAMPLE 72

(S)-2-(4,4-Difluoro-1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0608] The title compound was prepared following the experimental procedure described in Example 68 from (S)-2-(4,4-difluoropyrrolidin-2-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (100 mg, 0.32 mmol) and 6-bromo-9H-purine (126 mg, 0.63 mmol). The product was purified by flash chromatography (DCM to DCM/MeOH/NH₄OH, 100:8:1) to give 109 mg (85% yield) of the title compound as a white solid.

LRMS (m/z): 435 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ (mixture of 2 conformers) 13.18 (s, 1H), 8.40 - 8.01 (m, 3H), 7.80 - 7.36 (m, 5H), 6.99 - 6.82 (m, 1H), 6.58 - 6.38 (m, 1H), 5.02 - 3.98 (m, 3H), 3.11 - 2.85 (m, 2H).

EXAMPLE 73

(S)-4-Amino-6-(4,4-difluoro-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile

[0609] The title compound was prepared following the experimental procedure described in Example 68 from (S)-2-(4,4-difluoropyrrolidin-2-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (100 mg, 0.32 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (59 mg, 0.38 mmol). The product was purified by flash chromatography (0% to 7% MeOH/DCM) to give 57 mg (42% yield) of the title compound as a white solid.

LRMS (m/z): 435 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.14 (s, 1H), 7.80 - 7.40 (m, 7H), 6.95 (dd, 1H), 6.57 (dd, 1H), 4.91 (s, 1H), 4.53 - 4.21 (m, 2H), 3.01 - 2.74 (m, 1H), 2.48 - 2.40 (m, 1H).

EXAMPLE 74

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-6-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0610] The title compound (136 mg, 47% yield) was obtained from (S)-2-(1-aminoethyl)-6-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent.

LRMS (m/z): 391 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.83 (s, 1H), 8.14 (br s, 1H), 8.07 (s, 1H), 7.78 (d, 1H), 7.59-7.11 (m, 6H), 6.83 (d, 1H), 4.91 - 4.77 (m, 1H), 1.45 (d, 3H).

5 EXAMPLE 75

(S)-4-Amino-6-(1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

10 **[0611]** The title compound (241 mg, 75% yield) was obtained from (S)-2-(1-aminoethyl)-6-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent.

LRMS (m/z): 391 (M+1)⁺.

15 ¹H NMR (400 MHz, DMSO) δ 7.85 (d, 1H), 7.76 (s, 1H), 7.68 (d, 1H), 7.55 - 7.09 (m, 7H), 6.85 (d, 1H), 4.95 - 4.83 (m, 1H), 1.36 (d, 3H).

EXAMPLE 76

2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

20 **[0612]** The title compound (238 mg, 64% yield) was obtained from 2-((S)-1-aminoethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 401 (M+1)⁺.

25 ¹H NMR (400 MHz, DMSO) δ 12.98 (s, 1H), 8.45 - 8.03 (m, 3H), 8.13 (s, 1H), 7.53 (dd, 1H), 7.48 - 7.01 (m, 5H), 6.74 (dd, 1H), 6.51 (dd, 1H), 5.95 - 5.55 (m, 2H), 1.91 (d, 3H), 1.72 - 1.42 (m, 3H).

EXAMPLE 77

30 **4-Amino-6-((S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0613] The title compound (247 mg, 70% yield) was obtained from 2-((S)-1-aminoethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 401 (M+1)⁺.

35 ¹H NMR (400 MHz, DMSO) δ 8.18 - 7.88 (m, 2H), 7.60 (dd, 1H), 7.46 - 7.00 (m, 7H), 6.77 (dd, 1H), 6.54 (dd, 1H), 5.83 - 5.59 (m, 1H), 5.59 - 5.25 (m, 1H), 1.87 (d, 3H), 1.67 - 1.37 (m, 3H).

40 EXAMPLE 78

(S)-4-Amino-6-(1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

45 **[0614]** The title compound (27 mg, 22% yield) was obtained from (S)-2-(1-aminoethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 130°C during 15 hours.

LRMS (m/z): 401 (M+1)⁺.

50 ¹H NMR (400 MHz, DMSO) δ 7.87 (d, 1H), 7.79 (dd, 1H), 7.56 (s, 1H), 7.27 - 7.10 (m, 3H), 7.06 (t, 1H), 7.01 (dd, 1H), 6.83 (d, 1H), 6.66 (dd, 1H), 5.09 - 4.97 (m, 1H), 2.04 (s, 3H), 2.00 (s, 3H), 1.38 (d, 3H).

EXAMPLE 79

(S)-2-((9H-Purin-6-ylamino)methyl)-3-(1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

55 **[0615]** The title compound (116 mg, 41% yield) was obtained from (S)-2-(aminomethyl)-3-(1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 80°C during 15 hours.

LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.95 (s, 1 H), 8.38 - 7.94 (m, 3H), 7.54 (dd, 1 H), 7.44 - 7.06 (m, 5H), 6.77 (dd, 1 H), 6.51 (dd, 1 H), 5.99 - 5.53 (m, 1 H), 4.90 - 4.80 (bm, 2H), 1.86 (d, 3H).

5 EXAMPLE 80

(S)-4-Amino-6-((4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile

10 **[0616]** The title compound (107 mg, 38% yield) was obtained from (S)-2-(aminomethyl)-3-(1-phenylethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 50°C during 15 hours.

LRMS (m/z): 387 (M+1)⁺.

15 ¹H NMR (400 MHz, DMSO) δ 8.00 (s, 1H), 7.95 - 7.83 (m, 1H), 7.58 (dd, 1H), 7.47 - 7.18 (m, 7H), 6.79 (dd, 1H), 6.53 (dd, 1H), 5.83 - 5.49 (bm, 1H), 4.81 - 4.46 (bm, 2H), 1.85 (d, 3H).

EXAMPLE 81

(S)-2-(1-(5-Fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

20 **[0617]** The title compound (10 mg, 13% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (preparation 22) and 4-chloro-5-fluoro-7H-pyrrolo[2,3-d]pyrimidine (purchased from Matrix Scientific) following the experimental procedure described in example 26.

LRMS (m/z): 390 (M+1)⁺.

25 ¹H NMR (400 MHz, DMSO) δ 11.39 (s, 1H), 7.94 (s, 1 H), 7.68 - 7.64 (m, 1H), 7.52 (d, J = 8.4 Hz, 1H), 7.43 (m, 2H), 7.25 (m, 2H), 7.14 (td, J = 7.8, 0.8 Hz, 1H), 7.06 (t, J = 2.4 Hz, 1 H), 6.94 (dd, J = 4.3, 1.6 Hz, 1 H), 6.60 (dd, J = 4.2, 2.7 Hz, 1 H), 4.97 (p, J = 6.7 Hz, 1 H), 1.45 (d, J = 6.7 Hz, 3H)

EXAMPLE 82

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

30 **[0618]** The title compound (25 mg, 20% yield) was obtained from (S)-2-(1-aminoethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 130°C during 15 hours.

LRMS (m/z): 401 (M+1)⁺.

35 ¹H NMR (400 MHz, DMSO) δ 12.68 (s, 1 H), 8.16 - 7.80 (m, 3H), 7.77 (dd, 1 H), 7.24 - 7.11 (m, 1H), 7.09 - 6.96 (m, 2H), 6.66 (dd, 1H), 6.60 - 6.48 (m, 1H), 5.19 - 5.01 (m, 1 H), 2.05 (s, 3H), 1.90 (s, 3H), 1.45 (d, 3H).

40 EXAMPLE 83

(S)-4-Amino-6-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

45 **[0619]** The title compound (24 mg, 42% yield) was obtained from (S)-2-(1-aminoethyl)-5-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 391 (M+1)⁺.

50 ¹H NMR (400 MHz, CDCl₃) δ 10.73 (s, 1H), 8.33 (s, 1H), 7.98 (br s, 1H), 7.57 - 7.41 (m, 5H), 7.38 - 7.31 (m, 1H), 7.14 (dd, 1H), 6.52 (m, 1H), 5.18 (s, 1H), 1.50 (d, 3H).

EXAMPLE 84

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

55 **[0620]** The title compound (17 mg, 35% yield) was obtained from (S)-2-(1-aminoethyl)-5-fluoro-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 391 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 10.73 (s, 1H), 8.33 (s, 1H), 7.98 (br s, 1H), 7.57 - 7.41 (m, 5H), 7.38 - 7.31 (m, 1H), 7.14 (dd, 1H), 6.52 (m, 1H), 5.18 (s, 1H), 1.50 (d, 3H).

5 EXAMPLE 85

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

10 **[0621]** The title compound (84 mg, 54% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 434 (M+1)⁺.

15 ¹H NMR (400 MHz, DMSO) δ 7.96 (d, J = 3.0 Hz, 1H), 7.80 (s, 1H), 7.57 (d, J = 7.3 Hz, 1H), 7.47 (d, J = 8.5 Hz, 1H), 7.37 (m, 2H), 7.31 - 7.18 (m, 2H), 7.05 (d, J = 8.8 Hz, 1H), 5.10 (p, J = 6.5 Hz, 1H), 1.40 (d, J = 6.6 Hz, 3H)

EXAMPLE 86

20 **(S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroimidazo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

25 **[0622]** 75 mg (0.26 mmol) of (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)imidazo[1,2-*f*][1,2,4]triazin-4(3*H*)-one, 52 mg (0.33 mmol) of 4-amino-6-chloropyrimidine-5-carbonitrile and 134 μL (0.77 mmol) of diisopropylethylamine were heated at 120 °C in *tert*-butanol (3 mL) for 48 hours. The solvent was removed in vacuum and the residue was dissolved in dichloromethane. The organic solution was washed with water and brine, dried and the solvent was removed under reduced pressure. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 30 mg (28% yield) of the title compound.

LRMS (m/z): 410 (M+1)⁺.

30 ¹H NMR (400 MHz, DMSO) δ 8.19 (d, 1H), 8.19 (d, 1H), 7.80 (s, 1H), 7.63 (d, 1H), 7.57 (d, 1H), 7.48 (dd, 1H), 7.30 (s, 2H), 7.08 - 7.00 (m, 1H), 5.09 (m, 1H), 1.43 (d, 3H).

EXAMPLE 87

35 **(S)-2-(1-(9*H*-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

40 **[0623]** The title compound (31 mg, 20% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 6-bromo-9*H*-purine following the experimental procedure described in example 20.

LRMS (m/z): 434 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.96 (br s, 1H), 8.14 (m, 1H), 8.01 (m, 2H), 7.92 (d, J = 2.9 Hz, 1H), 7.50 (m, 1H), 7.23 (d, J = 3.0 Hz, 1H), 7.13 (t, J = 9.0 Hz, 1H), 7.04 (d, J = 8.9 Hz, 1H), 5.20 - 4.97 (m, 1H), 1.50 (d, J = 6.1 Hz, 3H).

45 EXAMPLE 88

4-Amino-6-((1*S*)-1-(5-(1,2-dihydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

50 **[0624]** The title compound (14 mg, 10% yield) was obtained from 2-((S)-1-aminoethyl)-5-(1,2-dihydroxyethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using *n*-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 433 (M+1)⁺.

55 ¹H NMR (400 MHz, DMSO) δ 7.78 (d, 1H), 7.73 - 7.67 (m, 1H), 7.61 - 7.56 (m, 1H), 7.52 - 7.16 (m, 7H), 6.66 - 6.57 (m, 1H), 5.18 - 5.04 (m, 2H), 4.93 - 4.81 (m, 1H), 4.69 - 4.60 (m, 1H), 1.34 (d, 3H).

EXAMPLE 89

(S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0625] The title compound (60 mg, 68% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 477 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.91 (d, J = 2.9 Hz, 1 H), 7.80 (s, 1 H), 7.57 (d, J = 7.3 Hz, 1 H), 7.49 (d, J = 9.2 Hz, 1 H), 7.30 (br s, 2H), 7.27 - 7.21 (m, 1 H), 7.06 (d, J = 3.0 Hz, 1H), 7.04 (d, J = 11.7 Hz, 1H), 5.10 (p, J = 6.5 Hz, 1H), 1.41 (d, J = 6.6 Hz, 3H).

EXAMPLE 90

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0626] The title compound (58 mg, 65% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20.

LRMS (m/z): 477 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.95 (br s, 1H), 8.15 (m, 1H), 8.02 (m, 2H), 7.87 (d, J = 2.8 Hz, 1H), 7.51 (d, J = 9.0 Hz, 1H), 7.11 (t, J = 8.9 Hz, 1H), 7.07 - 6.99 (m, 2H), 5.25 - 4.96 (m, 1H), 1.50 (d, J = 6.2 Hz, 3H).

EXAMPLE 91

(S)-4-Amino-6-(1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0627] The title compound (5 mg, 44% yield) was obtained from (S)-2-(1-aminoethyl)-5-(hydroxymethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 403 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.08 (s, 1 H), 7.64 - 7.53 (m, 3H), 7.51 - 7.45 (m, 1 H), 7.38 - 7.30 (m, 2H), 6.48 (d, 1 H), 5.76 (d, 1 H), 5.50 (br s, 2H), 5.11 - 4.98 (m, 1H), 4.81 (s, 2H), 1.42 (d, 3H).

EXAMPLE 92

(S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0628] The title compound (17 mg, 16% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (preparation 22) and 6-chloro-5-(trifluoromethyl)pyrimidine-4-amine (prepared as described at WO2005047279) following the experimental procedure described in example 26.

LRMS (m/z): 416 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.79 (s, 1 H), 7.68 - 7.62 (m, 1 H), 7.53 - 7.44 (m, 2H), 7.41 - 7.30 (m, 3H), 6.94 (dd, J = 4.2, 1.6 Hz, 1 H), 6.74 (br s, 2H), 6.65 (dd, J = 6.8, 1.7 Hz, 1 H), 6.60 (dd, J = 4.2, 2.7 Hz, 1 H), 5.02 - 4.85 (m, 1 H), 1.35 (d, J = 6.6 Hz, 3H).

EXAMPLE 93

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0629] To a solution of (S)-2-(1-aminoethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (39mg, 0.06mmols) in butan-1-ol (1.5ml), 4-amino-6-chloropyrimidine-5-carbonitrile (9mg, 0.06mmols) and DIEA (73 μl, 0.42mmols) were added. It was stirred at 120°C overnight. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound was obtained (3mg, 6%).

LRMS (m/z): 413 (M+1)+

EXAMPLE 94

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0630] The title compound (95 mg, 64% yield) was obtained from (S)-2-(1-aminoethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 459 (M+1)+.

¹H NMR (400 MHz, DMSO) δ 12.94 (s, 1 H), 8.15 (s, 1 H), 8.09 - 7.95 (m, 2H), 7.83 (d, 1H), 7.50 (d, 1H), 7.32 (t, 2H), 7.16 - 6.96 (m, 2H), 6.91 (d, 1H), 5.18 - 4.99 (m, 1 H), 1.50 (d, 3H).

EXAMPLE 95

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)imidazo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0631] The title compound was prepared following the experimental procedure described in Example 86 from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)imidazo[1,2-*f*][1,2,4]triazin-4(3H)-one (67 mg, 0.23 mmol) and 6-bromo-9H-purine (100 mg, 0.50 mmol). The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 28 mg (30% yield) of the title compound.

LRMS (m/z): 410 (M+1)+.

¹H NMR (400 MHz, DMSO) δ 12.91 (s, 1H), 8.36 - 6.86 (m, 7H), 5.07 (s, 1H), 1.51 (d, 3H).

EXAMPLE 96

(S)-4-Amino-6-(1-(5-(difluoromethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0632] The title compound (111 mg, 74% yield) was obtained from (S)-2-(1-aminoethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 459 (M+1)+.

¹H NMR (400 MHz, DMSO) δ 7.86 (d, 1H), 7.79 (s, 1H), 7.59 (d, 1H), 7.54 - 7.45 (m, 1 H), 7.39 - 7.14 (m, 4H), 7.03 (d, 1 H), 6.94 (d, 1 H), 5.16 - 5.03 (m, 1 H), 1.39 (d, 3H).

EXAMPLE 97

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0633] The title compound (44 mg, 28% yield) was obtained from (S)-2-(1-aminoethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 6-chloro-9H-purin-2-amine following the experimental procedure described in example 21 but heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 474 (M+1)+.

¹H NMR (400 MHz, DMSO) δ 12.07 (s, 1H), 8.17 (s, 1H), 7.80 (d, 1H), 7.68 (s, 1H), 7.57 - 7.41 (m, 2H), 7.32 (t, 1H), 7.17 (m, 1H), 6.90 (d, 1H), 5.59 (br s, 2H), 4.99 - 4.81 (m, 1 H), 1.44 (d, 3H).

EXAMPLE 98

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0634] The title compound (45 mg, 35% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 6-chloro-9H-purin-2-amine following the experimental procedure described in example 21.

LRMS (m/z): 449 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.07 (br s, 1H), 7.87 (d, J = 2.9 Hz, 1H), 7.67 (s, 1H), 7.46 (d, J = 7.7 Hz, 2H), 7.23 - 7.16 (m, 3H), 5.60 (bs, 2H), 4.88 (m, 1H), 1.44 (d, J = 6.6 Hz, 3H).

5 EXAMPLE 99

2-(1-(9H-Purin-6-ylamino)-2,2,2-trifluoroethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

10 **[0635]** 3-Phenyl-2-(2,2,2-trifluoro-1-(9-(4-methoxybenzyl)-9H-purin-6-ylamino)ethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (138 mg, 0.25 mmol) were dissolved in trifluoroacetic acid (2 mL). The solution was stirred at 60 °C overnight and at 80 °C for 4 hours. The volatiles were removed under reduced pressure and the residue was partitioned between ethyl acetate and a diluted aqueous solution of potassium carbonate. The organic layer was washed with water and brine, dried over magnesium sulphate, filtered and the solvent was removed in vacuum. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 17 mg (16% yield) of the title compound.

15 LRMS (m/z): 427 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 13.01 (s, 1H), 8.51 (s, 1H), 8.23 (s, 1H), 7.92 - 7.73 (m, 2H), 7.56 (s, 1H), 7.41 (s, 1H), 7.14 - 6.83 (m, 3H), 6.69 (dd, 1H), 6.53 (s, 1H), 6.07 (s, 1H).

20 EXAMPLE 100

(S)-4-Amino-6-(1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

25 **[0636]** The title compound was prepared following the experimental procedure described in Example 68 from (S)-2-(1-aminoethyl)-3-benzylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (obtained as a by-product of Preparation 22) (30 mg, 0.11 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (34 mg, 0.22 mmol). The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 19 mg (44% yield) of the title compound as a white solid.

30 LRMS (m/z): 387 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 11.49 (s, 1H), 8.04 (s, 1H), 7.45 (d, 1H), 7.38 (d, 3H), 7.26 (s, 2H), 7.25 (d, 2H), 7.15 (m, 1H), 6.32 (d, 1H), 5.13 (p, 1H), 4.20 (s, 2H), 1.52 (d, 3H).

EXAMPLE 101

35

(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0637] The title compound (20 mg, 37% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 5,6-difluoropyrimidin-4-amine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 24 hours.

40 LRMS (m/z): 366 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.70 - 7.29 (m, 7H), 7.18 (d, 1H), 6.94 (dd, 1H), 6.59 (dd, 1H), 6.36 (br s, 2H), 4.71 - 4.55 (m, 1H), 1.36 (d, 3H).

45 EXAMPLE 102

(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

50 **[0638]** The title compound (11 mg, 33% yield) was obtained from (S)-2-(1-aminoethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 5,6-difluoropyrimidin-4-amine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 6 hours.

LRMS (m/z): 452 (M+1)⁺.

55 ¹H NMR (400 MHz, DMSO) δ 7.86 (d, 1H), 7.37 (d, 1H), 7.26 (t, 2H), 7.23 - 7.16 (m, 1H), 7.06 - 6.97 (m, 1H), 6.95 - 6.88 (m, 1H), 6.86 - 6.81 (m, 1H), 5.00 - 4.90 (m, 1H), 4.74 (br s, 2H), 1.48 (d, 3H).

EXAMPLE 103**(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)propyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0639] The title compound (84 mg, 54% yield) was obtained from (S)-2-(1-aminopropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 448 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.93 (d, J = 2.8 Hz, 1 H), 7.78 (s, 1 H), 7.47 (d, J = 7.6 Hz, 1 H), 7.37 - 7.26 (m, 2H), 7.25 (d, J = 2.8 Hz, 1 H), 7.05 (d, J = 8.5 Hz, 1H), 4.91 (m, 1 H), 2.07 - 1.89 (m, 1 H), 1.80 (m, 1H), 0.84 (t, J = 7.1 Hz, 3H).

EXAMPLE 104**(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0640] The title compound was prepared following the experimental procedure described in Example 68 from (S)-2-(1-aminoethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (100 mg, 0.29 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (54 mg, 0.35 mmol). The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 91 mg (68% yield) of the title compound as a white solid.

LRMS (m/z): 467 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.97 (d, 1 H), 7.83 - 7.69 (m, 2H), 7.56 - 7.48 (m, 2H), 7.39 - 7.19 (m, 3H), 5.23 - 5.02 (m, 2H), 1.39 (d, 3H).

EXAMPLE 105**(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0641] The title compound (49 mg, 69% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 5,6-difluoropyrimidin-4-amine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 72 hours.

LRMS (m/z): 427 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.92 (s, 1 H), 7.59 - 7.40 (m, 2H), 7.34 - 7.02 (m, 4H), 6.43 (br s, 2H), 4.92 - 4.74 (m, 1 H), 1.41 (d, 3H).

EXAMPLE 106**(S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0642] The title compound (12 mg, 16% yield) was obtained from (S)-2-(1-aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (preparation 47) and 6-chloro-5-(trifluoromethyl)pyrimidine-4-amine (prepared as described at WO2005047279) following the experimental procedure described in example 26.

LRMS (m/z): 441 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.85 (d, J = 2.9 Hz, 1 H), 7.78 (s, 1 H), 7.55 (d, J = 7.8 Hz, 1 H), 7.49 (t, J = 7.0 Hz, 1 H), 7.36 (m, 3H), 7.20 (d, J = 2.9 Hz, 1 H), 6.76 (br s, 2H), 6.62 (d, J = 5.2 Hz, 1 H), 5.03 - 4.89 (m, 1 H), 1.36 (d, J = 6.5 Hz, 3H).

EXAMPLE 107**(R)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0643] The title compound was prepared following the experimental procedure described in Example 86 from (R)-2-(1-amino-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (71 mg, 0.21 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (40 mg, 0.26 mmol). The product was purified by flash chroma-

tography (0% to 10% MeOH/DCM) to give 32 mg (33% yield) of the title compound as a white solid.

LRMS (m/z): 450 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.92 (d, 1H), 7.83 (s, 1H), 7.66 - 7.51 (m, 1H), 7.46 - 7.28 (m, 4H), 7.24 (d, 1H), 7.15 (d, 1H), 5.04 - 4.90 (m, 2H), 3.94 - 3.80 (m, 1H), 3.70 - 3.54 (m, 1H).

EXAMPLE 108

(S)-2-(1-(6-Amino-5-carbamoylpyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide

[0644] A suspension of 30 mg (0.07 mmol) of (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile (Example 85) in sulphuric acid (2 ml) was heated to 60°C for 2h. The reaction mixture was slowly poured into a mixture of ice/water, neutralized with a 2N solution of NaHCO₃ and extracted with ethyl acetate. The organic layer was then washed with brine, dried over magnesium sulphate and concentrated. The title compound was obtained as a white solid (29mg, 88% yield).

LRMS (m/z): 470 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 9.16 (d, J = 2.2 Hz, 1H), 7.78 (d, J = 2.8 Hz, 1H), 7.77 (s, 1H), 7.62 (d, J = 7.5 Hz, 1H), 7.54 (d, J = 8.7 Hz, 1H), 7.46 (d, J = 2.2 Hz, 1H), 7.43 - 7.36 (m, 3H), 7.33 (d, J = 8.8 Hz, 1H), 7.09 (d, J = 2.9 Hz, 1H), 6.65 (s, 2H), 4.95 - 4.76 (m, 1H), 1.38 (d, J = 6.7 Hz, 3H).

EXAMPLE 109

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide

[0645] The title compound (25 mg, 49% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

LRMS (m/z): 452 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 9.16 (d, J = 2.4 Hz, 1H), 7.87 (d, J = 2.9 Hz, 1H), 7.79 (s, 1H), 7.60 (d, J = 7.4 Hz, 1H), 7.55 - 7.45 (m, 2H), 7.30 (br s, 2H), 7.25 (m, 1H), 7.13 (d, J = 2.8 Hz, 1H), 7.06 (d, J = 8.6 Hz, 1H), 5.10 (p, J = 6.6 Hz, 1H), 1.41 (d, J = 6.5 Hz, 3H).

EXAMPLE 110

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile

[0646] The title compound (6 mg, 36% yield) was obtained from (S)-2-(1-aminoethyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 446 (M+1)⁺.

EXAMPLE 111

2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile - first diastereomer

[0647] Same procedure as described in Example 93 was used from 2-((S)-1-aminoethyl)-4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile (118mg, 0.23mmols). The title compound (first eluting diastereomer) was obtained (64mg, 68%)

LRMS (m/z): 406 (M+1)⁺

¹H NMR (400 MHz, DMSO-d₆) δ ppm 0.88 - 1.03 (m, 1H) 1.46 (d, 3H) 1.52 - 1.64 (m, 1H) 1.67 - 1.86 (m, 1H) 2.55 - 2.78 (m, 1H) 2.57 - 2.75 (m, 1H) 3.09 - 3.30 (m, 1H) 3.60 - 3.78 (m, 1H) 3.87 (d, 1H) 4.10 - 4.31 (m, 1H) 5.64 - 5.92 (m, 1H) 7.05 - 7.22 (m, 1H) 7.40 (m, 1H) 7.80 (d, 1H) 8.00 (m, 1H) 8.17 (s, 1H) 8.40 (s, 1H)

EXAMPLE 112

(*R*)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile

[0648] Prepared from (*R*)-2-(1-amino-2-hydroxyethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (61 mg, 0.16 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (30 mg, 0.19 mmol) following the experimental procedure described in Example 86, but heating at 120 °C only for 16 hours. The product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1% v/v formic acid buffered] 0% to 100%) to furnish 8 mg (10% yield) of the title compound as a white solid.

LRMS (m/z): 493 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.87 (d, 1H), 7.84 (s, 1H), 7.48 - 7.25 (m, 5H), 7.15 (d, 1H), 7.04 (d, 1H), 5.05 - 4.84 (m, 2H), 4.00 - 3.79 (m, 2H), 3.72 - 3.52 (m, 1H).

EXAMPLE 113

(*S*)-2-(1-(2-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0649] The title compound (22 mg, 33% yield) was obtained from (*S*)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-chloro-5-fluoropyrimidin-2-amine (purchased from Ark Pharm) following the experimental procedure described in example 20.

LRMS (m/z): 427 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.91 (d, J = 3.0 Hz, 1H), 7.64 (d, J = 3.7 Hz, 1H), 7.48 (d, J = 9.0 Hz, 1H), 7.42 (d, J = 6.8 Hz, 1H), 7.29 (tt, J = 9.3, 2.3 Hz, 1H), 7.22 (m, 2H), 5.87 (s, 2H), 4.76 (p, J = 6.6 Hz, 1H), 1.39 (d, J = 6.7 Hz, 3H).

EXAMPLE 114

(*S*)-2-(1-(2-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0650] The title compound (34 mg, 50% yield) was obtained from (*S*)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 2-amino-4-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 20.

LRMS (m/z): 427 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.12 (s, 1H), 7.96 (d, J = 3.0 Hz, 1H), 7.55 (d, J = 7.2 Hz, 1H), 7.46 (d, J = 9.0 Hz, 1H), 7.26 (d, J = 3.0 Hz, 1H), 7.25-7.18 (m, 1H), 7.05 (m, 2H), 6.97 (br s, 1H), 5.05 (p, J = 6.5 Hz, 1H), 1.39 (d, J = 6.6 Hz, 3H).

EXAMPLE 115

((*S*)-2-(1-(9*H*-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(2*H*-tetrazol-5-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0651] 40 mg (0.1 mmol, Example 87) of (*S*)-2-(1-(9*H*-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 31 mg (0.15 mmol) of azidotrimethylstannane were stirred in toluene (2ml) at 100 °C for 3 days in a sealed reactor. Then the solvent was removed in vacuum and the residue was taken up in AcOEt, washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by normal phase chromatography (DCM-AcOEt, 0-100% in 40CV) to obtain the title compound (19 mg, 43% yield) as a white solid.

LRMS (m/z): 477(M+1)⁺.

¹H NMR (400 MHz, DMSO) δ ppm 1.53 (d, J=6.64Hz, 3H), 5.13 (q, 1H), 6.98-7.15 (m, 2H), 7.20 (s, 1H), 7.54 (d, 1H), 7.89-8.27 (m, 4H), 12.94 (s, 1H)

EXAMPLE 116

(S)-4-Amino-6-(1-(3-((5-methylisoxazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0652] The title compound (120 mg, 34% yield) was obtained from (S)-2-(1-aminoethyl)-3-((5-methylisoxazol-3-yl)methyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (preparation 59) and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (*m/z*): 392 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.15 (s, 1H), 7.35 (dd, 1H), 7.06 (dd, 1H), 6.54 (dd, 1H), 6.04 (d, 1H), 5.82 (d, 1H), 5.73 (m, 1H), 5.52 (d, 1H), 5.47 - 5.33 (m, 3H), 2.39 (d, 3H), 1.53 (d, 3H).

EXAMPLE 117

(S)-4-Amino-6-(1-(4-oxo-3-phenyl-7-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0653] Prepared following the experimental procedure described in Example 112 from 14.5 mg (0.04 mmol) of (S)-2-(1-aminoethyl)-3-phenyl-7-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one hydrochloric acid salt and 7 mg (0.05 mmol) of 4-amino-6-chloropyrimidine-5-carbonitrile. The product was purified first by flash chromatography (0% to 10% MeOH/DCM) and then preparative HPLC (Waters XBridge C18 OBD column, mixture of eluents A/B from 80% B to 100% B, in a 10 min. gradient) to give 10 mg (56% yield) of the title compound as a white solid.

LRMS (*m/z*): 441 (*M*+1)⁺.

EXAMPLE 118

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-7-carbonitrile

[0654] Prepared following the experimental procedure described in Example 112 from (S)-2-(1-aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-7-carbonitrile (27 mg, 0.10 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (33 mg, 0.21 mmol). The product was purified first by preparative HPLC (Waters SymmetryPrep C18 column, mixture of eluents A/B from 45% B to 75% B, in a 20 min. gradient) to give 15 mg (39% yield) of the title compound as a white solid.

LRMS (*m/z*): 398 (*M*+1)⁺.

EXAMPLE 119

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(1-(4-methoxybenzyl)-1*H*-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0655] The title compound (9 mg, 10% yield) was obtained from (S)-2-(1-aminoethyl)-3-(1-(4-methoxybenzyl)-1*H*-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (*m/z*): 508 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.00 (s, 1H), 7.69 (d, *J* = 0.7 Hz, 1H), 7.52 (d, *J* = 0.7 Hz, 1H), 7.31 (d, *J* = 2.9 Hz, 1H), 7.26 - 7.21 (m, 1H), 6.94 - 6.87 (m, 2H), 6.84 (d, *J* = 2.9 Hz, 1H), 5.68 (d, *J* = 8.0 Hz, 1H), 5.39 (s, 1H), 5.31 (s, 1H), 5.26 (dq, *J* = 13.8, 7.0 Hz, 1H), 3.81 (d, *J* = 3.0 Hz, 3H), 1.58 (s, 4H), 1.43 (d, *J* = 6.9 Hz, 2H).

EXAMPLE 120

(S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0656] Starting from (S)-2-(1-aminoethyl)-3-phenyl-5-(thiazol-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (26 mg, 0.08 mmol) and following the experimental procedure described in Example 23 were obtained 12 mg (34% yield) of the title compound of this example.

LRMS (*m/z*): 491 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ ppm 1.44 (d, *J*=7.03 Hz, 3 H) 5.04 (q, 1 H) 5.35 (s, 2 H) 5.73 (d, *J*=7.82 Hz, 1 H) 7.31 (d, *J*=3.13 Hz, 1 H) 7.33 - 7.39 (m, *J*=2.74 Hz, 2 H) 7.43 (d, *J*=2.74 Hz, 1 H) 7.47 - 7.63 (m, 4 H) 7.77 - 7.89 (m, *J*=3.13 Hz, 1 H) 8.11 (s, 1 H)

5 EXAMPLE 121

(S)-2-(1-(2,6-Diamino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

10 **[0657]** The title compound (20 mg, 29% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 2,4-diamino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 20.

LRMS (*m/z*): 449(*M*+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.94 (dd, *J* = 2.9, 1.7 Hz, 1 H), 7.44 (d, *J* = 9.1 Hz, 1 H), 7.30 - 7.19 (m, 2H), 7.10 (d, *J* = 9.1 Hz, 1 H), 6.93 (d, *J* = 6.7 Hz, 1 H), 6.53 (s, 2H), 6.28 (s, 2H), 5.05 - 4.89 (m, 1 H), 1.36 (d, *J* = 6.4 Hz, 3H).

EXAMPLE 122

20 **(S)-4-Amino-6-(1-(5-(morpholinomethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0658] The formate salt of the title compound (63 mg, 53% yield) was obtained from (S)-2-(1-aminoethyl)-5-(morpholinomethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23.

25 LRMS (*m/z*): 472 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.21 (s, 1H), 7.77 (s, 1H), 7.69 (d, *J* = 7.0 Hz, 1 H), 7.62 (s, 1 H), 7.49 (d, *J* = 8.1 Hz, 1 H), 7.42 (t, *J* = 7.1 Hz, 1 H), 7.28 (m, 4H), 6.56 (s, 1 H), 4.96 - 4.79 (m, 1 H), 3.78 (s, 2H), 3.55 (s, 4H), 2.39 (s, 4H), 1.36 (d, *J* = 6.4 Hz, 3H).

30 EXAMPLE 123

2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

35 **[0659]** The title compound (27 mg, 9% yield) was obtained from 2-((S)-1-aminoethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (*m/z*): 426 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.02 (s, 1H), 7.33 (s, 1H), 7.18 (s, 1H), 6.87 (s, 1 H), 5.62 (d, *J* = 26.3 Hz, 2H), 5.36 (s, 2H), 2.01 (d, *J* = 7.0 Hz, 3H), 1.60 (d, *J* = 6.2 Hz, 7H).

EXAMPLE 124

45 **(S)-4-Amino-6-(1-(4-oxo-3-(1*H*-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0660] The title compound (60 mg, 28% yield) was obtained from (S)-4-amino-6-(1-(3-(1-(4-methoxybenzyl)-1*H*-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (290 mg, 0.6 mmol) by heating with TFA (1.40 ml, 18.17 mmol) in dichloromethane (15 ml) and further purification.

50 LRMS (*m/z*): 363 (*M*+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 13.02 (br s, 1H), 7.95 (s, 1H), 7.90 (bs, 1H), 7.73 (d, 1 H), 7.65 (d, 1 H), 7.50 (bs, 1 H), 7.30 (bs, 2H), 6.95 (d, 1 H), 6.65 (d, 1 H), 5.10 (m, 1H), 1.40 (d, 3H).

55

EXAMPLE 125

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyrrolo[1,2-*f*]triazin-4(3H)-one

[0661]

a) 45 mg (0.1 mmol) of (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*]triazine-5-carbonitrile (Example 87), 14 mg (0.21 mmol) of hydroxylamine hydrochloride and 28 mg (0.21 mmol) of potassium carbonate were stirred in ethanol (2ml) at 80 °C for 48h. Then the precipitate is filtrated, washed with ethanol and diethylether obtaining (S,Z)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-N'-hydroxy-4-oxo-3,4-dihydropyrrolo[1,2-*f*]triazine-5-carboximidamide (70 mg, >100% yield) as a white solid.
LRMS (m/z): 467 (M+1)⁺.

b) 70 mg (0.1 mmol) of (S,Z)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-N'-hydroxy-4-oxo-3,4-dihydropyrrolo[1,2-*f*]triazine-5-carboximidamide in acetic acid (0.5 ml) was cooled to 0°C in an under pressure reactor. Then 507 µl of acetic anhydride (5.37 mmol) were added and the crude was heated at 90°C for 2h to obtain the acetylated title compound. The solvent was removed and methanol (2ml) and HCl 4M in dioxane (2 ml) is added and the resulting mixture is heated at 70°C for 5h. The solvent was removed in vacuum and the residue was purified directly by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%) obtaining (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyrrolo[1,2-*f*]triazin-4(3H)-one (10 mg, 13% yield).
LRMS (m/z): 491 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ ppm 1.25 (s, 3 H) 1.59 (d, *J*=6.25 Hz, 3 H) 5.06 - 5.35 (m, 1 H) 6.44 (s, 1 H) 6.88 - 6.99 (m, 2 H) 7.04 (s, 1 H) 7.10 - 7.25 (m, 1 H) 7.28 - 7.44 (m, 1 H) 8.01 (s, 1 H) 8.36 (s, 1 H) 12.22 - 12.76 (m, 1 H)

EXAMPLE 126

4-Amino-6-((1S)-1-(4-oxo-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile - first diastereomer

[0662] Same procedure as describes in Example 93 was used from 2-((S)-1-aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)pyrrolo[1,2-*f*]triazin-4(3H)-one (158mg, 0.20mmols). The title compound (first eluting diastereomer) was obtained (36mg, 41%)

LRMS (m/z): 449 (M+1)⁺ ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.82 (d, 1 H) 1.45 (d, 3 H) 1.50 - 1.64 (m, 1 H) 1.66 - 1.81 (m, 1 H) 2.60 - 2.77 (m, 1 H) 3.02 - 3.23 (m, 2 H) 3.60 - 3.75 (m, 1 H) 3.76 - 3.94 (m, 1 H) 4.11 - 4.30 (m, 1 H) 5.69 - 5.90 (m, 1 H) 6.95 (d, 1 H) 7.42 (s, 1 H) 7.74 (d, 1 H) 8.03 (s, 1 H) 8.12 - 8.23 (m, 1 H) 8.37 (s, 1 H)

EXAMPLE 127

(S)-4-Amino-6-(1-(3-(5-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0663] The title compound (9 mg, 12% yield) was obtained from (S)-2-(1-aminoethyl)-3-(5-methyl-1H-pyrazol-3-yl)pyrrolo[1,2-*f*]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 377 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.00 (s, 1H), 7.34 (dd, *J* = 2.6, 1.7 Hz, 1H), 7.04 (dd, *J* = 4.3, 1.7 Hz, 1 H), 6.49 (dd, *J* = 4.3, 2.7 Hz, 1 H), 6.09 (s, 1 H), 5.93 (d, *J* = 8.2 Hz, 1 H), 5.52 (s, 2H), 5.19 (dq, *J* = 13.7, 6.8 Hz, 1 H), 3.58 (q, *J* = 7.1 Hz, 1 H), 2.30 (s, 3H), 1.41 (d, *J* = 6.8 Hz, 3H).

EXAMPLE 128

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxylic acid

a) (S)-2-(1-Aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxylic acid

[0664] This compound was prepared starting from (S)-2-(1-(*tert*-butoxycarbonylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxylic acid (26 mg, 0.07 mmol, Preparation 103) and following the experimental procedure described in Preparation 46c to afford 28 mg (99% yield) of the title compound as a dihydrochloride salt that was used in the next step without any further purification.
LRMS (m/z): 299 (M+1)⁺.

b) (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxylic acid

[0665] The formate salt of the title compound (21 mg, 67% yield) was obtained from (S)-2-(1-aminoethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carboxylic acid (Example 128a) and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 417 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.78 (s, 1 H), 7.65 (d, J = 6.8 Hz, 1 H), 7.61 (d, J = 8.6 Hz, 1 H), 7.50 (td, J = 7.7, 1.6 Hz, 1 H), 7.44 - 7.38 (m, 1 H), 7.38 - 7.30 (m, 2H), 7.27 (br s, 2H), 7.18 (d, J = 2.9 Hz, 1 H), 4.95 (p, J = 6.6 Hz, 1 H), 1.41 (d, J = 6.6 Hz, 3H).

EXAMPLE 129

2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(tetrahydro-2*H*-pyran-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile - second diastereomer

[0666] Same procedure as described in Example 93 was used from 2-((S)-1-aminoethyl)-4-oxo-3-(tetrahydro-2*H*-pyran-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (118mg, 0.23mmols). The title compound (second eluting diastereomer) was obtained (3mg, 3%)

LRMS (m/z): 406 (M+1)⁺

EXAMPLE 130

(S)-4-Amino-6-(1-(3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0667] The title compound (39 mg, 39% yield) was obtained from (S)-2-(1-Aminoethyl)-3-(5-fluoropyridin-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 100°C during 5 hours.

LRMS (m/z): 392 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.53 (d, 1 H), 8.42 (d, 1 H), 8.17 (d, 1 H), 7.82 - 7.57 (m, 3H), 7.26 (br s, 2H), 7.02 (dd, 1H), 6.71 - 6.61 (dd, 1H), 5.13 - 4.95 (m, 1 H), 1.39 (d, 3H).

EXAMPLE 131

(S)-4-Amino-6-(1-(4-oxo-3-(1*H*-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0668] The title compound (8 mg, 10% yield) was obtained from (S)-2-(1-aminoethyl)-3-(1*H*-pyrazol-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 363 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.04 (s, 1 H), 7.69 (d, J = 2.4 Hz, 1 H), 7.43 (dd, J = 2.7, 1.7 Hz, 1 H), 7.12 (dd, J = 4.4, 1.7 Hz, 1 H), 6.58 (dd, J = 4.4, 2.7 Hz, 1 H), 6.45 (d, J = 2.4 Hz, 1 H), 5.90 (d, J = 8.3 Hz, 1 H), 5.76 (s, 1 H), 5.23 (dq, J = 13.6, 6.8 Hz, 1 H), 3.56 - 3.42 (m, 1H), 2.62 (s, 1 H), 1.49 (d, J = 6.8 Hz, 3H).

EXAMPLE 132**(S)-4-Amino-6-(1-(4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0669] The title compound (4 mg, 14% yield) was obtained from (S)-2-(1-aminoethyl)-3-(pyrimidin-5-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.
LRMS (m/z): 375 (M+1)⁺.

EXAMPLE 133**4-Amino-6-((1S)-1-(4-oxo-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile - second diastereomer**

[0670] Same procedure as describes in Example 93 was used from 2-((S)-1-aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (158mg, 0.20mmols). The title compound (second eluting diastereomer) was obtained (10mg, 11%)

LRMS (m/z): 449 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.46 (d, 3 H) 1.55 - 1.77 (m, 1 H) 1.83 - 2.03 (m, 1 H) 2.33 (d, 1 H) 2.53 - 2.79 (m, 2 H) 3.01 - 3.25 (m, 1 H) 3.55 - 3.78 (m, 1 H) 3.80 - 4.02 (m, 1 H) 4.24 (t, 1 H) 5.58 - 5.87 (m, 1 H) 6.95 (d, 1 H) 7.37 (s, 2 H) 7.74 (d, 1 H) 7.95 (d, 1 H) 8.05 - 8.21 (m, 1 H)

EXAMPLE 134**(S)-2,4-Diamino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0671] The title compound (17 mg, 33% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (preparation 22) and 2,4-diamino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 20.

LRMS (m/z): 388 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.67 (dd, J = 2.6, 1.7 Hz, 1 H), 7.46 (m, 3H), 7.35 (m, 2H), 6.99 - 6.90 (m, 2H), 6.60 (dd, J = 4.3, 2.7 Hz, 1 H), 6.49 (br s, 2H), 6.20 (bs, 2H), 4.80 (p, J = 6.7 Hz, 1 H), 1.31 (d, J = 6.7 Hz, 3H).

EXAMPLE 135**(S)-4-(1-(3-((1H-Pyrazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)-6-aminopyrimidine-5-carbonitrile**

[0672] The title compound (8 mg, 10% yield) was obtained from (S)-3-((1H-pyrazol-3-yl)methyl)-2-(1-aminoethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 377 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.22 (s, 1 H), 7.51 (d, J = 2.2 Hz, 1 H), 7.34 (dt, J = 16.1, 8.1 Hz, 1H), 7.05 (dd, J = 4.3, 1.6 Hz, 1H), 6.54 (dd, J = 4.4, 2.6 Hz, 1H), 6.39 (d, J = 2.2 Hz, 1 H), 5.82 (dq, J = 13.4, 6.7 Hz, 1 H), 5.58 (s, 1 H), 5.49 (d, J = 15.8 Hz, 1 H), 5.34 (d, J = 15.8 Hz, 1 H), 3.83 - 3.74 (m, 2H), 1.52 (d, J = 6.6 Hz, 3H).

EXAMPLE 136**(S)-4-Amino-6-(1-(4-oxo-3-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0673] To a solution of (S)-2-(1-aminoethyl)-3-(tetrahydro-2H-pyran-4-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (100mg, 0.38mmols, Preparation 133) in butan-1-ol (4ml), 4-amino-6-chloropyrimidine-5-carbonitrile (61mg, 0.39mmols) and DIEA (200 μl, 1.14mmols) were added. It was stirred at 120°C for 8h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound was obtained (70mg, 48% yield).

LRMS (m/z): 381 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.38 - 1.44 (m, 1 H) 1.62 - 1.75 (m, 1 H) 2.62 - 2.84 (m, 2 H) 3.03 - 3.17 (m, 1 H) 3.23 - 3.30 (m, 1 H) 3.71 - 3.83 (m, 1 H) 3.83 - 3.95 (m, 2 H) 5.81 (q, 1 H) 6.49 - 6.63 (m, 1 H) 6.79 - 6.91 (m, 1 H) 7.31 - 7.54 (s, 2 H) 7.52 - 7.66 (m, 1 H) 8.08 - 8.24 (m, 2 H)

5 EXAMPLE 137

(S)-4-Amino-6-(1-(4-oxo-3-(2,2,2-trifluoroethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

10 **[0674]** To a solution of (S)-2-(1-aminoethyl)-3-(2,2,2-trifluoroethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (100mg, 0.38mmols, preparation 134) in butan-1-ol (4ml), 4-amino-6-chloropyrimidine-5-carbonitrile (60mg, 0.38mmols) and DIEA (200 μl, 1.14mmols) were added. It was stirred at 120°C for 8h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound was obtained (81 mg, 56% yield).

LRMS (m/z): 379 (M+1)⁺

15 ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.52 (d, 3 H) 4.45 - 4.65 (m, 1 H) 5.05 - 5.25 (m, 1 H) 5.53 (q, 1 H) 6.64 (dd, 1 H) 7.03 (dd, 1 H) 7.34 - 7.56 (m, 2 H) 7.67 - 7.82 (m, 1 H) 7.95 (d, 1 H) 8.11 (s, 1 H)

EXAMPLE 138

20 **(S)-4-Amino-6-(1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0675] To a solution of (S)-2-(1-aminoethyl)-3-cyclobutylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (107mg, 0.38mmols, preparation 135) in butan-1-ol (4ml), 4-amino-6-chloropyrimidine-5-carbonitrile (59mg, 0.38mmols) and DIEA (466 μl, 2.68mmols) were added. It was stirred at 120°C for 12h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound was obtained (70mg, 53% yield).

LRMS (m/z): 351 (M+1)⁺

25 ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.26 - 1.56 (m, 4 H) 1.66 - 1.95 (m, 2 H) 1.97 - 2.22 (m, 1 H) 2.82 - 3.05 (m, 1 H) 3.04 - 3.22 (m, 1 H) 4.28 - 4.64 (m, 1 H) 5.56 - 5.79 (m, 1 H) 6.54 (dd, 1 H) 6.86 (dd, 1 H) 7.39 (s, 2 H) 7.46 - 7.64 (m, 1 H) 7.99 (d, 1 H) 8.15 (s, 1 H)

EXAMPLE 139

35 **(S)-2-Amino-4-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0676] The title compound (29 mg, 35% yield) was obtained from (S)-2-(1-aminoethyl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (preparation 22) and 2-amino-4-chloropyrimidine-5-carbonitrile (preparation 97) following the experimental procedure described in example 26.

40 LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ ppm 8.06 (s, 1 H), 7.73 - 7.64 (m, 2H), 7.48 (d, *J* = 7.7 Hz, 1 H), 7.43 (m, 1 H), 7.34 (m, 1 H), 7.30 - 7.21 (m, 2H), 6.97 (m, 2H), 6.76 (br s, 1 H), 6.65 - 6.58 (m, 1 H), 5.03 - 4.76 (m, 1 H), 1.36 (d, *J* = 6.5 Hz, 3H).

EXAMPLE 140

45 **4-Amino-6-(1-(5-(1-methyl-1*H*-pyrazol-4-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0677] Starting from (S)-2-(1-aminoethyl)-5-(1-methyl-1*H*-pyrazol-4-yl)-3-phenylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (78 mg, 0.16 mmol) and following the experimental procedure described in Example 23 were obtained 27 mg (36% yield) of the title compound of this example.

LRMS (m/z): 453 (M+1)⁺.

50 ¹H NMR (400 MHz, CDCl₃) δ ppm 1.63 (s, 3H) 3.85 (s, 3 H) 4.94 - 5.10 (m, 1 H) 5.37 (s, 2 H) 5.78 (d, *J*=7.82 Hz, 1 H) 6.74 (d, *J*=2.74 Hz, 1 H) 7.34 (d, *J*=7.82 Hz, 1 H) 7.38 (d, *J*=2.74 Hz, 1 H) 7.42 - 7.49 (m, 1 H) 7.49 - 7.64 (m, 3 H) 7.85 (s, 1 H) 8.09 (s, 1 H) 8.35 (s, 1 H)

EXAMPLE 141**(S)-4-Amino-6-(1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0678] To a solution of (S)-2-(1-aminoethyl)-3-cyclopropylpyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (311mg, 1.28mmols, preparation 136) in butan-1-ol (11.5ml), 4-amino-6-chloropyrimidine-5-carbonitrile (198mg, 1.28mmols) and DIEA (1.5ml, 8.61mmols) were added. It was stirred at 120°C for 6h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound was obtained (246mg, 57% yield).

LRMS (m/z): 337 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.75 - 1.06 (m, 3 H) 1.07 - 1.22 (m, 1 H) 2.70 - 2.84 (m, 1 H) 5.63 - 6.06 (m, 1 H) 6.51 (dd, 1 H) 6.83 (dd, 1 H) 7.33 (s, 2 H) 7.45 - 7.58 (m, 1 H) 7.81 (d, 1 H) 8.07 (s, 1 H)

EXAMPLE 142**(S)-4-Amino-6-(1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile****a) (S)-2-(1-Aminoethyl)-5-bromopyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**

[0679] 837 mg (2.34 mmol) of (S)-*tert*-butyl 1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (obtained as a by-product in Preparation 45b) was stirred in a 4M solution of hydrochloric acid in dioxane at room temperature for 4 hours. The volatiles were removed in vacuum and the residue was partitioned between ethyl acetate and a diluted aqueous solution of potassium carbonate. The organic layers were washed with water and brine, dried over magnesium sulphate, filtered and the solvent was removed.

LRMS (m/z): 258 (M+1)⁺.

b) (S)-4-Amino-6-(1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0680] Prepared from (S)-2-(1-Aminoethyl)-5-bromopyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (570 mg, 2.22 mmol) and 4-amino-6-chloropyrimidine-5-carbonitrile (377 mg, 2.44 mmol) following the experimental procedure described in Example 68 at a temperature of 120 °C. The product was purified by flash chromatography (0% to 10% MeOH/DCM) to give 563 mg (68% yield) of the title compound as a white solid.

LRMS (m/z): 376 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 11.79 (s, 1H), 8.03 (s, 1H), 7.62 (d, 1H), 7.49 - 7.18 (m, 3H), 6.66 (d, 1 H), 5.17 - 5.02 (m, 1 H), 1.51 (d, 3H).

EXAMPLE 143**4-Amino-6-((1S)-1-(4-oxo-3-(tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile - first diastereomer**

[0681] To a solution of 2-((S)-1-aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one (100 mg, 0.38 mmols) in butan-1-ol (4 ml), 4-amino-6-chloropyrimidine-5-carbonitrile (60mg, 0.38mmols) and DIEA (200 μl, 1.14 mmols) were added. It was stirred at 120°C for 12h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound (first eluting diastereomer) was obtained (34mg, 23%).

LRMS (m/z): 381 (M+1)⁺.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.77 - 0.94 (m, 1 H) 1.06 - 1.20 (m, 1 H) 1.46 (d, 3 H) 1.52 - 1.62 (m, 1 H) 1.70 - 1.82 (m, 1 H) 2.66 (d, 1 H) 3.07 - 3.24 (m, 1H) 3.68 (d, 1 H) 3.71 - 3.91 (m, 1 H) 4.17 - 4.34 (m, 1 H) 5.65 - 5.86 (m, 1 H) 6.56 (dd, 1 H) 6.84 (dd, 1 H) 7.41 (s, 2 H) 7.52 - 7.64 (m, 1 H) 8.06 - 8.22 (m, 2 H)

EXAMPLE 144**(S)-4-Amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0682] Starting from (S)-(1-aminoethyl)-bromo-3-phenylpyrrolon ,2-*f*[1,2,4]triazin-4(3H)-one (213mg, 0,58 mmol) and

following the experimental procedure described in Example 23 were obtained 203 mg (78% yield) of the title compound of this example.

LRMS (m/z): 451, 453 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.77 (s, 1 H), 7.74 (d, J = 2.9 Hz, 1 H), 7.64 (d, J = 6.9 Hz, 1 H), 7.50 (m, 1 H), 7.44 (m, 1 H), 7.37 - 7.32 (m, 1 H), 7.32 - 7.27 (m, 2H), 7.27 - 7.17 (br s, 2H), 6.76 (d, J = 2.9 Hz, 1 H), 4.85 (p, J = 6.6 Hz, 1 H), 1.35 (d, J = 6.7 Hz, 3H).

EXAMPLE 145

2-((3-Iodo-1*H*-pyrazolo[3,4-*d*]pyrimidin-4-ylamino)methyl)-5-methyl-3-*o*-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0683] To a solution of 317 mg (1.10 mmol) of 2-(chloromethyl)-5-methyl-3-*o*-tolylpyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one in DMF (10 ml), 3-iodo-1*H*-pyrazolo[3,4-*d*]pyrimidin-4-amine (316 mg, 1.21 mmol) and potassium carbonate (167 mg, 1.21 mmol) were added. The solution was stirred at room temperature for 2 h and the solvent was removed under vacuum. The residue was partitioned between water and dichloromethane and the organic layer was washed with water and brine, dried over magnesium sulphate, filtered and the solvent was evaporated. The product was purified first by flash chromatography (0% to 10% MeOH/DCM) and then by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1 % v/v formic acid buffered] 0% to 100%) to obtain 30 mg (6% yield) of the title compound, which was the minor isomer of the reaction.

LRMS (m/z): 513 (M+1)⁺.

EXAMPLE 146

(*S*)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0684] The title compound (15 mg, 10% yield) was obtained from (*S*)-2-(1-aminoethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using *n*-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 417 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.59 (d, 1H), 8.50 (d, 1H), 8.16 (d, 1H), 7.76 (d, 1H), 7.71 - 7.49 (m, 2H), 7.45 - 7.12 (m, 3H), 5.16 - 4.93 (m, 1 H), 1.40 (d, 3H).

EXAMPLE 147

4-Amino-6-((1*S*)-1-(4-oxo-3-(tetrahydro-2*H*-pyran-3-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile - second diastereomer

[0685] To a solution of 2-((*S*)-1-aminoethyl)-3-(tetrahydro-2*H*-pyran-3-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one (100mg, 0.38mmols) in butan-1-ol (4ml), 4-amino-6-chloropyrimidine-5-carbonitrile (60mg, 0.38mmols) and DIEA (200 μl, 1.14mmols) were added. It was stirred at 120°C for 12h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound (second eluting diastereomer) was obtained (14mg, 10%).

LRMS (m/z): 381 (M+1)⁺.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.47 (d, 3 H) 1.60 - 1.79 (m, 1 H) 1.81 - 1.97 (m, 1 H) 2.66 (d, 1 H) 3.06 - 3.21 (m, 2 H) 3.56 - 3.72 (m, 2 H) 3.73 - 3.91 (m, 1 H) 4.17 - 4.34 (m, 1 H) 5.60 - 5.81 (m, 1 H) 6.56 (dd, 1 H) 6.84 (d, 1 H) 7.36 (s, 2 H) 7.60 (s, 1 H) 8.02 - 8.20 (m, 2 H)

EXAMPLE 148

(*S*)-4-Amino-6-(1-(4-oxo-3-phenyl-5-(1*H*-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0686] 40 mg (0.09 mmol) of (*S*)-4-amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile, 33 mg (0.17 mmol, Example 144) of 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole, 6 mg (0.01 mmol) of tetrakis(triphenylphosphine)palladium(0) and 136 μl of sodium carbonate 2M in water under argon were stirred in dimethylformamide (1ml) at 120°C overnight. The crude was filtered over Celite washing

with ethyl acetate. Then the organic phase was washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The crude product was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%) to obtain the title compound (10 mg, 27% yield) as a white solid.

LRMS (m/z): 439 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ ppm 1.41 (d, J=6.64 Hz, 3 H) 2.00 (s, 1 H) 4.95 - 5.08 (m, 1 H) 5.47 (s, 2 H) 5.83 (d, J=8.21 Hz, 1 H) 6.74 (d, J=2.74 Hz, 1 H) 7.33 (d, J=7.42 Hz, 1 H) 7.39 (d, J=2.74 Hz, 1 H) 7.42 - 7.48 (m, 1 H) 7.49 - 7.62 (m, 3 H) 8.07 (s, 1 H) 8.23 (s, 2 H)

EXAMPLE 149

(S)-4-Amino-6-(1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0687] To a solution of (S)-2-(1-aminoethyl)-3-(isoxazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (188 mg, 0.77 mmols) in butan-1-ol (7 ml), 4-amino-6-chloropyrimidine-5-carbonitrile (118 mg, 0.77 mmols) and DIEA (0.94 ml, 5.37 mmols) were added. It was stirred at 120°C for 8h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound was obtained (82mg, 29%).

LRMS (m/z): 364 (M+1)⁺

¹H NMR (400 MHz, DMSO-d₆) δ ppm 1.43 (d, 3 H) 4.98 - 5.24 (m, 1 H) 6.58 - 6.83 (m, 2 H) 7.07 (dd, 1 H) 7.26 (s, 2 H) 7.67 - 7.85 (m, 2 H) 7.93 (s, 1 H) 8.98 (d, 1 H)

EXAMPLE 150

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-N,N-dimethyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide

[0688] The title compound (24 mg, 72% yield) was obtained from (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylic acid (30mg, 0.07mmol, Example 128) and dimethylamine (2M solution in Methanol, 0.09mmol) following the experimental procedure described in Preparation 20a.

LRMS (m/z): 444 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.77 (s, 1 H), 7.71 (d, J = 2.6 Hz, 1 H), 7.64 (d, J = 7.0 Hz, 1 H), 7.49 (d, J = 7.8 Hz, 1 H), 7.43 (m, 1 H), 7.30 (m, 3H), 7.19 (br s, 2H), 6.63 (d, J = 2.6 Hz, 1 H), 5.02 - 4.83 (m, 1 H), 2.92 (s, 3H), 2.83 (s, 3H), 1.38 (d, J = 6.6 Hz, 3H).

EXAMPLE 151

(S)-4-Amino-6-(1-(3-(1-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0689] The title compound (50 mg, 33% yield) was obtained from (S)-2-(1-aminoethyl)-3-(1-methyl-1H-pyrazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 377 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.10 (s, 1H), 7.50 (t, 1H), 7.40 (dd, 1H), 7.09 (dd, 1 H), 6.54 (dd, 1 H), 6.36 (d, 1 H), 6.08 (d, 1 H), 5.40 (s, 2H), 5.25 (dq, 1 H), 3.95 (s, 3H), 1.49 - 1.46 (m, 3H).

EXAMPLE 152

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-N-propyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide

[0690] The title compound (24 mg, 72% yield) was obtained from (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylic acid (30mg, 0.07mmol, Example 128) and propan-1-amine (7μl, 0.09mmol) following the experimental procedure described in Example 150.

LRMS (m/z): 458 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 9.87 (t, J = 5.5 Hz, 1 H), 7.81 (d, J = 2.9 Hz, 1 H), 7.77 (s, 1 H), 7.63 (d, J = 6.8 Hz, 1 H), 7.60 - 7.55 (m, 1 H), 7.50 - 7.43 (m, 1 H), 7.42 - 7.38 (m, 1 H), 7.37 - 7.29 (m, 2H), 7.22 (br s, 2H), 7.12 (d, J = 2.9 Hz, 1H), 4.90 (p, J = 6.6 Hz, 1H), 3.21 (q, J = 6.5 Hz, 2H), 1.46 (q, J = 7.1 Hz, 2H), 1.39 (d, J = 6.7 Hz, 3H) 0.84 (t, J = 7.4 Hz, 3H).

EXAMPLE 153**2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one - first diastereomer**

[0691] To a solution of 2-((S)-1-aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (89 mg, 0.34 mmols) in *tert*-butanol (4 ml), 6-bromo-9H-purine (68 mg, 0.34 mmols) and DIEA (415 μ l, 2.38 mmols) were added. It was stirred at 80°C for 18h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound (first eluting diastereomer) was obtained (37mg, 29%).

LRMS (m/z): 381 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.92 - 1.06 (m, 1 H) 1.15 - 1.30 (m, 1 H) 1.57 (d, 3 H) 1.60 - 1.75 (m, 1 H) 2.96 - 3.15 (m, 2 H) 3.49 - 3.66 (m, 1 H) 3.75 - 3.91 (m, 1 H) 3.93 - 4.11 (m, 1 H) 4.23 (d, 1 H) 5.75 - 5.97 (m, 1 H) 6.41 - 6.62 (m, 1 H) 6.83 (dd, 1 H) 7.58 (s, 1 H) 8.14 (s, 1 H) 8.24 (s, 1 H) 8.28 - 8.49 (m, 1 H)

EXAMPLE 154**2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one - second diastereomer**

[0692] To a solution of 2-((S)-1-aminoethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (89 mg, 0.34 mmols) in *tert*-butanol (4 ml), 6-bromo-9H-purine (68 mg, 0.34 mmols) and DIEA (415 μ l, 2.38 mmols) were added. It was stirred at 80°C for 18h. It was concentrated in vacuum and it was purified by reverse phase chromatography. The title compound (second eluting diastereomer) was obtained (13mg, 10%).

LRMS (m/z): 381 (M+1)⁺

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.34 - 1.75 (m, 4 H) 1.81 - 1.99 (m, 1 H) 2.22 - 2.39 (m, 1 H) 2.58 - 2.79 (m, 1 H) 2.86 - 3.17 (m, 2 H) 3.47 - 3.68 (m, 1 H) 4.00 - 4.33 (m, 2 H) 5.65 - 5.89 (m, 1 H) 6.54 (dd, 1 H) 6.75-6.85 (m, 1 H) 7.57 (s, 1 H) 8.14 (s, 1 H) 8.20 - 8.42 (m, 2 H)

EXAMPLE 155**(S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile**

[0693] The title compound (92 mg, 61% yield) was obtained from (S)-2-(1-Amino-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 20 but using *n*-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 403 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.84 (s, 1 H), 7.74 - 7.57 (m, 2H), 7.53 - 7.32 (m, 5H), 7.24 (br s, 2H), 6.94 (dd, 1 H), 6.60 (dd, 1 H), 4.92 - 4.79 (m, 1 H), 4.53 (t, 1 H), 3.46 - 3.34 (m, 2H), 2.12 - 1.99 (m, 1 H), 1.98 - 1.87 (m, 1 H).

EXAMPLE 156**(S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one**

[0694] The title compound (28 mg, 18% yield) was obtained from (S)-2-(1-Amino-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using *n*-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 403 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.89 (s, 1 H), 8.22 - 8.04 (m, 2H), 7.96 (s, 1 H), 7.60 (dd, 1 H), 7.55 - 7.13 (m, 5H), 6.93 (dd, 1 H), 6.57 (dd, 1 H), 5.00 - 4.78 (m, 1H), 4.51 (t, 1 H), 3.49 - 3.34 (m, 2H), 2.19 - 2.06 (m, 1 H), 2.06 - 1.93 (m, 1 H).

EXAMPLE 157**(R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile**

[0695] The title compound (9.2mg) was obtained as a by-product during the synthesis of Example 112.

LRMS (m/z): 425 (M+1)⁺.

EXAMPLE 158**4-Amino-6-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile**

[0696] The title compound (34 mg, 45% yield) was obtained from 2-(aminomethyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (preparation 9) and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 373 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ 8.09 (s, 1H), 7.49 - 7.34 (m, 4H), 7.20 (d, 1H), 7.11 (dd, 1H), 6.66 - 6.53 (m, 1H), 5.95 (s, 1H), 5.34 (s, 2H), 4.23 (ddd, 2H), 2.21 (s, 3H).

EXAMPLE 159**(S)-4-Amino-6-(1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0697] Starting from (S)-2-(1-aminoethyl)-5-(2-hydroxyethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one (35 mg, 0.08 mmol) and following the experimental procedure described in Example 23 were obtained 12 mg (32% yield) of the title compound of this example.

LRMS (m/z): 417 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ ppm 1.40 (d, J=7.03 Hz, 3H) 3.16 (t, J=6.06 Hz, 2H) 3.87 (t, J=5.86 Hz, 2H) 4.97 - 5.12 (m, 1H) 5.38 (s, 2H) 5.77 (d, J=8.21 Hz, 1H) 6.44 (d, J=2.74 Hz, 1H) 7.29 - 7.36 (m, 2H) 7.44 (d, J=7.03 Hz, 1H) 7.48 - 7.64 (m, 3H) 8.08 (s, 1H)

EXAMPLE 160**(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)-3-hydroxypropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile**

[0698] The title compound (41 mg, 25% yield) was obtained from (S)-2-(1-amino-3-hydroxypropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 464 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.89 (d, 1H), 7.87 (s, 1H), 7.58 (d, 1H), 7.43 (d, 1H), 7.39 - 7.27 (m, 3H), 7.22 (d, 1H), 7.18 (d, 1H), 5.04 - 4.95 (m, 1H), 4.49 (t, 1H), 3.46 - 3.37 (m, 2H), 2.16 - 2.04 (m, 1H), 2.01 - 1.90 (m, 1H).

EXAMPLE 161**(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile**

[0699] The title compound (2 mg, 3% yield) was obtained from (S)-2-(1-aminoethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 417 (M+1)⁺.

EXAMPLE 162**(S)-4-Amino-6-(1-(5-(2-methyloxazol-5-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0700] The title compound (18 mg, 34% yield) was obtained from (S)-4-amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (52 mg, 0.12 mmol, Example 144) and 2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)oxazole (48mg, 0.23mmol) following the experimental procedure described in Example 148.

LRMS (m/z): 454 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ ppm 1.43 (d, J=6.64 Hz, 3H) 2.50 (s, 3H) 4.98 - 5.08 (m, 1H) 5.35 (s, 2H) 5.75 (d, J=8.21

Hz, 1 H) 6.86 (d, $J=3.13$ Hz, 1 H) 7.34 (d, $J=7.42$ Hz, 1 H) 7.39 (d, $J=3.13$ Hz, 1 H) 7.44 - 7.51 (m, 1 H) 7.52 - 7.64 (m, 3 H) 7.85 (s, 1 H) 8.09 (s, 1 H)

EXAMPLE 163

(S)-4-Amino-6-(1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2- f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0701] Starting from (S)-2-(1-aminoethyl)-5-(2-methoxyethyl)-3-phenylpyrrolo[1,2- f][1,2,4]triazin-4(3H)-one (40 mg, 0.10 mmol) and following the experimental procedure described in Example 23 were obtained 8mg (18% yield) of the title compound.

LRMS (m/z): 431 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ ppm 1.40 (d, $J=7.03$ Hz, 3 H) 3.19 (t, $J=6.64$ Hz, 2 H) 3.34 (s, 3 H) 3.66 (t, $J=6.64$ Hz, 2 H) 4.99 - 5.08 (m, 1 H) 5.40 (s, 2 H) 5.80 (d, $J=7.82$ Hz, 1 H) 6.46 (d, $J=2.74$ Hz, 1 H) 7.29 - 7.34 (m, 2 H) 7.38 - 7.45 (m, 1 H) 7.46 - 7.59 (m, 3 H) 8.07 (s, 1 H)

EXAMPLE 164

(S)-Propyl 2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2- f][1,2,4]triazine-5-carboxylate

[0702] To a solution of (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2- f][1,2,4]triazine-5-carboxylic acid (55mg, 0.08mmol, Example 128) in DMF (2ml) was added potassium carbonate (22mg, 0.16mmol) and 1-bromopropane (14.4 μ l, 0.16 mmol). The reaction mixture was stirred overnight at 50°C. The crude product was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%) to obtain the title compound (15 mg, 40% yield) as a white solid.

LRMS (m/z): 459 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃) δ ppm 0.90 - 1.03 (m, 3 H) 1.41 (d, $J=6.64$ Hz, 3 H) 1.71 - 1.80 (m, 2 H) 4.24 (t, $J=6.84$ Hz, 2 H) 4.95 - 5.09 (m, 1 H) 5.38 (s, 2 H) 5.72 (d, $J=7.82$ Hz, 1 H) 7.04 (d, $J=3.13$ Hz, 1 H) 7.29 - 7.37 (m, 2 H) 7.43 - 7.60 (m, 4 H) 8.08 (s, 1 H)

EXAMPLE 165

(S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2- f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile

[0703] The title compound (16 mg, 27% yield) was obtained from (S)-2-(1-amino-3-hydroxypropyl)pyrrolo[1,2- f][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 327 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 11.68 (s, 1 H), 8.88 (br s, 2H), 8.02 (s, 1 H), 7.58 (dd, 1H), 7.36 (d, 1H), 6.86 (dd, 1H), 6.52 (dd, 1 H), 5.26 - 5.17 (m, 1H), 4.77 (t, 1 H), 3.55 - 3.45 (m, 2H), 2.13 - 1.99 (m, 2H).

EXAMPLE 166

(S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)pyrrolo[1,2- f][1,2,4]triazin-4(3H)-one

[0704] The title compound (11 mg, 18% yield) was obtained from (S)-2-(1-amino-3-hydroxypropyl)pyrrolo[1,2- f][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 327 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.94 (s, 1H), 11.93 (s, 1H), 8.21 (s, 1H), 8.18 (s, 1 H), 7.72 (d, 1 H), 7.54 (dd, 1 H), 6.82 (dd, 1 H), 6.49 (dd, 1 H), 5.50 - 5.24 (m, 1H), 4.71 (t, 1 H), 3.63 - 3.46 (m, 2H), 2.19 - 2.04 (m, 2H).

EXAMPLE 167

(S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)-3-hydroxypropylamino)pyrimidine-5-carbonitrile

[0705] The title compound (131 mg, 33% yield) was obtained from (S)-2-(1-amino-3-hydroxypropyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 507 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.86 (s, 1H), 7.85 (d, 1H), 7.57 (d, 1H), 7.46 (d, 1H), 7.41 - 7.23 (m, 3H), 7.15 (d, 1H), 7.03 (d, 1H), 5.07 - 4.93 (m, 1H), 4.49 (t, 1H), 3.48 - 3.37 (m, 2H), 2.18 - 2.06 (m, 1H), 2.01 - 1.89 (m, 1H).

EXAMPLE 168

(S)-4-Amino-6-(1-(4-oxo-3-(6-(trifluoromethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0706] The title compound (21 mg, 16% yield) was obtained from (S)-2-(1-aminoethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 442 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.22 - 8.09 (m, 1H), 7.83 (d, 1H), 7.79 (dd, 1H), 7.73 - 7.60 (m, 3H), 7.20 (br s, 2H), 7.06 (dd, 1H), 6.67 (dd, 1H), 5.28 - 5.15 (m, 1H), 1.42 (d, 3H).

EXAMPLE 169

(S)-4-Amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

a) (S)-2-(1-Aminoethyl)-5-bromo-3-(3-(trifluoromethyl)phenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one

[0707] This compound was prepared starting from ((S)-*tert*-butyl 1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (200 mg, 0.40 mmol) and following the experimental procedure described in Preparation 46c to afford 230 mg (99% yield) of the title compound as a dihydrochloride salt that was used in the next step without any further purification.

LRMS (m/z): 401, 403 (M+1)⁺.

b) (S)-4-Amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0708] The title compound (21 mg, 67% yield) was obtained from (S)-2-(1-aminoethyl)-5-bromo-3-(3-(trifluoromethyl)phenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3*H*)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 519, 521 (M+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 8.05 (s, 0.5H), 7.87 (d, J = 7.6 Hz, 0.5H), 7.79 (t, J = 3.3 Hz, 1H), 7.62 (m, 4.5H), 7.49 (t, J = 7.8 Hz, 0.5H), 7.17 (br s, 2H), 6.79 (dd, J = 3.4, 1.9 Hz, 1H), 5.08 - 4.88 (m, 1H), 1.36 (d, J = 6.5 Hz, 3H).

EXAMPLE 170

(S)-2-(2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-5-yl)ethyl acetate

[0709] A mixture of (S)-4-amino-6-(1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (11 mg, 0.03mmol, Example 159) in acetic acid was heated overnight at 100°C. Removal of the solvent yielded the title compound as a solid (13mg, 100% yield).

LRMS (m/z): 459 (M+1)⁺.

¹H NMR (400 MHz, CDCl₃-d) δ ppm 1.41 (d, 3 H) 2.01 (s, 3 H) 3.25 (t, J=6.64 Hz, 2 H) 4.31 (t, J=6.84 Hz, 2 H) 5.07 (d, J=6.25 Hz, 1 H) 5.45 (s, 2 H) 5.80 (d, J=8.21 Hz, 1 H) 6.43 (d, J=2.74 Hz, 1 H) 7.28 - 7.35 (m, 2 H) 7.42 (d, J=5.86 Hz, 1 H) 7.47 - 7.61 (m, 3 H) 8.08 (s, 1 H)

EXAMPLE 171

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0710] The title compound (7 mg, 5% yield) was obtained from (S)-2-(1-aminoethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 110°C during 15 hours.

LRMS (m/z): 442 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.87 (s, 1 H), 8.14 - 7.97 (m, 3H), 7.88 (br s, 2H), 7.82 - 7.74 (dd, 1 H), 7.74 - 7.65 (m, 1 H), 7.04 (dd, 1 H), 6.66 (dd, 1 H), 5.26 - 5.13 (m, 1H), 1.52 (d, 3H).

EXAMPLE 172

2-((2S,4R)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-hydroxypyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile

[0711] 2-((2S,4R)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-(benzyloxy)pyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile (20 mg, 0.04 mmol) was dissolved in methanol (2 ml) and 10 drops of concentrated acetic acid were added. This mixture was hydrogenated in an H-Cube apparatus (1ml/min, 30 °C, full H₂) using Pd/C 10% as catalyst. Then the solvent was removed under vacuum and the product was purified by reverse phase chromatography (C-18 silica from Waters®, water/1:1 acetonitrile-methanol as eluents [0.1 % v/v formic acid buffered] 0% to 100%) to obtain the title compound (7 mg, 42% yield) as a white solid.

LRMS (m/z): 476 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 8.06 (s, 1H), 7.28 (d, 1H), 7.26 - 7.24 (m, 1 H), 7.06 (ddd, 1 H), 6.86 - 6.82 (m, 2H), 5.48 (s, 2H), 4.95 (t, 1 H), 4.84 - 4.73 (m, 1H), 4.31 (dd, 1H), 4.07 (d, 1 H), 2.26 (ddd, 1 H), 2.18 (ddd, 1 H).

EXAMPLE 173

4-Amino-6-((2S,4R)-2-(5-(aminomethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-4-hydroxypyrrolidin-1-yl)pyrimidine-5-carbonitrile

[0712] The title compound was obtained and isolated as a by-product of the reaction described in Example 172.

LRMS (m/z): 480 (M+1)⁺.

¹H NMR (600 MHz, cdcl₃) δ 8.09 (s, 1H), 7.20 (d, 1H), 7.02 (t, 2H), 6.83 (d, 1 H), 6.46 (d, 1 H), 5.42 (s, 2H), 4.94 (t, 1 H), 4.79 (s, 1 H), 4.33 (dd, 2H), 4.04 (s, 3H), 2.28 - 2.20 (m, 2H), 2.18 - 2.10 (m, 2H).

EXAMPLE 174

(S)-4-Amino-6-(1-(5-(4-methyl-1H-imidazol-1-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0713] In a sealed tube a mixture of (S)-4-amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (50 mg, 0.11 mmol, Example 144), 4-methyl-1H-imidazole (14 mg, 0.17 mmol), L-proline (3 mg, 0.03mmol), copper iodide (2mg, 0.01mmol) and K₃PO₄ (59 mg, 0.28 mmol) in DMSO (2 ml) was heated at 140°C overnight. The same amount of 4-methyl-1H-imidazole, L-proline, copper iodide and K₃PO₄ were added and the mixture stirred at 140°C overnight. The crude product was purified by reverse phase chromatography (C-18 silica from Waters, water/1:1 acetonitrile-methanol as eluents 0% to 100%) to obtain the title compound (10 mg, 20% yield) as a white solid.

LRMS (m/z): 453 (M+1)⁺.

EXAMPLE 175

(S)-4-Amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

a) (S)-2-(1-Aminoethyl)-5-bromo-3-(3-methoxyphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one

[0714] This compound was prepared starting from (*S*)-*tert*-butyl 1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylcarbamate (453 mg, 0.98 mmol) and following the experimental procedure described in Preparation 46c to afford 414 mg (99% yield) of the title compound as a dihydrochloride salt that was used in the next step without any further purification.
LRMS (*m/z*): 363, 365 (*M*+1)⁺.

b) (S)-4-Amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0715] The title compound (178 mg, 75% yield) was obtained from (*S*)-2-(1-aminoethyl)-5-bromo-3-(3-methoxyphenyl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (*m/z*): 481, 483 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.79 (s, 0.5H), 7.77 - 7.70 (m, 1.5H), 7.59 (m, 1 H), 7.32 (t, *J* = 8.1 Hz, 0.5H), 7.25 - 7.13 (m, 2.5H), 7.05 (d, *J* = 7.8 Hz, 0.5H), 6.88 (m, 2.5H), 6.75 (s, 1 H), 5.05 - 4.86 (m, 1 H), 3.75 (s, 1.5H), 3.66 (s, 1.5H), 1.36 (t, *J* = 6.5 Hz, 3H).

EXAMPLE 176

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile

[0716] The title compound (85 mg, 71% yield) was obtained from (*S*)-2-(1-aminoethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (*m/z*): 466 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 8.07 (s, 0.5H), 7.95 (dd, *J* = 2.9, 2.2 Hz, 1 H), 7.90 (d, *J* = 7.6 Hz, 0.5H), 7.76 - 7.64 (m, 3H), 7.60 (m, 1 H), 7.57 - 7.50 (m, 1 H), 7.24 (d, *J* = 3.0 Hz, 1 H), 7.22 (br s, 2H), 5.08 - 4.90 (m, 1 H), 1.39 (d, *J* = 6.6 Hz, 3H).

EXAMPLE 177

(S)-4-Amino-6-(1-(5-bromo-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0717] To a mixture of (*S*)-4-amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (100mg, 0.21 mmol, Example 175) in anhydrous DCM (5ml) at 0°C, was added dropwise 1 M solution of boron tribromide in DCM (623 μl, 0.62 mmol). The reaction mixture was then stirred at room temperature overnight. Ethyl acetate was added and the organic phase washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The title compound (91 mg, 90% yield) was obtained as a beige solid.

LRMS (*m/z*): 467, 469 (*M*+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 9.67 (2s, 1H), 7.85 (2s, 1 H), 7.69 (2d, *J* = 5.6, 1 H), 7.59 (t, *J* = 6.9 Hz, 1 H), 7.23 (t, *J* = 8.0 Hz, 0.5H), 7.19 (br s, 2H), 7.07 (t, *J* = 8.0 Hz, 0.5H), 6.86 (m, 1 H), 6.81 - 6.67 (m, 3H), 4.96 - 4.75 (2q, *J* = 6.5 Hz, 1 H), 1.36 (2d, *J* = 6.5 Hz, 3H)

EXAMPLE 178

(S)-4-Amino-6-(1-(3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0718] To a mixture of (*S*)-4-amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-

2-yl)ethylamino)pyrimidine-5-carbonitrile (50 mg, 0.10 mmol, Example 175) and triethylamine (0.30 mmol) in MeOH (10 ml), was added Pd/C (10%) and the reaction mixture was hydrogenated at 4 psi for 2h. The catalyst was filtered off and the filtrate concentrated. Ethyl acetate was added and the organic phase washed with water and brine, dried over magnesium sulphate and the solvent evaporated. The title compound (30 mg, 71% yield) was obtained as a white solid.

LRMS (m/z): 403 (M+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 7.77 (2s, 1 H), 7.69 (dd, J = 4.5, 3.1 Hz, 1 H), 7.65 (dd, J = 9.5, 7.3 Hz, 1 H), 7.32 (t, J = 8.0 Hz, 0.5H), 7.23 - 7.12 (m, 3H), 7.04 (d, J = 7.6 Hz, 0.5H), 6.95 (dd, J = 4.2, 1.5 Hz, 1 H), 6.90 (m, 1 H), 6.85 (m, 1 H), 6.65 - 6.59 (m, 1 H), 5.09 - 4.94 (m, 1 H), 3.71 (2sd, 3H), 1.39 (2d, J = 6.6 Hz, 3H).

EXAMPLE 179

(S)-4-Amino-6-(1-(3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile

[0719] The title compound (35 mg, 82% yield) was obtained from (S)-4-amino-6-(1-(5-bromo-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (Example 177) following the experimental procedure described in Example 178.

LRMS (m/z): 389 (M+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 9.66 (2s, 1H), 7.84 (2s, 1H), 7.65 (m, 2H), 7.23 (t, J = 8.0 Hz, 0.5H), 7.18 (br s, 2H), 7.07 (t, J = 8.0 Hz, 0.5H), 6.93 (dd, J = 4.3, 1.6 Hz, 1 H), 6.86 (m, 0.5H), 6.82 (dt, J = 7.5, 2.0 Hz, 1 H), 6.76 (m, 1 H), 6.69 (m, 0.5H), 6.59 (2t, J = 4.4 Hz, 1 H), 5.00 - 4.93 (m, 0.5H), 4.92 - 4.85 (m, 0.5H), 1.38 (2d, J = 6.6 Hz, 3H).

EXAMPLE 180

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile

[0720] The title compound (177 mg, 87% yield) was obtained from (S)-2-(1-aminoethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 18.

LRMS (m/z): 428 (M+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 7.89 (2d, J = 2.8 Hz, 1 H), 7.77 (2s, 1H), 7.56 (m, 1H), 7.35 (m, 0.5H), 7.28 - 7.17 (m, 3.5H), 7.12 - 7.06 (m, 0.5H), 6.90 (m, 2.5H), 5.03 (2q, J = 6.6 Hz, 1H), 3.71 (2s, 3H), 1.39 (2d, J = 6.2 Hz, 3H).

EXAMPLE 181

4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)cyclopropylamino)pyrimidine-5-carbonitrile

[0721] The title compound (73 mg, 36% yield) was obtained from 2-(1-aminocyclopropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 4-amino-6-chloropyrimidine-5-carbonitrile following the experimental procedure described in example 23 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 385 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 7.99 (s, 1H), 7.68 (s, 1H), 7.59 - 7.45 (m, 3H), 7.39 - 7.15 (m, 4H), 6.90 (dd, 1 H), 6.58 (dd, 1 H), 5.38 (dd, 1 H), 1.83 - 1.69 (m, 2H), 1.15 - 1.03 (m, 2H).

EXAMPLE 182

2-(1-(9H-Purin-6-ylamino)cyclopropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one

[0722] The title compound (18 mg, 9% yield) was obtained from 2-(1-aminocyclopropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one and 6-bromo-9H-purine following the experimental procedure described in example 20 but using n-butanol as solvent and heating the reaction mixture at 120°C during 15 hours.

LRMS (m/z): 385 (M+1)⁺.

¹H NMR (400 MHz, DMSO) δ 12.96 (s, 1H), 8.17 (s, 1H), 8.06 (s, 1H), 7.71 (s, 1 H), 7.53 - 7.31 (m, 3H), 7.20 - 7.02 (m, 2H), 6.90 (dd, 1 H), 6.60 (dd, 1 H), 5.91 (dd, 1H), 1.87 - 1.75 (m, 2H), 1.16 - 1.05 (m, 2H).

EXAMPLE 183**(S)-4-Amino-6-(1-(4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile**

[0723] The title compound (35 mg, 82% yield) was obtained from (S)-4-amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile (Example 169) following the experimental procedure described in Example 178.

LRMS (m/z): 441 (M+1)⁺.

¹H NMR (600 MHz, DMSO, 1/1 mixture of isomers) δ 8.06 (s, 0.5H), 7.88 (d, J = 7.8 Hz, 0.5H), 7.76 (m, 1 H), 7.71 - 7.55 (m, 4.5H), 7.48 (t, J = 7.9 Hz, 0.5H), 7.20 (br s, 2H), 6.98 (m, 1 H), 6.65 (m, 1 H), 5.09 - 5.02 (p, J = 6.6 Hz, 0.5H), 4.99 (p, J = 6.6 Hz, 0.5H), 1.42 - 1.33 (m, 3H).

EXAMPLE 184**(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile**

[0724] The title compound (25 mg, 17% yield) was obtained from (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-*f*][1,2,4]triazine-5-carbonitrile (Example 180) following the experimental procedure described in example 177.

LRMS (m/z): 414 (M+1)⁺.

¹H NMR (400 MHz, DMSO, 1/1 mixture of isomers) δ 9.73 (2s, 1 H), 7.85 (m, 2H), 7.64 - 7.54 (m, 1 H), 7.29 - 7.06 (m, 4H), 6.91 (m, 1 H), 6.85 - 6.69 (m, 2H), 5.05 - 4.84 (m, 1 H), 1.39 (m, 3H).

[0725] Following a similar procedure to that described above, the following compounds were obtained:

EXAMPLE 185**(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(pyridin-2-yl)pyrrolo[1,2-*f*][1,2,4]triazin-4(3H)-one**EXAMPLE 186**(S)-2-(1-(9H-Purin-6-ylamino)propyl)-3-phenylimidazo[1,2-*f*][1,2,4]triazin-4(3H)-one**EXAMPLE 187**(S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-*f*][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile**REFERENCES**[0726]**

- 1.- Tehrani, A. K.; Borremans D.; De Kimpe N. Tetrahedron 1999, 55, 4133-4152
- 2.- Ohta, T.; Fukuda, T.; Ishibashi, F., Iwao, M. J. Org. Chem. 2009, 74, 8143-8153
- 3.- Leroy, J.; Porthiel, E.; Bondon, A. Tetrahedron 2002, 58, 6713-6722

PHARMACOLOGICAL ACTIVITY**PI3K α , β , δ and γ Enzymatic Inhibition Assays**

[0727] Compounds were screened for their ability to inhibit PI3K α (PI3K α), PI3K β (PI3K β), PI3K δ (PI3K δ) and PI3K γ (PI3K γ) using a cell-free based PI3K HTRF™ assay (Millipore, ref. #33-017) All reagents to perform the reactions were prepared according to the manufacturer protocol. All PI3K enzymes were recombinant and were purchased at Millipore.

[0728] All four assays were performed according to the following procedure:

- 1) Dilution curves of compounds were done in 100% DMSO and dispensed in a reservoir plate, typically from column 2 to 11. Column 1 and 12 were used for the negative controls (100% inhibition using a reference PI3K inhibitor for

EP 2 702 061 B1

the four isoforms) and the positive controls (0% inhibition using dimethyl sulfoxide (DMSO) only).

2) Mix of Phosphoinositide 3-kinase (PI3K) + Phosphatidylinositol 4,5-bisphosphate (PIP2) was diluted in buffer (supplied with the kit) and plated in a medium binding black 96-well plate (Greiner ref. #675076) PI3Ka was diluted at 0.25 nM with PIP2 at 2 μ M; PI3Kb was diluted at 0.50 nM with PIP2 at 5 μ M; PI3Kd was diluted at 0.60 nM with PIP2 at 2 μ M and PI3Kg was diluted at 0.30 nM with PIP2 at 10 μ M. These concentrations were final in the assay.

3) A reservoir plate containing Adenosine TriPhosphate (ATP) diluted in the Millipore kit buffer was prepared for each isoform. The final concentration of ATP in the assay was 10 μ M, 15 μ M, 20 μ M and 10 μ M for PI3Ka, PI3Kb, PI3Kd and PI3Kg respectively.

4) Reactions were started by addition to the PI3K+PIP2 plates of the ATP and compounds simultaneously. Incubation was done for 8 minutes (for all four isoforms) then the reaction was stopped by addition of the Stop solution. The Detection solution was added. These solutions were prepared previously according to the kit specifications. The plates were then incubated overnight at RT before reading the signal in an Envision (PerkinElmer), with excitation at 340 nm and emissions at 620 and 665 nm.

5) Data obtained from the compound curves were normalized in respect to the negative and positive controls, and then fitted by a 4-parameter log curve in ActivityBase (IDBS) in order to determine their potency.

[0729] The results are shown in Table 1.

Example	IC ₅₀ PI3Kd HTRF (nM)
1	120
11	119
15	18
17	9
20	11
21	5
23	4
25	347
29	376
31	6566
37	15
43	11
44	29
47	1
50	57
54	60
56	15
61	6
62	73
64	69
68	1
71	9
73	31
75	17
81	9
90	7

(continued)

Example	IC ₅₀ PI3Kd HTRF (nM)
102	42
104	36
122	2928
124	11
125	90
127	89
129	38
133	11
136	24
138	8
142	5
144	1
152	747
157	66
158	43
159	35
164	67
165	670
169	6
176	32
179	7

[0730] It can be seen from Table 1 that the compounds of formula (I) are potent inhibitors of Phosphoinositide 3-kinase delta (PI3kd). Preferred compounds of the invention possess an IC₅₀ value for the inhibition of PI3Kd (determined as defined above) of less than 10 μ M (10,000 nM), preferably less than 1 μ M (1,000 nM), even more preferably of less than 0.2 μ M (200 nM), most preferably less than 0.05 μ M (50 nM)

[0731] The invention is also directed to a compound of the invention as described herein for use in the treatment of the human or animal body by therapy. Compounds of the invention intended for pharmaceutical use may be administered as crystalline or amorphous products, or mixtures thereof. They may be obtained, for example, as solid plugs, powders, or films by methods such as precipitation, crystallization, freeze drying, spray drying, or evaporative drying. Microwave or radio frequency drying may be used for this purpose.

Combinations

[0732] The pyrrolotriazinone derivatives defined herein may also be combined with other active compounds in the treatment of a pathological condition or disease susceptible to amelioration by inhibition of PI3Ks.

[0733] The combinations of the invention can optionally comprise one or more additional active substances which are known to be useful in the treatment of respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors.

[0734] Particularly, the combinations of the invention can optionally comprise one or more additional active substances which are known to be useful in the treatment of neoplastic diseases (e.g. leukemia, lymphomas, solid tumors); transplant rejection, bone marrow transplant applications (e.g., graft- versus-host disease); autoimmune diseases (e.g. rheumatoid

arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis and blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa; respiratory inflammation diseases (e.g. asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis); skin inflammatory diseases (e.g., atopic dermatitis, contact dermatitis, eczema or psoriasis); premalignant and malignant skin conditions (e.g. basal cell carcinoma (BCC), squamous cell carcinoma (SCC) or actinic keratosis (AK)); neurological disorders and pain (such as pain associated with rheumatoid arthritis or osteoarthritis, back pain, general inflammatory pain, inflammatory neuropathic pain, trigeminal neuralgia or central pain).

[0735] Preferably, the combinations of the invention can optionally comprise one or more additional active substances which are known to be useful in the treatment of neoplastic diseases leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0736] In particular, the combinations of the invention can optionally comprise one or more additional active substances which are known to be useful in the treatment of neoplastic diseases leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis

[0737] The combinations of the invention comprise (i) a compound of the invention as defined above; and (ii) another compound selected from the group consisting of an Adenosine A_{2A} agonist, an agent for treating cardiovascular disorders, an agent for treating diabetes, and an agent for treating liver disease, an anti-allergic agent, an anti-cholinergic agent, an anti-inflammatory agent, an anti-infective agent, a β 2-adrenergic agonist, a Chemoattractant receptor homologous molecule expressed on TH₂ cells (CRTH2) inhibitor, a chemotherapeutic agent, a corticosteroid, an IKK β /IKBK β (IKB kinase beta or IKK2) inhibitor, an immunosuppressant, a Janus kinase (JAK) inhibitor, a topically acting p38 Mitogen-Activated Protein Kinase (p38 MAPK) inhibitor, a Phosphodiesterase (PDE) IV inhibitor, and a Spleen tyrosine kinase (Syk) inhibitor, for simultaneous, separate or sequential use in the treatment of the human or animal body.

[0738] In a particular embodiment, the combinations of the invention can optionally comprise one or more additional active substances selected from

- a) Dihydrofolate reductase inhibitors, such as Methotrexate or CH-1504;
- b) Dihydroorotate dehydrogenase (DHODH) inhibitors such as leflunomide, teriflunomide, or the compounds described in the International Patent Application Nos. WO2008/077639 and WO2009/021696;
- c) Immunomodulators such as Glatiramer acetate (Copaxone), Laquinimod or Imiquimod;
- d) Inhibitors of DNA synthesis and repair, such as Mitoxantrone or Cladribine;
- e) Immunosuppressants, such as Imuran (azathioprine) or Purinethol (6-mercaptopurine or 6-MP);
- f) Anti-alpha 4 integrin antibodies, such as Natalizumab (Tysabri) ;
- g) Alpha 4 integrin antagonists such as R-1295 , TBC-4746, CDP-323, ELND-002, Finategrast or TMC-2003;
- h) Corticoids and glucocorticoids such as prednisone or methylprednisolone, fluticasone, mometasone, budesonide, ciclesonide or beta-metasone;
- i) Fumaric acid esters, such as BG-12;
- j) Anti-tumor necrosis factor-alpha (Anti-TNF-alpha) monoclonal antibodies such as Infliximab, Adalimumab or Certolizumab pegol;
- k) Soluble Tumor necrosis factor-alpha (TNF-alpha) Antagonists such as Ethenarcept;
- l) Anti-CD20 (lymphocyte protein) monoclonal antibodies such as Rituximab, Ocrelizumab Ofatumumab or TRU-015;
- m) Anti-CD52 (lymphocyte protein) monoclonal antibodies such as alemtuzumab;
- n) Anti-CD25 (lymphocyte protein) such as daclizumab;
- o) Anti-CD88 (lymphocyte protein), such as eculizumab or pexelizumab;
- p) Anti-Interleukin 6 Receptor (IL-6R), such as tocilizumab;
- q) Anti-Interleukin 12 Receptor (IL-12R) / Interleukin 23 Receptor (IL-23R), such as ustekinumab;
- r) Calcineurin inhibitors such as cyclosporine A or tacrolimus;
- s) Inosine-monophosphate dehydrogenase (IMPDH) inhibitors, such as mycophenolate mophetyl, ribavirin, mizoribine or mycophenolic acid;
- t) Cannabinoid receptor agonists such as Sativex;
- u) Chemokine CCR1 antagonists such as MLN-3897 or PS-031291;

- v) Chemokine CCR2 antagonists such as INCB-8696;
- w) Necrosis factor-kappaB (NF-kappaB or NFKB) Activation Inhibitors such as Sulfasalazine, Igaratimod or MLN-0415;
- x) Adenosine A_{2A} agonists, such as ATL-313, ATL-146e, CGS-21680, Regadenoson or UK-432,097;
- 5 y) Sphingosine-1 (S1 P) phosphate receptor agonists such as fingolimod, BAF-312, or ACT128800;
- z) Sphingosine-1 (S1 P) liase inhibitors such as LX2931;
- aa) Spleen tyrosine kinase (Syk) inhibitors, such as R-112;
- bb) Protein Kinase Inhibitors (PKC) inhibitors, such as NVP-AEB071;
- cc) Anti-cholinergic agents such as tiotropium or aclidinium;
- 10 dd) Beta adrenergic agonists such as formoterol, indacaterol or LAS100977 (abediterol);
- ee) MABA (molecules with dual activity: beta-adrenergic agonists and muscarinic receptor antagonists)
- ff) Histamine 1 (H1) receptor antagonists, such as azelastine or ebastine;
- gg) Cysteinyl leukotriene (CysLT) receptor antagonists, such as montelukast;
- hh) Mast cell stabilizers, such as nedocromil or chromoglycate;
- 15 ii) 5-lipoxygenase-activating protein (FLAP) inhibitors, such as MK886 or BAY X 1005;
- jj) 5-lipoxygenase (5-LO) inhibitors, such as WY-50295T;
- kk) Chemoattractant receptor homologous molecule expressed on TH₂ cells (CRTH2) inhibitors, such as OC-459, AZD-1981, ACT-129968, QAV-680;
- ll) Vitamin D derivatives like calcipotriol (Daivonex) ;
- 20 mm) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (NSAIDs) or selective cyclooxygenase-2 (COX-2) inhibitors such as aceclofenac, diclofenac, ibuprofen, naproxen, apricoxib, celecoxib, cimicoxib, deracoxib, etoricoxib, lumiracoxib, parecoxib sodium, rofecoxib, selenocoxib-1 or valdecoxib;
- nn) Anti-allergic agents;
- oo) Anti-viral agents;
- 25 pp) Phosphodiesterase (PDE) III inhibitors;
- qq) Phosphodiesterase (PDE) IV inhibitors such as roflumilast or GRC-4039;
- rr) Dual Phosphodiesterase (PDE) III/IV inhibitors;
- ss) Xanthine derivatives, such as theophylline or theobromine;
- tt) p38 Mitogen-Activated Protein Kinase (p38 MAPK) Inhibitors such as ARRY-797;
- 30 uu) Mitogen-activated extracellular signal regulated kinase kinase (MEK) inhibitor, such as ARRY-142886 or ARRY-438162;
- vv) Janus kinase (JAK) inhibitors, such as tofacitinib (previously known as tasocitinib or CP-690,550) from Pfizer and INCB-18424, from Incyte;
- ww) Interferons comprising Interferon beta 1 a such as Avonex from Biogen Idec, CinnoVex from CinnaGen and Rebif from EMD Serono, and Interferon beta 1b such as Betaferon from Schering and Betaseron from Berlex;
- 35 xx) Interferon alpha such as Sumiferon MP;
- yy) Epidermal Growth Factor Receptor (EGFR) inhibitors such as erlotinib, Trastuzumab, Herceptin, Avastin, Platins (cisplatin, carboplatin) or Temazolamide;
- zz) Antineoplastic agents such as Docetaxel, Estramustine, Anthracyc lines, (doxorubicin (Adriamycin), epirubicin (Ellence), and liposomal doxorubicin (Doxil)), Taxanes (docetaxel (Taxotere), paclitaxel (Taxol), and protein-bound paclitaxel (Abraxane)), Cyclophosphamide (Cytosan), Capecitabine (Xeloda), 5 fluorouracil (5 FU), Gemcitabine (Gemzar) or Vinorelbine (Navelbine);
- 40

[0739] Specific examples of suitable corticoids and glucocorticoids that can be combined with the PI3K inhibitors of the present invention are prednisolone, methylprednisolone, dexamethasone, dexamethasone cipeclate, nafcort, deflazacort, halopredone acetate, budesonide, beclomethasone dipropionate, hydrocortisone, triamcinolone acetonide, fluocinolone acetonide, fluocinonide, clocortolone pivalate, methylprednisolone aceponate, dexamethasone palmitoate, tipredane, hydrocortisone aceponate, prednicarbate, alclometasone dipropionate, halometasone, methylprednisolone suleptanate, mometasone furoate, rimexolone, prednisolone farnesylate, ciclesonide, butixocort propionate, RPR-106541, deprodone propionate, fluticasone propionate, fluticasone furoate, halobetasol propionate, loteprednol etabonate, betamethasone butyrate propionate, flunisolide, prednisone, dexamethasone sodium phosphate, triamcinolone, betamethasone 17-valerate, betamethasone, betamethasone dipropionate, hydrocortisone acetate, hydrocortisone sodium succinate, prednisolone sodium phosphate and hydrocortisone probutate. Specific examples of suitable Syk kinase inhibitors that can be combined with the PI3K inhibitors of the present invention are fosfatinib (from Rigel), R-348 (from Rigel), R-343 (from Rigel), R-112 (from Rigel), piceatannol, 2-(2-Aminoethylamino)-4-[3-(trifluoromethyl)phenylamino] pyrimidine-5-carboxamide, R-091 (from Rigel), 6-[5-Fluoro-2-(3,4,5-trimethoxyphenylamino)pyrimidin-4-ylamino]-2,2-dimethyl-3,4-dihydro-2H-pyrido[3,2-b][1,4]oxazin-3-one benzenesulfonate (R-406 from Rigel), 1-(2,4,6-Trihydroxyphenyl)-2-(4-methoxyphenyl)ethan-1-one, N-[4-[6-(Cyclobutylamino)-9H-purin-2-ylamino]phenyl]-N-methyla-

cetamide (QAB-205 from Novartis), 2-[7-(3,4-Dimethoxyphenyl)imidazo[1,2-c]pyrimidin-5-ylamino]pyridine-3-carboxamide dihydrochloride (BAY-61-3606 from Bayer) and AVE-0950 (from Sanofi-Aventis).

[0740] Specific examples of suitable M3 antagonists (anticholinergics) that can be combined with the PI3K inhibitors of the present invention are tiotropium salts, oxitropium salts, flutropium salts, ipratropium salts, glycopyrronium salts, trospium salts, zamifenacin, revatropate, espatropate, darotroprum bromide, CI-923, NPC-14695, BEA-2108, 3-[2-Hydroxy-2,2-bis(2-thienyl)acetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane salts (in particular acridinium salts, more preferably acridinium bromide), 1-(2-Phenylethyl)-3-(9H-xanthen-9-ylcarbonyloxy)-1-azoniabicyclo[2.2.2]octane salts, 2-oxo-1,2,3,4-tetrahydroquinazoline-3-carboxylic acid endo-8-methyl-8-azabicyclo[3.2.1]oct-3-yl ester salts (DAU-5884), 3-(4-Benzylpiperazin-1-yl)-1-cyclobutyl-1-hydroxy-1-phenylpropan-2-one (NPC-14695), N-[1-(6-Aminopyridin-2-ylmethyl)piperidin-4-yl]-2(R)-[3,3-difluoro-1 (R)-cyclopentyl]-2-hydroxy-2-phenylacetamide (J-104135), 2(R)-Cyclopentyl-2-hydroxy-N-[1-[4(S)-methylhexyl]piperidin-4-yl]-2-phenylacetamide (J-106366), 2(R)-Cyclopentyl-2-hydroxy-N-[1-(4-methyl-3-pentenyl)-4-piperidinyl]-2-phenylacetamide (J-104129), 1-[4-(2-Aminoethyl)piperidin-1-yl]-2(R)-[3,3-difluorocyclopent-1 (R)-yl]-2-hydroxy-2-phenylethan-1-one (Banyu-280634), N-[N-[2-[N-[1-(Cyclohexylmethyl)piperidin-3(R)-ylmethyl]carbamoyl]ethyl]carbamoylmethyl]-3,3,3-triphenylpropionamide (Banyu CPTP), 2(R)-Cyclopentyl-2-hydroxy-2-phenylacetic acid 4-(3-azabicyclo[3.1.0]hex-3-yl)-2-butyryl ester (Ranbaxy 364057), 3(R)-[4,4-Bis(4-fluorophenyl)-2-oxoimidazolidin-1-yl]-1-methyl-1-[2-oxo-2-(3-thienyl)ethyl]pyrrolidinium iodide, N-[1-(3-Hydroxybenzyl)-1-methylpiperidinium-3(S)-yl]-N-[N-[4-(isopropoxycarbonyl)phenyl]carbamoyl]-L-tyrosinamide trifluoroacetate, UCB-101333, Merck's OrM3, 7-endo-(2-hydroxy-2,2-diphenylacetoxy)-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^(2,4)]nonane salts, 3(R)-[4,4-Bis(4-fluorophenyl)-2-oxoimidazolidin-1-yl]-1-methyl-1-(2-phenylethyl)pyrrolidinium iodide, trans-4-[2-[Hydroxy-2,2-(dithien-2-yl)acetoxy]-1-methyl-1-(2-phenoxyethyl)piperidinium bromide from Novartis (412682), 7-(2,2-diphenylpropionyloxy)-7,9,9-trimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^(2,4)]nonane salts, 7-hydroxy-7,9,9-trimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^(2,4)]nonane 9-methyl-9H-fluorene-9-carboxylic acid ester salts, all of them optionally in the form of their racemates, their enantiomers, their diastereomers and mixtures thereof, and optionally in the form of their pharmacologically-compatible acid addition salts. Among the salts chlorides, bromides, iodides and methanesulphonates are preferred.

[0741] Specific examples of suitable beta adrenergic agonists (β_2 -agonists) that can be combined with the PI3K inhibitors of the present invention are terbutaline sulphate, eformoterol fumarate, formoterol fumarate, bambuterol, ibuterol, isoprenaline hydrochloride, dopexamine, metaprotenerol, tulobuterol, procaterol hydrochloride, sibenadet hydrochloride, mabuterol hydrochloride, albuterol sulphate, salbutamol sulphate, salmefamol, salmeterol xinafoate, carmoterol hydrochloride, (R)-albuterol hydrochloride, Levalbuterol hydrochloride; Levosalbutamol hydrochloride; (-)-Salbutamol hydrochloride, formoterol, (R,R)-Formoterol tartrate; Arformoterol tartrate, sulfoneterol, Bedoradrine sulphate, Indacaterol, Trantinterol hydrochloride, Milveterol hydrochloride, Olodaterol, fenoterol hydrobromide, rimoterol hydrobromide, riproterol hydrochloride, Vilanterol broxaterol, pirbuterol hydrochloride, bitolterol mesylate, clenbuterol hydrochloride, AZD-3199, GSK-159802; GSK-597901, GSK-678007, GSK-961081; 4-[2-[3-(1H-Benzimidazol-1-yl)-1,1-dimethylpropylamino]-1-hydroxyethyl]-2-(4-methoxybenzylamino)phenol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-N,N-dimethylaminophenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-domethoxyphenyl)-2-methyl-2-propylamino]ethanol, 1-[2H-5-hydroxy-3-oxo-4H-1,4-benzoxazin-8-yl]-2-[3-(4-n-butyloxyphenyl)-2-methyl-2-propylamino]ethanol, KUL-1248, HOKU-81, SM-110444, RP-58802B, LAS100977 (abediterol) and compounds described in PCT patent applications Nos. WO 2007/124898, WO 2006/122788A1, WO 2008/046598, WO 2008095720, WO 2009/068177 and WO 2010/072354.

[0742] Specific examples of suitable anti-allergic agents that can be combined with the PI3K inhibitors of the present invention are anti-histamines (e.g. Methapyrilene, Mequitazine, Azelastine hydrochloride, Acrivastine, Emedastine difumarate, Emedastine fumarate, Loratadine, Cyproheptadine hydrochloride, Diphenhydramine hydrochloride, Doxepin hydrochloride, Promethazine hydrochloride, Levocabastine hydrochloride, Desloratadine, Cinnarizine, Setastine hydrochloride, Mizolastine, Ebastine, Cetirizine hydrochloride, Epinastine hydrochloride, Olopatadine hydrochloride, Bepotastine besilate, Triprolidine hydrochloride, Rupatadine fumarate, Fexofenadine hydrochloride, Levocetirizine dihydrochloride, Ketotifen, Azatadine maleate, Dimethindene maleate, Clemastine fumarate, Alcaftadine, Bilastine, Vapitadine hydrochloride, AZD-1744, GSK-1004723D, GSK-835726 or SUN-1334H.

[0743] Specific examples of suitable Phosphodiesterase IV (PDE IV) inhibitors that can be combined with the PI3K inhibitors of the present invention are benafentrine dimaleate, etazolate, denbufylline, rolipram, cipamfylline, zardaverine, arofylline, filaminast, tiplukast, tofimidast, piclamilast, tolafentrine, mesopram, drotaverine hydrochloride, lirimidast, roflumilast, cilomilast, oglemilast, apremilast, tetomilast, filaminast, (R)-(+)-4-[2-(3-Cyclopentylloxy-4-methoxyphenyl)-2-phenylethyl]pyridine (CDP-840), N-(3,5-Dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide (GSK-842470), 9-(2-Fluorobenzyl)-N6-methyl-2-(trifluoromethyl)adenine (NCS-613), N-(3,5-Dichloro-4-pyridinyl)-8-methoxyquinoline-5-carboxamide (D-4418), 3-[3-(Cyclopentylloxy)-4-methoxybenzyl]-6-(ethylamino)-8-isopropyl-3H-purine hydrochloride (V-11294A), 6-[3-(N,N-Dimethylcarbamoyl)phenylsulfonyl]-4-(3-methoxyphenylamino)-8-methylquinoline-3-carboxamide hydrochloride (GSK-256066), 4-[6,7-Diethoxy-2,3-bis(hydroxymethyl)naphthalen-1-yl]-1-(2-methoxyethyl)pyridin-2(1H)-one (T-440), (-)-trans-2-[3'-[3-(N-Cyclopropylcarbamoyl)-4-oxo-1,4-dihydro-1,8-naph-

thyridin-1-yl]-3-fluorobiphenyl-4-yl]cyclopropanecarboxylic acid, MK-0873, CDC-801, UK-500001, BLX-914, 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-one, cis [4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol, 5(S)-[3-(Cyclopentyloxy)-4-methoxyphenyl]-3(S)-(3-methylbenzyl)piperidin-2-one (IPL-455903), ONO-6126 (Eur Respir J 2003, 22(Suppl. 45): Abst 2557) and the compounds claimed in the PCT patent applications number WO 03/097613, WO 2004/058729, WO 2005/049581, WO 2005/123693, WO 2005/123692, and WO 2010/069504.

[0744] Specific examples of suitable immunosuppressants that can be combined with the PI3K inhibitors of the present invention are picremolimus, tacrolimus, cyclosporine A, leflunomide, teriflunomide, vidofludimus, laquinimod, methotrexate, 5-fluorouracil (5-FU), anti-TNF agents and compounds described in PCT patent applications Nos. WO 2008/077639, WO 2009/021696, WO 2009/153043, and WO2010083975 (in particular amino(iso)nicotinic acid derivatives selected from the group consisting of 2-(3'-ethoxy-3-(trifluoromethoxy)biphenyl-4-ylamino)nicotinic acid, 2-(3,5-difluoro-3'-methoxybiphenyl-4-ylamino)nicotinic acid and 2-(3,5-difluoro-2-methylbiphenyl-4-ylamino)nicotinic acid; and azabiphenylaminobenzoic acid derivatives selected from the group consisting of 5-cyclopropyl-2-(2-(2,6-difluorophenyl)pyrimidin-5-ylamino)benzoic acid, 5-cyclopropyl-2-((2-(2-(trifluoromethyl)phenyl)pyrimidin-5-yl)amino)benzoic acid and 5-methyl-2-((6-(2,3-difluorophenyl)pyridin-3-yl)amino)benzoic acid)

[0745] Specific examples of suitable anti-infectives that can be combined with the PI3K inhibitors of the present invention are aclarubicin, actinomycin D, amrubicin, annamycin, adhamycin, bleomycin, daunorubicin, doxorubicin, el-samitrucin, epirubicin, galarubicin, idarubicin, mitomycin C, mupiricin, nemorubicin, neocarzinostatin, peplomycin, pirarubicin, rebeccamycin, retapamulin, stimalamer, streptozocin, valrubicin, zinostatin, amphotericin B, bifonazole, caspofungin, clotrimazole, echinocandin B, econazole, fluconazole, flucytosine, itraconazole, ketoconazole, miconazole, posaconazole, ravuconazole, terbinafine, tioconazole, voriconazole and combinations thereof.

[0746] Particularly preferred combination products according to the invention comprise a compound of formula (I) and a therapeutically effective amount of one or more additional therapeutic agents selected from the group consisting of mometasone furoate, ciclesonide, budesonide, fluticasone propionate, fluticasone furoate, betamethasone valerate, clobetasol propionate, tiotropium salts, glycopyrronium salts, 3-[2-Hydroxy-2,2-bis(2-thienyl)acetoxy]-1-(3-phenoxypropyl)-1-azoniabicyclo[2.2.2]octane salts (in particular acridinium salts, preferably acridinium bromide), 1-(2-Phenylethyl)-3-(9H-xanthen-9-ylcarbonyloxy)-1-azoniabicyclo[2.2.2]octane salts, formoterol, salmeterol, indacaterol, carmoterol, LAS 100977 (abediterol), compounds described in PCT patent applications Nos. WO 2008/077639, WO 2009/021696, WO 2009/153043, and WO 2010/083975 (in particular amino(iso)nicotinic acid derivatives selected from the group consisting of 2-(3'-ethoxy-3-(trifluoromethoxy)biphenyl-4-ylamino)nicotinic acid, 2-(3,5-difluoro-3'-methoxybiphenyl-4-ylamino)nicotinic acid and 2-(3,5-difluoro-2-methylbiphenyl-4-ylamino)nicotinic acid; and azabiphenylaminobenzoic acid derivatives selected from the group consisting of 5-cyclopropyl-2-(2-(2,6-difluorophenyl)pyrimidin-5-ylamino)benzoic acid, 5-cyclopropyl-2-((2-(2-(trifluoromethyl)phenyl)pyrimidin-5-yl)amino)benzoic acid and 5-methyl-2-((6-(2,3-difluorophenyl)pyridin-3-yl)amino)benzoic acid), methapyrilene, cetirizine, loratadine, ebastine, desloratadine, fexofenadine, azelastine, levocabastine, olopatadine, Montelukast, picremolimus, tacrolimus, mupiricin, retapamulin, clotrimazole, ketoconazole and terbinafine.

[0747] The compounds of formula (I) and the combinations of the invention may be used in the treatment of respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs such as polycythemia vera, essential thrombocythemia or myelofibrosis); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors, wherein the use of a PI3K inhibitor is expected to have a beneficial effect, for example leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0748] In particular the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0749] The active compounds in the combination product may be administered together in the same pharmaceutical composition or in different compositions intended for separate, simultaneous, concomitant or sequential administration by the same or a different route.

[0750] It is contemplated that all active agents would be administered at the same time, or very close in time. Alternatively, one or two actives could be administered in the morning and the other (s) later in the day. Or in another scenario,

one or two actives could be administered twice daily and the other (s) once daily, either at the same time as one of the twice-a-day dosing occurred, or separately. Preferably at least two, and more preferably all, of the actives would be administered together at the same time. Preferably, at least two, and more preferably all actives would be administered as an admixture.

[0751] The invention is also directed to a combination product of the compounds of the invention together with one or more other therapeutic agents for use in the treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinases (PI3Ks), in particular wherein the pathological condition or disease is selected from respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs such as polycythemia vera, essential thrombocythemia or myelofibrosis); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0752] In particular the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0753] The invention also encompasses the use of a combination of the compounds of the invention together with one or more other therapeutic agents for the manufacture of a formulation or medicament for treating these diseases.

[0754] The active compounds in the combinations of the invention may be administered by any suitable route, depending on the nature of the disorder to be treated, e.g. orally (as syrups, tablets, capsules, lozenges, controlled-release preparations, fast-dissolving preparations, etc); topically (as creams, ointments, lotions, nasal sprays or aerosols, etc); by injection (subcutaneous, intradermic, intramuscular, intravenous, etc.) or by inhalation (as a dry powder, a solution, a dispersion, etc).

[0755] The active compounds in the combination, i.e. the pyrrolotriazinone derivatives of the invention, and the other optional active compounds may be administered together in the same pharmaceutical composition or in different compositions intended for separate, simultaneous, concomitant or sequential administration by the same or a different route.

[0756] One execution of the present invention consists of a kit of parts comprising a imidazopyridine derivative of the invention together with instructions for simultaneous, concurrent, separate or sequential use in combination with another active compound useful in the treatment of respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs such as polycythemia vera, essential thrombocythemia or myelofibrosis); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0757] In particular the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0758] Another execution of the present invention consists of a package comprising a imidazopyridine derivative of the invention and another active compound useful in the treatment of respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs such as polycythemia vera, essential thrombocythemia or myelofibrosis); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheu-

matoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0759] In particular the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

Pharmaceutical Compositions

[0760] Pharmaceutical compositions according to the present invention comprise the compounds of the invention in association with a pharmaceutically acceptable diluent or carrier.

[0761] As used herein, the term pharmaceutical composition refers to a mixture of one or more of the compounds described herein, or physiologically/pharmaceutically acceptable salts, solvates, N-oxides, stereoisomers, deuterated derivatives thereof or prodrugs thereof, with other chemical components, such as physiologically/pharmaceutically acceptable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

[0762] As used herein, a physiologically/pharmaceutically acceptable diluent or carrier refers to a carrier or diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound.

[0763] The invention further provides pharmaceutical compositions comprising the compounds of the invention in association with a pharmaceutically acceptable diluent or carrier together with one or more other therapeutic agents for use in the treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinases (PI3Ks), such as the ones previously described.

[0764] The invention is also directed to pharmaceutical compositions of the invention for use in the treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinases (PI3Ks), in particular wherein the pathological condition or disease is selected from respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated diseases; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs such as polycythemia vera, essential thrombocythemia or myelofibrosis); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors; more in particular wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.

[0765] In particular the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis. The invention also encompasses the use of a pharmaceutical composition of the invention for the manufacture of a medicament for treating these diseases.

[0766] The present invention also provides pharmaceutical compositions which comprise, as an active ingredient, at least a compound of formula (I) or a pharmaceutically acceptable salt thereof in association with a pharmaceutically acceptable excipient such as a carrier or diluent. The active ingredient may comprise 0.001% to 99% by weight, preferably 0.01% to 90% by weight, of the composition depending upon the nature of the formulation and whether further dilution is to be made prior to application. Preferably the compositions are made up in a form suitable for oral, inhalation, topical, nasal, rectal, percutaneous or injectable administration.

[0767] Pharmaceutical compositions suitable for the delivery of compounds of the invention and methods for their preparation will be readily apparent to those skilled in the art. Such compositions and methods for their preparation can be found, for example, in Remington: The Science and Practice of Pharmacy, 21 st Edition, Lippincott Williams & Wilkins, Philadelphia, Pa., 2001.

[0768] The pharmaceutically acceptable excipients which are admixed with the active compound or salts of such compound, to form the compositions of this invention are well-known per se and the actual excipients used depend inter alia on the intended method of administering the compositions. Examples, without limitation, of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

[0769] Additional suitable carriers for formulations of the compounds of the present invention can be found in Remington: The Science and Practice of Pharmacy, 21 st Edition, Lippincott Williams & Wilkins, Philadelphia, Pa., 2001.

i) Oral Administration

[0770] The compounds of the invention may be administered orally (peroral administration; *per os* (latin)). Oral administration involve swallowing, so that the compound is absorbed from the gut and delivered to the liver via the portal circulation (hepatic first pass metabolism) and finally enters the gastrointestinal (GI) tract.

[0771] Compositions for oral administration may take the form of tablets, retard tablets, sublingual tablets, capsules, inhalation aerosols, inhalation solutions, dry powder inhalation, or liquid preparations, such as mixtures, solutions, elixirs, syrups or suspensions, all containing the compound of the invention; such preparations may be made by methods well-known in the art. The active ingredient may also be presented as a bolus, electuary or paste.

[0772] Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include magnesium stearate, talc, gelatine, acacia, stearic acid, starch, lactose and sucrose.

[0773] A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, lubricating, surface active or dispersing agent.

[0774] Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein.

[0775] For tablet dosage forms, depending on dose, the drug may make up from 1 wt% to 80 wt% of the dosage form, more typically from 5 wt% to 60 wt% of the dosage form. In addition to the drug, tablets generally contain a disintegrant. Examples of disintegrants include sodium starch glycolate, sodium carboxymethyl cellulose, calcium carboxymethyl cellulose, croscarmellose sodium, crospovidone, polyvinylpyrrolidone, methyl cellulose, microcrystalline cellulose, lower alkyl- substituted hydroxypropyl cellulose, starch, pregelatinized starch and sodium alginate. Generally, the disintegrant will comprise from 1 wt% to 25 wt%, preferably from 5 wt% to 20 wt% of the dosage form.

[0776] Binders are generally used to impart cohesive qualities to a tablet formulation. Suitable binders include microcrystalline cellulose, gelatin, sugars, polyethylene glycol, natural and synthetic gums, polyvinylpyrrolidone, pregelatinized starch, hydroxypropyl cellulose and hydroxypropyl methylcellulose. Tablets may also contain diluents, such as lactose (monohydrate, spray-dried monohydrate, anhydrous and the like), mannitol, xylitol, dextrose, sucrose, sorbitol, microcrystalline cellulose, starch and dibasic calcium phosphate dihydrate. Tablets may also optionally include surface active agents, such as sodium lauryl sulfate and polysorbate 80, and glidants such as silicon dioxide and talc. When present, surface active agents are typically in amounts of from 0.2 wt% to 5 wt% of the tablet, and glidants typically from 0.2 wt% to 1 wt% of the tablet.

[0777] Tablets also generally contain lubricants such as magnesium stearate, calcium stearate, zinc stearate, sodium stearyl fumarate, and mixtures of magnesium stearate with sodium lauryl sulphate. Lubricants generally are present in amounts from 0.25 wt% to 10 wt%, preferably from 0.5 wt% to 3 wt% of the tablet. Other conventional ingredients include anti-oxidants, colorants, flavoring agents, preservatives and taste-masking agents.

[0778] Exemplary tablets contain up to about 80 wt% drug, from about 10 wt% to about 90 wt% binder, from about 0 wt% to about 85 wt% diluent, from about 2 wt% to about 10 wt% disintegrant, and from about 0.25 wt% to about 10 wt% lubricant. Tablet blends may be compressed directly or by roller to form tablets. Tablet blends or portions of blends may alternatively be wet-, dry-, or melt-granulated, melt congealed, or extruded before tableting. The final formulation may include one or more layers and may be coated or uncoated; or encapsulated.

[0779] The formulation of tablets is discussed in detail in "Pharmaceutical Dosage Forms: Tablets, Vol. 1 ", by H. Lieberman and L. Lachman, Marcel Dekker, N.Y., 1980.

[0780] Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example using the aforementioned carriers in a hard gelatine capsule. Where the composition is in the form of a soft gelatine capsule any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous gums, celluloses, silicates or oils, and are incorporated in a soft gelatine capsule.

[0781] Solid formulations for oral administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

[0782] Suitable modified release formulations are described in U.S. Patent No. 6,106,864. Details of other suitable release technologies such as high energy dispersions and osmotic and coated particles can be found in Verma et al, Pharmaceutical Technology On-line, 25(2), 1-14 (2001). The use of chewing gum to achieve controlled release is described in WO 00/35298. The disclosures of these references are incorporated herein by reference in their entireties.

[0783] Liquid formulations include suspensions, solutions, syrups and elixirs. Such formulations may be used as fillers in soft or hard capsules and typically include a carrier, for example, water, ethanol, polyethylene glycol, propylene glycol, methylcellulose, or a suitable oil, and one or more emulsifying agents and/or suspending agents. The solutions may be aqueous solutions of a soluble salt or other derivative of the active compound in association with, for example, sucrose to form a syrup. The suspensions may comprise an insoluble active compound of the invention or a pharmaceutically acceptable salt thereof in association with water, together with a suspending agent or flavouring agent. Liquid formulations may also be prepared by the reconstitution of a solid, for example, from a sachet.

ii) Oral mucosal administration

[0784] The compounds of the invention can also be administered via the oral mucosal. Within the oral mucosal cavity, delivery of drugs is classified into three categories: (a) sublingual delivery, which is systemic delivery of drugs through the mucosal membranes lining the floor of the mouth, (b) buccal delivery, which is drug administration through the mucosal membranes lining the cheeks (buccal mucosa), and (c) local delivery, which is drug delivery into the oral cavity.

[0785] Pharmaceutical products to be administered via the oral mucosal can be designed using mucoadhesive, quick dissolve tablets and solid lozenge formulations, which are formulated with one or more mucoadhesive (bioadhesive) polymers (such as hydroxy propyl cellulose, polyvinyl pyrrolidone, sodium carboxymethyl cellulose, hydroxy propyl methyl cellulose, hydroxy ethyl cellulose, polyvinyl alcohol, polyisobutylene or polyisoprene); and oral mucosal permeation enhancers (such as butanol, butyric acid, propranolol, sodium lauryl sulphate and others)

iii) Inhaled administration

[0786] The compounds of the invention can also be administered by inhalation, typically in the form of a dry powder (either alone, as a mixture, for example, in a dry blend with lactose, or as a mixed component particle, for example, mixed with phospholipids, such as phosphatidylcholine) from a dry powder inhaler or as an aerosol spray from a pressurized container, pump, spray, atomizer (preferably an atomizer using electrohydrodynamics to produce a fine mist), or nebulizer, with or without the use of a suitable propellant, such as 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-heptafluoropropane. For intranasal use, the powder may include a bioadhesive agent, for example, chitosan or cyclodextrin.

[0787] Dry powder compositions for topical delivery to the lung by inhalation may, for example, be presented in capsules and cartridges of for example gelatine or blisters of for example laminated aluminium foil, for use in an inhaler or insufflator. Formulations generally contain a powder mix for inhalation of the compound of the invention and a suitable powder base (carrier substance) such as lactose or starch. Use of lactose is preferred. Each capsule or cartridge may generally contain between 0.001-50 mg, more preferably 0.01-5 mg of active ingredient or the equivalent amount of a pharmaceutically acceptable salt thereof. Alternatively, the active ingredient (s) may be presented without excipients.

[0788] Packaging of the formulation may be suitable for unit dose or multi-dose delivery. In the case of multi-dose delivery, the formulation can be pre-metered or metered in use. Dry powder inhalers are thus classified into three groups: (a) single dose, (b) multiple unit dose and (c) multi dose devices.

[0789] For inhalers of the first type, single doses have been weighed by the manufacturer into small containers, which are mostly hard gelatine capsules. A capsule has to be taken from a separate box or container and inserted into a receptacle area of the inhaler. Next, the capsule has to be opened or perforated with pins or cutting blades in order to allow part of the inspiratory air stream to pass through the capsule for powder entrainment or to discharge the powder from the capsule through these perforations by means of centrifugal force during inhalation. After inhalation, the emptied capsule has to be removed from the inhaler again. Mostly, disassembling of the inhaler is necessary for inserting and removing the capsule, which is an operation that can be difficult and burdensome for some patients.

[0790] Other drawbacks related to the use of hard gelatine capsules for inhalation powders are (a) poor protection against moisture uptake from the ambient air, (b) problems with opening or perforation after the capsules have been exposed previously to extreme relative humidity, which causes fragmentation or indenture, and (c) possible inhalation of capsule fragments. Moreover, for a number of capsule inhalers, incomplete expulsion has been reported (e. g. Nielsen et al, 1997).

[0791] Some capsule inhalers have a magazine from which individual capsules can be transferred to a receiving chamber, in which perforation and emptying takes place, as described in WO 92/03175. Other capsule inhalers have revolving magazines with capsule chambers that can be brought in line with the air conduit for dose discharge (e. g. WO91/02558 and GB 2242134). They comprise the type of multiple unit dose inhalers together with blister inhalers,

which have a limited number of unit doses in supply on a disk or on a strip.

[0792] Blister inhalers provide better moisture protection of the medicament than capsule inhalers. Access to the powder is obtained by perforating the cover as well as the blister foil, or by peeling off the cover foil. When a blister strip is used instead of a disk, the number of doses can be increased, but it is inconvenient for the patient to replace an empty strip. Therefore, such devices are often disposable with the incorporated dose system, including the technique used to transport the strip and open the blister pockets.

[0793] Multi-dose inhalers do not contain pre-measured quantities of the powder formulation. They consist of a relatively large container and a dose measuring principle that has to be operated by the patient. The container bears multiple doses that are isolated individually from the bulk of powder by volumetric displacement. Various dose measuring principles exist, including rotatable membranes (Ex. EP0069715) or disks (Ex. GB 2041763; EP 0424790; DE 4239402 and EP 0674533), rotatable cylinders (Ex. EP 0166294; GB 2165159 and WO 92/09322) and rotatable frustums (Ex. WO 92/00771), all having cavities which have to be filled with powder from the container. Other multi dose devices have measuring slides (Ex. US 5201308 and WO 97/00703) or measuring plungers with a local or circumferential recess to displace a certain volume of powder from the container to a delivery chamber or an air conduit (Ex. EP 0505321, WO 92/04068 and WO 92/04928), or measuring slides such as the Genuair® (formerly known as Novolizer SD2FL), which is described the following patent applications Nos: WO97/000703, WO03/000325 and WO2006/008027.

[0794] Reproducible dose measuring is one of the major concerns for multi dose inhaler devices.

[0795] The powder formulation has to exhibit good and stable flow properties, because filling of the dose measuring cups or cavities is mostly under the influence of the force of gravity.

[0796] For reloaded single dose and multiple unit dose inhalers, the dose measuring accuracy and reproducibility can be guaranteed by the manufacturer. Multi dose inhalers on the other hand, can contain a much higher number of doses, whereas the number of handlings to prime a dose is generally lower.

[0797] Because the inspiratory air stream in multi-dose devices is often straight across the dose measuring cavity, and because the massive and rigid dose measuring systems of multi dose inhalers can not be agitated by this inspiratory air stream, the powder mass is simply entrained from the cavity and little de-agglomeration is obtained during discharge.

[0798] Consequently, separate disintegration means are necessary. However in practice, they are not always part of the inhaler design. Because of the high number of doses in multi-dose devices, powder adhesion onto the inner walls of the air conduits and the deagglomeration means must be minimized and/or regular cleaning of these parts must be possible, without affecting the residual doses in the device. Some multi dose inhalers have disposable drug containers that can be replaced after the prescribed number of doses has been taken (Ex. WO 97/000703). For such semi-permanent multi dose inhalers with disposable drug containers, the requirements to prevent drug accumulation are even more strict.

[0799] Apart from applications through dry powder inhalers the compositions of the invention can be administered in aerosols which operate via propellant gases or by means of so-called atomisers, via which solutions of pharmacologically-active substances can be sprayed under high pressure so that a mist of inhalable particles results. The advantage of these atomisers is that the use of propellant gases can be completely dispensed with. Such atomiser is the Respimat® which is described, for example, in PCT Patent Applications Nos. WO 91/14468 and WO 97/12687, reference here is being made to the contents thereof.

[0800] Spray compositions for topical delivery to the lung by inhalation may for example be formulated as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, such as a metered dose inhaler, with the use of a suitable liquefied propellant. Aerosol compositions suitable for inhalation can be either a suspension or a solution and generally contain the active ingredient (s) and a suitable propellant such as a fluorocarbon or hydrogen-containing chlorofluorocarbon or mixtures thereof, particularly hydrofluoroalkanes, e. g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetra-fluoroethane, especially 1,1, 1, 2-tetrafluoroethane, 1,1, 1,2, 3,3, 3-heptafluoro-n-propane or a mixture thereof. Carbon dioxide or other suitable gas may also be used as propellant.

[0801] The aerosol composition may be excipient free or may optionally contain additional formulation excipients well known in the art such as surfactants (eg oleic acid or lecithin) and cosolvents (eg ethanol). Pressurised formulations will generally be retained in a canister (eg an aluminium canister) closed with a valve (eg a metering valve) and fitted into an actuator provided with a mouthpiece.

[0802] Medicaments for administration by inhalation desirably have a controlled particle size. The optimum particle size for inhalation into the bronchial system is usually 1-10 µm, preferably 2-5 µm. Particles having a size above 20 µm are generally too large when inhaled to reach the small airways. To achieve these particle sizes the particles of the active ingredient as produced may be size reduced by conventional means eg by micronisation. The desired fraction may be separated out by air classification or sieving. Preferably, the particles will be crystalline.

[0803] Achieving high dose reproducibility with micronised powders is difficult because of their poor flowability and extreme agglomeration tendency. To improve the efficiency of dry powder compositions, the particles should be large while in the inhaler, but small when discharged into the respiratory tract. Thus, an excipient such as lactose or glucose is generally employed. The particle size of the excipient will usually be much greater than the inhaled medicament within the present invention. When the excipient is lactose it will typically be present as milled lactose, preferably crystalline

alpha lactose monohydrate.

[0804] Pressurized aerosol compositions will generally be filled into canisters fitted with a valve, especially a metering valve. Canisters may optionally be coated with a plastics material e. g. a fluorocarbon polymer as described in WO96/32150. Canisters will be fitted into an actuator adapted for buccal delivery.

iv) Nasal mucosal administration

[0805] The compounds of the invention may also be administered via the nasal mucosal. Typical compositions for nasal mucosa administration are typically applied by a metering, atomizing spray pump and are in the form of a solution or suspension in an inert vehicle such as water optionally in combination with conventional excipients such as buffers, anti-microbials, tonicity modifying agents and viscosity modifying agents.

v) Parenteral Administration

[0806] The compounds of the invention may also be administered directly into the blood stream, into muscle, or into an internal organ. Suitable means for parenteral administration include intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular and subcutaneous. Suitable devices for parenteral administration include needle (including microneedle) injectors, needle-free injectors and infusion techniques.

[0807] Parenteral formulations are typically aqueous solutions which may contain excipients such as salts, carbohydrates and buffering agents (preferably to a pH of from 3 to 9), but, for some applications, they may be more suitably formulated as a sterile nonaqueous solution or as a dried form to be used in conjunction with a suitable vehicle such as sterile, pyrogen-free water.

[0808] The preparation of parenteral formulations under sterile conditions, for example, by lyophilization, may readily be accomplished using standard pharmaceutical techniques well known to those skilled in the art. The solubility of compounds of the invention used in the preparation of parenteral solutions may be increased by the use of appropriate formulation techniques, such as the incorporation of solubility-enhancing agents.

[0809] Formulations for parenteral administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release. Thus compounds of the invention may be formulated as a solid, semi-solid, or thixotropic liquid for administration as an implanted depot providing modified release of the active compound. Examples of such formulations include drug-coated stents and PGLA microspheres.

vi) Topical Administration

[0810] The compounds of the invention may also be administered topically to the skin or mucosa, that is, dermally or transdermally. Typical formulations for this purpose include gels, hydrogels, lotions, solutions, creams, ointments, dusting powders, dressings, foams, films, skin patches, wafers, implants, sponges, fibers, bandages and microemulsions. Liposomes may also be used. Typical carriers include alcohol, water, mineral oil, liquid petrolatum, white petrolatum, glycerin, polyethylene glycol and propylene glycol. Penetration enhancers may be incorporated; see, for example, J Pharm Sci, 88 (10), 955-958 by Finnin and Morgan (October 1999). Other means of topical administration include delivery by electroporation, iontophoresis, phonophoresis, sonophoresis and microneedle or needle-free injection.

[0811] Formulations for topical administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

vii) Rectal/Intravaginal Administration

[0812] Compounds of the invention may be administered rectally or vaginally, for example, in the form of a suppository, pessary, or enema. Cocoa butter is a traditional suppository base, but various alternatives may be used as appropriate. Formulations for rectal/vaginal administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted and programmed release.

viii) Ocular Administration

[0813] Compounds of the invention may also be administered directly to the eye or ear, typically in the form of drops of a micronized suspension or solution in isotonic, pH-adjusted, sterile saline. Other formulations suitable for ocular and aural administration include ointments, biodegradable {e.g. absorbable gel sponges, collagen) and nonbiodegradable (e.g. silicone) implants, wafers, lenses and particulate or vesicular systems, such as niosomes or liposomes. A polymer such as crossed-linked polyacrylic acid, polyvinylalcohol, hyaluronic acid, a cellulosic polymer, for example, hydroxy-

propylmethylcellulose, hydroxyethylcellulose, or methyl cellulose, or a heteropolysaccharide polymer, for example, gelatin gum, may be incorporated together with a preservative, such as benzalkonium chloride. Such formulations may also be delivered by iontophoresis.

[0814] Formulations for ocular/aural administration may be formulated to be immediate and/or modified release. Modified release formulations include delayed-, sustained-, pulsed-, controlled-, targeted, or programmed release.

ix) Other Technologies

[0815] Compounds of the invention may be combined with soluble macromolecular entities, such as cyclodextrin and suitable derivatives thereof or polyethylene glycol-containing polymers, in order to improve their solubility, dissolution rate, taste-masking, bioavailability and/or stability for use in any of the aforementioned modes of administration.

[0816] The amount of the active compound administered will be dependent on the subject being treated, the severity of the disorder or condition, the rate of administration, the disposition of the compound and the discretion of the prescribing physician. However, an effective dosage is typically in the range of 0.01-3000 mg, more preferably 0.5-1000 mg of active ingredient or the equivalent amount of a pharmaceutically acceptable salt thereof per day. Daily dosage may be administered in one or more treatments, preferably from 1 to 4 treatments, per day.

[0817] Preferably, the pharmaceutical compositions of the invention are made up in a form suitable for oral, inhalation or topical administration, being particularly preferred oral or inhalation administration.

[0818] The pharmaceutical formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer a single dose.

[0819] The amount of each active which is required to achieve a therapeutic effect will, of course, vary with the particular active, the route of administration, the subject under treatment, and the particular disorder or disease being treated.

[0820] The following preparations forms are cited as formulation examples:

Formulation Examples

Formulation Example 1 (Oral suspension)

[0821]

Ingredient	Amount
Active Compound	3 mg
Citric acid	0,5 g
Sodium chloride	2,0 g
Methyl paraben	0,1 g
Granulated sugar	25 g
Sorbitol (70% solution)	11 g
Veegum K	1,0 g
Flavoring	0,02 g
Dye	0,5 mg
Distilled water	q.s. to 100 mL

Formulation Example 2 (Hard gelatine capsule for oral administration)

[0822]

Ingredient	Amount
Active Compound	1 mg
Lactose	150 mg
Magnesium stearate	3 mg

Formulation Example 3 (Gelatin cartridge for inhalation)

[0823]

Ingredient	Amount
Active Compound (micronized)	0,2 mg
Lactose	25 mg

Formulation Example 4 (Formulation for inhalation with a DPI)

[0824]

Ingredient	Amount
Active Compound (micronized)	15 mg
Lactose	3000 mg

Formulation Example 5 (Formulation for a MDI)

[0825]


Ingredient	Amount
Active Compound (micronized)	10 g
1,1,1,2,3,3,3-heptafluoro-n-propane	q.s. to 200 mL

[0826] In all the formulation examples, active compound is (S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one.

[0827] Modifications, which do not affect, alter, change or modify the essential aspects of the compounds, combinations or pharmaceutical compositions described, are included within the scope of the present invention.

Claims

1. A compound of formula (I), or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof:



Formula (I)

wherein,

X represents a nitrogen atom or a $-CR_6$ group;

n represents 0, 1, 2 or 3;

R_a and R_b each independently represent a hydrogen atom, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group or a linear or branched C_1 - C_4 alkyl group;

R_1 represents a hydrogen atom, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_3 - C_{10} cycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a monocyclic or bicyclic C_6 - C_{14} aryl group, a 5- to 14- membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N, or a 5- to 14- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N,

wherein the cycloalkyl, cycloalkenyl, aryl, heteroaryl, and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{1-3}CN$ group, a $-(CH_2)_{0-3}OR_8$ group, a $-(CH_2)_{0-3}NR_7R_8$ group,

a $-C(O)-(CH_2)_{1-3}CN$ group, a $-C(O)-(CH_2)_{0-3}R_8$ group, a $-C(O)-(CH_2)_{0-3}NR_7R_8$ group, a $-S(O)_2(CH_2)_{0-3}R_8$ group, a $-S(O)_2(CH_2)_{0-3}NR_7R_8$ group or a $-(CH_2)_{0-3}(phenyl)-OR_8$ group;

R_2 and R_3 each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_7 cycloalkyl group, a $-NR'R''$ group, or a linear or branched C_1 - C_6 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_7 cycloalkyl group;

R_4 represents a hydrogen atom, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_7 cycloalkyl group, a $-(CH_2)_{1-4}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group, a C_3 - C_4 cycloalkyl group, a $-C(O)-(CH_2)_{0-3}R$ group or a $-C(O)-(CH_2)_{0-3}NR'R''$ group;

R_6 represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C_1 - C_4 alkoxy group; a C_1 - C_4 haloalkyl group; a linear or branched C_1 - C_4 hydroxyalkyl group; a C_3 - C_7 cycloalkyl group; a $-(CH_2)_{0-3}NR'R''$ group; a $-(CH_2)_{1-3}O(C_1-C_4 \text{ alkyl group})$; a $-(CH_2)_{0-3}OC(O)-(C_1-C_4 \text{ alkyl group})$; a $-(CH_2)_{0-3}C(O)O-(C_1-C_4 \text{ alkyl group})$; a $-C(O)-(CH_2)_{0-3}NR'R''$ group; a $-(CH_2)_{0-3}C(O)OH$ group; a $-(CH_2)_{0-3}$ -(5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N); a $-(CH_2)_{0-3}$ -(5- to 14- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N); or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group;


wherein the heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1 - C_4 alkyl group or a C_1 - C_4 haloalkyl group,

R_7 and R_8 each independently represent a hydrogen atom, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group;

R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group.


R_5 represents a group selected from:

i) a group of formula (IIa)



formula (IIa)

ii) a group of formula (IIb)



formula (IIb)

and
iii) a group of formula (IIc)



formula (IIc)

wherein

Y represents a linker selected from a -NR'- group, -O- or -S-; wherein R' is as defined above;

(*) represents where R₅ is bonded to the carbon atom attached to R₄ and to the pyrrolotriazinone group;

W₁ represents a -CR₁₁ group and W₂ represents a nitrogen atom, or W₁ represents a nitrogen atom and W₂ represents a -CR₁₂ group;

G₁ represents a -CR₁₄ group and G₂ represents a nitrogen atom, or G₁ represents a nitrogen atom and G₂ represents a -CR₁₅ group, or G₁ represents a -CR₁₄ group and G₂ represents a -CR₁₅ group;

G₃ represents a nitrogen atom, or a -CR₁₆ group;


R₉, R₁₀, and R₁₂, R₁₃, R₁₄, and R₁₆ each independently represents a hydrogen atom; a halogen atom; a C₁-C₄ alkoxy; a C₁-C₄ haloalkyl group; a C₁-C₄ hydroxyalkyl group; a C₃-C₄ cycloalkyl group; a -(CH₂)₀₋₃CN group; a -C(O)-(CH₂)₁₋₃-CN group; a -C(O)-(CH₂)₀₋₃-R' group; a -C(O)-(CH₂)₀₋₃-NR'R'';

a (CH₂)₀₋₃NR'R'' group; a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group;

wherein R' and R'' are as defined above;


R₁₇ represents a group selected from

a) a group of formula (IIIa),



formula (IIIa)

b) a group of formula (IIIb),





formula (IIIb)

c) a group of formula (IIIc), and

d) a group of formula (IIId),

(continued)



wherein

G₄ represents a nitrogen atom or a -CR₂₂ group;

G₅ and G₆ each independently represents a nitrogen atom or a carbon atom, wherein when one of G₅ and G₆ represents a nitrogen atom the remaining represents a carbon atom;

G₇ represents a -NH group or a -CH group;

G₈ represents a nitrogen atom or a -CR₂₃ group;

G₉ represents a nitrogen atom or a -CR₂₄ group;

G₁₀ represents a nitrogen atom or a -CR₂₅ group;

G₁₁ represents a nitrogen atom or a -CR₂₆ group;

G₁₂ represents a nitrogen atom or a -CR₂₇ group;

G₁₃ represents a nitrogen atom or a -CR₂₈ group;

G₁₄ and G₁₅ each independently represents a nitrogen atom or a -carbon atom, wherein when one of G₁₄ and G₁₅ represents a nitrogen atom the remaining represents a carbon atom;

G₁₆ represents a -NH group or a -CH group;

G₁₇ represents a nitrogen atom or a -CR₂₉ group;

G₁₈ represents a nitrogen atom or a -CR₃₀ group;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉, and R₃₀ each independently represent a hydrogen atom, a halogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R" are as defined above; and wherein Y is as defined above;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.

2. A compound according to claim 1, wherein

R_a and R_b each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

n represents 0, 1 or 2;

R₁ represents a hydrogen atom, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a 5- to 10- membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, a tetrahydropyranyl group or a morpholinyl group;

wherein the cycloalkyl, phenyl, heteroaryl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydropyranyl or morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a linear or branched C₁-C₄ alkyl group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃OR₈ group, a -(CH₂)₀₋₃NR₇R₈ group, a -C(O)-(CH₂)₀₋₃-R₈ group, a -C(O)-(CH₂)₀₋₃-NR₇R₈ group or a -(CH₂)₀₋₃(phenyl)-OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a C₁-C₄ alkyl group;

R₂ represents a hydrogen atom, a halogen atom, a hydroxyl group, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted

or substituted by a C₁-C₂ alkoxy group;


R₃ represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group;

R₄ represents a hydrogen atom, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C₁-C₃ alkoxy group; a C₁-C₃ haloalkyl group; a linear or branched C₁-C₄ hydroxyalkyl group; a C₃-C₄ cycloalkyl group; a -(CH₂)₀₋₃NR'R" group; a -(CH₂)₁₋₃O(C₁-C₄ alkyl group); a -(CH₂)₀₋₃OC(O)-(C₁-C₄ alkyl group); a -(CH₂)₀₋₃C(O)O-(C₁-C₄ alkyl group); a -C(O)-(CH₂)₀₋₃-NR'R" group; a -(CH₂)₀₋₃C(O)OH group; a -(CH₂)₀₋₃-(5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N); a -(CH₂)₀₋₃-(5- to 14- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N);

or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group; wherein the heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C₁-C₄ alkyl group or a C₁-C₄ haloalkyl group,

R₅ represents a moiety of formula (IIa-2), (IIb-1), (IIb-2), (IIb-3), (IIIa-1), (IIIa-3), (IIIa-5) or (IIIa-14):




formula (IIa-2)

formula (IIb-1)

formula (IIb-2)

formula (IIb-3)



formula (IIIa-1)

formula (IIIa-3)

formula (IIIa-5)

formula (IIIa-14)

wherein:

• R₉, R₁₀, and R₁₂ each independently represent a hydrogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R" group, a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;

• R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a halogen atom, a C₁-C₄ haloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R",

• a -(CH₂)₀₋₃NR'R" group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;

• R₁₈, R₂₂, R₂₃ and R₂₄ each independently represent a hydrogen atom, a halogen atom, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R", a -(CH₂)₀₋₃NR'R" group, or a linear or branched C₁-C₄ alkyl group; wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group;

• Y represents a -NR'- group, -O- or -S-; wherein R' represents hydrogen or a linear or branched C₁-C₄ alkyl group; or in the case that Y represents a -NR'-group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7- membered, saturated N-containing heterocyclyl

group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a $-\text{CHF}_2$ group or a $-\text{CF}_3$ group.

3. A compound according to claim 1, wherein

5 X represents a nitrogen atom or a $-\text{CR}_6$ group;

n represents 0, 1, 2 or 3;

R_a and R_b each independently represent a hydrogen atom, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group or a linear or branched C_1 - C_4 alkyl group;

10 R_1 represents a linear or branched C_1 - C_4 alkyl group, a C_3 - C_{10} cycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a monocyclic or bicyclic C_6 - C_{14} aryl group, a 5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N, or a 5- to 14-membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N,

15 wherein the cycloalkyl, cycloalkenyl, aryl, heteroaryl, and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(\text{CH}_2)_{1-3}\text{CN}$ group, a $-(\text{CH}_2)_{0-3}\text{OR}_8$ group, a $-(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{1-3}\text{CN}$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}_8$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group, a $-\text{S}(\text{O})_2(\text{CH}_2)_{0-3}\text{R}_8$ group or a $-\text{S}(\text{O})_2(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group;

20 R_2 and R_3 each independently represent a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_7 cycloalkyl group, a $-\text{NR}'\text{R}''$ group, or a linear or branched C_1 - C_6 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_7 cycloalkyl group;

25 R_4 represents a hydrogen atom, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_7 cycloalkyl group, a $-(\text{CH}_2)_{1-4}\text{NR}'\text{R}''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group, a C_3 - C_4 cycloalkyl group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}$ group or a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ group;

30 R_6 represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_7 cycloalkyl group, a $-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group;

R_7 and R_8 each independently represent a hydrogen atom, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group;

35 R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group.

R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} and R_{16} each independently represent a hydrogen atom, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(\text{CH}_2)_{0-3}\text{CN}$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{1-3}\text{CN}$ group,

40 a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}'$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$, a $-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group; wherein R' and R'' are as defined above;

45 R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{28} , R_{29} , and R_{30} each independently represent a hydrogen atom, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(\text{CH}_2)_{0-3}\text{CN}$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{1-3}\text{CN}$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}'$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$, a $-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C_1 - C_4 alkoxy group, a cyano group or a C_3 - C_4 cycloalkyl group; wherein R' and R'' are as defined above; and wherein Y is as defined above.

4. A compound according to claim 3, wherein

50 R_a and R_b each independently represent a hydrogen atom or a linear or branched C_1 - C_3 alkyl group;

n represents 0, 1 or 2;

R_1 represents a linear or branched C_1 - C_4 alkyl group, a C_3 - C_7 cycloalkyl group, a phenyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing containing one, two or three heteroatoms selected from O, S and N, a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, a tetrahydropyranyl group or a morpholinyl group;

55 wherein the cycloalkyl, phenyl, heteroaryl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydropyranyl or morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(\text{CH}_2)_{0-3}\text{OR}_8$ group, a $-(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group, a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}_8$ group or a $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}_7\text{R}_8$ group; wherein

R_7 and R_8 each independently represent a hydrogen atom or a C_1 - C_4 alkyl group;


R_2 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_2 alkoxy group;

R_3 represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_2 alkoxy group;

R_4 represents a hydrogen atom, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, or a linear or branched C_1 - C_3 alkyl group;

R_6 represents a hydrogen atom, a halogen atom, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_2 alkoxy group;

R_5 represents a moiety of formula (II-a2), (IIb-1), (IIb-2) or (IIIa-1):



wherein:

• R_9 , R_{10} , and R_{12} each independently represent a hydrogen atom, a $-(CH_2)_{0-3}CN$ group, a $-C(O)-(CH_2)_{1-3}CN$ group, a $-C(O)-(CH_2)_{0-3}R'$ group, a $-C(O)-(CH_2)_{0-3}NR'R''$, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group;

• R_{13} , R_{14} , R_{15} and R_{16} each independently represent a hydrogen atom, a $-(CH_2)_{0-3}CN$ group, a $-C(O)-(CH_2)_{0-3}NR'R''$ or a $-(CH_2)_{0-3}NR'R''$ group; wherein R' and R'' each independently represent a hydrogen atom or a linear or branched C_1 - C_4 alkyl group;

• R_{18} and R_{23} each independently represent a hydrogen atom, a $-(CH_2)_{0-3}CN$ group, a $-C(O)-(CH_2)_{1-3}CN$ group, a $-C(O)-(CH_2)_{0-3}R'$ group, a $-C(O)-(CH_2)_{0-3}NR'R''$, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C_1 - C_4 alkoxy group or a linear or branched C_1 - C_4 alkyl group

• Y represents a $-NR'$ - group, $-O-$ or $-S-$; wherein R' represents hydrogen or a linear or branched C_1 - C_4 alkyl group; or in the case that Y represents a $-NR'$ -group, R_4 together with the $-NR'$ - group and the carbon atom to which both R_4 and the $-NR'$ - group are bonded form a 4- to 7- membered, saturated N-containing heterocycl group.

5. A compound according to claim 1, wherein R_1 represents a hydrogen atom, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_3 - C_{10} cycloalkyl group, a C_3 - C_{10} cycloalkenyl group, a monocyclic or bicyclic C_6 - C_{14} aryl group, a 5- to 14-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N, or a 5- to 14- membered monocyclic or bicyclic heterocycl group containing at least one heteroatom selected from O, S and N,

wherein the cycloalkyl, cycloalkenyl, aryl, heteroaryl, and heterocycl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{1-3}CN$ group, a $-(CH_2)_{0-3}OR_8$ group, a $-(CH_2)_{0-3}NR_7R_8$ group,

a $-C(O)-(CH_2)_{1-3}CN$ group, a $-C(O)-(CH_2)_{0-3}R_8$ group, a $-C(O)-(CH_2)_{0-3}NR_7R_8$ group, a $-S(O)_2(CH_2)_{0-3}R_8$ group, a $-S(O)_2(CH_2)_{0-3}NR_7R_8$ group or

a $-(CH_2)_{0-3}(\text{phenyl})-OR_8$ group;

and

wherein R_1 preferably represents a hydrogen atom, C_1 - C_3 alkyl group, a C_1 - C_3 haloalkyl group, a C_3 - C_7 cycloalkyl group, a phenyl group, a naphthyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, or a 5- to 10-membered monocyclic or bicyclic heterocyclyl group containing one, two or three heteroatoms selected from O, S and N,

wherein the cycloalkyl, phenyl, naphthyl, heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{0-3}OR_8$ group, a $-(CH_2)_{0-3}NR_7R_8$ group, a $-C(O)-(CH_2)_{0-3}R_8$ group, a $-C(O)-(CH_2)_{0-3}NR_7R_8$ group or a $-(CH_2)_{0-3}(phenyl)-OR_8$ group; wherein R_7 and R_8 are as defined in claim 1.

6. A compound according to any one of claims 1, 3 or 5, wherein:

(a) R_1 represents a C_1 - C_3 alkyl group, a C_3 - C_7 cycloalkyl group, a phenyl group, a naphthyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, or a 5- to 10-membered monocyclic or bicyclic heterocyclyl group containing one, two or three heteroatoms selected from O, S and N,

wherein the cycloalkyl, phenyl, naphthyl, heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{0-3}OR_8$ group, a $-(CH_2)_{0-3}NR_7R_8$ group, a $-C(O)-(CH_2)_{0-3}R_8$ group or a $-C(O)-(CH_2)_{0-3}NR_7R_8$ group; wherein R_7 and R_8 are as defined in claim 1;

and

wherein R_1 preferably represents a C_3 - C_7 cycloalkyl group, a phenyl group, a 5- to 10-membered monocyclic or bicyclic heteroaryl group containing one, two or three heteroatoms selected from O, S and N, a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, a tetrahydropyranyl group or a morpholinyl group, wherein the cycloalkyl, phenyl, heteroaryl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydropyranyl or morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C_1 - C_4 alkyl group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{0-3}OR_8$ group, a $-(CH_2)_{0-3}NR_7R_8$ group,

a $-C(O)-(CH_2)_{0-3}R_8$ group or a $-C(O)-(CH_2)_{0-3}NR_7R_8$ group; wherein R_7 and R_8 each independently represent a hydrogen atom or a C_1 - C_4 alkyl group;

and

wherein R_1 more preferably represents a phenyl group or a pyridinyl group, which phenyl or pyridinyl is unsubstituted or substituted by one, two or three substituents selected from a halogen atom, a linear or branched C_1 - C_3 alkyl group or a $-(CH_2)_{0-3}OCH_3$ group; and wherein preferably said phenyl and pyridinyl groups are directly bonded to the pyrrolotriazinone group; and/or

(b) R_2 represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_3 alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C_1 - C_3 alkyl group;

and

wherein R_2 preferably represents a hydrogen atom, a halogen atom, a hydroxyl group, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_2 alkoxy group; and/or

(c) R_3 represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1 - C_4 alkoxy group, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{0-3}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_3 alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxy group, or a linear or branched C_1 - C_3 alkyl group;

and

wherein R_3 preferably represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C_1 - C_3 alkoxy group, a C_1 - C_3 haloalkyl group, a C_3 - C_4 cycloalkyl group, a $-NH_2$ group, a $-N(CH_3)H$ group, a $-N(CH_3)_2$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or substituted by a C_1 - C_2 alkoxy group; and/or

(d) R_4 represents a hydrogen atom, a C_1 - C_4 haloalkyl group, a C_1 - C_4 hydroxyalkyl group, a C_3 - C_4 cycloalkyl group, a $-(CH_2)_{1-4}NR'R''$ group, or a linear or branched C_1 - C_4 alkyl group, which alkyl group is unsubstituted or

substituted by a C₁-C₃ alkoxy group, a -C(O)-(CH₂)₀₋₃-R group or a -C(O)-(CH₂)₀₋₃-NR'R'' group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group;

and

wherein R₄ preferably represents a hydrogen atom, a C₁-C₃ haloalkyl group, a C₁-C₃ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, or a linear or branched C₁-C₃ alkyl group;

and/or

(e) R₆ represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a linear or branched C₁-C₄ hydroxyalkyl group; a C₃-C₇ cycloalkyl group; a -(CH₂)₀₋₃NR'R'' group; a -(CH₂)₁₋₃O(C₁-C₄ alkyl group); a -(CH₂)₀₋₃OC(O)-(C₁-C₄ alkyl group); a -(CH₂)₀₋₃C(O)O-(C₁-C₄ alkyl group);

a -C(O)-(CH₂)₀₋₃-NR'R'' group; a -(CH₂)₀₋₃C(O)OH group; a -(CH₂)₀₋₃-(5- to 10-membered monocyclic or bicyclic heteroaryl group containing at least one heteroatom selected from O, S and N); a -(CH₂)₀₋₃-(5- to 10- membered monocyclic or bicyclic heterocyclyl group containing at least one heteroatom selected from O, S and N);

or a linear or branched C₁-C₃ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; and

wherein the heteroaryl and heterocyclyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a linear or branched C₁-C₃ alkyl group or a C₁-C₃ haloalkyl group; and/or

(f) R₆ represents a hydrogen atom, a halogen atom, a hydroxyl group, a cyano group, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₃ alkoxy group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group;


and

wherein R₆ preferably represents a hydrogen atom, a halogen atom, a C₁-C₃ alkoxy group, a C₁-C₃ haloalkyl group, a C₃-C₄ cycloalkyl group, a -NH₂ group, a -N(CH₃)H group, a -N(CH₃)₂ group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by a C₁-C₂ alkoxy group.


7. A compound according to any one of claims 1, 3, 5 or 6, wherein:

(a) R₅ represents a group selected from

i) a group of formula (IIa-1), and ii) a group of formula (IIa-2)



formula (IIa-1)




formula (IIa-2)

wherein

R₉, R₁₀, R₁₁ and R₁₂ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; or


(b) R₅ represents a group selected from

i) a group of formula (IIb-1),




formula (IIb-1)

ii) a group of formula (IIb-2),




formula (IIb-2)

iii) a group of formula (IIb-3),




formula (IIb-3)

iv) a group of formula (IIb-4), and



formula (IIb-4)

v) a group of formula (IIb-5),




formula (IIb-5)

wherein


R₁₃, R₁₄, R₁₅ and R₁₆ each independently represent a hydrogen atom, a halogen atom; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a C₁-C₄ hydroxyalkyl group; a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group; a -C(O)-(CH₂)₀₋₃-R' group; a -C(O)-(CH₂)₀₋₃-NR'R'' group; a -(CH₂)₀₋₃NR'R'' group; or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined in claim 1; or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group; or (c) R₅ represents a group selected from

i) a group of formula (IIIa-1),




formula (IIIa-1)

ii) a group of formula (IIIa-2),




formula (IIIa-2)

iii) a group of formula (IIIa-3),




formula (IIIa-3)

iv) a group of formula (IIIa-4),




formula (IIIa-4)

v) a group of formula (IIIa-5),



formula (IIIa-5)


vi) a group of formula (IIIa-6),



formula (IIIa-6)


(continued)

vii) a group of formula (IIIa-7),




formula (IIIa-7)

viii) a group of formula (IIIa-8),




formula (IIIa-8)

ix) a group of formula (IIIa-9),




formula (IIIa-9)

X) a group of formula (IIIa-10),




formula (IIIa-10)

xi) a group of formula (IIIa-11),




formula (IIIa-11)

xii) a group of formula (IIIa-12),




formula (IIIa-12)

xiii) a group of formula (IIIa-13),




formula (IIIa-13)

xiv) a group of formula (IIIa-14),




formula (IIIa-14)

xv) a group of formula (IIIa-15),




formula (IIIa-15)

xvi) a group of formula (IIIa-16),




formula (IIIa-16)

xvii) a group of formula (IIIa-17),




formula (IIIa-17)

xviii) a group of formula (IIIa-18),




formula (IIIa-18)

xix) a group of formula (IIIa-19),




formula (IIIa-19)

xx) a group of formula (IIIa-20),



formula (IIIa-20)


xxi) a group of formula (IIIa-21), and



formula (IIIa-21)

xxii) a group of formula (IIIa-22)

(continued)




wherein

R₁₈, R₂₂, R₂₃, and R₂₄ each independently represent a hydrogen atom, a halogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃-CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined in claim 1;


or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group; or

(d) R₅ represents a group selected from


i) a group of formula (IIIb-1),




ii) a group of formula (IIIb-2),




iii) a group of formula (IIIb-3),




iv) a group of formula (IIIb-4),




v) a group of formula (IIIb-5),




vi) a group of formula (IIIb-6),



vii) a group of formula (IIIb-7), and



viii) a group of formula (IIIb-8),




wherein

R₁₈, R₂₂, R₂₅, R₂₆, and R₂₇ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined in claim 1;


or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group; or

(e) R₅ represents a group selected from

i) a group of formula (IIIc-1), and ii) a group of formula (IIIc-2),



formula (IIIc-1)



formula (IIIc-2)


wherein

R₁₉, R₂₀, and R₂₈ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined in claim 1;


or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group; or

(f) R₅ represents a group selected from


i) a group of formula (IIId-1), ii) a group of formula (IIId-2), and iii) a group of formula (IIId-3),



formula (IIId-1)



formula (IIId-2)



formula (IIId-3)

wherein


R₂₁, R₂₈, R₂₉, and R₃₀ each independently represent a hydrogen atom, a C₁-C₄ alkoxy group, a C₁-C₄ haloalkyl group, a C₁-C₄ hydroxyalkyl group, a C₃-C₄ cycloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₁₋₃CN group, a -C(O)-(CH₂)₀₋₃-R' group, a -C(O)-(CH₂)₀₋₃-NR'R'' group, a -(CH₂)₀₋₃NR'R'' group, or a linear or branched C₁-C₄ alkyl group, which alkyl group is unsubstituted or substituted by one or more substituents selected from a C₁-C₄ alkoxy group, a cyano group or a C₃-C₄ cycloalkyl group; wherein R' and R'' each independently represent a

hydrogen atom, a hydroxyl group, a C₁-C₄ alkoxy group or a linear or branched C₁-C₄ alkyl group; and wherein Y is as defined in claim 1;

or in the case that Y represents a -NR'- group, R₄ together with the -NR'- group and the carbon atom to which both R₄ and the -NR'- group are bonded form a 4- to 7-membered, saturated N-containing heterocyclyl group, which heterocyclyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a cyano group, a -CHF₂ group or a -CF₃ group.


8. A compound of according to claim 1, which is:

(a) a compound of fomula (Ia):



Formula (Ia)

wherein R₁, R₂, R₃, R₄, R₅, R₆, R_a, R_b and n are as defined in any one of the preceding claims; or
(b) a compound of fomula (Ib):



Formula (Ib)

wherein R₁, R₂, R₃, R₄, R₅, R_a, R_b and n are as defined in any one of claims 1 to 7.

9. A compound according to claim 1, wherein:

X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom or a methyl group;

R₁ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a methyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a pyridinyl group, a pyrazolyl group, an isoxazolyl group, a piperidinyl group or a tetrahydropyranyl group;

wherein the cycloalkyl, phenyl, pyridinyl, pyrazolyl, isoxazolyl, piperidinyl or tetrahydropyranyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a hydroxyl group, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ alkyl group, a -(CH₂)-(phenyl)-O-(C₁-C₃ alkyl group), a -NR₇R₈

group or a -OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₂ and R₃ each independently represent a hydrogen atom, a halogen atom, a cyano group, a C₁-C₃ haloalkyl group or a linear or branched C₁-C₃ alkyl group;

R₄ represents a hydrogen atom, a C₁-C₃ haloalkyl group, a C₁-C₃ hydroxyalkyl group or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom; a halogen atom; a hydroxyl group; a cyano group; a C₁-C₄ alkoxy group; a C₁-C₄ haloalkyl group; a linear or branched C₁-C₄ hydroxyalkyl group; a C₃-C₇ cycloalkyl group; a linear or branched C₁-C₃ alkyl group; a -(CH₂)₀₋₃NR'R" group; a -(CH₂)₁₋₃O(C₁-C₃ alkyl group); a -(CH₂)₀₋₃OC(O)-(C₁-C₃ alkyl group); a -(CH₂)₀₋₃C(O)O-(C₁-C₃ alkyl group); a -C(O)-NR'R" group; a -(CH₂)₀₋₃C(O)OH group; a -(CH₂)₀₋₃-(imidazolyl group); a -(CH₂)₀₋₃-(oxazolyl group); a -(CH₂)₀₋₃-(oxadiazolyl group); a -(CH₂)₀₋₃-(pyrazolyl group) or a -(CH₂)₀₋₃-(morpholinyl group); wherein R' and R" each independently represent a hydrogen atom, a hydroxyl group, or a linear or branched C₁-C₃ alkyl group; and

wherein the imidazolyl, oxazolyl, oxadiazolyl, pyrazolyl and morpholinyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₃ alkyl group or a C₁-C₃ haloalkyl group.

R₅ represents a group selected from:

i) a group of formula (IIa), which group is a purinyl group unsubstituted or substituted by a -NR'R" group;
ii) a group of formula (IIb), which group is selected from a -NR'-pyridinyl group, a -S-pyridinyl group, a -NR'-pyrimidinyl group, a -S-pyrimidinyl group or

a -NR'-triazinyl group; wherein the pyridinyl, pyrimidinyl and triazinyl groups are unsubstituted or substituted by one, two or three substituents selected from a halogen atom, a C₁-C₃ haloalkyl group, a -(CH₂)₀₋₃CN group, a -C(O)-(CH₂)₀₋₃-NR'R", a -(CH₂)₀₋₃NR'R" group; and

iii) a group of formula (IIc), which group is selected from a -NR'-purinyl group, a -S-purinyl group, a -NR'-7H-pyrrolo[2,3-d]pyrimidinyl group, a -NR'-1 H-pyrazolo[3,4-d]pyrimidinyl group or a -NR'-pyrazolo[1,5-a]pyrimidinyl group;

wherein the purinyl, 7H-pyrrolo[2,3-d]pyrimidinyl, 1 H-pyrazolo[3,4-d]pyrimidinyl pyrazolo[1,5-a]pyrimidinyl and groups are unsubstituted or substituted by a halogen atom or a -(CH₂)₀₋₃NR'R" group; or

R₄ and R₅ together with the carbon atom to which they are attached form a pyrrolidinyl-purinyl group or a pyrrolidinyl-pyrimidinyl; wherein the pyrrolidinyl group is unsubstituted or substituted by one or more substituents selected from a halogen atom or a hydroxyl group; and wherein the purinyl group is unsubstituted or substituted by a -(CH₂)₀₋₃NR'R" group; and wherein the pyrimidinyl group is unsubstituted or substituted by one, two or three substituents selected from a -(CH₂)₀₋₃CN group or a -(CH₂)₀₋₃NR'R" group; and R' and R" each independently represent a hydrogen atom, a C₁-C₃ alkoxy group or a linear or branched C₁-C₃ alkyl group.

10. A compound according to claim 1 or claim 3, wherein:

X represents a nitrogen atom or a -CR₆ group;

R_a and R_b each independently represent a hydrogen atom or a methyl group;

R₁ represents a methyl group, a C₃-C₇ cycloalkyl group, a phenyl group, a pyridinyl group, a piperidinyl group or a tetrahydropyranyl group;

wherein the cycloalkyl, phenyl, pyridinyl, piperidinyl or tetrahydropyranyl groups are unsubstituted or substituted by one or more substituents selected from a halogen atom, a linear or branched C₁-C₃ alkyl group, a -NR₇R₈ group or a -OR₈ group; wherein R₇ and R₈ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₂ and R₃ each independently represent a hydrogen atom or a linear or branched C₁-C₃ alkyl group;

R₄ represents a hydrogen atom, a C₁-C₃ haloalkyl group, or a linear or branched C₁-C₃ alkyl group;

R₆ represents a hydrogen atom, a halogen atom, a C₁-C₃ haloalkyl group, a linear or branched C₁-C₃ alkyl group or a cyclopropyl group;

R₅ represents a group selected from:

i) a group of formula (IIa), which group is a purinyl group unsubstituted or substituted by a -NR'R" group;

ii) a group of formula (IIb), which group is selected from a -NH-pyridinyl group, a -S-pyridinyl group, a -NH-pyrimidinyl group or a -S-pyrimidinyl group; wherein the pyridinyl or pyrimidinyl groups are unsubstituted or substituted by one, two or three substituents selected from a -(CH₂)₀₋₃CN group, a

-C(O)-(CH₂)₀₋₃-NR'R" or a -(CH₂)₀₋₃NR'R" group; and

iii) a group of formula (IIc), which group is selected from a -NH-puriny group or a -S-puriny group; wherein the puriny group is unsubstituted or substituted by a -(CH₂)₀₋₃NR'R" group; or

5 R₄ and R₅ together with the carbon atom to which they are attached form a pyrrolidinyl-puriny group, wherein the puriny group is unsubstituted or substituted by a -(CH₂)₀₋₃NR'R" group;
R' and R" each independently represent a hydrogen atom, a C₁-C₃ alkoxy group or a linear or branched C₁-C₃ alkyl group.

10 **11.** A compound according to claim 1, which is one of:

2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Aminopyrimidin-4-ylamino)methyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
15 2-((6-amino-9H-purin-9-yl)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-aminopyrimidin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
4-((4-Oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)picolinamide;
2-((2-aminopyridin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((9H-purin-6-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
20 2-((6-Amino-9H-purin-9-yl)methyl)-3-cyclohexylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-amino-9H-purin-9-yl)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((9H-purin-6-ylthio)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-amino-9H-purin-9-yl)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((9H-purin-6-ylthio)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
25 2-(1-(6-amino-9H-purin-9-yl)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
(S)-2-(1-(6-aminopyrimidin-4-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
30 (R)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-(1-(6-aminopyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
35 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
2-(1-(6-amino-9H-purin-9-yl)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
40 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(3-methoxyphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(2,4-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-3-benzyl-5-chloropyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-amino-9H-purin-9-yl)methyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-amino-9H-purin-9-yl)methyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
45 2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(pyridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(tetrahydro-2H-pyran-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-methylpiperidin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-4-Amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
50 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-methylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
55 2-((6-Amino-9H-purin-9-yl)methyl)-3-((1r,4r)-4-aminocyclohexyl)-5-chloropyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(R)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
(S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenyl-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(5-(difluoromethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

2-(1-(9H-purin-6-ylamino)-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

4-amino-6-(3,3,3-trifluoro-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;

(S)-4-amino-6-(2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile;

(S)-3-phenyl-2-(1-(pyrazolo[1,5-a]pyrimidin-7-ylamino)ethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

2-((6-amino-9H-purin-9-yl)methyl)-5-(difluoromethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(2-amino-9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(4,6-diamino-1,3,5-triazin-2-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-((6-amino-9H-purin-9-yl)methyl)-5-chloro-3-(1-(5-fluoropyridin-2-yl)ethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(R)-2-(1-(9H-purin-6-ylamino)-2-hydroxyethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(R)-4-amino-6-(2-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(2-amino-9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(methyl(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethyl)amino)pyrimidine-5-carbonitrile;

(S)-2-(1-(methyl(9H-purin-6-yl)amino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-7-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(4,4-difluoro-1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(4,4-difluoro-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-6-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(6-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

2-((S)-1-(9H-purin-6-ylamino)ethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

4-amino-6-((S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-amino-6-(1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-((9H-purin-6-ylamino)methyl)-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-((4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-amino-6-(1-(5-fluoro-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-

carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-5-fluoro-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

4-amino-6-((1S)-1-(5-(1,2-dihydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(trifluoromethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-Amino-6-(1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)imidazo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-Amino-6-(1-(5-(difluoromethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

2-(1-(9H-Purin-6-ylamino)-2,2,2-trifluoroethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-4-Amino-6-(1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-5-(difluoromethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)propyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-(trifluoromethyl)pyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(R)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-carbamoylpyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(2-Amino-5-fluoropyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

(S)-2-(1-(2-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;

((S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(2H-tetrazol-5-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(1-(3-((5-methylisoxazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 5 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-7-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-7-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 10 (S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(2,6-Diamino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 15 (S)-4-Amino-6-(1-(5-(morpholinomethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 20 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 25 (S)-4-Amino-6-(1-(3-(5-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylic acid;
 2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 30 (S)-4-Amino-6-(1-(3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 35 (S)-4-Amino-6-(1-(4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2,4-Diamino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 40 (S)-4-(1-(3-((1H-Pyrazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)-6-aminopyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(4-oxo-3-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 45 (S)-4-Amino-6-(1-(4-oxo-3-(2,2,2-trifluoroethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-Amino-4-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 50 4-Amino-6-(1-(5-(1-methyl-1H-pyrazol-4-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 55 (S)-4-Amino-6-(1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-4-Amino-6-(1-(5-bromo-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 2-((3-Iodo-1H-pyrazolo[3,4-d]pyrimidin-4-ylamino)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 5 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 10 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-5-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-N,N-dimethyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;
 15 (S)-4-Amino-6-(1-(3-(1-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-N-propyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide;
 20 2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((S)-1-(9H-purin-6-ylamino)ethyl)-3-((S)-tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
 (S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 25 (R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidine-5-carbonitrile;
 4-Amino-6-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 30 S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)-3-hydroxypropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-4-Amino-6-(1-(5-(2-methyloxazol-5-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 35 (S)-4-Amino-6-(1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-Propyl 2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylate;
 (S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;
 40 (S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-4-amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluoromethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-3-hydroxypropylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(4-oxo-3-(6-(trifluoromethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 45 (S)-4-Amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-2-(2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-yl)ethyl acetate;
 50 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(6-(trifluoromethyl)pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 2-((2S,4R)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-hydroxypyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 4-Amino-6-((2S,4R)-2-(5-(aminomethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-4-hydroxypyrrolidin-1-yl)pyrimidine-5-carbonitrile;
 55 (S)-4-Amino-6-(1-(5-(4-methyl-1H-imidazol-1-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(5-bromo-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;

(S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-4-Amino-6-(1-(5-bromo-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 5 (S)-4-Amino-6-(1-(3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 (S)-4-Amino-6-(1-(3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 10 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)cyclopropylamino)pyrimidine-5-carbonitrile;
 2-(1-(9H-Purin-6-ylamino)cyclopropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one; (S)-4-Amino-6-(1-(4-oxo-3-(3-(trifluoromethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidine-5-carbonitrile;
 15 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile;
 (S)-2-(1-(9H-purin-6-ylamino)ethyl)-3-(pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one;
 (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-one;
 20 (S)-4-amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile;

or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof.

12. A compound according to claim 1 or claim 11, which is (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile, or a pharmaceutically acceptable salt, or solvate, or N-oxide, or stereoisomer or deuterated derivative thereof.

13. A compound according to any one of claims 1 to 12, for use in the treatment of a pathological condition or disease susceptible to amelioration by inhibition of Phosphoinositide 3-Kinase (PI3K), wherein the pathological condition or disease is selected from respiratory diseases; allergic diseases; inflammatory or autoimmune-mediated; function disorders and neurological disorders; cardiovascular diseases; viral infection; metabolism/endocrine function disorders; neurological disorders and pain; bone marrow and organ transplant rejection; myelo-dysplastic syndrome; myeloproliferative disorders (MPDs); cancer and hematologic malignancies, leukemia, lymphomas and solid tumors, or wherein the pathological condition or disease is selected from leukemia, lymphomas and solid tumors, rheumatoid arthritis, multiple sclerosis, amyotrophic lateral sclerosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, autoimmune hemolytic anemia, type I diabetes, cutaneous vasculitis, cutaneous lupus erythematosus, dermatomyositis, blistering diseases including but not limited to pemphigus vulgaris, bullous pemphigoid and epidermolysis bullosa, asthma, chronic obstructive pulmonary disease, cystic fibrosis, idiopathic pulmonary fibrosis, sarcoidosis, allergic rhinitis, atopic dermatitis, contact dermatitis, eczema, psoriasis, basal cell carcinoma, squamous cell carcinoma and actinic keratosis.


14. A pharmaceutical composition comprising a compound as defined in any one of claims 1 to 12 in association with a pharmaceutically acceptable diluent or carrier.

15. A combination product comprising (i) a compound as defined in any one of claims 1 to 12; and (ii) another compound selected from the group consisting of an Adenoside A_{2A} agonist, an agent for treating cardiovascular disorders, an agent for treating diabetes, and an agent for treating liver disease, an anti-allergic agent, an anti-cholinergic agent, an anti-inflammatory agent, an anti-infective agent, a β 2-adrenergic agonist, a Chemoattractant receptor homologous molecule expressed on TH₂ cells (CRTH2) inhibitor, a chemotherapeutic agent, a corticosteroid, an IKK β /IKBKB (I κ B kinase beta or IKK2) inhibitor, an immunosuppressant, a Janus kinase (JAK) inhibitor, a topically acting p38 Mitogen-Activated Protein Kinase (p38 MAPK) inhibitor, a Phosphodiesterase (PDE) IV inhibitor, and a Spleen tyrosine kinase (Syk) inhibitor, for simultaneous, separate or sequential use in the treatment of the human or animal body.

Patentansprüche

1. Verbindung der Formel (I) oder ein pharmazeutisch unbedenkliches Salz oder Solvat oder N-Oxid oder Stereoisomer

oder deuteriertes Derivat davon:



Formel (I)

wobei:

X ein Stickstoffatom oder eine $-CR_6$ -Gruppe darstellt;

n 0, 1, 2 oder 3 darstellt;

R_a und R_b jeweils unabhängig ein Wasserstoffatom, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen;

R_1 ein Wasserstoffatom, eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_3 - C_{10} -Cycloalkylgruppe, eine C_3 - C_{10} -Cycloalkenylgruppe, eine monocyclische oder bicyclische C_6 - C_{14} -Arylgruppe, eine 5- bis 14-gliedrige monocyclische oder bicyclische Heteroarylgruppe darstellt, die zumindest ein Heteroatom enthält, das aus der Gruppe ausgewählt ist, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist,

wobei die Cycloalkyl-, Cycloalkenyl-, Aryl-, Heteroaryl- und Heterocyclylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C_1 - C_4 -Alkylgruppe, einer C_1 - C_4 -Haloalkylgruppe, einer C_1 - C_4 -Hydroxyalkylgruppe, einer C_3 - C_4 -Cycloalkylgruppe, einer $-(CH_2)_{1-3}CN$ -Gruppe, einer $-(CH_2)_{0-3}OR_8$ -Gruppe, einer $-(CH_2)_{0-3}NR_7R_8$ -Gruppe, einer $-C(O)-(CH_2)_{1-3}CN$ -Gruppe, einer $-C(O)-(CH_2)_{0-3}R_8$ -Gruppe, einer $-C(O)-(CH_2)_{0-3}NR_7R_8$ -Gruppe, einer $-S(O)_2(CH_2)_{0-3}R_8$ -Gruppe, einer $-S(O)_2(CH_2)_{0-3}NR_7R_8$ -Gruppe oder einer $-(CH_2)_{0-3}(Phenyl)-OR_8$ -Gruppe ausgewählt sind;

R_2 und R_3 jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C_1 - C_4 -Alkoxygruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe, eine C_3 - C_7 -Cycloalkylgruppe, eine $-NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1 - C_6 -Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyanogruppe oder einer C_3 - C_7 -Cycloalkylgruppe ausgewählt sind;

R_4 ein Wasserstoffatom, eine C_1 - C_4 -Alkoxygruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe, eine C_3 - C_7 -Cycloalkylgruppe, eine $-(CH_2)_{1-4}NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyanogruppe, einer C_3 - C_4 -Cycloalkylgruppe, einer $-C(OH)-(CH_2)_{0-3}R'$ -Gruppe oder einer $-C(O)-(CH_2)_{0-3}NR'R''$ -Gruppe ausgewählt sind;

R_6 ein Wasserstoffatom; ein Halogenatom; eine Hydroxylgruppe; eine Cyanogruppe; eine C_1 - C_4 -Alkoxygruppe; eine C_1 - C_4 -Haloalkylgruppe; eine lineare oder verzweigte C_1 - C_4 -Hydroxyalkylgruppe; eine C_3 - C_7 -Cycloalkylgruppe; eine $-(CH_2)_{0-3}NR'R''$ -Gruppe; eine $-(CH_2)_{1-3}O(C_1-C_4-Alkylgruppe)$; eine $-(CH_2)_{0-3}OC(O)-(C_1-C_4-Alkylgruppe)$; eine $-(CH_2)_{0-3}C(O)O-(C_1-C_4-Alkylgruppe)$; eine $-C(O)-(CH_2)_{0-3}NR'R''$ -Gruppe; eine $-(CH_2)_{0-3}C(O)OH$ -Gruppe; eine $-(CH_2)_{0-3}$ -(5- bis 14-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist); eine $-(CH_2)_{0-3}$ -(5- bis 14-gliedrige monocyclische oder bicyclische Heterocyclylgruppe, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist); oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyanogruppe oder einer C_3 - C_4 -Cycloalkylgruppe ausgewählt ist;

wobei die Heteroaryl- und Heterocyclylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder


verzweigten C₁-C₄-Alkylgruppe oder einer C₁-C₄-Haloalkylgruppe ausgewählt sind,

R₇ und R₈ jeweils unabhängig ein Wasserstoffatom, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyano-

gruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind;
R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen.


R₅ eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIa)



Formel (IIa)

ii) einer Gruppe der Formel (IIb)



Formel (IIb)

und

iii) einer Gruppe der Formel (IIc)



Formel (IIc)

wobei

Y einen Linker darstellt, der aus einer -NR'-Gruppe, -O- oder -S- ausgewählt ist; wobei R' wie oben definiert ist; (*) darstellt, wo R₅ an das an R₄ angehaftete Kohlenstoffatom und an die Pyrrolotriazinongruppe gebunden ist; W₁ eine -CR₁₁-Gruppe darstellt und W₂ ein Stickstoffatom darstellt oder W₁ ein Stickstoffatom darstellt und W₂ eine -CR₁₂-Gruppe darstellt;

G₁ eine -CR₁₄-Gruppe darstellt und G₂ ein Stickstoffatom darstellt oder G₁ ein Stickstoffatom darstellt und G₂ eine -CR₁₅-Gruppe darstellt oder G₁ eine -CR₁₄-Gruppe darstellt und G₂ eine CR₁₅-Gruppe darstellt;

G₃ ein Stickstoffatom oder eine CR₁₆-Gruppe darstellt;


R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ und R₁₆ jeweils unabhängig ein Wasserstoffatom; ein Halogenatom; eine C₁-C₄-Alkoxygruppe; eine C₁-C₄-Haloalkylgruppe; eine C₁-C₄-Hydroxyalkylgruppe; eine C₃-C₄-Cycloalkylgrup-

pe; eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}CN$ -Gruppe; eine $-C(O)-(CH_2)_{0-3}R'$ -Gruppe; eine $-C(O)-(CH_2)_{0-3}NR'R''$ -Gruppe; eine $-(CH_2)_{0-3}NR'R''$ -Gruppe; oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyanogruppe oder einer C_3 - C_4 -Cycloalkylgruppe ausgewählt sind;


wobei R' und R'' wie oben definiert sind;

R_{17} eine Gruppe darstellt, ausgewählt aus:

a) einer Gruppe der Formel (IIIa), b) einer Gruppe der Formel (IIIb)




Formel (IIIa)




Formel (IIIb)

c) einer Gruppe der Formel (IIIc), und d) einer Gruppe der Formel (IIId)



Formel (IIIc)



Formel (IIId)

wobei:

G_4 ein Stickstoffatom oder eine $-CR_{22}$ -Gruppe darstellt;

G_5 und G_6 jeweils unabhängig ein Stickstoffatom oder ein Kohlenstoffatom darstellen, wobei, wenn eines von G_5 und G_6 ein Stickstoffatom darstellt, das andere ein Kohlenstoffatom darstellt;

G_7 eine $-NH$ -Gruppe oder eine $-CH$ -Gruppe darstellt;

G_8 ein Stickstoffatom oder eine $-CR_{23}$ -Gruppe darstellt;

G_9 ein Stickstoffatom oder eine $-CR_{24}$ -Gruppe darstellt;

G_{10} ein Stickstoffatom oder eine $-CR_{25}$ -Gruppe darstellt;

G_{11} ein Stickstoffatom oder eine $-CR_{26}$ -Gruppe darstellt;

G_{12} ein Stickstoffatom oder eine $-CR_{27}$ -Gruppe darstellt;

G_{13} ein Stickstoffatom oder eine $-CR_{28}$ -Gruppe darstellt;

G_{14} und G_{15} jeweils unabhängig ein Stickstoffatom oder ein Kohlenstoffatom darstellen, wobei, wenn eines von G_{14} und G_{15} ein Stickstoffatom darstellt, das andere ein Kohlenstoffatom darstellt;

G_{16} eine $-NH$ -Gruppe oder eine $-CH$ -Gruppe darstellt;

G_{17} ein Stickstoffatom oder eine $-CR_{29}$ -Gruppe darstellt;

G_{18} ein Stickstoffatom oder eine $-CR_{30}$ -Gruppe darstellt;

R_{18} , R_{19} , R_{20} , R_{21} , R_{22} , R_{23} , R_{24} , R_{25} , R_{26} , R_{27} , R_{28} , R_{29} und R_{30} jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine C_1 - C_4 -Alkoxygruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe, eine C_3 - C_4 -Cycloalkylgruppe, eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}R'$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}NR'R''$ -Gruppe, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe oder eine lineare oder verzweigte

C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R'' wie oben definiert sind; und wobei Y wie oben definiert ist;

oder wenn Y eine Gruppe -NR'- darstellt, R₄ zusammen mit der Gruppe -NR'- und dem Kohlenstoffatom, an das sowohl R₄ als auch die Gruppe -NR'- gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer -CHF₂-Gruppe- oder einer -CF₃-Gruppe ausgewählt sind.

2. Verbindung nach Anspruch 1, wobei:

R_a und R_b jeweils unabhängig ein Wasserstoffatom oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen;

n 0, 1 oder 2 darstellt;

R₁ ein Wasserstoffatom, eine lineare oder verzweigte C₁-C₄-Alkylgruppe, eine C₁-C₄-Haloalkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine Phenylgruppe, eine 5- bis 10-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind, eine Pyrrolidinylgruppe, eine Piperidinylgruppe, eine Piperazinylgruppe, eine Tetrahydropyranylgruppe oder eine Morpholinylgruppe darstellt;

wobei die Cycloalkyl-, Phenyl-, Heteroaryl-, Pyrrolidinyl-, Piperidinyl-, Piperazinyl-, Tetrahydropyranyl- oder Morpholinylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₀₋₃OR₈-Gruppe, einer -(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-(CH₂)₀₋₃R₈-Gruppe, einer -C(O)-(CH₂)₀₋₃NR₇R₈-Gruppe oder einer -(CH₂)₀₋₃(Phenyl)-OR₈-Gruppe ausgewählt sind; wobei R₇ und R₈ jeweils unabhängig ein Wasserstoffatom oder eine C₁-C₄-Alkylgruppe darstellen;

R₂ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist;


R₃ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist;

R₄ ein Wasserstoffatom, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellt;

R₆ ein Wasserstoffatom; ein Halogenatom; eine Hydroxylgruppe; eine Cyanogruppe; eine C₁-C₃-Alkoxygruppe; eine C₁-C₃-Haloalkylgruppe; eine lineare oder verzweigte C₁-C₄-Hydroxyalkylgruppe; eine C₃-C₄-Cycloalkylgruppe; eine -(CH₂)₀₋₃NR'R''-Gruppe; eine -(CH₂)₁₋₃O(C₁-C₄-Alkylgruppe); eine -(CH₂)₀₋₃OC(O)-(C₁-C₄-Alkylgruppe); eine -(CH₂)₀₋₃C(O)O-(C₁-C₄-Alkylgruppe); eine -C(O)-(CH₂)₀₋₃NR'R''-Gruppe; eine -(CH₂)₀₋₃C(O)OH-Gruppe; eine -(CH₂)₀₋₃-(5- bis 14-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist); eine -(CH₂)₀₋₃-(5- bis 14-gliedrige monocyclische oder bicyclische Heterocyclgruppe, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist); oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist;

wobei die Heteroaryl- und Heterocyclgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C₁-C₄-Alkylgruppe oder einer C₁-C₄-Haloalkylgruppe ausgewählt sind,

R₅ eine Einheit der Formel (II-a2), (IIb-1), (IIb-2), (IIb-3), (IIIa-1), (IIIa-3), (IIIa-5) oder (IIIa-14) darstellt:




Formel (IIa-2)

Formel (IIb-1)

Formel (IIb-2)

Formel (IIb-3)



Formel (IIIa-1)

Formel (IIIa-3)

Formel (IIIa-5)

Formel (IIIa-14)

wobei:

- R_9 , R_{10} und R_{12} jeweils unabhängig ein Wasserstoffatom, eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}R'$ -Gruppe, ein $-C(O)-(CH_2)_{0-3}NR'R''$, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellen; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C_1-C_4 -Alkoxygruppe oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellen;
- R_{13} , R_{14} , R_{15} und R_{16} jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine C_1-C_4 -Haloalkylgruppe, eine $-(CH_2)_{0-3}CN$ -Gruppe, ein $-C(O)-(CH_2)_{0-3}NR'R''$, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe darstellen; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C_1-C_4 -Alkoxygruppe oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellen;
- R_{18} , R_{22} , R_{23} und R_{24} jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}R'$ -Gruppe, ein $-C(O)-(CH_2)_{0-3}NR'R''$, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellen; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C_1-C_4 -Alkoxygruppe oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellen;
- Y eine Gruppe $-NR'$, $-O-$ oder $-S-$ darstellt, wobei R' einen Wasserstoff oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellt; oder wenn Y eine Gruppe $-NR'$ darstellt, R_4 zusammen mit der Gruppe $-NR'$ und dem Kohlenstoffatom, an das sowohl R_4 als auch die Gruppe $-NR'$ gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyano-Gruppe, einer $-CHF_2$ -Gruppe oder einer $-CF_3$ -Gruppe ausgewählt sind.

3. Verbindung nach Anspruch 1, wobei:

X ein Stickstoffatom oder eine $-CR_6$ -Gruppe;

n 0, 1, 2 oder 3 darstellt;

R_a und R_b jeweils unabhängig ein Wasserstoffatom, eine C_1-C_4 -Haloalkylgruppe, eine C_1-C_4 -Hydroxyalkylgruppe oder eine lineare oder verzweigte C_1-C_4 -Alkylgruppe darstellen;

R_1 eine lineare oder verzweigte C_1-C_4 -Alkylgruppe, eine C_3-C_{10} -Cycloalkylgruppe, eine C_3-C_{10} -Cycloalkenylgruppe, eine monocyclische oder bicyclische C_6-C_{14} -Arylgruppe, eine 5- bis 14-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die zumindest ein Heteroatom enthält, das aus der Gruppe ausgewählt ist, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist, oder eine 5- bis 14-gliedrige monocyclische oder bicyclische Heterocyclgruppe darstellt, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist,

wobei die Cycloalkyl-, Cycloalkenyl-, Aryl-, Heteroaryl- und Heterocyclylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₁₋₃CN-Gruppe, einer -(CH₂)₀₋₃OR₈-Gruppe, einer

-(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-(CH₂)₁₋₃CN-Gruppe, einer -C(O)-(CH₂)₀₋₃R₈-Gruppe, einer -C(O)-(CH₂)₀₋₃NR₇R₈-Gruppe, einer -S(O)₂(CH₂)₀₋₃R₈-Gruppe oder einer -S(O)₂(CH₂)₀₋₃NR₇R₈-Gruppe ausgewählt sind;

R₂ und R₃ jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine -NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₆-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₇-Cycloalkylgruppe ausgewählt sind;

R₄ ein Wasserstoffatom, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine -(CH₂)₁₋₄NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe, einer C₃-C₄-Cycloalkylgruppe, einer -C(O)-(CH₂)₀₋₃R'-Gruppe oder einer -C(O)-(CH₂)₀₋₃NR'R"-Gruppe ausgewählt sind;

R₆ ein Wasserstoffatom; ein Halogenatom; eine Hydroxylgruppe; eine Cyanogruppe; eine C₁-C₄-Alkoxygruppe; eine C₁-C₄-Haloalkylgruppe; eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine -(CH₂)₀₋₃NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt ist;

R₇ und R₈ jeweils unabhängig ein Wasserstoffatom, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind;

R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen.

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ und R₁₆ jeweils unabhängig ein Wasserstoffatom, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃CN-Gruppe, eine -C(O)-(CH₂)₀₋₃R'-Gruppe, eine -C(O)-(CH₂)₀₋₃NR'R"-Gruppe, eine -(CH₂)₀₋₃NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R" wie oben definiert sind;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉ und R₃₀ jeweils unabhängig ein Wasserstoffatom, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃CN-Gruppe, eine -C(O)-(CH₂)₀₋₃R'-Gruppe, eine -C(O)-(CH₂)₀₋₃NR'R"-Gruppe, eine -(CH₂)₀₋₃NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R" wie oben definiert sind; und wobei Y wie oben definiert ist;

4. Verbindung nach Anspruch 3, wobei:

R_a und R_b jeweils unabhängig ein Wasserstoffatom oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen;

n 0, 1 oder 2 darstellt;

R₁ eine lineare oder verzweigte C₁-C₄-Alkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine Phenylgruppe, eine 5- bis 10-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind, eine Pyrrolidinylgruppe, eine Piperidinylgruppe, eine Piperazinylgruppe, eine Tetrahydropyranylgruppe oder eine Morpholinylgruppe darstellt;

wobei die Cycloalkyl-, Phenyl-, Heteroaryl-, Pyrrolidinyl-, Piperidinyl-, Piperazinyl-, Tetrahydropyranyl- oder Morpholinylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₀₋₃OR₈-Gruppe, einer -(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-(CH₂)₀₋₃R₈-Gruppe oder einer -C(O)-(CH₂)₀₋₃NR₇R₈-Gruppe ausgewählt sind; wobei R₇ und R₈ jeweils unabhängig ein Wasserstoffatom oder eine C₁-C₄-Alkylgruppe darstellen;


R₂ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist;

R₃ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist;


R₄ ein Wasserstoffatom, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellt;

R₆ ein Wasserstoffatom; ein Halogenatom, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe ist, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist;


R₅ eine Einheit der Formel (IIa-2), (IIb-1), (IIb-2) oder (IIIa-1) darstellt:




Formel (IIa-2)



Formel (IIb-1)



Formel (IIb-2)



Formel (IIIa-1)

wobei:

- R₉, R₁₀ und R₁₂ jeweils unabhängig ein Wasserstoffatom, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃CN-Gruppe, eine -C(O)-(CH₂)₀₋₃-R'-Gruppe, ein -C(O)-(CH₂)₀₋₃-NRR"-Gruppe, eine -(CH₂)₀₋₃NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen;
- R₁₃, R₁₄, R₁₅ und R₁₆ jeweils unabhängig ein Wasserstoffatom, eine -(CH₂)₀₋₃CN-Gruppe, ein -C(O)-(CH₂)₀₋₃-NR'-R" oder eine -(CH₂)₀₋₃NR'R"-Gruppe darstellen; wobei R' und R" jeweils unabhängig ein Wasserstoffatom oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen;
- R₁₈ und R₂₃ jeweils unabhängig ein Wasserstoffatom, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃CN-Gruppe, eine C(O)-(CH₂)₀₋₃-R-Gruppe, ein -C(O)-CH₂₀₋₃-NR'R", eine -(CH₂)₀₋₃NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen;
- Y eine Gruppe -NR'-, -O- oder -S- darstellt; wobei R' einen Wasserstoff oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt; oder wenn Y eine Gruppe -NR'- darstellt, R₄ zusammen mit der Gruppe -NR'- und dem Kohlenstoffatom, an das sowohl R₄ als auch die Gruppe -NR'- gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden.

5. Verbindung nach Anspruch 1, wobei R₁ ein Wasserstoffatom, eine lineare oder verzweigte C₁-C₄-Alkylgruppe, eine C₁-C₄-Haloalkylgruppe, eine C₃-C₁₀-Cycloalkylgruppe, eine C₃-C₁₀-Cycloalkenylgruppe, eine monocyclische oder bicyclische C₆-C₁₄-Arylgruppe, eine 5- bis 14-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist, oder eine 5- bis 14-gliedrige monocyclische oder bicyclische Heterocyclgruppe darstellt, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist,

wobei die Cycloalkyl-, Cycloalkenyl-, Aryl-, Heteroaryl- und Heterocyclgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₁₋₃CN-Gruppe, einer -(CH₂)₀₋₃OR₈-Gruppe, einer -(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-CH₂₁₋₃-CN-Gruppe, einer -C(O)-(CH₂)₁₋₃-CN-Gruppe, einer -C(O)-(CH₂)₀₋₃-R₈-Gruppe, einer

-C(O)-(CH₂)₀₋₃-NR₇R₈-Gruppe, einer -S(O)₂(CH₂)₀₋₃R₈-Gruppe, einer -S(O)₂(CH₂)₀₋₃NR₇R₈-Gruppe oder einer -(CH₂)₀₋₃-(Phenyl)-OR₈-Gruppe ausgewählt sind;
und

wobei R₁ vorzugsweise ein Wasserstoffatom, eine C₁-C₃-Alkylgruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine Phenylgruppe, eine Naphthylgruppe, eine 5- bis 10-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind, oder eine 5- bis 10-gliedrige monocyclische oder bicyclische Heterocyclylgruppe darstellt, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind,

wobei die Cycloalkyl-, Phenyl-, Naphthyl-, Heteroaryl- und Heterocyclylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₀₋₃OR₈-Gruppe, einer -(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-CH₂)₀₋₃-R₈-Gruppe, einer -C(O)-(CH₂)₀₋₃-NR₇R₈-Gruppe, oder einer -(CH₂)₀₋₃-(Phenyl)-OR₈-Gruppe ausgewählt sind; wobei R₇ und R₈ wie in Anspruch 1 definiert sind.

6. Verbindung nach einem der Ansprüche 1, 3 oder 5, wobei:

(a) R₁ eine C₁-C₃-Alkylgruppe, eine C₃-C₇-Cycloalkylgruppe, eine Phenylgruppe, eine Naphthylgruppe, eine 5- bis 10-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind, oder eine 5- bis 10-gliedrige monocyclische oder bicyclische Heterocyclylgruppe darstellen, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind, wobei die Cycloalkyl-, Phenyl-, Naphthyl-, Heteroaryl- und Heterocyclylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₀₋₃OR₈-Gruppe, einer -(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-CH₂)₀₋₃-R₈-Gruppe oder einer -C(O)-(CH₂)₀₋₃-NR₇R₈-Gruppe ausgewählt sind; wobei R₇ und R₈ wie in Anspruch 1 definiert sind;

und

wobei R₁ vorzugsweise eine C₃-C₇-Cycloalkylgruppe, eine Phenylgruppe, eine 5- bis 10-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die ein, zwei oder drei Heteroatome enthält, die aus O, S und N ausgewählt sind, eine Pyrrolidinygruppe, eine Piperidinygruppe, eine Piperazinygruppe, eine Tetrahydropyranygruppe oder eine Morpholinylgruppe darstellt,

wobei die Cycloalkyl-, Phenyl-, - Heteroaryl- Pyrrolidiny-, Piperaziny-, Tetrahydropyrany- oder Morpholinylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom einer linearen oder verzweigten C₁-C₄-Alkylgruppe, einer C₁-C₄-Haloalkylgruppe, einer C₁-C₄-Hydroxyalkylgruppe, einer C₃-C₄-Cycloalkylgruppe, einer -(CH₂)₀₋₃CNOR₈-Gruppe, einer -(CH₂)₀₋₃NR₇R₈-Gruppe, einer -C(O)-CH₂)₀₋₃-R₈-Gruppe oder einer -C(O)-(CH₂)₀₋₃-NR₇R₈-Gruppe ausgewählt sind; wobei R₇ und R₈ jeweils

unabhängig ein Wasserstoffatom oder eine C₁-C₄-Alkylgruppe darstellen;

und

wobei R₁ mehr bevorzugt eine Phenylgruppe oder eine Pyridinygruppe darstellt, wobei das Phenyl oder Pyridiny unsubstituiert oder mit ein, zwei oder drei Substituenten substituiert ist, die aus einem Halogenatom, einer linearen oder verzweigten C₁-C₃-Alkylgruppe oder einer -(CH₂)₀₋₃OCH₃-Gruppe ausgewählt sind; und wobei vorzugsweise die Phenyl- oder Pyridinygruppen direkt an die Pyrrolotriazinongruppe gebunden sind; und/oder

(b) R₂ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃-NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₃-Alkoxygruppe substituiert ist; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen;

und

wobei R₂ vorzugsweise ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist; und/oder

(c) R₃ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃-NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₃-Alkoxygruppe substituiert ist; wobei R' und R" jeweils unabhängig ein Was-

serstoffatom, eine Hydroxygruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen;
und

wobei R₃ vorzugsweise ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist; und/oder

(d) R₄ ein Wasserstoffatom, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₁₋₄NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₃-Alkoxygruppe, einer -C(O)-(CH₂)₀₋₃-R'-Gruppe oder einer -C(O)-(CH₂)₀₋₃-NR'R"-Gruppe substituiert ist; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen;

und

wobei R₄ vorzugsweise ein Wasserstoffatom, eine C₁-C₃-Haloalkylgruppe, eine C₁-C₃-Hydroxylalkylgruppe, eine C₃-C₄-Cycloalkylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellt;

und/oder

(e) R₆ ein Wasserstoffatom; ein Halogenatom; eine Hydroxylgruppe; eine Cyanogruppe; eine C₁-C₄-Alkoxygruppe; eine C₁-C₄-Haloalkylgruppe; eine lineare oder verzweigte C₁-C₄-Hydroxyalkylgruppe; eine C₃-C₇-Cycloalkylgruppe; eine -(CH₂)₀₋₃-NR'R"-Gruppe; eine -(CH₂)₁₋₃O(C₁-C₄-Alkylgruppe); eine -(CH₂)₀₋₃OC(O)-(C₁-C₄-Alkylgruppe); eine -(CH₂)₀₋₃C(O)O-(C₁-C₄-Alkylgruppe); eine -C(O)-(CH₂)₀₋₃-NR'R"-Gruppe; eine -(CH₂)₀₋₃C(O)OH-Gruppe; eine -(CH₂)₀₋₃-(5- bis 14-gliedrige monocyclische oder bicyclische Heteroarylgruppe, die zumindest Heteroatom enthält, das aus O, S und N ausgewählt ist);

eine -(CH₂)₀₋₃-(5- bis 14-gliedrige monocyclische oder bicyclische Heterocyclylgruppe, die zumindest ein Heteroatom enthält, das aus O, S und N ausgewählt ist); oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einem der mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen; und

wobei die Heteroaryl- und Heterocyclylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer linearen oder verzweigten C₁-C₃-Alkylgruppe oder einer C₁-C₃-Haloalkylgruppe ausgewählt sind, und/oder

(f) R₆ ein Wasserstoffatom, ein Halogenatom, eine Hydroxylgruppe, eine Cyanogruppe, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₃-Alkoxygruppe substituiert ist; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe oder eine lineare oder verzweigte C₁-C₃-Alkylgruppe darstellen;

und


wobei R₆ vorzugsweise ein Wasserstoffatom, ein Halogenatom, eine C₁-C₃-Alkoxygruppe, eine C₁-C₃-Haloalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -NH₂-Gruppe, eine -N(CH₃)H-Gruppe, eine -N(CH₃)₂-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellt, wobei die Alkylgruppe unsubstituiert oder mit einer C₁-C₂-Alkoxygruppe substituiert ist.

7. Verbindung nach einem der Ansprüche 1, 3, 5 oder 6, wobei:


(a) R₅ eine Gruppe darstellt, ausgewählt aus:

- i) einer Gruppe der Formel (IIa-1) und ii) einer Gruppe der Formel (IIa-2)

(fortgesetzt)



Formel (IIa-1)



Formel (IIa-2)


wobei

R_9 , R_{10} , R_{11} und R_{12} jeweils unabhängig ein Wasserstoffatom, eine C_1 - C_4 -Alkoxygruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe, eine C_3 - C_4 -Cycloalkylgruppe, eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}R'$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}NR'R''$ -Gruppe, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyanogruppe oder einer C_3 - C_4 -Cycloalkylgruppe ausgewählt sind; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C_1 - C_4 -Alkoxygruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen; oder


(b) R_5 eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIb-1)

ii) einer Gruppe der Formel (IIb-2)




Formel (IIb-1)




Formel (IIb-2)

iii) einer Gruppe der Formel (IIb-3),

iv) einer Gruppe der Formel (IIb-4) und




Formel (IIb-3)



Formel (IIb-4)

v) einer Gruppe der Formel (IIb-5)

(fortgesetzt)



Formel (IIb-5)

wobei

R_{13} , R_{14} , R_{15} und R_{16} jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine C_1 - C_4 -Alkoxygruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe, eine C_3 - C_4 -Cycloalkylgruppe, eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}R'$ -Gruppe, eine $-C(O)-(CH_2)_{0-3}NR'R''$ -Gruppe, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyanogruppe oder einer C_3 - C_4 -Cycloalkylgruppe ausgewählt sind;


wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C_1 - C_4 -Alkoxygruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen; und

wobei Y wie in Anspruch 1 definiert ist;

oder wenn Y eine Gruppe $-NR'$ - darstellt, R_4 zusammen mit der Gruppe $-NR'$ - und dem Kohlenstoffatom, an das sowohl R_4 als auch die Gruppe $-NR'$ - gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer $-CHF_2$ -Gruppe oder einer $-CF_3$ -Gruppe ausgewählt sind; oder


(c) R_5 eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIIa-1)




Formel (IIIa-1)

ii) einer Gruppe der Formel (IIIa-2)




Formel (IIIa-2)

iii) einer Gruppe der Formel (IIIa-3),



Formel (IIIa-3)

iv) einer Gruppe der Formel (IIIa-4)




Formel (IIIa-4)


v) einer Gruppe der Formel (IIb-5)

vi) einer Gruppe der Formel (IIIa-6)

(fortgesetzt)




Formel (IIIa-5)




Formel (IIIa-6)

vii) einer Gruppe der Formel (IIIa-7),

viii) einer Gruppe der Formel (IIIa-8)




Formel (IIIa-7)




Formel (IIIa-8)

ix) einer Gruppe der Formel (IIIa-9),

x) einer Gruppe der Formel (IIIa-10)




Formel (IIIa-9)




Formel (IIIa-10)

xi) einer Gruppe der Formel (IIIa-11),

xii) einer Gruppe der Formel (IIIa-12)




Formel (IIIa-11)




Formel (IIIa-12)

xiii) einer Gruppe der Formel (IIIa-13),

xiv) einer Gruppe der Formel (IIIa-14)




Formel (IIIa-13)




Formel (IIIa-14)

(fortgesetzt)

xv) einer Gruppe der Formel (IIIa-15), xvi) einer Gruppe der Formel (IIIa-16)




Formel (IIIa-15)




Formel (IIIa-16)

xvii) einer Gruppe der Formel (IIIa-17), xviii) einer Gruppe der Formel (IIIa-18)




Formel (IIIa-17)




Formel (IIIa-18)

xix) einer Gruppe der Formel (IIIa-19), xx) einer Gruppe der Formel (IIIa-20)




Formel (IIIa-19)




Formel (IIIa-20)

xxi) einer Gruppe der Formel (IIIa-21), xxii) einer Gruppe der Formel (IIIa-22)



Formel (IIIa-21)



Formel (IIIa-22)


und

wobei


R₁₈, R₂₂, R₂₃ und R₂₄ jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃-CN-Gruppe, eine C(O)-(CH₂)₀₋₃-R'-Gruppe, ein -C(O)-CH₂)₀₋₃-NR'R'', eine -(CH₂)₀₋₃NR'R''-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyanogruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen; und wobei Y wie in Anspruch 1 definiert ist;

oder wenn Y eine Gruppe -NR'- darstellt, R₄ zusammen mit der Gruppe -NR'- und dem Kohlenstoffatom, an das sowohl R₄ als auch die Gruppe -NR'- gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer --CHF₂-Gruppe oder einer -CF₃-Gruppe ausgewählt sind; oder
 (d) R₅ eine Gruppe darstellt, ausgewählt aus:

- i) einer Gruppe der Formel (IIIb-1), ii) einer Gruppe der Formel (IIIb-2)




Formel (IIIb-1)




Formel (IIIb-2)

- iii) einer Gruppe der Formel (IIIb-3), iv) einer Gruppe der Formel (IIIb-4)




Formel (IIIa-3)




Formel (IIIa-4)

- v) einer Gruppe der Formel (IIIb-5), vi) einer Gruppe der Formel (IIIb-6)




Formel (IIIb-5)




Formel (IIIb-6)

- vii) einer Gruppe der Formel (IIIb-7), viii) einer Gruppe der Formel (IIIb-8)



Formel (IIIb-7) und




Formel (IIIb-8)


wobei

R_{18} , R_{22} , R_{25} , R_{26} und R_{27} jeweils unabhängig ein Wasserstoffatom, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkylgruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃CN-Gruppe, eine C(O)-(CH₂)₀₋₃-R'-Gruppe, ein -C(O)-CH₂)₀₋₃-NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyano-
 5 gruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen; und wobei Y wie in Anspruch 1 definiert ist;
 oder wenn Y eine Gruppe -NR'- darstellt, R₄ zusammen mit der Gruppe -NR'- und dem Kohlenstoffatom, an
 10 das sowohl R₄ als auch die Gruppe -NR'- gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer --CHF₂-Gruppe oder einer -CF₃-Gruppe ausgewählt sind; oder
 (e) R₅ eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIIc-1), ii) einer Gruppe der Formel (IIIc-2)



Formel (IIIc-1)




Formel (IIIc-2)


wobei

R_{19} , R_{20} und R_{28} jeweils unabhängig ein Wasserstoffatom, eine C₁-C₄-Alkoxygruppe, eine C₁-C₄-Haloalkyl-
 30 gruppe, eine C₁-C₄-Hydroxyalkylgruppe, eine C₃-C₄-Cycloalkylgruppe, eine -(CH₂)₀₋₃CN-Gruppe, eine -C(O)-(CH₂)₁₋₃CN-Gruppe, eine C(O)-(CH₂)₀₋₃-R'-Gruppe, ein -C(O)-CH₂)₀₋₃-NR'R"-Gruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert
 35 oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C₁-C₄-Alkoxygruppe, einer Cyano-
 gruppe oder einer C₃-C₄-Cycloalkylgruppe ausgewählt sind; wobei R' und R" jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C₁-C₄-Alkoxygruppe oder eine lineare oder verzweigte C₁-C₄-Alkylgruppe darstellen; und wobei Y wie in Anspruch 1 definiert ist;
 oder wenn Y eine Gruppe -NR'- darstellt, R₄ zusammen mit der Gruppe -NR'- und dem Kohlenstoffatom, an
 40 das sowohl R₄ als auch die Gruppe -NR'- gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer --CHF₂-Gruppe oder einer -CF₃-Gruppe ausgewählt sind; oder
 (f) R₅ eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIId-1), ii) einer Gruppe der Formel (IIId-2)



Formel (IIId-1)




Formel (IIId-2)

und

i) einer Gruppe der Formel (IIId-3)

(fortgesetzt)



Formel (IIIId-3)


wobei

R_{21} , R_{28} , R_{29} und R_{30} jeweils unabhängig ein Wasserstoffatom, eine C_1 - C_4 -Alkoxygruppe, eine C_1 - C_4 -Haloalkylgruppe, eine C_1 - C_4 -Hydroxyalkylgruppe, eine C_3 - C_4 -Cycloalkylgruppe, eine $-(CH_2)_{0-3}CN$ -Gruppe, eine $-C(O)-(CH_2)_{1-3}-CN$ -Gruppe, eine $C(O)-(CH_2)_{0-3}-R'$ -Gruppe, ein $-C(O)-CH_2)_{0-3}-NR'R''$, eine $-(CH_2)_{0-3}NR'R''$ -Gruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen, wobei die Alkylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einer C_1 - C_4 -Alkoxygruppe, einer Cyano-Gruppe oder einer C_3 - C_4 -Cycloalkylgruppe ausgewählt sind; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe, eine C_1 - C_4 -Alkoxygruppe oder eine lineare oder verzweigte C_1 - C_4 -Alkylgruppe darstellen; und wobei Y wie in Anspruch 1 definiert ist;

oder wenn Y eine Gruppe $-NR'$ - darstellt, R_4 zusammen mit der Gruppe $-NR'$ - und dem Kohlenstoffatom, an das sowohl R_4 als auch die Gruppe $-NR'$ - gebunden ist, eine 4- bis 7-gliedrige gesättigte N-haltige Heterocyclgruppe bilden, wobei die Heterocyclgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom, einer Hydroxylgruppe, einer Cyanogruppe, einer $-CHF_2$ -Gruppe oder einer $-CF_3$ -Gruppe ausgewählt sind.


8. Verbindung nach Anspruch 1, die ist:

(a) eine Verbindung der Formel (Ia):



Formel (Ia)

wobei R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_a , R_b und wie in einem der vorstehenden Ansprüche definiert sind; oder
(b) eine Verbindung der Formel (Ib)



Formel (Ib)

wobei R_1 , R_2 , R_3 , R_4 , R_5 , R_a , R_b und n wie in einem der Ansprüche 1 bis 7 definiert sind.

9. Verbindung nach Anspruch 1, wobei:

X ein Stickstoffatom oder eine $-CR_6$ -Gruppe darstellt;

R_a und R_b jeweils unabhängig ein Wasserstoffatom oder eine Methylgruppe darstellen;

R_1 ein Wasserstoffatom, ein Halogenatom, eine C_1 - C_3 -Haloalkylgruppe, eine Methylgruppe, eine C_3 - C_7 -Cycloalkylgruppe, eine Phenylgruppe, eine Pyridinylgruppe,

eine Pyrazolylgruppe, eine Isoxazolylgruppe, eine Piperidinylgruppe oder eine Tetrahydropyranylgruppe darstellt;

wobei die Cycloalkyl-, Phenyl-, -, Pyridinyl-, Pyrazolyl-, Isoxazolyl-, Piperidinyl-, -oder Tetrahydropyranylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer Hydroxylgruppe,

einer C_1 - C_3 -Haloalkylgruppe, einer linearen oder verzweigte C_1 - C_3 -Alkylgruppe, einer $-(CH_2)$ -Phenyl)-O- $(C_1$ - C_3 -Alkylgruppe), einer $-NR_7R_8$ -Gruppe oder einer $-OR_8$ -Gruppe ausgewählt sind; wobei R_7 und R_8 jeweils unabhängig ein Wasserstoffatom oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen;

R_2 und R_3 jeweils unabhängig ein Wasserstoffatom, ein Halogenatom, eine Cyanogruppe, eine C_1 - C_3 -Haloalkylgruppe oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen;

R_4 ein Wasserstoffatom, eine C_1 - C_3 -Haloalkylgruppe, eine C_1 - C_3 -Hydroxyalkylgruppe oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellt;

R_6 ein Wasserstoffatom; ein Halogenatom; eine Hydroxylgruppe; eine Cyanogruppe; eine C_1 - C_4 -Alkoxygruppe; eine C_1 - C_4 -Haloalkylgruppe; eine lineare oder verzweigte C_1 - C_4 -Hydroxyalkylgruppe; eine C_3 - C_7 -Cycloalkylgruppe; eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe; eine $-(CH_2)_{0-3}$ - $NR'R''$ -Gruppe; eine $-(CH_2)_{1-3}O(C_1$ - C_3 -Alkylgruppe); eine $-(CH_2)_{0-3}OC(O)-(C_1$ - C_3 -Alkylgruppe); eine $-(CH_2)_{0-3}C(O)-(C_1$ - C_3 -Alkylgruppe); eine $-C(O)-NR'R''$ -Gruppe; eine $-(CH_2)_{0-3}C(O)OH$ -Gruppe; eine $-(CH_2)_{0-3}$ -(Imidazolylgruppe); eine $-(CH_2)_{0-3}$ -(Oxazolylgruppe); eine $-(CH_2)_{0-3}$ -(Oxadiazolylgruppe); eine $-(CH_2)_{0-3}$ -(Pyrazolylgruppe) oder eine $-(CH_2)_{0-3}$ -(Morpholinylgruppe) darstellt; wobei R' und R'' jeweils unabhängig ein Wasserstoffatom, eine Hydroxylgruppe oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen; und

wobei die Imidazolyl-, Oxazolyl-, Oxadiazolyl-, Pyrazolyl- und Morpholinylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer linearen oder verzweigten C_1 - C_3 -Alkylgruppe oder einer C_1 - C_3 -Haloalkylgruppe ausgewählt sind,

R_5 eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIa), wobei die Gruppe eine Purinylgruppe ist, die unsubstituiert oder mit einer $-NR'R''$ -Gruppe substituiert ist;

ii) einer Gruppe der Formel (IIb), wobei die Gruppe aus einer $-NR'$ -Pyridinylgruppe, einer $-S$ -Pyridinylgruppe, einer $-NR'$ -Pyrimidinylgruppe, einer $-S$ -Pyrimidinylgruppe oder einer $-NR'$ -Triazinylgruppe ausgewählt ist; wobei die Pyridinyl-, Pyrimidinyl- und Triazinylgruppen unsubstituiert oder mit einem, zwei oder drei Substituenten substituiert sind, die aus einem Halogenatom, einer C_1 - C_3 -Haloalkylgruppe, einer $-(CH_2)_{0-3}CN$ -Gruppe, einem $-C(O)-(CH_2)_{0-3}-NR'R''$ oder einer $-(CH_2)_{0-3}NR'R''$ -Gruppe ausgewählt sind; und

iii) einer Gruppe der Formel (IIc), wobei die Gruppe aus einer $-NR'$ -Purinylgruppe, einer $-S$ -Purinylgruppe, einer $-NR'$ -7H-Pyrrolo[2,3-d]pyrimidinylgruppe, einer $-NR'$ -1H-Pyrazolo[3,4-d]pyrimidinylgruppe oder einer

-NR'-Pyrazolo[1,5-a]pyrimidinylgruppe ausgewählt ist;
wobei die Purinyl-, 7H-Pyrrolo[2,3-d]pyrimidinyl-, -1H-Pyrazolo[3,4-d]pyrimidinyl und Pyrazolo[1,5-a]pyrimidinyl- und -Gruppen unsubstituiert oder mit einem Halogenatom oder einer $-(CH_2)_{0-3}NR''$ -Gruppe substituiert sind; oder

R_4 und R_5 zusammen mit dem Kohlenstoffatom, an dem sie angehaftet sind, eine Pyrrolidinylpurinylgruppe oder ein Pyrrolidinylpyrimidinyl bilden; wobei die Pyrrolidinylgruppe unsubstituiert oder mit einem oder mehreren Substituenten substituiert ist, die aus einem Halogenatom oder einer Hydroxylgruppe ausgewählt sind; und wobei die Purinylgruppe unsubstituiert oder mit einer $-(CH_2)_{0-3}NR''$ -Gruppe substituiert ist; und wobei die Pyrimidinylgruppe unsubstituiert oder mit einem, zwei oder drei Substituenten substituiert ist, die aus einer $-(CH_2)_{0-3}CN$ -Gruppe oder einer $-(CH_2)_{0-3}NR''$ -Gruppe ausgewählt sind; und R' und R'' jeweils unabhängig ein Wasserstoffatom, eine C_1 - C_3 -Alkoxygruppe oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen.

10. Verbindung nach Anspruch 1 oder 3, wobei:

X ein Stickstoffatom oder eine $-CR_6$ -Gruppe darstellt;

R_a und R_b jeweils unabhängig ein Wasserstoffatom oder eine Methylgruppe darstellen;

R_1 eine Methylgruppe, eine C_3 - C_7 -Cycloalkylgruppe, eine Phenylgruppe, eine Pyridinylgruppe, eine Piperidinylgruppe oder eine Tetrahydropyranylgruppe darstellt;

wobei die Cycloalkyl-, Phenyl-, Pyridinyl-, Piperidinyl- oder Tetrahydropyranylgruppen unsubstituiert oder mit einem oder mehreren Substituenten substituiert sind, die aus einem Halogenatom, einer linearen oder verzweigten C_1 - C_3 -Alkylgruppe, einer $-NR_7R_8$ -Gruppe oder einer $-OR_8$ -Gruppe ausgewählt sind; wobei R_7 und R_8 jeweils unabhängig ein Wasserstoffatom oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen;

R_2 und R_3 jeweils unabhängig ein Wasserstoffatom oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen;

R_4 ein Wasserstoffatom, eine C_1 - C_3 -Haloalkylgruppe oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellt;

R_6 ein Wasserstoffatom, ein Halogenatom, eine C_1 - C_3 -Haloalkylgruppe, eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe oder eine Cyclopropylgruppe darstellt;

R_6 eine Gruppe darstellt, ausgewählt aus:

i) einer Gruppe der Formel (IIa), wobei die Gruppe eine Purinylgruppe, unsubstituiert oder mit einer $-NR'R''$ -Gruppe substituiert ist;

ii) einer Gruppe der Formel (IIb), wobei die Gruppe aus einer $-NH$ -Pyridinylgruppe, einer $-S$ -Pyridinylgruppe, einer $-NH$ -Pyrimidinylgruppe oder einer $-S$ -Pyrimidinylgruppe ausgewählt ist; wobei die Pyridinyl- oder Pyrimidinylgruppen unsubstituiert oder mit ein, zwei oder drei Substituenten substituiert sind, die aus einer $-(CH_2)_{0-3}CN$ -Gruppe, einem $-C(O)-(CH_2)_{0-3}NR'R''$ oder einer $-(CH_2)_{0-3}NR'R''$ -Gruppe ausgewählt sind; und

iii) einer Gruppe der Formel (IIc), wobei die Gruppe aus einer $-NH$ -Purinylgruppe oder einer $-S$ -Purinylgruppe ausgewählt ist; wobei die Purinylgruppe unsubstituiert oder mit einer $-(CH_2)_{0-3}NR'R''$ -Gruppe substituiert ist; oder

R_4 und R_6 zusammen mit dem Kohlenstoffatom, an dem sie angehaftet sind, eine Pyrrolidinylpurinylgruppe bilden, wobei die Purinylgruppe unsubstituiert oder durch eine $-(CH_2)_{0-3}NR'R''$ -Gruppe substituiert ist; R' und R'' jeweils unabhängig ein Wasserstoffatom, eine C_1 - C_3 -Alkoxygruppe oder eine lineare oder verzweigte C_1 - C_3 -Alkylgruppe darstellen.

11. Verbindung nach Anspruch 1, die eine des Folgenden ist:

2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
2-((6-Aminopyrimidin-4-ylamino)methyl)-5-chlor-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
2-((6-Amino-9H-purin-9-yl)methyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
2-((6-Aminopyrimidin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
4-((4-Oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)picolinamid;
2-((2-Aminopyridin-4-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
2-((9H-Purin-6-ylamino)methyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;

- 2-((6-Amino-9H-purin-9-yl)methyl)-3-cyclohexylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((9H-Purin-6-ylthio)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 5 2-((9H-Purin-6-ylthio)methyl)-6-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-(1-(6-Amino-9H-purin-9-yl)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Aminopyrimidin-4-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 10 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidin-5-carbonitril;
 (R)-2-(1-(9H-Purin-6-ylamino)propyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 15 (S)-2-(1-(6-Aminopyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 2-(1-(6-Amino-9H-purin-9-yl)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolyl-5-(trifluormethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 20 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(3-methoxyphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(2,4-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-benzyl-5-chlorpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
 25 2-((6-Amino-9H-purin-9-yl)methyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(pyridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(tetrahydro-2H-pyran-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(1-methylpiperidin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3-fluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 30 (S)-4-Amino-6-(1-(3-(3-fluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 35 2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-methylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-3-((1r,4r)-4-aminocyclohexyl)-5-chlorpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (R)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 40 (S)-2-(1-(9H-Purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-5-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-phenyl-5-(trifluormethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 45 (S)-4-Amino-6-(1-(5-(difluormethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluormethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 50 ril;
 2-(1-(9H-Purin-6-ylamino)-3,3,3-trifluoropropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 4-Amino-6-(3,3,3-trifluor-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidin-5-carbonitril;
 55 nitril;
 (S)-3-Phenyl-2-(1-(pyrazolo[1,5-a]pyrimidin-7-ylamino)ethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((6-Amino-9H-purin-9-yl)methyl)-5-(difluormethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(2-Amino-9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;

- (S)-2-(1-(4,6-Diamino-1,3,5-triazin-2-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-chlor-3-(1-(5-fluorpyridin-2-yl)ethyl)pyrrolo[[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 5 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (R)-2-(1-(9H-Purin-6-ylamino)-2-hydroxyethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (R)-4-Amino-6-(2-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 10 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(7H-Pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(methyl(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethyl)amino)pyrimidin-5-carbonitril;
 15 (S)-2-(1-(Methyl(9H-purin-6-yl)amino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(5-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-7-methyl-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 20 (S)-4-Amino-6-(1-(7-methyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(4,4-Difluor-1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(4,4-difluor-2-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidin-5-carbonitril;
 25 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-6-fluor-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(6-fluor-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 2-(S)-1-(9H-Purin-6-ylamino)ethyl)-3-((S)-1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 4-Amino-6-((S)-1-(4-oxo-3-((S)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 30 (S)-4-Amino-6-(1-(3-(2,6-dimethylphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-((9H-Purin-6-ylamino)methyl)-3-(1-phenylethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-((4-oxo-3-(1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidin-5-carbonitril;
 35 (S)-2-(1-(5-Fluor-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(2,6-dimethylphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(5-fluor-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 40 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-fluor-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-4-Amino-6-(1-(3-(3,5-difluorphenyl)-4-oxo-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 45 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 4-Amino-6-((1S)-1-(5-(1,2-dihydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(3-(3,5-difluorphenyl)-4-oxo-5-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 50 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorphenyl)-5-(trifluormethyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(5-(hydroxymethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-(trifluormethyl)pyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 55 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(pyridin-2-ylmethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-5-(difluormethyl)-3-(3,5-difluorphenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorphenyl)imidazo[1,2-f][1,2,4]triazin-4(3H)-on;

- (S)-4-Amino-6-(1-(5-(difluormethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-5-(difluormethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 5 (S)-2-(1-(2-Amino-9H-purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 2-(1-(9H-Purin-6-ylamino)-2,2,2-trifluorethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 10 (S)-2-(1-(6-Amino-5-fluorpyrimidin-4-ylamino)ethyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-fluorpyrimidin-4-ylamino)ethyl)-5-(difluormethyl)-3-(3,5-difluorophenyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)propyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 15 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-dichlorophenyl)-4-oxo-3,4-dihydropyrrolo [1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-fluorpyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-(trifluormethyl)pyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 20 (R)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)-2-hydroxyethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-carbamoylpyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carboxamid;
 25 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carboxamid;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(2-chlorbenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 30 (R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(2-Amino-5-fluorpyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 35 (S)-2-(1-(2-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 ((S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(2H-tetrazol-5-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(3-((5-methylisoxazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 40 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-7-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-7-carbonitril;
 45 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(1-(4-methoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-4-amino-6-(1-(4-oxo-3-phenyl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(2,6-Diamino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 50 (S)-4-Amino-6-(1-(5-(morpholinomethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-1-phenylethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 55 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(3,5-difluorophenyl)-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;

- 4-Amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-5-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(3-(5-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 5 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonsäure;
 2-((S)-1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 10 (S)-4-Amino-6-(1-(3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 15 4-Amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-5-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2,4-Diamino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-(1-(3-((1H-Pyrazol-3-yl)methyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)-6-aminopyrimidin-5-carbonitril;
 20 (S)-4-Amino-6-(1-(4-oxo-3-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(2,2,2-trifluorethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 25 (S)-4-Amino-6-(1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-Amino-4-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 4-Amino-6-(1-(5-(1-methyl-1H-pyrazol-4-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 30 (S)-4-Amino-6-(1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-brom-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 4-Amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 35 (S)-4-Amino-6-(1-(5-brom-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 2-((3-Iod-1H-pyrazolo[3,4-d]pyrimidin-4-ylamino)methyl)-5-methyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 40 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 4-Amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-5-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 45 (S)-4-Amino-6-(1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-N,N-dimethyl-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carboxamid;
 50 (S)-4-Amino-6-(1-(3-(1-methyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-N-propyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carboxamid;
 2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-(tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 55 2-((S)-1-(9H-Purin-6-ylamino)ethyl)-3-((S)-tetrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(R)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyethylamino)pyrimidin-5-carbonitril;
 4-Amino-6-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)methylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-(2-hydroxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)-3-hydroxypropyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(5-fluorpyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-(2-methyloxazol-5-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-(2-methoxyethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-Propyl-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carboxylat;
 (S)-4-Amino-6-(3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)-3-hydroxypropyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(3-(3,5-difluorophenyl)-4-oxo-5-(trifluormethyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-3-hydroxypropylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(6-(trifluormethyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-brom-4-oxo-3-(3-(trifluormethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-yl)ethylacetat;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(6-(trifluormethyl)pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 2-((2S,4R)-1-(6-Amino-5-cyanopyrimidin-4-yl)-4-hydroxypyrrolidin-2-yl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 4-Amino-6-((2S,4R)-2-(5-(aminomethyl)-3-(3,5-difluorophenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-4-hydroxypyrrolidin-1-yl)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-(4-methyl-1H-imidazol-1-yl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-brom-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-(3-(trifluormethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-4-Amino-6-(1-(5-brom-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-4-Amino-6-(1-(3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-methoxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)cyclopropylamino)pyrimidin-5-carbonitril;
 2-(1-(9H-Purin-6-ylamino)cyclopropyl)-3-phenylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(4-oxo-3-(3-(trifluormethyl)phenyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)ethylamino)pyrimidin-5-carbonitril;
 (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-3-(3-hydroxyphenyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril;
 (S)-2-(1-(9H-Purin-6-ylamino)ethyl)-3-(pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ylamino)propyl)-3-phenylimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(4-oxo-3-phenyl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidin-5-carbonitril;


oder ein pharmazeutisch unbedenkliches Salz oder Solvat oder N-Oxid oder Stereoisomer oder deuteriertes Derivat

davon.

12. Verbindung nach Anspruch 1 oder 11, die (S)-2-(1-(6-Amino-5-cyanopyrimidin-4-ylamino)ethyl)-4-oxo-3-phenyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-carbonitril ist, oder ein pharmazeutisch unbedenkliches Salz oder Solvat oder N-Oxid oder Stereoisomer oder deuteriertes Derivat davon.
13. Verbindung nach einem der Ansprüche 1 bis 12 zur Verwendung bei der Behandlung eines pathologischen Zustands oder einer pathologischen Krankheit, die für eine Besserung durch Hemmung von Phosphoinositid-3-Kinase (PI3K) zugänglich ist, wobei der pathologische Zustand oder die pathologische Krankheit aus Atemwegserkrankungen; allergischen Krankheiten; entzündungs- oder autoimmunvermittelt; Funktionsstörungen und neurologischen Störungen; Herz-Kreislauf-Krankheiten; Virusinfektionen; Stoffwechsel-/endokrinen Funktionsstörungen; neurologischen Störungen und Schmerzen; Knochenmarks- und Organtransplantatabstoß; myelodysplastischem Syndrom; myeloproliferativen Störungen (MPDs); Krebs und hämatologischen Malignitäten, Leukämie, Lymphomen und soliden Tumoren ausgewählt ist, oder wobei der pathologische Zustand oder die pathologische Krankheit aus Leukämie, Lymphomen und soliden Tumoren, rheumatoider Arthritis, multipler Sklerose, amyotropher Lateralsklerose, Morbus Crohn, ulzerativer Kolitis, systemischem Lupus erythematoses, autoimmuner hämolytischer Anämie, Diabetes Typ I, kutaner Vaskulitis, kutanem Lupus erythematoses, Dermatomyositis, bläschenbildenden Krankheiten, einschließlich, aber nicht ausschließlich Pemphigus vulgaris, bullösem Pemphigoid und Epidermolysis bullosa, Asthma, chronisch-obstruktiver Lungenkrankheit, zystischer Fibrose, idiopathischer Pulmonalfibrose, Sarkoidose, allergischer Rhinitis, atopischer Dermatitis, Kontaktdermatitis, Ekzem, Psoriasis, Basalzellkarziom, Plattenepithelzellkarzinom und aktinischer Keratose ausgewählt ist.
14. Pharmazeutische Zusammensetzung, die eine Verbindung nach einem der Ansprüche 1 bis 12 in Verbindung mit einem pharmazeutisch unbedenklichen Verdünnungsmittel oder Träger umfasst.
15. Kombinationsprodukt, das (i) eine Verbindung nach einem der Ansprüche 1 bis 12; und (ii) eine weitere Verbindung umfasst, die aus der Gruppe ausgewählt ist, bestehend aus einem Adenosid-A_{2A}-Agonisten, einem Mittel zur Behandlung von Herz-Kreislauf-Störungen, einem Mittel zur Behandlung von Diabetes und einem Mittel zur Behandlung von Leberkrankheiten, einem Antiallergikum, einem Anticholinergikum, einem Entzündungshemmer, einem Infektionshemmer, einem β 2-adrenergen Agonisten, einem Hemmer des auf TH₂-Zellen exprimierten homologen Chemoattraktanz-Rezeptor-Moleküls (CRTH2), einem chemotherapeutischen Mittel, einem Kortikosteroid, einem Hemmer von IKK β /IKKB (IkB-Kinase beta oder IKK2), einem Immunsuppressivum, einem Hemmer der Janus-Kinase (JAK), einem topisch wirkenden Hemmer von p38-mitogenaktivierter Proteinkinase (p38-MAPK), einem Hemmer der Phosphodiesterase (PDE) IV und einem Hemmer der Milz-Tyrosinkinase (Syk) ausgewählt ist, zur gleichzeitigen, separaten oder sequentiellen Verwendung bei der Behandlung des Körpers eines Menschen oder Tiers.

Revendications

1. Composé de formule (I), ou sel pharmaceutiquement acceptable, ou solvate ou N-oxyde, ou stéréoisomère ou dérivé deutéré de celui-ci :



Formule (I)

formule dans laquelle,

X représente un atome d'azote ou un groupe $-CR_6$;

n représente 0, 1, 2 ou 3 ;

R_a et R_b représente chacun indépendamment un atome hydrogène, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ;

R_1 représente un atome d'hydrogène, un groupe alkyle en C_1-C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_{10} , un groupe cycloalcényle en C_3-C_{10} , un groupe aryle en C_6-C_{14} monocyclique ou bicyclique, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N, ou un groupe hétérocyclyle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N,

où les groupes cycloalkyle, cycloalcényle, aryle, hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C_1-C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{1-3}CN$, un groupe $-(CH_2)_{0-3}OR_8$, un groupe $-(CH_2)_{0-3}NR_7R_8$, un groupe $-C(O)-(CH_2)_{1-3}CN$, un groupe $-C(O)-(CH_2)_{0-3}R_8$, un groupe $-C(O)-(CH_2)_{0-3}NR_7R_8$, un groupe $-S(O)_2(CH_2)_{0-3}R_8$, un groupe $-S(O)_2(CH_2)_{0-3}NR_7R_8$ ou un groupe $-(CH_2)_{0-3}(phényl)-OR_8$;

R_2 et R_3 représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C_1-C_4 , un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_7 , un groupe $-NR'R''$ ou un groupe alkyle en C_1-C_6 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1-C_4 , un groupe cyano ou un groupe cycloalkyle en C_3-C_7 ;

R_4 représente un atome d'hydrogène, un groupe alcoxy en C_1-C_4 , un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_7 , un groupe $-(CH_2)_{1-4}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1-C_4 , un groupe cyano, un groupe cycloalkyle en C_3-C_4 , un groupe $-C(O)-(CH_2)_{0-3}R$ ou un groupe $-C(O)-(CH_2)_{0-3}NR'R''$;

R_6 représente un atome d'hydrogène ; un atome d'halogène ; un groupe hydroxyle ; un groupe cyano ; un groupe alcoxy en C_1-C_4 ; un groupe halogénoalkyle en C_1-C_4 ; un groupe hydroxyalkyle en C_1-C_4 linéaire ou ramifié ; un groupe cycloalkyle en C_3-C_7 ; un groupe $-(CH_2)_{0-3}NR'R''$; un groupe $-(CH_2)_{1-3}O$ (alkyle en C_1-C_4) ; un groupe $-(CH_2)_{0-3}OC(O)$ (alkyle en C_1-C_4) ; un groupe $-(CH_2)_{0-3}C(O)O$ (alkyle en C_1-C_4) ; un groupe $-C(O)-(CH_2)_{0-3}NR'R''$; un groupe $-(CH_2)_{0-3}C(O)OH$; un groupe $-(CH_2)_{0-3}$ (hétéroaryle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N) ; un groupe $-(CH_2)_{0-3}$ (hétérocyclyle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N) ; ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1-C_4 , un groupe cyano ou un groupe cycloalkyle en C_3-C_4 ;


où les groupes hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C_1-C_4 linéaire ou ramifié ou un groupe halogénoalkyle en C_1-C_4 ,

R_7 et R_8 représentent chacun indépendamment un atome d'hydrogène, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1-C_4 , un groupe cyano ou un groupe cycloalkyle en C_3-C_4 ;

R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié,


R_5 représente un groupe choisi parmi :

i) un groupe de formule (IIa)



formule (IIa)

ii) un groupe de formule (IIb)



formule (IIb)

et

iii) un groupe de formule (IIc)



formule (IIc)

dans lesquelles

Y représente un lieu choisi parmi un groupe -NR', -O- ou -S-; où R' est tel que défini ci-avant ;

(*) représente l'endroit où R₅ est lié à l'atome de carbone attaché à R₄ et au groupe pyrrolotriazinone ;

W₁ représente un groupe -CR₁₁ et W₂ représente un atome d'azote, ou W₁ représente un atome d'azote et W₂ représente un groupe -CR₁₂ ;

G₁ représente un groupe -CR₁₄ et G₂ représente un atome d'azote, ou G₁ représente un atome d'azote et G₂ représente un groupe -CR₁₅, ou G₁ représente un groupe -CR₁₄ et G₂ représente un groupe -CR₁₅ ;


G₃ représente un atome d'azote ou un groupe -CR₁₆ ;

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ et R₁₆ représentent chacun indépendamment un atome d'hydrogène ; un atome d'halogène ; un groupe alcoxy en C₁-C₄ ; un groupe halogénoalkyle en C₁-C₄ ; un groupe hydroxyalkyle en C₁-C₄ ; un groupe cycloalkyle en C₃-C₄ ; un groupe -(CH₂)₀₋₃CN ; un groupe -C(O)-(CH₂)₁₋₃CN ; un groupe -C(O)-(CH₂)₀₋₃R' ; un groupe -C(O)-(CH₂)₀₋₃NR'' ; un groupe -(CH₂)₀₋₃NR''R'' ; ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' sont tels que définis ci-avant ;

R₁₇ représente un groupe choisi parmi


a) un groupe de formule (IIIa), b) un groupe de formule (IIIb),

(suite)




formule (IIIa)

c) un groupe de formule (IIIc),




formule (IIIb)

d) un groupe de formule (IIId),



formule (IIIc)



formule (IIId)

où

G₄ représente un atome d'hydrogène ou un groupe -CR₂₂ ;G₅ et G₆ représentent chacun indépendamment un atome d'azote ou un atome de carbone, où lorsqu'un de G₅ et G₆ représente un atome d'azote, l'autre représente un atome de carbone ;G₇ représente un groupe -NH ou un groupe -CH ;G₈ représente un atome d'azote ou un groupe -CR₂₃ ;G₉ représente un atome d'azote ou un groupe -CR₂₄ ;G₁₀ représente un atome d'azote ou un groupe -CR₂₅ ;G₁₁ représente un atome d'azote ou un groupe -CR₂₆ ;G₁₂ représente un atome d'azote ou un groupe -CR₂₇ ;G₁₃ représente un atome d'azote ou un groupe -CR₂₈ ;G₁₄ et G₁₅ représentent chacun indépendamment un atome d'azote ou un atome de carbone, l'un de G₁₄ et G₁₅ représentant un atome d'azote et le restant représentant un atome de carbone ;G₁₆ représente un groupe -NH ou un groupe -CH ;G₁₇ représente un atome d'azote ou un groupe -CR₂₉ ;G₁₈ représente un atome d'azote ou un groupe -CR₃₀ ;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉ et R₃₀ représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)(CH₂)₁₋₃-CN, un groupe -C(O)-(CH₂)₀₋₃-R', un groupe -C(O)-(CH₂)₀₋₃-NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ;

où R' et R'' sont tels que définis ci-avant ; et où Y est tel que défini ci-avant ;

ou dans le cas où Y représente un groupe -NR', R₄, conjointement au groupe -NR' et à l'atome de carbone auquel R₄ et le groupe -NR' sont tous deux liés, forme un groupe hétérocyclyle contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe -CHF₂ ou un groupe -CF₃.

2. Composé selon la revendication 1, dans lequel

R_a et R_b représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C_1 - C_3 linéaire ou ramifié ;

n représente 0, 1 ou 2 ;

R_1 représente un atome d'hydrogène, un groupe alkyle en C_1 - C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1 - C_4 , un groupe cycloalkyle en C_3 - C_7 , un groupe phényle, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S et N, un groupe pyrrolidinyle, un groupe pipéridinyle, un groupe pipérazinyle, un groupe tétrahydropyranyle ou un groupe morpholinyle ;

où les groupes cycloalkyle, phényle, hétéroaryle, pyrrolidinyle, pipéridinyle, pipérazinyle, tétrahydropyranyle ou morpholinyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, ou groupe hydroxyle, ou groupe alkyle en C_1 - C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1 - C_4 , un groupe hydroxyalkyle en C_1 - C_4 , un groupe cycloalkyle en C_3 - C_4 , un groupe $-(CH_2)_{0-3}OR_8$, un groupe $-(CH_2)_{0-3}NR_7R_8$, un groupe $-C(O)-(CH_2)_{0-3}R_8$, un groupe $-C(O)-(CH_2)_{0-3}NR_7R_8$ ou un groupe $-(CH_2)_{0-3}(\text{phényl})-OR_8$; R_7 et R_8 représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C_1 - C_4 ;

R_2 représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe alcoxy en C_1 - C_3 , un groupe halogénoalkyle en C_1 - C_3 , un groupe cycloalkyle en C_3 - C_4 , un groupe $-NH_2$, un groupe $-N(CH_3)H$, un groupe $-N(CH_3)_2$ ou un groupe alkyle en C_1 - C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1 - C_2 ;


R_3 représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C_1 - C_3 , un groupe halogénoalkyle en C_1 - C_3 , un groupe cycloalkyle en C_3 - C_4 , un groupe $-NH_2$, un groupe $-N(CH_3)H$, un groupe $-N(CH_3)_2$ ou un groupe alkyle en C_1 - C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1 - C_2 ;

R_4 représente un atome d'hydrogène, un groupe alcoxy en C_1 - C_3 , un groupe halogénoalkyle en C_1 - C_3 , un groupe hydroxyalkyle en C_1 - C_4 , un groupe cycloalkyle en C_3 - C_4 ou un groupe alkyle en C_1 - C_3 linéaire ou ramifié ;


R_6 représente un atome d'hydrogène ; un atome d'halogène ; un groupe hydroxyle ; un groupe cyano ; un groupe alcoxy en C_1 - C_3 ; un groupe halogénoalkyle en C_1 - C_3 ; un groupe hydroxyalkyle en C_1 - C_4 linéaire ou ramifié ; un groupe cycloalkyle en C_3 - C_4 ; un groupe $-(CH_2)_{0-3}NR'R''$ un groupe $-(CH_2)_{1-3}O$ (alkyle en C_1 - C_4) ; un groupe $-(CH_2)_{0-3}OC(O)-(alkyle\ en\ C_1-C_4)$; un groupe $-(CH_2)_{0-3}C(O)O-(alkyle\ en\ C_1-C_4)$; un groupe $-C(O)-(CH_2)_{0-3}NR'R''$; un groupe $-(CH_2)_{0-3}C(O)OH$; un groupe $-(CH_2)_{0-3}$ (hétéroaryle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N) ; un groupe $-(CH_2)_{0-3}$ (hétérocyclyle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N) ; ou un groupe alkyle en C_1 - C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1 - C_2 ;

où les groupes hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C_1 - C_4 linéaire ou ramifié ou un groupe halogénoalkyle en C_1 - C_4 ,


R_5 représente un fragment de formule (IIa-2), (IIb-1), (IIb-2), (IIb-3), (IIIa-1), (IIIa-3), (IIIa-5) ou (IIIa-14) :




formule (IIa-2)




formule (IIb-1)




formule (IIb-2)




formule (IIb-3)




formule (IIIa-1)



formule (IIIa-3)



formule (IIIa-5)



formule (IIIa-14)

où :

- R_9 , R_{10} et R_{12} représentent chacun indépendamment un atome d'hydrogène, un groupe $-(CH_2)_{0-3}CN$, un groupe $-C(O)-(CH_2)_{1-3}-CN$, un groupe $-C(O)-(CH_2)_{0-3}-R'$, un groupe $-C(O)-(CH_2)_{0-3}-NR'R''$, un groupe $-(CH_2)_{0-3}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ;
- R_{13} , R_{14} , R_{15} et R_{16} représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe halogénoalkyle en C_1-C_4 , un groupe $-(CH_2)_{0-3}CN$, un groupe $-C(O)-(CH_2)_{0-3}-NR'R''$, un groupe $-(CH_2)_{0-3}NR'R''$; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ;
- R_{18} , R_{22} , R_{23} et R_{24} représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe $-(CH_2)_{0-3}CN$, un groupe $-C(O)-(CH_2)_{1-3}-CN$, un groupe $-C(O)-(CH_2)_{0-3}-R'$, un groupe $-C(O)-(CH_2)_{0-3}-NRR''$, un groupe $-(CH_2)_{0-3}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ;
- Y représente un groupe $-NR'$, $-O-$ ou $-S-$; où R' représente un atome d'hydrogène ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ; ou dans le cas où Y représente un groupe $-NR'$, R_4 , conjointement au groupe $-NR'$ et à l'atome de carbone auquel R_4 et le groupe $-NR'$ sont tous deux liés, forme un groupe hétérocyclyle contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe $-CHF_2$ ou un groupe $-CF_3$.

3. Composé selon la revendication 1, dans lequel

X représente un atome d'azote ou un groupe $-CR_6$;

n représente 0, 1, 2 ou 3 ;

R_a et R_b représentent chacun indépendamment un atome d'hydrogène, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ;

R_1 représente un groupe alkyle en C_1-C_4 linéaire ou ramifié, un groupe cycloalkyle en C_3-C_{10} , un groupe cycloalcényle en C_3-C_{10} , un groupe aryle en C_6-C_{14} monocyclique ou bicyclique, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N, ou un groupe hétérocyclyle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N,

où les groupes cycloalkyle, cycloalcényle, aryle, hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C_1-C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{1-3}CN$, un groupe $-(CH_2)_{0-3}OR_8$, un groupe $-(CH_2)_{0-3}NR_7R_8$, un

groupe - C(O)-(CH₂)₁₋₃-CN, un groupe -C(O)-(CH₂)₀₋₃-R₈, un groupe - C(O)-(CH₂)₀₋₃-NR₇R₈, un groupe -S(O)₂(CH₂)₀₋₃R₈ ou un groupe - S(O)₂(CH₂)₀₋₃NR₇R₈ ;

R₂ et R₃ représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₇, un groupe -NR'R'' ou un groupe alkyle en C₁-C₆ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₇ ;

R₄ représente un atome d'hydrogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₇, un groupe -(CH₂)₁₋₄NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano, un groupe cycloalkyle en C₃-C₄, un groupe -C(O)-(CH₂)₀₋₃-R ou un groupe -C(O)-(CH₂)₀₋₃-NR'R'' ;

R₆ représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₇, un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ;

R₇ et R₈ représentent chacun indépendamment un atome d'hydrogène, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ;

R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié,

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ et R₁₆ représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₁₋₃-CN, un groupe -C(O)-(CH₂)₀₋₃-R, un groupe -C(O)-(CH₂)₀₋₃-NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' sont tels que définis ci-avant ;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉ et R₃₀ représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₁₋₃-CN, un groupe -C(O)-(CH₂)₀₋₃-R', un groupe -C(O)-(CH₂)₀₋₃-NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' sont tels que définis ci-avant ; et où Y est tel que défini ci-avant.

4. Composé selon la revendication 3, dans lequel

R_a and R_b représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C₁-C₃ linéaire ou ramifié ;

n représente 0, 1 ou 2 ;

R₁ représente un groupe alkyle en C₁-C₄ linéaire ou ramifié, un groupe cycloalkyle C₃-C₇, un groupe phényle, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S et N, un groupe pyrrolidinyle, un groupe pipéridinyle, un groupe pipérazinyle, un groupe tétrahydropyranyle ou un groupe morpholinyle ;

où les groupes cycloalkyle, phényle, hétéroaryle, pyrrolidinyle, pipéridinyle, pipérazinyle, tétrahydropyranyle ou morpholinyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, ou groupe alkyle en C₁-C₄ linéaire ou ramifié, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃OR₈, un groupe -(CH₂)₀₋₃NR₇R₈, un groupe -C(O)-(CH₂)₀₋₃-R₈ ou un groupe -C(O)-(CH₂)₀₋₃-NR₇R₈ ; où R₇ et R₈ représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C₁-C₄ ;

R₂ représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₃, un groupe halogénoalkyle en C₁-C₃, un groupe cycloalkyle en C₃-C₄, un groupe -NH₂, un groupe -N(CH₃)H, un groupe -N(CH₃)₂ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C₁-C₂ ;


R₃ représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₃, un groupe halogénoalkyle en C₁-C₃, un groupe cycloalkyle en C₃-C₄, un groupe -NH₂, un groupe -N(CH₃)H, un groupe -N(CH₃)₂ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par

un groupe alcoxy en C₁-C₂ ;

R₄ représente un atome d'hydrogène, un groupe alcoxy en C₁-C₃, un groupe halogénoalkyle en C₁-C₃, un groupe cycloalkyle en C₃-C₄ ou un groupe alkyle en C₁-C₃ linéaire ou ramifié ;

R₆ représente un atome d'hydrogène, un atome d'halogène, un groupe alcoxy en C₁-C₃, un groupe halogénoalkyle en C₁-C₃, un groupe cycloalkyle en C₃-C₄, un groupe -NH₂, un groupe -N(CH₃)H, un groupe -N(CH₃)₂ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C₁-C₂ ;

R₅ représente un fragment de formule (II-a2), (IIb-1), (IIb-2) ou (IIIa-1) :



formule IIa-2

formule IIb-1

formule IIb-2

formule IIIa-1

où :

• R₉, R₁₀ et R₁₂ représentent chacun indépendamment un atome d'hydrogène, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R, un groupe -C(O)-(CH₂)₀₋₃NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ;

• R₁₃, R₁₄, R₁₅ et R₁₆ représentent chacun indépendamment un atome d'hydrogène, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₀₋₃NR'R'' ou un groupe -(CH₂)₀₋₃NR'R'' ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ;

• R₁₈ et R₂₃ représentent chacun indépendamment un atome d'hydrogène, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R, un groupe -C(O)-(CH₂)₀₋₃NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ;

• Y représente un groupe -NR'-, -O- ou -S- ; où R' représente un atome d'hydrogène ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; ou dans le cas où Y représente un groupe -NR'-, R₄, conjointement au groupe -NR'- et à l'atome de carbone auquel R₄ et le groupe -NR'- sont tous deux liés, forme un groupe hétérocyclyle contenant N saturé à 4 à 7 chaînons.

5. Composé selon la revendication 1, dans lequel R₁ représente un atome d'hydrogène, un groupe alkyle en C₁-C₄ linéaire ou ramifié, un groupe halogénoalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₁₀, un groupe cycloalcényle en C₃-C₁₀, un groupe aryle en C₆-C₁₄ monocyclique ou bicyclique, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N ou un groupe hétérocyclyle monocyclique ou bicyclique à 5 à 14 chaînons contenant au moins un hétéroatome choisi parmi O, S et N,

où les groupes cycloalkyle, cycloalcényle, aryle, hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C₁-C₄ linéaire ou ramifié, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₁₋₃CN, un groupe -(CH₂)₀₋₃OR₈, un groupe -(CH₂)₀₋₃NR₇R₈, un groupe -C(O)-(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R₈, un groupe -C(O)-(CH₂)₀₋₃NR₇R₈, un groupe -S(O)₂(CH₂)O₃R₈, un groupe -S(O)₂(CH₂)₀₋₃NR₇R₈ ou un groupe -(CH₂)₀₋₃(phényl)-OR₈ ; et

où R₁ représente de préférence un atome d'hydrogène, un groupe alkyle en C₁-C₃, un groupe halogénoalkyle en C₁-C₃, un groupe cycloalkyle en C₃-C₇, un groupe phényle, un groupe naphthyle, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S et N, ou un groupe hétérocyclyle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S et N,

où les groupes cycloalkyle, phényle, naphthyle, hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C₁-C₄ linéaire ou ramifié, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃OR₈, un groupe -(CH₂)₀₋₃NR₇R₈, un groupe -C(O)-(CH₂)₀₋₃R₈, un groupe -C(O)-(CH₂)₀₋₃NR₇R₈ ou un groupe -(CH₂)₀₋₃(phényl)-OR₈ ; où R₇ et R₈ sont tels que définis dans la revendication 1.

6. Composé selon l'une quelconque des revendications 1, 3 ou 5, dans lequel :

(a) R_1 représente un groupe alkyle en C_1-C_3 , un groupe cycloalkyle en C_3-C_7 , un groupe phényle, un groupe naphthyle, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S et N, ou un groupe hétérocyclyle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S and N,

où les groupes cycloalkyle, phényle, naphthyle, hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C_1-C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1-C_4 , un hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{0-3}OR_8$, un groupe $-(CH_2)_{0-3}NR_7R_8$, un groupe $-C(O)-(CH_2)_{0-3}R_8$ ou un groupe $-C(O)-(CH_2)_{0-3}NR_7R_8$; où R_7 et R_8 sont tels que définis dans la revendication 1 ; et

où R_1 représente de préférence un groupe cycloalkyle en C_3-C_7 , un groupe phényle, un groupe hétéroaryle monocyclique ou bicyclique à 5 à 10 chaînons contenant un, deux ou trois hétéroatomes choisis parmi O, S et N, un groupe pyrrolidinyle, un groupe pipéridinyle, un groupe pipérazinyle, un groupe tétrahydropyranyle ou un groupe morpholinyle,

où les groupes cycloalkyle, phényle, hétéroaryle, pyrrolidinyle, pipéridinyle, pipérazinyle, tétrahydropyranyle ou morpholinyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe alkyle en C_1-C_4 linéaire ou ramifié, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{0-3}OR_8$, un groupe $-(CH_2)_{0-3}NR_7R_8$, un groupe $-C(O)-(CH_2)_{0-3}R_8$ ou un groupe $-C(O)-(CH_2)_{0-3}NR_7R_8$; où R_7 et R_8 représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C_1-C_4 ; et

où R_1 représente, de manière davantage préférée, un groupe phényle ou un groupe pyridinyle, ce groupe phényle ou pyridinyle étant non substitué ou substitué par un, deux ou trois substituants choisis parmi un atome d'halogène, ou groupe alkyle en C_1-C_3 linéaire ou ramifié ou un groupe $-(CH_2)_{0-3}OCH_3$; et où, de préférence, lesdits groupes phényle et pyridinyle sont directement liés au groupe pyrrolotriazinone ; et/ou

(b) R_2 représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C_1-C_4 , un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{0-3}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_3 ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle ou un groupe alkyle en C_1-C_3 linéaire ou ramifié ; et

où R_2 représente de préférence un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe alcoxy en C_1-C_3 , un groupe halogénoalkyle en C_1-C_3 , un groupe cycloalkyle en C_3-C_4 , un groupe $-NH_2$, un groupe $-N(CH_3)H$, un groupe $-N(CH_3)_2$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_2 ; et/ou

(c) R_3 représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C_1-C_4 , un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{0-3}NR'R''$, un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_3 ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxy ou un groupe alkyle en C_1-C_3 linéaire ou ramifié ; et

où R_3 représente de préférence un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C_1-C_3 , un groupe halogénoalkyle en C_1-C_3 , un groupe cycloalkyle en C_3-C_4 , un groupe $-NH_2$, un groupe $-N(CH_3)H$, un groupe $-N(CH_3)_2$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_2 ; et/ou

(d) R_4 représente un atome d'hydrogène, un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{1-4}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_3 , un groupe $-C(O)-(CH_2)_{0-3}R$ ou un groupe $-C(O)-(CH_2)_{0-3}NR'R''$; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle ou un groupe alkyle en C_1-C_3 linéaire ou ramifié ; et

où R_4 représente de préférence un atome d'hydrogène, un groupe halogénoalkyle en C_1-C_3 , un groupe hydroxyalkyle en C_1-C_3 , un groupe cycloalkyle en C_3-C_4 ou un groupe alkyle en C_1-C_3 linéaire ou ramifié ; et/ou



(e) R_6 représente un atome d'hydrogène ; un atome d'halogène ; un groupe hydroxyle ; un groupe cyano ; un groupe alcoxy en C_1-C_4 ; un groupe halogénoalkyle en C_1-C_4 ; un groupe hydroxyalkyle en C_1-C_4 linéaire ou ramifié ; un groupe cycloalkyle en C_3-C_7 ; un groupe $-(CH_2)_{0-3}NR'R''$;

un groupe $-(CH_2)_{1-3}O$ (alkyle en C_1-C_4) ; un groupe $-(CH_2)_{0-3}OC(O)-(alkyle en C_1-C_4)$; un groupe $-(CH_2)O-3C(O)O-(alkyle en C_1-C_4)$; un groupe $-C(O)-(CH_2)_{0-3}NR'R''$; un groupe $-(CH_2)_{0-3}C(O)OH$; un groupe $-(CH_2)_{0-3}$ (hétéroaryle monocyclique ou bicyclique à 5 à 10 chaînons contenant au moins un hétéroatome choisi parmi O, S et N) ; un groupe $-(CH_2)_{0-3}$ (hétérocyclyle monocyclique ou bicyclique à 5 à 10 chaînons contenant au moins un hétéroatome choisi parmi O, S et N) ; ou un groupe alkyle en C_1-C_3 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1-C_4 , un groupe cyano ou un groupe cycloalkyle en C_3-C_4 ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle ou un groupe alkyle en C_1-C_3 linéaire ou ramifié ; et où les groupes hétéroaryle et hétérocyclyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alkyle en C_1-C_3 linéaire ou ramifié ou un groupe halogénoalkyle en C_1-C_3 ; et/ou (f) R_6 représente un atome d'hydrogène, un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe alcoxy en C_1-C_4 , un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{0-3}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_3 ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle ou un groupe alkyle en C_1-C_3 linéaire ou ramifié ; et où R_6 représente de préférence un atome d'hydrogène, un atome d'halogène, un groupe alcoxy en C_1-C_3 , un groupe halogénoalkyle en C_1-C_3 , un groupe cycloalkyle en C_3-C_4 , un groupe $-NH_2$, un groupe $-N(CH_3)H$, un groupe $-N(CH_3)_2$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un groupe alcoxy en C_1-C_2 .

7. Composé selon l'une quelconque des revendications 1, 3, 5 ou 6, dans lequel :

(a) R_5 représente un groupe choisi parmi

i) un groupe de formule (IIa-1) et ii) un groupe de formule (IIa-2)



formule (IIa-1)

formule (IIa-2)


où

R_9 , R_{10} , R_{11} et R_{12} représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C_1-C_4 , un groupe halogénoalkyle en C_1-C_4 , un groupe hydroxyalkyle en C_1-C_4 , un groupe cycloalkyle en C_3-C_4 , un groupe $-(CH_2)_{0-3}CN$, un groupe $-C(O)-(CH_2)_{1-3}CN$, un groupe $-C(O)-(CH_2)_{0-3}R$, un groupe $-C(O)-(CH_2)_{0-3}NR'R''$, un groupe $-(CH_2)_{0-3}NR'R''$ ou un groupe alkyle en C_1-C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1-C_4 , un groupe cyano ou un groupe cycloalkyle en C_3-C_4 ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C_1-C_4 ou un groupe alkyle en C_1-C_4 linéaire ou ramifié ; ou


(b) R_5 représente un groupe choisi parmi

i) un groupe de ii) un groupe de 5 iii) un groupe de formule (IIb-1), formule (IIb-2), formule (IIb-3), formule (IIb-1) formule (IIb-2) formule (IIb-3)


iv) un groupe de formule (IIb-4), et v) un groupe de formule (IIb-5),




formule (IIb-1)




formule (IIb-2)



formule (IIb-3)



formule (IIb-4)



formule (IIb-5)


où R_{13} , R_{14} , R_{15} et R_{16} représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène ; un groupe alcoxy en C_1 - C_4 ; un groupe halogénoalkyle en C_1 - C_4 ; un groupe hydroxyle en C_1 - C_4 ; un groupe cycloalkyle en C_3 - C_4 , un groupe $-(CH_2)_{0-3}CN$, un groupe $-C(O)-(CH_2)_{1-3}CN$; un groupe $-C(O)-(CH_2)_{0-3}R'$; un groupe $-C(O)-(CH_2)_{0-3}NRR''$; un groupe $-(CH_2)_{0-3}NR'R''$; ou un groupe alkyle en C_1 - C_4 linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C_1 - C_4 , un groupe cyano ou un groupe cycloalkyle en C_3 - C_4 ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C_1 - C_4 ou un groupe alkyle en C_1 - C_4 linéaire ou ramifié ; et où Y est tel que défini dans la revendication 1 ; ou dans le cas où Y représente un groupe $-NR'$, R_4 , conjointement au groupe $-NR'$ et à l'atome de carbone auquel R_4 et le groupe $-NR'$ sont tous deux liés, forme un groupe hétérocyclyle contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe $-CHF_2$ ou un groupe $-CF_3$; ou

(c) R_5 représente un groupe choisi parmi


i) un groupe de formule (IIIa-1),

ii) un groupe de formule (IIIa-2),


iii) un groupe de formule (IIIa-3),



formule (IIIa-1)




formule (IIIa-2)



formule (IIIa-3)


(continued)

iv) un groupe de formule (IIIa-4),




formule (IIIa-4)

v) un groupe de formule (IIIa-5),




formule (IIIa-5)

5 vi) un groupe de formule (IIIa-6),




formule (IIIa-6)

vii) un groupe de formule (IIIa-7),




formule (IIIa-7)

viii) un groupe de formule (IIIa-8),




formule (IIIa-8)

ix) un groupe de formule (IIIa-9),




formule (IIIa-9)

x) un groupe de formule (IIIa-10),




formule (IIIa-10)

xi) un groupe de formule (IIIa-11),




formule (IIIa-11)

xii) un groupe de formule (IIIa-12),




formule (IIIa-12)

xiii) un groupe de formule (IIIa-13),




formule (IIIa-13)

xiv) un groupe de formule (IIIa-14),




formule (IIIa-14)

xv) un groupe de formule (IIIa-15),




formule (IIIa-15)

xvi) un groupe de formule (IIIa-16),




formule (IIIa-16)

xvii) un groupe de formule (IIIa-17),



formule (IIIa-17)

xviii) un groupe de formule (IIIa-18),




formule (IIIa-18)

xix) un groupe de formule (IIIa-19),


xx) un groupe de 5 formule (IIIa-20),

xxi) un groupe de formule (IIIa-21),


(continued)



formule (IIIa-19)




formule (IIIa-20)



formule (IIIa-21)

xxii) un groupe de formule (IIIa-22),




formule (IIIa-22)

où


R₁₈, R₂₂, R₂₃ et R₂₄ représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R', un groupe -C(O)-(CH₂)₀₋₃NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; et où Y est tel que défini dans la revendication 1 ;

ou dans le cas où Y représente un groupe -NR'-, R₄, conjointement au groupe -NR'- et à l'atome de carbone auquel R₄ et le groupe -NR'- sont tous deux liés, forme un groupe hétérocyclye contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclye étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe -CHF₂ ou un groupe -CF₃ ; ou (d) R₅ représente un groupe choisi parmi


i) un groupe de formule (IIIb-1), ii) un groupe de formule (IIIb-2), 5 iii) un groupe de formule (IIIb-3),



formule (IIIb-1)




formule (IIIb-2)




formule (IIIb-3)

iv) un groupe de formule (IIIb-4), v) un groupe de formule (IIIb-5), 5 vi) un groupe de formule (IIIb-6),


(continued)



formule (IIIb-4)




formule (IIIb-5)




formule (IIIb-6)

vii) un groupe de formule (IIIb-7), viii) un groupe de formule (IIIb-8),



formule (IIIb-7)




formule (IIIb-8)

où


R₁₈, R₂₂, R₂₅, R₂₆ et R₂₇ représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R, un groupe -C(O)-(CH₂)₀₋₃NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; et où Y est tel que défini dans la revendication 1 ;

ou dans le cas où Y représente un groupe -NR'-, R₄, conjointement au groupe -NR'- et à l'atome de carbone auquel R₄ et le groupe -NR'- sont tous deux liés, forme un groupe hétérocyclye contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclye étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe -CHF₂ ou un groupe -CF₃ ; ou (e) R₅ représente un groupe choisi parmi

i) un groupe de formule (IIIc-1), et ii) un groupe de formule (IIIc-2),



Formule (IIIc-1)



Formule (IIIc-2)


où

R₁₉, R₂₀ et R₂₈ représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R', un groupe -C(O)-(CH₂)₀₋₃NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; et où Y est tel que défini dans la revendication 1 ;


ou dans le cas où Y représente un groupe -NR'-, R₄, conjointement à l'atome de carbone auquel R₄ et le groupe -NR'-sont tous deux liés, forme un groupe hétérocyclyle contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe -CHF₂ ou un groupe -CF₃ ; ou

(f) R₅ représente un groupe choisi parmi


i) un groupe de formule (IIId-1), ii) un groupe de formule (IIId-2), et iii) un groupe de formule (IIId-3),



formule (IIId-1)



formule (IIId-2)



formule (IIId-3)


où

R₂₁, R₂₈, R₂₉ et R₃₀ représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₄, un groupe halogénoalkyle en C₁-C₄, un groupe hydroxyalkyle en C₁-C₄, un groupe cycloalkyle en C₃-C₄, un groupe -(CH₂)₀₋₃CN, un groupe -C(O)(CH₂)₁₋₃CN, un groupe -C(O)-(CH₂)₀₋₃R, un groupe -C(O)-(CH₂)₀₋₃NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ou un groupe alkyle en C₁-C₄ linéaire ou ramifié, ce groupe alkyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un groupe alcoxy en C₁-C₄, un groupe cyano ou un groupe cycloalkyle en C₃-C₄ ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle, un groupe alcoxy en C₁-C₄ ou un groupe alkyle en C₁-C₄ linéaire ou ramifié ; et où Y est tel que défini dans la revendication 1 ;

ou dans le cas où Y représente un groupe -NR'-, R₄, conjointement à l'atome de carbone auquel R₄ et le groupe -NR'-sont tous deux liés, forme un groupe hétérocyclyle contenant N saturé à 4 à 7 chaînons, ce groupe hétérocyclyle étant non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe cyano, un groupe -CHF₂ ou un groupe -CF₃.

8. Composé selon la revendication 1, qui est :


(a) un composé de formule (Ia) :



Formule (Ia)

dans laquelle R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_a , R_b et n sont tels que définis dans l'une quelconque des revendications précédentes ; ou

(b) un composé de formule (Ib) :



Formule (Ib)

formule dans laquelle R_1 , R_2 , R_3 , R_4 , R_5 , R_a , R_b et n sont tels que définis dans l'une quelconque des revendications 1 à 7.

9. Composé selon la revendication 1, dans lequel :

X représente un atome d'azote ou un groupe $-CR_6$;

R_a et R_b représentent chacun indépendamment un atome d'hydrogène ou un groupe méthyle ;

R_1 représente un atome d'hydrogène, un atome d'halogène, un groupe halogénoalkyle en C_1 - C_3 , un groupe méthyle, un groupe cycloalkyle en C_3 - C_7 , un groupe phényle, un groupe pyridinyle, un groupe pyrazolyle, un groupe isoxazolyle, un groupe pipéridinyle ou un groupe tétrahydropyranyle ;

où les groupes cycloalkyle, phényle, pyridinyle, pyrazolyle, isoxazolyle, pipéridinyle ou tétrahydropyranyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe hydroxyle, un groupe halogénoalkyle en C_1 - C_3 , un groupe alkyle en C_1 - C_3 linéaire ou ramifié, un groupe $-(CH_2)-(phényl)-O-(alkyle\ en\ C_1-C_3)$, un groupe $-NR_7R_8$ ou un groupe $-OR_8$; où R_7 et R_8 représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C_1 - C_3 linéaire ou ramifié ;

R_2 et R_3 représentent chacun indépendamment un atome d'hydrogène, un atome d'halogène, un groupe cyano, un groupe halogénoalkyle en C_1 - C_3 ou un groupe alkyle en C_1 - C_3 linéaire ou ramifié ;

R_4 représente un atome d'hydrogène, un groupe halogénoalkyle en C_1 - C_3 , un groupe hydroxyalkyle en C_1 - C_3 ou un groupe alkyle en C_1 - C_3 linéaire ou ramifié ;

R_6 représente un atome d'hydrogène ; un atome d'halogène ; un groupe hydroxyle ; un groupe cyano ; un groupe alcoxy en Ci - C_4 ; un groupe halogénoalkyle en C_1 - C_4 ; un groupe hydroxyalkyle en C_1 - C_4 linéaire ou ramifié ; un groupe cycloalkyle en C_3 - C_7 ; un groupe alkyle en C_1 - C_3 linéaire ou ramifié ; un groupe $-(CH_2)_{0-3}NR'R''$; un groupe $-(CH_2)_{1-3}O(alkyle\ en\ C_1-C_3)$; un groupe $-(CH_2)_{0-3}OC(O)-(alkyle\ en\ C_1-C_3)$; un groupe $-(CH_2)_{0-3}C(O)O-(alkyle\ en\ C_1-C_3)$; un groupe $-C(O)-NR'R''$; un groupe $-(CH_2)_{0-3}C(O)OH$; un groupe -

(CH₂)₀₋₃-(imidazolyle) ; un groupe - (CH₂)₀₋₃- (oxazolyle) ; un groupe - (CH₂)₀₋₃- (oxadiazolyle) ; un groupe - (CH₂)₀₋₃- (pyrazolyle) ou un groupe - (CH₂)₀₋₃- (morpholinyle) ; où R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe hydroxyle ou un groupe alkyle en C₁-C₃ linéaire ou ramifié ; et

Où les groupes imidazolyle, oxazolyle, oxadiazolyle, pyrazolyle et morpholinyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe alkyle en C₁-C₃ linéaire ou ramifié ou un groupe halogénoalkyle en C₁-C₃,

R₅ représente un groupe choisi parmi :

i) un groupe de formule (IIa), ce groupe étant un groupe purinyle non substitué ou substitué par un groupe -NR'R'' ;

ii) un groupe de formule (IIb), ce groupe étant choisi parmi un groupe -NR'-pyridinyle, un groupe -S-pyridinyle, un groupe -NR'-pyrimidinyle, un groupe -S-pyrimidinyle ou un groupe -NR'-triazinyle ; où les groupes pyridinyle, pyrimidinyle et triazinyle sont non substitués ou substitués par un, deux ou trois substituants choisis parmi un atome d'halogène, un groupe halogénoalkyle en C₁-C₃, un groupe - (CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₀₋₃-NR'R'', un groupe -(CH₂)₀₋₃NR'R'' ; et

iii) un groupe de formule (IIc), ce groupe étant choisi parmi un groupe -NR'-purinyle, un groupe -S-purinyle, un groupe -NR'-7H-pyrrolo[2,3-d]pyrimidinyle, un groupe -NR'-1H-pyrazolo[3,4-d]pyrimidinyle ou un groupe -NR'-pyrazolo[1,5-a]pyrimidinyle ; où les groupes purinyle, 7H-pyrrolo[2,3-d]pyrimidinyle, 1H-pyrazolo[3,4-d]pyrimidinyle, pyrazolo[1,5-a]pyrimidinyle sont non substitués ou substitués par un atome d'halogène ou un groupe -(CH₂)₀₋₃NR'R'' ; ou

R₄ et R₅, conjointement à l'atome de carbone auquel ils sont attachés, forment un groupe pyrrolidinyl-purinyle ou un groupe pyrrolidinyl-pyrimidinyle ; où les groupe pyrrolidinyle est non substitué ou substitué par un ou plusieurs substituants choisis parmi un atome d'halogène ou un groupe hydroxyle ; et où le groupe purinyle est non substitué ou substitué par un groupe -(CH₂)₀₋₃NR'R'' ; et où le groupe pyrimidinyle est non substitué ou substitué par un, deux ou trois substituants choisis parmi un groupe - (CH₂)₀₋₃CN ou un groupe - (CH₂)₀₋₃NR'R'' ; et

R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₃ ou un groupe alkyle en C₁-C₃ linéaire ou ramifié.

10. Composé selon la revendication 1 ou la revendication 3, dans lequel :

X représente un atome d'azote ou un groupe -CR₆ ;

R_a et R_b représentent chacun indépendamment un atome d'hydrogène ou un groupe méthyle ;

R₁ représente un groupe méthyle, un groupe cycloalkyle en C₃-C₇, un groupe phényle, un groupe pyridinyle, un groupe pipéridinyle ou un groupe tétrahydropyranyle ;

où les groupes cycloalkyle, phényle, pyridinyle, pipéridinyle ou tétrahydropyranyle sont non substitués ou substitués par un ou plusieurs substituants choisis parmi un atome d'halogène, un groupe alkyle en C₁-C₃ linéaire ou ramifié, un groupe -NR₇R₈ ou un groupe -OR₈ ; où R₇ et R₈ représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C₁-C₃ linéaire ou ramifié ;

R₂ et R₃ représentent chacun indépendamment un atome d'hydrogène ou un groupe alkyle en C₁-C₃ linéaire ou ramifié ;

R₄ représente un atome d'hydrogène, un groupe halogénoalkyle en C₁-C₃ ou un groupe alkyle en C₁-C₃ linéaire ou ramifié ;

R₆ représente un atome d'hydrogène, un atome d'halogène, un groupe halogénoalkyle en C₁-C₃, un groupe alkyle en C₁-C₃ linéaire ou ramifié ou un groupe cyclopropyle ;

R₅ représente un groupe choisi parmi :

i) un groupe de formule (IIa), ce groupe étant un groupe purinyle non substitué ou substitué par un groupe -NR'R'' ;

ii) un groupe de formule (IIb), ce groupe étant choisi parmi un groupe -NH-pyridinyle, un groupe -S-pyridinyle, un groupe -NH-pyrimidinyle ou un groupe -S-pyrimidinyle ; où les groupes pyridinyle ou pyrimidinyle sont non substitués ou substitués par un, deux ou trois substituants choisis parmi un groupe -(CH₂)₀₋₃CN, un groupe -C(O)-(CH₂)₀₋₃-NR'R'' ou un groupe -(CH₂)₀₋₃NR'R'' ; et

iii) un groupe de formule (IIc), ce groupe étant choisi parmi un groupe -NH-purinyle ou un groupe -S-purinyle ; où les groupe purinyle est non substitué ou substitué par un groupe -(CH₂)₀₋₃NR'R'' ; ou

R₄ et R₅, conjointement à l'atome de carbone auquel ils sont attachés, forment un groupe pyrrolidinyl-purinyle,

où le groupe purinyle est non substitué ou substitué par un groupe - (CH₂)₀₋₃NR'R'' ;

R' et R'' représentent chacun indépendamment un atome d'hydrogène, un groupe alcoxy en C₁-C₃ ou un groupe alkyle en C₁-C₃ linéaire ou ramifié.

5 11. Composé selon la revendication 1, qui est l'un des suivants :

- 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-aminopyrimidin-4-ylamino)méthyl)-5-chloro-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-5-cyclopropyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 10 2-((6-amino-9H-purin-9-yl)méthyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-aminopyrimidin-4-ylamino)méthyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 4-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)méthylamino)picolinamide ;
 2-((2-aminopyridin-4-ylamino)méthyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((9H-purin-6-ylamino)méthyl)-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 15 2-((6-amino-9H-purin-9-yl)méthyl)-3-cyclohexylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-5-méthyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((9H-purin-6-ylthio)méthyl)-5-méthyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-6-méthyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((9H-purin-6-ylthio)méthyl)-6-méthyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 20 2-(1-(6-amino-9H-purin-9-yl)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(6-aminopyrimidin-4-ylamino)propyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)propyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-
 25 carbonitrile ;
 (R)-2-(1-(9H-purin-6-ylamino)propyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(6-aminopyrimidin-4-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 30 (S)-4-amino-6-(1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-
 carbonitrile ;
 2-(1-(6-amino-9H-purin-9-yl)éthyl)-5-méthyl-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-3-o-tolyl-5-(trifluorométhyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 35 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(3-méthoxyphényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(2,4-difluorophényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-3-benzyl-5-chloropyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-3-phénylimidazo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-3-o-tolylimidazo[1,2-f][1,2,4]triazin-4(3H)-one ;
 40 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(pyridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(tétrahydro-2H-pyran-4-yl)pyrrolo[1,2-f][1,2,4]triazin-
 4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(1-méthylpipéridin-4-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3-fluorophényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 45 (S)-4-amino-6-(1-(3-(3-fluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-
 5-carbonitrile ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimi-
 dine-5-carbonitrile ;
 50 2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-méthylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((6-amino-9H-purin-9-yl)méthyl)-3-((1r,4r)-4-aminocyclohexyl)-5-chloropyrrolo[1,2-f][1,2,4]triazin-
 4(3H)-one ;
 (R)-2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(1-phényléthyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(1-phényléthyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 55 (S)-4-amino-6-(1-(4-oxo-3-(pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-
 carbonitrile ;
 (S)-2-(1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(4-oxo-3-phényl-5-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)py-

rimidine-5-carbonitrile ;

(S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-phényl-5-(trifluorométhyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(1-(5-(difluorométhyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

5 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-5-(difluorométhyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-phénylimidazo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(1-(4-oxo-3-phényl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

2-(1-(9H-purin-6-ylamino)-3,3,3-trifluoropropyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

10 4-amino-6-(3,3,3-trifluoro-1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(2-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile ;

(S)-3-phényl-2-(1-(pyrazolo[1,5-a]pyrimidin-7-ylamino)éthyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

15 2-((6-amino-9H-purin-9-yl)méthyl)-5-(difluorométhyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(2-amino-9H-purin-6-yl)pyrrolidin-2-yl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(4,6-diamino-1,3,5-triazin-2-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-((6-amino-9H-purin-9-yl)méthyl)-5-chloro-3-(1-(5-fluoropyridin-2-yl)éthyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

20 (S)-2-(1-(2-amino-9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

(S)-2-(1-(9H-purin-6-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

(R)-2-(1-(9H-purin-6-ylamino)-2-hydroxyéthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

25 (R)-4-amino-6-(2-hydroxy-1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-2-(1-(2-amino-9H-purin-6-ylamino)éthyl)-3-phénylimidazo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(méthyl(1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthyl)amino)pyrimidine-5-carbonitrile ;

30 (S)-2-(1-(méthyl(9H-purin-6-yl)amino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(9H-purin-6-ylamino)éthyl)-5-méthyl-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(1-(5-méthyl-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

35 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-7-méthyl-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(1-(7-méthyl-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-2-(4,4-difluoro-1-(9H-purin-6-yl)pyrrolidin-2-yl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(4,4-difluoro-2-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)pyrrolidin-1-yl)pyrimidine-5-carbonitrile ;

40 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-6-fluoro-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(1-(6-fluoro-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

2-((S)-1-(9H-purin-6-ylamino)éthyl)-3-((S)-1-phényléthyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

45 4-amino-6-((S)-1-(4-oxo-3-((S)-1-phényléthyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(3-(2,6-diméthylphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-2-((9H-purin-6-ylamino)méthyl)-3-(1-phényléthyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

50 (S)-4-amino-6-((4-oxo-3-(1-phényléthyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)méthylamino)pyrimidine-5-carbonitrile ;

(S)-2-(1-(5-fluoro-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(2,6-diméthylphényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-4-amino-6-(1-(5-fluoro-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

55 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-5-fluoro-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

(S)-4-amino-6-(1-(3-(3,5-difluorophényl)-4-oxo-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 5 4-amino-6-((1S)-1-(5-(1,2-dihydroxyéthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-4-amino-6-(1-(3-(3,5-difluorophényl)-4-oxo-5-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)-5-(trifluorométhyl)pyrrolo[1,2-f][1,2,4]triazin-
 10 4(3H)-one ;
 (S)-4-amino-6-(1-(5-(hydroxyméthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-(trifluorométhyl)pyrimidin-4-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-(pyridin-2-ylméthyl)-3,4-dihydropyrrolo[1,2-
 15 f][1,2,4]triazine-5-carbonitrile ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-5-(difluorométhyl)-3-(3,5-difluorophényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)imidazo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(5-(difluorométhyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-
 20 yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)éthyl)-5-(difluorométhyl)-3-(3,5-difluorophényl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(2-amino-9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 25 2-(1-(9H-purin-6-ylamino)-2,2,2-trifluoroéthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(3-benzyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-fluoropyrimidin-4-ylamino)éthyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(6-amino-5-fluoropyrimidin-4-ylamino)éthyl)-5-(difluorométhyl)-3-(3,5-difluorophényl)pyrrolo[1,2-
 30 f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)propyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3,5-dichlorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 35 (S)-2-(1-(6-amino-5-fluoropyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-(trifluorométhyl)pyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (R)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)-2-hydroxyéthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyr-
 40 rolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-carbamoylpyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide ;
 45 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(2-chlorobenzyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-((S)-tétrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (R)-4-amino-6-(1-(3-(3,5-difluorophényl)-4-oxo-5-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-
 50 2-hydroxyéthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(2-amino-5-fluoropyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (S)-2-(1-(2-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 55 ((S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)-5-(2H-tétrazol-5-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(3-((5-méthylisoxazol-3-yl)méthyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(4-oxo-3-phényl-7-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-7-carbonitrile ;

5 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(1-(4-méthoxybenzyl)-1H-pyrazol-4-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

(S)-4-amino-6-(1-(4-oxo-3-phényl-5-(thiazol-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

10 (S)-2-(1-(2,6-diamino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

(S)-4-amino-6-(1-(5-(morpholinométhyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-((R)-1-phényléthyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

15 (S)-4-amino-6-(1-(4-oxo-3-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(3,5-difluorophényl)-5-(5-méthyl-1,2,4-oxadiazol-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

20 4-amino-6-((S)-1-(4-oxo-3-((S)-tétrahydro-2H-pyran-3-yl)-5-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(3-(5-méthyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

Acide

25 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylique ;

2-((S)-1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-((R)-tétrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

(S)-4-amino-6-(1-(3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

30 (S)-4-amino-6-(1-(4-oxo-3-(1H-pyrazol-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(4-oxo-3-(pyrimidin-5-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

35 4-amino-6-((S)-1-(4-oxo-3-((R)-tétrahydro-2H-pyran-3-yl)-5-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-2,4-diamino-6-(1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ; (S)-4-(1-(3-((1H-Pyrazol-3-yl)méthyl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)-6-aminopyrimidine-5-carbonitrile ;

40 (S)-4-amino-6-(1-(4-oxo-3-(tétrahydro-2H-pyran-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(4-oxo-3-(2,2,2-trifluoroéthyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(3-cyclobutyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

45 (S)-2-amino-4-(1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

4-amino-6-(1-(5-(1-méthyl-1H-pyrazol-4-yl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

50 (S)-4-amino-6-(1-(3-cyclopropyl-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

(S)-4-amino-6-(1-(5-bromo-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

4-amino-6-((S)-1-(4-oxo-3-((R)-tétrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

55 (S)-4-amino-6-(1-(5-bromo-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;

2-((3-iodo-1H-pyrazolo[3,4-d]pyrimidin-4-ylamino)méthyl)-5-méthyl-3-o-tolylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;

(S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 4-amino-6-((S)-1-(4-oxo-3-((S)-tétrahydro-2H-pyran-3-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 5 (S)-4-amino-6-(1-(4-oxo-3-phényl-5-(1H-pyrazol-4-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-4-amino-6-(1-(3-(isoxazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-N,N-diméthyl-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide ; (S)-4-amino-6-(1-(3-(1-méthyl-1H-pyrazol-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ; (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-N-propyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxamide ; 2-((S)-1-(9H-purin-6-ylamino)éthyl)-3-(tétrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ; 2-((S)-1-(9H-purin-6-ylamino)éthyl)-3-((S)-tétrahydro-2H-pyran-3-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ; (S)-4-amino-6-(3-hydroxy-1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile ; (S)-2-(1-(9H-purin-6-ylamino)-3-hydroxypropyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ; (R)-4-amino-6-(1-(3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-2-hydroxyéthylamino)pyrimidine-5-carbonitrile ; 4-amino-6-((4-oxo-3-o-tolyl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)méthylamino)pyrimidine-5-carbonitrile ; (S)-4-amino-6-(1-(5-(2-hydroxyéthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ; (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)-3-hydroxypropyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ; (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(5-fluoropyridin-3-yl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 20 (S)-4-amino-6-(1-(5-(2-méthylloxazol-5-yl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 25 (S)-4-amino-6-(1-(5-(2-méthoxyéthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-propyl-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carboxylate ;
 (S)-4-amino-6-(3-hydroxy-1-(4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile ;
 30 (S)-2-(1-(9H-purin-6-ylamino)-3-hydroxypropyl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(3-(3,5-difluorophényl)-4-oxo-5-(trifluorométhyl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-3-hydroxypropylamino)pyrimidine-5-carbonitrile ;
 (S)-4-amino-6-(1-(4-oxo-3-(6-(trifluorométhyl)pyridin-2-yl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 35 (S)-4-amino-6-(1-(5-bromo-4-oxo-3-(3-(trifluorométhyl)phényl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 acétate de (S)-2-(2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-5-yl)éthyle ;
 40 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(6-(trifluorométhyl)pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 2-((2S,4R)-1-(6-amino-5-cyanopyrimidin-4-yl)-4-hydroxypyrrolidin-2-yl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 4-amino-6-((2S,4R)-2-(5-(aminométhyl)-3-(3,5-difluorophényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)-4-hydroxypyrrolidin-1-yl)pyrimidine-5-carbonitrile ;
 45 (S)-4-amino-6-(1-(5-(4-méthyl-1H-imidazol-1-yl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-4-amino-6-(1-(5-bromo-3-(3-méthoxyphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-(3-(trifluorométhyl)phényl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 50 (S)-4-amino-6-(1-(5-bromo-3-(3-hydroxyphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-4-amino-6-(1-(3-(3-méthoxyphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 55 (S)-4-amino-6-(1-(3-(3-hydroxyphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3-méthoxyphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;

4-amino-6-(1-(4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)cyclopropylamino)pyrimidine-5-carbonitrile ;
 2-(1-(9H-purin-6-ylamino)cyclopropyl)-3-phénylpyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(4-oxo-3-(trifluorométhyl)phényl)-3,4-dihydropyrrolo[1,2-f][1,2,4]triazin-2-yl)éthylamino)pyrimidine-5-carbonitrile ;
 (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-3-(3-hydroxyphényl)-4-oxo-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile ;
 (S)-2-(1-(9H-purin-6-ylamino)éthyl)-3-(pyridin-2-yl)pyrrolo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-2-(1-(9H-purin-6-ylamino)propyl)-3-phénylimidazo[1,2-f][1,2,4]triazin-4(3H)-one ;
 (S)-4-amino-6-(1-(4-oxo-3-phényl-3,4-dihydroimidazo[1,2-f][1,2,4]triazin-2-yl)propylamino)pyrimidine-5-carbonitrile ;

ou sel, solvate ou N-oxyde ou stéréoisomère et dérivé deutéré pharmaceutiquement acceptable de celui-ci.

12. Composé selon la revendication 1 ou la revendication 11, qui est le (S)-2-(1-(6-amino-5-cyanopyrimidin-4-ylamino)éthyl)-4-oxo-3-phényl-3,4-dihydropyrrolo[1,2-f][1,2,4]triazine-5-carbonitrile, ou sel ou solvate ou N-oxyde ou stéréoisomère ou dérivé deutéré pharmaceutiquement acceptable de celui-ci.

13. Composé selon l'une quelconque des revendications 1 à 12, à utiliser dans le traitement d'une affection pathologique ou d'une maladie susceptible d'amélioration par l'inhibition de la phosphoinositide 3-kinase (PI3K), l'affection pathologique ou maladie étant choisie parmi les maladies respiratoires ; les maladies allergiques ; les maladies inflammatoires ou auto-immunes ; les troubles fonctionnels et les troubles neurologiques ; les maladies cardiovasculaires ; les infections virales ; les troubles de la fonction métabolique/endocrinienne ; les troubles neurologiques et la douleur ; les rejets de greffes de moelle osseuse et d'organes ; le syndrome myélodysplasique ; le syndrome myéloprolifératif (SM) ; le cancer et les malignités hématologiques, la leucémie, les lymphomes et les tumeurs solides, où l'affection pathologique ou la maladie étant choisie parmi la leucémie, les lymphomes et les tumeurs solides, la polyarthrite rhumatoïde, la sclérose en plaques, la sclérose latérale amyotrophique, la maladie de Crohn, la rectocolite hémorragique, le lupus érythémateux disséminé, l'anémie hémolytique auto-immune, le diabète de type I, la vascularite cutanée, le lupus érythémateux cutané, la dermatomyosite, les maladies bulleuses, comprenant sans s'y limiter, le pemphigus vulgaire, la pemphigoïde bulleuse et l'épidermolyse bulleuse, l'asthme, la broncho-pneumopathie chronique obstructive, la mucoviscidose, la fibrose pulmonaire idiopathique, la sarcoïdose, la rhinite allergique, la dermatite atopique, la dermatite de contact, l'eczéma, le psoriasis, le carcinome basocellulaire, le carcinome épidermoïde et la kératose actinique.

14. Composition pharmaceutique comprenant un composé tel que défini dans les revendications 1 à 12 en association avec un diluant ou un support pharmaceutiquement acceptable.

15. Produit de combinaison comprenant (i) un composé tel que défini dans l'une quelconque des revendications 1 à 12 ; et (ii) un autre composé choisi dans le groupe constitué par un agoniste de l'adénosine A2A, un agent permettant de traiter les troubles cardiovasculaires, un agent permettant de traiter le diabète et un agent permettant de traiter une maladie du foie, un agent anti-allergique, un agent anticholinergique, un agent anti-inflammatoire, un agent anti-infectieux, un agent agoniste β 2-adrénérergique, une molécule homologue du récepteur chimiotactique exprimée sur l'inhibiteur des cellules TH₂ (CRTH2), un agent chimiothérapeutique, un corticostéroïde, un inhibiteur IKK β /IKBKB (IkB kinase bêta ou IKK2), un immunosuppresseur, un inhibiteur de Janus kinase (JAK), un inhibiteur de protéine kinase activée par le mitogène p38 (MAPK p38) à action topique, un inhibiteur de phosphodiesterase (PDE) IV et un inhibiteur de la tyrosine kinase de la rate (Syk), pour une utilisation simultanée, séparée ou séquentielle dans le traitement du corps humain ou animal.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2010151735 A2 [0534] [0553] [0559]
- WO 2005047279 A [0628] [0642]
- WO 2008077639 A [0738]
- WO 2009021696 A [0738] [0744] [0746]
- WO 2007124898 A [0741]
- WO 2006122788 A1 [0741]
- WO 2008046598 A [0741]
- WO 2008095720 A [0741]
- WO 2009068177 A [0741]
- WO 2010072354 A [0741]
- WO 03097613 PCT [0743]
- WO 2004058729 A [0743]
- WO 2005049581 A [0743]
- WO 2005123693 A [0743]
- WO 2005123692 A [0743]
- WO 2010069504 A [0743]
- WO 2008077639 PCT [0744] [0746]
- WO 2009153043 A [0744] [0746]
- WO 2010083975 A [0744] [0746]
- US 6106864 A [0782]
- WO 0035298 A [0782]
- WO 9203175 A [0791]
- WO 9102558 A [0791]
- GB 2242134 A [0791]
- EP 0069715 A [0793]
- GB 2041763 A [0793]
- EP 0424790 A [0793]
- DE 4239402 [0793]
- EP 0674533 A [0793]
- EP 0166294 A [0793]
- GB 2165159 A [0793]
- WO 9209322 A [0793]
- WO 9200771 A [0793]
- US 5201308 A [0793]
- WO 9700703 A [0793]
- EP 0505321 A [0793]
- WO 9204068 A [0793]
- WO 9204928 A [0793]
- WO 97000703 A [0793] [0798]
- WO 03000325 A [0793]
- WO 2006008027 A [0793]
- WO 9114468 A [0799]
- WO 9712687 A [0799]
- WO 9632150 A [0804]


Non-patent literature cited in the description

- VANHAESEBROECK B. *Nat Rev Mol Cell Biol*, 2010, vol. 5, 11381-6 [0001]
- KOK K. *Trends Biochem Science*, 2009, vol. 34, 115-127 [0003]
- JI H. *Blood*, 2007 [0004]
- OKKENHAUG K. *Science*, 2002 [0004]
- SOOND DR. *Blood* [0004]
- *Blood*, 03 June 2010 [0004]
- WILLIAM O. *Chemistry & Biology*, 2010, 17 [0004]
- SOOND DR. *Blood*, 2010 [0004]
- JARMIN SJ. *JCI*, 2008 [0004] [0007]
- BILANCIO A. *Blood*, 2006, 107 [0004]
- DIL N. *Mol Immunol*, 2009, 46 [0004]
- ALI K. *Nature*, 2004, vol. 431, 1007-1011 [0005]
- ALI K. *Journal of Immunology*, 2008, vol. 180, 2538-2544 [0005]
- LANNUTTI BJ. *Blood*, October 2002 [0005]
- SADHU C. *Jl*, 2003, 170 [0006]
- PURI KD. *Blood*, 2004, 103 [0006]
- SADHU C. *Biochem Biophys Res Comm*, 2003, 308 [0006]
- CONDLIFFE AM. *Blood*, 2005, 106 [0006] [0007]
- MARTIN AL. *Jl*, 2008, 180 [0007]
- THOMAS MS. *J Leukoc Biol*, 2008, 84 [0007]
- MATTHEW T. *Immunology*, 2008, 126 [0007]
- YUM HK. *Jl*, 2001, 167 [0007]
- PINHO V. *Jl*, 2007, 179 [0007]
- NASHED. *Eur J Immunol*, 2007, vol. 37, 416 [0009]
- LEE et al. *FASEB J*, 2006, vol. 20, 455 [0009]
- LEE KS et al. *J Allergy Clin Immunol*, 2006, vol. 118, 403 [0009]
- DOUKAS J. *JPET*, 2009, vol. 328, 758 [0009]
- PAR SJ. *ERJ*, 2010 [0009]
- PURI KD. *Blood*, 2004, vol. 103, 3448 [0009]
- MARWICK JA. *JRCCM*, 2009, vol. 179, 542-548 [0009]
- TO Y. *AJRCCM*, 2010, vol. 182, 897-904 [0009]
- TO Y. *AJRCCM*, 2010 [0009]
- MARWICK JA. *JACI*, 2010, vol. 125, 1146-53 [0009]
- SOOND DR. *Blood*, January 2010 [0010]
- HAYLOCK-JACOBS S. *J. Autoimmun*, 2010 [0010]
- RANDIS TM. *Eur J Immunol*, 2008, 38 [0010]
- PINHO JF. *Br. J. Pharmacol*, 2010, 161 [0010]
- HERMANN SE. *Blood*, 2010, vol. 116, 2078-88 [0011]
- LANNUTTI BJ. *Blood*, October 2010 [0011]

- **ERNEST L. ELIEL.** Stereochemistry of Organic Compounds. Wiley, 1994 [0050]
- **OKI, M.** *Topics in Stereochemistry*, 1983, 1 [0051]
- Pro-drugs as Novel Delivery Systems. ACS Symposium Series, vol. 14 [0069]
- Bioreversible Carriers in Drug Design. Pergamon Press, 1987 [0069]
- **H. BUNDGAARD.** Design of Prodrugs. Elsevier, 1985 [0070]
- **GRAY et al.** *Anal Biochem*, 2003, vol. 313, 234-45 [0075]
- **T. W. GREENE ; G. M. WUTS.** Protecting Groups in Organic Synthesis. Wiley, 1999 [0142] [0157]
- **TEHRANI, A. K. ; BORREMANS D ; DE KIMPE N.** *Tetrahedron*, 1999, vol. 55, 4133-4152 [0726]
- **OHTA, T. ; FUKUDA, T. ; ISHIBASHI, F. ; IWAO, M.** *J. Org. Chem.*, 2009, vol. 74, 8143-8153 [0726]
- **LEROY, J. ; PORTHIEL, E. ; BONDON, A.** *Tetrahedron*, 2002, vol. 58, 6713-6722 [0726]
- *Eur Respir J*, 2003, vol. 22 (45 [0743]
- Remington: The Science and Practice of Pharmacy. Lippincott Williams & Wilkins, 2001 [0767] [0769]
- **H. LIEBERMAN ; L. LACHMAN.** Pharmaceutical Dosage Forms: Tablets. Marcel Dekker, 1980, vol. 1 [0779]
- **VERMA et al.** *Pharmaceutical Technology On-line*, 2001, vol. 25 (2), 1-14 [0782]
- **FINNIN ; MORGAN.** *J Pharm Sci*, October 1999, vol. 88 (10), 955-958 [0810]

PIRROLO-TRIAZINON SZÁRMAZÉKOK PI3K INHIBITORKÉNT
SZABADALMI IGÉNYPONTOK

1. A formula (I) szerinti vegyület vagy annak egy gyógyszerészetileg elfogadható sója, szolvátja, N-oxidja, sztereoizomere vagy deuterizált származéka:



Formula (I)

ahol,

X nitrogénatomot vagy -CR₆ csoportot jelöl;

n 0-t, 1-et, 2-t vagy 3-at jelöl;

R₅ és R₆ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelölnek;

R₁ hidrogénatomot, lineáris vagy elágazó C₁-C₄ alkilcsoportot, C₁-C₄ haloalkil-csoportot, C₃-C₁₀ gyűrűs alkilcsoportot, C₃-C₁₀ cikloalkenil-csoportot, monociklusos vagy biciklusos C₆-C₁₄ arilcsoportot, 5-14 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely legalább egy heteroatomot tartalmaz O, S és N közül kiválasztva, vagy egy 5-14 tagú monociklusos vagy biciklusos heterociklikus csoportot jelöl, mely legalább egy heteroatomot tartalmaz O, S és N közül kiválasztva, ahol a gyűrűs alkil, cikloalkenil, aril, heteroaril és heterociklikus csoportok szubsztituátlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, hidroxilcsoportok, cianocsoportok, lineáris vagy elágazó C₁-C₄ alkilcsoportok, C₁-C₄ haloalkil-csoportok, C₁-C₄ hidroxialkil-csoportok, C₃-C₄ gyűrűs alkilcsoportok, -(CH₂)₁₋₃CN csoportok, -(CH₂)₀₋₃OR₈ csoportok, a -(CH₂)₀₋₃NR₇R₈ csoportok, -C(O)-(CH₂)₁₋₃CN csoportok, -C(O)-(CH₂)₀₋₃R₈ csoportok, -C(O)-(CH₂)₀₋₃NR₇R₈ csoportok, -S(O)₂(CH₂)₀₋₃R₈ csoportok, -S(O)₂(CH₂)₀₋₃NR₇R₈ csoportok vagy -(CH₂)₀₋₃(fenil)-OR₈ csoportok közül lettek kiválasztva;

R₂ és R₃ mindegyike egymástól függetlenül hidrogén atomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, -NR¹R² csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituátlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxycsoportok, cianocsoportok vagy C₃-C₇ gyűrűs alkilcsoportok közül lettek kiválasztva;

R₄ hidrogén atomot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, -(CH₂)₁₋₄NR¹R² csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituátlan vagy

szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoporthok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok, -C(O)-(CH₂)₀₋₃-R csoportok vagy -C(O)-(CH₂)₀₋₃-NR'R'' csoportok közül lettek kiválasztva;

R₆ hidrogénatomot; halogénatomot; hidroxilcsoportot; cianocsoportot; C₁-C₄ alkoxicsoporthot; a C₁-C₄ haloalkil-csoportot; lineáris vagy elágazó C₁-C₄ hidroxialkil-csoportot; C₃-C₇ gyűrűs alkilcsoportot; -(CH₂)₀₋₃NR'R'' csoportot; -(CH₂)₀₋₃O(C₁-C₄ alkilcsoportot); -(CH₂)₀₋₃OC(O)-(C₁-C₄ alkilcsoportot); -(CH₂)₀₋₃C(O)O-(C₁-C₄ alkilcsoportot); -C(O)-(CH₂)₀₋₃-NR'R'' csoportot; -(CH₂)₀₋₃C(O)OH csoportot jelöl vagy; olyan -(CH₂)₀₋₃-(5-14 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz legalább egy heteroatomot O, S és N közül kiválasztva) vagy; olyan -(CH₂)₀₋₃-(5-14 tagú monociklusos vagy biciklusos heterociklikus-csoportot jelöl, amely tartalmaz legalább egy heteroatomot O, S és N közül kiválasztva); vagy egy olyan lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több szubsztituenssel C₁-C₄ alkoxicsoporthok, cianocsoportot vagy C₃-C₄ gyűrűs alkilcsoportok közül kiválasztva;


ahol a heteroaril és heterociklikus csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, hidroxilcsoportok, cianocsoportok, lineáris vagy elágazó C₁-C₄ alkilcsoportok vagy C₁-C₄ haloalkil-csoportok közül lettek kiválasztva,

R₇ és R₈ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoporthok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva;

R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C₁-C₄ alkoxicsoporthot vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelölnek.


R₉ az alábbiak közül választott csoportot jelöl:

formula (IIa) csoport



formula (IIa)

formula (IIb) csoport



formula (IIb)

és

formula (IIc) csoport



formula (IIc)

ahol

Y olyan linkert jelöl, mely -NR'- csoport, -O- vagy -S- közül lett kiválasztva; ahol R' a fentiek szerint meghatározott;

(*) jelöli a helyet, ahol R₅ kapcsolódik az R₄-hez kapcsolódó szénatomhoz és a pirrolo-triazinon csoporthoz;

W₁ -CR₁₁ csoportot jelöl, és W₂ nitrogénatomot jelöl, vagy W₁ nitrogénatomot jelöl, és W₂ -CR₁₂ csoportot jelöl;

G₁ egy -CR₁₄ csoportot jelöl, és G₂ nitrogénatomot jelöl, vagy G₁ nitrogénatomot jelöl, és G₂ pedig -CR₁₅ csoportot jelöl; vagy G₁ egy -CR₁₄ csoportot jelöl, és G₂ pedig -CR₁₅ csoportot jelöl;


G₃ nitrogénatomot vagy -CR₁₆ csoportot jelöl;

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅ és R₁₆ mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroalkil csoportot, C₃-C₄ gyűrűs alkilcsoportot; -(CH₂)₀₋₃CN csoportot;


-C(O)-(CH₂)₀₋₃CN csoportot; -C(O)-(CH₂)₀₋₃R' csoportot; -C(O)-(CH₂)₀₋₃NR'R'' csoportot; -(CH₂)₀₋₃NR'R'' csoportot; vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituáltnak vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxycsoportok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' a fentiek szerint meghatározott;

R₁₇ az alábbiak közül választott csoportot jelöl

a) formula (IIla) csoport, b) formula (IIlb) csoport,




formula (IIlb)




formula (IIla)

a) formula (IIlc) csoport, b) formula (IIld) csoport,



formula (IIlc)



formula (IIld)

ahol

G₄ nitrogénatomot vagy -CR₂₂ csoportot jelöl;

G₅ és G₆ mindegyike egymástól függetlenül nitrogénatomot vagy szénatomot jelöl, ahol ha a G₅ és G₆ valamelyike nitrogénatomot jelöl, akkor a másik fennmaradó elem szénatomot jelöl.

G₇ -NH csoportot vagy -CH csoportot jelöl;

G₈ nitrogénatomot vagy -CR₂₃ csoportot jelöl;

G₉ nitrogénatomot vagy -CR₂₄ csoportot jelöl;

G₁₀ nitrogénatomot vagy -CR₂₅ csoportot jelöl;

G₁₁ nitrogénatomot vagy -CR₂₆ csoportot jelöl;

G₁₂ nitrogénatomot vagy -CR₂₇ csoportot jelöl;

G₁₃ nitrogénatomot vagy -CR₂₈ csoportot jelöl;

G₁₄ és G₁₅ mindegyike egymástól függetlenül nitrogénatomot vagy szénatomot jelöl, ahol ha a G₁₄ és G₁₅ valamelyike nitrogénatomot jelöl, akkor a másik fennmaradó elem szénatomot jelöl.

G₁₆ -NH csoportot vagy -CH csoportot jelöl;

G₁₇ nitrogénatomot vagy -CR₂₉ csoportot jelöl;

G₁₈ nitrogénatomot vagy -CR₃₀ csoportot jelöl;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉, és R₃₀ mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, -(CH₂)₀₋₃CN csoportot, -C(O)-(CH₂)₁₋₃CN csoportot, -C(O)-(CH₂)₀₋₃-R' csoportot, -C(O)-(CH₂)₀₋₃-NR'R'' csoportot, -(CH₂)₀₋₃NR'R'' csoportot jelöl, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituáltnak vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxycsoportok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' a fentiek szerint meghatározott; és ahol Y a fentiek szerint meghatározott;

vagy abban az esetben, ha Y egy -NR' csoportot jelöl, akkor R₄ az -NR'- csoporttal és a mind az R₄-hez, mind az -NR'- csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport szubsztituáltnak vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogén atomok, hidroxilcsoportok, cianocsoportok, -CHF₂ csoportok és -CF₃ csoportok közül lettek kiválasztva.

2. Vegyület az 1-es igénypont szerint, ahol

R₃ és R₆ mindegyike egymástól függetlenül hidrogénatomot, vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelölnek;

n 0-t, 1-et vagy 2-t jelöl;

R₁ hidrogénatomot, lineáris vagy elágazó C₁-C₄ alkilcsoportot, C₁-C₄ haloalkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, fenilcsoportot, 5-10 tagú monociklusos vagy biciklusos heteroanil-csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva, pirrolidinil-csoportot, piperidinil-csoportot, piperazinil-csoportot, tetrahidropiridinil-csoportot vagy morfolinil-csoportot;

ahol a gyűrűs alkil, fenil, heteroanil, pirrolidinil, piperidinil, piperazinil, tetrahidropiridinil vagy morfolinil-csoportok szubsztituáltnak vagy szubsztituált egy vagy több szubsztituenssel halogénatomok, hidroxilcsoportok, lineáris vagy elágazó C₁-C₄ alkilcsoportok, C₁-C₄ haloalkil-csoportok, C₁-C₄ hidroxialkil-csoportok, C₃-C₄ gyűrűs alkilcsoportok, -(CH₂)₀₋₃OR₈ csoportok, -(CH₂)₀₋₃NR₇R₈ csoportok, -C(O)-(CH₂)₀₋₃-R₈ csoportok, -C(O)-(CH₂)₀₋₃-NR₇R₈ csoportok vagy -(CH₂)₀₋₃(fenil)-OR₈ csoportok közül lettek kiválasztva; ahol R₇ és R₈ mindegyike egymástól függetlenül hidrogénatomot vagy C₁-C₄ alkilcsoportot jelöl,

R_2 jelölhet hidrogénatomot, halogén atomot, hidroxilcsoportot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-NH_2$ csoportot, $-N(CH_3)H$ csoportot, $-N(CH_3)_2$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot, mely alkilcsoport szubsztituálatlan vagy szubsztituált C_1 - C_2 alkoxycsoporttal.

R_3 jelölhet hidrogénatomot, halogén atomot, hidroxilcsoportot, cianocsoportot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-NH_2$ csoportot, $-N(CH_3)H$ csoportot, $-N(CH_3)_2$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot, mely alkilcsoport szubsztituálatlan vagy szubsztituált C_1 - C_2 alkoxycsoporttal.



R_4 hidrogénatomot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil-csoportot, C_1 - C_4 hidroxialkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot vagy egy lineáris vagy elágazó C_1 - C_3 alkilcsoportot jelöl.

R_5 hidrogénatomot; halogénatomot; hidroxilcsoportot; cianocsoportot; C_1 - C_3 alkoxycsoportot; a C_1 - C_3 haloalkil-csoportot; lineáris vagy elágazó C_1 - C_4 hidroxialkil-csoportot; C_3 - C_4 gyűrűs alkilcsoportot; $-(CH_2)_{0-3}NR'R''$ csoportot; $-(CH_2)_{1-3}O(C_1-C_4$ alkilcsoportot); $-(CH_2)_{0-3}OC(O)-(C_1-C_4$ alkilcsoportot); $-(CH_2)_{0-3}C(O)O-(C_1-C_4$ alkilcsoportot); $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportot; $-(CH_2)_{0-3}C(O)OH$ csoportot jelöl vagy; olyan $-(CH_2)_{0-3}$ -(5-14 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz legalább egy heteroatomot O, S és N közül kiválasztva) vagy; olyan $-(CH_2)_{0-3}$ -(5-14 tagú monociklusos vagy biciklusos heterociklikus-csoportot jelöl, amely tartalmaz legalább egy heteroatomot O, S és N közül kiválasztva);

vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált C_1 - C_2 alkoxycsoporttal;

ahol a heteroaril és heterociklikus csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, hidroxilcsoportok, cianocsoportok, lineáris vagy elágazó C_1 - C_4 alkilcsoportok vagy C_1 - C_4 haloalkil-csoportok közül lettek kiválasztva,

R_5 formula (II-a2), (IIb-1), (IIb-2), (IIb-3), (IIa-1), (IIa-3), (IIa-5) vagy (IIa-14) csoportját jelöli:



ahol:

R_9 , R_{10} és R_{12} mindegyike egymástól függetlenül hidrogénatomot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}R'$ csoportot, $-C(O)-(CH_2)_{0-3}NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelöl; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1-C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelöl;

R_{13} , R_{14} , R_{15} és R_{16} mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, C_1-C_4 haloalkil-csoportot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}NR'R''$ csoportot vagy,

$-(CH_2)_{0-3}NR'R''$ csoportot jelölnek; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1-C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelölnek.

R_{18} , R_{22} , R_{23} és R_{24} mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}R'$ csoportot, $-C(O)-(CH_2)_{0-3}NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelöl; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1-C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelöl;

Y egy $-NR'$ - csoportot, $-O$ - vagy $-S$ - kötést jelöl, ahol R' hidrogénatomot vagy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelöl, vagy abban az esetben, ha Y egy $-NR'$ - csoportot jelöl, akkor R_4 az $-NR'$ - csoporttal és a mind az R_4 -hez, mind az $-NR'$ - csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogén atomok, hidroxilcsoportok, cianocsoportok, $-CHF_2$ csoportok és $-CF_3$ csoportok közül lettek kiválasztva.

3. Vegyület az 1-es igénypont szerint, ahol

X nitrogénatomot vagy $-CR_5$ csoportot jelöl;

n 0-t, 1-et, 2-t vagy 3-at jelöl;

R_a és R_b mindegyike egymástól függetlenül hidrogénatomot, C_1-C_4 haloalkil-csoportot, C_1-C_4 hidroxialkil-csoportot vagy egy lineáris vagy elágazó C_1-C_4 alkilcsoportot jelölnek;

R_1 lineáris vagy elágazó C_1-C_4 alkilcsoportot, C_3-C_{10} gyűrűs alkilcsoportot, C_3-C_{10} cikloalkenil-csoportot, monociklusos vagy biciklusos C_5-C_{14} arilcsoportot, 5-14 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely legalább egy heteroatomot tartalmaz O , S és N közül kiválasztva, vagy egy 5-14 tagú monociklusos vagy biciklusos heterociklikus csoportot jelöl, mely legalább egy heteroatomot tartalmaz O , S és N közül kiválasztva,

ahol a gyűrűs alkil, cikloalkenil, aril, heteroaril és heterociklikus csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, hidroxilcsoportok, cianocsoportok, lineáris vagy elágazó C_1-C_4 alkilcsoportok, C_1-C_4 haloalkil-csoportok, C_1-C_4 hidroxialkil-csoportok, C_3-C_4 gyűrűs alkilcsoportok, $-(CH_2)_{1-3}CN$ csoportok, $-(CH_2)_{0-3}OR_a$ csoportok, a $-(CH_2)_{0-3}NR_7R_8$ csoportok, $-C(O)-(CH_2)_{1-3}CN$ csoportok, $-C(O)-(CH_2)_{0-3}R_a$ csoportok, $-C(O)-(CH_2)_{0-3}NR_7R_8$ csoportok, $-S(O)_2(CH_2)_{0-3}R_a$ csoportok vagy $-S(O)_2(CH_2)_{0-3}NR_7R_8$ csoportok közül lettek kiválasztva;

R_2 és R_3 mindegyike egymástól függetlenül hidrogén atomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C_1-C_4 alkoxycsoportot, C_1-C_4 haloalkil-csoportot, C_1-C_4 hidroxialkil-csoportot, C_3-C_7 gyűrűs alkilcsoportot, $-NR'R''$ csoportot, vagy egy lineáris vagy elágazó C_1-C_6 alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C_1-C_4 alkoxycsoportok, cianocsoportok vagy C_3-C_7 gyűrűs alkilcsoportok közül lettek kiválasztva;

R₄ hidrogénatomot, C₁-C₄ alkoxicsoprotot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, $-(CH_2)_{1-4}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoprotok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok, $-C(O)-(CH_2)_{0-3}-R'$ csoportok vagy $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportok közül lettek kiválasztva;

R₅ hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C₁-C₄ alkoxicsoprotot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoprotok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva;

R₇ és R₈ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoprotok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva;

R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C₁-C₄ alkoxicsoprotot vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelölnek.

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, és R₁₆ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ alkoxicsoprotot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}-R'$ csoportot, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot jelöl, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoprotok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' a fentiek szerint meghatározott;

R₁₈, R₁₉, R₂₀, R₂₁, R₂₂, R₂₃, R₂₄, R₂₅, R₂₆, R₂₇, R₂₈, R₂₉, és R₃₀ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ alkoxicsoprotot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}-R'$ csoportot, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot jelöl, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxicsoprotok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' a fentiek szerint meghatározott; és ahol Y a fentiek szerint meghatározott;

4. Vegyület az 3-as igénypont szerint, ahol

R_a és R_b mindegyike egymástól függetlenül hidrogénatomot, vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelölnek;

n 0-t, 1-et vagy 2-t jelöl;

R₁ lineáris vagy elágazó C₁-C₄ alkilcsoportot, C₃-C₇ gyűrűs alkilcsoportot, fenilcsoportot, 5-10 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva, pirrolidinil-csoportot, piperidinil-csoportot, piperazinil-csoportot, tetrahydro-piranilcsoportot vagy morfolinil-csoportot;

ahol a gyűrűs alkil, fenil, heteroaril, pirrolidinil, piperidinil, piperazinil, tetrahydro-piranil vagy morfolinil-csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több szubsztituenssel halogénatomok, lineáris vagy elágazó C₁-C₄

alkilcsoportok, C_1 - C_4 haloalkil-csoportok, C_1 - C_4 hidroxilalkil-csoportok, C_3 - C_4 gyűrűs alkilcsoportok, $-(CH_2)_{0-3}OR_8$ csoportok, $-(CH_2)_{0-3}NR_7R_8$ csoportok, $-C(O)-(CH_2)_{0-3}-R_8$ csoportok, $-C(O)-(CH_2)_{0-3}-NR_7R_8$ csoportok közül lettek kiválasztva; ahol R_7 és R_8 mindegyike egymástól függetlenül hidrogénatomot vagy C_1 - C_4 alkilcsoportot jelöl;


R_2 jelölhet hidrogénatomot, halogénatomot, hidroxilcsoportot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-NH_2$ csoportot, $-N(CH_3)H$ csoportot, $-N(CH_3)_2$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot, mely alkilcsoport szubsztituált vagy szubsztituált C_1 - C_2 alkoxycsoporttal.

R_3 jelölhet hidrogénatomot, halogénatomot, hidroxilcsoportot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-NH_2$ csoportot, $-N(CH_3)H$ csoportot, $-N(CH_3)_2$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot, mely alkilcsoport szubsztituált vagy szubsztituált C_1 - C_2 alkoxycsoporttal.

R_4 hidrogénatomot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot vagy egy lineáris vagy elágazó C_1 - C_3 alkilcsoportot jelöl;

R_5 jelölhet hidrogénatomot, halogén atomot, C_1 - C_3 alkoxycsoportot, C_1 - C_3 haloalkil csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-NH_2$ csoportot, $-N(CH_3)H$ csoportot, $-N(CH_3)_2$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot, mely alkilcsoport szubsztituált vagy szubsztituált C_1 - C_2 alkoxycsoporttal;

R_5 formula (IIa-2), (IIb-1), (IIb-2), vagy (IIa-1) csoportját jelöli:



ahol:

R_9 , R_{10} , és R_{12} mindegyike egymástól függetlenül hidrogénatomot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}-CN$ csoportot, $-C(O)-(CH_2)_{0-3}-R'$ csoportot, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1 - C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl;

R_{13} , R_{14} , R_{15} és R_{16} mindegyike egymástól függetlenül hidrogénatomot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot jelölnek; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelölnek;

R_{17} és R_{18} mindegyike egymástól függetlenül hidrogénatomot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}-CN$ csoportot, $-C(O)-(CH_2)_{0-3}-R'$ csoportot, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1 - C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl;

Y egy -NR'- csoportot, -O- vagy -S- kötést jelöl, ahol R' hidrogénatomot vagy lináris vagy elágazó C₁-C₄ alkilcsoportot jelöl, vagy abban az esetben, ha Y egy -NR'- csoportot jelöl, akkor R₄ az -NR'- csoporttal és a mind az R₄-hez, mind az -NR'- csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot.

5. Vegyület az 1-es igénypont szerint, ahol R₁ hidrogénatomot, lineáris vagy elágazó C₁-C₄ alkilcsoportot, C₁-C₄ haloalkil-csoportot, C₃-C₁₀ gyűrűs alkilcsoportot, C₂-C₁₀ cikloalkenil-csoportot, monociklusos vagy biciklusos C₆-C₁₄ arilcsoportot, 5-14 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely legalább egy heteroatomot tartalmaz O, S és N közül kiválasztva, vagy egy 5-14 tagú monociklusos vagy biciklusos heterociklikus csoportot jelöl, mely legalább egy heteroatomot tartalmaz O, S és N közül kiválasztva,

ahol a gyűrűs alkil, cikloalkenil, aril, heteroaril és heterociklikus csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogénatomok, hidroxilcsoportok, cianocsoportok, lináris vagy elágazó C₁-C₄ alkilcsoportok, C₁-C₄ haloalkil-csoportok, C₁-C₄ hidroxialkil-csoportok, C₃-C₄ gyűrűs alkilcsoportok, -(CH₂)₁₋₃CN csoportok, -(CH₂)₀₋₃OR₈ csoportok, a -(CH₂)₀₋₃NR₇R₈ csoportok, -C(O)-(CH₂)₁₋₃CN csoportok, -C(O)-(CH₂)₀₋₃R₈ csoportok, -C(O)-(CH₂)₀₋₃NR₇R₈ csoportok, -S(O)₂(CH₂)₀₋₃R₈ csoportok, -S(O)₂(CH₂)₀₋₃NR₇R₈ csoportok vagy -(CH₂)₀₋₃(fenil)-OR₈ csoportok közül lett kiválasztva;

és

ahol R₁ preferáltan hidrogénatomot, C₁-C₃ alkilcsoportot, C₁-C₃ haloalkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, fenilcsoportot, naftilcsoportot, 5-10 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva, vagy egy 5-10 tagú monociklusos vagy biciklusos heterociklikus csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva,

ahol a gyűrűs alkil, fenil, naftil, heteroaril és heterociklikus csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogénatomok, hidroxilcsoportok, cianocsoportok, lináris vagy elágazó C₁-C₄ alkilcsoportok, C₁-C₄ haloalkil-csoportok, C₁-C₄ hidroxialkil-csoportok, C₃-C₄ gyűrűs alkilcsoportok, -(CH₂)₀₋₃OR₈ csoportok, a -(CH₂)₀₋₃NR₇R₈ csoportok, -C(O)-(CH₂)₀₋₃R₈ csoportok, -C(O)-(CH₂)₀₋₃NR₇R₈ csoportok, vagy -(CH₂)₀₋₃(fenil)-OR₈ csoportok közül lett kiválasztva; ahol R₇ és R₈ az 1-es igénypont szerint meghatározott.

6. Vegyület az 1-es, 3-as vagy 5-ös igénypont bármelyike szerint, ahol:

R₁ egy C₁-C₃ alkilcsoportot, C₃-C₇ gyűrűs alkilcsoportot, fenilcsoportot, naftilcsoportot, 5-10 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva, vagy egy 5-10 tagú monociklusos vagy biciklusos heterociklikus csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva,

ahol a gyűrűs alkil, fenil, naftil, heteroaril és heterociklikus csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, hidroxilcsoportok, cianocsoportok, lináris vagy elágazó C₁-C₄ alkilcsoportok, C₁-C₄ haloalkil-csoportok, C₁-C₄ hidroxialkil-csoportok, C₃-C₄ gyűrűs alkilcsoportok, -(CH₂)₀₋₃OR₈ csoportok, a -(CH₂)₀₋₃NR₇R₈ csoportok, -C(O)-(CH₂)₀₋₃R₈ csoportok, -C(O)-(CH₂)₀₋₃NR₇R₈ csoportok közül lettek kiválasztva; ahol R₇ és R₈ az 1-es igénypont szerint meghatározott.

és

ahol R₁ preferáltan C₃-C₇ gyűrűs alkilcsoportot, fenilcsoportot, 5-10 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz egy, kettő vagy három heteroatomot O, S és N közül kiválasztva, pirrolidinil-csoportot, piperidinil-csoportot, piperazinil-csoportot, tetrahydro-piranilcsoportot vagy morfolinil-csoportot;

ahol a gyűrűs alkil, fenil, heteroaril, pirrolidinil, piperidinil, piperazinil, tetrahydro-piranil vagy morfolinil-csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több szubsztituenssel halogénatomok, lináris vagy elágazó C₁-C₄

alkilcsoportok, C₁-C₄ haloalkil-csoportok, C₁-C₄ hidroxialkil-csoportok, C₃-C₄ gyűrűs alkilcsoportok, -(CH₂)₀₋₃OR_a csoportok, -(CH₂)₀₋₃NR₇R_a csoportok, -C(O)-(CH₂)₀₋₃-R_a csoportok vagy -C(O)-(CH₂)₀₋₃-NR₇R_a csoportok közül lettek kiválasztva, ahol R₇ és R_a mindegyike egymástól függetlenül hidrogénatomot vagy C₁-C₄ alkilcsoportot jelöl;

és

ahol R₁ leginkább preferáltan fenilcsoportot vagy piridinilcsoportot jelöl, mely fenil vagy piridinil csoport szubsztituálatlan vagy szubsztituált egy, kettő vagy három szubsztituenssel halogénatomok, lineáris vagy elágazó C₁-C₃ alkilcsoportok vagy -(CH₂)₀₋₃OCH₃ csoportok közül kiválasztva; és ahol preferáltan a nevezett fenilcsoportok és piridinilcsoportok közvetlenül kapcsolódnak a pirrolo-triazinon csoporthoz; és/vagy

R₂ hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, -(CH₂)₀₋₃NR'R'' csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált C₁-C₃ alkoxycsoporttal; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot vagy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl;

és

ahol R₂ preferáltan hidrogénatomot, halogénatomot, hidroxilcsoportot, C₁-C₃ alkoxycsoportot, C₁-C₃ haloalkil csoportot, C₃-C₄ gyűrűs alkilcsoportot, -NH₂ csoportot, -N(CH₃)H csoportot, -N(CH₃)₂ csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált C₁-C₂ alkoxycsoporttal; és/vagy

R₃ hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, -(CH₂)₀₋₃NR'R'' csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált C₁-C₃ alkoxycsoporttal; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot vagy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl;

és

ahol R₃ preferáltan hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C₁-C₃ alkoxycsoportot, C₁-C₃ haloalkil csoportot, C₃-C₄ gyűrűs alkilcsoportot, -NH₂ csoportot, -N(CH₃)H csoportot, -N(CH₃)₂ csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált C₁-C₂ alkoxycsoporttal; és/vagy

R₄ hidrogénatomot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, -(CH₂)₁₋₄NR'R'' csoportot, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált C₁-C₃ alkoxycsoporttal, -C(O)-(CH₂)₀₋₃-R' csoporttal vagy -C(O)-(CH₂)₀₋₃-NR'R'' csoporttal; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot vagy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl;

és

ahol R₄ preferáltan hidrogénatomot, C₁-C₃ haloalkil-csoportot, C₁-C₃ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl;

R₅ hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, lineáris vagy elágazó C₁-C₄ hidroxialkil-csoportot, C₃-C₇ gyűrűs alkilcsoportot, -(CH₂)₀₋₃NR'R'' csoportot, -(CH₂)₁₋₃O(C₁-C₄ alkilcsoportot); -(CH₂)₀₋₃OC(O)-(C₁-C₄ alkilcsoportot); -(CH₂)₀₋₃C(O)O-(C₁-C₄ alkilcsoportot);

-C(O)-(CH₂)₀₋₃-NR'R'' csoportot, -(CH₂)₀₋₃C(O)OH csoportot jelöl vagy: olyan -(CH₂)₀₋₃-(5-10 tagú monociklusos vagy biciklusos heteroaril-csoportot jelöl, mely tartalmaz legalább egy heteroatomot O, S és N közül kiválasztva) vagy; olyan -(CH₂)₀₋₃-(5-10 tagú monociklusos vagy biciklusos heterociklikus-csoportot jelöl, amely tartalmaz legalább egy heteroatomot O, S és N közül kiválasztva); vagy egy olyan lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált

egy vagy több szubsztituenssel C_1 - C_4 alkoxicsoporthoz, cianocsoportot vagy C_3 - C_4 gyűrűs alkilcsoportok közül kiválasztva; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot vagy lineáris vagy elágazó C_1 - C_3 alkilcsoportot jelöl; és

ahol a heteroaril és heterociklikus csoportok szubsztituáltak vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, hidroxilcsoportok, cianocsoportok, lineáris vagy elágazó C_1 - C_3 alkilcsoportok vagy C_1 - C_3 haloalkil-csoportok közül lettek kiválasztva, és/vagy

R_8 hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C_1 - C_4 alkoxicsoporthoz, C_1 - C_4 haloalkil-csoportot, C_1 - C_4 hidroalkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituáltak vagy szubsztituáltak C_1 - C_3 alkoxicsoporthoz; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot vagy lineáris vagy elágazó C_1 - C_3 alkilcsoportot jelöl;


és

ahol R_8 preferáltan hidrogénatomot, halogén atomot, C_1 - C_3 alkoxicsoporthoz, C_1 - C_3 haloalkil csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-NH_2$ csoportot, $-N(CH_3)H$ csoportot, $-N(CH_3)_2$ csoportot, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituáltak vagy szubsztituáltak C_1 - C_3 alkoxicsoporthoz.


7. Vegyület az 1-es, 3-as 5-ös vagy 6-os igénypont bármelyike szerint, ahol:

R_6 az alábbiak közül választott csoportot jelöl

i) formula (IIa-1) csoport, ii) formula (IIa-2) csoport



formula (IIa-1)




formula (IIa-2)

ahol


R_9 , R_{10} , R_{11} , és R_{12} mindegyike egymástól függetlenül hidrogénatomot, C_1 - C_4 alkoxicsoporthoz, C_1 - C_4 haloalkil-csoportot, C_1 - C_4 hidroalkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}R'$ csoportot, $-C(O)-(CH_2)_{0-3}NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot jelöl, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituáltak vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek C_1 - C_4 alkoxicsoporthoz, cianocsoportok vagy C_3 - C_4 gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1 - C_4 alkoxicsoporthoz vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl; vagy

R_6 az alábbiak közül választott csoportot jelöl


i) formula (IIb-1) csoport, ii) formula (IIb-2) csoport, iii) formula (IIb-3) csoport



formula (IIb-1)




formula (IIb-2)




formula (IIb-3)

iv) formula (IIb-4) csoport, és

v) formula (IIb-5) csoport



formula (IIb-4)



formula (IIb-5)

ahol

R_{13} , R_{14} , R_{15} , és R_{16} mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, C_1 - C_4 alkoxycsoportot, C_1 - C_4 haloalkil-csoportot, C_1 - C_4 hidroalkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-(CH_2)_{3-5}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{3-5}R'$ csoportot, $-C(O)-(CH_2)_{3-5}NR'R''$ csoportot, $-(CH_2)_{3-5}NR'R''$ csoportot jelöl, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C_1 - C_4 alkoxycsoportok, cianocsoportok vagy C_3 - C_4 gyűrűs alkilcsoportok közül lettek kiválasztva;

ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxycsoportot, C_1 - C_3 alkoxycsoportot vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl; és

ahol Y az 1-es igényponi szerint meghatározott;


vagy abban az esetben, ha Y egy $-NR'$ - csoportot jelöl, akkor R_4 az $-NR'$ - csoporttal és a mind az R_4 -hez, mind az $-NR'$ -csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogénatomok, hidroxycsoportok, cianocsoportok, $-CHF_2$ csoportok és $-CF_3$ csoportok közül lettek kiválasztva; vagy

R_5 az alábbiak közül választott csoportot jelöl


i) formula (IIa-1) csoport,

ii) formula (IIa-2) csoport


iii) formula (IIa-3) csoport



formula (IIa-1)



formula (IIa-2)




formula (IIa-3)


iv) formula (IIa-4) csoport,

v) formula (IIa-5) csoport


vi) formula (IIa-6) csoport



formula (IIa-4)




formula (IIa-5)




formula (IIa-6)

vii) formula (IIa-7) csoport,




formula (IIa-7)

viii) formula (IIa-8) csoport




formula (IIa-8)

ix) formula (IIa-9) csoport




formula (IIa-9)

x) formula (IIa-10) csoport,




xiii) formula (IIa-10)

xi) formula (IIa-11) csoport




formula (IIa-11)

xii) formula (IIa-12) csoport




formula (IIa-12)

formula (IIa-13) csoport, xiv) formula (IIa-14) csoport




formula (IIa-13)




formula (IIa-14)

xv) formula (IIa-15) csoport




formula (IIa-15)

xvi) formula (IIa-16) csoport,




formula (IIa-16)

xvii) formula (IIa-17) csoport




formula (IIa-17)

xviii) formula (IIa-18) csoport




formula (IIa-18)

xix) formula (IIa-19) csoport,




formula (IIa-19)

xx) formula (IIa-20) csoport




formula (IIa-20)

xxi) formula (IIa-21) csoport



formula (IIa-21)

xxii) formula (IIa-19) csoport



formula (IIIa-22)

ahol


R_{18} , R_{22} , R_{23} , és R_{24} mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, C_1 - C_4 alkoxycsoportot, C_1 - C_4 haloalkil-csoportot, C_1 - C_4 hidroxiálkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-(CH_2)_{0-3}CN$ csoportot, $-C(O)-(CH_2)_{1-3}CN$ csoportot, $-C(O)-(CH_2)_{0-3}R'$ csoportot, $-C(O)-(CH_2)_{0-3}NR'R''$ csoportot, $-(CH_2)_{0-3}NR'R''$ csoportot jelöl, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C_1 - C_4 alkoxycsoportok, cianocsoportok vagy C_3 - C_4 gyűrűs alkilcsoportok közül lettek kiválasztva;

ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1 - C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl; és ahol Y az 1-es igénypont szerint meghatározott;

vagy abban az esetben, ha Y egy $-NR'$ - csoportot jelöl, akkor R_4 az $-NR'$ - csoporttal és a mind az R_4 -hez, mind az $-NR'$ -csoportoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogénatomok, hidroxilcsoportok, cianocsoportok, $-CHF_2$ csoportok és $-CF_3$ csoportok közül lettek kiválasztva; vagy


R_5 az alábbiak közül választott csoportot jelöl

i) formula (IIIb-1) csoport,




formula (IIIb-1)

ii) formula (IIIb-2) csoport




formula (IIIb-2)

iii) formula (IIIb-3) csoport




formula (IIIb-3)

iv) formula (IIIb-4) csoport,




formula (IIIb-4)

v) formula (IIIb-5) csoport




formula (IIIb-5)

vi) formula (IIIb-6) csoport




formula (IIIb-6)

vii) formula (IIIb-7) csoport,



viii) formula (IIIb-8) csoport




ahol

R₁₆, R₂₂, R₂₅, R₂₆, és R₂₇ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, -(CH₂)₀₋₃CN csoportot, -C(O)-(CH₂)₁₋₃-CN csoportot, -C(O)-(CH₂)₀₋₃-R' csoportot, -C(O)-(CH₂)₀₋₃-NR'R'' csoportot, -(CH₂)₀₋₃-NR'R'' csoportot jelöl, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxycsoportok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C₁-C₄ alkoxycsoportot vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl; és ahol Y az 1-es igénypont szerint meghatározott;


vagy abban az esetben, ha Y egy -NR'- csoportot jelöl, akkor R₄ az -NR'- csoporttal és a mind az R₄-hez, mind az -NR'- csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogénatomok, hidroxilcsoportok, cianocsoportok, -CHF₂ csoportok és -CF₃ csoportok közül lettek kiválasztva; vagy

R₅ az alábbiak közül választott csoportot jelöl

i) formula (IIIc-1) csoport,



ii) formula (IIIc-2) csoport



ahol


R₁₉, R₂₀, és R₂₈ mindegyike egymástól függetlenül hidrogénatomot, C₁-C₄ alkoxycsoportot, C₁-C₄ haloalkil-csoportot, C₁-C₄ hidroxialkil-csoportot, C₃-C₄ gyűrűs alkilcsoportot, -(CH₂)₀₋₃CN csoportot, -C(O)-(CH₂)₁₋₃-CN csoportot, -C(O)-(CH₂)₀₋₃-R' csoportot, -C(O)-(CH₂)₀₋₃-NR'R'' csoportot, -(CH₂)₀₋₃-NR'R'' csoportot jelöl, vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C₁-C₄ alkoxycsoportok, cianocsoportok vagy C₃-C₄ gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C₁-C₄ alkoxycsoportot vagy egy lineáris vagy elágazó C₁-C₄ alkilcsoportot jelöl; és ahol Y az 1-es igénypont szerint meghatározott;

vagy abban az esetben, ha Y egy -NR'- csoportot jelöl, akkor R₄ az -NR'- csoporttal és a mind az R₄-hez, mind az -NR'- csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport

szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogénatomok, hidroxilcsoportok, cianocsoportok, $-\text{CHF}_2$ csoportok és $-\text{CF}_3$ csoportok közül lettek kiválasztva; vagy


R_3 az alábbiak közül választott csoportot jelöl

i) formula (III-d-1) csoport,




formula (III-d-1)

ii) formula (III-d-2) csoport



formula (III-d-2)

iii) formula (III-d-3) csoport



formula (III-d-3)


ahol

R_{21} , R_{22} , R_{23} és R_{24} mindegyike egymástól függetlenül hidrogénatomot, C_1 - C_4 alkoxycsoportot, C_1 - C_4 haloalkil-csoportot, C_1 - C_4 hidroxialkil-csoportot, C_3 - C_4 gyűrűs alkilcsoportot, $-(\text{CH}_2)_{0-3}\text{CN}$ csoportot, $-\text{C}(\text{O})-(\text{CH}_2)_{1-3}\text{CN}$ csoportot, $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{R}'$ csoportot, $-\text{C}(\text{O})-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ csoportot, $-(\text{CH}_2)_{0-3}\text{NR}'\text{R}''$ csoportot jelöl, vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl, mely alkilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, melyek C_1 - C_4 alkoxycsoportok, cianocsoportok vagy C_3 - C_4 gyűrűs alkilcsoportok közül lettek kiválasztva; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot, C_1 - C_4 alkoxycsoportot vagy egy lineáris vagy elágazó C_1 - C_4 alkilcsoportot jelöl; és ahol Y az 1-es igénypont szerint meghatározott;

vagy abban az esetben, ha Y egy $-\text{NR}'$ - csoportot jelöl, akkor R_4 az $-\text{NR}'$ - csoporttal és a mind az R_4 -hez, mind az $-\text{NR}'$ -csoporthoz kötődő szénatommal egy 4-7 tagú telített, nitrogéntartalmú heterociklikus csoportot alkot, mely heterociklikus csoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogénatomok, hidroxilcsoportok, cianocsoportok, $-\text{CHF}_2$ csoportok és $-\text{CF}_3$ csoportok közül lettek kiválasztva.

8. Vegyület az 1-es igénypont szerint, amely:


(a) a formula (Ia) szerint épül fel:



Formula (Ia)

ahol R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 és n az eddigi igénypontok bármelyike szerint meghatározott; vagy

(b) a formula (Ib) szerint épül



fel:

ahol $R_1, R_2, R_3, R_4, R_5, R_6$ és n az 1-től 7-ig igénypontok bármelyike szerint meghatározott.

9. Vegyük az 1-es igénypont szerint, ahol:

X nitrogénatomot vagy $-CR_5$ csoportot jelöl;

R_5 és R_6 mindegyike egymástól függetlenül hidrogénatomot, metilcsoportot jelölnek;

R_1 hidrogénatomot, halogénatomot, C_1-C_3 haloalkil-csoportot, metilcsoportot, C_3-C_7 gyűrűs alkilcsoportot, fenilcsoportot, piridinilcsoportot, pirazolilcsoportot, izoxazolilcsoportot, piperidinil-csoportot vagy tetrahidro-piránil-csoportot jelöl;

ahol a gyűrűs alkil, fenil, piridinil, pirazolil, izoxazolil, piperidinil vagy tetrahidro-piránil-csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több szubsztituenssel halogénatomok, hidroxilcsoportok, C_1-C_3 haloalkil-csoportok, lineáris vagy elágazó C_1-C_3 alkilcsoportok, $-(CH_2)-(fenil)-O-(C_1-C_3)$ alkilcsoportok), $-NR_7R_8$ csoportok vagy $-OR_8$ csoportok közül kiválasztva; ahol R_7 és R_8 mindegyike egymástól függetlenül hidrogénatomot vagy lineáris vagy elágazó C_1-C_3 alkilcsoportot jelöl;

R_2 és R_3 mindegyike egymástól függetlenül hidrogénatomot, halogénatomot, cianocsoportot, C_1-C_3 haloalkil-csoportot, vagy egy lineáris vagy elágazó C_1-C_3 alkilcsoportot jelöl;

R_4 hidrogénatomot, C_1-C_3 haloalkil-csoportot, C_1-C_3 hidroxialkil-csoportot vagy egy lineáris vagy elágazó C_1-C_3 alkilcsoportot jelöl;

R_5 hidrogénatomot, halogénatomot, hidroxilcsoportot, cianocsoportot, C_1-C_4 alkoxicsoportot, C_1-C_4 haloalkil-csoportot, lineáris vagy elágazó C_1-C_4 hidroxialkil-csoportot, C_3-C_7 gyűrűs alkilcsoportot, lineáris vagy elágazó C_1-C_3 alkilcsoportot, $-(CH_2)_{0-3}NR'R''$ csoportot, $-(CH_2)_{1-3}O(C_1-C_3)$ alkilcsoportot, $-(CH_2)_{0-3}OC(O)-(C_1-C_3)$ alkilcsoportot, $-(CH_2)_{0-3}C(O)O-(C_1-C_3)$ alkilcsoportot, $-C(O)-NR'R''$ csoportot, $-(CH_2)_{0-3}C(O)OH$ csoportot, $-(CH_2)_{0-3}$ (imidazolil csoportot), $-(CH_2)_{0-3}$ (oxazolil csoportot), $-(CH_2)_{0-3}$ (oxadiazolil csoportot), $-(CH_2)_{0-3}$ (pirazolil csoportot) vagy $-(CH_2)_{0-3}$ (morfolinil csoportot) jelöl; ahol R' és R'' mindegyike egymástól függetlenül hidrogénatomot, hidroxilcsoportot vagy lineáris vagy elágazó C_1-C_3 alkilcsoportot jelöl; és

ahol az imidazolil, oxazolil, oxadiazolil, pirazolil és morfolinil csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több olyan szubsztituenssel, melyek halogén atomok, lineáris vagy elágazó C_1-C_3 alkilcsoportok vagy C_1-C_3 haloalkil-csoportok közül lettek kiválasztva.

R_6 az alábbiak közül választott csoportot jelöl:

formula (IIa) szerinti csoport, mely csoport egy purinilcsoport, ami szubsztituálatlan vagy szubsztituált egy $-NR'R''$ csoporttal;

formula (IIb) szerinti csoport, mely csoport $-NR'$ -piridinil-csoport, $-S$ -piridinil-csoport, $-NR'$ -pirimidinil-csoport, $-S$ -pirimidinil-csoport vagy $-NR'$ -triazinil-csoport közül lett kiválasztva; ahol a piridinil, pirimidinil és triazinilcsoportok szubsztituálatlanok vagy szubsztituáltak egy, kettő vagy három olyan szubsztituenssel, melyek halogénatomok, C_1-C_3 haloalkil-csoportok, $-(CH_2)_{0-3}CN$ csoportok, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportok, $-(CH_2)_{0-3}NR'R''$ csoportok közül lettek kiválasztva; és

formula (Iic) szerinti csoport, mely csoport -NR'-purinilcsoport, -S-purinilcsoport, -NR'-7H-pirroló[2,3-d]pirimidinil-csoport, -NR'-1H-pirazolo[3,4-d]pirimidinil-csoport vagy -NR'-pirazolo[1,5-a]pirimidinil-csoport közül lett kiválasztva;

ahol a purinil, 7H-pirroló[2,3-d]pirimidinil, 1H-pirazolo[3,4-d]pirimidinil, pirazolo[1,5-a]pirimidinil és a többi csoportok szubsztituálatlanok vagy szubsztituáltak halogénatommal vagy $-(CH_2)_{0-3}NR'R''$ csoporttal; vagy

R₄ és R₅ a szénatommal együtt, melyhez kapcsolódva pirrolidinil-purinilcsoportot vagy pirrolidinil-pirimidinil csoportot alkotnak, ahol a pirrolidinilcsoport szubsztituálatlan vagy szubsztituált egy vagy több olyan szubsztituenssel, mely halogénatomok, hidroxilcsoportok közül lett kiválasztva, és ahol a purinilcsoport szubsztituálatlan vagy szubsztituált $-(CH_2)_{0-3}NR'R''$ csoporttal; és ahol a pirimidinil csoport szubsztituálatlan vagy szubsztituált egy, kettő vagy három olyan szubsztituenssel, mely $-(CH_2)_{0-3}CN$ csoportok vagy $-(CH_2)_{0-3}NR'R''$ csoportok közül lett kiválasztva; és

R' és R'' mindegyike egymástól függetlenül hidrogénatomot, C₁-C₃ alkoxycsoportot vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelölnek.

10. Vegyület az 1-es vagy 3-as igénypont szerint, ahol:

X nitrogénatomot vagy -CR₆ csoportot jelöl;

R₆ és R₇ mindegyike egymástól függetlenül hidrogénatomot vagy metilcsoportot jelölnek;

R₁ metilcsoportot, C₃-C₇ gyűrűs alkilcsoportot, fenilcsoportot, piridinilcsoportot, piperidinil-csoportot vagy tetrahidro-piránil-csoportot jelöl;

ahol a gyűrűs alkil, fenil, piridinil, piperidinil vagy tetrahidro-piránil-csoportok szubsztituálatlanok vagy szubsztituáltak egy vagy több szubsztituenssel halogénatomok, lineáris vagy elágazó C₁-C₃ alkilcsoportok, -NR₇R₈ csoportok vagy -OR₈ csoportok közül kiválasztva; ahol R₇ és R₈ mindegyike egymástól függetlenül hidrogénatomot vagy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl;

R₂ és R₃ mindegyike egymástól függetlenül hidrogénatomot, vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelölnek;

R₄ hidrogénatomot, C₁-C₃ haloalkil-csoportot vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelöl;

R₅ hidrogénatomot, halogénatomot, C₁-C₃ haloalkil-csoportot vagy egy lineáris vagy elágazó C₁-C₃ alkilcsoportot vagy ciklopropil-csoportot jelöl;

R₉ az alábbiak közül választott csoportot jelöl:

formula (IIa) szerinti csoport, mely csoport egy purinilcsoport, ami szubsztituálatlan vagy szubsztituált egy -NR'R'' csoporttal;

formula (IIb) szerinti csoport, mely csoport -NH-piridinil-csoport, -S-piridinil-csoport, -NH-pirimidinil-csoport, vagy -S-pirimidinil-csoport közül lett kiválasztva; ahol a piridinil, pirimidinil-csoportok szubsztituálatlanok vagy szubsztituáltak egy, kettő vagy három olyan szubsztituenssel, melyek $-(CH_2)_{0-3}CN$ csoportok, $-C(O)-(CH_2)_{0-3}-NR'R''$ csoportok vagy $-(CH_2)_{0-3}NR'R''$ csoportok közül lettek kiválasztva; és

formula (IIc) szerinti csoport, mely csoport -NH-purinilcsoport, -S-purinilcsoport közül lett kiválasztva; ahol a purinilcsoport szubsztituálatlan vagy szubsztituált $-(CH_2)_{0-3}NR'R''$ csoporttal; vagy

R₄ és R₅ a szénatommal együtt, melyhez kapcsolódva pirrolidinil-purinilcsoportot alkotnak, ahol a purinilcsoport szubsztituálattal vagy szubsztituált $-(CH_2)_nNR'R''$ csoporttal;

R' és R'' mindegyike egymástól függetlenül hidrogénatomot, C₁-C₃ alkoxycsoportot vagy lineáris vagy elágazó C₁-C₃ alkilcsoportot jelölnek.

11. Vegyület az 1-es igénypont szerint, mely egyike a következőknek:

- 2-((6-Amino-9H-purin-9-il)metil)-5-klór-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Aminopirimidin-4-ilamino)metil)-5-klór-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-5-ciklopropil-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-amino-9H-purin-9-il)metil)-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-aminopirimidin-4-ilamino)metil)-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 4-((4-Oxo-3-o-tolil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)metilamino)picolinamid;
- 2-((2-aminopiridin-4-ilamino)metil)-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((9H-purin-6-ilamino)metil)-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-3-ciklohexilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-amino-9H-purin-9-il)metil)-5-metil-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((9H-purin-6-ilitio)metil)-5-metil-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-amino-9H-purin-9-il)metil)-5-metil-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((9H-purin-6-ilitio)metil)-6-metil-3-o-tolilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-(1-(6-amino-9H-purin-9-il)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-2-(1-(9H-purin-6-ilamino)propil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-2-(1-(6-aminopirimidin-4-ilamino)propil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-2-(1-(2-amino-9H-purin-6-ilamino)propil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-4-amino-6-(1-(4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)propilamino)pirimidin-5-karbonitril;
- (R)-2-(1-(9H-purin-6-ilamino)propil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-2-(1-(9H-purin-6-ilamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-2-(1-(2-amino-9H-purin-6-ilamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-2-(1-(6-aminopirimidin-4-ilamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- (S)-4-amino-6-(1-(4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
- 2-(1-(6-amino-9H-purin-9-il)etil)-5-metil-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-3-o-tolil-5-(trifluorometil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-5-klór-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-5-klór-3-(3-metoxifenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-5-klór-3-(2,4-difluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-3-benzil-5-klórpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-amino-9H-purin-9-il)metil)-3-fenilimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-amino-9H-purin-9-il)metil)-3-o-tolilimidazo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-5-klór-3-(piridin-4-il)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
- 2-((6-Amino-9H-purin-9-il)metil)-5-klór-3-(tetrahidro-2H-pirán-4-il)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

2-((6-Amino-9H-purin-9-yl)methyl)-5-klór-3-(1-metilpiperidin-4-yl)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(9H-Purin-6-ylamino)etil)-3-(3-fluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-Amino-6-(1-(3-(3-fluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-Purin-6-ylamino)etil)-3-(3,5-difluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-Amino-6-(1-(3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

2-((6-Amino-9H-purin-9-yl)methyl)-5-klór-3-metilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

2-((6-Amino-9H-purin-9-yl)methyl)-3-((1R,4R)-4-aminociklohexil)-5-klórpírolo[1,2-f][1,2,4]triazin-4(3H)-on;

(R)-2-((6-Amino-9H-purin-9-yl)methyl)-5-klór-3-(1-feniletíl)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-((6-Amino-9H-purin-9-yl)methyl)-5-klór-3-(1-feniletíl)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(4-oxo-3-(piridin-2-yl)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-purin-6-yl)pirrolidin-2-yl)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(4-oxo-3-fenil-5-(trifluormetil)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-purin-6-ylamino)etil)-3-fenil-5-(trifluormetil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(5-(difluormetil)-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-purin-6-ylamino)etil)-5-(difluormetil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(9H-purin-6-ylamino)etil)-3-fenilimidazo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(4-oxo-3-fenil-3,4-dihidroimidazo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

2-(1-(9H-purin-6-ylamino)-3,3,3-trifluorpropil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

4-amino-6-(3,3,3-trifluor-1-(4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)propilamino)pirimidin-5-karbonitril;

(S)-4-amino-6-(2-(4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)pirrolidin-1-yl)pirimidin-5-karbonitril;

(S)-3-fenil-2-(1-(pirazolo[1,5-a]pirimidin-7-ylamino)etil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

2-((6-amino-9H-purin-9-yl)methyl)-5-(difluormetil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(2-amino-9H-purin-6-yl)pirrolidin-2-yl)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(4,6-diamino-1,3,5-triazin-2-ylamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-((6-amino-9H-purin-9-yl)methyl)-5-klór-3-(1-(5-fluorpiridin-2-yl)etil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(2-amino-9H-purin-6-ylamino)etil)-3-(3,5-difluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(6-amino-5-cianopirimidin-4-ylamino)etil)-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;

(S)-2-(1-(9H-purin-6-ylamino)etil)-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;

(R)-2-(1-(9H-purin-6-ylamino)-2-hidroxi)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(R)-4-amino-6-(2-hidroxi-1-(4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(2-amino-9H-purin-6-ylamino)etil)-3-fenilimidazo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(7H-pirrolo[2,3-d]pirimidin-4-ylamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(metil(1-(4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etil)amino)pirimidin-5-karbonitril;

(S)-2-(1-(metil(9H-purin-6-yl)amino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(9H-purin-6-ylamino)etil)-5-metil-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(5-metil-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-purin-6-ylamino)etil)-7-metil-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(7-metil-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

(S)-2-(4,4-difluor-1-(9H-purin-6-yl)pirrolidin-2-yl)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(4,4-difluor-2-(4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)pirrolidin-1-yl)pirimidin-5-karbonitril;

(S)-2-(1-(9H-purin-6-ylamino)etil)-6-fluor-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(6-fluor-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-yl)etilamino)pirimidin-5-karbonitril;

2-((S)-1-(9H-purin-6-ilamino)etil)-3-((S)-1-feniletil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 4-amino-6-((S)-1-(4-oxo-3-((S)-1-feniletil)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-amino-6-(1-(3-(2,6-dimetilfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-((9H-purin-6-ilamino)etil)-3-(1-feniletil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-amino-6-((4-oxo-3-(1-feniletil)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(5-fluor-7H-pirrolo[2,3-d]pirimidin-4-ilamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-purin-6-ilamino)etil)-3-(2,6-dimetilfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-amino-6-(1-(5-fluor-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(9H-purin-6-ilamino)etil)-5-fluor-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-amino-5-cianopirimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-4-amino-6-(1-(3-(3,5-difluorfenil)-4-oxo-3,4-dihidroimidazo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(9H-purin-6-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 4-amino-6-((1S)-1-(5-(1,2-dihidroxi)etil)-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-amino-6-(1-(3-(3,5-difluorfenil)-4-oxo-5-(trifluorometil)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(9H-Purin-6-ilamino)etil)-3-(3,5-difluorfenil)-5-(trifluorometil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(5-(hidroximetil)-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-(trifluorometil)pirimidin-4-ilamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-(piridin-2-ilmetil)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-2-(1-(9H-Purin-6-ilamino)etil)-5-(difluorometil)-3-(3,5-difluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(9H-Purin-6-ilamino)etil)-3-(3,5-difluorfenil)imidazo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(5-(difluorometil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(2-Amino-9H-purin-6-ilamino)etil)-5-(difluorometil)-3-(3,5-difluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(2-Amino-9H-purin-6-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 2-(1-(9H-Purin-6-ilamino)-2,2,2-trifluoretil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(3-benzil-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-fluorpirimidin-4-ilamino)etil)-3-fenilpirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-fluorpirimidin-4-ilamino)etil)-5-(difluorometil)-3-(3,5-difluorfenil)pirrolo[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)propil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(3,5-diklorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-fluorpirimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-(trifluorometil)pirimidin-4-ilamino)etil)-4-oxo-3-fenil-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (R)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)-2-hidroxi)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-carbamoylpirimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karboxamid;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karboxamid;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(2-klorbenzil)-4-oxo-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 2-((S)-1-(6-amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-((S)-tetrahydro-2H-pirán-3-il)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-5-karbonitril;
 (R)-4-Amino-6-(1-(3-(3,5-difluorfenil)-4-oxo-5-(trifluorometil)-3,4-dihidropirrolo[1,2-f][1,2,4]triazin-2-il)-2-hidroxi)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(2-Amino-5-fluorimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-2-(1-(2-Amino-5-cianopirimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 ((S)-2-(1-(9H-Purin-6-ilamino)etil)-3-(3,5-difluorfenil)-5-(2H-tetrazol-5-il)pirrol[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-4-Amino-6-(1-(3-((5-metilizoxazol-3-il)metil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-fenil-7-(trifluometil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-7-karbonitril;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(1-(4-metoxibenzil)-1H-pirazol-4-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-4-amino-6-(1-(4-oxo-3-fenil-5-(tiazol-2-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(2,6-Diamino-5-cianopirimidin-4-ilamino)etil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-4-Amino-6-(1-(5-(morpholinometil)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 2-((S)-1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-((R)-1-feniletil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pirazol-4-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(9H-Purin-6-ilamino)etil)-3-(3,5-difluorfenil)-5-(5-metil-1,2,4-oxadiazol-3-il)pirrol[1,2-f][1,2,4]triazin-4(3H)-on;
 4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pirán-3-il)-5-(trifluometil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(3-(5-metil-1H-pirazol-3-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonsav;
 2-((S)-1-(6-amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-((R)-tetrahydro-2H-pirán-3-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 (S)-4-Amino-6-(1-(3-(5-fluorimidin-3-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(1H-pirazol-3-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(pirimidin-5-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pirán-3-il)-5-(trifluometil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2,4-Diamino-6-(1-(4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-(1-(3-((1H-Pirazol-3-il)metil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)-6-aminopirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(tetrahydro-2H-pirán-4-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-(2,2,2-trifluoretil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(3-ciklobutil-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-Amino-4-(1-(4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 4-Amino-6-(1-(5-(1-metil-1H-pirazol-4-il)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(3-ciklopropil-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(5-bróm-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 4-amino-6-((S)-1-(4-oxo-3-((R)-tetrahydro-2H-pirán-3-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(5-bróm-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 2-((3-Iodo-1H-pirazol[3,4-d]pirimidin-4-ilamino)metil)-5-metil-3-o-tolilpirrol[1,2-f][1,2,4]triazin-4(3H)-on;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(5-fluorimidin-3-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;
 4-amino-6-((S)-1-(4-oxo-3-((S)-tetrahydro-2H-pirán-3-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(4-oxo-3-fenil-5-(1H-pirazol-4-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-4-Amino-6-(1-(3-(izoxazol-3-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;
 (S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-N,N-dimetil-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karboxamid;

(S)-4-Amino-6-(1-(3-(1-metil-1H-pirazol-3-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril

(S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-fenil-N-propil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karboxamid;

2-((S)-1-(9H-Purin-6-ilamino)etil)-3-(tetrahidro-2H-pirán-3-il)pirolo[1,2-f][1,2,4]triazin-4(3H)-on;

2-((S)-1-(9H-purin-6-ilamino)etil)-3-((S)-tetrahidro-2H-pirán-3-il)pirolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-Amino-6-(3-hidroxi-1-(4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)propilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-Purin-6-ilamino)-3-hidroxiopropil)-3-fenilpirolo[1,2-f][1,2,4]triazin-4(3H)-on;

(R)-4-Amino-6-(1-(3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)-2-hidroxiethylamino)pirimidin-5-karbonitril;

4-Amino-6-((4-oxo-3-o-tolil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)metilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(5-(2-hidroxietyl)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(6-amino-5-cianopirimidin-4-ilamino)-3-hidroxiopropil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;

(S)-2-(1-(9H-Purin-6-ilamino)etil)-3-(5-fluorpiridin-3-il)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;

(S)-4-Amino-6-(1-(5-(2-metiloxazol-5-il)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(5-(2-metoxietyl)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-Propil-2-(1-(6-amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karboxilat;

(S)-4-Amino-6-(3-hidroxi-1-(4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)propilamino)pirimidin-5-karbonitril;

(S)-2-(1-(9H-Purin-6-ilamino)-3-hidroxiopropil)pirolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(3-(3,5-difluorfenil)-4-oxo-5-(trifluormetil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)-3-hidroxiopropilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(4-oxo-3-(6-(trifluormetil)piridin-2-il)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(5-bróm-4-oxo-3-(3-(trifluormetil)fenil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-2-(2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-il)etil-acetát;

(S)-2-(1-(9H-Purin-6-ilamino)etil)-3-(6-(trifluormetil)piridin-2-il)pirolo[1,2-f][1,2,4]triazin-4(3H)-on;

2-((2S,4R)-1-(6-Amino-5-cianopirimidin-4-il)-4-hidroxiopirrolidin-2-il)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;

4-Amino-6-((2S,4R)-2-(5-(aminometil)-3-(3,5-difluorfenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)-4-hidroxiopirrolidin-1-il)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(5-(4-metil-1H-imidazol-1-il)-4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(5-bróm-3-(3-metoxifenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-(3-(trifluormetil)fenil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;

(S)-4-Amino-6-(1-(5-bróm-3-(3-hidroxiifenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(3-(3-metoxifenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-4-Amino-6-(1-(3-(3-hidroxiifenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(3-metoxifenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;

4-Amino-6-(1-(4-oxo-3-fenil-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)ciklopropilamino)pirimidin-5-karbonitril;

2-(1-(9H-Purin-6-ilamino)ciklopropil)-3-fenilpirolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-Amino-6-(1-(4-oxo-3-(3-(trifluormetil)fenil)-3,4-dihidropirrol[1,2-f][1,2,4]triazin-2-il)etilamino)pirimidin-5-karbonitril;

(S)-2-(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-3-(3-hidroxiifenil)-4-oxo-3,4-dihidropirrol[1,2-f][1,2,4]triazin-5-karbonitril;

(S)-2-(1-(9H-purin-6-ilamino)etil)-3-(piridin-2-il)pirolo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-2-(1-(9H-purin-6-ilamino)propil)-3-fenilimidazo[1,2-f][1,2,4]triazin-4(3H)-on;

(S)-4-amino-6-(1-(4-oxo-3-fenil-3,4-dihidroimidazo[1,2-f][1,2,4]triazin-2-il)propilamino)pirimidin-5-karbonitril;

vagy annak egy gyógyszerészetileg elfogadható sója, szolvátja, N-oxidja, sztereoizomere, vagy deuterizált származéka.

12. Vegyület az 1-től 11-ig igénypontok bármelyike szerint, mely (S)-2(1-(6-Amino-5-cianopirimidin-4-ilamino)etil)-4-oxo-3-fenil-3,4-dihidropirroló[1,2-f][1,2,4]triazin-5-karbonitril, vagy annak egy gyógyszerészetileg elfogadható sója, szolvátja, N-oxidja, sztereoizomere, vagy deuterizált származéka.

13. Vegyület az 1-től 12-ig igénypontok bármelyike szerint, olyan patológiás állapot vagy betegség kezelésében történő felhasználásra, mely enyhülésre hajlamos foszfoinozitolid-3-kináz (PI3K) gátlása esetén, ahol a patológiás állapot vagy betegség légúti betegségek; allergiás megbetegedések; gyulladásos vagy autoimmun-közvetített működési rendellenességek és neurológiai rendellenességek; kardiovaszkuláris betegségek; vírusos fertőzés; metabolizmus / endokrin működési rendellenességek; neurológiai rendellenességek és fájdalom; csontvelő és az átültetett szerv kilökődése; myelo-displázias szindróma; myeloproliferatív betegségek (MPDs); rák és rosszindulatú hematológiai tumorkok, leukémia, limfómák és szilárd tumorkok közül lett kiválasztva, vagy ahol a patológiás állapot vagy betegség leukémia, limfómák és szilárd tumorkok, reumás ízületi gyulladás, sclerosis multiplex, amiotrófiás laterális szklerózis, Crohn-betegség, fekélyes vastagbélgyulladás, szisztémás lupus erythematosus, autoimmun hemolitikus anémia, I típusú cukorbetegség, cutan vasculitis, cutan lupus erythematosus, dermatomycosis, hólyagosodásos betegségek, beleértve, de nem kizárólagosan a pemphigus vulgaris, hólyagos pemphigoid és epidermolysis bullosa, asztma, krónikus obstruktív tüdőbetegség, cisztás fibrózis, idiopátiás tüdőfibrózis, szarkoidózis, allergiás nátha, atópiás dermatitisz, kontakt dermatitisz, ekcéma, psoriasis, bazális sejt karcinóma, pikkelyes sejt karcinóma és aktinikus keratózis közül lett kiválasztva.

14. Gyógyszerészeti összetétel, mely tartalmazza az 1-től 12-ig igénypontok bármelyike által meghatározott vegyületet együttesen egy gyógyszerészetileg elfogadható hígítóanyaggal vagy hordozóanyaggal.

15. Kombinált termék, mely tartalmaz (i) egy vegyületet az 1-től 12-ig igénypontok bármelyike által meghatározottan és egy másik vegyületet Adenozid A_{2A} agonistát, kardiovaszkuláris rendellenességek kezelésére szolgáló hatóanyagot, cukorbetegség kezelésére szolgáló hatóanyagot, májbetegség kezelésére szolgáló hatóanyagot, anti-allergiás hatóanyagot, antikilonerg hatóanyagot, gyulladáscsökkentő hatóanyagot, fertőzésgátló hatóanyagot, β_2 -adrenerg antagonistát, TH_2 sejteken (CRTH2) expresszált kemotaktáns receptor homológ molekulát, kemoterápiás hatóanyagot, kortikoszteroidot, IKK β /IKK γ (IKK kináz béta vagy IKK2) inhibitor, immunszuppresszánt, Janus kináz (JAK) inhibitor, egy helyileg ható p38 mitogén-aktivált protein kináz (p38 MAPK) inhibitor, foszfodiészteráz (PDE) IV inhibitor és lép tirozin-kináz (Syk) inhibitor tartalmazó csoportból választva, emberi vagy állati test kezelésére egyidejű, külön vagy egymás utáni alkalmazás által.