

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

18 November 2021 (18.11.2021)



(10) International Publication Number

WO 2021/231319 A1

(51) International Patent Classification:

A61K 38/20 (2006.01) C07K 16/28 (2006.01)

A61P 35/00 (2006.01)

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(21) International Application Number:

PCT/US2021/031616

(22) International Filing Date:

10 May 2021 (10.05.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/022,860 11 May 2020 (11.05.2020) US

(71) Applicant: **ALKERMES PHARMA IRELAND LIMITED** [IE/IE]; One Burlington Road, Connaught House, Dublin, 4 (IE).

(72) Inventor; and

(71) Applicant (for US only): **GBORMITTAH, Francisca O.** [GH/US]; 154 Butternut Hollow, Acton, Massachusetts 01718 (US).

(72) Inventor: **ZEIDAN, Tarek A.**; 499 Lowell St., Lexington, Massachusetts 02420 (US).

(74) Agent: **SPELLBERG, Michael** et al.; LathropGPM, 28 State Street, 7th Floor, Boston, Massachusetts 02109 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: IL-2 FUSION POLYPEPTIDE COMPOSITIONS AND METHODS OF MAKING AND USING THE SAME

(57) Abstract: Provided herein are compositions comprising polypeptides comprising a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain, and methods of making and using such compositions.

WO 2021/231319 A1

IL-2 FUSION POLYPEPTIDE COMPOSITIONS AND METHODS OF MAKING AND USING THE SAME

CROSS-REFERENCE TO RELATED APPLICATION

[001] This application claims the benefit of U.S. Provisional Application Serial No. 63/022,860, filed May 11, 2020, the entire disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION

[002] This disclosure relates to compositions comprising polypeptides comprising a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain, and methods of making and using such compositions.

BACKGROUND

[003] Polypeptides comprising a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain interleukin-2 (IL-2) interleukin-2 receptor alpha (IL-2R α) hold great promise as anti-cancer agents. These polypeptides retain full ability to signal through the intermediate-affinity IL-2R complex that is expressed on memory CD8 $^+$ T cells and Natural Killer (NK) cells, but are sterically prevented from binding to the high-affinity IL-2R complex that is preferentially expressed on CD4 $^+$ FOXP3 $^+$ regulatory T cells (CD4 $^+$ Tregs) and endothelial cells. As a result of this selective IL-2R binding, the polypeptides selectively activate CD8 $^+$ T cells and NK cells, thereby promoting tumor cell killing. The inability to activate the high-affinity IL-2R on endothelial cells may also reduce the risk of toxicity due to capillary leak syndrome, a known risk of IL-2 therapies.

[004] When used for the treatment of human subjects, the aforementioned polypeptides must be stored prior to use and transported to the point of administration. Reproducibly attaining a desired level of polypeptide in a subject requires that the polypeptide be stored in a formulation that maintains the bioactivity of the polypeptide. Accordingly, there is a need in the art for stable compositions of polypeptides. Preferably, such compositions will exhibit a long shelf-life, and be stable when stored and transported.

SUMMARY

[005] The present disclosure provides compositions comprising polypeptides comprising a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain, and methods of making and using such compositions. These compositions are specifically formulated to improve the stability and shelf-life of the polypeptides contained therein.

[006] In one aspect, the disclosure provides a composition comprising:

- a) about 1 mg to about 50 mg of a polypeptide comprising a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain;
- b) sucrose;
- c) mannitol;
- c) citrate buffer; and
- d) an emulsifier.

[007] In certain embodiments, the polypeptide comprises an amino acid sequence having at least 95% identity to SEQ ID NO: 1. In certain embodiments, the polypeptide comprises the amino acid sequence of SEQ ID NO: 1. In certain embodiments, the polypeptide consists of the amino acid sequence of SEQ ID NO: 1.

[008] In certain embodiments, the composition comprises about 1 mg to about 15 mg of the polypeptide. In certain embodiments, the composition comprises about 1 mg of the polypeptide. In certain embodiments, the composition comprises about 5 mg of the polypeptide. In certain embodiments, the composition comprises about 15 mg of the polypeptide. In certain embodiments, the composition comprises about 20 mg of the polypeptide. In certain embodiments, the composition comprises about 30 mg of the polypeptide.

[009] In certain embodiments, the composition comprises about 60 mg to about 72 mg sucrose. In certain embodiments, the composition comprises about 66 mg sucrose.

[010] In certain embodiments, the composition comprises about 60 mg to about 72 mg mannitol. In certain embodiments, the composition comprises about 66 mg mannitol.

[011] In certain embodiments, the composition comprises about 4.0 mg to about 6.0 mg citrate anion. In certain embodiments, the composition comprises about 5.0 mg citrate anion.

[012] In certain embodiments, the composition comprises citric acid and sodium citrate tribasic dihydrate in a mass ratio of citric acid:sodium citrate tribasic dihydrate of between

about 1:10 to about 1:2 (i.e., about 1:10, about 1:9, about 1:8, about 1:7, about 1:6, about 1:5, about 1:4, about 1:3, and about 1:2).

[013] In certain embodiments, the composition comprises citric acid and sodium citrate tribasic dihydrate in a mass ratio of citric acid:sodium citrate tribasic dihydrate of about 1:9. In certain embodiments, the composition comprises citric acid and sodium citrate tribasic dihydrate in a mass ratio of citric acid:sodium citrate tribasic dihydrate of about 1:2.

[014] In certain embodiments, the emulsifier comprises polysorbate 20. In certain embodiments, the composition comprises about 0.20 mg to about 0.24 mg polysorbate 20. In certain embodiments, the composition comprises about 0.22 mg polysorbate 20.

[015] In certain embodiments, the composition comprises about 0.10 mg to about 0.12 mg polysorbate 20. In certain embodiments, the composition comprises about 0.11 mg polysorbate 20.

[016] In certain embodiments, the composition is a lyophilized cake.

[017] In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with a pH of about 5.5 to about 6.5. In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with a pH of about 6.1.

[018] In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with an isotonic osmolality. In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 240 to about 340 mOsm/kg. In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 280 to about 320 mOsm/kg. In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 285 mOsm/kg. In certain embodiments, dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 300 mOsm/kg.

[019] In certain embodiments, the composition is an aqueous solution.

[020] In certain embodiments, the aqueous solution comprises about 0.03 mg/mL of the polypeptide to about 0.2 mg/mL of the polypeptide.

[021] In certain embodiments, the composition comprises about 0.5 mg/mL to about 30 mg/mL of the polypeptide.

[022] In certain embodiments, the composition comprises about 1 mg/mL of the polypeptide.

[023] In certain embodiments, the composition comprises about 5 mg/mL of the polypeptide.

[024] In certain embodiments, the composition comprises about 15 mg/mL of the polypeptide.

[025] In certain embodiments, the composition comprises about 20 mg/mL of the polypeptide.

In certain embodiments, the composition comprises about 30 mg/mL of the polypeptide.

[026] In certain embodiments, the composition comprises about 25 mg/mL to about 35 mg/mL sucrose. In certain embodiments, the composition comprises about 30 mg/mL sucrose.

[027] In certain embodiments, the composition comprises about 25 mg/mL to about 35 mg/mL mannitol. In certain embodiments, the composition comprises about 30 mg/mL mannitol.

[028] In certain embodiments, the composition comprises about 10 mM to about 20 mM citrate buffer. In certain embodiments, the composition comprises about 12 mM citrate buffer. In certain embodiments, the citrate buffer is formed by the combination of 2.03 mg/mL sodium citrate tribasic dihydrate and 0.97 mg/mL citric acid monohydrate in the aqueous solution. In certain embodiments, the citrate buffer is formed by the combination of 2.91 mg/mL sodium citrate tribasic dihydrate and 0.34 mg/mL citric acid monohydrate in the aqueous solution. In certain embodiments, the citrate buffer is formed by the combination of 2.96 mg/mL sodium citrate tribasic dihydrate and 0.30 mg/mL citric acid monohydrate in the aqueous solution.

[029] In certain embodiments, the composition comprises about 0.09 mg/mL to about 0.11 mg/mL polysorbate 20. In certain embodiments, the composition comprises about 0.1 mg/mL polysorbate 20.

[030] In certain embodiments, the pH of the composition is about 5.5 to about 6.5. In certain embodiments, the pH of the composition is about 6.1.

[031] In certain embodiments, the osmolality of the composition is about 240 to about 340 mOsm/kg. In certain embodiments, the osmolality of the composition is about 280 to about 320 mOsm/kg. In certain embodiments, the osmolality of the composition is about 285 mOsm/kg. In certain embodiments, the osmolality of the composition is about 300 mOsm/kg.

[032] In certain embodiments, the composition is a single unit dose of the polypeptide.

[033] In one aspect, the disclosure provides an aqueous composition comprising:

- a) about 1 mg/mL to about 30 mg/mL of a polypeptide comprising a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain;
- b) about 25 mg/mL to about 35 mg/mL sucrose;
- c) about 25 mg/mL to about 35 mg/mL mannitol;
- d) about 10 mM to about 20 mM citrate buffer; and
- e) about 0.09 mg/mL to about 0.11 mg/mL polysorbate 20,

wherein the pH of the solution is about 5.5 to about 6.5.

[034] In certain embodiments, the polypeptide comprises an amino acid sequence having at least 95% identity to SEQ ID NO: 1. In certain embodiments, the polypeptide comprises the amino acid sequence of SEQ ID NO: 1.

[035] In one aspect, the disclosure provides an aqueous composition comprising:

- a) about 1 mg/mL to about 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
- b) about 25 mg/mL to about 35 mg/mL sucrose;
- c) about 25 mg/mL to about 35 mg/mL mannitol;
- d) about 10 mM to about 20 mM citrate buffer; and
- e) about 0.09 mg/mL to about 0.11 mg/mL polysorbate 20,

wherein the pH of the solution is about 5.5 to about 6.5.

[036] In certain embodiments, the composition comprises about 30 mg/mL sucrose.

[037] In certain embodiments, the composition comprises about 30 mg/mL mannitol.

[038] In certain embodiments, the composition comprises about 12 mM citrate buffer.

[039] In certain embodiments, the composition comprises about 0.11 mg/mL polysorbate 20.

[040] In certain embodiments, the pH of the solution is about 6.1.

[041] In certain embodiments, the composition comprises about 1 mg/mL of the polypeptide. In certain embodiments, the composition comprises about 5 mg/mL of the polypeptide. In certain embodiments, the composition comprises about 15 mg/mL of the polypeptide. In certain embodiments, the composition comprises about 20 mg/mL of the polypeptide. In certain embodiments, the composition comprises about 30 mg/mL of the polypeptide.

[042] In another aspect the disclosure provides an aqueous composition comprising:

- a) about 1, 5, 15, or 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
 - b) about 25 mg/mL to about 35 mg/mL sucrose;
 - c) about 25 mg/mL to about 35 mg/mL sucrose;
 - d) about 8 mM citrate buffer to about 14 mM citrate buffer (e.g., 8 mM, 9 mM, 10 mM, 11 mM, 12 mM, 13 mM, or 14 mM); and
 - e) about 0.1 mg/mL polysorbate 20,
- wherein the pH of the composition is about 6.1.

[043] In another aspect the disclosure provides an aqueous composition comprising:

- a) about 1, 5, 15, or 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
 - b) about 25 mg/mL to about 35 mg/mL sucrose;
 - c) about 25 mg/mL to about 35 mg/mL sucrose;
 - d) about 12 mM citrate buffer; and
 - e) about 0.1 mg/mL polysorbate 20,
- wherein the pH of the composition is about 6.1.

[044] In another aspect the disclosure provides an aqueous composition comprising:

- a) about 1, 5, 15, or 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
 - b) about 25 mg/mL to about 35 mg/mL sucrose;
 - c) about 25 mg/mL to about 35 mg/mL sucrose;
 - d) about 2 mg/mL sodium citrate tribasic dihydrate;
 - e) about 1 mg/mL citric acid monohydrate; and
 - f) about 0.1 mg/mL polysorbate 20,
- wherein the pH of the composition is about 6.1.

[045] In another aspect the disclosure provides an aqueous composition comprising:

- a) about 1, 5, 15, or 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
- b) about 25 mg/mL to about 35 mg/mL sucrose;
- c) about 25 mg/mL to about 35 mg/mL sucrose;
- d) about 2.03 mg/mL sodium citrate tribasic dihydrate;
- e) about 0.97 mg/mL citric acid monohydrate; and
- f) about 0.1 mg/mL polysorbate 20,

wherein the pH of the composition is about 6.1.

[046] In another aspect the disclosure provides an aqueous composition comprising:

- a) about 1, 5, 15, or 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
- b) about 25 mg/mL to about 35 mg/mL sucrose;
- c) about 25 mg/mL to about 35 mg/mL sucrose;
- d) about 3 mg/mL sodium citrate tribasic dihydrate;
- e) about 0.3 mg/mL citric acid monohydrate; and
- f) about 0.1mg/mL polysorbate 20,

wherein the pH of the composition is about 6.1.

[047] In another aspect the disclosure provides an aqueous composition comprising:

- a) about 1, 5, 15, or 30 mg/mL of a polypeptide comprising the amino acid sequence of SEQ ID NO: 1;
- b) about 25 mg/mL to about 35 mg/mL sucrose;
- c) about 25 mg/mL to about 35 mg/mL sucrose;
- d) about 2.91 mg/mL sodium citrate tribasic dihydrate;
- e) about 0.34 mg/mL citric acid monohydrate; and
- f) about 0.1mg/mL polysorbate 20,

wherein the pH of the composition is about 6.1.

[048] In another aspect, the disclosure provides an article of manufacture comprising any of the foregoing compositions. In certain embodiments, the article is a glass vial.

[049] In another aspect, the disclosure provides a lyophilized composition made by lyophilizing any of the foregoing aqueous solutions.

[050] In another aspect, the disclosure provides a method of making a lyophilized composition, the method comprising lyophilizing any of the foregoing aqueous solutions.

[051] In another aspect, the disclosure provides a method of making an aqueous composition, the method comprising dissolving any of the foregoing lyophilized compositions in an aqueous solvent. In certain embodiments, the aqueous solvent is water for injection. In certain embodiments, the aqueous solvent is a sodium chloride solution.

[052] In certain embodiments, the pH of the aqueous composition is adjusted to about 6.1. In certain embodiments, the pH of the aqueous composition is adjusted to about 6.1 with a base. In certain embodiments, the base is sodium hydroxide.

[053] In certain embodiments, the aqueous composition is further diluted with an aqueous solution comprising about 1% (w/w) of a surfactant. In certain embodiments, the surfactant is polysorbate 20. In certain embodiments, the aqueous solution further comprises about 0.1% (w/w) citric acid monohydrate, 0.2% (w/w) sodium citrate tribasic dihydrate, and 98.7% (w/w) water for injection.

[054] In certain embodiments, the composition comprises a pharmaceutical composition.

[055] In another aspect, the disclosure provides a method of activating natural killer cells (NK) cells in a subject, the method comprising administering to the subject an effective amount of any of the foregoing compositions.

[056] In another aspect, the disclosure provides a method of treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of any of the foregoing compositions. In certain embodiments, the cancer is renal cell carcinoma, melanoma, ovarian cancer, or lung cancer. In certain embodiments, the cancer comprises a refractory solid tumor.

DETAILED DESCRIPTION

[057] Provided herein are compositions comprising polypeptides comprising a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain, and methods of making and using such compositions.

[058] The formulations disclosed herein provide improved stability and shelf-life of the polypeptides contained therein. In particular, the polypeptide product retains biological activity, including after being lyophilized in the recited formulations and reconstituted in water for injection (WFI) or a similarly acceptable diluent. Importantly, the formulations described herein have been designed to allow the lyophilized product to be reconstituted in WFI, which is readily available to a patient or healthcare provider. When reconstituted in WFI, the formulations described herein possess a physiologically-acceptable osmolality, allowing the reconstituted product to be administered subcutaneously. This eliminates the need for a specialized diluent to reconstitute the lyophilized product with an appropriate osmolality, making it easier for a patient or healthcare provider to use the drug and, therefore, improve drug use compliance.

[059] Subcutaneous administration has advantages for drug delivery as well. When delivered via the subcutaneous route, the drug may be delivered more quickly compared to other delivery routes (e.g., intravenous). Subcutaneous delivery may also be performed

by a patient in their home, rather than by a healthcare provider in a healthcare facility. This patient-directed delivery may also improve drug use compliance.

[060] The formulations provided herein also yield a lyophilized cake that has a preferred appearance. Specifically, the cake is intact (not fragmented), has little to no shrinkage from the container (e.g., a glass vial), and have an even, concave surface.

Selected Definitions

[061] Unless otherwise defined herein, scientific and technical terms used herein have the meanings that are commonly understood by those of ordinary skill in the art. In the event of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. The use of “or” means “and/or” unless stated otherwise. The use of the term “including,” as well as other forms, such as “includes” and “included,” is not limiting.

[062] As used herein, the terms “comprising,” “including,” “having,” and grammatical variants thereof are to be taken as specifying the stated features, integers, steps or components but do not preclude the addition of one or more additional features, integers, steps, components or groups thereof. These terms encompass the terms “consisting of” and “consisting essentially of.”

[063] As used herein, the terms “circular permutation” and “circularly permuted” refer to the process of taking a linear protein, or its cognate nucleic acid sequence, and fusing the native N- and C-termini (directly or through a linker, using protein or recombinant DNA methodologies) to form a circular molecule, and then cutting (opening) the circular molecule at a different location to form a new linear protein, or cognate nucleic acid molecule, with termini different from the termini in the original molecule. Circular permutation thus preserves the sequence, structure, and function of a protein, while generating new C- and N-termini at different locations that results in an improved orientation for fusing a desired polypeptide fusion partner as compared to the original molecule.

[064] As used herein, the term “about” will be understood by persons of ordinary skill in the art and will vary to some extent on the context in which it is used. As used herein when referring to a measurable value such as an amount, a temporal duration, and the like, the term “about” is meant to encompass variations of up to $\pm 5\%$, including $\pm 5\%$, $\pm 1\%$,

and $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods.

[065] As used herein, the terms “treat,” “treated,” “treating,” or “treatment” include the diminishment or alleviation of at least one symptom associated or caused by the state, disorder or disease being treated.

[066] As used herein, the term “effective amount” in the context of the administration of a therapy to a subject refers to the amount of a therapy that achieves a desired prophylactic or therapeutic effect.

[067] As used herein, the term “patient,” “individual” or “subject” refers to a human or a non-human mammal. Non-human mammals include, for example, livestock and pets, such as ovine, bovine, porcine, canine, feline and murine mammals. In certain embodiments, the subject is a human.

IL-2 Fusion Polypeptides

[068] In one aspect, the instant disclosure provides compositions of polypeptides comprising a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain. The polypeptides employed in the compositions disclosed herein exhibit preferential binding to the intermediate-affinity IL-2R complex comprising IL-2R β and the common gamma chain, IL-2R γ) relative to the high-affinity IL-2R complex (comprising IL-2R α , IL-2R β , and IL-2R γ), and behave as selective agonists of the intermediate-affinity IL-2R complex. The design and generation of such polypeptides is described in U.S. Patent No. 9,359,415, which is hereby incorporated by reference in its entirety.

[069] An exemplary polypeptide useful for inclusion in the compositions disclosed herein is set forth below in SEQ ID. NO:1:

SKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETATIVEFLNRWITFSQ
 SIISTLTGGSSSTKKTQLQLEHLLLDLQMILNGINNYKNPKLTRMLTFKFY
 MPKKATELKHLQCLEEELKPLEEVLNLAQGSGGGSELCCCCPPEIPHATF
 KAMAYKEGTMLNCECKRGFRRIKSGSLYMLCTGNSSHSSWDNQCQCTS
 SATRNTTKQVTPQPEEQKERKTTEMQSPMQPVDQASLPGHCREPPWEN
 EATERIYHFVVGQMVYYQCVQGYRALHRGPAESVCKMTHGKTRWTQP
 QLICTG (SEQ ID NO: 1)

[070] Accordingly, in certain embodiments, the amino acid sequence of the polypeptide comprises the amino acid sequence of SEQ ID. NO: 1. In certain embodiments, the amino acid sequence of the polypeptide consists of the amino acid sequence of SEQ ID. NO: 1.

[071] The skilled worker will appreciate that amino acid sequence variants of SEQ ID. NO: 1 can also be employed in the compositions disclosed herein. For example, in certain embodiments, the amino acid sequence of the polypeptide comprises or consists of an amino acid sequence having at least 80 % (e.g., 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 %) identity to the amino acid sequence of SEQ ID. NO:1. In certain embodiments, the amino acid sequence of the polypeptide comprises or consists of an amino acid sequence having at least 95 % identity to the amino acid sequence of SEQ ID. NO:1.

[072] The skilled worker will also appreciate that amino acid sequence of the polypeptides employed in the compositions disclosed herein can be derivatized or modified, e.g., pegylated, amidated, etc.

[073] In certain embodiments, the amount of the polypeptide in a formulation is about 1 mg to about 50 mg (e.g., about 1 mg, about 2 mg, about 3 mg, about 4 mg, about 5 mg, about 6 mg, about 7 mg, about 8 mg, about 9 mg, about 10 mg, about 11 mg, about 12 mg, about 13 mg, about 14 mg, about 15 mg, about 16 mg, about 17 mg, about 18 mg, about 19 mg, about 20 mg, about 25 mg, about 30 mg, about 40 mg, about 44 mg, about 45 mg, or about 50 mg). In certain embodiments, the amount of the polypeptide is about 1 mg to about 30 mg. In certain embodiments, the amount of the polypeptide is about 1 mg to about 15 mg. In certain embodiments, the amount of the polypeptide is about 1 mg. In certain embodiments, the amount of the polypeptide is about 2.2 mg. In certain embodiments, the amount of the polypeptide is about 5 mg. In certain embodiments, the amount of the polypeptide is about 11 mg. In certain embodiments, the amount of the polypeptide is about 15 mg. In certain embodiments, the amount of the polypeptide is about 20 mg. In certain embodiments, the amount of the polypeptide is about 30 mg. In certain embodiments, the amount of the polypeptide is about 44 mg.

[074] In certain embodiments, the concentration of the polypeptide in an aqueous formulation is about 0.5 mg/mL to about 50 mg/mL. In certain embodiments, the concentration of the polypeptide is about 0.5 mg/mL to about 20 mg/mL (e.g., about 0.5 mg/mL, about 1 mg/mL, about 2mg/mL, about 3 mg/mL, about 4 mg/mL, about 5 mg/mL, about 6 mg/mL, about 7 mg/mL, about 8 mg/mL, about 9 mg/mL, about 10 mg/mL, about

11 mg/mL, about 12 mg/mL, about 13 mg/mL, about 14 mg/mL, about 15 mg/mL, about 16 mg/mL, about 17 mg/mL, about 18 mg/mL, about 19 mg/mL, about 20 mg/mL, about 25 mg/mL, about 30 mg/mL, about 35 mg/mL, about 40 mg/mL, about 45 mg/mL, or about 50 mg/mL). In certain embodiments, the concentration of the polypeptide is about 1 mg/mL. In certain embodiments, the concentration of the polypeptide is about 5 mg/mL. In certain embodiments, the concentration of the polypeptide is about 15 mg/mL. In certain embodiments, the concentration of the polypeptide is about 20 mg/mL. In certain embodiments, the concentration of the polypeptide is about 30 mg/mL.

Excipients & Buffers

[075] In certain embodiments, the compositions disclosed herein comprise one or more excipients and/or buffers.

[076] As used herein, the term “excipient” refers to any non-therapeutic agent added to the composition or formulation to provide a desired consistency, viscosity, or stabilizing effect. Suitable excipients for use in the compositions disclosed herein can act, e.g., as viscosity enhancing agents, stabilizers, solubilizing agents, etc. The excipient can be ionic or non-ionic. Suitable ionic excipients include salts such as NaCl or amino acid components such as arginine-HCl. Suitable non-ionic excipients include sugars, for example, monosaccharides (e.g., fructose, maltose, galactose, glucose, D-mannose, sorbose, etc.); disaccharides (e.g., lactose, sucrose, trehalose, cellobiose, etc.); polysaccharides (e.g., raffinose, melezitose, maltodextrins, dextrans, starches, etc.); and sugar alcohols (e.g., mannitol, xylitol, maltitol, lactitol, xylitol sorbitol (glucitol), etc.). For example, the sugar may be sucrose, trehalose, raffinose, maltose, sorbitol or mannitol. Additionally or alternatively, the sugar may be a sugar alcohol or an amino sugar. In certain embodiments, the sugar is sucrose and mannitol.

[077] In certain embodiments, the amount of the excipient (e.g., sucrose and mannitol) in a formulation is about 1 mg to about 150 mg (e.g., about 1 mg, about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 90 mg, about 100 mg, about 110 mg, about 120 mg, about 130 mg, about 140 mg, or about 150 mg). In certain embodiments, the amount of the excipient (e.g., sucrose and mannitol) in a formulation is about 30 mg to about 90 mg. In certain embodiments, the amount of the excipient (e.g., sucrose and mannitol) in a formulation is about 60 mg to

about 72 mg. In certain embodiments, the amount of the excipient (e.g., sucrose and mannitol) in a formulation is about 66 mg.

[078] In certain embodiments, the concentration of the excipient (e.g., sucrose and mannitol) in an aqueous formulation is about 1 mg/mL to about 100 mg/mL (e.g., about 1 mg/mL, about 10 mg/mL, about 20 mg/mL, about 30 mg/mL, about 40 mg/mL, about 45 mg/mL, about 50 mg/mL, about 55 mg/mL, about 60 mg/mL, about 70 mg/mL, about 80 mg/mL, about 90 mg/mL, or about 100 mg/mL). In certain embodiments, the concentration of excipient (e.g., sucrose and mannitol) is about 10 mg/mL to about 50 mg/mL. In certain embodiments, the concentration of excipient (e.g., sucrose and mannitol) is about 25 mg/mL to about 35 mg/mL. In certain embodiments, the concentration of the excipient (e.g., sucrose and mannitol) is about 30 mg/mL.

[079] Suitable buffering agents for use in the compositions disclosed herein include organic acid and salts, such as salts of citric acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid or phthalic acid; Tris, tromethamine hydrochloride, or phosphate buffer. In addition, amino acid components can also be used as buffering agents. Such amino acid component includes glycine, histidine, and methionine. In certain embodiments, the buffer is a citrate buffer. As used herein, the term "citrate buffer" refers to a pH buffering system (in aqueous or lyophilized form) that utilizes citrate ions. Citrate buffer can be made using any art recognized methods, including, by combining: (i) citric acid, trisodium citrate dihydrate, and citric acid monohydrate; or (ii) citric acid monohydrate, sodium phosphate dibasic, and citric acid. In certain embodiments, citrate buffer is made using sodium citrate dihydrate and citric acid.

[080] In certain embodiments, the amount of the buffering agent (e.g., citrate) in the formulation is about 1mg to about 10 mg (e.g., about 1 mg, about 2 mg, about 3 mg, about 4 mg, about 5 mg, about 6 mg, about 7 mg, about 8 mg, about 9 mg, about 10 mg). In certain embodiments, the amount of the buffering agent (e.g., sodium citrate) is about 5.9 mg to about 7.2 mg (e.g., about 5.9 mg, about 6.0 mg, about 6.1 mg, about 6.2 mg, about 6.3 mg, about 6.4 mg, about 6.5 mg, about 6.6 mg about, 6.7 mg about 6.8 mg, about 6.9 mg, about 7.0 mg, about 7.1 mg, or about 7.2 mg). In certain embodiments, the amount of the buffering agent (e.g., citrate) is about 6.6 mg. In certain embodiments, the amount of the citrate anion in the buffering agent (e.g., citrate) is about 4.0 mg to about 6.0 mg. In

certain embodiments, the amount of the citrate anion in the buffering agent (e.g., citrate) is about 5.0 mg.

[081] In certain embodiments, the concentration of the buffering agent (e.g., citrate) in an aqueous formulation disclosed herein is about 1 mM to about 50 mM (e.g., about 1 mM, about 2 mM, about 3 mM, about 4 mM, about 5 mM, about 6 mM, about 7 mM, about 8 mM, about 9 mM, about 10 mM, about 11 mM, about 12 mM, about 13 mM, about 14 mM, about 15 mM, about 16 mM, about 17 mM, about 18 mM, about 19 mM, about 20 mM, about 25 mM, about 30 mM, about 35 mM, about 40 mM, about 45 mM, or about 50 mM). In certain embodiments, the concentration of the buffering agent (e.g., sodium citrate) is about 11 mM to about 13 mM (e.g., about 11.1 mM, 11.2 mM, 11.3 mM, 11.4 mM, 11.5 mM, 11.6 mM, 11.7 mM, 11.8 mM, 11.9 mM, 12.1 mM, 12.2 mM, 12.3 mM, 12.4 mM, 12.5 mM, 12.6 mM, 12.7 mM, 12.8 mM, or 12.9 mM). In certain embodiments, the concentration of the buffering agent (e.g., citrate) is about 12 mM. In certain embodiments, the concentration of the buffering agent (e.g., citrate) is about 11.95 mM. In certain embodiments, the concentration of the buffering agent (e.g., citrate) is about 11.67 mM. In certain embodiments, the citrate buffer contains 2.03 mg/mL (6.90 mM) sodium citrate tribasic dihydrate and 0.97 mg/mL (5.05 mM) citric acid. In certain embodiments, the citrate buffer contains 2.91 mg/mL (9.90 mM) sodium citrate tribasic dihydrate and 0.34 mg/mL (1.77 mM) citric acid.

[082] In certain embodiments, the compositions disclosed herein have a pH of about 5.0 to about 8.0, of about 5.5 to about 7.5, of about 5.0 to about 7.0, of about 6.0 to about 8.0, or of about 6.0 to about 7.0. In certain embodiments, the compositions have a pH of about 5.4 to about 6.5. In certain embodiments, the compositions have a pH of about 5.8 to about 6.4. In certain embodiments, the compositions have a pH of about 6.1. In certain embodiments, the pH of the composition is adjusted to a pH of about 6.1. In certain embodiment, the pH is adjusted with a base. In certain embodiments, the base is a hydroxide salt, such as sodium hydroxide (NaOH) or potassium hydroxide (KOH). In certain embodiments, the composition is an aqueous composition and the pH of the aqueous composition is adjusted to a pH of about 6.1.

[083] In certain embodiments, the compositions disclosed herein have an isotonic osmolality. In certain embodiments, osmolality of the composition is about 240 to about 340 mOsm/kg. In certain embodiments, osmolality of the composition is about 280 to about 320 mOsm/kg. In certain embodiments, the osmolality of the composition is about

285 mOsm/kg. In certain embodiments, the osmolality of the composition is about 300 mOsm/kg.

[084] As used herein, the term “surfactant” refers to organic substances having amphipathic structures; i.e., they are composed of groups of opposing solubility tendencies, typically an oil-soluble hydrocarbon chain and a water-soluble ionic group. Surfactants can be classified, depending on the charge of the surface-active moiety, into anionic, cationic and dispersing agents for various pharmaceutical compositions and preparations of biological materials. Suitable surfactants for use in the compositions disclosed herein include non-ionic surfactants, ionic surfactants and zwitterionic surfactants. Typical surfactants for use with the invention include sorbitan fatty acid esters (e.g., sorbitan monocaprylate, sorbitan monolaurate, sorbitan monopalmitate), sorbitan trioleate, glycerine fatty acid esters (e.g., glycerine monocaprylate, glycerine monomyristate, glycerine monostearate), polyglycerine fatty acid esters (e.g., decaglyceryl monostearate, decaglyceryl distearate, decaglyceryl monolinoleate), polyoxyethylene sorbitan fatty acid esters (e.g., polyoxyethylene sorbitan monolaurate, polyoxyethylene sorbitan monooleate, polyoxyethylene sorbitan monostearate, polyoxyethylene sorbitan monopalmitate, polyoxyethylene sorbitan trioleate, polyoxyethylene sorbitan tristearate), polyoxyethylene sorbitol fatty acid esters (e.g., polyoxyethylene sorbitol tetrastearate, polyoxyethylene sorbitol tetraoleate), polyoxyethylene glycerine fatty acid esters (e.g., polyoxyethylene glyceryl monostearate), polyethylene glycol fatty acid esters (e.g., polyethylene glycol distearate), polyoxyethylene alkyl ethers (e.g., polyoxyethylene lauryl ether), polyoxyethylene polyoxypropylene alkyl ethers (e.g., polyoxyethylene polyoxypropylene glycol, polyoxyethylene polyoxypropylene propyl ether, polyoxyethylene polyoxypropylene cetyl ether), polyoxyethylene alkylphenyl ethers (e.g., polyoxyethylene nonylphenyl ether), polyoxyethylene hydrogenated castor oils (e.g., polyoxyethylene castor oil, polyoxyethylene hydrogenated castor oil), polyoxyethylene beeswax derivatives (e.g., polyoxyethylene sorbitol beeswax), polyoxyethylene lanolin derivatives (e.g., polyoxyethylene lanolin), and polyoxyethylene fatty acid amides (e.g., polyoxyethylene stearic acid amide); C 10-C 18 alkyl sulfates (e.g., sodium cetyl sulfate, sodium lauryl sulfate, sodium oleyl sulfate), polyoxyethylene C 10-C 18 alkyl ether sulfate with an average of 2 to 4 moles of ethylene oxide units added (e.g., sodium polyoxyethylene lauryl sulfate), and C1-C 18 alkyl sulfosuccinate ester salts (e.g., sodium lauryl sulfosuccinate

ester); and natural surfactants such as lecithin, glycerophospholipid, sphingophospholipids (e.g., sphingomyelin), and sucrose esters of C 12-C 18 fatty acids. A composition may include one or more of these surfactants. In certain embodiments, the compositions disclosed herein comprise polyoxyethylene sorbitan fatty acid esters e.g., polysorbate 20, 40, 60 or 80. In certain embodiments, the compositions disclosed herein comprise polysorbate 20.

[085] In certain embodiments, the amount of the surfactant (e.g., polysorbate 20) in the formulation is about 0.1 mg to about 1 mg (e.g., about 0.1 mg, about 0.15 mg, about 0.2 mg, about 0.25 mg, about 0.3 mg, about 0.35 mg, about 0.4 mg, about 0.45 mg, about 0.5 mg, about 0.55 mg, about 0.6 mg, about 0.65 mg, about 0.7 mg, about 0.75 mg, about 0.8 mg, about 0.85 mg, about 0.9 mg, about 0.95 mg, or about 1 mg). In certain embodiments, the amount of the surfactant (e.g., polysorbate 20) is about 0.15 mg to about 0.3 mg (e.g., about 0.16 mg, about 0.17 mg, about 0.18 mg, about 0.19 mg, about 0.21 mg, about 0.22 mg, about 0.23 mg, about 0.24 mg, about 0.26 mg, about 0.27 mg, about 0.28 mg, or about 0.29 mg). In certain embodiments, the amount of the surfactant (e.g., polysorbate 20) is about 0.20 mg to about 0.24 mg. In certain embodiments, the amount of the surfactant (e.g., polysorbate 20) in an aqueous formulation is about 0.22 mg. In certain embodiments, the composition comprises about 0.10 mg to about 0.12 mg polysorbate 20. In certain embodiments, the composition comprises about 0.11 mg polysorbate 20.

[086] In certain embodiments, the concentration of the surfactant (e.g., polysorbate 20) in an aqueous formulation is about 0.01 mg/mL to about 1 mg/mL (e.g., about 0.01 mg/mL, about 0.1 mg/mL, about 0.2 mg/mL, about 0.3 mg/mL, about 0.4 mg/mL, about 0.5 mg/mL, about 0.6 mg/mL, about 0.7 mg/mL, about 0.8 mg/mL, about 0.9 mg/mL, or about 1 mg/mL). In certain embodiments, the concentration of the surfactant (e.g., polysorbate 20) is about 0.05 mg/mL to about 0.15 mg/mL (e.g., about 0.05 mg/mL, about 0.06 mg/mL, about 0.07 mg/mL, or about 0.08 mg/mL about 0.09 mg/mL, about 0.1 mg/mL, about 0.11 mg/mL, about 0.12 mg/mL, about 0.13 mg/mL, about 0.14 mg/mL, or about 0.15 mg/mL.). In certain embodiments, the concentration of the surfactant (e.g., polysorbate 20) is about 0.09 mg/mL to about 0.11 mg/mL. In certain embodiments, the concentration of the surfactant (e.g., polysorbate 20) in an aqueous formulation is about 0.1 mg/mL.

[087] It will be understood to those of skill in the art that the components of the compositions and compositions of the present invention may be described by units other

than mg/mL. For example, the components of the compositions and compositions of the present invention may be described in units of molarity. The components of the compositions and compositions of the present invention may be further described in units of weight or mass percent.

Lyophilization

[088] In one aspect, the instant disclosure provided lyophilized compositions (e.g., lyophilized cake) of the polypeptides disclosed herein, and methods of making the same.

[089] Lyophilization generally includes three main stages: freezing, primary drying and secondary drying. Freezing is necessary to convert water to ice or some amorphous formulation components to the crystalline form. Primary drying is the process step when ice is removed from the frozen product by direct sublimation at low pressure and temperature. Secondary drying is the process step when bounded water is removed from the product matrix utilizing the diffusion of residual water to the evaporation surface. Product temperature during secondary drying is normally higher than during primary drying. See, Tang X. et al. (2004) "Design of freeze-drying processes for pharmaceuticals: Practical advice," *Pharm. Res.*, 21:191-200; Nail S.L. et al. (2002) "Fundamentals of freeze-drying," in *Development and manufacture of protein pharmaceuticals*. Nail SL editors. New York: Kluwer Academic/Plenum Publishers, pp 281-353; Wang et al. (2000) "Lyophilization and development of solid protein pharmaceuticals," *M J Pharm.*, 203:1-60; Williams NA et al. (1984) "The lyophilization of pharmaceuticals; A literature review." *J. Parenteral Sci. Technol*, 38:48-59; and WO 2010/148337 A1.

[090] Because of the variations in temperature and pressure through the lyophilization process, an appropriate choice of excipients or other components such as stabilizers, buffering agents, bulking agents, and surfactants are needed to prevent the polypeptides disclosed herein from degradation (e.g., protein aggregation, deamidation, and/or oxidation) during freeze-drying and storage. The lyophilized compositions disclosed herein contain a particular combination of constituents allow for stable long-term storage of the polypeptides disclosed herein that comprise a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain.

[091] In another aspect, the disclosure provides a lyophilized composition made by lyophilizing any one of the aqueous compositions disclosed herein that comprise a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α

chain. In certain embodiments, the lyophilized composition is a lyophilized cake. In certain embodiments, the lyophilized composition is made by lyophilizing any one of the aqueous compositions disclosed herein following the lyophilization protocol recited in Table 11A or Table 11B.

[092] In another aspect, the disclosure provides a method of making a lyophilized composition, the method comprising lyophilizing any one of the aqueous compositions disclosed herein that comprise a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain. In certain embodiments, the method of making a lyophilized composition comprises following the lyophilization protocol recited in Table 11A or Table 11B.

[093] In another aspect, the disclosure provides a method of making an aqueous composition, the method comprising dissolving in an aqueous solvent any one of the lyophilized compositions disclosed herein that comprise a circularly permuted interleukin-2 (IL-2) fused to the extracellular portion of an IL-2R α chain. In certain embodiments, the lyophilized composition is a lyophilized cake. In certain embodiments, the lyophilized composition is dissolved in 1.1 ml of water. In certain embodiments, the lyophilized composition is dissolved in 2.2 ml of water.

Uses of Polypeptide Compositions

[094] The compositions disclosed herein are particularly useful for the treatment, prevention, or amelioration of any disease or disorder associated with Interleukin 2 receptor signaling.

[095] In one aspect, provided is a method of activating natural killer cells (NK) cells in a subject, the method comprising administering to the subject an effective amount of any one of the compositions disclosed herein that comprise a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain

[096] In another aspect, provided is a method of treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of any one of the compositions disclosed herein that comprise a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain. Cancers suitable for treatment using the composition disclosed herein include renal cell carcinoma, melanoma, ovarian cancer, and lung cancer. In certain embodiments, the cancer comprises a refractory solid tumor.

[097] In certain embodiments, the composition is administered subcutaneously.

[098] In certain embodiments, the composition is administered subcutaneously at a dose of about 1 mg to about 15 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 1 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 2 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 3 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 4 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 5 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 6 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 7 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 8 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 9 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 10 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 11 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 12 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 13 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 14 mg. In certain embodiments, the composition is administered subcutaneously at a dose of about 15 mg.

[099] In certain embodiments, the composition is administered subcutaneously once a week (Q1W), once every two weeks (Q2W), or once every three weeks (Q3W).

[0100] In certain embodiments, the composition is administered subcutaneously at a dose of about 1 mg to about 15 mg once a week (Q1W), once every two weeks (Q2W), or once every three weeks (Q3W).

[0101] In certain embodiments, the composition is administered subcutaneously at a dose of about 3 mg once a week (Q1W). In certain embodiments, the composition is administered subcutaneously at a dose of about 6 mg once every three weeks (Q3W).

[0102] In certain embodiments, the melanoma is one or both of mucosal melanoma or advanced cutaneous melanoma.

[0103] It will be readily apparent to those skilled in the art that other suitable modifications and adaptations of the methods described herein may be made using suitable equivalents without departing from the scope of the embodiments disclosed herein.

Having now described certain embodiments in detail, the same will be more clearly understood by reference to the following examples, which are included for purposes of illustration only and are not intended to be limiting.

EXAMPLES

[0104] The invention is further illustrated by the following examples, which should not be construed as further limiting. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of organic synthesis, cell biology, cell culture, molecular biology, transgenic biology, microbiology and immunology, which are within the skill of the art.

Example 1 – Design and Testing of Polypeptide Compositions

[0105] In an effort to determine the optimal subcutaneous formulation for Polypeptide A (a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain comprising the amino acid of SEQ ID NO: 1), several formulations of Polypeptide A were tested for their effects on protein stability, pH stability, physio-chemical behavior, lyophilized cake uniformity, and resistance to adhering to the storage vial post-lyophilization. Two main objectives were sought in this study. The first objective was to produce a lyophilized cake which, when reconstituted with off-the shelf diluent (i.e. water for injection), results in an isotonic solution ready for administration. A formulation that results in a non-isotonic solution when reconstituted with WFI would not be amenable to subcutaneous administration. The second objective was to produce a lyophilized cake with optimal cake appearance that has minimal cake shrinkage. An improved cake appearance may make the drug product more visually appealing to a patient or healthcare provider, thus potentially improving compliance in drug use. Table 1 below recites the specific components and their concentrations for the Polypeptide A formulation that was initially designed for intravenous administration.

Table 1: Polypeptide A intravenous administration formulation

Component	Function	Concentration (mg/mL)
Polypeptide A	Protein	1, 5 and 15
Sucrose	Protein Stabilizer	50
Sodium citrate dihydrate	Buffer	2.03
Citric acid monohydrate	Buffer	0.97
Polysorbate 20	Surfactant	0.1

	Buffer pH	6.1
	Osmolarity in WFI (mOsm/kg)	~125 mOsm/kg

Analytical Methods Used

Fourier Transform Infrared (FTIR) Spectroscopy

[0106] FTIR analysis was performed using PROTA FTIR Protein Analyzer equipped with a CaF₂ Biocell and ATR cell for solid sample analysis. Approximately 10 μ L of liquid sample or 10 mg of lyophilized powder sample was loaded for the analysis. Absorbance signals were processed by subtracting interfering signals from background lyophilized placebo and buffer where appropriate. The processed data were finally converted to second derivative signals to improve the resolution with parameters set as 100 scans with a 4 cm^{-1} resolution. To identify the percentage of the different structural elements of the Polypeptide A native protein, the subtracted spectrum was processed through the protein secondary structure database. The percent similarity between the native state and the dried state was calculated based on the area of overlap from 1700 to 1600 cm^{-1} or 1800 to 1400 cm^{-1} .

Osmolality Analysis

[0107] The osmolality of prepared formulations were determined using wescor vapro vapor pressure instrument. For each analysis, approximately 10 μ L of liquid sample was utilized.

Visual Inspection

[0108] All vials were inspected against a background that provided a clear picture when digital photos were captured.

Sub-ambient and Dry Powder Differential Scanning Calorimetry (DSC)

[0109] DSC analysis was performed using TA Q20 with a refrigerated cooling system I. For sub-ambient (frozen state) DSC, approximately 15 μ L of the formulated drug substance was loaded into DSC pan and hermetically sealed. The sample was then cooled to -90°C at 10°C/minute. The pans were held in the sample chamber for 2 minutes prior to warming to 30°C at 10°C/minute. During annealing, the sample was warmed to -10°C after holding at -90°C for 2 minutes, then cooled back to -90°C prior to warming to 30°C at 10°C/minute.

[0110] For dry powder high temperature DSC, vials were placed into a dry box that was purged to below 8% RH with dry air. Aliquots of the lyophilized samples were removed

and hermetically sealed in DSC pans. Thermal scanning was carried out from 20°C to 180°C at 1°C/minute using a modulation program of $\pm 1^\circ\text{C}$ every 120 seconds and the resulting reversing and non-reversing heat flow measured.

Screening Study 1

[0111] The first screening involved various formulations to understand how different combinations of stabilizers, bulking agents, etc. would impact the glass transition or collapse temperature (T_g') and the isotonicity. All formulations contained 12 mM sodium citrate buffer at pH 6.11 as the base formulation prior to the addition of the screening excipients. Tween 21 was used as a surface stabilizer. The formulations evaluated in this study with sub-ambient DSC and osmolality analyses are summarized in Table 2.

Table 2: Screening Study 1 formulations

Formulation #	Polypeptide A (mg/mL)	Sucrose (mg/mL)	Mannitol (mg/mL)	Osmolality (mOsm/kg)	T_g' °C
1	5	5	15	123	-28.81
2	5	5	10	Not measured	-27.91
3	5	10	10	66	-28.00
4	5	5	-	59	-25.70

[0112] The osmolality values for Formulations 1-4 were too low, a value closer to physiological (280-320 mOsm/kg) was desired. Formulation 1 showed signs of being metastable in the frozen phase; hence the lyophilization cycle was annealed at -10°C , which successfully converted this metastable state to a stable eutectic phase. In an attempt to increase the osmolality of these formulations, glycine was added to Formulation 4 at a concentration of 30 mg/mL. This resulted in an osmolality of 413 mOsm/kg. Based on this result, glycine formulations were prepared to replace Formulations 1 and 2 to target osmolality of 290 mOsm/kg, with a corresponding glass transition temperature of -28.23°C by DSC. Additionally, concentrations of sucrose and mannitol excipients in Formulations 3 and 4 were adjusted to increase the osmolality values. Four new formulations were lyophilized and analyzed in screening study 2.

Screening Study 2

[0113] The formulations tested in screening study 2 and their associated osmolality values are shown below in Table 3. All formulations contained 12 mM sodium citrate buffer at pH 6.11 as the base formulation prior to the addition of the screening excipients. Tween 21 was used as a surface stabilizer. Note that placebos were prepared for each formulation to be used for secondary structure analysis following lyophilization in a VirTis Genesis SQ Super XL – 70 freeze dryer.

Table 3: Screening Study 2 formulations with Osmolality and Percent Similarity of Secondary Structure by FTIR

Formulation #	Polypeptide A (mg/mL)	Sucrose (mg/mL)	Mannitol (mg/mL)	Glycine (mg/mL)	Osmolality (mOsm/kg)	% similarity to native
5	5	5	-	15	268	52.6
6	5	10	-	15	248	65.3
7	5	5	30	-	206	51.9
8	5	10	30	-	224	80.6

[0114] All samples were sterile filtered under aseptic conditions followed by filling into sterilized 2 cc vials at fill volumes of 0.5 mL. Table 4 details lyophilization cycle parameters used for screening study 2.

Table 4: Lyophilization Cycle Parameters for Screening Study 2

Step	Temperature (°C)	Time (Minutes)	Ramp Rate (°C /min)	Chamber Pressure (mTorr)
Loading	5	-	-	-
Freezing and Annealing	5 to -40	100	0.5	-
	Hold at -40	30	-	-
	-40 to -10	60	0.5	-
	Hold at -10	60	-	-
	-10 to -40	60	0.5	-
	Hold at -40	60	-	100
Primary Drying	-40 to -25	30	0.5	100
	Hold at -25	600	-	100
Secondary Drying	-25 to 10	100	0.5	100
	Hold at 10	60	-	100
Stoppering	10	Sealed under partial vacuum (nitrogen back fill)		

[0115] Following lyophilization, all lyophilized cakes showed no sign of melt-back or collapse (no images were taken). The samples were tested for secondary structure and compared to the native state structure collected in the liquid cell. FTIR spectra (second

derivative) of native Polypeptide A were overlaid with screening study 2 formulations. The percent similarity between the native and the dried state shown in Table 3 was calculated based on the second derivative area of overlap from 1800 to 1400 cm⁻¹. Formulation 8 showed the best retention of secondary structure after drying, however the tonicity was low.

Screening Study 3

[0116] Based on the above results, a third screening study was conducted to investigate different ratios of stabilizer vs. bulking agent. In addition, trehalose, a non-reducing disaccharide and polyvinyl pyrrolidone (PVP), a polymer, were introduced to evaluate their effect as stabilizers and also to evaluate its effectiveness to improve Tg'. Eight formulations were prepared for screening study 3 and freeze-dried using lyophilization cycle parameters described in Table 4. The formulation matrix for this study is detailed in Table 5 below. Tween 20 (PS20) was used in this study at 0.1 mg/mL for Formulations 9-16.

Table 5: Screening Study 3 formulations with Percent Similarity of Secondary Structure by FTIR

Formulation #	Polypeptide A (mg/mL)	Sucrose (mg/mL)	Trehalose (mg/mL)	Mannitol (mg/mL)	PVP (mg/mL)	% similarity to native
9	5	5	-	30	-	40.3
10	5	10	-	30	-	73.7
11	5	15	-	20	-	61.5
12	5	15	-	20	10	45.4
13	5	20	-	20	-	58.4
14	5	-	15	15	-	81.2
15	5	-	15	15	10	76.0
16	5	-	20	15	-	71.8

[0117] FTIR spectra were measured as described above to determine the percent similarity to native Polypeptide A. In addition to Formulation 8, the formulations showing the most promising results with >70% similarity were Formulations 10, 14, 15, and 16. Based on this observation, screening study 4 was designed with five formulations prepared with lower amounts of mannitol, and higher amounts of disaccharide. This combination appeared to favor a higher retention of secondary structure.

Screening Study 4

[0118] Table 6 depicts 5 formulations that were prepared as described above for screening study 3 and freeze-dried using lyophilization cycle parameters described in Table 4. No osmolality measurements were performed. Tween 20 (PS20) was used in this study at 0.1 mg/mL for Formulations 17-21.

Table 6: Screening Study 4 formulations with Percent Similarity of Secondary Structure by FTIR

Formulation #	Polypeptide A (mg/mL)	Sucrose (mg/mL)	Trehalose (mg/mL)	Mannitol (mg/mL)	% similarity to native
17	5	15	-	10	78.4
18	5	15	-	5	84.5
19	5	-	15	10	64.5
20	5	-	15	5	67.5
21	5	-	25	-	95.2

[0119] FTIR spectra were measured as described above to determine the percent similarity to native Polypeptide A. Based on the above results, a higher amount of disaccharide was confirmed to have a higher impact on the retention of secondary structure. The next screening study was designed to investigate higher concentrations of disaccharides in combination with bulking agents.

Fine Tuning the Tonicity

[0120] Prior to the start of the next screening, the isotonicity of various formulations were fine tuned. A number of formulations containing excipient combinations were prepared in 12 mM sodium citrate buffer at pH 6.1 with higher amounts of disaccharides and varying amounts of bulking agents, followed by osmolality measurement. The different formulations and the resulting osmolality values are shown in Table 7. The formulations all contain Polypeptide A at 5 mg/mL.

Table 7: Osmolality screening

Formulation #	Sucrose (mg/mL)	Trehalose (mg/mL)	Mannitol (mg/mL)	Glycine (mg/mL)	Osmolality (mOsm/kg)
22	20	-	-	40	612
23	25	-	-	15	303
24	30	-	-	10	258
25	-	25	10	-	164
26	-	25	20	-	209
27	-	30	30	-	289

28	-	30	-	20	372
29	-	20	-	20	293
30	-	25	-	15	303

Screening Study 5

[0121] Table 8 depicts 5 formulations that were prepared as described above for screening study 3. Tween 20 (PS20) was used in this study at 0.1 mg/mL for Formulations 31-35. Glycine in Formulations 33 and 34 interfered with the FTIR analysis, hence there is no percent similarity data reported. Moreover, generating an effective lyophilization cycle with glycine was more difficult than mannitol. Therefore, mannitol-containing formulations were pursued further.

Table 8: Screening Study 5 formulations with Osmolality and Percent Similarity of Secondary Structure by FTIR

Formulation #	Sucrose (mg/mL)	Trehalose (mg/mL)	Mannitol (mg/mL)	Glycine (mg/mL)	Osmolality (mOsm/kg)	% similarity to native
31	-	30	30	-	299	80.8
32	30	-	30	-	300	77.9
33	-	30	-	10	252	-
34	-	20	-	15	292	-
35	-	20	35	-	282	89.0

[0122] All samples were sterile filtered under aseptic conditions followed by filling into sterilized 2 cc vials at fill volumes of 0.5 mL. Table 9 details lyophilization cycle parameters used for screening study 5.

Table 9: Lyophilization Cycle Parameters for Screening Study 5

Step	Temperature (°C)	Time (Minutes)	Ramp Rate (°C /min)	Chamber Pressure (mTorr)
Loading	5	-	-	-
Freezing and Annealing	5 to -45	100	0.5	-
	Hold at -45	30	-	-
	-45 to -10	70	0.5	-
	Hold at -10	40	-	-
	-10 to -45	70	0.5	-
	Hold at -45	30	-	90
Primary Drying	-45 to -30	30	0.5	90
	Hold at -30	1260	-	90
	-30 to 10	100	0.5	90

Secondary Drying	Hold at 10	60	-	90
Stoppering	10	Sealed under partial vacuum (nitrogen back fill)		

[0123] After freeze-drying, the secondary structure was determined and compared to the native state as described above and their percent similarity calculated.

Leading Formulation Candidates

[0124] Based in part on FTIR and osmolality results, Formulations 31 and 32 were tested for appearance, glass transition temperature in the frozen state (T_g'), and glass transition temperature in the dried state (T_g). The intravenous administration formulation for Polypeptide A was used as a comparator. Compositions of the three formulations evaluated in this screen are described in Table 10.

Table 10: Formulations with Sub-Ambient Glass Transition Temperature

Formulation #	Polypeptide A (mg/mL)	Polysorbate 20 (mg/mL)	Sucrose (mg/mL)	Trehalose (mg/mL)	Mannitol (mg/mL)	T_g' (°C)
36	5	0.1	50	-	-	-29.42
37 (31 in Table 8)	5	0.1	30	-	30	-38.39
38 (32 in Table 8)	5	0.1	-	30	30	-37.83

[0125] All formulations (placebo and active) were prepared and sterile filtered under aseptic conditions followed by filling into sterilized 5 cc vials at fill volumes of 2.28 mL. An aliquot (15 μ L) of each formulation candidate was analyzed by sub-ambient DSC to assess their frozen state profiles. Based on the T_g' results, the formulations were lyophilized using cycle parameters listed in Table 11A. An annealing step is used to ensure complete crystallization of the bulking agent, mannitol. An alternative lyophilization cycle which was similarly effective at lyophilizing the formulations is recited below in Table 11B.

Table 11A: Lyophilization Cycle Parameters for Leading Formulation Candidates

Step	Temperature (°C)	Time (Minutes)	Ramp Rate (°C /min)	Chamber Pressure (mTorr)
Loading	5	-	-	-
Freezing and Annealing	5 to -42	94	0.5	-
	Hold at -42	30	-	-
	-42 to -10	64	0.5	-
	Hold at -10	60	-	-
	-10 to -40	20	0.5	-
Primary Drying	Hold at -40	60	-	100
	-45 to -30	20	0.5	100
Secondary Drying	Hold at -30	2580	-	100
	-30 to 25	110	0.5	100
Stopping	Hold at 25	540	-	100
	25	Sealed under partial vacuum (nitrogen back fill)		

Table 11B: Alternative Lyophilization Cycle Parameters

Step	Temperature (°C)	Time (Minutes)	Chamber Pressure (Torr)
Loading	5	-	760
Loading	5	30	550
Freezing and Annealing	5 to -45	150	550
	Hold at -45	120	550
	-45 to -10	105	550
	Hold at -10	180	550
	-10 to -45	105	550
Primary Drying	Hold at -45	120	550
	Hold at -45	30	0.1
	-45 to -25	40	0.1
Secondary Drying	Hold at -25	3000	0.1
	-25 to 30	220	0.1
Stopping	Hold at 30	1440	0.1
	30	-	600

[0126] Following lyophilization, all samples were evaluated based on cake appearance, glass transition temperature in the dried state (T_g), pH, reconstitution time, osmolality, concentration, moisture content, SEC, RP and potency assay.

Lyophilized Cake Appearance

[0127] The vials contained white intact cakes. The formulation containing 50 mg/mL sucrose (Formulation 36) displayed slight cake shrinkage. All other formulations were fully intact with no signs of shrinkage.

High Temperature DSC of Lyophilized Cake

[0128] Table 12 summarizes the results after high temperature DSC analysis. Glass transition temperature (T_g) values for Formulation 36 and 37 were determined at ~82 °C, attributed to sucrose. The glass transition of the formulation containing trehalose (Formulation 38) could not be detected under these conditions. The melting peak detected for Formulation 36 at 158 °C was attributed to the melting of sucrose. The melting peaks detected for Formulation 37 and 38 at 120-121 °C were attributed to the melting of mannitol.

Table 12: Summary of T_g and Melting Temperature

Formulation #	Sample weight (mg)	Glass Transition Temperature, T _g (°C)	Melting Temperature, T _m (°C)
36	4.5	81.86	157.86
37 (31 in Table 8)	5.2	81.86	119.69
38 (32 in Table 8)	4.9	-	120.80

pH, Reconstitution Time, Osmolality, Concentration, and Moisture Content for Leading Formulation Candidates

[0129] WFI was used to reconstitute the lyophilized cake. Table 13 summarizes the results for pH, Reconstitution Time, Osmolality, Concentration, and Moisture Content.

Table 13: Summary of T_g and Melting Temperature

Formulation #	pH	Osmolality (mOsm/kg)	Recon. Time with stirring and swirling (seconds)	Moisture Content (%)	Concentration (mg/mL)
36	6.18	188	20	1.55	5.127
37 (31 in Table 8)	6.20	296	22	1.74	5.058
38 (32 in Table 8)	6.17	289	16	1.79	4.566

Size Exclusion (SE) HPLC and Reverse Phase (RP) HPLC for Leading Formulation Candidates

[0130] SE-HPLC and RP-HPLC analysis of Formulations 36, 37, and 38 was performed. The integrated results for the SE-HPLC analysis showed the largest peak having a peak area of >98% for each formulation. The integrated results for the RP-HPLC analysis showed the largest peak having a peak area of >85% for each formulation.

Potency Assay for Leading Formulation Candidates

[0131] To ensure that the excipients would not impact bioactivity, Formulations 36-38 were compared using a pSTAT5 Activity Assay. The dose response curves were similar and comparable to that of a Polypeptide A Reference Standard. Data analysis in Table 14 showed EC50 values that were greater than 75% potency relative to the Reference Standard RT, which is acceptable.

Table 14: pSTAT5 activity assay results

Parameter	Formulation 36	Formulation 37	Formulation 38
Bottom Asymptote	0.116	0.109	0.072
Top Asymptote	0.757	0.786	0.755
Hill Slope	1.29	1.24	1.34
EC50	11.59	12.49	16.07
Relative Potency	84%	78%	78%

[0132] The pSTAT5 activity assay was performed by measuring binding of the formulations to HH cells (a human T lymphocyte cell line which have the $\beta\gamma$ IL2 receptor isoform present on their surface). Polypeptide A binding was measured by determining the amount of phosphorylated STAT5 (phospho-STAT5 or pSTAT5) present in the HH cells after contact with each formulation, using an ELISA assay. The Invitrogen InstantOne ELISA phosphor-STAT5 alpha/beta (pTyr694/pTyr699) kit was used to perform the ELISA assay.

[0133] The drug product sample was prepared by reconstituting the sample in 2.2 mL WFI. The sample was visually inspected to ensure that contents were free of visible particulates.

[0134] A sample diluent was prepared by adding 25 mL of Fetal Bovine Serum (FBS) to 500 mL of Hanks Balanced Salt Solution (HBSS) for a final concentration of 5% FBS and warmed to 37°C. A wash buffer comprising Phosphate Buffered Saline (PBS) with 0.05% Tween 20 was used.

[0135] Samples and standards were diluted to final protein concentrations in the assay of 750 ng/mL, 250 ng/mL, 83 ng/mL, 28 ng/mL, 9.3 ng/mL, 3.1 ng/mL, 1.0 ng/mL, and 0.3 ng/mL. A stock solution of HH cells at a density of approximately 1.2×10^6 cells/mL was prepared and 50 μ l of the cell stock solution was added to each well of a 96-well plate that contained the diluted sample or standard. The cells were incubated at 37 °C for 30 minutes. After incubation, the cells were lysed in a cell lysis buffer for 10 minutes. After lysis, 50 μ l of the lysed cell mix was transferred to an ELISA plate followed by 50 μ l of phospho-STAT5 A/B antibody cocktail. The mix was then incubated for 1 hour then washed 3 times with wash buffer. 100 μ l of a detection reagent was then added to each well and the plate was incubated for 15 minutes. 100 μ l of a stop solution was then added to each well and the plate was read at 450 nm on a microplate reader.

[0136] Individual EC50 values were measured and % relative standard deviation (RSD) was calculated for the reference standard EC50 values and for the control EC50 values.

[0137] The geometric mean of the three EC50 values for reference standard (Ref Std EC50GM) and the geometric mean of three EC50 values for the control (Control EC50) were calculated. The relative potency for the control was calculated using the following equation: $\text{Relative Potency} = (\text{RefStd EC50GM}) / (\text{Control EC50}) \times 100\%$.

[0138] The samples were calculated in the same manner. Results of the assay were determined by the following equation: $\text{Relative Potency} = (\text{RefStd EC50GM}) / (\text{Test sample EC50}) \times 100\%$.

Initial Conclusions

[0139] Based on all the results generated and its similarity to the intravenous administration Polypeptide A formulation, which also contains sucrose as the stabilizing agent, Formulation 37 was selected. The formulation composition is shown in Table 15. The selected formulation results in an isotonic solution when reconstituted using WFI, which will increase its usability for direct subcutaneous drug delivery.

[0140] The screening studies suggested that the preservation of Polypeptide A secondary structure is improved significantly when high amounts of disaccharides such as sucrose or trehalose is present in the formulation. Additionally, mannitol was determined to be a superior bulking agent and tonicity modifier compared to glycine.

Table 15: Polypeptide A Formulation 37

Component	Function	Concentration (mg/mL)
Polypeptide A	Protein	1 and 5
Sucrose	Protein Stabilizer	30
Mannitol	Bulking Agent	30
Sodium citrate tribasic dihydrate	Buffer	2.03
Citric acid monohydrate	Buffer	0.97
Polysorbate 20	Surfactant	0.1

[0141] Following identification of Formulation 37 recited above, the concentrations of sodium citrate tribasic dihydrate and citric acid monohydrate were optimized to avoid a pH titration step to achieve a pH of 6.1. Table 16 below describes altered Formulation 37-2.

Table 16: Polypeptide A Formulation 37-2

Component	Function	Concentration (mg/mL)
Polypeptide A	Protein	1 and 5
Sucrose	Protein Stabilizer	30
Mannitol	Bulking Agent	30
Sodium citrate tribasic dihydrate	Buffer	2.91
Citric acid monohydrate	Buffer	0.34
Polysorbate 20	Surfactant	0.1

Further Testing of Mannitol / Sucrose Levels

[0142] The amounts of sucrose and mannitol were altered from Formulation 37 as shown in Table 17 to test osmolality and monitor lyophilized cake appearance. Each formulation below in Table 17 had a sodium citrate buffer at 12 mM and a pH of 6.1. Following lyophilization, Formulation 37 had the best lyophilized cake appearance, showing more of a concave top appearance and no shrinkage on the walls compared to the other formulations. The improved cake appearance may make the drug product more visually appealing to a patient or healthcare provider, thus potentially improving compliance in drug use.

Table 17: Formulations with Altered Levels of Mannitol and Sucrose

Sample #	Formulation #	Polypeptide A	Polysorbate 20	Sucrose (mg/mL)	Mannitol (mg/mL)	Osmolality (mOsm/kg)
----------	---------------	---------------	----------------	-----------------	------------------	----------------------

		(mg/mL)	(mg/mL)			
1	37	5	0.1	30	30	285
2	39	5	0.1	20	40	311
3	40	5	0.1	13.5	40	292
4	41	5	0.1	10	40	281
5	42	5	0.1	90	-	296
6	36	5	0.1	50	-	179

What is claimed:

1. A composition comprising:

- a) about 1 mg to about 50 mg of a polypeptide comprising a circularly permuted IL-2 fused to the extracellular portion of an IL-2R α chain;
- b) sucrose;
- c) mannitol;
- c) citrate buffer; and
- d) an emulsifier.

2. The composition of claim 1, wherein the polypeptide comprises an amino acid sequence having at least 95% identity to SEQ ID NO: 1.

3. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 1.

4. The composition of any one of claims 1-3, wherein the composition comprises about 1 mg to about 15 mg of the polypeptide.

5. The composition of any one of claims 1-4, wherein the composition comprises about 1 mg of the polypeptide.

6. The composition of any one of claims 1-4, wherein the composition comprises about 5 mg of the polypeptide.

7. The composition of any one of claims 1-4, wherein the composition comprises about 15 mg of the polypeptide.

8. The composition of any one of claims 1-4, wherein the composition comprises about 20 mg of the polypeptide.

9. The composition of any one of claims 1-4, wherein the composition comprises about 30 mg of the polypeptide

10. The composition of any one of the preceding claims, wherein the composition comprises about 60 mg to about 72 mg sucrose.
11. The composition of any one of the preceding claims, wherein the composition comprises about 66 mg sucrose.
12. The composition of any one of the preceding claims, wherein the composition comprises about 60 mg to about 72 mg mannitol.
13. The composition of any one of the preceding claims, wherein the composition comprises about 66 mg mannitol.
14. The composition of any one of the preceding claims, wherein the composition comprises about 4.0 mg to about 6.0 mg citrate anion.
15. The composition of any one of the preceding claims, wherein the composition comprises about 5.0 mg citrate anion.
16. The composition of any one of the preceding claims, wherein the composition comprises citric acid and sodium citrate tribasic dihydrate in a mass ratio of citric acid:sodium citrate tribasic dihydrate of between about 1:10 to about 1:2.
17. The composition of any one of the preceding claims, wherein the composition comprises citric acid and sodium citrate tribasic dihydrate in a mass ratio of citric acid:sodium citrate tribasic dihydrate of about 1:9.
18. The composition of any one of the preceding claims, wherein the composition comprises citric acid and sodium citrate tribasic dihydrate in a mass ratio of citric acid:sodium citrate tribasic dihydrate of about 1:2.
19. The composition of any one of the preceding claims, wherein the emulsifier comprises polysorbate 20.

20. The composition of claim 19, wherein the composition comprises about 0.10 mg to about 0.12 mg polysorbate 20.

21. The composition of claim 19, wherein the composition comprises about 0.11 mg polysorbate 20.

22. The composition of any one of the preceding claims, wherein the composition is a lyophilized cake.

23. The composition of claim 22, wherein dissolution of the lyophilized cake in water results in an aqueous solution with a pH of about 5.5 to about 6.5.

24. The composition of claim 23, wherein dissolution of the lyophilized cake in water results in an aqueous solution with a pH of about 6.1.

25. The composition of any one of claims 22-24, wherein dissolution of the lyophilized cake in water results in an aqueous solution with an isotonic osmolality.

26. The composition of any one of claims 22-25, wherein dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 240 to about 340 mOsm/kg.

27. The composition of any one of claims 22-26, wherein dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 280 to about 320 mOsm/kg.

28. The composition of claim 26, wherein dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 285 mOsm/kg.

29. The composition of claim 26, wherein dissolution of the lyophilized cake in water results in an aqueous solution with an osmolality of about 300 mOsm/kg.

30. The composition of any one of claims 1-29, wherein the composition is an aqueous solution.
31. The composition of claim 30, wherein the composition comprises about 0.5 mg/mL to about 30 mg/mL of the polypeptide.
32. The composition of claim 31, wherein the composition comprises about 1 mg/mL of the polypeptide.
33. The composition of claim 32, wherein the composition is a 1.1 ml aqueous solution comprising about 1.1 mg of the polypeptide.
34. The composition of claim 31, wherein the composition comprises about 5 mg/mL of the polypeptide.
35. The composition of claim 34, wherein the composition is a 1.1 ml aqueous solution comprising about 15 mg of the polypeptide.
36. The composition of claim 31, wherein the composition comprises about 20 mg/mL of the polypeptide.
37. The composition of claim 31, wherein the composition comprises about 30 mg/mL of the polypeptide.
38. The composition of any one of claims 30-37, wherein the composition comprises about 25 mg/mL to about 35 mg/mL sucrose.
39. The composition of claim 38, wherein the composition comprises about 30 mg/mL sucrose.
40. The composition of any one of claims 30-37, wherein the composition comprises about 25 mg/mL to about 35 mg/mL mannitol.

41. The composition of claim 40, wherein the composition comprises about 30 mg/mL mannitol.

42. The composition of any one of claims 30-41, wherein the composition comprises about 10 mM to about 20 mM citrate buffer.

43. The composition of claim 42, wherein the composition comprises about 12 mM citrate buffer.

44. The composition of any one of claims 30-43, wherein the citrate buffer is formed by the combination of 2.03 mg/mL sodium citrate tribasic dihydrate and 0.97 mg/mL citric acid monohydrate in the aqueous solution.

45. The composition of any one of claims 30-43, wherein the citrate buffer is formed by the combination of 2.91 mg/mL sodium citrate tribasic dihydrate and 0.34 mg/mL citric acid monohydrate in the aqueous solution.

46. The composition of any one of claims 30-43, wherein the citrate buffer is formed by the combination of 2.96 mg/mL sodium citrate tribasic dihydrate and 0.30 mg/mL citric acid monohydrate in the aqueous solution.

47. The composition of any one of claims 30-46, wherein the composition comprises about 0.09 mg/mL to about 0.11 mg/mL polysorbate 20.

48. The composition of claim 47, wherein the composition comprises about 0.1 mg/mL polysorbate 20.

49. The composition of any one of claims 30-48, wherein the pH of the composition is about 5.5 to about 6.5.

50. The composition of claim 49, wherein the pH of the composition is about 6.1.

51. The composition of any one of claims 30-50, wherein the osmolality of the composition is about 240 to about 340 mOsm/kg.
52. The composition of any one of claims 30-50, wherein the osmolality of the composition is about 280 to about 320 mOsm/kg.
53. The composition of claim 51, wherein the osmolality of the composition is about 285 mOsm/kg.
54. The composition of claim 51, wherein the osmolality of the composition is about 300 mOsm/kg.
55. The composition of any one of claims 30-54, wherein the aqueous solution comprises about 0.03 mg/mL of the polypeptide to about 0.2 mg/mL of the polypeptide.
56. A lyophilized composition made by lyophilizing the composition of any one of claims 30-55.
57. The composition of any one of the preceding claims, wherein the composition is a single unit dose of the polypeptide.
58. An article of manufacture comprising the composition of any one of the preceding claims.
59. The article of claim 58, which is a glass vial.
60. A method of making a lyophilized composition, the method comprising lyophilizing the aqueous solution of any one of claims 30-55.
61. A method of making an aqueous composition, the method comprising dissolving the composition of claim 22 or 56 in an aqueous solvent.

62. The method of claim 61, wherein the pH of the aqueous composition is adjusted to about 6.1.

63. The method claim 61 or 62, wherein the pH of the aqueous composition is adjusted to about 6.1 with a base.

64. The method of claim 63, wherein the base is sodium hydroxide.

65. The method of any one of claims 61-64, wherein the aqueous composition is further diluted with an aqueous solution comprising about 1% (w/w) of a surfactant.

66. The method of claim 65, wherein the surfactant is polysorbate 20.

67. The method of claim 65, wherein the aqueous solution further comprises about 0.1% (w/w) citric acid monohydrate, 0.2% (w/w) sodium citrate tribasic dihydrate, and 98.7% (w/w) water for injection.

68. A method of activating natural killer cells (NK) cells in a subject, the method comprising administering to the subject an effective amount of the composition of any one of claims 30-55.

69. A method of treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of the composition of any one of claims 30-55.

70. The method claim 69, wherein the cancer is renal cell carcinoma, melanoma, ovarian cancer, or lung cancer.

71. The method claim 69 or 70, wherein the cancer comprises a refractory solid tumor.

72. The method of any one of claims 69-71, wherein the composition is administered subcutaneously.

73. The method of any one of claims 69-72, wherein the composition is administered subcutaneously at a dose of about 1 mg to about 15 mg.

74. The method of any one of claims 69-73, wherein the composition is administered subcutaneously at a dose of about 1 mg to about 15 mg once a week (Q1W), once every two weeks (Q2W), or once every three weeks (Q3W).

75. The method of any one of claims 69-73, wherein the melanoma is one or both of mucosal melanoma or advanced cutaneous melanoma.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US21/31616

A. CLASSIFICATION OF SUBJECT MATTER

IPC - A61K 38/20; A61P 35/00; C07K 16/28 (2021.01)

CPC - A61K 38/20; A61P 35/00; C07K 16/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2016/0175458 A1 (ALKERMES, INC.) 23 June 2016; paragraphs [0155], [0160], [0183], [0187]	1-4
Y	US 2019/0151448 A1 (AMGEN INC.) 23 May 2019; paragraph [0026], [0029], [0031], [0032], [0308], [0319], [0370]	1-4
P,Y	WO 2021/074689 A1 (ALKERMES PHARMA IRELAND LIMITED) 22 April 2021; entire document	1-4

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

27 August 2021 (27.08.2021)

Date of mailing of the international search report

SEP 27 2021

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450
Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US21/31616

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. forming part of the international application as filed:
 - in the form of an Annex C/ST.25 text file.
 - on paper or in the form of an image file.
 - b. furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. furnished subsequent to the international filing date for the purposes of international search only:
 - in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).
 - on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).
2. In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US21/31616

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 5-75
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.