

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 November 2004 (11.11.2004)

PCT

(10) International Publication Number
WO 2004/096180 A1

- (51) International Patent Classification⁷: **A61K 9/14**,
9/51, 31/496 (US). **PAPADOPOULOS, Pavlos** [GR/US]; 715 Creek
Bend Lane, Antioch, IL 60002 (US).
- (21) International Application Number: PCT/US2004/013268 (74) Agents: **MAYO, Michael, C.** et al.; Baxter International
Inc., One Baxter Parkway, Deerfield, IL 60015 (US).
- (22) International Filing Date: 29 April 2004 (29.04.2004) (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/466,354 29 April 2003 (29.04.2003) US
- (71) Applicant (for all designated States except US): **BAXTER
INTERNATIONAL INC.** [US/US]; One Baxter Parkway,
Deerfield, IL 60015 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **RABINOW, Bar-
rett, E.** [US/US]; 9119 Tripp Avenue, Skokie, IL 60076
(US). **WHITE, Randy** [US/US]; 3810 Tecoma Drive,
Crystal Lake, IL 60012-1888 (US). **SUN, Chong-Son**
[US/US]; 1051 Highland Avenue, Lake Forest, IL 60045
(US). **WONG, Joseph, Chung, Tak** [US/US]; 36141
N. Brindlewood Avenue, Gurnee, IL 60031 (US). **KIPP,
James, E.** [US/US]; 609 Hermosa Avenue, Wauconda, IL
60084 (US). **DOTY, Mark, J.** [US/US]; 610 Kenilworth
Street, Grayslake, IL 60030 (US). **REBBECK, Christine,
L.** [US/US]; 525 Eagle Ridge Lane, Algonquin, IL 60102
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FORMULATION TO RENDER AN ANTIMICROBIAL DRUG POTENT AGAINST ORGANISMS NORMALLY CONSIDERED TO BE RESISTANT TO THE DRUG

(57) Abstract: The present invention relates to compositions of submicron- to micron-size particles of antimicrobial agents. More particularly the invention relates to a composition of an antimicrobial agent that renders the agent potent against organisms normally considered to be resistant to the agent. The composition comprises an aqueous suspension of submicron-to-micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives. The particles have a volume-weighted mean particle size of less than 5 µm as measured by laser diffractometry.

WO 2004/096180 A1

**FORMULATION TO RENDER AN ANTIMICROBIAL DRUG POTENT AGAINST
ORGANISMS NORMALLY CONSIDERED
TO BE RESISTANT TO THE DRUG**

5 CROSS REFERENCE TO RELATED APPLICATIONS:

This application claims priority from provisional application Ser. No. 60/466,354, filed on April 29, 2003.

BACKGROUND OF THE INVENTION:

Technical Field

10 The present invention relates to compositions of antimicrobial agents. More particularly the invention relates to formulations of an antimicrobial agent which render the drug potent against organisms normally considered to be resistant to the agent.

Background of the Invention

Based upon *in vitro* microbicidal sensitivity tests, the level of an antimicrobial drug
15 considered effective against a particular organism may be determined. This is referred to as the MIC (minimum inhibitory concentration) of the drug. On the other hand, safety studies will determine the amount of drug that can be safely given to a patient or test animal. This maximal amount of drug that can be dosed will determine the maximal biological exposure to the host animal, normally measured by the area under the curve (AUC) of the plot of drug concentration
20 vs. time, the peak height of the plot of drug concentration vs. time, tissue levels vs. time, etc. The instantaneous tissue or plasma level of the *in vivo* experiment can be compared with the MIC value to determine relative efficacy of the attainable drug levels in the biological fluids. The actual comparison must be corrected for plasma protein binding, inasmuch as only the free drug level is the important parameter because it is in this state that the drug is freely diffusible to
25 cross biological membranes.

As a result of such analysis, clinical literature has been established specifying what drugs can be used generally for certain strains of organisms, or more precisely, for certain strains of organisms with MIC values below certain levels. As an example, the antifungal agent itraconazole is not considered effective for strains of *Candida albicans* with MIC>8 for this drug
30 (e.g., for *C. albicans* strain c43 (ATCC number 201794), MIC₈₀ = 16 µg/ml for SPORANOX®

- 2 -

itraconazole). These strains of *Candida albicans* are considered to be resistant to itraconazole. This presupposes the standard dosing level of this drug that can be administered.

However, if a method were available to substantially increase the amount of the antimicrobial drug (e.g., itraconazole) that could be administered, than it might be possible to
5 treat infections hithertofore considered untreatable by this agent. Such a method is available through formulation of the drug as a nanosuspension. Submicron sized drug crystals stabilized by a surfactant coating have been found, in some cases, not to dissolve immediately upon injection into the blood stream. Instead, they are captured by fixed macrophages of the spleen and liver. From this sanctuary, the drug will be slowly released over a prolonged period of days.
10 This is in contrast to conventionally solubilized drugs, which when injected, decrease in blood concentration at a much faster rate.

An example of an antimicrobial agent which is conventionally formulated to increase the solubility of the drug is the triazole antifungal agent itraconazole (FIG. 2). Itraconazole is effective against systemic mycoses, particularly aspergillosis and candidiasis. New oral and
15 intravenous preparations of itraconazole have been prepared in order to overcome bioavailability problems associated with a lack of solubility. For example, the bioavailability of itraconazole is increased when it is formulated in hydroxypropyl-beta-cyclodextrin, a carrier oligosaccharide that forms an inclusion complex with the drug, thereby increasing its aqueous solubility. The commercial preparation is known by the tradename SPORANOX[®] Injection and was originated
20 by JANSSEN PHARMACEUTICAL PRODUCTS, L.P. The drug is currently manufactured by Abbott Labs and distributed by Ortho Biotech, Inc.

Intravenous itraconazole may be useful in selected clinical situations. Examples are achlorhydria in AIDS patients, an inability to effectively absorb oral medications due to
25 concurrent treatments with other drugs, or in critical-care patients who cannot take oral medications. The current commercial product, SPORANOX[®] Injection, is made available in 25 mL glass vials that contain 250 mg of itraconazole, with 10 g of hydroxypropyl-beta-cyclodextrin (referenced as "HPBCD"). These vials are diluted prior to use in 50 mL of 0.9% saline. The resulting cyclodextrin concentration exceeds 10% (w/v) in the reconstituted product. Although HPBCD has been traditionally regarded as safe for injection, high concentrations, such as 10%,
30 have been reported in animal models to induce significant changes to endothelial tissues (Duncker G.; Reichelt J., Effects of the pharmaceutical cosolvent hydroxypropyl-beta-

cyclodextrin on porcine corneal endothelium. Graefe's Archive for Clinical and Experimental Ophthalmology (Germany) 1998, 236/5, 380-389).

Other excipients are often used to formulate poorly water-soluble drugs for intravenous injection. For example, paclitaxel (Taxol®, produced by Bristol-Myers Squibb) contains 52.7% (w/v) of Cremophor® EL (polyoxyethylated castor oil) and 49.7% (v/v) dehydrated alcohol, USP. Administration of Cremophor® EL can lead to undesired hypersensitivity reactions (Volcheck, G.W., Van Dellen, R.G. Anaphylaxis to intravenous cyclosporine and tolerance to oral cyclosporine: case report and review. *Annals of Allergy, Asthma, and Immunology*, 1998, 80, 159-163; Singla A.K.; Garg A.; Aggarwal D., Paclitaxel and its formulations. *International Journal of Pharmaceutics*, 2002, 235/ 1-2, 179-192).

The present invention discloses a composition which renders antimicrobial drugs more effective on the basis of their physical and biological properties than in their unformulated state or in their existing formulations. The approach used is to formulate the antimicrobial agents as nanosuspensions. This permits using of the improved formulation to treat microbes conventionally thought to be resistant to the unformulated drug. Conventional formulation approaches attempt to enhance solubility or bioavailability only. Such methods include pH change, modification of the salt form, use of organic modifiers, or cyclodextrin. The approach disclosed in the present invention involves altering the pharmacokinetic characteristic of the drug, permitting far greater dosing, resulting in improved efficacy over and above what can be accomplished by improving solubility and bioavailability only. Acute toxicity tests have demonstrated that much more drug, when formulated as a nanosuspension, can be administered to animals. More of the drug is therefore available at the target organ to exert efficacy.

SUMMARY OF THE INVENTION:

The present invention relates to a composition of an aqueous suspension of submicron- to micron-size particles of an antimicrobial agent that renders the agent potent against organisms normally considered to be resistant to the agent. The composition includes an aqueous suspension of submicron- to micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives. The particles have a volume-weighted mean particle size of less than 5 µm as measured by light scattering (HORIBA) or by microscopic measurements. More preferably the particles should be less than about 1

micron and most preferably from about 150 nm to about 1 micron or any range or combination of ranges therein.

The present invention is suitable for pharmaceutical use.

In an embodiment of the invention, the antimicrobial agent is an antifungal agent. In a preferred embodiment, the antifungal agent is a triazole antifungal agent. In yet another embodiment of the invention, the triazole antifungal agent is selected from itraconazole, ketoconazole, miconazole, fluconazole, ravuconazole, voriconazole, saperconazole, eberconazole, genaconazole, clotrimazole, econazole, oxiconazole, sulconazole, terconazole, tioconazole, and posaconazole. In a preferred embodiment of the invention, the antifungal agent is itraconazole.

Suitable surfactants for coating the particles in the present invention can be selected from ionic surfactants, nonionic surfactants, biologically derived surfactants, or amino acids and their derivatives.

In a further preferred embodiment, the composition of the present invention is prepared by a microprecipitation method which includes the steps of: (i) dissolving in the antifungal agent in a first water-miscible first solvent to form a solution; (ii) mixing the solution with a second solvent which is aqueous to define a pre-suspension; and (iii) adding energy to the pre-suspension to form particles having an average effective particle size of less than 5 μm ; more preferably less than about 1 micron, and most preferably from about 150 nm to about 1 micron or any range or combination of ranges therein, wherein the solubility of the antifungal agent is greater in the first solvent than in the second solvent, and the first solvent or the second solvent comprising one or more surfactants selected from the group consisting of: nonionic surfactants, ionic surfactants, biologically derived surfactants, and amino acids and their derivatives.

The present invention also relates to a method of rendering an antimicrobial agent potent against organisms normally considered to be resistant to the agent by formulating the agent as an aqueous suspension of submicron- to micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives.

The present invention further relates to a method of treating infection of a subject by organisms normally considered to be resistant to an antimicrobial agent by administering the agent to the subject formulated as an aqueous suspension of submicron- to micron-size particles

containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives.

5 These and other aspects and attributes of the present invention will be discussed with reference to the following drawings and accompanying specification.

BRIEF DESCRIPTION OF THE DRAWINGS:

FIG. 1 is the general molecular structure of a triazole antifungal agent;

FIG. 2 is the molecular structure of itraconazole;

10 FIG. 3 is a schematic diagram of Method A of the microprecipitation process used in the present invention to prepare the suspension;

FIG. 4 is a schematic diagram of Method B of the microprecipitation process used in the present invention to prepare the suspension;

15 FIG. 5 is a graph comparing the pharmacokinetics of SPORANOX® with Formulation 1 suspension of itraconazole of the present invention, wherein ITC = plasma concentration of itraconazole measured after bolus injection of Formulation 1 (80 mg/kg), ITC-OH = plasma concentration of primary metabolite, hydroxyitraconazole, measured after bolus injection of Formulation 1 (80 mg/kg), Total = combined concentration of itraconazole and hydroxyitraconazole (ITC + ITC-OH) measured after bolus injection of Formulation 1 (80 mg/kg), Sporanox-ITC = plasma concentration of itraconazole measured after bolus injection of 20 mg/kg Sporanox IV, Sporanox-ITC-OH = plasma concentration of primary metabolite, hydroxyitraconazole, measured after bolus injection of 20 mg/kg Sporanox IV, Sporanox - Total = combined concentration of itraconazole and hydroxyitraconazole (ITC + ITC-OH) measured after bolus injection of 20 mg/kg Sporanox IV;

25 FIG. 6 is a graph comparing the drug level for the rapidly dissolving formulation, Form A, and the slow dissolving (macrophage targeting) formulation, Form B, as determined in an in vitro dissolution experiment; the drug level for Form A is much higher than that attained by Form B;

FIG. 7 is a graph showing the comparison of results for body weight over time for immuno-suppressed rats treated with SPORANOX® Injection and Formulations 14288-1 and 14288-B;

FIG. 8 is a graph of kidney drug level vs. dose showing that the greater dosing that could be administered permitted greater drug levels to be manifested in the target organs, in this case, the kidney;

FIG. 9 is a graph of fungal counts vs. kidney drug level (N = nanosuspension; S = Sporanox IV solution) showing that the greater drug levels in the target organ (the kidney) led to a greater kill of the infectious organisms; and

FIG. 10 is a graph showing the mortality/moribundity profile after daily or every other day dosing with antifungal drugs for 10 days in rats systemically infected with itraconazole resistant *C. albicans*.

10 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

While this invention is susceptible of embodiment in many different forms, there are shown in the drawing, and will be described herein in detail, specific embodiments thereof with the understanding that the present disclosure is to be considered as an exemplification of the principles of the invention and is not intended to limit the invention to the specific embodiments illustrated.

The present invention relates to a composition of an antimicrobial agent that renders the agent potent against organisms normally considered to be resistant to the agent. The composition comprises an aqueous suspension of submicron- to micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives. The composition disclosed in the present invention involves altering the pharmacokinetic characteristic of the drug, permitting far greater dosing, resulting in improved efficacy over and above what can be accomplished by improving solubility and bioavailability only. Submicron sized drug crystals stabilized by a surfactant coating have been found, in some cases, not to dissolve immediately upon injection into the blood stream. Instead, they are captured by fixed macrophages of the spleen and liver. From this sanctuary, the drug can be slowly released over a prolonged period of days. Acute toxicity tests have demonstrated that much more drug, when formulated as a nanosuspension, can be administered to animals or human beings. More of the drug is therefore available at the target organ to exert efficacy.

The particles in the present invention have a volume-weighted mean particle size of less than 5 μm as measured by light scattering (HORIBA) or by microscopic measurements. More

preferably the particles should be less than about 1 micron and most preferably from about 150 nm to about 1 micron or any range or combination of ranges therein. The composition can be administered to a subject to treat infection by organisms normally considered to be resistant to the agent.

5 The antimicrobial agent is preferably a poorly water soluble organic compound. What is meant by "poorly water soluble" is that the water solubility of the compound is less than 10 mg/ml, and preferably, less than 1 mg/ml. A preferred class of antimicrobial agent is an antifungal agent. A preferred antifungal agent is the triazole antifungal agents having a general molecular structure as shown in FIG. 1. Examples of triazole antifungal agents include, but are
10 not limited to: itraconazole, ketoconazole, miconazole, fluconazole, ravuconazole, voriconazole, saperconazole, eberconazole, genaconazole, clotrimazole, econazole, oxiconazole, sulconazole, terconazole, tioconazole, and posaconazole. A preferred antifungal agent for the present invention is itraconazole. The molecular structure of itraconazole is shown in FIG. 2.

The present invention is suitable for pharmaceutical use. The compositions can be
15 administered by various routes, including but not limited to, intravenous, intracerebral, intrathecal, intralymphatic, pulmonary, intraarticular, and intraperitoneal. In an embodiment of the present invention, the aqueous medium of the composition is removed to form dry particles. The method to remove the aqueous medium can be any method known in the art. One example is evaporation. Another example is freeze drying or lyophilization. The dry particles may then be
20 formulated into any acceptable physical form including, but is not limited to, solutions, tablets, capsules, suspensions, creams, lotions, emulsions, aerosols, powders, incorporation into reservoir or matrix devices for sustained release (such as implants or transdermal patches), and the like.

If the particles do not have to be taken up by the macrophages, the particles can be larger than 5 μm (e.g., less than 50 μm , or less than 7 μm) or less than 150 nm (e.g., less than 100 μm).
25 These particles can be administered by various routes, including but not limited to parenteral, oral, buccal, periodontal, rectal, nasal, pulmonary, transdermal, or topical. Modes of parenteral administration include intravenous, intra arterial, intrathecal, intraperitoneal, intraocular, intra articular, intrathecal, intracerebral, intramuscular, subcutaneous, and the like.

The aqueous suspension of the present invention may also be frozen to improve stability
30 upon storage. Freezing of an aqueous suspension to improve stability is disclosed in the

commonly assigned and co-pending U.S. Patent Application Serial No. 60/347,548, which is incorporated herein by reference and made a part hereof.

In an embodiment of the present invention, the antimicrobial agent is present in an amount preferably from about 0.01% to about 50% weight to volume (w/v), more preferably from about 0.05% to about 30% w/v, and most preferably from about 0.1% to about 20% w/v.

Suitable surfactants for coating the particles in the present invention can be selected from ionic surfactants, nonionic surfactants, biologically derived surfactants or amino acids and their derivatives. Ionic surfactants can be anionic, cationic, or zwitterionic.

Suitable anionic surfactants include but are not limited to: alkyl sulfonates, alkyl phosphates, alkyl phosphonates, potassium laurate, triethanolamine stearate, sodium lauryl sulfate, sodium dodecylsulfate, alkyl polyoxyethylene sulfates, sodium alginate, dioctyl sodium sulfosuccinate, phosphatidylglycerol, phosphatidylinosine, phosphatidylinositol, diphosphatidylglycerol, phosphatidylserine, phosphatidic acid and their salts, sodium carboxymethylcellulose, cholic acid and other bile acids (e.g., cholic acid, deoxycholic acid, glycocholic acid, taurocholic acid, glycodeoxycholic acid) and salts thereof (e.g., sodium deoxycholate, etc.). As anionic surfactants, phospholipids may be used. Suitable phospholipids include, for example, phosphatidylserine, phosphatidylinositol, diphosphatidylglycerol, phosphatidylglycerol, or phosphatidic acid and its salts.

Zwitterionic surfactants are electrically neutral but possess local positive and negative charges within the same molecule. Suitable zwitterionic surfactants include but are not limited to zwitterionic phospholipids. Suitable phospholipids include phosphatidylcholine, phosphatidylethanolamine, diacyl-glycero-phosphoethanolamine (such as dimyristoyl-glycero-phosphoethanolamine (DMPE), dipalmitoyl-glycero-phosphoethanolamine (DPPE), distearoyl-glycero-phosphoethanolamine (DSPE), and dioleoyl-glycero-phosphoethanolamine (DOPE)). Mixtures of phospholipids that include anionic and zwitterionic phospholipids may be employed in this invention. Such mixtures include but are not limited to lysophospholipids, egg or soybean phospholipid or any combination thereof. The phospholipid, whether anionic, zwitterionic or a mixture of phospholipids, may be salted or desalted, hydrogenated or partially hydrogenated or natural semisynthetic or synthetic. The phospholipid may also be conjugated with a water-soluble or hydrophilic polymer to specifically target the delivery to macrophages in the present invention. However, conjugated phospholipids may be used to target other cells or tissue in other

applications. A preferred polymer is polyethylene glycol (PEG), which is also known as the monomethoxy polyethyleneglycol (mPEG). The molecule weights of the PEG can vary, for example, from 200 to 50,000. Some commonly used PEG's that are commercially available include PEG 350, PEG 550, PEG 750, PEG 1000, PEG 2000, PEG 3000, and PEG 5000. The phospholipid or the PEG-phospholipid conjugate may also incorporate a functional group which can covalently attach to a ligand including but not limited to proteins, peptides, carbohydrates, glycoproteins, antibodies, or pharmaceutically active agents. These functional groups may conjugate with the ligands through, for example, amide bond formation, disulfide or thioether formation, or biotin/streptavidin binding. Examples of the ligand-binding functional groups include but are not limited to hexanoylamine, dodecanylamine, 1,12-dodecanedicarboxylate, thioethanol, 4-(p-maleimidophenyl)butyramide (MPB), 4-(p-maleimidomethyl)cyclohexanecarboxamide (MCC), 3-(2-pyridyldithio)propionate (PDP), succinate, glutarate, dodecanoate, and biotin.

Suitable cationic surfactants include but are not limited to quaternary ammonium compounds, such as benzalkonium chloride, cetyltrimethylammonium bromide, lauryldimethylbenzylammonium chloride, acyl carnitine hydrochlorides, or alkyl pyridinium halides, or long-chain alkyl amines such as, for example, n-octylamine and oleylamine.

Suitable nonionic surfactants include: glyceryl esters, polyoxyethylene fatty alcohol ethers (Macrogol and Brij), polyoxyethylene sorbitan fatty acid esters (Polysorbates), polyoxyethylene fatty acid esters (Myrj), sorbitan esters (Span), glycerol monostearate, polyethylene glycols, polypropylene glycols, cetyl alcohol, cetostearyl alcohol, stearyl alcohol, aryl alkyl polyether alcohols, polyoxyethylene-polyoxypropylene copolymers (poloxamers), poloxamines, methylcellulose, hydroxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, noncrystalline cellulose, polysaccharides including starch and starch derivatives such as hydroxyethylstarch (HES), polyvinyl alcohol, and polyvinylpyrrolidone. In a preferred form of the invention, the nonionic surfactant is a polyoxyethylene and polyoxypropylene copolymer and preferably a block copolymer of propylene glycol and ethylene glycol. Such polymers are sold under the tradename POLOXAMER also sometimes referred to as PLURONIC®, and sold by several suppliers including Spectrum Chemical and Ruger. Among polyoxyethylene fatty acid esters is included

those having short alkyl chains. One example of such a surfactant is SOLUTOL® HS 15, polyethylene-660-hydroxystearate, manufactured by BASF Aktiengesellschaft.

Surface-active biological molecules include such molecules as albumin, casein, hirudin or other appropriate proteins. Polysaccharide biologics are also included, and consist of but are not limited to, starches, heparin and chitosans. Other suitable surfactants include any amino acids
5 such as leucine, alanine, valine, isoleucine, lysine, aspartic acid, glutamic acid, methionine, phenylalanine, or any derivatives of these amino acids such as, for example, amide or ester derivatives and polypeptides formed from these amino acids.

A preferred ionic surfactant is a bile salt, and a preferred bile salt is deoxycholate. A preferred nonionic surfactant is a polyalkoxyether, and a preferred polyalkoxyether is Poloxamer
10 188. Another preferred nonionic surfactant is Solutol HS 15 (polyethylene-660-hydroxystearate). Still yet another preferred nonionic surfactant is hydroxyethylstarch. A preferred biologically derived surfactant is albumin.

In another embodiment of the present invention, the surfactants are present in an amount
15 of preferably from about 0.001% to 5% w/v, more preferably from about 0.005% to about 5% w/v, and most preferably from about 0.01% to 5% w/v.

In a preferred embodiment of the present invention, the particles are suspended in an aqueous medium further including a pH adjusting agent. Suitable pH adjusting agents include, but are not limited to, hydrochloric acid, sulfuric acid, phosphoric acid, monocarboxylic acids
20 (such as, for example, acetic acid and lactic acid), dicarboxylic acids (such as, for example, succinic acid), tricarboxylic acids (such as, for example, citric acid), THAM (tris(hydroxymethyl)aminomethane), meglumine (N-methylglucosamine), sodium hydroxide, and amino acids such as glycine, arginine, lysine, alanine, histidine and leucine. The aqueous medium may additionally include an osmotic pressure adjusting agent, such as but not limited to
25 glycerin, a monosaccharide such as dextrose, a disaccharide such as sucrose, a trisaccharide such as raffinose, and sugar alcohols such as mannitol, xylitol and sorbitol.

In a preferred embodiment of the present invention, the composition comprises an aqueous suspension of particles of itraconazole present at 0.01 to 50% w/v, the particles are coated with 0.001 to 5% w/v of a bile salt (e.g., deoxycholate) and 0.001 to 5% w/v
30 polyalkoxyether (for example, Poloxamer 188), and glycerin added to adjust osmotic pressure of the formulation.

In another preferred embodiment of the present invention, the composition comprises an aqueous suspension of particles of itraconazole present at about 0.01 to 50% w/v, the particles coated with about 0.001 to 5% w/v of a bile salt (for example, deoxycholate) and 0.001 to 5% polyethylene-660-hydroxystearate w/v, and glycerin added to adjust osmotic pressure of the formulation.

In another preferred embodiment of the present invention, the composition comprises an aqueous suspension of itraconazole present at about 0.01 to 50% w/v, the particles are coated with about 0.001 to 5% of polyethylene-660-hydroxystearate w/v, and glycerin added to adjust osmotic pressure of the formulation.

In still yet another preferred embodiment of the present invention, the composition comprises an aqueous suspension of itraconazole present at 0.01 to 50% w/v, the particles are coated with about 0.001 to 5% albumin w/v.

The method for preparing the suspension in the present invention is disclosed in commonly assigned and co-pending U.S. Patent Applications Serial Nos. 60/258,160; 09/874,799; 09/874,637; 09/874,499; 09/964,273; 10/035,821, 60/347,548; 10/021,692; 10/183,035; 10/213,352; 10/246,802; 10/270,268; 10/270,267, and 10/390,333; which are incorporated herein by reference and made a part hereof. A general procedure for preparing the suspension useful in the practice of this invention follows.

The processes can be separated into three general categories. Each of the categories of processes share the steps of: (1) dissolving an antifungal agent in a water miscible first organic solvent to create a first solution; (2) mixing the first solution with a second solvent of water to precipitate the antifungal agent to create a pre-suspension; and (3) adding energy to the presuspension in the form of high-shear mixing or heat to provide a stable form of the antifungal agent having the desired size ranges defined above.

The three categories of processes are distinguished based upon the physical properties of the antifungal agent as determined through x-ray diffraction studies, differential scanning calorimetry (DSC) studies or other suitable study conducted prior to the energy-addition step and after the energy-addition step. In the first process category, prior to the energy-addition step the antifungal agent in the presuspension takes an amorphous form, a semi-crystalline form or a supercooled liquid form and has an average effective particle size. After the energy-addition

step, the antifungal agent is in a crystalline form having an average effective particle size essentially the same as that of the presuspension (i.e., from less than about 50 μm).

In the second process category, prior to the energy-addition step the antifungal agent is in a crystalline form and has an average effective particle size. After the energy-addition step, the
5 antifungal agent is in a crystalline form having essentially the same average effective particle size as prior to the energy-addition step but the crystals after the energy-addition step are less likely to aggregate.

The lower tendency of the organic compound to aggregate is observed by laser dynamic light scattering and light microscopy.

10 In the third process category, prior to the energy-addition step the antifungal agent is in a crystalline form that is friable and has an average effective particle size. What is meant by the term "friable" is that the particles are fragile and are more easily broken down into smaller particles. After the energy-addition step the organic compound is in a crystalline form having an average effective particle size smaller than the crystals of the pre-suspension. By taking the steps
15 necessary to place the organic compound in a crystalline form that is friable, the subsequent energy-addition step can be carried out more quickly and efficiently when compared to an organic compound in a less friable crystalline morphology.

The energy-addition step can be carried out in any fashion wherein the pre-suspension is exposed to cavitation, shearing or impact forces. In one preferred form of the invention, the
20 energy-addition step is an annealing step. Annealing is defined in this invention as the process of converting matter that is thermodynamically unstable into a more stable form by single or repeated application of energy (direct heat or mechanical stress), followed by thermal relaxation. This lowering of energy may be achieved by conversion of the solid form from a less ordered to a more ordered lattice structure. Alternatively, this stabilization may occur by a reordering of the
25 surfactant molecules at the solid-liquid interface.

These three process categories will be discussed separately below. It should be understood, however, that the process conditions such as choice of surfactants or combination of surfactants, amount of surfactant used, temperature of reaction, rate of mixing of solutions, rate of precipitation and the like can be selected to allow for any drug to be processed under any one
30 of the categories discussed next.

The first process category, as well as the second and third process categories, can be further divided into two subcategories, Method A, and B shown diagrammatically in FIG. 3 and FIG. 4, respectively.

The first solvent according to the present invention is a solvent or mixture of solvents in which the organic compound of interest is relatively soluble and which is miscible with the second solvent. Such solvents include, but are not limited to water-miscible protic compounds, in which a hydrogen atom in the molecule is bound to an electronegative atom such as oxygen, nitrogen, or other Group VA, VIA and VII A in the Periodic Table of elements. Examples of such solvents include, but are not limited to, alcohols, amines (primary or secondary), oximes, hydroxamic acids, carboxylic acids, sulfonic acids, phosphonic acids, phosphoric acids, amides and ureas.

Other examples of the first solvent also include aprotic organic solvents. Some of these aprotic solvents can form hydrogen bonds with water, but can only act as proton acceptors because they lack effective proton donating groups. One class of aprotic solvents is a dipolar aprotic solvent, as defined by the International Union of Pure and Applied Chemistry (IUPAC Compendium of Chemical Terminology, 2nd Ed., 1997):

A solvent with a comparatively high relative permittivity (or dielectric constant), greater than ca. 15, and a sizable permanent dipole moment, that cannot donate suitably labile hydrogen atoms to form strong hydrogen bonds, e.g. dimethyl sulfoxide.

Dipolar aprotic solvents can be selected from the group consisting of: amides (fully substituted, with nitrogen lacking attached hydrogen atoms), ureas (fully substituted, with no hydrogen atoms attached to nitrogen), ethers, cyclic ethers, nitriles, ketones, sulfones, sulfoxides, fully substituted phosphates, phosphonate esters, phosphoramides, nitro compounds, and the like. Dimethylsulfoxide (DMSO), N-methyl-2-pyrrolidinone (NMP), 2-pyrrolidinone, 1,3-dimethylimidazolidinone (DMI), dimethylacetamide (DMA), dimethylformamide (DMF), dioxane, acetone, tetrahydrofuran (THF), tetramethylenesulfone (sulfolane), acetonitrile, and hexamethylphosphoramide (HMPA), nitromethane, among others, are members of this class.

Solvents may also be chosen that are generally water-immiscible, but have sufficient water solubility at low volumes (less than 10%) to act as a water-miscible first solvent at these reduced volumes. Examples include aromatic hydrocarbons, alkenes, alkanes, and halogenated

aromatics, halogenated alkenes and halogenated alkanes. Aromatics include, but are not limited to, benzene (substituted or unsubstituted), and monocyclic or polycyclic arenes. Examples of substituted benzenes include, but are not limited to, xylenes (ortho, meta, or para), and toluene. Examples of alkanes include but are not limited to hexane, neopentane, heptane, isooctane, and cyclohexane. Examples of halogenated aromatics include, but are not restricted to, chlorobenzene, bromobenzene, and chlorotoluene. Examples of halogenated alkanes and alkenes include, but are not restricted to, trichloroethane, methylene chloride, ethylenedichloride (EDC), and the like.

Examples of the all of the above solvent classes include but are not limited to: N-methyl-2-pyrrolidinone (also called N-methyl-2-pyrrolidone), 2-pyrrolidinone (also called 2-pyrrolidone), 1,3-dimethyl-2-imidazolidinone (DMI), dimethylsulfoxide, dimethylacetamide, carboxylic acids (such as acetic acid and lactic acid), aliphatic alcohols (such as methanol, ethanol, isopropanol, 3-pentanol, and n-propanol), benzyl alcohol, glycerol, butylene glycol (butanediol), ethylene glycol, propylene glycol, mono- and diacylated monoglycerides (such as glyceryl caprylate), dimethyl isosorbide, acetone, dimethylsulfone, dimethylformamide, 1,4-dioxane, tetramethylenesulfone (sulfolane), acetonitrile, nitromethane, tetramethylurea, hexamethylphosphoramide (HMPA), tetrahydrofuran (THF), dioxane, diethylether, tert-butylmethyl ether (TBME), aromatic hydrocarbons, alkenes, alkanes, halogenated aromatics, halogenated alkenes, halogenated alkanes, xylene, toluene, benzene, substituted benzene, ethyl acetate, methyl acetate, butyl acetate, chlorobenzene, bromobenzene, chlorotoluene, trichloroethane, methylene chloride, ethylenedichloride (EDC), hexane, neopentane, heptane, isooctane, cyclohexane, polyethylene glycol (PEG, for example, PEG-4, PEG-8, PEG-9, PEG-12, PEG-14, PEG-16, PEG-120, PEG-75, PEG-150), polyethylene glycol esters (examples such as PEG-4 dilaurate, PEG-20 dilaurate, PEG-6 isostearate, PEG-8 palmitostearate, PEG-150 palmitostearate), polyethylene glycol sorbitans (such as PEG-20 sorbitan isostearate), polyethylene glycol monoalkyl ethers (examples such as PEG-3 dimethyl ether, PEG-4 dimethyl ether), polypropylene glycol (PPG), polypropylene alginate, PPG-10 butanediol, PPG-10 methyl glucose ether, PPG-20 methyl glucose ether, PPG-15 stearyl ether, propylene glycol dicaprylate/dicaprate, propylene glycol laurate, and glycofuroyl (tetrahydrofurfuryl alcohol polyethylene glycol ether). A preferred first solvent is N-methyl-2-pyrrolidinone. Another preferred first solvent is lactic acid.

The second solvent is an aqueous solvent. This aqueous solvent may be water by itself. This solvent may also contain buffers, salts, surfactant(s), water-soluble polymers, and combinations of these excipients.

Method A

5 In Method A (see FIG. 3), the antimicrobial agent is first dissolved in the first solvent to create a first solution. The antimicrobial agent can be added from about 0.01% (w/v) to about 50% (w/v) depending on the solubility of the antimicrobial agent in the first solvent. Heating of the concentrate from about 30°C to about 100°C may be necessary to ensure total dissolution of the antimicrobial agent in the first solvent.

10 A second aqueous solution is provided with one or more surfactants added thereto. The surfactants can be selected from an ionic surfactant, a nonionic surfactant or a biologically derived surfactant set forth above.

It may also be desirable to add a pH adjusting agent to the second solution such as sodium hydroxide, hydrochloric acid, tris buffer or citrate, acetate, lactate, meglumine, or the like. The
15 second solution should have a pH within the range of from about 3 to about 11.

In a preferred form of the invention, the method for preparing submicron sized particles of an antimicrobial agent includes the steps of adding the first solution to the second solution. The addition rate is dependent on the batch size, and precipitation kinetics for the antimicrobial agent. Typically, for a small-scale laboratory process (preparation of 1 liter), the addition rate is from
20 about 0.05 cc per minute to about 10 cc per minute. During the addition, the solutions should be under constant agitation. It has been observed using light microscopy that amorphous particles, semi-crystalline solids, or a supercooled liquid are formed to create a pre-suspension. The method further includes the step of subjecting the pre-suspension to an annealing step to convert the amorphous particles, supercooled liquid or semicrystalline solid to a crystalline more stable
25 solid state. The resulting particles will have an average effective particles size as measured by dynamic light scattering methods (e.g., photocorrelation spectroscopy, laser diffraction, low-angle laser light scattering (LALLS), medium-angle laser light scattering (MALLS), light obscuration methods (Coulter method, for example), rheology, or microscopy (light or electron) within the ranges set forth above).

30 The energy-addition step involves adding energy through sonication, homogenization, counter current flow homogenization (e.g., the Mini DeBEE 2000 homogenizer, available from

BEE Incorporated, NC, in which a jet of fluid is directed along a first path, and a structure is interposed in the first path to cause the fluid to be redirected in a controlled flow path along a new path to cause emulsification or mixing of the fluid), microfluidization, or other methods of providing impact, shear or cavitation forces. The sample may be cooled or heated during this stage. In one preferred form of the invention the annealing step is effected by homogenization. In another preferred form of the invention the annealing may be accomplished by ultrasonication. In yet another preferred form of the invention the annealing may be accomplished by use of an emulsification apparatus as described in U.S. Patent No. 5,720,551 which is incorporated herein by reference and made a part hereof.

Depending upon the rate of annealing, it may be desirable to adjust the temperature of the processed sample to within the range of from approximately -30°C to 100°C . Alternatively, in order to effect a desired phase change in the processed solid, it may also be necessary to adjust the temperature of the pre-suspension to a temperature within the range of from about -30°C to about 100°C during the annealing step.

15 Method B

Method B differs from Method A in the following respects. The first difference is a surfactant or combination of surfactants are added to the first solution. The surfactants may be selected from ionic surfactants, nonionic surfactants, or biologically derived as set forth above.

A drug suspension resulting from application of the processes described in this invention may be administered directly as an injectable solution, provided Water for Injection is used in formulation and an appropriate means for solution sterilization is applied. Sterilization may be accomplished by separate sterilization of the drug concentrate (drug, solvent, and optional surfactant) and the diluent medium (water, and optional buffers and surfactants) prior to mixing to form the pre-suspension. Sterilization methods include pre-filtration first through a 3.0 micron filter followed by filtration through a 0.45-micron particle filter, followed by steam or heat sterilization or sterile filtration through two redundant 0.2-micron membrane filters.

Optionally, a solvent-free suspension may be produced by solvent removal after precipitation. This can be accomplished by centrifugation, dialysis, diafiltration, force-field fractionation, high-pressure filtration or other separation techniques well known in the art. Complete removal of N-methyl-2-pyrrolidinone was typically carried out by one to three

successive centrifugation runs; after each centrifugation the supernatant was decanted and discarded. A fresh volume of the suspension vehicle without the organic solvent was added to the remaining solids and the mixture was dispersed by homogenization. It will be recognized by others skilled in the art that other high-shear mixing techniques could be applied in this reconstitution step.

Furthermore, any undesired excipients such as surfactants may be replaced by a more desirable excipient by use of the separation methods described in the above paragraph. The solvent and first excipient may be discarded with the supernatant after centrifugation or filtration.

A fresh volume of the suspension vehicle without the solvent and without the first excipient may then be added. Alternatively, a new surfactant may be added. For example, a suspension consisting of drug, N-methyl-2-pyrrolidinone (solvent), Poloxamer 188 (first excipient), sodium deoxycholate, glycerol and water may be replaced with phospholipids (new surfactant), glycerol and water after centrifugation and removal of the supernatant.

I. First Process Category

The methods of the first process category generally include the step of dissolving the antimicrobial agent in a water miscible first solvent followed by the step of mixing this solution with an aqueous solution to form a presuspension wherein the antimicrobial agent is in an amorphous form, a semicrystalline form or in a supercooled liquid form as determined by x-ray diffraction studies, DSC, light microscopy or other analytical techniques and has an average effective particle size within one of the effective particle size ranges set forth above. The mixing step is followed by an energy-addition step and, in a preferred form of the invention is an annealing step.

II. Second Process Category

The methods of the second processes category include essentially the same steps as in the steps of the first processes category but differ in the following respect. An x-ray diffraction, DSC or other suitable analytical techniques of the presuspension shows the antimicrobial agent in a crystalline form and having an average effective particle size. The antimicrobial agent after the energy-addition step has essentially the same average effective particle size as prior to the energy-addition step but has less of a tendency to aggregate into larger particles when compared to that of the particles of the presuspension. Without being bound to a theory, it is believed the

differences in the particle stability may be due to a reordering of the surfactant molecules at the solid-liquid interface.

III. Third Process Category

The methods of the third category modify the first two steps of those of the first and second processes categories to ensure the antimicrobial agent in the presuspension is in a friable form having an average effective particle size (e.g., such as slender needles and thin plates). Friable particles can be formed by selecting suitable solvents, surfactants or combination of surfactants, the temperature of the individual solutions, the rate of mixing and rate of precipitation and the like. Friability may also be enhanced by the introduction of lattice defects (e.g., cleavage planes) during the steps of mixing the first solution with the aqueous solution. This would arise by rapid crystallization such as that afforded in the precipitation step. In the energy-addition step these friable crystals are converted to crystals that are kinetically stabilized and having an average effective particle size smaller than those of the presuspension. Kinetically stabilized means particles have a reduced tendency to aggregate when compared to particles that are not kinetically stabilized. In such instance the energy-addition step results in a breaking up of the friable particles. By ensuring the particles of the presuspension are in a friable state, the organic compound can more easily and more quickly be prepared into a particle within the desired size ranges when compared to processing an organic compound where the steps have not been taken to render it in a friable form.

In addition to the microprecipitation methods described above, any other known precipitation methods for preparing submicron sized particles or nanoparticles in the art can be used in conjunction with the present invention. The following is a description of examples of other precipitation methods. The examples are for illustration purposes, and are not intended to limit the scope of the present invention.

25 Emulsion Precipitation Methods

One suitable emulsion precipitation technique is disclosed in the co-pending and commonly assigned U.S. Ser. No. 09/964,273, which is incorporated herein by reference and is made a part hereof. In this approach, the process includes the steps of: (1) providing a multiphase system having an organic phase and an aqueous phase, the organic phase having a pharmaceutically effective compound therein; and (2) sonicating the system to evaporate a

portion of the organic phase to cause precipitation of the compound in the aqueous phase and having an average effective particle size of less than about 2 μm . The step of providing a multiphase system includes the steps of: (1) mixing a water immiscible solvent with the pharmaceutically effective compound to define an organic solution, (2) preparing an aqueous based solution with one or more surface active compounds, and (3) mixing the organic solution with the aqueous solution to form the multiphase system. The step of mixing the organic phase and the aqueous phase can include the use of piston gap homogenizers, colloidal mills, high speed stirring equipment, extrusion equipment, manual agitation or shaking equipment, microfluidizer, or other equipment or techniques for providing high shear conditions. The crude emulsion will have oil droplets in the water of a size of approximately less than 1 μm in diameter. The crude emulsion is sonicated to define a microemulsion and eventually to define a submicron sized particle suspension.

Another approach to preparing submicron sized particles is disclosed in co-pending and commonly assigned U.S. Ser. No. 10/183,035, which is incorporated herein by reference and made a part hereof. The process includes the steps of: (1) providing a crude dispersion of a multiphase system having an organic phase and an aqueous phase, the organic phase having a pharmaceutical compound therein; (2) providing energy to the crude dispersion to form a fine dispersion; (3) freezing the fine dispersion; and (4) lyophilizing the fine dispersion to obtain submicron sized particles of the pharmaceutical compound. The step of providing a multiphase system includes the steps of: (1) mixing a water immiscible solvent with the pharmaceutically effective compound to define an organic solution; (2) preparing an aqueous based solution with one or more surface active compounds; and (3) mixing the organic solution with the aqueous solution to form the multiphase system. The step of mixing the organic phase and the aqueous phase includes the use of piston gap homogenizers, colloidal mills, high speed stirring equipment, extrusion equipment, manual agitation or shaking equipment, microfluidizer, or other equipment or techniques for providing high shear conditions.

Solvent Anti-Solvent Precipitation

Suitable solvent anti-solvent precipitation technique is disclosed in U.S. Pat. Nos. 5,118,528 and 5,100,591 which are incorporated herein by reference and made a part hereof. The process includes the steps of: (1) preparing a liquid phase of a biologically active substance in a solvent or a mixture of solvents to which may be added one or more surfactants; (2) preparing a

second liquid phase of a non-solvent or a mixture of non-solvents, the non-solvent is miscible with the solvent or mixture of solvents for the substance; (3) adding together the solutions of (1) and (2) with stirring; and (4) removing of unwanted solvents to produce a colloidal suspension of nanoparticles. The '528 Patent discloses that it produces particles of the substance smaller than
5 500 nm without the supply of energy.

Phase Inversion Precipitation

One suitable phase inversion precipitation is disclosed in U.S. Pat. Nos. 6,235,224, 6,143,211 and U.S. patent application No. 2001/0042932 which are incorporated herein by reference and made a part hereof. Phase inversion is a term used to describe the physical
10 phenomena by which a polymer dissolved in a continuous phase solvent system inverts into a solid macromolecular network in which the polymer is the continuous phase. One method to induce phase inversion is by the addition of a nonsolvent to the continuous phase. The polymer undergoes a transition from a single phase to an unstable two phase mixture: polymer rich and polymer poor fractions. Micellar droplets of nonsolvent in the polymer rich phase serve as
15 nucleation sites and become coated with polymer. The '224 patent discloses that phase inversion of polymer solutions under certain conditions can bring about spontaneous formation of discrete microparticles, including nanoparticles. The '224 patent discloses dissolving or dispersing a polymer in a solvent. A pharmaceutical agent is also dissolved or dispersed in the solvent. For the crystal seeding step to be effective in this process it is desirable the agent is dissolved in the
20 solvent. The polymer, the agent and the solvent together form a mixture having a continuous phase, wherein the solvent is the continuous phase. The mixture is then introduced into at least tenfold excess of a miscible nonsolvent to cause the spontaneous formation of the microencapsulated microparticles of the agent having an average particle size of between 10 nm and 10 μ m. The particle size is influenced by the solvent:nonsolvent volume ratio, polymer
25 concentration, the viscosity of the polymer-solvent solution, the molecular weight of the polymer, and the characteristics of the solvent-nonsolvent pair. The process eliminates the step of creating microdroplets, such as by forming an emulsion, of the solvent. The process also avoids the agitation and/or shear forces.

pH Shift Precipitation

pH shift precipitation techniques typically include a step of dissolving a drug in a solution having a pH where the drug is soluble, followed by the step of changing the pH to a point where the drug is no longer soluble. The pH can be acidic or basic, depending on the particular pharmaceutical compound. The solution is then neutralized to form a presuspension of submicron sized particles of the pharmaceutically active compound. One suitable pH shifting precipitation process is disclosed in U.S. Pat. No. 5,665,331, which is incorporated herein by reference and made a part hereof. The process includes the step of dissolving of the pharmaceutical agent together with a crystal growth modifier (CGM) in an alkaline solution and then neutralizing the solution with an acid in the presence of suitable surface-modifying surface-active agent or agents to form a fine particle dispersion of the pharmaceutical agent. The precipitation step can be followed by steps of diafiltration clean-up of the dispersion and then adjusting the concentration of the dispersion to a desired level. This process of reportedly leads to microcrystalline particles of Z-average diameters smaller than 400 nm as measured by photon correlation spectroscopy.

Other examples of pH shifting precipitation methods are disclosed in U.S. Pat. Nos. 5,716,642; 5,662,883; 5,560,932; and 4,608,278, which are incorporated herein by reference and are made a part hereof.

Infusion Precipitation Method

Suitable infusion precipitation techniques are disclosed in the U.S. Pat. Nos. 4,997,454 and 4,826,689, which are incorporated herein by reference and made a part hereof. First, a suitable solid compound is dissolved in a suitable organic solvent to form a solvent mixture. Then, a precipitating nonsolvent miscible with the organic solvent is infused into the solvent mixture at a temperature between about -10°C and about 100°C and at an infusion rate of from about 0.01 ml per minute to about 1000 ml per minute per volume of 50 ml to produce a suspension of precipitated non-aggregated solid particles of the compound with a substantially uniform mean diameter of less than 10 µm. Agitation (e.g., by stirring) of the solution being infused with the precipitating nonsolvent is preferred. The nonsolvent may contain a surfactant to stabilize the particles against aggregation. The particles are then separated from the solvent. Depending on the solid compound and the desired particle size, the parameters of temperature, ratio of nonsolvent to solvent, infusion rate, stir rate, and volume can be varied according to the invention. The particle size is proportional to the ratio of nonsolvent:solvent volumes and the

temperature of infusion and is inversely proportional to the infusion rate and the stirring rate. The precipitating nonsolvent may be aqueous or non-aqueous, depending upon the relative solubility of the compound and the desired suspending vehicle.

Temperature Shift Precipitation

5 Temperature shift precipitation technique, also known as the hot-melt technique, is disclosed in U.S. Pat. No. 5,188,837 to Domb, which is incorporated herein by reference and made a part hereof. In an embodiment of the invention, lipospheres are prepared by the steps of: (1) melting or dissolving a substance such as a drug to be delivered in a molten vehicle to form a liquid of the substance to be delivered; (2) adding a phospholipid along with an aqueous medium
10 to the melted substance or vehicle at a temperature higher than the melting temperature of the substance or vehicle; (3) mixing the suspension at a temperature above the melting temperature of the vehicle until a homogenous fine preparation is obtained; and then (4) rapidly cooling the preparation to room temperature or below.

Solvent Evaporation Precipitation

15 Solvent evaporation precipitation techniques are disclosed in U.S. Pat. No. 4,973,465 which is incorporated herein by reference and made a part hereof. The '465 Patent discloses methods for preparing microcrystals including the steps of: (1) providing a solution of a pharmaceutical composition and a phospholipid dissolved in a common organic solvent or combination of solvents, (2) evaporating the solvent or solvents and (3) suspending the film
20 obtained by evaporation of the solvent or solvents in an aqueous solution by vigorous stirring. The solvent can be removed by adding energy to the solution to evaporate a sufficient quantity of the solvent to cause precipitation of the compound. The solvent can also be removed by other well known techniques such as applying a vacuum to the solution or blowing nitrogen over the solution.

Reaction Precipitation

25 Reaction precipitation includes the steps of dissolving the pharmaceutical compound into a suitable solvent to form a solution. The compound should be added in an amount at or below the saturation point of the compound in the solvent. The compound is modified by reacting with a chemical agent or by modification in response to adding energy such as heat or UV light or the

like to such that the modified compound has a lower solubility in the solvent and precipitates from the solution.

Compressed Fluid Precipitation

A suitable technique for precipitating by compressed fluid is disclosed in WO 97/14407 to Johnston, which is incorporated herein by reference and made a part hereof. The method includes the steps of dissolving a water-insoluble drug in a solvent to form a solution. The solution is then sprayed into a compressed fluid, which can be a gas, liquid or supercritical fluid. The addition of the compressed fluid to a solution of a solute in a solvent causes the solute to attain or approach supersaturated state and to precipitate out as fine particles. In this case, the compressed fluid acts as an anti-solvent which lowers the cohesive energy density of the solvent in which the drug is dissolved.

Alternatively, the drug can be dissolved in the compressed fluid which is then sprayed into an aqueous phase. The rapid expansion of the compressed fluid reduces the solvent power of the fluid, which in turn causes the solute to precipitate out as fine particles in the aqueous phase. In this case, the compressed fluid acts as a solvent.

Other Methods for Preparing Particles

The particles of the present invention can also be prepared by mechanical grinding of the active agent. Mechanical grinding include such techniques as jet milling, pearl milling, ball milling, hammer milling, fluid energy milling or wet grinding techniques such as those disclosed in U.S. Pat. No. 5,145,684, which is incorporated herein by reference and made a part hereof.

Another method to prepare the particles of the present invention is by suspending an active agent. In this method, particles of the active agent are dispersed in an aqueous medium by adding the particles directly into the aqueous medium to derive a pre-suspension. The particles are normally coated with a surface modifier to inhibit the aggregation of the particles. One or more other excipients can be added either to the active agent or to the aqueous medium.

Example 1: Preparation of 1% Itraconazole Suspension with deoxycholic acid coating.

Each 100 mL of suspension contains:

Itraconazole	1.0 g	(1.0% w/v)
Deoxycholic Acid, Sodium Salt, Monohydrate	0.1 g	(0.1% w/v)
Poloxamer 188, NF	0.1 g	(0.1% w/v)

- 24 -

	Glycerin, USP	2.2 g (2.2% w/v)
	Sodium Hydroxide, NF (0.1 N or 1.0 N)	for pH Adjustment
	Hydrochloric Acid, NF (0.1 N or 1.0 N)	for pH Adjustment
	Sterile Water for Injection, USP	QS
5	Target pH (range)	8.0 (6 to 9)

Preparation of Surfactant Solution (2 Liters) for Microprecipitation

Fill a properly cleaned tank with Sterile Water for Injection and agitate. Add the required amount of glycerin and stir until dissolution. Add the required amount of deoxycholic acid, sodium salt monohydrate and agitate until dissolution. If necessary, adjust the pH of the surfactant solution with minimum amount of sodium hydroxide and/or hydrochloric acid to a pH of 8.0. Filter the surfactant solution through a 0.2 μ m filter. Quantitatively transfer the surfactant solution to the vessel supplying the homogenizer. Chill the surfactant solution in the hopper with mixing.

Preparation of Replacement Solution

Preparation of 4 liters of replacement solution. Fill a properly cleaned tank with WFI and agitate. Add the weighed Poloxamer 188 (Spectrum Chemical) to the measured volume of water. Begin mixing the Poloxamer 188/ water mixture until the Poloxamer 188 has completely dissolved. Add the required amount of glycerin and agitate until dissolved. Once the glycerin has completely dissolved, add the required amount of deoxycholic acid, sodium salt monohydrate and stir until dissolution. If necessary, adjust the pH of the wash solution with the minimum amount sodium hydroxide and/or hydrochloric acid to a pH of 8.0. Filter the replacement solution through a 0.2 μ m membrane filter.

Preparation of Drug Concentrate

For a 2-L batch, add 120.0 mL of N-methyl-2-pyrrolidinone into a 250-mL beaker. Weigh 2.0 g Poloxamer 188. Weigh 20.0 g of itraconazole (Wyckoff). Transfer the weighed Poloxamer 188 to the 250 mL beaker with N-methyl-2-pyrrolidinone. Stir until dissolved, then add the itraconazole. Heat and stir until dissolved. Cool the drug concentrate to room temperature and filter through a 0.2-micron filter.

Microprecipitation

Add sufficient WFI to the surfactant solution already in the vessel supplying the homogenizer so that the desired target concentration is reached. When the surfactant solution is cooled, start adding the drug concentrate into the surfactant solution with continuous mixing.

Homogenization

- 5 Slowly increase the pressure of the homogenizer until the operating pressure 10,000 psi has been reached. Homogenize the suspension with recirculation while mixing. For 2,000 mL of suspension at 50Hz, one pass should require approximately 54 seconds. Following homogenization, collect a 20-mL sample for particle size analysis. Cool the suspension.

Wash Replacement

- 10 The suspension is then divided and filled into 500-mL centrifuge bottles. Centrifuge until clean separation of sediment is observed. Measure the volume of supernatant and replace with fresh replacement solution, prepared earlier. Quantitatively transfer the precipitate from each centrifuge bottle into a properly cleaned and labeled container for resuspension (pooled sample). Resuspension of the pooled sample is performed with a high shear mixer until no visible clumps are observed. Collect a 20-mL sample for particle size analysis.

- 15 The suspension is then divided and filled into 500-mL centrifuge bottles. Centrifuge until clean separation of sediment is observed. Measure the volume of supernatant and replace with fresh replacement solution, prepared earlier. Quantitatively transfer the precipitate from each centrifuge bottle into a properly cleaned and labeled container for resuspension (pooled sample).
20 Resuspension of the pooled sample is performed with a high shear mixer until no visible clumps are observed. Collect a 20-mL sample for particle size analysis.

Second Homogenization

- 25 Transfer the above suspension to the hopper of the homogenizer and chill the suspension with mixing. Slowly increase the homogenizer pressure until an operating pressure 10,000 psi has been reached. Homogenize while monitoring the solution temperature. Following homogenization, cool the suspension and collect three 30-mL samples for particle analysis. Collect the remaining suspension in a 2-liter bottle.

Filling

- 26 -

Based on acceptable particle size determination testing (mean volume-weighted diameter of 50 nm to 5 microns), collect 30 mL samples in 50 mL glass vials with rubber stoppers.

Example 2: Preparation of 1% Itraconazole Nanosuspension with phospholipid coating.

Each 100 mL of suspension contains:

5	Itraconazole	1.0 g (1.0% w/v)
	Phospholipids (Lipoid E 80)	1.2 g (1.2% w/v)
	Glycerin, USP	2.2 g (2.2% w/v)
	Sodium Hydroxide, NF (0.1 N or 1.0 N)	for pH Adjustment
	Hydrochloric Acid, NF (0.1 N or 1.0 N)	for pH Adjustment
10	Sterile Water for Injection, USP	QS
	Target pH (range)	8.0 (7.5 to 8.5)

Preparation of Surfactant Solution (2 Liters) for Microprecipitation

The surfactant solution is prepared in two phases. Phase 1 is dispersed phospholipids, whereas Phase 2 includes filtered glycerin. The two fractions are combined prior to pH
15 adjustment.

Phase 1: Fill a properly cleaned vessel with approximately 700 mL of Sterile Water for Injection, USP (WFI) with agitation at 50 - 500 rpm. Increase the temperature of the filtrate to 50°C - 70°C and add the required amount of phospholipids with mixing at 50 - 500 rpm until complete dispersion is achieved. Document the time and temperature at which the phospholipids were
20 added and at which it was dispersed. Determine the total mixing time required to disperse the phospholipids. Cool the surfactant solution to 18°C - 30°C prior to the addition of glycerin.

Phase 2: Fill a properly cleaned vessel with approximately 700 mL of WFI with agitation at 50 - 500 rpm. Add the required amount of glycerin at 18°C - 30°C and agitate at 50 - 500 rpm until dissolution.

25 Combined Phases: Filter the glycerin solution through a 0.2 µm filter set-up into Phase 1 (at 18°C - 30°C) while mixing at 50 - 500 rpm. Volume is approximately 1.4 liters. Record the pH of the surfactant solution. If necessary, adjust the pH of the surfactant solution with a minimum

- 27 -

amount of sodium hydroxide and/or hydrochloric acid to a pH of 8.0 ± 0.5 . Measure the volume of the surfactant solution at $18^{\circ}\text{C} - 30^{\circ}\text{C}$ using a 2-L graduated cylinder.

Quantitatively transfer the surfactant solution to the vessel supplying the homogenizer (Avestin C-160). Chill the surfactant solution in the hopper with mixing at a speed with an observable solution vortex until the temperature is not more than 10°C .

Preparation of Replacement Solution (4 L)

The replacement solution is prepared in two phases. Phase 1 includes dispersed phospholipids, whereas Phase 2 includes filtered glycerin. The two fractions are combined prior to pH adjustment.

10 Phase 1: Fill a properly cleaned vessel with approximately 1.4 liters of WFI with agitation at 50 - 500 rpm. Increase the temperature of the water to $50^{\circ}\text{C} - 70^{\circ}\text{C}$ and add the required amount of phospholipids with mixing at 50 - 500 rpm until complete dispersion is achieved. Cool the surfactant solution to $18^{\circ}\text{C} - 30^{\circ}\text{C}$ prior to the addition of glycerin.

15 Phase 2: Fill a properly cleaned vessel with approximately 1.4 L of WFI with agitation at 50 - 500 rpm. Add the required amount of glycerin and agitate at 50 - 500 rpm until dissolution.

Combined Phases: Filter the glycerin solution through a $0.2\ \mu\text{m}$ filter set-up into Phase 1 (at $18^{\circ}\text{C} - 30^{\circ}\text{C}$) while mixing at 50 - 500 rpm. Dilute to volume with Water for Injection to 4.0 L using a graduated cylinder. Record the pH of the wash solution. If necessary, adjust the pH of the wash solution with the minimum amount sodium hydroxide and/or hydrochloric acid to a pH of 8.0 ± 0.5 .

Preparation of Drug Concentrate

25 For a 2-L batch, add 120.0 mL of N-methyl-2-pyrrolidinone (Pharmasolve®, ISP) into a 250-mL beaker. Weigh 20.0 g of itraconazole (Wyckoff). Transfer the weighed itraconazole to the 250-mL beaker with NMP at NMT 70°C . Maintain below 70°C and stir at 100 - 1000 rpm until dissolved. Cool the drug concentrate to $18^{\circ}\text{C} - 30^{\circ}\text{C}$. Filter the drug concentrate through a prefilter and filter set-up. Use one polypropylene prefilter SBPP and two $0.2\ \mu\text{m}$ filters at 15 psi and ambient temperature. Transfer the drug concentrate to three 60-mL syringes and attach

syringe needles to the luer connections of the syringes. Using the syringes, determine the volume of drug concentrate.

Microprecipitation

Add Water for Injection to the surfactant solution already in the vessel supplying the
5 homogenizer. The amount of water at 18°C - 30°C added should be calculated as:

$$V = 2,000 \text{ mL} - \text{Volume of Drug Concentrate} - \text{Volume of Surfactant Solution}$$

Mount each syringe needle assembly using a syringe pump. Position the outlet of the needle on top of the vessel. When the surfactant solution is not more than 10°C, start adding the drug concentrate into the surfactant solution with continuous mixing at a speed needed to create a
10 distinctive solution vortex. The concentrate should be added so that the drops hit the point of highest shear, at the bottom of the vortex. The rate of addition should be approximately 2.5 mL/min.

Homogenization

An Avestin C160 homogenizer was used. Slowly increase the pressure of the
15 homogenizer until the operating pressure 10,000 psi has been reached. Homogenize the suspension for 20 passes (18 minutes) with recirculation while mixing at 100 – 300 rpm and maintaining the suspension temperature below 70°C. For 2,000 mL of suspension at 50 Hz, one pass requires approximately 54 seconds. Following homogenization, collect a 20 mL sample in a 50 mL glass vial for particle size analysis. Cool the suspension to not more than 10°C.

20 Wash Replacement

The suspension is then divided and filled into 500-mL centrifuge bottles. Set the speed for the centrifuge at 11,000 rpm using the rotor SLA-3000, Superlite equivalent to approximately 20,434 g. The total centrifuge time is 60 min at not more than 10°C. Measure the volume of supernatant and replace with fresh replacement solution. Using spatula(s), quantitatively transfer
25 the precipitant from each centrifuge bottle into a properly cleaned and labeled container for resuspension (pooled sample). Resuspension of the pooled sample is performed with a high shear mixing until no visible clumps are observed.

Second Washing and Centrifuging Step

The suspension is then divided and filled into 500-mL centrifuge bottles. Set the speed for the centrifuge at 11,000 rpm using the rotor SLA-3000, Superlite equivalent to approximately 20,434 g. The total centrifuge time is 60 min at not more than 10°C. Measure the volume of supernatant and replace with fresh replacement solution. Using spatula(s), quantitatively transfer the precipitant from each centrifuge bottle into a properly cleaned and labeled container for resuspension (pooled sample). Resuspension of the pooled sample is performed under high-shear mixing until no visible clumps are observed. Record the pH of the suspension. If necessary, adjust the pH of the suspension with the minimum amount sodium hydroxide and/or hydrochloric acid to a pH of 8.0 ± 0.5 .

Second Homogenization

Transfer the above suspension to the hopper of the homogenizer. Chill the suspension with mixing until the temperature is less than 10°C. Slowly increase the pressure until an operating pressure of 10,000 psi has been reached. Homogenize for 20 passes (18 minutes) while maintaining the solution temperature below 70°C. Following homogenization, cool the suspension to less than 10°C and collect three 30-mL samples for particle-size analysis. Collect the remaining suspension in a 2-liter bottle. Sparge the suspension with nitrogen gas for 10 min. Ensure the nitrogen gas is filtered through a 0.2 μm filter.

Filling

Based on acceptable particle size determination testing (mean volume-weighted diameter of 50 to 1000 nm), collect 30-mL samples in 50-mL glass vials with PTFE®-coated stoppers. Purge the headspace of each vial with nitrogen prior to sealing.

Example 3: Other formulations of Itraconazole Suspensions

Other formulations of itraconazole suspensions with different combinations of the surfactants can also be prepared using the method described in Example 1 or Example 2. Table 1 summarizes the compositions of the surfactants of the various itraconazole suspensions.

Table 1: Summary of the compositions of the various 1% itraconazole suspensions

Formulation No.	Surfactants in the formulation	Amount*
1	Poloxamer 188	0.1%

	Deoxycholate	0.1%
	Glycerin	2.2%
2	Poloxamer 188	0.1%
	Deoxycholate	0.5%
	Glycerin	2.2%
3	Poloxamer 188	2.2%
	Deoxycholate	0.1%
	Glycerin	2.2%
4	Poloxamer 188	2.2%
	Deoxycholate	0.5%
	Glycerin	2.2%
9	Solutol	0.3%
	Deoxycholate	0.5%
	Glycerin	2.2%
14331-1	Solutol	1.5%
	Glycerin	2.2%
14443-1	Albumin	5%
14	Phospholipid	2.2%
	Deoxycholate	0.5%
	Glycerin	2.2%
	Na ₂ PO ₄	0.14%
A6	Phospholipid	1.2%
	Glycerin	2.2%
B	Phospholipid	1.2%
	Glycerin	2.2%
	N-methyl-2-pyrrolidinone	trace
C	Phospholipid	1.2%
	Glycerin	2.2%
	Lactic acid	trace
14412-3	Phospholipid	1.2%
	Hydroxyethyl starch	1.0%
	Glycerin	2.2%
	TRIS	0.06%

* % by weight of the final volume of the suspension (w/v)

Example 4: Comparison of the acute toxicity between commercially available itraconazole formulation (SPORANOX®) and the suspension compositions of the present invention.

The acute toxicity of the commercially available itraconazole formulation (SPORANOX®) is compared to that of the various 1% itraconazole formulations in the present invention as listed in Table 1. SPORANOX® is available from Janssen Pharmaceutical Products, L.P. It is available as a 1% intravenous (I.V.) solution solubilized by hydroxypropyl-β-cyclodextrin. The results are shown in Table 2 with the maximum tolerated dose (MTD) indicated for each formulation.

10 Table 2: Comparison of the acute toxicity of various formulations of itraconazole

Formulation Number	Results and Conclusions
SPORANOX® I.V.	LD ₁₀ =30 mg/kg

	MTD=20 mg/kg (slight ataxia)
1	MTD=320 mg/kg; NOEL=80 mg/kg Spleen obs ^b : 320 mg/kg Red ears/feet: ≥160 mg/kg
2	MTD=320 mg/kg Spleen obs ^b : 320 mg/kg Slight lethargy: 320 mg/kg Red urine: ≥80 mg/kg Tail obs ^c : ≥40mg/kg
3	MTD=160 mg/kg; NOEL=80 mg/kg Spleen obs ^b : 320 mg/kg Red ears/feet: ≥160 mg/kg
4	MTD=160 mg/kg LD ₂₀ = 320 mg/kg Spleen obs ^b : 320 mg/kg Slight lethargy: 320 mg/kg Red urine: ≥40 mg/kg Tail obs ^c : ≥40 mg/kg
9	LD ₆₀ =320 mg/kg; MTD=160 mg/kg Spleen obs ^b : 320 mg/kg Tail obs: 320 mg/kg Red ears/feet: ≥160 mg/kg Red urine: ≥40 mg/kg
14331-1	MTD=40 mg/kg; NOEL=40 mg/kg LD ₄₀ =80 mg/kg
14443-1	LD ₄₀ =80 mg/kg; NOEL=40 mg/kg
14	MTD=320 mg/kg; NOEL=40-80 mg/kg Spleen obs ^b : 320 mg/kg Ataxia=320 mg/kg Tail obs=320 mg/kg
A6	MTD=320 mg/kg; NOEL=160 mg/kg Spleen obs ^b : 320 mg/kg
B	MTD=320 mg/kg; NOEL=80 mg/kg Spleen obs ^b : 160 mg/kg Red ears/feet: ≥160 mg/kg
C	MTD=320 mg/kg; NOEL=80 mg/kg Spleen obs ^b : ≥160 mg/kg Red ears/feet: ≥160 mg/kg
14412-3	MTD=320 mg/kg; NOEL=80 mg/kg Spleen obs ^b : ≥160 mg/kg

^acyclodextrin = hydroxypropyl-β-cyclodextrin

^bSpleen obs = Enlarged and/or pale

^cTail obs = gray to black and/or necrosis

LD₁₀ = Lethal dose resulting in 10% mortality

5 LD₄₀ = Lethal dose resulting in 40% mortality

LD₅₀ = Lethal dose resulting in 50% mortality

NOEL = No effect level

MTD = Maximum tolerated dose

The data in Table 2 indicated that the animals tolerated a much higher level of the antifungal agent itraconazole when formulated in a nanosuspension than when formulated as a solution with cyclodextrin. It may be thought that the reason for the increased tolerability is associated with not using cyclodextrin. However, cyclodextrin, by itself, at the levels used in SporanoX would not cause the degree of toxicity observed. Rather, it is believe, the reason lies in alteration of the pharmacokinetic profile caused by the nanosuspension.

Example 5: Pharmacokinetic comparison of SPORANOX® vs. suspension formulation of itraconazole.

Young adult, male Sprague Dawley rats were treated intravenously (IV) via a caudal tail vein with a single injection at a rate of 1 ml/min. with either SPORANOX® Injection, or Formulations 1 and B at 20, 40, and 80 mg/kg, or Formulations 3, 14, A6 and C at 80 mg/kg.

Following administration, the animals were anesthetized and retro-orbital blood was collected at different time points (n=3). The time points were as follows: 0.03, 0.25, 0.5, 1, 2, 4, 6, 8, 24, 48, 96, 144, 192, 288, and 360 hours (SPORANOX® Injection only to 192 hours). Blood was collected into tubes with EDTA and centrifuged at 3200 rpm for 15 minutes to separate plasma. The plasma was stored frozen at -70°C until analysis. The concentration of the parent itraconazole and the metabolite hydroxy-itraconazole were determined by high-performance liquid chromatography (HPLC). Pharmacokinetic (PK) parameters for itraconazole (ITC) and hydroxy-itraconazole (OH-ITC) were derived using noncompartmental methods with WinNonlin® Professional Version 3.1 (Pharsight Corp., Mountain View, CA).

Table 3 provides a comparison of the plasma pharmacokinetic parameters determined for each itraconazole formulation. Plasma itraconazole was no longer detected at 24 hours for SPORANOX® Injection at 5 mg/kg, at 48 hours for SPORANOX® Injection at 20 mg/kg, and at 96 hours for Formulations 1 and B. Plasma hydroxy-itraconazole was initially detected at 0.25 hours for SPORANOX® Injection and Formulations 1 and B. Plasma hydroxy-itraconazole was initially detected at 0.25 hours for SPORANOX® Injection at 5 and 20 mg/kg and Formulations 1 and ;B at 20 mg/kg, Hydroxy-itraconazole was no longer detected at 48 hours for SPORANOX® Injection at 5mg/kg, at 96 hours for SPORANOX® Injection at 20 mg/kg, and at 144 hours for Formulations 1 and B.

Table 3. Comparison of Plasma Pharmacokinetic Parameters for Sporanox and a Suspension Formulation After IV Administration in Rats

Analyte	PK Parameters	Dosage/Formulation											
		5 mg/kg		20 mg/kg		40 mg/kg		80 mg/kg					
		Spor	I	Spor	I	B	I	B	I	B	A6	C	3
Itraconazole	C max (ug/ml)	2.42	30.41	13.12	119.16	9.10	10.20	446.33	15.20	39.72	53.19	365.09	68.15
	T max (h)	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
	AUC (0-∞) (ug.h/ml)	3.90	16.70	28.25	42.67	15.79	36.11	143.70	80.31	58.71	94.19	108.87	85.53
	T ½ (h)	2.75	14.36	5.36	23.95	14.54	20.49	25.89	28.63	54.02	33.75	38.46	31.17
	CL (ml/h)	320.17	299.35	176.97	234.38	316.67	276.90	139.18	249.04	340.64	212.33	183.71	233.83
Hydroxy-Itraconazole	MRT (h)	2.57	13.29	4.48	24.37	15.32	28.76	27.45	52.84	58.21	46.85	31.21	41.93
	C max (ug/ml)	0.38	0.40	0.78	0.61	0.44	0.69	1.03	0.48	0.32	0.56	0.52	0.51
	T max (h)	4.04	24	4.0	24	8	24	24	48	48	24.0	24.0	24.0
	AUC (0-∞) (ug.h/ml)	3.96	17.89	13.41	37.71	20.71	44.69	70.24	56.01	47.27	59.40	51.27	51.89
	T ½ (h)	7.98	15.50	5.89	22.27	18.06	28.12	23.21	36.45	60.87	38.84	50.29	25.50
MRT (h)	7.55	30.99	12.17	43.06	29.23	36.02	46.80	68.35	74.88	65.71	60.81	58.02	

**THIS PAGE WAS NOT FURNISHED UPON FILING
THE INTERNATIONAL APPLICATION**

FIG. 5 compares the pharmacokinetics (PK) of SPORANOX® with Formulation 1 suspension of itraconazole particles. Because, as shown above, the present suspension formulation is less toxic than SporanoX®, it was administered at higher amounts in this equitoxic experiment. SporanoX was dosed at 20 mg/kg and Formulation 1 at 80 mg/kg. The SPORANOX® decreases in plasma concentration relatively quickly, over 20 hours. The nanosuspension plasma levels remain elevated for approximately 3-4 times longer. The nanosuspension exhibits an initial minimum at 30 minutes in the plasma level. This corresponds to a nadir in plasma concentration due to sequestration of the drug nanocrystals by the macrophages of the spleen and liver, thus temporarily removing drug from circulation. However, the drug levels rebound quickly, as the macrophages apparently release the drug into the circulation. Furthermore, the nanosuspension drug is metabolized effectively, as is shown by the PK curve for the hydroxy itraconazole metabolite. The rate of appearance of the metabolite for the nanosuspension is delayed, compared with the PK curve for the metabolite for the SPORANOX® formulation. However, as with the case of the parent molecule for the nanosuspension, the metabolite persists in circulation for a much longer time than is the case with the metabolite for the SPORANOX® formulation. When the AUC (area under the blood concentration vs time curve) is normalized by the dose, the nanosuspension is at least as bioavailable as SPORANOX®.

Example 6: Acute Toxicity Of Fast Dissolving Nanosuspensions

Additional experiments were performed. Itraconazole nanosuspensions were formulated differently, so as to dissolve much more readily in blood. This was accomplished by making the particles either smaller or amorphous, or both. These acute toxicity of these formulations is described for formulation entries 14331-1 and 14443-1 in Table 1. In contrast to the slowly dissolving nanosuspensions, the fast dissolving nanosuspension caused death in the animals at much lower levels, similar to what was found with SPORANOX®. Since these fast dissolving nanosuspensions did not contain cyclodextrin, it is clear that this excipient was not responsible for the toxicity. Rather the rapid dissolution, resulting in immediate availability of the drug in the blood was the causative factor. The drug level for the rapidly dissolving formulation, Form A, is much higher than that attained by the slow dissolving (macrophage targeting) formulation, Form

B, as determined in an in vitro dissolution experiment. This involved a plasma simulating media consisting of 5% albumin/ Sorenson's buffer. Results are shown in Figure 6.

Example 7: Antifungal Efficacy Studies

Normal and immuno-suppressed (prednisolone administered twice daily on the day before and on the day of inoculation) rats inoculated with 9.5×10^6 or 3×10^6 cfu *C. albicans*/ml saline once intravenously were intravenously treated with SPORANOX[®] Injection once daily for ten consecutive days, with the first dose given 4 to 5 hours after inoculation. SPORANOX[®] Injection rats were dosed at 5 or 20 mg/kg for the first 2 days, then at 5 or 10 mg/kg for the remaining 8 days, due to toxicity at 20 mg/kg after 2 days of dosing. Similarly, immuno-suppressed rats inoculated with $1 \times 10^{6.5}$ cfu *C. albicans*/ml saline were intravenously treated with Formulation 1 or B each at 20, 40, or 80 mg/kg once every other day for ten days, beginning the day of inoculation. The SPORANOX[®] Injection, Formulation 1, and Formulation B treatment rats were terminated 11 days after the *C. albicans* inoculation and the kidneys were collected, weighed and cultured for determination of *C. albicans* colony counts and itraconazole and hydroxy-itraconazole concentration. Kidneys were collected from untreated control rats when a moribund condition was observed or when an animal had a 20% body weight. In addition, body weights were measured periodically during the course of each study.

Comparison of results for immuno-suppressed rats treated with SPORANOX[®] Injection and Formulations 1 and B are shown in Table 4 and Figure 7. Daily SPORANOX[®] Injection treatment at 10 – 20 mg/kg appeared to be slightly more effective than daily treatment with SPORANOX[®] Injection at 5 mg/kg. Based on kidney colony counts, every other day dosing at 20 mg/kg of Formulation 1 or B appeared to be as effective as every day dosing with SPORANOX[®] Injection at 20 mg/kg and possibly more effective than SPORANOX[®] Injection at 5 mg/kg (i.e., the recommended clinical dose), whereas the higher doses for both Formulation 1 and B appeared to most effective, based on kidney colony counts (i.e., *C. albicans* not detected) and increased kidney itraconazole concentration.

Table 4. Mean *C. albicans* Colony Count and Itraconazole and Hydroxy-Itraconazole Concentration in Kidney

Treatment	<i>C. albicans</i> Titer		Concentration in Kidney	
	Count (cfu/g)	Incidence	ITC (µg/g)	OH-ITC (µg/g)
No Treatment (3x 10 ⁶ cfu/ml)	6.9 x 10 ⁴	6/6	--	--
SPORANOX [®] , 5 mg/kg, (3x 10 ⁶ cfu/ml)	96.5	6/6	1.2	1.5
SPORANOX [®] , 10-20 mg/kg, (3 x 10 ⁶ cfu/ml)	12.4	4/6	8.5	8.0
No Treatment (2.5x 10 ⁶ cfu/ml)	3.5 x 10 ⁵	6/6	--	--
Formulation 1, 20 mg/kg, (2.5 x 10 ⁶ cfu/ml)	5.3	4/6	6.1	5.7
Formulation 1, 40 mg/kg, (2.5 x 10 ⁶ cfu/ml)	0	0/6	18.5	6.0
Formulation 1, 80 mg/kg, (2.5 x 10 ⁶ cfu/ml)	0	0/6	41.2	6.2
No Treatment (2.5x 10 ⁶ cfu/ml)	8.0 x 10 ⁴	6/6	--	--
Formulation B, 20 mg/kg, (2.5 x 10 ⁶ cfu/ml)	8.9	4/6	2.5	2.5
Formulation B, 40 mg/kg, (2.5 x 10 ⁶ cfu/ml)	0	0/6	7.8	4.0
Formulation B, 80 mg/kg, (2.5 x 10 ⁶ cfu/ml)	0	0/6	21.3	4.6

In the examples above, a nanosuspension formulation of an anti-fungal agent was shown to be less toxic than a conventional totally soluble formulation of the same drug. Thus, more of the drug could be administered without eliciting adverse effects. Because the nanoparticles of the drug did not immediately dissolve upon injection, they were trapped in a depot store in the liver and spleen. These acted as prolonged release sanctuaries, permitting less frequent dosing. The greater dosing that could be administered permitted greater drug levels to be manifested in the target organs, in this case, the kidney (Figure 8). The greater drug levels in this organ led to a greater kill of infectious organisms. (Figure 9).

Example 8: Resistant Strain Anti-fungal Efficacy Test

A lethal dose of a *C. albicans* strain c43 (ATCC number 201794) (MIC₈₀= 16 µg/ml for SPORANOX[®] itraconazole; 8-16 for Vfend, and 0.1 for Cancidas) was administered to an immunocompromised rat model (prednisolone qd). 24h later, test groups (n=6) were treated q2d with 20, 40, or 80 mg/kg NANOEDGE[™] itraconazole nanosuspension. Control groups included a no treatment arm, Sporanox[®] (10 mg/kg/d), Vfend[®] (10 mg/kg/d), and Cancidas[®] (1mg/kg/d). Treatment was continued for 10 days. Survival and kidney cfu/g were assessed.

The number of surviving animals after 6 and 10 days, were respectively: Sporanox (3,0), 20 and 40 mg/kg nanosuspension (5,3), 80 mg/kg nanosuspension (6,4), Vfend (0,0), Cancidas (0,0). FIG. 10.

5 It can be concluded that the greater dosing possible with the itraconazole nanosuspension can effectively treat infections of *C. albicans* strains, conventionally assumed to be resistant to itraconazole, resulting in increased survival in an immunocompromised rat model.

10 Current definitions of sensitive and resistant fungal strains presume a specified dose of itraconazole that is administered, using conventional dosage forms. Greater drug loading, attendant with nanosuspension injections, may permit treatment of what are currently considered itraconazole-resistant *C. albicans* infections.

Example 9: Prophetic examples of other triazole antifungal agents

15 The present invention contemplates preparing a 1% suspension of submicron- or micron size of a triazole antifungal agent using the method described in Example 1 or Example 2 and the formulations described in Example 3 with the exception that the antifungal agent is a triazole antifungal agent other than itraconazole. Examples of triazole antifungal agents that can be used include, but are not limited to, ketoconazole, miconazole, fluconazole, ravuconazole, voriconazole, saperconazole, eberconazole, genaconazole, clotrimazole, econazole, oxiconazole, sulconazole, terconazole, tioconazole, and posaconazole.

Example 10: Prophetic example of a non-triazole antifungal agent

20 The present invention contemplates preparing a 1% suspension of submicron- or micron size non-triazole antifungal agent using the method described in Example 1 or Example 2 and the formulations described in Example 3 with the exception that the antifungal agent is amphotericin B, nystatin, terbinafine, anidulafungin, or flucytosine instead of itraconazole.

25 From the foregoing, it will be observed that numerous variations and modifications may be effected without departing from the spirit and scope of the invention. It is to be understood that no limitation with respect to the specific apparatus illustrated herein is intended or should be inferred. It is, of course, intended to cover by the appended claims all such modifications as fall within the scope of the claims.

CLAIMS

What is claimed is:

1. A composition of an antimicrobial agent that renders the agent potent against organisms normally considered to be resistant to the agent, the composition comprising an aqueous suspension of submicron- to micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives, wherein the particles have a volume-weighted mean particle size of less than 5 μm as measured by laser diffractometry.
2. The composition of claim 1, wherein the particles have a volume-weighted mean particle size of less than 2 μm as measured by laser diffractometry.
3. The composition of claim 1, wherein the particles have a volume-weighted mean particle size of less than about 1 μm as measured by laser diffractometry.
4. The composition of claim 1, wherein the particles have a volume-weighted mean particle size of from about 150 nm to about 1 μm as measured by laser diffractometry.
5. The composition of claim 1, wherein the antimicrobicidal agent is an antifungal agent.
6. The composition of claim 5, wherein the antimicrobicidal agent is a triazole antifungal agent.
7. The composition of claim 6, wherein the triazole antifungal agent is selected from the group consisting of: itraconazole, ketoconazole, miconazole, fluconazole, ravuconazole, voriconazole, saperconazole, eberconazole, genaconazole, clotrimazole, econazole, oxiconazole, sulconazole, terconazole, tioconazole, and posaconazole.
8. The composition of claim 1, wherein the antimicrobicidal agent is itraconazole.
9. The composition of claim 1, wherein the ionic surfactant is selected from the group consisting of: anionic surfactants, cationic surfactants, zwitterionic surfactants, and combinations thereof.

10. The composition of claim 9, wherein the anionic surfactant is selected from the group consisting of: alkyl sulfonates, alkyl phosphates, alkyl phosphonates, potassium laurate, triethanolamine stearate, sodium lauryl sulfate, sodium dodecylsulfate, alkyl polyoxyethylene sulfates, sodium alginate, dioctyl sodium sulfosuccinate, phosphatidylserine, phosphatidylinositol, diphosphatidylglycerol, phosphatidylglycerol, phosphatidylinosine, phosphatidic acid and its salts, sodium carboxymethylcellulose, cholic acid and other bile acids and salts thereof.

11. The composition of claim 10, wherein the bile acid is selected from the group consisting of cholic acid, deoxycholic acid, glycocholic acid, taurocholic acid, and glycodeoxycholic acid.

12. The composition of claim 9, wherein the anionic surfactant is a phospholipid.

13. The composition of claim 12, wherein the phospholipid is natural or synthetic.

14. The composition of claim 12, wherein the phospholipid is pegylated.

15. The composition of claim 8, wherein the cationic surfactant is selected from the group consisting of: quaternary ammonium compounds, such as benzalkonium chloride, cetyltrimethylammonium bromide, lauryldimethylbenzylammonium chloride, acyl carnitine hydrochlorides, alkyl pyridinium halides, or aliphatic amines.

16. The composition of claim 9, wherein the zwitterionic surfactant is a phospholipid.

17. The composition of claim 16, wherein the phospholipid is natural or synthetic.

18. The composition of claim 16, wherein the phospholipid is pegylated.

19. The composition of claim 1, wherein the nonionic surfactant is selected from the group consisting of: glyceryl esters, polyoxyethylene fatty alcohol ethers (Macrogol and Brij), polyoxyethylene sorbitan fatty acid esters (Polysorbates), polyoxyethylene fatty acid esters (Myrj), sorbitan esters (Span), glycerol monostearate, polyethylene glycols, polypropylene glycols, cetyl alcohol, cetostearyl alcohol, stearyl alcohol, aryl alkyl polyether alcohols, polyoxyethylene-polyoxypropylene copolymers (poloxamers), poloxamines, methylcellulose,

hydroxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, noncrystalline cellulose, polysaccharides including starch and starch derivatives such as hydroxyethylstarch (HES), polyvinyl alcohol, and polyvinylpyrrolidone.

20. The composition of claim 1, wherein the biologically derived surfactant is selected from the group consisting of: albumin, casein, other proteins and polysaccharides.

21. The composition of claim 20, wherein the polysaccharide is selected from the group consisting of starches, heparin and chitosans.

22. The composition of claim 1, wherein the amino acid is selected from the group consisting of: leucine, alanine, valine, isoleucine, lysine, aspartic acid, glutamic acid, methionine, tyrosine and phenylalanine.

23. The composition of claim 1, wherein the amino acid derivative is an amide, an ester, or a polypeptide.

24. The composition of claim 1, wherein the surfactant is a bile salt.

25. The composition of claim 24, wherein the bile salt is deoxycholate.

26. The composition of claim 1, wherein the surfactant is a polyalkoxyether.

27. The composition of claim 26, wherein the polyalkoxyether is Poloxamer 188.

28. The composition of claim 1, wherein the surfactant is hydroxyethylstarch.

29. The composition of claim 1, wherein the surfactant is polyethylene-660-hydroxystearate.

30. The composition of claim 1, wherein the surfactant is albumin.

31. The composition of claim 1, wherein the surfactant is a phospholipid.

32. The composition of claim 1, wherein the aqueous medium further comprises a pH adjusting agent.

33. The composition of claim 32, wherein the pH adjusting agent is selected from the group consisting of: hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, lactic acid, succinic acid, citric acid, tris(hydroxymethyl)aminomethane, meglumine, sodium hydroxide, and amino acids.

34. The composition of claim 33, wherein the amino acid is selected from the group consisting of: glycine, arginine, lysine, alanine, methionine, valine, asparagine, tyrosine, proline, serine, isoleucine, tryptophan, phenylalanine, threonine, cysteine, glutamine, aspartic acid, glutamic acid, histidine, taurine and leucine.

35. The composition of claim 1, further comprising an osmotic pressure adjusting agent.

36. The composition of claim 35, wherein the osmotic pressure adjusting agent is selected from the group consisting of: glycerin, monosaccharides, disaccharides, trisaccharides, and sugar alcohols.

37. The composition of claim 36, wherein the monosaccharide is dextrose.

38. The composition of claim 36, wherein the disaccharide is selected from the group consisting of sucrose, maltose and trehalose.

39. The composition of claim 36, wherein the trisaccharide is raffinose.

40. The composition of claim 36, wherein the sugar alcohol is mannitol or sorbitol.

41. The composition of claim 1, wherein the antimicrobial agent is present in an amount of from about 0.01% to about 50% w/v.

42. The composition of claim 1, wherein the antimicrobial agent is present in an amount of from about 0.05% to about 30% w/v.

43. The composition of claim 1, wherein the antimicrobial agent is present in an amount of about 0.1% to about 20% w/v.

44. The composition of claim 1, wherein the surfactant is present in an amount of from about 0.001% to about 5% w/v.

45. The composition of claim 1, wherein the surfactant is present in an amount of from about 0.005% to about 5% w/v.

46. The composition of claim 1, wherein the surfactant is present in an amount of from about 0.01% to about 5% w/v.

47. The composition of claim 1 is administered by a route selected from the group consisting of: parenteral, oral, buccal, periodontal, rectal, nasal, pulmonary, and topical.

48. The composition of claim 1 is administered by a route selected from the group consisting of intravenous, intramuscular, intracerebral, subcutaneous, intradermal, intralymphatic, pulmonary, intraarticular, intrathecal, and intraperitoneal.

49. The composition of claim 1, wherein the aqueous medium is removed to form dry particles.

50. The composition of claim 49, wherein the method of removing the aqueous medium is selected from the group consisting of: evaporation and lyophilization.

51. The composition of claim 49, wherein the method of removing the aqueous medium is by lyophilization.

52. The composition of claim 49, wherein the dry particles are formulated into an acceptable pharmaceutical dosage form.

53. The composition of claim 52, wherein the pharmaceutical dosage form is selected from the group consisting of: parenteral solutions, tablets, capsules, suspensions, creams, lotions, emulsions, pulmonary formulations, topical formulations, controlled or sustained release formulations, and tissue specific targeted delivery formulations.

54. The composition of claim 1, wherein the composition is frozen.

55. A composition of an antimicrobial agent that renders the agent potent against organisms normally considered to be resistant to the agent, the composition comprising an aqueous suspension of submicron- to micron-size particles of itraconazole coated with at least one surfactant, and an osmotic pressure adjusting agent, wherein the nanoparticles having a volume-weighted mean particle size of less than 5 μm as measured by laser diffractometry, and wherein the itraconazole is present in an amount of from about 0.01% to about 50% w/v, and the surfactant is present in an amount of from about 0.001% to about 5%.

56. A composition of particles of an antimicrobial agent that renders the agent potent against organisms normally considered to be resistant to the agent, the composition prepared by a method comprising the steps of:

- (i) dissolving the antimicrobial agent in a water-miscible first solvent to form a solution;
- (ii) mixing the solution with a second solvent which is aqueous to define a pre-suspension; and
- (iii) adding energy to the pre-suspension to form particles having an average effective particle size of less than 5 μm ;

wherein the solubility of the antimicrobial agent is greater in the first solvent than in the second solvent, and the second solvent comprising one or more surfactants selected from the group consisting of: nonionic surfactants, ionic surfactants, biologically derived surfactants, and amino acids and their derivatives.

57. A method for rendering an antimicrobial agent potent against organisms normally considered to be resistant to the agent, the method comprising formulating the agent as an aqueous suspension of submicron- to micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives, wherein the particles have a volume-weighted mean particle size of less than 5 μm as measured by laser diffractometry.

58. A method for treating a subject infected with an organism normally considered to be resistant to an antimicrobicidal agent, the method comprising the step of administering the agent to the subject, wherein the agent is formulated as an aqueous suspension of submicron- to

micron-size particles containing the agent coated with at least one surfactant selected from the group consisting of: ionic surfactants, non-ionic surfactants, biologically derived surfactants, and amino acids and their derivatives, wherein the particles have a volume-weighted mean particle size of less than 5 μm as measured by laser diffractometry.




FIG. 1

2/10




FIG. 2




FIG. 3




FIG. 4





FIG. 5

FIG. 6



7/10







FIG. 7

FIG. 8






FIG. 9

Mortality/Morbidity Profile after Daily or Everyother Day Dosing with Antifungal Drugs for 10 Days in Rats Systemically Infected with ITC Resistant *C. albicans*




FIG. 10

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2004/013268

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K9/14 A61K9/51 A61K31/496

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2003/072807 A1 (PAPADOPOULOS PAVLOS ET AL) 17 April 2003 (2003-04-17) claim 10	1-58
X	US 2003/077329 A1 (REBECK CHRISTINE L ET AL) 24 April 2003 (2003-04-24) figures 2-7; example 1	1-58
X	WO 01/21154 A (RTP PHARMA INC) 29 March 2001 (2001-03-29) claims 1-128	1-58
X	WO 02/055059 A (BAXTER INT) 18 July 2002 (2002-07-18) figure 1; example 1	1-58
X	WO 99/61001 A (RTP PHARMA INC) 2 December 1999 (1999-12-02) examples 1-4	54

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

29 July 2004

06/08/2004

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

ESTANOL, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/013268

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
US 2003072807	A1	17-04-2003	US 2003096013 A1	22-05-2003
			US 2003044433 A1	06-03-2003
			US 2003031719 A1	13-02-2003
			US 2003003155 A1	02-01-2003
			US 2002127278 A1	12-09-2002
			WO 2004032902 A1	22-04-2004
			WO 2004054500 A2	01-07-2004
			US 2004022862 A1	05-02-2004
			CA 2431705 A1	18-07-2002
			EP 1347747 A2	01-10-2003
			NO 20032860 A	21-08-2003
			WO 02055059 A2	18-07-2002
			US 2003100568 A1	29-05-2003
			US 2003206959 A9	06-11-2003
			US 2002168402 A1	14-11-2002
US 2002176935 A1	28-11-2002			
US 2003077329	A1	24-04-2003	CA 2463313 A1	01-05-2003
			EP 1435909 A1	14-07-2004
			WO 03035031 A1	01-05-2003
WO 0121154	A	29-03-2001	AU 7984200 A	24-04-2001
			CA 2383233 A1	29-03-2001
			CN 1384734 T	11-12-2002
			EP 1214059 A2	19-06-2002
			JP 2003509453 T	11-03-2003
			WO 0121154 A2	29-03-2001
WO 02055059	A	18-07-2002	US 2002176935 A1	28-11-2002
			US 2002127278 A1	12-09-2002
			US 2002168402 A1	14-11-2002
			US 2003003155 A1	02-01-2003
			US 2003044433 A1	06-03-2003
			US 2003206959 A9	06-11-2003
			CA 2431705 A1	18-07-2002
			EP 1347747 A2	01-10-2003
			NO 20032860 A	21-08-2003
			WO 02055059 A2	18-07-2002
			US 2003031719 A1	13-02-2003
			US 2003100568 A1	29-05-2003
			US 2003096013 A1	22-05-2003
			US 2003072807 A1	17-04-2003
			US 2004022862 A1	05-02-2004
WO 9961001	A	02-12-1999	AT 259220 T	15-02-2004
			AU 764001 B2	07-08-2003
			AU 4217599 A	13-12-1999
			CA 2333648 A1	02-12-1999
			CN 1311666 T	05-09-2001
			DE 69914742 D1	18-03-2004
			DK 1079808 T3	07-06-2004
			EP 1079808 A1	07-03-2001
			JP 2002516267 T	04-06-2002
			SE 0004312 A	18-01-2001
			WO 9961001 A1	02-12-1999