

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number  
**WO 01/84929 A1**

(51) International Patent Classification<sup>7</sup>: **A01N 33/02**,  
C07C 215/00, 255/00

(74) Agent: **MYBECK, Richard, R.**; P.O. Box 5540, Scottsdale, AZ 85261-5540 (US).

(21) International Application Number: PCT/US01/14790

(81) Designated States (*national*): CA, JP, US.

(22) International Filing Date: 8 May 2001 (08.05.2001)

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/202,770 9 May 2000 (09.05.2000) US

**Declarations under Rule 4.17:**

- of inventorship (Rule 4.17(iv)) for US only
- of inventorship (Rule 4.17(iv)) for US only

(71) Applicant (*for all designated States except US*): **ARIZONA BOARD OF REGENTS, acting for and on behalf of ARIZONA STATE UNIVERSITY** [US/US]; Tempe, AZ 85287 (US).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **PETTIT, George, R.** [US/US]; 6232 Bret Hills Drive, Paradise Valley, AZ 85253 (US). **LIPPERT, John, W., III** [US/US]; 509 West Carmen, Tempe, AZ 85283 (US).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/84929 A1

(54) Title: ANTITUBULIN ASSEMBLY AND CELL GROWTH INHIBITOR DENOMINATED "DIOXOSTATIN"



(I)

(57) Abstract: A new inhibitor of microtubule assembly (IC<sub>50</sub> 0.59  $\mu$ M); with antineoplastic properties, denominated "dioxostatin", has been synthesized and its effectiveness against human cancer and murine P388 lymphocytic leukemia cell lines demonstrated. Dioxostatin has the following structure (I).

-1-

Antitubulin Assembly And Cell Growth Inhibitor Denominated "Dioxostatin"

## Introduction

This invention relates generally to the field of chemotherapy and more particularly to the recovery and structural elucidation of a new inhibitor of microtubule assembly ( $IC_{50}$  0.59 $\mu$ M) and cell growth denominated "dioxostatin".


This research was funded in part by the Outstanding Investigation Grant CA 44344-01-11 awarded by the Division of Cancer Research Treatment, National Cancer Institute, DHHS. Accordingly, the United States government may have certain rights to this invention.

## Background Of The Invention


The elucidation and isolation of agents from the African bush willow *combretum caffrum* first identified combretastatin A-4 as described in U.S. Patent No. 4,996,237 which issued to G.R. Pettit et al., on February 26, 1991. Other early efforts to develop a combretastatin A-4 prodrug are described in U.S. Patent No. 5,561,122, which issued to G.R. Pettit on October 1, 1996. The general background information from each of these patents is incorporated herein by this reference thereto.

The potent cancer cell growth and tubulin assembly inhibitor combretastatin A-4 was originally isolated from the African tree *Combretum caffrum* (Combretaceae) circa 1985 and has been undergoing preclinical development since that time. However, because of the very limited aqueous solubility of the phenol and its alkali metal salts, drug formulation attempts gave unsatisfactory results.


The African willow tree *Combretum caffrum* Kuntze (Combretaceae) has proven to be a very productive source of cancer cell growth (murine P388 lymphocytic leukemia) inhibitory stilbenes, bibenzyls and phenanthrenes. Since 1979 promising leads have been pursued which were focused on the three most active (inhibition of cancer cell growth and polymerization of tubulin (*Id.*)) constituents, namely combretastatin A-1 (1a) and A-4 (1b) (*Id.*) (See, Figure 1). Of these, combretastatin A-4 (1b) has reached the most advanced stage of preclinical development as the very soluble prodrug 1d. Meanwhile other research groups have also been further extending structure/activity relationships (hereinafter referred to as "SAR") among the combretastatins and related stilbenes.




1a,  $R_1=R_2=OH$   
**combretastatin A-1**  
 b,  $R_1=H, R_2=OH$   
**combretastatin A-4**  
 c,  $R_1=R_2=OP(O)(ONa)_2$   
**combretastatin A-1 prodrug**  
 d,  $R_1=H, R_2=OP(O)(ONa)_2$   
**combretastatin A-4 prodrug**




2a,  $R_1=R_2=TBDMS$   
 b,  $R_1=R_2=OH$   
**(E)-combretastatin A-1**  
 c,  $R_1=H, R_2=TBDMS$   
 d,  $R_1=H, R_2=OH$   
**(E)-combretastatin A-4**




3a,  $R=H$   
**(R,R)-combretastatin A-4 diol**  
 b,  $R=OH$   
**(R,R)-combretastatin A-1 diol**



4a,  $R=H$   
**(S,S)-combretastatin A-4 diol**  
 b,  $R=OH$   
**(S,S)-combretastatin A-1 diol**



(S,S)-9a,  $R=R_1=OH$   
 (S,S)- b,  $R=R_1=H$   
 (S,S)- c,  $R=H, R_1=OH$



(R,R)-10a,  $R=R_1=OH$   
 (R,R)- b,  $R=R_1=H$   
 (R,R)- c,  $R=H, R_1=OH$

Figure 1

In 1987 and 1989 respectively, combretastatin A-1 (**1a**) and A-4 (**1b**) were reported which structurally resembled the antimitotic and cell growth inhibitors colchicine and podophyllotoxin, and which represent two of the most promising antineoplastic constituents of *C. caffrum*. The monophenol **1b** proved to be a potent  
5 inhibitor of microtubule assembly ( $IC_{50}$  2-3  $\mu$ M) and exhibited an  $ED_{50}$  of 0.0003  $\mu$ g/mL against the murine P388 leukemia cell line, while the diphenol **1a**, while equally effective as an inhibitor of microtubule assembly ( $IC_{50}$  2-3  $\mu$ M), was much less potent as an inhibitor of P388 cell growth ( $ED_{50}$  0.2  $\mu$ g/mL).

Presumably the reduced cytotoxicity to cancer cells shown by the diphenol **1a**  
10 is due to the presence of its vicinal phenolic groups, which can be easily oxidized to the 1,2-quinone. Such an interpretation was supported by the fact that acetylation of the hydroxyl groups enhanced cytotoxicity 10-fold, while significantly reducing the in vitro inhibition of tubulin polymerization and colchicine binding. In spite of these superficial negative aspects, some biological properties exhibited by diphenol **1a** make  
15 it attractive. For example diphenol **1a** may be the most potent antagonist of colchicine binding known, with nearly a 99% inhibition at equal concentrations. In addition, diphenol **1a** was found to be more potent than monophenol **1b** in its ability to increase intracellular daunorubicin concentrations in MDR (multidrug resistant) cancer cell  
20 lines. Most importantly, tubulin-binding stilbenes **1a** and **1b** selectively elicit irreversible vascular shutdown within solid tumors. The degree of reduction ranged from 50% with **1a** to 70% with **1b**, while the combretastatin A-4 prodrug (**1d**), a sodium phosphate derivative of monophenol **1b**, induced a complete vascular shutdown within metastatic tumors at doses one-tenth of the maximum tolerated dose. Those  
25 very encouraging results indicated that the A-series of combretastatin stilbenes should be further investigated in respect to their antiangiogenic and other anticancer properties. Hence a structure- activity relationship (SAR) study of diphenol **1a** was initiated to parallel a long-term investigation of the combretastatin A-4 prodrug (**1d**).

The formation of new blood vessels, known as "angiogenesis" (neovascularisation), is controlled by a very complex series of biochemical interactions  
30 involving a large number of angiogenic factors ranging from various cytokines (e.g. IL-1) and growth factors (e.g., GM-CSF) to serine proteases (e.g. urokinase). In general, normal angiogenesis involves the activation and transport of endothelial cells

from already formed blood vessels to new locations. Normally that transition takes from three months to a year except in wound healing and in certain stages of female reproductive biology. When the angiogenic control mechanisms fail, the results lead to a wide range of human disease categories, such as cancer psoriasis, hemangioma,  
5 atherosclerotic plaque, diabetic and macular retinopathy, neovascular glaucoma, and vascular adhesions following surgery.

Because the African bush willow (*Combretum caffrum*) constituents combretastatins A-1 (1a) and A-4 (1b) were isolated and designated as well as their phosphate prodrug derivatives (1c,d) which displayed very promising antineoplastic,  
10 cancer antiangiogenesis, and other important biological activities. Recently extended SAR investigations of these *cis*-stilbenes have been conducted. Indeed, combretastatin A-4 prodrug (1d) has been undergoing a series of phase I human cancer clinical trials since November 1998.

Previous SAR analyses of the combretastatin A-4 series have indicated that  
15 the *cis* configuration of the stilbene unit is the most important factor for inhibition of cancer cell growth and inhibiting tubulin assembly. With the corresponding (*E*)-stilbenes, the cancer cell growth inhibitory and antitubulin activity is greatly reduced from that exhibited by the corresponding (*Z*)-isomers. Initially, both the *trans*-isomers 2b and 2d were found to be moderately active as cancer cell growth inhibitors. Later  
20 studies using *trans*-stilbene 2d revealed that freshly prepared solutions in dimethyl sulfoxide were inactive and only gained activity with the passage of time suggesting that the *trans*-isomers were slowly converted to the *cis* active isomer.

#### Brief Summary of the Invention

A high yield regioselective synthesis of *E*-combretastatin A-1 (2b) was  
25 completed using MOM protection and a Wadsworth-Emmons reaction as key steps. In turn, (*E*)-stilbene 15 was converted by convenient syntheses to both (*S,S*)- and (*R,R*)-1,3-dioxolanes 9a and 10a. A Sharpless asymmetric dihydroxylation reaction was employed for preparation of intermediates (*S,S*)-16 and (*R,R*)-17. The (4*S*,5*S*)-4-(2',3'-dihydroxy-4'-methoxyphenyl)-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane (9a) was  
30 found to be a highly potent inhibitor of microtubule assembly (IC<sub>50</sub> 0.59 μM) and was designated "dioxostatin". Conversion to sodium phosphate 21 (P388 lymphocytic leukemia cell line ED<sub>50</sub> 0.2 μg/mL) provided a very useful water-soluble prodrug.

Accordingly, a principal object of the present invention is to provide a new and unexpectedly active inhibitor of microtubule assembly and cell growth by the synthesis and structure determination of a derivative of combretastatin herein denominated "dioxostatin".

5        Another object of the present invention is the formulation of a water soluble chemotherapeutic agent which is effective in the treatment of leukemia and other cancerous disorders associateable positive results against the NCI P388 cell line.


These and still further objects as shall hereinafter appear are readily fulfilled by the present invention in a remarkably unexpected manner as will be readily  
10        discerned from the following detailed description of an exemplary embodiment thereof especially when read in conjunction with the accompanying schemes.

#### Description Of The Preferred Embodiment

Recently, the (*E*)-isomer of combretastatin A-4 (**2d**) was converted to chiral diols (*R,R*)-**3a** and (*S,S*)-**4a**, where the (*E*)-olefin unit was replaced by a freely rotating  
15        *sp*<sup>3</sup>-hybridized chiral ethanediol. These experiments have now been extended to include the (*E*)-isomer of combretastatin A-1 generating chiral diols (*R,R*)-**3b** and (*S,S*)-**4b**. Asymmetric dihydroxylation (See: Kolb, H.C.; VanNieuwenhze, M.S.; Sharpless, K.B. Chem. Rev. 1994, 94, 2483-2547); with AD-mix- $\alpha$  afforded (*S,S*)-diol  
5. Photoisomerization of the (*E*)-stilbene was prevented using an aluminum foil  
20        wrapped reaction vessel.

As a check that the correct chirality was being induced in the preparation of diols (*R,R*)-**3b** and (*S,S*)-**4b**, a single crystal X-ray structure determination was conducted on one of the intermediate silyl protected products formed in the AD-mix reactions. Thus, treatment of the *trans*-stilbene **2a** with AD-mix- $\alpha$  yielded diol **5**,  
25        which was expected to have 1*S*,2*S* absolute stereochemistry. X-ray structure determination of this product (which occurred as the hemihydrate in the solid state) confirmed the expected 1*S*,2*S* stereochemistry. A computer-generated perspective of the silylated diol **5** is shown in Figure 2, below. Although the absolute stereochemistry of **5** was certain (Flack  $\chi$  absolute structure parameter = 0.1080 with esd = 0.0492), it  
30        was apparent that considerable disorder was present in the solid state of this compound, as evidenced by multiple splitting occurrences for a number of the atoms present in the unit cell. This resulted in the rather poor standard crystallographic


residual ( $R_1$  of 0.1496) obtained for this structure determination. No attempt was made to further refine the molecules over the multiple occupancy sites, since our main objective of confirming the absolute stereochemistry had been met.



5      **Figure 2.** Crystal structure (excluding hydrogens) of (*S,S*)-diol (**5**).

Protected diol (*S,S*)-**5** was next desilylated with tetrabutylammonium fluoride (TBAF) to obtain the tetraol (*S,S*)-diol **4b** in moderate yield (52%). Reaction of silyl ether (*S,S*)-diol **5** with acetic anhydride afforded (*S,S*)-diacetate **7**. Selective  
10 desilylation of (*S,S*)-diacetate **7** with TBAF proved to be difficult, yielding several products and suggesting concurrent oxidation. Other methods for desilylation (See: Sinhababu, A.K.; Kawase, M.; Borchardt, R.T. *Synthesis* 1988, 710-711; Nelson, T.D.; Crouch, D. *Synthesis* 1996, 1031-1069) were investigated, but none were able to displace TBAF as the desilylation agent of choice based on its efficient reaction time  
15 (*ca.* 20 min). The (*R,R*)-diols silyl ether **6**, diacetate **8**, and tetraol **3b** were prepared analogously using AD-mix- $\beta$ . The relevant structures are illustrated in Scheme 1.

Scheme 1



Reagents and conditions: (a) AD-mix-β,  $\text{CH}_3\text{SO}_2\text{NH}_2$ ,  $t\text{-BuOH-H}_2\text{O}$ ,  $0^\circ\text{C}$ ; (b) AD-mix-α,  $\text{CH}_3\text{SO}_2\text{NH}_2$ ,  $t\text{-BuOH-H}_2\text{O}$ ,  $0^\circ\text{C}$ ; (c) TBAF, THF; (d)  $\text{Ac}_2\text{O}$ , pyridine, DMAP, DMF.

A comparison of the diols generated from (*E*)-olefin **2a** revealed a decrease in cancer cell growth inhibition from that of combretastatin A-1 (**1a**), which is consistent with results obtained from the parallel set of diols prepared from the (*E*)-isomer of combretastatin A-4. Interesting to note is the fact that (*S,S*)-diol **4a**, a derivative of the *trans*-isomer of combretastatin A-4, is significantly more active than (*R,R*)-diol **3a** in the cancer cell lines employed (Table 1). In the combretastatin A-1 series, on the other hand, the difference noted in cancer cell line inhibition was negligible, and both (*S,S*)-diol **4b** and (*R,R*)-diol **3b** were substantially less active than (*S,S*)-diol **4a** from the A-4 series.



**Table I. Human Cancer and Murine P388 Lymphocytic Leukemia Cell Line Inhibitory Activity of Dioxostatin and Related Substances.**

Compound	Leukemia P388 ED <sub>50</sub> μg/mL	Pancreas-a BXPC-3	Neuroblast SK-N-SH	Thyroid ca SW1736	Lung-NSC NCI-H460	Pharynx-sq FADU GI <sub>50</sub> μg/mL
(R,R)-3a	19.0	>10	1.5	6.1	4.4	2.1
(S,S)-4a	2.0	3.9	0.28	1.8	2.5	0.60
(R,R)-3b	3.9	>10	3.1	>10	7.9	4.9
(S,S)-4b	2.9	>10	3.8	>10	>10	>10
(S,S)-9a	1.5	5.4	-	-	>10	-
(R,R)-10a	1.6	>10	-	-	>10	-
(S,S)-21	0.2	>10	-	-	3.4	-

## 5 Table I. (Continued)

Compound	Prostate DU-145	Ovarian OVCAR-3	CNS SF-295	Renal A498	Colon KM20L2
(R,R)-3a	3.5	4.5	5.7	>10	8.2
(S,S)-4a	2.1	0.38	0.52	0.91	0.52
(R,R)-3b	>10	7.2	2.6	>10	7.0
(S,S)-4b	>10	>10	2.9	>10	>10
(S,S)-9a	3.0	4.3	2.9	-	5.0
(R,R)-10a	>10	>10	5.0	-	>10
(S,S)-21	2.9	2.6	3.2	-	>10

**Table II. Inhibition of tubulin polymerization by diols 4-7 and combretastatin A-4 (1b)**

Compound	Inhibition of tubulin Polymerization IC <sub>50</sub> (μM) ± SD	Inhibition of colchicine binding Inhibitor: colchicine % inhibition ± SD	
1a	1.1 ± 0.07	0.2:1 89 ± 3 81 ± 4	1:1 99.6 ± 0.7 98 ± 1

	<b>1b</b>	$1.0 \pm 0.05$
	<b>(<i>R,R</i>)-3a</b>	$> 40$
	<b>(<i>S,S</i>)-4a</b>	$14 \pm 2$
	<b>(<i>R,R</i>)-3b</b>	$20 \pm 2$
5	<b>(<i>S,S</i>)-4b</b>	$10 \pm 2$

---

**Footnote for Table II**

- 10 In the polymerization experiments, the tubulin concentration was 10  $\mu$ M.  
 In the colchicine binding experiments, the tubulin concentration was 1.0  $\mu$ M,  
 and the [3H] colchicine concentration was 5.0  $\mu$ M. See: Verdier-Pinard, P.;  
 Lai, J-Y.; Yoo, H-D.; Yu, J.; Marquez, B.; Nagle, D.G.; Nambu, M.; White,  
 J.D.; Falck, J.R.; Gerwick, W.H.; Day, B.W.; Hamel, E. Mol. Pharmacol.  
 15 1998, 53, 62-76 for further details. All data with **1a** were obtained with  
 solutions prepared immediately before the experiments were performed.

The data presented in Table II, above, demonstrate that all four diols are much  
 less potent than **1a** and **1b** as inhibitors of tubulin polymerization, in agreement with  
 their much reduced cytotoxicity towards human cancer cell lines. The apparent  
 20 preference of tubulin for (*S,S*)-diols **4a** and **4b** versus the analogous (*R,R*)-diols **3a** and  
**3b** is clear. The (*S,S*)-enantiomer is at least twice as inhibitory as the (*R,R*)-  
 enantiomer. Specifically addressing the question of the importance of the second  
 hydroxyl group in the combretastatin B-ring, the results with both the (*R,R*) pair and  
 the (*S,S*) pair support the concept that the strength of the combretastatin interaction  
 25 with tubulin is enhanced by the presence of the additional hydroxyl moiety. Curiously,  
 the addition of this hydroxyl enhances the interaction with tubulin of the (*R,R*)-diol  
 somewhat more (**3b** at least twice as active as **3a**) than of the (*S,S*)-diol (**4b** only 40%  
 more active than **4a**). The greater difference between **3a** and **4a** than between **3b** and  
**4b** in their interactions with tubulin may also account for the greater difference  
 30 between the former pair than the latter in the cytotoxicity results. However, relative  
 affinities of the four diols for tubulin does not account for the observation that **4a** is the  
 most cytotoxic of the four compounds.

In addition, the data in Table II confirm the previously observed relative  
 activities of **1a** and **1b**. **1a** appears to be more active as an inhibitor of colchicine  
 35 binding.

Against microbe panels, the (*E*)-isomer of combretastatin A-1 was the most  
 active compound, presumably due to its photoisomerization to the active *cis*-isomer

(Table III). The difference in antimicrobial activity between parent compounds **1a** and **1b** and their diol derivatives was unremarkable (Table II).

**Table III.** Antimicrobial activity of combretastatin A-1, A-4, B-1 and synthetic modifications

Compound	Microbe (s) initiated	Minimum inhibitory Concentration (µg/disk)
<b>1a</b>	<i>Staphylococcus aureus</i>	50-100
	<i>Neisseria gonorrhoeae</i>	25-50
<b>1b</b>	<i>N. gonorrhoeae</i>	25-50
	<i>N. gonorrhoeae</i>	50-100
	<i>N. gonorrhoeae</i>	12.5-25
<b>2b</b>	<i>S. aureus</i>	50-100
	<i>Streptococcus pneumoniae</i>	50-100
	<i>N. gonorrhoeae</i>	6.25-12.5
	<i>Stenotrophomonas maltophilia</i>	50-100
	<i>Candida albicans</i>	50-100
	<i>Cryptococcus neoformans</i>	50-100
<b>2d</b>	*	
<b>(R,R)-3a</b>	<i>S. pneumoniae</i>	50-100
<b>(S,S)-4a</b>	<i>S. pneumoniae</i>	25-50
<b>(R,R)-3b</b>	*	
<b>(S,S)-4b</b>	<i>N. gonorrhoeae</i>	25-50


\* at 100 µg/disk, no inhibition of the ten bacteria and fungi tested

Further structural modifications aimed at converting the cancer cell line inactive (*E*)-isomers **2a** and **2c** into more active compounds was extended as follows. Diols **3b** and **4b**, synthesized from *trans*-stilbene **2a**, were further transformed by modification of the aliphatic hydroxyl groups in order to restrict free rotation. The methylenedioxy derivative (*cf.* **9** and **10**) was the rigid-ring of choice owing to its sterically small and relatively stable nature. While exploring the chemistry of stilbene **2a**, during previous evaluations of diphenol **1a** derivatives, the *tert*-butyldimethylsilyl (TBDMS) group became a liability owing to oxidative decomposition of the diphenol following desilylation with tetrabutylammonium fluoride (TBAF). Isolation of the resulting desilylated diphenols proved difficult and resulted in low yields. Therefore, the methoxymethyl (MOM) group was employed in a resynthesis of the corresponding

(*E*)-isomer of 1a, owing to its general stability (See: Yardley, J. P.; Fletcher, H. III. Introduction of the Methoxymethyl Ether Protecting Group. Synthesis 1975, 244.), and easy cleavage by dilute acid.

- MOM protection of benzaldehyde 11 was accomplished in high yield with
- 5 MOMCl. (See: Yardley, J. P.; Fletcher, H. III. Introduction of the Methoxymethyl Ether Protecting Group. Synthesis 1975, 244.). A Wadsworth-Emmons; (See: Baker, R.; Sims, R. J. A Convenient Modification of the Wadsworth-Emmons Method of Olefin Formation Using a Crown Ether. Synthesis 1981, 117.), reaction was used in preference to the conventional Wittig reaction to favor formation of the (*E*)-stilbene.
- 10 Thus, phosphonate 14 was synthesized by reaction of benzyl bromide 13 with triethyl phosphite. Condensation of protected aldehyde 12 and phosphonate 14 in the presence of sodium hydride and 15-crown-5 afforded (*E*)-stilbene 15 in 70% yield. Asymmetric dihydroxylation of (*E*)-olefin 15 with AD-mix- $\alpha$  afforded the chiral diol (*S,S*)-16. The chirality of this diol was established by its relationship to the corresponding TBDMS
- 15 derivative when the X-ray crystal structure determination was completed and was further supported by optical rotation data. (See: Figure 2). Synthesis of (*S,S*)-1,3-dioxolane 18 (See: Yardley, J. P.; Fletcher, H. III. Introduction of the Methoxymethyl Ether Protecting Group. Synthesis 1975, 244.), was conducted using 50% aqueous sodium hydroxide, dibromomethane and dichloromethane in the presence of a phase
- 20 transfer catalyst. (See: Kim, K. S.; Sarek, W. A. Methylenation of Carbohydrates using Phase -Transfer Catalysis. Synthesis 1978, 48-50. Norman, D. G.; Reese, C. B.; Serafinowska, H. T. 2',3'-*O*-Methylene Derivatives of Ribonucleosides. Synthesis 1985, 751-754.). Removal of the MOM ether was accomplished by stirring 18 with
- 25 (*S,S*)-1,3-dioxolane 9a, later designated dioxostatin. The (*R,R*)-1,3-dioxolane 10a was synthesized analogously (15 $\rightarrow$ 19) using AD-mix- $\beta$ . The relevant structures are illustrated below in Scheme 2.

Scheme 2



Biological evaluation of dioxolanes (*S,S*)-1,3-dioxolane **9a** and (*R,R*)-1,3-dioxolane **10a** (Table I) using the P388 lymphocytic leukemia cell line showed that they had similar levels of activity ( $ED_{50}$  1.5  $\mu$ g/mL and  $ED_{50}$  1.6  $\mu$ g/mL, respectively) in this system. However, the human tumor cell line data consistently showed (*S,S*)-1,3-dioxolane **9a** to be the more active, suggesting that the (*S,S*)-chirality present in dioxolane **9a** was more favorable than that in its enantiomer, **10a**. The biological potential of (*S,S*)-1,3-dioxolane **9a** was more fully appreciated when the inhibition of tubulin assembly was evaluated because (*S,S*)-1,3-dioxolane **9a** was 10-fold more potent than (*R,R*)-1,3-dioxolane **10a** ( $0.59 \pm 0.1$  and  $5.7 \pm 0.8 \mu$ M, respectively, Table IV). In fact, although less cytotoxic than combretastatin A-1 (**1a**) and A-4 (**1b**), **9a** was twice as potent as the (*Z*)-stilbenes as an inhibitor of tubulin assembly. Its activity exceeded that of colchicine and was comparable to that of the potent antimitotic peptide dolastatin 10 (Table IV).

Table IV. Inhibition of Tubulin Assembly by Anticancer Drugs Acting on Areas Near the Colchicine or Vinca Sites<sup>a</sup>

Compound	Origin	IC <sub>50</sub> ( $\mu$ M $\pm$ SD) <sup>a</sup>	% Inhibition Colchicine binding $\pm$ SD ( $\mu$ M Inhibitor)
<b>1a</b>	Terrestrial Plant	$1.1 \pm 0.07$	$89 \pm 3$ (1); $99.6 \pm 0.7$ (5)
<b>1b</b>	Terrestrial Plant	$1.0 \pm 0.05$	$81 \pm 4$ (1); $98 \pm 1$ (5)
( <i>S,S</i> )- <b>9a</b>	Synthetic	$0.59 \pm 0.1$	$67 \pm 6$ (1); $91 \pm 1$ (5)
( <i>R,R</i> )- <b>10a</b>	Synthetic	$5.7 \pm 1$	$14 \pm 5$ (5); $57 \pm 0.8$ (50)

<sup>a</sup> SD, standard deviation.

Recently, dioxolanes **9b,c** and **10b,c** in the combretastatin series were synthesized and biologically evaluated by Shirai and co-workers. Initial investigations with these conformationally restricted combretastatin analogs dealt with the synthesis of 1,3-dioxolanes lacking a phenolic group on the aryl B-ring [(*S,S*)-**9b** and (*R,R*)-**10b**], which was earlier not considered essential for biological activity. The (*S,S*)-1,3-dioxolane **9b** was found to inhibit porcine brain tubulin polymerization with an IC<sub>50</sub> of


18  $\mu\text{M}$  while (*R,R*)-1,3-dioxolane **10b** was inactive exhibiting an  $\text{IC}_{50}$  of  $>100 \mu\text{M}$ . Eventually, Shirai and co-workers evaluated the series of 1,3-dioxolanes in which the 3-position on the B aryl ring contained the phenolic group [(*S,S*)-**9c** and (*R,R*)-**10c**]. They found that (*S,S*)-1,3-dioxolane **9c** showed even greater potency than did (*S,S*)-  
5 1,3-dioxolane **9b** with an  $\text{IC}_{50}$  of 4-6  $\mu\text{M}$ , while (*R,R*)-1,3-dioxolane **10c** was again found to be inactive exhibiting an  $\text{IC}_{50}$  of  $>50 \mu\text{M}$ . These data indicated that the (*S,S*)-1,3-dioxolane group present in **9b** and **9c** assumes a favorable conformation for tubulin binding while the (*R,R*)-1,3-dioxolanes **10a** and **10b** do not. Furthermore, the most active constituent, (*S,S*)-1,3-dioxolane **9c** derived from combretastatin A-4,  
10 inhibited the *in vitro* growth of PC-6 and PC-12 (multidrug resistant) cancer cell lines at a  $\text{GI}_{50}$  of 0.002  $\mu\text{g/mL}$ .

The greater potency exhibited by dioxostatin (**9a**) when compared to both (*S,S*)-1,3-dioxolanes **9b** and **9c** indicates that the vicinal hydroxyls at the 2-and 3-position of the B ring are essential for potent inhibition of tubulin polymerization. The  
15 apparently greater activity of **9a** relative to **9c** with tubulin seems to magnify the small but reproducible differences generally observed when combretastatin A-1 and combretastatin A-4 are compared (Table IV). (In most direct comparisons, the two (*Z*)-stilbenes have comparable effects on tubulin assembly, but combretastatin A-1 is slightly more potent as an inhibitor of colchicine binding). Also of interest is the fact  
20 that (*R,R*)-1,3-dioxolane **10a** exhibits activity nominally similar to that of (*S,S*)-1,3-dioxolane **9c** ( $\text{IC}_{50}$  of 5.7  $\mu\text{M}$  and  $\text{IC}_{50}$  of 4-6  $\mu\text{M}$  respectively, however, different reaction conditions and protein preparations were used), which further supports the conclusion that the vicinal hydroxyl unit at the 2-and 3-positions of the B-ring enhance the interaction of these compounds with tubulin.

25 The microtubule system of eukaryotic cells is an important target for the development of anticancer agents. Since dioxostatin (**9a**) represents a major advance in inhibition of tubulin assembly it became a target for preclinical development. Toward that objective, the synthesis of (*S,S*)-1,3-dioxolane prodrug **21** was initiated. Diphosphorylation (See: Pettit, G.R.; Rhodes, M.R. *Antineoplastic Agents* 389). New  
30 Synthesis of Combretastatin A-4 Prodrug. *Anti-Cancer Drug Des.* 1998, 13, 183-191; Pettit, G.R.; and Lippert, J.W. III. *Antineoplastic Agents* 429. Synthesis of Combretastatin A-1 and Combretastatin B-1 Prodrugs. *Anti-Cancer Drug Des.* 2000,

-15-

(accepted); and Silverberg, L. J.; Dillon, J. L.; and Vemisheti, P. A Simple, Rapid and Efficient Protocol for the Selective Phosphorylation of Phenols with Dibenzyl Phosphite. *Tetrahedron Lett.* 1996, 37, 771-774), of dioxostatin (**9a**) was performed with dibenzyl phosphite to create diphosphate **20**. Hydrogenolysis of the benzyl groups under a slight positive pressure of hydrogen followed by reaction of the diphosphoric acid with sodium methoxide in anhydrous methanol afforded the tetrasodium phosphate prodrug (*S,S*)-**21** in good yield. The diphosphate derivative was chosen owing to its biolability by phosphatases and favorable water solubility. Phosphate prodrugs have proven to be valuable improvements when deployed with certain anticancer and other types of drugs such as combretastatins A-1 and A-4, pancratistatin, tyrosine-containing peptides and etoposide. The relevant structures are illustrated below in Scheme 3.

**Scheme 3**

**Reagents and conditions:** (a)  $\text{HP(O)(OBn)}_2$ , DMAP, DIPEA,  $\text{CCl}_4$ ,  $\text{CH}_3\text{CN}$ ,  $-10^\circ\text{C}$ ; (b)  $\text{H}_2$ , 10% Pd/C, EtOH; (c)  $\text{NaOCH}_3$ , anhydrous MeOH.

The (*S,S*)-1,3-dioxolane prodrug (**21**) was found to exhibit an  $\text{ED}_{50}$  of 0.2  $\mu\text{g/mL}$  against the murine P388 lymphocytic leukemia cell line, an approximate 10-



fold increase in activity over the parent compound, dioxostatin **9a**. Also, diphosphate **21** was shown to have an aqueous solubility of >35 mg/mL. Prodrug **21** was inactive as an inhibitor of tubulin assembly, and has been the case with all other phosphorylated derivatives in this family of compounds.

## 5 Experimental Section

**General Experimental Procedures.** Ethyl acetate (EtOAc), hexane, and dichloromethane (DCM) were redistilled. Diisopropylethylamine (DIPEA) 4-dimethylaminopyridine (DMAP), *tert*-butyl alcohol, and dibromomethane were from the Fisher-Acros Chemical Company, and all other reagents including methoxymethyl chloride (MOMCl) were from the Sigma-Aldrich Chemical Company. Solvent  
10 extracts of aqueous solutions were dried over anhydrous sodium sulfate. Column chromatography was performed using either flash silica gel (230-400 mesh ASTM) or gravity silica gel (70-230 mesh ASTM). Analtech silica gel GHLF plates were used for TLC, and compounds were visible with fluorescent short-wave light (254 nm).

15 Melting points were determined using an Electrothermal 9100 apparatus and are uncorrected. The <sup>1</sup>H NMR spectra were prepared employing a Varian VXR-300 MHz or Unity 500 MHz instrument and referenced to TMS as the internal standard. Mass spectral data were recorded using a Varian MAT 312 instrument (EIMS), or Vestec Lasertec Research mass spectrometer incorporating a Laser Sciences nitrogen  
20 laser (337 nm light pulses of 3 ns duration with 4-hydroxybenzylidenemalononitrile as the matrix and cytochrome c as the external standard for calibration purposes, (TOFMS). (See: Brune, D.C.; Cheung, J. 4-Hydroxybenzylidenemalononitrile: A Useful matrix for MALDI of Water-Insoluble Analytes. Proc. 45<sup>th</sup> ASMS Conference on Mass Spectrometry and Allied Topics, Palm Springs, California, June 1-5, 1997).

25 The IR spectra were obtained with a Matteson Instruments 2020 Galaxy Series FTIR instrument. UV spectra were obtained employing a Hitachi U-2000 UV/VIS spectrophotometer. Optical rotation values were recorded employing a Perkin Elmer 241 polarimeter. Elemental analyses were determined by Galbraith Laboratories, Inc., Knoxville, TN.

30 (1*R*,2*R*)-1,2-Dihydroxy-1-[2',3'-di(*tert*-butyldimethylsilyloxy)-4'-methoxyphenyl]-2-(3'',4'',5''-trimethoxyphenyl)-ethane (**6**). To a 100-mL foil-wrapped round bottom flask, containing a magnetic stir bar, was added distilled water (14 mL), *tert*-butyl

-17-

alcohol (14 mL), AD-mix- $\beta$  (3.8 g; 1.40 g per mmol), and methanesulfonamide (0.27 g; 2.85 mmol). Before being cooled to 0 °C, the mixture was stirred at room temperature for several min. Some of the compounds began to precipitate and 2',3'-bis(*tert*-butyldimethylsilyloxy)-3,4,4',5-tetramethoxy-(*E*)-stilbene (**2a**) (See: Pettit, G.R.; Lippert, III, J.W.; Pettit, R.K. Anticancer Drug Design in preparation) (1.4 g; 2.54 mmol) was added at once, and the slurry was stirred vigorously at 0 °C for 8 hours and at room temperature for 24 hours. Sodium sulfite (3.9 g; 30.9 mmol; 12 eq) was added and the mixture stirred for an additional 0.5 hours. The product was extracted with EtOAc (4 x 50 mL), and the combined organic phase was washed with 2N KOH (aq) and dried. Evaporation of the solvent *in vacuo* produced an off-yellow solid that was absorbed onto silica gel and subjected to flash column chromatography (eluent gradient: 9:1 to 1:1 hexane-EtOAc) to afford a clear oil, which crystallized from hexane as a colorless fluffy solid (0.94 g; 74% based on 0.23 g of **2a** recovered): m.p. 122-123 °C;  $R_f$  0.50 (1:1, hexane-EtOAc);  $[\alpha]_D^{25} +46^\circ$  (c 1.2, CHCl<sub>3</sub>); IR (nujol)  $\nu_{max}$  3526, 1595, 1251, 1124, 1097, 831, 781 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  -0.16 (3H, s, SiCH<sub>3</sub>), -0.050 (3H, s, SiCH<sub>3</sub>), 0.040 (3H, s, SiCH<sub>3</sub>), 0.17 (3H, s, SiCH<sub>3</sub>), 0.89 (9H, s, C<sub>4</sub>H<sub>9</sub>), 1.01 (9H, s, C<sub>4</sub>H<sub>9</sub>), 2.45 (1H, d,  $J$  = 1.7 Hz, OH, D<sub>2</sub>O exchanged), 2.89 (1H, d,  $J$  = 1.2 Hz, OH, D<sub>2</sub>O exchanged), 3.72 (6H, s, 2 x OCH<sub>3</sub>), 3.75 (3H, s, OCH<sub>3</sub>), 3.76 (3H, s, OCH<sub>3</sub>), 4.66 (1H, dd,  $J$  = 7.5 Hz,  $J$  = 2.4 Hz, CH), 5.06 (1H, dd,  $J$  = 7.8 Hz,  $J$  = 3.3 Hz, CH), 6.33 (2H, s, H-2", H-6"), 6.59 (1H, d,  $J$  = 8.1 Hz, H-5'), 7.05 (1H, d,  $J$  = 8.7 Hz, H-6'); EIMS  $m/z$  594 ( $M^+$ , 5), 519 (95), 397 (100), 339 (10), 198 (95); *anal.* C 59.70%, H 8.50%, calcd for C<sub>30</sub>H<sub>50</sub>O<sub>8</sub>Si<sub>2</sub>•1/2H<sub>2</sub>O, C 59.80%, H 8.50%.

**(1*R*,2*R*)-1,2-Dihydroxy-1-(2',3'-dihydroxy-4'-methoxyphenyl)-2-(3'',4'',5''-trimethoxyphenyl)-ethane (3b).**

To a solution of (*R,R*)-diol **6** (0.24 g; 0.397 mmol) in anhydrous THF (4 mL), was added TBAF (0.84  $\mu$ L; 0.84 mmol; 2.1 eq; 1 M in THF soln) in a previously flame-dried flask. After 20 min the reaction was terminated with ice-cold 6N HCl and extracted with EtOAc (4 x 20 mL). The combined organic phase was washed with saturated NaCl (aq) and dried. Removal of the organic solvent yielded a dark oil, which was subjected to column chromatography (66:33:1 EtOAc-hexane-AcOH) to afford a light brown oil that crystallized from EtOAc-hexane as a colorless solid (75

-18-

mg; 52%); m.p. 64-66 °C;  $R_f$  0.36 (66:33:1, hexane-EtOAc-AcOH);  $[\alpha]_D^{25} +56^\circ$  (*c* 1.1, CHCl<sub>3</sub>); IR (film)  $\nu_{max}$  3425, 2939, 1593, 1454, 1327, 1234, 1145 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.70 (6H, s, 2 x OCH<sub>3</sub>), 3.79 (3H, s, OCH<sub>3</sub>), 3.80 (3H, s, OCH<sub>3</sub>), 4.67 (1H, d, *J* = 7.8 Hz, CH), 4.83 (1H, d, *J* = 7.2 Hz, CH), 6.12 (1H, d, *J* = 8.7 Hz, H-5'), 6.24 (1H, d, *J* = 9.0 Hz, H-6'), 6.33 (2H, s, H-2'', H-6''); EIMS *m/z* 348 (M<sup>+</sup>-H<sub>2</sub>O, 60), 319 (95), 196 (100), 167 (10), 153 (25); *anal.* C 57.59%, H 6.18%, calcd for C<sub>18</sub>H<sub>22</sub>O<sub>8</sub>•1/2H<sub>2</sub>O, C 57.41%, H 6.25%.

**(1*S*,2*S*)-1,2-Dihydroxy-1-[2',3'-di(*tert*-butyldimethylsilyloxy)-4'-methoxyphenyl]-2-(3'',4'',5''-trimethoxyphenyl)-ethane (5).** The asymmetric dihydroxylation reaction

conditions described above for preparation of (*R,R*)-diol 6 were applied to (*E*)-olefin 2a (1.56 g; 2.78 mmol) using AD-mix- $\alpha$  to afford (*S,S*)-diol 5 as a colorless fluffy solid (1.2 g; 75%); m.p. 122-123 °C;  $R_f$  0.50 (1:1, hexane-EtOAc);  $[\alpha]_D^{25} -46^\circ$  (*c* 1.3, CHCl<sub>3</sub>); IR (nujol)  $\nu_{max}$  3524, 2361, 1593, 1251, 1124, 1097, 954, 831 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  -0.16 (3H, s, SiCH<sub>3</sub>), -0.05 (3H, s, SiCH<sub>3</sub>), 0.04 (3H, s, SiCH<sub>3</sub>), 0.17 (3H, s, SiCH<sub>3</sub>), 0.89 (9H, s, C<sub>4</sub>H<sub>9</sub>), 1.01 (9H, s, C<sub>4</sub>H<sub>9</sub>), 2.44 (1H, d, *J* = 3.3 Hz, OH, D<sub>2</sub>O exchanged), 2.87 (1H, d, *J* = 1.2 Hz, OH, D<sub>2</sub>O exchanged), 3.72 (6H, s, 2 x OCH<sub>3</sub>), 3.75 (3H, s, OCH<sub>3</sub>), 3.76 (3H, s, OCH<sub>3</sub>), 4.66 (1H, dd, *J* = 7.8 Hz, *J* = 2.1 Hz, CH), 5.06 (1H, dd, *J* = 7.5 Hz, *J* = 3.3 Hz, CH), 6.33 (2H, s, H-2'', H-6''), 6.59 (1H, d, *J* = 8.7 Hz, H-5'), 7.06 (1H, d, *J* = 8.4 Hz, H-6'); EIMS *m/z* 594 (M<sup>+</sup>, 10), 519 (95), 397 (100), 339 (20), 198 (100); *anal.* C 59.70%, H 8.50%, calcd for C<sub>30</sub>H<sub>50</sub>O<sub>8</sub>Si<sub>2</sub>•1/2H<sub>2</sub>O, C 59.66%, H 8.61%.

**X-Ray Crystal Structure of (1*S*,2*S*)-1,2-dihydroxy-1-[2',3'-bis(*tert*-butyldimethyl-silanyloxy)-4'-methoxy-phenyl]-2-(3'',4'',5''-trimethoxy-phenyl)-ethane hemihydrate (5).** A thin plate, 0.40 x 0.40 x 0.06 mm, obtained from a methanol-

hexane solution, was mounted on the tip of a glass fiber with Super Glue. Data collection was performed at 27 °C for a monoclinic system, with all reflections corresponding to slightly more than a complete quadrant (2 $\theta$  < 130°) being measured using an  $\omega/2\theta$  scan technique. Friedel reflections were also collected, whenever possible, immediately after each original reflection. Subsequent statistical analysis of the complete reflection data set using the XPREP (See: *SHELXTL-PC, Version 5.1*, 1997, an integrated software system for the determination of crystal structures from diffraction data, Bruker Analytical X-ray Systems, Inc., Madison, WI 53719. This

package includes, among others: XPREP, an automatic space group determination program; XS, the Bruker SHELXS module for the solution of X-ray crystal structures from diffraction data; XL, the Bruker SHELXL module for structure refinement; XP, the Bruker interactive graphics display module for crystal structures) program

5 indicated the space group was P2<sub>1</sub>. Each asymmetric unit of the cell was found to contain four independent molecules of the parent molecule, as well as two molecules of water. Crystal data: C<sub>30</sub>H<sub>50</sub>O<sub>8</sub> • 1/2 H<sub>2</sub>O, a = 13.303(2), b = 32.475(7), c = 19.960(4) Å, V = 7179(3) Å<sup>3</sup>, λ (Cu Kα) = 1.54178 Å, ρ<sub>c</sub> = 1.117 g cm<sup>-3</sup> for Z = 8 and F.W. = 603.89, F(000) = 2616. After Lorentz and polarization corrections, merging of

10 equivalent reflections and rejection of systematic absences, 19017 unique reflections (R(int) = 0.1195) remained, of which 13433 were considered observed (I<sub>o</sub> > 2σ(I<sub>o</sub>)) and were used in the subsequent structure solution and refinement. Linear and anisotropic decay corrections were applied to the intensity data as well as an empirical absorption correction (based on a series of psi-scans), (See: Lin, C.M.; Singh, S.B.; chu, P.S.;

15 Dempcy, R.O.; Schmidt, J.M.; Pettit, G.R.; Hamel, E. Interactions of Tubulin with Potent Natural and Synthetic Analogs of the Antimitotic Agent Combretastatin: A Structure-Activity Study. Mol Pharmacol. 1998, 34, 200-208; Pettit, G.R.; Singh, S.B.; Schmidt, J.M. Isolation, Structure and Synthesis of Combretastatins B-3 and B-4 from *Combretum caffrum*. J. Nat. Prod. 1988, 51, 517-527; Nandy, P.; Samitendu, B.;

20 Gao, H.; Hui, M.B.V.; Lien, E.J. Quantitative Structure-Activity Relationship Analysis of combretastatins: A Class of Novel antimitotic agents. Pharm. Res. 1991, 8, 776-781; Roberson, R.W.; Tucker, B.; Pettit, G.R. Microtubule depolymerization in *Uromyces appendiculatus* by Three New Antineoplastic Drugs: Combretastatin A-4, Dolastatin 10, and Halichendrin B. Mycol. Res. 1998, 102, 378-382; Iwasakik, S.

25 Antimitotic Agents: Chemistry and Recognition of Tubulin Molecule. Medicinal Research Rev. 1993, 13, 183-198). Structure determination was accomplished with SHELXS (See: *SHELXTL-PC, Version 5.1*, 1997, an integrated software system for the determination of crystal structures from diffraction data, Bruker Analytical X-ray Systems, Inc., Madison, WI 53719. This package includes, among others: XPREP, an

30 automatic space group determination program; XS, the Bruker SHELXS module for the solution of X-ray crystal structures from diffraction data; XL, the Bruker SHELXL module for structure refinement; XP, the Bruker interactive graphics display

-20-

module for crystal structures). All non-hydrogen atoms for **5**, including the water solvate atoms, were located using the default settings of that program. The remaining hydrogen atom positions were calculated at optimum positions. The latter atoms were assigned thermal parameters equal to 1.2 or 1.5 (depending upon structural atom type) the Uiso value of the atom to which they were attached and then both coordinates and thermal values were forced to ride that atom during final cycles of refinement. All non-hydrogen atoms were refined anisotropically in a full-matrix least-squares refinement process with SHELXL (See: *SHELXTL-PC, Version 5.1, 1997*, an integrated software system for the determination of crystal structures from diffraction data, Bruker Analytical X-ray Systems, Inc., Madison, WI 53719. This package includes, among others: XPREP, an automatic space group determination program; XS, the Bruker SHELXS module for the solution of X-ray crystal structures from diffraction data; XL, the Bruker SHELXL module for structure refinement; XP, the Bruker interactive graphics display module for crystal structures) in the SHELXTL-PC software package. The final standard residual R value for the model shown in Figure 1 was 0.1484 for observed data (13433 reflections) and 0.1850 for all data (19017 reflections). The corresponding Sheldrick R values were  $wR_2$  of 0.3706 and 0.4031, respectively. The Flack absolute structure parameter  $\chi$  was 0.05 (5) for the model depicted in Figure 1, thus indicating that the absolute stereochemistry shown (and expected) for **5** is correct, i.e., 1*S*, 2*S*. A final difference Fourier map showed residual electron density attributed solely to the silicon atoms; the largest difference peak and hole being 0.61 and -0.55 e/Å<sup>3</sup>, respectively. Final bond distances and angles were all within acceptable limits.

**(1*S*,2*S*)-1,2-Dihydroxy-1-(2',3'-dihydroxy-4'-methoxyphenyl)-2-(3'',4'',5''-trimethoxyphenyl)-ethane (4b).** Silyl ether (*S,S*)-diol **5** (0.36 g; 0.610 mmol) was desilylated as described above for the synthesis of (*R,R*)-diol **3b** and afforded (*S,S*)-diol **4b** as a colorless solid (0.13 g; 59%): m.p. 64-66 °C;  $R_f$  0.36 (66:33:1 hexane-EtOAc-AcOH);  $[\alpha]_D^{25}$  -52° (c 0.99, CHCl<sub>3</sub>); IR (film)  $\nu_{max}$  3418, 2939, 1593, 1510, 1462, 1327, 1290, 1234, 1124 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.73 (6H, s, 2 x OCH<sub>3</sub>), 3.81 (3H, s, OCH<sub>3</sub>), 3.83 (3H, s, OCH<sub>3</sub>), 4.70 (1H, d,  $J$  = 7.2 Hz, CH), 4.88 (1H, d,  $J$  = 7.2 Hz, CH), 6.15 (1H, d,  $J$  = 8.1 Hz, H-5'), 6.27 (1H, d,  $J$  = 8.1 Hz, H-6'), 6.37 (2H, s, H-2'', H-6''); EIMS  $m/z$  348 (M<sup>+</sup>-H<sub>2</sub>O, 20), 319 (15), 196 (100), 168 (20), 153

- (10); *anal.* C 57.59%, H 6.18%, calcd for  $C_{18}H_{22}O_8 \cdot 1/2H_2O$ , C 57.61%, H 6.28%.  
**(1*R*,2*R*)-1,2-Diacetoxy-1-[2',3'-di-(*tert*-butyldimethylsilyloxy)-4'-methoxyphenyl]-2-(3'',4'',5''-trimethoxyphenyl)-ethane (8).** To a solution of (*R,R*)-diol 6 (0.21 g; 0.351 mmol) in anhydrous DCM (2 mL) was added acetic anhydride (26 mL; 2.7 mmol; 7.8 eq), pyridine (0.20 mL; 2.35 mmol; 6.7 eq), and a catalytic amount of DMAP (5 mg). After 3 hours, the reaction was terminated with ice-water and extracted with EtOAc (4 x 25 mL). The combined organic phase was washed with 2*N* HCl followed by 10%  $NaHCO_3$  (aq) and dried. Removal (reduced pressure) of solvent yielded a clear oil that was separated by flash column chromatography (1:1 hexane-EtOAc) to afford a crude product, which crystallized from hexane as a colorless solid (0.24 g; 99%); m.p. 128-129 °C;  $R_f$  0.63 (2:1, hexane-EtOAc);  $[\alpha]_D^{25}$  -22° (*c* 1.4,  $CHCl_3$ ); EIMS  $m/z$  678 ( $M^+$ , 5), 621 (40), 439 (60), 397 (100), 73 (30); *anal.* C 60.14%, H 8.01%, calcd for  $C_{34}H_{54}O_{10}Si_2$ , C 60.38%, H 8.09%.
- (1*S*,2*S*)-1,2-Diacetoxy-[2',3'-di-(*tert*-butyldimethylsilyloxy)-4'-methoxyphenyl]-2-(3'',4'',5''-trimethoxyphenyl)-ethane (7).** The silyl ether (*S,S*)-diol 5 (0.25 g; 0.42 mmol) was acylated as described above for the synthesis of (*R,R*)-diacetate 8 to afford a colorless solid (0.28 g; 98%); m.p. 128-129 °C;  $R_f$  0.63 (2:1, hexane-EtOAc);  $[\alpha]_D^{25}$  +20° (*c* 1.3,  $CHCl_3$ ); TOFMS  $m/z$  717, ( $M+K$ )<sup>+</sup>; *anal.* C 60.14%, H 8.01%, calcd for  $C_{34}H_{54}O_{10}Si_2$ , C 60.08%, H 8.08%.
- 4-Methoxy-2,3-di(methoxymethoxy)-benzaldehyde (12).** DIPEA (7.9 mL; 45 mmol; 3 eq) was added *via* syringe to a solution of 2,3-dihydroxy-4-methoxy-benzaldehyde (11) (2.5 g; 15 mmol) in DMF (20 mL) at 0 °C. After 5 minutes, MOMCl (2.6 g; 33 mmol; 2.5 eq) was added over 10 min and stirring was continued for 1 hour at 0 °C. The reaction was terminated with ice-water. The mixture was then extracted with EtOAc (4 x 50 mL), and the organic phase washed with 10%  $NaHCO_3$  (aq) and dried. Evaporation of the solvent (reduced pressure) yielded a brown oil, which was absorbed onto silica gel and subjected to flash column chromatography (5:1 hexane-EtOAc) to afford a clear oil, 3.5 g (90%); b.p. 164-165 °C/(0.11 mm Hg);  $R_f$  0.51 (3:1, hexane-EtOAc); EIMS  $m/z$  256 ( $M^+$ , 5%), 211 (10%), 180 (15%), 151 (10%), 46 (100%); IR (film)  $\nu_{max}$  2945, 1689, 1587, 1494, 1454, 1383, 1284, 1159, 1080, 808  $cm^{-1}$ ;  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  3.57 (3H, s,  $OCH_2OCH_3$ ), 3.61 (3H, s,  $OCH_2OCH_3$ ), 3.93 (3H, s,  $OCH_3$ ) 5.13 (2H, s,

-22-

OCH<sub>2</sub>OCH<sub>3</sub>), 5.27 (2H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 6.81 (1H, d,  $J = 8.7$  Hz, H-5), 7.66 (1H, d,  $J = 8.7$  Hz, H-6), 10.29 (1H, s, CHO). Anal. Calcd. For C<sub>12</sub>H<sub>16</sub>O<sub>6</sub>: C, 56.28%; H, 6.46%. Found: C, 56.29%; H, 6.42%.

**Diethyl 3,4,5-trimethoxybenzylphosphonate (14).** A solution of triethyl phosphite (28 mL; 163 mmol; 1 eq) in toluene (100 mL) was added to a stirred solution of 3,4,5-trimethoxybenzyl bromide (13, 43 g; 162 mmol) in toluene (100 mL), and the mixture was heated at reflux for 18 hours and stirred at RT for 6 hours. Evaporation (*in vacuo*) of the solvent resulted in a crude oil, which was distilled (34 g; 66%): b.p. 212-214 °C/(water aspirator);  $R_f$  0.14 (2:1, EtOAc-hexane); EIMS  $m/z$  318 ( $M^+$ , 100%), 303 (25%), 195 (10%), 181 (95%), 167 (20%); IR (film)  $\nu_{max}$  3462, 2982, 1591, 1504, 1454, 1334, 1244, 1124, 1026, 964 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  1.27 (6H, t,  $J = 7.2$  Hz, 2 x OCH<sub>2</sub>CH<sub>3</sub>), 3.09 (2H, d,  $J = 21$  Hz, Ph-CH<sub>2</sub>), 3.83 (3H, s, OCH<sub>3</sub>), 3.85 (6H, s, 2 x OCH<sub>3</sub>), 4.05 (4H, q,  $J = 7.2$  Hz, 2 x OCH<sub>2</sub>CH<sub>3</sub>), 6.53 (2H, d,  $J = 3.0$  Hz, H-2, H-6). Anal. Calcd. For C<sub>14</sub>H<sub>23</sub>O<sub>6</sub>P • ¼ H<sub>2</sub>O: C, 52.09%; H, 7.33%. Found: C, 52.30%; H, 7.32%.

**3,4,4',5'-Tetramethoxy-2',3'-di(methoxymethoxy)-(E)-stilbene (15).** A solution of protected aldehyde 12 (9.3 g; 36.3 mmol; 1 eq) and phosphonate 14 (12 g; 38.0 mmol; 1 eq) in anhydrous THF (50 mL) was added to a stirred slurry of NaH (1.5 g; 38.0 mmol, 60% dispersion in mineral oil; 1 eq), and 15-crown-5 (0.22 mL; 1.11 mmol; 0.03 eq) in anhydrous THF (50 mL) at 0 °C. After 12 hours of stirring, the reaction was stopped with H<sub>2</sub>O. The reaction mixture was then extracted with EtOAc (4 x 100 mL), and the organic phase washed with sat. NaCl (aq) and 10% NaHSO<sub>3</sub> (aq) and dried. Removal of the solvent under reduced pressure yielded a yellow oil, which was subjected to flash column chromatography (5:1 hexane-EtOAc) to afford a clear oil that crystallized from EtOAc-hexane (11 g; 70%): m.p. 96-97 °C;  $R_f$  0.38 (3:1, hexane-EtOAc); EIMS  $m/z$  420 ( $M^+$ , 100%), 375 (50%), 344 (100%), 329 (100%), 181 (40%); IR (film)  $\nu_{max}$  2941, 1581, 1504, 1464, 1240, 1126, 1010, 970, 927, 821 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.56 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.62 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.87 (3H, s, OCH<sub>3</sub>), 3.88 (3H, s, OCH<sub>3</sub>), 3.90 (6H, s, 2 x OCH<sub>3</sub>), 5.14 (2H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 5.18 (2H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 6.74 (2H, s, H-2, H-6), 6.74 (1H, d,  $J = 9.0$  Hz, H-5'), 6.93 (1H, d,  $J = 16$  Hz, -CH=CH-), 7.34 (1H, d,  $J = 16$  Hz, -CH=CH-), 7.34 (1H, d,  $J = 8.7$  Hz, H-6'). Anal. Calcd. For C<sub>22</sub>H<sub>28</sub>O<sub>8</sub>: C, 62.85%; H, 6.46%.

6.71%. Found: C, 62.95%; H, 6.71%.

(1*S*,2*S*)-1,2-Dihydroxy-1-[4'-methoxy-2',3'-di(methoxymethoxy)-phenyl]-2-(3'',4'',5''-trimethoxyphenyl)-ethane (16). The asymmetric dihydroxylation reaction conditions described below for obtaining (*R,R*)-diol 17 were applied to (*E*)-olefin 15 (1.4 g; 3.3 mmol) using AD-mix- $\alpha$  to afford diol (*S,S*)-diol 16 as a colorless fluffy light solid that recrystallized from hexane (1.3 g; 82%): m.p. 122-123 °C;  $R_f$  0.37 (2:1, hexane-EtOAc);  $[\alpha]_D^{25}$  -83° ( $c = 0.79$ , CHCl<sub>3</sub>); EIMS  $m/z$  436 ( $M^+ - H_2O$ , 5%), 331 (10%), 257 (10%), 197 (20%), 181 (60%); IR (film)  $\nu_{max}$  3454, 2941, 1593, 1498, 1456, 1236, 1157, 970, 927 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.53 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.56 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.73 (6H, s, 2 x OCH<sub>3</sub>), 3.78 (3H, s, OCH<sub>3</sub>), 3.81 (3H, s, OCH<sub>3</sub>), 4.66 (1H, d,  $J = 7.2$  Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 4.98 (1H, dd,  $J = 6.0$  Hz,  $J = 2.0$  Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 5.06 (2H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 5.09 (1H, d,  $J = 5.7$  Hz, CH), 5.21 (1H, d,  $J = 5.4$  Hz, CH), 6.42 (2H, s, H-2'', H-6''), 6.59 (1H, d,  $J = 8.7$  Hz, H-5'), 6.80 (1H, d,  $J = 8.7$  Hz, H-6'). Anal. Calcd. For C<sub>22</sub>H<sub>30</sub>O<sub>10</sub>: C, 58.14%; H, 6.65%. Found: C, 58.24%; H, 6.65%.

(1*R*,2*R*)-1,2-Dihydroxy-1-[4'-methoxy-2',3'-di(methoxymethoxy)phenyl]-2-(3'',4'',5''-trimethoxyphenyl)-ethane (17). To a 50 mL foil wrapped round bottom flask, equipped with a magnetic stirrer, was added distilled water (20 mL), *tert*-butyl alcohol (20 mL), AD-mix- $\beta$  (5 g; 1.40 g per mmol), and methanesulfonamide (0.34 g; 2.9 mmol). The mixture was stirred at room temperature for several min and cooled to 0 °C, whereupon some of the salts precipitated. Stilbene 15 (1.4 g; 3.3 mmol) was added at once, and the slurry stirred vigorously at 0 °C for 8 hours and room temperature for 24 hours. Sodium sulfite (3.9 g; 31 mmol; 12 eq) was added and the mixture was stirred for an additional 0.5 hours. The mixture was extracted with EtOAc (4 x 25 mL), and the combined organic extract washed with 2N KOH (aq) and dried. Evaporation *in vacuo* of the solvent yielded a straw-colored solid, that was subjected to flash column chromatography (eluent gradient: 3:1 to 1:4 hexane-EtOAc) to afford (*R,R*)-diol 17 as a colorless and fluffy solid which was recrystallized from hexane (1.4 g; 89%): m.p. 122-123 °C;  $R_f$  0.37 (2:1, hexane-EtOAc);  $[\alpha]_D^{25}$  +85° ( $c = 0.87$ , CHCl<sub>3</sub>); EIMS  $m/z$  454 ( $M^+$ , 5%), 257 (20%), 242 (20%), 198 (40%), 181 (85%); IR (film)  $\nu_{max}$  3445, 2941, 1593, 1498, 1456, 1236, 1157, 1126, 972, 927 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  3.53 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.55 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>),



- 3.73 (6H, s, 2 x OCH<sub>3</sub>), 3.78 (3H, s, OCH<sub>3</sub>), 3.82 (3H, s, OCH<sub>3</sub>), 4.66 (1H, d, *J* = 7.2 Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 4.97 (1H, dd, *J* = 7.5 Hz, *J* = 2.1 Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 5.06 (2H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 5.08 (1H, d, *J* = 5.4 Hz, CH), 5.21 (1H, d, *J* = 6.3 Hz, CH), 6.42 (2H, s, H-2", H-6"), 6.59 (1H, d, *J* = 9.0 Hz, H-5'), 6.80 (1H, d, *J* = 8.7 Hz, H-6'). Anal. Calcd. For C<sub>22</sub>H<sub>30</sub>O<sub>10</sub>: C, 58.14%; H, 6.65%. Found: C, 58.17%; H, 6.79%.
- 5 **(4*S*,5*S*)-4-[4'-Methoxy-2',3'-di(methoxymethoxy)phenyl]-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane (18).** The title compound was prepared by the same reaction conditions described for the synthesis of (*R,R*)-1,3-dioxolane 19, but (*S,S*)-diol 16 (2.1 g; 4.7 mmol) was used to yield (*S,S*)-1,3-dioxolane 18 as a clear oil (1.8 g; 79%); R<sub>f</sub> 0.54 (2:1, hexane-EtOAc); [α]<sub>D</sub><sup>25</sup> -100° (c = 0.46, CHCl<sub>3</sub>); TOFMS *m/z* 489 (M + Na)<sup>+</sup>; IR (film) ν<sub>max</sub> 2941, 1593, 1498, 1460, 1421, 1157, 1128, 1089, 968, 925 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.31 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.55 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.82 (9H, s, 3 x OCH<sub>3</sub>), 3.87 (3H, s, OCH<sub>3</sub>), 4.86 (1H, d, *J* = 6.0 Hz, CH), 4.91 (1H, d, *J* = 5.4 Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 5.02 (1H, d, *J* = 5.4 Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 5.07 (1H, d, *J* = 6.0 Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 5.10 (1H, d, *J* = 5.4 Hz, OCH<sub>2</sub>OCH<sub>3</sub>), 5.19 (1H, d, *J* = 6.6 Hz, CH), 5.39 (1H, s, CH bridgehead), 5.44 (1H, s, CH bridgehead), 6.57 (2H, s, H-2", H-6"), 6.78 (1H, d, *J* = 8.7 Hz, H-5'), 7.24 (1H, d, *J* = 9.0 Hz, H-6'). Anal. Calcd. For C<sub>23</sub>H<sub>30</sub>O<sub>10</sub>: C, 59.22%; H, 6.48%. Found: C, 59.30%; H, 6.31%.
- 10 **(4*R*,5*R*)-4-[4'-Methoxy-2',3'-di(methoxymethoxy)phenyl]-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane (19).** A solution of NaOH (9.2 g; 230 mmol; 75 eq) in H<sub>2</sub>O (10 mL) was added over a period of 10 min to a stirred solution of (*R,R*)-diol 17 (1.4 g; 3.1 mmol), dibromomethane (6.5 mL; 92 mmol; 30 eq), cetyltrimethylammonium bromide (0.11 g; 0.29 mmol; 0.09 eq) in CH<sub>2</sub>Cl<sub>2</sub> (50 mL), under gentle reflux. After 4.5 hours the light yellow solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 x 25 mL), and the combined extract washed with H<sub>2</sub>O and dried. The solvent was concentrated *in vacuo* to yield a yellow oil, that was separated by flash column chromatography (eluent 2:1 hexane-EtOAc) to provide dioxolane 19 as a clear oil (0.98 g; 69%); R<sub>f</sub> 0.51 (2:1, hexane-EtOAc); [α]<sub>D</sub><sup>25</sup> +91° (c = 0.63, CHCl<sub>3</sub>); TOFMS *m/z* 488 (M + Na)<sup>+</sup>; IR (film) ν<sub>max</sub> 2941, 1593, 1498, 1456, 1419, 1157, 1126, 1087, 966, 923 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.31 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.55 (3H, s, OCH<sub>2</sub>OCH<sub>3</sub>), 3.82 (9H, s, 3 x OCH<sub>3</sub>), 3.87 (3H, s, OCH<sub>3</sub>), 4.86 (1H, d, *J*
- 15
- 20
- 25
- 30

= 6.0 Hz, CH), 4.91 (1H, d,  $J = 5.4$  Hz,  $\text{OCH}_2\text{OCH}_3$ ), 5.02 (1H, d,  $J = 5.4$  Hz,  $\text{OCH}_2\text{OCH}_3$ ), 5.07 (1H, d,  $J = 6.0$  Hz,  $\text{OCH}_2\text{OCH}_3$ ), 5.10 (1H, d,  $J = 5.4$  Hz,  $\text{OCH}_2\text{OCH}_3$ ), 5.19 (1H, d,  $J = 6.6$  Hz, CH), 5.39 (1H, s, CH bridgehead), 5.44 (1H, s, CH bridgehead), 6.58 (2H, s, H-2", H-6"), 6.79 (1H, d,  $J = 8.7$  Hz, H-5'), 7.24 (1H, d,  $J = 9.0$  Hz, H-6'). Anal. Calcd. For  $\text{C}_{23}\text{H}_{30}\text{O}_{10}$ : C, 59.22%; H, 6.48%. Found: C, 59.19%; H, 6.60%.

**(4*S*,5*S*)-4-(2',3'-Dihydroxy-4'-methoxyphenyl)-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane [dioxostatin] (9a).** To a stirred solution of dimethoxymethyl ether 18 (1.7 g; 3.64 mmol) in THF (10 mL) and IPA (4 mL) was added 6N HCL (6 mL; 36 mmol; 9 eq). After 16 hours the light yellow mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (4 x 20 mL), and the combined organic extract washed with brine and dried. The solvent was concentrated *in vacuo* to yield a light brown oil that was subjected to column chromatography [gravity column (eluent 49:50:1 hexane-EtOAc-AcOH)] to afford a colorless fluffy solid which was recrystallized from EtOAc-hexane (0.82 g; 59%); m.p. 151-152 °C;  $R_f$  0.84 (66:33:1, EtOAc-hexane-AcOH);  $[\alpha]_D^{25}$  -68° ( $c = 1.16$ ,  $\text{CHCl}_3$ ); TOFMS  $m/z$  399 ( $\text{M} + \text{Na}$ )<sup>+</sup>; IR (film)  $\nu_{\text{max}}$  3452, 3288, 1593, 1510, 1464, 1327, 1292, 1236, 1124, 1093  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  3.81 (6H, s, 2 x  $\text{OCH}_3$ ), 3.84 (3H, s,  $\text{OCH}_3$ ), 3.89 (3H, s,  $\text{OCH}_3$ ), 4.91 (2H, s, 2 x CH), 5.40 (1H, s, CH bridgehead), 5.46 (1H, brs, OH,  $\text{D}_2\text{O}$  exchanged), 5.49 (1H, s, CH bridgehead), 6.46 (1H, d,  $J = 8.1$  Hz, H-5'), 6.47 (1H, brs, OH,  $\text{D}_2\text{O}$  exchanged), 6.53 (2H, s, H-2", H-6"), 6.60 (1H, d,  $J = 9.0$  Hz, H-6'). Anal. Calcd. For  $\text{C}_{19}\text{H}_{22}\text{O}_8 \cdot \frac{1}{2}\text{H}_2\text{O}$ : C, 59.60%; H, 5.92%. Found: C, 59.81%; H, 6.23%.

**(4*R*,5*R*)-4-(2',3'-Dihydroxy-4'-methoxyphenyl)-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane (10a).** Dimethoxymethyl ether 19 (0.90 g; 1.9 mmol) was deprotected (as described above for obtaining diphenol 9a) to yield 10a as a colorless fluffy solid (0.29 g; 40%); m.p. 151-152 °C;  $R_f$  0.89 (66:33:1, EtOAc-hexane-AcOH);  $[\alpha]_D^{25}$  +71° ( $c = 1.13$ ,  $\text{CHCl}_3$ ); TOFMS  $m/z$  399 ( $\text{M} + \text{Na}$ )<sup>+</sup>; IR (film)  $\nu_{\text{max}}$  3447, 3284, 1593, 1510, 1464, 1329, 1296, 1236, 1126, 1095  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  3.81 (6H, s, 2 x  $\text{OCH}_3$ ), 3.84 (3H, s,  $\text{OCH}_3$ ), 3.89 (3H, s,  $\text{OCH}_3$ ), 4.91 (2H, s, 2 x CH), 5.39 (1H, s, CH bridgehead), 5.46 (1H, brs, OH,  $\text{D}_2\text{O}$  exchanged), 5.49 (1H, s, CH bridgehead), 6.45 (1H, d,  $J = 8.1$  Hz, H-5'), 6.47 (1H, brs, OH,  $\text{D}_2\text{O}$  exchanged), 6.53

(2H, s, H-2", H-6"), 6.60 (1H, d,  $J = 9.0$  Hz, H-6'). Anal. Calcd. For  $C_{19}H_{22}O_8$ : C, 60.31%; H, 5.86%. Found: C, 59.96%; H, 6.15%.

**(4*S*,5*S*)-4-(2',3'-*O*-Di[bis(benzyl)phosphoryl]-4'-methoxyphenyl)-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane (20).** A solution of diphenol **9a** (0.2 g; 0.53 mmol) in acetonitrile (10 mL) was cooled to  $-20$  °C. Carbon tetrachloride (0.6 mL; 5.3 mmol; 10 eq) was added. The resulting solution was stirred for 10 min before the addition of DIPEA (0.4 mL; 2.2 mmol; 4.2 eq, *via* syringe) and DMAP (15 mg; 0.11 mmol; 0.2 eq). Approximately 1 min later, slow (dropwise) addition of dibenzyl phosphite (0.34 mL; 1.5 mmol; 2.9 eq) was begun at such a rate that the reaction temperature was kept below  $-20$  °C. After 45 min of stirring, 0.5 M  $KH_2PO_4$  (aq) was added and the reaction mixture was allowed to warm to room temperature. The mixture was extracted with EtOAc (4 x 15 mL), and the combined organic extract was successively washed with saturated NaCl (aq), water, and dried. Removal of the solvent under reduced pressure afforded a yellow oil that was separated by flash column chromatography (3:2 hexane-EtOAc) to afford 0.43 g (90%) of a clear oil:  $R_f$  0.24 (1:1, hexane-EtOAc);  $[\alpha]_D^{25} -55^\circ$  ( $c = 0.83$ ,  $CHCl_3$ ); EIMS  $m/z$  898 (5%,  $M^+$ ), 778 (10%), 546 (5%), 181 (20%), 91 (100%); IR (film)  $\nu_{max}$  2944, 1593, 1506, 1458, 1296, 1126, 1099, 1014, 898, 740  $cm^{-1}$ ;  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  3.70 (3H, s,  $OCH_3$ ), 3.76 (6H, s, 2 x  $OCH_3$ ), 3.81 (3H, s,  $OCH_3$ ), 4.75-5.11 (8H, m, 4 x  $CH_2$ -Ph), 5.16 (1H, d,  $J = 6.6$  Hz, CH), 5.29 (1H, s, CH bridgehead), 5.41 (1H, s, CH bridgehead), 5.50 (1H, d,  $J = 6.6$  Hz, CH), 6.57 (2H, s, H-2", H-6"), 6.91 (1H, d,  $J = 8.7$  Hz, H-5'), 7.43 (1H, d,  $J = 9.3$  Hz, H-6'), 7.07-7.25 (15H, m, 4 x  $C_6H_5$ ), 7.30 (5H, m,  $C_6H_5$ );  $^{13}C$  NMR (125 MHz,  $CDCl_3$ )  $\delta$  153.29, 152.13, 137.53, 135.74, 135.35, 133.54, 128.36, 127.84, 127.69, 127.61, 124.48, 123.65, 110.06, 102.94, 95.90, 84.69, 78.73, 77.62, 70.04, 69.67, 60.61, 56.24, 56.01;  $^{31}P$  NMR (202 MHz,  $CDCl_3$ )  $\delta$  -4.79, -4.86. Anal. Calcd. For  $C_{47}H_{48}O_{14}P_2$ : C, 62.81%; H, 5.38%. Found: C, 62.72%; H, 5.34%.

**Tetrasodium-(4*S*,5*S*)-4-(2',3'-*O*-diphosphonate-4'-methoxyphenyl)-5-(3'',4'',5''-trimethoxyphenyl)-1,3-dioxolane (21).** To a solution of tetrabenzyl diphosphate **20** (0.43 g; 0.45 mmol) in EtOH (5 mL in a hydrogenation flask) was added 10% Pd/C (0.14 g). The flask was evacuated and exposed (balloon) to a hydrogen atmosphere for 2 hours. Filtration of the solution through celite and subsequent evaporation (*in*

-27-

*vacuo*) of solvent afforded a clear oil. Anhydrous MeOH (5 mL) was added to the crude phosphoric acid followed by NaOCH<sub>3</sub> (0.11 g; 1.9 mmol; 4 eq). The mixture was stirred for 12 hours, at which point additional MeOH was added until the product dissolved. Collection (filtration) of the product in MeOH and subsequent  
5 concentration *in vacuo* afforded a crude colorless solid, which was reprecipitated from H<sub>2</sub>O/EtOH to yield a colorless powder (0.22 g; 72%): m.p. 166-168 °C dec.; [ $\alpha$ ]<sub>D</sub><sup>25</sup> -68° (c=1.05, H<sub>2</sub>O); UV  $\lambda_{\text{max}}$  (H<sub>2</sub>O) 285 nm (log  $\epsilon$ , 3.17); IR (KBr)  $\nu_{\text{max}}$  3375, 2939, 1591, 1496, 1454, 1296, 1168, 1120, 1095, 985 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, D<sub>2</sub>O)  $\delta$  3.59 (3H, s, OCH<sub>3</sub>), 3.68 (3H, s, OCH<sub>3</sub>), 3.71 (6H, s, 2 x OCH<sub>3</sub>), 5.04 (1H, d, *J* = 5.5  
10 Hz, CH), 5.15 (1H, s, CH bridgehead), 5.18 (1H, s, bridgehead), 5.85 (1H, d, *J* = 4.0 Hz, CH), 6.68 (1H, d, *J* = 8.5 Hz, H-5'), 6.71 (2H, s, H-2'', H-6''), 7.03 (1H, d, *J* = 8.5 Hz, H-6'); <sup>13</sup>C NMR (125 MHz, D<sub>2</sub>O/CD<sub>3</sub>OD)  $\delta$  154.69, 153.70, 147.42, 137.69, 135.86, 124.15, 122.09, 108.26, 105.53, 95.55, 83.19, 79.75, 61.68, 57.17, 56.74. <sup>31</sup>P NMR (202 MHz, D<sub>2</sub>O)  $\delta$  1.79, 1.97. LRFAB MS *m/z* 603 [(Anion + 3Na)<sup>+</sup>, 25%],  
15 581 [(Anion + 2Na + H)<sup>+</sup>, 70%], 559 [(Anion + Na + 2H)<sup>+</sup>, 100%].

**Tubulin assays.** The tubulin polymerization and colchicine binding experiments were performed as described previously (See: Verdier-Pinard, P.; Lai, J.-Y.; Yoo, H.-D.; Yu, J.; Marquez, B.; Nagle, D.G.; Nambu, M.; White, J.D.; Falck, J.R.; Gerwick, W.H.; Day, B.W.; Hamel, E. Mol. Pharmacol. 1998, 53, 62-76).  
20 However, in the polymerization assays Beckman DU7400/7500 spectrophotometers equipped with "high performance" temperature controllers were used. Unlike the manual control possible with the previously used Gilford spectrophotometers, the Beckman instruments are microprocessor controlled, and assays required use of programs provided by MDB Analytical Associates, South Plainfield, NJ. The  
25 Beckman instruments were unable to maintain 0EC, and the lower temperature in the assays fluctuated between 2 and 4EC. Temperature changes were, however, more rapid, with the jump from the lower temperature to 30EC taking about 20 sec and the reverse jump about 100 sec.

**Antimicrobial susceptibility testing.** Compounds were screened against the  
30 bacteria *Stenotrophomonas maltophilia*, *Micrococcus luteus*, *Staphylococcus aureus*, *Escherichia coli*, *Enterobacter cloacae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Neisseria gonorrhoeae*, and the fungi *Candida albicans* and

*Cryptococcus neoformans*, according to established disk susceptibility testing protocols (See: Oshumi, K.; Nakagawa, R.; Yumiko, F.; Hatanaka, T.; Morinaga, Y.; Nihei, Y.; Obishi, D.; Suga, Y.; Akiyama, Y.; Tsuji, T. Novel Combretastatin Analogues Effective Against Murine Solid Tumors: Design and Structure-Activity Relationships. J. Med. Chem. 1998, 41, 3022-3032; Madarde, M.; Ramos, a.; Caballero, E.; Pelaez-Lamamie de Clairac, R.; Lopez, J.L.; Gravalos, D.G.; Feliciano, A.S. Synthesis and Antineoplastic Activity of Combretastatin Analogues: Heterocombretastatins. Eur. J. Med. Chem. 1998, 33, 71-77).

#### Dosages

10           The dosage administered will be dependent upon the identity of the neoplastic disease; the type of host involved, including its age, health and weight; the kind of concurrent treatment, if any; the frequency of treatment and therapeutic ratio.

          Illustratively, dosage levels of the administered active ingredients are: intravenous, 0.1 to about 200 mg/kg; intramuscular, 1 to about 500 mg/kg; orally, 5 to  
15   about 1000 mg/kg; intranasal instillation, 5 to about 1000 mg/kg; and aerosol, 5 to about 1000 mg/k of host body weight.

          Expressed in terms of concentration, an active ingredient can be present in the compositions of the present invention for localized use about the cutis, intranasally, pharyngolaryngeally, bronchially, intravaginally, rectally, or ocularly in concentration  
20   of from about 0.01 to about 50% w/w of the composition; preferably about 1 to about 20% w/w of the composition; and for parenteral use in a concentration of from about 0.05 to about 50% w/v of the composition and preferably from about 5 to about 20% w/v.

          The compositions of the present invention are preferably presented for  
25   administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, suppositories, sterile parenteral solutions or suspensions, sterile non-parenteral solutions of suspensions, and oral solutions or suspensions and the like, containing suitable quantities of an active ingredient.

          For oral administration either solid or fluid unit dosage forms can be prepared.

30           Powders are prepared quite simply by comminuting the active ingredient to a suitably fine size and mixing with a similarly comminuted diluent. The diluent can be an edible carbohydrate material such as lactose or starch. Advantageously, a

sweetening agent or sugar is present as well as a flavoring oil.

Capsules are produced by preparing a powder mixture as hereinbefore described and filling into formed gelatin sheaths. Advantageously, as an adjuvant to the filling operation, a lubricant such as talc, magnesium stearate, calcium stearate and  
5 the like is added to the powder mixture before the filling operation.

Soft gelatin capsules are prepared by machine encapsulation of a slurry of active ingredients with an acceptable vegetable oil, light liquid petrolatum or other inert oil or triglyceride.

Tablets are made by preparing a powder mixture, granulating or slugging,  
10 adding a lubricant and pressing into tablets. The powder mixture is prepared by mixing an active ingredient, suitably comminuted, with a diluent or base such as starch, lactose, kaolin, dicalcium phosphate and the like. The powder mixture can be granulated by wetting with a binder such as corn syrup, gelatin solution, methylcellulose solution or acacia mucilage and forcing through a screen. As an  
15 alternative to granulating, the powder mixture can be slugged, i.e., run through the tablet machine and the resulting imperfectly formed tablets broken into pieces (slugs). The slugs can be lubricated to prevent sticking to the tablet-forming dies by means of the addition of stearic acid, a stearic salt, talc or mineral oil. The lubricated mixture is then compressed into tablets.

20 Advantageously, the tablet can be provided with a protective coating consisting of a sealing coat or enteric coat of shellac, a coating of sugar and methylcellulose and polish coating of carnauba wax.

Fluid unit dosage forms for oral administration such as in syrups, elixirs and suspensions can be prepared wherein each teaspoonful of composition contains a  
25 predetermined amount of an active ingredient for administration. The water-soluble forms can be dissolved in an aqueous vehicle together with sugar, flavoring agents and preservatives to form a syrup. An elixir is prepared by using a hydroalcoholic vehicle with suitable sweeteners together with a flavoring agent. Suspensions can be prepared of the insoluble forms with a suitable vehicle with the aid of a suspending agent such  
30 as acacia, tragacanth, methylcellulose and the like.

For parenteral administration, fluid unit dosage forms are prepared utilizing an active ingredient and a sterile vehicle, water being preferred. The active ingredient,

-30-

depending on the form and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the water-soluble active ingredient can be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampule and sealing. Advantageously, adjuvants such as a local anesthetic, preservative and buffering agents can be dissolved in the vehicle. Parenteral suspensions are prepared in substantially the same manner except that an active ingredient is suspended in the vehicle instead of being dissolved and sterilization cannot be accomplished by filtration. The active ingredient can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the active ingredient.

In addition to oral and parenteral administration, the rectal and vaginal routes can be utilized. An active ingredient can be administered by means of a suppository. A vehicle which has a melting point at about body temperature or one that is readily soluble can be utilized. For example, cocoa butter and various polyethylene glycols (Carbowaxes) can serve as the vehicle.

For intranasal instillation, a fluid unit dosage form is prepared utilizing an active ingredient and a suitable pharmaceutical vehicle, preferably P.F. water; a dry powder can be formulated when insufflation is the administration of choice.

For use as aerosols, the active ingredients can be packaged in a pressurized aerosol container together with a gaseous or liquefied propellant, for example, dichlorodifluoromethane, carbon dioxide, nitrogen, propane, and the like, with the usual adjuvants such as cosolvents and wetting agents, as may be necessary or desirable.

The term "unit dosage form" as used in the specification and claims refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical diluent, carrier or vehicle. The specifications for the novel unit dosage forms of this invention are dictated by and are directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitation inherent in the art of compounding such an active material for therapeutic use in humans, as disclosed in this specification, these being features of the present invention.

-31-

Examples of suitable unit dosage forms in accord with this invention are tablets, capsules, troches, suppositories, powder packets, wafers, cachets, teaspoonfuls, tablespoonfuls, dropperfuls, ampules, vials, segregated multiples of any of the foregoing, and other forms as herein described.

5           The active ingredients to be employed as antineoplastic agents can be easily prepared in such unit dosage form with the employment of pharmaceutical materials which themselves are available in the art and can be prepared by established procedures. The following preparations are illustrative of the preparation of the unit dosage forms of the present invention, and not as a limitation thereof. Several dosage forms were prepared embodying the present invention. They are shown in the following examples in which the notation "active ingredient" signifies either dioxostatin (9a) and/or dioxostatin prodrug or any other compound described herein.

#### COMPOSITION "A"

##### Hard-Gelatin Capsules

15           One thousand two-piece hard gelatin capsules for oral use, each capsule containing 200 mg of an active ingredient are prepared from the following types and amounts of ingredients:

	Active ingredient, micronized	200 g
	Corn Starch	20 g
20	Talc	20 g
	Magnesium stearate	2 g

The active ingredient, finely divided by means of an air micronizer, is added to the other finely powdered ingredients, mixed thoroughly and then encapsulated in the usual manner.

25           The foregoing capsules are useful for treating a neoplastic disease by the oral administration of one or two capsules one to four times a day.

Using the procedure above, capsules are similarly prepared containing an active ingredient in 50, 250 and 500 mg amounts by substituting 50 g, 250 g and 500 g of an active ingredient for the 200 g used above.

#### 30           COMPOSITION "B"

##### Soft Gelatin Capsules

One-piece soft gelatin capsules for oral use, each containing 200 mg of an



-32-

active ingredient, finely divided by means of an air micronizer, are prepared by first suspending the compound in 0.5 ml of corn oil to render the material capsulatable and then encapsulating in the above manner.

The foregoing capsules are useful for treating a neoplastic disease by the oral administration of one or two capsules one to four times a day.

### COMPOSITION "C"

#### Tablets

One thousand tablets, each containing 200 mg of an active ingredient, are prepared from the following types and amounts of ingredients:

10	Active ingredient, micronized	200 g
	Lactose	300 g
	Corn starch	50 g
	Magnesium stearate	4 g
	Light liquid petrolatum	5 g

15 The active ingredient, finely divided by means of an air micronizer, is added to the other ingredients and then thoroughly mixed and slugged. The slugs are broken down by forcing them through a Number Sixteen screen. The resulting granules are then compressed into tablets, each tablet containing 200 mg of the active ingredient.

20 The foregoing tablets are useful for treating a neoplastic disease by the oral administration of one or two tablets one to four times a day.

Using the procedure above, tablets are similarly prepared containing an active ingredient in 250 mg and 100 mg amounts by substituting 250 g and 100 g of an active ingredient for the 200 g used above.

### COMPOSITION "D"

#### 25 Oral Suspension

One liter of an aqueous suspension for oral use, containing in each teaspoonful (5 ml) dose, 50 mg of an active ingredient, is prepared from the following types and amounts of ingredients:

	Active ingredient, micronized	10 g
30	Citric acid	2 g
	Benzoic acid	1 g
	Sucrose	790 g

-33-

Tragacanth	5 g
Lemon Oil	2 g
Deionized water, q.s.	1000 ml

5 The citric acid, benzoic acid, sucrose, tragacanth and lemon oil are dispersed in sufficient water to make 850 ml of suspension. The active ingredient, finely divided by means of an air micronizer, is stirred into the syrup unit uniformly distributed. Sufficient water is added to make 1000 ml.

The composition so prepared is useful for treating a neoplastic disease at a dose of 1 teaspoonful (15 ml) three times a day.

10 **COMPOSITION "E"**  
**Parenteral Product**

A sterile aqueous suspension for parenteral injection, containing 30 mg of an active ingredient in each milliliter for treating a neoplastic disease, is prepared from the following types and amounts of ingredients:

15	Active ingredient, micronized	30 g
	POLYSORBATE 80	5 g
	Methylparaben	2.5 g
	Propylparaben	0.17 g
	Water for injection, q.s.	1000 ml.

20 All the ingredients, except the active ingredient, are dissolved in the water and the solution sterilized by filtration. To the sterile solution is added the sterilized active ingredient, finely divided by means of an air micronizer, and the final suspension is filled into sterile vials and the vials sealed.

The composition so prepared is useful for treating a neoplastic disease at a dose of 1 milliliter (1ml) three times a day.

25 **COMPOSITION "F"**  
**Suppository, Rectal and Vaginal**

One thousand suppositories, each weighing 2.5 g and containing 200 mg of an active ingredient are prepared from the following types and amounts of ingredients:

30	Active ingredient, micronized	15 g
	Propylene glycol	150 g
	Polyethylene glycol #4000, q.s.	2,500 g

-34-

The active ingredient is finely divided by means of an air micronizer and added to the propylene glycol and the mixture passed through a colloid mill until uniformly dispersed. The polyethylene glycol is melted and the propylene glycol dispersion is added slowly with stirring. The suspension is poured into unchilled molds at 40°C.

- 5 The composition is allowed to cool and solidify and then removed from the mold and each suppository foil wrapped.

The foregoing suppositories are inserted rectally or vaginally for treating a neoplastic disease.

#### COMPOSITION "G"

##### 10 Intranasal Suspension

One liter of a sterile aqueous suspension for intranasal instillation, containing 20 mg of an active ingredient in each milliliter, is prepared from the following types and amounts of ingredients:

	Active ingredient, micronized	15 g
15	POLYSORBATE 80	5 g
	Methylparaben	2.5 g
	Propylparaben	0.17 g
	Deionized water, q.s. 1000 ml.	

- 20 All the ingredients, except the active ingredient, are dissolved in the water and the solution sterilized by filtration. To the sterile solution is added the sterilized active ingredient, finely divided by means of an air micronizer, and the final suspension is aseptically filled into sterile containers.

The composition so prepared is useful for treating a neoplastic disease, by intranasal instillation of 0.2 to 0.5 ml given one to four times per day.

- 25 An active ingredient can also be present in the undiluted pure form for use locally about the cutis, intranasally, pharyngolaryngeally, bronchially, or orally.

#### COMPOSITION "H"

##### Powder

- 30 Five grams of an active ingredient in bulk form is finely divided by means of an air micronizer. The micronized powder is placed in a shaker-type container.

The foregoing composition is useful for treating a neoplastic disease, at localized sites by applying a powder one to four times per day.

-35-

**COMPOSITION "I"****Oral Powder**

One hundred grams of an active ingredient in bulk form is finely divided by means of an air micronizer. The micronized powder is divided into individual doses of  
5 200 mg and packaged.

The foregoing powders are useful for treating a neoplastic disease, by the oral administration of one or two powders suspended in a glass of water, one to four times per day.

**COMPOSITION "J"**

10

**Insufflation**

One hundred grams of an active ingredient in bulk form is finely divided by means of an air micronizer.

The foregoing composition is useful for treating a neoplastic disease, by the inhalation of 300 mg one to four times a day.

15


From the foregoing, it becomes readily apparent that a new and useful antineoplastic factor and new and useful antineoplastic preparations and microtubule inhibitors have been herein described and illustrated which fulfill all of the aforesaid objectives in a remarkably unexpected fashion. It is of course understood that such modifications, alterations and adaptations as will readily occur to the artisan

20

confronted with this disclosure are intended within the spirit of the present invention.

**Claims:**

1. A compound denominated "dioxostatin", having the general structure set forth below:



- 5 2. A method of treating cells afflicted with a neoplastic disease comprising administering to said cells an effective amount of the active ingredient dioxostatin in a pharmacological acceptable carrier.
3. A method according to claim 2 in which said carrier comprises an aqueous solution.
- 10 4. A method of inhibiting microtubule assembly comprising exposing said microtubules to an effective amount of the active ingredient dioxostatin in a pharmacologically acceptable carrier.
5. A method according to claim 4 in which said carrier comprises an aqueous solution.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/14790

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 33/02; C07C 215/00, 255/00

US CL : 546/646, 523; 564/443; 558/402

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/646, 523; 564/443; 558/402

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
MEDLINE, CAPLUS, EMBASE, EAST

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,525,632 A (OBSUMI K et al) 11 June 1996, see the entire document.	1-5
Y, P	WO 01/19794 A2 (BAYLOR UNIVERSITY) 22 March 2001, see the entire document.	1-5
Y, P	Dtabase Medline on STN, AN 2001256019. PETTIT GR et al. 'Antineoplastic agents 442. Synthesis and biological activities of dioxostatin'. Anti-Cancer Drug Design. October 2000, Vol. 15, No. 5, pages 361-371, see the abstract	1-5
Y	Database EMBASE ON STN, AN 1998162066. PETTIT GR et al. 'Antineoplastic agents 379. Synthesis of phenstatin phosphate'. Journal of Medicinal Chemistry. 07 May 1998, Vol. 41, No. 10, pages 1688-1695, see the abstract.	1-5
Y, P	Database BIOSIS on STN, AN 2000:393158. TAHIR SK et al. 'Rapid colchicine competition-binding scintillation proximity assay using biotin-labeled tubulin'. Biotechniques. July 2000, Vol. 29, No. 1, pages 156-160, see the abstract.	1-5

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

02 July 2001 (02.07.2001)

Date of mailing of the international search report

27 SEP 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Ravi R. Shukla

Telephone No. (703) 308-0196