



HU000030883T2



(19) **HU**

(11) Lajstromszám: **E 030 883**

(13) **T2**

MAGYARORSZÁG
Szellemi Tulajdon Nemzeti Hivatala

EURÓPAI SZABADALOM **SZÖVEGÉNEK FORDÍTÁSA**

(21) Magyar ügyszám: **E 12 748046**

(22) A bejelentés napja: **2012. 06. 19.**

(96) Az európai bejelentés bejelentési száma:
EP 20120748046

(97) Az európai bejelentés közzétételi adatai:
EP 2720989 A1 2012. 12. 27.

(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2720989 B1 2016. 08. 10.

(51) Int. Cl.: **C07B 59/00**

(2006.01)

A61K 314/95

(2006.01)

C07D 241/04

(2006.01)

A61P 25/00

(2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/IB 12/001386

(87) A nemzetközi közzétételi szám:

WO 12176066

(30) Elsőbbségi adatai:

201161498651 P 2011. 06. 20. US
201161537103 P 2011. 09. 21. US

(73) Jogosult(ak):

H. Lundbeck A/S, 2500 Valby, Copenhagen
(DK)

(72) Feltaláló(k):

JORGENSEN, Morten, DK-2880 Bagsvaerd (DK)
ANDERSEN, Peter, Hongaard, DK-3520 Farum (DK)
JENSEN, Klaus, Gjervig, DK-1418 Kobenhavn K (DK)
HVENEGAARD, Mette, Graulund, DK-2680 Solrod
Strand (DK)
BADOLO, Lassina, DK-2400 Kobenhavn Nv (DK)
JACOBSEN, Mikkel, Fog, DK-2000 Frederiksberg (DK)

(74) Képviselő:

SBGK Szabadalmi Ügyvivői Iroda, Budapest

(54)

Deuterált 1-piperazino-3-fenil-indánok, skizofrénia kezelésére

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



EP 2 720 989 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
10.08.2016 Bulletin 2016/32

(51) Int Cl.:
C07B 59/00 (2006.01) **C07D 241/04 (2006.01)**
A61K 31/495 (2006.01) **A61P 25/00 (2006.01)**

(21) Application number: **12748046.5**

(86) International application number:
PCT/IB2012/001386

(22) Date of filing: **19.06.2012**

(87) International publication number:
WO 2012/176066 (27.12.2012 Gazette 2012/52)

(54) DEUTERATED 1-PIPERAZINO-3-PHENYL INDANES FOR TREATMENT OF SCHIZOPHRENIA

DEUTERIERTE 1-PIPERAZIN-3-PHENYL-INDANE ZUR BEHANDLUNG VON SCHIZOPHRENIE
1-PIPERAZINO-3-PHENYLINDANES DEUTÉRÉS POUR LE TRAITEMENT DE LA SCHIZOPHRÉNIE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

(30) Priority: **20.06.2011 US 201161498651 P
21.09.2011 US 201161537103 P**

(43) Date of publication of application:
23.04.2014 Bulletin 2014/17

(73) Proprietor: **H. Lundbeck A/S
2500 Valby, Copenhagen (DK)**

(72) Inventors:
• **JORGENSEN, Morten
DK-2880 Bagsvaerd (DK)**

- **ANDERSEN, Peter, Hongaard
DK-3520 Farum (DK)**
- **JENSEN, Klaus, Gjervig
DK-1418 København K (DK)**
- **HVENEGAARD, Mette, Graulund
DK-2680 Solrod Strand (DK)**
- **BADOLØ, Lassina
DK-2400 København Nv (DK)**
- **JACOBSEN, Mikkel, Fog
DK-2000 Frederiksberg (DK)**

(74) Representative: **H. Lundbeck A/S
Ottiliavej 9
2500 Valby (DK)**

(56) References cited:
WO-A1-93/22293 WO-A1-2005/016901

EP 2 720 989 B1


Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**FIELD OF THE INVENTION**

5 [0001] The present invention relates to deuterated 1-piperazino-3-phenyl-indanes and salts thereof with activity at dopamine D₁ and D₂ receptors as well as the serotonin 5HT₂ receptors in the central nervous system, to medicaments comprising such compounds as active ingredients, and to the use of such compounds in the treatment of diseases in the central nervous system.

10 BACKGROUND OF THE INVENTION

15 [0002] 4-((1R,3S)-6-Chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine and salts thereof, pharmaceutical compositions containing these salts and the medical use thereof, including treatment of schizophrenia or other diseases involving psychotic symptoms, are disclosed in WO2005/01690. 4-((1R,3S)-6-Chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine has the general formula (X), hereinafter referred to as Compound (X)



35 [0003] EP 638 073 recites a group of trans isomers of 3-aryl-1-(1-piperazinyl)indanes substituted in the 2- and/or 3-position of the piperazine ring. The compounds are described as having high affinity for dopamine D₁ and D₂ receptors and the 5-HT₂ receptors and are suggested to be useful for treatment of several diseases in the central nervous system, including schizophrenia.

40 [0004] The enantiomer of formula (X) above has been described by Bøgesø et al. in J. Med. Chem., 1995, 38, page 4380-4392, in the form of the fumarate salt, see table 5, compound (-)-38. This publication concludes that the (-)-enantiomer of compound 38 is a potent D₁/D₂ antagonist showing some D₁ selectivity *in vitro*. The compound is also described as a potent 5-HT₂ antagonist. It is also mentioned that the compound does not induce catalepsy in rats.

45 [0005] The aetiology of schizophrenia is not known, but the dopamine hypothesis of schizophrenia (Carlsson, Am. J. Psychiatry 1978, 135, 164-173), formulated in the early 1960s, has provided a theoretical framework for understanding the biological mechanisms underlying this disorder. In its simplest form, the dopamine hypothesis states that schizophrenia is associated with a hyperdopaminergic state, a notion which is supported by the fact that all antipsychotic drugs on the market today exert some dopamine D₂ receptor antagonism (Seeman Science and Medicine 1995, 2, 28-37).

50 [0006] However, whereas it is generally accepted that antagonism of dopamine D₂ receptors in the limbic regions of the brain plays a key role in the treatment of positive symptoms of schizophrenia, the blockade of D₂ receptors in striatal regions of the brain causes extrapyramidal symptoms (EPS). As described in EP 638 073 a profile of mixed dopamine D₁/D₂ receptor inhibition has been observed with some so-called "atypical" antipsychotic compounds, in particular with clozapine (8-chloro-11-(4-methylpiperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine), used in treatment of schizophrenic patients.

55 [0007] Further, selective D₁ antagonists have been connected to treatment of sleep disorders and alcohol abuse (D.N. Eder, Current Opinion in Investigational Drugs, 2002 3(2):284-288).

[0008] Dopamine may also play an important role in the aetiology of affective disorders (P. Willner, Brain. Res. Rev. 1983, 6, 211-224, 225-236 and 237-246; Bøgesø et al, J. Med. Chem., 1985, 28, 1817-1828).

[0009] In EP 638 073 is described how compounds having affinity for 5-HT₂ receptors, in particular 5-HT_{2A} receptor antagonists, have been suggested for treatment of different diseases, such as schizophrenia including the negative symptoms in schizophrenic patients, depression, anxiety, sleep disturbance, migraine attacks and neuroleptic-induced parkinsonism. 5-HT_{2A} receptor antagonism has also been suggested to reduce the incidence of extrapyramidal side effects induced by classical neuroleptics (Balsara et al. Psychopharmacology 1979, 62, 67-69).

[0009] An isotopic substitution of one or more hydrogen atoms (H) by deuterium atoms (D) in a compound may give rise to a kinetic isotope effect which may influence the reaction rate, e.g. metabolism of the compound. This is particularly the case when the isotopic replacement is in a chemical bond that is broken or formed in a rate limiting step. In such a case, the change is termed a primary isotope effect. When the isotopic substitution(s) are not involved in one or more bonds that are broken a smaller rate change, termed the secondary isotope effect may be observed.

5

SUMMARY OF THE INVENTION

[0010] The present invention provides compounds wherein one or more hydrogen atoms atoms (H) in one or more of 10 the metabolic sites M1, M2 and M3 of Compound (X) have been substituted by deuterium atoms (D).

10


[0011] In one aspect, the invention provides a compound of formula Y:

15

20

25

30



wherein, R¹ - R¹⁰ are independently hydrogen or deuterium, and wherein at least one of R¹-R¹⁰ comprises at least about 35 50% deuterium, or a pharmaceutically acceptable acid addition salt thereof.

[0012] In another aspect, the invention provides pharmaceutical compositions comprising a compound of formula (Y) and one or more pharmaceutically acceptable carriers, diluents, or excipients.

[0013] In another aspect, the invention provides for uses of a compound of formula (Y) or a pharmaceutical composition comprising a compound of formula (Y) in the treatment of psychosis, other diseases involving psychotic symptoms, 40 psychotic disorders or diseases that present with psychotic symptoms.


[0014] In yet another aspect, the invention provides for the manufacture of a medicament comprising a compound of formula (Y) for treatment of psychosis, other diseases involving psychotic symptoms, psychotic disorders or diseases that present with psychotic symptoms. of formula (Y) to a subject in need thereof.

[0015] In still another aspect, the invention provides a compound of formula


45

50

55



[0016] In still another aspect, the invention provides a process for the preparation of compound



comprising treating compound (XIV) with [(S)-BINAP]Rh(I)BF₄.

[0017] In still another aspect, the invention provides a process for the preparation of compound (1*R*,3*S*)-(IV) tartrate comprising, treatment of racemic *trans*-1-(6-chloro-3-phenyl(*d*₅)-indan-1-yl)-1(*d*₃), 2, 2-trimethyl-piperazine with L-(+)-tartaric acid.

[0018] Still other objects and advantages of the invention will become apparent to those of skill in the art from the disclosure herein, which is simply illustrative and not restrictive.

20 BRIEF DESCRIPTION OF THE FIGURES

[0019]

25 FIG. 1 shows major metabolic sites of Compound (X).

FIG. 2 shows Compound (I) and Compound (XI), each as the (1*R*,3*S*)-enantiomer.

FIG. 3 shows NMR spectra of Compound (II) and Compound (V). Selected regions of the proton-decoupled and proton-and deuterium-decoupled ¹³C NMR spectra of Compound (II) [Fig. 3A] and Compound (V) [Fig. 3B] are shown.

FIG. 4 shows the mass spectrum of Compound (IV).

FIG. 5 shows formation of the metabolite Compound (XI) by metabolism of Compound (X) and Compound (I) (0.1 microM) in cryopreserved dog hepatocytes (n=2 the bars represent max and min results).

FIG. 6 shows formation of the metabolite Compound (XI) by metabolism of Compound (X) and Compound (I) (1 microM) in cryopreserved dog hepatocytes (n=2 the bars represent max and min results).

FIG. 7 shows formation of the desmethyl metabolite by metabolism of Compound (II), (IV) and (X) (1 micro M) in human liver microsomes (n=3, the bars represent standard deviation).

FIG. 8 shows formation of the desmethyl metabolite by metabolism of Compound (II), (IV) and (X) (10 micro M) in human liver microsomes (n=3, the bars represent standard deviation).

FIG. 9 shows formation of the desmethyl metabolite by metabolism of Compound (III) (10 micro M) in human liver microsomes (n=3, the bars represent standard deviation).

FIG. 10 shows formation of the desmethyl metabolite by metabolism of Compound (V) (10 micro M) in human liver microsomes (n=3, the bars represent standard deviation).

FIG. 11 shows formation of the desmethyl metabolite by metabolism of Compound (VI) (10 micro M) in human liver microsomes (n=3, the bars represent standard deviation).

FIG. 12 shows formation of the desmethyl metabolite by metabolism of Compound (VII) (10 micro M) in human liver microsomes (n=3, the bars represent standard deviation).

FIG. 13 shows the chemical structure of compounds (I)-(VII), (X)-(XI), and (XIX)-(XXI).

FIG. 14 shows formation of the desmethyl metabolite by metabolism of Compound (II) and (X) (10 micro M) by recombinant human liver CYP2C19 (n=3, the standard deviation).

FIG. 15 shows formation of the desmethyl metabolite by metabolism of Compound (IV) and Compound (X) (1 micro M) by recombinant human liver CYP2C19 (n=3, the bars represent standard deviation).

FIG. 16 shows PCP-induced hyperactivity in mice for compound (IV).

FIG. 17 shows cataleptic response in rats for compound (IV).

FIG. 18 shows X-ray diffractograms on two batches of hydrogen tartrate salt of Compound (IV).

DETAILED DESCRIPTION OF THE INVENTION

[0020] Atypical antipsychotics have been the subject of numerous studies by the pharmaceutical industry, and have shown promise in treating mental disorders such as schizophrenia, bipolar disorder, dementia, anxiety disorder and obsessive-compulsive disorder (OCD). The mechanism of action of these agents remains unknown; however all antip-

syotics work to some degree on the dopamine system. Most atypical antipsychotics exhibit activity at dopamine subtype receptors 1 and 2 (D_1 and D_2 , respectively), and at the serotonin receptors subtype 2 (5-HT₂). In some cases, the "atypical" designation was assigned to antipsychotics that did not induce extrapyramidal side effects; however it has been shown that some atypical antipsychotics still induce extrapyramidal side effects, albeit to a lesser degree than that observed with typical antipsychotics (Weiden, P.J., "EPS profiles: the atypical antipsychotics are not all the same" *J. Psychiatr. Pract.* 2007, 13(1): 13-24). Approved atypical antipsychotics include, for example, amisulpride (Solian), aripiprazole (Abilify), asenapine (Saphris), blonanserin (Lonasen), clozapine (Clozaril), iloperidone (Fanapt), lurasidone (Latuda), mosapramine (Cremin), olanzapine (Zyprexa), paliperidone (Invega), perospirone (Lullan), quetiapine (Seroquel), remoxipride (Roxiam), risperidone (Risperdal), sertindole (Serdolect), supiride (Supirid, Eglonyl), ziprasidone (Geodon, Zeldox), and zotepine (Nipolept). Several others are currently under development. Because the mechanism of atypical antipsychotics is not well understood, side effects associated with these drugs have been difficult to design around. Thus, there is a need for additional antipsychotic therapies with potential for reduced side effect and/or improved therapeutic profile relative to existing therapies.

[0021] In one aspect, the present invention provides compounds wherein one or more hydrogen atoms (H) in one or more of the metabolic sites M1, M2 and M3 of Compound (X) have been substituted by deuterium atoms (D). Compound (X) and variants thereof are described in, for example U.S. Patent Nos. 5,807,855; 7,648,991; 7,767,683; 7,772,240; 8,076,342; U.S. Patent Publication Nos. 2008/0269248; 2010/0069676; 2011/0178094; 2011/0207744; WO 2005/016900; EP 0 638 073; and *J. Med. Chem.* 1995, 38, 4380-4392.

[0022] The kinetic isotope effect may potentially influence the rate of metabolism at one or more of the metabolic sites M1, M2, and M3 indicated in Figure 1. The inventors of the present invention have identified three major metabolic sites of 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine (Compound (X)) denoted herein as M1, M2 and M3 and indicated in Figure 1.

[0023] Deuteration of a compound at a site subject to oxidative metabolism may, in some cases reduce the rate of metabolism for a compound due to the primary isotope effect. If the C-H bond cleavage step is rate limiting, a significant isotope effect may be observed. However, if other steps drive the rate of metabolism for a compound, the C-H bond cleavage step is not rate limiting and the isotope effect may be of little significance. Additionally, a negative isotope effect can be observed where reaction rate is increased upon substitution with deuterium. Thus, incorporation of deuterium at a site subject to oxidative enzymatic metabolism does not predictably impact pharmacokinetics (See, for example, U.S. Pat. No. 7,678,914; *Drug Metab. Dispos.* 1986, 14, 509; *Arch. Toxicol.* 1990, 64, 109; *Int. Arch. Occup. Environ. Health* 1993, 65(Suppl. 1): S139). The impact of deuterium incorporation is unpredictable does not work for many drugs or classes of drugs. Decreased metabolic clearance has been observed with some deuterated compounds relative to non-deuterated derivatives; whereas metabolism of other compounds has been unimpacted. Examples of studies indicating lack of predictability regarding deuterium incorporation include U.S. Patent No. 6,221,335; *J. Pharm. Sci.* 1975, 64, 367-391; *Adv. Drug. Res.* 1985, 14, 1-40; *J. Med. Chem.* 1991, 34, 2871-2876; *Can. J. Physiol. Pharmacol.* 1999, 79-88; Silverman, R. B., *The Organic Chemistry of Drug Design and Drug Action*, 2nd Ed. (2004), 422; *Curr. Opin. DrugDev.* 2006, 9, 101-109; *Chemical Res. Tox.* 2008, 1672; Harbeson, S.L and Tung, R.D. "Deuterium in Drug Discovery and Development," in *Ann. Rep. Med. Chem.* 2011, 46, 404-418. Even incorporation deuterium at known sites of metabolism has an unpredictable impact on metabolic profile. Metabolic switching may result wherein the metabolic profile of a particular drug is changed due to deuterium incorporation, thus leading to different proportions of (or different) metabolites than observed with a non-deuterated analog of the same drug. The new metabolic profile may result in a distinct toxicological profile of the deuterated analog. Adding to the potential complications of deuterium incorporation is the possibility of deuterium/hydrogen exchange in the physiological environment (*Adv. Drug. Res.* 1985, 14, 1-40).

[0024] In some embodiments, isotopic substitution of one or more hydrogen atoms in Compound (X) by deuterium atoms has given rise to a kinetic isotope effect that influences the rate of metabolism.

[0025] The isotopic substitution of hydrogen atoms in Compound (X) by deuterium atoms results in less metabolism of the deuterated compound as shown to occur in dog hepatocytes where for instance an approximately 50% decrease in formation of the desmethyl metabolite (Compound (XI)) from Compound (I) (Figure 2) was noted in comparison to the formation of Compound (XI) from the metabolism of Compound (X).

[0026] Deuteration of the free phenyl, optionally in combination with deuteration of the 1-methyl group (Compound (II) and (IV)), surprisingly reduces the amount of the desmethyl metabolite produced in human liver microsomes as compared to the non-deuterated compound (Compound (X)). Also surprisingly, deuteration of the 1-methyl group impacted metabolism in dog but not human hepatocytes, thus indicative of the unpredictability of deuteration on pharmacological properties.

[0027] The effect of the reduced metabolism is higher bioavailability of the deuterated, parent compound and less metabolite formation. Without being bound by theory, based on the results described in the experimental section of this application the same effect is expected to show up after multiple dosing in humans, allowing for lower doses to be administered to humans i.e. less burden to the entire body, e.g. the liver, and a less frequent dosing.

[0028] The desmethyl metabolite (Compound (XI)) is known to have hERG affinity and thus potentially contribute to

QTc prolongation. As mentioned above, deuteration of the free phenyl optionally in combination with deuteration of the 1-methyl group (Compound (II) and (IV)), surprisingly reduces the amount of the desmethyl metabolite produced in human liver microsomes as compared to the non-deuterated compound (Compound (X)). Accordingly and without being bound by theory, it is anticipated that there will be less interaction with the hERG channel and resultant lower burden on the heart when dosing the deuterated variants of Compound (X) [e.g., compounds of formula (Y)] compared to when dosing Compound (X).

5 [0029] The invention is further detailed in the exemplary embodiments provided herein.

10 Definitions

[0030] The term "compound(s) of the invention" as used herein means Compounds (Y), (I), (II), (III), (IV), (V), (VI), and/or (VII), and may include salts, hydrates and/or solvates thereof. The compounds of the present invention are prepared in different forms, such as salts, hydrates, and/or solvates, and the invention includes compositions and methods encompassing all variant forms of the compounds.

15 [0031] The term "composition(s) of the invention" as used herein means compositions comprising Compounds (Y), (I), (II), (III), (IV), (V), (VI), and/or (VII), or salts, hydrates, and solvates thereof. The compositions of the invention may further comprise one or more chemical components such as, for example, excipients, diluents, vehicles or carriers.

[0032] The term "method(s) of the invention" as used herein means methods comprising treatment with the compounds and/or compositions of the invention.

20 [0033] As used herein the term "about" is used herein to mean approximately, roughly, around, or in the region of. When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" is used herein to modify a numerical value above and below the stated value by a variance of 20 percent up or down (higher or lower).

25 [0034] An "effective amount", "sufficient amount" or "therapeutically effective amount" as used herein is an amount of a compound that is sufficient to effect beneficial or desired results, including clinical results. As such, the effective amount may be sufficient, for example, to reduce or ameliorate the severity and/or duration of an affliction or condition, or one or more symptoms thereof, prevent the advancement of conditions related to an affliction or condition, prevent the recurrence, development, or onset of one or more symptoms associated with an affliction or condition, or enhance or otherwise improve the prophylactic or therapeutic effect(s) of another therapy. An effective amount also includes the amount of the compound that avoids or substantially attenuates undesirable side effects.

30 [0035] As used herein and as well understood in the art, "treatment" is an approach for obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results may include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminution of extent of disease, a stabilized (*i.e.*, not worsening) state of disease, preventing spread of disease, delay or slowing of disease progression, amelioration or palliation of the disease state and remission (whether partial or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival as compared to expected survival if not receiving treatment.

35 [0036] The term "in need thereof" refers to the need for symptomatic or asymptomatic relief from a condition such as, for example, psychosis or a psychotic disorder. The subject in need thereof may or may not be undergoing treatment for conditions related to, for example, psychosis or a psychotic disorder.

40 [0037] The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which a compound is administered. Non-limiting examples of such pharmaceutical carriers include liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. The pharmaceutical carriers may also be saline, gum acacia, gelatin, starch paste, talc, keratin, colloidal silica, urea, and the like. In addition, auxiliary, stabilizing, thickening, lubricating and coloring agents may be used. Other examples of suitable pharmaceutical carriers are described in Remington: The Science and Practice of Pharmacy, 21st Edition (University of the Sciences in Philadelphia, ed., Lippincott Williams & Wilkins 2005).

45 [0038] The terms "animal," "subject" and "patient" as used herein include all members of the animal kingdom including, but not limited to, mammals, animals (*e.g.*, cats, dogs, horses, swine, etc.) and humans.

50 [0039] The term "isotopic variant" as used herein means a compound obtained by substituting one or more hydrogen in a parent compound not comprising deuterium atoms by deuterium atoms.

[0040] It is recognized that elements are present in natural isotopic abundances in most synthetic compounds, and result in inherent incorporation of deuterium. However, the natural isotopic abundance of hydrogen isotopes such as deuterium is immaterial (about 0.015%) relative to the degree of stable isotopic substitution of compounds indicated herein. Thus, as used herein, designation of an atom as deuterium at a position indicates that the abundance of deuterium is significantly greater than the natural abundance of deuterium. Any atom not designated as a particular isotope is intended to represent any stable isotope of that atom, as will be apparent to the ordinarily skilled artisan.

55 [0041] Compounds (Y) are isotopic variants of Compound (X).

[0042] In some embodiments, compounds (I), (II), (III), (IV), (V), (VI) and (VII) are isotopic variants of Compound (X).

[0043] M1 is a site of Compound (X) susceptible to metabolism; M1 consists of $-\text{CH}_2-$ in the 6-position of the piperazine of Compound (X).

[0044] M2 is a site of compound (X) susceptible to metabolism; M2 consists of the N-bound methyl of the piperazine of Compound (X).

5 [0045] M3 is a site of Compound (X) susceptible to metabolism; M3 consists of the phenyl group of Compound (X).

[0046] Parent compound is the chemical compound which is the basis for its derivatives obtained either by substitution or breakdown, e.g. metabolic breakdown. In the context of the present invention the parent compound is the Active Pharmaceutical Ingredient (API).

[0047] In some embodiments, any atom not designated as deuterium is present at its natural isotopic abundance. In some embodiments, any hydrogen atom not designated as deuterium is present at less than 1% isotopic abundance of deuterium.


[0048] In one aspect, the invention provides a compound of formula (Y):

15

20

25

30



35 wherein, R¹ - R¹⁰ are independently hydrogen or deuterium, wherein at least one of R¹-R¹⁰ comprises at least about 50% deuterium, or a pharmaceutically acceptable acid addition salt thereof.

[0049] In another aspect, the invention provides pharmaceutical compositions comprising a compound of formula (Y) and one or more pharmaceutically acceptable carriers, diluents, or excipients.

[0050] In yet another aspect, the invention provides for the manufacture of a medicament comprising a compound of formula (Y) for treatment of psychosis, other diseases involving psychotic symptoms, psychotic disorders or diseases that present with psychotic symptoms.


[0051] In some embodiments, the compound is racemic. In some embodiments, the compound is enantiomerically enriched.

[0052] In some embodiments, the compound is selected from the group consisting of

45

50


55



5

10

15



[0053] In some embodiments, R¹ and R² comprise deuterium, R³-R⁵ comprise deuterium, or R⁶-R¹⁰ comprise deuterium.

[0054] In some embodiments, R¹ and R² comprise deuterium. In some embodiments, R¹ and R² comprise deuterium and R³-R⁵ comprise hydrogen.

[0055] In some embodiments, R³-R⁵ comprise deuterium. In some embodiments, R³-R⁵ comprise hydrogen.

[0056] In some embodiments, R⁶-R¹⁰ comprise deuterium. In some embodiments, R⁶-R¹⁰ comprise deuterium and R³-R⁵ comprise hydrogen.

[0057] In some embodiments, R¹-R⁵ comprise deuterium.

[0058] In some embodiments, R¹, R², and R⁶-R¹⁰ comprise deuterium.

[0059] In some embodiments, R³-R¹⁰ comprise deuterium.


[0060] In some embodiments, R¹-R¹⁰ comprise deuterium.

[0061] In some embodiments, the compound is

35

40

45



or


50

55

5

10

15

(1*R*,3*S*)-(IV).


[0062] In some embodiments, the compound is

20

25

30

35


(1*R*,3*S*)-(II).

[0063] In some embodiments, the compound is

40

45

50


(1*R*,3*S*)-(III).

55 [0064] In some embodiments, the compound is

5

10

15




[0065] In some embodiments, the compound is

20

25

30

35




[0066] In some embodiments, the compound is

40


45

50

55



[0067] In some embodiments, the compound is



[0068] In some embodiments, at least about 75% of the compound has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.

[0069] In some embodiments, at least about 85% of the compound has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.

[0070] In some embodiments, at least about 90% of the compound has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.

[0071] In some embodiments, the compound is a salt selected from the group consisting of fumarate, maleate, succinate, and tartrate. In some embodiments, the compound is a fumarate salt. In some embodiments, the compound is a hydrogen fumarate salt. In some embodiments, the compound is a maleate salt. In some embodiments, the compound is a hydrogen maleate salt. In some embodiments, the compound is a succinate salt. In some embodiments, the compound is a hydrogen succinate salt. In some embodiments, the compound is a tartrate salt. In some embodiments, the compound is the hydrogen tartrate salt.

[0072] In some embodiments, the compound is the hydrogen tartrate salt of (1*R*,3*S*)-(IV).

[0073] In some embodiments, the psychosis or disease involving psychotic symptoms is schizophrenia, schizophrenia-form disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, bipolar disorder, or mania in bipolar disorder. In some embodiments, the psychosis or disease involving psychotic symptoms is schizophrenia.

[0074] In some embodiments, the methods further comprise administration of with one or more neuroleptic agents.

[0075] In some embodiments, the uses further comprise use of a one or more neuroleptic agents.

[0076] In some embodiments, the neuroleptic agent is selected from the group consisting of sertindole, olanzapine, risperidone, quetiapine, aripiprazole, haloperidol, clozapine, ziprasidone and osanetant.

[0077] In some embodiments, administration is oral, sublingual, or buccal. In some embodiments, administration is oral.

[0078] In some embodiments, the subject is a mammal. In some embodiments, the subject is a rodent, cat, dog, monkey, horse, swine, bovine, or human. In some embodiments, the subject is a rodent, cat, dog, monkey, bovine or human. In some embodiments, the subject is a mouse, rat, cat, dog, monkey, or human. In some embodiments, the subject is a mouse, rat, dog, monkey, or human. In some embodiments, the subject is a mouse, rat, dog, or human. In some embodiments, the subject is a mouse, rat or a human. In some embodiments, the subject is a dog or a human. In some embodiments, the subject is a human.

[0079] In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 40% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 50% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 60% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 65% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 70% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 75% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 80% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 85% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 90% at that position. In some embodiments,

designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 95% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 97% at that position. In some embodiments, designation of a position as "D" in a compound has a minimum deuterium incorporation of greater than about 99% at that position.

5

Pharmaceutically Acceptable Salts

[0080] The present invention also comprises salts of the compounds, typically, pharmaceutically acceptable salts. Such salts include pharmaceutically acceptable acid addition salts. Acid addition salts include salts of inorganic acids as well as organic acids.

10

[0081] Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, sulfamic, nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, itaconic, lactic, methanesulfonic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methane sulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanesulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, theophylline acetic acids, as well as the 8-halotheophyllines, for example 8-bromotheophylline and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in Berge, S.M. et al., *J. Pharm. Sci.* 1977, 66, 2, and Gould, P.L., *Int. J. Pharmaceutics* 1986, 33, 201-217.

15

[0082] Furthermore, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered comparable to the unsolvated forms for the purposes of this invention.

20

[0083] Headings and sub-headings are used herein for convenience only, and should not be construed as limiting the invention in any way.

25

[0084] The use of any and all examples, or exemplary language (including "for instance", "for example", "e.g.", and "as such") in the present specification is intended merely to better illuminate the invention, and does not pose a limitation on the scope of invention unless otherwise indicated.

30

[0085] The use of the terms "a" and "an" and "the" and similar referents in the context of describing the invention are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context.

[0086] Unless otherwise indicated, all exact values provided herein are representative of corresponding approximate values (e.g., all exact exemplary values provided with respect to a particular factor or measurement can be considered to also provide a corresponding approximate measurement, modified by "about," where appropriate).

35

[0087] The description herein of any aspect or aspect of the invention using terms such as "comprising", "having," "including," or "containing" with reference to an element or elements is intended to provide support for a similar aspect or aspect of the invention that "consists of", "consists essentially of", or "substantially comprises" that particular element or elements, unless otherwise stated or clearly contradicted by context.

40

[0088] Exemplary syntheses of the compounds of the invention can be readily achieved by methods described, for example, U.S. Patent Nos. 5,807,855; 7,648,991; 7,767,683; 7,772,240; 8,076,342; U.S. Patent Publication Nos. 2008/0269248; 2010/0069676; 2011/0178094; 2011/0207744; WO 2005/016900; EP 0 638 073; and *J. Med. Chem.* 1995, 38, 4380-4392. Such methods, and similar methods can be performed using deuterated reagents and/or intermediates, and/or introducing deuterium atoms to a chemical structure according to protocols known in the art.


45

[0089] Further exemplary methods of synthesis include conversion of indanone A to intermediate C via treatment of 3-bromo-6-chloro-indan-1-one (A; for references on this material, see: Bøgesø EP 35363 A1 19810909 and Kehler, Juhl, Püschl, WO 2008025361) with a base such as triethylamine in a solvent such as tetrahydrofuran at ambient temperature (Scheme 1). Removal of the precipitated amine hydrobromide salt by filtration and concentration of the filtrate will afford 6-chloro-inden-1-one (B). This material can be reacted with phenyl-d₅-boronic acid in the presence of approximately 1 equivalent of a base such as triethylamine and a catalytic amount of a 1:1 mixture of [Rh(ndb)₂]BF₄ (bis(norbornadiene)rhodium(I) tetrafluoroborate) and racemic BINAP (2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) in a suitable solvent (e.g. approximately 10:1 solvent mixture of 1,4-dioxane and water) under an atmosphere of argon at elevated temperature (e.g. about 100°C). Work-up will afford racemic 6-chloro-3-phenyl-d₅-inden-1-one (C).

50

Scheme 1. Exemplary synthesis of intermediate C.

55



catalog # 495859 or from CDN catalog # D-5340 or from Kanto catalog # 49132-27) the same procedure may lead to E (Scheme 4). As alternatives to the commercial sources, benzyl cyanide-*d*₇ and phenyl-*d*₅-acetonitrile can be prepared sodium cyanide and benzyl-*d*₇ chloride (commercially available from Aldrich, catalog # 217336) and benzyl-2,3,4,5,6-*d*₅ chloride (commercially available from Aldrich, catalog # 485764), respectively.

5

Scheme 4. Exemplary synthesis of intermediates E and E'.

10

15

20

25

30


35

40

45


50

55




[0093] Treatment of E with approximately 4 equivalents of di-iso-propylethylamine and approximately 2 equivalents methanesulphonic anhydride in tetrahydrofuran at approximately - 18 °C followed by slow heating to approximately -5 °C and subsequent treatment with approximately 4 equivalents 2,2-dimethyl-piperazine will lead to the formation of 1-((1R,3S)-6-chloro-3-phenyl-*d*₅-indan-1-yl)-3,3-dimethyl-piperazine (F) that can be purified after the reaction (Scheme 5). Alternatively, one can convert alcohol E to the corresponding chloride, predominantly with retention of configuration at C1 leading to (1S,3S)-1-chloro-3-*d*₅-phenyl-indan (E'); similarly E' can be converted to (1S,3S)-1-chloro-3-phenyl-indan (E''). Chloride E'' can be reacted with 2,2-dimethyl-piperazine to afford F. The final step can be performed as described for the preparation of Compound (I)•butanedioic acid salt by the use of iodomethane to give Compound (II) or *d*₃-iodomethane to give Compound (IV), respectively. Alternatively, as described below, the methyl group or *d*₃-methyl group can be installed by refluxing in HCHO/HCOOH or DCDO/DCOOD, respectively.

Scheme 5. Exemplary synthesis of intermediates F and Compounds (II) and (IV).




[0094] (2-Amino-2-methyl-propyl)-carbamic acid *tert*-butyl ester (G) can be prepared from 2-methyl-propane-1,2-diamine and di-*tert*-butyl dicarbonate (alternatively, G is claimed to be commercially available: Prime catalog # POI-1362-MB4; Rovathin catalog # NX45401). Reaction of G with a haloacetyl halide such as either chloroacetyl chloride or bromoacetyl bromide will give [2-(2-chloro-acetylamino)-2-methyl-propyl]-carbamic acid *tert*-butyl ester or [2-(2-bromoacetylamino)-2-methyl-propyl]-carbamic acid *tert*-butyl ester (H), respectively (Scheme 6). Treatment of either variant of H with acid followed by base will lead to the formation of 6,6-dimethyl-piperazine-2-one (I). This material can be reduced to 2,2-dimethyl-5,5-*d*₂-piperazine (J) by treatment with lithium aluminium deuteride.

Scheme 6. Exemplary synthesis of intermediate J.




[0095] Alternatively, J can be prepared from 2-amino-2-methyl-propionic acid. Reaction of 2-amino-2-methyl-propionic acid and di-*tert*-butyl dicarbonate will afford 2-*tert*-butoxycarbonylamino-2-methyl-propionic acid (K) (Scheme 7). The acid functionality can be converted to the corresponding Weinreb amide by reaction with O,N-dimethyl-hydroxylamine in the presence of a suitable coupling reagent such as 2-(1*H*-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium (HATU) or 1-methyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) to afford [1-(methoxy-methyl-carbamoyl)-1-methylethyl]-carbamic acid *tert*-butyl ester (L). Selective reduction of the Weinreb amide leads to (1,1-dimethyl-2-oxo-ethyl)-carbamic acid *tert*-butyl ester (M). Reductive amination involving aldehyde M and amino-acetic acid methyl ester can be used to prepare (2-*tert*-butoxycarbonylamino-2-methyl-propylamino)-acetic acid methyl ester (N). Treatment of carbamate-ester N with a suitable acid, such as trifluoroacetic acid, will lead to the formation of piperazinone I that upon treatment with lithium aluminium deuteride gives piperazine J.

Scheme 7. Alternative exemplary synthesis of intermediate J.



[0096] Using J instead of 2,2-dimethyl-piperazine as described for the conversion of E to Compounds (II) and (IV) will lead to Compounds (VI) and Compound (VII), respectively. Similarly, using E' and J instead of 2,2-dimethyl-piperazine and E will lead to Compound (III) and Compound (V).

[0097] In another aspect, the invention provides a process for the preparation of compound



comprising treating compound (XIV) with [(S)-BINAP]Rh(I)BF₄.

[0098] In another aspect, the invention provides a process of the preparation of compound (1*R*,3*S*)-(IV) tartrate comprising treatment of racemic *trans*-1-(6-chloro-3-phenyl(*d*₅)-indan-1-yl)-1(*d*₃), 2, 2-trimethyl-piperazine with L-(+)-tartric acid.

[0099] In some embodiments, racemic *trans*-1-(6-chloro-3-phenyl(*d*₅)-indan-1-yl)-1(*d*₃), 2, 2-trimethyl-piperazine is generated from the corresponding succinate salt thereof.

[0100] In some embodiments, racemic *trans*-1-(6-chloro-3-phenyl(*d*₅)-indan-1-yl)-1(*d*₃), 2, 2-trimethyl-piperazine succinate is generated from the maleate salt of racemic *trans*-1-(6-chloro-3-phenyl(*d*₅)-indan-1-yl)-3,3-dimethyl-piperazine.

[0101] In some embodiments, acetophenone-*d*₅ is converted to an enol ether. In some embodiments, the enol ether is a silyl enol ether. In some embodiments, the enol ether of acetophenone-*d*₅ is converted to the corresponding vinyl boronate. In some embodiments, the enol ether of acetophenone-*d*₅ is treated with bis(pinacolato)diboron. In some embodiments, the vinyl boronate is treated with 2-halo-5-chlorobenzaldehyde.

[0102] In some embodiments, the compounds exist as racemates. In some embodiments, the compounds exist in greater than about 70% enantiomeric excess. In some embodiments, the compounds exist in greater than about 75% enantiomeric excess. In some embodiments, the compounds exist in greater than about 80% enantiomeric excess. In some embodiments, the compounds exist in greater than about 85% enantiomeric excess. In some embodiments, the

compounds exist in greater than about 90% enantiomeric excess. In some embodiments, the compounds exist in greater than about 92% enantiomeric excess. In some embodiments, the compounds exist in greater than about 95% enantiomeric excess. In some embodiments, the compounds exist in greater than about 97% enantiomeric excess. In some embodiments, the compounds exist in greater than about 99% enantiomeric excess.

5

Pharmaceutical compositions

[0103] The present invention further provides pharmaceutical compositions comprising a therapeutically effective amount of the compounds of the present invention and a pharmaceutically acceptable carrier or diluent.

10

[0104] The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers, diluents or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 21st Edition (University of the Sciences in Philadelphia, ed., Lippincott Williams & Wilkins 2005)..

15

Further exemplary compositions of the compounds of the invention are described in, for example, U.S. Patent Nos. 5,807,855; 7,648,991; 7,767,683; 7,772,240; 8,076,342; U.S. Patent Publication Nos. 2008/0269248; 2010/0069676; 2011/0178094; 2011/0207744; WO 2005/016900; EP 0 638 073; and J. Med. Chem. 1995, 38, 4380-4392.

20

[0105] The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as oral, nasal, topical (including buccal and sublingual), and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) routes. It will be appreciated that the route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient.

25

[0106] The daily dose of the compounds of the invention, calculated as the free base, is suitably from about 1.0 to about 160 mg/day, more suitably from about 1 to about 100 mg, e.g. preferably from about 2 to about 55, such as from about 2 to about 15 mg, e.g. from about 3 to about 10 mg. In some embodiments, the daily dose is from about 0.1 mg to about 500 mg. In some embodiments, the daily dose is from about 1 mg to about 500 mg. In some embodiments, the daily dose is from about 1 mg to about 400 mg. In some embodiments, the daily dose is from about 1 mg to about 300 mg. In some embodiments, the daily dose is from about 1 mg to about 200 mg. In some embodiments, the daily dose is from about 1 mg to about 160 mg. In some embodiments, the daily dose is from about 1 mg to about 100 mg. In some embodiments, the daily dose is from about 1 mg to about 60 mg. In some embodiments, the daily dose is from about 2 mg to about 30 mg. In some embodiments, the daily dose is from about 2 mg to about 15 mg. In some embodiments, the daily dose is from about 3 mg to about 10 mg. In some embodiments, the daily dose is about 60 mg. In some embodiments, the daily dose is about 50 mg. In some embodiments, the daily dose is about 40 mg. In some embodiments, the daily dose is about 30 mg. In some embodiments, the daily dose is about 20 mg. In some embodiments, the daily dose is about 10 mg. In some embodiments, the daily dose is about 5 mg. In some embodiments, the daily dose is about 3 mg. In some embodiments, the daily dose is about 2 mg. In some embodiments, the daily dose is about 1 mg.

35

[0107] For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typical doses are in the order of half the dose employed for oral administration.

40

[0108] The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. Examples of suitable organic and inorganic acids are described herein.

45

[0109] In some embodiments, the composition comprises a cyclodextrin. In some embodiments, the composition comprises a cyclodextrin in water. In some embodiments, the cyclodextrin is hydroxypropyl- β -cyclodextrin. In some embodiments, the composition comprises hydroxypropyl- β -cyclodextrin in water.

Treatment of Disorders

50

[0110] The invention discloses the medical use of compounds of the present invention, such as for the treatment of a disease in the central nervous system, including psychosis, in particular schizophrenia or other diseases involving psychotic symptoms, such as, e.g., Schizophrenia, Schizophreniform Disorder, Schizoaffective Disorder, Delusional Disorder, Brief Psychotic Disorder, Shared Psychotic Disorder as well other psychotic disorders or diseases that present with psychotic symptoms, e.g. bipolar disorder, such as mania in bipolar disorder. Compounds and/or compositions of the invention can further be used in treatment of disorders such as those described in, for example, U.S. Patent Nos. 5,807,855; 7,648,991; 7,767,683; 7,772,240; 8,076,342; U.S. Patent Publication Nos. 2008/0269248; 2010/0069676; 2011/0178094; 2011/0207744; WO 2005/016900; EP 0 638 073; and J. Med. Chem. 1995, 38, 4380-4392. The invention also discloses the medical use of compounds of the present invention as combination therapy in conjunction with other therapeutic agents such as those described in, for example, U.S. Patent Nos. 5,807,855; 7,648,991; 7,767,683; 7,772,240; 8,076,342; U.S. Patent Publication Nos. 2008/0269248; 2010/0069676; 2011/0178094; 2011/0207744; WO 2005/016900; EP 0 638 073; and J. Med. Chem. 1995, 38, 4380-4392.

55

[0111] It will be recognized that one or more features of any embodiments disclosed herein may be combined and/or

rearranged within the scope of the invention to produce further embodiments that are also within the scope of the invention.
[0112] The invention is further described by the following non-limiting Examples.

EXAMPLES

[0113] Examples are provided below to facilitate a more complete understanding of the invention. The following examples illustrate the exemplary modes of making and practicing the invention. However, the scope of the invention is not limited to specific embodiments disclosed in these Examples, which are for purposes of illustration only, since alternative methods can be utilized to obtain similar results.

[0114] Purification of compounds by chromatography refers to the application of silica gel chromatography using either manual flash chromatography or automated flash chromatography, typically performed using eluent gradients from heptanes to ethyl acetate or mixtures of ethyl acetate, triethylamine and methanol.

Description of LCMS Methods.

[0115] Compounds (I), (II), (III), (IV), (V), (VI) and (VII) were characterized by LCMS using the following methods (Table 1):

Table 1: Methods for LCMS Analysis

Methods WXV-AB5, WXV-AB10, and WXV-AB30		
Equipment	Agilent 1100 LCMS system with ELS Detector [method WuXiAB25 Agilent 1200 LCMS system with ELS Detector]	
Pump	G1311A	
Degasser	G1379A	
Well-plate Autosampler	G1367A	
Column Oven	G1316A	
DAD	G1315B	
MSD	G1946C or G1956A [method WuXiAB25 6110]	
ELSD	Alltech ELSD 800 [method WuXiAB25 Algent1200]	
Column	YMC ODS-AQ [method WuXiAB25 Agilent TC-C18]	
35	Particle size	5 micrometer
	Pore size	12 nm
	Dimension	50 * 2.0 mm ID [method WuXiAB25 50*2.1 mm ID]
40	Injection volume	2 microL
	Column temperature	50°C
45	Flow	0.8 mL/min
	Mobile phases	A 0.1% TFA in water
50		B 0.05% TFA in acetonitrile
	Total run time	4.5 min
	Gradient	linear
55	UV Detection	Wavelength 254 nm
	ELSD Detection	Temperature: 50°C
	Gas Pressure:	3.2 bar
	Time	Gradient

(continued)

Methods WXV-AB5, WXV-AB10, and WXV-AB30

5	Equipment	Agilent 1100 LCMS system with ELS Detector	
WXV-AB05	0 min	95% A	5% B
	3.5 min	0% A	100% B
	3.55 min	95% A	6% B
WXV-AB10	0 min	90% A	10% B
	3.4 min	100% B	
	3.5 min	100% B	
WXV-AB30	3.51 min	90% A	10% B
	0 min	70% A	30% B
	3.2 min	0% A	100% B
WuXiAB25	3.5 min	0% A	100% B
	3.55 min	70% A	30% B
	0 min	75% A	25% B
20	3.4 min	0 % A	100% B
	4 min	0 % A	100% B
	4.01 min	75% A	25% B
25	4.5 min	75% A	25% B

Method 131

25	Equipment	Sciex API150EX equipped with APPI-source operating in positive ion mode	
	LC-MS	were run on a Sciex API150EX equipped with APPI-source operating in positive ion mode.	
30	The HPLC	consisted of Shimadzu LC10-ADvp LC pumps, SPD-M20A PDA detector (operating at 254 nm) and SCL-10A system controller. Autosampler was Gilson	
	Autosampler	Gilson 215	
35	Column	Column Oven	ELSD Waters Symmetry C-18
		Jones Chromatography 7990R Sedere	
		Sedex 85	
40	Injection volume	Particle size	3.5 micrometer
		Dimension	30 * 4.6 mm ID
45	Column temperature	10 microL	
	Flow	60°C	
50	Mobile phases	A	0.05% TFA in water
		B	0.05% TFA in methanol
	UV Detection	Total run time	2.8 min
		Gradient	non-linear
	ELSD Detection	Wavelength	254 nm
55		Temperature:	50°C
		Gas Pressure:	4.4 bar
		Time	Gradient
		0.01 min	17% B in A
		0.27 min	28% B in A

(continued)

	Time	Gradient
5	0.53 min	39% B in A
	0.80 min	50% B in A
	1.07 min	59% B in A
	1.34 min	68% B in A
	1.60 min	78% B in A
10	1.87 min	86% B in A
	2.14 min	93% B in A
	2.38 min	100% B
	2.40 min	17% B in A
15	2.80 min	17% B in A

Description of Chiral HPLC methods

[0116] The enantiomeric purity was assayed on a Hewlett Packard 1100 series system equipped with a diode array detector and using ChemStation for LC Rev. A.08.03[847]. The HPLC method parameters are described in the table below (Table 2). Compound (X) has a retention time around 13.6-13.7 min while its enantiomer, 4-((1S,3R)-6-chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine, elutes at 8.5-8.6 min.

Table 2: Methods for Chiral HPLC Analysis

25	Sample Preparation	1-3 mg/mL in hexane/2-propanol (80/20 v/v)
	Column:	Chiraldak ADH 5microm 250 x 4.6mm
	Column Temperature (°C):	30
30	Injection (microL):	5
	Detection: Wavelength, Bandwidth(nm):	240,8
	Total run-time	30 min
	Flow Rate (mL.min ⁻¹):	0.6
35	Mobile Phase	hexane/2-propanol/diethylamine/propionic acid 90/10/0.2/2

Example 1 Preparation of 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine•butanedioic acid (Compound (I)•butanedioic acid salt).


[0117]

45

50

55

Scheme 8. Synthesis of Compound (I).



[0118] 1-((1R,3S)-6-Chloro-3-phenyl-indan-1-yl)-3,3-dimethyl-piperazine hydrochloride (11.1 g) was dissolved in a mixture of toluene (74 mL) and water (74 mL). Preparation of 1-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-3,3-dimethyl-piperazine hydrochloride is disclosed in the patent literature (Dahl, Wøhlk Nielsen, Suteu, Robin, Brøsen WO2006/086984 A1; Bang-Andersen, Bøgesø, Jensen, Svane, Dahl, Howells, Lyngsø, Mow WO2005/016901 A1). 12.0 M of potassium hydroxide in water (5.38 mL), tetra-*N*-butylammonium bromide (1.42 g), and *d*₃-iodomethane (Aldrich catalog # 176036; 2.4 mL) were added and the mixture was stirred at room temperature for 18 hours (Scheme 8). The mixture was filtered through a glass filter into a separatory funnel. The solid on the filter was washed with toluene (50 mL) into the separatory funnel. The aqueous layer was extracted with toluene (100 mL) and the combined organic layers were washed with concentrated aqueous ammonia (100 mL) and subsequently with water (100 mL) before it was dried over sodium sulfate, filtered, and concentrated in vacuum affording a slightly yellow oil. The oil was cooled to -78°C under vacuum which solidified the oil. Upon warming to room temperature the oil became a semi-solid.

[0119] This material was dissolved in acetone (30 mL); in a separate flask butanedioic acid (3.46 g) was suspended in acetone (30 mL) and warmed to reflux (not all of the diacid went into solution). The acid suspension was added to the solution of the crude product and additional acetone (50 mL) was added to the butanedioic acid residue and then poured into solution. The mixture was stirred overnight. Partial precipitation had occurred overnight, and the mixture was concentrated in vacuum. The residue was re-dissolved in acetone (70 mL) and warmed to reflux and allowed to cool to room temperature and stirred for 2 hours.

[0120] The mixture was filtered affording 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine•butanedioic acid (Compound (I)•butanedioic acid salt; 7.61 g). LC-MS (method 131): RT(UV) 1.57 min; UV/ELS purity 100%/100%; mass observed 358.0. Incorporation of three deuterium atoms >99 %. The proton-decoupled ¹³C NMR spectrum showed a heptet around 36.4 ppm corresponding to the deuterated M2 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.

Example 2 Alternative method of preparation of 4-((1*R*,3*S*)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine•butanedioic acid (Compound (I)•butanedioic acid salt)


[0121] The free base of 1-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-3,3-dimethyl-piperazine was prepared from the corresponding hydrochloride salt by partitioning 23.4 g of the salt between a mixture of water (100 mL), concentrated aqueous potassium hydroxide (40 mL), and toluene (250 mL). The organic layer was washed with a mixture of water (50 mL) and concentrated aqueous potassium hydroxide (10 mL). The combined aqueous layers were extracted with toluene (75 mL). The combined organic layers were dried over sodium sulfate, filtered, and concentrated in vacuum affording the free base of 1-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-3,3-dimethyl-piperazine (21.0 g) as a colorless oil. This material was dissolved in a mixture of toluene (150 mL) and water (150 mL), before 12.0 M aqueous potassium hydroxide (11.3 mL), tetra-*N*-butylammonium bromide (2.98 g), and *d*₃-iodomethane (4.9 mL) were added and mixture was stirred at room temperature for 18 hours.

[0122] Work-up and purification was performed as described above and afforded 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₂-2,2-dimethyl-piperazine-butanedioic acid (Compound (I)[•]butanedioic acid salt; 14.34 g; 48.9%).

Example 3 Preparation of 4-((1R,3S)-6-chloro-3-phenyl-*d*₅-indan-1-yl)-1,2,2-trimethyl-piperazine (Compound (II)) and 4-((1R,3S)-6-chloro-3-phenyl-*d*₅-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine (Compound (IV)).

[0123] To a solution of compound A (57 g) in tetrahydrofuran (600 mL) was added triethylamine (30 mL) dropwise over 30 min. The reaction mixture was kept at room temperature for 3 hours. The precipitated solid was filtered and the filtrate was concentrated *in vacuo*. The residue was reprecipitated from diethyl ether to afford compound B (31 g) as a yellow solid. To a solution of compound phenyl-*d*₅-boronic acid (25 g) in 1,4-dioxane/water (900 mL/ 90 mL) was added [Rh(ndb)₂]BF₄ (1.3 g), racemic BINAP (2.1 g) and triethylamine (14 mL), then the reaction mixture was kept at room temperature for 2 hours under N₂. Then compound indenone (19 g) was added, and the resulting mixture was heated to 100 °C for 3 hours. The precipitated solid was filtered off. The filtrate was concentrated *in vacuo*. The residue was purified by chromatography to afford indanone C (10 g).

Scheme 9. Synthesis of Compound C.



[0124] 13.4 kg 3-Bromo-6-chloro-indan-1-one (A; for references on this material, see: Bøgesø EP 35363 A1 19810909 and Kehler, Juhl, Püschi, WO 2008025361) was dissolved in tetrahydrofuran (170.8 L), and the solution was cooled to 0-5 °C (Scheme 9). Triethylamine (9.1 L) was added over 0.5h. The mixture was stirred at 0-5 °C for 5 hours before an additional portion of triethylamine (2.48 L) was added over 0.5 hour, and stirring was continued for 2 hours. The mixture was filtered, and the filtrate was concentrated to 30 L before *n*-heptane (102 L) was added. The volume was reduced to 60 L. More *n*-heptane (203 L) was added, and the mixture was stirred for 1 hour. Silica gel (17.2 kg) was added. The mixture was filtered, and the residual solid was washed with *n*-heptane (100 L). The combined filtrates were concentrated to 30 L and stirred at 0-5 °C for 1 hour. The mixture was centrifuged, and the residual solid was dried to afford 6-chloro-inden-1-one (compound B; 2.42 kg) sufficiently pure for the next step.


[0125] 2-Methyl-tetrahydrofuran (85 L) and *N,N*-dimethyl acetamide (12.4 L) were added to a reactor followed by potassium acetate (10.9 kg) and bis(pinacolato)diboron (14.8 kg). The resulting mixture was stirred for 0.5 hour. Pd(dppf)Cl₂-DCM (0.91 kg) was added followed by bromobenzene-*d*₅ (9.0 kg) and 2-methyl-tetrahydrofuran (12.2 L). The mixture was heated to 80-85 °C for 3 hours, before the temperature was reduced to ambient temperature. The crude mixture was filtered via kieselguhr and silica gel. The filter-cake was washed with 2-methyl-tetrahydrofuran (31 L). The combined filtrates were concentrated to approximately 25 L while maintaining the temperature below 35 °C. *n*-Heptane (52 L) and 7% aqueous NaHCO₃ (31 L) were added, and the mixture was stirred for 0.5 hour. The organic layer was stirred with 7% aqueous NaHCO₃ (31 L) for 0.5 hour. The combined aqueous layers were extracted with *n*-heptane (22 L) over 0.5 hour. The combined organic extracts were washed with 25% aqueous NaCl (50 L) over 0.5 hour. The organic layer was concentrated while maintaining the temperature below 35 °C to afford 4,4,5,5-tetramethyl-2-*d*₅-phenyl-[1,3,2]dioxaborolane (compound B'; 10.5 kg) sufficiently pure for the next step.

[0126] To a reactor was added sequentially 1,4-dioxane (85 L), 6-chloro-inden-1-one (compound B; 9.09 kg prepared in a similar manner to the one described above), 1,5-cyclooctadiene (0.2 L), bis(norbornadiene)rhodium(I) tetrafluoroborate (0.52 kg), triethylamine (5.5 L), 4,4,5,5-tetramethyl-2-*d*₅-phenyl-[1,3,2]dioxaborolane (compound B'; 6.5 kg), and 1,4-dioxane (26 L). The mixture was heated to 48-53 °C and stirred at that temperature for 5 hours. The reaction was quenched by the addition of 2M aqueous HCl (13 kg). Then *n*-heptane (110 L), methyl *tert*-butyl ether (32 L), and water

(90 L) were added, and the resulting mixture was stirred for 0.3 hour. The organic layer was washed with water (90 L) over 0.3 hour. The combined aqueous layers were extracted with a mixture of methyl *tert*-butyl ether (30 L) and *n*-heptane (57 L) over 0.3 hour. The combined organic layers were filtered through silica gel (13 kg). The filter-cake was washed with a 2:1 mixture of *n*-heptane and methyl *tert*-butyl ether (19.5 kg). The filtrate was concentrated to approximately 25 L. *n*-Heptane (45 L) was added, and the volume was reduced to approximately 25 L. *n*-Heptane (45 L) was added, and the volume was reduced to approximately 35 L. The mixture was stirred at 0-5 °C for 3 hours. The mixture was centrifuged, and the residual solid was dried to afford racemic 6-chloro-3-*d*₅-phenyl-indan-1-one (compound C; 8.4 kg) sufficiently pure for the next step.

[0127] Tetrahydrofuran (90 L) was added to a reactor followed by water (10 L) and 6-chloro-3-*d*₅-phenyl-indan-1-one (compound C; 7.73 kg) (Scheme 10). The mixture was cooled to -35-30 °C. Sodium borohydride (1.5 kg) was added portion-wise while maintaining the temperature at -35 - -30 °C. The resulting mixture was stirred at -35 - -30 °C for 5 hours before it was allowed to warm to ambient temperature. Excess sodium borohydride was quenched by the addition of 2M aqueous HCl (7.6 kg) while maintaining the temperature below 45 °C. Water (17 L) and methyl *tert*-butyl ether (67 L) were added and the mixture was stirred for 0.3 hour. The aq layer was extracted with methyl *tert*-butyl ether (39 L) over 0.3 hour. The combined organic layers were washed with brine (36 kg) over 0.3 hour. The organic layer was filtered through silica gel (6.4 kg). The filter-cake was washed with methyl *tert*-butyl ether (20 L). The combined filtrates were concentrated to approximately 30 L while maintaining the temperature below 45 °C. *n*-Heptane (55 L) was added and the resulting mixture was concentrated to approximately 30 L while maintaining the temperature below 45 °C. The resulting mixture was stirred at 0-5 °C for 2 hours. The mixture was centrifuged, and the filter-cake was washed with *n*-heptane (12 L) before it was centrifuged again. The residual solid was dried to afford crude D. 4.87 kg of this material was dissolved in methyl *tert*-butyl ether (20 L) and dried over Na₂SO₄ (2 kg) over 0.25 hour. The mixture was filtered, and the filter-cake was washed with methyl *tert*-butyl ether (4.4 L). The combined filtrate was concentrated to approximately 20 L while maintaining the temperature below 45 °C. *n*-Heptane (32 L) was added and the mixture was to approximately 25 L while maintaining the temperature below 45 °C. *n*-Heptane (16 L) was added and the mixture was to approximately 20 L while maintaining the temperature below 45 °C. The solid was filtered off and dried to afford racemic *cis*-6-chloro-3-*d*₅-phenyl-indan-1-ol (compound D; 4.99 kg) sufficiently pure for the next step.

Scheme 10. Synthesis and resolution of Compound E.



[0128] To a solution of racemic *cis*-6-chloro-3-*d*₅-phenyl-indan-1-ol (compound D; 50 g) in 2-isopropoxypropane (200 mL) was added vinyl butyrate (120 mL) and Novozym-435 (15 g). The mixture was kept at ambient temperature for 2 days. The solid was filtered off. The filtrate was evaporated and purified by chromatography on silica gel to afford (1S,3S)-6-chloro-3-*d*₅-phenyl-indan-1-ol (compound E; 13 g) sufficiently pure for the next step.

[0129] To a solution of (1S,3S)-6-chloro-3-*d*₅-phenyl-indan-1-ol (compound E; 7 g) in THF (100 mL) was treated with SOCl₂ (6.6 g) at ambient temperature overnight. The mixture was poured into ice-cold water, and extracted with ethyl acetate. The organic layer was washed with brine. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo* to afford the intermediate chloride (7.5 g). 3.5 g of this material was dissolved in 2-butanone (50 mL) and reacted with 2,2-dimethyl-piperazine (1.7 g) in the presence of K₂CO₃ (2.7 g) at reflux overnight. The solid was filtered off. The filtrate was concentrated *in vacuo* and the residue was purified by preparative HPLC on a Shimadzu FRC-10A instrument fitted with a Synergi C18 column (250mm*50mm, 10 microm) using water and acetonitrile (containing 0.1%TFA, v/v) as the eluent to afford 1-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-3,3-dimethyl-piperazine (compound F; 2.6 g) sufficiently pure for the next step.

[0130] To a solution of 1-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-3,3-dimethyl-piperazine (compound F; 2.2 g) in HCHO/HCOOH (3 mL/3 mL) was refluxed overnight. The volatiles were removed *in vacuo*. The residue was partitioned between ethyl acetate and 10% aq NaOH. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*.

The residue was purified by chromatography on silica gel to afford 4-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine (Compound (II); 1.89 g). LC-MS (method WXV-AB05): RT(UV) 2.43 min; UV/ELS purity 95.1%/99.6%; mass observed 360.2. Incorporation of five deuterium atoms >95 %. The proton-decoupled ¹³C NMR spectrum showed three triplets around 126.1, 127.2, and 128.2 ppm corresponding to the deuterated M3 metabolic sites; these signal collapsed to three singlets in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.


[0131] To a solution of 1-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-3,3-dimethyl-piperazine (compound F; 3.0 g) in DCDO/DCOOD (4 mL/4 mL) was refluxed overnight. The volatiles were removed *in vacuo*. The residue was partitioned between ethyl acetate and 10% aq NaOH. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*.

The residue was purified by chromatography on silica gel to afford 4-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-1-*d*₃-methyl-2,2-diimethyl-piperazine (Compound (IV); 2.14 g). LC-MS (method WXV-AB10): RT(UV) 2.06 min; UV/ELS purity 98%/100%; mass observed 363.3. Incorporation of eight deuterium atoms >94 %. The proton-decoupled ¹³C NMR spectrum showed a heptet around 36.4 ppm corresponding to the deuterated M2 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. The proton-decoupled ¹³C NMR spectrum further showed three triplets around 126.1, 127.2, and 128.2 ppm corresponding to the deuterated M3 metabolic sites; these signal collapsed to three singlets in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.

Example 4: Preparation of 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine-6,6-*d*₂ (Compound (III)), 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine-6,6-*d*₂ (Compound (V)), 4-((1R,3S)-6-chloro-3-phenyl-*d*₅-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine-6,6-*d*₂ (Compound (VI)), and 4-((1R,3S)-6-chloro-3-phenyl-*d*₅-indan-1-yl)-1,2,2-trimethyl-piperazine-6,6-*d*₂ (Compound (VII)).

[0132] 2-Amino-2-methyl-propionic acid (50.0 g) was suspended in a mixture of methanol and triethylamine (9:1, 1.2 L) (Scheme 11). 1M aqueous NaOH (450 mL) was added with stirring until all solid was dissolved. Di-*tert*-butyl dicarbonate (Boc₂O; 214.0 g) was added, and the mixture was stirred at ambient temperature overnight. The organic volatiles were removed *in vacuo*. EtOAc (500 mL) was added. The organic layer was washed with brine and dried over Na₂SO₄, filtered, then concentrated to afford 2-*tert*-butoxycarbonylamino-2-methyl-propionic acid (compound K; 90 g) as a white solid which was used directly in next step directly.

Scheme 11. Synthesis of intermediate J.



[0133] A mixture of afford 2-*tert*-butoxycarbonylamino-2-methyl-propionic acid (compound K; 60.0 g) and 1-ethyl-3-(3-

dimethylaminopropyl) carbodiimide hydrochloride (EDC·HCl; 86.4 g) in dichloromethane (900 mL) was stirred at ambient temperature, then N,O-dimethyl hydroxylamine hydrochloride (35.3 g) and triethylamine (150 mL) were added. The resulting mixture was stirred at ambient temperature for 3 days. Water was added and most of volatiles were removed *in vacuo*. The residue was partitioned between DCM and aqueous NaHCO₃. The organic layer was washed with 3M aqueous HCl, subsequently with brine before it was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by silica gel chromatography to give [1-(methoxy-methyl-carbamoyl)-1-methyl-ethyl]-carbamic acid *tert*-butyl ester (compound L; 28.2 g) as a white solid sufficiently pure for the next step.

[0134] Lithium aluminum hydride (7.8 g) was added to a stirred solution of [1-(methoxy-methyl-carbamoyl)-1-methyl-ethyl]-carbamic acid *tert*-butyl ester (compound L; 42.0 g) in dry diethyl ether (1.5 L) at -40 °C. Then stirred at that temperature for about 5 min. Excess LiAlH₄ was quenched with a solution of potassium hydrogen sulfate in water. The resulting mixture was partitioned between EtOAc and 3M aqueous HCl. The organic layer was washed with sat. aqueous NaHCO₃, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to afford (1,1-dimethyl-2-oxo-ethyl)-carbamic acid *tert*-butyl ester (compound M; 29 g) sufficiently pure for the next step.

[0135] Amino-acetic acid methyl ester hydrochloride (80.6 g) and Et₃N (160 mL) were dissolved in DCM (1000 mL) and stirred for 15 min to liberate the amine from the salt. Then a solution of 1,1-dimethyl-2-oxo-ethyl)-carbamic acid *tert*-butyl ester (compound M; 29.0 g) in DCM (600 mL) was added. The resulting mixture was stirred for 0.5 hour at ambient temperature before NaBH(OAc)₃ (102 g) was added and the mixture was stirred at ambient temperature overnight. Sat. aqueous NaHCO₃ was added. The aqueous layer was extracted with DCM. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by silica gel chromatography to afford (2-*tert*-butoxycarbonylamino-2-methyl-propylamino)-acetic acid methyl ester (compound N; 26.5 g) as white solid which was used directly in the next step.

[0136] A mixture of (2-*tert*-butoxycarbonylamino-2-methyl-propylamino)-acetic acid methyl ester (compound N; 26.5 g) in DCM (800 mL) was stirred at ambient temperature, TFA (180 mL) was added drop-wise. The mixture was stirred at 30-40 °C for 5h before it was concentrated *in vacuo*. The residue was partitioned between dissolved toluene and water. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residual solid was dissolved in a mixture of ethanol (400 mL) and methanol (90 mL). K₂CO₃ (207 g) was added and the mixture was refluxed overnight. The mixture was cooled to room temperature. DCM (2500 mL) was added, and the mixture was stirred for 1 hour at ambient temperature. The solid was filtered off, and the filtrate was concentrated *in vacuo* to afford 6,6-dimethyl-piperazin-2-one (Compound I; 5.85 g) as a white solid sufficiently pure for the next step.

[0137] A solution of 6,6-dimethyl-piperazin-2-one (Compound I; 3.6 g) in THF (20 mL) was stirred at 0 °C. Lithium aluminum deuteride (LiAlD₄; 3.6 g) was added then the mixture was refluxed overnight. The mixture was cooled to ambient temperature and Na₂SO₄ was added. The mixture was stirred for 0.5h before most of the volatiles were removed *in vacuo*. The residue was suspended in a saturated solution of HCl in EtOAc at ambient temperature for 0.5 hour. The solid was filtered off and dried to afford to give 2,2-*d*₂-6,6-dimethyl-piperazine as the bis-hydrochloride salt (Compound J·2HCl; 5.3 g) sufficiently pure for the next step.


[0138] To a solution of compound E' (5 g) in THF (50 mL) was added SOCl₂ (4.7 g), and the resulting mixture was stirred overnight at ambient temperature (Scheme 12). The mixture was poured into ice-water and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to afford the corresponding chloride (5.3 g) which was used directly in the next step. 3.3 g of this material was dissolved in 2-butanone (50 mL) and reacted with 2,2-*d*₂-6,6-dimethyl-piperazine (Compound J; 3 g) in the presence of K₂CO₃ (8.28 g) at reflux overnight. The solid was filtered off. The filtrate was concentrated *in vacuo*. The residue was purified by preparative HPLC on a Shimadzu FRC-10A instrument fitted with a Synergy C18 column (250mm*50mm, 10 microm) using water and acetonitrile (containing 0.1%TFA, v/v) as the eluents to afford 1-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-3,3-*d*₂-5,5-dimethyl-piperazine (Compound O; 1.7 g).

45

50

55

Scheme 12. Synthesis of Compound (III) and Compound (V).



[0139] A solution of 1-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-3,3-*d*₂-5,5-dimethyl-piperazine (Compound O; 0.5 g) in HCHO/HCOOH (1 mL/1 mL) was refluxed overnight. The volatiles were removed *in vacuo*. The residue was partitioned between EtOAc and 10% aqueous NaOH. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel to afford 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine-6,6-*d*₂ (Compound (III); 0.33 g). LC-MS (method WXV-AB30): RT(UV) 1.42 min; UV/ELS purity 100%/100%; mass observed 357.2. Incorporation of two deuterium atoms >97 %. The proton-decoupled ¹³C NMR spectrum showed a quintet around 49.5 ppm corresponding to the deuterated M1 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. The proton-decoupled ¹³C NMR spectrum further showed three triplets around 126.1, 127.2, and 128.2 ppm corresponding to the deuterated M3 metabolic sites; these signal collapsed to three singlets in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.

[0140] A solution of 1-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-3,3-*d*₂-5,5-dimethyl-piperazine (Compound O; 0.7 g) in DCD₂/DCOOD (1 mL/1 mL) was refluxed overnight. The volatiles were removed *in vacuo*. The residue was partitioned between EtOAc and 10% aqueous NaOH. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel to afford 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine-6,6-*d*₂ (Compound (V); 0.49 g). LC-MS (method WXVAB25): RT(UV) 2.13 min; UV/ELS purity 100%/100%; mass observed 360.2. Incorporation of five deuterium atoms >95 %. The proton-decoupled ¹³C NMR spectrum showed a heptet around 36.4 ppm corresponding to the deuterated M2 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. The proton-decoupled ¹³C NMR spectrum further showed a quintet around 49.5 ppm corresponding to the deuterated M1 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.


[0141] To a solution of (1S,3S)-6-chloro-3-*d*₅-phenyl-indan-1-ol (compound E; 7 g) in THF (100 mL) was treated with SOCl₂ (6.6 g) at ambient temperature overnight (Scheme 13). The mixture was poured into ice-cold water, and extracted with ethyl acetate. The organic layer was washed with brine. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo* to afford the intermediate chloride (7.5 g).

45

50

55

Scheme 13. Synthesis of Compound (VI) and Compound (VII).



[0142] 1.8 g of this material was dissolved in 2-butanone (30 mL) and reacted with 2,2-*d*₂-6,6-dimethyl-piperazine (Compound J; 1.4 g) in the presence of K₂CO₃ (5.5 g) at reflux overnight. The solid was filtered off. The filtrate was concentrated *in vacuo*. The residue was purified by preparative HPLC on a Shimadzu FRC-10A instrument fitted with a Synergy C 18 column (250mm*50mm, 10 microm) using water and acetonitrile (containing 0.1 %TFA, v/v) as the eluents to afford 1-((1*R*,3*S*)-6-Chloro-3-*d*₅-phenyl-indan-1-yl)-3,3-*d*₂-5,5-dimethyl-piperazine (Compound P; 1.7 g).

[0143] A solution of 1-((1R,3S)-6-Chloro-3-*d*₅-phenyl-indan-1-yl)-3-*d*₂-5,5-dimethyl-piperazine (Compound P; 1 g) in DCDO/DCOOD (1.5 mL/1.5 mL) was refluxed overnight. The volatiles were removed *in vacuo*. The residue was partitioned between EtOAc and 10% aq NaOH. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel to afford 4-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-1-*d*₃-methyl-2,2-dimethyl-piperazine-6,6-*d*₂ (Compound (VI); 0.55 g). LC-MS (method WuXiAB25): RT(UV) 2.13 min; UV/ELS purity 98.2%/100%; mass observed 365.2. Incorporation of ten deuterium atoms >91 %. The proton-decoupled ¹³C NMR spectrum showed a heptet around 36.4 ppm corresponding to the deuterated M2 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. The proton-decoupled ¹³C NMR spectrum further showed a quintet around 49.5 ppm corresponding to the deuterated M1 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. The proton-decoupled ¹³C NMR spectrum further showed three triplets around 126.1, 127.2, and 128.2 ppm corresponding to the deuterated M3 metabolic sites; these signal collapsed to three singlets in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.

[0144] A solution of 1-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-3,3-*d*₂-5,5-dimethyl-piperazine (Compound P; 0.7 g) in HCHO/HCOOH (1 mL/1 mL) was refluxed overnight. The volatiles were removed *in vacuo*. The residue was partitioned between EtOAc and 10% aqueous NaOH. The organic layer was dried over Na₂SO₄, filtered, and concentrated *in vacuo*. The residue was purified by chromatography on silica gel to afford 4-((1R,3S)-6-chloro-3-*d*₅-phenyl-indan-1-yl)-1-methyl-2,2-dimethyl-piperazine-6,6-*d*₂ (Compound (VII); 0.47 g). LC-MS (method WXV-AB30): RT(UV) 1.33 min; UV/ELS purity 97.4%/100%; mass observed 362.3. Incorporation of seven deuterium atoms >93 %. The proton-decoupled ¹³C NMR spectrum showed a quintet around 49.5 ppm corresponding to the deuterated M1 metabolic site; this signal collapsed to a singlet in the proton- and deuterium-decoupled ¹³C NMR spectrum. The proton-decoupled ¹³C NMR spectrum further showed three triplets around 126.1, 127.2, and 128.2 ppm corresponding to the deuterated M3 metabolic sites; these signal collapsed to three singlets in the proton- and deuterium-decoupled ¹³C NMR spectrum. All other signals were singlets in both spectra. Optical purity >95% ee.

Example 5: Description of NMR determination of the position(s) bearing deuterium rather than hydrogen

[0145] NMR spectra were recorded on a Bruker 600-Avance-III spectrometer equipped with a 5 mm TCI cryoprobe operating at 150.91 MHz for ^{13}C . The solvent CDCl_3 was used as internal reference for the proton-decoupled experiments, while the proton- and inverse gated deuterium-decoupled spectra were recorded using gated lock. Difference(s) between the two spectra for the compounds of the invention determine(s) the position(s) of the deuterium atoms. When combining this information summarized in the table below (Table 3) with the electrospray mass spectrometry data that determined degree of deuteration, the structures of the compounds of the invention can be assigned unambiguously.

Table 3: Carbon NMR data for compounds.

Cmpd.	M2 (methyl group @ ~36.4 ppm)			M1 (methylene group @ ~49.5 ppm)			M3 (phenyl group @ ~126.1 ppm, ~127.2 (2C), and ~128.2 (2C))		
	¹³ C NMR proton-decoupled	¹³ C NMR proton- and deuterium-decoupled	¹³ C NMR proton-decoupled	¹³ C NMR proton- and deuterium-decoupled	¹³ C NMR proton-decoupled	¹³ C NMR proton-decoupled	¹³ C NMR proton- and deuterium-decoupled	¹³ C NMR proton- and deuterium-decoupled	¹³ C NMR proton- and deuterium-decoupled
(I)	heptet	singlet	singlet	singlet	singlet	singlet	singlets	singlets	singlets
(II)	singlet	singlet	singlet	singlet	singlet	singlet	3 triplets	3 triplets	3 singlets
(III)	singlet	singlet	quintet	quintet	quintet	quintet	3 singlets	3 singlets	3 singlets
(IV)	heptet	singlet	singlet	singlet	singlet	singlet	3 triplets	3 triplets	3 singlets
(V)	heptet	singlet	quintet	quintet	singlet	singlet	3 singlets	3 singlets	3 singlets
(VI)	heptet	singlet	quintet	quintet	singlet	singlet	3 triplets	3 triplets	3 singlets
(VII)	singlet	singlet	quintet	quintet	singlet	singlet	3 triplets	3 triplets	3 singlets

Only NMR signals that 'change' as a consequence of the presence of D rather than H in the compounds of the invention are included in the table.

[0146] Relevant regions of the ^{13}C proton-decoupled (lower spectrum) and ^{13}C proton- and deuterium-decoupled (upper spectrum) NMR spectra of Compound (II) and Compound (V) are shown in Figure 3 as representative examples. Selected regions of the proton-decoupled and proton- and deuterium-decoupled ^{13}C NMR spectra of Compound (II) [Fig. 3A] and Compound (V) [Fig. 3B].

5

Example 6: Description of the electrospray mass spectrometry to determine degree of deuteration

[0147] Instrumentation: Mass spectra of acidic, aqueous solutions of the compounds were obtained on a Hewlett Packard quadrupole mass spectrometer model 1100 LC-MSD. Liquid chromatography was performed on an Agilent 1100 HPLC-system coupled to the mass spectrometer.

10

[0148] Experimental: Solutions of the samples were made by dissolving approx. 2 mg substance in 2 mL methanol + 18 mL 10 mM ammonium formate pH 3.0. Subsequently the solutions were diluted 100-fold prior to analysis. In order to get a "clean" peak, the samples were chromatographed using a Waters X-bridge C 18, 3.5 microm (150x2.1mm) column, and 0.1 % trifluoroacetic acid / acetonitrile 50/50 as mobile phase. This procedure gave one peak of the compound of interest eluting at ca. 3.6 min, containing both the deuterated compounds of the inversion as well as small quantities of deuterium-deficient species. The mass spectra obtained from these peaks were used to evaluate the speciation of the target molecules. The results were analyzed in percent of the total amount of substance, adding up to 100%. The actual potency of the compounds were not analyzed, merely the relative content of the deuterium deficient species.

15

[0149] As a representative example, the mass spectrum of Compound (IV) is shown in Figure 4. The isotopic pattern of the protonated Compound (V) $[\text{M}+\text{H}]^+$ with mass 363.1u (362.1u + 1.0u) and the isotope ions 363.1u, 364.1u, 365.1u and 366.1u was in the ratio 100: 25.3 : 34.9 : 7.9; calculation for $\text{C}_{20}\text{H}_{22}\text{N}_2\text{ClD}_8$ gives the ratio 100: 25.2: 34.9: 8.3. Furthermore, D_7 -analogs and the D_3 -analogs were observed at masses 362.1u and 358.1u, respectively. The signals at 364u, 365u and 366u are primarily due to protonated molecules containing ^{13}C and/or ^{37}Cl isotopes instead of ^{12}C and ^{35}Cl (due to the natural distribution). This data shows that the incorporation of eight deuterium atoms was greater than 94 %.

25

Example 7: Experimental Binding Assays

Description of human D_2 binding assay

30

[0150] The assay was performed as a SPA-based competition-binding in a 50 mM Tris pH 7.4 assay buffer containing 120 mM NaCl, 5 mM KCl, 4 mM MgCl_2 , 1.5 mM CaCl_2 , 1 mM EDTA.

35

[0151] 1.5 nM ^3H -raclopride (Perkin Elmer, NET 975) was mixed with test compound before addition of 20 microg of a homogenised human D_2 receptor membrane-preparation and 0,25 mg SPA beads (WGA RPNQ 0001, Amersham) in a total volume of 90 microL. The assay plates were under agitation incubated for 60 minutes at room temperature and subsequently counted in a scintillation counter (TriLux, Wallac). The total binding, which comprised approximately 15 % of added radioligand, was defined using assay buffer, whereas the non-specific binding was defined in the presence of 10 microM haloperidol. The non-specific binding constituted approximately 10% of the total binding.

40

[0152] Data points were expressed in percent of the specific binding of ^3H -Raclopride and the IC_{50} values (concentration causing 50 percent inhibition of ^3H -raclopride specific binding) were determined by non-linear regression analysis using a sigmoidal variable slope curve fitting. The dissociation constant (K_i) was calculated from the Cheng Prusoff equation ($K_i = \text{IC}_{50}/(1+(L/K_d))$), where the concentration of free radioligand L is approximated to the concentration of added ^3H -raclopride in the assay. The K_D of ^3H -raclopride was determined to 1.5 nM from two independent saturation assays each performed with triplicate determinations.

45

Description of human D_1 binding assay

50

[0153] The assay was performed as a SPA-based competition-binding in a 50 mM Tris pH 7.4 assay buffer containing 120 mM NaCl, 5 mM KCl, 4 mM MgCl_2 , 1,5 mM CaCl_2 , 1 mM EDTA. Approximately 1 nM ^3H -SCH23390 (Perkin Elmer, NET 930) was mixed with test compound before addition of 2,5 microg of a homogenized human D_1 receptor membrane-preparation and 0,25 mg SPA beads (WGA RPNQ 0001, Amersham) in a total volume of 60 microL.

55

[0154] The assay plates were under agitation incubated for 60 minutes at room temperature before the plates were centrifuged and subsequently counted in a scintillation counter (TriLux, Wallac). The total binding, which comprised approximately 15 % of added radioligand, was defined using assay buffer whereas the non-specific binding was defined in the presence of 10 microM haloperidol.

[0155] Data points were expressed in percent of the specific binding and the IC_{50} values (concentration causing 50 percent inhibition of specific binding) and were determined by non-linear regression analysis using a sigmoidal variable slope curve fitting. The dissociation constant (K_i) was calculated from the Cheng Prusoff equation ($K_i = \text{IC}_{50}/(1+(L/K_D))$),

where the concentration of free radioligand L is approximated to the concentration of added radio-ligand in the assay.

Description of human 5-HT₂ binding

5 [0156] The experiment was carried out at Cerep Contract Laboratories (Cat. ref. # 471).

[0157] Compound (I) was also tested in an *in vivo* set up demonstrating central effects of the compound. By *in vivo* binding, the compound's *in vivo* affinity for D₂ receptors was assessed and occupancy of 60 % of the target was observed. Occupancy of D₂ receptors is closely linked to antipsychotic effects in animal models and in patients.

10 Description of *in vivo* binding to D₂ receptors in rat brain

[0158] *In vivo* binding was carried out according to Andersen et al (Eur J Pharmacol, (1987) 144:1-6) with a few modifications (Kapur S. et al, J Pharm Exp Ther, 2003, 305, 625 - 631). Briefly, 6 rats (male Wistar, 180-200 g) were treated with 20 mg/kg test compound subcutaneous 30 minutes before receiving 9.4 micro Ci [³H]-raclopride intravenously via the tail vein.

[0159] 15 minutes after the injection of the radio ligand the animals were killed by cervical dislocation, the brain quickly removed and striatum and cerebellum dissected out and homogenized in 5 mL (cerebellum in 20 mL) ice-cold buffer (50 mM K₃PO₄, pH 7.4). 1.0 mL of the homogenate was filtered through 0.1% PEI - soaked Whatman GF/C filters. This was completed within 60 seconds subsequent to the decapitation. Filters were washed 2 times with 5 mL ice-cold buffer and counted in a scintillation counter. A group of vehicle treated animals was used to determine [³H]-raclopride total binding in striatum and non-specific binding in cerebellum. The homogenate was measured for protein content by the BCA protein determination assay (Smith P.K. et al (1985) Anal. Biochem., 150: 6-85).

25 **Example 8:** Investigation of the metabolism of 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine (Compound (X)) and 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine (Compound (I))

[0160] Cryopreserved dog (male Beagle dog) hepatocytes (1 million cells/mL in suspension, 50 microL/well) were pre-incubated for 15 minutes in a 96 well plate at 37°C water bath in DMEM high glucose buffered with 1M HEPES. The cell suspension was added with 50 microL test compounds (final concentration 0.1 or 1 microM of 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1,2,2-trimethyl-piperazine (Compound (X)) or 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine (Compound (I)) and further incubated for 0, 15, 45, 75 and 120 minutes. The reaction was stopped by addition of 100 microL acetonitrile to the cell suspension, and the samples were then removed for LC-MS analysis of the desmethyl metabolite (Compound (XI)). Data were expressed as MS area relative to an internal standard.

[0161] 35 The results (Figure 5 and Figure 6) show that the amount of the desmethyl metabolite (Compound (XI)) produced in cryopreserved dog hepatocytes is lower from the deuterated form (Compound (I)) than from the parent compound (Compound (X)), both at a concentration of 0.1 micro M (Figure 5) and at a concentration of 1 micro M (Figure 6).

Example 9: Pharmacological testing of Compounds.

40 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-*d*₃-methyl-2,2-dimethyl-piperazine (Compound (I)):

[0162] 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-*d*₃-methyl-2,2-dimethyl-piperazine (Compound (I)) was tested in three *in vitro* assays for dopamine D₁, dopamine D₂ and serotonin 5-HT_{2A} affinity.

[0163] 45 The experiments were carried out as in the section Binding assays. The experimental results showed the following affinities for 4-((1R,3S)-6-chloro-3-phenyl-indan-1-yl)-1-methyl-*d*₃-2,2-dimethyl-piperazine:

$$D_1: Ki \text{ log mean} = 7.5 \text{ nM (pKi } 0.88 \text{ +/- } 0.15\text{)}$$

50 $D_2 : Ki \text{ log mean} = 34 \text{ nM (pKi } 1.54 \text{ +/- } 0.11\text{)}$

55 $5HT_{2A}: IC50 = 1.14 \text{ nM}$

[0164] 55 These binding affinities indicate that Compound (I) has biological activity likely to exert antipsychotic effect.

Pharmacological testing of Compound (II) and Compound (IV)

[0165] The experiments were carried out as described in the section "Binding assays". The experimental results for the two compounds are provided below.

5 [0166] Compounds (II) and Compound (IV) were tested in two *in vitro* assays for dopamine D₁ and dopamine D₂ affinity.

Compound (IV):

10 [0167]

D₁: Ki log mean = 26.1 nM (pKi 1.42 +/- 0.03)

15 D₂ : Ki log mean = 26.7 nM (pKi 1.43 +/- 0.04)

Compound (II):

20 [0168]

D₁: Ki log mean = 23.2 nM (pKi 1.37 +/- 0.03)

25 D₂ : Ki log mean = 26.5 nM (pKi 1.42 +/- 0.03)

[0169] These binding affinities indicate that Compound (II) and (IV) have biological activity likely to exert antipsychotic effect.

30 [0170] Compound (II) and (IV) were also tested in an *in vivo* set up demonstrating central effects of the compound. By *in vivo* binding, the compound's *in vivo* affinity for D₂ receptors was assessed and occupancy of 70% (Compound (IV)) and 75% (Compound (II)) of the target was observed. Occupancy of D₂ receptors is closely linked to antipsychotic effects in animal models and in patients.

35 [0171] Compounds (I) - (VII) and (X) were assayed in a side-by-side analysis at Cerep Contract Laboratories (Cat. Refs. # 44, 46 and 471). Results of receptor binding is listed in Table 4.

Table 4. Binding of Compounds to D₁, D₂ and 5-HT_{2A}.

Cmpd.	alternative human D ₁ receptor binding (K _i)	alternative human D ₂ receptor binding (K _i)	human 5-HT _{2A} (IC ₅₀)
(I)	0.10 nM	7.6 nM	0.37 nM; 1.14 nM*
(II)	0.20 nM	6.8 nM	1.1 nM
(III)	0.36 nM	7.6 nM	1.1 nM
(IV)	0.05 nM	10 nM	0.25 nM
(V)	0.10 nM	4.8 nM	0.61 nM
(VI)	0.10 nM	3.7 nM	0.24 nM
(VII)	0.14 nM	5.2 nM	0.33 nM
(X)	0.22 nM	7 nM	0.79 nM

* Compound (I) was tested twice in this assay.

Example 10: MetabolismInvestigations in pooled human liver microsomes (HLM)

55 [0172] Pooled human liver microsomes (50 donors, from Xenotech) were incubated with 1 microM or 10 microM of compound at 37 °C. The incubation mixture contained 50 mM Tris-HCl, 154 mM KCl, 5 mM MgCl₂ and a NADPH regenerating system (1 mM NADP⁺, 5 mM isocitric acid, 1 unit/mL isocitric dehydrogenase, from Sigma-Aldrich). The

protein concentration was 0.2 mg/mL and the final volume was 0.5 mL. Following a 10 minute pre-incubation, the reaction was initiated by adding Compound. After 0, 15, 30, 60, 90, 120 and 180 minutes, the reactions were terminated by transferring the subcellular fraction to 0.5 mL of stopping reagent containing internal standard. The incubations were carried out in triplicate. The samples were centrifuged at 4000 g (4 °C, 15 min) and the supernatants were analysed by HPLC-MS/MS. Data were expressed as MS area relative to an internal standard.

[0173] The results are shown as the mean of triplicate determinations \pm SD. Figure 7 and Figure 8 show that the amount of the desmethyl metabolite produced in human liver microsomes is lower from the deuterated form (Compound (II) and Compound (IV)) than from the non-deuterated compound (Compound (X)), both at a concentration of 1 microM (Figure 7) and at a concentration of 10 microM (Figure 8). Results for Compound (III) are shown in Figure 9. Results for Compounds (V) - (VII) are shown in Figs. 10-12, respectively. The desmethyl metabolites of compounds (II), (IV) and (X) are compounds (XX) and (XI), respectively (see Figure 13).

Investigations using recombinant human liver CYP2C19 and CYP3A4

[0174] Recombinant human liver CYP2C 19 or CYP3A4 isoenzymes (from BD biosciences) were incubated with 1 microM or 10 microM Compound (X), Compound (II) or Compound (IV) at 37 °C. The incubation mixture contained 50 mM Tris-HCl, 154 mM KCl, 5 mM MgCl₂ and a NADPH regenerating system (1 mM NADP⁺, 5 mM isocitric acid, 1 unit/mL isocitric dehydrogenase, from Sigma-Aldrich). The protein concentration was 0.5 mg/mL and the final volume was 0.5 mL. Following a 10 minutes pre-incubation, the reaction was initiated by adding Compound (X), Compound (II) and/or Compound (IV). After 0, 15, 30, 60, 90, 120 and 180 minutes the reactions were terminated by transferring the subcellular fraction to 0.5 mL of stopping reagent containing internal standard. The incubations were carried out in triplicate. The samples were centrifuged at 4000 g (4 °C, 15 minutes) and the supernatants were analyzed by HPLC-MS/MS. Data were expressed as MS area relative to an internal standard.

[0175] The results (Figure 14 and Figure 15) show that the amount of the desmethyl metabolite produced following incubation with recombinant human liver CYP2C19 enzymes is lower from the deuterated forms (Compound (II) and Compound (IV)) than from the non-deuterated compound (Compound (X)), both at a concentration of 10 micro M (Figure 14, Compound (II)) and at a concentration of 1 micro M (Figure 15, Compound (IV)). Corresponding results were obtained for Compound (II) at a concentration of 1 micro M and for Compound (IV) at a concentration of 10 micro M.

[0176] Correspondingly, the amount of the desmethyl metabolite produced by incubation with recombinant human liver CYP3A4 enzymes is lower from the deuterated forms (Compound (II) and (IV)) than from the non-deuterated compound (Compound (X)), both at a concentration of 1 micro M and 10 micro M.

Example 11: Pharmacology of Compound (IV).

PCP-Induced Hyperactivity

[0177] Compound (IV) dose-dependently reverses PCP-induced hyperactivity in mice, indicative of antipsychotic efficacy (Figure 16). Compound (IV) tartrate was administered subcutaneous (s.c.) 30 minutes before the test. PCP hydrochloride (2,3 mg/kg) was administered s.c. just before the test. Locomotor activity was measured for 60 minutes as number of beam breaks (counts). Eight to 16 male mice were used in each group. ## indicates P <0.01 versus Vehicle-PCP (One-way analysis of variance [ANOVA] followed by Bonferroni post-hoc test). PCP is blocking NMDA receptors and as such is used to model the hypo-glutamatergic state related to schizophrenia. PCP produces behavioural effects in animals reminiscent of positive, negative, and cognitive symptoms of schizophrenia patients (Jentsch, J.D. and Roth, R. H. *Neuropsychopharmacology* 1999; 20: 201-225). PCP-induced hyperactivity is commonly used as an assay for evaluation of antipsychotic compounds (Jackson, D.M. et al., *Pharmacol Biochem Behav*. 1994; 48: 465-471).

Catalepsy

[0178] Catalepsy is thought to reflect drug-induced suppression of the ability to initiate a behavioral response. The catalepsy test in rats is a common and widely used preclinical screening test for the EPS liability of potentially antipsychotic drugs. Although catalepsy is usually assessed following acute drug administration, the test has proven to be a reliable predictor for the propensity of an antipsychotic drug to induce EPS (that is, pseudo parkinsonism, dystonia) in humans (Elliott, P.J. et al., *J. Neural. Transm. Park. Dis.Dement. Sect.* 1990; 2: 79-89).

[0179] Compound (IV) dose-dependently induced catalepsy in rats suggestive of EPS liability. The minimal effective dose inducing catalepsy was 10 mg/kg (Figure 17). Compound (IV) tartrate was administered s.c. 30 minutes before the test. Eight male Sprague Dawley rats were used in each group. # indicates P <0.05, ## indicates P <0.01 versus vehicle (One-way ANOVA followed by Bonferroni post-hoc test). This dose is 100 times higher than the dose indicating antipsychotic activity (Figure 16).

Example 12: Human Pharmacokinetic Studies.

[0180] The pharmacokinetics of Compound (IV) and Compound (X) were compared in a multiple oral dose study in healthy young men. The study participants received daily doses of 3 mg Compound (IV) and 3 mg Compound (X) for 5 18 days and blood samples were collected for 24 hours (one dosing interval) after the last dose to measure the exposure of both compounds and their demethylated metabolites, Compound (XX) and Compound (XI), respectively.

[0181] For all study participants, the area under the time-plasma concentration curve for the dosing interval (AUC 0-24) for Compound (IV) was higher than that for Compound (X), mean 104 h*ng/mL vs 98 h*ng/mL. A consistent shift in the opposite direction was observed for the demethylated metabolites with mean AUC 0-24 of 117 h*ng/mL and 120 10 h*ng/ml for Compound (XX) and Compound (XI), respectively.

Example 13: Catalytic enantioselective synthesis of ketone intermediate.

[0182] This example discloses the synthesis of (S)-6-chloro-3-phenyl(d_5)-indan-1-one, Compound (XV), and (S)-6-chloro-3-phenyl-indan-1-one, Compound (XVIII).

[0183] (S)-6-chloro-3-phenyl(d_5)-indan-1-one, Compound (XV), has proven to be a valuable building block in the synthesis of deuterated variants of Compound (X) where the free phenyl group is deuterated.

General Experimental

[0184] Unless otherwise stated, all reactions were carried out under nitrogen. Reactions were monitored by thin-layer chromatography (TLC) analysis and LC-MS. All reagents were purchased and used without further purification. Spots were visualized by exposure to ultraviolet (UV) light (254 nm), or by staining with a 5 % solution of phosphomolybdenic acid (PMA) in ethanol or basic aqueous potassium permanganate ($KMnCO_4$) and then heating. Column chromatography was carried out using Merck C60 (40-63 μ m, 230-240 mesh) silica gel. NMR spectra were recorded at 500 or 600 MHz (1H NMR), and calibrated to the residual solvent peak. The following abbreviations are used for NMR data: s, singlet; d, doublet; t, triplet; m, multiplet. Coupling constants are rounded to nearest 0.5 Hz. Enantiomeric excess was determined by chiral HPLC.

LC-MS method:


[0185] Acuity UPLC BEH C 18 1.7 μ m column; 2.1 x 50 mm operating at 60°C with flow 1.2 mL/min of a binary gradient consisting of water + 0.1 % formic acid (A) and acetonitrile + 5% water + 0.1 % formic acid (B).

Chiral HPLC method:

[0186] Phenomenex Lux 5 μ Cellulose-2 column; 250 x 4.6 mm operating at 30°C with flow 0.6 mL/min of n-hexane:iso-propanol:diethylamine, 90:10:0.1.

Synthesis of (S)-6-chloro-3-phenyl(d_5)-indan-1-one (Compound (XV)) (Scheme 14)**[0187]**

Scheme 14. Synthesis of Compound (XV)



25 1-phenyl(d_5)-vinyl trifluoromethanesulfonate (XII):

[0188] To a solution of acetophenone- d_5 (1.56 g, 12.5 mmol) in CH_2Cl_2 (25.0 mL) was added trifluoromethanesulfonic anhydride (2.52 mL, 15.0 mmol) at room temperature. Then *N,N*-diisopropylethylamine (3.04 mL, 17.5 mmol) was added dropwise while the reaction mixture was cooled in an ice-water bath. The reaction mixture was allowed to warm to room temperature, and it was stirred for 1.5 h. Trifluoromethanesulfonic anhydride (0.63 mL, 3.74 mmol) was added followed by *N,N*-diisopropylethylamine (1.09 mL, 6.24 mmol). The reaction mixture was stirred for 2 hours at room temperature. Toluene (25 mL) and silica gel (5 g) was added. The mixture was concentrated *in vacuo*, and the resulting suspension was filtered through a pad of celite. The filter cake was washed with toluene (10 mL), and the filtrate was evaporated to dryness *in vacuo* to yield crude Compound (XII) (3.11 g, 82%, purity (NMR): approx. 85%) as a dark oil, that was used without further purification.

[0189] ^1H NMR (600 MHz, CDCl_3) δ _H 5.38 (d, 1H, J = 4.0 Hz), 5.62 (d, 1H, J = 4.0 Hz).

[0190] 5-chloro-2-(1-phenyl(d_5)-vinyl)benzaldehyde (XIV) (Takagi, J.; Takahashi, K.; Ishiyama, T.; Miyaura, N. *J. Am. Chem. Soc.* 2002, 124, 8001-8006; Simeone, J. P.; Sowa, J. R. Jr. *Tetrahedron* 2007, 63, 12646-12654).

[0191] To a solution of Compound (XII) (3.11 g, 10.3 mmol, purity (NMR): approx. 85%) in toluene was added triphenylphosphine (108 mg, 0.685 mmol), bis(pinacolato)diboron (2.61 g, 10.3 mmol), bis(triphenylphosphine)palladium(II) chloride (240 mg, 0.342 mmol) and potassium phenolate (1.92 g, 14.6 mmol). The reaction mixture was stirred at 50°C for 4 hours. This yielded Compound (XIII) in the mixture, which was not isolated. The mixture was cooled to room temperature, and ethanol (10 mL) and water (5 mL) was added, followed by tetrakis(triphenylphosphine)palladium(0) (495 mg, 0.428 mmol), potassium carbonate (4.73 g, 34.2 mmol) and 2-bromo-5-chlorobenzaldehyde (1.88 g, 8.56 mmol). The reaction mixture was stirred at 80°C for 16 hours. The mixture was cooled to room temperature, and partitioned between water (50 mL) and toluene (50 mL).

[0192] The organic phase was separated and washed with water (50 mL) twice, and brine. The organic phase was dried over MgSO_4 , filtered and evaporated to dryness *in vacuo*. The residue was subjected to purification by column chromatography eluting with 80:1 n-heptane:EtOAc mixture to afford Compound (XIV) (1.66 g, 74%) as an orange oil.

[0193] ^1H NMR (600 MHz, CDCl_3) δ _H 5.28 (d, 1H, J = 5 Hz), 6.00 (d, 1H, J = 0.5 Hz), 7.30 (d, 1H, J = 8.0 Hz), 7.56 (dd, 1H; J = 2.5, 8.0 Hz), 7.96 (d, 1H, J = 2.5 Hz); ^{13}C NMR (150 MHz, CDCl_3) δ _C 118.7, 126.6 (t, J = 24.0 Hz), 127.5, 128.2 (t, J = 24.0 Hz), 128.4 (t, J = 24.0 Hz), 132.5, 133.7, 134.7, 135.7, 140.3, 143.9, 144.8, 190.8; LC-MS (APPI): *m/e* calc. for $\text{C}_{15}\text{H}_7\text{D}_5\text{ClO}$ [$\text{M}+\text{H}]^+$ 248.1, found 248.1.

[0194] (S)-6-Chloro-3-phenyl(d_5)-indan-1-one (XV) (Kundu, K.; McCullagh, J. V.; Morehead, A. T. Jr. *J. Am. Chem. Soc.* 2005, 127, 16042-16043).

[0195] Hydrogen was bubbled through a N_2 -flushed solution of ((*R*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl)(norbornadiene)rhodium(I) tetrafluoroborate (37 mg, 0.0404 mmol) in acetone (7.5 mL) for 10 min at room temperature, during which the color of the solution changed from orange to more brownish red. The flask containing the solution was


subsequently flushed briefly with N_2 gas. Then a solution of (XIV) (526 mg, 2.02 mmol, purity (LC-MS): 95%) in acetone (7.5 mL) was added at room temperature. The reaction mixture was stirred for 24 hours at room temperature. The reaction mixture was mixed with silica gel and evaporated to dryness *in vacuo*. The obtained material was loaded onto a silica gel column and the product was eluted with 10:1 n-heptane:EtOAc mixture to obtain Compound (XV) (495 mg, 96%, 96.0% ee) as a solid.

[0196] 1H NMR (500 MHz, $CDCl_3$) δ_H 2.72 (dd, 1H, J = 4.0, 19.5 Hz), 3.27 (dd, 1H, J = 8.0, 19.5 Hz), 4.55 (dd, 1H, J = 4.0, 8.0 Hz), 7.21 (d, 1H, J = 8.0 Hz), 7.52 (dd, 1H, J = 2.0, 8.0 Hz), 7.77 (d, 1H, J = 2.0 Hz); ^{13}C NMR (125 MHz, $CDCl_3$) δ_C 44.0, 47.2, 123.2, 126.8 (t, J = 24.0 Hz), 127.3 (t, J = 24.0 Hz), 128.7 (t, J = 24.0 Hz), 134.4, 135.1, 138.2, 142.9, 156.0, 206.4; LC-MS (APPI): *m/e* calc. for $C_{15}H_7D_5ClO$ [M+H]⁺ 248.1, found 247.6.

Synthesis of (S)-6-chloro-3-phenyl-indan-1-one (XVIII) (Scheme 15)

[0197]

Scheme 15. Synthesis of Compound (XVIII)



(*E*)-1-(5-chloro-2-hydroxyphenyl)-3-phenylprop-2-en-1-one (XVI):

[0198] To an ice-cooled solution of sodium hydroxide (2.34 g, 58.6 mmol) in water (17.0 mL) was added benzaldehyde (0.746 g, 7.03 mmol) and then a solution of 5-chloro-2-hydroxyacetophenone (1.00 g, 5.86 mmol) in methanol (17.0 mL). The reaction mixture was allowed to warm to room temperature, and it was stirred for 24 hours. The bulk of the organic solvent was removed by evaporation *in vacuo*. The aqueous residue was extracted with EtOAc (3 x 30 mL). The combined extracts were washed with water (50 mL) and brine (50 mL), dried over $MgSO_4$, filtered and evaporated to dryness *in vacuo*. The residue was dissolved in a minimum volume of CH_2Cl_2 , and n-pentane was added which resulted in precipitation. The obtained suspension was filtered and the precipitate was washed with little cold pentane, and dried *in vacuo* to afford Compound (XVI) (695 mg, 46%) as an orange solid.

[0199] 1H NMR (500 MHz, $CDCl_3$) δ_H 6.22 (d, 1H, J = 9.0 Hz), 6.80 (dd, 1H, J = 3.0, 9.0 Hz), 7.33 (t, 1H, J = 7.5 Hz), 7.38-7.42 (m, 4H), 7.60 (d, 2H, J = 7.5 Hz); 8.63 (d, 1H, J = 16.0 Hz); ^{13}C NMR (125 MHz, $CDCl_3$) δ_C 110.6, 125.2, 127.8, 128.1, 128.8, 128.9, 129.4, 129.6, 133.0, 136.4, 137.1, 174.5, 188.2.

Trifluoromethanesulfonic acid 4-chloro-2-((*E*)-(3-phenyl-acryloyl))-phenyl ester (XVII):

[0200] To a solution of Compound (XVI) (517 mg, 2.00 mmol) in CH_2Cl_2 (10.0 mL) was added *N,N*-diisopropylethylamine (697 μ L, 4.00 mmol). Trifluoromethanesulfonic anhydride (437 μ L, 2.60 mmol) was added dropwise at 0°C. The reaction mixture was stirred for 45 min at 0°C. Sat. aq. NH_4Cl (5 mL) and water (10 mL) was added, and the mixture was stirred for 5 minutes. The organic phase was separated, and the aqueous phase was extracted with CH_2Cl_2 (10 mL). The combined extracts were dried over $MgSO_4$, filtered and evaporated to dryness *in vacuo*. The residue was purified by column chromatography eluting with 4:1 n-heptane:EtOAc to yield (XVII) (757 mg, 97%) as an oil.

[0201] 1H NMR (500 MHz, $CDCl_3$) δ_H 7.16 (d, 1H, J = 16.0 Hz), 7.34 (d, 1H, J = 9.0 Hz), 7.40-7.47 (m, 3H), 7.57 (dd, 1H, J = 2.5, 9.0 Hz), 7.60-7.62 (m, 2H), 7.69 (d, 1H, 16.0 Hz), 7.72 (d, 1H, J = 2.5 Hz); ^{13}C NMR (125 MHz, $CDCl_3$) δ_C 124.1, 124.2, 129.0, 129.2, 130.7, 131.5, 132.8, 134.1, 134.6, 145.2, 147.8, 188.4.

[0202] (S)-6-Chloro-3-phenyl-indan-1-one (XVIII) (Minatti, A.; Zheng, X.; Buchwald, S. L. J. Org. Chem. 2007, 72, 9253-9258).

[0203] To a solution of Compound (XVII) (195 mg, 0.500 mmol) in DMF (2.0 mL) was added proton-sponge (214 mg, 1.00 mmol), palladium acetate (6 mg, 0.025 mmol) and (R)-3,5-XylMeOBIPHEP (35 mg, 0.05 mmol) at rt. The reaction mixture was stirred at 85°C for 45 h. The mixture was cooled to rt, and diluted with TBME (15 mL). The mixture was washed three times with water (3 x 20 mL), and the organic phase was dried over MgSO₄, filtered and evaporated to dryness *in vacuo*. The residue was subjected to column chromatography eluting with 10:1 n-heptane:EtOAc to yield Compound (XVII) (94 mg, 77%, 64.0% ee).

[0204] ¹H NMR (600 MHz, CDCl₃) δ_H 2.71 (dd, 1H, *J* = 4.0, 19.5 Hz), 3.25 (dd, 1H, *J* = 8.0, 19.5 Hz), 4.54 (dd, 1H, *J* = 4.0, 8.0 Hz), 7.10 (d, 2H, *J* = 7.0 Hz), 7.20 (d, 1H, *J* = 8.0 Hz), 7.25 (t, 1H, *J* = 7.5 Hz), 7.31 (t, 2H, *J* = 7.5 Hz), 7.50 (dd, 1H, *J* = 2.0, 8.0 Hz), 7.75 (d, 2H, *J* = 2.0 Hz); ¹³C NMR (150 MHz, CDCl₃) δ_C 44.1, 47.2, 123.3, 127.3, 127.6, 128.3, 129.1, 134.4, 135.2, 138.3, 143.1, 156.1, 204.5.

Enantioenrichment of Compound (XVIII) by Reprecipitation


[0205] Compound (XVII) (940 mg, 3.87 mmol, 96% ee) was dissolved in a minimum volume of boiling ethanol (99% v/v). The resulting solution was allowed to cool slowly to rt by placing the glass flask containing the solution in the air. A precipitate formed that was filtered from solution to yield Compound (XVIII) (700 mg, 99.9% ee, 74%). A second crop of Compound (XVIII) could be obtained by cooling the filtrate in the freezer (-8°C) to yield Compound (XVIII) (80 mg, 98.6% ee, 9%).

[0206] Analytical data (NMR and LC-MS) for Compound (XVIII) were the same as those reported above.

Example 14: Large scale production of Compound (IV)

[0207] The following process was developed for the large scale production of the tartrate salt of Compound (IV)

Scheme 16: Synthesis of rac-trans-1-(6-Chloro-3-phenyl(*d*₅)-indan-1-yl)-3,3-dimethyl-piperazine maleate



Procedure:

[0208]

- 1.) 2.01 kg (16.9 mol) thionylchloride and 7.2 kg tetrahydrofuran are mixed and the reaction is cooled to 10-15 °C
- 2.) a solution of 2.76 kg (11.1 mol) (XXII) in 7.2 kg THF is slowly added and after completion 5.9 kg tetrahydrofuran is added
- 3.) the reaction is stirred at 15 °C for approximately 90 hours
- 4.) 16.7 kg water is cooled to 11 °C and added slowly to the reaction, afterwards 7.8 kg 27.7% aqueous sodium hydroxide is added slowly, followed by 10 kg ethylacetate
- 5.) the mixture is stirred for 20-40 minutes
- 6.) the phases are separated and the organic phase is reduced to a volume of approximately 6L
- 7.) 16 kg methyl isobutylketone is added and the volume is reduced to approximately 8 L
- 8.) 1.58 kg (11.4 mol) potassium carbonate, 1.69 kg (14.8 mol) 2,2-dimethylpiperazin and 13.6 kg methyl isobutyl

ketone are added

9.) the reaction is stirred 35 hours at 90-95 °C

10.) after cooling to room temperature 11 kg water is added and the mixture is stirred for 30 - 60 minutes

11.) the phases are separated. 13.7 kg water is added to the organic phase and the mixture is stirred slowly for 30 - 60 minutes

12.) the phases are separated and the organic phase is blank filtered

13.) 5 kg methyl isobutyl ketone, 7.8 kg water and 5.9 kg 36% aqueous hydrogen chloride are added and the mixture is stirred at 50 °C for 30 - 60 minutes.

14.) the phases are separated. 8 kg methyl isobutyl ketone is added to the water phase and the mixture is cooled to 10-15 °C

15.) a mixture of 3.5 kg methyl isobutyl ketone and 7.8 kg 25% aqueous ammonia are slowly added to the mixture and the reaction is stirred at 20-25 °C for 60 - 90 minutes

16.) the phases are separated and the organic phase is washed with 10.5 kg water

17.) the organic phase is reduced to 8 l

18.) 1.19 kg (10.25 mol) maleic acid and 9 kg methyl isobutyl ketone are added and the reaction is afterwards warmed to 75-80 °C.


19.) after cooling to 10-15 °C the precipitate is filtered off and washed with 10 kg methyl isobutyl ketone

20.) the solid is dried in a vacuum oven at 50 °C for approximately 20 hours to give 3.47 kg (68% yield) of (XXV) maleate.

[0209] NMR data for (XXV) maleate:

[0210] 1H-NMR (dmso-d6, 600 MHz, ppm): 8.60 (bs, 2H, maleic acid), 7.39 (d, 1H, J =1.6 Hz), 7.29 (dd, 1H, J =8.0 Hz, J =1.8 Hz), 6.98 (d, 1H, J =8.2 Hz), 6.04 (s, 2H, maleic acid), 4.56 (dd, 1H, J =8.1 Hz, J =4.9 Hz), 4.48 (dd, 1H, J =8.6 Hz, J =6.2 Hz), 3.37 (bs, 1H), 3.16 (bs, 2H), 2.77 (bs, 1H), 2.58-2.50 (m, 3H), 2.31 (d, 1H, J =12.0 Hz), 2.12 (ddd, 1H, J =13.8 Hz, J =8.0 Hz, J =6.0 Hz), 1.33 (s, 3H), 1.31 (s, 3H).

Scheme 17: Synthesis of rac-trans-1-(6-chloro-3-phenyl(d_5)-indan-1-yl)-1(d_3), 2,2-trimethyl-piperazine succinate



Procedure:

[0211]

1.) 1.1 kg (2.38 mol) (XXV) maleate, 11 L methyl tertbutyl ether, 1.8 L water and 1 L 25% aqueous ammonia are stirred for 1 - 2 hours.

2.) the phases are separated and the organic phase is washed with two times 2 l. water.

3.) a solution of 254 g (3.85 mol) 85% aqueous potassium hydroxide and 1.5 L water is added to the organic phase, followed by addition of 450 g (3.11 mol) methyl(^{d3})iodide (CD_3I)

4) the reaction is stirred at 20-25 °C for 16 - 24 hours.

5.) 2 L water are added and the precipitating by-product is filtered off.

6.1081 water and 0.2125% aqueous ammonia are added to the filtrate and the mixture is stirred for 20 - 40 minutes.

7.) the phases are separated and the organic phase is washed with 2 L water.

8.) the phases are separated and 38 g (0.48 mol) acetylchloride is added to the organic phase which is stirred for 20 - 40 minutes

9.) 0.8 L water and 0.2 L 25% aqueous ammonia are added and the mixture is stirred for 20 - 40 minutes

10.) the phases are separated and the organic phase is washed with 2 L water

5 11.) the organic phase is reduced to dryness and acetone is added

12.) 225 g (1.91 mol) succinic acid and acetone are added so that the reaction volume is approximately 6 - 6.5 L

13.) The reaction is warmed to reflux and afterwards cooled to 5-10 °C


14.) The precipitate is filtered off and washed with 1 L acetone

15.) the solid is dried in a vacuum oven at 50 °C for more than 16 hours to give 630 g (55% yield) of (XXVII) succinate

10 NMR data for (XXVII) succinate:

[0212] 1H-NMR (dmso-d6, 600 MHz, ppm): 7.33 (d, 1H, J=1.9 Hz), 7.26 (dd, 1H, J=8.1 Hz, J=2.0 Hz), 6.95 (d, 1H, J=8.0 Hz), 4.46 (dd, 1H, J=8.0 Hz, J=5.1 Hz), 4.46 (dd, 1H, J=8.8 Hz, J=5.8 Hz), 2.65-2.56 (m, 4H), 2.46-2.41 (m, 1H), 2.37 (s, 4H, succinic acid), 2.31 (bs, 1H), 2.13 (d, 1H, J=10.9 Hz), 2.02 (ddd, 1H, J=13.7 Hz, J=7.8 Hz, J=6.0 Hz), 1.04 (s, 3H), 1.02 (s, 3H).

20 Scheme 18: Synthesis of 4-((1R,3S)-6-chloro-3-phenyl(*d*5)-indan-1-yl)-1(*d*3),2,2-trimethyl-piperazine L(+)-tartrate



Procedure:

[0213]

40 1.) 1.0 kg (2.08 mol) (XXVII) succinate, 8 L ethyl acetate, 2L water and 1L 25% aqueous ammonia are stirred for 0.5 - 1 hours

2.) the phases are separated and the organic phase is washed with 2 L water

3.) the organic phase is reduced to approximately 1.5 L

45 4.) 10 L acetone and 312 g (2.08 mol) L(+)-tartaric acid are added

5.) the reaction is warmed to reflux and afterwards cooled to 5-10 °C

6.) the precipitate is filtered off, washed with 1.2 L acetone

7.) the wet filtercake is recharged and 11 L absolute ethanol are added

8.) the reaction is warmed to reflux and afterwards cooled to 5-10 °C

50 9.) the precipitate is filtered off and washed with 1.2 L absolute ethanol

10.) the solid is dried in a vacuum oven at 50 °C for more than 16 hours to give 395 g (37% yield) of (IV) L(+)-tartrate

NMR data for (IV) L(+)-tartrate:

55 [0214] 1H-NMR (dmso-d6, 600 MHz, ppm): 7.36 (s, 1H), 7.27 (d, 1H, J=8.2 Hz), 6.96 (d, 1H, J=8.2 Hz), 4.50 (dd, 1H, J=8.0 Hz, J=5.1 Hz), 4.45 (dd, 1H, J=8.5 Hz, J=5.8 Hz), 4.07 (s, 2H, tartrate), 2.95 (bs, 1H), 2.77 (bs, 1H), 2.61-2.50 (m, 3H), 2.31 (d, 1H, J=11.7 Hz), 2.04 (ddd, 1H, J=13.7 Hz, J=7.8 Hz, J=6.0 Hz) 1.21 (s, 3H), 1.18 (s, 3H).

Example 15: Physico-Chemical characterization of salts of Compound (IV)pK_a and log P/D of Compound (IV)

5 [0215] pK_a was determined by potentiometric titration of the base at ion strength 0.16 using MeOH as co-solvent. Three series of three repeated titrations on the same solution of the sample was performed in a conventional way from low to high pH and a difference curve was created from each of these titrations by blank subtraction. The apparent pK_a-value at each MeOH:water ratio is calculated from the difference curves, and the pK_a value is determined by extrapolation to zero MeOH content.

10 [0216] The lower pK_a value is too low to be determined by potentiometric titration as data only were found reliable down to ~3. The high pK_a was determined to be 8.9 + 0.1

[0217] The lower pK_a was determined by Dip Probe Absorption Spectroscopy detection during titration of the base at ion strength 0.16 using MeOH as co-solvent. The change in absorption spectra as a function of ionisation is used to calculate the pK_a-value. Two series of three repeated titrations on the same solution of the sample was performed from 15 low to high pH, with a photo diode array as additional detection. The apparent pK_a-value at each MeOH:water ratio is calculated by Target factor analysis on the change in absorption spectra, and the pK_a value is determined by extrapolation to zero MeOH content.

Result: The lower pK_a was determined to be 2.5+ 0.1

20 [0218] The logD profile was determined by titration at 27°C and ion strength of approx. 0.16. A series of three repeated titrations on the same sample in solution was performed, from low to high pH. The first titration was performed with a small amount of n-octanol present in the solution, the second and third with increasing amounts.

[0219] A difference curve was created from each of these titrations by blank subtraction, and from these difference curves, the apparent pK_a values (p_oK_a) were calculated. From the change in the apparent pK_a values (ΔpK_a) with the n-octanol:water ratio combined with the real pK_a value, the LogP value was calculated and the LogD profile was derived. The following values were determined: Log P = 5.4+ 0.4 and Log D_{7.4} = 3.9+ 0.4

Melting point determined by DSC

30 [0220] The melting point of the (R,R)-hydrogen tartrate salt of Compound (IV) was determined using differential scanning calorimetry (DSC), using a TA instruments DSC Q1000 heating the sample 5°/minute. The sample was placed in a covered pan with a punched pinhole.

[0221] The melting is characterized by onset and peak temperature of the melting endotherm, and the enthalpy of fusion is calculated from the area of the peak. Based on the DSC thermogram an onset temperature of 187.4°C and a peak maximum at 189.4°C was found. The enthalpy of fusion was 96 J/g corresponding to 49 kJ/mol, however the thermogram is indicative that the melting happens under decomposition meaning that the enthalpy probably contain energy other than melting.

Solubility

40 [0222] Solubility of the (R,R)-hydrogen tartrate salt of Compound (IV) was measured in aqueous solutions and in cyclodextrins with the following results (Table 5):

45 Table 5. Solubility of (R,R)-hydrogen tartrate salt of Compound (IV).

Solvent	Meas. conc. (mg base/ml)	pH
Hydrogen tartrate in water, 5 °C	3.1	3.25
Hydrogen tartrate in water, RT	4.0	3.15
Hydrogen tartrate in water, 37 °C	6.6	3.08
Solvent	Meas. conc. (mg base/ml)	pH
10%HPβCD	25.2	3.59
5%HPβCD, at RT	15.5	3.61

(continued)

Solvent	Meas. conc. (mg base/ml)	pH
5%HP β CD, at 5°C	12	

Polymorphism

[0223] One solvent free crystal form of the tartrate has been isolated. The XRPD of this form is shown in Figure 18, and designated herein as "polymorph A".

Salts of Compound (IV)

[0224] Four salts were prepared by precipitation of Compound (IV) from 99% EtOH.


[0225] Analytical data are given in the table below (Table 6).

Table 6. Data for salts of Compound (IV)

Salt	DSC (T_{onset} °C)	Solubility (mg/ml)	pH
Dihydrogen phosphate	Degradation at 250°C	1.4	4.67
Hydrogen fumarate	202.7°C	1.2	4.10
Hydrogen maleate	150.4°C	1.2	4.94
Hydrogen malonate	145.0°C followed by degradation	9.5	4.08
Hydrogen tartrate	187°C	4.0	3.15
Base	59.9	0.1	7.6

Claims

1. A compound of formula Y:



wherein,

R¹ - R¹⁰ are independently hydrogen or deuterium, wherein R⁶-R¹⁰ are each deuterium, wherein at least one of R¹-R¹⁰ comprises at least about 50% deuterium, or a pharmaceutically acceptable acid addition salt thereof.

2. The compound of claim 1, wherein R³-R⁵ are each hydrogen.

3. The compound of claim 1, wherein R³-R⁵ are each deuterium.


4. The compound of claim 2, wherein the compound is

5

10

15

20



(II).


25 5. The compound of claim 2, wherein the compound is

30

35

40

45



(VII).

6. The compound of claim 3, wherein the compound is

50


55

5

10

15

20




25

30

35

40



45

8. The compound of claim 1, wherein R¹ and R² are each deuterium.

9. The compound of claim 8, wherein R³-R⁵ are each deuterium.

50


10. The compound of claim 8, wherein R³-R⁵ are each hydrogen

11. The compound of any of claims 1-10, wherein at least about 85% of the compound has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.


55

12. The compound of any of claims 1-11, wherein at least about 90% of the compound has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.

13. The compound of claim 1, wherein the compound is the hydrogen tartrate salt of



14. The compound of claim 13, wherein the compound exists in polymorphic form having an XRPD diffraction pattern as indicated in Figure 18.
15. The compound of claim 13, wherein at least about 85% of the compound has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.
16. A pharmaceutical composition comprising the compound of any of claims 1 to 15 and one or more pharmaceutically acceptable carriers, diluents, or excipients.
17. The pharmaceutical composition of claim 16, wherein the compound is the hydrogen tartrate salt of



50 18. The composition of claims 16 or 17, wherein the carrier comprises hydroxypropyl- β -cyclodextrin in water, and wherein
at least about 85% of the compound has a deuterium atom at each position designated as deuterium, and any atom
not designated as deuterium is present at about its natural isotopic abundance.


55 19. Use of the compound of any of claims 1-15, or the composition of any of claims 16-18 for the manufacture of a
medicament for treatment of psychosis, other diseases involving psychotic symptoms, psychotic disorders or dis-
eases that present with psychotic symptoms.

20. Use according to claim 19, wherein the psychosis or disease involving psychotic symptoms is schizophrenia, schizopreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, bipolar disorder, or mania in bipolar disorder.

5 21. Use according to any of claims 19-20, further comprising a compound selected from the group consisting of sertindole, olanzapine, risperidone, quetiapine, aripiprazole, haloperidol, clozapine, ziprasidone and osanetant.

10 22. Use according to any of claims 19-20 wherein the psychosis or disease involving psychotic symptoms is schizophrenia.


15 23. Use according to any of claims 19-20, wherein the psychosis or disease involving psychotic symptoms is schizophrenia, wherein the pharmaceutical composition comprises an effective amount of hydrogen tartrate salt of



20 and hydroxypropyl- β -cyclodextrin in water, and wherein at least about 85% of (IV) has a deuterium atom at each position designated as deuterium, and any atom not designated as deuterium is present at about its natural isotopic abundance.

25


30 24. A compound of formula




35

40 25. A process for the preparation of compound

45




comprising treating compound (XIV) with [(S)-BINAP]Rh(I)BF₄.




26. The process of claim 25, wherein [(S)-BINAP]Rh(I)BF₄ is used in a catalytic amount.

30 27. A process for the preparation of compound (XIV) comprising a) treatment of



with bis(pinacolato) diboron, and b) treatment with 2-bromo-5-chlorobenzaldehyde.

45 28. The process of claim 27, wherein treatment of



with bis(pinacolato) diboron further comprises addition of Pd(II).


29. The process of claim 28, wherein treatment with 2-bromo-5-chlorobenzaldehyde further comprises addition of Pd(0).

30. A process of the preparation of compound (1*R*,3*S*)-(IV) tartrate comprising, treatment of racemic trans-1-(6-chloro-3-phenyl(d₅)-indan-1-yl)-1(*d*₃), 2, 2-trimethyl-piperazine with L-(+)-tartaric acid.

5 31. The process of claim 30, wherein racemic *trans*-1-(6-chloro-3-phenyl(d₅)-indan-1-yl)-1(*d*₃), 2, 2-trimethyl-piperazine is generated from the corresponding succinate salt thereof.

Patentansprüche

10 1. Eine Verbindung mit der Formel Y:



30 (Y)


wobei,

R¹ - R¹⁰ unabhängig voneinander Wasserstoff oder Deuterium sind, wobei R⁶ - R¹⁰ jeweils Deuterium sind, wobei mindestens ein von R¹ - R¹⁰ zumindest ungefähr 50% Deuterium umfassen, oder ein pharmazeutisch verträgliches Säureadditionssalz davon.

35 2. Die Verbindung nach Anspruch 1, wobei R³ - R⁵ jeweils Wasserstoff sind.

40 3. Die Verbindung nach Anspruch 1, wobei R³ - R⁵ jeweils Deuterium sind.

4. Die Verbindung nach Anspruch 2, wobei die Verbindung



55 (1*R*,3*S*) - (II)

ist.

5. Die Verbindung nach Anspruch 2, wobei die Verbindung

5

10

15

20

(1*R*, 3*S*) - (VII)

ist.

6. Die Verbindung nach Anspruch 3, wobei die Verbindung

25

30

35

40

(1*R*, 3*S*) - (IV)

ist


45 7 Die Verbindung nach Anspruch 3, wobei die Verbindung

50

55

5

10



15

(1*R*, 3*S*) - (VI)

ist.

8. Die Verbindung nach Anspruch 1, wobei R¹ und R² jeweils Deuterium sind.

9. Die Verbindung nach Anspruch 8, wobei R³ - R⁵ jeweils Deuterium sind.

10. Die Verbindung nach Anspruch 8, wobei R³ - R⁵ jeweils Wasserstoff sind.

11. Die Verbindung nach irgendeinem der Ansprüche 1 - 10, wobei mindestens etwa 85 % von der Verbindung ein Deuteriumatom an jeder als Deuterium bezeichneten Position hat, und irgendein Atom, welches nicht als Deuterium beschrieben ist, über seine natürliche Isotopenhäufigkeit vorhanden ist.

12. Die Verbindung nach irgendeinem der Ansprüche 1 - 11, wobei mindestens etwa 90 % von der Verbindung ein Deuteriumatom an jeder als Deuterium bezeichneten Position hat, und irgendein Atom, welches nicht als Deuterium beschrieben ist, über seine natürliche Isotopenhäufigkeit vorhanden ist.


13. Die Verbindung nach Anspruch 1, wobei die Verbindung das Hydrogentartrate Salz von

35

40

45

50

(1*R*, 3*S*) - (IV)

ist.


14. Die Verbindung nach Anspruch 13, wobei die Verbindung in einer polymorphen Form existiert, aufweisend ein XRPD

Beugungsmusters, wie in Figur 18 gezeigt ist.

5 15. Die Verbindung nach Anspruch 13, wobei mindestens etwa 85 % von der Verbindung ein Deuteriumatom an jeder als Deuterium bezeichneten Position hat, und irgendein Atom, welches nicht als Deuterium beschrieben ist, ist über seine natürliche Isotopenhäufigkeit vorhanden.

10 16. Eine pharmazeutische Zusammensetzung, umfassend die Verbindung nach irgendeinem der Ansprüche 1 bis 15 und ein oder mehrere pharmazeutisch verträgliche Träger, Verdünnungsmittel oder Arzneistoffträger.

10 17. Die pharmazeutische Zusammensetzung nach Anspruch 16, wobei die Verbindung das Hydrogentartrate Salz von



(1*R*, 3*S*) - (IV)

1

35 18. Die Zusammensetzung nach Anspruch 16 oder 17, wobei der Träger Hydroxypropyl- β -cyclodextrin in Wasser umfasst, und wobei mindestens etwa 85% von der Verbindung ein Deuteriumatom an jeder als Deuterium bezeichneten Position hat, und irgendein Atom, welches nicht als Deuterium beschrieben ist, ist über seine natürliche Isotopenhäufigkeit vorhanden.

40 19. Verwendung von der Verbindung nach irgendeinem der Ansprüche 1 - 15, oder der Zusammensetzung nach irgendeinem der Ansprüche 16 - 18 zur Herstellung eines Medikaments zur Behandlung von Psychose, anderer Erkrankungen, die psychotische Symptome involvieren, psychotischen Störungen oder Erkrankungen, die psychotische Symptome zeigen.

45 20. Verwendung gemäß Anspruch 19, wobei die Psychose oder Krankheit, welche psychotische Symptome involviert, Schizophrenie, schizophrene Störung, schizoaffektive Störung, wahnhafte Störung, kurze psychotische Störung, geteilte psychotische Störung, bipolare Störung oder Manie bei bipolare Störung ist.

50 21. Verwendung nach irgendeinem der Ansprüche 19 - 20, weiter umfassend eine Verbindung ausgewählt von der Gruppe bestehend aus Sertindol, Olanzapin, Risperidon, Quetiapin, Aripiprazol, Haloperidol, Clozapine, Ziprasidon und Olanzapin.


55 22. Verwendung nach irgendeinem der Ansprüche 19 - 20, wobei die Psychose oder Krankheit, welche psychotische Symptome involviert, Schizophrenie ist.

55 23. Verwendung nach irgendeinem der Ansprüche 19 - 20, wobei die Psychose oder Krankheit, welche psychotische Symptome involviert, Schizophrenie ist, wobei die pharmazeutische Zusammensetzung eine wirksame Menge an Hydrogentartrat Salz von

5

10

15

(1*R*, 3*S*) - (IV)

20


und Hydroxypropyl- β -Cyclodextrin in Wasser umfasst, und wobei mindestens etwa 85% von (IV) ein Deuteriumatom an jeder als Deuterium bezeichnete Position hat, und irgendein Atom, welches nicht als Deuterium beschrieben ist, über seine natürliche Isotopenhäufigkeit vorhanden ist.

24. Eine Verbindung von der Formel

25

30

35




(S) - (XV).

25. Ein Verfahren zur Herstellung einer Verbindung

40

45

50




(S) - (XV)

umfassend Behandeln der Verbindung (XIV) mit [(S)-BINAP]Rh(I)BF₄.

55

5

10



(XIV)


26. Das Verfahren nach Anspruch 25, wobei $[(S)\text{-BINAP}]\text{Rh(I)}\text{BF}_4$ in einer katalytischen Menge verwendet wird.

15

27. Ein Verfahren zur Herstellung der Verbindung (XIV), umfassend a) die Behandlung von

20

25




(XII)

30 mit Bis (pinacolato)diboron, und b) Behandlung mit 2-Brom-5-chlor-benzaldehyd.

28. Das Verfahren nach Anspruch 27, wobei die Behandlung des

35

40



(XII)

mit Bis (pinacolato)diboron weiter die Zugabe von Pd(II) umfasst.

45 29. Das Verfahren nach Anspruch 28, wobei die Behandlung mit 2-Brom-5-chlor-benzaldehyd weiter einen Zusatz von PD (0) umfasst.


30. Das Verfahren von der Herstellung der Verbindung (1*R*, 3*S*) - (IV) tartrat die Behandlung von racemischem *trans*-1-(6-chlor-3-phenyl (d_5)-indan-1-yl)-1 (d_3),2,2-trimethyl-piperazin mit L- (+) Weinsäure umfasst.

50

31. Das Verfahren nach Anspruch 30, wobei racemische *trans*-1-(6-chlor-3-phenyl (d_5)-indan-1-yl)-1 (d_3),2,2-trimethyl-piperazin aus dem entsprechenden Succinatsalz davon erzeugt ist.

55 Revendications

1. Composé de formule Y :




dans lequel :

R^1 à R^{10} sont indépendamment de l'hydrogène ou du deutérium, dans lequel R^6 à R^{10} sont chacun du deutérium, dans lequel au moins l'un de R^1 à R^{10} comprend au moins environ 50 % de deutérium, ou son sel d'addition avec un acide pharmaceutiquement acceptable.


2. Composé selon la revendication 1, dans lequel R^3 à R^5 sont chacun de l'hydrogène.

3. Composé selon la revendication 1, dans lequel R^3 à R^5 sont chacun du deutérium.


4. Composé selon la revendication 2, dans lequel le composé est le :




5. Composé selon la revendication 2, dans lequel le composé est le :



6. Composé selon la revendication 3, dans lequel le composé est le :



7. Composé selon la revendication 3, dans lequel le composé est le :



8. Composé selon la revendication 1, dans lequel R¹ et R² sont chacun du deutérium.


9. Composé selon la revendication 8, dans lequel R³ à R⁵ sont chacun du deutérium.

10. Composé selon la revendication 8, dans lequel R³ à R⁵ sont chacun de l'hydrogène.

5 11. Composé selon l'une quelconque des revendications 1 à 10, dans lequel au moins environ 85 % du composé ont une atome de deutérium dans chaque position désignée par deutérium et tout atome qui n'est pas désigné comme étant du deutérium est présent à peu près dans son abondance isotopique naturelle.

10 12. Composé selon l'une quelconque des revendications 1 à 11, dans lequel au moins environ 90 % du composé ont un atome de deutérium dans chaque position désignée par deutérium et tout atome qui n'est pas désigné par deutérium est présent à peu près dans son abondance isotopique naturelle.

13. Composé selon la revendication 1, dans lequel le composé est un sel d'hydrogénotartrate de :




14. Composé selon la revendication 13, dans lequel le composé existe sous forme polymorphe ayant un motif de diffraction XRPD comme indiqué sur la Fig. 18.

15. Composé selon la revendication 13, dans lequel au moins environ 85 % du composé ont un atome de deutérium dans chaque position désignée par deutérium et tout atome qui n'est pas désigné par deutérium est présent à peu près dans son abondance isotopique naturelle.

16. Composition pharmaceutique comprenant le composé selon l'une quelconque des revendications 1 à 15 et un ou plusieurs véhicules, diluants ou excipients pharmaceutiquement acceptables.

17. Composition pharmaceutique selon la revendication 16, dans lequel le composé est le sel d'hydrogénotartrate de :



18. Composition selon la revendication 16 ou la revendication 17, dans laquelle le véhicule comprend de l'hydroxypropyl-

β -cyclodextrine dans de l'eau et dans laquelle au moins environ 85 % du composé ont un atome de deutérium dans chaque position désignée par deutérium et tout atome non désigné par deutérium est présent à peu près dans son abondance isotopique naturelle.


5 **19.** Utilisation du composé selon l'une quelconque des revendications 1 à 15 ou composition selon l'une quelconque des revendications 16 à 18 pour la fabrication d'un médicament pour le traitement de la psychose, d'autres maladies impliquant des symptômes psychotiques, des troubles psychotiques ou des maladies qui se présentent avec des symptômes psychotiques.

10 **20.** Utilisation selon la revendication 19, dans laquelle la psychose ou la maladie impliquant des symptômes psychotiques est la schizophrénie, un trouble schizophréniforme, un trouble schizo-affectif, un trouble délirant, un trouble psychotique bref, un trouble psychotique partagé, un trouble bipolaire ou une manie en trouble bipolaire.

15 **21.** Utilisation selon l'une quelconque des revendications 19 à 20, comprenant en outre un composé choisi dans le groupe constitué du sertindole, de l'olanzapine, de la rispéridone, de la quétiapine, de l'aripiprazole, de l'halopéridol, de la clozapine, de la ziprasidone et de l'osanetant.


20 **22.** Utilisation selon l'une quelconque des revendications 19 à 20, dans laquelle la psychose ou la maladie impliquant des symptômes psychotiques est la schizophrénie.

25 **23.** Utilisation selon l'une quelconque des revendications 19 à 20, dans laquelle la psychose ou la maladie impliquant des symptômes psychotiques est la schizophrénie, dans laquelle la composition pharmaceutique comprend une quantité efficace de sel d'hydrogénotartrate de :




et de l'hydroxypropyl- β -cyclodextrine dans de l'eau et dans laquelle au moins environ 85 % de (IV) ont un atome de deutérium à chaque position désignée par deutérium et tout atome qui n'est pas désigné par deutérium est présent à peu près dans son abondance isotopique naturelle.

45 **24.** Composé de formule :



56 **25.** Procédé de préparation de composé :

5



10


(S)-(XV)

comportant le traitement du composé (XIV) avec du $[(S)\text{-BINAP}]\text{Rh(I)}\text{BF}_4$,

15

20

25



(XIV)

26. Procédé selon la revendication 25, dans lequel du $[(S)\text{-BINAP}]\text{Rh(I)}\text{BF}_4$ est utilisé en quantité catalytique.

30


27. Procédé de préparation du composé (XIV) comportant

a) le traitement de :

35

40

45




(XII)

avec du bis(pinacolato)dibore et b) un traitement avec du 2-bromo-5-chlorobenzaldéhyde.

50

28. Procédé selon la revendication 27, dans lequel le traitement de :

55



avec du bis(pinacolato)dibore comprend en outre l'addition de Pd(II).

15 29. Procédé selon la revendication 28, dans lequel le traitement avec du 2-bromo-5-chlorobenzaldéhyde comprend en outre l'addition de Pd(0).

30. Procédé de préparation du composé (1R,3S)-(IV)tartrate comprenant le traitement de *trans*-1-(6-chloro-3-phényleindan-1-yl)-1(d₃),2,2-triméthyl-pipérazine racémique avec de l'acide L-(+)tartrique.

20 31. Procédé selon la revendication 30, dans lequel la *trans*-1-(6-chloro-3-phényleindan-1-yl)-1(d₃), 2,2-triméthyl-pipérazine racémique est générée à partir de son sel de succinate correspondant.

25

30

35

40

45

50

55




Fig. 1




Fig. 2




Fig. 3A




Fig. 3B




Fig. 4




Fig. 5




Fig. 6

HLM incubated with 1 μ M
Compound (X) , Compound (II) and Compound (IV)




Fig. 7

HLM incubated with 10 μ M
Compound (X), Compound (II) or Compound (IV)




Fig. 8

HLM incubated with 10 μ M
Compound (X) and Compound (III)




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13

CYP2C19 incubated with 10 μ M
Compound (X) and Compound (II)




Fig. 14

CYP2C19 incubated with 1 μ M
Compound (X) and Compound (IV)




Fig. 15




Fig. 16




Fig. 17




Fig. 18

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 200501690 A [0002]
- EP 638073 A [0003] [0005] [0008]
- US 5807855 A [0021] [0088] [0104] [0110]
- US 7648991 B [0021] [0088] [0104] [0110]
- US 7767683 B [0021] [0088] [0104] [0110]
- US 7772240 B [0021] [0088] [0104] [0110]
- US 8076342 B [0021] [0088] [0104] [0110]
- US 20080269248 A [0021] [0088] [0104] [0110]
- US 20100069676 A [0021] [0088] [0104] [0110]
- US 20110178094 A [0021] [0088] [0104] [0110]
- US 20110207744 A [0021] [0088] [0104] [0110]
- WO 2005016900 A [0021] [0088] [0104] [0110]
- EP 0638073 A [0021] [0088] [0104] [0110]
- US 7678914 B [0023]
- US 6221335 B [0023]
- EP 35363 A1 [0089] [0124]
- WO 2008025361 A, Kehler, Juhl, Püschl [0089] [0124]
- WO 2006086984 A1, Dahl, Wøhlk Nielsen, Suteu, Robin, Brøsen [0092] [0118]
- WO 2005016901 A1, Bang-Andersen, Bøgesø, Jensen, Svane, Dahl, Howells, Lyngsø, Mow [0092] [0118]


Non-patent literature cited in the description

- BØGESØ et al. *J. Med. Chem.*, 1995, vol. 38, 4380-4392 [0004]
- CARLSSON. *Am. J. Psychiatry*, 1978, vol. 135, 164-173 [0005]
- SEEMAN. *Science and Medicine*, 1995, vol. 2, 28-37 [0005]
- D.N. EDER. *Current Opinion in Investigational Drugs*, 2002, vol. 3 (2), 284-288 [0006]
- P. WILLNER. *Brain. Res. Rev.*, 1983, vol. 6, 211-224, 225-236, 237-246 [0007]
- BØGESØ et al. *J. Med. Chem.*, 1985, vol. 28, 1817-1828 [0007]
- BALSARA et al. *Psychopharmacology*, 1979, vol. 62, 67-69 [0008]
- WEIDEN, P.J. EPS profiles: the atypical antipsychotics are not all the same. *J. Psychiatr. Pract.*, 2007, vol. 13 (1), 13-24 [0020]
- *J. Med. Chem.*, 1995, vol. 38, 4380-4392 [0021] [0088] [0104] [0110]
- *Drug Metab. Dispos.*, 1986, vol. 14, 509 [0023]
- *Arch. Toxicol.*, 1990, vol. 64, 109 [0023]
- *Int. Arch. Occup. Environ. Health*, 1993, vol. 65 (1), S139 [0023]
- *J. Pharm. Sci.*, 1975, vol. 64, 367-391 [0023]
- *Adv. Drug. Res.*, 1985, vol. 14, 1-40 [0023]
- *J. Med. Chem.*, 1991, vol. 34, 2871-2876 [0023]
- *Can. J. Physiol. Pharmacol.*, 1999, 79-88 [0023]
- SILVERMAN, R. B. *The Organic Chemistry of Drug Design and Drug Action*. 2004, 422 [0023]
- *Curr. Opin. DrugDev.*, 2006, vol. 9, 101-109 [0023]
- *Chemical Res. Tox.*, 2008, 1672 [0023]
- HARBESON, S.L ; TUNG, R.D. Deuterium in Drug Discovery and Development. *Ann. Rep. Med. Chem.*, 2011, vol. 46, 404-418 [0023]
- Remington: The Science and Practice of Pharmacy. Lippincott Williams & Wilkins, 2005 [0037] [0104]
- BERGE, S.M. et al. *J. Pharm. Sci.*, 1977, vol. 66, 2 [0081]
- GOULD, P.L. *Int. J. Pharmaceutics*, 1986, vol. 33, 201-217 [0081]
- ANDERSEN et al. *Eur J Pharmacol*, 1987, vol. 144, 1-6 [0158]
- KAPUR S. et al. *J Pharm Exp Ther*, 2003, vol. 305, 625-631 [0158]
- SMITH P.K. et al. *Anal. Biochem.*, 1985, vol. 150, 6-85 [0159]
- JENTSCH, J.D. ; ROTH, R. H. *Neuropsychopharmacology*, 1999, vol. 20, 201-225 [0177]
- JACKSON, D.M. et al. *Pharmacol Biochem Behav.*, 1994, vol. 48, 465-471 [0177]
- ELLIOTT, P.J. et al. *J. Neural. Transm. Park. Dis. Dement. Sect.*, 1990, vol. 2, 79-89 [0178]
- TAKAGI, J. ; TAKAHASHI, K. ; ISHIYAMA, T. ; MI-YAURA, N. *J. Am. Chem. Soc.*, 2002, vol. 124, 8001-8006 [0190]
- SIMEONE, J. P. ; SOWA, J. R. JR. *Tetrahedron*, 2007, vol. 63, 12646-12654 [0190]
- KUNDU, K. ; MCCULLAGH, J. V. ; MOREHEAD, A. T. JR. *J. Am. Chem. Soc.*, 2005, vol. 127, 16042-16043 [0194]
- MINATTI, A. ; ZHENG, X. ; BUCHWALD, S. L. J. *Org. Chem.*, 2007, vol. 72, 9253-9258 [0202]

Deuterált 1-piperazino-3-fenil-indánok, skizofrénia kezelésére

SZABADALMI IGÉNYPONTOK

1. Egy




képletű vegyület, amelyben R^1 - R^{10} jelentései egymástól függetlenül hidrogénatom vagy deutériumatom, ahol R^6 - R^{10} mindegyikének jelentései deutérium, ahol az R^1 - R^{10} legalább egyike legalább körülbelül 50% deutériumot tartalmaz, vagy ennek egy gyógyszerészeti leg elfogadható savaddíciós sója.


2. Az 1. igénypont szerinti vegyület, amelyben R^3 - R^6 mindegyikének jelentése hidrogénatom.

3. Az 1. igénypont szerinti vegyület, amelyben R^3 - R^6 mindegyikének jelentése deutériumatom.


4. A 2. igénypont szerinti vegyület, ahol a vegyület




5. A 2. igénypont szerinti vegyület, ahol a vegyület



6. A 3. igénypont szerinti vegyület, ahol a vegyület



7. A 3. igénypont szerinti vegyület, ahol a vegyület



8. Az 1. igénypont szerinti vegyület, amelyben R^1 és R^2 mindegyikének jelentése deutériumatom.


9. A 8. igénypont szerinti vegyület, amelyben R^3 - R^5 mindegyikének jelentése deutériumatom.

10. A 8. igénypont szerinti vegyület, amelyben R^3 - R^5 mindegyikének jelentése hidrogénatom.

11. Az 1-10. igénypontok bármelyike szerinti vegyület, amelyben a vegyület legalább körülbelül 85%-a rendelkezik egy deutériumatommal minden deutériumatomként megjelölt helyzetben, és bármely, deutériumatomként nem jelölt atom körülbelül ennek természetes izotópos gyakoriságában van jelen.

12. Az 1-11. igénypontok bármelyike szerinti vegyület, amelyben a vegyület legalább körülbelül 90%-a rendelkezik egy deutériumatommal minden deutériumatomként megjelölt helyzetben, és bármely, deutériumatomként nem jelölt atom körülbelül ennek természetes izotópos gyakoriságában van jelen.

13. Az 1. igénypont szerinti vegyület, ahol a vegyület a



hidrogén-tartarátja.


14. A 13. igénypont szerinti vegyület, ahol a vegyület ennek polimorf formájában van jelen, az XRPD diffrakciós mintázata pedig a 18. ábrán van meghatározva.

15. A 13. igénypontok bármelyike szerinti vegyület, amelyben a vegyület legalább körülbelül 85%-a rendelkezik egy deutériumatommal minden deutériumatomként megjelölt helyzetben, és bármely, deutériumatomként nem jelölt atom körülbelül ennek természetes izotópos gyakoriságában van jelen.

16. Gyógyszerkészítmény, amely tartalmazza az 1-15. igénypontok bármelyi-

ke szerinti vegyületet valamint egy vagy több gyógyszerészeti leg elfogadható vivőanyagot, oldószert vagy kötőanyagot.

17. A 16. igénypont szerinti gyógyszerkészítmény, ahol a vegyület a



hidrogén-tartarát-sója.

18. A 16. vagy 17. igénypont szerinti készítmény, amelyben a vívőanyag tartalmaz hidroxipropil- β -ciklodextrint vizben, és amelyben a vegyület legalább 85%-a rendelkezik egy deutériumatommal minden deutériumatomként megjelölt helyzetben, és bármely, deutériumatomként nem jelölt atom körülbelül ennek természetes izotópos gyakoriságában van jelen.

19. Az 1-15. igénypontok bármelyike szerinti vegyület vagy a 16-18. igénypontok bármelyike szerinti készítmény alkalmazása pszichózis, más, pszichotikus tüneteket, pszichotikus rendellenességeket magában foglaló betegség vagy olyan betegségek kezelésére szolgáló gyógyszer előállítására, amely pszichotikus tüneteket mutat.


20. A 19. igénypont szerinti alkalmazás, ahol a pszichózis vagy a pszichotikus tüneteket magában foglaló betegség skizofrénia, skizofrénia-jellegű rendellenesség, skizoaffektív rendellenesség, hallucinációs rendellenesség, rövid pszichotikus rendellenesség, osztott pszichotikus rendellenesség, bipoláris rendellenesség vagy bipoláris rendellenességeben jelentkező mania.

21. A 19-20. igénypontok bármelyike szerinti alkalmazás, amely továbbá mágában foglal egy, a sertindol, olanzapin, risperidon, quetiapin, aripiprazol, haloperidol, clozapine, ziprasidon és az osanetant által alkotott csoportból kiválasztott vegyületet.

22. A 19-20. igénypontok bármelyike szerinti alkalmazás, ahol a pszichózis


vagy a pszichotikus tüneteket magában foglaló betegség skizofrénia.

23. A 19-20. igénypontok bármelyike szerinti alkalmazás, ahol a pszichózis vagy a pszichotikus tüneteket magában foglaló betegség skizofrénia, ahol a gyógyszerkészítmény tartalmazza a:




hidrogén-tartarát-sójának egy hatékony mennyiségét valamint hidroxípropil- β -ciklodextrint vizben, és a (IV) legalább körülbelül 85%-a rendelkezik egy deutériumatommal minden deutériumatomként megjelölt helyzetben, és bármely, deutériumatomként nem jelölt atom körülbelül ennek természetes izotópos gyakoriságában van jelen.

24. A




képletű vegyület.

25. Eljárás a




képletű vegyület előállítására, amely magában foglalja a



vegyületnek $[(S)\text{-BINAP}]\text{Rh(I)}\text{BF}_4$ -el való kezelését.


26. A 25. igénypont szerinti eljárás, amelyben az $[(S)\text{-BINAP}]\text{Rh(I)}\text{BF}_4$ -et katalitikus mennyiségben alkalmazzuk.

27. Eljárás a (XIV) vegyület előállítására, amely magában foglalja a) a



vegyületnek bisz(pinakolát)dibor-al való kezelését, és b) 2-bróm-5-klór-benzaldehiddel való kezelését.

28. A 27. igénypont szerinti eljárás, amelyben a



vegyületnek bisz(pinakolát)dibor-al való kezelése továbbá magában foglalja Pd(II) hozzáadását is.


29. A 28. igénypont szerinti eljárás, amelyben a 2-bróm-5-klór-benzaldehiddel való kezelés továbbá magában foglalja Pd(0) hozzáadását is.

30. Eljárás (1*R*,3*S*)-(IV) tartarat vegyület előállítására, amely magában foglalja racém *transz*-1-(6-klór-3-fenil(d_5)-indan-1-il)-1(d_3)-2,2-trimetil-piperazinnak L-(+)-borostyánkósavval való kezelését.

31. A 30. igénypont szerinti eljárás, amelyben a racém *transz*-1-(6-klór-3-

-fenil(d₅)-indan-1-il)-1(d₃)-2,2-trimetil-piperazint ennek megfelelő szukcinát-sójából hozzuk létre.

A meghatalmazott:


Dr. Szőke Árpád
szekelyi@szekely.hu
Szekszárdi Gyógyfürdő Iroda
H-7700 Szekszárd, Andrássy ut 110.
tel: 052/410000 fax: 052/410001
szekgy@szekgy.hu