



(51) International Patent Classification:

A61P 3/06 (2006.01) A61P 9/06 (2006.01)
A61K 39/395 (2006.01) C07K 16/00 (2006.01)

(21) International Application Number:

PCT/US2020/064324

(22) International Filing Date:

10 December 2020 (10.12.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/946,268 10 December 2019 (10.12.2019) US
62/987,148 09 March 2020 (09.03.2020) US

(71) Applicants: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **SANOFI BIOTECHNOLOGY** [FR/FR]; 54 rue La Boétie, 75008 Paris (FR).

(72) Inventors: **ALI, Shazia**; c/o Regeneron Pharmaceuticals, INC., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **BACCARA-DINET, Marie**; c/o Sanofi Biotechnology, 54 rue La Boétie, 75008 Paris (FR). **DON-AHUE, Stephen**; c/o Regeneron Pharmaceuticals, INC., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **HANOTIN, Corinne**; c/o Sanofi Biotechnology, 54 rue La Boétie, 75008 Paris (FR). **LECORPS, Guillaume**; c/o Sanofi Biotechnology, 54 rue La Boétie, 75008 Paris (FR). **PORDY, Robert C.**; c/o Regeneron Pharmaceuticals, INC., 777 Old Saw Mill River Road, Tarrytown, Massachusetts 10591 (US).

(74) Agent: **DEGIULIO, James**; Lathrop GPM LLP, 28 State Street, Boston, Massachusetts 02109 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: USE OF A PCSK9 INHIBITOR TO TREAT HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA

(57) Abstract: The present disclosure provides methods for lowering LDL-C levels in patients with homozygous familial hypercholesterolemia (hoFH), the method comprising administering to the patient a pharmaceutical composition comprising a PCSK9 inhibitor. In certain embodiments, the PCSK9 inhibitor is an anti-PCSK9 antibody such as the exemplary antibody referred to herein as mAb316P.



USE OF A PCSK9 INHIBITOR TO TREAT HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA

RELATED APPLICATIONS

5 **[0001]** This application claims priority to U.S. Provisional Patent Application Nos. 62/946,268, filed December 10, 2019, and 62/987,148, filed March 9, 2020, the entire disclosures of which are hereby incorporated herein by reference in their entirety.

SEQUENCE LISTING

10 **[0002]** The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created December 10, 2020, is named 711599_SA9_291PC.txt and is 99,701 bytes in size.

FIELD

15 **[0003]** The present disclosure relates to the field of therapeutic treatments of diseases and disorders that are associated with elevated levels of lipids and lipoproteins. More specifically, the disclosure relates to the use of PCSK9 inhibitors to treat patients with homozygous familial hypercholesterolemia (hoFH) who are refractory to statin treatment, who are intolerant to statins, or who otherwise have a history of adverse reactions to statin therapy.

BACKGROUND

20 **[0004]** Familial hypercholesterolemia (FH) is an inherited disorder of lipid metabolism that predisposes a person to premature severe cardiovascular disease (CVD) (Kolansky, *et al.* 2008 *Am J Cardiol* 102(11):1438-1443). It can be either an autosomal dominant or an autosomal recessive disease that results from mutations in the low density lipoprotein receptor (LDLR), or in 3 associated genes: proprotein convertase subtilisin/kexin type 9 (PCSK9), apolipoprotein B (Apo B), and LDL receptor adaptor protein 1 (LDLRAP1), with a similar phenotype and varying severity.

25 **[0005]** Homozygous familial hypercholesterolemia (hoFH) is a rare, serious condition genetically defined to include individuals with the same mutation(s) in both LDLR, ApoB, or PCSK9 alleles (true homozygotes), different mutations in each allele of the same gene
30 (compound heterozygotes), or different mutations on different genes (double heterozygotes). Phenotypically, the severity of hoFH depends on the amount of residual LDLR activity, historically categorized as either receptor-negative (<2% of normal LDLR activity) or receptor-defective (2% to 25% of normal LDLR activity) based on the amount of activity in skin fibroblasts. The genetic definition used herein includes all individuals considered to be true

homozygotes, compound heterozygotes, or double heterozygotes. However, those individuals with null LDLR mutations in both alleles are excluded.

[0006] Patients with hoFH generally have severe hypercholesterolemia (500-1000 mg/dL, 12.95-25.9 mmol/L), resulting in lifelong exposure to high levels of plasma LDL-C and increased risk of developing atherosclerosis at a highly accelerated rate, often manifesting within the first 2 decades of life. Persistently high levels of LDL-C can also lead to cutaneous and tendon xanthomas, valvular and supraaortic stenosis (Kolansky, *et al.* 2008 *Am J Cardiol* 102(11):1438-1443). This accelerated atherosclerosis results in premature cardiovascular disease (CVD) and an increased risk of a cardiovascular (CV) event. A recent observational study of hoFH patients demonstrated that the mean age for first major CV event was 20 years (Kolansky 2008).

[0007] The initial goal of drug therapy in adult patients with FH is to achieve LDL-C reduction $\geq 50\%$ (Goldberg, *et al.* 2011 *J Clin Lipidol* 5(3 Supp):S1-S8). If this is achieved, therapy is escalated with an aim to achieve an LDL-C of <100 mg/dL (2.59 mmol/L) in the absence of coronary artery disease or other major risk factors or <70 mg/dL (<1.81 mmol/L) in the presence of coronary artery disease or other major risk factors (Watts, *et al.* 2014 *J Clin Lipidol* 8(2):148-172). However, management of elevated LDL-C in patients with hoFH is challenging with the current existing treatment options. Patients with hoFH generally have a poor response to conventional drug therapies, resulting in extremely elevated LDL-C levels that are often refractory to pharmacologic management thus requiring the need to initiate LDL apheresis.

[0008] Statins inhibit cholesterol synthesis by inhibiting 3-hydroxy-3-methyl-glutaryl coenzyme reductase and are used as first-line therapy in hoFH patients. Patients with hoFH tend to be refractory to statins because the mechanism of action generally lowers LDL-C levels through up-regulation of the hepatic LDL receptor. Nonetheless, despite the near total loss of functional LDL receptors in hoFH patients, statins are still used as first line therapy in order to maximize residual receptor activity (Raaij, *et al.* 2000 *Atherosclerosis* 150(2):421-428, Marais, *et al.* 2008 *Atherosclerosis* 197(1):400-406, Raaij, *et al.* 1997 *Atherosclerosis* 135(2):249-256).

[0009] Many patients treated with the combination of high dose statin and ezetimibe, a cholesterol absorption inhibitor, remain far from their target LDL-C. Newer therapies, *i.e.*, mipomersen and lomitapide, have been approved for use in patients with hoFH, but they are not commercially available in all countries and are associated with increases in hepatic fat content, elevated markers of liver injury, frequent injection site reactions that can be of severe intensity (mipomersen) or poorly tolerated gastrointestinal adverse effects (lomitapide) (Raaij, *et al.* 2010 *Lancet* 375(9719):998-1006, Cuchel, *et al.* 2013 *Lancet* 381(9860):40-46).

[0010] Mechanical removal of LDL-C using LDL apheresis is an option, but may lower the quality of life in patients and present other challenges (Schiel, *et al.* 1995 *Int J Artif Organs*

18(12):786-793). Low-density lipoprotein apheresis is a costly procedure that is invasive and burdensome for patients.

BRIEF SUMMARY

5 [0011] There remains a need for methods of effectively treating patients having homozygous familial hypercholesterolemia (hoFH). There also remains a need for methods of effectively treating patients having hoFH who are unable to achieve acceptable LDL-C levels despite treatment with statins, do not tolerate or experience adverse reactions to statin therapy, and/or must resort to LDL apheresis for treatment.

10 [0012] The present disclosure provides methods for treating homozygous familial hypercholesterolemia (hoFH). In particular, the methods of the present disclosure are useful for treating patients with hoFH, excluding those patients with null/null mutations in both LDLR alleles.

15 [0013] In one aspect, the disclosure provides a method for treating homozygous familial hypercholesterolemia (hoFH) in a patient in need thereof, the method comprising: (a) selecting a patient having hoFH who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy; and (b) administering one or more doses of a PCSK9 inhibitor to the patient.

20 [0014] In another aspect, the disclosure provides a method for reducing serum LDL-C levels in a patient having homozygous familial hypercholesterolemia (hoFH), the method comprising: (a) selecting a patient who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy; and (b) administering one or more doses of a PCSK9 inhibitor to the patient.

25 [0015] In still another aspect, the disclosure provides a method for treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH), the method comprising: (a) selecting a patient who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy; and (b) administering one or more doses of a PCSK9 inhibitor to the patient.

30 [0016] In one embodiment of the methods according to the disclosure, the patient has at least about 100 mg/dL LDL-C prior to or at the time of administration of the PCSK9 inhibitor. In one embodiment of the methods according to the disclosure, the patient has about 250 mg/dL to about 1000 mg/dL LDL-C prior to or at the time of administration of the PCSK9 inhibitor. In another embodiment of the methods according to the disclosure, the patient has about 500 mg/dL to about 1000 mg/dL LDL-C prior to or at the time of administration of the PCSK9 inhibitor.

[0017] In one embodiment of the methods according to the disclosure, the patient is receiving at least one lipid-modifying therapy (LMT) prior to or at the time of administration of the PCSK9 inhibitor. In further embodiments, the at least one LMT is at least one statin. In still further embodiments, the at least one LMT is LDL apheresis. In still further embodiments, the at least one LMT is ezetimibe.

[0018] In one embodiment of the methods according to the disclosure, the patient has an increased risk for premature cardiovascular disease and/or for a cardiovascular event.

[0019] In one embodiment of the methods according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof that specifically binds PCSK9. In further embodiments, the antibody or antigen binding fragment thereof that specifically binds PCSK9 is administered to the patient at a dose of about 150 mg at a frequency of once every two weeks. In still further embodiments, the antibody or antigen binding fragment thereof that specifically binds PCSK9 is administered to the patient subcutaneously.

[0020] In one embodiment of the methods according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof that comprises the heavy and light chain CDRs of a HCVR/LCVR amino acid sequence pair comprising SEQ ID NOs: 1/6. In further embodiments, the antibody or antigen-binding fragment thereof comprises heavy and light chain CDR amino acid sequences having SEQ ID NOs:2, 3, 4, 7, 8 and 10. In still further embodiments, the antibody or antigen-binding fragment thereof comprises an HCVR having the amino acid sequence of SEQ ID NO:1 and an LCVR having the amino acid sequence of SEQ ID NO:6.

[0021] In one embodiment of the methods according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof, wherein the antibody or antigen-binding fragment thereof binds to the same epitope on PCSK9 as an antibody comprising heavy and light chain CDR amino acid sequences having SEQ ID NOs:2, 3, 4, 7, 8 and 10.

[0022] In another embodiment of the methods according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof, wherein the antibody or antigen-binding fragment thereof competes for binding to PCSK9 with an antibody comprising heavy and light chain CDR amino acid sequences having SEQ ID NOs:2, 3, 4, 7, 8 and 10.

[0023] In one embodiment of the methods according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen-binding fragment thereof, wherein the antibody or antigen-binding fragment thereof is contained in a pre-filled pen delivery device.

[0024] In one embodiment of the methods according to the disclosure, the patient with hoFH has an LDL receptor genotype selected from the group consisting of: (a) homozygous non-null/non-null; (b) compound heterozygous non-null/-non-null; (c) compound heterozygous non-null/null; and (d) homozygous null/null. In another embodiment, the patient with hoFH has an

LDL receptor genotype selected from the group consisting of: (a) homozygous non-null/non-null; (b) compound heterozygous non-null-/non-null; and (c) compound heterozygous non-null/null.

[0025] In one embodiment of the methods according to the disclosure, about 12 weeks after administration of the one or more doses of the PCSK9 inhibitor, the patient exhibits one or more lipid parameter improvements selected from the group consisting of:

- i) a reduction in LDL-C level from baseline of about 35%;
- ii) a reduction in non-HDL-C level from baseline of about 33%;
- iii) a reduction in Apo B level from baseline of about 30%;
- iv) a reduction in total cholesterol level from baseline of about 27%;
- v) a reduction in (fasting) triglyceride level from baseline of about 11%; and/or
- vi) a reduction in Lp(a) level from baseline of about 28%.

[0026] In one aspect, the disclosure provides a pharmaceutical composition for use in treating homozygous familial hypercholesterolemia (hoFH) in a patient in need thereof, wherein the composition comprises a PCSK9 inhibitor and a pharmaceutically acceptable excipient.

[0027] In another aspect, the disclosure provides a pharmaceutical composition for use in reducing serum LDL-C levels in a patient having homozygous familial hypercholesterolemia (hoFH), wherein the composition comprises a PCSK9 inhibitor and a pharmaceutically acceptable excipient.

[0028] In still another aspect, the disclosure provides a pharmaceutical composition for use in treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH), wherein the composition comprises a PCSK9 inhibitor and a pharmaceutically acceptable excipient.

[0029] In one embodiment of the compositions according to the disclosure, the patient having hoFH is refractory to treatment with statins, is intolerant to statins, and/or has a history of adverse reaction(s) to statin therapy.

[0030] In another embodiment of the compositions according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof that specifically binds PCSK9. In further embodiments, the antibody or antigen binding fragment thereof comprises the heavy and light chain CDRs of a HCVR/LCVR amino acid sequence pair comprising SEQ ID NOs: 1/6.

[0031] In one embodiment of the compositions according to the disclosure, the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof that specifically binds PCSK9, wherein the antibody or antigen binding fragment thereof is administered to the patient at a dose of about 150 mg at a frequency of once every two weeks.

[0032] In one aspect, the disclosure provides a method for treating homozygous familial hypercholesterolemia (hoFH), the method comprising administering one or more doses of a

PCSK9 inhibitor to a patient who is diagnosed with hoFH and who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy.

[0033] The present disclosure also provides methods for treating homozygous familial hypercholesterolemia (hoFH) in a patient who is intolerant to statins or who has a history of adverse reactions to statin therapy by selecting a patient with moderate, high, or very high cardiovascular risk who has previously experienced skeletal muscle-related symptoms that began or increased while on a daily therapeutic statin regimen and administering one or more doses of a PCSK9 inhibitor to the patient. According to certain embodiments, the patient is selected on the basis of having previously experienced skeletal muscle-related symptoms that began or increased while on at least two separate daily therapeutic statin regimens (e.g., wherein at least one of the daily therapeutic statin regimens is the lowest approved daily dose of a statin).

[0034] The present disclosure also provides pharmaceutical compositions comprising a PCSK9 inhibitor for use in treating homozygous familial hypercholesterolemia (hoFH) in a patient in need thereof, pharmaceutical compositions comprising a PCSK9 inhibitor for use in reducing serum LDL-C levels in a patient having homozygous familial hypercholesterolemia (hoFH), and pharmaceutical compositions comprising a PCSK9 inhibitor for use in treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH). In certain embodiments of the pharmaceutical compositions of the disclosure, the patient has hoFH and is refractory to treatment with statins, is intolerant to statins, or has a history of adverse reactions to statin therapy.

[0035] Other embodiments of the present disclosure will become apparent from a review of the ensuing detailed description.

BRIEF DESCRIPTION OF THE FIGURES

[0036] **Figure 1** is a Study Flow Diagram illustrating the clinical trial described in Example 2 herein.

[0037] **Figure 2** graphically depicts the LDL-C LS Mean (+/-SE) percent change from baseline for the double-blind period over time. Least-squares (LS) means, standard errors (SE), and p-value taken from MMRM (mixed-effect model with repeated measures) analysis. The model includes the fixed treatment effect, randomization strata as per IVRS, time point, treatment-by-time point interaction, strata-by-time point interaction, as well as the continuous baseline LDL-C value by time-point interaction.

DETAILED DESCRIPTION

[0038] Before the present disclosure is described, it is to be understood that this disclosure is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present disclosure will be limited only by the appended claims.

[0039] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. As used herein, the term "about," when used in reference to a particular recited numerical value, means that the value may vary from the recited value by no more than 1%. For example, as used herein, the expression "about 100" includes 99 and 101 and all values in between (e.g., 99.1, 99.2, 99.3, 99.4, etc.).

[0040] Although any methods and materials similar or equivalent to those described herein can be used in the practice of the present disclosure, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to describe in their entirety.

Statin Inefficacy and Intolerance

[0041] In some embodiments, the present disclosure relates generally to methods and compositions that are useful for treating patients who are refractory to treatment with statins, intolerant to statins ("statin intolerant patients," also referred to herein as "a patient who is intolerant to statins"), or who has a history of adverse reactions to statin therapy. As used herein, a patient is regarded as "refractory to statins" if the patient has been subjected to statin therapy without sufficient improvement in the lipid parameters. For example, a patient who is refractory to statins has been on stable statin therapy but still has serum LDL-C levels of at least 70 mg/dL.

[0042] As used herein, a patient is regarded as "statin intolerant" or "intolerant to statins" if the patient, has a history of experiencing one or more adverse reactions that began or increased while on a daily statin therapeutic regimen and stopped when statin therapy was discontinued. In certain embodiments, the adverse reactions are musculoskeletal in nature, e.g., skeletal muscle pain, aches, weakness or cramping (e.g., myalgia, myopathy, rhabdomyolysis, etc.). In certain embodiments, the adverse reactions are skeletal muscle pain or aches that occur or are intensified following exercise or exertion. Statin-related adverse reactions also include hepatic, gastrointestinal and psychiatric symptoms that correlate with statin administration.

[0043] According to certain embodiments, a patient is deemed "statin intolerant" or "intolerant to statins" if the patient has a history of skeletal muscle-related symptoms associated with at

least two different and separate daily statin therapeutic regimens. According to certain embodiments, a patient is "statin intolerant" or "intolerant to statins" if the patient exhibits one or more statin-related adverse reaction(s) to the lowest approved daily doses of one or more statins. In certain embodiments, a patient is "statin intolerant" or "intolerant to statins" if the patient is unable to tolerate a cumulative weekly statin dose of seven times the lowest approved tablet size. According to other embodiments of the present disclosure, a patient is "statin intolerant" or "intolerant to statins" if the patient is able to tolerate a low dose statin therapy but develops symptoms when the dose is increased (e.g., to achieve a targeted LDL-C level).

[0044] According to the present disclosure, "a history of skeletal muscle-related symptoms associated with taking at least two different and separate statins" includes skeletal muscle-related pain, aches, weakness and/or cramping, that began or increased during statin therapy and stopped when statin therapy was discontinued. In the context of the present disclosure, exemplary statin therapies associated with statin intolerance may include daily therapeutic statin regimens selected from the group consisting of: 5 mg rosuvastatin daily, 10 mg atorvastatin daily, 10 mg simvastatin daily, 20 mg lovastatin daily, 40 mg pravastatin daily, 40 mg fluvastatin daily, and 2 mg pitavastatin daily.

Methods for Treating Homozygous Familial Hypercholesterolemia and Reducing Serum LDL-C Levels

[0045] According to certain embodiments, the patient who is treatable by the methods of the present disclosure has homozygous Familial Hypercholesterolemia (hoFH) (sometimes referred to herein as "a hypercholesterolemic patient"). Homozygous familial hypercholesterolemia (hoFH) can be characterized by high LDL-cholesterol levels and atherosclerotic cardiovascular disease, despite treatment with lipid-lowering therapies.

[0046] In some embodiments, a patient is diagnosed with hoFH based on genotype or clinical criteria. In some embodiments, patients diagnosed with hoFH include all individuals considered to be true homozygotes (same mutation(s) in both alleles of the same gene), compound heterozygotes (different mutations in each allele of the same gene), or double heterozygotes (different mutations in different genes) for mutations in the LDLR, ApoB, PCSK9, or LDLRAP1 genes. In some embodiments, the mutation is characterized as "null" or "non-null" based on the amount of residual LDLR activity. In some embodiments, the patient is diagnosed with hoFH based on a genotype including: (a) homozygous non-null/non-null; (b) compound heterozygous non-null/non-null; (c) compound heterozygous non-null/null; or (d) homozygous null/null. In some embodiments, a patient having a "null/null" mutation has residual LDLR activity < 2%. In some embodiments, the patient is diagnosed with hoFH based on one or more clinical criteria, including but not limited to: (a) untreated total cholesterol >500 mg/dL (12.93 mmol/L) and triglycerides (TG) <300 mg/dL (3.39 mmol/L), (b) both parents with history of total cholesterol

>250 mg/dL (6.46 mmol/L), and (c) cutaneous or tendinous xanthoma before age 10. In some embodiments, a patient having hoFH is selected for treatment with the methods and compositions disclosed herein.

[0047] The present disclosure includes methods for reducing serum LDL-C levels in a patient having hoFH. The subject may have hoFH and be refractory to treatment with statins, be intolerant to statins, and/or have a history of adverse reactions to statin therapy. Similarly, the present disclosure includes methods for reducing serum LDL-C levels in a patient having hoFH without inducing skeletal muscle pain, discomfort, weakness, or cramping. As used in this context, "reducing serum LDL-C levels" means causing the patient's serum LDL-C level to decrease by at least 10% (e.g., at least 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75% or more). The terms "subject" and "patient" are used interchangeably herein.

Methods for Eliminating or Reducing Statin Usage

[0048] The present disclosure includes methods and composition useful, *inter alia*, for eliminating or reducing statin usage in a patient having hoFH. In some embodiments, the patient having hoFH is refractory or intolerant to statins or who experiences adverse events upon treatment with statins. The methods according to this aspect of the disclosure comprise: (a) selecting a patient who is or was on a daily therapeutic statin regimen and who is refractory to statin therapy, intolerant to statins, or who has a history of adverse reactions to statin therapy; and (b) discontinuing or reducing the patient's daily therapeutic statin regimen; and (c) administering one or more doses of a PCSK9 inhibitor to the patient. According to certain embodiments of this aspect of the disclosure, the patient's daily therapeutic statin regimen may be completely discontinued at the time of or just prior to commencement of a therapeutic course of treatment comprising administration of one or more doses of a PCSK9 inhibitor to the patient. In other embodiments, the patient's daily therapeutic statin regimen may be gradually reduced at the time of or just prior to commencement of a therapeutic course of treatment comprising administration of one or more doses of a PCSK9 inhibitor to the patient. Gradual reduction of a statin regimen, in the context of this aspect of the disclosure, may comprise reducing the quantity of statin administered to a patient, and/or decreasing the frequency of administration of statin to the patient. Gradual reduction of a statin regimen, according to this aspect of the disclosure, may result in complete elimination of statin usage by the patient while the patient is receiving a PCSK9 inhibitor in place of the statin. In this respect, the adverse effects of statins on a patient are reduced or eliminated by reducing or eliminating statin usage by the patient, while still permitting adequate treatment of homozygous familial hypercholesterolemia in the patient by administration of a PCSK9 inhibitor.

Patient Selection

[0049] The present disclosure includes methods and composition useful, *inter alia*, for treating patients who have homozygous familial hypercholesterolemia (heFH), including for example patients who are "refractory to statins", "statin intolerant", or "intolerant to statins", and/or who experience adverse reactions upon treatment with statins (as defined elsewhere herein). The patients who are treatable by the methods of the present disclosure may also exhibit one or more additional selection criteria. For example, a patient may be selected for treatment with the methods of the present disclosure on the basis of having moderate, high, or very high CV risk. Degree of CV risk may be assessed and expressed in terms of a calculated 10-year fatal cardiovascular disease (CVD) risk SCORE value, as defined by The Task Force for the Management of Dislipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS), as set forth in the ESC/EAS Guidelines for the Management of Dislipidaemias, European Heart Journal, 2100; 32:1769-1818 (referred to herein as "ESC/EAS 2011"), the disclosure of which is incorporated by reference herein in its entirety.

[0050] As used herein, "moderate CV risk" means a calculated 10-year fatal CVD risk SCORE greater than or equal to 1% and less than 5%. As used herein, "high CV risk" means a calculated 10-year fatal CVD risk SCORE greater than or equal to 5%, and/or moderate kidney disease (CKD), and/or type 1 or type 2 diabetes mellitus without target organ damage, and/or heFH. As used herein, "very high CV risk" means a history of documented coronary heart disease (CHD), ischemic stroke, peripheral arterial disease (PAD), transient ischemic attack (TIA), abdominal aortic aneurysm, carotid artery occlusion greater than 50% without symptoms, carotid endarterectomy or carotid artery stent procedure, renal artery stenosis, renal artery stent procedure, and/or type 1 or type 2 diabetes mellitus with target organ damage.

[0051] According to certain embodiments, the patient may be selected on the basis of having a history of coronary heart disease (CHD). As used herein, a "history of CHD" (or "documented history of CHD") includes one or more of: (i) acute myocardial infarction (MI); (ii) silent MI; (iii) unstable angina; (iv) coronary revascularization procedure (e.g., percutaneous coronary intervention [PCI] or coronary artery bypass graft surgery [CABG]); and/or (v) clinically significant CHD diagnosed by invasive or non-invasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography or nuclear imaging).

[0052] According to certain embodiments, the patient may be selected on the basis of having one or more additional risk factors selected from the group consisting of age (e.g., older than 40, 45, 50, 55, 60, 65, 70, 75, or 80 years), race, national origin, gender (male or female), exercise habits (e.g., regular exerciser, non-exerciser), other preexisting medical conditions (e.g., type-II diabetes, high blood pressure, etc.), and current medication status (e.g., currently taking beta blockers, niacin, ezetimibe, fibrates, omega-3 fatty acids, bile acid resins, etc.).

[0053] The present disclosure includes methods and compositions useful, *inter alia*, for treating patients who have hoFH and are receiving treatment with maximally tolerated statin therapy. In some embodiments, the patient As used herein, “maximally tolerated statin therapy” or “maximum tolerated dose of statin therapy” are used interchangeably to mean a therapeutic regimen comprising the administration of a daily dose of a statin that is the highest dose of statin that can be administered to a particular patient without causing unacceptable adverse side effects in the patient. Maximally tolerated statin therapy includes, but is not limited to, high intensity statin therapy.

[0054] As used herein, “maximally tolerated lipid modifying therapy” or “maximum tolerated LMT” are used interchangeably to mean a therapeutic regimen comprising the administration of a daily, weekly, or monthly dose of a lipid modifying therapy (LMT) that is the highest dose of the LMT that can be administered to a particular patient without causing unacceptable adverse side effects in the patient. Maximally tolerated LMT includes, but is not limited to, high intensity statin therapy, ezetimibe, fibrates, bile acid sequestrants, cholesterol absorption inhibitors, nicotinic acid or derivatives, omega 3 fatty acids, probucol, lomitapide, and mipomersen.

[0055] As used herein, the term “premature cardiovascular disease” refers to cardiovascular disease in a patient before the age of 50 years old.

Therapeutic Efficacy

[0056] The methods of the present disclosure result in the reduction in serum levels of one or more lipid component selected from the group consisting of LDL-C, ApoB, non-HDL-C, total cholesterol (TC), triglycerides (TG), Lp(a), and/or remnant cholesterol. For example, according to certain embodiments of the present disclosure, administration of a pharmaceutical composition comprising a PCSK9 inhibitor to a patient with hoFH will result in a mean percent reduction from baseline in serum low density lipoprotein cholesterol (LDL-C) of at least about 25%, 30%, 40%, 45%, 50%, 60%, or greater; a mean percent reduction from baseline in ApoB of at least about 20%, 25%, 30%, 40%, 50%, 60%, or greater; a mean percent reduction from baseline in non-HDL-C of at least about 20%, 25%, 30%, 40%, 50%, 60%, or greater; a mean percent reduction from baseline in total cholesterol of at least about 10%, 15%, 20%, 25%, 30%, 35%, or greater; a mean percent reduction from baseline in triglycerides (e.g., fasting triglycerides) of at least about 5%, 10%, 15%, 20%, 25%, 30%, 35% or greater; and/or a mean percent reduction from baseline in Lp(a) of at least about 5%, 10%, 15%, 20%, 25%, 30%, or greater. The percent reductions in the various lipid parameters as set forth above may be achieved at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, or more weeks after the commencement of a therapeutic regimen comprising the administration of a PCSK9 inhibitor as disclosed herein (e.g., 150 mg mAb316P administered once every two weeks, or other similar administration regimens; see, e.g., Example 2 herein).

[0057] According to certain specific embodiments, the present disclosure includes methods for reducing serum LDL-C levels in a patient with hoFH. In some embodiments, the patient having hoFH is refractory to statins, intolerant to statins, or who has a history of adverse reactions to statin therapy. According to additional embodiments, the present disclosure includes methods for treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH). In some embodiments, the patient having hoFH is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy. The methods according to this aspect of the disclosure comprise: (a) selecting a patient with moderate, high, or very high cardiovascular risk who is refractory to statins, intolerant to statins, or has a history of adverse reactions to statin therapy; and (b) administering one or more doses of an anti-PCSK9 antibody to the patient at a dosing amount of about 150 mg per dose, and a dosing frequency of about once every two weeks. In further embodiments of the methods described herein, after about 12 weeks of treatment with the anti-PCSK9 antibody, the patient exhibits one or more lipid parameter improvements selected from the group consisting of: a reduction in LDL-C level from baseline of about 35%, a reduction in non-HDL-C level from baseline of about 33%, a reduction in Apo B level from baseline of about 30%, a reduction in total cholesterol level from baseline of about 27%, a reduction in (fasting) triglyceride level from baseline of about 11%, and/or a reduction in Lp(a) level from baseline of about 28%. Methods according to this aspect of the disclosure may comprise discontinuing the patient's background statin therapy prior to or concurrent with commencement of treatment with the anti-PCSK9 antibody.

PCSK9 Inhibitors

[0058] The methods of the present disclosure comprise administering to a patient a therapeutic composition comprising a PCSK9 inhibitor. As used herein, a "PCSK9 inhibitor" is any agent that binds to or interacts with human PCSK9 and inhibits the normal biological function of PCSK9 *in vitro* or *in vivo*. Non-limiting examples of categories of PCSK9 inhibitors include small molecule PCSK9 antagonists, peptide-based PCSK9 antagonists (e.g., "peptibody" molecules), and antibodies or antigen-binding fragments of antibodies that specifically bind human PCSK9.

[0059] The term "human proprotein convertase subtilisin/kexin type 9" or "human PCSK9" or "hPCSK9", as used herein, refers to PCSK9 encoded by the nucleic acid sequence shown in SEQ ID NO:197 and comprising the amino acid sequence of SEQ ID NO:198, or a biologically active fragment thereof.

[0060] The term "antibody", as used herein, is intended to refer to immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains interconnected by disulfide bonds, as well as multimers thereof (e.g., IgM). Each heavy chain

comprises a heavy chain variable region (abbreviated herein as HCVR or V_H) and a heavy chain constant region. The heavy chain constant region comprises three domains, C_H1 , C_H2 and C_H3 . Each light chain comprises a light chain variable region (abbreviated herein as LCVR or V_L) and a light chain constant region. The light chain constant region comprises one domain (C_L1). The V_H and V_L regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In different embodiments of the disclosure, the FRs of the anti-PCSK9 antibody (or antigen-binding portion thereof) may be identical to the human germline sequences, or may be naturally or artificially modified. An amino acid consensus sequence may be defined based on a side-by-side analysis of two or more CDRs.

[0061] The term "antibody," as used herein, also includes antigen-binding fragments of full antibody molecules. The terms "antigen-binding portion" of an antibody, "antigen-binding fragment" of an antibody, and the like, as used herein, include any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex. Antigen-binding fragments of an antibody may be derived, e.g., from full antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Such DNA is known and/or is readily available from, e.g., commercial sources, DNA libraries (including, e.g., phage-antibody libraries), or can be synthesized. The DNA may be sequenced and manipulated chemically or by using molecular biology techniques, for example, to arrange one or more variable and/or constant domains into a suitable configuration, or to introduce codons, create cysteine residues, modify, add or delete amino acids, etc.

[0062] Non-limiting examples of antigen-binding fragments include: (i) Fab fragments; (ii) $F(ab')_2$ fragments; (iii) Fd fragments; (iv) Fv fragments; (v) single-chain Fv (scFv) molecules; (vi) dAb fragments; and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (e.g., an isolated complementarity determining region (CDR) such as a CDR3 peptide), or a constrained FR3-CDR3-FR4 peptide. Other engineered molecules, such as domain-specific antibodies, single domain antibodies, domain-deleted antibodies, chimeric antibodies, CDR-grafted antibodies, diabodies, triabodies, tetrabodies, minibodies, nanobodies (e.g. monovalent nanobodies, bivalent nanobodies, etc.), small modular immunopharmaceuticals (SMIPs), and shark variable IgNAR domains, are also encompassed within the expression "antigen-binding fragment," as used herein.

[0063] An antigen-binding fragment of an antibody will typically comprise at least one variable

domain. The variable domain may be of any size or amino acid composition and will generally comprise at least one CDR, which is adjacent to or in frame with one or more framework sequences. In antigen-binding fragments having a V_H domain associated with a V_L domain, the V_H and V_L domains may be situated relative to one another in any suitable arrangement. For example, the variable region may be dimeric and contain V_H - V_H , V_H - V_L or V_L - V_L dimers. Alternatively, the antigen-binding fragment of an antibody may contain a monomeric V_H or V_L domain.

[0064] In certain embodiments, an antigen-binding fragment of an antibody may contain at least one variable domain covalently linked to at least one constant domain. Non-limiting, exemplary configurations of variable and constant domains that may be found within an antigen-binding fragment of an antibody of the present disclosure include: (i) V_H - C_H1 ; (ii) V_H - C_H2 ; (iii) V_H - C_H3 ; (iv) V_H - C_H1 - C_H2 ; (v) V_H - C_H1 - C_H2 - C_H3 ; (vi) V_H - C_H2 - C_H3 ; (vii) V_H - C_L ; (viii) V_L - C_H1 ; (ix) V_L - C_H2 ; (x) V_L - C_H3 ; (xi) V_L - C_H1 - C_H2 ; (xii) V_L - C_H1 - C_H2 - C_H3 ; (xiii) V_L - C_H2 - C_H3 ; and (xiv) V_L - C_L . In any configuration of variable and constant domains, including any of the exemplary configurations listed above, the variable and constant domains may be either directly linked to one another or may be linked by a full or partial hinge or linker region. A hinge region may consist of at least 2 (e.g., 5, 10, 15, 20, 40, 60 or more) amino acids which result in a flexible or semi-flexible linkage between adjacent variable and/or constant domains in a single polypeptide molecule. Moreover, an antigen-binding fragment of an antibody of the present disclosure may comprise a homo-dimer or hetero-dimer (or other multimer) of any of the variable and constant domain configurations listed above in non-covalent association with one another and/or with one or more monomeric V_H or V_L domain (e.g., by disulfide bond(s)).

[0065] As with full antibody molecules, antigen-binding fragments may be monospecific or multispecific (e.g., bispecific). A multispecific antigen-binding fragment of an antibody will typically comprise at least two different variable domains, wherein each variable domain is capable of specifically binding to a separate antigen or to a different epitope on the same antigen. Any multispecific antibody format, including the exemplary bispecific antibody formats disclosed herein, may be adapted for use in the context of an antigen-binding fragment of an antibody of the present disclosure using routine techniques available in the art.

[0066] The constant region of an antibody is important in the ability of an antibody to fix complement and mediate cell-dependent cytotoxicity. Thus, the isotype of an antibody may be selected on the basis of whether it is desirable for the antibody to mediate cytotoxicity.

[0067] The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the disclosure may nonetheless include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-

specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

5 **[0068]** The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further below), antibodies isolated from a recombinant, combinatorial human antibody library (described further below), antibodies isolated from an animal (e.g., a mouse)
10 that is transgenic for human immunoglobulin genes (see e.g., Taylor et al. (1992) Nucl. Acids Res. 20:6287-6295) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant
15 human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the V_H and V_L regions of the recombinant antibodies are sequences that, while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

20 **[0069]** Human antibodies can exist in two forms that are associated with hinge heterogeneity. In one form, an immunoglobulin molecule comprises a stable four chain construct of approximately 150-160 kDa in which the dimers are held together by an interchain heavy chain disulfide bond. In a second form, the dimers are not linked via inter-chain disulfide bonds and a molecule of about 75-80 kDa is formed composed of a covalently coupled light and heavy chain
25 (half-antibody). These forms have been extremely difficult to separate, even after affinity purification.

[0070] The frequency of appearance of the second form in various intact IgG isotypes is due to, but not limited to, structural differences associated with the hinge region isotype of the antibody. A single amino acid substitution in the hinge region of the human IgG4 hinge can
30 significantly reduce the appearance of the second form (Angal et al. (1993) Molecular Immunology 30:105) to levels typically observed using a human IgG1 hinge. The instant disclosure encompasses antibodies having one or more mutations in the hinge, C_H2 or C_H3 region, which may be desirable, for example, in production, to improve the yield of the desired antibody form.

35 **[0071]** An "isolated antibody," as used herein, means an antibody that has been identified and separated and/or recovered from at least one component of its natural environment. For

example, an antibody that has been separated or removed from at least one component of an organism, or from a tissue or cell in which the antibody naturally exists or is naturally produced, is an "isolated antibody" for purposes of the present disclosure. An isolated antibody also includes an antibody *in situ* within a recombinant cell. Isolated antibodies are antibodies that have been subjected to at least one purification or isolation step. According to certain embodiments, an isolated antibody may be substantially free of other cellular material and/or chemicals.

[0072] The term "specifically binds," or the like, means that an antibody or antigen-binding fragment thereof forms a complex with an antigen that is relatively stable under physiologic conditions. Methods for determining whether an antibody specifically binds to an antigen are well known in the art and include, for example, equilibrium dialysis, surface plasmon resonance, and the like. For example, an antibody that "specifically binds" PCSK9, as used in the context of the present disclosure, includes antibodies that bind PCSK9 or portion thereof with a K_D of less than about 1000 nM, less than about 500 nM, less than about 300 nM, less than about 200 nM, less than about 100 nM, less than about 90 nM, less than about 80 nM, less than about 70 nM, less than about 60 nM, less than about 50 nM, less than about 40 nM, less than about 30 nM, less than about 20 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, less than about 1 nM or less than about 0.5 nM, as measured in a surface plasmon resonance assay. An isolated antibody that specifically binds human PCSK9, however, have cross-reactivity to other antigens, such as PCSK9 molecules from other (non-human) species.

[0073] The anti-PCSK9 antibodies useful for the methods of the present disclosure may comprise one or more amino acid substitutions, insertions and/or deletions in the framework and/or CDR regions of the heavy and light chain variable domains as compared to the corresponding germline sequences from which the antibodies were derived. Such mutations can be readily ascertained by comparing the amino acid sequences disclosed herein to germline sequences available from, for example, public antibody sequence databases. The present disclosure includes methods involving the use of antibodies, and antigen-binding fragments thereof, which are derived from any of the amino acid sequences disclosed herein, wherein one or more amino acids within one or more framework and/or CDR regions are mutated to the corresponding residue(s) of the germline sequence from which the antibody was derived, or to the corresponding residue(s) of another human germline sequence, or to a conservative amino acid substitution of the corresponding germline residue(s) (such sequence changes are referred to herein collectively as "germline mutations"). A person of ordinary skill in the art, starting with the heavy and light chain variable region sequences disclosed herein, can easily produce numerous antibodies and antigen-binding fragments that comprise one or more individual germline mutations or combinations thereof. In certain embodiments, all of the framework

and/or CDR residues within the V_H and/or V_L domains are mutated back to the residues found in the original germline sequence from which the antibody was derived. In other embodiments, only certain residues are mutated back to the original germline sequence, e.g., only the mutated residues found within the first 8 amino acids of FR1 or within the last 8 amino acids of FR4, or only the mutated residues found within CDR1, CDR2 or CDR3. In other embodiments, one or more of the framework and/or CDR residue(s) are mutated to the corresponding residue(s) of a different germline sequence (*i.e.*, a germline sequence that is different from the germline sequence from which the antibody was originally derived). Furthermore, the antibodies of the present disclosure may contain any combination of two or more germline mutations within the framework and/or CDR regions, e.g., wherein certain individual residues are mutated to the corresponding residue of a particular germline sequence while certain other residues that differ from the original germline sequence are maintained or are mutated to the corresponding residue of a different germline sequence. Once obtained, antibodies and antigen-binding fragments that contain one or more germline mutations can be easily tested for one or more desired property such as, improved binding specificity, increased binding affinity, improved or enhanced antagonistic or agonistic biological properties (as the case may be), reduced immunogenicity, etc. The use of antibodies and antigen-binding fragments obtained in this general manner are encompassed within the present disclosure.

[0074] The present disclosure also includes methods involving the use of anti-PCSK9 antibodies comprising variants of any of the HCVR, LCVR, and/or CDR amino acid sequences disclosed herein having one or more conservative substitutions. For example, the present disclosure includes the use of anti-PCSK9 antibodies having HCVR, LCVR, and/or CDR amino acid sequences with, e.g., 10 or fewer, 8 or fewer, 6 or fewer, 4 or fewer, etc. conservative amino acid substitutions relative to any of the HCVR, LCVR, and/or CDR amino acid sequences disclosed herein.

[0075] The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore™ system (Biacore Life Sciences division of GE Healthcare, Piscataway, NJ).

[0076] The term " K_D ", as used herein, is intended to refer to the equilibrium dissociation constant of a particular antibody-antigen interaction.

[0077] The term "epitope" refers to an antigenic determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. A single antigen may have more than one epitope. Thus, different antibodies may bind to different areas on an antigen and may have different biological effects. Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino

acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain. In certain circumstance, an epitope may include moieties of saccharides, phosphoryl groups, or sulfonyl groups on the antigen.

[0078] According to certain embodiments, the anti-PCSK9 antibody used in the methods of the present disclosure is an antibody with pH-dependent binding characteristics. As used herein, the expression "pH-dependent binding" means that the antibody or antigen-binding fragment thereof exhibits "reduced binding to PCSK9 at acidic pH as compared to neutral pH" (for purposes of the present disclosure, both expressions may be used interchangeably). For the example, antibodies "with pH-dependent binding characteristics" includes antibodies and antigen-binding fragments thereof that bind PCSK9 with higher affinity at neutral pH than at acidic pH. In certain embodiments, the antibodies and antigen-binding fragments of the present disclosure bind PCSK9 with at least 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, or more times higher affinity at neutral pH than at acidic pH.

[0079] According to this aspect of the disclosure, the anti-PCSK9 antibodies with pH-dependent binding characteristics may possess one or more amino acid variations relative to the parental anti-PCSK9 antibody. For example, an anti-PCSK9 antibody with pH-dependent binding characteristics may contain one or more histidine substitutions or insertions, *e.g.*, in one or more CDRs of a parental anti-PCSK9 antibody. Thus, according to certain embodiments of the present disclosure, methods are provided comprising administering an anti-PCSK9 antibody which comprises CDR amino acid sequences (*e.g.*, heavy and light chain CDRs) which are identical to the CDR amino acid sequences of a parental anti-PCSK9 antibody, except for the substitution of one or more amino acids of one or more CDRs of the parental antibody with a histidine residue. The anti-PCSK9 antibodies with pH-dependent binding may possess, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or more histidine substitutions, either within a single CDR of a parental antibody or distributed throughout multiple (*e.g.*, 2, 3, 4, 5, or 6) CDRs of a parental anti-PCSK9 antibody. For example, the present disclosure includes the use of anti-PCSK9 antibodies with pH-dependent binding comprising one or more histidine substitutions in HCDR1, one or more histidine substitutions in HCDR2, one or more histidine substitutions in HCDR3, one or more histidine substitutions in LCDR1, one or more histidine substitutions in LCDR2, and/or one or more histidine substitutions in LCDR3, of a parental anti-PCSK9 antibody.

[0080] As used herein, the expression "acidic pH" means a pH of 6.0 or less (*e.g.*, less than about 6.0, less than about 5.5, less than about 5.0, etc.). The expression "acidic pH" includes pH values of about 6.0, 5.95, 5.90, 5.85, 5.8, 5.75, 5.7, 5.65, 5.6, 5.55, 5.5, 5.45, 5.4, 5.35, 5.3, 5.25, 5.2, 5.15, 5.1, 5.05, 5.0, or less. As used herein, the expression "neutral pH" means a pH of about 7.0 to about 7.4. The expression "neutral pH" includes pH values of about 7.0, 7.05, 7.1, 7.15, 7.2, 7.25, 7.3, 7.35, and 7.4.

[0081] The present disclosure includes anti-PCSK9 antibodies that bind to the same epitope as any of the specific exemplary antibodies described herein. Likewise, the present invention also includes anti-PCSK9 antibodies that compete for binding to PCSK9 or a PCSK9 fragment with any of the specific exemplary antibodies described herein.

5 **[0082]** Disclosed herein are anti-PCSK9 antibodies or antigen-binding fragments thereof which bind to the same epitope on PCSK9 as an antibody comprising heavy and light chain CDR amino acid sequences having SEQ ID NOs: 2, 3, 4, 7, 8, and 10. Also disclosed herein are anti-PCSK9 antibodies or antigen-binding fragments thereof which compete for binding to PCSK9 with an antibody comprising heavy and light chain CDR amino acid sequences having SEQ ID
10 NOs: 2, 3, 4, 7, 8, and 10.

[0083] One can easily determine whether an antibody binds to the same epitope as, or competes for binding with, a reference anti-PCSK9 antibody by using routine methods known in the art. For example, to determine if a test antibody binds to the same epitope as a reference anti-PCSK9 antibody of the invention, the reference antibody is allowed to bind to a PCSK9
15 protein or peptide under saturating conditions. Next, the ability of a test antibody to bind to the PCSK9 molecule is assessed. If the test antibody is able to bind to PCSK9 following saturation binding with the reference anti-PCSK9 antibody, it can be concluded that the test antibody binds to a different epitope than the reference anti-PCSK9 antibody. On the other hand, if the test antibody is not able to bind to the PCSK9 molecule following saturation binding with the
20 reference anti-PCSK9 antibody, then the test antibody may bind to the same epitope as the epitope bound by a reference anti-PCSK9 antibody described herein.

[0084] To determine if an antibody competes for binding with a reference anti-PCSK9 antibody, the above-described binding methodology is performed in two orientations: In a first orientation, the reference antibody is allowed to bind to a PCSK9 molecule under saturating
25 conditions followed by assessment of binding of the test antibody to the PCSK9 molecule. In a second orientation, the test antibody is allowed to bind to a PCSK9 molecule under saturating conditions followed by assessment of binding of the reference antibody to the PCSK9 molecule. If, in both orientations, only the first (saturating) antibody is capable of binding to the PCSK9 molecule, then it is concluded that the test antibody and the reference antibody compete for
30 binding to PCSK9. As will be appreciated by a person of ordinary skill in the art, an antibody that competes for binding with a reference antibody may not necessarily bind to the identical epitope as the reference antibody, but may sterically block binding of the reference antibody by binding an overlapping or adjacent epitope.

[0085] Two antibodies bind to the same or overlapping epitope if each competitively inhibits
35 (blocks) binding of the other to the antigen. That is, a 1-, 5-, 10-, 20- or 100-fold excess of one antibody inhibits binding of the other by at least 50% but preferably 75%, 90% or even 99% as

measured in a competitive binding assay (see, e.g., Junghans et al., Cancer Res, 1990:50:1495-1502). Alternatively, two antibodies have the same epitope if essentially all amino acid mutations in the antigen that reduce or eliminate binding of one antibody reduce or eliminate binding of the other. Two antibodies have overlapping epitopes if some amino acid mutations that reduce or eliminate binding of one antibody reduce or eliminate binding of the other.

[0086] Additional routine experimentation (e.g., peptide mutation and binding analyses) can then be carried out to confirm whether the observed lack of binding of the test antibody is in fact due to binding to the same epitope as the reference antibody or if steric blocking (or another phenomenon) is responsible for the lack of observed binding. Experiments of this sort can be performed using ELISA, RIA, surface plasmon resonance, flow cytometry or any other quantitative or qualitative antibody-binding assay available in the art.

[0087] Non-limiting examples of anti-PCSK9 antibodies that can be used in the context of the present disclosure include, e.g., alirocumab, bococizumab, or antigen-binding portions thereof.

Preparation of Human Antibodies

[0088] Methods for generating human antibodies in transgenic mice are known in the art. Any such known methods can be used in the context of the present disclosure to make human antibodies that specifically bind to human PCSK9.

[0089] Using VELOCIMMUNE™ technology (see, for example, US 6,596,541, Regeneron Pharmaceuticals) or any other known method for generating monoclonal antibodies, high affinity chimeric antibodies to PCSK9 are initially isolated having a human variable region and a mouse constant region. The VELOCIMMUNE® technology involves generation of a transgenic mouse having a genome comprising human heavy and light chain variable regions operably linked to endogenous mouse constant region loci such that the mouse produces an antibody comprising a human variable region and a mouse constant region in response to antigenic stimulation. The DNA encoding the variable regions of the heavy and light chains of the antibody are isolated and operably linked to DNA encoding the human heavy and light chain constant regions. The DNA is then expressed in a cell capable of expressing the fully human antibody.

[0090] Generally, a VELOCIMMUNE® mouse is challenged with the antigen of interest, and lymphatic cells (such as B-cells) are recovered from the mice that express antibodies. The lymphatic cells may be fused with a myeloma cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and selected to identify hybridoma cell lines that produce antibodies specific to the antigen of interest. DNA encoding the variable regions of the heavy chain and light chain may be isolated and linked to desirable isotypic constant regions of the heavy chain and light chain. Such an antibody protein may be produced in a cell, such as a CHO cell. Alternatively, DNA encoding the antigen-specific chimeric antibodies or the variable

domains of the light and heavy chains may be isolated directly from antigen-specific lymphocytes.

[0091] Initially, high affinity chimeric antibodies are isolated having a human variable region and a mouse constant region. The antibodies are characterized and selected for desirable characteristics, including affinity, selectivity, epitope, etc, using standard procedures known to those skilled in the art. The mouse constant regions are replaced with a desired human constant region to generate the fully human antibody of the disclosure, for example wild-type or modified IgG1 or IgG4. While the constant region selected may vary according to specific use, high affinity antigen-binding and target specificity characteristics reside in the variable region.

[0092] In general, the antibodies that can be used in the methods of the present disclosure possess high affinities, as described above, when measured by binding to antigen either immobilized on solid phase or in solution phase. The mouse constant regions are replaced with desired human constant regions to generate the fully human antibodies of the disclosure. While the constant region selected may vary according to specific use, high affinity antigen-binding and target specificity characteristics reside in the variable region.

[0093] Specific examples of human antibodies or antigen-binding fragments of antibodies that specifically bind PCSK9 which can be used in the context of the methods of the present disclosure include any antibody or antigen-binding fragment which comprises the three heavy chain CDRs (HCDR1, HCDR2 and HCDR3) contained within a heavy chain variable region (HCVR) having an amino acid sequence selected from the group consisting of SEQ ID NOs:1 and 11, or a substantially similar sequence thereof having at least 90%, at least 95%, at least 98% or at least 99% sequence identity. Alternatively, specific examples of human antibodies or antigen-binding fragments of antibodies that specifically bind PCSK9 which can be used in the context of the methods of the present disclosure include any antibody or antigen-binding fragment which comprises the three heavy chain CDRs (HCDR1, HCDR2 and HCDR3) contained within a heavy chain variable region (HCVR) having an amino acid sequence selected from the group consisting of SEQ ID NOs 37, 45, 53, 61, 69, 77, 85, 93, 101, 109, 117, 125, 133, 141, 149, 157, 165, 173, 181, and 189, or a substantially similar sequence thereof having at least 90%, at least 95%, at least 98% or at least 99% sequence identity. The antibody or antigen-binding fragment may comprise the three light chain CDRs (LCVR1, LCVR2, LCVR3) contained within a light chain variable region (LCVR) having an amino acid sequence selected from the group consisting of SEQ ID NOs 6 and 15, or a substantially similar sequence thereof having at least 90%, at least 95%, at least 98% or at least 99% sequence identity. Alternatively, the antibody or antigen-binding fragment may comprise the three light chain CDRs (LCVR1, LCVR2, LCVR3) contained within a light chain variable region (LCVR) having an amino acid sequence selected from the group consisting of SEQ ID NOs 41, 49, 57, 65, 73, 81, 89, 97, 105, 113, 121, 129, 137, 145, 153, 161, 169, 177, 185, and 193, or a substantially similar sequence

thereof having at least 90%, at least 95%, at least 98% or at least 99% sequence identity.

[0094] Sequence identity between two amino acids sequences is determined over the entire length of the reference amino acid sequence, i.e. the amino acid sequence identified with a SEQ ID NO, using the best sequence alignment and/or over the region of the best sequence

5 alignment between the two amino acid sequences, wherein the best sequence alignment can be obtained with art known tools, e.g. Align, using standard settings, preferably EMBOSS::needle, Matrix: Blosum62, Gap Open 10.0, Gap Extend 0.5.

[0095] In certain embodiments of the present disclosure, the antibody or antigen binding fragment thereof comprises the six CDRs (HCDR1, HCDR2, HCDR3, LCDR1, LCDR2 and

10 LCDR3) from the heavy and light chain variable region amino acid sequence pairs (HCVR/LCVR) selected from the group consisting of SEQ ID NOs:1/6 and 11/15. Alternatively, in certain embodiments of the present disclosure, the antibody or antigen binding fragment thereof comprises the six CDRs (HCDR1, HCDR2, HCDR3, LCDR1, LCDR2 and LCDR3) from the heavy and light chain variable region amino acid sequence pairs (HCVR/LCVR) selected

15 from the group consisting of SEQ ID NOs:37/41, 45/49, 53/57, 61/65, 69/73, 77/81, 85/89, 93/97, 101/105, 109/113, 117/121, 125/129, 133/137, 141/145, 149/153, 157/161, 165/169, 173/177, 181/185, and 189/193.

[0096] In certain embodiments of the present disclosure, the anti-PCSK9 antibody, or antigen binding fragment thereof, that can be used in the methods of the present disclosure has

20 HCDR1/HCDR2/HCDR3/LCDR1/LCDR2/LCDR3 amino acid sequences selected from SEQ ID NOs: 2/3/4/7/8/10 (mAb316P [also referred to as "REGN727," or "alirocumab"]) and 12/13/14/16/17/18 (mAb300N) (See US Patent App. Publ No. 2010/0166768) and 12/13/14/16/17/18, wherein SEQ ID NO:16 comprises a substitution of histidine for leucine at amino acid residue 30 (L30H).

25 **[0097]** In certain embodiments of the present disclosure, the antibody or antigen binding fragment thereof comprises HCVR/LCVR amino acid sequence pairs selected from the group consisting of SEQ ID NOs:1/6 and 11/15. In certain exemplary embodiments, the antibody or antigen binding fragment thereof comprises an HCVR amino acid sequence of SEQ ID NO:1 and an LCVR amino acid sequence of SEQ ID NO:6. In certain exemplary embodiments, the

30 antibody or antigen binding fragment thereof comprises an HCVR amino acid sequence of SEQ ID NO:11 and an LCVR amino acid sequence of SEQ ID NO:15. In certain exemplary embodiments, the antibody or antigen binding fragment thereof comprises an HCVR amino acid sequence of SEQ ID NO:11 and an LCVR amino acid sequence of SEQ ID NO:15 comprising a substitution of histidine for leucine at amino acid residue 30 (L30H).

35 **Pharmaceutical Compositions and Methods of Administration**

[0098] The present disclosure includes methods that comprise administering a PCSK9

inhibitor to a patient, wherein the PCSK9 inhibitor is contained within a pharmaceutical composition. The pharmaceutical compositions of the disclosure are formulated with suitable carriers, excipients, and other agents that provide suitable transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. See also Powell et al. "Compendium of excipients for parenteral formulations" PDA (1998) J Pharm Sci Technol 52:238-311.

[0099] Various delivery systems are known and can be used to administer the pharmaceutical composition of the disclosure, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the mutant viruses, receptor mediated endocytosis (see, e.g., Wu et al., 1987, J. Biol. Chem. 262:4429-4432). Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The composition may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents.

[00100] A pharmaceutical composition of the present disclosure can be delivered subcutaneously or intravenously with a standard needle and syringe. In addition, with respect to subcutaneous delivery, a pen delivery device readily has applications in delivering a pharmaceutical composition of the present disclosure. Such a pen delivery device can be reusable or disposable. Such a pen delivery device can be prefilled. A reusable pen delivery device generally utilizes a replaceable cartridge that contains a pharmaceutical composition. Once all of the pharmaceutical composition within the cartridge has been administered, and the cartridge is empty, the empty cartridge can readily be discarded and replaced with a new cartridge that contains the pharmaceutical composition. The pen delivery device can then be reused. In a disposable pen delivery device, there is no replaceable cartridge. Rather, the disposable pen delivery device comes prefilled with the pharmaceutical composition held in a reservoir within the device. Once the reservoir is emptied of the pharmaceutical composition, the entire device is discarded.

[00101] Numerous reusable pen and autoinjector delivery devices have applications in the subcutaneous delivery of a pharmaceutical composition of the present disclosure. Examples include, but are not limited to AUTOPEN™ (Owen Mumford, Inc., Woodstock, UK), DISETRONIC™ pen (Disetronic Medical Systems, Bergdorf, Switzerland), HUMALOG MIX

75/25™ pen, HUMALOG™ pen, HUMALIN 70/30™ pen (Eli Lilly and Co., Indianapolis, IN), NOVOPEN™ I, II and III (Novo Nordisk, Copenhagen, Denmark), NOVOPEN JUNIOR™ (Novo Nordisk, Copenhagen, Denmark), BD™ pen (Becton Dickinson, Franklin Lakes, NJ), OPTIPEN™, OPTIPEN PRO™, OPTIPEN STARLET™, and OPTICLIK™ (Sanofi-Aventis, Frankfurt, Germany), to name only a few. Examples of disposable pen delivery devices having applications in subcutaneous delivery of a pharmaceutical composition of the present disclosure include, but are not limited to the SOLOSTAR™ pen (Sanofi-Aventis), the FLEXPEN™ (Novo Nordisk), and the KWIKPEN™ (Eli Lilly), the SURECLICK™ Autoinjector (Amgen, Thousand Oaks, CA), the PENLET™ (Haselmeier, Stuttgart, Germany), the EPIPEN (Dey, L.P.), and the HUMIRA™ Pen (Abbott Labs, Abbott Park IL), to name only a few.

[00102] In certain situations, the pharmaceutical composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, supra; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:201). In another embodiment, polymeric materials can be used; see, Medical Applications of Controlled Release, Langer and Wise (eds.), 1974, CRC Pres., Boca Raton, Florida. In yet another embodiment, a controlled release system can be placed in proximity of the composition's target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, 1984, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138). In some embodiments, the pharmaceutical composition can be contained in a microinfusor. Other controlled release systems are discussed in the review by Langer, 1990, Science 249:1527-1533.

[00103] The injectable preparations may include dosage forms for intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, etc. These injectable preparations may be prepared by known methods. For example, the injectable preparations may be prepared, e.g., by dissolving, suspending or emulsifying the antibody or its salt described above in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared is preferably filled in an appropriate ampoule.

[00104] Advantageously, the pharmaceutical compositions for oral or parenteral use described above are prepared into dosage forms in a unit dose suited to fit a dose of the active ingredients. Such dosage forms in a unit dose include, for example, tablets, pills, capsules, injections (ampoules), suppositories, etc. In some embodiments, the composition is contained

in a glass vial.

[00105] Exemplary pharmaceutical formulations comprising an anti-PCSK9 antibody that can be used in the context of the methods of the present disclosure are set forth, e.g., in US 2013/0189277, the disclosure of which is hereby incorporated by reference in its entirety.

5 Dosage

[00106] The amount of PCSK9 inhibitor (e.g., anti-PCSK9 antibody) administered to a subject according to the methods of the present disclosure is, generally, a therapeutically effective amount. As used herein, the phrase "therapeutically effective amount" means a dose of PCSK9 inhibitor that results in a detectable reduction (at least about 5%, 10%, 15%, 20%, 25%, 30%, 10 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or more from baseline) in one or more parameters selected from the group consisting of LDL-C, ApoB100, non-HDL-C, total cholesterol, VLDL-C, triglycerides, Lp(a) and remnant cholesterol.

[00107] In the case of an anti-PCSK9 antibody, a therapeutically effective amount can be from about 0.05 mg to about 600 mg, e.g., about 0.05 mg, about 0.1 mg, about 1.0 mg, about 1.5 mg, 15 about 2.0 mg, about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 75 mg, about 80 mg, about 90 mg, about 100 mg, about 110 mg, about 120 mg, about 130 mg, about 140 mg, about 150 mg, about 160 mg, about 170 mg, about 180 mg, about 190 mg, about 200 mg, about 210 mg, about 220 mg, about 230 mg, about 240 mg, about 250 mg, about 260 mg, about 270 mg, about 280 mg, about 290 mg, about 300 mg, about 20 310 mg, about 320 mg, about 330 mg, about 340 mg, about 350 mg, about 360 mg, about 370 mg, about 380 mg, about 390 mg, about 400 mg, about 410 mg, about 420 mg, about 430 mg, about 440 mg, about 450 mg, about 460 mg, about 470 mg, about 480 mg, about 490 mg, about 500 mg, about 510 mg, about 520 mg, about 530 mg, about 540 mg, about 550 mg, about 560 mg, about 570 mg, about 580 mg, about 590 mg, or about 600 mg, of the anti-PCSK9 antibody.

25 [00108] The amount of anti-PCSK9 antibody contained within the individual doses may be expressed in terms of milligrams of antibody per kilogram of patient body weight (*i.e.*, mg/kg). For example, the anti-PCSK9 antibody may be administered to a patient at a dose of about 0.0001 to about 10 mg/kg of patient body weight.

Combination Therapies

30 [00109] According to certain embodiments of the present disclosure, additional therapeutic agents, besides a statin, may be administered to the patient in combination with a PCSK9 inhibitor. Examples of such additional therapeutic agents include e.g., (1) an agent which inhibits cholesterol uptake and or bile acid re-absorption (e.g., ezetimibe); (2) an agent which increase lipoprotein catabolism (such as niacin); and/or (3) activators of the LXR transcription 35 factor that plays a role in cholesterol elimination such as 22-hydroxycholesterol. According to

certain embodiments, an anti-ANGPTL3 antibody (such as evinacumab) is administered in combination with a PCSK9 inhibitor in the context of the methods of the present disclosure.

Administration Regimens

[00110] According to certain embodiments of the present disclosure, one or more doses of a PCSK9 inhibitor (*i.e.*, a pharmaceutical composition comprising a PCSK9 inhibitor) may be administered to a subject over a defined time course (*e.g.*, in place of a daily therapeutic statin regimen). The methods according to this aspect of the disclosure comprise sequentially administering to a subject one or more doses of a PCSK9 inhibitor. As used herein,

"sequentially administering" means that each dose of PCSK9 inhibitor is administered to the subject at a different point in time, *e.g.*, on different days separated by a predetermined interval (*e.g.*, hours, days, weeks or months). The present disclosure includes methods that comprise sequentially administering to the patient a single initial dose of a PCSK9 inhibitor, followed by one or more secondary doses of the PCSK9 inhibitor, and optionally followed by one or more tertiary doses of the PCSK9 inhibitor.

[00111] The terms "initial dose," "secondary doses," and "tertiary doses," refer to the temporal sequence of administration of the individual doses of a pharmaceutical composition comprising a PCSK9 inhibitor. Thus, the "initial dose" is the dose that is administered at the beginning of the treatment regimen (also referred to as the "baseline dose"); the "secondary doses" are the doses that are administered after the initial dose; and the "tertiary doses" are the doses that are administered after the secondary doses. The initial, secondary, and tertiary doses may all contain the same amount of the PCSK9 inhibitor, but generally may differ from one another in terms of frequency of administration. In certain embodiments, however, the amount of PCSK9 inhibitor contained in the initial, secondary and/or tertiary doses varies from one another (*e.g.*, adjusted up or down as appropriate) during the course of treatment. In certain embodiments, two or more (*e.g.*, 2, 3, 4, or 5) doses are administered at the beginning of the treatment regimen as "loading doses" followed by subsequent doses that are administered on a less frequent basis (*e.g.*, "maintenance doses").

[00112] According to exemplary embodiments of the present disclosure, each secondary and/or tertiary dose is administered 1 to 26 (*e.g.*, 1, 1½, 2, 2½, 3, 3½, 4, 4½, 5, 5½, 6, 6½, 7, 7½, 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12, 12½, 13, 13½, 14, 14½, 15, 15½, 16, 16½, 17, 17½, 18, 18½, 19, 19½, 20, 20½, 21, 21½, 22, 22½, 23, 23½, 24, 24½, 25, 25½, 26, 26½, or more) weeks after the immediately preceding dose. The phrase "the immediately preceding dose," as used herein, means, in a sequence of multiple administrations, the dose of antigen-binding molecule that is administered to a patient prior to the administration of the very next dose in the sequence with no intervening doses.

[00113] The methods according to this aspect of the disclosure may comprise administering to a patient any number of secondary and/or tertiary doses of a PCSK9 inhibitor. For example, in certain embodiments, only a single secondary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) secondary doses are administered to the patient. Likewise, in certain embodiments, only a single tertiary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) tertiary doses are administered to the patient.

[00114] In embodiments involving multiple secondary doses, each secondary dose may be administered at the same frequency as the other secondary doses. For example, each secondary dose may be administered to the patient 1 to 2, 4, 6, 8 or more weeks after the immediately preceding dose. Similarly, in embodiments involving multiple tertiary doses, each tertiary dose may be administered at the same frequency as the other tertiary doses. For example, each tertiary dose may be administered to the patient 1 to 2, 4, 6, 8 or more weeks after the immediately preceding dose. Alternatively, the frequency at which the secondary and/or tertiary doses are administered to a patient can vary over the course of the treatment regimen. The frequency of administration may also be adjusted during the course of treatment by a physician depending on the needs of the individual patient following clinical examination.

[00115] According to certain embodiments of the present disclosure, multiple doses of a pharmaceutical composition comprising about 75 mg of anti-PCSK9 antibody are administered to a patient at a frequency of once every two weeks.

[00116] According to certain embodiments of the present disclosure, multiple doses of a pharmaceutical composition comprising about 150 mg of anti-PCSK9 antibody are administered to a patient at a frequency of once every two weeks.

[00117] According to certain embodiments of the present disclosure, multiple doses of a pharmaceutical composition comprising about 75 mg of anti-PCSK9 antibody are administered to a patient at a frequency of once every four weeks.

[00118] According to certain embodiments of the present disclosure, multiple doses of a pharmaceutical composition comprising about 150 mg of anti-PCSK9 antibody are administered to a patient at a frequency of once every four weeks.

[00119] The present disclosure includes administration regimens comprising an up-titration option (also referred to herein as "dose modification"). As used herein, an "up-titration option" means that, after receiving a particular number of doses of a PCSK9 inhibitor, if a patient has not achieved a specified reduction in one or more defined therapeutic parameters, the dose of the PCSK9 inhibitor is thereafter increased. For example, in the case of a therapeutic regimen comprising administration of 75 mg doses of an anti-PCSK9 antibody to a patient at a frequency of once every two weeks, if after 8 weeks (i.e., 5 doses administered at Week 0, Week 2 and

Week 4, Week 6 and Week 8), the patient has not achieved a serum LDL-C concentration of less than 70 mg/dL, then the dose of anti-PCSK9 antibody is increased to e.g., 150 mg administered once every two weeks thereafter (e.g., starting at Week 10 or Week 12, or later).

5 [00120] In certain embodiments, the anti-PCSK9 antibody is administered to a subject at a dose of about 75 mg every two weeks, for example for at least three doses.

[00121] In certain embodiments, the anti-PCSK9 antibody is administered to a subject at a dose of about 150 mg every two weeks, for example for at least three doses.

10 [00122] In some embodiments, the antibody is administered to a subject at a dose of about 75 mg every two weeks for 12 weeks, and the dose remains at 75 mg every two weeks if, at week 8, the subject's LDL-C value was less than 100 mg/dl and a 30% reduction of LDL-C.

[00123] In other embodiments, the antibody is administered to a subject at a dose of about 75 mg every two weeks for 12 weeks, and the dose is titrated up to about 150 mg every two weeks if, at week 8, the subject's LDL-C value was greater than or equal to 100 mg/dl.

15 [00124] In some embodiments, the antibody is administered to a subject at a dose of about 75 mg every two weeks for 12 weeks, and the dose remains at 75 mg every two weeks if, at week 8, the subject's LDL-C value was less than 70 mg/dl and a 30% reduction of LDL-C.

[00125] In another embodiment, the antibody is administered to a subject at a dose of about 300 mg every four weeks.

20 [00126] In a further embodiment, the antibody is administered to a subject at a dose of about 300 mg every four weeks for a total of three doses, and the dose is changed to 150 mg every two weeks for another 36 weeks if, at week 8, the subject did not achieve a pre-determined treatment goal or the subject did not have at least a 30% reduction of LDL-C from baseline.

[00127] In certain embodiments, the anti-PCSK9 antibody is administered to a subject at a dose of about 150 mg every four weeks for at least three doses.

25 [00128] In some embodiments, the antibody is administered to a subject at a dose of about 150 mg every four weeks for 12 weeks, and the dose remains at 150 mg every four weeks if, at week 8, the subject's LDL-C value was less than 100 mg/dl and a 30% reduction of LDL-C.

30 [00129] In other embodiments, the antibody is administered to a subject at a dose of about 150 mg every four weeks for 12 weeks, and the dose is titrated up to about 300 mg every two weeks if, at week 8, the subject's LDL-C value was greater than or equal to 100 mg/dl.

[00130] In some embodiments, the antibody is administered to a subject at a dose of about 150 mg every four weeks for 12 weeks, and the dose remains at 150 mg every four weeks for another 12 weeks if, at week 8, the subject's LDL-C value was less than 70 mg/dl and a 30% reduction of LDL-C.

35 [00131] In another embodiment, the antibody is administered to a subject at a dose of about 300 mg every four weeks.

[00132] In a further embodiment, the antibody is administered to a subject at a dose of about 300 mg every four weeks for a total of three doses, and the dose is changed to 150 mg every two weeks for another 36 weeks if, at week 8, the subject did not achieve a pre-determined treatment goal or the subject did not have at least a 30% reduction of LDL-C from baseline.

EXAMPLES

[00133] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the disclosure, and are not intended to limit the scope of what the inventors regard as their disclosure. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1. Generation of Human Antibodies to Human PCSK9

[00134] Human anti-PCSK9 antibodies were generated as described in US Patent No. 8,062,640. The exemplary PCSK9 inhibitor used in the following Example is the human anti-PCSK9 antibody designated "mAb316P," also known as "REGN727," or "alirocumab."

mAb316P has the following amino acid sequence characteristics: a heavy chain comprising SEQ ID NO:5 and a light chain comprising SEQ ID NO:9; a heavy chain variable region (HCVR) comprising SEQ ID NO:1 and a light chain variable domain (LCVR) comprising SEQ ID NO:6; a heavy chain complementarity determining region 1 (HCDR1) comprising SEQ ID NO:2, a HCDR2 comprising SEQ ID NO:3, a HCDR3 comprising SEQ ID NO:4, a light chain complementarity determining region 1 (LCDR1) comprising SEQ ID NO:7, a LCDR2 comprising SEQ ID NO:8 and a LCDR3 comprising SEQ ID NO:10.

Example 2: A Randomized, Double-Blind, Placebo-controlled, Parallel-group Study to Evaluate the Efficacy and Safety of an Anti-PCSK9 Antibody ("alirocumab") in Patients with Homozygous Familial Hypercholesterolemia

[00135] The objective of the instant study was to evaluate the efficacy, safety and tolerability of an anti-PCSK9 antibody ("alirocumab") in patients with hoFH (excluding those patients with null/null mutations in both LDLR alleles). More specifically, an objective of the instant study was to demonstrate the reduction of LDL-C with alirocumab 150 mg subcutaneous (SC) every 2 weeks (Q2W) in comparison to placebo after 12 weeks of treatment. Secondary objectives of the study included: evaluating the effect of alirocumab 150 mg Q2W on other lipid parameters

(i.e., apolipoprotein [Apo] A-1 and B, non-high-density lipoprotein cholesterol [non HDL C], total cholesterol [TC], proportion of patients with 15%, 30%, and 50% LDL-C reductions, Lp(a), HDL-C, triglycerides [TG]) in patients with hoFH), evaluating the safety and tolerability of alirocumab 150 mg SC Q2W in patients with hoFH, assessing the pharmacokinetics (PK) of alirocumab 150 mg SC Q2W in patients with hoFH, and assessing the potential development of anti-drug (alirocumab) antibodies.

[00136] Other objectives of the instant study included: collecting genotype information for all patients to characterize hoFH mutation status in order to explore potential differences in efficacy and safety, assessing the effect of alirocumab on eligibility for apheresis (using German and US apheresis criteria), and assessing the effect of alirocumab on quality of life using the EQ-5D QOL questionnaire.

[00137] Patients with hoFH have persistently elevated LDL-C, which is a contributing factor to a number of health concerns, principally accelerated atherosclerosis resulting in premature CV disease. Despite treatment with LMTs such as pharmacological agents including statins, and mechanical removal of lipids by LDL apheresis, many patients with hoFH remain far from their LDL-C treatment goal. In a cohort study, the age of first CVD event was in the third decade of life in hoFH (Raaij, *et al.* 2011 *Circulation* 124(20):2202-2207). While this is later than the early teen years as seen before implementation of medical regimens such as statins, the need for more intensive treatments remain in order to delay the onset of CVD and occurrence of events.

[00138] In order to evaluate the efficacy and safety of alirocumab in patients with hoFH, the instant study was carried out. The study population included individuals ≥ 18 years of age. Diagnosis of hoFH was based on either genotyping data or clinical criteria. The genetic definition included all individuals considered to be true homozygotes, compound heterozygotes, or double heterozygotes for mutations in the LDLR, ApoB, PCSK9, or LDLRAP1 genes.

However, individuals with history null/null LDLR mutations were excluded.

[00139] Percent change in LDL-C from baseline was the primary endpoint. Low-density lipoprotein cholesterol is an accepted surrogate endpoint for CV risk and has repeatedly been used as the primary endpoint for approval of multiple other hoFH treatments. The instant study was designed as a placebo controlled trial, with the addition of alirocumab on top of patients' existing treatment regimens of maximally tolerated LMT, including lipid apheresis. An optional run-in period was utilized for those patients that have not yet achieved a stable background treatment regimen that would be required to be maintained throughout the double-blind treatment period. An add-on design was appropriate, because removal of any therapies from the patient's existing treatment regimen would lead to an increase in LDL-C and possibly contribute to the serious CV sequelae seen in this disease. Treatment duration of 12 weeks for the primary endpoint allowed alirocumab to achieve steady state and exert its full effect on LDL

C. An additional 12-week open-label treatment period in which all patients were administered alirocumab allowed further assessment of safety in this population.

Rationale for Dose Selection

[00140] Alirocumab 75 mg and 150 mg SC Q2W are the approved doses and is currently authorized in 40 countries worldwide (including the US, European Union, Canada, Norway, Iceland, Brazil, and Japan). Because patients with hoFH are proven to be a hard-to-treat population compared to non-FH and HeFH patients and will have a very high baseline LDL-C far from their target level, the dose of alirocumab proposed for the instant study is the highest approved dose, 150 mg SC Q2W.

[00141] In addition to being efficacious, alirocumab has a favorable safety profile. Overall, the most commonly occurring treatment-emergent adverse events (TEAEs), reported in a higher proportion of patients in the alirocumab group compared to placebo (i.e., incidence $\geq 2.0\%$ in the alirocumab group) were: injection site reaction (7.2% vs 5.1%), nasopharyngitis (11.3% vs 11.1%), influenza (5.7% vs 4.6%), myalgia (4.3% vs 3.4%), urinary tract infection (4.8% vs 4.6%), diarrhea (4.7% vs 4.4%) and bronchitis (4.3% vs 3.8%) (Praluent Product Insert). No differences in the safety profile have been observed between the two approved doses of 75 mg and 150 mg.

[00142] It was hypothesized that treatment with alirocumab 150 mg Q2W in patients with hoFH already receiving maximally tolerated LMT or LDL apheresis would be well-tolerated and have an acceptable safety profile while providing maximal LDL-C lowering effect.

Benefit / Risk Assessment

[00143] Patients with hoFH have extremely high LDL-C levels, are far from their target LDL-C, and require significant reductions in LDL-C. Despite the approval of newer treatments including lomitapide and mipomersen, the need for more intensive therapies remain. PCSK9 inhibitors are a new addition to the armamentarium of LMT that has proven to profoundly decrease LDL-C. The body of evidence from the statin literature shows that the relationship between LDL-C reduction and CV event reduction is approximately linear, and within the context of the instant study in the hoFH patient population, even a modest reduction in LDL-C would translate into significant benefit for these patients.

[00144] It was expected that treatment with alirocumab would be well tolerated and have an acceptable safety profile. The accumulated safety information showed that the most commonly occurring TEAEs with alirocumab were nasopharyngitis, injection site reactions, influenza, myalgia, musculoskeletal pain, and contusion. Moreover, the rates of these adverse events (AEs) were relatively low, ranging from 11.3% to 2.1% for the alirocumab treatment group (vs 11.1% - 1.6% in the placebo group) (Praluent Product Insert).

[00145] Taken together, these data indicated that the benefit/risk assessment of treatment with alirocumab in the hoFH populations could be favorable.

Demographic and Baseline Characteristics

[00146] Baseline characteristics included standard demography (age, race, weight, height, etc),
5 disease characteristics including lipid levels, mutation status, medical history, medication history
and apheresis schedule (if applicable) for each patient.

Primary and Secondary Endpoints

[00147] The primary efficacy endpoint was the percent change in LDL-C from baseline to week
12 in the ITT population for alirocumab 150 mg Q2W as compared with placebo in patients with
10 hoFH. The percent change in LDL-C from baseline to week 12 was defined as: $[100 \times (\text{LDL-C value at week 12} - \text{LDL-C value at baseline}) / \text{LDL-C value at baseline}]$. For LDL-C analysis, both
calculated and measured LDL-C values were taken into account. In case both calculated and
measured LDL-C values were available for the same sampling time point, the measured LDL-C
was considered. The baseline LDL-C value was the last LDL C value obtained before the first
15 dose of double blind-study drug. For randomized but not-treated patients, baseline was defined
as the last value before randomization. The LDL C at week 12 was the LDL-C value obtained
within the week 12 analysis window, regardless of adherence to treatment (ITT estimand).

[00148] All calculated and measured LDL-C values (scheduled or unscheduled, fasting or not
fasting) could be used for the primary efficacy endpoint, if appropriate, according to the above
20 definition. The analysis window used to allocate a time point to a measurement was defined in
the statistical analysis plan (SAP).

[00149] Key secondary efficacy endpoints included: the percent change in Apo B from baseline
to week 12 (ITT estimand), the percent change in non-HDL-C from baseline to week 12 (ITT
estimand), the percent change in total cholesterol from baseline to week 12 (ITT estimand),
25 proportion of patients with $\geq 15\%$ reduction in LDL-C at week 12 (ITT estimand), proportion of
patients with $\geq 30\%$ reduction in LDL-C at week 12 (ITT estimand), the percent change in Lp(a)
from baseline to week 12 (ITT estimand), proportion of patients with $\geq 50\%$ reduction in LDL-C at
week 12 (ITT estimand), the percent change in HDL-C from baseline to week 12 (ITT estimand),
the percent change in fasting TG from baseline to week 12 (ITT estimand), and the percent
30 change in Apo A-1 from baseline to week 12 (ITT estimand). The same definition and rules
applied to these key secondary efficacy endpoints as were applied to the primary efficacy
endpoint.

[00150] Other secondary efficacy endpoints included the percent change in LDL-C from
baseline to week 12 in the modified (m)ITT population (all randomized population who took at
35 least 1 dose or part of a dose of double-blind investigational study drug and has an evaluable
primary endpoint), using all LDL-C values within the week 12 analysis window and during the

efficacy treatment period (on-treatment estimand), the percent change in Apo B, non-HDL-C, TC, Lp(a), HDL-C, fasting TG and Apo A-1 from baseline to week 12 (on-treatment estimand), proportion of patients with $\geq 15\%$ reduction, $\geq 30\%$ reduction, and $\geq 50\%$ reduction in LDL-C at week 12 (on treatment estimand), and the absolute change in the ratio of Apo B/Apo A-1 from baseline to week 12 (ITT estimand). The efficacy treatment period was defined as the time from the first double-blind study drug injection up to 21 days after the last double-blind study drug injection, or the first open-label alirocumab injection (if applicable), whichever came first.

[00151] Safety endpoints constituted safety parameters (AEs, laboratory data, vital signs, and electrocardiogram [ECG]) assessed throughout the study. Other endpoints included exploratory relationships between hoFH genotype status and lipid parameters, the change in the proportion of patients who meet US apheresis eligibility criteria from baseline to week 12 (Goldberg, *et al.* 2011 *J Clin Lipidol* 5(3 Supp):S1-S8), the change in the proportion of patients who meet German apheresis eligibility criteria from baseline to week 12 (Schettler, *et al.* 2012 *Clin Res Cardiol Suppl* 7:15-19), and response of each EQ-5D item, index score, and change of index score from baseline through week 12.

Pharmacokinetic Variables

[00152] The pharmacokinetic (PK) variable was alirocumab serum concentration collected at specified sampling time.

Anti-Drug Antibody Variables

[00153] Anti-drug (alirocumab) antibody status were assessed: total patients negative in the ADA assay at all time points, pre-existing immunoreactivity (defined as either an anti-drug antibody (ADA) positive response in the assay at baseline with all post-dose ADA results negative OR a positive response at baseline with all post-treatment ADA responses less than 4-fold baseline titer levels), and/or treatment emergent (defined as either any post-dose positive ADA response when baseline results were negative OR any post-dose positive ADA response that was at least 4-fold over the baseline level when baseline was positive in the ADA assay).

[00154] Samples positive in the ADA assay were assessed for titer. The titer categories included low (titer $< 1,000$), moderate ($1,000 \leq \text{titer} \leq 10,000$), and high (titer $> 10,000$).

[00155] Samples positive in the ADA assay were assessed for neutralizing activity.

Study Design

[00156] The instant study was a randomized, double-blind, placebo-controlled, parallel-group study to evaluate the efficacy and safety of alirocumab in patients with hoFH.

[00157] Approximately 74 patients were randomized in a 2:1 ratio to receive either alirocumab 150 mg SC Q2W or matching placebo. Randomization was stratified by apheresis treatment status (Yes/No).

[00158] The study consisted of up to 4 periods: an optional 4-week run-in period (for patients whose background medical LMT regimen or apheresis schedule and/or apheresis settings were stable prior to screening), a 2-week screening period, a 12-week double-blind treatment period, and a mandatory 12-week open-label treatment period according to the following study flow

5 chart:

Optional Run-in		Screening		Open-Label Treatment	
		Baseline	End of Double-Blind	End of Open-Label	End of Study
(Day -42)	(Day -14)	(Day 1)	(Day 85)	(Day 169)	(Day 225)

Patients not continuing on to another lipid lowering study also underwent an 8-week follow-up period.

[00159] Optional run-ins included undergoing apheresis therapy and lipid-modifying therapy.

[00160] Patients who were undergoing apheresis therapy had to be on a stable weekly or every other week schedule. Patients whose schedule or apheresis settings had not been stable for at least 8 weeks before the screening visit entered a 4-week run-in period before the screening period. After the 4-week run-in period, patients whose lipid-apheresis schedule/settings remained stable (and were stable for at least 8 weeks in total) were eligible to enter the 2-week screening period. Additionally, all patients on LDL apheresis had to be diagnosed based on genotype and, if genotype information had not been determined previously, they could enter the run-in to allow time, if needed, to determine their mutation status.

[00161] Patients who were on background lipid-modifying therapy (LMT) that had not been stable for at least 4 weeks before the screening visit entered a 4-week run-in period to stabilize their LMT before entering the screening period. Patients who had not been on a stable dose of mipomersen for 6 months prior to screening or on a maximum tolerated dose of lomitapide for 12 weeks prior to screening were excluded.

Screening

[00162] Once on a stable background regimen as defined above, patients entered a 2-week screening period. Initial eligibility was determined during this screening period by standard screening procedures. A DNA sample was collected for hoFH mutation status.

[00163] Patients were to be on a stable low fat or heart-healthy diet throughout the duration of the study, starting at screening through the end of the double-blind treatment period and through the open label treatment period. Patients' exercise regimen was to remain stable throughout the duration of the study, from screening, through the end of the double-blind treatment period and through the open-label treatment period.

[00164] The patient or caregiver was trained to self-inject/inject using a dose of placebo during the screening period or at the first visit of the double-blind treatment period.

Double-blind Treatment

5 [00165] Patients who met all inclusion criteria AND who met none of the exclusion criteria were randomized in a 2:1 ratio to receive: alirocumab 150 mg SC Q2W OR matching placebo SC Q2W. What was received is also referred to herein as “study drug” and “investigational medical product”.

10 [00166] Study drug administration during the double-blind treatment period started on the day of randomization and was administered immediately after completion of the LDL apheresis procedure (if applicable). For those patients not undergoing LDL apheresis, administration of study drug was made after all samples for clinical laboratory evaluation were obtained. The last injection during the double-blind treatment period was on day 71/week 10.

15 [00167] For all patients undergoing LDL apheresis, all samples for clinical laboratory evaluation were obtained immediately prior to the LDL apheresis procedure and prior to administration of study drug. Given the impact of LDL apheresis on lipid parameters, it was important to match the time of the baseline activities with the timing of the week 12 activities. This meant that timing between the baseline sample collection relative to the most recently completed LDL apheresis procedure should match the timing of the week 12 sample collection relative to the most recently completed LDL apheresis procedure.

20 [00168] For all patients who were not undergoing apheresis, all samples for clinical laboratory evaluation were obtained prior to administration of investigational medical product. The efficacy of alirocumab was assessed by clinical laboratory evaluation of lipid levels at pre-specified time points throughout the study.

25 [00169] Patients who were receiving LMT or who were undergoing apheresis maintained stable LMT and a stable apheresis schedule (as applicable) throughout the duration of the study, from screening through the end of the open-label treatment period/end-of-study visit (week 24).

30 [00170] Patients who prematurely discontinued study drug during the double-blind treatment period were encouraged to remain in the study and undergo all double-blind study visits and procedures with the exception of dosing with study drug. At the time of study drug discontinuation, the patient was to have, as soon as possible, an unscheduled visit with assessments normally planned at the end of the double-blind treatment visit (this was to take place within 5 days of discontinuation of study drug, if possible).

Open-label Treatment

35 [00171] To provide further safety data in this rare patient population, all patients participated in an open-label treatment period. Regardless of treatment assignment in the double-blind treatment period, patients received open-label study drug (alirocumab 150 mg SC Q2W) starting

at week 12 (day 85) and continuing through week 24 (end of open-label treatment period, last injection of study drug on day 155/week 22). Patients who were receiving LMT or who were undergoing LDL apheresis continued a stable dose and regimen and a stable LDL apheresis schedule and settings (as applicable) throughout the duration of the open-label treatment period.

5 [00172] Upon completion of the open-label treatment period, patients either participated in an additional lipid-lowering clinical trial or directly underwent an 8-week follow-up period. A follow-up phone call was made at week 32 to collect AE and concomitant medication information.

[00173] The end of study for the instant study was defined as the last visit of the last patient.

Patient Selection

10 [00174] 69 patients with a 2:1 randomization to alirocumab and placebo were included in the study.

[00175] Patients who did not meet eligibility criteria during the initial screening were able to rescreen only once. Patients who were rescreened after the screening window ended had to re-consent for study participation and repeat all screening procedures. Patients who did not meet
15 all eligibility criteria during the initial screening and were still within the screening window were able to retest those assessments that did not meet eligibility criteria once.

Study Population

[00176] The study population will consist of male and female patients, ≥ 18 years of age, diagnosed with hoFH, except for patients known to have null/null mutations in both LDLR alleles.

20 *Inclusion Criteria*

[00177] In order to be eligible for the instant study, the patient had to meet the following criteria:

- i) males and females ≥ 18 years of age at the time of the screening visit,
- ii) diagnosis of hoFH by at least 1 of the following genotype or clinical criteria (all patients on LDL apheresis to be diagnosed based on genotype): a) documented
25 homozygous or compound heterozygous mutations in both LDLR alleles (note: patients with known null mutations in both LDLR alleles were excluded (see Exclusion Criteria, below), b) presence of homozygous or compound heterozygous mutations in Apo B, PCSK9 or LDLRAP1, c) presence of double heterozygous mutations, *i.e.*, mutations on different genes in the LDLR, Apo B or PCSK9 alleles,
30 d) untreated TC > 500 mg/dL (12.93 mmol/L) and TG < 300 mg/dL (3.39 mmol/L), and e) both parents with history of TC > 250 mg/dL (6.46 mmol/L) OR cutaneous or tendinous xanthoma before age 10,
- iii) receiving a stable dose of a statin at the screening visit (note: patients who were not able to tolerate a statin or if statins were found to be ineffective could be included in
35 the study, but the reason was to be documented in the case report form (CRF)),

- iv) if undergoing LDL apheresis, must have initiated LDL apheresis at least 3 months prior to screening and must have been on a stable weekly (every 7 days) or every other week (every 14 days) schedule or stable settings for at least 8 weeks,
- v) willing to maintain a stable low fat or heart-healthy diet for the duration of the study,
- 5 vi) willing and able to comply with clinic visits and study related procedures, and
- vii) provided signed informed consent.

[00178] A summary of the instant study's patient characteristics at baseline is provided in the following table:

10 **[00179] Table 1: Demographics and Patient Characteristics at Baseline – Randomized Population**

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
Age (years)				
Number	24	45	69	0.5081
Mean (SD)	45.4 (15.80)	42.3 (14.13)	43.4 (14.69)	
Median	43.0	42.0	42.0	
Min : Max	23 : 81	19 : 69	19 : 81	
Age group (years) [n (%)]				
Number	24	45	69	0.6210
<45	13 (54.2%)	27 (60.0%)	40 (58.0%)	
>=45 to <65	7 (29.2%)	14 (31.1%)	21 (30.4%)	
>=65 to <75	3 (12.5%)	4 (8.9%)	7 (10.1%)	
>=75	1 (4.2%)	0	1 (1.4%)	
Age group (years) [n (%)]				
Number	24	45	69	0.4354
<65	20 (83.3%)	41 (91.1%)	61 (88.4%)	
>=65	4 (16.7%)	4 (8.9%)	8 (11.6%)	
Sex [n (%)]				
Number	24	45	69	0.6183
Male	13 (54.2%)	21 (46.7%)	34 (49.3%)	
Female	11 (45.8%)	24 (53.3%)	35 (50.7%)	

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
Race [n (%)]				
Number	24	45	69	0.4188
White	18 (75.0%)	36 (80.0%)	54 (78.3%)	
Black or African American	0	2 (4.4%)	2 (2.9%)	
Asian	5 (20.8%)	7 (15.6%)	12 (17.4%)	
American Indian or Alaska Native	0	0	0	
Native Hawaiian or Other Pacific Islander	0	0	0	
Other	1 (4.2%)	0	1 (1.4%)	
Ethnicity [n (%)]				
Number	24	45	69	0.5396
Hispanic or Latino	0	2 (4.4%)	2 (2.9%)	
Not Hispanic or Latino	24 (100%)	43 (95.6%)	67 (97.1%)	
Weight (kg)				
Number	24	45	69	0.7913
Mean (SD)	70.1 (17.75)	71.6 (20.23)	71.1 (19.28)	
Median	69.8	71.0	70.0	
Min : Max	46 : 114	42 : 157	42 : 157	
Weight group (kg) [n (%)]				
Number	24	45	69	0.8480
<50	2 (8.3%)	6 (13.3%)	8 (11.6%)	
>=50 to <70	10 (41.7%)	14 (31.1%)	24 (34.8%)	
>=70 to <100	11 (45.8%)	22 (48.9%)	33 (47.8%)	
>=100	1 (4.2%)	3 (6.7%)	4 (5.8%)	
BMI (kg/m ²)				
Number	24	45	69	0.9899
Mean (SD)	25.1 (5.12)	25.1 (5.37)	25.1 (5.24)	
Median	24.8	25.1	24.8	
Min : Max	18 : 40	18 : 47	18 : 47	
BMI group (kg/m ²) [n (%)]				
Number	24	45	69	1.0000
<30	21 (87.5%)	40 (88.9%)	61 (88.4%)	
>=30	3 (12.5%)	5 (11.1%)	8 (11.6%)	
Note: p-values comparing baseline data between treatment groups are provided for descriptive purpose, as a screening tool, using Fisher exact test for qualitative data and the asymptotic one-way ANOVA test for Wilcoxon scores (Kruskal-Wallis test) for continuous data.				

Exclusion Criteria

[00180] A patient meeting any of the following criteria was excluded from the instant study:

- i) documented evidence of a null mutation in both LDLR alleles,
- ii) use of a PCSK9 inhibitor within 10 weeks from screening visit,
- iii) background medical LMT that has not been stable for at least 4 weeks (6 weeks for fibrates, 24 weeks for mipomersen, 12 weeks for maximum tolerated dose of lomitapide) before the screening visit; patients had the option to enter the optional run-in period; once the patient was stable on his/her background medical LMT for the appropriate amount of time, the patient could enter the screening period,
- iv) LDL apheresis schedule/apheresis settings that had not been stable for at least 8 weeks before the screening visit or an apheresis schedule/settings that was not anticipated to be stable over the next 24 weeks; patients had the option to enter the optional run-in period; once the patient was stable on his/her background lipid apheresis schedule/settings for the appropriate amount of time, the patient could enter the screening period,
- v) use of nutraceuticals or over-the-counter (OTC) therapies known to affect lipids, at a dose/amount that had not been stable for at least 4 weeks prior to the screening visit or between the screening and randomization visits; patients had the option to enter the optional run-in period; once the patient was stable on his/her nutraceuticals or OTC therapies for the appropriate amount of time, the patient could enter the screening period,
- vi) presence of any clinically significant uncontrolled endocrine disease known to influence serum lipids or lipoproteins; this could include newly diagnosed (within 3 months prior to randomization visit [week 0/day 1]) diabetes mellitus, or signs and symptoms of hypothyroidism; as a note, patients on thyroid replacement therapy could be included, if the dosage of replacement therapy had been stable for at least 12 weeks prior to screening and the thyroid stimulating hormone (TSH) level was within the normal range at the screening visit,
- vii) unstable weight (variation >5 kg) within 2 months prior to the screening visit (week - 2),
- viii) initiation of a new diet or major change to a previous diet within 4 weeks prior to screening,
- ix) chronic use of systemic corticosteroids, unless on a stable regimen of 10 mg daily prednisone equivalent or less for at least 6 weeks prior to randomization; as a note, topical, intra-articular, nasal, inhaled and ophthalmic steroid therapies were not considered as 'systemic' and were allowed,

- x) use of estrogen or testosterone therapy unless the regimen had been stable in the past 6 weeks prior to the screening visit and there were no plans to change the regimen during the study,
- 5 xi) systolic blood pressure >160 mmHg or diastolic blood pressure >100 mmHg at the screening visit (1 repeat measurement was allowed),
- xii) LDL-C level <70 mg/dL (1.81 mmol/L) at the screening visit,
- xiii) history of a MI, unstable angina leading to hospitalization, coronary artery bypass graft surgery, percutaneous coronary intervention, uncontrolled cardiac arrhythmia, carotid surgery or stenting, stroke, transient ischemic attack, valve replacement surgery, carotid revascularization, endovascular procedure or surgical intervention for peripheral vascular disease within 3 months prior to the screening visit,
- 10 xiv) history of New York Heart Association (NYHA) class IV heart failure within 12 months before screening,
- xv) history of cancer within the past 2 years, except for adequately treated basal cell skin cancer, squamous cell skin cancer, or *in situ* cervical cancer,
- 15 xvi) use of any active investigational drugs within 1 month or 5 half-lives prior to the screening visit, whichever is longer,
- xvii) conditions/situations such as any clinically significant abnormality identified at the time of screening that, in the judgment of the investigator or any sub-investigator, would preclude safe completion of the study or constrain endpoints assessment; e.g., major systemic diseases, patients with short life expectancy, OR considered by the investigator or any sub-investigator as inappropriate for the instant study for any reason, e.g., deemed unable to meet specific protocol requirements, such as scheduled visits, deemed unable to tolerate injections as per the patient or the investigator, investigator or any sub-investigator, pharmacist, study coordinator, other study staff or relative thereof directly involved in the conduct of the protocol, etc., or presence of any other conditions (e.g., geographic or social), either actual or anticipated, that the investigator feels would restrict or limit the patient's participation for the duration of the study,
- 20
- 25
- 30 xviii) laboratory findings during screening period (not including randomization labs): hepatitis B surface antigen and/or Hepatitis C antibody (associated with a positive HCV RNA polymerase chain reaction) at the screening visit, positive serum beta-human chorionic gonadotropin (hCG) or urine pregnancy test in women of childbearing potential, estimated glomerular filtration rate (eGFR) <30 mL/min/1.73 m² (calculated by central lab), alanine aminotransferase (ALT) or aspartate aminotransferase (AST) >3 x upper limit of normal (ULN) (1 repeat lab was allowed),
- 35

or unexplained serum creatine phosphokinase CPK >5 x ULN (1 repeat lab was allowed),

xix) known hypersensitivity to monoclonal antibody therapeutics,

xx) member of the clinical site study team and/or his/her immediate family,

5 xxi) pregnant or breastfeeding women, and

xxii) women of childbearing potential* who were unwilling to practice highly effective contraception prior to the initial dose/start of the first treatment and for the duration of the study; highly effective contraceptive measures included stable use of combined (estrogen and progestogen containing) hormonal contraception (oral, 10 intravaginal, transdermal) or progestogen-only hormonal contraception (oral, injectable, implantable) associated with inhibition of ovulation initiated 2 or more menstrual cycles prior to screening; intrauterine device (IUD); intrauterine hormone releasing system (IUS); bilateral tubal ligation; vasectomized partner; and or sexual abstinence**. *Postmenopausal women had to be amenorrheic for at least 12 15 months in order not to be considered of childbearing potential. Pregnancy testing and contraception were not required for women with documented hysterectomy or oophorectomy. **Sexual abstinence was considered a highly effective method only if defined as refraining from heterosexual intercourse during the entire period of risk associated with the study treatments.

20 *Premature Withdrawal from the Study*

[00181] Patients had the right to withdraw from the study at any time, for any reason, and without repercussion. The investigator and/or sponsor had the right to withdraw a patient from the study if it was no longer in the interest of the patient to continue in the study, or if the patient's continuation in the study placed the scientific outcome of the study at risk (e.g., if a 25 patient did not or could not follow study procedures). An excessive rate of withdrawals would render the study uninterpretable; therefore, unnecessary withdrawal of patients was to be avoided.

[00182] The investigator was to make the best effort to contact any patient (e.g., contacting patient's family or private physician, reviewing available registries or health care database) who 30 failed to return to the site and to determine health status, including vital status at a minimum. Attempts to contact such patients were to be documented in the patient's records (e.g., times and dates of attempted telephone contact, receipt for sending a registered letter).

[00183] Patients prematurely discontinued from the study were not replaced.

Study Treatments

35 [00184] The investigational study drug injections were provided in prefilled pens and were administered SC into the abdomen, thigh, or outer area of the upper arm. The patient or

caregiver used placebo for injection training at the clinical site. After study eligibility was confirmed, the patient or caregiver was trained to self-inject/inject using placebo.

Investigational and Reference Treatments

[00185] Double-blind-Treatment: study drug administration during the double-blind treatment period started on the day of randomization and was administered immediately after completion of the LDL apheresis procedure (if applicable). For those patients not undergoing LDL apheresis, administration of the investigational study drug was made after all samples for clinical laboratory evaluation had been obtained. The last injection of double-blind study drug occurred on day 71/week 10. If a dose was missed, the patient was instructed to administer the injection within 7 days from the missed dose. If the missed dose was not administered within 7 days, the patient was instructed to skip the dose and resume the original schedule.

[00186] Patients were randomized in a 2:1 ratio to receive: alirocumab 150 mg SC Q2W OR matching placebo SC Q2W. Sterile alirocumab drug product was supplied at a concentration of 150 mg/mL in a prefilled pen. Placebo was also supplied in a prefilled pen.

[00187] Open-label Treatment: to provide further safety data in this rare patient population, all patients received open-label investigational study drug (alirocumab 150 mg SC Q2W), starting at week 12 and continuing through week 24 (end-of open-label treatment period/EOS visit, last injection at week 22), regardless of treatment assignment in the double-blind treatment period. Patients who were receiving LMT or who were undergoing apheresis were to continue a stable dose and regimen and a stable apheresis schedule and settings (as applicable) throughout the duration of the open-label treatment period. Sterile alirocumab drug product was supplied at a concentration of 150 mg/mL in a prefilled pen.

Run-in (optional) and Background Treatment(s)

[00188] Apheresis therapy: patients who were undergoing apheresis therapy without a stable weekly or every other week schedule or stable settings for at least 8 weeks before the screening visit entered a 4-week optional run-in period before the screening period. After the 4-week run-in period, patients whose lipid apheresis schedule/settings remain stable were eligible to enter the 2-week screening period. Additionally, all patients on LDL apheresis had to be diagnosed based on genotype and, if genotype information had not been determined previously, they could enter the run-in to allow time, if needed, to determine their mutation status.

[00189] Lipid modifying therapy: patients who were on background LMT that had not been stable for at least 4 weeks before the screening visit entered a 4-week run-in period to stabilize their LMT. Patients who had not been on a stable dose of mipomersen within 6 months prior to screening or a maximum tolerated dose of lomitapide for 12 weeks prior to screening were excluded.

Dose Modification and Study Treatment Discontinuation

[00190] Dose modification for an individual patient was not allowed.

[00191] Study drug was to be continued whenever possible. In the event the investigational study drug dosing was stopped, it was to be determined if the stop could be made temporarily; permanent discontinuation was to be a last resort. In any case, the patient should remain in the study as long as possible.

[00192] Patients who permanently discontinued study drug during the double-blind treatment period were to remain in the study and undergo all double-blind study visits and procedures with the exception of dosing with study drug. At the time of study drug discontinuation, the patient was to have, as soon as possible, an unscheduled visit with assessments normally planned at end of double-blind treatment visit (within 5 days of discontinuation of study drug, if possible). Then, patients were to resume the original study schedule until the end of the double-blind treatment period and all efforts were to be made to perform the week 12 assessments at week 12. The original study schedule continued until the end of the study visit (*i.e.*, follow up phone call visit).

[00193] Patients who permanently discontinued study drug during the open-label period were to have, as soon as possible, an unscheduled visit with assessments normally planned at the end of the open label treatment period (within 5 days of discontinuation of the study drug, if possible). Upon completion of this visit, the original study schedule resumed until end of study (*i.e.*, follow-up phone visit).

[00194] Patients permanently discontinued study drug for the following reasons: for female patients, individuals that had become pregnant, were actively trying to become pregnant, or discontinued use of protocol-defined methods of effective birth control, acute injection reaction of clinical concern, at patient request, if, in the investigator's opinion, continuation of the investigational study drug dosing would be detrimental to the patient's well-being, intercurrent condition that required discontinuation of the investigational study drug, at the specific request of the sponsor, and/or patient received double-blind treatment before randomization.

[00195] Temporary discontinuation of the investigational study drug was considered by the investigator because of suspected AEs, including allergic events related to the dose of the investigational study drug. Reinitiating the investigational study drug dosing was done under close and appropriate clinical and/or laboratory monitoring. Temporary discontinuation of the investigational study drug is defined as 1 or more scheduled injections that were not administered to the patient as decided by the investigator.

Management of Acute Reactions

[00196] Acute systemic reactions following injection of the investigational study drug (subcutaneous [SC]) were to be treated using clinical judgment to determine the appropriate response according to typical clinical practice.

Method of Treatment Assignment

[00197] The randomized list of treatment kit numbers was generated centrally. An interactive voice response system (IVRS) and/or interactive web response system (IWRS) was used in the instant study. The investigational study drug was packaged in accordance with this list.

- 5 [00198] Patients were randomly assigned to receive alirocumab 150 mg or matching placebo in a 2:1 ratio, stratified by LDL apheresis treatment status (on vs off treatment).

[00199] The treatment kit numbers were allocated using the centralized treatment allocation system at the randomization visit, at weeks specified in Table 1, below, as re-supply visits, and at unscheduled visits, if needed.

10 *Blinding*

[00200] Study patients, the investigators, and study site personnel remained blinded to all randomization assignments throughout the instant study. The study director, medical monitor, study monitor, and any personnel in regular contact with the study site remained blinded to all patient randomization assignments.

- 15 [00201] Lipid results from blood samples collected after the randomization visit were not communicated to the sites, and the sponsor's operational team did not have access to these laboratory results until after the completion of the double-blind treatment period and the first step analysis.

- 20 [00202] Blinded investigational study drug kits coded with a medication numbering system were used. In order to maintain the blind, lists linking these codes with product lot numbers were not accessible to individuals involved in study conduct.

[00203] Anti-drug antibody (ADA) was not communicated to the sites, and the sponsor's operational team did not have access to results associated with patient identification until after the database lock after completion of the double-blind treatment period.

- 25 [00204] While the study was ongoing, it was anticipated that unblinded data, after the first step analysis (conducted as soon as all patients were randomized and all data through week 12 (double-blind period) were collected and validated; this consisted of the final analysis of the double-blind primary and secondary efficacy endpoints), would be submitted to health authorities. Sponsor representatives who conducted and reviewed such data analyses for
30 submission to the health authorities were not part of the study operational team from that point forward, and patient level results were not provided to the study sites.

Emergency Unblinding

- [00205] Unblinding of treatment assignment for a patient could become necessary due to a medical emergency or any other significant medical event (e.g., pregnancy). If unblinding was
35 required, only the investigator made the decision to unblind the treatment assignment, and only the affected patient was unblinded.

[00206] Treatment assignment was not provided to site personnel at any time during the conduct of the study, except in the case of a true emergency. In the event that there was no study pharmacist, the individual at the site fulfilling that role was the only unblinded member of the site personnel.

5 *Treatment Logistics and Accountability*

[00207] As to packaging, labeling, and storage, a medication numbering system was used to label blinded investigational study drug. Lists linking medication numbers with product lot numbers were maintained by the groups (or companies) responsible for the investigational study drug packaging. In order to maintain the blind, these lists were not accessible to individuals involved in study conduct. Training kits containing 1 placebo prefilled pen were provided to the sites for patient/caregiver injection training that was performed before randomization during the screening period or at the baseline visit. A second placebo prefilled pen could be used before randomization if the patient/caregiver required additional injection training. Study drug was refrigerated at the site at a temperature of 2°C to 8°C. Storage temperature was logged.

15 Detailed storage instructions were provided in the study manual.

[00208] As to supply and disposition of treatments, study drug was shipped at a temperature of 2°C to 8°C to the investigator or designee at regular intervals or as needed during the study. At specified time points during the study (e.g., interim site monitoring visits), at the site close-out visit, and following drug reconciliation and documentation by the site monitor, all opened and unopened investigational study drug were to be returned to the sponsor or designee. The investigational study drug was dispensed to each patient. The investigational study drug was stored, prepared, and administered by the patient/caregiver according to instructions provided to each patient/caregiver.

[00209] As to treatment accountability, all drug accountability records were kept current. The investigator had to be able to account for all opened and unopened investigational study drug. These records were to contain the dates, quantity, and study medication dispensed to each patient, returned from each patient (if applicable), and disposed of at the site or returned to the sponsor or designee.

[00210] Patients completed a dosing log to document compliance with the investigational study drug administration. Measures taken to ensure and document the investigational study drug accountability and compliance were:

- i) the investigator or designee obtained via IVRS/IWRS the treatment kit number(s) and dispensed the treatment kit(s) to the patient,
- ii) accountability was verified during the investigational study drug kit re-supply visits only; the used and unused kit(s) were to be brought to these visits for accountability purposes,

- 5 iii) all kits, including used and unused kits, were to be returned by the patient at the designated visit; an unused kit contained all of the unused prefilled pens; a used kit was one from which the patient had removed 1 or more prefilled pens; a used prefilled pen was one that had been removed from the kit with the intention of administration, including those injections that had been partially or fully injected; the patient was to discard all used prefilled pens into the sharps container and never put used prefilled pens back into the used kit,
- iv) all sharps containers were to be returned to the site by the patient,
- v) the investigator/study coordinator entered data in the appropriate CRF pages, according to data recorded in the treatment log form, and
- 10 vi) the monitor checked the data consistency among CRF pages, treatment log form, and returned unused prefilled pens of a corresponding kit.

15 **[00211]** All treatments kits were retrieved by the sponsor. A detailed treatment log of the returned investigational study drug was established with the investigator or designee and countersigned by the investigator and the monitoring team.

20 **[00212]** As to treatment compliance, all drug compliance records were to be kept current and made available for inspection by the sponsor and regulatory agency inspectors. Patients completed a dosing log to document compliance with the investigational study drug administration.

Concomitant Medications

25 **[00213]** Concomitant medications were to be kept to a minimum during the study. If considered necessary for the patient's welfare and unlikely to interfere with the investigational study drug, concomitant medications (other than those that were prohibited during the study) could be given at the discretion of the investigator, at a stable dose when possible. Any treatments administered from the time of informed consent/assent to the final study visit were considered concomitant medications. This included medications that were started before the study and were ongoing during the study.

30 **[00214]** Prohibited medications and procedures included: a) use of a PCSK9 inhibitor within 10 weeks from the screening visit, b) initiation of or changes to the LDL apheresis schedule and/or settings (if applicable) or background medical LMT from the initial screening visit until the end of study visit, c) use of continuous estrogen or testosterone hormone replacement therapy, unless the regimen had been stable in the past 6 weeks prior to the screening visit, and d) chronic use of systemic corticosteroids, unless on a stable regimen of 10 mg daily prednisone equivalent or less for at least 6 weeks prior to randomization. As a note, topical, intra articular, nasal, inhaled and ophthalmic steroid therapies were not considered as 'systemic' and were allowed.

35

[00215] Permitted medications and procedures included lipid modifying therapies, nutraceuticals, and over-the-counter therapies that may affect lipids, but only if they had been used at a stable dose and regimen for at least 4 weeks (6 months for mipomersen, 12 weeks for the maximum tolerated dose of lomitapide) before the screening visit. The dose and regimen had to remain stable until the end of study visit. Low-density lipoprotein apheresis was allowed only if the schedule/settings had been stable for at least 8 weeks before the screening visit and remained stable until the end of study visit.

[00216] The lipid-modifying therapy (LMT) history of the patients in the instant study are summarized in the following table:

10 **Table 2: LMT History – Randomized Population**

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)
Overall on a maximally tolerated LMT per investigator [n(%)]			
Yes	22 (91.7%)	44 (97.8%)	66 (95.7%)
No	2 (8.3%)	1 (2.2%)	3 (4.3%)
On statin at screening [n(%)]			
Yes	23 (95.8%)	44 (97.8%)	67 (97.1%)
No	1 (4.2%)	1 (2.2%)	2 (2.9%)
On High intensity HMG COA inhibitor (statin)			
Yes	23 (95.8%)	44 (97.8%)	67 (97.1%)
Patient on the maximum tolerated dose	23 (95.8%)	42 (93.3%)	65 (94.2%)
No	1 (4.2%)	1 (2.2%)	2 (2.9%)
Reason			
Intolerance - experienced muscle symptoms	0	1 (2.2%)	1 (1.4%)
Intolerance - other symptoms	0	0	0
Concern about drug interaction	0	0	0
Concern for potential adverse effects due to use (such as cognitive impairment)	0	0	0
Lack of efficacy	0	0	0
Not an approved medication	0	0	0
Regional practices/local prescribing information	0	0	0
Lack of access	0	0	0
Other	1 (4.2%)	0	1 (1.4%)
History of down titration of statin dose due to tolerability			
Yes	4 (16.7%)	12 (26.7%)	16 (23.2%)
No	20 (83.3%)	33 (73.3%)	53 (76.8%)

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)
History of change to different statin due to tolerability			
Yes	4 (16.7%)	9 (20.0%)	13 (18.8%)
No	20 (83.3%)	36 (80.0%)	56 (81.2%)
LLT at screening ^a [n (%)]			
High Intensity HMG COA inhibitor (Statin)	23 (95.8%)	44 (97.8%)	67 (97.1%)
Fibrates	1 (4.2%)	0	1 (1.4%)
Bile acid sequestrant	2 (8.3%)	3 (6.7%)	5 (7.2%)
Cholesterol absorption inhibitor	3 (12.5%)	6 (13.3%)	9 (13.0%)
Nicotinic acid and derivatives (Niacin)	1 (4.2%)	1 (2.2%)	2 (2.9%)
Omega 3 fatty acids (> 1000 mg/day)	0	6 (13.3%)	6 (8.7%)
Ezetimibe	19 (79.2%)	31 (68.9%)	50 (72.5%)
PCSK9 inhibitor	0	0	0
Probucol [1]	0	0	0
Apheresis	4 (16.7%)	6 (13.3%)	10 (14.5%)
Lomitapide	3 (12.5%)	7 (15.6%)	10 (14.5%)
Mipomersen	0	0	0
^a patient can be counted in several categories.			

[00217] The lipid efficacy parameters at baseline of the patients in the instant study are summarized in the following table (quantitative summary in conventional units – randomized population):

5

Table 3

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
LDL-C (mg/dL)				
Number	24	45	69	0.2338
Mean (SD)	259.6 (175.75)	295.0 (154.59)	282.7 (161.86)	
Median	232.0	248.0	240.0	
Q1 : Q3	152.5 : 296.5	186.0 : 366.0	183.0 : 346.0	
Min : Max	86 : 896	78 : 767	78 : 896	
Non-HDL-C (mg/dL)				
Number	24	45	69	0.2313
Mean (SD)	282.0 (177.41)	320.5 (160.36)	307.1 (166.22)	
Median	259.0	275.0	275.0	
Q1 : Q3	167.0 : 319.0	203.0 : 416.0	195.0 : 364.0	
Min : Max	102 : 928	92 : 791	92 : 928	

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
Total-C (mg/dL)				
Number	24	45	69	0.1966
Mean (SD)	325.1 (171.57)	364.3 (157.30)	350.7 (162.24)	
Median	305.5	326.0	315.0	
Q1 : Q3	209.0 : 362.0	253.0 : 444.0	235.0 : 400.0	
Min : Max	155 : 947	162 : 830	155 : 947	
HDL-C (mg/dL)				
Number	24	45	69	0.7526
Mean (SD)	43.2 (11.96)	43.8 (14.78)	43.6 (13.78)	
Median	46.0	40.0	40.0	
Q1 : Q3	35.0 : 53.5	32.0 : 50.0	32.0 : 53.0	
Min : Max	19 : 60	24 : 95	19 : 95	
Fasting TGs (mg/dL)				
Number	24	45	69	0.1403
Mean (SD)	111.7 (77.97)	128.0 (74.34)	122.3 (75.45)	
Median	80.5	110.0	97.0	
Q1 : Q3	61.0 : 128.5	79.0 : 160.0	72.0 : 148.0	
Min : Max	41 : 305	25 : 346	25 : 346	
Lp(a) (mg/dL)				
Number	24	45	69	0.7719
Mean (SD)	40.0 (36.41)	42.9 (36.34)	41.9 (36.12)	
Median	32.5	36.0	36.0	
Q1 : Q3	12.0 : 52.5	10.0 : 68.0	10.0 : 65.0	
Min : Max	4 : 144	4 : 157	4 : 157	
Apo-B (mg/dL)				
Number	24	45	69	0.2675
Mean (SD)	175.0 (95.12)	193.3 (87.59)	186.9 (90.01)	
Median	161.0	178.0	165.0	
Q1 : Q3	110.0 : 202.0	132.0 : 232.0	124.0 : 220.0	
Min : Max	82 : 525	66 : 468	66 : 525	
Apo-A1 (mg/dL)				
Number	24	45	69	0.7913
Mean (SD)	124.8 (24.59)	125.6 (28.57)	125.3 (27.07)	
Median	124.5	117.0	118.0	
Q1 : Q3	108.0 : 144.0	106.0 : 143.0	108.0 : 144.0	
Min : Max	67 : 164	80 : 206	67 : 206	
Apo-B/Apo-A1 (ratio)				
Number	24	45	69	0.2077
Mean (SD)	1.590 (1.4746)	1.635 (0.8693)	1.619 (1.1067)	
Median	1.235	1.400	1.380	
Q1 : Q3	0.860 : 1.630	1.020 : 1.990	0.970 : 1.930	
Min : Max	0.54 : 7.84	0.38 : 4.46	0.38 : 7.84	
Note: p-values comparing baseline data between treatment groups are provided for descriptive purpose, as a screening tool, using the asymptotic one-way ANOVA test for Wilcoxon scores (Kruskal-Wallis test).				

[00218] The lipid efficacy parameters at baseline of the patients in the instant study are summarized in the following table (qualitative summary– randomized population):

Table 4

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
LDL-C				
Number	24	45	69	0.2678
<70 mg/dL / <1.81 mmol/L	0	0	0	
>=70 to <100 mg/dL / >=1.81 to <2.59 mmol/L	4 (16.7%)	1 (2.2%)	5 (7.2%)	
>=100 to <130 mg/dL / >=2.59 to <3.37 mmol/L	1 (4.2%)	4 (8.9%)	5 (7.2%)	
>=130 to <160 mg/dL / >=3.37 to <4.14 mmol/L	1 (4.2%)	2 (4.4%)	3 (4.3%)	
>=160 to <190 mg/dL / >=4.14 to <4.91 mmol/L	3 (12.5%)	5 (11.1%)	8 (11.6%)	
>=190 mg/dL / >=4.91 mmol/L	15 (62.5%)	33 (73.3%)	48 (69.6%)	
HDL-C				
Number	24	45	69	0.6192
<40 mg/dL / <1.04 mmol/L	10 (41.7%)	22 (48.9%)	32 (46.4%)	
>=40 mg/dL / >=1.04 mmol/L	14 (58.3%)	23 (51.1%)	37 (53.6%)	
Fasting TGs				
Number	24	45	69	0.7709
<150 mg/dL / <1.7 mmol/L	19 (79.2%)	33 (73.3%)	52 (75.4%)	
>=150 mg/dL / >=1.7 mmol/L	5 (20.8%)	12 (26.7%)	17 (24.6%)	
>=150 to <200 mg/dL / >=1.7 to <2.3 mmol/L	2 (8.3%)	5 (11.1%)	7 (10.1%)	
>=200 mg/dL / >=2.3 mmol/L	3 (12.5%)	7 (15.6%)	10 (14.5%)	
Lp(a)				
Number	24	45	69	0.8006
<30 mg/dL / <0.3 g/L	12 (50.0%)	20 (44.4%)	32 (46.4%)	
>=30 mg/dL / >=0.3 g/L	12 (50.0%)	25 (55.6%)	37 (53.6%)	
>=30 to <50 mg/dL / >=0.3 to <0.5 g/L	5 (20.8%)	10 (22.2%)	15 (21.7%)	
>=50 mg/dL / >=0.5 g/L	7 (29.2%)	15 (33.3%)	22 (31.9%)	
Note: p-values comparing baseline data between treatment groups are provided for descriptive purpose, as a screening tool, using Fisher exact test.				

5 **[00219]** The summary of the patients in the instant study by mutation status are summarized in the following table (randomized population):

Table 5

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
Genotype State				
Homozygous	11 (45.8%)	19 (42.2%)	30 (43.5%)	0.6160
Compound Heterozygous	7 (29.2%)	11 (24.4%)	18 (26.1%)	
Double Heterozygous	0	4 (8.9%)	4 (5.8%)	
Other (Heterozygous, undetermined, or no mutation)	6 (25.0%)	11 (24.4%)	17 (24.6%)	
Homozygous (LDLR)				
Defective/Defective	10 (41.7%)	18 (40.0%)	28 (40.6%)	
Negative/Negative	0	2 (4.4%)	2 (2.9%)	
Homozygous (LDLRAP1)				
Negative/Negative	0	1 (2.2%)	1 (1.4%)	

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
Homozygous (PCSK9)	1 (4.2%)	0	1 (1.4%)	
Defective/Defective	1 (4.2%)	0	1 (1.4%)	
Compound Heterozygous (LDLR)	7 (29.2%)	11 (24.4%)	18 (26.1%)	
Defective/Defective	4 (16.7%)	7 (15.6%)	11 (15.9%)	
Defective/Negative	3 (12.5%)	4 (8.9%)	7 (10.1%)	
Negative/Negative	0	0	0	
Double Heterozygous (LDLR and APOB and PCSK9)	0	4 (8.9%)	4 (5.8%)	
Defective (LDLR)/Negative (APOB)	0	1 (2.2%)	1 (1.4%)	
Defective (LDLR)/Defective (APOB)	0	1 (2.2%)	1 (1.4%)	
Defective (LDLR)/Defective (PCSK9)	0	1 (2.2%)	1 (1.4%)	
Negative (LDLR)/Defective (PCSK9)	0	1 (2.2%)	1 (1.4%)	
Note: p-values comparing baseline data between treatment groups are provided for descriptive purpose, as a screening tool, using Fisher exact test.				

[00220] The summary of the patients in the instant study by mutation status are summarized in the following table (Null/Null vs Not Null/Null): LDLR Activity < 2% – Randomized Population:

5

Table 6

	Placebo (N=24)	Alirocumab 150 Q2W (N=45)	All (N=69)	P-Value vs. Placebo
Genotype State				
Homozygous (LDLR)				
Null/Null	1 (4.2%)	4 (8.9%)	5 (7.2%)	0.6264
Not Null/Null	9 (37.5%)	14 (31.1%)	23 (33.3%)	
Homozygous (LDLRAP1)				
Null/Null	0	0	0	
Compound Heterozygous (LDLR)				
Not Null/Null	7 (29.2%)	11 (24.4%)	18 (26.1%)	
Null/Null is defined as LDLR activity < 2%				
Note: p-values comparing baseline data between treatment groups are provided for descriptive purpose, as a screening tool, using Fisher exact test.				

Study Schedule of Events and Procedures

[00221] The study assessments and procedures are presented by study period in Table 7, below.

10

Table 7

	Optional Run-in	Screening Period	Double-Blind Treatment Period					Open-label Treatment Period		Follow-up ¹⁰
Study Procedure	Run-in Visit 1a	Screening Visit 1	Baseline Visit 2	Visit 3	Visit 4	Visit 5	End of Double-Blind Treatment Visit 6	Visit 7	End of Open-Label Treatment Visit 8	End of Study Phone Visit 9
Day	-42 to -14	-14 to -1	1(±1)	29(±5/±1 ⁹)	57(±5/±1 ⁹)	71(±5/±1 ⁹)	85(±3/±1 ⁹)	127(±7)	169(±3/±1 ⁹)	225(±5)
Week	-6 to -2	-2 to -1	0	4	8	10	12	18	24	32
Screening/Baseline:										
Inclusion/Exclusion	X	X								
Informed Consent/Assent ¹	X	X								
Pharmacogenomics Consent/Assent		X								
Medical/Surgical History, Alcohol habits, Smoking habits		X								
Demo-graphics		X								
Treatment:										
Injection training ²		X	X							
Investigational study drug kit dispensation			X	X			X	X		
Administer SC double-blind investigational study drug ³			X	X	X	X				
Administer open-label investigational study drug ³							X	X		
Kit return				X	X			X	X	
Review of dosing log				X	X			X	X	
Concomitant medications	X	X	X	X	X	X	X	X	X	X

	Optional Run-in	Screening Period	Double-Blind Treatment Period					Open-label Treatment Period		Follow-up ¹⁰
Study Procedure	Run-in Visit 1a	Screening Visit 1	Base-line Visit 2	Visit 3	Visit 4	Visit 5	End of Double-Blind Treatment Visit 6	Visit 7	End of Open-Label Treatment Visit 8	End of Study Phone Visit 9
Efficacy:										
Lipid panel ⁴		X	X	X	X		X	X	X	
Specialty lipid panel ⁴			X	X	X		X		X	
EQ-5D			X				X		X	
Safety										
Adverse events	X	X	X	X	X	X	X	X	X	X
Physical examination		X					X		X	
Body weight	X	X					X		X	
Vital signs	X	X	X	X	X	X	X	X	X	
Electrocardiogram ⁵		X					X		X	
Laboratory Testing:										
Hematology ⁶		X	X		X		X		X	
Blood chemistry ⁶		X	X		X		X		X	
Creatine Phosphokinase ⁶		X	X		X		X		X	
Hepatitis B surface antigen ⁶		X								
Hepatitis C antibody ⁶		X					X		X	
Serum pregnancy test		X								
Urine pregnancy test (done locally)	X						X		X	
Urinalysis		X	X		X		X		X	
TSH		X								
hs-CRP ⁶		X	X				X		X	
Research samples ⁶			X	X	X		X		X	
Mandatory DNA collection for hoFH Genetic Testing ⁷	X	X								

	Optional Run-in	Screening Period	Double-Blind Treatment Period					Open-label Treatment Period		Follow-up ¹⁰
Study Procedure	Run-in Visit 1a	Screening Visit 1	Base-line Visit 2	Visit 3	Visit 4	Visit 5	End of Double-Blind Treatment Visit 6	Visit 7	End of Open-Label Treatment Visit 8	End of Study Phone Visit 9
PK/Drug Concentration and ADA Samples:										
PK Sample ⁶			X	X	X	X	X		X	
ADA Sample ⁶			X				X		X	
Optional Genomic DNA Sample ⁸			X							

- [00222]** The footnotes from Table 6, above, signify the following: 1. Informed consent/assent was obtained either at Visit 1a (for patients who needed the optional run-in time) or at Visit 1 (for the patients who did not need optional run-in). 2. Injection training was performed with the patient and/or caregiver during the screening period or at baseline using placebo. 3. After the investigational study drug administration (double-blind and open-label), patients needed to be monitored for 30 minutes. 4. Lipid panel consisted of: total-C, LDL-C, HDL-C, TG, non-HDL-C. Specialty lipid panel consisted of: ApoB, Apo A-1, ratio Apo B/Apo A-1, and Lp(a). Lipid panels were to be collected after an approximately 8 hour fast. 5. ECG was to be performed before blood samples were collected. 6. On days when a clinic visit coincided with a dosing day, all blood samples (including ADA samples) were collected immediately prior to LDL apheresis (if applicable) and before the investigational study drug administration, but after study assessments were performed. PK samples were also used for free and total PCSK9 analysis. 7. Sample was to be obtained prior to randomization and was used to determine hoFH mutation status. 8. Patients on apheresis could collect this during visit 1a. 9. Optional DNA sample was to be collected on day 1; however, they could be collected at any visit during the course of the study. Genomic informed consent form (ICF) had to be signed prior to performing this assessment. 10. Visit window was ± 3 days for patients not on apheresis and +1 day for patients on apheresis. Every attempt was to be made to ensure all samples were collected immediately prior to LDL apheresis. The timing between the baseline sample collection relative to the most recently completed LDL apheresis procedure was to match the timing of the week 12 sample collection relative to the most recently completed LDL apheresis procedure. Depending on the duration between the LDL apheresis procedure and sample collection, the visit window might not apply. 10. This visit was only for patients who did not participate in another lipid-lowering study.

[00223] With respect to early termination visits, if for any reason the patient refused to continue the study, the patient was to undergo an unscheduled visit as soon as possible with assessments normally planned at the end of double blind treatment visit if the patient was in the double-blind treatment period (week 12 visit assessments); if the patient was in the open-label treatment visit, then the end of the open-label treatment assessments were to be used (week 24 visit assessments). This visit was to take place within 5 days of treatment discontinuation, if possible. The patient was to be followed for at least 70 days from the last dose of study drug or up to recovery or stabilization of any AE to be followed-up as specified, whichever came last.

[00224] As to unscheduled visits, all attempts were to be made to keep the patients on the study schedule. Unscheduled visits could be necessary to repeat testing following abnormal laboratory results, for follow-up of AEs, or for any other reason, as warranted.

Study Procedures

[00225] The following procedures were performed for the sole purpose of determining study eligibility or characterizing the baseline population: medical/surgical history, medication history, demographics, hepatitis B surface antigen, and serum pregnancy testing.

[00226] On day 1, after completion of study assessments, collection of blood samples, LDL apheresis (if applicable), the first dose of double-blind investigational study drug was administered. The patient was monitored at the clinical site for 30 minutes after the first dose. Subsequent doses of the investigational study drug were to be administered subcutaneously Q2W. Doses of the investigational study drug were to be administered at approximately the same time of day (based upon patient preference) throughout the study. After day 1, it was acceptable for dosing to fall within a window of ± 5 days, with the exception of the week 10 and week 22 doses, which had a ± 3 day dosing window for patients not on apheresis and a +1 day dosing window for patients on apheresis, since there were key efficacy assessments at week 12 and week 24.

[00227] In the event an injection was delayed by more than 7 days or completely missed, the patient was to return to the original schedule of the investigational study drug dosing without administering additional injections. If the delay was less than or equal to 7 days from the missed date, the patient was to administer the delayed injection and then resume the original dosing schedule. Site personnel provided the patient/caregiver with detailed instructions for transport, storage, preparation, and administration of the investigational study drug.

Efficacy Procedures

[00228] Total cholesterol, HDL-C, TG, Apo B, Apo A-1, and Lp(a) were directly measured by the central laboratory. Low-density lipoprotein cholesterol was calculated using the Friedewald formula. If TG values exceeded 400 mg/dL (4.52 mmol/L), or if calculated LDL-C values were below 15 mg/dL, then the central lab reflexively measured LDL-C using the beta quantification

method. Non-HDL-C was calculated by subtracting HDL-C from the Total C. The Apo B/Apo A-1 ratio was calculated.

[00229] Blood samples for the lipid panel and specialty lipid panel were collected at time points according to Table 7, above. What was included in the lipid panel and special lipid panel is described below.

Quality of Life Procedures

[00230] EuroQol-5 Questionnaire: the EQ-5D is a standardized measure of health status developed by the EuroQol Group in order to provide a simple, generic measure of health for clinical and economic appraisal. The EQ-5D as a measure of health related quality of life, defines health in terms of 5 dimensions: mobility, self-care, usual activities, pain/discomfort, anxiety/depression. Each dimension has 3 ordinal levels of severity: "no problem" (1), "some problems" (2), "severe problems" (3). Overall health state is defined as a 5-digit number. Health states defined by the 5-dimensional classification can be converted into corresponding index scores that quantify health status, where 0 represents "death" and 1 represents "perfect health."

Safety Procedures

[00231] Vital signs, including blood pressure and heart rate were collected at time points according to Table 7, above. A thorough and complete physical examination was performed at time points according to Table 7, above. Body weight was collected at time points according to Table 7, above. Care was to be taken to examine and assess any abnormalities that could be present, as indicated by the patient's medical history. Electrocardiogram was to be performed before blood was drawn during the visits requiring blood draws. A standard 12-lead ECG was performed with the patient in the supine position after resting quietly for 10 minutes at time points according to Table 7, above. Heart rate was recorded from the ventricular rate, and the PR, QRS, RR, and QT intervals were recorded. The ECG strips or reports were retained with the source.

[00232] All laboratory samples (including ADA samples) were collected after assessments were performed and before a dose of the investigational study drug was administered at visits that corresponded with a dosing day. Alcohol consumption within 48 hours or intense physical exercise within 24 hours preceding blood sampling was discouraged. Samples for laboratory testing were collected at time points according to Table 6, above, and analyzed by a central laboratory during the study. Lipid panel samples were collected after 8-hour fasting. Detailed instructions for blood sample collection were in the laboratory manual provided to study sites. Tests included:

Lipid Panel and Specialty Lipid Panel

Total cholesterol	Apo B
Triglyceride	Apo A-1
Calculated LDL-C	Apo B/Apo A-1 ratio
HDL-C	Lp(a)
Non-HDL-C	

Blood Chemistry

Sodium	Total protein, serum	Total bilirubin
Potassium	Creatinine	Uric acid
Chloride	Blood urea nitrogen (BUN)	Creatine phosphokinase (CPK)
Carbon dioxide	Aspartate aminotransferase (AST)	
Calcium	Alanine aminotransferase (ALT)	
Glucose	Alkaline phosphatase	
Albumin	Lactate dehydrogenase (LDH)	

Hematology

Hemoglobin	Differential:
Hematocrit	Neutrophils
Red blood cells (RBCs)	Lymphocytes
White blood cells (WBCs)	Monocytes
Red cell indices	Basophils
Platelet count	Eosinophils

5 ***Urinalysis***

Color	Glucose	RBC
Clarity	Blood	Hyaline and other casts
pH	Bilirubin	Bacteria
Specific gravity	Leukocyte esterase	Epithelial cells
Ketones	Nitrite	Crystals
Protein	WBC	Yeast

[00233] Other laboratory tests were performed as follows: pregnancy testing (serum and urine) was performed at time points according to Table 7, above, pregnancy testing (urine) was assessed locally at time points according to Table 7, above, samples for the liver panel (ALT, AST, alkaline phosphatase, and total bilirubin), high sensitivity C-reactive protein (hs-CRP) were collected at time points according to Table 7, above, and a sample for hepatitis B surface antigen, hepatitis C antibody, and TSH was collected at screening. Samples for hepatitis C antibody were collected at time points according to Table 6, above.

Abnormal Laboratory Values and Laboratory Adverse Events

[00234] All laboratory values had to be reviewed by the investigator or authorized designee. Significantly abnormal test results that occurred after start of treatment had to be repeated to confirm the nature and degree of the abnormality. When necessary, appropriate ancillary investigations was to be initiated. If the abnormality failed to resolve or could not be explained by events or conditions unrelated to the study medication or its administration, the medical monitor was consulted. The clinical significance of an abnormal test value, within the context of the disease under study, was to be determined by the investigator.

[00235] The criteria for determining whether an abnormal objective test finding should be reported as an AE included: the test result we associated with accompanying symptoms, and/or the test result required additional diagnostic testing or medical/surgical intervention, and/or the test result led to a change in dosing (outside of protocol-stipulated dose adjustments), discontinuation from the study, significant additional concomitant drug treatment, or other therapy.

Pharmacokinetic and Anti-Drug Antibody Procedures

[00236] Samples for drug concentration were collected at time points listed in Table 7, above. Any unused samples could be used for exploratory biomarker research.

[00237] Samples for anti-drug antibody (ADA) assessment were collected at time points listed in Table 6, above. At visits that took place on dosing days, all samples for ADA assessments were collected before a dose of the investigational study drug was administered. To maintain the blind of the study, ADA samples were collected from all patients, including those who received only placebo. Any unused samples could be used for exploratory biomarker research.

hoFH Genetic Testing

[00238] A sample was collected for mandatory hoFH genetic testing to characterize the mutation status of each patient as listed in Table 7, above.

Research Samples

[00239] Samples for exploratory research were collected as allowed by local regulations to study PCSK9 levels, PCSK9 function, effects of PCSK9 inhibition with a monoclonal antibody, and mechanisms of hyperlipidemia and heart disease. Research sampling was collected at time points according to Table 7, above. Research samples were coded to maintain patient confidentiality.

Biomarker Procedures

[00240] Biomarker samples were collected at time points according to Table 7, above, as part of the Research Samples. Biomarker measurements were performed in matrix, for example, serum samples to determine effects on biomarkers of indication or relevant physiological and pathogenic processes. The biomarkers studied were ones believed to be relevant to the

pathophysiology of indication target engagement, mechanism of action, and possible toxicities. Biomarkers studied could include, but were not limited to, PCSK9.

Adverse Events

[00241] An adverse event (AE) is any untoward medical occurrence in a patient administered an investigational study drug, which may or may not have a causal relationship with the investigational study drug. Therefore, an AE is any unfavorable and unintended sign (including abnormal laboratory finding), symptom, or disease temporally associated with the use of the investigational study drug, whether or not considered related to the investigational study drug. An AE also includes any worsening (*i.e.*, any clinically significant change in frequency and/or intensity) of a pre-existing condition that is temporally associated with the use of the investigational study drug.

[00242] An adverse event of special interest (serious or non-serious) is one of scientific and medical concern specific to the sponsor's product or program, for which ongoing monitoring and rapid communication by the investigator to the sponsor can be appropriate. Adverse events of special interest for this study include the following: a) increase in ALT: ALT $\geq 3 \times$ ULN (if baseline ALT < ULN), or ALT ≥ 2 times the baseline value (if baseline ALT $\geq$ ULN), b) allergic events and/or local injection site reactions that require consultation with another physician for further evaluation, c) pregnancy, d) symptomatic overdose with investigational medicinal product, e) neurologic events that require additional examinations/procedures and/or referral to a specialist, neurocognitive events, f) cataracts, g) new onset of diabetes (where the definition of new onset of diabetes (NOD) is: Type 1 or type 2 diabetes TEAE, and/or h) at least 2 values of HbA1c $\geq 6.5\%$ during the TEAE period (NOTE: for patients with only a single measurement available during the TEAE period, a single value $\geq 6.5\%$ will be considered and qualify the patient as NOD by default; for patients with several HbA1c measurements but only with the last one $\geq 6.5\%$, this single value $\geq 6.5\%$ will be considered and qualify the patient as NOD by default), and/or i) at least 2 values of fasting plasma glucose (FPG) ≥ 126 mg/dL (7.0 mmol/L) (NOTE: for patients with only a single measurement available during the TEAE period, a single value ≥ 126 mg/dL (7.0 mmol/L) will NOT be considered and will NOT qualify the patient as NOD; for patients with several FPG measurements but only with the last one ≥ 126 mg/dL (7.0 mmol/L), this single value ≥ 126 mg/dL (7.0 mmol/L) will NOT be considered and will NOT qualify the patient as NOD).

Serious Adverse Events

[00243] An SAE (serious adverse event) is any untoward medical occurrence that at any dose: a) results in death – includes all deaths, even those that appear to be completely unrelated to the investigational study drug (*e.g.*, a car accident in which a patient is a passenger), b) is life-threatening – in the view of the investigator, the patient is at immediate risk of death at the time

of the event; this does not include an AE that had it occurred in a more severe form, might have caused death, c) requires in-patient hospitalization or prolongation of existing hospitalization (where in patient hospitalization is defined as admission to a hospital or an emergency room for longer than 24 hours; prolongation of existing hospitalization is defined as a hospital stay that is longer than was originally anticipated for the event, or is prolonged due to the development of a new AE as determined by the investigator or treating physician), d) results in persistent or significant disability/incapacity (substantial disruption of one's ability to conduct normal life functions), e) is a congenital anomaly/birth defect, and/or f) is an important medical event - important medical events may not be immediately life-threatening or result in death or hospitalization, but may jeopardize the patient or may require intervention to prevent one of the other serious outcomes listed above (e.g., intensive treatment in an emergency room or at home for allergic bronchospasm; blood dyscrasias or convulsions that do not result in hospitalization; or development of drug dependency or drug abuse).

[00244] Thus, the instantly disclosed randomized, double-blind, placebo-controlled, parallel-group, phase 3 study evaluated the efficacy and safety of PCSK9 inhibitor, alirocumab 150 mg subcutaneous, every 2 weeks in reduction of LDL-C compared with placebo after 12 weeks of treatment in adult patients with hoFH. Secondary objectives included evaluation of 1) other lipid parameters (ie, apolipoprotein B [Apo B], non-high-density lipoprotein cholesterol [non-HDL-C], total-cholesterol [TC], proportion of patients with 15%, 30%, and 50% LDL-C reductions, lipoprotein(a) [Lp(a)], HDL-C, triglycerides [TG], Apo A-1); 2) the safety and tolerability of alirocumab; 3) the pharmacokinetics; 4) the potential development of anti-drug (alirocumab) antibodies. Finally, safety assessments included: adverse events (AEs), serious AEs, deaths, discontinuations due to AE.

RESULTS

Primary efficacy analysis in the ITT population

[00245] The primary efficacy analysis showed a statistically significant decrease in percent change from baseline LDL-C at Week 12 for the alirocumab treatment group (LSmean = -26.9%) as compared to the placebo group (LSmean = +8.6%). The LS mean difference between the alirocumab-treated patients and the placebo patients is -35.6% ($p < 0.0001$). The alirocumab LSmean reductions in percent change LDL-C from baseline could be seen as early as visit week 4, and alirocumab benefit was subsequently maintained throughout the 12 week double-blind treatment period.

[00246] The percent change from baseline in LDL-C at week 12 (MMRM - ITT Analysis – ITT Population) is shown in Table 8, below.

Table 8

LDL Cholesterol	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (mmol/L)		
Number	24	45
Mean (SD)	6.723 (4.5524)	7.640 (4.0040)
Median	6.010	6.420
Min : Max	2.23 : 23.21	2.02 : 19.87
Baseline (mg/dL)		
Number	24	45
Mean (SD)	259.6 (175.75)	295.0 (154.59)
Median	232.0	248.0
Min : Max	86 : 896	78 : 767
Week 12 percent change from baseline (%)		
LS mean (SE)	8.6 (6.3)	-26.9 (4.6)
LS mean difference (SE) vs Placebo		-35.6 (7.8)
95% CI		(-51.2 to -19.9)
p-value vs Placebo		<.0001
Note: Least-squares (LS) means, standard errors (SE) and p-value taken from MMRM (mixed-effect model with repeated measures) analysis. The model includes the fixed categorical effects of treatment group, randomization strata as per IVRS, time point, treatment-by-time point interaction, strata-by-time point interaction, as well as the continuous fixed covariates of baseline LDL-C value and baseline value by time-point interaction.		
MMRM model and baseline description run on patients with a baseline value and a post-baseline value in at least one of the analysis windows used in the model.		

[00247] The LDL-C LS Mean (+/-SE) percent change from baseline for the double-blind period: time profile (ITT analysis – ITT population) (data not shown) shows a statistically significant decrease in % change in LDL-C from baseline to week 12. For alirocumab, LSmean vs. baseline is -26.9% (in other words, a decrease in LDL-C). For placebo, LSmean vs. baseline is 8.6% (in other words, an increase in LDL-C).

[00248] The number of patients at the different timepoints (through week 12) were as follows:

Number of patients

Treatment	baseline	week 4	week 8	week 12
Placebo	24	24	24	24
Alirocumab	45	43	45	44

[00249] The LDL-C efficacy in the instant hoFH study is summarized as follows:

Timepoint	placebo (24 patients)	alirocumab (45 patients)	placebo adjusted difference
Wk 4	-0.7%	-31.9%	-31.3%
Wk 8	5.2%	-27.9%	-33.1%
Wk 12	8.6%	-26.9%	-35.6%

Key secondary efficacy variables

[00250] For easy reference, the following table (Table 9) summarizes analysis results on all key secondary endpoints in the hierarchical order for statistical testing at the 0.05 significance level.

The instant study achieved statistically significant results in favor of the alirocumab-treated

patients for the top 7 key efficacy endpoints. Statistical hypothesis testing terminates at the 8th endpoint of “Percent change from baseline in HDL-C at WK12” ($p = 0.3541$). Nominal p-values for the two remaining endpoints of “Percent change from baseline in fasting TG at WK12” and “Percent change from baseline in Apo A-1 at WK12” are provided for descriptive purposes.

5 **Table 9**

Order	Endpoint/Analysis	Placebo Result	Alirocumab Result	Comparison	P-value
1	Percent change from baseline in Apo B at WK12	LS mean: 7.2%	LS mean: -22.5%	Diff: -29.8%	<.0001
2	Percent change from baseline in non-HDL-C at WK12	LS mean: 8.0%	LS mean: -24.8%	Diff: -32.9%	<.0001
3	Percent change from baseline in total cholesterol at WK12	LS mean: 6.6%	LS mean: -19.8%	Diff: -26.5%	<.0001
4	≥ 15% reduction in LDL-C at WK12	Proportion: 12.5%	Proportion: 61.9%	Odds Ratio: 12.2	0.0004
5	≥ 30% reduction in LDL-C at WK12	Proportion: 4.2%	Proportion: 57.1%	Odds Ratio: 36.5	0.0010
6	Percent change from baseline in Lp(a) at WK12	LS mean: 8.8%	LS mean: -19.6%	Diff: -28.4%	<.0001
7	≥ 50% reduction in LDL-C at WK12	Proportion: 0%	Proportion: 26.7%	Odds Ratio*: 17.7	0.0017
8	Percent change from baseline in HDL-C at WK12	LS mean: 2.7%	LS mean: 6.3%	Diff: 3.6%	0.3541
9	Percent change from baseline in fasting TG at WK12	LS mean: 3.9%	LS mean: -7.4%	Diff: -11.3%	0.1112
10	Percent change from baseline in Apo A-1 at WK12	LS mean: 1.4%	LS mean: 5.0%	Diff: 3.6%	0.3212

[00251] More detailed efficacy endpoint statistical results are presented in the following tables. The percent change from baseline in Apo B at week 12 (MMRM - ITT Analysis – ITT Population) results are provided in Table 10, below.

10

15 **Table 10**

Apolipoprotein B	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (g/L)		
Number	24	45
Mean (SD)	1.750 (0.9512)	1.933 (0.8759)
Median	1.610	1.780
Min : Max	0.82 : 5.25	0.66 : 4.68
Baseline (mg/dL)		
Number	24	45
Mean (SD)	175.0 (95.12)	193.3 (87.59)
Median	161.0	178.0
Min : Max	82 : 525	66 : 468
Week 12 percent change from baseline (%)		
LS mean (SE)	7.2 (5.0)	-22.5 (3.7)
LS mean difference (SE) vs Placebo		-29.8 (6.3)
95% CI		(-42.3 to -17.3)
p-value vs Placebo		<.0001
Note: Least-squares (LS) means, standard errors (SE) and p-value taken from MMRM (mixed-effect model with repeated measures) analysis. The model includes the fixed categorical effects of treatment group, randomization strata as per IVRS, time point, treatment-by-time point interaction, strata-by-time point interaction, as well as the continuous fixed covariates of baseline ApoB value and baseline value by time-point interaction.		

[00252] The percent change from baseline in non-HDL-C at week 12 (MMRM - ITT Analysis – ITT Population) results are provided in Table 11, below.

Table 11

Non-HDL Cholesterol	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (mmol/L)		
Number	24	45
Mean (SD)	7.304 (4.5954)	8.302 (4.1535)
Median	6.710	7.120
Min : Max	2.64 : 24.04	2.38 : 20.49
Baseline (mg/dL)		
Number	24	45
Mean (SD)	282.0 (177.41)	320.5 (160.36)
Median	259.0	275.0
Min : Max	102 : 928	92 : 791
Week 12 percent change from baseline (%)		
LS mean (SE)	8.0 (5.9)	-24.8 (4.3)
LS mean difference (SE) vs Placebo		-32.9 (7.4)
95% CI		(-47.6 to -18.2)
p-value vs Placebo		<.0001
Note: Least-squares (LS) means, standard errors (SE) and p-value taken from MMRM (mixed-effect model with repeated measures) analysis. The model includes the fixed categorical effects of treatment group, randomization strata as per IVRS, time point, treatment-by-time point interaction, strata-by-time point interaction, as well as the continuous fixed covariates of baseline non-HDL-C value and baseline value by time-point interaction.		

[00253] The percent change from baseline in total cholesterol at week 12 (MMRM - ITT Analysis – ITT Population) results are provided in Table 12, below.

Table 12

Total Cholesterol	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (mmol/L)		
Number	24	45
Mean (SD)	8.422 (4.4443)	9.437 (4.0742)
Median	7.915	8.440
Min : Max	4.01 : 24.53	4.20 : 21.50
Baseline (mg/dL)		
Number	24	45
Mean (SD)	325.1 (171.57)	364.3 (157.30)
Median	305.5	326.0
Min : Max	155 : 947	162 : 830
Week 12 percent change from baseline (%)		
LS mean (SE)	6.6 (5.0)	-19.8 (3.7)
LS mean difference (SE) vs Placebo		-26.5 (6.2)
95% CI		(-38.9 to -14.0)
p-value vs Placebo		<.0001
Note: Least-squares (LS) means, standard errors (SE) and p-value taken from MMRM (mixed-effect model with repeated measures) analysis. The model includes the fixed categorical effects of treatment group, randomization strata as per IVRS, time point, treatment-by-time point interaction, strata-by-time point interaction, as well as the continuous fixed covariates of baseline total cholesterol value and baseline value by time-point interaction.		

- 5 **[00254]** The percent change from baseline in Lp(a) at week 12 (multiple imputation followed by robust regression - ITT Analysis – ITT Population) results are provided in Table 13, below.

10

Table 13

Lipoprotein-a	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (g/L)		
Combined estimate for mean (SE)	0.400 (0.074)	0.429 (0.054)
Baseline (mg/dL)		
Combined estimate for mean (SE)	40.0 (7.4)	42.9 (5.4)

Lipoprotein-a	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Week 12 percent change from baseline (%)		
Combined estimate for adjusted mean (SE)	8.8 (5.4)	-19.6 (4.0)
Combined estimate for adjusted mean difference (SE) (Alirocumab vs Comparator)		-28.4 (6.7)
95% CI		(-41.5 to -15.2)
p-value vs Placebo		<.0001
<p>Note: The two-step multiple imputation procedure is used to address missing values in the randomized population (seeds=1628 and 3256; number of imputations=100 and 1 in the two steps respectively). In the first step, the monotone missing pattern is induced in the multiply-imputed data. In the second step, the missing data at subsequent visits are imputed using the regression method for continuous variables. Combined estimates and standard errors (SE) are obtained by combining adjusted means and SE from robust regression model analyses of the different imputed data sets. The robust regression models include the fixed categorical effect of treatment group and randomization strata as per IVRS and the continuous fixed covariate of baseline Lp(a) value. Rubin's formulae are used to combine means and SE.</p>		

[00255] The percent change from baseline in HDL-C at week 12 (MMRM - ITT Analysis – ITT Population) results are provided in Table 14, below.

Table 14

HDL Cholesterol	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (mmol/L)		
Number	24	45
Mean (SD)	1.118 (0.3095)	1.134 (0.3824)
Median	1.195	1.040
Min : Max	0.49 : 1.55	0.62 : 2.46
Baseline (mg/dL)		
Number	24	45
Mean (SD)	43.2 (11.96)	43.8 (14.78)
Median	46.0	40.0
Min : Max	19 : 60	24 : 95
Week 12 percent change from baseline (%)		
LS mean (SE)	2.7 (3.1)	6.3 (2.3)
LS mean difference (SE) vs Placebo		3.6 (3.8)
95% CI		(-4.1 to 11.3)
p-value vs Placebo		0.3541
<p>Note: Least-squares (LS) means, standard errors (SE) and p-value taken from MMRM (mixed-effect model with repeated measures) analysis. The model includes the fixed categorical effects of treatment group, randomization strata as per IVRS, time point, treatment-by-time point interaction, strata-by-time point interaction, as well as the continuous fixed covariates of baseline HDL-C value and baseline value by time-point interaction.</p>		

5

[00256] The percent change from baseline in fasting triglycerides at week 12 (Multiple Imputation Followed by Robust Regression - ITT Analysis – ITT Population) results are provided in Table 15, below.

Table 15

Fasting triglycerides	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (mmol/L)		
Combined estimate for mean (SE)	1.263 (0.180)	1.446 (0.125)

Fasting triglycerides	Placebo (N=24)	Alirocumab 150 Q2W (N=45)
Baseline (mg/dL)		
Combined estimate for mean (SE)	111.7 (15.9)	128.0 (11.1)
Week 12 percent change from baseline (%)		
Combined estimate for adjusted mean (SE)	3.9 (5.7)	-7.4 (4.2)
Combined estimate for adjusted mean difference (SE) (Alirocumab vs Comparator)		-11.3 (7.1)
95% CI		(-25.2 to 2.6)
p-value vs Placebo		0.1112
<p>Note: The two-step multiple imputation procedure is used to address missing values in the randomized population (seeds=1628 and 3256; number of imputations=100 and 1 in the two steps respectively). In the first step, the monotone missing pattern is induced in the multiply-imputed data. In the second step, the missing data at subsequent visits are imputed using the regression method for continuous variables.</p> <p>Combined estimates and standard errors (SE) are obtained by combining adjusted means and SE from robust regression model analyses of the different imputed data sets. The robust regression models include the fixed categorical effect of treatment group and randomization strata as per IVRS and the continuous fixed covariate of baseline TG value. Rubin's formulae are used to combine means and SE.</p>		

[00257] Alirocumab treatment resulted in reductions in LDL-C in hoFH patients with various genotypes, including homozygous (LDLR), compound heterozygous (LDLR), double heterozygous (LDLR + APOB or PCSK9) and heterozygous (LDLR + other benign variants), with the expected minimal to no effect in null/null patients. No LDL-C reductions were observed in placebo-treated patients with any genotype.

[00258] In summary, after a 2-week screening period, 69 patients were randomized to study treatment (24 placebo; 45 alirocumab) with comparable demographics in both groups. At the time of randomization, 97% patients were on high intensity statin; 72% on ezetimibe; 17.4% on apheresis; mean baseline LDL-C was 259.6 mg/dL in placebo; 295.0 mg/dL in alirocumab. Difference in LDL-C change from baseline at week 12 was -35.6% (alirocumab [-26.9%] vs. placebo [8.6%]; $P<0.0001$). Secondary endpoints: ApoB -29.8% ($P<0.0001$); non-HDL-C -32.9% ($P<0.0001$); total cholesterol -26.5% ($P<0.0001$). No treatment-emergent SAEs occurred; no discontinuation due to a TEAE; no deaths reported.

[00259] Thus, treatment with alirocumab resulted in statistically significant and clinically meaningful reductions in LDL-C in patients with hoFH, observed early at visit week 4, and subsequently maintained throughout the 12-week double-blind treatment period. Treatment with alirocumab also resulted in significant reductions in other lipoprotein and lipid measures associated with elevated cardiovascular risk (Apo B, total cholesterol, non-HDL-C and Lp(a)). Finally, alirocumab was generally well tolerated with no clinically significant differences between treatment groups with regards to TEAEs, AESIs (adverse events of special interest), and laboratory parameters. Additionally, no safety concerns were identified from the open label data.

[00260] The present disclosure is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the disclosure in addition to those described

herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

Claims

What is claimed is:

1. A method for treating homozygous familial hypercholesterolemia (hoFH) in a patient in need thereof, the method comprising:

- 5 (a) selecting a patient having hoFH who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy; and
 (b) administering one or more doses of a PCSK9 inhibitor to the patient.

10 2. A method for reducing serum LDL-C levels in a patient having homozygous familial hypercholesterolemia (hoFH), the method comprising:

- (a) selecting a patient who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy; and
 (b) administering one or more doses of a PCSK9 inhibitor to the patient.

15 3. A method for treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH), the method comprising:

- (a) selecting a patient who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy; and
20 (b) administering one or more doses of a PCSK9 inhibitor to the patient.

4. A method for treating homozygous familial hypercholesterolemia (hoFH) in a patient in need thereof, the method comprising:

- (a) selecting a patient having hoFH; and
25 (b) administering a therapeutically effective amount of one or more doses of a PCSK9 inhibitor to the patient.

5. A method for reducing serum LDL-C levels in a patient having homozygous familial hypercholesterolemia (hoFH), the method comprising:

- 30 administering a therapeutically effective amount of one or more doses of a PCSK9 inhibitor to the patient in need thereof.

6. A method for treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH), the

35 method comprising:

administering a therapeutically effective amount of one or more doses of a PCSK9 inhibitor to the patient in need thereof.

5 7. The method of any one of claims 1-6, wherein the patient is diagnosed with hoFH based on genotype or clinical criteria.

8. The method of claim 7, wherein the genotype is selected from the group consisting of:

- 10 (a) homozygous non-null/non-null;
(b) compound heterozygous non-null/non-null;
(c) compound heterozygous non-null/null; and
(d) homozygous null/null.

15 9. The method of claim 7, wherein the clinical criteria is selected from the group consisting of:

- (a) untreated total cholesterol >500 mg/dL (12.93 mmol/L) and triglycerides <300 mg/dL (3.39 mmol/L),
(b) both parents with history of total cholesterol >250 mg/dL (6.46 mmol/L), and
(c) cutaneous or tendinous xanthoma before age 10.

20 10. The method of any one of claims 1-9, wherein the patient is undergoing LDL apheresis.

25 11. The method of any one of claims 1-10, wherein the patient is receiving at least one lipid-modifying therapy (LMT) prior to or at the time of administration of the PCSK9 inhibitor.

12. The method of claim 11, wherein the at least one LMT is at least one statin.

30 13. The method of claim 11, wherein the at least one LMT is LDL apheresis.

14. The method of claim 11, wherein the at least one LMT is ezetimibe.

35 15. The method of claim 11, wherein the at least one LMT is a fibrate, bile acid sequestrant, cholesterol absorption inhibitor, nicotinic acid or derivative, omega 3 fatty acid, probucol, lomitapide, or mipomersen.

16. The method of any one of claims 1-15, wherein the patient has at least about 100 mg/dL LDL-C prior to or at the time of administration of the PCSK9 inhibitor.

17. The method of any one of claims 1-15, wherein the patient has about 500 mg/dL to about 1000 mg/dL LDL-C prior to or at the time of administration of the PCSK9 inhibitor.

18. The method of claim 16 or 17, wherein the patient has an increased risk for premature cardiovascular disease and/or for a cardiovascular event.

19. The method of any one of claims 1-18, wherein the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof that specifically binds PCSK9.

20. The method of claim 19, wherein the antibody or antigen binding fragment thereof is administered to the patient at a dose of about 75 mg at a frequency of once every two weeks.

21. The method of claim 19, wherein the antibody or antigen binding fragment thereof is administered to the patient at a dose of about 150 mg at a frequency of once every two weeks.

22. The method of claim 19, wherein the antibody or antigen binding fragment thereof is administered to the patient at a dose of about 300 mg at a frequency of once every four weeks.

23. The method of any one of claims 19-22, wherein the antibody or antigen binding fragment thereof is administered to the patient subcutaneously.

24. The method of any one of claims 19-23, wherein the antibody or antigen binding fragment thereof comprises the heavy and light chain CDRs of a HCVR/LCVR amino acid sequence pair comprising SEQ ID NOs: 1/6.

25. The method of claim 12, wherein the antibody or antigen-binding fragment thereof comprises heavy and light chain CDR amino acid sequences having SEQ ID NOs: 2, 3, 4, 7, 8, and 10.

26. The method of claim 13, wherein the antibody or antigen-binding fragment thereof comprises an HCVR having the amino acid sequence of SEQ ID NO: 1 and an LCVR having the amino acid sequence of SEQ ID NO: 6.

27. The method of any one of claims 19-28, wherein the antibody or antigen-binding fragment thereof is contained in a pre-filled pen delivery device.

28. The method of any one of claims 1-27, wherein, about 12 weeks after administration of the one or more doses of the PCSK9 inhibitor, the patient exhibits one or more lipid parameter improvements selected from the group consisting of:

- i) a reduction in LDL-C level from baseline of about 35%;
- ii) a reduction in non-HDL-C level from baseline of about 33%;
- iii) a reduction in Apo B level from baseline of about 30%;
- iv) a reduction in total cholesterol level from baseline of about 27%;
- v) a reduction in (fasting) triglyceride level from baseline of about 11%; and
- vi) a reduction in Lp(a) level from baseline of about 28%.

29. The method of claim 28, wherein about 12 weeks after administration of the one or more doses of the PCSK9 inhibitor, the patient exhibits a reduction in LDL-C level from baseline of about 35%.

30. A pharmaceutical composition for use in treating homozygous familial hypercholesterolemia (hoFH) in a patient in need thereof, wherein the composition comprises a PCSK9 inhibitor and a pharmaceutically acceptable excipient.

31. A pharmaceutical composition for use in reducing serum LDL-C levels in a patient having homozygous familial hypercholesterolemia (hoFH), wherein the composition comprises a PCSK9 inhibitor and a pharmaceutically acceptable excipient.

32. A pharmaceutical composition for use in treating, delaying onset of, and/or reducing the risk of developing atherosclerosis in a patient having homozygous familial hypercholesterolemia (hoFH), wherein the composition comprises a PCSK9 inhibitor and a pharmaceutically acceptable excipient.

33. The pharmaceutical composition of any one of claims 30-32, wherein the patient having hoFH is refractory to treatment with statins, is intolerant to statins, and/or has a history of adverse reactions to statin therapy.

34. The pharmaceutical composition of any one of claims 30-33, wherein the PCSK9 inhibitor is an antibody or an antigen binding fragment thereof that specifically binds PCSK9.

35. The pharmaceutical composition of claim 34, wherein the antibody or antigen binding fragment thereof comprises the heavy and light chain CDRs of a HCVR/LCVR amino acid sequence pair comprising SEQ ID NOs: 1/6.

5 36. The pharmaceutical composition of claim 34 or 35, wherein the antibody or antigen binding fragment thereof is administered to the patient at a dose of about 75 mg at a frequency of once every two weeks.

10 37. The pharmaceutical composition of claim 34 or 35, wherein the antibody or antigen binding fragment thereof that specifically binds PCSK9 is administered to the patient at a dose of about 150 mg at a frequency of once every two weeks.

15 38. The pharmaceutical composition of claim 34 or 35, wherein the antibody or antigen binding fragment thereof is administered to the patient at a dose of about 300 mg at a frequency of once every four weeks.

20 39. A method for treating homozygous familial hypercholesterolemia (hoFH), the method comprising administering one or more doses of a PCSK9 inhibitor to a patient who is diagnosed with hoFH and who is refractory to treatment with statins, who is intolerant to statins, or who has a history of adverse reactions to statin therapy.




FIG. 1




FIG. 2

SEQUENCE LISTING

<110> Regeneron Pharmaceuticals, Inc.
Sanofi Biotechnology

<120> USE OF A PCSK9 INHIBITOR TO TREAT HOMOZYGOUS FAMILIAL
HYPERCHOLESTEROLEMIA

<130> 711599:SA9-291PC

<140> 62946268

<141> 2019-12-10

<140> 62987148

<141> 2020-03-09

<160> 198

<170> PatentIn version 3.5

<210> 1

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 1

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Asn	Asn	Tyr
			20					25					30		

Ala	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Asp	Trp	Val
		35					40					45			

Ser	Thr	Ile	Ser	Gly	Ser	Gly	Gly	Thr	Thr	Asn	Tyr	Ala	Asp	Ser	Val
	50					55					60				

Lys	Gly	Arg	Phe	Ile	Ile	Ser	Arg	Asp	Ser	Ser	Lys	His	Thr	Leu	Tyr
65					70					75					80

Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85					90					95	

Ala Lys Asp Ser Asn Trp Gly Asn Phe Asp Leu Trp Gly Arg Gly Thr
100 105 110

Leu Val Thr Val Ser Ser
115

<210> 2
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 2
Gly Phe Thr Phe Asn Asn Tyr Ala
1 5

<210> 3
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 3
Ile Ser Gly Ser Gly Gly Thr Thr
1 5

<210> 4
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 4
Ala Lys Asp Ser Asn Trp Gly Asn Phe Asp Leu
1 5 10

<210> 5
<211> 447
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
REGN727 heavy chain polypeptide

<400> 5

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Asn Tyr
20 25 30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Asp Trp Val
35 40 45

Ser Thr Ile Ser Gly Ser Gly Gly Thr Thr Asn Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Ile Ile Ser Arg Asp Ser Ser Lys His Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Lys Asp Ser Asn Trp Gly Asn Phe Asp Leu Trp Gly Arg Gly Thr
100 105 110

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
195 200 205

Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr
210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
340 345 350

Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
435 440 445

<210> 6

<211> 113

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 6

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Arg
20 25 30

Ser Asn Asn Arg Asn Phe Leu Gly Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Asn Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Thr Thr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
100 105 110

Lys

<210> 7

<211> 12

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 7

Gln	Ser	Val	Leu	Tyr	Arg	Ser	Asn	Asn	Arg	Asn	Phe
1			5					10			

<210> 8

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 8

Trp	Ala	Ser
1		

<210> 9

<211> 220

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic REGN727 light chain polypeptide

<400> 9

Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	Leu	Gly
1			5					10				15			

Glu	Arg	Ala	Thr	Ile	Asn	Cys	Lys	Ser	Ser	Gln	Ser	Val	Leu	Tyr	Arg
			20					25					30		

Ser	Asn	Asn	Arg	Asn	Phe	Leu	Gly	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln
		35					40					45			

Pro	Pro	Asn	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val
	50					55					60				

Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
65					70					75				80	

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Thr Thr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
100 105 110

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
115 120 125

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
130 135 140

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
145 150 155 160

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
165 170 175

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
180 185 190

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
195 200 205

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215 220

<210> 10

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 10

Gln Gln Tyr Tyr Thr Thr Pro Tyr Thr
1 5

<210> 11

<211> 127

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 11

Glu Met Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser His
 20 25 30

Trp Met Lys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Asn Ile Asn Gln Asp Gly Ser Glu Lys Tyr Tyr Val Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Phe
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Asp Ile Val Leu Met Val Tyr Asp Met Asp Tyr Tyr Tyr Tyr
 100 105 110

Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 115 120 125

<210> 12

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 12

Gly Phe Thr Phe Ser Ser His Trp
1 5

<210> 13
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 13
Ile Asn Gln Asp Gly Ser Glu Lys
1 5

<210> 14
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 14
Ala Arg Asp Ile Val Leu Met Val Tyr Asp Met Asp Tyr Tyr Tyr Tyr
1 5 10 15

Gly Met Asp Val
20

<210> 15
<211> 112
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 15
Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
20 25 30

Asn Gly Asn Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Thr
85 90 95

Leu Gln Thr Pro Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

<210> 16

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 16

Gln Ser Leu Leu His Ser Asn Gly Asn Asn Tyr
1 5 10

<210> 17

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 17

Leu Gly Ser
1

<210> 18

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 18

Met Gln Thr Leu Gln Thr Pro Leu Thr
1 5

<210> 19
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 19
Gly Phe Thr Phe Ser Ser His Trp
1 5

<210> 20
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 20
Ile Asn Gln Asp Gly Ser Glu Lys
1 5

<210> 21
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 21
Ala Arg Asp Ile Val Leu Met Val Tyr Asp Met Asp Tyr Tyr Tyr Tyr
1 5 10 15

Gly Met Asp Val
20

<210> 22
<211> 11
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 22

Gln Ser Leu His His Ser Asn Gly Asn Asn Tyr
1 5 10

<210> 23

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 23

Leu Gly Ser
1

<210> 24

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 24

Met Gln Thr Leu Gln Thr Pro Leu Thr
1 5

<210> 25

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 25

Gly Phe Thr Phe Ser Ser His Trp
1 5

<210> 26

<211> 8

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 26

Ile Asn Gln Asp Gly Ser Glu Lys
1 5

<210> 27

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 27

Ala Arg Asp Ile Val Leu Met Val Tyr His Met Asp Tyr Tyr Tyr Tyr
1 5 10 15

Gly Met Asp Val
20

<210> 28

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 28

Gln Ser Leu Leu His Ser Asn Gly Asn Asn Tyr
1 5 10

<210> 29

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 29

Leu Gly Ser
1

<210> 30
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 30
Met Gln Thr Leu Gln Thr Pro Leu Thr
1 5

<210> 31
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 31
Gly Phe Thr Phe Ser Ser His Trp
1 5

<210> 32
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 32
Ile Asn Gln Asp Gly Ser Glu Lys
1 5

<210> 33
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 33

Ala Arg Asp Ile Val Leu Met Val Tyr His Met Asp Tyr Tyr Tyr Tyr
1 5 10 15

Gly Met Asp Val
20

<210> 34

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 34

Gln Ser Leu His His Ser Asn Gly Asn Asn Tyr
1 5 10

<210> 35

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 35

Leu Gly Ser
1

<210> 36

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 36

Met Gln Thr Leu Gln Thr Pro Leu Thr
1 5

<210> 37

<211> 131

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VH; m2CX1D05 polypeptide

<400> 37

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Asn Ser His
20 25 30

Ala Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Gly Ile Asn Pro Ile Leu Gly Ile Ala Asn Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg His Tyr Glu Ile Gln Ile Gly Arg Tyr Gly Met Asn Val Tyr
100 105 110

Tyr Leu Met Tyr Arg Phe Ala Ser Trp Gly Gln Gly Thr Leu Val Thr
115 120 125

Val Ser Ser
130

<210> 38

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VH CDR1; m2CX1D05 peptide

<400> 38

Gly Gly Thr Phe Asn Ser His Ala Ile Ser
1 5 10

<210> 39
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
VH CDR2; m2CX1D05 peptide

<400> 39
Trp Met Gly Gly Ile Asn Pro Ile Leu Gly Ile Ala Asn Tyr Ala Gln
1 5 10 15

Lys Phe Gln Gly
20

<210> 40
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
VH CDR3; m2CX1D05 peptide

<400> 40
His Tyr Glu Ile Gln Ile Gly Arg Tyr Gly Met Asn Val Tyr Tyr Leu
1 5 10 15

Met Tyr Arg Phe Ala Ser
20

<210> 41
<211> 213
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
LC; m2CX1D05 polypeptide

<400> 41
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Ser Ala
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Asn Gly Ser Thr Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Phe Asp Gly Asp Pro Thr
85 90 95

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130 135 140

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145 150 155 160

Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
165 170 175

Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
180 185 190

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
195 200 205

Asn Arg Gly Glu Ala
210

<210> 42

<211> 11

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VL CDR 1; m2CX1D05 peptide

<400> 42
Arg Ala Ser Gln Gly Ile Arg Ser Ala Leu Asn
1 5 10

<210> 43
<211> 11
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VL CDR2; m2CX1D05 peptide

<400> 43
Leu Leu Ile Tyr Asn Gly Ser Thr Leu Gln Ser
1 5 10

<210> 44
<211> 7
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VL CDR3; m2CX1D05 peptide

<400> 44
Gln Gln Phe Asp Gly Asp Pro
1 5

<210> 45
<211> 119
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VH; 1B20 polypeptide

<400> 45
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Asn Tyr
20 25 30

Trp Ile Ser Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
35 40 45

Gly Ile Ile Tyr Pro Gly Asp Ser Tyr Thr Asn Tyr Ser Pro Ser Phe
50 55 60

Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80

Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ala Arg Asp Tyr Trp Tyr Lys Pro Leu Phe Asp Ile Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ser
115

<210> 46

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VH CDR1; 1B20 peptide

<400> 46

Gly Tyr Ser Phe Thr Asn Tyr Trp Ile Ser
1 5 10

<210> 47

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VH CDR2; 1B20 peptide

<400> 47

Trp Met Gly Ile Ile Tyr Pro Gly Asp Ser Tyr Thr Asn Tyr Ser Pro
1 5 10 15

Ser Phe Gln Gly
20

<210> 48

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VH CDR3; 1B20 peptide

<400> 48

Asp Tyr Trp Tyr Lys Pro Leu Phe Asp Ile
1 5 10

<210> 49

<211> 220

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
LC; 1B20 polypeptide

<400> 49

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Arg Ser Ser Gln Ser Val Leu Tyr Ser
20 25 30

Ser Asn Asn Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Ser Ser Phe Pro Ile Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100 105 110

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
115 120 125

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
130 135 140

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
145 150 155 160

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
165 170 175

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
180 185 190

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
195 200 205

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Ala
210 215 220

<210> 50

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VL CDR1; 1B20 peptide

<400> 50

Arg Ser Ser Gln Ser Val Leu Tyr Ser Ser Asn Asn Lys Asn Tyr Leu
1 5 10 15

Ala

<210> 51

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VL CDR2; 1B20 peptide

<400> 51

Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser
1 5 10

<210> 52

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
VL CDR3; 1B20 peptide

<400> 52

Gln Gln Tyr Ser Ser Phe Pro Ile
1 5

<210> 53

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable heavy antibody region polypeptide

<400> 53

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Ser Tyr
 20 25 30

Gly Met Tyr Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45

Gly Trp Ile Asp Pro Gly Ser Gly Gly Thr Lys Tyr Asn Glu Lys Phe
 50 55 60

Lys Gly Lys Ala Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Arg Tyr Gly Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser Ala Ser
115 120

<210> 54
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX132 heavy chain CDR1 antibody region peptide

<400> 54
Lys Ala Ser Gly Tyr Thr Phe Ser Ser Tyr Gly Met Tyr Trp Val Arg
1 5 10 15

<210> 55
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX132 heavy chain CDR2 antibody region peptide

<400> 55
Trp Ile Gly Trp Ile Asp Pro Gly Ser Gly Gly Thr Lys Tyr Asn Glu
1 5 10 15

Lys Phe Lys Gly Lys Ala Thr
20

<210> 56
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX132 heavy chain CDR3 antibody region peptide

<400> 56

Cys Ala Arg Glu Arg Tyr Gly Tyr Tyr Phe Asp Tyr Trp Gly Gln
1 5 10 15

<210> 57

<211> 108

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable light antibody region polypeptide

<400> 57

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Ile Thr Cys Arg Ala Ser Gln Tyr Val Gly Ser Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Val Trp Asp Ser Ser Pro Pro
85 90 95

Val Val Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 58

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX213 and AX132 light chain CDR1 antibody region peptide

<400> 58

Ile Thr Cys Arg Ala Ser Gln Tyr Val Gly Ser Tyr Leu Asn Trp Tyr
1 5 10 15

Gln

<210> 59

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX213 and AX132 light chain CDR2 antibody region peptide

<400> 59

Leu Ile Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro
1 5 10

<210> 60

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX132 & AX213 light chain CDR3 antibody region peptide

<400> 60

Tyr Tyr Cys Gln Val Trp Asp Ser Ser Pro Pro Val Val Phe Gly Gly
1 5 10 15

<210> 61

<211> 120

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable heavy antibody region polypeptide

<400> 61

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Arg Tyr
20 25 30

Gly Ile Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
35 40 45

Gly Arg Ile Asp Pro Gly Asn Gly Gly Thr Arg Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Lys Ala Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ala Asn Asp Gly Tyr Ser Phe Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser Ala Ser
115 120

<210> 62
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX213 heavy chain CDR1 antibody region peptide

<400> 62
Lys Ala Ser Gly Tyr Thr Phe Ser Arg Tyr Gly Ile Asn Trp Val Arg
1 5 10 15

<210> 63
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX213 heavy chain CDR2 antibody region peptide

<400> 63
Trp Ile Gly Arg Ile Asp Pro Gly Asn Gly Gly Thr Arg Tyr Asn Glu
1 5 10 15

Lys Phe Lys Gly Lys Ala Thr
20

<210> 64
<211> 15
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX213 heavy chain CDR3 antibody region peptide

<400> 64

Cys Ala Arg Ala Asn Asp Gly Tyr Ser Phe Asp Tyr Trp Gly Gln
1 5 10 15

<210> 65
<211> 108
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable light antibody region polypeptide

<400> 65

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Ile Thr Cys Arg Ala Ser Gln Tyr Val Gly Ser Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Asp Ala Ser Asn Arg Ala Thr Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Val Trp Asp Ser Ser Pro Pro
85 90 95

Val Val Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 66
<211> 17

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX213 and AX132 light chain CDR1 antibody region peptide

<400> 66

Ile	Thr	Cys	Arg	Ala	Ser	Gln	Tyr	Val	Gly	Ser	Tyr	Leu	Asn	Trp	Tyr
1				5					10					15	

Gln

<210> 67
<211> 13
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX213 and AX132 light chain CDR2 antibody region peptide

<400> 67

Leu	Ile	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro
1				5					10			

<210> 68
<211> 16
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX132 & AX213 light chain CDR3 antibody region peptide

<400> 68

Tyr	Tyr	Cys	Gln	Val	Trp	Asp	Ser	Ser	Pro	Pro	Val	Val	Phe	Gly	Gly
1				5					10					15	

<210> 69
<211> 119
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX1 VH antibody sequence polypeptide

<400> 69

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Ser Tyr
20 25 30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
35 40 45

Gly Arg Ile Asn Pro Asp Ser Gly Ser Thr Lys Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Ala Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Gly Arg Leu Ser Trp Asp Phe Asp Val Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ser
115

<210> 70

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX1 VH CDR1 antibody sequence peptide

<400> 70

Lys Ala Ser Gly Phe Thr Phe Thr Ser Tyr Tyr Met His Trp Val Arg
1 5 10 15

<210> 71

<211> 23

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX1 VH CDR2 antibody sequence peptide

<400> 71

Trp Ile Gly Arg Ile Asn Pro Asp Ser Gly Ser Thr Lys Tyr Asn Glu
1 5 10 15

Lys Phe Lys Gly Arg Ala Thr
20

<210> 72

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX1 VH CDR3 antibody sequence peptide

<400> 72

Cys Ala Arg Gly Gly Arg Leu Ser Trp Asp Phe Asp Val Trp Gly Gln
1 5 10 15

<210> 73

<211> 109

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX1 VL antibody sequence polypeptide

<400> 73

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Ser Arg Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Ala Ala Tyr Asp Tyr Ser Leu Gly
85 90 95

Gly Tyr Val Phe Gly Asp Gly Thr Lys Val Glu Ile Lys
100 105

<210> 74
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX1 VL CDR1 antibody sequence peptide

<400> 74
Arg Ala Ser Gln Asp Ile Ser Arg Tyr Leu Ala
1 5 10

<210> 75
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX1 AX9 AX189 VL CDR2 antibody sequence peptide

<400> 75
Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 76
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
AX1 VL CDR3 antibody sequence peptide

<400> 76
Ala Ala Tyr Asp Tyr Ser Leu Gly Gly Tyr Val
1 5 10

<210> 77
<211> 121
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX9 AX189 VH antibody sequence polypeptide

<400> 77

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Ser Tyr
20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
35 40 45

Gly Arg Ile Asp Pro Tyr Asn Gly Gly Thr Lys Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Lys Ala Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Tyr Gly Tyr Tyr Leu Gly Ser Tyr Ala Met Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 78

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX9 AX189 VH CDR1 antibody sequence peptide

<400> 78

Lys Ala Ser Gly Tyr Thr Phe Ser Ser Tyr Trp Met His Trp Val Arg
1 5 10 15

<210> 79

<211> 23

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX9 AX189 VH CDR2 antibody sequence peptide

<400> 79

Trp	Ile	Gly	Arg	Ile	Asp	Pro	Tyr	Asn	Gly	Gly	Thr	Lys	Tyr	Asn	Glu
1				5					10					15	

Lys	Phe	Lys	Gly	Lys	Ala	Thr
			20			

<210> 80

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX9 AX189 VH CDR3 antibody sequence peptide

<400> 80

Cys	Ala	Arg	Tyr	Gly	Tyr	Tyr	Leu	Gly	Ser	Tyr	Ala	Met	Asp	Tyr	Trp
1				5				10						15	

Gly Gln

<210> 81

<211> 109

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX189 VL antibody sequence polypeptide

<400> 81

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5					10					15	

Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Asp	Val	Ser	Arg	Tyr
			20					25					30		

Leu	Thr	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
		35					40				45				

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Ala Tyr Asp Tyr Ser Leu Ser
85 90 95

Gly Tyr Val Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 82

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX189 VL CDR1 antibody sequence peptide

<400> 82

Arg Ala Ser Gln Asp Val Ser Arg Tyr Leu Thr
1 5 10

<210> 83

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX1 AX9 AX189 VL CDR2 antibody sequence peptide

<400> 83

Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 84

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
AX189 VL CDR3 antibody sequence peptide

<400> 84

Gln Ala Tyr Asp Tyr Ser Leu Ser Gly Tyr Val
1 5 10

<210> 85

<211> 115

<212> PRT

<213> Homo sapiens

<400> 85

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Leu Thr Ser Tyr
20 25 30

Gly Ile Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Trp Val Ser Phe Tyr Asn Gly Asn Thr Asn Tyr Ala Gln Lys Leu
50 55 60

Gln Gly Arg Gly Thr Met Thr Thr Asp Pro Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr
100 105 110

Val Ser Ser
115

<210> 86

<211> 5

<212> PRT

<213> Homo sapiens

<400> 86

Ser Tyr Gly Ile Ser
1 5

<210> 87
<211> 17
<212> PRT
<213> Homo sapiens

<400> 87
Trp Val Ser Phe Tyr Asn Gly Asn Thr Asn Tyr Ala Gln Lys Leu Gln
1 5 10 15

Gly

<210> 88
<211> 6
<212> PRT
<213> Homo sapiens

<400> 88
Gly Tyr Gly Met Asp Val
1 5

<210> 89
<211> 109
<212> PRT
<213> Homo sapiens

<400> 89
Gln Ser Ala Leu Thr Gln Pro Ala Ser Val Ser Gly Ser Pro Gly Gln
1 5 10 15

Ser Ile Thr Ile Ser Cys Thr Gly Thr Ser Ser Asp Val Gly Gly Tyr
20 25 30

Asn Ser Val Ser Trp Tyr Gln Gln His Pro Gly Lys Ala Pro Lys Leu
35 40 45

Met Ile Tyr Glu Val Ser Asn Arg Pro Ser Gly Val Ser Asn Arg Phe
50 55 60

Ser Gly Ser Lys Ser Gly Asn Thr Ala Ser Leu Thr Ile Ser Gly Leu
65 70 75 80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Asn Ser Tyr Thr Ser Thr
85 90 95

Ser Met Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu
100 105

<210> 90
<211> 14
<212> PRT
<213> Homo sapiens

<400> 90
Thr Gly Thr Ser Ser Asp Val Gly Gly Tyr Asn Ser Val Ser
1 5 10

<210> 91
<211> 7
<212> PRT
<213> Homo sapiens

<400> 91
Glu Val Ser Asn Arg Pro Ser
1 5

<210> 92
<211> 9
<212> PRT
<213> Homo sapiens

<400> 92
Asn Ser Tyr Thr Ser Thr Ser Met Val
1 5

<210> 93
<211> 123
<212> PRT
<213> Homo sapiens

<400> 93
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Ser Ile Ser Ser Ser Ser Ser Tyr Ile Ser Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Phe Cys
85 90 95

Ala Arg Asp Tyr Asp Phe Trp Ser Ala Tyr Tyr Asp Ala Phe Asp Val
100 105 110

Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser
115 120

<210> 94
<211> 10
<212> PRT
<213> Homo sapiens

<400> 94
Gly Phe Thr Phe Ser Ser Tyr Ser Met Asn
1 5 10

<210> 95
<211> 17
<212> PRT
<213> Homo sapiens

<400> 95
Ser Ile Ser Ser Ser Ser Ser Tyr Ile Ser Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 96
<211> 14
<212> PRT
<213> Homo sapiens

<400> 96
Asp Tyr Asp Phe Trp Ser Ala Tyr Tyr Asp Ala Phe Asp Val
1 5 10

<210> 97
<211> 111

<212> PRT

<213> Homo sapiens

<400> 97

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30

Tyr Asp Val His Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu
35 40 45

Leu Ile Ser Gly Asn Ser Asn Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu
65 70 75 80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser
85 90 95

Leu Ser Gly Ser Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu
100 105 110

<210> 98

<211> 14

<212> PRT

<213> Homo sapiens

<400> 98

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Asp Val His
1 5 10

<210> 99

<211> 7

<212> PRT

<213> Homo sapiens

<400> 99

Gly Asn Ser Asn Arg Pro Ser
1 5

<210> 100

<211> 11

<212> PRT
<213> Homo sapiens

<400> 100
Gln Ser Tyr Asp Ser Ser Leu Ser Gly Ser Val
1 5 10

<210> 101
<211> 114
<212> PRT
<213> Homo sapiens

<400> 101
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Ala Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Val Ile Tyr Tyr Asp Gly Ile Asn Lys His Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Asp Arg Gly Leu Asp Trp Gly Gln Gly Thr Leu Val Thr Val
100 105 110

Ser Ser

<210> 102
<211> 10
<212> PRT
<213> Homo sapiens

<400> 102
Gly Phe Thr Phe Ser Ser Tyr Gly Met His
1 5 10

<210> 103
<211> 17
<212> PRT
<213> Homo sapiens

<400> 103
Val Ile Tyr Tyr Asp Gly Ile Asn Lys His Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 104
<211> 5
<212> PRT
<213> Homo sapiens

<400> 104
Asp Arg Gly Leu Asp
1 5

<210> 105
<211> 113
<212> PRT
<213> Homo sapiens

<400> 105
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Leu Tyr Ser
20 25 30

Ser Asn Ser Lys Asn Tyr Leu Val Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Thr Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100 105 110

Lys

<210> 106
<211> 17
<212> PRT
<213> Homo sapiens

<400> 106
Lys Ser Ser Gln Ser Val Leu Tyr Ser Ser Asn Ser Lys Asn Tyr Leu
1 5 10 15

Val

<210> 107
<211> 7
<212> PRT
<213> Homo sapiens

<400> 107
Trp Ala Ser Thr Arg Glu Ser
1 5

<210> 108
<211> 9
<212> PRT
<213> Homo sapiens

<400> 108
Gln Gln Tyr Tyr Ser Thr Pro Trp Thr
1 5

<210> 109
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody pJG04 (clones LGT-209 and
LGT-210) Vh heavy chain variable region (FR1-FR4)
polypeptide

<400> 109

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Thr Met
20 25 30

Tyr Met Ser Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Arg Ile Asp Pro Ala Asn Glu His Thr Asn Tyr Ala Gln Lys Phe
50 55 60

Gln Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Arg Leu Thr Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ser Tyr Tyr Tyr Tyr Asn Met Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser
115

<210> 110

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody clones LGT-209,
LGT-210 and LGT-211 heavy chain CDR1 peptide

<400> 110

Thr Met Tyr Met Ser
1 5

<210> 111

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody clones LGT-209,
LGT-210 and LGT-211 heavy chain CDR2 peptide

<400> 111

Arg Ile Asp Pro Ala Asn Glu His Thr Asn Tyr Ala Gln Lys Phe Gln
1 5 10 15

Gly

<210> 112

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody pJG04(clones
LGT-209 and LGT-210) Vh heavy chain complementarity
determining region 3 (CDR3) peptide

<400> 112

Ser Tyr Tyr Tyr Tyr Asn Met Asp Tyr
1 5

<210> 113

<211> 106

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody pJG10(clones
LGT-209 and LGT-211) Vk light chain variable
region (FR1-FR4) polypeptide

<400> 113

Gln Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Tyr
35 40 45

Gly Val Phe Arg Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Gly Arg Leu Glu Pro Glu
65 70 75 80

Asp Phe Ala Val Tyr Tyr Cys Leu Gln Trp Ser Ser Asp Pro Pro Thr
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 114

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody clones LGT-209,
LGT-210 and LGT-211 light chain CDR1 peptide

<400> 114

Arg Ala Ser Gln Ser Val Ser Tyr Met His
1 5 10

<210> 115

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
anti-PCSK9 monoclonal antibody clones LGT-209,
LGT-210 and LGT-211 light chain CDR1 peptide

<400> 115

Gly Val Phe Arg Arg Ala Thr
1 5

<210> 116

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
mouse anti-PCSK9 monoclonal antibody LFU720 and

anti-PCSK9 monoclonal antibody clones LGT-209,
LGT-210 and LGT-211 light chain CDR3 peptide

<400> 116

Leu Gln Trp Ser Ser Asp Pro Pro Thr
1 5

<210> 117

<211> 118

<212> PRT

<213> Homo sapiens

<400> 117

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile Ser Pro Phe Gly Gly Arg Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Ser Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Arg Pro Leu Tyr Ala Ser Asp Leu Trp Gly Gln Gly Thr
100 105 110

Thr Val Thr Val Ser Ser
115

<210> 118

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 118

Gly Tyr Thr Phe Thr Ser Tyr
1 5

<210> 119

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 119

Ser Pro Phe Gly Gly Arg
1 5

<210> 120

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable heavy chain CDR peptide

<400> 120

Glu Arg Pro Leu Tyr Ala Ser Asp Leu
1 5

<210> 121

<211> 107

<212> PRT

<213> Homo sapiens

<400> 121

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Ala
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Arg Tyr Ser Leu Trp Arg
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 122

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable light chain CDR peptide

<400> 122

Arg Ala Ser Gln Gly Ile Ser Ser Ala Leu Ala
1 5 10

<210> 123

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable light chain CDR peptide

<400> 123

Ser Ala Ser Tyr Arg Tyr Thr
1 5

<210> 124

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
variable light chain CDR peptide

<400> 124

Gln Gln Arg Tyr Ser Leu Trp Arg Thr
1 5

<210> 125
<211> 118
<212> PRT
<213> Homo sapiens

<400> 125
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile His Pro Ser Gly Gly Arg Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Ser Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Arg Pro Leu Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Thr Val Thr Val Ser Ser
115

<210> 126
<211> 10
<212> PRT
<213> Homo sapiens

<400> 126
Gly Tyr Thr Phe Thr Ser Tyr Tyr Met His
1 5 10

<210> 127
<211> 17
<212> PRT
<213> Homo sapiens

<400> 127

Glu Ile His Pro Ser Gly Gly Arg Thr Asn Tyr Asn Glu Lys Phe Lys
1 5 10 15

Ser

<210> 128

<211> 9

<212> PRT

<213> Homo sapiens

<400> 128

Glu Arg Pro Leu Tyr Ala Met Asp Tyr
1 5

<210> 129

<211> 107

<212> PRT

<213> Homo sapiens

<400> 129

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val His Thr Ala
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr His Ala Ser Tyr Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Arg Tyr Ser Leu Trp Arg
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 130
<211> 11
<212> PRT
<213> Homo sapiens

<400> 130
Lys Ala Ser Gln Asp Val His Thr Ala Val Ala
1 5 10

<210> 131
<211> 7
<212> PRT
<213> Homo sapiens

<400> 131
His Ala Ser Tyr Arg Tyr Thr
1 5

<210> 132
<211> 9
<212> PRT
<213> Homo sapiens

<400> 132
Gln Gln Arg Tyr Ser Leu Trp Arg Thr
1 5

<210> 133
<211> 118
<212> PRT
<213> Homo sapiens

<400> 133
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Tyr Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile His Pro Ser Gly Gly Arg Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Ser Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Arg Pro Leu Tyr Ala Ser Asp Leu Trp Gly Gln Gly Thr
100 105 110

Thr Val Thr Val Ser Ser
115

<210> 134
<211> 10
<212> PRT
<213> Homo sapiens

<400> 134
Gly Tyr Thr Phe Thr Ser Tyr Tyr Met His
1 5 10

<210> 135
<211> 17
<212> PRT
<213> Homo sapiens

<400> 135
Glu Ile His Pro Ser Gly Gly Arg Thr Asn Tyr Asn Glu Lys Phe Lys
1 5 10 15

Ser

<210> 136
<211> 9
<212> PRT
<213> Homo sapiens

<400> 136
Glu Arg Pro Leu Tyr Ala Ser Asp Leu
1 5

<210> 137
<211> 107
<212> PRT
<213> Homo sapiens

<400> 137

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val His Thr Ala
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr His Ala Ser Tyr Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Arg Tyr Ser Leu Trp Arg
85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 138
<211> 11
<212> PRT
<213> Homo sapiens

<400> 138
Lys Ala Ser Gln Asp Val His Thr Ala Val Ala
1 5 10

<210> 139
<211> 7
<212> PRT
<213> Homo sapiens

<400> 139
His Ala Ser Tyr Arg Tyr Thr
1 5

<210> 140
<211> 9
<212> PRT
<213> Homo sapiens

<400> 140

Gln Gln Arg Tyr Ser Leu Trp Arg Thr
1 5

<210> 141
<211> 118
<212> PRT
<213> Mus musculus

<400> 141
Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Glu Ile Asn Pro Ser Asn Gly Arg Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Arg Pro Leu Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Ser Val Thr Val Ser Ser
115

<210> 142
<211> 7
<212> PRT
<213> Mus musculus

<400> 142
Gly Tyr Thr Phe Thr Ser Tyr
1 5

<210> 143
<211> 6

<212> PRT
<213> Mus musculus

<400> 143
Asn Pro Ser Asn Gly Arg
1 5

<210> 144
<211> 9
<212> PRT
<213> Mus musculus

<400> 144
Glu Arg Pro Leu Tyr Ala Met Asp Tyr
1 5

<210> 145
<211> 108
<212> PRT
<213> Mus musculus

<400> 145
Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly
1 5 10 15

Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Thr Ala
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
35 40 45

Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala
65 70 75 80

Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln Arg Tyr Ser Thr Pro Arg
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> 146
<211> 11

<212> PRT
<213> Mus musculus

<400> 146
Lys Ala Ser Gln Asp Val Ser Thr Ala Val Ala
1 5 10

<210> 147
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
variable light chain CDR peptide

<400> 147
Ser Ala Ser Tyr Arg Tyr Thr
1 5

<210> 148
<211> 9
<212> PRT
<213> Mus musculus

<400> 148
Gln Gln Arg Tyr Ser Thr Pro Arg Thr
1 5

<210> 149
<211> 115
<212> PRT
<213> Mus musculus

<400> 149
Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Tyr Met Asn Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Asp Ile Asn Pro Asn Asn Gly Gly Thr Thr Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Tyr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Trp Leu Leu Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val Ser Ala
115

<210> 150
<211> 7
<212> PRT
<213> Mus musculus

<400> 150
Gly Tyr Thr Phe Thr Asp Tyr
1 5

<210> 151
<211> 6
<212> PRT
<213> Mus musculus

<400> 151
Asn Pro Asn Asn Gly Gly
1 5

<210> 152
<211> 6
<212> PRT
<213> Mus musculus

<400> 152
Trp Leu Leu Phe Ala Tyr
1 5

<210> 153
<211> 108
<212> PRT
<213> Mus musculus

<400> 153
Asp Ile Val Met Thr Gln Ser Gln Lys Phe Met Ser Thr Ser Val Gly
1 5 10 15

Asp Arg Val Ser Val Thr Cys Lys Ala Ser Gln Asn Val Gly Thr Asn
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Ala Leu Ile
35 40 45

Tyr Ser Ala Ser Tyr Arg Tyr Ser Gly Val Pro Asp Arg Phe Thr Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Asn Val Leu Ser
65 70 75 80

Glu Asp Leu Ala Glu Tyr Phe Cys Gln Gln Phe Tyr Ser Tyr Pro Tyr
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> 154
<211> 11
<212> PRT
<213> Mus musculus

<400> 154
Lys Ala Ser Gln Asn Val Gly Thr Asn Val Ala
1 5 10

<210> 155
<211> 7
<212> PRT
<213> Mus musculus

<400> 155
Ser Ala Ser Tyr Arg Tyr Ser
1 5

<210> 156
<211> 9
<212> PRT
<213> Mus musculus

<400> 156
Gln Gln Phe Tyr Ser Tyr Pro Tyr Thr
1 5

<210> 157
<211> 123
<212> PRT
<213> Mus musculus

<400> 157
Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Tyr Met Asn Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Asp Ile Asn Pro Asn Asn Gly Gly Thr Ser Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Gly Gly Gly Ile Tyr Tyr Arg Tyr Asp Arg Asn Tyr Phe Asp Tyr
100 105 110

Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
115 120

<210> 158
<211> 7
<212> PRT
<213> Mus musculus

<400> 158
Gly Tyr Thr Phe Thr Asp Tyr
1 5

<210> 159
<211> 6
<212> PRT
<213> Mus musculus

<400> 159

Asn Pro Asn Asn Gly Gly
1 5

<210> 160

<211> 14

<212> PRT

<213> Mus musculus

<400> 160

Gly Gly Ile Tyr Tyr Arg Tyr Asp Arg Asn Tyr Phe Asp Tyr
1 5 10

<210> 161

<211> 107

<212> PRT

<213> Mus musculus

<400> 161

Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly
1 5 10 15

Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Gly Ile Ser Asn Tyr
20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
35 40 45

Tyr Tyr Thr Ser Ser Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Lys Leu Pro Phe
85 90 95

Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 162

<211> 11

<212> PRT

<213> Mus musculus

<400> 162
Ser Ala Ser Gln Gly Ile Ser Asn Tyr Leu Asn
1 5 10

<210> 163
<211> 7
<212> PRT
<213> Mus musculus

<400> 163
Tyr Thr Ser Ser Leu His Ser
1 5

<210> 164
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 164
Gln Gln Tyr Ser Lys Leu Pro Phe Thr
1 5

<210> 165
<211> 117
<212> PRT
<213> Mus musculus

<400> 165
Glu Val Lys Leu Val Glu Ser Glu Gly Gly Leu Val Gln Pro Gly Ser
1 5 10 15

Ser Met Lys Leu Ser Cys Thr Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Tyr Met Ala Trp Val Arg Gln Val Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

Ala Asn Ile Asn Tyr Asp Gly Ser Asn Thr Ser Tyr Leu Asp Ser Leu
50 55 60

Lys Ser Arg Phe Ile Ile Ser Arg Asp Asn Ala Lys Asn Ile Leu Tyr
65 70 75 80

Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Thr Tyr Tyr Cys
85 90 95

Ala Arg Glu Lys Phe Ala Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser
100 105 110

Val Thr Val Ser Ser
115

<210> 166
<211> 7
<212> PRT
<213> Mus musculus

<400> 166
Gly Phe Thr Phe Ser Asp Tyr
1 5

<210> 167
<211> 6
<212> PRT
<213> Mus musculus

<400> 167
Asn Tyr Asp Gly Ser Asn
1 5

<210> 168
<211> 8
<212> PRT
<213> Mus musculus

<400> 168
Glu Lys Phe Ala Ala Met Asp Tyr
1 5

<210> 169
<211> 108
<212> PRT
<213> Mus musculus

<400> 169
Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Phe Gly
1 5 10 15

Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Asn Ala
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly His Ser Pro Lys Leu Leu Ile
35 40 45

Phe Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala
65 70 75 80

Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Ser Thr Pro Trp
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> 170
<211> 11
<212> PRT
<213> Mus musculus

<400> 170
Lys Ala Ser Gln Asp Val Ser Asn Ala Leu Ala
1 5 10

<210> 171
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
variable light chain CDR peptide

<400> 171
Ser Ala Ser Tyr Arg Tyr Thr
1 5

<210> 172
<211> 9
<212> PRT
<213> Mus musculus

<400> 172

Gln Gln His Tyr Ser Thr Pro Trp Thr
1 5

<210> 173

<211> 121

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 173

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr Arg His
20 25 30

Thr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Arg Ile Ser Pro Ala Asn Gly Asn Thr Asn Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Trp Ile Gly Ser Arg Glu Leu Tyr Ile Met Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 174

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 174

Gly Phe Thr Phe Thr Arg His Thr Ile His
1 5 10

<210> 175

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 175

Arg Ile Ser Pro Ala Asn Gly Asn Thr Asn Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 176

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 176

Trp Ile Gly Ser Arg Glu Leu Tyr Ile Met Asp Tyr
1 5 10

<210> 177

<211> 108

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 177

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser Thr Ala
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Arg Ile Gln Pro
85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 178
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 178
Arg Ala Ser Gln Asp Val Ser Thr Ala Val Ala
1 5 10

<210> 179
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
peptide

<400> 179
Ser Ala Ser Phe Leu Tyr Ser
1 5

<210> 180
<211> 9
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 180

Gln Gln Ser Tyr Arg Ile Gln Pro Thr
1 5

<210> 181

<211> 121

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 181

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Thr
20 25 30

Ala Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Arg Ile Ser Pro Ala Asn Gly Asn Thr Asn Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Trp Ile Gly Ser Arg Glu Leu Tyr Ile Met Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 182

<211> 10

<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 182

Gly	Phe	Thr	Phe	Ser	Ser	Thr	Ala	Ile	His
1				5				10	

<210> 183

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 183

Arg	Ile	Ser	Pro	Ala	Asn	Gly	Asn	Thr	Asn	Tyr	Ala	Asp	Ser	Val	Lys
1				5				10					15		

Gly

<210> 184

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 184

Trp	Ile	Gly	Ser	Arg	Glu	Leu	Tyr	Ile	Met	Asp	Tyr
1				5				10			

<210> 185

<211> 108

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 185

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Ser Thr Ala
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Pro Ala Leu His
85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg
100 105

<210> 186

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 186

Arg Ala Ser Gln Asp Val Ser Thr Ala Val Ala
1 5 10

<210> 187

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 187

Ser Ala Ser Phe Leu Tyr Ser
1 5

<210> 188
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 188
Gln Gln Ser Tyr Pro Ala Leu His Thr
1 5

<210> 189
<211> 125
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 189
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Pro Phe Ser Lys Leu
20 25 30

Gly Met Val Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Thr Ile Ser Ser Gly Gly Gly Tyr Thr Tyr Tyr Pro Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Gly Ile Ser Phe Gln Gly Gly Thr Tyr Thr Tyr Val Met
100 105 110

Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 190

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 190

Gly Phe Pro Phe Ser Lys Leu Gly Met Val
1 5 10

<210> 191

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 191

Thr Ile Ser Ser Gly Gly Gly Tyr Thr Tyr Tyr Pro Asp Ser Val Lys
1 5 10 15

Gly

<210> 192

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 192

Glu Gly Ile Ser Phe Gln Gly Gly Thr Tyr Thr Tyr Val Met Asp Tyr
1 5 10 15

<210> 193

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
polypeptide

<400> 193

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu His Arg
20 25 30

Asn Gly Ile Thr Tyr Ser Tyr Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Gln Leu Ser Asn Leu Ala Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Tyr Gln Asn
85 90 95

Leu Glu Leu Pro Leu Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
100 105 110

<210> 194

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide

<400> 194

Arg Ser Ser Lys Ser Leu Leu His Arg Asn Gly Ile Thr Tyr Ser Tyr
1 5 10 15

<210> 195

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 195

Gln Leu Ser Asn Leu Ala Ser
1 5

<210> 196

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 196

Tyr Gln Asn Leu Glu Leu Pro Leu Thr
1 5

<210> 197

<211> 2076

<212> DNA

<213> Homo sapiens

<400> 197

atgggcaccg tcagctccag gcggtcctgg tggccgctgc cactgctgct gctgctgctg	60
ctgctcctgg gtcccgcggg cgcccgtgcg caggaggacg aggacggcga ctacgaggag	120
ctggtgctag ccttgcgttc cgaggaggac ggcctggccg aagcaccgga gcacggaacc	180
acagccacct tccaccgctg cgccaaggat ccgtggagggt tgcctggcac ctacgtgggtg	240
gtgctgaagg aggagaccca cctctcgag tcagagcgca ctgcccgccg cctgcaggcc	300
caggctgccc gccggggata cctcaccaag atcctgcatg tcttccatgg ccttcttcct	360
ggcttcctgg tgaagatgag tggcgacctg ctggagctgg ccttgaagtt gccccatgtc	420
gactacatcg aggaggactc ctctgtcttt gccagagca tcccgtggaa cctggagcgg	480
attaccctc cacggtaccg ggcggatgaa taccagcccc ccgacggagg cagcctggtg	540
gaggtgtatc tcctagacac cagcatacag agtgaccacc gggaaatcga gggcagggtc	600
atggtcaccg acttcgagaa tgtgcccgag gaggacggga cccgcttcca cagacaggcc	660
agcaagtgtg acagtcatgg caccacctg gcaggggtgg tcagcggccg ggatgccggc	720
gtggccaagg gtgccagcat gcgcagcctg cgcgtgctca actgccaagg gaagggcacg	780

gtagcgga ccctcatagg cctggagttt attcgga aaa gccagctggg ccagcctgtg	840
gggccactgg tgggtgctgct gcccctggcg ggtgggtaca gccgcgtcct caacgccgcc	900
tgccagcgcc tggcgagggc tggggctctg ctgggtcaccg ctgccggcaa cttccgggac	960
gatgcctgcc tctactcccc agcctcagct cccgaggtca tcacagttgg ggccaccaat	1020
gccaagacc agccggtgac cctggggact ttggggacca actttggccg ctgtgtggac	1080
ctctttgccc caggggagga catcattggg gcctccagcg actgcagcac ctgctttgtg	1140
tcacagagtg ggacatcaca ggctgctgcc cacgtggctg gcattgcagc catgatgctg	1200
tctgccgagc cggagctcac cctggccgag ttgaggcaga gactgatcca cttctctgcc	1260
aaagatgtca tcaatgaggc ctggttcctt gaggaccagc gggtactgac ccccaacctg	1320
gtggccgccc tgccccccag caccatggg gcaggttggc agctgttttg caggactgta	1380
tggtcagcac actcggggcc tacacggatg gccacagccg tcgcccgtg cgccccagat	1440
gaggagctgc tgagctgctc cagtttctcc aggagtggga agcggcgggg cgagcgcatg	1500
gaggcccaag ggggcaagct ggtctgccgg gccacaacg cttttggggg tgagggtgtc	1560
tacgccattg ccaggtgctg cctgctaccc caggccaact gcagcgcca cacagctcca	1620
ccagctgagg ccagcatggg gaccctgtgc cactgccacc aacagggccg cgtcctcaca	1680
ggctgcagct cccactggga ggtggaggac cttggcacc acaagccgcc tgtgctgagg	1740
ccacgaggtc agcccaacca gtgcgtgggc cacagggagg ccagcatcca cgcttcctgc	1800
tgccatgccc caggtctgga atgcaaagtc aaggagcatg gaatcccggc ccctcaggag	1860
caggtgaccg tggcctgcga ggagggtg accctgactg gctgcagtgc cctccctggg	1920
acctcccacg tcctgggggc ctacgccgta gacaacacgt gtgtagtcag gagccgggac	1980
gtcagcacta caggcagcac cagcgaagg gccgtgacag ccgttgccat ctgctgccgg	2040
agccggcacc tggcgcaggc ctcccaggag ctccag	2076

<210> 198

<211> 692

<212> PRT

<213> Homo sapiens

<400> 198

Met	Gly	Thr	Val	Ser	Ser	Arg	Arg	Ser	Trp	Trp	Pro	Leu	Pro	Leu	Leu
1				5				10				15			

Leu Leu Leu Leu Leu Leu Leu Gly Pro Ala Gly Ala Arg Ala Gln Glu
20 25 30

Asp Glu Asp Gly Asp Tyr Glu Glu Leu Val Leu Ala Leu Arg Ser Glu
35 40 45

Glu Asp Gly Leu Ala Glu Ala Pro Glu His Gly Thr Thr Ala Thr Phe
50 55 60

His Arg Cys Ala Lys Asp Pro Trp Arg Leu Pro Gly Thr Tyr Val Val
65 70 75 80

Val Leu Lys Glu Glu Thr His Leu Ser Gln Ser Glu Arg Thr Ala Arg
85 90 95

Arg Leu Gln Ala Gln Ala Ala Arg Arg Gly Tyr Leu Thr Lys Ile Leu
100 105 110

His Val Phe His Gly Leu Leu Pro Gly Phe Leu Val Lys Met Ser Gly
115 120 125

Asp Leu Leu Glu Leu Ala Leu Lys Leu Pro His Val Asp Tyr Ile Glu
130 135 140

Glu Asp Ser Ser Val Phe Ala Gln Ser Ile Pro Trp Asn Leu Glu Arg
145 150 155 160

Ile Thr Pro Pro Arg Tyr Arg Ala Asp Glu Tyr Gln Pro Pro Asp Gly
165 170 175

Gly Ser Leu Val Glu Val Tyr Leu Leu Asp Thr Ser Ile Gln Ser Asp
180 185 190

His Arg Glu Ile Glu Gly Arg Val Met Val Thr Asp Phe Glu Asn Val
195 200 205

Pro Glu Glu Asp Gly Thr Arg Phe His Arg Gln Ala Ser Lys Cys Asp
210 215 220

Ser His Gly Thr His Leu Ala Gly Val Val Ser Gly Arg Asp Ala Gly

225		230		235		240									
Val	Ala	Lys	Gly	Ala	Ser	Met	Arg	Ser	Leu	Arg	Val	Leu	Asn	Cys	Gln
				245					250					255	
Gly	Lys	Gly	Thr	Val	Ser	Gly	Thr	Leu	Ile	Gly	Leu	Glu	Phe	Ile	Arg
			260					265					270		
Lys	Ser	Gln	Leu	Val	Gln	Pro	Val	Gly	Pro	Leu	Val	Val	Leu	Leu	Pro
		275					280					285			
Leu	Ala	Gly	Gly	Tyr	Ser	Arg	Val	Leu	Asn	Ala	Ala	Cys	Gln	Arg	Leu
	290					295					300				
Ala	Arg	Ala	Gly	Val	Val	Leu	Val	Thr	Ala	Ala	Gly	Asn	Phe	Arg	Asp
305					310					315					320
Asp	Ala	Cys	Leu	Tyr	Ser	Pro	Ala	Ser	Ala	Pro	Glu	Val	Ile	Thr	Val
			325						330					335	
Gly	Ala	Thr	Asn	Ala	Gln	Asp	Gln	Pro	Val	Thr	Leu	Gly	Thr	Leu	Gly
			340					345					350		
Thr	Asn	Phe	Gly	Arg	Cys	Val	Asp	Leu	Phe	Ala	Pro	Gly	Glu	Asp	Ile
	355						360					365			
Ile	Gly	Ala	Ser	Ser	Asp	Cys	Ser	Thr	Cys	Phe	Val	Ser	Gln	Ser	Gly
	370					375					380				
Thr	Ser	Gln	Ala	Ala	Ala	His	Val	Ala	Gly	Ile	Ala	Ala	Met	Met	Leu
385					390					395					400
Ser	Ala	Glu	Pro	Glu	Leu	Thr	Leu	Ala	Glu	Leu	Arg	Gln	Arg	Leu	Ile
				405					410					415	
His	Phe	Ser	Ala	Lys	Asp	Val	Ile	Asn	Glu	Ala	Trp	Phe	Pro	Glu	Asp
			420					425					430		
Gln	Arg	Val	Leu	Thr	Pro	Asn	Leu	Val	Ala	Ala	Leu	Pro	Pro	Ser	Thr
	435						440					445			

His Gly Ala Gly Trp Gln Leu Phe Cys Arg Thr Val Trp Ser Ala His
450 455 460

Ser Gly Pro Thr Arg Met Ala Thr Ala Val Ala Arg Cys Ala Pro Asp
465 470 475 480

Glu Glu Leu Leu Ser Cys Ser Ser Phe Ser Arg Ser Gly Lys Arg Arg
485 490 495

Gly Glu Arg Met Glu Ala Gln Gly Gly Lys Leu Val Cys Arg Ala His
500 505 510

Asn Ala Phe Gly Gly Glu Gly Val Tyr Ala Ile Ala Arg Cys Cys Leu
515 520 525

Leu Pro Gln Ala Asn Cys Ser Val His Thr Ala Pro Pro Ala Glu Ala
530 535 540

Ser Met Gly Thr Arg Val His Cys His Gln Gln Gly His Val Leu Thr
545 550 555 560

Gly Cys Ser Ser His Trp Glu Val Glu Asp Leu Gly Thr His Lys Pro
565 570 575

Pro Val Leu Arg Pro Arg Gly Gln Pro Asn Gln Cys Val Gly His Arg
580 585 590

Glu Ala Ser Ile His Ala Ser Cys Cys His Ala Pro Gly Leu Glu Cys
595 600 605

Lys Val Lys Glu His Gly Ile Pro Ala Pro Gln Glu Gln Val Thr Val
610 615 620

Ala Cys Glu Glu Gly Trp Thr Leu Thr Gly Cys Ser Ala Leu Pro Gly
625 630 635 640

Thr Ser His Val Leu Gly Ala Tyr Ala Val Asp Asn Thr Cys Val Val
645 650 655

Arg Ser Arg Asp Val Ser Thr Thr Gly Ser Thr Ser Glu Gly Ala Val

660

665

670

Thr Ala Val Ala Ile Cys Cys Arg Ser Arg His Leu Ala Gln Ala Ser
675 680 685

Gln Glu Leu Gln
690