

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Method and process for producing youthful-appearing, small-pored, and smooth skin

(51)⁶ International Patent Classification(s)
A61B 18/20 (2006.01) 20060101ALI2006112
A61B 17/00 (2006.01) 5BMEP **A61B**
A61B 17/50 (2006.01) 18/00
A61B 18/00 (2006.01) 20060101ALI2005111
A61K 31/203 0BMEP **A61K**
(2006.01) 31/203
A61K 31/74 (2006.01) 20060101ALI2005111
A61N 5/06 (2006.01) 0BMEP **A61K**
A61B 18/20 31/74
20060101AFI2005122 20060101ALI2006112
0BMJP **A61B** 5BMEP **A61N**
17/00 5/06
20060101ALI2005122 20060101ALI2005122
0BMJP **A61B** 0BMJP
17/50 PCT/US02/27570

(21) Application No: 2002332731 (22) Application Date: 2002.08.28

(87) WIPO No: WO03/059144

(30) Priority Data

(31) Number (32) Date (33) Country
10/047,335 2002.01.14 US

(43) Publication Date: 2003.07.30

(43) Publication Journal Date: 2003.09.04

(71) Applicant(s)
Diana Parsons

(72) Inventor(s)
Parsons, Diana J

(74) Agent/Attorney
FB Rice & Co, Level 23 200 Queen Street, Melbourne, VIC, 3000

(56) Related Art
Kye, Y.C. Resurfacing of pitted facial scars with a pulsed Er:YAG laser. Dermatological Surgery, October 1997, Vol. 23, No.10, pp. 880-883.
US 5423803
Ho et al., laser resurfacing in pigmented skin, Dermatological surgery, December 1995, Vol. 21, No.12, pp. 1035-1037.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
24 July 2003 (24.07.2003)

PCT

(10) International Publication Number
WO 2003/059144 A3

(51) International Patent Classification⁷: A61B 17/00, 17/50, A61K 31/74, 7/42

(21) International Application Number: PCT/US2002/027570

(22) International Filing Date: 28 August 2002 (28.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

10/047,335 14 January 2002 (14.01.2002) US

Published:

— with international search report

(71) Applicant and

(72) Inventor: PARSONS, Diana, J. [US/US]; 3031 Telegraph #130, Berkley, CA 94805 (US).

(88) Date of publication of the international search report:
22 January 2004

(74) Agent: WALSH, Gerald, M.; Reg. No. 34,946. Bush Intellectual Property Law Group LLC, P.O. Box 381146, Birmingham, AL 35238 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2003/059144 A3

(54) Title: METHOD AND PROCESS FOR PRODUCING YOUTHFUL-APPEARING, SMALL-PORED, AND SMOOTH SKIN

(57) Abstract: The method and process for producing youthful appearing, small-pored, and smooth skin comprises pretreating the skin topically with retinoic acid, applying a contaminant in oil to the skin, and removing the contaminant by exploding or rupturing it with laser light. When treating the face, removing the contaminant requires only about 4 minutes. The laser light itself does not injure the skin but the rupture of the contaminant produces thermal injury in the high dermis leaving the epidermis intact. Repeated intermittent laser treatments and applications of topical retinoic acid produces the desired results in the skin indefinitely as long as the treatments and applications are continued.

**METHOD AND PROCESS FOR PRODUCING
YOUTHFUL-APPEARING, SMALL-PORED,
AND SMOOTH SKIN**

5

BACKGROUND OF INVENTION

1. Field of the Invention

10 The present invention relates generally to laser treatment of the skin and, more particularly, to a laser treatment inducing a chronic wound in the high dermis, leaving the epidermis intact, with topical pretreatment and post-treatment of the skin with retinoic acid.

15 2. Technical Background

Laser light treatment of the skin is used to rejuvenate the skin, remove pigments and hair, and treat infection. In the field of dermatology and plastic surgery the use of lasers is principally based upon two types of mechanisms: a thermal effect where the 20 laser light energy is converted into heat energy, or a mechanical effect where the laser light energy is converted into shockwaves in the skin.

Most laser treatments use thermal-based laser therapy to vaporize the superficial wrinkles and top layers of the skin so that new collagen and skin can be naturally 25 provided in a healing response. The procedure is performed under either local or general anesthetic, takes a few hours to complete, and recovery takes one to two weeks. During the first week patients may suffer a sensation of intense heat on the skin. Severe burns can occur and result in permanent scarring. Bacterial and yeast infection have been reported and can also lead to scarring. Additional potential complications include 30 changes in pigmentation and herpes infection.

Retinoic acid (Retin-A, tretinoin) has also been used as a topical dermal treatment to improve the appearance and texture of the skin. Retinoic acid thins the stratum corneum, increases the thickness of the epidermis, and increases the production of

collagen in the dermis. However, retinoic acid causes skin redness and sensitivity to the sun. Repeated use can cause a loss of pigment, painful irritation, dryness, swelling of the skin, and contact dermatitis.

5 In an effort to overcome these drawbacks of laser therapy and to avoid the unwanted side effects of retinoic acid, *Tankovich et al.* (U.S. Patent No. 6,036,684) developed a laser method of photomechanical activation in the skin using a Q switched Nd:YAG laser with a wavelength of 1064nm and an exposure of 2.5 J/cm². Exposure of the skin to this type of laser treatment by itself has no effect on the skin because the skin
10 has no inherent target at the 1064nm wavelength. However, when an activating solution of graphite or carbon particles suspended in baby oil is applied to the skin, these particles become the target of this 1064nm wavelength, exploding when exposed to the laser light. Prior to application of the laser, the particles are forced into the skin, below the surface of the stratum corneum using ultrasound. Thereafter, the explosion of the carbon particles
15 by the laser light produces a localized mechanical injury in the hair follicles and pores of the skin. There is no significant injury to the skin tissue because the laser energy which is not absorbed in the carbon is harmlessly dissipated in the skin. The low fluence of 2.5 J/cm² leaves the epidermis intact and the typical adverse effects of laser treatment do not occur.

20

Although the photomechanical laser process of *Tankovich* is considerably safer than the standard photothermal laser treatments and leaves the epidermis intact, this photomechanical laser process is relatively ineffective in treating the skin because the extent of injury in the high dermis resulting from the exploding carbon particles in the
25 epidermis is insufficient to induce adequate collagen deposition during wound healing. What is needed, therefore, is means for enhancing the collagen deposition in the high dermis during the wound healing process.

30

SUMMARY OF THE INVENTION

The present invention provides a process and method for producing skin rejuvenation and therapy by pretreating the skin with retinoic acid, producing a chronic thermal injury wound in the high dermis using photomechanical laser therapy, maintaining skin rejuvenation and therapy chronically by repeating the thermal injury in the high dermis at least once per year, and by topically applying retinoic acid bi-weekly. A Q-switched Nd:YAG laser with a wave length of 1064 nm is used at 2.5 J/cm² to vaporize a topical activating solution of carbon particles suspended in baby oil. A single treatment of the face can be completed in 4 minutes without anesthesia, with no need for a period of recovery, and at a relatively low cost.

According to the invention there is also provided a method for treating the skin, comprising the steps of:

- 15 a) pretreating the skin with a topical collagen-promoting agent;
- b) producing a wound in the high dermis of the skin;
- c) treating the skin chronically with said topical collagen-promoting agent after producing said wound in the high dermis; and
- d) maintaining the high dermis in a chronically wounded condition,

20 wherein the step of producing a wound in the high dermis is performed by exploding a contaminant on the surface of the skin with laser light.

An embodiment provides a chronic wound in the high dermis with no damage to the epidermis.

25 Another embodiment provides enhancement of laser-induced collagen deposition in the high dermis with retinoic acid.

Another embodiment provides chronic life-time rejuvenation and therapy of the skin.

30 Another embodiment provides the removal and prevention of acne by suppression of sebaceous glands and reduction of skin pore size.

Another embodiment provides laser therapy at a low energy level.

35 Another embodiment provides the use of retinoic acid without side effects.

Another embodiment provides therapeutic removal of infections of the skin.

Another embodiment provides the chronic rejuvenation and therapy of the skin
5 at a relatively low cost.

Another embodiment provides a single treatment of the skin of the face can be
completed within four minutes.

10 Throughout this specification the word "comprise", or variations such as
"comprises" or "comprising", will be understood to imply the inclusion of a stated
element, integer or step, or group of elements, integers or steps, but not the exclusion of
any other element, integer or step, or group of elements, integers or steps.

15 Any discussion of documents, acts, materials, devices, articles or the like which
has been included in the present specification is solely for the purpose of providing a
context for the present invention. It is not to be taken as an admission that any or all of
these matters form part of the prior art base or were common general knowledge in the
field relevant to the present invention as it existed before the priority date of each claim
20 of this application.

BRIEF DESCRIPTION OF THE DRAWINGS

25 **Fig. 1** describes the method of the present invention.

Fig. 2 illustrates a cross sectional view of the skin and the role of heat and oil in
the method of the present invention.

30 **Fig. 3** describes the process of the present invention.

4A

Fig. 4 shows the effects of the laser method of the present invention on the restoration of the skin of a patient suffering from acne vulgaris.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

5

While the following description details the preferred embodiments of the present invention, it is to be understood that the invention is not limited in its application to the details of construction and arrangement of the parts illustrated in the accompanying drawings, since the invention is capable of other embodiments and of being practiced in 10 various ways.

Wound healing in the skin occurs over a year's time after an initial wounding, regardless of the means by which the wounding is induced. In the first five days the

15

wound is comprised of inflammatory cells and new blood vessels. Then an immature form of collagen is laid down parallel to the skin surface. For six months the body shifts this collagen around, trying to identify the strongest repair. Around six months after the initial wounding, this immature collagen is replaced with mature collagen protein oriented perpendicular to the skin surface. Three months into this process, or months ten, eleven, and twelve from the initial wounding, these protein strands cross-link. This effect is visible on the skin's surface as an apparent shrinkage of the size of the scar to one third its original size.

10 Twelve months from the initial injury, the wound becomes quiescent. However, any additional insult to the wound site during the first six months of repair causes a signal for a stronger permanent repair. Any additional insult to the wound site during the second six months results in additional mature collagen in the wound. As long as some injury is repeated before twelve months has elapsed, the site becomes a chronic wound
15 producing more collagen, undergoing more cross-linking, and more total surface area shrinkage.

20 The method and process of the present invention described herein outlines a series of treatments using the *Tankovich* laser and activating solution to maximize the initial wounding signal during the first six months of application. The *Tankovich* processes disclosed in U.S. Patent No. 5,423,803 and U.S. Patent No. 6,036,684 are incorporated herein by reference. However, in the present invention the activating solution does not need to be forced into the spaces of the skin with ultrasound as required by the *Tankovich* process. Beginning at the six month from the initial wounding, a series of booster laser
25 treatments is given until the desired skin resurfacing end point is reached. At a minimum, one booster laser treatment is given before twelve months has elapsed to keep the wound active. If the wound is not kept active, it will be necessary to repeat the initial wounding series of the first six months in order to continue the resurfacing process chronically.

The first step of the method of the present invention is shown in **Fig. 1, Step 10**, which is a topical pretreatment of the skin with a collagen inducing agent and an angiogenesis inducing agent, preferably, retinoic acid, and in particular its nongeneric form, such as Retin-A (tretinoin). Retinoic acid returns the architecture of the skin back 5 to its more youthful form. As skin ages, the hill and valley nature of the dermal-epidermal junction 21 flattens (see **Fig. 2**). Topical retinoic acid restores the up and down pattern at the junction 21. In youth, the basal cells which originate at the dermal-epidermal junction 21 create the layers of the epidermis every three weeks. With aging, this slows to four to six weeks. Topical retinoic acid on the skin returns the process to 10 every three weeks. The top most layer of epidermal cells, the stratum corneum, is comprised of dead cells held together by a cell glue. With age, this layer thickens making the skin less fresh-appearing and making the pores larger. Topical retinoic acid directly attacks the cell glue, thinning the stratum corneum and making the pores smaller by reducing the amount of debris in them. In my studies I have discovered that 15 0.05% to 1% concentration of retinoic acid in a topical formulation has the property of inducing new collagen and new blood vessel formation in the high dermis. The pre-treatment period is one to four weeks, preferably two weeks with retinoic acid being applied twice a week with at least two days in between each application. The laser therapy of the present invention will not be effective without the pre-treatment 20 application of topical retinoic acid.

The second step **11** in the method of the present invention shown in **Fig. 1** is the creation of a chronic wound in the high dermis. This is produced by a photomechanical laser treatment wherein the laser light does not interact directly with the skin but 25 instead interacts with a contaminant in the skin. The contaminant has the properties of absorbing the laser light and exploding. The preferred contaminant is carbon or graphite particles in oil 20 which is applied to the skin (see **Fig. 2**). Once the contaminant or activating solution is applied to the skin, the laser treatment can begin. The energy from the laser is adjusted to be just sufficient to cause the particles to 30 explode. As the particles explode, the mineral oil 20 penetrates into the epidermis producing hydration of the epidermis by retarding the evaporation of water

(see Fig. 2). The heat from the explosion of the contaminant particles will induce a photothermal injury in the rete peg area of the high dermis 22 initiating a normal wound healing process. In order to produce a sufficient degree of injury to the wound, the laser treatment is produced several times over a six month period, preferably six times over a 5 six month period. During this first six months the retinoic acid is applied topically twice per week as described above.

The build up of new collagen in the high dermis 22 in response to the laser treatment and retinoic acid treatment thickens the skin and increases its turgor resulting in 10 smaller skin pores. Broken blood vessels, angioma, and ice pick scars or suture marks are also minimized, being crowded out by the new collagen. Easy bruising of the skin is lessened by the collagen build up. Lips become redder, fine lines are minimized, and dark circles around the eyes are minimized. As the new collagen cross-links major 15 wrinkle lines, such as nasal labial folds and perioral and periocular lines (caused by the underlying attachment of muscles to the deep dermis), marionette lines, sunken-in corners of the mouth, and forehead, temporal, and glabellar lines, appear to melt away into the new skin. The new skin has a glowing quality resulting from the pulsating nature of the laser injury and the high content of mineral oil in the activating solution. Retinoic acid and laser treatment work together to minimize bound differential pigmentation 20 caused by sun damage and hormones (myelasma). Redness from inflammation, as in acne vulgaris, acne rosacea, or maturing scars or striae is also reduced.

Because of the closeness of the sebaceous glands to the skin surface and because 25 of the new collagen formation which reduces the size of the skin pores, the method of process of the present invention is useful in the treatment of sebaceous gland disorders such as acne. The secretion from the sebaceous glands is reduced and the smaller skin pore size prevents bacteria from entering the skin pores. In addition, heat generated from the explosion of the contaminant particles is sufficient to kill bacteria, fungi, and viruses on the skin's surface or in the high dermis. This also allows treatment of conditions such 30 as, for example, plantar warts, herpes cold sores, athletes foot, psoriasis, and excema.

The third step 12 in the method of the present invention, depicted in **Fig. 1**, is the long term continued chronic maintenance of the high dermis in a wounded condition in order to produce a sustained rejuvenation of the skin and treatment of skin conditions. This is accomplished by application of the laser treatment of the present invention at least 5 once a year and preferably twice, along with the continued application of topical retinoic acid to the skin twice per week as described above.

The process of the present invention for treating the skin is shown in **Fig. 3**. In the first step 40, pre-treatment is initiated with topical retinoic acid applied to the skin at 10 least twice per week. The concentration of retinoic acid in the topical formulation is between about 0.05% to 1%, preferably about 0.1%. The pretreatment period is, preferably, two weeks. In the next step 41, an activating solution of graphite in baby oil is applied to the skin. The graphite-oil ratio may range from about 1:1 to 1:9, preferably 1:4, *i.e.*, about 20 percent graphite suspended in about 80 percent oil by weight. In the 15 next steps 42 and 43 a laser beam is scanned over the area treated with the activating solution so as to clean substantially all of the mixture from the skin surface by exploding or fracturing the carbon or graphite particles in the oil. This scanning process takes from about 2 to 10 minutes to complete on the face, usually about 4 minutes.

20 It is preferred to use a Q-switched neodymium:yttrium-aluminum-garnet (Nd:YAG) laser for the process of the present invention. The wave length of the radiation may range from about 800nm to 2000nm, preferably about 1064nm. The frequency of the pulses from the laser range from about 1 to 20 per second, preferably, about 10 per second. The duration of each pulse ranges from about .001 to 1 25 microsecond, preferably about .01 microsecond. The fluence or exposure of the skin treated with the activating solution ranges from about 1 to 3 J/cm², preferably, about 2.5 J/cm².

In the next step 44 the laser process is repeated several times to create a chronic 30 wound in the high dermis 22 (see **Fig. 2**). The process may be repeated 2 to 12 times within a six month period, preferably six times. During this time retinoic acid is applied

to the skin 1 to 4 times per week, preferably 2 times per week, with 2 to 3 days between applications. The application of retinoic acid may be varied in an equivalent manner on a monthly basis, *i.e.*, 4 to 16 applications per month, preferably 8 applications per month, where the applications may all be applied within one or two weeks of a given month, but, 5 preferably, applied each week of a month.

In the next step 45 the laser process is repeated at least once in the next six months and thereafter at least once per year, but, preferably twice per year. The yearly treatments will maintain a chronic wound in the high dermis. At the same time, topical 10 retinoic acid is applied weekly to the skin as described above. The continued combination of intermittent laser therapy and topical retinoic acid will maintain the rejuvenation and therapeutic results in the skin indefinitely.

Fig. 4 shows, by way of example, the effects of the laser method of the present 15 invention on the skin of a patient suffering from acne vulgaris. The before picture presents the patient's skin prior to treatment and the after picture shows the patient's skin after the first six months of treatment of the skin with the method and process of the present invention. The lesions and scars are substantially reduced in severity and the process of acne formation has been completely prevented. Furthermore, this has been 20 accomplished leaving the epidermis intact and normal. Continued chronic treatment with the method of the present invention is expected to provide further improvement in the appearance of the skin.

The foregoing description has been limited to specific embodiments of this 25 invention. It will be apparent, however, that variations and modifications may be made by those skilled in the art to the disclosed embodiments of the invention, with the attainment of some or all of its advantages and without departing from the spirit and scope of the present invention. For example, agents other than retinoic acid which induce 30 collagen and blood vessel growth may be used in place of retinoic acid. Any laser system capable of exploding or rupturing a contaminant in the skin without injuring the skin

directly can be used in the present invention. The activating solution can be made with any suitable oil with any suitable contaminant.

It will be understood that various changes in the details, materials, and arrangements of the parts which have been described and illustrated above in order to explain the nature of this invention may be made by those skilled in the art without departing from the principle and scope of the invention as recited in the following claims.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for treating the skin, comprising the steps of :
 - a) pretreating the skin with a topical collagen-promoting agent;
 - b) producing a wound in the high dermis of the skin;
 - c) treating the skin chronically with said topical collagen-promoting agent after producing said wound in the high dermis; and
 - d) maintaining the high dermis in a chronically wounded condition, wherein the step of producing a wound in the high dermis is performed by exploding a contaminant on the surface of the skin with laser light.
2. The method of claim 1 further comprising the steps of pretreating and chronically treating the skin with a blood vessel promoting agent.
- 15 3. The method of either claim 1 or claim 2 wherein said collagen-promoting agent is retinoic acid.
4. The method of claim 2 wherein said blood vessel promoting agent is retinoic acid.
- 20
5. The method of any one of claims 1 - 4 wherein said production of a chronic wound in the high dermis leaves the epidermis intact.
- 25 6. The method of any one of claims 1 - 5 wherein the step of producing a chronic wound in the high dermis is performed by exploding a contaminant on the skin with laser light repeatedly within a six month period of said pretreating step.
7. The method of any one of claims 1 - 6 wherein the step of maintaining a chronic wound in the high dermis is performed by exploding a contaminant on the surface of the skin with laser light at least once every twelve months.
- 30
8. The method of any one of claims 1 - 7 wherein the steps of producing and maintaining a chronic wound in the high dermis are performed by exploding a contaminant on the skin with a laser light having a wave length of from about 800nM to 2000nM, a pulse frequency of from about 1 to 20 per second, a pulse
- 35

duration of from about 0.001 to 1 microsecond, and a fluence of from about 1 to 3 J/cm².

9. The method of claim 8 wherein said laser light has a wave length of about 5 1064nm, a pulse frequency of about 10 per second, a pulse duration of about .01 microsecond, and a fluence of about 2.5 J/cm².
10. The method of claim 3 wherein said retinoic acid is applied to the skin 4 to 16 times per month.
11. The method of claim 10 wherein said retinoic acid is applied to the skin 1 to 4 10 times per week.
12. The method of claim 11 wherein said retinoic acid is applied to the skin 2 times 15 per week.
13. The method of claim 3 wherein said retinoic acid is applied topically at a concentration of from about 0.05% to 1%.
- 20 14. The method of claim 13 wherein said retinoic acid is applied topically at a concentration of about 0.1%.
15. The method of claim 3 wherein the pretreating step is performed for two weeks.
- 25 16. The method of any one of claims 1 - 15 wherein producing a wound in the high dermis is completed in about two to ten minutes.
17. The method of claim 16 wherein producing a wound in the high dermis is completed in about four minutes.
- 30 19. The method of any one of claims 1 - 18 wherein said method for treating the skin is used to rejuvenate the skin and to remove broken blood vessels, angiomas, scars, lines, wrinkles, pigmentation, and redness in the skin.

20. The method of any one of claims 1 - 18 wherein said method for treating the skin is used to treat acne, warts, cold sores, eczema, psoriasis, and infections of the skin.
- 5 21. A method for treating the skin according to claim 1, substantially as herein exemplified with reference to any one or more of the accompanying Figures.




Fig. 1

SUBSTITUTE SHEET (RULE 26)

2/4




Fig. 2

SUBSTITUTE SHEET (RULE 26)





Fig. 3

SUBSTITUTE SHEET (RULE 26)

4/4



Before

After

Fig. 4

SUBSTITUTE SHEET (RULE 26)