(12) STANDARD PATENT (11) Application No. **AU 2012249493 B2**
(19) AUSTRALIAN PATENT OFFICE

(54) Title
Tolerogenic synthetic nanocarriers for inducing regulatory B cells

(51) International Patent Classification(s)
A61K 39/38 (2006.01) ***A61K 31/445*** (2006.01)
A61K 9/14 (2006.01) ***A61K 47/30*** (2006.01)
A61K 9/16 (2006.01)

(21) Application No: **2012249493** (22) Date of Filing: **2012.04.27**

(87) WIPO No: **WO12/149301**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/531,204	**2011.09.06**	**US**
61/531,209	**2011.09.06**	**US**
61/531,164	**2011.09.06**	**US**
61/531,180	**2011.09.06**	**US**
61/531,153	**2011.09.06**	**US**
61/513,514	**2011.07.29**	**US**
61/531,194	**2011.09.06**	**US**
61/531,215	**2011.09.06**	**US**
61/531,175	**2011.09.06**	**US**
61/531,147	**2011.09.06**	**US**
61/480,946	**2011.04.29**	**US**
61/531,168	**2011.09.06**	**US**

(43) Publication Date: **2012.11.01**
(44) Accepted Journal Date: **2017.06.15**

(71) Applicant(s)
Selecta Biosciences, Inc.

(72) Inventor(s)
Maldonado, Roberto A.

(74) Agent / Attorney
Pizzeys Patent and Trade Mark Attorneys Pty Ltd, PO Box 291, WODEN, ACT, 2606, AU

(56) Related Art
WO 2009/131712 A2
WO 2007/087341 A2
US 2010/0129439 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
1 November 2012 (01.11.2012)

WIPO | PCT

(10) International Publication Number
WO 2012/149301 A3

(51) International Patent Classification:
A61K 39/38 (2006.01) *A61K 9/14* (2006.01)
A61K 31/445 (2006.01) *A61K 47/48* (2006.01)
A61K 9/16 (2006.01) *A61K 47/30* (2006.01)

(21) International Application Number:
PCT/US2012/035431

(22) International Filing Date:
27 April 2012 (27.04.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

61/480,946	29 April 2011 (29.04.2011)	US
61/513,514	29 July 2011 (29.07.2011)	US
61/531,164	6 September 2011 (06.09.2011)	US
61/531,204	6 September 2011 (06.09.2011)	US
61/531,180	6 September 2011 (06.09.2011)	US
61/531,194	6 September 2011 (06.09.2011)	US
61/531,209	6 September 2011 (06.09.2011)	US
61/531,153	6 September 2011 (06.09.2011)	US
61/531,215	6 September 2011 (06.09.2011)	US
61/531,168	6 September 2011 (06.09.2011)	US
61/531,175	6 September 2011 (06.09.2011)	US
61/531,147	6 September 2011 (06.09.2011)	US

(71) Applicant *(for all designated States except US)*: **SELECTA BIOSCIENCES, INC.** [US/US]; 480 Arsenal Street, Building One, Watertown, MA 02472 (US).

(72) Inventor; and

(75) Inventor/Applicant *(for US only)*: **MALDONADO, Roberto, A.** [VE/US]; 38 Robeson Street #1, Jamaica Plain, MA 02130 (US).

(74) Agent: **VATLAND, Janice, A.**; Wolf, Greenfield & Sacks, P.c., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

[Continued on next page]

(54) Title: TOLEROGENIC SYNTHETIC NANOCARRIERS FOR INDUCING REGULATORY B CELLS



Fig. 2

(57) Abstract: Disclosed are synthetic nanocarrier methods, and related compositions, comprising B cell and/or MHC Class II-restricted epitopes and immunosuppressants in order to generate tolerogenic immune responses, such as the generation of antigen-specific regulatory B cells.

WO 2012/149301 A3

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

— *with sequence listing part of description (Rule 5.2(a))*

(88) Date of publication of the international search report:
31 January 2013

TOLEROGENIC SYNTHETIC NANOCARRIERS FOR INDUCING REGULATORY B CELLS

RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. §119 of United States provisional application 61/480,946, filed April 29, 2011, 61/513,514, filed July 29, 2011, 61/531,147, filed September 6, 2011, 61/531,153, filed September 6, 2011, 61/531,164, filed September 6, 2011, 61/531,168, filed September 6, 2011, 61/531,175, filed September 6, 2011, 61/531,180, filed September 6, 2011, 61/531,194, filed September 6, 2011, 61/531,204, filed September 6, 2011, 61/531,209, filed September 6, 2011, 61/531,215, filed September 6, 2011, the entire contents of each of which are incorporated herein by reference.

FIELD OF THE INVENTION

This invention relates to methods of administering synthetic nanocarrier compositions with immunosuppressants and B cell and/or MHC Class II-restricted epitopes of an antigen wherein the compositions can generate regulatory B cells, and related compositions. The methods allow for efficient uptake by APCs to shift the immune response in favor of regulatory B cell development specific to antigens.

BACKGROUND OF THE INVENTION

Conventional strategies for generating immunosuppression associated with an undesired immune response are based on broad-acting immunosuppressive drugs. Additionally, in order to maintain immunosuppression, immunosuppressant drug therapy is generally a life-long proposition. Unfortunately, the use of broad-acting immunosuppressants are associated with a risk of severe side effects, such as tumors, infections, nephrotoxicity and metabolic disorders. Accordingly, new immunosuppressant therapies would be beneficial.

SUMMARY OF THE INVENTION

In one aspect, a method comprising administering to a subject a composition that comprises: (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to comprise B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In another aspect, a method comprising

generating antigen-specific regulatory B cells in a subject by administering a composition that comprises (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In another aspect, a method comprising: administering to a subject a composition according to a protocol that was previously shown to generate antigen-specific regulatory B cells in one or more test subjects; wherein the composition comprises: (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In one embodiment, the composition is in an amount effective to generate antigen-specific regulatory B cells in the subject. In another embodiment, the composition is administered in an amount effective to generate antigen-specific regulatory B cells in the subject.

In one embodiment, the first population and the second population are the same population. In another embodiment, the first population and second population are different populations.

In another embodiment, the method further comprises providing or identifying the subject.

In another embodiment, the antigen is a therapeutic protein, an autoantigen or an allergen, or is associated with an inflammatory disease, an autoimmune disease, organ or tissue rejection or graft versus host disease.

In another embodiment, the method further comprises assessing the generation of antigen-specific regulatory B cells in the subject prior to and/or after the administration of the composition.

In another embodiment, the subject has or is at risk of having an inflammatory disease, an autoimmune disease, an allergy, organ or tissue rejection or graft versus host disease. In another embodiment, the subject has undergone or will undergo transplantation. In another embodiment, the subject has or is at risk of having an undesired immune response against a therapeutic protein that is being administered or will be administered to the subject.

In another embodiment, one or more maintenance doses of the composition comprising the first population and second population of synthetic nanocarriers is administered to the subject.

In another embodiment, the administering is by intravenous, intraperitoneal, transmucosal, oral, subcutaneous, pulmonary, intranasal, intradermal or intramuscular

administration. In another embodiment, the administering is by inhalation or intravenous, subcutaneous or transmucosal administration.

In another embodiment, the method further comprises administering a transplantable graft or therapeutic protein. In another embodiment, the administering of the transplantable graft or therapeutic protein, when the therapeutic protein is provided as one or more cells, is by parenteral, intraarterial, intranasal or intravenous administration or by injection to lymph nodes or anterior chamber of the eye or by local administration to an organ or tissue of interest.

In another embodiment, the method further comprises collecting the generated antigen-specific regulatory B cells.

In another embodiment, the immunosuppressants comprise a statin, an mTOR inhibitor, a TGF-β signaling agent, a corticosteroid, an inhibitor of mitochondrial function, a P38 inhibitor, an NF-κβ inhibitor, an adenosine receptor agonist, a prostaglandin E2 agonist, a phosphodiesterasse 4 inhibitor, an HDAC inhibitor or a proteasome inhibitor. In another embodiment, the mTOR inhibitor is rapamycin or a rapamycin analog.

In another embodiment, the load of the immunosuppressants and/or antigens on average across the first and/or second population of synthetic nanocarriers is between 0.0001% and 50% (weight/weight). In another embodiment, the load of the immunosuppressants and/or antigens on average across the first and/or second population of synthetic nanocarriers is between 0.1% and 10% (weight/weight).

In another embodiment, the synthetic nanocarriers of the first population and/or second population comprise lipid nanoparticles, polymeric nanoparticles, metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles or peptide or protein particles. In another embodiment, the synthetic nanocarriers of the first population and/or second population comprise lipid nanoparticles. In another embodiment, the synthetic nanocarriers of the first population and/or second population comprise liposomes. In another embodiment, the synthetic nanocarriers of the first population and/or second population comprise metallic nanoparticles. In another embodiment, the metallic nanoparticles comprise gold nanoparticles. In another embodiment, the synthetic nanocarriers of the first population and/or second population comprise polymeric nanoparticles. In another embodiment, the polymeric nanoparticle comprises polymer that is a non-methoxy-terminated, pluronic polymer. In another

embodiment, the polymeric nanoparticles comprise a polyester, a polyester coupled to a polyether, polyamino acid, polycarbonate, polyacetal, polyketal, polysaccharide, polyethyloxazoline or polyethyleneimine. In another embodiment, the polyester comprises a poly(lactic acid), poly(glycolic acid), poly(lactic-co-glycolic acid) or polycaprolactone. In another embodiment, the polymeric nanoparticles comprise a polyester and a polyester coupled to a polyether. In another embodiment, the polyether comprises polyethylene glycol or polypropylene glycol.

In another embodiment, the mean of a particle size distribution obtained using dynamic light scattering of the synthetic nanocarriers of the first and/or second population is a diameter greater than 100nm. In another embodiment, the diameter is greater than 150nm. In another embodiment, the diameter is greater than 200nm. In another embodiment, the diameter is greater than 250nm. In another embodiment, the diameter is greater than 300nm. In another embodiment, the aspect ratio of the synthetic nanocarriers of the first population and/or second population is greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7 or 1:10.

In another embodiment, the method further comprises making a dosage form comprising the collected antigen-specific regulatory B cells. In another embodiment, the method further comprises making the collected antigen-specific regulatory B cells available or dosage form available to a subject for administration.

In another embodiment, the second population of synthetic nanocarriers are also coupled to antigens that comprise MHC I-restricted epitopes.

In another aspect, a composition comprising the isolated antigen-specific regulatory B cells produced according to any of the methods provided herein is provided.

In another embodiment, the composition further comprises a pharmaceutically acceptable excipient.

In another aspect, a dosage form comprising the any of the compositions provided herein is provided.

In another aspect, a composition comprising isolated antigen-specific regulatory B cells for use in therapy or prophylaxis, said isolated antigen-specific regulatory B cells being obtainable by a process comprising the steps of: (a) generating antigen-specific regulatory B cells in a subject by administering a composition that comprises: (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen; and (b) collecting the generated antigen-specific regulatory B cells is provided. In

one embodiment, the composition is any of the compositions provided herein or used in any of the methods provided.

In another aspect, a composition comprising: (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen, for use in therapy or prophylaxis is provided. In one embodiment, the composition is any of the compositions provided herein or used in any of the methods provided.

In another aspect, any of the compositions provided may be for use in therapy or prophylaxis.

In another aspect, any of the compositions provided may be for use in promoting a tolerogenic immune response or generating antigen-specific regulatory B cells in a subject.

In another aspect, a use of any of the compositions or dosage forms provided for the manufacture of a medicament for use in promoting a tolerogenic immune response or generating antigen-specific regulatory B cells in a subject is provided.

In another embodiment, the composition further comprises a transplantable graft or therapeutic protein.

In another aspect, a dosage form comprising any of the compositions provided is provided.

In another aspect, a method comprising: (i) producing a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) producing a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In one embodiment, the composition is in an amount effective to generate antigen-specific regulatory B cells in a subject. In another embodiment, the first population and second population are the same population. In another embodiment, the first population and second population are different populations.

In another embodiment, the method further comprises making a dosage form comprising the first population and second population of synthetic nanocarriers.

In another embodiment, the method further comprises making a composition comprising the first population and second population of synthetic nanocarriers or the dosage form available to a subject for administration.

In another embodiment, the method further comprises assessing the generation of antigen-specific regulatory B cells with a composition comprising the first population and second population of synthetic nanocarriers.

In another embodiment, the first population and second population of synthetic nanocarriers that are produced are as defined in any of the compositions and methods provided herein.

In another aspect, a process for producing a composition or dosage form comprising the steps of: (i) coupling a first population of synthetic nanocarriers to immunosuppressants, and (ii) coupling a second population of synthetic nanocarriers to B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In one embodiment, the composition or dosage form is in an amount effective to generate regulatory B cells in a subject. In one embodiment, the steps are as defined in any of the methods provided herein.

In another aspect, a composition or dosage form obtainable by any of the methods or processes provided herein is provided.

In another embodiment, the methods further comprise including a transplantable graft or therapeutic protein with the collected antigen-specific regulatory B cells or dosage form.

In another embodiment, the first and/or second population of synthetic nanocarriers of any of the compositions or methods provided are also coupled to MHC Class I-restricted epitopes. In an embodiment of any of the compositions and methods provided herein, the antigens comprise substantially no MHC Class I-restricted epitopes.

In an embodiment of any of the compositions and methods provided herein, antigens that are proteins that comprise the aforementioned epitopes can be coupled to the synthetic nanocarriers. In another embodiment, polypeptides or peptides that comprise the aforementioned epitopes but additional amino acids that flank one or both ends of the epitope(s) can be coupled to the synthetic nanocarriers. In another embodiment, the epitopes themselves are coupled to the synthetic nanocarriers.

BRIEF DESCRIPTION OF FIGURES

Fig. 1 provides a representative example of a flow cytometric analysis of Treg cells.

Fig. 2 demonstrates that synthetic nanocarriers of the invention resulted in higher proportions of antigen-specific Bregs.

Fig. 3 shows a decrease in IgE production with synthetic nanocarriers of the invention.

DETAILED DESCRIPTION OF THE INVENTION

Before describing the present invention in detail, it is to be understood that this invention is not limited to particularly exemplified materials or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting of the use of alternative terminology to describe the present invention.

All publications, patents and patent applications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety for all purposes.

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the content clearly dictates otherwise. For example, reference to "a polymer" includes a mixture of two or more such molecules or a mixture of differing molecular weights of a single polymer species, reference to "a synthetic nanocarrier" includes a mixture of two or more such synthetic nanocarriers or a plurality of such synthetic nanocarriers, reference to “a DNA molecule” includes a mixture of two or more such DNA molecules or a plurality of such DNA molecules, reference to "an immunosuppressant" includes a mixture of two or more such materials or a plurality of an immunosuppressant molecules, and the like.

As used herein, the term “comprise” or variations thereof such as “comprises” or “comprising” are to be read to indicate the inclusion of any recited integer (e.g. a feature, element, characteristic, property, method/process step or limitation) or group of integers (e.g. features, element, characteristics, properties, method/process steps or limitations) but not the exclusion of any other integer or group of integers. Thus, as used herein, the term “comprising” is inclusive and does not exclude additional, unrecited integers or method/process steps.

In embodiments of any of the compositions and methods provided herein, “comprising” may be replaced with “consisting essentially of” or “consisting of”. The phrase “consisting essentially of” is used herein to require the specified integer(s) or steps as well as those which do not materially affect the character or function of the claimed invention. As used herein, the term “consisting” is used to indicate the presence of the recited integer (e.g. a feature, element, characteristic, property, method/process step or limitation) or group of integers (e.g. features, element, characteristics, properties, method/process steps or limitations) alone.

A. INTRODUCTION

Delivering immunosuppressants and antigens via synthetic nanocarrier compositions that comprise B cell and/or MHC Class II-restricted epitopes more directly at the sites of action in cells of interest, in particular APCs, can stimulate regulatory B cells and result in beneficial tolerogenic immune responses specific to the antigens. As shown in the examples, synthetic nanocarrier compositions were found to be effective in generating CD24+ B cells that produce IL-10 and TGF-β and also recognize antigen, which speaks to the antigen-specific nature of the response. Additionally, the same synthetic nanocarrier compositions were found to decrease IgE production, which suggests a downstream result of the regulatory B cell production. The compositions provided, without being bound to any particular theory, can provide tolerogenic immune effects directly as a result of the recognition of epitopes, which may cause the generation of the regulatory cells, the switching of B cells to a regulatory phenotype, etc. The tolerogenic immune responses may also be a result of the production of cytokines and/or the production of other regulatory immune cells stimulated by such cytokines. The generation of regulatory B cells can also lead to downstream tolerogenic effects. For example, regulatory B cells can reduce the proliferative capacity of effector T cells, such as CD4+ T cells, and/or enhance FoxP3 and CTLA-4 expression in regulatory T cells. Regulatory B cells can also produce the regulatory cytokine, IL-10. In embodiments, the antigen-specific itDCs provided herein may have these effects, for example, on a subject.

The generation of such regulatory B cells evidences the ability of the compositions of the invention to generate antigen-specific tolerogenic immune responses that can have utility in the treatment or prophylaxis of a variety of diseases, disorders or conditions. Therefore, this invention is useful, for example, to promote tolerogenic immune responses in subjects who have or are at risk of having an allergy, autoimmune disease, an inflammatory disease, organ or tissue rejection or graft versus host disease. This invention is also useful for promoting tolerogenic immune responses in subjects who have undergone or will undergo transplantation. This invention is also useful for promoting tolerogenic immune responses in subjects that have received, are receiving or will receive a therapeutic protein against which undesired immune responses are generated or are expected to be generated. The present invention, in some embodiments, prevents or suppresses undesired immune responses that may neutralize the beneficial effect of certain therapeutic treatments.

The inventors have unexpectedly and surprisingly discovered that the problems and limitations noted above can be overcome by practicing the invention disclosed herein. In particular, the inventors have unexpectedly discovered that it is possible to provide synthetic

nanocarrier compositions, and related methods, that induce a tolerogenic immune response through the the increase in the number and/or activity of regulatory B cells. The method described herein include administering to a subject a composition that comprises (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen. Preferably, the composition is in an amount effective to effect antigen-specific regulatory B cell proliferation and/or activity in the subject. In another aspect, a method comprising generating antigen-specific regulatory B cells in a subject by administering a composition that comprises (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In another aspect, a method comprising administering to a subject a composition according to a protocol that was previously shown to generate antigen-specific regulatory B cells in one or more test subjects wherein the composition comprises (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen is provided.

Transplantable grafts, therapeutic proteins, etc. may also be administered to the subjects as provided herein. Such compositions may be administered to a subject prior to, concomitantly with or after the administration of the first and second populations of synthetic nanocarriers. Such additional agents may or may not be coupled to the first or second population of synthetic nanocarriers or another population of synthetic nanocarriers. In embodiments, the compositions provided may also be administered as one or more maintenance doses to a subject. In such embodiments, the compositions provided are administered such that the generation of a desired immune response is maintained or an undesired immune response is reduced for a certain length of time. Examples of such lengths of time are provided elsewhere herein.

In another aspect, a composition comprising isolated antigen-specific regulatory B cells produced according to any of the methods provided is provided. In yet another aspect, the compositions comprising the first population and second population of synthetic nanocarriers are also provided. In embodiments, the compositions further comprise a transplantable graft or a therapeutic protein that may or may not be coupled to the first or second population of synthetic nanocarriers or another population of synthetic nanocarriers.

In another aspect, dosage forms of any of the compositions herein are provided. Such dosage forms may be administered to a subject, for example, one in need of antigen-specific regulatory B cell generation.

In yet another aspect, a method of (i) producing a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) producing a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen is provided. In one embodiment, the method further comprises producing a dosage form comprising the first and second populations of synthetic nanocarriers. In another embodiment, the method further comprises ensuring the second population of synthetic nanocarriers comprises B cell and/or MHC Class II-restricted epitopes of an antigen. In one embodiment, the presence of a B cell epitope and/or MHC Class II-restricted can be confirmed by the ability of such epitope to stimulate immune responses such as antibody production or B cell proliferation and/or activity for the B cell epitopes and CD4+ T cell immune responses for the MHC Class II-restricted epitopes. Methods for determining these activity are well known to those of ordinary skill in the art or are provided elsewhere herein. In still another embodiment, the method further comprises assessing the generation of regulatory B cells with a composition or dosage form comprising the first population and second population of synthetic nanocarriers. In one embodiment, this assessment is performed in vitro. In another embodiment, this assessment is performed in vivo, such as with a sample obtained from a subject. In yet another embodiment, the method further comprises making a composition comprising the first population and second population of synthetic nanocarriers or the dosage form available to a subject for administration.

In another aspect, the compositions or dosage forms produced by any of the methods provided herein are also provided.

The invention will now be described in more detail below.

B. DEFINITIONS

"Administering" or "administration" means providing a material to a subject in a manner that is pharmacologically useful.

“Allergens” are any substances that can cause an undesired (e.g., a Type 1 hypersensitive) immune response (i.e., an allergic response or reaction) in a subject. Allergens include, but are not limited to, plant allergens (e.g., pollen, ragweed allergen), insect allergens, insect sting allergens (e.g., bee sting allergens), animal allergens (e.g., pet

allergens, such as animal dander or cat Fel d 1 antigen), latex allergens, mold allergens, fungal allergens, cosmetic allergens, drug allergens, food allergens, dust, insect venom, viruses, bacteria, etc. Food allergens include, but are not limited to milk allergens, egg allergens, nut allergens (e.g., peanut or tree nut allergens, etc. (e.g., walnuts, cashews, etc.)), fish allergens, shellfish allergens, soy allergens, legume allergens, seed allergens and wheat allergens. Insect sting allergens include allergens that are or are associated with bee stings, wasp stings, hornet stings, yellow jacket stings, etc. Insect allergens also include house dust mite allergens (e.g., Der P1 antigen) and cockroach allergens. Drug allergens include allergens that are or are associated with antibiotics, NSAIDs, anaesthetics, etc. Pollen allergens include grass allergens, tree allergens, weed allergens, flower allergens, etc. Subjects that develop or are at risk of developing an undesired immune response to any of the allergens provided herein may be treated with any of the compositions and methods provided herein. Subjects that may be treated with any of the compositions and methods provided also include those who have or are at risk of having an allergy to any of the allergens provided.

An "allergy" also referred to herein as an "allergic condition," is any condition where there is an undesired (e.g., a Type 1 hypersensitive) immune response (i.e., allergic response or reaction) to a substance. Such substances are referred to herein as allergens. Allergies or allergic conditions include, but are not limited to, allergic asthma, hay fever, hives, eczema, plant allergies, bee sting allergies, pet allergies, latex allergies, mold allergies, cosmetic allergies, food allergies, allergic rhinitis or coryza, topic allergic reactions, anaphylaxis, atopic dermatitis, hypersensitivity reactions and other allergic conditions. The allergic reaction may be the result of an immune reaction to any allergen. In some embodiments, the allergy is a food allergy. Food allergies include, but are not limited to, milk allergies, egg allergies, nut allergies, fish allergies, shellfish allergies, soy allergies or wheat allergies.

"Amount effective" in the context of a composition or dosage form for administration to a subject refers to an amount of the composition or dosage form that produces one or more desired immune responses in the subject, for example, the generation of regulatory B cells. Therefore, in some embodiments, an amount effective is any amount of a composition provided herein that produces such a desired immune response. This amount can be for in vitro or in vivo purposes. For in vivo purposes, the amount can be one that a clinician would believe may have a clinical benefit for a subject in need of antigen-specific tolerization. Such subjects include those that have or are at risk of having an inflammatory disease, an autoimmune disease, an allergy, organ or tissue rejection or graft versus host disease. Such

subjects also include those that have undergone or will undergo transplantation. Such subjects further include those that have experienced, are experiencing or are expected to experience an undesired immune response against a therapeutic protein. Other subjects include those described elsewhere herein.

Amounts effective can involve only reducing the level of an undesired immune response, although in some embodiments, it involves preventing an undesired immune response altogether. Amounts effective can also involve delaying the occurrence of an undesired immune response. Amounts effective can also involve generating a desired immune response. An amount that is effective can also be an amount of a composition provided herein that produces a desired therapeutic endpoint or a desired therapeutic result. Amounts effective, preferably, result in a tolerogenic immune response in a subject to an antigen. Preferably, the amounts effective result in the increased proliferation and/or activity of regulatory B cells. The achievement of any of the foregoing can be monitored by routine methods.

In some embodiments of any of the compositions and methods provided, the amount effective is one in which the desired immune response persists in the subject for at least 1 week, at least 2 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, at least 9 months, at least 1 year, at least 2 years, at least 5 years, or longer. In other embodiments of any of the compositions and methods provided, the amount effective is one which produces a measurable desired immune response, for example, a measurable decrease in an immune response (e.g., to a specific antigen), for at least 1 week, at least 2 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, at least 9 months, at least 1 year, at least 2 years, at least 5 years, or longer.

Amounts effective will depend, of course, on the particular subject being treated; the severity of a condition, disease or disorder; the individual patient parameters including age, physical condition, size and weight; the duration of the treatment; the nature of concurrent therapy (if any); the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however,

that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reason.

In general, doses of the immunosuppressants and/or antigens in the compositions of the invention can range from about 10 μg/kg to about 100,000 μg/kg. In some embodiments, the doses can range from about 0.1 mg/kg to about 100 mg/kg. In still other embodiments, the doses can range from about 0.1 mg/kg to about 25 mg/kg, about 25 mg/kg to about 50 mg/kg, about 50 mg/kg to about 75 mg/kg or about 75 mg/kg to about 100 mg/kg. Alternatively, the dose can be administered based on the number of synthetic nanocarriers that provide the desired amount of immunosuppressants and/or antigens. For example, useful doses include greater than 10^6, 10^7, 10^8, 10^9 or 10^{10} synthetic nanocarriers per dose. Other examples of useful doses include from about $1x10^6$ to about $1x10^{10}$, about $1x10^7$ to about $1x10^9$ or about $1x10^8$ to about $1x10^9$ synthetic nanocarriers per dose.

"Antigen" means a B cell antigen or T cell antigen. "Type(s) of antigens" means molecules that share the same, or substantially the same, antigenic characteristics. In some embodiments, antigens may be proteins, polypeptides, peptides, lipoproteins, glycolipids, polynucleotides, polysaccharides or are contained or expressed in cells. In some embodiments, such as when the antigens are not well defined or characterized, the antigens may be contained within a cell or tissue preparation, cell debris, cell exosomes, conditioned media, etc. An antigen can be combined with the synthetic nanocarriers in the same form as what a subject is exposed to that causes an undesired immune response but may also be a fragment or derivative thereof. When a fragment or derivative, however, a desired immune response to the form encountered by such a subject is the preferable result with the compositions and methods provided. Preferably, the antigens comprise epitopes such that regulatory B cells are generated, recruited or activated with the compositions provided herein. Whether or not this occurs can be established through measurement of cytokine production (e.g., IL-10) and/or regulatory B cell proliferation using conventional techniques.

"Antigen-specific" refers to any immune response that results from the presence of the antigen, or portion thereof, or that generates molecules that specifically recognize or bind the antigen. For example, where the immune response is antigen-specific antibody production, antibodies are produced that specifically bind the antigen. As another example, where the immune response is antigen-specific regulatory B cell proliferation and/or activity, the proliferation and/or activity results from recognition of the antigen, or portion thereof, alone or in complex with MHC molecules, by B cells, etc.

"Antigens associated" with a disease, disorder or condition provided herein are antigens that can generate an undesired immune response against, as a result of, or in conjunction with the disease, disorder or condition; the cause of the disease, disorder or condition (or a symptom or effect thereof); and/or can generate an undesired immune response that is a symptom, result or effect of the disease, disorder or condition. Preferably, in some embodiments, the use of an antigen associated with a disease, disorder or condition, etc. in the compositions and methods provided herein will lead to a tolerogenic immune response against the antigen and/or the cells, by, on or in which the antigen is expressed. The antigens can be in the same form as expressed in a subject with the disease, disorder or condition but may also be a fragment or derivative thereof. When a fragment or derivative, however, a desired immune response to the form expressed in such a subject is the preferable result with the compositions and methods provided.

In one embodiment, the antigen is an antigen associated with an inflammatory disease, autoimmune disease, organ or tissue rejection or graft versus host disease. Such antigens include autoantigens, such as myelin basic protein, collagen (e.g., collagen type 11), human cartilage gp 39, chromogranin A, gp130-RAPS, proteolipid protein, fibrillarin, nuclear proteins, nucleolar proteins (e.g., small nucleolar protein), thyroid stimulating factor receptor, histones, glycoprotein gp 70, ribosomal proteins, pyruvate dehydrogenase dehydrolipoamide acetyltransferase, hair follicle antigens, human tropomyosin isoform 5, mitochondrial proteins, pancreatic β-cell proteins, myelin oligodendrocyte glycoprotein, insulin, glutamic acid decarboxylase (GAD), gluten, and fragments or derivatives thereof. Other autoantigens are provided in Table 1 below.

Antigens also include those associated with organ or tissue rejection. Examples of such antigens include, but are not limited to, antigens from allogeneic cells, e.g., antigens from an allogeneic cell extract and antigens from other cells, such as endothelial cell antigens.

Antigens also include those associated with an allergy. Such antigens include the allergens described elsewhere herein.

Antigens also include those associated with a transplantable graft. Such antigens are associated with a transplantable graft, or an undesired immune response in a recipient of a transplantable graft that is generated as a result of the introduction of the transplantable graft in the recipient, that can be presented for recognition by cells of the immune system and that can generate an undesired immune response. Transplant antigens include those associated

with organ or tissue rejection or graft versus host disease. Transplant antigens may be obtained or derived from cells of a biological material or from information related to a transplantable graft. Transplant antigens generally include proteins, polypeptides, peptides, lipoproteins, glycolipids, polynucleotides or are contained or expressed in cells. Information related to a transplantable graft is any information about a transplantable graft that can be used to obtain or derive transplant antigens. Such information includes information about antigens that would be expected to be present in or on cells of a transplantable graft such as, for example, sequence information, types or classes of antigens and/or their MHC Class I, MHC Class II or B cell presentation restrictions. Such information may also include information about the type of transplantable graft (e.g, autograft, allograft, xenograft), the molecular and cellular composition of the graft, the bodily location from which the graft is derived or to which the graft to be transplanted (e.g., whole or partial organ, skin, bone, nerves, tendon, neurons, blood vessels, fat, cornea, etc.).

Antigens also include antigens associated with a therapeutic protein that can be presented for recognition by cells of the immune system and that can generate an undesired immune response against the therapeutic protein. Therapeutic protein antigens generally include proteins, polypeptides, peptides, lipoproteins, or are contained or expressed in, by or on cells.

Antigens, can be antigens that are fully defined or characterized. However, in some embodiments, an antigen is not fully defined or characterized. Antigens, therefore, also include those that are contained within a cell or tissue preparation, cell debris, cell exosome or conditioned media and can be delivered in such form in some embodiments.

"Assessing an immune response" refers to any measurement or determination of the level, presence or absence, reduction, increase in, etc. of an immune response in vitro or in vivo. Such measurements or determinations may be performed on one or more samples obtained from a subject. Such assessing can be performed with any of the methods provided herein or otherwise known in the art.

An "at risk" subject is one in which a health practitioner believes has a chance of having a disease, disorder or condition as provided herein or is one a health practitioner believes has a chance of experiencing an undesired immune response as provided herein.

An "autoimmune disease" is any disease where the immune system mounts an undesired immune response against self (e.g., one or more autoantigens). In some embodiments, an autoimmune disease comprises an aberrant destruction of cells of the body

as part of the self-targeted immune response. In some embodiments, the destruction of self manifests in the malfunction of an organ, for example, the colon or pancreas. Examples of autoimmune diseases are described elsewhere herein. Additional autoimmune diseases will be known to those of skill in the art and the invention is not limited in this respect.

"Average", as used herein, refers to the arithmetic mean unless otherwise noted.

"B cell antigen" means any antigen that triggers an immune response in a B cell (e.g., an antigen that is specifically recognized by a B cell or a receptor thereon). In some embodiments, an antigen that is a T cell antigen is also a B cell antigen. In other embodiments, the T cell antigen is not also a B cell antigen. B cell antigens include, but are not limited to proteins, peptides, small molecules, and carbohydrates. In some embodiments, the B cell antigen comprises a non-protein antigen (i.e., not a protein or peptide antigen). In some embodiments, the B cell antigen comprises a autoantigen. In other embodiments, the B cell antigen is obtained or derived from an allergen, autoantigen, therapeutic protein, or transplantable graft.

"Concomitantly" means administering two or more substances to a subject in a manner that is correlated in time, preferably sufficiently correlated in time so as to provide a modulation in an immune response. In embodiments, concomitant administration may occur through administration of two or more substances in the same dosage form. In other embodiments, concomitant administration may encompass administration of two or more substances in different dosage forms, but within a specified period of time, preferably within 1 month, more preferably within 1 week, still more preferably within 1 day, and even more preferably within 1 hour.

"Couple" or "Coupled" or "Couples" (and the like) means to chemically associate one entity (for example a moiety) with another. In some embodiments, the coupling is covalent, meaning that the coupling occurs in the context of the presence of a covalent bond between the two entities. In non-covalent embodiments, the non-covalent coupling is mediated by non-covalent interactions including but not limited to charge interactions, affinity interactions, metal coordination, physical adsorption, host-guest interactions, hydrophobic interactions, TT stacking interactions, hydrogen bonding interactions, van der Waals interactions, magnetic interactions, electrostatic interactions, dipole-dipole interactions, and/or combinations thereof. In embodiments, encapsulation is a form of coupling.

"Derived" means prepared from a material or information related to a material but is not "obtained" from the material. Such materials may be substantially modified or processed

forms of materials taken directly from a biological material. Such materials also include materials produced from information related to a biological material.

"Dosage form" means a pharmacologically and/or immunologically active material in a medium, carrier, vehicle, or device suitable for administration to a subject.

"Encapsulate" means to enclose at least a portion of a substance within a synthetic nanocarrier. In some embodiments, a substance is enclosed completely within a synthetic nanocarrier. In other embodiments, most or all of a substance that is encapsulated is not exposed to the local environment external to the synthetic nanocarrier. In other embodiments, no more than 50%, 40%, 30%, 20%, 10% or 5% (weight/weight) is exposed to the local environment. Encapsulation is distinct from absorption, which places most or all of a substance on a surface of a synthetic nanocarrier, and leaves the substance exposed to the local environment external to the synthetic nanocarrier.

"Epitope", also known as an antigenic determinant, is the part of an antigen that is recognized by the immune system, specifically by, for example, antibodies, B cells, or T cells. As used herein, "MHC Class I-restricted epitopes" are epitopes that are presented to immune cells by MHC class I molecules found on nucleated cells. "MHC Class II-restricted epitopes" are epitopes that are presented to immune cells by MHC class II molecules found on antigen-presenting cells (APCs), for example, on professional antigen-presenting immune cells, such as on macrophages, B cells, and dendritic cells, or on non-hematopoietic cells, such as hepatocytes. "B cell epitopes" are molecular structures that are recognized by antibodies or B cells. In some embodiments, the epitope itself is an antigen.

A number of epitopes are known to those of skill in the art, and exemplary epitopes suitable according to some aspects of this invention include, but are not limited to those listed in the Immune Epitope Database (www.immuneepitope.org, Vita R, Zarebski L, Greenbaum JA, Emami H, Hoof I, Salimi N, Damle R, Sette A, Peters B. The immune epitope database 2.0. Nucleic Acids Res. 2010 Jan;38(Database issue):D854-62; the entire contents of which as well as all database entries of IEDB version 2.4, August 2011, and particularly all epitopes disclosed therein, are incorporated herein by reference). Epitopes can also be identified with publicly available algorithms, for example, the algorithms described in Wang P, Sidney J, Kim Y, Sette A, Lund O, Nielsen M, Peters B. 2010. peptide binding predictions for HLA DR, DP and DQ molecules. BMC Bioinformatics 2010, 11:568; Wang P, Sidney J, Dow C, Mothé B, Sette A, Peters B. 2008. A systematic assessment of MHC class II peptide binding predictions and evaluation of a consensus approach. PLoS Comput Biol. 4(4):e1000048;

Nielsen M, Lund O. 2009. NN-align. An artificial neural network-based alignment algorithm for MHC class II peptide binding prediction. BMC Bioinformatics. 10:296; Nielsen M, Lundegaard C, Lund O. 2007. Prediction of MHC class II binding affinity using SMM-align, a novel stabilization matrix alignment method. BMC Bioinformatics. 8:238; Bui HH, Sidney J, Peters B, Sathiamurthy M, Sinichi A, Purton KA, Mothé BR, Chisari FV, Watkins DI, Sette A. 2005. Immunogenetics. 57:304-314; Sturniolo T, Bono E, Ding J, Raddrizzani L, Tuereci O, Sahin U, Braxenthaler M, Gallazzi F, Protti MP, Sinigaglia F, Hammer J. 1999. Generation of tissue-specific and promiscuous HLA ligand databases using DNA microarrays and virtual HLA class II matrices. Nat Biotechnol. 17(6):555-561; Nielsen M, Lundegaard C, Worning P, Lauemoller SL, Lamberth K, Buus S, Brunak S, Lund O. 2003. Reliable prediction of T-cell epitopes using neural networks with novel sequence representations. Protein Sci 12:1007-1017; Bui HH, Sidney J, Peters B, Sathiamurthy M, Sinichi A, Purton KA, Mothe BR, Chisari FV, Watkins DI, Sette A. 2005. Automated generation and evaluation of specific MHC binding predictive tools: ARB matrix applications. Immunogenetics 57:304-314; Peters B, Sette A. 2005. Generating quantitative models describing the sequence specificity of biological processes with the stabilized matrix method. BMC Bioinformatics 6:132; Chou PY, Fasman GD. 1978. Prediction of the secondary structure of proteins from their amino acid sequence. Adv Enzymol Relat Areas Mol Biol 47:45-148; Emini EA, Hughes JV, Perlow DS, Boger J. 1985. Induction of hepatitis A virus-neutralizing antibody by a virus-specific synthetic peptide. J Virol 55:836-839; Karplus PA, Schulz GE. 1985. Prediction of chain flexibility in proteins. Naturwissenschaften 72:212-213; Kolaskar AS, Tongaonkar PC. 1990. A semi-empirical method for prediction of antigenic determinants on protein antigens. FEBS Lett276:172-174; Parker JM, Guo D, Hodges RS. 1986. New hydrophilicity scale derived from high-performance liquid chromatography peptide retention data: correlation of predicted surface residues with antigenicity and X-ray-derived accessible sites. Biochemistry 25:5425-5432; Larsen JE, Lund O, Nielsen M. 2006. Improved method for predicting linear B-cell epitopes. Immunome Res 2:2; Ponomarenko JV, Bourne PE. 2007. Antibody-protein interactions: benchmark datasets and prediction tools evaluation. BMC Struct Biol 7:64; Haste Andersen P, Nielsen M, Lund O. 2006. Prediction of residues in discontinuous B-cell epitopes using protein 3D structures. Protein Sci 15:2558-2567; Ponomarenko JV, Bui H, Li W, Fusseder N, Bourne PE, Sette A, Peters B. 2008. ElliPro: a new structure-based tool for the prediction of antibody epitopes. BMC Bioinformatics 9:514; Nielsen M, Lundegaard C, Blicher T, Peters B, Sette A, Justesen S,

Buus S, and Lund O. 2008. PLoS Comput Biol.4(7)e1000107. Quantitative predictions of peptide binding to any HLA-DR molecule of known sequence: NetMHCIIpan; the entire contents of each of which are incorporated herein by reference for disclosure of methods and algorithms for the identification of epitopes.

Other examples of epitopes include any of the MHC Class I-restricted, MHC Class II-restricted and B cell epitopes as provided as SEQ ID NOs: 1-943. Without wishing to being bound by any particular theory, MHC Class I-restricted epitopes include those set forth in SEQ ID NOs: 1-186, MHC Class II-restricted epitopes include those set forth in SEQ ID NOs: 187-537, and B cell epitopes include those set forth in SEQ ID NOs: 538-943. These epitopes include MHC Class I-restricted autoantigens, MHC Class II-restricted epitopes of allergens and B cell epitopes of autoantigens and allergens.

"Generating" means causing an action, such as an immune response (e.g., a tolerogenic immune response) to occur, either directly oneself or indirectly, such as, but not limited to, an unrelated third party that takes an action through reliance on one's words or deeds.

"Identifying" is any action or set of actions that allows a clinician to recognize a subject as one who may benefit from the methods and compositions provided herein. Preferably, the identified subject is one who is in need of a tolerogenic immune response as provided herein. The action or set of actions may be either directly oneself or indirectly, such as, but not limited to, an unrelated third party that takes an action through reliance on one's words or deeds.

"Immunosuppressant" means a compound that causes an APC to have an immunosuppressive (e.g., tolerogenic effect). An immunosuppressive effect generally refers to the production or expression of cytokines or other factors by the APC that reduces, inhibits or prevents an undesired immune response but can also include the production or expression of cytokines or other factors by the APC that increases, stimulates or promotes a desired immune response. When the APC results in an immunosuppressive effect on immune cells that recognize an antigen presented by the APC, the immunosuppressive effect is said to be specific to the presented antigen. Such effect is also referred to herein as a tolerogenic effect. Without being bound by any particular theory, it is thought that the immunosuppressive or tolerogenic effect is a result of the immunosuppressant being delivered to the APC, preferably in the presence of an antigen (e.g., an administered antigen or one that is already present in vivo). Accordingly, the immunosuppressant includes compounds that provide a

tolerogenic immune response to an antigen that may or may not be provided in the same composition or a different composition. In one embodiment, the immunosuppressant is one that causes an APC to promote a regulatory phenotype in one or more immune effector cells. For example, the regulatory phenotype may be characterized by the production, induction, stimulation or recruitment of Treg cells, etc. This may be the result of the conversion of B cells to a regulatory phenotype. This may also be the result of the stimulation, recruitment, etc. of regulatory B cells. This may also be the result of induction of FoxP3 in other immune cells, such as macrophages and iNKT cells. In one embodiment, the immunosuppressant is one that affects the response of the APC after it processes an antigen. In another embodiment, the immunosuppressant is not one that interferes with the processing of the antigen. In a further embodiment, the immunosuppressant is not an apoptotic-signaling molecule. In another embodiment, the immunosuppressant is not a phospholipid.

Immunosuppressants include, but are not limited to, statins; mTOR inhibitors, such as rapamycin or a rapamycin analog; TGF-β signaling agents; TGF-β receptor agonists; histone deacetylase inhibitors, such as Trichostatin A; corticosteroids; inhibitors of mitochondrial function, such as rotenone; P38 inhibitors; NF-κβ inhibitors, such as 6Bio, Dexamethasone, TCPA-1, IKK VII; adenosine receptor agonists; prostaglandin E2 agonists (PGE2), such as Misoprostol; phosphodiesterase inhibitors, such as phosphodiesterase 4 inhibitor (PDE4), such as Rolipram; proteasome inhibitors; kinase inhibitors; G-protein coupled receptor agonists; G-protein coupled receptor antagonists; glucocorticoids; retinoids; cytokine inhibitors; cytokine receptor inhibitors; cytokine receptor activators; peroxisome proliferator-activated receptor antagonists; peroxisome proliferator-activated receptor agonists; histone deacetylase inhibitors; calcineurin inhibitors; phosphatase inhibitors; PI3KB inhibitors, such as TGX-221; autophagy inhibitors, such as 3-Methyladenine; aryl hydrocarbon receptor inhibitors; proteasome inhibitor I (PSI); and oxidized ATPs, such as P2X receptor blockers. Immunosuppressants also include IDO, vitamin D3, cyclosporins, such as cyclosporine A, aryl hydrocarbon receptor inhibitors, resveratrol, azathiopurine (Aza), 6-mercaptopurine (6-MP), 6-thioguanine (6-TG), FK506, sanglifehrin A, salmeterol, mycophenolate mofetil (MMF), aspirin and other COX inhibitors, niflumic acid, estriol and triptolide. In embodiments, the immunosuppressant may comprise any of the agents provided herein.

The immunosuppressant can be a compound that directly provides the immunosuppressive (e.g., tolerogenic) effect on APCs or it can be a compound that provides

the immunosuppressive (e.g., tolerogenic) effect indirectly (i.e., after being processed in some way after administration). Immunosuppressants, therefore, include prodrug forms of any of the compounds provided herein.

Immunosuppressants also include nucleic acids that encode the peptides, polypeptides or proteins provided herein that result in an immunosuppressive (e.g., tolerogenic) immune response. In embodiments, therefore, the immunosuppressant is a nucleic acid that encodes a peptide, polypeptide or protein that results in an immunosuppressive (e.g., tolerogenic) immune response, and it is the nucleic acid that is coupled to the synthetic nanocarrier.

The nucleic acid may be DNA or RNA, such as mRNA. In embodiments, the inventive compositions comprise a complement, such as a full-length complement, or a degenerate (due to degeneracy of the genetic code) of any of the nucleic acids provided herein. In embodiments, the nucleic acid is an expression vector that can be transcribed when transfected into a cell line. In embodiments, the expression vector may comprise a plasmid, retrovirus, or an adenovirus amongst others. Nucleic acids can be isolated or synthesized using standard molecular biology approaches, for example by using a polymerase chain reaction to produce a nucleic acid fragment, which is then purified and cloned into an expression vector. Additional techniques useful in the practice of this invention may be found in Current Protocols in Molecular Biology 2007 by John Wiley and Sons, Inc.; Molecular Cloning: A Laboratory Manual (Third Edition) Joseph Sambrook, Peter MacCallum Cancer Institute, Melbourne, Australia; David Russell, University of Texas Southwestern Medical Center, Dallas, Cold Spring Harbor.

In embodiments, the immunosuppressants provided herein are coupled to synthetic nanocarriers. In preferable embodiments, the immunosuppressant is an element that is in addition to the material that makes up the structure of the synthetic nanocarrier. For example, in one embodiment, where the synthetic nanocarrier is made up of one or more polymers, the immunosuppressant is a compound that is in addition and coupled to the one or more polymers. As another example, in one embodiment, where the synthetic nanocarrier is made up of one or more lipids, the immunosuppressant is again in addition and coupled to the one or more lipids. In embodiments, such as where the material of the synthetic nanocarrier also results in an immunosuppressive (e.g., tolerogenic) effect, the immunosuppressant is an element present in addition to the material of the synthetic nanocarrier that results in an immunosuppressive (e.g., tolerogenic) effect.

Other exemplary immunosuppressants include, but are not limited, small molecule drugs, natural products, antibodies (e.g., antibodies against CD20, CD3, CD4), biologics-based drugs, carbohydrate-based drugs, nanoparticles, liposomes, RNAi, antisense nucleic acids, aptamers, methotrexate, NSAIDs; fingolimod; natalizumab; alemtuzumab; anti-CD3; tacrolimus (FK506), etc. Further immunosuppressants, are known to those of skill in the art, and the invention is not limited in this respect.

"Inflammatory disease" means any disease, disorder or condition in which undesired inflammation occurs.

"Load" of the immunosuppressant or antigen is the amount of the immunosuppressant or antigen coupled to a synthetic nanocarrier based on the total weight of materials in an entire synthetic nanocarrier (weight/weight). Generally, the load is calculated as an average across a population of synthetic nanocarriers. In one embodiment, the load of the immunosuppressant on average across the first population of synthetic nanocarriers is between 0.0001% and 50%. In another embodiment, the load of the antigen on average across the first and/or second population of synthetic nanocarriers is between 0.0001% and 50%. In yet another embodiment, the load of the immunosuppressant and/or antigen is between 0.01% and 20%. In a further embodiment, the load of the immunosuppressant and/or antigen is between 0.1% and 10%. In still a further embodiment, the load of the immunosuppressant and/or antigen is between 1% and 10%. In yet another embodiment, the load of the immunosuppressant and/or the antigen is at least 0.1%, at least 0.2%, at least 0.3%, at least 0.4%, at least 0.5%, at least 0.6%, at least 0.7%, at least 0.8%, at least 0.9%, at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, at least 6%, at least 7%, at least 8%, at least 9%, at least 10%, at least 11%, at least 12%, at least 13%, at least 14%, at least 15%, at least 16%, at least 17%, at least 18%, at least 19% or at least 20% on average across a population of synthetic nanocarriers. In yet a further embodiment, the load of the immunosuppressant and/or the antigen is 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% on average across a population of synthetic nanocarriers. In some embodiments of the above embodiments, the load of the immunosuppressant and/or the antigen is no more than 25% on average across a population of synthetic nanocarriers. In embodiments, the load is calculated as described in the Examples.

In embodiments of any of the compositions and methods provided, the load may be calculated as follows: Approximately 3 mg of synthetic nanocarriers are collected and

centrifuged to separate supernatant from synthetic nanocarrier pellet. Acetonitrile is added to the pellet, and the sample is sonicated and centrifuged to remove any insoluble material. The supernatant and pellet are injected on RP-HPLC and absorbance is read at 278nm. The μg found in the pellet is used to calculate % entrapped (load), μg in supernatant and pellet are used to calculate total μg recovered.

"Maintenance dose" refers to a dose that is administered to a subject, after an initial dose has resulted in an immunosuppressive (e.g., tolerogenic) response in a subject, to sustain a desired immunosuppressive (e.g., tolerogenic) response. A maintenance dose, for example, can be one that maintains the tolerogenic effect achieved after the initial dose, prevents an undesired immune response in the subject, or prevents the subject becoming a subject at risk of experiencing an undesired immune response, including an undesired level of an immune response. In some embodiments, the maintenance dose is one that is sufficient to sustain an appropriate level of a desired immune response. In some embodiments, the maintenance dose is one that is sufficient to sustain an appropriate antibody titre or level of regulatory B cell number and/or activity necessary to sustain a tolerogenic immune response or defend against a challenge with an agent that results in an undesired immune response.

"Maximum dimension of a synthetic nanocarrier" means the largest dimension of a nanocarrier measured along any axis of the synthetic nanocarrier. "Minimum dimension of a synthetic nanocarrier" means the smallest dimension of a synthetic nanocarrier measured along any axis of the synthetic nanocarrier. For example, for a spheroidal synthetic nanocarrier, the maximum and minimum dimension of a synthetic nanocarrier would be substantially identical, and would be the size of its diameter. Similarly, for a cuboidal synthetic nanocarrier, the minimum dimension of a synthetic nanocarrier would be the smallest of its height, width or length, while the maximum dimension of a synthetic nanocarrier would be the largest of its height, width or length. In an embodiment, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or greater than 100 nm. In an embodiment, a maximum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or less than 5 μm. Preferably, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is greater than 110 nm, more

preferably greater than 120 nm, more preferably greater than 130 nm, and more preferably still greater than 150 nm. Aspects ratios of the maximum and minimum dimensions of inventive synthetic nanocarriers may vary depending on the embodiment. For instance, aspect ratios of the maximum to minimum dimensions of the synthetic nanocarriers may vary from 1:1 to 1,000,000:1, preferably from 1:1 to 100,000:1, more preferably from 1:1 to 10,000:1, more preferably from 1:1 to 1000:1, still more preferably from 1:1 to 100:1, and yet more preferably from 1:1 to 10:1. Preferably, a maximum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample is equal to or less than 3 μm, more preferably equal to or less than 2 μm, more preferably equal to or less than 1 μm, more preferably equal to or less than 800 nm, more preferably equal to or less than 600 nm, and more preferably still equal to or less than 500 nm. In preferred embodiments, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or greater than 100nm, more preferably equal to or greater than 120 nm, more preferably equal to or greater than 130 nm, more preferably equal to or greater than 140 nm, and more preferably still equal to or greater than 150 nm. Measurement of synthetic nanocarrier dimensions (e.g., diameter) is obtained by suspending the synthetic nanocarriers in a liquid (usually aqueous) media and using dynamic light scattering (DLS) (e.g. using a Brookhaven ZetaPALS instrument). For example, a suspension of synthetic nanocarriers can be diluted from an aqueous buffer into purified water to achieve a final synthetic nanocarrier suspension concentration of approximately 0.01 to 0.1 mg/mL. The diluted suspension may be prepared directly inside, or transferred to, a suitable cuvette for DLS analysis. The cuvette may then be placed in the DLS, allowed to equilibrate to the controlled temperature, and then scanned for sufficient time to acquire a stable and reproducible distribution based on appropriate inputs for viscosity of the medium and refractive indicies of the sample. The effective diameter, or mean of the distribution, is then reported. "Dimension" or "size" or "diameter" of synthetic nanocarriers means the mean of a particle size distribution obtained using dynamic light scattering.

"MHC" refers to major histocompatibility complex, a large genomic region or gene family found in most vertebrates that encodes MHC molecules that display fragments or epitopes of processed proteins on the cell surface. The presentation of MHC:peptide on cell

surfaces allows for surveillance by immune cells, usually a T cell. There are two general classes of MHC molecules: Class I and Class II. Generally, Class I MHC molecules are found on nucleated cells and present peptides to cytotoxic T cells. Class II MHC molecules are found on certain immune cells, chiefly macrophages, B cells and dendritic cells, collectively known as professional APCs. The best-known genes in the MHC region are the subset that encodes antigen-presenting proteins on the cell surface. In humans, these genes are referred to as human leukocyte antigen (HLA) genes.

"Non-methoxy-terminated polymer" means a polymer that has at least one terminus that ends with a moiety other than methoxy. In some embodiments, the polymer has at least two termini that ends with a moiety other than methoxy. In other embodiments, the polymer has no termini that ends with methoxy. "Non-methoxy-terminated, pluronic polymer" means a polymer other than a linear pluronic polymer with methoxy at both termini. Polymeric nanoparticles as provided herein can comprise non-methoxy-terminated polymers or non-methoxy-terminated, pluronic polymers.

"Obtained" means taken directly from a material and used with substantially no modification and/or processing.

"Pharmaceutically acceptable excipient" means a pharmacologically inactive material used together with the recited synthetic nanocarriers to formulate the inventive compositions. Pharmaceutically acceptable excipients comprise a variety of materials known in the art, including but not limited to saccharides (such as glucose, lactose, and the like), preservatives such as antimicrobial agents, reconstitution aids, colorants, saline (such as phosphate buffered saline), and buffers.

"Protocol " refers to any dosing regimen of one or more substances to a subject. A dosing regimen may include the amount, frequency and/or mode of administration. In some embodiments, such a protocol may be used to administer one or more compositions of the invention to one or more test subjects. Immune responses in these test subject can then be assessed to determine whether or not the protocol was effective in reducing an undesired immune response or generating a desired immune response (e.g., the promotion of a tolerogenic effect). Any other therapeutic and/or prophylactic effect may also be assessed instead of or in addition to the aforementioned immune responses. Whether or not a protocol had a desired effect can be determined using any of the methods provided herein or otherwise known in the art. For example, a population of cells may be obtained from a subject to which a composition provided herein has been administered according to a specific protocol in order

to determine whether or not specific immune cells, cytokines, antibodies, etc. were reduced, generated, activated, etc. Useful methods for detecting the presence and/or number of immune cells include, but are not limited to, flow cytometric methods (e.g., FACS) and immunohistochemistry methods. Antibodies and other binding agents for specific staining of immune cell markers, are commercially available. Such kits typically include staining reagents for multiple antigens that allow for FACS-based detection, separation and/or quantitation of a desired cell population from a heterogeneous population of cells.

"Providing a subject" is any action or set of actions that causes a clinician to come in contact with a subject and administer a composition provided herein thereto or to perform a method provided herein thereupon. Preferably, the subject is one who is in need of a tolerogenic immune response as provided herein. The action or set of actions may be either directly oneself or indirectly, such as, but not limited to, an unrelated third party that takes an action through reliance on one's words or deeds.

"Regulatory B cells" are art-recognized and refer to a type of tolerogenic B cells or that have suppressive regulatory function. Surface markers and chemokine profiles characteristic of regulatory B cells, as well as subsets of regulatory B cells (e.g., IL-10 producing Bregs) are known to those of skill in the art (e.g., as described in DiLillo et al., B10 cells and regulatory B cells balance immune response during inflammation, autoimmunity, and cancer, Ann N. Y. Acad. Sci. 1183 (2010) 38-57, ISSN 0077-8923; the entire contents of which are incorporated herein by reference). The presence of regulatory B cells can be determined by intracellular staining for IL-10 by flow cytometry. For example, after treatment B cells can be stained for surface markers, then fixed and permeabilized and stained for intracellular IL-10 and analyzed by flow cytometry.

"Subject" means animals, including warm blooded mammals such as humans and primates; avians; domestic household or farm animals such as cats, dogs, sheep, goats, cattle, horses and pigs; laboratory animals such as mice, rats and guinea pigs; fish; reptiles; zoo and wild animals; and the like.

"Substantially no MHC Class I-restricted epitopes" refers to the absence of MHC Class I-restricted epitopes in an amount that by itself, within the context of the antigen, in conjunction with a carrier or in conjunction with an inventive composition stimulates substantial activation of an immune response specific to the epitope presented in the Class I context. In embodiments, a composition with substantially no MHC I-restricted epitopes does not contain a measurable amount of MHC Class I-restricted epitopes of an antigen. In

other embodiments, such a composition may comprise a measurable amount of MHC Class I-restricted epitopes of an antigen but said amount is not effective to generate a measurable cytotoxic T cell immune response (by itself, within the context of the antigen, in conjunction with a carrier or in conjunction with an inventive composition) or is not effective to generate a significant measurable cytotoxic T cell immune response (by itself, within the context of the antigen, in conjunction with a carrier or in conjunction with an inventive composition). In some embodiments, a significant measurable cytotoxic T cell immune response is one that produces or would be expected to produce an adverse clinical result in a subject. In other embodiments, a significant measurable cytotoxic T cell immune response is one that is greater than the level of the same type of immune response produced by a control antigen (e.g., one known not to comprise MHC Class I-restricted epitopes or to stimulate cytotoxic T cell immune responses).

In embodiments, the compositions do not comprise MHC Class I-restricted epitopes (by itself, within the context of the antigen, in conjunction with a carrier or in conjunction with an inventive composition) that generate antigen-specific cytotoxic T cell immune responses or an undesired level thereof. In some embodiments, to ensure that a composition does not comprise such epitopes, antigens are selected such that they do not comprise MHC Class I-restricted epitopes for coupling to the synthetic nanocarriers as provided herein. In other embodiments, to ensure that a composition does not comprise such epitopes, synthetic nanocarriers coupled to antigen are produced and tested for cytotoxic T cell immune responses. The appropriate synthetic nanocarriers may then be selected.

"Synthetic nanocarrier(s)" means a discrete object that is not found in nature, and that possesses at least one dimension that is less than or equal to 5 microns in size. Albumin nanoparticles are generally included as synthetic nanocarriers, however in certain embodiments the synthetic nanocarriers do not comprise albumin nanoparticles. In embodiments, inventive synthetic nanocarriers do not comprise chitosan. In other embodiments, inventive synthetic nanocarriers are not lipid-based nanoparticles. In further embodiments, inventive synthetic nanocarriers do not comprise a phospholipid.

A synthetic nanocarrier can be, but is not limited to, one or a plurality of lipid-based nanoparticles (also referred to herein as lipid nanoparticles, i.e., nanoparticles where the majority of the material that makes up their structure are lipids), polymeric nanoparticles, metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles (i.e., particles that are primarily made up of viral structural proteins but that are

not infectious or have low infectivity), peptide or protein-based particles (also referred to herein as protein particles, i.e., particles where the majority of the material that makes up their structure are peptides or proteins) (such as albumin nanoparticles) and/or nanoparticles that are developed using a combination of nanomaterials such as lipid-polymer nanoparticles. Synthetic nanocarriers may be a variety of different shapes, including but not limited to spheroidal, cuboidal, pyramidal, oblong, cylindrical, toroidal, and the like. Synthetic nanocarriers according to the invention comprise one or more surfaces. Exemplary synthetic nanocarriers that can be adapted for use in the practice of the present invention comprise: (1) the biodegradable nanoparticles disclosed in US Patent 5,543,158 to Gref et al., (2) the polymeric nanoparticles of Published US Patent Application 20060002852 to Saltzman et al., (3) the lithographically constructed nanoparticles of Published US Patent Application 20090028910 to DeSimone et al., (4) the disclosure of WO 2009/051837 to von Andrian et al., (5) the nanoparticles disclosed in Published US Patent Application 2008/0145441 to Penades et al., (6) the protein nanoparticles disclosed in Published US Patent Application 20090226525 to de los Rios et al., (7) the virus-like particles disclosed in published US Patent Application 20060222652 to Sebbel et al., (8) the nucleic acid coupled virus-like particles disclosed in published US Patent Application 20060251677 to Bachmann et al., (9) the virus-like particles disclosed in WO2010047839A1 or WO2009106999A2, (10) the nanoprecipitated nanoparticles disclosed in P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" Nanomedicine. 5(6):843-853 (2010), or (11) apoptotic cells, apoptotic bodies or the synthetic or semisynthetic mimics disclosed in U.S. Publication 2002/0086049. In embodiments, synthetic nanocarriers may possess an aspect ratio greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7, or greater than 1:10.

Synthetic nanocarriers according to the invention that have a minimum dimension of equal to or less than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface with hydroxyl groups that activate complement or alternatively comprise a surface that consists essentially of moieties that are not hydroxyl groups that activate complement. In a preferred embodiment, synthetic nanocarriers according to the invention that have a minimum dimension of equal to or less than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface that substantially activates complement or alternatively comprise a surface that consists essentially of moieties that do not substantially activate complement. In a more preferred embodiment, synthetic nanocarriers according to the

invention that have a minimum dimension of equal to or less than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface that activates complement or alternatively comprise a surface that consists essentially of moieties that do not activate complement. In embodiments, synthetic nanocarriers exclude virus-like particles. In embodiments, synthetic nanocarriers may possess an aspect ratio greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7, or greater than 1:10.

"T cell antigen" means a CD4+ T-cell antigen or CD8+ cell antigen. "CD4+ T-cell antigen" means any antigen that is recognized by and triggers an immune response in a CD4+ T-cell e.g., an antigen that is specifically recognized by a T-cell receptor on a CD4+T cell via presentation of the antigen or portion thereof bound to a Class II major histocompatability complex molecule (MHC). "CD8+ T cell antigen" means any antigen that is recognized by and triggers an immune response in a CD8+ T-cell e.g., an antigen that is specifically recognized by a T-cell receptor on a CD8+T cell via presentation of the antigen or portion thereof bound to a Class I major histocompatability complex molecule (MHC). In some embodiments, an antigen that is a T cell antigen is also a B cell antigen. In other embodiments, the T cell antigen is not also a B cell antigen. T cell antigens generally are proteins or peptides.

A "therapeutic protein" refers to any protein or protein-based therapy that may be administered to a subject and have a therapeutic effect. Such therapies include protein replacement and protein supplementation therapies. Such therapies also include the administration of exogenous or foreign protein, antibody therapies, and cell or cell-based therapies. Therapeutic proteins include enzymes, enzyme cofactors, hormones, blood clotting factors, cytokines, growth factors, monoclonal antibodies and polyclonal antibodies. Examples of other therapeutic proteins are provided elsewhere herein. Therapeutic proteins may be produced in, on or by cells and may be obtained from such cells or administered in the form of such cells. In embodiments, the therapeutic protein is produced in, on or by mammalian cells, insect cells, yeast cells, bacteria cells, plant cells, transgenic animal cells, transgenic plant cells, etc. The therapeutic protein may be recombinantly produced in such cells. The therapeutic protein may be produced in, on or by a virally transformed cell. The therapeutic protein may also be produced in, on or by autologous cells that have been transfected, transduced or otherwise manipulated to express it. Alternatively, the therapeutic protein may be administered as a nucleic acid or by introducing a nucleic acid into a virus, VLP, liposome, etc.. Alternatively, the therapeutic protein may be obtained from such forms

and administered as the therapeutic protein itself. Subjects, therefore, include any subject that has received, is receiving or will receive any of the foregoing. Such subject includes subjects that have received, is receiving or will receive gene therapy, autologous cells that have been transfected, transduced or otherwise manipulated to express a therapeutic protein, polypeptide or peptide; or cells that express a therapeutic protein, polypeptide or peptide.

"Therapeutic protein antigen" means an antigen that is associated with a therapeutic protein that can be, or a portion of which can be, presented for recognition by cells of the immune system and can generate an undesired immune response (e.g., the production of therapeutic protein-specific antibodies) against the therapeutic protein. Therapeutic protein antigens generally include proteins, polypeptides, peptides, lipoproteins, or are contained or expressed in, on or by cells.

"Tolerogenic immune response" means any immune response that can lead to immune suppression specific to an antigen or a cell, tissue, organ, etc. that expresses such an antigen. Such immune responses include any reduction, delay or inhibition in an undesired immune response specific to the antigen or cell, tissue, organ, etc. that expresses such antigen. Such immune responses also include any stimulation, production, induction, promotion or recruitment in a desired immune response specific to the antigen or cell, tissue, organ, etc. that expresses such antigen. Tolerogenic immune responses, therefore, include the absence of or reduction in an undesired immune response to an antigen that can be mediated by antigen reactive cells as well as the presence or promotion of suppressive cells. Tolerogenic immune responses as provided herein include immunological tolerance. To "generate a tolerogenic immune response" refers to the generation of any of the foregoing immune responses specific to an antigen or cell, tissue, organ, etc. that expresses such antigen. The tolerogenic immune response can be the result of MHC Class I-restricted presentation and/or MHC Class II-restricted presentation and/or B cell presentation and/or presentation by CD1d, etc. In embodiments, to ensure that a composition comprises desired epitopes, antigens are selected such that they comprise such epitopes for coupling to the synthetic nanocarriers as provided herein. In other embodiments, to ensure that a composition comprises such epitopes, synthetic nanocarriers coupled to an antigen are produced and tested for regulatory B cell immune responses, such as activation or generation. The appropriate synthetic nanocarriers may then be selected.

Tolerogenic immune responses include any reduction, delay or inhibition in CD4+ T cell, CD8+ T cell or B cell proliferation and/or activity. Tolerogenic immune responses also

include a reduction in antigen-specific antibody production. Tolerogenic immune responses can also include any response that leads to the stimulation, induction, production or recruitment of regulatory cells, such as CD4+ Treg cells, CD8+ Treg cells, Breg cells, etc. In some embodiments, the tolerogenic immune response, is one that results in the conversion to a regulatory phenotype characterized by the production, induction, stimulation or recruitment of regulatory cells.

Tolerogenic immune responses also include any response that leads to the stimulation, production or recruitment of CD4+ Treg cells and/or CD8+ Treg cells. CD4+ Treg cells can express the transcription factor FoxP3 and inhibit inflammatory responses and auto-immune inflammatory diseases (Human regulatory T cells in autoimmune diseases. Cvetanovich GL, Hafler DA. Curr Opin Immunol. 2010 Dec;22(6):753-60. Regulatory T cells and autoimmunity. Vila J, Isaacs JD, Anderson AE.Curr Opin Hematol. 2009 Jul;16(4):274-9). Such cells also suppress T-cell help to B-cells and induce tolerance to both self and foreign antigens (Therapeutic approaches to allergy and autoimmunity based on FoxP3+ regulatory T-cell activation and expansion. Miyara M, Wing K, Sakaguchi S. J Allergy Clin Immunol. 2009 Apr;123(4):749-55). CD4+ Treg cells recognize antigen when presented by Class II proteins on APCs. CD8+ Treg cells, which recognize antigen presented by Class I (and Qa-1), can also suppress T-cell help to B-cells and result in activation of antigen-specific suppression inducing tolerance to both self and foreign antigens. Disruption of the interaction of Qa-1 with CD8+ Treg cells has been shown to dysregulate immune responses and results in the development of auto-antibody formation and an auto-immune lethal systemic-lupus-erythematosus (Kim et al., Nature. 2010 Sep 16, 467 (7313): 328-32). CD8+ Treg cells have also been shown to inhibit models of autoimmune inflammatory diseases including rheumatoid arthritis and colitis (CD4+CD25+ regulatory T cells in autoimmune arthritis. Oh S, Rankin AL, Caton AJ. Immunol Rev. 2010 Jan;233(1):97-111. Regulatory T cells in inflammatory bowel disease. Boden EK, Snapper SB. Curr Opin Gastroenterol. 2008 Nov;24(6):733-41). In some embodiments, the compositions provided can effectively result in both types of responses (CD4+ Treg and CD8+ Treg). In other embodiments, FoxP3 can be induced in other immune cells, such as macrophages, iNKT cells, etc., and the compositions provided herein can result in one or more of these responses as well.

Tolerogenic immune responses also include, but are not limited to, the induction of regulatory cytokines, such as Treg cytokines; induction of inhibitory cytokines; the inhibition of inflammatory cytokines (e.g., IL-4, IL-1b, IL-5, TNF-α, IL-6, GM-CSF, IFN-γ, IL-2, IL-9,

IL-12, IL-17, IL-18, IL-21, IL-22, IL-23, M-CSF, C reactive protein, acute phase protein, chemokines (e.g., MCP-1, RANTES, MIP-1α, MIP-1β, MIG, ITAC or IP-10), the production of anti-inflammatory cytokines (e.g., IL-4, IL-13, IL-10, etc.), chemokines (e.g., CCL-2, CXCL8), proteases (e.g., MMP-3, MMP-9), leukotrienes (e.g., CysLT-1, CysLT-2), prostaglandins (e.g., PGE2) or histamines; the inhibition of polarization to a Th17, Th1 or Th2 immune response; the inhibition of effector cell-specific cytokines: Th17 (e.g., IL-17, IL-25), Th1 (IFN-γ), Th2 (e.g., IL-4, IL-13); the inhibition of Th1-, Th2- or TH17-specific transcription factors; the inhibition of proliferation of effector T cells; the induction of apoptosis of effector T cells; the induction of tolerogenic dendritic cell-specific genes, the induction of FoxP3 expression, the inhibition of IgE induction or IgE-mediated immune responses; the inhibition of antibody responses (e.g., antigen-specific antibody production); the inhibition of T helper cell response; the production of TGF-β and/or IL-10; the inhibition of effector function of autoantibodies (e.g., inhibition in the depletion of cells, cell or tissue damage or complement activation); etc.

Any of the foregoing may be measured *in vivo* in one or more animal models or may be measured *in vitro*. One of ordinary skill in the art is familiar with such *in vivo* or *in vitro* measurements. Undesired immune responses or tolerogenic immune responses can be monitored using, for example, methods of assessing immune cell number and/or function, tetramer analysis, ELISPOT, flow cytometry-based analysis of cytokine expression, cytokine secretion, cytokine expression profiling, gene expression profiling, protein expression profiling, analysis of cell surface markers, PCR-based detection of immune cell receptor gene usage (see T. Clay et al., "Assays for Monitoring Cellular Immune Response to Active Immunotherapy of Cancer" Clinical Cancer Research 7:1127-1135 (2001)), etc. Undesired immune responses or tolerogenic immune responses may also be monitored using, for example, methods of assessing protein levels in plasma or serum, immune cell proliferation and/or functional assays, etc. In some embodiments, tolerogenic immune responses can be monitored by assessing the induction of FoxP3. In addition, specific methods are described in more detail in the Examples.

Preferably, tolerogenic immune responses lead to the inhibition of the development, progression or pathology of the diseases, disorders or conditions described herein. Whether or not the inventive compositions can lead to the inhibition of the development, progression or pathology of the diseases, disorders or conditions described herein can be measured with

animal models of such diseases, disorders or conditions. In some embodiments, the reduction of an undesired immune response or generation of a tolerogenic immune response may be assessed by determining clinical endpoints, clinical efficacy, clinical symptoms, disease biomarkers and/or clinical scores. Undesired immune responses or tolerogenic immune responses can also be assessed with diagnostic tests to assess the presence or absence of a disease, disorder or condition as provided herein. Undesired or desired immune responses can further be assessed by methods of measuring therapeutic proteins levels and/or function in a subject. In embodiments, methods for monitoring or assessing undesired allergic responses include assessing an allergic response in a subject by skin reactivity and/or allergen-specific antibody production.

In some embodiments, monitoring or assessing the generation of an undesired immune response or a tolerogenic immune response in a subject can be prior to the administration of a composition of synthetic nanocarriers provided herein and/or prior to administration of a transplantable graft or therapeutic protein or exposure to an allergen. In other embodiments, assessing the generation of an undesired immune response or tolerogenic immune response can be after administration of a composition of synthetic nanocarriers provided herein and/or after administration of a transplantable graft or therapeutic protein or exposure to an allergen. In some embodiments, the assessment is done after administration of the composition of synthetic nanocarriers, but prior to administration of a transplantable graft or therapeutic protein or exposure to an allergen. In other embodiments, the assessment is done after administration of a transplantable graft or therapeutic protein or exposure to an allergen, but prior to administration of the composition. In still other embodiments, the assessment is performed prior to both the administration of the synthetic nanocarriers and administration of a transplantable graft or therapeutic protein or exposure to an allergen, while in yet other embodiments the assessment is performed after both the administration of synthetic nanocarriers and the administration of a transplantable graft or therapeutic protein or exposure to an allergen. In further embodiments, the assessment is performed both prior to and after the administration of the synthetic nanocarriers and/or administration of a transplantable graft or therapeutic protein or exposure to an allergen. In still other embodiments, the assessment is performed more than once on the subject to determine that a desirable immune state is maintained in the subject, such as a subject that has or is at risk of having an inflammatory disease, an autoimmune disease, an allergy, organ or tissue rejection or graft verus host disease. Other subjects include those that have undergone or will undergo

transplantation as well as those that have received, are receiving or will receive a therapeutic protein against which they have experienced, are experiencing or are expected to experience an undesired immune response.

An antibody response can be assessed by determining one or more antibody titers. "Antibody titer" means a measurable level of antibody production. Methods for measuring antibody titers are known in the art and include Enzyme-linked Immunosorbent Assay (ELISA). In embodiments, the antibody response can be quantitated, for example, as the number of antibodies, concentration of antibodies or titer. The values can be absolute or they can be relative. Assays for quantifying an antibody response include antibody capture assays, enzyme-linked immunosorbent assays (ELISAs), inhibition liquid phase absorption assays (ILPAAs), rocket immunoelectrophoresis (RIE) assays and line immunoelectrophoresis (LIE) assays. When an antibody response is compared to another antibody response the same type of quantitative value (e.g., titer) and method of measurement (e.g., ELISA) is preferably used to make the comparison.

An ELISA method for measuring an antibody titer, for example, a typical sandwich ELISA, may consist of the following steps (i) preparing an ELISA-plate coating material such that the antibody target of interest is coupled to a substrate polymer or other suitable material (ii) preparing the coating material in an aqueous solution (such as PBS) and delivering the coating material solution to the wells of a multiwell plate for overnight deposition of the coating onto the multiwell plate (iii) thoroughly washing the multiwell plate with wash buffer (such as 0.05% Tween-20 in PBS) to remove excess coating material (iv) blocking the plate for nonspecific binding by applying a diluent solution (such as 10% fetal bovine serum in PBS), (v) washing the blocking/diluent solution from the plate with wash buffer (vi) diluting the serum sample(s) containing antibodies and appropriate standards (positive controls) with diluent as required to obtain a concentration that suitably saturates the ELISA response (vii) serially diluting the plasma samples on the multiwell plate such to cover a range of concentrations suitable for generating an ELISA response curve (viii) incubating the plate to provide for antibody-target binding (ix) washing the plate with wash buffer to remove antibodies not bound to antigen (x) adding an appropriate concentration of a secondary detection antibody in same diluent such as a biotin-coupled detection antibody capable of binding the primary antibody (xi) incubating the plate with the applied detection antibody, followed by washing with wash buffer (xii) adding an enzyme such as streptavidin-HRP (horse radish peroxidase) that will bind to biotin found on biotinylated antibodies and

incubating (xiii) washing the multiwell plate (xiv) adding substrate(s) (such as TMB solution) to the plate (xv) applying a stop solution (such as 2N sulfuric acid) when color development is complete (xvi) reading optical density of the plate wells at a specific wavelength for the substrate (450 nm with subtraction of readings at 570 nm) (xvi) applying a suitable multiparameter curve fit to the data and defining half-maximal effective concentration (EC50) as the concentration on the curve at which half the maximum OD value for the plate standards is achieved.

A "transplantable graft" refers to a biological material, such as cells, tissues and organs (in whole or in part) that can be administered to a subject. Transplantable grafts may be autografts, allografts, or xenografts of, for example, a biological material such as an organ, tissue, skin, bone, nerves, tendon, neurons, blood vessels, fat, cornea, pluripotent cells, differentiated cells (obtained or derived in vivo or in vitro), etc. In some embodiments, a transplantable graft is formed, for example, from cartilage, bone, extracellular matrix, or collagen matrices. Transplantable grafts may also be single cells, suspensions of cells and cells in tissues and organs that can be transplanted. Transplantable cells typically have a therapeutic function, for example, a function that is lacking or diminished in a recipient subject. Some non-limiting examples of transplantable cells are β-cells, hepatocytes, hematopoietic stem cells, neuronal stem cells, neurons, glial cells, or myelinating cells. Transplantable cells can be cells that are unmodified, for example, cells obtained from a donor subject and usable in transplantation without any genetic or epigenetic modifications. In other embodiments, transplantable cells can be modified cells, for example, cells obtained from a subject having a genetic defect, in which the genetic defect has been corrected, or cells that are derived from reprogrammed cells, for example, differentiated cells derived from cells obtained from a subject.

"Transplantation" refers to the process of transferring (moving) a transplantable graft into a recipient subject (e.g., from a donor subject, from an in vitro source (e.g.,differentiated autologous or heterologous native or induced pluripotent cells)) and/or from one bodily location to another bodily location in the same subject.

"Undesired immune response" refers to any undesired immune response that results from exposure to an antigen, promotes or exacerbates a disease, disorder or condition provided herein (or a symptom thereof), or is symptomatic of a disease, disorder or condition provided herein. Such immune responses generally have a negative impact on a subject's health or is symptomatic of a negative impact on a subject's health.

C. INVENTIVE COMPOSITIONS

Provided herein are methods and compositions for generating regulatory B cells. Regulatory B cells are believed to modulate immune responses, for example, in that IL-10 release from regulatory B cells exerts anti-inflammatory and immunosuppressive effects on most hematopoietic cells. IL-10 also suppresses pro-inflammatory cytokine production by monocytes and macrophages and the proliferation of antigen-specific CD4+ T cells. Without wishing to be bound by any particular theory, it is believed that the compositions provided herein can affect the number and/or activity of regulatory B cells in vitro and/or in vivo by, for example, interacting with naïve B cells or regulatory B cell precursors, resulting in an induction of regulatory B cell maturation or an increase in regulatory B cell proliferation and/or activity. The effect on regulatory B cell number and/or activity may also be a result of the generation of other regulatory cell or tolerogenic immune responses, such as through regulatory T cell production and/or activation.

Provided herein are tolerogenic methods that include the administration of synthetic nanocarrier compositions comprising immunosuppressants and B cell and/or MHC Class II-restricted epitopes of an antigen and related compositions. Such methods and compositions are useful for generating, stimulating or recruiting regulatory B cells and promoting the generation of tolerogenic immune responses. The methods provided may also include the administration therapeutic proteins and/or transplantable grafts. The compositions provided, therefore, may also include the therapeutic proteins and/or transplantable grafts. The compositions may be administered to subjects in which a tolerogenic immune response is desired. Such subjects include those that have or are at risk of having an inflammatory disease, an autoimmune disease, an allergy, organ or tissue rejection or graft versus host disease. Such subjects also include those that have been, are being or will be administered a therapeutic protein against which the subject has experienced or is expected to experience an undesired immune response. Such subjects also include those that have undergone or will undergo transplantation.

Preferably, the compositions of the invention result in the generation, recruitment or activation of regulatory B cells. One hallmark of regulatory B cells is the ability to produce and secrete IL-10. Some regulatory B cells are CD1d+ or CD1dhigh. Some regulatory B cells are CD5+ and/or CD19+. For example, some regulatory B cells are CD1dhighCD5+CD19+. Some regulatory B cells are CD24+ or CD24high; and/or CD38+ or

CD38high. For example, some regulatory B cells are CD19+CD24highCD38high. Some regulatory B cells produce and secrete IL-10. Additional surface markers and chemokine secretion profiles that can be used to identify regulatory B cells are known to those of skill in the art. Based on the knowledge of surface markers useful for the identification of various regulatory B cell populations, those of skill in the art are able to identify and enumerate regulatory B cells in a heterogeneous population of cells, for example, in a population of cells in culture or in a population of cells obtained from a subject.

As mentioned above, the synthetic nanocarriers are designed to comprise B cell eptiopes and/or MHC Class II-restricted epitopes and/or immunosuppressants. A wide variety of synthetic nanocarriers can be used according to the invention. In some embodiments, synthetic nanocarriers are spheres or spheroids. In some embodiments, synthetic nanocarriers are flat or plate-shaped. In some embodiments, synthetic nanocarriers are cubes or cubic. In some embodiments, synthetic nanocarriers are ovals or ellipses. In some embodiments, synthetic nanocarriers are cylinders, cones, or pyramids.

In some embodiments, it is desirable to use a population of synthetic nanocarriers that is relatively uniform in terms of size, shape, and/or composition so that each synthetic nanocarrier has similar properties. For example, at least 80%, at least 90%, or at least 95% of the synthetic nanocarriers, based on the total number of synthetic nanocarriers, may have a minimum dimension or maximum dimension that falls within 5%, 10%, or 20% of the average diameter or average dimension of the synthetic nanocarriers. In some embodiments, a population of synthetic nanocarriers may be heterogeneous with respect to size, shape, and/or composition.

Synthetic nanocarriers can be solid or hollow and can comprise one or more layers. In some embodiments, each layer has a unique composition and unique properties relative to the other layer(s). To give but one example, synthetic nanocarriers may have a core/shell structure, wherein the core is one layer (e.g. a polymeric core) and the shell is a second layer (e.g. a lipid bilayer or monolayer). Synthetic nanocarriers may comprise a plurality of different layers.

In some embodiments, synthetic nanocarriers may optionally comprise one or more lipids. In some embodiments, a synthetic nanocarrier may comprise a liposome. In some embodiments, a synthetic nanocarrier may comprise a lipid bilayer. In some embodiments, a synthetic nanocarrier may comprise a lipid monolayer. In some embodiments, a synthetic nanocarrier may comprise a micelle. In some embodiments, a synthetic nanocarrier may

comprise a core comprising a polymeric matrix surrounded by a lipid layer (e.g., lipid bilayer, lipid monolayer, etc.). In some embodiments, a synthetic nanocarrier may comprise a non-polymeric core (e.g., metal particle, quantum dot, ceramic particle, bone particle, viral particle, proteins, nucleic acids, carbohydrates, etc.) surrounded by a lipid layer (e.g., lipid bilayer, lipid monolayer, etc.).

In other embodiments, synthetic nanocarriers may comprise metal particles, quantum dots, ceramic particles, etc. In some embodiments, a non-polymeric synthetic nanocarrier is an aggregate of non-polymeric components, such as an aggregate of metal atoms (e.g., gold atoms).

In some embodiments, synthetic nanocarriers may optionally comprise one or more amphiphilic entities. In some embodiments, an amphiphilic entity can promote the production of synthetic nanocarriers with increased stability, improved uniformity, or increased viscosity. In some embodiments, amphiphilic entities can be associated with the interior surface of a lipid membrane (e.g., lipid bilayer, lipid monolayer, etc.). Many amphiphilic entities known in the art are suitable for use in making synthetic nanocarriers in accordance with the present invention. Such amphiphilic entities include, but are not limited to, phosphoglycerides; phosphatidylcholines; dipalmitoyl phosphatidylcholine (DPPC); dioleylphosphatidyl ethanolamine (DOPE); dioleyloxypropyltriethylammonium (DOTMA); dioleoylphosphatidylcholine; cholesterol; cholesterol ester; diacylglycerol; diacylglycerolsuccinate; diphosphatidyl glycerol (DPPG); hexanedecanol; fatty alcohols such as polyethylene glycol (PEG); polyoxyethylene-9-lauryl ether; a surface active fatty acid, such as palmitic acid or oleic acid; fatty acids; fatty acid monoglycerides; fatty acid diglycerides; fatty acid amides; sorbitan trioleate (Span®85) glycocholate; sorbitan monolaurate (Span®20); polysorbate 20 (Tween®20); polysorbate 60 (Tween®60); polysorbate 65 (Tween®65); polysorbate 80 (Tween®80); polysorbate 85 (Tween®85); polyoxyethylene monostearate; surfactin; a poloxomer; a sorbitan fatty acid ester such as sorbitan trioleate; lecithin; lysolecithin; phosphatidylserine; phosphatidylinositol;sphingomyelin; phosphatidylethanolamine (cephalin); cardiolipin; phosphatidic acid; cerebrosides; dicetylphosphate; dipalmitoylphosphatidylglycerol; stearylamine; dodecylamine; hexadecyl-amine; acetyl palmitate; glycerol ricinoleate; hexadecyl sterate; isopropyl myristate; tyloxapol; poly(ethylene glycol)5000-phosphatidylethanolamine; poly(ethylene glycol)400-monostearate; phospholipids; synthetic and/or natural detergents having high surfactant properties; deoxycholates; cyclodextrins;

chaotropic salts; ion pairing agents; and combinations thereof. An amphiphilic entity component may be a mixture of different amphiphilic entities. Those skilled in the art will recognize that this is an exemplary, not comprehensive, list of substances with surfactant activity. Any amphiphilic entity may be used in the production of synthetic nanocarriers to be used in accordance with the present invention.

In some embodiments, synthetic nanocarriers may optionally comprise one or more carbohydrates. Carbohydrates may be natural or synthetic. A carbohydrate may be a derivatized natural carbohydrate. In certain embodiments, a carbohydrate comprises monosaccharide or disaccharide, including but not limited to glucose, fructose, galactose, ribose, lactose, sucrose, maltose, trehalose, cellbiose, mannose, xylose, arabinose, glucoronic acid, galactoronic acid, mannuronic acid, glucosamine, galatosamine, and neuramic acid. In certain embodiments, a carbohydrate is a polysaccharide, including but not limited to pullulan, cellulose, microcrystalline cellulose, hydroxypropyl methylcellulose (HPMC), hydroxycellulose (HC), methylcellulose (MC), dextran, cyclodextran, glycogen, hydroxyethylstarch, carageenan, glycon, amylose, chitosan, N,O-carboxylmethylchitosan, algin and alginic acid, starch, chitin, inulin, konjac, glucommannan, pustulan, heparin, hyaluronic acid, curdlan, and xanthan. In embodiments, the inventive synthetic nanocarriers do not comprise (or specifically exclude) carbohydrates, such as a polysaccharide. In certain embodiments, the carbohydrate may comprise a carbohydrate derivative such as a sugar alcohol, including but not limited to mannitol, sorbitol, xylitol, erythritol, maltitol, and lactitol.

In some embodiments, synthetic nanocarriers can comprise one or more polymers. In some embodiments, the synthetic nanocarriers comprise one or more polymers that is a nonmethoxy-terminated, pluronic polymer. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% (weight/weight) of the polymers that make up the synthetic nanocarriers are non-methoxy-terminated, pluronic polymers. In some embodiments, all of the polymers that make up the synthetic nanocarriers are non-methoxy-terminated, pluronic polymers. In some embodiments, the synthetic nanocarriers can comprise one or more polymers that is a non-methoxy-terminated polymer. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% (weight/weight) of the polymers that make up the synthetic nanocarriers are nonmethoxy-terminated polymers. In some embodiments, all of the

polymers that make up the synthetic nanocarriers are non-methoxy-terminated polymers. In some embodiments, the synthetic nanocarriers comprise one or more polymers that do not comprise pluronic polymer. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% (weight/weight) of the polymers that make up the synthetic nanocarriers do not comprise pluronic polymer. In some embodiments, all of the polymers that make up the synthetic nanocarriers do not comprise pluronic polymer. In some embodiments, such a polymer can be surrounded by a coating layer (e.g., liposome, lipid monolayer, micelle, etc.). In some embodiments, various elements of the synthetic nanocarriers can be coupled with the polymer.

The immunosuppressants and/or antigens can be coupled to the synthetic nanocarriers by any of a number of methods. Generally, the coupling can be a result of bonding between the immunosuppressants and/or antigens and the synthetic nanocarriers. This bonding can result in the immunosuppressants and/or antigens being attached to the surface of the synthetic nanocarrierss and/or contained within (encapsulated) the synthetic nanocarriers. In some embodiments, however, the immunosuppressants and/or antigens are encapsulated by the synthetic nanocarriers as a result of the structure of the synthetic nanocarriers rather than bonding to the synthetic nanocarriers. In preferable embodiments, the synthetic nanocarriers comprise a polymer as provided herein, and the immunosuppressants and/or antigens are coupled to the polymer.

When coupling occurs as a result of bonding between the immunosuppressants and/or antigens and synthetic nanocarriers, the coupling may occur via a coupling moiety. A coupling moiety can be any moiety through which an immunosuppressant and/or antigen is bonded to a synthetic nanocarrier. Such moieties include covalent bonds, such as an amide bond or ester bond, as well as separate molecules that bond (covalently or non-covalently) the immunosuppressant and/or antigen to the synthetic nanocarrier. Such molecules include linkers or polymers or a unit thereof. For example, the coupling moiety can comprise a charged polymer to which an immunosuppressant and/or antigen electrostatically binds. As another example, the coupling moiety can comprise a polymer or unit thereof to which it is covalently bonded.

In preferred embodiments, the synthetic nanocarriers comprise a polymer as provided herein. These synthetic nanocarriers can be completely polymeric or they can be a mix of polymers and other materials.

In some embodiments, the polymers of a synthetic nanocarrier associate to form a polymeric matrix. In some of these embodiments, a component, such as an immunosuppressant or antigen, can be covalently associated with one or more polymers of the polymeric matrix. In some embodiments, covalent association is mediated by a linker. In some embodiments, a component can be noncovalently associated with one or more polymers of the polymeric matrix. For example, in some embodiments a component can be encapsulated within, surrounded by, and/or dispersed throughout a polymeric matrix. Alternatively or additionally, a component can be associated with one or more polymers of a polymeric matrix by hydrophobic interactions, charge interactions, van der Waals forces, etc. A wide variety of polymers and methods for forming polymeric matrices therefrom are known conventionally.

Polymers may be natural or unnatural (synthetic) polymers. Polymers may be homopolymers or copolymers comprising two or more monomers. In terms of sequence, copolymers may be random, block, or comprise a combination of random and block sequences. Typically, polymers in accordance with the present invention are organic polymers.

In some embodiments, the polymer comprises a polyester, polycarbonate, polyamide, or polyether, or unit thereof. In other embodiments, the polymer comprises poly(ethylene glycol) (PEG), polypropylene glycol, poly(lactic acid), poly(glycolic acid), poly(lactic-co-glycolic acid), or a polycaprolactone, or unit thereof. In some embodiments, it is preferred that the polymer is biodegradable. Therefore, in these embodiments, it is preferred that if the polymer comprises a polyether, such as poly(ethylene glycol) or polypropylene glycol or unit thereof, the polymer comprises a block-co-polymer of a polyether and a biodegradable polymer such that the polymer is biodegradable. In other embodiments, the polymer does not solely comprise a polyether or unit thereof, such as poly(ethylene glycol) or polypropylene glycol or unit thereof.

Other examples of polymers suitable for use in the present invention include, but are not limited to polyethylenes, polycarbonates (e.g. poly(1,3-dioxan-2one)), polyanhydrides (e.g. poly(sebacic anhydride)), polypropylfumerates, polyamides (e.g. polycaprolactam), polyacetals, polyethers, polyesters (e.g., polylactide, polyglycolide, polylactide-co-glycolide, polycaprolactone, polyhydroxyacid (e.g. poly(β-hydroxyalkanoate))), poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates,

polymethacrylates, polyureas, polystyrenes, and polyamines, polylysine, polylysine-PEG copolymers, and poly(ethyleneimine), poly(ethylene imine)-PEG copolymers.

In some embodiments, polymers in accordance with the present invention include polymers which have been approved for use in humans by the U.S. Food and Drug Administration (FDA) under 21 C.F.R. § 177.2600, including but not limited to polyesters (e.g., polylactic acid, poly(lactic-co-glycolic acid), polycaprolactone, polyvalerolactone, poly(1,3-dioxan-2one)); polyanhydrides (e.g., poly(sebacic anhydride)); polyethers (e.g., polyethylene glycol); polyurethanes; polymethacrylates; polyacrylates; and polycyanoacrylates.

In some embodiments, polymers can be hydrophilic. For example, polymers may comprise anionic groups (e.g., phosphate group, sulphate group, carboxylate group); cationic groups (e.g., quaternary amine group); or polar groups (e.g., hydroxyl group, thiol group, amine group). In some embodiments, a synthetic nanocarrier comprising a hydrophilic polymeric matrix generates a hydrophilic environment within the synthetic nanocarrier. In some embodiments, polymers can be hydrophobic. In some embodiments, a synthetic nanocarrier comprising a hydrophobic polymeric matrix generates a hydrophobic environment within the synthetic nanocarrier. Selection of the hydrophilicity or hydrophobicity of the polymer may have an impact on the nature of materials that are incorporated (e.g. coupled) within the synthetic nanocarrier.

In some embodiments, polymers may be modified with one or more moieties and/or functional groups. A variety of moieties or functional groups can be used in accordance with the present invention. In some embodiments, polymers may be modified with polyethylene glycol (PEG), with a carbohydrate, and/or with acyclic polyacetals derived from polysaccharides (Papisov, 2001, ACS Symposium Series, 786:301). Certain embodiments may be made using the general teachings of US Patent No. 5543158 to Gref et al., or WO publication WO2009/051837 by Von Andrian et al.

In some embodiments, polymers may be modified with a lipid or fatty acid group. In some embodiments, a fatty acid group may be one or more of butyric, caproic, caprylic, capric, lauric, myristic, palmitic, stearic, arachidic, behenic, or lignoceric acid. In some embodiments, a fatty acid group may be one or more of palmitoleic, oleic, vaccenic, linoleic, alpha-linoleic, gamma-linoleic, arachidonic, gadoleic, arachidonic, eicosapentaenoic, docosahexaenoic, or erucic acid.

In some embodiments, polymers may be polyesters, including copolymers comprising lactic acid and glycolic acid units, such as poly(lactic acid-co-glycolic acid) and poly(lactide-co-glycolide), collectively referred to herein as "PLGA"; and homopolymers comprising glycolic acid units, referred to herein as "PGA," and lactic acid units, such as poly-L-lactic acid, poly-D-lactic acid, poly-D,L-lactic acid, poly-L-lactide, poly-D-lactide, and poly-D,L-lactide, collectively referred to herein as "PLA." In some embodiments, exemplary polyesters include, for example, polyhydroxyacids; PEG copolymers and copolymers of lactide and glycolide (e.g., PLA-PEG copolymers, PGA-PEG copolymers, PLGA-PEG copolymers, and derivatives thereof. In some embodiments, polyesters include, for example, poly(caprolactone), poly(caprolactone)-PEG copolymers, poly(L-lactide-co-L-lysine), poly(serine ester), poly(4-hydroxy-L-proline ester), poly[α-(4-aminobutyl)-L-glycolic acid], and derivatives thereof.

In some embodiments, a polymer may be PLGA. PLGA is a biocompatible and biodegradable co-polymer of lactic acid and glycolic acid, and various forms of PLGA are characterized by the ratio of lactic acid:glycolic acid. Lactic acid can be L-lactic acid, D-lactic acid, or D,L-lactic acid. The degradation rate of PLGA can be adjusted by altering the lactic acid:glycolic acid ratio. In some embodiments, PLGA to be used in accordance with the present invention is characterized by a lactic acid:glycolic acid ratio of approximately 85:15, approximately 75:25, approximately 60:40, approximately 50:50, approximately 40:60, approximately 25:75, or approximately 15:85.

In some embodiments, polymers may be one or more acrylic polymers. In certain embodiments, acrylic polymers include, for example, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, aminoalkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), methacrylic acid alkylamide copolymer, poly(methyl methacrylate), poly(methacrylic acid anhydride), methyl methacrylate, polymethacrylate, poly(methyl methacrylate) copolymer, polyacrylamide, aminoalkyl methacrylate copolymer, glycidyl methacrylate copolymers, polycyanoacrylates, and combinations comprising one or more of the foregoing polymers. The acrylic polymer may comprise fully-polymerized copolymers of acrylic and methacrylic acid esters with a low content of quaternary ammonium groups.

In some embodiments, polymers can be cationic polymers. In general, cationic polymers are able to condense and/or protect negatively charged strands of nucleic acids (e.g. DNA, or derivatives thereof). Amine-containing polymers such as poly(lysine) (Zauner et al.,

1998, Adv. Drug Del. Rev., 30:97; and Kabanov et al., 1995, Bioconjugate Chem., 6:7), poly(ethylene imine) (PEI; Boussif et al., 1995, Proc. Natl. Acad. Sci., USA, 1995, 92:7297), and poly(amidoamine) dendrimers (Kukowska-Latallo et al., 1996, Proc. Natl. Acad. Sci., USA, 93:4897; Tang et al., 1996, Bioconjugate Chem., 7:703; and Haensler et al., 1993, Bioconjugate Chem., 4:372) are positively-charged at physiological pH, form ion pairs with nucleic acids, and mediate transfection in a variety of cell lines. In embodiments, the inventive synthetic nanocarriers may not comprise (or may exclude) cationic polymers.

In some embodiments, polymers can be degradable polyesters bearing cationic side chains (Putnam et al., 1999, Macromolecules, 32:3658; Barrera et al., 1993, J. Am. Chem. Soc., 115:11010; Kwon et al., 1989, Macromolecules, 22:3250; Lim et al., 1999, J. Am. Chem. Soc., 121:5633; and Zhou et al., 1990, Macromolecules, 23:3399). Examples of these polyesters include poly(L-lactide-co-L-lysine) (Barrera et al., 1993, J. Am. Chem. Soc., 115:11010), poly(serine ester) (Zhou et al., 1990, Macromolecules, 23:3399), poly(4-hydroxy-L-proline ester) (Putnam et al., 1999, Macromolecules, 32:3658; and Lim et al., 1999, J. Am. Chem. Soc., 121:5633), and poly(4-hydroxy-L-proline ester) (Putnam et al., 1999, Macromolecules, 32:3658; and Lim et al., 1999, J. Am. Chem. Soc., 121:5633).

The properties of these and other polymers and methods for preparing them are well known in the art (see, for example, U.S. Patents 6,123,727; 5,804,178; 5,770,417; 5,736,372; 5,716,404; 6,095,148; 5,837,752; 5,902,599; 5,696,175; 5,514,378; 5,512,600; 5,399,665; 5,019,379; 5,010,167; 4,806,621; 4,638,045; and 4,946,929; Wang et al., 2001, J. Am. Chem. Soc., 123:9480; Lim et al., 2001, J. Am. Chem. Soc., 123:2460; Langer, 2000, Acc. Chem. Res., 33:94; Langer, 1999, J. Control. Release, 62:7; and Uhrich et al., 1999, Chem. Rev., 99:3181). More generally, a variety of methods for synthesizing certain suitable polymers are described in Concise Encyclopedia of Polymer Science and Polymeric Amines and Ammonium Salts, Ed. by Goethals, Pergamon Press, 1980; Principles of Polymerization by Odian, John Wiley & Sons, Fourth Edition, 2004; Contemporary Polymer Chemistry by Allcock et al., Prentice-Hall, 1981; Deming et al., 1997, Nature, 390:386; and in U.S. Patents 6,506,577, 6,632,922, 6,686,446, and 6,818,732.

In some embodiments, polymers can be linear or branched polymers. In some embodiments, polymers can be dendrimers. In some embodiments, polymers can be substantially cross-linked to one another. In some embodiments, polymers can be substantially free of cross-links. In some embodiments, polymers can be used in accordance with the present invention without undergoing a cross-linking step. It is further to be

understood that inventive synthetic nanocarriers may comprise block copolymers, graft copolymers, blends, mixtures, and/or adducts of any of the foregoing and other polymers. Those skilled in the art will recognize that the polymers listed herein represent an exemplary, not comprehensive, list of polymers that can be of use in accordance with the present invention.

Compositions according to the invention comprise synthetic nanocarriers in combination with pharmaceutically acceptable excipients, such as preservatives, buffers, saline, or phosphate buffered saline. The compositions may be made using conventional pharmaceutical manufacturing and compounding techniques to arrive at useful dosage forms. In an embodiment, inventive synthetic nanocarriers are suspended in sterile saline solution for injection together with a preservative.

In embodiments, when preparing synthetic nanocarriers as carriers, methods for coupling components to the synthetic nanocarriers may be useful. If the component is a small molecule it may be of advantage to attach the component to a polymer prior to the assembly of the synthetic nanocarriers. In embodiments, it may also be an advantage to prepare the synthetic nanocarriers with surface groups that are used to couple the component to the synthetic nanocarrier through the use of these surface groups rather than attaching the component to a polymer and then using this polymer conjugate in the construction of synthetic nanocarriers.

In certain embodiments, the coupling can be a covalent linker. In embodiments, peptides according to the invention can be covalently coupled to the external surface via a 1,2,3-triazole linker formed by the 1,3-dipolar cycloaddition reaction of azido groups on the surface of the nanocarrier with antigen or immunosuppressant containing an alkyne group or by the 1,3-dipolar cycloaddition reaction of alkynes on the surface of the nanocarrier with components containing an azido group. Such cycloaddition reactions are preferably performed in the presence of a Cu(I) catalyst along with a suitable Cu(I)-ligand and a reducing agent to reduce Cu(II) compound to catalytic active Cu(I) compound. This Cu(I)-catalyzed azide-alkyne cycloaddition (CuAAC) can also be referred as the click reaction.

Additionally, the covalent coupling may comprise a covalent linker that comprises an amide linker, a disulfide linker, a thioether linker, a hydrazone linker, a hydrazide linker, an imine or oxime linker, an urea or thiourea linker, an amidine linker, an amine linker, and a sulfonamide linker.

An amide linker is formed via an amide bond between an amine on one component with the carboxylic acid group of a second component such as the nanocarrier. The amide bond in the linker can be made using any of the conventional amide bond forming reactions with suitably protected amino acids or components and activated carboxylic acid such N-hydroxysuccinimide-activated ester.

A disulfide linker is made via the formation of a disulfide (S-S) bond between two sulfur atoms of the form, for instance, of R1-S-S-R2. A disulfide bond can be formed by thiol exchange of a component containing thiol/mercaptan group(-SH) with another activated thiol group on a polymer or nanocarrier or a nanocarrier containing thiol/mercaptan groups with a component containing activated thiol group.



A triazole linker, specifically a 1,2,3-triazole of the form , wherein R1 and R2 may be any chemical entities, is made by the 1,3-dipolar cycloaddition reaction of an azide attached to a first component such as the nanocarrier with a terminal alkyne attached to a second component such as the immunosuppressant or antigen. The 1,3-dipolar cycloaddition reaction is performed with or without a catalyst, preferably with Cu(I)-catalyst, which links the two components through a 1,2,3-triazole function. This chemistry is described in detail by Sharpless et al., Angew. Chem. Int. Ed. 41(14), 2596, (2002) and Meldal, et al, Chem. Rev., 2008, 108(8), 2952-3015 and is often referred to as a "click" reaction or CuAAC.

In embodiments, a polymer containing an azide or alkyne group, terminal to the polymer chain is prepared. This polymer is then used to prepare a synthetic nanocarrier in such a manner that a plurality of the alkyne or azide groups are positioned on the surface of that nanocarrier. Alternatively, the synthetic nanocarrier can be prepared by another route, and subsequently functionalized with alkyne or azide groups. The component is prepared with the presence of either an alkyne (if the polymer contains an azide) or an azide (if the polymer contains an alkyne) group. The component is then allowed to react with the nanocarrier via the 1,3-dipolar cycloaddition reaction with or without a catalyst which covalently couples the component to the particle through the 1,4-disubstituted 1,2,3-triazole linker.

A thioether linker is made by the formation of a sulfur-carbon (thioether) bond in the form, for instance, of R1-S-R2. Thioether can be made by either alkylation of a thiol/mercaptan (-SH) group on one component such as the component with an alkylating group such as halide or epoxide on a second component such as the nanocarrier. Thioether linkers can also be formed by Michael addition of a thiol/mercaptan group on one component to an electron-deficient alkene group on a second component such as a polymer containing a maleimide group or vinyl sulfone group as the Michael acceptor. In another way, thioether linkers can be prepared by the radical thiol-ene reaction of a thiol/mercaptan group on one component with an alkene group on a second component such as a polymer or nanocarrier.

A hydrazone linker is made by the reaction of a hydrazide group on one component with an aldehyde/ketone group on the second component such as the nanocarrier.

A hydrazide linker is formed by the reaction of a hydrazine group on one component with a carboxylic acid group on the second component such as the nanocarrier. Such reaction is generally performed using chemistry similar to the formation of amide bond where the carboxylic acid is activated with an activating reagent.

An imine or oxime linker is formed by the reaction of an amine or N-alkoxyamine (or aminooxy) group on one component with an aldehyde or ketone group on the second component such as the nanocarrier.

An urea or thiourea linker is prepared by the reaction of an amine group on one component with an isocyanate or thioisocyanate group on the second component such as the nanocarrier.

An amidine linker is prepared by the reaction of an amine group on one component with an imidoester group on the second component such as the nanocarrier.

An amine linker is made by the alkylation reaction of an amine group on one component with an alkylating group such as halide, epoxide, or sulfonate ester group on the second component such as the nanocarrier. Alternatively, an amine linker can also be made by reductive amination of an amine group on one component with an aldehyde or ketone group on the second component such as the nanocarrier with a suitable reducing reagent such as sodium cyanoborohydride or sodium triacetoxyborohydride.

A sulfonamide linker is made by the reaction of an amine group on one component with a sulfonyl halide (such as sulfonyl chloride) group on the second component such as the nanocarrier.

A sulfone linker is made by Michael addition of a nucleophile to a vinyl sulfone. Either the vinyl sulfone or the nucleophile may be on the surface of the nanocarrier or attached to a component.

The component can also be conjugated to the nanocarrier via non-covalent conjugation methods. For example, a negative charged antigen or immunosuppressant can be conjugated to a positive charged nanocarrier through electrostatic adsorption. A component containing a metal ligand can also be conjugated to a nanocarrier containing a metal complex via a metal-ligand complex.

In embodiments, the component can be attached to a polymer, for example polylactic acid-block-polyethylene glycol, prior to the assembly of the synthetic nanocarrier or the synthetic nanocarrier can be formed with reactive or activatible groups on its surface. In the latter case, the component may be prepared with a group which is compatible with the attachment chemistry that is presented by the synthetic nanocarriers' surface. In other embodiments, a peptide component can be attached to VLPs or liposomes using a suitable linker. A linker is a compound or reagent that capable of coupling two molecules together. In an embodiment, the linker can be a homobifuntional or heterobifunctional reagent as described in Hermanson 2008. For example, an VLP or liposome synthetic nanocarrier containing a carboxylic group on the surface can be treated with a homobifunctional linker, adipic dihydrazide (ADH), in the presence of EDC to form the corresponding synthetic nanocarrier with the ADH linker. The resulting ADH linked synthetic nanocarrier is then conjugated with a peptide component containing an acid group via the other end of the ADH linker on NC to produce the corresponding VLP or liposome peptide conjugate.

For detailed descriptions of available conjugation methods, see Hermanson G T "Bioconjugate Techniques", 2nd Edition Published by Academic Press, Inc., 2008. In addition to covalent attachment the component can be coupled by adsorption to a pre-formed synthetic nanocarrier or it can be coupled by encapsulation during the formation of the synthetic nanocarrier.

Any immunosuppressant as provided herein can be coupled to the synthetic nanocarrier. Immunosuppressants include, but are not limited to, statins; mTOR inhibitors, such as rapamycin or a rapamycin analog; TGF-β signaling agents; TGF-β receptor agonists; histone deacetylase (HDAC) inhibitors; corticosteroids; inhibitors of mitochondrial function, such as rotenone; P38 inhibitors; NF-κβ inhibitors; adenosine receptor agonists;

prostaglandin E2 agonists; phosphodiesterase inhibitors, such as phosphodiesterase 4 inhibitor; proteasome inhibitors; kinase inhibitors; G-protein coupled receptor agonists; G-protein coupled receptor antagonists; glucocorticoids; retinoids; cytokine inhibitors; cytokine receptor inhibitors; cytokine receptor activators; peroxisome proliferator-activated receptor antagonists; peroxisome proliferator-activated receptor agonists; histone deacetylase inhibitors; calcineurin inhibitors; phosphatase inhibitors and oxidized ATPs. Immunosuppressants also include IDO, vitamin D3, cyclosporine A, aryl hydrocarbon receptor inhibitors, resveratrol, azathiopurine, 6-mercaptopurine, aspirin, niflumic acid, estriol, tripolide, interleukins (e.g., IL-1, IL-10), cyclosporine A, siRNAs targeting cytokines or cytokine receptors and the like.

Examples of statins include atorvastatin (LIPITOR®, TORVAST®), cerivastatin, fluvastatin (LESCOL®, LESCOL® XL), lovastatin (MEVACOR®, ALTOCOR®, ALTOPREV®), mevastatin (COMPACTIN®), pitavastatin (LIVALO®, PIAVA®), rosuvastatin (PRAVACHOL®, SELEKTINE®, LIPOSTAT®), rosuvastatin (CRESTOR®), and simvastatin (ZOCOR®, LIPEX®).

Examples of mTOR inhibitors include rapamycin and analogs thereof (e.g., CCL-779, RAD001, AP23573, C20-methallylrapamycin (C20-Marap), C16-(S)-butylsulfonamidorapamycin (C16-BSrap), C16-(S)-3-methylindolerapamycin (C16-iRap) (Bayle et al. Chemistry & Biology 2006, 13:99-107)), AZD8055, BEZ235 (NVP-BEZ235), chrysophanic acid (chrysophanol), deforolimus (MK-8669), everolimus (RAD0001), KU-0063794, PI-103, PP242, temsirolimus, and WYE-354 (available from Selleck, Houston, TX, USA).

Examples of TGF-β signaling agents include TGF-β ligands (e.g., activin A, GDF1, GDF11, bone morphogenic proteins, nodal, TGF-βs) and their receptors (e.g., ACVR1B, ACVR1C, ACVR2A, ACVR2B, BMPR2, BMPR1A, BMPR1B, TGFβRI, TGFβRII), R-SMADS/co-SMADS (e.g., SMAD1, SMAD2, SMAD3, SMAD4, SMAD5, SMAD8), and ligand inhibitors (e.g, follistatin, noggin, chordin, DAN, lefty, LTBP1, THBS1, Decorin).

Examples of inhibitors of mitochondrial function include atractyloside (dipotassium salt), bongkrekic acid (triammonium salt), carbonyl cyanide m-chlorophenylhydrazone, carboxyatractyloside (e.g., from *Atractylis gummifera*), CGP-37157, (-)-Deguelin (e.g., from *Mundulea sericea*), F16, hexokinase II VDAC binding domain peptide, oligomycin,

rotenone, Ru360, SFK1, and valinomycin (e.g., from *Streptomyces fulvissimus*) (EMD4Biosciences, USA).

Examples of P38 inhibitors include SB-203580 (4-(4-Fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)1H-imidazole), SB-239063 (trans-1-(4hydroxycyclohexyl)-4-(fluorophenyl)-5-(2-methoxy-pyrimidin-4-yl) imidazole), SB-220025 (5-(2amino-4-pyrimidinyl)-4-(4-fluorophenyl)-1-(4-piperidinyl)imidazole)), and ARRY-797.

Examples of NF (e.g., NK-κβ) inhibitors include IFRD1, 2-(1,8-naphthyridin-2-yl)-Phenol, 5-aminosalicylic acid, BAY 11-7082, BAY 11-7085, CAPE (Caffeic Acid Phenethylester), diethylmaleate, IKK-2 Inhibitor IV, IMD 0354, lactacystin, MG-132 [Z-Leu-Leu-Leu-CHO], NFκB Activation Inhibitor III, NF-κB Activation Inhibitor II, JSH-23, parthenolide, Phenylarsine Oxide (PAO), PPM-18, pyrrolidinedithiocarbamic acid ammonium salt, QNZ, RO 106-9920, rocaglamide, rocaglamide AL, rocaglamide C, rocaglamide I, rocaglamide J, rocaglaol, (R)-MG-132, sodium salicylate, triptolide (PG490), wedelolactone.

Examples of adenosine receptor agonists include CGS-21680 and ATL-146e.

Examples of prostaglandin E2 agonists include E-Prostanoid 2 and E-Prostanoid 4.

Examples of phosphodiesterase inhibitors (non-selective and selective inhibitors) include caffeine, aminophylline, IBMX (3-isobutyl-1-methylxanthine), paraxanthine, pentoxifylline, theobromine, theophylline, methylated xanthines, vinpocetine, EHNA (erythro-9-(2-hydroxy-3-nonyl)adenine), anagrelide, enoximone (PERFAN™), milrinone, levosimendon, mesembrine, ibudilast, piclamilast, luteolin, drotaverine, roflumilast (DAXAS™, DALIRESP™), sildenafil (REVATION®, VIAGRA®), tadalafil (ADCIRCA®, CIALIS®), vardenafil (LEVITRA®, STAXYN®), udenafil, avanafil, icariin, 4-methylpiperazine, and pyrazolo pyrimidin-7-1.

Examples of proteasome inhibitors include bortezomib, disulfiram, epigallocatechin-3-gallate, and salinosporamide A.

Examples of kinase inhibitors include bevacizumab, BIBW 2992, cetuximab (ERBITUX®), imatinib (GLEEVEC®), trastuzumab (HERCEPTIN®), gefitinib (IRESSA®), ranibizumab (LUCENTIS®), pegaptanib, sorafenib, dasatinib, sunitinib, erlotinib, nilotinib, lapatinib, panitumumab, vandetanib, E7080, pazopanib, mubritinib.

Examples of glucocorticoids include hydrocortisone (cortisol), cortisone acetate, prednisone, prednisolone, methylprednisolone, dexamethasone, betamethasone,

triamcinolone, beclometasone, fludrocortisone acetate, deoxycorticosterone acetate (DOCA), and aldosterone.

Examples of retinoids include retinol, retinal, tretinoin (retinoic acid, RETIN-A®), isotretinoin (ACCUTANE®, AMNESTEEM®, CLARAVIS®, SOTRET®), alitretinoin (PANRETIN®), etretinate (TEGISON™) and its metabolite acitretin (SORIATANE®), tazarotene (TAZORAC®, AVAGE®, ZORAC®), bexarotene (TARGRETIN®), and adapalene (DIFFERIN®).

Examples of cytokine inhibitors include IL1ra, IL1 receptor antagonist, IGFBP, TNF-BF, uromodulin, Alpha-2-Macroglobulin, Cyclosporin A, Pentamidine, and Pentoxifylline (PENTOPAK®, PENTOXIL®, TRENTAL®).

Examples of peroxisome proliferator-activated receptor antagonists include GW9662, PPARγ antagonist III, G335, T0070907 (EMD4Biosciences, USA).

Examples of peroxisome proliferator-activated receptor agonists include pioglitazone, ciglitazone, clofibrate, GW1929, GW7647, L-165,041, LY 171883, PPARγ activator, Fmoc-Leu, troglitazone, and WY-14643 (EMD4Biosciences, USA).

Examples of histone deacetylase inhibitors include hydroxamic acids (or hydroxamates) such as trichostatin A, cyclic tetrapeptides (such as trapoxin B) and depsipeptides, benzamides, electrophilic ketones, aliphatic acid compounds such as phenylbutyrate and valproic acid, hydroxamic acids such as vorinostat (SAHA), belinostat (PXD101), LAQ824, and panobinostat (LBH589), benzamides such as entinostat (MS-275), CI994, and mocetinostat (MGCD0103), nicotinamide, derivatives of NAD, dihydrocoumarin, naphthopyranone, and 2-hydroxynaphaldehydes.

Examples of calcineurin inhibitors include cyclosporine, pimecrolimus, voclosporin, and tacrolimus.

Examples of phosphatase inhibitors include BN82002 hydrochloride, CP-91149, calyculin A, cantharidic acid, cantharidin, cypermethrin, ethyl-3,4-dephostatin, fostriecin sodium salt, MAZ51, methyl-3,4-dephostatin, NSC 95397, norcantharidin, okadaic acid ammonium salt from prorocentrum concavum, okadaic acid, okadaic acid potassium salt, okadaic acid sodium salt, phenylarsine oxide, various phosphatase inhibitor cocktails, protein phosphatase 1C, protein phosphatase 2A inhibitor protein, protein phosphatase 2A1, protein phosphatase 2A2, sodium orthovanadate.

In some embodiments, antigens as described herein are also coupled to synthetic nanocarriers. In some embodiments, the antigens are coupled to the same or different

synthetic nanocarriers as to which the immunosuppressants are coupled. In other embodiments, the antigens are not coupled to any synthetic nanocarriers. Antigens include any of the antigens provided herein, or fragments or derivatives thereof, such antigens are associated with inflammatory, autoimmune diseases, allergy, organ or tissue rejection, graft versus host disease, transplant antigens and therapeutic protein antigens. The epitopes, or proteins, polypeptides or peptides that comprise the epitopes, can be obtained or derived from any of the antigens provided or otherwise known in the art.

Therapeutic proteins include, but are not limited to, infusible therapeutic proteins, enzymes, enzyme cofactors, hormones, blood clotting factors, cytokines and interferons, growth factors, monoclonal antibodies, and polyclonal antibodies (e.g., that are administered to a subject as a replacement therapy), and proteins associated with Pompe's disease (e.g., alglucosidase alfa, rhGAA (e.g., Myozyme and Lumizyme (Genzyme)). Therapeutic proteins also include proteins involved in the blood coagulation cascade. Therapeutic proteins include, but are not limited to, Factor VIII, Factor VII, Factor IX, Factor V, von Willebrand Factor, von Heldebrant Factor, tissue plasminogen activator, insulin, growth hormone, erythropoietin alfa, VEGF, thrombopoietin, lysozyme, antithrombin and the like. Therapeutic proteins also include adipokines, such as leptin and adiponectin. Other examples of therapeutic proteins are as described below and elsewhere herein. Also included are fragments or derivatives of any of the therapeutic proteins provided as the antigen.

Examples of therapeutic proteins used in enzyme replacement therapy of subjects having a lysosomal storage disorder include, but are not limited to, imiglucerase for the treatment of Gaucher's disease (e.g., CEREZYME™), a-galactosidase A (a-gal A) for the treatment of Fabry disease (e.g., agalsidase beta, FABRYZYME™), acid a-glucosidase (GAA) for the treatment of Pompe disease (e.g., alglucosidase alfa, LUMIZYME™, MYOZYME™), arylsulfatase B for the treatment of Mucopolysaccharidoses (e.g., laronidase, ALDURAZYME™, idursulfase, ELAPRASE™, arylsulfatase B, NAGLAZYME™).

Examples of enzymes include oxidoreductases, transferases, hydrolases, lyases, isomerases, and ligases.

Examples of hormones include Melatonin (N-acetyl-5-methoxytryptamine), Serotonin, Thyroxine (or tetraiodothyronine) (a thyroid hormone), Triiodothyronine (a thyroid hormone), Epinephrine (or adrenaline), Norepinephrine (or noradrenaline), Dopamine (or prolactin inhibiting hormone), Antimullerian hormone (or mullerian inhibiting factor or

hormone), Adiponectin, Adrenocorticotropic hormone (or corticotropin), Angiotensinogen and angiotensin, Antidiuretic hormone (or vasopressin, arginine vasopressin), Atrial-natriuretic peptide (or atriopeptin), Calcitonin, Cholecystokinin, Corticotropin-releasing hormone, Erythropoietin, Follicle-stimulating hormone, Gastrin, Ghrelin, Glucagon, Glucagon-like peptide (GLP-1), GIP, Gonadotropin-releasing hormone, Growth hormone-releasing hormone, Human chorionic gonadotropin, Human placental lactogen, Growth hormone, Inhibin, Insulin, Insulin-like growth factor (or somatomedin), Leptin, Luteinizing hormone, Melanocyte stimulating hormone, Orexin, Oxytocin, Parathyroid hormone, Prolactin, Relaxin, Secretin, Somatostatin, Thrombopoietin, Thyroid-stimulating hormone (or thyrotropin), Thyrotropin-releasing hormone, Cortisol, Aldosterone, Testosterone, Dehydroepiandrosterone, Androstenedione, Dihydrotestosterone, Estradiol, Estrone, Estriol, Progesterone, Calcitriol (1,25-dihydroxyvitamin D3), Calcidiol (25-hydroxyvitamin D3), Prostaglandins, Leukotrienes, Prostacyclin, Thromboxane, Prolactin releasing hormone, Lipotropin, Brain natriuretic peptide, Neuropeptide Y, Histamine, Endothelin, Pancreatic polypeptide, Renin, and Enkephalin.

Examples of blood and blood coagulation factors include Factor I (fibrinogen), Factor II (prothrombin), tissue factor, Factor V (proaccelerin, labile factor), Factor VII (stable factor, proconvertin), Factor VIII (antihemophilic globulin), Factor IX (Christmas factor or plasma thromboplastin component), Factor X (Stuart-Prower factor), Factor Xa, Factor XI, Factor XII (Hageman factor), Factor XIII (fibrin-stabilizing factor), von Willebrand factor, prekallikrein (Fletcher factor), high-molecular weight kininogen (HMWK) (Fitzgerald factor), fibronectin, fibrin, thrombin, antithrombin III, heparin cofactor II, protein C, protein S, protein Z, protein Z-related protease inhibitot (ZPI), plasminogen, alpha 2-antiplasmin, tissue plasminogen activator (tPA), urokinase, plasminogen activator inhibitor-1 (PAI1), plasminogen activator inhibitor-2 (PAI2), cancer procoagulant, and epoetin alfa (Epogen, Procrit).

Examples of cytokines include lymphokines, interleukins, and chemokines, type 1 cytokines, such as IFN-γ, TGF-β, and type 2 cytokines, such as IL-4, IL-10, and IL-13.

Examples of growth factors include Adrenomedullin (AM), Angiopoietin (Ang), Autocrine motility factor, Bone morphogenetic proteins (BMPs), Brain-derived neurotrophic factor (BDNF), Epidermal growth factor (EGF), Erythropoietin (EPO), Fibroblast growth factor (FGF), Glial cell line-derived neurotrophic factor (GDNF), Granulocyte colony-stimulating factor (G-CSF), Granulocyte macrophage colony-stimulating factor (GM-CSF),

Growth differentiation factor-9 (GDF9), Hepatocyte growth factor (HGF), Hepatoma-derived growth factor (HDGF), Insulin-like growth factor (IGF), Migration-stimulating factor, Myostatin (GDF-8), Nerve growth factor (NGF) and other neurotrophins, Platelet-derived growth factor (PDGF), Thrombopoietin (TPO), Transforming growth factor alpha(TGF-α), Transforming growth factor beta(TGF-β), Tumour_necrosis_factor-alpha(TNF-α), Vascular endothelial growth factor (VEGF), Wnt Signaling Pathway, placental growth factor (PlGF), [(Foetal Bovine Somatotrophin)] (FBS), IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, and IL-7.

Examples of monoclonal antibodies include Abagovomab, Abciximab, Adalimumab, Adecatumumab, Afelimomab, Afutuzumab, Alacizumab pegol, ALD, Alemtuzumab, Altumomab pentetate, Anatumomab mafenatox, Anrukinzumab, Anti-thymocyte globin, Apolizumab, Arcitumomab, Aselizumab, Atlizumab (tocilizumab), Atorolimumab, Bapineuzumab, Basiliximab, Bavituximab, Bectumomab, Belimumab, Benralizumab, Bertilimumab, Besilesomab, Bevacizumab, Biciromab, Bivatuzumab mertansine, Blinatumomab, Brentuximab vedotin, Briakinumab, Canakinumab, Cantuzumab mertansine, Capromab pendetide, Catumaxomab, Cedelizumab, Certolizumab pegol, Cetuximab, Citatuzumab bogatox, Cixutumumab, Clenoliximab, Clivatuzumab tetraxetan, Conatumumab, Dacetuzumab, Daclizumab, Daratumumab, Denosumab, Detumomab, Dorlimomab aritox, Dorlixizumab, Ecromeximab, Eculizumab, Edobacomab, Edrecolomab, Efalizumab, Efungumab, Elotuzumab, Elsilimomab, Enlimomab pegol, Epitumomab cituxetan, Epratuzumab, Erlizumab, Ertumaxomab, Etaracizumab, Exbivirumab, Fanolesomab, Faralimomab, Farletuzumab, Felvizumab, Fezakinumab, Figitumumab, Fontolizumab , Foravirumab, Fresolimumab, Galiximab, Gantenerumab, Gavilimomab, Gemtuzumab ozogamicin, GC1008, Girentuximab, Glembatumumab vedotin, Golimumab, Gomiliximab, Ibalizumab, Ibritumomab tiuxetan, Igovomab, Imciromab, Infliximab, Intetumumab, Inolimomab, Inotuzumab ozogamicin, Ipilimumab, Iratumumab, Keliximab, Labetuzumab, Lebrikizumab, Lemalesomab, Lerdelimumab, Lexatumumab, Libivirumab, Lintuzumab, Lorvotuzumab mertansine, Lucatumumab, Lumiliximab, Mapatumumab, Maslimomab, Matuzumab, Mepolizumab, Metelimumab, Milatuzumab, Minretumomab, Mitumomab, Morolimumab, Motavizumab, Muromonab-CD3, Nacolomab tafenatox, Naptumomab estafenatox, Natalizumab, Nebacumab, Necitumumab, Nerelimomab, Nimotuzumab, Nofetumomab merpentan, Ocrelizumab, Odulimomab, Ofatumumab, Olaratumab, Omalizumab, Oportuzumab monatox, Oregovomab, Otelixizumab, Pagibaximab, Palivizumab, Panitumumab, Panobacumab, Pascolizumab, Pemtumomab,

Pertuzumab, Pexelizumab, Pintumomab, Priliximab, Pritumumab, Rafivirumab, Ramucirumab, Ranibizumab, Raxibacumab, Regavirumab Reslizumab, Rilotumumab, Rituximab, Robatumumab, Rontalizumab, Rovelizumab, Ruplizumab, Satumomab pendetide, Sevirumab, Sibrotuzumab, Sifalimumab, Siltuximab, Siplizumab, Solanezumab, Sonepcizumab, Sontuzumab, Stamulumab, Sulesomab, Tacatuzumab tetraxetan, Tadocizumab, Talizumab, Tanezumab, Taplitumomab paptox, Tefibazumab, Telimomab aritox, Tenatumomab, Teneliximab, Teplizumab, Ticilimumab (tremelimumab), Tigatuzumab, Tocilizumab (atlizumab), Toralizumab, Tositumomab, Trastuzumab, Tremelimumab, Tucotuzumab celmoleukin, Tuvirumab, Urtoxazumab, Ustekinumab, Vapaliximab, Vedolizumab, Veltuzumab, Vepalimomab, Visilizumab, Volociximab, Votumumab, Zalutumumab, Zanolimumab, Ziralimumab, and Zolimomab aritox.

Examples of infusion therapy or injectable therapeutic proteins include, for example, Tocilizumab (Roche/Actemra®), alpha-1 antitrypsin (Kamada/AAT), Hematide® (Affymax and Takeda, synthetic peptide), albinterferon alfa-2b (Novartis/Zalbin™), Rhucin® (Pharming Group, C1 inhibitor replacement therapy), tesamorelin (Theratechnologies/Egrifta, synthetic growth hormone-releasing factor), ocrelizumab (Genentech, Roche and Biogen), belimumab (GlaxoSmithKline/Benlysta®), pegloticase (Savient Pharmaceuticals/Krystexxa™), taliglucerase alfa (Protalix/Uplyso), agalsidase alfa (Shire/Replagal®), velaglucerase alfa (Shire).

Additional therapeutic proteins useful in accordance to aspects of this invention will be apparent to those of skill in the art, and the invention is not limited in this respect.

In some embodiments, a component, such as an antigen or immunosuppressant, may be isolated. Isolated refers to the element being separated from its native environment and present in sufficient quantities to permit its identification or use. This means, for example, the element may be (i) selectively produced by expression cloning or (ii) purified as by chromatography or electrophoresis. Isolated elements may be, but need not be, substantially pure. Because an isolated element may be admixed with a pharmaceutically acceptable excipient in a pharmaceutical preparation, the element may comprise only a small percentage by weight of the preparation. The element is nonetheless isolated in that it has been separated from the substances with which it may be associated in living systems, i.e., isolated from other lipids or proteins. Any of the elements provided herein can be included in the compositions in isolated form.

D. METHODS OF MAKING AND USING THE INVENTIVE COMPOSITIONS AND RELATED METHODS

Synthetic nanocarriers may be prepared using a wide variety of methods known in the art. For example, synthetic nanocarriers can be formed by methods as nanoprecipitation, flow focusing using fluidic channels, spray drying, single and double emulsion solvent evaporation, solvent extraction, phase separation, milling, microemulsion procedures, microfabrication, nanofabrication, sacrificial layers, simple and complex coacervation, and other methods well known to those of ordinary skill in the art. Alternatively or additionally, aqueous and organic solvent syntheses for monodisperse semiconductor, conductive, magnetic, organic, and other nanomaterials have been described (Pellegrino et al., 2005, Small, 1:48; Murray et al., 2000, Ann. Rev. Mat. Sci., 30:545; and Trindade et al., 2001, Chem. Mat., 13:3843). Additional methods have been described in the literature (see, e.g., Doubrow, Ed., "Microcapsules and Nanoparticles in Medicine and Pharmacy," CRC Press, Boca Raton, 1992; Mathiowitz et al., 1987, J. Control. Release, 5:13; Mathiowitz et al., 1987, Reactive Polymers, 6:275; and Mathiowitz et al., 1988, J. Appl. Polymer Sci., 35:755; US Patents 5578325 and 6007845; P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" Nanomedicine. 5(6):843-853 (2010)).

Various materials may be encapsulated into synthetic nanocarriers as desirable using a variety of methods including but not limited to C. Astete et al., "Synthesis and characterization of PLGA nanoparticles" J. Biomater. Sci. Polymer Edn, Vol. 17, No. 3, pp. 247–289 (2006); K. Avgoustakis "Pegylated Poly(Lactide) and Poly(Lactide-Co-Glycolide) Nanoparticles: Preparation, Properties and Possible Applications in Drug Delivery" Current Drug Delivery 1:321-333 (2004); C. Reis et al., "Nanoencapsulation I. Methods for preparation of drug-loaded polymeric nanoparticles" Nanomedicine 2:8– 21 (2006); P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" Nanomedicine. 5(6):843-853 (2010). Other methods suitable for encapsulating materials into synthetic nanocarriers may be used, including without limitation methods disclosed in United States Patent 6,632,671 to Unger October 14, 2003.

In certain embodiments, synthetic nanocarriers are prepared by a nanoprecipitation process or spray drying. Conditions used in preparing synthetic nanocarriers may be altered to yield particles of a desired size or property (e.g., hydrophobicity, hydrophilicity, external

morphology, "stickiness," shape, etc.). The method of preparing the synthetic nanocarriers and the conditions (e.g., solvent, temperature, concentration, air flow rate, etc.) used may depend on the materials to be coupled to the synthetic nanocarriers and/or the composition of the polymer matrix.

If particles prepared by any of the above methods have a size range outside of the desired range, particles can be sized, for example, using a sieve.

Elements (i.e., components) of the inventive synthetic nanocarriers (such as moieties of which an immunofeature surface is comprised, targeting moieties, polymeric matrices, antigens, immunosuppressants and the like) may be coupled to the overall synthetic nanocarrier, e.g., by one or more covalent bonds, or may be coupled by means of one or more linkers. Additional methods of functionalizing synthetic nanocarriers may be adapted from Published US Patent Application 2006/0002852 to Saltzman et al., Published US Patent Application 2009/0028910 to DeSimone et al., or Published International Patent Application WO/2008/127532 A1 to Murthy et al.

Alternatively or additionally, synthetic nanocarriers can be coupled to components directly or indirectly via non-covalent interactions. In non-covalent embodiments, the non-covalent coupling is mediated by non-covalent interactions including but not limited to charge interactions, affinity interactions, metal coordination, physical adsorption, host-guest interactions, hydrophobic interactions, TT stacking interactions, hydrogen bonding interactions, van der Waals interactions, magnetic interactions, electrostatic interactions, dipole-dipole interactions, and/or combinations thereof. Such couplings may be arranged to be on an external surface or an internal surface of an inventive synthetic nanocarrier. In embodiments, encapsulation and/or absorption is a form of coupling. In embodiments, the inventive synthetic nanocarriers can be combined with an antigen by admixing in the same vehicle or delivery system.

Populations of synthetic nanocarriers may be combined to form pharmaceutical dosage forms according to the present invention using traditional pharmaceutical mixing methods. These include liquid-liquid mixing in which two or more suspensions, each containing one or more subsets of nanocarriers, are directly combined or are brought together via one or more vessels containing diluent. As synthetic nanocarriers may also be produced or stored in a powder form, dry powder-powder mixing could be performed as could the re-suspension of two or more powders in a common media. Depending on the properties of the

nanocarriers and their interaction potentials, there may be advantages conferred to one or another route of mixing.

Typical inventive compositions that comprise synthetic nanocarriers may comprise inorganic or organic buffers (e.g., sodium or potassium salts of phosphate, carbonate, acetate, or citrate) and pH adjustment agents (e.g., hydrochloric acid, sodium or potassium hydroxide, salts of citrate or acetate, amino acids and their salts) antioxidants (e.g., ascorbic acid, alpha-tocopherol), surfactants (e.g., polysorbate 20, polysorbate 80, polyoxyethylene9-10 nonyl phenol, sodium desoxycholate), solution and/or cryo/lyo stabilizers (e.g., sucrose, lactose, mannitol, trehalose), osmotic adjustment agents (e.g., salts or sugars), antibacterial agents (e.g., benzoic acid, phenol, gentamicin), antifoaming agents (e.g., polydimethylsilozone), preservatives (e.g., thimerosal, 2-phenoxyethanol, EDTA), polymeric stabilizers and viscosity-adjustment agents (e.g., polyvinylpyrrolidone, poloxamer 488, carboxymethylcellulose) and co-solvents (e.g., glycerol, polyethylene glycol, ethanol).

Compositions according to the invention comprise inventive synthetic nanocarriers in combination with pharmaceutically acceptable excipients. The compositions may be made using conventional pharmaceutical manufacturing and compounding techniques to arrive at useful dosage forms. Techniques suitable for use in practicing the present invention may be found in Handbook of Industrial Mixing: Science and Practice, Edited by Edward L. Paul, Victor A. Atiemo-Obeng, and Suzanne M. Kresta, 2004 John Wiley & Sons, Inc.; and Pharmaceutics: The Science of Dosage Form Design, 2nd Ed. Edited by M. E. Auten, 2001, Churchill Livingstone. In an embodiment, inventive synthetic nanocarriers are suspended in sterile saline solution for injection together with a preservative.

It is to be understood that the compositions of the invention can be made in any suitable manner, and the invention is in no way limited to compositions that can be produced using the methods described herein. Selection of an appropriate method may require attention to the properties of the particular moieties being associated.

In some embodiments, inventive synthetic nanocarriers are manufactured under sterile conditions or are terminally sterilized. This can ensure that resulting compositions are sterile and non-infectious, thus improving safety when compared to non-sterile compositions. This provides a valuable safety measure, especially when subjects receiving synthetic nanocarriers have immune defects, are suffering from infection, and/or are susceptible to infection. In some embodiments, inventive synthetic nanocarriers may be lyophilized and stored in

suspension or as lyophilized powder depending on the formulation strategy for extended periods without losing activity.

The compositions of the invention can be administered by a variety of routes, including but not limited to subcutaneous, intranasal, oral, intravenous, intraperitoneal, intramuscular, transmucosal, transmucosal, sublingual, rectal, ophthalmic, pulmonary, intradermal, transdermal, transcutaneous or intradermal or by a combination of these routes. Routes of administration also include administration by inhalation or pulmonary aerosol. Techniques for preparing aerosol delivery systems are well known to those of skill in the art (see, for example, Sciarra and Cutie, "Aerosols," in Remington's Pharmaceutical Sciences, 18th edition, 1990, pp. 1694-1712; incorporated by reference).

The transplantable grafts or therapeutic proteins provided as a cell-based therapy of the invention may be administered by parenteral, intraarterial, intranasal or intravenous administration or by injection to lymph nodes or anterior chamber of the eye or by local administration to an organ or tissue of interest. The administration may be by subcutaneous, intrathecal, intraventricular, intramuscular, intraperitoneal, intracoronary, intrapancreatic, intrahepatic or bronchial injection.

The compositions of the invention can be administered in effective amounts, such as the effective amounts described elsewhere herein. Doses of dosage forms contain varying amounts of populations of synthetic nanocarriers and/or varying amounts of immunosuppressants and/or antigens, according to the invention. The amount of synthetic nanocarriers and/or immunosuppressants and/or antigens present in the inventive dosage forms can be varied according to the nature of the antigens, the therapeutic benefit to be accomplished, and other such parameters. In embodiments, dose ranging studies can be conducted to establish optimal therapeutic amount of the population of synthetic nanocarriers and the amount of immunosuppressants and/or antigens to be present in the dosage form. In embodiments, the synthetic nanocarriers and/or the immunosuppressants and/or antigens are present in the dosage form in an amount effective to generate a tolerogenic immune response to the antigens upon administration to a subject. It may be possible to determine amounts of the immunosuppressants and/or antigens effective to generate a tolerogenic immune response using conventional dose ranging studies and techniques in subjects. Inventive dosage forms may be administered at a variety of frequencies. In a preferred embodiment, at least one administration of the dosage form is sufficient to generate a pharmacologically relevant response. In more preferred embodiments, at least two administrations, at least three

administrations, or at least four administrations, of the dosage form are utilized to ensure a pharmacologically relevant response.

Prophylactic administration of the inventive compositions can be initiated prior to the onset of disease, disorder or condition or therapeutic administration can be initiated after a disorder, disorder or condition is established.

In some embodiments, administration of synthetic nanocarriers is undertaken e.g., prior to administration of a therapeutic protein or transplantable graft or exposure to an allergen. In exemplary embodiments, synthetic nanocarriers are administered at one or more times including, but not limited to, 30, 25, 20, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0 days prior to administration of a therapeutic protein or transplantable graft or exposure to an allergen. In addition or alternatively, synthetic nanocarriers can be administered to a subject following administration of a therapeutic protein or transplantable graft or exposure to an allergen. In exemplary embodiments, synthetic nanocarriers are administered at one or more times including, but not limited to, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, etc. days following administration of a therapeutic protein or transplantable graft or exposure to an allergen.

In some embodiments, a maintenance dose (e.g., of a synthetic nanocarrier composition provided herein) is administered to a subject after an initial administration has resulted in a tolerogenic response in the subject, for example to maintain the tolerogenic effect achieved after the initial dose, to prevent an undesired immune reaction in the subject, or to prevent the subject becoming a subject at risk of experiencing an undesired immune response or an undesired level of an immune response. In some embodiments, the maintenance dose is the same dose as the initial dose the subject received. In some embodiments, the maintenance dose is a lower dose than the initial dose. For example, in some embodiments, the maintenance dose is about 3/4, about 2/3, about 1/2, about 1/3, about 1/4, about 1/8, about 1/10, about 1/20, about 1/25, about 1/50, about 1/100, about 1/1,000, about 1/10,000, about 1/100,000, or about 1/1,000,000 (weight/weight) of the initial dose.

The compositions and methods described herein can be used to induce or enhance a tolerogenic immune response and/or to suppress, modulate, direct or redirect an undesired immune response for the purpose of immune suppression. The compositions and methods described herein can be used in the diagnosis, prophylaxis and/or treatment of diseases, disorders or conditions in which immune suppression (e.g., tolerogenic immune response) would confer a treatment benefit. Such diseases, disorders or conditions include

inflammatory diseases, autoimmune diseases, allergies, organ or tissue rejection and graft versus host disease. The compositions and methods described herein can also be used in subjects who have undergone or will undergo transplantation. The compositions and methods described herein can also be used in subjects who have received, are receiving or will receive a therapeutic protein against which they have generated or are expected to generate an undesired immune response.

Autoimmune diseases include, but are not limited to, rheumatoid arthritis, multiple sclerosis, immune-mediated or Type I diabetes mellitus, inflammatory bowel disease (e.g., Crohn's disease or ulcerative colitis), systemic lupus erythematosus, psoriasis, scleroderma, autoimmune thyroid disease, alopecia areata, Grave's disease, Guillain-Barré syndrome, celiac disease, Sjögren's syndrome, rheumatic fever, gastritis, autoimmune atrophic gastritis, autoimmune hepatitis, insulitis, oophoritis, orchitis, uveitis, phacogenic uveitis, myasthenia gravis, primary myxoedema, pernicious anemia, autoimmune haemolytic anemia, Addison's disease, scleroderma, Goodpasture's syndrome, nephritis, for example, glomerulonephritis, psoriasis, pemphigus vulgaris, pemphigoid, sympathetic opthalmia, idiopathic thrombocylopenic purpura, idiopathic feucopenia, Wegener's granulomatosis and poly/dermatomyositis.

Some additional exemplary autoimmune diseases, associated autoantigens, and autoantibodies, which are contemplated for use in the invention, are described in Table 1 below:

Autoantibody Type	Autoantibody	Autoantigen	Autoimmune disease or disorder
Antinuclear antibodies	Anti-SSA/Ro autoantibodies	ribonucleoproteins	Systemic lupus erythematosus, neonatal heart block, primary Sjögren's syndrome
	Anti-La/SS-B autoantibodies	ribonucleoproteins	Primary Sjögren's syndrome
	Anti-centromere antibodies	centromere	CREST syndrome
	Anti-neuronal nuclear antibody-2	Ri[disambiguation needed]	Opsoclonus
	Anti-dsDNA	double-stranded DNA	SLE
	Anti-Jo1	histidine-tRNA ligase	Inflammatory myopathy
	Anti-Smith	snRNP core proteins	SLE

	Anti-topoisomerase antibodies	Type I topoisomerase	Systemic sclerosis (anti-Scl-70 antibodies)
	Anti-histone antibodies	histones	SLE and Drug-induced LE[2]
	Anti-p62 antibodies[3]	nucleoporin 62	Primary biliary cirrhosis[3][4][5]
	Anti-sp100 antibodies [4]	Sp100 nuclear antigen	
	Anti-glycoprotein-210 antibodies[5]	nucleoporin 210kDa	
Anti-transglutaminase antibodies	Anti-tTG		Coeliac disease
	Anti-eTG		Dermatitis herpetiformis
Anti-ganglioside antibodies		ganglioside GQ1B	Miller-Fisher Syndrome
		ganglioside GD3	Acute motor axonal neuropathy (AMAN)
		ganglioside GM1	Multifocal motor neuropathy with conduction block (MMN)
Anti-actin antibodies		actin	Coeliac disease anti-actin antibodies correlated with the level of intestinal damage [6][7]
Liver kidney microsomal type 1 antibody			Autoimmune hepatitis.[8]
Lupus anticoagulant	Anti-thrombin antibodies	thrombin	Systemic lupus erythematosus
Anti-neutrophil cytoplasmic antibody		phospholipid	Antiphospholipid syndrome
	c-ANCA	proteins in neutrophil cytoplasm	Wegener's granulomatosis
	p-ANCA	neutrophil perinuclear	Microscopic polyangiitis, Churg-Strauss syndrome, systemic vasculitides (non-specific)
Rheumatoid factor		IgG	Rheumatoid arthritis
Anti-smooth muscle antibody		smooth muscle	Chronic autoimmune hepatitis
Anti-mitochondrial antibody		mitochondria	Primary biliary cirrhosis[9]

Anti-SRP		signal recognition particle	Polymyositis[10]
		exosome complex	Scleromyositis
		nicotinic acetylcholine receptor	Myasthenia gravis
		muscle-specific kinase (MUSK)	Myasthenia gravis
Anti-VGCC		voltage-gated calcium channel (P/Q-type)	Lambert-Eaton myasthenic syndrome
		thyroid peroxidase (microsomal)	Hashimoto's thyroiditis
		TSH receptor	Graves' disease
		Hu	Paraneoplastic cerebellar syndrome
		Yo (cerebellar Purkinje Cells)	Paraneoplastic cerebellar syndrome
		amphiphysin	Stiff person syndrome, paraneoplastic cerebellar syndrome
Anti-VGKC		voltage-gated potassium channel (VGKC)	Limbic encephalitis, Isaac's Syndrome (autoimmune neuromyotonia)
		basal ganglia neurons	Sydenham's chorea, paediatric autoimmune neuropsychiatric disease associated with Streptococcus (PANDAS)
		N-methyl-D-aspartate receptor (NMDA)	Encephalitis
		glutamic acid decarboxylase (GAD)	Diabetes mellitus type 1, stiff person syndrome
		aquaporin-4	Neuromyelitis optica (Devic's syndrome)

Inflammatory diseases include, but are not limited to, Alzheimer's, Ankylosing spondylitis, arthritis, asthma, atherosclerosis, Behcet's disease, chronic inflammatory demyelinating polyradiculoneuropathy, Crohn's disease, colitis, cystic fibrosis, dermatitis, diverticulitis, hepatitis, irritable bowel syndrome (IBS), lupus erythematous, muscular dystrophy, nephritis, Parkinson's, shingles and ulcerative colitis. Inflammatory diseases also

include, for example, cardiovascular disease, chronic obstructive pulmonary disease (COPD), bronchiectasis, chronic cholecystitis, tuberculosis, Hashimoto's thyroiditis, sepsis, sarcoidosis, silicosis and other pneumoconioses, and an implanted foreign body in a wound, but are not so limited. As used herein, the term "sepsis" refers to a well-recognized clinical syndrome associated with a host's systemic inflammatory response to microbial invasion. The term "sepsis" as used herein refers to a condition that is typically signaled by fever or hypothermia, tachycardia, and tachypnea, and in severe instances can progress to hypotension, organ dysfunction, and even death.

In some embodiments, the inflammatory disease is non-autoimmune inflammatory bowel disease, post-surgical adhesions, coronary artery disease, hepatic fibrosis, acute respiratory distress syndrome, acute inflammatory pancreatitis, endoscopic retrograde cholangiopancreatography-induced pancreatitis, burns, atherogenesis of coronary, cerebral and peripheral arteries, appendicitis, cholecystitis, diverticulitis, visceral fibrotic disorders, wound healing, skin scarring disorders (keloids, hidradenitis suppurativa), granulomatous disorders (sarcoidosis, primary biliary cirrhosis), asthma, pyoderma gandrenosum, Sweet's syndrome, Behcet's disease, primary sclerosing cholangitis or an abscess. In some preferred embodiment the inflammatory disease is inflammatory bowel disease (e.g., Crohn's disease or ulcerative colitis).

In other embodiments, the inflammatory disease is an autoimmune disease. The autoimmune disease in some embodiments is rheumatoid arthritis, rheumatic fever, ulcerative colitis, Crohn's disease, autoimmune inflammatory bowel disease, insulin-dependent diabetes mellitus, diabetes mellitus, juvenile diabetes, spontaneous autoimmune diabetes, gastritis, autoimmune atrophic gastritis, autoimmune hepatitis, thyroiditis, Hashimoto's thyroiditis, insulitis, oophoritis, orchitis, uveitis, phacogenic uveitis, multiple sclerosis, myasthenia gravis, primary myxoedema, thyrotoxicosis, pernicious anemia, autoimmune haemolytic anemia, Addison's disease, Anklosing spondylitis, sarcoidosis, scleroderma, Goodpasture's syndrome, Guillain-Barre syndrome, Graves' disease, glomerulonephritis, psoriasis, pemphigus vulgaris, pemphigoid, excema, bulous pemiphigous, sympathetic opthalmia, idiopathic thrombocylopenic purpura, idiopathic feucopenia, Sjogren's syndrome, systemic sclerosis, Wegener's granulomatosis, poly/dermatomyositis, primary biliary cirrhosis, primary sclerosing cholangitis, lupus or systemic lupus erythematosus.

Graft versus host disease (GVHD) is a complication that can occur after a pluripotent cell (e.g., stem cell) or bone marrow transplant in which the newly transplanted material

results in an attack on the transplant recipient's body. In some instances, GVHD takes place after a blood transfusion. Graft-versus-host-disease can be divided into acute and chronic forms. The acute or fulminant form of the disease (aGVHD) is normally observed within the first 100 days post-transplant, and is a major challenge to transplants owing to associated morbidity and mortality. The chronic form of graft-versus-host-disease (cGVHD) normally occurs after 100 days. The appearance of moderate to severe cases of cGVHD adversely influences long-term survival.

EXAMPLES

Example 1: Immune Response of Synthetic Nanocarriers with Coupled Rapamycin with and without Ovalbumin Peptide (323-339)

Materials

Ovalbumin peptide 323-339, a 17 amino acid peptide known to be a T and B cell epitope of Ovalbumin protein, was purchased from Bachem Americas Inc. (3132 Kashiwa Street, Torrance CA 90505; Part # 4065609). Rapamycin was purchased from TSZ CHEM (185 Wilson Street, Framingham, MA 01702; Product Catalogue # R1017). PLGA with a lactide:glycolide ratio of 3:1 and an inherent viscosity of 0.75 dL/g was purchased from SurModics Pharmaceuticals (756 Tom Martin Drive, Birmingham, AL 35211; Product Code 7525 DLG 7A). Polyvinyl alcohol (85-89% hydrolyzed) was purchased from EMD Chemicals (Product Number 1.41350.1001).

Solution 1: Ovalbumin peptide 323-339 @ 20 mg/mL in dilute hydrochloric acid aqueous solution. The solution was prepared by dissolving ovalbumin peptide in 0.13 M hydrochloric acid solution at room temperature.

Solution 2: Rapamycin @ 50 mg/mL in methylene chloride. The solution was prepared by dissolving rapamycin in pure methylene chloride.

Solution 3: PLGA @ 100 mg/mL in methylene chloride. The solution was prepared by dissolving PLGA in pure methylene chloride.

Solution 4: Polyvinyl alcohol @ 50 mg/mL in 100 mM pH 8 phosphate buffer.

Method for Preparing Synthetic Nanocarrier Containing Rapamycin and Ovalbumin (323-339)

A primary water-in-oil emulsion was prepared first. W1/O1 was prepared by combining solution 1 (0.2 mL), solution 2 (0.2 mL), and solution 3 (1.0 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) was then prepared by combining solution 4 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250.

The W1/O1/W2 emulsion was added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the synthetic nanocarriers to form. A portion of the synthetic nanocarriers were washed by transferring the synthetic nanocarrier suspension to a centrifuge tube and centrifuging at 21,000×g and 4 °C for one hour, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure was repeated, and the pellet was re-suspended in phosphate buffered saline for a final synthetic nanocarrier dispersion of about 10 mg/mL.

The amounts of peptide and rapamycin in the synthetic nanocarriers were determined by HPLC analysis. The total dry-synthetic nanocarrier mass per mL of suspension was determined by a gravimetric method.

Method for Synthetic Nanocarrier Containing Rapamycin

A primary water-in-oil emulsion was prepared first. W1/O1 was prepared by combining 0.13 M hydrochloric acid solution (0.2 mL), solution 2 (0.2 mL), and solution 3 (1.0 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) was then prepared by combining solution 4 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250.

The W1/O1/W2 emulsion was added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the synthetic nanocarriers to form. A portion of the synthetic nanocarriers were washed by transferring the synthetic nanocarrier suspension to a centrifuge tube and centrifuging at 21,000×g and 4 °C for one hour, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure was repeated, and the pellet was re-suspended in phosphate buffered saline for a final synthetic nanocarrier dispersion of about 10 mg/mL.

The amount of rapamycin in the synthetic nanocarrier was determined by HPLC analysis. The total dry-synthetic nanocarrier mass per mL of suspension was determined by a gravimetric method.

Method for Measuring Rapamycin Load

Approximately 3 mg of synthetic nanocarriers were collected and centrifuged to separate supernatant from synthetic nanocarrier pellet. Acetonitrile was added to the pellet, and the sample was sonicated and centrifuged to remove any insoluble material. The supernatant and pellet were injected on RP-HPLC and absorbance was read at 278nm. The µg found in the pellet were used to calculate % entrapped (load), µg in supernatant and pellet were used to calculate total µg recovered.

Method for Measuring Ovalbumin (323-339) Load

Approximately 3 mg of synthetic nanocarriers were collected and centrifuged to separate supernatant from synthetic nanocarrier pellet. Trifluoroethanol was added to the pellet and the sample was sonicated to dissolve the polymer, 0.2% trifluoroacetic acid was added and sample was sonicated and then centrifuged to remove any insoluble material. The supernatant and pellet were injected on RP-HPLC and absorbance was read at 215nm. The µg found in the pellet were used to calculate % entrapped (load), µg in supernatant and pellet were used to calculate total µg recovered.

Antigen-specific Tolerogenic Dendritic Cells (Tdc) Activity on Treg Cell Development

The assay included the use of OTII mice which have a transgenic T-cell receptor specific for an immune-dominant ovalbumin (323-339). In order to create antigen-specific tDCs, CD11c+ splenocytes were isolated, and the ovalbumin (323-339) peptide added in vitro at 1µg/ml or no antigen. Soluble or nanocarrier-encapsulated rapamycin was then added to the DCs for 2 hours which were then washed extensively to remove free rapamycin from the culture. Purified responder CD4+CD25- cells were isolated from OTII mice and added to tDC at a 10:1 T to DC ratio. The mixture of tDC and OTII T-cells were then cultured for 4-5 days, and the frequency of Treg cells (CD4+CD25highFoxP3+) were analyzed by flow cytometry as shown in **Fig. 1**. Regions were selected based on isotype controls.

Example 2: Mesoporous Silica Nanoparticles with Coupled Ibuprofen (Prophetic)

Mesoporous SiO2 nanoparticle cores are created through a sol-gel process. Hexadecyltrimethyl-ammonium bromide (CTAB) (0.5 g) is dissolved in deionized water (500 mL), and then 2 M aqueous NaOH solution (3.5 mL) is added to the CTAB solution. The solution is stirred for 30 min, and then Tetraethoxysilane (TEOS) (2.5 mL) is added to the solution. The resulting gel is stirred for 3 h at a temperature of 80 °C. The white precipitate which forms is captured by filtration, followed by washing with deionized water and drying at room temperature. The remaining surfactant is then extracted from the particles by suspension in an ethanolic solution of HCl overnight. The particles are washed with ethanol, centrifuged, and redispersed under ultrasonication. This wash procedure is repeated two additional times.

The SiO2 nanoparticles are then functionalized with amino groups using (3-aminopropyl)-triethoxysilane (APTMS). To do this, the particles are suspended in ethanol (30 mL), and APTMS (50 μL) is added to the suspension. The suspension is allowed to stand at room temperature for 2 h and then is boiled for 4 h, keeping the volume constant by periodically adding ethanol. Remaining reactants are removed by five cycles of washing by centrifugation and redispersing in pure ethanol.

In a separate reaction, 1-4 nm diameter gold seeds are created. All water used in this reaction is first deionized and then distilled from glass. Water (45.5 mL) is added to a 100 mL round-bottom flask. While stirring, 0.2 M aqueous NaOH (1.5 mL) is added, followed by a 1% aqueous solution of tetrakis(hydroxymethyl)phosphonium chloride (THPC) (1.0 mL). Two minutes after the addition of THPC solution, a 10 mg/mL aqueous solution of chloroauric acid (2 mL), which has been aged at least 15 min, is added. The gold seeds are purified through dialysis against water.

To form the core-shell nanocarriers, the amino-functionalized SiO2 nanoparticles formed above are first mixed with the gold seeds for 2 h at room temperature. The gold-decorated SiO2 particles are collected through centrifugation and mixed with an aqueous solution of chloroauric acid and potassium bicarbonate to form the gold shell. The particles are then washed by centrifugation and redispersed in water. Ibuprofen is loaded by suspending the particles in a solution of sodium ibuprofen (1 mg/L) for 72 h. Free ibuprofen is then washed from the particles by centrifugation and redispersing in water.

Example 3: Liposomes Containing Cyclosporine A (Prophetic)

The liposomes are formed using thin film hydration. 1,2-Dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) (32 μmol), cholesterol (32 μmol), and cyclosporin A (6.4 μmol) are dissolved in pure chloroform (3 mL). This lipid solution is added to a 50 mL round-bottom flask, and the solvent is evaporated on a rotary evaporator at a temperature of 60 °C. The flask is then flushed with nitrogen gas to remove remaining solvent. Phosphate buffered saline (2 mL) and five glass beads are added to the flask, and the lipid film is hydrated by shaking at 60 °C for 1 h to form a suspension. The suspension is transferred to a small pressure tube and sonicated at 60 °C for four cycles of 30s pulses with a 30 s delay between each pulse. The suspension is then left undisturbed at room temperature for 2 h to allow for complete hydration. The liposomes are washed by centrifugation followed by resuspension in fresh phosphate buffered saline.

Example 4: Polymeric Nanocarrier Containing Polymer-Rapamycin Conjugate (Prophetic)

Preparation of PLGA-rapamycin conjugate:

PLGA polymer with acid end group (7525 DLG1A, acid number 0.46 mmol/g, Lakeshore Biomaterials; 5 g, 2.3 mmol, 1.0 eq) is dissolved in 30 mL of dichloromethane (DCM). N,N-Dicyclohexylcarbodimide (1.2 eq, 2.8 mmol, 0.57 g) is added followed by rapamycin (1.0 eq, 2.3 mmol, 2.1 g) and 4-dimethylaminopyridine (DMAP) (2.0 eq, 4.6 mmol, 0.56 g). The mixture is stirred at rt for 2 days. The mixture is then filtered to remove insoluble dicyclohexylurea. The filtrate is concentrated to ca. 10 mL in volume and added to 100 mL of isopropyl alcohol (IPA) to precipitate out the PLGA-rapamycin conjugate. The IPA layer is removed and the polymer is then washed with 50 mL of IPA and 50 mL of methyl t-butyl ether (MTBE). The polymer is then dried under vacuum at 35 C for 2 days to give PLGA-rapamycin as a white solid (ca. 6.5 g).

Preparation of nanocarrier containing PLGA-rapamycin conjugate and ovalbumin peptide (323-339):

Nanocarrier containing PLGA-rapamycin is prepared according to the procedure described in Example 1 as follows:

Solutions for nanocarrier formation are prepared as follows:

Solution 1: Ovalbumin peptide 323-339 @ 20 mg/mL in dilute hydrochloric acid aqueous solution. The solution is prepared by dissolving ovalbumin peptide in 0.13 M hydrochloric acid solution at room temperature. Solution 2: PLGA-rapamycin @ 100 mg/mL

in methylene chloride. The solution is prepared by dissolving PLGA-rapamycin in pure methylene chloride. Solution 3: PLA-PEG @ 100 mg/mL in methylene chloride. The solution is prepared by dissolving PLA-PEG in pure methylene chloride. Solution 4: Polyvinyl alcohol @ 50 mg/mL in 100 mM pH 8 phosphate buffer.

A primary water-in-oil emulsion is prepared first. W1/O1 is prepared by combining solution 1 (0.2 mL), solution 2 (0.75 mL), and solution 3 (0.25 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) is then prepared by combining solution 4 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250. The W1/O1/W2 emulsion is added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the nanocarriers to form. A portion of the nanocarriers is washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,600×g and 4 °C for 35 min, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure is repeated, and the pellet is re-suspended in phosphate buffered saline for a final nanocarrier dispersion of about 10 mg/mL.

Example 5: Preparation of Gold Nanocarriers (AuNCs) Containing Rapamycin (Prophetic)

Preparation of HS-PEG-rapamycin:

A solution of PEG acid disulfide (1.0 eq), rapamycin (2.0-2.5 eq), DCC (2.5 eq) and DMAP (3.0 eq) in dry DMF is stirred at rt overnight. The insoluble dicyclohexylurea is removed by filtration and the filtrate is added to isopropyl alcohol (IPA) to precipitate out the PEG-disulfide-di-rapamycin ester and washed with IPA and dried. The polymer is then treated with tris(2-carboxyethyl)phosphine hydrochloride in DMF to reduce the PEG disulfide to thiol PEG rapamycin ester (HS-PEG-rapamycin). The resulting polymer is recovered by precipitation from IPA and dried as previously described and analyzed by H NMR and GPC.

Formation of Gold NCs (AuNCs):

An aq. solution of 500 mL of 1 mM HAuCl4 is heated to reflux for 10 min with vigorous stirring in a 1 L round-bottom flask equipped with a condenser. A solution of 50 mL of 40 mM of trisodium citrate is then rapidly added to the stirring solution. The resulting

deep wine red solution is kept at reflux for 25-30 min and the heat is withdrawn and the solution is cooled to room temperature. The solution is then filtered through a 0.8 μm membrane filter to give the AuNCs solution. The AuNCs are characterized using visible spectroscopy and transmission electron microscopy. The AuNCs are ca. 20 nm diameter capped by citrate with peak absorption at 520 nm.

AuNCs conjugate with HS-PEG-rapamycin:

A solution of 150 μl of HS-PEG-rapamycin (10 μM in 10 mM pH 9.0 carbonate buffer) is added to 1 mL of 20 nm diameter citrate-capped gold nanocarriers (1.16 nM) to produce a molar ratio of thiol to gold of 2500:1. The mixture is stirred at room temperature under argon for 1 hour to allow complete exchange of thiol with citrate on the gold nanocarriers. The AuNCs with PEG-rapamycin on the surface is then purified by centrifuge at 12,000g for 30 minutes. The supernatant is decanted and the pellet containing AuNC-S-PEG-rapamycin is then pellet washed with 1x PBS buffer. The purified Gold-PEG-rapamycin nanocarriers are then resuspend in suitable buffer for further analysis and bioassays.

Example 6: Mesoporous Silica-gold Core-shell Nanocarriers Containing Ovalbumin (Prophetic)

Mesoporous SiO2 nanoparticle cores are created through a sol-gel process. Hexadecyltrimethyl-ammonium bromide (CTAB) (0.5 g) is dissolved in deionized water (500 mL), and then 2 M aqueous NaOH solution (3.5 mL) is added to the CTAB solution. The solution is stirred for 30 min, and then Tetraethoxysilane (TEOS) (2.5 mL) is added to the solution. The resulting gel is stirred for 3 h at a temperature of 80 °C. The white precipitate which forms is captured by filtration, followed by washing with deionized water and drying at room temperature. The remaining surfactant is then extracted from the particles by suspension in an ethanolic solution of HCl overnight. The particles are washed with ethanol, centrifuged, and redispersed under ultrasonication. This wash procedure is repeated two additional times.

The SiO2 nanoparticles are then functionalized with amino groups using (3-aminopropyl)-triethoxysilane (APTMS). To do this, the particles are suspended in ethanol (30 mL), and APTMS (50 μL) is added to the suspension. The suspension is allowed to stand at room temperature for 2 h and then is boiled for 4 h, keeping the volume constant by periodically adding ethanol. Remaining reactants are removed by five cycles of washing by centrifugation and redispersing in pure ethanol.

In a separate reaction, 1-4 nm diameter gold seeds are created. All water used in this reaction is first deionized and then distilled from glass. Water (45.5 mL) is added to a 100 mL round-bottom flask. While stirring, 0.2 M aqueous NaOH (1.5 mL) is added, followed by a 1% aqueous solution of tetrakis(hydroxymethyl)phosphonium chloride (THPC) (1.0 mL). Two minutes after the addition of THPC solution, a 10 mg/mL aqueous solution of chloroauric acid (2 mL), which has been aged at least 15 min, is added. The gold seeds are purified through dialysis against water.

To form the core-shell nanocarriers, the amino-functionalized SiO2 nanoparticles formed above are first mixed with the gold seeds for 2 h at room temperature. The gold-decorated SiO2 particles are collected through centrifugation and mixed with an aqueous solution of chloroauric acid and potassium bicarbonate to form the gold shell. The particles are then washed by centrifugation and redispersed in water. Thiolated Ovalbumin (made by treating Ovalbumin with 2-iminothiolane hydrochloride) is loaded by suspending the particles in a solution of thiolated Ovalbumin (1 mg/L) for 72 h. The particles is then pellet washed with 1x PBS (pH 7.4) to remove free protein. The resulting silica-gold core-shell nanocarriers containing Ovalbumin are then re-suspended in 1x PBS for further analysis and assays.

Example 7: Liposomes Containing Rapamycin and Ovalbumin (Prophetic)

The liposomes are formed by thin film hydration. 1,2-Dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) (32 μmol), cholesterol (32 μmol), and rapamycin (6.4 μmol) are dissolved in pure chloroform (3 mL). This lipid solution is added to a 10 mL glass tube and the solvent is removed under nitrogen gas stream and desiccated for 6 hr. under vacuum. Multilamellar vesicles are obtained by hydration of the film with 2.0 ml of 25 mM MOPS buffer pH 8.5, containing excess amount of Ovalbumin. The tube is vortexed until the lipid film is peeled of from the tube surface. To break the multilamellar vesicles into monolamellar, ten cycles of freezing (liquid nitrogen) and thawing (30°C water bath) are applied. The sample is then diluted to 1 ml in 25 mM MOPS buffer pH 8.5. Size of the resulting liposome is homogenized by extrusion by passing the sample 10 fold through a 200 nm pore polycarbonate filters. The resulting liposomes are then used for further analysis and bioassays.

Example 8: Polymeric Nanocarriers Composed of Modified Polyamino Acid with Surface Conjugated Ovalbumin (Prophetic)

Step-1. Preparation of Poly(γ-glutamic acid) (γ-PGA) modified with L-phenylalanine ethyl ester (L-PAE): 4.7 unit mmol of γ-PGA (Mn= 300 kD) is dissolved in 0.3 N–NaHCO3 aqueous solution (50 mL). L-PAE (4.7 mmol) and EDC.HCl (4.7 mmol) are added to the solution and stirred for 30 min at 4 C. The solution is then maintained at room temperature with stirring for 24 h. Low-molecular-weight chemicals are removed by dialysis using dialysis membrane with MWCO 50 kD. The resulting γ-PGA-graft-L-PAE is obtained by freeze-drying.

Step-2. Preparation of nanoparticles from γ-PGA-graft-L-PAE polymer: Nanoparticles composed of γ-PGA-graft-L-PAE are prepared by a precipitation and dialysis method. γ-PGA-graft-L-PAE (20 mg) was dissolved in 2 ml of DMSO followed by addition of 2 mL of water to form a translucent solution. The solution is then dialyzed against distilled water using cellulose membrane tubing (50,000 MWCO) to form the nanoparticles and to remove the organic solvents for 72 h at room temperature. The distilled water is exchanged at intervals of 12 h. The resulting nanoparticle solution (10 mg/mL in water) is then used for antigen conjugation.

Step-3. Ovalbumin conjugation to γ-PGA nanoparticles: Surface carboxylic acid groups of the γ-PGA nanoparticles (10 mg/ml) are first activated by EDC and NHS (10 mg/mL each in phosphate buffer, pH 5.8) for 2 h at ambient temperature. After pellet washing to remove excess EDC/NHS, the activated nanoparticles are mixed with 1 mL of Ovalbumin (10 mg/ml) in phosphate-buffered saline (PBS, pH 7.4) and the mixture is incubated at 4-8 C for 24 h. The resulting Ovalbumin conjugated γ-PGA nanoparticles are washed twice with PBS and resuspended at 5 mg/mL in PBS for further analysis and bioassays.

Example 9: Erythropoietin (EPO)-encapsulated γ-PGA Nanoparticles (Prophetic)

To prepare the EPO-encapsulated γ-PGA nanoparticles, 0.25–4 mg of EPO is dissolved in 1 mL of PBS (pH 7.4) and 1 mL of the γ-PGA–graft-L-PAE (10 mg/mL in DMSO) is added to the EPO solution. The resulting solution is centrifuged at 14,000 x g for 15 min and repeatedly rinsed with PBS. The resulting EPO-encapsulated γ-PGA nanoparticles are then resuspended in PBS (5 mg/mL) for further analysis and bioassay.

Example 10: Preparation of Gold Nanocarriers (AuNCs) Containing Ovalbumin (Prophetic)

Step-1. Formation of Gold NCs (AuNCs): An aq. solution of 500 mL of 1 mM HAuCl4 is heated to reflux for 10 min with vigorous stirring in a 1 L round-bottom flask equipped with a condenser. A solution of 50 mL of 40 mM of trisodium citrate is then rapidly added to the stirring solution. The resulting deep wine red solution is kept at reflux for 25-30 min and the heat is withdrawn and the solution is cooled to room temperature. The solution is then filtered through a 0.8 μm membrane filter to give the AuNCs solution. The AuNCs are characterized using visible spectroscopy and transmission electron microscopy. The AuNCs are ca. 20 nm diameter capped by citrate with peak absorption at 520 nm.

Step-2. Conjugation of Ovalbumin to AuNCs: A solution of 150 μl of thiolated Ovalbumin (10 μM in 10 mM pH 9.0 carbonate buffer) is added to 1 mL of 20 nm diameter citrate-capped gold nanocarriers (1.16 nM) to produce a molar ratio of thiol to gold of 2500:1. The mixture is stirred at room temperature under argon for 1 hour to allow complete exchange of thiol with citrate on the gold nanocarriers. The AuNCs with Ovalbumin on the surface is then purified by centrifuge at 12,000g for 30 minutes. The supernatant is decanted and the pellet containing AuNC-Ovalbumin is then pellet washed with 1x PBS buffer. The purified Gold-Ovalbumin nanocarriers are then resuspend in suitable buffer for further analysis and bioassays.

Example 11: Evaluating Tolerogenic Immune Response to Antigen In Vivo (Prophetic)

Balb/c mice are immunized with an antigen in incomplete Freund's adjuvant, and the level of regulatory B cell proliferation is assessed. Subsequently, a composition of the invention is administered in a dose-dependent manner. The same mice are then again exposed to the antigen, and the level of regulatory B cell proliferation is again assessed. Changes in regulatory B cell proliferation are then monitored with an increase upon subsequent challenge with the antigen indicating a tolerogenic immune response.

Example 12: Evaluating In Vitro Induction of Regulatory B Cells (Prophetic)

A cell population comprising regulatory B cells or regulatory B cell precursors is contacted in vitro with a composition provided herein. After a time sufficient for the composition to interact with the regulatory B cells or regulatory B cell precursors in the cell

population, an increase in the number of regulatory B cells is expected. A time sufficient for the increase in the number of regulatory B cells in the population of cells is, in some embodiment, a period of about 1 day, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days, about 1 week, about 2 weeks, about 3 weeks, or about 4 weeks. In some embodiments, the number of regulatory B cells is increased after that time, for example, to at least about 5 times, at least about 10 times, at least about 20 times, at least about 50 times, at least about 100 times, at least about 1.000 times, at least about 10.000 times, at least about 100.000 times, or at least about 1.000.000 times as compared to the original total or relative number of regulatory B cells in the population. In some embodiments, the number of regulatory B cells is increased after that time, for example, to about 1%, about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 40%, about 50%, about 75%, about 90%, or about 95% of the total cells in the population or of the lymphocytes in the cell population. In some embodiments, regulatory B cells are absent from the cell population before the contacting, but are present after the contacting.

In some embodiments, the total and/or relative number of regulatory B cells in the population is determined before the population of cells is contacted with a composition provided herein to establish a baseline number of regulatory B cells in the population. In some embodiments, the population of cells is contacted once with a composition provided herein. In some embodiments, the population of cells is contacted repeatedly with a composition provided herein.

The number and/or presence of regulatory B cells in the population of cells is also determined after the contacting. In some embodiments, the number and/or presence of regulatory B cells is monitored over a period of time, for example, by performing a plurality of subsequent regulatory B cell detection assays. In some embodiments, the number and/or presence of regulatory B cells in the population of cells is determined by taking a sample from the cell population that is representative of the cell population, staining cells contained in that sample with antibodies or staining agents that specifically detect regulatory B cell markers, and detecting cells that express regulatory B cell markers in the sample, for example, by FACS or by immunohistochemistry. In some embodiments, the regulatory B cells are quantified. In some embodiments, the quantity of regulatory B cells determined after the contacting is compared to the quantity of regulatory B cells before the contacting, for example, to the baseline number of regulatory B cells, wherein, if the number of

regulatory B cells in the population of cells is higher after the contacting than the baseline number, then it is determined that a tolerogenic response to the composition has occurred.

Example 13: Evaluating In Vivo Induction of Regulatory B Cells (Prophetic)

A cell population comprising regulatory B cells or regulatory B cell precursors is contacted in vivo with a composition provided herein. In some embodiments, the composition is administered to a subject harboring the cell population comprising regulatory B cells or regulatory B cell precursors. After a time sufficient for the composition to interact with the regulatory B cells or regulatory B cell precursors in the subject, an increase in the number of regulatory B cells is expected. A time sufficient for the increase in the number of regulatory B cells in the population is, in some embodiment, a period of about 1 day, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days, about 1 week, about 2 weeks, about 3 weeks, or about 4 weeks. In some embodiments, the period of time sufficient to effect an increase in the number of regulatory B cells is longer than 4 weeks. In some embodiments, the number of regulatory B cells is increased after that time to at least about 5 times, at least about 10 times, at least about 20 times, at least about 50 times, at least about 100 times, at least about 1.000 times, at least about 10.000 times, at least about 100.000 times, or at least about 1.000.000 times as compared to the original total or relative number of regulatory B cells in the subject. In some embodiments, the number of regulatory B cells is increased after that time, for example, to about 1%, about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 40%, about 50%, about 75%, about 90%, or about 95% of the total cells in the subject, or of a lymphocyte population in the subject. In some embodiments, regulatory B cells are absent from the subject before the administration, but are present after the administration.

In some embodiments, the total and/or relative number of regulatory B cells in the subject is determined before the subject is administered a composition provided herein to establish a baseline number of regulatory B cells in the subject. In some embodiments, the composition provided herein is administered to a subject once. In some embodiments, a composition provided herein is administered to a subject multiple times, for example, until a desired increase or abundance of regulatory B cells is observed in the subject.

The number and/or presence of regulatory B cells in the subject is determined after the administration. In some embodiments, the number and/or presence of regulatory B cells is monitored over a period of time, for example, by performing a plurality of subsequent

regulatory B cell detection assays. In some embodiments, the number and/or presence of regulatory B cells in the subject is determined by taking a sample from the subject, for example, a peripheral blood sample, or a lymph sample, that is representative of a lymphocyte population in the subject, staining cells contained in that sample with antibodies or staining agents that specifically detect regulatory B cell markers, and detecting cells that express regulatory B cell markers in the sample, for example, by FACS or by immunohistochemistry. In some embodiments, the regulatory B cells are quantified. In some embodiments, the quantity of regulatory B cells determined after administration is compared to the quantity of regulatory B cells before administration, for example, to the baseline number of regulatory B cells, wherein, if the number of regulatory B cells in the subject is higher after the administration than the baseline number, then it is determined that a tolerogenic response to the composition has occurred.

Example 14: In Vivo Induction of a Desired Immune Response Against Transplantable Bone Marrow Cells (Prophetic)

A population of at least 10^6 synthetic nanocarriers comprising immunosuppressant and antigens obtained or derived from bone marrow cells are administered subcutaneously to a subject four weeks prior to the subject receiving a bone marrow transplant. After the transplant is received by the subject, the generation of a desired immune response in the subject is assessed once daily during the first week after receiving the transplant, and then weekly for the next three weeks, and then monthly for the next 11 months. As part of the assessment, regulatory B cell counts are taken and compared to regulatory B cell counts taken prior to administering the bone marrow transplant or the synthetic nanocarriers to the subject. During the first year, maintenance doses of the synthetic nanocarriers are administered bi-monthly to the subject. The subject is expected to exhibit higher or desired levels of regulatory B cells specific to the bone marrow transplant.

Example 15: Assessing the Effects of Nanocarriers with Antigens and Immunosuppressants

Nanocarriers

Ovalbumin peptide 323-339, a 17 amino acid peptide known to be a T and B cell epitope of Ovalbumin protein, was purchased from Bachem Americas Inc. (3132 Kashiwa

Street, Torrance CA 90505; Part # 4065609). PLGA with a lactide:glycolide ratio of 3:1 and an inherent viscosity of 0.75 dL/g was purchased from SurModics Pharmaceuticals (756 Tom Martin Drive, Birmingham, AL 35211; Product Code 7525 DLG 7A). PLA-PEG block co-polymer with a PEG block of approximately 5,000 Da and PLA block of approximately 20,000 Da was synthesized. Polyvinyl alcohol (85-89% hydrolyzed) was purchased from EMD Chemicals (Product Number 1.41350.1001).

Solutions were prepared as follows: Solution 1: Ovalbumin peptide 323-339 @ 20 mg/mL in dilute hydrochloric acid aqueous solution. The solution was prepared by dissolving ovalbumin peptide in 0.13 M hydrochloric acid solution at room temperature. Solution 2: PLGA @ 100 mg/mL in methylene chloride. The solution was prepared by dissolving PLGA in pure methylene chloride. Solution 3: PLA-PEG @ 100 mg/mL in methylene chloride. The solution was prepared by dissolving PLA-PEG in pure methylene chloride. Solution 4: Polyvinyl alcohol @ 50 mg/mL in 100 mM pH 8 phosphate buffer. A primary water-in-oil emulsion was prepared first. W1/O1 was prepared by combining solution 1 (0.2 mL), solution 2 (0.75 mL), and solution 3 (0.25 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) was then prepared by combining solution 4 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250.

The W1/O1/W2 emulsion was added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the nanocarriers to form. A portion of the nanocarriers were washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,600×g and 4 °C for 35 min, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure was repeated, and the pellet was re-suspended in phosphate buffered saline for a final nanocarrier dispersion of about 10 mg/mL.

Nanocarrier size was determined by dynamic light scattering. The amount of peptide in the nanocarrier was determined by HPLC analysis. The total dry-nanocarrier mass per mL of suspension was determined by a gravimetric method.

Nanocarrier	Effective Diameter (nm)	Peptide Content (% w/w)
1	234	2.1

Ovalbumin peptide 323-339, a 17 amino acid peptide known to be a T and B cell epitope of Ovalbumin protein, was purchased from Bachem Americas Inc. (3132 Kashiwa Street, Torrance CA 90505; Part # 4065609). Rapamycin was purchased from TSZ CHEM (185 Wilson Street, Framingham, MA 01702; Product Catalogue # R1017). PLGA with a lactide:glycolide ratio of 3:1 and an inherent viscosity of 0.75 dL/g was purchased from SurModics Pharmaceuticals (756 Tom Martin Drive, Birmingham, AL 35211; Product Code 7525 DLG 7A). PLA-PEG block co-polymer with a PEG block of approximately 5,000 Da and PLA block of approximately 20,000 Da was synthesized. Polyvinyl alcohol (85-89% hydrolyzed) was purchased from EMD Chemicals (Product Number 1.41350.1001).

Solutions were prepared as follows: Solution 1: Ovalbumin peptide 323-339 @ 20 mg/mL in dilute hydrochloric acid aqueous solution. The solution was prepared by dissolving ovalbumin peptide in 0.13 M hydrochloric acid solution at room temperature. Solution 2: Rapamycin @ 50 mg/mL in methylene chloride. The solution was prepared by dissolving rapamycin in pure methylene chloride. Solution 3: PLGA @ 100 mg/mL in methylene chloride. The solution was prepared by dissolving PLGA in pure methylene chloride. Solution 4: PLA-PEG @ 100 mg/mL in methylene chloride. The solution was prepared by dissolving PLA-PEG in pure methylene chloride. Solution 5: Polyvinyl alcohol @ 50 mg/mL in 100 mM pH 8 phosphate buffer.

A primary water-in-oil emulsion was prepared first. W1/O1 was prepared by combining solution 1 (0.2 mL), solution 2 (0.2 mL), solution 3 (0.75 mL), and solution 4 (0.25 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) was then prepared by combining solution 5 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250.

The W1/O1/W2 emulsion was added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the nanocarriers to form. A portion of the nanocarriers were washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 21,000×g and 4 °C for 45 min, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure was repeated, and the pellet was re-suspended in phosphate buffered saline for a final nanocarrier dispersion of about 10 mg/mL.

Nanocarrier size was determined by dynamic light scattering. The amounts of peptide and rapamycin in the nanocarrier were determined by HPLC analysis. The total dry-nanocarrier mass per mL of suspension was determined by a gravimetric method.

Nanocarrier ID	Effective Diameter (nm)	Rapamycin Content (% w/w)	Peptide Content (% w/w)
2	227	9.0	2.5

Measurement of synthetic nanocarrier dimensions was obtained by dynamic light scattering (DLS). A suspension of the synthetic nanocarriers was diluted with purified water to achieve a final synthetic nanocarrier suspension concentration of approximately 0.01 to 0.1 mg/mL. The diluted suspension was prepared directly inside a suitable cuvette for DLS analysis. The cuvette was then placed in a Brookhaven Instruments Corp. ZetaPALS, allowed to equilibrate to 25 °C, and then scanned for sufficient time to acquire a stable and reproducible distribution based on appropriate inputs for viscosity of the medium and refractive indicies of the sample. The effective diameter, or mean of the distribution, was then reported.

Immunization

The nanocarriers were thawed and equilibrated. Initial dilutions constituted a 10x stock solution, and were further diluted to a concentration of 100μg/ml in $OVA_{323-339}$, or a 1x solution. This 1x solution was used for injections at 200μl per i.v. injection. Animals were immunized with OVA protein (OVA) and treated with $OVA_{323-339}$ peptide. Immunization routes were as follows: 10μg of OVA+ 4mg Alum i.p. in 400μl per each Balb/C immunologically naïve female mouse. Experimental groups consisted of 5 animals each. Spleen cells were restimulated with antigen using CFSE or CTO to determine the amount of Ag-specific proliferation.

Measuring Regulatory B Cells

The frequency of Ovalbumin reactive IL-10 secreting B cells was calculated by way of flow cytometry. Splenocytes from experimental animals were stained with CFSE, a thiol-reactive Fluorescent Probe suitable for long-term cell labeling, and cultured in complete media at 37C, 5% CO2 with Ovalbumin protein for 3 days. On day 3 the cells were assayed for their potential to secrete different cytokines by intracellular staining using standard

methods and kits. Briefly, cells were restimulated with phorbol myristate acetate (PMA) and Ionomycin for 2 hours and protein transport was blocked for another 4 hours. Unspecific binding of antibodies was blocked with anti-CD16/32 antibody and then cells were stained with conjugated antibodies specific recognizing specifically B cells (CD45R (B220) and CD19). After fixation with paraformaldehyde cells were permeabilized to allow monoclonal antibodies into the cells and label intracellular epitopes (cytokines). Splenocytes that were B220+CD19+ were assessed for proliferation by comparing the differential CFSE staining and the proportion of IL-10-secreting cells determined.

Measurement of IgE Antibodies

IgE antibodies were measured using a Mouse OVA-IgE ELISA kit provided by MDBioproducts (Cat# M036005) consistent with the manufacturer's instructions.

Results

Fig. 2 demonstrate the effectiveness of the nanocarriers in the generation of regulatory B cells in lavage samples from animal subjects treated with synthetic nanocarriers comprising $OVA_{323-339}$ and immunosuppressant. The figure demonstrates that synthetic nanocarriers of the invention resulted in the production of IL-10 and TGF-β by antigen-specific CD24+ B cells. **Fig. 3** shows a decrease in IgE production with synthetic nanocarriers of the invention.

Example 16: Isolation of Regulatory B cell (Bregs) Cells from a Subject after Administration of Inventive Synthetic Nanocarriers (Prophetic)

Bregs are isolated from biological samples, for example, from peripheral blood, obtained from a subject after the subject is administered a synthetic nanocarrier composition as described herein. Typically, the biological sample is obtained from the subject after a time period sufficient for the administered itDCs to induce Bregs. Bregs are isolated from the biological sample, for example, from whole blood, by negative and/or positive selection. For example, the cellular fraction of whole blood is obtained by centrifugation, and erythrocytes are lysed using erythrocyte lysis buffer. After lysis, peripheral blood mononuclear cells are depleted for CD4+ cells and CD8+ T cells. Subsequently, Bregs are enriched for by positive selection for the markers provided elsewhere herein or otherwise known in the art. In some embodiments, the isolation of regulatory B cells is performed with

a cocktail of biotinylated antibodies and anti-biotin magnetic beads for the recognition of one or more of these markers.

2012249493 29 May 2017

What is claimed is:

CLAIMS

1. A method comprising:

administering to a subject a composition that comprises:

(i) a first population of synthetic nanocarriers coupled to immunosuppressants, and

(ii) a second population of synthetic nanocarriers coupled to antigens that comprise B cell epitopes and/or MHC Class II-restricted epitopes,

wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight), and

wherein the composition is in an amount effective to generate antigen-specific regulatory B cells in the subject.

2. A method comprising:

generating antigen-specific regulatory B cells in a subject by administering a composition that comprises:

(i) a first population of synthetic nanocarriers coupled to immunosuppressants, and

(ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen,

wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight).

3. A method comprising:

administering to a subject a composition according to a protocol that was previously shown to generate antigen-specific regulatory B cells in one or more test subjects;

wherein the composition comprises:

(i) a first population of synthetic nanocarriers coupled to immunosuppressants, and

29 May 2017 2012249493

(ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen,

wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight).

4. The method of claim 2 or 3, wherein the composition is in an amount effective to generate antigen-specific regulatory B cells in the subject

5. The method of any of claims 1-4, wherein the first population and the second population are the same population.

6. The method of any of claims 1-5, wherein the method further comprises providing or identifying the subject.

7. The method of any of claims 1-6, wherein the antigen is a therapeutic protein, an autoantigen or an allergen, or is associated with an inflammatory disease, an autoimmune disease, organ or tissue rejection or graft versus host disease.

8. The method of any of claims 1-7, wherein the method further comprises assessing the generation of antigen-specific regulatory B cells in the subject prior to and/or after the administration of the composition.

9. The method of any of claims 1-8, wherein the subject has or is at risk of having an inflammatory disease, an autoimmune disease, an allergy, organ or tissue rejection or graft versus host disease.

10. The method of any of claims 1-8, wherein the subject has undergone or will undergo transplantation.

29 May 2017 2012249493

11. The method of any of claims 1-8, wherein the subject has or is at risk of having an undesired immune response against a therapeutic protein that is being administered or will be administered to the subject.

12. The method of any of claims 1-11, wherein one or more maintenance doses of the composition comprising the first population and second population of synthetic nanocarriers is administered to the subject.

13. The method of any of claims 1-12, wherein the administering is by intravenous, intraperitoneal, transmucosal, oral, subcutaneous, pulmonary, intranasal, intradermal or intramuscular administration.

14. The method of any of claims 1-12, wherein the administering is by inhalation or intravenous, subcutaneous or transmucosal administration.

15. The method of any of claims 1-14, wherein the method further comprises collecting the generated antigen-specific regulatory B cells.

16. The method of any of claims 1-15, wherein the immunosuppressants comprise a statin, an mTOR inhibitor, a TGF-β signaling agent, a corticosteroid, an inhibitor of mitochondrial function, a P38 inhibitor, an NF-κβ inhibitor, an adenosine receptor agonist, a prostaglandin E2 agonist, a phosphodiesterasse 4 inhibitor, an HDAC inhibitor or a proteasome inhibitor.

17. The method of claim 16, wherein the mTOR inhibitor is rapamycin.

18. The method of claims 1-17, wherein at least 80%, at least 90%, or at least 95% of the first and/or second population of synthetic nanocarriers have a minimum dimension or maximum dimension that falls within 5%, 10%, or 20% of the mean diameter of the synthetic nanocarriers.

29 May 2017 2012249493

19. The method of any of claims 1-18, wherein the synthetic nanocarriers of the first population and/or second population comprise lipid nanoparticles, polymeric nanoparticles, metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles or peptide or protein particles.

20. The method of claim 19, wherein the synthetic nanocarriers of the first population and/or second population comprise lipid nanoparticles.

21. The method of claim 20, wherein the synthetic nanocarriers of the first population and/or second population comprise liposomes.

22. The method of claim 19, wherein the synthetic nanocarriers of the first population and/or second population comprise metallic nanoparticles.

23. The method of claim 22, wherein the metallic nanoparticles comprise gold nanoparticles.

24. The method of claim 19, wherein the synthetic nanocarriers of the first population and/or second population comprise polymeric nanoparticles.

25. The method of claim 24, wherein the polymeric nanoparticle comprises polymer that is a non-methoxy-terminated, pluronic polymer.

26. The method of claim 24 or 25, wherein the polymeric nanoparticles comprise a polyester, a polyester coupled to a polyether, polyamino acid, polycarbonate, polyacetal, polyketal, polysaccharide, polyethyloxazoline or polyethyleneimine.

27. The method of claim 26, wherein the polyester comprises a poly(lactic acid), poly(glycolic acid), poly(lactic-co-glycolic acid) or polycaprolactone.

28. The method of claim 26 or 27, wherein the polymeric nanoparticles comprise a polyester and a polyester coupled to a polyether.

29 May 2017 2012249493

29. The method of any of claims 26-28, wherein the polyether comprises polyethylene glycol or polypropylene glycol.

30. The method of any of claims 1-29, wherein the mean of a particle size distribution obtained using dynamic light scattering of the synthetic nanocarriers of the first and/or second population is a diameter greater than 100nm.

31. The method of claim 30, wherein the diameter is greater than 150nm.

32. The method of claim 31, wherein the diameter is greater than 200nm.

33. The method of claim 32, wherein the diameter is greater than 250nm.

34. The method of claim 33, wherein the diameter is greater than 300nm.

35. The method of any of claims 1-34, wherein the synthetic nanocarriers of the first and/or second population of synthetic nanocarriers have a maximum dimension, obtained using dynamic light scattering, that is equal to or less than 500 nm.

36. The method of any of claims 1-35, wherein the aspect ratio of the synthetic nanocarriers of the first population and/or second population is greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7 or 1:10.

37. The method of any of claims 15-36, wherein the method further comprises making a dosage form comprising the collected antigen-specific regulatory B cells.

38. The method of any of claims 15-37, wherein the method further comprises making the collected antigen-specific regulatory B cells available or dosage form available to a subject for administration.

39. The method of any of claims 1-38, wherein the second population of synthetic nanocarriers are also coupled to MHC Class I-restricted epitopes.

29 May 2017 2012249493

40. A composition comprising isolated antigen-specific regulatory B cells produced according to the method of any of claims 1-39.

41. The composition of claim 40, wherein the composition further comprises a pharmaceutically acceptable excipient.

42. A dosage form comprising the composition of claims 40 or 41.

43. A composition comprising isolated antigen-specific regulatory B cells when used in therapy or prophylaxis, said isolated antigen-specific regulatory B cells being obtained by a process comprising the steps of:
(a) generating antigen-specific regulatory B cells in a subject by administering a composition that comprises: (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen;
wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight), and
(b) collecting the generated antigen-specific regulatory B cells.

44. The composition of claim 43, which is as defined in any one of claims 40 or 41.

45. A composition comprising: (i) a first population of synthetic nanocarriers coupled to immunosuppressants, and (ii) a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen, wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight), for use in therapy or prophylaxis.

29 May 2017 2012249493

46. The composition of claim 45 wherein the composition is as defined in any one of claims 1-39.

47. The composition of claim 40 or 41 or as defined in any one of claims 1-39 or 43, or dosage form of claim 42 for use in therapy or prophylaxis.

48. The composition of claim 40 or 41 or as defined in any one of claims 1-39 or 43, or dosage form of claim 42 for use in promoting a tolerogenic immune response or generating antigen-specific regulatory B cells in a subject.

49. Use of the composition of claim 40 or 41 or as defined in any one of claims 1-39 or 43, or dosage form of claim 42 for the manufacture of a medicament for use in promoting a tolerogenic immune response or generating antigen-specific regulatory B cells in a subject.

50. A dosage form comprising the composition of any of claims 43-48.

51. A method comprising:
(i) producing a first population of synthetic nanocarriers coupled to immunosuppressants, and
(ii) producing a second population of synthetic nanocarriers coupled to B cell and/or MHC Class II-restricted epitopes of an antigen, wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight), and
wherein the composition is in an amount effective to generate antigen-specific regulatory B cells in a subject.

52. The method of claim 51, wherein the first population and second population are the same population.

53. The method of claim 51 or 52, wherein the method further comprises making a dosage form comprising the first population and second population of synthetic nanocarriers.

29 May 2017 2012249493

54. The method of any of claims 51-53, wherein the method further comprises making a composition comprising the first population and second population of synthetic nanocarriers or the dosage form available to a subject for administration.

55. The method of any of claims 51-54, wherein the method further comprises assessing the generation of antigen-specific regulatory B cells with a composition comprising the first population and second population of synthetic nanocarriers.

56. The method of any of claims 51-55, wherein the first population and second population of synthetic nanocarriers that are produced are as defined in any of claims 1-39.

57. A process for producing a composition or dosage form comprising the steps of:
(i) coupling a first population of synthetic nanocarriers to immunosuppressants, and
(ii) coupling a second population of synthetic nanocarriers to B cell and/or MHC Class II-restricted epitopes of an antigen, wherein the load of the immunosuppressants on average across the first population of synthetic nanocarriers is at least 2% but no more than 25% (weight/weight) and the load of the antigens on average across the second population of synthetic nanocarriers is between 1% and 10% (weight/weight), and
wherein the composition or dosage form in in an amount effective to generate regulatory B cells in a subject.

58. The process of claim 57, which comprises the steps as defined in the method of any one of claims 51-56.

59. A composition or dosage form obtained by the method or process of any one of claims 51-58.


SSC
FSC
lymphocyte
41.98
SSC
CD3
CD3
95.19
SSC
CD25
CD25
8.60
FoxP3
CD4
FOXP3
53.10
53%
FSC/SSC CD3 CD25 gated

Fig. 1


CD24+OVA- IL10+ TGFb+
CD24+OVA+ IL10+ TGFb+
CD24-OVA+ IL10+ TGFb+
CD24-OVA- IL10+ TGFb+
% of CD19+ B220+ cells
40
30
20
10
0
PBS
NPOVA323-339
t2SVP

Fig. 2


20000
15000
10000
5000
0
IgE conc. (ng/ml)
PBS
NPOVA323-339
t2SVP

Fig 3

S168170050WO00-SEQLISTING-JAV
SEQUENCE LISTING

<110> Selecta Biosciences, Inc.
Maldonado, Roberto

<120> TOLEROGENIC SYNTHETIC NANOCARRIERS FOR INDUCING REGULATORY B CELLS

<130> S1681.70050WO00

<150> US 61/480,946
<151> 2011-04-29

<150> US 61/513,514
<151> 2011-07-29

<150> US 61/531,147
<151> 2011-09-06

<150> US 61/531,153
<151> 2011-09-06

<150> US 61/531,164
<151> 2011-09-06

<150> US 61/531,168
<151> 2011-09-06

<150> US 61/531,175
<151> 2011-09-06

<150> US 61/531,180
<151> 2011-09-06

<150> US 61/531,194
<151> 2011-09-06

<150> US 61/531,204
<151> 2011-09-06

<150> US 61/531,209
<151> 2011-09-06

<150> US 61/531,215
<151> 2011-09-06

<160> 943

<170> PatentIn version 3.5

<210> 1
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Aggrecan core protein precursor epitope

<400> 1

Ala Gly Met Asp Met Cys Ser Ala Gly Trp Leu Ala Asp Arg Ser Val
1 5 10 15

Arg Tyr

<210> 2

<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Aggrecan core protein precursor epitope

<400> 2

Glu Asp Ser Glu Ala Thr Leu Glu Val Val Val Lys Gly Ile Val Phe
1 5 10 15

His Tyr

<210> 3
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Aggrecan core protein precursor epitope

<400> 3

Ser Arg Val Ser Lys Glu Lys Glu Val Val Leu Leu Val Ala Thr Glu
1 5 10 15

Gly Arg

<210> 4
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Aggrecan core protein precursor epitope

<400> 4

Val Val Leu Leu Val Ala Thr Glu Gly Arg Val Arg Val Asn Ser Ala
1 5 10 15

Tyr Gln

<210> 5
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Aggrecan core protein precursor epitope

<400> 5

Val Val Val Lys Gly Ile Val Phe His Tyr Arg Ala Ile Ser Thr Arg
1 5 10 15

Tyr Thr

<210> 6
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens alpha 2 type VI collagen isoform 2C2 precursor epitope

<400> 6

Asp Arg Ala Ser Phe Ile Lys Asn Leu
1 5

<210> 7
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 7

Ala Ser Ser Thr Ile Ile Lys Glu Gly Ile Asp Arg Thr Val Leu Gly
1 5 10 15

Ile Leu Val Ser
20

<210> 8
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 8

Ala Ser Thr Pro Thr Lys Leu Gln Glu Ser Leu Leu Lys Lys Leu Gly
1 5 10 15

Ser Asn Thr Tyr
20

<210> 9
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 9

Asp Arg Thr Val Leu Gly Ile Leu Val Ser Tyr Gln Ile Lys Val Lys
1 5 10 15

Leu Thr Val Ser
20

<210> 10
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 10

Glu Phe Ala Arg His Asn Leu Lys Asp Ala Gly Glu Ala Glu Glu Gly
1 5 10 15

Lys Arg Asp Lys
20

<210> 11
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 11

Glu Pro Asn His Val Ile Phe Lys Lys Ile Ser Arg Asp Lys Ser Val
1 5 10 15

Thr Ile Tyr Leu
20

<210> 12
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 12

Phe Glu Val Lys Ala Phe Ala Thr Asp Ser Thr Asp Ala Glu Glu Asp
1 5 10 15

Lys Ile Pro Lys
20

<210> 13
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 13

Gly Phe Leu Gly Glu Leu Thr Ser Ser Glu Val Ala Thr Glu Val Pro
1 5 10 15

Phe Arg Leu Met
20

<210> 14
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 14

Gly Lys Ile Lys His Glu Asp Thr Asn Leu Ala Ser Ser Thr Ile Ile
1 5 10 15

Lys Glu Gly Ile
20

<210> 15
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 15

Gly Asn Arg Asp Tyr Ile Asp His Val Ser Gln Val Gln Pro Val Asp
1 5 10 15

Gly Val Val Leu
20

<210> 16
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 16

Lys Pro Val Ala Met Glu Glu Ala Gln Glu Lys Val Pro Pro Asn Ser
1 5 10 15

Thr Leu Thr Lys
20

<210> 17
<211> 20
<212> PRT
<213> Artificial Sequence

<220>

<223> Homo sapiens arrestin epitope

<400> 17

Lys Val Pro Pro Asn Ser Thr Leu Thr Lys Thr Leu Thr Leu Leu Pro
1 5 10 15

Leu Leu Ala Asn
20

<210> 18
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 18

Leu Leu Lys Lys Leu Gly Ser Asn Thr Tyr Pro Phe Leu Leu Thr Phe
1 5 10 15

Pro Asp Tyr Leu
20

<210> 19
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 19

Leu Thr Phe Arg Arg Asp Leu Tyr Phe Ser Arg Val Gln Val Tyr Pro
1 5 10 15

Pro Val Gly Ala
20

<210> 20
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 20

Met Ala Ala Ser Gly Lys Thr Ser Lys Ser Glu Pro Asn His Val Ile
1 5 10 15

Phe Lys Lys Ile
20

<210> 21
<211> 20

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 21

Asn Arg Glu Arg Arg Gly Ile Ala Leu Asp Gly Lys Ile Lys His Glu
1 5 10 15

Asp Thr Asn Leu
20

<210> 22
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 22

Pro Cys Ser Val Met Leu Gln Pro Ala Pro Gln Asp Ser Gly Lys Ser
1 5 10 15

Cys Gly Val Asp
20

<210> 23
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 23

Pro Phe Leu Leu Thr Phe Pro Asp Tyr Leu Pro Cys Ser Val Met Leu
1 5 10 15

Gln Pro Ala Pro
20

<210> 24
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 24

Gln Asp Ser Gly Lys Ser Cys Gly Val Asp Phe Glu Val Lys Ala Phe
1 5 10 15

Ala Thr Asp Ser
20

<210> 25
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 25

Gln Val Gln Pro Val Asp Gly Val Val Leu Val Asp Pro Asp Leu Val
1 5 10 15

Lys Gly Lys Lys
20

<210> 26
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 26

Arg Val Gln Val Tyr Pro Pro Val Gly Ala Ala Ser Thr Pro Thr Lys
1 5 10 15

Leu Gln Glu Ser
20

<210> 27
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 27

Ser Arg Asp Lys Ser Val Thr Ile Tyr Leu Gly Asn Arg Asp Tyr Ile
1 5 10 15

Asp His Val Ser
20

<210> 28
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 28

Thr Leu Thr Leu Leu Pro Leu Leu Ala Asn Asn Arg Glu Arg Arg Gly
1 5 10 15

Ile Ala Leu Asp
20

<210> 29
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 29

Val Ala Thr Glu Val Pro Phe Arg Leu Met His Pro Gln Pro Glu Asp
1 5 10 15

Pro Ala Lys Glu
20

<210> 30
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 30

Val Asp Pro Asp Leu Val Lys Gly Lys Lys Val Tyr Val Thr Leu Thr
1 5 10 15

Cys Ala Phe Arg
20

<210> 31
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 31

Val Val Leu Tyr Ser Ser Asp Tyr Tyr Val Lys Pro Val Ala Met Glu
1 5 10 15

Glu Ala Gln Glu
20

<210> 32
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens arrestin epitope

<400> 32

Tyr Gln Ile Lys Val Lys Leu Thr Val Ser Gly Phe Leu Gly Glu Leu
1 5 10 15

Thr Ser Ser Glu
20

<210> 33
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Chain B, Structure Of Insulin epitope

<400> 33

Ala Leu Tyr Leu Val Cys Gly Glu Arg
1 5

<210> 34
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Chain B, Structure Of Insulin epitope

<400> 34

Ser His Leu Val Glu Ala Leu Tyr Leu Val
1 5 10

<210> 35
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens chaperonin (HSP60) epitope

<400> 35

Gln Met Arg Pro Val Ser Arg Val Leu
1 5

<210> 36
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Collagen alpha-3(IV) chain epitope

<400> 36

Gly Ser Pro Ala Thr Trp Thr Thr Arg
1 5

<210> 37
<211> 9

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens collagen, type II, alpha 1 isoform 1 precursor epitope

<400> 37

Ala Arg Gly Gln Pro Gly Val Met Gly
1 5

<210> 38
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens DNA topoisomerase 1 epitope

<400> 38

Lys Met Leu Asp His Glu Tyr Thr Thr
1 5

<210> 39
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens ezrin epitope

<400> 39

Glu Tyr Thr Ala Lys Ile Ala Leu Leu
1 5

<210> 40
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens ezrin epitope

<400> 40

Leu Asn Ile Tyr Glu Lys Asp Asp Lys Leu
1 5 10

<210> 41
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glial fibrillary acidic protein isoform 2 epitope

<400> 41

Asn Leu Ala Gln Asp Leu Ala Thr Val
1 5

<210> 42
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glial fibrillary acidic protein isoform 2 epitope

<400> 42

Gln Leu Ala Arg Gln Gln Val His Val
1 5

<210> 43
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glucagon receptor epitope

<400> 43

Arg Arg Arg Trp His Arg Trp Arg Leu
1 5

<210> 44
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glucose-6-phosphatase, catalytic, related epitope

<400> 44

Phe Leu Trp Ser Val Phe Trp Leu Ile
1 5

<210> 45
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 1 epitope

<400> 45

Asn Met Phe Thr Tyr Glu Ile Ala Pro Val Phe Val Leu Met Glu
1 5 10 15

<210> 46
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 46

Ile Ala Phe Thr Ser Glu His Ser His Phe Ser Leu Lys

1 5 10

<210> 47
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 47

Asn Phe Phe Arg Met Val Ile Ser Asn Pro Ala Ala Thr
1 5 10

<210> 48
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 48

Phe Leu Gln Asp Val Met Asn Ile Leu
1 5

<210> 49
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 49

Leu Leu Gln Glu Tyr Asn Trp Glu Leu
1 5

<210> 50
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 50

Arg Met Met Glu Tyr Gly Thr Thr Met Val
1 5 10

<210> 51
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 51

Val Met Asn Ile Leu Leu Gln Tyr Val Val
1 5 10

<210> 52
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 52

Ala Phe Thr Ser Glu His Ser His Phe Ser Leu
1 5 10

<210> 53
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 53

Ala Phe Thr Ser Glu His Ser His Phe Ser Leu Lys
1 5 10

<210> 54
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 54

Phe Lys Met Phe Pro Glu Val Lys Glu Lys Gly
1 5 10

<210> 55
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 55

Phe Thr Ser Glu His Ser His Phe Ser Leu
1 5 10

<210> 56
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 56

Met Ile Ala Arg Phe Lys Met Phe Pro Glu Val Lys Glu Lys Gly
1 5 10 15

<210> 57
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 57

Arg Phe Lys Met Phe Pro Glu Val Lys
1 5

<210> 58
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 58

Arg Phe Lys Met Phe Pro Glu Val Lys Glu
1 5 10

<210> 59
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 59

Arg Phe Lys Met Phe Pro Glu Val Lys Glu Lys
1 5 10

<210> 60
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 60

Thr Ser Glu His Ser His Phe Ser Leu
1 5

<210> 61
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 61

Val Met Asn Ile Leu Leu Gln Tyr Val
1 5

<210> 62
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 62

Glu Leu Ala Glu Tyr Leu Tyr Asn Ile
1 5

<210> 63
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 63

Ile Leu Met His Cys Gln Thr Thr Leu
1 5

<210> 64
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens heat shock 27kDa protein 1 epitope

<400> 64

Gln Leu Ser Ser Gly Val Ser Glu Ile Arg His
1 5 10

<210> 65
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA class I histocompatibility antigen, B-27 alpha chain precursor epitope

<400> 65

Leu Arg Arg Tyr Leu Glu Asn Gly Lys
1 5

<210> 66
<211> 9

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA class I histocompatibility antigen, B-7 alpha chain precursor epitope

<400> 66

Val Met Ala Pro Arg Thr Val Leu Leu
1 5

<210> 67
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA class I histocompatibility antigen, B-7 alpha chain precursor epitope

<400> 67

Ala Leu Asn Glu Asp Leu Arg Ser Trp Thr Ala Ala Asp Thr
1 5 10

<210> 68
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA-B27 epitope

<400> 68

Ala Leu Asn Glu Asp Leu Ser Ser Trp Thr Ala Ala Asp Thr
1 5 10

<210> 69
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA-B27 epitope

<400> 69

Leu Leu Arg Gly Tyr His Gln Asp Ala Tyr
1 5 10

<210> 70
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA-B27 epitope

<400> 70

Arg Val Ala Glu Gln Leu Arg Ala Tyr Leu Glu Gly Glu Cys Val
1 5 10 15

<210> 71
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA-B27 epitope

<400> 71

Trp Asp Arg Glu Thr Gln Ile Cys Lys Ala Lys Ala Gln
1 5 10

<210> 72
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens insulin epitope

<400> 72

Ala Leu Trp Gly Pro Asp Pro Ala Ala Ala Phe
1 5 10

<210> 73
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens insulin epitope

<400> 73

Leu Ala Leu Trp Gly Pro Asp Pro Ala Ala
1 5 10

<210> 74
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens insulin epitope

<400> 74

Arg Leu Leu Pro Leu Leu Ala Leu Leu Ala Leu
1 5 10

<210> 75
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 75

Ala Leu Trp Met Arg Leu Leu Pro Leu
1 5

<210> 76
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 76

His Leu Val Glu Ala Leu Tyr Leu Val
1 5

<210> 77
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 77

Ser Leu Gln Lys Arg Gly Ile Val Glu Gln
1 5 10

<210> 78
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 78

Ser Leu Gln Pro Leu Ala Leu Glu Gly
1 5

<210> 79
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 79

Ser Leu Tyr Gln Leu Glu Asn Tyr Cys
1 5

<210> 80
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 80

Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr
1 5 10

<210> 81
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 81

Trp Gly Pro Asp Pro Ala Ala Ala
1 5

<210> 82
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 82

Phe Tyr Thr Pro Lys Thr Arg Arg Glu
1 5

<210> 83
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 83

Gly Glu Arg Gly Phe Phe Tyr Thr
1 5

<210> 84
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 84

Glu Arg Gly Phe Phe Tyr Thr Pro Lys
1 5

<210> 85
<211> 10
<212> PRT
<213> Artificial Sequence

<220>

<223> Homo sapiens Insulin precursor epitope

<400> 85

Leu Cys Gly Ser His Leu Val Glu Ala Leu
1 5 10

<210> 86
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 86

Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr
1 5 10

<210> 87
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 87

Leu Tyr Leu Val Cys Gly Glu Arg Gly Phe
1 5 10

<210> 88
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Islet amyloid polypeptide precursor epitope

<400> 88

Phe Leu Ile Val Leu Ser Val Ala Leu
1 5

<210> 89
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Islet amyloid polypeptide precursor epitope

<400> 89

Lys Leu Gln Val Phe Leu Ile Val Leu
1 5

<210> 90
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens islet-specific glucose-6-phosphatase-related protein epitope

<400> 90

Phe Leu Trp Ser Val Phe Met Leu Ile
1 5

<210> 91
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens islet-specific glucose-6-phosphatase-related protein isoform 1 epitope

<400> 91

Phe Leu Phe Ala Val Gly Phe Tyr Leu
1 5

<210> 92
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens islet-specific glucose-6-phosphatase-related protein isoform 1 epitope

<400> 92

Leu Asn Ile Asp Leu Leu Trp Ser Val
1 5

<210> 93
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens islet-specific glucose-6-phosphatase-related protein isoform 1 epitope

<400> 93

Val Leu Phe Gly Leu Gly Phe Ala Ile
1 5

<210> 94
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens islet-specific glucose-6-phosphatase-related protein isoform 1 epitope

<400> 94

Asn Leu Phe Leu Phe Leu Phe Ala Val

1 5

<210> 95
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens islet-specific glucose-6-phosphatase-related protein isoform 1 epitope

<400> 95

Tyr Leu Leu Leu Arg Val Leu Asn Ile
1 5

<210> 96
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 96

Ala Leu Gln Lys Ala Lys Gln Asp Leu
1 5

<210> 97
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 97

Asp Ala Lys Asn Lys Leu Glu Gly Leu
1 5

<210> 98
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 98

Gly Ala Ser Gly Val Gly Ser Gly Leu
1 5

<210> 99
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 99

Lys Ala Lys Gln Asp Leu Ala Arg Leu
1 5

<210> 100
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 100

Lys Leu Glu Gly Leu Glu Asp Ala Leu
1 5

<210> 101
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 101

Asn Met Gln Asp Leu Val Glu Asp Leu
1 5

<210> 102
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 102

Arg Leu Leu Lys Glu Tyr Gln Glu Leu
1 5

<210> 103
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens keratin 6C epitope

<400> 103

Trp Tyr Gln Thr Lys Tyr Glu Glu Leu
1 5

<210> 104
<211> 20
<212> PRT
<213> Artificial Sequence

<220>

<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 104

Leu Arg Arg Val Leu Asp Glu Leu Thr Leu Ala Arg Thr Asp Leu Glu
1 5 10 15

Met Gln Ile Glu
20

<210> 105
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 105

Ala Leu Glu Glu Ala Asn Ala Asp Leu
1 5

<210> 106
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 106

Ala Asn Ala Asp Leu Glu Val Lys Ile
1 5

<210> 107
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 107

Ala Arg Thr Asp Leu Glu Met Gln Ile
1 5

<210> 108
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 108

Ala Ser Tyr Leu Asp Lys Val Arg Ala
1 5

<210> 109
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 109

Asp Val Asn Gly Leu Arg Arg Val Leu
1 5

<210> 110
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 110

Gly Leu Arg Arg Val Leu Asp Glu Leu
1 5

<210> 111
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 111

Ile Ser Ser Val Leu Ala Gly Ala Ser Cys Pro Ala
1 5 10

<210> 112
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 112

Leu Asp Lys Val Arg Ala Leu Glu Glu
1 5

<210> 113
<211> 9
<212> PRT

<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 113

Gln Ile Glu Gly Leu Lys Glu Glu Leu
1 5

<210> 114
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 114

Arg Ala Leu Glu Glu Ala Asn Ala Asp Leu Glu Val
1 5 10

<210> 115
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 115

Arg Leu Ala Ser Tyr Leu Asp Lys Val
1 5

<210> 116
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 116

Ser Tyr Leu Asp Lys Val Arg Ala
1 5

<210> 117
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Keratin, type I cytoskeletal 17 (Cytokeratin 17) (K17) (CK 17) (Version 2) epitope

<400> 117

Ser Tyr Leu Asp Lys Val Arg Ala Leu Glu Glu Ala Asn Ala Asp Leu
1 5 10 15

Glu Val Lys Ile
20

<210> 118
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens maspin epitope

<400> 118

Gly Leu Glu Lys Ile Glu Lys Gln Leu
1 5

<210> 119
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens maspin epitope

<400> 119

Met Gly Asn Ile Asp Ser Ile Asn Cys Lys
1 5 10

<210> 120
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens maspin epitope

<400> 120

Tyr Ser Leu Lys Leu Ile Lys Arg Leu
1 5

<210> 121
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 121

Ala Ser Gln Lys Arg Pro Ser Gln Arg His Gly Ser Lys Tyr Leu Ala
1 5 10 15

Thr Ala Ser Thr
20

<210> 122
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 122

Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
1 5 10 15

<210> 123
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 123

Val Val His Phe Phe Lys Asn Ile Val
1 5

<210> 124
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 124

Asp Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
1 5 10 15

Thr Pro Pro

<210> 125
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 125

His His Pro Ala Arg Thr Ala His Tyr Gly Ser Leu Pro Gln Lys Ser
1 5 10 15

His Gly Arg Thr
20

<210> 126
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 126

Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg Thr Pro Pro Pro
1 5 10 15

Ser Gln Gly Lys
20

<210> 127
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 127

Ala Ser Gln Lys Arg Pro Ser Gln Arg His Gly Ser Lys Tyr Leu Ala
1 5 10 15

Thr Ala Ser Thr Met
20

<210> 128
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 128

Phe Lys Gly Val Asp Ala Gln Gly Thr Leu Ser Lys Ile Phe Lys Leu
1 5 10 15

Gly Gly Arg Asp
20

<210> 129
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 129

Arg Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys Ser Ala
1 5 10 15

His Lys Gly

<210> 130
<211> 38
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 130

Ala Ser Gln Lys Arg Pro Ser Gln Arg His Gly Ser Lys Tyr Leu Ala
1 5 10 15

Thr Ala Ser Thr Met Asp His Ala Arg His Gly Phe Leu Pro Arg His
20 25 30

Arg Asp Thr Gly Ile Leu
35

<210> 131
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 131

Lys Tyr Leu Ala Thr Ala Ser Thr Met
1 5

<210> 132
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 132

Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln Arg Pro
1 5 10 15

Gly Phe Gly Tyr
20

<210> 133
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 133

Phe Gly Gly Asp Arg Gly Ala Pro Lys Arg Gly Ser Gly Lys Asp Ser
1 5 10 15

His His Pro Ala Arg Thr Ala His Tyr Gly Ser Leu Pro Gln Lys Ser
20 25 30

His Gly Arg Thr Gln Asp Glu Asn Pro Val Val
35 40

<210> 134
<211> 40
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 134

Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln Arg Pro
1 5 10 15

Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys Ser Ala His Lys
20 25 30

Gly Phe Lys Gly Val Asp Ala Gln
35 40

<210> 135
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MHC class I related protein A epitope

<400> 135

Ala Ala Ala Ala Ala Ile Phe Val Ile
1 5

<210> 136
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin basic protein epitope

<400> 136

Ser Leu Ser Arg Phe Ser Trp Gly Ala
1 5

<210> 137
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin basic protein epitope

<400> 137

Asp Tyr Lys Ser Ala His Lys Gly Phe
1 5

<210> 138
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin basic protein epitope

<400> 138

Ser Lys Ile Phe Lys Leu Gly Gly Arg Asp Ser Arg Ser Gly Ser Pro
1 5 10 15

Met Ala Arg

<210> 139
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin basic protein epitope

<400> 139

Thr Pro Arg Thr Pro Pro Pro Gln
1 5

<210> 140
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin proteolipid protein epitope

<400> 140

Phe Leu Tyr Gly Ala Leu Leu Leu Ala
1 5

<210> 141
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin proteolipid protein epitope

<400> 141

Lys Leu Ile Glu Thr Tyr Phe Ser Lys
1 5

<210> 142
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin-associated glycoprotein precursor epitope

<400> 142

Leu Met Trp Ala Lys Ile Gly Pro Val
1 5

<210> 143
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin-associated glycoprotein precursor epitope

<400> 143

Ser Leu Leu Leu Glu Leu Glu Glu Val
1 5

<210> 144
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin-associated glycoprotein precursor epitope

<400> 144

Val Leu Phe Ser Ser Asp Phe Arg Ile
1 5

<210> 145
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myosin heavy chain, skeletal muscle, adult 2 (Myosin heavy chain IIa) (MyHC-IIa) epitope

<400> 145

Glu Phe Gln Lys Met Arg Arg Asp Leu
1 5

<210> 146
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myosin heavy chain, skeletal muscle, adult 2 (Myosin heavy chain IIa) (MyHC-IIa) epitope

<400> 146

Lys Met Arg Arg Asp Leu Glu Glu Ala
1 5

<210> 147
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens peroxiredoxin-2 isoform a epitope

<400> 147

Glu Val Lys Leu Ser Asp Tyr Lys Gly Lys Tyr Val
1 5 10

<210> 148
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 148

His Leu Cys Gly Ser His Leu Val Glu Ala
1 5 10

<210> 149
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 149

Ala Leu Trp Gly Pro Asp Pro Ala Ala Ala
1 5 10

<210> 150
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 150

Arg Leu Leu Pro Leu Leu Ala Leu Leu
1 5

<210> 151
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 151

Ala Leu Trp Met Arg Leu Leu Pro Leu Leu
1 5 10

<210> 152
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 152

Trp Met Arg Leu Leu Pro Leu Leu Ala Leu
1 5 10

<210> 153
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 153

Pro Leu Ala Leu Glu Gly Ser Leu Gln Lys
1 5 10

<210> 154
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proinsulin precursor epitope

<400> 154

Pro Leu Leu Ala Leu Leu Ala Leu Trp Gly
1 5 10

<210> 155
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 155

Leu Leu Pro Pro Leu Leu Glu His Leu
1 5

<210> 156
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 156

Ser Leu Ala Ala Gly Val Lys Leu Leu
1 5

<210> 157
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 157

Ser Leu Ser Pro Leu Gln Ala Glu Leu
1 5

<210> 158
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 158

Ala Leu Thr Ala Val Ala Glu Glu Val
1 5

<210> 159
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 159

Ser Leu Tyr His Val Tyr Glu Val Asn Leu
1 5 10

<210> 160
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 160

Thr Ile Ala Asp Phe Trp Gln Met Val
1 5

<210> 161
<211> 9

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 161

Val Ile Val Met Leu Thr Pro Leu Val
1 5

<210> 162
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 162

Met Val Trp Glu Ser Gly Cys Thr Val
1 5

<210> 163
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 163

Phe Leu Gly Glu Leu Thr Ser Ser Glu Val Ala Thr Glu Val
1 5 10

<210> 164
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 164

Phe Met Ser Asp Lys Pro Leu His Leu Ala Val Ser Leu Asn Lys Glu
1 5 10 15

Ile Tyr Phe His
20

<210> 165
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 165

Gly Glu Ala Glu Glu Gly Lys Arg Asp Lys Asn Asp Val Asp Glu
1 5 10 15

<210> 166
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 166

Gly Glu Pro Ile Pro Val Thr Val Thr Val Thr Asn Asn Thr Glu Lys
1 5 10 15

Thr Val Lys Lys
20

<210> 167
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 167

His Pro Gln Pro Glu Asp Pro Ala Lys Glu Ser Tyr Gln Asp Ala Asn
1 5 10 15

Leu Val Phe Glu
20

<210> 168
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 168

Ile Lys Ala Phe Val Glu Gln Val Ala Asn Val Val Leu Tyr Ser Ser
1 5 10 15

Asp Tyr Tyr Val
20

<210> 169
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 169

Lys Ser Ser Val Arg Leu Leu Ile Arg Lys Val Gln His Ala Pro Leu
1 5 10 15

Glu Met Gly Pro
20

<210> 170
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 170

Gln Pro Arg Ala Glu Ala Ala Trp Gln Phe Phe Met Ser Asp Lys Pro
1 5 10 15

Leu His Leu Ala
20

<210> 171
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 171

Ser Tyr Gln Asp Ala Asn Leu Val Phe Glu Glu Phe Ala Arg His Asn
1 5 10 15

Leu Lys Asp Ala
20

<210> 172
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 172

Thr Asp Ala Glu Glu Asp Lys Ile Pro Lys Lys Ser Ser Val Arg Leu
1 5 10 15

Leu Ile Arg Lys
20

<210> 173
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 173

Thr Asn Asn Thr Glu Lys Thr Val Lys Lys Ile Lys Ala Phe Val Glu
1 5 10 15

Gln Val Ala Asn
20

<210> 174
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 174

Val Gln His Ala Pro Leu Glu Met Gly Pro Gln Pro Arg Ala Glu Ala
1 5 10 15

Ala Trp Gln Phe
20

<210> 175
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 175

Val Ser Leu Asn Lys Glu Ile Tyr Phe His Gly Glu Pro Ile Pro Val
1 5 10 15

Thr Val Thr Val
20

<210> 176
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 176

Val Tyr Val Thr Leu Thr Cys Ala Phe Arg Tyr Gly Gln Glu Asp Ile
1 5 10 15

Asp Val Ile Gly
20

<210> 177
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 177

Tyr Gly Gln Glu Asp Ile Asp Val Ile Gly Leu Thr Phe Arg Arg Asp
1 5 10 15

Leu Tyr Phe Ser
20

<210> 178
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens SSA protein SS-56 epitope

<400> 178

Tyr Thr Cys Pro Leu Cys Arg Ala Pro Val
1 5 10

<210> 179
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Steroid 21-hydroxylase epitope

<400> 179

Glu Pro Leu Ala Arg Leu Glu Leu
1 5

<210> 180
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Steroid 21-hydroxylase epitope

<400> 180

Glu Pro Leu Ala Arg Leu Glu Leu Phe Val Val Leu Thr Arg Leu Leu
1 5 10 15

Gln Ala Phe Thr
20

<210> 181
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Steroid 21-hydroxylase epitope

<400> 181

Ile Lys Asp Asp Asn Leu Met Pro Ala Tyr Tyr Lys Cys Ile Gln Glu
1 5 10 15

Val Leu Lys Thr
20

<210> 182
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Steroid 21-hydroxylase epitope

<400> 182

Ile Arg Asp Ser Met Glu Pro Val Val Glu Gln Leu Thr Gln Glu Phe
1 5 10 15

Cys Glu Arg Met
20

<210> 183
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T-cell receptor V beta chain 13.1 epitope

<400> 183

Leu Gly Arg Ala Gly Leu Thr Tyr
1 5

<210> 184
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens transaldolase 1 epitope

<400> 184

Leu Leu Phe Ser Phe Ala Gln Ala Val
1 5

<210> 185
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Vasoactive intestinal polypeptide receptor 1 precursor epitope

<400> 185

Arg Arg Lys Trp Arg Arg Trp His Leu
1 5

<210> 186
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Vasoactive intestinal polypeptide receptor 1 precursor epitope

<400> 186

Arg Arg Lys Trp Arg Arg Trp His Leu
1 5

<210> 187
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea 2S protein 1 epitope

<400> 187

Ala His Ala Ser Ala Arg Gln Gln Trp Glu Leu Gln Gly Asp Arg Arg
1 5 10 15

Cys Gln Ser Gln
20

<210> 188
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea 2S protein 1 epitope

<400> 188

Ala Lys Leu Thr Ile Leu Val Ala Leu Ala Leu Phe Leu Leu Ala Ala
1 5 10 15

His Ala Ser Ala
20

<210> 189
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea 2S protein 1 epitope

<400> 189

Ala Leu Gln Gln Ile Met Glu Asn Gln Ser Asp Arg Leu Gln Gly Arg
1 5 10 15

Gln Gln Glu

<210> 190
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea 2S protein 1 epitope

<400> 190

Ala Asn Leu Arg Pro Cys Glu Gln His Leu Met Gln Lys Ile Gln Arg
1 5 10 15

Asp Glu Asp Ser
20

<210> 191
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea 2S protein 1 epitope

<400> 191

Cys Asn Glu Leu Asn Glu Phe Glu Asn Asn Gln Arg Cys Met Cys Glu
1 5 10 15

Ala Leu Gln Gln
20

<210> 192
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 5-hydroxytryptamine receptor 2C (5-HT-2C) (Serotonin receptor 2C) (5-HT2C) (5-HTR2C) (5HT-1C) epitope

<400> 192

Pro Arg Gly Thr Met Gln Ala Ile Asn Asn Glu Arg Lys Ala Ser Lys
1 5 10 15

<210> 193
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 193

Asp Gln Gly Thr Cys Leu Leu Leu Thr Glu Val Ala
1 5 10

<210> 194
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 194

Glu Leu Glu Lys Tyr Gln Gln Leu Asn Ser Glu Arg Gly Val
1 5 10

<210> 195
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 195

Gly Glu Arg Ile Thr Lys Met Thr Glu Gly Leu Ala Lys
1 5 10

<210> 196
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 196

Pro Gly Glu Trp Arg Ile Ile Tyr Ala Ala Ala Asp Asn Lys
1 5 10

<210> 197
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 197

Arg Ile Glu Cys Ile Asn Asp Cys
1 5

<210> 198
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 198

Val Ala Lys Arg Gln Glu Gly Tyr Val Tyr Val Leu
1 5 10

<210> 199
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 199

Val Ser Glu Asn Met Leu Val Thr Tyr Val
1 5 10

<210> 200
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 200

Asp Gln Gly Thr Cys Leu Leu Leu Thr Glu Val Ala
1 5 10

<210> 201
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 201

Glu Leu Glu Lys Tyr Gln Gln Leu Asn Ser Glu Arg Gly Val
1 5 10

<210> 202
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 202

Glu Leu Glu Lys Tyr Gln Gln Leu Asn Ser Glu Arg Gly Val Pro Asn
1 5 10 15

<210> 203
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 203

Gly Glu Arg Ile Thr Lys Met Thr Glu Gly Leu Ala Lys
1 5 10

<210> 204
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 204

Pro Gly Glu Trp Arg Ile Ile Tyr Ala Ala Ala Asp Asn Lys
1 5 10

<210> 205
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 205

Arg Ile Glu Cys Ile Asn Asp Cys
1 5

<210> 206
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 206

Val Ala Lys Arg Gln Glu Gly Tyr Val Tyr Val Leu
1 5 10

<210> 207
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Allergen Bos d 2 precursor epitope

<400> 207

Val Ser Glu Asn Met Leu Val Thr Tyr Val
1 5 10

<210> 208
<211> 13
<212> PRT

<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Allergen Cry j 2 epitope

<400> 208

Asp Ile Phe Ala Ser Lys Asn Phe His Leu Gln Lys Asn
1 5 10

<210> 209
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Allergen Cry j 2 epitope

<400> 209

Gly Ile Ile Ala Ala Tyr Gln Asn Pro Ala Ser Trp Lys
1 5 10

<210> 210
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Allergen Cry j 2 epitope

<400> 210

Lys Leu Thr Ser Gly Lys Ile Ala Ser Cys Leu Asn
1 5 10

<210> 211
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Allergen Cry j 2 epitope

<400> 211

Gln Phe Ala Lys Leu Thr Gly Phe Thr Leu Met Gly
1 5 10

<210> 212
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 212

Ile Asn Gln Gln Leu Asn Pro Lys
1 5

<210> 213

<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 213

Ile Asn Gln Gln Leu Asn Pro Lys Thr Asn Lys Trp Glu Asp Lys
1 5 10 15

<210> 214
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 214

Leu Asn Pro Lys Thr Asn Lys Trp Glu Asp Lys
1 5 10

<210> 215
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 215

Ile Asn Gln Gln Leu Asn Pro Lys
1 5

<210> 216
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 216

Ile Asn Gln Gln Leu Asn Pro Lys Thr Asn Lys Trp Glu Asp Lys
1 5 10 15

<210> 217
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 217

Leu Asn Pro Lys Thr Asn Lys Trp Glu Asp Lys
1 5 10

<210> 218
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 218

Thr Asn Lys Trp Glu Asp Lys
1 5

<210> 219
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 219

Leu Asn Pro Lys Thr Asn Lys Trp Glu Asp Lys Arg
1 5 10

<210> 220
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Allergen Mag epitope

<400> 220

Pro Arg Leu Ser Trp His Gln Tyr Thr Lys Arg Asp Ser Arg Glu
1 5 10 15

<210> 221
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Allergen Mag epitope

<400> 221

Thr Val Asp Leu Ile Ser Pro Val Thr Lys Arg Ala Ser Leu Lys
1 5 10 15

<210> 222
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 222

Ala Trp Tyr Tyr Val Pro Leu Gly Thr Gln Tyr Thr Asp Ala Pro Ser

1 5 10 15

Phe Ser

<210> 223
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 223

Asp Ala Tyr Pro Ser Gly Ala Trp Tyr Tyr Val Pro Leu Gly Thr Gln
1 5 10 15

Tyr Thr

<210> 224
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 224

Asp Ile Gly Ser Glu Ser Thr Glu Asp Gln Ala Met Glu Asp Ile Lys
1 5 10 15

Gln Met

<210> 225
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 225

Glu Asp Ile Lys Gln Met
1 5

<210> 226
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 226

Glu Pro Met Ile Gly Val Asn Gln Glu Leu Ala Tyr

1 5 10

<210> 227
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 227

Glu Pro Met Ile Gly Val Asn Gln Glu Leu Ala Tyr Phe Tyr Pro Glu
1 5 10 15

Leu Phe

<210> 228
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea Ara h 2.01 allergen epitope

<400> 228

Glu Leu Asn Glu Phe Glu Asn Asn Gln Arg Cys Met Cys Glu Ala Leu
1 5 10 15

Gln

<210> 229
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea Ara h 2.01 allergen epitope

<400> 229

Ser Gln Leu Glu Arg Ala Asn Leu Arg Pro Cys Glu Gln His Leu Met
1 5 10 15

<210> 230
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Cry j 1 precursor epitope

<400> 230

Gly Ala Thr Arg Asp Arg Pro Leu Trp Ile Ile Phe Ser Gly Asn
1 5 10 15

<210> 231

<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Cry j 1 precursor epitope

<400> 231

Ile Phe Ser Gly Asn Met Asn Ile Lys Leu Lys Met Pro Met Tyr Ile
1 5 10 15

Ala Gly Tyr Lys
20

<210> 232
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Cry j 1 precursor epitope

<400> 232

Lys Met Pro Met Tyr Ile Ala Gly Tyr Lys Thr Phe Asp Gly Arg Gly
1 5 10 15

Ala Gln Val Tyr
20

<210> 233
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Cry j 1 precursor epitope

<400> 233

Leu Gly His Asp Asp Ala Tyr Ser Asp Asp Lys Ser Met Lys Val Thr
1 5 10 15

Val Ala Phe Asn
20

<210> 234
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Cry j 1 precursor epitope

<400> 234

Ser Gly Lys Tyr Glu Gly Gly Asn Ile Tyr Thr Lys Lys Glu Ala Phe
1 5 10 15

Asn Val Glu

<210> 235
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus Cytochrome c epitope

<400> 235

Glu Asn Pro Lys Lys Tyr Ile Pro Gly Thr Lys
1 5 10

<210> 236
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus Cytochrome c epitope

<400> 236

Gly Leu Phe Gly Arg Lys Thr Gly Ser Val Ala
1 5 10

<210> 237
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus Cytochrome c epitope

<400> 237

Lys Ile Gly Pro Glu Leu His Gly Leu
1 5

<210> 238
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus Cytochrome c epitope

<400> 238

Leu Lys Ala Gly Glu Gly Asn Lys Ile Gly Pro Glu
1 5 10

<210> 239
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus Cytochrome c epitope

<400> 239

Leu Lys Lys Pro Lys Asp Arg Asn Asp Leu Ile
1 5 10

<210> 240
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Der f 2 allergen epitope

<400> 240

Gly Leu Glu Ile Asp Val Pro Gly Ile Asp Thr Asn Ala Cys His Phe
1 5 10 15

Val Lys

<210> 241
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Der f 2 allergen epitope

<400> 241

Pro Gly Ile Asp Thr Asn Ala Cys His Phe Val Lys Cys Pro Leu Val
1 5 10 15

Lys Gly Gln Gln
20

<210> 242
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Der p 1 allergen epitope

<400> 242

Arg Phe Gly Ile Ser Asn Tyr Cys Gln Ile Tyr Pro Pro Asn Ala Asn
1 5 10 15

Lys Ile Arg

<210> 243
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Der p 1 allergen epitope

<400> 243

Ala Val Asn Ile Val Gly Tyr Ser Asn Ala Gln Gly Val Asp Tyr
1 5 10 15

<210> 244
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi globin Ctt 3-1 epitope

<400> 244

Phe Ala Gly Lys Asp Leu Glu Ser Ile Lys Gly Thr Ala Pro Phe Glu
1 5 10 15

Thr His Ala Asn
20

<210> 245
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi globin Ctt 3-1 epitope

<400> 245

Gly Thr Ala Pro Phe Glu Thr His Ala Asn Arg
1 5 10

<210> 246
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi globin Ctt 3-1 epitope

<400> 246

Lys Gly Thr Ala Pro Phe Glu Thr His Ala Asn Arg Ile Val Gly Phe
1 5 10 15

Phe Ser Lys Ile Ile
20

<210> 247
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III epitope

<400> 247

Ala His Thr Asp Phe Ala Gly Ala Glu Ala Ala Trp Gly Ala Thr Leu
1 5 10 15

Asp Thr Phe Phe Gly
            20

<210> 248
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III epitope

<400> 248

Phe Ala Gly Lys Asp Leu Glu Ser Ile Lys Gly Thr Ala Pro Phe Glu
1   5   10   15

Ile His Ala Asn
            20

<210> 249
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III epitope

<400> 249

Val Asn Thr Phe Val Ala Ser His Lys Pro Arg Gly Val Thr His Asp
1   5   10   15

Gln Leu Asn Asn Phe
            20

<210> 250
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III precursor epitope

<400> 250

Ala Asp Pro Ser Ile Met Ala Lys
1   5

<210> 251
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III precursor epitope

<400> 251

Ala Asp Pro Ser Ile Met Ala Lys Phe Thr Gln Phe Ala Gly Lys Asp
1   5   10   15

Leu Glu Ser Ile Lys
20

<210> 252
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III precursor epitope

<400> 252

Ala Glu Ala Ala Trp
1 5

<210> 253
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III precursor epitope

<400> 253

Ala Glu Ala Ala Trp Gly Ala Thr Leu Asp Thr Phe Phe Gly Met Ile
1 5 10 15

Phe Ser Lys Met
20

<210> 254
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III precursor epitope

<400> 254

Ala Gly Phe Val Ser Tyr Met Lys
1 5

<210> 255
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phaseolus vulgaris Glycine-rich cell wall structural protein 1.8 precursor epitope

<400> 255

Gly Gly Tyr Gly Asp Gly Gly Ala His Gly Gly Gly Tyr Gly Gly
1 5 10 15

<210> 256
<211> 15

<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Group V allergen Phl p 5 epitope

<400> 256

Ala Thr Pro Glu Ala Lys Tyr Asp Ala Tyr Val Ala Thr Leu Ser
1 5 10 15

<210> 257
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Group V allergen Phl p 5 epitope

<400> 257

Phe Thr Val Phe Glu Ala Ala Phe Asn Asn Ala Ile Lys Ala Gly
1 5 10 15

<210> 258
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Group V allergen Phl p 5 epitope

<400> 258

Lys Tyr Asp Ala Tyr Val Ala Thr Leu Ser Glu Ala Leu Arg Ile
1 5 10 15

<210> 259
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Group V allergen Phl p 5 epitope

<400> 259

Pro Ala Asn Asp Lys Phe Thr Val Phe Glu Ala Ala Phe Asn Asn
1 5 10 15

<210> 260
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Group V allergen Phl p 5 epitope

<400> 260

Pro Lys Gly Gly Ala Glu Ser Ser Ser Lys Ala Ala Leu Thr Ser
1 5 10 15

<210> 261
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens KIAA1224 protein epitope

<400> 261

Asp Leu Glu Ser Tyr Leu Gln Leu Asn Cys Glu Arg Gly Thr Trp Arg
1 5 10 15

<210> 262
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Lep D 2 precursor epitope

<400> 262

Lys Gly Glu Ala Leu Asp Phe Asn Tyr Gly Met Thr Ile Pro Ala
1 5 10 15

<210> 263
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana lipid transfer protein precursor epitope

<400> 263

Ala Gly Leu Pro Gly Lys Cys Gly Val Asn Ile Pro
1 5 10

<210> 264
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana lipid transfer protein precursor epitope

<400> 264

Ala Lys Gly Ile Ala Gly Leu Asn Pro Asn Leu Ala
1 5 10

<210> 265
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana lipid transfer protein precursor epitope

<400> 265

Cys Gly Val Asn Ile Pro Tyr Lys Ile Ser Pro Ser
1 5 10

<210> 266
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana lipid transfer protein precursor epitope

<400> 266

Cys Lys Gly Val Arg Ala Val Asn Asp Ala Ser Arg
1 5 10

<210> 267
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana lipid transfer protein precursor epitope

<400> 267

Cys Val Leu Tyr Leu Lys Asn Gly Gly Val Leu Pro
1 5 10

<210> 268
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Lipocalin 1 (tear prealbumin) epitope

<400> 268

Lys Pro Val Arg Gly Val Lys Leu Val Gly Arg Asp Pro Lys Asn Asn
1 5 10 15

<210> 269
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mag3 epitope

<400> 269

Glu Phe Asn Thr Glu Phe Thr Ile His Ala Asp Lys Asn Asn Leu
1 5 10 15

<210> 270
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mag3 epitope

<400> 270

Phe Thr Ile His Ala Asp Lys Asn Asn Leu Lys Met His Met Asp
1 5 10 15

<210> 271
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mag3 epitope

<400> 271

Lys Met His Met Asp Phe Pro Asn Val Phe Gln Ala Asp Leu Thr
1 5 10 15

<210> 272
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Apium graveolens Major allergen Api g 1 epitope

<400> 272

Ala Leu Phe Lys Ala Leu Glu Ala Tyr Leu Ile Ala Asn
1 5 10

<210> 273
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Apium graveolens Major allergen Api g 1 epitope

<400> 273

Asp Ala Val Val Pro Glu Glu Asn Ile Lys Tyr Ala
1 5 10

<210> 274
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Apium graveolens Major allergen Api g 1 epitope

<400> 274

Asp Ile Leu Leu Gly Phe Ile Glu Ser Ile Glu Asn
1 5 10

<210> 275
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Apium graveolens Major allergen Api g 1 epitope

<400> 275

Gly Gly Ser Ile Cys Lys Thr Thr Ala Ile Phe His
1 5 10

<210> 276
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Apium graveolens Major allergen Api g 1 epitope

<400> 276

Gly Val Gln Thr His Val Leu Glu Leu Thr Ser Ser
1 5 10

<210> 277
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Major allergen Asp f 2 precursor epitope

<400> 277

Phe Gly Asn Arg Pro Thr Met Glu Ala Val Gly Ala Tyr Asp Val
1 5 10 15

<210> 278
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Major allergen Asp f 2 precursor epitope

<400> 278

Met Glu Ala Val Gly Ala Tyr Asp Val Ile Val Asn Gly Asp Lys
1 5 10 15

<210> 279
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Canis lupus familiaris Major allergen Can f 1 precursor epitope

<400> 279

Ala Leu Glu Asp Phe Arg Glu Phe Ser Arg Ala Lys Gly Leu Asn Gln
1 5 10 15

<210> 280
<211> 16
<212> PRT
<213> Artificial Sequence

<220>

<223> Canis lupus familiaris Major allergen Can f 1 precursor epitope

<400> 280

Asp Gln Glu Val Pro Glu Lys Pro Asp Ser Val Thr Pro Met Ile Leu
1 5 10 15

<210> 281
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana major allergen Cor a 1.0401 epitope

<400> 281

Ala Gly Lys Glu Lys Ala Ala Gly Leu Phe Lys Ala
1 5 10

<210> 282
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana major allergen Cor a 1.0401 epitope

<400> 282

Ala Gly Leu Phe Lys Ala Val Glu Ala Tyr Leu Leu
1 5 10

<210> 283
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana major allergen Cor a 1.0401 epitope

<400> 283

Ala Pro Gln His Phe Thr Ser Ala Glu Asn Leu Glu
1 5 10

<210> 284
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana major allergen Cor a 1.0401 epitope

<400> 284

Ala Arg Leu Phe Lys Ser Phe Val Leu Asp Ala Asp
1 5 10

<210> 285
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana major allergen Cor a 1.0401 epitope

<400> 285

Glu Ile Asp His Ala Asn Phe Lys Tyr Cys Tyr Ser
1 5 10

<210> 286
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Daucus carota Major allergen Dau c 1 epitope

<400> 286

Ala Leu Phe Lys Ala Ile Glu Ala Tyr Leu Ile Ala Asn
1 5 10

<210> 287
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Equus caballus Major allergen Equ c 1 precursor epitope

<400> 287

Asp Gly Tyr Asn Val Phe Arg Ile Ser Glu Phe Glu Asn Asp Glu His
1 5 10 15

<210> 288
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Equus caballus Major allergen Equ c 1 precursor epitope

<400> 288

Asp Lys Asp Arg Pro Phe Gln Leu Phe Glu Phe Tyr Ala Arg Glu Pro
1 5 10 15

<210> 289
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Equus caballus Major allergen Equ c 1 precursor epitope

<400> 289

Asp Leu Thr Lys Ile Asp Arg Cys Phe Gln Leu Arg Gly Asn Gly Val
1 5 10 15

<210> 290
<211> 16

<212> PRT
<213> Artificial Sequence

<220>
<223> Equus caballus Major allergen Equ c 1 precursor epitope

<400> 290

Asp Arg Pro Phe Gln Leu Phe Glu Phe Tyr Ala Arg Glu Pro Asp Val
1 5 10 15

<210> 291
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Equus caballus Major allergen Equ c 1 precursor epitope

<400> 291

Asp Val Ser Pro Glu Ile Lys Glu Glu Phe Val Lys Ile Val Gln Lys
1 5 10 15

<210> 292
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus major allergen I epitope

<400> 292

Glu Asn Ala Arg Ile Leu Lys Asn Cys Val Asp Ala Lys Met Thr Glu
1 5 10 15

Glu

<210> 293
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus major allergen I epitope

<400> 293

Arg Asp Val Asp Leu Phe Leu Thr Gly Thr Pro Asp Glu Tyr Val Glu
1 5 10 15

Gln

<210> 294
<211> 17
<212> PRT
<213> Artificial Sequence

<220>

<223> Felis catus major allergen I epitope

<400> 294

Thr Gly Thr Pro Asp Glu Tyr Val Glu Gln Val Ala Gln Tyr Lys Ala
1 5 10 15

Leu

<210> 295
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 1 precursor epitope

<400> 295

Asp Val Asp Leu Phe Leu Thr Gly Thr Pro Asp Glu Tyr Val Glu Gln
1 5 10 15

Val

<210> 296
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 1 precursor epitope

<400> 296

Glu Ile Cys Pro Ala Val Lys Arg Asp Val Asp Leu Phe Leu Thr Gly
1 5 10 15

Thr

<210> 297
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 1 precursor epitope

<400> 297

Glu Gln Val Ala Gln Tyr Lys Ala Leu Pro Val Val Leu Glu Asn Ala
1 5 10 15

<210> 298
<211> 17
<212> PRT

<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 1 precursor epitope

<400> 298

Lys Ala Leu Pro Val Val Leu Glu Asn Ala Arg Ile Leu Lys Asn Cys
1 5 10 15

Val

<210> 299
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 1 precursor epitope

<400> 299

Leu Phe Leu Thr Gly Thr Pro Asp Glu Tyr Val Glu Gln Val Ala Gln
1 5 10 15

Tyr

<210> 300
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus major allergen I, polypeptide chain 1 epitope

<400> 300

Lys Glu Asn Ala Leu Ser Leu Leu Asp Lys Ile Tyr Thr Ser Pro Leu
1 5 10 15

<210> 301
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus major allergen I, polypeptide chain 1 epitope

<400> 301

Lys Met Thr Glu Glu Asp Lys Glu Asn Ala Leu Ser Leu Leu Asp Lys
1 5 10 15

<210> 302
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Malus x domestica Major allergen Mal d 1 epitope

<400> 302

Gly Leu Phe Lys Leu Ile Glu Ser Tyr Leu Lys Asp His Pro Asp
1 5 10 15

<210> 303
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus avium Major allergen Pru av 1 epitope

<400> 303

Asn Leu Phe Lys Leu Ile Glu Thr Tyr Leu Lys Gly His Pro Asp
1 5 10 15

<210> 304
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Major latex allergen Hev b 5 epitope

<400> 304

Ala Ala Pro Ala Glu Gly Glu Lys Pro Ala Glu Glu Glu Lys Pro Ile
1 5 10 15

Thr Glu Ala Ala
20

<210> 305
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Major latex allergen Hev b 5 epitope

<400> 305

Ala Glu Glu Glu Lys Pro Ile Thr Glu Ala Ala Glu Thr Ala Thr Thr
1 5 10 15

Glu Val Pro Val
20

<210> 306
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Major latex allergen Hev b 5 epitope

<400> 306

Ala Pro Ala Glu Pro Glu Ala Pro Ala Pro Glu Thr Glu Lys Ala Glu
1 5 10 15

Glu Val Glu Lys
20

<210> 307
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Major latex allergen Hev b 5 epitope

<400> 307

Ala Pro Glu Ala Asp Gln Thr Thr Pro Glu Glu Lys Pro Ala Glu Pro
1 5 10 15

Glu Pro Val Ala
20

<210> 308
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Major latex allergen Hev b 5 epitope

<400> 308

Ala Ser Glu Gln Glu Thr Ala Asp Ala Thr Pro Glu Lys Glu Glu Pro
1 5 10 15

Thr Ala Ala Pro
20

<210> 309
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Major mite fecal allergen Der p 1 epitope

<400> 309

Tyr Ala Tyr Val Ala Arg Glu Gln Ser Cys Arg
1 5 10

<210> 310
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Major mite fecal allergen Der p 1 epitope

<400> 310

Ala Leu Ala Gln Thr His Thr Ala Ile Ala Val Ile Ile Gly Ile Lys
1 5 10 15

Asp Leu Asp

<210> 311
<211> 35
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Major pollen allergen epitope

<400> 311

Glu Asp Ile Pro Gln Pro Pro Val Ser Gln Phe His Ile Gln Gly Gln
1 5 10 15

Val Tyr Cys Asp Thr Cys Arg Ala Gly Phe Ile Thr Glu Leu Ser Glu
20 25 30

Phe Ile Pro
35

<210> 312
<211> 31
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Major pollen allergen epitope

<400> 312

Gly Ala Ser Leu Arg Leu Gln Cys Lys Asp Lys Glu Asn Gly Asp Val
1 5 10 15

Thr Phe Thr Glu Val Gly Tyr Thr Arg Ala Glu Gly Leu Tyr Ser
20 25 30

<210> 313
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Major pollen allergen epitope

<400> 313

Gly Thr Thr Arg Thr Val Asn Pro Leu Gly Phe Phe Lys Lys Glu Ala
1 5 10 15

Leu Pro Lys Cys Ala Gln Val Tyr Asn Lys Leu Gly Met Tyr Pro Pro
20 25 30

Asn Met

<210> 314
<211> 53
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Major pollen allergen epitope

<400> 314

Leu Val Glu Arg Asp His Lys Asn Glu Phe Cys Glu Ile Thr Leu Ile
1 5 10 15

Ser Ser Gly Arg Lys Asp Cys Asn Glu Ile Pro Thr Glu Gly Trp Ala
20 25 30

Lys Pro Ser Leu Lys Phe Lys Leu Asn Thr Val Asn Gly Thr Thr Arg
35 40 45

Thr Val Asn Pro Leu
50

<210> 315
<211> 33
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Major pollen allergen epitope

<400> 315

Met Leu Val Glu Arg Asp His Lys Asn Glu Phe Cys Glu Ile Thr Leu
1 5 10 15

Ile Ser Ser Gly Arg Lys Asp Cys Asn Glu Ile Pro Thr Glu Gly Trp
20 25 30

Ala

<210> 316
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Artemisia vulgaris Major pollen allergen Art v 1 precursor epitope

<400> 316

Ala Gly Gly Ser Pro Ser Pro Pro Ala Asp Gly Gly
1 5 10

<210> 317
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Artemisia vulgaris Major pollen allergen Art v 1 precursor epitope

<400> 317

Ala Gly Ser Lys Leu Cys Glu Lys Thr Ser Lys Thr
1 5 10

<210> 318
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Artemisia vulgaris Major pollen allergen Art v 1 precursor epitope

<400> 318

Cys Asp Lys Lys Cys Ile Glu Trp Glu Lys Ala Gln
1 5 10

<210> 319
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Artemisia vulgaris Major pollen allergen Art v 1 precursor epitope

<400> 319

Asp Gly Gly Ser Pro Pro Pro Pro Ala Asp Gly Gly
1 5 10

<210> 320
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Artemisia vulgaris Major pollen allergen Art v 1 precursor epitope

<400> 320

Glu Lys Thr Ser Lys Thr Tyr Ser Gly Lys Cys Asp
1 5 10

<210> 321
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 321

Ala Ala Arg Leu Phe Lys Ala Phe Ile Leu Asp Gly
1 5 10

<210> 322
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 322

Ala Ala Arg Leu Phe Lys Ala Phe Ile Leu Asp Gly Asp Asn Leu
1 5 10 15

<210> 323
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 323

Ala Glu Gln Val Lys Ala Ser Lys Glu Met Gly Glu
1 5 10

<210> 324
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 324

Ala Phe Ile Leu Asp Gly Asp Asn Leu Phe Pro Lys Val Ala Pro Gln
1 5 10 15

Ala Ile Ser Ser Val
20

<210> 325
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 325

Ala Ile Ser Ser Val Glu Asn Ile Glu Gly Asn Gly
1 5 10

<210> 326
<211> 15

<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 326

Glu Thr Leu Leu Arg Ala Val Glu Ser Tyr Leu Leu Ala His Ser
1 5 10 15

<210> 327
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-F/I epitope

<400> 327

Gly Glu Thr Leu Leu Arg Ala Val Glu Ser Tyr Leu Leu Ala His Ser
1 5 10 15

<210> 328
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Major pollen allergen Cha o 1 precursor epitope

<400> 328

Ala Asn Asn Asn Tyr Asp Pro Trp Ser Ile Tyr Ala Ile Gly Gly Ser
1 5 10 15

Ser Asn Pro Thr
20

<210> 329
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Major pollen allergen Cha o 1 precursor epitope

<400> 329

Ala Ser Thr Gly Val Thr Ile Ser Asn Asn His Phe Phe Asn His His
1 5 10 15

Lys Val Met Leu
20

<210> 330
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Major pollen allergen Cha o 1 precursor epitope

<400> 330

Cys Ala Asn Trp Val Trp Arg Ser Thr Gln Asp Ser Phe Asn Asn Gly
1 5 10 15

Ala Tyr Phe Val
20

<210> 331
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Major pollen allergen Cha o 1 precursor epitope

<400> 331

Asp Ala Ile Thr Met Arg Asn Val Thr Asp Val Trp Ile Asp His Asn
1 5 10 15

Ser Leu Ser Asp
20

<210> 332
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Major pollen allergen Cha o 1 precursor epitope

<400> 332

Asp Ala Asn Trp Asp Gln Asn Arg Met Lys Leu Ala Asp Cys Ala Val
1 5 10 15

Gly Phe Gly Ser
20

<210> 333
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon Major pollen allergen Cyn d 1 epitope

<400> 333

Ala Ile Gly Asp Lys Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asn
1 5 10 15

Lys Trp Leu Glu

20

<210> 334
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon Major pollen allergen Cyn d 1 epitope

<400> 334

Cys Tyr Glu Ile Lys Cys Lys Glu Pro Val Glu Cys Ser Gly Glu Pro
1 5 10 15

Val Leu Val Lys
20

<210> 335
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon Major pollen allergen Cyn d 1 epitope

<400> 335

Asp His Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys Pro Pro Phe
1 5 10 15

Asp Gly Met Thr
20

<210> 336
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon Major pollen allergen Cyn d 1 epitope

<400> 336

Glu Gly Gly Ala His Leu Val Gln Asp Asp Val Ile Pro Ala Asn Trp
1 5 10 15

Lys Pro Asp Thr
20

<210> 337
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon Major pollen allergen Cyn d 1 epitope

<400> 337

Phe Lys Asp Gly Leu Gly Cys Gly Ala Cys Tyr Glu Ile Lys Cys Lys

1 5 10 15

Glu Pro Val Glu
20

<210> 338
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Major pollen allergen Phl p 4 precursor epitope

<400> 338

Phe Ala Glu Tyr Lys Ser Asp Tyr Val Tyr Gln Pro Phe Pro Lys
1 5 10 15

<210> 339
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Major pollen allergen Phl p 4 precursor epitope

<400> 339

Met Leu Leu Arg Lys Tyr Gly Ile Ala Ala Glu Asn Val Ile Asp
1 5 10 15

<210> 340
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Major pollen allergen Phl p 4 precursor epitope

<400> 340

Asn Ser Phe Lys Pro Phe Ala Glu Tyr Lys Ser Asp Tyr Val Tyr
1 5 10 15

<210> 341
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Rattus norvegicus Major urinary protein precursor epitope

<400> 341

Ala Ser Asn Lys Arg Glu Lys Ile Glu Glu Asn Gly Ser Met Arg Val
1 5 10 15

Phe Met Gln His
20

<210> 342

<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Rattus norvegicus Major urinary protein precursor epitope

<400> 342

Asp Ile Lys Glu Lys Phe Ala Lys Leu Cys Glu Ala His Gly Ile Thr
1 5 10 15

Arg Asp Asn Ile
20

<210> 343
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Rattus norvegicus Major urinary protein precursor epitope

<400> 343

Glu Glu Ala Ser Ser Thr Arg Gly Asn Leu Asp Val Ala Lys Leu Asn
1 5 10 15

Gly Asp Trp Phe
20

<210> 344
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Rattus norvegicus Major urinary protein precursor epitope

<400> 344

Glu Glu Asn Gly Ser Met Arg Val Phe Met Gln His Ile Asp Val Leu
1 5 10 15

Glu Asn Ser Leu
20

<210> 345
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Rattus norvegicus Major urinary protein precursor epitope

<400> 345

Glu Asn Ser Leu Gly Phe Lys Phe Arg Ile Lys Glu Asn Gly Glu Cys
1 5 10 15

Arg Glu Leu Tyr

20

<210> 346
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mite group 2 allergen Der f 2 precursor epitope

<400> 346

Asp Ile Lys Tyr Thr Trp Asn Val Pro Lys Ile Ala Pro Lys Ser Glu
1 5 10 15

Asn Val Val Val Thr
20

<210> 347
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mite group 2 allergen Der f 2 precursor epitope

<400> 347

Asp Asn Gly Val Leu Ala Cys Ala Ile Ala Thr His Gly Lys Ile Arg
1 5 10 15

Asp

<210> 348
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mite group 2 allergen Der f 2 precursor epitope

<400> 348

Glu Ala Leu Phe Asp Ala Asn Gln Asn Thr Lys Thr Ala Lys Ile Glu
1 5 10 15

Ile Lys Ala Ser Leu
20

<210> 349
<211> 45
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mite group 2 allergen Der f 2 precursor epitope

<400> 349

Gln Tyr Asp Ile Lys Tyr Thr Trp Asn Val Pro Lys Ile Ala Pro Lys
1 5 10 15

Ser Glu Asn Val Val Val Thr Val Lys Leu Ile Gly Asp Asn Gly Val
20 25 30

Leu Ala Cys Ala Ile Ala Thr His Gly Lys Ile Arg Asp
35 40 45

<210> 350
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mite group 2 allergen Der f 2 precursor epitope

<400> 350

Thr Lys Thr Ala Lys Ile Glu Ile Lys Ala Ser Leu Asp Gly Leu Glu
1 5 10 15

Ile Asp Val

<210> 351
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 epitope

<400> 351

Ala Ser Ile Asp Gly Leu Gly Val Asp Val Pro Gly Ile Asp
1 5 10

<210> 352
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 epitope

<400> 352

Phe Glu Ala Val Gln Asn Thr Lys Thr Ala Lys Ile Glu Ile Lys
1 5 10 15

<210> 353
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 epitope

<400> 353

Arg Gly Lys Pro Pro Gln Leu Glu Ala Val Phe Glu Ala Val Gln Asn
1 5 10 15

Thr

<210> 354
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 354

Cys His Gly Ser Glu Pro Cys Ile Ile His Arg Gly Lys Pro Phe
1 5 10 15

<210> 355
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 355

Cys Pro Leu Val Lys Gly Gln Gln Tyr Asp Ile Lys Tyr Thr Trp Asn
1 5 10 15

Val Pro Lys Ile Ala Pro Lys Ser Glu Asn Val
20 25

<210> 356
<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 356

Asp Ile Lys Tyr Thr Trp Asn Val Pro Lys Ile Ala Pro Lys Ser Glu
1 5 10 15

Asn Val Val Val Thr Val Lys Val Met Gly
20 25

<210> 357
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 357

Asp Gln Val Asp Val Lys Asp Cys Ala Asn His Glu Ile Lys Lys
1 5 10 15

<210> 358
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 358

Asp Gln Val Asp Val Lys Asp Cys Ala Asn His Glu Ile Lys Lys Val
1 5 10 15

Leu Val Pro Gly
20

<210> 359
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 359

Cys His Gly Ser Glu Pro Cys Ile Ile His Arg Gly Lys Pro Phe
1 5 10 15

<210> 360
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Mite group 2 allergen Lep d 2 precursor epitope

<400> 360

Asp His Gly Val Met Ala Cys Gly Thr Val His Gly Gln Val Glu
1 5 10 15

<210> 361
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Mite group 2 allergen Lep d 2 precursor epitope

<400> 361

Gly Cys Lys Phe Ile Lys Cys Pro Val Lys Lys Gly Glu Ala Leu
1 5 10 15

<210> 362
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Mite group 2 allergen Lep d 2 precursor epitope

<400> 362

Gly Glu Lys Met Thr Leu Glu Ala Lys Phe Ala Ala Asn Gln Asp
1 5 10 15

<210> 363
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Mite group 2 allergen Lep d 2 precursor epitope

<400> 363

Gly Glu Val Thr Glu Leu Asp Ile Thr Gly Cys Ser Gly Asp Thr
1 5 10 15

<210> 364
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Mite group 2 allergen Lep d 2 precursor epitope

<400> 364

Gly Lys Met Thr Phe Lys Asp Cys Gly His Gly Glu Val Thr Glu
1 5 10 15

<210> 365
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Neurofilament heavy polypeptide (NF-H) (Neurofilament triplet H protein) (200 kDa neurofilament protein) epitope

<400> 365

Tyr Gln Glu Ala Ile Gln Gln Leu Asp Ala Glu Leu Arg Asn Thr Lys
1 5 10 15

<210> 366
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica Non-specific lipid-transfer protein 1 epitope

<400> 366

Ala Ala Ala Leu Pro Gly Lys Cys Gly Val
1 5 10

<210> 367
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica Non-specific lipid-transfer protein 1 epitope

<400> 367

Ala Cys Cys Asn Gly Ile Arg Asn Val Asn
1 5 10

<210> 368
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica Non-specific lipid-transfer protein 1 epitope

<400> 368

Ala Pro Cys Ile Pro Tyr Val Arg Gly Gly
1 5 10

<210> 369
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica Non-specific lipid-transfer protein 1 epitope

<400> 369

Ile Arg Asn Val Asn Asn Leu Ala Arg Thr
1 5 10

<210> 370
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica Non-specific lipid-transfer protein 1 epitope

<400> 370

Ile Ser Ala Ser Thr Asn Cys Ala Thr Val Lys
1 5 10

<210> 371
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica Non-specific lipid-transfer protein 1 epitope

<400> 371

Asn Leu Ala Arg Thr Thr Pro Asp Arg Gln
1 5 10

<210> 372
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovalbumin epitope

<400> 372

Cys Phe Asp Val Phe Lys Glu Leu Lys Val
1 5 10

<210> 373
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovalbumin epitope

<400> 373

Gly Ser Ile Gly Ala Ala Ser Met Glu Phe
1 5 10

<210> 374
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovalbumin epitope

<400> 374

Ile Gly Leu Phe Arg Val Ala Ser Met Ala Ser Glu Lys Met Lys Ile
1 5 10 15

Leu Glu

<210> 375
<211> 18

<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovalbumin epitope

<400> 375

Ile Lys His Ile Ala Thr Asn Ala Val Leu Phe Phe Gly Arg Cys Val
1 5 10 15

Ser Pro

<210> 376
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovalbumin epitope

<400> 376

Ile Met Ser Ala Leu Ala Met Val Tyr Leu Gly Ala Lys
1 5 10

<210> 377
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid precursor epitope

<400> 377

Ala Glu Val Asp Cys Ser Arg Phe Pro Asn Ala Thr Asp Lys
1 5 10

<210> 378
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid precursor epitope

<400> 378

Ala Thr Asp Lys Glu Gly Lys Asp Val Leu Val Cys Asn Lys
1 5 10

<210> 379
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid precursor epitope

<400> 379

Ala Val Val Glu Ser Asn Gly Thr Leu Thr Leu Ser His Phe Gly Lys
1 5 10 15

Cys

<210> 380
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid precursor epitope

<400> 380

Cys Leu Leu Cys Ala Tyr Ser Ile Glu Phe Gly Thr Asn Ile Ser Lys
1 5 10 15

<210> 381
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid precursor epitope

<400> 381

Asp Asn Glu Cys Leu Leu Cys Ala His Lys Val Glu Gln Gly Ala Ser
1 5 10 15

Val Asp Lys Arg
20

<210> 382
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Musa acuminata pectate lyase epitope

<400> 382

Gly His Ser Asp Glu Leu Thr Ser Asp Lys Ser Met Gln Val Thr Ile
1 5 10 15

<210> 383
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Zinnia violacea Pectate lyase precursor epitope

<400> 383

Gly His Ser Asp Ser Tyr Thr Gln Asp Lys Asn Met Gln Val Thr Ile
1 5 10 15

<210> 384
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Peptidase 1 precursor (Major mite fecal allergen Der f 1) (Allergen Der f I) epitope

<400> 384

Asp Gly Arg Thr Ile Ile Gln His Asp Asn Gly Tyr Gln Pro Asn Tyr
1 5 10 15

His Ala Val Asn Ile
20

<210> 385
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Peptidase 1 precursor (Major mite fecal allergen Der f 1) (Allergen Der f I) epitope

<400> 385

Asp Leu Arg Ser Leu Arg Thr Val Thr Pro Ile Arg Met Gln Gly Gly
1 5 10 15

Cys Gly Ser

<210> 386
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Peptidase 1 precursor (Major mite fecal allergen Der f 1) (Allergen Der f I) epitope

<400> 386

Gly Cys Gly Ser Cys Trp Ala Phe Ser Gly Val Ala Ala Thr Glu Ser
1 5 10 15

Ala Tyr Leu

<210> 387
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Peptidase 1 precursor (Major mite fecal allergen Der f 1) (Allergen Der f I) epitope

<400> 387

Ile Arg Glu Ala Leu Thr Gln Thr His Thr Ala Ile Ala Val Ile Ile
1 5 10 15

Gly Ile Lys Asp Leu
20

<210> 388
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Peptidase 1 precursor (Major mite fecal allergen Der f 1) (Allergen Der f I) epitope

<400> 388

Ile Arg Met Gln Gly Gly Cys Gly Ser Cys Trp Ala Phe Ser Gly Val
1 5 10 15

Ala Ala Thr

<210> 389
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Euroglyphus maynei Peptidase 1 precursor (Mite group 1 allergen Eur m 1) (Allergen Eur m I) epitope

<400> 389

Phe Arg His Tyr Asp Gly Arg Thr Ile Met Gln His Asp Asn Gly Tyr
1 5 10 15

Gln Pro Asn

<210> 390
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Euroglyphus maynei Peptidase 1 precursor (Mite group 1 allergen Eur m 1) (Allergen Eur m I) epitope

<400> 390

Gly Arg Thr Ile Met Gln His Asp Asn Gly Tyr Gln Pro Asn Tyr His
1 5 10 15

Ala Val Asn

<210> 391
<211> 19
<212> PRT

<213> Artificial Sequence

<220>
<223> Euroglyphus maynei Peptidase 1 precursor (Mite group 1 allergen Eur m 1) (Allergen Eur m I) epitope

<400> 391

His Ala Val Asn Ile Val Gly Tyr Gly Asn Thr Gln Gly Val Asp Tyr
1 5 10 15

Trp Ile Val

<210> 392
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Euroglyphus maynei Peptidase 1 precursor (Mite group 1 allergen Eur m 1) (Allergen Eur m I) epitope

<400> 392

Asn Lys Ile Arg Gln Ala Leu Thr Gln Thr His Thr Ala Val Ala Val
1 5 10 15

Ile Ile Gly

<210> 393
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Euroglyphus maynei Peptidase 1 precursor (Mite group 1 allergen Eur m 1) (Allergen Eur m I) epitope

<400> 393

Pro Tyr Val Ala Arg Glu Gln Ser Cys His Arg Pro Asn Ala Gln Arg
1 5 10 15

Tyr Gly Leu

<210> 394
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Phl p 3 allergen epitope

<400> 394

Ala Val Gln Val Thr Phe Thr Val Gln Lys Gly Ser Asp Pro Lys
1 5 10 15

<210> 395
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Phl p 3 allergen epitope

<400> 395

Glu Glu Trp Glu Pro Leu Thr Lys Lys Gly Asn Val Trp Glu Val
1 5 10 15

<210> 396
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Phl p 3 allergen epitope

<400> 396

Phe Thr Val Gln Lys Gly Ser Asp Pro Lys Lys Leu Val Leu Asp
1 5 10 15

<210> 397
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Phl p 3 allergen epitope

<400> 397

Phe Thr Val Gln Lys Gly Ser Asp Pro Lys Lys Leu Val Leu Asn
1 5 10 15

<210> 398
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Phl p 3 allergen epitope

<400> 398

Gly Ser Asp Pro Lys Lys Leu Val Leu Asp Ile Lys Tyr Thr Arg
1 5 10 15

<210> 399
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Apis mellifera Phospholipase A2 precursor epitope

<400> 399

Cys Asp Cys Asp Asp Lys Phe Tyr Asp Cys Leu Lys Asn Ser Ala
1 5 10 15

<210> 400
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Apis mellifera Phospholipase A2 precursor epitope

<400> 400

Cys Leu His Tyr Thr Val Asp Lys Ser Lys Pro Lys
1 5 10

<210> 401
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Apis mellifera Phospholipase A2 precursor epitope

<400> 401

Cys Arg Thr His Asp Met Cys Pro Asp Val Met Ser Ala Gly Glu
1 5 10 15

<210> 402
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Apis mellifera Phospholipase A2 precursor epitope

<400> 402

Asp Thr Ile Ser Ser Tyr Phe Val Gly Lys Met Tyr Phe Asn Leu Ile
1 5 10 15

Asp Thr

<210> 403
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Apis mellifera Phospholipase A2 precursor epitope

<400> 403

Glu Arg Thr Glu Gly Arg Cys Leu His Tyr Thr Val Asp Lys Ser Lys
1 5 10 15

Pro Lys

<210> 404
<211> 16

<212> PRT
<213> Artificial Sequence

<220>
<223> Spiroplasma citri plectrovirus spv1-r8a2b orf 14 transmembrane protein epitope

<400> 404

His Val Ile Glu Val Gln Gln Ile Asn Ser Glu Arg Ser Trp Phe Phe
1 5 10 15

<210> 405
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne pollen allergen epitope

<400> 405

Cys Gly Tyr Lys Asp Val Asp Lys Ala Pro Phe Asn Gly Met Thr Gly
1 5 10 15

Cys Gly Asn Thr
20

<210> 406
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne pollen allergen epitope

<400> 406

Gly Ala Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val
1 5 10 15

Asp Lys Ala Pro
20

<210> 407
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne pollen allergen epitope

<400> 407

Ser Glu Val Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser
1 5 10 15

Tyr Ser Ala Lys
20

<210> 408

<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne pollen allergen epitope

<400> 408

Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys Tyr
1 5 10 15

Val Asp Gly Asp
20

<210> 409
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne pollen allergen epitope

<400> 409

Tyr Pro Asp Asp Thr Lys Pro Thr Phe His Val Glu Lys Gly Ser Asn
1 5 10 15

Pro Asn Tyr Leu
20

<210> 410
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia Pollen allergen Amb a 1.1 precursor epitope

<400> 410

Gly Ala Gly Asp Glu Asn Ile Glu Asp Arg Gly Met Leu Ala Thr Val
1 5 10 15

<210> 411
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia Pollen allergen Amb a 1.1 precursor epitope

<400> 411

Gly Ala Gly Asp Glu Asn Ile Glu Asp Arg Gly Met Leu Ala Thr Val
1 5 10 15

<210> 412
<211> 16
<212> PRT

<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia Pollen allergen Amb a 2 precursor epitope

<400> 412

Gly Ala Ser Asp Thr His Phe Gln Asp Leu Lys Met His Val Thr Leu
1 5 10 15

<210> 413
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia Pollen allergen Amb a 2 precursor epitope

<400> 413

Gly Ala Ser Asp Thr His Phe Gln Asp Leu Lys Met His Val Thr Leu
1 5 10 15

<210> 414
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 3 epitope

<400> 414

Glu Glu Ala Tyr His Ala Cys Asp Ile Lys Asp
1 5 10

<210> 415
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 3 epitope

<400> 415

Gly Lys Val Tyr Leu Val Gly Gly Pro Glu Leu Gly Gly Trp Lys
1 5 10 15

<210> 416
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 3 epitope

<400> 416

Leu Gly Gly Trp Lys Leu Gln Ser Asp Pro Arg Ala Tyr Ala Leu
1 5 10 15

<210> 417
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 3 epitope

<400> 417

Pro Gly Gly Pro Asp Arg Phe Thr Leu Leu Thr Pro Gly Ser His
1 5 10 15

<210> 418
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 418

Ala Tyr Cys Cys Ser Asp Pro Gly Arg Tyr Cys Pro Trp Gln Val
1 5 10 15

<210> 419
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 419

Cys Gly Glu Lys Arg Ala Tyr Cys Cys Ser Asp Pro Gly Arg Tyr Cys
1 5 10 15

Pro Trp Gln Val
20

<210> 420
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 420

Asp Pro Gly Arg Tyr Cys Pro Trp Gln Val Val Cys Tyr Glu Ser Ser
1 5 10 15

Glu

<210> 421
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 421

Asp Pro Gly Arg Tyr Cys Pro Trp Gln Val Val Cys Tyr Glu Ser Ser
1 5 10 15

Glu Ile Cys Ser
20

<210> 422
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 422

Gly Asn Val Cys Gly Glu Lys Arg Ala Tyr Cys Cys Ser Asp Pro
1 5 10 15

<210> 423
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 423

Leu Val Pro Cys Ala Trp Ala Gly Asn Val Cys Gly Glu Lys Arg
1 5 10 15

<210> 424
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 424

Leu Val Pro Cys Ala Trp Ala Gly Asn Val Cys Gly Glu Lys Arg Ala
1 5 10 15

Tyr Cys Cys Ser
20

<210> 425
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 5 epitope

<400> 425

Val Cys Tyr Glu Ser Ser Glu Ile Cys Ser Lys Lys Cys Gly Lys
1 5 10 15

<210> 426
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia trifida Pollen allergen Amb t 5 precursor epitope

<400> 426

Cys Gly Lys Val Gly Lys Tyr Cys Cys Ser Pro Ile Gly Lys Tyr Cys
1 5 10 15

Val Cys Tyr Asp
20

<210> 427
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia trifida Pollen allergen Amb t 5 precursor epitope

<400> 427

Asp Asp Gly Leu Cys Tyr Glu Gly Thr Asn Cys Gly Lys Val Gly Lys
1 5 10 15

Tyr Cys Cys Ser
20

<210> 428
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia trifida Pollen allergen Amb t 5 precursor epitope

<400> 428

Gly Lys Tyr Cys Val Cys Tyr Asp Ser Lys Ala Ile
1 5 10

<210> 429

<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia trifida Pollen allergen Amb t 5 precursor epitope

<400> 429

Pro Ile Gly Lys Tyr Cys Val Cys Tyr Asp Ser Lys Ala Ile
1 5 10

<210> 430
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia trifida Pollen allergen Amb t 5 precursor epitope

<400> 430

Pro Ile Gly Lys Tyr Cys Val Cys Tyr Asp Ser Lys Ala Ile Cys Asn
1 5 10 15

Lys Asn Cys Thr
20

<210> 431
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia trifida Pollen allergen Amb t 5 precursor epitope

<400> 431

Val Cys Tyr Asp Ser Lys Ala Ile Cys Asn Lys Asn Cys Thr
1 5 10

<210> 432
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula pollen allergen Bet v 1 epitope

<400> 432

His Glu Val Lys Ala Glu Gln Val Lys Ala Thr Lys Glu Met Gly
1 5 10 15

<210> 433
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Poa pratensis Pollen allergen KBG 60 precursor epitope

<400> 433

Ala Ala Asn Lys Tyr Lys Thr Phe Val Ala Thr Phe Gly Ala Ala Ser
1 5 10 15

Asn Lys Ala Phe
20

<210> 434
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Poa pratensis Pollen allergen KBG 60 precursor epitope

<400> 434

Ala Ala Pro Ala Asn Asp Lys Phe Thr Val Phe Glu Ala Ala Phe Asn
1 5 10 15

Asp Ala Ile Lys
20

<210> 435
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Poa pratensis Pollen allergen KBG 60 precursor epitope

<400> 435

Ala Ala Val Asp Ser Ser Lys Ala Ala Leu Thr Ser Lys Leu Asp Ala
1 5 10 15

Ala Tyr Lys Leu
20

<210> 436
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Poa pratensis Pollen allergen KBG 60 precursor epitope

<400> 436

Ala Glu Glu Val Lys Ala Thr Pro Ala Gly Glu Leu Gln Val Ile Asp
1 5 10 15

Lys Val Asp Ala
20

<210> 437
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Poa pratensis Pollen allergen KBG 60 precursor epitope

<400> 437

Ala Phe Lys Val Ala Ala Thr Ala Ala Asn Ala Ala Pro Ala Asn Asp
1 5 10 15

Lys Phe Thr Val
20

<210> 438
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 1 precursor epitope

<400> 438

Ala Phe Gly Ser Met Ala Lys Lys Gly Glu Glu Gln Asn Val Arg Ser
1 5 10 15

Ala Gly Glu Leu
20

<210> 439
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 1 precursor epitope

<400> 439

Ala Gly Glu Leu Glu Leu Gln Phe Arg Arg Val Lys Cys Lys Tyr Pro
1 5 10 15

Asp Asp Thr Lys
20

<210> 440
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 1 precursor epitope

<400> 440

Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly Pro Lys Asp
1 5 10 15

Asn Gly Gly Ala
20

<210> 441

<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 1 precursor epitope

<400> 441

Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ser Met Ala
1 5 10 15

Lys Lys Gly Glu
20

<210> 442
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 1 precursor epitope

<400> 442

Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala
1 5 10

<210> 443
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p VA precursor epitope

<400> 443

Ala Ala Leu Thr Lys Ala Ile Thr Ala Met Thr Gln Ala Gln Lys Ala
1 5 10 15

Gly Lys Pro Ala
20

<210> 444
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p VA precursor epitope

<400> 444

Ala Ala Asn Ala Ala Pro Thr Asn Asp Lys Phe Thr Val Phe Glu Ser
1 5 10 15

Ala Phe Asn Lys
20

<210> 445

<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p VA precursor epitope

<400> 445

Ala Asp Lys Phe Lys Ile Phe Glu Ala Ala Phe Ser Glu Ser Ser Lys
1 5 10 15

Gly Leu Leu Ala
20

<210> 446
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p VA precursor epitope

<400> 446

Ala Phe Ser Glu Ser Ser Lys Gly Leu Leu Ala Thr Ser Ala Ala Lys
1 5 10 15

Ala Pro Gly Leu
20

<210> 447
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p VA precursor epitope

<400> 447

Ala Tyr Ala Ala Thr Val Ala Ala Ala Pro Glu Val Lys Tyr Ala Val
1 5 10 15

Phe Glu Ala Ala
20

<210> 448
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 epitope

<400> 448

Ala Cys Ser Gly Glu Pro Val Val Val His Ile Thr
1 5 10

<210> 449

<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 epitope

<400> 449

Ala Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp
1 5 10

<210> 450
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 epitope

<400> 450

Ala Gly Glu Leu Glu Leu Gln Phe Arg Arg Val Lys
1 5 10

<210> 451
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 epitope

<400> 451

Asp Lys Trp Ile Glu Leu Lys Glu Ser Trp Gly Ala
1 5 10

<210> 452
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 epitope

<400> 452

Asp Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly
1 5 10

<210> 453
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 453

Phe Glu Ile Lys Cys Thr Lys Pro Glu Ala Cys Ser
1 5 10

<210> 454
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 454

Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ala
1 5 10

<210> 455
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 455

Glu Leu Lys Glu Ser Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro
1 5 10 15

<210> 456
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 456

Glu Pro Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe
1 5 10 15

<210> 457
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 457

Phe Glu Ile Lys Cys Thr Lys Pro Glu Ala Cys Ser Gly Glu Pro
1 5 10 15

<210> 458
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 458

Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly

1 5 10 15

<210> 459
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 11 epitope

<400> 459

Arg Tyr Ala Asn Pro Ile Ala Phe Phe Arg Lys Glu Pro Leu Lys
1 5 10 15

<210> 460
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 2 epitope

<400> 460

Glu His Gly Ser Asp Glu Trp Val Ala Met Thr Lys Gly Glu Gly
1 5 10 15

<210> 461
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 2 epitope

<400> 461

Glu Trp Val Ala Met Thr Lys Gly Glu Gly Gly Val Trp Thr Phe
1 5 10 15

<210> 462
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 2 epitope

<400> 462

Gly Val Trp Thr Phe Asp Ser Glu Glu Pro Leu Gln Gly Pro Phe
1 5 10 15

<210> 463
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 2 epitope

<400> 463

Lys Asn Val Phe Asp Asp Val Val Pro Glu Lys Tyr Thr Ile Gly
1 5 10 15

<210> 464
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 2 epitope

<400> 464

Leu Gln Gly Pro Phe Asn Phe Arg Phe Leu Thr Glu Lys Gly Met
1 5 10 15

<210> 465
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 4 epitope

<400> 465

Phe Lys Pro Phe Ala Glu Tyr Lys Ser Asp Tyr Val Tyr Glu Pro
1 5 10 15

<210> 466
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 4 epitope

<400> 466

Phe Pro Lys Glu Val Trp Glu Gln Ile Phe Ser Thr Trp Leu Leu
1 5 10 15

<210> 467
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 4 epitope

<400> 467

Phe Val His Leu Gly His Arg Asp Asn Ile Glu Asp Asp Leu Leu
1 5 10 15

<210> 468
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 4 epitope

<400> 468

Gly Ile Val Val Ala Trp Lys Val Arg Leu Leu Pro Val Pro Pro
1 5 10 15

<210> 469
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 4 epitope

<400> 469

Asn Arg Asn Asn Thr Phe Lys Pro Phe Ala Glu Tyr Lys Ser Asp
1 5 10 15

<210> 470
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5a epitope

<400> 470

Glu Val Lys Tyr Thr Val Phe Glu Thr Ala Leu Lys
1 5 10

<210> 471
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5a epitope

<400> 471

Asn Ala Gly Phe Lys Ala Ala Leu Ala Gly Ala Gly Val Gln Pro Ala
1 5 10 15

Asp Lys Tyr

<210> 472
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5b precursor epitope

<400> 472

Ala Ala Gly Lys Ala Thr Thr Glu Glu Gln Lys Leu Ile Glu Asp Ile
1 5 10 15

Asn Val Gly Phe Lys Ala Ala Val Ala Ala Ala

20 25

<210> 473
<211> 33
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5b precursor epitope

<400> 473

Ala Ala Gly Lys Ala Thr Thr Glu Glu Gln Lys Leu Ile Glu Asp Ile
1 5 10 15

Asn Val Gly Phe Lys Ala Ala Val Ala Ala Ala Ala Ser Val Pro Ala
20 25 30

Ala

<210> 474
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5b precursor epitope

<400> 474

Ala Ala Val Ala Ala Ala Ala Ser Val Pro Ala Ala Asp Lys Phe Lys
1 5 10 15

Thr Phe Glu

<210> 475
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5b precursor epitope

<400> 475

Ala Lys Phe Asp Ser Phe Val Ala Ser Leu Thr Glu Ala Leu Arg Val
1 5 10 15

Ile Ala Gly Ala Leu Glu Val His Ala Val Lys
20 25

<210> 476
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 5b precursor epitope

<400> 476

Ala Met Ser Glu Val Gln Lys Val Ser Gln Pro Ala Thr Gly Ala Ala
1 5 10 15

Thr Val Ala

<210> 477
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Polygalacturonase epitope

<400> 477

Ala Arg Trp Lys Asn Ser Lys Ile Trp Leu Gln Phe Ala Gln Leu Thr
1 5 10 15

Asp Phe Asn Leu
20

<210> 478
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Polygalacturonase epitope

<400> 478

Ala Val Leu Leu Val Pro Ala Asn Lys Lys Phe Phe Val Asn Asn Leu
1 5 10 15

Val Phe Arg Gly
20

<210> 479
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Polygalacturonase epitope

<400> 479

Asp Gly Thr Ile Val Ala Gln Pro Asp Pro Ala Arg Trp Lys Asn Ser
1 5 10 15

Lys Ile Trp Leu
20

<210> 480
<211> 20
<212> PRT

<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Polygalacturonase epitope

<400> 480

Phe Phe Val Asn Asn Leu Val Phe Arg Gly Pro Cys Gln Pro His Leu
1 5 10 15

Ser Phe Lys Val
20

<210> 481
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Polygalacturonase epitope

<400> 481

Phe Gly Glu Cys Glu Gly Val Lys Ile Gln Gly Leu Lys Ile Lys Ala
1 5 10 15

Pro Arg Asp Ser
20

<210> 482
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Polygalacturonase precursor epitope

<400> 482

Ala Ala Tyr Gln Asn Pro Ala Ser Trp Lys Asn Asn Arg Ile Trp
1 5 10 15

<210> 483
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Polygalacturonase precursor epitope

<400> 483

Ala Cys Lys Lys Pro Ser Ala Met Leu Leu Val Pro Gly Asn Lys
1 5 10 15

<210> 484
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Polygalacturonase precursor epitope

<400> 484

Ala Ile Lys Phe Asp Phe Ser Thr Gly Leu Ile Ile Gln Gly Leu
1 5 10 15

<210> 485
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Polygalacturonase precursor epitope

<400> 485

Ala Ile Asn Ile Phe Asn Val Glu Lys Tyr Gly Ala Val Gly Asp
1 5 10 15

<210> 486
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Polygalacturonase precursor epitope

<400> 486

Ala Asn Gly Tyr Phe Ser Gly His Val Ile Pro Ala Cys Lys Asn
1 5 10 15

<210> 487
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Arabidopsis thaliana Probable pectate lyase 18 precursor epitope

<400> 487

Gly His Ser Asp Thr Tyr Ser Arg Asp Lys Asn Met Gln Val Thr Ile
1 5 10 15

<210> 488
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Profilin-2/4 epitope

<400> 488

Leu Gly His Asp Gly Thr Val Trp Ala Gln Ser Ala Asp Phe Pro
1 5 10 15

<210> 489
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Pro-hevein precursor epitope

<400> 489

Asp Glu Tyr Cys Ser Pro Asp His Asn Cys Gln Ser Asn Cys Lys Asp
1 5 10 15

Ser Gly Glu Gly
20

<210> 490
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Pro-hevein precursor epitope

<400> 490

Glu Gln Cys Gly Arg Gln Ala Gly Gly Lys Leu Cys Pro Asn Asn Leu
1 5 10 15

Cys Cys Ser Gln
20

<210> 491
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Pro-hevein precursor epitope

<400> 491

Glu Gln Cys Gly Arg Gln Ala Gly Gly Lys Leu Cys Pro Asn Asn Leu
1 5 10 15

Cys Cys Ser Gln Trp Gly Trp Cys Gly Ser Thr Asp Glu Tyr Cys Ser
20 25 30

Pro Asp His Asn Cys Gln Ser Asn Cys Lys Asp
35 40

<210> 492
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Pro-hevein precursor epitope

<400> 492

Lys Leu Cys Pro Asn Asn Leu Cys Cys Ser Gln Trp Gly Trp Cys Gly
1 5 10 15

Ser Thr Asp Glu

20

<210> 493
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Pro-hevein precursor epitope

<400> 493

Asn Gly Gly Leu Asp Leu Asp Val Asn Val Phe Arg Gln Leu Asp Thr
1 5 10 15

Asp Gly Lys Gly
20

<210> 494
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica pru p 1 epitope

<400> 494

Gly Lys Cys Gly Val Ser Ile Pro Tyr Lys
1 5 10

<210> 495
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica pru p 1 epitope

<400> 495

Ile Thr Cys Gly Gln Val Ser Ser Ser Leu
1 5 10

<210> 496
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica pru p 1 epitope

<400> 496

Ser Ile Pro Tyr Lys Ile Ser Ala Ser Thr
1 5 10

<210> 497
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica pru p 1 epitope

<400> 497

Asp Arg Gln Ala Ala Cys Asn Cys Leu Lys Gln Leu Ser Ala Ser
1 5 10 15

<210> 498
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica pru p 1 epitope

<400> 498

Val Asn Pro Asn Asn Ala Ala Ala Leu Pro Gly Lys Cys Gly Val
1 5 10 15

<210> 499
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Arabidopsis thaliana Putative pectate lyase 17 precursor epitope

<400> 499

Gly His Asn Asp Asn Phe Val Lys Asp Val Lys Met Lys Val Thr Val
1 5 10 15

<210> 500
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens RAD51-like 1 isoform 1 epitope

<400> 500

Thr Arg Leu Ile Leu Gln Tyr Leu Asp Ser Glu Arg Arg Gln Ile Leu
1 5 10 15

<210> 501
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 501

Asp Pro Gly Pro Ala Arg Val Ile Tyr Thr Tyr Pro Asn Lys Val Phe
1 5 10 15

<210> 502
<211> 20
<212> PRT

<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 502

Ala Thr Trp Thr Cys Ile Asn Gln Gln Leu Asn Pro Lys Thr Asn Lys
1 5 10 15

Trp Glu Asp Lys
20

<210> 503
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 503

His Tyr Leu Leu Glu Phe Pro Thr Phe Pro Asp Gly His Asp Tyr Lys
1 5 10 15

Phe Asp Ser Lys
20

<210> 504
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 504

Lys Phe Asp Ser Lys Lys Pro Lys Glu Asp Pro Gly Pro Ala Arg Val
1 5 10 15

Ile Tyr Thr Tyr
20

<210> 505
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 505

Leu Ile Lys Gly Arg Thr Pro Ile Lys Phe Gly Lys Ala Asp Cys Asp
1 5 10 15

Arg Pro Pro Lys
20

<210> 506
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 506

Ser Tyr Pro His Trp Phe Thr Asn Gly Tyr Asp Gly Asn Gly Lys Leu
1 5 10 15

Ile Lys Gly Arg
20

<210> 507
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Rubber elongation factor protein epitope

<400> 507

Ala Glu Asp Glu Asp Asn Gln Gln Gly Gln Gly Glu Gly Leu Lys Tyr
1 5 10 15

Leu Gly Phe

<210> 508
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Rubber elongation factor protein epitope

<400> 508

Phe Ser Asn Val Tyr Leu Phe Ala Lys Asp Lys Ser Gly Pro Leu Gln
1 5 10 15

Pro Gly Val

<210> 509
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Rubber elongation factor protein epitope

<400> 509

Lys Phe Val Asp Ser Thr Val Val Ala Ser Val Thr Ile Ile Asp Arg
1 5 10 15

Ser Leu Pro

<210> 510
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Rubber elongation factor protein epitope

<400> 510

Gln Pro Gly Val Asp Ile Ile Glu Gly Pro Val Lys Asn Val Ala Val
1 5 10 15

Pro Leu Tyr

<210> 511
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Rubber elongation factor protein epitope

<400> 511

Arg Ser Leu Pro Pro Ile Val Lys Asp Ala Ser Ile Gln Val Val Ser
1 5 10 15

Ala Ile Arg

<210> 512
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Serum albumin precursor epitope

<400> 512

Asp Asp Ser Pro Asp Leu Pro Lys Leu Lys Pro Asp Pro Asn Thr Leu
1 5 10 15

Cys

<210> 513
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Serum albumin precursor epitope

<400> 513

Glu Lys Asp Ala Ile Pro Glu Asn Leu Pro Pro Leu Thr Ala Asp Phe
1 5 10 15

Ala Glu Asp Lys
20

<210> 514
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Serum albumin precursor epitope

<400> 514

Glu Ser His Ala Gly Cys Glu Lys Ser
1 5

<210> 515
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Serum albumin precursor epitope

<400> 515

His Pro Glu Tyr Ala Val Ser Val Leu Leu
1 5 10

<210> 516
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Serum albumin precursor epitope

<400> 516

Leu Ser Leu Ile Leu Asn Arg Leu Cys
1 5

<210> 517
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Small rubber particle protein epitope

<400> 517

Asp Phe Val Arg Ala Ala Gly Val Tyr Ala Val Asp
1 5 10

<210> 518
<211> 12
<212> PRT

<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Small rubber particle protein epitope

<400> 518

Lys Tyr Leu Asp Phe Val Arg Ala Ala Gly Val Tyr
1 5 10

<210> 519
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Small rubber particle protein epitope

<400> 519

Asn Val Val Lys Thr Val Val Thr Pro Val Tyr Tyr
1 5 10

<210> 520
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Small rubber particle protein epitope

<400> 520

Pro Arg Ile Val Leu Asp Val Ala Ser Ser Val Phe
1 5 10

<210> 521
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Small rubber particle protein epitope

<400> 521

Gln Gly Tyr Arg Val Ser Ser Tyr Leu Pro Leu Leu
1 5 10

<210> 522
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max Stress-induced protein SAM22 epitope

<400> 522

Ala Leu Phe Lys Ala Ile Glu Ala Tyr Leu Leu Ala His Pro Asp
1 5 10 15

<210> 523

<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Sugi basic protein precursor epitope

<400> 523

Ala Phe Asn Val Glu Asn Gly Asn Ala Thr Pro Gln Leu Thr Lys
1 5 10 15

<210> 524
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Sugi basic protein precursor epitope

<400> 524

Ala Asn Asn Asn Tyr Asp Pro Trp Thr Ile Tyr Ala Ile Gly Gly
1 5 10 15

<210> 525
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Sugi basic protein precursor epitope

<400> 525

Ala Tyr Ser Asp Asp Lys Ser Met Lys Val Thr Val Ala Phe Asn
1 5 10 15

<210> 526
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Sugi basic protein precursor epitope

<400> 526

Cys Gly Gln Arg Met Pro Arg Ala Arg Tyr Gly Leu Val His Val
1 5 10 15

<210> 527
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Sugi basic protein precursor epitope

<400> 527

Cys Ser Asn Trp Val Trp Gln Ser Thr Gln Asp Val Phe Tyr Asn
1 5 10 15

<210> 528
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Trichophyton rubrum Tri r 2 allergen epitope

<400> 528

Ala Asp Phe Ser Asn Tyr Gly Ala Val Val Asp Val Tyr Ala Pro Gly
1 5 10 15

Lys Asp Ile Thr
20

<210> 529
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Trichophyton rubrum Tri r 2 allergen epitope

<400> 529

Ala Lys Gly Val Ser Leu Val Ala Val Lys Val Leu Asp Cys Asp Gly
1 5 10 15

Ser Gly Ser Asn
20

<210> 530
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Trichophyton rubrum Tri r 2 allergen epitope

<400> 530

Ala Ser Asn Gln Ala Ala Lys Ala Ile Ser Asp Ala Gly Ile Phe Met
1 5 10 15

Ala Val Ala Ala
20

<210> 531
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Trichophyton rubrum Tri r 2 allergen epitope

<400> 531

Asp Cys Asn Gly His Gly Thr His Val Ala Gly Thr Val Gly Gly Thr
1 5 10 15

Lys Tyr Gly Leu
20

<210> 532
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Trichophyton rubrum Tri r 2 allergen epitope

<400> 532

Asp Pro Ser Ala Gly Lys Gly Val Thr Ala Tyr Ile Ile Asp Thr Gly
1 5 10 15

Ile Asp Ile Asp
20

<210> 533
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Vespula vulgaris Venom allergen 5 precursor epitope

<400> 533

Ala Cys Lys Tyr Gly Ser Leu Lys Pro Asn Cys Gly
1 5 10

<210> 534
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Vespula vulgaris Venom allergen 5 precursor epitope

<400> 534

Cys Asn Tyr Gly Pro Ser Gly Asn Phe Met Asn Glu
1 5 10

<210> 535
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Vespula vulgaris Venom allergen 5 precursor epitope

<400> 535

Asp Val Ala Lys Tyr Gln Val Gly Gln Asn Val Ala
1 5 10

<210> 536
<211> 12
<212> PRT

<213> Artificial Sequence

<220>
<223> Vespula vulgaris Venom allergen 5 precursor epitope

<400> 536

Glu Lys Trp His Lys His Tyr Leu Val Cys Asn Tyr
1 5 10

<210> 537
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Vespula vulgaris Venom allergen 5 precursor epitope

<400> 537

Glu Leu Ala Tyr Val Ala Gln Val Trp Ala Asn Gln
1 5 10

<210> 538
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana 11S globulin-like protein epitope

<400> 538

Ala Phe Gln Ile Ser Arg Glu Glu Ala Arg Arg Leu Lys Tyr Asn
1 5 10 15

<210> 539
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Carya illinoinensis 11S legumin protein epitope

<400> 539

Glu Glu Ser Gln Arg Gln Ser Gln Gln Gly Gln Arg
1 5 10

<210> 540
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Fagopyrum esculentum 13S globulin epitope

<400> 540

Asp Ala His Gln Pro Thr Arg Arg Val Arg Lys Gly Asp Val Val Ala
1 5 10 15

Leu Pro

<210> 541
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Fagopyrum esculentum 13S globulin seed storage protein 1 precursor (Legumin-like protein 1) epitope

<400> 541

Phe Lys Gln Asn Val Asn Arg Pro Ser Arg Ala Asp
1 5 10

<210> 542
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Fagopyrum esculentum 13S globulin seed storage protein 3 precursor (Legumin-like protein 3) (Allergen Fag e 1) epitope

<400> 542

Asp Ile Ser Thr Lys Glu Ala Phe Arg Leu Lys Asn
1 5 10

<210> 543
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Anacardium occidentale 2s albumin epitope

<400> 543

Cys Gln Arg Gln Phe Glu Glu Gln Gln Arg Phe Arg
1 5 10

<210> 544
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Sesamum indicum 2S seed storage protein 1 epitope

<400> 544

His Phe Arg Glu Cys Cys Asn Glu Ile Arg
1 5 10

<210> 545
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Sesamum indicum 2S seed storage protein 1 precursor epitope

<400> 545

Cys Met Gln Trp Met Arg Ser Met Arg Gly
1 5 10

<210> 546
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bertholletia excelsa 2S sulfur-rich seed storage protein precursor (Allergen Ber e 1) epitope

<400> 546

Cys Arg Cys Glu Gly Leu Arg Met Met Met Met Arg Met Gln
1 5 10

<210> 547
<211> 40
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 52 kDa Ro protein epitope

<400> 547

Leu Glu Lys Asp Glu Arg Glu Gln Leu Arg Ile Leu Gly Glu Lys Glu
1 5 10 15

Ala Lys Leu Ala Gln Gln Ser Gln Ala Leu Gln Glu Leu Ile Ser Glu
20 25 30

Leu Asp Arg Arg Cys His Ser Ser
35 40

<210> 548
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 52-kD SS-A/Ro autoantigen epitope

<400> 548

Gln Glu Lys Leu Gln Val Ala Leu Gly Glu
1 5 10

<210> 549
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 5-hydroxytryptamine (serotonin) receptor 4 epitope

<400> 549

Gly Ile Ile Asp Leu Ile Glu Lys Arg Lys Phe Asn Gln Asn Ser Asn
1 5 10 15

Ser Thr Tyr Cys Val
20

<210> 550
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 60 kDa heat shock protein, mitochondrial precursor epitope

<400> 550

Asp Gly Val Ala Val Leu Lys Val Gly Gly
1 5 10

<210> 551
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 60 kDa SS-A/Ro ribonucleoprotein epitope

<400> 551

Glu Leu Tyr Lys Glu Lys Ala Leu Ser Val Glu Thr Glu Lys Leu Leu
1 5 10 15

Lys Tyr Leu Glu Ala Val
20

<210> 552
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 60S acidic ribosomal protein P0 epitope

<400> 552

Ala Lys Val Glu Ala Lys Glu Glu Ser Glu Glu Ser Asp Glu Asp Met
1 5 10 15

Gly Phe Gly Leu Phe Asp
20

<210> 553
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 60S acidic ribosomal protein P2 epitope

<400> 553

Glu Glu Ser Asp Asp Asp Met Gly Phe Gly Leu Phe Asp
1 5 10

<210> 554
<211> 50
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 64 Kd autoantigen epitope

<400> 554

Ala Thr Lys Lys Glu Glu Glu Lys Lys Gly Gly Asp Arg Asn Thr Gly
1 5 10 15

Leu Ser Arg Asp Lys Asp Lys Lys Arg Glu Glu Met Lys Glu Val Ala
20 25 30

Lys Lys Glu Asp Asp Glu Lys Val Lys Gly Glu Arg Arg Asn Thr Asp
35 40 45

Thr Arg
50

<210> 555
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens 65 kDa heat shock protein epitope

<400> 555

Ala Leu Leu Arg Cys Ile Pro Ala Leu Asp Ser Leu Thr Pro Ala Asn
1 5 10 15

Glu Asp Cys

<210> 556
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Acetylcholine receptor subunit alpha precursor epitope

<400> 556

Ala Ile Asn Pro Glu Ser Asp Gln Pro Asp Leu Ser Asn Phe
1 5 10

<210> 557
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon acidic Cyn d 1 isoallergen isoform 1 precursor epitope

<400> 557

Gln Asp Asp Val Ile Pro Glu Asp Trp Lys Pro Asp Thr Val Tyr Lys
1 5 10 15

Ser Lys Ile Gln Phe
20

<210> 558
<211> 50
<212> PRT
<213> Artificial Sequence

<220>
<223> Cynodon dactylon acidic Cyn d 1 isoallergen isoform 3 precursor epitope

<400> 558

Glu Glu Asp Lys Leu Arg Lys Ala Gly Glu Leu Met Leu Gln Phe Arg
1 5 10 15

Arg Val Lys Cys Glu Tyr Pro Ser Asp Thr Lys Ile Thr Phe His Val
20 25 30

Glu Lys Gly Ser Ser Pro Asn Tyr Leu Ala Leu Leu Val Lys Tyr Ala
35 40 45

Ala Gly
50

<210> 559
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens acidic ribosomal phosphoprotein (P0) epitope

<400> 559

Ala Ala Ala Ala Ala Pro Ala Lys
1 5

<210> 560
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens acidic ribosomal phosphoprotein (P1) epitope

<400> 560

Glu Ser Glu Glu Ser Asp Asp Asp Met Gly Phe Gly Leu Phe Asp
1 5 10 15

<210> 561
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens acidic ribosomal phosphoprotein (P2) epitope

<400> 561

Ala Pro Ala Ala Gly Ser Ala Pro Ala Ala Ala Glu Glu Lys Lys
1 5 10 15

<210> 562
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Adrenergic, beta-2-, receptor, surface epitope

<400> 562

His Trp Tyr Arg Ala Thr His Gln Glu Ala Ile Asn Cys Tyr Ala Asn
1 5 10 15

<210> 563
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Alanyl-tRNA synthetase, cytoplasmic epitope

<400> 563

Phe Ile Asp Glu Pro Arg Arg Arg Pro Ile
1 5 10

<210> 564
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus albumin epitope

<400> 564

Pro Val Glu Ser Lys Val Thr
1 5

<210> 565
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Juglans regia Albumin seed storage protein epitope

<400> 565

Gly Leu Arg Gly Glu Glu Met Glu Glu Met Val Gln Ser
1 5 10

<210> 566
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus alcohol dehydrogenase epitope

<400> 566

Ala Val Asn Gly Asp Trp Pro Leu Pro Thr Lys Leu Pro Leu Val Gly
1 5 10 15

Gly His

<210> 567
<211> 37
<212> PRT
<213> Artificial Sequence

<220>
<223> Penicillium chrysogenum alkaline serine protease epitope

<400> 567

Ala Asn Val Val Gln Arg Asn Ala Pro Ser Trp Gly Leu Ser Arg Ile
1 5 10 15

Ser Ser Lys Lys Ser Gly Ala Thr Asp Tyr Val Tyr Asp Ser Thr Ala
20 25 30

Gly Glu Gly Ile Val
35

<210> 568
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea allergen epitope

<400> 568

Asp Asp Gln Cys Gln Arg Gln Leu Gln Arg
1 5 10

<210> 569
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Anacardium occidentale allergen Ana o 2 epitope

<400> 569

Glu Glu Ser Glu Asp Glu Lys Arg Arg Trp Gly Gln Arg Asp Asn
1 5 10 15

<210> 570
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea Allergen Ara h 1, clone P41B precursor epitope

<400> 570

Ala Lys Ser Ser Pro Tyr Gln Lys Lys Thr
1 5 10

<210> 571
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea allergen Arah3/Arah4 epitope

<400> 571

Ala Gly Val Ala Leu Ser Arg Leu Val Leu Arg Arg Asn Ala Leu
1 5 10 15

<210> 572
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea allergen Arah6 epitope

<400> 572

Asp Arg Gln Met Val Gln His Phe Lys Arg
1 5 10

<210> 573
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Periplaneta americana Allergen Cr-PI epitope

<400> 573

Ile Pro Lys Gly Lys Lys Gly Gly Gln Ala Tyr
1 5 10

<210> 574
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus allergen I/a; Asp f I/a epitope

<400> 574

Ile Asn Gln Gln Leu Asn Pro Lys
1 5

<210> 575
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea Allergen II epitope

<400> 575

Asp Arg Leu Gln Gly Arg Gln Gln Glu Gln
1 5 10

<210> 576
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Lens culinaris allergen Len c 1.0101 epitope

<400> 576

Ala Ile Asn Ala Ser Ser Asp Leu Asn Leu Ile Gly Phe Gly Ile
1 5 10 15

<210> 577
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Allergen Mag epitope

<400> 577

Asp Val Glu Leu Ser Leu Arg Ser Ser Asp Ile Ala
1 5 10

<210> 578
<211> 31
<212> PRT
<213> Artificial Sequence

<220>
<223> Penicillium chrysogenum Allergen Pen n 18 epitope

<400> 578

Ala His Ile Lys Lys Ser Lys Lys Gly Asp Lys Lys Phe Lys Gly Ser
1 5 10 15

Val Ala Asn Met Ser Leu Gly Gly Gly Ser Ser Arg Thr Leu Asp
20 25 30

<210> 579
<211> 14

<212> PRT
<213> Artificial Sequence

<220>
<223> Sinapis alba Allergen Sin a 1 epitope

<400> 579

Gln Gly Pro His Val Ile Ser Arg Ile Tyr Gln Thr Ala Thr
1 5 10

<210> 580
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Ziziphus mauritiana allergen Ziz m 1 epitope

<400> 580

Lys Thr Asn Tyr Ser Ser Ser Ile Ile Leu Glu Tyr
1 5 10

<210> 581
<211> 37
<212> PRT
<213> Artificial Sequence

<220>
<223> Fagopyrum tataricum allergenic protein epitope

<400> 581

Asp Ile Ser Thr Glu Glu Ala Tyr Lys Leu Lys Asn Gly Arg Gln Glu
1 5 10 15

Val Glu Val Phe Arg Pro Phe Gln Ser Arg Tyr Glu Lys Glu Glu Glu
20 25 30

Lys Glu Arg Glu Arg
35

<210> 582
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens alpha 2 interferon epitope

<400> 582

Glu Val Val Arg Ala Glu Ile Met Arg Ser Phe Ser Leu Ser
1 5 10

<210> 583
<211> 30
<212> PRT
<213> Artificial Sequence

<220>

<223> Bos taurus alpha S1 casein epitope

<400> 583

Glu Asp Gln Ala Met Glu Asp Ile Lys Gln Met Glu Ala Glu Ser Ile
1 5 10 15

Ser Ser Ser Glu Glu Ile Val Pro Asn Ser Val Glu Gln Lys
20 25 30

<210> 584
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Alpha/beta-gliadin A-II precursor epitope

<400> 584

Gln Val Ser Phe Gln Gln Pro Gln Gln Gln
1 5 10

<210> 585
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Alpha/beta-gliadin A-V epitope

<400> 585

Leu Ala Leu Gln Thr Leu Pro Ala Met Cys
1 5 10

<210> 586
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens alpha-1 type IV collagen epitope

<400> 586

Ser Arg Cys Gln Val Cys Met Arg Arg Thr
1 5 10

<210> 587
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens alpha1A-voltage-dependent calcium channel epitope

<400> 587

Glu Asp Ser Asp Glu Asp Glu Phe Gln Ile Thr Glu
1 5 10

<210> 588
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens alpha-2 type XI collagen epitope

<400> 588

Gly Ser Leu Asp Ser Leu Arg Arg Glu Ile Glu Gln Met Arg Arg
1 5 10 15

<210> 589
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus alpha2(I) collagen epitope

<400> 589

Leu Pro Gly Leu Lys Gly His Asn Gly Leu Gln Gly Leu Pro Gly Leu
1 5 10 15

Ala Gly His His
20

<210> 590
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Alpha-amylase inhibitor 0.28 precursor (CIII) (WMAI-1) epitope

<400> 590

Ala Tyr Pro Asp Val
1 5

<210> 591
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Alpha-enolase epitope

<400> 591

Lys Ile His Ala Arg Glu Ile Phe Asp Ser Arg Gly Asn Pro Thr Val
1 5 10 15

Glu

<210> 592
<211> 15
<212> PRT

<213> Artificial Sequence

<220>
<223> Homo sapiens alpha-fibrinogen precursor epitope

<400> 592

Gly Pro Arg Val Val Glu Arg His Gln Ser Ala Cys Lys Asp Ser
1 5 10 15

<210> 593
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Alpha-gliadin epitope

<400> 593

Leu Gly Gln Gly Ser Phe Arg Pro Ser Gln Gln Asn
1 5 10

<210> 594
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-lactalbumin epitope

<400> 594

Lys Asp Leu Lys Gly Tyr Gly Gly Val Ser
1 5 10

<210> 595
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-lactalbumin precursor epitope

<400> 595

Lys Cys Glu Val Phe Arg Glu Leu Lys Asp Leu Lys Gly Tyr
1 5 10

<210> 596
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus alpha-S1-casein epitope

<400> 596

Leu Asn Glu Asn Leu Leu Arg Phe Phe Val Ala Pro Phe Pro Gln Val
1 5 10 15

Phe Gly Lys Glu

20

<210> 597
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S1-casein precursor epitope

<400> 597

Ala Met Glu Asp Ile Lys Gln Met Glu Ala
1 5 10

<210> 598
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Alpha-S2-casein precursor epitope

<400> 598

Glu Asn Leu Cys Ser Thr Phe Cys Lys Glu
1 5 10

<210> 599
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens anti-beta-amyloid peptide immunoglobulin heavy chain variable region epitope

<400> 599

Ala His Ile Trp Trp Asn Asp
1 5

<210> 600
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Aquaporin-4 epitope

<400> 600

Phe Cys Pro Asp Val Glu Phe Lys Arg Arg Phe Lys Glu Ala Phe Ser
1 5 10 15

Lys Ala Ala Gln Gln Thr Lys Gly
20

<210> 601
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea Ara h 2.01 allergen epitope

<400> 601

Cys Cys Asn Glu Leu Asn Glu Phe Glu Asn Asn Gln Arg Cys Met
1 5 10 15

<210> 602
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens ATP-dependent DNA helicase 2 subunit 2 epitope

<400> 602

Glu Glu Ala Ser Gly Ser Ser Val Thr Ala Glu Glu Ala Lys Lys Phe
1 5 10 15

Leu Ala Pro Lys
20

<210> 603
<211> 28
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens autoantigen epitope

<400> 603

Glu Ile Arg Val Arg Leu Gln Ser Ala Ser Pro Ser Thr Arg Trp Thr
1 5 10 15

Glu Leu Asp Asp Val Lys Arg Leu Leu Lys Gly Ser
20 25

<210> 604
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Band 3 anion transport protein epitope

<400> 604

Leu Phe Lys Pro Pro Lys Tyr His Pro Asp Val Pro Tyr Val Lys Arg
1 5 10 15

<210> 605
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max Bd 30K (34 kDa maturing seed protein) epitope

<400> 605

Glu Asp Trp Gly Glu Asp Gly Tyr Ile Trp Ile Gln Arg Asn Thr
1 5 10 15

<210> 606
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Bence Jones protein HAG epitope

<400> 606

Ala Trp His Gln Gln Gln Pro
1 5

<210> 607
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Bet v 4 epitope

<400> 607

Phe Ala Arg Ala Asn Arg Gly Leu
1 5

<210> 608
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Musa acuminata beta-1, 3-glucananse epitope

<400> 608

Gly Leu Phe Tyr Pro Asn Lys Gln Pro
1 5

<210> 609
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis beta-1,3-glucanase epitope

<400> 609

Gly Leu Phe Phe Pro Asp Lys Arg Pro Lys Tyr Asn Leu Asn Phe
1 5 10 15

<210> 610
<211> 12
<212> PRT
<213> Artificial Sequence

<220>

<223> Olea europaea beta-1,3-glucanase-like protein epitope

<400> 610

Ala Gly Arg Asn Ser Trp Asn Cys Asp Phe Ser Gln
1 5 10

<210> 611
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens beta-2-glycoprotein 1 precursor epitope

<400> 611

Leu Lys Thr Pro Arg Val
1 5

<210> 612
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens beta-2-glycoprotein I epitope

<400> 612

Thr Leu Arg Val Tyr Lys
1 5

<210> 613
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus beta-casein epitope

<400> 613

Gln Ser Lys Val Leu Pro Val Pro Gln Lys Ala Val Pro
1 5 10

<210> 614
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Beta-casein precursor epitope

<400> 614

Asp Glu Leu Gln Asp Lys Ile His Pro Phe Ala Gln
1 5 10

<210> 615
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Beta-lactoglobulin epitope

<400> 615

Ala Gln Lys Lys Ile Ile Ala Glu Lys Thr
1 5 10

<210> 616
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Beta-lactoglobulin precursor epitope

<400> 616

Ala Ala Ser Asp Ile Ser Leu Leu Asp Ala Gln Ser Ala Pro Leu Arg
1 5 10 15

<210> 617
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Botulinum neurotoxin type E epitope

<400> 617

Trp Lys Ala Pro Ser Ser Pro
1 5

<210> 618
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens bullous pemphigoid antigen epitope

<400> 618

Lys Ser Thr Ala Lys Asp Cys Thr Phe Lys Pro Asp Phe Glu Met Thr
1 5 10 15

Val Lys Glu

<210> 619
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Bullous pemphigoid antigen 1, isoforms 1/2/3/4/5/8 epitope

<400> 619

Leu Thr Asp Thr Lys Thr Gly Leu His Phe Asn Ile Asn Glu Ala Ile

1 5 10 15

Glu Gln Gly Thr
20

<210> 620
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Fagopyrum esculentum BW 16kDa allergen epitope

<400> 620

Glu Gly Val Arg Asp Leu Lys Glu
1 5

<210> 621
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens calcium channel, alpha 1A subunit isoform 3 epitope

<400> 621

Gly Asn Ile Gly Ile Asp Val Glu Asp Glu Asp Ser Asp Glu Asp Glu
1 5 10 15

Phe

<210> 622
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Calpastatin epitope

<400> 622

Ala Val Cys Arg Thr Ser Met Cys Ser Ile Gln Ser Ala Pro Pro
1 5 10 15

<210> 623
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Calreticulin precursor epitope

<400> 623

Lys Glu Gln Phe Leu Asp Gly Asp Gly Trp Thr Ser Arg Trp Ile Glu
1 5 10 15

Ser Lys

<210> 624
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ca-sensing receptor epitope

<400> 624

Phe Val Ala Gln Asn Lys Ile Asp Ser Leu Asn Leu Asp
1 5 10

<210> 625
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Caspase-8 precursor epitope

<400> 625

Asp Arg Asn Gly Thr His Leu Asp Ala
1 5

<210> 626
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens centromere protein A isoform a epitope

<400> 626

Gly Pro Ser Arg Arg Gly Pro Ser Leu Gly Ala Ser Ser His
1 5 10

<210> 627
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens centromere protein B, 80kDa epitope

<400> 627

Met Gly Pro Lys Arg Arg Gln Leu Thr Phe
1 5 10

<210> 628
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens centromere protein-A epitope

<400> 628

Glu Ala Pro Arg Arg Ser Pro Ser Pro
1 5 10

<210> 629
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Chain A, Birch Pollen Profilin epitope

<400> 629

Ala Gln Ser Ser Ser Phe Pro Gln Phe Lys Pro Gln Glu Ile Thr Gly
1 5 10 15

Ile

<210> 630
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Chain A, Crystal Structure Of The Glycosylated Five-Domain Human Beta2-Glycoprotein I Purified From Blood Plasma epitope

<400> 630

Arg Gly Gly Met Arg
1 5

<210> 631
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Chain H, Three-Dimensional Structure Of A Human Immunoglobulin With A Hinge Deletion epitope

<400> 631

Ala Leu Pro Ala Pro Ile Glu
1 5

<210> 632
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens cholesterol side-chain cleavage enzyme P450scc (EC 1.14.15.67) epitope

<400> 632

Phe Asp Pro Glu Asn Phe Asp Pro Thr Arg Trp Leu Ser Lys Asp Lys
1 5 10 15

Asn Ile Thr Tyr Phe Arg Asn Leu Gly Phe Gly
20 25

<210> 633
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens citrate synthase epitope

<400> 633

Ala Leu Lys His Leu Pro Asn Asp Pro Met
1 5 10

<210> 634
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens claudin 11 epitope

<400> 634

Ala His Arg Glu Thr
1 5

<210> 635
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Coagulation factor VIII precursor epitope

<400> 635

Ala Pro Asp Asp Arg Ser Tyr Lys Ser Gln Tyr
1 5 10

<210> 636
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Oncorhynchus mykiss collagen a2(I) epitope

<400> 636

Met Lys Gly Leu Arg Gly His Gly Gly Leu Gln Gly Met Pro Gly Pro
1 5 10 15

Asn Gly Pro Ser
20

<210> 637
<211> 14
<212> PRT

<213> Artificial Sequence

<220>
<223> Homo sapiens Collagen alpha-1(II) chain epitope

<400> 637

Ala Arg Gly Ala Gln Gly Pro Pro Gly Ala Thr Gly Phe Pro
1 5 10

<210> 638
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens collagen alpha-1(VII) chain precursor epitope

<400> 638

Gly Thr Leu His Val Val Gln Arg
1 5

<210> 639
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Collagen alpha-1(XVII) chain epitope

<400> 639

Arg Ser Ile Leu Pro Tyr Gly Asp Ser Met Asp Arg Ile Glu Lys Asp
1 5 10 15

Arg Leu Gln Gly Met Ala Pro
20

<210> 640
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Collagen alpha-3(IV) chain epitope

<400> 640

Thr Ala Ile Pro Ser Cys Pro Glu Gly Thr Val Pro Leu Tyr Ser
1 5 10 15

<210> 641
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens collagen VII epitope

<400> 641

Ile Ile Trp Arg Ser Thr Gln Gly

1 5

<210> 642
<211> 46
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus collagen, type I, alpha 2 epitope

<400> 642

Ala Pro Gly Pro Asp Gly Asn Asn Gly Ala Gln Gly Pro Pro Gly Leu
1 5 10 15

Gln Gly Val Gln Gly Gly Lys Gly Glu Gln Gly Pro Ala Gly Pro Pro
20 25 30

Gly Phe Gln Gly Leu Pro Gly Pro Ala Gly Thr Ala Gly Glu
35 40 45

<210> 643
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens collagen, type II, alpha 1 epitope

<400> 643

Pro Pro Gly Pro Thr Gly Ala Ser Gly
1 5

<210> 644
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens collagen, type II, alpha 1 isoform 1 precursor epitope

<400> 644

Ala Arg Gly Leu Thr Gly Arg Pro Gly Asp Ala
1 5 10

<210> 645
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens collagen, type II, alpha 1 isoform 2 precursor epitope

<400> 645

Leu Val Gly Pro Arg Gly Glu Arg Gly Phe Pro
1 5 10

<210> 646
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Complement C1q subcomponent subunit A epitope

<400> 646

Lys Gly Glu Gln Gly Glu Pro Gly Ala
1 5

<210> 647
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Condensin-2 complex subunit D3 epitope

<400> 647

Pro Thr Pro Glu Thr Gly Pro Leu Gln Arg
1 5 10

<210> 648
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea Conglutin-7 precursor epitope

<400> 648

Ala Ala His Ala Ser Ala Arg Gln Gln Trp Glu Leu Gln Gly Asp
1 5 10 15

<210> 649
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Periplaneta americana Cr-PII allergen epitope

<400> 649

Ile Arg Ser Trp Phe Gly Leu Pro
1 5

<210> 650
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Cochliobolus lunatus Cytochrome c epitope

<400> 650

Glu Asn Pro Lys Lys Tyr Ile Pro Gly Thr Lys

1 5 10

<210> 651
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Rattus norvegicus Cytochrome P450 3A1 epitope

<400> 651

Asp Met Val Leu Asn Glu Thr Leu Arg Leu
1 5 10

<210> 652
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens cytoskeleton-associated protein 5 isoform b epitope

<400> 652

Cys Gln Ala Leu Val Arg Met Leu Ala Lys Lys Pro Gly Trp Lys
1 5 10 15

<210> 653
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Der f 2 epitope

<400> 653

Ile Ala Thr His Ala Lys Ile Arg Asp
1 5

<210> 654
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Der f 7 allergen epitope

<400> 654

His Ile Gly Gly Leu Ser Ile Leu Asp Pro Ile Phe Gly Val Leu
1 5 10 15

<210> 655
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Der p 1 allergen epitope

<400> 655

Ala Arg Glu Gln Ser Cys Arg Arg Pro Asn Ala Gln Arg Phe Gly Ile
1 5 10 15

Ser Asn Tyr Cys Gln Ile Tyr Pro Pro Asn Ala Asn Lys Ile Arg Glu
20 25 30

Ala Leu Ala Gln Thr His Ser Ala Ile Ala Val
35 40

<210> 656
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Der p 7 allergen polypeptide epitope

<400> 656

His Ile Gly Gly Leu Ser Ile Leu Asp Pro Ile Phe Ala Val Leu
1 5 10 15

<210> 657
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Desmoglein-1 epitope

<400> 657

Arg Glu Trp Ile Lys Phe Ala Ala Ala Cys Arg Glu
1 5 10

<210> 658
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Desmoglein-3 precursor epitope

<400> 658

Arg Glu Trp Val Lys Phe Ala Lys Pro Cys Arg Glu
1 5 10

<210> 659
<211> 30
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens desmoglein-3 preproprotein epitope

<400> 659

Ser Gln Glu Pro Ala Gly Thr Pro Met Phe Leu Leu Ser Arg Asn Thr
1 5 10 15

Gly Glu Val Arg Thr Leu Thr Asn Ser Leu Asp Arg Glu Gln
20 25 30

<210> 660
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens desmoplakin epitope

<400> 660

Gly Asn Ser Ser Tyr Ser Tyr Ser Tyr Ser Phe Ser
1 5 10

<210> 661
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens desmoplakin isoform II epitope

<400> 661

Leu Val Asp Arg Lys Thr Gly Ser Gln Tyr Asp Ile Gln Asp Ala Ile
1 5 10 15

Asp Lys Gly Leu
20

<210> 662
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens dihydrolipoamide S-acetyltransferase (E2 component of pyruvate dehydrogenase complex), isoform CRA_a epitope

<400> 662

Ala Glu Ile Glu Thr Asp Lys Ala Thr Ile Gly Phe Glu Val Gln Glu
1 5 10 15

Glu Gly

<210> 663
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens DNA topoisomerase 1 epitope

<400> 663

Gly Val Pro Ile Glu Lys Ile Tyr Asn Lys Thr Gln Arg Glu Lys Phe

1 5 10 15

Ala

<210> 664
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens DNA topoisomerase I epitope

<400> 664

Glu Leu Asp Gly Gln Glu Tyr Val Val Glu Phe Asp Phe Leu Gly Lys
1 5 10 15

Asp Ser Ile Arg
20

<210> 665
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens DNA topoisomerase II beta epitope

<400> 665

His Pro Met Leu Pro Asn Tyr Lys Asn Phe Lys Gly Thr Ile Gln Glu
1 5 10 15

Leu Gly Gln Asn
20

<210> 666
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens DNA-directed RNA polymerase II subunit RPB1 epitope

<400> 666

Tyr Ser Pro Thr Ser Pro Ser
1 5

<210> 667
<211> 36
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens E3 ubiquitin-protein ligase TRIM9 isoform 2 epitope

<400> 667

Ala Phe Asn Lys Thr Gly Val Ser Pro Tyr Ser Lys Thr Leu Val Leu

1 5 10 15

Gln Thr Ser Glu Gly Lys Ala Leu Gln Gln Tyr Pro Ser Glu Arg Glu
20 25 30

Leu Arg Gly Ile
35

<210> 668
<211> 39
<212> PRT
<213> Artificial Sequence

<220>
<223> Candida albicans Enolase 1 (2-phosphoglycerate dehydratase) (2-phospho-D-glycerate hydro-lyase) epitope

<400> 668

Gln Ala Ala Asn Asp Ser Tyr Ala Ala Gly Trp Gly Val Met Val Ser
1 5 10 15

His Arg Ser Gly Glu Thr Glu Asp Thr Phe Ile Ala Asp Leu Ser Val
20 25 30

Gly Leu Arg Ser Gly Gln Ile
35

<210> 669
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens enolase 1 variant epitope

<400> 669

Lys Ile His Ala Arg Glu Ile Phe Asp Ser Arg Gly Asn Pro Thr Val
1 5 10 15

<210> 670
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis ENSP-like protein epitope

<400> 670

Phe Pro Leu Ile Thr Cys Cys Gly Tyr Gly Gly Lys Tyr Asn Phe Ser
1 5 10 15

Val Thr Ala Pro Cys
20

<210> 671
<211> 20

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens envoplakin epitope

<400> 671

Ala Gly Glu Thr Lys Pro Ser Ser Ser Leu Ser Ile Gly Ser Ile Ile
1 5 10 15

Ser Lys Ser Pro
20

<210> 672
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Fagopyrum esculentum Fag e 1 epitope

<400> 672

Ala Val Val Leu Lys Ala Gly Asn Glu Gly Leu Glu
1 5 10

<210> 673
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Fas AMA epitope

<400> 673

Cys Val Pro
1

<210> 674
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens FGA protein epitope

<400> 674

Ser Arg Ala Leu Ala Arg Glu Val Asp Leu Lys Asp Tyr Glu Asp Gln
1 5 10 15

Gln Lys

<210> 675
<211> 20
<212> PRT
<213> Artificial Sequence

<220>

<223> Homo sapiens FGB protein epitope

<400> 675

Ala Arg Gly His Arg Pro Leu Asp Lys Lys Arg Glu Glu Ala Pro Ser
1 5 10 15

Leu Arg Pro Ala
20

<210> 676
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens fibrin beta epitope

<400> 676

Ala Asn Lys Tyr Gln Ile Ser Val Asn Lys Tyr Arg Gly Thr Ala Gly
1 5 10 15

Asn Ala Leu

<210> 677
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens fibrinogen alpha chain isoform alpha preproprotein epitope

<400> 677

Asp Ser Pro Gly Ser Gly Asn Ala Arg Pro Asn Asn Pro Asp Trp
1 5 10 15

<210> 678
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Fibrinogen alpha chain precursor epitope

<400> 678

Phe Leu Ala Glu Gly Gly Gly Val Arg Gly Pro Arg Val Val Glu Arg
1 5 10 15

His

<210> 679
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens fibrinogen alpha chain preproprotein, isoform alpha epitope

<400> 679

Asp His Glu Gly Thr His Ser Thr Lys Arg Gly His Ala Lys Ser Arg
1 5 10 15

Pro Val Arg Gly
20

<210> 680
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens fibrinogen beta chain epitope

<400> 680

Pro Arg Lys Gln Cys Ser Lys Glu Asp Gly Gly Gly Trp Trp Tyr
1 5 10 15

<210> 681
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens fibrinogen beta chain, isoform CRA_d epitope

<400> 681

Asn Glu Glu Gly Phe Phe Ser Ala Arg Gly His Arg Pro Leu Asp Lys
1 5 10 15

Lys

<210> 682
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens fibrinogen beta chain, isoform CRA_i epitope

<400> 682

Glu Glu Ala Pro Ser Leu Arg Pro Ala Pro Pro Pro Ile Ser Gly Gly
1 5 10 15

Gly Tyr Arg Ala Arg Pro Ala Lys
20

<210> 683
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Fibronectin precursor epitope

<400> 683

Leu Thr Ser Arg Pro Ala
1 5

<210> 684
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens filaggrin epitope

<400> 684

Asp Ser Gly His Arg Gly Tyr Ser Gly Ser Gln Ala Ser Asp Asn Glu
1 5 10 15

Gly His

<210> 685
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Follistatin-related protein 1 epitope

<400> 685

Leu Lys Phe Val Glu Gln Asn Glu
1 5

<210> 686
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Forkhead box protein E3 epitope

<400> 686

Pro Thr Pro Ala Pro Gly Pro Gly Arg Arg
1 5 10

<210> 687
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens GAD65 autoantigen glutamic acid decarboxylase epitope

<400> 687

Ala Pro Ala Met Ile Pro Pro

1 5

<210> 688
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Gamma-gliadin precursor epitope

<400> 688

Leu Gln Pro Gln Gln Pro Phe Pro Gln Gln
1 5 10

<210> 689
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III epitope

<400> 689

Ala His Thr Asp Phe Ala Gly Ala Glu Ala Ala Trp Gly Ala Thr Leu
1 5 10 15

Asp Thr Phe Phe Gly
20

<210> 690
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-III precursor epitope

<400> 690

Gly Val Thr His Asp Gln Leu Asn Asn Phe Arg
1 5 10

<210> 691
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-IV precursor epitope

<400> 691

Lys Ala His Thr Asp Phe Ala Gly Ala Glu Ala Ala Trp Gly Ala Thr
1 5 10 15

Leu Asp Ala Phe Phe Gly Met
20

<210> 692

<211> 35
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-VI precursor epitope

<400> 692

Ile Val Ser Phe Leu Ser Glu Val Ile Ser Leu Ala Gly Ser Asp Ala
1 5 10 15

Asn Ile Pro Ala Ile Gln Asn Leu Ala Lys Glu Leu Ala Thr Ser His
20 25 30

Lys Pro Arg
35

<210> 693
<211> 35
<212> PRT
<213> Artificial Sequence

<220>
<223> Chironomus thummi thummi Globin CTT-VIII epitope

<400> 693

Ile Val Gly Phe Phe Ser Glu Val Ile Gly Leu Ile Gly Asn Pro Glu
1 5 10 15

Asn Arg Pro Ala Leu Lys Thr Leu Ile Asp Gly Leu Ala Ser Ser His
20 25 30

Lys Ala Arg
35

<210> 694
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Glucan endo-1,3-beta-glucosidase, basic vacuolar isoform epitope

<400> 694

Ala Trp Leu Ala Gln Phe Val Leu Pro
1 5

<210> 695
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glutamate decarboxylase epitope

<400> 695

Phe Arg Glu Arg Gln Ser Ser Lys Asn Leu Leu Ser Cys Glu Asn Ser
1 5 10 15

Asp Arg Asp Ala
20

<210> 696
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 1 epitope

<400> 696

Met Ala Ser Ser Thr Pro Ser Ser Ser Ala Thr Ser Ser Asn Ala Gly
1 5 10 15

Ala Asp Pro Asn
20

<210> 697
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Glutamate decarboxylase 2 epitope

<400> 697

Pro Gly Ser Gly Phe Trp Ser Phe Gly Ser Glu Asp Gly Ser Gly Asp
1 5 10 15

Ser Glu Asn

<210> 698
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glutamate receptor, ionotropic, N-methyl D-aspartate 2A epitope

<400> 698

Ser Val Ser Tyr Asp Asp Trp Asp Tyr Ser Leu Glu Ala Arg Val
1 5 10 15

<210> 699
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens glutathione peroxidase-GI epitope

<400> 699

Asn Glu His Pro Val Phe Ala Tyr Leu Lys Asp Lys Leu Pro
1 5 10

<210> 700
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Glutenin, high molecular weight subunit DX5 epitope

<400> 700

Ala Gln Gly Gln Gln Pro Gly Gln Gly Gln Gln Gly Gln Gln
1 5 10

<210> 701
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Glutenin, high molecular weight subunit DX5 precursor epitope

<400> 701

Gln Gln Pro Gly Gln
1 5

<210> 702
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Glutenin, low molecular weight subunit precursor epitope

<400> 702

Gln Gln Gln Pro Pro
1 5

<210> 703
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phaseolus vulgaris Glycine-rich cell wall structural protein 1.8 precursor epitope

<400> 703

Gly Gly Tyr Gly Asp Gly Gly Ala His Gly Gly Gly Tyr Gly Gly
1 5 10 15

<210> 704
<211> 15
<212> PRT

<213> Artificial Sequence

<220>
<223> Arachis hypogaea Glycinin epitope

<400> 704

Ala Leu Ser Arg Leu Val Leu Arg Arg Asn Ala Leu Arg Arg Pro
1 5 10 15

<210> 705
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max Glycinin G1 precursor epitope

<400> 705

Gly Ala Ile Val Thr Val Lys Gly Gly Leu Ser Val Ile
1 5 10

<210> 706
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max Glycinin G2 precursor epitope

<400> 706

Ala Leu Ser Arg Cys Thr Leu Asn Arg Asn Ala Leu Arg Arg Pro
1 5 10 15

<210> 707
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Holcus lanatus group V allergen epitope

<400> 707

Ala Asn Val Pro Pro Ala Asp Lys Tyr Lys Thr Phe Glu Ala Ala
1 5 10 15

<210> 708
<211> 38
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Gu protein epitope

<400> 708

Ile Asp Ala Pro Lys Pro Lys Lys Met Lys Lys Glu Lys Glu Met Asn
1 5 10 15

Gly Glu Thr Arg Glu Lys Ser Pro Lys Leu Lys Asn Gly Phe Pro His

20 25 30

Pro Glu Pro Asp Cys Asn
35

<210> 709
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens H1 histone family, member 0 epitope

<400> 709

Lys Glu Ile Lys Lys Val Ala Thr Pro Lys Lys Ala Ser Lys Pro Lys
1 5 10 15

Lys

<210> 710
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens heat shock 60kDa protein 1 (chaperonin) epitope

<400> 710

Ala Tyr Ala Lys Asp Val Lys Phe Gly Ala Asp Ala
1 5 10

<210> 711
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Heat shock protein HSP 90-beta epitope

<400> 711

Gly Leu Glu Leu Pro Glu
1 5

<210> 712
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens high mobility group protein 17 epitope

<400> 712

Lys Lys Ala Pro Ala Lys Lys Gly Glu Lys Val Pro Lys Gly Lys
1 5 10 15

<210> 713

<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens High mobility group protein B1 epitope

<400> 713

Ala Lys Gly Lys Pro Asp Ala Ala Lys Lys Gly Val Val Lys Ala Glu
1 5 10 15

Lys Ser Lys Lys Lys Lys
20

<210> 714
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens high-mobility group box 2 epitope

<400> 714

Phe Glu Asp Met Ala Lys Ser Asp Lys Ala Arg Tyr Asp Arg Glu
1 5 10 15

<210> 715
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens histidyl-tRNA synthetase, cytoplasmic epitope

<400> 715

Ala Glu Arg Ala Ala Leu Glu Glu Leu Val Lys Leu Gln Gly Glu Arg
1 5 10 15

Val Arg Gly Leu Lys Gln Gln Lys Ala Ser Ala Glu Leu Ile Glu Glu
20 25 30

Glu Val Ala Lys Leu Leu Lys Leu Lys Ala Gln
35 40

<210> 716
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Histone H1.4 epitope

<400> 716

Ser Glu Thr Ala Pro Ala Ala Pro Ala Ala Pro Ala Pro Ala Glu Lys
1 5 10 15

<210> 717

<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens histone H1b epitope

<400> 717

Lys Pro Lys Ala Ala Lys Pro Lys Lys Ala Ala Ala Lys Lys Lys
1 5 10 15

<210> 718
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Histone H2A.Z epitope

<400> 718

Gly Lys Ala Lys Thr Lys Ala Val Ser Arg Ser Gln Arg Ala Gly Leu
1 5 10 15

Gln Phe Pro Val
20

<210> 719
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens histone H3 epitope

<400> 719

Leu Pro Phe Gln Arg Leu Val Arg Glu Ile Ala Gln Asp Phe Lys
1 5 10 15

<210> 720
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Histone H3-like centromeric protein A epitope

<400> 720

Lys Pro Glu Ala Pro Arg Arg Arg Ser Pro
1 5 10

<210> 721
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA class I histocompatibility antigen, B-27 alpha chain precursor epitope

<400> 721

Lys Ala Lys Ala Gln Thr Asp Arg
1 5

<210> 722
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA-B27 epitope

<400> 722

Ala Lys Ala Gln Thr Asp Arg Glu Asp Leu Arg Thr Leu Leu Arg Tyr
1 5 10 15

<210> 723
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HLA-DR3 epitope

<400> 723

Arg Pro Asp Ala Glu Tyr Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln
1 5 10 15

Lys Arg Gly Arg
20

<210> 724
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HMG-17 epitope

<400> 724

Asp Gly Lys Ala Lys Val Lys Asp Glu Pro Gln Arg Arg Ser Ala
1 5 10 15

<210> 725
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens HNRNPA2B1 protein epitope

<400> 725

Glu Thr Thr Glu Glu Ser Leu Arg Asn Tyr Tyr Glu Gln
1 5 10

<210> 726
<211> 35

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens hypothetical protein epitope

<400> 726

Ala Asn Glu Asp Ala Ala Gln Gly Ile Ala Asn Trp Asp Ala Val Gln
1 5 10 15

Asp Ile Ala Asn Glu Asp Gly Phe His Gly Ile Asp Ile Glu Asp Ala
20 25 30

Ala Gln Gly
35

<210> 727
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Oryza sativa Japonica Group hypothetical protein epitope

<400> 727

Ala Phe Asn His Phe Gly Ile Gln Leu Val Gln Arg
1 5 10

<210> 728
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ig alpha-1 chain C region epitope

<400> 728

Pro Val Pro Ser Thr Pro Pro Thr Pro Ser Pro Ser Thr Pro Pro Thr
1 5 10 15

Pro Ser Pro Ser
20

<210> 729
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ig gamma-1 chain C region epitope

<400> 729

Lys Phe Asn Trp Tyr Val Asp
1 5

<210> 730
<211> 7

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ig gamma-3 chain C region epitope

<400> 730

Asp Gly Ser Phe Phe Leu Tyr
1 5

<210> 731
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ig heavy chain V-III region (ART) epitope

<400> 731

Cys Ser Val Met His Glu Gly
1 5

<210> 732
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ig lambda chain V-II region MGC epitope

<400> 732

Ser Gly Leu Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Ser Ser Tyr
1 5 10 15

<210> 733
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ig L-chain V-region epitope

<400> 733

Ala Pro Ser Val Thr Leu Phe
1 5

<210> 734
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Immunoglobulin heavy chain epitope

<400> 734

Asp Lys Ser Arg Trp Gln Glu
1 5

<210> 735
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens immunoglobulin light chain epitope

<400> 735

Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly Ala Val
1 5 10 15

<210> 736
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens immunoglobulin light chain variable region epitope

<400> 736

Ala Gly Glu Lys Val Thr Met
1 5

<210> 737
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Insulin precursor epitope

<400> 737

Thr Ser Ile
1

<210> 738
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Integrin alpha-6 epitope

<400> 738

Leu Lys Arg Asp Met Lys Ser Ala His Leu Leu Pro Glu His
1 5 10

<210> 739
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Integrin beta-3 precursor epitope

<400> 739

Cys Ala Pro Glu Ser Ile Glu Phe Pro Val Ser Glu Ala Arg Val Leu
1 5 10 15

Glu Asp

<210> 740
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens interferon alpha 2 epitope

<400> 740

Cys Asp Leu Pro Gln Thr His Ser Leu Gly Ser Arg Arg Thr
1 5 10

<210> 741
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens interferon alpha A epitope

<400> 741

Glu Asp Ser Ile Leu Ala Val Arg Lys Tyr Phe Gln Arg Ile
1 5 10

<210> 742
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens interferon beta precursor epitope

<400> 742

His Leu Lys Arg Tyr Tyr Gly Arg Ile Leu His Tyr
1 5 10

<210> 743
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens interferon-alpha 2 epitope

<400> 743

Leu Met Leu Leu Ala Gln Met Arg Arg Ile Ser Leu Phe Ser
1 5 10

<210> 744
<211> 37
<212> PRT
<213> Artificial Sequence

<220>

<223> Homo sapiens Islet amyloid polypeptide precursor epitope

<400> 744

Met Gly Ile Leu Lys Leu Gln Val Phe Leu Ile Val Leu Ser Val Ala
1 5 10 15

Leu Asn His Leu Lys Ala Thr Pro Ile Glu Ser His Gln Val Glu Lys
20 25 30

Arg Lys Cys Asn Thr
35

<210> 745
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Kappa-casein precursor epitope

<400> 745

Ala Lys Tyr Ile Pro Ile Gln Tyr Val Leu Ser Arg Tyr Pro
1 5 10

<210> 746
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ku antigen epitope

<400> 746

Arg Gly Asp Gly Pro Phe Arg Leu Gly Gly
1 5 10

<210> 747
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens leukotriene B4 receptor 2 epitope

<400> 747

Gly Arg Gly Asn Gly Asp Pro Gly Gly Gly Met Glu Lys Asp Gly
1 5 10 15

<210> 748
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens liver histone H1e epitope

<400> 748

Ile Lys Lys Val Ala Thr Pro Lys Lys Ala Ser Pro Lys Lys
1 5 10

<210> 749
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Lupus La protein epitope

<400> 749

Ala Gln Pro Gly Ser Gly Lys Gly Lys Val Gln Phe Gln Gly Lys Lys
1 5 10 15

Thr Lys Phe Ala Ser Asp Asp
20

<210> 750
<211> 30
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens lymphocyte activation gene 3 protein precursor epitope

<400> 750

Gly Pro Pro Ala Ala Ala Pro Gly His Pro Leu Ala Pro Gly Pro His
1 5 10 15

Pro Ala Ala Pro Ser Ser Trp Gly Pro Arg Pro Arg Arg Tyr
20 25 30

<210> 751
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens m3 muscarinic cholinergic receptor epitope

<400> 751

Glu Pro Thr Ile Thr Phe Gly Thr Ala Ile
1 5 10

<210> 752
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Alternaria alternata Major allergen Alt a 1 precursor epitope

<400> 752

Ala Asp Pro Val Thr Thr Glu Gly Asp Tyr Val Val Lys Ile Ser Glu
1 5 10 15

Phe Tyr Gly Arg
20

<210> 753
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Anisakis simplex Major allergen Ani s 1 epitope

<400> 753

Cys Lys Met Pro Asp Arg Gly Ala Cys Ala Leu Gly Lys Lys Pro
1 5 10 15

<210> 754
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Major allergen Asp f 1 epitope

<400> 754

Leu Asn Pro Lys Thr Asn Lys Trp Glu Asp Lys Arg Tyr
1 5 10

<210> 755
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Major allergen Asp f 2 epitope

<400> 755

Ala His Ile Leu Arg Trp Gly Asn Glu Ser
1 5 10

<210> 756
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus major allergen beta-lactoglobulin epitope

<400> 756

Leu Gln Lys Trp Glu Asn Asp Glu Cys Ala Gln Lys Lys Ile Ile Ala
1 5 10 15

Glu Lys Thr Lys
20

<210> 757
<211> 14
<212> PRT

<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 1 precursor epitope

<400> 757

Asp Ala Lys Met Thr Glu Glu Asp Lys Glu Asn Ala Leu Ser
1 5 10

<210> 758
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus Major allergen I polypeptide chain 2 precursor epitope

<400> 758

Glu Pro Glu Arg Thr Ala Met Lys Lys Ile Gln Asp Cys Tyr
1 5 10

<210> 759
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Felis catus major allergen I, polypeptide chain 1 epitope

<400> 759

Leu Leu Asp Lys Ile Tyr Thr Ser Pro Leu Cys
1 5 10

<210> 760
<211> 29
<212> PRT
<213> Artificial Sequence

<220>
<223> Turbo cornutus major allergen Tur c1 - Turbo cornutus epitope

<400> 760

Leu Glu Asp Glu Leu Leu Ala Glu Lys Glu Lys Tyr Lys Ala Ile Ser
1 5 10 15

Asp Glu Leu Asp Gln Thr Phe Ala Glu Leu Ala Gly Tyr
20 25

<210> 761
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus major house dust allergen epitope

<400> 761

Leu Ala His Arg Asn Gln Ser Leu Asp Leu Ala Glu Gln Glu Leu Val
1 5 10 15

Asp Cys Ala Ser Gln His Gly Cys His
20 25

<210> 762
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Major latex allergen Hev b 5 epitope

<400> 762

Ala Pro Pro Ala Ser Glu Gln Glu Thr
1 5

<210> 763
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Major mite fecal allergen Der p 1 epitope

<400> 763

Ala Arg Glu Gln Ser Cys Arg Arg Pro Asn Ala Gln Arg Phe Gly Ile
1 5 10 15

Ser Asn Tyr Cys Gln Ile Tyr Pro Pro Asn Ala Asn Lys Ile Arg Glu
20 25 30

Ala Leu Ala Gln Pro Gln Arg Tyr Cys Arg His
35 40

<210> 764
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Major pollen allergen epitope

<400> 764

Phe Thr Glu Val Gly Tyr Thr Arg Ala Glu Gly Leu
1 5 10

<210> 765
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Major pollen allergen Bet v 1-A epitope

<400> 765

Asp Gly Asp Asn Leu Phe Pro Lys Val Ala
1 5 10

<210> 766
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Chamaecyparis obtusa Major pollen allergen Cha o 1 precursor epitope

<400> 766

Trp Arg Ser Thr Gln Asp Ser Phe Asn Asn Gly
1 5 10

<210> 767
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Corylus avellana Major pollen allergen Cor a 1 epitope

<400> 767

Tyr Val Leu Asp Gly Asp Lys Leu Leu Pro Lys Val Ala Pro Gln Ala
1 5 10 15

Leu

<210> 768
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> Holcus lanatus Major pollen allergen Hol l 1 precursor epitope

<400> 768

Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly Pro Lys Asp
1 5 10 15

Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp
20 25

<210> 769
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Juniperus ashei Major pollen allergen Jun a 1 precursor epitope

<400> 769

Ala Phe Asn Gln Phe Gly Pro Asn Ala Gly Gln Arg

1 5 10

<210> 770
<211> 34
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea major pollen allergen Ole e 1 epitope

<400> 770

Ser Gly Arg Lys Asp Cys Asn Glu Ile Pro Thr Glu Gly Trp Val Lys
1 5 10 15

Pro Ser Leu Lys Phe Ile Leu Asn Thr Val Asn Gly Thr Thr Arg Thr
20 25 30

Val Asn

<210> 771
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Malus x domestica mal d 3 epitope

<400> 771

Arg Thr Thr Ala Asp Arg Gln Thr Ala
1 5

<210> 772
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MBP protein epitope

<400> 772

Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
1 5 10 15

<210> 773
<211> 30
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens melanin-concentrating hormone receptor 1, isoform CRA_a epitope

<400> 773

Ala Glu His Ala Ser Arg Met Ser Val Leu Arg Ala Lys Pro Met Ser
1 5 10 15

Asn Ser Gln Arg Leu Leu Leu Leu Ser Pro Gly Ser Pro Pro
20 25 30

<210> 774
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Melanocyte protein Pmel 17 precursor epitope

<400> 774

Gln Val Pro Thr Thr Glu Val Val Gly Thr Thr Pro Gly Gln Ala Pro
1 5 10 15

<210> 775
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MHC classII HLA-DRB1 epitope

<400> 775

Glu Gln Arg Arg Ala Ala
1 5

<210> 776
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens MHC HLA-DR1-beta epitope

<400> 776

Arg Pro Asp Ala Glu Tyr Trp Asn Ser Gln Lys Asp Leu Leu Glu Gln
1 5 10 15

Arg Arg Ala Ala
20

<210> 777
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Blomia tropicalis Mite allergen Blo t 5 epitope

<400> 777

Glu Glu Ala Gln Thr Leu Ser Lys Ile Leu Leu Lys Asp Leu Lys Glu
1 5 10 15

<210> 778
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides farinae Mite group 2 allergen Der f 2 precursor epitope

<400> 778

Asp Pro Cys Ile Ile
1 5

<210> 779
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Mite group 2 allergen Der p 2 precursor epitope

<400> 779

Asp Gln Val Asp Val Lys Asp Cys Ala Asn His Glu Ile Lys Lys
1 5 10 15

<210> 780
<211> 33
<212> PRT
<213> Artificial Sequence

<220>
<223> Lepidoglyphus destructor Mite group 2 allergen Lep d 2 precursor epitope

<400> 780

Ala Ala Asn Gln Asp Thr Ala Lys Val Thr Ile Lys Val Leu Ala Lys
1 5 10 15

Val Ala Gly Thr Thr Ile Gln Val Pro Gly Leu Glu Thr Asp Gly Cys
20 25 30

Lys

<210> 781
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum monomeric alpha-amylase inhibitor epitope

<400> 781

Ala Ala Ser Val Pro Glu
1 5

<210> 782
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Muscarinic acetylcholine receptor M1 epitope

<400> 782

Gln Tyr Leu Val Gly Glu Arg Thr Val Leu Ala Gly Gln Cys Tyr Ile
1 5 10 15

Gln Phe Leu Ser Gln
20

<210> 783
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin associated glycoprotein epitope

<400> 783

Asp Ser Tyr Thr Leu Thr Glu Glu Leu Ala Tyr Ala Glu Ile Arg Val
1 5 10 15

Lys

<210> 784
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin basic protein epitope

<400> 784

Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys
1 5 10

<210> 785
<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin oligodendrocyte glycoprotein epitope

<400> 785

Ala Leu Val Gly Asp Glu Val Glu Leu Pro Cys Arg Ile Ser Pro Gly
1 5 10 15

Lys Asn Ala Thr Gly Met Glu Leu Gly Trp
20 25

<210> 786
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin oligodendrocyte glycoprotein isoform alpha6 precursor epitope

<400> 786

His Arg Thr Phe Glu
1 5

<210> 787
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens myelin proteolipid protein epitope

<400> 787

Ala Asp Ala Arg Met
1 5

<210> 788
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin-associated glycoprotein precursor epitope

<400> 788

Gly His Trp Gly Ala Trp Met Pro Ser Ser Ile Ser Ala Phe Glu Gly
1 5 10 15

Thr Cys Val Ser Ile
20

<210> 789
<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myelin-oligodendrocyte glycoprotein precursor epitope

<400> 789

Gly Gln Phe Arg Val Ile Gly Pro Arg His Pro Ile Arg Ala Leu Val
1 5 10 15

Gly Asp Glu Val Glu Leu Pro Cys Arg Ile
20 25

<210> 790
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myeloblastin precursor epitope

<400> 790

Ala His Arg Pro Pro Ser Pro Ala
1 5

<210> 791
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myeloperoxidase epitope

<400> 791

Gly Ser Ala Ser Pro Met Glu Leu Leu Ser
1 5 10

<210> 792
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Myosin-11 epitope

<400> 792

Ala Leu Lys Thr Glu Leu Glu Asp Thr Leu Asp Ser Thr Ala Thr Gln
1 5 10 15

Gln Glu Leu Arg
20

<210> 793
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Neurofilament heavy polypeptide (NF-H) (Neurofilament triplet H protein) (200 kDa neurofilament protein) epitope

<400> 793

Ala Lys Ser Pro Glu Lys Ala Lys
1 5

<210> 794
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens nicotinic acetylcholine receptor alpha subunit|AChR alpha subunit epitope

<400> 794

Glu Val Asn Gln Ile Val Thr Thr Asn Val Arg Leu Lys Gln Gln Trp
1 5 10 15

<210> 795
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Non-histone chromosomal protein HMG-17 epitope

<400> 795

Val Lys Asp Glu Pro Gln Arg Arg Ser Ala
1 5 10

<210> 796
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus armeniaca Non-specific lipid-transfer protein 1 epitope

<400> 796

Val Asn Pro Asn Asn Ala Ala Ala Leu
1 5

<210> 797
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus armeniaca Non-specific lipid-transfer protein 1 (LTP 1)
(Major allergen Pru ar 3) epitope

<400> 797

Leu Ala Arg Thr Thr Pro Asp Arg Arg Thr Ala Cys Asn Cys Leu
1 5 10 15

<210> 798
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus domestica Non-specific lipid-transfer protein 1 (LTP 1)
(Major allergen Pru d 3) epitope

<400> 798

Leu Ala Arg Thr Thr Ala Asp Arg Arg Ala Ala Cys Asn Cys Leu Lys
1 5 10 15

Gln Leu

<210> 799
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Malus x domestica Non-specific lipid-transfer protein precursor (LTP) (Allergen Mal d 3) epitope

<400> 799

Ala Asp Arg Gln Thr Ala Cys Asn Cys Leu Lys Asn Leu Ala Gly
1 5 10 15

<210> 800
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens NR2 subunit NMDA receptor epitope

<400> 800

Asp Trp Glu Tyr Ser Val Trp Leu Ser Asn
1 5 10

<210> 801
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens nuclear autoantigen Sp-100 isoform 1 epitope

<400> 801

Glu Val Phe Ile Ser Ala Pro Arg
1 5

<210> 802
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Ole e 1 protein epitope

<400> 802

Glu Asp Val Pro Gln Pro Pro Val Ser Gln Phe His
1 5 10

<210> 803
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Olea europaea Ole e 1.0102 protein epitope

<400> 803

Glu Asp Val Pro Gln Pro Pro Val Ser Gln Phe His Ile Gln Gly Gln
1 5 10 15

Val Tyr Cys Asp Thr Cys Arg Ala Gly

20 25

<210> 804
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Omega gliadin storage protein epitope

<400> 804

Gln Gln Pro Gln Gln Ser Phe Pro Gln Gln
1 5 10

<210> 805
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum omega-5 gliadin epitope

<400> 805

Gln Gln Phe His Gln Gln Gln
1 5

<210> 806
<211> 35
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Oryzin precursor epitope

<400> 806

Ala Ser Asn Thr Ser Pro Ala Ser Ala Pro Asn Ala Leu Thr Val Ala
1 5 10 15

Ala Ile Asn Lys Ser Asn Ala Arg Ala Ser Phe Ser Asn Tyr Gly Ser
20 25 30

Val Val Asp
35

<210> 807
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovalbumin epitope

<400> 807

Cys Phe Asp Val Phe Lys Glu Leu Lys
1 5

<210> 808

<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid epitope

<400> 808

Cys Asn Phe Cys Asn Ala Val Val Glu Ser
1 5 10

<210> 809
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus Ovomucoid precursor epitope

<400> 809

Ala Glu Val Asp Cys Ser Arg Phe Pro Asn Ala Thr Asp Lys
1 5 10

<210> 810
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max P34 probable thiol protease precursor epitope

<400> 810

Ala Ser Trp Asp Trp Arg Lys Lys Gly Val
1 5 10

<210> 811
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max P34 probable thiol protease precursor; Gly m 1
epitope

<400> 811

Pro Gln Glu Phe Ser Lys Lys Thr Tyr Gln
1 5 10

<210> 812
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens p70 autoantigen epitope

<400> 812

Glu Ala Leu Thr Lys His Phe Gln Asp
1 5

<210> 813
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens PADI-H protein epitope

<400> 813

Lys Ala Ala Ser Gly Ser Thr Gly Asp Gln Lys Val Gln Ile Ser Tyr
1 5 10 15

Tyr Gly Pro Lys
20

<210> 814
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Parietaria judaica Par j epitope

<400> 814

Gly Thr Ser Ser Cys Arg Leu Val Pro
1 5

<210> 815
<211> 47
<212> PRT
<213> Artificial Sequence

<220>
<223> Blomia tropicalis Paramyosin epitope

<400> 815

Glu Lys Leu Arg Asp Gln Lys Glu Ala Leu Ala Arg Glu Asn Lys Lys
1 5 10 15

Leu Ala Asp Asp Leu Ala Glu Ala Lys Ser Gln Leu Asn Asp Ala His
20 25 30

Arg Arg Ile His Glu Gln Glu Ile Glu Ile Lys Arg Leu Glu Asn
35 40 45

<210> 816
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Gadus morhua callarias Parvalbumin beta epitope

<400> 816

Ala Ala Glu Ala Ala Cys Phe Lys
1 5

<210> 817
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Salmo salar parvalbumin like 1 epitope

<400> 817

Ala Asp Ile Lys Thr Ala Leu Glu Ala Arg Lys Ala Ala Asp Thr
1 5 10 15

<210> 818
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Juniperus ashei Pathogenesis-related protein precursor epitope

<400> 818

Ala Asp Ile Asn Ala Val Cys Pro Ser Glu Leu Lys
1 5 10

<210> 819
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Nicotiana tabacum Pectate lyase epitope

<400> 819

Ala Tyr Asn His Phe Gly Lys Arg Leu Asp Gln Arg
1 5 10

<210> 820
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Musa acuminata AAA Group pectate lyase 2 epitope

<400> 820

Ala Phe Asn His Phe Gly Glu Gly Leu Ile Gln Arg
1 5 10

<210> 821
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Farfantepenaeus aztecus Pen a 1 allergen epitope

<400> 821

Ala Asn Ile Gln Leu Val Glu Lys Asp Lys Ala Leu Ser Asn Ala
1 5 10 15

<210> 822
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Dermatophagoides pteronyssinus Peptidase 1 precursor (Major mite fecal allergen Der p 1) (Allergen Der p I) epitope

<400> 822

Ala Arg Glu Gln Ser Cys Arg Arg Pro Asn Ala Gln Arg Phe Gly Ile
1 5 10 15

Ser Asn Tyr Cys Gln Ile Tyr Pro Pro Asn Val Asn Lys Ile Arg Glu
20 25 30

Ala Leu Ala Gln Thr His Ser Ala Ile Ala Val
35 40

<210> 823
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens pericentriolar material 1 protein epitope

<400> 823

Lys Asp Cys
1

<210> 824
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Periplakin epitope

<400> 824

Ile His Asp Arg Lys Ser Gly Lys Lys Phe Ser Ile Glu Glu Ala Leu
1 5 10 15

Gln Ser Gly Arg
20

<210> 825
<211> 45
<212> PRT
<213> Artificial Sequence

<220>
<223> Apis mellifera Phospholipase A2 precursor epitope

<400> 825

Leu Ile Asp Thr Lys Cys Tyr Lys Leu Glu His Pro Val Thr Gly Cys
1 5 10 15

Gly Glu Arg Thr Glu Gly Arg Cys Leu His Tyr Thr Val Asp Lys Ser
20 25 30

Lys Pro Lys Val Tyr Gln Trp Phe Asp Leu Arg Lys Tyr
35 40 45

<210> 826
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Myrmecia pilosula Pilosulin-1 precursor (Major allergen Myr p 1) (Myr p I) epitope

<400> 826

Lys Glu Ala Ile Pro Met Ala Val Glu Met Ala Lys Ser Gln
1 5 10

<210> 827
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens plasma protease C1 inhibitor precursor epitope

<400> 827

Ala Ser Ala Ile Ser Val Ala Arg
1 5

<210> 828
<211> 6
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens platelet glycoprotein IIIa epitope

<400> 828

Arg Ala Arg Ala Lys Trp
1 5

<210> 829
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens plexin domain containing 1, isoform CRA_b epitope

<400> 829

Asn Cys Ser Trp Cys His Val Leu Gln Arg Cys Ser
1 5 10

<210> 830
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens PM/Scl 100kD nucleolar protein epitope

<400> 830

Cys Ile Ala Ala Lys Lys Ile Lys Gln Ser Val Gly Asn Lys Ser
1 5 10 15

<210> 831
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula Polcalcin Bet v 4 epitope

<400> 831

Phe Gly Arg Ala Asn Arg Gly Leu
1 5

<210> 832
<211> 36
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Polcalcin Phl p 7 (Calcium-binding pollen allergen Phl p 7) (P7) epitope

<400> 832

Ala Asp Asp Met Glu Arg Ile Phe Lys Arg Phe Asp Thr Asn Gly Asp
1 5 10 15

Gly Lys Ile Ser Leu Ser Glu Leu Thr Asp Ala Leu Arg Thr Leu Gly
20 25 30

Ser Thr Ser Ala
35

<210> 833
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne pollen allergen epitope

<400> 833

Glu Gly Gly Thr Lys Ser Glu Val Glu Asp Val Ile Pro Glu Gly Trp
1 5 10 15

Lys Ala Asp Thr Ser Tyr Ser Ala Lys

20 25

<210> 834
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia Pollen allergen Amb a 1.4 epitope

<400> 834

Ala Phe Asn Lys Phe Thr Asp Asn Val Asp Gln Arg
1 5 10

<210> 835
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia Pollen allergen Amb a 2 precursor epitope

<400> 835

Met Pro Arg Cys Arg Phe Gly Phe
1 5

<210> 836
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Ambrosia artemisiifolia var. elatior Pollen allergen Amb a 3 epitope

<400> 836

Cys Asp Ile Lys Asp Pro Ile Arg Leu Glu Pro Gly Gly Pro Asp
1 5 10 15

<210> 837
<211> 31
<212> PRT
<213> Artificial Sequence

<220>
<223> Betula pendula pollen allergen Bet v 1 epitope

<400> 837

Lys Ala Glu Gln Val Lys Ala Ser Lys Glu Met Gly Glu Thr Leu Leu
1 5 10 15

Arg Ala Val Glu Ser Tyr Leu Leu Ala His Ser Asp Ala Tyr Asn
20 25 30

<210> 838
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Poa pratensis Pollen allergen KBG 60 precursor epitope

<400> 838

Ala Ala Asn Lys Tyr Lys Thr Phe Val Ala Thr Phe Gly Ala Ala Ser
1 5 10 15

Asn Lys Ala Phe
20

<210> 839
<211> 25
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 2-A (Lol p II-A) epitope

<400> 839

Glu Lys Gly Met Arg Asn Val Phe Asp Asp Val Val Pro Ala Asp Phe
1 5 10 15

Lys Val Gly Thr Thr Tyr Lys Pro Glu
20 25

<210> 840
<211> 27
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p 3 (Lol p III) epitope

<400> 840

Lys Gly Gly Met Lys Asn Val Phe Asp Glu Val Ile Pro Thr Ala Phe
1 5 10 15

Thr Val Gly Lys Thr Tyr Thr Pro Glu Tyr Asn
20 25

<210> 841
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Lolium perenne Pollen allergen Lol p VA precursor epitope

<400> 841

Ala Ala Glu Gly Ala Thr Pro Glu Ala Lys Tyr Asp
1 5 10

<210> 842
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense Pollen allergen Phl p 1 precursor epitope

<400> 842

Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ala Met
1 5 10 15

<210> 843
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Zea mays pollen allergen Phl p 11 epitope

<400> 843

Arg Asp Arg Ala Arg Val Pro Leu
1 5

<210> 844
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Phleum pratense pollen allergen Phl pI epitope

<400> 844

Ile Pro Lys Val Pro Pro Gly Pro Asn Ile Thr Ala
1 5 10

<210> 845
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Polygalacturonase precursor epitope

<400> 845

Gly Gln Cys Lys Trp Val Asn Gly Arg Glu Ile Cys Asn Asp Arg Asp
1 5 10 15

Arg Pro Thr Ala
20

<210> 846
<211> 30
<212> PRT
<213> Artificial Sequence

<220>
<223> Parietaria judaica Probable non-specific lipid-transfer protein epitope

<400> 846

Gln Glu Thr Cys Gly Thr Met Val Arg Ala Leu Met Pro Cys Leu Pro

1 5 10 15

Phe Val Gln Gly Lys Glu Lys Glu Pro Ser Lys Gly Cys Cys
20 25 30

<210> 847
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Parietaria judaica Probable non-specific lipid-transfer protein 2 epitope

<400> 847

Ala Glu Val Pro Lys Lys Cys Asp Ile Lys
1 5 10

<210> 848
<211> 30
<212> PRT
<213> Artificial Sequence

<220>
<223> Parietaria judaica Probable non-specific lipid-transfer protein 2 precursor epitope

<400> 848

Glu Ala Cys Gly Lys Val Val Gln Asp Ile Met Pro Cys Leu His Phe
1 5 10 15

Val Lys Gly Glu Glu Lys Glu Pro Ser Lys Glu Cys Cys Ser
20 25 30

<210> 849
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Solanum lycopersicum Probable pectate lyase P59 epitope

<400> 849

Ala Phe Asn His Phe Gly Lys Arg Leu Ile Gln Arg
1 5 10

<210> 850
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens profilaggrin epitope

<400> 850

Gly Gly Gln Gly Ser Arg His Gln Gln Ala Arg
1 5 10

<210> 851
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Cucumis melo profilin epitope

<400> 851

Ala Phe Arg Leu Glu Glu Ile Ala Ala Ile
1 5 10

<210> 852
<211> 56
<212> PRT
<213> Artificial Sequence

<220>
<223> Glycine max Profilin-1 epitope

<400> 852

Trp Ala Gln Ser Thr Asp Phe Pro Gln Phe Lys Pro Glu Glu Ile Thr
1 5 10 15

Ala Ile Met Asn Asp Phe Asn Glu Pro Gly Ser Leu Ala Pro Thr Gly
20 25 30

Leu Tyr Leu Gly Gly Thr Lys Tyr Met Val Ile Gln Gly Glu Pro Gly
35 40 45

Ala Val Ile Arg Gly Lys Lys Gly
50 55

<210> 853
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Pro-hevein precursor epitope

<400> 853

Glu Gln Cys Gly Arg Gln Ala Gly Gly Lys Leu Cys Pro Asn Asn Leu
1 5 10 15

Cys Cys Ser Gln Trp Gly Trp Cys Gly Ser Thr Asp Glu Tyr Cys Ser
20 25 30

Pro Asp His Asn Cys Gln Ser Asn Cys Lys Asp
35 40

<210> 854
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Proliferating cell nuclear antigen epitope

<400> 854

Leu Lys Tyr Tyr Leu Ala Pro Lys Ile Glu Asp Glu Glu Gly Ser
1 5 10 15

<210> 855
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Proline-rich transmembrane protein 2 epitope

<400> 855

His Ser Glu Ala Glu Thr Gly Pro Pro
1 5

<210> 856
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proteasome (prosome, macropain) activator subunit 3 (PA28 gamma; Ki), isoform CRA_a epitope

<400> 856

Leu Asp Gly Pro Thr Tyr Lys Arg Arg Leu Asp Glu Cys Glu Glu
1 5 10 15

<210> 857
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens protein tyrosine phosphatase-like autoantigen epitope

<400> 857

Gly Ala His Gly Asp Thr Thr Pro Glu Tyr Gln Asp Leu
1 5 10

<210> 858
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens protein-arginine deiminase type-4 epitope

<400> 858

Ala Phe Phe Pro Asn Met Val Asn Met Leu Val Leu Gly Lys His Leu
1 5 10 15

Gly Ile Pro Lys

<210> 859
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens proteinase 3 epitope

<400> 859

Cys Ala Thr Arg Leu Phe Pro Asp Phe Phe Thr Arg Val Ala Leu
1 5 10 15

<210> 860
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus persica pru p 1 epitope

<400> 860

Gly Lys Cys Gly Val Ser Ile Pro Tyr Lys
1 5 10

<210> 861
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus dulcis prunin 1 precursor epitope

<400> 861

Glu Glu Ser Gln Gln Ser Ser Gln Gln Gly Arg Gln Gln Glu Gln
1 5 10 15

<210> 862
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Prunus dulcis prunin 2 precursor epitope

<400> 862

Asp Ser Gln Pro Gln Gln Phe Gln Gln Gln Gln Gln Gln Gln Gln
1 5 10 15

<210> 863
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Hesperocyparis arizonica putative allergen Cup a 1 epitope

<400> 863

Trp Arg Phe Thr Arg Asp Ala Phe Thr Asn Gly
1 5 10

<210> 864
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Putative HTLV-1-related endogenous sequence (p25) epitope

<400> 864

Pro Thr Arg Ala Pro Ser Gly Pro Arg Pro Pro
1 5 10

<210> 865
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Putative small nuclear ribonucleoprotein polypeptide E-like protein 1 epitope

<400> 865

Glu Ile His Ser Lys Thr Lys Ser
1 5

<210> 866
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor tyrosine-protein kinase erbB-2 precursor epitope

<400> 866

Pro Glu Ser Phe Asp Gly Asp Pro Ala Ser Asn Thr Ala Pro Leu Gln
1 5 10 15

Pro Glu

<210> 867
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Receptor-type tyrosine-protein phosphatase-like N precursor epitope

<400> 867

Lys Glu Arg Leu Ala Ala Leu Gly Pro Glu Gly Ala His
1 5 10

<210> 868
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens recombinant IgG2 heavy chain epitope

<400> 868

Glu Pro Gln Val Val Thr Leu Pro Pro Ser Arg
1 5 10

<210> 869
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Replication protein A 32 kDa subunit epitope

<400> 869

Arg Ser Phe Gln Asn Lys Lys Ser Leu Val Ala Phe Lys Ile Met Pro
1 5 10 15

Leu Glu Asp Met
20

<210> 870
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Ribonuclease mitogillin precursor epitope

<400> 870

Phe Pro Thr Phe Pro Asp Gly His Asp Tyr
1 5 10

<210> 871
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens ribosomal protein L7 epitope

<400> 871

Glu Leu Lys Ile Lys Arg Leu Arg Lys Lys Phe Ala Gln Lys Met Leu
1 5 10 15

Arg Lys Ala Arg Arg Lys Leu
20

<210> 872
<211> 14
<212> PRT

<213> Artificial Sequence

<220>
<223> Homo sapiens ribosomal protein P2 epitope

<400> 872

Ser Glu Glu Ser Asp Asp Asp Met Gly Phe Gly Leu Phe Asp
1 5 10

<210> 873
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Mangifera indica ripening-related pectate lyase epitope

<400> 873

Ala Tyr Asn His Phe Gly Glu Gly Leu Ile Gln Arg
1 5 10

<210> 874
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens RNA binding protein, autoantigenic (hnRNP-associated with lethal yellow homolog (mouse)), isoform CRA_c epitope

<400> 874

Gly Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
1 5 10 15

Gly Gly Gly Ser Ser
20

<210> 875
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Ro ribonucleoprotein epitope

<400> 875

Asp Gly Tyr Val Trp Gln Val Thr Asp Met Asn Arg Leu His Arg Phe
1 5 10 15

Leu Cys Phe Gly Ser
20

<210> 876
<211> 10
<212> PRT
<213> Artificial Sequence

<220>

<223> Hevea brasiliensis Rubber elongation factor protein epitope

<400> 876

Ala Glu Asp Glu Asp Asn Gln Gln Gly Gln
1 5 10

<210> 877
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens S-arrestin epitope

<400> 877

Phe Leu Gly Glu Leu Thr Ser Ser Glu Val Ala Thr Glu Val
1 5 10

<210> 878
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Juglans regia seed storage protein epitope

<400> 878

Asp Asp Asn Gly Leu Glu Glu Thr Ile Cys Thr Leu Arg Leu Arg
1 5 10 15

<210> 879
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Arachis hypogaea seed storage protein SSP2 epitope

<400> 879

Cys Gly Leu Arg Ala Pro Gln Arg Cys Asp Leu Asp Val Glu Ser
1 5 10 15

<210> 880
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Gallus gallus serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 3 epitope

<400> 880

Arg Pro Asn Ala Thr Tyr Ser Leu
1 5

<210> 881
<211> 8
<212> PRT

<213> Artificial Sequence

<220>
<223> Gallus gallus Serum albumin epitope

<400> 881

Gln Ser Arg Ala Thr Leu Gly Ile
1 5

<210> 882
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> Bos taurus Serum albumin precursor epitope

<400> 882

Asp Asp Ser Pro Asp Leu Pro Lys Leu Lys Pro Asp Pro Asn Thr Leu
1 5 10 15

Cys

<210> 883
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein epitope

<400> 883

Pro Pro Pro Gly Ile Arg Gly Pro
1 5

<210> 884
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein B' epitope

<400> 884

Pro Pro Pro Gly Met Arg Gly Pro
1 5

<210> 885
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein D1 polypeptide epitope

<400> 885

Lys Met Thr Leu Lys Asn Arg Glu Pro Val Gln Leu Glu Thr Leu Ser
1 5 10 15

Ile Arg Gly Asn Arg Ile Arg Tyr
20

<210> 886
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein D2 isoform 1 epitope

<400> 886

Gly Lys Lys Lys Ser Lys Pro Val Asn Lys Asp Arg Tyr Ile Ser Lys
1 5 10 15

Met Phe Leu Arg Gly Asp Ser
20

<210> 887
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein F epitope

<400> 887

Glu Glu Glu Glu Asp Gly Glu Met
1 5

<210> 888
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein G epitope

<400> 888

Trp Ser Lys Ala His Pro Pro Glu
1 5

<210> 889
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein polypeptide A epitope

<400> 889

Ala Met Lys Ile Ser Phe Ala Lys Lys
1 5

<210> 890
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein polypeptide B epitope

<400> 890

Pro Pro Gly Met Arg Pro Pro
1 5

<210> 891
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein polypeptide B/B' isoform B epitope

<400> 891

Met Gly Arg Gly Ala Pro Pro Pro Gly Met Met Gly
1 5 10

<210> 892
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein polypeptide C, isoform CRA_b epitope

<400> 892

Ala Pro Gly Met Arg Pro Pro
1 5

<210> 893
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein polypeptide D3 epitope

<400> 893

Ala Ala Arg Gly Arg Gly Arg Gly Met Gly Arg Gly Asn Ile Phe
1 5 10 15

<210> 894
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens small nuclear ribonucleoprotein polypeptide N

variant epitope

<400> 894

Val Gly Arg Ala Thr Pro Pro Pro Gly Ile Met Ala
1 5 10

<210> 895
<211> 23
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Small nuclear ribonucleoprotein Sm D1 epitope

<400> 895

Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg Gly Arg
1 5 10 15

Gly Arg Gly Gly Pro Arg Arg
20

<210> 896
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Small nuclear ribonucleoprotein Sm D2 epitope

<400> 896

Glu Glu Leu Gln Lys Arg Glu Glu
1 5

<210> 897
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Small nuclear ribonucleoprotein-associated proteins B and B' epitope

<400> 897

Arg Gly Val Gly Gly Pro Ser Gln
1 5

<210> 898
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Hevea brasiliensis Small rubber particle protein epitope

<400> 898

Ala Glu Glu Val Glu Glu Glu Arg Leu Lys
1 5 10

<210> 899
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Smoothelin epitope

<400> 899

Gly Ser Thr Met Met Gln Thr Lys Thr Phe Ser Ser Ser Ser Ser Ser
1 5 10 15

Lys Lys Met Gly
20

<210> 900
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens snRNP polypeptide B epitope

<400> 900

Pro Pro Gly Met Arg Pro Pro Met Gly Pro Met Gly Ile Pro Pro
1 5 10 15

<210> 901
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens spectrin, alpha, non-erythrocytic 1 (alpha-fodrin), isoform CRA_e epitope

<400> 901

Phe Gln Phe Phe Gln Arg Asp Ala Glu Glu Leu Glu Lys Trp
1 5 10

<210> 902
<211> 43
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens steroid 17-alpha-hydroxylase/17,20 lyase epitope

<400> 902

Glu Val Pro Asp Asp Gly Gln Leu Pro Ser Leu Glu Gly Ile Pro Lys
1 5 10 15

Val Val Phe Leu Ile Asp Ser Phe Lys Val Lys Ile Lys Val Arg Gln
20 25 30

Ala Trp Arg Glu Ala Gln Ala Glu Gly Ser Thr
35 40

<210> 903
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Sucrase-isomaltase, intestinal epitope

<400> 903

Asp Phe Thr Tyr Asp Gln Val Ala Phe Asn Gly Leu Pro Gln Phe
1 5 10 15

<210> 904
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Cryptomeria japonica Sugi basic protein precursor epitope

<400> 904

Asp Ala Leu Thr Leu Arg Thr Ala Thr Asn Ile Trp
1 5 10

<210> 905
<211> 48
<212> PRT
<213> Artificial Sequence

<220>
<223> Aspergillus fumigatus Superoxide dismutase epitope

<400> 905

Tyr Thr Leu Pro Pro Leu Pro Tyr Pro Tyr Asp Ala Leu Gln Pro Tyr
1 5 10 15

Ile Ser Gln Gln Ile Met Glu Leu His His Lys Lys His His Gln Thr
20 25 30

Tyr Val Asn Gly Leu Asn Ala Ala Leu Glu Ala Gln Lys Lys Ala Ala
35 40 45

<210> 906
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T cell receptor beta variable 20 epitope

<400> 906

Arg Ser Leu Asp Phe Gln Ala Thr Thr Met Phe
1 5 10

<210> 907
<211> 18

<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T cell receptor beta variable 5 epitope

<400> 907

Ala Leu Gly Gln Gly Pro Gln Phe Ile Phe Gln Tyr Tyr Glu Glu Glu
1 5 10 15

Glu Arg

<210> 908
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Tax1-binding protein 1 epitope

<400> 908

Glu Phe Lys Lys Arg Phe Ser Asp Ala Thr Ser Lys Ala His Gln
1 5 10 15

<210> 909
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T-cell receptor beta chain epitope

<400> 909

Gln Pro Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu
1 5 10 15

<210> 910
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T-cell receptor beta chain C region epitope

<400> 910

Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys Gln Val
1 5 10 15

<210> 911
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T-cell receptor beta chain V region YT35 epitope

<400> 911

Cys Lys Pro Ile Ser Gly His Asn Ser Leu Phe Trp Tyr Arg Gln Thr
1 5 10 15

<210> 912
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens T-cell receptor beta-chain (V1-D-J-C) precursor epitope

<400> 912

Ser Pro Arg Ser Gly Asp Leu Ser Val Tyr
1 5 10

<210> 913
<211> 21
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens TCR V-beta 6.1 epitope

<400> 913

Leu Gly Gln Gly Pro Glu Phe Leu Ile Tyr Phe Gln Gly Thr Gly Ala
1 5 10 15

Ala Asp Asp Ser Gly
20

<210> 914
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens TCR V-beta 6.3 epitope

<400> 914

Asp Pro Ile Ser Gly His Val Ser Leu Phe
1 5 10

<210> 915
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Thyroglobulin epitope

<400> 915

Pro Pro Ala Arg Ala Leu Lys Arg Ser Leu Trp Val Glu Val Asp Leu
1 5 10 15

Leu Ile Gly Ser
20

<210> 916
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Thyroid peroxidase epitope

<400> 916

Gly Leu Pro Arg Leu Glu Thr Pro Ala Asp Leu Ser Thr Ala Ile Ala
1 5 10 15

Ser Arg Ser

<210> 917
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens thyroid stimulating hormone receptor epitope

<400> 917

Glu Ile Ile Gly Phe Gly Gln Glu Leu Lys Asn Pro Gln Glu Glu
1 5 10 15

<210> 918
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens thyroid stimulating hormone receptor variant epitope

<400> 918

Glu Glu Gln Glu Asp Glu Ile Ile Gly Phe Gly Gln Glu Leu Lys Asn
1 5 10 15

<210> 919
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Thyrotropin receptor epitope

<400> 919

Gly Gln Glu Leu Lys Asn Pro Gln Glu Glu Thr Leu Gln Ala Phe Asp
1 5 10 15

Ser His Tyr Asp
20

<210> 920
<211> 15
<212> PRT

<213> Artificial Sequence

<220>
<223> Homo sapiens transaldolase 1 epitope

<400> 920

Ala Ala Ala Gln Met Pro Ala Tyr Gln Glu Leu Val Glu Glu Ala
1 5 10 15

<210> 921
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Trichophyton rubrum Tri r 2 allergen epitope

<400> 921

Asp Cys Asn Gly His Gly Thr His Val Ala Gly Thr Val Gly Gly Thr
1 5 10 15

Lys Tyr Gly Leu
20

<210> 922
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens trinucleotide repeat containing 6A, isoform CRA_b epitope

<400> 922

Ala Phe Leu Ser Val Asp His Leu Gly Gly Gly Gly Glu Ser Met
1 5 10 15

<210> 923
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens trinucleotide repeat containing 6A, isoform CRA_c epitope

<400> 923

Trp Gly Ser Ser Ser Val Gly Pro Gln Ala Leu Ser Lys Ser Gly
1 5 10 15

<210> 924
<211> 36
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens tripartite motif-containing 67 epitope

<400> 924

Leu Gly Gly Gly Ala Gly Gly Gly Gly Asp His Ala Asp Lys Leu Ser
1 5 10 15

Leu Tyr Ser Glu Thr Asp Ser Gly Tyr Gly Ser Tyr Thr Pro Ser Leu
20 25 30

Lys Ser Pro Asn
35

<210> 925
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum Triticum aestivum proteins epitope

<400> 925

Leu Pro Gln Gln Gln Ile Pro Gln Gln Pro
1 5 10

<210> 926
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Penaeus tropomyosin epitope

<400> 926

Phe Leu Ala Glu Glu Ala Asp Arg Lys
1 5

<210> 927
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens TSHR protein epitope

<400> 927

Cys His Gln Glu Glu Asp Phe Arg Val Thr Cys Lys Asp Ile Gln Arg
1 5 10 15

Ile Pro Ser Leu
20

<210> 928
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens tubulin beta-6 chain epitope

<400> 928

Ala Ala Cys Asp Pro Arg His Gly Arg Tyr Leu Thr Val Ala
1 5 10

<210> 929
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens tumor necrosis factor ligand superfamily member 6 epitope

<400> 929

Glu Trp Glu Asp Thr Tyr Gly Ile Val Leu
1 5 10

<210> 930
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Paralichthys olivaceus type 1 collagen alpha 2 epitope

<400> 930

Met Lys Gly Leu Arg Gly His Pro Gly Leu Gln Gly Met Pro Gly Pro
1 5 10 15

Ser Gly Pro Ser
20

<210> 931
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Triticum aestivum type 1 non-specific lipid transfer protein precursor epitope

<400> 931

Ala Arg Gly Thr Pro Leu Lys Cys Gly Val
1 5 10

<210> 932
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens U1 small nuclear ribonucleoprotein 70 kDa epitope

<400> 932

Glu Arg Lys Arg Arg
1 5

<210> 933

<211> 26
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens U1 small nuclear ribonucleoprotein A epitope

<400> 933

Ala Gly Ala Ala Arg Asp Ala Leu Gln Gly Phe Lys Ile Thr Gln Asn
1 5 10 15

Asn Ala Met Lys Ile Ser Phe Ala Lys Lys
20 25

<210> 934
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens U1 small nuclear ribonucleoprotein C epitope

<400> 934

Pro Ala Pro Gly Met Arg Pro Pro
1 5

<210> 935
<211> 18
<212> PRT
<213> Artificial Sequence

<220>
<223> Anisakis simplex UA3-recognized allergen epitope

<400> 935

Met Cys Gln Cys Val Gln Lys Tyr Gly Thr Glu Phe Cys Lys Lys Arg
1 5 10 15

Leu Ala

<210> 936
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens unnamed protein product epitope

<400> 936

Ala Phe Gln Gln Gly Lys Ile Pro Pro
1 5

<210> 937
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Juglans nigra vicilin seed storage protein epitope

<400> 937

Ser Phe Glu Asp Gln Gly Arg Arg
1 5

<210> 938
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Anacardium occidentale Vicilin-like protein epitope

<400> 938

Ala Ile Met Gly Pro Pro Thr Lys Phe Ser Phe Ser Leu Phe Leu
1 5 10 15

<210> 939
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Juglans regia vicilin-like protein precursor epitope

<400> 939

Asp Gln Arg Ser Gln Glu Glu Arg Glu Arg
1 5 10

<210> 940
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens Vimentin epitope

<400> 940

Arg Leu Arg Ser Ser Val Pro Gly Val Arg Leu Leu Gln Asp Ser Val
1 5 10 15

Asp Phe Ser Leu
20

<210> 941
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens von Willebrand factor epitope

<400> 941

His Cys Gln Ile Cys His Cys Asp Val Val Asn Leu Thr Cys Glu
1 5 10 15

<210> 942
<211> 45
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens von Willebrand factor-cleaving protease precursor epitope

<400> 942

Pro Ser His Phe Gln Gln Ser Cys Leu Gln Ala Leu Glu Pro Gln Ala
1 5 10 15

Val Ser Ser Tyr Leu Ser Pro Gly Ala Pro Leu Lys Gly Arg Pro Pro
20 25 30

Ser Pro Gly Phe Gln Arg Gln Arg Gln Arg Gln Arg Arg
35 40 45

<210> 943
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Homo sapiens XRCC4 protein epitope

<400> 943

Val Ser Lys Asp Asp Ser Ile Ile Ser Ser Leu Asp Val Thr Asp
1 5 10 15