



US 20050053981A1

(19) **United States**

(12) **Patent Application Publication**
Swayze et al.

(10) **Pub. No.: US 2005/0053981 A1**

(43) **Pub. Date: Mar. 10, 2005**

(54) **GAPPED OLIGOMERIC COMPOUNDS
HAVING LINKED BICYCLIC SUGAR
MOIETIES AT THE TERMINI**

(76) Inventors: **Eric E. Swayze**, Carlsbad, CA (US);
Michael T. Migawa, San Marcos, CA
(US); **Tadeusz Krzysztof**
Wyrzykiewicz, Carlsbad, CA (US)

Correspondence Address:
COZEN O'CONNOR, P.C.
1900 MARKET STREET
PHILADELPHIA, PA 19103-3508 (US)

(21) Appl. No.: **10/872,106**

(22) Filed: **Jun. 18, 2004**

Related U.S. Application Data

(60) Provisional application No. 60/501,719, filed on Sep. 9, 2003. Provisional application No. 60/568,039, filed on May 3, 2004. Provisional application No. 60/568,489, filed on May 6, 2004.

Publication Classification

(51) **Int. Cl.⁷** **C12Q 1/68; C07H 21/04**
(52) **U.S. Cl.** **435/6; 536/23.1; 536/25.32**

(57) **ABSTRACT**

The present invention relates to bicyclic nucleosides and oligomeric compounds comprising at least one such nucleoside. These oligomeric compounds typically have enhanced binding affinity and nuclease resistance properties compared to unmodified oligomeric compounds. The oligomeric compounds are useful, for example, for investigative and therapeutic purposes.

GAPPED OLIGOMERIC COMPOUNDS HAVING LINKED BICYCLIC SUGAR MOIETIES AT THE TERMINI

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. provisional application Ser. No. 60/501,719 filed Sep. 9, 2003, U.S. provisional application Ser. No. 60/568,039 filed May 3, 2004, and U.S. provisional application Ser. No. 60/568,489 filed May 6, 2004, each of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] The present invention relates to nucleoside compositions comprising novel bicyclic sugar moieties and oligomeric compounds comprising at least one such nucleoside. The oligomeric compounds of the present invention typically will have enhanced binding affinity properties compared to unmodified oligomeric compounds. The oligomeric compounds are useful, for example, for investigative and therapeutic purposes.

BACKGROUND OF THE INVENTION

[0003] Nearly all disease states in multicellular organisms involve the action of proteins. Classic therapeutic approaches have focused on the interaction of proteins with other molecules in efforts to moderate the proteins' disease-causing or disease-potentiating activities. In newer therapeutic approaches, modulation of the production of proteins has been sought. A general object of some current therapeutic approaches is to interfere with or otherwise modulate gene expression.

[0004] One method for inhibiting the expression of specific genes involves the use of oligonucleotides, particularly oligonucleotides that are complementary to a specific target messenger RNA (mRNA) sequence. Due to promising research results in recent years, oligonucleotides and oligonucleotide analogs are now accepted as therapeutic agents holding great promise for therapeutic and diagnostic methods.

[0005] Oligonucleotides and their analogs can be designed to have particular properties. A number of chemical modifications have been introduced into oligomeric compounds to increase their usefulness as therapeutic agents. Such modifications include those designed to increase binding affinity to a target strand, to increase cell penetration, to stabilize against nucleases and other enzymes that degrade or interfere with the structure or activity of the oligonucleotide, to provide a mode of disruption (terminating event) once the oligonucleotide is bound to a target, and to improve the pharmacokinetic properties of the oligonucleotide.

[0006] One group of bicyclic nucleoside compounds having bicyclic sugar moieties that are conformationally locked is locked nucleic acids or LNA (Koshkin et al., Tetrahedron, 1998, 54, 3607-3630; U.S. Pat. No. 6,268,490). These compounds are also referred to in the literature as bicyclic nucleotide analogs (International Patent Application WO 98/39352), but this term is also applicable to a genus of compounds that includes other analogs in addition to LNAs. LNAs have been used in numerous studies where ribo-

nucleoside mimics are desired. Such modified nucleosides mimic the 3'-endo sugar conformation of native ribonucleosides with the advantage of having enhanced binding affinity and increased resistance to nucleases. LNAs are discussed more thoroughly below.

[0007] One group has added an additional methylene group to the LNA 2',4'-bridging group (e.g. 4'-CH₂—CH₂—O-2' (ENA), Kaneko et al., U.S. Patent Application Publication No.: U.S. 2002/0147332, also see Japanese Patent Application HEI-11-33863, Feb. 12, 1999; U.S. Patent Application Publication Nos. 2003/0207841 and 2002/0147332).

[0008] Another publication reports a large genus of nucleosides having a variety of bicyclic sugar moieties with the various bridges creating a bicyclic sugar having a variety of configurations and chemical composition (U.S. Patent Application Publication No.: US 2002/0068708).

[0009] Despite these advances, a need exists in the art for the development of means to improve the binding affinity and nuclease resistance properties of oligomeric compounds.

BRIEF SUMMARY OF THE INVENTION

[0010] The present invention provides oligomeric compounds comprising the structure T₁-(Nu₁-L₁)_{n1}-(Nu₂-L₂)_{n2}-(Nu₃-L₃)_{n3}-T₂ wherein each Nu₁ and Nu₃ is, independently, a high affinity modified nucleoside, wherein at least one nucleoside of Nu₁ and/or at least one nucleoside of Nu₃ is a bicyclic sugar modified nucleoside comprising a 4'-CH₂—O-2' bridge or a 4'-(CH₂)₂—O-2' bridge. Each Nu₂ is a 2'-deoxy nucleoside and each L₁, L₂ and L₃ is, independently, an internucleoside linking group. Each T₁ and T₂ is, independently, H, a hydroxyl protecting group, an optionally linked conjugate group, or a covalent attachment to a solid support medium. n1 is from 1 to about 6, n2 is from 11 to about 18, and n3 is from 2 to about 6.

[0011] In some embodiments, each of the high affinity modified nucleosides is, independently, a bicyclic sugar modified nucleoside, a 2'-O—(CH₂)₂—O—CH₃ modified nucleoside, a 2'-F modified nucleoside, or a 2'-O—CH₂—C(=O)—NR₁R₂ modified nucleoside, where each R₁ and R₂ is, independently, H, a nitrogen protecting group, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, wherein the substitution is OR₃, SR₃, NH₃⁺, NR₃R₄, guanidino or acyl, wherein the acyl is acid amide or an ester, or R₁ and R₂, together, are a nitrogen protecting group, or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O.

[0012] In some embodiments, each of the hydroxyl protecting groups is, independently, 4,4'-dimethoxytrityl, monomethoxytrityl, 9-phenylxanthen-9-yl, 9-(p-methoxyphenyl)xanthen-9-yl, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,p-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triphenylsilyl, benzoylformate, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, pivaloyl, benzoyl, p-phenylbenzoyl, mesyl, tosyl, 4,4',4"-tris-(benzyloxy)trityl, 4,4',4"-tris-(4,5-dichlorophthalimido)trityl, 4,4',4"-tris(levulinyllox-

y)trityl, 3-(imidazolylmethyl)-4,4'-dimethoxytrityl, 4-decyloxytrityl, 4-hexadecyloxytrityl, 9-(4-octadecyloxyphenyl)xanthene-9-yl, 1,1-bis-(4-methoxyphenyl)-1'-pyrenyl methyl, p-phenylazophenylloxycarbonyl, 9-fluorenylmethoxycarbonyl, 2,4-dinitrophenylethoxycarbonyl, 4-(methylthiomethoxy)butyryl, 2-(methylthiomethoxymethyl)-benzoyl, 2-(isopropyl-thiomethoxymethyl)benzoyl, 2-(2,4-dinitrobenzenesulphenyloxymethyl)benzoyl, or levulinyl group.

[0013] In some embodiments, the support medium is a controlled pore glass, oxalyl-controlled pore glass, silica-containing particles, polymers of polystyrene, copolymers of polystyrene, copolymers of styrene and divinylbenzene, copolymers of dimethylacrylamide and N,N'-bisacryloyl ethylenediamine, soluble support medium, or PEPS.


[0014] In some embodiments, each L_1 , L_2 and L_3 is, independently, phosphodiester, phosphorothioate, chiral phosphorothioate, phosphorodithioate, phosphotriester, aminoalkylphosphotriester, methyl phosphonate, alkyl phosphonate, 5'-alkylene phosphonate, chiral phosphonate, phosphinate, phosphoramidate, 3'-amino phosphoramidate, aminoalkylphosphoramidate, thionophosphoramidate, thionoalkylphosphonate, thionoalkylphosphotriester, selenophosphate, boranophosphate, siloxane, sulfide, sulfoxide,

5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-substituted adenines and guanines, 5-substituted uracils and cytosines, 7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, or 3-deazaadenine.

[0016] In some embodiments, n_1 is from 1 to about 5, or from 1 to about 3, or from 2 to about 3. In some embodiments, n_3 is from 2 to about 5, or from 2 to about 3. In some embodiments, n_2 is from 12 to about 18, or from 12 to about 16, or from 14 to about 16. In some embodiments, the total of n_1 , n_2 and n_3 is from 14 to about 30, or from 14 to 24, or from 14 to 21, or from 16 to 21. In some embodiments, n_1 is from 1 to about 3, n_2 is 12 or 13, and n_3 is 2 or 3; or n_1 is 3, n_2 is 12 and n_3 is 3; or n_1 is from 1 to about 3, n_2 is 14 or 15 and n_3 is 2 or 3; or n_1 is 2, n_2 is 14 and n_3 is 2; or n_1 is from 1 to about 3, n_2 is 16 or 17 and n_3 is 2 or 3; or n_1 is 2, n_2 is 16 and n_3 is 2.

[0017] In some embodiments, at least one Nu_1 nucleoside and at least one Nu_3 nucleoside is an LNA or ENA. In some embodiments, at least one of the 5'-most or 3'-most terminal affinity modified nucleosides is an LNA or ENA.

[0018] The present invention also provides oligomeric compounds comprising the structure:



sulfone, formacetyl, thioformacetyl, methylene formacetyl, thioformacetyl, sulfamate, methyleneimino, methylenehydrazino, sulfonate, sulfonamide, or amide. In some embodiments, each of the internucleoside linking groups is, independently, $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$ or $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$, or $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$.

[0015] In some embodiments, each nucleoside comprises a heterocyclic base moiety that is, independently, adenine, guanine, thymine, cytosine, uracil, 5-methylcytosine, 5-hydroxy-methyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, 5-halocytosine,

[0019] wherein each Bx is a heterocyclic base moiety; each X is, independently, O or S; T₁ and T₂ are each, independently, H, a hydroxy protecting group, an optionally linked conjugate group, or a covalent attachment to a solid support medium; each m is, independently, 1 or 2; na is from 1 to about 6; nb is from 11 to about 18; and nc is from 2 to about 6.

[0020] In some embodiments, each m is 1 or 2. In some embodiments, at least one of T₁ and T₂ is H or a hydroxyl protecting group, such as, 4,4'-dimethoxytrityl, monomethoxytrityl, 9-phenylxanthene-9-yl, 9-(p-methoxyphenyl)xanthene-9-yl, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl,

p,p-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butylidimethylsilyl, t-butylidiphenylsilyl, triphenylsilyl, benzoylformate, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, pivaloyl, benzoyl, p-phenylbenzoyl, mesyl, tosyl, 4,4',4''-tris-(benzyloxy)trityl, 4,4',4''-tris-(4,5-dichlorophthalimido)trityl, 4,4',4''-tris(levulinyl)trityl, 3-(imidazolylmethyl)-4,4'-dimethoxytrityl, 4-decyloxytrityl, 4-hexadecyloxytrityl, 9-(4-octadecyloxyphenyl)xanthene-9-yl, 1,1-bis-(4-methoxyphenyl)-1'-pyrenyl methyl, p-phenylazophenylloxycarbonyl, 9-fluorenylmethoxycarbonyl, 2,4-dinitrophenylethoxycarbonyl, 4-(methylthiomethoxy)butyryl, 2-(methylthiomethoxymethyl)-benzoyl, 2-(isopropylthiomethoxymethyl)benzoyl, 2-(2,4-dinitrobenzenesulphenyl-oxy)methyl)benzoyl, or levulinyl group.

[0021] In some embodiments, the support medium is a controlled pore glass, oxalyl-controlled pore glass, silica-containing particles, polymers of polystyrene, copolymers of polystyrene, copolymers of styrene and divinylbenzene, copolymers of dimethylacrylamide and N,N'-bisacryloyl ethylenediamine, soluble support medium, or PEPS.

[0022] In some embodiments, each Bx is, independently, adenine, guanine, thymine, cytosine, uracil, 5-methylcytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, 5-halocytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-substituted adenines and guanines, 5-substituted uracils and cytosines, 7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, or 3-deazaadenine.

[0023] In some embodiments, na is from 1 to about 5, or from 1 to about 3, or from 2 to about 3. In some embodiments, nc is from 2 to about 5, or from 2 to about 3. In some embodiments, nb is from 12 to about 18, or from 12 to about 16, or from 14 to about 16. In some embodiments, the total of na, nb and nc is from 14 to about 30, or from 14 to 24, or from 14 to 21, or from 16 to 21. In some embodiments, na is from 1 to about 3, nb is 12 or 13 and nc is 2 or 3; or na is 3, nb is 12 and nc is 3; or na is from 1 to about 3, nb is 14 or 15 and nc is 2 or 3; or na is 2, nb is 14 and nc is 2; or na is from 1 to about 3, nb is 16 or 17 and nc is 2 or 3; or na is 2, nb is 16 and nc is 2.

[0024] The present invention also provides methods of inhibiting gene expression comprising contacting one or more cells, a tissue, or an animal with any oligomeric compound described herein.

DETAILED DESCRIPTION OF THE INVENTION

[0025] The present invention provides bicyclic nucleoside analogs and oligomeric compounds having at least one of these bicyclic nucleoside analogs. Each of the bicyclic nucleoside analogs has a bridge between the 2' and 4'-positions of the ribofuranose sugar moiety. Oligomeric compounds having at least one of these bicyclic nucleoside analogs will be useful in the modulation of gene expression. More specifically, oligomeric compounds of the invention will modulate gene expression by hybridizing to a nucleic acid target resulting in loss of normal function of the target nucleic acid.

[0026] As used herein, the term "target nucleic acid" or "nucleic acid target" is used for convenience to encompass any nucleic acid capable of being targeted including without limitation DNA, RNA (including pre-mRNA and mRNA or portions thereof) transcribed from such DNA, and also cDNA derived from such RNA. In some embodiment of the invention, the target nucleic acid is a messenger RNA that is degraded by a mechanism involving a nuclease such as or RNaseH. The hybridization of an oligomeric compound of this invention with its target nucleic acid is generally referred to as "antisense." Consequently, one mechanism believed to be included in the practice of some embodiments of the invention is referred to herein as "antisense inhibition." Such antisense inhibition is typically based upon hydrogen bonding-based hybridization of oligonucleotide strands or segments such that at least one strand or segment is cleaved, degraded, or otherwise rendered inoperable. In this regard, it is presently suitable to target specific nucleic acid molecules and their functions for such antisense inhibition.

[0027] The functions of DNA to be interfered include, but are not limited to, replication and transcription. Replication and transcription, for example, can be from an endogenous cellular template, a vector, a plasmid construct or otherwise. The functions of RNA to be interfered with include, but are not limited to, functions such as translocation of the RNA to a site of protein translation, translocation of the RNA to sites within the cell which are distant from the site of RNA synthesis, translation of protein from the RNA, splicing of the RNA to yield one or more RNA species, and catalytic activity or complex formation involving the RNA which may be engaged in or facilitated by the RNA.

[0028] In the context of the present invention, "modulation" and "modulation of expression" mean either an increase (stimulation) or a decrease (inhibition) in the amount or levels of a nucleic acid molecule encoding the gene, e.g., DNA or RNA. Inhibition is often the desired form of modulation of expression and mRNA is often a suitable target nucleic acid.

[0029] Compounds of the Invention

[0030] In the context of the present invention, the term "oligomeric compound" refers to a polymeric structure capable of hybridizing a region of a nucleic acid molecule. This term includes oligonucleotides, oligonucleosides, oligonucleotide analogs, oligonucleotide mimetics and combinations of these. Oligomeric compounds routinely prepared linearly but can be joined or otherwise prepared to be circular and may also include branching. Oligomeric compounds can hybridized to form double stranded compounds which can be blunt ended or may include overhangs. In general an oligomeric compound comprises a backbone of linked monomeric subunits where each linked monomeric subunit is directly or indirectly attached to a heterocyclic base moiety. The linkages joining the monomeric subunits, the sugar moieties or surrogates and the heterocyclic base moieties can be independently modified giving rise to a plurality of motifs for the resulting oligomeric compounds including hemimers, gapmers and chimeras.

[0031] As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base moiety. The two most common classes of such heterocyclic bases are purines and pyrimidines. Nucle-

otides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. The respective ends of this linear polymeric structure can be joined to form a circular structure by hybridization or by formation of a covalent bond, however, open linear structures are generally suitable. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside linkages of the oligonucleotide. The normal internucleoside linkage of RNA and DNA is a 3' to 5' phosphodiester linkage.

[0032] In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA). This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside linkages. The term "oligonucleotide analog" refers to oligonucleotides that have one or more non-naturally occurring portions which function in a similar manner to oligonucleotides. Such non-naturally occurring oligonucleotides are often desired over the naturally occurring forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

[0033] In the context of this invention, the term "oligonucleoside" refers to nucleosides that are joined by internucleoside linkages that do not have phosphorus atoms. Internucleoside linkages of this type include short chain alkyl, cycloalkyl, mixed heteroatom alkyl, mixed heteroatom cycloalkyl, one or more short chain heteroatomic and one or more short chain heterocyclic. These internucleoside linkages include, but are not limited to, siloxane, sulfide, sulfoxide, sulfone, acetyl, formacetyl, thioformacetyl, methylene formacetyl, thioformacetyl, alkenyl, sulfamate; methyleneimino, methylenehydrazino, sulfonate, sulfonamide, amide and others having mixed N, O, S and CH₂ component parts.

[0034] Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439.

[0035] Further included in the present invention are oligomeric compounds such as antisense oligomeric compounds, antisense oligonucleotides, ribozymes, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds which hybridize to at least a portion of the target nucleic acid. As such, these oligomeric compounds may be introduced in the form of single-stranded, double-stranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges or loops. Once introduced

to a system, the oligomeric compounds of the invention may elicit the action of one or more enzymes or structural proteins to effect modification of the target nucleic acid.


[0036] One non-limiting example of such an enzyme is RNase H, a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. It is known in the art that single-stranded antisense oligomeric compounds which are "DNA-like" elicit RNase H. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide-mediated inhibition of gene expression. Similar roles have been postulated for other ribonucleases such as those in the RNase III and ribonuclease L family of enzymes.

[0037] While one form of antisense oligomeric compound is a single-stranded antisense oligonucleotide, in many species the introduction of double-stranded structures, such as double-stranded RNA (dsRNA) molecules, has been shown to induce potent and specific antisense-mediated reduction of the function of a gene or its associated gene products. This phenomenon occurs in both plants and animals and is believed to have an evolutionary connection to viral defense and transposon silencing.

[0038] In addition to the modifications described above, the nucleosides of the oligomeric compounds of the invention can have a variety of other modification so long as these other modifications either alone or in combination with other nucleosides enhance one or more of the desired properties described above. Thus, for nucleotides that are incorporated into oligonucleotides of the invention, these nucleotides can have sugar portions that correspond to naturally-occurring sugars or modified sugars. Representative modified sugars include carbocyclic or acyclic sugars, sugars having substituent groups at one or more of their 2', 3' or 4' positions and sugars having substituents in place of one or more hydrogen atoms of the sugar. Additional nucleosides amenable to the present invention having altered base moieties and or altered sugar moieties are disclosed in U.S. Pat. No. 3,687,808 and PCT application PCT/US89/02323.

[0039] Altered base moieties or altered sugar moieties also include other modifications consistent with the spirit of this invention. Such oligonucleotides are best described as being structurally distinguishable from, yet functionally interchangeable with, naturally occurring or synthetic wild type oligonucleotides. All such oligonucleotides are comprehended by this invention so long as they function effectively to mimic the structure of a desired RNA or DNA strand. A class of representative base modifications include tricyclic cytosine analog, termed "G clamp" (Lin et al., J. Am. Chem. Soc., 1998, 120, 8531). This analog makes four hydrogen bonds to a complementary guanine (G) within a helix by simultaneously recognizing the Watson-Crick and Hoogsteen faces of the targeted G. This G clamp modification when incorporated into phosphorothioate oligonucleotides, dramatically enhances antisense potencies in cell culture. The oligonucleotides of the invention also can include phenoxazine-substituted bases of the type disclosed by Flanagan et al., Nat. Biotechnol., 1999, 17, 48-52.

[0040] The present invention provides oligomeric compounds comprising the structure:



[0041] wherein:

[0042] each Nu₁ and Nu₃ is, independently, a high affinity modified nucleoside, wherein at least one of Nu₁ and Nu₃ is a bicyclic sugar modified nucleoside comprising a 4'-CH₂-O-2' bridge or a 4'-(CH₂)₂-O-2' bridge;

[0043] each Nu₂ is a 2'-deoxy nucleoside;

[0044] each L₁, L₂ and L₃ is, independently, an internucleoside linking group;

[0045] each T₁ and T₂ is, independently, H, a hydroxy protecting group, an optionally linked conjugate group, or a covalent attachment to a solid support medium;

[0046] n₁ is from 1 to about 6;

[0047] n₂ is from 11 to about 18; and

[0048] n₃ is from 2 to about 6.

[0049] In some embodiments, each of the high affinity modified nucleosides is, independently, a bicyclic sugar modified nucleoside, a 2'-O-(CH₂)₂-O-CH₃ modified nucleoside, a 2'-F modified nucleoside, or a 2'-O-CH₂-C(=O)-NR₁R₂ modified nucleoside, where each R₁ and R₂ is, independently, H, a nitrogen protecting group, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, wherein the substitution is OR₃, SR₃, NH₃⁺, NR₃R₄, guanidino or acyl, wherein the acyl is acid amide or an ester, or R₁ and R₂, together, are a nitrogen protecting group, or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O. In some embodiments, each R₁ and R₂ each is, independently, H, a nitrogen protecting group, or C₁-C₁₀ alkyl. In some embodiments, T₁ is H or a hydroxyl protecting group. In some embodiments, T₂ is H or a hydroxyl protecting group.

[0050] In some embodiments, each of the hydroxyl protecting groups is, independently, 4,4'-dimethoxytrityl, monomethoxytrityl, 9-phenylxanthen-9-yl, 9-(p-methoxyphenyl)xanthen-9-yl, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,p-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triphenylsilyl, benzoylformate, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, pivaloyl, benzoyl, p-phenylbenzoyl, mesyl, tosyl, 4,4',4''-tris-(benzyloxy)trityl, 4,4',4''-tris-(4,5-dichlorophthalimido)trityl, 4,4',4''-tris(levulinyloxy)trityl, 3-(imidazolylmethyl)-4,4'-dimethoxytrityl, 4-decyloxytrityl, 4-hexadecyloxytrityl, 9-(4-octadecyloxyphenyl)xanthen-9-yl, 1,1-bis-(4-methoxyphenyl)-1'-pyrenyl methyl, p-phenylazophenylloxycarbonyl, 9-fluorenylmethoxycarbonyl, 2,4-dinitrophenylethoxycarbonyl, 4-(methylthiomethoxy)butyryl, 2-(methylthiomethoxymethyl)-benzoyl, 2-(isopropylthiomethoxymethyl)benzoyl, 2-(2,4-dinitrobenzenesulphenyloxymethyl)benzoyl, or levulinyl group.

[0051] In some embodiments, one of T₁ and T₂ is a covalent attachment to a support medium, such as, for example, a controlled pore glass, oxalyl-controlled pore glass, silica-containing particles, polymers of polystyrene,

copolymers of polystyrene, copolymers of styrene and divinylbenzene, copolymers of dimethylacrylamide and N,N'-bisacryloyl-ethylenediamine, soluble support medium, or PEPS.

[0052] In some embodiments, each L₁, L₂ and L₃ is, independently, phosphodiester, phosphorothioate, chiral phosphorothioate, phosphorodithioate, phosphotriester, aminoalkylphosphotriester, methyl phosphonate, alkyl phosphonate, 5'-alkylene phosphonate, chiral phosphonate, phosphinate, phosphoramidate, 3'-amino phosphoramidate, aminoalkylphosphoramidate, thionophosphoramidate, thionoalkylphosphonate, thionoalkylphosphotriester, selenophosphate, or boranophosphate. In some embodiments, each L₁, L₂ and L₃ is, independently, a phosphodiester or a phosphorothioate internucleoside linking group. In some embodiments, each L₁, L₂ and L₃ is a phosphodiester internucleoside linking group. In some embodiments, each L₁, L₂ and L₃ is, independently, siloxane, sulfide, sulfoxide, sulfone, formacetyl, thioformacetyl, methylene formacetyl, thioformacetyl, sulfamate, methyleneimino, methylenehydrazino, sulfonate, sulfonamide, or amide. In some embodiments, each of the internucleoside linking groups is, independently, -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- or -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂-, or -O-N(CH₃)-CH₂-CH₂-.


[0053] In some embodiments, each nucleoside comprises a heterocyclic base moiety that is, independently, adenine, guanine, thymine, cytosine, uracil, 5-methylcytosine, 5-hydroxy-methyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, 5-halocytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-substituted adenines and guanines, 5-substituted uracils and cytosines, 7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, or 3-deazaadenine.

[0054] In some embodiments, n₁ is from 1 to about 5, or from 1 to about 3, or from 2 to about 3. In some embodiments, n₃ is from 2 to about 5, or from 2 to about 3. In some embodiments, n₂ is from 12 to about 18, or from 12 to about 16, or from 14 to about 16. In some embodiments, the total of n₁, n₂ and n₃ is from 14 to about 30; or the total of n₁, n₂ and n₃ is from 14 to 24; or the total of n₁, n₂ and n₃ is from 14 to 21; or the total of n₁, n₂ and n₃ is from 16 to 21. In some embodiments, n₁ is from 1 to about 3, n₂ is 12 or 13 and n₃ is 2 or 3; or n₁ is 3, n₂ is 12 and n₃ is 3; or n₁ is from 1 to about 3, n₂ is 14 or 15 and n₃ is 2 or 3; or n₁ is 2, n₂ is 14 and n₃ is 2; or n₁ is from 1 to about 3, n₂ is 16 or 17 and n₃ is 2 or 3; or n₁ is 2, n₂ is 16 and n₃ is 2.

[0055] In some embodiments, at least one Nu₁ nucleoside and at least one Nu₃ nucleoside is an LNA or ENA. In some embodiments, at least one of the 5'-most or 3'-most terminal affinity modified nucleosides is an LNA or ENA. Thus, the terminal most 5' and/or 3' nucleoside can, independently, be either LNA or ENA.

[0056] The present invention also provides oligomeric compounds comprising the structure:

omethoxymethyl)benzoyl, 2-(2,4-dinitrobenzenesulphenyl-oxymethyl)benzoyl, or levulinyl group.



[0057] wherein:

[0058] each Bx is a heterocyclic base moiety;

[0059] each X is, independently, O or S;

[0060] T_1 and T_2 are each, independently, H, a hydroxyl protecting group, an optionally linked conjugate group, or a covalent attachment to a solid support medium;

[0061] each m is, independently, 1 or 2;

[0062] na is from 1 to about 6;

[0063] nb is from 11 to about 18; and

[0064] nc is from 2 to about 6.

[0065] In some embodiments, each m is 1 or 2. In some embodiments, at least one of T_1 and T_2 is H or a hydroxyl protecting group. In some embodiments, each of the hydroxyl protecting groups is, independently, 4,4'-dimethoxytrityl, monomethoxytrityl, 9-phenylxanthen-9-yl, 9-(p-methoxyphenyl)xanthen-9-yl, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,p-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triphenylsilyl, benzoylformate, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, pivaloyl, benzoyl, p-phenylbenzoyl, mesyl, tosyl, 4,4',4''-tris-(benzyloxy)trityl, 4,4',4''-tris-(4,5-dichlorophthalimido)trityl, 4,4',4''-tris(levulinyl)oxytrityl, 3-(imidazolylmethyl)-4,4'-dimethoxytrityl, 4-decyloxytrityl, 4-hexadecyloxytrityl, 9-(4-octadecyloxyphenyl)xanthen-9-yl, 1,1-bis-(4-methoxyphenyl)-1'-pyrenyl methyl, p-phenylazophenylloxycarbonyl, 9-fluorenylmethoxycarbonyl, 2,4-dinitrophenylethoxycarbonyl, 4-(methylthiomethoxy)butyryl, 2-(methylthiomethoxymethyl)-benzoyl, 2-(isopropylthi-

[0066] In some embodiments, one of T_1 and T_2 is a covalent attachment to a support medium such as, for example, a controlled pore glass, oxalyl-controlled pore glass, silica-containing particles, polymers of polystyrene, copolymers of polystyrene, copolymers of styrene and divinylbenzene, copolymers of dimethylacrylamide and N,N'-bisacryloyl-ethylenediamine, soluble support medium, or PEPS.

[0067] In some embodiments, each L_1 , L_2 and L_3 is, independently, a phosphodiester or a phosphorothioate internucleoside linking group. In some embodiments, each L_1 , L_2 and L_3 is a phosphodiester internucleoside linking group.

[0068] In some embodiments, each Bx is, independently, adenine, guanine, thymine, cytosine, uracil, 5-methylcytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, 5-halocytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-substituted adenines and guanines, 5-substituted uracils and cytosines, 7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, or 3-deazaadenine.

[0069] In some embodiments, n1 is from 1 to about 5 (e.g., 1, 2, 3, 4, 5), or from 1 to about 3 (e.g., 1, 2, 3), or from 2 to about 3. In some embodiments, n3 is from 2 to about 5 (e.g., 2, 3, 4, 5), or from 2 to about 3. In some embodiments, n2 is from 12 to about 18 (e.g., 12, 13, 14, 15, 16, 17, 18), or from 12 to about 16 (e.g., 12, 13, 14, 15, 16), or from 14 to about 16 (e.g., 14, 15, 16). In some embodiments, the total of n1, n2 and n3 is from 14 to about 30 (e.g., 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30); or the total of n1, n2 and n3 is from 14 to 24 (e.g., 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24); or the total of n1, n2 and n3 is from 14 to 21 (e.g., 14, 15, 16, 17, 18, 19, 20, 21); or the total of n1, n2 and n3 is from 16 to 21 (e.g., 16, 17, 18, 19, 20, 21). In some embodiments, n1 is from 1 to about 3, n2 is 12 or

13 and n3 is 2 or 3; or n1 is 3, n2 is 12 and n3 is 3; or n1 is from 1 to about 3, n2 is 14 or 15 and n3 is 2 or 3; or n1 is 2, n2 is 14 and n3 is 2; or n1 is from 1 to about 3, n2 is 16 or 17 and n3 is 2 or 3; or n1 is 2, n2 is 16 and n3 is 2.

[0070] Oligomer Synthesis

[0071] Oligomerization of modified and unmodified nucleosides is performed according to literature procedures for DNA (Protocols for Oligonucleotides and Analogs, Ed. Agrawal (1993), Humana Press) and/or RNA (Scaringe, Methods (2001), 23, 206-217. Gait et al., Applications of Chemically synthesized RNA in RNA: Protein Interactions, Ed. Smith (1998), 1-36. Gallo et al., Tetrahedron (2001), 57, 5707-5713) synthesis as appropriate. In addition specific protocols for the synthesis of oligomeric compounds of the invention are illustrated in the examples below.

[0072] The oligomeric compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

[0073] The present invention is also useful for the preparation of oligomeric compounds incorporating at least one 2'-O-protected nucleoside. After incorporation and appropriate deprotection the 2'-O-protected nucleoside will be converted to a ribonucleoside at the position of incorporation. The number and position of the 2-ribonucleoside units in the final oligomeric compound can vary from one at any site or the strategy can be used to prepare up to a full 2'-OH modified oligomeric compound. All 2'-O-protecting groups amenable to the synthesis of oligomeric compounds are included in the present invention. In general a protected nucleoside is attached to a solid support by for example a succinate linker. Then the oligonucleotide is elongated by repeated cycles of deprotecting the 5'-terminal hydroxyl group, coupling of a further nucleoside unit, capping and oxidation (alternatively sulfurization). In a more frequently used method of synthesis the completed oligonucleotide is cleaved from the solid support with the removal of phosphate protecting groups and exocyclic amino protecting groups by treatment with an ammonia solution. Then a further deprotection step is normally required for the more specialized protecting groups used for the protection of 2'-hydroxyl groups which will give the fully deprotected oligonucleotide.

[0074] A large number of 2'-O-protecting groups have been used for the synthesis of oligoribonucleotides but over the years more effective groups have been discovered. The key to an effective 2'-O-protecting group is that it is capable of selectively being introduced at the 2'-O-position and that it can be removed easily after synthesis without the formation of unwanted side products. The protecting group also needs to be inert to the normal deprotecting, coupling, and capping steps required for oligoribonucleotide synthesis. Some of the protecting groups used initially for oligoribonucleotide synthesis included tetrahydropyran-1-yl and 4-methoxytetrahydropyran-4-yl. These two groups are not compatible with all 5'-O-protecting groups so modified versions were used with 5'-DMT groups such as 1-(2-

fluorophenyl)-4-methoxypiperidin-4-yl (Fpmp). Reese has identified a number of piperidine derivatives (like Fpmp) that are useful in the synthesis of oligoribonucleotides including 1-((chloro-4-methylphenyl)-4'-methoxypiperidin-4-yl (Reese et al., Tetrahedron Lett., 1986, 27, 2291). Another approach was to replace the standard 5'-DMT (dimethoxytrityl) group with protecting groups that were removed under non-acidic conditions such as levulinyl and 9-fluorenylmethoxycarbonyl. Such groups enable the use of acid labile 2'-protecting groups for oligoribonucleotide synthesis. Another more widely used protecting group initially used for the synthesis of oligoribonucleotides was the t-butyl dimethylsilyl group (Ogilvie et al., Tetrahedron Lett., 1974, 2861; Hakimelahi et al., Tetrahedron Lett., 1981, (22), 2543; and Jones et al., J. Chem. Soc. Perkin I., 2762). The 2'-O-protecting groups can require special reagents for their removal such as for example the t-butyl dimethylsilyl group is normally removed after all other cleaving/deprotecting steps by treatment of the oligomeric compound with tetrabutylammonium fluoride (TBAF).

[0075] One group of researchers examined a number of 2'-O-protecting groups (Pitsch, Chimia, 2001, (55), 320-324.) The group examined fluoride labile and photolabile protecting groups that are removed using moderate conditions. One photolabile group that was examined was the (2-(nitrobenzyl)oxy)methyl (nbm) protecting group (Schwartz et al., Bioorg. Med. Chem. Lett., 1992, 2, 1019). Other groups examined included a number structurally related formaldehyde acetal-derived, 2'-O-protecting groups. Also prepared were a number of related protecting groups for preparing 2'-O-alkylated nucleoside phosphoramidites including 2'-O-((triisopropylsilyl)oxy)methyl (2'-O-CH₂-O-Si(iPr)₃, TOM). One 2'-O-protecting group that was prepared to be used orthogonally to the TOM group was 2'-O-((R)-1-(2-nitrophenyl)ethoxy)methyl ((R)-mnbm).

[0076] Another strategy using a fluoride labile 5'-O-protecting group (non-acid labile) and an acid labile 2'-O-protecting group has been reported (Scaringe, Stephen A., Methods, 2001, 23, 206-217). A number of possible silyl ethers were examined for 5'-O-protection and a number of acetals and orthoesters were examined for 2'-O-protection. The protection scheme that gave the best results was 5'-O-silyl ether-2'-ACE (5'-O-bis(trimethylsiloxy)cyclododecyloxysilyl ether (DOD)-2'-O-bis(2-acetoxyethoxy)methyl (ACE). This approach uses a modified phosphoramidite synthesis approach in that some different reagents are required that are not routinely used for RNA/DNA synthesis.

[0077] Although a lot of research has focused on the synthesis of oligoribonucleotides the main RNA synthesis strategies that are presently being used commercially include 5'-O-DMT-2'-O-t-butyl dimethylsilyl (TBDMS), 5'-O-DMT-2'-O-(1 (2-fluorophenyl)-4-methoxypiperidin-4-yl) (FPMP), 2'-O-((triisopropylsilyl)oxy)methyl (2'-O-CH₂-O-Si(iPr)₃, TOM), and the 5'-O-silyl ether-2'-ACE (5'-O-bis(trimethylsiloxy)cyclododecyloxysilyl ether (DOD)-2'-O-bis(2-acetoxyethoxy)methyl (ACE). A current list of some of the major companies currently offering RNA products include Pierce Nucleic Acid Technologies, Dharmacon Research Inc., Ameri Biotechnologies Inc., and Integrated DNA Technologies, Inc. One company, Princeton Separations, is marketing an RNA synthesis activator advertised to reduce coupling times especially with TOM and

TBDMS chemistries. Such an activator would also be amenable to the present invention.

[0078] The primary groups being used for commercial RNA synthesis are:

[0079] TBDMS=5'-O-DMT-2'-O-t-butyl dimethylsilyl;

[0080] TOM=2'-O-((triisopropylsilyl)oxy)methyl;

[0081] DOD/ACE=(5'-O-bis(trimethylsilyloxy)cyclododecyloxysilyl ether-2'-O-bis(2-acetoxyethoxy)methyl

[0082] FPMP=5'-O-DMT-2'-O-(1 (2-fluorophenyl)-4-methoxypiperidin-4-yl).

[0083] All of the aforementioned RNA synthesis strategies are amenable to the present invention. Strategies that would be a hybrid of the above e.g. using a 5'-protecting group from one strategy with a 2'-O-protecting from another strategy is also amenable to the present invention.

[0084] The preparation of ribonucleotides and oligomeric compounds having at least one ribonucleoside incorporated and all the possible configurations falling in between these two extremes are encompassed by the present invention. The corresponding oligomeric compounds can be hybridized to further oligomeric compounds including oligoribonucleotides having regions of complementarity to form double-stranded (duplexed) oligomeric compounds. Such double stranded oligonucleotide moieties have been shown in the art to modulate target expression and regulate translation as well as RNA processing via an antisense mechanism. Moreover, the double-stranded moieties may be subject to chemical modifications (Fire et al., *Nature*, 1998, 391, 806-811; Timmons and Fire, *Nature* 1998, 395, 854; Timmons et al., *Gene*, 2001, 263, 103-112; Tabara et al., *Science*, 1998, 282, 430-431; Montgomery et al., *Proc. Natl. Acad. Sci. USA*, 1998, 95, 15502-15507; Tuschl et al., *Genes Dev.*, 1999, 13, 3191-3197; Elbashir et al., *Nature*, 2001, 411, 494-498; Elbashir et al., *Genes Dev.* 2001, 15, 188-200). For example, such double-stranded moieties have been shown to inhibit the target by the classical hybridization of antisense strand of the duplex to the target, thereby triggering enzymatic degradation of the target (Tijsterman et al., *Science*, 2002, 295, 694-697).

[0085] The methods of preparing oligomeric compounds of the present invention can also be applied in the areas of drug discovery and target validation. The present invention comprehends the use of the oligomeric compounds and suitable targets identified herein in drug discovery efforts to elucidate relationships that exist between proteins and a disease state, phenotype, or condition. These methods include detecting or modulating a target peptide comprising contacting a sample, tissue, cell, or organism with the oligomeric compounds of the present invention, measuring the nucleic acid or protein level of the target and/or a related phenotypic or chemical endpoint at some time after treatment, and optionally comparing the measured value to a non-treated sample or sample treated with a further oligomeric compound of the invention. These methods can also be performed in parallel or in combination with other experiments to determine the function of unknown genes for the process of target validation or to determine the validity

of a particular gene product as a target for treatment or prevention of a particular disease, condition, or phenotype.

[0086] Effect of nucleoside modifications on RNAi activity is evaluated according to existing literature (Elbashir et al., *Nature* (2001), 411, 494-498; Nishikura et al., *Cell* (2001), 107, 415-416; and Bass et al., *Cell* (2000), 101, 235-238.)

[0087] A number of chemical functional groups can be introduced into compounds of the invention in a blocked form and subsequently deblocked to form a final, desired compound. Such as groups directly or indirectly attached at the heterocyclic bases, the internucleoside linkages and the sugar substituent groups at the 2', 3' and 5'-positions. Protecting groups can be selected to block functional groups located in a growing oligomeric compound during iterative oligonucleotide synthesis while other positions can be selectively deblocked as needed. In general, a blocking group renders a chemical functionality of a larger molecule inert to specific reaction conditions and can later be removed from such functionality without substantially damaging the remainder of the molecule (Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd ed, John Wiley & Sons, New York, 1999). For example, the nitrogen atom of amino groups can be blocked as phthalimido groups, as 9-fluorenylmethoxycarbonyl (FMOC) groups, and with triphenylmethylsulfenyl, t-BOC or benzyl groups. Carboxyl groups can be blocked as acetyl groups. Representative hydroxyl protecting groups are described by Beaucage et al., *Tetrahedron* 1992, 48, 2223. Suitable hydroxyl protecting groups are acid-labile, such as the trityl, monomethoxytrityl, dimethoxytrityl, trimethoxytrityl, 9-phenylxanthine-9-yl (Pixyl) and 9-(p-methoxyphenyl)xanthine-9-yl (MOX).

[0088] Chemical functional groups can also be "blocked" by including them in a precursor form. Thus, an azido group can be used considered as a "blocked" form of an amine since the azido group is easily converted to the amine. Further representative protecting groups utilized in oligonucleotide synthesis are discussed in Agrawal, et al., *Protocols for Oligonucleotide Conjugates*, Eds, Humana Press; New Jersey, 1994; Vol. 26 pp. 1-72.

[0089] Examples of hydroxyl protecting groups include, but are not limited to, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,=dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triphenylsilyl, benzoylformate, acetate, chloroacetate, trichloroacetate, trifluoroacetate, pivaloate, benzoate, p-phenylbenzoate, 9-fluorenylmethyl carbonate, mesylate and tosylate.

[0090] Amino-protecting groups stable to acid treatment are selectively removed with base treatment, and are used to make reactive amino groups selectively available for substitution. Examples of such groups are the Fmoc (E. Atherton and R. C. Sheppard in *The Peptides*, S. Udenfriend, J. Meienhofer, Eds., Academic Press, Orlando, 1987, volume 9, p.1), and various substituted sulfonylethyl carbamates exemplified by the Nsc group (Samukov et al., *Tetrahedron Lett*, 1994, 35:7821; Verhart and Tesser, *Rec. Trav. Chim. Pays-Bas*, 1987, 107, 621).

[0091] Additional amino-protecting groups include but are not limited to, carbamate-protecting groups, such as 2-tri-

methylsilylethoxycarbonyl (Teoc), 1-methyl-1-(4-biphenyl)ethoxycarbonyl (Bpoc), t-butoxycarbonyl (BOC), allyloxycarbonyl (Alloc), 9-fluorenylmethyloxycarbonyl (Fmoc), and benzyloxycarbonyl (Cbz); amide-protecting groups, such as formyl, acetyl, trihaloacetyl, benzoyl, and nitrophenylacetyl; sulfonamide-protecting groups, such as 2-nitrobenzenesulfonyl; and imine- and cyclic imide-protecting groups, such as phthalimido and dithiasuccinoyl.

[0092] The current method of choice for the preparation of oligomeric compounds utilizes support media. Support media is used for attachment of a first nucleoside or other synthon which is then iteratively elongated to give a final oligomeric compound or other polymer such as a polypeptide. Support media can be selected to be insoluble or have variable solubility in different solvents to allow the growing support bound polymer to be either in or out of solution as desired. Traditional support media such as solid supports are for the most part insoluble and are routinely placed in a reaction vessel while reagents and solvents react and or wash the growing chain until cleavage the final polymeric compound. More recent approaches have introduced soluble supports including soluble polymer supports to allow precipitating and dissolving the iteratively synthesized product at desired points in the synthesis (Gravert et al., *Chem. Rev.*, 1997, 97, 489-510).

[0093] The term support media is intended to include all forms of support known to the art skilled for the synthesis of oligomeric compounds and related compounds such as peptides. Some representative support media that are amenable to the methods of the present invention include but are not limited to the following: controlled pore glass (CPG); oxalyl-controlled pore glass (see, e.g., Alul, et al., *Nucleic Acids Research* 1991, 19, 1527); silica-containing particles, such as porous glass beads and silica gel such as that formed by the reaction of trichloro-(3-(4-chloromethyl)phenyl)propylsilane and porous glass beads (see Parr and Grohmann, *Angew. Chem. Internat. Ed.* 1972, 11, 314, sold under the trademark "PORASIL E" by Waters Associates, Framingham, Mass., USA); the mono ester of 1,4-dihydroxymethylbenzene and silica (see Bayer and Jung, *Tetrahedron Lett.*, 1970, 4503, sold under the trademark "BIOPAK" by Waters Associates); TENTAGEL (see, e.g., Wright, et al., *Tetrahedron Letters* 1993, 34, 3373); cross-linked styrene/divinylbenzene copolymer beaded matrix or POROS, a copolymer of polystyrene/divinylbenzene (available from Perceptive Biosystems); soluble support media, polyethylene glycol PEG's (see Bonora et al., *Organic Process Research & Development*, 2000, 4, 225-231).

[0094] Further support media amenable to the present invention include without limitation PEPS support a polyethylene (PE) film with pendant long-chain polystyrene (PS) grafts (molecular weight on the order of 106, (see Berg, et al., *J. Am. Chem. Soc.*, 1989, 111, 8024 and International Patent Application WO 90/02749). The loading capacity of the film is as high as that of a beaded matrix with the additional flexibility to accommodate multiple syntheses simultaneously. The PEPS film may be fashioned in the form of discrete, labeled sheets, each serving as an individual compartment. During all the identical steps of the synthetic cycles, the sheets are kept together in a single reaction vessel to permit concurrent preparation of a multitude of peptides at a rate close to that of a single peptide by conventional methods. Also, experiments with other geometries of the

PEPS polymer such as, for example, non-woven felt, knitted net, sticks or microwellplates have not indicated any limitations of the synthetic efficacy.

[0095] Further support media amenable to the present invention include without limitation particles based upon copolymers of dimethylacrylamide cross-linked with N,N'-bisacryloylethylenediamine, including a known amount of N-tert-butoxycarbonyl-beta-alanyl-N'-acryloylhexamethylenediamine. Several spacer molecules are typically added via the beta alanyl group, followed thereafter by the amino acid residue subunits. Also, the beta alanyl-containing monomer can be replaced with an acryloyl safcosine monomer during polymerization to form resin beads. The polymerization is followed by reaction of the beads with ethylenediamine to form resin particles that contain primary amines as the covalently linked functionality. The polyacrylamide-based supports are relatively more hydrophilic than are the polystyrene-based supports and are usually used with polar aprotic solvents including dimethylformamide, dimethylacetamide, N-methylpyrrolidone and the like (see Atherton, et al., *J. Am. Chem. Soc.*, 1975, 97, 6584, *Bioorg. Chem.* 1979, 8, 351, and *J. C. S. Perkin* 1538 (1981)).

[0096] Further support media amenable to the present invention include without limitation a composite of a resin and another material that is also substantially inert to the organic synthesis reaction conditions employed. One exemplary composite (see Scott, et al., *J. Chrom. Sci.*, 1971, 9, 577) utilizes glass particles coated with a hydrophobic, cross-linked styrene polymer containing reactive chloromethyl groups, and is supplied by Northgate Laboratories, Inc., of Hamden, Conn., USA. Another exemplary composite contains a core of fluorinated ethylene polymer onto which has been grafted polystyrene (see Kent and Merrifield, *Israel J. Chem.* 1978, 17, 243 and van Rietschoten in *Peptides* 1974, Y. Wolman, Ed., Wiley and Sons, New York, 1975, pp. 113-116). Contiguous solid supports other than PEPS, such as cotton sheets (Lebl and Eichler, *Peptide Res.* 1989, 2, 232) and hydroxypropylacrylate-coated polypropylene membranes (Daniels, et al., *Tetrahedron Lett.* 1989, 4345). Acrylic acid-grafted polyethylene-rods and 96-microtiter wells to immobilize the growing peptide chains and to perform the compartmentalized synthesis. (Geysen, et al., *Proc. Natl. Acad. Sci. USA*, 1984, 81, 3998). A "tea bag" containing traditionally-used polymer beads. (Houghten, *Proc. Natl. Acad. Sci. USA*, 1985, 82, 5131). Simultaneous use of two different supports with different densities (Tregear, *Chemistry and Biology of Peptides*, J. Meienhofer, ed., Ann Arbor Sci. Publ., Ann Arbor, 1972 pp. 175-178). Combining of reaction vessels via a manifold (Gorman, *Anal. Biochem.*, 1984, 136, 397). Multicolumn solid-phase synthesis (e.g., Krchnak, et al., *Int. J. Peptide Protein Res.*, 1989, 33, 209), and Holm and Meldal, in "Proceedings of the 20th European Peptide Symposium", G. Jung and E. Bayer, eds., Walter de Gruyter & Co., Berlin, 1989 pp. 208-210). Cellulose paper (Eichler, et al., *Collect. Czech. Chem. Commun.*, 1989, 54, 1746). Support mediated synthesis of peptides have also been reported (see, *Synthetic Peptides: A User's Guide*, Gregory A. Grant, Ed. Oxford University Press 1992; U.S. Pat. Nos. 4,415,732; 4,458,066; 4,500,707; 4,668,777; 4,973,679; 5,132,418; 4,725,677 and Re-34, 069.)

[0097] Support bound oligonucleotide synthesis relies on sequential addition of nucleotides to one end of a growing

chain. Typically, a first nucleoside (having protecting groups on any exocyclic amine functionalities present) is attached to an appropriate glass bead support and activated phosphite compounds (typically nucleotide phosphoramidites, also bearing appropriate protecting groups) are added stepwise to elongate the growing oligonucleotide. Additional methods for solid-phase synthesis may be found in Caruthers U.S. Pat. Nos. 4,415,732; 4,458,066; 4,500,707; 4,668,777; 4,973,679; and 5,132,418; and Koster U.S. Pat. Nos. 4,725,677 and Re. 34,069.

[0098] Commercially available equipment routinely used for the support media based synthesis of oligomeric compounds and related compounds is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). Any other means for such synthesis known in the art may additionally or alternatively be employed. Suitable solid phase techniques, including automated synthesis techniques, are described in F. Eckstein (ed.), *Oligonucleotides and Analogues, a Practical Approach*, Oxford University Press, New York (1991).

[0099] Oligomer and Monomer Modifications

[0100] As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn, the respective ends of this linear polymeric compound can be further joined to form a circular compound, however, linear compounds are generally suitable. In addition, linear compounds may have internal nucleobase complementarity and may therefore fold in a manner as to produce a fully or partially double-stranded compound. Within oligonucleotides, the phosphate groups are commonly referred to as forming the internucleoside linkage or in conjunction with the sugar ring the backbone of the oligonucleotide. The normal internucleoside linkage that makes up the backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

[0101] Modified Internucleoside Linkages

[0102] Specific examples of suitable oligomeric compounds useful in this invention include oligonucleotides containing modified e.g. non-naturally occurring internucleoside linkages. As defined in this specification, oligonucleotides having modified internucleoside linkages include internucleoside linkages that retain a phosphorus atom and internucleoside linkages that do not have a phosphorus atom. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

[0103] In the *C. elegans* system, modification of the internucleotide linkage (phosphorothioate) did not significantly interfere with RNAi activity. Based on this observation, it is suggested that certain oligomeric compounds of the invention can also have one or more modified internucleo-

side linkages. A suitable phosphorus containing modified internucleoside linkage is the phosphorothioate internucleoside linkage.

[0104] Suitable modified oligonucleotide backbones containing a phosphorus atom therein include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thiono-phosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Suitable oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

[0105] Representative U.S. patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050.

[0106] In some embodiments of the invention, oligomeric compounds have one or more phosphorothioate and/or heteroatom internucleoside linkages, in particular $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$ (known as a methylene (methylimino) or MMI backbone), $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$ and $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$ (wherein the native phosphodiester internucleotide linkage is represented as $-\text{O}-\text{P}(=\text{O})(\text{OH})-\text{O}-\text{CH}_2-$). The MMI type internucleoside linkages are disclosed in the above referenced U.S. Pat. No. 5,489,677. Suitable amide internucleoside linkages are disclosed in the above referenced U.S. Pat. No. 5,602,240.

[0107] Suitable modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH_2 component parts.

[0108] Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,

444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439.


[0109] In some embodiments, the nucleoside components of the oligomeric compounds are connected to each other by optionally protected phosphorothioate internucleoside linkages. Representative protecting groups for phosphorus containing internucleoside linkages such as phosphite, phosphodiester and phosphorothioate linkages include β -cyanoethyl, diphenylsilylethyl, δ -cyanobutenyl, cyano p-xylyl (CPX), N-methyl-N-trifluoroacetyl ethyl (META), acetoxo phenoxy ethyl (APE) and butene-4-yl groups. See for example U.S. Pat. Nos. 4,725,677 and Re. 34,069 (β -cyanoethyl); Beaucage, S. L. and Iyer, R. P., Tetrahedron, 49 No. 10, pp. 1925-1963 (1993); Beaucage, S. L. and Iyer, R. P., Tetrahedron, 49 No. 46, pp. 10441-10488 (1993); Beaucage, S. L. and Iyer, R. P., Tetrahedron, 48 No. 12, pp. 2223-2311 (1992).

[0110] Oligomer Mimetics (Oligonucleotide Mimics)

[0111] Another suitable group of oligomeric compounds amenable to the present invention includes oligonucleotide mimetics. The term mimetic as it is applied to oligonucleotides is intended to include oligomeric compounds wherein only the furanose ring or both the furanose ring and the internucleotide linkage are replaced with novel groups, replacement of only the furanose ring is also referred to in the art as being a sugar surrogate. The heterocyclic base moiety or a modified heterocyclic base moiety is maintained for hybridization with an appropriate target nucleic acid. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA oligomeric compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA oligomeric compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262. Further teaching of PNA oligomeric compounds can be found in Nielsen et al., Science, 1991, 254, 1497-1500.

[0112] One oligonucleotide mimetic that has been reported to have excellent hybridization properties, is peptide nucleic acids (PNA). The backbone in PNA compounds is two or more linked aminoethylglycine units which gives PNA an amide containing backbone. The heterocyclic base moieties are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262. Further teaching of PNA compounds can be found in Nielsen et al., Science, 1991, 254, 1497-1500.

[0113] PNA has been modified to incorporate numerous modifications since the basic PNA structure was first prepared. The basic structure is shown below:



[0114] wherein

[0115] Bx is a heterocyclic base moiety;

[0116] T₄ is hydrogen, an amino protecting group, —C(O)R₅, substituted or unsubstituted C₁-C₁₂ alkyl, substituted or unsubstituted C₂-C₁₂ alkenyl, substituted or unsubstituted C₂-C₁₂ alkynyl, alkylsulfonyl, arylsulfonyl, a chemical functional group, a reporter group, a conjugate group, a D or L α -amino acid linked via the α -carboxyl group or optionally through the ω -carboxyl group when the amino acid is aspartic acid or glutamic acid or a peptide derived from D, L or mixed D and L amino acids linked through a carboxyl group, wherein the substituent groups are selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl;

[0117] T₅ is —OH, —N(Z₁)Z₂, R₅, D or L α -amino acid linked via the α -amino group or optionally through the ϵ -amino group when the amino acid is lysine or ornithine or a peptide derived from D, L or mixed D and L amino acids linked through an amino group, a chemical functional group, a reporter group or a conjugate group;

[0118] Z₁ is hydrogen, C₁-C₆ alkyl, or an amino protecting group;

[0119] Z₂ is hydrogen, C₁-C₆ alkyl, an amino protecting group, —C(=O)—(CH₂)_n-J-Z₃, a D or L α -amino acid linked via the α -carboxyl group or optionally through the ω -carboxyl group when the amino acid is aspartic acid or glutamic acid or a peptide derived from D, L or mixed D and L amino acids linked through a carboxyl group;

[0120] Z₃ is hydrogen, an amino protecting group, —C-C₆ alkyl, —C(=O)—CH₃, benzyl, benzoyl, or —(CH₂)_n-N(H)Z₁;

[0121] each J is O, S or NH;


[0122] R₅ is a carbonyl protecting group; and

[0123] n is from 2 to about 50.

[0124] Another class of oligonucleotide mimetic that has been studied is based on linked morpholino units (morpholino nucleic acid) having heterocyclic bases attached to the morpholino ring. A number of linking groups have been reported that link the morpholino monomeric units in a morpholino nucleic acid. One class of linking groups have been selected to give a non-ionic oligomeric compound. The non-ionic morpholino-based oligomeric compounds are less

likely to have undesired interactions with cellular proteins. Morpholino-based oligomeric compounds are non-ionic mimics of oligonucleotides which are less likely to form undesired interactions with cellular proteins (Dwayne A. Braasch and David R. Corey, *Biochemistry*, 2002, 41(14), 4503-4510). Morpholino-based oligomeric compounds are disclosed in U.S. Pat. No. 5,034,506, issued Jul. 23, 1991. The morpholino class of oligomeric compounds have been prepared having a variety of different linking groups joining the monomeric subunits.

[0125] Morpholino nucleic acids have been prepared having a variety of different linking groups (L_2) joining the monomeric subunits. The basic formula is shown below:



[0126] wherein:

[0127] T_1 is hydroxyl or a protected hydroxyl;


[0128] T_5 is hydrogen or a phosphate or phosphate derivative;

[0129] L_2 is a linking group; and

[0130] n is from 2 to about 50.

[0131] A further class of oligonucleotide mimetic is referred to as cyclohexenyl nucleic acids (CeNA). The furanose ring normally present in a DNA/RNA molecule is replaced with a cyclohexenyl ring. CeNA DMT protected phosphoramidite monomers have been prepared and used for oligomeric compound synthesis following classical phosphoramidite chemistry. Fully modified CeNA oligomeric compounds and oligonucleotides having specific positions modified with CeNA have been prepared and studied (see Wang et al., *J. Am. Chem. Soc.*, 2000, 122, 8595-8602). In general the incorporation of CeNA monomers into a DNA chain increases its stability of a DNA/RNA hybrid. CeNA oligoadenylates formed complexes with RNA and DNA complements with similar stability to the native complexes. The study of incorporating CeNA structures into natural nucleic acid structures was shown by NMR and circular dichroism to proceed with easy conformational adaptation. Furthermore the incorporation of CeNA into a sequence targeting RNA was stable to serum and able to activate *E. Coli* RNase resulting in cleavage of the target RNA strand.

[0132] The general formula of CeNA is shown below:




[0133] wherein:

[0134] each Bx is a heterocyclic base moiety;


[0135] T_1 is hydroxyl or a protected hydroxyl; and

[0136] T_2 is hydroxyl or a protected hydroxyl.

[0137] Another class of oligonucleotide mimetic (anhydrohexitol nucleic acid) can be prepared from one or more anhydrohexitol nucleosides (see, Wouters and Herdewijn, *Bioorg. Med. Chem. Lett.*, 1999, 9, 1563-1566) and would have the general formula:



[0138] Another modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 4' carbon atom of the sugar ring thereby forming a 2'-C,4'-C-oxymethylene linkage thereby forming a bicyclic sugar moiety. The linkage can be a methylene ($-\text{CH}_2-$) $_n$ group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 for LNA (or 2 for ENA, Singh et al., *Chem. Commun.*, 1998, 4, 455-456). LNA and LNA analogs display very high duplex thermal stabilities with complementary DNA and RNA ($T_m = +3$ to $+10$ C), stability towards 3'-exonucleolytic degradation and good solubility properties. The basic structure of LNA showing the bicyclic ring system is shown below:



[0139] The conformations of LNAs determined by 2D NMR spectroscopy have shown that the locked orientation of the LNA nucleotides, both in single-stranded LNA and in duplexes, constrains the phosphate backbone in such a way as to introduce a higher population of the N-type conformation (Petersen et al., *J. Mol. Recognit.*, 2000, 13, 44-53). These conformations are associated with improved stacking of the nucleobases (Wengel et al., *Nucleosides Nucleotides*, 1999, 18, 1365-1370).

[0140] LNA has been shown to form exceedingly stable LNA:LNA duplexes (Koshkin et al., *J. Am. Chem. Soc.*, 1998, 120, 13252-13253). LNA:LNA hybridization was shown to be the most thermally stable nucleic acid type duplex system, and the RNA-mimicking character of LNA was established at the duplex level. Introduction of 3 LNA monomers (T or A) significantly increased melting points ($T_m = +15/+11$) toward DNA complements. The universality of LNA-mediated hybridization has been stressed by the formation of exceedingly stable LNA:LNA duplexes. The RNA-mimicking of LNA was reflected with regard to the N-type conformational restriction of the monomers and to the secondary structure of the LNA:RNA duplex.

[0141] LNAs also form duplexes with complementary DNA, RNA or LNA with high thermal affinities. Circular dichroism (CD) spectra show that duplexes involving fully modified LNA (esp. LNA:RNA) structurally resemble an A-form RNA:RNA duplex. Nuclear magnetic resonance (NMR) examination of an LNA:DNA duplex confirmed the 3'-endo conformation of an LNA monomer. Recognition of double-stranded DNA has also been demonstrated suggesting strand invasion by LNA. Studies of mismatched sequences show that LNAs obey the Watson-Crick base pairing rules with generally improved selectivity compared to the corresponding unmodified reference strands.

[0142] Novel types of LNA-oligomeric compounds, as well as the LNAs, are useful in a wide range of diagnostic and therapeutic applications. Among these are antisense applications, PCR applications, strand-displacement oligomers, substrates for nucleic acid polymerases and generally as nucleotide based drugs.

[0143] Potent and nontoxic antisense oligonucleotides containing LNAs have been described (Wahlestedt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2000, 97, 5633-5638.) The authors have demonstrated that LNAs confer several desired properties to antisense agents. LNA/DNA copolymers were not degraded readily in blood serum and cell extracts. LNA/DNA copolymers exhibited potent antisense activity in assay systems as disparate as G-protein-coupled receptor signaling in living rat brain and detection of reporter genes in *Escherichia coli*. Lipofectin-mediated efficient delivery of LNA into living human breast cancer cells has also been accomplished.


[0144] The synthesis and preparation of the LNA monomers adenine, cytosine, guanine, 5-methyl-cytosine, thymine and uracil, along with their oligomerization, and nucleic acid recognition properties have been described (Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630). LNAs and preparation thereof are also described in WO 98/39352 and WO 99/14226.

[0145] The first analogs of LNA, phosphorothioate-LNA and 2'-thio-LNAs, have also been prepared (Kumar et al.,

Bioorg. Med. Chem. Lett., 1998, 8, 2219-2222). Preparation of locked nucleoside analogs containing oligodeoxyribo-nucleotide duplexes as substrates for nucleic acid polymerases has also been described (Wengel et al., PCT International Application WO 98-DK393 19980914). Furthermore, synthesis of 2'-amino-LNA, a novel conformationally restricted high-affinity oligonucleotide analog with a handle has been described in the art (Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039). In addition, 2'-Amino- and 2'-methylamino-LNA's have been prepared and the thermal stability of their duplexes with complementary RNA and DNA strands has been previously reported.

[0146] One group has added an additional methylene group to the LNA 2',4'-bridging group (e.g. 4'-CH₂—CH₂—O-2' (ENA), Kaneko et al., United States Patent Application Publication No.: US 2002/0147332, also see Japanese Patent Application HEI-11-33863, Feb. 12, 1999).

[0147] Further oligonucleotide mimetics have been prepared to include bicyclic and tricyclic nucleoside analogs having the formulas (amidite monomers shown):




[0148] (see Steffens et al., *Helv. Chim. Acta*, 1997, 80, 2426-2439; Steffens et al., *J. Am. Chem. Soc.*, 1999, 121, 3249-3255; and Renneberg et al., *J. Am. Chem. Soc.*, 2002, 124, 5993-6002). These modified nucleoside analogs have been oligomerized using the phosphoramidite approach and the resulting oligomeric compounds containing tricyclic nucleoside analogs have shown increased thermal stabilities (T_m 's) when hybridized to DNA, RNA and itself. Oligomeric compounds containing bicyclic nucleoside analogs have shown thermal stabilities approaching that of DNA duplexes.

[0149] Another class of oligonucleotide mimetic is referred to as phosphonomonoester nucleic acids incorporate a phosphorus group in a backbone the backbone. This class of oligonucleotide mimetic is reported to have useful physical and biological and pharmacological properties in the areas of inhibiting gene expression (antisense oligonucleotides, ribozymes, sense oligonucleotides and triplex-

forming oligonucleotides), as probes for the detection of nucleic acids and as auxiliaries for use in molecular biology.

[0150] The general formula (for definitions of Markush variables see: U.S. Pat. Nos. 5,874,553 and 6,127,346) is shown below.



[0151] Another oligonucleotide mimetic has been reported wherein the furanosyl ring has been replaced by a cyclobutyl moiety.

[0152] Modified Sugars


[0153] Oligomeric compounds of the invention may also contain one or more substituted sugar moieties. Suitable oligomeric compounds comprise a sugar substituent group selected from: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₂ alkyl or C₂ to C₁₂ alkenyl and alkynyl. Particularly suitable are O((CH₂)_nO)_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON((CH₂)_nCH₃)₂, where n and m are from 1 to about 10. Other suitable oligonucleotides comprise a sugar substituent group selected from: C₁ to C₁₂ lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. One modification includes 2'-methoxyethoxy (2'-O—CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. Another modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylamino-ethoxy-ethyl or 2'-DMAEOE), i.e., 2'-O—CH₂—O—CH₂—N(CH₃)₂.

[0154] Other sugar substituent groups include methoxy (—O—CH₃), aminopropoxy (—OCH₂CH₂CH₂NH₂), allyl (—CH₂—CH=CH₂), —O-allyl (—O—CH₂—CH=CH₂) and fluoro (F). 2'-Sugar substituent groups may be in the arabino (up) position or ribo (down) position. One 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligomeric compound, particularly the 3' position of the sugar on the 3' terminal nucleoside or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligomeric compounds may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents


that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application.

[0155] Further representative sugar substituent groups include groups of formula I_a or II_a:

I_a



II_a




[0156] wherein:

[0157] R_b is O, S or NH;

[0158] R_d is a single bond, O, S or C(=O);

[0159] R_e is C₁-C₁₂ alkyl, N(R_k)(R_m), N(R_k)(R_n), N=C(R_p)(R_q), N=C(R_p)(R_r) or has formula III_a;

III_a



[0160] R_p and R_q are each independently hydrogen or C₁-C₁₂ alkyl;

[0161] R_r is —R_x—R_y;


[0162] each R_s, R_t, R_u and R_v is, independently, hydrogen, C(O)R_w, substituted or unsubstituted C₁-C₁₂ alkyl, substituted or unsubstituted C₂-C₁₂ alkenyl, substituted or unsubstituted C₂-C₁₂ alkynyl, alkylsulfonyl, arylsulfonyl, a chemical functional group or a conjugate group, wherein the substituent groups are selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl;

[0163] or optionally, R_u and R_v, together form a phthalimido moiety with the nitrogen atom to which they are attached;

[0164] each R_w is, independently, substituted or unsubstituted C₁-C₁₂alkyl, trifluoromethyl, cyanoethoxy, methoxy, ethoxy, t-butoxy, allyloxy, 9-fluorenylmethoxy, 2-(trimethylsilyl)-ethoxy, 2,2,2-trichloroethoxy, benzyloxy, butyryl, iso-butyryl, phenyl or aryl;

- [0165] R_k is hydrogen, a nitrogen protecting group or $-R_x-R_y$;
- [0166] R_p is hydrogen, a nitrogen protecting group or $-R_x-R_y$;
- [0167] R_x is a bond or a linking moiety;
- [0168] R_y is a chemical functional group, a conjugate group or a solid support medium;
- [0169] each R_m and R_n is, independently, H, a nitrogen protecting group, substituted or unsubstituted C_1-C_{12} alkyl, substituted or unsubstituted C_2-C_{12} alkenyl, substituted or unsubstituted C_2-C_{12} alkynyl, wherein the substituent groups are selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, alkynyl; NH_3^+ , $N(R_u)(R_v)$ guanidino and acyl where said acyl is an acid amide or an ester;
- [0170] R_m and R_n , together, are a nitrogen protecting group, are joined in a ring structure that optionally includes an additional heteroatom selected from N and O or are a chemical functional group;
- [0171] R_i is OR_z , SR_z , or $N(R_z)_2$;
- [0172] each R_z is, independently, H, C_1-C_8 alkyl, C_1-C_8 haloalkyl, $C(=NH)N(H)R_u$, $C(=O)N(H)R_u$ or $OC(=O)N(H)R_u$;
- [0173] R_f , R_g and R_h comprise a ring system having from about 4 to about 7 carbon atoms or having from about 3 to about 6 carbon atoms and 1 or 2 heteroatoms wherein said heteroatoms are selected from oxygen, nitrogen and sulfur and wherein said ring system is aliphatic, unsaturated aliphatic, aromatic, or saturated or unsaturated heterocyclic;
- [0174] R_j is alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, aryl having 6 to about 14 carbon atoms, $N(R_k)(R_m)OR_k$, halo, SR_k or CN;
- [0175] m_a is 1 to about 10;
- [0176] each m_b is, independently, 0 or 1;
- [0177] m_c is 0 or an integer from 1 to 10;
- [0178] m_d is an integer from 1 to 10;
- [0179] m_e is from 0, 1 or 2; and
- [0180] provided that when m_c is 0, m_d is greater than 1.
- [0181] Representative substituents groups of Formula I are disclosed in U.S. patent application Ser. No. 09/130,973, filed Aug. 7, 1998, entitled "Capped 2'-Oxyethoxy Oligonucleotides."
- [0182] Representative cyclic substituent groups of Formula II are disclosed in U.S. patent application Ser. No. 09/123,108, filed Jul. 27, 1998, entitled "RNA Targeted 2'-Oligomeric compounds that are Conformationally Preorganized."
- [0183] Particularly suitable sugar substituent groups include $O((CH_2)_nO)_mCH_3$, $O(CH_2)_nOCH_3$, $O(CH_2)_nNH_2$, $O(CH_2)_nCH_3$, $O(CH_2)_nONH_2$ and $O(CH_2)_nON((CH_2)_nCH_3)_2$, where n and m are from 1 to about 10.
- [0184] Representative guanidino substituent groups that are shown in formula III and IV are disclosed in co-owned U.S. patent application Ser. No. 09/349,040, entitled "Functionalized Oligomers" filed Jul. 7, 1999.
- [0185] Representative acetamido substituent groups are disclosed in U.S. Pat. No. 6,147,200.
- [0186] Representative dimethylaminoethoxyethyl substituent groups are disclosed in International Patent Application PCT/US99/17895, entitled "2'-O-Dimethylaminoethoxy-ethyl-Oligomeric compounds", filed Aug. 6, 1999.
- [0187] Modified Nucleobases/Naturally Occurring Nucleobases
- [0188] Oligomeric compounds may also include nucleobase (often referred to in the art simply as "base" or "heterocyclic base moiety") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases also referred herein as heterocyclic base moieties include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ($-C\equiv C-CH_3$) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine.
- [0189] Heterocyclic base moieties may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J. I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., Angewandte Chemie, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y. S., Chapter 15, Antisense Research and Applications, pages 289-302, Crooke, S. T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2° C. (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., eds., Antisense Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently suitable base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0190] In one aspect of the present invention oligomeric compounds are prepared having polycyclic heterocyclic compounds in place of one or more heterocyclic base moieties. A number of tricyclic heterocyclic compounds have been previously reported. These compounds are routinely used in antisense applications to increase the binding properties of the modified strand to a target strand. The most studied modifications are targeted to guanosines hence they have been termed G-clamps or cytidine analogs. Many of these polycyclic heterocyclic compounds have the general formula:



[0191] Representative cytosine analogs that make 3 hydrogen bonds with a guanosine in a second strand include 1,3-diazaphenoxazine-2-one ($R_{10}=O$, $R_{11}-R_{14}=H$) (Kurchavov, et al., *Nucleosides and Nucleotides*, 1997, 16, 1837-1846), 1,3-diazaphenothiazine-2-one ($R_{10}=S$, $R_{11}-R_{14}=H$) (Lin, K.-Y.; Jones, R. J.; Matteucci, M. J. *Am. Chem. Soc.* 1995, 117, 3873-3874) and 6,7,8,9-tetrafluoro-1,3-diazaphenoxazine-2-one ($R_{10}=O$, $R_{11}-R_{14}=F$) (Wang, J.; Lin, K.-Y.; Matteucci, M. *Tetrahedron Lett.* 1998, 39, 8385-8388). Incorporated into oligonucleotides these base modifications were shown to hybridize with complementary guanine and the latter was also shown to hybridize with adenine and to enhance helical thermal stability by extended stacking interactions (also see U.S. Patent Application entitled "Modified Peptide Nucleic Acids" filed May 24, 2002, Ser. No. 10/155,920; and U.S. Patent Application entitled "Nuclease Resistant Chimeric Oligonucleotides" filed May 24, 2002, Ser. No. 10/013,295).

[0192] Further helix-stabilizing properties have been observed when a cytosine analog/substitute has an aminoethoxy moiety attached to the rigid 1,3-diazaphenoxazine-2-one scaffold ($R_{10}=O$, $R_1=-O-(CH_2)_2-NH_2$, $R_{12-14}=H$) (Lin, K.-Y.; Matteucci, M. J. *Am. Chem. Soc.* 1998, 120, 8531-8532). Binding studies demonstrated that a single incorporation could enhance the binding affinity of a model oligonucleotide to its complementary target DNA or RNA with a ΔT_m of up to 18° relative to 5-methyl cytosine ($dC5^{me}$), which is the highest known affinity enhancement for a single modification, yet. On the other hand, the gain in helical stability does not compromise the specificity of the oligonucleotides. The T_m data indicate an even greater discrimination between the perfect match and mismatched sequences compared to $dC5^{me}$. It was suggested that the tethered amino group serves as an additional hydrogen bond donor to interact with the Hoogsteen face, namely the O6, of a complementary guanine thereby forming 4 hydrogen bonds. This means that the increased affinity of G-clamp is

mediated by the combination of extended base stacking and additional specific hydrogen bonding.

[0193] Further tricyclic heterocyclic compounds and methods of using them that are amenable to the present invention are disclosed in U.S. Pat. No. 6,028,183, which issued on May 22, 2000, and U.S. Pat. No. 6,007,992, which issued on Dec. 28, 1999.

[0194] The enhanced binding affinity of the phenoxazine derivatives together with their uncompromised sequence specificity makes them valuable nucleobase analogs for the development of more potent antisense-based drugs. In fact, promising data have been derived from in vitro experiments demonstrating that heptanucleotides containing phenoxazine substitutions are capable to activate RNaseH, enhance cellular uptake and exhibit an increased antisense activity (Lin et al., *J. Am. Chem. Soc.*, 1998, 120, 8531-8532). The activity enhancement was even more pronounced in case of G-clamp, as a single substitution was shown to significantly improve the in vitro potency of a 20mer 2'-deoxyphosphorothioate oligonucleotides (Flanagan et al., *Proc. Natl. Acad. Sci. USA*, 1999, 96, 3513-3518). Nevertheless, to optimize oligonucleotide design and to better understand the impact of these heterocyclic modifications on the biological activity, it is important to evaluate their effect on the nuclease stability of the oligomers.

[0195] Further modified polycyclic heterocyclic compounds useful as heterocyclic bases are disclosed in but not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos. 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,434,257; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,645,985; 5,646,269; 5,750,692; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, and U.S. patent application Ser. No. 09/996,292 filed Nov. 28, 2001.

[0196] Conjugates

[0197] Another substitution that can be appended to the oligomeric compounds of the invention involves the linkage of one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the resulting oligomeric compounds. In one embodiment such modified oligomeric compounds are prepared by covalently attaching conjugate groups to functional groups such as hydroxyl or amino groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, poly-ethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include cholesterol, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed Oct. 23, 1992.

[0198] Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al.,

Proc. Natl. Acad. Sci. USA, 1989, 86, 6553-6556), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxcholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

[0199] The oligomeric compounds of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansyl-sarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folic acid, a benzothiazide, chlorothiazide, a diazepam, indomethacin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in U.S. patent application Ser. No. 09/334,130 (filed Jun. 15, 1999).

[0200] Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

[0201] Chimeric Oligomeric Compounds

[0202] It is not necessary for all positions in a oligomeric compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single oligomeric compound or even at a single monomeric subunit such as a nucleoside within a oligomeric compound. The present invention also includes oligomeric compounds which are chimeric oligomeric compounds. "Chimeric" oligomeric compounds or "chimeras," in the context of this invention, are oligomeric compounds which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of a nucleic acid based oligomer.

[0203] Chimeric oligomeric compounds typically contain at least one region modified so as to confer increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target


nucleic acid. An additional region of the oligomeric compound may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligomeric compounds when chimeras are used, compared to for example phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

[0204] Chimeric oligomeric compounds of the invention may be formed as composite structures of two or more oligonucleotides, oligonucleotide analogs, oligonucleosides and/or oligonucleotide mimetics as described above. Such oligomeric compounds have also been referred to in the art as hybrids hemimers, gapmers or inverted gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922.


[0205] 3'-endo Modifications

[0206] In one aspect of the present invention oligomeric compounds include nucleosides synthetically modified to induce a 3'-endo sugar conformation. A nucleoside can incorporate synthetic modifications of the heterocyclic base, the sugar moiety or both to induce a desired 3'-endo sugar conformation. These modified nucleosides are used to mimic RNA like nucleosides so that particular properties of an oligomeric compound can be enhanced while maintaining the desirable 3'-endo conformational geometry. There is an apparent preference for an RNA type duplex (A form helix, predominantly 3'-endo) as a requirement (e.g. trigger) of RNA interference which is supported in part by the fact that duplexes composed of 2'-deoxy-2'-F-nucleosides appears efficient in triggering RNAi response in the *C. elegans* system. Properties that are enhanced by using more stable 3'-endo nucleosides include but aren't limited to modulation of pharmacokinetic properties through modification of protein binding, protein off-rate, absorption and clearance; modulation of nuclease stability as well as chemical stability; modulation of the binding affinity and specificity of the oligomer (affinity and specificity for enzymes as well as for complementary sequences); and increasing efficacy of RNA cleavage. The present invention provides oligomeric triggers of RNAi having one or more nucleosides modified in such a way as to favor a C3'-endo conformation.

Scheme 1



-continued



[0207] Nucleoside conformation is influenced by various factors including substitution at the 2', 3' or 4'-positions of the pentofuranosyl sugar. Electronegative substituents generally prefer the axial positions, while sterically demanding substituents generally prefer the equatorial positions (Principles of Nucleic Acid Structure, Wolfgang Sanger, 1984, Springer-Verlag.) Modification of the 2' position to favor the 3'-endo conformation can be achieved while maintaining the 2'-OH as a recognition element, as illustrated in FIG. 2, below (Gallo et al., *Tetrahedron* (2001), 57, 5707-5713. Harry-O'kuru et al., *J. Org. Chem.*, (1997), 62(6), 1754-1759 and Tang et al., *J. Org. Chem.* (1999), 64, 747-754.) Alternatively, preference for the 3'-endo conformation can be achieved by deletion of the 2'-OH as exemplified by 2'-deoxy-2' F-nucleosides (Kawasaki et al., *J. Med. Chem.* (1993), 36, 831-841), which adopts the 3'-endo conformation positioning the electronegative fluorine atom in the axial position. Other modifications of the ribose ring, for example substitution at the 4'-position to give 4'-F modified nucleosides (Guillem et al., *Bioorganic and Medicinal Chemistry Letters* (1995), 5, 1455-1460 and Owen et al., *J. Org. Chem.* (1976), 41, 3010-3017), or for example modification to yield methanocarba nucleoside analogs (Jacobson et al., *J. Med. Chem. Lett.* (2000), 43, 2196-2203 and Lee et al., *Bioorganic and Medicinal Chemistry Letters* (2001), 11, 1333-1337) also induce preference for the 3'-endo conformation. Along similar lines, oligomeric triggers of RNAi response might be composed of one or more nucleosides modified in such a way that conformation is locked into a C3'-endo type conformation, i.e. Locked Nucleic Acid (LNA, Singh et al, *Chem. Commun.* (1998), 4, 455-456), and ethylene bridged Nucleic Acids (ENA, Morita et al, *Bioorganic & Medicinal Chemistry Letters* (2002), 12, 73-76.) Examples of modified nucleosides amenable to the present invention are shown below in Table I. These examples are meant to be representative and not exhaustive.

TABLE I





TABLE I-continued



[0208] A preferred conformation of modified nucleosides and their oligomers can be estimated by various methods such as molecular dynamics calculations, nuclear magnetic resonance spectroscopy and CD measurements. Hence, modifications predicted to induce RNA like conformations, A-form duplex geometry in an oligomeric context, are selected for use in the modified oligonucleotides of the present invention. The synthesis of numerous of the modified nucleosides amenable to the present invention are known in the art (see for example, *Chemistry of Nucleosides and Nucleotides Vol 1-3*, ed. Leroy B. Townsend, 1988, Plenum press., and the examples section below.) Nucleosides known to be inhibitors/substrates for RNA dependent RNA polymerases (for example HCV NS5B

[0209] In one aspect, the present invention is directed to oligonucleotides that are prepared having enhanced properties compared to native RNA against nucleic acid targets. A target is identified and an oligonucleotide is selected having an effective length and sequence that is complementary to a portion of the target sequence. Each nucleoside of the

selected sequence is scrutinized for possible enhancing modifications. Another modification would be the replacement of one or more RNA nucleosides with nucleosides that have the same 3'-endo conformational geometry. Such modifications can enhance chemical and nuclease stability relative to native RNA while at the same time being much cheaper and easier to synthesize and/or incorporate into an oligonucleotide. The selected sequence can be further divided into regions and the nucleosides of each region evaluated for enhancing modifications that can be the result of a chimeric configuration. Consideration is also given to the 5' and 3'-termini as there are often advantageous modifications that can be made to one or more of the terminal nucleosides. The oligomeric compounds of the present invention include at least one 5'-modified phosphate group on a single strand or on at least one 5'-position of a double stranded sequence or sequences. Further modifications are also considered such as internucleoside linkages, conjugate groups, substitute sugars or bases, substitution of one or more nucleosides with nucleoside mimetics and any other modification that can enhance the selected sequence for its intended target.

[0210] The terms used to describe the conformational geometry of homoduplex nucleic acids are "A Form" for RNA and "B Form" for DNA. The respective conformational geometry for RNA and DNA duplexes was determined from X-ray diffraction analysis of nucleic acid fibers (Arnott and Hukins, *Biochem. Biophys. Res. Comm.*, 1970, 47, 1504.) In general, RNA:RNA duplexes are more stable and have higher melting temperatures (T_m 's) than DNA:DNA duplexes (Sanger et al., *Principles of Nucleic Acid Structure*, 1984, Springer-Verlag; New York, N.Y.; Lesnik et al., *Biochemistry*, 1995, 34, 10807-10815; Conte et al., *Nucleic Acids Res.*, 1997, 25, 2627-2634). The increased stability of RNA has been attributed to several structural features, most notably the improved base stacking interactions that result from an A-form geometry (Searle et al., *Nucleic Acids Res.*, 1993, 21, 2051-2056). The presence of the 2' hydroxyl in RNA biases the sugar toward a C3' endo pucker, i.e., also designated as Northern pucker, which causes the duplex to favor the A-form geometry. In addition, the 2' hydroxyl groups of RNA can form a network of water mediated hydrogen bonds that help stabilize the RNA duplex (Egli et al., *Biochemistry*, 1996, 35, 8489-8494). On the other hand, deoxy nucleic acids prefer a C2' endo sugar pucker, i.e., also known as Southern pucker, which is thought to impart a less stable B-form geometry (Sanger, W. (1984) *Principles of Nucleic Acid Structure*, Springer-Verlag, New York, N.Y.). As used herein, B-form geometry is inclusive of both C2'-endo pucker and O4'-endo pucker. This is consistent with Berger, et. al., *Nucleic Acids Research*, 1998, 26, 2473-2480, who pointed out that in considering the furanose conformations which give rise to B-form duplexes consideration should also be given to a O4'-endo pucker contribution.

[0211] DNA:RNA hybrid duplexes, however, are usually less stable than pure RNA:RNA duplexes, and depending on their sequence may be either more or less stable than DNA:DNA duplexes (Searle et al., *Nucleic Acids Res.*, 1993, 21, 2051-2056). The structure of a hybrid duplex is intermediate between A- and B-form geometries, which may result in poor stacking interactions (Lane et al., *Eur. J. Biochem.*, 1993, 215, 297-306; Fedoroff et al., *J. Mol. Biol.*, 1993, 233, 509-523; Gonzalez et al., *Biochemistry*, 1995, 34,

4969-4982; Horton et al., *J. Mol. Biol.*, 1996, 264, 521-533). The stability of the duplex formed between a target RNA and a synthetic sequence is central to therapies such as but not limited to antisense and RNA interference as these mechanisms require the binding of a synthetic oligonucleotide strand to an RNA target strand. In the case of antisense, effective inhibition of the mRNA requires that the antisense DNA have a very high binding affinity with the mRNA. Otherwise the desired interaction between the synthetic oligonucleotide strand and target mRNA strand will occur infrequently, resulting in decreased efficacy.

[0212] One routinely used method of modifying the sugar pucker is the substitution of the sugar at the 2'-position with a substituent group that influences the sugar geometry. The influence on ring conformation is dependant on the nature of the substituent at the 2'-position. A number of different substituents have been studied to determine their sugar pucker effect. For example, 2'-halogens have been studied showing that the 2'-fluoro derivative exhibits the largest population (65%) of the C3'-endo form, and the 2'-iodo exhibits the lowest population (7%). The populations of adenosine (2'-OH) versus deoxyadenosine (2'-H) are 36% and 19%, respectively. Furthermore, the effect of the 2'-fluoro group of adenosine dimers (2'-deoxy-2'-fluoroadenosine-2'-deoxy-2'-fluoro-adenosine) is further correlated to the stabilization of the stacked conformation.

[0213] As expected, the relative duplex stability can be enhanced by replacement of 2'-OH groups with 2'-F groups thereby increasing the C3'-endo population. It is assumed that the highly polar nature of the 2'-F bond and the extreme preference for C3'-endo pucker may stabilize the stacked conformation in an A-form duplex. Data from UV hypochromicity, circular dichroism, and ^1H NMR also indicate that the degree of stacking decreases as the electronegativity of the halo substituent decreases. Furthermore, steric bulk at the 2'-position of the sugar moiety is better accommodated in an A-form duplex than a B-form duplex. Thus, a 2'-substituent on the 3'-terminus of a dinucleoside monophosphate is thought to exert a number of effects on the stacking conformation: steric repulsion, furanose pucker preference, electrostatic repulsion, hydrophobic attraction, and hydrogen bonding capabilities. These substituent effects are thought to be determined by the molecular size, electronegativity, and hydrophobicity of the substituent. Melting temperatures of complementary strands is also increased with the 2'-substituted adenosine diphosphates. It is not clear whether the 3'-endo preference of the conformation or the presence of the substituent is responsible for the increased binding. However, greater overlap of adjacent bases (stacking) can be achieved with the 3'-endo conformation.

[0214] One synthetic 2'-modification that imparts increased nuclease resistance and a very high binding affinity to nucleotides is the 2-methoxyethoxy (2'-MOE, 2'-OCH₂CH₂OCH₃) side chain (Baker et al., *J. Biol. Chem.*, 1997, 272, 11944-12000). One of the immediate advantages of the 2'-MOE substitution is the improvement in binding affinity, which is greater than many similar 2' modifications such as O-methyl, O-propyl, and O-aminopropyl. Oligonucleotides having the 2'-O-methoxyethyl substituent also have been shown to be antisense inhibitors of gene expression with promising features for in vivo use (Martin, P., *Helv. Chim. Acta*, 1995, 78, 486-504; Altmann et al., *Chimia*, 1996, 50, 168-176; Altmann et al., *Biochem. Soc.*

Trans., 1996, 24, 630-637; and Altmann et al., Nucleosides Nucleotides, 1997, 16, 917-926). Relative to DNA, the oligonucleotides having the 2'-MOE modification displayed improved RNA affinity and higher nuclease resistance. Chimeric oligonucleotides having 2'-MOE substituents in the wing nucleosides and an internal region of deoxy-phosphorothioate nucleotides (also termed a gapped oligonucleotide or gapmer) have shown effective reduction in the growth of tumors in animal models at low doses. 2'-MOE substituted oligonucleotides have also shown outstanding promise as antisense agents in several disease states. One such MOE substituted oligonucleotide is presently being investigated in clinical trials for the treatment of CMV retinitis.

[0215] Chemistries Defined

[0216] Unless otherwise defined herein, alkyl means C₁-C₁₂, C₁-C₈, or C₁-C₆, straight or (where possible) branched chain aliphatic hydrocarbyl.

[0217] Unless otherwise defined herein, heteroalkyl means C₁-C₁₂, C₁-C₈, or C₁-C₆, straight or (where possible) branched chain aliphatic hydrocarbyl containing at least one or about 1 to about 3 hetero atoms in the chain, including the terminal portion of the chain. Suitable heteroatoms include N, O and S.

[0218] Unless otherwise defined herein, cycloalkyl means C₃-C₁₂, C₃-C₈, or C₃-C₆, aliphatic hydrocarbyl ring.

[0219] Unless otherwise defined herein, alkenyl means C₂-C₁₂, C₂-C₈, or C₂-C₆ alkenyl, which may be straight or (where possible) branched hydrocarbyl moiety, which contains at least one carbon-carbon double bond.

[0220] Unless otherwise defined herein, alkynyl means C₂-C₁₂, C₂-C₈, or C₂-C₆ alkynyl, which may be straight or (where possible) branched hydrocarbyl moiety, which contains at least one carbon-carbon triple bond.

[0221] Unless otherwise defined herein, heterocycloalkyl means a ring moiety containing at least three ring members, at least one of which is carbon, and of which 1, 2 or three ring members are other than carbon. The number of carbon atoms can vary from 1 to about 12, or 1 to about 6, and the total number of ring members can vary from three to about 15, or from about 3 to about 8. Suitable ring heteroatoms are N, O and S. Suitable heterocycloalkyl groups include morpholino, thiomorpholino, piperidinyl, piperazinyl, homopiperidinyl, homopiperazinyl, homomorpholino, homothiomorpholino, pyrrolodinyl, tetrahydrooxazolyl, tetrahydroimidazolyl, tetrahydrothiazolyl, tetrahydroisoxazolyl, tetrahydropyrazolyl, furanyl, pyranyl, and tetrahydroisothiazolyl.

[0222] Unless otherwise defined herein, aryl means any hydrocarbon ring structure containing at least one aryl ring. Suitable aryl rings have about 6 to about 20 ring carbons. Suitable aryl rings include phenyl, naphthyl, anthracenyl, and phenanthrenyl.

[0223] Unless otherwise defined herein, hetaryl means a ring moiety containing at least one fully unsaturated ring, the ring consisting of carbon and non-carbon atoms. The ring system can contain about 1 to about 4 rings. The number of carbon atoms can vary from 1 to about 12, or 1 to about 6, and the total number of ring members can vary from three

to about 15, or from about 3 to about 8. Suitable ring heteroatoms are N, O and S. Suitable hetaryl moieties include, but are not limited to, pyrazolyl, thiophenyl, pyridyl, imidazolyl, tetrazolyl, pyridyl, pyrimidinyl, purinyl, quinazoliny, quinoxaliny, benzimidazolyl, benzothiofenyl, and the like.

[0224] Unless otherwise defined herein, where a moiety is defined as a compound moiety, such as hetarylalkyl (hetaryl and alkyl), aralkyl (aryl and alkyl), etc., each of the submoieties is as defined herein.

[0225] Unless otherwise defined herein, an electron withdrawing group is a group, such as the cyano or isocyanato group that draws electronic charge away from the carbon to which it is attached. Other electron withdrawing groups of note include those whose electronegativities exceed that of carbon, for example halogen, nitro, or phenyl substituted in the ortho- or para-position with one or more cyano, isothiocyanato, nitro or halo groups.


[0226] Unless otherwise defined herein, the terms halogen and halo have their ordinary meanings. Suitable halo (halogen) substituents are Cl, Br, and I.

[0227] The aforementioned optional substituents are, unless otherwise herein defined, suitable substituents depending upon desired properties. Included are halogens (Cl, Br, I), alkyl, alkenyl, and alkynyl moieties, NO₂, NH₃ (substituted and unsubstituted), acid moieties (e.g.—CO₂H, —OSO₃H₂, etc.), heterocycloalkyl moieties, hetaryl moieties, aryl moieties, etc. In all the preceding formulae, the squiggle (~) indicates a bond to an oxygen or sulfur of the 5'-phosphate.

[0228] Phosphate protecting groups include those described in U.S. Pat. No. 5,760,209, U.S. Pat. No. 5,614,621, U.S. Pat. No. 6,051,699, U.S. Pat. No. 6,020,475, U.S. Pat. No. 6,326,478, U.S. Pat. No. 6,169,177, U.S. Pat. No. 6,121,437, U.S. Pat. No. 6,465,628.

[0229] The present invention discloses novel nucleosides comprising bicyclic sugar moieties and oligomeric compounds comprising at least one such nucleoside. The bicyclic sugar moieties have a locked 3'-endo sugar conformation which provides nucleosides having A form, RNA like without having some of the undesirable properties associated with native RNA nucleosides. One of the immediate advantages is the nuclease stability gained by replacing RNA nucleosides with locked e.g. bicyclic sugar nucleosides. The bicyclic sugar modified nucleosides are also expected to have enhanced binding affinity that has been previously reported for LNA (3-8° C. per modification).

[0230] The nucleosides of the present invention have a bridge from the 2', to the 4'-position defined by -Q₁-Q₂-Q₃- as shown below in structure I.



I

[0231] wherein:

[0232] Bx is a heterocyclic base moiety;

[0233] each T_a and T_b is, independently, H, a hydroxyl protecting group, an activated phosphorus moiety, a conjugate group or a covalent attachment to a support medium;

[0234] -Q₁-Q₂-Q₃- is —CH₂—N(R₁)-CH₂—, —C(=O)—N(R₁)-CH₂—, —CH₂—O—N(R₁)- or N(R₁)—O—CH₂—; and


[0235] R₁ is C₁-C₁₂ alkyl or an amino protecting group.

[0236] The present invention also provides for oligomeric compounds having at least one nucleoside having a bicyclic sugar moiety of structure I. The nucleosides of structure I can be used to modify the properties an oligomeric compound. Such a nucleoside can be put into an oligonucleotide or an oligonucleoside in a single position or at multiple positions to create a hemimer, blockmer, gapmer or a more complicated alternating or chimeric oligomeric compound. The nucleosides having structure I can also be used to modify properties of oligomeric compounds that comprise more complicated chemistries to prepare oligomeric compounds that are essentially oligonucleotide mimics such as, for example, peptide nucleic acids (PNA), morpholino nucleic acids, cyclohexenyl nucleic acids (CeNA), anhydrohexitol nucleic acids, locked nucleic acids (LNA and ENA), bicyclic and tricyclic nucleic acids, phosphonomonoester nucleic acids and cyclobutyl nucleic acids.

[0237] The compositions of the present invention illustrate the use of activated phosphorus compositions (e.g. compounds having activated phosphorus-containing substituent groups) in coupling reactions. As used herein, the term "activated phosphorus composition" includes monomers and oligomers that have an activated phosphorus-containing substituent group that is reactive with a hydroxyl group of another monomeric or oligomeric compound to form a phosphorus-containing internucleotide linkage. Such activated phosphorus groups contain activated phosphorus atoms in P^{III} valence state and are known in the art and include, but are not limited to, phosphoramidite, H-phosphonate, phosphate triesters and chiral auxiliaries. One synthetic solid phase synthesis utilizes phosphoramidites as activated phosphates. The phosphoramidites utilize P^{III} chemistry. The intermediate phosphite compounds are subsequently oxidized to the P^V state using known methods to yield, in some embodiments, phosphodiester or phosphorothioate internucleotide linkages. Additional activated phosphates and phosphites are disclosed in Tetrahedron Report Number 309 (Beaucage and Iyer, Tetrahedron, 1992, 48, 2223-2311).

[0238] Activated phosphorus groups are useful in the preparation of a wide range of oligomeric compounds including but not limited to oligonucleosides and oligonucleotides as well as oligonucleotides that have been modified or conjugated with other groups at the base or sugar or both. Also included are oligonucleotide mimetics including but not limited to peptide nucleic acids (PNA), morpholino nucleic acids, cyclohexenyl nucleic acids (CeNA), anhydrohexitol nucleic acids, locked nucleic acids (LNA and ENA), bicyclic and tricyclic nucleic acids, phosphonomonoester nucleic acids and cyclobutyl nucleic

acids. A representative example of one type of oligomer synthesis that utilizes the coupling of an activated phosphorus group with a reactive hydroxyl group is the widely used phosphoramidite approach. A phosphoramidite synthon is reacted under appropriate conditions with a reactive hydroxyl group to form a phosphite linkage that is further oxidized to a phosphodiester or phosphorothioate linkage. This approach commonly utilizes nucleoside phosphoramidites of the formula:



[0239] wherein:

[0240] Bx' is an optionally protected heterocyclic base moiety;

[0241] R₁ is, independently, H or an optionally protected sugar substituent group;

[0242] T₃ is H, a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

[0243] L₁ is N(R₁)R₂;

[0244] R₁ and R₂ is, independently, C₁-C₁₂ straight or branched chain alkyl;

[0245] or R₁ and R₂ are joined together to form a 4- to 7-membered heterocyclic ring system including the nitrogen atom to which R₁ and R₂ are attached, wherein the ring system optionally includes at least one additional heteroatom selected from O, N and S;

[0246] L₂ is Pg-O—, Pg-S—, C₁-C₁₂ straight or branched chain alkyl, CH₃(CH₂)₀₋₁₀-O— or —NR₅R₆;

[0247] Pg is a protecting/blocking group; and

[0248] R₅ and R₆ is, independently, hydrogen, C₁-C₁₂ straight or branched chain alkyl, cycloalkyl or aryl;

[0249] or optionally, R₅ and R₆, together with the nitrogen atom to which they are attached form a cyclic moiety that may include an additional heteroatom selected from O, S and N; or

[0250] L₁ and L₂ together with the phosphorus atom to which L₁ and L₂ are attached form a chiral auxiliary.

[0251] Groups that are attached to the phosphorus atom of internucleotide linkages before and after oxidation (L₁ and L₂) can include nitrogen containing cyclic moieties such as morpholine. Such oxidized internucleoside linkages include a phosphoromorpholidothioate linkage (Wilk et al., Nucleosides and nucleotides, 1991, 10, 319-322). Further cyclic moieties amenable to the present invention include mono-, bi- or tricyclic ring moieties which may be substituted with groups such as oxo, acyl, alkoxy, alkoxy carbonyl, alkyl,

alkenyl, alkynyl, amino, amido, azido, aryl, heteroaryl, carboxylic acid, cyano, guanidino, halo, haloalkyl, haloalkoxy, hydrazino, ODMT, alkylsulfonyl, nitro, sulfide, sulfone, sulfonamide, thiol and thioalkoxy. A suitable bicyclic ring structure that includes nitrogen is phthalimido.

[0252] Hybridization

[0253] In the context of this invention, "hybridization" means the pairing of complementary strands of oligomeric compounds. In the present invention, one mechanism of pairing involves hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases (nucleobases) of the strands of oligomeric compounds. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. Hybridization can occur under varying circumstances.

[0254] An antisense oligomeric compound is specifically hybridizable when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a loss of activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense oligomeric compound to non-target nucleic acid sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays.

[0255] In the present invention the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which an oligomeric compound of the invention will hybridize to its target sequence, but to a minimal number of other sequences. Stringent conditions are sequence-dependent and will vary with different circumstances and in the context of this invention, "stringent conditions" under which oligomeric compounds hybridize to a target sequence are determined by the nature and composition of the oligomeric compounds and the assays in which they are being investigated.

[0256] "Complementary," as used herein, refers to the capacity for precise pairing of two nucleobases regardless of where the two are located. For example, if a nucleobase at a certain position of an oligomeric compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, the target nucleic acid being a DNA, RNA, or oligonucleotide molecule, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be a complementary position. The oligomeric compound and the further DNA, RNA, or oligonucleotide molecule are complementary to each other when a sufficient number of complementary positions in each molecule are occupied by nucleobases which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of precise pairing or complementarity over a sufficient number of nucleobases such that stable and specific binding occurs between the oligonucleotide and a target nucleic acid.

[0257] It is understood in the art that the sequence of an antisense oligomeric compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, an oligonucleotide may hybridize

over one or more segments such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure or hairpin structure). The oligomeric compounds of the present invention can comprise at least 70%, at least 80%, at least 90%, at least 95%, or at least 99% sequence complementarity to a target region within the target nucleic acid sequence to which they are targeted. For example, an antisense oligomeric compound in which 18 of 20 nucleobases of the antisense oligomeric compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleobases may be clustered or interspersed with complementary nucleobases and need not be contiguous to each other or to complementary nucleobases. As such, an antisense oligomeric compound which is 18 nucleobases in length having 4 (four) noncomplementary nucleobases which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid and would thus fall within the scope of the present invention. Percent complementarity of an antisense oligomeric compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., *J. Mol. Biol.*, 1990, 215, 403-410; Zhang and Madden, *Genome Res.*, 1997, 7, 649-656).

[0258] Targets of the Invention

[0259] "Targeting" an antisense oligomeric compound to a particular nucleic acid molecule, in the context of this invention, can be a multistep process. The process usually begins with the identification of a target nucleic acid whose function is to be modulated. This target nucleic acid may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent.

[0260] The targeting process usually also includes determination of at least one target region, segment, or site within the target nucleic acid for the antisense interaction to occur such that the desired effect, e.g., modulation of expression, will result. Within the context of the present invention, the term "region" is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or characteristic. Within regions of target nucleic acids are segments. "Segments" are defined as smaller or sub-portions of regions within a target nucleic acid. "Sites," as used in the present invention, are defined as positions within a target nucleic acid. The terms region, segment, and site can also be used to describe an oligomeric compound of the invention such as for example a gapped oligomeric compound having 3 separate segments.

[0261] Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon." A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even

though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA transcribed from a gene encoding a nucleic acid target, regardless of the sequence(s) of such codons. It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, *i.e.*, 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively).

[0262] The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (*i.e.*, 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (*i.e.*, 5' or 3') from a translation termination codon. Consequently, the "start codon region" (or "translation initiation codon region") and the "stop codon region" (or "translation termination codon region") are all regions which may be targeted effectively with the oligomeric compounds of the present invention.

[0263] The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Within the context of the present invention, a suitable region is the intragenic region encompassing the translation initiation or termination codon of the open reading frame (ORF) of a gene.

[0264] Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA (or corresponding nucleotides on the gene), and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA (or corresponding nucleotides on the gene). The 5' cap site of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap site. It is also suitable to target the 5' cap region.

[0265] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. Targeting splice sites, *i.e.*, intron-exon junctions or exon-intron junctions,

may also be particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also suitable target sites. mRNA transcripts produced via the process of splicing of two (or more) mRNAs from different gene sources are known as "fusion transcripts." It is also known that introns can be effectively targeted using antisense oligomeric compounds targeted to, for example, DNA or pre-mRNA.

[0266] It is also known in the art that alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants." More specifically, "pre-mRNA variants" are transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and exonic sequences.

[0267] Upon excision of one or more exon or intron regions, or portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants." Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative splice variants." If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

[0268] It is also known in the art that variants can be produced through the use of alternative signals to start or stop transcription and that pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate at unique polyA sites. Within the context of the invention, the types of variants described herein are also suitable target nucleic acids.

[0269] The locations on the target nucleic acid to which the oligomeric compounds hybridize are hereinbelow referred to as "suitable target segments." As used herein the term "suitable target segment" is defined as at least an 8-nucleobase portion of a target region to which an active oligomeric compound is targeted. While not wishing to be bound by theory, it is presently believed that these target segments represent portions of the target nucleic acid which are accessible for hybridization.

[0270] Exemplary suitable oligomeric compounds include oligomeric compounds that comprise at least the 8 consecutive nucleobases from the 5'-terminus of a targeted nucleic acid *e.g.* a cellular gene or mRNA transcribed from the gene (the remaining nucleobases being a consecutive stretch of the same oligonucleotide beginning immediately upstream of the 5'-terminus of the compound which is specifically hybridizable to the target nucleic acid and continuing until the oligonucleotide contains from about 8 to about 80 nucleobases). Additional suitable oligomeric compounds are

represented by oligonucleotide sequences that comprise at least the 8 consecutive nucleobases from the 3'-terminus of one of the illustrative compounds (the remaining nucleobases being a consecutive stretch of the same oligonucleotide beginning immediately downstream of the 3'-terminus of the compound which is specifically hybridizable to the target nucleic acid and continuing until the oligonucleotide contains from about 8 to about 80 nucleobases). One having skill in the art armed with the suitable compounds illustrated herein will be able, without undue experimentation, to identify further oligomeric compounds.

[0271] Once one or more target regions, segments or sites have been identified, oligomeric compounds are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

[0272] In accordance with one embodiment of the present invention, a series of nucleic acid duplexes comprising the oligomeric compounds of the present invention and their complements can be designed for a specific target or targets. The ends of the strands may be modified by the addition of one or more natural or modified nucleobases to form an overhang. The sense strand of the duplex is then designed and synthesized as the complement of the antisense strand and may also contain modifications or additions to either terminus. For example, in one embodiment, both strands of the duplex would be complementary over the central nucleobases, each having overhangs at one or both termini.

[0273] For example, a duplex comprising an antisense oligomeric compound having the sequence CGAGAGGCG-GACGGGACCG (SEQ ID NO:1) and having a two-nucleobase overhang of deoxythymidine(dT) would have the following structure:

```

cgagagggcggacgggaccgTT      Antisense Strand (SEQ ID NO:1)
|||||
TTgctctccgcctgcctggc      Complement Strand (SEQ ID NO:2)

```

[0274] RNA strands of the duplex can be synthesized by methods disclosed herein or purchased from various RNA synthesis companies such as for example Dharmacon Research Inc., (Lafayette, Colo.) (see also the section on RNA synthesis below). Once synthesized, the complementary strands are annealed. The single strands are aliquoted and diluted to a concentration of 50 μ M. Once diluted, 30 μ L of each strand is combined with 15 μ L of a 5 \times solution of annealing buffer. The final concentration of the buffer is 100 mM potassium acetate, 30 mM HEPES-KOH pH 7.4, and 2 mM magnesium acetate. The final volume is 75 μ L. This solution is incubated for 1 minute at 90 $^{\circ}$ C. and then centrifuged for 15 seconds. The tube is allowed to sit for 1 hour at 37 $^{\circ}$ C. at which time the dsRNA duplexes are used in experimentation. The final concentration of the dsRNA compound is 20 μ M. This solution can be stored frozen (-20 $^{\circ}$ C.) and freeze-thawed up to 5 times.

[0275] Once prepared, the desired synthetic duplexes are evaluated for their ability to modulate target expression. When cells reach 80% confluency, they are treated with synthetic duplexes comprising at least one oligomeric compound of the invention. For cells grown in 96-well plates, wells are washed once with 200 μ L OPTI-MEM-1 reduced-

serum medium (Gibco BRL) and then treated with 130 μ L of OPTI-MEM-1 containing 12 μ g/mL LIPOFECTIN (Gibco BRL) and the desired dsRNA compound at a final concentration of 200 nM. After 5 hours of treatment, the medium is replaced with fresh medium. Cells are harvested 16 hours after treatment, at which time RNA is isolated and target reduction measured by RT-PCR.

[0276] In another embodiment, the "suitable target segments" identified herein may be employed in a screen for additional oligomeric compounds that modulate the expression of a target. "Modulators" are those oligomeric compounds that decrease or increase the expression of a nucleic acid molecule encoding a target and which comprise at least an 8-nucleobase portion which is complementary to a suitable target segment. The screening method comprises the steps of contacting a suitable target segment of a nucleic acid molecule encoding a target with one or more candidate modulators, and selecting for one or more candidate modulators which decrease or increase the expression of a nucleic acid molecule encoding a target. Once it is shown that the candidate modulator or modulators are capable of modulating (e.g. either decreasing or increasing) the expression of a nucleic acid molecule encoding a target, the modulator may then be employed in further investigative studies of the function of a target, or for use as a research, diagnostic, or therapeutic agent in accordance with the present invention.

[0277] The suitable target segments of the present invention may also be combined with their respective complementary antisense oligomeric compounds of the present invention to form stabilized double-stranded (duplexed) oligonucleotides.

[0278] Screening and Target Validation

[0279] In a further embodiment, "suitable target segments" may be employed in a screen for additional oligomeric compounds that modulate the expression of a selected protein. "Modulators" are those oligomeric compounds that decrease or increase the expression of a nucleic acid molecule encoding a protein and which comprise at least an 8-nucleobase portion which is complementary to a suitable target segment. The screening method comprises the steps of contacting a suitable target segment of a nucleic acid molecule encoding a protein with one or more candidate modulators, and selecting for one or more candidate modulators which decrease or increase the expression of a nucleic acid molecule encoding a protein. Once it is shown that the candidate modulator or modulators are capable of modulating (e.g. either decreasing or increasing) the expression of a nucleic acid molecule encoding a peptide, the modulator may then be employed in further investigative studies of the function of the peptide, or for use as a research, diagnostic, or therapeutic agent in accordance with the present invention.

[0280] The oligomeric compounds of the present invention can also be applied in the areas of drug discovery and target validation. The present invention comprehends the use

of the oligomeric compounds and suitable targets identified herein in drug discovery efforts to elucidate relationships that exist between proteins and a disease state, phenotype, or condition. These methods include detecting or modulating a target peptide comprising contacting a sample, tissue, cell, or organism with the oligomeric compounds of the present invention, measuring the nucleic acid or protein level of the target and/or a related phenotypic or chemical endpoint at some time after treatment, and optionally comparing the measured value to a non-treated sample or sample treated with a further oligomeric compound of the invention. These methods can also be performed in parallel or in combination with other experiments to determine the function of unknown genes for the process of target validation or to determine the validity of a particular gene product as a target for treatment or prevention of a particular disease, condition, or phenotype.

[0281] Effect of nucleoside modifications on RNAi activity is evaluated according to existing literature (Elbashir et al., *Nature* (2001), 411, 494-498; Nishikura et al., *Cell* (2001), 107, 415-416; and Bass et al., *Cell* (2000), 101, 235-238.)

[0282] Kits, Research Reagents, Diagnostics, and Therapeutics

[0283] The oligomeric compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. Furthermore, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes or to distinguish between functions of various members of a biological pathway.

[0284] For use in kits and diagnostics, the oligomeric compounds of the present invention, either alone or in combination with other oligomeric compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

[0285] As one nonlimiting example, expression patterns within cells or tissues treated with one or more antisense oligomeric compounds are compared to control cells or tissues not treated with antisense oligomeric compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds and or oligomeric compounds which affect expression patterns.

[0286] Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, 2000, 480, 17-24; Celis, et al., *FEBS Lett.*, 2000, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., *Drug Discov. Today*, 2000, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, 1999, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2000, 97, 1976-81), protein arrays and proteomics (Celis, et

al., *FEBS Lett.*, 2000, 480, 2-16; Jungblut, et al., *Electrophoresis*, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, 2000, 480, 2-16; Larsson, et al., *J. Biotechnol.*, 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., *Anal. Biochem.*, 2000, 286, 91-98; Larson, et al., *Cytometry*, 2000, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, 2000, 3, 316-21), comparative genomic hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*, 1998, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, *Eur. J. Cancer*, 1999, 35, 1895-904) and mass spectrometry methods (To, *Comb. Chem. High Throughput Screen*, 2000, 3, 235-41).

[0287] The oligomeric compounds of the invention are useful for research and diagnostics, because these oligomeric compounds hybridize to nucleic acids encoding proteins. For example, oligonucleotides that are shown to hybridize with such efficiency and under such conditions as disclosed herein as to be effective protein inhibitors will also be effective primers or probes under conditions favoring gene amplification or detection, respectively. These primers and probes are useful in methods requiring the specific detection of nucleic acid molecules encoding proteins and in the amplification of the nucleic acid molecules for detection or for use in further studies. Hybridization of the antisense oligonucleotides, particularly the primers and probes, of the invention with a nucleic acid can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of selected proteins in a sample may also be prepared.

[0288] The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligomeric compounds have been employed as therapeutic moieties in the treatment of disease states in animals, including humans. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that antisense oligomeric compounds can be useful therapeutic modalities that can be configured to be useful in treatment regimes for the treatment of cells, tissues and animals, especially humans.

[0289] For therapeutics, an animal, such as a human, suspected of having a disease or disorder which can be treated by modulating the expression of a selected protein is treated by administering antisense oligomeric compounds in accordance with this invention. For example, in one non-limiting embodiment, the methods comprise the step of administering to the animal in need of treatment, a therapeutically effective amount of a protein inhibitor. The protein inhibitors of the present invention effectively inhibit the activity of the protein or inhibit the expression of the protein. In some embodiments, the activity or expression of a protein in an animal or in vitro is inhibited by at least 10%, by at least 20%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 80%, by at least 90%, by at least 95%, by at least 99%, or by 100%.

[0290] For example, the reduction of the expression of a protein may be measured in serum, adipose tissue, liver or

any other body fluid, tissue or organ of the animal. The cells contained within the fluids, tissues or organs being analyzed can contain a nucleic acid molecule encoding a protein and/or the protein itself.

[0291] The oligomeric compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an oligomeric compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the oligomeric compounds and methods of the invention may also be useful prophylactically.

[0292] Formulations

[0293] The oligomeric compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption-assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756.

[0294] The antisense oligomeric compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the oligomeric compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents. The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE ((S-acetyl-2-thioethyl) phosphate) derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published Dec. 9, 1993 or in WO 94/26764 and U.S. Pat. No. 5,770,713 to Imbach et al.

[0295] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the oligomeric compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto. For oligonucleotides, examples of pharmaceutically acceptable salts and their uses are further described in U.S. Pat. No. 6,287,860.

[0296] The present invention also includes pharmaceutical compositions and formulations which include the antisense oligomeric compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery),

pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

[0297] The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0298] The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0299] Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, foams and liposome-containing formulations. The pharmaceutical compositions and formulations of the present invention may comprise one or more penetration enhancers, carriers, excipients or other active or inactive ingredients.

[0300] Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter. Emulsions may contain additional components in addition to the dispersed phases, and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Microemulsions are included as an embodiment of the present invention. Emulsions and their uses are well known in the art and are further described in U.S. Pat. No. 6,287,860.

[0301] Formulations of the present invention include liposomal formulations. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers. Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior that contains the composition to be delivered. Cationic liposomes are positively charged liposomes which are believed to interact with negatively charged DNA molecules to form a stable complex. Liposomes that are pH-sensitive or negatively-charged are believed to entrap DNA rather

than complex with it. Both cationic and noncationic liposomes have been used to deliver DNA to cells.

[0302] Liposomes also include “sterically stabilized” liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome comprises one or more glycolipids or is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. Liposomes and their uses are further described in U.S. Pat. No. 6,287,860.

[0303] The pharmaceutical formulations and compositions of the present invention may also include surfactants. The use of surfactants in drug products, formulations and in emulsions is well known in the art. Surfactants and their uses are further described in U.S. Pat. No. 6,287,860.

[0304] In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants. Penetration enhancers and their uses are further described in U.S. Pat. No. 6,287,860.

[0305] One of skill in the art will recognize that formulations are routinely designed according to their intended use, i.e. route of administration.

[0306] Suitable formulations for topical administration include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Suitable lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA).

[0307] For topical or other administration, oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Suitable fatty acids and esters, pharmaceutically acceptable salts thereof, and their uses are further described in U.S. Pat. No. 6,287,860. Topical formulations are described in detail in U.S. patent application Ser. No. 09/315,298 filed on May 20, 1999.

[0308] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitabets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Suitable oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Suitable

surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Suitable bile acids/salts and fatty acids and their uses are further described in U.S. Pat. No. 6,287,860. Also suitable are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A suitable combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents and their uses are further described in U.S. Pat. No. 6,287,860. Oral formulations for oligonucleotides and their preparation are described in detail in U.S. application Ser. Nos. 09/108,673 (filed Jul. 1, 1998), 09/315,298 (filed May 20, 1999) and 10/071,822, filed Feb. 8, 2002.

[0309] Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[0310] Certain embodiments of the invention provide pharmaceutical compositions containing one or more oligomeric compounds and one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to cancer chemotherapeutic drugs such as daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). When used with the oligomeric compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to non-steroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. Combinations of antisense oligomeric compounds and other non-antisense drugs are also within the scope of this invention. Two or more combined compounds such as two oligomeric compounds or one oligomeric compound combined further compounds may be used together or sequentially.

[0311] In another related embodiment, compositions of the invention may contain one or more antisense oligomeric

compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense oligomeric compounds targeted to a second nucleic acid target. Alternatively, compositions of the invention may contain two or more antisense oligomeric compounds targeted to different regions of the same nucleic acid target.

[0312] Numerous examples of antisense oligomeric compounds are known in the art. Two or more combined compounds may be used together or sequentially

[0313] Dosing

[0314] The formulation of therapeutic compositions and their subsequent administration (dosing) is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC₅₀s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 μ g to 100 g per kg of body weight, from 0.1 μ g to 10 g per kg of body weight, from 1 μ g to 1 g per kg of body weight, from 10 μ g to 100 mg per kg of body weight, from 100 μ g to 10 mg per kg of body weight, or from 100 μ g to 1 mg per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 μ g to 100 g per kg of body weight, once or more daily, to once every 20 years.

[0315] While the present invention has been described with specificity in accordance with certain of its embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

EXAMPLES

Example 1

LNA Gappers Targeted to Mouse TRADD: In Vivo Study

[0316] Six-week old male Balb/c mice (Jackson Laboratory, Bar Harbor, Me.) were injected with compounds targeted to TRADD. Each treatment group was comprised of four animals. Animals were dosed via intraperitoneal injection twice per week for three weeks. The LNA gappers ISIS 335385, ISIS 353878, and ISIS 353879 were evaluated at doses of 40, 12, or 4 mg/kg (4.5, 1.5, or 0.5 μ mol/kg, respectively). The 2'-MOE gappers ISIS 325585, ISIS325587, and ISIS 325588 were similarly administered. Saline-injected animals and animals injected with ISIS 141923, a control 2'-MOE gapper, served as control groups.

Animals were sacrificed after the last dose of oligonucleotide was administered, and liver, kidney, and fat tissues were harvested.

[0317] Target reduction in liver was measured at the conclusion of the study using real-time PCR. Results are presented in Table 2 for animals treated with 40 mg/kg (4.5 μ mol/kg) of the compounds as described. The data were normalized to saline-treated controls and are expressed as percent of control, wherein a percentage less than 100 is indicative of target mRNA reduction.

TABLE 2

Target reduction by LNA compounds targeted to TRADD in mouse liver				
ISIS #	SEQUENCE	CHEMISTRY	SEQ ID NO	% Control
141923	CCTTCCCTGAAGGTTCTCTCC	5-10-5 MOE	3	94
335385	CTCATACTCGTAGGCC	3-10-3 LNA	4	20
353878	GCTCATACTCGTAGGCCA	3-12-3 LNA	5	17
353879	GCTCATACTCGTAGGCCA	2-14-2 LNA	6	30
325585	CTCATACTCGTAGGCC	3-10-3 MOE	7	65
325588	GCTCATACTCGTAGGCCA	3-12-3 MOE	8	19
325587	GCTCATACTCGTAGGCCA	2-14-2 MOE	9	17

[0318] As shown in Table 2, all of the compounds targeted to TRADD were effective at reducing mRNA levels in mouse liver, whereas ISIS 141923, a control oligonucleotide, did not cause substantial target reduction. ISIS 335385, an oligomeric compound comprised of a ten 2'-deoxynucleotide gap flanked by three LNAs on both the 3' and 5' ends, substantially reduced TRADD mRNA levels as compared to ISIS 325585, an oligomeric compound comprised of a ten 2'-deoxynucleotide gap flanked by three 2'-MOE nucleotides on both the 3' and 5' ends. ISIS 353878, an oligomeric compound comprised of a twelve 2'-deoxynucleotide gap flanked by three LNAs on both the 3' and 5' ends, caused reduction in TRADD mRNA levels comparable to that produced by ISIS 325588, an oligomeric compound comprised of a twelve 2'-deoxynucleotide gap flanked by three 2'-MOE nucleotides on both the 3' and 5' ends.

Example 2

Synthesis of Nucleoside Phosphoramidites

[0319] The following compounds, including amidites and their intermediates were prepared as described in U.S. Pat. No. 6,426,220 and published PCT WO 02/36743; 5'-O-Dimethoxytrityl-thymidine intermediate for 5-methyl dC amidite, 5'-O-Dimethoxytrityl-2'-deoxy-5-methylcytidine intermediate for 5-methyl-dC amidite, 5'-O-Dimethoxytrityl-2'-deoxy-N⁴-benzoyl-5-methylcytidine penultimate intermediate for 5-methyl dC amidite, (5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-deoxy-N-4-benzoyl-5-methylcytidin-3'-O-yl)-2'-cyanoethyl-N,N-diisopropylphosphoramidite (5-methyl dC amidite), 2'-Fluorodeoxyadenosine, 2'-Fluorodeoxyguanosine, 2'-Fluorouridine,

2'-Fluorodeoxycytidine, 2'-O-(2-Methoxyethyl) modified amidites, 2'-O-(2-methoxyethyl)-5-methyluridine intermediate, 5'-O-DMT-2'-O-(2-methoxyethyl)-5-methyluridine penultimate intermediate, (5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-5-methyluridin-3'-O-yl)-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE T amidite), 5'-O-Dimethoxytrityl-2'-O-(2-methoxyethyl)-5-methylcytidine intermediate, 5'-O-dimethoxytrityl-2'-O-(2-methoxyethyl)-N⁴-benzoyl-5-methyl-cytidine penultimate intermediate, (5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N⁴-benzoyl-5-methylcytidin-3'-O-yl)-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE 5-Me-C amidite), (5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N⁶-benzoyladenoin-3'-O-yl)-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE A amidite), (5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N₄-isobutrylguanosin-3'-O-yl)-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE G amidite), 2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminoxyethyl) nucleoside amidites, 2'-(Dimethylaminoxyethoxy) nucleoside amidites, 5'-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine, 5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine, 2'-O-((2-phthalimidoxy)ethyl)-5'-t-butylidiphenylsilyl-5-methyluridine, 5'-O-tert-butylidiphenylsilyl-2'-O-((2-formadoximinoxy)ethyl)-5-methyluridine, 5'-O-tert-Butyldiphenylsilyl-2'-O-(N,N dimethylaminoxyethyl)-5-methyluridine, 2'-O-(dimethyl-aminoxyethyl)-5-methyluridine, 5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine, 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-((2-cyanoethyl)-N,N-diisopropylphosphoramidite), 2'-(Aminooxyethoxy) nucleoside amidites, N₂-isobutryl-6-O-diphenylcarbonyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-((2-cyanoethyl)-N,N-diisopropylphosphoramidite), 2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites, 2'-O-(2(2-N,N-dimethylaminoethoxy)ethyl)-5-methyl uridine, 5'-O-dimethoxytrityl-2'-O-(2(2-N,N-dimethylaminoethoxy)-ethyl))-5-methyl uridine and 5'-O-Dimethoxytrityl-2'-O-(2(2-N,N-dimethylaminoethoxy)-ethyl))-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite.

Example 3

Oligonucleotide and Oligonucleoside Synthesis

[0320] The antisense oligomeric compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

[0321] Oligonucleotides: Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 394) using standard phosphoramidite chemistry with oxidation by iodine.

[0322] Phosphorothioates (P=S) are synthesized similar to phosphodiester oligonucleotides with the following

exceptions: thiation was effected by utilizing a 10% w/v solution of 3,4,5-trimethyl-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the oxidation of the phosphite linkages. The thiation reaction step time was increased to 180 sec and preceded by the normal capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55° C. (12-16 hr), the oligonucleotides were recovered by precipitating with >3 volumes of ethanol from a 1 M NH₄OAc solution. Phosphinate oligonucleotides are prepared as described in U.S. Pat. No. 5,508,270.

[0323] Alkyl phosphonate oligonucleotides are prepared as described in U.S. Pat. No. 4,469,863.

[0324] 3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Pat. No. 5,610,289 or 5,625,050.

[0325] Phosphoramidite oligonucleotides are prepared as described in U.S. Pat. No. 5,256,775 or U.S. Pat. No. 5,366,878.

[0326] Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively).

[0327] 3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Pat. No. 5,476,925.

[0328] Phosphotriester oligonucleotides are prepared as described in U.S. Pat. No. 5,023,243.

[0329] Borano phosphate oligonucleotides are prepared as described in U.S. Pat. Nos. 5,130,302 and 5,177,198.

[0330] Oligonucleosides: Methyleneethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone oligomeric compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Pat. Nos. 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289.

[0331] Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Pat. Nos. 5,264,562 and 5,264,564.

[0332] Ethylene oxide linked oligonucleosides are prepared as described in U.S. Pat. No. 5,223,618.

Example 4

RNA Synthesis

[0333] In general, RNA synthesis chemistry is based on the selective incorporation of various protecting groups at strategic intermediary reactions. Although one of ordinary skill in the art will understand the use of protecting groups in organic synthesis, a useful class of protecting groups includes silyl ethers. In particular bulky silyl ethers are used to protect the 5'-hydroxyl in combination with an acid-labile orthoester protecting group on the 2'-hydroxyl. This set of protecting groups is then used with standard solid-phase synthesis technology. It is important to lastly remove the

acid labile orthoester protecting group after all other synthetic steps. Moreover, the early use of the silyl protecting groups during synthesis ensures facile removal when desired, without undesired deprotection of 2' hydroxyl.

[0334] Following this procedure for the sequential protection of the 5'-hydroxyl in combination with protection of the 2'-hydroxyl by protecting groups that are differentially removed and are differentially chemically labile, RNA oligonucleotides were synthesized.

[0335] RNA oligonucleotides are synthesized in a step-wise fashion. Each nucleotide is added sequentially (3' to 5'-direction) to a solid support-bound oligonucleotide. The first nucleoside at the 3'-end of the chain is covalently attached to a solid support. The nucleotide precursor, a ribonucleoside phosphoramidite, and activator are added, coupling the second base onto the 5'-end of the first nucleoside. The support is washed and any unreacted 5'-hydroxyl groups are capped with acetic anhydride to yield 5'-acetyl moieties. The linkage is then oxidized to the more stable and ultimately desired P(V) linkage. At the end of the nucleotide addition cycle, the 5'-silyl group is cleaved with fluoride. The cycle is repeated for each subsequent nucleotide.

[0336] Following synthesis, the methyl protecting groups on the phosphates are cleaved in 30 minutes utilizing 1 M disodium-2-carbamoyl-2-cyanoethylene-1,1-dithiolate trihydrate (S₂Na₂) in DMF. The deprotection solution is washed from the solid support-bound oligonucleotide using water. The support is then treated with 40% methylamine in water for 10 minutes at 55° C. This releases the RNA oligonucleotides into solution, deprotects the exocyclic amines, and modifies the 2'-groups. The oligonucleotides can be analyzed by anion exchange HPLC at this stage.

[0337] The 2'-orthoester groups are the last protecting groups to be removed. The ethylene glycol monoacetate orthoester protecting group developed by Dharmacon Research, Inc. (Lafayette, Colo.), is one example of a useful orthoester protecting group which, has the following important properties. It is stable to the conditions of nucleoside phosphoramidite synthesis and oligonucleotide synthesis. However, after oligonucleotide synthesis the oligonucleotide is treated with methylamine which not only cleaves the oligonucleotide from the solid support but also removes the acetyl groups from the orthoesters. The resulting 2-ethyl-hydroxyl substituents on the orthoester are less electron withdrawing than the acetylated precursor. As a result, the modified orthoester becomes more labile to acid-catalyzed hydrolysis. Specifically, the rate of cleavage is approximately 10 times faster after the acetyl groups are removed. Therefore, this orthoester possesses sufficient stability in order to be compatible with oligonucleotide synthesis and yet, when subsequently modified, permits deprotection to be carried out under relatively mild aqueous conditions compatible with the final RNA oligonucleotide product.

[0338] Additionally, methods of RNA synthesis are well known in the art (Scaringe, S. A. Ph.D. Thesis, University of Colorado, 1996; Scaringe, S. A., et al., J. Am. Chem. Soc., 1998, 120, 11820-11821; Matteucci, M. D. and Caruthers, M. H. J. Am. Chem. Soc., 1981, 103, 3185-3191; Beaucage, S. L. and Caruthers, M. H. Tetrahedron Lett., 1981, 22, 1859-1862; Dahl, B. J., et al., Acta Chem. Scand, 1990, 44, 639-641; Reddy, M. P., et al., Tetrahedron Lett., 1994, 25, 4311-4314; Wincott, F. et al., Nucleic Acids Res., 1995, 23,

2677-2684; Griffin, B. E., et al., Tetrahedron, 1967, 23, 2301-2313; Griffin, B. E., et al., Tetrahedron, 1967, 23, 2315-2331).

[0339] RNA antisense oligomeric compounds (RNA oligonucleotides) of the present invention can be synthesized by the methods herein or purchased from Dharmacon Research, Inc (Lafayette, Colo.). Once synthesized, complementary RNA antisense oligomeric compounds can then be annealed by methods known in the art to form double stranded (duplexed) antisense oligomeric compounds. For example, duplexes can be formed by combining 30 μ l of each of the complementary strands of RNA oligonucleotides (50 μ M RNA oligonucleotide solution) and 15 μ l of 5 \times annealing buffer (100 mM potassium acetate, 30 mM HEPES-KOH pH 7.4, 2 mM magnesium acetate) followed by heating for 1 minute at 90° C., then 1 hour at 37° C. The resulting duplexed antisense oligomeric compounds can be used in kits, assays, screens, or other methods to investigate the role of a target nucleic acid.

Example 5

Synthesis of Chimeric Oligonucleotides

[0340] Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3'"wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers."

[0341] (2'-O-Me)—(2'-deoxy)—(2'-O-Me) Chimeric Phosphorothioate Oligonucleotides

[0342] Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 394, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by incorporating coupling steps with increased reaction times for the 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite. The fully protected oligonucleotide is cleaved from the support and deprotected in concentrated ammonia (NH₄OH) for 12-16 hr at 55° C. The deprotected oligo is then recovered by an appropriate method (precipitation, column chromatography, volume reduced in vacuo and analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[0343] (2'-O-(2-Methoxyethyl))—(2'-deoxy)—(2'-O-(Methoxyethyl)) Chimeric Phosphorothioate Oligonucleotides

[0344] (2'-O-(2-methoxyethyl))—(2'-deoxy)—(2'-O-(methoxyethyl)) chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

[0345] (2'-O-(2-Methoxyethyl)Phosphodiester)—(2'-deoxy Phosphorothioate)—(2'-O-(2-Methoxyethyl) Phosphodiester) Chimeric Oligonucleotides

[0346] (2'-O-(2-methoxyethyl phosphodiester)—(2'-deoxy phosphorothioate)—(2'-O-(methoxyethyl) phosphodiester) chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidation with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfuration utilizing 3,4-dithiothiopyran-2-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

[0347] Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to U.S. Pat. No. 5,623,065.

Example 6

Oligonucleotide Isolation

[0348] After cleavage from the controlled pore glass solid support and deblocking in concentrated ammonium hydroxide at 55° C. for 12-16 hours, the oligonucleotides or oligonucleosides are recovered by precipitation out of 1 M NH₄OAc with >3 volumes of ethanol. Synthesized oligonucleotides were analyzed by electrospray mass spectroscopy (molecular weight determination) and by capillary gel electrophoresis and judged to be at least 70% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in the synthesis was determined by the ratio of correct molecular weight relative to the -16 amu product (+/-32+/-48). For some studies oligonucleotides were purified by HPLC, as described by Chiang et al., J. Biol. Chem. 1991, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

Example 7

Oligonucleotide Synthesis —96 Well Plate Format

[0349] Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a 96-well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfuration utilizing 3,4-dithiothiopyran-2-one 1,1 dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, Calif., or Pharmacia, Piscataway, N.J.). Non-standard nucleosides are synthesized as per standard or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

[0350] Oligonucleotides were cleaved from support and deprotected with concentrated NH₄OH at elevated temperature (55-60° C.) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

Example 8

Oligonucleotide Analysis 96 Well Plate Format

[0351] The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96-well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the oligomeric compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the oligomeric compounds on the plate were at least 85% full length.

Example 9

Cell Culture and Oligonucleotide Treatment

[0352] The effect of oligomeric compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, ribonuclease protection assays, or RT-PCR.

[0353] T-24 Cells:

[0354] The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, Va.). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Invitrogen Corporation, Carlsbad, Calif.) supplemented with 10% fetal calf serum (Invitrogen Corporation, Carlsbad, Calif.), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Invitrogen Corporation, Carlsbad, Calif.). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #353872) at a density of 7000 cells/well for use in RT-PCR analysis.

[0355] For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

[0356] A549 Cells:

[0357] The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, Va.). A549 cells were routinely cultured in DMEM basal media (Invitrogen Corporation, Carlsbad, Calif.) supplemented with 10% fetal calf serum (Invitrogen Corporation, Carlsbad, Calif.), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Invitrogen Corporation, Carlsbad, Calif.). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

[0358] NHDF Cells:

[0359] Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville, Md.). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville, Md.) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

[0360] HEK Cells:

[0361] Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville, Md.). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville, Md.) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

[0362] MCF-7 Cells

[0363] The human breast carcinoma cell line MCF-7 is obtained from the American Type Culture Collection (Manassas, Va.). These cells contain a wild-type p53 gene. MCF-7 cells are routinely cultured in DMEM low glucose (Gibco/Life Technologies, Gaithersburg, Md.) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, Md.). Cells are routinely passaged by trypsinization and dilution when they reach 90% confluence. Cells are seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for treatment with the oligomeric compounds of the invention.

[0364] HepB3 Cells

[0365] The human hepatoma cell line HepB3 (Hep3B2.1-7) is obtained from the American Type Culture Collection (ATCC-ATCC Catalog # HB-8064) (Manassas, Va.). This cell line was initially derived from a hepatocellular carcinoma of an 8-yr-old black male. The cells are epithelial in morphology and are tumorigenic in nude mice. HepB3 cells are routinely cultured in Minimum Essential Medium (MEM) with Earle's Balanced Salt Solution, 2 mM L-glutamine, 1.5 g/L sodium bicarbonate, 0.1 mM nonessential amino acids, 1.0 mM sodium pyruvate (ATCC #20-2003, Manassas, Va.) and with 10% heat-inactivated fetal bovine serum (Gibco/Life Technologies, Gaithersburg, Md.). Cells are routinely passaged by trypsinization and dilution when they reach 90% confluence.

[0366] Primary Mouse Hepatocytes

[0367] Primary mouse hepatocytes are prepared from CD-1 mice purchased from Charles River Labs. Primary mouse hepatocytes are routinely cultured in Hepatocyte Attachment Media (Invitrogen Life Technologies, Carlsbad, Calif.) supplemented with 10% Fetal Bovine Serum (Invitrogen Life Technologies, Carlsbad, Calif.), 250 nM dexamethasone (Sigma-Aldrich Corporation, St. Louis, Mo.), 10 nM bovine insulin (Sigma-Aldrich Corporation, St. Louis, Mo.). Cells are seeded into 96-well plates (Falcon-Primaria #353872, BD Biosciences, Bedford, Mass.) at a density of 4000-6000 cells/well for treatment with the oligomeric compounds of the invention.

[0368] Treatment with Oligomeric Compounds:

[0369] When cells reached 65-75% confluency, they were treated with oligonucleotide. For cells grown in 96-well

plates, wells were washed once with 100 μ L OPTI-MEM™-1 reduced-serum medium (Invitrogen Corporation, Carlsbad, Calif.) and then treated with 130 μ L of OPTI-MEM™-1 containing 3.75 μ g/mL LIPOFECTIN™ (Invitrogen Corporation, Carlsbad, Calif.) and the desired concentration of oligonucleotide. Cells are treated and data are obtained in triplicate. After 4-7 hours of treatment at 37° C., the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

[0370] The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is selected from either ISIS 13920 (TCCGT-CATCGCTCCTCAGGG, SEQ ID NO:10) which is targeted to human H-ras, or ISIS 18078, (GTGCGCGGAGC-CCGAAATC, SEQ ID NO:11) which is targeted to human Jun-N-terminal kinase-2 (JNK2). Both controls are 2'-O-methoxyethyl gapmers (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, ATGCAT-TCTGCCCCCAAGGA, SEQ ID NO:12, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-H-ras (for ISIS 13920), JNK2 (for ISIS 18078) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of c-H-ras, JNK2 or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments. The concentrations of antisense oligonucleotides used herein are from 50 nM to 300 mM.

Example 10

Analysis of Oligonucleotide Inhibition of a Target Expression

[0371] Antisense modulation of a target expression can be assayed in a variety of ways known in the art. For example, a target mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently suitable. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. One method of RNA analysis of the present invention is the use of total cellular RNA as described in other examples herein. Methods of RNA isolation are well known in the art. Northern blot analysis is also routine in the art. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7600, 7700, or 7900 Sequence Detection System, available from PE-Applied Biosystems, Foster City, Calif. and used according to manufacturer's instructions.

[0372] Protein levels of a target can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), enzyme-

linked immunosorbent assay (ELISA) or fluorescence-activated cell sorting (FACS). Antibodies directed to a target can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, Mich.), or can be prepared via conventional monoclonal or polyclonal antibody generation methods well known in the art.

Example 11

Design of Phenotypic Assays and In Vivo Studies for the Use of a Target Inhibitors

[0373] Phenotypic Assays

[0374] Once a target inhibitors have been identified by the methods disclosed herein, the oligomeric compounds are further investigated in one or more phenotypic assays, each having measurable endpoints predictive of efficacy in the treatment of a particular disease state or condition.

[0375] Phenotypic assays, kits and reagents for their use are well known to those skilled in the art and are herein used to investigate the role and/or association of a target in health and disease. Representative phenotypic assays, which can be purchased from any one of several commercial vendors, include those for determining cell viability, cytotoxicity, proliferation or cell survival (Molecular Probes, Eugene, Oreg.; PerkinElmer, Boston, Mass.), protein-based assays including enzymatic assays (Panvera, LLC, Madison, Wis.; BD Biosciences, Franklin Lakes, N.J.; Oncogene Research Products, San Diego, Calif.), cell regulation, signal transduction, inflammation, oxidative processes and apoptosis (Assay Designs Inc., Ann Arbor, Mich.), triglyceride accumulation (Sigma-Aldrich, St. Louis, Mo.), angiogenesis assays, tube formation assays, cytokine and hormone assays and metabolic assays (Chemicon International Inc., Temecula, Calif.; Amersham Biosciences, Piscataway, N.J.).

[0376] In one non-limiting example, cells determined to be appropriate for a particular phenotypic assay (i.e., MCF-7 cells selected for breast cancer studies; adipocytes for obesity studies) are treated with a target inhibitors identified from the in vitro studies as well as control compounds at optimal concentrations which are determined by the methods described above. At the end of the treatment period, treated and untreated cells are analyzed by one or more methods specific for the assay to determine phenotypic outcomes and endpoints. Phenotypic endpoints include changes in cell morphology over time or treatment dose as well as changes in levels of cellular components such as proteins, lipids, nucleic acids, hormones, saccharides or metals. Measurements of cellular status which include pH, stage of the cell cycle, intake or excretion of biological indicators by the cell, are also endpoints of interest.

[0377] Analysis of the genotype of the cell (measurement of the expression of one or more of the genes of the cell) after treatment is also used as an indicator of the efficacy or potency of the a target inhibitors. Hallmark genes, or those genes suspected to be associated with a specific disease state, condition, or phenotype, are measured in both treated and untreated cells.

[0378] In Vivo Studies

[0379] The individual subjects of the in vivo studies described herein are warm-blooded vertebrate animals, which includes humans.

[0380] The clinical trial is subjected to rigorous controls to ensure that individuals are not unnecessarily put at risk and that they are fully informed about their role in the study.

[0381] To account for the psychological effects of receiving treatments, volunteers are randomly given placebo or a target inhibitor. Furthermore, to prevent the doctors from being biased in treatments, they are not informed as to whether the medication they are administering is a target inhibitor or a placebo. Using this randomization approach, each volunteer has the same chance of being given either the new treatment or the placebo.

[0382] Volunteers receive either the a target inhibitor or placebo for eight week period with biological parameters associated with the indicated disease state or condition being measured at the beginning (baseline measurements before any treatment), end (after the final treatment), and at regular intervals during the study period. Such measurements include the levels of nucleic acid molecules encoding a target or a target protein levels in body fluids, tissues or organs compared to pre-treatment levels. Other measurements include, but are not limited to, indices of the disease state or condition being treated, body weight, blood pressure, serum titers of pharmacologic indicators of disease or toxicity as well as ADME (absorption, distribution, metabolism and excretion) measurements. Information recorded for each patient includes age (years), gender, height (cm), family history of disease state or condition (yes/no), motivation rating (some/moderate/great) and number and type of previous treatment regimens for the indicated disease or condition.

[0383] Volunteers taking part in this study are healthy adults (age 18 to 65 years) and roughly an equal number of males and females participate in the study. Volunteers with certain characteristics are equally distributed for placebo and a target inhibitor treatment. In general, the volunteers treated with placebo have little or no response to treatment, whereas the volunteers treated with the a target inhibitor show positive trends in their disease state or condition index at the conclusion of the study.

Example 12

RNA Isolation

[0384] Poly(A)+ mRNA Isolation

[0385] Poly(A)+ mRNA was isolated according to Miura et al., (Clin. Chem., 1996, 42, 1758-1764). Other methods for poly(A)+ mRNA isolation are routine in the art. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 60 μ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55 μ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine Calif.). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200 μ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60 μ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70° C., was added to

each well, the plate was incubated on a 90° C. hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

[0386] Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

[0387] Total RNA Isolation

[0388] Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia, Calif.) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 150 μ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 150 μ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 1 minute. 500 μ L of Buffer RW1 was added to each well of the RNEASY 96™ plate and incubated for 15 minutes and the vacuum was again applied for 1 minute. An additional 500 μ L of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum was applied for 2 minutes. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 90 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 3 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 140 μ L of RNase free water into each well, incubating 1 minute, and then applying the vacuum for 3 minutes.

[0389] The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia Calif.). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

Example 13

Real-Time Quantitative PCR Analysis of a Target mRNA Levels

[0390] Quantitation of a target mRNA levels was accomplished by real-time quantitative PCR using the ABI PRISM™ 7600, 7700, or 7900 Sequence Detection System (PE-Applied Biosystems, Foster City, Calif.) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., FAM or JOE, obtained from either PE-Applied Biosystems, Foster City, Calif., Operon Technologies Inc.,

Alameda, Calif. or Integrated DNA Technologies Inc., Coralville, Iowa) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either PE-Applied Biosystems, Foster City, Calif., Operon Technologies Inc., Alameda, Calif. or Integrated DNA Technologies Inc., Coralville, Iowa) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

[0391] Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

[0392] PCR reagents were obtained from Invitrogen Corporation, (Carlsbad, Calif.). RT-PCR reactions were carried out by adding 20 μ L PCR cocktail (2.5 \times PCR buffer minus $MgCl_2$, 6.6 mM $MgCl_2$, 375 μ M each of dATP, dCTP, dCTP and dGTP, 375 nM each of forward primer and reverse primer, 125 nM of probe, 4 Units RNase inhibitor, 1.25 Units PLATINUM® Taq, 5 Units MULV reverse transcriptase, and 2.5 \times ROX dye) to 96-well plates containing 30 μ L total RNA solution (20-200 ng). The RT reaction was carried out by incubation for 30 minutes at 48° C. Following a 10 minute incubation at 95° C. to activate the PLATINUM® Taq, 40 cycles of a two-step PCR protocol were carried out: 95° C. for 15 seconds (denaturation) followed by 60° C. for 1.5 minutes (annealing/extension).

[0393] Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreen™ (Molecular Probes, Inc. Eugene, Oreg.). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen™ RNA quantification reagent (Molecular

Probes, Inc. Eugene, Oreg.). Methods of RNA quantification by RiboGreen™ are taught in Jones, L. J., et al, (Analytical Biochemistry, 1998, 265, 368-374).

[0394] In this assay, 170 μ L of RiboGreen™ working reagent (RiboGreen™ reagent diluted 1:350 in 10 mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 30 μ L purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 485 nm and emission at 530 nm.

[0395] Probes and are designed to hybridize to a human a target sequence, using published sequence information.

Example 14

Northern Blot Analysis of a Target mRNA Levels

[0396] Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST “B” Inc., Friendswood, Tex.). Total RNA was prepared following manufacturer’s recommended protocols. Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, Ohio). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, N.J.) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST “B” Inc., Friendswood, Tex.). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, Calif.) and then probed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, Calif.) using manufacturer’s recommendations for stringent conditions.

[0397] To detect human a target, a human a target specific primer probe set is prepared by PCR To normalize for variations in loading and transfer efficiency membranes are stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, Calif.).

[0398] Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, Calif.). Data was normalized to GAPDH levels in untreated controls.

Example 15

Antisense Inhibition of Human a Target Expression by Oligonucleotides

[0399] In accordance with the present invention, a series of oligomeric compounds are designed to target different

regions of the human target RNA. The oligomeric compounds are analyzed for their effect on human target mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from three experiments. The target regions to which these suitable sequences are complementary are herein referred to as “suitable target segments” and are therefore suitable for targeting by oligomeric compounds of the present invention. The sequences represent the reverse complement of the suitable antisense oligomeric compounds.

[0400] As these “suitable target segments” have been found by experimentation to be open to, and accessible for, hybridization with the antisense oligomeric compounds of the present invention, one of skill in the art will recognize or be able to ascertain, using no more than routine experimentation, further embodiments of the invention that encompass other oligomeric compounds that specifically hybridize to these suitable target segments and consequently inhibit the expression of a target.

[0401] According to the present invention, antisense oligomeric compounds include antisense oligomeric compounds, antisense oligonucleotides, ribozymes, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other short oligomeric compounds which hybridize to at least a portion of the target nucleic acid.

Example 16

Western Blot Analysis of a Target Protein Levels

[0402] Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 μ L/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to a target is used, with a radiolabeled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale Calif.).

[0403] Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. Each reference (including, but not limited to, journal articles, U.S. and non-U.S. patents, patent application publications, international patent application publications, gene bank accession numbers, and the like) cited in the present application is incorporated herein by reference in its entirety.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 12

<210> SEQ ID NO 1

<211> LENGTH: 19

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 1

cgagaggcgg acgggaccg 19

<210> SEQ ID NO 2
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 2

cgggtcccgtc cgctctcgt t 21

<210> SEQ ID NO 3
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 3

ccttcctga aggttcctcc 20

<210> SEQ ID NO 4
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 4

ctcactactcg taggcc 16

<210> SEQ ID NO 5
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 5

gctcactac gtaggcca 18

<210> SEQ ID NO 6
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 6

gctcactac gtaggcca 18

<210> SEQ ID NO 7
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

-continued

```

<400> SEQUENCE: 7
ctcactactcg taggcc                               16

<210> SEQ ID NO 8
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 8
gctcactactc gtaggcca                             18

<210> SEQ ID NO 9
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 9
gctcactactc gtaggcca                             18

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 10
tccgctcatcg ctctcaggg                           20

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 11
gtgcgcgcgga gccccaagtc                           20


<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Oligomer

<400> SEQUENCE: 12
atgcattctg cccccaagga                             20

```

What is claimed is:

1. An oligomeric compound comprising the structure:



wherein:

each Nu_1 and Nu_3 is, independently, a high affinity modified nucleoside, wherein at least one of Nu_1 and Nu_3 is a bicyclic sugar modified nucleoside comprising a 4'-CH₂-O-2' bridge or a 4'-(CH₂)₂-O-2' bridge;

each Nu_2 is a 2'-deoxy nucleoside;

each L_1 , L_2 and L_3 is, independently, an internucleoside linking group;

each T_1 and T_2 is, independently, H, a hydroxy protecting group, an optionally linked conjugate group, or a covalent attachment to a solid support medium;

$n1$ is from 1 to about 6;

n2 is from 11 to about 18; and

n3 is from 2 to about 6.

2. The oligomeric compound of claim 1 wherein each of the high affinity modified nucleosides is, independently, a bicyclic sugar modified nucleoside, a 2'-O-(CH₂)₂-O-CH₃ modified nucleoside, a 2'-F modified nucleoside, or a 2'-O-CH₂-C(=O)-NR₁R₂ modified nucleoside, where each R₁ and R₂ is, independently, H, a nitrogen protecting group, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, wherein the substitution is OR₃, SR₃, NH₃⁺, NR₃R₄, guanidino or acyl, wherein the acyl is acid amide or an ester, or R₁ and R₂, together, are a nitrogen protecting group, or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O.

3. The oligomeric compound of claim 2 wherein each of the bicyclic sugar modified nucleosides independently has a 4'-CH₂-O-2' bridge or a 4'-(CH₂)₂-O-2' bridge.

4. The oligomeric compound of claim 2 wherein each R₁ and R₂ each is, independently, H, a nitrogen protecting group, or C₁-C₁₀ alkyl.

5. The oligomeric compound of claim 1 wherein T₁ is H or a hydroxyl protecting group.

6. The oligomeric compound of claim 1 wherein T₂ is H or a hydroxyl protecting group.

7. The oligomeric compound of claim 1 wherein each of the hydroxyl protecting groups is, independently, 4,4'-dimethoxytrityl, monomethoxytrityl, 9-phenylxanthene-9-yl, 9-(p-methoxyphenyl)xanthene-9-yl, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl, 2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,p-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triphenylsilyl, benzoylformate, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, pivaloyl, benzoyl, p-phenylbenzoyl, mesyl, tosyl, 4,4',4''-tris-(benzyloxy)trityl, 4,4',4''-tris-(4,5-dichlorophthalimido)trityl, 4,4',4''-tris(levulinyloxy)trityl, 3-(imidazolymethyl)-4,4'-dimethoxytrityl, 4-decyloxytrityl, 4-hexadecyloxytrityl, 9-(4-octadecyloxyphenyl)xanthene-9-yl, 1,1-bis-(4-methoxyphenyl)-1'-pyrenyl methyl, p-phenylazophenylloxycarbonyl, 9-fluorenylmethoxycarbonyl, 2,4-dinitrophenylethoxycarbonyl, 4-(methylthiomethoxy)butyl, 2-(methylthiomethoxymethyl)-benzoyl, 2-(isopropylthiomethoxymethyl)benzoyl, 2-(2,4-dinitrobenzenesulphenylloxymethyl)benzoyl, or levulinyl group.

8. The oligomeric compound of claim 1 wherein one of T₁ and T₂ is a covalent attachment to a support medium.

9. The oligomeric compound of claim 8 wherein the support medium is a controlled pore glass, oxalyl-controlled pore glass, silica-containing particles, polymers of polystyrene, copolymers of polystyrene, copolymers of styrene and divinylbenzene, copolymers of dimethylacrylamide and N,N'-bisacryloyl ethylenediamine, soluble support medium, or PEPS.

10. The oligomeric compound of claim 1 wherein each L₁, L₂ and L₃ is, independently, phosphodiester, phosphorothioate, chiral phosphorothioate, phosphorodithioate, phosphotriester, aminoalkylphosphotriester, methyl phosphonate, alkyl phosphonate, 5'-alkylene phosphonate, chiral phosphonate, phosphinate, phosphoramidate, 3'-amino phosphoramidate, aminoalkylphosphoramidate, thionophosphoramidate,

thionoalkylphosphonate, thionoalkylphosphotriester, selenophosphate, or boranophosphate.

11. The oligomeric compound of claim 10 wherein each L₁, L₂ and L₃ is, independently, a phosphodiester or a phosphorothioate internucleoside linking group.

12. The oligomeric compound of claim 10 wherein each L₁, L₂ and L₃ is a phosphodiester internucleoside linking group.

13. The oligomeric compound of claim 1 wherein each L₁, L₂ and L₃ is, independently, siloxane, sulfide, sulfoxide, sulfone, formacetyl, thioformacetyl, methylene formacetyl, thioformacetyl, sulfamate, methyleneimino, methylenehydrazino, sulfonate, sulfonamide, or amide.

14. The oligomeric compound of claim 13 wherein each of the internucleoside linking groups is, independently, —CH₂—NH—O—CH₂—, —CH₂—N(CH₃)—O—CH₂— or —CH₂—O—N(CH₃)—CH₂—, —CH₂—N(CH₃)—N(CH₃)—CH₂—, or —O—N(CH₃)—CH₂—CH₂—.

15. The oligomeric compound of claim 1 wherein each nucleoside comprises a heterocyclic base moiety that is, independently, adenine, guanine, thymine, cytosine, uracil, 5-methylcytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, 5-halocytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-substituted adenines and guanines, 5-substituted uracils and cytosines, 7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, or 3-deazaadenine.

16. The oligomeric compound of claim 1 wherein n1 is from 1 to about 5.

17. The oligomeric compound of claim 1 wherein n1 is from 1 to about 3.

18. The oligomeric compound of claim 1 wherein n1 is from 2 to about 3.

19. The oligomeric compound of claim 1 wherein n3 is from 2 to about 5.

20. The oligomeric compound of claim 1 wherein n3 is from 2 to about 3.

21. The oligomeric compound of claim 1 wherein n2 is from 12 to about 18.

22. The oligomeric compound of claim 1 wherein n2 is from 12 to about 16.

23. The oligomeric compound of claim 1 wherein n2 is from 14 to about 16.

24. The oligomeric compound of claim 1 wherein the total of n1, n2 and n3 is from 14 to about 30.

25. The oligomeric compound of claim 1 wherein the total of n1, n2 and n3 is from 14 to 24.

26. The oligomeric compound of claim 1 wherein the total of n1, n2 and n3 is from 14 to 21.

27. The oligomeric compound of claim 1 wherein the total of n1, n2 and n3 is from 16 to 21.

28. The oligomeric compound of claim 1 wherein n1 is from 1 to about 3, n2 is 12 or 13 and n3 is 2 or 3.

29. The oligomeric compound of claim 1 wherein n1 is 3, n2 is 12 and n3 is 3.

30. The oligomeric compound of claim 1 wherein n1 is from 1 to about 3, n2 is 14 or 15 and n3 is 2 or 3.

31. The oligomeric compound of claim 1 wherein n1 is 2, n2 is 14 and n3 is 2.

32. The oligomeric compound of claim 1 wherein n_1 is from 1 to about 3, n_2 is 16 or 17 and n_3 is 2 or 3.


33. The oligomeric compound of claim 1 wherein n_1 is 2, n_2 is 16 and n_3 is 2.

34. The oligomeric compound of claim 1 wherein at least one Nu_1 nucleoside and at least one Nu_3 nucleoside is an LNA or ENA.

35. The oligomeric compound of claim 1 wherein at least one of the 5'-most or 3'-most terminal affinity modified nucleosides is an LNA or ENA.

36. A method of inhibiting gene expression comprising contacting one or more cells, a tissue, or an animal with an oligomeric compound of claim 1.

37. An oligomeric compound comprising the structure:



wherein:

each Bx is a heterocyclic base moiety;

each X is, independently, O or S;

T_1 and T_2 are each, independently, H, a hydroxy protecting group, an optionally linked conjugate group, or a covalent attachment to a solid support medium;

each m is, independently, 1 or 2;

n_a is from 1 to about 6;

n_b is from 11 to about 18; and

n_c is from 2 to about 6.

38. The oligomeric compound of claim 37 wherein each m is 1.

39. The oligomeric compound of claim 37 wherein each m is 2.

40. The oligomeric compound of claim 37 wherein at least one of T_1 and T_2 is H or a hydroxyl protecting group.

41. The oligomeric compound of claim 40 wherein each of the hydroxyl protecting groups is, independently, 4,4'-dimethoxytrityl, monomethoxytrityl, 9-phenylxanthene-9-yl, 9-(p-methoxyphenyl)xanthene-9-yl, t-butyl, t-butoxymethyl, methoxymethyl, tetrahydropyranyl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-trimethylsilylethyl, p-chlorophenyl,

2,4-dinitrophenyl, benzyl, 2,6-dichlorobenzyl, diphenylmethyl, p,p-dinitrobenzhydryl, p-nitrobenzyl, triphenylmethyl, trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, triphenylsilyl, benzoylformate, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, pivaloyl, benzoyl, p-phenylbenzoyl, mesyl, tosyl, 4,4',4''-tris-(benzyloxy)trityl, 4,4',4''-tris-(4,5-dichlorophthalimido)trityl, 4,4',4''-tris(levulinyl)trityl, 3-(imidazolylmethyl)-4,4'-dimethoxytrityl, 4-decyloxytrityl, 4-hexadecyloxytrityl, 9-(4-octadecyloxyphenyl)xanthene-9-yl, 1,1-bis-(4-methoxyphenyl)-1'-pyrenyl methyl, p-phenylazophenylloxycarbonyl, 9-fluorenylmethoxycarbonyl, 2,4-dinitrophenylethoxycarbonyl, 4-(methylthiomethoxy)butyryl, 2-(methylthiomethoxymethyl)-benzoyl, 2-(isopropyl-thi-

omethoxymethyl)benzoyl, 2-(2,4-dinitrobenzenesulphenyloxymethyl)benzoyl, or levulinyl group.

42. The oligomeric compound of claim 37 wherein one of T_1 and T_2 is a covalent attachment to a support medium.

43. The oligomeric compound of claim 42 wherein the support medium is a controlled pore glass, oxalyl-controlled pore glass, silica-containing particles, polymers of polystyrene, copolymers of polystyrene, copolymers of styrene and divinylbenzene, copolymers of dimethylacrylamide and N,N'-bisacryloylethylenediamine, soluble support medium, or PEPS.

44. The oligomeric compound of claim 37 wherein each L_1 , L_2 and L_3 is, independently, a phosphodiester or a phosphorothioate internucleoside linking group.

45. The oligomeric compound of claim 44 wherein each L_1 , L_2 and L_3 is a phosphodiester internucleoside linking group.

46. The oligomeric compound of claim 37 wherein each Bx is, independently, adenine, guanine, thymine, cytosine, uracil, 5-methylcytosine, 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine, 2-thiocytosine, 5-halouracil, 5-halocytosine, 5-propynyl uracil, 5-propynyl cytosine, 6-azo uracil, 6-azo cytosine, 6-azo thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-substituted adenines and guanines, 5-substituted uracils and cytosines,

7-methylguanine, 7-methyladenine, 8-azaguanine, 8-azaadenine, 7-deazaguanine, 7-deazaadenine, 3-deazaguanine, or 3-deazaadenine.

47. The oligomeric compound of claim 37 wherein na is from 1 to about 5.

48. The oligomeric compound of claim 37 wherein na is from 1 to about 3.

49. The oligomeric compound of claim 37 wherein na is from 2 to about 3.

50. The oligomeric compound of claim 37 wherein nc is from 2 to about 5.

51. The oligomeric compound of claim 37 wherein nc is from 2 to about 3.

52. The oligomeric compound of claim 37 wherein nb is from 12 to about 18.

53. The oligomeric compound of claim 37 wherein nb is from 12 to about 16.

54. The oligomeric compound of claim 37 wherein nb is from 14 to about 16.

55. The oligomeric compound of claim 37 wherein the total of na, nb and nc is from 14 to about 30.

56. The oligomeric compound of claim 37 wherein the total of na, nb and nc is from 14 to 24.

57. The oligomeric compound of claim 37 wherein the total of na, nb and nc is from 14 to 21.

58. The oligomeric compound of claim 37 wherein the total of na, nb and nc is from 16 to 21.

59. The oligomeric compound of claim 37 wherein na is from 1 to about 3, nb is 12 or 13 and nc is 2 or 3.

60. The oligomeric compound of claim 37 wherein na is 3, nb is 12 and nc is 3.

61. The oligomeric compound of claim 37 wherein na is from 1 to about 3, nb is 14 or 15 and nc is 2 or 3.

62. The oligomeric compound of claim 37 wherein na is 2, nb is 14 and nc is 2.

63. The oligomeric compound of claim 37 wherein na is from 1 to about 3, nb is 16 or 17 and nc is 2 or 3.

64. The oligomeric compound of claim 37 wherein na is 2, nb is 16 and nc is 2.

65. A method of inhibiting gene expression comprising contacting one or more cells, a tissue, or an animal with an oligomeric compound of claim 37.

* * * * *