

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/074847 A1

- (51) International Patent Classification⁷: **C08K 5/00**,
5/3435 (US). **HORSEY, Douglas, Wayne** [US/US]; 5 Simpson
Road, Briarcliff Manor, NY 10510 (US).
- (21) International Application Number: PCT/EP02/02706 (74) **Common Representative: CIBA SPECIALITY
CHEMICALS HOLDING INC.**; Klybeckstrasse 141,
CH-4057 Basle (CH).
- (22) International Filing Date: 12 March 2002 (12.03.2002)
- (25) Filing Language: English (81) **Designated States (national):** AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.
- (26) Publication Language: English (84) **Designated States (regional):** ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).
- (30) Priority Data:
60/277,222 20 March 2001 (20.03.2001) US
- (63) **Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:**
US 60/277,222 (CON)
Filed on 20 March 2001 (20.03.2001)
- (71) **Applicant (for all designated States except US): CIBA
SPECIALITY CHEMICALS HOLDING INC.**
[CH/CH]; Klybeckstrasse 141, CH-4057 Basle (CH).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only): TROUTMAN, Mal-
isa, V.** [US/US]; 308 East 89th St. #3B, New York, NY
10128 (US). **RAVICHANDRAN, Ramanathan** [US/US];
1 Caddy Lane, Suffern, NY 10901 (US). **SRINIVSAN,
Rangarajan** [IN/US]; 177 White Plains Road, #83A,
Tarrytown, NY 10591 (US). **KING, Roswell, Easton**
[US/US]; 700 Bear Ridge Road, Pleasantville, NY 10570
- Published:**
— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 02/074847 A1

(54) **Title:** FLAME RETARDANT COMPOSITIONS

(57) **Abstract:** Polymeric substrates, for example polyolefins such as polypropylene, can be made flame retardant by the incorporation therein of a synergistic mixture of (i.) at least one compound selected from the group consisting of nitroxyl stabilizers, hydroxylamine stabilizers, nitron stabilizers, substituted hydroxylamine stabilizers, amine oxide stabilizers, benzofuranone stabilizers, phosphite and phosphonite stabilizers, quinone methide stabilizers and monoacrylate esters of 2,2'-alkylidenebisphenol stabilizers and (ii.) at least one compound selected from the group consisting of brominated flame retardants, phosphorus containing flame retardants and inorganic flame retardants such as ammonium polyphosphate or decabromodiphenyl oxide; wherein the amount of organic or inorganic flame retardant of component (ii.) required to achieve an acceptable level of flame retardancy is significantly reduced compared to that needed when component (i.) is not present.

FLAME RETARDANT COMPOSITIONS

The instant invention pertains to a novel method of flame retarding a polymeric substrate by adding thereto an effective flame retarding amount of a mixture of a synergist compound and a known organic or inorganic flame retardant. The synergists are selected from the group consisting of the nitroxyl stabilizers, hydroxylamine stabilizers, nitron stabilizers, substituted hydroxylamine stabilizers, amine oxide stabilizers, benzofuranone stabilizers, phosphite and phosphonite stabilizers, quinone methide stabilizers and monoacrylate esters of 2,2'-alkylidenebisphenol stabilizers.

Background of the Invention

United States Patent No. 5,393,812 describes polyolefin compositions which are made flame retardant by a combination of a halogenated hydrocarbyl phosphate or phosphonate ester flame retardant in combination with a alkoxyamine functional hindered amine, but there is no suggestion that the hindered amine itself is responsible for the flame retardancy, but rather that the hindered amine is preventing delustering and other undesirable effects from occurring in these polyolefin compositions.

European Application No. 0 792 911 A2, discloses that alkoxyamine functional hindered amines are effective when used to enhance the flame retarding efficacy of tris(trihalogenopentyl) phosphate flame retardants.

U.S. Pat. No. 6,117,995 discloses that certain N-alkoxy hindered amines may be used as flame retardants for organic polymers. Copending U.S. application Nos. 09/502,239, filed Nov. 3, 1999, and 09/714,717, filed Nov. 16, 2000, disclose the use of certain N-alkoxy hindered amines as flame retardants.

The flame retardant (FR) market today is comprised of products which function to interfere with the combustion process by chemical and/or physical means. Mechanistically these FRs have been proposed to function during combustion of an article in either the gas phase, the condensed phase or both. The organohalogenes are proposed to generate halogen species (e.g. HX) which interferes in the gas phase with free radical organic "fuel"

from the polymer substrate. Synergists are proposed to react with HX to form additional chemical species which interfere with combustion in the gas phase, such as reaction of antimony oxide with HX to form antimony halide and water vapor. Still other flame retardant classes are proposed to impart efficacy in the "condensed" phase such as forming a protective char layer on the polyester, or forming an intumescent or foaming on the polymer surface. The char or intumescent layer is thought either to prevent organic fuel from migrating from the polymer into the vapor phase where it can fuel combustion, or the char can act as a thermal shield to protect the underlying polymer article from thermally induced decomposition and generation of fuel. Phosphorus compounds of various classes (e.g. halo- or non-halogenated) are an example. Further still, other classes of compounds are proposed to function in the condensed and/or vapor phase. Metal hydrates or metal oxides are proposed to generate water vapor under thermal conditions, the water acting to dilute the fuel mix in the combustion zone and to remove heat from the flame zone via conversion of water to vapor. Alumina trihydrate, magnesium hydroxide or oxide, and other compounds are reported to function in this way.

These state of the art chemistries described above have various detrimental aspects in addition to the effective flame retarding attributes mentioned. Certain organobrominated compounds are under governmental scrutiny for the generation of toxic by-products during the production or combustion such as dioxanes from polybrominated diphenyl oxides. Certain metal-containing flame retardants, notably antimony oxides, are under scrutiny for worker exposure and toxicity reasons. Antimony oxides often contain trace amounts of arsenic compounds which are suspected carcinogens. Overall, a growing concern has arisen regarding the generation of smoke and toxic gases which are evolved from these flame retardants during a fire. While the classic FRs may be effective combustion suppressants, the toxic gases they form pose a threat to human exposure.

The instant invention alleviates some of the detrimental aspects of the current state of the art which the use of large amounts of commercial flame retardants pose. The present synergist compounds are non-halogenated and free of heavy metals, thus avoiding generation of corrosive HX gases and avoiding exposure to toxic metals. In some applications, the instant invention provides a direct replacement for current FR systems where the instant synergist compounds provide a complimentary enhancement or synergistic

system (e.g. antimony oxide replacement in ABS) where good flame retardancy can be achieved by using less classic FR agent in the presence of the instant synergist compounds.

Hydroxylamine stabilizers are disclosed in U.S. Patent Nos. 4,590,231, 4,612,393, 4,649,221, 4,668,721, 4,691,015, 4,696,964, 4,703,073, 4,720,517, 4,757,102, 4,782,105, 4,831,134, 4,876,300, 5,006,577, 5,019,285, 5,064,883, 5,185,448 and 5,235,056.

Nitrone stabilizers of component are described in U.S. Pat. No. 4,898,901.

U.S. Pat. Nos. 4,666,962, 4,666,963, 4,678,826, 4,753,972, 4,757,102, 4,760,179, 4,929,657, 5,057,563, 5,021,479, 5,045,583 and 5,185,448 disclose the use of various substituted hydroxylamine stabilizers towards the stabilization of organic materials.

U.S. Patent Nos. 5,081,300, 5,162,408, 5,844,029, 5,880,191 and 5,922,794 disclose the use of saturated hydrocarbon amine oxides towards the stabilization of thermoplastic resins.

Benzofuranone stabilizers are disclosed for example in U.S. Patent Nos. 4,325,863; 4,338,244; 5,175,312; 5,216,052; 5,252,643; 5,369,159; 5,488,117; 5,356,966; 5,367,008; 5,428,162; 5,428,177; 5,614,572; 5,883,165 and 5,516,920.

Quinone methide stabilizers are disclosed for example in U.S. Pat. Nos. 5,583,247, 5,616,774, 5,670,692 and 5,750,765.

O-alkenyl substituted hydroxylamine stabilizers are disclosed in U.S. Pat. No. 5,045,583.

Non-hindered alkoxyamine stabilizers are disclosed in U.S. Pat. No. 5,185,448.

Detailed Disclosure

The instant invention pertains to a flame retardant composition which comprises

(A) a polymer substrate, and

(B) an effective flame retarding amount of a synergistic mixture of


(i.) at least one compound selected from the group consisting of

- (a) nitroxyl stabilizers,
- (b) hydroxylamine stabilizers,
- (c) nitron stabilizers,
- (d) substituted hydroxylamine stabilizers,
- (e) amine oxide stabilizers,
- (f) benzofuranone stabilizers,
- (g) phosphite and phosphonite stabilizers,
- (h) quinone methide stabilizers and
- (j) monoacrylate esters of 2,2'-alkylidenebisphenol stabilizers and

(ii.) at least one compound selected from the group consisting of brominated flame retardants, phosphorus containing flame retardants and inorganic flame retardants.


Component (a)

The nitroxyl stabilizers of component (a) useful in this invention have the generic structure




or are compounds that contain one or more groups of the formula

- 5 -



where each R is alkyl and T is a group required to complete a 5- or 6-membered ring.

Two or more nitroxyl groups may be present in the same molecule by being linked through the T moiety as exemplified below where E is a linking group.



Typical nitroxyls of component (a) include bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) sebacate, 4-hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine, 4-ethoxy-1-oxyl-2,2,6,6-tetramethylpiperidine, 4-propoxy-1-oxyl-2,2,6,6-tetramethylpiperidine, 4-acetamido-1-oxyl-2,2,6,6-tetramethylpiperidine, 1-oxyl-2,2,6,6-tetramethylpiperidine, 1-oxyl-2,2,6,6-tetramethylpiperidin-4-one, 1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl acetate, 1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl 2-ethylhexanoate, 1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl stearate, 1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl benzoate, 1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl 4-*t*-butylbenzoate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) succinate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) adipate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) *n*-butylmalonate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) phthalate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) isophthalate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) terephthalate, bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) hexahydroterephthalate, N,N'-

bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl)adipamide, N-(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl)caprolactam, N-(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl)dodecylsuccinimide, 2,4,6-tris-[N-butyl-N-(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl)]-s-triazine, 4,4'-ethylenebis(1-oxyl-2,2,6,6-tetramethylpiperazin-3-one), 2-oxyl-1,1,3,3-tetramethyl-2-isobenzazole, 1-oxyl-2,2,5,5-tetramethylpyrrolidine, and N,N-bis-(1,1,3,3-tetramethylbutyl)nitroxide.


Nitroxyl stabilizers of component (a) are for example bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) sebacate, 4-hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine, 4-ethoxy-1-oxyl-2,2,6,6-tetramethylpiperidine, 4-propoxy-1-oxyl-2,2,6,6-tetramethylpiperidine, 4-acetamido-1-oxyl-2,2,6,6-tetramethylpiperidine, 1-oxyl-2,2,6,6-tetramethylpiperidine, and 1-oxyl-2,2,6,6-tetramethylpiperidin-4-one.

A specific embodiment is where the nitroxyl stabilizers of component (a) are bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) sebacate and 4-hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine.

Component (b)

Hydroxylamine stabilizers of component (b) are for example those disclosed in U.S. Patent Nos. 4,590,231, 4,612,393, 4,649,221, 4,668,721, 4,691,015, 4,696,964, 4,703,073, 4,720,517, 4,757,102, 4,782,105, 4,831,134, 4,876,300, 5,006,577, 5,019,285, 5,064,883, 5,185,448 and 5,235,056, the relevant parts of which are incorporated herein by reference.

The hydroxylamine stabilizers of component (b) employed in the novel compositions and methods are for example of the formula (I)




wherein

T₁ is straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms; and

T₂ is hydrogen, or independently has the same meaning as T₁.

Alternatively, the hydroxylamine stabilizers of component (b) of the present invention are compounds that contain one or more of the groups of the formula (II)



wherein

T is a group forming a five- or six-membered ring; and

R₁ and R₂ are independently hydrogen, alkyl of 1 to 4 carbon atoms or phenyl.

In the present invention the compounds of component (b) are, for example, N,N-dihydrocarbylhydroxylamines of formula (I) wherein T₁ and T₂ are independently benzyl, methyl, ethyl, octyl, lauryl, dodecyl, tetradecyl, hexadecyl, heptadecyl or octadecyl, or wherein T₁ and T₂ are each the alkyl mixture found in hydrogenated tallow amine.


The compounds of component (b) in the present compositions and methods are, for example, N,N-dihydrocarbylhydroxylamines selected from the group consisting of N,N-dibenzylhydroxylamine, N,N-diethylhydroxylamine, N,N-dioctylhydroxylamine, N,N-dilaurylhydroxylamine, N,N-didodecylhydroxylamine, N,N-ditetradecylhydroxylamine, N,N-dihexadecylhydroxylamine, N,N-dioctadecylhydroxylamine, N-hexadecyl-N-tetradecylhydroxylamine, N-hexadecyl-N-heptadecylhydroxylamine, N-hexadecyl-N-octadecylhydroxylamine, N-heptadecyl-N-octadecylhydroxylamine, N-methyl-N-octadecylhydroxylamine and N,N-di(hydrogenated tallow)hydroxylamine.

Component (b) in the present invention may be for example the N,N-di(alkyl)hydroxylamine produced by the direct oxidation of N,N-di(hydrogenated tallow)amine (Irgastab[®] FS-042, Ciba Specialty Chemicals Corp.).

Component (c)

The nitrones of component (c) may be for example as described in U.S. Pat. No. 4,898,901, which is hereby incorporated by reference.

The nitrones of component (c) are for example of the formula (III)



wherein

L₁ is straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms;

L₂ and L₃ are independently hydrogen, straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms;

or L₁ and L₂ together form a five- or six-membered ring including the nitrogen atom.


The nitrones of component (c) may be the corresponding oxidation products of the hydroxylamines of component (b). That is to say, the nitrones of component (c) may be nitronium analogues of the hydroxylamines of component (b). The nitrones may be for example, N-benzyl- α -phenylnitronium, N-ethyl- α -methylnitronium, N-octyl- α -heptylnitronium, N-lauryl- α -undecylnitronium, N-tetradecyl- α -tridcylnitronium, N-hexadecyl- α -pentadecylnitronium, N-octadecyl- α -heptadecylnitronium, N-hexadecyl- α -heptadecylnitronium, N-octadecyl- α -pentadecylnitronium, N-heptadecyl- α -heptadecylnitronium, N-octadecyl- α -hexadecylnitronium, N-methyl- α -heptadecylnitronium and the nitronium derived from N,N-di(hydrogenated tallow)hydroxylamine.

Component (d)

The substituted hydroxylamine stabilizers of component (d) are for example those described in U.S. Pat. Nos. 4,666,962, 4,666,963, 4,678,826, 4,753,972, 4,757,102, 4,760,179, 4,929,657, 5,057,563, 5,021,479, 5,045,583 and 5,185,448 the disclosures of which are hereby incorporated by reference. Component (d) includes the Michael addition products from the reaction of the hydroxylamines of component (b) with any α,β -unsaturated ketone, ester, amide, or phosphonate. Component (d) also includes Mannich-type condensation products from the reaction of the hydroxylamines of component (b) with formaldehyde and secondary amines. Component (d) also includes O-alkenyl substituted analogues of the present hydroxylamines of component (b) as disclosed in U.S. Pat. No. 5,045,583. Component (d) also includes non-hindered substituted hydroxylamine stabilizers as disclosed in U.S. Pat. No. 5,185,448. Component (d) also includes acyl derivatives of the hydroxylamine stabilizers of component (b), for example such as those disclosed in U.S. Pat. No. 5,021,479.

The substituted hydroxylamines of component (d) may be derivatives of the above-described hydroxylamines of formulae (I) and (II), provided that if they are derivatives of hydroxylamines of formula (II), that they are limited to derivatives of hydroxylamines as described in U.S. Pat. Nos. 5,185,448 and 5,235,056.

The present substituted hydroxylamines may be for example of the formula (IIIb) or (IIIa)



wherein

T_1 is straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms;




T_2 is hydrogen, or independently has the same meaning as T_1 ; and

T_3 is allyl, straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 18 carbon atoms, cycloalkenyl of 5 to 18 carbon atoms or a straight or branched chain alkyl

of 1 to 4 carbon atoms substituted by phenyl or by phenyl substituted by one or two alkyl groups of 1 to 4 carbon atoms or by 1 or 2 halogen atoms.

The substituted hydroxylamines of component (d) may be for example O-allyl-N,N-dioctadecylhydroxylamine or O-n-propyl-N,N-dioctadecylhydroxylamine or N,N-di(hydrogenated tallow)acetoxylamine.

The compounds of component (d) include the following general structures:



wherein

in the compounds of formula (IV)


a is 1 to 4;

R' and R'' are independently hydrogen, alkyl of 1 to 36 carbon atoms, cycloalkyl of 5

to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms or said aralkyl substituted by alkyl of 1 to 36 carbon atoms,

R_b and R_c are independently hydrogen or alkyl of 1 to 12 carbon atoms;


A, when $a = 1$, is a group NHR_d , wherein R_d is hydrogen, amino, alkyl of 1 to 18 carbon atoms or



where B is a direct bond or alkylene of 1 to 10 carbon atoms;


A, when $a = 2$, is a bivalent radical of a 5-7 membered heterocyclic compound containing two nitrogen atoms in the ring, with the free valencies on the nitrogen atoms, or is a group $-N(R_d)-B-N(R_d)-$ where B and R_d have the meanings given above;

A, when $a = 3$, is a group of the formula



where n is 2 to 6 and R_e is hydrogen or alkyl of 1 to 4 carbon atoms; and

A, when $a = 4$, is a group of the formula




where R_e and n have the meanings given above.


The R' and R'' groups are for example hydrogen, straight-chain or branched alkyl with 1 to 18 carbon atoms such as methyl, ethyl, n-propyl, n-butyl, tert-butyl, n-pentyl, n-octyl, 2-ethylhexyl, decyl, dodecyl and octadecyl; cyclopentyl and cyclohexyl; and benzyl,

α -methylbenzyl and α,α -dimethyl-benzyl. A specific embodiment is where R_b and R_c are hydrogen or alkyl of 1 to 4 carbon atoms.

Other specific groups include, for $a = 1$, R_d as C_1 - C_{18} alkyl (see list for R' and R'') and B as C_1 - C_6 alkylene; for $a = 2$,



R_d as hydrogen and B as straight chain C_1 - C_6 alkylene; and for $a = 3$, A as



In compounds of formula (V),


R_f , R_g , R_h and R_i independently are hydrogen, alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms or said aralkyl substituted by alkyl of 1 to 36 carbon atoms; and the alkali metal and alkaline earth metal salts thereof.

Specific embodiments of the R_f - R_i groups are where they are independently straight-chain or branched alkyl with 1 to 18 carbon atoms, for instance with 1 to 12 carbon atoms such as methyl, ethyl, n-propyl, n-butyl, tert-butyl, n-pentyl, n-octyl, 2-ethylhexyl, decyl, dodecyl, tetradecyl and octadecyl; cyclopentyl and cyclohexyl; and benzyl, α -methylbenzyl and α,α -dimethylbenzyl. Another specific embodiment are compounds having R_f and R_g as benzyl and R_h and R_i as alkyl of 1 to 8 carbon atoms. Typical salts include the sodium,

potassium, calcium and magnesium salts.

In compounds of formula (VI),

R_j and R_k are independently alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms or said aralkyl substituted by alkyl of 1 to 36 carbon atoms, or R_j and R_k together with the nitrogen can form a 5-7 member heterocyclic ring, or R_j and R_k are independently a group of the formula



R_i , R_m , R_n , R_o and R_p are independently hydrogen, alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, allyl, aryl, aralkyl of 7 to 9 carbon atoms, said aralkyl substituted by alkyl of 1 to 36 carbon atoms, bornyl, norbornyl or isobornyl, or R_m and R_n together with nitrogen can form a 5-7 member heterocyclic ring;

b is 1-4;

x is 0-5;

T_a is R_i or $-\text{CH}(R_i)-\text{O}-\text{N}(R_j)(R_k)$;

when $b = 1$, T_b is R_i or 5-7 member unsaturated heterocyclic-substituted (C_1 - C_4)alkyl, or T_a and T_b together with the nitrogen can form a 5-7 member saturated heterocyclic ring;

when $b = 2$, T_b is alkylene of 2 to 12 carbon atoms, cycloalkylene of 6 to 10 carbon atoms, arylene of 6 to 10 carbon atoms or alkylenearylenealkylene of 8 to 10 carbon atoms or T_a and T_b together with the two nitrogens can form a 5-7 member saturated heterocyclic ring;


when $b = 3$, T_b is alkanetriyl of 3 to 6 carbon atoms or T_a and T_b together with the three nitrogens can form a 5-6 member saturated heterocyclic ring; and

when $b = 4$, T_b is alkanetetriyl of 4 to 6 carbon atoms.

Representative R_j - R_p groups are straight-chain or branched alkyl with 1 to 18 carbon atoms such as methyl, ethyl, n-propyl, n-butyl, tert-butyl, n-pentyl, n-octyl, 2-ethylhexyl, decyl, dodecyl and octadecyl; cyclopentyl or cyclohexyl; and benzyl, α -methylbenzyl and α,α -dimethylbenzyl. Specific embodiments are where R_j and R_k as benzyl or substituted benzyl

and R_l as hydrogen. R_j/R_k and R_m/R_n heterocyclic groups include piperidyl, pyrrol, morpholino or pyrrolidino. T_a is for example one of the above noted R_j-R_p representative groups or the substituted aminoxy group, while T_b when $b = 1$, may also be one of the above noted R_j-R_p representative groups or pyridin-2-yl alkyl. T_a/T_b ($b = 1$) includes pyrrolidino, piperidyl and morpholino. T_b when $b = 2$ is, for example, ethylene, propylene, hexamethylene, phenylene and xylylene, and T_a/T_b heterocyclic is for example, piperazine or pyrazolidine. T_b when $b = 3$ is, for example, trimethylpropane and T_a/T_b heterocyclic is fully saturated triazine or triazole. T_b when $b = 4$ is, for example, pentaerythryl.

Typical $b = 2$ and 3 heterocyclic compounds are




A specific example of a compound of formula (V) is diethyl (N,N-dioctadecylaminoxy)ethyl phosphonate.

Component (e)

The amine oxide stabilizers of component (e) are for example those disclosed in U.S. Pat. Nos. 5,081,300, 5,162,408, 5,844,029, 5,880,191 and 5,922,794, the relevant parts of each incorporated herein by reference.

The amine oxide stabilizers of component (e) are for example saturated tertiary amine oxides as represented by general formula (VII):



wherein

G_1 and G_2 are independently a straight or branched chain alkyl of 6 to 36 carbon atoms, aryl of 6 to 12 carbon atoms, aralkyl of 7 to 36 carbon atoms, alkaryl of 7 to 36 carbon atoms, cycloalkyl of 5 to 36 carbon atoms, alkylcycloalkyl of 6 to 36 carbon atoms or cycloalkylalkyl of 6 to 36 carbon atoms;

G_3 is a straight or branched chain alkyl of 1 to 36 carbon atoms, aryl of 6 to 12 carbon atoms, aralkyl of 7 to 36 carbon atoms, alkaryl of 7 to 36 carbon atoms, cycloalkyl of 5 to 36 carbon atoms, alkylcycloalkyl of 6 to 36 carbon atoms or cycloalkylalkyl of 6 to 36 carbon atoms; and

wherein said aryl groups may be substituted by one to three halogen, alkyl of 1 to 8 carbon atoms, alkoxy of 1 to 8 carbon atoms or combinations thereof; and

wherein said alkyl, aralkyl, alkaryl, cycloalkyl, alkylcycloalkyl and cycloalkylalkyl groups may be interrupted by -O-, -S-, -SO-, -SO₂-, -COO-, -OCO-, -CO-, -NG₄-, -CONG₄- and -NG₄CO- groups, or wherein said alkyl, aralkyl, alkaryl, cycloalkyl, alkylcycloalkyl and cycloalkylalkyl groups may be substituted by -OG₄-, -SG₄-, -COOG₄-, -OCOG₄-, -COG₄-, -N(G₄)₂-,

-CON(G₄)₂, -NG₄COG₄ and 5- and 6-membered rings containing the -C(CH₃)(CH₂R_x)NL(CH₂R_x)(CH₃)C- group or wherein said alkyl, aralkyl, alkaryl, cycloalkyl, alkycycloalkyl and cycloalkylalkyl groups are both interrupted and substituted by the groups mentioned above; and

wherein

G₄ is independently hydrogen or alkyl of 1 to 8 carbon atoms;

R_x is hydrogen or methyl;

L is hydrogen, hydroxy, C₁₋₃₀ straight or branched chain alkyl moiety, a -C(O)R moiety where R is a C₁₋₃₀ straight or branched chain alkyl group, or a -OR_y moiety; and

R_y is C₁₋₃₀ straight or branched chain alkyl, C₂-C₃₀ alkenyl, C₂-C₃₀ alkynyl, C₅-C₁₂ cycloalkyl, C₆-C₁₀ bicycloalkyl, C₅-C₈ cycloalkenyl, C₆-C₁₀ aryl, C₇-C₉ aralkyl, C₇-C₉ aralkyl substituted by alkyl or aryl, or -CO(D), where D is C₁-C₁₈ alkyl, C₁-C₁₈ alkoxy, phenyl, phenyl substituted by hydroxy, alkyl or alkoxy, or amino or amino mono- or di-substituted by alkyl or phenyl.

Within compounds of the formula (VII), at least one of G₁, G₂ and G₃ contains a β-carbon-hydrogen bond, i.e. a carbon-hydrogen bond on the second carbon atom from the nitrogen.


The number of substituents or interrupting groups in G₁, G₂ or G₃ as alkyl, aralkyl, alkaryl, cycloalkyl, alkycycloalkyl and cycloalkylalkyl defined above, if present, are from one to sixteen, within the limits given.

Examples of structures of formula (VII) are where G₁ and G₂ are independently benzyl or substituted benzyl. It is also possible for each of G₁, G₂, and G₃ to be the same residue. G₁ and G₂ may also independently be alkyl groups of 8 to 26 carbon atoms, for example alkyl groups of 10 to 26 carbon atoms. G₃ may be an alkyl group of 1 to 22 carbon atoms, for example methyl or substituted methyl. Also, the present amine oxides include those wherein G₁, G₂, and G₃ are the same alkyl groups of 6 to 36 carbon atoms. The aforementioned residues for G₁, G₂, and G₃ are, for instance, saturated hydrocarbon residues or saturated hydrocarbon residues containing at least one of the aforementioned -O-, -S-, -SO-, -CO₂-, -CO-, or -CON- moieties. Those skilled in the art will be able to envision other useful residues for each of G₁, G₂, and G₃ without detracting from the present invention.

The saturated amine oxides of component (e) may also include poly(amine oxides). By poly(amine oxides) is meant tertiary amine oxides containing at least two tertiary amine oxides per molecule. Illustrative poly(amine oxides), also called "poly(tertiary amine oxides)", include the tertiary amine oxide analogues of aliphatic and alicyclic diamines such as, for example, 1,4-diaminobutane; 1,6-diaminohexane; 1,10-diaminodecane; and 1,4-diaminocyclohexane, and aromatic based diamines such as, for example, diamino anthraquinones and diaminoanisoles.

Also included as component (e) are tertiary amine oxides derived from oligomers and polymers of the aforementioned diamines. Useful amine oxides also include amine oxides attached to polymers, for example, polyolefins, polyacrylates, polyesters, polyamides, polystyrenes, and the like. When the amine oxide is attached to a polymer, the average number of amine oxides per polymer can vary widely as not all polymer chains need to contain an amine oxide. All of the aforementioned amine oxides may optionally contain at least one -O-, -S-, -SO-, -CO₂-, -CO-, or -CONG₄- moiety. For instance, each tertiary amine oxide of the polymeric tertiary amine oxide may contain a C₁ residue.

The groups G₁, G₂ and G₃ of formula (VII) may be attached to a molecule containing a hindered amine. Hindered amines are known in the art and the amine oxide of the present invention may be attached to the hindered amine in any manner and structural position of the hindered amine. Useful hindered amines when part of a compound of component (e) include those of the general formulas:




wherein L and R_x are as described above. Also included are amine oxides containing more than one hindered amine and more than one saturated amine oxide per molecule. The hindered amine may be attached to a poly(tertiary amine oxide) or attached to a polymeric substrate, as discussed above.


Component (f)

The benzofuranone stabilizers of component (f) are those disclosed for example in U.S. Patent Nos. 4,325,863; 4,338,244; 5,175,312; 5,216,052; 5,252,643; 5,369,159; 5,488,117; 5,356,966; 5,367,008; 5,428,162; 5,428,177; 5,614,572; 5,883,165 or 5,516,920, all incorporated herein by reference, or 3-(4-(2-acetoxyethoxy)phenyl)-5,7-di-tert-butyl-benzofuran-2-one, 5,7-di-tert-butyl-3-(4-(2-stearoyloxyethoxy)phenyl)benzofuran-2-one, 3,3'-bis(5,7-di-tert-butyl-3-(4-(2-hydroxyethoxy)phenyl)benzofuran-2-one), 5,7-di-tert-butyl-3-(4-ethoxyphenyl)benzofuran-2-one, 3-(4-acetoxy-3,5-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one, 3-(3,5-dimethyl-4-pivaloyloxyphenyl)-5,7-di-tert-butyl-benzofuran-2-one, 3-(3,4-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one, 3-(2,3-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one.

For example, component (f) is one or more benzofuranone stabilizers selected from:







(BF1) and





(BF2).

Component (g)

Component (g) is at least one compound selected from the formulae (1), (2), (3), (4), (5), (6) and (7)






in which the indices are integral and


n is 2, 3 or 4; p is 1 or 2; q is 2 or 3; r is 4 to 12; y is 1, 2 or 3; and z is 1 to 6;

A₁, if n is 2, is C₂-C₁₈ alkylene; C₂-C₁₂ alkylene interrupted by oxygen, sulfur or -NR₄-;



phenylene;

A₁, if n is 3, is a radical of the formula -C_rH_{2r-1}-;



A₂ is as defined for A₁ if n is 2;

- 21 -

B is a direct bond, $-\text{CH}_2-$, $-\text{CHR}_4-$, $-\text{CR}_1\text{R}_4-$, sulfur, C_5 - C_7 cycloalkylidene, or cyclohexylidene which is substituted by from 1 to 4 C_1 - C_4 alkyl radicals in position 3, 4 and/or 5;

D_1 , if p is 1, is C_1 - C_4 alkyl and, if p is 2, is $-\text{CH}_2\text{OCH}_2-$;

D_2 , if p is 1, is C_1 - C_4 alkyl;


E, if y is 1, is C_1 - C_{18} alkyl, $-\text{OR}_1$ or halogen;

E, if y is 2, is $-\text{O}-\text{A}_2-\text{O}-$,

E, if y is 3, is a radical of the formula $\text{R}_4\text{C}(\text{CH}_2\text{O})_3$ or $\text{N}(\text{CH}_2\text{CH}_2\text{O})_3$;

Q is the radical of an at least z-valent alcohol or phenol, this radical being attached via the oxygen atom to the phosphorus atom;

R_1 , R_2 and R_3 independently of one another are C_1 - C_{18} alkyl which is unsubstituted or substituted by halogen, $-\text{COOR}_4$, $-\text{CN}$ or $-\text{CONR}_4\text{R}_4$; C_2 - C_{18} alkyl interrupted by oxygen, sulfur or $-\text{NR}_4-$; C_7 - C_9 phenylalkyl; C_5 - C_{12} cycloalkyl, phenyl or naphthyl; naphthyl or phenyl substituted by halogen, 1 to 3 alkyl radicals or alkoxy radicals having a total of 1 to 18 carbon

atoms or by C_7 - C_9 phenylalkyl; or a radical of the formula  in which

m is an integer from the range 3 to 6;

R_4 is hydrogen, C_1 - C_{18} alkyl, C_5 - C_{12} cycloalkyl or C_7 - C_9 phenylalkyl,

R_5 and R_6 independently of one another are hydrogen, C_1 - C_8 alkyl or C_5 - C_6 cycloalkyl,

R_7 and R_8 , if q is 2, independently of one another are C_1 - C_4 alkyl or together are a 2,3-dehydropentamethylene radical; and

- 22 -

R₇ and R₈, if q is 3, are methyl;

R₁₄ is hydrogen, C₁-C₉ alkyl or cyclohexyl,

R₁₅ is hydrogen or methyl and, if two or more radicals R₁₄ and R₁₅ are present, these radicals are identical or different,

X and Y are each a direct bond or oxygen,

Z is a direct bond, methylene, -C(R₁₆)₂- or sulfur, and

R₁₆ is C₁-C₈ alkyl.

Of interest are compositions and methods wherein component (g) is at least one compound selected from the formulae (1), (2), (5) and (6), in which

n is the number 2, and y is the number 1, 2 or 3;

A₁ is C₂-C₁₈ alkylene, p-phenylene or p-biphenylene,

E, if y is 1, is C₁-C₁₈ alkyl, -OR₁ or fluorine;

E, if y is 2, is p-biphenylene,

E, if y is 3, is N(CH₂CH₂O-)₃,

R₁, R₂ and R₃ independently of one another are C₁-C₁₈ alkyl, C₇-C₉ phenylalkyl, cyclohexyl, phenyl, or phenyl substituted by 1 to 3 alkyl radicals having a total of 1 to 18 carbon atoms;

R₁₄ is hydrogen or C₁-C₉ alkyl,

R₁₅ is hydrogen or methyl;

X is a direct bond,

Y is oxygen,

Z is a direct bond or $-\text{CH}(\text{R}_{16})-$, and

R_{16} is C_1 - C_4 alkyl.

Likewise of interest are compositions and methods wherein component (g) is at least one compound selected from the formulae (1), (2), (5) and (6), in which

n is the number 2 and y is the number 1 or 3;

A_1 is p-biphenylene,

E, if y is 1, is C_1 - C_{18} alkoxy or fluorine,

E, if y is 3, is $\text{N}(\text{CH}_2\text{CH}_2\text{O})_3$,

R_1 , R_2 and R_3 independently of one another are C_1 - C_{18} alkyl, or phenyl substituted by 2 or 3 alkyl radicals having a total of 2 to 12 carbon atoms;

R_{14} is methyl or tert-butyl;

R_{15} is hydrogen;


X is a direct bond;

Y is oxygen; and

Z is a direct bond, methylene or $-\text{CH}(\text{CH}_3)-$.

Component (g) in the present invention may be at least one compound selected from the formulae (1), (2) and (6).

Component (g) in the present invention may be at least one compound of the formula



in which

R_{17} and R_{18} independently of one another are hydrogen, C_1 - C_8 alkyl, cyclohexyl or phenyl, and





R_{19} and R_{20} independently of one another are hydrogen or C_1 - C_4 alkyl.






The following compounds are examples of organic phosphites and phosphonites which are suitable for use in component (g):

Triphenyl phosphite, diphenyl alkyl phosphites, phenyl dialkyl phosphites, tris(nonylphenyl) phosphite (TNPP[®], GE), trilauryl phosphite, trioctadecyl phosphite, distearyl pentaerythritol diphosphite, tris(2,4-di-tert-butylphenyl) phosphite (Irgafos[®] 168, Ciba Specialty Chemicals Corp.), diisodecyl pentaerythritol diphosphite, bis(2,4-di-tert-butylphenyl) pentaerythritol diphosphite (formula (D)), bis(2,6-di-tert-butyl-4-methylphenyl) pentaerythritol diphosphite (formula (E)), bisisodecyloxy-pentaerythritol diphosphite, bis(2,4-di-tert-butyl-6-methylphenyl) pentaerythritol diphosphite, bis(2,4,6-tri-tert-butylphenyl) pentaerythritol diphosphite, tristearyl sorbitol triphosphite, tetrakis(2,4-di-tert-butylphenyl) 4,4'-biphenylene-diphosphonite (Irgafos[®] P-EPQ, Ciba Specialty Chemicals Corp., formula (H)), 6-isooctyloxy-2,4,8,10-tetra-tert-butyl-dibenzo[d,f][1,3,2]dioxaphosphepin (formula (C)), 6-fluoro-2,4,8,10-tetra-tert-butyl-12-methyl-dibenzo[d,g][1,3,2]dioxaphosphocin (formula (A)), bis(2,4-di-tert-butyl-6-methylphenyl) methyl phosphite, bis(2,4-di-tert-butyl-6-methylphenyl) ethyl phosphite (formula (G)).

The following phosphites and phosphonites may be used in component (g) in the novel compositions and methods:

Tris(2,4-di-tert-butylphenyl) phosphite (Irgafos[®] 168, Ciba Specialty Chemicals Corp.),
tris(nonylphenyl) phosphite (TNPP[®], GE),





Component (g) in the novel compositions and methods of the present invention may for example be: Tris(2,4-di-tert-butylphenyl) phosphite (Irgafos® 168, Ciba Specialty Chemicals


Corp.), 2,2'-ethylidenebis(2,4-di-tert-butylphenyl) fluorophosphite, (ETHANOX[®] 398, Ethyl Corp.), bis(2,4-di-tert-butyl-6-methylphenyl) ethyl phosphite (Irgafos[®] 38, Ciba Specialty Chemicals Corp., formula (G)), bis(2,4-di-tert-butylphenyl) pentaerythritol diphosphite (Ultrinox[®] 626, GE Chemicals, formula (D)), tetrakis(2,4-di-tert-butylphenyl)4,4'-biphenylene-diphosphonite (Irgafos[®] P-EPQ, Ciba Specialty Chemicals Corp., formula (H)), 2,2',2''-nitrilo[triethyltris(3,3',5,5'-tetra-tert-butyl-1,1'-biphenyl-2,2'-diyl)phosphite] (Irgafos[®] 12, Ciba Specialty Chemicals Corp., formula (B)), Ultrinox[®] 641 (GE Chemicals, formula (J)), Doverphos[®] S9228 (Dover Chemicals, formula (K)) or Mark[®] HP10 (Adeka Argus, formula (L)).

These organic phosphites and phosphonites are known compounds; many of them are commercially available.

Component (h)

The quinone methide stabilizers of component (h) of the present invention are those disclosed for example in U.S. Pat. Nos. 5,583,247, 5,616,774, 5,670,692 and 5,750,765, all incorporated herein by reference.

For example, the quinone methides of component (h) are of formula (VIII)



wherein

R_1' and R_2' are independently straight or branched chain alkyl of 4 to 18 carbon atoms, cycloalkyl of 5 to 12 carbon atoms or phenylalkyl of 7 to 15 carbon atoms, and

R_3' is 2-, 3- or 4-pyridyl; 2- or 3-thienyl; 2- or 3-pyrrolyl; 2- or 3-furyl; aryl of 6 to 10 carbon atoms; or said aryl substituted by one to three alkyl of 1 to 8 carbon atoms, alkoxy of 1 to 8 carbon atoms, alkylthio of 1 to 8 carbon atoms, alkylamino of 1 to 8 carbon atoms, dialkylamino of 2 to 8 carbon atoms, alkoxy carbonyl of 2 to 8 carbon atoms, hydroxy, nitro, amino, cyano, carboxy, aminocarbonyl, chloro or mixtures of said substituents; or R_3' is -CN, -COOH, -COOR_{4'}, -COR_{5'}, -OCOR_{6'}, -CONR_{7'}R_{8'} or -PO(OR_{9'})₂ where

R_4' is alkyl of 1 to 18 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, phenyl or benzyl,

R_5' is alkyl of 1 to 18 carbon atoms, aryl of 6 to 10 carbon atoms or said aryl substituted by 1 or 2 alkyl of 1 to 4 carbon atoms or by hydroxyl,

R_6' is alkyl of 1 to 18 carbon atoms, aryl of 6 to 10 carbon atoms or said aryl substituted by 1 or 2 alkyl of 1 to 4 carbon atoms or by hydroxyl,

R_7' and R_8' are independently hydrogen, alkyl of 1 to 18 carbon atoms or said alkyl substituted by alkylamino of 1 to 4 carbon atoms, by dialkylamino of 2 to 8 carbon atoms or by hydroxyl; benzyl, aryl of 6 to 10 carbon atoms or said aryl substituted by alkyl of 1 to 4 carbon atoms, by alkylamino of 1 to 4 carbon atoms, by dialkylamino of 2 to 8 carbon atoms, by phenylamino or by hydroxyl, or

-NR_{7'}R_{8'} is morpholino, piperidino or pyrrolidino, and

R_9' is hydrogen or alkyl of 1 to 18 carbon atoms.

For example, R_1' and R_2' are tert-butyl, tert-amyl, tert-octyl, cyclohexyl, α -methylbenzyl or α,α -dimethylbenzyl.

A specific embodiment is where R_1' and R_2' are tert-butyl, tert-amyl or tert-octyl.

In the compound of formula I, R_3' is for example phenyl or phenyl substituted by nitro, cyano, dimethylamino, methoxy, alkyl of 1 to 4 carbon atoms, hydroxy or mixtures of said substituents; specific embodiments are where R_3' is phenyl, or where R_3' is -CN, -COOH, -COOR_{4'}, -COR_{5'}, -OCOR_{6'}, -CONR_{7'}R_{8'} or -PO(OR_{9'})₂ where

R_4' is alkyl of 1 to 8 carbon atoms,

R_5' is methyl or phenyl,

R_6' is alkyl of 1 to 18 carbon atoms or phenyl,

R_7' and R_8' are independently hydrogen or alkyl of 1 to 18 carbon atoms, or

-NR_{7'}R_{8'} is morpholino or piperidino, and

R_9' is alkyl of 1 to 4 carbon atoms.

R_3' is for instance -CN, -COOH, -COOR_{4'}, -COR_{5'}, -CONR_{7'}R_{8'} or -PO(OR_{9'})₂ where

R_4' is alkyl of 1 to 4 carbon atoms,

R_5' is methyl or phenyl,

R_7' and R_8' are independently hydrogen or alkyl of 1 to 4 carbon atoms, or

-NR_{7'}R_{8'} is morpholino, and

R_9' is alkyl of 1 to 4 carbon atoms.


Specific examples of compounds of formula (VIII) are

(3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetonitrile,
(3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetic acid,
(3,5-di-tert-amyl-4-oxocyclohexa-2,5-dienylidene)acetic acid,
methyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetate,

ethyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetate,
 n-butyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetate,
 2,6-di-tert-butyl-4-(2-oxopropylidene)-cyclohexa-2,5-dienone,
 diethyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methanephosphonate,
 (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methyl acetate,
 (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methyl pivalate,
 (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methyl benzoate, and
 N,N-diethyl-2-(3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetamide,
 2,6-di-tert-butyl-4-benzylidene-cyclohexa-2,5-dienone,
 2,6-di-tert-butyl-4-(4-nitrobenzylidene)-cyclohexa-2,5-dienone,
 2,6-di-tert-butyl-4-(3-nitrobenzylidene)-cyclohexa-2,5-dienone,
 2,6-di-tert-butyl-4-(4-cyanobenzylidene)-cyclohexa-2,5-dienone,
 2,6-di-tert-butyl-4-(4-dimethylaminobenzylidene)-cyclohexa-2,5-dienone,
 2,6-di-tert-amyl-4-benzylidene-cyclohexa-2,5-dienone,
 2,6-di-tert-butyl-4-(4-methoxybenzylidene)-cyclohexa-2,5-dienone, and
 2,6-di-tert-butyl-4-(3,5-di-tert-butyl-4-hydroxybenzylidene)-cyclohexa-2,5-dienone;
 especially 2,6-di-tert-butyl-4-benzylidene-cyclohexa-2,5-dienone.

Component (j)

The monoacrylate esters of 2,2'-alkylidenebisphenol stabilizers are for example of the formula (IX)




wherein

R₄₀, R₄₁, R₄₂, R₄₃ and R₄₄ are independently straight or branched chain alkyl of 1 to 18 carbon atoms, cycloalkyl of 5 to 12 carbon atoms or phenylalkyl of 7 to 15 carbon atoms.

The present stabilizers of component (j) are for example monoacrylate esters of compounds selected from the group consisting of 2,2'-methylenebis(6-tert-butyl-4-methylphenol), 2,2'-methylenebis(6-tert-butyl-4-ethylphenol), 2,2'-methylenebis[4-methyl-6-(α -methylcyclohexyl)phenol], 2,2'-methylenebis(4-methyl-6-cyclohexylphenol), 2,2'-methylenebis(6-nonyl-4-methylphenol), 2,2'-methylenebis(4,6-di-tert-butylphenol), 2,2'-ethylidenebis(4,6-di-tert-butylphenol), 2,2'-ethylidenebis(6-tert-butyl-4-isobutylphenol), 2,2'-methylenebis[6-(α -methylbenzyl)-4-nonylphenol], 2,2'-methylenebis[6-(α,α -dimethylbenzyl)-4-nonylphenol], 2,6-bis(3-tert-butyl-5-methyl-2-hydroxybenzyl)-4-methylphenol and 1,1-bis-(3,5-dimethyl-2-hydroxyphenyl)butane.

For example, component (j) may be the monoacrylate ester of 2,2'-methylenebis(6-tert-butyl-4-methylphenol), available from Ciba Specialty Chemicals as Irganox[®] 3052:



In the structures of the present compounds of component (i.) if any substituents are alkyl, they are for example methyl, ethyl, n-propyl, n-butyl, sec-butyl, tert-butyl, n-hexyl, n-octyl, 2-ethylhexyl, n-nonyl, n-decyl, n-undecyl, n-dodecyl, n-tridecyl, n-tetradecyl, n-hexadecyl or n-octadecyl. Typical cycloalkyl groups include cyclopentyl and cyclohexyl; typical cycloalkenyl groups include cyclohexenyl; while typical aralkyl groups include benzyl, α -methyl-benzyl, α,α -dimethylbenzyl or phenethyl.

For example, component (i.) is at least one compound selected from the group consisting of

(a) bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) sebacate and 4-hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine,

(b) an N,N-di(alkyl)hydroxylamine produced by the direct oxidation of N,N-di(hydrogenated tallow)amine (Irgastab[®] FS-042),

(c) N-octadecyl- α -heptadecylnitrone,

(d) O-n-propyl-N,N-dioctadecylhydroxylamine,

(e) Genox[™] EP, a di(C₁₆-C₁₈)alkyl methyl amine oxide, CAS# 204933-93-7,

(f) 3-(3,4-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one, Irganox[®] HP-136, (BF1),

(g) tris(2,4-di-tert-butylphenyl) phosphite (Irgafos[®] 168),

(h) 2,6-di-tert-butyl-4-benzylidene-cyclohexa-2,5-dienone (Prostab[™] 6007)

and

(j) the monoacrylate ester of 2,2'-methylenebis(6-tert-butyl-4-methylphenol), Irganox[®] 3052.

Irganox, Irgafos, Prostab and Irgastab are trademarks of Ciba Specialty Chemicals. Genox[™] EP is available from GE Chemicals. N-octadecyl- α -heptadecylnitrone is as prepared in Example 3 of U.S. Pat. No. 4,898,901.

The polymeric substrate of component (A) is any of a wide variety of polymeric types including polyolefins, polystyrenics, and PVC. For example, the polymer substrate may be selected from the group of resins consisting of the polyolefins, the thermoplastic olefins, styrenic polymers and copolymers, ABS and polymers which contain hetero atoms, double

bonds or aromatic rings. Specific embodiments are where component (A) is polypropylene, polyethylene, thermoplastic olefin (TPO), ABS or high impact polystyrene.

For example, the polymer substrate is selected from the group of resins consisting of the polyolefins, the thermoplastic olefins, styrenic polymers and copolymers, and ABS.

Another embodiment of the present invention is where the polymer substrate is selected from the group consisting of polypropylene, polyethylene, thermoplastic olefin (TPO), ABS and high impact polystyrene.

For instance, the polymer substrate is polypropylene, polyethylene or thermoplastic olefin (TPO).

Examples for polyolefins are:

1. Polymers of monoolefins and diolefins, for example polypropylene, polyisobutylene, polybut-1-ene, poly-4-methylpent-1-ene, polyisoprene or polybutadiene, as well as polymers of cycloolefins, for instance of cyclopentene or norbornene, polyethylene (which optionally can be crosslinked), for example high density polyethylene (HDPE), high density and high molecular weight polyethylene (HDPE-HMW), high density and ultrahigh molecular weight polyethylene (HDPE-UHMW), medium density polyethylene (MDPE), low density polyethylene (LDPE), linear low density polyethylene (LLDPE), (VLDPE) and (ULDPE).

Polyolefins, i.e. the polymers of monoolefins exemplified in the preceding paragraph, for example polyethylene and polypropylene, can be prepared by different, and especially by the following, methods:

i) radical polymerization (normally under high pressure and at elevated temperature).

ii) catalytic polymerization using a catalyst that normally contains one or more than one metal of groups IVb, Vb, Vlb or VIII of the Periodic Table. These metals usually have one or more than one ligand, typically oxides, halides, alcoholates, esters, ethers, amines, alkyls, alkenyls and/or aryls that may be either p- or s-coordinated. These metal complexes

may be in the free form or fixed on substrates, typically on activated magnesium chloride, titanium(III) chloride, alumina or silicon oxide. These catalysts may be soluble or insoluble in the polymerization medium. The catalysts can be used by themselves in the polymerization or further activators may be used, typically metal alkyls, metal hydrides, metal alkyl halides, metal alkyl oxides or metal alkyloxanes, said metals being elements of groups Ia, IIa and/or IIIa of the Periodic Table. The activators may be modified conveniently with further ester, ether, amine or silyl ether groups. These catalyst systems are usually termed Phillips, Standard Oil Indiana, Ziegler (-Natta), TNZ (DuPont), metallocene or single site catalysts (SSC).

2. Mixtures of the polymers mentioned under 1.), for example mixtures of polypropylene with polyisobutylene, polypropylene with polyethylene (for example PP/HDPE, PP/LDPE) and mixtures of different types of polyethylene (for example LDPE/HDPE).

3. Copolymers of monoolefins and diolefins with each other or with other vinyl monomers, for example ethylene/propylene copolymers, linear low density polyethylene (LLDPE) and mixtures thereof with low density polyethylene (LDPE), propylene/but-1-ene copolymers, propylene/isobutylene copolymers, ethylene/but-1-ene copolymers, ethylene/hexene copolymers, ethylene/methylpentene copolymers, ethylene/heptene copolymers, ethylene/octene copolymers, propylene/butadiene copolymers, isobutylene/isoprene copolymers, ethylene/alkyl acrylate copolymers, ethylene/alkyl methacrylate copolymers, ethylene/vinyl acetate copolymers and their copolymers with carbon monoxide or ethylene/acrylic acid copolymers and their salts (ionomers) as well as terpolymers of ethylene with propylene and a diene such as hexadiene, dicyclopentadiene or ethylidene-norbornene; and mixtures of such copolymers with one another and with polymers mentioned in 1) above, for example polypropylene/ethylene-propylene copolymers, LDPE/ethylene-vinyl acetate copolymers (EVA), LDPE/ethylene-acrylic acid copolymers (EAA), LLDPE/EVA, LLDPE/EAA and alternating or random polyalkylene/carbon monoxide copolymers and mixtures thereof with other polymers, for example polyamides.

The effective flame retarding amount of component (B) is that needed to show flame retarding efficacy as measured by one of the standard methods used to assess flame retardancy. These include the NFPA 701 Standard Methods of Fire Tests for Flame-Resistant Textiles and Films, 1989 and 1996 editions; the UL 94 Test for Flammability of

Plastic Materials for Parts in Devices and Appliances, 5th Edition, October 29, 1996; Limiting Oxygen Index (LOI), ASTM D-2863; and Cone Calorimetry, ASTM E-1354.

In the synergistic mixture (B), component (i.) is present from about 0.25 to about 10 % by weight based on component (A); for example from about 0.5 to about 8 % by weight based on component (A); for example from about 0.5 to about 5 % by weight based on component (A); for example from about 0.5 to about 2 % by weight based on component (A); for example from 5.1 % to about 10 % by weight based on component (A); for example from 5.1 % to about 8 % based on component (A).

In the synergistic mixture (B), component (ii.) is present from about 0.25 to about 60 % by weight based on component (A); for example from about 0.25 to about 30 % by weight based on component (A); for example from about 0.5 % to about 15 % by weight based on component (A); for example from about 1 % to about 10 % by weight based on component (A); for example from about 1 % to about 5 % by weight based on component (A).

The flame retardant compounds of component (ii.) useful in the instant invention are for instance selected from the group consisting of

tetraphenyl resorcinol diphosphite (FYROLFLEX[®] RDP, Akzo Nobel),
chloroalkyl phosphate esters (ANTIBLAZE[®] AB-100, Albright & Wilson; FYROL[®] FR-2, Akzo Nobel)
polybrominated diphenyl oxide (DE-60F, Great Lakes Corp.)
decabromodiphenyl oxide (DBDPO; SAYTEX[®] 102E)
antimony trioxide (Sb₂O₃),
antimony pentoxide (Sb₂O₅),
tris[3-bromo-2,2-bis(bromomethyl)propyl] phosphate (PB 370[®], FMC Corp.),
triphenyl phosphate,
bis(2,3-dibromopropyl ether) of bisphenol A (PE68),
ammonium polyphosphate (APP) or (HOSTAFLAM[®] AP750),
resorcinol diphosphate oligomer (RDP),
brominated epoxy resin,
ethylene-bis(tetrabromophthalimide) (SAYTEX[®] BT-93),
bis(hexachlorocyclopentadieno)cyclooctane (DECLORANE PLUS[®]),

- 36 -

calcium sulfate
chlorinated paraffins,
magnesium carbonate,
melamine phosphates,
melamine pyrophosphates,
molybdenum trioxide,
zinc oxide,
1,2-bis(tribromophenoxy)ethane (FF680),
tetrabromo-bisphenol A (SAYTEX[®] RB100),
magnesium hydroxide,
alumina trihydrate,
zinc borate,
ethylenediamine diphosphate (EDAP)
silica,
silicones,
calcium silicate,
magnesium silicate and
ethylene bis-(dibromo-norbornanedicarboximide), (SAYTEX[®] BN-451).

Coadditives found useful for use with the instant compounds of components (a)-(j) in flame retardant compositions are as follows:

UV absorbers:

2-(2-hydroxy-3,5-di- α -cumylphenyl)-2H-benzotriazole, (TINUVIN[®] 234, Ciba Specialty Chemicals Corp.);

2-(2-hydroxy-5-methylphenyl)-2H-benzotriazole, (TINUVIN[®] P, Ciba Specialty Chemicals Corp.);

5-chloro-2-(2-hydroxy-3,5-di-tert-butylphenyl)-2H-benzotriazole, (TINUVIN[®] 327, Ciba Specialty Chemicals Corp.);

2-(2-hydroxy-3,5-di-tert-amylphenyl)-2H-benzotriazole, (TINUVIN[®] 328, Ciba Specialty Chemicals Corp.);

2-(2-hydroxy-3- α -cumyl-5-tert-octylphenyl)-2H-benzotriazole, (TINUVIN[®] 928, Ciba Specialty Chemicals Corp.);

2,4-di-tert-butylphenyl 3,5-di-tert-butyl-4-hydroxybenzoate, (TINUVIN[®] 120, Ciba Specialty Chemicals Corp.);

2-hydroxy-4-n-octyloxybenzophenone, (CHIMASSORB[®] 81, Ciba Specialty Chemicals Corp.);

2,4-bis(2,4-dimethylphenyl)-6-(2-hydroxy-4-octyloxyphenyl)-s-triazine, (CYASORB[®] 1164, Cytac).

It is contemplated that in addition to components (i.) and (ii.) of the present compositions, that hindered amines as disclosed in U.S. Pat. No. 6,117,995 and copending U.S. application Nos. 09/502,239, filed Nov. 3, 1999, and 09/714,717, filed Nov. 16, 2000, may also be present. The disclosures of this U.S. patent and applications are hereby incorporated by reference.

Certain compounds selected from components (a)-(j) are effective alone towards providing flame retardancy to plastic substrates, for example polyolefins, polystyrenics and PVC. In particular, compounds of component (d), certain substituted hydroxylamine stabilizers, are effective alone as flame retardants. In particular, O-n-propyl-N,N-dioctadecylhydroxylamine is an effective flame retardant alone, that is without the presence of known flame retardants such as brominated flame retardants, phosphorus containing flame retardants and inorganic flame retardants.

The following examples are meant for illustrative purposes only and are not to be construed to limit the scope of this invention in any manner whatsoever.

Test Methods

NFPA 701 Standard Methods of Fire Tests for Flame-Resistant Textiles and Films, 1989 and 1996 editions;

UL 94 Test for Flammability of Plastic Materials for Parts in Devices and Appliances, 5th Edition, October 29, 1996;

- 38 -

Limiting Oxygen Index (LOI), ASTM D-2863;

Cone Calorimetry, ASTM E-1 or ASTM E 1354;

ASTM D 2633-82, burn test.

Test Compounds

NO•1 is bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) sebacate, (Prostab™ 5415, Ciba Specialty Chemicals Corp.),

NOH-1 is bis(1-hydroxy-2,2,6,6-tetramethylpiperidin-4-yl) sebacate,

DBDPO is decabromodiphenyl oxide,

FR-1 is tris[3-bromo-2,2-bis(bromomethyl)propyl] phosphate, (PB 370®, FMC Corp.).

FR-2 is ammonium polyphosphate (APP).

FR-3 is bis(2,3-dibromopropyl) ether of tetrabromobisphenol A (PE68).

FR-4 is ammonium polyphosphate/synergist blend, HOSTAFLAM® AP750.

FR-5 is decabromodiphenyl oxide, SAYTEX® 102E.

FR-6 is ethylene bis-(tetrabromophthalimide), (SAYTEX® BT-93).

FR-7 is melamine phosphate, MELAPUR® P 46.

FR-8 is ammonium polyphosphate, EXOLIT® AP752.

Example 1

Molding grade polypropylene is dry blended with the test additives and then melt compounded at 425°F (218°C). Base stabilization is 500 ppm N,N-di(alkyl)hydroxylamine produced by the direct oxidation of N,N-di(hydrogenated tallow)amine (Irgastab® FS-042) and 500 ppm calcium stearate. Plaques (125 mil) are prepared by injection molding from the formulations using a Boy Injection Molder at 475°F (246°C). The specimens are tested for

flame retardancy according to the UL-94 vertical burn test specifications. The results are shown below.

The plaques are tested for flame retardancy by the UL 94V thick section test. The time in seconds for the plaques to extinguish after the insult flame is removed is reported as "After Flame". The time in seconds that the plaques glow after the flame incurred from the second insult flame is extinguished is reported as "Afterglow". Efficacy as a flame retardant is demonstrated when low After Flame times are observed relative to a blank sample containing no flame retardant. The average time for 5 samples is reported in the table below. Additive levels are reported in weight percent based on polymer.

Formulation	Additive	1 st After-flame	2 nd After-flame	Afterglow
1 (control)	none	>100	xxx	xxx
2	15 % DBDPO 3 % Sb ₂ O ₃	11.6	8.2	0.0
3	10 % DBDPO	30.8	22.1	0.0
4	5 % DBDPO	>100	xxx	xxx
5	5 % DBDPO 0.25 % NOH-1	14.4	48.7	0.0

The entry "xxx" means that the sample is completely consumed and could not be reignited.

DBDPO at the 5 % level does not provide adequate flame retardancy. However, when this level of DBDPO is augmented with only 0.25 % NOH-1, the flame retardancy is greatly improved.

These data show that decabromodiphenyl oxide can be replaced with a small amount of an instant hydroxylamine stabilizer in order to achieve good flame retardancy.

Example 2

Polyethylene fibers are prepared from fiber grade polyethylene by dry blending with test additives and melt compounding at 400°F. Fibers are extruded from this formulation using a Hills laboratory scale fiber extruder. Socks are knitted from the fibers and are tested for flame retardancy according to NFPA 701 vertical burn method. Polyethylene fibers contain an additive of present components (a)-(j) in combination with a classic brominated flame retardant decabromodiphenyl oxide (DBDPO); bis(2,3-dibromopropyl) ether of tetrabromobis phenol A (PE68); or ethylene bis-tetrabromophthalimide (SAYTEX® BT-93). These formulated fibers are tested for flame retardancy according to NFPA 701.

The fibers containing both an additive of components (a)-(j) of the present invention and a classic brominated flame retardant exhibit enhanced flame retardancy compared to the classic flame retardant alone.

Example 3

Polyethylene (LDPE) is melt compounded on a twin screw extruder at 450°F (232°C) with an additive of present components (a)-(j) in combination with a classic brominated flame retardant decabromodiphenyl oxide (DBDPO); bis(2,3-dibromopropyl) ether of tetrabromobis phenol A (PE68); or ethylene bis-tetrabromophthalimide (SAYTEX® BT-93). 125 mil plaques are compression molded at 400°F(204°C).

The plaques are tested for flame retardancy by the UL 94V thick section test.

The plaques containing both an additive of components (a)-(j) of the present invention and a classic brominated flame retardant exhibit enhanced flame retardancy compared to the classic flame retardant alone.

Example 4

Foam grade polyethylene is dry blended with test additives and then melt compounded into pellets. The pelletized fully formulated resin is then blown into foam.

The polyethylene foam prepared contains an instant additive of components (a)-(j) in combination with a classic brominated flame retardant. The formulated foam is tested for flame retardancy according to the UL-94 burn test method.

The foam containing both an additive of present components (a)-(j) and a classic brominated flame retardant exhibits enhanced flame retardancy compared to foam containing the classic halogenated flame retardant alone.

Example 5

Wire & cable grade polyethylene is dry blended with test additives and then melt compounded into pellets. The pelletized fully formulated resin is then extruded onto wire.

Test specimens are tested for flame retardancy using the ASTM D 2633-82 burn test conditions. The formulations containing both a compound of the present invention selected from components (a)-(j) and a classic brominated flame retardant exhibits enhanced flame retardancy compared to the classic halogenated flame retardant alone.

Example 6

Fiber grade polyethylene is dry-blended with test additives. In addition to a compound of present components (a)-(j), selected brominated flame retardants are also included in the various formulations. Non-woven fabrics are produced from the polymer blend formulations by a spun-bonded or melt-blown process.

The non-woven fabrics made thereby are tested for flame retardancy according to the NFPA 701 vertical burn test specifications. The fabrics containing the present compounds of components (a)-(j) and selected brominated flame retardants exhibit flame retardancy.

Example 7

Fiber grade polypropylene is dry-blended with test additives. In addition to a present compound of components (a)-(j), selected brominated flame retardants are also included in the various formulations. Non-woven fabrics are produced from the polymer blend formulations by a spun-bonded or melt-blown process.

The non-woven fabrics made thereby are tested for flame retardancy according to the NFPA 701 vertical burn test specifications. The fabrics containing an additive selected from present components (a)-(j) and selected brominated flame retardants exhibit flame retardancy.

Example 8

Molding grade polystyrene is dry-blended with test additives and then melt compounded. In addition to an additive selected from present components (a)-(j), selected brominated flame retardants are also included in the test formulations. Specimens are injection molded from these test formulations.

The specimens are tested for flame retardancy according to the UL-94 burn test specifications. The molded specimens containing the present compounds of components (a)-(j) and selected brominated flame retardants exhibit flame retardancy.

Example 9

Foam grade polystyrene is dry-blended with test additives and then melt compounded. In addition to an additive selected from present components (a)-(j), selected brominated

flame retardants are also included in these test formulations. Foam polystyrene specimens are prepared from these test formulations.

The specimens are tested for flame retardancy according to the UL-94 burn test specifications. The foam specimens containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Example 10

Molding grade ABS is dry blended with the an additive selected from present components (a)-(j) and selected brominated flame retardants, then melt compounded at 425°F (218°C). Specimens 125 mil (1/8") thick are then injection molded from this formulation using a Boy Injection Molder at 450°F (232°C). The specimens are tested for flame retardancy according to the UL-94 vertical burn test specifications.

The specimens containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Example 11

Fiber grade polypropylene is dry blended with an additive selected from the present components (a)-(j) and a select brominated flame retardant and then melt compounded at 234°C (450°F) into pellets. The pelletized fully formulated resin is then spun at 246°C (475°F) into fiber using a Hills laboratory model fiber extruder. The spun tow of 41 filaments is stretched at a ratio of 1:3.2 to give a final denier of 615/41.

Socks are knitted from the stabilized polypropylene fiber on a Lawson-Hemphill Analysis Knitter and tested under NFPA 701 vertical burn procedure. The time in seconds for the knitted sock to extinguish after the insult flame is removed is reported as "After Flame". Both the maximum time for any one replicate and the total time for ten replicates are measured. Efficacy as a flame retardant is demonstrated when low After Flame times are observed relative to a blank sample containing no flame retardant.

The specimens containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Example 12

Film grade polyethylene is dry blended with the with an additive selected from the present components (a)-(j) and a select brominated flame retardant and then melt compounded into pellets. The pelletized fully formulated resin is then blown at 205°C using a MPM Superior Blown film extruder.

The films are tested for flame retardancy under NFPA 701 test conditions. The specimens containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Film grade polypropylene is handled in a similar fashion and polypropylene films containing the instant compounds and brominated flame retardants also show flame retardancy.

Example 13

Molded test specimens are prepared by injection molding thermoplastic olefin (TPO) pellets containing a present test compound selected from components (a)-(j) and a select brominated flame retardant. The TPO formulations may also contain a pigment, a phosphite, a phenolic antioxidant or hydroxylamine, a metal stearate, a UV absorber or a hindered amine stabilizer or a mixture of hindered amine and UV absorber.

Pigmented TPO formulation composed of polypropylene blended with a rubber modifier where the rubber modifier is an in-situ reacted copolymer or blended product containing copolymers of propylene and ethylene with or without a ternary component such as ethylidene norbornene are stabilized with a base stabilization system consisting of an N,N-

dialkylhydroxylamine or a mixture of hindered phenolic antioxidant and an organophosphorus compound.

The TPO plaques are tested for flame retardancy using the UL-94 Vertical Burn conditions. A minimum of three replicates are tested. Efficacy as a flame retardant is measured relative to a blank sample containing no flame retardant.

The specimens containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Example 14

Film grade ethylene/vinyl acetate (EVA) copolymers containing 20 weight percent or less of vinyl acetate are dry blended with test additives and then melt compounded into pellets. The pelletized fully formulated resin is then blown into a film at 205°C using a MPM Superior Blown-film extruder.

The films are tested for flame retardancy under NFPA 701 test conditions. The films containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Film grade low density polyethylene (LDPE) which contains some linear low density polyethylene (LLDPE) and/or ethylene/vinyl acetate (EVA) are dry blended with test additives and blown into film as described above for EVA copolymer resin. The films are tested for flame retardancy under NFPA 701 test conditions and those containing the present compounds selected from components (a)-(j) and brominated flame retardants exhibit flame retardancy.

Example 15

High impact polystyrene (HIPS) polymer (STYRON[®] 484C, Dow Chemical Co.) is compounded with a present compound of components (a)-(j) and a select brominated flame

retardant, pelletized and then injection or compression molded into plaques. These plaques are tested for flame retardant efficacy using cone calorimetry, LOI or UL-94 test method.

The plaques containing an instant compound of component (a)-(j) and a select brominated flame retardant exhibit flame retardancy. Flame retardant HIPS polymers find application in housings for business machines.

Example 16

This Example shows the efficacy of the present compounds in PVC formulations. Such formulations are useful in flexible or rigid PVC and in wire and cable applications.

Typical formulations are seen below:

<u>Component</u>	<u>parts</u>	<u>parts</u>	<u>parts</u>	<u>parts</u>
PVC resin	100	100	100	100
tin mercaptide	1.5	--	2.0	--
tin carboxyate	--	2.5	--	2.0
process aid	1.5	1.5	2.0	2.0
impact mod.	6.0	6.0	7.0	7.0
paraffin wax	1.0	0.3	1.0	1.0
polyethyl wax	0.1	0.1	0.2	0.2
Ca stearate	1.0	--	0.8	--
pigment	1.0	0.9	5.0	5.0

Fully formulated PVC containing the present compounds of components (a)-(j) and a known organic or inorganic flame retardant is pelletized and then injection molded into test plaques for examination of flame retardancy using the UL-94 or LOI test method.

The PVC plaques containing the instant compounds of components (a)-(j) and a classic organic or inorganic flame retardant demonstrate flame retardancy.

Example 17

Fiber grade poly(ethylene terephthalate) (PET) is dry blended with test additives of components (a)-(j) and a classic flame retardant, then melt compounded at 550°F and then pelletized. The polyester pellets are dried at 175°F for 24 hours under vacuum. The dried pellets are extruded into fibers using a Hills laboratory scale fiber extruder at 550°F. Socks are knitted from these fibers and tested for flame retardancy according to NFPA 701 test method.

The fibers containing both a present compound of components (a)-(j) and a classic halogenated or phosphorus flame retardant exhibit enhanced flame retardancy compared to the classic flame retardant alone.

Example 18

Thermoplastic resins including polypropylene, polyethylene homopolymer, polyolefin copolymer or thermoplastic olefins (TPO), high impact polystyrene (HIPS) and ABS are dry blended with the instant compounds of components (a)-(j) and a classic organic or inorganic flame retardant and then melt compounded into pellets. The pelletized fully formulated resin is then processed into a useful article such as extrusion into fiber; blown or cast extrusion into film; blow molded into bottles; injection molded into molded articles, thermoformed into molded articles, extruded into wire and cable housing or rotation molded into hollow articles.

The articles containing the instant compounds of components (a)-(j) and a known organic or inorganic flame retardant exhibit flame retardancy when tested by a known standard test method.

Polyethylene wire and cable applications are tested for flame retardancy according to ASTM D-2633-82 burn test method. The materials containing the instant compounds of components (a)-(j) and a known organic or inorganic flame retardant show flame retardancy.

Example 19

Articles prepared according to Example 18 which additionally contain an organophosphorus stabilizer selected from the group consisting of tris(2,4-di-tert-butylphenyl) phosphite, bis(2,4-di-tert-butyl-6-methylphenyl) ethyl phosphite, 2,2',2"-nitrido[triethyl-tris-(3,3',5,5'-tetra-tert-butyl-1,1'-biphenyl-2,2'-diyl) phosphite], tetrakis(2,4-di-butylphenyl) 4,4'-biphenylenediphosphonite, tris(nonylphenyl) phosphite, bis(2,4-di-tert-butylphenyl) pentaerythryl diphosphite, 2,2'-ethylidenebis(2,4-di-tert-butylphenyl) fluorophosphite and 2-butyl-2-ethylpropan-1,3-diyl 2,4,6-tri-tert-butylphenyl phosphite exhibit good flame retardancy properties.

Example 20

Articles prepared according to Example 18 which additionally contain a o-hydroxyphenyl-2H-benzotriazole, a hydroxyphenyl benzophenone or a o-hydroxyphenyl-s-triazine UV absorber selected from the group consisting of 2-(2-hydroxy-3,5-di- α -cumylphenyl)-2H-benzotriazole, 2-(2-hydroxy-5-methylphenyl)-2H-benzotriazole, 5-chloro-2-(2-hydroxy-3,5-di-tert-butylphenyl)-2H-benzotriazole, 2-(2-hydroxy-3,5-di-tert-amylphenyl)-2H-benzotriazole, 2-(2-hydroxy-3- α -cumyl-5-tert-octylphenyl)-2H-benzotriazole, 2,4-di-tert-butylphenyl 3,5-di-tert-butyl-4-hydroxybenzoate, 2-hydroxy-4-n-octyloxybenzophenone and 2,4-bis(2,4-dimethylphenyl)-6-(2-hydroxy-4-octyloxyphenyl)-s-triazine exhibit good flame retardancy.

Example 21

Articles prepared according to Example 18 which additionally contain a o-hydroxyphenyl-2H-benzotriazole, a hydroxyphenyl benzophenone or a o-hydroxyphenyl-s-triazine UV absorber selected from the group consisting of 2-(2-hydroxy-3,5-di- α -cumylphenyl)-2H-benzotriazole, 2-(2-hydroxy-5-methylphenyl)-2H-benzotriazole, 5-chloro-2-(2-hydroxy-3,5-di-tert-butylphenyl)-2H-benzotriazole, 2-(2-hydroxy-3,5-di-tert-amylphenyl)-2H-benzotriazole, 2-(2-hydroxy-3- α -cumyl-5-tert-octylphenyl)-2H-benzotriazole, 2,4-di-tert-butylphenyl 3,5-di-tert-butyl-4-hydroxybenzoate, 2-hydroxy-4-n-octyloxybenzophenone and 2,4-bis(2,4-dimethylphenyl)-6-(2-hydroxy-4-octyloxyphenyl)-s-triazine exhibit good flame retardancy properties.

Example 22

Examples 1-21 are repeated where the compound of components (a)-(j) is selected from

- (a) bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) sebacate, 4-hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine,
 - (b) an N,N-di(alkyl)hydroxylamine produced by the direct oxidation of N,N-di(hydrogenated tallow)amine (Irgastab[®] FS-042),
 - (c) N-octadecyl- α -heptadecylnitrone,
 - (d) O-n-propyl-N,N-dioctadecylhydroxylamine,
 - (e) Genox[™] EP, a di(C₁₆-C₁₈)alkyl methyl amine oxide, CAS# 204933-93-7,
 - (f) 3-(3,4-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one, Irganox[®] HP-136, (BF1),
 - (g) tris(2,4-di-tert-butylphenyl) phosphite (Irgafos[®] 168),
 - (h) 2,6-di-tert-butyl-4-benzylidene-cyclohexa-2,5-dienone (Prostab[™] 6007)
- and
- (j) the monoacrylate ester of 2,2'-methylenebis(6-tert-butyl-4-methylphenol), Irganox[®] 3052.

Formulations containing an instant stabilizer of components (a)-(j) and a brominated flame retardant achieve good flame retardancy.

Example 23

Examples 1-22 are repeated where the classic flame retardant is a flame retardant such as bis(2,3-dibromopropyl) ether of bisphenol A (PE68), tris[3-bromo-2,2-bis(bromomethyl)propyl] phosphate (PB370), ethylene bis-tetrabromophthalimide (SAYTEX BT-93) or ethylene bis-dibromonorbornanedicarboximide (SAYTEX BN-451). Other formulations may contain antimony trioxide in addition to brominated flame retardants. Other formulations may contain phosphorus based flame retardants such as ethylenediamine diphosphate (EDAP) or ammonium polyphosphate (APP) instead of a brominated flame retardant.

The samples containing the present compounds selected from components (a)-(j) and classic flame retardants exhibit flame retardancy.

Example 24

Example 1 is repeated with the test additives of the table below. Additive levels are reported in weight percent based on polymer. The average of 2 or 3 samples are reported.

Formulation	Additive	1 st After-flame	2 nd After-flame
1 (control)	none	burns	xxx
6	1% NO•1	0	burns
7	1% NOH-1	1.2	burns
8	1% NOR-1	burns	xxx
9	3% FR-1	0	burns
10	4% FR-1	0	burns
11	5% DBDPO	2.2	burns
12	6% DBDPO	2.1	burns
13	3% FR-1 1% NOR-1	0	50.5
14	5% DBDPO 1% NO•1	0.2	6.8
15	5% DBDPO 1% NOH-1	0.4	19.5

The entry "xxx" means that the sample is completely consumed and could not be reignited. "Burns" means the plaque burned entirely to the clamp. NOR-1 is $(\text{H}_{37}\text{C}_{18})_2\text{NOCH}_2\text{CH}_2\text{CH}_3$. It is clear that combinations of nitroxyl stabilizers, hydroxylamine stabilizers or substituted hydroxylamine stabilizers with conventional flame retardants are synergistic towards providing flame retardancy to plastic substrates.

WHAT IS CLAIMED IS:

1. A flame retardant composition which comprises

(A) a polymer substrate, and

(B) an effective flame retarding amount of a synergistic mixture of


(i.) at least one compound selected from the group consisting of

- (a) nitroxyl stabilizers,
- (b) hydroxylamine stabilizers,
- (c) nitron stabilizers,
- (d) substituted hydroxylamine stabilizers,
- (e) amine oxide stabilizers,
- (f) benzofuranone stabilizers,
- (g) phosphite and phosphonite stabilizers,
- (h) quinone methide stabilizers and
- (j) monoacrylate esters of 2,2'-alkylidenebisphenol stabilizers and


(ii.) at least one compound selected from the group consisting of brominated flame retardants, phosphorus containing flame retardants and inorganic flame retardants.

2. A composition according to claim 1 wherein component (i.) of the synergistic mixture (B) is selected from

a) nitroxyl stabilizers of the formula




or nitroxyl stabilizers that contain one or more groups of the formula



where each R is alkyl of 1 to 4 carbon atoms and T is a group required to complete a 5- or 6-membered ring;

b) hydroxylamine stabilizers of the formula (I)




wherein

T₁ is straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms; and

T₂ is hydrogen, or independently has the same meaning as T₁;

or hydroxylamine stabilizers that contain one or more of the groups of the formula (II)




wherein

T is a group forming a five- or six-membered ring; and

R₁ and R₂ are independently hydrogen, alkyl of 1 to 4 carbon atoms or phenyl;

c) nitrene stabilizers of the formula (III)




wherein

L₁ is straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms;

L₂ and L₃ are independently hydrogen, straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms;

or L₁ and L₂ together form a five- or six-membered ring including the nitrogen atom;

d) hydroxylamines of the formula (IIIb) or (IIIa)



wherein


T₁ is straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms, or said aralkyl substituted by one or two alkyl of 1 to 12 carbon atoms or by one or two halogen atoms;

T₂ is hydrogen, or independently has the same meaning as T₁; and

T₃ is allyl, straight or branched chain alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 18 carbon atoms, cycloalkenyl of 5 to 18 carbon atoms or a straight or branched chain alkyl of 1 to 4 carbon atoms substituted by phenyl or by phenyl substituted by one or two alkyl groups of 1 to 4 carbon atoms or by 1 or 2 halogen atoms;

e) amine oxide stabilizers of formula (VII)

- 54 -



wherein

G_1 and G_2 are independently a straight or branched chain alkyl of 6 to 36 carbon atoms, aryl of 6 to 12 carbon atoms, aralkyl of 7 to 36 carbon atoms, alkaryl of 7 to 36 carbon atoms, cycloalkyl of 5 to 36 carbon atoms, alkylcycloalkyl of 6 to 36 carbon atoms or cycloalkylalkyl of 6 to 36 carbon atoms;

G_3 is a straight or branched chain alkyl of 1 to 36 carbon atoms, aryl of 6 to 12 carbon atoms, aralkyl of 7 to 36 carbon atoms, alkaryl of 7 to 36 carbon atoms, cycloalkyl of 5 to 36 carbon atoms, alkylcycloalkyl of 6 to 36 carbon atoms or cycloalkylalkyl of 6 to 36 carbon atoms; and

wherein said aryl groups may be substituted by one to three halogen, alkyl of 1 to 8 carbon atoms, alkoxy of 1 to 8 carbon atoms or combinations thereof; and

wherein said alkyl, aralkyl, alkaryl, cycloalkyl, alkylcycloalkyl and cycloalkylalkyl groups may be interrupted by -O-, -S-, -SO-, -SO₂-, -COO-, -OCO-, -CO-, -NG₄-, -CONG₄- and -NG₄CO- groups, or wherein said alkyl, aralkyl, alkaryl, cycloalkyl, alkylcycloalkyl and cycloalkylalkyl groups may be substituted by -OG₄-, -SG₄-, -COOG₄-, -OCOG₄-, -COG₄-, -N(G₄)₂-, -CON(G₄)₂-, -NG₄COG₄ and 5- and 6-membered rings containing the -C(CH₃)(CH₂R_x)NL(CH₂R_x)(CH₃)C- group or wherein said alkyl, aralkyl, alkaryl, cycloalkyl, alkylcycloalkyl and cycloalkylalkyl groups are both interrupted and substituted by the groups mentioned above; and

wherein

G_4 is independently hydrogen or alkyl of 1 to 8 carbon atoms;




R_x is hydrogen or methyl;

L is hydrogen, hydroxy, C₁₋₃₀ straight or branched chain alkyl moiety, a -C(O)R moiety where R is a C₁₋₃₀ straight or branched chain alkyl group, or a -OR_y moiety; and

R_y is C₁₋₃₀ straight or branched chain alkyl, C₂-C₃₀ alkenyl, C₂-C₃₀ alkynyl, C₅-C₁₂ cycloalkyl, C₆-C₁₀ bicycloalkyl, C₅-C₈ cycloalkenyl, C₆-C₁₀ aryl, C₇-C₉ aralkyl, C₇-C₉ alkaryl substituted by alkyl or aryl, or -CO(D), where D is C₁-C₁₈ alkyl, C₁-C₁₈ alkoxy, phenyl, phenyl substituted by hydroxy, alkyl or alkoxy, or amino or amino mono- or di-substituted by alkyl or phenyl;

f) benzofuranone stabilizers;


g) phosphite and/or phosphonite stabilizers of the formulae (1), (2), (3), (4), (5), (6) and (7)



in which the indices are integral and


n is 2, 3 or 4; p is 1 or 2; q is 2 or 3; r is 4 to 12; y is 1, 2 or 3; and z is 1 to 6;

A₁, if n is 2, is C₂-C₁₈ alkylene; C₂-C₁₂ alkylene interrupted by oxygen, sulfur or -NR₄;



phenylene;

A₁, if n is 3, is a radical of the formula -C_rH_{2r-1}-;



A₂ is as defined for A₁ if n is 2;

B is a direct bond, -CH₂-, -CHR₄-, -CR₁₁R₄-, sulfur, C₅-C₇ cycloalkylidene, or cyclohexylidene which is substituted by from 1 to 4 C₁-C₄ alkyl radicals in position 3, 4 and/or 5;

D₁, if p is 1, is C₁-C₄ alkyl and, if p is 2, is -CH₂OCH₂-;

D₂, if p is 1, is C₁-C₄ alkyl;

E, if y is 1, is C₁-C₁₈ alkyl, -OR₁₁ or halogen;

E, if y is 2, is -O-A₂-O-,

E, if y is 3, is a radical of the formula R₄C(CH₂O-)₃ or N(CH₂CH₂O-)₃;

Q is the radical of an at least z-valent alcohol or phenol, this radical being attached via the oxygen atom to the phosphorus atom;

R₁₁, R₁₂ and R₃ independently of one another are C₁-C₁₈ alkyl which is unsubstituted or substituted by halogen, -COOR₄, -CN or -CONR₄R₄; C₂-C₁₈ alkyl interrupted by oxygen, sulfur or -NR₄-; C₇-C₉ phenylalkyl; C₅-C₁₂ cycloalkyl, phenyl or naphthyl; naphthyl or phenyl substituted by halogen, 1 to 3 alkyl radicals or alkoxy radicals having a total of 1 to 18 carbon

atoms or by C₇-C₉ phenylalkyl; or a radical of the formula
$$\text{---}(\text{CH}_2)_m\text{---}\begin{array}{c} \text{R}_5 \\ | \\ \text{---} \text{C}_6\text{H}_3 \text{---} \\ | \\ \text{R}_6 \end{array}\text{---OH}$$
 in which

m is an integer from the range 3 to 6;

R₄ is hydrogen, C₁-C₁₈ alkyl, C₅-C₁₂ cycloalkyl or C₇-C₉ phenylalkyl,

R₅ and R₆ independently of one another are hydrogen, C₁-C₈ alkyl or C₅-C₆ cycloalkyl,

R₇ and R₈, if q is 2, independently of one another are C₁-C₄ alkyl or together are a 2,3-dehydropentamethylene radical; and

R₇ and R₈, if q is 3, are methyl;

R₁₄ is hydrogen, C₁-C₉ alkyl or cyclohexyl,


R₁₅ is hydrogen or methyl and, if two or more radicals R₁₄ and R₁₅ are present, these radicals are identical or different,

X and Y are each a direct bond or oxygen,

Z is a direct bond, methylene, -C(R₁₆)₂- or sulfur, and

R₁₆ is C₁-C₈ alkyl;

h) quinone methide stabilizers of formula (VIII)



wherein

R₁' and R₂' are independently straight or branched chain alkyl of 4 to 18 carbon atoms, cycloalkyl of 5 to 12 carbon atoms or phenylalkyl of 7 to 15 carbon atoms, and

R₃' is 2-, 3- or 4-pyridyl; 2- or 3-thienyl; 2- or 3-pyrrolyl; 2- or 3-furyl; aryl of 6 to 10 carbon atoms; or said aryl substituted by one to three alkyl of 1 to 8 carbon atoms, alkoxy of 1 to 8 carbon atoms, alkylthio of 1 to 8 carbon atoms, alkylamino of 1 to 8 carbon atoms, dialkylamino of 2 to 8 carbon atoms, alkoxy carbonyl of 2 to 8 carbon atoms, hydroxy, nitro, amino, cyano, carboxy, aminocarbonyl, chloro or mixtures of said substituents; or R₃' is -CN, -COOH, -COOR₄', -COR₅', -OCOR₆', -CONR₇'R₈' or -PO(OR₉')₂ where

R₄' is alkyl of 1 to 18 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, phenyl or benzyl,

R₅' is alkyl of 1 to 18 carbon atoms, aryl of 6 to 10 carbon atoms or said aryl substituted by 1 or 2 alkyl of 1 to 4 carbon atoms or by hydroxyl,


R₆' is alkyl of 1 to 18 carbon atoms, aryl of 6 to 10 carbon atoms or said aryl substituted by 1 or 2 alkyl of 1 to 4 carbon atoms or by hydroxyl,

R₇' and R₈' are independently hydrogen, alkyl of 1 to 18 carbon atoms or said alkyl substituted by alkylamino of 1 to 4 carbon atoms, by dialkylamino of 2 to 8 carbon atoms or by hydroxyl; benzyl, aryl of 6 to 10 carbon atoms or said aryl substituted by alkyl of 1 to 4 carbon atoms, by alkylamino of 1 to 4 carbon atoms, by dialkylamino of 2 to 8 carbon atoms, by phenylamino or by hydroxyl, or

-NR₇'R₈' is morpholino, piperidino or pyrrolidino, and

R₉' is hydrogen or alkyl of 1 to 18 carbon atoms;

j) monoacrylate esters of the formula (IX)



wherein

R_{40} , R_{41} , R_{42} , R_{43} and R_{44} are independently straight or branched chain alkyl of 1 to 18 carbon atoms, cycloalkyl of 5 to 12 carbon atoms or phenylalkyl of 7 to 15 carbon atoms.




3. A composition according to claim 2 wherein component (i.) of the synergistic mixture (B) is selected from

a) nitroxyl stabilizers selected from the group consisting of bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) sebacate, 4-hydroxy-1-oxy-2,2,6,6-tetramethylpiperidine, 4-ethoxy-1-oxy-2,2,6,6-tetramethylpiperidine, 4-propoxy-1-oxy-2,2,6,6-tetramethylpiperidine, 4-acetamido-1-oxy-2,2,6,6-tetramethylpiperidine, 1-oxy-2,2,6,6-tetramethylpiperidine, 1-oxy-2,2,6,6-tetramethylpiperidin-4-one, 1-oxy-2,2,6,6-tetramethylpiperidin-4-yl acetate, 1-oxy-2,2,6,6-tetramethylpiperidin-4-yl 2-ethylhexanoate, 1-oxy-2,2,6,6-tetramethylpiperidin-4-yl stearate, 1-oxy-2,2,6,6-tetramethylpiperidin-4-yl benzoate, 1-oxy-2,2,6,6-tetramethylpiperidin-4-yl 4-t-butyl-benzoate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) succinate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) adipate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) n-butylmalonate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) phthalate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) isophthalate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) terephthalate, bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl) hexahydroterephthalate, N,N'-bis(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl)adipamide, N-(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl)caprolactam, N-(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl)dodecylsuccinimide, 2,4,6-tris-[N-butyl-N-(1-oxy-2,2,6,6-tetramethylpiperidin-4-yl)]-s-triazine, 4,4'-ethylenebis(1-oxy-2,2,6,6-tetramethylpiperazin-3-one), 2-oxy-1,1,3,3-tetramethyl-2-isobenzazole, 1-oxy-2,2,5,5-tetramethylpyrrolidine, and N,N-bis-(1,1,3,3-tetramethylbutyl)nitroxide;

b) hydroxylamine stabilizers of the formula (I), wherein T_1 and T_2 are independently benzyl, methyl, ethyl, octyl, lauryl, dodecyl, tetradecyl, hexadecyl, heptadecyl or octadecyl, or wherein T_1 and T_2 are each the alkyl mixture found in hydrogenated tallow amine;

c) nitrones selected from the group consisting of N-benzyl- α -phenylnitronone, N-ethyl- α -methylnitronone, N-octyl- α -heptylnitronone, N-lauryl- α -undecylnitronone, N-tetradecyl- α -tridcylnitronone, N-hexadecyl- α -pentadecylnitronone, N-octadecyl- α -heptadecylnitronone, N-hexadecyl- α -heptadecylnitronone, N-ocatadecyl- α -pentadecylnitronone, N-heptadecyl- α -heptadecylnitronone, N-octadecyl- α -hexadecylnitronone, N-methyl- α -heptadecylnitronone and the nitronone derived from N,N-di(hydrogenated tallow)hydroxylamine;

d) hydroxylamines selected from the group consisting of compounds of the formula (IV), (V) and (VI)



wherein


in the compounds of formula (IV)

a is 1 to 4;

R' and R'' are independently hydrogen, alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms or said aralkyl substituted by alkyl of 1 to 36 carbon atoms,

R_b and R_c are independently hydrogen or alkyl of 1 to 12 carbon atoms;

A, when a = 1, is a group NHR_d, wherein R_d is hydrogen, amino, alkyl of 1 to 18 carbon atoms or




where B is a direct bond or alkylene of 1 to 10 carbon atoms;

A, when a = 2, is a bivalent radical of a 5-7 membered heterocyclic compound containing two nitrogen atoms in the ring, with the free valencies on the nitrogen atoms, or is a group $-\text{N}(\text{R}_d)-\text{B}-\text{N}(\text{R}_d)-$


where B and R_d have the meanings given above;

A, when a = 3, is a group of the formula



where n is 2 to 6 and R_e is hydrogen or alkyl of 1 to 4 carbon atoms; and

A, when a = 4, is a group of the formula




in compounds of formula (V),

R_f , R_g , R_h and R_i independently are hydrogen, alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms or said aralkyl substituted by alkyl of 1 to 36 carbon atoms; and the alkali metal and alkaline earth metal salts thereof;

in compounds of formula (VI),

R_j and R_k are independently alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, aralkyl of 7 to 9 carbon atoms or said aralkyl substituted by alkyl of 1 to 36 carbon atoms, or R_j and R_k together with the nitrogen can form a 5-7 member heterocyclic ring, or R_j and R_k are independently a group of the formula



R_l , R_m , R_n , R_o and R_p are independently hydrogen, alkyl of 1 to 36 carbon atoms, cycloalkyl of 5 to 12 carbon atoms, allyl, aryl, aralkyl of 7 to 9 carbon atoms, said aralkyl substituted by alkyl of 1 to 36 carbon atoms, bornyl, norbornyl or isobornyl, or R_m and R_n together with nitrogen can form a 5-7 member heterocyclic ring;

b is 1-4;

x is 0-5;

T_a is R_l or $-\text{CH}(R_l)-\text{O}-\text{N}(R_j)(R_k)$;

when $b = 1$, T_b is R_l or 5-7 member unsaturated heterocyclic-substituted (C_1 - C_4)alkyl, or T_a and T_b together with the nitrogen can form a 5-7 member saturated heterocyclic ring;

when $b = 2$, T_b is alkylene of 2 to 12 carbon atoms, cycloalkylene of 6 to 10 carbon atoms, arylene of 6 to 10 carbon atoms or alkylenearylenealkylene of 8 to 10 carbon atoms or T_a and T_b together with the two nitrogens can form a 5-7 member saturated heterocyclic ring;



when $b = 3$, T_b is alkanetriyl of 3 to 6 carbon atoms or T_a and T_b together with the three nitrogens can form a 5-6 member saturated heterocyclic ring; and

when $b = 4$, T_b is alkanetetriyl of 4 to 6 carbon atoms;

e) amine oxide stabilizers of formula (VII) wherein G_1 and G_2 are independently alkyl groups of 8 to 26 carbon atoms and G_3 is methyl;

f) benzofuranone stabilizers selected from the group consisting of 3-(4-(2-acetoxyethoxy)phenyl)-5,7-di-tert-butyl-benzofuran-2-one, 5,7-di-tert-butyl-3-(4-(2-stearoyloxyethoxy)phenyl)benzofuran-2-one, 3,3'-bis(5,7-di-tert-butyl-3-(4-(2-hydroxyethoxy)phenyl)benzofuran-2-one), 5,7-di-tert-butyl-3-(4-ethoxyphenyl)benzofuran-2-one, 3-(4-acetoxy-3,5-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one, 3-(3,5-dimethyl-4-pivaloyloxyphenyl)-5,7-di-tert-butyl-benzofuran-2-one, 3-(3,4-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one and 3-(2,3-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one;

g) phosphites and phosphonites selected from the group consisting of triphenyl phosphite, tris(nonylphenyl) phosphite, trilauryl phosphite, trioctadecyl phosphite, distearyl pentaerythritol diphosphite, tris(2,4-di-tert-butylphenyl) phosphite, diisodecyl pentaerythritol diphosphite, bis(2,4-di-tert-butylphenyl) pentaerythritol diphosphite, bis(2,6-di-tert-butyl-4-methylphenyl) pentaerythritol diphosphite, bisisodecylpentaerythritol diphosphite, bis(2,4-di-tert-butyl-6-methylphenyl) pentaerythritol diphosphite, bis(2,4,6-tri-tert-butylphenyl) pentaerythritol diphosphite, tristearyl sorbitol triphosphite, tetrakis (2,4-di-tert-butylphenyl) 4,4'-biphenylene-diphosphonite, 6-isoctyloxy-2,4,8,10-tetra-tert-butyl-dibenzo[d,f][1,3,2]dioxaphosphepin, 6-fluoro-2,4,8,10-tetra-tert-butyl-12-methyl-dibenzo[d,g][1,3,2]dioxaphosphocin, bis(2,4-di-tert-butyl-6-methylphenyl) methyl phosphite, bis(2,4-di-tert-butyl-6-methylphenyl) ethyl phosphite and 2,2',2''-nitrido[triethyltris(3,3',5,5'-tetra-tert-butyl-1,1'-biphenyl-2,2'-diyl)phosphite],



h) quinone methide stabilizers selected from the group consisting of (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetonitrile, (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetic acid, (3,5-di-tert-amyl-4-oxocyclohexa-2,5-dienylidene)acetic acid, methyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetate, ethyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetate, n-butyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetate, 2,6-di-tert-butyl-4-(2-oxopropylidene)-cyclohexa-2,5-dienone, diethyl (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methanephosphonate, (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methyl acetate, (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methyl pivalate, (3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)methyl benzoate, and N,N-diethyl-2-(3,5-di-tert-butyl-4-oxocyclohexa-2,5-dienylidene)acetamide, 2,6-di-tert-butyl-4-benzylidene-cyclohexa-2,5-dienone, 2,6-di-tert-butyl-4-(4-nitrobenzylidene)-cyclohexa-2,5-dienone, 2,6-di-tert-butyl-4-(3-nitrobenzylidene)-cyclohexa-2,5-dienone, 2,6-di-tert-butyl-4-(4-cyanobenzylidene)-cyclohexa-2,5-dienone, 2,6-di-tert-butyl-4-(4-dimethylaminobenzylidene)-cyclohexa-2,5-dienone, 2,6-di-tert-amyl-4-benzylidene-cyclohexa-2,5-dienone, 2,6-di-tert-butyl-4-(4-methoxybenzylidene)-cyclohexa-2,5-dienone, and 2,6-di-tert-butyl-4-(3,5-di-tert-butyl-4-hydroxybenzylidene)-cyclohexa-2,5-dienone;


j) monoacrylate esters selected from the group consisting of 2,2'-methylenebis(6-tert-butyl-4-methylphenol), 2,2'-methylenebis(6-tert-butyl-4-ethylphenol), 2,2'-methylenebis[4-methyl-6-(α -methylcyclohexyl)phenol], 2,2'-methylenebis(4-methyl-6-cyclohexylphenol), 2,2'-methylenebis(6-nonyl-4-methylphenol), 2,2'-methylenebis(4,6-di-tert-butylphenol), 2,2'-ethylidenebis(4,6-di-tert-butylphenol), 2,2'-ethylidenebis(6-tert-butyl-4-isobutylphenol), 2,2'-methylenebis[6-(α -methylbenzyl)-4-nonylphenol], 2,2'-methylenebis[6-(α,α -dimethylbenzyl)-4-nonylphenol], 2,6-bis(3-tert-butyl-5-methyl-2-hydroxybenzyl)-4-methylphenol and 1,1-bis-(3,5-dimethyl-2-hydroxyphenyl)butane.

4. A composition according to claim 3 wherein component (i.) of the synergistic mixture (B) is selected from


a) nitroxyl stabilizers;

b) N,N-dihydrocarbylhydroxylamines selected from the group consisting of N,N-dibenzylhydroxylamine, N,N-diethylhydroxylamine, N,N-dioctylhydroxylamine, N,N-dilaurylhydroxylamine, N,N-didodecylhydroxylamine, N,N-ditetradecylhydroxylamine, N,N-dihexadecylhydroxylamine, N,N-dioctadecylhydroxylamine, N-hexadecyl-N-tetradecylhydroxylamine, N-hexadecyl-N-heptadecylhydroxylamine, N-hexadecyl-N-octadecylhydroxylamine, N-heptadecyl-N-octadecylhydroxylamine, N-methyl-N-octadecylhydroxylamine and N,N-di(hydrogenated tallow)hydroxylamine;

f) benzofuranone stabilizers selected from the group consisting of



(BF1) and



(BF2).

5. A composition according to claim 1 in which component (ii.) of the synergistic mixture (B) is at least one compound selected from the group consisting of

- tetraphenyl resorcinol diphosphite,
- chloroalkyl phosphate esters,
- polybrominated diphenyl oxide,
- decabromodiphenyl oxide,

- 65 -

antimony trioxide,
antimony pentoxide,
tris[3-bromo-2,2-bis(bromomethyl)propyl] phosphate,
triphenyl phosphate,
bis(2,3-dibromopropyl ether) of bisphenol A,
ammonium polyphosphate,
resorcinol diphosphate oligomer,
brominated epoxy resin,
ethylene-bis(tetrabromophthalimide),
bis(hexachlorocyclopentadieno)cyclooctane,
calcium sulfate,
chlorinated paraffins,
magnesium carbonate,
melamine phosphates,
melamine pyrophosphates,
molybdenum trioxide,
zinc oxide,
1,2-bis(tribromophenoxy)ethane,
tetrabromo-bisphenol A,
magnesium hydroxide,
alumina trihydrate,
zinc borate,
ethylenediamine diphosphate,
silica,
silicones,
calcium silicate,
magnesium silicate and
ethylene bis-(dibromo-norbornanedicarboximide).

6. A composition according to claim 1 in which component (i.) of the synergistic mixture (B) is at least one compound selected from the group consisting of

(a) bis(1-oxyl-2,2,6,6-tetramethylpiperidin-4-yl) sebacate, 4-hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine,

- 66 -

- (b) an N,N-di(alkyl)hydroxylamine produced by the direct oxidation of N,N-di(hydrogenated tallow)amine,
- (c) N-octadecyl- α -heptadecylnitrone,
- (d) O-n-propyl-N,N-dioctadecylhydroxylamine,
- (e) di(C₁₆-C₁₈)alkyl methyl amine oxide,
- (f) 3-(3,4-dimethylphenyl)-5,7-di-tert-butyl-benzofuran-2-one,
- (g) tris(2,4-di-tert-butylphenyl) phosphite,
- (h) 2,6-di-tert-butyl-4-benzylidene-cyclohexa-2,5-dienone and
- (j) monoacrylate ester of 2,2'-methylenebis(6-tert-butyl-4-methylphenol) and

component (ii.) of the synergistic mixture (B) is at least one compound selected from the group consisting of

- polybrominated diphenyl oxide,
- decabromodiphenyl oxide,
- bis(2,3-dibromopropyl ether) of bisphenol A,
- ammonium polyphosphate,
- resorcinol diphosphate oligomer,
- brominated epoxy resin,
- ethylene-bis(tetrabromophthalimide),
- 1,2-bis(tribromophenoxy)ethane and
- tetrabromo-bisphenol A.

7. A composition according to claim 1 or 6 in which component (ii.) of the synergistic mixture (B) is the flame retardant ammonium polyphosphate or decabromodiphenyl oxide.

8. A composition according to claim 1 wherein of the synergistic mixture (B), component (i.) is present from 0.25 to 10 % by weight, based on component (A); and component (ii.) is present from 0.25 to 60 % by weight, based on component (A).

9. A composition according to claim 1 further comprising a UV absorber.

10. A composition according to claim 9 comprising a UV absorber selected from the group consisting of 2-(2-hydroxy-3,5-di- α -cumylphenyl)-2H-benzotriazole, 2-(2-hydroxy-5-methylphenyl)-2H-benzotriazole, 5-chloro-2-(2-hydroxy-3,5-di-tert-butylphenyl)-2H-benzo-

triazole, 2-(2-hydroxy-3,5-di-tert-amylphenyl)-2H-benzotriazole, 2-(2-hydroxy-3- α -cumyl-5-tert-octylphenyl)-2H-benzotriazole, 2,4-di-tert-butylphenyl 3,5-di-tert-butyl-4-hydroxybenzoate, 2-hydroxy-4-n-octyloxybenzophenone and 2,4-bis(2,4-dimethylphenyl)-6-(2-hydroxy-4-octyloxyphenyl)-s-triazine.

11. A composition according to claim 1 wherein the polymer substrate (A) is selected from the group of resins consisting of the polyolefins, the thermoplastic olefins, styrenic polymers and copolymers.

12. A composition according to claim 11 wherein the polymer substrate (A) is selected from the group consisting of polypropylene, polyethylene, thermoplastic olefin, acrylonitrile/butadiene/styrene copolymer and high impact polystyrene.

13. A composition according to claim 1 wherein component (i.) of the synergistic mixture (B) is present in an amount from 5.1 to 10 % by weight based on the polymer substrate (A).

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/02706

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C08K5/00 C08K5/3435

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C08K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 00450 A (CIBA GEIGY AG ;DYAS DARRELL DAVID JR (US); PUGLISI JOSEPH STEVEN () 7 January 1999 (1999-01-07) Exemples ---	1-13
X	US 5 326 805 A (SICKEN MARTIN ET AL) 5 July 1994 (1994-07-05) claims 1,9,10 ---	1-3,5-13
X	WO 98 28361 A (CIBA GEIGY AG ;GILG BERNARD (FR)) 2 July 1998 (1998-07-02) claims 1,5,11 ---	1-6,8-13
X	US 5 369 159 A (NESVADBA PETER) 29 November 1994 (1994-11-29) claim 1 -----	1-13

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

22 July 2002

Date of mailing of the international search report

30/07/2002

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rose, E

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 02/02706

Patent document cited in search report		Publication date	Patent family member(s)	Publication date			
WO 9900450	A	07-01-1999	AU 735643 B2	12-07-2001			
			AU 8174998 A	19-01-1999			
			BR 9810957 A	26-09-2000			
			CN 1261904 T	02-08-2000			
			EP 0996669 A1	03-05-2000			
			JP 2002507238 T	05-03-2002			
			WO 9900450 A1	07-01-1999			
			US 2002010235 A1	24-01-2002			
			US 5326805	A	05-07-1994	AT 162836 T	15-02-1998
AU 4428593 A	03-02-1994						
CA 2101655 A1	02-02-1994						
DE 59308059 D1	05-03-1998						
DK 584567 T3	21-09-1998						
EP 0584567 A2	02-03-1994						
ES 2114577 T3	01-06-1998						
FI 933402 A	02-02-1994						
JP 6172663 A	21-06-1994						
NO 932740 A	02-02-1994						
ZA 9305519 A	24-02-1994						
WO 9828361	A	02-07-1998				AT 202587 T	15-07-2001
						AU 5661398 A	17-07-1998
			DE 69705426 D1	02-08-2001			
			DE 69705426 T2	23-05-2002			
			WO 9828361 A1	02-07-1998			
			EP 0946632 A1	06-10-1999			
			ES 2158613 T3	01-09-2001			
			JP 2001507053 T	29-05-2001			
			SK 85399 A3	13-03-2000			
			TW 455603 B	21-09-2001			
			US 6228911 B1	08-05-2001			
US 5369159	A	29-11-1994	BE 1007037 A3	28-02-1995			
			CA 2096326 A1	23-11-1993			
			CZ 9300917 A3	19-01-1994			
			DE 4316611 A1	25-11-1993			
			FR 2691468 A1	26-11-1993			
			GB 2267491 A , B	08-12-1993			
			IT 1264452 B1	23-09-1996			
			JP 3250055 B2	28-01-2002			
			JP 6041109 A	15-02-1994			
			NL 9300801 A	16-12-1993			
			SK 49393 A3	06-04-1994			
			US 5488117 A	30-01-1996			