



US009402886B2

(12) **United States Patent**
Caggiano et al.

(10) **Patent No.:** **US 9,402,886 B2**
(45) **Date of Patent:** ***Aug. 2, 2016**

(54) **COMPOSITIONS AND METHODS OF USING CHONDROITINASE ABCI MUTANTS**

- (71) Applicant: **ACORDA THERAPEUTICS, INC.**, Ardsley, NY (US)
- (72) Inventors: **Anthony O. Caggiano**, Larchmont, NY (US); **Jennifer Iaci**, Boonton, NJ (US); **Andrea Vecchione**, Mt. Vernon, NY (US); **Elizabeth Hunter**, Manchester (GB)
- (73) Assignee: **Acorda Therapeutics, Inc.**, Ardsley, NY (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **14/480,486**

(22) Filed: **Sep. 8, 2014**

(65) **Prior Publication Data**

US 2015/0190483 A1 Jul. 9, 2015

Related U.S. Application Data

(60) Continuation of application No. 13/543,749, filed on Jul. 6, 2013, now Pat. No. 8,852,583, which is a continuation of application No. 12/757,006, filed on Jun. 4, 2010, now Pat. No. 8,236,302, which is a continuation of application No. 12/338,207, filed on Dec. 18, 2008, now Pat. No. 7,731,956, which is a division of application No. 11/527,318, filed on Sep. 26, 2006, now Pat. No. 7,485,295.

(60) Provisional application No. 60/720,628, filed on Sep. 26, 2005.

(51) **Int. Cl.**

- A61K 38/00** (2006.01)
- A61K 38/47** (2006.01)
- A61K 38/46** (2006.01)
- C12N 9/88** (2006.01)
- C12N 9/26** (2006.01)
- C12N 9/24** (2006.01)
- C12N 9/14** (2006.01)
- C07H 21/04** (2006.01)
- C07K 1/00** (2006.01)
- A61K 38/51** (2006.01)

(52) **U.S. Cl.**

CPC . **A61K 38/51** (2013.01); **C12N 9/88** (2013.01); **A61K 38/00** (2013.01); **C12Y 402/02004** (2013.01)

(58) **Field of Classification Search**

CPC **A61K 38/00**; **A61K 38/51**; **C12Y 402/02004**; **C12N 9/88**
USPC **424/94.5**, **94.61**, **94.62**; **435/232**, **201**, **435/200**, **195**; **536/23.2**; **530/350**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,262,522 A	11/1993	Gearing
5,270,194 A	12/1993	D'Alterio et al.
5,496,718 A	3/1996	Hashimoto
5,498,536 A	3/1996	Khandke
5,578,480 A	11/1996	Khandke
5,652,122 A	7/1997	Frankel et al.
5,670,617 A	9/1997	Frankel et al.
5,674,980 A	10/1997	Frankel et al.
5,747,641 A	5/1998	Frankel et al.
5,763,205 A	6/1998	Hashimoto et al.
5,792,743 A	8/1998	Schachner
5,804,604 A	9/1998	Frankel et al.
5,869,301 A	2/1999	Nghiem et al.
5,997,863 A	12/1999	Zimmermann et al.
6,007,810 A	12/1999	Ishikawa et al.
6,063,378 A	5/2000	Nohara et al.
6,093,563 A	7/2000	Bennett et al.
6,153,187 A	11/2000	Yacoby-Zeevi
6,171,575 B1	1/2001	Okuyama
6,184,023 B1	2/2001	Hashimoto et al.
6,200,564 B1	3/2001	Lamont et al.
6,248,562 B1	6/2001	Dunn et al.
6,313,265 B1	11/2001	Phillips et al.
6,326,166 B1	12/2001	Pomerantz et al.
6,972,168 B2	12/2005	Muir
7,008,783 B1	3/2006	Sato et al.
7,074,581 B2	7/2006	Yamashita et al.
7,163,545 B2	1/2007	Yaszemski et al.
7,465,705 B2	12/2008	Lee et al.
7,485,295 B2*	2/2009	Caggiano A61K 38/51 424/94.61

(Continued)

FOREIGN PATENT DOCUMENTS

AU	2003/208466 B2	9/2003
AU	2003/265561 A1	3/2004

(Continued)

OTHER PUBLICATIONS

Accession P59807, Aug. 15, 2003 UniProtKB/Swiss-Prot.

Aldrich "Enzymer Explorer" 2009, <http://www.sigmaaldrich.com/life-science/metabolomics/enzyme-explorer/learning-center/carbohydrate-analysis/carbohydrate-analysis-iii>.

Anderson et al. "Tumor Cell Retention of Antibody Fab Fragments is Enhanced by an Attached HIV TAT Protein-Derived Peptide" 1993, Biochem. & Biophys. Res. Commun. 194(2):876-884.

Appel et al. "Several Extracellular Domains of the Neural Cell Adhesion Molecule L1 are Involved in Neurite Outgrowth and Cell Body Adhesion" 1993, J. Neurosci. 13(11): 4764-4775.

Avrameas et al. "Polyreactive anti-DNA monoclonal antibodies and a derived peptide as vectors for the intracytoplasmic and intranuclear translocation of macromolecules" 1998, Proc. Natl. Acad. Sci. USA 95:5601-5606.

(Continued)

Primary Examiner — Ganapathirama Raghu
(74) Attorney, Agent, or Firm — Pepper Hamilton LLP

(57) **ABSTRACT**

One aspect of the present invention relates to mutants of chondroitinase ABCI. Such chondroitinase ABCI mutants exhibit altered chondroitin lyase activity or increased resistance to inactivation from stressors including exposure to UV light or heat. Methods of using chondroitinase ABCI mutant enzymes are also provided.

8 Claims, 2 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

7,507,570	B2	3/2009	Prabhakar et al.
7,560,106	B2	7/2009	Sasisekharan et al.
7,731,956	B2 *	6/2010	Caggiano A61K 38/51 424/94.5
8,236,302	B2 *	8/2012	Caggiano A61K 38/51 424/94.5
8,852,583	B2 *	10/2014	Caggiano A61K 38/51 424/94.5
2003/0040112	A1	2/2003	Muir et al.
2003/0072749	A1	4/2003	Muir et al.
2003/0077258	A1	4/2003	Muir
2004/0033221	A1	2/2004	Masuda et al.
2004/0265297	A1	12/2004	Gruskin et al.
2005/0118157	A1	6/2005	McMahon et al.
2005/0233419	A1	10/2005	Pojasek et al.
2006/0078959	A1	4/2006	Prabhakar et al.
2006/0153827	A1	7/2006	Gruskin et al.
2006/0233782	A1	10/2006	Gruskin et al.
2007/0104703	A1	5/2007	Caggiano et al.
2007/0274979	A1	11/2007	Gruskin et al.
2009/0104176	A1	4/2009	Caggiano et al.
2010/0239557	A1	9/2010	Caggiano et al.
2011/0250631	A1	10/2011	Gruskin et al.
2011/0262413	A1	10/2011	Gruskin et al.
2012/0207732	A1	8/2012	Gruskin et al.
2012/0308547	A1	12/2012	Caggiano et al.
2013/0210082	A1	8/2013	Caggiano et al.
2013/0243765	A1	9/2013	Gruskin et al.
2014/0193387	A1	7/2014	Gruskin et al.
2014/0248253	A1	9/2014	Gruskin et al.
2014/0322192	A1	10/2014	Gruskin et al.
2015/0023942	A1	1/2015	Gruskin et al.
2015/0299687	A1	10/2015	Gruskin et al.

FOREIGN PATENT DOCUMENTS

AU	2004/241088	A1	12/2004
AU	2006/294755	B2	4/2012
CA	2623635	C	4/2013
EP	0704532	A2	4/1996
EP	1631234	A2	3/2006
EP	1646353	A2	4/2006
EP	2353606	A2	8/2011
EP	2354155	A2	8/2011
JP	H06 (1994)-153947		6/1994
JP	H10 (1998)-506263		6/1998
JP	H10-174598		6/1998
JP	H11 (1999)-500308		1/1999
JP	H11 (1999) 236336		8/1999
JP	2002-505873		2/2002
JP	2002-526028		8/2002
JP	2003-500016		1/2003
JP	2004-89191		3/2004
JP	2004-113166		4/2004
JP	2013-5391069		10/2013
JP	5452820	B2	1/2014
WO	WO 91/06303	A	5/1991
WO	WO 94/25567	A1	11/1994
WO	WO 95/13091	A1	5/1995
WO	WO 95/14478	A1	6/1995
WO	WO 99/46368	A2	9/1999
WO	WO 00/52149	A1	9/2000
WO	WO 00/62067	A1	10/2000
WO	WO 00/64482	A1	11/2000
WO	WO 00/75319	A1	12/2000
WO	WO 01/39795	A1	6/2001
WO	WO 02/08285	A2	1/2002
WO	WO 02/055684	A	7/2002
WO	WO 02/065136	A2	8/2002
WO	WO 02/083179	A2	10/2002
WO	WO 03/000901	A2	1/2003
WO	WO 03/015612	A2	2/2003
WO	WO 03/022882	A2	3/2003
WO	WO 03/031578	A2	4/2003

WO	WO 03/074080	A1	9/2003
WO	WO 03/100031	A2	12/2003
WO	WO 03/102160	A2	12/2003
WO	WO 2004/017044	A2	2/2004
WO	WO 2004/103299	A2	12/2004
WO	WO 2004/108069	A2	12/2004
WO	WO 2004/110359	A2	12/2004
WO	WO 2004/110360	A2	12/2004
WO	WO 2005/112986	A2	12/2004
WO	WO 2005/087920	A2	9/2005
WO	WO 2007/038548	A2	4/2007
WO	WO 2008/045970	A2	4/2008

OTHER PUBLICATIONS

- Banker et al. "Modern Pharmaceutics" 1979, Marcel Dekker, Inc. (TOC).
- Banker et al. "Modern Pharmaceutics" 2002, 4th Ed., Informa Healthcare, New York (TOC).
- Bao et al. "A Functional Dermatan Sulfate Epitope Containing Iduronate (2-O-sulfate) α 1-3GalNAc (6-O-sulfate) Disaccharide in the Mouse Brain" 2005, J. of Bio. Chem. 280(24):23184-23193.
- Basso et al. "A Sensitive and Reliable Locomotor Rating Scale for Open Field Testing in Rats" 1995, J. of Neurotrauma 12(1):1-21.
- Ben-Bassat et al. "Processing of the Initiation Methionine from Proteins: Properties of the *Escherichia coli* Methionine Aminopeptidase and Its Gene Structure" 1987, J. Bacteriol. 169(2):751-757.
- Bixby et al. "Neurite outgrowth on muscle cell surfaces involves extracellular matrix receptors as well as Ca²⁺-dependent and -independent cell adhesion molecules" 1987, Proc. Natl. Acad. Sci. USA 84:2555-2559.
- Blight et al. "Animal models of spinal cord injury" 2002, Top Spinal Cord Inj. Rehabi. 6(2):1-13.
- Bowie et al. "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions" 1990, Science 247:1306-1319.
- Bradbury et al. "Chondroitinase ABC Promotes Functional Recovery After Spinal Cord Injury" 2002, Nature 416:636-640.
- Bradbury et al. "Chondroitinase ABC Promotes Regeneration and Functional Recovery Following Spinal Cord Injury" 2001, Soc. for Neuroscience Abstracts 27(2):1835.
- Bradbury et al. "NT-3 Promotes Growth of Lesioned Adult Rat Sensory Axons Ascending in the Dorsal Columns of the Spinal Cord" 1999, Eur. J. Neurosci. 11(11):3873-3783.
- Bray et al. "Neuronal and Nonneuronal Influences on Retinal Ganglion Cell Survival, Axonal Regrowth, and Connectivity after Axotomy" 1991, Ann. NY Acad. Sci., 214-228.
- Broach et al. "Experimental Manipulation of Gene Expression" M. Inouye ed., Academic Press, New York, 1983, pp. 83-117.
- Burgess et al. "Possible Dissociation of the Heparin-binding and Mitogenic Activities of Heparin-binding (Acidic Fibroblast) Growth Factor-1 From Its Receptor-binding Activities by Site-directed Mutagenesis of a Single Lysine Residue" 1990, J. of Cell. Bio. 111:2129-2138.
- Cadelli et al. "Oligodendrocyte- and Myelin-Associated Inhibitors of Neurite Outgrowth: Their Involvement in the Lack of CNS Regeneration" 1992, Exp. Neur. 115:189-192.
- Caggiano et al., Chondroitinase ABCI Improves Locomotion and Bladder Function following Contusion Injury of the Rat Spinal Cord, Feb. 1, 2005, J. Neurotrauma 22(2):226-239.
- Cajal "Degeneration & Regeneration of the Nervous System" May 1959 ed., Hafner Publ. Co., New York (TOC).
- Chang et al. "Extension of Neurites on Axons is Impaired by Antibodies against Specific Neural Cell Surface Glycoproteins" 1987, J. Cell. Biol. 104:355-362.
- Chau et al. "Chondroitinase ABC Enhances Axonal Regrowth Through Schwann Cell-seeded Guidance Channels After Spinal Cord Injury" Nov. 20, 2003 FASEB J. 18(1):1-24.
- Chen et al. "Peripheral nerve regeneration using silicone rubber chambers filled with collagen, laminin and fibronectin" 2000, Biomat. 21:1541-1547.
- Crespo et al. "How Does Chondroitinase Promote Functional Recovery in the Damaged CNS?" 2007, Ex. Neurology 206:159-171.
- Curinga et al. "Mammalian-produced Chondroitinase AC Mitigates Axon Inhibition by Chondroitin Sulfate Proteoglycans" 2007, J. of Neurochemistry 102:275-288.

(56)

References Cited

OTHER PUBLICATIONS

- Daichi "Text Book of Physiology" 2000, 3rd Ed. 81.
- Degrendele et al. "Requirement for CD44 in Activated T Cell Extravassation into an Inflammatory Site" 1997, *Science* 278:672-675.
- Denuziere et al. "Chitosan-Chondroitin sulfate and chitosan-hyaluronate polyelectrolyte complexes: biological properties" 1998, *Biomaterials* 19:1275-1285.
- Derossi et al. "Cell Internalization of the Third Helix of the Antennapedia Homeodomain is Receptorindependent" 1996, *J. Biol. Chem* 271:18188-18193.
- Dimayuga et al. "The Neuregulin GGF2 Attenuates Free Radical Release from Activated Microglial Cells" Mar. 2003, *J. Neuroim.* 136(1-2):67-74.
- Doppenberg et al. "Clinical Trials in Traumatic Brain Injury" 1998, *Ann. NY Acad. Sci.* 305-319.
- Edelman "Cell Adhesion Molecules" 1983, *Science* 219:450-457.
- Edelman et al. "Morphoregulatory Molecules" 1990, Wiley, New York (TOC).
- Efthymiadis et al. "The HIV-1 Tat Nuclear Localization Sequence Confers Novel Nuclear Import Properties" 1998, *J. Biol. Chem.*, Jan. 16,273(3):1623-1628.
- Ellioit et al. "Intercellular Trafficking and Protein Delivery by a Herpesvirus Structural Protein" 1997, *Cell* 88:223-233.
- European Search Report and Written Opinion for EP04752310 dated Oct. 7, 2008.
- European Search Report and Written Opinion for EP06815505 dated Feb. 22, 2010.
- European Search Report and Written Opinion for EP10183555 dated Jan. 20, 2011.
- European Search Report and Written Opinion for EP10184697 dated Jul. 12, 2011.
- European Search Report and Written Opinion for EP11152626 dated Jul. 21, 2011.
- Fahraeus et al. "Inhibition of pRb phosphorylation and cell-cycle progression by a 20-residue peptide derived from p16CDKN2IINK4A" 1996, *Curr. Biol.* 6(1):84-91.
- Favre et al. "Hyaluronidase enhances recombinant adeno-associated virus (Raav)-mediated gene transfer in the rat skeletal muscle" 2000, *Gene Ther.* 7(16):1417-1420.
- Fawcett et al. "The glial scar and central nervous system repair" 1999, *Brain Res. Bull.* 49(6):377-391.
- Fawell et al. "Tat-mediated delivery of heterologous proteins into cells" 1994, *Proc. Natl. Acad. Sci. USA* 91:664-668.
- Fethiere et al. "Crystal Structure of Chondroitin AC Lyase, a Representative of a family of Glycosaminoglycan Degrading Enzymes" 1999, *J. Mol. Biol.* 288:635-647.
- Fongmoon et al. "Chondroitinase-mediated Degradation of Rare 3-) Sulfated Glucuronic Acid in Functional Oversulfated Chondroitin Sulfate K and E" 2007, *J. of Bio. Chem.* 282(51):36895-39904.
- Frankel et al. "Tat Protein from Human Immunodeficiency Virus Forms a Metal-Linked Dimer" 1988, *Science* 240:70-73.
- Frankish et al. "Spinal-cord Repair Moves a Step Closer" 2002, *The Lancet* 359(9314):1317.
- Gennaro "Remington's Pharmaceutical Sciences" 1985, Mack Publishing Company (PA) 17th Ed. (TOC).
- Goodman et al. "The Pharmacological Basis of Therapeutics" 1980, 6th ed., MacMillan Pub., New York (TOC).
- Goodman et al. "The Pharmacological Basis of Therapeutics" 2001, 10th ed., McGraw Hill, New York (TOC).
- Grandpre et al. "Nogo-66 Receptor Antagonist Peptide Promotes Axonal Regeneration" May 30, 2002, *Nature* 417(6888):547-551.
- Hamai et al. "Two Distinct Chondroitin Sulfate ABC Lyases" 1997, *J. Biol. Chem.* 272(14):9123-9130.
- Hirschberg et al. "Inflammation after axonal injury has conflicting consequences for recovery of function: rescue of spared axons is impaired but regeneration is supported" 1994, *J. Neuroimmunol.* 50(1):9-16 (Abstract).
- Hiyama et al. "Crystallization and Some Properties of Chondroitinase from *Arthrobacter Aureus*" 1975, *J. Biol. Chem.* 250:1824-1828.
- Hlavín et al. "Molecular Structure and Functional Testing of Human LICAM: An Interspecies Comparison" 1991, *Genomics* 11:416-423.
- Hoffman et al. "Chondroitin Sulfates" 1958, *Federation Proc.* 17:1078-1082.
- Horstkorte et al. "The Fourth Immunoglobulin-like Domain of NCAM Contains a Carbohydrate Recognition Domain for Oligomannosidic Glycans Implicated in Associated with L1 and Neurite Outgrowth" 1993, *J. Cell Biol.* 121(6):1409-1421.
- Hou et al. "Endotoxin Removal by Anion-Exchange Polymeric Matrix" 1990, *Biotech. Appl. Biochem.* 12:315-324.
- Huang et al. "Active Site of Chondroitin AC Lyase Revealed by the Structure of Enzyme-Oligosaccharide Complexes and Mutagenesis" Jan. 1, 2001, *Biochemistry*, 40(8):2359-2372.
- Huang et al. "Crystal Structure of Chondroitinase B from *Flavobacterium heparinum* and its Complex with a Disaccharide Product at 107 Å Resolution" 1999, *J. Mol. Biol.* 294:1257-1269.
- Huang et al. "Crystal Structure of *Proteus vulgaris* Chondroitin Sulfate ABC Lyase I at 1.9 Å Resolution" May 2, 2003, *J. Mol. Biol.* 328(3):623-634.
- Hunt et al. "The Nogo Receptor, Its Ligands and Axonal Regeneration in the Spinal Cord; a Review" Feb. 2002, *J. Neurocytology* 31(2):93-120.
- Iida et al. "Cell Surface Chondroitin Sulfate Proteoglycans in Tumor Cell Adhesion, Motility and Invasion" 1996, *Seminars in Cancer Biology* 7:155-162.
- International Search Report for PCT/US2006/037554 dated Aug. 15, 2008.
- Iwai et al. "Axon Patterning Requires DN-cadherin, a Novel Neuronal Adhesion Receptor, in the *Drosophila* Embryonic CNS" 1997, *Neuron* 19:77-89.
- Jones "Taking a new TAK on Tat transactivation" 1997, *Genes & Dev.* 11:2593-2599.
- Jung et al. "Transit time of leucocytes rolling through venules controls cytokine-induced inflammatory cell recruitment in vivo" 1998, *J. Clin. Invest.* 102(8):1526-1533.
- Kadmon et al. "Functional Cooperation between the Neural Adhesion Molecules L1 and N-CAM is Carbohydrate Dependent" 1990, *J. Cell Biol.* 110:209-218.
- Kadmon et al. "The Neural Cell Adhesion Molecule N-CAM Enhances L1-dependent Cell-Cell Interactions" 1990, *J. Cell Biol.* 110:193-208.
- Khan et al. "Animal Models of Spinal Cord Contusion Injuries" 1999, *Laboratory Animal Science* 49(2): 161-172.
- Kim et al. "Insertion and Deletion Mutants of *FoII* Restriction Endonuclease" 1994, *J. Biol. Chem.* 269(50):31978-31982.
- Korn, 1957 "The Degradation of Heparin by Bacterial Enzymes" *J. Biol. Chem.* 226:841-844.
- Krekoski et al. "Axonal Regeneration into Acellular Nerve Grafts is Enhanced by Degradation of Chondroitin Sulfate Proteoglycan" 2001, *J. Neurosci.* 15:21(16):6206-6213.
- Kubota et al. "Functional Similarity of HIV-1 Rev and HTLV-1 Rex Proteins: Identification of a New Nucleolar-Targeting Signal in Rev Protein" Aug. 15, 1989, *Biochem. Biophys. Res. Commun.* 162(3):963-970.
- Kwon et al. "Animal Models Used in Spinal Cord Regeneration Research" 2002, *Spine* 27(14):1504-1510.
- Lagenaur et al. "An L1-like molecule, the 8D9 antigen, is a potent substrate for neurite extension" 1987, *Proc. Natl. Acad. Sci. USA* 84:7753-7757.
- Lemons et al. "Chondroitin Sulfate Preteoglycan Immunoreactivity Increases Following Spinal Cord Injury and Transplantation" 1999, *Exper. Neurology* 160:51-65.
- Lesley et al. "Variant Cell Lines Selected for Alterations in the Function of the Hyaluronan Receptor CD44 Show Differences in Glycosylation" 1995, *J. Exp. Med.* 182:431-437.
- Li et al. "Delayed systemic Nogo-66 Receptor Antagonist Promotes Recovery from Spinal Cord Injury" 2003, *J. Neuroscience* 23(10):4219-4227.

(56)

References Cited

OTHER PUBLICATIONS

- Lindner et al. "L1 mono- and polyclonal antibodies modify cell migration in early postnatal mouse cerebellum" 1983, *Nature* 305:427-430.
- Lodish et al. "Integrating cells into tissue" 2000, *Mol. Cell Biology*, 5th Ed., Chapter 6.
- Mahanthappa et al. "Glial Growth Factor 2, a Soluble Neuregulin, Directly Increases Schwann Cell Motility and Indirectly Promotes Neurite Outgrowth" 1996, *J. Neuroscience* 16(15):4673-4683.
- Maniatis et al. "Molecular Cloning: A Laboratory Manual" 1982, Cold Spring Harbor Lab. (TOC).
- Mann et al. "Endocytosis and Targeting of Exogenous HIV-1 Tat Protein" 1991, *EMBO J.*, July, 10(7):1733-1739.
- Martinez et al. "Purification and Properties of the Enzyme Chondroitinase" 1959, *J. Biol. Chem.* 234(9):2236-2239.
- Martini et al. "Restricted Localization of L1 and N-CAM Sites of Contact Between Schwann Cells and Neurites in Culture" 1994, *GLIA* 10:70-74.
- Matynyn "Restoration of functions due to Enzyme Therapy After Complete Transection of the Spinal Cord" 1965, *ZH EK SP KLIN MED* 5(3):3-13.
- Matsumoto et al. "Peripheral nerve regeneration across an 80-mm gap bridged by a polyglycolic acid (PGA)-collagen tube filled with laminin-coated collagen fibers: a histological and electrophysiological evaluation of regenerated nerves" 2000, *Brain Res.* 868:315-328.
- Matteuci et al. "Synthesis of Deoxyoligonucleotides on a Polymer Support" 1981, *J. Am. Chem. Soc.* 103:3185-3191.
- McGee et al. "The Nogo-66 Receptor: Focusing Myelin Inhibition of Axon Regeneration" Apr. 2003, *Trends in Neuroscience* 26(4):193-198.
- Michelacci et al. "A Comparative Study Between a Chondroitinase B and a Chondroitinase AC from *Flavobacterium heparinum*" 1975, *Biochem. J.* 151:121-129.
- Michelacci et al. "Chondroitinase C from *Flavobacterium heparinum*" 1976, *J. Biol. Chem.* 251(4):1154-1158.
- Michelacci et al. "Isolation and characterization of an induced Chondroitinase ABC" 1987, *Biochem. Biophys. Acta* 923:291-301.
- Michelacci et al. "Isolation and Partial Characterization of an Induced Chondroitinase β from *Flavobacterium Heparium*" 1974, *Biochem. & Biophys. Res. Comm.* 56(4):973-980.
- Miller et al. "N-terminal methionine-specific peptidase in *Salmonella typhimurium*" 1987, *PNAS* 84:2718-2722.
- Miura et al. "Analysis of Glycosaminoglycan-Degrading Enzymes by Substrate Gel Electrophoresis (Zymography)" 1995, *Anal. Biochem.* 225:333-340.
- Modena et al. "Hyaluronidase-injectable microparticles intended for the treatment of extravasation" 1998, *J. Microencapsulation* 15(1):85-92.
- Moon et al. "Regeneration of CNS axons back to their target following treatment of adult rat brain with chondroitinase ABC" 2001, *Nature Neurosc.* 4(5): 465-466.
- Moos et al. "Neural adhesion molecule L1 as a member of the immunoglobulin superfamily with binding domains similar to fibronectin" 1988, *Nature* 334:701-703.
- Nagahara et al. "Transduction of full-length TAT fusion proteins into mammalian cells: TAT_{p27K1p1} induces cell migration" 1998, *Nat. Med.* 4(12):1449-1452.
- Netti et al. "Role of Extracellular Matrix Assembly in Interstitial Transport in Solid Tumors" 2000, *Cancer Res.* 60(9):2497-2503.
- Nieke et al. "Expression of the neural cell adhesion molecules L1 and N-CAM and their common carbohydrate epitope L2/HNK-1 during development and after transection of the mouse sciatic nerve" 1985, *Differentiation* 30:141-151.
- Oermann et al. "The Use of Anti-inflammatory Medications in Cystic Fibrosis" 1999, *Chest* 115:1053-1058.
- Olmaker et al. "Chondroitinase ABC (Pharmaceutical Grade) for Chemonucleolysis" 1996, *Spine* 21(17):1952-1956.
- Pawson et al. "Assembly of Cell Regulatory systems Through Protein Interaction Domains" 2003, *Science* 300:445-452.
- Pillwein et al. "Hyaluronidase Additional to Standard Chemotherapy Improves Outcome for Children with Malignant Tumors" 1998, *Cancer Letters* 131:101-108.
- Pojasek et al. "Biochemical Characterization of the Chondroitinase B Active Site" Aug. 23, 2002, *J. Biol. Chem.*, 277(34):31179-31186.
- Pojasek et al. "Recombinant Expression, Purification, and Kinetic Characterization of Chondroitinase AC and Chondroitinase B from *Flavobacterium heparinum*" 2001, *Biochem, Biophys. Res. Commun.* 286:343-351.
- Prabhakar et al. "Biochemical Characterization of the Chondroitinase ABC I Active Site" Aug. 23, 2005, *Biochem. J.* 390(2):395-405.
- Priestley et al. "Stimulating regeneration in the damaged spinal cord" 2002, *J. Phyl.* 96:123-133.
- Rathjen et al. "Immunocytological and biochemical characterization of a new neuronal cell surface component (L1 antigen) which is involved in cell adhesion" 1984, *EMBO J.* 3(1):1-10.
- Ratjen et al. "Cystic Fibrosis" 2003, *The Lancet* 361(9358):681-689 (Presentation).
- Reich et al. "Small interfering RNA (siRNA) targeting VEGF effectively inhibits ocular neovascularization in a mouse model" 2003, *Molecular Vision* 9:210-216 (Abstract).
- Reid et al. "Variants of Human L1 Cell Adhesion Molecule Arise through Alternate Splicing of RNA" 1992, *J. Mol. Neurosc.* 3:127-135.
- Roy et al. "Generation of Substantially Smaller Deletion Mutants of Chondroitinase AC and B Those are Biologically Active" Nov. 8-12, 2003, Society for Neuroscience Abstract Viewer and Itinerary Planner, 33rd Annual Meeting of the Society of Neuroscience, New Orleans, LA, Database Biosis, (Abstract).
- Roy et al. "Treatment with Recombinant Chondroitinases AC and B Permits Neuronal Outgrowth Over Inhibitory Chondroitin Sulfate Proteoglycans (CSPGs)" Nov. 7, 2002, Society for Neuroscience Abstract Archives 2000-2005 (Abstract).
- Saito et al. "Enzymatic Methods for the Determination of Small Quantities of Isomeric Chondroitin Sulfates" 1968, *J. Biol. Chem.* 243(7):1536-1542.
- Sambrook et al. "Molecular Cloning" 1989, 2nd ed., Cold Spring Harbor Laboratory Press, Ch. 16 and 17.
- Sambrook et al. "Molecular Cloning" 1989, 2nd ed., Cold Spring Harbor Laboratory Press, TOC.
- Sato et al. "Cloning and expression in *Escherichia coli* of the gene encoding the *Proteus vulgaris* chondroitin ABC-lyase" Jan. 1, 1994, *Appl. Microbiol. Biotechnol.* 41:39-46.
- Sato, et al. "Subunit Structure of Chondroitinase ABC from *Proteus Vulgaris*" 1986 *Agric. Biol. Chem.* 50(4):1057-1059.
- Schachner "Functional implications of glial cell recognition molecules" 1990, *Neurosc.* 2:497-507.
- Schwab "Nerve fibre regeneration after traumatic lesions of the CNS; progress and problems" 1991, *Phil. Trans. R. Soc. Lond.* 331:303-306.
- Schwarze et al. "In Vivo Protein Transduction: Delivery of a Biologically Active Protein into the Mouse" 1999, *Science* 285:1569-1572.
- Seikagaku Biobus. Corp. "Chondroitinase AC II pamphlet" 2009, http://www.seikagakubb.co.jp/bio/cgi-bin/search/tenpu_pdf/100335.pdf.
- Seilheimer et al. "Studies of Adhesion Molecules Mediating Interactions between Cells of Peripheral Nervous System Indicate a Major Role for L1 in Mediating Sensory Neuron Growth on Schwann Cells in Culture" 1988, *J. Cell Biol.* 107:341-351.
- Silver et al. "Postnatally induced formation of the corpus callosum in acallosal mice on glia-coated cellulose bridges" 1983, *Science* 220:1067-1069.
- Smiseth et al. "Effect of Hyaluronidase on Substrate Exchange and Blood Flow in the Ischaemic Myocardium of the Dog" 1982, *Clinical Physiology* 2(1):39-50.
- Smith-Thomas et al. "Increased Axon Regeneration in Astrocytes Grown in the Presence of Proteoglycan Synthesis Inhibitors" 1995, *J. of Cell Science* 108(3):1307-1315.
- Southern "Detection of Specific Sequences Among DNA Fragments Separated by Gel Electrophoresis" 1975, *J. Mol. Biol.* 98:503-517.
- Stedman's Medical Dictionary 2000, Lippincott Williams & Wilkins, 27th Ed.

(56)

References Cited

OTHER PUBLICATIONS

Sterne et al. "Neurotrophin-3 Delivered Locally via Fibronectin Mats Enhances Peripheral Nerve Regeneration" 1997, *Eur. J. Neurosc.* 9:1388-1396.

Tona et al. "Effect of Hyaluronidase on Brain Extracellular Matrix in Vivo and Optic Nerve Regeneration" 1993, *J. Neurosc. Res.* 36:191-199.

Trigg et al. "Peripheral Nerve Regeneration: Comparison of Laminin and Acidic Fibroblast Growth Factor" 1998, *Am. J. Otolaryngology* 19(1):29-32.

Tsuda et al. "Substrate Specificity Studies of Flavobacterium Chondroitinase C and Heparitinases Towards the Glycosaminoglycan-protein Linkage region" 1999, *European J. of Biochem.* 262:127-133.

Vives et al. "A Truncated HIV-1 Tat Protein Basic Domain Rapidly Translocates through the Plasma Membrane and Accumulates in the Cell Nucleus" 1997, *J. Biol. Chem.* 272(25):16010-16017.

Vives et al. "Effects of the Tat Basic Domain on Human Immunodeficiency Virus Type 1 Transactivation, Using Chemically Synthesized Tat Protein and Tat Peptides" May 1994, *J. Virol.* 68(5):3343-3353.

Williams et al. "Calcium Influx into Neurons Can Solely Account for Cell Contact-dependent Neurite Outgrowth Stimulated by Transfected L1" 1992, *J. Cell Biol.* 119(4):883-892.

Wood et al. "Inhibition of Schwann Cell Myelination in vitro by Antibody to the L1 Adhesion Molecule" 1990, *J. Neurosc.* 10(11):3635-3645.

Yamagata et al. "Purification and Properties of Bacterial Chondroitinases and Chondrosulfatases" 1968, *J. Biol. Chem.* 243(7):1523-1535.

Yamagata et al. "Repression of a Malignant Cell-Substratum Adhesion Phenotype by Inhibiting the Production of the Anti-Adhesive Proteoglycan, PG-M/Versican" 1994, *J. of Cell Science* 1007:2581-2590.

Yang et al. "Developmental Regulation of a Matrix Metalloproteinase during Regeneration of Axolotl Appendages" 1994, *Dev. Biol.* 166:696-703.

Yang et al. "Expression of *Mmp-9* and Related Matrix Metalloproteinase Genes During Axolotl Limb Regeneration" 1999, *Dev. Dyn.* 216:2-9.

Yasuda et al. "Effect of Hyaluronidase on Experimental Cerebral Infarct Size and Mortality" 1982, *Lab. Invest.* 46(4):400-404.

Yick et al. "Chondroitinase ABC promotes axonal regeneration of Clarke's neurons after spinal cord injury" 2000, *Regeneration and Transpl.* 11(5):1063-1067.

Yick et al. "Chondroitinase ABC Promotes Axonal Regrowth of Clarke's Neurons Into Peripheral Nerve Graft After Hemisection of the Spinal Cord" 1999, *Soc. for Neuroscience Abstracts* 25:747.

Zuo et al. "Degradation of Chondroitin Sulfate Proteoglycan Enhances the Neurite-Promoting Potential of Spinal Cord Tissue" 1998, *Exp. Neurol.* 154(2):654-662.

Zuo et al. "Regeneration of Axons After Nerve Transection Repair is Enhanced by Degradation of Chondroitin Sulfate Proteoglycan" 2002, *Exp. Neurology* 176:221-228.

* cited by examiner




Fig. 1




Fig. 2

COMPOSITIONS AND METHODS OF USING CHONDROITINASE ABCI MUTANTS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation application of U.S. application Ser. No. 13/543,749, filed Jul. 6, 2012, now U.S. Pat. No. 8,852,583, issued Oct. 7, 2014, which is a continuation application of U.S. application Ser. No. 12/757,006, filed Jun. 4, 2010, now U.S. Pat. No. 8,236,302, issued Aug. 7, 2012, which is a continuation application of U.S. application Ser. No. 12/338,207, filed Dec. 18, 2008, now U.S. Pat. No. 7,731,956, issued Jun. 8, 2010, which is a divisional application of U.S. application Ser. No. 11/527,318, filed Sep. 26, 2006, now U.S. Pat. No. 7,485,295, issued Feb. 3, 2009, which claims priority from U.S. Provisional Application No. 60/720,628, filed Sep. 26, 2005; all aforementioned applications being herein incorporated by reference in their entirety.

BACKGROUND

Proteoglycans, major constituents of the extracellular matrix, are known to be present in large amounts in glial scar tissue and to inhibit recovery following spinal cord injuries (Fawcett & Asher, 1999). Enzymes that are capable of digesting glial scar tissue are an important target for the development of spinal cord injury (SCI) therapeutics. Chondroitinase ABCI (EC 4.2.2.4; cABCI) is a bacterial enzyme that catalyzes the digestion of sulfated chondroitin and dermatan side chains of proteoglycans. This enzyme has been shown to promote functional recovery after spinal cord injury (Bradbury et al., 2002; Caggiano et al., 2005).

The spinal cord is made up of nerve fibers. Damage to the central nervous system, including the spinal cord, results in a loss of function. Depending upon the type of injury to the central nervous system, the loss of function may manifest itself in loss of sensory, motor or autonomic function or a combination thereof. Sensory functions include the ability to feel sensations, like pain. Motor functions include the ability to voluntarily move your body. Autonomic functions include involuntary body functions, for example the ability to sweat and breathe.

The most common types of spinal cord injuries (SCI) include contusions (bruising of the spinal cord) and compression injuries (caused by prolonged pressure on the spinal cord). In contusion and compression injuries, a cavity or hole often forms in the center of the spinal cord. Unlike nerve cells, or neurons of the peripheral nervous system (PNS), neurons of the central nervous system (CNS) do not regenerate after injury.

Spinal cord injury can be characterized by contusion of the neural tissue with a resultant decrease or loss of the ability of nerve tissue to properly transmit nerve impulses. The usual cause is due to an impact injury of some nature, but it may also occur during the manipulation of the spinal cord in certain surgical procedures. After a spinal cord injury in the adult mammal, the inability of axons to regenerate may lead to loss of sensation, loss of motor function and/or loss of autonomic function, as well as permanent paralysis. One reason that neurons fail to regenerate is their inability to traverse the glial scar that develops following a spinal cord injury. The injury-induced lesion will develop glial scarring, which contains extracellular matrix molecules including chondroitin sulfate proteoglycans (CSPGs). CSPGs inhibit nerve tissue growth in vitro and nerve tissue regeneration at CSPGs rich regions in vivo.

A number of molecules, and specified regions thereof, have been implicated in the ability to support the sprouting of neurites from a neuronal cell, a process also referred to as neurite outgrowth. The term neurite refers to both axon and dendrite structures. The process of sprouting neurites is essential in neural development and regeneration, especially after physical injury or disease has damaged neuronal cells. Neurites elongate profusely during development both in the central and peripheral nervous systems of all animal species. This phenomenon pertains to both axons and dendrites.

Various polypeptides, especially cell adhesion molecules (CAMs), have been known to promote neural cell growth. While early efforts in this area of research concentrated on the adhesion-promoting extracellular matrix protein fibronectin (FN), other polypeptides have also been found to promote neural growth. For example, U.S. Pat. No. 5,792,743 discloses novel polypeptides and methods for promoting neural growth in the CNS of a mammal by administering a soluble neural CAM, a fragment thereof, or a Fc-fusion product thereof. U.S. Pat. No. 6,313,265 discloses synthetic polypeptides containing the pharmacologically active regions of CAMs that can be used in promoting nerve regeneration and repair in both peripheral nerve injuries as well as lesions in the CNS. While helpful, the use of regenerative proteins alone may not be sufficient to effect repair of a damaged nervous system.

During approximately the past two decades, knowledge of cell adhesion and migration in extracellular matrices (ECMs) at the molecular level has expanded rapidly. The action of enzymes and other polypeptides which degrade components of the extracellular matrix and basement membranes may facilitate the events of neural repair by a variety of mechanisms, including the release of bound cytokines and by increasing the permeability of the matrix, thereby enhancing the mobility of mediator molecules, growth factors and chemotactic agents, as well as the cells involved in the healing process. For example, U.S. Pat. No. 5,997,863 discloses the use of glycosaminoglycans to manipulate cell proliferation and promote wound healing.

Components of the inhibitory CSPGs have been identified as the glycosaminoglycans, chondroitin sulfate (CS) and dermatan sulfate (DS). Removal of these inhibitory molecules would allow neurites to regenerate and reinnervate an area after physical injury or disease, as well as to allow for the recovery of sensory, motor and autonomic functions.

Previous studies have found that chondroitinases can lyse and degrade CSPGs including, CS and DS. One study found that chondroitinase ABC removed glycosaminoglycan (GAG) chains in and around lesioned areas of rat CNS in vivo. The degradation of GAGs promoted expression of a growth-associated protein, GAP-43, indicating an increase in the ability of treated cells to regenerate. However, this growth-associated protein is associated with regeneration in peripheral, but not central, nerve injuries.

Chondroitin sulfates (CS) are sulfated polysaccharides in linear chains of a repeated disaccharides. They range in molecular weight from about 10,000 to over 100,000 Da. Chondroitin sulfate substrates exist in different isomers designated by the appended letters A, B, and C (Hoffman et al., 1958). The repeating units are composed of uronic acid (GlcA or IdoA) and galactosamine, and are called galactosaminoglycans, and are one example of the glycosaminoglycans, typically abbreviated as GAG. Although these GAG chain species have different repeating disaccharide regions, they are covalently bound through the so-called linkage region tetrasaccharide sequence (see below) to the serine residue in the GAG attachment consensus sequence (Glu/Asp-X-Ser-

Gly) of respective core proteins. Chondroitin A and C sulfates (ChS-A, ChS-C) are the most abundant GAGs and are found in cartilage, bone and heart valves. Chondroitin B (ChS-B, or, alternatively, dermatan sulfate) is expressed mostly in skin, blood vessels, and heart valves.

When chondroitinase bacterial preparations were characterized against different chondroitin sulfate (ChS) substrates, a series of distinct chondroitinases were discovered: Chondroitinase AC that degrades mostly chondroitin A (ChA) and chondroitin C (ChC) (Yamagata et al., 1968), Chondroitinase B that degrades chondroitin B (ChB) (Michelacci and Dietrich, 1976), Chondroitinase C that acts mostly on ChC (Michelacci Y M & Dietrich C P, 1976) and Chondroitinase ABC exhibits specificity against all three substrates—ChS-A, ChS-B and ChS-C (Yamagata et al., 1968, Michelacci et al., 1987).

SUMMARY OF THE INVENTION

One aspect of the present invention provides mutants of chondroitinase ABCI.

In preferred embodiments, such chondroitinase ABCI mutants exhibit enhanced activity. In other preferred embodiments, such chondroitinase ABCI mutants exhibit enhanced resistance to inactivation, including inactivation from UV or heat exposure. More preferably, the chondroitinase ABCI mutant enzymes are selected from BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), BF4 (SEQ ID NO:3). In another preferred embodiment, the chondroitinase ABCI mutant enzymes are selected from BC9 (SEQ ID NO:4), BC7 (SEQ ID NO:5), RD4 (SEQ ID NO:6) and BE11 (SEQ ID NO: 7).

Another embodiment of the present invention is a method of designing mutants of chondroitinase ABCI having altered activity.

Other embodiments of the present invention relate to methods for promoting neurological functional recovery, including sensory, motor and autonomic function, after central nervous system ("CNS") injury or disease.

Further embodiments relate to methods of promoting neuronal outgrowth and the use in treating spinal cord injuries and related disorders of the CNS by administering such chondroitinase ABCI mutants.

DESCRIPTION OF THE DRAWINGS

In part, other aspects, features, benefits and advantages of the embodiments of the present invention will be apparent with regard to the following description, appended claims and accompanying drawings where:

FIG. 1 is a bar graph of the chondroitin lyase activity of wild-type, not-inactivated chondroitinase ABCI (normal), wild-type, inactivated (No Enz) and chondroitinase ABCI mutant enzymes of the present invention following UV exposure.

FIG. 2 is a bar graph of the chondroitin lyase activity of wild-type, not-inactivated chondroitinase ABCI (normal), wild-type, inactivated (No Enz) and chondroitinase ABCI mutant enzymes of the present invention.

DETAILED DESCRIPTION

Before the present compositions and methods are described, it is to be understood that this invention is not limited to the particular molecules, compositions, methodologies or protocols described, as these may vary. It is also to be understood that the terminology used in the description is for the purpose of describing the particular versions or

embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must also be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to a "cell" is a reference to one or more cells and equivalents thereof known to those skilled in the art, and so forth. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are incorporated by reference. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

As used herein, the term "about" means plus or minus 10% of the numerical value of the number with which it is being used. Therefore, about 50% means in the range of 45%-55%.

"Administering" when used in conjunction with a therapeutic means to administer a therapeutic directly into or onto a target tissue or to administer a therapeutic to a patient whereby the therapeutic positively impacts the tissue to which it is targeted. Thus, as used herein, the term "administering", can include, but is not limited to, providing an enzyme into the CNS or onto the target tissue; providing an enzyme systemically to a patient by, e.g., intravenous injection whereby the therapeutic reaches the target tissue; providing an enzyme in the form of the encoding sequence thereof to the target tissue (e.g., by so-called gene-therapy techniques). "Administering" a composition may be accomplished by injection, topical administration, or by either method in combination with other known techniques.

The term "animal" as used herein includes, but is not limited to, humans and non-human vertebrates such as wild, domestic and farm animals.

The term "improves" is used to convey that the present invention changes either the appearance, form, characteristics and/or the physical attributes of the target to which it is being provided, applied or administered. The change may be demonstrated by any of the following alone or in combination, including degradation of the CSPGs of the lesioned area of the spinal cord or within the CNS or restoring, in whole or in part, motor, sensory or autonomic function of the mammal.

The term "inhibiting" includes the administration of a compound of the present invention to prevent the onset of the symptoms, alleviating the symptoms, or eliminating the disease, condition or disorder.

By "pharmaceutically acceptable", it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The term "recombinant protein" refers to a polypeptide of the present invention which is produced by recombinant DNA techniques, wherein generally, DNA encoding a polypeptide is inserted into a suitable expression vector which is in turn used to transform a host cell to produce the protein. Moreover, the phrase "derived from", with respect to a recombinant gene, is meant to include within the meaning of "recombinant protein" those proteins having an amino acid sequence of a native protein, or an amino acid sequence similar thereto which is generated by mutations including substitutions and deletions (including truncation) of a naturally occurring form of the protein.

As used herein, the term “therapeutic” means an agent utilized to treat, combat, ameliorate, prevent or improve an unwanted condition or disease of a patient. In part, embodiments of the present invention are directed to the treatment of the central nervous system, such as degradation of the CSPGs of the lesioned area of the spinal cord or within the CNS or restoration, in whole or in part, motor, sensory or autonomic function of the mammal.

The terms “therapeutically effective amount” or “effective amount”, as used herein, may be used interchangeably and refer to an amount of a therapeutic compound component of the present invention. For example, a therapeutically effective amount of a therapeutic compound is a predetermined amount calculated to achieve the desired effect, i.e., to effectively treat an injury to the central nervous system. For example, a therapeutic compound comprising a therapeutically effective amount of chondroitinase which may be purified by a method of the present invention and formulated to provide a stable, active enzyme, is sufficient to degrade the CSPGs of the lesioned area of the spinal cord or an amount sufficient to restore, in whole or in part, motor, sensory or autonomic function of the mammal and may result in a regeneration of neurons in a central nervous system, such as by promoting axonal growth into an injured area.

The terms “treat,” “treated,” or “treating” as used herein refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) an undesired physiological condition, disorder or disease, or to obtain beneficial or desired clinical results. For the purposes of this invention, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms; diminishment of the extent of the condition, disorder or disease; stabilization (i.e., not worsening) of the state of the condition, disorder or disease; delay in onset or slowing of the progression of the condition, disorder or disease; amelioration of the condition, disorder or disease state; and remission (whether partial or total), whether detectable or undetectable, or enhancement or improvement of the condition, disorder or disease. Treatment includes eliciting a clinically significant response without excessive levels of side effects. Treatment also includes prolonging survival as compared to expected survival if not receiving treatment.

The term “vector” refers to a vehicle which can transport the nucleic acid molecules. The nucleic acid molecules encoding the chondroitinase polypeptide are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector can be a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

Chondroitinase may be obtained from a microorganism that naturally expresses a chondroitinase; for example, but not limited to, *E. coli*, *Proteus vulgaris* or from the expression of a recombinant protein in a host cell. The host cell can be a prokaryotic cell (such as *E. coli*) or a eukaryotic cell (such as yeast, a mammalian cell or an insect cell).

The nucleotide sequence of chondroitinase ABCI is set forth as SEQ ID NO. 8 and the amino acid sequence of chondroitinase ABCI is set forth as SEQ ID NO. 9.

One aspect of the present invention provides mutants of chondroitinase ABCI. In a preferred embodiment, the chondroitinase ABCI mutant enzymes are selected from BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), BF4 (SEQ ID NO:3). In another preferred embodiment, the chondroitinase ABCI mutant enzymes are selected from BC9 (SEQ ID NO:4), BC7 (SEQ ID NO:5), RD4 (SEQ ID NO:6) and BE11 (SEQ ID NO: 7).

Such enzymes may be formulated into pharmaceutical compositions and formulations. Suitable stable formulations and methods of purification are set forth in co-pending PCT Application No. US2005/017464 filed May 18, 2005 entitled “Methods of Purifying Chondroitinase and Stable Formulations Thereof” herein incorporated by reference in its entirety.

One aspect of the present invention provides mutants of chondroitinase ABCI. In preferred embodiments, such chondroitinase ABCI mutants exhibit enhanced activity.

In other preferred embodiments, such chondroitinase ABCI mutants exhibit enhanced resistance to inactivation. More preferably, the chondroitinase ABCI mutant enzymes are selected from BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), BF4 (SEQ ID NO:3). In another preferred embodiment, the chondroitinase ABCI mutant enzymes are selected from BC9 (SEQ ID NO:4), BC7 (SEQ ID NO:5), RD4 (SEQ ID NO:6) and BE11 (SEQ ID NO: 7).

Another embodiment of the present invention is a method of designing mutants of chondroitinase ABCI having altered activity. The method comprises altering the nucleotide sequence or amino acid sequence of chondroitinase ABCI, expressing the chondroitinase ABCI in a suitable vector and measuring the activity of the mutant enzyme.

In a further embodiment, a stable chondroitinase ABCI enzyme is provided. The enzyme may exhibit increased resistance to inactivation under stressed conditions, including exposure to UV light or heat. In a preferred embodiment, the enzyme exhibits increased stability compared to wild-type chondroitinase ABCI enzyme following a challenge by a stress.

A further embodiment of the present invention is a method of treating central nervous system injuries comprising administering a chondroitinase ABCI mutant enzyme. In preferred embodiments, the chondroitinase ABCI mutant enzyme is administered in a therapeutically effective amount. In a preferred embodiment, the chondroitinase ABCI mutant enzyme is selected from the group consisting of BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), BF4 (SEQ ID NO:3), BC9 (SEQ ID NO:4), BC7 (SEQ ID NO:5), RD4 (SEQ ID NO:6) and BE11 (SEQ ID NO: 7), more preferably, the enzyme is selected from the group consisting of BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), and BF4 (SEQ ID NO:3). Such central nervous system injuries may include, but are not limited to, spinal cord injuries.

Another embodiment of the present invention is a method promoting neuronal outgrowth comprising administering a chondroitinase ABCI mutant enzyme. In preferred embodiments, the chondroitinase ABCI mutant enzyme is administered in a therapeutically effective amount. In a preferred embodiment, the chondroitinase ABCI mutant enzyme is selected from the group consisting of BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), BF4 (SEQ ID NO:3), BC9 (SEQ ID NO:4), BC7 (SEQ ID NO:5), RD4 (SEQ ID NO:6) and BE11 (SEQ ID NO: 7), more preferably, the enzyme is selected from the group consisting of BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2), and BF4 (SEQ ID NO:3).

Other embodiments of the present invention relate to methods for promoting neurological functional recovery after central nervous system (“CNS”) injury or disease. In preferred embodiments, the chondroitinase ABCI mutant enzyme is administered in a therapeutically effective amount. In particular, the present invention is directed to a method of utilizing chondroitinase to promote sensory, motor or autonomic neurological functional recovery following injury in or to the spinal cord. Compositions useful in this method include acceptable formulations of chondroitinase, more particularly sustained release formulations of chondroitinase. The present

invention is also directed to a method of promoting neurological functional recovery after a contusion injury to the spinal cord. The most common types of spinal cord injuries (SCI) include contusions (bruising of the spinal cord) and compression injuries (caused by pressure on the spinal cord). In contusion injuries, the most common type of injury, a cavity or hole often forms in the center of the spinal cord.

The treatments of the present disclosure deliver an effective amount of the mutant or other optional therapeutic agent to the CNS or the injured site of the CNS. Such methods may include optionally administering other chondroitin sulfate proteoglycans, including, but not limited to chondroitinase ABC_{TypeI}, chondroitinase ABC_{TypeII}, chondroitinase AC and chondroitinase B or mammalian enzymes with chondroitinase-like activity such as Hyal1, Hyal2, Hyal3, and Hyal4, preferably to the CNS, and more preferably to the lesions of the injured area of the CNS.

As is known in the art, chondroitinase polypeptides can be produced by standard biological techniques or by chemical synthesis. For example, a host cell transfected with a nucleic acid vector directing expression of a nucleotide sequence encoding the subject polypeptides can be cultured under appropriate conditions to allow expression of the peptide to occur. The chondroitinase polypeptide may be secreted and isolated and from a mixture of cells and medium containing the recombinant chondroitinase polypeptide. Aspects of the invention described herein provide purification methods wherein the chondroitinase is isolated in a pure form that is more stable and active than those methods currently used.

Alternatively, the peptide may be retained cytoplasmically by removing the signal peptide sequence from the recombinant chondroitinase gene and the cells harvested, lysed and the protein isolated by the purification methods described herein.

Chondroitinase may be administered topically, locally or systemically. Topical or local administration is preferable for greater control of application. The chondroitinases, singularly or in combination, can be mixed with an appropriate pharmaceutical carrier prior to administration. Examples of generally used pharmaceutical carriers and additives are conventional diluents, binders, lubricants, coloring agents, disintegrating agents, buffer agents, isotonicizing fatty acids, isotonicizing agents, preservatives, anesthetics, surfactants and the like, and are known to those skilled in the art. Specifically pharmaceutical carriers that may be used are dextran, sucrose, lactose, maltose, xylose, trehalose, mannitol, xylitol, sorbitol, inositol, serum albumin, gelatin, creatinine, polyethylene glycol, non-ionic surfactants (e.g. polyoxyethylene sorbitan fatty acid esters, polyoxyethylene hardened castor oil, sucrose fatty acid esters, polyoxyethylene polyoxypropylene glycol) and similar compounds. Pharmaceutical carriers may also be used in combination, such as polyethylene glycol and/or sucrose, or polyoxyethylene sorbitan fatty acid esters, polyoxyethylene sorbitan monooleate (20 E. 0.) is particularly preferred.

A treatment regimen according to the invention may be carried out by a means of administering a mutant chondroitinase ABCI enzyme of the present invention. The treatment regimen may further comprise administering chondroitinase ABCII, chondroitinase AC and chondroitinase B or mammalian enzymes with chondroitinase-like activity such as Hyal1, Hyal2, Hyal3, Hyal4 and PH2O to the lesions of the injured area of the CNS. The mode of administration, the timing of administration and the dosage are carried out such that the functional recovery from impairment of the CNS is enhanced by the promotion of neurite outgrowth.

The effective amount of chondroitinase can be administered in a single dosage, two dosages or a plurality of dosages. Although it is to be understood that the dosage may be administered at any time, in one embodiment, the dosage is administered within 12 hours after injury, or as soon as is feasible. In another embodiment, the dosage is administered to an injured mammal in one, two or a plurality of dosages; such dosages would be dependant on the severity of the injury and the amount of CSPGs present in the glial scarring. Where a plurality of dosages is administered, they may be delivered on a daily, weekly, or bi-weekly basis. The delivery of the dosages may be by means of catheter or syringe. Alternatively, the treatment can be administered during surgery to allow direct application to the glial scar.

For example, in some aspects, the invention is directed to a pharmaceutical composition comprising a compound, as defined above, and a pharmaceutically acceptable carrier or diluent, or an effective amount of a pharmaceutical composition comprising a compound as defined above.

The compounds of the present invention can be administered in the conventional manner by any route where they are active. Administration can be systemic, topical, or oral. For example, administration can be, but is not limited to, parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, oral, buccal, or ocular routes, or intravaginally, by inhalation, by depot injections, or by implants. Thus, modes of administration for the compounds of the present invention (either alone or in combination with other pharmaceuticals) can be, but are not limited to, sublingual, injectable (including short-acting, depot, implant and pellet forms injected subcutaneously or intramuscularly), or by use of vaginal creams, suppositories, pessaries, vaginal rings, rectal suppositories, intrauterine devices, and transdermal forms such as patches and creams.

Specific modes of administration will depend on the indication. The selection of the specific route of administration and the dose regimen is to be adjusted or titrated by the clinician according to methods known to the clinician in order to obtain the optimal clinical response. The amount of compound to be administered is that amount which is therapeutically effective. The dosage to be administered will depend on the characteristics of the subject being treated, e.g., the particular animal treated, age, weight, health, types of concurrent treatment, if any, and frequency of treatments, and can be easily determined by one of skill in the art (e.g., by the clinician).

Pharmaceutical formulations containing the compounds of the present invention and a suitable carrier can be solid dosage forms which include, but are not limited to, tablets, capsules, cachets, pellets, pills, powders and granules; topical dosage forms which include, but are not limited to, solutions, powders, fluid emulsions, fluid suspensions, semi-solids, ointments, pastes, creams, gels and jellies, and foams; and parenteral dosage forms which include, but are not limited to, solutions, suspensions, emulsions, and dry powder; comprising an effective amount of a polymer or copolymer of the present invention. It is also known in the art that the active ingredients can be contained in such formulations with pharmaceutically acceptable diluents, fillers, disintegrants, binders, lubricants, surfactants, hydrophobic vehicles, water soluble vehicles, emulsifiers, buffers, humectants, moisturizers, solubilizers, preservatives and the like. The means and methods for administration are known in the art and an artisan can refer to various pharmacologic references for guidance. For example, *Modern Pharmaceutics*, Banker & Rhodes, Marcel Dekker, Inc. (1979); and *Goodman & Gilman's The*

Pharmaceutical Basis of Therapeutics, 6th Edition, Mac-Millan Publishing Co., New York (1980) can be consulted.

The compounds of the present invention can be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. The compounds can be administered by continuous infusion subcutaneously over a period of about 15 minutes to about 24 hours. Formulations for injection can be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

For oral administration, the compounds can be formulated readily by combining these compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by adding a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, but are not limited to, fillers such as sugars, including, but not limited to, lactose, sucrose, mannitol, and sorbitol; cellulose preparations such as, but not limited to, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and polyvinylpyrrolidone (PVP). If desired, disintegrating agents can be added, such as, but not limited to, the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores can be provided with suitable coatings. For this purpose, concentrated sugar solutions can be used, which can optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include, but are not limited to, push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as, e.g., lactose, binders such as, e.g., starches, and/or lubricants such as, e.g., talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers can be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions can take the form of, e.g., tablets or lozenges formulated in a conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or

insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds of the present invention can also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds of the present invention can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection.

Depot injections can be administered at about 1 to about 6 months or longer intervals. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

In transdermal administration, the compounds of the present invention, for example, can be applied to a plaster, or can be applied by transdermal, therapeutic systems that are consequently supplied to the organism.

Pharmaceutical compositions of the compounds also can comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as, e.g., polyethylene glycols.

The compounds of the present invention can also be administered in combination with other active ingredients, such as, for example, adjuvants, protease inhibitors, or other compatible drugs or compounds where such combination is seen to be desirable or advantageous in achieving the desired effects of the methods described herein.

The following methods are used to illustrate the various embodiments of the present invention. The methods are exemplary methods and are not meant to limit the invention.

Example 1

The present example illustrates exemplary chondroitinase mutant enzymes of the present invention. All nucleotide and amino acids are indicated as the wild-type and then the mutant version (Wild-type to Mutant).

Mutant ABCI enzyme	Nucleotide sequence	Amino Acid sequence
BC6 (SEQ ID NO. 1)	T1206 to C1206 C1114 to A1114	E403 to G403 W372 to C372
BE7 (SEQ ID NO. 2)	G1925 to T1925 T2226 to G2226	S642 to I642 I742 to M742
BF4 (SEQ ID NO. 3)	T2160 to A2160	N720 to K720e
BC9 (SEQ ID NO. 4)	G1238 to A1238	S413 to N413
BC7 (SEQ ID NO. 5)	A1468 to G1468	K490 to E490
RD4 (SEQ ID NO. 6)	T1661 to A1661	L554 to H554
BE11 (SEQ ID NO. 7)	A1901 to T1901 C1935 to T1935 (in wobble position of codon- does not result in AA change)	D634 to V634

Example 2

The present example illustrates the chondroitin lyase activity of exemplary chondroitinase ABCI mutants according to the present invention following UV exposure. Mutant chondroitinase ABCI genes were generated and transformed into

11

bacteria. Bacteria were grown and the mutagenized chondroitinase expressed. The chondroitinase were then exposed to UV light and their chondroitin lyase activity measured. As depicted in FIG. 1, clone BC6 (SEQ ID NO:1), BE7 (SEQ ID NO:2) and BF4 (SEQ ID NO:3) exhibited greater chondroitin lyase activity following exposure to UV light as compared to control.

Example 3

The present example illustrates the chondroitin lyase activity of exemplary chondroitinase ABCI mutants according to

12

the present invention. The chondroitinase lyase activity of Clone BC9, Clone BC7, Clone RD4 and Clone BE11 under normal (i.e., non-stressed) conditions was measured and exhibited decreased activity as compared to control an wild-type chondroitinase ABCI, as depicted in FIG. 2.

Although the present invention has been described in considerable detail with reference to certain preferred embodiments thereof, other versions are possible. Therefore the spirit and scope of the appended claims should not be limited to the description and the preferred versions contained within this specification.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1

<211> LENGTH: 1027

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Chondroitinase ABCI

<400> SEQUENCE: 1

```

Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
1           5           10           15
Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
20           25           30
Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
35           40           45
Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
50           55           60
Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
65           70           75           80
Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
85           90           95
Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
100          105          110
Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
115          120          125
Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
130          135          140
Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
145          150          155          160
Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
165          170          175
Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
180          185          190
Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
195          200          205
Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
210          215          220
Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
225          230          235          240
Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
245          250          255
Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
260          265          270

```

-continued

Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
 275 280 285
 Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
 290 295 300
 Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
 305 310 315 320
 Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
 325 330 335
 Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
 340 345 350
 Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
 355 360 365
 Ser Ser Arg Cys Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380
 Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400
 Ser Arg Gly Phe Lys Ser Ser Phe Asp Met Lys Val Ser Ala Asp Ser
 405 410 415
 Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430
 Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445
 Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460
 Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480
 Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Lys Asn Ala Ser Gln Leu Ile
 485 490 495
 Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510
 Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525
 Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu
 530 535 540
 Lys Ser Val Ala Gln Gly Tyr Tyr Trp Leu Ala Met Ser Ala Lys Ser
 545 550 555 560
 Ser Pro Asp Lys Thr Leu Ala Ser Ile Tyr Leu Ala Ile Ser Asp Lys
 565 570 575
 Thr Gln Asn Glu Ser Thr Ala Ile Phe Gly Glu Thr Ile Thr Pro Ala
 580 585 590
 Ser Leu Pro Gln Gly Phe Tyr Ala Phe Asn Gly Gly Ala Phe Gly Ile
 595 600 605
 His Arg Trp Gln Asp Lys Met Val Thr Leu Lys Ala Tyr Asn Thr Asn
 610 615 620
 Val Trp Ser Ser Glu Ile Tyr Asn Lys Asp Asn Arg Tyr Gly Arg Tyr
 625 630 635 640
 Gln Ser His Gly Val Ala Gln Ile Val Ser Asn Gly Ser Gln Leu Ser
 645 650 655
 Gln Gly Tyr Gln Gln Glu Gly Trp Asp Trp Asn Arg Met Pro Gly Ala
 660 665 670
 Thr Thr Ile His Leu Pro Leu Lys Asp Leu Asp Ser Pro Lys Pro His
 675 680 685
 Thr Leu Met Gln Arg Gly Glu Arg Gly Phe Ser Gly Thr Ser Ser Leu

-continued

690			695			700									
Glu	Gly	Gln	Tyr	Gly	Met	Met	Ala	Phe	Asp	Leu	Ile	Tyr	Pro	Ala	Asn
705					710					715					720
Leu	Glu	Arg	Phe	Asp	Pro	Asn	Phe	Thr	Ala	Lys	Lys	Ser	Val	Leu	Ala
				725					730						735
Ala	Asp	Asn	His	Leu	Ile	Phe	Ile	Gly	Ser	Asn	Ile	Asn	Ser	Ser	Asp
			740					745						750	
Lys	Asn	Lys	Asn	Val	Glu	Thr	Thr	Leu	Phe	Gln	His	Ala	Ile	Thr	Pro
		755					760					765			
Thr	Leu	Asn	Thr	Leu	Trp	Ile	Asn	Gly	Gln	Lys	Ile	Glu	Asn	Met	Pro
	770					775						780			
Tyr	Gln	Thr	Thr	Leu	Gln	Gln	Gly	Asp	Trp	Leu	Ile	Asp	Ser	Asn	Gly
785					790					795					800
Asn	Gly	Tyr	Leu	Ile	Thr	Gln	Ala	Glu	Lys	Val	Asn	Val	Ser	Arg	Gln
				805					810						815
His	Gln	Val	Ser	Ala	Glu	Asn	Lys	Asn	Arg	Gln	Pro	Thr	Glu	Gly	Asn
			820						825					830	
Phe	Ser	Ser	Ala	Trp	Ile	Asp	His	Ser	Thr	Arg	Pro	Lys	Asp	Ala	Ser
		835					840						845		
Tyr	Glu	Tyr	Met	Val	Phe	Leu	Asp	Ala	Thr	Pro	Glu	Lys	Met	Gly	Glu
	850					855					860				
Met	Ala	Gln	Lys	Phe	Arg	Glu	Asn	Asn	Gly	Leu	Tyr	Gln	Val	Leu	Arg
865					870					875					880
Lys	Asp	Lys	Asp	Val	His	Ile	Ile	Leu	Asp	Lys	Leu	Ser	Asn	Val	Thr
				885					890						895
Gly	Tyr	Ala	Phe	Tyr	Gln	Pro	Ala	Ser	Ile	Glu	Asp	Lys	Trp	Ile	Lys
			900						905						910
Lys	Val	Asn	Lys	Pro	Ala	Ile	Val	Met	Thr	His	Arg	Gln	Lys	Asp	Thr
		915					920							925	
Leu	Ile	Val	Ser	Ala	Val	Thr	Pro	Asp	Leu	Asn	Met	Thr	Arg	Gln	Lys
	930					935						940			
Ala	Ala	Thr	Pro	Val	Thr	Ile	Asn	Val	Thr	Ile	Asn	Gly	Lys	Trp	Gln
945					950					955					960
Ser	Ala	Asp	Lys	Asn	Ser	Glu	Val	Lys	Tyr	Gln	Val	Ser	Gly	Asp	Asn
				965						970					975
Thr	Glu	Leu	Thr	Phe	Thr	Ser	Tyr	Phe	Gly	Ile	Pro	Gln	Glu	Ile	Lys
			980					985							990
Leu	Ser	Pro	Leu	Pro	Ala	Cys	Arg	Asp	Ala	Thr	His	Glu	Arg	Ala	Pro
		995					1000							1005	
Glu	Thr	Ile	Cys	Ser	Ile	Asn	Cys	Cys	Asn	Phe	Ile	Asp	Glu	Asn	
	1010					1015						1020			
Thr	Ile	Ala	Leu												
	1025														

<210> SEQ ID NO 2

<211> LENGTH: 1027

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Chondroitinase ABCI

<400> SEQUENCE: 2

Ala	Thr	Ser	Asn	Pro	Ala	Phe	Asp	Pro	Lys	Asn	Leu	Met	Gln	Ser	Glu
1			5						10					15	

Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp

-continued

20			25			30									
Lys	Asn	Ser	Ile	Leu	Thr	Leu	Ser	Asp	Lys	Arg	Ser	Ile	Met	Gly	Asn
	35						40					45			
Gln	Ser	Leu	Leu	Trp	Lys	Trp	Lys	Gly	Gly	Ser	Ser	Phe	Thr	Leu	His
	50					55						60			
Lys	Lys	Leu	Ile	Val	Pro	Thr	Asp	Lys	Glu	Ala	Ser	Lys	Ala	Trp	Gly
65					70					75					80
Arg	Ser	Ser	Thr	Pro	Val	Phe	Ser	Phe	Trp	Leu	Tyr	Asn	Glu	Lys	Pro
			85						90						95
Ile	Asp	Gly	Tyr	Leu	Thr	Ile	Asp	Phe	Gly	Glu	Lys	Leu	Ile	Ser	Thr
			100						105						110
Ser	Glu	Ala	Gln	Ala	Gly	Phe	Lys	Val	Lys	Leu	Asp	Phe	Thr	Gly	Trp
		115					120					125			
Arg	Thr	Val	Gly	Val	Ser	Leu	Asn	Asn	Asp	Leu	Glu	Asn	Arg	Glu	Met
	130					135						140			
Thr	Leu	Asn	Ala	Thr	Asn	Thr	Ser	Ser	Asp	Gly	Thr	Gln	Asp	Ser	Ile
145					150					155					160
Gly	Arg	Ser	Leu	Gly	Ala	Lys	Val	Asp	Ser	Ile	Arg	Phe	Lys	Ala	Pro
			165						170						175
Ser	Asn	Val	Ser	Gln	Gly	Glu	Ile	Tyr	Ile	Asp	Arg	Ile	Met	Phe	Ser
		180							185						190
Val	Asp	Asp	Ala	Arg	Tyr	Gln	Trp	Ser	Asp	Tyr	Gln	Val	Lys	Thr	Arg
	195						200					205			
Leu	Ser	Glu	Pro	Glu	Ile	Gln	Phe	His	Asn	Val	Lys	Pro	Gln	Leu	Pro
	210					215						220			
Val	Thr	Pro	Glu	Asn	Leu	Ala	Ala	Ile	Asp	Leu	Ile	Arg	Gln	Arg	Leu
225					230					235					240
Ile	Asn	Glu	Phe	Val	Gly	Gly	Glu	Lys	Glu	Thr	Asn	Leu	Ala	Leu	Glu
			245						250						255
Glu	Asn	Ile	Ser	Lys	Leu	Lys	Ser	Asp	Phe	Asp	Ala	Leu	Asn	Thr	His
		260							265						270
Thr	Leu	Ala	Asn	Gly	Gly	Thr	Gln	Gly	Arg	His	Leu	Ile	Thr	Asp	Lys
	275						280						285		
Gln	Ile	Ile	Ile	Tyr	Gln	Pro	Glu	Asn	Leu	Asn	Ser	Gln	Asp	Lys	Gln
	290					295						300			
Leu	Phe	Asp	Asn	Tyr	Val	Ile	Leu	Gly	Asn	Tyr	Thr	Thr	Leu	Met	Phe
305					310					315					320
Asn	Ile	Ser	Arg	Ala	Tyr	Val	Leu	Glu	Lys	Asp	Pro	Thr	Gln	Lys	Ala
			325							330					335
Gln	Leu	Lys	Gln	Met	Tyr	Leu	Leu	Met	Thr	Lys	His	Leu	Leu	Asp	Gln
			340						345						350
Gly	Phe	Val	Lys	Gly	Ser	Ala	Leu	Val	Thr	Thr	His	His	Trp	Gly	Tyr
	355						360						365		
Ser	Ser	Arg	Trp	Trp	Tyr	Ile	Ser	Thr	Leu	Leu	Met	Ser	Asp	Ala	Leu
	370					375						380			
Lys	Glu	Ala	Asn	Leu	Gln	Thr	Gln	Val	Tyr	Asp	Ser	Leu	Leu	Trp	Tyr
385					390					395					400
Ser	Arg	Glu	Phe	Lys	Ser	Ser	Phe	Asp	Met	Lys	Val	Ser	Ala	Asp	Ser
			405							410					415
Ser	Asp	Leu	Asp	Tyr	Phe	Asn	Thr	Leu	Ser	Arg	Gln	His	Leu	Ala	Leu
		420							425						430
Leu	Leu	Leu	Glu	Pro	Asp	Asp	Gln	Lys	Arg	Ile	Asn	Leu	Val	Asn	Thr
		435							440						445

-continued

Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460

Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480

Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Lys Asn Ala Ser Gln Leu Ile
 485 490 495

Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510

Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525

Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu
 530 535 540

Lys Ser Val Ala Gln Gly Tyr Tyr Trp Leu Ala Met Ser Ala Lys Ser
 545 550 555 560

Ser Pro Asp Lys Thr Leu Ala Ser Ile Tyr Leu Ala Ile Ser Asp Lys
 565 570 575

Thr Gln Asn Glu Ser Thr Ala Ile Phe Gly Glu Thr Ile Thr Pro Ala
 580 585 590

Ser Leu Pro Gln Gly Phe Tyr Ala Phe Asn Gly Gly Ala Phe Gly Ile
 595 600 605

His Arg Trp Gln Asp Lys Met Val Thr Leu Lys Ala Tyr Asn Thr Asn
 610 615 620

Val Trp Ser Ser Glu Ile Tyr Asn Lys Asp Asn Arg Tyr Gly Arg Tyr
 625 630 635 640

Gln Ile His Gly Val Ala Gln Ile Val Ser Asn Gly Ser Gln Leu Ser
 645 650 655

Gln Gly Tyr Gln Gln Glu Gly Trp Asp Trp Asn Arg Met Pro Gly Ala
 660 665 670

Thr Thr Ile His Leu Pro Leu Lys Asp Leu Asp Ser Pro Lys Pro His
 675 680 685

Thr Leu Met Gln Arg Gly Glu Arg Gly Phe Ser Gly Thr Ser Ser Leu
 690 695 700

Glu Gly Gln Tyr Gly Met Met Ala Phe Asp Leu Ile Tyr Pro Ala Asn
 705 710 715 720

Leu Glu Arg Phe Asp Pro Asn Phe Thr Ala Lys Lys Ser Val Leu Ala
 725 730 735

Ala Asp Asn His Leu Met Phe Ile Gly Ser Asn Ile Asn Ser Ser Asp
 740 745 750

Lys Asn Lys Asn Val Glu Thr Thr Leu Phe Gln His Ala Ile Thr Pro
 755 760 765

Thr Leu Asn Thr Leu Trp Ile Asn Gly Gln Lys Ile Glu Asn Met Pro
 770 775 780

Tyr Gln Thr Thr Leu Gln Gln Gly Asp Trp Leu Ile Asp Ser Asn Gly
 785 790 795 800

Asn Gly Tyr Leu Ile Thr Gln Ala Glu Lys Val Asn Val Ser Arg Gln
 805 810 815

His Gln Val Ser Ala Glu Asn Lys Asn Arg Gln Pro Thr Glu Gly Asn
 820 825 830

Phe Ser Ser Ala Trp Ile Asp His Ser Thr Arg Pro Lys Asp Ala Ser
 835 840 845

Tyr Glu Tyr Met Val Phe Leu Asp Ala Thr Pro Glu Lys Met Gly Glu
 850 855 860

-continued

Met Ala Gln Lys Phe Arg Glu Asn Asn Gly Leu Tyr Gln Val Leu Arg
865 870 875 880

Lys Asp Lys Asp Val His Ile Ile Leu Asp Lys Leu Ser Asn Val Thr
885 890 895

Gly Tyr Ala Phe Tyr Gln Pro Ala Ser Ile Glu Asp Lys Trp Ile Lys
900 905 910

Lys Val Asn Lys Pro Ala Ile Val Met Thr His Arg Gln Lys Asp Thr
915 920 925

Leu Ile Val Ser Ala Val Thr Pro Asp Leu Asn Met Thr Arg Gln Lys
930 935 940

Ala Ala Thr Pro Val Thr Ile Asn Val Thr Ile Asn Gly Lys Trp Gln
945 950 955 960

Ser Ala Asp Lys Asn Ser Glu Val Lys Tyr Gln Val Ser Gly Asp Asn
965 970 975

Thr Glu Leu Thr Phe Thr Ser Tyr Phe Gly Ile Pro Gln Glu Ile Lys
980 985 990

Leu Ser Pro Leu Pro Ala Cys Arg Asp Ala Thr His Glu Arg Ala Pro
995 1000 1005

Glu Thr Ile Cys Ser Ile Asn Cys Cys Asn Phe Ile Asp Glu Asn
1010 1015 1020

Thr Ile Ala Leu
1025

<210> SEQ ID NO 3
 <211> LENGTH: 1027
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chondroitinase ABCI

<400> SEQUENCE: 3

Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
1 5 10 15

Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
20 25 30

Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
35 40 45

Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
50 55 60

Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
65 70 75 80

Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
85 90 95

Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
100 105 110

Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
115 120 125

Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
130 135 140

Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
145 150 155 160

Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
165 170 175

Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
180 185 190

-continued

Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
 195 200 205
 Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
 210 215 220
 Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
 225 230 235 240
 Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
 245 250 255
 Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
 260 265 270
 Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
 275 280 285
 Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
 290 295 300
 Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
 305 310 315 320
 Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
 325 330 335
 Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
 340 345 350
 Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
 355 360 365
 Ser Ser Arg Trp Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380
 Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400
 Ser Arg Glu Phe Lys Ser Ser Phe Asp Met Lys Val Ser Ala Asp Ser
 405 410 415
 Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430
 Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445
 Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460
 Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480
 Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Lys Asn Ala Ser Gln Leu Ile
 485 490 495
 Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510
 Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525
 Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu
 530 535 540
 Lys Ser Val Ala Gln Gly Tyr Tyr Trp Leu Ala Met Ser Ala Lys Ser
 545 550 555 560
 Ser Pro Asp Lys Thr Leu Ala Ser Ile Tyr Leu Ala Ile Ser Asp Lys
 565 570 575
 Thr Gln Asn Glu Ser Thr Ala Ile Phe Gly Glu Thr Ile Thr Pro Ala
 580 585 590
 Ser Leu Pro Gln Gly Phe Tyr Ala Phe Asn Gly Gly Ala Phe Gly Ile
 595 600 605
 His Arg Trp Gln Asp Lys Met Val Thr Leu Lys Ala Tyr Asn Thr Asn

-continued

```

<210> SEQ ID NO 4
<211> LENGTH: 1027
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: CHONDROITINASE ABCI

<400> SEQUENCE: 4
Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
1          5          10          15
Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
20          25          30
Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
35          40          45
Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
50          55          60
Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
65          70          75          80
Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
85          90          95
Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
100         105         110
Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
115         120         125
Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
130         135         140
Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
145         150         155         160
Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
165         170         175
Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
180         185         190
Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
195         200         205
Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
210         215         220
Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
225         230         235         240
Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
245         250         255
Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
260         265         270
Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
275         280         285
Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
290         295         300
Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
305         310         315         320
Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
325         330         335
Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
340         345         350
Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
355         360         365

```

-continued

Ser Ser Arg Trp Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380

Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400

Ser Arg Glu Phe Lys Ser Ser Phe Asp Met Lys Val Asn Ala Asp Ser
 405 410 415

Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430

Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445

Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460

Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480

Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Lys Asn Ala Ser Gln Leu Ile
 485 490 495

Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510

Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525

Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu
 530 535 540

Lys Ser Val Ala Gln Gly Tyr Tyr Trp Leu Ala Met Ser Ala Lys Ser
 545 550 555 560

Ser Pro Asp Lys Thr Leu Ala Ser Ile Tyr Leu Ala Ile Ser Asp Lys
 565 570 575

Thr Gln Asn Glu Ser Thr Ala Ile Phe Gly Glu Thr Ile Thr Pro Ala
 580 585 590

Ser Leu Pro Gln Gly Phe Tyr Ala Phe Asn Gly Gly Ala Phe Gly Ile
 595 600 605

His Arg Trp Gln Asp Lys Met Val Thr Leu Lys Ala Tyr Asn Thr Asn
 610 615 620

Val Trp Ser Ser Glu Ile Tyr Asn Lys Asp Asn Arg Tyr Gly Arg Tyr
 625 630 635 640

Gln Ser His Gly Val Ala Gln Ile Val Ser Asn Gly Ser Gln Leu Ser
 645 650 655

Gln Gly Tyr Gln Gln Glu Gly Trp Asp Trp Asn Arg Met Pro Gly Ala
 660 665 670

Thr Thr Ile His Leu Pro Leu Lys Asp Leu Asp Ser Pro Lys Pro His
 675 680 685

Thr Leu Met Gln Arg Gly Glu Arg Gly Phe Ser Gly Thr Ser Ser Leu
 690 695 700

Glu Gly Gln Tyr Gly Met Met Ala Phe Asp Leu Ile Tyr Pro Ala Asn
 705 710 715 720

Leu Glu Arg Phe Asp Pro Asn Phe Thr Ala Lys Lys Ser Val Leu Ala
 725 730 735

Ala Asp Asn His Leu Ile Phe Ile Gly Ser Asn Ile Asn Ser Ser Asp
 740 745 750

Lys Asn Lys Asn Val Glu Thr Thr Leu Phe Gln His Ala Ile Thr Pro
 755 760 765

Thr Leu Asn Thr Leu Trp Ile Asn Gly Gln Lys Ile Glu Asn Met Pro
 770 775 780

-continued

Tyr Gln Thr Thr Leu Gln Gln Gly Asp Trp Leu Ile Asp Ser Asn Gly
 785 790 795 800
 Asn Gly Tyr Leu Ile Thr Gln Ala Glu Lys Val Asn Val Ser Arg Gln
 805 810 815
 His Gln Val Ser Ala Trp Ile Asp His Ser Thr Arg Pro Thr Glu Gly Asn
 820 825 830
 Phe Ser Ser Ala Trp Ile Asp His Ser Thr Arg Pro Lys Asp Ala Ser
 835 840 845
 Tyr Glu Tyr Met Val Phe Leu Asp Ala Thr Pro Glu Lys Met Gly Glu
 850 855 860
 Met Ala Gln Lys Phe Arg Glu Asn Asn Gly Leu Tyr Gln Val Leu Arg
 865 870 875 880
 Lys Asp Lys Asp Val His Ile Ile Leu Asp Lys Leu Ser Asn Val Thr
 885 890 895
 Gly Tyr Ala Phe Tyr Gln Pro Ala Ser Ile Glu Asp Lys Trp Ile Lys
 900 905 910
 Lys Val Asn Lys Pro Ala Ile Val Met Thr His Arg Gln Lys Asp Thr
 915 920 925
 Leu Ile Val Ser Ala Val Thr Pro Asp Leu Asn Met Thr Arg Gln Lys
 930 935 940
 Ala Ala Thr Pro Val Thr Ile Asn Val Thr Ile Asn Gly Lys Trp Gln
 945 950 955 960
 Ser Ala Asp Lys Asn Ser Glu Val Lys Tyr Gln Val Ser Gly Asp Asn
 965 970 975
 Thr Glu Leu Thr Phe Thr Ser Tyr Phe Gly Ile Pro Gln Glu Ile Lys
 980 985 990
 Leu Ser Pro Leu Pro Ala Cys Arg Asp Ala Thr His Glu Arg Ala Pro
 995 1000 1005
 Glu Thr Ile Cys Ser Ile Asn Cys Cys Asn Phe Ile Asp Glu Asn
 1010 1015 1020
 Thr Ile Ala Leu
 1025

 <210> SEQ ID NO 5
 <211> LENGTH: 1027
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: CHONDROITINASE ABCI

 <400> SEQUENCE: 5
 Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
 1 5 10 15
 Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
 20 25 30
 Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
 35 40 45
 Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
 50 55 60
 Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
 65 70 75 80
 Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
 85 90 95
 Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
 100 105 110

-continued

Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
 115 120 125
 Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
 130 135 140
 Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
 145 150 155 160
 Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
 165 170 175
 Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
 180 185 190
 Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
 195 200 205
 Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
 210 215 220
 Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
 225 230 235 240
 Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
 245 250 255
 Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
 260 265 270
 Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
 275 280 285
 Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
 290 295 300
 Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
 305 310 315 320
 Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
 325 330 335
 Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
 340 345 350
 Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
 355 360 365
 Ser Ser Arg Trp Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380
 Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400
 Ser Arg Glu Phe Lys Ser Ser Phe Asp Met Lys Val Ser Ala Asp Ser
 405 410 415
 Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430
 Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445
 Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460
 Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480
 Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Glu Asn Ala Ser Gln Leu Ile
 485 490 495
 Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510
 Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525
 Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu

-continued

530			535			540									
Lys	Ser	Val	Ala	Gln	Gly	Tyr	Trp	Trp	Leu	Ala	Met	Ser	Ala	Lys	Ser
545					550					555					560
Ser	Pro	Asp	Lys	Thr	Leu	Ala	Ser	Ile	Tyr	Leu	Ala	Ile	Ser	Asp	Lys
			565						570					575	
Thr	Gln	Asn	Glu	Ser	Thr	Ala	Ile	Phe	Gly	Glu	Thr	Ile	Thr	Pro	Ala
			580					585						590	
Ser	Leu	Pro	Gln	Gly	Phe	Tyr	Ala	Phe	Asn	Gly	Gly	Ala	Phe	Gly	Ile
		595					600					605			
His	Arg	Trp	Gln	Asp	Lys	Met	Val	Thr	Leu	Lys	Ala	Tyr	Asn	Thr	Asn
610						615					620				
Val	Trp	Ser	Ser	Glu	Ile	Tyr	Asn	Lys	Asp	Asn	Arg	Tyr	Gly	Arg	Tyr
625				630						635					640
Gln	Ser	His	Gly	Val	Ala	Gln	Ile	Val	Ser	Asn	Gly	Ser	Gln	Leu	Ser
			645						650					655	
Gln	Gly	Tyr	Gln	Gln	Glu	Gly	Trp	Asp	Trp	Asn	Arg	Met	Pro	Gly	Ala
			660					665					670		
Thr	Thr	Ile	His	Leu	Pro	Leu	Lys	Asp	Leu	Asp	Ser	Pro	Lys	Pro	His
		675					680					685			
Thr	Leu	Met	Gln	Arg	Gly	Glu	Arg	Gly	Phe	Ser	Gly	Thr	Ser	Ser	Leu
	690					695					700				
Glu	Gly	Gln	Tyr	Gly	Met	Met	Ala	Phe	Asp	Leu	Ile	Tyr	Pro	Ala	Asn
705				710						715					720
Leu	Glu	Arg	Phe	Asp	Pro	Asn	Phe	Thr	Ala	Lys	Lys	Ser	Val	Leu	Ala
			725						730					735	
Ala	Asp	Asn	His	Leu	Ile	Phe	Ile	Gly	Ser	Asn	Ile	Asn	Ser	Ser	Asp
			740					745					750		
Lys	Asn	Lys	Asn	Val	Glu	Thr	Thr	Leu	Phe	Gln	His	Ala	Ile	Thr	Pro
		755					760					765			
Thr	Leu	Asn	Thr	Leu	Trp	Ile	Asn	Gly	Gln	Lys	Ile	Glu	Asn	Met	Pro
	770					775						780			
Tyr	Gln	Thr	Thr	Leu	Gln	Gln	Gly	Asp	Trp	Leu	Ile	Asp	Ser	Asn	Gly
785					790					795					800
Asn	Gly	Tyr	Leu	Ile	Thr	Gln	Ala	Glu	Lys	Val	Asn	Val	Ser	Arg	Gln
			805						810					815	
His	Gln	Val	Ser	Ala	Glu	Asn	Lys	Asn	Arg	Gln	Pro	Thr	Glu	Gly	Asn
		820						825						830	
Phe	Ser	Ser	Ala	Trp	Ile	Asp	His	Ser	Thr	Arg	Pro	Lys	Asp	Ala	Ser
		835					840					845			
Tyr	Glu	Tyr	Met	Val	Phe	Leu	Asp	Ala	Thr	Pro	Glu	Lys	Met	Gly	Glu
	850					855					860				
Met	Ala	Gln	Lys	Phe	Arg	Glu	Asn	Asn	Gly	Leu	Tyr	Gln	Val	Leu	Arg
865				870						875					880
Lys	Asp	Lys	Asp	Val	His	Ile	Ile	Leu	Asp	Lys	Leu	Ser	Asn	Val	Thr
			885						890					895	
Gly	Tyr	Ala	Phe	Tyr	Gln	Pro	Ala	Ser	Ile	Glu	Asp	Lys	Trp	Ile	Lys
		900						905						910	
Lys	Val	Asn	Lys	Pro	Ala	Ile	Val	Met	Thr	His	Arg	Gln	Lys	Asp	Thr
		915						920						925	
Leu	Ile	Val	Ser	Ala	Val	Thr	Pro	Asp	Leu	Asn	Met	Thr	Arg	Gln	Lys
	930					935					940				
Ala	Ala	Thr	Pro	Val	Thr	Ile	Asn	Val	Thr	Ile	Asn	Gly	Lys	Trp	Gln
945				950					955						960

-continued

Ser Ala Asp Lys Asn Ser Glu Val Lys Tyr Gln Val Ser Gly Asp Asn
 965 970 975

Thr Glu Leu Thr Phe Thr Ser Tyr Phe Gly Ile Pro Gln Glu Ile Lys
 980 985 990

Leu Ser Pro Leu Pro Ala Cys Arg Asp Ala Thr His Glu Arg Ala Pro
 995 1000 1005

Glu Thr Ile Cys Ser Ile Asn Cys Cys Asn Phe Ile Asp Glu Asn
 1010 1015 1020

Thr Ile Ala Leu
 1025

<210> SEQ ID NO 6
 <211> LENGTH: 1027
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: CHONDROITINASE ABCI

<400> SEQUENCE: 6

Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
 1 5 10 15

Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
 20 25 30

Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
 35 40 45

Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
 50 55 60

Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
 65 70 75 80

Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
 85 90 95

Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
 100 105 110

Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
 115 120 125

Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
 130 135 140

Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
 145 150 155 160

Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
 165 170 175

Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
 180 185 190

Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
 195 200 205

Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
 210 215 220

Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
 225 230 235 240

Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
 245 250 255

Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
 260 265 270

Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
 275 280 285

-continued

Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
 290 295 300
 Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
 305 310 315 320
 Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
 325 330 335
 Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
 340 345 350
 Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
 355 360 365
 Ser Ser Arg Trp Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380
 Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400
 Ser Arg Glu Phe Lys Ser Ser Phe Asp Met Lys Val Ser Ala Asp Ser
 405 410 415
 Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430
 Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445
 Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460
 Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480
 Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Lys Asn Ala Ser Gln Leu Ile
 485 490 495
 Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510
 Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525
 Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu
 530 535 540
 Lys Ser Val Ala Gln Gly Tyr Tyr Trp His Ala Met Ser Ala Lys Ser
 545 550 555 560
 Ser Pro Asp Lys Thr Leu Ala Ser Ile Tyr Leu Ala Ile Ser Asp Lys
 565 570 575
 Thr Gln Asn Glu Ser Thr Ala Ile Phe Gly Glu Thr Ile Thr Pro Ala
 580 585 590
 Ser Leu Pro Gln Gly Phe Tyr Ala Phe Asn Gly Gly Ala Phe Gly Ile
 595 600 605
 His Arg Trp Gln Asp Lys Met Val Thr Leu Lys Ala Tyr Asn Thr Asn
 610 615 620
 Val Trp Ser Ser Glu Ile Tyr Asn Lys Asp Asn Arg Tyr Gly Arg Tyr
 625 630 635 640
 Gln Ser His Gly Val Ala Gln Ile Val Ser Asn Gly Ser Gln Leu Ser
 645 650 655
 Gln Gly Tyr Gln Gln Glu Gly Trp Asp Trp Asn Arg Met Pro Gly Ala
 660 665 670
 Thr Thr Ile His Leu Pro Leu Lys Asp Leu Asp Ser Pro Lys Pro His
 675 680 685
 Thr Leu Met Gln Arg Gly Glu Arg Gly Phe Ser Gly Thr Ser Ser Leu
 690 695 700

-continued

Glu Gly Gln Tyr Gly Met Met Ala Phe Asp Leu Ile Tyr Pro Ala Asn
 705 710 715 720
 Leu Glu Arg Phe Asp Pro Asn Phe Thr Ala Lys Lys Ser Val Leu Ala
 725 730 735
 Ala Asp Asn His Leu Ile Phe Ile Gly Ser Asn Ile Asn Ser Ser Asp
 740 745 750
 Lys Asn Lys Asn Val Glu Thr Thr Leu Phe Gln His Ala Ile Thr Pro
 755 760 765
 Thr Leu Asn Thr Leu Trp Ile Asn Gly Gln Lys Ile Glu Asn Met Pro
 770 775 780
 Tyr Gln Thr Thr Leu Gln Gln Gly Asp Trp Leu Ile Asp Ser Asn Gly
 785 790 795 800
 Asn Gly Tyr Leu Ile Thr Gln Ala Glu Lys Val Asn Val Ser Arg Gln
 805 810 815
 His Gln Val Ser Ala Glu Asn Lys Asn Arg Gln Pro Thr Glu Gly Asn
 820 825 830
 Phe Ser Ser Ala Trp Ile Asp His Ser Thr Arg Pro Lys Asp Ala Ser
 835 840 845
 Tyr Glu Tyr Met Val Phe Leu Asp Ala Thr Pro Glu Lys Met Gly Glu
 850 855 860
 Met Ala Gln Lys Phe Arg Glu Asn Asn Gly Leu Tyr Gln Val Leu Arg
 865 870 875 880
 Lys Asp Lys Asp Val His Ile Ile Leu Asp Lys Leu Ser Asn Val Thr
 885 890 895
 Gly Tyr Ala Phe Tyr Gln Pro Ala Ser Ile Glu Asp Lys Trp Ile Lys
 900 905 910
 Lys Val Asn Lys Pro Ala Ile Val Met Thr His Arg Gln Lys Asp Thr
 915 920 925
 Leu Ile Val Ser Ala Val Thr Pro Asp Leu Asn Met Thr Arg Gln Lys
 930 935 940
 Ala Ala Thr Pro Val Thr Ile Asn Val Thr Ile Asn Gly Lys Trp Gln
 945 950 955 960
 Ser Ala Asp Lys Asn Ser Glu Val Lys Tyr Gln Val Ser Gly Asp Asn
 965 970 975
 Thr Glu Leu Thr Phe Thr Ser Tyr Phe Gly Ile Pro Gln Glu Ile Lys
 980 985 990
 Leu Ser Pro Leu Pro Ala Cys Arg Asp Ala Thr His Glu Arg Ala Pro
 995 1000 1005
 Glu Thr Ile Cys Ser Ile Asn Cys Cys Asn Phe Ile Asp Glu Asn
 1010 1015 1020
 Thr Ile Ala Leu
 1025

<210> SEQ ID NO 7
 <211> LENGTH: 1027
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: CHONDROITINASE ABCI

<400> SEQUENCE: 7

Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
 1 5 10 15
 Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
 20 25 30

-continued

Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
 35 40 45
 Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
 50 55 60
 Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
 65 70 75 80
 Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
 85 90 95
 Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
 100 105 110
 Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
 115 120 125
 Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
 130 135 140
 Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
 145 150 155 160
 Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
 165 170 175
 Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
 180 185 190
 Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
 195 200 205
 Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
 210 215 220
 Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
 225 230 235 240
 Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
 245 250 255
 Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
 260 265 270
 Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
 275 280 285
 Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
 290 295 300
 Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
 305 310 315 320
 Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
 325 330 335
 Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
 340 345 350
 Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
 355 360 365
 Ser Ser Arg Trp Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380
 Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400
 Ser Arg Glu Phe Lys Ser Ser Phe Asp Met Lys Val Ser Ala Asp Ser
 405 410 415
 Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430
 Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445
 Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly

-continued

450			455			460									
Lys	Asp	Gly	Leu	Arg	Pro	Asp	Gly	Thr	Ala	Trp	Arg	His	Glu	Gly	Asn
465					470					475					480
Tyr	Pro	Gly	Tyr	Ser	Phe	Pro	Ala	Phe	Lys	Asn	Ala	Ser	Gln	Leu	Ile
				485						490					495
Tyr	Leu	Leu	Arg	Asp	Thr	Pro	Phe	Ser	Val	Gly	Glu	Ser	Gly	Trp	Asn
			500					505					510		
Ser	Leu	Lys	Lys	Ala	Met	Val	Ser	Ala	Trp	Ile	Tyr	Ser	Asn	Pro	Glu
		515					520					525			
Val	Gly	Leu	Pro	Leu	Ala	Gly	Arg	His	Pro	Leu	Asn	Ser	Pro	Ser	Leu
	530						535				540				
Lys	Ser	Val	Ala	Gln	Gly	Tyr	Tyr	Trp	Leu	Ala	Met	Ser	Ala	Lys	Ser
545					550					555					560
Ser	Pro	Asp	Lys	Thr	Leu	Ala	Ser	Ile	Tyr	Leu	Ala	Ile	Ser	Asp	Lys
				565						570					575
Thr	Gln	Asn	Glu	Ser	Thr	Ala	Ile	Phe	Gly	Glu	Thr	Ile	Thr	Pro	Ala
			580					585						590	
Ser	Leu	Pro	Gln	Gly	Phe	Tyr	Ala	Phe	Asn	Gly	Gly	Ala	Phe	Gly	Ile
		595					600					605			
His	Arg	Trp	Gln	Asp	Lys	Met	Val	Thr	Leu	Lys	Ala	Tyr	Asn	Thr	Asn
	610						615				620				
Val	Trp	Ser	Ser	Glu	Ile	Tyr	Asn	Lys	Val	Asn	Arg	Tyr	Gly	Arg	Tyr
625					630					635					640
Gln	Ser	His	Gly	Val	Ala	Gln	Ile	Val	Ser	Asn	Gly	Ser	Gln	Leu	Ser
				645						650					655
Gln	Gly	Tyr	Gln	Gln	Glu	Gly	Trp	Asp	Trp	Asn	Arg	Met	Pro	Gly	Ala
			660					665					670		
Thr	Thr	Ile	His	Leu	Pro	Leu	Lys	Asp	Leu	Asp	Ser	Pro	Lys	Pro	His
		675					680						685		
Thr	Leu	Met	Gln	Arg	Gly	Glu	Arg	Gly	Phe	Ser	Gly	Thr	Ser	Ser	Leu
	690						695				700				
Glu	Gly	Gln	Tyr	Gly	Met	Met	Ala	Phe	Asp	Leu	Ile	Tyr	Pro	Ala	Asn
705					710					715					720
Leu	Glu	Arg	Phe	Asp	Pro	Asn	Phe	Thr	Ala	Lys	Lys	Ser	Val	Leu	Ala
				725						730					735
Ala	Asp	Asn	His	Leu	Ile	Phe	Ile	Gly	Ser	Asn	Ile	Asn	Ser	Ser	Asp
			740					745					750		
Lys	Asn	Lys	Asn	Val	Glu	Thr	Thr	Leu	Phe	Gln	His	Ala	Ile	Thr	Pro
		755						760					765		
Thr	Leu	Asn	Thr	Leu	Trp	Ile	Asn	Gly	Gln	Lys	Ile	Glu	Asn	Met	Pro
	770						775				780				
Tyr	Gln	Thr	Thr	Leu	Gln	Gln	Gly	Asp	Trp	Leu	Ile	Asp	Ser	Asn	Gly
785					790					795					800
Asn	Gly	Tyr	Leu	Ile	Thr	Gln	Ala	Glu	Lys	Val	Asn	Val	Ser	Arg	Gln
				805						810					815
His	Gln	Val	Ser	Ala	Glu	Asn	Lys	Asn	Arg	Gln	Pro	Thr	Glu	Gly	Asn
			820					825					830		
Phe	Ser	Ser	Ala	Trp	Ile	Asp	His	Ser	Thr	Arg	Pro	Lys	Asp	Ala	Ser
		835					840						845		
Tyr	Glu	Tyr	Met	Val	Phe	Leu	Asp	Ala	Thr	Pro	Glu	Lys	Met	Gly	Glu
	850						855				860				
Met	Ala	Gln	Lys	Phe	Arg	Glu	Asn	Asn	Gly	Leu	Tyr	Gln	Val	Leu	Arg
865					870					875					880

-continued

Lys Asp Lys Asp Val His Ile Ile Leu Asp Lys Leu Ser Asn Val Thr
 885 890 895
 Gly Tyr Ala Phe Tyr Gln Pro Ala Ser Ile Glu Asp Lys Trp Ile Lys
 900 905 910
 Lys Val Asn Lys Pro Ala Ile Val Met Thr His Arg Gln Lys Asp Thr
 915 920 925
 Leu Ile Val Ser Ala Val Thr Pro Asp Leu Asn Met Thr Arg Gln Lys
 930 935 940
 Ala Ala Thr Pro Val Thr Ile Asn Val Thr Ile Asn Gly Lys Trp Gln
 945 950 955 960
 Ser Ala Asp Lys Asn Ser Glu Val Lys Tyr Gln Val Ser Gly Asp Asn
 965 970 975
 Thr Glu Leu Thr Phe Thr Ser Tyr Phe Gly Ile Pro Gln Glu Ile Lys
 980 985 990
 Leu Ser Pro Leu Pro Ala Cys Arg Asp Ala Thr His Glu Arg Ala Pro
 995 1000 1005
 Glu Thr Ile Cys Ser Ile Asn Cys Cys Asn Phe Ile Asp Glu Asn
 1010 1015 1020
 Thr Ile Ala Leu
 1025

<210> SEQ ID NO 8
 <211> LENGTH: 2994
 <212> TYPE: DNA
 <213> ORGANISM: *Proteus vulgaris*

<400> SEQUENCE: 8

gccaccagca atcctgcatt tgatcctaaa aatctgatgc agtcagaaat ttaccatttt 60
 gcacaaaata acccattagc agacttttca tcagataaaa actcaatact aacgttatct 120
 gataaacgta gcattatggg aaaccaatct cttttatgga aatggaaagg tggtagtagc 180
 tttactttac ataaaaaact gattgtcccc accgataaag aagcatctaa agcatgggga 240
 cgctcatcca cccccgtttt ctcattttgg ctttacaatg aaaaaccgat tgatggttat 300
 cttactatcg atttcggaga aaaactcatt tcaaccagtg aggctcaggc aggctttaa 360
 gtaaaattag atttcactgg ctggcgctact gtgggagtct ctttaaataa cgatcttgaa 420
 aatcgagaga tgaccttaaa tgcaaccaat acctcctctg atggtactca agacagcatt 480
 gggcggttctt taggtgctaa agtcgatagt attcgtttta aagcgccttc taatgtgagt 540
 cagggtgaaa tctatatcga ccgtattatg ttttctgtcg atgatgctcg ctaccaatgg 600
 tctgattatc aagtaaaaac tcgcttatca gaacctgaaa ttcaatttca caacgtaaag 660
 ccacaactac ctgtaacacc tgaaaattta gcggccattg atcttattcg ccaacgtcta 720
 attaatgaat ttgtcggagg tgaaaaagag acaaacctcg cattagaaga gaatatcagc 780
 aaattaaana gtgatttcga tgctcttaat actcacactt tagcaaatgg tggaacgcaa 840
 ggcagacatc tgatcactga taaacaaatc attattttatc aaccagagaa tcttaactct 900
 caagataaac aactatttga taattatggt attttaggta attacacgac attaattggtt 960
 aatattagcc gtgcttatgt gctggaaaaa gatccacac aaaaggcgca actaaagcag 1020
 atgtacttat taatgacaaa gcatttatta gatcaaggct ttgttaaagg gagtgcttta 1080
 gtgacaacc atcaactggg atacagttct cgttgggtgt atatttcac gttattaatg 1140
 tctgatgcac taaaagaagc gaacctacaa actcaagttt atgattcatt actgtgggat 1200

-continued

```

tcacgtgagt ttaaaagtag ttttgatag aaagtaagtg ctgatagctc tgatctagat 1260
tatttcaata ctttatctcg ccaacattta gccttattac tactagagcc tgatgatcaa 1320
aagcgtatca acttagttaa tactttcagc cattatatca ctggcgcatc aacgcaagtg 1380
ccaccgggtg gtaaagatgg tttacgcct gatggtacag catggcgaca tgaaggaac 1440
tatccgggct actctttccc agccttataa aatgcctctc agcttattta tttattacgc 1500
gatacaccat tttcagtggg tgaaagtggg tggaatagcc tgaaaaaagc gatggtttca 1560
gcgtggatct acagtaatcc agaagtggga ttaccgcttg caggaagaca ccctctaac 1620
tcacctctcg taaaatcagt cgctcaaggc tattactggc ttgccatgct tgcaaatca 1680
tcgctgata aaacacttgc atctatttat cttgcgatta gtgataaaac acaaaatgaa 1740
tcaactgcta tttttggaga aactattaca ccagcgtctt tacctcaagg tttctatgcc 1800
ttaatggcg gtgcttttgg tattcatcgt tggcaagata aaatgggtgac actgaaagct 1860
tataacacca atgtttggtc atctgaaatt tataacaag ataaccgta tggccgttac 1920
caaagtcagt gtgctgctca aatagtgagt aatggctcgc agctttcaca gggctatcag 1980
caagaaggtt gggattggaa tagaatgcca ggggcaacca ctatccacct tcctcttaa 2040
gacttagaca gtccaaacc tcatacctta atgcaacgtg gagagcgtgg atttagcgga 2100
acatcatccc ttgaaggtca atatggcatg atggcattcg atcttattta tcccgccaat 2160
cttgagcgtt ttgatcctaa tttcactgcg aaaaagagtg tattagcgc tgataatcac 2220
ttaattttta ttggtagcaa tataaatagt agtgataaaa ataaaaatgt tgaacgacc 2280
ttattccaac atgccattac tccaacatta aatacccttt ggattaatgg acaaaagata 2340
gaaaacatgc cttatcaaac aacacttcaa caaggtgatt ggtaattga tagcaatggc 2400
aatggttact taattactca agcagaaaaa gtaaatgtaa gtcgccaaca tcaggtttca 2460
gcggaaaata aaaatcgcca accgacagaa ggaaacttta gctcggcatg gatcgatcac 2520
agcactcgcc ccaaagatgc cagttatgag tatatggtct ttttagatgc gacacctgaa 2580
aaaaatggag agatggcaca aaaattccgt gaaaataatg ggttatatca ggttctctgt 2640
aaggataaag acgttcatat tattctcgat aaactcagca atgtaacggg atatgccttt 2700
tatcagccag catcaattga agacaatgg atcaaaaagg ttaataaacc tgcaattgtg 2760
atgactcacc gacaaaaaga cactcttatt gtcagtgcag ttacacctga tttaaatag 2820
actcgccaaa aagcagcaac tcctgtcacc atcaatgtca cgattaatgg caaatggcaa 2880
tctgctgata aaaatagtga agtgaatat caggtttctg gtgataaacac tgaactgacg 2940
tttacgagtt actttgggat tccacaagaa atcaaaactct cgccactccc ttga 2994

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1027
<212> TYPE: PRT
<213> ORGANISM: Proteus vulgaris

```

```

<400> SEQUENCE: 9

```

```

Ala Thr Ser Asn Pro Ala Phe Asp Pro Lys Asn Leu Met Gln Ser Glu
1           5           10           15
Ile Tyr His Phe Ala Gln Asn Asn Pro Leu Ala Asp Phe Ser Ser Asp
20           25           30
Lys Asn Ser Ile Leu Thr Leu Ser Asp Lys Arg Ser Ile Met Gly Asn
35           40           45
Gln Ser Leu Leu Trp Lys Trp Lys Gly Gly Ser Ser Phe Thr Leu His
50           55           60

```


-continued

Lys Lys Leu Ile Val Pro Thr Asp Lys Glu Ala Ser Lys Ala Trp Gly
 65 70 75 80
 Arg Ser Ser Thr Pro Val Phe Ser Phe Trp Leu Tyr Asn Glu Lys Pro
 85 90 95
 Ile Asp Gly Tyr Leu Thr Ile Asp Phe Gly Glu Lys Leu Ile Ser Thr
 100 105 110
 Ser Glu Ala Gln Ala Gly Phe Lys Val Lys Leu Asp Phe Thr Gly Trp
 115 120 125
 Arg Thr Val Gly Val Ser Leu Asn Asn Asp Leu Glu Asn Arg Glu Met
 130 135 140
 Thr Leu Asn Ala Thr Asn Thr Ser Ser Asp Gly Thr Gln Asp Ser Ile
 145 150 155 160
 Gly Arg Ser Leu Gly Ala Lys Val Asp Ser Ile Arg Phe Lys Ala Pro
 165 170 175
 Ser Asn Val Ser Gln Gly Glu Ile Tyr Ile Asp Arg Ile Met Phe Ser
 180 185 190
 Val Asp Asp Ala Arg Tyr Gln Trp Ser Asp Tyr Gln Val Lys Thr Arg
 195 200 205
 Leu Ser Glu Pro Glu Ile Gln Phe His Asn Val Lys Pro Gln Leu Pro
 210 215 220
 Val Thr Pro Glu Asn Leu Ala Ala Ile Asp Leu Ile Arg Gln Arg Leu
 225 230 235 240
 Ile Asn Glu Phe Val Gly Gly Glu Lys Glu Thr Asn Leu Ala Leu Glu
 245 250 255
 Glu Asn Ile Ser Lys Leu Lys Ser Asp Phe Asp Ala Leu Asn Thr His
 260 265 270
 Thr Leu Ala Asn Gly Gly Thr Gln Gly Arg His Leu Ile Thr Asp Lys
 275 280 285
 Gln Ile Ile Ile Tyr Gln Pro Glu Asn Leu Asn Ser Gln Asp Lys Gln
 290 295 300
 Leu Phe Asp Asn Tyr Val Ile Leu Gly Asn Tyr Thr Thr Leu Met Phe
 305 310 315 320
 Asn Ile Ser Arg Ala Tyr Val Leu Glu Lys Asp Pro Thr Gln Lys Ala
 325 330 335
 Gln Leu Lys Gln Met Tyr Leu Leu Met Thr Lys His Leu Leu Asp Gln
 340 345 350
 Gly Phe Val Lys Gly Ser Ala Leu Val Thr Thr His His Trp Gly Tyr
 355 360 365
 Ser Ser Arg Trp Trp Tyr Ile Ser Thr Leu Leu Met Ser Asp Ala Leu
 370 375 380
 Lys Glu Ala Asn Leu Gln Thr Gln Val Tyr Asp Ser Leu Leu Trp Tyr
 385 390 395 400
 Ser Arg Glu Phe Lys Ser Ser Phe Asp Met Lys Val Ser Ala Asp Ser
 405 410 415
 Ser Asp Leu Asp Tyr Phe Asn Thr Leu Ser Arg Gln His Leu Ala Leu
 420 425 430
 Leu Leu Leu Glu Pro Asp Asp Gln Lys Arg Ile Asn Leu Val Asn Thr
 435 440 445
 Phe Ser His Tyr Ile Thr Gly Ala Leu Thr Gln Val Pro Pro Gly Gly
 450 455 460
 Lys Asp Gly Leu Arg Pro Asp Gly Thr Ala Trp Arg His Glu Gly Asn
 465 470 475 480

-continued

Tyr Pro Gly Tyr Ser Phe Pro Ala Phe Lys Asn Ala Ser Gln Leu Ile
 485 490 495
 Tyr Leu Leu Arg Asp Thr Pro Phe Ser Val Gly Glu Ser Gly Trp Asn
 500 505 510
 Ser Leu Lys Lys Ala Met Val Ser Ala Trp Ile Tyr Ser Asn Pro Glu
 515 520 525
 Val Gly Leu Pro Leu Ala Gly Arg His Pro Leu Asn Ser Pro Ser Leu
 530 535 540
 Lys Ser Val Ala Gln Gly Tyr Tyr Trp Leu Ala Met Ser Ala Lys Ser
 545 550 555 560
 Ser Pro Asp Lys Thr Leu Ala Ser Ile Tyr Leu Ala Ile Ser Asp Lys
 565 570 575
 Thr Gln Asn Glu Ser Thr Ala Ile Phe Gly Glu Thr Ile Thr Pro Ala
 580 585 590
 Ser Leu Pro Gln Gly Phe Tyr Ala Phe Asn Gly Gly Ala Phe Gly Ile
 595 600 605
 His Arg Trp Gln Asp Lys Met Val Thr Leu Lys Ala Tyr Asn Thr Asn
 610 615 620
 Val Trp Ser Ser Glu Ile Tyr Asn Lys Asp Asn Arg Tyr Gly Arg Tyr
 625 630 635 640
 Gln Ser His Gly Val Ala Gln Ile Val Ser Asn Gly Ser Gln Leu Ser
 645 650 655
 Gln Gly Tyr Gln Gln Glu Gly Trp Asp Trp Asn Arg Met Pro Gly Ala
 660 665 670
 Thr Thr Ile His Leu Pro Leu Lys Asp Leu Asp Ser Pro Lys Pro His
 675 680 685
 Thr Leu Met Gln Arg Gly Glu Arg Gly Phe Ser Gly Thr Ser Ser Leu
 690 695 700
 Glu Gly Gln Tyr Gly Met Met Ala Phe Asp Leu Ile Tyr Pro Ala Asn
 705 710 715 720
 Leu Glu Arg Phe Asp Pro Asn Phe Thr Ala Lys Lys Ser Val Leu Ala
 725 730 735
 Ala Asp Asn His Leu Ile Phe Ile Gly Ser Asn Ile Asn Ser Ser Asp
 740 745 750
 Lys Asn Lys Asn Val Glu Thr Thr Leu Phe Gln His Ala Ile Thr Pro
 755 760 765
 Thr Leu Asn Thr Leu Trp Ile Asn Gly Gln Lys Ile Glu Asn Met Pro
 770 775 780
 Tyr Gln Thr Thr Leu Gln Gln Gly Asp Trp Leu Ile Asp Ser Asn Gly
 785 790 795 800
 Asn Gly Tyr Leu Ile Thr Gln Ala Glu Lys Val Asn Val Ser Arg Gln
 805 810 815
 His Gln Val Ser Ala Glu Asn Lys Asn Arg Gln Pro Thr Glu Gly Asn
 820 825 830
 Phe Ser Ser Ala Trp Ile Asp His Ser Thr Arg Pro Lys Asp Ala Ser
 835 840 845
 Tyr Glu Tyr Met Val Phe Leu Asp Ala Thr Pro Glu Lys Met Gly Glu
 850 855 860
 Met Ala Gln Lys Phe Arg Glu Asn Asn Gly Leu Tyr Gln Val Leu Arg
 865 870 875 880
 Lys Asp Lys Asp Val His Ile Ile Leu Asp Lys Leu Ser Asn Val Thr
 885 890 895
 Gly Tyr Ala Phe Tyr Gln Pro Ala Ser Ile Glu Asp Lys Trp Ile Lys

-continued

900	905	910
Lys Val Asn Lys Pro Ala Ile Val Met Thr His Arg Gln Lys Asp Thr		
915	920	925
Leu Ile Val Ser Ala Val Thr Pro Asp Leu Asn Met Thr Arg Gln Lys		
930	935	940
Ala Ala Thr Pro Val Thr Ile Asn Val Thr Ile Asn Gly Lys Trp Gln		
945	950	955
Ser Ala Asp Lys Asn Ser Glu Val Lys Tyr Gln Val Ser Gly Asp Asn		
965	970	975
Thr Glu Leu Thr Phe Thr Ser Tyr Phe Gly Ile Pro Gln Glu Ile Lys		
980	985	990
Leu Ser Pro Leu Pro Ala Cys Arg Asp Ala Thr His Glu Arg Ala Pro		
995	1000	1005
Glu Thr Ile Cys Ser Ile Asn Cys Cys Asn Phe Ile Asp Glu Asn		
1010	1015	1020
Thr Ile Ala Leu		
1025		

What is claimed is:

1. A pharmaceutical composition comprising a therapeutically effective amount of mutant chondroitinase ABC I enzyme selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, and SEQ ID NO: 7, and a pharmaceutically acceptable excipient.

2. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 1, and a pharmaceutically acceptable excipient.

3. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 2, and a pharmaceutically acceptable excipient.

4. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 3, and a pharmaceutically acceptable excipient.

25 5. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 4, and a pharmaceutically acceptable excipient.

30 6. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 5, and a pharmaceutically acceptable excipient.

35 7. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 6, and a pharmaceutically acceptable excipient.

40 8. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a therapeutically effective amount of mutant chondroitinase ABC I enzyme SEQ ID NO: 7, and a pharmaceutically acceptable excipient.

* * * * *