




(86) Date de dépôt PCT/PCT Filing Date: 2014/03/12  
 (87) Date publication PCT/PCT Publication Date: 2014/09/18  
 (45) Date de délivrance/Issue Date: 2021/02/16  
 (85) Entrée phase nationale/National Entry: 2015/09/10  
 (86) N° demande PCT/PCT Application No.: JP 2014/056416  
 (87) N° publication PCT/PCT Publication No.: 2014/142146  
 (30) Priorité/Priority: 2013/03/13 (JP2013-050766)

(51) Cl.Int./Int.Cl. *A61K 31/453* (2006.01),  
*A61K 9/10* (2006.01), *A61K 9/107* (2006.01),  
*A61P 27/02* (2006.01)  
 (72) Inventeurs/Inventors:  
 MIYAKE, HIDEKI, JP;  
 ODA, TOMOKO, JP;  
 SHII, DAISUKE, JP  
 (73) Propriétaire/Owner:  
 SANTEN PHARMACEUTICAL CO., LTD., JP  
 (74) Agent: MARKS & CLERK

(54) Titre : AGENTS THERAPEUTIQUES POUR UN DYSFONCTIONNEMENT MEIBOMIEN  
 (54) Title: THERAPEUTIC AGENT FOR MEIBOMIAN GLAND DYSFUNCTION

[Formula 1]




(57) Abrégé/Abstract:

A compound represented by the formula (1): (see formula 1) wherein R<sup>1</sup> represents a hydroxyl group, a dimethylphosphoryloxy group, etc.; and R<sup>2</sup> and R<sup>3</sup> represent a methoxy group, etc., or a pharmaceutically acceptable salt thereof can reduce a score of telangiectasia around the meibomian gland orifices and a number of obstruction at the orifices, so that it is useful as a prophylactic and/or therapeutic agent for meibomian gland dysfunction.

## ABSTRACT

A compound represented by the formula (1):




- 5            wherein R<sup>1</sup> represents a hydroxyl group, a dimethylphosphinoxyloxy group, etc.; and R<sup>2</sup> and R<sup>3</sup> represent a methoxy group, etc., or a pharmaceutically acceptable salt thereof can reduce a score of telangiectasia around the meibomian gland orifices and a number of obstruction at the orifices, so that it is useful as a prophylactic and/or therapeutic agent for meibomian gland dysfunction.

## DESCRIPTION

## THERAPEUTIC AGENT FOR MEIBOMIAN GLAND DYSFUNCTION

## 5 TECHNICAL FIELD

[0001] The present invention relates to a prophylactic and/or therapeutic agent for meibomian gland dysfunction containing a compound represented by the formula (1):  
[Formula 1]



10 wherein

$R^1$  represents a hydroxyl group, a methoxy group, a hydroxymethoxy group, an ethoxy group, a 1-hydroxyethoxy group, a 2-hydroxyethoxy group, a 2-methoxyethoxy group, a 2-ethoxyethoxy group, a formyloxy group, a carboxyloxy group, an acetoxy group, a hydroxyacetoxy group, a propionyloxy group, a 2-hydroxypropionyloxy group, a 3-hydroxypropionyloxy group, a 2-methylpropionyloxy group, a 2-(hydroxymethyl)propionyloxy group, a 3-hydroxy-2-(hydroxymethyl)propionyloxy group, a 2,2-dimethylpropionyloxy group, a 2-(hydroxymethyl)-2-methylpropionyloxy group, a 2,2-bis(hydroxymethyl)propionyloxy group, a methylphosphynoyloxy group, a dimethylphosphynoyloxy group or a 1H-tetrazol-1-yl group;

20  $R^2$  represents a hydrogen atom, a hydroxyl group, a methoxy group, a mercapto group or a methylthio group;

$R^3$  represents a hydrogen atom, a hydroxyl group or a methoxy group;

a wavy line represents that the carbon atom bonded to  $R^1$ ,  $R^2$  or  $R^3$  can take either configuration of S or R;

25 (hereinafter sometimes referred to as "the present compound") or a pharmaceutically acceptable salt thereof as an active ingredient.

## BACKGROUND ART

[0002] The meibomian gland is present in the tarsus, is a sebaceous gland having orifices at the upper and lower eyelid margins, and a lipid secreted by the meibomian gland plays a variety of roles in the external eye.

5 [0003] In Non-Patent Document 1, it has been disclosed that among the patients visiting ophthalmology who appeal symptoms such as ocular discomfort, as a main complaint, meibomian gland dysfunction (hereinafter also referred to as “MGD”) becomes a cause thereof in a significant proportion, and lowering in quality of life is caused in many patients.

10 [0004] However, it has never been known an effective treatment method of MGD, and yet, clear definition for MGD or diagnostic criteria itself has never been present in the first place. Thus, in recent years, there are worldwide movements to define MGD as an independent disease and to prepare its diagnostic criteria, and in Japan, MGD has been defined to be a disease “which is a state in which the function of the meibomian  
15 gland causes abnormality in diffuse by various causes, and is accompanied by chronic ocular discomfort” (see Non-Patent Document 1).

[0005] Also, in Non-Patent Document 1, it has been disclosed that MGD is classified into a decreased secretion type MGD and an increased secretion type MGD, and in the decreased secretion type MGD, it has also been disclosed that secretion of meibomian  
20 gland lipids is reduced by obstruction of the meibomian gland orifices and the like. Further, in Non-Patent Document 1, it has been also disclosed that three of <1> subjective symptom such as ocular discomfort, <2> signs/findings on abnormalities around the meibomian gland orifices such as vasodilation, and <3> signs/findings on obstruction at the meibomian gland orifices are positive is diagnosed to be a decreased  
25 secretion type MGD.

[0006] Incidentally, as a disease generating at the meibomian gland, chalazion, internal hordeolum, etc., have been known other than MGD, and in Non-Patent Document 1, there are disclosed that these diseases are topical diseases, and are different diseases from MGD in which the meibomian gland is obstructed in diffuse.  
30 In addition, MGD sometimes causes an evaporation promoting type dry eye, in Non-Patent Document 1, it has been disclosed that there is a case where no dry eye is accompanied depending on an amount of tear fluid, a term of the disease or a degree of severity.

[0007] In Non-Patent Document 2, a classification method of MGD in the U.S. has  
35 been disclosed, and it has also been disclosed that the decreased secretion type MGD (low-delivery state MGD) can be further classified into two of “hyposecretory MGD

(meibomian hyposecretion)” and “obstructive MGD (meibomian gland obstruction)”. Further, in Non-Patent Document 2, it has been suggested that MGD is one of the diseases which cause posterior blepharitis.

5 [0008] Sirolimus (also referred to as rapamycin) has been known to be an immunosuppressant, and has been used as an oral preparation in the U.S., etc. It has also been known that a derivative (hereinafter sometimes referred to as “sirolimus derivative”) in which a part of the structure of sirolimus has been modified has the similar activity to that of the sirolimus, and in the sirolimus derivative, deforolimus, everolimus, temsirolimus, zotarolimus, biolimus, novolimus, etc., are contained.

10 [0009] In Patent Document 1, it has been disclosed an ophthalmic composition comprising an mTOR inhibitor such as sirolimus, everolimus, temsirolimus, etc., a first surfactant having an HLB index exceeding about 10, and a second surfactant having an HLB index exceeding about 13. However, in Patent Document 1, it has never been disclosed whether the mTOR inhibitor has a treatment effect on MGD or not.

15 [0010] Also, in Patent Document 2, it has been suggested that a pipercolic acid derivative such as rapamycin, can treat vision disorder, and in the vision disorder, meibomian gland carcinoma and internal hordeolum (meibomian gland sty) are contained. However, the meibomian gland carcinoma is a disease clearly different from MGD, and as stated above, the internal hordeolum is a disease also different from  
20 MGD.

[0011] As mentioned above, in Patent Document 1 and 2, there is neither description nor suggestion on whether the present compound has a treatment effect on MGD or not.

[0012] In addition, it is also not clear for those skilled in the art whether or not a medicament which can treat dry eye or posterior blepharitis can prevent and/or treat  
25 MGD itself.

#### PRIOR ART DOCUMENT

##### Patent Documents

[0013] Patent Document 1: JP 2010-540682A  
30 Patent Document 2: JP 2002-522485A

##### Non-Patent Documents

[0014] Non-Patent Document 1: Journal of the Eye, 27(5), 627-631 (2010)  
Non-Patent Document 2: Investigative Ophthalmology & Visual Science, 52(4),  
35 1930-1937 (2011)

## DISCLOSURE OF THE INVENTION

## Problems to be Solved by the Invention


[0015] An object of the present invention is to provide a prophylactic and/or therapeutic agent for meibomian gland dysfunction (MGD).

5

## Means to Solve the Problems

[0016] The present inventors have intensively studied to search a prophylactic and/or therapeutic agent for meibomian gland dysfunction (MGD), and as a result, they have found that a compound represented by the formula (1):

10 [Formula 2]



wherein

$R^1$  represents a hydroxyl group, a methoxy group, hydroxymethoxy group, ethoxy group, 1-hydroxyethoxy group, 2-hydroxyethoxy group, 2-methoxyethoxy group, 2-ethoxyethoxy group, a formyloxy group, a carboxyloxy group, an acetoxy group, a hydroxyacetoxy group, a propionyloxy group, a 2-hydroxypropionyloxy group, a 3-hydroxypropionyloxy group, a 2-methylpropionyloxy group, a 2-(hydroxymethyl)propionyloxy group, a 3-hydroxy-2-(hydroxymethyl)propionyloxy group, a 2,2-dimethylpropionyloxy group, a 2-(hydroxymethyl)-2-methylpropionyloxy group, a 2,2-bis(hydroxymethyl)propionyloxy group, a methylphosphynoyloxy group, a dimethylphosphynoyloxy group or a 1H-tetrazol-1-yl group;

$R^2$  represents a hydrogen atom, a hydroxyl group, a methoxy group, a mercapto group or a methylthio group;

$R^3$  represents a hydrogen atom, a hydroxyl group or a methoxy group;

25 a wavy line represents that the carbon atom bonded to  $R^1$ ,  $R^2$  or  $R^3$  can take either configuration of S or R,

or a pharmaceutically acceptable salt thereof can reduce a number of obstructions at the meibomian gland orifices, and the compound of the above-mentioned formula (1) or a pharmaceutically acceptable salt thereof has a treatment effect on MGD and suppresses obstruction of the meibomian gland, whereby the present invention has been  
5 accomplished.

[0017] That is, the present invention relates to a prophylactic and/or therapeutic agent for meibomian gland dysfunction (MGD), which comprises the compound of the above-mentioned formula (1) (the present compound) or a pharmaceutically acceptable salt thereof as an active ingredient (hereinafter sometimes referred to as “the present  
10 agent”).

[0018] Also, the present compound is preferably a compound, in the above-mentioned formula (1), wherein  $R^1$  represents a hydroxyl group, a 2-hydroxyethoxy group, a 2-ethoxyethoxy group, a 2,2-bis(hydroxymethyl)propionyloxy group, a dimethylphosphinyloxy group or a 1H-tetrazol-1-yl group;  $R^2$  represents a hydrogen atom, a  
15 hydroxyl group, a methoxy group or a methylthio group; and  $R^3$  represents a hydrogen atom or a methoxy group, or a pharmaceutically acceptable salt thereof.

[0019] Further, the present compound is preferably a compound, in the above-mentioned formula (1), wherein  $R^1$  represents a hydroxyl group or a dimethylphosphinyloxy group;  $R^2$  represents a methoxy group;  $R^3$  represents a methoxy group,  
20 or a pharmaceutically acceptable salt thereof.

[0020] Moreover, the present compound is preferably sirolimus, deforolimus, everolimus, temsirolimus, zotarolimus, biolimus, novolimus, 7-epi-rapamycin, 7-thiomethyl-rapamycin, 7-epi-thiomethyl-rapamycin, 7-demethoxy-rapamycin, 32-demethoxy-rapamycin or their pharmaceutically acceptable salts, particularly  
25 preferably sirolimus, deforolimus or a salt thereof.

[0021] Furthermore, an administration form of the present agent is preferably instillation administration or administration to the eyelid skin.

[0022] Also, a dosage form of the present agent is preferably an eye drop, an ophthalmic ointment or an ointment (excluding an ophthalmic ointment), and  
30 characteristics of the eye drop is preferably a suspension or an emulsion.

[0023] Further, other embodiments of the present invention relate to a composition for suppressing obstruction of the meibomian gland which comprises the present compound or a pharmaceutically acceptable salt thereof as an active ingredient (hereinafter sometimes referred to as “the present composition”).

[0024] Moreover, an active ingredient of the present composition is preferably  
35 sirolimus, deforolimus or a salt thereof.

[0025] Furthermore, other embodiments of the present invention relate to a pharmaceutical composition for prophylaxis and/or treatment of meibomian gland dysfunction (MGD) which comprises the compound of the above-mentioned formula (1) (the present compound) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable additive.

[0026] Also, other embodiments of the present invention relate to the compound of the above-mentioned formula (1) (the present compound) for use in the suppression of obstruction of the meibomian gland.

[0027] Further, other embodiments of the present invention relate to the compound of the above-mentioned formula (1) (the present compound) for use in the suppression of telangiectasia around the meibomian gland orifices.

[0028] Moreover, other embodiments of the present invention relate to the compound of the above-mentioned formula (1) (the present compound) for use in the prophylaxis and/or treatment of meibomian gland dysfunction (MGD).

[0029] Furthermore, other embodiments of the present invention relate to a use of the compound of the above-mentioned formula (1) (the present compound) for the manufacture of a composition for suppressing obstruction of the meibomian gland.

[0030] Also, other embodiments of the present invention relate to a use of the compound of the above-mentioned formula (1) (the present compound) for the manufacture of a composition for suppressing telangiectasia around the meibomian gland orifices.

[0031] Further, other embodiments of the present invention relate to a use of the compound of the above-mentioned formula (1) (the present compound) for the manufacture of a medicine for prophylaxis and/or treatment of meibomian gland dysfunction (MGD).

[0032] Moreover, other embodiments of the present invention relate to a method for suppressing obstruction of meibomian gland, which comprises administering the compound of the above-mentioned formula (1) (the present compound) to a human or an animal.

[0033] Furthermore, other embodiments of the present invention relate to a method for suppressing telangiectasia around the meibomian gland orifices, which comprises administering the compound of the above-mentioned formula (1) (the present compound) to a human or an animal.

[0034] Still further, other embodiments of the present invention relate to a method for prophylaxis and/or treatment of meibomian gland dysfunction (MGD), which comprises administering the compound of the above-mentioned formula (1) (the present

compound) to a human or an animal.


#### Effects of the Invention

[0035] The compound of the above-mentioned formula (1) or a pharmaceutically acceptable salt thereof is useful as a prophylactic and/or therapeutic agent for meibomian gland dysfunction (MGD).

#### BEST MODE TO CARRY OUT THE INVENTION

[0036] (A) The present compound is a compound in which the respective groups are the groups shown below in the compound represented by the formula (1):

[Formula 3]



[0037] (A1) R<sup>1</sup> represents a hydroxyl group, a methoxy group, a hydroxymethoxy group, an ethoxy group, a 1-hydroxyethoxy group, a 2-hydroxyethoxy group, a 2-methoxyethoxy group, a 2-ethoxyethoxy group, a formyloxy group, a carboxyoxo group, an acetoxy group, a hydroxyacetoxy group, a propionyloxy group, a 2-hydroxypropionyloxy group, a 3-hydroxypropionyloxy group, a 2-methylpropionyloxy group, a 2-(hydroxymethyl)propionyloxy group, a 3-hydroxy-2-(hydroxymethyl)propionyloxy group, a 2,2-dimethylpropionyloxy group, a 2-(hydroxymethyl)-2-methylpropionyloxy group, a 2,2-bis(hydroxymethyl)propionyloxy group, a methylphosphynoyloxy group, a dimethylphosphynoyloxy group or a 1H-tetrazol-1-yl group;

(A2) R<sup>2</sup> represents a hydrogen atom, a hydroxyl group, a methoxy group, a mercapto group, a methylthio group, a phenyl group, a 2,4,6-trihydroxyphenyl group or a 2,4,6-trimethoxyphenyl group; and

(A3) R<sup>3</sup> represents a hydrogen atom, a hydroxyl group or a methoxy group.

That is, the present compound is a material in which the respective group enumerated in the above-mentioned (A1), (A2) and (A3) are combined in the compound represented by the above-mentioned formula (1).

[0038] (B) As a preferred example of the present compound, there may be mentioned a compound wherein the respective groups are the groups shown below in the compound represented by the above-mentioned formula (1).

(B1) R<sup>1</sup> represents a hydroxyl group, a 2-hydroxyethoxy group, a 2-ethoxyethoxy group, a 2,2-bis(hydroxymethyl)propionyloxy group, a dimethylphosphinoxyloxy group or a 1H-tetrazol-1-yl group;

10 (B2) R<sup>2</sup> represents a hydrogen atom, a hydroxyl group, a methoxy group, a methylthio group or a 2,4,6-trimethoxyphenyl group; and

(B3) R<sup>3</sup> represents a hydrogen atom or a methoxy group.

That is, a preferred example of the present compound is a material in which the respective group enumerated in the above-mentioned (B1), (B2) and (B3) are combined in the above-mentioned formula (1).

[0039] (C) As a more preferred example of the present compound, there may be mentioned a compound in which the respective groups are groups mentioned below in the above-mentioned formula (1).

(C1) R<sup>1</sup> represents a hydroxyl group, a 2-hydroxyethoxy group, a 2-ethoxyethoxy group, a 2,2-bis(hydroxymethyl)propionyloxy group or a dimethylphosphinoxyloxy group;

(C2) R<sup>2</sup> represents a methoxy group;

(C3) R<sup>3</sup> represents a methoxy group;


That is, a more preferred example of the present compound is a material in which the respective groups enumerated in the above-mentioned (C1), (C2) and (C3) are combined in the compound represented by the above-mentioned formula (1).

[0040] In the above-mentioned formula (1), a wavy line represents that the carbon atom bonded to R<sup>1</sup>, R<sup>2</sup> or R<sup>3</sup> can take either configuration of S or R, and in the present invention, the carbon atom bonded to R<sup>1</sup> preferably has an R-configuration, and when R<sup>2</sup> or R<sup>3</sup> is not a hydrogen atom, the carbon atom bonded to R<sup>2</sup> preferably has an S-configuration, and the carbon atom bonded to R<sup>3</sup> preferably has an R-configuration.

[0041] Specific examples of the present compound may be mentioned the following compounds.


[0042] · Sirolimus

35 [Formula 4]




[0043] · Deforolimus

[Formula 5]




5 [0044] · Everolimus

[Formula 6]




[0045] · Temsirolimus

[Formula 7]




5 [0046] · Zotarolimus

[Formula 8]



[0047] · Biolimus


[Formula 9]



5 [0048] · Novolimus


[Formula 10]

- 12 -




[0049] · 7-Epi-rapamycin

[Formula 11]




5 [0050] · 7-Thiomethyl-rapamycin

[Formula 12]




[0051] · 7-Epi-thiomethyl-rapamycin

[Formula 13]




5 [0052] · 7-Demethoxy-rapamycin

[Formula 14]



[0053] · 32-Demethoxy-rapamycin

[Formula 15]



5 [0054] Preferred specific examples of the present compound are sirolimus or deforolimus, and sirolimus is particularly preferred.

[0055] The present compound or a pharmaceutically acceptable salt thereof can be prepared in accordance with the usual method in the field of the organic synthetic chemistry, and in particular, with regard to sirolimus, that commercially available from  
 10 LKT laboratories, Inc., (Catalog number: R0161) may be used. In addition, with regard to deforolimus, it may be prepared in accordance with the method disclosed in JP 2005-516065A.

[0056] The pharmaceutically acceptable salt of the present compound may be mentioned, for example, a salt with an inorganic acid such as hydrochloric acid,  
 15 hydrobromic acid, hydroiodic acid, nitric acid, sulfuric acid and phosphoric acid; a salt with an organic acid such as acetic acid, fumaric acid, maleic acid, succinic acid, citric

acid, tartaric acid, adipic acid, gluconic acid, glucoheptonic acid, glucuronic acid, terephthalic acid, methanesulfonic acid, lactic acid, hippuric acid, 1,2-ethanedisulfonic acid, isethionic acid, lactobionic acid, oleic acid, pamoic acid, polygalacturonic acid, stearic acid, tannic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, lauryl sulfate, methyl sulfate, naphthalenesulfonic acid and sulfosalicylic acid; a quaternary ammonium salt with methyl bromide and methyl iodide; and a salt with a halogen ion including a bromine ion, a chlorine ion and an iodine ion.

5  
10 [0057] Also, the present compound or a pharmaceutically acceptable salt may take a form of a hydrate or a solvate.

[0058] When a geometric isomer or an optical isomer exists in the present compound or a pharmaceutically acceptable salt thereof, the isomer(s) or a salt thereof is/are also included in the scope of the present invention. In addition, when a proton tautomerism exists in the present compound or a pharmaceutically acceptable salt thereof, the tautomer(s) or a salt thereof is/are also included in the scope of the present invention.

15 [0059] When crystal polymorphism and a crystal polymorphism group (crystal polymorphism system) exist in the present compound or a pharmaceutically acceptable salt (including a hydrate or a solvate), these crystal polymorphs and crystal polymorphism group (crystal polymorphism system) are also included in the scope of the present invention. Here, the crystal polymorphism group (crystal polymorphism system) means a crystal form at the respective stages when the crystal form is changed by the conditions and states of preparation, crystallization, preservation, etc., of these crystals (incidentally, the state after formulation is also contained in the above states), and the whole processes.

25 [0060] The present compound or a pharmaceutically acceptable salt thereof can be used for a prophylaxis and/or a treatment of meibomian gland dysfunction (MGD).

[0061] The definition of the meibomian gland dysfunction (MGD) is, for example, “a state in which the function of the meibomian gland causes abnormality in diffuse by various causes, which is accompanied by chronic ocular discomfort.” Here, “the state in which the function of the meibomian gland causes abnormality in diffuse” means that, for example, not topical meibomian gland abnormality recognized in chalazion, internal hordeolum, etc., but meibomian gland abnormality such as telangiectasia, obstruction at the meibomian gland orifices, etc., is recognized in diffuse. Also, MGD is classified into a decreased secretion type MGD (low-delivery state) and an increased secretion type MGD (high delivery state), further as the decreased secretion type MGD (low-delivery state MGD), there may be mentioned “hyposecretory MGD (meibomian

30  
35

hyosecretion)” and “obstructive MGD (meibomian gland obstruction)”.

[0062] In the decreased secretion type MGD, secretion of meibomian gland lipids is reduced by obstruction at the meibomian gland orifices, etc. Also, in the increased secretion type MGD, secretion of meibomian gland lipids is increased by various causes.

[0063] On the other hand, a state in which obstruction of the meibomian gland orifices can be admitted, but no subjective symptom is accompanied is sometimes called as, for example, “meibomian gland infarction (meibomian gland concretion)”, and the meibomian gland infarction is also included and in MGD of the present invention.

10 [0064] As MGD which can be treated by the present compound or a pharmaceutically acceptable salt thereof, particularly preferred is the decreased secretion type MGD.

[0065] By the way, as explained at the column of the background art, MGD sometimes becomes a cause of the dry eye, and has a possibility of causing posterior blepharitis.

15 [0066] The MGD contains “MGD which is accompanied by dry eye and/or posterior blepharitis (complicated)”, “MGD which becomes a cause of dry eye and/or posterior blepharitis”, “MGD which is not accompanied by dry eye (not complicated)”, “MGD which does not become a cause of dry eye”, “MGD which is not accompanied by posterior blepharitis (not complicated)”, and “MGD which does not become a cause of posterior blepharitis”.

20 [0067] In the present invention, “prophylactic and/or therapeutic agent for MGD” means a medicine which prevents and/or treats MGD. Here, “to treat and/or prevent MGD” means that, among the three indexes admitted by the MGD patient disclosed in the above-mentioned Non-Patent Document 1 (<1> subjective symptom such as ocular discomfort, <2> signs/findings on abnormalities around the meibomian gland orifices such as vasodilation, and <3> signs/findings on obstruction at the meibomian gland orifices), at least an improvement in “signs/findings on obstruction at the meibomian gland orifices” can be recognized, preferably improvements in “signs/findings on abnormalities around the meibomian gland orifices such as vasodilation” and “signs/findings on obstruction at the meibomian gland orifices” can be recognized.

[0068] Improvement in “signs/findings on abnormalities around the meibomian gland orifices such as vasodilation” means that, for example, telangiectasia around the meibomian gland orifices is suppressed, etc.

35 [0069] In the present invention, “suppressing telangiectasia around the meibomian gland orifices” means that, for example, “signs/findings on abnormalities around the meibomian gland orifices signs/findings of vasodilation” is to be resolved.

[0070] Improvement in “signs/findings on obstruction at the meibomian gland orifices” means that, for example, obstruction of the meibomian gland orifices is suppressed.

[0071] In the present invention, “suppressing obstruction of the meibomian gland” means that, for example, “signs/findings on obstruction at the meibomian gland orifices” is to be resolved.

[0072] In the present invention, the present agent or the present composition may contain an active ingredient other than the present compound or a pharmaceutically acceptable salt thereof, and may contain the present compound or a pharmaceutically acceptable salt thereof as a sole active ingredient.

[0073] In the present invention, the present agent or the present composition can be administered to the patient by, for example, orally or parenterally, and preferably administered by parenterally. As a parenteral administration form, there may be mentioned instillation administration (including instillation of an ophthalmic ointment), subconjunctival administration, administration to the interior of the conjunctival sac, administration under the Tenon’s capsule, etc., and instillation administration is particularly preferred.

[0074] In addition, the parenteral administration form may also contain, for example, dermal administration, and in the dermal administration according to the present invention, administration to the eyelid skin is particularly preferred.

[0075] In the present invention, the preparation containing the compound of the present invention or pharmaceutically acceptable salt thereof as an active ingredient is formulated into a dosage form suitable for administration with a pharmaceutically acceptable additive, if necessary. The dosage form suitable for the oral administration may be mentioned, for example, capsules, fine granules, granules, powders, pills, tablets, etc. Also, the dosage form suitable for parenteral administration may be mentioned, for example, eye drops, ophthalmic ointments, ointments (excluding ophthalmic ointments), injections, a preparation for intraocular implant (including punctal plug), intercalating agents, plasters, gels, etc. Incidentally, these can be prepared by using usual techniques generally used in the field of the art. Moreover, the present compound or a pharmaceutically acceptable salt thereof can be made a formulation which is made like DDS (drug delivery system) such as a microsphere.

[0076] As stated above, the present agent or the present composition is preferably subjected to instillation administration, so that the preferred dosage form of the present agent or the present composition is eye drops or ophthalmic ointments, and eye drops are particularly preferred. Incidentally, when the present compound or a

pharmaceutically acceptable salt thereof is prepared as the eye drops, characteristics of the eye drops may be a dissolved type solution (dissolved type eye drops), a suspension (suspended type eye drops), or an emulsion (emulsion eye drops), preferably a suspension or an emulsion.

5 [0077] Also, as stated above, the present agent or the present composition is preferably subjected to administration to the eyelid skin, so that the preferred dosage form of the present agent or the present composition is ointments (excluding ophthalmic ointments).

[0078] To the present compound or a pharmaceutically acceptable salt thereof may be added by optionally selecting, for example, an excipient such as microcrystalline  
10 cellulose, lactose, glucose, D-mannitol, anhydrous dibasic calcium phosphate, starch and sucrose; a disintegrator such as carboxymethyl cellulose, calcium carboxymethyl cellulose, croscarmellose sodium, crospovidone, starch, partially pregelatinized starch and low substituted degree hydroxypropyl cellulose; a binder such as hydroxypropyl cellulose, ethyl cellulose, gum Arabic, starch, partially pregelatinized starch,  
15 polyvinylpyrrolidone and polyvinyl alcohol; a lubricant such as magnesium stearate, calcium stearate, talc, hydrated silicon dioxide and hydrogenated oil; a coating agent such as refined white sugar, hydroxypropylmethyl cellulose, hydroxypropyl cellulose, methyl cellulose, ethyl cellulose and polyvinylpyrrolidone; a corrigent such as citric acid, aspartame, ascorbic acid and menthol, to prepare capsules, fine granules, granules,  
20 powders, pills or tablets.

[0079] In the present invention, the eye drops can be prepared by selecting and using, for example, an oil component such as medium chain fatty acid triglyceride (MCT); an isotonicifier such as sodium chloride, potassium chloride and glycerin; a buffering agent such as sodium phosphate, sodium acetate and  $\epsilon$ -aminocaproic acid; a surfactant such as  
25 tyloxapol, poloxamer 188, polyoxyethylene sorbitan monooleate, polyoxyl 40 stearate and polyoxyethylene hardened castor oil; a stabilizer such as sodium citrate and sodium edetate; a preservative such as benzalkonium chloride and paraben, depending on necessity. A pH of the eye drops may be within the range acceptable for an ophthalmic preparation, and it is usually preferred in the range of 4 to 8.

30 [0080] In the present invention, ophthalmic ointments or ointments (excluding ophthalmic ointments) can be prepared by using a generally used base, for example, white petrolatum, liquid paraffin, etc.

[0081] In the present invention, the injections can be prepared by selecting and using, for example, an isotonicifier such as sodium chloride; a buffering agent such as sodium  
35 phosphate; a surfactant such as polyoxyethylene sorbitan monooleate; a thickener such as methyl cellulose, depending on necessity.

[0082] In the present invention, the preparation for intraocular implant can be prepared by using, for example, a biodegradable polymer such as a polylactic acid, a polyglycolic acid, a lactic acid-glycolic acid copolymer and a hydroxypropyl cellulose.

5 [0083] In the present invention, the intercalating agents can be prepared by pulverizing and mixing, for example, a biodegradable polymer such as hydroxypropyl cellulose, hydroxypropylmethyl cellulose, carboxyvinyl polymer and polyacrylic acid, with the present compound, and compression-molding the powder, and if necessary, an excipient, a binder, a stabilizer and/or a pH adjuster may be used.

10 [0084] In the present invention, an administration dose of the present compound or a pharmaceutically acceptable salt thereof may be optionally changed depending on a dosage form, severity of the symptoms, an age or a body weight of a patient to be administered, and a judgment of a doctor, etc., and, for example, it can be generally administered to an adult person per a day of 0.000001 to 1,000 mg once or divided into several times.

15 [0085] When the present compound or a pharmaceutically acceptable salt thereof is to be applied to the eyes as eye drops, for example, the eye drops with a concentration of the active ingredient of 0.0001 to 1% (w/v), preferably 0.001 to 1% (w/v), more preferably 0.01 to 0.5% (w/v) can be applied to the eyes of an adult person once a day or divided into several times, preferably 1 to 2 times per a day, more preferably once a  
20 day.

[0086] When the present compound or a pharmaceutically acceptable salt thereof is to be applied to the eyes (instillation) as ophthalmic ointments, or to be administered to the eyelid skin as ointments (excluding ophthalmic ointments), for example, the ophthalmic ointment with a concentration of the active ingredient of 0.0001 to 1% (w/w), preferably  
25 0.001 to 1% (w/w), more preferably 0.01 to 0.5% (w/w) can be applied to the eyes (instillation) or administered to the eyelid skin of an adult person once a day or divided into several times.

[0087] When the present compound or a pharmaceutically acceptable salt thereof is administered as the injections, for example, the injection containing 0.000001 to 1,000  
30 mg of an active ingredient can be administered to an adult person per a day once or divided into several times.

[0088] When the present compound or a pharmaceutically acceptable salt thereof is administered as the preparation for intraocular implant, for example, the preparation for intraocular implant containing 0.000001 to 1,000 mg of an active ingredient can be  
35 implanted to an adult person.

[0089] When the present compound or a pharmaceutically acceptable salt thereof is

administered as the intercalating agents, for example, the intercalating agent containing 0.000001 to 1,000 mg of an active ingredient can be intercalated to an adult person.

[0090] In the following, the results of Pharmacological tests and Preparation examples are shown, and these examples are intended to better understand the present invention and not to limit the scope of the present invention.

#### EXAMPLES

[0091] [Test using complete Freund's adjuvant administered rabbit]

In a complete Freund's adjuvant administered rabbit, telangiectasia around the meibomian gland orifices, and obstruction at the meibomian gland orifices, which are similar to the signs/findings of MGD can be recognized. Effects of sirolimus which was a representative example of the present compound on the telangiectasia and obstruction were investigated (see JP 2014-024835A).

[0092] (Sample preparation)

0.1% (w/v) sirolimus suspension: it was prepared by suspending in 0.01% (w/v) hydroxypropylmethyl cellulose. Incidentally, with regard to sirolimus, that purchased from LKT Laboratories, Inc., (Catalog number: R0161) was used (which is the same in the following Examples).

[0093] (Test method)

10  $\mu$ L of the complete Freund's adjuvant was administered to the right upper eyelid (3 portions) of about 2 kg of male Japanese white rabbits, respectively. After 4 days from provocation, around the meibomian gland orifices of the right upper eyelid was observed by using slit lamp, a score of telangiectasia and a score of obstruction at the meibomian gland orifices were judged.

[0094] Incidentally, with regard to the score of telangiectasia, in accordance with the criteria in the following Tables 1 and 2, the eyelid margin of the upper eyelid was equally divided to three fractions of the ear side, the center portion and the nose side, the score of telangiectasia around the meibomian gland orifices was judged with regard to the respective fractions, and the sum of the scores of the three fractions was calculated as a score per one eye. Here, the presence or absence of dilation of the capillaries was judged by the state whether the capillaries which cannot be usually recognized with naked eyes can be recognized or not as a result of dilation of the diameter of the blood vessel. Also, the presence or absence of obstruction of the meibomian gland orifices was judged by the state whether the meibomian gland orifices are in a turbid or not.

[0095] [Table 1]

Score	State of telangiectasia around meibomian gland orifices
0	No telangiectasia around meibomian gland orifices was recognized in opened state of eyelid
1	Dilation of several capillaries around meibomian gland orifices was recognized in opened state of eyelid
2	Moderate dilation of capillaries or slight redness around meibomian gland orifices was recognized in opened state of eyelid
3	Redness and advanced dilation of capillaries around meibomian gland orifices were recognized in opened state of eyelid

[0096] [Table 2]

Score	Grade	State of obstruction at meibomian gland orifices
0	None	No signs/findings
1	Slight	Obstructions of 3 or less
2	Moderate	Obstructions of 4 to 6
3	Advanced	Obstructions of 7 or more

[0097] The rabbits were grouped into a physiological saline solution-administered group and a 0.1% sirolimus suspension-administered group, and tested with 7 or 8 eyes in each group so that fluctuation of the average value of the respective scores became small. After 5<sup>th</sup> day from provocation, a physiological saline solution (50  $\mu$ L/eye, twice a day) or a 0.1% sirolimus suspension (50  $\mu$ L/eye, twice a day) was dropped into the right eye for 10 days. After 11<sup>th</sup> and 15<sup>th</sup> days from provocation, around the meibomian gland orifices of the right upper eyelid was observed by using a slit lamp, whereby the score of telangiectasia was judged and the number of the obstructive orifices was measured.

[0098] (Test results)

Test results (score of telangiectasia and average value of number of obstructive orifices) are shown in Table 3.

[0099] [Table 3]

	Score of telangiectasia		Score of obstruction of orifices	
	11 <sup>th</sup> day after	15 <sup>th</sup> day after	11 <sup>th</sup> day after	15 <sup>th</sup> day after
Physiological saline-administered group (N=8)	5.0	5.1	3.8	4.4
0.1% Sirolimus suspension-administered group (N=7)	2.1	2.1	2.3	1.3

[0100] (Consideration)

0.1% sirolimus suspension decreased the score of telangiectasia around the meibomian gland orifices and the number of the obstructive orifices. That is, since

sirolimus resolved these meibomian gland abnormalities, it could be shown that the present compound or a pharmaceutically acceptable salt thereof has a treatment effect to MGD.

[0101] [Test using HR-AD feed fed hairless mouse]

5 It has been known that a HR-AD feed is fed to Hos: HR-1 series hairless mouse, obstruction of the meibomian gland orifices similar to the signs/findings of MGD could be recognized (Japanese Patent Application No. 2012-137778). Thus, effects of sirolimus and deforolimus which are representative examples of the present compounds on the obstruction were investigated.

10 [0102] (Sample preparation)

0.1% (w/v) sirolimus suspension: it was prepared by suspending in 0.01% (w/v) hydroxypropylmethyl cellulose.

0.1% (w/v) deforolimus suspension: it was prepared by suspending in 0.01% (w/v) hydroxypropylmethyl cellulose. Incidentally, deforolimus used in the present  
15 test was prepared according to the method described in JP 2005-516065A.

[0103] (Test method)

Six-weeks old male Hos: HR-1 series hairless mice were grouped into 5 mice of a normal feed fed group (CRF-1 feed, available from Oriental Yeast Co., Ltd.) and 15 mice of a HR-AD feed fed group (available from Nosan Corporation), and fed the  
20 normal feed or the HR-AD feed, respectively, which were spontaneously taken. After 28<sup>th</sup> day from initiation of the feeding, the meibomian gland orifices were observed by using a slit lamp, and a number of obstructive orifices among the eight meibomian gland orifices at the center of the upper eyelid was measured. Incidentally, the presence or absence of obstruction of the meibomian gland orifices was judged by the  
25 state whether the meibomian gland orifices were in a turbid and raised or not (see JP 2014-023526A).

[0104] Fifteen mice of the HR-AD feed fed group were grouped into a physiological saline solution-administered group, a 0.1% sirolimus suspension-administered group and a 0.1% deforolimus suspension-administered group (5 mice in each group), and  
30 from 29<sup>th</sup> day from initiation of the feeding, the physiological saline solution (2  $\mu$ L/eye, twice a day), the 0.1% sirolimus suspension (2  $\mu$ L/eye, twice a day) or the 0.1% (w/v) deforolimus suspension (2  $\mu$ L/eye, twice a day) were dropped to the eye for 28 days. At 42<sup>nd</sup> day and 56<sup>th</sup> day after initiation of the feeding, a number of obstructions of the meibomian gland orifices was measured by using a slit lamp.

35 [0105] (Test results)

Test results are shown in Table 4.

[0106] [Table 4]

	Number of obstructive orifices	
	42 <sup>nd</sup> day after initiation of the feeding	56 <sup>th</sup> day after initiation of the feeding
Non-treatment group (Normal feed fed group)	1.2	0.7
Physiological saline solution-administered group	5.8	6.3
0.1% Sirolimus suspension-administered group	0.7	0.0
0.1% Deforolimus suspension-administered group	2.7	2.2

## [0107] (Consideration)

0.1% sirolimus suspension and 0.1% deforolimus suspension decreased the number of the obstructions of the meibomian gland orifices. That is, since sirolimus and deforolimus resolved meibomian gland abnormalities, it could be shown that the present compounds or a pharmaceutically acceptable salt thereof have a treatment effect to MGD.

## [0108] [Preparation examples]

The drugs of the present invention are more specifically explained by referring to Preparation examples, but the present invention is not limited by these Preparation examples alone.

## [0109] Prescription example 1: Eye drop (0.1% (w/v))

	In 100 ml	
	Sirolimus	0.1 g
15	Medium chain fatty acid triglyceride (MCT)	7.5 g
	Benzalkonium chloride	0.2 g
	Tyloxapol	1.2 g
	Poloxamer 188	1 g
	Glycerin	22.5 g
20	Sterile purified water	q.s.

[0110] To sterile purified water are added sirolimus and the above-mentioned components other than these, and the mixture is thoroughly mixed to prepare an eye drop. Characteristics of the present eye drop is an emulsion. By changing the formulation amount of the sirolimus, eye drops with the concentration of the sirolimus of 0.05% (w/v), 0.5% (w/v) or 1% (w/v) can be prepared.

## [0111] Prescription example 2: Eye drop (0.1% (w/v))

	In 100 ml	
	Deforolimus	0.1 g
	Medium chain fatty acid triglyceride (MCT)	7.5 g

	Benzalkonium chloride	0.2 g
	Tyloxapol	1.2 g
	Poloxamer 188	1 g
	Glycerin	22.5 g
5	Sterile purified water	Suitable amount

[0112] To sterile purified water are added deforolimus and the above-mentioned components other than these, and the mixture is thoroughly mixed to prepare an eye drop. Characteristics of the present eye drop is an emulsion. By changing the formulation amount of the deforolimus, eye drops with the concentration of the deforolimus of 0.05% (w/v), 0.5% (w/v) or 1% (w/v) can be prepared.

10 [0113] Prescription example 3: Ophthalmic ointment or ointment (excluding ophthalmic ointment) (1% (w/w))

	In 100 g	
	Sirolimus	1 g
15	Liquid paraffin	q.s.
	White petrolatum	q.s.

[0114] Sirolimus is added to uniformly melt white petrolatum and liquid paraffin, and the mixture is thoroughly mixed and then gradually cooled to prepare an ophthalmic ointment or an ointment (excluding an ophthalmic ointment). By changing the formulation amount of the sirolimus, an ophthalmic ointment or an ointment (excluding an ophthalmic ointment) with the concentration of the sirolimus of 0.05% (w/w), 0.1% (w/w) or 0.5% (w/w) can be prepared.

[0115] Prescription example 4: Ophthalmic ointment or ointment (excluding ophthalmic ointment) (1% (w/w))

25	In 100 g	
	Deforolimus	1 g
	Liquid paraffin	q.s.
	White petrolatum	q.s.

[0116] Deforolimus is added to uniformly melt white petrolatum and liquid paraffin, and the mixture is thoroughly mixed and then gradually cooled to prepare an ophthalmic ointment or an ointment (excluding an ophthalmic ointment). By changing the formulation amount of the deforolimus, an ophthalmic ointment or an ointment (excluding an ophthalmic ointment) with the concentration of the deforolimus of 0.05% (w/w), 0.1% (w/w) or 0.5% (w/w) can be prepared.

35

UTILIZABILITY IN INDUSTRY

[0117] As can be clearly seen from the results of the tests using complete Freund's adjuvant administered rats and HR-AD feed fed hairless mice, sirolimus and deforolimus suppressed obstruction of the meibomian gland, so that the present compounds or a pharmaceutically acceptable salt thereof are useful as a prophylactic and/or therapeutic agent for MGD.

5

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. Use of sirolimus, or a pharmaceutically acceptable salt thereof, as the sole active ingredient in the manufacture of a medicament for suppressing obstruction of meibomian gland and/or telangiectasia around meibomian gland orifices, wherein the medicament is formulated for instillation administration in a form of an eye drop.
2. The use according to claim 1, wherein the obstruction of meibomian gland and/or the telangiectasia around meibomian gland orifices is a cause of dry eye.
3. The use according to claim 1 or 2, wherein the eye drop is a suspension or an emulsion.
4. The use according to any one of claims 1 to 3, wherein the medicament is formulated for instillation administration 1 to 2 times per day.
5. Use of deforolimus, or a pharmaceutically acceptable salt thereof, as the sole active ingredient in the manufacture of a medicament for suppressing obstruction of meibomian gland and/or telangiectasia around meibomian gland orifices, wherein the medicament is formulated for instillation administration in a form of an eye drop.

6. The use according to claim 5, wherein the obstruction of meibomian gland and/or the telangiectasia around meibomian gland orifices is a cause of dry eye.

7. The use according to claim 5 or 6, wherein the eye drop is a suspension or an emulsion.

8. The use according to any one of claims 5 to 7, wherein the medicament is formulated for instillation administration 1 to 2 times per day.

9. A pharmaceutical composition for suppressing obstruction of meibomian gland and/or telangiectasia around meibomian gland orifices comprising sirolimus, or a pharmaceutically acceptable salt thereof, as the sole active ingredient, wherein the composition is formulated for instillation administration in a form of an eye drop.

10. The pharmaceutical composition according to claim 9, wherein the obstruction of meibomian gland and/or the telangiectasia around meibomian gland orifices is a cause of dry eye.

11. The pharmaceutical composition according to claim 9 or 10, wherein the eye drop is a suspension or an emulsion.

12. The pharmaceutical composition according to any one of claims 9 to 11, wherein the composition is formulated for instillation administration 1 to 2 times per day.

13. A pharmaceutical composition for suppressing obstruction of meibomian gland and/or telangiectasia around meibomian gland orifices comprising deforolimus, or a pharmaceutically acceptable salt thereof, as the sole active ingredient, wherein the composition is formulated for instillation administration in a form of an eye drop.

14. The pharmaceutical composition according to claim 13, wherein the obstruction of meibomian gland and/or the telangiectasia around meibomian gland orifices is a cause of dry eye.

15. The pharmaceutical composition according to claim 13 or 14, wherein the eye drop is a suspension or an emulsion.

16. The pharmaceutical composition according to any one of claims 13 to 15, wherein the composition is formulated for instillation administration 1 to 2 times per day.

[Formula 1]

