

(19)



INTELLECTUAL PROPERTY
OFFICE OF SINGAPORE

(11) Publication number:

SG 179402 A1

(43) Publication date:

27.04.2012

(51) Int. Cl:

;

(12)

Patent Application

(21) Application number: 2012012126

(71) Applicant:

ONCOTHERAPY SCIENCE, INC. 2-1,
SAKADO 3-CHOME, TAKATSU-KU,
KAWASAKI-SHI, KANAGAWA 2130012
JP

(22) Date of filing: 21.02.2008

(72) Inventor:

TAKUYA TSUNODA C/O
ONCOTHERAPY SCIENCE, INC., 2-1,
SAKADO 3-CHOME, TAKATSU-KU,
KAWASAKI-SHI, KANAGAWA 2130012
JP

(30) Priority: US 60/902,949 21.02.2007

RYUJI OHSAWA C/O ONCOTHERAPY
SCIENCE, INC., 2-1, SAKADO 3-
CHOME, TAKATSU-KU, KAWASAKI-SHI,
KANAGAWA 2130012 JP

(54) Title:

PEPTIDE VACCINES FOR CANCERS EXPRESSING
TUMOR-ASSOCIATED ANTIGENS

(57) Abstract:

83 Peptide Vaccines For Cancers Expressing Tumor-Associated Antigens Abstract The present invention provides peptides having an amino acid sequence as set forth in SEQ ID NO: SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288, as well as peptides having the above-mentioned amino acid sequences in which 1, 2, or several (e.g., up to 5) amino acids are substituted, deleted, or added, provided the peptides possess cytotoxic T cell inducibility. The present invention also provides drugs for treating or preventing a disease associated with over-expression of the CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers containing as an active ingredient one or more of these peptides. The peptides of the present invention find further utility as vaccines. No suitable figure.

Peptide Vaccines For Cancers Expressing Tumor-Associated Antigens**Abstract**

The present invention provides peptides having an amino acid sequence as set forth in SEQ ID NO: SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288, as well as peptides having the above-mentioned amino acid sequences in which 1, 2, or several (e.g., up to 5) amino acids are substituted, deleted, or added, provided the peptides possess cytotoxic T cell inducibility. The present invention also provides drugs for treating or preventing a disease associated with over-expression of the CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers containing as an active ingredient one or more of these peptides. The peptides of the present invention find further utility as vaccines.

No suitable figure.

Description

PEPTIDE VACCINES FOR CANCERS EXPRESSING TUMOR-ASSOCIATED ANTIGENS

Technical Field

[0001] The present application claims the benefit of U.S. Provisional Application No. 60/902,949, filed February 21, 2007, the entire disclosure of which is hereby incorporated herein by reference for all purposes.

[0002] The present invention relates to the field of biological science, more specifically to the field of cancer therapy. In particular, the present invention relates to novel immunogenic peptides that serve as extremely effective as cancer vaccines, and drugs for treating and preventing tumors containing such peptides.

Background Art

[0003] It has been demonstrated that CD8⁺ cytotoxic T lymphocytes (CTLs) recognize epitope peptides derived from tumor-associated antigens (TAAs) presented on MHC class I molecules, and subsequently lyse the tumor cells. Since the discovery of the MAGE family as the first example of TAAs, many other TAAs have been discovered using immunological approaches (Boon T. (1993) *Int J Cancer* 54: 177-80.; Boon T. et al., (1996) *J Exp Med* 183: 725-9.; van der Bruggen P et al., (1991) *Science* 254: 1643-7.; Brichard V et al., (1993) *J Exp Med* 178: 489-95.; Kawakami Y et al., (1994) *J Exp Med* 180: 347-52.). Some of them are now in clinical development as targets of immunotherapy. TAAs discovered to date include MAGE (van der Bruggen P et al., (1991) *Science* 254: 1643-7.), gp100 (Kawakami Y et al., (1994) *J Exp Med* 180: 347-52.), SART (Shichijo S et al., (1998) *J Exp Med* 187:277-88.), and NY-ESO-1 (Chen Y.T. et al., (1997) *Proc. Natl. Acad. Sci. USA*, 94: 1914-8.). On the other hand, certain gene products demonstrated to be somewhat specifically over-expressed in tumor cells have been shown to be recognized as targets for inducing cellular immune responses. Such gene products include p53 (Umano Y et al., (2001) *Br J Cancer*, 84:1052-7.), HER2/neu (Tanaka H et al., (2001) *Br J Cancer*, 84: 94-9.), CEA (Nukaya I et al., (1999) *Int. J. Cancer* 80, 92-7.) and the like.

[0004] Despite significant progress in basic and clinical research concerning TAAs (Rosenberg SA et al., (1998) *Nature Med*, 4: 321-7.; Mukherji B. et al., (1995) *Proc Natl Acad Sci USA*, 92: 8078-82.; Hu X et al., (1996) *Cancer Res*, 56: 2479-83.), only a very limited number of candidate TAAs suitable for treatment of cancers are presently available. TAAs that are abundantly expressed in cancer cells, and whose expression is restricted to cancer cells, would be promising candidates as immuno-therapeutic targets.

[0005] Both HLA-A24 and HLA-A0201 are common HLA alleles in the Japanese and Caucasian populations (Date Y et al., (1996) *Tissue Antigens* 47: 93-101.; Kondo A et al., (1995) *J Immunol* 155: 4307-12.; Kubo RT et al., (1994) *J Immunol* 152: 3913-24.; Imanishi et al., Proceeding of the eleventh International Histocompatibility Workshop and Conference Oxford University Press, Oxford, 1065 (1992); Williams F et al., (1997) *Tissue Antigen* 49: 129-33.). Thus, antigenic peptides of cancers presented by these HLA alleles may find particular utility in the treatment of cancers among Japanese and Caucasian patients. Further, it is known that the induction of low-affinity CTL in vitro usually results from exposure to high concentrations of peptides, generating a high level of specific peptide/MHC complexes on antigen-presenting cells (APCs), which will effectively activate these CTL (Alexander-Miller et al., (1996) *Proc Natl Acad Sci USA* 93: 4102-7.).

[0006] Recently, HLA class I-binding peptide sequence can be expected using algorithms (Journal of Immunological Methods, (1995), Vol.185, pp.181-190, J. Immunol., (1994), Vol.152, pp.163-175, protein science, (2000), Vol.9, pp.1838-1846). However, it is hard to say that the expected epitope peptide can be cut to the size and expressed on the target cell surface with HLA molecule and recognized by CTL. Moreover, the algorithm, for example BIMAS (http://bimas.dcrf.nih.gov/cgi-bin/molbio/ken_parker_comboform) (Parker KC, et al., (1994) *J Immunol*;152(1):163-75.; Kuzushima K, et al., (2001) *Blood*;98(6):1872-81.) can suggest the HLA molecule-binding peptide, but the suggested peptide is not so rigorous (Bachinsky MM, et. al., *Cancer Immun*. 2005 Mar 22;5:6.). Thus TAA screening still remains a lot of challenges and difficulties.

[0007] Recent developments in cDNA microarray technologies have enabled the construction of comprehensive profiles of gene expression in malignant cells as compared to normal cells (Okabe, H. et al., (2001) *Cancer Res.*, 61, 2129-37.; Lin YM. et al., (2002) *Oncogene*, 21;4120-8.; Hasegawa S. et al., (2002) *Cancer Res* 62:7012-7.). This approach enables a more thorough understanding of the complex nature of cancer cells and the mechanisms of carcinogenesis and facilitates the identification of genes whose expression is deregulated in tumors (Bienz M. et al., (2000) *Cell* 103, 311-20.). Among the transcripts identified as up-regulated in cancers, CDH3 (GenBank Accession No. NM_001793; SEQ ID Nos.1, 2), EPHA4 (GenBank Accession No. L36645; SEQ ID Nos.3, 4), ECT2 (GenBank Accession No. AY376439; SEQ ID Nos.5, 6), HIG2 (GenBank Accession No. NM_013332; SEQ ID Nos.7, 8) INHBB (GenBank Accession No. NM_002193; SEQ ID Nos.9, 10), KIF20A (GenBank Accession No. NM_005733; SEQ ID Nos.11, 12), KNTC2 (GenBank Accession No. AF017790; SEQ ID Nos.13, 14), TTK (GenBank Accession No. NM_003318; SEQ ID Nos.15, 16) and URLC10 (GenBank Accession No. NM_017527; SEQ ID Nos.17, 18) have been

recently discovered. The entire contents of the references are incorporated by reference herein. These genes are of particular interest to the present inventors, being specifically up-regulated in tumor cells of the various cancer tissues of the cases analyzed (see below). Thus, immunogenic peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 may find utility in selectively killing tumor cells that express such antigens. The present invention addresses these and other needs.

[0008] Since cytotoxic drugs, such as M-VAC, often cause severe adverse reactions, it is clear that thoughtful selection of novel target molecules on the basis of well-characterized mechanisms of action should be very helpful in the development of effective anti-cancer drugs having a minimized risk of side effects. Toward this goal, expression profile analyses were previously performed on various cancers and normal human tissue. Such studies led to the discovery of multiple genes that are specifically over-expressed in cancer (Lin YM, et al., *Oncogene*. 2002 Jun 13;21:4120-8.; Kitahara O, et al., *Cancer Res.* 2001 May 1;61:3544-9.; Suzuki C, et al., *Cancer Res.* 2003 Nov 1;63:7038-41.; Ashida S, *Cancer Res.* 2004 Sep 1;64:5963-72.; Ochi K, et al., *Int J Oncol.* 2004 Mar;24(3):647-55.; Kaneta Y, et al., *Int J Oncol.* 2003 Sep;23:681-91.; Obama K, *Hepatology*. 2005 Jun;41:1339-48.; Kato T, et al., *Cancer Res.* 2005 Jul 1;65:5638-46.; Kitahara O, et al., *Neoplasia*. 2002 Jul-Aug;4:295-303.; Saito-Hisaminato A et al., *DNA Res* 2002, 9: 35-45.). Examples of such genes identified as over-expressed in various cancers include, but are not limited to, CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10. CDH3 has been previously identified as over-expressed in bladder cancer, cervical cancer, cholangiocellular carcinoma, colorectal cancer, endometriosis, gastric cancer, diffuse-type gastric cancer, non-small cell lung cancer (NSCLC), pancreatic cancer, soft tissue tumor and testicular tumor. EPHA4 has been identified in bladder cancer, cervical cancer, cholangiocellular carcinoma, endometriosis, diffuse-type gastric cancer, ovarian cancer, pancreatic cancer, prostate cancer and soft tissue tumor. ECT2 has been identified in bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, chronic myeloid leukemia (CML), colorectal cancer, esophageal cancer, NSCLC, lymphoma, prostate cancer, renal carcinoma and small cell lung cancer (SCLC). HIG2 has been identified in renal carcinoma and SCLC. INHBB has been identified in cholangiocellular carcinoma, esophageal cancer, NSCLC, renal carcinoma, SCLC and soft tissue tumor. KIF20A has been identified in bladder cancer, breast cancer, cholangiocellular carcinoma, esophageal cancer, NSCLC, pancreatic cancer, prostate cancer, renal carcinoma and SCLC. KNTC2 has been identified in bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, esophageal cancer, NSCLC, lymphoma, osteosarcoma, ovarian

cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC and soft tissue tumor. TTK has been identified in bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, esophageal cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, prostate cancer, SCLC and soft tissue tumor. URLC10 has been identified in bladder cancer, cervical cancer, cholangiocellular carcinoma, esophageal cancer, gastric cancer, NSCLC, osteosarcoma, pancreatic cancer and SCLC.

[0009] Summary of the Invention

The present invention is based in part on the discovery of the applicable targets of immunotherapy. Because TAAs have often no immunogenicity, the discovery of appropriate targets is of extreme importance. As noted above, CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 have been identified as up-regulated in various cancers. More particularly, these genes were identified using gene expression profiling with a genome-wide cDNA microarray. As discussed above, expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 has been shown to be specifically up-regulated in various tumor cells, from pancreatic cancer cells to renal cell carcinomas. As described in Table 1, CDH3 expression is validly elevated in 26 out of 34 bladder cancer, 17 out of 19 cervical cancer, all of 19 cholangiocellular carcinoma, 30 out of 34 colorectal cancer, 20 out of 21 endometriosis, 13 out of 20 gastric cancer, 7 out of 8 diffuse-type gastric cancer, 36 out of 37 NSCLC, all of 16 pancreatic cancer, all of 21 soft tissue tumor and all of 10 testicular tumor.

[0010] Table 1 further demonstrates that:

EPHA4 expression is validly elevated in 14 out of 34 bladder cancer, 8 out of 14 cervical cancer, 10 out of 25 cholangiocellular carcinoma, 5 out of 15 endometriosis, 5 out of 8 diffuse-type gastric cancer, all of 5 ovarian cancer, all 14 pancreatic cancer, 20 out of 51 prostate cancer and 14 out of 23 soft tissue tumor.

ECT2 expression is validly elevated in 17 out of 19 bladder cancer, 5 out of 12 breast cancer, all of 14 cervical cancer, all of 13 cholangiocellular carcinoma, all of 5 CML, 7 out of 8 colorectal cancer, 12 out of 16 esophageal cancer, 6 out of 16 NSCLC, 8 out of 10 lymphoma, 1 out of 1 pancreatic cancer, 10 out of 13 prostate cancer, 3 out of 6 renal carcinoma and 12 out of 13 SCLC cancer.

HIG2 expression is validly elevated in 19 out of 20 renal cancer and 7 out of 9 soft tissue tumor.

INHBB expression is validly elevated in 10 out of 21 cholangiocellular carcinoma, all of 12 esophageal cancer, 10 out of 13 NSCLC, 22 out of 24 renal carcinoma, 8 out of 14 SCLC cancer and 45 out of 49 soft tissue tumor.

KIF20A expression is validly elevated in all of 31 bladder cancer, 38 out of 61 breast

cancer, 10 out of 11 cholangiocellular carcinoma, 7 out of 19 esophageal cancer, 21 out of 22 NSCLC, all of 6 ovarian cancer, 17 out of 36 prostate cancer, 6 out of 11 renal carcinoma and all of 15 SCLC.

KNTC2 expression is validly elevated in 30 out of 32 bladder cancer, 47 out of 56 breast cancer, all of 10 cervical cancer, 16 out of 22 cholangiocellular carcinoma, 17 out of 37 CML, 3 out of 10 colorectal cancer, 11 out of 46 esophagus cancer, 15 out of 19 NSCLC, 7 out of 8 lymphoma, 20 out of 24 osteosarcoma, 3 out of 5 ovarian cancer, all of 2 pancreatic cancer, 15 out of 37 prostate cancer, 14 out of 19 renal carcinoma, all of 15 SCLC and 40 out of 59 soft tissue tumor.

TTK expression is validly elevated in all of 27 bladder cancer, 25 out of 30 breast cancer, 15 out of 16 cervical cancer, all of 10 cholangiocellular carcinoma, 5 out of 7 CML, 6 out of 10 colorectal cancer, 24 out of 44 esophageal cancer, 8 out of 15 liver cancer, all of 12 NSCLC, all of 6 lymphoma, 13 out of 16 osteoblastoma, 12 out of 17 prostate cancer, all of 15 SCLC and 16 out of 33 soft tissue tumor.

URLC10 expression is validly elevated in all of 29 bladder cancer, 15 out of 16 cervical cancer, all of 7 cholangiocellular carcinoma, 7 out of 19 esophageal cancer, all of 3 gastric cancer, 24 out of 27 NSCLC, 15 out of 19 osteosarcoma, 4 out of 5 pancreatic cancer, 33 out of 43 soft tissue tumor.

[0011] The present invention is based, at least in part, on the identification of specific epitope peptides of the gene products of these genes (CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10) which possess the ability to induce cytotoxic T lymphocytes (CTLs) specific to the corresponding molecules. As discussed in detail below, Peripheral Blood Mononuclear Cells (PBMC) of healthy donor were stimulated using HLA-A*2402 or HLA-A*0201 binding candidate peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10. CTL clones and/or lines were then established with specific cytotoxicity against the HLA-A24 or HLA-A2 positive target cells pulsed with each of the candidate peptides. These results demonstrate that these peptides are HLA-A24 or HLA-A2 restricted epitope peptides that can induce potent and specific immune responses against cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10.

[0012] Accordingly, the present invention provides methods for treating or preventing a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10, e.g. cancer. Such methods involve the step of administering to a subject in need thereof a CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10 polypeptides of the invention. Administration of such peptide(s) results in the induction of anti-tumor immunity is induced by the administration of these polypeptides. Thus, the present invention provides methods for inducing anti-tumor immunity in a subject, such methods involving the step of admin-

istering to the subject the CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10 polypeptides, as well as pharmaceutical compositions for treating or preventing a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancer, that include the CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 polypeptides. Examples of such cancers include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0013]

The present invention further provides methods for preventing post-surgery recurrence of the disease mentioned above.

Regarding the specific aims and objectives recited above, it will be understood by those skilled in the art that one or more aspects of this invention can meet certain objectives, while one or more other aspects can meet certain other objectives. Each objective may not apply equally, in all its respects, to every aspect of this invention. As such, the objects herein can be viewed in the alternative with respect to any one aspect of this invention.

[0014]

Additional objects and features of the invention will become more fully apparent when the following detailed description is read in conjunction with the accompanying figures and examples. However, it is to be understood that both the foregoing summary of the invention and the following detailed description are of preferred embodiments, and not restrictive of the invention or other alternate embodiments of the invention. In particular, while the invention is described herein with reference to a number of specific embodiments, it will be appreciated that the description is illustrative of the invention and is not construed as limiting of the invention. Various modifications and applications may occur to those who are skilled in the art, without departing from the spirit and the scope of the invention, as described by the appended claims. Likewise, other objects, features, benefits and advantages of the present invention will be apparent from this summary and certain embodiments described below, and will be readily apparent to those skilled in the art. Such objects, features, benefits and advantages will be apparent from the above in conjunction with the accompanying examples, data, figures and all reasonable inferences to be drawn therefrom, alone or with consideration of the references incorporated herein.

Brief Description of Drawings

[0015]

Various aspects and applications of the present invention will become apparent to the skilled artisan upon consideration of the brief description of the figures and the

detailed description of the present invention and its preferred embodiments which follows:

[fig. 1-1]Figure 1 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that CDH3-A24-10-332 (SEQ ID NO: 34), CDH3-A24-10-470 (SEQ ID NO: 358), CDH3-A24-9-513 (SEQ ID NO: 19), CDH3-A24-9-406 (SEQ ID NO: 22), CDH3-A24-10-807 (SEQ ID NO: 30) and CDH3-A24-10-655 (SEQ ID NO: 344) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA-A*2402. "b" depicts the CTL-inducing ability of CDH3-A24-10-332 (SEQ ID NO: 34). CDH3-A24-10-332 (SEQ ID NO: 34) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line that was established from the positive well #4 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "c" depicts the CTL-inducing ability of CDH3-A24-10-470 (SEQ ID NO: 358). CDH3-A24-10-470 (SEQ ID NO: 358) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line that was established from the positive well #4 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "d" depicts the CTL-inducing ability of CDH3-A24-9-513 (SEQ ID NO: 19). CDH3-A24-9-513 (SEQ ID NO: 19) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay. The well #6 shown in boxed wells in left panel demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL line that was established from the positive well #5 shown in boxed wells in middle panel, demonstrated the specific response against the target cells pulsed with the epitope peptide. "e" depicts the CTL-inducing ability of CDH3-A24-9-406 (SEQ ID NO: 22). CDH3-A24-9-406 (SEQ ID NO: 22) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line that was established from the positive well #2 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide.

[fig. 1-2]Figure 1 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that CDH3-A24-10-332 (SEQ ID NO: 34), CDH3-A24-10-470 (SEQ ID NO: 358), CDH3-A24-9-513 (SEQ ID NO: 19), CDH3-A24-9-406 (SEQ ID NO: 22), CDH3-A24-10-807 (SEQ ID NO: 30) and CDH3-A24-10-655 (SEQ ID NO: 344) show potent IFN-gamma production. "f" depicts the CTL-inducing ability of CDH3-A24-10-807 (SEQ ID NO: 30). CDH3-A24-10-807 (SEQ ID NO: 30) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and the clone were established from the positive

well #5 shown in boxed wells. The established CTL clone raised against the peptide demonstrated the specific CTL activity against COS7 transfected both full length of CDH3 gene and HLA-A24 molecule (lower right graph). On the other hand, COS7 transfected full length of CDH3 but not HLA-A24 and COS7 transfected HLA-A24 but not full length of CDH3 were prepared for the negative control. The CTL clone showed high specific CTL activity against COS7 that transfected both CDH3 and HLA-A24. "g" depicts the CTL-inducing ability of CDH3-A24-10-655 (SEQ ID NO: 344). CDH3-A24-10-655 (SEQ ID NO: 344) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and the clone were established from the positive well #1 shown in boxed wells. The established CTL clone raised against the peptide demonstrated the specific CTL activity against COS7 transfected both full length of CDH3 gene and HLA-A24 molecule (lower right graph). On the other hand, COS7 transfected full length of CDH3 but not HLA-A24 and COS7 transfected HLA-A24 but not full length of CDH3 were prepared for the negative control. The CTL clone showed high specific CTL activity against COS7 that transfected both CDH3 and HLA-A24.

[fig.2]Figure 2 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that Eph4-A24-9-453 (SEQ ID NO: 41), Eph4-A24-9-5 (SEQ ID NO: 44), Eph4-A24-9-420 (SEQ ID NO: 48), Eph4-A24-9-869 (SEQ ID NO: 46), Eph4-A24-10-24 (SEQ ID NO: 78) Eph4-A02-9-501 (SEQ ID NO: 376) and Eph4-A02-9-165 (SEQ ID NO: 379) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of Eph4-A24-9-453 (SEQ ID NO: 41). Eph4-A24-9-453 (SEQ ID NO: 41) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line that was established from the positive well #3 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "c" depicts the CTL-inducing ability of Eph4-A24-9-5 (SEQ ID NO: 44). Eph4-A24-9-5 (SEQ ID NO: 44) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #2 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "d" depicts the CTL-inducing ability of Eph4-A24-9-420 (SEQ ID NO: 48). Eph4-A24-9-420 (SEQ ID NO: 48) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay. The well #6 shown in boxed wells in upper panel demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover CTL line that was established from the positive well #6 shown in boxed wells in middle panel, demonstrated the

specific response against the target cells pulsed with the epitope peptide. "e" depicts the CTL-inducing ability of EphA4-A24-9-869 (SEQ ID NO: 46). EphA4-A24-9-869 (SEQ ID NO: 46) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line that was established from the positive well #5 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "f" depicts the CTL-inducing ability of EphA4-A24-10-24 (SEQ ID NO: 78). EphA4-A24-10-24 (SEQ ID NO: 78) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line that was established from the positive well #4 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "g" depicts the CTL-inducing ability of EphA4-A02-9-501 (SEQ ID NO: 376). EphA4-A02-9-501 (SEQ ID NO: 376) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone was established from the positive well #8 shown in boxed wells. Cytotoxic activity of the established CTL line against the target cells pulsed with the peptide was measured by Cr-release assay (CRA) (lower graph), and the CTL line had very potent specific cytotoxic activity against the target cells pulsed with the peptides. "h" depicts the CTL-inducing ability of EphA4-A02-9-165 (SEQ ID NO: 379). EphA4-A02-9-165 (SEQ ID NO: 379) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line was established from the positive well #3 shown in boxed wells. Cytotoxic activity of the established CTL line against target cells pulsed with peptide was measured by Cr-release assay (CRA) (right graph), and the CTL line had very potent specific cytotoxic activity against the target cells pulsed with the peptides.

[fig.3]Figure 3 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that ECT2-A24-9-515 (SEQ ID NO: 80), ECT2-A24-10-40 (SEQ ID NO: 100) and ECT2-A24-10-101 (SEQ ID NO: 101) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of ECT2-A24-9-515 (SEQ ID NO: 80). ECT2-A24-9-515 (SEQ ID NO: 80) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay. The well #5 and #7 shown in boxed wells in left panel demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL line that was established from the positive well #7 shown in boxed wells in second panel, demonstrated the specific response against the target cells pulsed with the epitope peptide. Cytotoxic activity of the CTL line against cancer cell line, TE6 endogenously expressing ECT2 and HLA-A24 was measured by Cr-release

assay (CRA), and the CTL clone had very potent cytotoxic activity against TE6. On the other hand, the effector cells did not demonstrate the cytotoxic activity of the CTL line against cancer cell line, TE5 expressing only ECT2 was not detected. "c" depicts the CTL-inducing ability of ECT2-A24-10-40 (SEQ ID NO: 100). ECT2-A24-10-40 (SEQ ID NO: 100) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and the clone were established from the positive well #2 shown in boxed wells. The established CTL clone raised against the peptide demonstrated specific CTL activity against COS7 transfected both full length of ECT2 gene and HLA-A24 molecule. On the other hand, COS7 transfected full length of ECT2 but not HLA-A24, COS7 transfected HLA-A24 and URLC10 gene as a substitute for full length of ECT2 and COS7 transfected HLA-A24 and pulsed with ECT2-10-101 were prepared for the negative control. The CTL clone showed high specific CTL activity against COS7 that transfected both ECT2 and HLA-A24. "d" depicts the CTL-inducing ability of ECT2-A24-10-101 (SEQ ID NO: 101). ECT2-A24-10-101 (SEQ ID NO: 101) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line were established from the positive well #1 shown in boxed wells. The established CTL line raised against the peptide demonstrated specific CTL activity against COS7 transfected both full length of ECT2 gene and HLA-A24 molecule. COS7 transfected full length of ECT2 but not HLA-A24, COS7 transfected HLA-A24 and URLC10 gene as substitute for full length of ECT2 and COS7 transfected HLA-A24 and pulsed with ECT2-10-40 were prepared for the negative control. The CTL clone showed high specific CTL activity against COS7 that transfected both ECT2 and HLA-A24.

[fig.4-1]Figure 4 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that HIG2-A24-9-19 (SEQ ID NO: 110), HIG2-A24-9-22 (SEQ ID NO: 111), HIG2-A24-9-8 (SEQ ID NO: 387), HIG2-A24-10-7 (SEQ ID NO: 112), HIG2-A24-10-18 (SEQ ID NO: 394), HIG2-A02-9-15 (SEQ ID NO: 116), HIG2-A02-9-4 (SEQ ID NO: 117) and HIG2-A02-10-8 (SEQ ID NO: 121) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of HIG2-A24-9-19 (SEQ ID NO: 110). HIG2-A24-9-19 (SEQ ID NO: 110) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #6 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "c" depicts the CTL-inducing ability of HIG2-A24-9-22 (SEQ ID NO: 111). HIG2-A24-9-22 (SEQ ID NO: 111) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone, that was established from the

positive well #7 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "d" depicts the CTL-inducing ability of HIG2-A24-9-8 (SEQ ID NO: 387). HIG2-A24-9-8 (SEQ ID NO: 387) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone, that were established from the positive well #5 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "e" depicts the CTL-inducing ability of HIG2-A02-9-8 (SEQ ID NO: 114). HIG2-A02-9-8 (SEQ ID NO: 114) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line was established from the positive well #10 shown in boxed wells. The established CTL line raised against the peptide demonstrate specific CTL activity against 293T transfected both full length of HIG2 gene and HLA-A02 molecule. 293T transfected full length of HIG2 but not HLA-A02, 293Ts transfected HLA-A02 and FoxP3 gene as substitute dor full length of HIG2 and 293Ts transfected HLA-A02 and pulsed with HIG2-9-15 were prepared for the negative control. The CTL line showed high specific CTL activity against 293T that transfected both HIG2 and HLA-A02.

[fig.4-2]Figure 4 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that HIG2-A24-9-19 (SEQ ID NO: 110), HIG2-A24-9-22 (SEQ ID NO: 111), HIG2-A24-9-8 (SEQ ID NO: 387), HIG2-A24-10-7 (SEQ ID NO: 112), HIG2-A24-10-18 (SEQ ID NO: 394), HIG2-A02-9-15 (SEQ ID NO: 116), HIG2-A02-9-4 (SEQ ID NO: 117) and HIG2-A02-10-8 (SEQ ID NO: 121) show potent IFN-gamma production. "f" depicts the CTL-inducing ability of HIG2-A24-10-7 (SEQ ID NO: 112). HIG2-A24-10-7 (SEQ ID NO: 112) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL lines or clone, that were established from the positive well #1 and #7 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "g" depicts the CTL-inducing ability of HIG2-A24-10-18 (SEQ ID NO: 394). HIG2-A24-10-18 (SEQ ID NO: 394) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone, that were established from the positive well #7 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "h" depicts the CTL-inducing ability of HIG2-A02-9-15 (SEQ ID NO: 116). HIG2-A02-9-15 (SEQ ID NO: 116) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line was established from the positive well #10 shown in boxed wells. The established CTL line raised against the peptide demonstrated specific CTL activity against COS7 transfected both full length of HIG2 gene and HLA-A02 molecule. COS7 transfected full length of HIG2 but not HLA-A02 and COS7s transfected HLA-

A02 and pulsed with HIG2-9-8 peptide were prepared for the negative control. The CTL line showed high specific CTL activity against COS7 that transfected both HIG2 and HLA-A02.

[fig.4-3]Figure 4 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that HIG2-A24-9-19 (SEQ ID NO: 110), HIG2-A24-9-22 (SEQ ID NO: 111), HIG2-A24-9-8 (SEQ ID NO: 387), HIG2-A24-10-7 (SEQ ID NO: 112), HIG2-A24-10-18 (SEQ ID NO: 394), HIG2-A02-9-15 (SEQ ID NO: 116), HIG2-A02-9-4 (SEQ ID NO: 117) and HIG2-A02-10-8 (SEQ ID NO: 121) show potent IFN-gamma production. "i" depicts the CTL-inducing ability of HIG2-A02-9-4 (SEQ ID NO: 117). HIG2-A02-9-4 (SEQ ID NO: 117) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone were established from the positive well #10 shown in boxed wells. The established CTL line raised against the peptide demonstrated specific CTL activity against COS7 transfected both full length of HIG2 gene and HLA-A02 molecule (middle graph). Also, COS7 transfected full length of HIG2 but not HLA-A02, COS7s transfected HLA-A02 and TTK gene as substitute for full length of HIG2 and COS7s transfected HLA-A02 and pulsed with HIG2-9-8 were prepared for the negative control. Cytotoxic activity of the CTL clone against 293T, transfected both full length of HIG2 gene and HLA-A02 molecule, and cancer cell line, Caki-1 endogenously expressing HIG2 and HLA-A02 was measured by Cr-release assay (CRA) (lower graphs), and the CTL clone had very potent cytotoxic activity against the transfectant with both of HIG2 gene and HLA-A02, and Caki-1. On the other hand, the effector cells did not demonstrate the cytotoxic activity of the CTL line against 293T, transfected only HIG2 or only HLA-A02, and cancer cell line, A498 expressing only HIG2 was not detected. "j" depicts the CTL-inducing ability of HIG2-A02-10-8 (SEQ ID NO: 121). HIG2-A02-10-8 (SEQ ID NO: 121) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #9 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide.

[fig.5-1]Figure 5 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that INHBB-A24-9-180 (SEQ ID NO: 395), INHBB-A24-10-180 (SEQ ID NO: 133), INHBB-A24-10-305 (SEQ ID NO: 135), INHBB-A24-10-7 (SEQ ID NO: 137) and INHBB-A24-10-212 (SEQ ID NO: 426) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of INHBB-A24-9-180 (SEQ ID NO: 395). INHBB-A24-9-180 (SEQ ID NO: 395) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone was established from the

positive well #7 shown in boxed wells. Cytotoxic activity of the established CTL clone against tumor cells, Miapaca2 expressing both of INHBB and HLA-A02 was measured by Cr-release assay (CRA), and the effector cells showed high specific cytotoxic activity against Miapaca2. On the other hand, it did not show significant specific cytotoxic activity against Caki-1 expressing INHBB but not HLA-A02. "c" depicts the CTL-inducing ability of INHBB-A24-10-180 (SEQ ID NO: 133). INHBB-A24-10-180 (SEQ ID NO: 133) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line was established from the positive well #3 shown in boxed wells. The established CTL line raised against the peptide demonstrated high specific CTL activity against 293T transfected both of full length of INHBB gene and HLA-A24 molecule. Also, 293T transfected full length of INHBB but not HLA-A24 and 293Ts transfected HLA-A24 and pulsed with INHBB-10-305 peptide were prepared for the negative control.

[fig.5-2]Figure 5 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that INHBB-A24-9-180 (SEQ ID NO: 395), INHBB-A24-10-180 (SEQ ID NO: 133), INHBB-A24-10-305 (SEQ ID NO: 135), INHBB-A24-10-7 (SEQ ID NO: 137) and INHBB-A24-10-212 (SEQ ID NO: 426) show potent IFN-gamma production. "d" depicts the CTL-inducing ability of INHBB-A24-10-305 (SEQ ID NO: 135). INHBB-A24-10-305 (SEQ ID NO: 135) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone were established from the positive well #2 shown in boxed wells. The established CTL clone raised against the peptide demonstrated high specific CTL activity against 293T transfected both full length of INHBB gene and HLA-A24 molecule. Also, 293T transfected full length of INHBB but HLA-A24 and 293Ts transfected HLA-A24 and pulsed with INHBB-10-180 peptide were prepared for the negative control. "e" depicts the CTL-inducing ability of INHBB-A24-10-7 (SEQ ID NO: 137). INHBB-A24-10-7 (SEQ ID NO: 137) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL lines were established from the positive well #8 shown in boxed wells in upper panel and #2 shown in boxed wells in lower panel. The CTL line from #8 well demonstrated specific CTL activity against 293T transfected both full length of INHBB gene and HLA-A24 molecule. Also, 293T transfected full length of INHBB but not HLA-A24 and 293Ts transfected HLA-A24 and pulsed with INHBB-10-40 peptide were prepared for the negative control. "f" depicts the CTL-inducing ability of INHBB-A24-10-212 (SEQ ID NO: 426). INHBB-A24-10-212 (SEQ ID NO: 426) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #1 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide.

[fig.6-1]Figure 6 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that KIF20A-A24-10-304 (SEQ ID NO: 186), KIF20A-A24-9-383 (SEQ ID NO: 178), KIF20A-A24-10-66 (SEQ ID NO: 194) and KIF20A-A24-9-305 (SEQ ID NO: 174) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of KIF20A-A24-10-304 (SEQ ID NO: 186). KIF20A-A24-10-304 (SEQ ID NO: 186) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay. The well #5 shown in boxed wells in lower right panel demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL line and clone, that were established from the positive well #5 shown in boxed wells in upper left panel, also demonstrated the specific response against the target cells pulsed with the epitope peptide. The established CTL clone raised against the peptide demonstrated specific CTL activity against 24-LCL transfected full length of KIF20A gene. Also, A24-LCL transfected mock vector was prepared for the negative control. Cytotoxic activity of the CTL clone against tumor cells, Miapaca2 expressing both of KIF20A and HLA-A24 was measured by Cr-release assay (CRA), and the CTL clone had very potent specific cytotoxic activity against Miapaca2 (lower right graph). On the other hand, it did not show significant specific cytotoxic activity against PK59 expressing KIF20A but not HLA-A24. "c" depicts the CTL-inducing ability of KIF20A-A24-9-383 (SEQ ID NO: 178). KIF20A-A24-9-383 (SEQ ID NO: 178) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay. The well #3 and 4 shown in boxed wells in right panel demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL line, that was established from the positive well #3 shown in boxed wells in left panel, also demonstrated the specific response against the target cells pulsed with the epitope peptide. The established CTL line demonstrated high specific CTL activity against COS7 transfected both full length of KIF20A gene and HLA-A24 molecule. Also, COS7 transfected full length of KIF20A but not HLA-A24 and COS7s transfected HLA-A24 and pulsed with KIF20A-9-621 peptide were prepared for the negative control.

[fig.6-2]Figure 6 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that KIF20A-A24-10-304 (SEQ ID NO: 186), KIF20A-A24-9-383 (SEQ ID NO: 178), KIF20A-A24-10-66 (SEQ ID NO: 194) and KIF20A-A24-9-305 (SEQ ID NO: 174) show potent IFN-gamma production. "d" depicts the CTL-inducing ability of KIF20A-A24-10-66 (SEQ ID NO: 194). KIF20A-A24-10-66 (SEQ ID NO: 194) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL lines, that were established from the positive well

#6 shown in boxed wells in upper left panel and #3 shown in boxed wells in lower middle panel demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL clone selected from CTL line from #6 well by limiting dilution demonstrated specific CTL activity against the target cells. The established CTL clone showed specific CTL activity against COS7 transfected both full length of KIF20A gene and HLA-A24 molecule. Also, COS7 transfected full length of KIF20A but not HLA-A24, COS7s transfected HLA-A24 and URLC10 gene as substitute for full length of KIF20A and COS7 transfected HLA-A24 and pulsed with KIF20A-10-308 peptide were prepared for the negative control. "e" depicts the CTL-inducing ability of KIF20A-A24-9-305 (SEQ ID NO: 174). KIF20A-A24-9-305 (SEQ ID NO: 174) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL lines, that were established from the positive well #2 shown in boxed wells in upper left panel and #6 shown in boxed wells in lower middle panel, demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL clone selected from CTL line from #2 well by limiting dilution demonstrated specific CTL activity against the target cells. Cytotoxic activity of the CTL clone against tumor cells, PK45P expressing both of KIF20A and HLA-A24 was measured by Cr-release assay (CRA), and the CTL clone had very potent cytotoxic activity against PK45P. On the other hand, it did not show significant specific cytotoxic activity against PK59 expressing KIF20A but not HLA-A24.

[Fig. 7-1]

Figure 7 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that KNTC2-A24-9-309 (SEQ ID NO: 196), KNTC2-A24-9-124 (SEQ ID NO: 202), KNTC2-A24-9-154 (SEQ ID NO: 210) KNTC2-A24-9-150 (SEQ ID NO: 213), KNTC2-A24-10-452 (SEQ ID NO: 214), KNTC2-A24-10-227 (SEQ ID NO: 217) and KNTC2-A24-10-273 (SEQ ID NO: 223) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of KNTC2-A24-9-309 (SEQ ID NO: 196). KNTC2-A24-9-309 (SEQ ID NO: 196) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #8 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide.. "c" depicts the CTL-inducing ability of KNTC2-A24-9-124 (SEQ ID NO: 202). KNTC2-A24-9-124 (SEQ ID NO: 202) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #5 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "d" depicts the CTL-inducing ability of

KNTC2-A24-9-154 (SEQ ID NO: 210). KNTC2-A24-9-154 (SEQ ID NO: 210) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clone, that were established from the positive well #5 shown in boxed wells demonstrated the specific response against the target cells pulsed with the epitope peptide. "e" depicts the CTL-inducing ability of KNTC2-A24-9-150 (SEQ ID NO: 213). KNTC2-A24-9-150 (SEQ ID NO: 213) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #7 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide.

[Fig. 7-2]

Figure 7 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that KNTC2-A24-9-309 (SEQ ID NO: 196), KNTC2-A24-9-124 (SEQ ID NO: 202), KNTC2-A24-9-154 (SEQ ID NO: 210) KNTC2-A24-9-150 (SEQ ID NO: 213), KNTC2-A24-10-452 (SEQ ID NO: 214), KNTC2-A24-10-227 (SEQ ID NO: 217) and KNTC2-A24-10-273 (SEQ ID NO: 223) show potent IFN-gamma production. "f" depicts the CTL-inducing ability of KNTC2-A24-10-452 (SEQ ID NO: 214). KNTC2-A24-10-452 (SEQ ID NO: 214) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL lines and clone, that were established from the positive well #4 shown in boxed wells in upper left panel and #5 shown in boxed wells in middle panel, demonstrated the specific response against the target cells pulsed with the epitope peptide. Moreover, CTL clone selected from CTL line from #5 well by limiting dilution demonstrated specific CTL activity against the target cells. The established CTL line from #4 well showed specific CTL activity against HEK293 transfected both full length of KNTC2 gene and HLA-A24 molecule. Also, HEK293 transfected full length of KNTC2 but not HLA-A24, HEK293 transfected HLA-A24 but full length of KNTC2 and HEK293 transfected HLA-A24 pulsed with KNTC-9-309 peptide were prepared for the negative control. "g" depicts the CTL-inducing ability of KNTC2-A24-10-227 (SEQ ID NO: 217). KNTC2-A24-10-227 (SEQ ID NO: 217) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #1 shown in boxed well s, demonstrated the specific response against the target cells pulsed with the epitope peptide. "h" depicts the CTL-inducing ability of KNTC2-A24-10-273 (SEQ ID NO: 223). KNTC2-A24-10-273 (SEQ ID NO: 223) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #8 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide.

[Fig. 8-1]

Figure 8 depicts the results of the screening of epitope peptides, which, in turn,

demonstrate that TTK-A02-9-462 (SEQ ID NO: 227), TTK-A02-9-719 (SEQ ID NO: 233), TTK-A02-9-547 (SEQ ID NO: 228) and TTK-A02-10-462 (SEQ ID NO: 254), show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of TTK-A02-9-462 (SEQ ID NO: 227). TTK-A02-9-462 (SEQ ID NO: 227) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and two clones, that were established from the positive well #4 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. The established CTL clone showed high specific CTL activity against COS7 transfected both full length of TTK gene and HLA-A02 molecule. Also, COS7 transfected full length of TTK but not HLA-A02, COS7s transfected HLA-A02 but not full length of TTK and COS7s transfected HLA-A02 pulsed with TTK-9-547 peptide were prepared for the negative control. "c" depicts the CTL-inducing ability of TTK-A02-9-719 (SEQ ID NO: 233). TTK-A02-9-719 (SEQ ID NO: 233) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clones were established from the positive well #1 shown in boxed wells. The established CTL line showed high specific CTL activity against COS7 transfected both full length of TTK gene and HLA-A02 molecule. Also, COS7 transfected full length of TTK but not HLA-A02 and COS7s transfected HLA-A02 and HIG2 gene as substitute for full length of TTK were prepared for the negative control. [fig.8-2]Figure 8 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that TTK-A02-9-462 (SEQ ID NO: 227), TTK-A02-9-719 (SEQ ID NO: 233), TTK-A02-9-547 (SEQ ID NO: 228) and TTK-A02-10-462 (SEQ ID NO: 254), show potent IFN-gamma production. "d" depicts the CTL-inducing ability of TTK-A02-9-547 (SEQ ID NO: 228). TTK-A02-9-547 (SEQ ID NO: 228) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clones were established from the positive well #2 shown in boxed wells. The established CTL line showed specific CTL activity against COS7 transfected both full length of TTK gene and HLA-A02 molecule. Also, COS7 transfected full length of TTK but not HLA-A02, COS7s transfected HLA-A02 but not full length of TTK and COS7s transfected HLA-A02 and pulsed with TTK-10-462 were prepared for the negative control.

[fig.8-3]Figure 8 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that TTK-A02-9-462 (SEQ ID NO: 227), TTK-A02-9-719 (SEQ ID NO: 233), TTK-A02-9-547 (SEQ ID NO: 228) and TTK-A02-10-462 (SEQ ID NO: 254), show potent IFN-gamma production. "e" depicts the CTL-inducing ability of TTK-A02-10-462 (SEQ ID NO: 254). TTK-A02-10-462 (SEQ ID NO: 254) demonstrated

potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and three clones were established from the positive well #8 shown in boxed wells. The established CTL clone showed specific CTL activity against COS7 transfected both full length of TTK gene and HLA-A02 molecule. Also, COS7 transfected full length of TTK but not HLA-A02, COS7s transfected HLA-A02 but not full length of TTK and COS7s transfected HLA-A02 and pulsed with TTK-9-547 peptide were prepared for the negative control.

[fig.9-1]Figure 9 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that URLC10-A02-9-206 (SEQ ID NO: 271), URLC10-A02-9-212 (SEQ ID NO: 272) and URLC10-A02-10-211 (SEQ ID NO: 288) show potent IFN-gamma production. "a" depicts the example of negative peptides which could not be detected CTL-inducing ability despite possible binding activity with HLA. "b" depicts the CTL-inducing ability of URLC10-A02-9-206 (SEQ ID NO: 271). URLC10-A02-9-206 (SEQ ID NO: 271) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #7 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "c" depicts the CTL-inducing ability of URLC10-A02-9-212 (SEQ ID NO: 272). URLC10-A02-9-212 (SEQ ID NO: 272) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line, that was established from the positive well #3 shown in boxed wells, demonstrated the specific response against the target cells pulsed with the epitope peptide. "d" depicts the CTL-inducing ability of URLC10-A02-10-211 (SEQ ID NO: 288). URLC10-A02-10-211 (SEQ ID NO: 288) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and CTL line and clones, that were established from the positive well #5 shown in boxed wells.

[fig.9-2]Figure 9 depicts the results of the screening of epitope peptides, which, in turn, demonstrate that URLC10-A02-9-206 (SEQ ID NO: 271), URLC10-A02-9-212 (SEQ ID NO: 272) and URLC10-A02-10-211 (SEQ ID NO: 288) show potent IFN-gamma production. "Continuation of d" The established CTL clone showed high specific CTL activity against COS7, Hek293 and 293T which were transfected both full length of URLC10 gene and HLA-A02 molecule. Also, COS7, Hek293 or 293T which were transfected full length of URLC10 but not HLA-A02 and COS7s, Hek293s or 293Ts, which were transfected HLA-A02 and pulsed with URLC10-10-64, were prepared for the negative control. In this drawings, "+" means the peptide pulsed target, "-" means the no peptide pulsed target, "R" means Responder, "S" means Stimulator, "E" means Effector, and "T" means Target.

The words "a", "an", and "the" as used herein mean "at least one" unless otherwise specifically indicated.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0017] The present invention is based in part on the discovery of applicable targets of immunotherapy. Identification of new TAAs, particularly those that induce potent and specific anti-tumor immune responses, warrants further development of the clinical application of the peptide vaccination strategy in various types of cancer (Boon T et al., (1996) *J Exp Med* 183: 725-9.; van der Bruggen P et al., (1991) *Science* 254: 1643-7.; Brichard V et al., (1993) *J Exp Med* 178: 489-95.; Kawakami Y et al., (1994) *J Exp Med* 180: 347-52.; Shichijo S et al., (1998) *J Exp Med* 187:277-88.; Chen YT et al., (1997) *Proc.Natl.Acd. Sci.USA*, 94: 1914-8.; Harris CC, (1996) *J Natl Cancer Inst* 88:1442-55.; Butterfield LH et al., (1999) *Cancer Res* 59:3134-42.; Vissers JL et al., (1999) *Cancer Res* 59: 5554-9.; van der Burg SH et al., (1996) *J. Immunol* 156:3308-14.; Tanaka F et al., (1997) *Cancer Res* 57:4465-8.; Fujie T et al., (1999) *Int J Cancer* 80:169-72.; Kikuchi M et al., (1999) *Int J Cancer* 81 : 459-66.; Oiso M et al., (1999) *Int J Cancer* 81:387-94.). Because TAAs have often no immunogenicity, discovery of fitting targets is extremely important issue.

[0018] As noted above,
 CDH3 (GenBank Accession No. NM_001793; SEQ ID Nos.1, 2),
 EPHA4 (GenBank Accession No. L36645; SEQ ID Nos.3, 4),
 ECT2 (GenBank Accession No. AY376439; SEQ ID Nos.5, 6),
 HIG2 (GenBank Accession No. NM_013332; SEQ ID Nos.7, 8)
 INHBB (GenBank Accession No. NM_002193; SEQ ID Nos.9, 10),
 KIF20A (GenBank Accession No. NM_005733; SEQ ID Nos.11, 12),
 KNTC2 (GenBank Accession No. AF017790; SEQ ID Nos.13, 14),
 TTK (GenBank Accession No. NM_003318; SEQ ID Nos.15, 16) and
 URLC10 (GenBank Accession No. NM_017527; SEQ ID Nos.17, 18) were previously identified as over-expressed in various cancers using cDNA microarray technologies.

[0019] In the present invention, peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10 are shown to be TAA epitopes restricted by HLA-A24 and HLA-A2, an HLA allele commonly found in the Japanese and Caucasian populations. Specifically, using their binding affinities to HLA-A24 or HLA-A2, candidates of HLA-A24 or HLA-A2 binding peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10 were identified. After the in vitro stimulation of T-cells by dendritic cells (DCs) loaded with these

peptides, CTLs were successfully established using the following peptides.

CDH3-A24-9-513 (SEQ ID NO: 19),
CDH3-A24-9-406 (SEQ ID NO: 22),
CDH3-A24-10-807 (SEQ ID NO: 30),
CDH3-A24-10-332 (SEQ ID NO: 34),
CDH3-A24-10-655 (SEQ ID NO: 344),
CDH3-A24-10-470 (SEQ ID NO: 358),
EphA4-A24-9-453 (SEQ ID NO: 41),
EphA4-A24-9-5 (SEQ ID NO: 44),
EphA4-A24-9-869 (SEQ ID NO: 46),
EphA4-A24-9-420 (SEQ ID NO: 48),
EphA4-A24-10-24 (SEQ ID NO: 78),
EphA4-A02-9-501 (SEQ ID NO: 376),
EphA4-A02-9-165 (SEQ ID NO: 379),
ECT2-A24-9-515 (SEQ ID NO: 80),
ECT2-A24-10-40 (SEQ ID NO: 100),
ECT2-A24-10-101 (SEQ ID NO: 101),
HIG2-A24-9-19 (SEQ ID NO: 110),
HIG2-A24-9-22 (SEQ ID NO: 111),
HIG2-A24-9-8 (SEQ ID NO: 387),
HIG2-A24-10-7 (SEQ ID NO: 112),
HIG2-A24-10-18 (SEQ ID NO: 394),
HIG2-A02-9-8 (SEQ ID NO: 114),
HIG2-A02-9-15 (SEQ ID NO: 116),
HIG2-A02-9-4 (SEQ ID NO: 117),
HIG2-A02-10-8 (SEQ ID NO: 121),
INHBB-A24-9-180 (SEQ ID NO: 395),
INHBB-A24-10-180 (SEQ ID NO: 133),
INHBB-A24-10-305 (SEQ ID NO: 135),
INHBB-A24-10-7 (SEQ ID NO: 137),
INHBB-A24-10-212 (SEQ ID NO: 426),
KIF20A-A24-9-305 (SEQ ID NO: 174),
KIF20A-A24-9-383 (SEQ ID NO: 178),
KIF20A-A24-10-304 (SEQ ID NO: 186),
KIF20A-A24-10-66 (SEQ ID NO: 194),
KNTC2-A24-9-309 (SEQ ID NO: 196),
KNTC2-A24-9-124 (SEQ ID NO: 202),
KNTC2-A24-9-154 (SEQ ID NO: 210),

KNTC2-A24-9-150 (SEQ ID NO: 213),
 KNTC2-A24-10-452 (SEQ ID NO: 214),
 KNTC2-A24-10-227 (SEQ ID NO: 217),
 KNTC2-A24-10-273 (SEQ ID NO: 223),
 TTK-A02-9-462 (SEQ ID NO: 227),
 TTK-A02-9-547 (SEQ ID NO: 228),
 TTK-A02-9-719 (SEQ ID NO: 233),
 TTK-A02-10-462 (SEQ ID NO: 254),
 URLC-A02-9-206 (SEQ ID NO: 271),
 URLC-A02-9-212 (SEQ ID NO: 272) and
 URLC-A02-10-211 (SEQ ID NO: 288)

[0020]

These peptides are epitope peptides of each TAA restricted by HLA-A24 or HLA-A2. Since these antigens are over-expressed in most cancers and are associated with tumor cell proliferation, they find utility as immunotherapeutic targets against cancers. Exemplary cancers include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0021]

Accordingly, the present invention further provides methods of treating or preventing a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers in a subject, such methods including the steps of administering to the subject an immunogenic peptide of less than about 40 amino acids, often less than about 20 amino acids, usually less than about 15 amino acids and having the amino acid sequence of SEQ ID NOs: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288.

[0022]

Alternatively, the immunogenic peptide may have an amino acid sequence as set forth in SEQ ID NOs: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 in which 1, 2, or several (e.g., up to 5) amino acids are substituted, deleted or added, provided the resulting variant peptide retains the immunogenic activity (i.e., the ability to induce CTLs specific to cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers).

[0023]

The number of residues to be substituted, deleted, or added is generally 5 amino acids or less, preferably 4 amino acids or less, more preferably 3 amino acids or less,

even more preferably one or two amino acids. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor. Furthermore the present invention provides methods for preventing post-surgery recurrence of these diseases mentioned above.

[0024]

Variant peptides (i.e., peptides having an amino acid sequence modified by substituting, deleting, or adding one, two or several amino acid residues to an original amino acid sequence) are known to retain the original biological activity (Mark DF et al., (1984) Proc Natl Acad Sci USA 81: 5662-6.; Zoller MJ and Smith M, (1982) Nucleic Acids Res 10:6487-500.; Dalbadie-McFarland G et al., (1982) Proc Natl Acad Sci USA 79: 6409-13.). In the context of the present invention, it is preferable that the amino acid modification results in conservation of the properties of the original amino acid side-chain (a process known as conservative amino acid substitution). Examples of properties of amino acid side chains include hydrophobic amino acids (A, I, L, M, F, P, W, Y, V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S, T), and side chains having the following functional groups or characteristics in common: an aliphatic side-chain (G, A, V, L, I, P); a hydroxyl group containing side-chain (S, T, Y); a sulfur atom containing side-chain (C, M); a carboxylic acid and amide containing side-chain (D, N, E, Q); a base containing side-chain (R, K, H); and an aromatic containing side-chain (H, F, Y, W). Note, the parenthetic letters indicate the one-letter codes of amino acids.

[0025]

In preferred embodiments, the immunogenic peptide is a nonapeptide (9-mer) or a decapeptide (10-mer).

The present invention further provides a method of inducing anti-tumor immunity for a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, in a subject, such a method including the steps of administering to the subject an immunogenic peptide of the present invention, namely one having the amino acid sequence of SEQ ID NOS: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288, or a variant thereof (i.e., including 1, 2, or several (e.g., up to 5) amino acid substitutions, deletions, or additions) to the subject in need thereof. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer,

pancreatic cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0026]

In the context of the present invention, the subject is preferably a mammal. Exemplary mammals include, but are not limited to, e.g., a human, non-human primate, mouse, rat, dog, cat, horse, or cow.

In the present invention, the peptide can be administered to a subject via an in vivo or ex vivo protocol. Furthermore, the present invention also provides use of nonapeptide or decapeptide selected from peptides having the amino acid sequence of SEQ ID NOS: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 (and variants thereof) for manufacturing an immunogenic composition for treating or preventing a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0027]

Homology analyses of the following peptides demonstrate that they do not have significant homology with the peptides derived from any known human gene products.

CDH3-A24-9-513 (SEQ ID NO: 19),
CDH3-A24-9-406 (SEQ ID NO: 22),
CDH3-A24-10-807 (SEQ ID NO: 30),
CDH3-A24-10-332 (SEQ ID NO: 34),
CDH3-A24-10-655 (SEQ ID NO: 344),
CDH3-A24-10-470 (SEQ ID NO: 358),
EphA4-A24-9-453 (SEQ ID NO: 41),
EphA4-A24-9-5 (SEQ ID NO: 44),
EphA4-A24-9-869 (SEQ ID NO: 46),
EphA4-A24-9-420 (SEQ ID NO: 48),
EphA4-A24-10-24 (SEQ ID NO: 78),
EphA4-A02-9-501 (SEQ ID NO: 376),
EphA4-A02-9-165 (SEQ ID NO: 379),
ECT2-A24-9-515 (SEQ ID NO: 80),
ECT2-A24-10-40 (SEQ ID NO: 100),
ECT2-A24-10-101 (SEQ ID NO: 101),
HIG2-A24-9-19 (SEQ ID NO: 110),

HIG2-A24-9-22 (SEQ ID NO: 111),
HIG2-A24-9-8 (SEQ ID NO: 387),
HIG2-A24-10-7 (SEQ ID NO: 112),
HIG2-A24-10-18 (SEQ ID NO: 394),
HIG2-A02-9-8 (SEQ ID NO: 114),
HIG2-A02-9-15 (SEQ ID NO: 116),
HIG2-A02-9-4 (SEQ ID NO: 117),
HIG2-A02-10-8 (SEQ ID NO: 121),
INHBB-A24-9-180 (SEQ ID NO: 395),
INHBB-A24-10-180 (SEQ ID NO: 133),
INHBB-A24-10-305 (SEQ ID NO: 135),
INHBB-A24-10-7 (SEQ ID NO: 137),
INHBB-A24-10-212 (SEQ ID NO: 426),
KIF20A-A24-9-305 (SEQ ID NO: 174),
KIF20A-A24-9-383 (SEQ ID NO: 178),
KIF20A-A24-10-304 (SEQ ID NO: 186),
KIF20A-A24-10-66 (SEQ ID NO: 194),
KNTC2-A24-9-309 (SEQ ID NO: 196),
KNTC2-A24-9-124 (SEQ ID NO: 202),
KNTC2-A24-9-154 (SEQ ID NO: 210),
KNTC2-A24-9-150 (SEQ ID NO: 213),
KNTC2-A24-10-452 (SEQ ID NO: 214),
KNTC2-A24-10-227 (SEQ ID NO: 217),
KNTC2-A24-10-273 (SEQ ID NO: 223),
TTK-A02-9-462 (SEQ ID NO: 227),
TTK-A02-9-547 (SEQ ID NO: 228),
TTK-A02-9-719 (SEQ ID NO: 233),
TTK-A02-10-462 (SEQ ID NO: 254),
URLC-A02-9-206 (SEQ ID NO: 271),
URLC-A02-9-212 (SEQ ID NO: 272) and
URLC-A02-10-211 (SEQ ID NO: 288)

[0028] Accordingly, the possibility of unknown or undesirable immune responses with immunotherapy against these molecules is significantly reduced.

[0029] Regarding HLA antigens, the data presented here demonstrate that the uses of A-24 type or A-2 type antigens (which are said to be highly expressed among the Japanese) are favorable for obtaining effective results. The uses of subtypes such as A-2402 and A-0201 are even more preferable. Typically, in the clinic, the type of HLA antigen of the patient requiring treatment is investigated in advance, which, in turn, enables the

selection of appropriate peptides having high levels of binding affinity to the patient antigen, or having cytotoxic T cell (CTL) inducibility by antigen presentation. Furthermore, in order to obtain peptides having high binding affinity and CTL inducibility, substitution, deletion, or addition of 1, 2, or several (e.g., up to 5) amino acids may be performed based on the amino acid sequence of the naturally occurring CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 partial peptide. Herein, the term "several" means refers to 5 or less, more preferably 3 or less. Furthermore, in addition to peptides that are naturally displayed, since the regularity of the sequences of peptides displayed by binding to HLA antigens is already known (Kubo RT, et al., (1994) *J. Immunol.*, 152, 3913-24.; Rammensee HG, et al., (1995) *Immunogenetics*, 41:178-228.; Kondo A, et al., (1995) *J. Immunol.* 155:4307-12.), modifications based on such regularity can be performed on the immunogenic peptides of the invention. For example, peptides possessing high HLA-24 binding affinity in which the second amino acid from the N terminus substituted with phenylalanine, tyrosine, methionine, or tryptophan may be favorably used. Likewise, peptides whose C-terminal amino acid is substituted with phenylalanine, leucine, isoleucine, tryptophan, or methionine may also be used favorably. On the other hand, peptides possessing high HLA-A2 binding affinity in which the second amino acid from the N terminus substituted with leucine or methionine, and peptides whose C-terminal amino acid is substituted with valine or leucine may be used favorably. The substitution is performed not only at the terminus amino acids but also at the position of potential TCR recognition of peptides. Several studies have demonstrated that amino acid substitutions in a peptide can be equal to or better than the original, for example CAP1, p53₍₂₆₄₋₂₇₂₎, Her-2/neu₍₃₆₉₋₃₇₇₎ or gp100₍₂₀₉₋₂₁₇₎ (Zaremba et al. *Cancer Res.* 57, 4570-4577, 1997, T. K. Hoffmann et al. *J Immunol.* (2002) Feb 1;168(3):1338-47, S. O. Dionne et al. *Cancer Immunol Immunother.* (2003) 52: 199-206 and S. O. Dionne et al. *Cancer Immunology, Immunotherapy* (2004) 53, 307-314). Furthermore, 1 to 2 amino acids may be added to the N terminus and/or C terminus of the peptide.

[0030] However, when the peptide sequence is identical to a portion of the amino acid sequence of an endogenous or exogenous protein having a different function, side effects such as autoimmune disorders or allergic symptoms against specific substances may be induced. Therefore, it is preferable to avoid the situation wherein the immunogenic sequence matches the amino acid sequence of a known protein. This situation may be avoided by performing a homology search using available databases. If homology searches confirm that peptides in which 1, 2 or several different amino acids do not exist in nature, then the danger that modifications of the above-mentioned amino acid sequence that, for example, increase the binding affinity with HLA antigens, and/or increase the CTL inducibility can be avoided.

[0031] Although peptides having high binding affinity to the HLA antigens as described above are expected to be highly effective as cancer vaccines, the candidate peptides, which are selected according to the presence of high binding affinity as an indicator, must be examined for the actual presence of CTL inducibility. CTL inducibility may be routinely confirmed by inducing antigen-presenting cells carrying human MHC antigens (for example, B-lymphocytes, macrophages, and dendritic cells), or more specifically dendritic cells derived from human peripheral blood mononuclear leukocytes, and, after stimulation with the peptide of interest, mixing with CD8-positive cells and measuring the cytotoxic activity against the target cells. As the reaction system, transgenic animals produced to express a human HLA antigen (for example, those described in BenMohamed L, et al., (2000) *Hum. Immunol.*; 61(8):764-79 Related Articles, Books, Linkout.) may be used. For example, the target cells can be radio-labeled with ^{51}Cr and such, and cytotoxic activity can be calculated from radioactivity released from the target cells. Alternatively, it can be examined by measuring IFN-gamma produced and released by CTL in the presence of antigen-presenting cells that carry immobilized peptides, and visualizing the inhibition zone on the media using anti-IFN-gamma monoclonal antibodies.

[0032] As a result of examining the CTL inducibility of peptides as described above, it was discovered that those peptides having high binding affinity to an HLA antigen did not necessarily have high inducibility. However, nonapeptides or decapeptides selected from the group of peptides having the amino acid sequences indicated by the following peptides showed particularly high CTL inducibility.

CDH3-A24-9-513 (SEQ ID NO: 19),
CDH3-A24-9-406 (SEQ ID NO: 22),
CDH3-A24-10-807 (SEQ ID NO: 30),
CDH3-A24-10-332 (SEQ ID NO: 34),
CDH3-A24-10-655 (SEQ ID NO: 344),
CDH3-A24-10-470 (SEQ ID NO: 358),
EphA4-A24-9-453 (SEQ ID NO: 41),
EphA4-A24-9-5 (SEQ ID NO: 44),
EphA4-A24-9-869 (SEQ ID NO: 46),
EphA4-A24-9-420 (SEQ ID NO: 48),
EphA4-A24-10-24 (SEQ ID NO: 78),
EphA4-A02-9-501 (SEQ ID NO: 376),
EphA4-A02-9-165 (SEQ ID NO: 379),
ECT2-A24-9-515 (SEQ ID NO: 80),
ECT2-A24-10-40 (SEQ ID NO: 100),
ECT2-A24-10-101 (SEQ ID NO: 101),

HIG2-A24-9-19 (SEQ ID NO: 110),
HIG2-A24-9-22 (SEQ ID NO: 111),
HIG2-A24-9-8 (SEQ ID NO: 387),
HIG2-A24-10-7 (SEQ ID NO: 112),
HIG2-A24-10-18 (SEQ ID NO: 394),
HIG2-A02-9-8 (SEQ ID NO: 114),
HIG2-A02-9-15 (SEQ ID NO: 116),
HIG2-A02-9-4 (SEQ ID NO: 117),
HIG2-A02-10-8 (SEQ ID NO: 121),
INHBB-A24-9-180 (SEQ ID NO: 395),
INHBB-A24-10-180 (SEQ ID NO: 133),
INHBB-A24-10-305 (SEQ ID NO: 135),
INHBB-A24-10-7 (SEQ ID NO: 137),
INHBB-A24-10-212 (SEQ ID NO: 426),
KIF20A-A24-9-305 (SEQ ID NO: 174),
KIF20A-A24-9-383 (SEQ ID NO: 178),
KIF20A-A24-10-304 (SEQ ID NO: 186),
KIF20A-A24-10-66 (SEQ ID NO: 194),
KNTC2-A24-9-309 (SEQ ID NO: 196),
KNTC2-A24-9-124 (SEQ ID NO: 202),
KNTC2-A24-9-154 (SEQ ID NO: 210),
KNTC2-A24-9-150 (SEQ ID NO: 213),
KNTC2-A24-10-452 (SEQ ID NO: 214),
KNTC2-A24-10-227 (SEQ ID NO: 217),
KNTC2-A24-10-273 (SEQ ID NO: 223),
TTK-A02-9-462 (SEQ ID NO: 227),
TTK-A02-9-547 (SEQ ID NO: 228),
TTK-A02-9-719 (SEQ ID NO: 233),
TTK-A02-10-462 (SEQ ID NO: 254),
URLC-A02-9-206 (SEQ ID NO: 271),
URLC-A02-9-212 (SEQ ID NO: 272) and
URLC-A02-10-211 (SEQ ID NO: 288)

[0033] As noted above, the present invention provides peptides having cytotoxic T cell inducibility, namely those having the amino acid sequence of SEQ ID NOs: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 or a variant thereof (i.e., those in which 1, 2, or several amino acids are substituted, deleted, or added).

[0034] It is preferable that the amino acid sequences composed of 9 or 10 amino acids indicated in SEQ ID NOs: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 or a variant thereof do not match an amino acid sequence associated with another endogenous protein.

[0035] In particular, amino acid substitution to leucine or methionine at the second amino acid from the N terminus, amino acid substitution to valine or leucine at the C-terminal amino acid, and amino acid addition of 1 to 2 amino acids at the N terminus and/or C terminus are examples of preferred variants.

[0036] One of skill in the art will recognize that in addition to amino acid substitutions and additions, immunologically active fragments of the peptides may also be used in the methods of the invention. Methods for determining active fragments are well known in the art. CTL clones obtained by stimulation by these modified peptides can recognize the original peptides and cause damage for cells expressing the original peptides.

[0037] Peptides of the present invention can be prepared using well known techniques. For example, the peptides can be prepared synthetically, using either recombinant DNA technology or chemical synthesis. Peptides of the present invention may be synthesized individually or as longer polypeptides composed of two or more peptides. The peptides of the present invention are preferably isolated, i.e., substantially free of other naturally occurring host cell proteins and fragments thereof.

[0038] The peptides of the present invention may contain modifications, such as glycosylation, side chain oxidation, or phosphorylation; so long as the modifications do not destroy the biological activity of the peptides as described herein, namely the ability to bind to an HLA antigen and induce CTL. Other modifications include incorporation of D-amino acids or other amino acid mimetics that can be used, for example, to increase the serum half life of the peptides.

[0039] Moreover, this invention may contain a method of screening for a peptide which 1, 2, or several amino acids are substituted, wherein said peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 80, 100, 101, 110, 111, 387, 112, 394, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217 or 223, said method comprising the steps of:

- (a) conforming no significant sequence homology to the entire sequence of 1, 2 or several amino acids substitute;
- (b) measuring the CTL inducibility of the candidate substitute peptide; and
- (c) selecting the peptide which CTL inducibility is same to or higher than the original peptide.

[0040]

For example, in preferred embodiments, the present invention provides a method of identifying for a peptide having an ability to induce CTL against cells expressing at least one tumor-associated antigen, wherein the tumor-associated antigen is antigen selected from the group consisting of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10, said method comprising the steps of:

- (i) providing or generating at least one candidate sequence which consists of an amino acid sequence modified by substituting, deleting, or adding one, two or several amino acid residues to an original amino acid sequence, wherein the original amino acid sequence is selected from the group consisting of SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 80, 100, 101, 110, 111, 387, 112, 394, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217 or 223;
- (ii) selecting the candidate sequence that does not have substantial significant homology with the peptides derived from any known human gene products other than said tumor-associated antigens;
- (iii) contacting a peptide consisting of the candidate sequence selected in step (ii) with antigen presenting cells;
- (iv) contacting the antigen presenting cells of step (iii) with T-cells to evaluate the ability of the peptide to stimulate the T-cells; and
- (v) identifying the peptide of which CTL inducibility is same to or higher than a peptide consisting of the original amino acid sequence.

[0041]

Preferably, the amino acid is substituted for a different amino acid in which the properties of the amino acid side-chain are conserved (a process known as conservative amino acid substitution). Examples of properties of amino acid side chains are hydrophobic amino acids (A, I, L, M, F, P, W, Y, V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S, T), and side chains having the following functional groups or characteristics in common: an aliphatic side-chain (G, A, V, L, I, P); a hydroxyl group containing side-chain (S, T, Y); a sulfur atom containing side-chain (C, M); a carboxylic acid and amide containing side-chain (D, N, E, Q); a base containing side-chain (R, K, H); and an aromatic containing side-chain (H, F, Y, W). Note, the parenthetic letters indicate the one-letter codes of amino acids. In the present invention, substantial significant homology is, for example, more than 90%, preferably 95%, more preferably 99% or 100% identity with a known human gene product to be compared.

[0042]

The peptides of this invention can be prepared as a combination, which includes two or more of peptides of the invention, for use as a vaccine for a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, such a vaccine inducing CTL in vivo. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer,

cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor. The peptides may be in a cocktail or may be conjugated to each other using standard techniques. For example, the peptides can be expressed as a single polypeptide sequence. The peptides in the combination may be the same or different.

[0043]

By administering the peptides of this invention, the peptides are presented at a high density on the HLA antigens of antigen-presenting cells, which, in turn, induces CTLs that specifically react toward the complex formed between the displayed peptide and the HLA antigen. Alternatively, antigen-presenting cells having immobilized the peptides of this invention on their cell surface, obtained by removing dendritic cells from the subjects, may be stimulated by the peptides of this invention. Re-administration of these cells to the respective subjects induces CTL, and, as a result, aggressiveness towards the target cells can be increased.

[0044]

More specifically, the present invention provides drugs for treating and/or preventing proliferation, metastasis, and such of a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, which include one or more of peptides of the present invention, or a polynucleotide encoding the peptides. The peptides or polynucleotides of the present invention find particular utility in the treatment of a disease associating CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0045]

The peptides of this invention can be administered to a subject directly, as a pharmaceutical composition that has been formulated by conventional formulation methods. In such cases, in addition to the peptides of this invention, carriers, excipients, and such that are ordinarily used for drugs can be included as appropriate, without particular limitations. The immunogenic compositions of this invention may be used for treatment and prevention of a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer,

renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0046]

The immunogenic compositions for treatment and/or prevention of a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, which include as the active ingredient one or more peptides of the present invention, can further include an adjuvant so that cellular immunity will be established effectively. Alternatively, they may be administered with other active ingredients, such as anti-cancer agents.

[0047]

The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor. Suitable formulations include granules. Suitable adjuvants are described in the literature (Johnson AG. (1994) C1in. Microbiol. Rev., 7:277-89.).

[0048]

Exemplary adjuvants include, but are not limited to, aluminum phosphate, aluminum hydroxide, and alum. Furthermore, liposome formulations, granular formulations in which the drug is bound to few-micrometer diameter beads, and formulations in which a lipid is bound to the peptide may be conveniently used. The method of administration may be oral, intradermal, subcutaneous, intravenous injection, or such, and may include systemic administration or local administration to the vicinity of the targeted tumor.

[0049]

The dose of the peptide(s) of this invention can be adjusted appropriately according to the disease to be treated, age of the patient, weight, method of administration, and such. Though the dosage is ordinarily 0.001 mg to 1000 mg, preferably 0.01 mg to 100 mg, more preferably 0.1 mg to 10 mg, preferably administered once in a few days to few months, one skilled in the art can readily select the appropriate dose and method of administration, as, the selection and optimization of these parameters is well within routine skill.

[0050]

The present invention further provides intracellular vesicles called exosomes, which present complexes formed between the peptides of this invention and HLA antigens on their surface. Exosomes can be prepared, for example, by using the methods described in detail in Published Japanese Translation of International Publication Nos. Hei 11-510507 and 2000-512161, and are preferably prepared using antigen-presenting cells obtained from subjects who are targets of treatment and/or prevention. The exosomes of this invention can be inoculated as cancer vaccines, similarly to the peptides of this invention.

[0051]

The type of HLA antigens used must match that of the subject requiring treatment and/or prevention. For example, in the Japanese population, HLA-A24 or HLA-A2,

particularly HLA-A2402 or HLA-A0201, is often appropriate.

[0052] In some embodiments, the vaccine compositions of the present invention include a component which primes cytotoxic T lymphocytes. Lipids have been identified as agents capable of priming CTL in vivo against viral antigens. For example, palmitic acid residues can be attached to the epsilon-and alpha-amino groups of a lysine residue and then linked to an immunogenic peptide of the invention. The lipidated peptide can then be administered either directly, in a micelle or particle, incorporated into a liposome, or emulsified in an adjuvant. As another example of a lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P3CSS), can be used to prime CTL when covalently attached to an appropriate peptide (see, e.g., Deres K, et al., (1989) *Nature* 342:561-4.).

[0053] The immunogenic compositions of the present invention may also include nucleic acids encoding one or more of the immunogenic peptides disclosed here. See, e.g., Wolff JA et al., (1990) *Science* 247:1465-8; U.S. Patent Nos. 5,580,859; 5,589,466; 5,804,566; 5,739,118; 5,736,524; 5,679,647; and WO 98/04720. Examples of DNA-based delivery technologies include "naked DNA", facilitated (bupivacaine, polymers, peptide-mediated) delivery, cationic lipid complexes, and particle-mediated ("gene gun") or pressure-mediated delivery (see, e.g., U.S. Patent No. 5,922,687).

[0054] The immunogenic peptides of the invention can also be expressed by viral or bacterial vectors. Examples of suitable expression vectors include attenuated viral hosts, such as vaccinia or fowlpox. This approach involves the use of vaccinia virus, e.g., as a vector to express nucleotide sequences that encode the peptide. Upon introduction into a host, the recombinant vaccinia virus expresses the immunogenic peptide, and thereby elicits an immune response. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Patent No. 4,722,848. Another suitable vector is BCG (Bacille Calmette Guerin). BCG vectors are described in Stover CK, et al., (1991) *Nature* 351:456-60. A wide variety of other vectors useful for therapeutic administration or immunization e.g., adeno and adeno-associated virus vectors, retroviral vectors, *Salmonella typhi* vectors, detoxified anthrax toxin vectors, and the like, are known in the art. See, e.g., Shata MT, et al., (2000) *Mol. Med. Today* 6:66-71; Shedlock DJ and Weiner DB., et al., (2000) *J. Leukoc. Biol.* 68:793-806; and Hipp JD, et al., (2000) *In Vivo* 14:571-85.

[0055] The present invention also provides methods of inducing antigen-presenting cells using one or more peptides of this invention. The antigen-presenting cells can be induced by inducing dendritic cells from the peripheral blood monocytes and then contacting (stimulating) them with one or more peptides of this invention *in vitro*, *ex vivo* or *in vivo*. When peptides of the present invention are administered to the subjects, antigen-presenting cells that have the peptides of this invention immobilized

to them are induced in the body of the subject. Alternatively, after immobilizing the peptides of this invention to the antigen-presenting cells, the cells can be administered to the subject as a vaccine. For example, the ex vivo administration may include the steps of:

- a: collecting antigen-presenting cells from a subject, and
- b: contacting the antigen-presenting cells of step a with a peptide of the present invention.

[0056]

Alternatively, according to the present invention, use of the peptides of this invention for manufacturing a pharmaceutical composition inducing antigen-presenting cells is provided. Further, the present invention also provides the peptide of the present invention for inducing antigen-presenting cells. The antigen-presenting cells obtained by step b can be administered to the subject as a vaccine.

[0057]

This invention also provides a method for inducing antigen-presenting cells having a high level of cytotoxic T cell inducibility, in which the method includes the step of transferring genes composed of polynucleotide(s) encoding one or more peptides of this invention to antigen-presenting cells in vitro. The introduced genes may be in the form of DNAs or RNAs. For the method of introduction, without particular limitations, various methods conventionally performed in this field, such as lipofection, electroporation, and calcium phosphate method may be suitably used. More specifically, transfection may be performed as described in Reeves ME, et al., (1996) *Cancer Res.*, 56:5672-7.; Butterfield LH, et al., (1998) *J. Immunol.*, 161:5607-13.; Boczkowski D, et al., (1996) *J. Exp. Med.*, 184:465-72.; Published Japanese Translation of International Publication No. 2000-509281. By transferring the gene into antigen-presenting cells, the gene undergoes transcription, translation, and such in the cell, and then the obtained protein is processed by MHC Class I or Class II, and proceeds through a presentation pathway to present partial peptides.

[0058]

The present invention further provides methods for inducing CTL using one or more peptides of this invention. When the peptides of this invention are administered to a subject, CTL are induced in the body of the subject, and the strength of the immune system targeting the cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancer cells in the tumor tissues is thereby enhanced.

[0059]

The cancers contemplated include, but are not limited to bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor. Alternatively, the peptides of the present invention may be used in the context of an ex vivo therapeutic

method, in which subject-derived antigen-presenting cells and CD8-positive cells or peripheral blood mononuclear leukocytes are contacted (stimulated) with one or more peptides of this invention in vitro, and, after inducing CTL, the cells are returned to the subject. For example, the method may include the steps of:

- a: collecting antigen-presenting cells from a subject,
- b: contacting the antigen-presenting cells of step a with a peptide of the present invention,
- c: mixing the antigen-presenting cells of step b with CD⁸⁺ T cells and co-culturing so as to induce cytotoxic T-cells; and
- d: collecting CD⁸⁺ T cells from the co-culture of step c.

[0060] Alternatively, according to the present invention, use of the peptides of this invention for manufacturing a pharmaceutical composition inducing CTLs is provided. Further, the present invention also provides the peptide of the present invention for inducing CTLs. The CD⁸⁺ T cells having cytotoxic activity obtained by step d can be administered to the subject as a vaccine.

[0061] The present invention further provides isolated cytotoxic T cells induced using the peptides of this invention. The cytotoxic T cells, induced by stimulation with an antigen-presenting cell presenting one or more peptides of this invention, are preferably derived from subjects who are the target of treatment and/or prevention, and can be administered alone or in combination with other drugs, including one or more peptides of this invention or exosomes having anti-tumor activity. The obtained cytotoxic T cells act specifically against target cells presenting the peptides of this invention, or preferably the same peptide(s) used for induction. The target cells may be cells that express CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10 endogenously, or cells that are transfected with CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10 genes. Cells that present the peptides of this invention on the cell surface, due to stimulation with these peptides, can also become targets of attack.

[0062] The present invention also provides antigen-presenting cells presenting complexes formed between HLA antigens and one or more peptides of this invention. The antigen-presenting cells, obtained through contact with the peptides of this invention or the nucleotides encoding such peptides, are preferably derived from subjects who are the target of treatment and/or prevention, and can be administered as vaccines, alone or in combination with other drugs, including the peptides, exosomes, or cytotoxic T cells of the present invention.

[0063] The present invention also provides a composition composed of nucleic acids encoding polypeptides that are capable of forming a subunit of a T cell receptor (TCR), and methods of using the same. The TCR subunits have the ability to form TCRs that

confer specificity to T cells for tumor cells presenting CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10. By using the known method in the art, the nucleic acids of alpha- and beta-chain as the TCR subunits of the CTL induced with one or more peptides of this invention may be identified (WO2007/032255 and Morgan et al., J Immunol, 171, 3288 (2003)). The derivative TCRs preferably bind target cells displaying the CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10 peptide with high avidity, and optionally mediate efficient killing of target cells presenting the CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10 peptide in vivo and in vitro.

- [0064] The nucleic acids encoding the TCR subunits can be incorporated into suitable vectors e.g. retroviral vectors. These vectors are well known in the art. The nucleic acids or the vectors containing them usefully can be transferred into a T cell, which T cell is preferably from a patient. Advantageously, the invention provides an off-the-shelf composition allowing rapid modification of a patient's own T cells (or those of another mammal) to rapidly and easily produce modified T cells having excellent cancer cell killing properties.
- [0065] Also, the present invention provides CTLs which are prepared by transduction with the nucleic acids encoding the TCR subunits polypeptides binding with CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10 peptide e.g. SEQ ID NOs: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 in the context of HLA-A24 or HLA-A2. The transduced CTLs are capable of homing to cancer cells in vivo, and expanded by well known culturing method in vitro (e.g., Kawakami et al., J Immunol., 142, 3452-3461 (1989)). The T cells of the invention can be used to form an immunogenic composition useful in treating or preventing cancer in a patient in need of therapy or protection (WO2006/031221).
- [0066] In the context of the present invention, the term "vaccine" (also referred to as an immunogenic composition) refers to a substance that induces anti-tumor immunity or suppresses cancers upon inoculation into animals. According to the present invention, polypeptides having the amino acid sequence of SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 80, 100, 101, 110, 111, 387, 112, 394, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217 or 223 were suggested to be HLA-A24 restricted epitope peptides and those of SEQ ID NO: 376, 379, 114, 116, 117, 121, 227, 228, 233, 254, 271, 272 or 288 were suggested to be HLA-A2 restricted epitope peptides that may induce potent and specific immune response against cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancer cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK

and/or URLC10. The cancers contemplated include, but are not limited to bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0067]

Thus, the present invention also encompasses a method of inducing anti-tumor immunity using polypeptides having the amino acid sequence of SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 or a variant thereof (i.e., including 1, 2, or several (e.g., up to 5) amino acid substitutions, deletions, or additions). In general, anti-tumor immunity includes immune responses such as follows:

- an induction of cytotoxic lymphocytes against tumors containing cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10,
- an induction of antibodies that recognize tumors containing cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, and
- an induction of anti-tumor cytokine production.

[0068]

Therefore, when a certain peptide induces any one of these immune responses upon inoculation into an animal, the peptide is decided to have anti-tumor immunity inducing effect. The induction of the anti-tumor immunity by a peptide can be detected by observing in vivo or in vitro the response of the immune system in the host against the peptide.

[0069]

For example, a method for detecting the induction of cytotoxic T lymphocytes is well known. A foreign substance that enters the living body is presented to T cells and B cells by the action of antigen-presenting cells (APCs). T cells that respond to the antigen presented by APC in antigen specific manner differentiate into cytotoxic T cells (also referred to as cytotoxic T lymphocytes or CTLs) due to stimulation by the antigen, and then proliferate; this process is referred to herein as "activation" of T cells. Therefore, CTL induction by a certain peptide can be evaluated by presenting the peptide to a T cell by APC, and detecting the induction of CTL. Furthermore, APCs have the effect of activating CD4+ T cells, CD8+ T cells, macrophages, eosinophils and NK cells. Since CD4+ T cells are also important in anti-tumor immunity, the anti-tumor immunity inducing action of the peptide can be evaluated using the activation effect of these cells as indicators.

[0070]

A method for evaluating the inducing action of CTL using dendritic cells (DCs) as APC is well known in the art. DC is a representative APC having the strongest CTL inducing action among APCs. In this method, the test polypeptide is initially contacted with DC and then this DC is contacted with T cells. Detection of T cells having

cytotoxic effects against the cells of interest after the contact with DC shows that the test polypeptide has an activity of inducing the cytotoxic T cells. Activity of CTL against tumors can be detected, for example, using the lysis of ^{51}Cr -labeled tumor cells as the indicator. Alternatively, it is well known to evaluate the degree of tumor cell damage using ^3H -thymidine uptake activity or LDH (lactose dehydrogenase)-release as the indicator. Furthermore, it can be also examined by measuring IFN-gamma produced and released by CTL in the presence of antigen-presenting cells that carry immobilized peptides by visualizing using anti-IFN-gamma antibodies, such as an ELISPOT assay. [0071]

Apart from DC, peripheral blood mononuclear cells (PBMCs) may also be used as the APC. The induction of CTL is reported to be enhanced by culturing PBMC in the presence of GM-CSF and IL-4. Similarly, CTL has been shown to be induced by culturing PBMC in the presence of keyhole limpet hemocyanin (KLH) and IL-7. [0072]

The test polypeptides confirmed to possess CTL inducing activity by these methods are polypeptides having DC activation effect and subsequent CTL inducing activity. Therefore, polypeptides that induce CTL against cells expressed CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10 are useful as vaccines against diseases associating CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers. Furthermore, APC that have acquired the ability to induce CTL against a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, by contacting with the polypeptides are useful as vaccines against the disease. Furthermore, CTL that have acquired cytotoxicity due to presentation of the polypeptide antigens by APC can be also used as vaccines against a disease associating CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers. Such therapeutic methods for a disease associating CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, using anti-tumor immunity due to APC and CTL, are referred to as cellular immunotherapy. The cancers contemplated include, but are not limited to, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[0073]

Generally, when using a polypeptide for cellular immunotherapy, efficiency of the CTL-induction can be increased by combining a plurality of polypeptides having different structures and contacting them with DC. Therefore, when stimulating DC with protein fragments, it is advantageous to use a mixture of multiple types of fragments.

[0074] The induction of anti-tumor immunity by a polypeptide can be further confirmed by observing the induction of antibody production against tumors. For example, when antibodies against a polypeptide are induced in a laboratory animal immunized with the polypeptide, and when growth, proliferation and/or metastasis of tumor cells is suppressed by those antibodies, the polypeptide is determined to induce anti-tumor immunity.

[0075] Anti-tumor immunity can be induced by administering a vaccine of this invention, and the induction of anti-tumor immunity enables treatment and prevention of a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers. Therapy against or prevention of the onset of a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, may include inhibition of the growth of cells expressing CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancer cells, involution of these cells and suppression of occurrence of these cells, e.g. cancer cells. Decrease in mortality of individuals having a disease associating CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, decrease of the disease markers in the blood, alleviation of detectable symptoms accompanying the disease and such are also included in the therapy or prevention of the disease, e.g. cancers. Such therapeutic and preventive effects are preferably statistically significant, for example, observed at a significance level of 5% or less, wherein the therapeutic or preventive effect of a vaccine against a disease associating CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, is compared to a control without vaccine administration. For example, Student's t-test, the Mann-Whitney U-test or ANOVA may be used for determining statistical significance.

[0076] In that the present invention provides a method for treating, or preventing a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, the therapeutic compounds or compositions may be administered prophylactically or therapeutically to subjects suffering from or at risk of (or susceptible to) developing the disease. Such subjects may be identified using standard clinical methods. In the context of the present invention, prophylactic administration occurs prior to the manifestation of overt clinical symptoms of disease, such that a disease or disorder is prevented or alternatively delayed in its progression. In the context of the field of medicine, the term "prevent" encompasses any activity which reduces the burden of mortality or morbidity from disease. Prevention can occur at primary, secondary and tertiary prevention levels. While primary prevention avoids the development of a disease, secondary and tertiary levels of prevention encompass activities aimed at preventing the progression

of a disease and the emergence of symptoms as well as reducing the negative impact of an already established disease by restoring function and reducing disease-related complications.

[0077] In the context of cancer treatment, the term "efficacious" refers to a treatment that leads to a decrease in size, prevalence or metastatic potential of cancer in a subject. When a treatment is applied prophylactically, "efficacious" means that the treatment retards or prevents occurrence of non cancer or alleviates a clinical symptom of cancer. The assessment of cancer can be made using standard clinical protocols. Furthermore, the efficaciousness of a treatment may be determined in association with any known method for diagnosing or treating cancer. For example, cancer can be diagnosed histopathologically or by identifying symptomatic anomalies.

[0078] The above-mentioned peptide, having immunological activity, or a polynucleotide or vector encoding such a peptide, may be combined with an adjuvant. An adjuvant refers to a compound that enhances the immune response against the peptide when administered together (or successively) with the peptide having immunological activity. Examples of suitable adjuvants include cholera toxin, salmonella toxin, alum and such, but are not limited thereto. Furthermore, a vaccine of this invention may be combined appropriately with a pharmaceutically acceptable carrier. Examples of such carriers are sterilized water, physiological saline, phosphate buffer, culture fluid and such. Furthermore, the vaccine may contain as necessary, stabilizers, suspensions, preservatives, surfactants and such. The vaccine is administered systemically or locally. Vaccine administration may be performed by single administration or boosted by multiple administrations.

[0079] When using APC or CTL as the vaccine of this invention, a disease associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or URLC10, e.g. cancers, can be treated or prevented, for example, by the ex vivo method. More specifically, PBMCs of the subject receiving treatment or prevention are collected, contacted ex vivo with a peptide of the present invention. Following the induction of APC or CTL, the cells may be administered to the subject. APC can be also induced by introducing a vector encoding the peptide into PBMCs ex vivo. APC or CTL induced in vitro can be cloned prior to administration. By cloning and growing cells having high activity of damaging target cells, cellular immunotherapy can be performed more effectively. Furthermore, APC and CTL isolated in this manner may be used for cellular immunotherapy not only against individuals from whom the cells are derived, but also against similar types of diseases in other individuals.

[0080] Aspects of the present invention are described in the following examples, which are presented only to illustrate the present invention and to assist one of ordinary skill in making and using the same. The examples are not intended in any way to otherwise

limit the scope of the invention.

[0081] Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below.

[0082] EXAMPLES

Hereinafter, the present invention is exemplified, but not restricted, by the following Examples. However, materials, methods and such described herein only illustrate aspects of the invention and in no way are intended to limit the scope of the present invention. As such, materials, methods and such similar or equivalent to those described therein may be used in the practice or testing of the present invention.

[0083] MATERIALS AND METHODS

Cell lines

A24-LCL cells (HLA-A24), human B-lymphoblastoid cell line, was established by transforming with Epstein-bar virus. T2 cell, COS7, A498, Caki-2 and HEK 293 were purchased from ATCC. Caki-1 and MIAPaca-2 were purchased from JCRB. PK-45P, PK-59, TE-5 and TE-6 were purchased from TKG. 293 T was purchased from GenHunter.

[0084] Candidate selection of peptide derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10

9-mer and 10-mer peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10 that bind to HLA-A*2402 or HLA-A*0201 molecule were predicted using the binding prediction software "BIMAS" (http://bimas.dcrt.nih.gov/cgi-bin/molbio/ken_parker_comboform) (Parker KC, et al., (1994) J Immunol.;152(1):163-75.; Kuzushima K, et al., (2001)

Blood.;98(6):1872-81.). These peptides were synthesized by Sigma (Sapporo, Japan) according to the standard solid phase synthesis method and purified by reversed phase HPLC. The purity (>90%) and the identity of the peptides were determined by analytical HPLC and mass spectrometry analysis, respectively. Peptides were dissolved in dimethylsulfoxide (DMSO) at 20 mg/ml and stored at -80 degrees C.

[0085] In vitro CTL Induction

Monocyte-derived dendritic cells (DCs) were used as antigen-presenting cells (APCs) to induce CTL responses against peptides presented on HLA. DCs were generated in vitro as described elsewhere (Nukaya I et al., (1999) Int. J. Cancer 80, 92-7., Tsai V et al., (1997) J. Immunol 158:1796-802.). Briefly, peripheral blood mononuclear cells (PBMCs) isolated from a normal volunteer (HLA-A*2402 and/or HLA-A*0201) by Ficoll-Paque (Pharmacia) solution were separated by adherence to a plastic tissue culture flask (Becton Dickinson) so as to enrich them for the monocyte fraction. The monocyte-enriched population was cultured in the presence of 1000 U/ml

of GM-CSF (Genzyme) and 1000 U/ml of IL-4 (Genzyme) in AIM-V (Invitrogen) containing 2% heat-inactivated autologous serum (AS). After 7 days in the culture, the cytokine-generated DCs were pulsed with 20 micro g/ml of the synthesized peptides in the presence of 3 micro g/ml of beta 2-microglobulin for 4 hrs at 20 degrees C in AIM-V. These peptide-pulsed DCs were then inactivated by MMC (30 micro g/ml for 30 mins) and mixed at a 1:20 ratio with autologous CD8⁺ T cells, obtained by positive selection with Dynabeads M-450 CD8 (Dynal) and DETACHa BEAD™ (Dynal). These cultures were set up in 48-well plates (Corning); each well contained 1.5x10⁴ peptide-pulsed DCs, 3x10⁵ CD8⁺ T cells and 10 ng/ml of IL-7 (Genzyme) in 0.5 ml of AIM-V/2% AS. Three days later, these cultures were supplemented with IL-2 (CHIRON) to a final concentration of 20 IU/ml. On day 7 and 14, the T cells were further restimulated with the autologous peptide-pulsed DCs. The DCs were prepared each time by the same way described above. CTL was tested against peptide-pulsed A24-LCL cells or T2 cells after the 3rd round of peptide stimulation on day 21.

[0086] CTL Expansion Procedure

CTLs were expanded in culture using the method similar to that described by Riddell SR, et al., (Walter EA et al., (1995) N Engl J Med 333:1038-44.; Riddel SR, et al., (1996) Nature Med. 2:216-23.). A total 5x10⁴ of CTLs were resuspended in 25 ml of AIM-V/5% AS with 2 kinds of human B-lymphoblastoid cell lines, inactivated by MMC, in the presence of 40 ng/ml of anti-CD3 monoclonal antibody (Pharmingen). One day after initiating the cultures, 120 IU/ml of IL-2 were added to the cultures. The cultures were fed with fresh AIM-V/5% AS containing 30 IU/ml of IL-2 on days 5, 8 and 11.

[0087] Establishment of CTL clones

The dilutions were made to have 0.3, 1, and 3 CTLs/well in 96 round-bottomed micro titer plate (Nalge Nunc International). CTLs were cultured with 7x10⁴ cells/well of 2 kinds of human B-lymphoblastoid cell lines, 30ng/ml of anti-CD3 antibody, and 125 U/ml of IL-2 in total of 150micro l/well of AIM-V containing 5%AS. 50 micro l / well of IL-2 was added to the medium 10 days later so that IL-2 became 125 U/ml in the final concentration. CTL activity of CTLs was tested on the 14th day, and CTL clones were expanded using the same method above.

[0088] Specific CTL activity

To examine the specific CTL activity, IFN-gamma ELISPOT assay and IFN-gamma ELISA assay were performed.

Briefly, peptide-pulsed A24-LCL or T2 cell (1x10⁴/well) was prepared as stimulator cells. Cultured Cells in 48 wells or CTL clones after limiting dilution were used as responder cells. IFN-gamma ELISPOT assay and ELISA assay were performed under

manufacture procedure.

[0089]

Establishment of the cells forcibly expressing either or both of the target gene and HLA-A02 or HLA-A24

The cDNA encoding an open reading frame of target genes or HLA-A02 or HLA-A24 was amplified by PCR. The PCR-amplified product was cloned into pcDNA3.1 myc-His vector (Invitrogen). The plasmids were transfected into the target cells, HLA-A02 and HLA-A24-null nomal human cell line COS7 or 293T using lipofectamine (Invitrogen) according to the manufacturer's recommended procedures. Alternatively, the plasmid contained the target genes were transfected into A24-LCL by electroporation using GenePulserII (Biorad). Briefly, 2.5×10^6 A24-LCL cells were pulsed with 10 mcg prasmid at 140V and 1000 micro F. After 2days from transfection, the transfected cells were treated with Cell dissociation solution and used as the target cells for CTL activity assay.

[0090]

Cytotoxicity Assay

Cytotoxic activity was evaluated by a four-hour ^{51}Cr release assay. The target cells were pulsed with a 20 micro g/mL concentration of peptide overnight. The target cells were labeled with 100 micro Ci of $\text{Na}_2^{51}\text{CrO}_4$ at 37 degrees C for one hour, and then washed three times with RPMI1640. The target cells (1×10^4 /100 micro L) and 100 micro L of effector cells at various numbers with a total volume of 200 micro L were placed into a round-bottomed 96-well microtiter plate (Corning), and cultured at 37 degrees C in a CO_2 incubator for four hours. After culturing, 100 micro L of the supernatant was collected from each well, and measured the radioactivity using a gamma counter. Spontaneous release was the radioactivity from the target cells with medium in the absence of effector cells, and maximum releasewas the radioactivity from the target cells with 1 M HCl.

The Percentage of specific cytotoxicity was determined by calculating as following formula:

% Specific lysis = [(experimental release - spontaneous release) / (maximum release - spontaneous release)] x100.

[0091]

RESULTS

Enhanced CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 expression in cancers

The global gene expression profile data obtained from various cancers using cDNA-microarray revealed that the expression of the following genes was elevated.

CDH3 (GenBank Accession No. NM_001793; SEQ ID Nos.1, 2),

EPHA4 (GenBank Accession No. L36645; SEQ ID Nos.3, 4),

ECT2 (GenBank Accession No. AY376439; SEQ ID Nos.5, 6),

HIG2 (GenBank Accession No. NM_013332; SEQ ID Nos.7, 8),

INHBB (GenBank Accession No. NM_002193; SEQ ID Nos.9, 10),
KIF20A (GenBank Accession No. NM_005733; SEQ ID Nos.11, 12),
KNTC2 (GenBank Accession No. AF017790; SEQ ID Nos.13, 14),
TTK (GenBank Accession No. NM_003318; SEQ ID Nos.15, 16) and
URLC10 (GenBank Accession No. NM_017527; SEQ ID Nos.17, 18)

CDH3 expression was validly elevated in the following cancers in comparison with corresponding normal tissue.

26 out of 34 bladder cancer,
17 out of 19 cervical cancer,
19 out of 19 cholangiocellular carcinoma,
30 out of 34 colorectal cancer,
20 out of 21 endometriosis,
13 out of 20 gastric cancer,
7 out of 8 diffuse-type gastric cancer,
36 out of 37 NSCLC,
16 out of 16 pancreatic cancer,
21 out of 21 soft tissue tumor and
10 out of 10 testicular tumor

[0092] EPHA4 expression was validly elevated in the following cancers in comparison with corresponding normal tissue.

14 out of 34 bladder cancer,
8 out of 14 cervical cancer,
10 out of 25 cholangiocellular carcinoma,
5 out of 15 endometriosis,
5 out of 8 diffuse-type gastric cancer,
5 out of 5 ovarian cancer,
14 out of 14 pancreatic cancer,
20 out of 51 prostate cancer and
14 out of 23 soft tissue tumor

[0093] ECT2 expression was validly elevated in the following cancers in comparing with corresponding normal tissue.

17 out of 19 bladder cancer,
5 out of 12 breast cancer,
14 out of 14 cervical cancer,
13 out of 13 cholangiocellular carcinoma,
5 out of 5 CML,
7 out of 8 colorectal cancer,
12 out of 16 esophageal cancer,

6 out of 16 NSCLC,
8 out of 10 lymphoma,
1 out of 1 pancreatic cancer,
10 out of 13 prostate cancer,
3 out of 6 renal carcinoma and
12 out of 13 SCLC cancer

HIG2 expression was validly elevated in 19 out of 20 renal cancer and 7 out of 9 soft tissue tumor in comparing with corresponding normal tissue.

INHBB expression was validly elevated in the following cancers in comparing with corresponding normal tissue.

10 out of 21 cholangiocellular carcinoma,
12 out of 12 esophageal cancer,
10 out of 13 NSCLC,
22 out of 24 renal carcinoma,
8 out of 14 SCLC cancer and
45 out of 49 soft tissue tumor

[0094] KIF20A expression was validly elevated in the following cancers in comparing with corresponding normal tissue.

31 out of 31 bladder cancer,
38 out of 61 breast cancer,
10 out of 11 cholangiocellular carcinoma,
7 out of 19 esophageal cancer,
21 out of 22 NSCLC,
6 out of 6 ovarian cancer,
17 out of 36 prostate cancer,
6 out of 11 renal carcinoma and
15 out of 15 SCLC

[0095] KNTC2 expression was validly elevated in the following cancers in comparing with corresponding normal tissue.

30 out of 32 bladder cancer,
47 out of 56 breast cancer,
10 out of 10 cervical cancer,
16 out of 22 cholangiocellular carcinoma,
17 out of 37 CML,
3 out of 10 colorectal cancer,
11 out of 46 esophagus cancer,
15 out of 19 NSCLC,
7 out of 8 lymphoma,

20 out of 24 osteosarcoma,
3 out of 5 ovarian cancer,
2 out of 2 pancreatic cancer,
15 out of 37 prostate cancer,
14 out of 19 renal carcinoma,
15 out of 15 SCLC and
40 out of 59 soft tissue tumor

[0096] TTK expression was validly elevated in the following cancers in comparing with corresponding normal tissue.

27 out of 27 bladder cancer,
25 out of 30 breast cancer,
15 out of 16 cervical cancer,
10 out of 10 cholangiocellular carcinoma,
5 out of 7 CML,
6 out of 10 colorectal cancer,
24 out of 44 esophageal cancer,
8 out of 15 liver cancer,
12 out of 12 NSCLC,
6 out of 6 lymphoma,
13 out of 16 osteoblastoma,
12 out of 17 prostate cancer,
15 out of 15 SCLC and
16 out of 33 soft tissue tumor

[0097] URLC10 expression was validly elevated in the following cancers in comparing with corresponding normal tissue

29 out of 29 bladder cancer,
15 out of 16 cervical cancer,
7 out of 7 cholangiocellular carcinoma,
7 out of 19 esophageal cancer,
3 out of 3 gastric cancer, 24 out of 27 NSCLC,
15 out of 19 osteosarcoma,
4 out of 5 pancreatic cancer,
33 out of 43 soft tissue tumor.

[0098]

[Table 1]

Ratio of cases observed up-regulation of *CDH3*, *EPHA4*, *ECT2*, *HIG2*, *INHBB*, *KIF20A*, *KNTC2*, *TTK* or *URLC10* in cancerous tissue as compared to normal corresponding tissue

	CDH3	EPHA4	ECT2	HIG2	INHBB
Bladder cancer	26/34	14/34	17/19	-	-
Breast cancer	-	-	5/12	-	-
Cervical cancer	17/19	8/14	14/14	-	-
Cholangiocellular carcinoma	19/19	10/25	13/13	-	10/21
CML	-	-	5/5	-	-
Colectal cancer	30/34	-	7/8	-	-
Endometriosis	20/21	5/15	-	-	-
Esophageal cancer	-	-	12/16	-	12/12
Gastric cancer	13/20	-	-	-	-
Diffuse-type Gastric cancer	7/8	5/8	-	-	-
Liver cancer	-	-	-	-	-
non-small cell lung cancer	36/37	-	6/16	-	10/13
Lymphoma	-	-	8/10	-	-
Osteosarcoma	-	-	-	-	-
Ovarian cancer	-	5/5	-	-	-
Pancreatic cancer	16/16	14/14	1/1	-	-
Prostate cancer	-	20/51	10/13	-	-
Renal carcinoma	-	-	3/6	19/20	22/24
Small cell lung cancer	-	-	12/13	-	8/14
Soft tissue tumor	21/21	14/23	-	7/9	45/49
Testicular tumor	10/10	-	-	-	-

KIF20A KNTC2 TTK URLC10

Bladder cancer	31/31	30/32	27/27	29/29
Breast cancer	38/61	47/56	25/30	-
Cervical cancer	-	10/10	15/16	15/16
Cholangiocellular carcinoma	10/11	16/22	10/10	7/7
CML	-	17/37	5/7	-
Colectal cancer	-	3/10	6/10	-
Endometriosis	-	-	-	-
Esophageal cancer	7/19	11/46	24/44	7/19
Gastric cancer	-	-	-	3/3
Diffuse-type Gastric cancer	-	-	-	-
Liver cancer	-	-	8/15	-
non-small cell lung cancer	21/22	15/19	12/12	24/27
Lymphoma	-	7/8	6/6	-
Osteosarcoma	-	20/24	13/16	15/19
Ovarian cancer	-	3/5	-	-
Pancreatic cancer	6/6	2/2	-	4/5
Prostate cancer	17/36	15/37	12/17	-
Renal carcinoma	6/11	14/19	-	-
Small cell lung cancer	15/15	15/15	15/15	-
Soft tissue tumor	-	40/59	16/33	33/43
Testicular tumor	-	-	-	-

[0099] Prediction of HLA-A24 or HLA-A2 binding peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK or URLC10

Table 2 sets forth the HLA-A*2402 binding peptides for CDH3 in order of binding affinity. Table 2A sets forth 9-mer peptides derived from CDH3 and Table 2B sets forth 10-mer peptides derived from CDH3.

Table 3 sets forth the HLA-A*2402 and HLA-A*0201 binding peptides for EPHA4 in order of binding affinity. Table 3A sets forth the HLA-A*2402 binding 9-mer peptides derived from EPHA4, Table 3B shows the HLA-A*2402 binding 10-mer peptides derived from EPHA4 and Table 3C sets forth the HLA-A*0201 binding 9-mer peptides derived from EPHA4.

Table 4 sets forth the HLA-A*2402 binding peptides for ECT2 in order of binding affinity. Table 4A sets forth 9-mer peptides derived from ECT2 and Table 4B shows 10-mer peptides derived from ECT2.

Table 5 sets forth the HLA-A*2402 and HLA-A*0201 binding peptides for HIG2, Table 5A sets forth the HLA-A*2402 binding 9-mer peptides derived from HIG2, Table 5B sets forth the HLA-A*2402 binding 10-mer peptides derived from HIG2, Table 5C sets forth the HLA-A*0201 binding 9-mer peptides derived from HIG2, and Table 5D sets forth HLA-A*0201 binding 10-mer peptides derived from HIG2.

Table 6 sets forth the HLA-A*2402 and HLA-A*0201 binding peptides for INHBB, Table 6A shows the HLA-A*2402 binding 9-mer peptides derived from INHBB, Table 6B sets forth the HLA-A*2402 binding 10-mer peptides derived from INHBB, Table 6C sets forth the HLA-A*0201 binding 9-mer peptides derived from INHBB, and Table 6D sets forth HLA-A*0201 binding 10-mer peptides derived from INHBB.

Table 7 sets forth the HLA-A*2402 binding peptides for KIF20A in order of binding affinity. Table 7A sets forth 9-mer peptides derived from KIF20A and Table 7B sets forth 10-mer peptides derived from KIF20A.

Table 8 sets forth the HLA-A*2402 binding peptides for KNTC2 in order of binding affinity. Table 8A sets forth 9-mer peptides derived from KNTC2 and Table 8B sets forth 10-mer peptides derived from KNTC2.

Table 9 sets forth the HLA-A*0201 binding peptides for TTK in order of binding affinity. Table 9A sets forth 9-mer peptides derived from TTK and Table 9B sets forth 10-mer peptides derived from TTK.

Table 10 sets forth the HLA-A*0201 binding peptides for URLC10 in order of binding affinity. Table 10A sets forth 9-mer peptides derived from URLC10 and Table 10B sets forth 10-mer peptides derived from URLC10.

[0100] Explanation and definition about the terms in tables

Start position indicates the number of amino acid from N-terminal.

Binding score is derived from "BIMAS" described in Materials and Methods.

Positive donor number indicates the number of donor whoes CD8+ T-cells can be induced to the specific CTL by the ex vivo stimulation with antigen-presenting cells. This is shown as (positive donor number / hole donor nember).

Positive well number indicates the number of wells where specific IFN-gamma production can be detected by IFN-gamma ELISPOT assay. 4 to 8 wells can be prepared from one donor. This is shown as (positive wells number / the number of hole wells tested by IFN-gamma ELISPOT assay).

Positve CTL line indicates the number of CTL line established from positive wells. The generation of CTL line is determined by ELISA. This is shown as (established CTL line number / the number of positive wells tested by IFN-gamma ELISPOT assay).

No positive donor is not defined by no detectable positive wells, but by no established CTL line.

The peptides showed by bold character in tables possesses the stimulation activity of the T cells.

No data at positive donor number, positive well nember and positive CTL line indicating "-" means that the peptides can't be synthesized for any reason.

[0101]

[Table 2A-1]

HLA-A*2402 binding 9-mer peptides derived from *CDH3*

Strat position	Amino acid sequence	Binding Score	Positive donor number	Positive well number	Positive CTL line	SEQ ID NO.
513	IYEVMLAM	37.5	1/3			19
667	LFLLLVL L	36	-	-	-	20
30	VFREAEVTL	24	0/3	1/22	0/1	21
406	LYVEVTNEA	16.632	1/3			22
332	KYEAHVPEN	16.5	0/3	1/22	0/1	23
180	KYELFGH A V	15	0/3	1/22	0/1	24
85	RSLKERNPL	14.4	0/3	1/22	0/1	25
5	RGPLASLLL	12	0/3	2/22	0/2	26
652	KGGFILPVL	11.2	0/3	0/22	-	27
248	TYNGVVAYS	10.5	0/3	2/22	0/2	28
65	LFSTDNDDF	10	0/3	0/22	-	29
94	KIFPSKRIL	9.6	0/1	0/8	-	306
221	RGSVLEGVL	9.6	0/1	0/8	-	307
668	FLLLVL L LL	8.4	-	-	-	308
754	IGNFIENL	8.4	-	-	-	309
311	TAVAVVEIL	8.4	0/1	0/8	-	310
557	NQSPVRQVL	8.064	0/1	0/8	-	311
611	KQDTYDVHL	8	0/1	0/8	-	312
781	DYEGSGSDA	7.5	0/1	0/8	-	313
165	GWLLL N KPL	7.2	0/1	0/8	-	314

[Table 2A-2]

656	ILPVLGAVL	7.2	0/1	0/8	-	315
770	TAPPYDTLL	7.2	0/1	0/8	-	316
602	VVLSLKKFL	7.2	0/1	0/8	-	317
665	ALLFLLLVL	7.2	-	-	-	318
410	VTNEAPFVL	7.2	0/1	0/8	-	319
662	AVLALLFLL	7.2	-	-	-	320
613	DTYDVHLSL	6.72	0/1	0/8	-	321
6	GPLASLLLL	6	0/1	0/8	-	322
564	VLNITDKDL	6	0/1	0/8	-	323
159	AVEKETGWL	6	0/1	0/8	-	324
511	NNIYEVMVL	6	0/1	0/8	-	325
11	LLLLQVCWL	6	-	-	-	326
57	GCPGQEPAL	6	0/1	0/8	-	327
293	EYTLTIQAT	6	0/1	0/8	-	328
79	ETVQERRSL	6	0/1	0/8	-	329
475	SYRILRDPA	6	0/1	0/8	-	330
493	GQVTAVGTL	6	0/1	0/8	-	331
661	GAVLALLFL	6	0/1	0/8	-	332
388	GILTTRKGL	6	0/1	0/8	-	333
382	HPESNQGIL	6	0/1	0/8	-	334
663	VLALLFLL	5.76	-	-	-	335
598	EGDTVVLSL	5.6	0/1	0/8	-	336
278	TISVISSGL	5.6	0/1	2/8	0/2	337
659	VLGAVLALL	5.6	0/1	0/8	-	338
811	EWGSRFKKL	5.28	0/1	0/8	-	339
445	KVVEVQEGI	5.04	0/1	0/8	-	340
614	TYDVHLSLS	5	0/1	0/8	-	341
142	FYSITGPGA	5	0/1	0/8	-	342
246	IYTNYNGVVA	5	0/1	0/8	-	343

[0102]

[Table 2B-1]

HLA-A*2402 binding 10-mer peptides derived from *CDH3*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
807	DYLN ^c WGSRF	150	1/3			30
248	TYNG ^v VAYSI	105	0/3	4/22	0/4	31
667	LFLLIVLLL	42	-	-	-	32
397	DFEAKNQHTL	30	0/3	2/22	0/2	33
332	KYEAbVPENA	21	1/3			34
180	KYELFGHAVS	15	0/3	2/22	0/2	35
510	RNNIYEVMVL	12	0/3	4/22	0/4	36
5	RGPLASLLL	12	0/3	1/22	0/1	37
477	RILRD ^v PAGWL	12	0/3	1/22	0/1	38
556	CNQSPVRQVL	10.08	0/3	2/22	0/2	39
655	FILP ^v LGA ^v VL	8.64	1/3			344
662	AVLAILFLLL	8.64	-	-	-	345

[Table 2B-2]

277	GTISvISSGL	8.4	0/3	0/20	-	346
781	DYEGsGSDAA	7.5	0/3	0/20	-	347
601	TVVLsLKKFL	7.2	0/3	3/20	0/3	348
158	FAVEKETGWL	7.2	0/3	0/20	-	349
665	ALLFILLVLL	7.2	-	-	-	350
259	SQEPKDPHDL	7.2	0/3	0/20	-	351
664	LALLFLLLVL	7.2	-	-	-	352
42	GAEQePGQAL	7.2	0/3	1/20	0/1	353
661	GAVLaLLFLL	7.2	-	-	-	354
595	VNEEgDTVVL	7.2	0/2	0/12	-	355
340	NAVGhEVQRL	7.2	0/2	0/12	-	356
411	TNEApFVLKL	6.6	0/2	0/12	-	357
470	ENQKISYRIL	6	1/2			358
10	SLLLIQVCWL	6	0/2	1/12	0/1	359
721	GLEAPPEVVL	6	0/2	2/12	0/2	360
345	EVQRITVTDL	6	0/2	4/12	0/4	361
2	GLPRgPLASL	6	0/2	3/12	0/3	362
657	LPVLgAVLAL	6	-	-	-	363
563	QVLNITDKDL	6	0/2	1/12	0/1	364
159	AVEKETGWLL	6	0/2	2/12	0/2	365
492	SGQVtAVGTL	6	0/2	-	-	366
387	QGILITRKGL	6	0/2	-	-	367
525	SPPTtGTGTL	6	0/2	2/12	0/2	368
358	NSPAwRATYL	6	0/2	2/12	0/2	369
122	GPFPqRLNQL	5.6	0/2	3/12	0/3	370
753	EIGNHIIENL	5.6	0/2	1/12	0/1	371
310	TTAVaVVEIL	5.6	-	-	-	372
246	IYTYnGVVAY	5	0/2	2/12	0/2	373
805	DYDYINEWGS	5	0/2	0/12	-	374

[0103]

[Table 3A]

HLA-A*2402 binding 9-mer peptides derived from *EPHA4*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
97	VYIEIKFTL	504	0/2	1/16	0/1	40
453	RYSVALAWL	400	2/3			41
25	VYPANEVTL	300	0/3	0/22	-	42
384	HYTPQQNGL	288	0/3	1/22	0/1	43
5	FYFALFSCL	288	1/2			44
519	GYGDFSEPL	240	0/3	3/22	0/3	45
869	KFGQIVNML	67.2	1/3			46
777	AYTTRGGKI	55	0/3	1/22	0/1	47
420	KYNPNPDQS	18	1/3			48
749	RNILVNSNL	16.8	0/3	1/22	0/1	49
734	KYLSDMSYV	15	0/3	0/22	-	50
879	KLIRNPNSL	14.4	0/3	0/22	-	51
926	RYKDNFTAA	14.4	0/3	0/22	-	52
834	KAJEEGYRL	14.4	0/3	0/22	-	53
574	KYSKAKQEA	13.2	0/3	0/22	-	54
184	AFQDVGACI	12.6	0/3	1/22	0/1	55
252	WLVPIGNCL	12.096	0/3	0/22	-	56
326	RPPSAPLNL	12	0/3	0/22	-	57
203	KCPLTVRNL	12	0/3	0/22	-	58
360	SYNVVCKKC	11.55	0/3	0/22	-	59

[0104]

[Table 3B]

HLA-A*2402 binding 10-mer peptides derived from *EPHA4*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
25	VYPANEVTL	300	0/3	0/22	-	60
244	MYCGADGEWL	200	0/3	1/22	0/1	61
657	GYTDKQRRDF	120	0/3	1/22	0/1	62
5	FYFAIFSCLF	100	-	-	-	63
102	KFTLRDCNSL	48	0/3	1/22	0/1	64
818	SYGERPYWDM	30	0/3	2/22	0/2	65
4	IFYFALFSCL	28.8	-	-	-	66
808	SYGIVMWEVM	25	-	-	-	67
630	EFGEVCSGRL	24	0/3	0/22	-	68
420	KYNPNPDQSV	21.6	0/3	0/22	-	69
930	NFTAAGYTTL	20	0/2	0/16	-	70
675	QFDHPNIIHL	20	0/3	0/22	-	71
708	AFLRKNDGRF	15	0/3	0/22	-	72
579	KQEADEEKHL	12	0/3	1/22	0/1	73
727	RGIGSGMKYL	12	0/3	0/22	-	74
96	RVYIEIKFTL	11.2	0/2	1/16	0/1	75
507	SYVFHVRART	10.5	0/3	1/22	0/1	76
251	EWLVPIGNCL	10.08	0/3	0/22	-	77
24	RVYPANEVTL	9.6	1/3	-	-	78
699	EYMENGSLDA	9	0/3	0/22	-	79

[0105] [Table 3C]

HLA-A*0201 binding 9-mer peptides derived from *EPHA4*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
8	ALFSCLFGI	514.942	-	-	-	375
501	GLNPLTSYV	382.536	1/1	-	-	376
12	CLFGICDAV	126.098	0/1	1/5	0/1	377
977	QMHGRMVPV	115.534	0/1	1/5	0/1	378
165	KLNTEIRDV	111.979	1/1	-	-	379
252	WLVPIGNCL	98.267	0/1	1/5	0/1	380
879	KLIRNPNSL	74.768	0/1	1/5	0/1	381
559	VVILIAAFV	56.902	-	-	-	382
812	VMWEVMSYG	39.386	0/1	0/5	-	383
728	GIGSGMKYL	37.157	0/1	0/5	-	384
750	NILVNSNLV	35.385	0/1	1/5	0/1	385
937	TTLEAVVHV	33.705	0/1	1/5	0/1	386

[0106]

[Table 4A]

HLA-A*2402 binding 9-mer peptides derived from ECT2

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
515	TYPPFVNFF	216	1/1			80
140	LYCTSMMNL	200	0/1	0/8	-	81
298	LYVVKQEWF	150	0/1	0/8	-	82
435	NYVNILATI	105	0/1	0/8	-	83
773	IYTADPESF	100	0/1	0/8	-	84
110	LYKADCRVI	50	0/1	0/8	-	85
739	SFQMTSDEL	33	0/1	0/8	-	86
504	IFLKYSKDL	30	0/1	0/8	-	87
867	FFERRSHTL	30	0/1	0/8	-	88
178	DFNSKVTHL	30	0/1	0/8	-	89
61	KQEELIKAL	17.28	0/1	0/8	-	90
657	RGEQVTLFL	16.8	0/1	2/8	0/2	91
568	RLPSVALLL	16.8	0/1	0/8	-	92
550	KPECGRQSL	14.4	0/1	0/8	-	93
470	IFGSIPDIF	14	0/1	0/8	-	94
116	RVIGPPVVL	12	0/1	0/8	-	95
507	KYSKDLVKT	11	0/1	0/8	-	96
223	DFYAAVDDF	10	0/1	0/8	-	97

[0107] [Table 4B]

HLA-A*2402 binding 10-mer peptides derived from ECT2

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
322	LYEKaNTPEL	330	0/1	0/8	-	98
435	NYVNiLATH	90	0/1	0/8	-	99
40	SYVEeEMPOQI	90	1/1			100
101	DFQDsVFNDL	72.576	1/1			101
866	SFFEiRSHTL	24	0/1	0/8	-	102
811	SFSKtPKRAL	20	0/1	1/8	0/1	103
268	KYLPiGDERC	18	0/1	0/8	-	104
84	EFEGLIDSPEF	16.5	0/1	1/8	0/1	105
236	KVPPfQDCIL	14.4	0/1	0/8	-	106
728	RPPTeQANVL	14.4	0/1	0/8	-	107
507	KYSKdLVKTY	12	0/1	0/8	-	108
281	VVEEnIVKDL	10.08	0/1	0/8	-	109

[0108]

[Table 5A]

HLA-A*2402 binding 9-mer peptides derived from *HIG2*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
19	IFVRVMESL	42	1/3			110
22	RVMESLEGL	14.4	1/3			111
8	YLLGVVLTL	8.4	1/3			387
7	LYLLGVVLT	7.5	0/2	3/15	0/3	388
23	VMESLEGLL	7.2	0/2	0/16	-	389
9	LLGVVLTL	5.6	-	-	-	390

[0109] [Table 5B]

HLA-A*2402 binding 10-mer peptides derived from *HIG2*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
7	LYLLGVVLTL	420	1/3			112
22	RVMESLEGLL	17.28	0/3	4/24	0/4	113
8	YLLGVVLTL	8.4	-	-	-	391
5	LNLYLLGVVL	7.2	0/2	0/12	-	392
46	LANTEPTKGL	6	0/2	0/14	-	393
18	SIFVRVMESL	5.6	1/2			394

[0110] [Table 5C]

HLA-A*0201 binding 9-mer peptides derived from *HIG2*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
8	YLLGVVLTL	836.253	1/1			114
13	VLTLLSIFV	650.311	0/1	0/12	-	115
15	TLLSIFVRV	488.951	1/1			116
4	VLNLYLLGV	271.948	1/1			117
9	LLGVVLTL	83.527	0/1	0/12	-	118
22	RVMESLEGL	31.957	0/1	0/12	-	119
6	NLYLLGVVL	28.027	0/1	0/12	-	120

[0111]

[Table 5D]

HLA-A*0201 binding 10-mer peptides derived from *HIG2*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
8	YLLGvVLTL	836.253	1/1			121
12	VVLTILSIFV	210.538	-	-	-	122
29	GLLEsPSPGT	113.047	0/1	0/12	-	123
6	NLYLIGVVLT	54.847	-	-	-	124
4	VLNLyLLGVV	14.495	0/1	0/12	-	125
15	TLLSiFVRVM	13.174	0/1	0/12	-	126
18	SIFVrVMESL	12.248	0/1	0/12	-	127
14	LTLLsIFVRV	11.545	-	-	-	128

[0112] [Table 6A]

HLA-A*2402 binding 9-mer peptides derived from *INHBB*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
383	LYFDDEYNI	60	0/3	0/20	-	129
238	LFERGERRL	30	0/3	1/19	0/1	130
7	RALGAACLL	12	0/3	0/21	-	131
388	EYNIVKRDV	10.5	0/3	0/18	-	132
180	LYLKLLPYV	9	1/2			395
163	ISNEGNQNL	8.64	0/1	0/8	-	396
223	RSGWHTFPL	8	0/1	0/6	-	397
176	ASLWLYLKL	7.92	0/1	0/7	-	398
338	AYLAGVPGS	7.5	0/1	1/7	0/1	399
213	NMVEKRVDL	7.2	0/1	0/8	-	400
102	AMVTALRKL	6.6	0/1	0/8	-	401
250	VQCDSCQEL	6.336	0/1	0/8	-	402
369	NSCCIPTKL	6.16	0/1	0/8	-	403
330	NYCEGSCPA	6	0/1	0/7	-	404
172	FVVQASLWL	6	0/1	0/8	-	405
355	VNQYRMRGL	6	0/1	0/8	-	406
307	QFFIDFRLI	6	0/1	0/7	-	407
14	LLLLAAGWL	6	-	-	-	408
306	QQFFIDFRL	5.6	0/1	0/6	-	409
170	NLFVVQASL	5.6	0/1	0/7	-	410
327	YYGNYCEGS	5	0/1	1/8	0/1	411

[0113]

[Table 6B]

HLA-A*2402 binding 10-mer peptides derived from *INHBB*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
180	LYLKLLPYVL	360	1/3			133
171	LFVVQASLWL	30	-	-	-	134
305	RQQFFIDFRL	16.8	1/3			135
73	DFLEAVKRHI	12.6	0/3	4/20	0/4	136
7	RALGAAACLLL	12	1/3			137
273	RPFVVVQARL	11.2	0/3	1/20	0/1	138
338	AYLAGVPGSA	10	0/3	2/20	0/2	139
169	QNLFVVQASL	8.4	0/1	1/6	0/1	412
249	DVQCdSCQEL	7.92	0/1	4/6	0/4	413
173	VVQAsLWLYL	7.2	0/1	0/6	-	414
383	LYFDdEYNIV	7.2	0/1	0/6	-	415
229	FPLTeAIQAL	7.2	0/1	1/6	0/1	416
299	RTNLcCRQQF	7.2	0/1	5/6	0/5	417
101	AAMVIALRKL	6.6	0/1	2/6	0/2	418
368	VNSCCeIPTKL	6.16	0/1	2/6	0/2	419
13	CLLLIAAGWL	6	-	-	-	420
354	VVNQyRMRGL	6	0/1	0/6	-	421
150	DGLAsSRVRL	6	0/1	2/6	0/2	422
293	GLECdGRTNL	6	0/1	0/6	-	423
330	NYCEgSCPAY	6	0/1	1/6	0/1	424
176	ASLWIYLKLL	6	0/1	1/6	0/1	425
212	WNMVeKRVDL	6	1/1			426
74	FLEAvKRHIL	6	0/1	2/6	0/2	427
331	YCEGgCPAYL	6	0/1	1/6	0/1	428
77	AVKRhILSRL	5.6	0/1	1/6	0/1	429
175	QASLwLYLKL	5.28	0/1	2/6	0/2	430
326	GYYGnYCEGS	5	0/1	1/6	0/1	431
159	LYFFFiSNEGN	5	0/1	4/6	0/4	432
327	YYGNyCEGSC	5	0/1	1/6	0/1	433

[0114]

[Table 6C]

HLA-A*0201 binding 9-mer peptides derived from *INHBB*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
177	SLWLYLKLL	407.808	0/1	0/8		140
14	LLLLAAGWL	96.074	-	-	-	141
170	NLFVVQASL	79.041	0/1	0/8		142
213	NMVEKRVDL	63.256	0/1	0/8		143
172	FVHQASLWL	47.291	0/1	0/8		144
306	QQFFIDFRL	46.48	0/1	0/8		145
281	RLGDSRHRI	42.774	0/1	0/8		146
174	VQASLWLYL	34.427	0/1	0/8		147
257	ELAVVPVFV	28.69	0/1	1/8	0/1	148
313	RLIGWNDWI	28.116	0/1	1/8	0/1	149
139	RVSEIISFA	22.546	0/1	3/8	0/3	150
151	GLASSRVRL	21.362	0/1	0/8		151
8	ALGAACLLL	21.362	0/1	1/8	0/1	152
250	VQCDSCQEL	15.096	0/1	1/8	0/1	153

[0115] [Table 6D]

HLA-A*0201 binding 10-mer peptides derived from *INHBB*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
179	WLYLKLLPYV	12951.1	0/1	1/8	0/1	154
301	NLCCRQQFFI	332.806	0/1	0/8		155
237	ALFERGERRL	64.814	0/1	0/8		156
382	MLYFDDEYNI	56.754	0/1	0/8		157
13	CLLLAAGWL	56.514	-	-	-	158
8	ALGAACLLL	49.134	-	-	-	159
313	RLIGWNDWII	32.081	0/1	0/8		160
173	VVQASLWLYL	29.711	0/1	2/8	0/2	161
256	QELAVVPVFV	27.521	0/1	0/8		162
162	FISNEGNQNL	13.512	0/1	1/8	0/1	163
305	RQQFFIDFRL	12.562	0/1	0/8		164
362	GLNPGTVNSC	11.426	0/1	0/7		165
85	RLQMRGRPNI	10.433	0/1	1/8	0/1	166
69	RVDGDFLEAV	10.425	0/1	0/8		167

[0116]

[Table 7A]

HLA-A*2402 binding 9-mer peptides derived from *KIF20A*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
308	IYNELLYDL	432	0/2	0/14	-	168
621	MYEEKLNIL	432	0/2	0/14	-	169
67	VYLRVRPLL	420	0/2	0/14	-	170
499	KFSAIASQL	56	0/2	0/14	-	171
304	SFFEIYNEL	44.352	0/2	0/14	-	172
187	IFNSLQGQL	36	0/2	0/14	-	173
305	FFEIYNELL	30	1/2			174
23	MFESTAADL	30	0/2	0/14	-	175
256	SFDSGIAGL	20	0/2	0/14	-	176
298	RFSTIWISFF	20	-	-	-	177
383	IFSIRILHL	20	1/2			178
647	KIEELEALL	17.28	0/2	0/14	-	179
625	KLNILKESL	14.4	0/2	0/14	-	180
695	KLQQCKAEL	13.2	0/2	0/14	-	181
726	FTIDVDKKL	11.088	0/2	0/14	-	182
688	QLQEVKAKL	11.088	0/2	0/14	-	183

[0117] [Table 7B]

HLA-A*2402 binding 10-mer peptides derived from *KIF20A*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
308	IYNELLYDLL	432	0/2	0/14	-	184
182	RSLAIIIFNSL	24.192	0/2	1/14	0/1	185
304	SFFEIYNELL	24	1/2			186
742	RLLRRELQKL	15.84	0/2	0/14	-	187
739	KNIRILRTEL	15.84	0/2	0/14	-	188
218	RQEEmKKLSL	14.4	0/2	2/14	0/2	189
70	RVRPILPSEL	12.672	0/2	0/14	-	190
871	RILRsRRSPL	12	0/2	0/14	-	191
89	RIENvETLVL	12	0/2	1/14	0/1	192
364	KNQSFASTHL	12	0/2	0/14	-	193
66	KVYLRVRPLL	11.2	1/2			194
60	DSMEmVKVYL	10.08	0/2	0/14	-	195

[0118]

[Table 8A]

HLA-A*2402 binding 9-mer peptides derived from *KNTC2*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
309	KYQAYMSNL	600	1/3			196
457	VYVPLKELL	432	0/3	0/18	-	197
414	EYHKLARKL	264	0/3	0/18	-	198
139	SYELPDTKF	165	0/3	0/18	-	199
629	KYEKKATLI	150	0/3	0/18	-	200
400	KYARGKEAI	100	0/3	1/18	0/1	201
124	DFLKIFTFL	50.4	1/3			202
134	GFLCPSYEL	33	0/3	0/18	-	203
257	LFNVDAFKL	33	0/3	0/18	-	204
242	SFDEMNAEL	26.4	0/3	0/18	-	205
128	IFTFLYGFL	24	0/3	0/18	-	206
146	KFEEEVPRI	18	0/3	1/18	0/1	207
368	RINHERNEL	15.84	0/3	1/18	0/1	208
235	SFMSGADSF	15	0/3	0/18	-	209
154	IFKDLGYPF	14.4	1/3			210
563	EYQLVVQTT	12.6	0/3	0/18	-	211
474	KALNKKMGL	12	0/3	1/18	0/1	212
150	EVPRIFKDL	10.08	1/3			213

[0119] [Table 8B]

HLA-A*2402 binding 10-mer peptides derived from *KNTC2*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
452	KYRAQVYVPL	560	2/3			214
610	EYEECMSEDL	360	0/3	1/18	0/1	215
360	KYSVADIERI	100	0/3	0/18	-	216
227	DYTIKCYESF	100	1/3			217
146	KFEEEVPRI	50.4	0/3	0/18	-	218
90	AFIQQCIRQL	30	0/3	0/18	-	219
20	RSQDVNKQGL	17.28	0/3	1/18	0/1	220
501	RTLKEEVQKL	15.84	0/3	0/18	-	221
403	RGKEAIETQL	13.44	0/3	1/18	0/1	222
273	RALNEQIARL	12	1/3			223
563	EYQLVVQTTT	10.5	0/3	3/22	0/3	224
467	ETEEEINKAL	10.08	0/3	1/22	0/1	225
541	LLESTVNQGL	10.08	0/3	1/22	0/1	226

[0120]

[Table 9A]

HLA-A*0201 binding 9-mer peptides derived from *TTK*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
462	YMSCFRTPV	878.055	1/1			227
547	KQIYAIKYV	312.218	1/1			228
630	NMLEAVHTI	262.897	0/1	1/8	0/1	229
278	LLNSPDCDV	118.238	0/1	1/8	0/1	230
498	ILATPLQNL	83.527	0/1	0/8	-	231
811	YVLGQLVGL	73.172	0/1	0/8	-	232
719	SLGCILYYM	62.845	1/2			233
670	QMQPDTTSV	50.232	0/1	0/8	-	234
804	GTTEEMKYV	50.102	0/1	0/8	-	235
654	LIVDGMLKL	47.088	0/1	1/8	0/1	236
363	SLLAKLEET	31.074	0/1	0/8	-	237
790	YVQIQTHPV	27.995	0/1	0/8	-	238
785	LLAHPYVQI	26.604	0/1	0/8	-	239
86	KLIGRYSQA	26.082	0/1	0/8	-	240
186	NLNQKKQL	21.362	0/1	0/8	-	241
671	MQPDITTSVV	20.152	0/1	0/8	-	242
577	KLQQHSDKI	17.892	0/1	0/8	-	243
142	FAFVHISFA	14.856	0/1	0/8	-	244
322	CELRLNLKSV	11.509	0/1	0/8	-	245
824	SILKAAKTL	10.868	0/1	0/8	-	246

[0121]

[Table 9B]

HLA-A*0201 binding 10-mer peptides derived from *ITK*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
68	LLLKLEKNSV	437.482	0/1	0/8	-	247
277	NLLNSPDCDV	257.342	0/1	0/8	-	248
653	FLIVDGMLKL	226.014	0/1	0/8	-	249
423	TTFEQPVFSV	195.487	0/1	0/8	-	250
542	VLNEKKQIYA	190.448	0/1	0/8	-	251
658	GMLKLIDFGI	161.697	0/1	0/8	-	252
194	LLSEEEKKNL	148.896	0/1	0/8	-	253
462	YMSCFRTPVV	94.738	1/1			254
57	MMANNPEDWL	70.685	0/1	0/8	-	255
600	MVMECGNIDL	48.205	0/1	0/8	-	256
689	YMPPEAIKDM	37.961	0/1	0/8	-	257
86	KLIGRYSQAI	36.515	0/1	0/8	-	258
669	NQMQPDTTSV	26.092	0/1	1/8	0/1	259
497	QILATPLQNL	24.997	0/1	0/8	-	260
654	LIVDGMLKLI	22.997	0/1	0/8	-	261
186	NLNQKKQLL	21.362	0/1	1/8	0/1	262
670	QMQPDTTSVV	20.595	0/1	0/8	-	263
803	KGTTEEMKYV	20.102	0/1	0/8	-	264
11	LTIDSIMNKV	15.486	0/1	0/8	-	265
577	KLQQHSDKII	14.971	0/1	0/8	-	266

[0122]

[Table 10A]

HLA-A*0201 binding 9-mer peptides derived from *URLC10*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
131	KIFPRFFMV	1364.78	0/1	0/8	-	267
204	GLWLAILL	407.808	0/1	0/8	-	268
65	LLVVALPRV	271.948	0/1	0/8	-	269
60	ALLALLLVV	242.674	-	-	-	270
206	WLAILLLA	52.561	1/1			271
212	LLASIAAGL	36.316	1/1			272
210	LLLLASIAA	31.249	0/1	0/8	-	273
137	FMVAKQCSA	16.505	0/1	2/8	0/2	274
58	TMALLALLL	15.428	0/1	2/8	0/2	275
59	MALLALLLV	13.975	0/1	2/8	0/2	276
209	ILLLASIA	12.812	0/1	0/8	-	434
208	AILLLLASI	12.208	-	-	-	277
69	ALPRVWTDA	8.446	0/1	0/8	-	278
197	SMGESCGGL	8.223	0/1	0/8	-	279
61	LLALLLVVA	7.964	-	-	-	280
67	VVALPRVWT	6.097	0/1	0/8	-	281
72	RVWT DANLT	5.412	0/1	0/8	-	282
160	FLLEEPMPF	5.2	0/1	1/8	0/1	283
62	LALLLVVAL	4.292	0/1	0/8	-	284
57	GTMALLALL	2.525	0/1	1/8	0/1	285

[0123]

[Table 10B]

HLA-A*0201 binding 10-mer peptides derived from *URLC10*

strat position	sequence	Binding Score	positive donor number	positive well number	positive CTL line	SEQ ID NO
64	LLLVVVALPRV	1006.21	0/1	0/8	-	286
204	GLWLAILLL	407.808	0/1	1/8	0/1	287
211	LLLASIAAGL	134.369	1/1			288
258	TMALLALLLV	115.534	-	-	-	289
61	LLALLLVVAL	83.527	-	-	-	290
160	FLLEEPMPFF	65.782	0/1	0/8	-	291
209	ILLLLASIAA	31.249	0/1	0/8	-	292
131	KIFPRFFMVA	26.186	0/1	0/8	-	293
60	ALLALLLVVA	17.334	-	-	-	294
66	LVVALPRVWT	6.097	0/1	0/8	-	295
59	MALLALLLVV	5.73	-	-	-	296
2	RLQRPRQAPA	4.968	0/1	1/8	0/1	297
112	CQNPRRCKWT	4.156	0/1	0/8	-	298
72	RVWTDANLTA	3.608	0/1	0/8	-	299
53	WAPLGTMALL	3.139	0/1	0/8	-	300
121	TEPYCVIAAV	3.111	0/1	0/8	-	301
162	LEEPMPFFYL	2.739	0/1	1/8	0/1	302
181	LEGPPINSSV	2.299	0/1	2/8	0/2	303
170	YLKCCCKIRYC	2.024	0/1	0/8	-	304
130	VKIFPRFFMV	1.81	0/1	0/8	-	305

[0124] Stimulation of the T cells using the predicted peptides from CDH3 restricted with HLA-A*2402 and establishment for CTL lines stimulated with CDH3 derived peptides

CTLs for those peptides derived from CDH3 were generated according to the protocols set forth in "Materials and Methods" section above. Resulting that CTLs having detectable specific CTL activity, as determined by IFN-gamma ELISPOT assay, are shown in Figure 1. In particular, CDH3-A24-9-513 (SEQ ID NO: 19), CDH3-A24-9-406 (SEQ ID NO: 22), CDH3-A24-10-807 (SEQ ID NO: 30), CDH3-A24-10-332 (SEQ ID NO: 34), CDH3-A24-10-655 (SEQ ID NO: 344) and CDH3-A24-10-470 (SEQ ID NO: 358) demonstrated potent IFN-gamma production as compared to the control by IFN-gamma ELISPOT assay, and the cells in the positive well number #5 stimulated with SEQ ID NO: 19, #2 with SEQ ID NO: 22, #5 with SEQ ID NO: 30, #4 with SEQ ID NO: 34, #1 with SEQ ID NO: 344 and #4 with SEQ ID NO: 358 were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Results are shown in Figure 1. While, other peptides shown in table 2 could not establish the CTL

lines despite possible binding activity with HLA-A*2402. For example, the typical negative peptide (CDH3-A24-10-248) were shown in Figure 1a. In this invention, the peptiedes which could establish CTL line were selected as potent CTL stimulation peputide.

[0125] Establishment for CTL clones stimulated with CDH3 derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from CDH3-A24-10-807 (SEQ ID NO: 30) #5 and CDH3-A24-10-655 (SEQ ID NO: 344) #1 CTL line are shown in Figure 1f and g. CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0126] Specific CTL activity against the target cells expressing CDH3 and HLA-A*2402

The established CTL line raised against these peptides were examined for their ability to recognize the target cells expressing CDH3 and HLA-A*2402. Specific CTL activity against COS7 transfected with both full length CDH3 gene and the HLA-A*2402 molecule, which serves as a specific model for the target cells endogenously express CDH3 and HLA-A*2402, was tested using as effector cells the CTL lines raised by CDH3-A24-10-807 (SEQ ID NO: 30) and CDH3-A24-10-655 (SEQ ID NO: 344). COS7 transfected with full length CDH3 but not HLA-A*2402 and COS7 transfected with HLA-A*2402 but not full length CDH3 were prepared as controls. The CTL clones demonstrating the highest specific CTL activity against COS7 was that transfected with both CDH3 and HLA-A2402 (Figure 1f and g).

These results clearly demonstrate that CDH3-A24-10-807 (SEQ ID NO: 30) and CDH3-A24-10-655 (SEQ ID NO: 344) are naturally expressed on the target cell surface with HLA-A2402 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which may serve as cancer vaccines targeting CDH3 expressed tumors.

[0127] Stimulation of the T cells using the predicted peptides from EPHA4 restricted with HLA-A*2402 or HLA-A*0201, and establishment for CTL lines stimulated with EPHA4 derived peptides

CTLs for those peptides derived from EphA4 were generated by IFN-gamma ELISPOT assay. Resulting that CTLs having detectable specific CTL activity , as determined by IFN-gamma ELISPOT assay, are shown in Figure 2. In particular, EphA4-A24-9-453 (SEQ ID NO: 41), EphA4-A24-9-5 (SEQ ID NO: 44), EphA4-A24-9-869 (SEQ ID NO: 46), EphA4-A24-9-420 (SEQ ID NO: 48), EphA4-A24-10-24 (SEQ ID NO: 78), EphA4-A02-9-501 (SEQ ID NO: 376) and EphA4-A02-9-165 (SEQ ID NO: 379) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #3 stimultaed

with EphA4-A24-9-453 (SEQ ID NO: 41), #2 with EphA4-A24-9-5 (SEQ ID NO: 44), #5 with EphA4-A24-9-869 (SEQ ID NO: 46), #6 with EphA4-A24-9-420 (SEQ ID NO: 48), #4 with EphA4-A24-10-24 (SEQ ID NO: 78), #8 with EphA4-A02-9-501 (SEQ ID NO: 376) and #3 with EphA4-A02-9-165 (SEQ ID NO: 379) were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Especially, CTL lines stimulated with EphA4-A02-9-501 (SEQ ID NO: 376) and EphA4-A02-9-165 (SEQ ID NO: 379) were tested by ⁵¹Cr-release assay according to the protocols set forth in the "Materials and Methods" section above. Results are shown in Figure 2a-h. While, other peptides shown in table 3 could not establish the CTL lines despite possible binding activity with HLA-A*2402 or HLA-A*0201. For example, the typical negative peptide (EphA4-A24-9-384) were shown in Figure 2a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptides.

[0128]

Stimulation of the T cells using the predicted peptides from ECT2 restricted with HLA-A*2402, and establishment for CTL lines stimulated with ECT2 derived peptides
CTLs for those peptides derived from ECT2 were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by an IFN-gamma ELISPOT assay are shown in Figure 3. In particular, ECT2-A24-9-515 (SEQ ID NO: 80), ECT2-A24-10-40 (SEQ ID NO: 100) and ECT2-A24-10-101 (SEQ ID NO: 101) showed potent IFN-gamma production, and the cells in the positive well number #7 stimulated with ECT2-A24-9-515 (SEQ ID NO: 80), #2 with ECT2-A24-10-40 (SEQ ID NO: 100) and #1 with ECT2-A24-10-101 (SEQ ID NO: 101) were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Results are shown in Figure 3a-d. While, other peptides shown in table 4 could not establish the CTL lines despite possible binding activity with HLA-A*2402. For example, the typical negative peptide (ECT2-A24-10-322, ECT2-A24-9-657 and ECT2-A24-10-811) were shown in Figure 3a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptide.

[0129]

Establishment for CTL clones stimulated with ECT2 derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from ECT2-A24-10-40 (SEQ ID NO: 100) #2 CTL line are shown in Figure 3c. CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0130] Specific CTL activity against the target cells expressing ECT2 and HLA-A*2402

The established CTL line raised against these peptides were examined for their ability to recognize the target cells expressing ECT2 and HLA-A*2402. Specific CTL activity against COS7 transfected with both full length ECT2 gene and the HLA-A*2402 molecule, which serves as a specific model for the target cells endogenously express ECT2 and HLA-A*2402, was tested using as effector cells the CTL clone raised by ECT2-A24-10-40 (SEQ ID NO: 100) and the CTL line raised by ECT2-A24-10-101 (SEQ ID NO: 101). COS7 transfected with full length ECT2 but not HLA-A*2402 and COS7 transfected with HLA-A*2402 but not full length ECT2 (replaced other gene e.g. URLC10 or INHBB) were prepared as controls. The CTL line demonstrating the highest specific CTL activity against COS7 was that transfected with both ECT2 and HLA-A2402 (Figure 3c and d).

These results clearly demonstrate that ECT2-A24-10-40 (SEQ ID NO: 100) and ECT2-A24-10-101 (SEQ ID NO: 101) are naturally expressed on the target cell surface with HLA-A2402 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which may serve as cancer vaccines targeting ECT2 expressed tumors.

[0131] Cytotoxic activity against cancer cell line endogenously expressing HLA-A*2402 and ECT2

Furthermore, Cytotoxic activity was performed by cytotoxicity assay according to the protocols set forth in the "Materials and Methods" section above. As a result, as shown in Fig. 3b, CTL clone stimulated with ECT2-A24-9-515 (SEQ ID NO: 80) showed remarkably high cytotoxic effect towards HLA-A24-positive and ECT-positive cancer cell lines TE6, compared to that towards HLA-A24-negative and ECT-positive cancer cell lines TE5.

[0132] Stimulation of the T cells using the predicted peptides from HIG2 restricted with HLA-A*2402 or HLA-A*0201, and establishment for CTL lines stimulated with HIG2 derived peptides

CTLs for those peptides derived from HIG2 were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by an IFN-gamma ELISPOT assay are shown in Figure 4. In particular, HIG2-A24-9-19 (SEQ ID NO: 110), HIG2-A24-9-22 (SEQ ID NO: 111), HIG2-A24-9-8 (SEQ ID NO: 387), HIG2-A24-10-7 (SEQ ID NO: 112), HIG2-A24-10-18 (SEQ ID NO: 394), HIG2-A02-9-8 (SEQ ID NO: 114), HIG2-A02-9-15 (SEQ ID NO: 116), HIG2-A02-9-4 (SEQ ID NO: 117) and HIG2-A02-10-8 (SEQ ID NO: 121) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #6 stimulated with HIG2-A24-9-19 (SEQ ID NO:

110), #7 with HIG2-A24-9-22 (SEQ ID NO: 111), #5 with HIG2-A24-9-8 (SEQ ID NO: 387), #1 with HIG2-A24-10-7 (SEQ ID NO: 112), #7 with HIG2-A24-10-18 (SEQ ID NO: 394), #10 with HIG2-A02-9-8 (SEQ ID NO: 114), #10 with HIG2-A02-9-15 (SEQ ID NO: 116), #10 with HIG2-A02-9-4 (SEQ ID NO: 117) and #9 with HIG2-A02-10-8 (SEQ ID NO: 121) were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Results are shown in Figure 4a-j. While, other peptides shown in table 5 could not establish the CTL lines despite possible binding activity with HLA-A*2402. For example, the typical negative peptide (HIG2-A24-9-7) were shown in Figure 4a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptide.

[0133] Establishment for CTL clones stimulated with HIG2 derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from HIG2-A24-9-22 (SEQ ID NO: 111) #7 CTL line, HIG2-A24-9-8 (SEQ ID NO: 387) #5 CTL line, HIG2-A24-10-7 (SEQ ID NO: 112) #1 CTL line, HIG2-A24-10-18 (SEQ ID NO: 394) #7 CTL line and HIG2-A02-9-4 (SEQ ID NO: 117) #10 CTL line are shown in Figure 4c, e, f, g and i. CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0134] Specific CTL activity against the target cells expressing HIG2 and HLA-A*0201

The established CTL line raised against these peptides were examined for their ability to recognize the target cells expressing HIG2 and HLA-A*0201. Specific CTL activity against 293T or COS7 transfected with both full length HIG2 gene and the HLA-A*0201 molecule, which serves as a specific model for the target cells endogenously express HIG2 and HLA-A*0201, was tested using as effector cells the CTL lines raised by HIG2-A02-9-8 (SEQ ID NO: 114), HIG2-A02-9-15 (SEQ ID NO: 116) and the CTL clone raised by HIG2-A02-9-4 (SEQ ID NO: 117). 293T or COS7 transfected with full length ECT2 but not HLA-A*0201 and 293T or COS7 transfected with HLA-A*0201 but not full length ECT2 (or replaced other gene e.g. FoxP3 or TTK) were prepared as controls. The CTL line demonstrating the highest specific CTL activity against 293T or COS7 was that transfected with both ECT2 and HLA-A*0201 (Figure 4e, h and i).

[0135] These results clearly demonstrate that HIG2-A02-9-8 (SEQ ID NO: 114), HIG2-A02-9-15 (SEQ ID NO: 116) and HIG2-A02-9-4 (SEQ ID NO: 117) are naturally expressed on the target cell surface with HLA-A2402 or HLA-A0201 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which

may serve as cancer vaccines targeting HIG2 expressed tumors.

[0136]

Cytotoxic activity against cancer cell line endogenously expressing HLA-A*0201 and HIG2

Furthermore, Cytotoxic activity was performed by cytotoxicity assay according to the protocols set forth in the "Materials and Methods" section above. As a result, as shown in Fig. 4i, CTL clone stimulated with HIG2-A02-9-4 (SEQ ID NO: 117) showed remarkably high cytotoxic effect towards HLA-A02-positive and HIG2-positive cancer cell lines CAki-1, compared to that towards HLA-A02-negative and HIG2-positive cancer cell lines A498.

[0137]

Stimulation of the T cells using the predicted peptides from INHBB restricted with HLA-A*2402 or HLA-A*0201, and establishment for CTL lines stimulated with INHBB derived peptides

CTLs for those peptides derived from INHBB were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by an IFN-gamma ELISPOT assay are shown in Figure 5. In particular, INHBB-A24-9-180 (SEQ ID NO: 395), INHBB-A24-10-180 (SEQ ID NO: 133), INHBB-A24-10-305 (SEQ ID NO: 135), INHBB-A24-10-7 (SEQ ID NO: 137) and INHBB-A24-10-212 (SEQ ID NO: 426) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #7 stimulated with INHBB-A24-9-180 (SEQ ID NO: 395), #3 with INHBB-A24-10-180 (SEQ ID NO: 133), #2 with INHBB-A24-10-305 (SEQ ID NO: 135), #8 and #2 with INHBB-A24-10-7 (SEQ ID NO: 137) and #1 with INHBB-A24-10-212 (SEQ ID NO: 426) were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Results are shown in Figure 5b-e. While, other peptides shown in table 6 could not establish the CTL lines despite possible binding activity with HLA-A*2402 and HLA*0201. For example, the typical negative peptide (INHBB-A24-9-238) were shown in Figure 5a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptide.

[0138]

Establishment for CTL clones stimulated with INHBB derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from INHBB-A24-9-180 (SEQ ID NO: 395) #7 CTL line, and INHBB-A24-10-305 (SEQ ID NO: 135) #2 CTL line are shown in Figure 5b and d . CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0139]

Specific CTL activity against the target cells expressing INHBB and HLA-A*2402

The established CTL line raised against these peptides were examined for their ability to recognize the target cells expressing INHBB and HLA-A*2402. Specific CTL activity against 293T transfected with both full length INHBB gene and the HLA-A*2402 molecule, which serves as a specific model for the target cells endogenously express INHBB and HLA-A*2402, was tested using as effector cells the CTL lines raised by INHBB-A24-10-180 (SEQ ID NO: 133) and INHBB-A24-10-7 (SEQ ID NO: 137) and the CTL clone raised by INHBB-A24-10-305 (SEQ ID NO: 135)., 293T transfected with full length INHBB but not HLA-A*2402 and 293T transfected with HLA-A*2402 but not full length INHBB were prepared as controls. The CTL line demonstrating the highest specific CTL activity against 293T was that transfected with both INHBB and HLA-A*2402 (Figure 5c, d and e).

[0140] These results clearly demonstrate that INHBB-A24-10-305 (SEQ ID NO: 135), INHBB-A24-10-180 (SEQ ID NO: 133) and INHBB-A24-10-7 (SEQ ID NO: 137) are naturally expressed on the target cell surface with HLA-A2402 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which may serve as cancer vaccines targeting INHBB expressed tumors.

[0141] Cytotoxic activity against cancer cell line endogenously expressing HLA-A*2402 and INHBB

Furthermore, Cytotoxic activity was performed by cytotoxicity assay according to the protocols set forth in the "Materials and Methods" section above. As a result, as shown in Fig. 5b, CTL clone stimulated with INHBB-A24-9-180 (SEQ ID NO: 395) showed remarkably high cytotoxic effect towards HLA-A24-positive and INHBB - positive cancer cell lines MIAPaca2, compared to that towards HLA-A24-negative and INHBB -positive cancer cell lines CAki-2.

[0142] Stimulation of the T cells using the predicted peptides from KIF20A restricted with HLA-A*2402, and establishment for CTL lines stimulated with KIF20A derived peptides

CTLs for those peptides derived from KIF20A were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by an IFN-gamma ELISPOT assay are shown in Figure 6. In particular, KIF20A-A24-9-305 (SEQ ID NO: 174), KIF20A-A24-9-383 (SEQ ID NO: 178), KIF20A-A24-10-304 (SEQ ID NO: 186) and KIF20A-A24-10-66 (SEQ ID NO: 194) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #2 stimultaed with KIF20A-A24-9-305 (SEQ ID NO: 174), #3 with KIF20A-A24-9-383 (SEQ ID NO: 178), #5 with KIF20A-A24-10-304 (SEQ ID NO: 186) and #6 with KIF20A-A24-10-66 (SEQ ID NO: 194) were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-

pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Results are shown in Figure 6a-e. While, other peptides shown in table 7 could not establish the CTL lines despite possible binding activity with HLA-A*2402. For example, the typical negative peptide (KIF20A -A24-9-647 and KIF20A -A24-10-182) were shown in Figure 6a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptide.

[0143] Establishment for CTL clones stimulated with KIF20A derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from KIF20A-A24-9-305 (SEQ ID NO: 174) #2 CTL line, KIF20A-A24-10-304 (SEQ ID NO: 186) #5 CTL line and KIF20A-A24-10-66 (SEQ ID NO: 194) #6 CTL line are shown in Figure 6b, d and e. CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0144] Specific CTL activity against the target cells expressing KIF20A and HLA-A*2402

The established CTL line raised against these peptides were examined for their ability to recognize the target cells expressing KIF20A and HLA-A*2402. Specific CTL activity against COS7 transfected with both full length KIF20A gene and the HLA-A*2402 molecule and A24-LCL transfected by electroporation with full length KIF20A gene, which serve as a specific model for the target cells endogenously express KIF20A and HLA-A*2402, was tested using as effector cells the CTL lines raised by KIF20A-A24-9-383 (SEQ ID NO: 178) and KIF20A-A24-10-304 (SEQ ID NO: 186) and the CTL clone raised by KIF20A-A24-10-66 (SEQ ID NO: 194). COS7 transfected with full length KIF20A but not HLA-A*2402 and COS7 transfected with HLA-A*2402 but not full length KIF20A (or replaced full length URLC10 gene), COS7 transfected with HLA-A*2402 and pulsed with KIF20A-10-308, and A24-LCL transfected with mock vector were prepared as controls. The CTL line demonstrating the highest specific CTL activity against COS7 was that transfected with both KIF20A and HLA-A*2402 (Figure 6b, c and d). Alternatively, the CTL line stimulated with KIF20A-A24-10-304 (SEQ ID NO: 186) demonstrated against A24-LCL transfected with KIF20A.

[0145] These results clearly demonstrate that KIF20A-A24-9-383 (SEQ ID NO: 178), KIF20A-A24-10-304 (SEQ ID NO: 186) and KIF20A-A24-10-66 (SEQ ID NO: 194) is naturally expressed on the target cell surface with HLA-A2402 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which may serve as cancer vaccines targeting KIF20A expressed tumors.

[0146] Cytotoxic activity against cancer cell line endogenously expressing HLA-A*2402 and KIF20A

Furthermore, Cytotoxic activity was performed by cytotoxicity assay according to the protocols set forth in the "Materials and Methods" section above. As a result, as shown in Fig. 6b and e, CTL clone stimulated with KIF20A-A24-9-305 (SEQ ID NO: 174) or KIF20A-A24-10-304 (SEQ ID NO: 186) showed remarkably high cytotoxic effect towards HLA-A24-positive and KIF20A-positive cancer cell lines PK45P or MIAPaca2 respectively, compared to that towards HLA-A24-negative and KIF20A-positive cancer cell lines PK59.

[0147] Stimulation of the T cells using the predicted peptides from KNTC2 restricted with HLA-A*2402, and establishment for CTL lines stimulated with KNTC2 derived peptides

CTLs for those peptides derived from KNTC2 were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by an IFN-gamma ELISPOT assay are shown in Figure 7. In particular, KNTC2-A24-9-309 (SEQ ID NO: 196), KNTC2-A24-9-124 (SEQ ID NO: 202), KNTC2-A24-9-154 (SEQ ID NO: 210), KNTC2-A24-9-150 (SEQ ID NO: 213), KNTC2-A24-10-452 (SEQ ID NO: 214), KNTC2-A24-10-227 (SEQ ID NO: 217) and KNTC2-A24-10-273 (SEQ ID NO: 223) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #8 stimulated with KNTC2-A24-9-309 (SEQ ID NO: 196), #5 with KNTC2-A24-9-124 (SEQ ID NO: 202), #5 with KNTC2-A24-9-154 (SEQ ID NO: 210), #7 with KNTC2-A24-9-150 (SEQ ID NO: 213), #4 and #5 with KNTC2-A24-10-452 (SEQ ID NO: 214), #1 with KNTC2-A24-10-227 (SEQ ID NO: 217) and #8 with KNTC2-A24-10-273 (SEQ ID NO: 223) were expanded and CTL lines were established. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. Results are shown in Figure 7a-h. While, other peptides shown in table 8 could not establish the CTL lines despite possible binding activity with HLA-A*2402. For example, the typical negative peptide (KNTC2-A24-10-610) were shown in Figure 7a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptide.

[0148] Establishment for CTL clones stimulated with KNTC2 derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from KNTC2-A24-9-154 (SEQ ID NO: 210) #5 CTL line and KNTC2-A24-10-452 (SEQ ID NO: 214) #5 CTL line are shown in Figure 7d and f. CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0149] Specific CTL activity against the target cells expressing KNTC2 and HLA-A*2402

The established CTL line raised against these peptides were examined for their ability to recognize the target cells expressing KNTC2 and HLA-A*2402. Specific CTL activity against HEK293 transfected with both full length KNTC2 gene and the HLA-A*2402 molecule which serves as a specific model for the target cells endogenously express KNTC2 and HLA-A*2402, was tested using as effector cells the CTL clones raised by KNTC2-A24-10-452 (SEQ ID NO: 214). HEK293 transfected with full length KNTC2 but not HLA-A*2402, HEK293 transfected with HLA-A*2402 but not full length KNTC2 and HEK293 transfected with HLA-A*2402 and pulsed with KNTC2-9-309 were prepared as controls. The CTL line demonstrating the highest specific CTL activity against HEK293 was that transfected with both KNTC2 and HLA-A*2402 (Figure 7f).

[0150] These results clearly demonstrate that KNTC2-A24-10-452 (SEQ ID NO: 214) is naturally expressed on the target cell surface with HLA-A2402 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which may serve as cancer vaccines targeting KNTC2 expressed tumors.

[0151] Stimulation of the T cells using the predicted peptides from TTK restricted with HLA-A*0201, and establishment for CTL lines stimulated with TTK derived peptides

CTLs for those peptides derived from TTK were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by an IFN-gamma ELISPOT assay are shown in Figure 8. As depicted in Figure 8b-d, TTK-A2-9-462 (SEQ ID NO: 227), TTK-A2-9-547 (SEQ ID NO: 228), TTK-A2-9-719 (SEQ ID NO: 233) and TTK-A2-10-462 (SEQ ID NO: 254) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #4 stimulated with TTK-A2-9-462 (SEQ ID NO: 227), #2 with TTK-A2-9-547 (SEQ ID NO: 228), #1 with TTK-A2-9-719 (SEQ ID NO: 233) and #8 with TTK-A2-10-462 (SEQ ID NO: 254) were expanded. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. While, other peptides shown in table 9 could not establish the CTL lines despite possible binding activity with HLA-A*0201. For example, the typical negative peptide (TTK-A2-9-278) were shown in Figure 8a. In this invention, the peptides which could establish CTL line were selected as potent CTL stimulation peptide.

[0152] Establishment for CTL clones stimulated with TTK derived peptides

Furthermore, the limiting dilution from these CTL lines was performed according to the protocols set forth in the "Materials and Methods" section above. The establishment of CTL clones from TTK-A2-9-462 (SEQ ID NO: 227) #4 CTL line, TTK-A2-9-547 (SEQ ID NO: 228) #2 CTL line, TTK-A2-9-719 (SEQ ID NO: 233) #1 CTL

line and TTK-A2-10-462 (SEQ ID NO: 254) #8 CTL line were shown in Figure 8d, c, d and e. CTL clones had potent and specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse.

[0153] Specific CTL activity against the target cells expressing TTK and HLA-A*0201

The established CTL clone raised against these peptides were examined for their ability to recognize the target cells endogenously expressing TTK and HLA-A*0201. Specific CTL activity against COS7 transfected with both the full length TTK gene and the HLA-A*0201 molecule, which is a specific model for the target cells endogenously express TTK and HLA-A*0201, was tested using as effector cells the CTL clones raised by TTK-A2-9-462 (SEQ ID NO: 227), TTK-A02-9-547 (SEQ ID NO: 228), TTK-A2-9-719 (SEQ ID NO: 233) and TTK-A2-10-462 (SEQ ID NO: 254). COS7 transfected with full length TTK but HLA-A*0201, COS7 transfected HLA-A*0201 but not full length of TTK (or replaced full length HIG2 gene) and COS7 transfected with HLA-A*0201 and pulsed with different target epitope peptide, were prepared as controls. The CTL Clone having the highest specific CTL activity against COS7 was that transfected with both TTK and HLA-A*0201 (Figure 8b, c, d and e).

[0154] These results clearly demonstrate that TTK-A2-9-462 (SEQ ID NO: 227), TTK-A02-9-547 (SEQ ID NO: 228), TTK-A2-9-719 (SEQ ID NO: 233) and TTK-A02-10-462 (SEQ ID NO: 254) are naturally expressed on the target cell surface with HLA-A2 molecule and recognize CTL. Furthermore, these peptides are epitope peptides, which may serve as cancer vaccines targeting TTK expressed tumors.

[0155] Stimulation of the T cells using the predicted peptides from URLC10 restricted with HLA-A*0201, and establishment for CTL lines stimulated with URLC10 derived peptides

CTLs for those peptides derived from URLC10 were generated according to the protocols set forth in the "Materials and Methods" section above. Resulting CTLs having detectable specific CTL activity as determined by IFN-gamma ELISPOT assay are shown in Figure 9. As shown in Figure 9b-d, URLC-A2-9-206 (SEQ ID NO: 271), URLC-A2-9-212 (SEQ ID NO: 272) and URLC-A2-10-211 (SEQ ID NO: 288) demonstrated potent IFN-gamma production by IFN-gamma ELISPOT assay, and the cells in the positive well number #7 stimulated with URLC-A2-9-206 (SEQ ID NO: 271), #3 with URLC-A2-9-212 (SEQ ID NO: 272) and #5 with URLC-A2-10-211 (SEQ ID NO: 288) were expanded. Those CTL lines having higher specific CTL activities against the peptide-pulsed target as compared to the activities against target without peptide pulse were determined by ELISA. While, other peptides shown in table 10 could not establish the CTL lines despite possible binding activity with HLA-A*0201. For example, the typical negative peptide (URLC-A2-9-58) were shown in Figure 9a. In this invention, the peptide which could establish CTL line were selected

as potent CTL stimulation peptide.

[0156] Specific CTL activity against the target cells expressing URLC10 and HLA-A*0201

The established CTL line raised against these peptides were examined for their ability to recognize the target cells endogenously expressing URLC10 and HLA-A*0201. Specific CTL activity against COS7, Hek293 and 293T transfected with both full length URLC10 gene and the HLA-A*0201 molecule, which serves as a specific model for the target cells endogenously express URLC10 and HLA-A*0201, was tested using as effector cells the CTL line raised by URLC10-A02-10-211. COS7, Hek293 or 293T transfected with full length URLC10 but not HLA-A*0201 (replaced HLA-A*2402), COS7, Hek293 or 293T transfected with HLA-A*0201 but not full length URLC10 and COS7 transfected with HLA-A*0201 and pulsed with different target epitope peptide (URLC10-A02-10-64) were prepared as controls. The CTL line demonstrating the highest specific CTL activity against COS7, Hek293 or 293T was that transfected with both URLC10 and HLA-A*0201 (Figure 9-2).

[0157] These results clearly demonstrate that URLC10-A02-10-211 is naturally expressed on the target cell surface with HLA-A*0201 molecule and recognizes CTL. Furthermore, this peptide was epitope peptides, which may utilize cancer vaccine targeting URLC10 expressed tumors.

[0158] Homology analysis of the antigen peptides

The CTL clones established against the following peptides showed potent specific CTL activity.

CDH3-A24-9-513 (SEQ ID NO: 19),
 CDH3-A24-9-406 (SEQ ID NO: 22),
 CDH3-A24-10-807 (SEQ ID NO: 30),
 CDH3-A24-10-332 (SEQ ID NO: 34),
 CDH3-A24-10-655 (SEQ ID NO: 344),
 CDH3-A24-10-470 (SEQ ID NO: 358),
 EphA4-A24-9-453 (SEQ ID NO: 41),
 EphA4-A24-9-5 (SEQ ID NO: 44),
 EphA4-A24-9-869 (SEQ ID NO: 46),
 EphA4-A24-9-420 (SEQ ID NO: 48),
 EphA4-A24-10-24 (SEQ ID NO: 78),
 EphA4-A02-9-501 (SEQ ID NO: 376),
 EphA4-A02-9-165 (SEQ ID NO: 379),
 ECT2-A24-9-515 (SEQ ID NO: 80),
 ECT2-A24-10-40 (SEQ ID NO: 100),
 ECT2-A24-10-101 (SEQ ID NO: 101),
 HIG2-A24-9-19 (SEQ ID NO: 110),

HIG2-A24-9-22 (SEQ ID NO: 111),
HIG2-A24-9-8 (SEQ ID NO: 387),
HIG2-A24-10-7 (SEQ ID NO: 112),
HIG2-A24-10-18 (SEQ ID NO: 394),
HIG2-A02-9-8 (SEQ ID NO: 114),
HIG2-A02-9-15 (SEQ ID NO: 116),
HIG2-A02-9-4 (SEQ ID NO: 117),
HIG2-A02-10-8 (SEQ ID NO: 121),
INHBB-A24-9-180 (SEQ ID NO: 395),
INHBB-A24-10-180 (SEQ ID NO: 133),
INHBB-A24-10-305 (SEQ ID NO: 135),
INHBB-A24-10-7 (SEQ ID NO: 137),
INHBB-A24-10-212 (SEQ ID NO: 426),
KIF20A-A24-9-305 (SEQ ID NO: 174),
KIF20A-A24-9-383 (SEQ ID NO: 178),
KIF20A-A24-10-304 (SEQ ID NO: 186),
KIF20A-A24-10-66 (SEQ ID NO: 194),
KNTC2-A24-9-309 (SEQ ID NO: 196),
KNTC2-A24-9-124 (SEQ ID NO: 202),
KNTC2-A24-9-154 (SEQ ID NO: 210),
KNTC2-A24-9-150 (SEQ ID NO: 213),
KNTC2-A24-10-452 (SEQ ID NO: 214),
KNTC2-A24-10-227 (SEQ ID NO: 217),
KNTC2-A24-10-273 (SEQ ID NO: 223),
TTK-A02-9-462 (SEQ ID NO: 227),
TTK-A02-9-547 (SEQ ID NO: 228),
TTK-A02-9-719 (SEQ ID NO: 233),
TTK-A02-10-462 (SEQ ID NO: 254),
URLC-A02-9-206 (SEQ ID NO: 271),
URLC-A02-9-212 (SEQ ID NO: 272) and
URLC-A02-10-211 (SEQ ID NO: 288)

[0159] This suggests that the sequences of SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 are homologous to the peptides derived from other molecules, which are known to sensitize human immune system.

[0160] To exclude this possibility, homology analysis was performed with the peptide sequences as queries using BLAST algorithm

(<http://www.ncbi.nlm.nih.gov/blast/blast.cgi>). No significant sequence homology was revealed.

These results suggest that the sequences of SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 376, 379, 80, 100, 101, 110, 111, 387, 112, 394, 114, 116, 117, 121, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217, 223, 227, 228, 233, 254, 271, 272 or 288 are unique and thus possess a low risk of raising unintended immunologic response to any unrelated molecule.

[0161] **DISCUSSION**

Identification of new TAAs, particularly those that induce potent and specific anti-tumor immune responses, warrants further development of the clinical application of peptide vaccination strategies in various types of cancer (Boon T. et al., (1996) *J Exp Med* 183: 725-9.; van der Bruggen P et al., (1991) *Science* 254: 1643-7.; Brichard V et al., (1993) *J Exp Med* 178: 489-95.; Kawakami Y et al., (1994) *J Exp Med* 180: 347-52.; Shichijo S et al., (1998) *J Exp Med* 187:277-88.; Chen YT et al., (1997) *Proc.Natl.Acad. Sci. USA*, 94: 1914-8.; Harris CC., (1996) *J Natl Cancer Inst* 88:1442-5.; Butterfield LH et al., (1999) *Cancer Res* 59:3134-42.; Vissers JL et al., (1999) *Cancer Res* 59: 5554-9.; van der Burg SH et al., (1996) *J. Immunol* 156:3308-14.; Tanaka F et al., (1997) *Cancer Res* 57:4465-8.; Fujie T et al., (1999) *Int J Cancer* 80:169-72.; Kikuchi M et al., (1999) *Int J Cancer* 81 : 459-66.; Oiso M et al., (1999) *Int J Cancer* 81:387-94.).

[0162] cDNA microarray technologies can disclose comprehensive profiles of gene expression of malignant cells (Lin YM, et al., *Oncogene*. 2002 Jun 13;21:4120-8.; Kitahara O, et al., *Cancer Res*. 2001 May 1;61:3544-9.; Suzuki C, et al., *Cancer Res*. 2003 Nov 1;63:7038-41.; Ashida S, *Cancer Res*. 2004 Sep 1;64:5963-72.; Ochi K, et al., *Int J Oncol*. 2004 Mar;24(3):647-55.; Kaneta Y, et al., *Int J Oncol*. 2003 Sep;23:681-91.; Obama K, *Hepatology*. 2005 Jun;41:1339-48.; Kato T, et al., *Cancer Res*. 2005 Jul 1;65:5638-46.; Kitahara O, et al., *Neoplasia*. 2002 Jul-Aug;4:295-303.; Saito-Hisamimoto A et al., *DNA Res* 2002, 9: 35-45.) and, find utility in the identification of potential TAAs. Among the transcripts that are up-regulated in various cancers, novel human genes, termed CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10, were identified using these technologies.

[0163] As demonstrated above, CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10, are over-expressed in various cancers but show minimal expression in normal tissues. In addition, these genes have been shown to have a significant function related to cell proliferation. Thus, peptides derived from CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 can serve as TAA epitopes, which, in turn, can be used to induce significant and specific immune responses against cancer cells.

Thus, as CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and URLC10 are novel TAAs, vaccines using these epitope peptides find utility as immuno-therapeutics against various carcinomas or other disease expressing these molecules.

Industrial Applicability

[0164] The present invention identifies new TAAs, particularly those which induce potent and specific anti-tumor immune responses. Such TAAs warrants further development as peptide vaccines against diseases associated with the over-expression of CDH3, EPHA4, ECT2, HIG2, INHBB, KIF20A, KNTC2, TTK and/or ,URLC10 e.g. cancers.

All patents, patent applications, and publications cited herein are incorporated by reference.

[0165] While the invention has been described in detail and with reference to specific embodiments thereof, it is to be understood that the foregoing description is exemplary and explanatory in nature and is intended to illustrate the invention and its preferred embodiments. Through routine experimentation, one skilled in the art will readily recognize that various changes and modifications can be made therein without departing from the spirit and scope of the invention. Thus, the invention is intended to be defined not by the above description, but by the following claims and their equivalents.

CLAIMS

[Claim 1]

An isolated peptide of less than about 15 amino acids selected from the group consisting of peptides comprising the amino acid sequences of SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 80, 100, 101, 110, 111, 387, 112, 394, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217 or 223.

[Claim 2]

A peptide having cytotoxic T cell inducibility, wherein said peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 80, 100, 101, 110, 111, 387, 112, 394, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217 or 223, wherein 1, 2, or several amino acids are substituted, deleted, or added.

[Claim 3]

The peptide of claim 2, wherein the second amino acid from the N-terminus is phenylalanine, tyrosine, methionine, or tryptophan.

[Claim 4]

The peptide of claim 2 or 3, wherein the C-terminal amino acid is phenylalanine, leucine, isoleucine, tryptophan, or methionine.

[Claim 5]

An isolated peptide of less than about 15 amino acids selected from the group consisting of peptides comprising the amino acid sequences of SEQ ID NO: 376, 379, 114, 116, 117, 121, 227, 228, 233, 254, 271, 272 or 288.

[Claim 6]

A peptide having cytotoxic T cell inducibility, wherein said peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: SEQ ID NO: 376, 379, 114, 116, 117, 121, 227, 228, 233, 254, 271, 272 or 288, wherein 1, 2, or several amino acids are substituted, deleted, or added.

[Claim 7]

The peptide of claim 6, wherein the second amino acid from the N-terminus is leucine or methionine.

[Claim 8]

The peptide of claim 6 or 7, wherein the C-terminal amino acid is valine or leucine.

[Claim 9]

A pharmaceutical composition for treating or preventing a disease associated with over-expression of the genes of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and/or 17 said composition comprising one or more peptides of claim 1 to 8 or a polynucleotide encoding the peptides.

[Claim 10]

A pharmaceutical composition of claim 9, wherein the disease associating genes of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and/or 17 include cancers.

[Claim 11]

A pharmaceutical composition of claim 10, wherein the cancers include bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and

testicular tumor.

[Claim 12]

An exosome that presents on its surface a complex comprising a peptide of claim 1-4 and an HLA antigen.

[Claim 13]

The exosome of claim 12, wherein the HLA antigen is HLA-A24.

[Claim 14]

The exosome of claim 13, wherein the HLA antigen is HLA-A2402.

[Claim 15]

An exosome that presents on its surface a complex comprising a peptide of claim 5-8 and an HLA antigen.

[Claim 16]

The exosome of claim 15, wherein the HLA antigen is HLA-A2.

[Claim 17]

The exosome of claim 16, wherein the HLA antigen is HLA-A0201.

[Claim 18]

A method of inducing antigen-presenting cells having a high cytotoxic T cell inducibility comprising the step of contacting an antigen-presenting cell with a peptide of claim 1-8.

[Claim 19]

A method of inducing cytotoxic T cells by contacting a T cell with a peptide of claim 1-8.

[Claim 20]

A method of inducing antigen-presenting cells having high cytotoxic T cell inducibility, said method comprising the step of transferring a gene comprising a polynucleotide encoding a peptide of claim 1 to 8 to an antigen-presenting cell.

[Claim 21]

An isolated cytotoxic T cell, which is induced by contacting a T cell with a peptide of claim 1-8 or which is transduced with the nucleic acids encoding the TCR subunits polypeptides binding with a peptide of any one claims 1 to 8 in the context of HLA-A24 or HLA-A2.

[Claim 22]

An antigen-presenting cell, which comprises a complex formed between an HLA antigen and a peptide of claim 1 to 8.

[Claim 23]

An antigen-presenting cell, induced by the method of claim 18-20.

[Claim 24]

A vaccine for inhibiting proliferation of cells expressing genes of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and/or 17 wherein the vaccine comprises a peptide of claim 1-8 as the active ingredient.

[Claim 25]

The vaccine of claim 24, wherein the cells expressing genes of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and/or 17 include cancers.

[Claim 26]

The vaccine of claim 25, wherein the cancers include bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[Claim 27]

The vaccine of claim 26, formulated for administration to a subject whose HLA antigen is HLA-A24 or HLA-A2.

[Claim 28]

A method of treating or preventing a disease associated with over-expression of the genes of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and/or 17 in a subject comprising admin-

istering to said subject a vaccine comprising a peptide of claim 1-8, an immunologically active fragment thereof, or a polynucleotide encoding said peptide or immunologically active fragment.

[Claim 29]

The method of claim 28, wherein the disease associating genes of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, 15 and/or 17 include cancers.

[Claim 30]






A method of treating or preventing cancers of claim 29, wherein the cancer cells include bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophageal cancer, gastric cancer, diffused type gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma, SCLC, soft tissue tumor and testicular tumor.

[Claim 31]



A method of screening for a peptide which 1, 2, or several amino acids are substituted, wherein said peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: SEQ ID NO: 19, 22, 30, 34, 344, 358, 41, 44, 46, 48, 78, 80, 100, 101, 110, 111, 387, 112, 394, 395, 133, 135, 137, 426, 174, 178, 186, 194, 196, 202, 210, 213, 214, 217 or 223, said method comprising the steps of:

- (a) conforming no significant sequence homology to the entire sequence of 1, 2 or several amino acids substitute;
- (b) measuring the CTL inducibility of the candidate substitute peptide; and
- (c) selecting the peptide which CTL inducibility is same to or higher than the original peptide.


[Fig. 1-1]

a CDH3-A24-10-248**b CDH3-A24-10-332****c CDH3-A24-10-470****d CDH3-A24-9-513****e CDH3-A24-9-406**


[Fig. 1-2]

f CDH3-A24-10-807**g CDH3-A24-10-655**


[Fig. 2]




[Fig. 3]




[Fig. 4-1]




[Fig. 4-2]




[Fig. 4-3]




[Fig. 5-1]




[Fig. 5-2]




[Fig. 6-1]




[Fig. 6-2]




[Fig. 7-1]





[Fig. 7-2]




[Fig. 8-1]




[Fig. 8-2]

d TTK-A02-9-547

[Fig. 8-3]



[Fig. 9-1]



[Fig. 9-2]

Continuation of d

CTL line stimulated by URLC10-A02-10-211 recognized endogenously URLC10 expressed target cells with HLA-A02

