



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2344761 C 2012/08/21

(11)(21) **2 344 761**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2000/08/23
(87) Date publication PCT/PCT Publication Date: 2001/03/01
(45) Date de délivrance/Issue Date: 2012/08/21
(85) Entrée phase nationale/National Entry: 2001/03/27
(86) N° demande PCT/PCT Application No.: US 2000/023156
(87) N° publication PCT/PCT Publication No.: 2001/013731
(30) Priorité/Priority: 1999/08/23 (US60/150,319)

(51) Cl.Int./Int.Cl. *A01N 63/00* (2006.01),
A01N 63/02 (2006.01)
(72) Inventeurs/Inventors:
STOCKHOFF, BRIAN A., US;
CONLAN, CHRISTOPHER, US
(73) Propriétaire/Owner:
MYCOGEN CORPORATION, US
(74) Agent: MACRAE & CO.

(54) Titre : PROCEDES DESTINES A LA LUTTE CONTRE LE VER-GRIS
(54) Title: METHODS OF CONTROLLING CUTWORM PESTS

(57) **Abrégé/Abstract:**

The subject invention concerns methods, and materials, useful for controlling cutworm pests. In one preferred embodiment, Cry1Fa proteins are used for controlling black cutworm.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 March 2001 (01.03.2001)

PCT

(10) International Publication Number
WO 01/13731 A1

(51) International Patent Classification⁷: A01N 63/00, 63/02

(21) International Application Number: PCT/US00/23156

(22) International Filing Date: 23 August 2000 (23.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/150,319 23 August 1999 (23.08.1999) US

(71) Applicant: MYCOGEN CORPORATION [US/US];
5501 Oberlin Drive, San Diego, CA 92121 (US).

(72) Inventors: STOCKHOFF, Brian, A.; 11771 Ramsdell
Court, San Diego, CA 92131 (US). CONLAN, Christo-
pher; 11513 Hadar Drive, San Diego, CA 92126 (US).

(74) Agents: SANDERS, Jay, M. et al.; Saliwanchik, Lloyd &
Saliwanchik, Suite A-1, 2421 N.W. 41st Street, Gainesville,
FL 32606-6669 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS OF CONTROLLING CUTWORM PESTS

(57) Abstract: The subject invention concerns methods, and materials, useful for controlling cutworm pests. In one preferred embodiment, Cry1Fa proteins are used for controlling black cutworm.

WO 01/13731 A1

DESCRIPTIONMETHODS OF CONTROLLING CUTWORM PESTS

5

Background of the Invention

10

Insects and other pests cost farmers billions of dollars annually in crop losses and in the expense of keeping these pests under control. The losses caused by insect pests in agricultural production environments include decrease in crop yield and reduced crop quality. Increased harvesting costs are also a result of insect infestations.

15

Insect problems have been partially addressed by cultivation methods, such as crop rotation, and by fertilizing with high levels of phosphate to stimulate the growth of strong root systems by the plants. However, crop rotation can be disrupted by, for example, an emerging two-year diapause (or overwintering) trait of northern corn rootworms. Chemical insecticides are relied upon most heavily to guarantee the desired level of control.

20

Chemical pesticides have provided an effective method of pest control. However, the public has become concerned about the amount of residual chemicals which might be found in food, ground water, and elsewhere in the environment. Therefore, synthetic chemical pesticides are being increasingly scrutinized for their potential adverse environmental consequences. Some synthetic chemical pesticides can poison the soil and underlying aquifers, pollute surface waters as a result of runoff, and destroy non-target life forms. Some synthetic chemical control agents have the further disadvantage of presenting public safety hazards when they are applied in areas where pets, farm animals, or children may come into contact with them. They can also pose health hazards to the people applying them, especially if the proper application techniques are not followed. Regulatory agencies around the world are restricting and/or banning the uses of many synthetic pesticides, particularly those that are persistent in the environment and that enter the food chain. Stringent new restrictions on the use of pesticides and the elimination of some effective pesticides from the market place could limit economical and effective options for controlling costly pests.

25

30

Because of the problems associated with the use of many synthetic chemical pesticides, there exists a clear need to limit the use of these agents and to identify alternative control agents.

The replacement of synthetic chemical pesticides, or the combination of these agents with biological pesticides, could reduce the levels of toxic chemicals in the environment.

A biological pesticidal agent that is being used with increasing popularity is the soil microbe *Bacillus thuringiensis* (*B.t.*). The soil microbe *Bacillus thuringiensis* (*B.t.*) is a Gram-positive, spore-forming bacterium. Most strains of *B.t.* do not exhibit pesticidal activity. Some *B.t.* strains produce pesticidal parasporal protein inclusions. These protein inclusions often appear microscopically as distinctively shaped crystals. The shape and type of the crystal inclusions can be used to characterize *B.t.* strains. The “ δ -endotoxins” present in the crystalline inclusions, which typically have specific pesticidal activity, are different from exotoxins, which have a non-specific host range.

Commercial use of *B.t.* pesticides was originally restricted to a narrow range of lepidopteran (caterpillar) pests. For example, preparations of the spores and crystals of *B. thuringiensis* subsp. *kurstaki* have been used for many years as commercial insecticides for lepidopteran pests. More recently, however, investigators have discovered *B.t.* pesticides with specificities for a much broader range of pests. For example, *B.t. israelensis* and *morrisoni* have been used commercially to control insects of the orders Diptera and Coleoptera, respectively (Gaertner, F.H. [1989] “Cellular Delivery Systems for Insecticidal Proteins: Living and Non-Living Microorganisms,” in *Controlled Delivery of Crop Protection Agents*, R.M. Wilkins, ed., Taylor and Francis, New York and London, 1990, pp. 245-255).

New subspecies of *B.t.* have now been identified, and genes responsible for active δ -endotoxin proteins have been isolated and sequenced (Höfte, H., H.R. Whiteley [1989] *Microbiological Reviews* 52(2):242-255). Höfte and Whiteley classified *B.t.* crystal protein genes into four major classes. The classes were *cryI* (Lepidoptera-specific), *cryII* (Lepidoptera- and Diptera-specific), *cryIII* (Coleoptera-specific), and *cryIV* (Diptera-specific). The discovery of strains specifically toxic to other pests has been reported (Feitelson, J.S., J. Payne, L. Kim [1992] *Bio/Technology* 10:271-275). For example, the designations CryV and CryVI have been proposed for two new groups of nematode-active toxins.

Many *Bacillus thuringiensis* δ -endotoxin proteins are composed of two functional segments. These “full-length” proteins are comprised of a protease-resistant core toxin that corresponds to about the first half (the N-terminal portion) of the protein molecule. The three-dimensional structure of a core segment of a CryIIIA *B.t.* δ -endotoxin is known, and it was proposed that all related toxins have that same overall structure (Li, J., J. Carroll, D.J. Ellar [1991] *Nature* 353:815-821). The second half (the C-terminal portion) of the molecule is often referred to as the “protoxin segment.” The protoxin segment is believed to participate in toxin

crystal formation (Arvidson, H., P.E. Dunn, S. strand, A.I. Aronson [1989] *Molecular Microbiology* 3:1533-1534; Choma, C.T., W.K. Surewicz, P.R. Cary, M. Pozsgay, T. Raynor, H. Kaplan [1990] *Eur. J. Biochem* 189:523-527). The full 130 kDa toxin molecule is typically processed to the resistant core segment by proteases in the insect gut. The protoxin segment may thus convey a partial insect specificity for the toxin by limiting the accessibility of the core to the insect by reducing the protease processing of the toxin molecule (Haider, M.Z., B.H. Klawns, D.J. Ellar [1986] *Eur. J. Biochem* 156:531-540) or by reducing toxin solubility (Aronson, A.I., E.S. Han, W. McGaughey, D. Johnson [1991] *Appl. Environ. Microbiol.* 57:981-986).

The 1989 nomenclature and classification scheme of Höfte and Whiteley was based on both the deduced amino acid sequence and the host range of the toxin. That system was adapted to cover 14 different types of toxin genes which were divided into five major classes. A revised nomenclature scheme has been proposed which is based solely on amino acid identity (Crickmore *et al.* [1996] Society for Invertebrate Pathology, 29th Annual Meeting, IIIrd International Colloquium on *Bacillus thuringiensis*, University of Cordoba, Cordoba, Spain, September 1-6, 1996, abstract). The mnemonic “cry” has been retained for all of the toxin genes except cytA and cytB, which remain a separate class. Roman numerals have been exchanged for Arabic numerals in the primary rank, and the parentheses in the tertiary rank have been removed. Current boundaries represent approximately 95% (tertiary rank), 75% (secondary rank), and 48% (primary rank) sequence identity. Many of the original names have been retained, although a number have been reclassified. *See also* “Revisions of the Nomenclature for the *Bacillus thuringiensis* Pesticidal Crystal Proteins.” N. Crickmore, D.R. Zeigler, J. Feitelson, E. Schnepf, J. Van Rie, D. Lereclus, J. Baum, and D.H. Dean, *Microbiology and Molecular Biology Reviews* (1998) Vol. 62:807-813. That system uses the freely available software applications CLUSTAL™W and PHYLIP™. The NEIGHBOR™ application within the PHYLIP package uses an arithmetic averages (UPGMA) algorithm.

Isolating novel B.t. isolates and their toxin genes, as well as determining the precise pesticidal range of the toxins, has been a slow empirical process. As a result of extensive research and resource investment, patents have issued for new B.t. isolates, toxins, and genes, and for new uses of known B.t. toxins and isolates. *See Feitelson et al., supra*, for a review. However, the discovery of new B.t. isolates and new uses of known B.t. isolates and toxins remains an empirical, unpredictable art.

Smulevitch *et al.* ([1991] *FEBS Lett.* 293:25-26), Gleave *et al.* ([1991] *JGM* 138:55-62), and Shevelev *et al.* ([1993] *FEBS Lett.* 336:79-82) describe the characterization of *Cry9* toxins active against lepidopterans. Lambert *et al.* (Lambert, B., L. Buysse, C. Decock, S. Jansens, C. Piens, B. Saey, J. Seurinck, K. van Audenhove, J. Van Rie, A. Van Vliet, M. Peferoen [1996] *Appl. Environ. Microbiol* 62(1):80-86), and Published PCT applications WO 94/05771 and WO 94/24264 also describe *B.t.* isolates and *Cry9C* toxins active against lepidopteran pests.

U.S. Patent Nos. 5,126,133; 5,188,960; 5,246,852; and 5,691,308 disclose a *Cry1Fa* toxin and gene (81IA) from *B.t.* isolate PS81I. U.S. Patent Nos. 5,527,883; 5,508,264; 5,827,514; and 5,840,554 relate to *CryIF* chimeric toxins. WO 99/24581 discloses various plant-optimized *cry1F* genes. U.S. Patent No. 5,686,069 discloses a *Cry1Fb* toxin and gene from *B.t.* isolate PS91C2.

With the use of genetic engineering techniques, various approaches for delivering *B.t.* toxins to agricultural environments are under development. The cloning and expression of a *B.t.* crystal protein gene in *Escherichia coli* was described in the published literature more than 15 years ago (Schnepf, H.E., H.R. Whiteley [1981] *Proc. Natl. Acad. Sci. USA* 78:2893-2897.). U.S. Patent No. 4,448,885 and U.S. Patent No. 4,467,036 both disclose the expression of *B.t.* crystal protein in *E. coli*. Recombinant DNA-based *B.t.* products have been produced and approved for use, including the use of plants genetically engineered with *B.t.* genes for insect resistance and the use of stabilized, microbial cells as delivery vehicles of *B.t.* proteins. Various improvements have been achieved by modifying *B.t.* toxins and/or their genes. For example, U.S. Patent Nos. 5,380,831 and 5,567,862 relate to the production of synthetic insecticidal crystal protein genes having improved expression in plants. Thus, isolated *B.t.* endotoxin genes are becoming commercially valuable.

Obstacles to the successful agricultural use of *B.t.* toxins include the development of resistance to *B.t.* toxins by insects. In addition, certain insects can be refractory to the effects of *B.t.* The latter includes insects such as boll weevils and cutworms, which heretofore have demonstrated no apparent sensitivity to most *B.t.* δ -endotoxins.

Cutworms progress through several instars as larvae. Although seedling cutting by later instar larvae produces the most obvious damage and economic loss, leaf feeding commonly results in yield loss in crops such as maize. Upon reaching later instars (third to fourth, for example), larvae begin to cut plants and plant parts, especially seedlings. Because of the shift in feeding behavior, economically damaging populations may build up unexpectedly with few early warning signs. Large cutworms can destroy several seedlings per day, and a heavy

infestation can remove entire stands of crops. Cutworms' nocturnal habit and behavior of burrowing into the ground also makes detection and treatment problematic.

The black cutworm (*Agrotis ipsilon* (Hufnagel); Lepidoptera: Noctuidae) is a serious pest of many crops including maize, cotton, cole crops (*Brassica*, broccoli, cabbages, Chinese cabbages), and turf. Secondary host plants include beetroots, *Capsicum* (peppers), chickpeas, faba beans, lettuces, lucerne, onions, potatoes, radishes, rape (canola), rice, soybeans, strawberries, sugarbeet, tobacco, tomatoes, and forest trees. In North America, pests of the genus *Agrotis* feed on clover, corn, tobacco, hemp, onion, strawberries, blackberries, raspberries, alfalfa, barley, beans, cabbage, oats, peas, potatoes, sweetpotatoes, tomato, garden flowers, grasses, lucerne, maize, asparagus, grapes, almost any kind of leaf, weeds, and many other crops and garden plants. Other cutworms in the Tribe Agrotini are pests, in particular those in the genus *Feltia* (e.g., *F. jaculifera* (Guenée); equivalent to *ducens subgothica*) and *Euxoa* (e.g., *E. messoria* (Harris), *E. scandens* (Riley), *E. auxiliaris* Smith, *E. detersa* (Walker), *E. tessellata* (Harris), *E. ochrogaster* (Guenée)). Cutworms such as *Peridroma saucia* can also be significant pests. Citrus plants, for example, can also be the target of cutworm attacks (the "citrus cutworm" *Xylomyges curialis* is an example).

Cutworms are also pests outside North America, and the more economically significant pests attack chickpeas, wheat, vegetables, sugarbeet, lucerne, maize, potatoes, turnips, rape, lettuces, strawberries, loganberries, flax, cotton, soybeans, tobacco, beetroots, Chinese cabbages, tomatoes, aubergines, sugarcane, pastures, cabbages, groundnuts, *Cucurbita*, turnips, sunflowers, *Brassica*, onions, leeks, celery, sesame, asparagus, rhubarb, chicory, greenhouse crops, and spinach. The black cutworm *A. ipsilon* occurs as a pest outside North America, including Central America, Europe, Asia, Australia, Africa, India, Taiwan, Mexico, Egypt, and New Zealand.

The main cutworm species in Argentina are *Agrotis malefida*, *Porosagrotis gypaetiana*, and *Agrotis ipsilon* (also spelled *ypsilon*). These cutworms attack corn, soybean, and sunflower cultivars, for example. These insects can seriously reduce the population of seedlings, and in cases of severe attacks, can totally destroy entire plots.

Cultural controls for *A. ipsilon* such as peripheral weed control can help prevent heavy infestations; however, such methods are not always feasible or effective. Infestations are very sporadic, and applying an insecticide prior to planting or at planting has not been effective in the past. Some baits are available for control of cutworms in crops. To protect turfgrass such as creeping bentgrass, chemical insecticides have been employed. Use of chemical pesticides is a particular concern in turf-covered areas (e.g., golf greens, athletic fields, parks and other

recreational areas, professional landscaping, and home lawns) because of the close contact the public has with these areas. Natural products (*e.g.*, nematodes and azadirachtin) generally perform poorly.

To date, *Bacillus thuringiensis* products for control black cutworm have not been widely used to due to their heretofore lack of sufficient effectiveness. The rapid feeding and burrowing behaviors of the cutworm have made them particularly difficult to control with current biologically based pest solutions. For example, *Cry1A(b)* toxins are basically ineffective against cutworms.

Brief Summary of the Invention

The subject invention relates to the surprising discovery that *Cry1F* proteins are active against cutworms such as the black cutworm (*Agrotis ipsilon*). Thus, the subject invention provides methods for controlling these pests wherein said method comprises contacting said pest with a pesticidal amount of a *Bacillus thuringiensis* toxin comprising at least a pesticidal portion of a *Cry1F* toxin. Therefore, the use of full-length, truncated, and chimeric *Cry1F* proteins and genes is included in the subject invention. In preferred embodiments, the *Cry1F* toxin is a *Cry1Fa* toxin. Wild-type and synthetic *Cry1F* proteins can be used according to the subject invention. Thus, the use of polynucleotides (and/or their complements, preferably their full complements) that hybridize with known *cry1F* genes, preferably the core-toxin encoding portions, are included in this invention. Plant-optimized polynucleotides are used in preferred embodiments.

The subject invention includes the use of transgenic hosts including plant hosts such as corn, cotton, and sunflower. In a preferred embodiment, the subject invention concerns plants cells transformed with at least one polynucleotide sequence of the subject invention such that the transformed plant cells express pesticidal proteins in tissues consumed by the target pests. Such transformation of plants can be accomplished using techniques well known to those skilled in the art and would typically involve modification of the gene to optimize expression of the toxin in plants.

Brief Description of the Drawings

Figure 1 shows mass of intact wheat seedlings and relative damage ratings 24 hours after infestation with 4th instar black cutworm larvae and after various treatments.

Figure 2 shows the fresh weights of wheat seedlings infested with 3rd instar black cutworm larvae 48 hours after infestation and after various treatments.

Figures 3 and 4 illustrate the results discussed in Example 4.

Brief Description of the Sequences

SEQ ID NO. 1 is a polynucleotide sequence for a full-length, plant-optimized *cryIF/cryIA(b)* hybrid gene designated 1F1AB-PO.

5 SEQ ID NO. 2 is an amino acid sequence for a full-length, plant-optimized CryIF/CryIA(b) chimeric toxin. The 1F1AB-PO gene encodes this toxin.

SEQ ID NO. 3 is a polynucleotide sequence for a truncated, plant-optimized *cryIF* gene designated 1F-T-PO.

10 SEQ ID NO. 4 is an amino acid sequence for a truncated, plant-optimized CryIF toxin. The genes designated 1F-T-PO, 1F-7G-PO, and 1F-7Z-PO encode this toxin.

SEQ ID NO. 5 is the native polynucleotide sequence of the wild-type, full length *B.t.* toxin gene designated 81IA (*cryIFa*).

SEQ ID NO. 6 is the amino acid sequence of the full length, wild-type *B.t.* toxin designated 81IA (CryIFa).

15 SEQ ID NO. 7 is a polynucleotide sequence for a gene designated 1F-7G-PO, which is optimized for expression in cotton.

SEQ ID NO. 8 is a polynucleotide sequence for a gene designated 1F-7Z-PO, which is optimized for expression in maize.

20

Detailed Disclosure of the Invention

The subject invention relates to the surprising discovery that *Cry1F* proteins are active against cutworms such as the black cutworm (*Agrotis ipsilon*). Particularly surprising are the findings that 3rd instar, and greater, larvae are controlled by the toxins and genes according to the subject invention. Controlling first and second instar cutworm larvae is much less important than controlling later instars. While it was known that *CryI* toxins are active against lepidopterans generally, cutworms were thought to be recalcitrant to *B.t.* toxins generally, as discussed in more detail above in the Background section.

25

The subject invention includes the use of recombinant hosts and provides plant-optimized polynucleotide sequences. These polynucleotide sequences include plant-optimized genes designated 1F1AB-PO, 1F-T-PO, 1F-7G-PO, and 1F-7Z-PO. These genes are disclosed in WO 99/24581. Preferred plant hosts include corn, soybeans, cotton, wheat, canola, and sunflower.

30

In some embodiments of the subject invention, genes encode a *CryIF* toxin that is truncated compared to the full length *CryIF* toxin. The truncated toxins of the subject invention are typically missing all or a portion of the protoxin segment. Also, the truncated genes of the subject invention can be used for the production of chimeric genes and proteins. One example is the plant-optimized gene comprising a *cryIF* portion and a *cryIA(b)* portion, wherein the hybrid gene encodes a chimeric toxin. Preferred chimeric genes and toxins are disclosed in U.S. Patent Nos. 5,527,883 and 5,840,554. Other chimeric genes and toxins, which can be used according to the subject invention, are disclosed in U.S. Patent Nos. 5,508,264 and 5,827,514. In a preferred embodiment, the *CryIF* portion of the chimeric toxin is itself pesticidal. Fusion toxins can also be used according to the subject invention.

In a preferred embodiment, the subject invention concerns plants cells transformed with at least one polynucleotide sequence such that the transformed plant cells express pesticidal toxins in tissues consumed by the target pests which thereby contact the pesticidal protein. Such transformation of plants can be accomplished using techniques well known to those skilled in the art and would typically involve modification of the gene to optimize expression of the toxin in plants.

It should be apparent to a person skilled in this art that, given the sequences of the genes as set forth herein, the genes of the subject invention can be obtained through several means. In preferred embodiments, the subject genes may be constructed synthetically by using a gene synthesizer, for example. The specific genes exemplified herein can also be obtained by modifying, according to the teachings of the subject invention, certain wild-type genes (for example, by point-mutation techniques) from certain isolates deposited at a culture depository as discussed below. For example, a wild-type *cryIF* gene can be obtained from *B.t.* isolate PS81I. Likewise, the *cryIA(b)* portions of the hybrid genes of the subject invention can be produced synthetically or can be derived by modifying wild-type genes. *CryIA(b)* toxins and genes have been described in, for example, Höfte *et al.* (1986) *Eur. J. Biochem.* 161:273; Geiser *et al.* (1986) *Gene* 48:109; and Haider *et al.* (1988) *Nucleic Acids Res.* 16:10927. Clones and additional wild-type isolates are discussed in more detail, above, in the section entitled "Background of the Invention" and in the list, below.

Cultures discussed in this application have been deposited, in accordance with the Budapest Treaty, in the Agricultural Research Service Patent Culture Collection (NRRL), Northern Regional Research Center, 1815 North University Street, Peoria, Illinois 61604, USA. The deposited strains listed below are disclosed in the patent references as discussed above in the section entitled "Background of the Invention."

Subculture	Accession Number	Deposit Date
<i>B.t.</i> PS81I	NRRL B-18484	April 19, 1989
<i>E. coli</i> (NM522) (pMYC1603) (81IA)	NRRL B-18517	June 30, 1989

It should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

Genes and toxins. The polynucleotides of the subject invention can be used to form complete "genes" to encode proteins or peptides in a desired host cell. For example, as the skilled artisan would readily recognize, some of the polynucleotides in the attached sequence listing are shown without stop codons. Also, the subject polynucleotides can be appropriately placed under the control of a promoter in a host of interest, as is readily known in the art.

As the skilled artisan would readily recognize, DNA typically exists in a double-stranded form. In this arrangement, one strand is complementary to the other strand and vice versa. As DNA is replicated in a plant (for example) additional, complementary strands of DNA are produced. Thus, the subject invention includes the use of the exemplified polynucleotides shown in the attached sequence listing and the complementary strands thereof. The "coding strand" is often used in the art to refer to the strand that binds with the anti-sense strand. In order to express a protein *in vivo*, a strand of DNA is typically transcribed into a complementary strand of mRNA which is used as the template for translating the protein. The mRNA is actually transcribed from the "anti-sense" strand of DNA. The "sense" or "coding" strand has a series of codons (a codon is three nucleotides that can be read three-at-a-time to yield a particular amino acid) that can be read as an open reading frame (ORF) to form a protein or peptide of interest. RNA and PNA (peptide nucleic acids) that are functionally equivalent to the exemplified DNA are included in the subject invention.

Genes and toxins of the subject invention can be identified, obtained, and characterized by using oligonucleotide probes, for example. Probes are detectable nucleotide sequences. The specifically exemplified polynucleotides of the subject invention, including portions thereof that are sufficient to encode an active toxin, can, themselves, be used as probes. Probes may be DNA, RNA, or PNA (peptide nucleic acid). These sequences may be detectable by virtue of an appropriate label including or may be made inherently fluorescent, for example, as described in International Application No. WO93/16094. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong bond between the two molecules, it can be reasonably assumed that the probe and sample have substantial homology. Preferably,

hybridization is conducted under stringent conditions by techniques well-known in the art, as described, for example, in Keller, G.H., M.M. Manak (1987) *DNA Probes*, Stockton Press, New York, NY., pp. 169-170. For example, as stated therein, hybridization conditions of high stringency can be achieved by first washing with 2x SSC (Standard Saline Citrate)/0.1% SDS (Sodium Dodecyl Sulfate) for 15 minutes at room temperature. Two washes are typically performed. Higher stringency can be achieved by lowering the salt concentration and/or by raising the temperature. For example, the above hybridization step can be followed by washing with 0.1x SSC/0.1% SDS for 15 minutes at room temperature, which in turn can be followed by washing with 0.1x SSC/0.1% SDS for 30 minutes at 55°C. The temperatures used for these steps can also be used with other protocols discussed herein (where SSPE is used in place of SSC, for example) as would be known in the art. The 2x SSC/0.1% SDS can be prepared by adding 50 ml of 20x SSC and 5 ml of 10% SDS to 445 ml of water. 20x SSC can be prepared by combining NaCl (175.3 g / 0.150 M), sodium citrate (88.2 g / 0.015 M), and water to 1 liter, followed by adjusting pH to 7.0 with 10 N NaOH. 10% SDS can be prepared by dissolving 10 g of SDS in 50 ml of autoclaved water, diluting to 100 ml, and aliquotting.

Detection of the probe provides a means for determining in a known manner whether hybridization has occurred. Such a probe analysis provides a rapid method for identifying toxin-encoding genes of the subject invention. The nucleotide segments which are used as probes according to the invention can be synthesized using a DNA synthesizer and standard procedures.

As used herein "stringent" conditions for hybridization refers to conditions which achieve the same, or about the same, degree of specificity of hybridization as the conditions employed by the current applicants. Specifically, hybridization of immobilized DNA on Southern blots with ³²P-labeled gene-specific probes was performed by standard methods (Maniatis, T., E.F. Fritsch, J. Sambrook [1982] *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.). In general, hybridization and subsequent washes were carried out under stringent conditions that allowed for detection of target sequences. For double-stranded DNA gene probes, hybridization was carried out overnight at 20-25° C below the melting temperature (T_m) of the DNA hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula (Beltz, G.A., K.A. Jacobs, T.H. Eickbush, P.T. Cherbas, and F.C. Kafatos [1983] *Methods of Enzymology*, R. Wu, L. Grossman and K. Moldave [eds.] Academic Press, New York 100:266-285).

$$T_m = 81.5^{\circ} \text{C} + 16.6 \log[\text{Na}^+] + 0.41(\% \text{G} + \text{C}) - 0.61(\% \text{formamide}) - 600 / \text{length of duplex in base pairs.}$$

Washes are typically carried out as follows:

- (1) Twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at $T_m - 20^\circ\text{C}$ for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

For oligonucleotide probes, hybridization was carried out overnight at $10-20^\circ\text{C}$ below the melting temperature (T_m) of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. T_m for oligonucleotide probes was determined by the following formula:

$$T_m (^{\circ}\text{C}) = 2(\text{number T/A base pairs}) + 4(\text{number G/C base pairs})$$

(Suggs, S.V., T. Miyake, E.H. Kawashime, M.J. Johnson, K. Itakura, and R.B. Wallace [1981] *ICN-UCLA Symp. Dev. Biol. Using Purified Genes*, D.D. Brown [ed.], Academic Press, New York, 23:683-693).

Washes were typically carried out as follows:

- (1) Twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

Modification of genes and toxins. The genes and toxins useful according to the subject invention include not only the specifically exemplified sequences, but also include portions and/or fragments (including internal and/or terminal deletions compared to the full-length proteins), variants, mutants, substitutes (proteins having substituted amino acids), chimerics, and fusion proteins which retain the characteristic pesticidal activity of the proteins specifically exemplified herein. As used herein, the terms "variants" or "variations" of genes refer to nucleotide sequences which encode the same toxins or which encode equivalent toxins having pesticidal activity. As used herein, the term "equivalent toxins" refers to toxins having the same or essentially the same biological activity against the target pests as the claimed toxins.

Genes can be modified, and variations of genes may be readily constructed, using standard techniques. For example, techniques for making point mutations are well known in the art. Also, U.S. Patent No. 5,605,793, for example, describes methods for generating additional molecular diversity by using DNA reassembly after random fragmentation. Fragments of full-length genes can be made using commercially available endonucleases, and exonucleases can be used according to standard procedures. For example, enzymes such as *Bal31* or site-directed mutagenesis can be used to systematically cut off nucleotides from the ends of these genes.

Also, genes which encode active fragments may be obtained using a variety of restriction enzymes. Proteases may be used to directly obtain active fragments of these toxins.

Equivalent toxins and/or genes encoding these equivalent toxins can be derived from *B.t.* isolates and/or DNA libraries using the teachings provided herein. There are a number of methods for obtaining the pesticidal toxins of the instant invention. For example, antibodies to the pesticidal toxins disclosed and claimed herein can be used to identify and isolate other toxins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the toxins which are most constant and most distinct from other *B.t.* toxins. These antibodies can then be used to specifically identify equivalent toxins with the characteristic activity by immunoprecipitation, enzyme linked immunosorbent assay (ELISA), or western blotting. Antibodies to the toxins disclosed herein, or to equivalent toxins, or fragments of these toxins, can readily be prepared using standard procedures in this art. The genes which encode these toxins can then be obtained from the microorganism.

Because of the redundancy of the genetic code, a variety of different DNA sequences can encode the same amino acid sequences. It is well within the skill of a person trained in the art to create these alternative DNA sequences encoding the same, or essentially the same, toxins. These variant DNA sequences are within the scope of the subject invention. As used herein, reference to "essentially the same" sequence refers to sequences which have amino acid substitutions, deletions, additions, or insertions which do not materially affect pesticidal activity. Fragments retaining pesticidal activity are also included in this definition.

Equivalent toxins will have amino acid similarity (and/or homology) with an exemplified toxin. The amino acid similarity/identity will typically be greater than 60%, preferably greater than 75%, more preferably greater than 80%, even more preferably greater than 90%, and can be greater than 95%. Preferred polynucleotides and proteins of the subject invention can also be defined in terms of more particular identity and/or similarity ranges. For example, the identity and/or similarity can be 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% as compared to a sequence exemplified herein. Unless otherwise specified, as used herein percent sequence identity and/or similarity of two nucleic acids is determined using the algorithm of Karlin and Altschul (1990), *Proc. Natl. Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul (1993), *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLASTTM and XBLASTTM programs of Altschul *et al.* (1990), *J. Mol. Biol.* 215:402-410. BLAST nucleotide searches are performed with the NBLAST program, score = 100, wordlength = 12, to obtain nucleotide sequences with

the desired percent sequence identity. To obtain gapped alignments for comparison purposes, Gapped BLAST is used as described in Altschul *et al.* (1997), *Nucl. Acids Res.* 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (NBLAST and XBLAST) are used. The identity scores can also be calculated using the methods and algorithms of Crickmore *et al.* as described in the Background section, above.

The amino acid homology will be highest in critical regions of the toxin which account for biological activity or are involved in the determination of three-dimensional configuration which ultimately is responsible for the biological activity. In this regard, certain amino acid substitutions are acceptable and can be expected if these substitutions are in regions which are not critical to activity or are conservative amino acid substitutions which do not affect the three-dimensional configuration of the molecule. For example, amino acids may be placed in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type fall within the scope of the subject invention so long as the substitution does not materially alter the biological activity of the compound. Table 1 provides a listing of examples of amino acids belonging to each class.

Table 1.

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gin
Acidic	Asp, Glu
Basic	Lys, Arg, His

In some instances, non-conservative substitutions can also be made. The critical factor is that these substitutions must not significantly detract from the ability of plants to express the subject DNA sequences or from the biological activity of the toxin.

As used herein, reference to "isolated" polynucleotides and / or "purified" toxins refers to these molecules when they are not associated with the other molecules with which they would be found in nature; this would include their use in plants. Thus, reference to "isolated" and/or "purified" signifies the involvement of the "hand of man" as described herein.

While the subject invention provides specific embodiments of synthetic genes, other genes that are functionally equivalent to the genes exemplified herein can also be used to transform hosts, preferably plant hosts. Additional guidance for the production of synthetic genes can be found in, for example, U.S. Patent No. 5,380,831.

5 Recombinant hosts. The toxin-encoding genes of the subject invention can be introduced into a wide variety of microbial or plant hosts. In preferred embodiments, expression of the toxin gene results, directly or indirectly, in the intracellular production and maintenance of the pesticide. When transgenic/recombinant/transformed host cells are ingested by the pest, the pests will ingest the toxin. This is the preferred manner in which to cause contact of the pest
10 with the toxin. The result is a control (killing or making sick) of the pest.

In some embodiments of the subject invention, transformed microbial hosts can be used in preliminary steps for preparing precursors, for example, that will eventually be used to transform, in preferred embodiments, plant cells and plants so that they express the toxins encoded by the genes of the subject invention. Microbes transformed and used in this manner
15 are within the scope of the subject invention. Recombinant microbes may be, for example, *B.t.*, *E. coli*, or *Pseudomonas*.

The production of various recombinant organisms can be made by those skilled in the art using standard techniques. Materials necessary for these transformations are disclosed herein or are otherwise readily available to the skilled artisan. A wide variety of methods are available
20 for introducing a *B.t.* gene encoding a toxin into the target host under conditions which allow for stable maintenance and expression of the gene. These methods are well known to those skilled in the art and are described, for example, in United States Patent No. 5,135,867.

As contemplated herein, the following:

25 The toxin-encoding genes of the subject invention can be introduced, via a suitable vector, into a wide variety of microbial and plant hosts. There are many crops of interest, such as corn, wheat, rice, cotton, soybeans, and sunflowers. Certain genes of the subject invention are particularly well suited for providing stable maintenance and expression, in the transformed plant, of the gene expressing the polypeptide pesticide, and, desirably, provide for improved protection of the pesticide from environmental degradation and inactivation. Expression of the
30 toxin gene results, directly or indirectly, in the intracellular production and maintenance of the pesticide proteins. Thus, the target pest can contact the pesticidal proteins by ingesting plant tissue containing the pesticidal proteins, which are toxic to the pest. The result is control of the pest. Alternatively, suitable microbial hosts, *e.g.*, *Pseudomonas fluorescens*, can be applied to the site of the pest, where some of which can proliferate, and are ingested by the target pests.

The microbe hosting the toxin gene can be treated under conditions that prolong the activity of the toxin and stabilize the cell. The treated cell, which retains the toxic activity, then can be applied to the environment of the target pest.

Where the *B.t.* toxin gene is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, certain host microbes should be used. Microorganism hosts are selected which are known to occupy the "phytosphere" (phylloplane, phyllosphere, rhizosphere, and/or rhizoplane) of one or more crops of interest. These microorganisms are selected so as to be capable of successfully competing in the particular environment (crop and other insect habitats) with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing the polypeptide pesticide, and, desirably, provide for improved protection of the pesticide from environmental degradation and inactivation.

A large number of microorganisms are known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, *e.g.*, genera *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, *e.g.*, genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Serratia marcescens*, *Acetobacter xylinum*, *Agrobacterium tumefaciens*, *Rhodopseudomonas spheroides*, *Xanthomonas campestris*, *Rhizobium melioli*, *Alcaligenes entrophus*, and *Azotobacter vinlandii*; and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces roseus*, *S. odoratus*, *Kluyveromyces veronae*, and *Aureobasidium pollulans*. Of particular interest are the pigmented microorganisms.

A wide variety of ways are available for introducing a *B.t.* gene encoding a toxin into the target host under conditions which allow for stable maintenance and expression of the gene. These methods are well known to those skilled in the art and are described, for example, in United States Patent No. 5,135,867, *entomopathogenic bacteria*, issued to the applicant.

Treatment of cells. As mentioned above, *B.t.* or recombinant cells expressing a *B.t.* toxin can be treated to prolong the toxin activity and stabilize the cell. The pesticide microcapsule that is formed comprises the *B.t.* toxin within a cellular structure that has been

stabilized and will protect the toxin when the microcapsule is applied to the environment of the target pest. Suitable host cells may include either prokaryotes or eukaryotes, normally being limited to those cells which do not produce substances toxic to higher organisms, such as mammals. However, organisms which produce substances toxic to higher organisms could be used, where the toxic substances are unstable or the level of application sufficiently low as to avoid any possibility of toxicity to a mammalian host. As hosts, of particular interest will be the prokaryotes and the lower eukaryotes, such as fungi.

The cell will usually be intact and be substantially in the proliferative form when treated, rather than in a spore form, although in some instances spores may be employed.

Treatment of the microbial cell, *e.g.*, a microbe containing the *B.t.* toxin gene, can be by chemical or physical means, or by a combination of chemical and/or physical means, so long as the technique does not deleteriously affect the properties of the toxin, nor diminish the cellular capability of protecting the toxin. Examples of chemical reagents are halogenating agents, particularly halogens of atomic no. 17-80. More particularly, iodine can be used under mild conditions and for sufficient time to achieve the desired results. Other suitable techniques include treatment with aldehydes, such as glutaraldehyde; anti-infectives, such as zephiran chloride and cetylpyridinium chloride; alcohols, such as isopropyl and ethanol; various histologic fixatives, such as Lugol iodine, Bouin's fixative, various acids and Helly's fixative (See: Humason, Gretchen L., *Animal Tissue Techniques*, W.H. Freeman and Company, 1967); or a combination of physical (heat) and chemical agents that preserve and prolong the activity of the toxin produced in the cell when the cell is administered to the host environment. Examples of physical means are short wavelength radiation such as gamma-radiation and X-radiation, freezing, UV irradiation, lyophilization, and the like. Methods for treatment of microbial cells are disclosed in United States Patent Nos. 4,695,455 and 4,695,462.

The cells generally will have enhanced structural stability which will enhance resistance to environmental conditions. Where the pesticide is in a proform, the method of cell treatment should be selected so as not to inhibit processing of the proform to the mature form of the pesticide by the target pest pathogen. For example, formaldehyde will crosslink proteins and could inhibit processing of the proform of a polypeptide pesticide. The method of treatment should retain at least a substantial portion of the bio-availability or bioactivity of the toxin.

Characteristics of particular interest in selecting a host cell for purposes of production include ease of introducing the *B.t.* gene into the host, availability of expression systems, efficiency of expression, stability of the pesticide in the host, and the presence of auxiliary

genetic capabilities. Characteristics of interest for use as a pesticide microcapsule include protective qualities for the pesticide, such as thick cell walls, pigmentation, and intracellular packaging or formation of inclusion bodies; survival in aqueous environments; lack of mammalian toxicity; attractiveness to pests for ingestion; ease of killing and fixing without damage to the toxin; and the like. Other considerations include ease of formulation and handling, economics, storage stability, and the like.

Growth of cells. The cellular host containing the *B.t.* insecticidal gene may be grown in any convenient nutrient medium, preferably where the DNA construct provides a selective advantage, providing for a selective medium so that substantially all or all of the cells retain the *B.t.* gene. These cells may then be harvested in accordance with conventional ways. Alternatively, the cells can be treated prior to harvesting.

The *B.t.* cells of the invention can be cultured using standard art media and fermentation techniques. Upon completion of the fermentation cycle the bacteria can be harvested by first separating the *B.t.* spores and crystals from the fermentation broth by means well known in the art. The recovered *B.t.* spores and crystals can be formulated into a wettable powder, liquid concentrate, granules or other formulations by the addition of surfactants, dispersants, inert carriers, and other components to facilitate handling and application for particular target pests. These formulations and application procedures are all well known in the art.

Formulations. Formulated bait granules containing an attractant and spores and crystals of the *B.t.* isolates, or recombinant microbes comprising the genes obtainable from the *B.t.* isolates disclosed herein, can be applied to the soil. Formulated product can also be applied as a seed-coating or root treatment or total plant treatment at later stages of the crop cycle. Plant and soil treatments of *B.t.* cells may be employed as wettable powders, granules or dusts, by mixing with various inert materials, such as inorganic minerals (phyllosilicates, carbonates, sulfates, phosphates, and the like) or botanical materials (powdered corncobs, rice hulls, walnut shells, and the like). The formulations may include spreader-sticker adjuvants, stabilizing agents, other pesticidal additives, or surfactants. Liquid formulations may be aqueous-based or non-aqueous and employed as foams, gels, suspensions, emulsifiable concentrates, or the like. The ingredients may include rheological agents, surfactants, emulsifiers, dispersants, or polymers.

As would be appreciated by a person skilled in the art, the pesticidal concentration will vary widely depending upon the nature of the particular formulation, particularly whether it is a concentrate or to be used directly. The pesticide will be present in at least 1% by weight and may be 100% by weight. The dry formulations will have from about 1-95% by weight of the

pesticide while the liquid formulations will generally be from about 1-60% by weight of the solids in the liquid phase. The formulations will generally have from about 10^2 to about 10^4 cells/mg. These formulations will be administered at about 50 mg (liquid or dry) to 1 kg or more per hectare.

5 The formulations can be applied to the environment of the pest, *e.g.*, soil and foliage, by spraying, dusting, sprinkling, or the like.

Mutants. Mutants of the isolates of the invention can be made by procedures well known in the art. For example, an asporogenous mutant can be obtained through ethylmethane sulfonate (EMS) mutagenesis of an isolate. The mutants can be made using ultraviolet light and
10 nitrosoguanidine by procedures well known in the art.

 A smaller percentage of the asporogenous mutants will remain intact and not lyse for extended fermentation periods; these strains are designated lysis minus (-). Lysis minus strains can be identified by screening asporogenous mutants in shake flask media and selecting those mutants that are still intact and contain toxin crystals at the end of the fermentation. Lysis minus
15 strains are suitable for a cell treatment process that will yield a protected, encapsulated toxin protein.

 To prepare a phage resistant variant of said asporogenous mutant, an aliquot of the phage lysate is spread onto nutrient agar and allowed to dry. An aliquot of the phage sensitive bacterial strain is then plated directly over the dried lysate and allowed to dry. The plates are
20 incubated at 30°C. The plates are incubated for 2 days and, at that time, numerous colonies could be seen growing on the agar. Some of these colonies are picked and subcultured onto nutrient agar plates. These apparent resistant cultures are tested for resistance by cross streaking with the phage lysate. A line of the phage lysate is streaked on the plate and allowed to dry. The presumptive resistant cultures are then streaked across the phage line. Resistant bacterial
25 cultures show no lysis anywhere in the streak across the phage line after overnight incubation at 30°C. The resistance to phage is then reconfirmed by plating a lawn of the resistant culture onto a nutrient agar plate. The sensitive strain is also plated in the same manner to serve as the positive control. After drying, a drop of the phage lysate is placed in the center of the plate and allowed to dry. Resistant cultures showed no lysis in the area where the phage lysate has been
30 placed after incubation at 30°C for 24 hours.

Following are examples which illustrates procedures for practicing the invention. These examples should not be construed as limiting.

Example 1 - Bioassay of Cry1F against black cutworm (*Agrotis ipsilon*; Lepidoptera: Noctuidae) larvae

Biological activity of Cry1F against black cutworm (BCW) larvae was determined using standard bioassay procedures using MR872 (a *Pseudomonas fluorescens* clone expressing a cry1Fa/cry1Ab chimeric gene disclosed in U.S. Patent No. 5,840,554) lyophilized powder preparations. BCW bioassays were conducted by incorporating test samples into artificial diet and challenging larvae with treated and untreated diet. All assays were conducted with BCW artificial diet (BioServ Corporation, Frenchtown, NJ). Diet incorporation tests were conducted by mixing the samples with artificial diet (cooled first to 55°C or below) at a rate of 6 mL suspension plus 54 mL diet. Multiple treatment concentrations were generated by serial dilution. After vortexing, this mixture was poured into plastic trays with compartmentalized 3-ml (Nutrend Container Corporation, Jacksonville, FL) at a rate of approximately 30 ml diet per 24-well tray (each well filled halfway). NutrendTM trays with larger wells were used for bioassays with insects in the third instar or older; these also are filled approximately halfway with artificial diet. A water blank containing no test material served as the control. Following at least 30 minutes to allow diet to cool and set, larvae (French Ag Resources, Lamberton, MN) were placed onto the diet mixture, one larva per well. Wells were then sealed with MylarTM sheeting (ClearLam Packaging, IL) using a tacking iron, and several pinholes were made in each well to provide gas exchange. Larvae were held at 25°C for 6 days in a 14:10 (light:dark) holding room. Mortality and stunting were recorded after approximately six days. For LC₅₀ bioassays, a minimum of 3 replicates were run, each with 5-7 doses and approximately 24 insects/dose. Results are shown in Table 2 (Bioassays against first- and third-instar *Agrotis ipsilon* larvae with MR872 Cry1F lyophilized powder preparation).

Table 2

Instar	# test dates	LC ₁₀	LC ₅₀	LC ₉₀	slope	# test insects	# control insects	% control mortality
1	5	9	46	249	1.8	790	220	6
3	3	4	17	67	2.2	249	75	0

Example 2 - Efficacy of Cry1F bait against black cutworm larvae

Cry1F was tested for activity against black cutworm larvae using a bait formulation. MR872 clones were mixed with commercial bait (SoilServ, Salinas, CA) and the resulting mixture was spread across the soil surface of potted wheat seedlings (Feekes growth stage 1.5) at a rate of 50 lb bait/acre. Two concentrations of Cry1F were tested by adding different amounts of toxin to the bait, yielding rates equivalent to 6.25g and 12.5 g of Cry1F toxin/acre. The plants were then infested with 3rd or 4th instar larvae at a rate of approximately 1 insect per 5 plants, and plant damage was measured after 24-48 hours. The damage rating scale runs from 0-4 with 0 being plants cut totally to the ground, and 4 being no visible damage. Controls included plants with no bait added ("no bait"), bait added but without toxin ("blank bait"), and plants with no bait or insects ("control"). Results are shown in Figures 1 and 2.

Example 3 — Efficacy of Transgenic Cry1F Sunflower Plants Against *Agrotis ypsilon* as Measured by Insect Growth and Mortality, and Plant Damage

- Plant treatments:
- Control seedlings at V2 stage
 - *B.t.* seedlings at V2 stage (donors with fixed gene pattern); these plants expressed a gene substantially as shown in SEQ ID NO:3.
- Insect: *Agrotis ypsilon*. Second instar larvae (third laboratory generation).
- Trial: *B.t.* seedlings: five repetitions, five seedlings per repetition, one larva per seedling. Control: five repetitions, five seedlings per repetition, one larva per seedling.
- Protocol:
- Each repetition consisted of five seeds that were planted in transparent rectangular trays (15cm x 25cm) and covered to allow confinement of larval insects.
 - Each repetition was infested with five larvae. Trays were kept in a chamber at 25°C and 75% humidity. Evaluation was done at 24, 48, 72, and 96 hours after infestation.
 - Evaluation
 1. Plants — categorized as undamaged, damaged (bitten stem, cotyledons or leaves) or cut below the cotyledons. A rating was also given to each repetition: 1 (undamaged plants or with the stem with some bites) to 9 (all plants cut and more than 80% of eaten tissue).
 2. Insects — 96 hours after infestation dead larvae were counted, and surviving larvae were weighed.

3. Data were analyzed via contingency table analysis.

Results: Pictures are available that show repetition four (48 and 96 hours after infestation). Data are reported in the following tables, which show evaluation of repetitions 24, 48, 72 and 96 hours after infestation with *Agrotis ypsilon*.

5

10

Table 3. 24 hours after infestation.								
Rep.	No. Undamaged Plants		No. Damaged Plants		No. Cut Plants		Rating	
	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control
1	2	2	2	0	1	3	1	5
2	3	3	0	1	2	1	3	4
3	3	2	2	0	0	3	2	5
4	3	1	0	2	0	2	2	3
5	4	2	0	2	1	1	2	3

15

20

Table 4. 48 hours after infestation.								
Rep.	No. Undamaged Plants		No. Damaged Plants		No. Cut Plants		Rating	
	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control
1	2	1	1	1	2	3	3	7
2	2	0	1	2	2	3	3	7
3	1	1	1	0	3	4	3	7
4	2	1	0	0	3	4	4	7
5	2	1	1	1	2	3	3	7

25

30

Table 5. 72 hours after infestation.								
Rep.	No. Undamaged Plants		No. Damaged Plants		No. Cut Plants		Rating	
	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control
1	2	0	1	0	3	5	3	9
2	0	0	3	0	2	5	4	9
3	1	0	1	0	3	5	5	9
4	1	0	0	0	4	5	6	9
5	2	0	0	0	5	5	5	9

35

Table 6. 96 hours after infestation.									
Rep.	No. Undamaged Plants		No. Damaged Plants		No. Cut Plants		Rating		No. Dead Larvae
	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control	<i>B.t.</i>	control	
1	3	0	0	0	2	5	3	9	4
2	0	0	2	0	3	5	7	9	0
3	1	0	0	0	4	5	5	9	0
4	0	0	0	0	5	5	6	9	0
5	1	0	0	0	4	5	6	9	0

Table 7. Larval weight (mg) 96 hours after infestation*							
Repetition		Larvae per Repetition					
		1	2	3	4	5	Mean
2	<i>B.t.</i>	100	95	65	115	75	90
	Control	230	170	205	175	175	191
3	<i>B.t.</i>	150	135	85	115	120	121
	Control	215	200	90	120	160	157
4	<i>B.t.</i>	110	80	110	85	150	107
	Control	170	160	185	200	155	174
5	<i>B.t.</i>	120	65	95	60	45	77
	Control	120	190	210	255	110	177

*Four repetitions because in one repetition larvae died

Conclusion: The *B.t.* seedlings show activity towards *Agrotis ypsilon* at a high level of infestation (1 larva/seedling). This activity is sufficient to adversely affect the growth of the cutworms so that control is achieved.

Example 4 — Black Cutworm (BCW) Control with New Cry1F Maize Events

Summary. The ability of ten Cry1F maize events (expressing substantially the polynucleotide of SEQ ID NO:3) to control Black Cutworm (*Agrotis ipsilon* (Hufnagel)) under field conditions was compared to two non-Bt hybrids (Control 1 and Control 2), a hybrid expressing *Cry1Ab*, and a hybrid expressing the *Cry9C* protein for purposes of comparison. All the *Cry1F* events provided good protection against stand loss in comparison to the non-Bt hybrids.

Materials and methods. Twenty-five seeds for each of the entries were planted using an AlmacoTM cone planter on Day 1. Four replications were planted — each randomized separately. Rows were in pairs, with an unplanted walkway between each pair of rows. Each plot was enclosed in a galvanized steel barrier that is 8 inches high, 2.5 feet wide and 6 feet long. The barriers were pounded into the ground about 2 inches deep on Days 13 and 14. The rims of the barriers were treated with petroleum jelly to discourage larval escape. Wheat straw was scattered inside the barriers to provide shelter for the cutworms. Shortly after emergence, entry plants with *Cry1F* were tested with ELISA strips and the stands were thinned to 10 plants by the removal of non-expressing plants or surplus expressing plants. In three of the plots, the final stand was 8 or 9 plants. The non-*Cry1F*'s were also thinned to 10 plants. A small plastic stake was placed behind each plant so that it would be apparent that a plant had disappeared, if this occurred. On day 15, when the plants reached late V1 or early V2, three third instar larval black cutworms were placed by each plant. A second infestation of 2 fourth instar BCW per plant was made on Day 19.

Plots were examined each day from Day 16 to Day 26. An additional rating was made on Day 29, and the final rating was made on Day 34. Any plant that received damage was marked with a small plastic stake in front of it and the type of damage was recorded on a data sheet. At the end of the trial the final stand count was taken.

Results. A moderate level of black cutworm pressure was achieved in this trial, resulting in stand reduction in some of the non-Bt entries. For unknown reasons, there was a tendency in this trial for the BCW to dead-heart the plants instead of cutting them. For this reason, the data on damaged plants were also compared.

Table 8 shows plant type and the percent Stand Reduction (%SR) and percent damaged plants (%Dam) for each type of plant. Stand Reduction means that the plant was killed. Damaged plants had obvious feeding damage but did not lose their growing point.

Table 8.		
Treatment/ <i>Cry</i> 1F event	%SR	%Dam
Non- <i>B.t.</i> Control 1	47.5	60
Non- <i>B.t.</i> Control 2	32.5	45
<i>Cry</i> 9C	7.5	20
<i>Cry</i> 1Ab	15	43
308	2.5	5
188	0	5
218	7.5	15
386	7.5	15
538	2.5	18
778	2.5	15
366	3.1	11
663	5	18
058	2.8	9
227	0	5

These results are shown graphically in Figures 3 and 4.

Although no entries were without damage, this is not surprising because the caterpillars have to eat the plants to die. However, in summary, photographs are available that clearly show slightly affected *Cry*1F plants next to completely cut control plants.

Example 5 – Insertion of Toxin Genes Into Plants

One aspect of the subject invention is the transformation of plants with the subject polynucleotide sequences encoding insecticidal toxins. The transformed plants are resistant to attack by the target pest. Preferred genes for use according to the subject invention are optimized for use in plants. Examples of cutworm-active toxins and genes for use according to the subject invention are shown in SEQ ID NOS. 1-8. The protein and gene of SEQ ID NOS. 1 and 2 are preferred.

Obviously, a promoter region capable of expressing the gene in a plant is needed. Thus, for *in planta* expression, the DNA of the subject invention is under the control of and operably linked to an appropriate promoter region. Techniques for obtaining *in planta* expression by using such constructs is known in the art.

Genes encoding pesticidal toxins, as disclosed herein, can be inserted into plant cells using a variety of techniques which are well known in the art. For example, a large number of

cloning vectors comprising a replication system in *E. coli* and a marker that permits selection of the transformed cells are available for preparation for the insertion of foreign genes into higher plants. The vectors comprise, for example, pBR322, pUC series, M13mp series, pACYC184, etc. Accordingly, the sequence encoding the *B.t.* toxin can be inserted into the vector at a suitable restriction site. The resulting plasmid is used for transformation into *E. coli*. The *E. coli* cells are cultivated in a suitable nutrient medium, then harvested and lysed. The plasmid is recovered. Sequence analysis, restriction analysis, electrophoresis, and other biochemical-molecular biological methods are generally carried out as methods of analysis. After each manipulation, the DNA sequence used can be cleaved and joined to the next DNA sequence. Each plasmid sequence can be cloned in the same or other plasmids.

Depending on the method of inserting desired genes into the plant, other DNA sequences may be necessary. If, for example, the Ti or Ri plasmid is used for the transformation of the plant cell, then at least the right border, but often the right and the left border of the Ti or Ri plasmid T-DNA, has to be joined as the flanking region of the genes to be inserted. The use of T-DNA for the transformation of plant cells has been intensively researched and sufficiently described in EP 120 516; Hoekema (1985) In: *The Binary Plant Vector System*, Offset-durkkerij Kanters B.V., Alblasterdam, Chapter 5; Fraley (1986) *et al.*, *Crit. Rev. Plant Sci.* 4:1-46; and An *et al.* (1985) *EMBO J.* 4:277-287.

Once the inserted DNA has been integrated in the genome, it is relatively stable there and, as a rule, does not come out again. It normally contains a selection marker that confers on the transformed plant cells resistance to a biocide or an antibiotic, such as kanamycin, G 418, bleomycin, hygromycin, or chloramphenicol, *inter alia*. The individually employed marker should accordingly permit the selection of transformed cells rather than cells that do not contain the inserted DNA.

A large number of techniques are available for inserting DNA into a plant host cell. Those techniques include transformation with T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation agent, fusion, injection, biolistics (microparticle bombardment), or electroporation as well as other possible methods. If Agrobacteria are used for the transformation, the DNA to be inserted has to be cloned into special plasmids, namely either into an intermediate vector or into a binary vector. The intermediate vectors can be integrated into the Ti or Ri plasmid by homologous recombination owing to sequences that are homologous to sequences in the T-DNA. The Ti or Ri plasmid also comprises the *vir* region necessary for the transfer of the T-DNA. Intermediate vectors cannot replicate themselves in Agrobacteria. The intermediate vector can be transferred into *Agrobacterium tumefaciens* by

means of a helper plasmid (conjugation). Binary vectors can replicate themselves both in *E. coli* and in *Agrobacteria*. They comprise a selection marker gene and a linker or polylinker which are framed by the right and left T-DNA border regions. They can be transformed directly into *Agrobacteria* (Holsters *et al.* [1978] *Mol. Gen. Genet.* 163:181-187). The *Agrobacterium* used as host cell is to comprise a plasmid carrying a *vir* region. The *vir* region is necessary for the transfer of the T-DNA into the plant cell. Additional T-DNA may be contained. The bacterium so transformed is used for the transformation of plant cells. Plant explants can advantageously be cultivated with *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* for the transfer of the DNA into the plant cell. Whole plants can then be regenerated from the infected plant material (for example, pieces of leaf, segments of stalk, roots, but also protoplasts or suspension-cultivated cells) in a suitable medium, which may contain antibiotics or biocides for selection. The plants so obtained can then be tested for the presence of the inserted DNA. No special demands are made of the plasmids in the case of injection and electroporation. It is possible to use ordinary plasmids, such as, for example, pUC derivatives.

The transformed cells grow inside the plants in the usual manner. They can form germ cells and transmit the transformed trait(s) to progeny plants. Such plants can be grown in the normal manner and crossed with plants that have the same transformed hereditary factors or other hereditary factors. The resulting hybrid individuals have the corresponding phenotypic properties.

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art. The scope of the claims should not be limited by the preferred embodiments set forth in the examples, but should be given the broadest interpretation consistent with the description as a whole.

WO 01/13731

PCT/US00/23156

1

SEQUENCE LISTING

<110> Mycogen Corporation

<120> Methods of Controlling Cutworm Pests

<130> MA-731XC1

<140>

<141>

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 3444

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic B.t. toxin gene

<400> 1

```

atggagaaca acatacagaa tcagtgcgtc ccctacaact gcctcaacaa tcctgaagta 60
gagattctca acgaagagag gtcgactggc agattgccgt tagacatctc cctgtccctt 120
acacgtttcc tgttgtctga gtttgttcca ggtgtgggag ttgcgttttg cctcttcgac 180
ctcatctggg gcttcatcac tccatctgat tggagcctct ttcttctcca gattgaacag 240
ttgattgaac aaaggattga gaccttgga aggaatcggg ccatcactac ccttcgtggc 300
ttagcagaca gctatgagat ctacattgaa gcactaagag agtgggaagc caatcctaac 360
aatgcccac tgagagaaga tgtgcgtata cgctttgcta acacagatga tgctttgatc 420
acagccatca acaacttcac cttaccagc ttcgagatcc ctcttctctc ggtctatgtt 480
caagctgcta acctgcactt gtcactactg cgcgacgctg tgtcgttttg gcaagggttg 540
ggactggaca tagctactgt caacaatcac tacaacagac tcatcaatct gattcatcga 600
tacacgaaac attgtttgga tacctacaat cagggattgg agaacctgag aggtactaac 660
actcgccaat gggccagggt caatcagttc aggagagacc ttacacttac tgtgttagac 720
atagttgctc tctttccgaa ctacgatgtt cgtacctatc cgattcaaac gtcatcccaa 780
cttacaaggg agatctacac cagttcagtc attgaagact ctccagtttc tgcgaacata 840
cccaatgggt tcaacagggc tgagtttgga gtcagaccac cccatctcat ggacttcatg 900
aactctttgt ttgtgactgc agagactgtt agatcccaaa ctgtgtgggg aggacactta 960
gttagctcac gcaacacggc tggcaatcgt atcaactttc ctagttacgg ggtcttcaat 1020
cccgggggcg ccatctggat tgcagatgaa gatccacgct ctttctatcg gaccttgtca 1080
gatcctgtct tcgtccgagg aggctttggc aatcctcact atgtactcgg tcttagggga 1140
gtggcctttc aacaaactgg tacgaatcac acccgcatat tcaggaactc cgggaccatt 1200
gactctctag atgagatacc acctcaagac aacagcggcg caccttgga tgactactcc 1260
catgtgctga atcatgttac ctttgtgcgc tggccagggt agatctcagg ttccgactca 1320
tggagagcac caatgttctc ttggacgcat cgtagcgcta ccccccacaa caccattgat 1380
ccagagagaa tcactcagat tcccttggtg aaggcacaca cacttcagtc aggaactaca 1440
gttgtaagag ggccgggggt cacgggagga gacattcttc gacgcactag tggaggacca 1500
ttcgcgtaca ccattgtcaa catcaatggg caacttcccc aaaggatatc tgccaggata 1560
cgctatgcct ctactaccaa tctaagaatc tacgttacgg ttgcagggtg acggatcttt 1620
gctggtcagt tcaacaagac aatggatacc ggtgatccac ttacattcca atctttctcc 1680
tacgccacta tcaacaccgc gttcaccttt ccaatgagcc agagcagttt cacagtaggt 1740

```

```

gctgatacct tcagttcagg caacgaagtg tacattgaca ggtttgagtt gattccagtt 1800
actgccacac tcgaggcaga gtctgacttg gaaagagcac agaaggcggg gaatgctctg 1860
ttcacttcgt ccaatcagat tgggctcaag acagatgtga ctgactatca catcgatcgc 1920
gtttccaacc ttgttgagtg cctctctgat gagttctgtt tggatgagaa gaaggagttg 1980
tccgagaagg tcaaacatgc taagcgactt agtgatgagc ggaacttgct tcaagatccc 2040
aactttcgcg ggatcaacag gcaactagat cgtggatgga ggggaagtac ggacatcacc 2100
attcaaggag gtgatgatgt gttcaaggag aactatgtta cgctcttggg tacctttgat 2160
gagtgcctatc caacatacct gtaccagaag atagatgaat cgaaactcaa agcctacaca 2220
agataccagt tgagagggtta catcgaggac agtcaagacc ttgagatcta cctcatcaga 2280
tacaacgcca aacatgagac agtcaatgtg cctgggacgg gttcactctg gccactttca 2340
gccccaaagt ccatcggcaa gtgtgcccac cactcacacc acttctcctt ggacatagac 2400
gttggtctga ccgacctgaa cgaagacctc ggtgtgtggg tgatcttcaa gatcaagact 2460
caagatggcc atgccaggct aggcaatctg gagtttctag aagagaaacc acttgttgga 2520
gaagccctcg ctagagtgaag gagggctgag aagaagtgga gggacaagag agagaagttg 2580
gaatgggaaa caaacattgt gtacaaagaa gccaaagaaa gcgttgacgc tctgtttgtg 2640
aactctcagt atgataggct ccaagctgat accaacatag ctatgattca tgctgcagac 2700
aaacgcgttc atagcattcg ggaagcttac cttcctgaac ttagcgtgat tccgggtgtc 2760
aatgctgcta tctttgaaga gttagaaggg cgcattctca ctgcattctc cttgtatgat 2820
gcgaggaatg tcatcaagaa tggtgacttc aacaatggcc tatcctgctg gaatgtgaaa 2880
gggcacgtag atgtagaaga acagaacaat caccgctctg tccttgttgt tcctgagtgg 2940
gaagcagaag tttcacaaga agttcgtgtc tgtcctggtc gtggctacat tcttcgtgtt 3000
accgcgtaca aagaaggata cggagaagggt tgcgtcacca tacacgagat tgagaacaac 3060
accgacgagc tgaagttcag caactgcgtc gaggaggaag tctacccaaa caacaccgta 3120
acttgcaatg actacactgc gactcaagag gagtatgagg gtacttacac ttctcgcaat 3180
cgaggatacg atggagccta tgagagcaac tcttctgtac ccgctgacta tgcacagcc 3240
tatgaggaga aggcttacac cgatggacgt agggacaatc cttgcgaatc taacagaggc 3300
tatggggact acacaccgtt accagccggc tatgtcacca aagagttaga gtactttcca 3360
gaaaccgaca aggtttggat tgagattgga gaaacggaag gaacattcat tgttgatagc 3420
gtggagttac ttctgatgga ggaa 3444

```

<210> 2

<211> 1148

<212> PRT

<213> Artificial Sequence

<220>

<223> Toxin encoded by synthetic B.t. gene

<400> 2

```

Met Glu Asn Asn Ile Gln Asn Gln Cys Val Pro Tyr Asn Cys Leu Asn
 1              5              10              15

Asn Pro Glu Val Glu Ile Leu Asn Glu Glu Arg Ser Thr Gly Arg Leu
          20              25              30

Pro Leu Asp Ile Ser Leu Ser Leu Thr Arg Phe Leu Leu Ser Glu Phe
          35              40              45

Val Pro Gly Val Gly Val Ala Phe Gly Leu Phe Asp Leu Ile Trp Gly
          50              55              60

Phe Ile Thr Pro Ser Asp Trp Ser Leu Phe Leu Leu Gln Ile Glu Gln
          65              70              75              80

```

WO 01/13731

PCT/US00/23156

3

Leu Ile Glu Gln Arg Ile Glu Thr Leu Glu Arg Asn Arg Ala Ile Thr
 85 90 95

Thr Leu Arg Gly Leu Ala Asp Ser Tyr Glu Ile Tyr Ile Glu Ala Leu
 100 105 110

Arg Glu Trp Glu Ala Asn Pro Asn Asn Ala Gln Leu Arg Glu Asp Val
 115 120 125

Arg Ile Arg Phe Ala Asn Thr Asp Asp Ala Leu Ile Thr Ala Ile Asn
 130 135 140

Asn Phe Thr Leu Thr Ser Phe Glu Ile Pro Leu Leu Ser Val Tyr Val
 145 150 155 160

Gln Ala Ala Asn Leu His Leu Ser Leu Leu Arg Asp Ala Val Ser Phe
 165 170 175

Gly Gln Gly Trp Gly Leu Asp Ile Ala Thr Val Asn Asn His Tyr Asn
 180 185 190

Arg Leu Ile Asn Leu Ile His Arg Tyr Thr Lys His Cys Leu Asp Thr
 195 200 205

Tyr Asn Gln Gly Leu Glu Asn Leu Arg Gly Thr Asn Thr Arg Gln Trp
 210 215 220

Ala Arg Phe Asn Gln Phe Arg Arg Asp Leu Thr Leu Thr Val Leu Asp
 225 230 235 240

Ile Val Ala Leu Phe Pro Asn Tyr Asp Val Arg Thr Tyr Pro Ile Gln
 245 250 255

Thr Ser Ser Gln Leu Thr Arg Glu Ile Tyr Thr Ser Ser Val Ile Glu
 260 265 270

Asp Ser Pro Val Ser Ala Asn Ile Pro Asn Gly Phe Asn Arg Ala Glu
 275 280 285

Phe Gly Val Arg Pro Pro His Leu Met Asp Phe Met Asn Ser Leu Phe
 290 295 300

Val Thr Ala Glu Thr Val Arg Ser Gln Thr Val Trp Gly Gly His Leu
 305 310 315 320

Val Ser Ser Arg Asn Thr Ala Gly Asn Arg Ile Asn Phe Pro Ser Tyr
 325 330 335

Gly Val Phe Asn Pro Gly Gly Ala Ile Trp Ile Ala Asp Glu Asp Pro
 340 345 350

Arg Pro Phe Tyr Arg Thr Leu Ser Asp Pro Val Phe Val Arg Gly Gly
 355 360 365

WO 01/13731

PCT/US00/23156

4

Phe Gly Asn Pro His Tyr Val Leu Gly Leu Arg Gly Val Ala Phe Gln
 370 375 380

Gln Thr Gly Thr Asn His Thr Arg Thr Phe Arg Asn Ser Gly Thr Ile
 385 390 395 400

Asp Ser Leu Asp Glu Ile Pro Pro Gln Asp Asn Ser Gly Ala Pro Trp
 405 410 415

Asn Asp Tyr Ser His Val Leu Asn His Val Thr Phe Val Arg Trp Pro
 420 425 430

Gly Glu Ile Ser Gly Ser Asp Ser Trp Arg Ala Pro Met Phe Ser Trp
 435 440 445

Thr His Arg Ser Ala Thr Pro Thr Asn Thr Ile Asp Pro Glu Arg Ile
 450 455 460

Thr Gln Ile Pro Leu Val Lys Ala His Thr Leu Gln Ser Gly Thr Thr
 465 470 475 480

Val Val Arg Gly Pro Gly Phe Thr Gly Gly Asp Ile Leu Arg Arg Thr
 485 490 495

Ser Gly Gly Pro Phe Ala Tyr Thr Ile Val Asn Ile Asn Gly Gln Leu
 500 505 510

Pro Gln Arg Tyr Arg Ala Arg Ile Arg Tyr Ala Ser Thr Thr Asn Leu
 515 520 525

Arg Ile Tyr Val Thr Val Ala Gly Glu Arg Ile Phe Ala Gly Gln Phe
 530 535 540

Asn Lys Thr Met Asp Thr Gly Asp Pro Leu Thr Phe Gln Ser Phe Ser
 545 550 555 560

Tyr Ala Thr Ile Asn Thr Ala Phe Thr Phe Pro Met Ser Gln Ser Ser
 565 570 575

Phe Thr Val Gly Ala Asp Thr Phe Ser Ser Gly Asn Glu Val Tyr Ile
 580 585 590

Asp Arg Phe Glu Leu Ile Pro Val Thr Ala Thr Leu Glu Ala Glu Ser
 595 600 605

Asp Leu Glu Arg Ala Gln Lys Ala Val Asn Ala Leu Phe Thr Ser Ser
 610 615 620

Asn Gln Ile Gly Leu Lys Thr Asp Val Thr Asp Tyr His Ile Asp Arg
 625 630 635 640

Val Ser Asn Leu Val Glu Cys Leu Ser Asp Glu Phe Cys Leu Asp Glu
 645 650 655

WO 01/13731

PCT/US00/23156

5

Lys Lys Glu Leu Ser Glu Lys Val Lys His Ala Lys Arg Leu Ser Asp
 660 665 670

Glu Arg Asn Leu Leu Gln Asp Pro Asn Phe Arg Gly Ile Asn Arg Gln
 675 680 685

Leu Asp Arg Gly Trp Arg Gly Ser Thr Asp Ile Thr Ile Gln Gly Gly
 690 695 700

Asp Asp Val Phe Lys Glu Asn Tyr Val Thr Leu Leu Gly Thr Phe Asp
 705 710 715 720

Glu Cys Tyr Pro Thr Tyr Leu Tyr Gln Lys Ile Asp Glu Ser Lys Leu
 725 730 735

Lys Ala Tyr Thr Arg Tyr Gln Leu Arg Gly Tyr Ile Glu Asp Ser Gln
 740 745 750

Asp Leu Glu Ile Tyr Leu Ile Arg Tyr Asn Ala Lys His Glu Thr Val
 755 760 765

Asn Val Pro Gly Thr Gly Ser Leu Trp Pro Leu Ser Ala Pro Ser Pro
 770 775 780

Ile Gly Lys Cys Ala His His Ser His His Phe Ser Leu Asp Ile Asp
 785 790 795 800

Val Gly Cys Thr Asp Leu Asn Glu Asp Leu Gly Val Trp Val Ile Phe
 805 810 815

Lys Ile Lys Thr Gln Asp Gly His Ala Arg Leu Gly Asn Leu Glu Phe
 820 825 830

Leu Glu Glu Lys Pro Leu Val Gly Glu Ala Leu Ala Arg Val Lys Arg
 835 840 845

Ala Glu Lys Lys Trp Arg Asp Lys Arg Glu Lys Leu Glu Trp Glu Thr
 850 855 860

Asn Ile Val Tyr Lys Glu Ala Lys Glu Ser Val Asp Ala Leu Phe Val
 865 870 875 880

Asn Ser Gln Tyr Asp Arg Leu Gln Ala Asp Thr Asn Ile Ala Met Ile
 885 890 895

His Ala Ala Asp Lys Arg Val His Ser Ile Arg Glu Ala Tyr Leu Pro
 900 905 910

Glu Leu Ser Val Ile Pro Gly Val Asn Ala Ala Ile Phe Glu Glu Leu
 915 920 925

Glu Gly Arg Ile Phe Thr Ala Phe Ser Leu Tyr Asp Ala Arg Asn Val
 930 935 940

WO 01/13731

PCT/US00/23156

6

Ile Lys Asn Gly Asp Phe Asn Asn Gly Leu Ser Cys Trp Asn Val Lys
 945 950 955 960
 Gly His Val Asp Val Glu Glu Gln Asn Asn His Arg Ser Val Leu Val
 965 970 975
 Val Pro Glu Trp Glu Ala Glu Val Ser Gln Glu Val Arg Val Cys Pro
 980 985 990
 Gly Arg Gly Tyr Ile Leu Arg Val Thr Ala Tyr Lys Glu Gly Tyr Gly
 995 1000 1005
 Glu Gly Cys Val Thr Ile His Glu Ile Glu Asn Asn Thr Asp Glu Leu
 1010 1015 1020
 Lys Phe Ser Asn Cys Val Glu Glu Glu Val Tyr Pro Asn Asn Thr Val
 1025 1030 1035 1040
 Thr Cys Asn Asp Tyr Thr Ala Thr Gln Glu Glu Tyr Glu Gly Thr Tyr
 1045 1050 1055
 Thr Ser Arg Asn Arg Gly Tyr Asp Gly Ala Tyr Glu Ser Asn Ser Ser
 1060 1065 1070
 Val Pro Ala Asp Tyr Ala Ser Ala Tyr Glu Glu Lys Ala Tyr Thr Asp
 1075 1080 1085
 Gly Arg Arg Asp Asn Pro Cys Glu Ser Asn Arg Gly Tyr Gly Asp Tyr
 1090 1095 1100
 Thr Pro Leu Pro Ala Gly Tyr Val Thr Lys Glu Leu Glu Tyr Phe Pro
 1105 1110 1115 1120
 Glu Thr Asp Lys Val Trp Ile Glu Ile Gly Glu Thr Glu Gly Thr Phe
 1125 1130 1135
 Ile Val Asp Ser Val Glu Leu Leu Leu Met Glu Glu
 1140 1145

<210> 3

<211> 1815

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic B.t. toxin gene

<400> 3

atggagaaca acatacagaa tcagtgcgtc ccctacaact gcctcaacaa tcctgaagta 60
 gagattctca acgaagagag gtcgactggc agattgccgt tagacatctc cctgtccctt 120
 acacgtttcc tggtgtctga gtttgttcca ggtgtgggag ttgcgtttgg cctcttcgac 180
 ctcattctggg gcttcatcac tccattctgat tggagcctct ttcttctcca gattgaacag 240
 ttgattgaac aaaggattga gaccttggaagggaatcggg ccatacactac ccttcgtggc 300
 ttagcagaca gctatgagat ctacattgaa gcactaagag agtgggaagc caatcctaac 360

WO 01/13731

PCT/US00/23156

7

```

aatgccaac tgagagaaga tgtgcgtata cgctttgcta acacagatga tgctttgatc 420
acagccatca acaacttcac ccttaccagc ttcgagatcc ctcttctctc ggtctatggt 480
caagctgcta acctgcactt gtcactactg cgcgacgctg tgctgtttgg gcaagggttg 540
ggactggaca tagctactgt caacaatcac tacaacagac tcatcaatct gattcatcga 600
tacacgaaac attgttttga tacctacaat cagggattgg agaacctgag aggtactaac 660
actcgccaat gggccagggt caatcagttc aggagagacc ttacacttac tgtgttagac 720
atagttgctc tctttccgaa ctacgatgtt cgtacctatc cgattcaaac gtcatcccaa 780
cttacaaggg agatctacac cagttcagtc attgaagact ctccagtttc tgcgaacata 840
cccaatgggt tcaacagggc tgagtttggg gtcagaccac cccatctcat ggacttcatg 900
aactctttgt ttgtgactgc agagactggt agatcccaaa ctgtgtgggg aggacactta 960
gtagctcac gcaacacggc tggcaatcgt atcaactttc ctagttacgg ggtcttcaat 1020
cccgggggcg ccatctggat tgcagatgaa gatccacgtc ctttctatcg gaccttgtca 1080
gactctgtct tcgtccgagg aggctttggc aatcctcact atgtactcgg tcttagggga 1140
gtggcctttc aacaaactgg tacgaatcac acccgcacat tcaggaactc cgggaccatt 1200
gactctctag atgagatacc acctcaagac aacagcggcg caccttgga tgaactactcc 1260
catgtgctga atcatgttac ctttgtgcgc tggccagggt agatctcagg ttccgactca 1320
tggagagcac caatgttctc ttggacgcat cgtagcgcta ccccccacaa caccattgat 1380
ccagagagaa tcaactcagat tcccttggtg aaggcacaca cacttcagtc aggaactaca 1440
gttgtaagag ggccgggggt cacgggagga gacattcttc gacgcactag tggaggacca 1500
ttcgcgtaca ccattgtcaa catcaatggg caacttcccc aaaggatatc tgccaggata 1560
cgctatgcct ctactaccaa tctaagaatc tacgttacgg ttgcagggtg acggatcttt 1620
gctgggtcagt tcaacaagac aatggatacc ggtgatccac ttacattcca atctttctcc 1680
tacgccacta tcaacaccgc gttcaccttt ccaatgagcc agagcagttt cacagtaggt 1740
gctgatacct tcagttcagg caacgaagtg tacattgaca ggtttgagtt gattccagtt 1800
actgccacac tcgag                                     1815

```

<210> 4

<211> 605

<212> PRT

<213> Artificial Sequence

<220>

<223> Toxin encoded by synthetic B.t. gene

<400> 4

```

Met Glu Asn Asn Ile Gln Asn Gln Cys Val Pro Tyr Asn Cys Leu Asn
 1              5              10              15

```

```

Asn Pro Glu Val Glu Ile Leu Asn Glu Glu Arg Ser Thr Gly Arg Leu
      20              25              30

```

```

Pro Leu Asp Ile Ser Leu Ser Leu Thr Arg Phe Leu Leu Ser Glu Phe
 35              40              45

```

```

Val Pro Gly Val Gly Val Ala Phe Gly Leu Phe Asp Leu Ile Trp Gly
 50              55              60

```

```

Phe Ile Thr Pro Ser Asp Trp Ser Leu Phe Leu Leu Gln Ile Glu Gln
 65              70              75              80

```

```

Leu Ile Glu Gln Arg Ile Glu Thr Leu Glu Arg Asn Arg Ala Ile Thr
      85              90              95

```

```

Thr Leu Arg Gly Leu Ala Asp Ser Tyr Glu Ile Tyr Ile Glu Ala Leu
100              105              110

```

WO 01/13731

PCT/US00/23156

8

Arg	Glu	Trp	Glu	Ala	Asn	Pro	Asn	Asn	Ala	Gln	Leu	Arg	Glu	Asp	Val	115	120	125	
Arg	Ile	Arg	Phe	Ala	Asn	Thr	Asp	Asp	Ala	Leu	Ile	Thr	Ala	Ile	Asn	130	135	140	
Asn	Phe	Thr	Leu	Thr	Ser	Phe	Glu	Ile	Pro	Leu	Leu	Ser	Val	Tyr	Val	145	150	155	160
Gln	Ala	Ala	Asn	Leu	His	Leu	Ser	Leu	Leu	Arg	Asp	Ala	Val	Ser	Phe	165	170	175	
Gly	Gln	Gly	Trp	Gly	Leu	Asp	Ile	Ala	Thr	Val	Asn	Asn	His	Tyr	Asn	180	185	190	
Arg	Leu	Ile	Asn	Leu	Ile	His	Arg	Tyr	Thr	Lys	His	Cys	Leu	Asp	Thr	195	200	205	
Tyr	Asn	Gln	Gly	Leu	Glu	Asn	Leu	Arg	Gly	Thr	Asn	Thr	Arg	Gln	Trp	210	215	220	
Ala	Arg	Phe	Asn	Gln	Phe	Arg	Arg	Asp	Leu	Thr	Leu	Thr	Val	Leu	Asp	225	230	235	240
Ile	Val	Ala	Leu	Phe	Pro	Asn	Tyr	Asp	Val	Arg	Thr	Tyr	Pro	Ile	Gln	245	250	255	
Thr	Ser	Ser	Gln	Leu	Thr	Arg	Glu	Ile	Tyr	Thr	Ser	Ser	Val	Ile	Glu	260	265	270	
Asp	Ser	Pro	Val	Ser	Ala	Asn	Ile	Pro	Asn	Gly	Phe	Asn	Arg	Ala	Glu	275	280	285	
Phe	Gly	Val	Arg	Pro	Pro	His	Leu	Met	Asp	Phe	Met	Asn	Ser	Leu	Phe	290	295	300	
Val	Thr	Ala	Glu	Thr	Val	Arg	Ser	Gln	Thr	Val	Trp	Gly	Gly	His	Leu	305	310	315	320
Val	Ser	Ser	Arg	Asn	Thr	Ala	Gly	Asn	Arg	Ile	Asn	Phe	Pro	Ser	Tyr	325	330	335	
Gly	Val	Phe	Asn	Pro	Gly	Gly	Ala	Ile	Trp	Ile	Ala	Asp	Glu	Asp	Pro	340	345	350	
Arg	Pro	Phe	Tyr	Arg	Thr	Leu	Ser	Asp	Pro	Val	Phe	Val	Arg	Gly	Gly	355	360	365	
Phe	Gly	Asn	Pro	His	Tyr	Val	Leu	Gly	Leu	Arg	Gly	Val	Ala	Phe	Gln	370	375	380	
Gln	Thr	Gly	Thr	Asn	His	Thr	Arg	Thr	Phe	Arg	Asn	Ser	Gly	Thr	Ile	385	390	395	400

WO 01/13731

PCT/US00/23156

9

Asp Ser Leu Asp Glu Ile Pro Pro Gln Asp Asn Ser Gly Ala Pro Trp
 405 410 415
 Asn Asp Tyr Ser His Val Leu Asn His Val Thr Phe Val Arg Trp Pro
 420 425 430
 Gly Glu Ile Ser Gly Ser Asp Ser Trp Arg Ala Pro Met Phe Ser Trp
 435 440 445
 Thr His Arg Ser Ala Thr Pro Thr Asn Thr Ile Asp Pro Glu Arg Ile
 450 455 460
 Thr Gln Ile Pro Leu Val Lys Ala His Thr Leu Gln Ser Gly Thr Thr
 465 470 475 480
 Val Val Arg Gly Pro Gly Phe Thr Gly Gly Asp Ile Leu Arg Arg Thr
 485 490 495
 Ser Gly Gly Pro Phe Ala Tyr Thr Ile Val Asn Ile Asn Gly Gln Leu
 500 505 510
 Pro Gln Arg Tyr Arg Ala Arg Ile Arg Tyr Ala Ser Thr Thr Asn Leu
 515 520 525
 Arg Ile Tyr Val Thr Val Ala Gly Glu Arg Ile Phe Ala Gly Gln Phe
 530 535 540
 Asn Lys Thr Met Asp Thr Gly Asp Pro Leu Thr Phe Gln Ser Phe Ser
 545 550 555 560
 Tyr Ala Thr Ile Asn Thr Ala Phe Thr Phe Pro Met Ser Gln Ser Ser
 565 570 575
 Phe Thr Val Gly Ala Asp Thr Phe Ser Ser Gly Asn Glu Val Tyr Ile
 580 585 590
 Asp Arg Phe Glu Leu Ile Pro Val Thr Ala Thr Leu Glu
 595 600 605

<210> 5

<211> 3522

<212> DNA

<213> *Bacillus thuringiensis*

<400> 5

atggagaata atattcaaaa tcaatgcgta ccttacaatt gtttaaataa tcctgaagta 60
 gaaatattaa atgaagaaag aagtactggc agattaccgt tagatatatc cttatcgctt 120
 acacgtttcc ttttgagtga atttgttcca ggtgtgggag ttgcgtttgg attatttgat 180
 ttaatatggg gttttataac tccttctgat tggagcttat ttcttttaca gattgaacaa 240
 ttgattgagc aaagaataga aacattggaa aggaaccggg caattactac attacgaggg 300
 ttagcagata gctatgaaat ttatattgaa gcactaagag agtgggaagc aaatcctaata 360
 aatgcacaat taagggaaga tgtgcgtatt cgatttgcta atacagacga cgctttaata 420
 acagcaataa ataattttac acttacaagt tttgaaatcc ctcttttatc ggtctatggt 480
 caagcggcga atttacattt atcactatta agagacgctg tatcgtttgg gcagggttgg 540

WO 01/13731

PCT/US00/23156

10

```

ggactggata tagctactgt taataatcat tataatagat taataaatct tattcataga 600
tatacgaaac attgttttga cacatacaat caaggattag aaaactttaag aggtactaat 660
actcgacaat gggcaagatt caatcagttt aggagagatt taacacttac tgtattagat 720
atcgttgctc tttttccgaa ctacgatgtt agaacatata caattcaaac gtcattccaa 780
ttaacaaggg aaattttatac aagttcagta attgaggatt ctccagtttc tgctaataata 840
cctaattggtt ttaatagggc ggaatttga gttagaccgc cccatcttat ggactttatg 900
aattctttgt ttgtaactgc agagactgtt agaagtcaaa ctgtgtgggg aggacactta 960
gttagttcac gaaatacggc tggtaaccgt ataaatttcc ctagttacgg ggtcttcaat 1020
cctggtggcg ccatttggat tgcagatgag gatccacgtc ctttttatcg gacattatca 1080
gatcctgttt ttgtccgagg aggatttggg aatcctcatt atgtactggg gcttagggga 1140
gtagcatttc aacaaactgg tacgaaccac acccgaacat ttagaaatag tgggaccata 1200
gattctctag atgaaatccc acctcaggat aatagtgggg caccttggaa tgattatagt 1260
catgtattaa atcatgttac atttgtacga tggccagggt agatttcagg aagtgattca 1320
tggagagctc caatgttttc ttggacgcac cgtagtgcaa cccctacaaa tacaattgat 1380
ccggagagga ttactcaaat accattggta aaagcacata cacttcagtc aggtactact 1440
gttgtaagag ggcccgggtt tacgggagga gatattcttc gacgaacaag tggaggacca 1500
tttgcttata ctattgttaa tataaatggg caattacccc aaaggatatc tgcaagaata 1560
cgctatgcct ctactacaaa tctaagaatt tacgtaacgg ttgcagggtga acggattttt 1620
gctggtcaat ttaacaaaac aatggatacc ggtgaccat taacattcca atcttttagt 1680
tacgcaacta ttaatacagc ttttacattc ccaatgagcc agagtagttt cacagtaggt 1740
gctgatactt ttagttcagg gaatgaagtt tatatagaca gatttgaatt gattccagtt 1800
actgcaacat ttgaagcaga atatgattta gaaagagcac aaaaggcggg gaatgcgctg 1860
tttacttcta taaaccaa ataggataaaa acagatgtga cggattatca tattgatcaa 1920
gtatccaatt tagtggattg tttatcagat gaattttgtc tggatgaaaa gcgagaattg 1980
tccgagaaag tcaaacatgc gaagcgactc agtgatgagc ggaatttact tcaagatcca 2040
aacttcaaag gcatcaatag gcaactagac cgtggttga gaggaagtac ggatattacc 2100
atccaaagag gagatgacgt attcaaagaa aattatgtca cactaccagg tacctttgat 2160
gagtgtctat caacgtattt atatcaaaaa atagatgagt cgaaattaaa accctatact 2220
cgttatcaat taagagggtat tatcgaggat agtcaagact tagaaatcta tttgatccgc 2280
tataatgcaa aacacgaaac agtaaattgt ctaggtacgg gttcttttatg gccgctttca 2340
gtccaaagtc caatcagaaa gtgtggagaa ccgaatcgat gcgcgccaca ccttgaatgg 2400
aatcctgatc tagattgttc ctgcagagac ggggaaaaat gtgcacatca ttcgcatcat 2460
ttctccttgg acattgatgt tggatgtaca gacttaaatt aggacttaga tgtatgggtg 2520
atattcaaga ttaagacgca agatggccat gcaagactag gaaatctaga gtttctcgaa 2580
gagaaacat tagtcgggga agcactagct cgtgtgaaaa gagcagagaa aaaatggaga 2640
gataaacgtg aaaaattgga attggaaaca aatattgttt ataaagaggc aaaagaatct 2700
gtagatgctt tatttgtaaa ctctcaatat gatcaattac aagcggatac gaatattgcc 2760
atgattcatg cggcagataa acgtgttcat agaattcggg aagcgtatct tccagagtta 2820
tctgtgattc cgggtgtaaa tgtagacatt ttcgaagaat taaaagggcg tattttcact 2880
gcattcttcc tatatgatgc gagaaatgtc attaaaaacg gtgatttcaa taatggctta 2940
tcatgctgga acgtgaaagg gcatgtagat gtagaagaac aaaacaacca ccgttcggtc 3000
cttgttggtc cggaatggga agcagaagtg tcacaagaag ttcgtgtctg tccgggtcgt 3060
ggctatatcc ttcgtgtcac agcgtacaag gagggatatg gagaagggtg cgtaaccatt 3120
catgagatcg agaacaatac agacgaactg aagtttagca actgcgtaga agaggaagtc 3180
tatccaaaca acacggtaac gtgtaattgat tatactgcaa atcaagaaga atacgggggt 3240
gcgtacactt cccgtaatcg tggatatgac gaaacttatg gaagcaattc ttctgtacca 3300
gctgattatg cgtcagtcta tgaagaaaaa tcgtatacag atggacgaag agacaatcct 3360
tgtgaatcta acagaggata tggggattac acaccactac cagctggcta tgtgacaaaa 3420
gaattagagt acttcccaga aaccgataag gtatggattg agatcggaga aacggaagga 3480
acattcatcg tggacagcgt ggaattactc cttatggagg aa 3522

```

<210> 6

<211> 1174

<212> PRT

<213> *Bacillus thuringiensis*

WO 01/13731

PCT/US00/23156

11

<400> 6

```

Met Glu Asn Asn Ile Gln Asn Gln Cys Val Pro Tyr Asn Cys Leu Asn
 1           5           10           15

Asn Pro Glu Val Glu Ile Leu Asn Glu Glu Arg Ser Thr Gly Arg Leu
          20           25           30

Pro Leu Asp Ile Ser Leu Ser Leu Thr Arg Phe Leu Leu Ser Glu Phe
      35           40           45

Val Pro Gly Val Gly Val Ala Phe Gly Leu Phe Asp Leu Ile Trp Gly
      50           55           60

Phe Ile Thr Pro Ser Asp Trp Ser Leu Phe Leu Leu Gln Ile Glu Gln
      65           70           75           80

Leu Ile Glu Gln Arg Ile Glu Thr Leu Glu Arg Asn Arg Ala Ile Thr
          85           90           95

Thr Leu Arg Gly Leu Ala Asp Ser Tyr Glu Ile Tyr Ile Glu Ala Leu
          100          105          110

Arg Glu Trp Glu Ala Asn Pro Asn Asn Ala Gln Leu Arg Glu Asp Val
          115          120          125

Arg Ile Arg Phe Ala Asn Thr Asp Asp Ala Leu Ile Thr Ala Ile Asn
          130          135          140

Asn Phe Thr Leu Thr Ser Phe Glu Ile Pro Leu Leu Ser Val Tyr Val
          145          150          155          160

Gln Ala Ala Asn Leu His Leu Ser Leu Leu Arg Asp Ala Val Ser Phe
          165          170          175

Gly Gln Gly Trp Gly Leu Asp Ile Ala Thr Val Asn Asn His Tyr Asn
          180          185          190

Arg Leu Ile Asn Leu Ile His Arg Tyr Thr Lys His Cys Leu Asp Thr
          195          200          205

Tyr Asn Gln Gly Leu Glu Asn Leu Arg Gly Thr Asn Thr Arg Gln Trp
          210          215          220

Ala Arg Phe Asn Gln Phe Arg Arg Asp Leu Thr Leu Thr Val Leu Asp
          225          230          235          240

Ile Val Ala Leu Phe Pro Asn Tyr Asp Val Arg Thr Tyr Pro Ile Gln
          245          250          255

Thr Ser Ser Gln Leu Thr Arg Glu Ile Tyr Thr Ser Ser Val Ile Glu
          260          265          270

Asp Ser Pro Val Ser Ala Asn Ile Pro Asn Gly Phe Asn Arg Ala Glu
          275          280          285

```

WO 01/13731

PCT/US00/23156

12

Phe Gly Val Arg Pro Pro His Leu Met Asp Phe Met Asn Ser Leu Phe
 290 295 300

Val Thr Ala Glu Thr Val Arg Ser Gln Thr Val Trp Gly Gly His Leu
 305 310 315 320

Val Ser Ser Arg Asn Thr Ala Gly Asn Arg Ile Asn Phe Pro Ser Tyr
 325 330 335

Gly Val Phe Asn Pro Gly Gly Ala Ile Trp Ile Ala Asp Glu Asp Pro
 340 345 350

Arg Pro Phe Tyr Arg Thr Leu Ser Asp Pro Val Phe Val Arg Gly Gly
 355 360 365

Phe Gly Asn Pro His Tyr Val Leu Gly Leu Arg Gly Val Ala Phe Gln
 370 375 380

Gln Thr Gly Thr Asn His Thr Arg Thr Phe Arg Asn Ser Gly Thr Ile
 385 390 395 400

Asp Ser Leu Asp Glu Ile Pro Pro Gln Asp Asn Ser Gly Ala Pro Trp
 405 410 415

Asn Asp Tyr Ser His Val Leu Asn His Val Thr Phe Val Arg Trp Pro
 420 425 430

Gly Glu Ile Ser Gly Ser Asp Ser Trp Arg Ala Pro Met Phe Ser Trp
 435 440 445

Thr His Arg Ser Ala Thr Pro Thr Asn Thr Ile Asp Pro Glu Arg Ile
 450 455 460

Thr Gln Ile Pro Leu Val Lys Ala His Thr Leu Gln Ser Gly Thr Thr
 465 470 475 480

Val Val Arg Gly Pro Gly Phe Thr Gly Gly Asp Ile Leu Arg Arg Thr
 485 490 495

Ser Gly Gly Pro Phe Ala Tyr Thr Ile Val Asn Ile Asn Gly Gln Leu
 500 505 510

Pro Gln Arg Tyr Arg Ala Arg Ile Arg Tyr Ala Ser Thr Thr Asn Leu
 515 520 525

Arg Ile Tyr Val Thr Val Ala Gly Glu Arg Ile Phe Ala Gly Gln Phe
 530 535 540

Asn Lys Thr Met Asp Thr Gly Asp Pro Leu Thr Phe Gln Ser Phe Ser
 545 550 555 560

Tyr Ala Thr Ile Asn Thr Ala Phe Thr Phe Pro Met Ser Gln Ser Ser
 565 570 575

WO 01/13731

PCT/US00/23156

13

Phe Thr Val Gly Ala Asp Thr Phe Ser Ser Gly Asn Glu Val Tyr Ile
 580 585 590

Asp Arg Phe Glu Leu Ile Pro Val Thr Ala Thr Phe Glu Ala Glu Tyr
 595 600 605

Asp Leu Glu Arg Ala Gln Lys Ala Val Asn Ala Leu Phe Thr Ser Ile
 610 615 620

Asn Gln Ile Gly Ile Lys Thr Asp Val Thr Asp Tyr His Ile Asp Gln
 625 630 635 640

Val Ser Asn Leu Val Asp Cys Leu Ser Asp Glu Phe Cys Leu Asp Glu
 645 650 655

Lys Arg Glu Leu Ser Glu Lys Val Lys His Ala Lys Arg Leu Ser Asp
 660 665 670

Glu Arg Asn Leu Leu Gln Asp Pro Asn Phe Lys Gly Ile Asn Arg Gln
 675 680 685

Leu Asp Arg Gly Trp Arg Gly Ser Thr Asp Ile Thr Ile Gln Arg Gly
 690 695 700

Asp Asp Val Phe Lys Glu Asn Tyr Val Thr Leu Pro Gly Thr Phe Asp
 705 710 715 720

Glu Cys Tyr Pro Thr Tyr Leu Tyr Gln Lys Ile Asp Glu Ser Lys Leu
 725 730 735

Lys Pro Tyr Thr Arg Tyr Gln Leu Arg Gly Tyr Ile Glu Asp Ser Gln
 740 745 750

Asp Leu Glu Ile Tyr Leu Ile Arg Tyr Asn Ala Lys His Glu Thr Val
 755 760 765

Asn Val Leu Gly Thr Gly Ser Leu Trp Pro Leu Ser Val Gln Ser Pro
 770 775 780

Ile Arg Lys Cys Gly Glu Pro Asn Arg Cys Ala Pro His Leu Glu Trp
 785 790 795 800

Asn Pro Asp Leu Asp Cys Ser Cys Arg Asp Gly Glu Lys Cys Ala His
 805 810 815

His Ser His His Phe Ser Leu Asp Ile Asp Val Gly Cys Thr Asp Leu
 820 825 830

Asn Glu Asp Leu Asp Val Trp Val Ile Phe Lys Ile Lys Thr Gln Asp
 835 840 845

Gly His Ala Arg Leu Gly Asn Leu Glu Phe Leu Glu Glu Lys Pro Leu
 850 855 860

WO 01/13731

PCT/US00/23156

14

Val	Gly	Glu	Ala	Leu	Ala	Arg	Val	Lys	Arg	Ala	Glu	Lys	Lys	Trp	Arg	865	870	875	880
Asp	Lys	Arg	Glu	Lys	Leu	Glu	Leu	Glu	Thr	Asn	Ile	Val	Tyr	Lys	Glu	885	890	895	
Ala	Lys	Glu	Ser	Val	Asp	Ala	Leu	Phe	Val	Asn	Ser	Gln	Tyr	Asp	Gln	900	905	910	
Leu	Gln	Ala	Asp	Thr	Asn	Ile	Ala	Met	Ile	His	Ala	Ala	Asp	Lys	Arg	915	920	925	
Val	His	Arg	Ile	Arg	Glu	Ala	Tyr	Leu	Pro	Glu	Leu	Ser	Val	Ile	Pro	930	935	940	
Gly	Val	Asn	Val	Asp	Ile	Phe	Glu	Glu	Leu	Lys	Gly	Arg	Ile	Phe	Thr	945	950	955	960
Ala	Phe	Phe	Leu	Tyr	Asp	Ala	Arg	Asn	Val	Ile	Lys	Asn	Gly	Asp	Phe	965	970	975	
Asn	Asn	Gly	Leu	Ser	Cys	Trp	Asn	Val	Lys	Gly	His	Val	Asp	Val	Glu	980	985	990	
Glu	Gln	Asn	Asn	His	Arg	Ser	Val	Leu	Val	Val	Pro	Glu	Trp	Glu	Ala	995	1000	1005	
Glu	Val	Ser	Gln	Glu	Val	Arg	Val	Cys	Pro	Gly	Arg	Gly	Tyr	Ile	Leu	1010	1015	1020	
Arg	Val	Thr	Ala	Tyr	Lys	Glu	Gly	Tyr	Gly	Glu	Gly	Cys	Val	Thr	Ile	1025	1030	1035	1040
His	Glu	Ile	Glu	Asn	Asn	Thr	Asp	Glu	Leu	Lys	Phe	Ser	Asn	Cys	Val	1045	1050	1055	
Glu	Glu	Glu	Val	Tyr	Pro	Asn	Asn	Thr	Val	Thr	Cys	Asn	Asp	Tyr	Thr	1060	1065	1070	
Ala	Asn	Gln	Glu	Glu	Tyr	Gly	Gly	Ala	Tyr	Thr	Ser	Arg	Asn	Arg	Gly	1075	1080	1085	
Tyr	Asp	Glu	Thr	Tyr	Gly	Ser	Asn	Ser	Ser	Val	Pro	Ala	Asp	Tyr	Ala	1090	1095	1100	
Ser	Val	Tyr	Glu	Glu	Lys	Ser	Tyr	Thr	Asp	Gly	Arg	Arg	Asp	Asn	Pro	1105	1110	1115	1120
Cys	Glu	Ser	Asn	Arg	Gly	Tyr	Gly	Asp	Tyr	Thr	Pro	Leu	Pro	Ala	Gly	1125	1130	1135	
Tyr	Val	Thr	Lys	Glu	Leu	Glu	Tyr	Phe	Pro	Glu	Thr	Asp	Lys	Val	Trp	1140	1145	1150	

WO 01/13731

PCT/US00/23156

15

Ile Glu Ile Gly Glu Thr Glu Gly Thr Phe Ile Val Asp Ser Val Glu
 1155 1160 1165

Leu Leu Leu Met Glu Glu
 1170

<210> 7

<211> 1815

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic B.t. toxin gene

<400> 7

```

atggagaaca acatacagaa tcagtgtgtc ccctacaact gcctcaacaa tcctgaagta 60
gagattctca acgaagaaag gtcgactggc agattgccgt tagacatctc cctgtccctt 120
acacgattcc tgttgtctga gttcgttcct ggtgtgggtg ttgcgtttgg cctcttcgat 180
ctcatctggg gggtcatcac tccatctgat tggagcctct ttcttctaca gattgaacag 240
ttgattgaac aaaggattga gaccttagaa aggaatcggg ccatcactac acttcgtggg 300
ttagcagaca gctatgagat ctacattgaa gcactaagag agtgggaagc caatcctaac 360
aatgcacaac tgagagaaga tgtgcgcata cgctttgcta acacagatga tgctttgatc 420
acagccatca acaacttcac acttaccagc ttcgagattc ctcttctctc ggtctatgtt 480
caagctgcta accttcactt gtcactactg agggatgctg tgtcgtttgg ccaagggttg 540
ggactggaca tagctactgt caacaatcac tacaacagac tcatcaatct gattcatcga 600
tacacgaaac attgtttgga tacctacaat cagggttgg agaactctgag aggtactaac 660
actcgtcaat gggctaggtt caatcagttc aggagagacc ttacacttac tgtgttagac 720
atagttgctc tctttccgaa ctatgatgtt cgtacctatc cgattcaaac gtcatcccaa 780
cttacaaggg agatctacac cagttcagtc attgaagact ctccagtttc tgcgaaacata 840
ccgaatggtt tcaacagggc tgagtttgga gtcagacctc cccatctcat ggacttcatg 900
aactctttgt ttgtgactgc agaaactgtt agatcgcaaa ctgtgtgggg aggacactta 960
gttagctcaa ggaacacggc tggcaatcgt atcaactttc ctagttacgg ggtcttcaat 1020
cccgggggtg ccatctggat tgcagatgaa gatccacgtc ctttctatcg gaccttgtca 1080
gatcctgtct tcgttcgagg aggctttggc aatcctcact atgtactagg tcttagggga 1140
gtggcctttc aacaaactgg tacgaatcac acacgcacat tcaggaactc cgggaccatt 1200
gactctctag atgagatacc acctcaagac aacagcggcg caccttgga tgactactcg 1260
catgtgctga atcatgttac ctttgtgcgc tggccagggt agatctctgg ttcgactca 1320
tggagagcac ctatgttctc ttggacgcac cgtagcgcta cacctacaaa caccattgat 1380
ccagaaagaa tcactcagat tcccttgggt aaggcacaca cacttcagtc aggaactaca 1440
gttgtaagag ggccgggggt cacgggagga gacattcttc gaaggactag tggaggacca 1500
ttcgcgta ca cattgtcaa catcaatggg caacttcccc aaaggatatg tgctaggata 1560
cgctatgcct ctactaccaa tctacgaatc tatgttacgg ttgcaggtga acggatcttt 1620
gctggtcagt tcaacaagac aatggatacc ggtgatccac ttacattcca atctttctcc 1680
tacgccacta tcaacaccgc gttcaccttt ccaatgagcc agagcagttt cacagtaggt 1740
gctgatacct tcagttcagg gaacgaagtg tacattgata ggtttgagtt gattccagtt 1800
actgctacac tcgag
  
```

1815

<210> 8

<211> 1815

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic B.t. toxin gene

WO 01/13731

PCT/US00/23156

16

<400> 8

```

atggagaaca acatacagaa tcagtgcgtc ccctacaact gcctcaacaa tcctgaggta 60
gagattctca acgaagagag gtcgacgggc agactgccgc tggacatctc cctgtccctc 120
acacgctttc tcctgtctga gttcgttcca ggtgtgggag tcgcgtttgg cctgttcgac 180
ctcatctggg gcttcatcac tccgtcggat tggagcctct ttcttctcca gatcgagcag 240
ttgattgaac agaggattga gaccttggag aggaaccggg ccatcactac ccttcgtggc 300
ttagcagaca gctacgagat ctacattgaa gccctacggg agtgggaggc caatcccaac 360
aatgcccac tgcggggaaga tgtgcgtatc cgcttcgcga acaccgatga cgctctgac 420
accgccatca acaacttcac ccttaccagc ttcgagatac ctctcctctc ggtctatgtt 480
caagctgcga acctgcactt gtcactactg cgcgacgctg tgtcgtttgg gcaaggggtg 540
ggcctggaca tcgctacggt caacaaccac tacaaccgcc tcatcaatct gattcatcga 600
tacacgaaac actgtctgga tacctacaat cagggttgg agaacctgag aggtacgaac 660
actcgccagt gggccagggt caaccagttc aggcgcgacc ttacacttac tgtgctggac 720
atagtcgctc tctttccgaa ctacgacgtt cgtacctatc cgatccaaac gaggttcccag 780
cttaccaggg agatctacac cagctccgtc attgaagact ctccagtgtc ggcgaacata 840
cccaatggct tcaacagggc tgagtccgga gtccgccac cccatctcat ggacttcatg 900
aactctctgt tcgtgactgc agagactgtt agatcccaaa cgggtgtggg aggccactta 960
gtcagctcac gcaacacggc gggcaatcgg atcaactttc ctagctacgg ggtgttcaat 1020
cccgggggcg ccatctggat tgcagatgaa gatccgcggc ccttctatcg gaccttgtcc 1080
gatectgtct tcgtccgagg aggctttggc aacctcact acgtactcgg tctcaggggc 1140
gtggccttcc aacagactgg tacgaatcac accgcacat tcaggaactc cgggaccatc 1200
gactctctag acgagatccc gcctcaagac aacagcggcg caccttggaa tgactactcc 1260
cacgtgctga atcatgttac ctttgtgcgc tggccagggtg agatctcagg ctccgactca 1320
tggcgcgcac caatgttctc gtggacgcat cgtagcgcta ccccaacaaa caccattgat 1380
ccggagagaa tactcagat tcccttgggt aaggcccaca cacttcagtc aggcacgaca 1440
gtggtcagag ggccgggggt cacgggagga gacatccttc gacgcactag tggcggacca 1500
ttcgcgtaca ccattgtcaa catcaacggg cagcttcccc aaaggatatc tgccaggata 1560
cgctatgcct ctactaccaa tctacgcac tacgttacgg tggcaggcga gcgcatcttc 1620
gcgggtcagt tcaacaagac catggacacc ggtgatccac tcacattcca gtctttctcc 1680
tacgccacga tcaacaccgc gttcaccttt ccgatgagcc agagcagctt cacagtaggt 1740
gctgatacct tcagttccgg caacgaagtg tacattgaca ggtttgagtt gatcccagtt 1800
actgccacac tcgag                                     1815

```

CLAIMS:

1. A method for controlling a cutworm pest in an environment wherein said method comprises contacting said pest with a *Cry1F* toxin by delivering said toxin to the environment wherein said delivery is effected by expression of a gene or polynucleotide sequence which encodes the *Cry1F* toxin by a transgenic plant.
- 5 2. The method of claim 1 wherein said toxin is a *Cry1Fa* toxin.
3. The method of claim 1 wherein said toxin is a truncated toxin.
4. The method of claim 1 wherein said toxin is applied to a plant.
5. The method of claim 1 wherein said cutworm pest is of the genus *Agrotis*.
6. The method of claim 1 wherein said cutworm pest is an *Agrotis ipsilon*.
- 10 7. The method of claim 1 wherein said cutworm pest is an *Agrotis malefida*.
8. The method of claim 1 wherein said cutworm pest is of the genus *Porosagrotis*.
9. The method of claim 1 wherein said cutworm pest is a *Porosagrotis gypaetiana*.
10. The method of claim 1 wherein said cutworm pest is of the genus *Xylomyges*.
11. The method of claim 1 wherein said cutworm pest is a *Xylomyges curialis*.
- 15 12. The method of claim 1 wherein said cutworm pest is of the Tribe Agrotini.
13. The method of claim 1 wherein said cutworm pest is of the genus *Feltia*.

14. The method of claim 1 wherein said cutworm pest is a *Feltia jaculifera*.

15. The method of claim 1 wherein said cutworm pest is of the genus *Euxoa*.

16. The method of claim 1 wherein said cutworm pest is an *Euxoa messoria*.

17. The method of claim 1 wherein said cutworm pest is an *Euxoa scandens*.

5 18. The method of claim 1 wherein said cutworm pest is an *Euxoa auxiliaris*.

19. The method of claim 1 wherein said cutworm pest is an *Euxoa detersa*.

20. The method of claim 1 wherein said cutworm pest is an *Euxoa tessellata*.

21. The method of claim 1 wherein said cutworm pest is an *Euxoa ochrogaster*.

22. The method of claim 1 wherein said cutworm pest is of the genus *Peridroma*.

10 23. The method of claim 1 wherein said cutworm pest is a *Peridroma saucia*.

24. The method of claim 4 wherein said plant is a corn plant.

25. The method of claim 4 wherein said plant is a sunflower plant.

26. The method of claim 4 wherein said plant is a soybean plant.

27. The method of claim 4 wherein said plant is a canola plant.

15 28. The method of claim 4 wherein said plant is a cotton plant.

29. The method of claim 1 wherein said environment is soil.




FIG. 1

2/4




FIG. 2

3/4




FIG. 3

4/4




FIG. 4