



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2484975 C 2010/04/20

(11)(21) **2 484 975**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2003/06/04
(87) Date publication PCT/PCT Publication Date: 2003/12/11
(45) Date de délivrance/Issue Date: 2010/04/20
(85) Entrée phase nationale/National Entry: 2004/11/15
(86) N° demande PCT/PCT Application No.: US 2003/017554
(87) N° publication PCT/PCT Publication No.: 2003/101418
(30) Priorité/Priority: 2002/06/04 (US60/385,641)

(51) Cl.Int./Int.Cl. *A61K 8/41* (2006.01),
A61K 8/34 (2006.01), *A61Q 5/02* (2006.01)

(72) Inventeurs/Inventors:
WELLS, ROBERT LEE, US;
ROYCE, DOUGLAS ALLAN, US;
RICCI, NICOLE CRISTINE, US

(73) Propriétaire/Owner:
THE PROCTER & GAMBLE COMPANY, US

(74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(54) Titre : SHAMPOOING CONTENANT UN RESEAU DE GEL
(54) Title: SHAMPOO CONTAINING A GEL NETWORK

(57) **Abrégé/Abstract:**

The compositions of the present invention relate to improved shampoo compositions having from about from about 5 to about 50 weight percent of a deterative surfactant, at least about 0.05 weight percent of a fatty alcohol gel network and at least about 20.0 weight percent of an aqueous carrier.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 December 2003 (11.12.2003)

PCT
(10) International Publication Number
WO 03/101418 A1

(51) International Patent Classification⁷: **A61K 7/50**

(21) International Application Number: PCT/US03/17554

(22) International Filing Date: 4 June 2003 (04.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/385,641 4 June 2002 (04.06.2002) US

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, OH
45202 (US).

(72) Inventors: **WELLS, Robert, Lee**; 4245 Rose Hill venue,
Cincinnati, OH 45229 (US). **ROYCE, Douglas, Allan**;
10424 US 50, Aurora, IN 47001 (US). **RICCI, Nicole,**
Cristine; 9641 Waterford Place #309, Loveland, OH 45140
(US).

(74) Agents: **REED, T., David** et al.; The Procter & Gamble
Company, 6110 Center Hill Road, Cincinnati, OH 45224
(US).

(81) Designated States (*national*): AE, AG, AL, AM, AT (util-
ity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (util-
ity model), DE, DK (utility model), DK, DM, DZ, EC, EE
(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK (utility model), SK, SL, TJ, TM, TN,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: SHAMPOO CONTAINING A GEL NETWORK

(57) Abstract: The compositions of the present invention relate to improved shampoo compositions having from about from about 5 to about 50 weight percent of a deterative surfactant, at least about 0.05 weight percent of a fatty alcohol gel network and at least about 20.0 weight percent of an aqueous carrier.



WO 03/101418 A1

SHAMPOO CONTAINING A GEL NETWORK

FIELD

The present invention relates to a hair cleansing shampoo containing a fatty alcohol gel network.

BACKGROUND

Human hair becomes soiled due to its contact with the surrounding environment and from the sebum secreted by the scalp. The soiling of hair causes it to have a dirty feel and an unattractive appearance. The soiling of the hair necessitates shampooing with frequent regularity.

Shampooing cleans the hair by removing excess soil and sebum. However, shampooing can leave the hair in a wet, tangled, and generally unmanageable state. Once the hair dries, it is often left in a dry, rough, lusterless, or frizzy condition due to removal of the hair's natural oils and other natural conditioning and moisturizing components. The hair can further be left with increased levels of static upon drying, which can interfere with combing and result in a condition commonly referred to as "fly-away hair."

A variety of approaches have been developed to alleviate these after-shampoo problems. These approaches range from post-shampoo application of hair conditioners such as leave-on and rinse-off products, to hair conditioning shampoos which attempt to both cleanse and condition the hair from a single product.

In order to provide hair conditioning benefits in a cleansing shampoo base, a wide variety of conditioning actives have been proposed. However, many of these actives have the disadvantage of leaving the hair feeling soiled or coated and of interfering with the cleansing efficacy of the shampoo.

Coacervate formation in a shampoo composition is known to be advantageous for providing conditioning benefits to the hair. The use of cationic polymers to form coacervates are known in the art, such as in PCT publications WO93/08787 and WO95/01152. However, these shampoo compositions are good for delivering wet hair conditioning but are not capable of delivering satisfactory dry hair smooth feel.

Based on the foregoing, there is a need for a conditioning shampoo which can provide improved conditioning benefit for dry hair, while not interfering with the cleansing efficacy, nor providing negative feel to the hair when it is dried. Specifically, there is a need to provide long lasting moisturized feel, smooth feel, and manageability control to the hair when the hair is dried, yet not leave the hair feeling greasy, as well as to provide softness and ease of combing when the hair is wet.

None of the existing art provides all of the advantages and benefits of the present invention.

SUMMARY OF THE INVENTION

An object of the present invention is to provide a shampoo providing a gel network. In accordance with an aspect of the present invention, there is provided a shampoo composition comprising:

- a) from about 5 to about 50 weight percent of a deterative surfactant,
- b) at least about 0.05 weight percent of a fatty alcohol gel network, and
- c) at least about 20.0 weight percent of an aqueous carrier.

In accordance with another aspect of the invention, there is provided a process for preparing a cleansing composition comprising:

- a) combining a fatty alcohol and a surfactant in a premix at a temperature sufficient to allow partitioning of the surfactant into the fatty alcohol,
- b) cooling the mixture below the chain melt temperature of the premix to form a gel network,
- c) adding the gel network to a deterative surfactant and an aqueous carrier to form a cleansing composition.

The present invention is further directed to a method of using the shampoo composition.

These and other features, aspects, and advantages of the present invention will become evident to those skilled in the art from a reading of the present disclosure.

DETAILED DESCRIPTION

While the specification concludes with claims which particularly point out and distinctly claim the invention, it is believed the present invention will be better understood from the following description.

The shampoo compositions of the present invention include deterative surfactant, a fatty alcohol gel network and an aqueous carrier. Each of these essential components, as well as preferred or optional components, are described in detail hereinafter.

All percentages, parts and ratios are based upon the total weight of the compositions of the present invention, unless otherwise specified. All such weights as they pertain to listed ingredients are based on the active level and, therefore, do not include solvents or by-products that may be included in commercially available materials, unless otherwise specified.

All molecular weights as used herein are weight average molecular weights expressed as grams/mole, unless otherwise specified.

The term "charge density", as used herein, refers to the ratio of the number of positive charges on a polymer to the molecular weight of said polymer.

Herein, "comprising" means that other steps and other ingredients which do not affect the end result can be added. This term encompasses the terms "consisting of" and "consisting essentially of". The compositions and methods/processes of the present invention can comprise, consist of, and consist essentially of the essential elements and limitations of the invention described herein, as well as any of the additional or optional ingredients, components, steps, or limitations described herein.

The term "polymer" as used herein shall include materials whether made by polymerization of one type of monomer or made by two (*i.e.*, copolymers) or more types of monomers.

The term "suitable for application to human hair" as used herein, means that the compositions or components thereof so described are suitable for use in contact with human hair and the scalp and skin without undue toxicity, incompatibility, instability, allergic response, and the like.

The term "water soluble" as used herein, means that the material is soluble in water in the present composition. In general, the material should be soluble at 25° C at a concentration of 0.1% by weight of the water solvent, preferably at 1%, more preferably at 5%, more preferably at 15%.

Citation of any reference is not an admission regarding any determination as to its availability as prior art to the claimed invention.

A. Deterstive Surfactant

The composition of the present invention includes a deterstive surfactant. The deterstive surfactant component is included to provide cleaning performance to the composition. The deterstive surfactant component in turn comprises anionic deterstive surfactant, zwitterionic or amphoteric deterstive surfactant, or a combination thereof. Such surfactants should be physically and chemically compatible with the essential components described, herein, or should not otherwise unduly impair product stability, aesthetics or performance.

Suitable anionic deterstive surfactant components for use in the composition herein include those which are known for use in hair care or other personal care cleansing compositions. The concentration of the anionic surfactant component in the composition should be sufficient to provide the desired cleaning and lather performance, and generally range from about 5% to about 50%, preferably from about 8% to about 30%, more preferably from about 10% to about 25%, even more preferably from about 12% to about 22%, by weight of the composition.

Preferred anionic surfactants suitable for use in the compositions are the alkyl and alkyl ether sulfates. These materials have the respective formulae ROSO_3M and $\text{RO}(\text{C}_2\text{H}_4\text{O})_x\text{SO}_3\text{M}$, wherein R is alkyl or alkenyl of from about 8 to about 18 carbon atoms, x is an integer having a value of from 1 to 10, and M is a cation such as ammonium, alkanolamines, such as triethanolamine, monovalent metals, such as sodium and potassium, and polyvalent metal cations, such as magnesium, and calcium.

Preferably, R has from about 8 to about 18 carbon atoms, more preferably from about 10 to about 16 carbon atoms, even more preferably from about 12 to about 14 carbon atoms, in both the alkyl and alkyl ether sulfates. The alkyl ether sulfates are typically made as condensation products of ethylene oxide and monohydric alcohols having from about 8 to about 24 carbon atoms. The alcohols can be synthetic or they can be derived from fats, e.g., coconut oil, palm

kernel oil, tallow. Lauryl alcohol and straight chain alcohols derived from coconut oil or palm kernel oil are preferred. Such alcohols are reacted with between about 0 and about 10, preferably from about 2 to about 5, more preferably about 3, molar proportions of ethylene oxide, and the resulting mixture of molecular species having, for example, an average of 3 moles of ethylene oxide per mole of alcohol, is sulfated and neutralized.


Other suitable anionic deterative surfactants are the water-soluble salts of organic, sulfuric acid reaction products conforming to the formula $[R^1-SO_3-M]$ where R^1 is a straight or branched chain, saturated, aliphatic hydrocarbon radical having from about 8 to about 24, preferably about 10 to about 18, carbon atoms; and M is a cation described hereinbefore.

Still other suitable anionic deterative surfactants are the reaction products of fatty acids esterified with isethionic acid and neutralized with sodium hydroxide where, for example, the fatty acids are derived from coconut oil or palm kernel oil; sodium or potassium salts of fatty acid amides of methyl tauride in which the fatty acids, for example, are derived from coconut oil or palm kernel oil. Other similar anionic surfactants are described in U.S. Pat. Nos. 2,486,921; 2,486,922; and 2,396,278.

Other anionic deterative surfactants suitable for use in the compositions are the succinates, examples of which include disodium N-octadecylsulfosuccinate; disodium lauryl sulfosuccinate; diammonium lauryl sulfosuccinate; tetrasodium N-(1,2-dicarboxyethyl)-N-octadecylsulfosuccinate; diamyl ester of sodium sulfosuccinic acid; dihexyl ester of sodium sulfosuccinic acid; and dioctyl esters of sodium sulfosuccinic acid.

Other suitable anionic deterative surfactants include olefin sulfonates having about 10 to about 24 carbon atoms. In addition to the true alkene sulfonates and a proportion of hydroxy-alkanesulfonates, the olefin sulfonates can contain minor amounts of other materials, such as alkene disulfonates depending upon the reaction conditions, proportion of reactants, the nature of the starting olefins and impurities in the olefin stock and side reactions during the sulfonation process. A non limiting example of such an alpha-olefin sulfonate mixture is described in U.S. Patent 3,332,880.

Another class of anionic deterative surfactants suitable for use in the compositions are the beta-alkyloxy alkane sulfonates. These surfactants conform to the formula



where R¹ is a straight chain alkyl group having from about 6 to about 20 carbon atoms, R² is a lower alkyl group having from about 1 to about 3 carbon atoms, preferably 1 carbon atom, and M is a water-soluble cation as described hereinbefore.

Preferred anionic deterative surfactants for use in the compositions include ammonium lauryl sulfate, ammonium laureth sulfate, triethylamine lauryl sulfate, triethylamine laureth sulfate, triethanolamine lauryl sulfate, triethanolamine laureth sulfate, monoethanolamine lauryl sulfate, monoethanolamine laureth sulfate, diethanolamine lauryl sulfate, diethanolamine laureth sulfate, lauric monoglyceride sodium sulfate, sodium lauryl sulfate, sodium laureth sulfate, potassium lauryl sulfate, potassium laureth sulfate, sodium lauryl sarcosinate, sodium lauroyl sarcosinate, lauryl sarcosine, cocoyl sarcosine, ammonium cocoyl sulfate, ammonium lauroyl sulfate, sodium cocoyl sulfate, sodium lauroyl sulfate, potassium cocoyl sulfate, potassium lauryl sulfate, triethanolamine lauryl sulfate, triethanolamine lauryl sulfate, monoethanolamine cocoyl sulfate, monoethanolamine lauryl sulfate, sodium tridecyl benzene sulfonate, sodium dodecyl benzene sulfonate, sodium cocoyl isethionate and combinations thereof.

Suitable amphoteric or zwitterionic deterative surfactants for use in the composition herein include those which are known for use in hair care or other personal care cleansing. Concentration of such amphoteric deterative surfactants preferably ranges from about 0.5% to about 20%, preferably from about 1% to about 10%, by weight of the composition. Non limiting examples of suitable zwitterionic or amphoteric surfactants are described in U.S. Pat. Nos. 5,104,646 (Bolich Jr. et al.), 5,106,609 (Bolich Jr. et al.).

Amphoteric deterative surfactants suitable for use in the composition are well known in the art, and include those surfactants broadly described as derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight or branched chain and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic group such as carboxy, sulfonate, sulfate, phosphate, or phosphonate. Preferred amphoteric deterative surfactants for use in the present invention include cocoamphoacetate, cocoamphodiaceate, lauroamphoacetate, lauroamphodiaceate, and mixtures thereof.

Zwitterionic deterative surfactants suitable for use in the composition are well known in the art, and include those surfactants broadly described as derivatives of aliphatic quaternary

ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight or branched chain, and wherein one of the aliphatic substituents contains from about 8 to about 18 carbon atoms and one contains an anionic group such as carboxy, sulfonate, sulfate, phosphate or phosphonate. Zwitterionics such as betaines are preferred.

The compositions of the present invention may further comprise additional surfactants for use in combination with the anionic deterative surfactant component described hereinbefore. Suitable optional surfactants include nonionic and cationic surfactants. Any such surfactant known in the art for use in hair or personal care products may be used, provided that the optional additional surfactant is also chemically and physically compatible with the essential components of the composition, or does not otherwise unduly impair product performance, aesthetics or stability. The concentration of the optional additional surfactants in the composition may vary with the cleansing or lather performance desired, the optional surfactant selected, the desired product concentration, the presence of other components in the composition, and other factors well known in the art.

Non limiting examples of other anionic, zwitterionic, amphoteric or optional additional surfactants suitable for use in the compositions are described in McCutcheon's, Emulsifiers and Detergents, 1989 Annual, published by M. C. Publishing Co., and U.S. Pat. Nos. 3,929,678, 2,658,072; 2,438,091; 2,528,378.

B. Fatty Alcohol Gel Network


Fatty alcohol gel networks have been used for years in cosmetic creams and hair conditioners. These gel networks are formed by combining fatty alcohols and surfactants in the ratio of 1:1 to 40:1 (preferably 2:1 to 20:1, more preferably 3:1 to 10:1). The formation of a gel network involves heating a dispersion of the fatty alcohol in water with the surfactant to a temperature above the melting point of the fatty alcohol. During the mixing process, the fatty alcohol melts, allowing the surfactant to partition into the fatty alcohol droplets. The surfactant brings water along with it into the fatty alcohol. This changes the isotropic fatty alcohol drops into liquid crystalline phase drops. When the mixture is cooled below the chain melt temperature, the liquid crystal phase is converted into a solid crystalline gel network. The gel network contributes a stabilizing benefit to cosmetic creams and hair conditioners. In addition, they deliver conditioned feel benefits for hair conditioners.

The fatty alcohol is preferably included in the composition at a level by weight of from about .05% to about 14%, more preferably from about 1% to about 10%, and more preferably from about 6% to about 8%.

The fatty alcohols useful herein are those having from about 10 to about 40 carbon atoms, preferably from about 12 to about 22 carbon atoms, more preferably from about 16 to about 22

carbon atoms, and more preferably about 16 to about 18 carbon atoms. These fatty alcohols can be straight or branched chain alcohols and can be saturated or unsaturated. Nonlimiting examples of fatty alcohols include, cetyl alcohol, stearyl alcohol, behenyl alcohol, and mixtures thereof. Mixtures of cetyl and stearyl alcohol in a ratio of from about 20:80 to about 80:20, are a preferred embodiment.

Useful surfactants include anionic, zwitterionic and amphoteric surfactants as described above. Other useful surfactants include cationic and nonionic surfactants. Among the cationic surfactants useful herein are those corresponding to the general formula (XI):



wherein at least one of R^{71} , R^{72} , R^{73} and R^{74} is selected from an aliphatic group of from 8 to 30 carbon atoms or an aromatic, alkoxy, polyoxyalkylene, alkylamido, hydroxyalkyl, aryl or alkylaryl group having up to about 22 carbon atoms, the remainder of R^{71} , R^{72} , R^{73} and R^{74} are independently selected from an aliphatic group of from 1 to about 22 carbon atoms or an aromatic, alkoxy, polyoxyalkylene, alkylamido, hydroxyalkyl, aryl or alkylaryl group having up to about 22 carbon atoms; and X is a salt-forming anion such as those selected from halogen, (e.g. chloride, bromide), acetate, citrate, lactate, glycolate, phosphate, nitrate, sulfonate, sulfate, alkylsulfate, and alkyl sulfonate radicals. The aliphatic groups can contain, in addition to carbon and hydrogen atoms, ether linkages, and other groups such as amino groups. The longer chain aliphatic groups, e.g., those of about 12 carbon atoms, or higher, can be saturated or unsaturated. Preferred is when R^{71} , R^{72} , R^{73} and R^{74} are independently selected from C_1 to about C_{22} branched or straight alkyl or alkenyl. Nonlimiting examples of cationic surfactants useful in the present invention include the materials having the following CTFA designations: quaternium-8, quaternium-14, quaternium-18, quaternium-18 methosulfate, quaternium-24, and mixtures thereof.

Among the cationic surfactants of general formula (XI), preferred are those containing in the molecule at least one alkyl chain having at least 16 carbon atoms. Nonlimiting examples of such preferred cationic surfactants include: behenyl trimethyl ammonium chloride available, for example, with tradename INCROQUAT TMC-80 from Croda and ECONOL TM22 from Sanyo Kasei; cetyl trimethyl ammonium chloride available, for example, with tradename CA-2350 from Nikko Chemicals, hydrogenated tallow alkyl trimethyl ammonium chloride, dialkyl (14-18) dimethyl ammonium chloride, ditallow alkyl dimethyl ammonium chloride, dihydrogenated tallow alkyl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dicetyl

dimethyl ammonium chloride, di(behenyl/arachidyl) dimethyl ammonium chloride, dibehenyl dimethyl ammonium chloride, stearyl dimethyl benzyl ammonium chloride, stearyl propyleneglycol phosphate dimethyl ammonium chloride, stearoyl amidopropyl dimethyl benzyl ammonium chloride, stearoyl amidopropyl dimethyl (myristylacetate) ammonium chloride, and N-(stearoyl colamino formyl methy) pyridinium chloride.

Amines are suitable as cationic surfactants. Primary, secondary, and tertiary fatty amines are useful. Particularly useful are tertiary amido amines having an alkyl group of from about 12 to about 22 carbon atoms. Exemplary tertiary amido amines include: stearamidopropyldimethylamine, stearamidopropyldiethylamine, stearamidoethyldiethylamine, stearamidoethyldimethylamine, palmitamidopropyldimethylamine, palmitamidopropyldiethylamine, palmitamidoethyldiethylamine, palmitamidoethyldimethylamine, behenamidopropyldimethylamine, behenamidopropyldiethylamine, behenamidoethyldiethylamine, behenamidoethyldimethylamine, arachidamidopropyldimethylamine, arachidamidopropyldiethylamine, arachidamidoethyldiethylamine, arachidamidoethyldimethylamine, diethylaminoethylstearamide. Also useful are dimethylstearamine, dimethylsoyamine, soyamine, myristylamine, tridecylamine, ethylstearylamine, N-tallowpropane diamine, ethoxylated (with 5 moles of ethylene oxide) stearylamine, dihydroxyethylstearylamine, isostearylamidopropyldimethylamine, oleamidopropyldimethylamine, cocamidopropyldimethylamine and arachidylbehenylamine. Useful amines in the present invention are disclosed in U.S. Patent 4,275,055, Nachtigal, et al.

These amines can also be used in combination with acids such as *l*-glutamic acid, lactic acid, hydrochloric acid, malic acid, succinic acid, acetic acid, fumaric acid, tartaric acid, citric acid, maleic acid, and mixtures thereof; more preferably *l*-glutamic acid, lactic acid, citric acid. The amines herein are preferably partially neutralized with any of the acids at a molar ratio of the amine to the acid of from about 1:0.3 to about 1:3, more preferably from about 1:0.4 to about 1:2.

More preferable cationic surfactants are tertiary amines or quaternary amines. More preferable for the cationic surfactant is mono-alkyl or alkenyl amidoamine, mono-alkyl or alkenyl ammonium salt, di-alkyl ammonium salt, PEG(n) alkylamine, or any combination thereof. The cationic alkyl or alkenyl chain length is between about 10 and about 40 carbon atoms branched or straight, preferably between about 12 and about 22 carbon atoms, more preferably between about 16 and about 18 carbon atoms.

The cationic surfactant comprises between about 0.2% to about 5% by weight of the hair care composition, preferably between about 0.5% and about 4% by weight of the hair care

composition, and more preferably between about 1% and about 3% by weight of the hair care composition.

Nonionic deterative surfactants which are useful in the present invention include those broadly defined as compounds produced by the condensation of alkylene oxide groups (hydrophilic in nature) with an organic hydrophobic compound, which may be aliphatic or alkyl aromatic in nature. Examples of preferred classes of nonionic deterative surfactants are:


1. The polyethylene oxide condensates of alkyl phenols, e.g., the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 20 carbon atoms in either a straight chain or branched chain configuration, with ethylene oxide, the said ethylene oxide being present in amounts equal to from about 10 to about 60 moles of ethylene oxide per mole of alkyl phenol.
2. Those derived from the condensation of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylene diamine products.
3. The condensation product of aliphatic alcohols having from about 8 to about 18 carbon atoms, in either straight chain or branched chain configuration, with ethylene oxide, e.g., a coconut alcohol ethylene oxide condensate having from about 10 to about 30 moles of ethylene oxide per mole of coconut alcohol, the coconut alcohol fraction having from about 10 to about 14 carbon atoms.

4. Long chain tertiary amine oxides corresponding to the following general formula:



wherein R_1 contains an alkyl, alkenyl or monohydroxy alkyl radical of from about 8 to about 18 carbon atoms, from 0 to about 10 ethylene oxide moieties, and from 0 to about 1 glyceryl moiety, and R_2 and R_3 contain from about 1 to about 3 carbon atoms and from 0 to about 1 hydroxy group, e.g., methyl, ethyl, propyl, hydroxyethyl, or hydroxypropyl radicals. The arrow in the formula is a conventional representation of a semipolar bond.

5. Long chain tertiary phosphine oxides corresponding to the following general formula:



wherein R contains an alkyl, alkenyl or monohydroxyalkyl radical ranging from about 8 to about 18 carbon atoms in chain length, from 0 to about 10 ethylene oxide moieties and from 0 to about 1 glyceryl moiety and R' and R'' are each alkyl or monohydroxyalkyl groups containing from about 1 to about 3 carbon atoms.

6. Long chain dialkyl sulfoxides containing one short chain alkyl or hydroxy alkyl radical of from about 1 to about 3 carbon atoms (usually methyl) and one long hydrophobic chain which include alkyl, alkenyl, hydroxy alkyl, or keto alkyl radicals containing from about 8 to about 20 carbon atoms, from 0 to about 10 ethylene oxide moieties and from 0 to about 1 glyceryl moiety.

7. Alkyl polysaccharide (APS) surfactants such as the alkyl polyglycosides. Such surfactants are described in U.S. Patent 4,565,647, Llenado, issued January 21, 1986,

which discloses APS surfactants having a hydrophobic group with about 6 to about 30 carbon atoms and polysaccharide (e.g., polyglycoside) as the hydrophilic group. Optionally, there can be a polyalkylene-oxide group joining the hydrophobic and hydrophilic moieties. The alkyl group (i.e., the hydrophobic moiety) can be saturated or unsaturated, branched or unbranched, and unsubstituted or substituted (e.g., with hydroxy or cyclic rings).

8. Polyethylene glycol (PEG) glyceryl fatty esters, such as those of the formula $R(O)OCH_2CH(OH)CH_2(OCH_2CH_2)_nOH$ wherein n is from about 5 to about 200, preferably from about 20 to about 100, and R is an aliphatic hydrocarbyl having from about 8 to about 20 carbon atoms.

In a preferred embodiment, the gel networks are made from mixtures of cetyl alcohol, stearyl alcohol and cationic and/or anionic surfactant.

C. Aqueous Carrier

The compositions of the present invention are typically in the form of pourable liquids (under ambient conditions). The compositions will therefore typically comprise an aqueous carrier, which is present at a level of from about 20% to about 95%, preferably from about 60% to about 85%, by weight of the compositions. The aqueous carrier may comprise water, or a miscible mixture of water and organic solvent, but preferably comprises water with minimal or no significant concentrations of organic solvent, except as otherwise incidentally incorporated into the composition as minor ingredients of other essential or optional components.

Additional Components

The compositions of the present invention may further comprise one or more optional components known for use in hair care or personal care products, provided that the optional components are physically and chemically compatible with the essential components described herein, or do not otherwise unduly impair product stability, aesthetics or performance. Individual concentrations of such optional components may range from about 0.001% to about 10% by weight of the compositions.

Non-limiting examples of optional components for use in the composition include cationic polymers, conditioning agents (hydrocarbon oils, fatty esters, silicones), anti dandruff agents, suspending agents, viscosity modifiers, dyes, nonvolatile solvents or diluents (water soluble and insoluble), pearlescent aids, foam boosters, additional surfactants or nonionic cosurfactants, pediculocides, pH adjusting agents, perfumes, preservatives, chelants, proteins, skin active agents, sunscreens, UV absorbers, and vitamins.

Deposition Aid

The shampoo compositions of the present invention may include a deposition aid. The deposition aid is included to effectively enhance deposition of the fatty alcohol gel network component described previously. The deposition aid can comprise any material that enhances the deposition of the fatty alcohol gel network from the shampoo onto the hair and/or scalp. Preferably, the deposition aids are cationic polymers. The deposition aid should be physically and chemically compatible with the essential components described herein, or should not otherwise unduly impair product stability, aesthetics or performance. The concentration of the deposition aid in the shampoo composition should be sufficient to effectively enhance the deposition of the fatty alcohol gel network component and typically range from about 0.05% to about 5%, preferably from about 0.075% to about 2.5%, more preferably from about 0.1% to about 1.0%, by weight of the shampoo composition.

Cationic Polymers

The compositions of the present invention may contain a cationic polymer. Concentrations of the cationic polymer in the composition typically range from about 0.05% to about 3%, preferably from about 0.075% to about 2.0%, more preferably from about 0.1% to about 1.0%, by weight of the composition. Preferred cationic polymers will have cationic charge densities of at least about 0.9 meq/gm, preferably at least about 1.2 meq/gm, more preferably at least about 1.5 meq/gm, but also preferably less than about 7 meq/gm, more preferably less than about 5 meq/gm, at the pH of intended use of the composition, which pH will generally range from about pH 3 to about pH 9, preferably between about pH 4 and about pH 8. The "cationic charge density" of a polymer, as that term is used herein, refers to the ratio of the number of positive charges on the polymer to the molecular weight of the polymer. The average molecular weight of such suitable cationic polymers will generally be between about 10,000 and 10 million, preferably between about 50,000 and about 5 million, more preferably between about 100,000 and about 3 million.


Suitable cationic polymers for use in the compositions of the present invention contain cationic nitrogen-containing moieties such as quaternary ammonium or cationic protonated amino moieties. The cationic protonated amines can be primary, secondary, or tertiary amines (preferably secondary or tertiary), depending upon the particular species and the selected pH of the composition. Any anionic counterions can be used in association with the cationic polymers so long as the polymers remain soluble in water, in the composition, or in a coacervate phase of the composition, and so long as the counterions are physically and chemically compatible with the essential components of the composition or do not otherwise unduly impair product performance, stability or aesthetics. Non limiting examples of such counterions include halides (e.g., chloride, fluoride, bromide, iodide), sulfate and methylsulfate.

Non limiting examples of such polymers are described in the CTFA Cosmetic Ingredient Dictionary, 3rd edition, edited by Estrin, Crosley, and Haynes, (The Cosmetic, Toiletry, and Fragrance Association, Inc., Washington, D.C. (1982)).

Non limiting examples of suitable cationic polymers include copolymers of vinyl monomers having cationic protonated amine or quaternary ammonium functionalities with water soluble spacer monomers such as acrylamide, methacrylamide, alkyl and dialkyl acrylamides, alkyl and dialkyl methacrylamides, alkyl acrylate, alkyl methacrylate, vinyl caprolactone or vinyl pyrrolidone.


Suitable cationic protonated amino and quaternary ammonium monomers, for inclusion in the cationic polymers of the composition herein, include vinyl compounds substituted with dialkylaminoalkyl acrylate, dialkylaminoalkyl methacrylate, monoalkylaminoalkyl acrylate, monoalkylaminoalkyl methacrylate, trialkyl methacryloxyalkyl ammonium salt, trialkyl acryloxyalkyl ammonium salt, diallyl quaternary ammonium salts, and vinyl quaternary ammonium monomers having cyclic cationic nitrogen-containing rings such as pyridinium, imidazolium, and quaternized pyrrolidone, e.g., alkyl vinyl imidazolium, alkyl vinyl pyridinium, alkyl vinyl pyrrolidone salts.

Other suitable cationic polymers for use in the compositions include copolymers of 1-vinyl-2-pyrrolidone and 1-vinyl-3-methylimidazolium salt (e.g., chloride salt) (referred to in the industry by the Cosmetic, Toiletry, and Fragrance Association, "CTFA", as Polyquaternium-16); copolymers of 1-vinyl-2-pyrrolidone and dimethylaminoethyl methacrylate (referred to in the industry by CTFA as Polyquaternium-11); cationic diallyl quaternary ammonium-containing polymers, including, for example, dimethyldiallylammonium chloride homopolymer, copolymers of acrylamide and dimethyldiallylammonium chloride (referred to in the industry by CTFA as Polyquaternium 6 and Polyquaternium 7, respectively); amphoteric copolymers of acrylic acid including copolymers of acrylic acid and dimethyldiallylammonium chloride (referred to in the industry by CTFA as Polyquaternium 22), terpolymers of acrylic acid with dimethyldiallylammonium chloride and acrylamide (referred to in the industry by CTFA as Polyquaternium 39), and terpolymers of acrylic acid with methacrylamidopropyl trimethylammonium chloride and methylacrylate (referred to in the industry by CTFA as Polyquaternium 47). Preferred cationic substituted monomers are the cationic substituted dialkylaminoalkyl acrylamides, dialkylaminoalkyl methacrylamides, and combinations thereof. These preferred monomers conform the to the formula



wherein R^1 is hydrogen, methyl or ethyl; each of R^2 , R^3 and R^4 are independently hydrogen or a short chain alkyl having from about 1 to about 8 carbon atoms, preferably from about 1 to about 5 carbon atoms, more preferably from about 1 to about 2 carbon atoms; n is an integer having a value of from about 1 to about 8, preferably from about 1 to about 4; and X is a counterion. The nitrogen attached to R^2 , R^3 and R^4 may be a protonated amine (primary, secondary or tertiary), but is preferably a quaternary ammonium wherein each of R^2 , R^3 and R^4 are alkyl groups a non limiting example of which is polymethacrylamidopropyl trimonium chloride, available under the trade name Polycare 133, from Rhone-Poulenc, Cranberry, N.J., U.S.A. Also preferred are copolymers of the above cationic monomer with nonionic monomers such that the charge density of the total copolymer is from about 2.0 to about 4.5 meq/gm.

Other suitable cationic polymers for use in the composition include polysaccharide polymers, such as cationic cellulose derivatives and cationic starch derivatives. Suitable cationic polysaccharide polymers include those which conform to the formula



wherein A is an anhydroglucose residual group, such as a starch or cellulose anhydroglucose residual; R is an alkylene oxyalkylene, polyoxyalkylene, or hydroxyalkylene group, or combination thereof; R^1 , R^2 , and R^3 independently are alkyl, aryl, alkylaryl, arylalkyl, alkoxyalkyl, or alkoxyaryl groups, each group containing up to about 18 carbon atoms, and the total number of carbon atoms for each cationic moiety (i.e., the sum of carbon atoms in R^1 , R^2 and R^3) preferably being about 20 or less; and X is an anionic counterion as described in hereinbefore.

Preferred cationic cellulose polymers are salts of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide, referred to in the industry (CTFA) as Polyquaternium

10 and available from Amerchol Corp. (Edison, N.J., USA) in their Polymer LR, JR, and KG series of polymers. Other suitable types of cationic cellulose includes the polymeric quaternary ammonium salts of hydroxyethyl cellulose reacted with lauryl dimethyl ammonium-substituted epoxide referred to in the industry (CTFA) as Polyquaternium 24. These materials are available from Amerchol Corp. under the tradename Polymer LM-200.

Other suitable cationic polymers include cationic guar gum derivatives, such as guar hydroxypropyltrimonium chloride, specific examples of which include the Jaguar series commercially available from Rhone-Poulenc Incorporated and the N-Hance series commercially available from Aqualon Division of Hercules, Inc. Other suitable cationic polymers include quaternary nitrogen-containing cellulose ethers, some examples of which are described in U.S. Pat. No. 3,962,418.

Other suitable cationic polymers include copolymers of etherified cellulose, guar and starch, some examples of which are described in U.S. Pat. No. 3,958,581.

When used, the cationic polymers herein are either soluble in the composition or are soluble in a complex coacervate phase in the composition formed by the cationic polymer and the anionic, amphoteric and/or zwitterionic deterative surfactant component described hereinbefore. Complex coacervates of the cationic polymer can also be formed with other charged materials in the composition.


Techniques for analysis of formation of complex coacervates are known in the art. For example, microscopic analyses of the compositions, at any chosen stage of dilution, can be utilized to identify whether a coacervate phase has formed. Such coacervate phase will be identifiable as an additional emulsified phase in the composition. The use of dyes can aid in distinguishing the coacervate phase from other insoluble phases dispersed in the composition.

Dispersed Particles

The composition of the present invention may include dispersed particles. In the compositions of the present invention, it is preferable to incorporate at least 0.025% by weight of the dispersed particles, more preferably at least 0.05%, still more preferably at least 0.1%, even more preferably at least 0.25%, and yet more preferably at least 0.5% by weight of the dispersed particles. In the compositions of the present invention, it is preferable to incorporate no more than about 20% by weight of the dispersed particles, more preferably no more than about 10%, still more preferably no more than 5%, even more preferably no more than 3%, and yet more preferably no more than 2% by weight of the dispersed particles.

Nonionic polymers

Polyalkylene glycols having a molecular weight of more than about 1000 are useful herein. Useful are those having the following general formula:



wherein R⁹⁵ is selected from the group consisting of H, methyl, and mixtures thereof. Polyethylene glycol polymers useful herein are PEG-2M (also known as Polyox WSR[®] N-10, which is available from Union Carbide and as PEG-2,000); PEG-5M (also known as Polyox WSR[®] N-35 and Polyox WSR[®] N-80, available from Union Carbide and as PEG-5,000 and Polyethylene Glycol 300,000); PEG-7M (also known as Polyox WSR[®] N-750 available from Union Carbide); PEG-9M (also known as Polyox WSR[®] N-3333 available from Union Carbide); and PEG-14 M (also known as Polyox WSR[®] N-3000 available from Union Carbide).

Conditioning agents

Conditioning agents include any material which is used to give a particular conditioning benefit to hair and/or skin. In hair treatment compositions, suitable conditioning agents are those which deliver one or more benefits relating to shine, softness, combability, antistatic properties, wet-handling, damage, manageability, body, and greasiness. The conditioning agents useful in the compositions of the present invention typically comprise a water insoluble, water dispersible, non-volatile, liquid that forms emulsified, liquid particles or are solubilized by the surfactant micelles, in the anionic deterative surfactant component (described herein). Suitable conditioning agents for use in the composition are those conditioning agents characterized generally as silicones (e.g., silicone oils, cationic silicones, silicone gums, high refractive silicones, and silicone resins), organic conditioning oils (e.g., hydrocarbon oils, polyolefins, and fatty esters) or combinations thereof, or those conditioning agents which otherwise form liquid, dispersed particles in the aqueous surfactant matrix herein. Such conditioning agents should be physically and chemically compatible with the essential components of the composition, and should not otherwise unduly impair product stability, aesthetics or performance.

The concentration of the conditioning agent in the composition should be sufficient to provide the desired conditioning benefits, and as will be apparent to one of ordinary skill in the art. Such concentration can vary with the conditioning agent, the conditioning performance desired, the average size of the conditioning agent particles, the type and concentration of other components, and other like factors.

1. Silicones

The conditioning agent of the compositions of the present invention is preferably an insoluble silicone conditioning agent. The silicone conditioning agent particles may comprise

volatile silicone, non-volatile silicone, or combinations thereof. Preferred are non-volatile silicone conditioning agents. If volatile silicones are present, it will typically be incidental to their use as a solvent or carrier for commercially available forms of non-volatile silicone materials ingredients, such as silicone gums and resins. The silicone conditioning agent particles may comprise a silicone fluid conditioning agent and may also comprise other ingredients, such as a silicone resin to improve silicone fluid deposition efficiency or enhance glossiness of the hair.

The concentration of the silicone conditioning agent typically ranges from about 0.01% to about 10%, by weight of the composition, preferably from about 0.1% to about 8%, more preferably from about 0.1% to about 5%, more preferably from about 0.2% to about 3%. Non-limiting examples of suitable silicone conditioning agents, and optional suspending agents for the silicone, are described in U.S. Reissue Pat. No. 34,584, U.S. Pat. No. 5,104,646, and U.S. Pat. No. 5,106,609.

The silicone conditioning agents for use in the compositions of the present invention preferably have a viscosity, as measured at 25°C, from about 20 to about 2,000,000 centistokes ("csk"), more preferably from about 1,000 to about 1,800,000 csk, even more preferably from about 50,000 to about 1,500,000 csk, more preferably from about 100,000 to about 1,500,000 csk.

The dispersed silicone conditioning agent particles typically have a volume average particle diameter ranging from about 0.01 μ m to about 50 μ m. For small particle application to hair, the volume average particle diameters typically range from about 0.01 μ m to about 4 μ m, preferably from about 0.01 μ m to about 2 μ m, more preferably from about 0.01 μ m to about 0.5 μ m. For larger particle application to hair, the volume average particle diameters typically range from about 5 μ m to about 125 μ m, preferably from about 10 μ m to about 90 μ m, more preferably from about 15 μ m to about 70 μ m, more preferably from about 20 μ m to about 50 μ m.


Background material on silicones including sections discussing silicone fluids, gums, and resins, as well as manufacture of silicones, are found in *Encyclopedia of Polymer Science and Engineering*, vol. 15, 2d ed., pp 204-308, John Wiley & Sons, Inc. (1989).

a. Silicone oils

Silicone fluids include silicone oils, which are flowable silicone materials having a viscosity, as measured at 25°C, less than 1,000,000 csk, preferably from about 5 csk to about

1,000,000 csk, more preferably from about 100 csk to about 600,000 csk. Suitable silicone oils for use in the compositions of the present invention include polyalkyl siloxanes, polyaryl siloxanes, polyalkylaryl siloxanes, polyether siloxane copolymers, and mixtures thereof. Other insoluble, non-volatile silicone fluids having hair conditioning properties may also be used.

Silicone oils include polyalkyl or polyaryl siloxanes which conform to the following Formula (I):



wherein R is aliphatic, preferably alkyl or alkenyl, or aryl, R can be substituted or unsubstituted, and x is an integer from 1 to about 8,000. Suitable R groups for use in the compositions of the present invention include, but are not limited to: alkoxy, aryloxy, alkaryl, arylalkyl, arylalkenyl, alkamino, and ether-substituted, hydroxyl-substituted, and halogen-substituted aliphatic and aryl groups. Suitable R groups also include cationic amines and quaternary ammonium groups.

Preferred alkyl and alkenyl substituents are C₁ to C₅ alkyls and alkenyls, more preferably from C₁ to C₄, more preferably from C₁ to C₂. The aliphatic portions of other alkyl-, alkenyl-, or alkynyl-containing groups (such as alkoxy, alkaryl, and alkamino) can be straight or branched chains, and are preferably from C₁ to C₅, more preferably from C₁ to C₄, even more preferably from C₁ to C₃, more preferably from C₁ to C₂. As discussed above, the R substituents can also contain amino functionalities (e.g. alkamino groups), which can be primary, secondary or tertiary amines or quaternary ammonium. These include mono-, di- and tri- alkylamino and alkoxyamino groups, wherein the aliphatic portion chain length is preferably as described herein.

b. Amino and Cationic silicones

Cationic silicone fluids suitable for use in the compositions of the present invention include, but are not limited to, those which conform to the general formula (II):



wherein G is hydrogen, phenyl, hydroxy, or C₁-C₈ alkyl, preferably methyl; a is 0 or an integer having a value from 1 to 3, preferably 0; b is 0 or 1, preferably 1; n is a number from 0 to 1,999, preferably from 49 to 499; m is an integer from 1 to 2,000, preferably from 1 to 10; the sum of n and m is a number from 1 to 2,000, preferably from 50 to 500; R₁ is a monovalent radical

$$-\text{N}(\text{R}_2)\text{CH}_2-\text{CH}_2-\text{NR}_2\text{H}_2\text{A}^-$$
$$\begin{array}{c} \text{(CH}_3\text{)}_3\text{Si} - \text{O} - \text{Si} \begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3 \end{array} \left[\begin{array}{c} \text{---} \end{array} \right]_n \text{O} - \text{Si} \begin{array}{c} \text{CH}_3 \\ | \\ \text{(CH}_2\text{)}_3 \\ | \\ \text{NH} \\ | \\ \text{(CH}_2\text{)}_2 \\ | \\ \text{NH}_2 \end{array} \left[\begin{array}{c} \text{---} \end{array} \right]_m \text{OSi(CH}_3\text{)}_3 \end{array}$$
$$\begin{array}{c} \text{R}^4\text{CH}_2\text{---CHOH---CH}_2\text{---N}^+(\text{R}^3)_3\text{Q}^- \\ | \\ (\text{R}^3)_3\text{Si---O---}\left[\text{Si}\text{---O}\right]_r\text{---}\left[\text{Si}\text{---O}\right]_s\text{---Si---O---Si(R}^3)_3 \\ | \qquad \qquad | \\ \text{R}^3 \qquad \qquad \text{R}^3 \end{array}$$

wherein R³ is a monovalent hydrocarbon radical from C₁ to C₁₈, preferably an alkyl or alkenyl radical, such as methyl; R₄ is a hydrocarbon radical, preferably a C₁ to C₁₈ alkylene radical or a C₁₀ to C₁₈ alkyleneoxy radical, more preferably a C₁ to C₈ alkyleneoxy radical; Q⁻ is a halide ion, preferably chloride; r is an average statistical value from 2 to 20, preferably from 2 to 8; s is an

average statistical value from 20 to 200, preferably from 20 to 50. A preferred polymer of this class is known as UCARE SILICONE ALE 56™, available from Union Carbide.

c. Silicone gums


Other silicone fluids suitable for use in the compositions of the present invention are the insoluble silicone gums. These gums are polyorganosiloxane materials having a viscosity, as measured at 25°C, of greater than or equal to 1,000,000 csk. Silicone gums are described in U.S. Pat. No. 4,152,416; Noll and Walter, *Chemistry and Technology of Silicones*, New York: Academic Press (1968); and in General Electric Silicone Rubber Product Data Sheets SE 30, SE 33, SE 54 and SE 76.

Specific non-limiting examples of silicone gums for use in the compositions of the present invention include polydimethylsiloxane, (polydimethylsiloxane) (methylvinylsiloxane) copolymer, poly(dimethylsiloxane) (diphenyl siloxane)(methylvinylsiloxane) copolymer and mixtures thereof.

d. High refractive index silicones

Other non-volatile, insoluble silicone fluid conditioning agents that are suitable for use in the compositions of the present invention are those known as "high refractive index silicones," having a refractive index of at least about 1.46, preferably at least about 1.48, more preferably at least about 1.52, more preferably at least about 1.55. The refractive index of the polysiloxane fluid will generally be less than about 1.70, typically less than about 1.60. In this context, polysiloxane "fluid" includes oils as well as gums.

The high refractive index polysiloxane fluid includes those represented by general Formula (I) above, as well as cyclic polysiloxanes such as those represented by Formula (V) below:



wherein R is as defined above, and n is a number from about 3 to about 7, preferably from about 3 to about 5.

The high refractive index polysiloxane fluids contain an amount of aryl-containing R substituents sufficient to increase the refractive index to the desired level, which is described herein. Additionally, R and n must be selected so that the material is non-volatile.

Aryl-containing substituents include those which contain alicyclic and heterocyclic five and six member aryl rings and those which contain fused five or six member rings. The aryl rings themselves can be substituted or unsubstituted.

Generally, the high refractive index polysiloxane fluids will have a degree of aryl-containing substituents of at least about 15%, preferably at least about 20%, more preferably at least about 25%, even more preferably at least about 35%, more preferably at least about 50%. Typically, the degree of aryl substitution will be less than about 90%, more generally less than about 85%, preferably from about 55% to about 80%.

Preferred high refractive index polysiloxane fluids have a combination of phenyl or phenyl derivative substituents (more preferably phenyl), with alkyl substituents, preferably C₁-C₄ alkyl (more preferably methyl), hydroxy, or C₁-C₄ alkylamino (especially -R¹NHR²NH₂ wherein each R¹ and R² independently is a C₁-C₃ alkyl, alkenyl, and/or alkoxy).

When high refractive index silicones are used in the compositions of the present invention, they are preferably used in solution with a spreading agent, such as a silicone resin or a surfactant, to reduce the surface tension by a sufficient amount to enhance spreading and thereby enhance the glossiness (subsequent to drying) of hair treated with the compositions.

Silicone fluids suitable for use in the compositions of the present invention are disclosed in U.S. Pat. No. 2,826,551, U.S. Pat. No. 3,964,500, U.S. Pat. No. 4,364,837, British Pat. No. 849,433, and *Silicon Compounds*, Petrarch Systems, Inc. (1984).

e. Silicone resins

Silicone resins may be included in the silicone conditioning agent of the compositions of the present invention. These resins are highly cross-linked polymeric siloxane systems. The cross-linking is introduced through the incorporation of trifunctional and tetrafunctional silanes with monofunctional or difunctional, or both, silanes during manufacture of the silicone resin.

Silicone materials and silicone resins in particular, can conveniently be identified according to a shorthand nomenclature system known to those of ordinary skill in the art as "MDTQ" nomenclature. Under this system, the silicone is described according to presence of various siloxane monomer units which make up the silicone. Briefly, the symbol M denotes the monofunctional unit (CH₃)₃SiO_{0.5}; D denotes the difunctional unit (CH₃)₂SiO; T denotes the trifunctional unit (CH₃)SiO_{1.5}; and Q denotes the quadra- or tetra-functional unit SiO₂. Primes of the unit symbols (e.g. M', D', T', and Q') denote substituents other than methyl, and must be specifically defined for each occurrence.

Preferred silicone resins for use in the compositions of the present invention include, but are not limited to MQ, MT, MTQ, MDT and MDTQ resins. Methyl is a preferred silicone

substituent. Especially preferred silicone resins are MQ resins, wherein the M:Q ratio is from about 0.5:1.0 to about 1.5:1.0 and the average molecular weight of the silicone resin is from about 1000 to about 10,000.

The weight ratio of the non-volatile silicone fluid, having refractive index below 1.46, to the silicone resin component, when used, is preferably from about 4:1 to about 400:1, more preferably from about 9:1 to about 200:1, more preferably from about 19:1 to about 100:1, particularly when the silicone fluid component is a polydimethylsiloxane fluid or a mixture of polydimethylsiloxane fluid and polydimethylsiloxane gum as described herein. Insofar as the silicone resin forms a part of the same phase in the compositions hereof as the silicone fluid, i.e. the conditioning active, the sum of the fluid and resin should be included in determining the level of silicone conditioning agent in the composition.

2. Organic conditioning oils

The conditioning component of the compositions of the present invention may also comprise from about 0.05% to about 3%, by weight of the composition, preferably from about 0.08% to about 1.5%, more preferably from about 0.1% to about 1%, of at least one organic conditioning oil as the conditioning agent, either alone or in combination with other conditioning agents, such as the silicones (described herein).

a. Hydrocarbon oils

Suitable organic conditioning oils for use as conditioning agents in the compositions of the present invention include, but are not limited to, hydrocarbon oils having at least about 10 carbon atoms, such as cyclic hydrocarbons, straight chain aliphatic hydrocarbons (saturated or unsaturated), and branched chain aliphatic hydrocarbons (saturated or unsaturated), including polymers and mixtures thereof. Straight chain hydrocarbon oils preferably are from about C₁₂ to about C₁₉. Branched chain hydrocarbon oils, including hydrocarbon polymers, typically will contain more than 19 carbon atoms.

Specific non-limiting examples of these hydrocarbon oils include paraffin oil, mineral oil, saturated and unsaturated dodecane, saturated and unsaturated tridecane, saturated and unsaturated tetradecane, saturated and unsaturated pentadecane, saturated and unsaturated hexadecane, polybutene, polydecene, and mixtures thereof. Branched-chain isomers of these compounds, as well as of higher chain length hydrocarbons, can also be used, examples of which include highly branched, saturated or unsaturated, alkanes such as the permethyl-substituted isomers, e.g., the permethyl-substituted isomers of hexadecane and eicosane, such as 2, 2, 4, 4, 6, 6, 8, 8-dimethyl-10-methylundecane and 2, 2, 4, 4, 6, 6-dimethyl-8-methylnonane, available from Permethyl Corporation. Hydrocarbon polymers such as polybutene and polydecene. A preferred hydrocarbon polymer is polybutene, such as the copolymer of isobutylene and butene. A

commercially available material of this type is L-14 polybutene from Amoco Chemical Corporation. The concentration of such hydrocarbon oils in the composition preferably range from about 0.05% to about 20%, more preferably from about 0.08% to about 1.5%, and even more preferably from about 0.1% to about 1%, by weight of the composition.

b. Polyolefins

Organic conditioning oils for use in the compositions of the present invention can also include liquid polyolefins, more preferably liquid poly- α -olefins, more preferably hydrogenated liquid poly- α -olefins. Polyolefins for use herein are prepared by polymerization of C_4 to about C_{14} olefinic monomers, preferably from about C_6 to about C_{12} .

Non-limiting examples of olefinic monomers for use in preparing the polyolefin liquids herein include ethylene, propylene, 1-butene, 1-pentene, 1-hexene, 1-octene, 1-decene, 1-dodecene, 1-tetradecene, branched chain isomers such as 4-methyl-1-pentene, and mixtures thereof. Also suitable for preparing the polyolefin liquids are olefin-containing refinery feedstocks or effluents. Preferred hydrogenated α -olefin monomers include, but are not limited to: 1-hexene to 1-hexadecenes, 1-octene to 1-tetradecene, and mixtures thereof.

c. Fatty Esters

Other suitable organic conditioning oils for use as the conditioning agent in the compositions of the present invention include, but are not limited to, fatty esters having at least 10 carbon atoms. These fatty esters include esters with hydrocarbyl chains derived from fatty acids or alcohols (e.g. mono-esters, polyhydric alcohol esters, and di- and tri-carboxylic acid esters). The hydrocarbyl radicals of the fatty esters hereof may include or have covalently bonded thereto other compatible functionalities, such as amides and alkoxy moieties (e.g., ethoxy or ether linkages, etc.).

Specific examples of preferred fatty esters include, but are not limited to: isopropyl isostearate, hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate, dihexyldecyl adipate, lauryl lactate, myristyl lactate, cetyl lactate, oleyl stearate, oleyl oleate, oleyl myristate, lauryl acetate, cetyl propionate, and oleyl adipate.

Other fatty esters suitable for use in the compositions of the present invention are mono-carboxylic acid esters of the general formula $R'COOR$, wherein R' and R are alkyl or alkenyl radicals, and the sum of carbon atoms in R' and R is at least 10, preferably at least 22.


Still other fatty esters suitable for use in the compositions of the present invention are di- and tri-alkyl and alkenyl esters of carboxylic acids, such as esters of C_4 to C_8 dicarboxylic acids (e.g. C_1 to C_{22} esters, preferably C_1 to C_6 , of succinic acid, glutaric acid, and adipic acid). Specific

non-limiting examples of di- and tri- alkyl and alkenyl esters of carboxylic acids include isocetyl stearyl stearate, diisopropyl adipate, and tristearyl citrate.


Other fatty esters suitable for use in the compositions of the present invention are those known as polyhydric alcohol esters. Such polyhydric alcohol esters include alkylene glycol esters, such as ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty acid esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters.

Still other fatty esters suitable for use in the compositions of the present invention are glycerides, including, but not limited to, mono-, di-, and tri-glycerides, preferably di- and tri-glycerides, more preferably triglycerides. For use in the compositions described herein, the glycerides are preferably the mono-, di-, and tri-esters of glycerol and long chain carboxylic acids, such as C₁₀ to C₂₂ carboxylic acids. A variety of these types of materials can be obtained from vegetable and animal fats and oils, such as castor oil, safflower oil, cottonseed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, lanolin and soybean oil. Synthetic oils include, but are not limited to, triolein and tristearin glyceryl dilaurate.

Other fatty esters suitable for use in the compositions of the present invention are water insoluble synthetic fatty esters. Some preferred synthetic esters conform to the general Formula (VI):



wherein R¹ is a C₇ to C₉ alkyl, alkenyl, hydroxyalkyl or hydroxyalkenyl group, preferably a saturated alkyl group, more preferably a saturated, linear, alkyl group; n is a positive integer having a value from 2 to 4, preferably 3; and Y is an alkyl, alkenyl, hydroxy or carboxy substituted alkyl or alkenyl, having from about 2 to about 20 carbon atoms, preferably from about 3 to about 14 carbon atoms. Other preferred synthetic esters conform to the general Formula (VII):



wherein R^2 is a C_8 to C_{10} alkyl, alkenyl, hydroxyalkyl or hydroxyalkenyl group; preferably a saturated alkyl group, more preferably a saturated, linear, alkyl group; n and Y are as defined above in Formula (VII).

Specific non-limiting examples of suitable synthetic fatty esters for use in the compositions of the present invention include: P-43 (C_8 - C_{10} triester of trimethylolpropane), MCP-684 (tetraester of 3,3 diethanol-1,5 pentadiol), MCP 121 (C_8 - C_{10} diester of adipic acid), all of which are available from Mobil Chemical Company.

3. Other conditioning agents

Also suitable for use in the compositions herein are the conditioning agents described by the Procter & Gamble Company in U.S. Pat. Nos. 5,674,478, and 5,750,122.

Also suitable for use herein are those conditioning agents described in U.S. Pat. Nos. 4,529,586 (Clairol), 4,507,280 (Clairol), 4,663,158 (Clairol), 4,197,865 (L'Oreal), 4,217, 914 (L'Oreal), 4,381,919 (L'Oreal), and 4,422, 853 (L'Oreal).

Anti-dandruff Actives

The compositions of the present invention may also contain an anti-dandruff agent. Suitable, non-limiting examples of anti-dandruff particulates include: pyridinethione salts, azoles, selenium sulfide, particulate sulfur, and mixtures thereof. Preferred are pyridinethione salts. Such anti-dandruff particulate should be physically and chemically compatible with the essential components of the composition, and should not otherwise unduly impair product stability, aesthetics or performance.

Pyridinethione salts

Pyridinethione anti-dandruff particulates, especially 1-hydroxy-2-pyridinethione salts, are highly preferred particulate anti-dandruff agents for use in compositions of the present invention. The concentration of pyridinethione anti-dandruff particulate typically ranges from about 0.1% to about 4%, by weight of the composition, preferably from about 0.1% to about 3%, more preferably from about 0.3% to about 2%. Preferred pyridinethione salts include those formed from heavy metals such as zinc, tin, cadmium, magnesium, aluminum and zirconium, preferably zinc, more preferably the zinc salt of 1-hydroxy-2-pyridinethione (known as "zinc pyridinethione" or "ZPT"), more preferably 1-hydroxy-2-pyridinethione salts in platelet particle form, wherein the particles have an average size of up to about 20 μ , preferably up to about 5 μ , more preferably up to about 2.5 μ . Salts formed from other cations, such as sodium, may also be suitable. Pyridinethione anti-dandruff agents are described, for example, in U.S. Pat. No. 2,809,971; U.S. Pat. No. 3,236,733; U.S. Pat. No. 3,753,196; U.S. Pat. No. 3,761,418; U.S. Pat. No. 4,345,080; U.S. Pat. No. 4,323,683; U.S. Pat. No. 4,379,753; and U.S. Pat. No. 4,470,982.

It is contemplated that when ZPT is used as the anti-dandruff particulate in the compositions herein, that the growth or re-growth of hair may be stimulated or regulated, or both, or that hair loss may be reduced or inhibited, or that hair may appear thicker or fuller.

Other Anti-microbial Actives - In addition to the anti-dandruff active selected from polyvalent metal salts of pyrithione, the present invention may further comprise one or more anti-fungal or anti-microbial actives in addition to the metal pyrithione salt actives. Suitable anti-microbial actives include coal tar, sulfur, whitfield's ointment, castellani's paint, aluminum chloride, gentian violet, octopirox (piroctone olamine), ciclopirox olamine, undecylenic acid and its metal salts, potassium permanganate, selenium sulphide, sodium thiosulfate, propylene glycol, oil of bitter orange, urea preparations, griseofulvin, 8-Hydroxyquinoline ciloquinol, thiobendazole, thiocarbamates, haloprogin, polyenes, hydroxypyridone, morpholine, benzylamine, allylamines (such as terbinafine), tea tree oil, clove leaf oil, coriander, palmarosa, berberine, thyme red, cinnamon oil, cinnamic aldehyde, citronellic acid, hinokitol, ichthyol pale, Sensiva SC-50, Elestab HP-100, azelaic acid, lyticase, iodopropynyl butylcarbamate (IPBC), isothiazalinones such as octyl isothiazalinone and azoles, and combinations thereof. Preferred anti-microbials include itraconazole, ketoconazole, selenium sulphide and coal tar.

Azoles

Azole anti-microbials include imidazoles such as benzimidazole, benzothiazole, bifonazole, butaconazole nitrate, climbazole, clotrimazole, croconazole, eberconazole, econazole, elubiol, fenticonazole, fluconazole, flutimazole, isoconazole, ketoconazole, lanoconazole, metronidazole, miconazole, neticonazole, omoconazole, oxiconazole nitrate, sertaconazole, sulconazole nitrate, tioconazole, thiazole, and triazoles such as terconazole and itraconazole, and combinations thereof. When present in the composition, the azole anti-microbial active is included in an amount from about 0.01% to about 5%, preferably from about 0.1% to about 3%, and more preferably from about 0.3% to about 2%, by weight of the composition. Especially preferred herein is ketoconazole.

Selenium Sulfide

Selenium sulfide is a particulate anti-dandruff agent suitable for use in the anti-microbial compositions of the present invention, effective concentrations of which range from about 0.1% to about 4%, by weight of the composition, preferably from about 0.3% to about 2.5%, more preferably from about 0.5% to about 1.5%. Selenium sulfide is generally regarded as a compound having one mole of selenium and two moles of sulfur, although it may also be a cyclic structure that conforms to the general formula Se_xS_y , wherein $x + y = 8$. Average particle diameters for the selenium sulfide are typically less than $15\mu\text{m}$, as measured by forward laser

light scattering device (e.g. Malvern 3600 instrument), preferably less than 10 μm . Selenium sulfide compounds are described, for example, in U.S. Pat. No. 2,694,668; U.S. Pat. No. 3,152,046; U.S. Pat. No. 4,089,945; and U.S. Pat. No. 4,885,107.

Sulfur

Sulfur may also be used as a particulate anti-microbial/anti-dandruff agent in the anti-microbial compositions of the present invention. Effective concentrations of the particulate sulfur are typically from about 1% to about 4%, by weight of the composition, preferably from about 2% to about 4%.

Keratolytic Agents

The present invention may further comprise one or more keratolytic agents such as Salicylic Acid.

Additional anti-microbial actives of the present invention may include extracts of melaleuca (tea tree) and charcoal. The present invention may also comprise combinations of anti-microbial actives. Such combinations may include octopirox and zinc pyrithione combinations, pine tar and sulfur combinations, salicylic acid and zinc pyrithione combinations, octopirox and climbazole combinations, and salicylic acid and octopirox combinations, and mixtures thereof.

Humectant

The compositions of the present invention may contain a humectant. The humectants herein are selected from the group consisting of polyhydric alcohols, water soluble alkoxyated nonionic polymers, and mixtures thereof. The humectants, when used herein, are preferably used at levels by weight of the composition of from about 0.1% to about 20%, more preferably from about 0.5% to about 5%.

Polyhydric alcohols useful herein include glycerin, sorbitol, propylene glycol, butylene glycol, hexylene glycol, ethoxylated glucose, 1, 2-hexane diol, hexanetriol, dipropylene glycol, erythritol, trehalose, diglycerin, xylitol, maltitol, maltose, glucose, fructose, sodium chondroitin sulfate, sodium hyaluronate, sodium adenosine phosphate, sodium lactate, pyrrolidone carbonate, glucosamine, cyclodextrin, and mixtures thereof.

Water soluble alkoxyated nonionic polymers useful herein include polyethylene glycols and polypropylene glycols having a molecular weight of up to about 1000 such as those with CTA names PEG-200, PEG-400, PEG-600, PEG-1000, and mixtures thereof.

Suspending Agent

The compositions of the present invention may further comprise a suspending agent at concentrations effective for suspending water-insoluble material in dispersed form in the compositions or for modifying the viscosity of the composition. Such concentrations range from

about 0.1% to about 10%, preferably from about 0.3% to about 5.0%, by weight of the compositions.

Suspending agents useful herein include anionic polymers and nonionic polymers. Useful herein are vinyl polymers such as cross linked acrylic acid polymers with the CTFA name Carbomer, cellulose derivatives and modified cellulose polymers such as methyl cellulose, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, nitro cellulose, sodium cellulose sulfate, sodium carboxymethyl cellulose, crystalline cellulose, cellulose powder, polyvinylpyrrolidone, polyvinyl alcohol, guar gum, hydroxypropyl guar gum, xanthan gum, arabia gum, tragacanth, galactan, carob gum, guar gum, karaya gum, carragheenin, pectin, agar, quince seed (*Cydonia oblonga* Mill), starch (rice, corn, potato, wheat), algae colloids (algae extract), microbiological polymers such as dextran, succinoglucan, pulleran, starch-based polymers such as carboxymethyl starch, methylhydroxypropyl starch, alginic acid-based polymers such as sodium alginate, alginic acid propylene glycol esters, acrylate polymers such as sodium polyacrylate, polyethylacrylate, polyacrylamide, polyethyleneimine, and inorganic water soluble material such as bentonite, aluminum magnesium silicate, laponite, hectonite, and anhydrous silicic acid.

Commercially available viscosity modifiers highly useful herein include Carbomers with tradenames Carbopol 934, Carbopol 940, Carbopol 950, Carbopol 980, and Carbopol 981, all available from B. F. Goodrich Company, acrylates/stearate-20 methacrylate copolymer with tradename ACRY SOL 22 available from Rohm and Hass, nonoxynyl hydroxyethylcellulose with tradename AMERCELL POLYMER HM-1500 available from Amerchol, methylcellulose with tradename BENECEL, hydroxyethyl cellulose with tradename NATROSOL, hydroxypropyl cellulose with tradename KLUCEL, cetyl hydroxyethyl cellulose with tradename POLYSURF 67, all supplied by Hercules, ethylene oxide and/or propylene oxide based polymers with tradenames CARBOWAX PEGs, POLYOX WASRs, and UCON FLUIDS, all supplied by Amerchol.

Other optional suspending agents include crystalline suspending agents which can be categorized as acyl derivatives, long chain amine oxides, and mixtures thereof. These suspending agents are described in U.S. Pat. No. 4,741,855.

These preferred suspending agents include ethylene glycol esters of fatty acids preferably having from about 16 to about 22 carbon atoms. More preferred are the ethylene glycol stearates, both mono and distearate, but particularly the distearate containing less than about 7% of the mono stearate. Other suitable suspending agents include alkanol amides of fatty acids, preferably having from about 16 to about 22 carbon atoms, more preferably about 16 to 18 carbon atoms, preferred examples of which include stearic monoethanolamide, stearic diethanolamide, stearic monoisopropanolamide and stearic monoethanolamide stearate. Other long chain acyl

derivatives include long chain esters of long chain fatty acids (e.g., stearyl stearate, cetyl palmitate, etc.); long chain esters of long chain alkanol amides (e.g., stearamide diethanolamide distearate, stearamide monoethanolamide stearate); and glyceryl esters (e.g., glyceryl distearate, trihydroxystearin, tribehenin) a commercial example of which is Thixin R available from Rheox, Inc. Long chain acyl derivatives, ethylene glycol esters of long chain carboxylic acids, long chain amine oxides, and alkanol amides of long chain carboxylic acids in addition to the preferred materials listed above may be used as suspending agents.

Other long chain acyl derivatives suitable for use as suspending agents include N,N-dihydrocarbyl amido benzoic acid and soluble salts thereof (e.g., Na, K), particularly N,N-di(hydrogenated) C.sub.16, C.sub.18 and tallow amido benzoic acid species of this family, which are commercially available from Stepan Company (Northfield, Ill., USA).

Examples of suitable long chain amine oxides for use as suspending agents include alkyl dimethyl amine oxides, e.g., stearyl dimethyl amine oxide.

Other suitable suspending agents include primary amines having a fatty alkyl moiety having at least about 16 carbon atoms, examples of which include palmitamine or stearamine, and secondary amines having two fatty alkyl moieties each having at least about 12 carbon atoms, examples of which include dipalmitoylamine or di(hydrogenated tallow)amine. Still other suitable suspending agents include di(hydrogenated tallow)phthalic acid amide, and crosslinked maleic anhydride-methyl vinyl ether copolymer.

Other Optional Components

The compositions of the present invention may contain also vitamins and amino acids such as: water soluble vitamins such as vitamin B1, B2, B6, B12, C, pantothenic acid, pantothenyl ethyl ether, panthenol, biotin, and their derivatives, water soluble amino acids such as asparagine, alanin, indole, glutamic acid and their salts, water insoluble vitamins such as vitamin A, D, E, and their derivatives, water insoluble amino acids such as tyrosine, tryptamine, and their salts.

The compositions of the present invention may also contain pigment materials such as inorganic, nitroso, monoazo, disazo, carotenoid, triphenyl methane, triaryl methane, xanthene, quinoline, oxazine, azine, anthraquinone, indigoid, thionindigoid, quinacridone, phthalocianine, botanical, natural colors, including: water soluble components such as those having C. I. Names.

The compositions of the present invention may also contain antimicrobial agents which are useful as cosmetic biocides and antidandruff agents including: water soluble components such as piroctone olamine, water insoluble components such as 3,4,4'- trichlorocarbanilide (trichlosan), triclocarban and zinc pyrithione.

The compositions of the present invention may also contain chelating agents.

METHOD OF USE

The shampoo compositions of the present invention are used in a conventional manner for cleansing hair or skin and providing enhanced deposition of silicone and other benefits of the present invention. An effective amount of the composition for cleansing the hair or skin is applied to the hair or skin that has preferably been wetted with water, and then rinsed off. Such effective amounts generally range from about 1g to about 50g, preferably from about 1g to about 20g. Application to the hair typically includes working the composition through the hair such that most or all of the hair is contacted with the composition. These steps can be repeated as many times as desired to achieve the desired cleansing and particle deposition benefits.

The silicones of the present invention may also be useful in a hair conditioning composition, which may require no deterative surfactant or a lower level of deterative surfactant than that needed for a shampoo composition.

The following examples further describe and demonstrate the preferred embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration, and are not to be construed as limitations of the present invention since many variations thereof are possible without departing from its scope.

Examples

The shampoo compositions illustrated in the following Examples illustrate specific embodiments of the shampoo compositions of the present invention, but are not intended to be limiting thereof. Other modifications can be undertaken by the skilled artisan without departing from the spirit and scope of this invention. These exemplified embodiments of the shampoo composition of the present invention provide enhanced conditioning benefits to the hair.

The shampoo compositions illustrated in the following Examples are prepared by conventional formulation and mixing methods, an example of which is set forth hereinbelow. All exemplified amounts are listed as weight percents and exclude minor materials such as diluents, preservatives, color solutions, imagery ingredients, botanicals, and so forth, unless otherwise specified.

The compositions illustrated in the examples were prepared in the following manner (all percentages are based on weight unless otherwise specified).

For each of the compositions, 6-9% of ammonium laureth-3 sulfate, P43 oil, PureSyn6 oil, cationic polymers, 0-1.5% Ammonium Xylene Sulfonate, and 0-5% water was added to a jacketed mix tank and heated to about 74°C with agitation to form a solution. Citric Acid, Sodium Citrate, Sodium Benzoate, Disodium EDTA, Cocamide MEA and 0.6-0.9% Cetyl alcohol, were added to the tank and allowed to disperse. Ethylene glycol distearate (EGDS) was then added to the mixing vessel, and melted. After the EGDS was well dispersed (after about 10 minutes) preservative was

added and mixed into the surfactant solution. This mixture was passed through a heat exchanger where it was cooled to about 35°C and collected in a finishing tank. As a result of this cooling step, the ethylene glycol distearate crystallized to form a crystalline network in the product.

Separately about 20% of the water was heated to about 74°C and the remainder of the Cetyl Alcohol, Stearyl Alcohol, and the Cationic Surfactant were added to it. After incorporation, this mixture was passed through a heat exchanger where it was cooled to about 35°C. As a result of this cooling step, the Fatty Alcohols and surfactant crystallized to form a crystalline gel network.

These two premixes are the mixed together and the remainder of the surfactants, perfume, Dimethicone, Sodium Chloride or Ammonium Xylene Sulfonate for viscosity adjustment and the remainder of the water were added with ample agitation to insure a homogeneous mixture.

Preferred viscosities range from about 5000 to about 9000 centipoise at 27°C (as measured by a Wells-Brookfield model RVTDCP viscometer using a CP-41 cone and plate at 2/s at 3 minutes).

EXAMPLE COMPOSITION	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Ammonium Laureth-3 Sulfate	10.00	10.00	10.00	10.00	10.00	11.70	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00
Ammonium Lauryl Sulfate	6.00	6.00	6.00	2.00	2.00	2.30	2.00	2.00	2.00	6.00	2.00	2.00	2.00	2.00
Cocamidopropyl Betaine FB				2.00			2.00	2.00	2.00		2.00	2.00	2.00	2.00
Sodium Lauraoamphoacetate				2.00			2.00	2.00	2.00		2.00	2.00	2.00	2.00
Cocaminopropionic Acid					4.00	2.00								
Cetyl Alcohol	1.600	1.300	1.300	1.300	1.300	1.300	1.300	1.300	1.300	1.600	1.30	1.300	1.300	1.300
Stearyl Alcohol	1.25	1.25	1.25	1.25	1.25	1.25	1.00	1.00	1.00	1.25	1.25	1.00	1.00	1.00
Cocamide MEA	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80	0.80
Polyquat 10 (1)	0.50	0.50		0.50	0.50	0.50	0.50	0.50	0.50		0.50	0.50	0.50	0.50
Guar Hydroxypropyl trimonium Chloride (2)										0.50				
Polymethacrylamidopropyl trimonium Chloride (3)			0.50											
Ethylene Glycol Distearate	1.50	1.50	0.75	1.50	1.50	1.50	0.75	1.50	1.50	1.50	1.50	0.75	0.75	0.75
Dimethicone (4)	2.00	2.00	2.00							2.00				
Dimethicone (5)				1.00	2.00	2.00	2.00	2.00	2.00			2.00	2.00	2.00
Dimethicone (11)											2.00			
Behentrimonium Chloride(6)	0.55	0.55				0.55	0.50			0.55	0.55			
Cetrimonium Chloride (7)			0.60					0.50						
Distearyldimonium Chloride (8)				0.50	0.60				0.50					
Oleamidopropyl dimethylamine												0.55		
Isostearylamidopropyl dimethylamine													.55	
Dicetyl dimethyl ammonium chloride														.55
Mobil P43 Synthetic Oil (9)	0.10									0.10				
Puresyn 6 (MCP-1812) (10)	0.40			0.25		0.25	0.25		0.25	0.40	0.25	0.25	0.25	0.25
Perfume Solution	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70	0.70
Sodium Citrate	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40
Citric Acid	0.04	0.40	0.40	0.40	0.04	0.04	0.40	0.04	0.04	0.04	0.40	0.40	0.40	0.40
Ammonium Xylene Sulsonate				1.50			1.50	1.50	1.50		1.50	1.50	1.50	1.50
Sodium Chloride	0.50	0.50	0.50		1.50	1.00				0.75				
Water and Minors (QS to 100%)														

- (1) Polymer KG30M available from Amerchol/Dow Chemical
- (2) Jaguar C17 available from Rhodia
- (3) Polycare 133 available from Rhodia
- (4) Viscasil 330M available from General Electric Silicones
- (5) DC1664 available from Dow Corning Silicones
- (6) Varisoft BT-85 available from Degussa
- (7) Varisoft 300 available from Degussa
- (8) Varisoft TA 100 available from Degussa
- (9) P43 oil available from Exxon/Mobil Chemical
- (10) Puresyn 6 available from Exxon/Mobil Chemical
- (11) DC 2-1865 available from Dow Corning

The fatty alcohol deposition of these products is measured by treating a switch of hair with 3 cycles (2 lather/rinse steps per cycle, 0.1gm shampoo/gm hair on each lather/rinse step) with the shampoo. Four switches were treated with each shampoo. The switches were then extracted with solvent and the level of adsorbed fatty alcohol measured by gas chromatographic-mass spectrophotometric analysis of the extracts.

In accordance with one embodiment of the present invention, the shampoo composition deposits at least 200 parts per million of fatty alcohol as measured by the following method: a) treat four individual switches of hair with 3 cycles each of the shampoo composition, wherein a "cycle" involves lathering the switch with 0.1 grams of shampoo per gram of hair, rinsing, lathering the switch with another 0.1 grams of shampoo per gram of hair, and rinsing again; b) extract the switches with solvent; and c) measure the level of adsorbed fatty alcohol by gas chromatographic-mass spectrophotometric analysis of the extracts. In one embodiment, the composition deposits at least 400 parts per million of fatty alcohol, preferably at least 600 parts per million of fatty alcohol.

In accordance with one embodiment of the present invention, the shampoo composition deposits at least 20 parts per million of cationic surfactant as measured by the following method: a) treat four individual switches of hair with 3 cycles each of the shampoo composition, wherein a "cycle" involves lathering the switch with 0.1 grams of shampoo per gram of hair, rinsing, lathering the switch with another 0.1 grams of shampoo per gram of hair, and rinsing again; b) extract the switches with solvent; and c) measure the level of adsorbed cationic surfactant by gas chromatographic-mass spectrophotometric analysis of the extracts. In one embodiment, the composition deposits at least 50 parts per million, preferably at least 100 parts per million, of cationic surfactant.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to one skilled in the art without departing from the scope of the present invention.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A shampoo composition comprising:
 - a) from about 5 to about 50 weight percent of a deterative surfactant,
 - b) a fatty alcohol gel network comprising a surfactant and a fatty alcohol, wherein said fatty alcohol gel network is a solid crystalline gel network, wherein said fatty alcohol is present in an amount of at least 0.05 weight percent of the composition, and wherein said fatty alcohol and said surfactant are present in a weight ratio of 1:1 to 40:1, and
 - c) at least 20.0 weight percent of an aqueous carrier.
2. The composition of Claim 1, further comprising a deposition aid.
3. The composition of Claim 2, wherein said deposition aid is a cationic polymer.
4. The composition of Claim 3, wherein said cationic polymer has a molecular weight of from about 10,000 to about 10,000,000 and a charge density from about 0.9 meq/gm to about 7.0 meq/gm.
5. The composition of any one of Claims 1 to 4, wherein said fatty alcohol is present in an amount from about 0.05 weight percent to about 14 weight percent.
6. The composition of any one of Claims 1 to 4, wherein said fatty alcohol is present in an amount from about 1 weight percent to about 10 weight percent.
7. The composition of any one of Claims 1 to 6, wherein said fatty alcohol gel network is formed using a cationic surfactant.
8. The composition of Claim 7, wherein said cationic surfactant is selected from the group consisting of quaternium-8, quaternium-14, quaternium-18, quaternium-18 methosulfate, quaternium-24, behenyltrimethylammonium chloride, and mixtures thereof.

9. A method of treating hair by administering a safe and effective amount of the composition according to any one of Claims 1 to 8.

10. A process for preparing a cleansing composition comprising:

a) combining a fatty alcohol and a surfactant in a premix at a temperature sufficient to allow partitioning of the surfactant into the fatty alcohol, wherein the premix has a chain melt temperature,

b) cooling the mixture below the chain melt temperature of the premix to form a gel network, and

c) adding the gel network to a deterative surfactant and an aqueous carrier to form a cleansing composition,

wherein the gel network comprises said fatty alcohol and said surfactant in a weight ratio of 1:1 to 40:1.

11. The process of Claim 10, wherein the surfactant of step a) is quaternium-8, quaternium-14, quaternium-18, quaternium-18 methosulfate, quaternium-24, behenyltrimethylammonium chloride, or mixtures thereof.

12. A shampoo composition prepared by the process of Claim 10.

13. The shampoo composition of Claim 1, wherein said composition deposits at least 200 parts per million of fatty alcohol as measured by the following method:

a) treat four individual switches of hair with 3 cycles each of said shampoo composition, wherein a "cycle" involves lathering the switch with 0.1 grams of shampoo per gram of hair, rinsing, lathering the switch with another 0.1 grams of shampoo per gram of hair, and rinsing again;

b) extract the switches with solvent; and

c) measure the level of adsorbed fatty alcohol by gas chromatographic-mass spectrophotometric analysis of the extracts.

14. The shampoo composition of Claim 13, wherein said composition deposits at least 400 parts per million of fatty alcohol.

15. The shampoo composition of Claim 13, wherein said composition deposits at least 600 parts per million of fatty alcohol.

16. The shampoo composition of Claim 7, wherein said composition deposits at least 20 parts per million of cationic surfactant as measured by the following method:

a) treat four individual switches of hair with 3 cycles each of said shampoo composition, wherein a "cycle" involves lathering the switch with 0.1 grams of shampoo per gram of hair, rinsing, lathering the switch with another 0.1 grams of shampoo per gram of hair, and rinsing again;

b) extract the switches with solvent; and

c) measure the level of adsorbed cationic surfactant by gas chromatographic-mass spectrophotometric analysis of the extracts.

17. The shampoo composition of Claim 16, wherein said composition deposits at least 50 parts per million of cationic surfactant.

18. The shampoo composition of Claim 16, wherein said composition deposits at least 100 parts per million of cationic surfactant.