



(86) Date de dépôt PCT/PCT Filing Date: 2000/04/28
(87) Date publication PCT/PCT Publication Date: 2000/11/09
(45) Date de délivrance/Issue Date: 2012/03/13
(85) Entrée phase nationale/National Entry: 2001/10/25
(86) N° demande PCT/PCT Application No.: US 2000/011473
(87) N° publication PCT/PCT Publication No.: 2000/066067
(30) Priorité/Priority: 1999/04/30 (US09/303,120)

(51) Cl.Int./Int.Cl. *C07K 7/00* (2006.01),
A61K 38/00 (2006.01), *A61K 39/00* (2006.01),
A61K 39/38 (2006.01), *C07H 21/02* (2006.01),
C07H 21/04 (2006.01), *C07K 14/33* (2006.01),
C07K 14/35 (2006.01), *C07K 16/00* (2006.01),
C07K 17/00 (2006.01), *C07K 5/00* (2006.01),
G01N 33/53 (2006.01)
(72) Inventeurs/Inventors:
BRAUN, JONATHAN, US;
SUTTON, CHRISTOPHER, US
(73) Propriétaire/Owner:
THE REGENTS OF THE UNIVERSITY OF CALIFORNIA,
US
(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : ANTIGENES MICROBIENS ASSOCIES AUX AFFECTIONS INTESTINALES INFLAMMATOIRES ET
PROCEDES POUR L'UTILISATION DE CES ANTIGENES
(54) Title: IBD-ASSOCIATED MICROBIAL ANTIGENS AND METHODS OF USING SAME

(57) **Abrégé/Abstract:**

The present invention provides nucleic acid and amino acid sequence of the novel I-1 and I-2 polypeptides, which are associated with human inflammatory bowel disease (IBD). Methods of diagnosing and treating inflammatory bowel disease using the IBD-associated I-1 and I-2 antigens also are provided.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number
WO 00/66067 A3

(51) International Patent Classification⁷: **G01N 33/53**,
A61K 39/00, 39/38, 38/00, C07K 16/00, 17/00, 5/00,
7/00, C07H 21/04, 21/02

(74) Agents: **GASHLER, Andrea, L.** et al.; Campbell & Flores LLP, 7th floor, 4370 La Jolla Village Drive, San Diego, CA 92122 (US).

(21) International Application Number: PCT/US00/11473

(22) International Filing Date: 28 April 2000 (28.04.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/303,120 30 April 1999 (30.04.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/303,120 (CIP)
Filed on 30 April 1999 (30.04.1999)

(71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th floor, Oakland, CA 94607 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BRAUN, Jonathan** [—/US]; 3924 Hilton Head Way, Tarzana, CA 91356 (US).
SUTTON, Christopher [—/US]; 1625 Eolus Avenue, Encinitas, CA 92024 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (utility model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— With international search report.

(88) Date of publication of the international search report:
15 February 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IBD-ASSOCIATED MICROBIAL ANTIGENS AND METHODS OF USING SAME

(57) Abstract: The present invention provides nucleic acid and amino acid sequence of the novel I-1 and I-2 polypeptides, which are associated with human inflammatory bowel disease (IBD). Methods of diagnosing and treating inflammatory bowel disease using the IBD-associated I-1 and I-2 antigens also are provided.



WO 00/66067 A3

**IBD-ASSOCIATED MICROBIAL ANTIGENS AND
METHODS OF USING SAME**

ACKNOWLEDGMENT

This work was supported by grant number DK46763
5 awarded by the National Institutes of Health. The United
States government has certain rights in this invention.

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The invention relates generally to the fields
10 of immunology, microbiology and inflammatory bowel
disease and more specifically to the diagnosis and
treatment of inflammatory bowel disease using microbial
antigens.

BACKGROUND INFORMATION

15 Inflammatory bowel disease (IBD) is the
collective term used to describe two gastrointestinal
disorders of unknown etiology: Crohn's disease (CD) and
ulcerative colitis (UC). The course and prognosis of
IBD, which occurs world-wide and is reported to afflict
20 as many as two million people, varies widely. Onset of
IBD is predominantly in young adulthood with diarrhea,
abdominal pain, and fever the three most common
presenting symptoms. The diarrhea may range from mild to
severe, and anemia and weight loss are additional common
25 signs of IBD. Ten percent to fifteen percent of all
patients with IBD will require surgery over a ten year
period. In addition, patients with IBD are at increased
risk for the development of intestinal cancer. Reports
of an increased occurrence of psychological problems,

including anxiety and depression, are perhaps not surprising symptoms of what is often a debilitating disease that strikes people in the prime of life.

Unfortunately, the available therapies for inflammatory bowel disease are few, and both diagnosis and treatment have been hampered by a lack of knowledge regarding the etiology of the disease. What is clear, however, is that a combination of genetic factors, exogenous triggers and endogenous microflora can contribute to the immune-mediated damage to the intestinal mucosa seen in inflammatory bowel disease. In Crohn's disease, bacteria have been implicated in initiation and progression of the disease: the intestinal inflammation in Crohn's disease is notable for its frequent responsiveness to antibiotics and susceptibility to bacterial fecal flow. Common intestinal colonists and novel pathogens have been implicated in Crohn's by direct detection or by disease associated anti-microbial immune responses. Furthermore, in many genetically susceptible animal models of chronic colitis, lumenal micro-organisms are a necessary cofactor for disease; animals housed in a germ-free environment do not develop colitis. However, despite much direct and indirect evidence for a role for enteric microorganisms in Crohn's disease, the pathogenic organisms and antigens contributing to the immune dysregulation seen in this disease have not been identified.

Current diagnostic assays for Crohn's disease are unable to detect all patients with the disease. Thus, identification of novel microbial antigens associated with Crohn's disease would provide reagents that can increase the sensitivity of current diagnostic assays. In addition, such microbial antigens can bear a

disease related T-cell epitope and, as original or contributing inducers of the disease-related immune response, can be effective tolerogenic antigens for treating inflammatory bowel disease. Identification of IBD-associated microbial antigens also would facilitate isolation of the involved microbial species, paving the way for the discovery of new antibiotics or drugs for treating inflammatory bowel disease, such drugs ameliorating disease by eliminating the microbial inducers of disease.

Thus, there is a need for identification and isolation of microbial IBD-associated antigens for diagnosing and treating the many individuals suffering from inflammatory bowel disease. The present invention satisfies this need by providing the IBD-associated I-1 and I-2 microbial antigens. Related advantages are provided as well.

SUMMARY OF THE INVENTION

The present invention provides an isolated inflammatory bowel disease-associated I-2 polypeptide having substantially the same amino acid sequence as SEQ ID NO: 2. The invention also provides an isolated immunoreactive fragment of an I-2 polypeptide having substantially the same amino acid sequence as a portion of SEQ ID NO: 2. An isolated immunoreactive fragment of an I-2 polypeptide can have, for example, at least ten contiguous amino acids of SEQ ID NO: 2.

Also provided by the present invention is substantially purified antibody material that selectively binds an I-2 polypeptide having SEQ ID NO: 2. Such a substantially purified antibody material can be, for

example, substantially purified polyclonal or monoclonal antibody material.

The invention further provides an isolated nucleic acid molecule having a nucleic acid sequence
5 encoding substantially the same amino acid sequence as SEQ ID NO: 2. An isolated nucleic acid molecule of the invention can have, for example, the nucleic acid sequence SEQ ID NO: 1.

Also provided by the invention is a method of
10 diagnosing inflammatory bowel disease (IBD) in a subject. The method includes the steps of obtaining a sample from the subject; contacting the sample with an I-2 polypeptide, or immunoreactive fragment thereof, under conditions suitable to form a complex of the I-2
15 polypeptide, or the immunoreactive fragment thereof, and antibody to the I-2 polypeptide; and detecting the presence or absence of the complex, where the presence of the complex indicates that the subject has IBD. A method of the invention for diagnosing inflammatory bowel
20 disease can be useful, for example, for diagnosing Crohn's disease. In a method of the invention for diagnosing inflammatory bowel disease, the presence or absence of the complex can be detected, for example, with a detectable secondary antibody that has specificity for
25 a class determining portion of the antibody to the I-2 polypeptide.

Further provided by the invention is a method of inducing tolerance in a patient with inflammatory bowel disease by administering an effective dose of an
30 I-2 polypeptide, or tolerogenic fragment thereof, to the patient with IBD. The methods of the invention can be particularly useful for treating a patient having Crohn's

disease. In a method of the invention for inducing tolerance, the I-2 polypeptide to be administered can have, for example, the amino acid sequence of SEQ ID NO: 2.

5 The invention also provides a composition including an I-2 polypeptide having substantially the same amino acid sequence as SEQ ID NO: 2, or tolerogenic fragment thereof, combined with a tolerogizing molecule. In a composition of the invention, the I-2 polypeptide
10 can have, for example, the amino acid sequence SEQ ID NO: 2. A tolerogenic fragment useful in a composition of the invention can have, for example, at least ten contiguous amino acids of SEQ ID NO: 2.

15 The present invention also provides a method of identifying an agent useful in treating inflammatory bowel disease. The method includes the steps of obtaining a specimen of an enteric microbe from a patient with inflammatory bowel disease; isolating from the
20 specimen a microbial species that includes a nucleic acid molecule encoding an I-2 polypeptide; contacting the microbial species with an agent; and assaying for reduced growth or viability of the microbial species as compared to the growth or viability in the absence of the agent,
25 where the reduced growth or viability of the microbial species indicates that the agent is an agent useful in treating inflammatory bowel disease. A method of the invention can be useful, for example, for identifying an agent for treating Crohn's disease. The methods of the
30 invention can be particularly useful for screening agents which are antibiotics.

The invention additionally provides a method of identifying an agent useful in treating inflammatory bowel disease using a novel animal model. The method includes the steps of administering an I-2 polypeptide to
5 a non-human animal, whereby one or more symptoms of IBD are exhibited; administering an agent to the non-human animal; and assaying the level of the one or more symptoms characteristic of IBD, where a reduction in the level of the one or more symptoms as compared to a
10 control level indicates that the agent is an agent useful in treating IBD. The methods of the invention can be applied, for example, to identification of agents useful in treating Crohn's disease. The I-2 polypeptide administered can have, for example, the amino acid
15 sequence SEQ ID NO: 2. A non-human animal particularly useful in the methods of the invention can be, for example, a mouse deficient in *Gxi2*, *TCR α* or *IL-10*.

BRIEF DESCRIPTION OF THE DRAWINGS

20 Figure 1 shows the I-1 and I-2 nucleic acid sequences and encoded polypeptides. A. Shown are the I-2 nucleic acid sequence (SEQ ID NO: 1) and predicted amino acid sequence (SEQ ID NO: 2). B. Shown are the I-1 nucleic acid sequence (SEQ ID NO: 3) and I-1 open
25 reading frame 1 (top; SEQ ID NO: 4) and open reading frame 2 (bottom; SEQ ID NO: 5).

Figure 2 shows an alignment of the I-2 amino acid sequence SEQ ID NO: 2 with homologous sequences. Sequences were identified by tBLASTx search against the
30 non-redundant database. Alignments were generated by CLUSTALW, with modifications by Genedoc. "Clostridium" is the predicted protein 4 from *Clostridium pasteurianum* (SEQ ID NO: 6; accession number 481591); "Mycobacterium"

is the predicted protein Rv3557c from *Mycobacterium tuberculosis* (SEQ ID NO: 7; accession number 1877288); and "Aquifex" is a transcriptional regulator from *Aquifex aeolicus* (SEQ ID NO: 8; accession number 2984362).

5 Residues identical or conservatively substituted among four polypeptides are shown in black; residues identical or conservatively substituted among three polypeptides are shown in dark gray; and residues identical or conservatively substituted between two polypeptides are
10 shown in light gray. Consensus residues are shown beneath the alignment, with one-letter codes indicating conserved amino acids; "6" indicating a conserved non-polar (hydrophobic) residue such as leucine, isoleucine, alanine, valine or methionine; and "5"
15 indicating the conserved aromatic residue phenylalanine or tyrosine.

Figure 3 shows the results of PCR analysis with I-2 (SEQ ID NO: 1) specific primers using paraffin embedded colonic samples from CD patients, UC patients
20 and control individuals. "N" designates the number of samples assayed, and "N+" designates the number of positive samples. "Ca" designates cancer resection samples; "Divertic" designates diverticulitis samples; "append" designates appendicitis samples; and "I-col "
25 designates ischemic colitis.

Figure 4 shows IgG and IgA reactivities to the I-2 polypeptide (SEQ ID NO: 2) in various populations. A. Sera from Crohn's disease patients and normal individuals was assayed by ELISA at the indicated
30 dilutions for IgG reactivity to the GST-I-2 fusion polypeptide (SEQ ID NO: 2). The absorbance values were calculated by subtracting absorbance to GST alone.

B. Sera from Crohn's disease, ulcerative colitis and normal individuals was assayed by ELISA for IgA reactivity to the GST-I-2 fusion polypeptide (SEQ ID NO: 2). The absorbance values were calculated by subtracting absorbance to GST alone. The cut-off (dotted line) was set as two standard deviations above the mean value for the normal population.

Figure 5 shows T cell proliferation assays using CD4⁺ T cells derived from normal C57BL/6J mice. T cells were stimulated with the indicated concentration of GST-I-2 fusion protein, shown on the Y axis. The X axis represents counts of incorporated [³H]-thymidine. SEB represents T cells stimulated with the positive control *Staphylococcus aureus* enterotoxin B.

Figure 6 shows T cell cytokine expression in response to challenge with the GST-I-2 fusion protein.

Figure 7 shows a variety of conditions for culturing microbial organisms isolated from IBD patients.

Figure 8 shows activation of naive CD4⁺ T cells upon I-2 stimulation. (A) [³H]-thymidine proliferation assay of native CD4⁺ T cells. (B) [³H]-thymidine proliferation assay of memory CD4⁺ T cells.

Figure 9 shows that recombinant I-2 is a superantigen. (A) Splenic C57 BL/6J CD4⁺ T cells were cultured with antigen-presenting cells from C57 BL/6 that were pulsed with I-2-GST, or fixed with paraformaldehyde and then pulsed with I-2-GST overnight. (B) Pigeon cytochrome C (PPC) specific CD4⁺ T cells were cultured with syngenic antigen-presenting cells pulsed with PPC, or fixed with paraformaldehyde and then pulsed with PPC.

DETAILED DESCRIPTION OF THE INVENTION

The pathogenesis of inflammatory bowel disease, although poorly understood, ultimately involves immune-mediated tissue damage. Similar to autoimmune disorders such as diabetes mellitus and multiple sclerosis, inflammatory bowel disease is associated with various immunologic abnormalities and can represent a process of immune dysfunction. However, unlike the other disorders, inflammatory bowel disease occurs in a mucosal site interfacing with the intestinal lumen, and, therefore, a primary immune target in inflammatory bowel disease can be extrinsic agent such as a chronic microbial colonist. In this case, the mucosal injury characteristic of inflammatory bowel disease is a consequence of inflammatory bystander damage to resident parenchymal cells.

The present invention is directed to the exciting discovery that several microbial DNA sequences are found preferentially in involved Crohn's disease (CD) mucosa as compared to uninvolved mucosa. As disclosed herein, representational difference analysis (RDA), a PCR driven subtractive cloning approach for identifying DNA sequences found preferentially in an infected area, was used to isolate DNA sequences from a Crohn's disease patient that were differentially present in mononuclear cells from the lamina propria in an area with ulcerations as compared to an area macroscopically free of disease (see Example I). As disclosed herein, two IBD-associated sequences were of microbial origin. The nucleic acid and amino acid sequences, designated I-1 (SEQ ID NOS: 3 to 5) and I-2 (SEQ ID NOS: 1 and 2) are shown in Figure 1. As further disclosed herein, PCR analysis of colonic samples from CD, UC and non-IBD patients revealed that the I-2

sequence (SEQ ID NO: 1) was more often found in involved CD tissue than in UC or non-IBD samples (Example I and Figure 3). Thus, novel microbial sequences have been identified that are associated with inflammatory bowel disease, in particular, with inflamed CD lesions. Isolation of microbial sequences associated with IBD implicates microbes in the pathogenesis of IBD and provides valuable reagents for diagnosing or ameliorating inflammatory bowel disease.

Thus, the present invention provides an isolated inflammatory bowel disease associated I-2 polypeptide having substantially the same amino acid sequence as SEQ ID NO: 2. The invention also provides an isolated immunoreactive fragment of an I-2 polypeptide having substantially the same amino acid sequence as a portion of SEQ ID NO: 2. An isolated immunoreactive fragment of an I-2 polypeptide can have, for example, at least ten contiguous amino acids of SEQ ID NO: 2.

The term "isolated," as used herein in reference to a polypeptide means a polypeptide that is in a form that is relatively free from contaminating lipids, polypeptides, nucleic acids or other cellular material normally associated with the polypeptide in a cell.

As used herein, the term "I-2 polypeptide" means a polypeptide having substantially the same amino acid sequence as the microbial I-2 polypeptide (SEQ ID NO: 2) shown in Figure 1A. The microbial I-2 polypeptide (SEQ ID NO: 2) is a polypeptide of 100 amino acids sharing some similarity to bacterial transcriptional regulators, with the greatest similarity in the amino-terminal 30 amino acids. As illustrated in Figure 2, the I-2 polypeptide shares weak homology with

the predicted protein 4 from *C. pasteurianum* (SEQ ID NO: 6); Rv3557c from *Mycobacterium tuberculosis* (SEQ ID NO: 7); and a transcriptional regulator from *Aquifex aeolicus* (SEQ ID NO: 8). As disclosed in Example I, the I-2 encoding nucleic acid (SEQ ID NO: 1) is differentially present in involved Crohn's disease tissue, as compared to mucosa macroscopically free of disease.

An I-2 polypeptide having substantially the same amino acid sequence as SEQ ID NO: 2 can be the naturally occurring I-2 polypeptide (SEQ ID NO: 2) or a related polypeptide having substantial amino acid sequence similarity to this sequence. Such related polypeptides exhibit greater sequence similarity to the I-2 polypeptide SEQ ID NO: 2 than to the *C. pasteurianum* sequence SEQ ID NO: 6 and include isotype variants or homologs of the amino acid sequence shown in Figure 1A. As used herein, the term I-2 polypeptide generally describes polypeptides generally having an amino acid sequence with greater than about 50% identity, preferably greater than about 60% identity, more preferably greater than about 70% identity, and can be a polypeptide having greater than about 80%, 90%, 95%, 97%, or 99% amino acid sequence identity with SEQ ID NO: 2, said amino acid identity determined with CLUSTALW using the BLOSUM 62 matrix with default parameters. The *C. pasteurianum* protein 4 (SEQ ID NO: 6) has about 21% amino acid identity with the I-2 polypeptide SEQ ID NO: 2 and, therefore, is not an I-2 polypeptide as defined herein.

The I-1 nucleic acid sequence (SEQ ID NO: 3) contains two open reading frames (ORFs) on opposite strands. The top strand I-1 ORF shown in Figure 1B encodes a predicted polypeptide of 115 amino acids (SEQ

ID NO: 4) with homology to the prokaryotic transcription factor ptxR. The bottom strand ORF encodes a predicted polypeptide of 114 amino acids (SEQ ID NO: 5). The I-1 encoding nucleic acid (SEQ ID NO: 3) also was isolated by
5 RDA analysis as a sequence preferentially found in involved CD mucosa (see Example I).

An I-1 polypeptide can have substantially the same amino acid sequence as SEQ ID NO: 4 or SEQ ID NO: 5. Such an I-1 polypeptide can be a naturally occurring I-1
10 polypeptide (SEQ ID NOS: 4 or 5) or a related polypeptide, for example, an isotype variant or homologous sequence from a different bacterial species having substantial amino acid sequence similarity to one of these sequences. As used herein, the term I-1
15 polypeptide generally describes polypeptides generally having an amino acid sequence with greater than about 50% identity, preferably greater than about 60% identity, more preferably greater than about 70% identity, and can be a polypeptide having greater than about 80%, 90%, 95%,
20 97%, or 99% amino acid sequence identity with SEQ ID NOS: 4 or 5, said amino acid identity determined with CLUSTALW using the BLOSUM 62 matrix with default parameters.

25 As used herein, the term "substantially the same amino acid sequence," when used in reference to an I-1 or I-2 polypeptide, is intended to mean a sequence as shown in Figure 1A or Figure 1B, or a similar, non-identical sequence that is considered by those
30 skilled in the art to be a functionally equivalent amino acid sequence. For example, an amino acid sequence that has substantially the same amino acid sequence as an I-1 polypeptide (SEQ ID NOS: 4 or 5) or I-2 polypeptide (SEQ ID NO: 2) can have one or more modifications such as

amino acid additions, deletions or substitutions relative to the amino acid sequence of SEQ ID NO: 4 or 5, or SEQ ID NO: 2, respectively, provided that the modified polypeptide retains substantially at least one biological activity of I-1 or I-2 such as immunoreactivity or tolerogenic activity, described further below. Comparison for substantial similarity between amino acid sequences is usually performed with sequences between about 6 and 100 residues, preferably between about 10 and 100 residues and more preferably between about 25 and 35 residues.

Thus, it is understood that limited modifications can be made to an I-1 or I-2 polypeptide, or to an immunoreactive or tolerogenic fragment thereof, as described further below, without destroying its biological function. A modification of an I-1 or I-2 polypeptide that does not destroy immunoreactivity or a modification of an I-1 or I-2 polypeptide that does not destroy tolerogenic activity is encompassed within the meaning of the term I-1 polypeptide, or I-2 polypeptide, as used herein. A modification can be, for example, an addition, deletion, or substitution of one or more amino acid residues; substitution of a compound that mimics amino acid structure or function; or addition of chemical moieties such as amino or acetyl groups. The activity of a modified I-1 or I-2 polypeptide or fragment thereof can be assayed, for example, using one of the assays for immunoreactivity or tolerogenic activity disclosed herein (see below).

A particularly useful modification of a polypeptide of the invention, or fragment thereof, is a modification that confers, for example, increased stability. Incorporation of one or more D-amino acids is

a modification useful in increasing stability of a polypeptide or polypeptide fragment. Similarly, deletion or substitution of lysine can increase stability by protecting against degradation. For example, such a
5 substitution can increase stability and, thus, bioavailability of one of the polypeptide antigens disclosed herein.

The I-2 polypeptide, and fragments thereof, can
10 be useful to prepare substantially purified antibody material that selectively binds an I-2 polypeptide (SEQ ID NO: 2). The antibody material can be, for example, substantially purified polyclonal antiserum or monoclonal antibody material. The antibody material of the
15 invention be useful, for example, in determining the presence and location of I-2 polypeptide within the mucosa of afflicted patients and in diagnosing inflammatory bowel disease. The substantially purified antibody material of the invention can be useful, for
20 example, in an ELISA or immunohistopathological assay for diagnosing Crohn's disease.

As used herein, the term "antibody material" is used in its broadest sense to include polyclonal and monoclonal antibodies, as well as polypeptide fragments
25 of antibodies that retain a selective binding activity for an I-2 polypeptide of at least about $1 \times 10^5 \text{ M}^{-1}$. One skilled in the art would know that anti-I-2 antibody fragments such as Fab, F(ab')_2 and Fv fragments can retain selective binding activity for an I-2 polypeptide and,
30 thus, are included within the definition of an antibody. In addition, the term antibody material as used herein encompasses non-naturally occurring antibodies and fragments containing, at a minimum, one V_H and one V_L domain, such as chimeric antibodies, humanized

antibodies, single chain Fv fragments (scFv) that selectively bind an I-2 polypeptide. Such non-naturally occurring antibodies can be constructed using solid phase peptide synthesis, produced recombinantly or obtained, 5 for example, by screening combinatorial libraries consisting of variable heavy chains and variable light chains as described by Borrebaeck (Ed.), Antibody Engineering (Second edition) New York: Oxford University Press (1995), which is incorporated herein by reference.

10 Antibody material "selective for" an I-2 polypeptide, or that "selectively binds" an I-2 polypeptide, binds with substantially higher affinity to that polypeptide than to an unrelated polypeptide. The substantially purified antibody material of the invention 15 also can be specific for an I-2 polypeptide, whereby its binding affinity is significantly higher for an I-2 polypeptide than for related polypeptides such as SEQ ID NOS: 6 to 8.

 Anti-I-2 antibody material can be prepared, for 20 example, using an I-2 fusion protein or a synthetic peptide encoding a portion of the I-2 polypeptide (SEQ ID NO: 2) as an immunogen. One skilled in the art would know that purified I-2 polypeptide, which can be produced recombinantly, or fragments of I-2, including peptide 25 portions of I-2 such as synthetic peptides, can be used as an immunogen. Non-immunogenic fragments or synthetic peptides of I-2 can be made immunogenic by coupling the hapten to a carrier molecule such as bovine serum albumin (BSA) or keyhole limpet hemocyanin (KLH). In addition, 30 various other carrier molecules and methods for coupling a hapten to a carrier molecule are well known in the art as described, for example, by Harlow and Lane, Antibodies: A Laboratory Manual (Cold Spring Harbor

Laboratory Press, 1988).

The term "substantially purified," as used herein in reference to antibody material, means that the antibody material is substantially devoid of polypeptides, nucleic acids and other cellular material which with an antibody is normally associated in a cell. The claimed antibody material that selectively binds an I-2 polypeptide (SEQ ID NO: 2) further is substantially devoid of antibody material of unrelated specificities, i.e. that does not selectively bind an I-2 polypeptide. The antibody material of the invention can be prepared in substantially purified form, for example, by I-2 affinity purification of polyclonal anti-I-2 antisera, by screening phage displayed antibodies against the I-2 polypeptide (SEQ ID NO: 2), or as monoclonal antibodies prepared from hybridomas.

As disclosed herein, the I-2 sequence is present in mice housed under conventional conditions and induces a dose-dependent proliferative response as well as IL-10 secretion (Example V). As further disclosed herein, both naive and memory CD4⁺ T cell populations spontaneously proliferate and produce IL-10 in response to I-2 (see Figure 8). Furthermore, the I-2 response is dependent on MHC class II-mediated recognition but does not require antigen processing, as indicated by the ability of paraformaldehyde-fixed cells to induce an I-2 response (see Figure 9). These results demonstrate that I-2 is a novel superantigen and indicate that methods of blocking an I-2 superantigen response can prevent or ameliorate Crohn's disease.

Thus, the present invention provides a method of treating a patient with IBD by administering to the patient an effective dose of an agent that selectively binds an I-2 polypeptide having SEQ ID NO: 2. In a
5 method of the invention, a useful agent can be, for example, substantially purified antibody material, such as polyclonal or monoclonal antibody material, that selectively binds to an I-2 polypeptide having SEQ ID NO: 2. The methods of the invention are useful in
10 treating patients with IBD such as patients having Crohn's disease.

Further provided by the invention is a method of treating a patient with IBD by inhibiting a superantigen response induced by an I-2 polypeptide
15 having SEQ ID NO: 2. Such a superantigen response can be, for example, T cell proliferation or IL-10 secretion. A method of the invention can be particularly useful for inhibiting a superantigen response in a patient with Crohn's disease.

20 The term superantigen response as used herein refers to the response of a T cell to a superantigen and generally is a proliferative response. Superantigens, which are among the most potent T-cell mitogens known, bind residues in the V β domain of the T-cell receptor and
25 residues in class II MHC molecules outside of the antigen-binding cleft. Thus, a superantigen can act to cross-link a T cell to a class II MHC molecule in an antigen-independent manner and activate all T cells expressing T-cell receptors with a particular V β domain.

30 The term inhibiting, as used herein in reference to a superantigen response, refers to a partial or complete reduction of a specific response induced by

the I-2 superantigen having SEQ ID NO: 2. As disclosed herein in Example V, the I-2 superantigen results in a proliferative response as well as IL-10 secretion. Thus, a partial or complete reduction in the
5 proliferative response or IL-10 secretion can be used to identify an agent useful in a method of the invention.

The invention further provides an isolated nucleic acid molecule having a nucleic acid sequence encoding substantially the same amino acid sequence as
10 SEQ ID NO: 2. An isolated nucleic acid molecule of the invention can have, for example, the nucleic acid sequence SEQ ID NO: 1. These nucleic acid molecules are useful, for example, in producing recombinant polypeptides and as probes for detecting I-2 mRNA
15 expression. Nucleotide portions of SEQ ID NO: 1 also are useful, for example, as primers for PCR analysis (see Example I).

Isolated nucleic acid molecules of the
20 invention include, for example, nucleic acid molecules encoding I-2 polypeptide homologs, nucleic acid molecules that are related, but different and encode the polypeptide of SEQ ID NO: 2 due to the degeneracy of the genetic code, and nucleic acid molecules that are
25 related, but different and encode an I-2 polypeptide different from SEQ ID NO: 2 that exhibits immunoreactivity or tolerogenic activity.

The methods of the invention relate to the diagnosis and treatment of inflammatory bowel disease,
30 which is a designation that encompasses the broad categories of Crohn's disease and ulcerative colitis. Crohn's disease (regional enteritis) is a disease of chronic inflammation that can involve any part of the

gastrointestinal tract. Commonly the distal portion of the small intestine (ileum) and cecum are affected. In other cases, the disease is confined to the small intestine, colon or anorectal region. Crohn's disease
5 occasionally involves the duodenum and stomach, and more rarely the esophagus and oral cavity.

The variable clinical manifestations of Crohn's disease are, in part, a result of the varying anatomic localization of the disease. The most frequent symptoms
10 of CD are abdominal pain, diarrhea and recurrent fever. CD is commonly associated with intestinal obstruction or fistula, which is an abnormal passage between diseased loops of bowel, for example. Crohn's disease also includes complications such as inflammation of the eye,
15 joints and skin; liver disease; kidney stones or amyloidosis. In addition, CD is associated with an increased risk of intestinal cancer.

Several features are characteristic of the pathology of Crohn's disease. The inflammation
20 associated with CD, known as transmural inflammation, involves all layers of the bowel wall. Thickening and edema, for example, typically also appear throughout the bowel wall, with fibrosis also present in long-standing disease. The inflammation characteristic of CD also is
25 discontinuous in that segments of inflamed tissue, known as "skip lesions," are separated by apparently normal intestine. Furthermore, linear ulcerations, edema, and inflammation of the intervening tissue lead to a
"cobblestone" appearance of the intestinal mucosa, which
30 is distinctive of CD.

A hallmark of Crohn's disease is the presence of discrete aggregations of inflammatory cells, known as

granulomas, which are generally found in the submucosa. Some Crohn's disease cases display the typical discrete granulomas, while others show a diffuse granulomatous reaction or nonspecific transmural inflammation. As a
5 result, the presence of discrete granulomas is indicative of CD, although the absence of granulomas also is consistent with the disease. Thus, transmural or discontinuous inflammation, rather than the presence of granulomas, is a preferred diagnostic indicator of
10 Crohn's disease (Rubin and Farber, Pathology (Second Edition) Philadelphia: J.B. Lippincott Company (1994)).

Ulcerative colitis (UC) is a disease of the large intestine characterized by chronic diarrhea with
15 cramping abdominal pain, rectal bleeding, and loose discharges of blood, pus and mucus. The manifestations of ulcerative colitis vary widely. A pattern of exacerbations and remissions typifies the clinical course of most UC patients (70%), although continuous symptoms
20 without remission are present in some patients with UC. Local and systemic complications of UC include arthritis, eye inflammation such as uveitis, skin ulcers and liver disease. In addition, ulcerative colitis and especially long-standing, extensive disease is associated with an
25 increased risk of colon carcinoma.

Several pathologic features characterize UC in distinction to other inflammatory bowel diseases. Ulcerative colitis is a diffuse disease that usually extends from the most distal part of the rectum for a
30 variable distance proximally. The term left-sided colitis describes an inflammation that involves the distal portion of the colon, extending as far as the splenic flexure. Sparing of the rectum or involvement of

the right side (proximal portion) of the colon alone is unusual in ulcerative colitis. The inflammatory process of ulcerative colitis is limited to the colon and does not involve, for example, the small intestine, stomach or esophagus. In addition, ulcerative colitis is distinguished by a superficial inflammation of the mucosa that generally spares the deeper layers of the bowel wall. Crypt abscesses, in which degenerated intestinal crypts are filled with neutrophils, also are typical of ulcerative colitis (Rubin and Farber, *supra*, 1994).

In comparison with Crohn's disease, which is a patchy disease with frequent sparing of the rectum, ulcerative colitis is characterized by a continuous inflammation of the colon that usually is more severe distally than proximally. The inflammation in ulcerative colitis is superficial in that it is usually limited to the mucosal layer and is characterized by an acute inflammatory infiltrate with neutrophils and crypt abscesses. In contrast, Crohn's disease affects the entire thickness of the bowel wall with granulomas often, although not always, present. Disease that terminates at the ileocecal valve, or in the colon distal to it, is indicative of ulcerative colitis, while involvement of the terminal ileum, a cobblestone-like appearance, discrete ulcers or fistulas suggest Crohn's disease. Characteristics that serve to distinguish Crohn's disease from ulcerative colitis are summarized in Table 1 (Rubin and Farber, *supra*, 1994).

Table 1		
Characteristic Features of Crohn's disease and ulcerative colitis		
Feature	Crohn's Disease	Ulcerative Colitis
Macroscopic		
Thickened bowel wall	Typical	Uncommon
Luminal narrowing	Typical	Uncommon
"Skip" lesions	Common	Absent
Right colon predominance	Typical	Absent
Fissures and fistulas	Common	Absent
Circumscribed ulcers	Common	Absent
Confluent linear ulcers	Common	Absent
Pseudopolyps	Absent	Common
Microscopic		
Transmural inflammation	Typical	Uncommon
Submucosal fibrosis	Typical	Absent
Fissures	Typical	Rare
Granulomas	Common	Absent
Crypt abscesses	Uncommon	Typical

20 Certain immune-mediated disorders, including
 systemic lupus erythematosus, primary biliary cirrhosis
 and autoimmune hepatitis, are closely associated with
 distinctive patterns of autoantibody production.
 Disease-specific marker antibodies also have been
 25 observed in IBD. In the case of ulcerative colitis,
 anti-neutrophil cytoplasmic antibodies that produce a
 perinuclear staining pattern (pANCA), for example, upon
 indirect immunofluorescence microscopy of alcohol-fixed
 neutrophils, are elevated in 68-80% of UC patients and
 30 less frequently in other disorders of the colon. In

Crohn's disease, serum reactivity to the cell wall mannan polysaccharide of *Saccharomyces uvarum* (brewer's yeast) is a serologic marker for a majority of individuals with Crohn's disease (Sendid et al., Clin. Diag. Lab. Immunol., 3:219-226 (1996)).

As disclosed herein, ELISA analysis showed increased IgG serum reactivity to a GST-I-2 fusion polypeptide (SEQ ID NO: 2) in patients with Crohn's disease as compared to normal individuals (see Figure 4A). As shown in Figure 4B, 9 of 10 Crohn's disease patients had IgA serum reactivity to the GST-I-2 fusion polypeptide greater than two standard deviations above the mean value for a normal population. In contrast, no normal serum samples contained anti-I-2 IgA reactivity above this cutoff. The anti-I-2 serum IgA reactivity also was significantly higher on average in samples from Crohn's disease patients as compared to ulcerative colitis patients. These results indicate that reactivity to the I-2 polypeptide (SEQ ID NO: 2) can be used to differentiate Crohn's disease from normal individuals and those with UC.

Based on the above findings, the present invention provides methods of diagnosing inflammatory bowel disease (IBD) in a subject. The methods include the steps of obtaining a sample from the subject; contacting the sample with an I-2 polypeptide, or immunoreactive fragment thereof, under conditions suitable to form a complex of the I-2 polypeptide, or the immunoreactive fragment thereof, and antibody to the I-2 polypeptide; and detecting the presence or absence of the complex, where the presence of the complex indicates that the subject has IBD. A method of the invention for

diagnosing inflammatory bowel disease can be useful, for example, for diagnosing Crohn's disease. The presence or absence of the complex can be detected, for example, with a detectable secondary antibody that has specificity for a class determining portion of the antibody to the I-2 polypeptide.

As used herein, the term "subject" means any animal capable of having inflammatory bowel disease, including a human, non-human primate, rabbit, rat or mouse, especially a human. A subject generally has one or more symptoms of ulcerative colitis or Crohn's disease.

A sample useful in the methods of the invention can be obtained from any biological fluid having antibodies such as, for example, whole blood, plasma, saliva, or other bodily fluid or tissue, preferably serum. A sample to be assayed according to the methods of the invention can be obtained from any such subject.

As used herein, the term "complex" is synonymous with "immune complex" and means an aggregate of two or more molecules that results from specific binding between an antigen, such as a protein or peptide, and an antibody. In the methods of the invention, a complex is formed by specific binding of an I-2 polypeptide to an antibody.

In the methods of the invention, a complex can be detected with a detectable secondary antibody that has specificity for a class determining portion of the antibody to the I-2 polypeptide. Such a secondary antibody can be, for example, an anti-IgA secondary antibody, an anti-IgG secondary antibody, or a

combination of anti-IgA and anti-IgG secondary antibodies.

As used herein, the term "secondary antibody" means an antibody or combination of antibodies, which
5 binds an antibody that specifically binds an I-2 polypeptide having substantially the amino acid sequence SEQ ID NO: 2. One skilled in the art understands that, preferably, a secondary antibody does not compete with the I-2 antigen for binding to the primary antibody. A
10 secondary antibody can bind any epitope of the antibody that specifically binds the I-2 polypeptide. A particularly useful secondary antibody is an anti-IgA or anti-IgG antibody having specificity for the class determining portion of the primary antibody. A useful
15 secondary antibody is specific for the species from which the sample was obtained. For example, if human serum is the sample to be assayed, mouse anti-human IgA or IgG can be a useful secondary antibody. A combination of different antibodies, which can be useful in the methods
20 of the invention, also is encompassed within the meaning of the term secondary antibody, provided that at least one antibody of the combination reacts with an antibody that specifically binds an I-2 polypeptide.

As used herein, the term "class determining
25 portion," when used in reference to a secondary antibody, means the heavy chain constant-region sequence of an antibody that determines the isotype, such as IgA, IgD, IgE, IgG or IgM. Thus, a secondary antibody that has specificity for the class determining portion of an IgA
30 molecule, for example, binds IgA in preference to other antibody isotypes.

A secondary antibody useful in the invention can be obtained commercially or by techniques well known in the art. Such an antibody can be a polyclonal or, preferably, monoclonal antibody. For example, IgG
5 reactive polyclonal antibodies can be prepared using IgG or Fc fragments of IgG as an immunogen to stimulate the production of antibodies in the antisera of an animal such as a rabbit, goat, sheep or rodent, as described in Harlow and Lane, Antibodies: A Laboratory Manual New
10 York: Cold Spring Harbor Laboratory (1988). Monoclonal secondary antibodies, which are a population of antibody molecules that contain only one species of idiotope capable of binding a particular antigen epitope also can be produced by routine methods (see, for example, Harlow
15 and Lane, *supra*, 1988) or obtained commercially.

The term "detectable secondary antibody" means a secondary antibody, as defined above, that can be detected or measured by analytical methods. Thus, the
20 term secondary antibody includes an antibody labeled directly or indirectly with a detectable marker that can be detected or measured and used in a convenient assay such as an enzyme-linked immunosorbent assay, radioimmunoassay, radial immunodiffusion assay or Western
25 blotting assay. A secondary antibody can be labeled, for example, with an enzyme, radioisotope, fluorochrome or chemiluminescent marker. In addition, a secondary antibody can be rendered detectable using a biotin-avidin linkage such that a detectable marker is associated with
30 the secondary antibody. Labeling of the secondary antibody, however, should not impair binding of the secondary antibody to the I-2 polypeptide. If desired, a multiple antibody system can be used as the secondary antibody as discussed above. In such a system, at least

one of the antibodies is capable of binding the primary anti-I-2 antibody and at least one of the antibodies can be readily detected or measured by analytical methods.

A secondary antibody can be rendered detectable
5 by labeling with an enzyme such as horseradish peroxidase (HRP), alkaline phosphatase (AP), β -galactosidase or urease, for example. A horseradish-peroxidase detection system can be used, for example, with the chromogenic substrate tetramethylbenzidine (TMB), which yields a
10 soluble product in the presence of hydrogen peroxide that is detectable by measuring absorbance at 450 nm. An alkaline phosphatase detection system can be used with the chromogenic substrate *p*-nitrophenyl phosphate, for example, which yields a soluble product readily
15 detectable by measuring absorbance at 405 nm. Similarly, a β -galactosidase detection system can be used with the chromogenic substrate *o*-nitrophenyl- β -D-galactopyranoside (ONPG), which yields a soluble product detectable by measuring absorbance at 410 nm, or a urease detection
20 system can be used with a substrate such as urea-bromocresol purple (Sigma Immunochemicals, St. Louis, MO). A secondary antibody can be linked to an enzyme by methods well known in the art (Harlow and Lane, *supra*, 1988) or can be obtained from a number of
25 commercial sources. For example, goat F(ab')₂ anti-human IgG-alkaline phosphatase is a useful detectable secondary antibody that can be purchased from Jackson Immuno-Research (West Grove, PA).

A secondary antibody also can be rendered
30 detectable by labeling with a fluorochrome. Such a fluorochrome emits light of ultraviolet or visible wavelength after excitation by light or another energy source. DAPI, fluorescein, Hoechst 33258, R-phycocyanin,

B-phycoerythrin, R-phycoerythrin, rhodamine, Texas red or lissamine, for example, is a fluorochrome that can be linked to a secondary antibody and used to detect the presence or absence of a complex. A particularly useful
5 fluorochrome is fluorescein or rhodamine. Methods of conjugating and using these and other suitable fluorochromes are described, for example, in Van Vunakis and Langone, Methods in Enzymology, Volume 74, Part C (1991). A secondary antibody linked to a fluorochrome
10 also can be obtained from commercial sources. For example, goat F(ab')₂ anti-human IgG-FITC is available from Tago Immunologicals (Burlingame, CA).

A secondary antibody also can be labeled with a
15 chemiluminescent marker. Such a chemiluminescent secondary antibody is convenient for sensitive, non-radioactive detection of a complex containing an I-2 polypeptide and can be obtained commercially from various sources such as Amersham Lifesciences, Inc. (Arlington
20 Heights, IL).

A secondary antibody further can be rendered detectable by labeling with a radioisotope. An iodine-125 labeled secondary antibody is a particularly useful detectable secondary antibody (see, for example,
25 Harlow and Lane, *supra*, 1988).

A signal from a detectable secondary antibody can be analyzed, for example, using a spectrophotometer to detect color from a chromogenic substrate; a fluorometer to detect fluorescence in the presence of
30 light of a certain wavelength; or a radiation counter to detect radiation, such as a gamma counter for detection of iodine-125. For detection of an enzyme-linked

secondary antibody, for example, a quantitative analysis can be made using a spectrophotometer such as an EMAX Microplate Reader (Molecular Devices, Menlo Park, CA) in accordance with the manufacturer's instructions. If
5 desired, the assays of the invention can be automated or performed robotically, and the signal from multiple samples can be detected simultaneously.

The assays of the present invention can be forward, reverse or simultaneous as described in U.S.
10 Patent No. 4,376,110, issued March 8, 1983, to David et al. In the forward assay, each reagent is sequentially contacted with an I-2 polypeptide of the invention. If desired, separation of bound from unbound reagent can be performed before the addition of the next reagent. In a
15 reverse assay, all reagents are pre-mixed prior to contacting with I-2 polypeptide. A modified reverse assay is described in U.S. Patent No. 4,778,751 issued October 18, 1988, to El Shami et al. In a simultaneous assay, all reagents are separately but contemporaneously
20 contacted with an I-2 polypeptide of the invention. A reagent is any component useful in performing the assays of the present invention, for example, the sample, I-2 polypeptide, detectable secondary antibody, washing buffer or other solutions.

25 Separation steps for the various assay formats described herein, including the removal of unbound secondary antibody from the complex, can be performed by methods known in the art (Harlow and Lane, *supra*, 1988). For example, washing with a suitable buffer can be
30 followed by filtration, aspiration or magnetic separation. If the I-2 polypeptide or an immunoreactive

fragment thereof is immobilized on a particulate support, such as on microparticles, the particulate material can be centrifuged, if desired, followed by removal of wash liquid. If the I-2 polypeptide or an immunoreactive
5 fragment thereof is immobilized on a membrane, filter or well, a vacuum or liquid absorbing apparatus can be applied to the opposite side of the membrane, filter or well to draw the wash liquid away from the complex.

The invention also provides methods of
10 determining susceptibility to IBD in an individual by obtaining a sample from the individual; contacting the sample with an I-2 polypeptide, or immunoreactive fragment thereof, under conditions suitable to form a complex of the I-2 polypeptide, or the immunoreactive
15 fragment thereof, and antibody to the I-2 polypeptide; and detecting the presence or absence of the complex, where the presence of the complex indicates that the individual is susceptible to IBD.

The term "individual," as used herein, means
20 any animal capable of having inflammatory bowel disease, including a human, non-human primate, rabbit, rat or mouse, provided that the animal does not have inflammatory bowel disease as defined by the clinical, endoscopic and histopathologic parameters disclosed
25 herein. A sample to be assayed according to the methods of the invention can be obtained from any such individual.

As used herein, the term "increased susceptibility to IBD" as indicated by the presence of a
30 complex of I-2 polypeptide and antibody to I-2 polypeptide, means a reduced ability to resist IBD-causing factors, as compared with an individual from

whom a sample is obtained that does not form a complex when contacted with I-2 polypeptide or immunoreactive fragment thereof. Increased susceptibility to IBD in an individual does not mean the individual will necessarily
5 develop IBD. However, increased susceptibility to IBD in an individual is associated with an increased probability of having IBD in the future.

The term "immunoreactive fragment," as used in reference to an I-2 polypeptide, means a peptide or
10 polypeptide portion of an I-2 polypeptide that has immunoreactivity as defined by the ability of an anti-I-2 antibody-positive sample to form a complex with the I-2 polypeptide. Thus, the term "immunoreactive fragment of an I-2 polypeptide" as used herein, means a peptide or
15 polypeptide that has an amino acid sequence that is substantially the same as a portion of the amino acid sequence provided as SEQ ID NO: 2 and immunoreactivity as defined by the ability to form a complex with an anti-I-2 antibody-positive sample such as an I-2 reactive serum
20 sample from a Crohn's disease patient. In general, an immunoreactive fragment has from about three amino acids to the full-length of an I-2 polypeptide. An immunoreactive fragment of an I-2 polypeptide can have, for example, at least 5, 8, 10, 12, 15, 18, 20 or 25
25 amino acids. For example, an immunoreactive fragment of an I-2 polypeptide can be from five to fifty amino acids, from eight to fifty amino acids, or from ten to fifty amino acids. More preferably, an immunoreactive fragment has from eight to twenty amino acids or from ten to
30 twenty amino acids. Most preferably, an immunoreactive fragment has from twelve to twenty amino acids or from fifteen to twenty amino acids.

An immunoreactive fragment of an I-2 polypeptide can be identified by the ability to form a complex with an I-2 reactive sample, for example, an I-2 reactive CD patient serum sample. For example, an immunoreactive fragment of an I-2 polypeptide can be identified by its ability to form a complex when contacted with I-2 reactive CD sera. Assays for the formation of a complex between an antigen and anti-I-2 serum sample are disclosed herein. An ELISA assay can be particularly useful in identifying an immunoreactive fragment of an I-2 polypeptide (see Example IIB).

Identification of a microbial sequence associated with IBD implicates microbes in the pathogenesis of IBD and provides a valuable reagent for ameliorating inflammatory bowel disease. Furthermore, as disclosed herein, the I-2 polypeptide can elicit a T cell response, as demonstrated by the proliferation of murine T cells in response to a GST-I-2 fusion polypeptide (SEQ ID NO: 2), indicating that the I-2 polypeptide antigen can contribute to the etiology of inflammatory bowel disease. Based on identification and isolation of the IBD-associated I-2 sequence, there are provided methods of inducing tolerance in a patient with IBD as well as methods of preventing IBD in a healthy individual.

Thus, the present invention provides methods of inducing tolerance in a patient with inflammatory bowel disease by administering an effective dose of I-2 polypeptide, or tolerogenic fragment thereof, to the patient with IBD. The methods of the invention can be particularly useful for treating a patient having Crohn's disease. In a method of the invention for inducing tolerance, the I-2 polypeptide to be administered can

have, for example, the amino acid sequence of SEQ ID NO:
2.

As used herein, the term "patient with
inflammatory bowel disease" means a patient having
5 Crohn's disease or ulcerative colitis.

As used herein, the term "effective dose"
means the amount of an I-2 polypeptide, or a tolerogenic
fragment thereof, useful for inducing tolerance in a
patient with IBD. For induction of oral tolerance, for
10 example, multiple smaller oral doses can be administered
or a large dose can be administered. Such doses can be
extrapolated, for example, from the induction of
tolerance in animal models (see, for example, Trentham
et al., Science 261:1727-1730 (1993)).

15 An effective dose of an I-2 polypeptide or a
tolerogenic fragment thereof for inducing tolerance can
be administered by methods well known in the art. Oral
tolerance is well-recognized in the art as a method of
treating autoimmune disease (see, for example, Weiner,
20 Hospital Practice, pp. 53-58 (Sept. 15, 1995)). For
example, orally administered autoantigens suppress
several experimental autoimmune models in a disease- and
antigen-specific fashion; the diseases include
experimental autoimmune encephalomyelitis, uveitis, and
25 myasthenia, collagen- and adjuvant-induced arthritis,
and diabetes in the NOD mouse (see, for example, Weiner
et al., Ann. Rev. Immunol. 12:809-837 (1994)).
Furthermore, clinical trials of oral tolerance have
produced positive results in treating multiple
30 sclerosis, rheumatoid arthritis and uveitis. In

addition, parenteral administration of an I-2 polypeptide, or a tolerogenic fragment thereof, can be used to induce tolerance. Subcutaneous injection, for example, can be used to deliver an I-2 polypeptide, or a tolerogenic fragment thereof, to an IBD patient, for example, a patient having Crohn's disease (Johnson, Ann. Neurology 36(suppl.):S115-S117 (1994), which is incorporated herein by reference).

The term "tolerogenic fragment," as used in reference to an I-2 polypeptide of the invention, means a peptide or polypeptide portion of the polypeptide that has tolerogenic activity as defined by its ability either alone, or in combination with another molecule, to produce a decreased immunological response. Thus, a tolerogenic fragment of an I-2 polypeptide is a peptide or polypeptide that has substantially the same amino acid sequence as a portion of SEQ ID NO: 2 and tolerogenic activity as defined by its ability either alone, or in combination with another molecule, to produce a decreased immunological response. A tolerogenic fragment of an I-2 polypeptide can have from about three amino acids to about 90 amino acids. A tolerogenic fragment of an I-2 polypeptide can have, for example, at least 5, 8, 10, 12, 15, 18, 20 or 25 amino acids. For example, a tolerogenic fragment of an I-2 polypeptide can have from five to fifty amino acids, from eight to fifty amino acids, or from ten to fifty amino acids. More preferably, a tolerogenic fragment has from eight to twenty amino acids or from ten to twenty amino acids. Most preferably, a tolerogenic fragment has from twelve to twenty amino acids or from fifteen to twenty amino acids.

A tolerogenic fragment of an I-2 polypeptide can be identified using a variety of assays, including in

vitro assays such as T-cell proliferation or cytokine secretion assays and *in vivo* assays such as the induction of tolerance in murine models of inflammatory bowel disease. T-cell proliferation assays, for example, are well recognized in the art as predictive of tolerogenic activity (see, for example, Miyahara et al., Immunol. 86:110-115 (1995) or Lundin et al., J. Exp. Med. 178:187-196 (1993)). A T-cell proliferation assay can be performed by culturing T-cells with irradiated antigen-presenting cells, such as normal spleen cells, in microtiter wells for 3 days with varying concentrations of the fragment of an I-2 polypeptide to be assayed; adding ³H-thymidine; and measuring incorporation of ³H-thymidine into DNA. In such an assay, a fragment of an I-2 polypeptide can be tested for activity, for example, at concentrations of 20 µg/ml and 40 µg/ml.

A tolerogenic fragment of an I-2 polypeptide can be identified using a T-cell cytokine secretion assay known in the art. For example, T-cells can be cultured with irradiated antigen-presenting cells in microtiter wells with varying concentrations of the fragment of interest and, after three days, the culture supernatants can be assayed for IL-2, IL-4 or IFN-γ as described in Czerinsky et al., Immunol. Rev. 119:5-22 (1991).

Primary T-cells for use in a T-cell proliferation assay or cytokine secretion assay, for example, can be isolated from lamina propria or peripheral blood. In addition, a convenient source of T-cells for use in an *in vitro* assay for tolerogenic activity can be a T-cell line established from an IBD patient such as a Crohn's disease patient, from a murine model of IBD or from a healthy animal immunized with an I-2 polypeptide of the invention. A preferred source of T-cells for use in

identifying a tolerogenic fragment of an I-2 polypeptide is a Crohn's disease patient.

A T-cell line can be established from a patient with CD or UC, for example, by culturing T lymphocytes with about 1 $\mu\text{g/ml}$ recombinant I-2 polypeptide or GST-I-2, in the presence of low concentrations of growth-supporting IL-2 (about 10 $\mu\text{g/ml}$). A T-cell line can be established by culturing T lymphocytes with antigen-presenting cells and feeding the cells on an alternating four to five day cycle with either IL-2 and I-2 polypeptide or IL-2 alone as described in Nanda et al., J. Exp. Med. 176:297-302 (1992). A cell line that develops into a reliably proliferating cell line dependent on the presence of I-2 polypeptide is particularly useful in identifying tolerogenic fragments of I-2. The establishment of T-cell lines from small intestinal mucosa is described, for example, in Lundin et al., *supra*, 1993. T-cell lines dependent upon the presence of an I-2 polypeptide and useful for identifying I-2 tolerogenic fragments can be prepared similarly.

A tolerogenic fragment of an I-2 polypeptide also can be identified by its ability to induce tolerance *in vivo*, as indicated by a decreased immunological response, which can be a decreased T-cell response, such as a decreased proliferative response or cytokine secretion response as described above, or a decreased antibody titer to the antigen. A neonatal or adult mouse can be tolerized with a fragment of an I-2 polypeptide, for example, and a T-cell response or anti-I-2 polypeptide antibody titer can be assayed after

challenging by immunization. For example, a neonatal mouse can be tolerized within 48 hours of birth by intraperitoneal administration of about 100 µg of a fragment of an I-2 polypeptide emulsified with incomplete Freund's adjuvant and subsequently immunized with I-2 polypeptide at about 8 weeks of age (see, for example, Miyahara, *supra*, 1995). An adult mouse can be tolerized intravenously with about 0.33 mg of a fragment of an I-2 polypeptide, administered daily for three days (total dose 1 mg), and immunized one week later with an I-2 polypeptide. A decreased T-cell response, such as decreased proliferation or cytokine secretion, which indicates tolerogenic activity, can be measured using T-cells harvested 10 days after immunization. In addition, a decreased anti-I-2 polypeptide antibody titer, which also indicates tolerogenic activity, can be assayed using blood harvested 4-8 weeks after immunization. Methods for assaying a T-cell response or anti-I-2 polypeptide antibody titer are described above and are well known in the art.

A tolerogenic fragment of an I-2 polypeptide also can be identified using a murine model of inflammatory bowel disease. Neonatal or adult mice having IBD-like disease can be tolerized with a fragment of an I-2 polypeptide as described above, and the T-cell response or anti-I-2 polypeptide antibody titer assayed. A decreased T-cell response or decreased antibody titer to the antigen indicates a decreased immunological response and, thus, serves to identify a tolerogenic fragment of an I-2 polypeptide. In addition, a tolerogenic fragment of an I-2 polypeptide can be identified by the ability to reduce the frequency, time of onset or severity of colitis in a murine model of IBD.

Several well-accepted murine models of inflammatory bowel disease can be useful in identifying a tolerogenic fragment of an I-2 polypeptide of the invention. For example, mice with target disruption of the genes encoding the alpha subunit of the G-protein Gi2, are a well known model exhibiting features of human bowel disease (Hornquist et al., J. Immunol. 158:1068-1077 (1997); Rudolph et al., Nat. Genet. 10:143-150 (1995)). Mice deficient in IL-10 as described in Kühn et al., Cell 75:263-274 (1993), and mice deficient in IL-2 as described in Sadlack et al., Cell 75:253-261 (1993), also have colitis like disease and are useful in identifying a tolerogenic fragment of an I-2 polypeptide of the invention. Furthermore, mice with mutations in T-cell receptor (TCR) α , TCR β , TCR $\beta \times \delta$, or the class II major histocompatibility complex (MHC) as described in Mombaerts et al., Cell 75:275-282 (1993), develop inflammatory bowel disease and, thus, are useful in identifying a tolerogenic fragment of an I-2 polypeptide. Similarly, a fragment can be assayed for tolerogenic activity in a SCID mouse reconstituted with CD45RB CD4⁺ T-cells, which is a well-accepted model of inflammatory bowel disease, as described in Powrie et al., Immunity 1:553-562 (1994). Additional animal models of IBD also are well known in the art (see, for example, Podolsky, Acta Gastroenterol. Belg. 60:163-165 (1997); and Bregenholt et al., APMIS 105:655-662 (1997)). Thus, a tolerogenic fragment of an I-2 polypeptide can be readily identified by an *in vitro* or

in vivo assay disclosed herein or by another assay well known in the art.

An immunoreactive or tolerogenic fragment of an I-2 polypeptide can be identified by screening a
5 large collection, or library, of peptides of interest or random peptides for immunoreactivity or tolerogenic activity. For example, a panel of peptides spanning the entire sequence of an I-2 polypeptide can be screened for immunoreactivity or tolerogenic activity as
10 described above. Such a panel can be a panel of 15-mer peptides spanning the sequence of the I-2 polypeptide (SEQ ID NO: 2), each overlapping by three or five residue shifts using the Mimotope cleavable pin technology (Cambridge Research Biochemicals, Wilmington,
15 DE), as described by Geysen et al., Science 235:1184 (1987). The panel is subsequently screened for immunoreactivity or tolerogenic activity using one of the assays described above (see, for example, Miyahara et al., *supra*, 1995). A library of peptides to be
20 screened also can be a population of peptides related in amino acid sequence to SEQ ID NO: 2 but having one or more amino acids that differ from SEQ ID NO: 2.

Additional peptides to be screened include, for example, tagged chemical libraries of peptides and
25 peptidomimetic molecules. Peptide libraries also comprise those generated by phage display technology. Phage display technology includes the expression of peptide molecules on the surface of phage as well as other methodologies by which a protein ligand is or can
30 be associated with the nucleic acid which encodes it. Methods for production of phage display libraries,

including vectors and methods of diversifying the population of peptides which are expressed, are well known in the art (see, for example, Smith and Scott, Methods Enzymol. 217:228-257 (1993); Scott and Smith, Science 249:386-390 (1990); and Huse, WO 91/07141 and WO 91/07149). These or other well known methods can be used to produce a phage display library which can be screened, for example, with one of the disclosed assays for immunoreactivity or tolerogenic activity. If desired, a population of peptides can be assayed for activity *en masse*. For example, to identify an immunoreactive fragment of an I-2 polypeptide, a population of peptides can be assayed for the ability to form a complex with a sample containing anti-I-2 polypeptide reactivity; the active population can be subdivided and the assay repeated in order to isolate the immunoreactive fragment from the population.

An immunoreactive or tolerogenic fragment of an I-2 polypeptide also can be identified by screening, for example, fragments of the polypeptide produced by chemical or proteolytic cleavage. Methods for chemical and proteolytic cleavage and for purification of the resultant protein fragments are well known in the art (see, for example, Deutscher, Methods in Enzymology, Vol. 182, "Guide to Protein Purification", San Diego: Academic Press, Inc. (1990)). For example, a chemical such as cyanogen bromide or a protease such as trypsin, chymotrypsin, V8 protease, endoproteinase Lys-C, endoproteinase Arg-C or endoproteinase Asp-N can be used to produce convenient fragments of an I-2 polypeptide that can be screened for immunoreactivity or tolerogenic activity using one of the assays disclosed herein.

As used herein, the term "fragment" means a peptide, polypeptide or compound containing naturally occurring amino acids, non-naturally occurring amino acids or chemically modified amino acids. An
5 immunoreactive or tolerogenic fragment of an I-2 polypeptide also can be a peptide mimetic, which is a non-amino acid chemical structure that mimics the structure of a peptide having an amino acid sequence, provided that the peptidomimetic retains immunoreactivity
10 or tolerogenic activity, as defined herein. Such a mimetic generally is characterized as exhibiting similar physical characteristics such as size, charge or hydrophobicity in the same spatial arrangement found in its peptide counterpart. A specific example of a peptide
15 mimetic is a compound in which the amide bond between one or more of the amino acids is replaced, for example, by a carbon-carbon bond or other bond well known in the art (see, for example, Sawyer, Peptide Based Drug Design, ACS, Washington (1995)).

20

As used herein, the term "amino acid" refers to one of the twenty naturally occurring amino acids, including, unless stated otherwise, L-amino acids and D-amino acids. The term amino acid also refers to
25 compounds such as chemically modified amino acids including amino acid analogs, naturally occurring amino acids that are not usually incorporated into proteins such as norleucine, and chemically synthesized compounds having properties known in the art to be characteristic
30 of an amino acid, provided that the compound can be substituted within a peptide such that it retains immunoreactivity or tolerogenic activity. Examples of amino acids and amino acids analogs are listed in Gross and Meienhofer, The Peptides: Analysis, Synthesis,

Biology, Academic Press, Inc., New York (1983). An amino acid also can be an amino acid mimetic, which is a structure that exhibits substantially the same spatial arrangement of functional groups as an amino acid but
5 does not necessarily have both the α -amino and α -carboxyl groups characteristic of an amino acid.

An immunoreactive or tolerogenic fragment of an I-2 polypeptide can be produced or synthesized using methods well known in the art. Such methods include
10 recombinant DNA methods and chemical synthesis methods for production of a peptide. Recombinant methods of producing a peptide through expression of a nucleic acid sequence encoding the peptide in a suitable host cell are well known in the art and are described, for
15 example, in Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed, Vols 1 to 3, Cold Spring Harbor Laboratory Press, New York (1989). The sequence of a nucleic acid molecule encoding an I-2 polypeptide is disclosed herein as SEQ ID NO: 1.

20 An immunoreactive or tolerogenic fragment of an I-2 polypeptide also can be produced by chemical synthesis, for example, by the solid phase peptide synthesis method of Merrifield et al., J. Am. Chem. Soc. 85:2149 (1964). Standard solution methods well known in
25 the art also can be used to synthesize an immunoreactive or tolerogenic fragment useful in the invention (see, for example, Bodanszky, Principles of Peptide Synthesis, Springer-Verlag, Berlin (1984) and Bodanszky, Peptide Chemistry, Springer-Verlag, Berlin (1993)). A newly

synthesized peptide can be purified, for example, by high performance liquid chromatography (HPLC), and can be characterized using, for example, mass spectrometry or amino acid sequence analysis.

5 The present invention also provides tolerogenic compositions that contain an I-2 polypeptide and are useful in inducing tolerance in a patient with IBD. In particular, the invention provides compositions including an I-2 polypeptide having substantially the
10 same amino acid sequence as SEQ ID NO: 2, or tolerogenic fragment thereof, combined with a tolerogizing molecule. In a composition of the invention, the I-2 polypeptide can have, for example, the amino acid sequence SEQ ID NO: 2. A tolerogenic fragment useful in a composition
15 of the invention can have, for example, at least ten contiguous amino acids of SEQ ID NO: 2.

 Various molecules are known in the art to cause, promote or enhance tolerance. See, for example, U.S. Patent No. 5,268,454, and citations therein. As
20 used herein, the term "tolerogizing molecule" means a molecule, compound or polymer that causes, promotes or enhances tolerogenic activity when combined with an I-2 polypeptide of the invention, or fragment thereof. A tolerogizing molecule can be, for example, conjugated to
25 an I-2 polypeptide. Such tolerogizing molecules include, for example, polyethylene glycol and are well known in the art (see, for example, U.S. Patent No. 5,268,454, *supra*).

 The invention also provides methods of
30 preventing IBD in an individual by administering an effective dose of an I-2 polypeptide, or tolerogenic

fragment thereof, to the individual. The methods of the invention are particularly useful for preventing IBD, for example, Crohn's disease in an individual having increased susceptibility to IBD. Such methods can be particularly useful for preventing IBD when an effective dose of the antigen or tolerogenic fragment is administered to a newborn individual.

Enteric bacteria have been shown to play a role in the pathogenesis of several diseases. For example, *H. pylori* has been implicated in pathogenesis of peptic ulcer disease, and antibiotics against *H. pylori* can be used to effectively treat this disease (see, for example, Sontag, Am. J. Gastroenterol. 92:1255-1261 (1997); and Pipkin et al., Helicobacter 2:159-171 (1997)). In Crohn's disease, intestinal inflammation is notable for its frequent responsiveness to antibiotics and susceptibility to bacterial fecal flow (Gui et al., J. Antimicrob. Chemother. 39:393-400 (1997); Prantera et al., Am. J. Gastroenterol. 91:328-332 (1996); and Janowitz et al., Inflamm. Bowel. Dis. 4:29-39 (1998)). Common intestinal colonists and novel pathogens have been implicated in CD by direct detection or by disease-associated anti-microbial immune responses (Blaser et al., Gastroenterology 87:888-894 (1984); Elsaghier et al., Clin. Exp. Immunol. 90:503-508 (1992); Del Prete et al., J. Microbial. Methods 33:105-114 (1998); Metcalf, British Medical Journal 316:166 (1998); and Sendid et al., *supra*, 1996). Furthermore, in most animal models of chronic colitis, lumenal micro-organisms are a necessary co-factor for disease. Dietary antigens also have been implicated in IBD pathogenesis (Sonnenberg, Gut

31:1037-1040 (1990); Davidson et al., Clin. Exp. Immunol.
35:147-148 (1979); and Knoflach et al., Gastroenterology
92:479-485 (1987). Despite extensive research, the
pathogenic organism remains to be identified.

5

As disclosed herein, colonic microbes harbored
in inflamed lesions in Crohn's disease patients contain a
nucleic acid sequence encoding the I-2 polypeptide
antigen (SEQ ID NO: 2). Based on this finding, the
10 microbial organism can be isolated and used for discovery
of agents that reduce the viability or growth of the
organism, thereby diminishing the immune stimulus
contributing to Crohn's and ameliorating symptoms of the
disease.

15 Thus, the invention provides a method of
identifying an agent useful in treating inflammatory
bowel disease. The method includes the steps of
obtaining a specimen of an enteric microbe from a patient
with inflammatory bowel disease; isolating from the
20 specimen a microbial species that includes a nucleic acid
molecule encoding an I-2 polypeptide; contacting the
microbial species with an agent; and assaying for reduced
growth or viability of the microbial species as compared
to the growth or viability in the absence of the agent,
25 where the reduced growth or viability of the microbial
species indicates that the agent is an agent useful in
treating inflammatory bowel disease. The methods of the
invention can be particularly useful for screening agents
which are antibiotics and for identifying agents for the
30 treatment of Crohn's disease. One skilled in the art
understands that the microbial species which is contacted
with an agent in the methods of the invention can be a
single microbial species or can be a mixture of two or

more microbial species, where at least one species contains a nucleic acid sequence encoding an I-2 polypeptide.

Isolation of a microbial species that includes
5 a nucleic acid molecule encoding an I-2 polypeptide can be performed by culturing the I-2 positive specimen on a variety of mediums and under aerobic and anaerobic conditions as described in Example IV; isolates are subsequently screened for the presence of the I-2
10 sequence using, for example, PCR analysis. Exemplary culture conditions are set forth in Figure 7.

As used herein, the term "agent" means a biological or chemical compound such as a simple or complex organic molecule, a peptide, a protein, an
15 antibody, a lipid or an oligonucleotide.

An agent identified by the methods of the invention reduces the viability or growth of a microbial species that contains a nucleic acid molecule encoding an I-2 polypeptide. Thus, an agent useful in treating IBD
20 can be an agent that functions to kill or slow the growth of a microbial species that contains a nucleic acid molecule encoding an I-2 polypeptide. An agent useful in treating IBD can be a bacteriostatic or bacteriocidal agent such as a bacterial antibiotic, which is a molecule
25 that is produced by a microorganism or a plant, or a chemical derivative of such a molecule, that can reduce the growth or viability of a bacterial species that contains a nucleic acid molecule encoding an I-2 polypeptide. One skilled in the art understands that an
30 agent useful in treating IBD can function by a variety of mechanisms, for example, by inhibiting microbial protein synthesis, inhibiting microbial DNA synthesis, inhibiting

microbial cell wall synthesis or inhibiting synthesis of an essential nutrient of a microbial species that contains a nucleic acid molecule encoding an I-2 polypeptide. Such an agent can selectively reduce the viability or growth of a particular microbial species that contains a nucleic acid molecule encoding an I-2 polypeptide. An agent useful in treating IBD also can have activity in reducing the growth or viability of a broad spectrum of microbes. One skilled in the art understands that, preferably, an agent useful in treating IBD reduces the growth or viability of a microbial species that contains a nucleic acid molecule encoding an I-2 polypeptide without significantly altering the growth or viability of mammalian cells, especially human cells.

As used herein, the term "agent useful in treating IBD" means an agent that reduces the severity, frequency, or time of onset of one or more symptoms of inflammatory bowel disease.

Although animal models of inflammatory bowel disease are known, these models do not involve an antigen associated with human IBD. Based on the disclosed isolation of the IBD-associated I-2 polypeptide antigen (SEQ ID NO: 2) and the demonstration that murine T cells are responsive to this antigen (see Example III), the invention provides novel animal models for IBD, in which disease is initiated with the I-2 antigen associated with human disease. Thus, superior IBD animal models are provided for identifying new IBD therapeutics, including antibiotics, anti-inflammatories and other drugs.

In particular, the invention provides a method of identifying an agent useful in treating inflammatory bowel disease by administering an I-2 polypeptide to a

non-human animal, whereby one or symptoms of IBD are exhibited; administering an agent to the non-human animal; and assaying the level of the one or more symptoms characteristic of IBD, where a reduction in the level of the one or more symptoms as compared to a control level indicates that the agent is an agent useful in treating IBD. The methods of the invention can be applied, for example, to identification of agents useful in treating Crohn's disease. The I-2 polypeptide administered can have, for example, the amino acid sequence SEQ ID NO: 2. A non-human animal particularly useful in the methods of the invention can be, for example, a mouse deficient in $G\alpha i2$, $TCR\alpha$ or IL-10.

As set forth above, the term "agent" means a biological or chemical compound such as a simple or complex organic molecule, a peptide, a protein, an antibody, a lipid or an oligonucleotide. An agent can be, for example, an antibiotic or anti-inflammatory compound.

The methods of the invention rely on the use of a non-human animal, which can exhibit one or more symptoms of inflammatory bowel disease in response to an antigenic trigger. Any non-human animal model of disease can be useful in the methods of the invention including non-human primate, rat or mouse model of IBD. A variety of mouse models of IBD well known in the art can be particularly useful in the invention (see, for example, Podolsky, *supra*, 1997; Bregenholt et al., *supra*, 1997). Such mouse models include those described hereinabove, for example, mice deficient in $G\alpha i2$ ($G\alpha i2^{-/-}$; Rudolph et al., *supra*, 1995); mice deficient in IL-10 ($IL-10^{-/-}$; Kühn et al., *supra*, 1993); mice deficient in $TCR\alpha$ ($TCR\alpha^{-/-}$); mice deficient in $TCR\beta$ ($TCR\beta^{-/-}$); mice deficient

in keratin 8; mice deficient in IL-2 (IL-2 $-/-$; Sadlack, *supra*, 1993); and SCID mice reconstituted with CD45RB CD4⁺ T cells (Powrie et al., *supra*, 1994). Non-human animal models useful in the methods of the invention also
5 include animals expressing transgenes for human HLA-B27 (with β 2-microglobulin) or animals expressing a dominant negative construct that functionally blocks N-cadherin (Podolsky, *supra*, 1997).

One skilled in the art understands that, in
10 order to initiate one or more symptoms of IBD in a non-human animal model, an I-2 polypeptide of the invention can be administered by a variety of routes. Preferred administration of an I-2 polypeptide to initiate one or more symptoms of IBD in a non-human
15 animal model of the disease is by enema administration or by administering a cellular composition in which the I-2 polypeptide is expressed using *E. coli*.

Local administration of free antigen can be achieved by enema, for example, as described in Mahler et
20 al., Am. J. Physiol. 274:G544-G551 (1998). An I-2 polypeptide also can be administered by colonization of the mouse colon with antigen-bearing microorganisms, for example, *E. coli* (Cahill et al., Infect Immuno., 65:3126-3131 (1997)) or another bacterial species (Kullberg et
25 al., Infect. Immun. 66:5157-5166 (1998); Li et al., Infect. Immun. 66:5477-5484 (1998); and Cahill, *supra*, 1997). In a non-human animal model of IBD, one or more symptoms of IBD also can be induced using a T cell line reactive with a bacterial antigen, where the T cell line
30 is derived from a colitis susceptible mouse strain. Such a T cell line can be, for example, an *E. coli* reactive T cell line produced from C3H/HeJBir mice (Cong et al., J. Exp. Med. 187:855-864 (1998)). A C.B-17Scid mouse

also can be colonized with antigen-bearing bacteria,
either naturally or by intra-intestinal infection
(Kullberg et al., *supra*, 1998; Li et al., *supra*, 1998;
and Cong et al., *supra*, 1998). Transfer of a T cell line
5 into such a mouse results in a robust antigen-specific
inflammatory bowel disease model (Kullberg et al., *supra*,
1998; and Cong et al., *supra*, 1998).

The following examples are intended to
10 illustrate but not limit the present invention.

EXAMPLE I

IDENTIFICATION OF THE I-2 and I-1 TARGET ANTIGENS

This example describes isolation of novel
microbial nucleic acid sequences encoding the I-1 and I-2
15 polypeptides.

A. Representational difference analysis

Representational difference analysis (RDA),
which uses PCR driven subtractive cloning to identify DNA
sequences found preferentially in an infected area, was
20 utilized to examine sequences differentially present in
involved versus uninvolved CD mucosa.

RDA was performed essentially as described in
Chang et al., Science 266:1865-1869 (1994). Briefly,
mononuclear cells from the lamina propria of a Crohn's
25 disease patient were isolated from an area with
ulcerations (tester cells) and from an area
macroscopically free of disease (driver cells), and DNA
purified from each cell population. Using RDA, the DNA
sequences from the area free of disease (driver DNA)
30 were subtracted from the DNA

found at the site of ulceration (tester DNA), leaving sequences preferentially found in the inflamed area.

Four sequences were obtained, two of which were human sequences. The remaining two sequences, designated I-1 and I-2, showed no significant homology to any human genes. The I-2 nucleic acid sequence (SEQ ID NO: 1) and encoded polypeptide (SEQ ID NO: 2) are shown in Figure 1A. The I-1 nucleic acid sequence (SEQ ID NO: 3) and two encoded open reading frames (SEQ ID NOS: 4 and 5) are shown in Figure 1B.

B. PCR analysis of archived paraffin embedded tissue samples

Paraffin embedded tissue samples from CD patients, UC patients, or non-IBD controls were subject to PCR analysis. Briefly, sample was amplified with the I-2 specific primers 5'-CCGTGGGCATCCAGTCCG-3' (SEQ ID NO: 9) and 5'-TCTGCTCATACACGTCACG-3' (SEQ ID NO: 10) using standard PCR conditions with a final concentration of 4 mM MgCl₂. In particular, the reactions consisted of 26.75 µl H₂O; 5.0 µl 10X PCR Buffer; 5.0 µl 25 mM MgCl₂; 1.0 µl 10 mM each dNTP; 1.0 µl each of SEQ ID NOS: 9 and 10; and 0.25 µl Taq polymerase (5 units/µl). The reactions were incubated at 94°C for 5 minutes; followed by 39 cycles of 94°C for 30 seconds/65°C for 30 seconds and 72°C for 30 seconds, followed by a final 5 minute extension at 72°C. Reactions were analyzed on 2% agarose, with the presence of the I-2 sequence indicated by the expected 285 bp fragment.

As shown in Figure 3, the I-2 sequence (SEQ ID NO: 1) was present significantly more frequently in involved CD samples than in UC samples or non-IBD

samples. These results indicate that the I-2 polypeptide antigen (SEQ ID NO: 2) is associated with inflammatory bowel disease, in particular, with Crohn's disease.

EXAMPLE II

5 DIFFERENTIAL REACTIVITY OF IBD PATIENT AND NORMAL SERA TO THE I-2 POLYPEPTIDE

This example demonstrates that the I-2 polypeptide is differentially reactive with Crohn's disease patient sera as compared to normal sera.

10 A. GST-I-2 fusion protein

The full-length I-2 encoding nucleic acid sequence (SEQ ID NO: 1) was cloned into the GST expression vector pGEX. After expression in *E. coli*, the protein was purified on a GST column. The purified
15 protein was shown to be of the expected molecular weight by silver staining, and had anti-GST reactivity upon western analysis.

B. ELISA analysis

ELISA analysis was performed with GST-I-2 (SEQ
20 ID NO: 2) fusion polypeptide using diluted patient or normal serum. Reactivity was determined after subtracting reactivity to GST alone. Varying dilutions of CD sera and sera from normal individuals were assayed for IgG reactivity to the GST-I-2 fusion polypeptide. As
25 shown in Figure 4A, dilutions of 1:100 to 1:1000 resulted in significantly higher anti-I-2 polypeptide reactivity for the CD sera as compared to normal sera. These results indicate that the I-2 polypeptide (SEQ ID NO: 2)

is differentially reactive with Crohn's disease sera as compared to normal sera.

Serum IgA reactivity of UC, CD and normal sera to the I-2 polypeptide (SEQ ID NO: 2) was assayed as described below. As shown in Figure 4B, using a cutoff that is two standard deviations above the mean value for the normal population, nine of ten CD values were positive, while none of the normal serum samples were positive. Furthermore, seven of ten Crohn's disease patients showed an OD₄₀₅ greater than 0.3, while none of the UC or normal samples were positive by this measure. These results indicate that immunoreactivity to the I-2 polypeptide, in particular, IgA immunoreactivity, can be used to diagnose Crohn's disease.

Human IgA and IgG antibodies that bind the I-2 polypeptide (SEQ ID NO: 2) were detected by direct ELISA assays essentially as follows. Plates (Immulon 3; DYNEX Technologies; Chantilly, VA) were coated overnight at 4°C with 100 µl/well GST-I-2 fusion polypeptide (5 µg/ml in borate buffered saline, pH 8.5). After three washes in 0.05% Tween 20 in phosphate buffered saline (PBS), the plates were blocked with 150 µl/well of 0.5% bovine serum albumin in PBS, pH 7.4 (BSA-PBS) for 30 minutes at room temperature. The blocking solution was then replaced with 100 µl/well of Crohn's disease serum, ulcerative colitis serum, or normal control serum, diluted 1:100. The plates were then incubated for 2 hours at room temperature and washed as before. Alkaline phosphatase conjugated secondary antibody [goat anti-human IgA (α-chain specific), Jackson ImmunoResearch, West Grove, PA] was added to the IgA plates at a dilution of 1:1000 in BSA-PBS. For IgG reactivity, alkaline phosphatase conjugated secondary antibody (goat anti-human IgG

(γ -chain specific), Jackson ImmunoResearch) was added. The plates were incubated for 2 hours at room temperature before washing three times with 0.05% TweenTM 20/PBS followed by another three washes with Tris buffered normal saline, pH 7.5. Substrate solution (1.5 mg/ml disodium P-nitrophenol phosphate (Aresco; Solon, OH) in 2.5 mM MgCl₂, 0.01 M Tris, pH 8.6) was added at 100 μ l/well, and color allowed to develop for one hour. The plates were then analyzed at 405 nm.

10 C. Histological analysis

Rabbit anti-I-2 antibodies are prepared using purified GST-I-2 fusion protein as the immunogen. GST binding antibodies are removed by adherence to GST bound to an agarose support (Pierce), and the rabbit sera
15 validated for anti-I-2 immunoreactivity by ELISA analysis.

Slides are prepared from paraffin embedded biopsy specimens from CD, UC and normal controls. Hematoxylin and eosin staining are performed, followed by
20 incubation with I-2 specific antiserum. Binding of antibodies is detected with peroxidase labeled anti-rabbit secondary antibodies (Pierce). The assay is optimized to maximize the signal to background and the distinction between CD and control populations.

25

EXAMPLE III

REACTIVITY OF CD4⁺ T CELLS WITH THE I-2 POLYPEPTIDE

This example demonstrates that T cells derived from normal mice proliferate in response to the I-2 antigen (SEQ ID NO: 2).

CD4⁺ T cells were isolated from normal C57BL/6J mice, and T cell proliferation assayed in response to the I-2 antigen as follows. To prepare the CD4⁺ T cells, spleens from 8-10 week old female mice were removed and placed into cell suspension. The cells were depleted for B cells and antigen presenting cells (APCs) using nylon wool and then depleted for CD8⁺ T cells using anti-CD8 magnetic beads. Flow cytometry was used to determine the purity of the CD4⁺ T cell suspension.

10 For preparing APCs, spleens from 8-10 week old females are removed, placed into cell suspension, and pulsed overnight with 0, 2, 10, or 15 µg/ml of the I-2 polypeptide. The APCs are irradiated with 3,000 rads before being added to the T cell cultures. To assay for
15 antigen specific proliferation of T cells, 4 x 10⁵ CD4⁺ T cells/well were incubated with 4 x 10⁵ antigen-pulsed APCs/well in a 96-well flat bottomed tissue culture plate at 37 C in 5% CO₂ humidified air. After varying periods of incubation, 0.5 µCi of [³H]-thymidine was added to each
20 culture for the last 18 hours of incubation. The cells were harvested and proliferation assessed as the amount of [³H]-thymidine into cell DNA by scintillation counting.

As shown in Figure 5, the I-2 polypeptide antigen (SEQ ID NO: 2) stimulates T cell proliferation at
25 concentrations of 5 µg/ml and 25 µg/ml. These results demonstrate that the I-2 polypeptide (SEQ ID NO: 2) associated with human inflammatory bowel disease can elicit a T cell response by murine cells and indicate that an I-2 polypeptide can contribute to the etiology of
30 inflammatory bowel disease.

B. Cytokine expression in response to the I-2 polypeptide

A role for regulatory T cells is supported by studies with CD45RB^{low}, CD28+CD45RB^{low} and CD25+CD45RB^{low} cells. Cytokine expression was therefore analyzed in response to the I-2 polypeptide. T cell proliferation cultures were used for cytokine assays. On day 4 of the T cell proliferation assay, supernatants were analyzed for IL-10 and interferon- γ expression. As shown in Figure 6, expression of interferon- γ , a Th1 cytokine regulator, is induced by the GST-I-2 polypeptide (SEQ ID NO: 2). Furthermore, Figure 6 shows that IL-10 expression, which can act to down regulate a Th1 cytokine response, is induced by exposure of CD4⁺ T cells to as little as 2 μ g/ml GST-I-2 polypeptide (SEQ ID NO: 2). These results indicate that an animal model of IBD in which disease is initiated by the I-2 polypeptide antigen (SEQ ID NO: 2) can be a useful model of human disease.

EXAMPLE IV

ISOLATION AND IDENTIFICATION OF THE ORGANISM CONTAINING THE I-2 SEQUENCE

This example describes isolation and identification of the organism containing the I-2 sequence.

Tissue resections from CD patients are obtained from Cedars-Sinai under anaerobic conditions. DNA is extracted from a small piece of tissue using the Qiagen tissue prep kit, and tested for the presence of the I-2 nucleic acid sequence SEQ ID NO: 1 using the PCR assay described above. Cells from positive scoring sections

are cultured under a variety of conditions (see Figure 7), and the isolates catalogued.

After assaying the isolates by PCR for 16S RNA to assure that the target DNA is accessible to
5 amplification, isolates are assayed by PCR for the I-2 nucleic acid sequence (SEQ ID NO: 1) as described in Example I. The 16S RNA of isolates containing the I-2 nucleic acid are sequenced to identify the organism essentially as described in Wilson and Blitchington,
10 Applied and Environ. Microbiol. 62:2273-2278. (1996).

EXAMPLE V

DEMONSTRATION OF I-2 ENTERIC, T CELL SUPERANTIGEN ACTIVITY

15 This example demonstrates that an I-2-containing microorganism present in normal C57BL/6J mice is a target of a CD4⁺ T cell response. This example further demonstrates that the I-2 response is dependent on MHC class II-mediated recognition but does not require
20 antigen processing, indicating that I-2 represents a new class of T cell superantigen.

A. Mouse tissues contain the I-2 sequence

To determine if the I-2 sequence is present in normal mouse tissue, genomic DNA was prepared from the
25 distal colon, proximal colon, cecum, distal small intestine, proximal small intestine and stomach of normal C57BL/6J mice. PCR analysis detected the 285 bp fragment characteristic of the I-2 sequence in DNA prepared from the distal small intestine, cecum and proximal colon, a
30 pattern identical to that seen in human samples. In contrast, no detectable I-2 fragment was amplified from

genomic DNA isolated from similar tissues of C57BL/6J mice harboring only defined flora, including a non-pathogenic strain of *Clostridium*. These results indicate that an I-2 expressing microbe is present in 5 mice under conventional conditions, but not in mice containing defined flora.

Mice were maintained as follows. C57BL/6J mice were purchased from The Jackson Laboratory (Bar Harbor, ME). Normal C57BL/6J mice greater than eight weeks old 10 were housed in conventional facilities unless otherwise indicated. Both male and female mice were used. Rag-/- mice containing defined flora crossed onto a C57BL/6J background were obtained from the UCLA breeding colony of Dr. W. McBride and housed in specific pathogen free 15 conditions.

I-2 and mouse glyceraldehyde 3-phosphate dehydrogenase (GAPDH) PCR was performed as follows. Mouse genomic DNA was prepared from various tissues by NucleoSpin Tissue kit (Clontech). The native I-2 and 20 positive control mimic I-2 PCR reactions yielded 285 and 606 bp products, respectively, when amplified with primers SEQ ID NOS: 9 and 10. The negative control PCR reaction (buffer only) did not produce any detectable product. PCR conditions were specific for I-2, as shown 25 by amplification of a 606 bp fragment from a synthetic I-2 template. Similar quantities of DNA were present in each sample as shown by amplification of a 340 bp mouse GAPDH fragment from each genomic DNA sample template. The GAPDH primers were 5'-GCCAAGATGTCATTTGAATGG G-3' (SEQ 30 ID NO: 11) and 5'-GAATCCATATAGAGGCTGGG-3' (SEQ ID NO: 12). All reactions were performed at 94° C, 30 seconds for 1 cycle; 94° C, 30 seconds, 65° C, 30 seconds,

72° C, 40 seconds for 39 cycles; and 72° C, 10 minute for 1 cycle.

B. CD4⁺ T cells from normal C57BL/6J and 129Sv mice
5 spontaneously proliferate and produce IL-10 in response
to I-2 pulsed antigen-presenting cells

The native response to the I-2 antigen was evaluated in normal mouse strains housed in conventional
10 conditions. Splenic CD4⁺ T cells were isolated from non-immunized C57BL/6J and 129Sv mice, and cultured with syngeneic splenic antigen-presenting cells pretreated with various concentrations of GST-I-2, GST alone, or *Staphylococcus* enterotoxin (SEB). Proliferation was
15 assessed by [³H]-thymidine incorporation, and secretion of various cytokines (IL-10, IFN- γ , TNF- α , and IL-4) was quantitated by ELISA.

CD4⁺ T cells from C57BL/6J mice displayed a dose-dependent proliferative response to GST-I-2. This
20 response was specific for the I-2 sequence, as indicated by a minimal response to GST alone. The positive control for the T cell proliferation assay, *Staphylococcus* enterotoxin (SEB), yielded responses similar in magnitude to the maximal I-2 response. Furthermore, C57BL/6J CD4⁺ T
25 cells secreted the cytokine IL-10 in response to GST-I-2, while several other cytokines were not detectably secreted, including IFN- γ , IL-4, and TNF- α . GST alone did not stimulate secretion of any cytokines, indicating that the response was specific to the I-2 antigen. These
30 results indicate that CD4⁺ T cells from conventionally housed C57BL/6J mice spontaneously proliferate and produce IL-10 in response to antigen-presenting cells that have been previously exposed to I-2.

The proliferation and cytokine secretion assays were performed as follows. Splenic CD4⁺ T cells were isolated from non-immunized C57BL/6J and 129Sv mice essentially as described in Example III.

5 Antigen-presenting cells were isolated from spleens removed from C57BL/6J or 129Sv mice and placed into cell suspension. The nucleated spleen cells were treated with various concentrations of GST-I-2, GST, or *Staphylococcus* enterotoxin at 2×10^7 cells per 4 ml in 15 ml conical
10 tubes (Corning, Acton, MA) for 12 to 16 hours at 37°C. After washing, the cells were resuspended at 4×10^6 cells/ml in complete media containing RPMI 1640 5% fetal calf serum (FCS), 0.05 mM 2-mercaptoethanol, penicillin, and streptomycin. Antigen-presenting cells were
15 irradiated at 3000 rads (567.7cGy/minute for 5.5 minutes) before addition to T cell cultures.

Syngeneic antigen-presenting cells (4×10^5) were antigen-pulsed with GST alone or GST-I-2 at 0, 1 and 5 µg/ml and SEB (SIGMA) at 1 µg/mL and cultured with $4 \times$
20 10^5 splenic CD4⁺ T cells from C57BL/6J mice in microtiter wells. GST and GST-I-2 proteins were produced as described in Example II. After 48 hours, cells were assayed for cytokine production by collecting culture supernatants and ELISA assays were performed for IL-4,
25 IL-10, TNF-α, and IFN-γ according to the manufacturer's instructions (R&D; Minneapolis, MN); values are the means of duplicate samples. After 72 hours, cells were assayed for proliferation by pulsing with [³H]-thymidine, and incorporated cpm (mean ± SD) were determined for
30 triplicate cultures essentially as described in Example III.

C. CD4⁺ T cells from mice containing defined flora spontaneously proliferate, but do not produce IL-10 in response to I-2

5 The response of normal C57BL/6J mice to GST-I-2 was compared to that of C57BL/6J mice containing defined flora and housed under specific pathogen-free conditions. CD4⁺ T cells isolated from C57BL/6J mice containing defined flora had a proliferative response to the GST-I-2
10 protein. However, in contrast to the GST-I-2 response observed in the conventionally housed normal C57BL/6J mice, no cytokine response was observed. The cells were proliferation competent as shown with the SEB positive control, and, furthermore, the response was I-2 specific
15 since no proliferative or cytokine response was observed in response to GST alone.

 The response of mice containing defined flora to GST-I-2 indicates that the I-2 antigen was able to stimulate CD4⁺ T cells without prior exposure to antigen.
20 Coupled with the demonstration that the I-2 sequence was not present in C57BL/6J mice containing defined flora, these results indicate that the GST-I-2 response can be a result of superantigen or mitogen stimulation.

25 D. Both naïve and memory CD4⁺ T cell populations from normal C57BL/6J mice spontaneously proliferate and produce IL-10 in response to I-2

 The I-2 response was evaluated using a naïve CD4⁺ T cell population. CD4⁺ T cells from conventionally
30 housed C57BL/6J mice were fractionated into CD62L^{hi}CD44^{low} naïve T cells and CD62L^{low}CD44^{hi} memory T cells, and subsequently cultured with GST-I-2, GST alone or SEB pulsed antigen-presenting cells. Both naïve and memory

CD4⁺ T cells had a strong proliferative and IL-10 response to GST-I-2 (Figures 8A and 8B). The response was specific to I-2 since neither population was responsive to GST alone (Figures 8A and 8B), and both populations
5 were proliferation competent as shown with the SEB positive control. These results demonstrate that GST-I-2 antigen is not a conventional MHC class II processed antigen, but a superantigen or mitogen.

10 Naïve and memory subpopulations of CD4⁺ T cells from conventionally housed normal C57BL/6J mice were obtained and analyzed as follows. Naïve CD4⁺ T cells were obtained from CD4⁺ T cells, isolated as described in section VB above, and incubated with rat anti-mouse CD44
15 for 30 minutes on ice. Memory CD4⁺ T cells were obtained from CD4⁺ T cells, incubated with rat anti-mouse CD45RB and rat-anti mouse CD62L for 30 minutes on ice. The cells were washed and treated with sheep anti-rat-IgG coated Dynal magnetic beads for 30 minutes on ice. After
20 depletion, the unbound T cells were negatively selected and resuspended at 4×10^6 cells/ml in complete RPMI media containing 5% FCS. Fractionated T cells were assessed for purity by flow cytometry with FITC-conjugated anti-CD4, CD44, CD45RB and CD62L (Pharmingen; San Diego,
25 CA). Populations were typically greater than 85% memory or naïve CD4⁺ T cells.

Fractionated naïve and memory CD4⁺ T cells from C57BL/6J mice (4×10^5) were cultured in microtiter wells with 4×10^5 syngeneic antigen-presenting cells previously
30 antigen-pulsed with antigen (0, 1, 5 µg/mL; GST; or GST-I-2) or 1 µg/ml positive control SEB as described above. Proliferation and cytokine production were assayed as described above.

E. MHC class II restriction of CD4⁺ T cell response to I-2

The I-2 response was analyzed for dependence on MHC class II processing. CD4⁺ T cells were cultured with I-2 pulsed antigen-presenting cells with either monoclonal antibody to I-A^{-b} or I-A^{-d} MHC class II molecules. Anti-I-A^{-b} antibody blocked the T cell proliferative response to both SEB and GST-I-2 while anti-I-A^{-d} antibody had minimal effect on the proliferative response of SEB or GST-I-2. These results indicate that the I-2 response is dependent on MHC II-mediated recognition and, thus, is not a mitogen response.

15

MHC class II blocking of T cell proliferation was performed as follows. CD4⁺ T cells (4×10^5) were incubated with 4×10^5 antigen-pulsed antigen-presenting cells as described in section VB above, in the presence and absence of 10 µg/ml monoclonal anti-I-A^b or anti-I-A^d B (Pharmingen). After 48 hours of incubation, 0.5 µCi [³H]-thymidine was added to each culture for the last 18 hours of the incubation period and collected as before. Results are expressed as mean CPM +/- SD of triplicate cultures.

25

F. The CD4⁺ T cell response to I-2 does not require antigen processing

The results set forth above indicate that the I-2 antigen is not a mitogen, but rather either a conventional MHC II processed antigen or a superantigen, which is an antigen characterized, in part, as able to stimulate a proliferative response in the absence of antigen processing. To determine whether the I-2

30

response requires antigen processing, antigen-presenting cells from normal C57BL/6J mice were fixed with paraformaldehyde to eliminate antigen-processing and then pulsed with GST-I-2 or SEB. As shown in Figure 9A, CD4⁺ T cells still responded strongly to GST-I-2 and SEB. Furthermore, the response was specific to I-2 since GST alone did not induce a proliferative response. In contrast, fixed antigen-presenting cells pulsed with pigeon cytochrome C (PPC) overnight did not induce PPC specific CD4⁺ T cells to proliferate (Figure 9B). These results demonstrate that the GST-I-2 proliferative response does not require antigen processing and, therefore, that GST-I-2 acts as a superantigen.

Paraformaldehyde-fixed antigen-presenting cells were prepared and analyzed as follows. The splenic C57 BL/6J CD4⁺ T cells were cultured with antigen-presenting cells from C57 BL/6J mice that were pulsed with GST-I-2 overnight and for 30 minutes, or fixed with paraformaldehyde for 30 minutes, washed with PBS, incubated with PBS for 30 minutes, and washed twice with PBS. The fixed cells were pulsed with GST-I-2 overnight at 1 µg/ml and 5 µg/ml. GST and SEB were used as the negative and positive control antigens, respectively. As the control for fixation experiment, pigeon cytochrome C (PPC) specific CD4⁺ T cells were cultured with syngeneic antigen-presenting cells pulsed with PPC overnight, or fixed with paraformaldehyde and then pulsed with PPC overnight. 1 µCi of [³H]TdR was added to the wells for the last 18 hour of a 3 day culture. Results are expressed as mean CPM ± SD of triplicate cultures.

G. TCR V β expression is restricted in the responding T cell population

Most superantigens are characterized by a specific TCR V β . From a panel of 15 V β s, V β 9 and V β 17
5 were expanded in CD4⁺ T cells after GST-I-2 injection. These results confirm that GST-I-2 is a superantigen.

V β restriction was characterized as follows. GST-I-2 protein (5 μ g) was injected i.v. into normal 8 week old female C57BL/6J mice. One day after injection,
10 spleens were removed and placed into cell suspension. Red blood cells were lysed, and the nucleated spleen cells were resuspended into 5 ml of complete RPMI media containing 5% FCS. Cells (2×10^6) were double stained
15 with PERCP-conjugated anti-CD4⁺ (Pharmingen) and various FITC-conjugated anti-V β (Pharmingen). Stained cells were assessed for CD4⁺ and V β positivity by flow cytometry.

All journal article, reference and patent citations provided above, in parentheses or otherwise,
whether previously stated or not.

20

Although the invention has been described with reference to the examples above, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the
25 invention is limited only by the following claims.

65/1

SEQUENCE LISTING

<110> The Regents of the University of California

<120> IBD-Associated Microbial Antigens and Methods of Using Same

<130> 08-893043CA

<140> 2,365,898

<141> 2000-04-28

<150> US 09/303,120

<151> 1999-04-30

<160> 12

<170> PatentIn Ver. 2.0

<210> 1

<211> 302

<212> DNA

<213> Unknown

<220>

<221> CDS

<222> (2) .. (301)

<220>

<223> Description of Unknown Organism: Microbial
Organism from the human gut

<400> 1

```

a gat ctg gcc agc gcc gtg ggc atc cag tcc ggc agc atc ttt cat cac 49
  Asp Leu Ala Ser Ala Val Gly Ile Gln Ser Gly Ser Ile Phe His His
    1             5             10             15

```

```

ttc aag agc aag gat gag ata ttg cgt gcc gtg atg gag gaa acc atc 97
Phe Lys Ser Lys Asp Glu Ile Leu Arg Ala Val Met Glu Glu Thr Ile
      20             25             30

```

```

cat tac aac acc gcg atg atg cgc gct tca ctg gag gag gcg agc acg 145
His Tyr Asn Thr Ala Met Met Arg Ala Ser Leu Glu Glu Ala Ser Thr
      35             40             45

```

```

gtg cgc gaa cgc gtg ctg gcg ctg atc cgc tgc gag ttg cag tcg atc 193
Val Arg Glu Arg Val Leu Ala Leu Ile Arg Cys Glu Leu Gln Ser Ile
    50             55             60

```

```

atg ggc ggc agt ggc gag gcc atg gcg gtg ctg gtc tac gaa tgg cgc 241
Met Gly Gly Ser Gly Glu Ala Met Ala Val Leu Val Tyr Glu Trp Arg
    65             70             75             80

```

```

tcg ctg tcg gcc gaa ggc cag gcg cac gtg ctg gcc ctg cgt gac gtg 289

```

65/2

Ser Leu Ser Ala Glu Gly Gln Ala His Val Leu Ala Leu Arg Asp Val

85

90

95

tat gag cag atc t

302

Tyr Glu Gln Ile

100

<210> 2

<211> 100

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Microbial Organism from the human gut

<400> 2

Asp Leu Ala Ser Ala Val Gly Ile Gln Ser Gly Ser Ile Phe His His

1

5

10

15

Phe Lys Ser Lys Asp Glu Ile Leu Arg Ala Val Met Glu Glu Thr Ile

20

25

30

His Tyr Asn Thr Ala Met Met Arg Ala Ser Leu Glu Glu Ala Ser Thr

35

40

45

Val Arg Glu Arg Val Leu Ala Leu Ile Arg Cys Glu Leu Gln Ser Ile

50

55

60

Met Gly Gly Ser Gly Glu Ala Met Ala Val Leu Val Tyr Glu Trp Arg

65

70

75

80

Ser Leu Ser Ala Glu Gly Gln Ala His Val Leu Ala Leu Arg Asp Val

85

90

95

Tyr Glu Gln Ile

100

<210> 3

<211> 392

<212> DNA

<213> Unknown

<220>

<221> CDS

<222> (2)..(346)

<220>

<223> Description of Unknown Organism: Microbial
Organism from the human gut

<220>

<221> misc_feature

65/3

<222> (378)

<223> n=a, c, t or g

<400> 3

a gat ctt gag cgt cat gag tgc ctg ggg tac gcc ttt tca tcg cgt ccg 49

Asp Leu Glu Arg His Glu Cys Leu Gly Tyr Ala Phe Ser Ser Arg Pro

1

5

10

15

gcg gat cga gag tgg gtg ttt ttt cag ggc acg gtt tcc tac aag gta 97

Ala Asp Arg Glu Trp Val Phe Phe Gln Gly Thr Val Ser Tyr Lys Val

20

25

30

cga gtg gcc agc cgt ttg ctc atc aat gaa agc cgg gca ttg atg tcg 145

Arg Val Ala Ser Arg Leu Leu Ile Asn Glu Ser Arg Ala Leu Met Ser

35

40

45

gcg gca ttg gat ggt ttt ggc ata gtg ctc ggc ccg caa gac ttc ctg 193

Ala Ala Leu Asp Gly Phe Gly Ile Val Leu Gly Pro Gln Asp Phe Leu

50

55

60

cga acg gcg ttg gcg agt ggc gag ttg gtg cgg gtg ttg ccg gag ttt 241

Arg Thr Ala Leu Ala Ser Gly Glu Leu Val Arg Val Leu Pro Glu Phe

65

70

75

80

gag gct ccg agt cgg tcg atg cat ttg gtc tac acc gca aac cgc cag 289

Glu Ala Pro Ser Arg Ser Met His Leu Val Tyr Thr Ala Asn Arg Gln

85

90

95

cgt acc gcc aag ttg cgc tgc ttt gtc gag act gtg ctg gga cgt ttt 337

Arg Thr Ala Lys Leu Arg Cys Phe Val Glu Thr Val Leu Gly Arg Phe

100

105

110

ggt ccg gta tgaaggagca ccaccgtggc ggtcgccggg angcacctaa agatct 392

Gly Pro Val

115

<210> 4

<211> 115

<212> PRT

<213> Unknown

<220>

<223> Description of Unknown Organism: Microbial Organism from the human gut

65/4

<400> 4

Asp Leu Glu Arg His Glu Cys Leu Gly Tyr Ala Phe Ser Ser Arg Pro

1 5 10 15

Ala Asp Arg Glu Trp Val Phe Phe Gln Gly Thr Val Ser Tyr Lys Val

20 25 30

Arg Val Ala Ser Arg Leu Leu Ile Asn Glu Ser Arg Ala Leu Met Ser

35 40 45

Ala Ala Leu Asp Gly Phe Gly Ile Val Leu Gly Pro Gln Asp Phe Leu

50 55 60

Arg Thr Ala Leu Ala Ser Gly Glu Leu Val Arg Val Leu Pro Glu Phe

65 70 75 80

Glu Ala Pro Ser Arg Ser Met His Leu Val Tyr Thr Ala Asn Arg Gln

85 90 95

Arg Thr Ala Lys Leu Arg Cys Phe Val Glu Thr Val Leu Gly Arg Phe

100 105 110

Gly Pro Val

115

<210> 5

<211> 114

<212> PRT

<213> Unknown

<220>

<221> SITE

<222> (113)

<223> Any amino acid

<220>

<223> Description of Unknown Organism: Microbial
Organism from the human gut

<400> 5

Arg Thr Arg Arg Ile Ser Leu Pro His Lys Lys Leu Ala Arg Asn Gly

1 5 10 15

Val Leu Tyr Ser His Gly Ala Thr Gln Glu Asp Ile Phe Ala Pro Cys

20 25 30

Gln His Arg Arg Cys Gln Ile Thr Lys Ala Tyr His Glu Ala Arg Leu

35 40 45

Val Glu Gln Ser Arg Arg Gln Arg Thr Ala Leu Gln His Pro His Gln

65/5

50		55		60											
Arg	Leu	Lys	Leu	Ser	Arg	Thr	Pro	Arg	His	Met	Gln	Asp	Val	Gly	Cys
65			70					75						80	
Val	Ala	Leu	Thr	Gly	Gly	Leu	Gln	Ala	Ala	Lys	Asp	Leu	Ser	His	Gln
			85					90						95	
Ser	Thr	Lys	Thr	Arg	Tyr	Ser	Pro	Ala	Gly	Gly	His	Arg	Asp	Gly	Pro
			100					105						110	

Xaa Val

<210> 6
 <211> 190
 <212> PRT
 <213> Clostridium pasteurianum

<400> 6

Met	Asn	Lys	Thr	Lys	Asp	Asn	Ile	Phe	Tyr	Ser	Ala	Ile	Lys	Val	Phe
1				5					10					15	
Ser	Asn	Asn	Gly	Tyr	Asn	Gly	Ala	Thr	Met	Asp	Glu	Ile	Ala	Ser	Asn
			20					25					30		
Ala	Gly	Val	Ala	Lys	Gly	Thr	Leu	Tyr	Tyr	His	Phe	Lys	Ser	Lys	Glu
			35				40					45			
Glu	Ile	Phe	Lys	Tyr	Ile	Ile	Glu	Glu	Gly	Val	Asn	Leu	Met	Lys	Asn
			50				55				60				
Glu	Ile	Asp	Glu	Ala	Thr	Asp	Lys	Glu	Lys	Thr	Ala	Leu	Glu	Lys	Leu
65				70					75					80	
Lys	Ala	Val	Cys	Arg	Val	Gln	Leu	Asn	Leu	Ile	Tyr	Lys	Asn	Arg	Asp
				85					90					95	
Phe	Phe	Lys	Val	Ile	Ala	Ser	Gln	Leu	Trp	Gly	Lys	Glu	Leu	Arg	Gln
			100					105						110	
Leu	Glu	Leu	Arg	Asp	Ile	Met	Arg	Asn	Tyr	Val	Val	His	Ile	Glu	Glu
			115					120					125		
Phe	Val	Lys	Asp	Ala	Met	Glu	Ala	Gly	Ser	Ile	Lys	Lys	Gly	Asn	Ser
			130					135					140		
Leu	Phe	Val	Ala	Tyr	Ala	Phe	Leu	Gly	Thr	Leu	Cys	Ser	Val	Ser	Leu

Met	Asp	Arg	Val	Ala	Gly	Gln	Val	Asn	Ser	Arg	Arg	Gly	Glu	Leu	Leu	
1				5					10					15		
Glu	Leu	Ala	Ala	Ala	Met	Phe	Ala	Glu	Arg	Gly	Leu	Arg	Ala	Thr	Thr	
			20					25					30			
Val	Arg	Asp	Ile	Ala	Asp	Gly	Ala	Gly	Ile	Leu	Ser	Gly	Ser	Leu	Tyr	
		35					40					45				
His	His	Phe	Ala	Ser	Lys	Glu	Glu	Met	Val	Asp	Glu	Leu	Leu	Arg	Gly	
	50					55					60					
Phe	Leu	Asp	Trp	Leu	Phe	Ala	Arg	Tyr	Arg	Asp	Ile	Val	Asp	Ser	Thr	
65					70					75					80	
Ala	Asn	Pro	Leu	Glu	Arg	Leu	Gln	Gly	Leu	Phe	Met	Ala	Ser	Phe	Glu	
				85					90					95		
Ala	Ile	Glu	His	His	His	Ala	Gln	Val	Val	Ile	Tyr	Gln	Asp	Glu	Ala	
			100					105					110			
Gln	Arg	Leu	Ala	Ser	Gln	Pro	Arg	Phe	Ser	Tyr	Ile	Glu	Asp	Arg	Asn	
		115					120					125				
Lys	Gln	Gln	Arg	Lys	Met	Trp	Val	Asp	Val	Leu	Asn	Gln	Gly	Ile	Glu	
	130					135					140					
Glu	Gly	Tyr	Phe	Arg	Pro	Asp	Leu	Asp	Val	Asp	Leu	Val	Tyr	Arg	Phe	
145					150					155					160	
Ile	Arg	Asp	Thr	Thr	Trp	Val	Ser	Val	Arg	Trp	Tyr	Arg	Pro	Gly	Gly	
				165					170					175		

65/7

Pro Leu Thr Ala Gln Gln Val Gly Gln Gln Tyr Leu Ala Ile Val Leu

180

185

190

Gly Gly Ile Thr Lys Glu Gly Val

195

200

<210> 8

<211> 192

<212> PRT

<213> Aquifex aeolicus

<400> 8

Met Tyr Ile Leu Leu Phe Met Gly Glu Lys Arg Ser Asp Thr Lys Glu

1

5

10

15

Lys Ile Leu Ser Ser Ala Leu Lys Leu Phe Ser Lys Lys Gly Phe Lys

20

25

30

Glu Thr Thr Ile Lys Asp Ile Ala Lys Glu Val Gly Ile Thr Glu Gly

35

40

45

Ala Ile Tyr Arg His Phe Thr Ser Lys Glu Glu Ile Ile Lys Ser Leu

50

55

60

Leu Glu Ser Ile Thr Lys Glu Leu Arg His Lys Leu Glu Val Ala Leu

65

70

75

80

Gln Arg Gly Glu Thr Asp Glu Glu Ile Leu Glu Ser Ile Val Asp Thr

85

90

95

Leu Ile Asp Tyr Ala Phe Ser Asn Pro Glu Ser Phe Arg Phe Leu Asn

100

105

110

Leu Tyr His Leu Leu Lys Glu Tyr Gly Glu Val Lys Asn Leu Pro Gly

115

120

125

Glu Leu Ile Leu Lys Phe Leu Asn Gly Leu Tyr Leu Lys Arg Lys Leu

130

135

140

Lys Thr Tyr Pro Glu Ile Ala Leu Ala Val Val Thr Gly Ser Val Glu

145

150

155

160

Arg Val Phe Ile Phe Lys Glu Arg Asn Phe Leu Asp Tyr Asp Glu Glu

165

170

175

Thr Ile Lys Lys Glu Leu Lys Lys Val Leu Lys Ser Ala Ile Leu Ala

65/8

180	185	190
<210> 9 <211> 18 <212> DNA <213> Unknown		
<220> <223> Description of Unknown Organism: Microbial organism from the human gut		
<400> 9		
ccgtgggcat ccagtccg		18
<210> 10 <211> 19 <212> DNA <213> Unknown		
<220> <223> Description of Unknown Organism: Microbial organism from the human gut		
<400> 10		
tctgctcata cacgtcacg		19
<210> 11 <211> 22 <212> DNA <213> Mus musculus		
<400> 11		
gccaaagatgt catttgaatg gg		22
<210> 12 <211> 20 <212> DNA <213> Mus musculus		
<400> 12		
gaatccatat agaggctggg		20

CLAIMS

1. An isolated inflammatory bowel disease (IBD) associated polypeptide (I-2 polypeptide), comprising an amino acid sequence which is greater than 80% identical to the amino acid sequence defined by SEQ ID NO:2, wherein said I-2 polypeptide retains substantially the immunoreactivity or tolerogenic activity of a polypeptide comprising the amino acid sequence defined by SEQ ID NO:2, wherein said immunoreactivity is defined by the ability of said I-2 polypeptide to form a complex with antibody from a sample wherein the antibody is against the polypeptide comprising the amino acid sequence defined by SEQ ID NO:2, and wherein said tolerogenic activity is defined by the ability of said I-2 polypeptide, either alone, or in combination with another molecule, to produce a decreased immunological response to the polypeptide comprising the amino acid sequence defined by SEQ ID NO:2.

2. The isolated I-2 polypeptide of claim 1, wherein said I-2 polypeptide comprises an amino acid sequence which is greater than 90% identical to the amino acid sequence defined by SEQ ID NO:2.

3. The isolated I-2 polypeptide of claim 1, wherein said I-2 polypeptide comprises an amino acid sequence which is greater than 95% identical to the amino acid sequence defined by SEQ ID NO:2.

4. The isolated I-2 polypeptide of claim 1, 2 or 3, wherein said I-2 polypeptide retains substantially the immunoreactivity of a polypeptide comprising the amino acid sequence defined by SEQ ID NO:2 and said immunoreactivity is defined by the ability of said polypeptide to form a

complex with an antibody against the polypeptide comprising the amino acid sequence defined by SEQ ID NO:2.

5. The isolated I-2 polypeptide of claim 1, consisting of the amino acid sequence defined by SEQ ID NO:2.

6. An isolated immunoreactive fragment of an I-2 polypeptide of any one of claims 1 to 5, said fragment having at least 10 contiguous amino acids of said I-2 polypeptide, wherein said immunoreactivity is defined by the ability of the polypeptide to form a complex with an antibody in a sample wherein the antibody is against the polypeptide comprising the amino acid sequence defined by SEQ ID NO:2.

7. The isolated immunoreactive fragment of claim 6, wherein said fragment is at least 15 amino acids in length.

8. The isolated immunoreactive fragment of claim 6, wherein said fragment is from 10 to 50 amino acids in length.

9. An isolated tolerogenic fragment of an I-2 polypeptide of any one of claims 1 to 5, said fragment having at least 8 contiguous amino acids of said I-2 polypeptide; wherein said fragment tolerogenic activity is defined by the ability of said fragment, either alone, or in combination with another molecule, to produce a decreased immunological response.

10. The isolated tolerogenic fragment of claim 9, wherein said fragment is at least 15 amino acids in length.

11. The isolated tolerogenic fragment of claim 9, wherein said fragment is from 10 to 50 amino acids in length.

12. The isolated tolerogenic fragment of claim 9, wherein said decreased immunological response is a decreased T-cell proliferative response.

13. An antibody material that selectively binds an I-2 polypeptide of any one of claims 1 to 5, wherein the antibody material is substantially devoid of antibody material that does not bind the I-2 polypeptide.

14. An isolated nucleic acid molecule, comprising a nucleic acid sequence encoding the amino acid sequence of an I-2 polypeptide of any one of claims 1 to 5.

15. The isolated nucleic acid molecule of claim 14, comprising the nucleic acid sequence defined by SEQ ID NO:1.

16. A pharmaceutical composition comprising an I-2 polypeptide of any one of claims 1 to 5, or a tolerogenic fragment of any one of claims 9 to 12, combined with a tolerogenizing molecule.

17. A method of diagnosing inflammatory bowel disease (IBD) in a subject, comprising:

(a) contacting an I-2 polypeptide of any one of claims 1 to 5, or an immunoreactive fragment of any one of claims 6 to 8, under conditions suitable to form a complex of said I-2 polypeptide, or said immunoreactive fragment thereof, and an antibody to said I-2 polypeptide; and

(b) detecting the presence or absence of said complex,

wherein the presence of said complex indicates that said subject has IBD.

18. The method of claim 17, wherein the presence or absence of said complex is detected with a detectable secondary antibody that has specificity for a class determining portion of said antibody to said I-2 polypeptide.

19. The method of claim 17, wherein said IBD is Crohn's disease.

20. The method of any one of claims 18 to 19, wherein said I-2 polypeptide comprises the sequence defined by SEQ ID NO:2.

21. Use of an I-2 polypeptide of any one of claims 1 to 5, or a tolerogenic fragment of any one of claims 9 to 12, for preparing a pharmaceutical composition for inducing tolerance in a patient with IBD.

22. The use of claim 21, wherein said IBD is Crohn's disease.

23. The use of claim 21 or 22, wherein said I-2 polypeptide comprises the sequence defined by SEQ ID NO:2.

24. A method of identifying an agent useful in treating inflammatory bowel disease (IBD), comprising:

(a) isolating from a specimen of an enteric bacteria from a patient with IBD, a microbial species that

comprises a nucleic acid molecule encoding an I-2 polypeptide of any one of claims 1 to 5;

(b) contacting said microbial species with the agent; and

(c) assaying for reduced growth or viability of said microbial species as compared to the growth or viability in the absence of said agent,

wherein said reduced growth or viability of said microbial species indicates that said agent is an agent useful in treating IBD.

25. The method of claim 24, wherein said IBD is Crohn's disease.

26. The method of claim 24, wherein said agent is an antibiotic.

27. A method of identifying an agent useful in treating inflammatory bowel disease (IBD), comprising:

(a) administering an I-2 polypeptide of any one of claims 1 to 5 to a non-human animal, whereby one or more symptoms of IBD are exhibited;

(b) administering an agent to said non-human animal; and

(c) assaying the level of said one or more symptoms characteristic of IBD,

wherein a reduction in the level of said one or more symptoms as compared to a control level indicates that said agent is an agent useful in treating IBD.

28. The method of claim 27, wherein said IBD is Crohn's disease.

29. The method of claim 27 or 28, wherein said I-2 polypeptide comprises the amino acid sequence defined by SEQ ID NO:2.

30. The method of any one of claims 27 to 29, wherein said non-human animal is a mouse selected from the group consisting of a *Gai2* ^{-/-} mouse; a *TCRα* ^{-/-} mouse and a *IL-10* ^{-/-} mouse.

31. Use of an immunoreactive fragment of any one of claims 6 to 8 in an *in vitro* method of diagnosing IBD.

32. Use of an isolated colonic microbe containing a nucleic acid sequence encoding an I-2 polypeptide of any one of claims 1 to 5 in the identification of agents useful for treating IBD, wherein the agent is contacted with the microbe and the agent reduces the viability of growth of the microbe.

33. Use of an effective dose of I-2 polypeptide of any one of claims 1 to 5, or tolerogenic fragments thereof, for treating a patient with IBD.

FIGURE 1A

A GAT CTG GCC AGC GCC GTG GGC ATC CAG TCC GGC AGC ATC TTT CAT CAC TTC AAG AGC AAG
▶ D L A S A V G I Q S G S I F H H F K S K
GAT GAG ATA TTG OGT GCC GTG ATG GAG GAA ACC ATC CAT TAC AAC ACC GCG ATG ATG OGC
▶ D E I L R A V M E E T I H Y N T A M M R
GCT TCA CTG GAG GAG GCG AGC ACG GTG GCG GAA OGC GTG CTG GCG ATC OGC TGC GAG
▶ A S L E E A S T V R E R V L A L I R C E
TTG CAG TCG ATC ATG GGC GGC AGT GGC GAG GGC ATG GCG GTG CTG GTC TAC GAA TGG OGC
▶ L Q S I M G G S G E A M A V L V Y E W R
TOG CTG TOG GGC GAA GGC CAG GCG CAC GTG CTG GCG CIG OGT GAC GTG TAT GAG CAG ATC T
▶ S L S A E G Q A H V L A L R D V Y E Q I

FIGURE 1B

AGATCTGAGCGTCATGAGTCCCTGGGTACGCCCTTTTCATCGCGTCCGGGATCGAGAGTGGTGT
▶ D L E R H E C L G Y A F S S R P A D R E W V
TTTTCAGGGCACGGTTTCCTACAAGGTACGAGTGGCCAGCGGTTTGCTCATCAATGAAGCGGGCA
▶ F F Q G T V S Y K V R V A S R L L I N E S R A
▶ K K L A R N G V L Y S H G A T Q E D I F A P C
TTGATGTCCGCGCATTTGGATGGTTTGGCATAGTGTCTGCGCCGCGCAAGACTTCTCGGAACGGCGTT
▶ L M S A A L D G F G I V L G P Q D F L R T A L
▶ Q H R R C Q I T K A Y H E A R L V E Q S R R Q
GGCGAGTGGCGAGTTGGTCCGGGTGTTCGCGAGTTTGAGGCTCCGAGTGGTGGATTTGGTCT
▶ A S G E L V R V L P E F E A P S R S M H L V
▶ R T A L Q H P H Q R L K L S R T P R H M Q D
ACACCGCAACCGCCAGCGTACCGCCAGTTGGCTGTCTGTTCGAGACTGTGCTGGGACGTTTGGT
▶ Y T A N R Q R T A K L R C F V E T V L G R F G
▶ V G C V A L T G G L Q A A K D L S H Q S T K T
CCGTTATGAGGACCAACCGTGGCGGTCCCGGGANGCACCTAAAGATCT
▶ P V
▶ R Y S P A G G H R D G P . V

2/10

I-2	:	-----D	:	1
Clostridium	:	-----MKT KDNIFYSAIVFNNNGNGATMDE	:	28
Mycobacterium	:	---MDRVAGQVNSRRGELLEAAAMFAEGLRAITVRD	:	35
Aquifex	:	MYIILFMGEKRSDKEKILSSALILFKKGGETTIKD	:	38
		a f g t d		
I-2	:	LASAVGIQSGSIFTHFKSKDEILRAVMEETLHNTAMM	:	39
Clostridium	:	IASNAGVAKGTLYYHFKSKEEIFKYIIEEGVNLKNEI	:	66
Mycobacterium	:	IADGAGILSGSLYHFKSKEEMVDELLRGFLDWLFARY	:	73
Aquifex	:	IAKEVGITEGAIYRHFTSKEEIIKSLLSITKELRHKL	:	76
		6A G6 G 65 HF SKE6 66e		
I-2	:	RASLEEASIVRERVLAALRCCELOSMGGSGEAMAVLWY	:	77
Clostridium	:	DEATREKTALEKLVKAVCRVOLNLIYKTRDFEKVIASQ	:	104
Mycobacterium	:	RDIVSTANPEERLGLFMASFEATEHHHAQVVYQDE	:	111
Aquifex	:	EVALORGEIDEEILESIMDTLIIYASNPESERFLNLY	:	114
		t E 6 6 i		
I-2	:	EWRSLSAEGGAHVIAIR--DVMEQI-----	:	100
Clostridium	:	LWGKELROTELRLDMRN--YVHIEEIVKDAMEAGSIK	:	140
Mycobacterium	:	AQRLASQPRFSYIEDRNKQQRKMWDVNLNOGTIEGYFR	:	149
Aquifex	:	HLLKEYGEVKNLPGETI--LKEINGLYLKRKLKT---Y	:	147
I-2	:	-----	:	-
Clostridium	:	KGNSLFVAMAFEGTICSVSLEYEVINAENDNINNTIENE	:	178
Mycobacterium	:	EDLDVDLYYRFTIRDTTWVSVRWYRPGGPLTAQVGOOY	:	187
Aquifex	:	EEALAVMT---SSVERVFIEKERNFLEYDEETKKEL	:	182
		v		
I-2	:	-----	:	-
Clostridium	:	MNYLNLGIGLON-	:	190
Mycobacterium	:	LAIVLGGITKEGV	:	200
Aquifex	:	KKMLKSAILA---	:	192
		1		

FIGURE 2

	Diagnosis	Tissue		N	N(+)	% Pos
	CD	Ileum	Involved	46	25	54.3
	CD	Colon	Involved	35	15	42.9
	CD	Ileum	Uninvolved	14	6	42.9
	CD	Colon	Uninvolved	26	5	19.2
	UC	Colon	Involved	22	2	9.1
	UC	Colon	Uninvolved	20	3	15.0
	Ca	Colon	Uninvolved	15	2	13.3
	Ca	Ileum	Uninvolved	7	3	42.9
	Divertic	Colon	Uninvolved	7	0	0.0
	Append	Colon	Uninvolved	5	0	0.0
	I-Col	Colon	Uninvolved	15	0	0.0
Total				212	61	

FIGURE 3

4/10




FIGURE 4A

5/10




FIGURE 4B




FIGURE 5




FIGURE 6

Medium	Incubation	Organism
Trypticase soy agar	O ₂	Aerobes
McConkey	O ₂	<i>Enterobacteriae</i>
Sabouraud dextrose with Chloamphenical and gentamycin	O ₂	Yeast
Bile eculin agar	O ₂	<i>Enterococcus</i>
Chocolate	CO ₂	<i>Haemophilus</i>
CDC	An O ₂	Anaerobes
Brucella	An O ₂	Anaerobes
EYA+neomycin	An O ₂	<i>Clostridium</i>
EYA+heat treatment	An O ₂	<i>Clostridium</i>
CDC+heat treatments	An O ₂	<i>Clostridium</i>
CCFA+heat treatment	An O ₂	<i>C.difficile/Clostridium</i>
EYA+ ethanol treatment	An O ₂	<i>Clostridium</i>
CDC+ ethanol treatment	An O ₂	<i>Clostridium</i>
CCFA+ethanol treatment	An O ₂	<i>C.difficile/Clostridium</i>
BBE	An O ₂	BFG
LKV	An O ₂	Pigmenters
<i>Fusobacterium</i> selective medium	An O ₂	<i>Fusobacterium</i>
PEA	An O ₂	Gpc
CFA	An O ₂	<i>C. difficile</i>
LAMVAB	An O ₂	<i>Lactobacillus</i>
RB	An O ₂	<i>Bifidobacterium</i>
BBE+vancomycin	An O ₂	<i>Bilophila, Sutterella</i>
Modified BGSA for Campyobacter	An O ₂ , 6%O ₂ (37C), 6%O ₂ (42C)	<i>Campyobacter</i>
Campy CVA(CSL)	An O ₂ , 6%O ₂ (37C), 6%O ₂ (42C)	<i>Camplobacter</i>
Modified Skirrow	An O ₂ , 6%O ₂ (37C), 6%O ₂ (42C)	<i>Helicobacter pylori</i>

FIGURE 7

Figure 8




A.**Naïve CD4 T cells**
(CD44^{low}; CD62L^{high})**B.****Memory CD4 T cells**
(CD44^{high}; CD62L^{low}; CD 45RB^{low})

Figure 9

A. I2 (splenic CD4⁺ T cells)



B. PCC (A.E7 cell line)

