




(86) Date de dépôt PCT/PCT Filing Date: 2012/10/25
(87) Date publication PCT/PCT Publication Date: 2013/05/02
(45) Date de délivrance/Issue Date: 2019/12/03
(85) Entrée phase nationale/National Entry: 2014/04/24
(86) N° demande PCT/PCT Application No.: US 2012/061847
(87) N° publication PCT/PCT Publication No.: 2013/063221
(30) Priorité/Priority: 2011/10/25 (US61/551,395)

(51) Cl.Int./Int.Cl. *C07D 403/04* (2006.01),
A61K 31/4184 (2006.01)
(72) Inventeurs/Inventors:
SEPASSI, KIA, US;
RIZZOLIO, MICHELE C., US
(73) Propriétaire/Owner:
JANSSEN PHARMACEUTICA NV, BE
(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : FORMULATIONS DE SEL DE MEGLUMINE DE L'ACIDE 1-(5,6-DICHLORO-1H-BENZO[D]IMIDAZOL-2-YL)-1H-PYRAZOLE-4-CARBOXYLIQUE
(54) Title: MEGLUMINE SALT FORMULATIONS OF 1-(5,6-DICHLORO-1H-BENZO[D]IMIDAZOL-2-YL)-1H-PYRAZOLE-4-CARBOXYLIC ACID



(57) **Abrégé/Abstract:**

The meglumine salt of 1-(5,6-dichloro-1H-benzo[d]imidazol-2-yl)-1H-pyrazole-4-carboxylic acid (compound (1)) and pharmaceutically acceptable formulations thereof are described. Such compounds may be used in pharmaceutical compositions and methods for the treatment of disease states, disorders, and conditions mediated by prolyl hydroxylase activity.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number

WO 2013/063221 A1

(43) International Publication Date
2 May 2013 (02.05.2013)

(51) International Patent Classification:

A61K 31/4184 (2006.01) *C07D 403/04* (2006.01)
A61P 3/10 (2006.01)

(21) International Application Number:

PCT/US2012/061847

(22) International Filing Date:

25 October 2012 (25.10.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/551,395 25 October 2011 (25.10.2011) US

(71) Applicant: **JANSSEN PHARMACEUTICA NV**
[BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).

(72) Inventors; and

(71) Applicants : **SEPASSI, Kia** [US/US]; 924 Hornblend
Street #308, San Diego, California 92109 (US). **RIZZO-**
LIO, Michele C. [CH/US]; 8525 Celtic Court, San Diego,
California 92129 (US).

(74) Agents: **JOHNSON, Philip S.** et al.; JOHNSON & JOHN-
SON, One Johnson & Johnson Plaza, New Brunswick,
New Jersey 08933 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: MEGLUMINE SALT FORMULATIONS OF 1-(5,6-DICHLORO-1H-BENZO[D]IMIDAZOL-2-YL)-1H-PYRAZOLE-4-CARBOXYLIC ACID



compound (1)

(57) Abstract: The meglumine salt of 1-(5,6-dichloro-1H-benzo[d]imidazol-2-yl)-1H-pyrazole-4-carboxylic acid (compound (1)) and pharmaceutically acceptable formulations thereof are described. Such compounds may be used in pharmaceutical compositions and methods for the treatment of disease states, disorders, and conditions mediated by prolyl hydroxylase activity.



WO 2013/063221 A1

5 MEGLUMINE SALT FORMULATIONS OF 1-(5,6-DICHLORO-1H-BENZO[D]IMIDAZOL-2-YL)-1H-PYRAZOLE-4-CARBOXYLIC ACID

CROSS REFERENCE TO RELATED APPLICATION

10 This application claims the benefit of US provisional patent application serial number 61/551,395, filed on October 25, 2011.

FIELD OF THE INVENTION

15 The present invention is directed to the meglumine salt of 1-(5,6-dichloro-1H-benzo[d]imidazol-2-yl)-1H-pyrazole-4-carboxylic acid and related methods of manufacture.

BACKGROUND

20 A family of highly conserved oxygen, iron, and 2-oxoglutarate-dependent prolyl hydroxylase (PHD) enzymes mediate the cells response to hypoxia via post-translational modification of hypoxia-inducible factors (HIF) (Ivan et al., 2001, *Science*, 292:464-68; Jaakkola et al., 2001, *Science*, 292:468-72). Under normoxic conditions, PHD catalyzes the hydroxylation of two conserved proline residues within HIF. As the affinity of PHD for oxygen is within the physiological range of oxygen and oxygen is a necessary co-factor for modifying hydroxylated HIF, PHD is
25 inactivated when oxygen tension is reduced. In this way, HIF is rapidly degraded under normoxic conditions but accumulates in cells under hypoxic conditions or when PHD is inhibited.

30 Four isotypes of PHD have been described: PHD1, PHD2, PHD3, and PHD4 (Epstein et al., 2001, *Cell*, 107:43-54; Kaelin, 2005, *Annu Rev Biochem.*, 74:115-28; Schmid et al., 2004, *J Cell Mol Med.*, 8:423-31). The different isotypes are ubiquitously expressed but are differentially regulated and have distinct physiological roles in the cellular response to hypoxia. There is evidence that the various isotypes have different selectivity for the three different HIF- α sub-types (Epstein et al.,
35 *supra*). In terms of cellular localization, PHD1 is primarily nuclear, PHD2 is primarily cytoplasmic, and PHD3 appears to be both cytoplasmic and nuclear (Metzen E, et al.

5 2003, *J Cell Sci.*, 116(7):1319–26). PHD2 appears to be the predominant HIF- α prolyl hydroxylase under normoxic conditions (Ivan et al., 2002, *Proc Natl Acad Sci. USA*, 99(21):13459–64; Berra et al., 2003, *EMBO J.*, 22:4082–90). The three isotypes have a high degree of amino-acid homology and the active site of the enzyme is highly conserved.

10

Targeted disruption of the PHD enzyme activity by small molecules has potential utility in the treatment of disorders of oxygen sensing and distribution. Examples include but are not limited to: anemia; sickle cell anemia; peripheral vascular disease; coronary artery disease; heart failure; protection of tissue from
15 ischemia in conditions such as myocardial ischemia, myocardial infarction and stroke; preservation of organs for transplant; treatment of tissue ischemia by regulating and/or restoring blood flow, oxygen delivery and/or energy utilization; acceleration of wound healing particularly in diabetic and aged patients; treatment of burns; treatment of infection; bone healing, and bone growth. In addition, targeted
20 disruption of PHD is expected to have utility in treating metabolic disorders such as diabetes, obesity, ulcerative colitis, inflammatory bowel disease and related disorders such as Crohn's disease. (*Recent Patents on Inflammation & Allergy Drug Discovery*, 2009, 3:1-16).

25 HIF has been shown to be the primary transcriptional factor that leads to increased erythropoietin production under conditions of hypoxia (Wang et al., 1993, *supra*). While treatment with recombinant human erythropoietin has been demonstrated to be an effective method of treating anemia, small molecule mediated PHD inhibition can be expected to offer advantages over treatment with
30 erythropoietin. Specifically, the function of other HIF gene products is necessary for hematopoiesis and regulation of these factors increases the efficiency of hematopoiesis. Examples of HIF target gene products that are critical for hematopoiesis include: transferrin (Rolfs et al., 1997, *J Biol Chem.*, 272(32):20055-62), transferrin receptor (Lok et al., 1999, *J Biol Chem.*, 274(34):24147-52; Tacchini
35 et al., 1999, *J Biol Chem.*, 274(34):24142-46) and ceruloplasmin (Mukhopadhyay et al., 2000, *J Biol Chem.*, 275(28):21048-54). Hpcidin expression is also suppressed

5 by HIF (Peyssonnaud et al., 2007, *J Clin Invest.*, 117(7):1926-32) and small molecule inhibitors of PHD have been shown to reduce hepcidin production (Braliou et al., 2008, *J Hepatol.*, 48:801-10). Hepcidin is a negative regulator of the availability of the iron that is necessary for hematopoiesis, so a reduction in hepcidin production is expected to be beneficial to the treatment of anemia. PHD inhibition
10 may also be useful when used in conjunction with other treatments for anemia including iron supplementation and/or exogenous erythropoietin. Studies of mutations in the PHD2 gene occurring naturally in the human population provide further evidence for the use of PHD inhibitors to treat anemia. Two recent reports have shown that patients with dysfunctional mutations in the PHD2 gene display
15 increased erythrocytosis and elevated blood hemoglobin (Percy et al., 2007, *PNAS*, 103(3):654-59; Al-Sheikh et al., 2008, *Blood Cells Mol Dis.*, 40:160-65). In addition, a small molecule PHD inhibitor has been evaluated in healthy volunteers and patients with chronic kidney disease (U.S. Pat. App. No. US2006/0276477, December 7, 2006). Plasma erythropoietin was increased in a dose-dependent
20 fashion and blood hemoglobin concentrations were increased in the chronic kidney disease patients.

Overall accumulation of HIF under hypoxic conditions governs an adaptive up-regulation of glycolysis, a reduction in oxidative phosphorylation resulting in a
25 reduction in the production of hydrogen peroxide and superoxide, optimization of oxidative phosphorylation protecting cells against ischemic damage. Thus, PHD inhibitors are expected to be useful in organ and tissue transplant preservation (Bernhardt et al., 2007, *Methods Enzymol.*, 435:221-45). While benefit may be achieved by administering PHD inhibitors before harvesting organs for transplant,
30 administration of an inhibitor to the organ/tissue after harvest, either in storage (e.g., cardioplegia solution) or post-transplant, may also be of therapeutic benefit.

PHD inhibitors are expected to be effective in preserving tissue from regional ischemia and/or hypoxia. This includes ischemia/hypoxia associated with inter alia:
35 angina, myocardial ischemia, stroke, ischemia of skeletal muscle. Recently, ischemic pre-conditioning has been demonstrated to be a HIF-dependent

5 phenomenon (Cai et al., 2008, *Cardiovasc Res.*, 77(3):463-70). While the concept of pre-conditioning is best known for its protective effects in the heart, it also applies to other tissues including but not limited to: liver, skeletal muscle, liver, lung, kidney, intestine and brain (Pasupathy et al., 2005, *Eur J Vasc Endovasc Surg.*, 29:106-15; Mallick et al., 2004, *Dig Dis Sci.*, 49(9):1359-77). Experimental evidence for the
10 tissue protective effects of PHD inhibition and elevation of HIF have been obtained in a number of animal models including: germ-line knock out of PHD1 which conferred protection of the skeletal muscle from ischemic insult (Aragonés et al., 2008, *Nat Genet.*, 40(2):170-80), silencing of PHD2 through the use of siRNA which protected the heart from ischemic insult (Natarajan et al., 2006, *Circ Res.*, 98(1):133-40),
15 inhibition of PHD by administering carbon monoxide which protected the myocardium from ischemic injury (Chin et al., 2007, *Proc Natl Acad Sci. U.S.A.*, 104(12):5109-14), hypoxia in the brain which increased the tolerance to ischemia (Bernaudin et al., 2002, *J Cereb Blood Flow Metab.*, 22(4):393-403). In addition, small molecule inhibitors of PHD protect the brain in experimental stroke models
20 (Siddiq et al., 2005, *J Biol Chem.*, 280(50):41732-43). Moreover, HIF up-regulation has also been shown to protect the heart of diabetic mice, where outcomes are generally worse (Natarajan et al., 2008, *J Cardiovasc Pharmacol.*, 51(2):178-187). The tissue protective effects may also be observed in Buerger's disease, Raynaud's disease, and acrocyanosis.

25

The reduced reliance on aerobic metabolism via the Krebs's cycle in the mitochondria and an increased reliance on anaerobic glycolysis produced by PHD inhibition may have beneficial effects in normoxic tissues. It is important to note that PHD inhibition has also been shown to elevate HIF under normoxic conditions.
30 Thus, PHD inhibition produces a pseudohypoxia associated with the hypoxic response being initiated through HIF but with tissue oxygenation remaining normal. The alteration of metabolism produced by PHD inhibition can also be expected to provide a treatment paradigm for diabetes, obesity and related disorders, including co-morbidities.

35

5 Globally, the collection of gene expression changes produced by PHD inhibition reduce the amount of energy generated per unit of glucose and will stimulate the body to burn more fat to maintain energy balance. The mechanisms for the increase in glycolysis are discussed above. Other observations link the hypoxic response to effects that are expected to be beneficial for the treatment of
10 diabetes and obesity. Hypoxia and hypoxia mimetics such as desferrioxamine have been shown to prevent adipocyte differentiation (Lin et al., 2006, *J Biol Chem.*, 281(41):30678-83; Carrière et al., 2004, *J Biol Chem.*, 279(39):40462-69). Inhibition of PHD activity during the initial stages of adipogenesis inhibits the formation of new adipocytes (Floyd et al., 2007, *J Cell Biochem.*, 101:1545-57). Hypoxia, cobalt
15 chloride and desferrioxamine elevated HIF and inhibited PPAR gamma 2 nuclear hormone receptor transcription (Yun et al., 2002, *Dev Cell.*, 2:331-41). As PPAR gamma 2 is an important signal for adipocyte differentiation, PHD inhibition can be expected to inhibit adipocyte differentiation. These effects were shown to be mediated by the HIF-regulated gene DEC1/Stra13 (Yun et al., *supra*).

20

Small molecular inhibitors of PHD have been demonstrated to have beneficial effects in animal models of diabetes and obesity (Intl. Pat. App. Pub. No. WO2004/052284, June 24, 2004; WO2004/052285, June 24, 2004). Among the effects demonstrated for PHD inhibitors in mouse diet-induced obesity, db/db mouse
25 and Zucker fa/fa rat models were lowering of: blood glucose concentration, fat mass in both abdominal and visceral fat pads, hemoglobin A1c, plasma triglycerides, body weight as well as changes in established disease bio-markers such as increases in the levels of adrenomedullin and leptin. Leptin is a known HIF target gene product (Grosfeld et al., 2002, *J Biol Chem.*, 277(45):42953-57). Gene products involved in
30 the metabolism in fat cells were demonstrated to be regulated by PHD inhibition in a HIF-dependent fashion (Intl. Pat. App. Pub. No. WO2004/052285, *supra*). These include apolipoprotein A-IV, acyl CoA thioesterase, carnitine acetyl transferase, and insulin-like growth factor binding protein (IGFBP)-1.

35 PHD inhibitors are expected to be therapeutically useful as stimulants of vasculogenesis, angiogenesis, and arteriogenesis. These processes establish or

5 restore blood flow and oxygenation to the tissues under ischemia and/or hypoxia conditions (Semenza et al., 2007, *J Cell Biochem.*, 102:840-47; Semenza, 2007, *Exp Physiol.*, 92(6):988-91). It has been shown that physical exercise increases HIF-1 and vascular endothelial growth factor in experimental animal models and in humans (Gustafsson et al. 2001, *Front Biosci.*, 6:D75-89) and consequently the number of
10 blood vessels in skeletal muscle. VEGF is a well-known HIF target gene product that is a key driver of angiogenesis (Liu et al., *supra*). PHD inhibition offers a potential advantage over other angiogenic therapies in that it stimulates a controlled expression of multiple angiogenic growth factors in a HIF-dependent fashion including but not limited to: placental growth factor (PLGF), angiopoietin-1
15 (ANGPT1), angiopoietin-2 (ANGPT2), platelet-derived growth factor beta (PDGFB) (Carmeliet, 2004, *J Intern Med.*, 255:538-61; Kelly et al., 2003, *Circ Res.*, 93:1074-81) and stromal cell derived factor 1 (SDF-1) (Ceradini et al., 2004, *Nat Med.*, 10(8):858-64). Expression of angiopoietin-1 during angiogenesis produces leakage-resistant blood vessels, in contrast to the vessels produced by administration of
20 VEGF alone (Thurston et al., 1999, *Science*, 286:2511-14; Thurston et al., 2000, *Nat Med.*, 6(4):460-3; Elson et al., 2001, *Genes Dev.*, 15(19):2520-32). Stromal cell derived factor 1 (SDF-1) has been shown to be critical to the process of recruiting endothelial progenitor cells to the sites of tissue injury. SDF-1 expression increased the adhesion, migration and homing of circulating CXCR4-positive progenitor cells to
25 ischemic tissue. Furthermore inhibition of SDF-1 in ischemic tissue or blockade of CXCR4 on circulating cells prevents progenitor cell recruitment to sites of injury (Ceradini et al., 2004, *supra*; Ceradini et al., 2005, *Trends Cardiovasc Med.*, 15(2):57-63). Importantly, the recruitment of endothelial progenitor cells to sites of injury is reduced in aged mice and this is corrected by interventions that increase
30 HIF at the wound site (Chang et al., 2007, *Circulation*, 116(24):2818-29). PHD inhibition offers the advantage not only of increasing the expression of a number of angiogenic factors but also a co-ordination in their expression throughout the angiogenesis process and recruitment of endothelial progenitor cells to ischemic tissue.

35

5 PHD inhibitors are useful in pro-angiogenic therapies, too. Adenovirus-mediated over-expression of HIF has been demonstrated to induce angiogenesis in non-ischemic tissue of an adult animal (Kelly et al., 2003, *Circ Res.*, 93(11):1074-81) providing evidence that therapies that elevate HIF, such as PHD inhibition, will induce angiogenesis. Placental growth factor (PLGF), also a HIF target gene, has
10 been shown to play a critical role in angiogenesis in ischemic tissue (Carmeliet, 2004, *J Intern Med.*, 255(5):538-61; Luttun et al., 2002, *Ann N Y Acad Sci.*, 979:80-93). The potent pro-angiogenic effects of therapies that elevate HIF have been demonstrated, via HIF over-expression, in skeletal muscle (Pajusola et al., 2005, *FASEB J.*, 19(10):1365-7; Vincent et al., 2000, *Circulation*, 102:2255-61) and in the
15 myocardium (Shyu et al., 2002, *Cardiovasc Res.*, 54:576-83). The recruitment of endothelial progenitor cells to the ischemic myocardium by the HIF target gene SDF-1 has also been demonstrated (Abbott et al., 2004, *Circulation*, 110(21):3300-05). Thus, PHD inhibitors will likely be effective in stimulating angiogenesis in the setting of tissue ischemia, particularly muscle ischemia. Therapeutic angiogenesis
20 produced by PHD inhibitors will likely lead to restoring blood flow to tissues and therefore meliorate such diseases as but not limited to angina pectoris, myocardial ischemia and infarction, peripheral ischemic disease, claudication, gastric and duodenal ulcers, ulcerative colitis, and inflammatory bowel disease.

25 PHD and HIF play a central role in tissue repair and regeneration including healing of wounds and ulcers. Recent studies have demonstrated that an increased expression of all three PHDs at wound sites in aged mice with a resulting reduction in HIF accumulation (Chang et al., *supra*). Thus, elevation of HIF in aged mice by administering desferrioxamine increased the degree of wound healing back to levels
30 observed in young mice. Similarly, in a diabetic mouse model, HIF elevation was suppressed compared to non-diabetic litter mates (Mace et al., 2007, *Wound Repair Regen.*, 15(5):636-45). Topical administration of cobalt chloride, a hypoxia mimetic, or over-expression of a murine HIF that lacks the oxygen-dependent degradation domain and thus provides for a constitutively active form of HIF, resulted in
35 increased HIF at the wound site, increased expression of HIF target genes such as VEGF, Nos2, and Hmox1 and accelerated wound healing. The beneficial effect of

5 PHD inhibition is not restricted to the skin and small molecule inhibitors of PHD have recently been demonstrated to provide benefit in a mouse model of colitis (Robinson et al., 2008, *Gastroenterology*, 134(1):145-55).

In summary, PHD inhibition resulting in accumulation of HIF likely acts by at
10 least four mechanisms to contribute to accelerated and more complete healing of wounds: 1) protection of tissue jeopardized by hypoxia and/or ischemia, 2) stimulation of angiogenesis to establish or restore appropriate blood flow to the site, 3) recruitment of endothelial progenitor cells to wound sites, 4) stimulation of the release of growth factors that specifically stimulate healing and regeneration.

15

As PDGF is a HIF gene target (Schultz et al., 2006, *Am J Physiol Heart Circ Physiol.*, 290(6):H2528-34; Yoshida et al., 2006, *J Neurooncol.*, 76(1):13-21), PHD inhibition likely increases the expression of endogenous PDGF and produces a similar or more beneficial effect to those produced with PDGF alone. Studies in
20 animals have shown that topical application of PDGF results in increased wound DNA, protein, and hydroxyproline amounts; formation of thicker granulation and epidermal tissue; and increased cellular repopulation of wound sites. PDGF exerts a local effect on enhancing the formation of new connective tissue. The effectiveness of PHD inhibition is likely greater than that produced by PDGF due to the additional
25 tissue protective and pro-angiogenic effects mediated by HIF.

The beneficial effects of inhibition of PHD extends not only to accelerated wound healing in the skin and colon but also to the healing of other tissue damage including but not limited to gastrointestinal ulcers, skin graft replacements, burns,
30 chronic wounds and frost bite.

Stem cells and progenitor cells are found in hypoxic niches within the body and hypoxia regulates their differentiation and cell fate (Simon et al., 2008, *Nat Rev Mol Cell Biol.*, 9:285-96). Thus, PHD inhibitors may be useful to maintain stem cells
35 and progenitor cells in a pluripotent state and to drive differentiation to desired cell types. Stem cells may be useful in culturing and expanding stem cell populations

5 and may hold cells in a pluripotent state while hormones and other factors are administered to the cells to influence the differentiation and cell fate.

A further use of PHD inhibitors in the area of stem cell and progenitor cell therapeutics relates to the use of PHD inhibitors to condition these cells to withstand
10 the process of implantation into the body and to generate an appropriate response to the body to make the stem cell and progenitor cell implantation viable (Hu et al., 2008, *J Thorac Cardiovasc Surg.*, 135(4):799-808). More specifically PHD inhibitors may facilitate the integration of stem cells and draw in an appropriate blood supply to sustain the stem cells once they are integrated. This blood vessel formation will also
15 function to carry hormones and other factors released from these cells to the rest of the body.

PHD inhibitors may also be useful in the treatment of infection (Peyssonnaud et al., 2005, *J Invest Dermatol.*, 115(7):1806-15; Peyssonnaud et al., 2008 *J Invest*
20 *Dermatol.*, 2008 Aug;128(8):1964-8). HIF elevation has been demonstrated to increase the innate immune response to infection in phagocytes and in keratinocytes. Phagocytes in which HIF is elevated show increased bacteriicidal activity, increased nitric oxide production and increased expressed of the anti-bacterial peptide cathelicidin. These effects may also be useful in treating infection
25 from burns.

HIF has also been shown to be involved in bone growth and healing (Pfander D et al., 2003 *J Cell Sci.*, 116(Pt 9):1819-26., Wang et al., 2007 *J Clin Invest.*, 117(6):1616-26.) and may therefore be used to heal or prevent fractures. HIF
30 stimulates glycolysis to provide energy to allow the synthesis of extracellular matrix of the epiphyseal chondrocytes under a hypoxic environment. HIF also plays a role in driving the release of VEGF and angiogenesis in bone healing process. The growth of blood vessels into growing or healing bone can be the rate limiting step in the process.

35

5 Small molecules inhibitors of PHD have been described in the literature,
which include, but are not limited to, imidazo[1,2-*a*]pyridine derivatives (Warshakoon
et al., 2006, *Bioorg Med Chem Lett.*, 16(21):5598-601), substituted pyridine
derivatives (Warshakoon et al., 2006, *Bioorg Med Chem Lett.*, 16(21):5616-20),
pyrazolopyridines (Warshakoon et al., 2006, *Bioorg Med Chem Lett.*, 16(21):5687-
10 90), bicyclic heteroaromatic N-substituted glycine derivatives (Intl. Pat. App. Pub. No.
WO2007/103905, September 13, 2007), quinoline based compounds (Intl. Pat. App.
Pub. No. WO2007/070359, June 21, 2007), pyrimidinetrione N-substituted glycine
derivatives (Intl. Pat. App. Pub. No. WO2007/150011, December 27, 2007),
substituted aryl or heteroaryl amide compounds (U.S. Pat. App. Pub. No. US
15 2007/0299086, December 27, 2007) and substituted 4-hydroxypyrimidine-5-
carboxamides (Intl. Pat. App. Pub. No. WO2009/117269, September 24, 2009).


SUMMARY OF THE INVENTION

The invention is directed to the general and preferred embodiments defined, as
20 set forth herein. Preferred and exemplary features of the invention will be apparent
from the detailed description below and with reference to the drawing figures.

In its many embodiments, the present invention relates to a novel salt of an
inhibitor of prolyl hydroxylase (PHD) enzymes, and a method of treatment,
25 prevention, inhibition or amelioration of one or more diseases disorders associated
with PHD enzymes is provided.

More particularly, the present invention relates to the meglumine salt of a
compound of the following formula:

30



compound (1)

and related methods of preparation or manufacture of the compound.

5

In another embodiment, the present invention relates to the hydrated form of the meglumine salt of compound (1).

10 In another embodiment, the present invention relates to a formulation comprising the meglumine salt of 1-(5,6-dichloro-1H-benzo[d]imidazol-2-yl)-1H-pyrazole-4-carboxylic acid, and a pharmaceutically acceptable excipient; wherein said meglumine salt is in the form of a dihydrate.

15 In another embodiment, the present invention relates to compound (1) in the form of a meglumine salt; wherein said compound is in the form of a dihydrate.

Additional embodiments and advantages of the invention will become apparent from the detailed discussion, schemes, examples, and claims below.

20

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the pH-dependency of saturation of compound (1) in solution.

25 Figure 2 shows PXRD data of the meglumine salt of compound (1).

Figure 3 shows DSC, TGA and x-ray data of the meglumine salt of compound (1).

30 Figure 4 shows: (A) the single crystal structure of the meglumine salt of compound (1); and (B) the experimental and stimulated powder pattern of a single crystal for the meglumine salt of compound (1).

Figure 5 shows the percentage stimulation of HIF1- α upon exposure of formulations of the meglumine salt of compound (1).

35

5 Figure 6 shows plasma levels (systemic burden) in wounded mice after topical application of formulations of the meglumine salt of compound (1).


 Figure 7 shows the correlation of the flux of compound (1) across skin (human dermis) and an artificial membrane using a Franz diffusion cell upon application of a
10 formulation of meglumine salt of compound (1).

 Figure 8 shows no to very low irritation caused by application of a formulation of the meglumine salt of compound (1) as tested in a HET-CAM assay.

5

DETAILED DESCRIPTION OF THE INVENTION

This invention relates to a novel salt of a compound of the following formula:



compound (1)

10 that is a inhibitor of prolyl hydroxylase (PHD) enzymes, and compositions thereof for the treatment, amelioration or inhibition of disorders and diseases related to the modulation of a prolyl hydroxylase enzyme. The present invention also relates to methods of making such a compound, pharmaceutical compositions, pharmaceutically acceptable salts, pharmaceutically acceptable prodrugs, and
15 pharmaceutically active metabolites thereof.

A) Terms

The present invention is best understood by reference to the following definitions, the drawings and exemplary disclosure provided herein.

20

The terms “**comprising**”, “**containing**”, and “**including**,” are used herein in their open, non-limiting sense.

“**Administering**” or “**administration**” means providing a drug to a patient in a
25 manner that is pharmacologically useful.

“**Composition**” means a product containing a compound of the present invention (such as a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from such
30 combinations of the specified ingredients in the specified amounts).

“**Compound**” or “**drug**” means a compound of Formula (1) or pharmaceutically acceptable forms thereof.

5

“**Forms**” means various isomers and mixtures of one or more compounds of Formula (1) and salts or hydrates thereof. The term “**isomer**” refers to compounds that have the same composition and molecular weight but differ in physical and/or chemical properties. Such substances have the same number and kind of atoms but
10 differ in structure. The structural difference may be in constitution (geometric isomers) or in an ability to rotate the plane of polarized light (stereoisomers). The term “**stereoisomer**” refers to isomers of identical constitution that differ in the arrangement of their atoms in space. Enantiomers and diastereomers are stereoisomers wherein an asymmetrically substituted carbon atom acts as a chiral
15 center. The term “**chiral**” refers to a molecule that is not superposable on its mirror image, implying the absence of an axis and a plane or center of symmetry.

The term “**hypoxia**” or “**hypoxic disorder**” refers to a condition where there is an insufficient level of oxygen provided in the blood or to tissues and organs.
20 Hypoxic disorders can occur through a variety of mechanisms including where there is an insufficient capacity of the blood to carry oxygen (i.e. anemia), where there is an inadequate flow of blood to the tissue and/or organ caused by either heart failure or blockage of blood vessels and/or arteries (i.e. ischemia), where there is reduced barometric pressure (i.e. elevation sickness at high altitudes), or where dysfunctional
25 cells are unable to properly make use of oxygen (i.e. hystotoxic conditions). Accordingly, one of skill in the art would readily appreciate the present invention to be useful in the treatment of a variety of hypoxic conditions including anemia, heart failure, coronary artery disease, thromboembolism, stroke, angina and the like.

30 “**Patient**” or “**subject**” means an animal, preferably a mammal, more preferably a human, in need of therapeutic intervention.

“**Pharmaceutically acceptable**” means molecular entities and compositions that are of sufficient purity and quality for use in the formulation of a composition or
35 medicament of the present invention. Since both human use (clinical and over-the-counter) and veterinary use are equally included within the scope of the present

5 invention, a formulation would include a composition or medicament for either human or veterinary use.

“**Pharmaceutically acceptable excipient**” refers to a substance that is non-toxic, biologically tolerable, and otherwise biologically suitable for administration to a
10 subject, such as an inert substance, added to a pharmacological composition or otherwise used as a vehicle, carrier, or diluent to facilitate administration of an agent and that is compatible therewith. Examples of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils, and polyethylene glycols.

15

“**Pharmaceutically acceptable salt**” means an acid or base salt of the compounds of the invention that is of sufficient purity and quality for use in the formulation of a composition or medicament of the present invention and are tolerated and sufficiently non-toxic to be used in a pharmaceutical preparation.
20 Suitable pharmaceutically acceptable salts include acid addition salts which may, for example, be formed by reacting the drug compound with a suitable pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

25

The term “**solvates**” means those compounds that are formed from the interaction or complexation of such compounds with one or more solvent molecule, either in solution or in solid or crystalline form. The term “**hydrates**” mean solvates, wherein the solvent is water.

30

“**Therapeutically effective amount**” means that amount of compound that elicits the biological or medicinal response in a tissue system, animal or human, that is being sought by a researcher, veterinarian, medical doctor, or other clinician, which includes therapeutic alleviation of the symptoms of the disease or disorder
35 being treated.

5 The term "**treating**" as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, lessening the severity of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "**treatment**", as used herein, unless otherwise indicated, refers to the act of treating.

10

B) Compounds

 The present invention relates to novel salts of compound of Formula (1). In particular, the invention relates to the meglumine salt of compound of Formula (1). In general, the invention relates to all compounds that upon administration to
15 patients in need of treatment of disorders and diseases related to the modulation of a prolyl hydroxylase enzyme.

 Some embodiments of the invention include hydrates, solvates or polymorphs of such compounds, and mixtures thereof, even if such forms are not explicitly stated
20 in the present specification. Preferably, some embodiments of compounds of Formula (1) or pharmaceutically acceptable salts thereof include solvates. More preferably, some embodiments of compounds of Formula (1) or pharmaceutically acceptable salts thereof include hydrates.

25 Yet another embodiment of the invention includes crystalline forms of compounds of Formula (1) or pharmaceutically acceptable salts of compounds of Formula (1) may be obtained as co-crystals.

 In certain embodiments of the invention, compounds of Formula (1) were
30 obtained in a crystalline form. In other embodiments, crystalline forms of compounds of Formula (1) were cubic in nature. In other embodiments, pharmaceutically acceptable salts of compounds of Formula (1) were obtained in a crystalline form. In still other embodiments, compounds of Formula (1) were obtained in one of several polymorphic forms, as a mixture of crystalline forms, as a polymorphic form, or as an
35 amorphous form. In other embodiments, compounds of Formula (1) convert in solution between one or more crystalline forms and/or polymorphic forms.

5 Drug compounds of the present invention also include a mixture of stereoisomers, or each pure or substantially pure isomer. For example, the present compound may optionally have one or more asymmetric centers at a carbon atom containing any one substituent. Therefore, the compound may exist in the form of enantiomer or diastereomer, or a mixture thereof. When the present compound
10 contains a double bond, the present compound may exist in the form of geometric isomerism (cis-compound, trans-compound), and when the present compound contains an unsaturated bond such as carbonyl, then the present compound may exist in the form of a tautomer, and the present compound also includes these isomers or a mixture thereof. The starting compound in the form of a racemic
15 mixture, enantiomer or diastereomer may be used in the processes for preparing the present compound. When the present compound is obtained in the form of a diastereomer or enantiomer, they can be separated by a conventional method such as chromatography or fractional crystallization. In addition, the present compound includes an intramolecular salt, hydrate, solvate or polymorphism thereof. Suitable
20 drug compounds are those that exert a local physiological effect, or a systemic effect, either after penetrating the mucosa, dermis or – in the case of oral administration – after transport to the gastrointestinal tract with saliva.

The invention further relates to pharmaceutically acceptable salts of
25 compounds of Formula (1) and methods of using such salts. A pharmaceutically acceptable salt refers to a salt of a free acid or base of the compound that is non-toxic, biologically tolerable, or otherwise biologically suitable for administration to the subject. See, generally, S.M. Berge, et al., "Pharmaceutical Salts", J. Pharm. Sci., 1977, 66:1-19, and Handbook of Pharmaceutical Salts, Properties, Selection, and
30 Use, 2002, Stahl and Wermuth, Eds., Wiley-VCH and VHCA, Zürich. Preferred pharmaceutically acceptable salts are those that are pharmacologically effective and suitable for contact with the tissues of patients without undue toxicity, irritation, or allergic response. A compound may possess a sufficiently acidic group, a sufficiently basic group, or both types of functional groups, and accordingly react with
35 a number of inorganic or organic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt. Examples of pharmaceutically acceptable salts

5 include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrogen-phosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates, succinates, suberates, sebacates, fumarates, maleates,
10 butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates, γ -hydroxybutyrates, glycolates, tartrates, methane-sulfonates, propanesulfonates, naphthalene-1-sulfonates, naphthalene-2-
15 sulfonates, and mandelates.

In the presence of a basic nitrogen, the desired pharmaceutically acceptable salt may be prepared by any suitable method available in the art, for example, treatment of the free base with an inorganic acid, such as hydrochloric acid,
20 hydrobromic acid, hydriodic acid, perchloric acid, sulfuric acid, sulfamic acid, nitric acid, boric acid, phosphoric acid, and the like, or with an organic acid, such as acetic acid, trifluoroacetic acid, phenylacetic acid, propionic acid, stearic acid, lactic acid, ascorbic acid, maleic acid, hydroxymaleic acid, malic acid, pantoic acid, isethionic acid, succinic acid, valeric acid, fumaric acid, saccharinic acid, malonic acid, pyruvic
25 acid, oxalic acid, glycolic acid, salicylic acid, oleic acid, palmitic acid, lauric acid, a pyranosidyl acid, such as glucuronic acid or galacturonic acid, an α -hydroxy acid, such as mandelic acid, citric acid, or tartaric acid, an amino acid, such as aspartic acid or glutamic acid, an aromatic acid, such as benzoic acid, 2-acetoxybenzoic acid, naphthoic acid, or cinnamic acid, a sulfonic acid, such as
30 laurylsulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, p-toluenesulfonic acid, methanesulfonic acid, ethanesulfonic acid, hydroxyethanesulfonic, a cyclohexanesulfamic acid, any compatible mixture of acids such as those given as examples herein, and any other acid and mixture thereof that are regarded as equivalents or acceptable substitutes in light of the ordinary level of
35 skill in this technology.

5 In the presence of an acid group, such as a carboxylic acid or sulfonic acid, the desired pharmaceutically acceptable salt may be prepared by any suitable method, for example, treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary or tertiary), an alkali metal hydroxide, alkaline earth metal hydroxide, any compatible mixture of bases such as those given as
10 examples herein, and any other base and mixture thereof that are regarded as equivalents or acceptable substitutes in light of the ordinary level of skill in this technology. Illustrative examples of suitable salts include organic salts derived from amino acids, such as glycine and arginine, ammonia, carbonates, bicarbonates, primary, secondary, and tertiary amines, and cyclic amines, such as benzylamines, pyrrolidines, piperidine, morpholine, and piperazine, and inorganic salts derived from
15 sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum, and lithium. Representative organic or inorganic bases further include benzathine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine, and procaine.

20 The invention also relates to pharmaceutically acceptable prodrugs of the compounds, and treatment methods employing such pharmaceutically acceptable prodrugs. The term "prodrug" means a precursor of a designated compound that, following administration to a subject yields the compound *in vivo* via a chemical or physiological process such as solvolysis or enzymatic cleavage, or under
25 physiological conditions. A "pharmaceutically acceptable prodrug" is a prodrug that is non-toxic, biologically tolerable, and otherwise biologically suitable for administration to the subject. Illustrative procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, 1985, Elsevier.

30

Additional types of prodrugs may be produced, for instance, by derivatizing free carboxyl groups of structures of the compound as amides or alkyl esters. Examples of amides include those derived from ammonia, primary alkyl amines and
35 secondary di-alkyl amines. Secondary amines include 5- or 6-membered heterocycloalkyl or heteroaryl ring moieties. Examples of amides include those that are derived from ammonia, alkyl primary amines, and di-alkyl amines. Examples of

5 esters of the invention include alkyl, cycloalkyl, phenyl, and phenyl-alkyl esters. Preferred esters include methyl esters. Prodrugs may also be prepared by derivatizing free hydroxy groups using groups including hemisuccinates, phosphate esters, dimethylaminoacetates, and phosphoryloxymethyloxycarbonyls, following procedures such as those outlined in Fleisher et al., *Adv. Drug Delivery Rev.*, 1996,
10 19:115-130. Carbamate derivatives of hydroxy and amino groups may also yield prodrugs. Carbonate derivatives, sulfonate esters, and sulfate esters of hydroxy groups may also provide prodrugs. Derivatization of hydroxy groups as acyloxy-methyl and acyloxy-ethyl ethers, wherein the acyl group may be an alkyl ester, optionally substituted with one or more ether, amine, or carboxylic acid
15 functionalities, or where the acyl group is an amino acid ester as described above, is also useful to yield prodrugs. Prodrugs of this type may be prepared as described in Greenwald, et al., *J. Med. Chem.*, 1996, 39 (10):1938-40. Free amines can also be derivatized as amides, sulfonamides or phosphonamides. All of these prodrug moieties may incorporate groups including ether, amine, and carboxylic acid
20 functionalities.

The present invention also relates to pharmaceutically active metabolites of the compounds of Formula (1), which may also be used in the methods of the invention. A "pharmaceutically active metabolite" means a pharmacologically active
25 product of metabolism in the body of the compound or salt thereof. Prodrugs and active metabolites of a compound may be determined using routine techniques known or available in the art. See, e.g., Bertolini, et al., *J. Med. Chem.*, 1997, 40:2011-2016; Shan, et al., *J. Pharm. Sci.*, 1997, 86 (7):765-767; Bagshawe, *Drug Dev. Res.*, 1995, 34:220-230; Bodor, *Adv. Drug Res.*, 1984, 13:224-331; Bundgaard,
30 Design of Prodrugs, 1985, Elsevier Press; and Larsen, Design and Application of Prodrugs, Drug Design and Development, 1991, Krogsgaard-Larsen, et al., eds., Harwood Academic Publishers.

C) Pharmaceutical Compositions

35 In particular embodiments of the invention, the salts of compounds of Formula (1), more particularly the meglumine salt, are used alone, or in combination with one

5 or more additional ingredients, to formulate pharmaceutical compositions. A pharmaceutical composition comprises an effective amount of at least one compound in accordance with the invention.

The disclosure also provides compositions (including pharmaceutical
10 compositions) comprising a compound or derivatives described herein, and one or more of pharmaceutically acceptable carrier, excipient, and diluent. In certain embodiments of the invention, a composition may also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. In a specific embodiment, the pharmaceutical composition is pharmaceutically acceptable for administration to a
15 human. In certain embodiments, the pharmaceutical composition comprises a therapeutically or prophylactically effective amount of a compound or derivative described herein. The amount of a compound or derivative of the invention that will be therapeutically or prophylactically effective can be determined by standard clinical techniques. Exemplary effective amounts are described in more detail in below
20 sections. In certain embodiments of the invention, a composition may also contain a stabilizer. A stabilizer is a compound that reduces the rate of chemical degradation of the composition of compound (1). Suitable stabilizers include, but are not limited to, antioxidants, such as ascorbic acid, pH buffers, or salt buffers.

25 The pharmaceutical compositions can be in any form suitable for administration to a subject, preferably a human subject. In certain embodiments, the compositions are in the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, and sustained-release formulations. The compositions may also be in particular unit dosage forms. Examples of unit dosage forms include, but are
30 not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; ointments; cataplasms (poultices); pastes; powders; dressings; creams; plasters; solutions; patches; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non
35 aqueous liquid suspensions, oil in water emulsions, or a water in oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral

5 administration to a subject; and sterile solids (e.g., crystalline or amorphous solids)
that can be reconstituted to provide liquid dosage forms suitable for parenteral
administration to a subject.

In a specific embodiment, the subject is a mammal such as a cow, horse,
10 sheep, pig, fowl, cat, dog, mouse, rat, rabbit, or guinea pig. In a preferred
embodiment, the subject is a human. Preferably, the pharmaceutical composition is
suitable for veterinary and/or human administration. In accordance with this
embodiment, the term "pharmaceutically acceptable" means approved by a
regulatory agency of the Federal or a state government or listed in the U.S.
15 Pharmacopeia or other generally recognized pharmacopeia for use in animals, and
more particularly for use in humans.

Suitable pharmaceutical carriers for use in the compositions are sterile liquids,
such as water and oils, including those of petroleum, animal, vegetable or synthetic
20 origin. In a specific embodiment, the oil is peanut oil, soybean oil, mineral oil, or
sesame oil. Water is a preferred carrier when the pharmaceutical composition is
administered intravenously. Saline solutions and aqueous dextrose and glycerol
solutions can also be employed as liquid carriers, particularly for injectable solutions.
Further examples of suitable pharmaceutical carriers are known in the art, e.g., as
25 described in Remington's Pharmaceutical Sciences (1990) 18th ed. (Mack
Publishing, Easton Pa.).

Suitable excipients for use in the compositions include starch, glucose,
lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol
30 monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol,
water, and ethanol. Whether a particular excipient is suitable for incorporation into a
pharmaceutical composition depends on a variety of factors well known in the art
including, but not limited to, the route of administration and the specific active
ingredients in the composition.

35

5 Pharmaceutical compositions comprising the compounds or derivatives described herein, or their pharmaceutically acceptable salts and solvates, are formulated to be compatible with the intended route of administration. The formulations are preferably for topical administration, but can be for administration by other means such as by inhalation or insufflation (either through the mouth or the
10 nose), intradermal, oral, subcutaneous, buccal, parenteral, vaginal, or rectal. Preferably, the compositions are also formulated to provide increased chemical stability of the compound during storage and transportation. The formulations may be lyophilized or liquid formulations.

15 **D) Administration**

 A compound or derivative described herein, or a pharmaceutically acceptable salt thereof, is preferably administered as a component of a composition that optionally comprises a pharmaceutically acceptable vehicle. The compound or derivative is preferably administered orally. Another preferred method of
20 administration is via topical application of the compound or derivative.

 In certain embodiments, the compound or derivative is administered by any other convenient route, for example, by absorption through skin, epithelial or mucocutaneous linings (e.g., (epi-)dermis, oral mucosa, rectal, and intestinal
25 mucosa). Methods of administration include but are not limited to parenteral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intranasal, intracerebral, intravaginal, transdermal, rectally, by inhalation, or topically, particularly to the ears, nose, eyes, or skin. In most instances, administration will result in the release of the compound or derivative into
30 the bloodstream. In preferred embodiments, the compound or derivative is delivered orally.

 Furthermore, the invention relates to methods of using the compounds described herein to treat subjects diagnosed with or suffering from a disease,
35 disorder, or condition mediated by prolyl hydroxylase, such as: anemia, vascular disorders, metabolic disorders, and wound healing.

5

In a preferred embodiment, compounds of the present invention are useful in the treatment or prevention of anemia comprising treatment of anemic conditions associated with chronic kidney disease, polycystic kidney disease, aplastic anemia, autoimmune hemolytic anemia, bone marrow transplantation anemia, Churg-Strauss syndrome, Diamond Blackfan anemia, Fanconi's anemia, Felty syndrome, graft versus host disease, hematopoietic stem cell transplantation, hemolytic uremic syndrome, myelodysplastic syndrome, nocturnal paroxysmal hemoglobinuria, osteomyelofibrosis, pancytopenia, pure red-cell aplasia, purpura Schoenlein-Henoch, refractory anemia with excess of blasts, rheumatoid arthritis, Shwachman syndrome, sickle cell disease, thalassemia major, thalassemia minor, thrombocytopenic purpura, anemic or non-anemic patients undergoing surgery, anemia associated with or secondary to trauma, sideroblastic anemia, anemic secondary to other treatment including: reverse transcriptase inhibitors to treat HIV, corticosteroid hormones, cyclic cisplatin or non-cisplatin-containing chemotherapeutics, vinca alkaloids, mitotic inhibitors, topoisomerase II inhibitors, anthracyclines, alkylating agents, particularly anemia secondary to inflammatory, aging and/or chronic diseases. PHD inhibition may also be used to treat symptoms of anemia including chronic fatigue, pallor and dizziness.

25 In another preferred embodiment, molecules of the present invention are useful for the treatment or prevention of diseases of metabolic disorders, including but not limited to diabetes and obesity. In another preferred embodiment, molecules of the present invention are useful for the treatment or prevention of vascular disorders. These include but are not limited to hypoxic or wound healing related diseases requiring pro-angiogenic mediators for vasculogenesis, angiogenesis, and arteriogenesis

In treatment methods according to the invention, an effective amount of a pharmaceutical agent according to the invention is administered to a subject suffering from or diagnosed as having such a disease, disorder, or condition. An "effective amount" means an amount or dose sufficient to generally bring about the

5 desired therapeutic or prophylactic benefit in patients in need of such treatment for the designated disease, disorder, or condition. Effective amounts or doses of the compounds of the present invention may be ascertained by routine methods such as modeling, dose escalation studies or clinical trials, and by taking into consideration routine factors, e.g., the mode or route of administration or drug delivery, the
10 pharmacokinetics of the compound, the severity and course of the disease, disorder, or condition, the subject's previous or ongoing therapy, the subject's health status and response to drugs, and the judgment of the treating physician. An example of a dose is in the range of from about 0.001 to about 200 mg of compound per kg of subject's body weight per day, preferably about 0.05 to 100 mg/kg/day, or about 1 to
15 35 mg/kg/day, in single or divided dosage units (e.g., BID, TID, QID). For a 70-kg human, an illustrative range for a suitable dosage amount is from about 0.05 to about 7 g/day, or about 0.2 to about 2.5 g/day.

Oral tablets may include a compound according to the invention mixed with
20 pharmaceutically acceptable excipients such as inert diluents, disintegrating agents, binding agents, lubricating agents, sweetening agents, flavoring agents, coloring agents and preservative agents. Suitable inert fillers include sodium and calcium carbonate, sodium and calcium phosphate, lactose, starch, sugar, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol, and the like. Exemplary liquid oral
25 excipients include ethanol, glycerol, water, and the like. Starch, polyvinyl-pyrrolidone (PVP), sodium starch glycolate, microcrystalline cellulose, and alginic acid are suitable disintegrating agents. Binding agents may include starch and gelatin. The lubricating agent, if present, may be magnesium stearate, stearic acid or talc. If desired, the tablets may be coated with a material such as glyceryl monostearate or
30 glyceryl distearate to delay absorption in the gastrointestinal tract, or may be coated with an enteric coating.

Capsules for oral administration include hard and soft gelatin capsules. To prepare hard gelatin capsules, compounds of the invention may be mixed with a
35 solid, semi-solid, or liquid diluent. Soft gelatin capsules may be prepared by mixing the compound of the invention with water, an oil such as peanut oil or olive oil, liquid

- 5 paraffin, a mixture of mono and di-glycerides of short chain fatty acids, polyethylene glycol 400, or propylene glycol.

Liquids for oral administration may be in the form of suspensions, solutions, emulsions or syrups or may be presented as a dry product for reconstitution with
10 water or other suitable vehicle before use. Such liquid compositions may optionally contain: pharmaceutically-acceptable excipients such as suspending agents (for example, sorbitol, methyl cellulose, sodium alginate, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminum stearate gel and the like); non-aqueous vehicles, e.g., oil (for example, almond oil or fractionated coconut oil), propylene glycol, ethyl
15 alcohol, or water; preservatives (for example, methyl or propyl p-hydroxybenzoate or sorbic acid); wetting agents such as lecithin; and, if desired, flavoring or coloring agents.

The active agents of this invention may also be administered by non-oral
20 routes. For example, the compositions may be formulated for rectal administration as a suppository. For parenteral use, including intravenous, intramuscular, intraperitoneal, or subcutaneous routes, the compounds of the invention may be provided in sterile aqueous solutions or suspensions, buffered to an appropriate pH and isotonicity or in parenterally acceptable oil. Suitable aqueous vehicles include
25 Ringer's solution and isotonic sodium chloride. Such forms will be presented in unit-dose form such as ampules or disposable injection devices, in multi-dose forms such as vials from which the appropriate dose may be withdrawn, or in a solid form or pre-concentrate that can be used to prepare an injectable formulation. Illustrative infusion doses may range from about 1 to 1000 $\mu\text{g/kg/minute}$ of compound, admixed
30 with a pharmaceutical carrier over a period ranging from several minutes to several days.

For topical administration, the compounds may be mixed with a pharmaceutical carrier at a concentration of about 0.1% to about 10% of drug to
35 vehicle. Examples include lotions, creams, ointments and the like and can be

5 formulated by known methods. Another mode of administering the compounds of the invention may utilize a patch formulation to affect transdermal delivery.

A salt-selection evaluation was carried to identify a salt of compound (1) with properties most suitable for development. The criteria considered essential for the
10 selection process were crystallinity, form reproducibility from a recrystallization process, chemical and physical stability under accelerated conditions and adequate solubility to support both drug substance and drug product development.

In embodiments, the compound is formulated into dosage forms suitable for
15 administration to patients in need thereof. The processes and equipment for preparing drug and carrier particles are disclosed in Pharmaceutical Sciences, Remington, 1985, 17th Ed., 1585-1594; Chemical Engineers Handbook, Perry, 1984, 6th Ed., pp. 21-13 to 21-19 (1984); Parrot et al., 1974, *J. Pharm.Sci.*, 61(6): 813-829; and Hixon et al., 1990, *Chem. Engineering*, pp. 94-103.

20

The amount of compound incorporated in the dosage forms of the present invention may generally vary from about 10% to about 90% by weight of the composition depending upon the therapeutic indication and the desired administration period, e.g., every 12 hours, every 24 hours, and the like. Depending
25 on the dose of compound desired to be administered, one or more of the dosage forms can be administered. Depending upon the formulation, the compound will preferably be in the form of an acetate salt or free base form.

Further, this invention also relates to a pharmaceutical composition or a
30 pharmaceutical dosage form as described hereinbefore for use in a method of therapy or diagnosis of the human or non-human animal body.

This invention also relates to a pharmaceutical composition for use in the manufacture of a pharmaceutical dosage form for oral administration to a mammal in
35 need of treatment, characterized in that said dosage form can be administered at any time of the day independently of the food taken in by said mammal.

5

This invention also relates to a method of therapy or diagnosis of the human or non-human animal body that comprises administering to said body a therapeutically or diagnostically effective dose of a pharmaceutical composition described herein.

10

This invention also relates to a pharmaceutical package suitable for commercial sale comprising a container, a dosage form as described herein, and associated with said package written matter non-limited as to whether the dosage form can be administered with or without food.

15

The following formulation examples are illustrative only and are not intended to limit the scope of the inventions in any way.

EXAMPLES

Five versions of compound (1), namely the free acid, sodium, potassium, tromethamine and meglumine salts were produced and their physical properties and manufacturability potential guided the selection of an preferred form of the compound.

20

E) Example Synthesis

To obtain the compounds described in the examples below and their corresponding analytical data, the following experimental and analytical protocols were adhered to unless otherwise indicated. Unless otherwise stated, reaction mixtures were magnetically stirred at room temperature (rt), solutions were generally “dried” over a drying agent such as Na_2SO_4 or MgSO_4 , and mixtures, solutions, and extracts were typically “concentrated” on a rotary evaporator under reduced pressure.

Data Analysis Setup

Thin-layer chromatography (TLC) was performed using Merck silica gel 60 F₂₅₄ 2.5 cm x 7.5 cm 250 μm or 5.0 cm x 10.0 cm 250 μm pre-coated silica gel


5 plates. Preparative thin-layer chromatography was performed using EM Science silica gel 60 F₂₅₄ 20 cm x 20 cm 0.5 mm pre-coated plates with a 20 cm x 4 cm concentrating zone.

Normal-phase flash column chromatography (FCC) was performed on silica
10 gel (SiO₂) eluting with hexanes/ethyl acetate, unless otherwise noted, whereas reversed-phase HPLC was performed on a Hewlett Packard HPLC Series 1100, with a Phenomenex Luna C₁₈ (5 μ m, 4.6x150 mm) column, and detection was done at λ = 230, 254 and 280 nm with a gradient of 10 to 99% acetonitrile/water (0.05% trifluoroacetic acid) over 5.0 min with a flow rate of 1 mL/min. Alternately,
15 preparative HPLC purification was performed on a Gilson automated HPLC system running Gilson Unipoint LC software with UV peak detection done at λ = 220 nm and fitted with a reverse phase YMC-Pack ODS-A (5 μ m, 30 x 250 mm) column; mobile gradient of 10-99% of acetonitrile/water (0.05% trifluoroacetic acid) over 15-20 min and flow rates of 10-20 mL/min.

20

Mass spectra (MS) were obtained on an Agilent series 1100 MSD equipped with a ESI/APCI positive and negative multimode source unless otherwise indicated, and nuclear magnetic resonance (NMR) spectra were obtained on Bruker model DRX spectrometers with the ¹H NMR data showing chemical shifts in ppm downfield
25 of the tetramethylsilane reference (apparent multiplicity, coupling constant *J* in Hz, integration).

Example 1: Free acid of 1-(5,6-Dichloro-1H-benzoimidazol-2-yl)-1H-pyrazole-4-carboxylic acid (compound (1))



30

Method A:

The free acid of compound (1) was prepared by using 2,5,6-trichloro-1H-benzoimidazole and 1H-pyrazole-4-carboxylic acid. MS (ESI/CI): mass calculated

- 5 for $C_{11}H_6Cl_2N_4O_2$, 297.1; m/z found, 296.0 $[M-H]^-$. 1H NMR (500 MHz, $DMSO-d_6$): 14.18-12.52 (br s, 2H), 8.89 (d, $J = 0.5$ Hz, 1H), 8.31 (d, $J = 0.5$ Hz, 1H), 7.80 (s, 2H).

Method B:

- 10 Step A: 5,6-Dichloro-1,3-dihydro-benzoimidazol-2-one: To the solution of 4,5-dichloro-benzene-1,2-diamine (25 g, 0.14 mol) in dry DMF (200 mL), was added CDI (23 g, 0.14 mol) as the solid. The reaction solution was stirred at room temperature for 1 hour, then water (500 mL) was added. The precipitated solid was collected by filtration, washed with water, dried thoroughly to afford the titled
15 compound (26.0 g, 90%). The crude product was used in the following reaction without further purification.

- Step B: 2,5,6-Trichloro-1H-benzoimidazole: Thoroughly dried 5,6-dichloro-1,3-dihydro-benzoimidazol-2-one (28.4 g, 0.14 mol) was suspended in $POCl_3$ (75 mL). The reaction solution was heated to reflux temperature for 3 hours and cooled
20 to room temperature. The solution was poured into crushed ice/water (1.5 L) slowly with sufficient stirring. The solution was neutralized to pH = 7.0 with NaOH. The precipitated solid was collected by filtration, washed with water, and dried to afford the title compound (27.9 g, 90%). The crude product was used in the following reaction without further purification.

- 25 Step C: 1-(5,6-Dichloro-1-dimethylsulfamoyl-1H-benzoimidazol-2-yl)-1H-pyrazole-4-carboxylic acid ethyl ester. 2,5,6-Trichloro-1H-benzoimidazole 2 (27.6 g, 0.125 mol) was dissolved in dry DMF (200 mL) and then K_2CO_3 (20.7 g, 0.15 mol) and dimethylsulfamoyl chloride (17.9 g, 0.125 mol) were added. The reaction mixture was stirred at room temperature for 16 hours. HPLC analysis showed the
30 complete formation of 2,5,6-trichloro-benzoimidazole-1-sulfonic acid dimethylamide. In the same pot, without isolation of 2,5,6-trichloro-benzoimidazole-1-sulfonic acid dimethylamide, was added 1H-pyrazole-4-carboxylic acid ethyl ester (17.5 g, 0.125 mol) and K_2CO_3 (20.7 g, 0.15 mol). The reaction mixture was stirred at 70 °C for 4 hours and water (500 mL) was added while the reaction solution was still hot. The
35 reaction solution was cooled to room temperature. The precipitated solid was

5 collected via filtration, washed with water and dried. The crude product was used in the following reaction without further purification.

Step D: 1-(5,6-Dichloro-1H-benzoimidazol-2-yl)-1H-pyrazole-4-carboxylic acid. Crude 1-(5,6-Dichloro-1-dimethylsulfamoyl-1H-benzoimidazol-2-yl)-1H-pyrazole-4-carboxylic acid ethyl ester was dissolved in THF (125 mL) and LiOH·H₂O
 10 (21 g, 0.5 mol) in water (250 mL) was added. The reaction mixture was stirred at reflux temperature for 2 hours and cooled to room temperature. Concentrated HCl was added to adjust pH to 2.0. The solid precipitated was collected by filtration, washed with water and dried. The solid was triturated in hot EtOAc (1L). After cooling to room temperature and filtration, the compound of Formula (I) was obtained
 15 as a tan solid (18.5 g, 50%). MS [M+H]⁺ found 297.0. ¹H NMR (500 MHz, DMSO-*d*₆): 13.71 (s, 1H), 12.99 (s, 1H), 8.90 (s, 1H), 8.32 (s, 1H), 7.94 (s, 1H), 7.67 (s, 1H).

The thermal properties, crystalline nature, apparent purity and moisture uptake of a 6.0 g batch of the free acid of compound (1) are summarized in Table 1.
 20 Saturation data for compound (1) is shown in Figure 1.

Table 1

Apparent Purity (HPLC)	Crystallinity (PXRD)	Melting Point (DSC)	Adsorption (40-90% RH)	Desorption (90-0% RH)
99.8%	Weakly crystalline	343 °C ^a	+0.59 %	-0.96%

^a decomposition

25 Example 2: Potassium salt of compound (1)

The potassium salt of 1-(5,6-dichloro-1H-benzoimidazol-2-yl)-1H-pyrazole-4-carboxylic acid was prepared by suspending the free acid (55 g, 1.7 mol) in EtOH (1.5 L) at reflux temperature with K₂CO₃ (12.79 g, 0.85 mol) in 20 mL water added dropwise over 5 min. Strong mechanic stirring was required to ensure proper
 30 agitation. The suspension was stirred at reflux temperature for eight hours and then cooled to room temperature over five hours. The precipitated solid was collected by filtration and quickly washed with 100 mL of water followed by EtOH. The potassium salt was obtained as a white solid (38 g, 65%). Subsequently, the mother liquor was concentrated and the above process was repeated once to give the second crop of

- 5 the potassium salt (13 g, 22%). MS $[M+H]^+ = 297.0$. ^1H NMR (500 MHz, DMSO- d_6): 8.65 (s, 1H), 7.96 (s, 1H), 7.57 (s, 2H).

The potassium salt as prepared by the above re-slurry methodology is non-hygroscopic and consistent with a poorly crystalline hydrate as seen by PXRD and thermal analysis. Two broad endothermic peaks of the potassium salt of compound (1) were seen by DSC that can be associated with a dehydration event and melt/decomposition, respectively (Table 2).

Table 2

Apparent Purity (HPLC)	Crystallinity (PXRD)	Melting Point (DSC)	Adsorption (40-90% RH)	Desorption (90-0% RH)
100.0 %	crystalline monohydrate	277 °C ^d	+0.53%	-0.89%

^ddecomposition

Example 3: Sodium salt of compound (1)

The sodium salt of compound (1) is a poorly crystalline, hydrated solid as shown by PXRD and thermal analysis. The DSC reveals two broad endothermic peaks; the first event is associated with a loss of water (~9% by TGA), while the second endotherm is caused by melting/decomposition of the salt. The sodium salt was prepared in a method similar to that used to prepare the potassium salt (slurry method).

Example 4: Crystallization procedure of the tromethamine salt of compound (1)

Two forms of the tromethamine salt have been produced to date. The first form was obtained from the slurry of compound (1) and tromethamine in aqueous ethanol (14% water). Although not a salt, this physical mixture was not pursued. The second form was produced from an aqueous workup containing excess amounts of counterion. This form was a hydrated salt that was observed to have a lower apparent aqueous solubility than the potassium salt. This compound also exhibited poor bulk properties.

5

Example 5: Crystallization procedure of the meglumine salt of compound (1)

A clear solution (30 mg/mL) of the free acid of compound (1) and 1.2 molar equivalent of meglumine was produced in aqueous methanol (12% water) following slight heating. Room temperature stirring with seeding or refrigeration with or
10 without seeding consistently led to crystallization of the meglumine salt, which was collected via filtration. This methodology was used to produce a 2 g batch of the salt. The solvent composition in the above procedure was modified to aqueous ethanol and used by the PDMS API SM Development team to produce 8.7-kg of GMP-grade material in support of FIH-enabling and FIH studies.

15

The thermal properties, crystalline nature, and apparent purity of the meglumine salt of compound (1) are summarized in Table 3. Figures 2 and 3 show salt bulk properties of the meglumine salt of compound (1), including PXRD data and DSC, TGA and x-ray data, respectively. The single crystal data confirmed the
20 meglumine salt of compound (1) to be a dihydrate with the simulated powder pattern in excellent agreement with the experimental powder pattern as shown in Figures 4A and 4B. The meglumine salt of compound (1) was observed to have improved bulk properties, solubility, and enhanced processability compared to the free acid or the potassium salt of compound (1).

25

Table 3

Sample ID	Apparent Purity (HPLC)	Crystallinity (PXRD)	Melting Point (DSC)
Meglumine salt of compound (1)	≥ 99.9 %	crystalline dihydrate	80 °C

Example 6: Topical formulations of meglumine salt of compound (1)

Materials and excipients that used in the development of topical formulations
30 of a meglumine salt of compound (1) are listed in Table 4.

5

Table 4

Materials
Meglumine salt of compound (1)
Hydroxypropyl Methylcellulose (HPMC K15M)
Poloxamer 407*
Methylcellulose (MC)
PEG4000
Meglumine (NMDG)
Vitamin E-TPGS
Carbomer 941
Carbomer 934P
Carboxymethylcellulose (Na-CMC)
HP-β-CD
Sterile Water for Irrigation

*solubilization at 5 °C due to the thermo-reversible property of the excipient.

Table 5 lists the physical and chemical stability results performed on selected formulations of the meglumine salt of compound (1) after 4 weeks of storage.

10

Table 5

Experiment	Formulation Composition	Meglumine salt of compound (1) remaining (%)		
		2 °C	20 °C	40 °C
1	0.5% HPMC K15M (pH 8.33)	99.89	101.14	101.44
2	1.0% HPMC K15M (pH 8.33)	99.59	99.21	99.08
3	2.0% Methylcellulose (pH 8.32)	98.49	99.71	97.20

4	15% Poloxamer 407 (pH 8.10)	99.19	100.03	87.29 [*]
5	20% Poloxamer 407 (pH 8.03)	97.60	99.57	96.43 [*]
6	VitE- TPGS/PEG4000 /water (20:20:60; pH 8.06)	100.27	100.32	98.87

5

*Precipitation observed in the vial

HPLC analysis showed that the six formulation compositions were chemically stable for four weeks under the studied storage conditions. The apparent loss of mass balance for the poloxamer-based formulation (Experiment 4) was attributed to the precipitation of the free acid of compound (1) at 40° Celsius and not degradation. At 40° Celsius, the polymers degraded that resulted in the formation of acetic acid, aldehydes, and a concurrent loss of viscosity and drop in pH. The resulting acidic environment caused precipitation of the insoluble free acid of compound (1) and discoloration of the formulation. The instability of poloxamer (or lutrol) is known and has been disclosed in Erlandsson, B., 2002, *Polym. Degrad. and Stab.*, 78:571-575. Such a degradation was not observed in samples stored at room temperature or refrigerated.

The results of the solubility screen of formulations of the meglumine salt of compound (1) are shown in Table 6. Four of the ten vehicles investigated, namely 1% Na-CMC (Experiment 1), 1% Carbomer 941 (Experiment 2), 1% Carbomer 934P (Experiment 3) and 20% HP- β -CD (Experiment 4), did not meet the targeted solubility criterion of 10 mg/mL (free acid equivalent) or were toxic to Hela cells. The 1% MC vehicle did not have notable advantages over the product containing 2% MC. The meglumine salt of compound (1) was sufficiently soluble within an acceptable pH range (6-8.5) in Experiments 5 and 7-11.

Table 6

Experiment	Formulation Composition	Compound (1) Solubility
------------	-------------------------	-------------------------

1	1% Na-CMC	< 10 mg/mL
2	1% Carbomer 941	< 10 mg/mL
3	1% Carbomer 934P	< 10 mg/mL
4	0.5% HPMC K15M	≥ 10 mg/mL
5	1.0% HPMC K15M	≥ 10 mg/mL
6	1% Methylcellulose	≥ 10 mg/mL
7	2% Methylcellulose	≥ 10 mg/mL
8	15% Poloxamer 407	≥ 10 mg/mL
9	20% Poloxamer 407	≥ 10 mg/mL
10	VitE-TPGS/PEG4000/water (20:20:60)	≥ 10 mg/mL
11	20% HP-β -CD*	≥ 10 mg/mL

5

*Formulation was toxic to the Hela cells

F) Biological Examples

Cellular Assay for HIF1- α

Hela cells (ATCC, Manassas, VA) were plated in 96-well plates at 20,000 cells per well in 100 μ l of DMEM containing 10% fetal bovine serum, 1% non-essential amino acids, 50 IU/mL of penicillin and 50 μ g/mL of streptomycin (all cell culture reagents from Invitrogen, Carlsbad, CA). 24 hours after plating, changed media to 100 μ l of DMEM without 10% fetal bovine serum, 1.1 μ l of the stock solution for each compound was added and incubated for six hours. All compounds were tested with a final compound concentration of 100 μ M. The supernatant was removed and the cells were lysed in 55 μ l of MSD lysis buffer containing protease inhibitors. 50 μ l of the cell lysate was then transferred to a blocked MSD human HIF-1 α detection plate (Meso-Scale Discovery, Gaithersburg, MD, as per manufacturers protocol), and incubated at room temperature on an orbital shaker for two hour. After three washes in PBS, 25 μ l of 20nM anti-HIF1 α detection antibody was added and incubated for 1 hour at room temperature on an orbital shaker. After three washes in PBS, 150 μ l of 1X read buffer was added and the plate was then read on

10
15
20

5 a MSD SECTOR instrument. Data was analyzed by determining the percent of HIF stimulation in the presence of 100 μ M compound relative to an assay control compound, 7-[(4-Chloro-phenyl)-(5-methyl-isoxazol-3-ylamino)-methyl]-quinolin-8-ol. This biological data for the meglumine salt of compound (1) is presented in Figure 5.


10 Additional biological data for formulations of the meglumine salt of compound (1) is presented in Figures 6 through 8.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood
15 that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

5 What is claimed is:

1. A formulation comprising the meglumine salt of 1-(5,6-dichloro-1H-benzo[d]imidazol-2-yl)-1H-pyrazole-4-carboxylic acid, and a pharmaceutically acceptable excipient;
- 10 wherein said meglumine salt is in the form of a dihydrate.

2. A compound of the formula




- 15 in the form of a meglumine salt;
- wherein said compound is in the form of a dihydrate.

3. A pharmaceutical composition comprising the compound of claim 2 and a pharmaceutically acceptable excipient.

20

4. A topical ointment comprising the formulation of claim 1.

Figure 1**Compound 1**

Properties	Values
pKa	3.67 and 8.57 (exp.)
log D _{7.4}	0.28 (exp.)
MW	297.10




Figure 2




Figure 3






Figure 4**A****B**

Figure 5




Figure 6




Figure 7






Figure 8



Remove Egg Membrane



Treatment

