



(51) International Patent Classification:
C12N 1/20 (2006.01)

(21) International Application Number:
PCT/EE2012/000001

(22) International Filing Date:
23 February 2012 (23.02.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
P201100012 25 February 2011 (25.02.2011) EE

(71) Applicant (for all designated States except US): **OÜ TER-VISLIKU PIIMA BIOTEHNOLOOGIATE ARENDUSKESKUS** [EE/EE]; (Bio-competence Centre Of Healthy Dairy Products), Kreutzwaldi 1, EE-51014 Tartu (EE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KULLISAAR, Tiit** [EE/EE]; Ropka 12A-45, EE-50111 Tartu (EE). **ZILMER, Mihkel** [EE/EE]; Puusepa 37, EE-50406 Tartu (EE). **TAMMSAAR, Ene** [EE/EE]; Sõpruse põik, 'Prillimäe, EE-79702 Rapla Maakond (EE). **VESKIOJA, ANDRE** [EE/EE]; Tammemäe tee 1, Rosma Küla, EE-63221 Põlva Maakond (EE). **SONGISEPP, Epp** [EE/EE]; Tähe 105-7, EE-50101 Tartu (EE). **ZILMER, Kersti** [EE/EE]; Puusepa 37, EE-50406 Tartu (EE). **BOBROVSKI, Lauri** [EE/EE]; Väinjärve Küla, EE-73021 Järvamaa (EE). **EHR-LICH, Kersti** [EE/EE]; Pargi 2-1, EE-50103 Tartu (EE). **RÄTSEP, Merle** [EE/EE]; Alasi 31-49, EE-50109 Tartu

(EE). **PUNAB, Margus** [EE/EE]; Hiie 30, EE-51006 Tartu (EE). **VASAR, Maire** [EE/EE]; J.Hurda 27, EE-51005 Tartu (EE). **MIKELSAAR, Marika** [EE/EE]; Jakobsoni 11-4, EE-51005 Tartu (EE).

(74) Agent: **KAHU, Sirje**; Patendibüroo Ustervall OÜ, Kivi 21-6, EE-51009 Tartu (EE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: ISOLATED MICROORGANISM STRAINS LACTOBACILLUS PLANTARUM MCC1 DSM 23881 AND LACTOBACILLUS GASSERI MCC2 DSM 23882 AND THEIR USE

(57) Abstract: The present invention relates to new microorganism strains *L.plantarum* MCC1 DSM 23881 and *L.gasseri* MCC2 DSM 23882 and their use as antioxidant proteolytic ingredients for the production of food products and dietary supplements. Food products and dietary supplements containing *L.plantarum* MCC1 and *L.gasseri* MCC2 are hypoallergenic, reduce milk allergy and lower urinary tract irritation symptoms accompanying benign prostatic hyperplasia and oxidative stress and inflammation associated with these.



ISOLATED MICROORGANISM STRAINS *LACTOBACILLUS PLANTARUM* MCC1
DSM 23881 AND *LACTOBACILLUS GASSERI* MCC2 DSM 23882 AND THEIR
USE

5

TECHNICAL FIELD

The invention belongs to the field of biotechnology and will
be used in food industry. More precisely the invention deals
10 with the microorganism strains *L. plantarum* MCC1 and *L.*
gasseri MCC2 and their use in the reduction of milk allergy
and irritation symptoms of lower urinary tract accompanying
benign prostate hyperplasia and oxidative stress and
inflammation associated with these.

15

BACKGROUND ART

Lactobacilli have for a long time been widely used to make
healthy food products, for example as starter cultures. The
most common way is the use of lactobacilli in functional
20 food. Functional food is a food product that in addition to
ordinary dietary values has additional natural components
that beneficially influence some functions of the organism or
reduce the risk of diseases.

Methods which include purposefully bioselected live
25 lactobacilli that in the following technological stage are
killed with pasteurization have been less used in preparing
food products. This means that the end product does not
contain live lactic acid bacteria, but with pretreatment,
suitable technological scheme and necessary additives the
30 created product has purposeful bioeffect.

Milk allergy and relieving it

Infections, inflammation-, allergy- and oxidative stress-
related problems, for example food allergies, urinary tract
35 symptoms (e.g. micturition disorders associated with prostate

ailments) are becoming economically more burdensome for the society. For example up to 4% of population is suffering from food allergies and in the USA 100-200 deaths are directly caused by food allergy. Food allergies (including allergic reaction to cow's milk) are an especially serious problem for infants. The use of milk products in infants is essential to guarantee their normal growth and development. The solving of the problem of food allergenicity is very complicated as it is not possible to create products that are absolutely allergic reaction-free and at the same time with very high good bioquality. The last discussions in UN, FAO and WHO have declared - *allergen-free: time to clarity; statement: there is no actual legislation to declare free from all allergens*. Thus: any technological ideas/solutions which result in more hypoallergenic products that would reduce the number of infants with allergic response by at least some percent would be highly rated. Every percent of the reduction of allergic responses is very important in conclusion/longer perspective, as any success in the field of allergy to reduce the allergenicity of food products has significant socio-economic output (including improvement of life quality in children and adults and reduction of treatment costs).

The allergenicity of milk products is determined by proteins, but it is difficult to bring out one so-called main allergen (Encyclopedia of Food Sciences and Nutrition (EFSN) AP, 2003, Wal, J. M. (2004) Bovine milk allergenicity. *Ann.Allergy Asthma Immunol.*, 93, S2-11). Nevertheless both casein and beta-lactoglobulin cause allergic reactions in equal proportion. Cow's milk allergy is mainly associated with α -casein and β -lactoglobulin (Host A. (2002). Frequency of cow's milk allergy in childhood. *Ann Allergy Immunol* 89(Suppl 1):33-7; EFSN, 2003. Many children with milk allergy have a reaction to alpha-casein (Hill D.J, Firer M.A, Shelton M.J et

al. (1986) Manifestation of milk allergy in infancy: clinical and immunologic findings. J Pediatr 109: 270-6; EFSN, 2003, Vandenplas Y, Brueton M, Dupont C, Hill D, Isolauri E, Koletzko S, Oranje AP, Staiano A. (2007) Guidelines for the
5 diagnosis and management of cow's milk protein allergy in infants. Arch Dis Child. 92:902-908).

Casein consists of the following subunits: α -S1 (main allergen), α -S2, β - and κ -casein, whereat α -S1 and β -casein
10 are predominant. Enzymatic hydrolysis of milk proteins reduces their allergenicity. Large peptides are formed during partial hydrolysis of milk proteins, more complete hydrolysis results in a mixture of large and small peptides and amino acids. Even a thorough pepsin-trypsin hydrolysis does not
15 give an allergy-free result as allergic reaction may be caused by negligible amounts of immunoreactive components of native protein. Thus only a thorough hydrolysis-based approach has created some possibilities, but does not offer a final solution, even in case of hydrolyzates of whey proteins
20 (**Businco L**, Cantani A, Longhi MA, Giampietro PG. (1989) Anaphylactic reactions to a cow's milk whey protein hydrolysate (Alfa-Ré, Nestlé) in infants with cow's milk allergy. Ann Allergy. 62(4):333-5.; Sampson HA, James MJ, Bernhisel-Broadbent J. (1992) Safety of an amino-derived
25 infant formula in children allergic to cow milk. Pediatrics. 90,463-465). Thus new approaches to reduce cow's milk allergy are needed, wherefore directed hydrolysis of allergy-causing milk proteins with the strain(s) of microorganism(s) giving additional values to the products, is a promising trend for
30 significant reduction of allergy problems.

The hydrolysis of milk proteins with the help of lactic acid bacteria and additives has been described previously. In the patent EE03424 (patent application W09700078, Valio Oy) the

use of *L. rhamnosus* ATCC 53103 (LGG) strain is described for the production of protein hydrolysate-preparation, where the proteins are hydrolyzed with pepsin and trypsin. The use of LGG for the reduction of allergy risk is considered in several inventions of Nestec SA. In patent EP1296694 the use of LGG for reduction of atopic diseases risk in children is described. In patent application WO2008116907 the use of bifidobacteria and LGG, *L. rhamnosus* CGMCC 1.3724, *L. reuteri* ATCC 55730 and *L. paracasei* CNCM I-2116 for the reduction of allergy risk in infants born with Caesarean section is described, including with partially (2-20%) hydrolyzed proteins. In patent application WO03099037 by Nestec SA a proteolytic system of lactic acid bacteria and a food product containing *Bifidobacterium lactis* ATCC 27536 is described for use among other things to reduce food allergy while the bacteria may be inactivated or dead. In patent EE04724 (patent application WO9929833, Arla Foods) the use of *L. paracasei subsp. paracasei* is described, including for alleviating atopic problems in children. In patent EP1175156 (Bioferme OY) a pasteurized cereal drink containing microorganisms is described. In the patent claim CN101427782 (Min Zhao) bifidobacterium-containing maize drink is described, which is produced with the help of fermentation and where the bacteria are inactivated in the fermented product. In a whey protein concentrate (WPC) the maximum achieved hydrolysis of β -lactoglobulin has been 21% (M.Pescuma et al (2007), Hydrolysis of whey proteins by *L. acidophilus*, *Streptococcus thermophilus* and *L. delbrueckii ssp. bulgaricus* grown in a chemically defined medium, *Journal of Appl Microbiol*, 103,5,1738-1746). Milk protein hydrolysis (2-32%) by two microorganisms with the aim to receive a product with good gustatory properties, is described in the European patent EP1383394 (New Zealand Dairy Board), where one proposed microorganism is *Macroccoccus*, *Microccoccus*,

Enterococcus, *Staphylococcus*, *Brevibacterium*, *Anthrobacter* or *Corynebacterium*, preferably *Macrococcus caseolyticus*, and the other microorganism lactic acid bacterium *Lactococcus*, *Lactobacillus*, *Pediococcus* or *Leuconostoc*, and the
5 fermentation is stopped by removal or killing of the microorganisms. In Japanese patent application JP7203844 (Morinaga Milk Industry Co) partial hydrolysis of milk whey (30%) with the help of an enzyme derived from microorganism *Bacillus subtilis*, trypsin and papain is described.

10

Prostate complaints and relieving them

Prostate may cause several complaints: urinary obstruction and irritation and pain or discomfort in the region of minor pelvis. Prostate complaints, especially those associated with
15 urination, are very common and reduce the quality of life of many men. It is estimated that prostate hyperplasia (Benign Prostatic Hyperplasia - BPH) occurs in about half of 50-year old men and in about 70% of men over 70. The lower urinary tract symptoms (LUTS) accompanying benign prostatic
20 hyperplasia include irritating and obstructive (voiding) symptoms which are assessed using the International Prostate Symptom Score (IPSS). Every non-pharmaceutical possibility to relieve urinary difficulties of men would be valued and it would be the best if a novel food product would have such
25 effect.

In several patent applications the use of lactic acid bacteria has been described in therapeutic compositions for the treatment and alleviation of urogenital infections,
30 including prostate infection, whereas the compositions contain one or several species of lactic acid bacteria (US2008274162, Nessa Jeffrey Bryan et al., 2007).

The combination of lactic acid bacteria, where one component contains *L. crispatus*, *L. salivarius* and *L. casei* and the

other component contains *L. brevis*, *L. gasseri* and *L. fermentum*, may be used as a dietary supplement or pharmaceutical composition for the treatment and prophylaxis of infections and inflammations, including urethritis
5 (EE04620, *Actial Farmaceutica Lda*, 2000). Lactic acid bacterium *L. coprophilus* PL9001 (KCCM-10245) has been used to prevent and treat urogenital infections (KR20040067161, PL BIO Co Ltd., 2003). Pharmaceutical compositions used in urology for local treatment may contain lactic acid bacteria
10 *L. casei*, *L. gasseri* as an active substance (EP353581, Silvana Tosi et al., 1993).

This shows that milk allergy and prostate complaints alleviation, including with the help of partial hydrolysis
15 and lactic acid bacteria, has been studied for a long time.

Irrespective of this there are not sufficient amount of food products and dietary supplements that reduce milk allergy and prostate complaints available on the market. Thus there is a
20 need for production and use of reliable, test-proven food products and dietary supplements with healthy properties.

DISCLOSURE OF THE INVENTION

This invention relates to new isolated antioxidant
25 microorganism strains *L. plantarum* MCC1 DSM 23881 and *L. gasseri* MCC2 DSM 23882, compositions containing one or both strains, the use of the mentioned microorganisms as an antioxidant proteolytic ingredient for production of food product and dietary supplement, the use of microorganisms for
30 production of food product and dietary supplement reducing milk allergy and lower urinary tract irritation symptoms accompanying benign prostatic enlargement and oxidative stress and inflammation associated with these, and method for

producing food product and dietary supplement reducing abovementioned complaints.

Lactic acid bacteria partly hydrolyzing milk proteins, that
5 are the objects of this invention, were found by gradual
bioselection. The aim was to find strains with multivalent
biopotential, the multivalency of which would allow the
synergistic use of the beneficial bioeffects (more
10 hypoallergenic, less allergic responses, inflammation and
oxidative stress reducing effect) to obtain the food products
with desired properties. The approach ideology of the
invention is not the achievement of complete hydrolysis of
proteins. Complete hydrolysis has been attempted, but this
creates new problems including a very severe organoleptic
15 unpleasantness of the product. The use of multibiopotent
strains with required technical procedures and the use of
appropriate healthy additional raw products were chosen as
the ideology of the invention. *L. plantarum* MCC1 and *L.*
gasseri MCC2 are multibiopotent and with synergistic effect -
20 they are able to purposefully (partially) hydrolyze milk
proteins (caseins), they have significant oxidation
resistance, they produce conjugated linoleic acid, NO and
other components. Thus the invention makes it possible with
the help of a biotechnologic integral solution to create food
25 products that would have a potential to reduce milk allergy
and urinary tract complaints.

Description of *L. plantarum* MCC1 and *L. gasseri* MCC2

The objects of the invention, *L. plantarum* MCC1 DSM 23881 and
30 *L. gasseri* MCC2 DSM 23882 were isolated from the faeces of a
healthy child during an Estonian-Swedish comparative study of
children's microflora. Strain *L. plantarum* MCC1 and strain *L.*
gasseri MCC2 were isolated by plating serial dilutions of
faeces of a 1-year old child (10^{-2} - 10^{-7} in phosphate buffer

with 0.04% thioglycolic acid, pH 7.2). The dilutions were plated on a freshly prepared MRS agar culture media (Oxoid, UK) and cultivated at 37°C in microaerobic environment. The strains that were the objects of the invention were isolated on the basis of colony and cell morphology characteristic to *Lactobacillus* ssp. This was followed by provisional and thereafter more precise identification described below.

L. plantarum MCC1 and *L. gasseri* MCC2 were deposited in culture collection Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH under numbers DSM 23881 and DSM 23882, respectively, on August 5, 2010.

Culture-morphological characteristics

Culture-morphological characteristics were determined after cultivation in MRS agar and broth (Oxoid) media.

For the cultivation of *L. plantarum* MCC1 MRS broth during 24-48 hours in microaerobic environment was suitable, after what a uniformly turbid growth was seen in the broth. After cultivating for 48 hours on MRS agar culture media at 37°C in microaerobic environment the colonies of *L. plantarum* MCC1 are with a diameter of 2-2.5 mm, white, convex, shiny and with regular margin. The cells are short rods in a chain with partially parallel position. The optimum growth temperature is 37°C; the strain multiplies also at 15°C and 45°C. The pH of optimum growth environment is 6.5. *L. plantarum* MCC1 is catalase- and oxidase-negative, facultatively heterofermentative, does not hydrolyze arginine or produce gas during glucose fermentation.

L. plantarum MCC1 was identified on the basis of API 50CHL System (bioMérieux, France) test-kit as *L. plantarum* (coincidence with type strain: excellent, ID %-99.9, T index 0.83).

Molecular identification of *L. plantarum* MCC1 on the basis of 16S rRNA sequence showed homology with type strain *L. plantarum* Lp-01 (BLAST <http://www.ncbi.nlm.nih.gov/guide/sequence-analysis/>).

5 Carbohydrate fermentation profile of *L. plantarum* MCC1 was the following on the basis of API CHL 50. The strain ferments: ribose, galactose, D-glucose, D-fructose, D-mannose, N-acetyl-glucosamine, amygdalin, arbutin, esculin, salicin, cellobiose, maltose, lactose, saccharose,
10 trechalose, melecitose, D-raffinose, L-arabinose, methyl- α D-mannopyranoside, D-turanose, melibiose, sorbitol and mannitol.

According to API ZYM (bioMérieux, France) test-kit *L.*
15 *plantarum* MCC1 has got leucin arylamidase, valine arylamidase, α -glucosidase, β -glucosidase activity.

MRS broth during 24-48 hours in anaerobic environment was suitable for the cultivation of *L. gasseri* MCC2, thereafter a
20 uniformly turbid growth was seen in the broth. After cultivating for 48 hours on MRS agar culture media at 37°C in anaerobic environment (CO₂/H₂/N₂:5/5/90) the colonies of *L. gasseri* MCC2 are with a diameter of 1.5-2 mm, off-white, rough and with irregular margin. The cells are short to
25 coccoid rods with medium thickness. The optimum growth temperature is 37°C; the strain multiplies also at 45°C. The pH of optimum growth environment is 6.5. *L. gasseri* MCC2 is catalase- and oxidase-negative, obligately homo-fermentative (OHOL), does not hydrolyze arginine and does not produce gas
30 during glucose fermentation.

L. gasseri MCC2 was identified on the basis of API 50CHL System (bioMérieux, France) test-kit as *L. delbrueckii* ssp. *delbrueckii* (coincidence with type strain: good, ID %-94.6, T

index 0.35). Molecular identification of *L. gasseri* MCC2 on the basis of 16S rRNA sequence showed homology with type strain *L. gasseri* IDCC 3102 (BLAST <http://www.ncbi.nlm.nih.gov/guide/sequence-analysis/>).

5

Carbohydrate fermentation profile of *L. gasseri* MCC2 was the following on the basis of API CHL 50. The strain ferments: lactose, saccharose, trechalose and esculine.

According to API ZYM (bioMérieux, France) test-kit *L. gasseri* MCC2 has got leucin arylamidase, valine arylamidase, α -glucosidase, and β -glucosidase activity.

Safety

The origin of the microbe strains from the digestive tract of a healthy child proves their GRAS (*generally recognised as safe*) status or the microbe strains are safe for humans and suitable for oral administration. Also the species of lactobacilli *L. plantarum* and *L. gasseri* have been entered in the list of taxonomic units with QPS (*Qualified Presumption of Safety*) status by EFSA (EFSA. Introduction of a Qualified Presumption of Safety (QPS) approach for assessment of selected microorganisms referred to EFSA. *EFSA J* 2007; 587, 1-16).

25 Haemolytic activity

Methodology: The haemolytic activity of *L. plantarum* MCC1 and *L. gasseri* MCC2 was determined after incubation in microaerobic environment on blood agar plates (Blood Agar Base nr 2, Oxoid +7% human or equine blood).

30 Result. The strains *L. plantarum* MCC1 and *L. gasseri* MCC2 had no haemolytic activity.

Resistance to antibiotics

Methodology: The susceptibility of *L. plantarum* MCC1 and *L. gasser*i MCC2 to antibiotics was tested with the help of E-test (AB Biodisk, Solna). The minimum inhibitory concentration was determined according to the epidemiological breakpoints recommended by the European Commission.

Table 1. Antibacterial susceptibility of *L. plantarum* MCC1 DSM 23881 and *L. gasser*i MCC2 DSM 23882 (MIC, mg/L)

Antibiotic	MIC* values (µg/mL)				
	<i>L. plan- tarum</i> MCC1	<i>L. gasse- ri</i> MCC2	<i>Lacto- bacillus</i> spp**	<i>L. plan- tarum</i> #	OHOL#
Ampicillin	0.094	0.031	2	2	1
Gentamicin	1.5	2	1	16	16
Streptomycin	16	ND	16	n.r.**	16
Erythromycin	0.125	0.047	4	1	1
Clindamycin	0.064	ND	n.r.	1	1
Tetracycline	2	0.5	16	32	4
Chloramphenicol	1.5	0.75	16	8	4
Quinupristin/ dalfopristin	1	ND	4	4	4
Cefoxitin	256	1.5	4	n.r.	n.r.
Cefuroxim	0.5	0.19	-	n.r.	n.r.
Imipenem	ND	0.047	-	n.r.	n.r.
Ciprofloxacin	0.38	>32	4	n.r.	n.r.
Rifampin	ND	0.064	32	n.r.	n.r.
Trim/sulfam	ND	>32	32	n.r.	n.r.
Vancomycin	256	0,75	4	n.r.	2

*MIC - minimum inhibitory concentration, breakpoint for resistance, OHOL - Obligately homofermentative lactobacilli, n.r. - not required#

** European Commission (EC), 2002. Opinion of the Scientific Committee on Animal Nutrition on the criteria for assessing the safety of micro-organisms resistant to antibiotics of human clinical and veterinary importance. European Commission, Health and Consumer Protection

Directorate General, Directorate C, Scientific Opinions, Brussels, Belgium. http://ec.europa.eu/food/fs/sc/scan/out64_en.pdf

FEEDAP Panel EFSA Technical guidance prepared by the Panel on Additives and Products or Substances used in Animal Feed (FEEDAP) on the update of the criteria used in the assessment of bacterial resistance to antibiotics of human or veterinary importance. *The EFSA Journal* (2008) 732, 1-15

L. plantarum MCC1 turned out to be resistant to cefoxitin and vancomycin according to the indicators established for lactobacilli by the European Commission (EC, 2002. Opinion of the Scientific Committee on Animal Nutrition on the criteria for assessing the safety of micro-organisms resistant to antibiotics of human clinical and veterinary importance. European Commission, Health and Consumer Protection Directorate General, Directorate C, Scientific Opinions, Brussels, Belgium. http://ec.europa.eu/food/fs/sc/scan/out64_en.pdf).

It is known that several species/strains of lactobacilli have natural resistance to the mentioned antibiotics and thus horizontal transfer of antibiotic resistance genes is not expected with the use of this microbe. Resistance to other studied antibiotics did not occur.

The strain *L. gasseri* MCC2 did not have resistance to tested antibiotics.

Functional properties of *L. plantarum* MCC1 and *L. gasseri* MCC2

30 Profile of metabolites

Methodology: The metabolite profile has been established with gas chromatography (HP 6890) following 24 h and 48 h of incubation in microaerobic growth environment (Table 2). Lactobacilli were grown for 48 hours on MRS agar culture media in 10% CO₂ environment, then suspension was prepared into 0.9% NaCl solution with a density 10⁹ microbes/mL

according to McFarland, and 1.0 mL of obtained suspension was inoculated into 9.0 mL of MRS liquid culture media. The amount of metabolites mmol/L was determined using capillary column HP-INNOWax (15 m x 0.25 mm; 0.15 μ m). Column temperature program 60°C 1 min, 20°C/min 120°C 10 min; detector (FID) 350°C.

Table 2. Concentration of acetic acid, lactic acid and succinic acid (mmol/L) in MRS culture media following 24 h and 48 h microaerobic cultivation of *L. plantarum* MCC1 and *L. gasserii* MCC2 in MRS culture media.

Species	Acetic acid (mmol/L)		Lactic acid (mmol/L)		Succinic acid (mmol/L)	
	24h	48h	24h	48h	24h	48h
<i>L. plantarum</i> MCC1	0.9	1.3	100.0	137.2	0.4	0.5
<i>L. gasserii</i> MCC2	0.7	0.7	39.7	48.4	0.3	0.4

15 Determination of *L. plantarum* MCC1 and *L. gasserii* MCC2 aggregation and milk coagulation ability

Methodology: Milk coagulation ability. The ability to coagulate milk was studied after incubating the strains for 48 h in microaerobic conditions in UHT fat-free milk at 37°C. Coagulation ability was assessed according to the following scheme:

1. Strong (+) - milk coagulates during 24 h,
2. Slow (+/-) - milk coagulates during 48 h,
3. Absent (-) - milk doesn't coagulate.

25

Autoaggregation. To assess the ability of autoaggregation the strains previously pre-cultivated for 24 h on MRS agar culture media were suspended in physiological solution. The obtained suspension was kept for 15 minutes at room temperature. Aggregation ability was assessed according to the following scheme: 0 - absent, suspension homogenously

30

turbid; 1 - some flakes in the solution; 2 - lots of flakes in the suspension, slight deposit on the bottom of the test tube; 3 - intensive, no suspension develops, deposit on the bottom, solution almost transparent.

5

L. plantarum MCC1 and *L. gasseri* MCC2 coagulated milk during 24 hours, had strong autoaggregative properties (Table 3).

Table 3. Aggregation, general proteolysis and milk
10 coagulation ability of the strains

Strain	Autoaggregation	Milk coagulation
<i>L. plantarum</i> MCC1	3	+
<i>L. gasseri</i> MCC2	3	+

Fermentation ability of prebiotics

15 To assess the fermentation of Tagatose (Arla Food Ingredients), Raftilose P95 (Beneo Orafiti), Raftilose Synergy 1 (Beneo Orafiti) and Raftilose L60/75 (Orafiti), 0.05 mL of 48 h old culture of the tested *Lactobacillus* sp. was inoculated into MRS culture media, which contained 0.5% of sorbite,
20 tagatose or melecitose instead of glucose. Chlorophenol red (0.004%) was added to the culture media as an indicator. Fermentation was assessed on the basis of colour change (from red to yellow) after 48 h.

25 Table 4. The ability to use prebiotics by *L. plantarum* MCC1 and *L. gasseri* MCC2

	<i>L. plantarum</i> MCC1	<i>L. gasseri</i> MCC2
Tagatose	-	+
L60/75	+	+
Raftilose Synergy 1	+	-
Raftilose P95	+	+

Antimicrobial activity against pathogens

Methodology: For the assessment of antimicrobial properties
30 streak-line method was used (Hütt P, Shchepetova J, Loivukene

K, Kullisaar T, Mikelsaar M. Antagonistic activity of probiotic lactobacilli and bifidobacteria against entero- and uropathogens. J Appl Microbiol. 2006; 100(6):1324-32).

To determine the inhibition of pathogens the growth-free zone was measured in mm. Similarly to Hütt et al (2006) the arithmetic mean and standard error were calculated on the basis of the results of the used sample and the antagonistic activity of the strains (mm) during incubation at 37°C was assessed as follows: - weak <9.7; medium 9.7-12.2; strong >12.2. All tests were repeated in parallel at least three times.

Table 5. Antimicrobial activity of *L. plantarum* MCC1 and *L. gasseri* MCC2 against pathogens with streak-line method during cultivation in anaerobic and microaerobic environments (inhibition of pathogen growth, mm)

Pathogen	<i>L. plantarum</i> MCC1		<i>L. gasseri</i> MCC2	
	Anaerobic	Micro-aerobic	Anaerobic	Micro-aerobic
<i>Shigella sonnei</i>	20.3±2.3	20.3±2.1	9.5±2.9	7.7±1.5
<i>Staphylococcus aureus</i>	15.7±2.9	15.0±1.0	3±3.2	2.3±2.5
<i>Escherichia coli</i>	17.3±2.5	18.0±1.0	6.3±3.4	6.3±1.2
<i>Enterococcus faecalis</i>	16.7±2.1	14.7±2.7	4.5±2.9	2.0±1.0
<i>Salmonella enterica</i> serovar Typhimurium	18.8±2.1	17.8±1.8	7.15±1.6	4.3±2.1
<i>S. enteritidis</i>	nd	7.1±2.8	nd	2.2 ±2.6
<i>Listeria monocytogenes</i>	nd	12.8±1.7	nd	2.8±1.0
<i>Yersinia enterocolitica</i>	nd	12.8±1.0	nd	3.5±2.6

Weak <9.7; medium 9.7-12.2; strong >12.2; nd - not determined

The antimicrobial activity of *L. plantarum* MCC1 in streak line test (killed cells and the effect of their excreted metabolites) was strong against most tested pathogens (except *L. monocytogenes* and *S. enteritidis*) in both environments.

L.gasseri MCC2 turned out to be a weak antagonist in both environments.

General proteolytic properties

- 5 General proteolytic properties of *L. plantarum* MCC1 and *L. gasseri* MCC2 were studied on SMA culture media (skim milk-agar media) and on Ca-caseinate agar, streaking the strains previously pregrown during 48 h in MRS broth onto the test culture media. These were incubated during 7 days in
10 microaerobic environment at 37°C. The strength of proteolysis was assessed (6 tests, arithmetic mean) on the basis of rating system: 4 - strong, 3 - medium, 2 - weak, 1 - very weak, 0 - absent.
- 15 Results: *L. plantarum* MCC1 and *L. gasseri* MCC2 have sufficiently good general proteolytic effect (in a 4-point system *L. plantarum* MCC1 was three points and *L. gasseri* MCC2 two points).
- 20 Bioselection on the basis of more specific proteolytic ability (mass-spectrometric and RP-HPLC technological tests to identify the extent of proteolysis of α -S1-casein and β -casein)
- 25 It was not known for *L. plantarum* MCC1 and *L. gasseri* MCC2 how they hydrolyze different milk proteins. So at first the changes in molecular mass spectra occurring during hydrolysis were ascertained using individual pure milk proteins (β -lactoglobulin, α -S1 casein, β -casein) to analyse the protein
30 specter changes of unskimmed milk later. The molecular masses of milk protein hydrolysis products were determined on the MALDI-TOF (*Matrix-assisted laser desorption/ionization time-of-flight*) mass-spectrometer (Marsilio, R. Catinella, S. Seraglia R. Traldi P. (1995) *Matrix-Assisted Laser-Desorption*

Ionization Mass-Spectrometry for the Rapid Evaluation of Thermal-Damage in Milk. *Rapid Communications in Mass Spectrometry*, 9, 550-552). A suspension with a density of 10^9 CFU/mL according to McFarland 4 Standard was prepared from the cultures into KCl physiological solution. 10 μ l of milk protein fraction was added to obtained suspension. The survival of the strains in the solutions of these fractions was determined with streaking the dilution series before and after the test. The strains were incubated in corresponding protein fraction environment for 2-6 days at 37°C. Then the samples were centrifuged and protein profile was determined on mass-spectrometer (*Voyager DE Pro, Applied Biosystems*) Figure 1 and Figure 2). It was found that the strains that are the objects of the invention are able to adjust metabolism to nutrient-poor environment. For example in the test the only nutrient source was either α -S1 casein, β -lactoglobulin, β -casein or the substances obtained from the strain cells themselves. The strains were detectable at the end of the test in all three protein fraction environments (α -S1 casein, β -lactoglobulin, β -casein).

Then the proteolytic properties of the strains were investigated in marketed milk with 2.5% fat content with the help of RP-HPLC (*Reversed-Phase High-Performance Liquid Chromatography*) method. The extent of proteolysis of α -S1 casein and β -casein was measured on the 2nd and 4th day of incubation (Figure 3). For this the RP-HPLC chromatogram peak area corresponding to investigated milk protein was compared to the start moment (control, i.e. milk before the addition of lactobacilli, was equated with 100%) and the percentage amount of remaining milk protein was calculated. In addition experiments were carried out with longer than 21-day incubation times. Also combinations of the strains that are the objects of this invention were investigated to improve

proteolytic efficacy and proteolytic activity also with whey (to reduce the amount of allergic caseins) in case of prediluted milk. The preparation of milk samples for RP-HPLC analysis and the conditions of chromatographing have been developed along the lines of methodology described by Bobe et al. (Bobe G, Beitz DC, Freeman AE, Lindberg GL. (1998) Separation and Quantification of Bovine Milk Proteins By reversed-Phase High-Performance Liquid Chromatography. J. Agric. Food Chem., 46, 458-463). Marketed milk with 2.5% fat content where 10^7 CFU/mL of investigated strain was added and incubated at 37°C, was compared to milk without the strain. High-performance liquid chromatograph (HPLC) (Hewlett Packard 1100) and reverse phase column Zorbax 300SB-C18 filled with porous silica were used for the separation of milk proteins. The mobile phase was an eluting system with a changing gradient, consisting of: Solution A - acetonitrile, water, TFA in ratio 900:100:1 (v/v/v); Solution B - acetonitrile, water, TFA in ratio 100:900:1 (v/v/v). The gradient starts with solution B and the proportion of solution A starts to increase immediately after the sample is injected into the system. During separation phase, when the milk protein separation occurs, the content of solution A increases by 0.47% per minute. The obtained fractions were identified by comparing the retention times with the retention times of the main milk protein standards and determining the molecular mass with MALDI-TOF mass-spectrometer. Milk proteins elute in the next sequence: κ -CN, α_{s2} -CN, α_{s1} -CN, β -CN, α -LA and β -LG. *L. plantarum* MCC1 and *L. gasseri* MCC2 had appropriate effects in relation to α -S1 casein and β -casein (see summary data in Figure 1 and Figure 2), the amount of proteins dominating in the children's allergy reply to cow's milk, α -S1 casein and β -casein, was reduced on the 2nd day. On the basis of RP-HPLC analysis the best directed partial proteolyzer was *L. gasseri* MCC2 (Figure 3). Combining the aforementioned lactobacilli

between themselves, the proteolytic effect in milk was somewhat reduced compared to the individual use of the same strains. *L. plantarum* MCC1 was also incubated for 4 days in a mixture consisting of 40% milk and 60% whey: the content of α -casein was on the fourth day 81% and the content of β -casein 83% compared to the starting moment.

The set objective was achieved: the strains *L. plantarum* MCC1 and *L. gasseri* MCC2 and their combination reduced the amount of α -S1 casein and/or β -casein by 20-40% (i.e. directed moderate hydrolysis took place).

L. plantarum MCC1 and *L. gasseri* MCC2 hydrolyze the proteins causing milk allergy and they have a good general proteolytic activity. Their use separately or together reduces milk allergy and therefore they can be used to produce a milk allergy reducing food product and dietary supplement with healthy properties.

Bioselection on the basis of conjugated linoleic acid (CLA) production

CLA stands for a group of 18-carbon fatty acid, linoleic acid (LA, *cis*-9, *cis*-12-18:2) isomers, where the double bonds are conjugated. CLA is formed naturally during the processes of biohydrogenation and oxidation. CLA is thought to have anti-adipogenous, anti-cancerogenous, anti-diabetogenous and anti-inflammatory effect, regulatory effect on immune system, production of cytokines and immunoglobulins and an ability to modulate the expression of certain genes directly or indirectly through specific transcription factors (Wahle, K. W. J., Heys, S. D. and Rotondo, D. (2004). Conjugated linoleic acids: are they beneficial or detrimental to health? Prog. Lipid Res. 43: 553-587). As several lactic acid bacteria are able to change linoleic acid into conjugated linoleic acid, then it would be possible to increase the CLA

content in fermented milk products (Sieber, R., Collomb, M., Aeschlimann, A., Jelen, P. and Eyer, H. (2004). Impact of microbial cultures on CLA in dairy products- a review. Int. Dairy J. 14: 1-15). For this it should be tested, which

5 lactic acid bacteria are CLA producers. It is known that free fatty acids occurring in the growth environment of microbes have bacteriostatic effect, influencing microbe growth and cell membrane permeability. The strength of the effect depends on the fatty acid. Long-chain fatty acids with higher

10 level of saturation have stronger antimicrobial effect. The stereochemistry of double bonds also influences the antimicrobial activity: a fatty acid with *cis*-configuration is a stronger inhibitor than *trans*-form (Kankaanpää, P. E., Salminen, S. J., Isolauri, E., Lee, Y. K. (2001). The

15 influence of polyunsaturated fatty acids on probiotic growth and adhesion. FEMS Microbiol. Lett. 194: 149-153). The microbes have an established defence mechanism for counteracting a compound inhibiting its vital functions: polyunsaturated fatty acids are isomerized to less inhibiting

20 variants, for example linoleic acid (LA) with stronger bacteriostatic effect is converted to CLA (Sieber, R., Collomb, M., Aeschlimann, A., Jelen, P. and Eyer, H. (2004). Impact of microbial cultures on conjugated linoleic acid in dairy products- a review. Int. Dairy J. 14: 1-15). Most of

25 the emerged CLA stays in the environment; one part is added to the lipids of cell membrane. It has been found that CLA may in small quantities be located in the microbial cell. The ability to conjugate LA and other fatty acids (oleic acid, ricinolic acid) has been found in the representatives of

30 several microbial genera, including *Lactobacillus* sp. The ability of the strains to produce CLA on the basis of linoleic acid added to the growth environment (MRS, skim milk) was assessed quantitatively using CLA determination method developed by us. It became clear that *L. plantarum*

MCC1 and *L. gasseri* MCC2 that had the needed purposeful proteolytic effect produced CLA significantly (Table 7). At the same time these strains were not practically susceptible to the bacteriostatic effect of LA.

5

Bioselection on the basis of oxidation resistance
(antioxidativity)

It was investigated if the strains *L. plantarum* MCC1 and *L. gasseri* MCC2 that have specific limited proteolytic properties, also CLA-productivity, have physiologically significant oxidation resistance. Oxidation resistance should be determined with at least two methods.

10

Linoleic acid test (LA-test) which makes possible to determine the ability of cell suspensions of lactobacilli to inhibit peroxidation of linoleic acid, was used as the first method.

15

A commercial test (Total Antioxidant Status - TAS, Randox Laboratories Ltd., UK) based on the inhibition of the formation of ferryl metmyoglobin radical generated by hydrogen peroxide by cell suspension of lactobacilli, was used as the second method (Rice-Evans and Miller 1994). TAS value was reported in trolox (soluble form of vitamin E) units (mmol/L). Both methods have been published (Kullisaar T, Zilmer M, Mikelsaar M, Vihalemm T, Annuk H, Kairane C, Kilk A. (2002). Two antioxidative lactobacilli strains as promising probiotics. *Int. J. Food Microbiol.* 72, 215-224; Kullisaar, T., Songisepp, E., Mikelsaar, M., Zilmer, K., Vihalemm, T., Zilmer, M. (2003) Antioxidative probiotic fermented goats milk decreases oxidative stress-mediated atherogenicity in human subjects. *British Journal of Nutrition* 90, 449-456).

20

25

30

For the determination of TAA and TAS the strains were incubated in MRS broth (Oxoid) for 24 h at 37°C. The microbial cells were centrifuged at 4°C and 1500 rpm for 10 minutes, washed with isotonic saline (4°C) and suspended in 1.15% KCl (Sigma, USA). The suspension density at OD₂₆₀ 1.1 was 10⁹ microbial cells/mL. *L. plantarum* MCC1 and *L. gasseri* MCC2 have physiologically significant total antioxidativity (Table 6).

Table 6. Oxidation resistance with TAA and TAS methods (n=7-9; M±SD)

Strain	TAA (%)	TAS (mmol/l)
<i>L. plantarum</i> Inducia DSM 21379	26±12	0.13±0.04
<i>L. plantarum</i> MCC1 DSM 23881	20±8	0.05±0.02
<i>L. gasseri</i> MCC2 DSM 23882	25±6	0.03±0.03
<i>L. plantarum</i> Tensia DSM 21380	22±5	0.05±0.02

Bioselection on the basis of nitrogen oxide and hydrogen peroxide production ability

To obtain purposeful information, the preselected strains (specific limited proteolytic properties, CLA production, antioxidativity) were also investigated in respect to NO and H₂O₂ production. According to the literature there are no previous studies about concurrent production of NO and H₂O₂ by microbial strains. The measurements were carried out with live cells using electrochemical measuring (Apollo 4000 free radical analyzer WPI, Berlin, Germany), using ISO-HPO2 and ISO-NOP type electrodes. The electrodes were placed into culture grown in MRS broth (Oxoid), the signals were recorded concurrently during 5-7 minutes and mean signal strength during the measuring time was calculated. Each experiment point was measured as 4 independent parallels and each parallel was measured twice. The NO and H₂O₂ concentrations in the investigated sample were determined by comparison of sample signal with the standard curve.

L. plantarum MCC1 has significant ability to produce NO and hydrogen peroxide, both compounds were also produced by *L. gasseri* MCC2 (Figures 4 and 5).

Therefore the strains *L. plantarum* MCC1 and *L. gasseri* MCC2 are suitable as antioxidant additives for the production of food product and dietary supplement.

Table 7 shows the summary of functional properties of abovementioned strains and their combination.

Table 7. Summary of functional properties of the strains *L. plantarum* MCC1 and *L. gasseri* MCC2 and their combination

FUNCTIONAL PROPERTIES		MCC1	MCC2
1	<i>In vitro</i> antagonism:#	2	1
2	Production of bactericidal H ₂ O ₂ 1	3	2
3	Production of bactericidal and relaxing factor NO 2	3	2
4	CLA production*** MRS Skim milk	3 3	2 1
5	General proteolytic effect	3	2
	Oxidation resistance & TAA TAS	3 1	3 1
7	Fermentation of prebiotics 4	3	1
8	Production level of biogenic amines 3	3	3
9	Proteolyticity of caseins 5	2	3
	Sum of positive scores	29	21
SAFETY AND POSITIVE EFFECTS		MCC1	MCC1+MCC2
1	Haemolytic activity <i>in vitro</i>	0	
2	Animal test (NIH line mice) Translocation of strain Histological changes	0 0	
3	Safety studies in volunteers, 2 weeks ## Lower abdomen pain Flatulence Summary score	n = 10 0/30SC 30 0/30SC 30 3	n = 9 0/27SC 27 0/27SC 27 3
4	BMI, blood pressure, blood glucose, blood lipids, cholesterol, lipoproteins, liver function, inflammation and OxS markers did not differ from endemic reference values	3	3

5	Statistically true positive changes in systemic inflammation and oxidative stress markers	3 MPO 3 Isoprostanes	3 hsCRP, 3 IL-10 3 Isoprostanes
6	Significant other trends (number of other markers)	3 (oxLDL, B2GPI-LDL, sdLDL)	5 (IgE, BP, oxLDL, b2GPI-LDL, sdLDL)
Sum of positive scores		15	20

#SCORE 3 $\geq$ 20 mm; SCORE 2 = 15-20 mm; SCORE 1 = < 15 mm, ## SCORE 3 - no complaints; SCORE 2 - some complaints; SCORE 1 - changes, complaints, ¹ SCORE 3 - 400-500; SCORE 2- 300 -399; SCORE 1 - 100-299 units, ² SCORE 3 - 2.0 -3.0; SCORE 2 - 1.0 -1.99; SCORE 1- <1.0 units, ⁴ SCORE 3 - Inulin, its derivatives, tagatose SCORE 2 - only inulin or tagatose - no inulin or tagatose, ⁵ SCORE 3 - 30-45%, SCORE 2 - 15-30%, SCORE 1 - < 15%, ³ SCORE 3 - only traces; SCORE 2 - < 10; SCORE 1- 11-30 units (during long-term effect), ⁶ SCORE 3 TAA > 20, TAS >0.1; SCORE 2 - TAA >15, TAS 0.05-0.09; SCORE 1 - TAA <15, TAS < 0.05 units, *** SCORE: CLA production 3 - 40-31, SCORE 2 30 - 20-15, SCORE 1 14 -10 units

Afore-cited described the microorganism strains being the objects of the invention.

15

The next objects of the invention are compositions containing one or both strains *L. plantarum* MCC1 and *L. gasseri* MCC2.

If the composition contains only one strain, the strain content is in the range of 0.8-1.2 volume percent, whereat the germ count of *L. plantarum* MCC1 strain is 0.25×10^8 - 4×10^8 CFU/mL and the germ count of *L. gasseri* MCC2 strain is 0.5×10^7 - 1×10^8 CFU/mL.

When one strain is used the content of the strain is preferably 1.0 volume percent, whereat the germ count of *L. plantarum* MCC1 strain is in the range 0.4×10^8 - 4.8×10^8 CFU/mL and the germ count of *L. gasseri* MCC2 strain is 0.8×10^7 - 1.2×10^8 CFU/mL.

If the composition contains both strains *L. plantarum* MCC1 and *L. gasseri* MCC2 the content of both is in the range of 0.4-0.6 volume percent, whereat the germ count of *L. plantarum* MCC1 strain is in the range 0.2×10^8 - 2.4×10^8 CFU/mL and the germ count of *L. gasseri* MCC2 strain is 0.4×10^7 - 0.6×10^8 CFU/mL.

When both strains are used the content of both strains is preferably at least 0.5 volume percent, whereat the germ count of *L. plantarum* MCC1 strain is in the range 0.25×10^8 - 2×10^8 CFU/mL and the germ count of *L. gasseri* MCC2 strain is
5 0.5×10^7 - 0.5×10^8 CFU/mL.

The aforementioned strains may be live or killed.

The next objects of the invention is separate or concurrent use of the mentioned strains

- as antioxidant proteolytic ingredient for the production
10 of food product and/or dietary supplement,
- for the production of hypoallergenic food product and/or dietary supplement reducing milk allergy,
- for the production of food product and/or dietary
15 supplement reducing lower urinary tract irritation symptoms accompanying benign prostatic hyperplasia and oxidative stress and inflammation associated with these.

The next object of the invention is the method for the production of food product and/or dietary supplement reducing milk allergy and lower urinary tract irritation symptoms and
20 oxidative stress and inflammation associated with these.

The method according to the invention includes the next stages:

A) The initial mixture containing milk protein is warmed to
25 $37 \pm 2^\circ\text{C}$;

B) *L. plantarum* MCC1 and/or *L. gasseri* MCC2 or both together are added. If *L. plantarum* MCC1 is used alone, it will be added up to 1.2 volume percent, preferably 1 volume percent with germ count 0.4×10^8 - 4.8×10^8 CFU/mL. If *L. gasseri* MCC2 is
30 used alone, it will be added up to 1.2 volume percent, preferably 1 volume percent with germ count 0.8×10^7 - 1.2×10^8 CFU/mL. When both abovementioned strains are used together, they will be added up to 1.2 volume percent, preferably 1 volume percent, whereat *L. plantarum* MCC1 will be added

0.25x10⁸-2x10⁸ CFU/mL and *L. gasseri* MCC2 0.5x10⁷-0.5x10⁸ CFU/mL. The end-pH of fermenting is 4.2±0.25.

C) The mixture is fermented by *L. plantarum* MCC1 or *L. gasseri* MCC2 or by *L. plantarum* MCC1 + *L. gasseri* MCC2 at
5 37±2°C at least 18-26 hours.

D) The mixture is pasteurized in the range of 80-85°C (preferably 82°C±2) for 30-35 minutes (preferably 30 minutes) and cooled down to temperature 20-30°C.

E) The obtained mixture is flavoured with additives and
10 cooled down to temperature 2-6°C.

F) The obtained mixture is used for the production of food product and/or dietary supplement. In case of food product the mixture is bottled, stored (at 2-6°C) and all required analyses are performed (determination of oxidation resistance
15 etc.). The mixture may also be used in powder (capsules, lozenges, tablets, powder sachets etc.) or liquid (ampoules) form for the production of marketed dietary supplement.

During the preparation of the dietary supplement in powder form water is removed from the mixture, for example by
20 lyophilization (freeze-drying), spray-drying, mill-drying or other known method (until the powder consistency is obtained). During the preparation of liquid dietary supplement water is removed from the product partially until the desired consistency is obtained). The dietary supplement
25 may be prepared with appropriate additive (e.g. antioxidants, sweeteners, prebiotics) or without.

In the aforementioned method the source of milk protein is milk, milk powder, milk protein concentrate, whey or other
30 milk protein source. Fruit or berry juice, -concentrate, syrup or juice drink or fruit or berry jam, preferably sea buckthorn, blueberry or raspberry juice or other juice, syrup, concentrate are suitable additives.

DESCRIPTION OF THE DRAWINGS

Figure 1 - MALDI-TOF mass-spectrometric analysis (illustrative typical experiment). Beta-casein 30 minutes after mixing with strain *L. gasseri* MCC2.

- 5 Figure 2 - MALDI-TOF mass-spectrometric analysis (illustrative typical experiment). After 48 h incubation with the strain *L. gasseri* MCC2 the mass specter peak of beta-casein 24,0 kDa was substantially eliminated.

Figure 3 - Effect of strains *L. plantarum* MCC1 and *L. gasseri*
10 MCC2 in case of 48 and 96-hour incubation on the level of beta-casein (the basal level was 100%, on the basis of MALDI-TOF mass-spectrometric and RP-HPLC analysis data).

Figure 4 - NO production by strains *L. plantarum* MCC1 and *L. gasseri* MCC2.

- 15 Figure 5 - H₂O₂ production by strains *L. plantarum* MCC1 and *L. gasseri* MCC2.

Figure 6 - The scheme of the production of food product and dietary supplement.

20 DESCRIPTION OF THE EMBODIMENTS

Example 1. Method for production of milk allergy responses and urinary tract irritation symptoms and associated oxidative stress and inflammation reducing food product and dietary supplement with abovementioned healthy properties,
25 containing microorganisms *L. plantarum* MCC1 or *L. gasseri* MCC2 or their combination.

Below is presented the preferable embodiment.

- For the production of food product and/or dietary supplement reducing milk allergy responses and urinary tract complaints
30 microbial cultures were used that were pre-grown in fermenter during 24 h and then lyophilized.

For the production of food product the lyophilized microbial cultures *L. plantarum* MCC1 and/or *L. gasseri* MCC2 were activated in small amount of warm (at least 20°C) whey, milk

or solution made from milk powder/milk protein concentrate or other milk protein source for at least 4 hours.

Method for production of food product or dietary supplement with healthy properties included the next stages (see scheme in figure 6):

a) The initial mixture containing milk protein, whey obtained from cheese industry (chemical-physical properties in table 8), in which fat phase was reduced by separation and cheese dust was removed, was warmed to fermentation temperature $37\pm 2^{\circ}\text{C}$;

b) 1 volume percent of strain *L. plantarum* MCC1 (with germ count 0.5×10^8 – 4×10^8 CFU/mL) or strain *L. gasseri* MCC2 (with germ count 1×10^7 – 1×10^8 CFU/mL) or both together (both 0.5 volume percent containing live cells of strain *L. plantarum* MCC1 with a density in a range 0.25 – 2×10^8 CFU/mL and live cells of strain *L. gasseri* MCC2 with a density in a range 0.5×10^7 – 0.5×10^8 CFU/mL).

Table 8. Chemical-physical properties of whey

Ingredient/parameter	Content
Fat (%)	< 0.1
Protein (%)	0.66 ± 0.10
Lactose (%)	4.07 ± 0.59
Dry substance (%)	5.37 ± 0.27
Density (g/cm^3)	1.0199 ± 0.0002
pH	6.14 ± 0.41

c) The mixture was fermented for 18–26 h at temperature $37\pm 2^{\circ}\text{C}$, during which a partial hydrolysis of milk protein took place due to the effect of strain(s) *L. plantarum* MCC1 and/or *L. gasseri* MCC2. The fermentation lasted until acidity reached the value of $\text{pH } 4.20 \pm 0.25$;

d) To inactivate (kill) the strain(s) *L. plantarum* MCC1 and/or *L. gasseri* MCC2 the mixture was pasteurized at $82\pm 2^{\circ}\text{C}$ for 30–35 minutes and cooled to temperature 20 – 30°C ;

- e) The obtained mixture was flavoured with selected additives, which could be fruit or berry juice, - concentrate, -syrup, juice drink or fruit or berry jam. For example 16-17 volume percent of sea-buckthorn jam or 4.7-5 volume percent of concentrated cherry juice drink and sea-buckthorn juice and 9.8-10.2 volume percent of raspberry-blueberry concentrated juice drink were used.
- f) The obtained mixture was cooled to 2-6°C, the food product was bottled, organoleptic and oxidation resistance analysis performed and stored (2-6°C) and clinical studies were performed (see below).

To produce a powder dietary supplement water was removed from the obtained mixture using methods known in food processing technology (for example with the help of lyophilization or drying (spray-drying, mill drying etc.)). Preferably a one-stage spray drier (air inflow temperature 160°C and air outflow temperature 80°C) was used to produce a powder dietary supplement.

To the mixture produced by the abovementioned way a suitable filling agent was added that had a melting point higher than lactose, for example maltodextrin (in 1:1 ratio with whey dry substance).

Water, sugar (lactose) and salt separation with ultrafiltration was used as another choice, which was followed by concentrating with vacuum machine and drying (spray-drying), and so a powder dietary supplement with low lactose content and high protein content was obtained.

As a third choice concentrating with vacuum device followed by spray-drying was used for water removal. The final result was a powder dietary supplement containing lactose.

Table 9. Chemical-physical properties of food product (whey drink) containing strains MCC1+MCC2, flavoured with sea-

buckthorn jam, and of dietary supplement containing strains MCC1+MCC2, flavoured with sea-buckthorn jam

Ingredient/parameter	Content	
	Whey drink	Dietary supplement
Calculated energetic value of food		
kcal/100g	66	390
kJ/100g	219	1654
Fat (%)	0.41	3.46
Protein (%)	0.90	5.96
Ash (%)	0.46	3.4 1
Dry substance (%)	16.36	96.58
Calculated carbohydrate content (%)	14.6	83.8
Glucose (%)	<0.5(0.1)	1.1
Fructose	<0.5(0.02)	<0.5(0.2)
Saccharose	<0.3(0)	<0.3(0.1)

Food products FP-I - FP-V containing the microorganisms being the objects of the invention and additives were produced in the abovementioned way as follows:

Food product I (or FP-I): Milk protein containing mixture + *L. plantarum* MCC1 + *L. gasseri* MCC2; additive sea-buckthorn juice (BT);

Food product II (or FP-II): Milk protein containing mixture + *L. plantarum* MCC1; additive raspberry-blueberry concentrated juice drink (RB).

Food product III (or FP-III): Milk protein containing mixture + *L. plantarum* MCC1 + *L. gasseri* MCC2; additive blackcurrant-raspberry concentrated juice drink (BR).

Food product IV (or FP-IV): Milk protein containing mixture + *L. plantarum* MCC1; additive blueberry concentrate (B).

Food product V (or FP-V): Milk protein containing mixture + *L. plantarum* MCC1; additive cowberry concentrated juice drink (C).

Dietary supplement I (or DS/FP-I): Milk protein containing mixture + *L. plantarum* MCC1 + *L. gasseri* MCC2; additive sea-buckthorn juice (BT);

Dietary supplement II (or DS/FP-II): Milk protein containing mixture + *L. plantarum* MCC1; additive raspberry-blueberry concentrated juice drink (RB).

Dietary supplement III (or DS/FP-III): Milk protein
5 containing mixture + *L. gasseri* MCC2; additive.

Table 10.

Example of typical organoleptical test of food product.

Degustators (n = 15). The next parameters were asked to be
10 evaluated: consistency, colour, smell, taste and acidity on the basis of the following score: 1 - repulsive, 2 - not too tolerable, 3 - tolerable, 4 - good, 5 - excellent/inviting.

	FP-V	FP-IV	FP-III	FP-I
Consistency	2.3±1.2	3.6±1.0	3.3±1.3	4±1.1
Colour	2.6±0.9	3.8±0.8	3±1	4.1±1.2
Taste	2.8±0.9	3.2±1.0	3.8±0.8	3.5±1.0
Smell	2.6±0.9	2.8±1.0	3.4±0.8	3.4±1.2
Acidity	2.8±0.9	3.1±0.8	3.7±0.6	3.2±1.2
Total	2.6±1.0	3.3±0.9	3.4±0.9	3.7±1.2

15 Example 2. Additives for production of milk allergy responses and urinary tract irritation symptoms and associated oxidative stress and inflammation reducing food product and dietary supplement containing microorganisms *L. plantarum* MCC1 or *L. gasseri* MCC2 or their combination.

20

As product additives sea-buckthorn juice (BT) (made from sea-buckthorn berries, without preservatives, cold-pressed and pasteurized), raspberry-blueberry concentrated juice drink (composition: concentrated raspberry-blueberry juice (RB),
25 sugar, acidity regulator citric acid, preservative potassium sorbate, was diluted 10 times), cowberry concentrated juice drink (C) and blueberry concentrate (B) (see Table 10) were used. Antioxidancy (TAA and DPPH tests) and contents of bioelements with atom absorption spectrophotometric (AAS)

method Spektra AAFS and 220Z were determined for the additives. (Varian, Australia), Cu, Zn, Fe, Mn contents were determined with AAS flame method, K content with emission method in air-acetylen mixture. Organoleptical tests of the end-product were carried out according to valid requirements with special questionnaires. The results are presented in Table 10.

Example 3. Clinical trial no. 1.

Safety and beneficial effects of the products, biochemical-clinical and organoleptic tests.

With the use of multipotent strains obtained by versatile bioselection and additives and application of suitable technological scheme (see example 1) food products FP-I (MCC1+ *L. gasser*i MCC2, BT) and FP-II (MCC1, RB) were obtained. Subsequently clinical trials (in accordance to Declaration of Helsinki of the World Medical Association, approved by the Tartu University Ethics Review Committee) were performed to clarify the safety and healthy effects of food products FP-I and FP-II.

The main scheme of the clinical trial was the following: 25 volunteers (11 women, 14 men), 40-65 years old, were randomly assigned to two groups on the basis of the next study inclusion criteria: persons without clinical problems, chronic diseases, special diets, use of vitamins, mineral preparations, the participants do not have to change their physical activity, usual alcohol use, smoking habits or eating habits. Before and after two-week everyday use of 200 g of FP-I (n=10) or FP-II (n=15) the next biochemical-clinical parameters were determined: inflammation markers (hsCRP, MPO, IL-6, IL-10), lipid and lipoprotein markers (TG, cholesterol, LDL-cholesterol, HDL-cholesterol, sdLDL), immunoglobulins (IgM, IgA, IgG, IgE), oxidative stress (OxS) markers (isoprostanes, oxLDL, LDL-b2GPI), blood glucose, blood pressure and urine creatinine.

Results

1) Safety

Scientific analysis of everyday questionnaire showed that no subjects complained of discomfort or intestinal problems during the two-week use and no negative symptoms occurred. The use of FP-I or FP-II did not have effects that would drive their biochemical and clinical parameters outside the limits endemic reference values. They did not increase the amount of IgE antibodies as one of the important allergy markers compared to the pre-trial period, did not cause inflammatory reaction (hsCRP oxLDL, MPO, IL-6) or change the values of blood lipids, lipoproteins (TG, LDL, HDL) and glucose nor increase the level of OxS in the body (isoprostanes, oxLDL, MPO, sdLDL, LDL b2GPI) (Table 10) and did not damage the renal functions (urine creatinine). Thus: FP-I and FP-II obtained on the basis of the ideology of the invention don't cause inflammatory reactions, general allergic sensibilization, OxS and therefore do not damage the organism in healthy persons.

20

2) Health promoting effects

The analysis of the results of the clinical trial showed that the use of FP-I or FP-II induced some statistically significant and somewhat different positive effects (Table 10). FP-I exerted the effect reducing the systemic OxS load of the whole body (decrease of urine isoprostanes and blood MPO level) and anti-inflammatory effect (decrease of blood MPO). FP-II reduced the systemic OxS load of the whole body (decrease of urine isoprostane level) and exerted anti-inflammatory effect (decrease of hc-CRP and increase of IL-10). At the same time the use of both food products did not change the numerical values of immunoglobulins IgM, IgA and IgM.

30

Thus: the first clinical trial showed that the ideology of the invention (multivariant purposeful bioselection of the strains, bioselection of additives and technological solution) resulted in food products that are safe and are more hypoallergenic having several inflammation and oxidative stress-related effects (cause less allergic responses compared to cow's milk) and have a positive effect on biochemical and clinical parameters also in persons with prostate complaints (including urination disorders).

10

Table 11. Biochemical and clinical parameters (clinical trial no. 1; M± SD)

	GROUP I (FP-II)*		GROUP II (FP-I)**	
	Before	After	Before	After
Blood pressure, mmHg	123/74±7/5	125/70±9/5	118/71±12/5	
Total cholesterol mmol/L	5.8±1.3	5.9± 1.4	6.3±1.1	6.6±0.9
HDL-cholesterol mmol/L	1.6±0.4	1.6±0.5	1.7±0.5	1.7± 0.5
LDL-cholesterol	4.0±1.1	4.1±1.2	4.6± 1.1	4.9±1.1
triglycerides mmol/L	1.4±1.3	1.4± 1.3	1.4± 0.5	1.5±0.7
Glucose mmol/L	5.2±0.4	5.2±0.7	5.2± 0.5	5.3±0.5
hs-CRP mg/L	0.72±0.45	0.90±0.59	3.1±1.8***	1.9±1.5*** p = 0.049
IgM g/L	1.26±0.48	1.29±0.49	1.09±0.37	1.10±0.35
IgA g/L	2.42±0.85	2.38±0.87	3.06±1.46	3.08±1.48
IgE g/L	52±56	53±59	88±95	77±81
MPO pmol/L	67±22	51± 13 p=0.018	53±14	61±16
IL-6 pg/mL	2.6±1.2	2.6±0.9	3.9±2.1	2.9±0.9
IL-10 pg/mL	6.8±2.9	6.1±2.6	6.4± 2.5***	7.8±2.9*** p= 0.04
oxLDL U/L	77±33	75±33	74±24***	69± 21*** p = 0.039
LDL□2GPI	1.4±1.6	1.1±1.2	1.2±1.2***	1.0± 0.7***
sdLDL	38.6±15.7	37.1±16.3	39.6±16.1	38.8±16.0
8-isoprostanes ng/mL	36.7±6.9	31.3±12.5 p =0.034	49.6± 8.9***	37.7± 6.7*** p=0.021

*n = 10, ** n= 9, ***n = 15; blood pressure lowered in four persons in group II

Example 4. Clinical trial no. 2. Cow's milk allergy studies in children

Reactions caused by cow's milk protein occur in 5-15% of infants (Host A. (2002). Frequency of cow's milk allergy in childhood. *Ann Allergy Immunol* 89(Suppl 1):33-7), but cow's milk protein allergy (CMPA) occurs in 2-7.5% (Hill DJ, Firer MA, Shelton MJ et al. (1986) Manifestation of milk allergy in infancy: clinical and immunologic findings. *J Pediatr* 109: 270-6). CMPA may be IgE- or non-IgE-mediated. As a rule the early reactions are urticaria, angioedema, vomiting or exacerbation of atopic dermatitis. Exacerbation of atopic dermatitis or gastrointestinal complaints occur as late reactions (Vandenplas Y, Brueton M, Dupont C, Hill D, Isolauri E, Koletzko S, Oranje AP, Staiano A. (2007) Guidelines for the diagnosis and management of cow's milk protein allergy in infants. *Arch Dis Child*. 92:902-908). None of the existing diagnostic tests does prove or deny if the child has CMPA or not (Vanto T, Juntunen-Backman K, Kalimo K et al. (1999) The patch test, skin prick test, and serum milk-specific IgE as diagnostic tools in cows' milk allergy in infants. *Allergy* 54:837-42). Therefore the diagnostic gold standard of CMPA is elimination diet with subsequent provocation (Vandenplas Y, Brueton M, Dupont C, Hill D, Isolauri E, Koletzko S, Oranje AP, Staiano A. (2007) Guidelines for the diagnosis and management of cow's milk protein allergy in infants. *Arch Dis Child*. 92:902-908). As a protocol for the provocation test an internationally acknowledged scheme for such investigations described by Isolauri et al. was used (Isolauri E, Turjanmaa K. (1996) Combined skin prick and patch testing enhances identification of food allergy in infants with atopic dermatitis. *J Allergy Clin Immunol*;97:9-15; Sütas Y, Kekki O-M, Isolauri E. (2000) Late onset reactions to oral food challenge are linked to low

serum interleukin-10 concentrations in patients with atopic dermatitis and food allergy. Clin Exp Allergy 30:1121-8).

Methodology

Open provocation with food products FP-II and FP-I was performed in the Centre of Allergic Diseases of the Children's Clinic of the Tartu University Clinics Foundation (in accordance to Declaration of Helsinki of the World Medical Association, approved by the Tartu University Ethics Review Committee). The testing was performed 20 times (part of subjects participated in the trials of both food products) in children aged 9 months to 4 years 8 months (mean age 25.1 months). Pre-trial background was following: all children were diagnosed with cow's milk allergy and/or intolerance by a doctor at the Centre of Allergic Diseases of the Children's Clinic of the Tartu University Clinic Foundation (IgE test to milk, NTT to milk) relying on history and ancillary investigations. All children in the study group were on milk-free elimination diet. After the use of milk 9 children developed skin rash, usually exacerbation of atopic dermatitis, one child developed acute urticaria, one child diarrhoea. One child developed breathing difficulties and skin rash only after eating processed milk products. 3 children had been concurrently diagnosed with asthma, one of them also allergic rhinitis and one pollen allergy (pollinosis). One child had asthma and pollen allergy as concurrent diseases. None of the children received antihistamine or corticosteroid treatment. In three children the concurrent asthma was in remission, therefore they did not receive permanent asthma treatment when the provocation test was performed.

Open oral provocation was performed on the first trial day in the Centre of Allergic Diseases of the Children's Clinic of the TUCF. The child was administered the investigated product (FP-II or FP-I) in increasing amounts (a drop on lower lip, 2

mL, 10 mL, 50 mL, 100 mL). After each amount the child's condition and the development of allergic reaction was evaluated during 20 minutes. If rash, cough, breathing difficulties or abdominal pain/diarrhoea developed, the administration of the product was immediately discontinued. If the child did not have reaction on the first trial day after up to 100 ml of product, the child then received the investigated product at home 100 mL/day on the second, third and fourth day. The parent was asked to evaluate the development of late-onset reactions. The test was considered positive if the child had: a) early reaction - on the first provocation day after administration of small amount of product pruritus, rash, hives or vomiting developed; b) late-onset reaction - dermatitis exacerbation, vomiting, abdominal pain, and diarrhoea developed > 24 hours after initiating the investigated product provocation. The open provocation test was negative if during the provocation (four days) and during subsequent week no reaction developed. Thus the subject tolerated the product.

Summary of results

The solving of the problem of food allergenicity is very complicated and it is not possible to create products that are absolutely allergic reaction-free but with very high bio-quality. The problem becomes more difficult as currently a child can have an allergic reaction to several food products (but also to other common factors) at the same time. This was also observed in our trials (see Table 5), which makes it difficult to exclude that at the given moment the positive response rash did not manifest due to initiation by other causes. Therefore the focus of the investigation was directed so that less allergic responses would result, i.e. the obtained food products would be more hypoallergenic than cow's milk. Each percent of children who are allergy-free or

less allergic to given food products would increase the number of children who would get sufficient biologically high-quality food for their development during their most important period of biological development. The provocation test showed that both food product FP-II as well as FP-I were more hypoallergenic than cow's milk (Table 5). At least 3-4 children from 12 did not develop any allergic reaction after FP-I use on the basis of the provocation test. Three subjects out of eight did not develop any allergic reaction after FP-II use. Out of five subjects in whom FP-I caused some reaction, three tolerated FP-II (60%). FP-II was thus in terms of cases even more hypoallergenic than FP-I. Most of the children also preferred FP-II due to taste properties.

Table 12. Investigation of allergic response to cow's milk (hypoallergenicity trial)

B/ G	Diagnosis Clinical manifestation :	NTT to milk	NTT to FP-I	Provocation test to FP-I	Other NTT	Age
G1	CMA, AD skin rash	neg	neg	neg	egg yolk egg white	23m
B2	CMA, BA, AD (30m) skin rash, breathing difficulty	pos	neg	neg	-	36m
B3	CMA, AD, BA (24m), pollinosis, skin rash	pos	pos	positive, (from 50 mL) rash on cheeks	egg, nuts, cat, pollen, house dust mite etc	46m
B4	CMA, AD skin rash	pos	pos	pos (from 10 mL, rash)	egg yolk egg white	16m
B5	CMA, AD, BA (27m) diarrhoea	neg	neg	pos (on the second day, rash)	egg white	28m
G6	CMA, AD skin rash	neg	neg	pos (on the third day, rash)	egg yolk egg white	9m
G7	AD skin rash	neg	neg	pos (on the second day, rash, abdominal pain)	Neg	18m
B8	CMA, AD skin rash	pos	neg	neg	Neg	40m
B9	CMA, AD skin rash	pos	pos	pos (from 10 mL, rash)	egg yolk, egg white	14m

B1 0	CMA, AD skin rash	pos	neg	neg (on the first day, on the second didn't want to drink)	Neg	31m
B1 1	Urticaria hives	pos	neg	neg/pos (days 2- 4 diarrhoea)	egg yolk, egg white	14m
B1 2	CMA, AD pruritus	pos	pos	pos (on the second day pruritus)	egg yolk egg white	33m
B/ G	Diagnosis Clinical manifestation :	NTT to milk	NTT to FP-II	Provocation test to FP-II	Other NTT	Age
B9	CMA, AD skin rash	neg	pos	neg	egg yolk, egg white	23m
B3	CMA, AD, BA (24m), pollinosis, skin rash	pos	pos	pos (on the second day pruritus, days 3 and 4 overexcitability)	egg, nuts, cat, pollen, house dust mite etc	56m
B1 1	Urticaria hives	pos	neg	neg (on the first day diarr- hoea after 100 mL, days 2-4 without problems)	egg yolk, egg white	16m
B1 2	CMA, AD pruritus	pos	pos	pos (on the first day itching rash)	egg yolk egg white	34m
G7	AD skin rash	neg	neg	neg	neg	27m
B1 3	CMA, AD skin rash	pos	pos	neg	egg yolk egg white	12m
B1 4	CMA, AD skin rash	pos	neg	pos (on the second day, rash)	egg yolk, egg white	18m
B1 5	AD skin rash	pos	neg	neg	egg yolk, egg white	17m

CMA - cow's milk allergy, AD - atopic dermatitis, BA - bronchial asthma

Example 5. Clinical trial no. 3. The effect of food products FP-I and FP-II on the lower urinary tract irritation symptoms accompanying benign prostatic hyperplasia and on oxidative stress and inflammation indicators.

Abovementioned clinical trials showed that the ideology of the invention (multivariant purposeful bioselection of the strains, bioselection of additives and technological solution) resulted in food products FP-I and FP-II that are safe for use and are more hypoallergenic having several inflammation and oxidative stress-related effects and have a

positive effect on urinary tract irritation symptoms and OxS and inflammation indicators.

Methodology

In the trial participated 55 men (administered FP-I) and 5
5 men (administered FP-II) who had mild or moderate IPSS
(*International Prostate Symptom Score*) < 8. The IPSS
questionnaire contains 3 questions assessing the lower
urinary tract irritation symptoms and 4 questions assessing
urinary obstruction (obstruction symptoms). Also required
10 prior tests (PSA, prostate secretion) were carried out for
accurate selection of the study group. It means that with the
help of prostate secretion investigations and NIH-CPSI
questionnaire active prostatitis (prostate inflammation) was
ruled out and by using PSA test and digital palpation the
15 clinical possibility of prostate cancer was ruled out. Thus
we included in the study group men with mild or medium
urinary disorders in whom other causes of complaints
(prostatitis, prostate cancer) were ruled out. The patients
(with mild and moderate urination disorder, without
20 inflammation reaction) used food product FP-I or FP-II 200 g
daily for two weeks.

Results

25 FP-I. Two test series were carried out. In the first 23
patients participated and in the second (where placebo
control was also performed) 32 patients (Table 13). On both
occasions statistically significant reduction of inflammation
marker HsCRP and OxS marker isoprostanes was observed. The
30 second test series also pointed out statistically positive
changes of anti-inflammatory IL-10 and inflammation/OxS
marker oxLDL. In the first test series (n=23) also the trend
of positive changes of this marker was observed. In the

second test series also the level of glycated hemoglobin was determined, the decrease of which was statistical.

When summarizing the results of both test series (Table 13),
5 then hsCRP, isoprostanes and oxLDL decreased statistically
and there was a positive trend of IgE decrease and IL-10
increased statistically. There was a correlation between
hsCRP and oxLDL after the use of FP-I. In control group the
statistically significant changes were not seen for any of
10 the mentioned parameters.

FP-II. One test series was performed and the next parameters
determined: IgE kU/L, oxLDL, IL-10, 8-isoprostanes, hsCRP and
MPO. It was found that in the group of RP-II users
15 statistically significant reduction of inflammation and Oxs
marker MPO (before 77 ± 13 and after 53 ± 9 ng/mL, $p=0.0056$) and
the reduction of the frequency of urinary disorders in more
than 40% of users occurred.

20 Summary of results

The administration of FP-I as well as FP-II reduced the level
of Oxs and inflammation in subjects who had moderate lower
urinary tract complaints (IPSS score <8) and had leading
irritation symptoms. In case of FP-I in 66% of study group
25 men (55), who had moderate lower urinary tract complaints
(IPSS score <8) and leading irritation symptoms, the
reduction of complaints for at least 20% occurred and in 30%
of study group reduction of complaints for an average of 40%
occurred. The clinical and biochemical parameters of the
30 patients (hs-CRP, IL-10, oxLD, 8-isoprostanes) improved
statistically possibly. No changes occurred in the control
group.

Table 13. Test series of FP-I (M $\pm$ SD)

	First test series (n=23)	Second test series (n=32)	Total (n=55)
	Before After	Before After	Before After
hsCRP mg/L	2.09 $\pm$ 1.06 1.04 $\pm$ 0.90 p=0.016	1.92 $\pm$ 1.24 1.57 $\pm$ 1.10 p=0.060	1.99 $\pm$ 1.17 1.50 $\pm$ 0.96 p=0.002
B-Hb A1c, % total Hb 4.5-5.7	Not determined	5.87 $\pm$ 0.35 5.79 $\pm$ 0.36 p=0.0013	
IgE, kU/L	98.3 $\pm$ 67.5 86.6 $\pm$ 56.1	129.8 $\pm$ 133.4 130.6 $\pm$ 129.3	116.2 $\pm$ 107.0 112.5 $\pm$ 99.8
oxLDL, U/L	64 $\pm$ 12 63 $\pm$ 13	72 $\pm$ 16 70 $\pm$ 16 p=0.047	68 $\pm$ 15 65 $\pm$ 15 p=0.045
IL-10	5.0 $\pm$ 1.7 5.6 $\pm$ 2.5	8.6 $\pm$ 2.2 10.0 $\pm$ 2.8 p=0.014	7.1 $\pm$ 2.6 8.2 $\pm$ 3.3 p=0.012
8-iso- prosta- nes to ng/mmol creatini ne	50.8 $\pm$ 10.6 43.3 $\pm$ 0.9 p=0.03	47.8 $\pm$ 15.9 37.2 $\pm$ 10.1 p= 0.0007	48.2 $\pm$ 14.2 39.8 $\pm$ 10.2 p=0.005

- 5 Example 6. The effect of dietary supplement (DS/FP-I) on the lower urinary tract irritation symptoms accompanying benign prostatic hyperplasia and on oxidative stress and inflammation indicators.
- 10 10 men who had mild or moderate urination disorder - IPSS (International Prostate Symptom Score) were included in the trial, whereas on the basis of prior prostate secretion investigations and NIH-CPSI questionnaire active prostatitis was excluded and by using PSA test and digital palpation the
- 15 possibility of clinically significant prostate cancer was excluded. The dietary supplement was used 45 g a day for 14 days. In all 10 patients with mild (IPSS score 2-7) and

moderate (IPSS score 8-19) urinary disorders, 3/7 respectively, the urinary disorders improved after the use of dietary supplement ($p=0.003$) according to the "International Prostate Symptom Score" by 50-66%, whereas it was found that
5 significant positive change occurred in reduction of the need to urinate during the night (average 87.5%).

Overall summary of all clinical trials

Food products and dietary supplements containing *L. plantarum*
10 MCC1 and *L. gasseri* MCC2 are hypoallergenic, reduce the level of Oxs and have statistically valid effect on prostate complaints in those who have moderate lower urinary tract complaints and leading irritation symptoms.

CLAIMS

1. Isolated microorganism strain *L. plantarum* MCC1 DSM 23881.
2. Isolated microorganism strain *L. gasserii* MCC2 DSM 23882.
3. Microorganism according to claims 1 or 2 in killed form.
4. A composition containing one or more microorganisms according to claims 1-3.
5. Composition according to claim 4, where in case of occurrence of one microorganism the content of the microorganism is in the range of 0.8-1.2 volume percent, whereas the germ count in case of *L. plantarum* MCC1 is 0.25×10^8 - 4×10^8 CFU/mL and in case of *L. gasserii* MCC2 0.5×10^7 - 1×10^8 CFU/mL and in case of the occurrence of both microorganisms the content of both is in the range of 0.4-0.6 volume percent, whereas the germ count of *L. plantarum* MCC1 is in the range 0.2×10^8 - 2.4×10^8 CFU/mL and the germ count of *L. gasserii* MCC2 in the range 0.4×10^7 - 0.6×10^8 CFU/mL.
6. The use of microorganism according to claim 1 and/or 2 together or separately as antioxidant proteolytic ingredient for the production of food product and/or dietary supplement.
7. The use of microorganism according to claims 1-3 together or separately for the production of hypoallergenic, milk allergy reducing food product and/or dietary supplement.
8. The use of microorganism according to claims 1-3 together or separately for the production of food product and/or dietary supplement reducing lower urinary tract irritation symptoms accompanying benign prostatic hyperplasia and oxidative stress and inflammation associated with these.

9. Method for the production of food product and dietary supplement reducing milk allergy and urinary tract complaints and oxidative stress and inflammation associated with these that includes the next stages:

5 A) the initial mixture containing milk protein is warmed ($37\pm 2^{\circ}\text{C}$);

B) 1 volume percent of microorganism according to claim 1 with germ count 0.4×10^8 - 4.8×10^8 CFU/mL and/or microorganism according to claim 2 with germ count
10 0.8×10^7 - 1.2×10^8 CFU/mL or microorganism according to claim 1 and claim 2 together respectively 0.25×10^8 - 2×10^8 CFU/mL and 0.5×10^7 - 0.5×10^8 CFU/mL is added;

C) the obtained mixture is fermented at temperature $37\pm 2^{\circ}\text{C}$ for 18-26 hours;

15 D) the mixture is pasteurized at temperature $82\pm 2^{\circ}\text{C}$ for 30-35 minutes and cooled down to temperature 20 - 30°C ;

E) the mixture is flavoured with an additive and the obtained food product is cooled to 2 - 6°C ;

F) for powder dietary supplement production, water is
20 removed from the mixture until powder consistency is achieved and for liquid dietary supplement production, water is removed partially from the mixture until desired consistency is achieved.

10. Method according to claim 9, where the additive is fruit
25 or berry juice, concentrate, syrup or juice drink or fruit or berry jam, preferably sea-buckthorn, blueberry or raspberry juice.

11. Method according to claim 9, where the milk protein
30 source is whey, concentrated whey, whey powder, milk, milk powder or milk protein concentrate.

1/6




FIG.1

2/6




FIG. 2

3/6




FIG.3

4/6




FIG. 4

5/6




FIG. 5

6/6




FIG. 6

INTERNATIONAL SEARCH REPORT

International application No

PCT/EE2012/000001

A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N1/20

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2006/041930 A2 (NUTRACEA INC [US]; CHERUVANKY RUKMINI [US]; CHERUKURI REDDY SASTRY [US] 20 April 2006 (2006-04-20) paragraph [0026] -----	1-11
A	WO 2005/058335 A1 (NUTRICIA NV [NL]; M RABET LAURA [NL]; SPEELMANS GELSKE [NL]; VRIESEMA) 30 June 2005 (2005-06-30) page 14, line 33 -----	1-11



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

16 May 2012

Date of mailing of the international search report

29/05/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Stoyanov, Borislav

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EE2012/000001

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-11(partially)

Isolated microorganism strain L. plantarum MCC1 DSM 23881
and the uses thereof.

2. claims: 1-11(partially)

Isolated microorganism strain L. gasseri MCC1 DSM 23882 and
the uses thereof.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EE2012/000001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2006041930	A2	20-04-2006	
		AU 2005294369 A1	20-04-2006
		CA 2583668 A1	20-04-2006
		EP 1796473 A2	20-06-2007
		JP 2008515809 A	15-05-2008
		WO 2006041930 A2	20-04-2006

WO 2005058335	A1	30-06-2005	
		AU 2004298384 A1	30-06-2005
		CA 2550106 A1	30-06-2005
		CN 1913906 A	14-02-2007
		EP 1696941 A1	06-09-2006
		JP 4850715 B2	11-01-2012
		JP 2007514738 A	07-06-2007
		JP 2012025760 A	09-02-2012
		RU 2010129069 A	20-01-2012
		US 2009257993 A1	15-10-2009
		WO 2005058335 A1	30-06-2005
