



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/08/28
 (87) Date publication PCT/PCT Publication Date: 2021/03/04
 (85) Entrée phase nationale/National Entry: 2022/02/24
 (86) N° demande PCT/PCT Application No.: JP 2020/032514
 (87) N° publication PCT/PCT Publication No.: 2021/039943
 (30) Priorités/Priorities: 2019/08/29 (JP2019-156537);
 2020/01/29 (JP2020-012333)

(51) Cl.Int./Int.Cl. *C12N 5/0775* (2010.01),
C07K 14/78 (2006.01)
 (71) Demandeur/Applicant:
 AJINOMOTO CO., INC., JP
 (72) Inventeurs/Inventors:
 HARATA, IKUE, JP;
 HARADA, ERI, JP;
 TAKAHASHI, KAZUMA, JP;
 HONKAWA, FUMIE, JP
 (74) Agent: NORTON ROSE FULBRIGHT CANADA
 LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : PROCÉDE DE PRODUCTION DE CELLULES SOUCHES MESENCHYMATEUSES A PARTIR D'UN
 ECHANTILLON DE CELLULES DERIVEES D'UN CORPS VIVANT CONTENANT DES CELLULES SOUCHES
 MESENCHYMATEUSES
 (54) Title: METHOD FOR PRODUCING MESENCHYMAL STEM CELLS FROM LIVING BODY-DERIVED CELL SAMPLE
 CONTAINING MESENCHYMAL STEM CELLS

(57) **Abrégé/Abstract:**

The present invention provides a method for producing mesenchymal stem cells from a living body-derived cell sample containing mesenchymal stem cells, the method comprising: (1) a step for culturing a living body-derived cell sample containing mesenchymal stem cells in a serum-free medium in the presence of vitronectin or a partial peptide of the vitronectin capable of bonding mesenchymal stem cells; and (2) a step for collecting a cell aggregate of the mesenchymal stem cells.

ABSTRACT

A method for producing a mesenchymal stem cell from a biological cell sample containing mesenchymal stem cells is provided by the present invention, which includes the following
5 steps:

- (1) a step of culturing the biological cell sample containing mesenchymal stem cells in a serum-free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells,
- 10 (2) a step of collecting a cell aggregate of the mesenchymal stem cells.

[DESCRIPTION]

[Title of Invention]

METHOD FOR PRODUCING MESENCHYMAL STEM CELLS FROM LIVING BODY-
DERIVED CELL SAMPLE CONTAINING MESENCHYMAL STEM CELLS

5 [Technical Field]

[0001]

The present invention relates to a method for producing a
mesenchymal stem cell from a biological cell sample containing
mesenchymal stem cells, including a step of culturing the
10 biological cell sample containing mesenchymal stem cells in a
serum-free medium or a xeno-free medium in the presence of
vitronectin or a partial peptide thereof capable of adhering
mesenchymal stem cells.

[Background Art]

15 [0002]

In recent years, the development of pharmaceutical products
using biological cells or tissues and the research on regenerative
medicine have progressed and are attracting attention. Of these,
research using embryonic stem cells and induced pluripotent stem
20 cells provided with pluripotency as organ regeneration technique or
drug discovery screening tool is accelerating. However, the
production of embryonic stem cells requires the destruction of
embryos developed from fertilized eggs, which poses an ethical
problem. Induced pluripotent stem cells are obtained by
25 reprogramming somatic cells, and the above-mentioned problems do not
occur. However, there is a concern that cancer cells may be
generated from induced pluripotent stem cells due to the use of c-
myc as a reprogramming factor, random gene transfer into chromosomes
by a retrovirus vector, undifferentiated cells remaining after
30 differentiation, and the like. On the other hand, mesenchymal stem
cells have pluripotency permitting differentiation into not only

multiple cells of mesenchymal lineage (osteoblast, adipocyte, chondrocyte) but also cells of non-mesenchymal lineage (neural progenitor cell, hepatocyte), and are free of problems caused by embryonic stem cells or induced pluripotent stem cells. Thus, they
5 are expected to be used as a cell source for regenerative medicine and cell therapy.

[0003]

Mesenchymal stem cells can be produced not only from tissues of adults such as bone marrow, fat, synovial membrane, alveolar bone,
10 periodontal ligament, and the like but also from various tissues such as placenta, umbilical cord blood, and umbilical cord, and can be cultured and amplified in vitro. As a conventional method for obtaining mesenchymal stem cells, since bone marrow mononuclear cells produced from bone marrow contain a small amount of
15 mesenchymal stem cells, the mesenchymal stem cells were produced by culturing bone marrow mononuclear cells produced from bone marrow in a medium containing bovine fetal serum (FBS), and utilizing the adhesiveness of the mesenchymal stem cells to a culture container. However, when mesenchymal stem cells are used as a cell source for
20 regenerative medicine, contamination of the mesenchymal stem cells with xenogeneic components was inconvenient. Therefore, a method for culturing mesenchymal stem cells using a serum-free medium has been devised (Patent Literature 1). However, in order to achieve proliferation of mesenchymal stem cells, it is necessary to produce
25 mesenchymal stem cells from bone marrow mononuclear cells containing mesenchymal stem cells, and a method for efficiently producing mesenchymal stem cells from bone marrow mononuclear cells containing mesenchymal stem cells by using a serum-free medium has not yet been developed.

30 [Citation List]

[Patent Literature]

[0004]

[PTL 1]

WO 2011/111787

[Summary of Invention]

5 [Technical Problem]

[0005]

An object of the present invention is to provide a method for efficiently producing a mesenchymal stem cell from a biological cell sample containing mesenchymal stem cells.

10 [Solution to Problem]

[0006]

The present inventors have conducted intensive studies in an attempt to achieve the above-mentioned object and succeeded in producing cell aggregates formed by mesenchymal stem cells adhered
15 to a culture container via vitronectin, by culturing bone marrow mononuclear cells containing mesenchymal stem cells in a serum-free medium on a culture container coated with vitronectin. The produced aggregate was dissociated to obtain a population of single mesenchymal stem cells. Then, the obtained mesenchymal
20 stem cells were cultured again in the serum-free medium in the presence of vitronectin, and it was confirmed that the number of cells was remarkably higher than that of mesenchymal stem cells cultured under the same conditions except that fibronectin was used instead of the first vitronectin. It was also confirmed that
25 the TGF β receptor inhibitor further increases the production efficiency of mesenchymal stem cells. Furthermore, it was found that mesenchymal stem cells can be produced from adipocytes containing mesenchymal stem cells by using vitronectin and a TGF β receptor inhibitor. From the above findings, the present
30 invention has been completed.

[0007]

Accordingly, the present invention provides the following.

[1] A method for producing a mesenchymal stem cell from a biological cell sample comprising mesenchymal stem cells, comprising the following steps:

5 (1) a step of culturing the biological cell sample comprising mesenchymal stem cells in a serum-free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells,

(2) a step of collecting a cell aggregate of the mesenchymal
10 stem cells.

[2] The method of [1], wherein the culture in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a culture container on which vitronectin or a partial peptide thereof
15 capable of adhering mesenchymal stem cells has been immobilized.

[3] The method of [1] or [2], further comprising the following steps:

(3) a step of dissociating the collected cell aggregate,

(4) a step of culturing the dissociated mesenchymal stem
20 cells in a serum-free medium in the presence of an extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells,

(5) a step of collecting the mesenchymal stem cells proliferated on the culture container via the extracellular matrix
25 protein or a partial peptide thereof capable of adhering mesenchymal stem cells.

[4] The method of [3], wherein the culture in the presence of the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a
30 culture container on which the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells

has been immobilized.

[5] The method of any one of [1] to [4], wherein the partial peptide of vitronectin comprises an RGD domain.

[6] The method of [5], wherein the partial peptide of vitronectin
5 further comprises a somatomedin B domain.

[7] The method of [6], wherein the partial peptide of vitronectin is a polypeptide consisting of amino acid Nos. 1 - 379 of the amino acid sequence shown in SEQ ID NO: 1.

[8] The method of any one of [1] to [7], wherein the serum-free
10 medium in step (1) comprises a TGF- β receptor inhibitor.

[9] A method for producing a mesenchymal stem cell from a biological cell sample comprising mesenchymal stem cells, comprising the following steps:

(1) a step of culturing the biological cell sample
15 comprising mesenchymal stem cells in a xeno -free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells,

(2) a step of collecting a cell aggregate of the mesenchymal stem cells.

[10] The method of [9], wherein the culture in the presence of
20 vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a culture container on which vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells has been immobilized.

[11] The method of [9] or [10], further comprising the following
25 steps:

(3) a step of dissociating the collected cell aggregate,

(4) a step of culturing the dissociated mesenchymal stem cells in a xeno-free medium in the presence of an extracellular
30 matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells,

(5) a step of collecting the mesenchymal stem cells proliferated on the culture container via the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells.

5 [12] The method of [11], wherein the culture in the presence of the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a culture container on which the extracellular matrix protein or a partial peptide thereof capable of adhering
10 mesenchymal stem cells has been immobilized.

[13] The method of any one of [9] to [12], wherein the partial peptide of vitronectin comprises an RGD domain.

[14] The method of [13], wherein the partial peptide of vitronectin further comprises a somatomedin B domain.

15 [15] The method of [14], wherein the partial peptide of vitronectin is a polypeptide consisting of amino acid Nos. 1 - 379 of the amino acid sequence shown in SEQ ID NO: 1.

[16] The method of any one of [9] to [15], wherein the xeno-free medium in step (1) comprises a TGF- β receptor inhibitor.

20 [17] The method of any one of [9] to [16], wherein the xeno-free medium comprises an allogeneic serum.

[18] The method of [17], wherein the allogeneic serum is an autologous serum.

[19] The method of any one of [1] to [18], wherein the biological
25 cell sample comprising mesenchymal stem cells is a bone marrow-derived cell.

[20] The method of [19], wherein the number of the cells derived from bone marrow and to be cultured is $0.5 \times 10^5 - 25 \times 10^5$ cells/cm².

[21] The method of [19] or [20], wherein the cells derived from
30 bone marrow are cultured for a period of 4 days to 14 days.

[22] The method of any one of [1] to [18], wherein the biological

cell sample comprising mesenchymal stem cells is a cell derived from adipose tissue.

[23] The method of [22], wherein the number of the cells derived from adipose tissue and to be cultured is $1 \times 10^3 - 1 \times 10^6$ cells/cm².

5 [24] The method of [22] or [23], wherein the cells derived from adipose tissue are cultured for a period of 1 day to 14 days.

[Advantageous Effects of Invention]

[0008]

Mesenchymal stem cells can be efficiently produced from a
10 biological cell sample containing mesenchymal stem cells by culturing the biological cell sample containing mesenchymal stem cells in a serum-free medium or a xeno-free medium in the presence of vitronectin or a partial peptide thereof. By adopting this method, the obtained mesenchymal stem cells can be directly used
15 as a cell source in regenerative medicine.

[Brief Description of Drawings]

[0009]

[Fig. 1]

Fig. 1 shows photographs of MSC aggregates 5 days after MNC
20 seeding.

[Fig. 2]

Fig. 2 shows the measurement results of the number of cells 13 days after MNC seeding.

[Fig. 3]

25 Fig. 3 shows photographs of cells 13 days after MNC seeding.

[Fig. 4]

Fig. 4 shows the measurement results of the number of MSC 13 days after MNC seeding when different Vitronectin was used.

[Fig. 5]

30 Fig. 5 shows the measurement results of the number of MSC 12 days after MNC seeding when different Vitronectin was used.

[Fig. 6]

Fig. 6 shows the measurement results of the number of MSC 15 days after MNC seeding when a TGF β inhibitor was used.

[Fig. 7]

5 Fig. 7 shows the measurement results of the number of MSC 12 days after MNC seeding when a different TGF β receptor inhibitor was used.

[Fig. 8]

10 Fig. 8 shows the measurement results of the number of cells isolated from mouse adipose tissue and at 5 days after seeding.

[Fig. 9]

Fig. 9 shows photographs of cells isolated from mouse adipose tissue and at 5 days after seeding.

[Description of Embodiments]

15 [0010]

The present invention provides a method for producing a mesenchymal stem cell from a biological cell sample containing mesenchymal stem cells (hereinafter, the production method of the present invention).

20 [0011]

In the present specification, the biological cell sample containing mesenchymal stem cells is a cell sample separated from a biological tissue containing mesenchymal stem cells. As the biological tissue containing mesenchymal stem cells, tissues of
25 bone marrow, fat, synovium, alveolar bone, periodontal membrane, placenta, cord blood, umbilical cord and the like can be mentioned.

[0012]

In the present specification, the cell sample is a cell population contained in a biological tissue. The cell population
30 means two or more cells of the same type or different types. In addition, the cell population also means a clump (mass) of cells

of the same or different types. The cell population may be a primary cell directly separated from a biological tissue, or may be a cell that is passage cultured from the primary cell. As used herein, directly means not going through the step of culturing
5 and/or proliferating in vitro.

[0013]

In the present specification, the mesenchymal stem cell is a somatic stem cell derived from a mesodermal tissue (mesenchyma). The mesenchymal stem cell expresses positive markers on the cell
10 surface and does not express negative markers. By detecting both markers on the cell surface, whether or not the cell is a mesenchymal stem cell can be determined. Positive markers include CD73, CD90, and CD105. Negative markers include CD11b, CD14, CD19, CD34, CD45, CD79a, HLA-Class II(DR). The expression of these
15 markers can be examined by a known immunological method (e.g., flow cytometry using an antibody) or the like.

[0014]

In one embodiment, the production method of the present invention includes the following steps:

20 (1a) a step of culturing the biological cell sample containing mesenchymal stem cells in a serum-free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells (step (1a) of the present invention),
(2a) a step of collecting a cell aggregate of the mesenchymal stem
25 cells (step (2a) of the present invention).

In another embodiment, the production method of the present invention includes the following steps:

(1b) a step of culturing the biological cell sample containing mesenchymal stem cells in a xeno-free medium in the presence of
30 vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells (step (1b) of the present invention),

(2b) a step of collecting a cell aggregate of the mesenchymal stem cells (step (2b) of the present invention).

[0015]

In steps (1a) and (1b) of the present invention, culture is
5 performed in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells (hereinafter to be referred to as "the partial peptide of vitronectin").

Vitronectin may be, for example, a protein isolated and purified from the cells of mammals (e.g., human, mouse, rat,
10 rabbit, sheep, swine, bovine, horse, cat, dog, monkey, chimpanzee and the like), or any tissue or organ in which those cells are present. In addition, it may also be a protein chemically synthesized or biochemically synthesized by a cell-free translation system, or a recombinant protein produced from a
15 transformant into which a nucleic acid having a base sequence encoding vitronectin has been introduced.

[0016]

The amino acid sequence of vitronectin is disclosed in a known database, and for example, NP_000629 (human vitronectin),
20 NP_035837 (mouse vitronectin) and the like are disclosed as NCBI Reference Sequence Nos. Since the mesenchymal stem cells produced by the production method of the present invention preferably do not contain xenogeneic components, the vitronectin is preferably derived from a living body from which the cell sample to be
25 cultured is derived. Therefore, when the cell sample to be cultured is derived from human, the vitronectin used in the production method of the present invention is preferably a protein containing the same or substantially the same amino acid sequence as SEQ ID NO: 1.

30 [0017]

Examples of the amino acid sequence substantially the same

as the amino acid sequence shown in SEQ ID NO: 1 include amino acid sequences having a homology of not less than about 60%, preferably not less than about 70%, further preferably not less than about 80%, particularly preferably not less than about 90%,
5 with the amino acid sequence shown in SEQ ID NO: 1, and the like. As used herein, the "homology" means the proportion (%) of the same amino acid residues and similar amino acid residues relative to the total overlapping amino acid residues, in an optimal alignment (preferably, the algorithm is capable of considering
10 introduction of gap into one of or both of the sequences for optimal alignment), when two amino acid sequences are aligned using a mathematical algorithm known in the technical field.

The homology of the amino acid sequences in the present specification can be calculated using homology calculation
15 algorithm NCBI BLAST (National Center for Biotechnology Information Basic Local Alignment Search Tool) and under the following conditions (expectancy =10; gap allowed; matrix =BLOSUM62; filtering=OFF).

More preferably, an amino acid sequence substantially the
20 same as the amino acid sequence shown in SEQ ID NO: 1 is an amino acid sequence having identity of not less than about 60%, preferably not less than about 70%, further preferably not less than about 80%, particularly preferably not less than about 90%, with the amino acid sequence shown in SEQ ID NO: 1.

25 [0018]

As the protein containing an amino acid sequence substantially the same as the amino acid sequence shown in SEQ ID NO: 1, for example, a protein containing an amino acid sequence substantially the same as the aforementioned amino acid sequence
30 shown in SEQ ID NO: 1 and having activities substantially equivalent to those of a protein containing the amino acid

sequence shown in SEQ ID NO: 1 and the like is preferable.

[0019]

Examples of the substantially equivalent activity include mesenchymal stem cell adhesion activity. Being "substantially
5 equivalent" means that the activities thereof are qualitatively (e.g., physiological or pharmacologically) the same. Therefore, while the mesenchymal stem cell adhesion activity is preferably equivalent (e.g., about 0.5- to about 2-fold), the levels of these activities, and quantitative factors such as molecular weight of
10 protein, and the like may be different.

[0020]

The human vitronectin also encompasses, for example, proteins containing (1) an amino acid sequence resulting from deletion of one or two or more (preferably 1 - about 10) amino
15 acids from the amino acid sequence shown in SEQ ID NO: 1, (2) an amino acid sequence resulting from addition of one or two or more (preferably 1 - about 10) amino acids to the amino acid sequence shown in SEQ ID NO: 1, (3) an amino acid sequence resulting from insertion of one or two or more (preferably 1 - about 10) amino
20 acids in the amino acid sequence shown in SEQ ID NO: 1, (4) an amino acid sequence resulting from substitution of one or two or more (preferably 1 - about 10) amino acids with other amino acids in the amino acid sequence shown in SEQ ID NO: 1, (5) an amino acid sequence which is a combination thereof, and the like.

25 As mentioned above, when the amino acid sequence is inserted, deleted, or substituted, the position of the insertion, deletion, or substitution is not particularly limited as long as the activity of the protein is maintained.

[0021]

30 The partial peptide of vitronectin is a peptide having the above-mentioned partial amino acid sequence of vitronectin, and

may be any as long as it has activity substantially equivalent to that of vitronectin. As used herein, the "substantially equivalent activity" has the same meaning as described above. The "substantially equivalent activity" can be measured in the same
5 manner as in the case of vitronectin. Such partial peptide of vitronectin includes proteins containing the RGD domain. More preferably, the partial peptide of vitronectin is a protein containing a somatomedin B domain and an RGD domain.

Specifically, as the somatomedin B domain, for example, the
10 region shown by amino acid No. 1 - 40 in the amino acid sequence shown in SEQ ID NO: 1 is used. As the RGD domain, for example, the region shown by amino acid No. 41 - 52 in the amino acid sequence shown in SEQ ID NO: 1 is used. The size of the partial peptide of vitronectin is not particularly limited as long as it
15 has mesenchymal stem cell adhesion activity. It preferably contains not less than 100 partial amino acid sequences, more preferably not less than 200 partial amino acid sequences, further preferably not less than 300 partial amino acid sequences. The partial amino acid sequence may be one continuous partial amino
20 acid sequence, or may be a plurality of discontinuous partial amino acid sequences linked together. The most preferable partial peptide of vitronectin satisfying such conditions includes a polypeptide consisting of amino acid Nos. 1 - 379 of the amino acid sequence shown in SEQ ID NO: 1.

25 In addition, as the partial peptide of vitronectin, a commercially available partial peptide of vitronectin may be used. Examples of the commercially available partial peptide of vitronectin include Vitronectin (20-398 aa) (wako), Vitronectin (VTN-N, 62-478 aa) (manufactured by Thermo Fisher Scientific),
30 Vitronectin (Full length, 20-478 aa) (Sigma), synthemax II (manufactured by Corning Incorporated), and the like.

[0022]

In step (1a) or (1b) in the present invention, the biological cell sample containing mesenchymal stem cells is cultured in the presence of vitronectin or a partial peptide of vitronectin by any method that brings mesenchymal stem cells into contact with vitronectin or a partial peptide of vitronectin. For example, a method of culturing in a state where vitronectin or a partial peptide of vitronectin is present in a culture medium or the surface of a culture container can be mentioned. The presence of vitronectin or a partial peptide of vitronectin in a culture medium means an embodiment in which it is directly contained in the culture medium. When vitronectin or a partial peptide of vitronectin is contained in a culture medium, the concentration of the vitronectin or the partial peptide of vitronectin in the culture medium is 0.1 $\mu\text{g/ml}$ - 4.0 $\mu\text{g/ml}$, preferably 1.0 $\mu\text{g/ml}$ - 4.0 $\mu\text{g/ml}$.

[0023]

The presence of vitronectin or a partial peptide of vitronectin on the surface of a culture container means an embodiment in which it is immobilized on the surface of the culture container. When vitronectin or a partial peptide of vitronectin is immobilized on the surface of a culture container, for example, a container or a carrier (microbeads, etc.) used for cell culture is used as the culture container. As the culture container, any material with any shape can be used as long as it does not inhibit maintenance, survival, differentiation, maturation, and self-replication of the cells. Examples of the material of the culture container include glass, synthetic resin including non-woven fabric, natural resin, metal and the like. The shape of the culture container includes polygonal prisms such as triangular prism, cube, cuboid and the like, polygonal pyramids

such as circular cylinder, triangular pyramid, quadrangular pyramid and the like, arbitrary shapes such as circular cone, gourd and the like, globular shape, hemisphere shape, circular shape, ellipse shape, semicircle shape, and the like.

5 Commercially available culture flask, culture dish (culture dish), culture bag, hollow thread type culture device, and the like can also be used. A culture bag having gas permeability is preferred. When a large number of cells is required, a large fermenter may also be used.

10 The culture can be performed in either an open system or a closed system. When the purpose is to administer the obtained mesenchymal stem cells to human or the like, it is preferable to perform the culture in a closed system.

[0024]

15 Vitronectin or a partial peptide of vitronectin can be immobilized in a culture container based on a known means. For example, vitronectin or a partial peptide of vitronectin is dissolved in a solvent (e.g., sterile distilled water, buffer, saline, etc.), added to a culture container, and then allowed to
20 stand overnight at 4°C, whereby vitronectin or partial peptide of vitronectin can be immobilized on a culture container. When vitronectin or a partial peptide of vitronectin is immobilized on a culture container, those of ordinary skill in the art can appropriately determine the concentration of a vitronectin or
25 partial peptide of vitronectin solution. For example, the concentration may be set such that 0.5 µg - 10.0 µg of vitronectin or a partial peptide of vitronectin is generally immobilized per unit area of the culture container.

[0025]

30 The culture container on which vitronectin or a partial peptide of vitronectin is immobilized can be stored at a low

temperature, for example, 4°C until use. Immediately before use, a solution containing vitronectin or a partial peptide of vitronectin is removed by suction from these culture containers, washed once with PBS and then once with a culture medium, and then subjected to
5 culture.

[0026]

In step (1a) of the present invention, the serum-free medium is not particularly limited as long as it does not contain serum. Therefore, as long as serum is not contained, the serum-free medium
10 may contain a component derived from the same species as the species from which the biological cell sample containing mesenchymal stem cells to be cultured is derived (allogeneic component) or a component derived from a different species (xenogeneic component). Examples of the allogeneic component include platelet lysate, serum-
15 derived protein (e.g., albumin, etc.), and the like. Examples of the xenogeneic component include animal-derived lipid, and the like.

[0027]

In step (1b) of the present invention, the xeno-free medium is not particularly limited as long as it does not contain a
20 xenogeneic component. Therefore, as long as a xenogeneic component is not contained, the xeno-free medium may contain an allogeneic serum. As the allogeneic serum, an autologous serum is preferred. As used herein, the autologous serum, and the below-mentioned autologous plasma respectively mean serum and plasma obtained from
25 blood collected from the same donor as the biological cell sample to be cultured.

[0028]

Since plasma contains serum components, a medium containing autologous plasma may be used. Preferably, inactivated autologous
30 plasma is added to the medium. For example, cells are cultured in a culture medium containing not more than 10% (V/V), preferably not

more than 5%(V/V), further preferably not more than 2%(V/V), of inactivated autologous plasma. By using autologous plasma, xenogeneic components are excluded from the production method of the present invention, and a highly safe method for producing
5 mesenchymal stem cells is provided.

[0029]

A serum-free medium or xeno-free medium can be prepared using a medium generally used for culturing animal cells as a basal medium. Examples of the basal medium include, but are not
10 limited to, Dulbecco's Medium (e.g., IMDM), Eagle's medium (e.g., DMEM, EMEM, BME, MEM, α MEM), Ham's medium (e.g., F10 medium, F12 medium), RPMI medium (e.g., RPMI-1640 medium, RPMI-1630 medium), MCDB medium (e.g., MCDB104, 107, 131, 151, 153 medium), Fischer's medium, 199 medium, culture medium for primate ES cell (culture
15 medium for primate ES/iPS cell, Reprorcell), medium for mouse ES cell (TX-WES culture medium, Thromb-X), serum-free medium (mTeSR, Stemcell Technologies), ReproFF, StemSpan (registered trade mark) SFEM, StemSpan (registered trade mark) H3000, StemlineII, ESF-B medium, ESF-C medium, CSTI-7 medium, Neurobasal medium (Life
20 Technologies, Inc.), StemPro-34 medium, StemFit (registered trade mark) (e.g., StemFit AK03N, StemFit AK02N) and the like. Furthermore, these media can be mixed as necessary and used and, for example, DMEM/F12 medium and the like can be mentioned. As a serum-free medium or a xeno-free medium, a known medium or a
25 commercially available medium may be used as it is or after modification. As the commercially available xeno-free medium, for example, DEF-CS500 XF (manufactured by Cellartis) or DXF (manufactured by PromoCell) can be used.

[0030]

30 In step (1a) or (1b) in the present invention, a serum-free medium or a xeno-free medium may contain a TGF- β receptor

inhibitor. TGF- β is a peptide factor that is secreted as a non-active form from almost all normal cells, activated under specific conditions, and exhibits various functions such as suppression of the proliferation of epithelial cells and lymphocytes, and the like. Furthermore, osteogenic factor (BMP) that induces differentiation of osteoblast, activin that promotes secretion of follicle-stimulating hormone and differentiation of erythrocyte, and the like can be mentioned as TGF- β family molecules having a structure similar to TGF- β . In the present specification, TGF- β also includes TGF- β family molecules. Specifically, as TGF- β , TGF- β , activin, Nodal, BMP, GDF (growth/differentiation factor), AMH (anti-Mullerian hormone), and MIS (Mullerian inhibitory substance) can be mentioned, with preference given to TGF- β . The TGF- β receptor is composed of type I and type II receptors present on the cell membrane. Both type I and type II receptors have serine/threonine kinase activity, and the substrate for type II receptor is type I receptor. When the TGF- β family molecule binds to TGF- β receptor, type I receptor is phosphorylated by type II receptor, and the activated type I receptor further phosphorylates intracellular signal transduction molecule Smad, and transmits signals into the cell. Specifically, as the combination of type I receptor and type II receptor of TGF- β receptor, a combination of TGF- β type I receptor (TGFB1, activin receptor-like kinase (ALK5)) or ALK1, and TGF- β type II receptor (TGFB2) can be mentioned for TGF- β ; a combination of ALK4 or ALK7, and ActR-II or ActR-IIB can be mentioned for activin, Nodal; a combination of ALK2, ALK3 or ALK6, and BMPR-II can be mentioned for BMP; a combination of ALK2, ALK3 or ALK6, and ActR-II or ActR-IIB can be mentioned for GDF; and a combination of ALK2, ALK3 or ALK6, and ActR-II or ActR-IIB can be mentioned for AMH or MIS. As a preferable combination of type I receptor and type II receptor of

TGF- β receptor, a combination of ALK5 or ALK1, and TGFBR2 can be mentioned. The above-mentioned inhibitor of TGF- β receptor may be any as long as it suppresses the above-mentioned functions of the TGF- β receptor and, for example, a substance that inhibits
5 formation of TGF- β and TGF- β receptor complex, and the like can be mentioned.

[0031]

To be specific, as the TGF- β receptor inhibitor, for example, neutralizing antibodies against TGF- β receptor can be mentioned.
10 The antibody may be either a polyclonal antibody or a monoclonal antibody. These antibodies can be produced according to a production method of antibody or antiserum known per se. While the isotype of the antibody is not particularly limited, IgG, IgM or IgA is preferred, and IgG is particularly preferred. The
15 antibody is not particularly limited as long as it has at least a complementarity determining region (CDR) for specifically recognizing and binding to a target antigen, and may be a complete antibody molecule, a fragment such as Fab, Fab', F(ab')₂ or the like, a genetically engineered conjugate molecule such as scFv,
20 scFv-Fc, minibody, diabody or the like, a derivative thereof modified with a molecule having a protein stabilizing action such as polyethylene glycol (PEG) or the like, or the like. Since the neutralizing antibody is a neutralizing antibody contained in a serum-free medium or a xeno-free medium, when the biological cell
25 sample is derived from human, it is preferable to (i) obtain a human antibody by immunizing a human antibody-producing animal (e.g., mouse), (ii) produce a chimeric antibody, humanized antibody or complete human antibody, or (iii) obtain a human antibody by combining ex-vivo immunization method and cell
30 immortalization by virus, human-human (or mouse) hybridoma production technique, a phage display method, and the like. The

concentration of a neutralizing antibody against TGF- β receptors in a serum-free medium or a xeno-free medium is not particularly limited as long as it can inhibit intracellular signal transduction of the TGF- β receptor, and is, for example, 0.01 $\mu\text{g/mL}$ - 10 $\mu\text{g/mL}$, preferably 0.05 $\mu\text{g/mL}$ - 5 $\mu\text{g/mL}$, more preferably 0.1 $\mu\text{g/mL}$ - 2.5 $\mu\text{g/mL}$.

[0032]

In another preferred embodiment, the TGF- β receptor inhibitor is a low-molecular-weight compound showing an antagonist activity against TGF- β receptors. As used herein, the "antagonist activity" means an activity to bind to TGF- β receptor to inhibit the binding between TGF- β and TGF- β receptor. Examples of such compound include SB431542 (Stemgent), sc-203294, RepSox, Vactosertib (TEW-7197), SB525334, GW788388, SB505124, SD-208, LDN-193189, Galunisertib (LY2157299), LY2109761, LY364947, K02288, LDN-214117, ML347, LDN-212854, DMH1, Pirfenidone, LY 3200882, Alantolactone, SIS3, Hesperetin, A-83-01 and the like. The concentration of a low-molecular-weight compound showing an antagonist activity against TGF- β receptors in a serum-free medium or a xeno-free medium is not particularly limited as long as it can inhibit intracellular signal transduction of the TGF- β receptor, and is, for example, 0.1 μM - 100 μM , preferably 1 μM - 50 μM , more preferably 5 μM - 25 μM .

[0033]

The serum-free medium or xeno-free medium may be further appropriately supplemented with insulin, transferrin, selenium, various vitamins, L-glutamine, various amino acids such as non-essential amino acid and the like, 2-mercaptoethanol, various cytokines (interleukins (IL-2, IL-7, IL-15 etc.), SCF (Stem cell factor), activin and the like), various hormones, various growth factors (Leukemia inhibitory factor (LIF), basic fibroblast growth

factor (bFGF), etc.), antibiotics such as penicillin/streptomycin, puromycin and the like, pH indicator such as phenol red and the like, and the like.

[0034]

5 The biological cell sample containing mesenchymal stem cells which is cultured in step (1a) or (1b) of the present invention is not particularly limited as long as it is a cell sample containing mesenchymal stem cells. In one embodiment of the present invention, the biological cell sample containing mesenchymal stem
10 cells is a bone marrow-derived cell. The method for separating the cell derived from bone marrow may follow a known means. For example, it can be performed by removing interstitial cells from the collected bone marrow fluid by a density gradient centrifugation method using a separation medium with adjusted
15 density. Specifically, a layer of bone marrow-derived cells containing mesenchymal stem cells and mononuclear cells at the interface between the separation medium and the cerebrospinal fluid can be obtained by layering a bone marrow fluid diluted with physiological saline on the upper part of the separation medium in
20 a tube and centrifuging same. The thus-obtained bone marrow-derived cells contain a trace amount of mesenchymal stem cells. The proportion of the mesenchymal stem cells contained in the bone marrow-derived cells is not particularly limited, and is about 0.01% - about 1%, preferably about 0.01% - about 0.1%, of the
25 number of cells derived from the bone marrow.

[0035]

 The number of the bone marrow-derived cells including the mesenchymal stem cells cultured in step (1a) or (1b) of the present invention is not particularly limited, and may be
30 generally 0.5×10^5 cells/cm² - 25×10^5 cells/cm², preferably 2×10^5 cells/cm² - 13×10^5 cells/cm², per a culture container.

[0036]

The culture conditions for the bone marrow-derived cells containing mesenchymal stem cells are not particularly limited, and general cell culture conditions can be adopted. While the
5 aforementioned culture conditions include culturing at temperature 37°C, humidity 95%, and CO₂ concentration 5%, but the present invention is not limited to such conditions. For example, culturing at temperature 30 - 40°C, humidity 90 - 98%, CO₂ concentration 3 - 7% is exemplified. However, as long as
10 proliferation of desired cells can be achieved, temperature, humidity, and CO₂ concentration outside the ranges mentioned above may also be employed.

[0037]

It is preferable to change the medium at appropriate intervals
15 during the culture. The exchange of the medium includes the exchange of the whole amount of the medium, the exchange of a part of the medium, the addition of the medium, a combination thereof, and the like. In a preferred embodiment of the present invention, the whole amount of the medium is exchanged with a medium having the
20 same composition the day after the start of the culture, and the cells are cultured with the addition of 20% of the medium on day 3 and day 5 from the start of the culture.

[0038]

The culture period is, for example, 4 to 14 days, preferably
25 7 days. By this culture, mesenchymal stem cells contained in bone marrow-derived cells can be selectively adhered onto a culture container via vitronectin or a partial peptide of vitronectin. Adhered mesenchymal stem cells form cell aggregates. Here, the cell aggregate includes any cell populations of a cell population
30 that has proliferated so as to spread parallel to the adhesive surface of the culture container, a cell population that has

proliferated so as to vertically overlap the adhesive surface of the culture container, and a cell population having the characteristics of the both. When the culture period is less than 4 days, the number of cell aggregates that can be formed is small, and the number of cells for culturing in step (4) in the present invention to be described later cannot be secured. Furthermore, when the culture period exceeds 14 days, the cell aggregates collapse and the number of cells decreases. As a result, the number of cells for culturing in step (4) in the present invention cannot be secured.

[0039]

In another embodiment, the biological cell sample containing mesenchymal stem cells which is cultured in step (1a) or (1b) of the present invention is a cell derived from an adipose tissue. The method for separating the cell derived from an adipose tissue from the adipose tissue may follow a known means. For example, as a method for separating adipose tissue-derived cells including mesenchymal stem cells from adipose tissue, the collected adipose tissue is shredded, incubated in a collagenase solution, and filtered with a mesh sheet, whereby adipose tissue-derived cells including mesenchymal stem cells can be obtained. The thus-obtained cell derived from the adipose tissue contains a trace amount of mesenchymal stem cells. The proportion of the mesenchymal stem cells contained in the cells derived from the adipose tissue is not particularly limited, and is about 0.01% - about 1%, preferably about 0.1% - about 1%, of the number of cells derived from the adipose tissue.

[0040]

The number of adipose tissue-derived cells including the mesenchymal stem cells cultured in step (1a) or (1b) of the present invention is not particularly limited, and may be

generally 1×10^3 cells/cm² - 1×10^6 cells/cm², preferably 1×10^4 cells/cm² - 1×10^5 cells/cm², per a culture container.

[0041]

The culture conditions for the adipose tissue-derived cells
5 containing mesenchymal stem cells are not particularly limited,
and culture conditions similar to those for bone marrow-derived
cells containing mesenchymal stem cells can be adopted.

[0042]

It is preferable to change the medium at appropriate intervals
10 during the culture. The exchange of the medium includes the
exchange of the whole amount of the medium, the exchange of a part
of the medium, the addition of the medium, a combination thereof,
and the like. In a preferred embodiment of the present invention,
the whole amount of the medium is exchanged with a medium having the
15 same composition the day after the start of the culture and on day 2.

[0043]

The culture period is, for example, 1 to 14 days, preferably
5 days. By this culture, mesenchymal stem cells contained in
adipose tissue-derived cells can be selectively adhered onto a
20 culture container via vitronectin or a partial peptide of
vitronectin. Adhered mesenchymal stem cells form cell aggregates,
particularly, a cell population that has proliferated so as to
spread parallel to the adhesive surface of the culture container.

[0044]

25 The cell aggregate that is formed by mesenchymal stem cells
adhered onto a culture container via vitronectin or a partial
peptide of vitronectin in step (2a) or (2b) in the present
invention can be collected by a known means. For example, cells
other than the mesenchymal stem cells contained in the biological
30 cell sample do not adhere onto a culture container via vitronectin
or a partial peptide of vitronectin. Therefore, they are removed

from the culture container together with the serum-free medium or xeno-free medium by total amount medium change. As a result, cell aggregates of mesenchymal stem cells remain in the culture container in the culture after total medium exchange. The cell aggregates of
5 mesenchymal stem cells adhere to the culture container via vitronectin or a partial peptide of vitronectin. However, the cell aggregates also show weak cell-cell adhesion and easily separate from the cell aggregates and float in the medium. Therefore, the recovery of cell aggregates of mesenchymal stem cells may contain
10 2 steps of (i) recovery of cell aggregates of mesenchymal stem cells suspended in a serum-free medium or a xeno-free medium and (ii) recovery of cell aggregates of mesenchymal stem cells adhered on the culture container. The (i) recovery of cell aggregates of mesenchymal stem cells suspended in a serum-free medium or a xeno-
15 free medium can be performed, for example, by recovering the entire amount of the serum-free medium or xeno-free medium and centrifuging same. The (ii) recovery of cell aggregates of mesenchymal stem cells adhered on the culture container can be performed, for example, by easily detaching the aggregates from the culture
20 container only by pipetting, collecting the entire amount together with the medium or PBS, and centrifuging same. In another embodiment, for the recovery of cell aggregates of mesenchymal stem cells adhered on the culture container, adhesion between vitronectin or a partial peptide of vitronectin, and the cell
25 aggregates of mesenchymal stem cells is decomposed by a treatment with a detaching agent, and monocelled mesenchymal stem cells can be recovered. As the detaching agent, a mixed solution of trypsin and EDTA (generally 0.001-0.5% trypsin/0.1-5 mM EDTA, preferably about 0.1% trypsin/1 mM EDTA) may also be used, or a commercially
30 available product (e.g., TrypLE (Thermo Fisher Scientific)) may also be used.

[0045]

Vitronectin and a partial peptide of vitronectin show high adhesion activity to mesenchymal stem cells as compared with other extracellular matrices, and can efficiently adhere to mesenchymal stem cells contained in biological cell samples. Therefore, mesenchymal stem cells can be efficiently produced from a biological cell sample by doing as described above. However, when the cell aggregates of the mesenchymal stem cells adhered in step (1a) or (1b) in the present invention are continuously cultured for a long period of time, the proliferation of the mesenchymal stem cells is not observed depending on the origin of the cell sample. For example, when bone marrow-derived cells are cultured in step (1a) or (1b) in the present invention and cell aggregates of adhered mesenchymal stem cells are continuously cultured for a long period of time, proliferation of the mesenchymal stem cells is not observed. On the other hand, when the adipose tissue-derived cells are cultured in step (1a) or (1b) in the present invention, the cell aggregates of the adhered mesenchymal stem cells show confirmed proliferation. This is considered to be due to the presence of cells in the bone marrow-derived cells that prevent the proliferation of mesenchymal stem cells. Thus, when a large amount of mesenchymal stem cells is desired to be produced from biological cell samples containing mesenchymal stem cells by steps (1a) and (2a) or steps (1b) and (2b) in the present invention, it is preferable to re-seed the produced mesenchymal stem cells, proliferate them, and recover them again. Therefore, the production method of the present invention may further contain the following steps.

That is, a production method containing steps (1a) and (2a) of the present invention may further contain the following steps: (3a) a step of dissociating the collected cell aggregate (step

(3a) of the present invention),
(4a) a step of culturing the dissociated mesenchymal stem cells in a serum-free medium in the presence of an extracellular matrix protein or a partial peptide thereof capable of adhering
5 mesenchymal stem cells (step (4a) of the present invention),
(5a) a step of collecting the mesenchymal stem cells proliferated on the culture container via the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells (step (5a) of the present invention).

10 In addition, a production method containing steps (1b) and (2b) of the present invention may further contain the following steps:
(3b) a step of dissociating the collected cell aggregate (step (3b) of the present invention),
(4b) a step of culturing the dissociated mesenchymal stem cells in
15 a xeno-free medium in the presence of an extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells (step (4b) of the present invention),
(5b) a step of collecting the mesenchymal stem cells proliferated on the culture container via the extracellular matrix protein or a
20 partial peptide thereof capable of adhering mesenchymal stem cells (step (5b) of the present invention).

[0046]

In step (3a) or (3b) in the present invention, the dissociation of the recovered cell aggregate is performed by a
25 known means. Since cell aggregates show weak cell-cell adhesion, for example, cell-cell adhesion of cell aggregates can be easily eliminated by pipetting alone, and a cell population of single mesenchymal stem cells can be prepared. Alternatively, it can also be performed by treating with the above-mentioned detaching
30 agent.

[0047]

In step (4a) or (4b) in the present invention, the culture container used for culturing, serum-free medium, xeno-free medium, the mode of contact between mesenchymal stem cells and extracellular matrix proteins during culture, and the like may be
5 the same as those in step (1a) or (1b) in the present invention.
[0048]

In step (4a) or (4b) in the present invention, culture is performed in the presence of an extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells
10 (hereinafter to be indicated as "partial peptide of extracellular matrix protein"). The extracellular matrix protein is not particularly limited as long as it can cause adhesion of mesenchymal stem cells to a culture container. Examples of such extracellular matrix protein include vitronectin, fibronectin,
15 laminin, collagen and the like. In addition, examples of the partial peptide of an extracellular matrix protein include iMatrix-511 (partial peptide of laminin-511) and the like.
[0049]

Like vitronectin, the extracellular matrix protein may be
20 any of a protein isolated and purified from mammalian cells, etc., a biochemically-synthesized protein, and a recombinant protein produced from a transformant incorporating a nucleic acid having a base sequence encoding an extracellular matrix protein.
[0050]

25 In addition, the partial peptide of the extracellular matrix protein may be any as long as it is a peptide having a partial amino acid sequence of the extracellular matrix protein and has mesenchymal stem cell adhesion activity. Examples of such partial peptide of extracellular matrix protein include proteins
30 containing at least one domain selected from the group consisting of RGD domain and heparin binding domain.

[0051]

The tissue from which the mesenchymal stem cells seeded in step (4a) or (4b) in the present invention is derived is not particularly limited. In step (1a) or (1b) in the present
5 invention, a tissue in which mesenchymal stem cells of cell aggregates do not grow sufficiently is preferred. Examples of such tissue include bone marrow, cord blood and the like. Even when the mesenchymal stem cells of the cell aggregates proliferate in step (1a) or (1b) of the present invention, mesenchymal stem
10 cells may be cultured in step (4a) or (4b) in the present invention for the purpose of further increasing the number of mesenchymal stem cells.

[0052]

The number of the mesenchymal stem cells seeded in step (4a)
15 or (4b) in the present invention is not particularly limited, and may be generally 2×10^5 cells/cm² - 26×10^5 cells/cm², preferably 8×10^5 cells/cm² - 13×10^5 cells/cm², per a culture container.

[0053]

The culture conditions for the mesenchymal stem cells are
20 not particularly limited, and may be the same as those for the biological cell sample containing mesenchymal stem cells. General cell culture conditions can be adopted.

[0054]

It is preferable to change the medium at appropriate intervals
25 during the culture. The exchange of the medium includes the exchange of the whole amount of the medium, the exchange of a part of the medium, the addition of the medium, a combination thereof, and the like. In a preferred embodiment of the present invention, the whole amount of the medium is exchanged with a medium having the
30 same composition every 2 or 3 days from the day when the culture was started.

[0055]

The culture period is, for example, 1 to 14 days, preferably 1 to 8 days. By this culture, mesenchymal stem cells start to proliferate.

5 [0056]

The mesenchymal stem cells that have proliferated on a culture container via an extracellular matrix protein or a partial peptide of the extracellular matrix protein in step (5a) or (5b) in the present invention are collected by a known means. The
10 means of collection may be the same as the method described in step (2a) or (2b) in the present invention.

[0057]

The present invention is explained in more detail in the following by referring to Examples. These are mere
15 exemplifications and the present invention is not limited thereby.

[Example]

[0058]

Example 1: Study of mesenchymal stem cell (MSC) production promoting effect by Vitronectin

20 Due to the limitations of purification techniques, bone marrow mononuclear cells (MNC) separated from bone marrow are slightly contained with mesenchymal stem cells (MSC). In this Example, a method for producing MSC from MNC by using a serum-free medium is verified. Bone marrow mononuclear cells (MNC) (Lonza) were thawed
25 using the following medium for seeding. The above-mentioned cells were seeded at a concentration of 2.6×10^6 cells/well in a 24-well plate coated with Fibronectin (Sigma) or Vitronectin (wako) each at a concentration of $1.5 \mu\text{g}/\text{cm}^2$, and cultured under the conditions of
30 37°C , 5% CO_2 . The next day of seeding, the entire amount of the medium in the plate was replaced with a medium for seeding, and the medium for seeding in an amount corresponding to 20% of the amount

of the medium in the plate was further added on day 3 and day 5 after seeding.

Medium for seeding: StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), 10 μ M SB431542 (Stemgent), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

10 Fig. 1 shows photographs of the cells on day 5 after seeding. More cell aggregates were formed in a well plate coated with Vitronectin than in a well plate coated with Fibronectin.

Aggregates were collected day 7 after seeding and the cells were re-seeded using the following medium for proliferation.

15 Specifically, after collecting the culture supernatant, DPBS (Nacalai Tesque) was added to the plate, and the aggregates were detached from the plate by pipetting, and all aggregates were collected together with DPBS. Thereafter, the collected culture supernatant and DPBS were combined and centrifuged, and only the 20 aggregates were collected. The aggregates were dissociated into single cells by re-suspending the collected aggregates in the proliferation medium. Using Fibronectin (Sigma), Vitronectin (wako) and iMatrix-511 (Nippi, Inc.), the entire amount of the collected cells was seeded in 24-well plates coated with 25 Fibronectin or Vitronectin at a concentration of 1.5 μ g/cm², or iMatrix-511 at a concentration of 0.5 μ g/cm², and cultured under the conditions of 37°C, 5% CO₂. Thereafter, the entire amount of the medium in the plate was replaced with the medium for proliferation every 2-3 days until the cells became sub-confluent. 30 Medium for proliferation: StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered

trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION C, 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

After confirming that the cells became sub-confluent, the cells were passaged on day 13 after seeding, and the number of cells was measured. Fig. 2 shows the measurement results of the cell number. In addition, Fig. 3 shows photographs of the cells. It was found that the difference in the extracellular matrix for coating the culture container in re-seeding does not affect the number of cells obtained.

From the above results, it was found that the number of cell aggregates obtained and the number of cells obtained by the subsequent re-seeding were larger when a culture container coated with Vitronectin was used at the time of seeding of the cells than when a culture container coated with Fibronectin was used.

Furthermore, the cells detached from the plate coated with Vitronectin were expansion cultured, and surface antigen analysis was performed. Using FACS, surface antigen analysis of 3 kinds of MSC positive markers (CD105, CD90, CD73) and 2 kinds of MSC negative markers (CD45, CD34) was performed. Table 1 shows the analysis results. It could be confirmed that the obtained cells were positive for CD105, CD90, CD73, negative for CD45, CD34, and were MSCs.

[0059]

[Table 1]

		MSC isolated with serum-free medium + Vitronectin	general MSC
MSC markers	CD105	+	+
	CD90	+	+

	CD73	+	+
blood cell markers	CD34	-	-
	CD45	-	-

[0060]

Example 2: Study of difference in MSC production promoting effect by the kind of Vitronectin

5 MNC (Lonza) was thawed using the following medium for seeding (serum-free). The above-mentioned cells were seeded at a concentration of 2.6×10^6 cells/well in 24-well plates coated with Vitronectin (20-398 aa) (wako) (corresponding to amino acid Nos. 1 - 379 of SEQ ID NO: 1), Vitronectin (VTN-N, 62-478 aa) (Life
10 Technologies) (corresponding to amino acid Nos. 43 - 459 of SEQ ID NO: 1) or Vitronectin (Full length, 20-478 aa) (Sigma) (corresponding to SEQ ID NO: 1) at a concentration of $1.5 \mu\text{g}/\text{cm}^2$, and cultured under the conditions of 37°C , 5% CO_2 . In addition, the cells were seeded at a concentration of 1.6×10^6 cells/well in a
15 24-well plate coated with Vitronectin (20-398 aa) (wako) or Synthemax II (CORNING) at concentrations of $1.5 \mu\text{g}/\text{cm}^2$ and $5.0 \mu\text{g}/\text{cm}^2$, respectively, and cultured under the conditions of 37°C , 5% CO_2 . Synthemax II is a vitronectin-based synthetic peptide containing RGD motif and flanking sequence. The next day of
20 seeding, the entire amount of the medium in the plate was replaced with a medium for seeding, and the medium for seeding in an amount corresponding to 20% of the amount of the medium in the plate was further added on day 3 and day 5 after seeding.

Medium for seeding: StemFit (registered trade mark) AK03N medium
25 (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), $10 \mu\text{M}$ SB431542 (Stemgent), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM

Lithium Chloride (Sigma-Aldrich)

Aggregates were collected day 7 after seeding and the cells were re-seeded using the following medium for proliferation. Specifically, after collecting the culture supernatant, DPBS (Nacalai Tesque) was added to the plate, and the aggregates were detached from the plate by pipetting, and all aggregates were collected together with DPBS. Thereafter, the collected culture supernatant and DPBS were combined and centrifuged, and only the aggregates were collected. The aggregates were dissociated into single cells by resuspending the collected aggregates in the proliferation medium. the entire amount of the collected cells was seeded in a 24-well plate, and cultured under the conditions of 37°C, 5% CO₂. Thereafter, the entire amount of the medium in the plate was replaced with the medium for proliferation every 2-3 days until the cells became sub-confluent.

Medium for proliferation: StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION C, 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich), 0.2 µg/mL iMatrix 511 (Nippi, Inc.)

After confirming that the cells became sub-confluent, the cells were passaged on days 12 and 13 after seeding, and the number of cells was measured. Fig. 4 shows the measurement results of the cell number when Vitronectin (20-398 aa) (wako), Vitronectin (VTN-N, 62-478aa) (Life Technologies) or Vitronectin (Full length, 20-478 aa) (Sigma) were used. While MSC could be produced using any vitronectin when seeding MNC, Vitronectin (20-398 aa) (wako) could produce MSC most efficiently from MNC. Fig. 5 shows the

measurement results of the cell number when Vitronectin (20-398 aa) (wako) or Synthemax II (CORNING) were used. MSC could be efficiently produced from MNC even when Synthemax II (CORNING) was used when seeding MNC.

5 [0061]

Example 3: Study of MSC production promoting effect by TGF β receptor inhibitor

MNC (Lonza) was thawed using the following medium for seeding (1) or (2). The above-mentioned cells were seeded at a
10 concentration of 2.6×10^6 cells/well in a 24-well plate coated with Vitronectin (VTN-N, 62-478 aa) (Life Technologies) at a concentration of $1.5 \mu\text{g}/\text{cm}^2$, and cultured under the conditions of 37°C , 5% CO_2 . The next day of seeding, the entire amount of the medium in the plate was replaced with a medium for seeding (1) or
15 (2), and the medium for seeding in an amount corresponding to 20% of the amount of the medium (1) or (2) in the plate was further added on day 3 and day 5 after seeding.

Medium for seeding (1) (TGF β inhibitor (-)): StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, StemFit
20 (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

25 Medium for seeding (2) (TGF β inhibitor (+)): StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), 10 μM SB431542 (Stemgent), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone
30 (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

Aggregates were collected day 7 after seeding and the cells were re-seeded using the following medium for proliferation. Specifically, after collecting the culture supernatant, DPBS (Nacalai Tesque) was added to the plate, and the aggregates were detached from the plate by pipetting, and all aggregates were collected together with DPBS. Thereafter, the collected culture supernatant was combined and centrifuged, and only the aggregates were collected. The aggregates were dissociated into single cells by resuspending the collected aggregates in the proliferation medium. The entire amount of the collected cells was seeded in a 24-well plate, and cultured under the conditions of 37°C, 5% CO₂. Thereafter, the entire amount of the medium in the plate was replaced with the medium for proliferation every 2-3 days until the cells became sub-confluent.

Medium for proliferation: StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION C, 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich), 0.2 µg/mL iMatrix 511 (Nippi, Inc.)

After confirming that the cells became sub-confluent, the cells were passaged on day 15 after seeding, and the number of cells was measured. Fig. 6 shows the measurement results of the cell number. MSC could be efficiently produced from MNC when the medium for seeding contained a TGFβ inhibitor at the time of seeding MNC.

[0062]

Example 4: Study of difference in MSC production promoting effect by the kind of TGFβ receptor inhibitor

MNC (Lonza) was thawed using the following medium for seeding

(1), (2) or (3). The above-mentioned cells were seeded at a concentration of 2.6×10^6 cells/well in a 24-well plate coated with Vitronectin (20-398 aa) (wako) at a concentration of $1.5 \mu\text{g}/\text{cm}^2$, and cultured under the conditions of 37°C , 5% CO_2 . The next day of seeding, the entire amount of the medium in the plate was replaced with a medium for seeding (1), (2) or (3), and the medium for seeding (1), (2) or (3) in an amount corresponding to 20% of the amount of the medium in the plate was further added on day 3 and day 5 after seeding.

10 Medium for seeding (1): StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), 10 μM SB431542 (Stemgent), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-
15 Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

Medium for seeding (2): StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL
20 bFGF (peprotech), 0.5 μM A-83-01 (wako), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

Medium for seeding (3): StemFit (registered trade mark) AK03N
25 medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), 0.5 μM LDN-193189 (Stemgent), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-
Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries,
30 Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)

Aggregates were collected day 7 after seeding and the cells

were re-seeded using the following medium for proliferation. Specifically, after collecting the culture supernatant, DPBS (Nacalai Tesque) was added to the plate, and the aggregates were detached from the plate by pipetting, and all aggregates were collected together with DPBS. Thereafter, the collected culture supernatant and DPBS were combined and centrifuged, and only the aggregates were collected. The aggregates were dissociated into single cells by resuspending the collected aggregates in the proliferation medium. the entire amount of the collected cells was seeded in a 24-well plate, and cultured under the conditions of 37°C, 5% CO₂. Thereafter, the entire amount of the medium in the plate was replaced with the medium for proliferation every 2-3 days until the cells became sub-confluent.

Medium for proliferation: StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION C, 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich), 0.2 µg/mL iMatrix 511 (Nippi, Inc.)

After confirming that the cells became sub-confluent, the cells were passaged on day 12 after seeding, and the number of cells was measured. Fig. 7 shows the measurement results of the cell number. MSC could be efficiently produced from MNC by using any of TGFβ receptor inhibitors SB431542 (ALK5 inhibition), A-83-01 (ALK4, ALK5, ALK7 inhibition) and LDN-193189 (ALK2, ALK3 inhibition).

[0063]

Example 5: Study of MSC production from adipose tissue

Adipose tissue surrounding epididymis collected from C57BL/6J mice (11-week-old, male) was treated with collagenase,

and cells were obtained using the following medium for seeding (1) or (2). The above-mentioned cells were seeded at a concentration of 6.0×10^4 cells/well in a 24-well plate, and cultured under the conditions of 37°C, 5% CO₂. A 24-well plate coated with Vitronectin (VTN-N, 62-478 aa) (Life Technologies) at a concentration of 1.5 $\mu\text{g}/\text{cm}^2$ was used for the cells seeded in medium for seeding (1). The next day of seeding and day 2 after seeding, the entire amount of the medium in the plate was replaced with medium for seeding (1) or (2).

10 Medium for seeding (1): StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION A, 1/4 StemFit (registered trade mark) AK03N medium (Ajinomoto Co., Inc.) SOLUTION B, 3 ng/mL bFGF (peprotech), 10 μM SB431542 (Stemgent), 1/100 Lipid Concentrate (Life Technologies), 10 nM Dexamethasone (Sigma-
15 Aldrich), 10 ng/mL PDGF-BB (FUJIFILM Wako Pure Chemical Industries, Ltd.), 1 mM Lithium Chloride (Sigma-Aldrich)
Medium for seeding (2): DMEM medium (sigma), 10% fetal calf serum (Life Technologies)

After confirming that the cells became sub-confluent, the
20 cells were passaged on day 5 after seeding, and the number of cells was measured. Fig. 8 shows the measurement results of the cell number. Fig. 9 shows photographs of the cells. MSC could be efficiently produced from adipose tissue by using a medium containing a TGF β receptor inhibitor and vitronectin.

25 [Industrial Applicability]
[0064]

Mesenchymal stem cells can be efficiently produced from a biological cell sample by culturing the biological cell sample containing mesenchymal stem cells in a serum-free medium or a
30 xeno-free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells. By

adopting this method, the obtained mesenchymal stem cells can be directly used as a cell source in regenerative medicine.

This application is based on a patent application No. 2019-156537 filed in Japan (filing date: August 29, 2019) and a patent
5 application No. 2020-012333 filed in Japan (filing date: January 29, 2020), the contents of which are incorporated in full herein.

CLAIMS

[Claim 1]

A method for producing a mesenchymal stem cell from a biological cell sample comprising mesenchymal stem cells,
5 comprising the following steps:

- (1) a step of culturing the biological cell sample comprising mesenchymal stem cells in a serum-free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells,
- 10 (2) a step of collecting a cell aggregate of the mesenchymal stem cells.

[Claim 2]

The method according to claim 1, wherein the culture in the presence of vitronectin or a partial peptide thereof capable of
15 adhering mesenchymal stem cells is performed by culturing on a culture container on which vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells has been immobilized.

[Claim 3]

20 The method according to claim 1 or 2, further comprising the following steps:

- (3) a step of dissociating the collected cell aggregate,
- (4) a step of culturing the dissociated mesenchymal stem cells in a serum-free medium in the presence of an extracellular
25 matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells,
- (5) a step of collecting the mesenchymal stem cells proliferated on the culture container via the extracellular matrix protein or a partial peptide thereof capable of adhering
30 mesenchymal stem cells.

[Claim 4]

The method according to claim 3, wherein the culture in the presence of the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a culture container on which the extracellular matrix
5 protein or a partial peptide thereof capable of adhering mesenchymal stem cells has been immobilized.

[Claim 5]

The method according to any one of claims 1 to 4, wherein the partial peptide of vitronectin comprises an RGD domain.

10 [Claim 6]

The method according to claim 5, wherein the partial peptide of vitronectin further comprises a somatomedin B domain.

[Claim 7]

The method according to claim 6, wherein the partial peptide
15 of vitronectin is a polypeptide consisting of amino acid Nos. 1 - 379 of the amino acid sequence shown in SEQ ID NO: 1.

[Claim 8]

The method according to any one of claims 1 to 7, wherein the serum-free medium in step (1) comprises a TGF- β receptor
20 inhibitor.

[Claim 9]

A method for producing a mesenchymal stem cell from a biological cell sample comprising mesenchymal stem cells, comprising the following steps:

25 (1) a step of culturing the biological cell sample comprising mesenchymal stem cells in a xeno-free medium in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells,

(2) a step of collecting a cell aggregate of the mesenchymal
30 stem cells.

[Claim 10]

The method according to claim 9, wherein the culture in the presence of vitronectin or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a culture container on which vitronectin or a partial peptide
5 thereof capable of adhering mesenchymal stem cells has been immobilized.

[Claim 11]

The method according to claim 9 or 10, further comprising the following steps:

10 (3) a step of dissociating the collected cell aggregate,
(4) a step of culturing the dissociated mesenchymal stem cells in a xeno-free medium in the presence of an extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells,

15 (5) a step of collecting the mesenchymal stem cells proliferated on the culture container via the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells.

[Claim 12]

20 The method according to claim 11, wherein the culture in the presence of the extracellular matrix protein or a partial peptide thereof capable of adhering mesenchymal stem cells is performed by culturing on a culture container on which the extracellular matrix protein or a partial peptide thereof capable of adhering
25 mesenchymal stem cells has been immobilized.

[Claim 13]

The method according to any one of claims 9 to 12, wherein the partial peptide of vitronectin comprises an RGD domain.

[Claim 14]

30 The method according to claim 13, wherein the partial peptide of vitronectin further comprises a somatomedin B domain.

[Claim 15]

The method according to claim 14, wherein the partial peptide of vitronectin is a polypeptide consisting of amino acid Nos. 1 - 379 of the amino acid sequence shown in SEQ ID NO: 1.

5 [Claim 16]

The method according to any one of claims 9 to 15, wherein the xeno-free medium in step (1) comprises a TGF- β receptor inhibitor.

[Claim 17]

10 The method according to any one of claims 9 to 16, wherein the xeno-free medium comprises an allogeneic serum.

[Claim 18]

The method according to claim 17, wherein the allogeneic serum is an autologous serum.

15 [Claim 19]

The method according to any one of claims 1 to 18, wherein the biological cell sample comprising mesenchymal stem cells is a bone marrow-derived cell.

[Claim 20]

20 The method according to claim 19, wherein the number of the cells derived from bone marrow and to be cultured is 0.5×10^5 - 25×10^5 cells/cm².

[Claim 21]

25 The method according to claim 19 or 20, wherein the cells derived from bone marrow are cultured for a period of 4 days to 14 days.

[Claim 22]

The method according to any one of claims 1 to 18, wherein the biological cell sample comprising mesenchymal stem cells is a
30 cell derived from adipose tissue.

[Claim 23]


The method according to claim 22, wherein the number of the cells derived from adipose tissue and to be cultured is 1×10^3 - 1×10^6 cells/cm².

[Claim 24]


5 The method according to claim 22 or 23, wherein the cells derived from adipose tissue are cultured for a period of 1 day to 14 days.

[DRAWINGS]


[Fig. 1]




[Fig. 2]




[Fig. 3]




[Fig. 4]




[Fig. 5]




[Fig. 6]



[Fig. 7]



[Fig. 8]



[Fig. 9]

