

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199667123 B2
(10) Patent No. 727480

(54) Title
Human CTLA-8 and uses of CTLA-8-related proteins

(51)⁶ International Patent Classification(s)
C12N 015/12 C07K 014/725
A61K 038/17 C07K 016/28
A61K 038/20 C12N 005/10
C07K 014/54 C12N 015/24

(21) Application No: 199667123 (22) Application Date: 1996.07.18

(87) WIPO No: W097/04097

(30) Priority Data

(31) Number	(32) Date	(33) Country
08/504032	1995.07.19	US
08/514014	1995.08.11	US

(43) Publication Date : 1997.02.18
(43) Publication Journal Date : 1997.04.17
(44) Accepted Journal Date : 2000.12.14

(71) Applicant(s)
Genetics Institute, Inc.

(72) Inventor(s)
Kenneth Jacobs; Kerry Kelleher; McKeough Carlin; Samuel Goldman;
Debra Pittman; Sha Mi; Steven Neben; Joann Giannotti; Margaret
Golden-Fleet

(74) Agent/Attorney
F.B. RICE and CO.,139 Rathdowne Street,CARLTON VIC 3053

(56) Related Art
WO 95/18826
ROOVIES ET AL J. IMMUNOL. 1993 150:5445-5456

**CORRECTED
VERSION***

pages 1/7-7/7, drawings, replaced by new pages bearing the same number, due to late transmittal by the receiving Office

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



67123/26

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/19, 15/12, 15/24, C07K 14/52, 14/725, 14/54, A61K 38/17, 38/19, 38/20, C12N 5/10, C07K 16/24, 16/28, A61K 48/00		A3	(11) International Publication Number: WO 97/04097 (43) International Publication Date: 6 February 1997 (06.02.97)
(21) International Application Number: PCT/US96/11889 (22) International Filing Date: 18 July 1996 (18.07.96) (30) Priority Data: 08/504,032 19 July 1995 (19.07.95) US 08/514,014 11 August 1995 (11.08.95) US (71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 Cambridge Park Drive, Cambridge, MA 02140 (US). (72) Inventors: JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). KELLEHER, Kerry; 50 Hurley Circle, Marlborough, MA 01752 (US). CARLIN, McKeough; 16 Chauncy Street #2, Cambridge, MA 02138 (US). GOLDMAN, Samuel; 9 Mohawk Drive, Acton, MA 01720 (US). PITTMAN, Debra; 20 N. Shore Road, Windham, NH 03087 (US). MI, Sha; 4 Vernon Road, Belmont, MA 02178 (US). NEBEN, Steven; 13 Duggan Road, Acton, MA 01720 (US). GIANNOTTI, JoAnn; 409 Arlington Street, Acton, MA 01720 (US). GOLDEN-FLEET, Margaret; 19 Usher Road, Medford, MA 02155 (US). (74) Agent: BROWN, Scott, A.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).		(81) Designated States: AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 12 September 1997 (12.09.97)	
(54) Title: HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS (57) Abstract Polynucleotides encoding human CTLA-8 and related proteins are disclosed. Human CTLA-8 proteins and methods for their production are also disclosed. Methods of treatment using human CTLA-8 proteins, rat CTLA-8 proteins and herpesvirus herpes CTLA-8 proteins are also provided.			

* (Referred to in PCT Gazette No. 45/1997, Section II)

HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS

This application is a continuation-in-part of application Ser. No. 08/504,032, filed July 19, 1995, and a continuation-in-part of application Ser. No. 08/514,014, filed August 11, 1995.

Field of the Invention

- 5 The present invention relates to human CTLA-8 proteins, nucleic acids encoding such proteins, methods of treatment using such proteins. The invention also relates to the use of rat CTLA-8 proteins and herpesvirus *Saimiri* ORF13 proteins in methods of treatment.

10 Background of the Invention

- Cytokines are secreted proteins which act on specific hematopoietic target cells to cause a differentiation event or on other target cells to induce a particular physiological response, such as secretion of proteins characteristic of inflammation. Cytokines, also variously known as lymphokines, hematopoietins, interleukins, colony stimulating factors, and the like, can be important therapeutic agents, especially for
15 diseases or conditions in which a specific cell population is depleted. For example, erythropoietin, G-CSF, and GM-CSF, have all become important for treatment of anemia and leukopenia, respectively. Other cytokines such as interleukin-3, interleukin-6, interleukin-11 and interleukin-12 show promise in treatment of
20 conditions such as thrombocytopenia and modulation of immune response.

- For these reasons a significant research effort has been expended in searching for novel cytokines and cloning the DNAs which encode them. In the past, novel cytokines were identified by assaying a particular cell such as a bone marrow cell, for a measurable response, such as proliferation. The search for novel cytokines has thus
5 been limited by the assays available, and if a novel cytokine has an activity which is unmeasurable by a known assay, the cytokine remains undetectable. In a newer approach, cDNAs encoding cytokines have been detected using the polymerase chain

reaction (PCR) and oligonucleotide primers having homology to share motifs of known cytokines or their receptors. The PCR approach is also limited by the necessity for knowledge of previously cloned cytokines in the same protein family. Cytokines have also been cloned using subtractive hybridization to construct and screen cDNA libraries, or they can potentially be cloned using PCR followed by gene electrophoresis to detect differentially expressed genes. The subtractive hybridization methods are based on the assumption that cytokine mRNAs are those that are differentially expressed, and these methods do not require any prior knowledge of the sequence of interest. However, many cytokines may be encoded by mRNAs which are not differentially expressed, and thus are undetectable using these methods.

It would be desirable to develop new methods for identifying novel cytokines and other secreted factors and to isolate polynucleotides encoding them.

Summary of the Invention

In developing the present invention, methods were employed which selectively identify polynucleotides which encode secreted proteins. One such polynucleotide was isolated which encodes "human CTLA-8". In accordance with the present invention, polynucleotides encoding human CTLA-8 and active fragments thereof are disclosed. "CTLA-8" is used throughout the present specification to refer to both proteins and polynucleotides encoding those proteins and to refer to proteins and polynucleotides from all mammalian species.

In a first aspect, the present invention provides an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544;
- (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 85 to nucleotide 544;
- (c) the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544;
- (d) the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;
- (e) the nucleotide sequence of the full length protein coding sequence of clone B18 deposited under accession number ATCC 69868;



(f) a nucleotide encoding the full length protein encoded by the cDNA insert of clone B18 deposited under accession number ATCC 69868;

(g) a nucleotide sequence that specifically hybridizes to a nucleic acid sequence specified in (a) to (f);

5 (h) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) to (f) as a result of degeneracy of the genetic code; and

(i) an allelic variant of the nucleotide specified in (a) to (f).

In a second aspect, the present invention provides an isolated human CTLA-8 protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:2;

(b) the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;

15 (c) the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;

(d) the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163;

(e) a fragment of SEQ ID NO:2, the fragment comprising amino acids 11 to 163 of SEQ ID NO:2;

20 (f) a fragment of SEQ ID NO:2, the fragment comprising amino acids 29 to 163 of SEQ ID NO:2;

(g) a fragment of SEQ ID NO:2, the fragment comprising amino acids 31 to 163 of SEQ ID NO:2; and

25 (h) fragments of (a), (b), (c), (d), (e), (f) or (g) having a CTLA-8 activity selected from the group consisting of: biological activity in a factor-dependent cell proliferation assay; induction of expression or secretion of γ -IFN; chemoattractant or chemotactic activity in a chemoattraction or chemotaxis assay; and induction or expression of secretion of IL-3 or GM-CSF.

30 In a third aspect, the present invention provides a human CTLA-8 protein produced according to a process which comprises the steps of:

(a) growing a culture of a cell in a suitable culture medium, wherein the cell is transformed with a polynucleotide operably linked to an expression control sequence, wherein the polynucleotide is selected from the group consisting of:

35 (aa) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544;



(ab) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;

(ac) a polynucleotide comprising the nucleotide sequence of the full-length coding sequence of clone B18 deposited under accession number ATCC 69868;

(ad) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone B18 deposited under accession number ATCC 69868; and

(ae) a polynucleotide comprising a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (aa) to (ad) as a result of degeneracy of the genetic code; and

(b) purifying the human CTLA-8 protein encoded by said polynucleotide from the culture.

In certain embodiments, the present invention provides an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;

(b) a nucleotide sequence capable of hybridizing to a nucleic acid sequence specified in (a);

(c) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) as a result of degeneracy of the genetic code; and

(d) an allelic variant of the nucleotide sequence specified in (a).

Preferably, the polynucleotide of the invention encodes a protein having CTLA-8 activity. In other embodiments the polynucleotide is operably linked to an expression control sequence. In other preferred embodiments, the polynucleotide is contained in a vector suitable for *in vivo* expression in a mammalian subject. Polynucleotides comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544, the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544 or the nucleotide sequence of SEQ ID NO:1 from nucleotide 86 to nucleotide 544 are particularly preferred.

Host cells transformed with the polynucleotides of the invention are also provided, comprising mammalian cells.

Processes are also provided for producing a human CTLA-8 protein, said processes comprising:



(a) growing a culture of the host cell of the invention in a suitable culture medium; and

(b) purifying the human CTLA-8 protein from the culture.

Isolated human CTLA-8 protein is also provided comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:2;

(b) the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;

(c) the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;

(d) the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163; and

(e) fragments of (a), (b), (c) or (d) having CTLA-8 activity.

Proteins comprising the amino acid sequence of SEQ ID NO:2 and comprising the sequence from amino acids 29 to 163, from amino acid 31 to 163, or from amino acids 11 to 163 of SEQ ID NO:2 are particularly preferred. Preferably, the protein has CTLA-8 activity. Pharmaceutical compositions comprising a human CTLA-8 protein of the invention and a pharmaceutically acceptable carrier are also provided.

Compositions are also disclosed which comprise an antibody which specifically reacts with a human CTLA-8 protein of the invention.

Methods of treating a mammalian subject are also provided which comprise administering a therapeutically effective amount of a pharmaceutical composition comprising a human CTLA-8 protein.

Rat CTLA-8 and active (*i.e.*, having CTLA-8 activity) fragments thereof may also be used in such methods of treatment. Preferably the rat protein is administered as a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

(b) the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150; and

(c) fragments of (a) or (b) having CTLA-8 activity.

Herpesvirus *Saimiri* ORF13, referred to herein as "herpes CTLA-8", and active (*i.e.*, having CTLA-8 activity) fragments thereof and active fragments thereof may also be used in such methods of treatment. Preferably the herpes




CTLA-8 protein is administered as a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
- 5 (b) the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151; and
- (c) fragments of (a) or (b) having CTLA-8 activity.

The invention also provides a method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition
10 comprising a pharmaceutically acceptable carrier and IL-17 or an active fragment thereof.

In methods of treatment provided by the present invention, preferably the subject is treated to produce an effect selected from the group consisting of inhibition of angiogenesis, inhibition of growth or proliferation of vascular
15 endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, proliferation of myeloid cells or progenitors, proliferation of erythroid cells or progenitors, proliferation of lymphoid cells or progenitors, induction of IFN γ production, induction of IL-3 production and induction of GM-CSF production.

20 Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.



Brief Description of the Figures

Fig. 1 is a comparison of homologous regions of the amino acid sequences of human CTLA-8 (indicated as "B18_F1"), rat CTLA-8 (indicated as "Musctla8") and herpes CTLA-8 (indicated as "Hsvie_2").

5 Fig. 2 depicts autoradiographs demonstrating expression of human CTLA-8 in COS cells.

Fig. 3 presents data relating to the ability of human CTLA-8 to inhibit angiogenesis.

10 Figs. 4 and 5 present data relating to the ability of human CTLA-8 to produce or induce hematopoietic activity.

Figs. 6 and 7 present data demonstrating the ability of human CTLA-8 to induce production of IL-6 and IL-8.

Detailed Description of Preferred Embodiments

15 The inventors of the present application have identified and provided a polynucleotide encoding a human CTLA-8 protein. SEQ ID NO:1 provides the nucleotide sequence of a cDNA encoding the human CTLA-8 protein. SEQ ID NO:2 provides the amino acid sequence of the human CTLA-8 protein. Alternatively, the initiating methionine may be at amino acid 11 of SEQ ID NO:2. On the basis of
20 amino terminal sequencing, the mature protein sequence is believed to begin at amino acid 31 of SEQ ID NO:2 (encoded by the sequence beginning with nucleotide 146 of SEQ ID NO:1).

The region from amino acid 29 to amino acid 163 of human CTLA-8 (SEQ ID NO:2) shows marked homology to portions of rat CTLA-8 (amino acids 18 to 150 of
25 SEQ ID NO:4) and herpesvirus *Saimiri* ORF13 ("herpes CTLA-8") (amino acids 19 to 151 of SEQ ID NO:5). A cDNA sequence encoding rat CTLA-8 is listed at SEQ ID NO:3 and its corresponding amino acid sequence is reported at SEQ ID NO:4. A cDNA sequence encoding herpes CTLA-8 is listed at SEQ ID NO:5 and its corresponding amino acid sequence is reported at SEQ ID NO:6. Homology between
30 rat CTLA-8 and herpes CTLA-8 was reported by Rouvier et al., J. Immunol. 1993, 150, 5445-5456.

Applicants had previously incorrectly identified the rat sequences of SEQ ID NO:3 and SEQ ID NO:4 as applying to murine CTLA-8. Applicants' human CTLA-8 (B18) does also show homolgy to the true murine CTLA-8 sequence.

5 Golstein et al. (WO95/18826; Fossiez et al., Microbial Evasion and Subversion of Immunity 544:3222 (Abstract)) have also reported a species they initially identified as "human CTLA-8." However, examination of the sequence of the Golstein et al. species and the human CTLA-8 (B18) sequence of the present invention readily reveals that they are two different proteins, although they are homologous with each other and with the rat CTLA-8 and herpes CTLA-8 identified herein. The Golstein
10 et al. species has now been renamed as interleukin-17 (IL-17). Because of the homology between applicants' human CTLA-8 (B18) and IL-17, these proteins are expected to share some activities.

It has also been preliminarily determined that human CTLA-8 (B18) forms homodimers when expressed. As a result, human CTLA-8 proteins may possess
15 activity in either monomeric or dimeric forms. Human CTLA-8 proteins can also be produced as heterodimers with rat and herpes CTLA-8 proteins and with human IL-17. These heterodimers are also expected to have activities of the proteins of which they are comprised.

Forms of human CTLA-8 protein of less than full length are encompassed
20 within the present invention and may be produced by expressing a corresponding fragment of the polynucleotide encoding the human CTLA-8 protein (SEQ ID NO:1). These corresponding polynucleotide fragments are also part of the present invention. Modified polynucleotides as described above may be made by standard molecular biology techniques, including site-directed mutagenesis methods which are known in
25 the art or by the polymerase chain reaction using appropriate oligonucleotide primers.

For the purposes of the present invention, a protein has "CTLA-8 activity" if it either (1) displays biological activity in a factor-dependent cell proliferation assay (preferably an assay in which full-length the corresponding species full-length CTLA-8 is active) (including without limitation those assays described below), or (2) induces
30 expression or secretion of γ -IFN, or (3) displays chemoattractant or chemotactic activity in a chemoattraction or chemotaxis assay (preferably as assay in which full-

length the corresponding species full-length CTLA-8 is active) or (4) induces expression of secretion of IL-3 or GM-CSF.

Human CTLA-8 protein or fragments thereof having CTLA-8 activity may be fused to carrier molecules such as immunoglobulins. For example, human CTLA-8 protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin.

The invention also encompasses allelic variations of the nucleotide sequence as set forth in SEQ ID NO:1, that is, naturally-occurring alternative forms of the isolated polynucleotide of SEQ ID NO:1 which also encode human CTLA-8 or CTLA-8 proteins having CTLA-8 activity. Also included in the invention are isolated polynucleotides which hybridize to the nucleotide sequence set forth in SEQ ID NO:1 under highly stringent (0.2xSSC at 65°C), stringent (e.g. 4xSSC at 65°C or 50% formamide and 4xSSC at 42°C), or relaxed (4xSSC at 50°C or 30-40% formamide and 4xSSC at 42°C) conditions. Isolated polynucleotides which encode human CTLA-8 protein but which differ from the nucleotide sequence set forth in SEQ ID NO:1 by virtue of the degeneracy of the genetic code are also encompassed by the present invention. Variations in the nucleotide sequence as set forth in SEQ ID NO:1 which are caused by point mutations or by induced modifications which enhance CTLA-8 activity, half-life or production level are also included in the invention.

The isolated polynucleotides of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., Nucleic Acids Res. 19, 4485-4490 (1991), in order to produce the CTLA-8 protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" means enzymatically or chemically ligated to form a covalent bond between the isolated polynucleotide of the invention and the expression control sequence, in such a way that the CTLA-8 protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the human CTLA-8 protein. Any cell type capable of expressing functional human CTLA-8 protein may be used. Suitable mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other
5 transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK, Rat2, BaF3, 32D, FDCEP-1, PC12 or C2C12 cells.

The human CTLA-8 protein may also be produced by operably linking the
10 isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas
15 Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. Soluble forms of the human CTLA-8 protein may also be produced in insect cells using appropriate isolated polynucleotides as described above.

Alternatively, the human CTLA-8 protein may be produced in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Suitable yeast strains
20 include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins.

The human CTLA-8 protein of the invention may also be expressed as a
25 product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a polynucleotide sequence encoding the human CTLA-8 protein.

The human CTLA-8 protein of the invention may be prepared by growing a culture of transformed host cells under culture conditions necessary to express the
30 desired protein. The resulting expressed protein may then be purified from the culture medium or cell extracts. Soluble forms of the human CTLA-8 protein of the invention

can be purified from conditioned media. Membrane-bound forms of human CTLA-8 protein of the invention can be purified by preparing a total membrane fraction from the expressing cell and extracting the membranes with a non-ionic detergent such as Triton X-100.

5 The human CTLA-8 protein can be purified using methods known to those skilled in the art. For example, the human CTLA-8 protein of the invention can be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium.

10 Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or

15 carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-Sepharose® columns). The purification of the human CTLA-8 protein from culture supernatant may also include one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether,

20 or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the human CTLA-8 protein. Some or all of the foregoing purification steps, in various combinations or with other known

25 methods, can also be employed to provide a substantially purified isolated recombinant protein.

 Preferably, the human CTLA-8 protein is purified so that it is substantially free of other mammalian proteins.

 It is believed that human CTLA-8, active fragments and variants thereof, and

30 CTLA-8 related proteins (such as, for example, rat CTLA-8 and herpes CTLA-8) (collectively "CTLA-8 proteins") possess or induce cytokine activities. Human

CTLA-8 expression correlated with γ -IFN expression in induced primary cells and can induce the expression of IL-3 and/or GM-CSF, which expression can in turn produce effects associated with the induced cytokine. Therefore, human CTLA-8 and CTLA-8 related proteins may have an effect on proliferation or function of myeloid cells, erythroid cells, lymphoid cells and their progenitors. Human CTLA-8 proteins may also play a role in formation of platelets or their progenitors.

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon γ , Schreiber, R.D.

In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of
5 Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human
10 interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto.
15 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by
20 measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober
Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors;
25 Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

A protein of the present invention may also exhibit immune stimulating or
30 immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various

immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g.,
5 HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpes viruses, mycobacteria, leishmania, malaria and various fungal infections such as candida. Of course, in this regard, a protein of
10 the present invention may also be useful where a boost to the immune system generally would be indicated, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome,
15 autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, asthma and related respiratory
20 conditions), may also be treatable using a protein of the present invention.

A protein of the present invention may also suppress chronic or acute inflammation, such as, for example, that associated with infection (such as septic shock or systemic inflammatory response syndrome (SIRS)), inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF
25 or IL-1 (such as the effect demonstrated by IL-11).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan,
30 A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte

- Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988;
- 5 Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.
- 10 Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current*
- 15 *Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto, 1994.
- Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan,
- 20 A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.
- 25 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., *Journal of Experimental Medicine* 173:549-559, 1991; Macatonia et al., *Journal of Immunology* 154:5071-5079, 1995; Porgador et al., *Journal of Experimental Medicine*
- 30 182:255-260, 1995; Nair et al., *Journal of Virology* 67:4062-4069, 1993; Huang et al., *Science* 264:961-965, 1994; Macatonia et al., *Journal of Experimental Medicine*

169:1255-1264, 1989; Bhardwaj et al., *Journal of Clinical Investigation* 94:797-807, 1994; and Inaba et al., *Journal of Experimental Medicine* 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in:

5 Darzynkiewicz et al., *Cytometry* 13:795-808, 1992; Gorczyca et al., *Leukemia* 7:659-670, 1993; Gorczyca et al., *Cancer Research* 53:1945-1951, 1993; Itoh et al., *Cell* 66:233-243, 1991; Zacharchuk, *Journal of Immunology* 145:4037-4045, 1990; Zamai et al., *Cytometry* 14:891-897, 1993; Gorczyca et al., *International Journal of*

10 *Oncology* 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., *Blood* 84:111-117, 1994; Fine et al., *Cellular Immunology* 155:111-122, 1994; Galy et al., *Blood* 85:2770-2778, 1995; Toki et al., *Proc. Nat. Acad. Sci. USA* 88:7548-7551,

15 1991.

A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the

20 growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional

25 CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complementarily to platelet transfusions; and/or in supporting the growth

30 and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility

in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e. in conjunction with bone marrow transplantation) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

- 10 Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.
- 15 Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911,
- 20 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp.
- 25 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162,
- 30 Wiley-Liss, Inc., New York, NY. 1994.

CTLA-8 proteins are useful in the treatment of various immune deficiencies and disorders (including SCID), *e.g.*, in regulating (up or down) growth, proliferation and/or activity of T and/or B lymphocytes, as well as the cytolytic activity of NK cells. These immune deficiencies may be caused by viral (*e.g.*, HIV) as well as bacterial infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using CTLA-8 proteins, including infections by HIV, hepatitis, influenza, CMV, herpes, mycobacterium, leishmaniasis, malaria and various fungal infections (such as candida). Of course, in this regard, the CTLA-8 proteins may also be useful where a boost to the immune system generally would be indicated, *i.e.*, in the treatment of cancer or as an adjuvant to vaccines. Autoimmune disorders which may be treated using factors of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes melitis and autoimmune inflammatory eye disease. The CTLA-8 proteins are also expected to be useful in the treatment of allergic reactions and conditions.

CTLA-8 proteins are also expected to have chemotactic activity. A protein or peptide has "chemotactic activity," as used herein, if it can stimulate, directly or indirectly, the directed orientation or movement of cells, including myeloid and lymphoid cells. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells (particularly T-cells). Whether a particular protein or peptide has chemotactic activity for cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

CTLA-8 proteins also inhibit growth and proliferation of vascular endothelial cells. As a result, human CTLA-8 proteins are effective in inhibiting angiogenesis (*i.e.*, vascular formation). This activity will also be useful in the treatment of tumors and other conditions in which angiogenesis is involved. Inhibition of angiogenesis by human CTLA-8 proteins will also result in inhibition or prevention of the condition to which normal angiogenesis would contribute.

Isolated CTLA-8 proteins, purified from cells or recombinantly produced, may be used as a pharmaceutical composition when combined with a pharmaceutically

acceptable carrier. Such a composition may contain, in addition to CTLA-8 protein and carrier, diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-15, G-CSF, γ -IFN, stem cell factor, and erythropoietin. The pharmaceutical composition may contain thrombolytic or anti-thrombotic factors such as plasminogen activator and Factor VIII. The pharmaceutical composition may further contain other anti-inflammatory agents. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with CTLA-8 protein, or to minimize side effects caused by the CTLA-8 protein. Conversely, CTLA-8 protein may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

The pharmaceutical composition of the invention may be in the form of a liposome in which CTLA-8 protein is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, e.g., amelioration of symptoms of,

healing of, or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in
5 combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of CTLA-8 protein is administered to a mammal. CTLA-8 protein may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing
10 cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines, other hematopoietic factors or vaccine components (such as antigens or other adjuvants), CTLA-8 protein may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If
15 administered sequentially, the attending physician will decide on the appropriate sequence of administering CTLA-8 protein in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of CTLA-8 protein used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of
20 conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of CTLA-8 protein is administered orally, CTLA-8 protein will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the
25 invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% CTLA-8 protein, and preferably from about 25 to 90% CTLA-8 protein. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid
30 form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol,

propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of CTLA-8 protein, and preferably from about 1 to 50% CTLA-8 protein.

5 When a therapeutically effective amount of CTLA-8 protein is administered by intravenous, cutaneous or subcutaneous injection, CTLA-8 protein will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in
10 addition to CTLA-8 protein an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additive known to those of skill in the art.

15 The amount of CTLA-8 protein in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of CTLA-8 protein with which to treat each individual patient. Initially, the attending physician will
20 administer low doses of CTLA-8 protein and observe the patient's response. Larger doses of CTLA-8 protein may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1 μ g to about 100 mg of
25 CTLA-8 protein per kg body weight, preferably about 0.1 μ g to about 10 mg of CTLA-8 protein per kg body weight, more preferably about 0.1 μ g to about 100 μ g of CTLA-8 protein per kg body weight, most preferably preferably about 0.1 μ g to about 10 μ g of CTLA-8 protein per kg body weight.

30 The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It

is contemplated that the duration of each application of the CTLA-8 protein will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

5 CTLA-8 protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the CTLA-8 protein and which may inhibit CTLA-8 binding to its receptor. Such antibodies are also useful for performing diagnostics assays for CTLA-8 in accordance with known methods. Such antibodies may be obtained using the entire CTLA-8 protein as an
10 immunogen, or by using fragments of human CTLA-8 protein. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Additional peptide immunogens may be generated by replacing tyrosine residues with sulfated tyrosine residues. Methods for synthesizing such peptides are known in the art, for
15 example, as in R.P. Merrifield, J.Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, et al., FEBS Lett. 211, 10 (1987).

Neutralizing or non-neutralizing antibodies (preferably monoclonal antibodies) binding to human CTLA-8 protein may also be useful therapeutics for certain tumors and also in the treatment of conditions described above. These neutralizing
20 monoclonal antibodies are capable of blocking the ligand binding to the human CTLA-8 protein or mayh promote clearance of protein from the patient.

Because of their homology to human CTLA-8, rat CTLA-8 proteins, herpes CTLA-8 proteins and IL-17 proteins (the "human CTLA-8" of Golstein et al., supra) will also possess CTLA-8 activity as described above. As a result, rat and herpes
25 CTLA-8 proteins and IL-17 proteins, as well as active fragments and variants thereof, can be used in preparation of pharmaceutical compositions and in methods of treatment as described for human CTLA-8. Rat and herpes CTLA-8 proteins, and active fragments and variants thereof, can be produced as described above using the polynucleotides (or fragments or variants thereof) described in SEQ ID NO:3 and SEQ
30 ID NO:5, respectively. Rat and herpes CTLA-8 may also be produced as described in Rouvier et al., J. Immunol. 1993, 150, 5445-5456. CTLA-8 proteins of other

species can also be used as described herein. cDNAs encoding rat CTLA-8 and herpes CTLA-8 were deposited with the American Type Culture Collection on July 6, 1995 and assigned accession numbers ATCC 69867 and ATCC 69866, respectively. IL-17 proteins may also be produced as described in Golstein et al., supra.

5 Because of its homology to IL-17, the human CTLA-8 (B18) proteins of the present invention may also share some activities with IL-17.

For the purposes of treatment or therapy, any of the proteins discussed or disclosed herein may be administered by *in vivo* expression of the protein in a mammalian subject. In such instances, a polynucleotide encoding the desired protein
10 is administered to the subject in manner allowing expression in accordance with known methods, including without limitation the adenovirus methods disclosed herein.

Example 1

Isolation of Human CTLA-8 cDNA

15 A partial clone for human CTLA-8 was isolated from a cDNA library made from RNA isolated from stimulated human peripheral blood mononuclear cells. This partial was identified as "B18." B18 is sometimes used herein to refer to the human CTLA-8 of the present invention. Homology searches identified this partial clone as
20 being related to the herpes and rat CTLA-8 genes. DNA sequence of this partial clone was used to isolate the full-length clone.

In order to isolate a full-length cDNA for B18, a directional, full-length cDNA library by standard means in the COS expression vector pMV2. The cDNA library was transformed into *E. coli* by electroporation. The bulk of the original transformed
25 cDNA library was frozen in glycerol at -80°C. An aliquot was titered to measure the concentration of transformed *E. coli*. The *E. coli* were thawed, diluted to 76,000/0.1 ml in media containing ampicillin, and 0.1 ml was distributed into the wells of a microtiter dish in an 8 x 8 array. The microtiter dish was placed at 37°C overnight to grow the *E. coli*.

30 To prepare DNA for PCR, 20 µl aliquots of culture from each well were withdrawn and pooled separately for each row and column of eight wells, giving 16

pools of 160 μ l each. The *E. coli* were pelleted, resuspended in 160 μ l of standard lysis buffer consisting of 10 mM TrisHCl pH8, 1 mM EDTA, 0.01% Triton X-100, and lysed by heating to 95°C for 10 minutes.

5 To identify which of the wells contained *E. coli* transformed with B18, PCR was performed first on the DNA preps corresponding to the eight columns. The PCR consisted of two sequential reactions with nested oligonucleotides using standard conditions. The oligonucleotides used for the PCR reaction were derived from the sequence of the partial B18 clone. They were:

10 B185: CACAGGCATACACAGGAAGATACATTCA (SEQ ID NO:7)

B183: TCTTGCTGGATGGGAACGGAATTCA (SEQ ID NO:8)

B18N: ATACATTACAGAAGAGCTTCCTGCACA (SEQ ID NO:9)

15 The PCR conditions were 2.5 mM MgCl₂ and 95°C x 2 min for one cycle, 95°C x 1 min plus 68°C x 1 min for 30 cycles, and 68°C x 10 for one cycle. Each reaction was 20 μ l. The first reaction contained oligonucleotides B185 and B183 and 1 μ l of the DNA preparations. The second reaction contained oligonucleotides B183 and B18N and 1 μ l of the first reaction.

20 DNA preps that potentially contained a full-length B18 cDNA clone were identified by agarose gel electrophoresis on an aliquot of the second PCR reaction. A DNA band of the correct mobility was assumed to be derived from a B18 cDNA. Next the same sequence of PCR reactions and gel analysis was done on the DNA preps corresponding to the eight rows. The intersection of a row and a column identified well A2 as potentially containing B18, narrowing it down to the 76,000 *E. coli* originally seeded into that well.

25 To further purify the individual *E. coli* containing the putative full-length B18 cDNA clone, the concentration of *E. coli* in well A2 was measured by titering and plating dilutions of the well. Then 7600 *E. coli* were seeded into the wells of a second microtiter plate in an 8 x 8 array. The *E. coli* were grown overnight; wells were 30 pooled, and DNA was prepared as described above. To identify which of these wells contained *E. coli* transformed with B18, sequential PCR reactions were performed

essentially as described above. Agarose gel electrophoresis identified well B2 as potentially containing a B18 cDNA.

5 The *E. coli* containing this cDNA was further purified by seeding wells of a microtiter plate with 253 *E. coli* per well and proceeding as for the purification of the *E. coli* in well A2. Well C3 was identified as containing a putative full-length B18 cDNA clone. The exact *E. coli* was identified by plating the contents of the well onto bacterial culture media and then screening the *E. coli* colonies following established protocols. The probe for these hybridizations was a PCR fragment generated by doing a PCR reaction on the B18 clone using as primers the oligonucleotides described
10 above (SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9). When a single colony was identified, DNA was prepared and sequenced by standard methods. Comparison of this sequence to the sequence of the original partial clone confirmed identity and that the isolated cDNA was full-length.

15 The full-length clone was deposited with the American Type Culture Collection on July 6, 1995 and assigned accession number ATCC 69868.

Example 2

Expression of Human CTLA-8

20 The full-length B18 clone for human CTLA-8 was transfected into COS cells which were then labelled with ³⁵S-methionine. An aliquot of conditioned medium from the transfected cell culture was reduced, denatured and electrophoresed on polyacrylamide gels. Autoradiographs of those gels are reproduced in Fig. 2. The band indicated by the arrow demonstrates expression of human CTLA-8.

Example 3

Inhibition of Angiogenesis by Human CTLA-8

25 The ability of human CTLA-8 to inhibit angiogenesis was examined in an angiostatic activity assay (endothelial cell proliferation assay). The assay was done in a 96 well plate. Primary human umbilical cells (HUVECs) were seeded to 2×10^3
30 cells per well in EGM medium (Clonetics)/20% FCS and incubated at 37°C for 24 hr. The cells were then starved in M199 medium (GIBCO BRL) containing 10% charcoal

5 treated serum (M199-CS) for 48 hr at 37°C. Conditioned media containing B18 (human CTLA-8) was obtained from transfected COS or stably expressing CHO cells and 1:10, 1:50, 1:250, and 1:1250 dilutions prepared in M199-CS medium containing 100 ng/ml FGF. The dilutions of B18 were added to the starved cells and incubated
10 for 72 hr at 37°C. The cells were then radiolabeled by [³H]-thymidine for 6 hr. Radiolabeled cells were washed with PBS and trypsinized for liquid scintillation counting. Results were plotted using Kaleidograph software. The results are shown in Fig. 3. In the figure, "Med" is the mock control, "B18" and "B18-1" were conditioned medium from two independent transfections of COS with DNA encoding human CTLA-8 (B18). IFN γ was used as a positive control angiostatic (i.e., angiogenesis inhibition) activity. These data demonstrate that human CTLA-8 (B18) inhibits angiogenesis.

Example 4

15 Hematopoietic Activity of Human CTLA-8

The hematopoietic activity of human CTLA-8 (B18) expressed *in vivo* was examined by construction of a recombinant adenovirus vector.

The B18 cDNA in the expression plasmid Adori 2-12 B18 was driven by the cytomegalovirus(CMV) immediate early promoter and enhancer.

20 The Adori 2-12 vector was created by addition of an SV40 origin and enhancer to a known adenovirus vector (Barr et al., Gene Therapy 1:51 (1994); Davidson et al., Nature Genetics 3:219 (1993)). The HindIII/BamHI fragment encoding the SV40 origin and enhancer was isolated from the pMT2 mammalian expression vector, blunted with Klenow and cloned into the NotI site (blunted with Klenow) of the Ad5
25 expression vector.

The vector was derived by digesting pNOT-B18 cDNA with *SalI*, filling in the 5' overhang with Klenow to generate a blunt end and digesting with *EcoRI* to isolate the B18 cDNA. The blunted- *EcoRI* B18 fragment was inserted into the restriction sites *EcoRV-EcoRI* of the adenovirus vector Adori 2-12. The CMV-B18 expression
30 cassette was located downstream of the SV40 origin and enhancer, and 0-1 map units of the left hand end of the adenovirus type 5(Ad5). The SV40 splice donor and

acceptor were located between the CMV promoter and B18 cDNA. Following the insert was SV40 poly A site, 9-16 map units of Ad5 and the puc 19 origin.

5 A recombinant adenovirus was generated by homologous recombination in 293 cells. *AscI* linearized Adori 2-12 B18 and *ClaI* digested AdCMVlacZ were introduced into the 293 cells using lipofectamine. Recombinant adenovirus virus was isolated and amplified on 293 cells. The virus was released from infected 293 cells by three cycles of freeze-thawing. The virus was further purified by two cesium chloride centrifugation gradients and dialyzed against PBS 4°C. Following dialysis of the virus glycerol was added to a concentration of 10 % and the virus was stored at - 70 °C until
10 use. The virus was characterized by expression of the transgene, plaque forming units on 293 cells, particles/ml and Southern analysis of the virus.

A single dose of 5×10^{10} particles of recombinant adenovirus encoding B18 was injected into the tail vein of male C57/bl6 mice, age 7-8 weeks. Control mice received an adenovirus encoding B-galactosidase. Four mice from each experimental
15 group were killed on day 7 and 14. Blood was collected and automated hematologic analysis was performed using a Baker 9000. Differential counts were performed on blood smears. Tissue was harvested, fixed in formalin, and stained with hematoxylin and eosin for histopathology. In the first set of experiments, serum and tissues were analyzed 7 and 14 days post injection. A slight increase in peripheral platelet counts
20 were observed. The animals that received B18 exhibited a slight increase in spleen size. Macroscopic analysis of the spleen showed an increase in splenic extramedullary hematopoiesis on day 7 compared to the control. These results showed a hematopoietic growth activity associated with B18.

In a second set of experiments 5×10^{10} particles of recombinant adenovirus
25 encoding B18 were injected into the tail vein of male C57/bl6 mice, age 17-18 weeks. Control mice received an adenovirus encoding B-galactosidase. Blood samples were collected via retro-orbital sinus on days 2, 5, 7, 10, 14, and 21. The hematologic analyses were performed on the Baker 9000 automated cell counter with murine-specific settings. Analyses included WBC, RBC, HCT and PLT counts. Blood smears
30 were prepared and stained with Wright-Geimsa for WBC differentials based on a 100 cellcount. Reticulocytes and reticulated platelets were quantitated using flow

cytometry. Four mice from each group were killed on days 7, 14, and 21. In addition to peripheral blood analysis, serum was collected via cardiac puncture for quantitation of systemic IL-6 using a commercial kit (Endogen). Spleen and liver were collected for histopathology, spleen and bone marrow hematopoietic progenitors were quantitated, and bone marrow smears were prepared and stained with Wright-Geimsa for cell counts.

Administration of adenovirus encoding B18 resulted in a marked increase in peripheral blood neutrophils and WBC (Fig. 4). Maximum increases in neutrophils were observed at day 5 and day 7. The control mice showed little difference at day 5 and day 7. Peripheral blood neutrophils were similar in the control mice and mice that received B18 at day 21. In both the B18 and control groups an increase in white blood cells was also observed. The mice that received B18 had a greater increase in WBC between day 2 and day 7. By Day 21 a more pronounced increase was observed in the B-gal group. No other changes in cellular chemistries were observed (Table I).

Bone marrow cellularity was calculated from pooled femurs in each group (Table III). No significant differences were observed in either group. No significant changes were observed in bone marrow hematopoietic progenitors from day 7, 14, and 21. The CFU-GM, BFU-E and CFU-MEG in the B18 mice were similar to the B-gal control (Table II).

Administration of the adenovirus encoding B18 resulted in an increase in CFU-GM (myeloid) and BFU-E (erythroid) progenitors in the spleen compared to animals that received the B-gal virus on day 7. The increase in progenitors in the B18 mice was 11-fold in CFU-GM and a 52-fold in BFU-E (Table II). There was a 2-fold increase in CFU-MEG at day 7 for the B18 mice. By day 21 no significant differences were observed in splenic CFU-MEG or BFU-E between the groups (Table II). A 3-fold decrease in CFU-GM was observed in mice that received adenovirus encoding B18. A slight increase in spleen size at day 7 was observed in the B18 group. This is consistent with an increase in splenic cellularity. By day 14 and day 21 spleen weights were similar to the control group (Table III). Macroscopic analysis of the spleen showed an increase in splenic extramedullary hematopoiesis of the B18 mice on day 7 compared to the control.

The bone marrow myeloid: erythroid ratios (Table IV) suggest a granulocytic hyperplasia with a possible erythroid hypoplasia in mice that received adenovirus B18 on day 7. By day 21 the ratio in the B-gal group was higher. No changes were observed in the IL6 serum levels.

- 5 These results show a hematopoietic activity associated with the administration of adenovirus encoding B18 (human CTLA-8). Increases in neutrophils and white blood cells were observed at day 7 in animals that received B18 adenovirus. The data showed that B18 resulted in increase in splenic CFU-GM and BFU-E 7 days post administration compared to the control animals. Splenic extramedullary
- 10 hematopoiesis on day 7 support that B18 exhibits a hematopoietic growth activity. These data suggest that B18 may mobilize early hematopoietic precursors.

Table I: Peripheral hematology for day 2, 5, 7, 10, 14, and 21.

Study AS4-IB - B18 (Platelets) Day 2, 4-25-95														
Group A	WBC x10 ⁹ /μL	Neutro %	ANC x10 ⁹ /μL	Lympho %	ALC x10 ⁹ /μL	Eos %	Monos %	RBC x10 ⁶ /μL	Hct %	Abn Retic x10 ⁶ /μL	HCT %	PLT x10 ⁹ /μL	RPR %	Abn RPR x10 ⁹ /μL
B-Qul #1	5.4	40	2.16	54	2.95	0	5	10.85	3.65	0.46	49.0	830	11.94	99.62
B-Qul #2	7.4	25	1.85	65	4.81	3	7	12.34	2.04	0.25	56.6	900	10.10	90.80
B-Qul #3	6.6	40	2.72	52	3.54	2	5	11.26	3.28	0.37	51.6	894	9.77	87.34
B-Qul #4	8.8	23	2.02	64	5.63	1	12	12.00	2.55	0.31	54.8	840	10.03	89.29
AVG	7.1	32.0	2.19	58.8	4.22	1.5	7.8	11.82	2.88	0.33	52.8	868	10.61	91.84
SEM	0.7	4.6	0.19	3.4	0.61	0.6	1.4	0.33	0.36	0.03	1.3	17	0.45	2.76
B18 #1	11.4	59	6.73	31	3.53	1	9	11.16	4.88	0.54	52.4	1242	14.82	185.31
B18 #2	9.2	30	2.78	62	5.70	3	5	10.14	3.97	0.40	46.8	828	10.90	68.89
B18 #3	5.0	51	2.55	40	2.00	0	9	11.16	3.23	0.38	51.2	832	11.19	63.02
B18 #4	6.4	41	2.62	65	3.52	0	4	10.80	3.09	0.33	49.2	904	12.31	156.48
AVG	8.0	45.3	3.87	47.0	3.69	1.0	6.8	10.82	3.79	0.41	50.2	903	13.56	125.82
SEM	1.4	6.3	1.01	7.0	0.76	0.7	1.3	0.24	0.41	0.05	1.2	127	1.55	27.07
Study AS4-IB - B18 (Platelets) Day 5, 4-28-96														
Group B	WBC x10 ⁹ /μL	Neutro %	ANC x10 ⁹ /μL	Lympho %	ALC x10 ⁹ /μL	Eos %	Monos %	RBC x10 ⁶ /μL	Hct %	Abn Retic x10 ⁶ /μL	HCT %	PLT x10 ⁹ /μL	RPR %	Abn RPR x10 ⁹ /μL
B-Qul #1	7.6	14	1.06	76	5.83	3	5	11.28	3.25	0.39	52.4	1062	15.31	167.82
B-Qul #2	10.6	20	2.12	78	8.27	1	1	10.72	4.71	0.50	48.4	954	17.37	172.66
B-Qul #3	8.6	18	1.51	69	6.07	2	11	11.12	3.40	0.38	51.2	916	9.55	87.40
B-Qul #4	10.6	38	4.10	66	8.26	0	4	10.22	6.21	0.53	47.0	1052	13.83	152.12
AVG	9.6	22.5	2.30	70.8	6.83	1.5	5.3	10.83	4.89	0.53	50.8	1021	14.09	145.82
SEM	0.8	6.3	0.87	4.8	0.55	0.6	2.1	0.23	0.59	0.06	1.2	41	1.67	19.87
B18 #1	14.8	16	2.36	71	10.51	1	10	12.56	2.31	0.29	57.0	1204	7.57	61.14
B18 #2	14.2	37	5.85	53	7.53	2	8	9.80	3.22	0.33	44.6	828	14.30	127.25
B18 #3	12.8	50	3.84	59	7.55	1	10	12.12	4.12	0.50	53.6	1134	10.15	116.10
B18 #4	16.0	58	6.28	37	5.92	0	5	11.04	3.93	0.43	60.8	1166	15.75	163.65
AVG	14.5	35.8	5.28	55.0	7.88	1.0	6.3	11.41	3.42	0.38	52.0	1099	11.85	129.38
SEM	0.7	8.4	1.44	7.1	0.96	0.4	1.2	0.53	0.41	0.05	2.8	71	1.88	19.91
Study AS4-IB - B18 (Platelets) Day 7, 4-28-96														
Group C	WBC x10 ⁹ /μL	Neutro %	ANC x10 ⁹ /μL	Lympho %	ALC x10 ⁹ /μL	Eos %	Monos %	RBC x10 ⁶ /μL	Hct %	Abn Retic x10 ⁶ /μL	HCT %	PLT x10 ⁹ /μL	RPR %	Abn RPR x10 ⁹ /μL
B-Qul #1	15.2	14	2.13	69	10.48	1	16	11.04	3.54	0.39	50.8	1062	12.40	107.41
B-Qul #2	14.0	12	1.68	81	11.84	0	7	11.38	5.05	0.57	52.6	1104	14.81	164.81
B-Qul #3	14.8	14	2.07	73	10.50	1	12	10.92	5.42	0.59	49.8	952	11.49	109.30
B-Qul #4	14.7	13.3	1.86	74.3	10.88	0.7	11.7	11.11	4.87	0.52	51.0	879	12.86	127.43
AVG	14.7	13.3	1.86	74.3	10.88	0.7	11.7	11.11	4.87	0.52	51.0	879	12.86	127.43
SEM	0.4	0.7	0.14	3.5	0.25	0.3	2.6	0.14	0.58	0.06	0.9	71	1.02	18.78
B18 #1	18.4	33	6.40	62	12.63	0	5	10.14	2.93	0.30	45.2	964	12.80	110.58
B18 #2	25.4	39	9.91	53	13.46	0	8	9.46	6.05	0.57	43.6	1288	12.49	160.87
B18 #3	23.6	44	10.38	50	11.80	0	6	9.74	5.17	0.50	44.4	1076	15.41	165.81
B18 #4	12.8	15	1.92	75	9.00	0	19	9.54	6.35	0.50	45.4	1136	15.86	180.40
AVG	20.8	32.8	7.16	60.0	11.72	0.8	7.3	9.78	6.10	0.48	44.2	1091	14.15	154.42
SEM	2.8	6.3	1.96	5.6	0.90	0.2	1.1	0.15	0.76	0.07	0.4	89	0.87	15.19
Study AS4-IB - B18 (Platelets) Day 10, 5-3-96														
Group A	WBC x10 ⁹ /μL	Neutro %	ANC x10 ⁹ /μL	Lympho %	ALC x10 ⁹ /μL	Eos %	Monos %	RBC x10 ⁶ /μL	Hct %	Abn Retic x10 ⁶ /μL	HCT %	PLT x10 ⁹ /μL	RPR %	Abn RPR x10 ⁹ /μL
B-Qul #1	18.8	17	3.18	69	12.63	3	11	10.22	12.41	1.27	46.8	1460	16.30	236.52
B-Qul #2	13.2	18	2.11	79	10.43	1	4	10.48	6.00	0.63	48.8	1128	14.48	163.33
B-Qul #3	18.8	18	3.14	74	14.80	0	10	10.72	6.25	0.87	49.4	1338	16.58	221.84
B-Qul #4	18.6	21	3.91	72	13.39	3	4	10.44	7.59	0.76	46.4	1036	14.35	153.28
AVG	17.5	17.5	3.08	73.5	12.79	1.8	7.3	10.47	8.06	0.84	46.4	1249	15.40	193.74
SEM	1.5	1.2	0.27	2.1	0.86	0.8	1.9	0.10	1.49	0.15	0.6	81	0.68	20.78
B18 #1	14.2	33	4.69	56	7.95	5	6	8.70	11.87	1.04	39.2	1760	14.49	255.02
B18 #2	17.8	35	6.18	57	10.03	1	7	9.04	9.48	0.88	42.0	1104	18.88	208.44
B18 #3	16.2	39	6.32	57	9.23	1	3	4.74	18.77	0.79	22.4	894	28.19	380.96
B18 #4	14.2	28	3.55	66	9.27	1	8	9.30	8.93	0.82	42.0	1415	16.81	238.03
AVG	15.6	33.0	6.18	69.0	9.15	2.8	6.0	7.95	12.04	0.90	36.4	1294	19.84	240.81
SEM	0.8	2.3	0.66	2.3	0.63	1.0	1.1	1.88	1.67	0.05	4.7	183	3.24	11.77
Study AS4-IB - B18 (Platelets) Day 14, 5-7-96														
Group B	WBC x10 ⁹ /μL	Neutro %	ANC x10 ⁹ /μL	Lympho %	ALC x10 ⁹ /μL	Eos %	Monos %	RBC x10 ⁶ /μL	Hct %	Abn Retic x10 ⁶ /μL	HCT %	PLT x10 ⁹ /μL	RPR %	Abn RPR x10 ⁹ /μL
B-Qul #1	17.8	18	3.20	74	13.17	0	8	10.50	5.87	0.85	50.8	1360	11.03	150.01
B-Qul #2	20.4	28	5.90	68	13.46	1	7	10.82	7.07	0.77	50.8	1616	8.18	132.19
B-Qul #3	18.0	7	1.12	90	14.40	1	3	11.28	8.41	0.73	52.8	1298	7.50	85.53
B-Qul #4	18.0	36	6.40	57	10.38	1	8	9.30	7.22	0.71	43.0	1672	10.05	158.04
AVG	18.1	21.8	4.03	71.3	12.82	0.8	6.0	10.61	6.77	0.71	49.4	1487	8.18	138.44
SEM	0.9	6.1	1.18	7.0	0.89	0.3	1.1	0.45	0.36	0.03	2.1	83	0.84	15.48
B18 #1	15.4	9	1.29	81	12.47	1	9	10.52	6.74	0.81	48.2	1282	9.51	120.02
B18 #2	15.4	31	4.77	58	8.63	2	9	9.76	10.33	1.01	44.8	1092	14.29	152.05
B18 #3	13.4	42	5.63	39	5.23	0	19	10.34	4.89	0.52	48.8	1376	15.79	217.27
B18 #4	11.5	57	6.61	34	3.94	2	7	9.38	8.57	0.52	43.0	1092	16.56	181.83
AVG	14.9	34.5	4.59	53.0	7.54	1.3	11.0	10.03	8.58	0.66	45.6	1206	14.86	158.82
SEM	0.9	10.1	1.14	10.7	1.83	0.5	2.7	0.29	1.23	0.12	1.1	79	1.69	20.54
Study AS4-IB - B18 (Platelets) Day 21, 5-14-96														
Group A	WBC x10 ⁹ /μL	Neutro %	ANC x10 ⁹ /μL	Lympho %	ALC x10 ⁹ /μL	Eos %	Monos %	RBC x10 ⁶ /μL	Hct %	Abn Retic x10 ⁶ /μL	HCT %	PLT x10 ⁹ /μL	RPR %	Abn RPR x10 ⁹ /μL
B-Qul #1	23.4	23	5.34	67	17.02	0	10	8.22	8.15	0.75	42.8	1776	9.61	170.87
B-Qul #2	19.8	19	3.72	68	13.82	0	12	8.50	9.85	0.95	44.4	1682	9.44	158.89
B-Qul #3	27.8	11	3.04	82	22.53	3	4	9.74	8.84	0.86	45.8	1894	11.45	182.82
B-Qul #4	28.0	13	3.64	63	23.24	0	4	9.04	7.54	0.69	41.8	1346	10.40	141.06
AVG	25.2	16.5	4.06	73.3	18.10	0.8	7.5	8.38	8.62	0.81	43.7	1617	10.25	168.36
SEM	1.9	2.8	0.61	4.2	2.33	0.5	2.1	0.15	0.52	0.06	0.3	94	0.46	10.97
B18 #1	18.8	9	1.67	83	15.44	1	7	9.54	7.40	0.73	43.8	1642	8.37	137.44
B18 #2	16.2	48	7.78	45	7.29	1	5	11.10	7.87	0.88	51.4	1870	10.78	212.37
B18 #3	14.8	53	7.84	42	6.22	0	5	7.52	20.25	1.82	38.0	1568	10.50	164.64
B18 #4	15.2	18	2.74	74	11.25	2	6	9.64	7.02	0.68	43.0	1422	7.60	108.07
AVG	16.2	32.0	5.01	61.0	10.65	1.0	6.0	9.53	10.66	0.85	44.1	1651	9.31	155.63
SEM	0.9	10.9	1.63	10.3	2.10	0.4	0.4	0.74	3.21	0.23	2.8	116	0.78	22.18

Table II: Bone marrow and Splenic Hematopoietic Progenitors

	CFU-MEG		CFU-GM		BFU-E	
	B-Gal	B18	B-Gal	B18	B-Gal	B18
Bone Marrow*						
Day 7	16.0 ± 3.5	15.7 ± 3.1	307 ± 117	241 ± 78	51 ± 19	25 ± 11
Day 14	10.7 ± 2.3	15.3 ± 1.2	233 ± 15	373 ± 35	30 ± 10	60 ± 30
Day 21	5.7 ± 0.6	6.7 ± 3.1	170 ± 17	160 ± 27	40 ± 10	27 ± 6
Spleen**						
Day 7	9.3 ± 1.6	19.5 ± 1.5	27 ± 3	298 ± 6	1.3 ± 1.2	68 ± 10
Day 14	9.7 ± 0.6	12.7 ± 0.6	267 ± 32	197 ± 21	33 ± 6	10 ± 10
Day 21	17.0 ± 1.0	19.3 ± 2.5	187 ± 6	73 ± 15	23 ± 6	23 ± 6

Hematopoietic precursors were determined from pooled spleen and bone marrow samples from four animals in each group. For quantitation of CFU-GM and BFU-E, either 1×10^4 bone marrow cells or 1×10^5 spleen cells were added to complete alpha methylcellulose medium (0.9% methylcellulose in alpha medium, 30% fetal bovine serum, 1% bovine serum albumin, 10-4M 2-mercaptoethanol, 2 mM L-glutamine, 2% murine spleen cell conditioned medium, and 3 U/mL erythropoietin) and aliquoted into 35 mm tissue culture dishes in a final volume of 1.0 mL. Cultures were incubated for 7 days at 37°C, and 5% CO₂. Microscopic colonies were defined as clusters of 50 or more cells. For quantitation of CFU-MEG, either 1×10^5 bone marrow cells or 1×10^6 spleen cells were added to complete alpha methylcellulose medium and incubated as described above. Megakaryocyte colonies were defined as a group of 3 or more cells.

*Bone marrow progenitors are represented as mean ± sd number of colonies per 10^5 cells.

**Spleen progenitors are represented as mean ± sd number of colonies per 10^6 cells.

Table III: Spleen Weights and Femur Cellularity

	Spleen Wt. (Mg)	B-Gal	B18		Femur Cellularity ($\times 10^6$)	B-Gal	B18
	Day 7	187 $\pm$ 19	224 $\pm$ 29		Day 7	28	23
	Day 14	175 $\pm$ 13	170 $\pm$ 10		Day 14	28	27
5	Day 21	174 $\pm$ 21	151 $\pm$ 27		Day 21	28	26

Spleen weights were determined at time of sacrifice are represented as means $\pm$ sd from four animals.

10 Table IV: Bone Marrow Myeloid:Erythroid Ratios

	Group	Mouse #	Day 7	Day 14	Day 21
	B-gal	1	1.43	2.12	5.78
		2	0.91	2.46	5.83
15		3	1.62	1.03	3.66
		4		5.44	4.82
	AVG		1.32	2.76	5.02
	SD		0.37	0.37	1.89
	B-gal	1	5.59	2.01	2.02
20		2	6.51	1.25	2.13
		3	5.49	1.58	1.81
		4	0.50	2.51	2.92
	AVG		4.52	1.86	2.22
	SD-		1.29	2.72	0.56
25					

All entries represent the number of myeloid cells per 1 erythroid cell. Normal mouse ratios are approximately 1:1 to 2:1.

Example 5

30 Additional Experiments Relating to Hematopoietic Activity of Human CTLA-8

B18 (human CTLA-8) was tested for the ability to induce production of factors having hematopoietic activity in a factor-dependent cell proliferation assay using the

human erythroleukemic cell line, TF-1 (Kitamura et al., J. Cell Physiol. 140:323 (1989)). The cells were initially grown in the presence of rhGMCSF (100 U/ml). The cells were fed three days prior to setting up the assay. The assay conditions were as follows:

5	cells/well	5000/200 μ l
	incubation time	3 days
	pulse time	4 hours
	amount of tritiated thymidine	0.5 μ Ci/well
	counting time	1 minute
10	replicates	2

B18 alone, conditioned medium (CM) from B18 induced HS-5 cells were assayed. Buffer alone, CM from HS-5 cells induced with buffer and CM from uninduced HS-5 cells were assayed as controls. Results are shown in Fig. 5. B18 (human CTLA-8) demonstrated an ability to induce production of factors which
 15 induced TF-1 proliferation. This activity was substantially eliminated by the addition of anti-GMCSF antibodies. These data demonstrate that human CTLA-8 (B18) is able to induce hematopoiesis. Particularly, without being bound by any theory, it appears that human CTLA-8 (B18) induces production of GM-CSF and/or IL-3.

20 Example 6

Ability of Human CTLA-8 to Induce Production of IL-6 and IL-8

MRC5 cells were incubated in the presence of human CTLA-8 (B18) and production of IL-6 and IL-8 were measured. Herpes CTLA-8 (IL-17) was used as a positive control. Applicants' human CTLA-8 (B18) demonstrated titratable
 25 production of both IL-6 and IL-8 (see Figs. 6 and 7).

All patent and literature references cited herein are incorporated by reference as if fully set forth.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Jacobs, Kenneth
Kelleher, Kerry
Carlin, McKeough
Goldman, Samuel
Pittman, Debra
Mi, Sha
Neben, Steven
Giannotti, JoAnn
Golden Fleet, Margaret
- (ii) TITLE OF INVENTION: Human CTLA-8 and Uses of CTLA-8-Related Proteins
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Genetics Institute, Inc.
 - (B) STREET: 87 CambridgePark Drive
 - (C) CITY: Cambridge
 - (D) STATE: Massachusetts
 - (E) COUNTRY: USA
 - (F) ZIP: 02140
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Brown, Scott A.
 - (B) REGISTRATION NUMBER: 32,724
 - (C) REFERENCE/DOCKET NUMBER: GI5262
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (617) 498-8224
 - (B) TELEFAX: (617) 876-5851

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 813 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 56..544

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

GGGAAGATAC ATTACAGAA AGAGCTTCCT GCACAAAGTA AGCCACCAGC GCAAC ATG      58
                                         Met
                                         1
ACA GTG AAG ACC CTG CAT GGC CCA GCC ATG GTC AAG TAC TTG CTG CTG      106
Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu Leu Leu
      5                               10                15
TCG ATA TTG GGG CTT GCC TTT CTG AGT GAG GCG GCA GCT CGG AAA ATC      154
Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Arg Lys Ile
      20                25                30
CCC AAA GTA GGA CAT ACT TTT TTC CAA AAG CCT GAG AGT TGC CCG CCT      202
Pro Lys Val Gly His Thr Phe Phe Gln Lys Pro Glu Ser Cys Pro Pro
      35                40                45
GTG CCA GGA GGT AGT ATG AAG CTT GAC ATT GGC ATC ATC AAT GAA AAC      250
Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu Asn
      50                55                60                65
CAG CGC GTT TCC ATG TCA CGT AAC ATC GAG AGC CGC TCC ACC TCC CCC      298
Gln Arg Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser Pro
      70                75                80
TGG AAT TAC ACT GTC ACT TGG GAC CCC AAC CGG TAC CCC TCG GAA GTT      346
Trp Asn Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu Val
      85                90                95
GTA CAG GCC CAG TGT AGG AAC TTG GGC TGC ATC AAT GCT CAA GGA AAG      394
Val Gln Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly Lys
      100                105                110
GAA GAC ATC TCC ATG AAT TCC GTT CCC ATC CAG CAA GAG ACC CTG GTC      442
Glu Asp Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu Val
      115                120                125
GTC CGG AGG AAG CAC CAA GGC TGC TCT GTT TCT TTC CAG TTG GAG AAG      490
Val Arg Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu Lys
      130                135                140                145
GTG CTG GTG ACT GTT GGC TGC ACC TGC GTC ACC CCT GTC ATC CAC CAT      538
Val Leu Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His His
      150                155                160
GTG CAG TAAGAGGTGC ATATCCACTC AGCTGAAGAA GCTGTAGAAA TGCCACTCCT      594
Val Gln

TACCCAGTGC TCTGCAACAA GTCCTGTCTG ACCCCCAATT CCTTCCACTT CACAGGACTC      654
TTAATAAGAC CTGCACGGAT GGAAACAGAA AATATTCACA ATGTATGTGT GTATGTACTA      714
CACTTTATAT TTGATATCTA AAATGTTAGG AGAAAAATTA ATATATTCAG TGCTAATATA      774
ATAAAGTATT AATAATTTAA AAATAAAAAA AAAAAAAA      813

```

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 163 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu Leu
 1           5           10           15
Leu Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Ala Arg Lys
          20           25           30
Ile Pro Lys Val Gly His Thr Phe Phe Gln Lys Pro Glu Ser Cys Pro
          35           40           45
Pro Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu
          50           55           60
Asn Gln Arg Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser
          65           70           75           80
Pro Trp Asn Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu
          85           90           95
Val Val Gln Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly
          100          105          110
Lys Glu Asp Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu
          115          120          125
Val Val Arg Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu
          130          135          140
Lys Val Leu Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His
          145          150          155          160
His Val Gln

```

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 461 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 6..455

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

CCACC ATG TGC CTG ATG CTG TTG CTG CTA CTG AAC CTG GAG GCT ACA      47
      Met Cys Leu Met Leu Leu Leu Leu Asn Leu Glu Ala Thr
        1           5           10
GTG AAG GCA GCG GTA CTC ATC CCT CAA AGT TCA GTG TGT CCA AAC GCC      95
Val Lys Ala Ala Val Leu Ile Pro Gln Ser Ser Val Cys Pro Asn Ala
  15           20           25           30
GAG GCC AAT AAC TTT CTC CAG AAC GTG AAG GTC AAC CTG AAA GTC ATC     143
Glu Ala Asn Asn Phe Leu Gln Asn Val Lys Val Asn Leu Lys Val Ile
          35           40           45

```

AAC TCC CTT AGC TCA AAA GCG AGC TCG AGA AGG CCC TCA GAT TAC CTC Asn Ser Leu Ser Ser Lys Ala Ser Ser Arg Arg Pro Ser Asp Tyr Leu 50 55 60	191
AAC CGT TCC ACT TCA CCC TGG ACT CTG AGC CGC AAT GAG GAC CCT GAT Asn Arg Ser Thr Ser Pro Trp Thr Leu Ser Arg Asn Glu Asp Pro Asp 65 70 75	239
AGA TAT CCT TCT GTG ATC TGG GAG GCA CAG TGC CGC CAC CAG CGC TGT Arg Tyr Pro Ser Val Ile Trp Glu Ala Gln Cys Arg His Gln Arg Cys 80 85 90	287
GTC AAC GCT GAG GGG AAG TTG GAC CAC CAC ATG AAT TCT GTT CTC ATC Val Asn Ala Glu Gly Lys Leu Asp His His Met Asn Ser Val Leu Ile 95 100 105 110	335
CAG CAA GAG ATA CTA GTC CTG AAG AGG GAG CCT GAG AAG TGC CCC TTC Gln Gln Glu Ile Leu Val Leu Lys Arg Glu Pro Glu Lys Cys Pro Phe 115 120 125	383
ACT TTC CGG GTG GAG AAG ATG CTG GTG GGC GTG GGC TGC ACC TGC GTT Thr Phe Arg Val Glu Lys Met Leu Val Gly Val Gly Cys Thr Cys Val 130 135 140	431
TCC TCT ATT GTC CGC CAT GCG TCC TAATAA Ser Ser Ile Val Arg His Ala Ser 145 150	461

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 150 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Cys Leu Met Leu Leu Leu Leu Asn Leu Glu Ala Thr Val Lys 1 5 10 15
Ala Ala Val Leu Ile Pro Gln Ser Ser Val Cys Pro Asn Ala Glu Ala 20 25 30
Asn Asn Phe Leu Gln Asn Val Lys Val Asn Leu Lys Val Ile Asn Ser 35 40 45
Leu Ser Ser Lys Ala Ser Ser Arg Arg Pro Ser Asp Tyr Leu Asn Arg 50 55 60
Ser Thr Ser Pro Trp Thr Leu Ser Arg Asn Glu Asp Pro Asp Arg Tyr 65 70 75 80
Pro Ser Val Ile Trp Glu Ala Gln Cys Arg His Gln Arg Cys Val Asn 85 90 95
Ala Glu Gly Lys Leu Asp His His Met Asn Ser Val Leu Ile Gln Gln 100 105 110
Glu Ile Leu Val Leu Lys Arg Glu Pro Glu Lys Cys Pro Phe Thr Phe 115 120 125
Arg Val Glu Lys Met Leu Val Gly Val Gly Cys Thr Cys Val Ser Ser 130 135 140

Ile Val Arg His Ala Ser
145 150

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 459 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..453

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG ACA TTT AGA ATG ACT TCA CTT GTG TTA CTT CTG CTG CTG AGC ATA	48
Met Thr Phe Arg Met Thr Ser Leu Val Leu Leu Leu Leu Ser Ile	
1 5 10 15	
GAT TGT ATA GTA AAG TCA GAA ATA ACT AGT GCA CAA ACC CCA AGA TGC	96
Asp Cys Ile Val Lys Ser Glu Ile Thr Ser Ala Gln Thr Pro Arg Cys	
20 25 30	
TTA GCT GCT AAC AAT AGC TTT CCA CGG TCT GTG ATG GTT ACT TTG AGC	144
Leu Ala Ala Asn Asn Ser Phe Pro Arg Ser Val Met Val Thr Leu Ser	
35 40 45	
ATC CGT AAC TGG AAT ACC AGT TCT AAA AGG GCT TCA GAC TAC TAC AAT	192
Ile Arg Asn Trp Asn Thr Ser Ser Lys Arg Ala Ser Asp Tyr Tyr Asn	
50 55 60	
AGA TCT ACG TCT CCT TGG ACT CTC CAT CGC AAT GAA GAT CAA GAT AGA	240
Arg Ser Thr Ser Pro Trp Thr Leu His Arg Asn Glu Asp Gln Asp Arg	
65 70 75 80	
TAT CCC TCT GTG ATT TGG GAA GCA AAG TGT CGC TAC TTA GGA TGT GTT	288
Tyr Pro Ser Val Ile Trp Glu Ala Lys Cys Arg Tyr Leu Gly Cys Val	
85 90 95	
AAT GCT GAT GGG AAT GTA GAC TAC CAC ATG AAC TCA GTC CCT ATC CAA	336
Asn Ala Asp Gly Asn Val Asp Tyr His Met Asn Ser Val Pro Ile Gln	
100 105 110	
CAA GAG ATT CTA GTG GTG CGC AAA GGG CAT CAA CCC TGC CCT AAT TCA	384
Gln Glu Ile Leu Val Val Arg Lys Gly His Gln Pro Cys Pro Asn Ser	
115 120 125	
TTT AGG CTA GAG AAG ATG CTA GTG ACT GTA GGC TGC ACA TGC GTT ACT	432
Phe Arg Leu Glu Lys Met Leu Val Thr Val Gly Cys Thr Cys Val Thr	
130 135 140	
CCC ATT GTT CAC AAT GTA GAC TAAAAG	459
Pro Ile Val His Asn Val Asp	
145 150	

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 151 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Thr Phe Arg Met Thr Ser Leu Val Leu Leu Leu Leu Ser Ile
 1           5           10           15
Asp Cys Ile Val Lys Ser Glu Ile Thr Ser Ala Gln Thr Pro Arg Cys
          20           25           30
Leu Ala Ala Asn Asn Ser Phe Pro Arg Ser Val Met Val Thr Leu Ser
          35           40           45
Ile Arg Asn Trp Asn Thr Ser Ser Lys Arg Ala Ser Asp Tyr Tyr Asn
          50           55           60
Arg Ser Thr Ser Pro Trp Thr Leu His Arg Asn Glu Asp Gln Asp Arg
          65           70           75           80
Tyr Pro Ser Val Ile Trp Glu Ala Lys Cys Arg Tyr Leu Gly Cys Val
          85           90           95
Asn Ala Asp Gly Asn Val Asp Tyr His Met Asn Ser Val Pro Ile Gln
          100          105          110
Gln Glu Ile Leu Val Val Arg Lys Gly His Gln Pro Cys Pro Asn Ser
          115          120          125
Phe Arg Leu Glu Lys Met Leu Val Thr Val Gly Cys Thr Cys Val Thr
          130          135          140
Pro Ile Val His Asn Val Asp
          145          150
  
```

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CACAGGCATA CACAGGAAGA TACATTCA

28

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCTTGCTGGA TGGGAACGGA ATTCA

25

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATACATTCAC AGAAGAGCTT CCTGCACA

28

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - 5 (a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544;
 - (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 85 to nucleotide 544;
 - (c) the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to
 - 10 nucleotide 544;
 - (d) the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;
 - (e) the nucleotide sequence of the full length protein coding sequence of clone B18 deposited under accession number ATCC 69868;
 - 15 (f) a nucleotide encoding the full length protein encoded by the cDNA insert of clone B18 deposited under accession number ATCC 69868;
 - (g) a nucleotide sequence that specifically hybridizes to a nucleic acid sequence specified in (a) to (f);
 - (h) a nucleotide sequence varying from the sequence of the nucleotide
 - 20 sequence specified in (a) to (f) as a result of degeneracy of the genetic code; and
 - (i) an allelic variant of the nucleotide specified in (a) to (f).
2. The polynucleotide of claim 1 wherein said nucleotide sequence encodes a protein having CTLA-8 activity.
- 25 3. The polynucleotide of claim 1 or claim 2 wherein said nucleotide sequence is operably linked to an expression control sequence.
4. The polynucleotide according to any one of claims 1 to 3 comprising the
- 30 nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544.
5. A host cell transformed with the polynucleotide according to any one of claims 1 to 4.
- 35 6. The host cell of claim 5, wherein said cell is a mammalian cell.



7. A process for producing a human CTLA-8 protein, said process comprising:

(a) growing a culture of the host cell of claim 5 or claim 6 in a suitable culture medium; and

5 (b) purifying the human CTLA-8 protein from the culture.

8. An isolated human CTLA-8 protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:2;

10 (b) the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;

(c) the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;

15 (d) the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163;

(e) a fragment of SEQ ID NO:2, the fragment comprising amino acids 11 to 163 of SEQ ID NO:2;

(f) a fragment of SEQ ID NO:2, the fragment comprising amino acids 29 to 163 of SEQ ID NO:2;

20 (g) a fragment of SEQ ID NO:2, the fragment comprising amino acids 31 to 163 of SEQ ID NO:2; and

(h) fragments of (a), (b), (c), (d), (e), (f) or (g) having a CTLA-8 activity selected from the group consisting of: biological activity in a factor-dependent cell proliferation assay; induction of expression or secretion of γ -IFN; chemoattractant or chemotactic activity in a chemoattraction or chemotaxis assay; and induction or expression of secretion of IL-3 or GM-CSF.

25

9. The protein of claim 8 comprising the amino acid sequence of SEQ ID NO:2.

30

10. The protein of claim 8 comprising the sequence from amino acid 29 to 163 of SEQ ID NO:2.

11. A pharmaceutical composition comprising a human CTLA-8 protein according to any one of claims 8 to 10 and a pharmaceutically acceptable carrier.

35



12. A human CTLA-8 protein produced according to a process which comprises the steps of:

(a) growing a culture of a cell in a suitable culture medium, wherein the cell is transformed with a polynucleotide operably linked to an expression control sequence, wherein the polynucleotide is selected from the group consisting of:

(aa) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544;

(ab) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;

(ac) a polynucleotide comprising the nucleotide sequence of the full-length coding sequence of clone B18 deposited under accession number ATCC 69868;

(ad) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone B18 deposited under accession number ATCC 69868; and

(ae) a polynucleotide comprising a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (aa) to (ad) as a result of degeneracy of the genetic code; and

(b) purifying the human CTLA-8 protein encoded by said polynucleotide from the culture.

13. A composition comprising an antibody which specifically reacts with a human CTLA-8 protein according to any one of claims 8 to 10.

14. A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition of claim 11.

15. The polynucleotide according to any one of claims 1 to 4 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 86 to nucleotide 544.

16. The protein of claim 8 comprising the sequence from amino acid 11 to 163 of SEQ ID NO:2.

17. The method of claim 14, wherein said subject is treated to produce an effect selected from the group consisting of inhibition of angiogenesis, inhibition



of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, proliferation of myeloid cells or progenitors, proliferation of erythroid cells or progenitors, proliferation of lymphoid cells or progenitors, induction of IFN γ production,
 5 induction of IL-3 production and induction of GM-CSF production:

18. The polynucleotide of claim 3 wherein said polynucleotide is contained in a vector suitable for *in vivo* expression in a mammalian subject.
- 10 19. The polynucleotide according to any one of claims 1 to 4 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544.
20. The protein of claim 8 comprising the sequence from amino acid 31 to 163 of SEQ ID NO:2.
- 15 21. The protein of claim 8 comprising a fragment of SEQ ID NO:2, the fragment comprising amino acids 11 to 163 of SEQ ID NO:2.
22. The protein of claim 8 comprising a fragment of SEQ ID NO:2, the
 20 fragment comprising amino acids 29 to 163 of SEQ ID NO:2.
23. The protein of claim 8 comprising a fragment of SEQ ID NO:2, the fragment comprising amino acids 31 to 163 of SEQ ID NO:2.
- 25 24. The human CTLA-8 protein of claim 12, wherein said polynucleotide comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544.
25. The human CTLA-8 protein of claim 12, wherein said polynucleotide
 30 comprises the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544.
26. The human CTLA-8 protein of claim 12, wherein said polynucleotide comprises the nucleotide sequence of the full-length coding sequence of clone
 35 B18 deposited under accession number ATCC 69868.

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000



27. The human CTLA-8 protein of claim 12, wherein said polynucleotide comprises a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone B18 deposited under accession number ATCC 69868.
- 5 28. The polynucleotide of claim 1 comprising the nucleotide sequence of the full-length protein coding sequence of clone B18 deposited under accession number ATCC 69868.
- 10 29. The polynucleotide of claim 1 comprising the nucleotide encoding the full length protein encoded by the cDNA insert of clone B18 deposited under accession number ATCC 69868.

Dated this second day of May 2000

GENETICS INSTITUTE, INC.

Patent Attorneys for the Applicant:

F B RICE & CO


5
6
8
8
4



Hsv1e-2	-----[L]N[S]E[T]E[A]Q-T[P]R[C]L-[A]A[N]S[P]P[R]S[V]M[V]T[L]S[I]R[N]-----[Q]N[T]S[S]R[A]S[D]YV[N]R[S]T[S]P	51
Musct1a8	-----[A]V[L]I[P]Q-S[V]C[P]N[A]E[A]M[N]E[L]Q[N]V[K]U[N]L[K]V[N]S[L]S[K]A[S]S[R]P[S]D[Y]L[N]R[S]T[S]P	51
B18_F1	A[A]R[K]E[F]K[V]G[H]T[P]F[F]Q[K]R[P]E[S]C[P]F[V]P[G]G-----[S]M[K]L[D]I[G]T[I]N[E]N[Q]R[V]S[M]S[R]-----N[I]E[S]R[S]T[S]P	53
Hsv1e-2	[M]T[L]H[E]N[E]D[Q]D[R]Y[P]S[V]I[E]A[K]C[R]V[L]G[C]V[N]A[D]G[N]C[D]V[H]M[N]S[V]P[I]Q[Q]E[I]L[V]V[R]K[G]H[Q]P[C]P[M]S[F]	111
Musct1a8	[M]T[L]S[E]N[E]D[F]D[R]Y[P]S[V]I[E]A[K]C[R]H[Q]E[C]V[N]A[E]G[K]L[D]H[H]M[N]S[V]L[I]Q[Q]E[I]L[V]L[K]R[E]P[E]K[C]P[F]T[F]	111
B18_F1	[M]N[Y]T[V]T[W]D[F]N[R]Y[P]S[E]V[V]Q[A]C[R]N[L]G[C]I[N]A[Q]K[E]D[I]S[M]N[S]V[P]I[Q]Q[E]T[L]V[V]R[R]K[H]Q[G]C[S]V[S]F	113
Hsv1e-2	[R]L[E]K[M]L[V]T[V]G[C]T[C]V[T]P[I]V[H]N[V]D	133
Musct1a8	[R]V[E]K[M]L[V]G[V]G[C]T[C]V[S]S[I]V[R]H[A]S	133
B18_F1	[Q]L[E]K[V]L[V]T[V]G[C]T[C]V[T]P[V]I[H]V[Q]	135

Fig. 1

Figure 2



B18 M B18 M
R R NR NR

Supernatant Cell Lysate

27




Fig. 3

3/7

Fig. 4A




Fig. 4B




Fig. 4C





Fig. 4D






Fig. 5

5/7

SUBSTITUTE SHEET (RULE 26)




Fig. 6

