



US009755163B2

(12) **United States Patent**
Li et al.

(10) **Patent No.:** **US 9,755,163 B2**
(45) **Date of Patent:** **Sep. 5, 2017**

(54) **SYNTHESIS OF FOUR COORDINATED PALLADIUM COMPLEXES AND THEIR APPLICATIONS IN LIGHT EMITTING DEVICES THEREOF**

(71) Applicant: **Arizona Board of Regents Acting for and on Behalf of Arizona State University**, Scottsdale, AZ (US)

(72) Inventors: **Jian Li**, Phoenix, AZ (US); **Eric Turner**, Phoenix, AZ (US)

(73) Assignee: **ARIZONA BOARD OF REGENTS ACTING FOR OR ON BEHALF OF ARIZONA STATE UNIVERSITY**, Scottsdale, AZ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/202,058**

(22) Filed: **Jul. 5, 2016**

(65) **Prior Publication Data**
US 2017/0005278 A1 Jan. 5, 2017

Related U.S. Application Data

(63) Continuation of application No. 14/145,461, filed on Dec. 31, 2013, now Pat. No. 9,382,273, which is a (Continued)

(51) **Int. Cl.**
H05B 33/14 (2006.01)
H01L 51/42 (2006.01)
C09K 11/06 (2006.01)
C07F 7/24 (2006.01)
H01L 51/00 (2006.01)
C07D 213/643 (2006.01)
(Continued)

(52) **U.S. Cl.**
CPC **H01L 51/0084** (2013.01); **C07D 213/643** (2013.01); **C07F 7/24** (2013.01); **C07F 15/006** (2013.01); **C09K 11/06** (2013.01); **H05B 33/14** (2013.01); **C09K 2211/1007** (2013.01); **C09K 2211/1014** (2013.01); **C09K 2211/1029** (2013.01); **C09K 2211/1044** (2013.01); **C09K 2211/185** (2013.01); **H01L 51/424** (2013.01); **H01L 51/5016** (2013.01); **Y02E 10/549** (2013.01)

(58) **Field of Classification Search**
CPC H01L 51/0084; C07D 213/643
USPC 546/4
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,769,292 A 9/1988 Tang et al.
5,707,745 A 1/1998 Forrest et al.
(Continued)

FOREIGN PATENT DOCUMENTS

CN 1777663 5/2006
CN 1894269 1/2007
(Continued)


OTHER PUBLICATIONS

Wong; Journal of Organometallic Chemistry 2009, 694, 2644-2647.*
(Continued)

Primary Examiner — Daniel Carcanague
(74) *Attorney, Agent, or Firm* — Fish & Richardson P.C.

(57) **ABSTRACT**
Synthesis of four coordinated palladium complexes and their applications in light emitting devices thereof.

23 Claims, 6 Drawing Sheets



Related U.S. Application Data

continuation of application No. 13/695,337, filed as application No. PCT/US2011/034776 on May 2, 2011, now abandoned.

(60) Provisional application No. 61/329,684, filed on Apr. 30, 2010.

(51) **Int. Cl.**

H01L 51/50 (2006.01)
C07F 15/00 (2006.01)

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,844,363	A	12/1998	Gu et al.
6,200,695	B1	3/2001	Arai et al.
6,303,238	B1	10/2001	Thompson et al.
6,780,528	B2	8/2004	Tsuboyama et al.
7,002,013	B1	2/2006	Chi et al.
7,037,599	B2	5/2006	Culligan et al.
7,279,704	B2	10/2007	Walters et al.
7,332,232	B2	2/2008	Ma et al.
7,442,797	B2	10/2008	Itoh et al.
7,501,190	B2	3/2009	Ise
7,655,322	B2	2/2010	Forrest et al.
7,947,383	B2	5/2011	Ise et al.
8,389,725	B2	3/2013	Li et al.
8,617,723	B2	12/2013	Stoessel
8,816,080	B2	8/2014	Li et al.
8,871,361	B2	10/2014	Xia et al.
8,927,713	B2	1/2015	Li et al.
8,946,417	B2	2/2015	Li et al.
9,059,412	B2	6/2015	Zeng et al.
9,224,963	B2	12/2015	Li et al.
9,238,668	B2	1/2016	Li et al.
9,312,505	B2	4/2016	Brooks et al.
9,324,957	B2	4/2016	Li et al.
9,382,273	B2	7/2016	Li
9,385,329	B2	7/2016	Li et al.
9,425,415	B2	8/2016	Li et al.
9,461,254	B2	10/2016	Tsai et al.
9,550,801	B2	1/2017	Li et al.
9,617,291	B2	4/2017	Li et al.
2002/0068190	A1	6/2002	Tsuboyama et al.
2003/0062519	A1	4/2003	Yamazaki et al.
2003/0186077	A1	10/2003	Chen
2005/0170207	A1	8/2005	Ma et al.
2005/0260446	A1	11/2005	Mackenzie et al.
2006/0073359	A1	4/2006	Ise et al.
2006/0094875	A1	5/2006	Itoh et al.
2006/0202197	A1	9/2006	Nakayama et al.
2006/0210831	A1	9/2006	Sano et al.
2006/0263635	A1	11/2006	Ise
2006/0286406	A1	12/2006	Igarashi et al.
2007/0057630	A1	3/2007	Nishita et al.
2007/0059551	A1	3/2007	Yamazaki
2007/0082284	A1	4/2007	Stoessel et al.
2007/0103060	A1	5/2007	Itoh et al.
2008/0001530	A1	1/2008	Ise et al.
2008/0036373	A1	2/2008	Itoh et al.
2008/0054799	A1	3/2008	Satou
2008/0079358	A1	4/2008	Satou
2008/0241518	A1	10/2008	Satou et al.
2008/0241589	A1	10/2008	Fukunaga et al.
2009/0026936	A1	1/2009	Satou et al.
2009/0026939	A1	1/2009	Kinoshita et al.
2009/0032989	A1	2/2009	Karim et al.
2009/0039768	A1	2/2009	Igarashi et al.
2009/0079340	A1	3/2009	Kinoshita et al.
2009/0128008	A1	5/2009	Ise et al.
2009/0218561	A1	9/2009	Kitamura et al.
2009/0261721	A1	10/2009	Murakami et al.
2009/0267500	A1	10/2009	Kinoshita et al.
2010/0000606	A1	1/2010	Thompson et al.
2010/0013386	A1	1/2010	Thompson et al.

2010/0171111	A1	7/2010	Takada et al.
2012/0095232	A1	4/2012	Li et al.
2012/0181528	A1	7/2012	Takada et al.
2012/0215001	A1	8/2012	Li et al.
2012/0223634	A1	9/2012	Xia et al.
2012/0302753	A1	11/2012	Li
2013/0048963	A1	2/2013	Beers et al.
2013/0168656	A1	7/2013	Tsai et al.
2013/0203996	A1	8/2013	Li et al.
2013/0237706	A1	9/2013	Li
2013/0341600	A1	12/2013	Lin et al.
2014/0014922	A1	1/2014	Lin et al.
2014/0027733	A1	1/2014	Zeng et al.
2014/0084261	A1	3/2014	Brooks et al.
2014/0114072	A1	4/2014	Li et al.
2014/0203248	A1	7/2014	Zhou et al.
2014/0330019	A1	11/2014	Li et al.
2014/0364605	A1	12/2014	Li et al.
2015/0008419	A1	1/2015	Li
2015/0028323	A1	1/2015	Xia et al.
2015/0069334	A1	3/2015	Xia et al.
2015/0105556	A1	4/2015	Li et al.
2015/0162552	A1	6/2015	Li et al.
2015/0194616	A1	7/2015	Li et al.
2015/0228914	A1	8/2015	Li et al.
2015/0287938	A1	10/2015	Li et al.
2015/0318500	A1	11/2015	Li
2015/0349279	A1	12/2015	Li et al.
2016/0028028	A1	1/2016	Li et al.
2016/0043331	A1	2/2016	Li et al.
2016/0072082	A1	3/2016	Brooks et al.
2016/0133862	A1	5/2016	Li et al.
2016/0197291	A1	7/2016	Li
2016/0285015	A1	9/2016	Li et al.
2016/0359120	A1	12/2016	Li
2016/0359125	A1	12/2016	Li et al.
2017/0012224	A1	1/2017	Li et al.
2017/0047533	A1	2/2017	Li et al.
2017/0066792	A1	3/2017	Li et al.
2017/0069855	A1	3/2017	Li

FOREIGN PATENT DOCUMENTS

CN	101142223	3/2008
CN	101667626	3/2010
CN	102449108 A	5/2012
CN	102892860	1/2013
CN	102971396	3/2013
CN	104232076	12/2014
CN	104693243	6/2015
CN	105367605	3/2016
CN	105418591	3/2016
EP	1808052	7/2007
EP	1874893	1/2008
EP	1874894	1/2008
EP	1919928	5/2008
EP	2036907	3/2009
EP	2096690 A2	9/2009
EP	2417217	2/2012
EP	2112213	7/2012
EP	2711999	3/2014
JP	2005267557	9/2005
JP	200531073	11/2005
JP	2006047240	2/2006
JP	2006232784	9/2006
JP	2006242080	9/2006
JP	2006242081	9/2006
JP	2006256999	9/2006
JP	2006257238	9/2006
JP	2006261623	9/2006
JP	2006290988	10/2006
JP	2006313796	11/2006
JP	2006332622	12/2006
JP	2006351638	12/2006
JP	2007019462	1/2007
JP	2007042875	2/2007
JP	2007053132	3/2007
JP	2007066581	3/2007
JP	2007073620	3/2007

(56)

References Cited

FOREIGN PATENT DOCUMENTS

JP	2007073845	3/2007
JP	2007073900	3/2007
JP	2007080593	3/2007
JP	2007080677	3/2007
JP	2007088105	4/2007
JP	2007088164	4/2007
JP	2007096259	4/2007
JP	2007110067	4/2007
JP	2007110102	4/2007
JP	2007258550	10/2007
JP	2007324309	12/2007
JP	2008010353	1/2008
JP	2008091860	4/2008
JP	2008103535	5/2008
JP	2008108617	5/2008
JP	2008109085	5/2008
JP	2008109103	5/2008
JP	2008160087	7/2008
JP	2008198801	8/2008
JP	2008270729	11/2008
JP	2008270736	11/2008
JP	2009016184	1/2009
JP	2009016579	1/2009
JP	2009032977	2/2009
JP	2009032988	2/2009
JP	2009266943	11/2009
JP	2009267171	11/2009
JP	2009267244	11/2009
JP	2009272339	11/2009
JP	2009283891	12/2009
JP	2010135689	6/2010
JP	2012222255	11/2012
JP	2013525436	6/2013
JP	5604505	10/2014
JP	2014221807	11/2014
JP	2015081257	4/2015
KR	1020060115371	11/2006
KR	2007061830	6/2007
KR	2007112465	11/2007
KR	1020130043460	4/2013
TW	200701835	1/2007
TW	201307365	2/2013
TW	201710277	3/2017
WO	WO0070655	11/2000
WO	WO2004003108 A1	1/2004
WO	WO2004108857	12/2004
WO	WO2005042444	5/2005
WO	WO2005042550	5/2005
WO	WO2006033440	3/2006
WO	WO2006098505	9/2006
WO	WO2006115299	11/2006
WO	WO2006115301	11/2006
WO	WO2007034985	3/2007
WO	WO2007069498	6/2007
WO	WO2008066192	6/2008
WO	WO2008066195	6/2008
WO	WO2008066196	6/2008
WO	WO2008117889	10/2008
WO	WO2008123540	10/2008
WO	WO2009017211	2/2009
WO	WO2010118026	10/2010
WO	WO2011137429	11/2011
WO	WO2011137431	11/2011
WO	WO2011137439	11/2011
WO	WO2012112853	8/2012
WO	WO2012142387	10/2012
WO	WO2012162488	11/2012
WO	WO2012163471	12/2012
WO	WO2013130483	9/2013
WO	WO2014016611 A1	1/2014
WO	WO2014031977	2/2014
WO	WO2014047616	3/2014
WO	WO2014109814	7/2014
WO	WO2015027060	2/2015

WO	WO2015131158	9/2015
WO	WO2016025921	2/2016
WO	WO2016029186	2/2016

OTHER PUBLICATIONS

Marc Lepeltier et al., "Efficient blue green organic light-emitting devices based on a monofluorinated heteroleptic iridium(III) complex," *Synthetic Metals*, vol. 199, 2015, pp. 139-146.

Stefan Bernhard, "The First Six Years: A Report," Department of Chemistry, Princeton University, May 2008, 11 pages.

Non-Final Rejection dated Jul. 1, 2013 for U.S. Appl. No. 13/695,337, filed May 2, 2011 (Applicants—Arizona Technology Enterprises (AZTE); Inventors—Li et al.; (28 pages).

Preliminary Amendment filed on Oct. 30, 2012 for U.S. Appl. No. 13/695,337, filed May 2, 2011 (Applicants—Arizona Technology Enterprises (AZTE); Inventors—Li et al.; (4 pages).

International Preliminary Report on Patentability dated Nov. 6, 2012 for Intl. Pat. App. No. PCT/US2011/034776 filed May 2, 2011 and published as WO 2011/137429 dated Nov. 3, 2011 (Applicants—Arizona Board of Regents Acting for and on behalf of Arizona State University; Inventors—Li et al.; (6 pages).

International Search Report dated Feb. 9, 2012 for Intl. Pat. App. No. PCT/US2011/034776 filed May 2, 2011 and published as WO 2011/137429 dated Nov. 3, 2011 (Applicants—Arizona Board of Regents Acting for and on behalf of Arizona State University; Inventors—Li et al.; (3 pages).

Written Opinion dated Feb. 9, 2012 for Intl. Pat. App. No. PCT/US2011/034776 filed May 2, 2011 and published as WO 2011/137429 dated Nov. 3, 2011 (Applicants—Arizona Board of Regents Acting for and on behalf of Arizona State University; Inventors—Li et al.; (5 pages).

First Office Action (and English Translation) for Chinese Application No. 201180023966.1, dated Feb. 7, 2014, 14 pages.

Second Office Action (and English Translation) for Chinese Application No. 201180023966.1, dated Dec. 22, 2014, 8 pages.

Official Action (and English Translation) issued by the Japanese Patent Office dated Mar. 11, 2015 for Pat. App. No. 2013-508082 filed May 2, 2011, 16 pages.

Third Office Action (and English Translation) for Chinese Application No. 201180023966.1, dated Sep. 1, 2015, 7 pages.

Final Rejection (and English Translation) issued by the Japanese Patent Office dated Oct. 30, 2015 for Pat. App. No. 2013-508082, 6 pages.

Wong; Challenges in organometallic research—Great opportunity for solar cells and OLEDs, *Journal of Organometallic Chemistry*, 2009, 694, 2644-2647.

JP2009267244, English Translation from EPO, dated Nov. 2009, 80 pages.

JP2010135689, English translation from EPO, dated Jun. 2010, 95 pages.

Chi et al.; Transition-metal phosphors with cyclometalating ligands: fundamentals and applications, *Chemical Society Reviews*, vol. 39, No. 2, Feb. 2010, pp. 638-655.

Dorwald; Side Reactions in Organic Synthesis: A Guide to Successful Synthesis Design, 2005, Wiley-VCH.

Satake et al., "Interconvertible Cationic and Neutral Pyridinylimidazole η³-Allylpalladium Complexes. Structural Assignment by 1H, 13C, and 15N NMR and X-ray Diffraction", *Organometallics*, vol. 18, No. 24, 1999, pp. 5108-5111.

Baldo et al., "Highly Efficient Phosphorescent Emission from Organic Electroluminescent Devices," *Nature*, vol. 395, Sep. 10, 1998, pp. 151-154.

Baldo et al., "Very high-efficiency green organic light-emitting devices based on electrophosphorescence," *Applied Physics Letters*, vol. 75, No. 1, Jul. 5, 1999, pp. 4-6.

Ying Yang et al., "Induction of Circularly Polarized Electroluminescence from an Achiral Light-Emitting Polymer via a Chiral Small-Molecule Dopant," *Advanced Materials*, vol. 25, Issue 18, May 14, 2013, pp. 2624-2628.

(56)

References Cited

OTHER PUBLICATIONS

Ayan Maity et al., "Room-temperature synthesis of cyclometalated iridium(III) complexes; kinetic isomers and reactive functionalities" *Chem. Sci.*, vol. 4, pp. 1175-1181 (2013).

Shiro Koseki et al., "Spin-orbit coupling analyses of the geometrical effects on phosphorescence in Ir(ppy)₃ and its derivatives", *J. Phys. Chem. C*, vol. 117, pp. 5314-5327 (2013).

Ji Hyun Seo et al., "Efficient blue-green organic light-emitting diodes based on heteroleptic tris-cyclometalated iridium (III) complexes". *Thin Solid Films*, vol. 517, pp. 1807-1810 (2009).

Barry O'Brien et al.: White organic light emitting diodes using Pt-based red, green and blue phosphorescent dopants. *Proc. SPIE*, vol. 8829, pp. 1-6, Aug. 25, 2013.

Vanessa Wood et al., "Colloidal quantum dot light-emitting devices," *Nano Reviews*, vol. 1, 2010, 8 pages.

Glauco Ponterini et al., "Comparison of Radiationless Decay Processes in Osmium and Platinum Porphyrins," *J. Am. Chem. Soc.*, vol. 105, No. 14, 1983, pp. 4639-4645.

Shizuo Tokito et al., "Confinement of triplet energy on phosphorescent molecules for highly-efficient organic blue-light-emitting devices," *Applied Physics Letters*, vol. 83, No. 3, Jul. 21, 2003, pp. 569-571.

Brian W. D'Andrade et al., "Controlling Exciton Diffusion in Multilayer White Phosphorescent Organic Light Emitting Devices," *Adv. Mater.*, vol. 14, No. 2, Jan. 16, 2002, pp. 147-151.

Dileep A. K. Vezzu et al., "Highly Luminescent Tetradentate Bis-Cyclometalated Platinum Complexes: Design, Synthesis, Structure, Photophysics, and Electroluminescence Application," *Inorg. Chem.*, vol. 49, 2010, pp. 5107-5119.

Evan L. Williams et al., "Excimer-Based White Phosphorescent Organic Light Emitting Diodes with Nearly 100% Internal Quantum Efficiency," *Adv. Mater.*, vol. 19, 2007, pp. 197-202.

Shih-Chun Lo et al., "High-Triplet-Energy Dendrons: Enhancing the Luminescence of Deep Blue Phosphorescent Iridium(III) Complexes," *J. Am. Chem. Soc.*, vol. 131, 2009, pp. 16681-16688.

Jan Kalinowski et al., "Light-emitting devices based on organometallic platinum complexes as emitters," *Coordination Chemistry Reviews*, vol. 255, 2011, pp. 2401-2425.

Ke Feng et al., "Norbornene-Based Copolymers Containing Platinum Complexes and Bis(carbazolyl)benzene Groups in Their Side-Chains," *Macromolecules*, vol. 42, 2009, pp. 6855-6864.

Chi-Ming Che et al., "Photophysical Properties and OLED Applications of Phosphorescent Platinum(II) Schiff Base Complexes," *Chem. Eur. J.*, vol. 16, 2010, pp. 233-247.

Stephen R. Forrest, "The path to ubiquitous and low-cost organic electronic appliances on plastic," *Nature*, vol. 428, Apr. 29, 2004, pp. 911-918.

Nicholas R. Evans et al., "Triplet Energy Back Transfer in Conjugated Polymers with Pendant Phosphorescent Iridium Complexes," *J. Am. Chem. Soc.*, vol. 128, 2006, pp. 6647-6656.

Xiaofan Ren et al., "Ultrahigh Energy Gap Hosts in Deep Blue Organic Electrophosphorescent Devices," *Chem. Mater.*, vol. 16, 2004, pp. 4743-4747.

Dorwald; "Side Reactions in Organic Synthesis: A Guide to Successful Synthesis Design," Chapter 1, 2005 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 32 pages.

Murakami; JP 2007258550, English machine translation from EPO, dated Oct. 4, 2007, 80 pages.

Murakami; JP 2007324309, English machine translation from EPO, dated Dec. 13, 2007, 89 pages.

* cited by examiner




FIG. 1




FIG. 2






FIG. 3




(A)




(B)

FIG. 4




(A)




(B)

FIG. 5



(C)

FIG. 5



(D)

FIG. 5

FIG. 2 is a photoluminescence spectrum produced from a specific embodiment, 2-(3-(pyridin-2-yl)phenoxy)-6-(pyridin-2-yloxy)phenyl]Pd(II) (Pd003) taken in degassed dichloromethane at room temperature, wherein the quantum yield was about 0.1 and the luminescence lifetime was about 11 μ sec. The chemical structure of Pd003 is also illustrated in the inset.

FIG. 3 is a photoluminescence spectrum produced from a specific embodiment, 2-(3-(pyridin-2-yl)phenoxy)-6-(pyridin-2-yloxy)phenyl]Pd(II) (Pd003) doped into a polymethylmethacrylate (PMMA) thin film at room temperature, which shows a thin film photoluminescence quantum yield over 70%.

FIG. 4 illustrates (A) a specific embodiment, and (B) a photoluminescence spectrum of the same in a solution of dichloromethane, obtained at room temperature wherein the quantum yield was about 0.16 and the luminescent lifetime was about 30 μ sec.

FIG. 5 illustrates (A) a specific embodiment, (B) an electroluminescent spectrum from a OLED device comprising the specific embodiment, (C) a plot of external quantum efficiency vs. current density for a Pd003 based device, and (D) a current vs. voltage plot at a 1% doping level.

Additional aspects of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or can be learned by practice of the invention. The advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DESCRIPTION

The present invention can be understood more readily by reference to the following detailed description of the invention and the Examples included therein.

Before the present compounds, devices, and/or methods are disclosed and described, it is to be understood that they are not limited to specific synthetic methods unless otherwise specified, or to particular reagents unless otherwise specified, as such can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular aspects only and is not intended to be limiting. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, example methods and materials are now described.

As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a component" includes mixtures of two or more components.

Ranges can be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another aspect includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another aspect. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It is also understood that there are a number of values disclosed herein, and that each value is also herein disclosed as "about" that particular

value in addition to the value itself. For example, if the value "10" is disclosed, then "about 10" is also disclosed. It is also understood that each unit between two particular units are also disclosed. For example, if 10 and 15 are disclosed, then 11, 12, 13, and 14 are also disclosed.

As used herein, the terms "optional" or "optionally" means that the subsequently described event or circumstance can or can not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

The term "alkyl" as used herein is a branched or unbranched saturated hydrocarbon group of 1 to 24 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, s-butyl, t-butyl, n-pentyl, isopentyl, s-pentyl, neopentyl, hexyl, heptyl, octyl, nonyl, decyl, dodecyl, tetradecyl, hexadecyl, eicosyl, tetracosyl, and the like. The alkyl group can be cyclic or acyclic. The alkyl group can be branched or unbranched. The alkyl group can also be substituted or unsubstituted. For example, the alkyl group can be substituted with one or more groups including, but not limited to, optionally substituted alkyl, cycloalkyl, alkoxy, amino, ether, halide, hydroxy, nitro, silyl, sulfo-oxo, or thiol, as described herein. A "lower alkyl" group is an alkyl group containing from one to six (e.g., from one to four) carbon atoms.

The terms "amine" or "amino" as used herein are represented by the formula $NA^1A^2A^3$, where A^1 , A^2 , and A^3 can be, independently, hydrogen or optionally substituted alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, aryl, or heteroaryl group as described herein.

The term "halide" as used herein refers to the halogens fluorine, chlorine, bromine, and iodine.

The term "hydroxyl" as used herein is represented by the formula $-OH$.

The term "nitro" as used herein is represented by the formula $-NO_2$.

The term "nitrile" as used herein is represented by the formula $-CN$.

The term "thiol" as used herein is represented by the formula $-SH$.

Disclosed are the components to be used to prepare the compositions of the invention as well as the compositions themselves to be used within the methods disclosed herein. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each various individual and collective combinations and permutation of these compounds can not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a particular compound is disclosed and discussed and a number of modifications that can be made to a number of molecules including the compounds are discussed, specifically contemplated is each and every combination and permutation of the compound and the modifications that are possible unless specifically indicated to the contrary. Thus, if a class of molecules A, B, and C are disclosed as well as a class of molecules D, E, and F and an example of a combination molecule, A-D is disclosed, then even if each is not individually recited each is individually and collectively contemplated meaning combinations, A-E, A-F, B-D, B-E, B-F, C-D, C-E, and C-F are considered disclosed. Likewise, any subset or combination of these is also disclosed. Thus, for example, the sub-group of A-E, B-F, and C-E would be considered disclosed. This concept applies to all aspects of this application including, but not limited to, steps in methods of making and using the compositions of the invention. Thus, if there are a variety of


5

additional steps that can be performed it is understood that each of these additional steps can be performed with any specific embodiment or combination of embodiments of the methods of the invention.

Throughout the specification, it should be understood that where letters and/or symbols are utilized to represent atoms or functional groups, and wherein multiple instances of the same letter and/or symbol are present, that each individual instance can represent the same or different species (e.g., atom and/or functional group) than any other instance using the same letter and/or symbol. Similarly, when a metal is depicted as a portion of a chemical structure, the notation can refer to a single metal atom and/or to a plurality of atoms. In one aspect, a notation for a metal refers to a single metal atom. In another aspect, a notation for a metal refers to a plurality of metal atoms.


In one aspect, the invention comprises phosphorescent multidentate, for example, tridentate and/or tetradentate, palladium (II) complexes. In another aspect, such multidentate palladium complexes can be modified and/or specifically tailored to tune the emission spectra from, for example, ultraviolet to near-infrared emission. In yet another aspect, the inventive compositions can provide improved stability and efficiency over conventional light emitting materials. In yet other aspects, the inventive compositions can be useful as luminescent labels, absorbers, emitters, or a combination thereof.

In one aspect, the inventive compositions are represented by the general formula:




wherein Ar1, Ar2, Ar3, and Ar4, if present, represent aromatic groups, wherein each E represents an optional linking atom, such as, for example, carbon or oxygen, and wherein R, if present, represents an ancillary ligand

In various aspects, an ancillary ligand can comprise one or more of the following:




6

-continued




7

-continued



8

-continued

U = O, S, N—R³

10

15

20

25

30

35

40

45

50

55

60


65

wherein, R¹-R¹⁰ of the ancillary ligand each independently represent a hydrogen atom, an alkyl group, a haloalkyl group, an aralkyl group, an alkenyl group, an alkynyl group, an aryl group, an amino group, a mono- or di-alkylamino group, a mono- or diaryl amino group, an alkoxy group, an aryloxy group, a heteroaryloxy group, an alkoxy-carbonyl group, an acyloxy group, an acylamino group, an alkoxy-carbonylamino group, an aryloxy-carbonylamino group, a sulfonylamino group, a sulfamoyl group, a carbamoyl group, an alkylthio group, a sulfinyl group, a ureido group, a phosphoramidate group, a hydroxyl group, a mercapto group, a halogen atom, a cyano group, a sulfo group, a carboxyl group, a nitro group, a hydrazino group, a substituted silyl group, or a polymerizable group; further, wherein a plurality of Rs exist, the number of R should be from 0 to about 4, and each R can be the same or different from any other R. In one aspect, an ancillary ligand can comprise a hydrogen atom. In another aspect, an ancillary ligand can comprise an alkyl group. In another aspect, an ancillary ligand can comprise a haloalkyl group. In another aspect, an ancillary ligand can comprise an aralkyl group. In another aspect, an ancillary ligand can comprise an alkenyl group. In another aspect, an ancillary ligand can comprise an alkynyl group. In another aspect, an ancillary ligand can comprise an aryl group. In another aspect, an ancillary ligand can comprise an amino group. In another aspect, an ancillary ligand can comprise an alkoxy group, an aryloxy group, a heteroaryloxy group, an alkoxy-carbonyl group, an acyloxy group, or a combination thereof. In other aspects, an ancillary ligand can comprise an acylamino group, an alkoxy-carbonylamino group, an aryloxy-carbonylamino group, a sulfonylamino group, a sulfamoyl group, a carbamoyl group, an alkylthio group, a sulfinyl group, a ureido group, a phosphoramidate group, a hydroxyl group, a mercapto group, a halogen atom, a cyano group, a sulfo group, a carboxyl group, a nitro group, a hydrazino group, a substituted silyl group, a polymerizable group, or a combination thereof.


In still other aspects, an ancillary ligand can comprise a group or groups different from those specifically recited herein, and the present invention is not intended to be limited to any particular ancillary ligand.

In various aspects, specific non-limiting examples of the inventive composition can be grouped and illustrated by ligand class. In one aspect, the inventive composition can be represented by the general formula:

9




wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, X represents a halogen or other electronegative group, and C represents an aromatic group. Specific examples of inventive compositions within this ligand class can comprise:




or a combination thereof.

In one aspect, the inventive composition can be represented by the general formula:




wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, X represents a halogen or other electronegative group, and C represents an aromatic group. Specific examples of inventive composition within this ligand class can comprise:

10




or a combination thereof.

In one aspect, the inventive composition can be represented by the general formula:




wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, and X represents a halogen or other electronegative group. Specific examples of inventive composition within this ligand class can comprise:




or a combination thereof.

11

In one aspect, the inventive composition can be represented by the general formula:




wherein Pd represents palladium, N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, X represents a halogen or other electronegative group, and each C represents an aromatic group. Specific examples of inventive composition within this ligand class can comprise:




or a combination thereof.

In one aspect, the inventive composition can be represented by the general formula:




wherein Pd represents palladium, each E represents an optional linking atom, such as, for example, carbon or oxygen, R represents an ancillary ligand, and each C represents an aromatic group. Specific examples of inventive composition within this ligand class can comprise:

12




or a combination thereof.

In one aspect, the inventive composition can be represented by the general formula:



45


wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, and each C represents an aromatic group. Specific examples of inventive composition within this ligand class can comprise:




or a combination thereof.

13

In one aspect, the inventive composition can be represented by the general formula:




wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, and each C represents an aromatic group. Specific examples of inventive composition within this ligand class can comprise:



or a combination thereof.

In one aspect, the inventive composition can be represented by the general formula:



wherein Pd represents palladium, N represents a nitrogen substituted aromatic group, each E represents an optional

14

linking atom, such as, for example, carbon or oxygen, and each C represents an aromatic group. Specific examples of inventive composition within this ligand class can comprise:


5

10

15

20

25



or a combination thereof.


In one aspect, the inventive composition can be represented by the general formula:

30

35

40

45




50

wherein Pd represents palladium, N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, and each C represents an aromatic group. A specific example of an inventive composition within this ligand class can comprise:

55


60

65




In one aspect, the inventive composition can be represented by the general formula:

15




wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or oxygen, and C represents an aromatic group. Specific examples of inventive compositions within this ligand class can comprise:



or a combination thereof.


In one aspect, the inventive composition can be represented by the general formula:




wherein Pd represents palladium, each N represents a nitrogen substituted aromatic group, each E represents an optional linking atom, such as, for example, carbon or

16

oxygen, and C represents an aromatic group. A specific example of an inventive composition within this ligand class can comprise:




In another aspect, the compounds of the present invention are represented by the formula:



In the above formula, each R^1 and R^2 in $(R^1)_2$ and $(R^2)_2$ independently represents hydrogen, optionally substituted C_1 - C_4 alkyl, halogen, hydroxyl, amino, nitro, or thiol; Y^{1a} represents O, S, NR^{4a} , wherein R^{4a} represents optionally substituted C_1 - C_4 alkyl; $Si(R^{4b})_2$, wherein each R^{4b} in $(R^{4b})_2$ independently represents optionally substituted C_1 - C_4 alkyl; or $C(R^{4c})_2$, wherein each R^{4c} in $(R^{4c})_2$ represents hydrogen or optionally substituted C_1 - C_4 alkyl; n is an integer 0 or 1; Y^{1b} , when present, represents O, S, NR^{5a} , wherein R^{5a} represents optionally substituted C_1 - C_4 alkyl; $Si(R^{5b})_2$, wherein each R^{5b} in $(R^{5b})_2$ independently represents optionally substituted C_1 - C_4 alkyl; or $C(R^{5c})_2$, wherein each R^{5c} in $(R^{5c})_2$ represents hydrogen or optionally substituted C_1 - C_4 alkyl; each of Y^{2a} , Y^{2b} , Y^{2c} , and Y^{2d} independently represents N, NR^{6a} , or CR^{6b} , wherein each of R^{6a} and R^{6b} independently represents hydrogen, optionally substituted C_1 - C_4 alkyl, halogen, hydroxyl, amino, nitro, or thiol; each of Y^{3a} , Y^{3b} , Y^{3c} , Y^{3d} , Y^{3e} , Y^{4a} , Y^{4b} , Y^{4c} , and Y^{4d} independently represents N, O, S, NR^{6a} , CR^{6b} , wherein each of R^{6a} and R^{6b} independently represents hydrogen or optionally substituted C_1 - C_4 alkyl; or $Z(R^{6c})_2$, wherein Z is C or Si, and wherein each R^{6c} in $(R^{6c})_2$ independently represents hydrogen or optionally substituted C_1 - C_4 alkyl; m is an integer 1 or 2; wherein the open dotted circle () indicates partial or full unsaturation of the ring with which it is associated.


In one embodiment of the formula above, if m is 1, each of Y^{2a} and Y^{2d} is CH and each of Y^{2b} and Y^{2c} is N, then at least one of Y^{4a} , Y^{4b} , Y^{3a} , or Y^{3d} is not N. For example, according to this embodiment, the following compound is not included in the above formula:

17




As can be seen in the preceding example above, m is 1, each of Y^{2a} and Y^{2d} is CH and each of Y^{2b} and Y^{2c} is N; however, each of Y^{4a} , Y^{4b} , Y^{3a} , or Y^{3d} is N. It follows that the preceding example, according to this embodiment, is not included within the general formula above. In the practice of this embodiment, similar analysis can be used to determine whether or not a compound is or is not included within the general formula above.

In a further embodiment of the general formula above, if n is 0, m is 2, each of Y^{2a} and Y^{2d} is CH, and each of Y^{2b} and Y^{2c} is N, then at least one of Y^{3b} or Y^{3c} is not N. For example, according to this embodiment, the following compound is not included in the above formula:




As can be seen in the preceding example above, n is 0, m is 2, each of Y^{2a} and Y^{2d} is CH, and each of Y^{2b} and Y^{2c} is N. However, each of Y^{3b} and Y^{3c} is N. It follows that the preceding example, according to this embodiment, is not included within the general formula above. Once more, in the practice of this embodiment, similar analysis can be used to determine whether or not a compound is or is not included within the general formula above.

In one embodiment of the general formula above, the compound is represented by the formula:




Non-limiting examples of specific embodiments within this formula include:


18



In another embodiment of the general formula above, the compound is represented by the formula:




Non-limiting examples of specific embodiments within this formula include:




19

-continued




20

-continued




21


-continued



In another embodiment of the general formula above, the compound is represented by the formula:




Non-limiting examples of specific embodiments within this formula include:




22

-continued



In another embodiment of the general formula above, the compound is represented by the formula:




Non-limiting examples of specific embodiments within this formula include:

23


24

-continued




25

-continued




26

-continued




In another embodiment of the general formula above, the compound is represented by the formula:

40
45
50
55




Non-limiting examples of specific embodiments within this formula include:




27

-continued




28




-continued




In other aspects, any one or more of the general formulas and/or specific examples recited herein can be excluded from the invention. For example, in one aspect, the formula



In another embodiment of the general formula above, the compound is represented by the formula:




Non-limiting examples of specific embodiments within these formula include:



is not included in the present invention.

The compounds of the invention can be made using a variety of methods. In one embodiment, wherein Y^{1a} is O, the compounds can be provided according to Scheme 1.

Scheme 1.





50

55

60

65






With reference to Scheme 1, step "a" can be accomplished, for example, by using a catalytic amount of a coupling reagent, such as Cu_2O , which couples alcohols, particularly phenols, with halogenated phenyl groups. The variable "X" in Scheme 1 above represents a halogen (i.e., Cl, F, I, Br), and is preferably I when used in conjunction with Scheme 1.

Each side of a ligand which complexes a metal can be made independently using a variety of methods, which, in one aspect, depend on whether Y^{3a} is N or C. With reference to Scheme 2 below, when Y^{4a} is N, the precursor can, in one aspect, be provided according to Scheme 2(A), wherein a halogenated phenyl compound is reacted with a pyrazole, imadazole, 1H-1,2,3-triazole, 1H-tetrazole, or 2H-pentazole. In such an aspect, the halogenated phenyl compound can comprise any halogen (X), including Cl, Br, F, or I, but is preferably I, which is typically more reactive in a coupling reaction. The halogenated phenyl compound and corresponding pyrazole, imadazole, 1H-1,2,3-triazole, 1H-tetrazole, or 2H-pentazole can be coupled using a metallic and/or organometallic coupling agent, such as, for example, Cu_2O . During such a coupling reaction, it can, in one aspect, be advantageous to include an acid scavenger, such as, for example, syn-2-pyridinealdoxime, in a small molar ratio, for example, about 20 mol %.

In another aspect, when Y^{4a} is C, a different protocol can be used to provide the precursor. With reference to Scheme 2(B) below, a halogenated phenyl, as discussed above can be reacted with a tetrazole, 1,2,3-triazole, pyrazole, or pyrrole to achieve a carbon-carbon bond coupling, as opposed to a carbon-nitrogen bond coupling as shown in Scheme 2(A). In one aspect, the carbon-carbon bond coupling can also be achieved using an organometallic catalyst, such as, for example, a Pd(II) catalyst (e.g., $\text{Pd}(\text{OAc})_2$) in a small molar ratio. In one aspect, such an organometallic catalyst can optionally be used together with an excess of a salt mixture, such as KI and/or CuI. As one of skill in the art can appreciate, when employing each of the coupling reactions shown in Scheme 2, it can, in various aspects, be advantageous to perform the reactions in a dry atmosphere, for example under argon, or even in a dry box to minimize and/or avoid moisture or oxygen inclusion.

30

Scheme 2.



In one aspect, the compounds of the invention can be useful in a variety of optical applications. When utilized as light emitting materials, the inventive compounds can be useful in organic light emitting diodes (OLED)s, luminescent devices and displays, and/or other light emitting devices. With reference to FIG. 3, for example, a specific embodiment, 2-(3-(pyridin-2-yl)phenoxy)-6-(pyridin-2-yloxy)phenyl]Pd(II) exhibits photoluminescence (absorption of light followed by emission of light) across a range of wavelengths, including a narrow blue emission and other emission bands out to the red to near-IR region of the spectrum.

The emission (and absorption) profile of the compounds can be tuned by varying the structure of the ligand surrounding the metal center. For example, compounds having a ligand with electron withdrawing substituents can, in one aspect, generally exhibit different optical properties, including emission and absorption, than compounds having a ligand with electron donating substituents. Generally, a chemical structural change can affect the electronic structure of the compound, thereby affecting the absorption and emission properties of the compound. Thus, the compounds of the present invention can be tailored or tuned to a specific application that desires a particular emission or absorption characteristic. One of skill in the art, in possession of this disclosure, could readily design and/or select an appropriate multidentate palladium compound, in accordance with the various aspects described herein, to use in a particular application.

In one embodiment, the compounds can be used in an OLED. FIG. 1 illustrates a cross-sectional view of an exemplary OLED 100, which includes substrate 102 with an anode 104, which is typically a transparent material, such as indium tin oxide, a layer of hole-transporting material(s) (HTL) 106, a layer of light processing material 108, such as an emissive material (EML) including an emitter and a host, a layer of electron-transporting material(s) (ETL) 110, and a metal cathode layer 112.

In such an embodiment, the layer of light processing material 108 can comprise one or more compounds of the present invention optionally together with a host material.

31

The host material, if present, can be any suitable host material known in the art. The emission color of an OLED can be determined by the emission energy (optical energy gap) of the light processing material **108**, which as discussed above can be tuned by tuning the electronic structure of the emitting compounds and/or the host material. Both the hole-transporting material in the HTL layer **106** and the electron-transporting material(s) in the ETL layer **110** can comprise any suitable hole-transporter known in the art. A selection of which is well within the purview of those skilled in the art.


It will be apparent that the compounds of the present invention can exhibit phosphorescence. Phosphorescent OLEDs (i.e., OLEDs with phosphorescent emitters) typically have higher device efficiencies than other OLEDs, such as fluorescent OLEDs. Light emitting devices based on electrophosphorescent emitters are described in more detail in WO2000/070655 to Baldo et al., which is incorporated herein by this reference for its teaching of OLEDs, and in particular phosphorescent OLEDs.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices and/or methods claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.), but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in ° C. or is at ambient temperature, and pressure is at or near atmospheric.

Example 1. Preparation of Specific Embodiment
2-(3-(pyridin-2-yl)phenoxy)-6-(pyridin-2-yloxy)
phenyl]Pd(II)


Synthesis of 3-(pyridin-2-yloxy)phenol



Under a nitrogen atmosphere, a pressure vessel was charged with a magnetic stir bar, resorcinol (110 mmol), 2-bromopyridine (100 mmol), 1-methylimidazole (5 mmol), and potassium carbonate (200 mmol). Pyridine (80 mL) was added and bubbled with nitrogen for 20 minutes before copper(I) iodide (10 mmol) was added and bubbled 10 minutes further. The vessel was sealed and heated to 140° C. while stirring. After 2 days, the solution was allowed to cool. The solids were filtered off and rinsed with a 50:50 mixture of toluene and methanol. The filtrate was reduced by rotary evaporation and 150 ml of water containing 10 mL glacial acetic acid was added and shaken vigorously. The water was decanted off and 50 mL of DCM was added, forming an off white precipitate which was collected by vacuum filtration and dried with ether, resulting in the pure product 3-(pyridin-2-yloxy)phenol with a 55% yield. ¹H NMR (CDCl₃): 5.98 (s, 1H), 6.59 (s, 1H), 6.62-6.69 (m, 2H), 6.94 (d, 1H), 7.02 (dd, 1H), 7.23 (vt, 1H), 7.70 (dd, 1H), 8.23 (b, 1H)


32

Synthesis of
2-(3-(3-bromophenoxy)phenoxy)pyridine



Under a nitrogen atmosphere, a pressure vessel was charged with a magnetic stir bar, 3-(pyridin-2-yloxy)phenol (50 mmol), 2,6-dibromopyridine (50 mmol), 1-methylimidazole (25 mmol), and potassium carbonate (100 mmol). Toluene (80 mL) was added and bubbled with nitrogen for 20 minutes before copper(I) iodide (5 mmol) was added and the solution bubbled for 10 minutes further. The vessel was sealed and heated to 140° C. while stirring. After 2 days, the solution was allowed to cool and the solids were filtered off and rinsed with dichloromethane. The filtrate was added to a separatory funnel containing DCM and water. The water phase was washed 3 times with 75 mL DCM, and the combined organic layers were washed once with pure water. The organic layer was collected, dried with magnesium sulfate, filtered, and the filtrate reduced by rotary evaporation. The resulting oil was purified by column chromatography using DCM over silica resulting in the pure product 2-(3-(3-bromophenoxy)phenoxy)pyridine with a 60% yield. ¹H NMR (CDCl₃): 6.80-6.85 (m, 2H), 6.91 (s, 1H), 6.94 (s, 1H), 6.97-7.03 (m, 2H), 7.19 (vt, 1H), 7.21-7.24 (m, 2H), 7.36 (vt, 1H), 7.70 (dd, 1H), 8.21 (dd, 1H).


Synthesis of 2-(3-(3-(pyridin-2-yl)phenoxy)phenoxy)pyridine



Under a nitrogen atmosphere, an oven dried three neck flask was charged with a magnetic stir bar, 2-(3-(3-bromophenoxy)phenoxy) (10 mmol), and 2-(tripropylstannyl)pyridine (10 mmol). Dry toluene (100 mL) was added and bubbled with nitrogen for 20 minutes before Tetrakis(triphenylphosphine)palladium(0) (0.5 mmol) was added, bubbled 10 minutes further, and brought to reflux for 2 days. After cooling, the contents of the flask were filtered, the liquid reduced by rotary evaporation, and the resulting oil was purified by column chromatography using DCM over silica to yield the pure product 2-(3-(3-(pyridin-2-yl)phenoxy)phenoxy)pyridine with a 65% yield. ¹H NMR (CDCl₃): 6.84 (vt, 1H), 6.85-6.89 (m, 2H), 6.91 (d, 1H), 6.98 (dd, 1H), 7.11 (dd, 1H), 7.24 (dd, 1H), 7.34 (vt, 1H), 7.44 (vt, 1H), 7.66-7.78 (m, 5H), 8.19 (dd, 1H), 8.67 (dd, 1H).

33


Synthesis of [2-(3-(pyridin-2-yl)phenoxy)-6-(pyridin-2-yloxy)phenyl]Pd(II)



A mixture of 2-(3-(3-(pyridin-2-yl)phenoxy)phenoxy)pyridine (1 mmol), K_2PdCl_4 (1 mmol), and acetic acid (10 mL) was refluxed for 3 days. The mixture was allowed to cool to room temperature. The resulting white complex was filtered off and washed with H_2O , MeOH, and Et_2O , and dried under vacuum to 2-(3-(pyridin-2-yl)phenoxy)-6-(pyridin-2-yloxy)phenyl]Pd(II). The product was purified by sublimation for further testing.

What is claimed is:

1. A compound represented by the formula:



wherein:

each R^1 and R^2 in $(R^1)_2$ and $(R^2)_2$ independently represents hydrogen;

Y^{1a} represents O;

n is an integer 0;

each of Y^{2a} , Y^{2b} , Y^{2c} and Y^{2d} independently represents N, C, or CH;

each of Y^{3a} , Y^{3b} , Y^{3c} , Y^{3d} , and Y^{3e} independently represents N, NR^{6a} , C or CR^{6b} , wherein each of R^{6a} and R^{6b} independently represents hydrogen or optionally substituted C_1 - C_4 alkyl, and the ring which comprises the Y^{3a} , Y^{3b} , Y^{3c} , Y^{3d} , and Y^{3e} groups is a heteroaryl group having one or two nitrogen atoms, wherein the heteroaryl group is unsubstituted or substituted with one or more optionally substituted C_1 - C_4 alkyl;

each of Y^{4a} , Y^{4b} , Y^{4c} , and Y^{4d} independently represents N, NR^{6a} , C or CR^{6b} , wherein each of R^{6a} and R^{6b} independently represents hydrogen or optionally substituted C_1 - C_4 alkyl, and the ring which comprises the Y^{4a} , Y^{4b} , Y^{4c} , and Y^{4d} groups is a heteroaryl group having one or two nitrogen atoms, wherein the heteroaryl group is unsubstituted or substituted with one or more optionally substituted C_1 - C_4 alkyl;


m is an integer 1.

34


2. An organic light-emitting diode (OLED) comprising, as an emissive material, a compound represented by the formula of claim 1.

3. An organic photovoltaic device comprising, as a donor or acceptor material, a compound represented by the formula of claim 1.

4. The compound of claim 1, which is represented by the formula:




5. The compound of claim 1, which is represented by the formula:




35

-continued




36

-continued




37

-continued




38

-continued



39


-continued



6. An organic light-emitting diode (OLED) comprising, as an emissive material, a compound represented by the formula of claim 4.

7. An organic photovoltaic device comprising, as a donor or acceptor material, a compound represented by the formula of claim 4.

8. The compound of claim 1, wherein the compound is represented by the formula:



9. The compound of claim 1, wherein Y^{2a} , Y^{2b} , Y^{2c} , and Y^{2d} are N.

10. The compound of claim 1, wherein Y^{2a} , Y^{2b} , Y^{2c} , and Y^{2d} are C or CH.

11. The compound of claim 1, wherein one of Y^{2a} , Y^{2b} , Y^{2c} , and Y^{2d} is N and the others are C or CH.

12. The compound of claim 1, wherein two of Y^{2a} , Y^{2b} , Y^{2c} , and Y^{2d} are N and the others are C or CH.

13. The compound of claim 1, wherein three of Y^{2a} , Y^{2b} , Y^{2c} , and Y^{2d} are N and the other is C or CH.

14. The compound of claim 1, wherein the ring which comprises Y^{3a} , Y^{3b} , Y^{3c} , Y^{3d} , and Y^{3e} groups is a heteroaryl group having one nitrogen atom.

15. The compound of claim 1, wherein the ring which comprises Y^{3a} , Y^{3b} , Y^{3c} , Y^{3d} , and Y^{3e} groups is a heteroaryl group having two nitrogen atoms.

40


16. The compound of claim 1, wherein the ring which comprises Y^{3a} , Y^{3b} , Y^{3c} , Y^{3d} , and Y^{3e} groups is a pyrazolyl group.

17. The compound of claim 1, wherein the ring which comprises Y^{4a} , Y^{4b} , Y^{4c} , and Y^{4d} groups is a heteroaryl group having one nitrogen atom.


18. The compound of claim 1, wherein the ring which comprises Y^{4a} , Y^{4b} , Y^{4c} , and Y^{4d} groups is a heteroaryl group having two nitrogen atoms.

19. The compound of claim 1, wherein the ring which comprises Y^{4a} , Y^{4b} , Y^{4c} , and Y^{4d} groups is a pyrazolyl group.

20. The compound of claim 1, which is represented by the formula:




21. The compound of claim 1, which is represented by the formula:




41

-continued



42

-continued




22. The compound of claim 1, which is represented by the formula:

5
10
15
20
25
30
35
40
45
50
55
60
65


43

-continued




44

-continued




23. The compound of claim 1, which is represented by the formula:




45

-continued



or



* * * * *