



(86) Date de dépôt PCT/PCT Filing Date: 2008/09/29
(87) Date publication PCT/PCT Publication Date: 2009/04/02
(45) Date de délivrance/Issue Date: 2017/06/20
(85) Entrée phase nationale/National Entry: 2010/03/24
(86) N° demande PCT/PCT Application No.: GB 2008/003305
(87) N° publication PCT/PCT Publication No.: 2009/040556
(30) Priorités/Priorities: 2007/09/28 (GB0719038.2);
2008/04/15 (GB0806844.7)


(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01),
A61K 31/551 (2006.01), *A61P 35/00* (2006.01),
C07D 519/00 (2006.01)

(72) Inventeurs/Inventors:
HOLLICK, JONATHAN JAMES, GB;
JONES, STUART DONALD, GB;
FLYNN, CLAIRE JUNE, GB;
THOMAS, MICHEAL GEORGE, GB

(73) Propriétaire/Owner:
CYCLACEL LIMITED, GB

(74) Agent: SIM & MCBURNEY

(54) Titre : DERIVES DE PYRIMIDINE COMME INHIBITEURS DE PROTEINE KINASES
(54) Title: PYRIMIDINE DERIVATIVES AS PROTEIN KINASE INHIBITORS



(VIII)

(57) **Abrégé/Abstract:**

The present invention relates to a compound of formula (VII), or a pharmaceutically acceptable salt or ester thereof wherein: X is NR⁷; Y is O or N-(CH₂)_n R¹⁹; n is 1, 2 or 3; m is 1 or 2; R¹ and R² are each independently H, alkyl or cycloalkyl; R⁴ and R^{4'} are each independently H or alkyl; or R⁴ and R^{4'} together form a spiro cycloalkyl group; R¹⁹ is H, alkyl, aryl or a cycloalkyl group; R⁶ is OR⁸ or halogen; and R⁷ and R⁸ are each independently H or alkyl. Further aspects relate to pharmaceutical compositions comprising said compounds and use therefore in the treatment of proliferative disorders and the like.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 April 2009 (02.04.2009)

PCT

(10) International Publication Number
WO 2009/040556 A1

(51) International Patent Classification:

C07D 487/04 (2006.01) A61P 35/00 (2006.01)
C07D 519/00 (2006.01) A61K 31/551 (2006.01)

(21) International Application Number:

PCT/GB2008/003305

(22) International Filing Date:

29 September 2008 (29.09.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0719038.2 28 September 2007 (28.09.2007) GB
0806844.7 15 April 2008 (15.04.2008) GB

(71) Applicant (for all designated States except US): **CYCLACEL LIMITED** [GB/GB]; 6-8 Underwood Street, London N1 7JQ (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HOLLICK, Jonathan, James** [GB/GB]; 6 Quarryknowe Crescent, Inchtute, Perth PH14 9RH (GB). **JONES, Stuart, Donald** [GB/GB]; 17 Oakwood Drive, Prestbury, Cheshire SK10 4HG (GB). **FLYNN, Claire, June** [GB/GB]; 33 Esk Gardens, Carnoustie, Angus DD7 6GH (GB). **THOMAS, Micheal, George** [GB/GB]; 7 Midlane Close, Braco FK15 9PQ (GB).

(74) Agents: **CLYDE-WATSON, Zoe** et al.; D. Young & Co., 120 Holborn, London EC1N 2DY (GB).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(54) Title: PYRIMIDINE DERIVATIVES AS PROTEIN KINASE INHIBITORS



(VIII)

(57) Abstract: The present invention relates to a compound of formula (VII)I, or a pharmaceutically acceptable salt or ester thereof, wherein: X is NR⁷; Y is O or N-(CH₂)_nR¹⁹; n is 1, 2 or 3; m is 1 or 2; R¹ and R² are each independently H, alkyl or cycloalkyl; R⁴ and R^{4'} are each independently H or alkyl; or R⁴ and R^{4'} together form a spiro cycloalkyl group; R¹⁹ is H, alkyl, aryl or a cycloalkyl group; R⁶ is OR⁸ or halogen; and R⁷ and R⁸ are each independently H or alkyl. Further aspects relate to pharmaceutical compositions comprising said compounds and use therefore in the treatment of proliferative disorders and the like.

PYRIMIDINE DERIVATIVES AS PROTEIN KINASE INHIBITORS

The present invention relates to substituted pyrimidine derivatives and their use in therapy. More specifically, but not exclusively, the invention relates to compounds that
5 are capable of inhibiting one or more protein kinases, particularly polo-like kinases.

BACKGROUND TO THE INVENTION

The Polo-like kinase family consists of key cell cycle regulatory enzymes with integral roles in controlling entry into and progression through mitosis. Many tumour cells
10 express high levels of PLK1 and are responsive to antisense oligonucleotides targeting this protein.

Initiation of mitosis requires activation of M-phase promoting factor (MPF), *i.e.* the complex between CDK1 and B-type cyclins [Nurse, P. (1990) *Nature*, 344, 503-508].
15 The latter accumulate during the S and G2 phases of the cell cycle and promote the inhibitory phosphorylation of the MPF complex by WEE1, MIK1, and MYT1 kinases. At the end of the G2 phase, corresponding dephosphorylation by the dual-specificity phosphatase CDC25C triggers the activation of MPF [Nigg, E.A. (2001) *Nat. Rev. Mol. Cell Biol.*, 2, 21-32]. In interphase, cyclin B localizes to the cytoplasm and becomes
20 phosphorylated during prophase, followed by nuclear translocation. The nuclear accumulation of active MPF during prophase is thought to be important for initiating M-phase events [Takizawa, C.G. and Morgan, D.O. (2000) *Curr. Opin. Cell Biol.*, 12, 658-665]. However, nuclear MPF is kept inactive by WEE1 unless counteracted by CDC25C. The phosphatase CDC25C itself, localized to the cytoplasm during
25 interphase, accumulates in the nucleus in prophase. The nuclear entry of both cyclin B and CDC25C are promoted through phosphorylation by PLK1 [Roshak, A.K., Capper, E.A., Imburgia, C., Fornwald, J., Scott, G. and Marshall, L.A. (2000) *Cell. Signalling*, 12, 405-411]. This kinase is thus an important regulator of M-phase initiation.

In humans, there exist three closely related polo-like kinases (PLKs) [Glover, D.M., Hagan, I.M. and Tavares, A.A. (1998) *Genes Dev.*, 12, 3777-3787]. They contain a highly homologous N-terminal catalytic kinase domain and their C-termini contain two or three conserved regions, the polo boxes. The function of the polo boxes remains incompletely understood but polo box-dependent PLK1 activity is required for proper metaphase/anaphase transition and cytokinesis [Seong, Y.-S., Kamijo, K., Lee, J.-S., Fernandez, E., Kuriyama, R., Miki, T. and Lee, K.S. (2002) *J. Biol. Chem.*, 277, 32282-32293]. Of the three PLKs, PLK1 is the best characterized; it regulates a number of cell division cycle effects, including the onset of mitosis, DNA-damage checkpoint activation, regulation of the anaphase promoting complex, phosphorylation of the proteasome, and centrosome duplication and maturation. Mammalian PLK2 (also known as SNK) and PLK3 (also known as PRK and FNK) were originally shown to be immediate early gene products. PLK3 kinase activity appears to peak during late S and G2 phase. It is also activated during DNA damage checkpoint activation and severe oxidative stress. PLK3 also plays an important role in the regulation of microtubule dynamics and centrosome function in the cell and deregulated PLK3 expression results in cell cycle arrest and apoptosis [Wang, Q., Xie, S., Chen, J., Fukusawa, K., Naik, U., Traganos, F., Darzynkiewicz, Z., Jhanwar-Uniyal, M. and Dai, W. (2002) *Mol. Cell. Biol.*, 22, 3450-3459]. PLK2 is the least-well understood homologue of the three PLKs. Both PLK2 and PLK3 may have additional important post-mitotic functions [Kauselmann, G., Weiler, M., Wulff, P., Jessberger, S., Konietzko, U., Scafidi, J., Staubli, U., Bereiter-Hahn, J., Strebhardt, K. and Kuhl, D. (1999) *EMBO J.*, 18, 5528-5539].

The fact that human PLKs regulate some fundamental aspects of mitosis was shown by anti-PLK1 antibody microinjection of human tumour cells [Lane, H.A. and Nigg, E.A. (1996) *J. Cell. Biol.*, 135, 1701-1713]. This treatment had no effect on DNA replication but impaired cell division. Cells were arrested in mitosis and showed abnormal distribution of condensed chromatin and monoastal microtubules nucleated from duplicated but unseparated centrosomes. By contrast, non-immortalized human cells

arrested as single, mononucleated cells in G2. Moreover, when PLK1 function was blocked through adenovirus-mediated delivery of a dominant-negative gene, tumour-selective apoptosis in many tumour cell lines was observed, whereas again normal epithelial cells, although arrested in mitosis, escaped the mitotic catastrophe seen in
5 tumour cells [Cogswell, J.P., Brown, C.E., Bisi, J.E. and Neill, S.D. (2000) *Cell Growth Differ.*, 11, 615-623]. PLK1 activity is thus necessary for the functional maturation of centrosomes in late G2/early prophase and subsequent establishment of a bipolar spindle. Furthermore, these results suggest the presence in normal cells of a centrosome-maturation checkpoint that is sensitive to PLK1 impairment. Depletion of
10 cellular PLK1 through the small interfering RNA (siRNA) technique also confirmed that this protein is required for multiple mitotic processes and completion of cytokinesis [Liu, X. and Erikson, R.L. (2003) *Proc. Natl. Acad. Sci. USA*, 100, 5789–5794]. A potential therapeutic rationale for PLK inhibition is also suggested by work with PLK1-specific antisense oligonucleotides, which were shown to induce growth
15 inhibition in cancer cells both *in vitro* and *in vivo* [Spankuch-Schmitt, B., Wolf, G., Solbach, C., Loibl, S., Knecht, R., Stegmüller, M., von Minckwitz, G., Kaufmann, M. and Strebhardt, K. (2002) *Oncogene*, 21, 3162-3171]. Constitutive expression of PLK1 in mammalian cells was shown to lead to malignant transformation [Smith, M.R., Wilson, M.L., Hamanaka, R., Chase, D., Kung, H., Longo, D.L. and Ferris, D.K.
20 (1997) *Biochem. Biophys. Res. Commun.*, 234, 397-405]. Furthermore, overexpression of PLK1 is frequently observed in human tumours and PLK1 expression is of prognostic value for patients suffering from various types of tumours [Takahashi, T., Sano, B., Nagata, T., Kato, H., Sugiyama, Y., Kunieda, K., Kimura, M., Okano, Y. and Saji, S. (2003) *Cancer Science*, 94, 148-152; Tokumitsu, Y., Mori, M., Tanaka, S.,
25 Akazawa, K., Nakano, S. and Niho, Y. (1999) *Int. J. Oncol.*, 15, 687-692; Wolf, G., Elez, R., Doermer, A., Holtrich, U., Ackermann, H., Stutte, H.J., Altmannsberger, H.-M., Rübsamen-Waigmann, H. and Strebhardt, K. (1997) *Oncogene*, 14, 543-549].

Although the therapeutic potential of pharmacological PLK inhibition has been
30 appreciated [Kraker, A.J. and Booher, R.N. (1999) In *Annual Reports in Medicinal*


Chemistry (Vol. 34) (Doherty, A.M., ed.), pp. 247-256, Academic Press], very little has been reported to date concerning small molecule PLK inhibitors that may be useful as drugs. One of the few biochemical PLK1 inhibitors characterized to date is scytonemin, a symmetric indolic marine natural product [Stevenson, C.S., Capper, E.A., Roshak, A.K., Marquez, B., Eichman, C., Jackson, J.R., Mattern, M., Gerwick, W.H., Jacobs, R.S. and Marshall, L.A. (2002) *J. Pharmacol. Exp. Ther.*, 303, 858-866; Stevenson, C.S., Capper, E.A., Roshak, A.K., Marquez, B., Grace, K., Gerwick, W.H., Jacobs, R.S. and Marshall, L.A. (2002) *Inflammation Research*, 51, 112-114]. Scytonemin inhibits phosphorylation of CDC25C by recombinant PLK1 with an IC₅₀ value of about 2 µM (at an ATP concentration of 10 µM). Inhibition is apparently reversible and the mechanism with respect to ATP of mixed-competitive mode. Similar potency against other protein serine/threonine- and dual specificity cell-cycle kinases, including MYT1, CHK1, CDK1/cyclin B, and PKC, was observed. Scytonemin showed pronounced anti-proliferative effects on various human cell lines in vitro. Further small molecule PLK inhibitors and their use in the treatment of proliferative disorders are described in International patent application WO2004/067000 in the name of Cyclacel Limited.

The present invention seeks to elucidate new small molecule PLK inhibitors. More specifically, the invention seeks to provide small molecule PLK inhibitors that have therapeutic applications in the treatment of a range of proliferative disorders.

STATEMENT OF INVENTION

A first aspect of the invention relates to a compound of formula VIII, or a pharmaceutically acceptable salt or ester thereof,

5



(VIII)

wherein:

5 X is NR^7 ;

Y is O or $\text{N}-(\text{CH}_2)_n\text{R}^{19}$;

n is 1, 2 or 3;

m is 1 or 2;

R^1 and R^2 are each independently H, alkyl or cycloalkyl;

10 R^4 and $\text{R}^{4'}$ are each independently H or alkyl; or

R^4 and $\text{R}^{4'}$ together form a spiro cycloalkyl group;


R^{19} is H, alkyl, aryl or a cycloalkyl group;

R^6 is OR^8 or halogen; and

R^7 and R^8 are each independently H or alkyl.

15

A second aspect of the invention relates to a compound of formula VI, or a pharmaceutically acceptable salt or ester thereof,



(VI)

20 wherein:

X is NR^7 ;

6

R^1 and R^2 are each independently H, alkyl or cycloalkyl;

R^4 and $R^{4'}$ are each independently H or alkyl; or

R^4 and $R^{4'}$ together form a spiro cycloalkyl group;


R^{18} is H or alkyl, wherein said alkyl group is optionally substituted by R^6 ;

5 each R^6 is independently OR^8 or halogen; and

R^7 and R^8 are each independently H or alkyl.

A third aspect of the invention relates to a compound of formula VII, or a pharmaceutically acceptable salt or ester thereof,

10



(VII)

wherein:

X is NR^7 ;

15 R^1 and R^2 are each independently H, alkyl or cycloalkyl;

R^4 and $R^{4'}$ are each independently H or alkyl; or

R^4 and $R^{4'}$ together form a spiro cycloalkyl group;

R^6 is OR^8 or halogen; and

R^7 and R^8 are each independently H or alkyl.

20

A fourth aspect of the invention relates to a compound selected from the following:

Compound No.	Name
254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide

218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
345	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide
373	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide
374	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-

	ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
375	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide
376	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
379	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
380	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
381	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-

	ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide
382	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide
383	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
384	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
385	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
386	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
387	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
388	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide
389	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide
390	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

	tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide
391	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide

A fifth aspect of the invention relates to a pharmaceutical composition comprising a compound as described above admixed with a pharmaceutically acceptable diluent, excipient or carrier.

5

A sixth aspect of the invention relates to the use of a compound as described above in the preparation of a medicament for treating a proliferative disorder.

A seventh aspect of the invention relates to a method of treating a proliferative disorder, said method comprising administering to a subject a therapeutically effective amount of a compound as described above.

An eighth aspect of the invention relates to a method of treating a PLK-dependent disorder, said method comprising administering to a subject a therapeutically effective amount of a compound as described above.

A ninth aspect of the invention relates to a process for preparing compounds as defined above.

Another aspect of the invention relates to a compound selected from the group consisting of:

Compound No.	Name
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;

10a

194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;

a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable ester thereof.

- 5 Another aspect of the invention relates to a compound which is selected from the group consisting of:

Compound No.	Name
254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide;
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide;
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide;

371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
345	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide;
373	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide;
374	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide;
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
375	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide;
376	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide;
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

	tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
379	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
380	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
381	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide;
382	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide;
383	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
384	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;

10d

385	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
386	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
387	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
388	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide;
389	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide;
390	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide;
391	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide;

a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable ester thereof.

5 DETAILED DESCRIPTION

As used herein, the term “alkyl” includes both saturated straight chain and branched alkyl groups which may be substituted (mono- or poly-) or unsubstituted. Preferably, the alkyl group is a C₁₋₂₀ alkyl group, more preferably a C₁₋₁₅, more preferably still a C₁₋₁₂ alkyl group, more preferably still, a C₁₋₆ alkyl group, more preferably a C₁₋₃ alkyl group. Particularly preferred alkyl groups include, for example, methyl, ethyl, propyl,

isopropyl, butyl, isobutyl, tert-butyl, pentyl and hexyl. Suitable substituents include, for example, one or more R^6 groups. Preferably, the alkyl group is unsubstituted.


As used herein, the term “cycloalkyl” refers to a cyclic alkyl group which may be substituted (mono- or poly-) or unsubstituted. Preferably, the cycloalkyl group is a C_{3-12} cycloalkyl group, more preferably a C_{3-6} cycloalkyl group. Suitable substituents include, for example, one or more R^6 groups.

As used herein, the term “aryl” refers to a C_{6-12} aromatic group which may be substituted (mono- or poly-) or unsubstituted. Typical examples include phenyl and naphthyl etc. Suitable substituents include, for example, one or more R^6 groups.

COMPOUNDS OF FORMULA VIII

As mentioned above, one aspect of the invention relates to a compound of formula VIII, or a pharmaceutically acceptable salt or ester thereof.

In one particularly preferred embodiment, the invention relates to a compound of formula VIIIa, or a pharmaceutically acceptable salt or ester thereof,



(VIIIa)

wherein:

X is NR^7 ;

R^1 and R^2 are each independently H, alkyl or cycloalkyl;

12

R^4 and $R^{4'}$ are each independently H or alkyl; or
 R^4 and $R^{4'}$ together form a spiro cycloalkyl group;
 R^{19} is H, alkyl, aryl or a cycloalkyl group;
 R^6 is OR^8 or halogen; and
5 R^7 and R^8 are each independently H or alkyl.

In one preferred embodiment, R^{19} is H, alkyl, or a cycloalkyl group.

Preferably, R^1 is H or alkyl, more preferably, alkyl. Even more preferably, R^1 is methyl
10 or ethyl, more preferably still, Me.

Preferably, R^2 is a cycloalkyl group, more preferably a C_{3-6} cycloalkyl group. Even
more preferably, R^2 is a cyclopentyl or cyclohexyl group, more preferably still,
cyclopentyl.

15

Preferably, R^7 is H or alkyl, more preferably, H or methyl, even more preferably, H.

In one preferred embodiment, one of R^4 and $R^{4'}$ is alkyl, and the other is H or alkyl.

20 In another preferred embodiment, R^4 and $R^{4'}$ are each independently alkyl. More
preferably, R^4 and $R^{4'}$ are both methyl.

In another preferred embodiment, R^4 and $R^{4'}$ are both H.

25 In another preferred embodiment, R^4 and $R^{4'}$ together form a spiro cycloalkyl group,
more preferably, a spiro C_{3-6} cycloalkyl group. Even more preferably, R^4 and $R^{4'}$
together form a spiro C_3 or C_4 cycloalkyl group, more preferably still, a C_3 cycloalkyl
group.


30 In one preferred embodiment, R^6 is OR^8 , even more preferably, OMe.

13

Preferably, R¹⁹ is cyclopropyl.

In another particularly preferred embodiment, the invention relates to a compound of formula VIIIb, or a pharmaceutically acceptable salt or ester thereof,

5




(VIIIb)

wherein R¹, R², R⁶, R⁴, R^{4'} and X are as defined above.

10

In another particularly preferred embodiment, the invention relates to a compound of formula VIIIc, or a pharmaceutically acceptable salt or ester thereof,



(VIIIc)

wherein X, R¹, R², R⁶, R⁴, R^{4'} and R¹⁹ are as defined above.

15

In one particularly preferred embodiment, said compound of formula VIII is selected from the following:

20

Compound No.	Name
371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
379	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
380	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
381	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide
382	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-

	ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide
383	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
384	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
385	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
386	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
387	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
388	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide
389	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide
390	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide
391	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-

	ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide
--	--

and pharmaceutically acceptable salts and esters thereof.

More preferably, the compound of formula VIII is selected from the following:

Compound No.	Name
371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide

5


or a pharmaceutically acceptable salt or ester thereof.

Even more preferably, the compound of the invention is compound [371].

10 COMPOUNDS OF FORMULA VII

One aspect of the invention relates to a compound of formula VII, or a pharmaceutically acceptable salt or ester thereof,

17



(VII)

wherein:

X is NR⁷;

5 R¹ and R² are each independently H, alkyl or cycloalkyl;

R⁴ and R^{4'} are each independently H or alkyl; or

R⁴ and R^{4'} together form a spiro cycloalkyl group; and

R⁶ is OR⁸ or halogen; and

R⁷ and R⁸ are each independently H or alkyl.

10

Preferably, R¹ is alkyl, more preferably, methyl.

Preferably, R² is a cycloalkyl group, more preferably a C₃₋₆ cycloalkyl group. Even more preferably, R² is a cyclopentyl or cyclohexyl group, more preferably still,
15 cyclopentyl.

Preferably, R⁷ is H or alkyl, more preferably, H or methyl, even more preferably, H.

In one preferred embodiment, one of R⁴ and R^{4'} is alkyl, and the other is H or alkyl.

20

In another preferred embodiment, R⁴ and R^{4'} are each independently alkyl. More preferably, R⁴ and R^{4'} are both methyl.

In another preferred embodiment, R⁴ and R^{4'} are both H.

25

In another preferred embodiment, R⁴ and R^{4'} together form a spiro cycloalkyl group, more preferably, a spiro C₃₋₆ cycloalkyl group. More preferably, R⁴ and R^{4'} together form a spiro C₃ or C₄ cycloalkyl group, even more preferably, a C₃ cycloalkyl group.

- 5 Preferably, R⁶ is OR⁸, even more preferably, OMe.

In one preferred embodiment, the compound of formula VII is selected from the following:


Compound No.	Name
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide

- 10 and pharmaceutically acceptable salts and esters thereof.

COMPOUNDS OF FORMULA VI

One aspect of the invention relates to a compound of formula VI, or a pharmaceutically acceptable salt or ester thereof,

19



(VI)

wherein:

X is NR⁷;

5 R¹ and R² are each independently H, alkyl or cycloalkyl;

one of R⁴ and R^{4'} is alkyl, and the other is H or alkyl; or

R⁴ and R^{4'} together form a spiro cycloalkyl group;

R¹⁸ is H or alkyl, wherein said alkyl group is optionally substituted by R⁶;

each R⁶ is independently OR⁸ or halogen; and

10 R⁷ and R⁸ are each independently H or alkyl.

Preferably, R¹ is alkyl, more preferably methyl.

15 Preferably, R² is a cycloalkyl group, more preferably, a C₃₋₆ cycloalkyl group. Even more preferably, R² is cyclopentyl or cyclohexyl, more preferably still, cyclopentyl.

Preferably, R⁷ is H or alkyl, more preferably, H or methyl, even more preferably, H.

In one preferred embodiment, one of R⁴ and R^{4'} is alkyl, and the other is H or alkyl.

20

In another preferred embodiment, R⁴ and R^{4'} are each independently alkyl. More preferably, R⁴ and R^{4'} are both methyl.

In another preferred embodiment, R⁴ and R^{4'} are both H.

25

In another preferred embodiment, R⁴ and R^{4'} together form a spiro cycloalkyl group, more preferably, a spiro C₃₋₆ cycloalkyl group. Even more preferably, R⁴ and R^{4'} together form a spiro C₃ or C₄ cycloalkyl group, even more preferably, a C₃ cycloalkyl group.

5

Preferably, R⁶ is OMe or F.

Preferably, R¹⁸ is methyl or CH₂CH₂OH.

10 In one preferred embodiment, the compound of formula VI is selected from the following:

254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
373	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide
374	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide
375	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide

and pharmaceutically acceptable salts and esters thereof.

In one especially preferred embodiment, the compound is [254], or a pharmaceutically acceptable salt or ester thereof.

In one highly preferred embodiment, the compound is the HCl salt of compound [254],
 5 i.e. 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro [cyclopropane-1,7'-pyrimido [4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide.HCl.

10 Another aspect of the invention relates to a compound which is selected from the following:

Compound No.	Name
254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-

	(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
345	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide
373	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide
374	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
375	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide
376	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-

	ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
379	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
380	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
381	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide
382	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide
383	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
384	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
385	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-

	ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
386	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
387	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide
388	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide
389	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide
390	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide
391	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide

and pharmaceutically acceptable salts and esters thereof.

Preferably, the compound is selected from the following:

Compound No.	Name
254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

	tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide
345	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide

and pharmaceutically acceptable salts and esters thereof.

Advantageously, certain compounds of the invention exhibit improved selectivity for one or more kinases compared to structurally related compounds previously known in the art

In one preferred embodiment, the compound of the invention exhibits selectivity for one or more kinases over one or more other kinases.

For example, in one preferred embodiment, the compound of the invention is capable of preferentially inhibiting PLK family kinases over one or more other protein kinases; thus the compound is capable of “selectively” inhibiting PLK family kinases over one or more other protein kinases. As used herein, the term “selectively” refers to the compounds that are selective for PLK over one or more other protein kinases. Preferably, the selectivity ratio for PLK family kinases over one or more other protein kinases is greater than about 2 to 1, more preferably greater than about 5 to 1 or about 10 to 1, even more preferably greater than about 20 to 1, or 50 to 1 or 100 to 1. Selectivity ratios may be determined by the skilled person in the art.

In one particularly preferred embodiment, the compound of the invention displays selectivity for PLK1. Thus, in one preferred embodiment, the compound of the invention is capable of preferentially inhibiting PLK1 over one or more other protein kinases. Preferably, the selectivity ratio for PLK1 over one or more other protein kinases is greater than about 2 to 1, more preferably greater than about 5 to 1 or about 10 to 1, even more preferably greater than about 20 to 1, or 50 to 1 or 100 to 1.

20

In an even more preferred embodiment, the compound of the invention displays selectivity for PLK1 over PLK2 and/or PLK3. Thus, in one preferred embodiment, the compound of the invention is capable of preferentially inhibiting PLK1 over PLK2 and/or PLK3. Preferably, the selectivity ratio for PLK1 over PLK2 and/or PLK3 is greater than about 2 to 1, more preferably greater than about 5 to 1 or about 10 to 1, even more preferably greater than about 20 to 1, or 50 to 1 or 100 to 1.

25

By way of illustration, compound [218] displays selectivity for PLK1 over PLK2 and PLK3 compared with selected compounds of the prior art. For example, compound [218] of the invention is twice as selective for PLK1 versus PLK2 compared to compound [I-64] of WO 07/095188, *ca* 6-fold more selective than compound [I-76]

30

and 7-fold more selective than compound [I-4] of WO 07/095188. Similarly, compound [218] is 5-fold more selective for PLK1 versus PLK3 compared to compound [I-64] of WO 07/095188, *ca* 12-fold more selective than compound [I-76] and 5-fold more selective than compound [I-4] of WO 07/095188. Further details may
5 be found in the accompanying Examples.

In another preferred embodiment, the compounds of the invention exhibit superior solubility and/or pharmacokinetic properties when compared to structurally related compounds already known in the art.

10 For Example, compound [254] displays superior solubility and/or pharmacokinetic properties to Examples [I-76] and [I-253] of WO 07/095188, thereby rendering the compound more suitable for the treatment of diseases. Again, further details of these comparative studies may be found in the accompanying Examples.

15 Similarly, compound [371] displays advantageous pharmacokinetic properties in terms of its systemic exposure and oral bioavailability, compared to structurally related compounds known in the art.

20 One preferred embodiment of the invention relates to a subset of compounds that are characterised by low cellular efflux by the ATP-binding cassette (ABC) transporters such as the drug-efflux pump P-glycoprotein (P-gp). It is known that P-gp efflux substrates commonly show poor oral absorption due to the action of P-gp which can transport absorbed drug back into the gut [Ambudkar S.V., et al. (2003) *Oncogene*, 22,
25 7468–7485]. P-gp can also reduce the systemic exposure of substrates through promoting excretion into bile and urine. In addition to being drug efflux pumps, ABC transporters can confer drug resistance to tumour cells through elevated expression. Expression of P-gp is widely known to be a mechanism by which cancer cells can acquire multi-drug resistance (MDR) [Gottesman M.M., et al. (2002) *Nat. Rev. Cancer*,
30 2, 48–58]. Certain tumours derived from cells that normally express P-gp can exhibit intrinsic drug resistance [Ambudkar S.V., et al. (2005) *Trends Pharmacol. Sci.*, 26,

385-387]. Drugs that are readily transported out of cells do not accumulate in high levels in transporter-expressing cells, limiting the success of the therapy due to reduced drug-target interaction.

- 5 Compounds exhibiting low cellular efflux therefore potentially have applications in the oral treatment of cancer, treatment of epithelial cell derived tumours and drug-resistant malignancies, as well as having a lower potential for inducing drug resistance through increased efflux pump expression. Particularly preferred compounds of the invention that exhibit low cellular efflux include compounds [371], [372] and [377]-[391].
- 10 Further details of cell viability assays and cellular efflux studies may be found in the Example 9 and Figure 3.

In addition, compounds of the invention which exhibit low MDR/parental cell asymmetry in cell viability assays are also more effective at inhibiting PLK1 within

15 MDR tumour cells than compounds of the prior art. Thus, these compounds have greater potential as therapeutic PLK inhibitors. Details of experiments measuring PLK1 inhibition in A2780 - A2780/ADR cell line pairs are set forth in the accompanying examples.

20 THERAPEUTIC USE

The compounds of the invention have been found to possess anti-proliferative activity and are therefore believed to be of use in the treatment of proliferative disorders such as cancers, leukaemias and other disorders associated with uncontrolled cellular proliferation such as psoriasis and restenosis.

25

Thus, one aspect of the invention relates to the use of a compound of the invention, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a proliferative disorder.

- 30 As used herein the phrase "preparation of a medicament" includes the use of one or more of the above described compounds directly as the medicament in addition to its

use in a screening programme for further antiproliferative agents or in any stage of the manufacture of such a medicament.

As defined herein, an anti-proliferative effect within the scope of the present invention may be demonstrated by the ability to inhibit cell proliferation in an *in vitro* whole cell assay, for example using any of the cell lines AGS, H1299 or SJSA-1. Using such assays it may be determined whether a compound is anti-proliferative in the context of the present invention.

One preferred embodiment relates to the use of one or more compounds of the invention in the treatment of proliferative disorders. Preferably, the proliferative disorder is a cancer or leukaemia. The term proliferative disorder is used herein in a broad sense to include any disorder that requires control of the cell cycle, for example cardiovascular disorders such as restenosis and cardiomyopathy, auto-immune disorders such as glomerulonephritis and rheumatoid arthritis, dermatological disorders such as psoriasis, anti-inflammatory, anti-fungal, antiparasitic disorders such as malaria, emphysema and alopecia. In these disorders, the compounds of the present invention may induce apoptosis or maintain stasis within the desired cells as required.

In one preferred embodiment of the invention, the proliferative disorder is cancer or leukaemia, more preferably cancer.

In one preferred embodiment, the proliferative disorder is a solid tumour.

In another preferred embodiment, the proliferative disorder is a hematological cancer. Preferably, the haematological cancer is leukaemia, more preferably, advanced leukemias or myelodysplastic syndromes (MDS). Other examples include acute myelogenous leukemia (AML), acute lymphocytic leukemia (ALL) or chronic lymphocytic leukemia (CLL).

30

In another preferred embodiment, the proliferative disorder is glomerulonephritis.

In yet another preferred embodiment, the proliferative disorder is rheumatoid arthritis.

In another preferred embodiment, the proliferative disorder is psoriasis.

- 5 In another preferred embodiment, the proliferative disorder is a chronic obstructive pulmonary disorder.

The compounds of the invention may inhibit any of the steps or stages in the cell cycle, for example, formation of the nuclear envelope, exit from the quiescent phase of the
10 cell cycle (G0), G1 progression, chromosome decondensation, nuclear envelope breakdown, START, initiation of DNA replication, progression of DNA replication, termination of DNA replication, centrosome duplication, G2 progression, activation of mitotic or meiotic functions, chromosome condensation, centrosome separation, microtubule nucleation, spindle formation and function, interactions with microtubule
15 motor proteins, chromatid separation and segregation, inactivation of mitotic functions, formation of contractile ring, and cytokinesis functions. In particular, the compounds of the invention may influence certain gene functions such as chromatin binding, formation of replication complexes, replication licensing, phosphorylation or other secondary modification activity, proteolytic degradation, microtubule binding, actin
20 binding, septin binding, microtubule organising centre nucleation activity and binding to components of cell cycle signalling pathways.

In one embodiment of the invention, the compound of the invention is administered in an amount sufficient to inhibit at least one PLK enzyme.

25

A further aspect of the invention relates to a method of treating a PLK-dependent disorder, said method comprising administering to a subject in need thereof, a compound of the invention or a pharmaceutically acceptable salt thereof, as defined above in an amount sufficient to inhibit PLK.

30

The polo-like kinases (PLKs) constitute a family of serine/threonine protein kinases. Mitotic *Drosophila melanogaster* mutants at the polo locus display spindle abnormalities [Sunkel *et al.*, *J. Cell Sci.*, 1988, 89, 25] and polo was found to encode a mitotic kinase [Llamazares *et al.*, *Genes Dev.*, 1991, 5, 2153]. In humans, there exist
5 three closely related PLKs [Glover *et al.*, *Genes Dev.*, 1998, 12, 3777]. They contain a highly homologous amino-terminal catalytic kinase domain and their carboxyl termini contain two or three conserved regions, the polo boxes. The function of the polo boxes remains incompletely understood but they are implicated in the targeting of PLKs to subcellular compartments [Lee *et al.*, *Proc. Natl. Acad. Sci. USA*, 1998, 95, 9301;
10 Leung *et al.*, *Nat. Struct. Biol.*, 2002, 9, 719], mediation of interactions with other proteins [Kauselmann *et al.*, *EMBO J.*, 1999, 18, 5528], or may constitute part of an autoregulatory domain [Nigg, *Curr. Opin. Cell Biol.*, 1998, 10, 776]. Furthermore, the polo box-dependent PLK1 activity is required for proper metaphase/anaphase transition and cytokinesis [Yuan *et al.*, *Cancer Res.*, 2002, 62, 4186; Seong *et al.*, *J. Biol. Chem.*,
15 2002, 277, 32282].

Studies have shown that human PLKs regulate some fundamental aspects of mitosis [Lane *et al.*, *J. Cell. Biol.*, 1996, 135, 1701; Cogswell *et al.*, *Cell Growth Differ.*, 2000, 11, 615]. In particular, PLK1 activity is believed to be necessary for the functional
20 maturation of centrosomes in late G2/early prophase and subsequent establishment of a bipolar spindle. Depletion of cellular PLK1 through the small interfering RNA (siRNA) technique has also confirmed that this protein is required for multiple mitotic processes and completion of cytokinesis [Liu *et al.*, *Proc. Natl. Acad. Sci. USA*, 2002, 99, 8672].

25 In a more preferred embodiment of the invention, the compound of the invention is administered in an amount sufficient to inhibit PLK1.

Of the three human PLKs, PLK1 is the best characterized; it regulates a number of cell division cycle effects, including the onset of mitosis [Toyoshima-Morimoto *et al.*,
30 *Nature*, 2001, 410, 215; Roshak *et al.*, *Cell. Signalling*, 2000, 12, 405], DNA-damage checkpoint activation [Smits *et al.*, *Nat. Cell Biol.*, 2000, 2, 672; van Vugt *et al.*, *J. Biol.*

Chem., 2001, 276, 41656], regulation of the anaphase promoting complex [Sumara *et al.*, *Mol. Cell*, 2002, 9, 515; Golan *et al.*, *J. Biol. Chem.*, 2002, 277, 15552; Kotani *et al.*, *Mol. Cell*, 1998, 1, 371], phosphorylation of the proteasome [Feng *et al.*, *Cell Growth Differ.*, 2001, 12, 29], and centrosome duplication and maturation [Dai *et al.*,
5 *Oncogene*, 2002, 21, 6195].

In one particularly preferred embodiment, the compounds of the invention are ATP-antagonistic inhibitors of PLK1.

10 In the present context ATP antagonism refers to the ability of an inhibitor compound to diminish or prevent PLK catalytic activity, i.e. phosphotransfer from ATP to a macromolecular PLK substrate, by virtue of reversibly or irreversibly binding at the enzyme's active site in such a manner as to impair or abolish ATP binding.

15 In another preferred embodiment, the compound of the invention is administered in an amount sufficient to inhibit PLK2 and/or PLK3.

A further aspect of the invention relates to a method of inhibiting PLK in a cell comprising contacting said cell with an amount of a compound of formula I, or a
20 pharmaceutically acceptable salt thereof, such that PLK is inhibited in said cell.

Yet another aspect of the invention relates to a method of treating a proliferative disorder comprising inhibiting PLK by administering to a subject in need thereof, a therapeutically effective amount of a compound of formula I, or a pharmaceutically
25 acceptable salt thereof, such that treatment of said proliferative disorder occurs.

In another preferred embodiment, the compound of the invention is administered in an amount sufficient to inhibit at least one aurora kinase. Preferably, the aurora kinase is aurora kinase A, aurora kinase B or aurora kinase C.

A further aspect of the invention relates to a method of treating an aurora kinase-dependent disorder, said method comprising administering to a subject in need thereof, a compound of the invention or a pharmaceutically acceptable salt thereof, as defined above in an amount sufficient to inhibit an aurora kinase.

5

Another aspect relates to the use of a compound of the invention for inhibiting a protein kinase.

A further aspect of the invention relates to a method of inhibiting a protein kinase, said method comprising contacting said protein kinase with a compound of the invention.

10

Preferably, the protein kinase is selected from an aurora kinase and a PLK. More preferably, the protein kinase is a PLK, more preferably PLK1.

15 PHARMACEUTICAL COMPOSITIONS

A further aspect of the invention relates to a pharmaceutical composition comprising a compound of the invention admixed with one or more pharmaceutically acceptable diluents, excipients or carriers. Even though the compounds of the present invention (including their pharmaceutically acceptable salts, esters and pharmaceutically acceptable solvates) can be administered alone, they will generally be administered in admixture with a pharmaceutical carrier, excipient or diluent, particularly for human therapy. The pharmaceutical compositions may be for human or animal usage in human and veterinary medicine.

Examples of such suitable excipients for the various different forms of pharmaceutical compositions described herein may be found in the "Handbook of Pharmaceutical Excipients, 2nd Edition, (1994), Edited by A Wade and PJ Weller.

25

Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A. R. Gennaro edit. 1985).

30

Examples of suitable carriers include lactose, starch, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol and the like. Examples of suitable diluents include ethanol, glycerol and water.

- 5 The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as, or in addition to, the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s).

10

Examples of suitable binders include starch, gelatin, natural sugars such as glucose, anhydrous lactose, free-flow lactose, beta-lactose, corn sweeteners, natural and synthetic gums, such as acacia, tragacanth or sodium alginate, carboxymethyl cellulose and polyethylene glycol.

15

Examples of suitable lubricants include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like.

- 20 Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. Examples of preservatives include sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid. Antioxidants and suspending agents may be also used.

SALTS/ESTERS

- 25 The compounds of the invention can be present as salts or esters, in particular pharmaceutically acceptable salts or esters.

30 Pharmaceutically acceptable salts of the compounds of the invention include suitable acid addition or base salts thereof. A review of suitable pharmaceutical salts may be found in Berge et al, J Pharm Sci, 66, 1-19 (1977). Salts are formed, for example with strong inorganic acids such as mineral acids, e.g. sulphuric acid, phosphoric acid or

hydrohalic acids; with strong organic carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted (e.g., by halogen), such as acetic acid; with saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or tetraphthalic; with hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid; with aminoacids, for example aspartic or glutamic acid; with benzoic acid; or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are unsubstituted or substituted (for example, by a halogen) such as methane- or p-toluene sulfonic acid. Preferably, the salt is an HCl salt.

10

Esters are formed either using organic acids or alcohols/hydroxides, depending on the functional group being esterified. Organic acids include carboxylic acids, such as alkanecarboxylic acids of 1 to 12 carbon atoms which are unsubstituted or substituted (e.g., by halogen), such as acetic acid; with saturated or unsaturated dicarboxylic acid, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or tetraphthalic; with hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid; with aminoacids, for example aspartic or glutamic acid; with benzoic acid; or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are unsubstituted or substituted (for example, by a halogen) such as methane- or p-toluene sulfonic acid. Suitable hydroxides include inorganic hydroxides, such as sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminium hydroxide. Alcohols include alkanealcohols of 1-12 carbon atoms which may be unsubstituted or substituted, e.g. by a halogen).

15

20

25 ENANTIOMERS/TAUTOMERS

In all aspects of the present invention previously discussed, the invention includes, where appropriate all enantiomers and tautomers of the compounds of the invention. The person skilled in the art will recognise compounds that possess an optical properties (one or more chiral carbon atoms) or tautomeric characteristics. The

corresponding enantiomers and/or tautomers may be isolated/prepared by methods known in the art.

STEREO AND GEOMETRIC ISOMERS

5 Some of the compounds of the invention may exist as stereoisomers and/or geometric isomers – e.g. they may possess one or more asymmetric and/or geometric centres and so may exist in two or more stereoisomeric and/or geometric forms. The present invention contemplates the use of all the individual stereoisomers and geometric isomers of those inhibitor agents, and mixtures thereof. The terms used in the claims
10 encompass these forms, provided said forms retain the appropriate functional activity (though not necessarily to the same degree).

The present invention also includes all suitable isotopic variations of the agent or a pharmaceutically acceptable salt thereof. An isotopic variation of an agent of the
15 present invention or a pharmaceutically acceptable salt thereof is defined as one in which at least one atom is replaced by an atom having the same atomic number but an atomic mass different from the atomic mass usually found in nature. Examples of isotopes that can be incorporated into the agent and pharmaceutically acceptable salts thereof include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulphur,
20 fluorine and chlorine such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. Certain isotopic variations of the agent and pharmaceutically acceptable salts thereof, for example, those in which a radioactive isotope such as ^3H or ^{14}C is incorporated, are useful in drug and/or substrate tissue distribution studies. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of
25 preparation and detectability. Further, substitution with isotopes such as deuterium, i.e., ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements and hence may be preferred in some circumstances. Isotopic variations of the agent of the present invention and pharmaceutically acceptable salts thereof of this invention can generally
30 be prepared by conventional procedures using appropriate isotopic variations of suitable reagents.

SOLVATES

The present invention also includes solvate forms of the compounds of the present invention. The terms used in the claims encompass these forms.

5 POLYMORPHS

The invention furthermore relates to the compounds of the present invention in their various crystalline forms, polymorphic forms and (an)hydrous forms. It is well established within the pharmaceutical industry that chemical compounds may be isolated in any of such forms by slightly varying the method of purification and or
10 isolation form the solvents used in the synthetic preparation of such compounds.

PRODRUGS

The invention further includes the compounds of the present invention in prodrug form. Such prodrugs are generally compounds of the invention wherein one or more
15 appropriate groups have been modified such that the modification may be reversed upon administration to a human or mammalian subject. Such reversion is usually performed by an enzyme naturally present in such subject, though it is possible for a second agent to be administered together with such a prodrug in order to perform the reversion in vivo. Examples of such modifications include ester (for example, any of
20 those described above), wherein the reversion may be carried out by an esterase etc. Other such systems will be well known to those skilled in the art.

ADMINISTRATION

The pharmaceutical compositions of the present invention may be adapted for oral,
25 rectal, vaginal, parenteral, intramuscular, intraperitoneal, intraarterial, intrathecal, intrabronchial, subcutaneous, intradermal, intravenous, nasal, buccal or sublingual routes of administration.

For oral administration, particular use is made of compressed tablets, pills, tablets, gellules, drops, and capsules. Preferably, these compositions contain from 1 to 250 mg and more preferably from 10-100 mg, of active ingredient per dose.

- 5 Other forms of administration comprise solutions or emulsions which may be injected intravenously, intraarterially, intrathecally, subcutaneously, intradermally, intraperitoneally or intramuscularly, and which are prepared from sterile or sterilisable solutions. The pharmaceutical compositions of the present invention may also be in form of suppositories, pessaries, suspensions, emulsions, lotions, ointments, creams,
10 gels, sprays, solutions or dusting powders.

An alternative means of transdermal administration is by use of a skin patch. For example, the active ingredient can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin. The active ingredient can
15 also be incorporated, at a concentration of between 1 and 10% by weight, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives as may be required.

Injectable forms may contain between 10 - 1000 mg, preferably between 10 - 250 mg,
20 of active ingredient per dose.

Compositions may be formulated in unit dosage form, i.e., in the form of discrete portions containing a unit dose, or a multiple or sub-unit of a unit dose.

DOSAGE

25 A person of ordinary skill in the art can easily determine an appropriate dose of one of the instant compositions to administer to a subject without undue experimentation. Typically, a physician will determine the actual dosage which will be most suitable for an individual patient and it will depend on a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that
30 compound, the age, body weight, general health, sex, diet, mode and time of

administration, rate of excretion, drug combination, the severity of the particular condition, and the individual undergoing therapy. The dosages disclosed herein are exemplary of the average case. There can of course be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

Depending upon the need, the agent may be administered at a dose of from 0.01 to 30 mg/kg body weight, such as from 0.1 to 10 mg/kg, more preferably from 0.1 to 1 mg/kg body weight.

In an exemplary embodiment, one or more doses of 10 to 150 mg/day will be administered to the patient for the treatment of malignancy.

COMBINATIONS

In a particularly preferred embodiment, the one or more compounds of the invention are administered in combination with one or more other active agents, for example, existing anticancer drugs available on the market. In such cases, the compounds of the invention may be administered consecutively, simultaneously or sequentially with the one or more other active agents.

Anticancer drugs in general are more effective when used in combination. In particular, combination therapy is desirable in order to avoid an overlap of major toxicities, mechanism of action and resistance mechanism(s). Furthermore, it is also desirable to administer most drugs at their maximum tolerated doses with minimum time intervals between such doses. The major advantages of combining chemotherapeutic drugs are that it may promote additive or possible synergistic effects through biochemical interactions and also may decrease the emergence of resistance in early tumor cells which would have been otherwise responsive to initial chemotherapy with a single agent. An example of the use of biochemical interactions in selecting drug combinations is demonstrated by the administration of leucovorin to increase the

binding of an active intracellular metabolite of 5-fluorouracil to its target, thymidylate synthase, thus increasing its cytotoxic effects.


Numerous combinations are used in current treatments of cancer and leukemia. A more
5 extensive review of medical practices may be found in "Oncologic Therapies" edited by
E. E. Vokes and H. M. Golomb, published by Springer.

Beneficial combinations may be suggested by studying the growth inhibitory activity of the test compounds with agents known or suspected of being valuable in the treatment of a particular cancer initially or cell lines derived from that cancer. This procedure can also be used to determine the order of administration of the agents, i.e. before, simultaneously, or after delivery. Such scheduling may be a feature of all the cycle acting agents identified herein.

15 PROCESS

Another aspect relates to a process for preparing a compound of the invention, said process comprising the steps of:

- (i) converting a compound of formula (II) to a compound of formula (IV) either directly or via isolation of a compound of formula (III); and
- 20 (ii) converting said compound of formula (IV) to a compound of formula (I)



25 Preferably, step (i) comprises heating said compound of formula (II) with NH_4Cl and Fe in EtOH/ H_2O .

The present invention is further illustrated by way of the following non-limiting examples, and with reference to the following figures, wherein:

Figure 1 shows the area under the curve (AUC) for the plasma concentration of compounds A-H versus time for each compound tested. The dosing vehicle across these experiments remained constant (intravenous = citrate buffer pH 3, 1 mL/kg; oral = DMA/PEG400/10mM Tartrate buffer, pH 4 (1:3:6), 5 ml/kg);

Figure 2 represents the same data as Figure 1, but in the form of oral bioavailability (% F) for each compound.

Figure 3 shows a comparison of selected compounds versus prior art compounds A'-J' (defined in table 8) in parental and MDR tumor cells. Asymmetry ratios are presented relative to compound [378].

EXAMPLES

General experimental

Chemicals and solvents were purchased from commercial sources and were used as received unless otherwise stated. Anhydrous MgSO_4 was used as a standard drying agent for organic solutions unless otherwise stated. NMR spectra were recorded using a Varian INOVA-500 instrument. Chemical shifts are reported in parts per million relative to internal tetramethyl silane standard. Coupling constants (J) are quoted to the nearest 0.1 Hz. The following abbreviations are used: s, singlet; d, doublet; t, triplet; q, quartet; qu, quintuplet; m, multiplet and br, broad. Mass spectra were obtained using a Waters ZQ2000 single quadrupole mass spectrometer with electro spray ionisation (ESI). Analytical and preparative RP-HPLC was performed using either Vydac 218TP54 (250 x 4.6 mm) and Vydac 218TP1022 (250 x 22mm) columns using a linear gradient elution of water/acetonitrile systems (containing 0.1% trifluoroacetic acid) at flow rates of 1 mL/min (analytical) and 9mL/min (preparative). Analytical gradients used were as follows:

Method	% Acetonitrile Time=0min	% Acetonitrile Time=20min	% Acetonitrile Time=25min
Vydac 1	0	60	100
Vydac 2	10	70	100
Vydac 3	20	80	100

Alternatively Xbridge (100 x 4.6 mm) and Xbridge (100 x 19mm) columns using a linear gradient of water/ acetonitrile systems (containing 0.1% ammonium hydroxide) at flow rates of 1 mL/min (analytical) and 20mL/min (preparative).

Analytical gradients used were as follows:

Method	% Acetonitrile Time=0min	% Acetonitrile Time=5min	% Acetonitrile Time=10min	% Acetonitrile Time=12min
XBridge1	0		50	100
XBridge2	10			100
XBridge3	20	70		100

Silica gel (EM Kieselgel 60, 0.040-0.063 mm, Merck) or ISOLUTE pre-packed columns (Biotage) were used for flash chromatography.

Abbreviations:

DIPEA diisopropylethyl amine
 DCM dichloromethane
 DMF dimethylformamide
 DMSO dimethyl sulfoxide
 EtOAc ethyl acetate
 EtOH ethanol
 HCl hydrochloric acid
 K₂CO₃ potassium carbonate
 MeI methyl iodide
 MeOH methanol
 MgSO₄ magnesium sulfate
 NaH sodium hydride

NaCl sodium chloride

NH₄Cl ammonium chloride

RM reaction mixture

rt room temperature

5 TBTU O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate


TFA trifluoroacetic acid

TFE 2,2,2-trifluoroethanol

THF tetrahydrofuran

10 EXAMPLE 1

Compound [2]: 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide



15

1. Thionyl chloride (2.1 eq), MeOH, 0 °C for addition, reflux 2h; 2. cyclopentanone (0.77 eq), sodium acetate (0.77 eq), sodium triacetoxyborohydride (1.11 eq), DCM, rt 16h; 3. 2,4-dichloro-5-nitropyrimidine (1.1 eq), K₂CO₃ (1 eq), acetone, 0 °C – rt 16h; 4. NH₄Cl (8.5 eq), Fe (8 eq), EtOH/H₂O (4:1), reflux 2.5h; - not isolated 5. MeI (1.18 eq), NaH (1.07 eq), DMF, -10 °C – rt, 3h; 6. 4-amino-3-methoxybenzoic acid (1.5 eq), conc.

20

HCl, H₂O/EtOH (4:1), reflux 48h; 7. DIPEA (2 eq), TBTU (1.1 eq), 4-aminomethylpiperidine (1.2 eq), DCM, rt 16h.

Step 1: Methyl 3-aminopropanoate

β-Alanine (9.37 g, 0.105 mol), was added to MeOH (50 ml) and the mixture cooled to
5 0 °C using an ice-bath before the dropwise addition of thionyl chloride [caution: exothermic addition]. Once the addition was complete the reaction was warmed to rt, then refluxed for 2h. The solvent was next evaporated under reduced pressure and the resulting oil treated with t-butylmethyl ester and the resulting crystals filtered off and further dried *in vacuo* to give a white crystalline solid (11g, quant.); ¹H (DMSO-*d*₆): δ
10 2.73 (dd, *J* = 7 Hz, 2H, CH₂), 2.98 (dd, *J* = 7 Hz, 2H, CH₂), 3.61 (s, 3H, CH₃), 8.28 (bs, 2H, NH₂); MS+ve: 104.1.

Step 2: Methyl 3-(cyclopentylamino)propanoate

Methyl 3-aminopropanoate (9.37 g, 0.09 mol) was solubilised in DCM (200 ml) and
15 cyclopentanone (6.43 ml, 0.07 mol), sodium acetate (5.96 g, 0.07 mol) and sodium triacetoxyborohydride (22 g, 0.10 mol) were added. The reaction was stirred at rt for 16h. 20% sodium bicarbonate (100 ml) and 2M sodium hydroxide (50 ml) were then added and the product extracted using DCM/H₂O. The organic extracts were combined, washed with sat NaCl, dried using MgSO₄, filtered and the filtrate evaporated under
20 reduced pressure and further dried *in vacuo* to give the product as a pale yellow oil (8.90 g, 55%); MS+ve: 172.4.

Step 3: Methyl 3-[cyclopentyl(2-chloro-5-nitropyrimidin-4-yl)amino]propanoate

Methyl 3-(cyclopentylamino)propanoate (838 mg, 0.005 mol) and K₂CO₃ (676 mg,
25 0.005mol) were added to acetone (5 ml) and the resulting mixture cooled to 0 °C using an ice-bath before the addition of 2,4-dichloro-5-nitropyrimidine (1.044 g, 1.1 eq). The reaction mixture (RM) was then warmed to rt and stirring continued for an additional 16h before the addition of a further 0.12 eq of the pyrimidine. Stirring was then continued for a further 3h. The RM was then evaporated under reduced pressure and
30 the product extracted using EtOAc/H₂O. The organic extracts were combined, washed

with sat NaCl, dried using MgSO₄, filtered and the filtrate evaporated under reduced pressure and further dried *in vacuo* to give the product as a brown oily residue (1.04g, 65%); R_t = 16 min (Vydac 1); ME+ve: 329.1.

5 *Step 4: 2-Chloro-9-cyclopentyl-5,7,8,9-tetrahydro-6H-pyrimido[4,5-b][1,4]diazepin-6-one*

Methyl 3-[cyclopentyl(2-chloro-5-nitropyrimidin-4-yl)amino]propanoate (1.00 g, 0.003 mol) and NH₄Cl (1.38 g, 0.025 mol, 8.5 eq) were added to EtOH/H₂O (4:1 ml, 10 ml) and the mixture heated to reflux before the portion-wise addition of iron powder (1.36 g, 10 0.024 mol, 8 eq). The RM was then refluxed for an additional 2 h. Progress of the reaction was monitored by HPLC and when no SM remained the RM was filtered hot through celiteTM. The celiteTM was washed with EtOAc (10 ml) and EtOH (10 ml) [both hot] and the filtrate evaporated under reduced pressure and further dried *in vacuo* to give the product as a brown solid (350 mg, 43%); R_t = 12 min (Vydac 1); MS+ve: 15 267.2.

Step 5: 2-Chloro-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-6H-pyrimido[4,5-b][1,4]diazepin-6-one

2-Chloro-9-cyclopentyl-5,7,8,9-tetrahydro-6H-pyrimido[4,5-b][1,4]diazepin-6-one 20 (318 mg, 0.0012 mol) and MeI (88 µl, 0.0014 mol, 1.18 eq) were added to DMF (5 ml) and the solution cooled to -10 °C using acetone/dry ice before the addition of NaH (30 mg, 0.0013 mol, 1.07 eq). The RM was then stirred at 0 °C for 30 min and rt for 30 min. The RM was concentrated and the product extracted using EtOAc/H₂O. The organic extracts were combined, washed with sat NaCl, dried using MgSO₄, filtered and the 25 filtrate evaporated under reduced pressure and further dried *in vacuo* to give the product as a purple oily residue (252 mg, 75%); R_t = 13 min (Vydac 1); MS+ve: 281.2.

Step 6: (Compound [1]) 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid.

Coupling Method A

5 2-Chloro-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-6H-pyrimido[4,5-b][1,4]diazepin-6-one (248 mg, 0.0009 mol) and 4-amino-3-methoxybenzoic acid (221 mg, 0.0013 mol, 1.5 eq), conc. HCl (152 μ l) and H₂O/EtOH (8:2 ml) were added to a RBF and the resulting RM heated to reflux for 4h. Progress of the reaction was monitored by HPLC and when no starting material remained (4h reaction time), the RM was concentrated
10 under reduced pressure and the product extracted using DCM/H₂O. The organic extracts were combined, washed with sat NaCl, dried using MgSO₄, filtered and the filtrate evaporated under reduced pressure and further dried *in vacuo* to give a brown oily residue. A few drops of MeOH were added to the residue and the solid ppt formed was collected by suction filtration, washed with MeOH and further dried *in vacuo* to
15 give the product as a purple solid (51 mg, 14%); R_t = 11.2 min (0_60_20 min, purity 100%); ¹H NMR (DMSO-*d*₆): δ 1.61 (bs, 4H, cyclopent-*H*), 1.72 (bs, 2H, cyclopent-*H*), 1.94 (bs, 2H, cyclopent-*H*), 2.59 (dd, J = 4.5 Hz, 2H, CH₂), 3.18 (s, 3H, CH₃), 3.63 (dd, J = 4.5 Hz, 2H, CH₂), 3.95 (s, 3H, CH₃), 7.51 (s, 1H), 7.56 (d, J = 8 Hz, 1H, phe-*H*), 7.82 (s, 1H, 8.10 (s, 1H, 8.47 (d, J = 8.5 Hz, 1H, phe-*H*), 12.64 (bs, 1H, NH); MS+ve:
20 412.2.

Step 7: (Compound [2]): 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide
4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-
25 2-ylamino)-benzoic acid (35 mg, 0.085 mmol), DIPEA (28 μ l, 0.17 mmol, 2 eq) and TBTU (30 mg, 0.093 mmol, 1.1 eq) were added to 3 ml DCM and the resulting solution stirred at rt for 30 min before the addition of 4-aminomethylpiperidine (13 μ l, 0.10 mmol, 1.2 eq). The RM was then stirred at rt for 16h. The RM was concentrated and the product extracted using DCM/H₂O. The organic extracts were combined, washed
30 with sat NaCl, dried using MgSO₄, filtered and the filtrate evaporated under reduced pressure and further dried *in vacuo* to give a yellow oily residue (21 mg, 49%); R_t =

47


9.86 min (0_60_20 min, purity 100%); ^1H NMR (CD_3OD): δ 1.69 – 1.75 (m, 6H), 1.82 (bs, 2H), 1.96 – 1.99 (m, 2H), 2.03 – 2.05 (bs, 2H), 2.18 – 2.23 (m, 2H), 2.34 (s, 3H, CH_3), 2.68 (dd, $J = 4.5$ Hz, 2H, CH_2), 2.95 – 2.97 (m, 2H), 3.28 (s, 3H, CH_3), 3.73 (dd, $J = 4.5$ Hz, 2H, CH_2), 3.89 – 3.94 (m, 1H, CH), 4.01 (s, 3H, CH_3), 4.91 – 4.96 (m, 1H, CH), 7.49 – 7.51 (m, 2H), 8.02 (bs, 1H), 8.50 (d, $J = 8.5$ Hz, 1H, phe- H); MS+ve: 508.2.

Aminoester intermediates

The following intermediates were prepared by the method described in Example 1, step


10 2:

3-Cyclopentylamino-butyric acid methyl ester




MS+ve: 186.3

15 *3-Cyclohexylamino-butyric acid methyl ester*



MS+ve: 200.3

3-Cyclopentylamino-2-methyl-propionic acid methyl ester




20

^1H NMR (CDCl_3): 1.18 (3H, d, J 7Hz, CH_3), 1.33 (2H, m, CH_2), 1.53 (2H, m, CH_2), 1.67 (2H, m, CH_2), 1.82 (2H, m, CH_2), 2.63 (2H, m, CH_2), 2.87 (1H, m, CH), 3.06 (1H, m, CH), 3.69 (3H, s, OCH_3)


25

3-Cyclohexylamino-propionic acid methyl ester


48



MS+ve: 186.4.


3-Cyclopentylamino-4-methyl-pentanoic acid methyl ester

¹H NMR (CDCl₃): 1.18 (3H, d, J 7Hz, CH₃), 1.33 (2H, m, CH₂), 1.53 (2H, m, CH₂), 1.67 (2H, m, CH₂), 1.82 (2H, m, CH₂), 2.63 (2H, m, CH₂), 2.87 (1H, m, CH), 3.06 (1H, m, CH), 3.69 (3H, s, OCH₃)


10 *3-(1-Ethyl-propylamino)-propionic acid methyl ester*

¹H NMR (CDCl₃): 0.89 (6H, t, J 8Hz, CH₃), 1.44 (4H, m, CH₂CH₃), 1.77 (1H, bs, NH), 2.39 (1H, m, CH), 2.53 (2H, m, CH₂), 2.89 (2H, m, CH₂), 3.69 (3H, s, OCH₃)

15

3-(Tetrahydro-pyran-4-ylamino)-propionic acid methyl ester


¹H NMR (CDCl₃): 1.41 (2H, m, CH₂), 1.86 (2H, m, CH₂), 2.54 (2H, m, CH₂), 2.71 (1H, m, CH), 2.95 (2H, m, CH₂), 3.44 (2H, t, J 11.5Hz, CH₂), 3.71 (3H, s, OCH₃), 3.99 (2H, d, J 11Hz, CH₂)

1-Cyclopentylaminomethyl-cyclopropanecarboxylic acid methyl ester

49

¹H NMR (CDCl₃): 0.84 (3H, t, J 7Hz, CH₃), 1.25 (4H, m, CH₂), 1.37 (2H, m, CH), 1.54 (2H, m, CH), 1.70 (2H, m, CH), 1.83 (2H, m, CH), 2.71 (2H, s, CH₂), 3.10 (1H, m, CH), 4.16 (2H, q, J 7Hz, CH₂CH₃)

5 *3-Cyclopentylamino-2,2-dimethyl-propionic acid ethyl ester*



¹H NMR (CDCl₃): 1.19 (6H, s, CH₃), 1.25 (3H, t, J 7Hz, CH₃), 1.28 (2H, m, CH), 1.49 (2H, m, CH), 1.65 (2H, m, CH), 2.64 (1H, s, CH₂), 3.01 (1H, m, CH), 4.10 (2H, q, J 7.5Hz, CH₂)

10


Alternative method Step 2a: Methyl 3-aminopropanoate



15 Methyl acrylate (4.50 ml, 4.302 g, 49.97 mmole, 0.99 eq) was added dropwise to a solution of cyclopentylamine (5.00 ml, 4.315 g, 50.68 mmole) in methanol (120 ml) at -60°C (dry ice, acetone). The reaction mixture was allowed to warm to room temperature and stirred overnight. The solvent was removed (reduced pressure, vacuo) to afford the product as an oil (8.236 g, 96% crude yield). Proton NMR indicated that
20 the product contained ~25% of a dialkylated byproduct.

The following compounds were prepared by a similar method.

3-Phenylamino-propionic acid ethyl ester



25


¹H NMR (CDCl₃): 1.29 (3H, t, J 7.5Hz, CH₃), 2.64 (2H, t, J 6.5Hz, CH₂), 3.48 (2H, t, J 6.5Hz, CH₂), 4.18 (2H, q, J 7Hz, CH₂), 6.65 – 6.74 (5H, m, Ar-H)

50

Nitropyrimidine intermediates

The following intermediates were also prepared by the method described in Example 1 step 3:


3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-butyric acid methyl ester



5


$R_t = 18.0$ min (0_60_20 min); MS+ve: 343.20.

3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclohexyl-amino]-butyric acid methyl ester




10 MS+ve: 357.2.

3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-2-methyl-propionic acid methyl ester



15 ^1H NMR (CDCl_3): 1.22 (3H, d, J 12Hz, CH_3), 1.57 (2H, m, CH_2), 1.76 (2H, m, CH_2), 1.93 (2H, m, CH_2), 1.98 (2H, m, CH_2), 3.18 (1H, s, CH), 3.61 (3H, m, CH + CH_2), 3.68 (3H, s, OCH_3), 8.74 (1H, s, Pyr-H)

3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclohexyl-amino]-propionic acid methyl ester




20

51

Rt = 17.6 min (Vydac 1); MS+ve: 343.1.


3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-4-methyl-pentanoic acid methyl ester



5


¹H NMR (CDCl₃): 0.94 (6H, t, J 7Hz), 1.45 – 1.82 (7H, m, 3 x CH₂ + CH), 2.17 (1H, m, CH), 2.69 (2H, m, 2 x CH), 3.35 (2H, m, CH₂), 2.58 (1H, m, CH), 3.73 (3H, s, OCH₃), 8.64 (1H, s, Pyr-H)

10 *3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-(1-ethyl-propyl)-amino]-propionic acid methyl ester*




MS (+ve) 331.1, 333.1

15 *3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-(tetrahydro-pyran-4-yl)-amino]-propionic acid methyl ester*



MS (+ve) 345.1, 347.0; tR = 12.62 min (Vydac 1).


3-[Benzyl-(2-chloro-5-nitro-pyrimidin-4-yl)-amino]-propionic acid ethyl ester



20


¹H NMR (CDCl₃): 1.26 (3H, t, J 7Hz, CH₃), 2.73 (2H, t, J 7Hz, CH₂), 3.85 (2H, m, CH₂), 4.16 (2H, q, J 7Hz, CH₂), 4.71 (2H, s, CH₂), 7.20 (2H, m, Ar-H), 7.37 (3H, m, Ar-H), 8.69 (1H, s, Pyr-H).

5 *3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-phenyl-amino]-propionic acid ethyl ester*




¹H NMR (CDCl₃): 1.22 (3H, t, J 7.5Hz, CH₃), 2.73 (2H, t, J 7Hz, CH₂), 4.09 (2H, q, J 7Hz, CH₂), 4.43 (2H, t, J 7Hz, CH₂), 7.14 – 7.48 (5H, m, Ar-H), 8.57 (1H, s, Pyr-H).

10 *1-{[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-methyl} cyclopropanecarboxylic acid ethyl ester*



MS (+ve) 381.1, 383.3; tR = 4.80 min (XBridge 1).

15 *3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-2,2-dimethyl-propionic acid ethyl ester*




MS (+ve) 371.1, 373.1; tR = 4.67 min (XBridge 1).

20 Aminopyrimidine-esters leading to chloropyrimidine-diazepines

The following compounds were isolated from the reduction reaction as shown in Example 1, Step 4 as uncyclised intermediates.

3-[(5-Amino-2-chloro-pyrimidin-4-yl)-benzyl-amino]-propionic acid ethyl ester


53



^1H NMR (CDCl_3): 1.26 (3H, t, J 7.5Hz, CH_3), 2.73 (2H, t, J 7Hz, CH_2), 3.86 (2H, t, J 7Hz, CH_2), 4.14 (2H, q, J 7Hz, CH_2), 4.86 (2H, s, CH_2), 7.26 – 7.39 (5H, m, Ar-H), 7.94 (1H, s, Pyr-H)


5

3-[(5-Amino-2-chloro-pyrimidin-4-yl)-phenyl-amino]-propionic acid ethyl ester




MS (+ve) 321.2, 323.2; tR = 3.54 min (XBridge 1).

10 1-{[(5-Amino-2-chloro-pyrimidin-4-yl)-cyclopentyl-amino]-methyl}-cyclopropanecarboxylic acid ethyl ester



MS (+ve) 339.2, 341.2; tR = 4.15 min (XBridge 1).

15 3-[(5-Amino-2-chloro-pyrimidin-4-yl)-cyclopentyl-amino]-2,2-dimethyl-propionic acid ethyl ester




MS (+ve) 341.2, 343.2; tR = 4.25 min (XBridge 1).

The above compounds were then cyclised using the following method.

20 The relevant compound (5 mmol) in DMF (10 ml) was heated to 140°C for 2 hours. Solvent was evaporated *in vacuo*, ethyl acetate (10 ml) added, and the resulting solid filtered and dried under vacuum to give the compounds listed below.

Note: This reaction may also be carried out with DMSO/NaH


9-Benzyl-2-chloro-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one



5

^1H NMR (DMSO): 2.72 (2H, m, CH₂), 3.67 (2H, m, CH₂), 4.86 (2H, s, CH₂), 7.26 – 7.37 (5H, m, Ar-H), 7.88 (1H, s, Pyr-H), 9.77 (1H, s, NH)


2-Chloro-9-phenyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one



10

MS (+ve) 275.2, 277.2; tR = 2.80 min (XBridge 1).


2-Chloro-9-cyclopentyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-7,1'-cyclopropan]-6-one



15

MS (+ve) 293.2, 295.2; tR = 3.48 min (XBridge 1).

2-Chloro-9-cyclopentyl-7,7-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one




20

MS (+ve) 295.1, 297.2; tR = 3.58 min (XBridge 1).

Alkylated diazepinones


The following intermediates were also prepared by the method described in Example 1, step 5:

- 5 *2-Chloro-9-cyclopentyl-5,8-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-*b*][1,4]diazepin-6-one*




$R_t = 14.6$ min (Vydac 1); MS+ve: 295.2.

- 10 *2-Chloro-9-cyclopentyl-5,7-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-*b*][1,4]diazepin-6-one*




MS (+ve) 295.1, 297.2; $t_R = 9.11$ min (XBridge 2).

- 15 *2-Chloro-9-cyclopentyl-8-isopropyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-*b*][1,4]diazepin-6-one*




MS (+ve) 323.1, 325.1; $t_R = 10.17$ min (XBridge 2).

- 20 *2-Chloro-9-cyclopentyl-5-ethyl-7-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-*b*][1,4]diazepin-6-one*




¹H NMR (CDCl₃): 1.08 (6H, m, 2 x CH₃), 1.38 (1H, m, CH), 1.55 – 1.70 (5H, m, CH + CH₂), 1.70 (1H, m, CH), 2.10 (1H, m, CH), 2.73 (1H, m, CH), 3.48 (2H, m, CH₂), 4.03 (1H, m, CH), 4.72 (1H, m, CH), 7.98 (1H, s, Pyr-H)

- 5 *2-Chloro-9-(1-ethyl-propyl)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one*




MS (+ve) 283.2, 285.1; tR = 8.44 min (XBridge 2).

- 10 *2-Chloro-9-(1-ethyl-propyl)-5-ethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one*




MS (+ve) 297.12, 299.14; tR = 3.67 min (XBridge 2).

- 15 *2-Chloro-5-methyl-9-(tetrahydro-pyran-4-yl)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one*



MS (+ve) 297.1, 299.2; tR = 5.74 min (XBridge 2).


- 20 *9-Benzyl-2-chloro-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one*



MS (+ve) 303.1, 305.2; tR = 13.57 min (Vydac 1).


2-Chloro-5-methyl-9-phenyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one

57




MS (+ve) 289.1, 291.1; tR = 2.93 min (XBridge 2).

5 *2-Chloro-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-7,1'-cyclopropan]-6-one*



MS (+ve) 307.1, 309.2; tR = 3.67 min (XBridge 2).

2-Chloro-9-cyclopentyl-5,7,7-trimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one



10


MS (+ve) 309.1, 311.2; tR = 3.90 min (XBridge 2).

Coupled pyrimidinediazepinones

The following were also prepared by the method described in Example 1, step 6,

15 Coupling Method A:

Compound [3]: 4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid




20 Purple solid (29%) ¹H NMR (DMSO-*d*₆): δ 1.33 (d, *J* = Hz, 3H, CH₃), 1.52 – 1.58 (m, 3H, cyclopent-*H*), 1.70 – 1.78 (m, 3H, cyclopent-*H*), 1.87 - 1.93 (m, 2H, cyclopent-*H*), 2.97 – 3.00 (m, 2H, CH₂), 3.94 (s, 3H, CH₃), 4.15 – 4.17 (m, 1H, CH), 4.93 – 4.97 (m,

58

1H, CH), 7.59 – 7.60 (m, 2H), 7.82 (t, 1H), 8.01 (d, $J = 9$ Hz, 1H, phe-*H*), 9.55 (bs, 1H, OH), 9.96 (s, 1H); MS+ve: 426.2. $R_t = 11.08$ min (purity 100%, (Vydac 1))


5 *Compound [4]: 4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-*b*][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid*



Purple solid (67%). ^1H NMR (DMSO- d_6): δ 1.26 (d, $J = \text{Hz}$, 3H, CH₃), 1.43 – 1.93 (m, 8H, cyclopent-H), 2.52 (s, 3H, CH₃), 2.99 – 3.02 (m, 2H, CH₂), 3.96 (s, 3H, CH₃), 4.13 – 4.16 (m, 1H, CH), 4.65 – 4.69 (m, 1H, CH), 7.60 – 7.61 (m, 2H), 8.06 (d, $J = 9$ Hz, 1H, phe-H), 8.17 (s, 1H), 9.35 (bs, 1H); MS+ve: 426.2. $R_t = 11.59$ min (purity 96%, (Vydac 1))

15


*Compound [5]: 4-(9-Cyclohexyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-*b*][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid*



Brown solid (12%); ^1H NMR (DMSO- d_6): δ 1.27(d, $J = 6.5$ Hz, 3H, CH₃), 1.58 – 1.87 (m, 10H, cyclohex-H), 2.85 – 2.88 (m, 2H, CH₂), 3.94 (s, 3H, CH₃), 4.21 – 4.23 (m, 1H, CH), 4.77 (bs, 1H, CH), 7.03 (s, 1H), 7.13 (s, 1H), 7.23 (s, 1H), 7.57 – 7.58 (m, 2H), 7.086 (s, 1H), 8.07 – 8.08 (m, 1H), 9.89 (bs, 1H, NH); ME+ve: 426.2. $R_t = 11.58$ min (purity 90%, (Vydac 1))


*Compound [6]: 4-(9-Cyclopentyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-*b*][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid*

59




Purple solid (13%); ^1H NMR ($\text{DMSO}-d_6$): δ 1.55 – 1.86 (m, 8H, cyclopent-*H*), 2.70 – 2.73 (m, 2H, CH_2), 3.70 – 3.72 (m, 2H, CH_2), 5.02 – 5.09 (m, 1H, *CH*), 7.57 – 7.59 (m, 2H), 7.77 (s, 1H, pyrimid-*H*), 8.06 (d, $J = 8.5$ Hz, 1H, phe-*H*), 9.54 (bs, 1H, OH), 9.75 (s, 1H, NH); MS+ve: 398.2. $R_t = 10.55$ min (purity 96%, (Vydac 1))

Compound [7]: 4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid




Off-white solid (10%); ^1H NMR ($\text{DMSO}-d_6$): δ 1.27 (d, $J = 6.5$ Hz, 3H, CH_3), 1.46 – 1.90 (m, 8H, cyclopent-*H*), 2.83 – 2.86 (m, 2H, CH_2), 4.07 – 4.09 (m, 1H, *CH*), 5.04 – 5.09 (m, 1H, *CH*), 7.20 (dd, $J = 3.5$ and 8.5 Hz, 2H, phe-*H*), 7.79 (d, $J = 2.0$ Hz, 1H, pyrimid-*H*), 7.86 (d, $J = 8.5$ Hz, 2H, phe-*H*), 9.78 (bs, 1H, NH); MS+ve: 382.2. $R_t = 10.46$ min (purity 90%, (Vydac 1))

Compound [8]: 4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid




Off-white solid (25%); ^1H NMR ($\text{DMSO}-d_6$): δ 1.19 (d, $J = 6.0$ Hz, 3H, CH_3), 1.32 – 2.10 (m, 8H, cyclopent-*H*), 2.73 – 2.76 (m, 2H, CH_2), 3.22 (s, 3H, CH_3), 4.04 (m, 1H, *CH*), 4.70 – 4.75 (m, 1H, *CH*), 7.78 – 7.85 (m, 4H, phe-*H*), 8.12 (s, 1H, pyrimid-*H*), 9.68 (bs, 1H, NH); MS+ve: 394.0. $R_t = 11.05$ min (purity 96%, (Vydac 1))

Compound [9]: 4-(9-Cyclohexyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid




- 5 Brown solid (15%); ^1H NMR (DMSO- d_6): δ 1.15 – 1.86 (m, 10H, cyclohex- H), 2.72 (dd, $J = 4.5$ Hz, 2H, CH_2), 3.75 (dd, $J = 4.5$ Hz, 2H, CH_2), 3.96 (s, 3H, CH_3), 4.58 – 4.63 (m, 1H, CH), 7.58 – 7.60 (m, 2H), 7.79 (s, 1H, pyrimid- H), 8.16 (d, $J = 8.5$ Hz, 1H, phe- H), 9.74 (bs, 1H, NH); MS+ve: 412.2. $R_t = 11.19$ min (purity 100%, (Vydac 1))

10 *Compound [10]: 9-Cyclohexyl-2-(4-hydroxy-phenylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one*



- Purple solid (7%); ^1H NMR (DMSO- d_6): δ 1.17 – 1.87 (m, 10H, cyclohex- H), 2.72 (bs, 2H, CH_2), 3.72 (bs, 2H, CH_2), 4.62 – 4.66 (m, 1H, CH), 6.82 (d, $J = 8.5$ Hz, 2H, phe- H),
15 7.13 (s, 1H), 7.23 (s, 1H), 7.34 – 7.37 (m, 3H, 2 x phe- H and NH); MS+ve: 354.2. $R_t = 10.63$ min (purity 100%, (Vydac 1))

Compound [11]: 9-Cyclopentyl-2-(4-hydroxy-phenylamino)-7-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one



20


^1H NMR (CD $_3$ OD): 1.18 (3H, d, J 7Hz, CH_3), 1.52 (1H, m, CH), 1.61 – 1.84 (6H, m, CH), 1.99 (1H, m, CH), 2.85 (1H, m, CH), 3.53 (2H, m, CH), 5.14 (1H, m, CH), 6.74

61

(2H, m, Ar-H), 7.34 (2H, m, Ar-H), 7.64 (1H, s, Pyr-H) ; MS(+ve): 354.2; tR = 10.85 min (Vydac 1).

Compound [12]: 9-Cyclopentyl-2-(4-hydroxy-phenylamino)-8-isopropyl-5-methyl-


5 *5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one*



¹H NMR (DMSO): 0.72 (3H, d, J 7Hz, CHCH₃), 0.85 (2H, d, J 7Hz, CHCH₃), 1.29 (1H, m, CH), 1.51 (4H, m, CH), 1.60 (2H, m, CH), 1.93 (1H, m, CH), 2.08 (1H, m, CH), 2.29 (2H, m, 2 x CH), 3.13 (3H, s, N-CH₃), 3.82 (1H, m, CH), 4.15 (1H, m, CH), 6.65 (2H, d, J 6Hz, Ar-H), 7.47 (2H, d, J 6Hz, Ar-H), 8.14 (1H, s, Pyr-H), 8.99 (1H, s, NH); MS(+ve): 396.1; tR = 12.54 min (Vydac 1).

Compound [13]: 9-Cyclopentyl-2-(4-hydroxy-phenylamino)-5,8-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one


15



White solid (27%); ¹H NMR (DMSO-*d*₆): δ 1.22 (d, J = 6.5 Hz, 3H, CH₃), 1.38 – 1.85 (m, 8H, cyclopent-H), 2.49 – 2.53 (m, 2H, CH₂), 3.18 (s, 3H, CH₃), 4.05 – 4.10 (m, 1H, CH), 4.51 – 4.60 (m, 1H, CH), 6.80 (d, J = 9.0 Hz, 2H, phe-H), 7.22 (d, J = 9.0 Hz, 2H, phe-H), 7.98 (s, 1H, pyrimid-H), 9.53 (bs, 1H, OH), 10.05 (bs, 1H, NH); MS+ve: 368.3. R_t = 10.81 min (purity 93%, (Vydac 1))

Compound [14]: 4-(9-Benzyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid


62



¹H NMR (DMSO): 2.72 (2H, m, CH₂), 3.68 (2H, m, CH₂), 3.90 (3H, s, OCH₃), 4.88 (2H, s, CH₂), 7.22 – 7.44 (6H, m, Ar-H), 7.44 (1H, s, Ar-H), 7.77 (1H, s, Pyr-H), 8.10 (1H, d, J 7Hz), 8.18 (1H, s, NH); MS(+ve): 434.2; tR = 10.97 min (Vydac 1).


5

Compound [15]: 4-(5-Methyl-6-oxo-9-phenyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid



¹H NMR (DMSO): 2.84 (2H, m, CH₂), 3.28 (3H, s, CH₃), 4.08 (2H, m, CH₂), 7.24 (2H, d, J 8.5Hz, Ar-H), 7.35 – 7.37 (3H, m, Ar-H), 7.47 – 7.52 (4H, m, Ar-H), 8.29 (1H, s, Pyr-H), 9.77 (1H, s, NH); MS(+ve): 390.2; tR = 1.93 min (XBridge 2).

Compound [16]: 4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid




¹H NMR (DMSO): 1.09 (6H, s, CH₃), 1.60 (4H, m, CH), 1.74 (2H, m, CH), 1.88 (2H, m, CH), 3.19 (3H, s, N-CH₃), 3.37 (2H, s, CH₂), 5.23 (1H, s, CH), 7.82 (4H, m, Ar-H), 7.99 (1H, s, Pyr-H), 9.57 (1H, s, NH); MS(+ve): 410.25; tR = 2.25 min (XBridge 2).

20 Preparation and reaction of amino-benzamides

4-Amino-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide

63




To 4-amino-3-methoxybenzoic acid (1.064 g mg, 6.37 mmol) in dichloromethane (50 ml) was added DIPEA (2.22 ul, 12.74 mmol) and TBTU (2.25 g, 7.0 mmol) and the mixture stirred for 10 minutes. 4-amino-1-methylpiperidine (0.872 g, 7.65 mmol) was added and stirring continued for 16 hours. The resulting solid was filtered and dried under vacuum (1.04 g, 3.95 mmol, 62%).

¹H NMR (DMSO): 1.54 (2H, m, CH), 1.68 (2H, m, CH), 1.90 (2H, m, CH), 2.15 (3H, s, N-CH₃), 2.76 (2H, d, J 11.5Hz, CH), 3.66 (1H, m, CH), 3.77 (3H, s, OCH₃), 6.56 (1H, m, Ar-H), 7.26 (1H, m, Ar-H), 7.75 (1H, d, J 7.5Hz); MS (+ve) 264.4; tR = 7.45 min (Vydac 2).

The following amino-benzamides were prepared by a similar method:

15 *4-Amino-N-(1-methyl-piperidin-4-yl)-benzamide* via [4-(1-Methyl-piperidin-4-ylcarbamoyl)-phenyl]-carbamic acid tert-butyl ester




[4-(1-Methyl-piperidin-4-ylcarbamoyl)-phenyl]-carbamic acid tert-butyl ester ¹H NMR (DMSO): 1.44 (9H, s, C(CH₃)₃), 1.52 (2H, m, CH), 1.71 (2H, m, CH), 1.92 (2H, m, CH), 1.92 (3H, s, N-CH₃), 2.74 (2H, m, CH), 3.68 (1H, m, N-CH), 7.47 (2H, d, J 8.5Hz, Ar-H), 7.72 (2H, d, J 8.5Hz, Ar-H), 8.00 (1H, d, J 7.5Hz, NH), 9.54 (1H, s, NH)

This compound was used without prior deprotection when using Coupling Method B (below)

Coupling Method B

Compound [17]: 4-(9-Cyclopentyl-5,7-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



5

The aniline (149 mg, 0.57 mmol), pyrimidine (53 mg, 0.19 mmol) and TFA (109 μ l, 0.95 mmol) in 2,2,2-trifluoroethanol (5 ml) were heated to reflux for 18 hours. Solvent was evaporated, Ethyl acetate (10 ml) added, washed with sat. NaHCO₃, brine and the solvent evaporated. Ethyl acetate (10 ml) was added and the resulting solid filtered and dried under vacuum (30 mg, 0.06 mmol, 30%).

10


¹H NMR (DMSO): 1.06 (3H, d, J 7Hz, CHCH₃), 1.52 – 1.78 (10H, m, alkyl CH + CH₂), 2.01 (2H, m, CH), 2.36 (3H, s, N-CH₃), 2.73 (1H, m, CH), 2.85 (2H, m, CH), 3.45 (1H, m, CH), 3.71 (1H, m, CH), 3.94 (3H, s, OCH₃), 5.06 (1H, m, CH), 7.46 (2H, m, Ar_h), 7.61 (1H, s, Ar-H), 7.77 (1H, s, NH), 8.10 (1H, d, J 8Hz, Ar-H), 8.36 (1H, d, J 8Hz), Ar-H), 9.44 (1H, s, NH); MS(+ve): 508.2; tR = 7.30 min (XBridge 2).

15

The following compounds were prepared in a similar manner to the above example:

Compound [18]: 4-(9-Cyclopentyl-7-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide

20




¹H NMR (DMSO): 1.08 (3H, d, J 7Hz, CHCH₃), 1.53 – 1.80 (10H, m, alkyl CH + CH₂), 2.01 (2H, m, CH), 2.12 (2H, m, CH), 2.29 (3H, s, N-CH₃), 2.75 (1H, m, CH), 2.90 (2H, m, CH), 3.48 (1H, m, CH), 3.77 (1H, m, CH), 5.11 (1H, m, CH), 7.78 (4H, m,

25

65


Ar-H), 8.04 (1H, d, J 7.5Hz, NH), 9.35 (1H, s, CONH), 9.43 (1H, s, CONH); MS(+ve): 478.2; tR = 6.74 min (XBridge 2).

Compound [19]: 4-(9-Cyclopentyl-8-isopropyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



¹H NMR (CD₃OD): 0.92 (3H, d, J 6.5Hz, CHCH₃), 1.04 (3H, d, J 6Hz, CHCH₃), 1.65 – 1.97 (14H, m, alkyl CH + CH₂), 2.25 (2H, dd, J 11.5Hz, 11.5Hz, CH), 2.36 (3H, s, N-CH₃), 2.67 (1H, m, CH), 3.03 (3H, m, CH + CH₂), 3.51 (1H, m, CH), 3.95 (1H, m, CH), 4.02 (3H, s, OCH₃), 5.09 (1H, m, CH), 7.59 (2H, m, Ar-H), 7.87 (1H, s, pyr-H-H), 8.46 (1H, d, J 7Hz, Ar-H); MS(+ve): 536.3; tR = 7.68 min (XBridge 2).


Compound [20]: 4-(9-Cyclopentyl-5,7-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



¹H NMR (DMSO): 1.02 (3H, d, J 6.5Hz, CHCH₃), 1.54 – 1.80 (12H, m, alkyl CH + CH₂), 2.10 (3H, m, (CH + CH₂), 2.27 (3H, s, N-CH₃), 2.84 (3H, m, CH₃), 3.15 (3H, s, CH₃), 3.46 (1H, m, CH), 3.80 (1H, m, N-CH), 3.95 (3H, s, OCH₃), 4.76 (1H, m, CH), 7.50 (2H, m, Ar-H), 7.75 (1H, s, Ar-H), 8.10 (1H, s, NH), 8.14 (1H, d, J 7.5Hz, Ar-H), 8.40 (1H, d, J 8Hz, Ar-H); MS(+ve): 522.3; tR = 7.76 min (XBridge 2).


Compound [21]: 4-[9-(1-Ethyl-propyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide

66




¹H NMR (DMSO): 0.85 (6H, d, J 7Hz, CH₂CH₃), 1.59 (4H, m, CH₂CH₃), , 1.66 (2H, m, CH), 1.83 (2H, m, CH), 2.37 (3H, s, N-CH₃), 2.65 (2H, m, CH₂), 2.98 (2H, m, CH), 3.50 (2H, m, CH₂), 3.83 (1H, m, N-CH), 3.94 (3H, s, OCH₃), 4.93 (1H, s, CH),
 5 4.;49(2H, m, Ar-H + NH), 7.57 (1H, s, NH), 7.79 (1H, s, Ar-H), 8.15 (1H, d, J 6.5Hz, Ar-H), 8.37 (1H, d, J 8.5Hz, Ar-H), 9.41 (1H, s, NH); MS(+ve): 496.2; tR = 6.73 min (XBridge 2).

Compound [22]: 4-[9-(1-Ethyl-propyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-
 10 b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide



¹H NMR (DMSO): 0.86 (6H, d, J 7Hz, CH₂CH₃), 1.56 (4H, m, CH₂CH₃), 1.62 (2H, m, CH), 1.76 (2H, m, CH), 2.52 (3H, s, N-CH₃), 2.63 (2H, m, CH₂), 3.22 (2H, m, CH), 3.49 (2H, m, CH₂), 3.92 (1H, m, CH), 4.97 (1H, m, CH), 7.78 (4H, m, Ar-H), 8.15 (1H, s, Pyr-H), 9.31 (1H, s, NH), 9.39 (1H, s, NH); MS(+ve): 466.3; tR = 6.26 min (XBridge 2).

Compound [23]: 4-(9-Cyclopentyl-8-isopropyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-
 20 benzamide




¹H NMR (CD₃OD): 0.83 (3H, d, J 7Hz, CHCH₃), 0.93 (3H, d, J 7Hz, CHCH₃), 1.41 (2H, m, CH), 1.65 (4H, m, CH), 1.79 (2H, M, CH), 1.91 (2H, m, CH), 2.17 (2H, m,

67

CH), 2.41 (2H, m, 2 x CH), 2.71 (4H, m, CH₃ + CH), 2.91 (2H, m, CH), 3.35 (3H, s, CH₃), 3.90 (1H, m, CH), 3.94 (3H, s, OCH₃), 4.21 (1H, m, CH), 4.27 (1H, m, CH), 7.18 (2H, m, Ar-H), 8.23 (1H, s, Pyr-H), 8.56 (1H, d, J 8.5Hz, Ar-H); MS(+ve): 550.3; tR = 8.40 min (XBridge 2).


5

Compound [24]: 4-[9-(1-Ethyl-propyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



10 ¹H NMR (DMSO): 0.85 (6H, t, J 7.5, CH₂CH₃), 1.62 (8H, m, CH + CH₂), 1.76 (2H, d, J 7Hz, CH), 1.94 (2H, dd, J 10Hz, 10 Hz, CH), 2.17 (3H, s, N-CH₃), 2.64 (2H, m, CH₂), 2.80 (2H, d, J 11Hz, CH), 3.18 (3H, s, N-CH₃), 3.49 (2H, m, CH₂), 3.75 (1H, m, CH), 3.95 (3H, s, OCH₃), 4.65 (1H, m, CH), 7.49 (2H, m, Ar-H), 7.69 (1H, s, Pyr-H), 8.07 (1H, s, NH), 8.10 (1H, d, J 9Hz, Ar-H), 8.37 (1H, d, J 9Hz, Ar-H); MS(+ve):
15 510.3; tR = 8.35 min (XBridge 2).

Compound [25]: 3-Methoxy-N-(1-methyl-piperidin-4-yl)-4-[6-oxo-9-(tetrahydro-pyran-4-yl)-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzamide




20

¹H NMR (CD₃OD): 1.60 (2H, m, CH), 1.69 (2H, d, J 11Hz, CH), 1.93 (2H, d, J 12Hz, CH), 1.97 (2H, dd, J 12Hz, 12Hz, CH), 2.18 (3H, s, N-CH₃), 2.61 (2H, m, CH₂), 2.80 (2H, d, J 10.5Hz, CH), 3.48 (2H, dd, J 11Hz, CH), 3.62 (2H, m, CH₂), 3.75 (1H, m, CH), 3.95 (3H, s, OCH₃), 4.04 (2H, d, J 8Hz, CH), 4.85 (1H, s, CH), 7.49 (2H, m, Ar-

68


H), 7.66 (1H, s, NH), 7.81 (1H, s, Pyr-H), 8.11 (1H, d, J 7.5Hz, Ar-H), 8.33 (1H, d, J 8Hz, Ar-H), 9.45 (1H, s, NH); MS(+ve): 510.1; tR = 5.34 min (XBridge 2).

Compound [26]: 3-Methoxy-4-[5-methyl-6-oxo-9-(tetrahydro-pyran-4-yl)-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide




¹H NMR (CD₃OD): 1.73 (2H, ddd, J 3Hz, 12 Hz, 18 Hz, CH), 1.84 (2H, d, J 11Hz, CH), 1.98 (4H, m, alkyl CH), 2.19 (2H, dd, J 10.5Hz, 10.5Hz, CH), 2.33 (3H, s, N-CH₃), 2.71 (2H, m, CH₂), 2.97 (2H, d, J 12Hz, CH), 3.65 (2H, dd, J 11Hz, CH), 3.75 (2H, m, CH₂), 3.92 (1H, m, CH), 4.02 (3H, s, OCH₃), 4.12 (1H, dd, J 4Hz, 11Hz, CH), 4.78 (1H, m, CH), 7.53 (2H, m, Ar-H), 8.04 (1H, s, Pyr-H), 8.45 (1H, d, J 9Hz, Ar-H); MS(+ve): 524.3; tR = 5.71 min (XBridge 2).

Compound [27]: 4-[5-Ethyl-9-(1-ethyl-propyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide




¹H NMR (DMSO): 0.86 (6H, t, J 7.5Hz, CH₃), 0.94 (3H, d, J 7Hz, CH₃), 1.63 (6H, m, alkyl CH + CH₂), 1.78 (2H, d, J 11Hz, CH), 1.94 (2H, dd, J 11.5Hz, CH), 2.18 (3H, s, N-CH₃), 2.57 (2H, m, CH₂), 2.80 (2H, m, CH₂), 3.47 (2H, m, CH₂), 3.77 (3H, m, CH + CH₂), 3.96 (3H, s, OCH₃), 4.59 (1H, m, CH), 7.50 (2H, m, Ar-H), 7.24 (1H, s, NH), 8.01 (2H, m, Ar-H + Pyr-H), 8.39 (1H, d, J 9Hz, Ar-H); MS(+ve): 524.3; tR = 7.56 min (XBridge 2).

Compound [29]: 4-(9-Benzyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



¹H NMR (DMSO): 1.58 (2H, ddd, J 4Hz, 12.5Hz, 24.5Hz, CH), 1.73 (2H, d, J 11.5Hz, CH), 1.93 (2H, dd, J 15Hz, CH), 2.17 (3H, s, N-CH₃), 2.69 (2H, m, CH₂), 2.78 (2H, d, J 11.5Hz, CH), 3.22 (3H, s, N-CH₃), 3.63 (2H, m, CH₂), 3.72 (1H, m, CH), 3.91 (3H, s, OCH₃), 4.87 (2H, s, CH₂), 7.23 – 7.32 (4H, m, Ar-H), 7.35 – 7.43 (2H, m, Ar-H), 7.44 (1H, d, J 2Hz, Ar-H), 7.72 (1H, s, Pyr-H), 8.04 (1H, d, J 7.5Hz, Ar-H), 8.12 (1H, d, J 8.5Hz), 8.16 (1H, s, NH); MS(+ve): 530.4; t_R = 4.55 min (XBridge 2).


Compound [30]: 3-Methoxy-4-(5-methyl-6-oxo-9-phenyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide



¹H NMR (DMSO): 1.56 (2H, dd, J 11.5Hz, 20.5Hz, CH), 1.75 (2H, d, J 10Hz, CH), 1.95 (2H, m, CH), 2.18 (3H, s, N-CH₃), 2.83 (3H, m, CH + CH₂), 3.16 (1H, s, CH), 3.27 (3H, s, N-CH₃), 3.71 (1H, m, N-CH), 3.88 (3H, s, OCH₃), 4.06 (2H, m, CH₂); MS(+ve): 516.3; t_R = 2.72 min (XBridge 2).


Compound [31]: 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide

70



¹H NMR (DMSO): 0.66 (2H, d, J 6Hz, cyclopropyl-CH), 0.91 (2H, d, J 6Hz, cyclopropyl-CH), 1.50 (2H, m, CH), 1.58 (4H, m, CH), 1.69 (2H, m, CH), 1.77 (2H, m, CH), 1.95 (4H, m, CH), 2.16 (3H, s, N-CH₃), 2.73 (1H, s, CH), 2.79 (2H, d, J 11Hz, CH), 2.89 (1H, s, CH), 3.16 (3H, s, N-CH₃), 3.47 (2H, s, CH₂), 3.72 (1H, m, CH), 3.94 (3H, s, OCH₃), 4.86 (1H, m, CH), 7.46 (2H, m, Ar-H), 7.67 (1H, s, NH), 7.95 (1H, s, Pyr-H), 8.07 (1H, d, J 8Hz, Ar-H), 8.38 (1H, d, J 8Hz, Ar-H); MS(+ve): 534.3; tR = 3.05 min (XBridge 2).

10 *Compound [32]: 4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide*




15 ¹H NMR (DMSO): 1.09 6H, s, CH₃), 1.58 (6H, m, CH₂), 1.74 (4H, m, CH), 1.95 (4H, m, CH), 2.16 (3H, s, NCH₃), 2.73 (2H, d, J 11.5Hz, CH), 3.18 (3H, s, NCH₃), 3.37 (2H, s, CH₂), 3.74 (1H, m, CH), 3.94 (3H, s, OCH₃), 5.19 (1H, m, CH), 7.46 (2H, m, Ar-H), 7.67 (1H, s, NH), 7.98 (1H, s, Pyr-H), 8.07 ~ (1H, d, J 8Hz, Ar-H), 8.36 (1H, d, J 8Hz); MS(+ve): 536.40; tR = 3.32 min (XBridge 2).

20 Amide couplings

The following were also prepared by the method described in Example 1, step 7:


Compound [33]: 4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



5 Yellow solid (31%). ¹H NMR (CD₃OD): δ 1.24 (d, *J* = 6.5 Hz, 3H, CH₃), 1.36 – 2.17 (m, 16H, 8 x cyclopent-*H* and 4 x piperid-*H*), 2.51 (s, 3H, CH₃), 3.14 – 3.16 (m, 2H, CH₂), 3.28 (s, 3H, CH₃), 4.00 (s, 3H, CH₃), 4.04 – 4.06 (m, 1H, CH), 4.71 – 4.74 (m, 1H, CH), 7.49 – 7.52 (m, 2H), 7.99 (s, 1H), 8.41 (d, *J* = 8.5 Hz, 1H, phe-*H*); MS+ve: 522.4. R_t = 10.03 min (purity 97%, Vydac 1)

10

Compound [34]: 4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide




15

Yellow solid (38%); MS+ve: 508.3. R_t = 9.53 min (purity 97%, Vydac 1)

Compound [35]: 4-(9-Cyclohexyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide


20

72




Off-white solid (33%); ^1H NMR (CD_3OD): δ 1.34 – 1.41 (m, 2H), 1.44 (d, $J = 6.5$ Hz, 3H, CH_3), 1.60 – 1.99 (m, 8H), 2.22 – 2.26 (m, 2H), 2.73 – 2.77 (m, 1H), 2.91 (s, 3H, CH_3), 2.96 – 3.25 (m, 5H), 3.61 – 3.63 (m, 2H, CH_2), 4.01 (s, 3H, CH_3), 4.17 – 4.21 (m, 1H, CH), 4.33 – 4.36 (m, 1H, CH), 7.59 (dd, $J = 2.0$ and 8.5 Hz, 1H, phe- H), 7.63 (d, $J = 2.0$ Hz, 1H, phe- H), 7.69 (s, 1H, pyrimid- H), 8.00 (d, $J = 8.5$ Hz, 1H, phe- H); MS+ve: 522.2. $R_t = 10.17$ min (purity 100%, Vydac 1)

Compound [36]: 4-(9-Cyclopentyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



Purple solid (40%); ^1H NMR (CD_3OD): δ 1.65 – 2.25 (m, 8H, cyclopent- H), 2.34 (s, 3H, CH_3), 2.73 – 2.75 (m, 2H, CH_2), 3.69 – 3.71 (m, 2H, CH_2), 3.90 – 3.94 (m, 1H, CH), 4.01 (s, 3H, CH_3), 5.22 – 5.25 (m, 1H, CH), 7.48 – 7.51 (m, 2H, 2 x phe- H), 7.76 (s, 1H, pyrimid- H), 8.47 (s, 1H, NH); MS+ve: 493.4. $R_t = 9.21$ min (purity 92%, Vydac 1)

Compound [37]: 4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide




Off-white solid (45%); ^1H NMR ($\text{DMSO}-d_6$): δ 1.17 (d, $J = 6.0$ Hz, 3H, CH_3), 1.29 – 2.10 (m, 16H, 8 x cyclopent- H and 8 x piperid- H), 2.21 (bs, 3H, CH_3), 2.81 – 2.83 (m,

73


2H, CH₂), 3.21 (s, 3H, CH₃), 3.73 – 3.76 (m, 1H, CH), 4.03 – 4.05 (m, 1H, CH), 4.68 – 4.72 (m, 1H, CH), 7.74 – 7.79 (m, 4H, phe-H), 8.11 (s, 1H, pyrimid-H), 9.47 (bs, 1H, NH); MS+ve: 492.3. R_t = 9.70 min (purity 92%, Vydac 1)

Compound [38]: 4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide



Off-white solid (15%); ¹H NMR (DMSO-*d*₆): δ 1.25 (d, *J* = 7.0 Hz, 3H, CH₃), 1.51 – 2.04 (m, 10H), 2.09 (s, 3H, CH₃), 2.55 – 2.66 (m, 6H), 2.78 – 2.79 (m, 2H, CH₂), 3.05 – 3.15 (m, 1H, CH), 3.95 – 4.04 (m, 1H, CH), 5.13 – 5.15 (m, 1H, CH), 7.75 (dd, *J* = 8.5 Hz, 4H, phe-H), 7.81 (s, 1H, pyrimid-H), 8.24 (d, *J* = 7 Hz, 1H, NH), 9.48 (s, 1H, NH), 9.66 (bs, 1H, NH); MS+ve: 478.3. R_t = 9.28 min (purity 100%, Vydac 1)

Compound [39]: 4-(9-Cyclohexyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide



Off-white solid (19%); ¹H NMR (DMSO-*d*₆): δ 1.16 – 1.98 (m, 18H, 10 x cyclohex-H) and 8 x piperid-H), 2.18 (s, 3H, CH₃), 2.60 (dd, *J* = 5 Hz, 2H, CH₂), 3.60 (dd, *J* = 5 Hz, 2H, CH₂), 3.73 – 3.76 (m, 1H, CH), 3.95 (s, 3H, CH₃), 4.58 – 4.63 (m, 1H, CH), 7.46 (dd, *J* = 2.0 and 8.0 Hz, 1H, phe-H), 7.49 (d, *J* = 2 Hz, 1H, phe-H), 7.60 (s, 1H), 7.78 (s, 1H), 8.10 (d, *J* = 8.0 Hz, 1H, NH), 8.37 (d, *J* = 8.0 Hz, 1H, phe-H), 9.41 (bs, 1H, NH); MS+ve: 508.4. R_t = 9.70 min (purity 95%, Vydac 1)


EXAMPLE 2General experimental

Analytical and preparative RP-HPLC-MS was performed using Waters XBridge (50 x 4.6 mm C18 3.5 μ m or 100 x 4.6 mm C18 3.5 μ m) and XBridge (100 x 19mm C18 5 μ m) columns using a linear gradient of solvent A (water containing 0.1% ammonium hydroxide and 5% acetonitrile) / solvent B (acetonitrile) systems. Preparative RP-HPLC was performed using Apex Prepsil ODS 10 μ column (22 x 250 mm) using a linear gradient of solvent A (water containing 0.1% TFA) / solvent B (acetonitrile). Mass spectra were obtained using a Waters ZQ2000 single quadrupole mass spectrometer with electro spray ionisation (ESI).

Gradients used were as follows:

Mode	Method	Column	Flow rate mL/min	% solvent B							
				0 min	0.5 min	4 min	6 min	8.5 min	10 min	12 min	40 min
Analytical	Analytical_1	4.6 x 50	1.6	10		100	100				
Analytical	Analytical_2	4.6 x 100	1	10					50	100	
Analytical	Analytical_3	4.6 x 100	1								
Preparative	Preparative_1	19 x 100	20	10	10			100		100	
Preparative	Preparative_2	19 x 100	20	30	30			80		100	
Preparative	Preparative_3	19 x 100	20	10	10			55		100	
Preparative	Preparative_4	22 x 250	9	0							60
Preparative	Preparative_5	19 x 100	20	55	55			100		100	

Intermediate 1: 2-chloro-9-cyclopentyl-5-methyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one.



1. Thionyl chloride (2.1 eq), MeOH, 0 °C for addition, reflux 2h; 2. Cyclopentanone (0.77 eq), sodium acetate (0.77 eq), sodium triacetoxyborohydride (1.11 eq), DCM, rt

16h; **3**. 2,4-Dichloro-5-nitropyrimidine (1.1 eq), K₂CO₃ (1 eq), acetone, 0 °C – rt 16h; **4**. NH₄Cl (8.5 eq), Fe (8 eq), EtOH/H₂O (4:1), reflux 2.5h; **5**. MeI (1.18 eq), NaH (1.07 eq), DMF, -10 °C – rt, 3h.

Step 1: Methyl 3-aminopropanoate

5 β-Alanine (9.37 g, 0.105 mol, 1.0 eq), was added to MeOH (50 mL) and the mixture cooled to 0 °C before the dropwise addition of thionyl chloride (16 mL, 0.221 mol, 2.1 eq) [caution: exothermic addition]. The reaction was allowed to warm to rt then heated at reflux for 2h. The solution was concentrated *in vacuo*, treated with *t*-butylmethyl ester and the resulting crystals removed by filtration. The product was a white
10 crystalline solid (11 g, 100%); ¹H (DMSO-*d*₆): δ 2.73 (2H, dd, *J* = 7 Hz, CH₂), 2.98 (2H, dd, *J* = 7 Hz, CH₂), 3.61 (3H, s, CH₃), 8.28 (2H, bs, NH₂); MS+ve: 104.1.

Step 2: Methyl 3-(cyclopentylamino)propanoate

Methyl 3-aminopropanoate (9.37 g, 0.09 mol) was dissolved in DCM (200 mL).
15 Cyclopentanone (6.43 mL, 0.07 mol, 0.77 eq), sodium acetate (5.96 g, 0.07 mol, 0.77 eq) and sodium triacetoxyborohydride (22 g, 0.10 mol, 1.11 eq) were added then the reaction was stirred at rt for 16h. 20% Sodium bicarbonate (100 mL) and 2 M sodium hydroxide (50 mL) were added and the product extracted using DCM/H₂O. The organic extracts were combined, washed with sat. NaCl, dried (MgSO₄) and evaporated under
20 reduced pressure to give the product as a pale yellow oil (8.90 g, 55%); MS+ve: 172.4.

Step 3: Methyl 3-[cyclopentyl(2-chloro-5-nitropyrimidin-4-yl)amino]propanoate

Methyl 3-(cyclopentylamino)propanoate (0.838 g, 0.005 mol, 1 eq) and K₂CO₃ (0.676 g, 0.005mol, 1 eq) were added to acetone (5 mL) and the resulting mixture cooled to 0
25 °C before the dropwise addition of 2,4-dichloro-5-nitropyrimidine (1.04 g, 0.0055 mol, 1.1 eq). The reaction mixture (RM) was warmed to rt and stirring continued for an additional 16h before the addition of a further 0.12 eq of the pyrimidine. Stirring continued for a further 3h. The RM was evaporated under reduced pressure and the product extracted using EtOAc/H₂O. The organic extracts were washed with sat NaCl,

76

dried (MgSO₄) and evaporated under reduced pressure to give the product as a brown oily residue (1.04 g, 65%); MS+ve: 329.2; R_t = 3.78 min (Analytical_1).

Step 4: 2-Chloro-9-cyclopentyl-5,7,8,9-tetrahydro-6H-pyrimido[4,5-b][1,4]diazepin-6-one

Methyl 3-[cyclopentyl(2-chloro-5-nitropyrimidin-4-yl)amino]propanoate (1.0 g, 0.003 mol, 1 eq) and NH₄Cl (1.38 g, 0.025 mol, 8.5 eq) were added to EtOH/H₂O (4:1, 10 mL) and the mixture heated to reflux. Iron powder (1.36 g, 0.024 mol, 8 eq) was added portionwise and after 2 h the RM was hot filtered through celite washing through with EtOAc (10 mL) and EtOH (10 mL) [both hot]. The solvent was evaporated under reduced pressure to give the product as a brown solid (0.35 g, 43%); ¹H (DMSO-*d*₆): δ 1.55 (4H, m, 2CH₂), 1.7 (2H, m, CH₂), 1.83 (2H, m, CH₂), 2.66 (2H, t, *J* = 5 Hz, CH₂), 3.57 (2H, t, *J* = 5 Hz, CH₂), 5.01 (1H, m, CH), 7.83 (1H, s, CH), 9.71 (1H, s, NH); MS+ve: 267.2; R_t = 2.87 min (Analytical_1).

Step 5: 2-chloro-9-cyclopentyl-5-methyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one


2-Chloro-9-cyclopentyl-5,7,8,9-tetrahydro-6H-pyrimido[4,5-b][1,4]diazepin-6-one (0.318 g, 0.0012 mol) and MeI (0.088 mL, 0.0014 mol, 1.18 eq) were added to DMF (5 mL) and the solution cooled to -10 °C. NaH (0.03 g, 0.0013 mol, 1.07 eq) was added and the RM was stirred at 0 °C for 30 min and rt for 30 min. The solvent was evaporated under reduced pressure and the product extracted using EtOAc/H₂O. The organic extracts were combined, washed with sat NaCl, dried (MgSO₄) and evaporated under reduced pressure to give the product as a purple oily residue (0.25 g, 75%); ¹H (DMSO-*d*₆): δ 1.55 (4H, m, 2CH₂), 1.7 (2H, m, CH₂), 1.9 (2H, m, CH₂), 2.63 (2H, t, *J* = 5 Hz, CH₂), 3.17 (3H, s, CH₃), 3.65 (2H, t, *J* = 5 Hz, CH₂), 4.74 (1H, m, CH), 8.14 (1H, s, CH); MS+ve: 281.2; R_t = 3.11 min (Analytical_1).

Using similar methods to those described above for Intermediate 1, the following intermediates 2-4 were prepared:

77


Intermediate 2: 2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one

Reaction of 1-aminomethyl-cyclopropanecarboxylic acid ethyl ester with cyclopentanone, sodium acetate and sodium triacetoxyborohydride in DCM gave 1-cyclopentylaminomethyl-cyclopropanecarboxylic acid ethyl ester




¹H NMR (CDCl₃): 0.84 (3H, t, J 7Hz, CH₃), 1.25 (4H, m, CH₂), 1.37 (2H, m, CH), 1.54 (2H, m, CH), 1.70 (2H, m, CH), 1.83 (2H, m, CH), 2.71 (2H, s, CH₂), 3.10 (1H, m, CH), 4.16 (2H, q, J 7Hz, CH₂CH₃)

Reaction of 1-cyclopentylaminomethyl-cyclopropanecarboxylic acid methyl ester with 2,4-dichloro-5-nitropyrimidine and K₂CO₃ in acetone gave 1-[(2-Chloro-5-nitropyrimidin-4-yl)-cyclopentyl-amino]-methyl} cyclopropanecarboxylic acid ethyl ester. Treatment with NH₄Cl and iron powder in ethanol gave 1-[(5-Amino-2-chloropyrimidin-4-yl)-cyclopentyl-amino]-methyl}-cyclopropanecarboxylic acid ethyl ester




MS (+ve): 339.2, 341.2; Rt = 4.15 min (Analytical₂).

The above compound in DMF was heated to 140°C for 2 hours. Solvent was evaporated *in vacuo*, ethyl acetate added, and the resulting solid filtered and dried under vacuum to give 2'-chloro-9'-cyclopentyl-8',9'-dihydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one



MS (+ve): 293.2, 295.2; Rt = 3.48 min (Analytical_2).


Reaction of 2-chloro-9-cyclopentyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-7,1'-cyclopropan]-6-one with MeI and NaH in DMF gave 2'-chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one



MS (+ve): 307.1, 309.2; Rt = 3.67 min (Analytical_3).

Intermediate 3: 2-chloro-9-cyclopentyl-5,7,7-trimethyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one


Reaction of 3-amino-2,2-dimethyl-propionic acid ethyl ester with cyclopentanone, sodium acetate and sodium triacetoxyborohydride in DCM gave 3-Cyclopentylamino-2,2-dimethyl-propionic acid ethyl ester



¹H NMR (CDCl₃): 1.19 (6H, s, CH₃), 1.25 (3H, t, J 7Hz, CH₃), 1.28 (2H, m, CH), 1.49 (2H, m, CH), 1.65 (2H, m, CH), 2.64 (1H, s, CH₂), 3.01 (1H, m, CH), 4.10 (2H, q, J 7.5Hz, CH₂)


Reaction of 3-cyclopentylamino-2,2-dimethyl-propionic acid ethyl ester with 2,4-dichloro-5-nitropyrimidine and K₂CO₃ in acetone gave 3-[(2-Chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-2,2-dimethyl-propionic acid ethyl ester

79




MS (+ve): 371.1, 373.1; Rt = 4.67 min (Analytical_2).

- 5 Reaction of 3-[(2-chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-2,2-dimethyl-propionic acid ethyl ester with NH_4Cl and iron powder in ethanol gave 3-[(5-Amino-2-chloro-pyrimidin-4-yl)-cyclopentyl-amino]-2,2-dimethyl-propionic acid ethyl ester



- 10 MS (+ve): 341.2, 343.2; Rt = 4.25 min (Analytical_2).

- 3-[(5-Amino-2-chloro-pyrimidin-4-yl)-cyclopentyl-amino]-2,2-dimethyl-propionic acid ethyl ester in DMF was heated to 140°C for 2 hours. Solvent was evaporated *in vacuo*, ethyl acetate added, and the resulting solid filtered and dried under vacuum to give 2-Chloro-9-cyclopentyl-7,7-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one
- 15




MS (+ve): 295.1, 297.2; Rt = 3.58 min (Analytical_2).

20

Reaction of 2-chloro-9-cyclopentyl-7,7-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one with MeI and NaH in DMF gave 2-chloro-9-cyclopentyl-5,7,7-trimethyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one

80




MS (+ve): 309.1, 311.2; Rt = 3.90 min (Analytical_3).

5 Intermediate 4: 2'-chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one

Reaction of 1-aminomethyl-cyclobutanecarboxylic acid methyl ester with cyclopentanone, sodium acetate and sodium triacetoxyborohydride in DCM gave 1-cyclopentylaminomethyl-cyclobutanecarboxylic acid methyl ester

10




¹H NMR (CDCl₃): 1.36 (2H, m, CH₂), 1.51 (2H, m, CH₂), 1.66 (2H, m, CH₂), 1.95 (4H, m, 2 x CH₂), 2.15 (2H, m, CH₂), 2.43 (4H, m, 2 x CH₂), 2.93 (2H, s, CH₂), 3.21 (1H, m, CH), 3.76 (3H, s, CH₃)

15

Reaction of 1-cyclopentylaminomethyl-cyclobutanecarboxylic acid methyl ester with 2,4-dichloro-5-nitropyrimidine and K₂CO₃ in acetone gave 1-{[(2-chloro-5-nitropyrimidin-4-yl)-cyclopentyl-amino]-methyl}-cyclobutanecarboxylic acid methyl ester


20



MS(+ve):369.2, 371.2; Rt = 3.48 min (Analytical_1).

81


Reaction of 1-{[(2-chloro-5-nitro-pyrimidin-4-yl)-cyclopentyl-amino]-methyl}-cyclobutanecarboxylic acid methyl ester with NH_4Cl and iron powder in ethanol gave 2'-chloro-9'-cyclopentyl-8',9'-dihydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one, isolated directly.



5

MS(+ve): 307.2, 309.2; R_t = 3.48 min (Analytical_1).

Reaction of 2-chloro-9-cyclopentyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-7,1'-cyclobutan]-6-one with MeI and NaH in DMF gave 2'-chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one




15 MS(+ve);:321.2, 323.3; R_t = 3.77 min (Analytical_1).

Intermediate 5: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid

20 2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one (Intermediate 2) (0.70 g, 2.5 mmol, 1 eq), 4-amino-3-methoxybenzoic acid (1.25 g, 7.5 mmol, 3 eq) and TFA (1.43 g, 12.5 mmol, 5 eq) were heated to reflux in TFE (25 mL) for 18 hours. Solvent was evaporated *in vacuo* and the resulting residue was triturated with MeOH to give 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (0.81 g, 74%).

25

82




Rt = 1.88 min (Analytical_1) MS(+ve): 438.4, MS (-ve): 436.5.

Intermediate 6: 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-

pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid

2-Chloro-9-cyclopentyl-5,7,7-trimethyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one (Intermediate 3) (0.35 g, 1.14 mmol, 1 eq), 4-amino-3-methoxybenzoic acid (0.57 g, 3.42 mmol, 3 eq) and TFA (0.42 mL, 5.7 mmol, 5 eq) were heated to reflux in TFE (10 mL) for 36 hours. Solvent was evaporated *in vacuo* and the resulting residue was triturated with EtOAc to give 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid (0.38 g, 76%).




MS(+ve): 440.3, MS(-ve): 438.5.

Intermediate 7: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid

2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-6'(5'H)-one (Intermediate 4) (1.02 g, 3.18 mmol, 1 eq), 4-amino-3-methoxybenzoic acid (1.6 g, 9.5 mmol, 3 eq) and TFA (1.2 mL, 15.9 mmol, 5 eq) were heated to reflux in TFE (40 mL) for 24 hours. Solvent was evaporated *in vacuo* and the resulting residue was triturated with EtOAc to give 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (1.08 g, 75%).


83



Rt = 2.10 min (Analytical_1); MS(+ve): 452.3, MS (-ve): 450.4.

5 Intermediate 8: 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid

2-Chloro-9-cyclopentyl-5-methyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one (Intermediate 1) (0.70 g, 2.5 mmol, 1 eq), 4-amino-3-methoxybenzoic acid (1.25 g, 7.5 mmol, 3 eq) and TFA (0.93 mL, 12.5 mmol, 5 eq) were heated to reflux in TFE (25 mL) for 20 hours. Solvent was evaporated *in vacuo* and the resulting residue was triturated with MeOH to give 4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid (0.66 g, 64%).




15 Rt = 1.88 min (Analytical_1); MS(+ve): 412.4, MS (-ve): 410.5.

Intermediate 9: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluorobenzoic acid

20 2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one (Intermediate 2) (0.42 g, 1.36 mmol, 1 eq), 4-amino-3-fluorobenzoic acid (0.63 g, 4.1 mmol, 3 eq) and TFA (0.51 mL, 6.8 mmol, 5 eq) were heated to reflux in TFE (10 mL) for 36 hours. Further 4-amino-3-fluorobenzoic acid (0.63 g, 4.1 mmol, 3 eq) and TFA (0.51 mL, 6.8 mmol, 5 eq) were added and the reaction was heated to reflux again for 4 days. Solvent was evaporated *in vacuo* and the

84

resulting residue was triturated with EtOAc to give 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluorobenzoic acid (0.41 g, 71%).




5

Rt = 1.93 min (Analytical_1); MS(+ve): 426.3

Intermediate 10: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

10 tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluorobenzoic acid


2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one (Intermediate 4) (0.20 g, 0.63 mmol, 1 eq), 4-amino-3-fluorobenzoic acid (0.30 g, 1.9 mmol, 3 eq) and TFA (0.24 mL, 3.2 mmol, 5 eq) were
 15 heated to reflux in TFE for 18 hours. Further 4-amino-3-fluorobenzoic acid (0.20 g) and TFA (0.1 mL) were added and the reaction was heated to reflux again for 2 days. Further TFA (0.24 mL) was added and the reaction was heated to reflux again for 3 days. Solvent was evaporated *in vacuo* and the resulting residue was triturated with EtOAc to give 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro
 20 [cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluorobenzoic acid (0.15 g, 55%).



Rt = 2.02 min (Analytical_1); MS(+ve): 440.3; MS(-ve): 438.4

25

Intermediate 11: 4-Amino-N-(trans-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide



5

4-Amino-N-(trans-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide is a compound known in the art and was prepared by the methods described in WO 2007/090844 A1.


10 Intermediate 12: 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(piperidin-4-yl)benzamide

4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid (Intermediate 8) (0.49 g, 1.2 mmol, 1 eq), TBTU (0.42 g, 1.3 mmol, 1.1 eq) and DIPEA (0.40 mL, 2.4 mmol, 2 eq) were stirred together
 15 in DCM (25 mL) for 30 min before addition of 4-amino-1-BOC-piperidine (0.29 g, 1.4 mmol, 1.2 eq). The RM was stirred at rt for 20 hours before diluting with DCM (30 mL) and washing sequentially with water (2 x 15 mL), 50% saturated aqueous sodium hydrogen carbonate (1 x 20 mL), saturated aqueous potassium carbonate (1 x 20 mL) and saturated brine (1 x 20 mL). Concentrated *in vacuo* to isolate 4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-BOC-piperidin-4-yl)benzamide that was used without further
 20 purification (0.71 g). *Rt* = 3.58 min (Analytical_1).

Methanolic hydrochloric acid was added and the solution was stirred for 4 hours at rt.
 25 Concentration *in vacuo* provided the hydrochloride salt of 4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(piperidin-4-yl)benzamide. This salt was partitioned between DCM and aqueous potassium carbonate solution at pH 8. The DCM layer was separated and the aqueous layer extracted with DCM and EtOAc. The combined organic layers were dried

86

(MgSO₄) and concentrated *in vacuo* to provide 4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(piperidin-4-yl)benzamide as free base (0.51 g).




Rt = 2.78 min (Analytical_1); MS(+ve): 494; MS(-ve): 492.

Compound [254]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide


4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (22 mg, 0.05 mmol, 1 eq), DIPEA (17 μ l, 0.10 mmol, 2 eq) and TBTU (18 mg, 0.055 mmol, 1.1 eq) were added to 0.5 mL DMF and the resulting solution stirred at rt for 30 min before the addition of 1-amino-4-methylpiperazine (12 μ l, 0.1 mmol, 2 eq). The RM was then stirred at rt for 4 hours before purifying by preparative RP-HPLC-MS (Preparative_1) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide (white solid, 8 mg, 30%).



Rt = 2.63 min (Analytical_1) MS(+ve): 535.5; MS(-ve): 533.6; ¹H NMR (DMSO-d₆) δ ppm: 0.63 - 0.70 (2 H, m), 0.87 - 0.94 (2 H, m), 1.43 - 1.54 (2 H, m), 1.55 - 1.63 (2 H,

m), 1.64 - 1.73 (2 H, m), 1.83 - 1.94 (2 H, m), 2.18 (3 H, s), 2.31 - 2.48 (4 H, m), 2.92 (4 H, m), 3.16 (3 H, s), 3.47 (2 H, s), 3.94 (3 H, s), 4.78 - 4.91 (1 H, m), 7.35 - 7.48 (2 H, m), 7.68 (1 H, s), 7.98 (1 H, s), 8.39 (1 H, d, J=8.3 Hz), 9.31 (1 H, s).


- 5 *Compound [218]: (±)-4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide*
- 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 7)
- 10 (15 mg, 0.033 mmol), DIPEA (12 µl, 0.066 mmol, 2 eq) and TBTU (12 mg, 0.036 mmol, 1.1 eq) were added to 0.5 mL DMF and the resulting solution stirred at rt for 5 min before the addition of 3-aminoquinuclidine dihydrochloride (8 mg, 0.04 mmol, 1.2 eq) and DIPEA (13 µL). The RM was then stirred at rt for 3 hours before purifying by preparative RP-HPLC-MS (Preparative_2) to provide (±)-4-(9'-cyclopentyl-5'-methyl-
- 15 6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide.



- 20 Rt = 3.26 min (Analytical_1); MS(+ve): 560.4; ¹H NMR (DMSO-d₆) δ ppm: 8.37 (d, J = 8.3 Hz, 1H), 8.08 (d, J = 6.8 Hz, 1H), 8.05 (s, 1H), 7.73 (s, 1H), 7.45 - 7.53 (m, 2H), 4.82 (quin, J = 8.3 Hz, 1H), 3.84 - 4.01 (m, 4H), 3.64 (s, 2H), 3.18 (s, 3H), 3.04 - 3.14 (m, 1H), 2.88 (t, J = 9.8 Hz, 1H), 2.58 - 2.75 (m, 4H), 2.21 - 2.33 (m, 2H), 2.07 (s, 2H), 1.97 (br. s., 2H), 1.87 (br. s., 2H), 1.76 (d, J = 4.9 Hz, 2H), 1.56 - 1.70 (m, 8H), 1.30
- 25 (br. s., 1H)

Compound [195]: 4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methyl-piperazin-1-yl)-benzamide

4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid (Intermediate 6) (20 mg, 0.046 mmol, 1 eq), DIPEA (16 μ l, 0.091 mmol, 2 eq) and TBTU (16 mg, 0.05 mmol, 1.1 eq) were added to 0.5 mL DMF and the resulting solution stirred at rt for 5 min before the addition of 1-amino-4-methylpiperazine (7 mg). The RM was then stirred at rt for 16 hours before purifying by preparative RP-HPLC-MS (Preparative_1) to provide 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methyl-piperazin-1-yl)-benzamide.




Rt = 2.81 min (Analytical_1); MS(+ve): 537.5; ^1H NMR (DMSO- d_6) δ ppm: 1.09 (6H, s), 1.61 (4H, m), 1.73 (2H, m), 1.87 (2H, m), 2.19 (3H, s), 2.42 (2H, m), 2.92 (4H, m), 3.29 (3H, s), 3.37 (3H, m), 3.93 (3H, s), 3.17 (1H, m), 7.40 (2H, m), 7.68 (1H, s), 7.98 (1H, s), 8.37 (1H, d, J 7Hz), 9.30 (1H, s).

Compound [221]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 7) (15 mg, 0.033 mmol, 1 eq), DIPEA (12 μ l, 0.066 mmol, 2 eq) and TBTU (12 mg, 0.036 mmol, 1.1 eq) were added to 0.5 mL DMF and the resulting solution stirred at rt for 5 min before the addition of 1-amino-4-methylpiperazine (5 mg) and DIPEA (13 μ L). The RM was then stirred at rt for 3 hours before purifying by preparative RP-HPLC-

89

MS (Preparative_2) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide.




5

Rt = 2.79 min (Analytical_1); MS(+ve): 549.5; ¹H NMR (DMSO-d₆) δ ppm: 9.31 (s, 1H), 8.37 (d, J = 8.3 Hz, 1H), 8.05 (s, 1H), 7.72 (s, 1H), 7.37 - 7.47 (m, 2H), 4.73 - 4.86 (m, 1H), 3.93 (s, 3H), 3.64 (s, 2H), 3.18 (s, 3H), 2.92 (br. s., 4H), 2.36 (br. s., 4H), 2.27 (d, J = 9.3 Hz, 2H), 2.18 (s, 3H), 1.97 (br. s., 2H), 1.79 - 1.91 (m, 1H), 1.76 (br. s., 2H), 1.54 - 1.72 (m, 7H).


Compound [371]: -(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
 2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one (Intermediate 2) (0.14 g, 0.45 mmol, 1 eq), 4-amino-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (Intermediate 11) (0.26 g, 0.67 mmol, 1.5 eq) and TFA (0.17 mL, 2.2 mmol, 5 eq) in TFE (3 mL) were heated together at 80°C for 18 hours. Further quantity of Intermediate 2 (0.14 g) was added and the reaction further heated for 48 hours. RM was concentrated *in vacuo* and the residue was purified by flash column chromatography on silica eluting with 0→20% ammonia/MeOH in DCM gradient followed by preparative HPLC (Preparative_4) then finally preparative RP-HPLC-MS (Preparative_1) to give 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (0.11 g, 25%).

90



Rt = 3.18 min (Analytical_1); ES(+ve): 657.6; ES(-ve): 655.7; ¹H NMR (DMSO-d₆) δ ppm: 0.05 (d, J = 4.20 Hz, 3H), 0.44 (d, J = 7.74 Hz, 2H), 0.67 (d, J = 1.61 Hz, 2H), 0.74 - 0.85 (m, 1H), 0.90 (s, 2H), 1.23 - 1.44 (m, 3H), 1.44 - 1.55 (m, 2H), 1.54 - 1.64 (m, 2H), 1.64 - 1.74 (m, 2H), 1.88 (br. s., 5H), 2.08 (s, 3H), 2.13 (d, J = 6.45 Hz, 2H), 2.17 - 2.28 (m, 1H), 2.29 - 2.47 (m, 3H), 2.60 - 2.67 (m, 1H), 3.06 - 3.23 (m, 5H), 3.47 (s, 2H), 3.65 - 3.79 (m, 1H), 3.94 (s, 2H), 4.05 - 4.13 (m, 1H), 4.78 - 4.91 (m, 1H), 7.48 (s, 2H), 7.67 (s, 1H), 7.98 (s, 1H), 8.00 - 8.07 (m, 1H), 8.31 - 8.45 (m, 1H).

Compound [372]: 4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
2-Chloro-9-cyclopentyl-5,7,7-trimethyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one (Intermediate 3) (0.13 g, 0.43 mmol, 1 eq), 4-amino-N-(trans-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (Intermediate 11) (0.26 g, 0.67 mmol, 1.5 eq) and TFA (0.17 mL, 2.2 mmol, 5 eq) in TFE (3 mL) were heated together at 80°C for 18 hours. Further quantity of Intermediate 6 (0.13 g) was added and the reaction further heated for 48 hours. RM was concentrated *in vacuo* and the residue was purified by flash column chromatography on silica eluting with 0→20% ammonia/MeOH in DCM gradient followed by preparative HPLC (Preparative_4) then finally preparative RP-HPLC-MS (Preparative_1) to give 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (0.03 g, 7%).




Rt = 3.38 min (Analytical_1); ES(+ve): 659.6; ES(-ve): 657.7; ¹H NMR (DMSO-d₆) δ ppm: 0.05 (d, J = 3.86 Hz, 3H), 0.40 - 0.48 (m, 3H), 0.73 - 0.84 (m, 2H), 1.03 - 1.13 (m, 10H), 1.18 (s, 1H), 1.24 - 1.43 (m, 6H), 1.61 (br. s., 3H), 1.74 (d, J = 9.65 Hz, 2H), 1.88 (br. s., 5H), 2.08 (s, 5H), 2.13 (d, J = 6.75 Hz, 2H), 2.17 - 2.25 (m, 1H), 2.36 (br. s., 3H), 2.60 - 2.67 (m, 1H), 3.14 - 3.22 (m, 2H), 3.37 (s, 1H), 3.51 (s, 1H), 3.94 (s, 2H), 5.18 (t, J = 8.52 Hz, 1H), 7.43 - 7.50 (m, 1H), 7.67 (s, 1H), 7.98 (s, 1H), 8.03 (d, J = 7.72 Hz, 1H), 8.35 (d, J = 8.36 Hz, 1H).

Alternatively, the skilled person would appreciate that Examples 5, 6 could be synthesised by reaction of Intermediates 5 or 6 with the known compound trans-4-(4-cyclopropylmethyl-piperazin-1-yl)-cyclohexylamine (described in US 6,861,422 B2) using standard amide bond formation conditions such as those described in Example 1.


Compound [345]: 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide
 4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(piperidin-4-yl)benzamide (Intermediate 12) (35 mg, 0.07 mmol, 1 eq) and sodium triacetoxyborohydride (18 mg, 0.084 mmol, 1.2 eq) in DCM (0.5 mL) were reacted with tetrahydropyran-4-one (6.5 µL, 0.07 mmol, 1 eq) at rt for 2 days. Further and sodium triacetoxyborohydride (18 mg, 1.2 eq), tetrahydropyran-4-one (6.5 µL, 1 eq) were added and acetic acid (4 µL, 1 eq). After two days the RM was diluted with DCM, washed with water and the organic layer concentrated *in vacuo*.
 The resulting residue was purified by preparative RP-HPLC-MS (Preparative_1) to provide *4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide* (10 mg, 25%).

92



Rt = 2.74 min (Analytical_1); ES(+ve): 578.5; ES(-ve): 576.7.


- 5 *Compound [373]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide*
- 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluorobenzoic acid (Intermediate 10)
- 10 (100 mg, 0.23 mmol, 1 eq), DIPEA (80 μ l, 0.46 mmol, 2 eq) and TBTU (80 mg, 0.25 mmol, 1.1 eq) were added to 1 mL DMF and the resulting solution stirred at rt for 10 min before the addition of 1-amino-4-methylpiperazine (33 μ L, 0.27 mmol, 1.2 eq). The RM was then stirred at rt for 1 hour before splitting into two equal batches and purifying by preparative RP-HPLC-MS (Preparative_3) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-
- 15 *b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide* (0.05 g, 41%).



- 20 Rt = 2.74 min (Analytical_1); MS(+ve): 537.5; ^1H NMR (DMSO- d_6) δ ppm: 1.55 – 1.65 (8H, m), 1.71 (2H, m), 1.88 (2H, m), 2.18 (3H, s), 2.28 (2H, q, J 10.5 Hz), 2.41 – 2.52 (4H, m), 2.89 (4H, m), 3.18 (3H, s), 3.61 (2H, s), 4.76 (1H, quintet, 8.5 Hz), 7.61 (2H, m), 8.03 (1H, s), 8.12 (1H, dd, J 8Hz, 8.5 Hz), 8.76 (1H, s), 9.36 (1H, s).

Compound [374]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluorobenzoic acid (Intermediate 9) (0.27 g, 0.63 mmol, 1 eq), DIPEA (0.21 mL, 1.26 mmol, 2 eq) and TBTU (0.22 g, 0.69 mmol, 1.1 eq) were added to 5 mL DCM and the resulting solution stirred at rt for 30 min before the addition of 1-amino-4-methylpiperazine (91 μ l, 0.75 mmol, 1.2 eq). The RM was then stirred at rt for 4 hours before washing with water, concentrating *in vacuo* and purifying the residue by preparative RP-HPLC-MS (Preparative_1) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide (0.14 g, 44%).



15

Rt = 2.58 min (Analytical_1); MS(+ve): 523.4; ^1H NMR (DMSO- d_6) δ ppm: 0.65 – 0.67 (2H, s, CH₂), 0.85 – 0.95 (2H, m, CH₂), 1.23 – 1.82 (8H, m, alkyl-H), 2.18 (3H, s, CH₃), 2.36 – 2.41 (4H, m, alkyl-H), 2.88 (4H, s, alkyl-H), 3.16 (3H, s, CH₃), 3.44 (2H, s, alkyl-H), 4.77 – 4.80 (1H, m, CH), 7.6 (2H, m, aryl-H), 7.96 (1H, s, aryl-H), 8.16 (1H, t, J = 8 Hz, aryl-H), 8.70 (1H, s, NH), 9.35 (1H, s, NH).

20


Compound [194]: ($\pm$)-4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide

4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid (Intermediate 6) (20 mg, 0.046 mmol, 1 eq), DIPEA (16 μ l, 0.091 mmol, 2 eq) and TBTU (16 mg, 0.05 mmol, 1.1 eq) were added to 0.5 mL DMF and the resulting solution stirred at rt for 5 min before the addition of 3-aminoquinuclidine dihydrochloride (11 mg, 0.055 mmol, 1.1 eq). The RM

25

94


was then stirred at rt for 16 hours before purifying by preparative RP-HPLC-MS (Preparative_1) to provide ($\pm$)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide (8 mg, 32%).



Rt = 3.28 min (Analytical_1); MS(+ve): 548.5; ¹H NMR (DMSO-d₆) δ ppm: 8.37 (1H, d, J 8 Hz), 8.07 (1H, d, J 6.5 Hz), 7.98 (1H, s), 7.69 (1H, s), 7.47 – 7.50 (2H, m), 5.19 (1H, quint, J 8 Hz), 3.95 (4H, m), 3.37 (2H, s), 3.18 (3H, s), 3.11 (1H, m), 2.89 (1H, m), 2.65 – 2.72 (4H, m), 1.88 (2H, m), 1.73 – 1.87 (4H, m), 1.57 – 1.61 (6H, m), 1.31 (1H, m), 1.09 (6H, s).

Compound [186]: ($\pm$)-4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (25 mg, 0.057 mmol, 1 eq), DIPEA (20 μ L, 0.11 mmol, 2 eq) and TBTU (20 mg, 0.063 mmol, 1.1 eq) were added to 0.5 mL DMF and the resulting solution stirred at rt for 15 min before the addition of 3-aminoquinuclidine dihydrochloride (11 mg, 0.055 mmol, 1.1 eq) and DIPEA (40 μ L). The RM was then stirred at rt for 16 hours before purifying by preparative RP-HPLC-MS (Preparative_1) to provide ($\pm$)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide.




Rt = 3.02 min (Analytical_1); MS(+ve): 546.4; ¹H NMR (DMSO-d₆) δ ppm: 8.40 (1H, d, J 8 Hz), 8.08 (1H, d, J 7 Hz), 7.98 (1H, s), 7.69 (1H, s), 7.47 – 7.50 (2H, m), 4.87 (1H, quint, J 9 Hz), 4.10 (1H, q, J 5.5 Hz), 3.95 (4H, m), 3.47 (2H, s), 3.27 (3H, s), 3.07 (1H, m), 2.08 (1H, m), 2.63 – 2.86 (4H, m), 1.88 (3H, m), 1.78 (1H, m), 1.68 (2H, m),
 5 1.50 – 1.62 (5H, m), 1.32 (1H, m), 0.90 (2H, m), 0.66 (2H, m).

Compound [375]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide

10 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (48 mg, 0.11 mmol, 1 eq), DIPEA (36 μL, 0.22 mmol, 2 eq) and TBTU (39 mg, 0.12 mmol, 1.1 eq) were added to 5 mL DCM and the resulting solution stirred at rt for 15 min before the addition of 2-(4-aminopiperazin-1-yl)ethanol (19 mg, 0.13 mmol, 1.2
 15 eq). The RM was then stirred at rt for 16 hours before concentrating *in vacuo* and purifying by preparative RP-HPLC-MS (Preparative_1) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide (24 mg, 39%).

20




Rt = 2.48 min (Analytical_1); MS(+ve): 565.4; ¹H NMR (DMSO) δ ppm: 0.66 – 0.67 (2H, m, CH₂), 0.88 – 0.90 (2H, m, CH₂), 1.23 – 1.90 (8H, m, alkyl-H), 2.34-2.41 (4H, m, alkyl-H), 2.91 (4H, t, J = 4 Hz, alkyl-H), 3.16 (3H, s, CH₃), 3.46-3.51 (4H, m), 3.93 (3H, s, CH₃), 4.38-4.45 (1H, m,) 4.81–4.87 (1H, m, CH), 7.40 (2H, d, J = 10.5 Hz, aryl-H), 7.67 (1H, s), 7.98 (1H, s), 8.39 (1H, d, J = 8 Hz, aryl-H), 9.30 (1H, s, NH).

25

Compound [376]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (50 mg, 0.11 mmol, 1 eq), DIPEA (38 μ L, 0.23 mmol, 2 eq) and TBTU (40 mg, 0.12 mmol, 1.1 eq) were added to 5 mL DCM and the resulting solution stirred at rt for 15 min before the addition of 4-aminomorpholine (12 mg, 0.14 mmol, 1.2 eq). The RM was then stirred at rt for 16 hours before concentrating *in vacuo* and purifying by preparative RP-HPLC-MS (Preparative_1) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide (26 mg, 44%).



15


Rt = 2.72 min (Analytical_1); MS(+ve): 522.4; ¹H NMR (DMSO) δ ppm: 0.65 – 0.71 (2H, m, CH₂), 0.85 – 0.91 (2H, m, CH₂), 1.23 – 1.88 (8H, m, alkyl-H), 2.92 (4H, t, J = 4 Hz, alkyl-H), 3.16 (3H, s, CH₃), 3.47 (2H, s, alkyl-H), 3.66 (4H, t, J = 4.5 Hz), 3.94 (3H, s, CH₃), 4.81 – 4.87 (1H, m, CH), 7.41 (2H, d, J = 9.5 Hz), 7.69 (1H, s), 7.98 (1H, s), 8.40 (1H, d, J = 8 Hz), 9.41 (1H, s, NH).

20

Compound [347]: (R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide

25 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (0.11 g, 0.25 mmol, 1 eq), DIPEA (0.17 mL) and TBTU (88 mg, 0.28 mmol, 1.1 eq) were added to 5 mL DCM and the resulting solution stirred at rt for 30 min before the

addition of (R)-(+)-3-aminoquinuclidine dihydrochloride (60 mg, 0.30 mmol, 1.2 eq). The RM was then stirred at rt for 16 hours before diluting with DCM and washing sequentially with water (x2), 50% saturated aqueous sodium hydrogen carbonate, saturated aqueous sodium hydrogen carbonate and saturated brine. The organic layer
 5 was dried (MgSO₄) and concentrated *in vacuo*. The residue was dissolved in the minimum quantity of EtOAc before addition of n-heptane to precipitate (R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide (white solid, 0.13 g, 96%).




Rt = 3.07 min (Analytical_1); MS(+ve): 546.5; MS(-ve): 544.6; ¹H NMR (DMSO-d₆)
 δ ppm 0.62 - 0.72 (2 H, m), 0.87 - 0.94 (2 H, m), 1.20 - 1.38 (1 H, m), 1.44 - 1.54 (2 H,
 m), 1.54 - 1.64 (4 H, m), 1.64 - 1.74 (2 H, m), 1.75 - 1.85 (1 H, m), 1.85 - 1.95 (3 H, m),
 15 2.65 - 2.79 (4 H, m), 2.83 - 3.01 (1 H, m), 3.11 - 3.21 (4 H, m), 3.47 (2 H, s), 3.95 (4 H,
 s), 4.77 - 4.93 (1 H, m), 7.42 - 7.54 (2 H, m), 7.69 (1 H, s), 7.98 (1 H, s), 8.10 (1 H, d,
 J=6.3 Hz), 8.39 (1 H, d, J=8.3 Hz).

Compound [348]: (S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-
 20 *tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-*
methoxy-N-(quinuclidin-3-yl)benzamide

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-
 pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5)
 (0.11 g, 0.25 mmol, 1 eq), DIPEA (0.17 mL) and TBTU (88 mg, 0.28 mmol, 1.1 eq)
 25 were added to 5 mL DCM and the resulting solution stirred at rt for 30 min before the
 addition of (S)-(-)-3-aminoquinuclidine dihydrochloride (60 mg, 0.30 mmol, 1.2 eq).
 The RM was then stirred at rt for 16 hours before diluting with DCM and washing
 sequentially with water (x2), 50% saturated aqueous sodium hydrogen carbonate,


saturated aqueous sodium hydrogen carbonate and saturated brine. The organic layer was dried (MgSO₄) and concentrated *in vacuo*. The residue was dissolved in the minimum quantity of EtOAc before addition of n-heptane to precipitate (S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide (white solid, 0.075 g, 55%).



Rt = 3.04 min (Analytical_1); MS(+ve): 546.5; MS(-ve): 544.6; ¹H NMR (DMSO-d₆) δ ppm: 0.59 - 0.74 (2 H, m), 0.86 - 0.97 (2 H, m), 1.20 - 1.43 (1 H, m), 1.45 - 1.75 (7 H, m), 1.76 - 1.98 (4 H, m), 2.66 - 2.86 (3 H, m), 2.89 - 3.07 (1 H, m), 3.17 (3 H, s), 3.47 (3 H, s), 3.89 - 4.02 (4 H, m), 4.76 - 4.96 (1 H, m), 7.42 - 7.59 (2 H, m), 7.70 (1 H, s), 7.98 (1 H, s), 8.14 (1 H, d, J=6.3 Hz), 8.40 (1 H, d, J=7.8 Hz).


Compound [377]: 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide 2'-Chloro-9'-cyclopentyl-5'-methyl-8',9'-dihydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepin]-6'(5'H)-one (Intermediate 4) (0.065 g, 0.2 mmol, 1 eq), 4-amino-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (Intermediate 11) (0.24 g, 0.6 mmol, 3 eq) and TFA (75 μL, 1 mmol, 5 eq) in TFE (6 mL) were heated to reflux together 40 hours. RM was concentrated *in vacuo* and the residue was partitioned between EtOAc and saturated sodium hydrogen carbonate. The organic layer was separated, washed with saturated brine and concentrated *in vacuo* to a residue that was purified by preparative RP-HPLC-MS (Preparative_1) to give 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide.

99




Rt = 3.41 min (Analytical_1); ES(+ve): 671.6; ES(-ve): 669.7; ¹H NMR (DMSO-d₆) δ ppm: 0.03 (2H, dd, J 9.5 Hz, 5 Hz), 0.41 (2H, dd, J 10 Hz, 5 Hz), 0.79 (1H, m), 1.20 – 1.37 (4H, m), 1.55 – 1.94 (18H, m), 2.04 (2H, d, J 6.5 Hz), 2.16 – 2.46 (8H, m), 3.15 (3H, s), 3.32 (2H, s), 3.69 (1H, m), 3.90 (3H, s), 4.79 (1H, m), 7.44 (2H, m), 7.68 (1H, s), 8.01 (2H, m), 8.34 (1H, d, J 8.5 Hz).

Compound [378]: 4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide
2-Chloro-9-cyclopentyl-5-methyl-8,9-dihydro-5H-pyrimido[4,5-b][1,4]diazepin-6(7H)-one (Intermediate 1) (0.13 g, 0.43 mmol, 1 eq), 4-amino-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (Intermediate 11) (0.26 g, 0.67 mmol, 1.5 eq) and TFA (0.17 mL, 2.2 mmol, 5 eq) in TFE (3 mL) were heated together at 80°C for 18 hours. Further quantity of Intermediate 1 (0.10 g) was added and the reaction further heated for 48 hours. RM was concentrated *in vacuo* and the residue was purified by flash column chromatography on silica eluting with 0→20% ammonia/MeOH in DCM gradient followed by preparative HPLC (Preparative_4) then finally preparative RP-HPLC-MS (Preparative_1) to give 4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide (0.11 g, 26%).



Rt = 2.88 min (Analytical_1); ES(+ve): 631.5; ES(-ve): 629.6; ¹H NMR (DMSO-d₆) δ ppm: -0.15 - 0.11 (m, 2 H), 0.40 (d, *J* = 8.06 Hz, 2 H), 0.63 - 0.85 (m, 1 H), 1.11 - 1.41 (m, 3 H), 1.57 (br. s., 2 H), 1.69 (d, *J* = 17.73 Hz, 2 H), 1.74 - 1.95 (m, 6 H), 2.04 (s, 2 H), 2.09 (d, *J* = 6.45 Hz, 2 H), 2.16 (d, *J* = 9.67 Hz, 1 H), 2.32 (br. s., 2 H), 2.49 (br. s., 1 H), 2.51 - 2.59 (m, 2 H), 2.59 (br. s., 1 H), 3.07 - 3.17 (m, 3 H), 3.27 (s, 5 H), 3.53 - 3.62 (m, 2 H), 3.67 (dd, *J* = 7.41, 3.55 Hz, 1 H), 3.89 (s, 2 H), 4.04 (q, *J* = 5.37 Hz, 1 H), 4.76 (quin, *J* = 8.22 Hz, 1 H), 7.38 - 7.48 (m, 2 H), 7.68 (s, 1 H), 7.99 (d, *J* = 7.74 Hz, 1 H), 8.03 (s, 1 H), 8.33 (d, *J* = 8.06 Hz, 1 H).

Intermediates 13-19: 4-(amine-substituted)cyclohexanamines



1. [Analogous to Method III in Abdel-Magid A.F. et al. (1996) *J. Org. Chem.*, 61, 3849-3862] sodium triacetoxyborohydride (1.5 eq), acetic acid (1 eq), amine (1.1 eq), THF, rt 20h; 2. H₂, 10% Pd/C, 60 °C, methanol, flow hydrogenation; 3. sodium iodide (4 eq), chlorotrimethylsilane (4 eq), acetonitrile, 0 °C -rt, 18h.

Step 1:

N-benzyloxycarbonyl-4-aminocyclohexanone, a compound known in the art [WO2007/002181 A2] (typically 1 mmol scale), was added to a reaction tube containing THF (5 mL), the appropriate substituted-piperazine, -homopiperazine, or morpholine, along with acetic acid and sodium triacetoxyborohydride. The reaction was stirred at ambient temperature for 20 hours. The reaction was quenched with NaHCO₃ solution (2 mL) before acidifying to pH2 with 1N HCl solution. The mixture was washed with EtOAc before separating the aqueous layer and basifying to pH10 with 2N NaOH solution. The product was extracted into EtOAc, which was washed with sat. NaCl, dried (MgSO₄) and evaporated under reduced pressure to provide the required benzyl 4-(amine-substituted)cyclohexylcarbamate.








Step 2:

Each benzyl 4-(amine-substituted)cyclohexylcarbamate were dissolved in methanol to a concentration of 0.05 M. Hydrogenation was conducted using an H-CubeTM (ThalesNano Inc.) flow reactor at 1 mL/min flow rate over 10% Pd/C catalyst heated to 60 °C under full hydrogen mode. Concentration under reduced pressure provided the desired 4-(amine-substituted)cyclohexanamine product as oils. Solid hydrochloride salts could be obtained by stirring with ethereal HCl.

Where it was required to selectively deprotect the benzylcarbamates in the presence of an *N*-benzyl group (intermediates 16 & 18), step 3 was employed instead of step 2.

Step 3:

Sodium iodide (1.5 mmol, 4eq) was dissolved in anhydrous acetonitrile (5 mL) and was stirred whilst chlorotrimethylsilane (1.5 mmol, 4 eq) was added. After 5 minutes this mixture was added slowly to a solution of the 4-(amine-substituted)cyclohexanamine dissolved in anhydrous acetonitrile (2 mL) cooled in an ice-water bath. After 1 hour the cooling bath was removed and the RM was stirred at ambient temperature for 18 hours. RM was concentrated under reduced pressure and redissolved in 1% water in methanol solution. The mixture was absorbed onto an SCX II column, washed with methanol and eluted with ammonia in methanol. Concentrated in vacuo to provide the desired 4-(amine-substituted)cyclohexanamine.


Intermediate	Structure	ES(+ve) [M+H] ⁺	Analytical_1 Rt (min)
13		212.3	2.05
14		252.3	2.42
15		nd	-
16		nd	-
17		nd	-
18		274.3	3.27
19		185.2	1.47

nd = no data (ionization not detected under standard conditions)

Compound [379]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-
5 tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-
((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide
4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-
pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5)
(66 mg, 0.15 mmol, 1 eq), DIPEA (52 μ l, 0.3 mmol, 2 eq) and TBTU (54 mg, 0.17

103

mmol, 1.1 eq) were added to 1.5 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-(4-ethylpiperazin-1-yl)cyclohexanamine (Intermediate 13) as the trihydrochloride salt (58 mg, 0.18 mmol, 1.2 eq) and DIPEA (78 μ L, 0.45 mmol, 3 eq). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_2) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide (colourless glass, 26 mg, 28%).




Rt = 3.00 min (Analytical_1) MS(+ve) 631.6; MS(-ve) 629.7; Rt = 3.00 min (Analytical_1) MS(+ve) 631.6; MS(-ve) 629.7; ¹H NMR (DMSO-d₆) δ ppm: 0.59 - 0.72 (2 H, m), 0.81 - 0.92 (2 H, m), 0.97 (3 H, t, J=7.1 Hz), 1.20 - 1.42 (4 H, m), 1.42 - 1.55 (2 H, m), 1.54 - 1.64 (2 H, m), 1.64 - 1.73 (2 H, m), 1.83 (2 H, d, J=11.7 Hz), 1.89 (4 H, d, J=7.8 Hz), 2.14 - 2.42 (7 H, m), 3.16 (3 H, s), 3.47 (2 H, s), 3.64 - 3.79 (1 H, m), 3.94 (3 H, s), 4.84 (1 H, quin, J=8.7 Hz), 7.40 - 7.54 (2 H, m), 7.67 (1 H, s), 7.98 (1 H, s), 8.03 (1 H, d, J=7.8 Hz), 8.38 (1 H, d, J=8.3 Hz).

Compound [380]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide


From the synthesis and purification of Compound [379] was isolated as a separate compound 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide (colourless glass, 27 mg, 29%).

104



Rt = 3.13 min (Analytical_1) MS(+ve) 631.6; MS(-ve) 629.7; ^1H NMR (DMSO- d_6) δ ppm: 0.60 - 0.72 (2 H, m), 0.83 - 0.93 (2 H, m), 0.98 (3 H, t, $J=7.1$ Hz), 1.38 - 1.55 (6 H, m), 1.55 - 1.63 (2 H, m), 1.63 - 1.80 (4 H, m), 1.80 - 1.96 (4 H, m), 2.02 - 2.17 (1 H, m), 2.16 - 2.44 (7 H, m), 3.16 (3 H, s), 3.47 (2 H, s), 3.79 - 3.92 (1 H, m), 3.94 (3 H, s), 4.84 (1 H, quin, $J=8.5$ Hz), 7.44 - 7.58 (2 H, m), 7.67 (1 H, s), 7.98 (1 H, s), 8.03 (1 H, d, $J=7.3$ Hz), 8.37 (2 H, d, $J=8.8$ Hz).


- 10 *Compound [381]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide*
- 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5)
- 15 (88 mg, 0.2 mmol, 1 eq), DIPEA (70 μL , 0.4 mmol, 2 eq) and TBTU (72 mg, 0.24 mmol, 1.2 eq) were added to 2 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexanamine (Intermediate 14) (60 mg, 0.24 mmol, 1.2 eq). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_2)
- 20 to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide as a mixture cis and trans isomers (off-white solid, 67 mg, 50%).



105

Rt = 4.12 min (broad peak) (Analytical_1) MS(+ve) 671.6; MS(-ve) 669.7; ¹H NMR (DMSO-*d*₆) δ ppm: 0.01 (2 H, d, *J*=4.4 Hz), 0.40 (2 H, d, *J*=7.3 Hz), 0.63 (2 H, br. s.), 0.71 - 0.83 (1 H, m), 0.87 (2 H, br. s.), 1.02 (1 H, t, *J*=7.1 Hz), 1.25 - 1.38 (2.5 H, m), 1.39 - 1.52 (3.5 H, m), 1.52 - 1.61 (2 H, m), 1.65 (4 H, d, *J*=5.9 Hz), 1.70 - 1.80 (3 H, m), 1.85 (3 H, d, *J*=4.4 Hz), 2.21 - 2.31 (1 H, m), 2.55 - 2.83 (8 H, m), 3.13 (3.5 H, s), 3.35 - 3.55 (3 H, m), 3.69 (0.5 H, br. s.), 3.91 (3.5 H, s), 4.31 (0.5 H, t, *J*=5.1 Hz), 4.81 (1 H, t, *J*=8.5 Hz), 7.25 - 7.54 (2 H, m), 7.64 (1 H, s), 7.84 - 8.16 (2 H, m), 8.35 (1 H, d, *J*=8.3 Hz).

- 10 *Compound [382]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide*
- 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5)
- 15 (44 mg, 0.1 mmol, 1 eq), DIPEA (35 μL, 0.2 mmol, 2 eq) and TBTU (36 mg, 0.12 mmol, 1.2 eq) were added to 1 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-(4-ethyl-1,4-diazepan-1-yl)cyclohexanamine (Intermediate 15) as the trihydrochloride salt (32 mg, 0.1 mmol, 1 eq) and DIPEA (53 μL, 0.3 mmol, 3 eq). The RM was then stirred at rt for 2 hours before purifying by
- 20 preparative RP-HPLC-MS (Preparative_2) to provide *4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide* as a mixture cis and trans isomers (white solid, 10 mg, 16%).




25

Rt = 3.98 min, 4.18 min (broad peaks) (Analytical_1) MS(+ve) 645.6; MS(-ve) 643.7; ¹H NMR (DMSO-*d*₆) δ ppm: 0.57 - 0.74 (2 H, m), 0.82 - 0.93 (2 H, m), 0.95 - 1.09 (3

H, m), 1.30 - 1.42 (2 H, m), 1.42 - 1.54 (3.5 H, m), 1.58 (2 H, d, $J=4.4$ Hz), 1.63 - 1.75 (3.5 H, m), 1.75 - 1.83 (1.5 H, m), 1.88 (3.5 H, br. s.), 1.98 - 2.16 (1.5 H, m), 2.59 - 2.86 (4 H, m), 3.17 (4 H, s), 3.41 - 3.53 (3.5 H, m), 3.78 (2.5 H, m), 3.88 - 4.02 (3.5 H, m), 4.06 - 4.38 (1 H, m), 4.76 - 4.91 (1 H, m), 7.39 - 7.57 (2 H, m), 7.59 - 7.78 (1 H, m),
 5 7.98 (2 H, s), 8.31 - 8.45 (1 H, m)

Compound [383]: N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide


10 4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (66 mg, 0.15 mmol, 1 eq), DIPEA (52 μ l, 0.3 mmol, 2 eq) and TBTU (54 mg, 0.17 mmol, 1.1 eq) were added to 1 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-(4-benzyl-1,4-diazepan-1-yl)cyclohexanamine
 15 (Intermediate 16) (54 mg, 0.19 mmol, 1.27 eq) dissolved in DMF (0.5 mL). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_5) to provide *N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide* as a mixture cis and
 20 trans isomers (off-white solid, 33 mg, 31%).



Rt = 4.40 min (Analytical_1) MS(+ve) 707.6; MS(-ve) 705.7; ^1H NMR (DMSO- d_6) δ
 25 ppm: 0.73 (2 H, br. s.), 0.96 (2 H, br. s.), 1.22 - 2.22 (19 H, m), 2.64 - 2.91 (4 H, m), 3.23 (5 H, s), 3.53 (8 H, s), 3.89 - 4.24 (4 H, m), 4.91 (1 H, t, $J=8.5$ Hz), 5.82 (0 H, s), 7.23 - 8.54 (10 H, m).

Compound [384]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide


4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (66 mg, 0.15 mmol, 1 eq), DIPEA (52 μ l, 0.3 mmol, 2 eq) and TBTU (54 mg, 0.17 mmol, 1.1 eq) were added to 1 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-(4-methylpiperazin-1-yl)cyclohexanamine (Intermediate 17) (35 mg, 0.18 mmol, 1.2 eq) dissolved in DMF (0.5 mL). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_2 and Preparative_3) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide (white solid, 17 mg, 18%).



Rt = 2.85 min (Analytical_1) MS(+ve) 617.5; MS(-ve) 615.6; ^1H NMR (DMSO- d_6) δ ppm: 0.61 - 0.76 (2 H, m), 0.89 (1 H, t, $J=6.8$ Hz), 0.92 - 0.98 (2 H, m), 1.22 - 1.47 (5 H, m), 1.47 - 1.79 (6 H, m), 1.81 - 2.02 (5 H, m), 2.07 - 2.20 (3 H, m), 2.20 - 2.42 (4 H, m), 3.20 (4 H, s), 3.34 (3 H, s), 3.51 (2 H, s), 3.67 - 3.82 (1 H, m), 3.98 (3 H, s), 4.88 (1 H, quin, $J=8.5$ Hz), 7.42 - 7.59 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.08 (1 H, d, $J=7.8$ Hz), 8.42 (1 H, d, $J=8.3$ Hz).

Compound [385]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide

From the synthesis and purification of Compound [384] was isolated as a separate compound 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide (white solid, 34 mg, 37%).




Rt = 2.93 min (Analytical_1) MS(+ve) 617.5; MS(-ve) 615.6; ¹H NMR (DMSO-*d*₆) δ ppm: 0.62 - 0.80 (2 H, m), 0.84 - 1.02 (4 H, m), 1.18 - 1.36 (4 H, m), 1.43 - 1.84 (12 H, m), 1.84 - 2.03 (4 H, m), 2.09 - 2.22 (4 H, m), 2.41 (4 H, br. s.), 3.21 (3 H, s), 3.52 (2 H, s), 3.89 - 3.97 (1 H, m), 3.99 (3 H, s), 4.89 (1 H, quin, *J*=8.5 Hz), 7.47 - 7.61 (2 H, m), 7.72 (1 H, s), 8.03 (1 H, s), 8.08 (1 H, d, *J*=7.3 Hz), 8.43 (1 H, d, *J*=9.3 Hz)

Compound [386]: N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (66 mg, 0.15 mmol, 1 eq), DIPEA (52 μl, 0.3 mmol, 2 eq) and TBTU (54 mg, 0.17 mmol, 1.1 eq) were added to 1 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-(4-benzylpiperazin-1-yl)cyclohexanamine (Intermediate 18) (35 mg, 0.18 mmol, 1.2 eq) dissolved in DMF (0.5 mL). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_2) to provide N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide (white solid, 13 mg, 13%).


109



Rt = 3.64 min (Analytical_1) MS(+ve) 693.6; MS(-ve) 691.7; ^1H NMR (DMSO- d_6) δ ppm: 0.62 - 0.79 (2 H, m), 0.85 - 1.02 (2 H, m), 1.20 - 1.46 (4 H, m), 1.46 - 1.58 (2 H, m), 1.59 - 1.67 (2 H, m), 1.67 - 1.78 (2 H, m), 1.80 - 2.02 (6 H, m), 2.12 (1 H, s), 2.18 - 2.46 (6 H, m), 3.20 (3 H, s), 3.43 - 3.56 (5 H, m), 3.67 - 3.84 (1 H, m), 3.97 (3 H, s), 4.88 (1 H, quin, $J=8.4$ Hz), 7.22 - 7.42 (6 H, m), 7.44 - 7.59 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.08 (1 H, d, $J=7.8$ Hz), 8.42 (1 H, d, $J=7.8$ Hz)

10 *Compound [387]: N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide*


From the synthesis and purification of Compound [386] was isolated as a separate compound *N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide* (white solid, 20 mg, 19%).



20 Rt = 3.85 min (Analytical_1) MS(+ve) 693.6; MS(-ve) 691.7; ^1H NMR (DMSO- d_6) δ ppm: 0.62 - 0.79 (2 H, m), 0.87 - 1.01 (2 H, m), 1.43 - 1.59 (5 H, m), 1.59 - 1.67 (2 H, m), 1.66 - 1.74 (2 H, m), 1.75 - 1.84 (2 H, m), 1.92 (4 H, br. s.), 2.07 - 2.29 (2 H, m), 2.41 (4 H, br. s.), 3.20 (3 H, s), 3.51 (4 H, s), 3.87 - 4.05 (4 H, m), 4.74 - 4.99 (1 H, m), 7.24 - 7.39 (5 H, m), 7.50 - 7.56 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.07 (1 H, d, $J=6.8$ Hz), 8.42 (1 H, d, $J=8.8$ Hz)

Compound [388]: 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide

4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid (Intermediate 6) (66 mg, 0.15 mmol, 1 eq), DIPEA (52 μ l, 0.3 mmol, 2 eq) and TBTU (54 mg, 0.17 mmol, 1.1 eq) were added to 1.5 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-morpholinocyclohexanamine (Intermediate 16) (33 mg, 0.18 mmol, 1.2 eq). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_2) to provide 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide (off-white solid, 10 mg, 11%).



15

Rt = 3.16 min (Analytical_1) MS(+ve) 606.4; MS(-ve) 604.5; ^1H NMR (DMSO- d_6) δ ppm: 1.07 - 1.18 (7 H, m), 1.24 - 1.49 (4 H, m), 1.58 - 1.71 (4 H, m), 1.71 - 1.84 (2 H, m), 1.83 - 2.03 (6 H, m), 2.23 (1 H, t, $J=10.5$ Hz), 2.43 - 2.51 (3 H, m), 3.17 - 3.28 (4 H, m), 3.42 (3 H, br. s.), 3.60 (4 H, br. s.), 3.64 (1 H, br. s.), 3.71 - 3.83 (1 H, m), 3.98 (3 H, s), 5.23 (1 H, t, $J=8.3$ Hz), 7.44 - 7.60 (2 H, m), 7.72 (1 H, s), 8.02 (1 H, s), 8.09 (1 H, d, $J=7.8$ Hz), 8.40 (1 H, d, $J=8.3$ Hz)

20


Compound [389]: 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide

25

From the synthesis and purification of Compound [388] was isolated as a separate compound 4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-

111

b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide
(white solid, 22 mg, 24%).



5

Rt = 3.31 min (Analytical_1) MS(+ve) 606.4; MS(-ve) 604.5; ¹H NMR (DMSO-*d*₆) δ ppm: 1.13 (6 H, s), 1.39 - 1.60 (4 H, m), 1.60 - 1.71 (4 H, m), 1.71 - 1.85 (4 H, m), 1.85 - 2.01 (4 H, m), 2.07 - 2.22 (2 H, m), 2.42 - 2.48 (4 H, m), 3.22 (4 H, s), 3.64 (4 H, br. s.), 3.87 - 4.05 (4 H, m), 5.14 - 5.31 (1 H, m), 7.45 - 7.62 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.09 (1 H, d, *J*=7.3 Hz), 8.39 (1 H, d, *J*=8.3 Hz)

10

*Compound [390]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-*b*][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide*


15

4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-*b*][1,4]diazepine]-2'-ylamino)-3-methoxybenzoic acid (Intermediate 5) (66 mg, 0.15 mmol, 1 eq), DIPEA (52 μl, 0.3 mmol, 2 eq) and TBTU (54 mg, 0.17 mmol, 1.1 eq) were added to 1.5 mL DMF and the resulting solution was stirred at rt for 20 min before the addition of 4-morpholinocyclohexanamine (Intermediate 16) (33 mg, 0.18 mmol, 1.2 eq). The RM was then stirred at rt for 2 hours before purifying by preparative RP-HPLC-MS (Preparative_2) to provide 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-*b*][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide (off-white solid, 13 mg, 14%).

20

25


112



Rt = 2.96 min (Analytical_1) MS(+ve) 604.4; MS(-ve) 602.5; ^1H NMR (DMSO- d_6) δ ppm: 0.72 (2 H, s), 0.95 (2 H, s), 1.24 - 1.49 (4 H, m), 1.49 - 1.60 (2 H, m), 1.60 - 1.69 (2 H, m), 1.69 - 1.82 (2 H, m), 1.82 - 2.03 (6 H, m), 2.14 (1 H, s), 2.17 - 2.32 (1 H, m), 3.22 (4 H, s), 3.53 (2 H, s), 3.61 (4 H, br. s.), 3.71 - 3.84 (1 H, m), 4.00 (3 H, s), 4.90 (1 H, quin, $J=8.5$ Hz), 7.43 - 7.61 (2 H, m), 7.73 (1 H, s), 8.04 (1 H, s), 8.10 (1 H, d, $J=7.8$ Hz), 8.44 (1 H, d, $J=8.3$ Hz)

10 *Compound [391]: 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide*

From the synthesis and purification of Compound [390] was isolated as a separate compound 4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide (white solid, 27 mg, 30%).



20 Rt = 3.10 min (Analytical_1) MS(+ve) 604.4; MS(-ve) 602.5; ^1H NMR (DMSO- d_6) δ ppm: 0.66 - 0.81 (2 H, m), 0.88 - 1.03 (2 H, m), 1.45 - 1.86 (12 H, m), 1.87 - 2.02 (4 H, m), 2.10 - 2.21 (1 H, m), 2.47 (4 H, br. s.), 3.22 (3 H, s), 3.52 (2 H, s), 3.65 (4 H, br. s.), 3.89 - 4.05 (4 H, m), 4.90 (1 H, quin, $J=8.5$ Hz), 7.50 - 7.63 (2 H, m), 7.73 (1 H, s), 8.03 (1 H, s), 8.10 (1 H, d, $J=7.3$ Hz), 8.43 (1 H,

EXAMPLE 3

A full-length human CDC25C clone was isolated by PCR from HeLa mRNA and was inserted on a BamHI-HindIII fragment into pRsetA. The amino terminal fragment of CDC25C (encoding residues 1-300) was excised from this vector and inserted into pET28a (between the NcoI and BamHI sites). Expression was under the control of the T7 promoter, and the encoded protein contained a His₆ tag at the carboxyl terminus. The vector was transformed into *E. coli* strain BRL(DE3) pLysS for expression experiments. CDC25C was expressed in BL21(DE3) RIL bacteria cells, grown in LB media at 37 °C until OD_{600nm} of 0.6 was reached. The expression was induced with 1 mM IPTG and the bacterial culture was grown further for 3 h. The bacteria were harvested by centrifugation and the cell pellet was re-suspended in 50 mM Tris pH 7.5 and 10 % sucrose, flash frozen, and stored at -70 °C until used. CDC25C protein was purified from *E. coli* inclusion bodies. The inclusion bodies were isolated in a buffer (50 mM Tris pH 8.0, 2 mM EDTA, 100 mM NaCl, 0.5 % triton X-100). After denaturation in the presence of 6 M urea, the protein was refolded via slow dialysis of the urea. The protein was stored in 25 mM Tris pH 8.0, 100 mM NaCl, 1 mM DTT, 1 mM EDTA and 10 % glycerol at -70 °C until used.

A full-length human PLK1 (XM_047240) (amino acids 1-603) clone was amplified from a foetal lung cDNA library using primers incorporating restriction enzyme sites. The 5' primer (gccgctagcgacgatgacgataagatgagtgcctgcagtgcagggaagc) had an NheI site prior to the ATG start codon. The 3' primer (ggaattcttaggagccttgagacgg) incorporated a stop codon prior to the EcoRI site. The PCR product was cloned into the NheI/EcoRI sites of a baculovirus expression vector, pSSP1. Cloning into this vector resulted in a His₆ tag fusion at the amino terminus of the PLK1 construct. Sf9 cells of a passage number less than 20 were split back to give a 300 mL culture volume, at a cell density of 1.5 x 10⁶ cells/mL. Cells were only used for expression in logarithmic growth phase. PLK1 baculovirus (from P2 amplification) was added to give a multiplicity of infection of 3, this is equivalent to 3 virus particles for each insect cell. The flasks were incubated at 27 °C, with shaking at 100 r.p.m. for 48 h. On harvest, cell density and viability was determined, the cultures spun down at 2500 r.p.m. for 5 min

and washed with ice-cold phosphate-buffered saline. The wash was re-spun at the same speed and the pellet was snap frozen. PLK1 protein was purified on a metal affinity column. The insect cell pellet was lysed in a buffer (10 mM Tris-HCl pH 8.0, 150 mM NaCl, 5 mM β -mercaptoethanol, 1 mM PMSF, 1 mM benzamidine, 20 mM imidazole and protease inhibitor cocktail (Sigma) and the pre-cleared supernatant was loaded onto NiNTA-agarose (Qiagen). The affinity column was washed with the lysis buffer and the bound protein was eluted with 250 mM imidazole in the same buffer. After overnight dialysis against 25 mM Tris HCl, pH 7.5, 100 mM NaCl, 1 mM DTT, 1 mM PMSF, 1 mM benzamidine, protease inhibitors cocktail (Sigma) and 10 % glycerol, the purified protein was stored at -70°C until used.

EXAMPLE 4

PLK1 protein kinase assays were carried out using a 96-well plate format by incubating CDC25C (2 $\mu\text{g}/\text{well}$) with PLK1 (1 $\mu\text{g}/\text{well}$) in 20 mM Tris/HCl buffer pH 7.5, supplemented with 25 mM β -glycerophosphate, 5 mM EGTA, 1 mM DTT and 1 mM NaVO_3 . Serial dilutions of test compound in assay buffer were added. Reaction was initiated by the addition of 100 μM ATP and 0.5 μCi of $[\gamma\text{-}^{32}\text{P}]\text{-ATP}$. The reaction mixture was incubated at 30°C for 1 h, then stopped with 75 mM aq orthophosphoric acid, transferred onto a 96-well P81 filter plate (Whatman), dried, and the extent of CDC25C phosphorylation was assessed by scintillation counting using a Packard TopCount plate reader. The raw assay data was analysed by non-linear regression analysis parameters and IC_{50} values were determined using the equation: $y = A + ((B - A)/(1 + ((C/x)^D)))$, where y is % inhibition, A is minimum inhibition, B is maximum inhibition, C is EC_{50} , and D is the slope factor.

25

Cellular proliferation assays using human tumour cell lines (obtained from the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, USA) were carried out. Standard 72-h MTT (thiazolyl blue; 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) assays were performed (Loveland et al, Biochem. Int., 1992, 27, 501; Haselsberger et al, Anti Cancer Drugs, 1996, 7, 331). In short: cells were seeded into 96-well plates according to doubling time

30

and incubated overnight at 37 °C. Test compounds were made up in DMSO and a 1/3 dilution series prepared in 100 µL cell media, added to cells (in triplicates) and incubated for 72 h at 37 °C. MTT was made up as a stock of 5 mg/mL in cell media and filter-sterilised. Media was removed from cells followed by a wash with 200 µL PBS. MTT solution was then added at 20 µL per well and incubated in the dark at 37 °C for 4 h. MTT solution was removed and cells again washed with 200 µL PBS. MTT dye was solubilised with 200 µL per well of DMSO with agitation. Absorbance was read at 540 nm and data analysed using curve-fitting software (GraphPad Prism version 3.00 for Windows, GraphPad Software, San Diego California USA) to determine IC₅₀ values (concentration of test compound which inhibits cell growth by 50%).

EXAMPLE 5

PLK Selectivity


Table 4 shows the selectivity of compound [218] for PLK1 over PLK2 and PLK3 compared with selected compounds of the prior art. By way of illustration, compound [218] of the invention is twice as selective for PLK1 versus PLK 2 compared to compound [I-64] of WO 07/095188, *ca* 6-fold more selective than compound [I-76] and 7-fold more selective than compound [I-4] of WO 07/095188. Similarly, compound [218] is 5-fold more selective for PLK1 versus PLK3 compared to compound [I-64] of WO 07/095188, *ca* 12-fold more selective than compound [I-76] and 5-fold more selective than compound [I-4] of WO 07/095188.

EXAMPLE 6

Solubility Studies

Table 5 shows that compound [254] (boxed structure below) has superior solubility and/or pharmacokinetic properties when compared to structurally related compounds already known in the art.

116



Nephelometry Studies

If a DMSO solution of a water soluble compound is introduced into aqueous buffer, the mixture remains clear unless the aqueous solubility limit is reached. Above the solubility limit, precipitation occurs and light travelling through the turbid suspension is scattered. The measurement of the intensity of the scattered light is the basis of nephelometric solubility determination.

Selected compounds were screened using a nephelometer to identify the concentration at which turbidity occurs, thereby giving an indication of kinetic solubility. By nephelometry (2% DMSO : 98% phosphate buffered saline pH 7.5) compound [254] was freely soluble at 200 μ M (>0.11 mg/mL) which is the highest concentration tested in this assay. In contrast, in the same nephelometry assay, the maximum soluble concentration of prior art Example [I-253] of WO 07/095188 was determined to be 75 μ M (Table 6). This shows that compound [254] is more soluble in aqueous buffer in the concentration range used to study compound activity *in vitro*.

Pharmacokinetic Studies

In order to study compounds *in vivo*, higher concentrations are often required. To achieve these higher concentrations, salt forms of compounds are normally utilised.

Compound [254] and Example [I-76] of WO 07/095188 were studied as the free base and hydrochloride salts in various pharmaceutically relevant solvent mixtures that can be used for administration of compounds to living organisms. Having solubilized the compound by heating in the solvent, the solutions were cooled to 25 °C and observed.

Rather than forming a precipitate, supersaturated mixtures tend to form a gelatinous mesophase after cooling that is not suitable or convenient for dosing. It is clear from the data presented in Table 5 that compound [254] and its HCl salt have consistently better solution profiles than Example [I-76] and its corresponding HCl salt.

5

The HCl salt of Example [I-76] was administered to mice as a solution in 3% DMA 97% water by intravenous injection. The maximum tolerated dose appears to be dictated by reaching the maximum concentration of compound in stable solution. At the maximum tolerated dose no xenograft antitumour efficacy was observed (qd x 7 dosing) as insufficient compound concentration could be administered. In contrast, compound [254] and its salt, having significantly higher water solubility is able to be injected at concentrations more likely to produce efficacy in the treatment of diseases.

10

EXAMPLE 8

15 Pharmacokinetic Properties of compound [371]

(a) Methodology

Preliminary investigations of the pharmacokinetics of selected compounds were carried out in fasted conscious CD-1 mice (25-30 g). Compounds were dosed as 1 mg/kg by intravenous administration or 5-10 mg/kg by oral route. Twenty one male mice per administration route (n=3 mice per time point) received dose solution by single bolus injection in the tail vein or by oral gavage. Blood samples were collected by cardiac puncture at 5, 15, 30, 60, 120, 240 and 360 min following iv administration and at 30, 60, 120, 240, 360, 480 and 1440 min following oral administration. Blood samples were collected into tubes containing the anti-coagulant lithium heparin, mixed and placed promptly on crushed ice. Plasma was derived from whole blood by centrifugation, frozen and stored at -80 °C prior to analysis. Plasma concentrations were determined by liquid chromatography-tandem mass spectrometry in electrospray positive mode (ESI LC/MS/MS) using calibration curves prepared in the appropriate matrix. The lower level of quantification (LLOQ) in all cases was ca. 5 ng/mL. Plasma concentration-time data was processed using the computer program

20

25

30

WinNonLin version 5.2 from which areas under the plasma concentration-time curve (AUC_{all}) were calculated by non-compartmental analysis using Model 201 (IV-bolus Input) or Model 200 (Extravascular Input).

5 (b) Comparison of compounds [371] and [378] with selected prior art compounds

Further pharmacokinetic studies were carried to compare compounds [371] and [378] with a series of related analogues (see Table 7; compound [371] corresponds to compound H; compound [378] corresponds to compound I).

10 Figure 1 plots the area under the curve (AUC) for plasma concentration vs time for each compound tested. The dosing vehicle across these experiments remained constant (intravenous = citrate buffer pH 3, 1 mL/kg; oral = DMA/PEG400/10mM Tartrate buffer, pH 4 (1:3:6), 5 ml/kg). Figure 2 represents the same data in the form of oral bioavailability (%F) for each compound.

15

The results indicate that compounds H and I exhibit markedly superior systemic exposure and oral bioavailability compared to compounds A-G.

EXAMPLE 9

20 (a) Cellular Efflux Data

In vitro cell viability assays have been used to measure the effect of drug accumulation in paired parental and drug resistant tumour cell lines. The pairs used are the human ovarian carcinoma A2780 line with its MDR counterpart A2780/ADR and the human uterine sarcoma MES-SA line with its MDR counterpart MES-SA/Dx5 [Wesolowska
25 O., et al. (2005) *Anticancer Res.*, 25, 383-389].

Method: Cultivated human tumour cells of the ovarian cell line A2780 and its doxorubicin-resistant pair, A2780/ADR and the uterine cell line MES-SA and its doxorubicin-resistant pair, MES-SA/DX5 were seeded at 3000 cells per well in Nunc
30 96-well tissue culture plates, 100µl per well in DMEM media containing 10% foetal calf serum and penicillin/streptomycin. The wells of the first column in each plate

were filled with 100µl DMEM media instead of cells to provide a blank control for the Alamar Blue. The cells were incubated overnight at 37°C and 5% CO₂. The compounds of interest, dissolved in DMSO, were added on to the cells at various concentrations. Dilutions of the compounds were first prepared in DMEM at twice the
5 final desired concentration and 100µl of the dilutions added on to the cells. Each set of compounds were tested in triplicate in each cell line. After 72 hours incubation, 20µl Alamar Blue reagent (AbD Serotec) were added to each well and incubated for 3 hours. The plates were then read at 544/595nm.

10 The amount of Alamar Blue reagent reacted represents the metabolic activity of the cells. The relative cell activity was calculated as a percentage of the control (cells without compound) and the active compound concentration which inhibited the cell activity by 50% (IC₅₀) was derived. The values were calculated from the average of three individual measurements with blank correction (medium-only control).

15 The assay was performed on the Biomek FxP, automation platform from Beckman Coulter. The plates were read on the Paradigm plate reader supplied by Beckman Coulter. Asymmetry ratios were calculated as the IC₅₀ value of the compound in the drug resistant cells divided by the IC₅₀ value in the parental cells. Figure 3 shows a
20 comparison of selected compounds versus prior art compounds A'-J' (defined in Table 8) in parental and MDR tumor cells. Asymmetry ratios are presented relative to compound [378].

(b) Inhibition of PLK1

25 In addition, compounds of this invention which exhibit low MDR/parental cell asymmetry in cell viability assays are also more effective at inhibiting PLK1 within MDR tumour cells than compounds of the prior art. Thus, they have greater potential as therapeutic PLK inhibitors. The intermediate filament protein vimentin is a cellular target of PLK1 phosphorylation. Inhibition of PLK1 is associated with a decrease in
30 phospho-vimentin. This has been measured in the A2780 - A2780/ADR cell line pair.

Method: A2780 cells and A2780/ADR cells were plated in separate 96-well plates (Perkin Elmer) at a density of 20 000 cells/well and incubated overnight at 37°C and 5% CO₂. Test compounds were added to cells at a range of concentrations, with triplicate wells for each concentration (top concentration of DMSO on cells was 0.1%).

5 After 7hrs incubation, the cells were fixed in ice-cold 3.7% formaldehyde for 10mins. The cells were washed in PBS, then permeabilised in cold methanol for 10mins. The cells were washed again in PBS then incubated with PBS containing 0.1% Triton X-100 for 5mins. Cells were washed once with PBS containing 1% bovine serum albumin (Sigma) then incubated for 3hrs with antibodies to phospho-histone H3-Ser10

10 (1:4000; Millipore) and phospho-vimentin-Ser82 (1:1000; MBL), in 1% bovine serum albumin. Antibodies were detected with goat anti-mouse IgG conjugated to Alexafluor 488 (Invitrogen) and goat anti-rabbit IgG conjugated to Alexafluor 546 (Invitrogen), in 1% bovine serum albumin with 300nM DAPI (Cambridge Biosciences). Plates were scanned on a Cellomics Arrayscan II HCS Reader (Cellomics) using a 10X objective.

15 The Cell Health Profiling V2 Bioapplication (Cellomics) was used to acquire and analyse the images. Phospho-vimentin staining, at intensities above a user-defined threshold, was measured in mitotic cells only (defined by the presence of phospho-histone H3) and the average staining intensity per mitotic cell for each well was reported. IC₅₀ values were derived using average values from three replicate wells.

20 Asymmetry ratios were calculated as the IC₅₀ value of the compound in the drug resistant A2780/ADR cells divided by the IC₅₀ value in the parental A2780 cells. Figure 3 shows a comparison of selected compounds versus prior art compounds A'-J' (defined in Table 8) in parental and MDR tumor cells. As before, asymmetry ratios are presented relative to compound [378].












25

Various modifications and variations of the described aspects of the invention will be apparent to those skilled in the art without departing from the scope of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly

30 limited to such specific embodiments. Indeed, various modifications of the described modes of carrying out the invention which are obvious to those skilled in the

relevant fields are intended to be within the scope of the following claims.

TABLE 1: PLK IC₅₀ values (μM) for selected compounds of the invention; ****denotes <0.1 μM IC₅₀; ** denotes <1.0 μM IC₅₀; * denotes <10.0 μM IC₅₀*

		AA	PLK1	
1		*	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
2		***		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
3		*		4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
4		*		4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
5		*		4-(9-Cyclohexyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
6				4-(9-Cyclopentyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
7		*	**	4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid
8		***	**	4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid
9		*		4-(9-Cyclohexyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
10		*		9-Cyclohexyl-2-(4-hydroxy-phenylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one
11		*		9-Cyclopentyl-2-(4-hydroxy-phenylamino)-7-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one

123

AA PLK1

12				9-Cyclopentyl-2-(4-hydroxy-phenylamino)-8-isopropyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one
13		*		9-Cyclopentyl-2-(4-hydroxy-phenylamino)-5,8-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one
14		-		4-(9-Benzyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoic acid
15		**	*	4-(5-Methyl-6-oxo-9-phenyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid
16				4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid
17		*		4-(9-Cyclopentyl-7-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
18		*	*	4-(9-Cyclopentyl-7-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide
19		*		4-(9-Cyclopentyl-8-isopropyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
20		***		4-(9-Cyclopentyl-5,7-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
21				4-[9-(1-Ethyl-propyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide

124

AA PLK1

22			4-[9-(1-Ethyl-propyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide
23		*	4-(9-Cyclopentyl-8-isopropyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
24		*	4-[9-(1-Ethyl-propyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
25			3-Methoxy-N-(1-methyl-piperidin-4-yl)-4-[6-oxo-9-(tetrahydro-pyran-4-yl)-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzamide
26		**	3-Methoxy-4-[5-methyl-6-oxo-9-(tetrahydro-pyran-4-yl)-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide
27			4-[5-Ethyl-9-(1-ethyl-propyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
28		**	4-(9-Cyclopentyl-5-ethyl-7-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
29		- *	4-(9-Benzyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
30		**	3-Methoxy-4-(5-methyl-6-oxo-9-phenyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide
31		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide

125

AA PLK1
























32		4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
33		** 4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
34		** 4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
35		** 4-(9-Cyclohexyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
36		* 4-(9-Cyclopentyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide
37		** ** 4-(9-Cyclopentyl-5,8-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide
38		** 4-(9-Cyclopentyl-8-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide
39		* 4-(9-Cyclohexyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide




TABLE 2




Cpd		AurA IC50	PLK1 IC50	Name	Data
40			***	4-(9-Cyclopentyl-7-ethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	MS(+ve): 536.3; tR = 3.33 min (xbridge 4).
41		***	**	4-(9-Cyclopentyl-7-ethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid	MS(+ve): 410.2; tR = 2.28 min (xbridge 4).
42		**	*	4-(9-tert-Butyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	MS(+ve): 496.3; tR = 2.88 min (xbridge 4).
43		**		4-(9-tert-Butyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid	MS(+ve): 370.1; tR = 2.08 min (xbridge 4).





44		*		4-[9-(3,4-Dimethoxy-benzyl)-5,7-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	MS(+ve): 604.4; tR = 2.87 min (xbridge 4).
45		*		4-[9-(3,4-Dimethoxy-benzyl)-5,7-dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzoic acid	MS(+ve): 478.2; tR = 2.12 min (xbridge 4).
46		***	*	9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.56 (4H, m, CH ₂), 1.70 (2H, m, CH ₂), 1.89 (2H, m, CH ₂), 2.65 (2H, m, CH ₂), 3.16 (3H, s, NCH ₃), 3.67 (2H, m, CH ₂), 4.45 (4H, d, J 10Hz, CH ₂), 4.84 (1H, m, CH), 7.33 (1H, d, J 8Hz, Ar-H), 7.57 (1H, d, J 8Hz, Ar-H), 7.71 (1H, s, Ar-H), 8.05 (1H, s, Ar-H), 9.96 (1H, s, NH); MS (+ve): 428.3; tR = 6.54 min (XBridge 2).




47		***	*	9-Cyclopentyl-5-methyl-2-[4-(morpholine-4-sulfonyl)-phenylamino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.60 – 1.99 (8H, m, alkyl-H), 2.50 (4H, bs, morph-H), 2.58 – 2.60 (2H, m, CH ₂), 2.83 (4H, bs, morph-H), 3.18 (3H, s, CH ₃), 3.31 (2H, J 4.5 Hz, CH ₂), 4.79 – 4.84 (1H, m, CH), 7.60 (2H, d, J 8 Hz, Ar-H), 7.98 (2H, d, J 8 Hz, Ar-H), 8.11 (1H, s, Ar-H), 9.77 (1H, bs, NH); MS(+ve): 487.31; tR = 3.07 min (Xbridge 4).
48			*	3-Methoxy-4-[9-(3-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.58 (2H, dd, J 12.5 Hz, 22Hz, CH), 1.74 (2H, d, J 12Hz, CH), 1.90 (2H, dd, J 11.5, 11.5Hz, CH), 2.16 (3H, s, N-CH ₃), 2.79 (4H, m, CH + CH ₂), 3.29 (3H, s, N-CH ₃), 3.69 (1H, m, CH), 3.70 (3H, s, OCH ₃), 3.88 (3H, s, OCH ₃), 4.04 (2H, m, CH ₂), 6.88 (2H, m, Ar-H), 6.99 (1H, d, J 8Hz, Ar-H), 7.37 (2H, m, Ar-H), 7.54 (1H, d, J 8Hz, Ar-H), 7.56 (1H, s, NH), 8.00 (1H, d, J 7.5 Hz, Ar-H), 8.29 (1H, s, Pyr-H); MS(+ve): 546.4; tR = 2.58 min (xbridge 4).
49		*	*	4-[9-(3-Methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzoic acid	MS(+ve); 420.22; tR = 1.44 min (xbridge 4).
50		***	*	4-[9-(2-Fluoro-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzoic acid	MS(+ve); 408.2; tR = 2.59 min (xbridge 4).




51		**	**	9-Cyclopentyl-5-methyl-2-(4-morpholin-4-yl-phenylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.56 (4H, m, CH ₂), 1.68 (2H, m, CH ₂), 1.91 (2H, m, CH ₂), 2.55 (2H, m, CH ₂), 3.01 (4H, m, CH ₂), 3.14 (3H, s, NCH ₃), 3.57 (2H, m, CH ₂), 3.73 (4H, m, CH ₂), 4.77 (1H, m, CH), 6.85 (1H, d, J 9Hz, Ar-H), 7.55 (1H, d, J 9Hz, Ar-H), 7.99 (1H, s, Ar-H), 8.92 (1H, s, NH); MS (+ve): 423.3; tR = 7.09 min (XBridge 2); tR = 10.51 min (Vydac 1).
52		**		4-[9-(2-Fluoro-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.58 (2H, dd, J 9.5Hz, 21Hz, CH), 1.74 (2H, d, J 11.5Hz, CH), 1.92 (2H, dd, J 11Hz, 11Hz, CH), 2.16 (3H, s, N-CH ₃), 2.76 (2H, d, J 11Hz, CH), 2.86 (2H, m, CH ₂), 3.70 (1H, m, CH), 3.82 (3H, s, OCH ₃), 4.02 (2H, m, CH ₂), 6.92 (2H, d, J 8.5Hz, Ar-H), 7.32 (1H, d, J 8.5Hz, Ar-H), 7.37 (2H, m, Ar-H), 7.46 (1H, m, Ar-H), 7.56 (1H, m, Ar-H), 7.66 (1H, s, NH), 7.99 (1H, d, J 7.5Hz, Ar-H), 8.28 (1H, s, Ar-H); MS(+ve): 534.3; tR = 2.59 min (xbridge 4).
53		**	**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzenesulfonamide	¹ H NMR (DMSO): 1.41 – 1.75 (8H, alkyl-H), 2.45 (2H, bs, CH ₂), 2.99 (3H, s, CH ₃), 3.47 (2H, bs, CH ₂), 4.74 – 4.75 (1H, m, CH), 7.13 (2H, bs, NH ₂), 7.29 (2H, bs, Ar-H), 7.59 (1H, bs, Ar-H), 7.89 (1H, bs, Ar-H), 8.06 (1H, bs, Ar-H), 9.72 (1H, bs, NH); MS(+ve): 417.22; tR = 2.62 (Xbridge 4).
54		***	***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid	¹ H NMR (DMSO): 1.58 – 1.72 (6H, m, alkyl-H), 1.93 – 1.95 (2H, m, alkyl-H), 2.64 (2H, dd, J 4.5 Hz, CH ₂), 3.17 (3H, s, CH ₃), 3.67 (2H, dd, J 4.5 Hz, CH ₂), 4.83 – 4.89 (1H, m, CH), 7.78 (2H, dd, J 1.5 and 8 Hz, Ar-H), 7.87 (2H, d, J 8.5

						Hz, Ar-H), 8.09 (1H, d, J 1.5 Hz, Ar-H), 10.02 (1H, s, NH); MS(+ve): 382.22; tR = 1.60 (Xbridge 4).
55		***	*		9-Cyclopentyl-2-(4-methanesulfonylphenylamino)-5-methyl-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.61 – 1.99 (8H, m, alkyl-H), 2.59 (2H, dd, J 5 Hz, CH ₂), 3.14 (3H, s, CH ₃), 3.17 (3H, s, CH ₃), 3.63 (2H, dd, J 5 Hz, CH ₂), 4.82 – 4.86 (1H, m, CH), 7.76 (2H, d, J 9 Hz, Ar-H), 7.98 (2H, d, J 9 Hz, Ar-H), 8.11 (1H, s, Ar-H), 9.77 (1H, bs, NH); MS(+ve): 416.26; tR = 2.83 (Xbridge 4).
56		**	**		3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzoic acid	¹ H NMR (DMSO): 1.39 – 1.73 (8H, m, alkyl-H), 2.48 (2H, bs, CH ₂), 2.99 (3H, s, CH ₃), 3.49 (2H, bs, CH ₂), 4.77 (1H, bs, CH), 7.26 (1H, bs, Ar-H), 7.44 (1H, bs, Ar-H), 7.52 (1H, bs, Ar-H), 7.89 (1H, bs, Ar-H), 8.23 (1H, bs, Ar-H), 9.83 (1H, bs, NH); MS(+ve): 382.28; tR = 1.93 (Xbridge 4).
57		**	*		3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.54 – 1.64 (8H, m, alkyl-H), 1.74 – 1.76 (2H, m, alkyl-H), 1.90 – 2.01 (4H, m, alkyl-H), 2.18 (3H, s, CH ₃), 2.56 (2H, dd, J 5 Hz, CH ₂), 2.77 – 2.80 (2H, m, alkyl-H), 3.16 (3H, s, CH ₃), 3.59 (2H, dd, J 5 Hz, CH ₂), 3.69 – 3.73 (1H, m, CH), 4.79 – 4.84 (1H, m, CH), 7.27 – 7.33 (2H, m, Ar-H), 7.74 (1H, d, J 8 Hz, Ar-H), 8.04 (1H, 1H, J 1 Hz, Ar-H), 8.11 (1H, d, J 8 Hz, Ar-H), 8.17 (1H, s, Ar-H), 9.28 (1H, s, NH); MS(+ve): 478.33; tR = 2.60 (Xbridge 4).





58		***	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.50 (6H, m, CH), 1.70 (2H, m, CH), 1.81 (2H, m, CH), 1.98 (2H, m, CH), 2.07 (2H, m, CH), 2.16 (3H, s, NCH ₃), 2.57 (2H, m, CH ₂), 2.63 (2H, m, CH), 3.17 (3H, s, NCH ₃), 3.61 (2H, m, CH ₂), 3.75 (1H, m, CH), 3.89 (3H, s, OCH ₃), 4.86 (1H, , CH), 7.48 (1H, s, ArH), 7.54 (1H, dd, J 2Hz, 8.5Hz, NH), 7.73 (1H, d, J 8.5Hz, ArH), 7.83 (1H, d, J 7.5Hz, ArH), 8.09 (1H, s, pyrH), 9.44 (1H, s, NH); MS(+ve): 508.4; tR = 2.68 min (xbridge 4).
59			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-ethoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.44 (3H, t, J 6.5Hz), 1.59 (4H, m, CH), 1.76 (2H, m, CH), 1.93 (2H, m, CH), 2.16 (3H, s, N-CH ₃), 2.59 (2H, m, CH ₂), 2.78 (2H, d, J 11.5Hz, CH ₂), 3.17 (3H, s, CH ₃), 3.63 (2H, m, CH ₂), 3.73 (1H, m, CH), 4.20 (2H, q, J 7Hz, CH ₂ CH ₃), 4.79 (1H, m, CH), 7.48 (2H, m, Ar-H), 7.70 (1H, s, NH), 8.06 (2H, m, Ar-H), 8.38 (1H, d, J 8.5Hz, NH); MS(+ve): 522.4; tR = 2.88 min (xbridge 4).
60			**	9-Cyclopentyl-2-(2-methoxy-phenylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.57 (4H, m, CH), 1.68 (2H, m, CH), 1.90 (2H, m, CH), 2.57 (2H, m, CH ₂), 3.16 (3H, s, N-CH ₃), 3.62 (2H, m, CH ₂), 3.86 (3H, s, OCH ₃), 4.75 (1H, m, CH), 6.90 (1H, dd, J 8Hz, 8Hz, Ar-H), 6.95 (1H, dd, J 8Hz, 8Hz, Ar-H), 7.02 (1H, d, J 8Hz, Ar-H), 7.60 (1H, s, NH), 8.03 (1H, s, pyr-H), 8.20 (1H, d, J 8Hz, Ar-H); MS(+ve): 368.3; tR = 3.74 min (xbridge 4).




61		**	*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzenesulfonamide	¹ H NMR (DMSO): 1.57 (4H, m, CH ₂), 1.71 (2H, m, CH ₂), 1.93 (2H, m, CH ₂), 2.55 (2H, m, CH ₂), 2.70 (4H, m, CH ₂), 2.91 (4H, m, CH ₂), 3.17 (3H, s, NCH ₃), 3.63 (2H, m, CH ₂), 4.81 (1H, m, CH), 7.50 (1H, d, J 7 Hz, NH), 7.66 (2H, d, J 9 Hz, Ar-H), 7.89 (2H, d, J 9 Hz, Ar-H), 8.09 (1H, s, Ar-H), 9.67 (1H, s, NH); MS (+ve) 514.4; tR = 6.29 min (XBridge 2).
62			**	2-Benzylamino-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
63		***	*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzenesulfonamide	
64		***	***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.61 – 2.20 (16H, alkyl-H), 2.58 (2H, dd, J 5 Hz, CH ₂), 2.69 (3H, s, CH ₃), 3.31 (3H, s, CH ₃), 3.62 (2H, dd, J 5 Hz, CH ₂), 4.81 – 4.84 (1H, m, CH), 7.74 – 7.80 (2H, m, Ar-H), 8.08 (1H, s, Ar-H), 9.46 (1H, s, NH); MS(+ve): 478.40; tR = 2.56 (Xbridge 4).





65			***	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 0.29 (2H, d, J 4Hz, cyclopropyl CH), 0.52 (2H, d, J 4Hz, cyclopropyl CH), 1.16 (1H, m, CH), 1.58 (2H, dd, J 11.5, 20.5, CH), 1.75 (2H, m, CH), 1.95 (2H, dd, J 10Hz, 10Hz, CH), 2.16 (3H, s, NCH₃), 2.64 (2H, m, CH₂), 2.79 (2H, d, J 11.5Hz, CH), 3.18 (3H, s, NCH₃), 3.50 (2H, d, J 6.5Hz, CH₂), 3.73 (1H, m, CH), 3.79 (2H, m, CH₂), 3.94 (3H, s, OCH₃), 7.48 (2H, m, ArH), 7.71 (1H, s, ArH), 8.07 (2H, m, ArH), 8.40 (1H, d, J 9Hz, ArH); MS(+ve); 494.38; tR = 2.60 min (xbridge 4).
66			**	4-(9-Butyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 0.91 (3H, t, J 7Hz, CH₃), 1.34 (2H, m, CH₂), 1.62 (4H, m, CH₂), 1.75 (2H, m, CH₂), 1.93 (2H, d, J 11Hz, CH₂), 2.16 (3H, s, NCH₃), 2.61 (2H, m, CH₂), 2.77 (2H, d, J 11.5Hz, CH₂), 3.16 (3H, s, NCH₃), 3.56 (2H, m, CH₂), 3.70 (3H, m, CH₂ + CH), 3.94 (3H, s, OCH₃), 7.48 (2H, m, ArH), 7.70 (1H, s, ArH), 8.06 (1H, s, ArH), 8.08 (1H, d, J 8Hz, ArH), 8.34 (1H, d, J 8.5Hz, ArH); MS(+ve); 496.4; tR = 2.69 min (xbridge 4).
67			***	4-(9-Isopropyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 1.24 (6H, d, J 7Hz, CH₃), 1.61 (2H, ddd, J 3.5Hz, 12Hz, 18Hz, CH), 1.76 (2H, d, J 9.5Hz, CH), 1.93 (dd, J 9.5Hz, 12Hz, CH), 2.16 (3H, s, CH₂), 2.59 (2H, m, CH₂), 2.78 (2H, d, J 12Hz, CH), 3.16 (3H, s, NCH₃), 3.60 (2H, m, CH₂), 3.73 (1H, m, CH), 3.94 (3H, s, OCH₃), 4.80 (1H, m, CH), 7.48 (1H, d, J 1.5Hz, ArH), 7.51 (1H, d, J 8.5Hz, ArH), 7.68 (1H, s, ArH), 8.06 (2H, m, ArH), 8.1 (1H, d, J 8.5Hz, ArH);





						MS(+ve); 482.3; tR = 2.55 min (xbridge 4).
68				**	3-Methoxy-4-[5-methyl-9-(3-methyl-butyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 0.92 (6H, d, J 7Hz, CH₃), 1.54 (4H, m, CH₂), 1.62 (2H, d, J 9.5Hz, CH), 1.93 (2H, dd, J 9.5Hz, 12Hz, CH), 2.16 (3H, s, NCH₃), 2.60 (2H, m, CH₂), 2.78 (2H, d, J 11.5Hz, CH), 3.16 (3H, s, NCH₃), 3.67 (2H, m, CH₂), 3.71 (2H, m, CH₂), 3.73 (1H, m, CH), 3.94 (3H, s, OCH₃); MS(+ve); 510.4; tR = 2.85 min (xbridge 4).
69				**	4-(9-Cyclopentylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 1.23 (2H, m, CH₂), 1.49 (2H, m, CH₂), 1.59 (4H, m, 2 x CH₂), 1.69 (2H, m, CH₂), 1.75 (2H, m, CH₂), 1.93 (2H, dd, J 10Hz, 12Hz, CH₂), 2.16 (3H, s, NCH₃), 2.36 (1H, m, CH), 2.62 (2H, m, CH₂), 2.79 (2H, d, J 12Hz, CH₂), 3.29 (3H, s, NCH₃), 3.60 (2H, d, J 7.5Hz, CH₂), 3.73 (2H, m, CH₂), 3.94 (3H, s, OCH₃), 7.49 (2H, M, ArH), 7.70 (1H, S, ArH), 8.08 (2H, M, ArH), 8.32 (1H, d, J 8.5Hz, ArH); MS(+ve); 522.4; tR = 2.89 min (xbridge 4).
70		**		*	9-Cyclopentyl-5-methyl-2-phenylamino-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹H NMR (DMSO): 1.62 – 2.00 (8H, m, alkyl-H), 2.76 (2H, dd, J 5 Hz, CH₂), 3.27 (3H, s, CH₃), 3.76 (2H, dd, J 5 Hz, CH₂), 4.96 – 4.99 (1H, m, CH), 7.01 – 7.04 (1H, m, Ar-H), 7.16 (1H, t, J 7.5 Hz, Ar-H), 7.29 (1H, t, J 7.5 Hz, Ar-H), 7.37 (1H, t, J 7.5 Hz, Ar-H), 7.54 – 7.55 (2H, m, Ar-H), 7.89 (1H, bs, NH); MS(+ve); 338.26; tR = 3.52 (Xbridge 4).





71		***	9-Cyclopentyl-2-(2,4-dimethoxyphenylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.53 (4H, m, CH), 1.65 (2H, m, CH), 1.83 (2H, m, CH), 2.54 (2H, m, CH ₂), 3.14 (3H, s, NCH ₃), 3.55 (2H, m, CH ₂), 3.73 (3H, s, OCH ₃), 3.81 (3H, s, OCH ₃), 4.64 (9H, m, CH), 6.47 (1H, dd, J 3Hz, 9Hz, ArH), 6.61 (1H, d, J 3Hz, ArH), 7.51 (1H, s, NH), 7.86 (1H, d, J 9Hz, ArH), 7.97 (1H, s, PyrH); MS(+ve); 398.3; tR = 3.51 min (xbridge 4).
72		**	2-(4-Chloro-3-methyl-phenylamino)-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (CD ₃ OD): 1.62 – 1.99 (8H, m, alkyl-H), 2.39 (3H, s, CH ₃), 2.76 (2H, dd, J 5.5 Hz, CH ₂), 3.23 (3H, s, CH ₃), 3.77 (2H, dd, J 5.5 Hz, CH ₂), 4.96 – 5.00 (1H, m, CH), 7.33 – 7.34 (2H, m, Ar-H), 7.56 (1H, s, Ar-H), 7.90 (1H, s, Ar-H); MS(+ve); 386.23; tR = 4.13 (Xbridge 4).
73		***	2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-9-(2-fluorophenyl)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.82 (2H, m, CH ₂), 3.27 (3H, s, CH ₃), 4.0 (2H, m, CH ₂), 4.12 (2H, s, CH ₂), 4.30 (2H, s, CH ₂), 6.91 (1H, d, J 8.5Hz, CH), 7.16 (2H, m, 2CH), 7.36 (2H, m, 2CH), 7.43 (1H, m, CH), 7.56 (1H, m, CH), 8.28 (1H, s, CH), 9.50 (1H, s, NH); MS(+ve): 454.3; tR = 2.67 min (XBridge 4).
74		***	9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-7-ethyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 0.87 (3H, t, J 7.5Hz, CH ₂ CH ₃), 1.29-1.79 (9H, m, 4CH ₂ , CH), 2.08 (1H, m, CH), 2.59 (1H, m, CH ₂), 3.18 (3H, s, CH ₃), 3.38 (2H, d, J 6.5Hz, NCH ₂), 4.4 (4H, m, 2CH ₂), 4.77 (1H, m, CH), 7.24 (1H, d, J 8.5Hz, CH), 7.63 (1H, d, J 8.5Hz, CH), 7.83 (1H, s, CH), 8.08 (1H, s, CH), 9.38 (1H, s, NH); MS(+ve): 456.3; tR = 3.31 min (XBridge 4).





75		***		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-9-phenyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.78 (2H, m, CH ₂), 3.27 (3H, s, CH ₃), 4.03 (2H, m, CH ₂), 4.09 (2H, s, CH ₂), 4.30 (2H, s, CH ₂), 6.93 (1H, d, J 8.5Hz, CH), 7.19 (1H, d, J 8.5Hz, CH), 7.29 (4H, m, 4CH), 7.46 (2H, m, 2CH), 8.3 (1H, s, CH), 9.49 (1H, s, NH); MS(+ve): 436.3; tR = 2.66 min (XBridge 4).
76		***	**	9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5,7-dimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.01 (3H, d, J 6.5Hz, CH ₃), 1.4-1.85 (6H, m, 3CH ₂), 2.07 (1H, m, CH), 2.83 (1H, m, CH), 3.18 (3H, s, CH ₃), 3.3 (2H, d, CH ₂), 3.4 (1H, t, J 11.5Hz, CH), 4.3-4.5 (4H, m, 2CH ₂), 4.73 (1H, m, CH), 7.24 (1H, d, J 8.5Hz, CH), 7.64 (1H, dd, J 9, 2Hz, CH), 7.84 (1H, s, CH), 8.06 (1H, s, CH), 9.37 (1H, s, NH); MS(+ve): 442.3; tR = 3.06 min (XBridge 4).
77		**	**	9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5,7,7-trimethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.09 (6H, s, 2CH ₃), 1.58 (4H, m, 2CH ₂), 1.72 (2H, m, CH ₂), 1.86 (2H, m, CH ₂), 3.18 (3H, s, CH ₃), 3.35 (2H, s, CH ₂), 4.39 (2H, s, CH ₂), 4.43 (2H, s, CH ₂), 5.21 (1H, m, CHN), 7.23 (1H, d, J 8.5Hz, CH), 7.6 (1H, dd, J 8.5, 2Hz, CH), 7.83 (1H, s, CH), 7.95 (1H, s, CH), 9.30 (1H, s, NH); MS(+ve): 456.3; tR = 3.25 min (XBridge 4).
78		***	**	(9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane])	¹ H NMR (DMSO): 0.66 (2H, m, CH ₂), 0.89 (2H, m, CH ₂), 1.47 (2H, m, CH ₂), 1.56 (2H, m, CH ₂), 1.67 (2H, m, CH ₂), 1.88 (2H, m, CH ₂), 3.16 (3H, s, CH ₃), 3.45 (2H, s, CH ₂), 4.39 (2H, s, CH ₂), 4.42 (2H, s, CH ₂), 4.86 (1H, m, CHN), 7.23 (1H, d, J 8.5Hz, CH), 7.63 (1H, dd, J 8.5, 2Hz, CH), 7.84 (1H, s, CH), 7.95 (1H, s, CH), 9.31 (1H, s, NH); MS(+ve):




79		***	2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-9-(1-ethylpropyl)-5-methyl-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	454.3; tR = 3.03 min (XBridge 4). ¹H NMR (DMSO): 0.83 (6H, t, J 7.5Hz, 2CH₃), 1.58 (4H, m, 2CH₂), 2.61 (2H, m, CH₂), 3.17 (3H, s, CH₃), 3.46 (2H, m, CH₂), 4.39 (2H, s, CH₂), 4.44 (2H, s, CH₂), 4.66 (1H, m, CHN), 7.23 (1H, d, J 8.5Hz, CH), 7.68 (1H, dd, J 8.5, 2Hz, CH), 7.75 (1H, s, CH), 8.03 (1H, s, CH), 9.29 (1H, s, NH); MS(+ve): 430.3; tR = 2.88 min (XBridge 4).
80		**	9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-8-isopropyl-5-methyl-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	¹H NMR (DMSO): 0.72 (3H, d, J 7Hz, CH₃), 0.84 (3H, d, J 7Hz, CH₃), 1.26-2.4 (10H, m, 5CH₂), 2.54 (1H, m, CH), 3.14 (3H, s, CH₃), 3.85 (1H, m, CH), 4.2 (1H, m, CH), 4.39-4.5 (4H, m, 2CH₂), 7.25 (1H, d, J 8.5Hz, CH), 7.68 (1H, dd, J 8.5, 2Hz, CH), 7.84 (1H, s, CH), 8.24 (1H, s, CH), 9.49 (1H, s, NH); MS(+ve): 470.3; tR = 3.30 min (XBridge 4).
81		**	2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-9-(tetrahydro-pyran-4-yl)-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	¹H NMR (DMSO): 1.7 (2H, m, CH₂), 1.85 (2H, m, CH₂), 2.57 (2H, m, CH₂), 3.16 (3H, s, CH₃), 3.43 (2H, t, J 11Hz, CH₂), 3.62 (2H, m, CH₂), 3.97 (2H, m, CH₂), 4.39 (2H, s, CH₂), 4.46 (2H, s, CH₂), 4.58 (1H, m, CHN), 7.26 (1H, d, J 8.5Hz, CH), 7.63 (1H, s, CH), 7.73 (1H, dd, J 8.5, 2Hz, CH), 8.06 (1H, s, CH), 9.36 (1H, s, NH); MS(+ve): 444.3; tR = 2.35 min (XBridge 4).




82		**		9-Benzyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.69 (2H, m, CH ₂), 3.22 (3H, s, CH ₃), 3.62 (2H, m, CH ₂), 4.01 (2H, s, CH ₂), 4.31 (2H, s, CH ₂), 4.86 (2H, s, CH ₂ Ph), 7.04 (1H, d, J 8.5Hz, CH), 7.42-7.27 (6H, m, 6CH), 7.69 (1H, s, CH), 8.13 (1H, s, CH), 9.40 (1H, s, NH); MS(+ve): 450.3; tR = 2.78 min (XBridge 4).
83		***	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzamide	¹ H NMR (DMSO): 1.61 (4H, m, CH), 1.72 (2H, m, CH), 1.93 (2H, m, CH), 2.63 (2H, m, CH ₂), 3.17 (3H, s, N-CH ₃), 3.66 (2H, m, CH ₂), 4.85 (1H, m, CH), 7.18 (1H, bs, NH), 7.72 (2H, d, J 8.5Hz, Ar-H), 7.83 (2H, d, J 8.5Hz, Ar-H), 8.06 (1H, s, pyr-H), 9.77 (1H, bs, NH); MS(+ve): 381.3; tR = 2.45 min (xbridge 4).
84		*	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-benzoic acid	¹ H NMR (DMSO): 0.66 (2H, m, cyclopropyl-CH), 0.90 (2H, m, cyclopropyl-CH), 1.50 (2H, m, CH), 1.58 (2H, m, CH), 1.69 (2H, m, CH), 1.91 (2H, m, CH), 3.17 (3H, s, N-CH ₃), 3.47 (2H, s, CH ₂), 4.88 (1H, m, CH), 7.81 (4H, m, Ar-H), 7.99 (1H, s, NH), 9.56 (1H, s, pyr-H); MS(+ve): 408.3; tR = 1.81 min (xbridge 4).
85		***	*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N,N-dimethyl-benzamide	¹ H NMR (DMSO): 1.59 (4H, m, CH), 1.70 (2H, m, CH), 1.94 (2H, m, CH), 2.58 (2H, m, CH ₂), 2.96 (6H, s, CH ₃), 3.16 (3H, s, N-CH ₃), 3.62 (2H, m, CH ₂), 4.81 (1H, m, CH), 7.31 (2H, d, J 8.5Hz, Ar-H), 7.78 (2H, d, J 8.5Hz, Ar-H), 8.06 (1H, s, N-H), 9.41 (1H, s, pyr-H); MS(+ve): 409.3; tR = 2.71 min (xbridge 4).





86		***	*	N-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzyl]-acetamide	¹ H NMR (CD ₃ OD): 1.62 – 1.98 (8H, m, alkyl-H), 2.00 (3H, s, CH ₃), 2.80 (2H, dd, J 5 Hz, CH ₂), 3.27 (3H, s, CH ₃), 3.80 (2H, dd, J 5 Hz, CH ₂), 4.37 – 4.38 (2H, m, CH ₂), 4.98 – 5.02 (1H, m, CH), 7.36 (2H, d, J 8.5 Hz, Ar-H), 7.47 (2H, d, J 8.5 Hz, Ar-H), 7.83 (1H, s, Ar-H), 8.50 (1H, bs, NH); MS(+ve): 409.33; tR = 2.53 (Xbridge 4).
87		*	***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-methyl-benzamide	
88		*	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N,N-dimethyl-benzamide	
89		*	***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide	





90		**	*	N-[3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzyl]-acetamide	
91		**	*	2-(3-Aminomethyl-phenylamino)-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
92			*	9-Cyclopentyl-2-(4-methoxy-benzylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
93		***	*	N-Benzyl-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzenesulfonamide	1H NMR (CD ₃ OD): 1.68 – 1.81 (8H, m, alkyl-H), 2.69 – 2.71 (2H, m, CH ₂), 3.29 (3H, s, CH ₃), 3.73 – 3.75 (2H, m, CH ₂), 4.06 (2H, s, CH ₂), 7.21 – 7.27 (5H, m, Ar-H), 7.73 (2H, d, J 9 Hz, Ar-H), 7.84 (2H, d, J 9 Hz, Ar-H), 8.04 (1H, bs, NH); MS(+ve): 507.28; tR = 3.39 (Xbridge 4).





94			*	9-Cyclopentyl-5-methyl-2-[(thiophen-2-ylmethyl)-amino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (CD ₃ OD): 1.61 (4H, m, CH), 1.73 (2H, m, CH), 1.92 (2H, m, CH), 2.63 (2H, m, CH ₂), 3.23 (3H, s, N-CH ₃), 3.66 (2H, m, CH ₂), 4.71 (2H, s, CH ₂), 6.93 (6.93 (1H, m, Ar-H), 6.97 (1H, m, Ar-H), 7.23 (1H, m, Ar-H), 7.84 (1H, s, pyr-H); MS(+ve): 358.3; tR = 3.32 min (xbridge 4).
95			*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-phenyl-benzamide	1H NMR (CD ₃ OD): 1.51 – 2.00 (8H, m, alkyl-H), 3.28 (3H, s, CH ₃), 2.69 (2H, dd, J 5.5 Hz, CH ₂), 3.72 (2H, dd, J 5.5 Hz, CH ₂), 4.03 (3H, s, CH ₃), 5.04 – 5.08 (1H, m, CH) 7.12 – 7.16 (2H, m, Ar-H), 7.34 – 7.37 (3H, m, Ar-H), 7.65 (1H, dd, J 2 and 8.5 Hz, Ar-H), 7.70 – 7.72 (2H, m, Ar-H), 7.80 (1H, s, Ar-H), 8.82 (1H, d, J 2 Hz, Ar-H); MS(+ve): 487.35; tR = 3.46 (Xbridge 4).
96		***	*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-methyl-benzenesulfonamide	1H NMR (CD ₃ OD): 1.67 – 2.01 (8H, m, alkyl-H), 2.55 (3H, s, CH ₃), 2.80 (2H, dd, J 4.5 Hz, CH ₂), 3.29 (3H, s, CH ₃), 5.01 – 5.04 (1H, m, CH), 7.78 – 7.81 (2H, m, Ar-H), 7.84 – 7.86 (2H, m, Ar-H), 7.96 – 8.00 (2H, m, Ar-H and NH); MS(+ve): 431.32; tR = 2.81 (Xbridge 4).
97		***	*	N-Acetyl-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-methyl-benzenesulfonamide	1H NMR (CD ₃ OD): 1.68 – 2.04 (8H, m, alkyl-H), 2.34 (3H, s, CH ₃), 2.76 (2H, dd, J 5 Hz, CH ₂), 3.29 (3H, s, CH ₃), 3.31 (3H, s, CH ₃), 3.79 (2H, dd, J 5 Hz, CH ₂), 4.94 – 5.00 (1H, m, CH), 7.86 (2H, d, J 9 Hz, Ar-H), 7.91 (2H, d, J 9 Hz, Ar-H), 8.01 (1H, s, Ar-H); MS(+ve): 473.35; tR = 3.15 (Xbridge 4).





98		**	**	9-Cyclopentyl-2-(1H-indol-7-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹H NMR (DMSO): 1.50 – 1.86 (8H, m, alkyl-H), 2.77 (2H, dd, J 4.5 Hz, CH₂), 3.23 (3H, s, CH₃), 3.74 (2H, dd, J 4.5 Hz, CH₂), 4.87 – 4.91 (1H, m, CH), 6.63 (1H, s, Ar-H), 7.16 (1H, t, J 8 Hz, Ar-H), 7.33 (1H, d, J 8 Hz, Ar-H), 7.42 – 7.47 (3H, m, Ar-H), 8.03 – 8.04 (1H, m, Ar-H), 9.94 (1H, bs, NH), 11.31 (1H, bs, NH); MS(+ve): 377.28; tR = 3.13 (Xbridge 4).
99		**		4-(9-Isopropyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 0.68 (2H, m, cyclopropyl CH), 0.92 (2H, m, cyclopropyl CH), 1.17 (6H, d, J 6.5 Hz, CH₃), 1.57 (2H, ddd, J 3.5 Hz, 12 Hz, 15 Hz, CH), 1.77 (2H, d, J 10 Hz, CH), 1.95 (2H, dd, J 10 Hz, 10 Hz, CH), 2.16 (3H, s, NCH₃), 2.79 (2H, d, J 11.5 Hz, CH), 3.18 (3H, s, NCH₃), 3.46 (2H, s, CH₂), 3.77 (1H, m, CH), 3.46 (3H, s, OCH₃), 4.84 (1H, m, CH), 7.48 (1H, s, Ar-H), 7.51 (1H, d, J 8.5 Hz, Ar-H), 7.97 (1H, s, Pyr-H), 8.08 (1H, d, J 8 Hz, Ar-H), 8.41 (1H, d, J 8.5 Hz, Ar-H); MS(+ve); 508.4; tR = 2.74 min (xbridge 4).
100		***		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 1.60 (8H, m, CH + CH₂), 1.75 (4H, m, CH), 1.87 – 1.99 (6H, m, CH + CH₂), 2.16 (3H, s, NCH₃), 2.29 (2H, m, CH), 2.79 (2H, d, J 11 Hz, CH), 3.18 (3H, s, NCH₃), 3.64 (2H, s, CH₂), 3.73 (1H, m, CH), 3.94 (3H, s, OCH₃), 4.82 (1H, m, CH), 7.48 (2H, m, Ar-H), 7.71 (1H, s, NH), 8.05 (1H, s, pyrH), 8.09 (1H, d, J 8 Hz, Ar-H), 8.36 (1H, d, J 8 Hz, Ar-H); MS(+ve); 548.4; tR = 3.20 min (xbridge 4).





101			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopentane]-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 1.30 (2H, m, CH₂), 1.51-1.63 (10H, m, 4 x CH₂), 1.88 (4H, m, 2 x CH₂), 1.91 - 2.17 (6H, m, 3 x CH₂), 2.17 (3H, s, NCH₃), 2.89 (2H, d, J 8.5Hz, CH), 3.19 (3H, s, NCH₃), 3.41 (2H, s, CH₂), 3.71 (1H, m, CH), 3.94 (3H, s, OCH₃), 5.02 (1H, m, CH), 7.48 (2H, m, Ar-H), 7.69 (1H, s, Ar-H), 8.01 (1H, s, Ar-H), 8.09 (1H, d, J 7.5Hz, Ar-H), 8.38 (1H, d, J 8.5Hz, Ar-H); MS(+ve); 562.5; tR = 3.45 min (xbridge 4).
102			**	4-(9-Allyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (CD₃OD): 1.72 (2H, m, CH₂), 1.96 (2H, m, CH₂), 2.23 (2H, dd, J 10.5Hz, 10.5Hz, CH), 2.33 (3H, s, NCH₃), 2.73 (2H, m, CH₂), 2.94 (2H, d, J 12Hz, CH₂), 3.76 (2H, m, CH₂), 3.91 (1H, m, CH), 4.01 (3H, s, OCH₃), 4.30 (2H, d, J 5.5Hz, CH₂), 5.30 (2H, m, =CH₂), 6.06 (1H, m, =CH), 7.46 (1H, dd, J 8.5Hz, 2Hz, ArH), 7.50 (1H, d, J 2 Hz, ArH), 8.06 (1H, s, pyrH), 8.49 (1H, d, J 8.5Hz, ArH); MS(+ve); 480.3; tR = 2.49 min (xbridge 4).
103		**	*	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzenesulfonamide	





104		***	*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-methoxy-ethyl)-benzenesulfonamide	¹ H NMR (CD ₃ OD): 1.67 – 1.99 (8H, m, alkyl-H), 2.80 (2H, dd, J 7.5 Hz, CH ₂), 3.04 – 3.06 (2H, m, CH ₂), 3.26 (3H, s, CH ₃), 3.38 – 3.41 (2H, m, CH ₂), 3.82 (2H, dd, J 4.5 Hz, CH ₂), 5.01 – 5.04 (1H, m, CH), 7.76 – 7.79 (2H, m, Ar-H), 7.85 – 7.87 (2H, m, Ar-H), 7.96 (1H, s, Ar-H); MS(+ve): 475.33; tR = 2.87 (Xbridge 4).
105		**		9-Cyclopentyl-5-methyl-2-[4-(4-methylpiperazine-1-sulfonyl)-phenylamino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	MS(+ve): 500.33; tR = 2.97 (Xbridge 4).
106			**	9-Cyclopentyl-5-methyl-2-[(thiophen-3-ylmethyl)-amino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
107			*	9-Cyclopentyl-2-[(furan-3-ylmethyl)-amino]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	





108		**	*	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-methoxy-ethyl)-benzenesulfonamide	
109		**	**	9-Cyclopentyl-5-methyl-2-[3-(pyrrolidine-1-sulfonyl)-phenylamino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
110		**	*	N-Benzyl-3-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzenesulfonamide	
111			**	4-(5,9-Dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.63 (2H, m, CH), 1.79 (2H, m, CH), 2.26 (2H, m, CH), 2.60 (2H, m, CH ₂), 2.89 (2H, m, CH), 3.06 (3H, s, NCH ₃), 3.17 (3H, s, NCH ₃), 3.29 (3H, s, NCH ₃), 3.68 (2H, m, CH ₂), 3.78 (1H, m, CH), 3.94 (3H, s, OCH ₃), 7.47 (1H, s, ArH), 7.53 (1H, d, J 9Hz, ArH), 8.10 (2H, m, ArH), 8.46 (1H, d, J 8.5Hz, ArH); MS(+ve); 454.4; tR = 2.33 min (xbridge 4).





112			*	4-(5,9-Dimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-methyl-benzamide	¹ H NMR (DMSO): 2.60 (2H, m, CH ₂), 2.78 (3H, m, NCH ₃), 3.06 (3H, s, NCH ₃), 3.17 (3H, s, NCH ₃), 3.68 (2H, m, CH ₂), 3.93 (3H, s, OCH ₃), 7.49 (2H, m, ArH), 7.73 (1H, m, ArH), 8.10 (1H, d, J 2.5 Hz, ArH), 8.29 (1H, bs, NH), 8.45 (1H, dd, J 9 Hz, 2.5 Hz, ArH); MS(+ve): 371.3; tR = 2.29 min (xbridge 4).
113			**	4-(9-Isopropyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-methyl-benzamide	¹ H NMR (DMSO): 0.67 (2H, m, cyclopropyl H), 0.92 (2H, m, cyclopropyl H), 1.17 (6H, d, J 6.5 Hz), 2.78 (3H, d, J 5 Hz, NCH ₃), 3.16 (3H, s, NCH ₃), 3.46 (2H, s, CH ₂), 3.93 (3H, s, OCH ₃), 4.83 (1H, m, CH), 7.47 (2H, m, ArH), 7.63 (1H, s, NH), 7.97 (1H, s, pyrH), 8.29 (1H, d, J 4.5 Hz, NH), 8.42 (1H, d, J 8.5 Hz, ArH); MS(+ve): 425.3; tR = 2.70 min (xbridge 4).
114		***		9-Cyclopropylmethyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	MS(+ve): 414.2; tR = 2.64 min (XBridge 4).
115		***	*	2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-9-isopropyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.21 (6H, d, J 7 Hz, 2CH ₃), 2.58 (2H, t, J 5 Hz, CH ₂), 3.16 (3H, s, CH ₃), 3.59 (2H, t, J 5 Hz, CH ₂), 4.39 (2H, s, CH ₂), 4.46 (2H, s, CH ₂), 4.8 (1H, m, CH(CH ₃) ₂), 7.24 (1H, d, J 8.5 Hz, CH), 7.59 (1H, dd, J 8.5, 2 Hz, CH), 7.87 (1H, s, CH), 8.03 (1H, s, CH), 9.39 (1H, s, NH); MS(+ve): 402.3; tR = 2.58 min (XBridge 4).





116		**		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-9-(3-methyl-butyl)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 0.90 (3H, s, CH ₃), 0.91 (3H, s, CH ₃), 1.51 (2H, m, CH ₂), 1.60 (1H, m, CH(CH ₃) ₂), 2.60 (2H, t, J 5.0Hz, C(O)CH ₂), 3.15 (3H, s, CH ₃ N), 3.59 (2H, t, J 7.5Hz, CH ₂ N), 3.67 (2H, t, J 5.0Hz, CH ₂ N), 4.39 (2H, s, CH ₂), 4.43 (2H, s, CH ₂), 7.22 (1H, d, J 8Hz, CH), 7.69 (1H, s, CH), 7.74 (1H, d, J 8Hz, CH), 8.04 (1H, s, CH), 9.37 (1H, s, NH); MS(+ve): 430.3; tR = 3.01 min (XBridge 4).
117		**	**	(9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]	¹ H NMR (DMSO): 1.5-2.0 (12H, m, 6CH ₂), 2.27 (2H, m, CH ₂), 3.18 (3H, s, CH ₃), 3.62 (2H, s, CH ₂ N), 4.40 (2H, s, CH ₂), 4.43 (2H, s, CH ₂), 4.82 (1H, m, CHN), 7.24 (1H, d, J 8Hz, CH), 7.62 (1H, d, J 8Hz, CH), 7.83 (1H, s, CH), 8.04 (1H, s, CH), 9.35 (1H, s, NH); MS(+ve): 468.3; tR = 3.32 min (XBridge 4).
118		**		(9-Cyclopentyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopentane]	¹ H NMR (DMSO): 1.33 (2H, m, CH ₂), 1.45-1.65 (8H, m, 4CH ₂), 1.71 (2H, m, CH ₂), 1.89 (2H, m, CH ₂), 2.01 (2H, m, CH ₂), 3.18 (3H, s, CH ₃), 3.40 (2H, m, CH ₂), 4.39 (2H, s, CH ₂), 4.43 (2H, s, CH ₂), 5.02 (1H, m, CHN), 7.24 (1H, d, J 8.5Hz, CH), 7.61 (1H, d, J 8.5Hz, CH), 7.82 (1H, s, CH), 7.98 (1H, s, CH), 9.37 (1H, s, NH); MS(+ve): 482.3; tR = 3.57 min (XBridge 2).
119		***	*	9-Allyl-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.64 (2H, m, CH ₂), 3.18 (3H, s, CH ₃), 3.63 (2H, m, CH ₂), 4.17 (2H, m, CH ₂), 4.39 (2H, s, CH ₂), 4.42 (2H, s, CH ₂), 5.22 (2H, m, CH=CH ₂), 5.97 (1H, m, CH=CH ₂), 7.22 (1H, d, J 8.5Hz, CH), 7.59 (1H, dd, J 8.5, 2Hz, CH), 7.85 (1H, s, CH), 8.08 (1H, s, CH).




120		***			CH), 9.43 (1H, s, NH); MS(+ve): 400.3; tR = 2.60 min (XBridge 4). 1H NMR (DMSO): 0.83 (6H, t, J 7.5Hz, 2CH3), 0.93 (3H, t, J 7Hz, CH2CH3), 1.59 (4H, m, 2CH2), 2.57 (2H, m, CH2), 3.45 (2H, m, CH2), 3.74 (2H, q, J 7.0Hz, CH2CH3), 4.4 (2H, m, CH2), 4.45 (2H, s, CH2), 4.58 (1H, m, CHN), 7.25 (1H, d, J 8Hz, CH), 7.69 (1H, d, J 8Hz, CH), 7.73 (1H, s, CH), 8.08 (1H, s, CH), 9.38 (1H, s, NH); MS(+ve): 442.3; tR = 3.08 min (XBridge 4).
121		**			1H NMR (DMSO): 2.60 (2H, m, CH2), 3.19 (3H, s, CH3), 3.54 (2H, m, CH2), 4.38 (2H, s, CH2), 4.43 (2H, s, CH2), 7.21 (1H, d, J 8.5Hz, CH), 7.45 (1H, s, NH), 7.68 (1H, dd, J 8.5, 2Hz, CH), 7.89 (1H, s, CH), 8.06 (1H, s, CH), 9.3 (1H, s, NH); MS(+ve): 360.2; tR = 2.18 min (XBridge 4).
122		**			1H NMR (DMSO): 0.91 (3H, t, J 7.0Hz, CH2CH3), 1.41 (9H, s, C(CH3)3), 2.39 (2H, t, J 6Hz, CH2), 3.60 (2H, t, J 6Hz, CH2), 3.67 (2H, q, J 7.0Hz, CH2), 4.4 (2H, s, CH2), 4.45 (2H, s, CH2), 7.26 (1H, d, J 8Hz, CH), 7.62 (1H, d, J 8Hz, CH), 7.83 (1H, s, CH), 8.22 (1H, s, CH), 9.41 (1H, s, NH); MS(+ve): 430.3; tR = 2.94 min (XBridge 4).
123		**	**		1H NMR (DMSO): 1.61 (4H, m, 2CH2), 1.68 (2H, m, CH2), 1.92 (2H, m, CH2), 2.57 (2H, t, J 5Hz, CH2), 3.16 (3H, s, CH3), 3.60 (2H, t, J 5Hz, CH2), 4.78 (1H, m, CHN), 7.28 (2H, d, J 9Hz, 2CH), 7.74 (2H, d, J 9Hz, 2CH), 8.04 (1H, s, CH), 9.35 (1H, s, NH); MS(+ve): 372.3; tR =




						3.93 min (XBridge 4).
124		*	*	*	9-Cyclopentyl-2-[(furan-2-ylmethyl)-amino]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (CD ₃ OD): 2.38 – 2.80 (8H, m, alkyl-H), 3.15 (3H, s, CH ₃), 3.36 (2H, dd, J 5 Hz, CH ₂), 3.96 (3H, s, CH ₃), 4.39 (2H, dd, J 5 Hz, CH ₂), 5.60 – 5.64 (1H, m, CH), 6.78 (1H, s, Ar-H), 7.91 – 7.98 (2H, m, Ar-H), 8.68 (1H, s, Ar-H), 8.80 (1H, s, Ar-H), 9.65 (1H, s, Ar-H), 11.47 (1H, bs, NH); 391.35; tR = 3.21 (Xbridge 4).
125		***		**	9-Cyclopentyl-5-methyl-2-(2-methyl-1H-indol-5-ylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.02 (3H, t, J 7 Hz, CH ₂ CH ₃), 2.55 (2H, m, CH ₂), 3.53 (2H, m, CH ₂), 3.74 (2H, q, J 7 Hz, CH ₂ CH ₃), 4.4 (2H, s, CH ₂), 4.43 (2H, s, CH ₂), 7.22 (1H, d, J 8.5 Hz, CH), 7.42 (1H, s, NH), 7.68 (1H, dd, J 8.5, 1.5 Hz, CH), 7.89 (1H, s, CH), 8.10 (1H, s, CH), 9.35 (1H, s, NH); MS(+ve): 374.2; tR = 2.2 min (XBridge 4).
126		*			2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-ethyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.50 – 1.78 (8H, m, alkyl-H), 2.48 (2H, dd, J 5.5 Hz, CH ₂), 3.09 (3H, s, CH ₃), 3.50 (2H, dd, J 5.5 Hz, CH ₂), 4.17 – 4.19 (4H, m, 2 x CH ₂), 4.27 (2H, d, J 6 Hz, CH ₂ NH), 4.57 (1H, bs, CH), 6.75 – 6.78 (3H, m, Ar-H), 7.22 (1H, bs, NH), 7.89 (1H, s, Ar-H); MS(+ve): 410.35; tR = 3.22 (Xbridge 4).
127				*	9-Cyclopentyl-2-[(2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-amino]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.50 – 1.78 (8H, m, alkyl-H), 2.48 (2H, dd, J 5.5 Hz, CH ₂), 3.09 (3H, s, CH ₃), 3.50 (2H, dd, J 5.5 Hz, CH ₂), 4.17 – 4.19 (4H, m, 2 x CH ₂), 4.27 (2H, d, J 6 Hz, CH ₂ NH), 4.57 (1H, bs, CH), 6.75 – 6.78 (3H, m, Ar-H), 7.22 (1H, bs, NH), 7.89 (1H, s, Ar-H); MS(+ve): 410.35; tR = 3.22 (Xbridge 4).





128		*	**	9-Cyclopentyl-2-[(2,3-dihydrobenzo[1,4]dioxin-5-ylmethyl)-amino]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.46 – 1.99 (8H, m, alkyl-H), 2.61 (2H, dd, J 5.5 Hz, CH ₂), 3.23 (3H, s, CH ₃), 3.63 (2H, dd, J 5.5 Hz, CH ₂), 4.24 – 4.27 (2H, m, CH ₂), 4.29 – 4.32 (2H, m, CH ₂), 4.50 (2H, s, CH ₂), 4.66 – 4.69 (1H, m, CH), 6.71 – 6.72 (m, 2H, Ar-H), 6.77 – 6.79 (1H, m, Ar-H), 7.83 (1H, s, Ar-H); MS(+ve): 410.35; tR = 3.34 (Xbridge 4).
129			*	2-[(Benzo[b]thiophen-2-ylmethyl)-amino]-9-cyclopentyl-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.17 – 1.60 (8H, m, alkyl-H), 2.48 (2H, dd, J 4.5 Hz, 2H, CH ₂), 3.10 (3H, s, CH ₃), 3.49 (2H, dd, J 4.5 Hz, CH ₂), 4.65 (2H, d, J 6 Hz, CH ₂ NH), 7.35 – 7.42 (4H, m, Ar-H), 7.89 (1H, s, Ar-H), 7.95 – 7.97 (1H, m, Ar-H); MS(+ve): 408.30; tR = 3.78 (Xbridge 4).
130		**	*	9-Cyclopentyl-2-(1H-indol-6-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (CD ₃ OD): 1.60 – 1.80 (8H, m, alkyl-H), 2.02 (3H, s, CH ₃), 2.65 (2H, dd, J 5.5 Hz, CH ₂), 3.68 (2H, dd, J 5.5 Hz, CH ₂), 6.37 (1H, d, J 2.5 Hz, Ar-H), 6.88 (1H, s, Ar-H), 7.12 – 7.18 (1H, m, Ar-H), 7.21 (1H, d, J 3 Hz, Ar-H), 7.84 (1H, d, J 1.5 Hz, Ar-H), 7.90 (1H, s, Ar-H); MS(+ve): 377.32; tR = 3.02 (Xbridge 4).
131			*	9-Cyclopentyl-5-methyl-2-(3-methylbenzylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.48 – 1.77 (8H, m, alkyl-H), 2.27 (3H, s, CH ₃), 2.47 (2H, dd, J 4.5 Hz, CH ₂), 3.09 (3H, s, CH ₃), 3.50 (2H, dd, J 4.5 Hz, CH ₂), 4.36 (2H, d, J 6 Hz, CH ₂), 4.56 (1H, bs, CH), 7.00 (1H, d, J 7.5 Hz, Ar-H), 7.08 – 7.11 (2H, m, Ar-H), 7.16 (1H, t, J 7.5 Hz, Ar-H), 7.27 (1H, bs, NH), 7.86 (1H, s, Ar-H); MS(+ve): 366.29; tR = 3.65 (Xbridge 4).






132		*	*	9-Cyclopentyl-5-methyl-2-(4-[1,2,3]thiadiazol-4-yl-benzylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (CD ₃ OD): 1.59 – 1.79 (8H, m, alkyl-H), 2.75 (2H, dd, J 5 Hz, CH ₂), 3.24 (3H, s, CH ₃), 3.73 (2H, dd, J 5 Hz, CH ₂), 4.71 (2H, s, CH ₂), 7.53 (2H, d, J 8 Hz, Ar-H), 7.83 (1H, s, Ar-H), 8.12 (2H, d, J 8 Hz, Ar-H), 9.26 (1H, s, Ar-H); MS(+ve): 436.29; tR = 3.28 (Xbridge 4).
133		**		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-9-(2-methoxyphenyl)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.80 (2H, m, CH ₂), 3.26 (3H, s, CH ₃), 3.69 (3H, s, CH ₃), 3.87 (2H, m, CH ₂), 4.13 (2H, s, CH ₂), 4.29 (2H, s, CH ₂), 6.88 (1H, d, J 8 Hz, CH), 7.08 (1H, m, CH), 7.12 (2H, m, 2CH), 7.22 (1H, d, J 8 Hz, CH), 7.31 (1H, m, CH), 7.41 (1H, m, CH), 8.17 (1H, s, CH), 9.36 (1H, s, NH); MS(+ve): 466.3; tR = 2.68 min (XBridge 4).
134		**		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-9-o-tolyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.06 (3H, s, CH ₃), 2.8 (1H, s, CH), 2.93 (1H, s, CH), 3.28 (3H, s, CH ₃), 3.82 (1H, s, CH), 4.03 (1H, s, CH), 4.09 (2H, d, J 8.5 Hz, CH ₂), 4.28 (2H, s, CH ₂), 6.85 (1H, d, J 9 Hz, CH), 7.09 (2H, m, 2CH), 7.35 (4H, m, 4CH), 8.19 (1H, s, CH), 9.38 (1H, s, NH); MS(+ve): 450.25; tR = 2.76 min (XBridge 4).
135		**		9-(2-Chloro-phenyl)-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	MS(+ve): 470.21; tR = 2.76 min (XBridge 4).





136		**		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-9-(3-methoxyphenyl)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 2.77 (2H, m, CH ₂), 3.27 (3H, s, CH ₃), 3.74 (3H, s, CH ₃), 4.02 (2H, m, CH ₂), 4.10 (2H, s, CH ₂), 4.32 (2H, s, CH ₂), 6.89 (3H, m, 3CH), 6.95 (1H, d, J 8.5Hz, CH), 7.25 (1H, d, J 9Hz, CH), 7.36 (2H, m, 2CH), 8.3 (1H, s, CH), 9.51 (1H, s, NH); MS(+ve): 466.3; tR = 2.68 min (XBridge 4).
137		***	*	9-Cyclopentyl-5-methyl-2-m-tolylamino-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.58 (4H, m, 2CH ₂), 1.7 (2H, m, CH ₂), 1.95 (2H, m, CH ₂), 2.26 (3H, s, CH ₃), 2.56 (2H, m, CH ₂), 3.16 (3H, s, CH ₃), 3.60 (2H, m, CH ₂), 4.84 (1H, m, CHN), 6.71 (1H, d, J 7.5Hz, CH), 7.10 (1H, m, CH), 7.40 (1H, d, J 8Hz, CH), 7.67 (1H, s, CH), 8.03 (1H, s, CH), 9.11 (1H, s, NH); MS(+ve): 352.3; tR = 3.72 min (XBridge 4).
138		**	*	9-Cyclopentyl-2-(3-methoxy-phenylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.57 (4H, m, 2CH ₂), 1.69 (2H, m, CH ₂), 1.95 (2H, m, CH ₂), 2.56 (2H, m, CH ₂), 3.16 (3H, s, CH ₃), 3.60 (2H, m, CH ₂), 3.72 (3H, s, CH ₃), 4.82 (1H, m, CHN), 6.48 (1H, dd, J 8, 2Hz, CH), 7.12 (1H, t, J 8Hz, CH), 7.25 (1H, dd, J 8, 1Hz, CH), 7.45 (1H, s, CH), 8.04 (1H, s, CH), 9.15 (1H, s, NH); MS(+ve): 368.3; tR = 3.47 min (XBridge 4).
139		***	**	9-Cyclopentyl-2-(1H-indazol-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.58 – 1.95 (8H, m, alkyl-H), 2.54 (2H, dd, J 5 Hz, CH ₂), 3.16 (3H, s, CH ₃), 3.60 (2H, dd, J 5 Hz, CH ₂), 4.80 – 4.84 (1H, m, CH), 7.41 (1H, d, J 9 Hz, Ar-H), 7.52 (1H, d, J 9 Hz, Ar-H), 7.89 (1H, s, Ar-H), 8.04 (1H, s, Ar-H), 8.21 (1H, s, Ar-H), 9.14 (1H, bs, NH), 12.86 (1H, bs, NH); MS(+ve): 376.30; tR = 2.59 (Xbridge 4).





140		*	*	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-methyl-N-(1-methyl-piperidin-4-yl)-benzamide	¹ H NMR (DMSO): 1.47 (6H, m, CH), 1.62 (2H, m, CH), 1.81 (4H, m, CH), 1.93 (2H, dd, J 10Hz, 10Hz, CH), 2.14 (3H, s, CH ₃), 2.19 (3H, s, CH ₃), 2.54 (2H, m, CH ₂), 2.74 (2H, m, CH ₂), 3.14 (3H, s, CH ₃), 3.54 (2H, m, CH ₂), 3.69 (1H, m, CH), 4.65 (1H, m, CH), 6.96 (1H, d, J 7.5Hz, Ar-H), 7.15 (1H, dd, J 7.5Hz, 7.5Hz, Ar-H), 7.63 (1H, d, J 7Hz, Ar-H), 7.96 (1H, s, NH), 8.15 (1H, d, J 8Hz, Ar-H), 8.35 (1H, s, Pyr-H); MS(+ve); 492.4; tR = 2.57 min (xbridge 4).
141		*		3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2,N-dimethyl-benzamide	¹ H NMR (DMSO): 1.45 (4H, m, CH), 1.62 (2H, m, CH), 1.80 (2H, m, CH), 2.19 (3H, s, N-CH ₃), 2.55 (2H, m, CH ₂), 2.74 (3H, d, J 5Hz, N-CH ₃), 3.29 (3H, s, CH ₃), 3.56 (2H, m, CH ₂), 4.61 (1H, s, CH), 6.99 (1H, d, J 8Hz, Ar-H), 7.16 (1H, dd, J 7.5Hz, 8Hz, Ar-H), 7.61 (1H, d, J 8Hz, Ar-H), 7.96 (1H, s, NH), 8.13 (1H, m, Ar-H), 8.35 (1H, s, Pyr-H); MS(+ve); 409.3; tR = 2.58 min (xbridge 4).
142		*		3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-hydroxy-ethyl)-2-methyl-benzamide	¹ H NMR (DMSO): 1.47 (4H, m, CH), 1.62 (2H, m, CH), 1.79 (2H, m, CH), 2.14 (3H, s, CH ₃), 2.20 (3H, s, CH ₃), 2.55 (2H, m, CH ₂), 3.14 (3H, s, CH ₃), 3.27 (2H, m, CH ₂), 3.49 (2H, m, CH ₂), 3.56 (2H, m, CH ₂), 4.63 (1H, m, CH), 4.67 (1H, bs, OH), 7.01 (1H, d, J 7Hz, Ar-H), 7.16 (1H, dd, J 7.5Hz, 7.5Hz, Ar-H), 7.63 (1H, d, J 7Hz, Ar-H), 7.96 (1H, s, NH), 8.13 (1H, m, Ar-H), 8.35 (1H, s, Pyr-H); MS(+ve); 439.3; tR = 2.39 min (xbridge 4).





143		**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	¹H NMR (DMSO): 1.49 – 1.57 (8H, m, CH), 1.74 (2H, m, CH), 1.84 (2H, m, CH), 1.94 (2H, dd, J 10.5Hz, 10.5Hz, CH), 2.15 (3H, s, CH₃), 2.53 (2H, m, CH₂), 2.76 (2H, d, J 11.5Hz, CH), 3.16 (3H, s, NCH₃), 3.58 (2H, m, CH₂), 3.68 (1H, m, CH), 3.89 (3H, s, OCH₃), 4.81 (1H, m, CH), 7.04 (1H, d, J 8.5Hz, Ar-H), 7.50 (1H, d, J 8.5Hz, Ar-H), 7.70 (1H, s, NH), 8.00 (1H, d, J 7Hz, Ar-H), 8.03 (1H, s, NH), 8.56 (1H, s, pyr-H): MS(+ve); 508.4; tR = 2.64 min (xbridge 4).
144		**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-N-methyl-benzamide	¹H NMR (DMSO): 1.60 (6H, m, CH), 1.86 (2H, m, CH), 2.56 (2H, m, CH₂), 2.74 (3H, d, J 4.5Hz, NCH₃), 3.16 (3H, s, NCH₃), 3.58 (2H, m, CH₂), 3.90 (3H, s, CH₃), 4.42 (1H, m, CH), 7.04 (1H, d, J 8.5Hz, Ar-H), 7.43 (1H, d, J 8.5Hz, Ar-H), 7.68 (1H, s, NH), 8.04 (1H, s, N-H), 8.20 (1H, m, Ar-H), 8.60 (1H, s, pyr-H): MS(+ve); 425.3; tR = 2.64 min (xbridge 4).
145		**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-hydroxy-ethyl)-4-methoxy-benzamide	¹H NMR (DMSO): 1.55 (6H, m, CH), 1.86 (2H, m, CH), 2.56 (2H, m, CH₂), 3.16 (3H, s, CH₃), 3.30 (2H, m, CH₂), 3.49 (2H, m, CH₂), 3.59 (2H, m, CH₂), 3.90 (3H, s, OCH₃), 4.69 (1H, m, CH), 4.84 (1H, bs, OH), 7.04 (1H, d, J 8.5Hz, Ar-H), 7.48 (1H, d, J 8Hz, Ar-H), 7.61 (1H, s, NH), 8.03 (1H, s, NH), 8.21 (1H, d, J 5.5Hz, Ar-H), 8.61 (1H, s, pyr-H): MS(+ve); 455.3; tR = 2.46 min (xbridge 4).





146		*	*	9-Cyclopentyl-5-methyl-2-[3-(piperazine-1-sulfonyl)-phenylamino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
147		**		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-propyl)-3-methoxy-N-methyl-benzamide	
148		***		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-diethylamino-ethyl)-3-methoxy-benzamide	
149		**		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-propyl)-3-methoxy-benzamide	





150			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((1S,2R)-2-hydroxy-indan-1-yl)-3-methoxy-benzamide	
151		*	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(4-hydroxy-butyl)-3-methoxy-benzamide	
152			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((R)-1-hydroxymethyl-2-methyl-propyl)-3-methoxy-benzamide	
153			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-hydroxymethyl-propyl)-3-methoxy-benzamide	
154			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-hydroxy-propyl)-3-methoxy-benzamide	





159			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-hydroxy-propyl)-3-methoxybenzamide	
160			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(4-methyl-piperazin-1-yl)-3-methoxybenzamide	
161			***	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzamide	
162			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1,2,2,6,6-pentamethyl-piperidin-4-yl)-3-methoxybenzamide	





163			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-3-yl)-benzamide	
164			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-hydroxy-butyl)-3-methoxy-benzamide	
165			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-2,2-dimethyl-propyl)-3-methoxy-benzamide	
166			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-hydroxy-ethyl)-3-methoxy-benzamide	





167			*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N,N-bis-(2-hydroxy-ethyl)-3-methoxy-benzamide	
168			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-[2-(2-hydroxy-ethoxy)-ethyl]-3-methoxy-benzamide	
169			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3,N-dimethoxy-benzamide	
170		***	*	3-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-propionic acid	





171		***	*	N-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-acetamide	
172		***	*	[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-acetic acid	
173		***	*	9-Cyclopentyl-5-methyl-2-(2-oxo-2,3-dihydro-1H-indol-5-ylamino)-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.51 – 1.69 (8H, m, alkyl-H), 1.85 (2H, bs, CH2), 2.66 (2H, dd, J 4.5 Hz, CH2), 3.15 (3H, s, CH3), 3.66 (2H, dd, J 4.5 Hz, CH2), 4.78 – 4.83 (1H, m, CH), 6.79 (1H, d, J 8 Hz, Ar-H), 7.30 (1H, d, J 8 Hz, Ar-H), 7.47 (1H, s, Ar-H), 7.96 (1H, s, Ar-H), 9.86 (1H, bs, NH), 10.35 (1H, bs, NH); MS(+ve): 393.32; tR = 2.46 (Xbridge 4).
174		*	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methyl-benzoic acid	1H NMR (DMSO): 1.49 – 1.83 (8H, m, alkyl-H), 2.31 (3H, s, CH3), 2.57 (2H, dd, J 4.5 Hz, CH2), 3.14 (3H, s, CH3), 3.58 (2H, dd, J 4.5 Hz, CH2), 4.68 – 4.71 (1H, m, CH), 7.70 (1H, d, J 8.5 Hz, Ar-H), 7.75 (1H, s, Ar-H), 7.98 (1H, d, J 8.5 Hz, Ar-H), 9.04 (1H, s, Ar-H), 8.35 (s, 1H, Ar-H), 12.55 (1H, bs, NH); MS(+ve): 396.32; tR = 1.78 (Xbridge 4).





175		*		9-Cyclopentyl-2-[2-(1H-indol-3-yl)-1-methylethylamino]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (CD ₃ OD): 1.22 (3H, d, J 5.5 Hz, CH ₃), 1.52 – 1.97 (8H, m, alkyl-H), 2.57 (2H, dd, J 6 Hz, CH ₂), 2.90 – 3.04 (2H, m, CH ₂), 3.19 (3H, s, CH ₃), 3.58 (2H, dd, J 6 Hz, CH ₂), 4.35 – 4.39 (1H, m, CH), 4.68 – 4.74 (1H, m, CH), 6.97 (1H, t, J 8 Hz, Ar-H), 7.04 – 7.08 (2H, m, Ar-H), 7.31 (1H, d, J 8 Hz, Ar-H), 7.54 (1H, d, J 8 Hz, Ar-H), 7.75 (1H, s, Ar-H); MS(+ve): 419.36; tR = 3.60 (XBridge 4).
176		***	**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-cyclopropyl-benzamide	1H NMR (DMSO): 0.55 (2H, m, CH ₂), 0.66 (2H, m, CH ₂), 1.61 (4H, m, 2CH ₂), 1.72 (2H, m, CH ₂), 1.95 (2H, m, CH ₂), 2.58 (2H, m, CH ₂), 2.81 (1H, m, CH), 3.17 (3H, s, CH ₃), 3.62 (2H, m, CH ₂), 4.82 (1H, m, CHN), 7.72 (2H, d, J 9 Hz, 2CH), 7.78 (2H, d, J 9 Hz, 2CH), 8.08 (1H, s, CH), 8.21 (1H, d, J 4 Hz, NH) 9.47 (1H, s, NH); MS(+ve): 421.4; tR = 2.75 min (XBridge 4).
177		***	**	9-Cyclopentyl-2-[3-(1,1-dioxo-thiomorpholin-4-ylmethyl)-phenylamino]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.6 (4H, m, 2CH ₂), 1.71 (2H, m, CH ₂), 1.97 (2H, m, CH ₂), 2.56 (2H, m, CH ₂), 2.86 (4H, m, 2CH ₂), 3.11 (4H, m, 2CH ₂), 3.16 (3H, s, CH ₃), 3.61 (4H, m, 2CH ₂), 4.84 (1H, m, CHN), 6.88 (1H, d, J 7.5 Hz, CH), 7.2 (1H, t, J 8 Hz, CH), 7.61 (1H, d, J 8.5 Hz, CH), 7.71 (1H, s, CH), 8.04 (1H, s, CH), 9.17 (1H, s, NH); MS(+ve): 485.4; tR = 3.0 min (XBridge 4).
178		***	**	Cyclopropanecarboxylic acid [4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-amide	1H NMR (DMSO): 0.76 (4H, m, 2CH ₂), 1.57 (5H, m, 2CH ₂ + CH), 1.72 (2H, m, CH ₂), 1.91 (2H, m, CH ₂), 2.56 (2H, m, CH ₂), 3.15 (3H, s, CH ₃), 3.59 (2H, m, CH ₂), 4.77 (1H, m, CHN), 7.44 (2H, d, J 9 Hz, 2CH), 7.61 (2H, d, J 9 Hz, 2CH),





					8.01 (1H, s, CH), 9.09 (1H, s, NH) 10.0 (1H, s, NH); MS(+ve): 421.4; tR = 2.9 min (XBridge 4).
179		***	*	Cyclopropanecarboxylic acid [4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-methyl-phenyl]-amide	¹ H NMR (DMSO): 0.75 (4H, m, 2CH ₂), 1.57 (4H, m, 2CH ₂), 1.7 (2H, m, CH ₂), 1.8 (1H, m, CH), 1.96 (2H, m, CH ₂), 2.51 (3H, s, CH ₃), 2.56 (2H, m, CH ₂), 3.15 (3H, s, CH ₃ N), 3.60 (2H, m, CH ₂), 4.83 (1H, m, CHN), 7.17 (1H, d, J 8.5Hz, CH), 7.38 (1H, d, J 8.5Hz, CH), 7.64 (1H, s, CH), 8.03 (1H, s, CH), 9.01 (1H, s, NH) 9.38 (1H, s, NH); MS(+ve): 435.4; tR = 2.89 min (XBridge 4).
180		***	*	3-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-N-(1-methyl-piperidin-4-yl)-propionamide	
181		***	*	2-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-N-(1-methyl-piperidin-4-yl)-acetamide	
182			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(2-diethylamino-ethyl)-3-methoxybenzamide	MS(+ve); 536.5; tR = 3.42 min (xbridge 4).





183		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(3-methylamino-propyl)-3-methoxybenzamide	MS(+ve); 522.5; tR = 3.12 min (xbridge 4).
184		**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-((1S,2R)-2-hydroxy-indan-1-yl)-3-methoxybenzamide	MS(+ve); 569.5; tR = 3.50 min (xbridge 4).
185		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(4-hydroxy-butyl)-3-methoxybenzamide	MS(+ve); 509.3; tR = 2.77 min (xbridge 4).
186		***	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxybenzamide	MS(+ve); 546.4; tR = 3.02 min (xbridge 4).

187		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-3-yl)-benzamide	MS(+ve); 534.4; tR = 3.12 min (xbridge 4).
188		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(2-hydroxy-ethyl)-3-methoxy-benzamide	MS(+ve); 481.3; tR = 2.67 min (xbridge 4).
189		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-[2-(2-hydroxy-ethoxy)-ethyl]-3-methoxy-benzamide	MS(+ve); 525.4; tR = 2.68 min (xbridge 4).
190		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3,N-dimethoxy-benzamide	MS(+ve); 467.4; tR = 2.51 min (xbridge 4).





191					N-Acetyl-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzenesulfonamide	1H NMR (DMSO): 1.6 (4H, m, 2CH ₂), 1.72 (2H, m, CH ₂), 1.9 (3H, s, CH ₃), 1.95 (2H, m, CH ₂), 2.59 (2H, m, CH ₂), 3.17 (3H, s, CH ₃), 3.62 (2H, m, CH ₂), 4.83 (1H, m, CHN), 7.45 (2H, d, J 8.5 Hz, 2CH), 7.92 (2H, d, J 8.5 Hz, 2CH), 8.10 (1H, s, CH), 9.76 (1H, s, NH), 11.86 (1H, s, NH); MS(+ve): 459.3; tR = 1.63 min (XBridge 4).
192		*	*		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-ethyl benzoic acid	1H NMR (CD ₃ OD): 1.28 (3H, t, J 7.5 Hz, CH ₂ CH ₃), 1.51 – 1.54 (2H, m, alkyl-H), 1.63 – 1.65 (m, 2H, alkyl-H), 1.71 – 1.75 (2H, m, alkyl-H), 1.83 (2H, bs, alkyl-H), 2.76 (2H, q, J 7.5 Hz, CH ₂ CH ₃), 2.79 (2H, dd, J 5 Hz, CH ₂), 3.27 (3H, s, CH ₃), 3.78 (2H, dd, J 5 Hz, CH ₂), 7.63 (1H, d, J 8 Hz, Ar-H), 7.84 (1H, s, Ar-H), 7.95 (1H, dd, J 1.5 and 8 Hz, Ar-H), 8.05 (1H, d, J 1.5 Hz, Ar-H); MS(+ve): 410.36; tR = 1.89 (Xbridge 4).
193		*			N-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-yl)-benzamide	
194		***			N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide	





195			***	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methyl-piperazin-1-yl)-benzamide	
196			***	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-ylmethyl)-benzamide	
197			***	N-(1-Benzyl-piperidin-4-yl)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide	
198			**	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(2-methoxy-ethyl)-benzamide	





199			***	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-pyridin-4-ylmethylbenzamide	
200			***	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-[2-(1H-imidazol-4-yl)-ethyl]-3-methoxybenzamide	
201			***	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-furan-2-ylmethyl-3-methoxybenzamide	
202			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-2-ylmethyl)-benzamide	





203		**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(2-morpholin-4-ylethyl)-benzamide	
204		***	N-Cyclopentyl-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzamide	
205		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-1H-pyrrol-2-ylmethyl)-benzamide	
206		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-piperidin-4-ylbenzamide	





207			*	4-(9-Cyclopropyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
208			**	4-[9-(2-Hydroxy-ethyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
209			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methanesulfonyl-piperidin-4-yl)-3-methoxy-benzamide	
210			**	N-(1-Acetyl-piperidin-4-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide	
211			**	{4-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoylamino]-piperidin-1-yl}-acetic acid ethyl ester	





212			**		N-(1-Benzoyl-piperidin-4-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide				
213		*			9-Cyclopentyl-2-[4-(4-methoxy-phenyl)-piperazin-1-yl]-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one				
214			***		3-Chloro-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide				
215		*	***		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-fluoro-N-(1-methyl-piperidin-4-yl)-benzamide				

216			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-yl)-3-trifluoromethoxy-benzamide	
217			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-3-methoxy-N-(1-benzylpiperidin-4-yl)-benzamide	
218			***	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-3-methoxy-benzamide	
219			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-N-(2-diethylamino-ethyl)-3-methoxy-benzamide	

220		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-N-(2-dimethylamino-propyl)-3-methoxybenzamide	
221		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-N-(4-methyl-piperazin-1-yl)-3-methoxybenzamide	
222		**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclobutane]-2-ylamino)-N-furan-2-ylmethyl-3-methoxybenzamide	
223		*	4-{9-[2-(2-Hydroxy-ethyl-carbamoyl)-ethyl]-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino}-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	





224			**	3-Methoxy-4-[5-methyl-9-(2-methylcarbamoyl-ethyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide	
225			**	4-[9-(2-Dimethylcarbamoyl-ethyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
226			**	4-[9-[2-(3-Dimethylamino-propylcarbamoyl)-ethyl]-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
227		**	***	9-Cyclopentyl-2-(1H-indazol-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-6-one	






228		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-hydroxy-N-(1-methyl-piperidin-4-yl)-benzamide	
229		**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-ethyl-benzoic acid	
230		**	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-benzoic acid	
231		**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-diethylamino-ethyl)-4-methoxybenzamide	





232			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-propyl)-4-methoxy-benzamide	
233			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-hydroxy-propyl)-4-methoxy-benzamide	
234			*	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1,1-dimethyl-2-piperidin-1-yl-ethyl)-4-methoxy-benzamide	
235			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-4-ylmethyl)-benzamide	





236					3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-2,2-dimethylpropyl)-4-methoxy-benzamide	
237		***			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-ethyl-piperidin-4-yl)-3-methoxy-benzamide	
238		***			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-cyclopropylmethyl-piperidin-4-yl)-3-methoxy-benzamide	
239		**			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-benzoic acid	





240			***	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(2-diethylaminoethyl)-3-methoxybenzamide	
241			**	N-(1-Benzylpiperidin-4-yl)-4-(9-cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzamide	
242			**	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methylpiperidin-4-ylmethyl)-benzamide	
243			**	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzamide	





244			**	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1,2,2,6,6-pentamethyl-piperidin-4-yl)-benzamide	
245			**	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-3-yl)-benzamide	
246			**	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-2,2-dimethyl-propyl)-3-methoxy-benzamide	
247			**	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(3-dimethylamino-propyl)-3-methoxy-benzamide	





248			**	N-Cyclopentyl-4-(9-cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide	
249			**	4-(9-Cyclopentylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-1H-pyrrol-2-ylmethyl)-benzamide	
250			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-N-(1,1-dimethyl-2-pyrrolidin-1-yl-ethyl)-3-methoxy-benzamide	
251			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-N-(1,1-dimethyl-2-piperidin-1-yl-ethyl)-3-methoxy-benzamide	
252			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-(1-benzylpiperidin-4-yl)-benzamide	





253		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-ylmethyl)-benzamide	
254		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-N-(4-methyl-piperazin-1-yl)-benzamide	
255		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-(1,2,2,6,6-pentamethyl-piperidin-4-yl)-benzamide	
256		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-N-(3-dimethylamino-2,2-dimethyl-propyl)-3-methoxy-benzamide	





257					4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-pyridin-4-yl-benzamide	
258		***			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-N-pyridin-4-ylmethyl-benzamide	
259		***			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-N-(1-methyl-1H-pyrrol-2-ylmethyl)-benzamide	
260		***			N-Cyclopentyl-4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-benzamide	





261			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-ethyl-N-(1-methyl-piperidin-4-yl)-benzamide	
262			***	4-(9-Cyclopropylmethyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
263			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-[2-(2-hydroxy-ethoxy)-ethyl]-4-methoxy-benzamide	
264			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4,N-dimethoxy-benzamide	





265			**	N-(1-Benzyl-pyrrolidin-3-yl)-3-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-benzamide	
266			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-N-(2-morpholin-4-ylethyl)-benzamide	
267			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-[2-(1H-imidazol-4-yl)-ethyl]-4-methoxy-benzamide	
268			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-N-(1-methyl-1H-pyrrol-2-ylmethyl)-benzamide	




269			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-N-(1-methyl-piperidin-2-ylmethyl)-benzamide	
270			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-N-(2-methoxy-ethyl)-benzamide	
271			**	N-(1-Benzyl-piperidin-4-yl)-3-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-4-methoxy-benzamide	
272			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(4-hydroxy-butyl)-4-methoxy-benzamide	





273			**	3-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-furan-2-ylmethyl-4-methoxybenzamide	
274			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-hydroxy-N-(1-methyl-piperidin-4-yl)-benzamide	
275			**	4-(9-Cyclopropylmethyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxybenzoic acid	
276			**	4-(9-Cyclopropylmethyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	





277			*	3-Methoxy-4-[9-(2-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide	
278			*	3-Methoxy-4-[9-(2-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1,2,2,6,6-pentamethyl-piperidin-4-yl)-benzamide	
279			*	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-3-methoxy-4-[9-(2-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzamide	
280			*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-methylbenzoic acid	






281					4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane-2-ylamino)-3-trifluoromethoxy-benzoic acid	
282					4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-fluorobenzoic acid	
283		***			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-fluorobenzoic acid	
284		***			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-trifluoromethoxy-N-(1-methyl-piperidin-4-yl)-benzamide	





285		***	*	N-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-methyl-phenyl]-acetamide	¹ H NMR (DMSO): 1.57 (4H, m, 2CH ₂), 1.7 (2H, m, CH ₂), 1.93 (2H, m, CH ₂), 2.01 (3H, s, CH ₃), 2.14 (3H, s, CH ₃), 2.56 (2H, m, CH ₂), 3.16 (3H, s, CH ₃), 3.6 (2H, m, CH ₂), 4.83 (1H, m, CHN), 7.15 (1H, d, J 8.5Hz, CH), 7.39 (1H, d, J 8.5Hz, CH), 7.64 (1H, s, CH), 8.02 (1H, s, CH), 9.10 (1H, s, NH), 9.16 (1H, s, NH); MS(+ve): 409.4; tR = 2.58 min (XBridge 4).
286		**		N-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-trifluoromethyl-phenyl]-acetamide	¹ H NMR (DMSO): 1.56 (4H, m, 2CH ₂), 1.69 (2H, m, CH ₂), 1.92 (2H, m, CH ₂), 2.01 (3H, s, CH ₃), 2.58 (2H, m, CH ₂), 3.16 (3H, s, CH ₃), 3.61 (2H, m, CH ₂), 4.84 (1H, m, CHN), 7.27 (1H, d, J 8.5Hz, CH), 7.8 (1H, d, J 8.5Hz, CH), 8.08 (1H, s, CH), 8.24 (1H, s, CH), 9.39 (1H, s, NH), 9.54 (1H, s, NH); MS(+ve): 463.34; tR = 2.85 min (XBridge 4).
287		***	*	N-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-methyl-phenyl]-2,2-dimethylpropionamide	¹ H NMR (DMSO): 1.22 (9H, s, C(CH ₃) ₃), 1.58 (4H, m, 2CH ₂), 1.70 (2H, m, CH ₂), 1.94 (2H, m, CH ₂), 2.10 (3H, s, CH ₃), 2.57 (2H, m, CH ₂), 3.16 (3H, s, CH ₃), 3.60 (2H, m, CH ₂), 4.84 (1H, m, CHN), 6.98 (1H, d, J 8.5Hz, CH), 7.41 (1H, d, J 8.5Hz, CH), 7.67 (1H, s, CH), 8.03 (1H, s, CH), 8.76 (1H, s, NH), 9.14 (1H, s, NH); MS(+ve): 451.42; tR = 3.29 min (XBridge 4).
288		***		N-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-2-methyl-phenyl]-methanesulfonimide	¹ H NMR (DMSO): 1.6 (4H, m, 2CH ₂), 1.71 (2H, m, CH ₂), 1.95 (2H, m, CH ₂), 2.3 (3H, s, CH ₃), 2.58 (2H, m, CH ₂), 3.17 (3H, s, CH ₃), 3.5 (6H, s, 2CH ₃), 3.62 (2H, m, CH ₂), 4.86 (1H, m, CHN), 7.27 (1H, d, J 8.5Hz, CH), 7.51 (1H, d, J 8.5Hz, CH), 7.86 (1H, s, CH), 8.07 (1H, s, CH).





						s, CH), 9.42 (1H, s, NH); MS(+ve): 523.28; tR = 3.12 min (XBridge 4).
289		***	**		9-Cyclopentyl-2-(1,1-dioxo-1H-1lambda*6*-benzo[b]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹H NMR (DMSO): 1.59 (4H, m, 2CH₂), 1.7 (2H, m, CH₂), 1.94 (2H, m, CH₂), 2.59 (2H, m, CH₂), 3.18 (3H, s, CH₃), 3.62 (2H, m, CH₂), 4.83 (1H, m, CHN), 7.3 (1H, d, J 6.5Hz, CH), 7.49 (1H, d, J 6.5Hz, CH), 7.69 (1H, d, J 8.5Hz, CH), 7.84 (1H, d, J 8.5Hz, CH), 7.98 (1H, s, CH), 8.11 (1H, s, CH), 9.82 (1H, s, NH); MS(+ve): 426.31; tR = 2.95 min (XBridge 4).
290		***	*		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-9-p-tolyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹H NMR (DMSO): 2.38 (3H, s, CH₃), 2.78 (2H, m, CH₂), 3.26 (3H, s, CH₃), 4.0 (2H, m, CH₂), 4.03 (2H, s, CH₂), 4.31 (2H, s, CH₂), 6.93 (1H, d, J 8.0Hz, CH), 7.16 (1H, d, J 8.0Hz, CH), 7.2 (2H, d, J 8.0Hz, 2CH), 7.27 (2H, d, J 8.0Hz, 2CH), 7.32 (1H, s, CH), 8.25 (1H, s, CH), 9.45 (1H, s, NH); MS(+ve): 450.33; tR = 2.82 min (XBridge 4).
291		***	*		Cyclopropanecarboxylic acid [2-methyl-4-(5-methyl-6-oxo-9-p-tolyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-amide	¹H NMR (DMSO): 0.72 (4H, m, 2CH₂), 1.78 (1H, m, CH), 1.87 (3H, s, CH₃), 2.33 (3H, s, CH₃), 2.78 (2H, m, CH₂), 3.25 (3H, s, CH₃), 3.99 (2H, m, CH₂), 6.87 (1H, m, CH), 6.96 (1H, m, CH), 7.16 (3H, m, 3CH), 7.21 (2H, d, J 8.0Hz, 2CH), 8.25 (1H, s, CH), 9.17 (1H, s, NH), 9.24 (1H, s, NH); MS(+ve): 457.41; tR = 2.89 min (XBridge 4).






292		***	**	5-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzo[b]thiophene-2-carboxylic acid	1H NMR (DMSO): 1.60 (4H, m, 2CH ₂), 1.72 (2H, m, CH ₂), 1.97 (2H, m, CH ₂), 2.58 (2H, m, CH ₂), 3.17 (3H, s, CH ₃), 3.61 (2H, m, CH ₂), 4.87 (1H, m, CHN), 7.53 (1H, s, CH), 7.55 (1H, d, J 8.5Hz, CH), 7.73 (1H, d, J 8.5Hz, CH), 8.06 (1H, s, CH), 8.40 (1H, s, CH), 9.29 (1H, s, NH); MS(+ve): 438.28; tR = 1.90 min (XBridge 4).
293		***	*	9-Cyclopentyl-5-methyl-2-[4-(morpholine-4-carbonyl)-phenylamino]-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.59 (4H, m, 2CH ₂), 1.70 (2H, m, CH ₂), 1.94 (2H, m, CH ₂), 2.57 (2H, m, CH ₂), 3.17 (3H, s, CH ₃), 3.5 (4H, m, 2CH ₂), 3.60 (6H, m, 3CH ₂), 4.81 (1H, m, CHN), 7.32 (2H, d, J 8.5Hz, 2CH), 7.79 (2H, d, J 8.5Hz, 2CH), 8.06 (1H, s, CH), 9.43 (1H, s, NH); MS(+ve): 451.40; tR = 2.63 min (XBridge 4).
294		**	**	9-Cyclopentyl-5-methyl-2-(3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-ylamino)-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.58 (4H, m, 2CH ₂), 1.69 (2H, m, CH ₂), 1.93 (2H, m, CH ₂), 2.54 (2H, m, CH ₂), 3.15 (3H, s, CH ₃), 3.58 (2H, m, CH ₂), 4.51 (2H, s, CH ₂), 4.77 (1H, m, CHN), 6.75 (1H, d, J 8.5Hz, CH), 7.2 (1H, d, J 8.5Hz, CH), 7.50 (1H, s, CH), 8.02 (1H, s, CH), 9.13 (1H, s, NH), 10.5 (1H, s, NH); MS(+ve): 409.33; tR = 2.60 min (XBridge 4).
295		***	*	9-Cyclopentyl-5-methyl-2-(3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-6-ylamino)-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	1H NMR (DMSO): 1.54 (4H, m, 2CH ₂), 1.66 (2H, m, CH ₂), 1.91 (2H, m, CH ₂), 2.55 (2H, m, CH ₂), 3.15 (3H, s, CH ₃), 3.58 (2H, m, CH ₂), 4.48 (2H, s, CH ₂), 4.74 (1H, m, CHN), 6.82 (1H, d, J 8.5Hz, CH), 7.22 (2H, m, 2CH), 7.99 (1H, s, CH), 9.05 (1H, s, NH), 10.65 (1H, s, NH); MS(+ve): 409.33; tR = 2.75 min (XBridge 4).





296		***	**	5-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzo[b]thiophene-2-carboxylic acid	
297		***	*	9-Cyclopentyl-5-methyl-2-[4-(morpholine-4-carbonyl)-phenylamino]-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
298		**	**	9-Cyclopentyl-5-methyl-2-(3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-ylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	
299		***	*	9-Cyclopentyl-5-methyl-2-(3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-6-ylamino)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	





300		***	*	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N,N-dimethyl-benzenesulfonamide	1H NMR (DMSO): 1.59 (4H, m, 2CH ₂), 1.70 (2H, m, CH ₂), 1.95 (2H, m, CH ₂), 2.57 (8H, m, 2CH ₃ , CH ₂), 3.17 (3H, s, CH ₃), 3.62 (2H, m, CH ₂), 4.82 (1H, m, CHN), 7.6 (2H, d, J 8Hz, 2CH), 7.96 (2H, d, J 8Hz, 2CH), 8.10 (1H, s, CH), 9.73 (1H, s, NH); MS(+ve): 445.36; tR = 3.16 min (XBridge 4).
301			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pirimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methyl-N-(1-methyl-piperidin-4-yl)-benzamide	
302			**	4-(9-Isobutyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
303			**	4-(9-Isobutyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(1-methyl-piperidin-4-yl)-benzamide	
304			**	N-(1-Benzyl-piperidin-4-yl)-4-(9-isobutyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzamide	





309			*	3-Methoxy-4-(5-methyl-6-oxo-9-o-tolyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-methyl-piperidin-3-yl)-benzamide	
310			*	N-(1-Benzyl-piperidin-4-yl)-3-methoxy-4-(5-methyl-6-oxo-9-o-tolyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzamide	
311			**	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-3-methoxy-4-(5-methyl-6-oxo-9-o-tolyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzamide	
312			*	3-Methoxy-4-(5-methyl-6-oxo-9-o-tolyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1,2,2,6,6-pentamethyl-piperidin-4-yl)-benzamide	





313			*	N-(2-Diethylamino-ethyl)-3-methoxy-4-(5-methyl-6-oxo-9-o-tolyl-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-benzamide	
314			**	3-Methoxy-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-4-yl)-benzamide	
315			**	3-Methoxy-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1-methyl-piperidin-3-yl)-benzamide	
316			**	N-(1-Benzyl-piperidin-4-yl)-3-methoxy-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzamide	





317			**	N-(1-Aza-bicyclo[2.2.2]oct-3-yl)-3-methoxy-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzamide	
318			**	3-Methoxy-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-N-(1,2,2,6,6-pentamethyl-piperidin-4-yl)-benzamide	
319			*	N-(2-Diethylamino-ethyl)-3-methoxy-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzamide	
320			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(1-ethyl-pyrrolidin-3-yl)-3-methoxy-N-benzamide	
321			**	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-[2-(4-iso	





322				***	N-(1-Benzyl-pyrrolidin-3-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-N-benzamide	
323				***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-fluoro-N-(1-methyl-piperidin-4-yl)-benzamide	1H NMR (DMSO): 1.60 (4H, m, 2CH2), 1.70 (2H, m, CH2), 1.96 (2H, m, CH2), 2.04 (3H, s, CH3), 2.58 (2H, m, CH2), 3.16 (3H, s, CH3), 3.61 (2H, m, CH2), 4.83 (1H, m, CHN), 7.42 (2H, s, 2CH), 8.06 (1H, s, CH), 8.15 (1H, s, CH), 9.38 (1H, s, NH), 9.40 (1H, s, NH); MS(+ve): 429.28; tR = 2.76 min (XBridge 4).
324		**	*	*	N-[2-Chloro-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-phenyl]-acetamide	MS(+ve): 535.27; tR = 2.54 min (XBridge 4).
325		**			4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1,1-dioxo-tetrahydro-1lambda*6*-thiophen-3-yl)-benzenesulfonamide	






326		***	*		MS(+ve): 431.25; tR = 2.80 min (XBridge 4).
327		***			1H NMR (DMSO): 0.73 (4H, m, 2CH2), 1.78 (1H, m, CH), 1.89 (3H, s, CH3), 2.77 (2H, m, CH2), 3.25 (3H, s, CH3), 3.79 (3H, s, CH3), 3.98 (2H, m, CH2), 6.85 (1H, d, J 8.5Hz, CH), 6.96 (1H, m, CH), 6.98 (2H, d, J 8.5Hz, 2CH), 7.11 (1H, s, CH), 7.23 (2H, d, J 8.5Hz, 2CH), 8.19 (1H, s, CH), 9.11 (1H, s, NH), 9.25 (1H, s, NH); MS(+ve): 473.35; tR = 2.73 min (XBridge 4).
328		*			1H NMR (DMSO): 0.73 (4H, m, 2CH2), 1.05 (3H, t, J 7Hz, CH3), 1.78 (1H, m, CH), 1.87 (3H, s, CH3), 2.32 (3H, s, CH3), 2.68 (2H, m, CH2), 3.78 (2H, q, J 7Hz, CH2), 3.98 (2H, m, CH2), 6.93 (1H, m, CH), 7.0 (1H, m, CH), 7.09 (2H, d, J 8Hz, 2CH), 7.19 (2H, d, J 8.0Hz, 2CH), 7.23 (1H, s, CH), 8.32 (1H, s, CH), 9.28 (2H, m, 2NH); MS(+ve): 471.34; tR = 3.05 min (XBridge 4).
329		**			1H NMR (DMSO): 0.73 (4H, m, 2CH2), 1.77 (1H, m, CH), 1.96 (3H, s, CH3), 2.86 (2H, m, CH2), 3.27 (3H, s, CH3), 3.89 (2H, m, CH2), 6.77 (1H, d, J 9Hz, CH), 6.90 (1H, d, J 9Hz, CH), 6.99 (1H, s, CH), 7.29 (2H, m, 2CH), 7.52 (1H, m, CH), 8.24 (1H, s, CH), 9.22 (1H, s, NH), 9.3 (1H, s, NH); MS(+ve): 479.27; tR = 2.78 min (XBridge 4).

330		*		Cyclopropanecarboxylic acid {4-[5-ethyl-9-(4-methoxy-phenyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-2-methyl-phenyl}-amide	¹ H NMR (DMSO): 0.73 (4H, m, 2CH ₂), 1.06 (3H, t, J 7Hz, CH ₃), 1.78 (1H, m, CH), 1.89 (3H, s, CH ₃), 2.70 (2H, m, CH ₂), 3.78 (5H, m, CH ₂ , CH ₃), 3.98 (2H, m, CH ₂), 6.90 (1H, d, J 8.5Hz, CH), 6.97 (3H, m, 3CH), 7.17 (3H, m, 3CH), 8.25 (1H, s, CH), 9.20 (1H, s, NH), 9.26 (1H, s, NH); MS(+ve): 487.31; tR = 2.86 min (XBridge 4).
331		***		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-ethyl-9-p-tolyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.06 (3H, t, J 7Hz, CH ₃), 2.36 (3H, s, CH ₃), 2.70 (2H, m, CH ₂), 3.78 (2H, q, J 7Hz, CH ₂), 3.99 (2H, m, CH ₂), 4.04 (2H, s, CH ₂), 4.32 (2H, s, CH ₂), 6.97 (1H, d, J 8.0Hz, CH), 7.13 (2H, d, J 8.0Hz, 2CH), 7.2 (1H, d, J 8.0Hz, CH), 7.25 (2H, d, J 8.0Hz, 2CH), 7.38 (1H, s, CH), 8.33 (1H, s, CH), 9.53 (1H, s, NH); MS(+ve): 464.29; tR = 2.96 min (XBridge 4).
332		***		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-9-(4-methoxy-phenyl)-5-methyl-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.8 (2H, m, CH ₂), 3.26 (3H, s, CH ₃), 3.82 (3H, s, CH ₃), 3.99 (2H, m, CH ₂), 4.02 (2H, s, CH ₂), 4.31 (2H, s, CH ₂), 6.94 (1H, d, J 8.5Hz, CH), 7.05 (2H, d, J 8Hz, 2CH), 7.15 (1H, d, J 8.5Hz, CH), 7.27 (3H, m, 3CH), 8.20 (1H, s, CH), 9.41 (1H, s, NH); MS(+ve): 466.23; tR = 2.67 min (XBridge 4).
333		***		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-ethyl-9-(4-methoxy-phenyl)-5,7,8,9-tetrahydro-pyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.07 (3H, t, J 7Hz, CH ₃), 2.72 (2H, m, CH ₂), 3.78 (2H, q, J 7Hz, CH ₂), 3.81 (3H, s, CH ₃), 3.99 (2H, m, CH ₂), 4.03 (2H, s, CH ₂), 4.32 (2H, s, CH ₂), 6.97 (1H, d, J 8.5Hz, CH), 7.04 (2H, d, J 8.5Hz, 2CH), 7.18 (1H, m, CH), 7.21 (2H, d, J 8.5Hz, 2CH), 7.32 (1H, s, CH), 8.26 (1H, s, CH), 9.48 (1H, s, NH); MS(+ve): 480.22; tR = 2.82 min (XBridge 4).

334		*		9-(4-tert-Butyl-phenyl)-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.33 (9H, s, C(CH ₃) ₃), 2.76 (2H, m, CH ₂), 3.26 (3H, s, CH ₃), 3.99 (2H, m, CH ₂), 4.12 (2H, s, CH ₂), 4.29 (2H, s, CH ₂), 6.92 (1H, d, J 8Hz, CH), 7.20 (2H, d, J 8.5Hz, 2CH), 7.28 (1H, d, J 8Hz, CH), 7.44 (3H, m, 3CH), 8.31 (1H, s, CH), 9.5 (1H, s, NH); MS(+ve): 492.36; tR = 3.32 min (XBridge 4).
335		*		9-(4-tert-Butyl-phenyl)-2-(2,2-dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-ethyl-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 1.04 (3H, t, J 7Hz, CH ₃), 1.32 (9H, s, C(CH ₃) ₃), 2.68 (2H, m, CH ₂), 3.78 (2H, q, J 7Hz, CH ₂), 3.98 (2H, m, CH ₂), 4.12 (2H, s, CH ₂), 4.31 (2H, s, CH ₂), 6.98 (1H, d, J 8.5Hz, CH), 7.12 (2H, d, J 8.5Hz, 2CH), 7.31 (1H, d, J 8.5Hz, CH), 7.41 (2H, d, J 8.5Hz, 2CH), 7.52 (1H, s, CH), 8.4 (1H, s, CH), 9.6 (1H, s, NH); MS(+ve): 506.32; tR = 3.45 min (XBridge 4).
336		***		2-(2,2-Dioxo-2,3-dihydro-1H-2lambda*6*-benzo[c]thiophen-5-ylamino)-5-methyl-9-(4-morpholin-4-yl-phenyl)-5,7,8,9-tetrahydropyrimido[4,5-b][1,4]diazepin-6-one	¹ H NMR (DMSO): 2.79 (2H, m, CH ₂), 3.19 (4H, m, 2CH ₂), 3.26 (3H, s, CH ₃), 3.79 (4H, m, 2CH ₂), 3.98 (2H, m, CH ₂), 4.04 (2H, s, CH ₂), 4.30 (2H, s, CH ₂), 7.01 (1H, d, J 8.5Hz, CH), 7.05 (2H, d, J 8.5Hz, 2CH), 7.10 (1H, d, J 8.5Hz, CH), 7.20 (2H, d, J 8.5Hz, 2CH), 7.31 (1H, s, CH), 8.18 (1H, s, CH), 9.39 (1H, s, NH); MS(+ve): 521.30; tR = 2.60 min (XBridge 4).
337		***		Cyclopropanecarboxylic acid {2-methyl-4-[5-methyl-9-(4-morpholin-4-yl-phenyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-phenyl}-amide	¹ H NMR (DMSO): 0.73 (4H, m, 2CH ₂), 1.79 (1H, m, CH), 1.92 (3H, s, CH ₃), 2.77 (2H, m, CH ₂), 3.13 (4H, m, 2CH ₂), 3.25 (3H, s, CH ₃), 3.75 (4H, m, 2CH ₂), 3.97 (2H, m, CH ₂), 6.89 (1H, d, J 8Hz, CH), 6.98 (3H, m, 3CH), 7.10 (1H, s,

					CH), 7.16 (2H, d, J 8.5Hz, 2CH), 8.17 (1H, s, CH), 9.09 (1H, s, NH), 9.21 (1H, s, NH); MS(+ve): 528.33; tR = 2.60 min (XBridge 4).
338		**			1H NMR (DMSO): 1.07 (3H, t, J 7Hz, CH3), 2.71 (2H, m, CH2), 3.18 (4H, m, 2CH2), 3.79 (6H, m, 3CH2), 3.97 (2H, m, CH2), 4.05 (2H, s, CH2), 4.30 (2H, s, CH2), 7.03 (3H, m, 3CH), 7.14 (3H, m, 3CH), 7.35 (1H, s, CH), 8.24 (1H, s, CH), 9.46 (1H, s, NH); MS(+ve): 535.27; tR = 2.72 min (XBridge 4).
339		***			1H NMR (DMSO): 2.87 (2H, m, CH2), 3.28 (3H, s, CH3), 3.91 (2H, m, CH2), 4.18 (2H, s, CH2), 4.30 (2H, s, CH2), 6.91 (1H, d, J 8Hz, CH), 7.13 (2H, m, 2CH), 7.35 (2H, m, 2CH), 7.57 (1H, m, CH), 8.27 (1H, s, CH), 9.53 (1H, s, NH); MS(+ve): 472.18; tR = 2.72 min (XBridge 4).
340		**			1H NMR (DMSO): 1.03 (3H, t, J 7Hz, CH3), 2.80 (2H, m, CH2), 3.83 (2H, q, J 7Hz, CH2), 3.9 (2H, m, CH2), 4.18 (2H, s, CH2), 4.31 (2H, s, CH2), 6.93 (1H, d, J 9Hz, CH), 7.16 (2H, m, 2CH), 7.34 (2H, m, 2CH), 7.57 (1H, m, CH), 8.32 (1H, s, CH), 9.57 (1H, s, NH); MS(+ve): 486.22; tR = 2.87 min (XBridge 4).
341			***		

342		*		Cyclopropanecarboxylic acid {4-[9-(4-tert-butyl-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-2-methyl-phenyl}-amide	¹ H NMR (DMSO): 0.71 (4H, m, 2CH ₂), 1.30 (9H, s, C(CH ₃) ₃), 1.77 (1H, m, CH), 1.91 (3H, s, CH ₃), 2.75 (2H, m, CH ₂), 3.25 (3H, s, CH ₃), 3.99 (2H, m, CH ₂), 6.88 (1H, d, J 8.5Hz, CH), 7.05 (1H, d, J 8.5Hz, CH), 7.17 (2H, d, J 8.5Hz, 2CH), 7.19 (1H, s, CH), 7.41 (2H, d, J 8.5Hz, 2CH), 8.28 (1H, s, CH), 9.22 (2H, s, 2NH); MS(+ve): 499.45; tR = 3.37 min (XBridge 4).
343			**	{4-[4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-benzoylamino]-piperidin-1-yl}-acetic acid	
344			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-(1-isopropyl-piperidin-4-yl)-3-methoxy-benzamide	
345			***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide	

346					4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxy-benzoyl acid tert-butyl ester	
347		***			N-(R)-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxybenzamide	
348		***			N-(S)-(1-Aza-bicyclo[2.2.2]oct-3-yl)-4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-3-methoxybenzamide	
349		**	*		4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-spiro[pyrimido[4,5-b][1,4]diazepin-3,1'-cyclopropane]-2-ylamino)-N-(1-methyl-piperidin-4-yl)-benzamide	
350		**			N-Acetyl-4-[9-(4-methoxy-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]benzenesulfonamide	


















351		***				N-Acetyl-4-[9-(2,6-difluoro-phenyl)-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzenesulfonamide		
352		**				N-Acetyl-4-[5-methyl-9-(4-morpholin-4-yl-phenyl)-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino]-benzenesulfonamide		

TABLE 3: PLK IC₅₀ values and characterization data for selected compounds of the invention; *** denotes <0.1 μ M IC₅₀; ** denotes between 1 μ M and 0.1 μ M IC₅₀; * denotes between 10 μ M and 1 μ M IC₅₀




No.	Structure	PLK 1	Name	Data
254		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide	Rt = 2.63 min (Analytical_1); MS(+ve): 535.5; MS(-ve): 533.6; ¹ H NMR (DMSO-d ₆) δ ppm: 0.63 - 0.70 (2 H, m), 0.87 - 0.94 (2 H, m), 1.43 - 1.54 (2 H, m), 1.55 - 1.63 (2 H, m), 1.64 - 1.73 (2 H, m), 1.83 - 1.94 (2 H, m), 2.18 (3 H, s), 2.31 - 2.48 (4 H, m), 2.92 (4 H, m), 3.16 (3 H, s), 3.47 (2 H, s), 3.94 (3 H, s), 4.78 - 4.91 (1 H, m), 7.35 - 7.48 (2 H, m), 7.68 (1 H, s), 7.98 (1 H, s), 8.39 (1 H, d, J=8.3 Hz), 9.31 (1 H, s).
218		***	($\pm$)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide	Rt = 3.26 min (Analytical_1); MS(+ve): 560.4; ¹ H NMR (DMSO-d ₆) δ ppm: 8.37 (d, J = 8.3 Hz, 1H), 8.08 (d, J = 6.8 Hz, 1H), 8.05 (s, 1H), 7.73 (s, 1H), 7.45 - 7.53 (m, 2H), 4.82 (quin, J = 8.3 Hz, 1H), 3.84 - 4.01 (m, 4H), 3.64 (s, 2H), 3.18 (s, 3H), 3.04 - 3.14 (m, 1H), 2.88 (t, J = 9.8 Hz, 1H), 2.58 - 2.75 (m, 4H), 2.21 - 2.33 (m, 2H), 2.07 (s, 2H), 1.97 (br. s., 2H), 1.87 (br. s., 2H), 1.76 (d, J = 4.9 Hz, 2H), 1.56 - 1.70 (m, 8H), 1.30 (br. s., 1H)
195		***	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide	Rt = 2.81 min (Analytical_1); MS(+ve): 537.5; ¹ H NMR (DMSO-d ₆) δ ppm: 1.09 (6H, s), 1.61 (4H, m), 1.73 (2H, m), 1.87 (2H, m), 2.19 (3H, s), 2.42 (2H, m), 2.92 (4H, m), 3.29 (3H, s), 3.37 (3H, m), 3.93 (3H, s), 3.17 (1H, m), 7.40 (2H, m), 7.68 (1H, s), 7.98 (1H, s), 8.37 (1H, d, J 7Hz), 9.30 (1H, s).





221		***	4-(9-cyclopentyl-5-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide	Rt = 2.79 min (Analytical_1); MS(+ve): 549.5; 1H NMR (DMSO-d6) δ ppm: 9.31 (s, 1H), 8.37 (d, J = 8.3 Hz, 1H), 8.05 (s, 1H), 7.72 (s, 1H), 7.37 - 7.47 (m, 2H), 4.73 - 4.86 (m, 1H), 3.93 (s, 3H), 3.64 (s, 2H), 3.18 (s, 3H), 2.92 (br. s., 4H), 2.36 (br. s., 4H), 2.27 (d, J = 9.3 Hz, 2H), 2.18 (s, 3H), 1.97 (br. s., 2H), 1.79 - 1.91 (m, 1H), 1.76 (br. s., 2H), 1.54 - 1.72 (m, 7H).
371		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 3.18 min (Analytical_1); ES(+ve): 657.6; ES(-ve): 655.7; 1H NMR (DMSO-d6) δ ppm: 0.05 (d, J = 4.20 Hz, 3H), 0.44 (d, J = 7.74 Hz, 2H), 0.67 (d, J = 1.61 Hz, 2H), 0.74 - 0.85 (m, 1H), 0.90 (s, 2H), 1.23 - 1.44 (m, 3H), 1.44 - 1.55 (m, 2H), 1.54 - 1.64 (m, 2H), 1.64 - 1.74 (m, 2H), 1.88 (br. s., 5H), 2.08 (s, 3H), 2.13 (d, J = 6.45 Hz, 2H), 2.17 - 2.28 (m, 1H), 2.29 - 2.47 (m, 3H), 2.60 - 2.67 (m, 1H), 3.06 - 3.23 (m, 5H), 3.47 (s, 2H), 3.65 - 3.79 (m, 1H), 3.94 (s, 2H), 4.05 - 4.13 (m, 1H), 4.78 - 4.91 (m, 1H), 7.48 (s, 2H), 7.67 (s, 1H), 7.98 (s, 1H), 8.00 - 8.07 (m, 1H), 8.31 - 8.45 (m, 1H).
372		***	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 3.38 min (Analytical_1); ES(+ve): 659.6; ES(-ve): 657.7; 1H NMR (DMSO-d6) δ ppm: 0.05 (d, J = 3.86 Hz, 3H), 0.40 - 0.48 (m, 3H), 0.73 - 0.84 (m, 2H), 1.03 - 1.13 (m, 10H), 1.18 (s, 1H), 1.24 - 1.43 (m, 6H), 1.61 (br. s., 3H), 1.74 (d, J = 9.65 Hz, 2H), 1.88 (br. s., 5H), 2.08 (s, 5H), 2.13 (d, J = 6.75 Hz, 2H), 2.17 - 2.25 (m, 1H), 2.36 (br. s., 3H), 2.60 - 2.67 (m, 1H), 3.14 - 3.22 (m, 2H), 3.37 (s, 1H), 3.51 (s, 1H), 3.94 (s, 2H), 5.18 (t, J = 8.52 Hz, 1H), 7.43 - 7.50 (m, 1H), 7.67 (s, 1H), 7.98 (s, 1H), 8.03 (d, J = 7.72 Hz, 1H), 8.35 (d, J = 8.36 Hz, 1H).
345		***	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide	Rt = 2.74 min (Analytical_1); ES(+ve): 578.5; ES(-ve): 576.7.

373		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide	Rt = 2.74 min (Analytical_1); MS(+ve): 537.5; 1H NMR (DMSO-d6) δ ppm: 1.55 – 1.65 (8H, m), 1.71 (2H, m), 1.88 (2H, m), 2.18 (3H, s), 2.28 (2H, q, J 10.5 Hz), 2.41 – 2.52 (4H, m), 2.89 (4H, m), 3.18 (3H, s), 3.61 (2H, s), 4.76 (1H, quintet, 8.5 Hz), 7.61 (2H, m), 8.03 (1H, s), 8.12 (1H, dd, J 8 Hz, 8.5 Hz), 8.76 (1H, s), 9.36 (1H, s).
374		***	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide	Rt = 2.58 min (Analytical_1); MS(+ve): 523.4; 1H NMR (DMSO-d6) δ ppm: 0.65 – 0.67 (2H, s, CH2), 0.85 – 0.95 (2H, m, CH2), 1.23 – 1.82 (8H, m, alkyl-H), 2.18 (3H, s, CH3), 2.36 – 2.41 (4H, m, alkyl-H), 2.88 (4H, s, alkyl-H), 3.16 (3H, s, CH3), 3.44 (2H, s, alkyl-H), 4.77 – 4.80 (1H, m, CH), 7.6 (2H, m, aryl-H), 7.96 (1H, s, aryl-H), 8.16 (1H, t, J = 8 Hz, aryl-H), 8.70 (1H, s, NH), 9.35 (1H, s, NH).
194		***	(±)-4-(9'-cyclopentyl-5',7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepine-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide	Rt = 3.28 min (Analytical_1); MS(+ve): 548.5; 1H NMR (DMSO-d6) δ ppm: 8.37 (1H, d, J 8 Hz), 8.07 (1H, d, J 6.5 Hz), 7.98 (1H, s), 7.69 (1H, s), 7.47 – 7.50 (2H, m), 5.19 (1H, quintet, J 8 Hz), 3.95 (4H, m), 3.37 (2H, s), 3.18 (3H, s), 3.11 (1H, m), 2.89 (1H, m), 2.65 – 2.72 (4H, m), 1.88 (2H, m), 1.73 – 1.87 (4H, m), 1.57 – 1.61 (6H, m), 1.31 (1H, m), 1.09 (6H, s).
186		***	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide	Rt = 3.02 min (Analytical_1); MS(+ve): 546.4; 1H NMR (DMSO-d6) δ ppm: 8.40 (1H, d, J 8 Hz), 8.08 (1H, d, J 7 Hz), 7.98 (1H, s), 7.69 (1H, s), 7.47 – 7.50 (2H, m), 4.87 (1H, quintet, J 9 Hz), 4.10 (1H, q, J 5.5 Hz), 3.95 (4H, m), 3.47 (2H, s), 3.27 (3H, s), 3.07 (1H, m), 2.08 (1H, m), 2.63 – 2.86 (4H, m), 1.88 (3H, m), 1.78 (1H, m), 1.68 (2H, m), 1.50 – 1.62 (5H, m), 1.32 (1H, m), 0.90 (2H, m), 0.66 (2H, m).

375		***	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide	Rt = 2.48 min (Analytical_1); MS(+ve): 565.4; ¹ H NMR (DMSO) δ ppm: 0.66 – 0.67 (2H, m, CH ₂), 0.88 – 0.90 (2H, m, CH ₂), 1.23 – 1.90 (8H, m, alkyl-H), 2.34-2.41 (4H, m, alkyl-H), 2.91 (4H, t, J = 4 Hz, alkyl-H), 3.16 (3H, s, CH ₃), 3.46-3.51 (4H, m), 3.93 (3H, s, CH ₃), 4.38-4.45 (1H, m), 4.81-4.87 (1H, m, CH), 7.40 (2H, d, J = 10.5 Hz, aryl-H), 7.67 (1H, s), 7.98 (1H, s), 8.39 (1H, d, J = 8 Hz, aryl-H), 9.30 (1H, s, NH).
376		***	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide	Rt = 2.72 min (Analytical_1); MS(+ve): 522.4; ¹ H NMR (DMSO) δ ppm: 0.65 – 0.71 (2H, m, CH ₂), 0.85 – 0.91 (2H, m, CH ₂), 1.23 – 1.88 (8H, m, alkyl-H), 2.92 (4H, t, J = 4 Hz, alkyl-H), 3.16 (3H, s, CH ₃), 3.47 (2H, s, alkyl-H), 3.66 (4H, t, J = 4.5 Hz), 3.94 (3H, s, CH ₃), 4.81 – 4.87 (1H, m, CH), 7.41 (2H, d, J = 9.5 Hz), 7.69 (1H, s), 7.98 (1H, s), 8.40 (1H, d, J = 8 Hz), 9.41 (1H, s, NH).
347		***	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide	Rt = 3.07 min (Analytical_1); MS(+ve): 546.5; MS(-ve): 544.6; ¹ H NMR (DMSO-d ₆) δ ppm: 0.62 – 0.72 (2H, m), 0.87 – 0.94 (2H, m), 1.20 – 1.38 (1H, m), 1.44 – 1.54 (2H, m), 1.54 – 1.64 (4H, m), 1.64 – 1.74 (2H, m), 1.75 – 1.85 (1H, m), 1.85 – 1.95 (3H, m), 2.65 – 2.79 (4H, m), 2.83 – 3.01 (1H, m), 3.11 – 3.21 (4H, m), 3.47 (2H, s), 3.95 (4H, s), 4.77 – 4.93 (1H, m), 7.42 – 7.54 (2H, m), 7.69 (1H, s), 7.98 (1H, s), 8.10 (1H, d, J=6.3 Hz), 8.39 (1H, d, J=8.3 Hz).
348		***	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide	Rt = 3.04 min (Analytical_1); MS(+ve): 546.5; MS(-ve): 544.6; ¹ H NMR (DMSO-d ₆) δ ppm: 0.59 – 0.74 (2H, m), 0.86 – 0.97 (2H, m), 1.20 – 1.43 (1H, m), 1.45 – 1.75 (7H, m), 1.76 – 1.98 (4H, m), 2.66 – 2.86 (3H, m), 2.89 – 3.07 (1H, m), 3.17 (3H, s), 3.47 (3H, s), 3.89 – 4.02 (4H, m), 4.76 – 4.96 (1H, m), 7.42 – 7.59 (2H, m), 7.70 (1H, s), 7.98 (1H, s), 8.14 (1H, d, J=6.3 Hz), 8.40 (1H, d, J=7.8 Hz).

377		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 3.41 min (Analytical_1); ES(+ve): 671.6; ES(-ve): 669.7; 1H NMR (DMSO-d ₆) δ ppm: 0.03 (2H, dd, J 9.5 Hz, 5 Hz), 0.41 (2H, dd, J 10 Hz, 5 Hz), 0.79 (1H, m), 1.20 - 1.37 (4H, m), 1.55 - 1.94 (18H, m), 2.04 (2H, d, J 6.5 Hz), 2.16 - 2.46 (8H, m), 3.15 (3H, s), 3.32 (2H, s), 3.69 (1H, m), 3.90 (3H, s), 4.79 (1H, m), 7.44 (2H, m), 7.68 (1H, s), 8.01 (2H, m), 8.34 (1H, d, J 8.5 Hz)
378		***	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2'-ylamino)-N-((trans)-4-(4-cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 2.88 min (Analytical_1); ES(+ve): 631.5; ES(-ve): 629.6; 1H NMR (DMSO-d ₆) δ ppm: -0.15 - 0.11 (m, 2 H), 0.40 (d, J = 8.06 Hz, 2 H), 0.63 - 0.85 (m, 1 H), 1.11 - 1.41 (m, 3 H), 1.57 (br. s., 2 H), 1.69 (d, J = 17.73 Hz, 2 H), 1.74 - 1.95 (m, 6 H), 2.04 (s, 2 H), 2.09 (d, J = 6.45 Hz, 2 H), 2.16 (d, J = 9.67 Hz, 1 H), 2.32 (br. s., 2 H), 2.49 (br. s., 1 H), 2.51 - 2.59 (m, 2 H), 2.59 (br. s., 1 H), 3.07 - 3.17 (m, 3 H), 3.27 (s, 5 H), 3.53 - 3.62 (m, 2 H), 3.67 (dd, J = 7.41, 3.55 Hz, 1 H), 3.89 (s, 2 H), 4.04 (q, J = 5.37 Hz, 1 H), 4.76 (quin, J = 8.22 Hz, 1 H), 7.38 - 7.48 (m, 2 H), 7.68 (s, 1 H), 7.99 (d, J = 7.74 Hz, 1 H), 8.03 (s, 1 H), 8.33 (d, J = 8.06 Hz, 1 H)
379		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 3.00 min (Analytical_1) MS(+ve) 631.6; MS(-ve) 629.7; 1H NMR (DMSO-d ₆) δ ppm: 0.59 - 0.72 (2 H, m), 0.81 - 0.92 (2 H, m), 0.97 (3 H, t, J=7.1 Hz), 1.20 - 1.42 (4 H, m), 1.42 - 1.55 (2 H, m), 1.54 - 1.64 (2 H, m), 1.64 - 1.73 (2 H, m), 1.83 (2 H, d, J=11.7 Hz), 1.89 (4 H, d, J=7.8 Hz), 2.14 - 2.42 (7 H, m), 3.16 (3 H, s), 3.47 (2 H, s), 3.64 - 3.79 (1 H, m), 3.94 (3 H, s), 4.84 (1 H, quin, J=8.7 Hz), 7.40 - 7.54 (2 H, m), 7.67 (1 H, s), 7.98 (1 H, s), 8.03 (1 H, d, J=7.8 Hz), 8.38 (1 H, d, J=8.3 Hz)
380		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 3.13 min (Analytical_1) MS(+ve) 631.6; MS(-ve) 629.7; 1H NMR (DMSO-d ₆) δ ppm: 0.60 - 0.72 (2 H, m), 0.83 - 0.93 (2 H, m), 0.98 (3 H, t, J=7.1 Hz), 1.38 - 1.55 (6 H, m), 1.55 - 1.63 (2 H, m), 1.63 - 1.80 (4 H, m), 1.80 - 1.96 (4 H, m), 2.02 - 2.17 (1 H, m), 2.16 - 2.44 (7 H, m), 3.16 (3 H, s), 3.47 (2 H, s), 3.79 - 3.92 (1 H, m), 3.94 (3 H, s), 4.84 (1 H, quin, J=8.5 Hz), 7.44 - 7.58 (2 H, m), 7.67 (1 H, s), 7.98 (1 H, s)

			methoxybenzamide	H, s), 8.03 (1 H, d, J=7.3 Hz), 8.37 (2 H, d, J=8.8 Hz)
381		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 4.12 min (Analytical_1) MS(+ve) 671.6; MS(-ve) 669.7; ¹ H NMR (DMSO-d ₆) δ ppm: 0.01 (2 H, d, J=4.4 Hz), 0.40 (2 H, d, J=7.3 Hz), 0.63 (2 H, br. s.), 0.71 - 0.83 (1 H, m), 0.87 (2 H, br. s.), 1.02 (1 H, t, J=7.1 Hz), 1.25 - 1.38 (2.5 H, m), 1.39 - 1.52 (3.5 H, m), 1.52 - 1.61 (2 H, m), 1.65 (4 H, d, J=5.9 Hz), 1.70 - 1.80 (3 H, m), 1.85 (3 H, d, J=4.4 Hz), 2.21 - 2.31 (1 H, m), 2.55 - 2.83 (8 H, m), 3.13 (3.5 H, s), 3.35 - 3.55 (3 H, m), 3.69 (0.5 H, br. s.), 3.91 (3.5 H, s), 4.31 (0.5 H, t, J=5.1 Hz), 4.81 (1 H, t, J=8.5 Hz), 7.25 - 7.54 (2 H, m), 7.64 (1 H, s), 7.84 - 8.16 (2 H, m), 8.35 (1 H, d, J=8.3 Hz)
382		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 3.98 min, 4.18 min (Analytical_1) MS(+ve) 645.6; MS(-ve) 643.7; ¹ H NMR (DMSO-d ₆) δ ppm: 0.57 - 0.74 (2 H, m), 0.82 - 0.93 (2 H, m), 0.95 - 1.09 (3 H, m), 1.30 - 1.42 (2 H, m), 1.42 - 1.54 (3.5 H, m), 1.58 (2 H, d, J=4.4 Hz), 1.63 - 1.75 (3.5 H, m), 1.75 - 1.83 (1.5 H, m), 1.88 (3.5 H, br. s.), 1.98 - 2.16 (1.5 H, m), 2.59 - 2.86 (4 H, m), 3.17 (4 H, s), 3.41 - 3.53 (3.5 H, m), 3.78 (2.5 H, m), 3.88 - 4.02 (3.5 H, m), 4.06 - 4.38 (1 H, m), 4.76 - 4.91 (1 H, m), 7.39 - 7.57 (2 H, m), 7.59 - 7.78 (1 H, m), 7.98 (2 H, s), 8.31 - 8.45 (1 H, m)
383		**	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide	Rt = 4.40 min (Analytical_1) MS(+ve) 707.6; MS(-ve) 705.7; ¹ H NMR (DMSO-d ₆) δ ppm: 0.73 (2 H, br. s.), 0.96 (2 H, br. s.), 1.22 - 2.22 (19 H, m), 2.64 - 2.91 (4 H, m), 3.23 (5 H, s), 3.53 (8 H, s), 3.89 - 4.24 (4 H, m), 4.91 (1 H, t, J=8.5 Hz), 5.82 (0 H, s), 7.23 - 8.54 (10 H, m)

384		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 2.85 min (Analytical_1) MS(+ve) 617.5; MS(-ve) 615.6; 1H NMR (DMSO-d6) δ ppm: 0.61 - 0.76 (2 H, m), 0.89 (1 H, t, J=6.8 Hz), 0.92 - 0.98 (2 H, m), 1.22 - 1.47 (5 H, m), 1.47 - 1.79 (6 H, m), 1.81 - 2.02 (5 H, m), 2.07 - 2.20 (3 H, m), 2.20 - 2.42 (4 H, m), 3.20 (4 H, s), 3.34 (3 H, s), 3.51 (2 H, s), 3.67 - 3.82 (1 H, m), 3.98 (3 H, s), 4.88 (1 H, quin, J=8.5 Hz), 7.42 - 7.59 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.08 (1 H, d, J=7.8 Hz), 8.42 (1 H, d, J=8.3 Hz)
385		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide	Rt = 2.93 min (Analytical_1) MS(+ve) 617.5; MS(-ve) 615.6; 1H NMR (DMSO-d6) δ ppm: 0.62 - 0.80 (2 H, m), 0.84 - 1.02 (4 H, m), 1.18 - 1.36 (4 H, m), 1.43 - 1.84 (12 H, m), 1.84 - 2.03 (4 H, m), 2.09 - 2.22 (4 H, m), 2.41 (4 H, br. s.), 3.21 (3 H, s), 3.52 (2 H, s), 3.89 - 3.97 (1 H, m), 3.99 (3 H, s), 4.89 (1 H, quin, J=8.5 Hz), 7.47 - 7.61 (2 H, m), 7.72 (1 H, s), 8.03 (1 H, s), 8.08 (1 H, d, J=7.3 Hz), 8.43 (1 H, d, J=9.3 Hz)
386		***	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide	Rt = 3.64 min (Analytical_1) MS(+ve) 693.6; MS(-ve) 691.7; 1H NMR (DMSO-d6) δ ppm: 0.62 - 0.79 (2 H, m), 0.85 - 1.02 (2 H, m), 1.20 - 1.46 (4 H, m), 1.46 - 1.58 (2 H, m), 1.59 - 1.67 (2 H, m), 1.67 - 1.78 (2 H, m), 1.80 - 2.02 (6 H, m), 2.12 (1 H, s), 2.18 - 2.46 (6 H, m), 3.20 (3 H, s), 3.43 - 3.56 (5 H, m), 3.67 - 3.84 (1 H, m), 3.97 (3 H, s), 4.88 (1 H, quin, J=8.4 Hz), 7.22 - 7.42 (6 H, m), 7.44 - 7.59 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.08 (1 H, d, J=7.8 Hz), 8.42 (1 H, d, J=7.8 Hz)
387		**	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide	Rt = 3.85 min (Analytical_1) MS(+ve) 693.6; MS(-ve) 691.7; 1H NMR (DMSO-d6) δ ppm: 0.62 - 0.79 (2 H, m), 0.87 - 1.01 (2 H, m), 1.43 - 1.59 (5 H, m), 1.59 - 1.67 (2 H, m), 1.66 - 1.74 (2 H, m), 1.75 - 1.84 (2 H, m), 1.92 (4 H, br. s.), 2.07 - 2.29 (2 H, m), 2.41 (4 H, br. s.), 3.20 (3 H, s), 3.51 (4 H, s), 3.87 - 4.05 (4 H, m), 4.74 - 4.99 (1 H, m), 7.24 - 7.39 (5 H, m), 7.50 - 7.56 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.07 (1 H, d, J=6.8 Hz), 8.42 (1 H, d, J=8.8 Hz)









388		***	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide	Rt = 3.16 min (Analytical_1) MS(+ve) 606.4; MS(-ve) 604.5; 1H NMR (DMSO-d6) δ ppm: 1.07 - 1.18 (7 H, m), 1.24 - 1.49 (4 H, m), 1.58 - 1.71 (4 H, m), 1.71 - 1.84 (2 H, m), 1.83 - 2.03 (6 H, m), 2.23 (1 H, t, J=10.5 Hz), 2.43 - 2.51 (3 H, m), 3.17 - 3.28 (4 H, m), 3.42 (3 H, br. s.), 3.60 (4 H, br. s.), 3.64 (1 H, br. s.), 3.71 - 3.83 (1 H, m), 3.98 (3 H, s), 5.23 (1 H, t, J=8.3 Hz), 7.44 - 7.60 (2 H, m), 7.72 (1 H, s), 8.02 (1 H, s), 8.09 (1 H, d, J=7.8 Hz), 8.40 (1 H, d, J=8.3 Hz)
389		**	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide	Rt = 3.31 min (Analytical_1) MS(+ve) 606.4; MS(-ve) 604.5; 1H NMR (DMSO-d6) δ ppm: 1.13 (6 H, s), 1.39 - 1.60 (4 H, m), 1.60 - 1.71 (4 H, m), 1.71 - 1.85 (4 H, m), 1.85 - 2.01 (4 H, m), 2.07 - 2.22 (2 H, m), 2.42 - 2.48 (4 H, m), 3.22 (4 H, s), 3.64 (4 H, br. s.), 3.87 - 4.05 (4 H, m), 5.14 - 5.31 (1 H, m), 7.45 - 7.62 (2 H, m), 7.71 (1 H, s), 8.02 (1 H, s), 8.09 (1 H, d, J=7.3 Hz), 8.39 (1 H, d, J=8.3 Hz)
390		***	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide	Rt = 2.96 min (Analytical_1) MS(+ve) 604.4; MS(-ve) 602.5; 1H NMR (DMSO-d6) δ ppm: 0.72 (2 H, s), 0.95 (2 H, s), 1.24 - 1.49 (4 H, m), 1.49 - 1.60 (2 H, m), 1.60 - 1.69 (2 H, m), 1.69 - 1.82 (2 H, m), 1.82 - 2.03 (6 H, m), 2.14 (1 H, s), 2.17 - 2.32 (1 H, m), 3.22 (4 H, s), 3.53 (2 H, s), 3.61 (4 H, br. s.), 3.71 - 3.84 (1 H, m), 4.00 (3 H, s), 4.90 (1 H, quin, J=8.5 Hz), 7.43 - 7.61 (2 H, m), 7.73 (1 H, s), 8.04 (1 H, s), 8.10 (1 H, d, J=7.8 Hz), 8.44 (1 H, d, J=8.3 Hz)
391		**	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide	Rt = 3.10 min (Analytical_1) MS(+ve) 604.4; MS(-ve) 602.5; 1H NMR (DMSO-d6) δ ppm: 0.66 - 0.81 (2 H, m), 0.88 - 1.03 (2 H, m), 1.45 - 1.86 (12 H, m), 1.87 - 2.02 (4 H, m), 2.10 - 2.21 (1 H, m), 2.47 (4 H, br. s.), 3.22 (3 H, s), 3.52 (2 H, s), 3.65 (4 H, br. s.), 3.89 - 4.05 (4 H, m), 4.90 (1 H, quin, J=8.5 Hz), 7.50 - 7.63 (2 H, m), 7.73 (1 H, s), 8.03 (1 H, s), 8.10 (1 H, d, J=7.3 Hz), 8.43 (1 H, d, J=8.3 Hz)

TABLE 4: PLK activity data for compound [218] versus selected prior art compounds


Compound	Structure	PLK1	PLK2 IC50 / PLK1 IC50	PLK3 IC50 / PLK1 IC51
[218]		***	35	330
WO 2007/095188 [I-64]		***	17	67
WO 2007/095188 [I-76]		***	6	26
WO 2007/095188 [I-4]		***	5	64

*** denotes $<0.1 \mu\text{M}$ IC₅₀


215

TABLE 5: Solubility data for compound [254] and its HCl salt versus Example [I-76] of WO 07/095188 and its HCl salt

Compound	Description	Liquid phase	Concentration based on free base (mg/mL)	Forms stable solution	Forms gel?
[254]	free base	DMA / PEG400 / 10mM pH4 Tartrate buffer (1:3:6)	8	y	
Example [I-76] of WO 07/095188	free base	DMA / PEG400 / 10mM pH4 Tartrate buffer (1:3:6)	8	n	y
[254].HCl	HCl salt of [254]	40% Captisol / water	20	y	
[254].HCl	HCl salt of [254]	30% Captisol / water	20	y	
[254].HCl	HCl salt of [254]	20% HP β CD / water	20	y	
[I-76].HCl	HCl salt of [I-76]	40% Captisol / water	20	n	y
[I-76].HCl	HCl salt of [I-76]	30% Captisol / water	15	n	y
[I-76].HCl	HCl salt of [I-76]	20% HP β CD / water	10	n	y





Compound [I-76] of WO 07/095188
(= Example 246 of WO 08/003958)



Compound [254]

TABLE 6: Nephelometry maximum soluble concentration for compound [254] of the invention versus compound [I-253] of WO 07/095188

Invention versus compound [I-253] of WO 07/095188	Compound	PLK1 Ki (nM)	Nephelometry maximum soluble concentration (μM)
	Compound [254]	<10	>200
	Example [I-253] of WO 07/095188	<10	75
[254].HCl		<10	>200

217

TABLE 7: Compounds tested in pharmacokinetic studies; compound H = compound [371]; compound I = compound [378]; compounds A-G are for comparative purposes





















Structure	Compound
	A
	B
	C
	D
	E
	F
	G
	H
	I

TABLE 8: Comparator compounds tested in cell assays; letters correspond to labels in Figure 3.

Structure	Figure label
 Example 76 in WO 07/095188	A'
 B (cf. Example 59 in WO 07/095188)	B'
 Example 28 in WO 07/095188	C'
 D (cf. Example 28 in WO 07/095188)	D'
 E (cf. Example 360 in WO 08/003958)	E'
 F (cf. Example 360 in WO 08/003958)	F'
 G (cf. Example 394 in WO 08/003958)	G'
 H (cf. Example 394 in WO 08/003958)	H'
 Example 360 in WO 08/003958 (trans)	I'
 Example 360 in WO 08/003958 (cis)	J'

WE CLAIM:

1. A compound of formula VIII, or a pharmaceutically acceptable salt or ester thereof,



(VIII)

wherein:

X is NR⁷;

Y is O or N-(CH₂)_nR¹⁹;

n is 1, 2 or 3;

m is 1 or 2;

R¹ and R² are each independently H, alkyl or cycloalkyl;

R⁴ and R^{4'} are each independently H or alkyl; or

R⁴ and R^{4'} together form a spiro cycloalkyl group;


R¹⁹ is H, alkyl, aryl or a cycloalkyl group;

R⁶ is OR⁸ or halogen; and

R⁷ and R⁸ are each independently H or alkyl.

2. A compound according to claim 1 which is of formula VIIIa, or a pharmaceutically acceptable salt or ester thereof,

220



(VIIIa)

wherein:

X is NR⁷;

R¹ and R² are each independently H, alkyl or cycloalkyl;

R⁴ and R^{4'} are each independently H or alkyl; or

R⁴ and R^{4'} together form a spiro cycloalkyl group;

R¹⁹ is H, alkyl, aryl or a cycloalkyl group;

R⁶ is OR⁸ or halogen; and

R⁷ and R⁸ are each independently H or alkyl.

3. A compound according to claim 1 or claim 2 wherein R¹ is H or alkyl.
4. A compound according to claim 3 wherein R¹ is methyl.
5. A compound according to any one of claims 1 to 4 wherein R² is cycloalkyl.
6. A compound according to claim 5 wherein R² is cyclopentyl or cyclohexyl.
7. A compound according to any one of claims 1 to 6 wherein R⁷ is H or alkyl.
8. A compound according to claim 7 wherein R⁷ is H.
9. A compound according to any one of claims 1 to 8 wherein R⁴ and R^{4'} are each independently alkyl.

10. A compound according to claim 9 wherein R^4 and $R^{4'}$ are each independently methyl.
11. A compound according to any one of claims 1 to 8 wherein R^4 and $R^{4'}$ together form a spiro cycloalkyl group.
12. A compound according to any one of claims 1 to 8 wherein R^4 and $R^{4'}$ together form a spiro C_3 cycloalkyl group.
13. A compound according to any one of claims 1 to 12 wherein R^6 is OR^8 .
14. A compound according to claim 13 wherein R^6 is OMe.
15. A compound according to any one of claims 1 to 14 wherein R^{19} is cyclopropyl.
16. A compound according to claim 1 which is selected from the group consisting of:


Compound No.	Name
371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-

	(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
379	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
380	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
381	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide;
382	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide;
383	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
384	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
385	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
386	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
387	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-

	methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
388	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide;
389	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide;
390	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide;
391	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide;

a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable ester thereof.

17. A compound of formula VI, or a pharmaceutically acceptable salt or ester thereof,



wherein:

X is NR⁷;

R¹ and R² are each independently H, alkyl or cycloalkyl;

R⁴ and R^{4'} are each independently H or alkyl; or

R⁴ and R^{4'} together form a spiro cycloalkyl group;

R¹⁸ is H or alkyl, wherein said alkyl group is optionally substituted by R⁶;
 each R⁶ is independently OR⁸ or halogen; and
 R⁷ and R⁸ are each independently H or alkyl.

18. A compound according to claim 17 wherein R¹, R², R⁴, R^{4'}, R⁶, R⁷ and R⁸ are as defined in any one of claims 3 to 14.

19. A compound according to claim 17 which is selected from the group consisting of:

Compound No.	Name
254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide;
195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide;
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide;
373	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide;
374	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide;
375	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide;

a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable ester thereof.

20. A compound selected from the group consisting of:

Compound No.	Name
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;

a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable ester thereof.

21. A compound which is selected from the group consisting of:

Compound No.	Name
254	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide;
218	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;

195	4-(9-Cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)-benzamide;
221	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(4-methylpiperazin-1-yl)benzamide;
371	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
372	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
345	4-(9-Cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-[1-(tetrahydro-pyran-4-yl)-piperidin-4-yl]-benzamide;
373	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide;
374	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-fluoro-N-(4-methylpiperazin-1-yl)benzamide;
194	(±)-4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
186	(±)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
375	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(2-hydroxyethyl)piperazin-1-yl)-3-methoxybenzamide;
376	4-(9'-Cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-

	tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-morpholinobenzamide;
347	(R)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
348	(S)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-(quinuclidin-3-yl)benzamide;
377	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclobutane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
378	4-(9-cyclopentyl-5-methyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-N-((trans)-4-(4-(cyclopropylmethyl)piperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
379	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
380	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-ethylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
381	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-(cyclopropylmethyl)-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide;
382	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-(4-(4-ethyl-1,4-diazepan-1-yl)cyclohexyl)-3-methoxybenzamide;
383	N-(4-(4-benzyl-1,4-diazepan-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-

	methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
384	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((trans)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
385	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-N-((cis)-4-(4-methylpiperazin-1-yl)cyclohexyl)-3-methoxybenzamide;
386	N-((trans)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
387	N-((cis)-4-(4-benzylpiperazin-1-yl)cyclohexyl)-4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxybenzamide;
388	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide;
389	4-(9-cyclopentyl-5,7,7-trimethyl-6-oxo-6,7,8,9-tetrahydro-5H-pyrimido[4,5-b][1,4]diazepin-2-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide;
390	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((trans)-4-morpholinocyclohexyl)benzamide;
391	4-(9'-cyclopentyl-5'-methyl-6'-oxo-5',6',8',9'-tetrahydrospiro[cyclopropane-1,7'-pyrimido[4,5-b][1,4]diazepine]-2'-ylamino)-3-methoxy-N-((cis)-4-morpholinocyclohexyl)benzamide;

a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable ester thereof.

22. A pharmaceutical composition comprising a compound according to any one of claims 1 to 21 admixed with a pharmaceutically acceptable diluent, excipient or carrier.
23. Use of a compound according to any one of claims 1 to 21 in the manufacture of a medicament for treating a proliferative disorder.
24. Use of a compound according to any one of claims 1 to 21 for treating a proliferative disorder.
25. Use of the compound according to claim 23 or claim 24 wherein the proliferative disorder is cancer or leukemia.
26. Use of the compound according to claim 23 or claim 24 wherein the proliferative disorder is glomerulonephritis, rheumatoid arthritis, psoriasis or chronic obstructive pulmonary disorder.
27. Use of the compound according to any one of claims 23 to 26, wherein the compound is administrable in a therapeutically effective amount for treating a proliferative disorder in a subject.
28. Use of a therapeutically effective amount of a compound as defined in any one of claims 1 to 21 for treating a PLK-dependent disorder in a subject.
29. A pharmaceutical composition for treating a proliferative disorder, said composition comprising a therapeutically effective amount of a compound as defined in any one of claims 1 to 21.
30. A pharmaceutical composition for treating a PLK-dependent disorder, said composition comprising a therapeutically effective amount of a compound as defined in any one of claims 1 to 21.

31. A compound according to any one of claims 1 to 21 for treating a proliferative disorder.

FIGURE 1







FIGURE 2

FIGURE 3





CIII