



(51) International Patent Classification:

A61K 31/485 (2006.01) C07D 489/08 (2006.01)

(21) International Application Number:

PCT/US2017/032792

(22) International Filing Date:

16 May 2017 (16.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/338,636 19 May 2016 (19.05.2016) US

(71) Applicant: VIRGINIA COMMONWEALTH UNIVERSITY [US/US]; 800 East Leigh Street, Suite 3000, Richmond, VA 23298 (US).

(72) Inventors: ZHANG, Yan; 800 East Leigh Street, Suite 3000, Richmond, VA 23298 (US). SELLEY, Dana, E.; 800 East Leigh Street, Suite 3000, Richmond, VA 23298 (US). DEWEY, William; 800 East Leigh Street, Suite 3000, Richmond, VA 23298 (US). ABKARALI, Hamid; 800 East Leigh Street, Suite 3000, Richmond, VA 23298 (US).

(74) Agent: WHITHAM, Michael, E. et al.; 11491 Sunset Hills Road, Suite 340, Reston, VA 20190 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: POTENT AND SELECTIVE MU OPIOID RECEPTOR MODULATORS

(57) Abstract: Analogues of 6 α/β -naltrexamine (NAQ) are provided. The analogues are selective, reversible antagonists of the mu opioid receptor (MOR) that exhibit good blood brain barrier penetration. The compounds are used in the treatment of opioid addiction and other diseases and conditions, including for the treatment of pain.



POTENT AND SELECTIVE MU OPIOID RECEPTOR MODULATORS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims benefit of United States provisional patent application
5 62/388,636, filed May 19, 2016, the complete contents of which is hereby incorporated by
reference.

STATEMENT OF FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

This invention was made with government support under grant number DA024022
10 awarded by the National Institutes of Health/National Institutes on Drug Abuse. The United
States government has certain rights in the invention.

BACKGROUND OF THE INVENTION

Field of the Invention

15 The invention generally relates to Mu opioid receptor (MOR) modulators and
methods of their use. In particular, the invention provides potent non-peptidyl selective and
reversible MOR antagonists that are naltrexamine analogues, and methods of using them to
treat opioid addiction, pain and other conditions.

Background

20 It is estimated that between 26.4 million and 36 million people abuse opioids such as
heroin, morphine, and other prescription pain relievers worldwide, and this abuse affects the
health, social, and economic welfare of every demographic group. An alarming observation
is that opioid misuse is on the rise. For example, the number of unintentional overdose
25 deaths from opioid prescription analgesics has soared in the United States, more than
quadrupling since 1999. Therefore, opioid addiction has become a global epidemic and has
affected the economies of many countries. Effective drug addiction treatment has been
shown to reduce associated health and social cost far more than the cost of treatment itself.
Therefore, treating opioid addicts rather than imprisoning them would drive down the huge
30 cost associated with substance abuse.

Opioid antagonists such as naltrexone and naloxone target opioid receptors and
have been shown to curb drug craving and prevent relapse. However, severe side effects (e.g.
depression, dysphoria, pulmonary edema and cardiac arrhythmias) have been reported when

these drugs are used as long-term treatments for opioid addiction. The side effects are likely due to the lack of selectivity for the Mu opioid receptor (MOR) over other opioid receptors, namely the delta-opioid receptor (DOR) and the kappa-opioid receptor (KOR). For example, DOR activation has antidepressant and anxiolytic effects and also plays an important role in neuroprotection and cell proliferation, and KOR agonists and partial agonists have been shown to produce psychotomimetic and dysphoric effects. Therefore, an ideal agent to treat opioid addiction without the severe side effects would be a highly selective MOR antagonist.

Some selective MOR antagonist such as β -FNA and clocinnamox have been developed, but these agents bind irreversibly to MOR, which limits their use as drugs. Cyprodime is a reversible MOR antagonist that has been intensively studied. However, it has a lower affinity for MOR than naltrexone and naloxone and it has only moderate selectivity for MOR (KOR/MOR $\approx$ 10, DOR/MOR $\approx$ 39). The highly selective MOR antagonists that have been identified are conformationally constrained peptides such as D-Phe-Cys-Tyr-D-Trp-Orn-Thr-Pen-Thr-NH₂ (CTOP; SEQ ID NO: 1) and D-Phe-Cys-Tyr-D-Trp-Arg-Thr-Pen-Thr-NH₂ (CTAP; SEQ ID NO: 2). However, CTOP and CTAP have limited bioavailability and have poor blood-brain barrier penetration capacity, making them unsuitable as drug development candidates.

There is thus a pressing need to develop new opioid addiction treatment medications with fewer side effects. In particular, it would be beneficial to have available potent, highly selective, and reversible antagonists for MOR that exhibit high bioavailability and good blood-brain barrier penetration capacity.

SUMMARY OF THE INVENTION


Other features and advantages of the present invention are set forth in the description of invention that follows, and in part will be apparent from the description or may be learned by practice of the invention. The invention is realized and attained by the compositions and methods particularly pointed out in the written description and claims.

Provided herein are novel non-peptidyl, selective, and reversible MOR antagonists with high blood-brain barrier penetration capacity. Based on homology modeling studies of the three opioid receptor types and binding mode analyses of naltrexone in these models, two exemplary series of novel α/β -naltrexamine (NAQ) analogues were designed, synthesized and analyzed. The results of in vitro competition binding assays and in vivo functional studies show that the compounds are potent, selective and reversible MOR

antagonists. Significantly, the compounds have less efficacy at DOR than NAQ but they retain NAQ's low efficacy at KOR and high selectivity for MOR. The compounds thus have high therapeutic value in treating opioid addiction.

It is an object of this invention to provide compound with generic Formula I:


5



Formula I

wherein * indicates a chiral C that can be either an R configuration or an S configuration in terms of chirality; and R of Formula I =


10




and is bonded to Formula I at X₂, X₃, X₄, X₅, X₆ or X₇;

and wherein: X₁ = N, O or S; one or more of X₄, X₅, X₆ and X₇ are independently substituted by R' where R' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group, and one or both of X₂ and X₃ are independently substituted by R'' where R'' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group; and salts thereof; with the caveat that R is not:


15






In some aspects, the straight chain or branched alkyl group comprises from about 1 to about 20 carbon atoms. In other aspects, the straight chain or branched alkyl group is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl, octyl, nonyl, decyl, undecyl and dodecyl, and isomers (e.g. stereoisomers) thereof. In other aspects, the electron withdrawing group is selected from the group consisting of F, CN, NO₂, SO₃ and CF₃. In further aspects, the electron donating group is OCH₃, phenyl or substituted phenyl.

The invention also provides methods of treating opioid addiction in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of Formula I:




Formula I


wherein * indicates a chiral C that can be either an R configuration or an S configuration in terms of chirality; and R of Formula I =




and is bonded to Formula I at X₂, X₃, X₄, X₅, X₆ or X₇; and wherein X₁ = N, O or S; one or more of X₄, X₅, X₆ and X₇ are independently substituted by R' where R' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group, and one or both of X₂ and X₃ are independently substituted by R'' where R'' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group; or a physiologically acceptable salt thereof. In some aspects, the straight chain or branched alkyl group comprises from 1 to 20 carbon atoms. In other aspects, the straight chain or branched alkyl group is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl, octyl, nonyl, decyl, undecyl and dodecyl, and isomers thereof. In further aspects, the electron withdrawing group is selected from the group consisting of F, CN, NO₂, SO₃ and CF₃. In additional aspects, the electron donating group is OCH₃, phenyl or substituted phenyl. In some aspects, the compound is



or




The invention further provides methods of treating pain in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of Formula I:





Formula I

wherein * indicates a chiral C that can be either an R configuration or an S configuration in terms of chirality; and R of Formula I =




- and is bonded to Formula I at X₂, X₃, X₄, X₅, X₆ or X₇; and wherein X₁ = N, O or S; one or more of X₄, X₅, X₆ and X₇ are independently substituted by R' where R' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group, and one or both of X₂ and X₃ are independently substituted by R'' where R'' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group; or a physiologically acceptable salt thereof; with the caveat that R is not:





In some aspects, the straight chain or branched alkyl group comprises from 1 to 20 carbon atoms. In other aspects, the straight chain or branched alkyl group is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl, octyl, nonyl, decyl, undecyl and dodecyl, and isomers thereof. In further aspects, the electron withdrawing group is selected from the group consisting of F, CN, NO₂, SO₃ and CF₃. In additional aspects, the electron donating group is OCH₃, phenyl or substituted phenyl. In some aspects, the compound is



BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A-D. Warm-water tail immersion assay in mice (n=5) at 56 ± 0.1 °C. All tested compounds were administered subcutaneously (s.c.). Antinociceptive effects of (A) indole analogues of 6 α -naltrexamine (B) indole analogues of 6 β -naltrexamine. Compounds (10 mg/kg) were injected at Time 0. Twenty minutes after injection, tail flick was assessed with hot water. Blockage of the antinociceptive effect of morphine by (C) indole analogues of 6 α -naltrexamine (D) indole analogues of 6 β -naltrexamine. Tested compounds (10 mg/kg) were injected at Time 0. Five minutes later, morphine (10 mg/kg) was administered. Twenty minutes after morphine injection, tail flick was tested using hot water.

Figure 2A-C. Dose dependent studies on compound **6** (A), **17** (B), and **18** (C) for opioid antagonist effect. Compounds 6, 17, and 18 had AD₅₀ values of 2.39 (0.46-12.47), 7.30 (3.38-15.74), and 9.64 (3.17-29.29) mg/kg (95% CL) respectively.

Figure 3A-C. Compound **6** (s.c.) in opioid-withdrawal assays in chronic morphine-exposed mice (n=5): (A) Wet-dog shakes, (B) Escape jumps, and (C) Paw tremors. The first column in each figure represents placebo-pelleted mice while the second to the fourth represent morphine-pelleted mice. *** Indicates $P < 0.05$, compared to 1 mg/kg naltrexone (NTX, s.c.), whilst * indicates $P < 0.1$, compared to 1 mg/kg naltrexone (NTX, s.c.).


DETAILED DESCRIPTION

Described herein are non-peptidyl compounds that are potent MOR antagonists. The antagonists are highly selective and reversible, and thus in some aspects, the compounds are used to treat opioid addiction. In some aspects, certain of the compounds are also used to treat other conditions such as pain.

By “antagonist” we mean a substance that interferes with or inhibits the physiological action of another. In this case, the substance is one or more compounds as described herein, and the “another” is one or more of the MOR, the KOR and/or the DOR, especially the MOR.

As used herein “chiral center” or “stereocenter” refers to an atom in a compound at which the spatial arrangement of atoms or atomic groups bonded to the chiral atom is such that the compound is not superposable on its mirror image. Chiral centers are assigned a configuration of either R (rectus) or S (sinister) according to well defined Cahn–Ingold–Prelog rules.


The compounds disclosed herein have the general formula shown in Formula I:



Formula I


where * indicates a chiral C where the arrangement of atoms bonded to C can be either R or S, in terms of chirality;

- 5 and where the R substituent (variable group or component) of Formula I (i.e. the R that is bonded to C of NHC=O) is




wherein:

- 10 the substituent R is bonded to Formula I at X_2 , X_3 , X_4 , X_5 , X_6 or X_7 of R;
 X_1 of substituent R is nitrogen (N), oxygen (O) or sulfur (S);
 one or more of X_4 , X_5 , X_6 and X_7 are independently substituted by R' where $R' =$ a straight or branched, saturated or unsaturated carbon chain, an electron withdrawing group, or an electron donating group; and
 15 one or more of X_2 and X_3 are independently substituted by R'' where $R'' =$ a straight or branched, saturated or unsaturated carbon chain, an electron withdrawing group, or an electron donating group,
 with the caveat that, in the compound of Formula I, R is not:




as well as salts thereof, which may be physiologically acceptable or compatible salts. However, this caveat does not apply for the use of the compounds of the invention to treat e.g. drug addiction such as opioid addiction.

Accordingly, the invention also provides methods of treating opioid addiction by administering a therapeutically effective amount of at least one compound having the general formula shown in Formula I:



Formula I

where * indicates a chiral C where the arrangement of atoms bonded to C can be either R or S, in terms of chirality;
and where the R substituent (variable group or component) of Formula I (i.e. the R that is bonded to C of NHC=O) is



wherein:

the substituent R is bonded to Formula I at X_2 , X_3 , X_4 , X_5 , X_6 or X_7 of R;

X_1 of substituent R is nitrogen (N), oxygen (O) or sulfur (S);

one or more of X_4 , X_5 , X_6 and X_7 are independently substituted by R' where R' = a straight or branched, saturated or unsaturated carbon chain, an electron withdrawing group, or an electron donating group; and

one or more of X_2 and X_3 are independently substituted by R'' where R'' = a straight

or branched, saturated or unsaturated carbon chain, an electron withdrawing group, or an electron donating group.

For both R' and R'', suitable straight or branched, saturated or unsaturated carbon chains include but are not limited to those that contain from about 1 to about 20 carbon atoms. In some aspects, the carbon chain is saturated, i.e. the carbon chain is an alkyl chain. Examples of suitable alkyl groups include but are not limited to methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl, octyl, nonyl, decyl, undecyl and dodecyl, as well as branched isomers thereof, and substituted variants thereof. Alternatively, the branched or unbranched carbon chain may be unsaturated alkenes, e.g. containing one or more (e.g. 1, 2, 3, 4, or 5 or more) double bonds. For example, suitable lower alkenes containing 1 double bond include but are not limited to e.g. ethane, propene, butene, etc.

For both R' and R'', suitable electron withdrawing groups include but are not limited to, e.g. halogens such as F, CN, NO₂, SO₃, CF₃, NR₂, OR, NHCOR (where R is H or a lower alkyl with from about 1-5 carbon atoms such as methyl, ethyl, propyl, butyl, pentyl, etc., or an alkene equivalent).

For both R' and R'', suitable electron donating groups include but are not limited to, e.g. alkyl groups; alkoxy groups (e.g. OR where R = H or a lower alkyl); alcohol groups (methanol, ethanol, propanol, etc.); amino and substituted amino groups, etc. Exemplary functional groups include but are not limited to OCH₃, phenyl and substituted phenyl.

The compounds described herein generally have a binding affinity (K_i) for MOR in the range of from about 1.0 to about 0.05, and preferably in the range of from about 0.8 to about 0.1, e.g. about 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2 or 0.1. Generally, the compounds have a K_i for the KOR in the range of from about 0.15 to about 4.0, or about 0.2 to about 3.0, or about 0.25 to about 2.0 or even about 0.3 to about 1.0, including all intervening integers and decimals to two decimal places. Further, the compounds generally have a K_i for the DOR in the range of from about 50 to about 0.5, e.g. about 40 to about 0.75, or about 30 to about 1.0, or even from about 20 to about 5.0, including all intervening integers and decimals to two decimal places. Typically, the ratio of KOR/MOR affinity ranges from about 15.0 to about 0.1, e.g. from about 12.0 to about 0.5, for example from about 5.0 to about 1.0, such as about 5.0, 4.0, 3.0, 2.0 or 1.0, including all intervening integers and decimals to two decimal places (5.0, 4.99, 4.98, etc.). Similarly, the DOR/MOR affinity ratio is generally in the range of from about 200 to about 5, e.g. about 100 to about 10, such as about 100, 90, 80, 70, 60, 50, 40, 30, 20 or 10, including all intervening integers and decimals to two decimal places.

The compounds described herein are generally delivered (administered) as a pharmaceutical composition. Such pharmaceutical compositions generally comprise at least one of the disclosed compounds, i.e. one or more than one (a plurality) of different compounds (e.g. 2 or more such as 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) may be included in a single formulation. Accordingly, the present invention encompasses such formulations and compositions. The compositions generally include one or more substantially purified compounds as described herein, and a pharmacologically suitable (physiologically compatible) carrier, which may be aqueous or oil-based. In some aspects, such compositions are prepared as liquid solutions or suspensions, or as solid forms such as tablets, pills, powders and the like. Solid forms suitable for solution in, or suspension in, liquids prior to administration are also contemplated (e.g. lyophilized forms of the compounds), as are emulsified preparations. In some aspects, the liquid formulations are aqueous or oil-based suspensions or solutions. In some aspects, the active ingredients are mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredients. Suitable excipients include, for example, water, saline, dextrose, glycerol, ethanol and the like, or combinations thereof. In addition, the composition may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, preservatives, pharmaceutically acceptable salts and the like. In some aspects, various thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders and the like are added. The compositions of the present invention may contain any such additional ingredients so as to provide the composition in a form suitable for administration. The final amount of compound in the formulations varies, but is generally from about 1-99%. Still other suitable formulations for use in the present invention are found, for example in Remington's Pharmaceutical Sciences, 22nd ed. (2012; eds. Allen, Adejare Desselle and Felton).

Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins (such as human serum albumin), buffer substances (such as twin 80, phosphates, glycine, sorbic acid, or potassium sorbate), partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes (such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, or zinc salts), colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, methylcellulose, hydroxypropyl methylcellulose, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato

starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

"Pharmaceutically acceptable salts" refers to the relatively non-toxic, inorganic and organic acid addition salts, and base addition salts, of compounds of the present invention. These salts can be prepared in situ during the final isolation and purification of the compounds. In particular, acid addition salts can be prepared by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed. Exemplary acid addition salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactiobionate, sulfamates, malonates, salicylates, propionates, methylene-bis-.beta.-hydroxynaphthoates, gentisates, isethionates, di-p-toluoyltartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, p-toluenesulfonates, cyclohexylsulfamates and laurylsulfonate salts, and the like. See, for example S. M. Berge, et al., "Pharmaceutical Salts," J. Pharm. Sci., 66, 1-19 (1977) which is incorporated herein by reference. Base addition salts can also be prepared by separately reacting the purified compound in its acid form with a suitable organic or inorganic base and isolating the salt thus formed. Base addition salts include pharmaceutically acceptable metal and amine salts. Suitable metal salts include the sodium, potassium, calcium, barium, zinc, magnesium, and aluminum salts. The sodium and potassium salts are preferred. Suitable inorganic base addition salts are prepared from metal bases which include sodium hydride, sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminum hydroxide, lithium hydroxide, magnesium hydroxide, zinc hydroxide and the like. Suitable amine base addition salts are prepared from amines which have

sufficient basicity to form a stable salt, and preferably include those amines which are frequently used in medicinal chemistry because of their low toxicity and acceptability for medical use. ammonia, ethylenediamine, N-methyl-glucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chlorprocaine, diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, triethylamine, dibenzylamine, ephenamine, dehydroabietylamine, N-ethylpiperidine, benzylamine, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, ethylamine, basic amino acids, e.g., lysine and arginine, and dicyclohexylamine, and the like.

The compositions are administered in vivo by any suitable route including but not limited to: inoculation or injection (e.g. intravenous, intraperitoneal, intramuscular, subcutaneous, intra-aural, intra-ocular, intraarticular, intramammary, and the like); or by absorption through epithelial or mucocutaneous linings. Other suitable means include but are not limited to: inhalation (e.g. as a mist or spray) and orally (e.g. as a pill, capsule, liquid, etc.). In preferred embodiments, the mode of administration is oral. In addition, the compositions may be administered in conjunction with other treatment modalities.

The amount of a compound that is administered depends on several factors, e.g. the weight, gender, age, overall health, etc. of the recipient, and is best determined by a skilled medical practitioner such as a physician. However, the amount is generally in the range of from about 1-1000 mg/kg of body weight, e.g. from about 5-500 or about 10-250 or about 15-125 mg/kg of body weight, including all integers and fractional values within these ranges.

The compounds described herein generally are used to target diseases and conditions which can be prevented or treated by antagonizing the MOR. Exemplary conditions and symptoms thereof that are prevented and/or treated using the compounds described herein include but are not limited to: pain of any type, including neuropathic pain; substance abuse and addiction (e.g. opioid, cocaine, alcohol, amphetamine, methamphetamine, bath salts, etc.) in which case the present non-addictive compounds can be substituted for the addictive substance; Parkinson's disease; obesity; epilepsy; inflammation; gastrointestinal tract disorders such as constipation and irritable bowel syndrome (IBS); AIDs; various psychological disorders such as depression, schizophrenia, bipolar disorder, schizoaffective disorder, gambling addiction, etc.; and others. The addictions (dependencies) that may be treated include both recreational addictions which occur as the result of illegal substance

abuse, and addictions that result from the legal administration of drugs under the care of a medical professional, e.g. to treat pain.

However, varying degrees of affinity for the KOR and the DOR are exhibited by the compounds, and some may be suitable for preventing or treating diseases and conditions which are treatable by antagonizing the KOR or the DOR (alone or on combination with the MOR), or the KOR and the DOR (alone or on combination with the MOR).

Before exemplary embodiments of the present invention are described in greater detail, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

Where a range of values is provided, it is understood that each intervening value between the upper and lower limit of that range (to a tenth of the unit of the lower limit) is included in the range and encompassed within the invention, unless the context or description clearly dictates otherwise. In addition, smaller ranges between any two values in the range are encompassed, unless the context or description clearly indicates otherwise.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Representative illustrative methods and materials are herein described; methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention.

All publications and patents cited in this specification are herein incorporated by reference as if each individual publication or patent were specifically and individually indicated to be incorporated by reference, and are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual dates of public availability and may need to be independently confirmed.

It is noted that, as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural referents unless the context clearly dictates otherwise. It is further noted that the claims may be drafted to exclude any optional element. As such, this

statement is intended to serve as support for the recitation in the claims of such exclusive terminology as "solely," "only" and the like in connection with the recitation of claim elements, or use of a "negative" limitations, such as "wherein [a particular feature or element] is absent", or "except for [a particular feature or element]", or "wherein [a particular feature or element] is not present (included, etc.)..."

As will be apparent to those of skill in the art upon reading this disclosure, each of the individual embodiments described and illustrated herein has discrete components and features which may be readily separated from or combined with the features of any of the other several embodiments without departing from the scope or spirit of the present invention. Any recited method can be carried out in the order of events recited or in any other order which is logically possible.

EXAMPLES

EXAMPLE 1. Design, Synthesis and Pharmacological Characterization of Indole Analogues of 6 α/β -naltrexamine (NAQ)

INTRODUCTION

Naltrexamine (17-cyclopropylmethyl-3,14 β -dihydroxy- 4,5 α -epoxy-6 α (isoquinoline-3-carboxamido)morphinan or NAQ) is a MOR selective antagonist that acts in the central nervous system and has been shown to hold tremendous promise as a candidate or lead compound to treat opioid addiction (Schiff, P. L. Opium and Its Alkaloids. Am. J. Pharm. Educ. 2002, 66, 186–194; Katz, N. P. et al. Clin. J. Pain 2007, 23 (8), 648–660). However, NAQ's selectivity for MOR over DOR (146 fold) is lower than that of naltrexone (435 fold) (Meldrum, M. L. A Capsule History of Pain Management. JAMA 2003, 290 (18), 2470–2475). Even though NAQ has low efficacy at KOR, it has high efficacy and moderate potency at DOR. Studies have demonstrated that DOR activation is involved in the development of morphine dependence and also, deletion of the genes for DOR inhibits the development of morphine tolerance. Thus, new compounds having less efficacy at DOR in addition to retaining NAQ's low efficacy at KOR with high selectivity for MOR would have more therapeutic value in treating opioid addiction.

Through molecular modeling and mutagenesis studies, it was demonstrated that the selectivity of NAQ for MOR is as a result of its π - π stacking with W318 of MOR. To explore whether other heterocyclic ring systems might form π - π interactions with W318,

new compounds were designed in which the isoquinoline ring of NAQ was replaced with an indole ring. It was theorized that introducing other heterocyclic ring systems might yield compounds with improved pharmacologic (increased selectivity and reduced partial agonist effect) and drug-like properties (improved absorption, distribution, metabolism, and excretion properties) and might lead to the identification of new lead compounds.



MATERIALS AND METHODS

SYNTHESIS

Naltrexone was used as the starting material in the synthesis of both 6 α -naltrexamine (22) and 6 β -naltrexamine (25). In the synthesis of 6 α -naltrexamine HCl, (**Scheme 1, route A**) a reductive amination was conducted between naltrexone and benzyl amine. The Schiff base formed (22) was reduced and the benzyl group cleaved off using hydrogenation. However, dibenzyl amine was refluxed with naltrexone to synthesize 6 β -naltrexamine HCl (Courtwright, D. T.; Joseph, H.; DesJarlais, D. Addicts Who Survived; University of Tennessee Press: Knoxville, TN, 1989). The reduction of Schiff base of benzyl naltrexamine yielded the 6 α -isomer because the ring adopts a chair-like conformation. Therefore, hydride transfer occurs on the β -face leading to the formation of the 6 α -isomer. On the other hand, the Schiff base of dibenzyl naltrexamine adopts a boat conformation. Therefore, hydride transfer from NaCNBH₃, occurs exclusively on the more accessible α face, thereby leading to the 6 β isomer. Details of the syntheses and characterizations are given in Example 2 below.

Synthesis of indole analogues substituted at 6 position of naltrexamine


















6 α / β naltrexamine HCl were coupled to their respective indole carboxylic acids using EDCI coupling (**Scheme 2**). This coupling reaction also results in the formation of an ester at position 3. The ester group was then hydrolyzed since esters are more susceptible to hydrolysis than amides. The final compounds synthesized were converted to the HCl salt and the percentage yield obtained for each compound is shown in **Table 1**. The percentage yield obtained for the 6 α -naltrexamine derivatives ranged from 22-81% while that for the 6 β -naltrexamine derivatives ranged from 23-61%. Details of the syntheses and characterizations are given in Example 2 below.

Scheme 1. Synthesis of 6 α /6 β naltrexamine HCl

Scheme 2. Coupling of naltrexamine to indole carboxylic acids

The newly designed compounds were indole analogues of NAQ and are depicted in **Table 1**. A total of 18 exemplary compounds were synthesized (additional examples can be made by this synthetic route). In Table 1, the compound IDs for the 6 α -naltrexamine derivatives are in black while the compound IDs for the 6 β -naltrexamine derivatives are in grey. Percentages are percent yields. The final compounds were characterized by ^1H NMR, ^{13}C NMR, HRMS and IR and the purity determined by HPLC.

Table 1. Indole analogues of NAQ

 6α-naltrexamine analogues  6β-naltrexamine analogues	1		81%	6		22%
	10		32%	15		56%
	2		29%	7		80%
	11		23%	16		49%
	3		80%	8		55%
	12		31%	17		51%
	4		50%	9		34%
	13		61%	18		39%
	5		75%			
	14		29%			

PHARMACOLOGICAL CHARACTERIZATION

5 Pharmacological assays were conducted to determine the binding affinity, selectivity and functional activity of the synthesized compounds.

Cell culture

Chinese hamster ovary (CHO) cells expressing MOR, KOR and DOR were cultured for use in this assay. The culture media used for MOR CHO cells contained 500 mL
 10 DMEM/F12 media, 10% FBS, 1% penicillin/streptomycin, and 2.75 mL G418. The DOR and KOR CHO cells were cultured in the same media except that 5% FBS was used instead of 10%. The cells are allowed to grow until they are 80-95% confluent after which they are split. The cells are continuously cultured and split until there were sufficient cells for the studies.

15 *Cell membrane harvesting*

Once the cells reached confluence, cells were harvested, centrifuged at 1000 x g for 10 minutes, and suspended in membrane buffer (50 mM Tris, 3mM MgCl₂, and 1 mM EGTA, pH 7.4). The cells were then homogenized, centrifuged at 50000 x g for 10 minutes, and re-homogenized in membrane buffer. A Bradford assay was conducted to determine membrane protein concentrations, and membrane protein preps (3 µg/aliquot) were stored at -80 °C.

Competition binding assay

The competition binding assay was conducted to determine the affinity and selectivity of the synthesized compounds to MOR over KOR and DOR. The K_d and B_{max} values for [³H]NLX at MOR and [³H]DPN at KOR had been determined previously. The K_d and B_{max} values for [³H]DPN at DOR were determined using varying concentrations of [³H]DPN and fixed a concentration of 30 µg DOR membrane protein and 5 µM SNC80. [³H]NLX was used to label MOR whilst [³H]DPN was used to label both DOR and KOR. The potency of the new compounds in displacing the specific binding of the radioligand was determined by linear regression analysis of Hill plots. Specific (i.e., opioid receptor-related) binding at MOR, KOR and DOR was determined as the difference in binding obtained in the absence and presence of 5 µM naltrexone, U50,488 and SNC80, respectively. The IC₅₀ values were determined and converted to K_i values using the Cheng–Prusoff equation (Rosenblum, A. et al.. Exp Clin Psychopharmacol. 2008, 16 (5), 405–416).

[³⁵S]GTPγS functional assay

The [³⁵S]GTPγS functional assay was used to determine whether the synthesized compounds are agonists. The agonist's activity of the compounds was measured relative to 3 µM DAMGO at MOR. In this assay, upon receptor activation due to agonist binding, [³⁵S]GTPγS binds to the Gα subunit. Thus, [³⁵S]GTPγS replaces endogenous GTP and since the γ-thiophosphate bond is resistant to hydrolysis by GTPase, [³⁵S]GTPγS labelled Gα subunits accumulate and can be quantified by measuring the amount of incorporated [³⁵S].

Briefly, after thawing, membrane protein preparations were suspended in membrane (TME) buffer (50 mM Tris, 3mM MgCl₂, 0.2 mM EGTA, and 100 mM NaCl adjusted to pH 7.7), homogenized, and centrifuged at 50000 x g for 10 mins and then re-homogenized. The Bradford assay was then conducted to determine the concentration of the membrane protein using BSA as the standard. Suitable dilutions of membrane protein, [³⁵S]GTPγS, cold GTPγS, DAMGO and GDP were made in TME. The desired amount of the protein per tube

was 10 µg. All the test-tubes were vortexed and incubated in the shaking water bath at 30 °C for 90 minutes, after which bound radioligand was filtered and rinsed three times with cold Tris buffer. The filtered bound radioligand was then transferred into scintillation tubes with 4 mL of scintillation fluid and radioligand was quantified. However, counting was delayed for 9 hrs to allow the radioactivity to enter the solution. Percent DAMGO stimulation was defined as (net stimulated binding of test compound/ net stimulated binding of DAMGO) x 100%. The normalized data were subjected to nonlinear regression analysis to determine EC₅₀ and E_{max} values using Prism 6.0 software (GraphPad Software, San Diego, CA). The assay was performed in triplicate and repeated at least three times.

IN VIVO STUDIES

For the in vivo studies, male Swiss Webster mice were used for both the warm-water immersion and opioid withdrawal assays. The mice weighed 23-35 g and were housed five to a cage in animal care quarters maintained at 22 °C on a 12 hour light-dark cycle with food and water available ad libitum. The mice were transferred to the test room and the tests were then conducted 18 hours later. Protocols and procedures were approved by the Institutional Animal Care and Use Committee at Virginia Commonwealth University Medical Center and complied with the recommendations of the International Association for the Study of Pain.

A 75 mg morphine pellet was implanted in to the base of the neck of the mice and the mice were then given time to recover in their home cages before the test was conducted.

The mice were then allowed for 30 min habituation to an open-topped, square, clear Plexiglas observation chamber (26 x 26 x 26 cm³) with lines partitioning the bottom into quadrants before they were given antagonist. All drugs and test compounds were administered subcutaneously (s.c.). Withdrawal was precipitated at 72 h from pellet implantation with naltrexone (1.0 mg/kg, s.c.), and the test compounds at indicated doses. Withdrawal commenced within 3 min after antagonist administration. Escape jumps, paw tremors and wet dog shakes were quantified by counting their occurrences over 20 min for each mouse using five mice per drug. The data is given as the mean ± SEM. One-way ANOVA followed by the post hoc Dunnett test were performed to assess significance using the Prism 6.0 software.

Warm-water immersion assay


Each indole analogue of naltrexamine was tested for its ability to produce antinociception and/or to antagonize the antinociceptive effects of morphine in mice using










the warm-water tail immersion assay described by Coderre and Rollman. The a water bath temperature was maintained at 56 ± 0.1 °C. The baseline latency (control) was determined before injecting the compounds into the mice. The average baseline latency obtained for this experiment was 3.0 ± 0.1 s and only mice with a baseline latency of 2 to 4 s were used. To test for agonism, the tail immersion was conducted 20 min (time that morphine effect starts to peak) after injecting the indole analogues of $6\alpha/\beta$ -naltrexamine. To prevent tissue damage, a 10 s maximum cut off time was imposed. Antinociceptive response was calculated as the percentage maximum possible effect (%MPE), where $\%MPE = [(test - control) / (10 - control)] \times 100$. In the antagonism study, the $6\alpha/\beta$ -naltrexamine derivatives were given 5 min before morphine. The tail immersion test was conducted 20 min after giving morphine. %MPE was calculated for each mouse using at least five mice per drug. AD_{50} values were calculated using the least-squares linear regression analysis followed by calculation of 95% confidence interval by Bliss method. A higher % MPE indicates a stronger antinociception effect by the ligand.

RESULTS

The K_i values obtained for the indole analogues of 6α -naltrexamine at MOR, KOR and DOR and the selectivity for MOR over KOR and DOR are shown in **Table 2** whilst those for the indole analogues of 6β -naltrexamine are shown in **Table 3**.









Table 2. Binding affinity and selectivity of indole derivatives of 6α -naltrexamine



						
		K_i (nM) $\pm$ SEM			Selectivity Ratio	
Compd	R	μ	κ	δ	κ/μ	δ/μ
NTX ^a		0.33 ± 0.02	1.44 ± 0.11	143.5 ± 13.7	4.4	435

NAQ ^b		1.11 ± 0.07	13.3 ± 1.1	161.9 ± 15.0	12	146
1		0.363 ± 0.027	0.926 ± 0.125	14.244 ± 2.783	2.6	39.2
2		0.288 ± 0.042	0.984 ± 0.125	10.538 ± 2.888	3.4	37.0
3		0.264 ± 0.035	1.488 ± 0.349	9.260 ± 2.826	5.6	35.1
4		0.763 ± 0.112	3.447 ± 0.994	26.715 ± 7.722	4.5	35.0
5		0.434 ± 0.048	1.633 ± 0.277	12.775 ± 3.360	3.8	29.4
6		0.229 ± 0.015	1.693 ± 0.350	10.918 ± 2.938	7.4	47.7
7		0.837 ± 0.124	3.135 ± 0.449	9.169 ± 2.670	3.7	11.0
8		0.437 ± 0.039	1.362 ± 0.178	31.231 ± 8.121	3.1	71.5
9		0.293 ± 0.038	0.149 ± 0.033	6.731 ± 0.396	0.5	23.0

^a The K_i values of NTX and ^b NAQ were obtained from previous experiments.

Table 3. Binding affinity and selectivity of indole derivatives of 6 β -naltrexamine

						
Compd	R	K_i (nM) $\pm$ SEM			Selectivity Ratio	
		μ	κ	δ	κ/μ	δ/μ
NTX ^a		0.33 $\pm$ 0.02	1.44 $\pm$ 0.11	143.5 $\pm$ 13.7	4.4	435
NAQ ^b		1.11 $\pm$ 0.07	13.3 $\pm$ 1.1	161.9 $\pm$ 15.0	12	146
10		0.287 $\pm$ 0.037	0.178 $\pm$ 0.002	1.544 $\pm$ 0.473	0.6	5.4
11		0.192 $\pm$ 0.016	0.160 $\pm$ 0.005	7.169 $\pm$ 1.232	0.83	37.3
12		0.235 $\pm$ 0.031	1.936 $\pm$ 0.299	45.878 $\pm$ 16.292	8.2	195.2
13		0.259 $\pm$ 0.037	0.512 $\pm$ 0.038	17.673 $\pm$ 4.780	2.0	68.2
14		0.250 $\pm$ 0.044	0.166 $\pm$ 0.012	6.231 $\pm$ 0.825	0.7	25
15		0.188 $\pm$ 0.011	0.520 $\pm$ 0.094	30.449 $\pm$ 9.494	2.8	162
16		0.241 $\pm$ 0.029	0.947 $\pm$ 0.114	37.019 $\pm$ 1.209	3.9	153.6

17		0.167 ± 0.012	0.387 ± 0.045	49.010 ± 4.717	2.3	293.5
18		0.203 ± 0.027	0.175 ± 0.037	25.299 ± 4.344	0.86	124.6

^a The K_i values of NTX and ^b NAQ were obtained from previous experiments.

MOR binding for indole analogues of 6 α -naltrexamine

From the results obtained, it was observed that the position of the substitution on the indole ring did not affect the binding affinities significantly. Among the indole analogues of 6 α -naltrexamine, compound **4** had the least binding affinity which was 3.3 fold worse than the compound **6** which had the best binding affinity at MOR. Also, increasing alkyl chain at position 3 did not affect binding affinity at MOR significantly. Compound **7** with the one carbon linker had the least binding affinity which was approximately 3 fold worse than compound **3** which has its carbonyl group directly attached to position 3 and compound **9** with the three carbon linker. The binding affinities of the compounds at MOR were similar to the binding affinities of NTX at MOR.

MOR binding for indole analogues of 6 β -naltrexamine

The binding affinities at MOR for the indole analogues of 6 β -naltrexamine were higher than their respective 6 α -naltrexamine analogues. The K_i of all the indole analogues of 6 β -naltrexamine at MOR was approximately 0.2nM. Thus, neither the substitution position nor the length of the alkyl group at position 3 had any significant effect on binding affinity at MOR.

KOR binding for indole analogues of 6 α -naltrexamine

The binding affinities for the indole analogues of 6 α -naltrexamine was lower at KOR compared to MOR. Compound **4** with substitution at position 5 had the least binding to KOR. The binding affinities at KOR for the indole analogues of 6 α -naltrexamine were similar to that of naltrexone at KOR. The position of substitution on the indole ring did not have a significant effect on the binding of the compounds at KOR. Also, increasing alkyl chain length at position 3 on the indole ring did not have any significant effect on KOR

binding; however, compound **9** which had a three carbon linker had a greater binding affinity at KOR than MOR. Compound **9** was the only indole derivatives of 6 α -naltrexamine that had a greater affinity at KOR than MOR.

KOR binding for indole analogues of 6 β -naltrexamine

5 Again, the binding affinities at KOR for the indole analogues of 6 β -naltrexamine were higher than their respective 6 α -naltrexamine analogues. Among the indole analogues of 6 β -naltrexamine, compound **12** (with substitution at position 4) had the least affinity for KOR. Also, increasing the alkyl chain at position 3 did not significantly alter the binding affinity. Most of the indole analogues of 6 β -naltrexamine might either be dual MOR and
10 KOR agonist or antagonists.

DOR binding for indole analogues of 6 α -naltrexamine

The indole analogues of 6 α -naltrexamine had the least binding affinities at DOR compared to MOR and KOR. Compound **8** with substitution at position 3 and a two carbon linker had the least binding to DOR. The position of substitution on the indole ring did not
15 have a significant effect on the binding of the compounds at DOR. However, it was observed that the analogue that was substituted at position 4 (which had the best binding affinity compared to substitutions at other positions on the indole ring) had a binding affinity that was 3 fold better than compounds substituted at position 5 (which had the worst binding affinity compared to substitutions at other positions on the indole ring). Interestingly, it was
20 observed that when the alkyl chain at position 3 was only a 1 carbon chain, there was little change in the binding affinity. However, upon increasing the chain length to 2 carbon atoms, the binding affinity decreased by three fold. On the other hand, when the chain length was increased to 3 carbon atoms, the binding affinity improved.

DOR binding for indole analogues of 6 β -naltrexamine

25 Among the indole analogues of 6 β -naltrexamine, compound **10** (with substitution at position 1) had the best binding affinity for DOR whilst compound **17** (with 2 carbon chain linker substituted at position 3) had the least binding affinity for DOR. It was also observed that, increasing the alkyl chain length at position 3 resulted in a reduction in DOR binding.

Selectivity for MOR over KOR and DOR of synthesized compounds

30 Compound **6** and compound **12** had the best selectivity for MOR over KOR whilst compound **17** had the greatest selectivity for MOR over DOR. However, it's worth noting that among the 6 α -naltrexamine analogues, compound **6** and compound **8** had good


selectivity for MOR over DOR. Thus, compound **6** had a good selectivity when considering selectivity for MOR over both KOR and DOR.










Results of [³⁵S]GTPγS functional assay

The results from the [³⁵S]GTPγS functional assay are shown in **Table 4** (indole analogues of 6α-naltrexamine) and **Table 5** (indole analogues of 6β-naltrexamine). As can be seen, the position of substitution on the indole ring did not significantly affect potency and efficacy. All the compounds had partial agonist effect at MOR. Compound **6** (substitution at position 7) had the highest potency but the lowest efficacy at MOR. Compound **6** was more potent than NAQ but had similar efficacy as NAQ. Compound **1** had the highest efficacy at MOR. Increasing the alkyl chain at position 3 did not have a significant effect on potency and efficacy. Compound **9** with the three carbon linker was the most potent and efficacious.

Compounds **10**, **11** and **14** were the most potent compounds and these compounds were the most efficacious having E_{max} values greater than 70%. In fact, compound **10** with substitution at position 2 had the greatest efficacy among both the indole analogues of 6 α and β naltrexamine, with an E_{max} value of 92%. From the results obtained it was observed that substitution at position 2, 3 and 6 had a significant effect on potency and efficacy. It was also observed that increasing the alkyl chain length at position 3 did not significantly affect potency. However, as the alkyl chain increased, the efficacy decreased correspondingly.








Table 4. MOR [³⁵S]GTPγS binding functional assay results of indole derivatives of 6α-naltrexamine




			
Compound	R	EC ₅₀ (nM) ± SEM	E _{max} (% DAMGO) ± SEM
NTX^a		0.16 ± 0.04	5.4 ± 0.8
NAQ^a		3.3 ± 0.4	20.8 ± 1.2

1		3.29 ± 1.32	52.0 ± 13.5
2		1.11 ± 0.09	35.4 ± 4.8
3		4.99 ± 2.19	28.9 ± 1
4		1.55 ± 0.08	32.2 ± 1.3
5		5.41 ± 1.52	27.0 ± 0.31
6		1.0 ± 0.1	20.95 ± 4.43
7		4.92 ± 0.42	33.88 ± 1.43
8		1.87 ± 0.42	41.10 ± 6.54
9		0.62 ± 0.13	48.36 ± 5.17

^a The EC_{50} and E_{max} (%DAMGO) values of NTX and ^b NAQ were obtained from previous experiments.

Table 5. MOR [35 S]GTP γ S binding functional assay results of indole derivatives of 6 β -naltrexamine

			
Compound	R	EC ₅₀ (nM) ± SEM	E _{max} (% DAMGO) ± SEM
NTX ^a		0.16 ± 0.04	5.4 ± 0.8
NAQ ^a		3.3 ± 0.4	20.8 ± 1.2
10		0.21 ± 0.01	92.4 ± 2.8
11		0.30 ± 0.03	70.45 ± 0.9
12		1.97 ± 0.12	15.21 ± 3.13
13		2.28 ± 0.19	20.15 ± 4.37
14		0.19 ± 0.02	71.8 ± 4.4
15		0.75 ± 0.20	32.0 ± 4.6

16		0.56 ± 0.18	50.17 ± 7.75
17		0.39 ± 0.11	40.43 ± 5.28
18		1.04 ± 0.17	36.29 ± 4.11

^a The EC_{50} and E_{max} (%DAMGO) values of NTX and ^b NAQ were obtained from previous experiments.

Results of warm-water immersion assay

5 **Figure 1A** shows the antinociceptive effects of the indole analogues of 6α -naltrexamine, while **Figure 1B** shows the antinociceptive effects of the indole analogues of 6β -naltrexamine. Not surprisingly, compound **10**, which was identified as a potent and efficacious MOR agonist, was also identified as an opioid agonist in the warm-water immersion assay. Compounds **15** and **16** were identified as partial opioid agonists. Figure

10 **1C and D** show the antinociceptive effect of morphine at 10 mg/kg in the presence of each indole analogue of 6α -naltrexamine (**Figure 1C**) and 6β -naltrexamine (**Figure 1D**) at 10 mg/kg. Compounds **6**, **15**, **17**, and **18** effectively blocked morphine-mediated antinociception. Not surprisingly, compound **6**, which had a high potency and the least efficacy in the [35 S]GTP γ S functional assay, was identified as the most potent opioid

15 antagonist in the warm-water immersion assay. The opioid antagonist effects of compound **6**, **17**, and **18** were dose dependent and their AD_{50} values were 2.39 (0.46-12.47), 7.30 (3.38-15.74), and 9.64 (3.17-29.29) mg/kg (95% CL) respectively (**Figure 2A-C**). The slight increase in the %MPE for compound **6** at 8 mg/kg in the dose response study could be due to the partial agonist effects of compound **6** or could be due to variability in the mice.


20 Results of Opioid Withdrawal Assay

Compound **6** had the most promising pharmacological characteristics in the invitro

studies conducted compared to the other compounds synthesized. Compound **6** was also identified as the most potent opioid antagonist in the warm-water immersion assay. Therefore further in vivo studies were conducted on compound **6** using morphine-pelleted mice to determine if compound **6** had opioid withdrawal. The results showed that compound **6** itself did not induce any significant withdrawal symptoms in placebo-pelleted mice at 50 mg/kg (**Figure 3A, B and C**, first columns). On the other hand, a 1mg/kg dose of naltrexone induced significant withdrawal signs immediately after injection in morphine-pelleted mice and gave 18.6 ± 1.2 times wet-dog shakes, 43.4 ± 15.7 paw tremors and 39.0 ± 14.2 times escape jumps (**Figure 3A, B and C**, second columns). Secondly, compound **6** produced significantly less wet-dog shakes than naltrexone in morphine-pelleted mice even at dose of 50 mg/kg (**Figure 3A**, column 4). Again, compound **6** produced significantly less paw tremors than naltrexone in morphine pelleted mice even at a high dose of 50 mg/kg (**Figure 3B**, column 4). On the other hand, compound **6** at dose of 1mg/kg produced significantly less escape jumps than naltrexone at a dose of 1mg/kg in morphine pelleted mice (**Figure 3C**, column 3). However, compound **6** produced more escape jumps at 50mg/kg in morphine-pelleted mice than naltrexone at 1mg/kg (**Figure 3C**, column 5), but these results were not significantly different. Thus, compound **6** had less opioid withdrawal effects at a dose 26 times its AD_{50} value and produced less significant withdrawal symptoms than naltrexone overall. It was also observed that both compound **6** and naltrexone at 1mg/kg produced diarrhea in morphine pelleted mice but not in the placebo pelleted mice.

EXAMPLE 2. DETAILED CHEMICAL SYNTHESSES


Synthesis of 6 α -benzyl naltrexamine



A benzene solution (150 mL) in a three-necked flask containing naltrexone base (**19**), benzyl amine (1.2 eq.) and a trace of p-toluene sulfonic acid was refluxed for 10 hours using a Dean-Stark trap for azeotropic removal of water. The mixture was then concentrated (30 mL) and NaBH₄, a-solute ethanol and 4 Å molecular sieves were added to the reaction mixture and stirred under N₂ overnight. When the reaction was completed, the solvent was


removed using a rotavapor and the product purified using column chromatography with a Dichloromethane:Methanol:Ammonium hydroxide (40: 1: 1 %) mobile phase system to obtain 6 α -benzyl naltrexamine (**21**) in 88 % yield (Courtwright, D. T.; Joseph, H.; DesJarlais, D. Addicts Who Survived; University of Tennessee Press: Knoxville, TN, 1989).

5 Synthesis of 6 α -naltrexamine HCl




6 α -benzyl naltrexamine (**21**) was dissolved in 35-60 mL of methanol and then concentrated HCl was added to obtain a pH of 2. 10 % Pd/C (approximately 10% of the starting material) was added and hydrogenation was carried out at 60 psi for 6 days using a Parr hydrogenator. The Pd/C was changed every three days. When the reaction was completed, the Pd/C was filtered using celite to obtain pure 6 α -naltrexamine HCl (**22**) with a yield of 76 % (ibid).

Synthesis of 6 β -dibenzyl naltrexamine




A toluene solution (150 mL) in a three-necked flask containing naltrexone base (**19**), benzoic acid (1.3 eq.), a trace of p-toluene sulfonic acid was stirred for 30 minutes at room temperature. Dibenzyl amine (1.3 eq.) was then added and the mixture refluxed with a Dean-Stark apparatus for 24 hours under N₂. The mixture was then concentrated to 30 mL, cooled and then 50 mL anhydrous ethanol and NaCNBH₃ (0.8 eq) were added and the reaction mixture was stirred under N₂ overnight. The mixture was then filtered with celite and the solvent was removed using a rotavapor apparatus. The solid obtained was refluxed with methanol for 1 hour and then cooled and filtered. The residue obtained was washed with cold methanol to obtain 6 β -dibenzyl naltrexamine (**24**) in 71 % yield (ibid).

Synthesis of 6 β -naltrexamine HCl



Dibenzyl naltrexamine (**24**) was dissolved in 35-60 mL of methanol and then concentrated HCl was added to obtain a pH of 2. 10% Pd/C (approximately 10 % of the starting material) was added and hydrogenation was carried out at 60 psi for 6 days using a Parr hydrogenator. The Pd/C was changed every three days. When the reaction was completed, the Pd/C was filtered using celite to obtain pure 6β-naltrexamine HCl (**25**) with a yield of 82 % (ibid).


General method of coupling of naltrexamine to carboxylic acid derivatives



The carboxylic acid derivatives (3 eq.), hydrobenzotriazole (HoBt) (3eq.), 4 Å molecular sieves triethylamine (9 eq.), *N*-(3-Dimethylaminopropyl)-*N*'-ethylcarbodiimide hydrochloride (EDCI) (3 eq.), and DMF (2 mL) were added into a three-necked flask on an ice-water bath and stirred for 15 minutes. Naltrexamine (1 eq.) suspended in 2 mL DMF was added dropwise and then stirred on the water bath for 24 hours. When the reaction was completed, the reaction mixture was filtered with celite and the filtrate concentrated to remove DMF. MeOH (7 mL) and K₂CO₃ (3 eq.) were added to the concentrate and stirred at ambient temperature. When hydrolysis of the ester at position 3 was complete, the mixture was filtered through celite again and concentrated to remove MeOH. The mixture was then purified using column chromatography with dichloromethane: MeOH (20:1) and NH₄OH to give the free base (Meldrum, M. L. A Capsule History of Pain Management. JAMA 2003, 290 (18), 2470–2475).

Characterization of compounds


17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6α-(indole-2-carboxamido)morphinan (1)



The title compound was prepared by following the general procedure in 81% yield.

¹H NMR (400 MHz, DMSO-*d*₆) δ 11.65 (s, 1H, Exchangeable), 9.25 (s, 1H, Exchangeable), 8.87 (s, 1H, Exchangeable), 8.09 (d, *J* = 7.76 Hz, 1H, Exchangeable), 7.62 (d, *J* = 7.96 Hz, 1H), 7.45 (d, *J* = 8.2 Hz, 1H), 7.23 (dd, *J* = 7.96 Hz, 1.44 Hz, 1H), 7.18 (d, *J* = 8.2 Hz, 1H) 7.05 (t, *J* = 7.88 Hz, 1H), 6.72 (d, *J* = 8.08 Hz, 1H), 6.59 (d, *J* = 8.16 Hz, 1H), 6.34 (s, 1H, Exchangeable), 4.79 (d, *J* = 3.8 Hz, 1H) 4.62 (m, 1H), 3.93 (d, *J* = 7 Hz, 1H), 3.40-3.26 (m, 2H), 3.12-3.05 (m, 2H), 2.96 (m, 1H), 2.73 (m, 1H), 1.92 (m, 1H) 1.66 (d, *J* = 10.56, 1 H) 1.56-1.45 (m, 2H), 1.2 (m, 1H) 1.09 (m, 1H), 0.71-0.61 (m, 2H), 0.51-0.40 (m, 2H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 160.45, 146.16, 138.79, 136.50, 131.46, 128.74, 126.95, 123.40, 122.16, 121.44, 119.73, 119.15, 118.55, 112.26, 103.77, 87.41, 69.41, 61.07, 59.70, 57.08, 45.57, 45.28, 30.23, 29.27, 23.57, 19.56, 5.70, 5.16, 2.60. HRMS *m/z* found 486.2414 [M + H]⁺. Calculated 485.2315. IR (ATR, cm⁻¹) $\nu_{\max}$ 3146, 1633, 1544.97, 1505, 1457, 1341, 1312, 1235, 1173, 1116, 989, 810, 746. % Purity 97.69%.

17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6α-(indole-3-carboxamido)morphinan (2)




The title compound was prepared by following the general procedure in 29% yield.

¹H NMR (400 MHz, DMSO-*d*₆) δ 11.65 (s, 1H, exchangeable), 9.21 (s, 1H, exchangeable), 8.85 (s, 1H, exchangeable), 8.14 (d, *J* = 2.88, 1H, exchangeable), 8.11 (d, *J* = 7.56, 1H), 7.45 (t, *J* = 6.68, 2H), 7.18-7.09 (m, 2H), 6.71 (d, *J* = 8.08, 1 H), 6.58 (d, *J* = 8.12, 1H), 6.27 (s, 1H, exchangeable), 4.80 (d, *J* = 3.8, 1H), 4.66-4.60 (m, 1H), 3.91 (d, *J* = 6.68, 1H), 3.12-3.05 (m, 2H), 2.95 (m, 1H), 2.74 (m, 1H), 1.90 (m, 1H), 1.66 (d, *J* = 10.64, 1H), 1.56-1.44 (m, 2H), 1.11-1.06 (m, 2H), 0.71-0.61 (m, 2H), 0.51-0.40 (m, 2H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ

164.32, 145.79, 138.41, 135.91, 128.82, 128.09, 125.73, 122.18, 122.07, 120.62, 119.31, 118.06, 111.90, 110.04, 87.85, 69.23, 60.97, 56.99, 45.13, 44.82, 30.12, 29.16, 23.36, 19.59, 5.55, 5.19, 2.41, 0.012. HRMS m/z found 486.2357 $[M + H]^+$. Calculated 585.2315. IR (ATR, cm^{-1}) ν_{max} 3269, 1667, 1540, 1504, 1459, 1318, 1174, 1118.


5 **17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(indole-4-carboxamido)morphinan (3)**



The title compound was prepared by following the general procedure in 81% yield.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.37 (s, 1H, exchangeable), 9.18 (s, 1H, exchangeable)
 10 8.92 (s, 1H, exchangeable), 7.65 (d, $J=8$, 1H, exchangeable), 7.57 (d, $J=8$, 1H), 7.46-7.44 (m, 2H), 7.16 (t, $J=7.72$, 1H), 6.85 (s, 1H), 6.71 (d, $J=8.04$, 1H), 6.58 (d, $J=8$, 1H), 6.33 (s, 1H, exchangeable), 4.83 (d, $J=3.24$, 1H), 4.66 (m, 1H), 3.94 (d, $J=6.08$, 1H), 3.32 (s, 1H), 3.10 (m, 2H), 3.05 (m, 2H), 2.74 (m, 2H), 1.94 (m, 1H), 1.67 (d, $J=12.92$, 1H), 1.58 (m, 1H), 1.47 (m, 1H), 1.09 (m, 2H), 0.69 (m, 1H), 0.62 (m, 1H), 0.49 (m, 1H), 0.41 (m, 1H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 167, 145.62, 137.71, 136.30, 130.81, 126.53, 126.19, 125.36, 125.20, 120.34, 118.93, 118.77, 117.22, 114.36, 101.16, 88.85, 69.20, 61.27, 58.73, 54.59, 46.52, 45.99, 42.74, 29.27, 22.31, 20.32, 9.03, 3.80, 3.34. HRMS m/z found 486.2391 $[M + H]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3212, 2152, 1980, 1606, 1505, 1462, 1318, 1118, 771. %Purity 98.96%.

20 **17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(indole-5-carboxamido)morphinan (4)**




The title compound was prepared by following the general procedure in 85% yield.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.39 (s, 1H, exchangeable), 9.21 (s, 1H, exchangeable),

8.89 (s, 1H, exchangeable), 8.16 (s, 1H), 7.80 (d, J=7.76, 1H), 7.65 (d, J=8.44, 1H), 7.44 (d, J=8.84, 1H, exchangeable), 7.43 (s, 1H), 6.72 (d, J=8.04, 1H), 6.58 (d, J=8.08, 1H), 6.55 (s, 1H), 6.37 (s, 1H, exchangeable), 4.79 (d, J=3.52, 1H), 4.66-4.61 (m, 1H), 3.95 (d, J=6.24, 1H), 3.35 (s, 2H), 3.09-2.98 (m, 4H), 2.73 (m, 2H), 1.97-1.93 (m, 1H), 1.65 (d, J=12.04, 1H), 1.56-1.43 (m, 2H), 1.18-1.09 (m, 2H), 0.69-0.62 (m, 2H), 0.50 (m, 1H), 0.41 (m, 1H).

^{13}C NMR (100 MHz, DMSO- d_6) δ 167.01, 146.10, 138.81, 137.43, 128.80, 126.92, 126.64, 125.33, 122.09, 120.66, 120.08, 119.06, 118.34, 110.85, 102.03, 87.55, 69.41, 61.02, 57.02, 45.79, 45.21, 39.54, 30.23, 29.29, 23.54, 19.61, 5.71, 5.17, 2.58. HRMS m/z found 486.2421 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3147, 1627, 1603, 1522, 1502, 1464, 1351, 1323, 1118, 1036, 771, 755, 726. % Purity 99.99%.


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(indole-6-carboxamido)morphinan (5)



The title compound was prepared by following the general procedure in 75% yield.

^1H NMR (400 MHz, DMSO- d_6) δ 11.43 (s, 1H, exchangeable), 9.21 (s, 1H, exchangeable), 8.88 (s, 1H, exchangeable), 7.98 (s, 1H), 7.85 (d, J=7.72, 1H, exchangeable), 7.61 (d, J=8.28, 1H), 7.54 (dd, J=1.32, 8.28, 1H), 7.50 (t, J=2.72, 1H), 6.73 (d, J=8.04, 1H), 6.58 (d, J=8.12, 1H), 6.49 (d, J=1.88, 1H), 6.35 (s, 1H, exchangeable), 4.81 (d, J=3.76, 1H), 4.63 (m, 1H), 3.94 (d, J=6.76, 1H), 3.3 (m, 2H), 3.11-3.05 (m, 2H), 2.98 (m, 1H), 2.72 (m, 1H), 1.94 (m, 1H), 1.65 (d, J=10.96, 1H), 1.57-1.43 (m, 2H), 1.23-1.09 (m, 3H), 0.71-0.59 (m, 2H), 0.50 (m, 1H), 0.40 (m, 1H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 167.16, 145.92, 138.53, 134.98, 129.89, 128.76, 127.88, 127.01, 122.15, 119.46, 119.24, 118.21, 118.10, 111.31, 101.26, 87.53, 69.29, 61.04, 57.03, 45.73, 45.17, 39.28, 30.17, 29.16, 23.44, 19.44, 5.60, 5.18, 2.49. HRMS m/z found 486.2371 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3225, 2162, 1975, 1605, 1540, 1503, 1318, 1119, 1066, 780, 735. % purity 99.93.


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -(indole-7-carboxamido)morphinan (6)



The title compound was prepared by following the general procedure in 22% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 11.13 (s, 1H, exchangeable), 9.25 (s, 1H, exchangeable), 8.91 (s, 1H, exchangeable), 8.14 (d, *J*=7.52, 1H, exchangeable), 7.74 (m, 2H), 7.36 (t, *J*=2.72, 1H), 7.09 (t, *J*=7.6, 1H), 6.71 (d, *J*=8.04, 1H), 6.58 (d, *J*=8.12, 1H), 6.50 (m, 1H), 6.39 (s, 1H, exchangeable), 4.88 (d, *J*=3.68, 1H), 14.70 (m, 1H), 3.95 (d, *J*=6.64, 1H), 3.12-3.04 (m, 3H), 2.94 (m, 1H), 2.73 (m, 2H), 2.00-1.92 (m, 1H), 1.68 (d, *J*=11.2, 1H), 1.56-1.45 (m, 3H), 1.24-1.18 (m, 1H), 1.08 (bs, 1H), 0.70 (m, 1H), 0.62 (m, 1H), 0.50 (m, 1H), 0.41 (m, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ 166.63, 145.93, 138.61, 133.95, 129.03, 128.69, 126.40, 123.86, 122.10, 120.24, 119.20, 118.17, 118.06, 116.80, 101.06, 87.30, 69.31, 69.01, 57.00, 45.61, 45.18, 30.22, 29.15, 23.45, 19.27, 5.62, 5.16, 2.53. HRMS *m/z* found 486.2408 [M + H]⁺. Calculated 485.2315. IR (ATR, cm⁻¹) $\nu_{\max}$ 3060, 2166, 1634, 1585, 1504, 1456, 1312, 1280, 1123, 1030, 984.

17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6α-[2-(indol-3-yl)acetamido)morphinan (7)




The title compound was prepared by following the general procedure in 80% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 10.85 (s, 1H, exchangeable), 9.20 (s, 1H, exchangeable), 8.78 (s, 1H, exchangeable), 7.80 (d, *J*=7.96, 1H, exchangeable), 7.58 (d, *J*=7.92, 1H), 7.33 (d, *J*=8.08, 1H), 7.20 (d, *J*=2.28, 1H), 7.06 (t, *J*=7.50, 7.58, 1H), 6.96 (t, *J*=7.88, 7.04, 1H), 6.71 (d, *J*=8.08, 1H), 6.56 (d, *J*=8.16, 1H), 6.15 (s, 1H, exchangeable), 4.60 (d, *J*=3.88, 1H), 4.40 (m, 1H), 3.84 (d, *J*=6.84, 1H), 3.58 (s, 2H), 3.37 (s, 2H), 3.07-3.00 (m, 2H), 2.91 (m, 1H), 2.69 (m, 1H), 1.18 (m, 1H), 1.61 (d, *J*=10.76, 1H), 1.40 (q, *J*=10.52, 4.4, 9.92, 2H), 1.10 (m, 2H), 0.68 (m, 1H), 0.59 (m, 1H), 0.46 (m, 1H), 0.39 (m, 1H). ¹³CNMR

(100 MHz, MeOD-*d*₆) δ 174.63, 185.37, 146.81, 139.41, 138.03, 131.58, 128.34, 126.14, 125.26, 122.75, 120.82, 120.16, 119.49, 118.91, 112.80, 109.25, 90.01, 71.39, 63.20, 60.22, 45.06, 37.47, 34.11, 33.81, 32.11, 29.98, 23.93, 21.85, 9.43, 5.13, 4.01. HRMS *m/z* found 500.2539 [M + H]⁺. Calculated 499.25. IR (ATR, cm⁻¹) $\nu_{\max}$ 3218, 1640, 1506, 1317, 1234, 117, 1032, 746. % Purity 95.99%.


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -[3-(indol-3-yl) propanamido)morphinan (8)



The title compound was prepared by following the general procedure in 55% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 10.78 (s, 1H, exchangeable), 9.20 (s, 1H, exchangeable), 8.85 (s, 1H, exchangeable), 7.71 (d, J=8.08, 1H exchangeable), 7.53 (d, J=7.88, 1H), 7.32 (d, J=8.04, 1H), 7.12 (d, J=2.16, 1H), 7.05 (t, J=7.08, 1H), 6.97 (t, J=7.88, 1H), 6.71 (d, J=8.08, 1H), 6.55 (d, J=8.16, 1H), 6.28 (s, 1H, exchangeable), 4.57 (d, J=3.84, 1H), 4.42 (m, 1H), 3.90 (d, J=6.8, 1H), 3.25 (m, 1H), 3.17 (s, 2H), 3.07-3.01 (m, 2H), 2.93 (t, J=7.72, 3H), 2.74-2.66 (m, 1H), 2.53 (m, 1H), 2.44 (dd, J=4.76, 13.36, 1H), 1.90-1.82 (m, 1H), 1.60 (d, J=10.92, 1H), 1.42-1.36 (m, 2H), 1.11-1.06 (m, 1H), 0.96-0.90 (m, 1H), 0.70-0.59 (m, 2H), 0.50-0.46 (m, 1H), 0.42-0.37 (m, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ 169.37, 169.29, 143.31, 136.02, 133.54, 126.26, 124.50, 119.45, 118.46, 116.73, 115.85, 115.57, 111.26, 108.78, 85.09, 66.74, 58.46, 54.49, 45.97, 42.61, 42.29, 33.51, 27.59, 26.55, 20.90, 18.48, 17.08, 3.12, 2.70. HRMS *m/z* found 514.2691, [M + H]⁺. Calculated 513.2628. IR (ATR, cm⁻¹) $\nu_{\max}$ 3267, 3146, 2157, 1670, 1616, 1543, 1504, 1463, 1427, 1343, 1316, 1172, 1118, 1031


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 α -[4-(indol-3-yl) butanamido)morphinan (9)



The title compound was prepared by following the general procedure in 34% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 10.79 (s, 1H, exchangeable), 9.17 (s, 1H, exchangeable), 8.86 (s, 1H, exchangeable), 7.64 (d, *J*=8.04, 1H, exchangeable), 7.51 (d, *J*=7.84, 1H), 7.32 (d, *J*=8.04, 1H), 7.11 (d, *J*=2.16, 1H), 7.05 (t, *J*=7.04, 1H), 7.96 (t, *J*=7.88, 1H), 6.71 (d, *J*=8.08, 1H), 6.54 (d, *J*=8.12, 1H), 6.29 (s, 1H, exchangeable), 4.59 (d, *J*=3.88, 1H), 4.45-4.38 (m, 1H), 3.91 (d, *J*=6.8, 1H), 3.32 (d, *J*=19.72, 1H), 3.25 (m, 1H), 3.02 (m, 2H), 2.96 (m, 1H), 2.69 (t, *J*=7.4, 3H), 2.43 (dd, *J*=4.92, 13.32, 1H), 2.22 (t, *J*=7.32, 2H), 1.92-1.83 (m, 1H), 1.60 (d, *J*=13.16, 1H), 1.38 (m, 2H), 1.07 (m, 1H), 0.99-0.87 (m, 1H), 0.69-0.59 (m, 2H), 0.50-0.45 (m, 1H), 0.41-0.36 (m, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ 173.08, 146.69, 139.36, 137.00, 129.64, 127.98, 123.04, 122.95, 121.78, 120.15, 119.20, 119.07, 118.97, 115.05, 112.19, 88.45, 70.11, 61.85, 57.90, 45.98, 45.66, 35.98, 30.97, 29.95, 27.11, 25.11, 24.28, 20.41, 15.92, 6.46, 6.08, 3.34. HRMS *m/z* found 528.2876, 556.2742, 1055.5772 [*M* + *H*]⁺. Calculated 527.2784. IR (ATR, cm⁻¹) $\nu_{\max}$ 3267, 3124, 2162, 1671, 1623, 1506, 1456, 1373, 1119

17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6β-(indole-2-carboxamido)morphinan (10)




The title compound was prepared by following the general procedure in 32% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 11.60 (s, 1H, exchangeable), 9.33 (s, 1H, exchangeable), 8.88 (s, 1H, exchangeable), 8.73 (d, *J*=7.92, 1H, exchangeable), 7.62 (d, *J*=7.96, 1H), 7.42 (d, *J*=8.24, 1H), 7.18 (m, 2H), 7.04 (t, *J*=7.52, 1H), 6.73 (d, *J*=8.08, 1H), 6.66 (d, *J*=7.96, 1H), 6.24 (s, 1H, exchangeable), 4.84 (d, *J*=7.72, 1H), 3.89 (s, 1H), 3.71 (m, 1H), 3.33 (m, 2H), 3.11-3.06 (m, 2H), 2.88 (m, 1H), 2.50 (m, 2H), 1.96-1.78 (m, 2H), 1.63 (m, 1H), 1.49-1.40


(m, 2H), 1.09 (s, 1H), 0.68 (s, 1H), 0.60 (s, 1H), 0.52 (s, 1H), 0.42 (s, 1H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 160.73, 142.02, 141.09, 136.23, 131.30, 129.62, 126.97, 125.26, 123.46, 121.52, 120.64, 119.83, 119.39, 117.87, 112.23, 102.63, 89.85, 69.62, 61.67, 56.69, 50.66, 46.37, 45.61, 29.37, 23.78, 22.91, 5.62, 5.07, 2.55. HRMS m/z found 486.2411 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3212, 2167, 1975, 1617, 1551, 1506, 1315, 1239, 1125, 1034, 815, 747, 668. % Purity 96.75%.

17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 β -(indole-3-carboxamido)morphinan (11)



The title compound was prepared by following the general procedure in 27% yield. ^1H NMR (400 MHz, DMSO- d_6) δ 11.64 (s, 1H, exchangeable), 9.34 (bs, 1H, exchangeable), 8.89 (s, 1H, exchangeable), 8.13-8.10 (m, 3H, 1 proton exchangeable), 7.42 (d, $J=7.88$, 1H), 7.16-7.06 (m, 2H), 6.73 (d, $J=8.08$, 1H), 6.65 (d, $J=8.12$, 1H), 6.24 (s, 1H, exchangeable), 4.82 (d, $J=7.76$, 1H), 3.90 (s, $J=4.28$, 1H), 3.70 (m, 1H), 3.36 (s, 1H), 3.12-3.05 (m, 3H), 2.88 (m, 2H), 1.89 (q, $J=12.32$, 1H), 1.78 (d, $J=13.2$, 1H), 1.63 (m, 1H), 1.48-1.41 (m, 2H), 1.22 (bs, 1H), 0.68 (m, 1H), 0.50 (m, 1H), 0.52 (m, 1H), 0.42 (m, 1H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 164.46, 142.18, 141.03, 135.90, 129.70, 127.60, 125.86, 121.98, 120.89, 120.63, 120.43, 119.29, 117.89, 111.79, 110.30, 90.17, 69.69, 61.70, 56.70, 50.21, 46.38, 45.60, 29.40, 27.30, 23.99, 22.93, 5.61, 5.12, 2.53. HRMS m/z found 486.2411 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3267, 1671, 1621, 1539, 1505, 1455, 1313, 1176, 1119


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 β -(indole-4-carboxamido)morphinan (12)



The title compound was prepared by following the general procedure in 31% yield.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.31 (s, 1H, exchangeable), 9.34 (s, 1H, exchangeable), 8.87 (bs, 1H, exchangeable), 8.39 (d, $J=8.12$, 1H, exchangeable), 7.55 (d, $J=8.04$, 1H), 7.48 (d, $J=7.16$, 1H), 7.43 (m, 1H), 7.15 (t, $J=7.72$, 1H), 6.88 (s, 1H), 6.73 (d, $J=8.12$, 1H), 6.66 (d, $J=8.16$, 1H), 6.19 (s, 1H, exchangeable), 4.89 (d, $J=7.76$, 1H), 3.88 (d, $J=4.92$, 1H), 3.77-3.72 (m, 1H), 3.32 (m, 1H), 3.13-3.04 (m, 2H), 2.87 (m, 1H), 1.93 (q, $J=12.08$, 1H), 1.78 (d, $J=13.68$, 1H), 1.65 (m, 1H), 1.46 (m, 2H), 1.09 (t, $J=6.96$, 2H), 0.70-0.68 (m, 1H), 0.60 (m, 1H), 0.53-0.50 (m, 1H), 0.43-0.41 (m, 1H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 167.77, 142.15, 141.02, 136.35, 129.67, 126.30, 126.09, 125.75, 120.64, 120.19, 119.34, 118.50, 117.90, 114.31, 101.78, 89.92, 69.68, 61.68, 56.68, 50.89, 46.37, 45.61, 29.43, 27.29, 23.66, 22.90, 5.58, 5.11, 2.50. HRMS m/z found 486.2434 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3207, 2166, 1639, 1503, 1455, 1319, 1237, 1195, 1123, 1037, 917.


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 β -(indole-5-carboxamido)morphinan (13)



The title compound was prepared by following the general procedure in 61% yield.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.36 (s, 1H, exchangeable), 9.32 (s, 1H, exchangeable), 8.88 (s, 1H, exchangeable), 8.48 (d, $J=8.08$, 1H, exchangeable), 8.18 (s, 1H), 7.66 (dd, $J=1.56$, 8.56, 1H), 7.42 (m, 2H), 6.73 (d, $J=8.12$, 1H), 6.65 (d, $J=8.16$, 1H), 6.53 (s, 1H), 6.20 (s, 1H, exchangeable), 4.89 (d, $J=7.8$, 1H), 3.88 (s, 1H), 3.71 (m, 1H), 3.35 (s, 2H), 3.11-3.04 (m, 2H), 2.87 (m, 1H), 1.90 (q, $J=12.6$, 1H), 1.78 (d, $J=13.76$, 1H), 1.62 (m, 1H), 1.49-1.39 (m, 2H), 1.09 (m, 1H), 0.68 (m, 1H), 0.59 (m, 1H), 0.52 (m, 1H), 0.42 (m, 1H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 166.77, 142.30, 141.30, 137.41, 129.77, 126.99, 126.60, 125.29, 120.56, 120.49, 119.91, 119.16, 117.92, 110.80, 102.05, 90.06, 69.80, 61.81, 56.74, 51.11, 46.52, 45.62, 29.44, 27.41, 23.93, 23.06, 5.72, 5.09, 2.65. HRMS m/z found 486.2434 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3125, 2162, 1980, 1622, 1595, 1547, 1360, 1311, 1249, 1124, 1037, 919, 760, 672. % Purity 98.51%.


17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6β-(indole-6-carboxamido)morphinan (14)



The title compound was prepared by following the general procedure in 29% yield.

5 ¹HNMR (400 MHz, DMSO-*d*₆) δ 11.43 (s, 1H, exchangeable), 9.33 (s, 1H, exchangeable), 8.87 (s, 1H, exchangeable), 8.54(d, J=7.4, 1H, exchangeable), 7.99 (s, 1H), 7.58 (s, 2H), 7.50 (s, 1H), 6.73 (d, J=6.36, 1H), 6.66 (d, J=7.68, 1H), 6.49 (s, 1H), 6.20 (s, 1H, exchangeable), 4.90 (d, J=5.92, 1H), 3.88 (s, 1H), 3.71 (s, 1H), 3.17-3.06 (m, 2H), 2.89 (m, 2H), 2.5 (s, 1H), 1.90 (m, 1H), 1.78 (d, J=12.32, 1H), 1.62 (d, J=10.56, 1H), 1.49-1.39 (m, 2H), 1.09 (t, J=6.52, 2H), 0.68 (s, 1H), 0.60 (s, 1H), 0.52 (s, 1H), 0.42 (s, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ 166.85, 142.16, 140.23, 135.05, 129.76, 127.74, 127.20, 119.34, 118.39, 117.95, 116.98, 111.16, 101.20, 90.74, 69.62, 61.76, 58.35, 51.50, 49.99, 39.96, 39.76, 30.60, 29.99, 24.65, 22.25, 3.72, 3.46. HRMS *m/z* found 486.2396 [M + H]⁺. Calculated 485.2315. IR (ATR, cm⁻¹) $\nu_{\max}$ 3223, 2152, 1970, 1608, 1551, 1312, 1035, 920, 820, 740. %
15 Purity 99.86%.

17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6β-(indole-7-carboxamido)morphinan (15)




The title compound was prepared by following the general procedure in 56% yield.

20 ¹HNMR (400 MHz, DMSO-*d*₆) δ 11.07 (s, 1H, exchangeable), 9.37 (s, 1H, exchangeable), 8.90 (s, 1H, exchangeable), 8.75 (d, J=8.04, 1H, exchangeable), 7.78-7.75 (dd, J=4.24, 3.12, 2H), 7.35 (t, J=2.6 1H), 7.10 (t, J=7.6, 1H), 6.74 (d, J=8.12, 1H), 6.67 (d, J=8.16, 1H), 6.49 (m, 1H), 6.26 (bs, 1H, exchangeable), 4.93 (d, J=7.8, 1H), 3.91 (d, J=4.84, 1H), 3.82-3.78 (m, 2H), 3.18-3.06 (m, 3H), 2.91-2.86 (m, 1H), 2.02-1.93 (q, J=11.56, 1H), 1.65 (m, 1H),


1.56-1.41 (m, 2H), 1.10 (t, J=7, 2H), 0.71-0.68 (m, 1H), 0.62-0.60 (m, 1H), 0.55-0.52 (m, 1H), 0.44-0.42 (m, 1H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 164.50, 139.51, 138.46, 131.48, 127.15, 126.67, 123.95, 121.72, 118.20, 117.28, 117.02, 115.79, 115.40, 113.77, 98.68, 87.35, 67.21, 62.45, 59.20, 54.25, 48.23, 43.86, 43.16, 26.96, 24.75, 21.14, 20.39, 12.43, 3.07, 2.65. HRMS m/z found 486.2411 $[\text{M} + \text{H}]^+$. Calculated 485.2315. IR (ATR, cm^{-1}) ν_{max} 3090, 2162, 1979, 1633, 1540, 1327, 1248, 1040

17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 β -[2-(indol-3-yl)acetamido)morphinan (16)



The title compound was prepared by following the general procedure in 49% yield. ^1H NMR (400 MHz, DMSO- d_6) δ 10.89 (s, 1H, exchangeable), 9.32 (s, 1H, exchangeable), 8.82 (s, 1H, exchangeable), 8.20 (d, J=7.92, 1H, exchangeable), 7.54 (d, J=7.76, 1H), 7.34 (d, J=8.04, 1H), 7.18 (d, J=2.08, 1H), 7.06 (t, J=7.08, 1H), 6.98 (t, J=7.24, 1H), 6.70 (d, J=8.12, 1H), 6.62 (d, J=8.16, 1H), 6.14 (s, 1H, exchangeable), 4.60 (d, J=7.84, 1H), 3.82 (d, J=5.2, 1H), 3.50 (q, J=14.96, 2H), 3.40 (m, 1H), 3.32 (m, 1H), 3.28 (s, 1H), 3.07-3.01 (dd, J=5.96, 19.24, 2H), 2.84 (m, 1H), 2.43-2.39 (m, 2H), 1.68 (m, 2H), 1.49-1.28 (m, 3H), 1.07 (m, 1H), 0.67-0.66 (m, 1H), 0.60-0.56 (m, 1H), 0.51-0.48 (m, 1H), 0.41-0.39 (m, 1H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 170.94, 170.86, 141.99, 140.99, 135.88, 129.55, 127.03, 123.57, 121.02, 120.58, 119.36, 118.53, 118.42, 117.83, 111.28, 108.49, 99.49, 69.56, 61.54, 56.62, 46.32, 45.49, 32.78, 29.17, 27.21, 23.45, 22.82, 5.55, 5.08, 2.46. HRMS m/z found 500.2539 $[\text{M} + \text{H}]^+$. Calculated 499.2471. IR (ATR, cm^{-1}) ν_{max} 3270, 1666, 1549, 1463, 1303, 1274, 1232, 1174, 1124, 1030, 898


17-Cyclopropylmethyl-3,14 β -dihydroxy-4,5 α -epoxy-6 β -[3-(indol-3-yl) propanamido)morphinan (17)



The title compound was prepared by following the general procedure in 70% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 10.77 (s, 1H, exchangeable), 9.34 (s, 1H, exchangeable), 8.83 (s, 1H, exchangeable), 8.14 (d, *J*=7.8, 1H, exchangeable), 7.52 (d, *J*=7.8, 1H), 7.32 (d, *J*=8.04, 1H), 7.10 (s, 1H), 7.06 (t, *J*=7.96, 1H), 6.97 (t, *J*=7.88, 1H), 6.72 (d, *J*=8.12), 6.63 (d, *J*=8.16, 1H), 6.18 (s, 1H, exchangeable), 4.55 (d, *J*=7.8, 1H), 3.85 (s, 1H), 3.45 (m, 1H), 3.34-3.29 (m, 2H), 3.09-3.00 (m, 2H), 2.92-2.83 (m, 3H), 2.45-2.42 (m, 4H), 1.71-1.68 (m, 2H), 1.53-1.50 (m, 1H), 1.42 (d, *J*=9.76, 1H), 1.33 (t, *J*=12.32, 1H), 1.08 (m, 1H), 0.68-0.63 (m, 1H), 0.60-0.57 (m, 1H), 0.52-0.48 (m, 1H), 0.42-0.38 (m, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ 141.98, 140.93, 135.97, 129.53, 126.86, 121.96, 120.98, 120.64, 119.41, 118.26, 117.89, 113.57, 111.27, 89.91, 69.58, 61.58, 56.67, 50.47, 46.31, 45.49, 39.62, 29.21, 27.21, 23.43, 22.83, 20.85, 5.55, 5.11, 2.47, 0.012. HRMS *m/z* found 514.2714, [M + H]⁺. Calculated 513.2628. IR (ATR, cm⁻¹) $\nu_{\max}$ 3187, 2160, 1640, 1541, 1505, 1450, 1325, 1271, 1234, 1124, 1032, 854

17-Cyclopropylmethyl-3,14β-dihydroxy-4,5α-epoxy-6β-[4-(indol-3-yl) butanamido]morphinan 18




The title compound was prepared by following the general procedure in 39% yield.

¹HNMR (400 MHz, DMSO-*d*₆) δ 10.79 (s, 1H, exchangeable), 9.35 (bs, 1H, exchangeable), 8.84 (s, 1H, exchangeable), 8.09 (d, *J*=7.88 1H, exchangeable), 7.51 (d, *J*=7.84, 1H), 7.33 (d, *J*=8.04, 1H), 7.12 (d, *J*=2.08, 1H), 7.08 (t, *J*=7.46, 1H), 6.97 (t, *J*=7.42, 1H), 6.72 (d, *J*=8.12, 1H), 6.64 (d, *J*=8.16, 1H), 6.18 (bs, *J*=1H, exchangeable), 4.56 (d, *J*=7.84, 1H), 3.85 (d, *J*=4.84, 1H), 3.09-3.03 (dd, *J*=5.88, 19.48, 2H), 2.88-2.84 (m, 1H), 2.68 (t, *J*=7.48, 2H), 2.44-2.41 (m, 2H), 2.16 (t, *J*=7.32, 2H), 1.89-1.85 (m, 2H), 1.75-1.70 (m, 2H), 1.56-1.53 (m,

1H), 1.43 (d, J=9.68, 1H), 1.38-1.31 (m, 1H), 1.12-1.05 (m, 2H), 0.69-0.67 (m, 1H),
 0.61-0.58 (m, 1H), 0.53-0.50 (m, 1H), 0.43-0.40 (m, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ
 142.02, 140.99, 136.07, 129.56, 127.05, 122.03, 120.88, 120.60, 119.35, 118.28, 118.18,
 117.87, 114.11, 111.27, 89.88, 69.59, 67.88, 61.59, 56.65, 50.44, 46.33, 45.49, 29.24, 27.22,
 26.15, 24.05, 23.53, 22.85, 15.00, 5.56, 5.10, 2.48. HRMS *m/z* found 528.2874 [M + H]⁺.
 Calculated 527.2784. IR (ATR, cm⁻¹) $\nu_{\max}$ 3268, 1671, 1633, 1505, 1455, 1423, 1316, 1178,
 1127, 1033

**17-Cyclopropylmethyl-3,14 β -Dihydroxy-4,5 α -Epoxy-6 α -(Isoquinoline-3-Carboxamido)
 morphinan (NAQ)**



¹HNMR (400 MHz, DMSO-*d*₆) δ 9.44 (s, 1H), 8.98 (bs, 1H, exchangeable), 8.65 (s,
 1H), 8.57 (d, J=8.76, exchangeable), 8.28 (d, J=8.08, 1H), 8.24 (d, J=8.08, 1H), 7.94 (t,
 J=8.04, 1H), 7.85 (d, J=8, 1H), 6.80 (d, J=8.04, 1H), 6.60 (d, J=8.16, 1H), 4.80 (d, J=3.76,
 1H), 4.77-4.70 (m, 1H), 4.03 (d, J=6.68, 1H), 3.38 (d, J=19.68, 1H), 3.31-3.25 (m, 1H),
 3.13-3.02 (m, 3H), 2.78-2.72 (m, 1H), 2.57 (dd, J=4.76, 13.6, 1H), 2.04-1.98 (m, 1H),
 1.68-1.65 (m, 2H), 1.50-1.44 (m, 1H), 1.14-1.02 (m, 2H), 0.72-0.66 (m, 1H), 0.65-0.62 (m,
 1H), 0.53-0.50 (m, 1H), 0.43-0.41 (m, 1H). ¹³CNMR (100 MHz, DMSO-*d*₆) δ 163.46,
 151.54, 145.32, 141.94, 138.54, 135.30, 132.08, 129.72, 129.12, 128.66, 128.07, 127.91,
 122.03, 120.21, 119.73, 118.12, 87.44, 69.21, 57.05, 48.43, 45.27, 45.17, 45.07, 29.92,
 28.87, 23.31, 19.82, 5.50, 5.19, 2.38. HRMS *m/z* found 498.2461 [M + H]⁺. Calculated
 497.2315. IR (ATR, cm⁻¹) $\nu_{\max}$

Instruments used for characterization

The Bruker 400 MHz NMR was used to determine the ¹H and ¹³C NMR spectra of
 the compounds. The masspec and IR spectra were determined using the Perkin Elmer ToF
 mass spectrometer and Thermo Scientific smart iTR instruments respectively. The purity of
 the compounds was determined using Varian Prostar HPLC instrument. The HPLC
 parameters used were: column= C18, injection volume=5 μ L, sample concentration= 3mM,
 mobile phase=60 H₂O : 40 MeCN, flowrate=1 mL/min, detector= UV detection at 210 and

254 nm.

EXAMPLE 3. MOLECULAR MODELING STUDIES

Molecular modeling studies were conducted to observe the binding interactions of compound **6** with MOR. Chemical structures of the compounds were sketched in SybylX-2.1, and Gasteiger-Hückel charges were assigned before energy minimization (100 000 iterations) with the Tripos Force Fields. The X-ray crystal structure for MOR (4DKL) was retrieved from the Protein Data Bank (PDB) and prepared for docking by adding hydrogens and deleting water molecules and bound ligands inside the binding pocket.

Molecular docking study

Automated docking on MOR was done utilizing genetic algorithm docking program GOLD 5.4. The binding site was defined to include all atoms within 10 Å of the γ -carbon atom of Asp147 along with a hydrogen bond constraint between the 17-N of the ligand's morphinan skeleton and the carboxylate group of Asp147. Pictures were generated using PyMOL Molecular Graphics System, version 1.7.4.5

Compound **6** was first sketched with sybylx2.1 and then docked 50 times into the crystal structure of MOR (4dkl). From the docking results obtained, it was observed that compound **6** docked in two sites (not shown). The morphinan part of the compound was docked in the same position in both sites, however, the indole ring was docked differently. In binding site 1, the nitrogen of the indole ring was hydrogen bonded to K303 and the indole ring formed pi-pi interactions with W318. In binding site 2, the nitrogen of the indole ring formed hydrogen bonds with K233. The orientation of the indole ring was away from W318 and pi-pi stacking was not observed. The CHEM-PLP scores for the two binding sites were quite similar, however binding pose 1 had better GOLD and HINT scores than binding pose 2 (not shown).

Molecular dynamics simulation

The best CHEM-PLP-scored solutions for binding pose 1 and 2 were chosen for molecular dynamics (MD) studies. Force field parameter and topology files for NNQ were generated utilizing CGenFF. Coordinates for the spatial arrangement of the receptors within the lipid bilayer were retrieved from the Orientations of Proteins in Membranes (OPM) database. The simulation system, consisting of the receptor-ligand complex embedded in a lipid (POPC) bilayer surrounded with saline solution (0.15 MNaCl) was created in VMD 1.9.3 using the CHARMM force field topology file. All simulations were performed under

hybrid CHARMM force field parameters that included protein, lipids and ligand with a time-step of 2 femtoseconds (fs). Periodic boundary conditions were employed, and Particle Mesh Ewald (PME) summation was used to calculate long-range electrostatic interactions. Non-bonded interactions were calculated with a smooth cutoff at 14 Å with a frequency of 2 fs. The temperature was maintained at 310 K via Langevin dynamics. All molecular modeling simulations were performed using NAMD 2.8. MD simulations were carried out in four stages. In the first stage, equilibration of the fluid-like lipid bilayer was performed via minimization (1000 iterations) followed by NPT equilibration (pressure equilibration, 0.5 ns) of the lipid tails only. In the second stage, an NPT equilibration of the system was run for a period of 1 ns with harmonic constraints placed on protein and NNQ atoms (5 kcal/(mol-Å)). The harmonic restraint was released in stage 3 and the entire system was equilibrated using the NVT canonical ensemble for a further 1 ns. The final production run was conducted for 10 ns using an NVT ensemble. Energy landscape analysis was performed using the NAMD Energy 1.4 plug-in; non-bonded interaction analyses were performed at various distances with a dielectric constant of 6.5. The best-scored poses based on the NAMD non-bonded interactions were selected for further analysis.

For binding pose 1, it was observed that after 1 ns of simulation, the indole ring of compound 6 still formed pi-pi interactions with W318. The nitrogen of the indole ring was hydrogen bonded to K303. However, it was observed that after 10 ns of simulation the indole ring moved into the binding pocket 2. In the case of binding pose 2), it was observed that there was no change in the binding pose after 1 ns of MD simulation nor after 10 ns. It was observed that the RMSD for binding pose 2 was lower than that of binding pose 1 during the simulation. The HINT score for binding pose 2 improved after 10 ns of simulation but that of binding pose 1 worsened. Thus, compound 6 binds to MOR by interacting with K233, D147, Y148, and Y326 (binding pose 2).

SUMMARY

These Examples describe new morphinan analogues which are highly selective and reversible MOR antagonists. The MOR binding affinities of several compounds in this series were determined and the results showed that all compounds tested bound with high affinity (≤ 5 nM). In fact, several bound with sub-nM affinity (in the range of 100-500 pM). These results indicate that the molecular design strategy was successful in producing very high affinity MOR ligands.


While the invention has been described in terms of its several exemplary embodiments, those skilled in the art will recognize that the invention can be practiced with modification within the spirit and scope of the appended claims. Accordingly, the present invention should not be limited to the embodiments as described above, but should further
5 include all modifications and equivalents thereof within the spirit and scope of the description provided herein.

CLAIMS

We claim:

1. A compound with generic Formula I:

5




Formula I

wherein * indicates a chiral C that can be either an R configuration or an S configuration in terms of chirality;

10

and R of Formula I =



and is bonded to Formula I at X₂, X₃, X₄, X₅, X₆ or X₇;

and wherein

15

X₁ = N, O or S;


one or more of X₄, X₅, X₆ and X₇ are independently substituted by R' where R' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group, and

one or both of X₂ and X₃ are independently substituted by R'' where R'' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group;

20

and salts thereof;

with the caveat that R is not:



5

2. The compound of claim 1, wherein the straight chain or branched alkyl group comprises from 1 to 20 carbon atoms.

3. The compound of claim 1, wherein the straight chain or branched alkyl group is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl, octyl, nonyl, decyl, undecyl and dodecyl, and isomers thereof.


15

4. The compound of claim 1, wherein the electron withdrawing group is selected from the group consisting of F, CN, NO₂, SO₃ and CF₃.

5. The compound of claim 1, wherein the electron donating group is OCH₃, phenyl or substituted phenyl.

6. A method of treating opioid addiction in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of a compound of Formula I:


20



Formula I

wherein * indicates a chiral C that can be either an R configuration or an S configuration in terms of chirality;

and R of Formula I =



and is bonded to Formula I at X₂, X₃, X₄, X₅, X₆ or X₇;

and wherein

X₁ = N, O or S;

one or more of X₄, X₅, X₆ and X₇ are independently substituted by R' where R' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group, and

one or both of X₂ and X₃ are independently substituted by R'' where R'' = a straight chain or branched alkyl group, a straight chain or branched alkene group, an electron withdrawing group, or an electron donating group;

or a physiologically acceptable salt thereof.

7. The method of claim 6, wherein the straight chain or branched alkyl group comprises from 1 to 20 carbon atoms.



8. The method of claim 6, wherein the straight chain or branched alkyl group is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl, octyl, nonyl, decyl, undecyl and dodecyl, and isomers thereof.

9. The method of claim 6, wherein the electron withdrawing group is selected from the group consisting of F, CN, NO₂, SO₃ and CF₃.


5 10. The method of claim 6, wherein the electron donating group is OCH₃, phenyl or substituted phenyl.

11. The method of claim 6, wherein the compound is

10




or



15


12. A method of treating pain in a subject in need thereof, comprising
administering to the subject a therapeutically effective amount of a compound of
Formula I:



Formula I

wherein * indicates a chiral C that can be either an R configuration or an S configuration in
terms of chirality;

and R of Formula I =



and is bonded to Formula I at X₂, X₃, X₄, X₅, X₆ or X₇;

and wherein


X₁ = N, O or S;

one or more of X₄, X₅, X₆ and X₇ are independently substituted by R' where R' = a
straight chain or branched alkyl group, a straight chain or branched alkene group, an electron
withdrawing group, or an electron donating group, and

one or both of X₂ and X₃ are independently substituted by R'' where R'' = a straight
chain or branched alkyl group, a straight chain or branched alkene group, an electron
withdrawing group, or an electron donating group;

or a physiologically acceptable salt thereof;

with the caveat that R is not:




5 13. The method of claim 12, wherein the straight chain or branched alkyl group comprises from 1 to 20 carbon atoms.

14. The method of claim 12, wherein the straight chain or branched alkyl group is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, pentyl, heptyl, septyl,
 10 octyl, nonyl, decyl, undecyl and dodecyl, and isomers thereof.

15. The method of claim 12, wherein the electron withdrawing group is selected from the group consisting of F, CN, NO₂, SO₃ and CF₃.

15 16. The method of claim 12, wherein the electron donating group is OCH₃, phenyl or substituted phenyl.

17. The method of claim 12, wherein the compound is






Figure 1A




Figure 1B




Figure 1C




Figure 1D




Figure 2A




Figure 2B




Figure 2C




Figure 3A




Figure 3B




Figure 3C

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 17/32792

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(8) - A61K 31/485; C07D 489/08 (2017.01)
 CPC - C07D 403/12; C07D 405/12; C07D 491/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History Document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History Document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History Document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2014/0371255 A1 (ZHANG et al.) 18 December 2014 (18.12.2014) Fig 14; pg 31, Table 8; para [0012]-[0014], [0027], [0045], [0073], [0170], [0463]	1-17
Y	US 5,330,986 A (SHUTSKE) 19 July 1994 (19.07.1994) col 4, ln 60-68; col 5, Table 1	1-5, 12-17
Y	US 2013/0281698 A1 (SCHUETZ et al.) 24 October 2013 (24.10.2013) para [0161]-[0162], [0319], [0377]	6-11
A	US 2015/0203504 A1 (GOEHRING et al.) 23 July 2015 (23.07.2015) Entire Document	1-17
A	US 2013/0224151 A1 (PEARSON et al.) 29 August 2013 (29.08.2013) Entire Document	1-17
A	WO 2012/166891 A2 (ALGYNOMICS INC.) 06 December 2012 (06.12.2012) Entire Document	1-17
A	US 7,320,984 B2 (IZUMIMOTO et al.) 22 January 2008 (22.01.2008) Entire Document	1-17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 July 2017

Date of mailing of the international search report

25 AUG 2017

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
 P.O. Box 1450, Alexandria, Virginia 22313-1450
 Facsimile No. 571-273-8300

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
 PCT OSP: 571-272-7774