

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 3442977 T3**

-
- (51) Int.Cl.: **C 07 D 487/04 (2006.01)** **A 61 K 31/5025 (2006.01)** **A 61 P 19/00 (2006.01)**
C 07 D 519/00 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2023-10-09**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-06-28**
- (86) Europæisk ansøgning nr.: **17733108.9**
- (86) Europæisk indleveringsdag: **2017-04-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-02-20**
- (86) International ansøgning nr.: **US2017027775**
- (87) Internationalt publikationsnr.: **WO2017181117**
- (30) Prioritet: **2016-04-15 US 201662322948 P** **2016-10-21 US 201662411172 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Blueprint Medicines Corporation, 45 Sidney Street, Cambridge, MA 02139, USA**
- (72) Opfinder: **BROOIJMANS, Natasja, 43 Concord Square Apt. 2, Boston, MA 02118, USA**
BRUBAKER, Jason, D., 26 Sciarappa St Unit 1, Cambridge, MA 02141, USA
FLEMING, Paul, E., 7 Sunnyside Ave, Wellesley, MA 02482, USA
HODOUS, Brian, L., 22 Lawrence Lane, Arlington, MA 02474, USA
KIM, Joseph, L., 20 Green Way, Wayland, MA 01778, USA
WAETZIG, Josh, 30 Livermore Road, Waltham, MA 02451, USA
WILLIAMS, Brett, 24 Castleton St., Boston, MA 02130, USA
WILSON, Kevin, J., 27 Beryl Street, Boston, MA 02131, USA
CRONIN, Mark, "--", deceased, USA
WILSON, Douglas, Blueprint Medicines Corporation, 45 Sidney Street, Cambridge MA 02139, USA
- (74) Fuldmægtig i Danmark: **Inspicos P/S, Agern Allé 24, 2970 Hørsholm, Danmark**
- (54) Benævnelse: **INHIBITORER AF ACTIVINRECEPTORLIGNENDE KINASE**
- (56) Fremdragne publikationer:
WO-A1-2014/138088

INHIBITORS OF ACTIVIN RECEPTOR-LIKE KINASE

Description

[0001] This invention relates to inhibitors of Activin receptor-like kinase-2 (ALK2).

5 BACKGROUND

[0002] Activin receptor-like kinase-2 (ALK2) is encoded by the Activin A receptor, type I gene (ACVR1). ALK2 is a serine/threonine kinase in the bone morphogenetic protein (BMP) pathway (Shore et al., Nature Genetics 2006, 38: 525-27). It binds to complexes comprising bone morphogenetic proteins (BMPs) and is responsible for transducing BMP signals. Certain mutations in ALK2 cause the kinase to be constitutively active and are associated with various diseases. Fibrodysplasia ossificans progressiva (FOP) is a rare, severely debilitating heritable disorder characterized by progressive heterotopic ossification in extraskeletal sites. Individuals with this disease experience significantly reduced mobility and shortened lifespan. Current therapy is limited to ameliorating swellings (flare-ups) that characterize the disease.

[0003] All FOP patients carry heterozygous, activating mutations in the ACVR1 gene. Further, the vast majority of FOP patients harbor the same ALK2 mutation, R206H. Transgenic mice that express ALK2-R206H recapitulate the key features of the human disease, including malformation of the first digit in the hind limbs and inflammatory infiltration and muscle cell apoptosis followed by formation of heterotopic bone through an endochondral pathway (Chakkalakal et al., J Bone Miner Res. 2012, 27(8): 1746-1756). A second engineered mouse strain has been developed that expresses the activated ALK2-Q207D variant in muscle and phenocopies key features of human FOP. Treatment of these mice with an inhibitor of BMP receptor type 1 kinases resulted in inhibition of SMAD signaling and reduction in ectopic ossification and associated functional impairment (Fukuda et al., Genesis 2006, 44, 159-167). Other mutations in ALK2 that have been associated with FOP include but are not limited to L196P, PF197-8L, R202I, R258S, R258G, G328A, G328W, G328E, G328R, G356D, and R375P (Kaplan et al., Hum Mutat. 2009, 30(3): 379 - 390; Gregson et al., Bone 2011, 48:654-658; Kaplan et al., Am J Med Genet 2015, 167: 2265-2271; Petrie et al., PLoS One 2009, 4(3): e5005; Bocciardi et al., Eur J Hum Genetics 2009, 17:311-318; Pacifici and Shore, Cytokine & Growth Factor Reviews 2016, 27:93-104).

[0004] In certain circumstances, heterotopic ossification (HO) can also be induced in people who are wild-type ALK2. These circumstances can include major surgical interventions, trauma (such as head or blast injuries), protracted immobilization, or severe burns. An ALK2 inhibitor could potentially be an effective therapy for the treatment of FOP and other conditions caused by HO.

[0005] Diffuse intrinsic pontine glioma (DIPG) is a rare, aggressive and typically fatal pediatric brain stem cancer with no effective treatment options. Due to its anatomical

location and diffuse nature, DIPG cannot be treated by surgery. DIPG arises exclusively in young children and the two year survival rate is approximately less than 10%. Because of their location in the brainstem, DIPGs cause pressure on cranial nerves leading to double vision, difficulty in controlling eye movement, difficulty chewing/ swallowing, weakness in the arms/ legs leading to loss of movement and difficulty speaking. As the tumor progresses there is increasing pressure inside the skull causing severe headaches, nausea vomiting and fatigue. Unlike many other pediatric cancers, there has been virtually no progress in improving treatments for DIPG over the last few decades. Historically, the lack of understanding regarding the drivers of DIPG has hindered the identification of potential new treatment options. Consequently, the medical need for DIPG treatments is exceedingly high. Recent genomic characterization has demonstrated that ~25% of DIPG tumors possess somatic, heterozygous ALK2 activating mutations. Mutations in ALK2 associated with DIPG include, but are not limited to R206H, G328V, G328W, G328E, and G356D (Jones and Baker, Nature Rev Cancer 2014, 14:651-661).


[0006] Notably, the ALK2 mutations found in DIPG overlap with those found in FOP, suggesting a potential synergy between inhibitor development efforts for the two diseases (e.g., via overlapping screening funnels and chemistry efforts). The finding that a significant proportion of DIPG contain activating ALK2 mutations suggests that ALK2 inhibitors may be of clinical benefit for DIPG patients.

[0007] Anemia of chronic disease, inflammation or cancer can develop in settings of chronic inflammatory, infectious, or neoplastic disease. In this form of anemia, inflammatory cytokines, induce hepatic expression of hepcidin, which negatively regulates iron bioavailability by inactivating ferroportin. Hepcidin is transcriptionally regulated by amongst other things bone morphogenetic protein (BMP) signaling. Inhibition of BMP phosphorylation through inhibition of ALK2 can modulate BMP-mediated signaling, thus reducing hepcidin expression. Reduced hepcidin expression may be an effective strategy for the treatment of anemia of chronic disease, inflammation, or cancer.

SUMMARY

[0008] The present invention and disclosure provides inhibitors of ALK2 and ALK2 mutants, e.g., ALK2 mutants as defined herein, for example, inhibitors of structural formula (I) and formula (Ia) and pharmaceutically acceptable salts and compositions thereof. The present invention further provides the compounds of the invention, and pharmaceutically acceptable salts and compositions thereof, for use to inhibit the activity of ALK2 or ALK2 mutants in a cell or in a patient. The present invention further provides the compounds of the invention and pharmaceutically acceptable salts and compositions thereof, for use to treat a subject or patient suffering from a condition mediated by aberrant ALK2 activity, e.g., at least one of fibrodysplasia ossificans progressiva (FOP) or heterotopic ossification or diffuse intrinsic pontine glioma (DIPG) or anemia of chronic disease or anemia of inflammation or anemia of cancer.

[0009] In one aspect, the invention features a compound of structural formula (I) or at least one pharmaceutically acceptable salt thereof:



(I)

5 wherein each of ring A, R¹, R², R³, and n is defined as described in the claims.

[0010] In another aspect, the present invention provides pharmaceutical compositions comprising a compound of structural formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

[0011] In another aspect, the present invention provides a compound for use in a method for treating or ameliorating fibrodysplasia ossificans progressiva in a subject comprising administering to the subject a therapeutically effective amount of a compound of structural formula (I) or a pharmaceutically acceptable salt or composition thereof. In an embodiment, the subject has a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification selected from one or more of L196P, PF197-8L, R202I, R206H, Q207E, R258S, R258G, G328A, G328W, G328E, G328R, G356D, and R375P.

[0012] In another aspect, the present invention provides a compound for use in a method of treating or ameliorating diffuse intrinsic pontine glioma in a subject comprising administering to the subject a therapeutically effective amount of a compound of structural formula (I) or a pharmaceutically acceptable salt or composition thereof. In an embodiment, the subject has a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification selected from one or more of R206H, G328V, G328W, G328E, and G356D.

[0013] In another aspect, the present disclosure provides a compound for use in a method of inhibiting aberrant ALK2 activity in a subject. In an embodiment, said method comprises administering to the subject a therapeutically effective amount of a compound of structural formula (I) or a pharmaceutically acceptable salt or composition thereof. In an embodiment, the subject has a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification selected from one or more of L196P, PF197-8L, R202I, R206H, Q207E, R258S, R258G, G328A, G328V, G328W, G328E, G328R, G356D, and R375P.

[0014] Various evaluation steps may be carried out prior to, during, and/or following treatment with a compound of the invention. For example, prior to, during and/or following treatment with a compound of the invention, the subject may be evaluated e.g., by visualizing heterotopic ossification in the subject. This may be achieved by spectroscopic analysis, e.g., magnetic resonance-based analysis, e.g.,

MRI, positron emission tomography (PET), micro computed tomography (μ CT), or by histology.

[0015] The evaluation may comprise evaluating a pre-treatment or baseline level of the heterotopic ossification in a subject, *e.g.*, using spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI, positron emission tomography (PET), micro computed tomography (μ CT), or by histology. The evaluation may alternatively comprise evaluating the post-treatment level of heterotopic ossification after administering to the subject a compound of the invention, *e.g.*, by using spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI, positron emission tomography (PET), micro computed tomography (μ CT), or by histology; comparing the post-treatment level of heterotopic ossification in the subject with the pre-treatment or baseline level of heterotopic ossification; and determining whether to continue treatment, *e.g.*, using spectroscopic analysis, *e.g.*, magnetic resonance-based analysis, *e.g.*, MRI, positron emission tomography (PET), micro computed tomography (μ CT), or by histology. The heterotopic ossification may be preceded by edema, *e.g.*, sustained edema.

EMBODIMENTS OF THE INVENTION AND DISCLOSURE

Definitions

[0016] As used herein, the terms a "patient," "subject," "individual," and "host" refer to either a human or a non-human animal suffering from or suspected of suffering from a disease or disorder associated with aberrant ALK2 activity (*i.e.*, aberrant ALK2 activity due to a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification) or aberrant ALK2 biological activity.

[0017] "Treat", "treatment" and "treating" such a disease or disorder refers to ameliorating at least one symptom of the disease or disorder described herein. These terms, when used in connection with a condition such as fibrodysplasia ossificans progressiva, refer to one or more of: controlling the rate of heterotopic bone growth; relieving pain and inflammation associated with development of new bone; extending the expected survival time of the patient; reducing the size or the number of heterotopic bone growth lesions; maintaining or improving mobility; preventing or treating new flare ups; inhibiting the development of new heterotopic bone lesions; enabling surgery to remove existing heterotopic ossifications to restore limb function and/or mobility; prolonging survival; prolonging progression-free survival; prolonging time to progression; inhibiting FOP related injury induced edema, and/or enhancing quality of life. When used in connection with a condition such as diffuse intrinsic pontine glioma, these terms refer to one or more of: impeding growth of the glioma, causing the glioma to shrink by weight or volume, extending the expected survival time of the patient, inhibiting glial tissue growth, reducing glial tumor mass, reducing size or number of metastatic lesions, inhibiting the development of new metastatic lesions, prolonging survival, prolonging progression-free survival, prolonging time to progression, and/or enhancing quality of life.

[0018] The term "therapeutic effect" refers to a beneficial local or systemic effect in animals, particularly mammals, and more particularly humans, caused by administration of a compound or composition of the disclosure. The phrase "therapeutically effective amount" means that amount of a compound or composition of the disclosure that is effective to treat a disease or condition associated with aberrant ALK2 activity at a reasonable benefit/risk ratio. The therapeutically effective amount of such substance will vary, for example, depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration, etc., which can readily be determined by one of skill in the art.

[0019] "Alkylene" refers to a divalent radical of an alkyl group, e.g., -CH₂-, -CH₂CH₂-, and -CH₂CH₂CH₂-.

[0020] "Alkyl" or "alkyl group" refers to a monovalent radical of a saturated straight or branched hydrocarbon, such as a straight or branched group of 1-12, 1-10, or 1-6 carbon atoms, referred to herein as C₁-C₁₂ alkyl, C₁-C₁₀ alkyl, and C₁-C₆ alkyl, respectively. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, etc.

[0021] "Aromatic" when referring to a ring is art-recognized, and refers to a fully conjugated, unsaturated ring that has $4n + 2$ π electrons and is often characterized by structural formulae showing alternating double and single bonds. Aromatic rings include both benzene and rings containing one or more heteroatoms selected from N, O and S.

[0022] "Aryl" refers to a ring system is art-recognized and refers to a monocyclic, bicyclic or polycyclic hydrocarbon ring system wherein at least one ring is aromatic.

[0023] "Halo" refers to a radical of any halogen, e.g., -F, -Cl, -Br, or -I.

[0024] "Carbocyclic ring system" refers to a monocyclic, bicyclic or polycyclic hydrocarbon ring system, wherein each ring is either completely saturated or contains one or more units of unsaturation, but where no ring is aromatic.

[0025] "Carbocyclyl" refers to a monovalent radical of a carbocyclic ring system. Representative carbocyclyl groups include cycloalkyl groups (e.g., cyclopentyl, cyclobutyl, cyclopentyl, cyclohexyl and the like), and cycloalkenyl groups (e.g., cyclopentenyl, cyclohexenyl, cyclopentadienyl, and the like).

[0026] "Cycloalkyl" refers to a cyclic, bicyclic, tricyclic, or polycyclic non-aromatic hydrocarbon groups having 3 to 12 carbons. Any substitutable ring atom may be substituted (e.g., by one or more substituents). The cycloalkyl groups can contain

fused or spiro rings. Fused rings are rings that share at least two common (carbon) atoms. Examples of cycloalkyl moieties include, but are not limited to, cyclopropyl, cyclohexyl, methylcyclohexyl, adamantyl, and norbornyl.

[0027] "Heteroalkyl" refers to a monovalent, straight or branched alkyl chain where one methylene unit other than the methylene unit bound to the rest of the molecule is replaced with -O-, -S-, or -N(R^d), wherein R^d is defined below. For the sake of clarity, the moiety -CH₂-NH-CH₃ would be a heteroalkyl, but -NH-CH₂-CH₃ would not because the -NH group is bound to the rest of the molecule.

[0028] "Heteroalkylene" refers to a divalent radical of a heteroalkyl group.

[0029] "Heteroaromatic ring system" is art-recognized and refers to a monocyclic, bicyclic or polycyclic ring system wherein at least one ring is both aromatic and comprises at least one heteroatom (e.g., N, O, or S); and wherein no other rings are heterocyclyl (as defined below). In certain instances, a ring which is aromatic and comprises a heteroatom contains 1, 2, 3, or 4 ring heteroatoms in such ring.

[0030] "Heteroaryl" refers to a monovalent radical of a heteroaromatic ring system. Representative heteroaryl groups include ring systems where (i) each ring comprises a heteroatom and is aromatic, e.g., imidazolyl, oxazolyl, thiazolyl, triazolyl, pyrrolyl, furanyl, thiophenyl, pyrazolyl, pyridinyl, pyrazinyl, pyridazinyl, pyrimidinyl, indoliziny, purinyl, naphthyridinyl, and pteridinyl; (ii) each ring is aromatic or carbocyclyl, at least one aromatic ring comprises a heteroatom and at least one other ring is a hydrocarbon ring or e.g., indolyl, isoindolyl, benzothienyl, benzofuranyl, dibenzofuranyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, isoquinolyl, cinnoliny, phthalazinyl, quinazolinyl, quinoxalinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, pyrido[2,3-b]-1,4-oxazin-3-(4H)-one, 5,6,7,8-tetrahydroquinolinyl and 5,6,7,8-tetrahydroisoquinolinyl; and (iii) each ring is aromatic or carbocyclyl, and at least one aromatic ring shares a bridgehead heteroatom with another aromatic ring, e.g., 4H-quinolizinyl.

[0031] "Heterocyclic ring system" refers to monocyclic, bicyclic and polycyclic ring systems where at least one ring is saturated or partially unsaturated (but not aromatic) and that ring comprises at least one heteroatom. A heterocyclic ring system can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure and any of the ring atoms can be optionally substituted. Heterocyclic ring systems may be fused rings.

[0032] "Heterocyclyl" refers to a monovalent radical of a heterocyclic ring system. Representative heterocyclyls include ring systems in which (i) every ring is non-aromatic and at least one ring comprises a heteroatom, e.g., tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothienyl, pyrrolidinyl, pyrrolidonyl, piperidinyl, pyrrolinyl, decahydroquinolinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholinyl, and quinuclidinyl; (ii) at least one ring is non-aromatic and comprises a heteroatom and at least one other ring is an aromatic carbon ring, e.g., 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl; and (iii) at least one ring is non-aromatic and comprises a heteroatom and at least one

other ring is aromatic and comprises a heteroatom, e.g., 3,4-dihydro-1H-pyrano[4,3-c]pyridine, and 1,2,3,4-tetrahydro-2,6-naphthyridine.

[0033] "Cyano" refers to a -CN radical.

[0034] "Hydroxy" or "hydroxyl" refers to -OH.

[0035] Certain compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present disclosure contemplates all such compounds, including cis- and trans-isomers, R- and S-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the disclosure. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this disclosure. Thus, when a disclosed compound is named or depicted by a structure without specifying the stereochemistry and has one or more chiral centers, it is understood to represent all possible stereoisomers of the compound, as well as enantiomeric mixtures thereof. When a disclosed compound is named or depicted by a structure specifying stereochemistry at each chiral center, it is understood to represent only the compound having the designated stereochemistry at such chiral centers. However, when a disclosed compound specifies stereochemistry at some, but not all chiral centers, it is understood to represent all possible stereoisomers at the non-specified chiral centers of the compound, as well as enantiomeric mixtures thereof.

[0036] If, for instance, a particular enantiomer of compound of the present disclosure is desired, it may be prepared by asymmetric synthesis, or by derivation with a chiral auxiliary, where the resulting diastereomeric mixture is separated and the auxiliary group cleaved to provide the pure desired enantiomers. Alternatively, where the molecule contains a basic functional group, such as amino, or an acidic functional group, such as carboxyl, diastereomeric salts are formed with an appropriate optically-active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means well known in the art, and subsequent recovery of the pure enantiomers.

[0037] The "enantiomeric excess" or "% enantiomeric excess" of a composition can be calculated using the equation shown below. In the example shown below a composition contains 90% of one enantiomer, e.g., the S enantiomer, and 10% of the other enantiomer, i.e., the R enantiomer.

$$ee = (90-10)/100 = 80\%.$$

[0038] Thus, a composition containing 90% of one enantiomer and 10% of the other enantiomer is said to have an enantiomeric excess of 80%.

[0039] The compounds or compositions described herein may contain an enantiomeric excess of at least 50%, 75%, 90%, 95%, or 99% of one form of the compound, e.g., the S-enantiomer. In other words such compounds or compositions contain an enantiomeric excess of the S enantiomer over the R enantiomer.

[0040] The compounds described herein may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example deuterium (^2H), tritium (^3H), carbon-13 (^{13}C), or carbon-14 (^{14}C). All isotopic variations of the compounds disclosed herein, whether radioactive or not, are intended to be encompassed within the scope of the present disclosure. In addition, all tautomeric forms of the compounds described herein are intended to be within the scope of the claimed disclosure.

[0041] The compound can be useful as the free base or as a salt. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthalate, mesylate, glucoheptonate, lactobionate, and laurylsulfonate salts and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", J. Pharm. Sci. 66:1-19.)


[0042] As described herein, compounds of the invention may contain "optionally substituted" moieties. In general, the term "substituted", whether preceded by the term "optionally" or not, means that one or more hydrogens of the designated moiety are replaced with a suitable substituent. Unless otherwise indicated, an "optionally substituted" group may have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at each position. Combinations of substituents envisioned under this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and use for one or more of the purposes disclosed herein.

[0043] Suitable substituents for an optionally substituted alkyl, alkylene, carbocyclyl, heterocyclyl, aryl group and heteroaryl group include halogen, =O, -CN, -OR^c, -NR^dR^e, -S(O)_kR^c, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -OC(=O)OR^c, -OC(=O)R^c, -OC(=S)OR^c, -C(=S)OR^c, -O(C=S)R^c, -C(=O)NR^dR^e, -NR^cC(=O)R^c, -C(=S)NR^dR^e, -NR^cC(=S)R^c, -NR^c(C=O)OR^c, -O(C=O)NR^dR^e, -NR^c(C=S)OR^c, -O(C=S)NR^dR^e, -NR^c(C=O)NR^dR^e, -NR^c(C=S)NR^dR^e, -C(=S)R^c, -C(=O)R^c, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ heteroalkyl, carbocyclyl, (C₁-C₆-alkylene)-carbocyclyl, (C₁-C₆-heteroalkylene)-carbocyclyl, heterocyclyl, (C₁-C₆-alkylene)-heterocyclyl, (C₁-C₆-heteroalkylene)-heterocyclyl, aryl, (C₁-C₆-alkylene)-aryl, (C₁-C₆-heteroalkylene)-aryl, heteroaryl, (C₁-C₆-alkylene)-heteroaryl, or (C₁-C₆-heteroalkylene)-heteroaryl, wherein each of said alkyl, alkylene, heteroalkyl, heteroalkylene, carbocyclyl, heterocyclyl, aryl, and heteroaryl are optionally substituted with one or more of halogen, OR^c, -NO₂, -CN, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆ alkyl, C₁-C₆ haloalkyl, or C₁-C₆ heteroalkyl, and wherein R^c is hydrogen, hydroxy, C₁-C₆ alkyl, C₁-C₆ heteroalkyl, carbocyclyl, (C₁-C₆-alkylene)-carbocyclyl, (C₁-C₆-heteroalkylene)-carbocyclyl, heterocyclyl, (C₁-C₆-alkylene)-heterocyclyl, (C₁-C₆-heteroalkylene)-heterocyclyl, aryl, (C₁-C₆-alkylene)-aryl, (C₁-C₆-

heteroalkylene)-aryl, heteroaryl, (C₁-C₆-alkylene)-heteroaryl, or (C₁-C₆-heteroalkylene)-heteroaryl, each of which is optionally substituted with one or more of halogen, hydroxy, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ heteroalkyl, carbocyclyl, heterocyclyl, aryl, or heteroaryl; R^d and R^e are each independently selected from hydrogen, C₁-C₆ alkyl, or C₁-C₆ heteroalkyl; and k is 0, 1, or 2. The claimed disclosure is not intended to be limited in any manner by the above exemplary listing of substituents.

Compounds

[0044] In one aspect, the present invention features a compound having the structural formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

ring A is phenyl or heteroaryl, wherein ring A is optionally substituted with 1, 2, or 3 substituents independently selected from the list given in claim 1 in addition to R²;

R¹ is selected from NH(C₁-C₆ alkyl), N(C₁-C₆ alkyl)₂, C₁-C₆ alkyl, -O-C₁-C₆ alkyl, -C(O)-C₁-C₄ alkyl, carbocyclyl, heterocyclyl, -O-(C₀-C₄ alkylene)-carbocyclyl, -O-(C₀-C₄ alkylene)-heterocyclyl, -NH-(C₀-C₄ alkylene)-carbocyclyl, -NH-aryl, NH-O-(C₁-C₄ alkyl), -S-heterocyclyl, -S-(C₀-C₃ alkylene)-(O-containing heterocyclyl), -NH-(C₀-C₄ alkylene)-heterocyclyl, wherein each alkyl, alkylene, carbocyclyl, aryl and heterocyclyl portion of R¹ is optionally substituted with 1, 2, 3, or 4 independently selected substituents independently selected from the list in claim 1; or

R¹ is taken together with one R³ to form a saturated ring fused to the piperazine ring in formula (I), and wherein the ring formed by R¹ and R³ is optionally substituted with 1, 2, or 3 substituents independently selected from the list in claim 1;

R² is selected from halo, C₁-C₆ alkyl, heterocyclyl, cycloalkyl, -NH-(C₀-C₄ alkylene)-heterocyclyl, -(C₁-C₄ alkylene)-heterocyclyl, -(C₁-C₄ alkylene)-NH-heterocyclyl, and -O-(C₀-C₄ alkylene)-heterocyclyl, wherein any heterocyclyl, cycloalkyl, alkyl, or alkylene portion of R² is optionally substituted with 1, 2, 3, or 4 substituents independently selected from the list in claim 1;

or

R² is taken together with any ring atom in ring A to form a cycloalkyl or saturated heterocyclyl ring that is fused to ring A, and wherein the ring formed by R² and the ring atom in ring A is optionally substituted with 1, 2 or 3 substituents independently selected from the list in claim 1; or

R² is taken together with any saturated ring atom in ring A to form a cycloalkyl or saturated heterocyclyl ring that is fused, spirofused, or bridged to ring A, and wherein the ring formed by R² and the ring atom in ring A is optionally

substituted with 1, 2, or 3 substituents independently selected from the list in claim 1;
 each R³ is independently selected from C₁-C₄ alkyl and C₁-C₄ haloalkyl;
 and
 n is 0, 1, 2, or 3.

[0045] In an embodiment, the compound is a compound of Formula (I) or a pharmaceutically acceptable salt thereof, wherein:

ring A is phenyl or heteroaryl, wherein ring A is optionally substituted with 1, 2, or 3 substituents independently selected from the list in claim 1, in addition to R²;

R¹ is selected from NH(C₁-C₆ alkyl), N(C₁-C₆ alkyl)₂, C₁-C₆ alkyl, -O-C₁-C₆ alkyl, -C(O)-C₁-C₄ alkyl, carbocyclyl, heterocyclyl, -O-(C₀-C₄ alkylene)-carbocyclyl, -O-(C₀-C₄ alkylene)-heterocyclyl, -NH-(C₀-C₄ alkylene)-carbocyclyl, and -NH-(C₀-C₄ alkylene)-heterocyclyl, wherein each alkyl, alkylene, carbocyclyl, and heterocyclyl portion of R¹ is optionally substituted with 1, 2, 3, or 4 substituents independently selected from the list in claim 1;

or

R¹ is taken together with one R³ to form a saturated ring fused to the piperazine ring in formula (I), and wherein the ring formed by R¹ and R³ is optionally substituted with 1, 2, or 3 substituents independently selected from the list in claim 1;

R² is selected from halo, C₁-C₆ alkyl, heterocyclyl, cycloalkyl, -NH-(C₀-C₄ alkylene)-heterocyclyl, and -O-(C₀-C₄ alkylene)-heterocyclyl, wherein any heterocyclyl, cycloalkyl, alkyl, or alkylene portion of R² is optionally substituted with 1, 2, 3, or 4 substituents independently selected from the list in claim 1;


or


R² is taken together with any saturated ring atom in ring A to form a cycloalkyl or saturated heterocyclyl ring that is fused, spirofused, or bridged to ring A, and wherein the ring formed by R² and the ring atom in ring A is optionally substituted with 1, 2, or 3 substituents independently selected from the list in claim 1;

each R³ is independently selected from C₁-C₄ alkyl and C₁-C₄ haloalkyl; and
 n is 0, 1, 2, or 3.

[0046] In an embodiment, n is 0; or n is 1 and R³ is selected from methyl, ethyl, and -CHF₂.

[0047] In an embodiment, ring A is selected from:







wherein:

- 5 "1" represents a portion of ring A bound to a pyrrolo[1,2-b]pyridazine moiety;
 "2" represents a portion of ring A bound to R²; and
 ring A is optionally substituted with 1, 2, or 3 substituents independently selected
 from the list in claim 3 in addition to R².

- 10 **[0048]** In an embodiment, ring A is selected from:



and



, wherein:

- 15 "1" represents a portion of ring A bound to a pyrrolo[1,2-b]pyridazine moiety;
 "2" represents a portion of ring A bound to R²; and
 ring A is substituted with 0, 1, 2, or 3 independently selected substituents in addition
 20 to R².

[0049] In an embodiment, ring A is selected from phenyl and pyridinyl. In one aspect
 of this embodiment, ring A is selected from phenyl and pyridin-2-yl.

- 25 **[0050]** In an embodiment, ring A is optionally substituted with 1, or 2 substituents in
 addition to R², wherein each optional substituent is independently selected from
 halo, methyl, -cyano, and -OCHF₂.

- 30 **[0051]** In an embodiment, R¹ is selected from -C(O)-(C₁-C₃ alkyl), C₁-C₃ alkyl, -O-
 (C₁-C₅ alkyl), -NH(C₁-C₅ alkyl), -N(C₁-C₄ alkyl)₂, -NH-(C₃-C₆ cycloalkyl), C₃-
 C₆ cycloalkyl, -O-(C₃-C₆ cycloalkyl), -O-(C₁-C₃ alkylene)-(C₃-C₆ cycloalkyl), -(C₁-
 C₃ alkylene)-(C₃-C₆ cycloalkyl), -O-(C₀-C₃ alkylene)-(O-containing heterocyclyl), -NH-
 (C₀-C₃ alkylene)-(O-containing heterocyclyl), an O-containing heterocyclyl, and an N-
 containing heterocyclyl, -O-(C₀-C₃ alkylene)-(N-containing heterocyclyl), -S-(C₀-
 35 C₃ alkylene)-(O-containing heterocyclyl), -NH-O-(C₁-C₃ alkyl), and -NH-phenyl,
 wherein any alkyl, alkylene, cycloalkyl, phenyl or heterocyclyl portion of R¹ is

optionally substituted with 1, 2, or 3 substituents independently selected from deuterium, halo, cyano, acetyl, C₁-C₄ alkyl, C₁-C₄ haloalkyl, -O-C₁-C₄ alkyl, -C₁-C₄ alkylene-O-C₁-C₄ alkyl, heteroaryl, phenyl, cycloalkyl, -COOH and hydroxy, wherein each of said heteroaryl, phenyl, and cycloalkyl are optionally substituted with one or more of halo, OR^c, -NO₂, cyano, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆ alkyl, C₁-C₆ haloalkyl, or C₁-C₆ heteroalkyl; or R¹ is taken together with any ring atom in the piperazine moiety of formula (I) to form a carbocyclyl or heterocyclyl ring fused to the piperazine moiety.

[0052] In some embodiments, R¹ is selected from -O-(C₀-C₃ alkylene)-(N-containing heterocyclyl), -S-(C₀-C₃ alkylene)-(O-containing heterocyclyl), -NH-O-(C₁-C₃ alkyl), and -NH-phenyl, wherein any alkyl, alkylene, phenyl, or heterocyclyl portion of R¹ is unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from deuterium, halo, cyano, acetyl, C₁-C₄ alkyl, C₁-C₄ haloalkyl, -O-C₁-C₄ alkyl, -C₁-C₄ alkylene-O-C₁-C₄ alkyl, optionally substituted heteroaryl, optionally substituted phenyl, optionally substituted cycloalkyl, -COOH, and -OH.

[0053] In an embodiment, R¹ is selected from -C(O)-(C₁-C₃ alkyl), C₁-C₃ alkyl, -O-(C₁-C₃ alkyl), -NH(C₁-C₃ alkyl), -N(C₁-C₄ alkyl)₂, -NH-(C₃-C₆ cycloalkyl), C₃-C₆ cycloalkyl, -O-(C₃-C₆ cycloalkyl), -O-(C₁-C₃ alkylene)-(C₃-C₆ cycloalkyl), -(C₁-C₃ alkylene)-(C₃-C₆ cycloalkyl), and an N-containing heterocyclyl, wherein any alkyl, cycloalkyl, or heterocyclyl portion of R¹ is optionally substituted with 1, 2, or 3 substituents independently selected from halo, cyano, acetyl, C₁-C₄ alkyl, C₁-C₄ haloalkyl, -O-C₁-C₄ alkyl, -C₁-C₄ alkylene-O-C₁-C₄ alkyl, heteroaryl, phenyl, cycloalkyl, and hydroxy; wherein each of said heteroaryl, phenyl and cycloalkyl are optionally substituted with one or more of the substituents listed in claim 4; or R¹ is taken together with any ring atom in the piperazine moiety of formula (I) to form a carbocyclyl or heterocyclyl ring fused to the piperazine moiety.

[0054] In an embodiment, R¹ is selected from 1-(3-chlorophenyl)cyclopropyl, 1-(3-fluorophenyl)cyclopropyl, 1-acetylcyclopropyl, 1-cyclopropylcyclopropyl, 1-difluoromethylcyclopropyl, 1-fluorocyclopropyl, 1-methylpropylamino, 1-pyridin-3-ylcyclopropyl, 1-thiazol-2-ylcyclopropyl, 1-thien-2-ylcyclopropyl, 1-trifluoromethylcyclopropyl, 2-(4-chlorophenyl)cyclopropyl, 2,2,2-trifluoroethoxy, 2,2-difluorocyclopropyl, 2,2-dimethylcyclopropyl, 2-cyanocyclopropyl, 2-cyanoethyl, 2-cyanoethylamino, 2-cyclobutylcyclopropyl, 2-fluorocyclopropyl, 2-fluoroethoxy, 2-hydroxyethylamino, 2-methoxyethoxy, 2-methylcyclopropyl, 2-oxa-6-azaspiro[3.3]heptan-6-yl, 3,3-difluorocyclobutyl, 3-cyanoazetidin-1-yl, 3-cyanocyclobutyl, 3-fluorocyclobutyl, 3-hydroxy-3-methylcyclobutyl, 3-hydroxy-3-trifluoromethylcyclobutyl, 3-hydroxyazetidin-1-yl, 3-hydroxycyclobutyl, 3-methoxyazetidin-1-yl, 3-methoxymethylazetidin-1-yl, 3-phenyl-3-hydroxycyclobutyl, 4-cyanocyclohexyl, 4-cyanoypiperidin-1-yl, 4-hydroxy-4-methylcyclohexyl, 4-hydroxycyclohexyl, 4-hydroxypiperidin-1-yl, 4-methylcyclohexyl, acetyl, azetidin-1-yl, cyclobutoxy, cyclobutyl, cyclobutylamino, cyclopentylamino, cyclopropyl, cyclopropylmethyl, diethylamino, ethoxy, ethyl, ethylamino, isobutoxy, isopropoxy, isopropyl, isopropylamino, methoxymethyl, N-ethyl-N-methylamino, pentylamino, piperidin-1-yl, propylamino, propyloxypyrrolidin-1-yl, t-butylamino, t-butoxy, 2,2-dimethylpropoxy, 2,2-difluoroethoxy, N-(2,2-dimethylpropyl)amino, N-(1,2-

dimethylpropyl)amino, 2,2,2-trifluoroethylamino, N-(methoxymethyl)amino, oxetan-3-yloxy, oxetan-3-yl, oxetan-3-ylamino, oxetan-3-ylmethoxy, N-(oxetan-3-ylmethyl)amino, tetrahydrofuran-3-yloxy, tetrahydropyran-4-yloxy, and 3-cyanocyclobutoxy, or R¹ is taken together with a ring atom in the piperazine moiety to form a 6-oxohexahydropyrrolo[1,2-a]pyrazin-2-yl, or 2-ethyl-3-oxohexahydroimidazo[1,5-a]pyrazin-7-yl.

[0055] In an embodiment, R¹ is selected from 1,3-dihydroxypropan-2-yloxy, 1-acetylazetidin-3-yloxy, 1-hydroxy-2-hydroxycarbonylethan-2-yloxy, 1-methyl-2-fluoroethoxy, 2,2-difluoroethylamino, 2-cyanoethan-1-yloxy, 2-fluoroethylamino, 2-fluorophenylamino, 2-fluoropropoxy, 2-methyloxetan-3-yloxy, 3-cyano-oxetan-3-yloxy, 3-deutero-oxetan-3-yloxy, 6-oxa-1-azaspiro[3.3]heptan-1-yl, cyclopropoxy, ethoxyamino, oxetan-3-ylthio, perdeuteroethoxy, phenylamino, tetrahydrofuran-2-yloxy, and tetrahydrofuran-3-yl. In an embodiment, R¹ is selected from 1-(3-chlorophenyl)cyclopropyl, 1-(3-fluorophenyl)cyclopropyl, 1-acetylcyclopropyl, 1-cyclopropylcyclopropyl, 1-difluoromethylcyclopropyl, 1-fluorocyclopropyl, 1-methylpropylamino-1-pyridin-3-ylcyclopropyl, 1-thiazol-2-ylcyclopropyl, 1-thien-2-ylcyclopropyl, 1-trifluoromethylcyclopropyl, 2-(4-chlorophenyl)cyclopropyl, 2,2,2-trifluoroethoxy, 2,2-difluorocyclopropyl, 2,2-dimethylcyclopropyl, 2-cyanocyclopropyl, 2-cyanoethyl, 2-cyanoethylamino, 2-cyclobutylcyclopropyl, 2-fluorocyclopropyl, 2-fluoroethoxy, 2-hydroxyethylamino, 2-methoxyethoxy, 2-methylcyclopropyl, 2-oxa-6-azaspiro[3.3]heptan-6-yl, 3,3-difluorocyclobutyl, 3-cyanoazetidin-1-yl, 3-cyanocyclobutyl, 3-fluorocyclobutyl, 3-hydroxy-3-methylcyclobutyl, 3-hydroxy-3-trifluoromethylcyclobutyl, 3-hydroxyazetidin-1-yl, 3-hydroxycyclobutyl, 3-methoxyazetidin-1-yl, 3-methoxymethylazetidin-1-yl, 3-phenyl-3-hydroxycyclobutyl, 4-cyanocyclohexyl, 4-cyanoypiperidin-1-yl, 4-hydroxy-4-methylcyclohexyl, 4-hydroxycyclohexyl, 4-hydroxypiperidin-1-yl, 4-methylcyclohexyl, acetyl, azetidin-1-yl, cyclobutoxy, cyclobutyl, cyclobutylamino, cyclopentylamino, cyclopropyl, cyclopropylmethyl, diethylamino, ethoxy, ethyl, ethylamino, isobutoxy, isopropoxy, isopropyl, isopropylamino, methoxymethyl, N-ethyl-N-methylamino, pentylamino, piperidin-1-yl, propylamino, propyloxypyrrolidin-1-yl, and t-butylamino, or R¹ is taken together with a ring atom in the piperazine moiety to form a 6-oxohexahydropyrrolo[1,2-a]pyrazin-2-yl, or 2-ethyl-3-oxohexahydroimidazo[1,5-a]pyrazin-7-yl.

[0056] In an embodiment, R² is selected from halo, cycloalkyl, heterocyclyl, -O-(C₀-C₄ alkylene)-(heterocyclyl), -(C₁-C₃ alkylene)-heterocyclyl, -(C₁-C₃ alkylene)-NH-(C₁-C₃ alkyl), -(hydroxy-substituted C₁-C₃ alkylene)-NH-(C₁-C₃ alkyl), C₁-C₃ alkyl substituted with both hydroxy and amino, and cyano-substituted C₁-C₄ alkyl, C₁-C₄ alkyl substituted with both hydroxy and one or more of amino, C₁-C₄ alkylamino or di-C₁-C₄ alkylamino cyano-substituted C₁-C₄ alkyl, hydroxy, -S(O)₂-C₁-C₄ alkyl, and -(amino substituted C₁-C₃ alkylene)-heterocyclyl, or R² is taken together with a ring atom in ring A to form a heterocyclyl or a carbocyclyl that is fused to ring A, wherein any heterocyclyl, or carbocyclyl is optionally substituted with 1, 2 or 3 substituents independently selected from halo, cyano, hydroxy, -NH₂, -NH(C₁-C₄ alkyl), -NH-C(O)-O-(C₁-C₄ alkyl), =O, -OH, -C(O)-C₁-C₄ alkyl, -C₁-C₄ alkyl, deuterated C₁-C₄ alkyl, -C₁-C₄ haloalkyl, hydroxy-substituted -C₁-C₄ alkyl, -O-C₁-C₄ alkyl, -(C₁-C₄ alkylene)-O-(C₁-C₄ alkyl), -(C₁-C₄ alkylene)-O-(C₁-C₄ haloalkyl), -C(O)-O-C₁-

C₄ alkyl, -COOH, C₃-C₆ cycloalkyl, heterocyclyl, or -NH heterocyclyl, wherein each said heterocycle is optionally substituted with one or more of halo, -OR^c, NO₂, cyano, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆ alkyl, C₁-C₆ haloalkyl, or C₁-C₆ heteroalkyl.

5

[0057] In an embodiment R² is selected from -OH, -S(O)₂-C₁-C₄ alkyl, -(amino substituted C₁-C₃ alkylene)-heterocyclyl, C₄ alkyl, C₁-C₃ alkyl substituted with both hydroxy and either C₁-C₄ alkylamino or di-C₁-C₄ alkylamino.

10

[0058] In an embodiment, R² is selected from halo, cycloalkyl, heterocyclyl, -O-(C₀-C₄ alkylene)-(heterocyclyl), and cyano-substituted C₁-C₄ alkyl, or R² is taken together with a ring atom in ring A to form a heterocyclyl that is fused to ring A, wherein any heterocyclyl is optionally substituted with 1, 2 or 3 substituents independently selected from halo, -NH₂, =O, -OH, -C(O)-C₁-C₄ alkyl, -C₁-C₄ alkyl, deuterated C₁-C₄ alkyl, -C₁-C₄ haloalkyl, hydroxy-substituted -C₁-C₄ alkyl, -O-C₁-C₄ alkyl, -(C₁-C₄ alkyl)-O-(C₁-C₄ alkyl), C₃-C₆ cycloalkyl, an optionally substituted second heterocyclyl, or -NH-(optionally substituted second heterocyclyl).

15

[0059] In an embodiment, R² is selected from 1-(1-hydroxypropan-2-yl)-4-methoxypiperidin-4-yl, 1-(3-difluoromethoxy)propan-2-yl-4-methoxypiperidin-4-yl, 1-(3-methoxy)propan-2-yl-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)piperidin-4-yl, 1-(propan-2-yl)piperidin-3-yl, 1-(propan-2-yl)piperidin-4-yl, 1-(pyrrolidin-1-yl)ethan-1-yl, 1,2,3,6-tetrahydropyridin-4-yl, 1,4-diazabicyclo[4.2.0]octan-4-yl, 1-acetylpiperidin-4-yl, 1-cyclobutylpiperidin-3-yl, 1-ethyl-3,3-difluoropiperidin-4-yl, 1-ethyl-3-fluoropiperidin-4-yl, 1-ethyl-3-hydroxyazetidin-3-yl, 1-ethyl-4-fluoropyrrolidin-3-yl, 1-ethylazetidin-3-yl, 1-ethylazetidin-3-yloxy, 1-ethylpiperidin-3-yl, 1-ethylpiperidin-3-yloxy, 1-ethylpiperidin-4-yl, 1-ethylpiperidin-4-yloxy, 1-ethylpyrrolidin-3-yl, 1-ethylpyrrolidin-3-ylmethoxy, 1-ethylpyrrolidin-3-yloxy, 1H-pyrrolidin-2-yl, 1-hydroxy-2-aminoprop-2-yl, 1-isopropyl-1,2,3,6-tetrahydropyridin-4-yl, 1-isopropyl-2-methylpyrrolidin-2-yl, 1-isopropyl-3,4-dimethylpiperazin-3-yl, 1-isopropyl-3-ethoxypiperidin-3-yl, 1-isopropyl-3-fluoropiperidin-3-yl, 1-isopropyl-3-hydroxyazetidin-3-yl, 1-isopropyl-3-hydroxypiperidin-3-yl, 1-isopropyl-3-hydroxypyrrolidin-3-yl, 1-isopropyl-3-methoxyazetidin-3-yl, 1-isopropyl-3-methoxypiperidin-3-yl, 1-isopropyl-3-methoxypyrrolidin-3-yl, 1-isopropyl-3-methylpiperazin-3-yl, 1-isopropyl-4-cyanopiperidin-4-yl, 1-isopropyl-4-ethoxypiperidin-4-yl, 1-isopropyl-4-fluoropiperidin-4-yl, 1-isopropyl-4-fluoropyrrolidin-3-yl, 1-isopropyl-4-hydroxypiperidin-3-yl, 1-isopropyl-4-hydroxypiperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-isopropyl-4-methylpiperazin-3-yl, 1-isopropyl-4-methylpiperidin-4-yl, 1-isopropyl-4-trifluoromethylpiperidin-4-yl, 1-isopropyl-5-methylpyrrolidin-3-yl, 1-isopropylazetidin-2-ylmethoxy, 1-isopropylazetidin-3-yl, 1-isopropylazetidin-3-ylmethoxy, 1-isopropylazetidin-3-yloxy, 1-isopropylpiperazin-3-yl, 1-isopropylpiperazin-4-yl, 1-isopropylpiperidin-2-yl, 1-isopropylpiperidin-3-yl, 1-isopropylpiperidin-4-yl, 1-isopropylpyrrolidin-2-yl, 1-isopropylpyrrolidin-3-yl, 1-methyl-1-cyanoethyl, 1-sec-butylpiperidin-4-yl, 1-t-butoxycarbonyl-4-aminopiperidin-4-yl, 2-(isopropylamino)-3-hydroxypropan-2-yl, 2-(isopropylamino)-propan-2-yl, 2,3,5,6-tetrahydroimidazo[2,1-b]thiazol-6-yl, 2,6-diazaspiro[3.3]heptan-2-yl, 2,6-diazaspiro[3.4]octan-2-yl, 2,6-diazaspiro[3.4]octan-6-yl, 2,7-diazaspiro[4.4]nonan-2-yl, 2-difluoromethylpiperazin-1-

45

yl, 2-isopropyl-2,6-diazaspiro[3.3]heptan-6-yl, 2-methyl-1H-pyrrolidin-2-yl, 2-oxa-5,8-diazaspiro[3.5]nonan-8-yl, 2-oxo-4-ethylpiperazin-1-yl, 2-oxopiperazin-1-yl, 2-trifluoromethylpiperazin-1-yl, 3,3-difluoropiperidin-4-yl, 3,3-dimethyl-4-ethylpiperazin-1-yl, 3-aminopyrrolidin-1-yl, 3-fluoropiperidin-3-yl, 3-fluoropiperidin-4-yl, 3-hydroxyazetidin-3-yl, 3-hydroxyquinuclidin-3-yl, 3-methyl-4-ethylpiperazin-1-yl, 3-methylpiperazin-1-yl, 3-trifluoromethylpiperazin-1-yl, 4-(1,1,2,2,2-pentadeuteroethyl)piperazin-1-yl, 4-(2,2-difluoroethyl)piperazin-1-yl, 4-(2-hydroxyethyl)piperazin-1-yl, 4-(2-methoxyethyl)piperazin-1-yl, 4-(methoxycarbonylamino)piperidin-4-yl, 4-(oxetan-3-yl)piperazin-1-yl, 4,5-dihydro-1H-imidazol-2-yl, 4,5-dihydro-1H-imidazol-2-ylamino, 4-aminopiperidin-1-yl, 4-cyanopiperidin-4-yl, 4-ethoxypiperidin-4-yl, 4-ethylmorpholin-2-yl, 4-ethylpiperazin-1-yl, 4-ethylpiperazin-1-ylethoxy, 4-fluoropiperidin-4-yl, 4-fluoropyrrolidin-3-yl, 4-hydroxy-tetrahydro-2H-pyran-4-yl, 4-isopropylmorpholin-3-yl, 4-isopropylpiperazin-1-yl, 4-methoxypiperidin-4-yl, 4-methylpiperazin-1-yl, 4-methylpiperidin-4-yl, 5,5-difluoropiperidin-3-yl, 5-ethyl-2,5-diazabicyclo[2.2.1]heptan-2-yl, 6-ethyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-isopropyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-methylmorpholin-2-yl, azetidin-2-ylmethoxy, azetidin-3-yl, bromo, cyclopentyl, hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl, hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, morpholin-2-yl, morpholin-3-yl, octahydro-2H-pyrido[1,2-a]pyrazin-2-yl, piperazin-1-yl, piperazin-1-ylethoxy, piperidin-4-yl, piperidin-2-yl, piperidin-3-yl, piperidin-3-yloxy, piperidin-4-yloxy, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrrolidin-3-ylmethoxy, pyrrolidin-3-yloxy, quinuclidin-4-yl, tetrahydro-2H-pyran-4-yl, or R² is taken together with ring A to form 3'H-spiro[azetidine-3,1'-isobenzofuran]-5'-yl, 6-isopropyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[4,5-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 5,6,7,8-tetrahydroimidazo[1,5-a]pyrazin-3-yl, 7-methyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyrazin-3-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, or 5-isopropyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 1-amino-2,3-dihydro-1H-inden-5-yl, or 1-(isopropylamino)-2,3-dihydro-1H-inden-5-yl.

30


[0060] In an embodiment, R² is selected from 1-(1-fluoropropan-2-yl)-4-ethoxypiperidin-4-yl, 1-(1-fluoropropan-2-yl)-4-methoxypiperidin-4-yl, 1-(2-fluoropropyl)-4-ethoxypiperidin-4-yl, 1-(2-fluoropropyl)-4-methoxypiperidin-4-yl, 1-(3-difluoromethoxy)propan-2-yl)-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)-4-ethoxypiperidin-4-yl, 1-(tetrahydrofuran-2-yl)-1-aminomethyl, 1-amino-2-hydroxy-2-methylpropyl, 1-amino-2-methoxyethyl, 1-azabicyclo[2.2.1]heptan-4-yl, 1-cyclopropyl-4-ethoxypiperidin-4-yl, 1-diethylamino-2-hydroxyethyl, 1-ethylamino-2-hydroxyethyl, 1-isopropyl-4-difluoromethoxypiperidin-3-yl, 1-isopropyl-4-difluoromethoxypiperidin-4-yl, 1-isopropyl-4-hydroxymethylpiperidin-4-yl, 1-isopropyl-4-methoxycarbonylpiperidin-4-yl, 1-isopropyl-4-(methoxymethyl)piperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-methyl-1-isopropylamino-2-hydroxyethyl, 2,2,5,5-tetramethyl-4-hydroxypiperidin-4-yl, 2,2-dimethyl-4-methoxypiperidin-4-yl, 2-amino-1-hydroxyethyl, 2-amino-3-hydroxypropyl, 2-azaspiro[3.3]heptan-6-yl, 2-hydroxy-1-aminoethyl, 2-hydroxy-1-isopropylaminoethyl, 2-hydroxyethylaminomethyl, 3-amino-oxetan-3-yl, 3-ethoxypiperidin-3-yl, 3-methoxypiperidin-3-yl, 4-amino-tetrahydropyran-4-yl, 4-ethoxytetrahydropyran-4-yl, 4-hydroxycarbonylpiperidin-4-yl, 4-hydroxymethylpiperidin-4-yl, 4-

45

methoxycarbonylpiperidin-4-yl, 4-methoxytetrahydropyran-4-yl, 4-trifluoromethylpiperidin-4-yl, ethylsulfonyl, and oxetan-3-ylaminomethyl.


[0061] In an embodiment, R² is selected from 1-(oxetan-3-yl)piperidin-4-yl, 1-acetylpiperidin-4-yl, 1-cyclobutylpiperidin-3-yl, 1-ethyl-3-fluoropiperidin-4-yl, 1-ethyl-3-hydroxyazetidin-3-yl, 1-ethyl-4-fluoropyrrolidin-3-yl, 1-ethylazetidin-3-yl, 1-ethylpiperidin-3-yl, 1-ethylpiperidin-4-yl, 1-ethylpyrrolidin-3-yl, 1-isopropyl-3-hydroxyazetidin-3-yl, 1-isopropyl-3-hydroxypiperidin-3-yl, 1-isopropylazetidin-3-yl, 1-isopropylpiperidin-3-yl, 1-isopropylpiperidin-4-yl, 1-isopropylpyrrolidin-2-yl, 1-isopropylpyrrolidin-3-yl, 1-methyl-1-cyanoethyl, 2-difluoromethylpiperazin-1-yl, 2-oxo-4-ethylpiperazin-1-yl, 2-oxopiperazin-1-yl, 2-trifluoromethylpiperazin-1-yl, 3-fluoropiperidin-3-yl, 3-fluoropiperidin-4-yl, 3-hydroxyazetidin-3-yl, 3-methyl-4-ethylpiperazin-1-yl, 3-methylpiperazin-1-yl, 3-trifluoromethylpiperazin-1-yl, 4-(2-hydroxyethyl)-piperazin-1-yl, 4-(oxetan-3-yl)piperazin-1-yl, 4-aminopiperidin-1-yl, 4-ethylmorpholin-2-yl, 4-ethylpiperazin-1-yl, 4-fluoropyrrolidin-3-yl, 4-hydroxy-tetrahydro-2H-pyran-4-yl, 4-isopropylpiperazin-1-yl, 4-(2-methoxyethyl)piperazin-1-yl, 4-methylpiperazin-1-yl, 6-methylmorpholin-2-yl, azetidin-3-yl, morpholin-2-yl, piperazin-1-yl, piperidin-4-yl, piperidin-3-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, tetrahydro-2H-pyran-4-yl, azetidin-2-ylmethoxy, pyrrolidin-3-yloxy, piperidin-2-yl, piperidin-3-yloxy, piperidin-4-yloxy, pyrrolidin-3-ylmethoxy, 1-ethylazetidin-3-yloxy, 1-isopropylazetidin-3-yloxy, 1-ethylpyrrolidin-3-yloxy, 1-isopropylazetidin-3-ylmethoxy, 1-isopropylazetidin-2-ylmethoxy, piperazin-1-ylethoxy, 4-(2-hydroxyethyl)piperazin-1-yl, 1-isopropyl-4-fluoropyrrolidin-3-yl, 1-ethylpiperidin-4-yloxy, 1-ethylpiperidin-3-yloxy, 1-ethylpyrrolidin-3-ylmethoxy, 1-isopropyl-3-hydroxypyrrolidin-3-yl, 1-isopropylpiperidin-2-yl, 1-isopropyl-4-hydroxypiperidin-4-yl, 4-ethylpiperazin-1-ylethoxy, 1-sec-butylpiperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-isopropyl-3-methoxypiperidin-3-yl, bromo, cyclopentyl, 1,4-diazabicyclo[4.2.0]octan-4-yl, 5-ethyl-2,5-diazabicyclo[2.2.1]heptan-2-yl, 4-(1,1,2,2-pentadeuteroethyl)piperazin-1-yl, 1-ethyl-3,3-difluoropiperidin-4-yl, hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, octahydro-2H-pyrido[1,2-a]pyrazin-2-yl, hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl, 4-(2,2-difluoroethyl)piperazin-1-yl, 2,6-diazaspiro[3.3]heptan-2-yl, 3,3-difluoropiperidin-4-yl, 3,3-dimethyl-4-ethylpiperazin-1-yl, 6-ethyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-isopropyl-2,6-diazaspiro[3.3]heptan-2-yl, 2,6-diazaspiro[3.4]octan-6-yl, 2,6-diazaspiro[3.4]octan-2-yl, 2-oxa-5,8-diazaspiro[3.5]nonan-8-yl, 2,7-diazaspiro[4.4]nonan-2-yl, 4,5-dihydro-1H-imidazol-2-yl, 4,5-dihydro-1H-imidazol-2-ylamino, and 5,5-difluoropiperidin-3-yl, or R² is taken together with ring A to form 3'H-spiro[azetidine-3,1'-isobenzofuran]-5'-yl, 6-isopropyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[4,5-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 5,6,7,8-tetrahydroimidazo[1,5-a]pyrazin-3-yl, 7-methyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyrazin-3-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, or 5-isopropyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl.

[0062] In another aspect, the compound has formula (Ia):



or a pharmaceutically acceptable salt thereof, wherein ring A, R¹ and R² are as defined as for Formula (I).


5 **[0063]** In an embodiment, ring A is selected from:




wherein:

- 10 "1" represents a portion of ring A bound to a pyrrolo[1,2-b]pyridazine moiety;
 "2" represents a portion of ring A bound to R²; and
 ring A is substituted with 0, 1, 2, or 3 independently selected substituents in addition to R².
 R¹ is selected from C₁-C₃ alkyl, -O-(C₁-C₅ alkyl), -NH(C₁-C₅ alkyl), -NH-(C₃-C₆ cycloalkyl), C₃-C₆ cycloalkyl, -O-(C₃-C₆ cycloalkyl), an N-containing heterocyclyl, -O-(C₀-C₃ alkylene)-(O-containing heterocyclyl), -NH-(C₀-C₃ alkylene)-(O-containing heterocyclyl), and an O-containing heterocyclyl,
 15 wherein any alkyl, cycloalkyl, or heterocyclyl portion of R¹ is unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from halo, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, -O-C₁-C₄ alkyl, and -OH;
 R² is selected from heterocyclyl, -O-(C₀-C₄ alkylene)-(heterocyclyl), -(C₁-C₃ alkylene)-heterocyclyl, -(C₁-C₃ alkylene)-NH-(C₁-C₃ alkyl), -(hydroxy-substituted C₁-C₃ alkylene)-NH-(C₁-C₃ alkyl), and C₁-C₃ alkyl substituted with both hydroxy and amino, or R² is taken together with a ring atom in ring A to form a heterocyclyl or a carbocyclyl that is fused to ring A, wherein any heterocyclyl is unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from halo, cyano, -NH₂, -OH, -C₁-C₄ alkyl, deuterated C₁-C₄ alkyl, -C₁-C₄ haloalkyl, hydroxy-substituted -C₁-C₄ alkyl, -O-C₁-C₄ alkyl, C₃-C₆ cycloalkyl, -NH(C₁-C₄ alkyl), -NH-C(O)-O-(C₁-C₄ alkyl), -(C₁-C₄ alkylene)-O-(C₁-C₄ alkyl), -(C₁-C₄ alkylene)-O-(C₁-C₄ haloalkyl), -C(O)-O-C₁-C₄ alkyl, and an optionally substituted second heterocyclyl; and
 20 R^{3a} is selected from hydrogen and C₁-C₄ alkyl.

35 **[0064]** In an embodiment of Formula Ia, ring A is selected from:



and



wherein:

"1" represents a portion of ring A bound to a pyrrolo[1,2-b]pyridazine moiety;

"2" represents a portion of ring A bound to R²; and

ring A is substituted with 0, 1, 2, or 3 independently selected substituents in addition to R².


R¹ is selected from C₁-C₃ alkyl, -O-(C₁-C₃ alkyl), -NH(C₁-C₃ alkyl), -NH-(C₃-C₆ cycloalkyl), C₃-C₆ cycloalkyl, -O-(C₃-C₆ cycloalkyl), and an N-containing heterocyclyl, wherein any alkyl, cycloalkyl, or heterocyclyl portion of R¹ is unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from halo, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, and -OH;

R² is selected from heterocyclyl, and -O-(C₀-C₄ alkylene)-(heterocyclyl), or


R² is taken together with a ring atom in ring A to form a heterocyclyl that is fused to ring A, wherein any heterocyclyl is unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from halo, -OH, -C₁-C₄ alkyl, deuterated C₁-C₄ alkyl, -C₁-C₄ haloalkyl, -O-C₁-C₄ alkyl, and C₃-C₆ cycloalkyl; and

R^{3a} is selected from hydrogen and C₁-C₄ alkyl.

[0065] In another embodiment of Formula Ia, ring A is selected from:



and



wherein:

"1" represents a portion of ring A bound to a pyrrolo[1,2-b]pyridazine moiety;

"2" represents a portion of ring A bound to R²; and

ring A is substituted with 0, 1, 2, or 3 independently selected substituents in addition to R².

R¹ is selected from C₁-C₃ alkyl, -O-(C₁-C₃ alkyl), -NH(C₁-C₃ alkyl), -NH-(C₃-C₆ cycloalkyl), C₃-C₆ cycloalkyl, -O-(C₃-C₆ cycloalkyl), an N-containing heterocyclyl, -O-(O-containing heterocycle), and -O-(N-containing heterocycle), wherein any alkyl, cycloalkyl, or heterocyclyl portion of R¹ is

unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from deuterium, halo, cyano, C₁-C₄ alkyl, C₁-C₄ haloalkyl, and -OH;

R² is selected from heterocyclyl, and -O-(C₀-C₄ alkylene)-(heterocyclyl), or

R² is taken together with a ring atom in ring A to form a heterocyclyl that is fused to ring A, wherein any heterocyclyl portion of R² is unsubstituted, or is substituted with 1, 2, or 3 substituents independently selected from halo, -CN, -OH, -C₁-C₅ alkyl optionally substituted with one or more -OH and/or one or

more -NH₂, deuterated C₁-C₅ alkyl, -C₁-C₅ haloalkyl, -O-C₁-C₄ alkyl, and C₃-C₆ cycloalkyl; and
 R^{3a} is selected from hydrogen and C₁-C₄ alkyl.

5

[0066] In an embodiment, ring A is optionally substituted with up to 1 substituent in addition to R², wherein the substituent, if present, is halo or methyl. In an embodiment, ring A is optionally substituted with up to 1 substituent in addition to R², wherein the substituent, if present, is halo.

10

[0067] In an embodiment, R¹ is selected from 2,2,2-trifluoroethoxy, 2,2,2-trifluoroethylamino, 2,2-difluoroethoxy, 2,2-dimethylcyclopropyl, 2,2-dimethylpropoxy, 2-cyanocyclopropyl, 2-cyanoethyl, 2-cyanoethylamino, 2-fluorocyclopropyl, 2-methylcyclopropyl, 3-cyanoazetidin-1-yl, 3-cyanocyclobutoxy, 3-cyanocyclobutyl, 3-fluorocyclobutyl, 3-hydroxy-3-methylcyclobutyl, 3-hydroxy-3-trifluoromethylcyclobutyl, 3-hydroxyazetidin-1-yl, 3-hydroxycyclobutyl, 4-cyanocyclohexyl, 4-hydroxycyclohexyl, 4-methylcyclohexyl, cyclobutoxy, cyclobutyl, cyclobutylamino, cyclopropyl, ethoxy, ethylamino, isopropoxy, isopropylamino, N-(1,2-dimethylpropyl)amino, N-(2,2-dimethylpropyl)amino, N-(methoxymethyl)amino, N-(oxetan-3-ylmethyl)amino, oxetan-3-yl, oxetan-3-ylamino, oxetan-3-ylmethoxy, oxetan-3-yloxy, propylamino, t-butoxy, tetrahydrofuran-3-yloxy, and tetrahydropyran-4-yloxy.

20

[0068] In an embodiment, R¹ is selected from 2,2,2-trifluoroethoxy, 2,2-dimethylcyclopropyl, 2-cyanocyclopropyl, 2-cyanoethyl, 2-fluorocyclopropyl, 2-methylcyclopropyl, 3-cyanoazetidin-1-yl, 3-cyanocyclobutyl, 3-fluorocyclobutyl, 3-hydroxy-3-methylcyclobutyl, 3-hydroxy-3-trifluoromethylcyclobutyl, 3-hydroxyazetidin-1-yl, 3-hydroxycyclobutyl, 4-cyanocyclohexyl, 4-hydroxycyclohexyl, 4-methylcyclohexyl, cyclobutoxy, cyclobutyl, cyclobutylamino, cyclopropyl, ethoxy, ethylamino, isopropoxy, isopropylamino, and propylamino.

25

30

[0069] In an embodiment, R² is selected from 1-(1-hydroxypropan-2-yl)-4-methoxypiperidin-4-yl, 1-(3-difluoromethoxy)propan-2-yl-4-methoxypiperidin-4-yl, 1-(3-methoxy)propan-2-yl-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)-4-methoxypiperidin-4-yl, 1-(propan-2-yl)piperidin-3-yl, 1-(propan-2-yl)piperidin-4-yl, 1-(pyrrolidin-1-yl)ethan-1-yl, 1,2,3,6-tetrahydropyridin-4-yl, 1-cyclobutylpiperidin-3-yl, 1-ethyl-3-fluoropiperidin-4-yl, 1-ethyl-3-hydroxyazetidin-3-yl, 1-ethyl-4-fluoropyrrolidin-3-yl, 1-ethylazetidin-3-yl, 1-ethylazetidin-3-yloxy, 1-ethylpiperidin-3-yl, 1-ethylpiperidin-3-yloxy, 1-ethylpiperidin-4-yl, 1-ethylpyrrolidin-3-yl, 1-ethylpyrrolidin-3-ylmethoxy, 1-ethylpyrrolidin-3-yloxy, 1H-pyrrolidin-2-yl, 1-hydroxy-2-aminoprop-2-yl, 1-isopropyl-1,2,3,6-tetrahydropyridin-4-yl, 1-isopropyl-2-methylpyrrolidin-2-yl, 1-isopropyl-3,4-dimethylpiperazin-3-yl, 1-isopropyl-3-ethoxypiperidin-3-yl, 1-isopropyl-3-fluoropiperidin-3-yl, 1-isopropyl-3-hydroxyazetidin-3-yl, 1-isopropyl-3-hydroxypiperidin-3-yl, 1-isopropyl-3-hydroxypyrrolidin-3-yl, 1-isopropyl-3-methoxyazetidin-3-yl, 1-isopropyl-3-methoxypiperidin-3-yl, 1-isopropyl-3-methoxypyrrolidin-3-yl, 1-isopropyl-3-methylpiperazin-3-yl, 1-isopropyl-4-cyanopiperidin-4-yl, 1-isopropyl-4-ethoxypiperidin-4-yl, 1-isopropyl-4-fluoropiperidin-4-yl, 1-isopropyl-4-fluoropyrrolidin-3-yl, 1-isopropyl-4-hydroxypiperidin-3-yl, 1-

35

40

45

isopropyl-4-hydroxypiperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-isopropyl-4-methylpiperazin-3-yl, 1-isopropyl-4-methylpiperidin-4-yl, 1-isopropyl-4-trifluoromethylpiperidin-4-yl, 1-isopropyl-5-methylpyrrolidin-3-yl, 1-isopropylazetidin-3-yl, 1-isopropylpiperazin-3-yl, 1-isopropylpiperazin-4-yl, 1-isopropylpiperidin-2-yl, 1-isopropylpiperidin-3-yl, 1-isopropylpiperidin-4-yl, 1-isopropylpyrrolidin-2-yl, 1-isopropylpyrrolidin-3-yl, 1-sec-butylpiperidin-4-yl, 1-t-butoxycarbonyl-4-aminopiperidin-4-yl, 2-(isopropylamino)-3-hydroxypropan-2-yl, 2-(isopropylamino)-propan-2-yl, 2,3,5,6-tetrahydroimidazo[2,1-b]thiazol-6-yl, 2-difluoromethylpiperazin-1-yl, 2-isopropyl-2,6-diazaspiro[3.3]heptan-6-yl, 2-methyl-1H-pyrrolidin-2-yl, 3-aminopyrrolidin-1-yl, 3-fluoropiperidin-3-yl, 3-hydroxyquinuclidin-3-yl, 3-methyl-4-ethylpiperazin-1-yl, 3-methylpiperazin-1-yl, 4-(1,1,2,2-pentadeuteroethyl)piperazin-1-yl, 4-(methoxycarbonylamino)piperidin-4-yl, 4-cyanopiperidin-4-yl, 4-ethoxypiperidin-4-yl, 4-ethylpiperazin-1-yl, 4-fluoropiperidin-4-yl, 4-fluoropyrrolidin-3-yl, 4-isopropylmorpholin-3-yl, 4-isopropylpiperazin-1-yl, 4-methoxypiperidin-4-yl, 4-methylpiperidin-4-yl, 6-ethyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-isopropyl-2,6-diazaspiro[3.3]heptan-2-yl, azetidin-2-ylmethoxy, hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, morpholin-2-yl, morpholin-3-yl, piperazin-1-yl, piperdin-4-yl, piperidin-2-yl, piperidin-3-yl, piperidin-3-yloxy, pyrrolidin-2-yl, pyrrolidin-3-yloxy, and quinuclidin-4-yl, or

R² is taken together with ring A to form 6-isopropyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 5-isopropyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 1-amino-2,3-dihydro-1H-inden-5-yl, or 1-(isopropylamino)-2,3-dihydro-1H-inden-5-yl.


[0070] In an embodiment, R² is selected from 1-cyclobutylpiperidin-3-yl, 1-ethyl-3-fluoropiperidin-4-yl, 1-ethyl-3-hydroxyazetidin-3-yl, 1-ethyl-4-fluoropyrrolidin-3-yl, 1-ethylazetidin-3-yl, 1-ethylazetidin-3-yloxy, 1-ethylpiperidin-3-yl, 1-ethylpiperidin-3-yloxy, 1-ethylpiperidin-4-yl, 1-ethylpyrrolidin-3-yl, 1-ethylpyrrolidin-3-ylmethoxy, 1-ethylpyrrolidin-3-yloxy, 1-isopropyl-3-hydroxyazetidin-3-yl, 1-isopropyl-3-hydroxypiperidin-3-yl, 1-isopropyl-3-methoxypiperidin-3-yl, 1-isopropyl-4-fluoropyrrolidin-3-yl, 1-isopropyl-4-hydroxypiperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-isopropylazetidin-3-yl, 1-isopropylpiperidin-2-yl, 1-isopropylpiperidin-3-yl, 1-isopropylpiperidin-4-yl, 1-isopropylpyrrolidin-2-yl, 1-isopropylpyrrolidin-3-yl, 1-sec-butylpiperidin-4-yl, 2-difluoromethylpiperazin-1-yl, 3-fluoropiperidin-3-yl, 3-methyl-4-ethylpiperazin-1-yl, 3-methylpiperazin-1-yl, 4-(1,1,2,2,2-pentadeuteroethyl)piperazin-1-yl, 4-ethylpiperazin-1-yl, 4-fluoropyrrolidin-3-yl, 4-isopropylpiperazin-1-yl, 6-ethyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-isopropyl-2,6-diazaspiro[3.3]heptan-2-yl, azetidin-2-ylmethoxy, hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, morpholin-2-yl, piperazin-1-yl, piperdin-4-yl, piperidin-2-yl, piperidin-3-yl, piperidin-3-yloxy, pyrrolidin-2-yl, and pyrrolidin-3-yloxy, or R² is taken together with ring A to form 6-isopropyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, or 5-isopropyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl.

[0071] In an embodiment, R^{3a} is selected from hydrogen and methyl.

[0072] In an embodiment, ring A is substituted with 0 or 1 substituent in addition to R^2 , wherein the substituent, if present, is halo.

- 5 **[0073]** In an alternate embodiment, ring A is substituted with 0 or 1 substituent in addition to R^2 , wherein the substituent, if present, is selected from chloro, fluoro, and methyl.

[0074] In yet another embodiment, the compound is a compound of formula (II):



(II),

or a pharmaceutically acceptable salt thereof, wherein:

X is $C(R^{13})$ or N;


R^{11} is selected from $-NH-(C_3-C_4 \text{ cycloalkyl})$; $-NH-C_1-C_3 \text{ alkyl}$; $-O-C_3-C_4 \text{ cycloalkyl}$; $-O-C_1-C_3 \text{ alkyl}$ optionally substituted with one or more substituents selected from fluoro, hydroxy, cyano, and deuterium; and $-O-(O-$ containing heterocycle);

R^{12} is selected from piperidin-3-yl optionally 3-substituted with $C_1-C_3 \text{ alkoxy}$, fluoro, $C_1-C_3 \text{ alkyl}$, or cyano; and piperidin-4-yl optionally 4-substituted with $C_1-C_3 \text{ alkoxy}$, fluoro, $C_1-C_3 \text{ alkyl}$, cyano, wherein R^{12} is additionally optionally 1-substituted with $C_1-C_5 \text{ alkyl}$ optionally substituted with one or more hydroxy and/or one or more $-NH_2$;

R^{13} is selected from hydrogen, cyano and fluoro; and

R^{14} is fluoro.

[0075] In certain embodiments of Formula II, the compound is a compound of Formula IIa:



(IIa),


or a pharmaceutically acceptable salt thereof, wherein:

X, R^{11} , R^{14} , and subvariables thereof are as defined in Formula II;

The term "subvariables" as used herein means the variables that are used to define a variable. For example, X is $C(R^{13})$; R^{13} is a subvariable of X.

R^{15} is selected from hydrogen, C_1 - C_3 alkoxy, fluoro, C_1 - C_3 alkyl, and cyano;
and

R^{16} is C_1 - C_5 alkyl optionally substituted with one or more -OH and/or one or more NH_2 . In certain embodiments of Formula II, the compound is a compound of Formula lib:




(IIb),

or a pharmaceutically acceptable salt thereof, wherein:

X, R¹¹, R¹⁴, and subvariables thereof are as defined in Formula II;

R^{15} is selected from hydrogen, C_1 - C_3 alkoxy, fluoro, C_1 - C_3 alkyl, and cyano;


and R^{16} is C_1 - C_5 alkyl optionally substituted with one or more -OH and/or one or more -NH₂. In certain embodiments of a compound of Formula IIb, the compound is a compound of Formula IIb-1:



(IIb-1),

or a pharmaceutically acceptable salt thereof, wherein X, R¹¹, R¹², R¹⁴, R¹⁵, R¹⁶, and subvariables thereof are as defined in Formula lib.

[0076] In certain embodiments of a compound of Formula IIb, the compound is a compound of Formula IIb-2:



(IIb-2),

or a pharmaceutically acceptable salt thereof, wherein X, R¹¹, R¹², R¹⁴, R¹⁵, R¹⁶, and subvariables thereof are as defined in Formula lib.

[0077] In certain embodiments of Formula II, IIa, IIb, IIb-1 and IIb-2, R¹⁴ is absent.

[0078] In certain embodiments of Formula II, IIa, IIb, IIb-1 and IIb-2, R¹³ is hydrogen.

[0079] In certain embodiments of Formula II, IIa, IIb, IIb-1 and IIb-2, R¹¹ is selected from -NH-C₁-C₃ alkyl; -O-C₁-C₃ alkyl optionally substituted with one or more substituents selected from fluoro, hydroxy, cyano, and deuterium; oxetan-3-yl and tetrahydrofuran-3-yloxy. In some aspects of these embodiments, R¹¹ is selected from -OCH₂CH₃, -NHCH(CH₃)₂, oxetan-3-yl and tetrahydrofuran-3 -yloxy.





[0080] In certain embodiments the compound is a compound of Formula I or Formula Ia that is not a compound of any of Formula II, IIa, IIb, IIb-1 or IIb-2.

[0081] In an embodiment, the compound is a compound of any of Formulae I, Ia, II, IIb, IIb-1, or IIb-2 selected from a compound in Table 1.






[0082] In another aspect, the present disclosure features a pharmaceutical composition comprising a compound of any of Formulae I, Ia, II, IIb, IIb-1, or IIb-2 described herein (e.g., a compound in Table 1) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.






[0083] Table 1 below shows the structures of compounds described herein.






Table 1.






#	Structure	LCMS (M+1)	NMR
100		392	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (s, 1H), 7.94 (s, 1H), 7.88 (s, 1H), 7.86 (d, 1H, J = 5.2 Hz), 6.85 (s, 1H), 5.96 (d, 1H, J = 5.2 Hz), 5.18-5.13 (m, 1H), 3.92-3.88 (m, 4H), 3.74 (t, 2H, J = 8.0 Hz), 3.70-3.68 (m, 2H), 3.55-3.45 (m, 5H), 2.02-2.00 (m, 1H), 0.77-0.73 (m, 4H).
101		392	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 1.6 Hz), 7.95 (d, 1H, J = 5.2 Hz), 6.92 (d, 1H, J = 1.6 Hz), 6.70 (s, 1H), 6.01 (d, 1H, J = 5.2 Hz), 4.09 (t, 2H, J = 5.2 Hz), 3.93-3.90 (m, 4H), 3.72-3.68 (m, 2H), 3.57-3.55 (m, 2H), 3.52-3.47 (m, 2H), 3.06 (t, 2H, J = 5.2 Hz), 2.04-1.99 (m, 1H), 0.78-0.72 (m, 4H).
102		402	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.93 (s, 1H), 7.85 (d, 1H, J = 5.2 Hz), 7.63 (d, 2H, J = 8.4 Hz), 7.36 (d, 2H, J = 8.4 Hz), 6.72 (s, 1H), 5.85 (d, 1H, J = 5.2 Hz), 4.00-3.90 (m, 4H), 3.90-3.80 (m, 4H), 3.57-3.50 (m, 2H), 3.50-3.40 (m, 2H), 1.85-1.80 (m, 1H), 1.79-1.73 (m, 1H), 1.06-1.04 (m, 2H), 0.85-0.81 (m, 2H).
103		405	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.12 (s, 1H), 7.94 (s, 1H), 7.86 (d, 1H, J = 5.2 Hz), 7.80 (s, 1H), 6.83 (s, 1H), 5.96 (d, 1H, J = 5.2 Hz), 4.68 (quintet, 1H, J = 9.0 Hz), 3.92-3.43 (m, 8H), 2.11-2.06 (m, 2H), 2.04-2.02 (m, 1H), 1.97-1.93 (m, 2H), 1.83-1.78 (m, 2H), 1.68-1.64 (m, 2H), 0.79-0.73 (m, 4H).






#	Structure	LCMS (M+1)	NMR
104		406	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (s, 1H), 7.95 (s, 1H), 7.91 (s, 1H), 7.86 (d, 1H, J = 5.2 Hz), 6.83 (s, 1H), 5.96 (d, 1H, J = 5.2 Hz), 5.09-5.05 (br. s, 1H), 3.94-3.86 (m, 2H), 3.72-3.66 (m, 2H), 3.55-3.42 (m, 6H), 3.27-3.20 (m, 2H), 2.38-2.32 (m, 1H), 2.25-2.21 (m, 1H), 2.10-2.00 (m, 1H), 0.77-0.73 (m, 4H).
105		406	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.98 (s, 1H), 7.89 (d, 1H, J = 5.2 Hz), 7.78 (s, 1H), 6.86 (s, 1H), 6.61 (s, 1H), 5.97 (d, 1H, J = 5.2 Hz), 4.91-4.77 (m, 1H), 3.96-3.90 (m, 2H), 3.74-3.68 (m, 2H), 3.57-3.53 (m, 2H), 3.47-3.43 (m, 2H), 3.18-3.11 (m, 1H), 3.10-3.02 (m, 1H), 3.01-2.94 (m, 1H), 2.90-2.80 (m, 1H), 2.34-2.23 (m, 1H), 2.22-2.14 (m, 1H), 2.08-1.95 (m, 2H), 0.83-0.68 (m, 4H).
106		406	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 2.0 Hz), 7.95 (d, 1H, J = 5.6 Hz), 6.93 (d, 1H, J = 2.0 Hz), 6.72 (s, 1H), 6.02 (d, 1H, J = 5.6 Hz), 4.21 (t, 2H, J = 5.6 Hz), 3.93-3.90 (m, 2H), 3.72-3.68 (m, 2H), 3.60-3.52 (m, 4H), 3.49-3.45 (m, 2H), 2.76 (t, 2H, J = 5.6 Hz), 2.38 (s, 3H), 2.04-1.99 (m, 1H), 0.79-0.74 (m, 4H).
107		408	¹ H-NMR (400 MHz, MeOD) δ ppm 7.86 (d, 1H, J = 5.6 Hz), 7.83 (d, 1H, J = 1.6 Hz), 7.09 (s, 1H), 6.77 (d, 1H, J = 1.6 Hz), 6.02 (d, 1H, J = 5.6 Hz), 4.24 (s, 2H), 4.05-3.92 (m, 2H), 3.90-3.77 (m, 2H), 3.70-3.60 (m, 2H), 3.59-3.49 (m, 2H), 3.35 (t, 2H, J = 6.0 Hz), 2.89 (t, 2H, J = 6.0 Hz), 2.05-1.97 (m, 1H), 0.95-0.91 (m, 2H), 0.89-0.85 (m, 2H).
108		409	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.32 (s, 1H), 7.98 (d, 1H, J = 5.6 Hz), 7.18 (s, 1H), 6.03 (d, 1H, J = 5.6 Hz), 4.05-3.80 (m, 2H), 3.75-3.65 (m, 2H), 3.65-3.55 (m, 2H), 3.50-3.40 (m, 2H), 3.27-3.24 (m, 2H), 2.81-2.76 (m, 2H), 2.05-1.97 (m, 1H), 1.97-1.8 (m, 2H), 0.79-0.74 (m, 4H).
109		409	¹ H-NMR (400 MHz, MeOD) δ ppm 8.10 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.04 (d, 1H, J = 1.6 Hz), 6.04 (d, 1H, J = 6.0 Hz), 4.10-3.95 (m, 4H), 3.95-3.75 (m, 2H), 3.70-3.60 (m, 2H), 3.60-3.50 (m, 2H), 3.19 (t, 2H, J = 6.0 Hz), 2.89 (t, 2H, J = 6.0 Hz), 2.05-1.97 (m, 1H), 0.99-0.85 (m, 2H), 0.85-0.80 (m, 2H).






#	Structure	LCMS (M+1)	NMR
110		414	
111		415	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.32 (d, 1H, J = 1.2 Hz), 7.97 (d, 2H, J = 8.0 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.88 (d, 2H, J = 8.0 Hz), 7.18 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.93-3.92 (m, 2H), 3.73-3.67 (m, 2H), 3.57-3.53 (m, 2H), 3.51-3.47 (m, 2H), 3.49-3.33 (m, 4H), 2.04-1.99 (m, 1H), 0.78-0.72 (m, 4H).
112		416	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.89 (d, 1H, J = 6.0 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.31 (d, 2H, J = 7.6 Hz), 7.04 (s, 1H), 5.97 (d, 1H, J = 5.2 Hz), 3.92-3.91 (m, 2H), 3.69-3.68 (m, 2H), 3.53-3.47 (m, 4H), 3.43-3.41 (m, 1H), 3.41-3.38 (m, 1H), 3.30-3.28 (m, 1H), 3.09-3.06 (m, 1H), 2.89-2.84 (m, 1H, J = 10.0 Hz), 2.25-2.22 (m, 1H), 2.03-2.00 (m, 1H), 1.83-1.82 (m, 1H), 0.78-0.74 (m, 4H).
113		417	
114		420	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (d, 1H, J = 2.0 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.88 (d, 2H, J = 8.4 Hz), 7.57 (d, 2H, J = 8.4 Hz), 7.10 (d, 1H, J = 1.6 Hz), 6.58 (br s, 1H), 5.99 (d, 1H, J = 5.6 Hz), 4.25 (d, 2H, J = 10.8 Hz), 4.02 (d, 2H, J = 10.4 Hz), 3.95-3.90 (m, 2H), 3.73-3.68 (m, 1H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 2.06-2.00 (m, 1H), 0.79-0.75 (m, 4H).






#	Structure	LCMS (M+1)	NMR
115		418	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.11 (s, 1H), 7.89 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.4 Hz), 6.98 (s, 1H), 6.84 (d, 2H, J = 8.4 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.05-4.98 (m, 1H), 3.97-3.92 (m, 3H), 3.75-3.70 (m, 2H), 3.68-3.61 (m, 2H), 3.53-3.49 (m, 2H), 3.45-3.41 (m, 2H), 2.04-1.98 (m, 1H), 0.78-0.74 (m, 4H).
116		420	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.83 (d, 1H, J = 5.6 Hz), 7.76 (d, 1H, J = 1.6 Hz), 7.74 (s, 1H), 7.68 (s, 1H), 6.53 (d, 1H, J = 1.6 Hz), 5.84 (d, 1H, J = 5.6 Hz), 4.33-4.23 (m, 1H), 4.00-3.83 (m, 4H), 3.60-3.40 (m, 5H), 3.13-3.03 (m, 2H), 2.80-2.70 (m, 1H), 2.33-2.23 (m, 1H), 2.10-1.98 (m, 1H), 1.82-1.76 (m, 2H), 1.72-1.64 (m, 1H), 1.09-1.03 (m, 2H), 0.88-0.80 (m, 2H).
117		420	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.11 (s, 1H), 7.92 (d, 1H, J = 1.6 Hz), 7.85 (d, 2H, J = 5.2 Hz), 7.80 (s, 1H), 6.82 (d, 1H, J = 1.6 Hz), 5.95 (d, 1H, J = 5.2 Hz), 4.18-4.13 (m, 1H), 3.91-3.90 (m, 2H), 3.69-3.68 (m, 2H), 3.50-3.43 (m, 3H), 3.05-3.02 (m, 2H), 2.61-2.55 (m, 2H), 2.04-1.96 (m, 3H), 1.81-1.73 (m, 2H), 0.78-0.72 (m, 4H).
118		420	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26-8.24 (m, 2H), 7.96 (s, 1H), 7.91 (s, 1H), 7.88 (d, 1H, J = 5.2 Hz), 6.86 (s, 1H), 5.97 (d, 1H, J = 4.8 Hz), 5.00 (br. s., 1H), 4.30-3.80 (m, 4H), 3.97-3.91 (m, 2H), 3.85-3.75 (m, 2H), 3.70-3.65 (m, 2H), 3.52-3.50 (m, 2H), 2.59 (q, 2H, J = 6.4 Hz), 2.03-2.01 (m, 1H), 0.95 (t, 3H, J = 6.4 Hz), 0.77-0.74 (m, 4H).
119		425	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.92 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.53-7.51 (m, 4H), 6.69 (d, 1H, J = 1.6 Hz), 5.85 (d, 1H, J = 5.6 Hz), 3.94-3.90 (m, 4H), 3.56-3.49 (m, 4H), 1.81-1.74 (m, 1H), 1.06-1.03 (m, 2H), 0.86-0.82 (m, 2H).






#	Structure	LCMS (M+1)	NMR
120		430	¹ H-NMR (400 MHz, MeOD) δ ppm 8.01 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.83 (d, 1H, J = 8.0 Hz), 7.72 (d, 1H, J = 8.0 Hz), 7.67 (s, 1H), 6.96 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.17 (s, 2H), 4.40-4.30 (m, 4H), 4.10-3.95 (m, 2H), 3.85-3.75 (m, 2H), 3.66-3.57 (m, 2H), 3.55-3.50 (m, 2H), 2.05-1.97 (m, 1H), 0.95-0.87 (m, 2H), 0.85-0.80 (m, 2H).
121		430	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (s, 1H), 7.88 (d, 1H, J = 4.8 Hz), 7.62 (d, 2H, J = 8.4 Hz), 7.05-6.95 (m, 2H), 6.94 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.75-3.65 (m, 2H), 3.58-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.45-3.20 (m, 4H), 2.05-1.95 (m, 1H), 0.78-0.74 (m, 4H).
122		430	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.23 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.2 Hz), 3.91 (br. s., 2H), 3.70 (br. s., 2H), 3.53-3.46 (m, 4H), 3.05-3.02 (m, 2H), 2.62-2.49 (m, 3H), 2.05-1.99 (m, 1H), 1.76-1.70 (m, 2H), 1.58-1.48 (m, 2H), 0.80-0.70 (m, 4H).
123		430	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.23 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.00-3.80 (m, 2H), 3.78-3.62 (m, 2H), 3.60-3.42 (m, 4H), 3.40-3.20 (m, 2H), 3.00-2.90 (m, 2H), 2.60-2.54 (m, 2H), 2.10-1.93 (m, 1H), 1.90-1.80 (m, 1H), 1.70-1.40 (m, 3H), 1.80-0.60 (m, 4H).
124		430	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.61 (s, 1H), 7.94 (s, 1H), 7.87 (d, 1H, J = 4.8 Hz), 7.66 (d, 2H, J = 8.0 Hz), 7.31 (d, 2H, J = 8.0 Hz), 6.71 (s, 1H), 5.86 (d, 1H, J = 4.8 Hz), 4.41-4.39 (m, 2H), 4.21-4.17 (m, 1H), 3.96-3.90 (m, 2H), 3.85-3.90 (m, 2H), 3.76-3.75 (m, 2H), 3.57-3.54 (m, 2H), 3.51-3.48 (m, 2H), 3.07 (q, 2H, J = 6.4 Hz), 1.79-1.78 (m, 1H), 1.25 (t, 3H, J = 6.4 Hz), 1.11-1.05 (m, 2H), 0.85-0.83 (m, 2H).






#	Structure	LCMS (M+1)	NMR
125		431	¹ H NMR (400 MHz, Chloroform-d) δ 7.92 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.3 Hz, 1H), 7.61 (d, J = 8.1 Hz, 2H), 7.41 (d, J = 8.0 Hz, 2H), 6.71 (d, J = 1.9 Hz, 1H), 5.84 (d, J = 5.4 Hz, 1H), 3.91 (d, J = 19.0 Hz, 5H), 3.68 – 3.42 (m, 7H), 3.21 (d, J = 11.7 Hz, 2H), 2.82 (t, J = 11.1 Hz, 1H), 1.94 – 1.75 (m, 3H), 1.04 (dt, J = 6.5, 3.2 Hz, 2H), 0.82 (dq, J = 7.1, 3.8 Hz, 2H).
126		431	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 7.6 Hz), 7.03 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.97-3.95 (m, 4H), 3.71-3.70 (m, 2H), 3.55-3.52 (m, 2H), 3.48-3.41 (m, 4H), 2.78-2.77 (m, 1H), 2.04-2.00 (m, 1H), 1.73-1.70 (m, 4H), 0.79-0.75 (m, 4H).
127		431	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.06 (d, 1H, J = 1.2 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.64 (d, 2H, J = 8.4 Hz), 7.00-6.85 (m, 3H), 5.95 (d, 1H, J = 5.6 Hz), 3.91 (br. s, 2H), 3.69 (br. s, 2H), 3.52 (br. s, 2H), 3.44 (br. s, 2H), 3.06-3.04 (m, 4H), 2.82 (br. s, 4H), 2.05-1.95 (m, 1H), 0.80-0.65 (m, 4H).
128		432	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.71-7.69 (m, 2H), 7.34 (t, 1H, J = 7.6 Hz), 7.21 (d, 1H, J = 7.6 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.43-4.41 (m, 1H), 3.91-3.89 (m, 3H), 3.71 (br. s., 2H), 3.64-3.32 (m, 5H), 3.00-2.96 (m, 1H), 2.76-2.75 (m, 2H), 2.65-2.55 (m, 1H), 2.05-1.99 (m, 1H), 0.78-0.74 (m, 4H).
129		432	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.0 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.95 (d, 1H, J = 5.6 Hz), 4.40-4.36 (m, 1H), 3.89-3.87 (m, 3H), 3.69-3.46 (m, 8H), 2.94-2.90 (m, 1H), 2.75-2.73 (m, 2H), 2.55-2.49 (m, 1H), 2.05-1.97 (m, 1H), 0.77-0.71 (m, 4H).

#	Structure	LCMS (M+1)	NMR
130		432	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.38 (dd, 1H, J = 9.6, 1.6 Hz), 3.92-3.87 (m, 3H), 3.75-3.65 (m, 2H), 3.64-3.40 (m, 5H), 3.38-3.35 (m, 1H), 2.92 (dd, 1H, J = 12.0, 2.0 Hz), 2.76-2.70 (m, 2H), 2.52-2.50 (m, 1H), 2.05-1.99 (m, 1H), 0.80-0.72 (m, 4H).
131		432	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.40-4.35 (m, 1H), 3.95-3.87 (m, 3H), 3.80-3.63 (m, 2H), 3.61-3.40 (m, 5H), 2.94-2.91 (m, 1H), 2.75-2.73 (m, 2H), 2.52-2.50 (m, 1H), 2.05-1.99 (m, 1H), 0.80-0.72 (m, 4H).
132		432	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.11 (s, 1H), 7.88 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.4 Hz), 6.99 (s, 1H), 6.96 (d, 2H, J = 8.4 Hz), 5.96 (d, 1H, J = 5.2 Hz), 4.19-4.12 (m, 1H), 4.02-3.97 (m, 2H), 3.95-3.90 (m, 2H), 3.75-3.68 (m, 2H), 3.54-3.48 (m, 4H), 3.31-3.29 (m, 2H), 2.30-2.25 (m, 1H), 2.14-2.07 (m, 1H), 2.04-1.97 (m, 1H), 0.78-0.74 (m, 4H).
133		432	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.09 (s, 1H), 7.87 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.4 Hz), 6.97 (s, 1H), 6.91 (d, 2H, J = 8.4 Hz), 5.96 (d, 1H, J = 6.0 Hz), 4.90-4.85 (m, 1H), 3.94-3.89 (m, 2H), 3.73-3.65 (m, 2H), 3.55-3.50 (m, 2H), 3.10-3.05 (m, 2H), 2.92-2.85 (m, 2H), 2.85-2.76 (m, 2H), 2.04-1.99 (m, 2H), 1.82-1.74 (m, 1H), 0.77-0.74 (m, 4H).
134		432	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.12 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.8 Hz), 6.98 (d, 3H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.14 (d, 2H, J = 6.8 Hz), 3.95-3.90 (m, 2H), 3.75-3.70 (m, 2H), 3.66 (d, 2H, J = 8.0 Hz), 3.55-3.50 (m, 2H), 3.49-3.45 (m, 2H), 3.44-3.40 (m, 2H), 3.09-2.99 (m, 1H), 2.05-2.01 (m, 1H), 0.78-0.75 (m, 4H).





#	Structure	LCMS (M+1)	NMR
135		432	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.52 (d, 1H, J = 2.4 Hz), 7.87-7.84 (m, 2H), 7.74 (dd, 1H, J = 2.4, 8.8 Hz), 6.71 (d, 1H, J = 8.8 Hz), 6.63 (d, 1H, J = 2.0 Hz), 5.85 (d, 1H, J = 5.6 Hz), 3.94-3.87 (m, 4H), 3.56-3.47 (m, 8H), 3.03-3.00 (m, 4H), 1.80-1.76 (m, 1H), 1.07-1.03 (m, 2H), 0.85-0.81 (m, 2H).
136		433	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.2 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.91 (d, 1H, J = 1.2 Hz), 5.94 (d, 1H, J = 5.2 Hz), 3.66-3.65 (m, 4H), 3.49-3.48 (m, 2H), 3.42-3.41 (m, 4H), 3.15-3.14 (m, 4H), 2.52-2.51 (m, 2H), 2.38 (q, 2H, J = 7.2 Hz), 2.06 (s, 3H), 1.03 (t, 3H, J = 7.2 Hz).
137		434	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.8 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.30-5.00 (m, 1H), 4.00-3.85 (m, 2H), 3.75-3.65 (m, 2H), 3.60-3.40 (m, 6H), 3.20-3.10 (m, 1H), 3.08-3.00 (m, 1H), 2.75-2.65 (m, 1H), 2.05-1.95 (m, 1H), 0.80-0.70 (m, 4H).
138		434	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.97 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.2 Hz), 7.78 (d, 1H, J = 2.4 Hz), 6.86 (d, 1H, J = 1.6 Hz), 7.96 (d, 1H, J = 5.2 Hz), 6.62 (d, 1H, J = 2.4 Hz), 5.96 (d, 1H, J = 5.2 Hz), 4.92-4.85 (m, 1H), 3.92 (br. s., 2H), 3.69 (br. s., 2H), 3.53-3.44 (m, 4H), 2.85 (q, 2H, J = 7.2 Hz), 2.807-2.77 (m, 1H), 2.48-2.42 (m, 3H), 2.39-2.30 (m, 1H), 2.13-2.06 (m, 1H), 2.04-1.97 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.78-0.73 (m, 4H).
139		434	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (s, 1H), 7.93 (d, 1H, J = 1.2 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.81 (s, 1H), 6.83 (d, 1H, J = 1.2 Hz), 5.95 (d, 1H, J = 5.2 Hz), 4.87-4.85 (m, 1H), 3.91-3.90 (m, 2H), 3.68-3.67 (m, 2H), 3.50-3.49 (m, 4H), 3.43-3.34 (m, 4H), 2.91-2.90 (m, 1H), 2.76-2.74 (m, 2H), 2.08-2.07 (m, 1H), 2.02-1.98 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.77-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
140		434	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.94 (d, 2H, J = 8.4 Hz), 6.91 (d, 1H, J = 1.6 Hz), 6.60 (t, 1H, J = 5.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.52-3.51 (m, 4H), 3.43-3.41 (m, 4H), 3.12-3.09 (m, 2H), 3.10-3.05 (m, 4H), 2.86-2.83 (m, 4H), 1.04 (t, 3H, J = 6.8 Hz).
141		435	¹ H-NMR (400 MHz, MeOD) δ ppm 7.99 (s, 1H), 7.86 (m, 1H), 7.78 (d, 2H, J = 8.0 Hz), 6.93 (s, 1H), 7.45 (d, 2H, J = 8.0 Hz), 6.96 (s, 1H), 6.00 (d, 1H, J = 5.6 Hz), 4.79-4.70 (m, 1H), 4.29-4.25 (m, 1H), 4.04-3.97 (m, 1H), 3.67-3.64 (m, 4H), 3.56-3.54 (m, 4H), 3.50-3.47 (m, 1H), 3.39-3.34 (m, 2H), 3.25-3.21 (m, 2H), 3.18-3.14 (m, 1H), 1.54 (t, 3H, J = 7.2 Hz).
142		435	¹ H-NMR (400 MHz, MeOD) δ ppm 7.99 (s, 1H), 7.85 (m, 1H), 7.77 (d, 2H, J = 8.4 Hz), 7.44 (d, 2H, J = 8.0 Hz), 6.93 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 4.79-4.77 (m, 1H), 4.28-4.25 (m, 1H), 4.05-3.98 (m, 1H), 3.66-3.64 (m, 4H), 3.55-3.53 (m, 4H), 3.49-3.47 (m, 1H), 3.39-3.40 (m, 2H), 3.27-3.22 (m, 2H), 3.16-3.14 (m, 1H), 1.53 (t, 3H, J = 7.2 Hz).
143		435	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.29 (br. s., 1H), 8.07 (d, 1H, J = 1.2 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.66 (d, 2H, J = 8.4 Hz), 6.96 (d, 2H, J = 8.4 Hz), 6.92 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.08 (q, 2H, J = 6.8 Hz), 3.65-3.55 (m, 4H), 3.50-3.40 (m, 4H), 3.20-3.10 (m, 4H), 3.05-2.90 (m, 4H), 1.21 (t, 3H, J = 6.8 Hz).
144		436	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 0.8 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.32 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 0.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.40-4.35 (m, 1H), 4.08 (q, 2H, J = 7.2 Hz), 3.90-3.80 (m, 1H), 3.65-3.60 (m, 1H), 3.61-3.56 (m, 4H), 3.48-3.40 (m, 4H), 3.00-2.90 (m, 1H), 2.80-2.70 (m, 2H), 2.60-2.40 (m, 1H), 1.21 (t, 3H, J = 7.2 Hz).






#	Structure	LCMS (M+1)	NMR
145		438	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.40-5.00 (m, 1H), 4.08 (q, 2H, J = 7.2 Hz), 3.70-3.50 (m, 4H), 3.50-3.47 (m, 2H), 3.45-3.40 (m, 4H), 3.20-3.15 (m, 1H), 3.12-3.05 (m, 1H), 2.85-2.75 (m, 1H), 1.21 (t, 3H, J = 7.2 Hz).
146		443	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.31 (s, 1H), 8.04 (d, 1H, J = 1.2 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.62 (d, 2H, J = 8.4 Hz), 6.91 (d, 1H, J = 1.2 Hz), 6.47 (d, 2H, J = 8.4 Hz), 5.95 (d, 1H, J = 5.2 Hz), 4.17-4.07 (m, 4H), 4.01-3.95 (m, 4H), 3.95-3.90 (m, 2H), 3.73-3.65 (m, 2H), 3.52-3.51 (m, 2H), 3.46-3.44 (m, 2H), 2.05-1.99 (m, 1H), 0.80-0.72 (m, 4H).
147		444	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.0 Hz), 7.31 (d, 2H, J = 8.0 Hz), 7.03 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.93-3.91 (m, 2H), 3.70-3.68 (m, 2H), 3.54-3.48 (m, 5H), 3.11-3.10 (m, 1H), 2.84-2.83 (m, 2H), 2.67-2.62 (m, 3H), 2.28-2.25 (m, 1H), 2.04-2.01 (m, 1H), 1.86-1.81 (m, 1H), 1.11 (t, 3H, J = 6.8 Hz), 0.79-0.73 (m, 4H).
148		444	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.70-3.69 (m, 2H), 3.53-3.47 (m, 4H), 3.29-3.25 (m, 1H), 2.93-2.89 (m, 1H), 2.68-2.62 (m, 2H), 2.47-2.40 (m, 2H), 2.24-2.20 (m, 2H), 2.03-1.98 (m, 1H), 1.78-1.73 (m, 1H), 1.05 (t, 3H, J = 7.2 Hz), 0.78-0.72 (m, 4H).
149		444	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.70-3.69 (m, 2H), 3.53-3.47 (m, 5H), 2.95-2.91 (m, 1H), 2.70-2.63 (m, 2H), 2.47-2.40 (m, 2H), 2.24-2.20 (m, 2H), 2.03-1.98 (m, 1H), 1.77-1.75 (m, 1H), 1.05 (t, 3H, J = 7.6 Hz), 0.78-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
150		444	¹ H-NMR (400 MHz, MeOD) δ ppm 7.99 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.78 (d, 2H, J = 8.4 Hz), 7.41 (d, 2H, J = 8.4 Hz), 6.95 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.40-4.32 (m, 2H), 4.10-4.02 (m, 5H), 3.88-3.82 (m, 2H), 3.65-3.60 (m, 2H), 3.58-3.52 (m, 2H), 3.38-3.32 (m, 1H), 2.07-2.00 (m, 1H), 1.24 (d, 6H, J = 6.4 Hz), 0.96-0.92 (m, 2H), 0.92-0.84 (m, 2H).
151		445	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.81 (d, 2H, J = 8.4 Hz), 7.31 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.92 (br. s, 2H), 3.70-3.65 (m, 4H), 3.55-3.50 (m, 4H), 3.34 (s, 2H), 3.01 (t, 2H, J = 5.2 Hz), 2.00 (quintet, 1H, J = 5.2 Hz), 0.79-0.75 (m, 4H).
152		445	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.06 (s, 1H), 7.85 (d, 1H, J = 5.2 Hz), 7.62 (d, 2H, J = 8.4 Hz), 6.93 (d, 2H, J = 8.4 Hz), 6.86 (s, 1H), 5.92 (d, 1H, J = 5.2 Hz), 4.65 (br. s., 1H), 4.25-4.10 (m, 1H), 4.00-3.80 (m, 2H), 3.40-3.20 (m, 4H), 3.10-3.00 (m, 4H), 2.90-2.70 (m, 4H), 2.05-1.90 (m, 1H), 1.45-1.20 (m, 3H), 0.75 (br. s., 4H).
153		445	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 1.6 Hz), 7.83 (d, 1H, J = 5.6 Hz), 7.57 (d, 2H, J = 8.8 Hz), 6.98 (d, 2H, J = 8.8 Hz), 6.66 (d, 1H, J = 0.8 Hz), 5.83 (d, 1H, J = 5.2 Hz), 3.95-3.89 (m, 4H), 3.56-3.47 (m, 4H), 3.29-3.27 (m, 4H), 2.66-2.23 (m, 4H), 2.40 (s, 3H), 1.82-1.75 (m, 1H), 1.06-1.04 (m, 2H), 0.84-0.82 (m, 2H).
154		445	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.08 (s, 1H), 7.87 (d, 1H, J = 5.6 Hz), 7.67 (d, 2H, J = 8.4 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.94 (s, 1H), 5.96 (d, 1H, J = 5.2 Hz), 3.68-3.03 (m, 17H), 2.23-2.11 (m, 4H), 1.93-1.90 (m, 1H), 1.78-1.77 (m, 1H).





#	Structure	LCMS (M+1)	NMR
155		445	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.95 (d, 2H, J = 8.4 Hz), 6.94 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 3.94-3.92 (m, 2H), 3.71-3.69 (m, 2H), 3.55-3.53 (m, 4H), 3.47-3.45 (m, 3H), 3.00-2.97 (m, 1H), 2.84-2.81 (m, 2H), 2.57-2.56 (m, 1H), 2.25-2.20 (m, 1H), 2.02-2.01 (m, 1H), 1.05 (d, 3H, J = 6.4 Hz), 0.78-0.75 (m, 4H).
156		445	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.94 (d, 2H, J = 8.8 Hz), 6.93 (s, 1H), 5.94 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.71-3.70 (m, 2H), 3.58-3.45 (m, 7H), 3.01-2.98 (m, 1H), 2.86-2.85 (m, 2H), 2.58-2.57 (m, 1H), 2.27-2.21 (m, 1H), 2.01-1.99 (m, 1H), 1.05 (d, 3H, J = 6.4 Hz), 0.78-0.74 (m, 4H).
157		445	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.07 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.91 (d, 1H, J = 1.6 Hz), 6.03 (d, 1H, J = 5.6 Hz), 4.14-4.05 (m, 1H), 4.04-3.96 (m, 1H), 3.95-3.88 (m, 1H), 3.86-3.74 (m, 1H), 3.50-3.38 (m, 4H), 3.20-3.10 (m, 4H), 3.09-3.00 (m, 1H), 2.85-2.75 (m, 1H), 2.70-2.60 (m, 1H), 2.36 (q, 2H, J = 7.2 Hz), 2.33-2.25 (m, 2H), 2.20-2.10 (m, 1H), 1.70-1.55 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz).
158		445	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.63 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.94-6.92 (m, 1H), 5.95 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.67-3.64 (m, 4H), 3.54-3.40 (m, 4H), 3.40-3.35 (m, 1H), 2.75-2.69 (m, 4H), 2.02-2.01 (m, 1H), 1.80-1.77 (m, 2H), 1.35-1.22 (m, 2H), 0.77-0.73 (m, 4H).
159		446	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 2.0 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.66 (d, 1H, J = 2.0 Hz), 5.84 (d, 1H, J = 5.2 Hz), 4.46 (heptet, 1H, J = 3.6 Hz), 3.99-3.90 (m, 2H), 3.90-3.81 (m, 2H), 3.61-3.50 (m, 2H), 3.50-3.39 (m, 2H), 3.23-3.18 (m, 1H), 2.92-2.85 (m, 2H), 2.82-.75 (m, 1H), 2.07-2.00 (m, 1H), 1.88-1.75 (m, 3H), 1.58-1.50 (m, 1H), 1.07-1.02 (m, 2H), 0.86-0.80 (m, 2H).






#	Structure	LCMS (M+1)	NMR
160		446	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.88 (d, 1H, J = 1.6 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.59 (d, 2H, J = 8.8 Hz), 6.96 (d, 2H, J = 8.8 Hz), 6.65 (d, 1H, J = 1.6 Hz), 5.85 (d, 1H, J = 5.2 Hz), 4.67-4.61 (m, 1H), 3.99-3.90 (m, 2H), 3.90-3.75 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.41-3.33 (m, 2H), 3.19-3.15 (m, 2H), 2.25-2.08 (m, 4H), 1.79-1.75 (m, 1H), 1.06-1.02 (m, 2H), 0.85-0.80 (m, 2H).
161		446	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 2.0 Hz), 8.17 (s, 1H), 7.91 (d, 1H, J = 5.2 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.61 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.2 Hz), 6.09 (br s, 1H), 5.99 (d, 1H, J = 5.2 Hz), 3.96-3.92 (m, 2H), 3.79-3.77 (m, 2H), 3.73-3.68 (m, 2H), 3.57-3.52 (m, 2H), 3.47-3.45 (m, 4H), 2.72 (q, 2H, J = 6.8 Hz), 2.06-1.98 (m, 1H), 0.98 (t, 3H, J = 6.8 Hz), 0.79-0.74 (m, 4H).
162		446	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 2.0 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.66 (d, 1H, J = 2.0 Hz), 5.84 (d, 1H, J = 5.2 Hz), 4.46 (heptet, 1H, J = 3.6 Hz), 3.99-3.90 (m, 2H), 3.90-3.81 (m, 2H), 3.61-3.50 (m, 2H), 3.50-3.39 (m, 2H), 3.23-3.18 (m, 1H), 2.92-2.85 (m, 2H), 2.82-.75 (m, 1H), 2.07-2.00 (m, 1H), 1.88-1.75 (m, 3H), 1.58-1.50 (m, 1H), 1.07-1.02 (m, 2H), 0.86-0.80 (m, 2H).
163		446	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 2.0 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.66 (d, 1H, J = 2.0 Hz), 5.84 (d, 1H, J = 5.2 Hz), 4.46 (heptet, 1H, J = 3.6 Hz), 3.99-3.90 (m, 2H), 3.90-3.81 (m, 2H), 3.61-3.50 (m, 2H), 3.50-3.39 (m, 2H), 3.23-3.18 (m, 1H), 2.92-2.85 (m, 2H), 2.82-.75 (m, 1H), 2.07-2.00 (m, 1H), 1.88-1.75 (m, 3H), 1.58-1.50 (m, 1H), 1.07-1.02 (m, 2H), 0.86-0.80 (m, 2H).

#	Structure	LCMS (M+1)	NMR
164		446	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.77 (d, 1H, J = 8.0 Hz), 7.33 (d, 1H, J = 8.0 Hz), 7.05 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.44 (d, 1H, J = 9.2 Hz), 3.95-3.90 (m, 2H), 3.73-3.69 (m, 2H), 3.69-3.68 (m, 1H), 3.57-3.54 (m, 2H), 3.49-3.46 (m, 2H), 2.90 (d, 1H, J = 14 Hz), 2.80 (d, 1H, J = 14 Hz), 2.50-2.45 (m, 1H), 2.35 (t, 1H, J = 14 Hz), 2.05-2.00 (m, 1H), 1.11 (d, 3H, J = 6.0 Hz), 0.78-0.74 (m, 4H).
165		446	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.77 (d, 1H, J = 8.0 Hz), 7.33 (d, 1H, J = 8.0 Hz), 7.05 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.44 (d, 1H, J = 9.2 Hz), 3.95-3.90 (m, 2H), 3.73-3.69 (m, 2H), 3.69-3.68 (m, 1H), 3.57-3.54 (m, 2H), 3.49-3.46 (m, 2H), 2.90 (d, 1H, J = 14 Hz), 2.80 (d, 1H, J = 14 Hz), 2.50-2.45 (m, 1H), 2.35 (t, 1H, J = 14 Hz), 2.05-2.00 (m, 1H), 1.11 (d, 3H, J = 6.0 Hz), 0.78-0.74 (m, 4H).
166		446	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.34 (s, 1H), 8.12 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.4 Hz), 6.99 (s, 1H), 6.97 (d, 2H, J = 8.4 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.04-4.01 (m, 2H), 3.99-3.92 (m, 2H), 3.71-3.70 (m, 2H), 3.55-3.51 (m, 2H), 3.49-3.45 (m, 2H), 3.33-3.31 (m, 1H), 3.29-3.23 (m, 1H), 3.15-3.14 (m, 1H), 3.00-2.99 (m, 1H), 2.73-2.71 (m, 1H), 2.04-2.02 (m, 1H), 2.02-2.00 (m, 1H), 1.80-1.75 (m, 1H), 0.79-0.75 (m, 4H).
167		446	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (s, 1H), 8.12 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.8 Hz), 6.98 (d, 1H, J = 1.6 Hz), 6.87 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.89-3.86 (m, 1 H), 3.94-3.92 (m, 4H), 3.56-3.51 (m, 4H), 3.47-3.45 (m, 2H), 3.22-3.20 (m, 2H), 2.64 (q, 2H, J = 7.2 Hz), 2.04-2.00 (m, 1H), 0.95 (t, 3H, J = 7.2 Hz), 0.79-0.73 (m, 4H).






#	Structure	LCMS (M+1)	NMR
168		446	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.38 (d, 1H, J = 8.4 Hz), 7.05 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.75-4.70 (m, 1H), 3.95-3.90 (m, 2H), 3.73-3.68 (m, 2H), 3.57-3.52 (m, 2H), 3.50-3.45 (m, 2H), 2.95-2.90 (m, 1H), 2.85-2.80 (m, 1H), 2.79-2.70 (m, 1H), 2.50-2.34 (m, 1H), 2.03-1.99 (m, 1H), 1.27 (d, 3H, J = 6.0 Hz), 0.78-0.74 (m, 4H).
169		446	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.40 (d, 2H, J = 8.4 Hz), 7.07 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.75-4.70 (m, 1H), 3.95-3.90 (m, 2H), 3.73-3.69 (m, 2H), 3.58-3.54 (m, 2H), 3.50-3.45 (m, 2H), 3.23-3.19 (m, 1H), 3.08-3.05 (m, 1H), 3.04-3.01 (m, 1H), 2.84-2.80 (m, 1H), 2.10-2.01 (m, 1H), 2.01-1.98 (m, 1H), 1.33 (d, 3H, J = 7.2 Hz), 0.78-0.74 (m, 4H).
170		446	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.63 (d, 1H, J = 2.0 Hz), 8.16 (d, 1H, J = 2.0 Hz), 8.01 (dd, 1H, J = 8.8, 2.4 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.00-6.90 (m, 2H), 5.95 (d, 1H, J = 5.2 Hz), 4.63 (br. s., 1H), 4.25-4.10 (m, 1H), 4.00-3.80 (m, 2H), 3.65 (br. s., 4H), 3.27 (br. s., 4H), 3.08 (br. s., 4H), 2.05-1.90 (m, 1H), 1.45-1.20 (m, 3H), 0.80-0.65 (m, 4H).
171		447	
172		447	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.0 Hz), 7.50 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.02 (s, 1H), 3.92-3.93 (m, 2H), 3.80-3.70 (m, 6H), 3.55-3.48 (m, 4H), 2.03-1.94 (m, 3H), 1.57-1.54 (m, 2H), 0.79-0.75 (m, 4H).

#	Structure	LCMS (M+1)	NMR
173		447	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.06 (br. s., 1H), 7.87 (br. s., 1H), 7.64-7.53 (m, 3H), 6.94 (br. s., 2H), 5.94 (br. s., 1H), 3.67 (br. s., 4H), 3.47-3.33 (m, 4H), 3.16 (br. s., 4H), 2.50-2.48 (m, 4H), 2.31 (br. s., 4H), 1.03 (d, 6H, J = 4.0 Hz).
174		448	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.16 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.30 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.2 Hz), 4.68-4.50 (m, 1H), 3.91 (br. s., 2H), 3.70 (br. s., 2H), 3.54 (br. s., 2H), 3.43 (br. s., 2H), 3.29-3.25 (m, 1H), 2.90-2.87 (m, 1H), 2.77-2.64 (m, 1H), 2.53-2.51 (m, 1H), 2.48-2.45 (m, 1H), 2.05-1.99 (m, 1H), 1.77-1.74 (m, 1H), 1.67-1.57 (m, 1H), 0.79-0.71 (m, 4H).
175		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.94 (d, 1H, J = 2.0 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.0 Hz), 7.33 (d, 2H, J = 8.0 Hz), 6.72 (d, 1H, J = 2.0 Hz), 5.85 (d, 1H, J = 5.6 Hz), 4.71-4.52 (m, 1H), 3.94-3.90 (m, 4H), 3.57-3.52 (m, 4H), 3.13-3.10 (m, 1H), 2.86-2.67 (m, 3H), 1.98-1.95 (m, 1H), 1.27 (s, 1H), 1.08-1.02 (m, 2H), 0.86-0.82 (m, 2H).
176		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.94 (d, 1H, J = 2.0 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.0 Hz), 7.33 (d, 2H, J = 8.0 Hz), 6.72 (d, 1H, J = 2.0 Hz), 5.85 (d, 1H, J = 5.2 Hz), 4.73-4.52 (m, 1H), 3.94-3.90 (m, 4H), 3.57-3.52 (m, 4H), 3.13-3.10 (m, 1H), 2.86-2.67 (m, 3H), 2.05-1.96 (m, 2H), 1.86-1.72 (m, 2H), 1.08-1.02 (m, 2H), 0.86-0.82 (m, 2H).
177		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.07 (d, 1H, J = 2.0 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.73-3.68 (m, 2H), 3.58-3.50 (m, 2H), 3.49-3.40 (m, 2H), 2.99-2.78 (m, 3H), 2.65-2.60 (m, 1H), 2.22-2.15 (m, 1H), 2.10-1.95 (m, 2H), 1.82-1.67 (m, 1H), 1.60-1.49 (m, 1H), 0.80-0.73 (m, 4H).






#	Structure	LCMS (M+1)	NMR
178		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.21 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.07 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.73-3.68 (m, 2H), 3.57-3.53 (m, 2H), 3.50-3.42 (m, 2H), 2.99-2.95 (m, 2H), 2.94-2.90 (m, 1H), 2.55-2.51 (m, 1H), 2.22-2.10 (m, 1H), 2.05-1.95 (m, 2H), 1.82-1.65 (m, 1H), 1.60-1.45 (m, 1H), 0.80-0.73 (m, 4H).
179		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.21 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.07 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.73-3.68 (m, 2H), 3.57-3.53 (m, 2H), 3.50-3.42 (m, 2H), 2.99-2.95 (m, 2H), 2.94-2.90 (m, 1H), 2.55-2.51 (m, 1H), 2.22-2.10 (m, 1H), 2.05-1.95 (m, 2H), 1.82-1.65 (m, 1H), 1.60-1.45 (m, 1H), 0.80-0.73 (m, 4H).
180		464	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.65 (s, 1H), 7.92 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.4 Hz), 7.31 (d, 2H, J = 8.4 Hz), 6.68 (d, 1H, J = 1.2 Hz), 5.85 (d, 1H, J = 4.8 Hz), 4.20 (q, 2H, J = 7.2 Hz), 3.81-3.70 (m, 4H), 3.67-3.55 (m, 2H), 3.49-3.40 (m, 4H), 3.39-3.35 (m, 1H), 3.26-3.14 (m, 1H), 3.10-3.00 (m, 2H), 2.99-2.95 (m, 1H), 2.51-2.42 (m, 1H), 2.21-2.08 (m, 1H), 1.34 (t, 3H, J = 7.2 Hz), 1.31 (t, 3H, J = 7.2 Hz).
181		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (s, 1H), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.28 (d, 2H, J = 8.0 Hz), 6.98 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.70-3.55 (m, 4H), 3.50-3.40 (m, 4H), 3.30-3.20 (m, 1H), 2.91 (t, 1H, J = 8.4 Hz), 2.75-2.68 (m, 1H), 2.68-2.57 (m, 1H), 2.50-2.47 (m, 1H), 2.44 (q, 2H, J = 7.2 Hz), 2.25-2.10 (m, 1H), 1.80-1.65 (m, 1H), 1.21 (t, 3H, J = 7.2 Hz), 1.04 (t, 3H, J = 7.2 Hz).






#	Structure	LCMS (M+1)	NMR
182		448	¹ H-NMR (400 MHz, MeOD) δ ppm 7.97 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.36 (d, 2H, J = 8.0 Hz), 6.90 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 4.18 (q, 2H, J = 7.2 Hz), 4.06-4.00 (m, 2H), 3.90-3.80 (m, 1H), 3.77-3.71 (m, 4H), 3.56-3.52 (m, 2H), 3.52-3.47 (m, 4H), 2.90-2.80 (m, 1H), 1.30 (t, 3H, J = 7.2 Hz), 1.10 (d, 6H, J = 6.0 Hz).
183		448	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (s, 1H), 7.84-7.81 (m, 3H), 6.80 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 4.0 Hz), 4.23-4.21 (m, 1H), 4.04-4.03 (m, 2H), 3.86-3.85 (m, 4H), 3.62-3.53 (m, 4H), 3.17-3.14 (m, 2H), 2.54 (q, 2H, J = 7.2 Hz), 2.25-2.12 (m, 2H), 2.12-2.04 (m, 7H), 1.17 (t, 3H, J = 7.2 Hz), 0.95-0.92 (m, 2H), 0.89-0.86 (m, 2H).
184		449	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.07 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.95 (s, 1H), 6.94 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.98-4.68 (m, 1H), 4.00-3.86 (m, 2H), 3.78-3.61 (m, 2H), 3.61-3.48 (m, 2H), 3.47-3.40 (m, 2H), 3.14-2.97 (m, 4H), 2.91-2.78 (m, 4H), 2.72-2.59 (m, 1H), 2.35-2.29 (m, 1H), 1.52-1.34 (m, 1H), 1.25-1.11 (m, 1H).
185		449	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.07 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.96 (s, 1H), 6.94 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.08-4.83 (m, 1H), 4.00-3.82 (m, 2H), 3.80-3.66 (m, 2H), 3.63-3.54 (m, 2H), 3.53-3.45 (m, 2H), 3.16-3.04 (m, 4H), 2.93-2.80 (m, 4H), 2.27-2.16 (m, 1H), 1.62-1.48 (m, 1H), 1.12-0.98 (m, 1H).
186		450	¹ H-NMR (500 MHz, CD ₃ OD) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.01 (dd, 1H, J = 8.5 Hz, 2.5 Hz), 7.96 (d, 1H, J = 2.0 Hz), 7.87 (d, 1H, J = 5.5 Hz), 7.00 (d, 1H, J = 8.5 Hz), 6.91 (d, 1H, J = 2.0 Hz), 6.03 (d, 1H, J = 5.5 Hz), 4.07-4.01 (m, 2H), 3.89-3.79 (m, 6H), 3.70-3.60 (m, 3H), 3.57-3.51 (m, 2H), 3.39-3.35 (m, 4H), 2.61-2.52 (m, 1H), 1.55-1.45 (m, 1H), 1.35-1.28 (m, 1H).

#	Structure	LCMS (M+1)	NMR
187		450	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.92 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.11 (s, 1H), 6.80 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.90-3.89 (m, 2H), 3.69-3.68 (m, 2H), 3.51 (s, 2H), 3.45-3.40 (m, 2H), 3.35-3.30 (m, 2H), 2.90-2.85 (m, 1H), 2.73-2.71 (m, 2H), 2.62-2.61 (m, 2H), 2.02-2.00 (m, 1H), 1.05 (d, 6H, J = 6.4 Hz), 0.77-0.71 (m, 4H).
188		451	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 6.62 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.60-3.50 (m, 4H), 3.50-3.40 (m, 4H), 3.11-3.05 (m, 2H), 3.03-2.95 (m, 2H), 2.75-2.60 (m, 1H), 2.50-2.44 (m, 2H), 2.25-2.20 (m, 1H), 2.08-1.95 (m, 2H), 1.83-1.74 (m, 1H), 1.60-1.53 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz).
189		451	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.88 (d, 2H, J = 8.4 Hz), 7.47 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 2.0 Hz), 6.62 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.60-3.52 (m, 4H), 3.48-3.42 (m, 4H), 3.42-3.38 (m, 1H), 3.23-3.18 (m, 1H), 3.12-3.05 (m, 2H), 2.98-2.90 (m, 2H), 2.30-2.20 (m, 1H), 2.15-2.00 (m, 2H), 1.98-1.85 (m, 1H), 1.83-1.75 (m, 1H), 1.04 (t, 3H, J = 7.6 Hz).
190		451	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 9.15-9.00 (m, 1H), 8.90-8.80 (m, 1H), 7.96 (d, 1H, J = 5.6 Hz), 7.95 (d, 1H, J = 1.2 Hz), 6.73 (d, 1H, J = 1.2 Hz), 6.03 (d, 1H, J = 5.6 Hz), 4.00-3.80 (m, 2H), 3.69-3.55 (m, 2H), 3.58-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.40-3.25 (m, 3H), 3.15-2.90 (m, 2H), 2.49 (s, 3H), 2.22-2.16 (m, 2H), 2.05-1.98 (m, 1H), 1.98-1.90 (m, 2H), 0.79-0.73 (m, 4H).
191		451	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.96 (d, 1H, J = 5.6 Hz), 6.97 (d, 1H, J = 1.6 Hz), 6.02 (d, 1H, J = 5.6 Hz), 3.95-3.92 (m, 2H), 3.73 (s, 2H), 3.73-3.68 (m, 2H), 3.58-3.54 (m, 2H), 3.50-3.45 (m, 2H), 2.97-2.92 (m, 1H), 2.81 (d, 2H, J = 4.4 Hz), 2.77 (d, 2H, J = 4.4 Hz), 2.02-1.98 (m, 1H), 1.07 (d, 6H, J = 6.8 Hz), 0.79-0.73 (m, 4H).






#	Structure	LCMS (M+1)	NMR
192		453	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.91 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.10 (s, 1H), 6.77 (d, 1H, J = 1.6 Hz), 6.59 (t, 1H, J = 5.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.64-3.62 (m, 2H), 3.51-3.48 (m, 4H), 3.40-3.38 (m, 4H), 3.08-3.05 (m, 2H), 2.90-2.88 (m, 1H), 2.61-2.58 (m, 2H), 2.50-2.48 (m, 2H), 1.05-1.00 (m, 9H).
193		454	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.95 (d, 1H, J = 5.6 Hz), 6.94 (s, 1H), 6.59 (t, 1H, J = 5.6 Hz), 6.02 (d, 1H, J = 5.6 Hz), 3.73 (s, 2H), 3.53-3.50 (m, 4H), 3.46-3.44 (m, 4H), 3.10-3.06 (quintet, 2H, J = 6.4 Hz), 2.93 (heptet, 1H, J = 6.8 Hz), 2.82-2.79 (m, 2H), 2.77-2.72 (m, 2H), 1.07 (d, 6H, J = 6.8 Hz), 1.03 (t, 3H, J = 7.2 Hz).
194		456	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.83 (d, 1H, J = 5.6 Hz), 7.75 (s, 2H), 7.63 (s, 1H), 6.52 (d, 1H, J = 1.6 Hz), 5.84 (d, 1H, J = 5.6 Hz), 4.49-4.40 (m, 1H), 3.93-3.88 (m, 4H), 3.53-3.45 (m, 4H), 3.41-3.22 (m, 2H), 3.10-2.89 (m, 2H), 2.70-2.45 (m, 2H), 1.80-1.75 (m, 1H), 1.06-1.03 (m, 2H), 0.89-0.80 (m, 2H).
195		457	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.45-4.35 (m, 1H), 4.01-3.92 (m, 2H), 3.90-3.80 (m, 1H), 3.75-3.65 (m, 2H), 3.62-3.52 (m, 3H), 3.50-3.40 (m, 2H), 2.95-2.85 (m, 2H), 2.80-2.70 (m, 2H), 2.55-2.51 (m, 2H), 2.15-2.00 (m, 1H), 1.50-1.40 (m, 1H), 1.37-1.30 (m, 1H).
196		457	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.33-8.29 (br., 1H), 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.93 (d, 1H, J = 1.6 Hz), 6.85 (d, 2H, J = 8.8 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.93-3.87 (m, 2H), 3.84-3.71 (m, 2H), 3.59-3.52 (m, 4H), 3.36-3.26 (m, 4H), 3.04-2.97 (m, 2H), 2.74-2.67 (m, 1H), 2.40-2.32 (m, 2H), 2.04-2.00 (m, 1H), 1.82-1.77 (m, 1H), 1.61-1.55 (m, 1H), 0.78-0.74 (m, 4H).






#	Structure	LCMS (M+1)	NMR
197		457	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.80-8.60 (br., 1H), 8.05 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.92 (d, 1H, J = 1.6 Hz), 6.49 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.83 (dd, 4H, J = 12.8, 7.2 Hz), 3.71-3.70 (m, 2H), 3.52-3.50 (m, 2H), 3.48-3.44 (m, 2H), 3.39-3.37 (m, 2H), 3.22 (t, 2H, J = 7.2 Hz), 2.22 (t, 2H, J = 7.2 Hz), 2.03-2.02 (m, 1H), 0.79-0.75 (m, 4H).
198		457	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.86 (s, 1H), 7.82 (d, 1H, J = 5.2 Hz), 7.54 (d, 2H, J = 8.4 Hz), 6.64 (s, 1H), 6.60 (d, 2H, J = 8.4 Hz), 5.83 (d, 1H, J = 5.2 Hz), 3.95-3.90 (m, 2H), 3.90-3.85 (m, 2H), 3.60-3.53 (m, 4H), 3.53-3.42 (m, 2H), 3.41-3.37 (m, 2H), 2.62-2.40 (br., 4H), 2.30-2.25 (m, 2H), 1.80-1.77 (m, 1H), 1.05-1.03 (m, 2H), 0.84-0.81 (m, 2H).
199		458	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.06 (s, 1H), 7.90-7.80 (m, 1H), 7.63 (d, 2H, J = 8.0 Hz), 6.94 (d, 2H, J = 8.0 Hz), 6.85 (s, 1H), 6.00-5.88 (m, 1H), 4.66 (br. s., 1H), 4.40-4.20 (m, 1H), 4.00-3.40 (m, 4H), 3.26-3.12 (m, 2H), 3.10-3.02 (m, 4H), 2.98-2.76 (m, 5H), 2.74-2.60 (m, 3H), 1.40-1.20 (m, 3H).
200		458	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.22 (s, 1H), 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 3.94-3.90 (m, 4H), 3.79-3.62 (m, 4H), 3.60-3.47 (m, 3H), 3.10-3.07 (m, 2H), 2.18-2.13 (m, 2H), 2.01-2.00 (m, 1H), 1.81-1.69 (m, 4H), 1.06 (t, 3H, J = 7.2 Hz), 0.80-0.73 (m, 4H).
201		458	





#	Structure	LCMS (M+1)	NMR
202		458	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.74 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.00-3.80 (m, 2H), 3.78-3.62 (m, 2H), 3.60-3.42 (m, 4H), 3.30-3.20 (m, 2H), 3.10-2.90 (m, 2H), 2.85-2.70 (m, 1H), 2.48-2.45 (m, 1H), 2.10-1.95 (m, 2H), 1.90-1.70 (m, 2H), 1.68-1.40 (m, 2H), 1.15-0.90 (m, 3H), 0.80-0.60 (m, 4H).
203		458	
204		458	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.5 Hz, 1H), 7.76 – 7.67 (m, 2H), 7.35 – 7.22 (m, 2H), 7.01 (d, J = 1.9 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 3.91 (s, 2H), 3.70 (s, 2H), 3.50 (d, J = 27.8 Hz, 5H), 3.25 (dt, J = 9.4, 7.5 Hz, 1H), 2.98 (t, J = 8.4 Hz, 1H), 2.83 – 2.62 (m, 2H), 2.44 – 2.31 (m, 1H), 2.26 – 2.11 (m, 0H), 2.01 (tt, J = 7.7, 4.9 Hz, 1H), 1.81 – 1.68 (m, 1H), 1.05 (t, J = 6.3 Hz, 6H), 0.76 (ddt, J = 9.8, 5.0, 2.5 Hz, 4H).
205		458	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (s, 1H), 7.92 (d, 1H, J = 5.6 Hz), 7.82 (d, 2H, J = 8.0 Hz), 7.40-7.35 (m, 2H), 7.05 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 3.99-3.97 (m, 1H), 3.69-3.38 (m, 6H), 3.63-3.62 (m, 5H), 3.50-3.49 (m, 2H), 3.30-3.25 (m, 1H), 2.34-2.32 (m, 3H), 2.22-2.20 (m, 1H), 1.25 (t, 3H, J = 7.2 Hz), 1.01-0.99 (m, 1H), 0.48-0.45 (m, 2H), 0.15-0.14 (m, 2H).
206		458	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.80 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.0 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 2H), 3.80-3.65 (m, 2H), 3.60-3.40 (m, 6H), 3.20-2.90 (m, 4H), 2.10-1.80 (m, 5H), 1.70-1.50 (m, 1H), 1.30-1.10 (m, 3H), 0.80-0.60 (m, 4H).

#	Structure	LCMS (M+1)	NMR
207		458	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.80 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.0 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 2H), 3.80-3.65 (m, 2H), 3.60-3.40 (m, 6H), 3.20-2.90 (m, 4H), 2.10-1.80 (m, 5H), 1.70-1.50 (m, 1H), 1.30-1.10 (m, 3H), 0.80-0.60 (m, 4H).
208		458	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.28 (d, 2H, J = 8.0 Hz), 7.01 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.90-3.88 (m, 2H), 3.68-3.66 (m, 2H), 3.54-3.50 (m, 2H), 3.49-3.44 (m, 2H), 3.31-3.27 (m, 1H), 2.93-2.89 (m, 1H), 2.68-2.62 (m, 2H), 2.50-2.48 (m, 1H), 2.46-2.42 (m, 2H), 2.26-2.18 (m, 1H), 1.76-1.72 (m, 2H), 1.18-1.12 (m, 1H), 1.09-1.03 (m, 6H), 0.99-0.95 (m, 1H), 0.60-0.56 (m, 1H).
209		459	
210		459	
211		459	

#	Structure	LCMS (M+1)	NMR
212		459	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.06 (s, 1H), 7.87 (d, 1H, J = 5.2 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.93 (s, 1H), 5.95 (d, 1H, J = 5.6 Hz), 3.68-3.36 (m, 9H), 3.17-3.15 (m, 4H), 2.50-2.47 (m, 4H), 2.23-2.11 (m, 7H), 1.93-1.90 (m, 1H), 1.78-1.76 (m, 1H).
213		459	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.27 (br. s., 1H), 8.07 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.2 Hz), 7.66 (d, 2H, J = 8.4 Hz), 6.97-6.95 (m, 3H), 5.96 (d, 1H, J = 5.6 Hz), 3.92-3.70 (m, 6H), 3.52-3.45 (m, 6H), 3.17-3.15 (m, 4H), 2.41-2.36 (q, 2H, J = 7.2 Hz), 2.03-2.00 (m, 1H), 1.06-1.02 (m, 3H), 0.78-0.72 (m, 4H).
214		460	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.0 Hz), 7.36 (d, 2H, J = 8.0 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.49-4.47 (m, 1H), 3.96-3.93 (m, 3H), 3.70-3.64 (m, 3H), 3.54-3.47 (m, 4H), 2.94-2.91 (m, 1H), 2.80-2.77 (m, 1H), 2.50-2.35 (m, 2H), 2.08-2.00 (m, 2H), 1.91-1.88 (m, 1H), 1.02 (t, 3H, J = 7.2 Hz), 0.78-0.72 (m, 4H).
215		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 7.03 (s, 1H), 6.00 (d, 1H, J = 5.6 Hz), 3.77-3.68 (m, 2H), 3.66-3.58 (m, 2H), 3.56-3.44 (m, 4H), 3.06-2.94 (m, 2H), 2.42 (s, 3H), 2.40-2.29 (m, 1H), 2.36 (q, 2H, J = 7.2 Hz), 2.08-1.90 (m, 2H), 1.83-1.72 (m, 2H), 1.72-1.58 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
216		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.8 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.96 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.07-5.03 (m, 1H), 3.95-3.90 (m, 2H), 3.73-3.69 (m, 2H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 3.35-3.12 (m, 4H), 2.88-2.82 (m, 2H), 2.42-2.33 (m, 1H), 2.05-2.03 (m, 1H), 2.00-1.97 (m, 1H), 1.14 (t, 3H, J = 7.2 Hz), 0.79-0.75 (m, 4H).





#	Structure	LCMS (M+1)	NMR
217		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.12 (d, 1H, J = 2.0 Hz), 7.89(d, 1H, J = 5.2 Hz), 7.76 (d, 2H, J = 8.8 Hz), 7.01 (d, 2H, J = 8.8 Hz), 6.99 (s, 1H), 5.97 (d, 1H, J = 5.2 Hz), 4.17 (d, 2H, J = 5.2 Hz), 4.05-3.99 (m, 2H), 3.95-3.90 (m, 2H), 3.90-3.78 (m, 2H), 3.74-3.65 (m, 2H), 3.58-3.48 (m, 2H), 3.47-3.42 (m, 2H), 3.20-3.10 (m, 1H), 3.10-3.04 (m, 2H), 2.04-2.0 (m, 1H), 1.06 (t, 3H, J = 6.8 Hz), 0.78-0.74 (m, 4H).
218		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (s, 1H), 8.11 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.97 (d, 2H, J = 8.4 Hz), 5.98(d, 1H, J = 5.6 Hz), 4.05-3.93 (m, 2H), 3.87-3.85 (m, 2H), 3.70-3.66 (m, 4H), 3.48-3.45 (m, 4H), 2.93-2.89 (m, 1H), 2.85-2.78 (m, 1H), 2.11-2.07 (m, 1H), 2.03-1.97 (m, 2H), 0.94 (t, 3H, J = 7.2 Hz), 0.78-0.75 (m, 4H).
219		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 10.50-9.84 (m, 1H), 8.28 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.90 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.11 (d, 1H, J = 2.0 Hz), 6.76 (s, 1H), 5.99 (d, 1H, J = 5.6 Hz), 4.66-4.22 (m, 2H), 4.27-4.11 (m, 2H), 3.95-3.90 (m, 2H), 3.74-3.68 (m, 2H), 3.55-3.50 (m, 5H), 2.06-1.99 (m, 1H), 1.20-1.16 (m, 6H), 0.79-0.75 (m, 4H).
220		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.10 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 6.97 (d, 1H, J = 1.6 Hz), 6.86 (d, 2H, J = 8.4 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.75-4.74 (m, 1H), 3.91-3.90 (m, 1H), 3.69 (dd, 4H, J = 7.6, 6.0 Hz), 3.53-3.48 (m, 2H), 3.46-3.41 (m, 2H), 2.93-2.89 (m, 2H), 2.32-2.29 (m, 2H), 2.02-2.01 (m, 1H), 0.87 (d, 6H, J = 6.0 Hz), 0.78-0.72 (m, 4H).
221		460	¹ H-NMR (400 MHz, DMSO-d6) δ ppm 8.61 (s, 1H), 8.13 (s, 1H), 7.98-7.96 (dd, 1H, J = 8.8, 1.6 Hz), 7.88 (d, 1H, J = 5.2 Hz), 7.01 (s, 1H), 6.88 (d, 1H, J = 5.2 Hz), 3.92 (br. s., 2H), 3.70 (br. s., 2H), 3.52-3.46 (m, 8H), 3.33 (br. s., 2H), 2.44-2.36 (m, 2H), 2.04-1.99 (m, 1H), 1.05 (t, 3H, J = 6.4 Hz), 0.78-0.74 (m, 4H).





#	Structure	LCMS (M+1)	NMR
222		460	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.29 (s, 1H), 8.16 (s, 1H), 7.90 (d, 1H, J = 4.8 Hz), 7.76 (d, 1H, J = 8.8 Hz), 7.36 (d, 1H, J = 8.8 Hz), 7.08 (s, 1H), 5.97 (d, 1H, J = 4.8 Hz), 3.94-3.91 (m, 2H), 3.73-3.68 (m, 2H), 3.56-3.52 (m, 2H), 3.49-3.45 (m, 2H), 3.27-3.19 (m, 4H), 2.50-2.40 (m, 4H), 2.39 (q, 2H, J = 7.2 Hz), 2.03-1.97 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.80-0.73 (m, 4H).
223		461	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.97 (s, 1H), 6.95 (d, 2H, J = 8.4 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.74-3.69 (m, 4H), 3.59-3.57 (m, 8H), 3.17-3.15 (m, 4H), 2.90-2.80 (m, 1H), 2.37 (q, 2H, J = 7.2 Hz), 1.04 (d, 6H, J = 7.2 Hz), 1.03 (t, 3H, J = 7.2 Hz).
224		461	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.60-3.47 (m, 4H), 3.46-3.38 (m, 4H), 3.30-3.20 (m, 1H), 3.12-3.03 (m, 2H), 3.02-2.94 (m, 1H), 2.80-2.64 (m, 2H), 2.49-2.44 (m, 1H), 2.43-2.35 (m, 1H), 2.26-2.15 (m, 1H), 1.80-1.70 (m, 1H), 1.10-1.00 (m, 9H).
225		461	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.52-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.31-3.25 (m, 1H), 3.12-3.05 (m, 2H), 3.01-2.97 (m, 1H), 2.74-2.68 (m, 2H), 2.47-2.38 (m, 2H), 2.23-2.19 (m, 1H), 1.78-1.73 (m, 1H), 1.07-1.01 (m, 9H).
226		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.04 Hz), 7.30 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 5.05-4.88 (m, 1H), 3.93-3.90 (m, 2H), 3.74-3.73 (m, 2H), 3.48-3.28 (m, 4H), 2.93-2.92 (m, 1H), 2.68-2.65 (m, 2H), 2.46-2.42 (m, 3H), 2.24-2.20 (m, 2H), 1.76-1.75 (m, 1H), 1.52-1.08 (m, 1H), 1.06 (t, 3H, J = 6.8 Hz).





#	Structure	LCMS (M+1)	NMR
227		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (s, 1H), 8.17 (s, 1H), 7.91 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.0 Hz), 7.05 (s, 1H), 5.99 (d, 1H, J = 5.2 Hz), 5.09-4.83 (m, 1H), 4.01-3.82 (m, 3H), 3.79-3.68 (m, 3H), 3.62-3.58 (m, 1H), 3.53-3.41 (m, 3H), 3.12 (t, 1H, J = 8.4 Hz), 2.90-2.80 (m, 2H), 2.72-2.59 (m, 3H), 1.89-1.78 (m, 1H), 1.62-1.50 (m, 1H), 1.11 (t, 3H, J = 7.2 Hz).
228		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 2.0 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 2.0 Hz), 5.97 (d, 1H, J = 5.2 Hz), 5.30-5.00 (m, 1H), 4.00-3.83 (m, 2H), 3.80-3.62 (m, 2H), 3.57-3.50 (m, 2H), 3.49-3.42 (m, 2H), 3.40-3.35 (m, 2H), 3.30-3.20 (m, 2H), 3.18-3.00 (m, 1H), 2.80-2.60 (m, 1H), 2.35-2.20 (m, 1H), 2.10-1.95 (m, 1H), 1.06 (t, 3H, J = 7.2 Hz), 0.80-0.60 (m, 4H).
229		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.21 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.44 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.72-3.60 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.38 (m, 4H), 3.37-3.34 (m, 1H), 3.08-3.05 (m, 1H), 3.04-2.97 (m, 2H), 2.70-2.61 (m, 1H), 2.27-2.20 (m, 2H), 2.20-2.08 (m, 4H), 2.03-1.98 (m, 1H), 1.95-1.90 (m, 1H), 1.85-1.80 (m, 1H), 1.80-1.74 (m, 1H), 1.59-1.54 (m, 1H).
230		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.66-3.55 (m, 4H), 3.49-3.41 (m, 4H), 3.03-2.93 (m, 2H), 2.35 (q, 2H, J = 7.2 Hz), 2.33-2.29 (m, 1H), 2.00-1.90 (m, 2H), 1.80-1.72 (m, 2H), 1.72-1.59 (m, 2H), 1.22 (t, 3H, J = 7.2 Hz), 1.02 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
231		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.28 (d, 2H, J = 8.0 Hz), 6.98 (d, 1H, J = 1.2 Hz), 5.95 (d, 1H, J = 5.2 Hz), 4.07 (q, 2H, J = 6.8 Hz), 3.65-3.55 (m, 4H), 3.50-3.42 (m, 4H), 3.30-3.20 (m, 1H), 2.98 (t, 1H, J = 8.4 Hz), 2.78-2.74 (m, 1H), 2.70-2.65 (m, 1H), 2.48 (t, 1H, J = 8.4 Hz), 2.45-2.37 (m, 1H), 2.25-2.10 (m, 1H), 1.80-1.65 (m, 1H), 1.20 (t, 3H, J = 6.8 Hz), 1.10-1.00 (m, 6H).
232		463	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 8.13 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.1 Hz, 2H), 7.24 (d, J = 8.1 Hz, 2H), 6.98 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.64 (s, 3H), 3.60 (t, J = 5.1 Hz, 4H), 3.45 (t, J = 5.1 Hz, 4H), 2.88 (d, J = 10.9 Hz, 3H), 2.70 (p, J = 6.6 Hz, 1H), 2.21 (t, J = 11.2 Hz, 2H), 1.76 (d, J = 12.4 Hz, 2H), 1.61 (tt, J = 12.2, 6.3 Hz, 2H), 0.99 (d, J = 6.5 Hz, 6H).
233		478	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.68 (s, 1H), 7.92 (s, 1H), 7.85 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.0 Hz), 6.68 (s, 1H), 5.85 (d, 1H, J = 5.6 Hz), 4.20 (q, 2H, J = 7.2 Hz), 3.77-3.71 (m, 4H), 3.60-3.55 (m, 1H), 3.54-3.50 (m, 1H), 3.49-3.41 (m, 4H), 3.40-3.38 (m, 1H), 3.21-3.28 (m, 1H), 3.18-3.11 (m, 1H), 3.05-2.94 (m, 1H), 2.40-2.50 (m, 1H), 2.20-2.10 (m, 1H), 1.36 (d, 6H, J = 6.4 Hz), 1.31 (t, 3H, J = 7.2 Hz).
234		464	
235		463	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.2 Hz), 8.15 (s, 1H), 7.89 (d, 1H, J = 5.2 Hz), 7.80 (d, 2H, J = 8.0 Hz), 7.61 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.6 Hz), 6.60 (t, 1H, J = 5.6 Hz), 6.00-5.97 (m, 2H), 3.72-3.70 (m, 2H), 3.52-3.51 (m, 4H), 3.41-3.44 (m, 6H), 3.11-3.04 (m, 2H), 2.66-2.50 (m, 1H), 1.03 (t, 3H, J = 7.6 Hz), 0.96 (d, 6H, J = 6.0 Hz).

#	Structure	LCMS (M+1)	NMR
236		464	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.10 (d, 1H, J = 2.0 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 6.97 (d, 1H, J = 2.0 Hz), 6.86 (d, 2H, J = 8.4 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.75-4.74 (m, 1H), 4.08 (q, 2H, J = 7.2 Hz), 3.71-3.68 (m, 2H), 3.59-3.58 (m, 4H), 3.43-3.40 (m, 4H), 3.35-3.30 (m, 1H), 2.95-2.90 (m, 2H), 1.21 (d, 3H, J = 7.2 Hz), 0.88 (d, 2H, J = 6.4 Hz).
237		464	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (s, 1H), 7.87 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.4 Hz), 7.00-6.90 (m, 3H), 5.95 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 2H), 3.75-3.60 (m, 2H), 3.58-3.48 (m, 2H), 3.47-3.40 (m, 2H), 3.20-3.00 (m, 4H), 2.48-2.40 (m, 4H), 2.10-1.90 (m, 1H), 0.80-0.65 (m, 4H).
238		466	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 4.2 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.0 Hz), 7.05 (s, 1H), 5.96 (d, 1H, J = 4.2 Hz), 3.92 (br. s., 2H), 3.70 (br. s., 2H), 3.54 (br. s., 2H), 3.47 (br. s., 2H), 3.26-3.08 (m, 3H), 3.01-2.98 (m, 1H), 2.89-2.77 (m, 1H), 2.65-2.59 (m, 1H), 2.05-1.96 (m, 2H), 1.79-1.75 (m, 1H), 0.77-0.74 (m, 4H).
239		467	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.94 (d, 1H, J = 5.6 Hz), 7.89 (d, 1H, J = 1.2 Hz), 6.83 (d, 1H, J = 1.2 Hz), 6.73 (d, 2H, J = 12.8 Hz), 3.95-3.90 (m, 2H), 3.70-3.65 (m, 2H), 3.57-3.52 (m, 2H), 3.47-3.43 (m, 2H), 3.15-3.12 (m, 4H), 2.82-2.79 (m, 4H), 2.03-1.99 (m, 1H), 0.78-0.73 (m, 4H).
240		467	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.4 Hz), 6.94 (d, 2H, J = 8.4 Hz), 6.91 (d, 1H, J = 1.6 Hz), 6.60 (t, 1H, J = 5.2 Hz), 5.96 (d, 1H, J = 5.2 Hz), 3.60-3.46 (m, 4H), 3.44-3.36 (m, 4H), 3.20-3.12 (m, 4H), 3.06 (quintet, 2H, J = 6.8 Hz), 2.48-2.40 (m, 4H), 1.02 (d, 3H, J = 7.2 Hz).






#	Structure	LCMS (M+1)	NMR
241		471	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.85 (d, 1H, J = 2.0 Hz), 7.82 (d, 1H, J = 5.2 Hz), 7.52 (d, 2H, J = 8.4 Hz), 6.63 (d, 1H, J = 2.0 Hz), 6.50 (d, 2H, J = 8.8 Hz), 5.83 (d, 1H, J = 5.2 Hz), 4.07-4.02 (m, 4H), 3.95-3.92 (m, 2H), 3.90-3.85 (m, 2H), 3.80-3.75 (m, 4H), 3.56-3.52 (m, 2H), 3.48-3.44 (m, 2H), 2.79 (q, 2H, J = 7.2 Hz), 1.80-1.75 (m, 1H), 1.16 (t, 3H, J = 7.2 Hz), 1.06-1.03 (m, 2H), 0.84-0.81 (m, 2H).
242		471	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.01 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.57 (d, 2H, J = 8.8 Hz), 6.88 (d, 1H, J = 1.6 Hz), 6.59 (d, 2H, J = 8.8 Hz), 5.95 (d, 1H, J = 5.6 Hz), 4.35-4.30 (m, 1H), 3.95-3.85 (m, 2H), 3.75-3.65 (m, 2H), 3.60-4.0 (m, 5H), 3.16-3.14 (m, 1H), 2.84-2.82 (m, 1H), 2.45-2.35 (m, 4H), 2.05-1.95 (m, 1H), 1.80 (q, 2H, J = 7.2 Hz), 0.93 (t, 3H, J = 7.2 Hz), 0.80-0.71 (m, 4H).
243		471	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (br s., 1H), 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.93 (d, 1H, J = 1.6 Hz), 5.95 (d, 1H, J = 5.2 Hz), 3.96-3.90 (m, 2H), 3.82-3.64 (m, 8H), 3.08-3.00 (m, 2H), 2.78-2.71 (m, 1H), 2.41 (t, 1H, J = 10.4 Hz), 2.27-2.21 (m, 1H), 2.11-2.00 (m, 3H), 1.86-1.80 (m, 1H), 1.76-1.66 (m, 2H), 1.43-1.32 (m, 1H), 0.78-0.74 (m, 4H).
244		471	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.37 (s, 1H), 8.02 (d, 1H, J = 2.0 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.63 (d, 2H, J = 8.8 Hz), 6.90 (d, 1H, J = 2.0 Hz), 6.54 (d, 2H, J = 8.8 Hz), 5.96 (d, 1H, J = 5.60 Hz), 3.92-3.91 (m, 2H), 3.71-3.70 (m, 2H), 3.52-3.49 (m, 2H), 3.47-3.44 (m, 2H), 3.35 (t, 2H, J = 7.2 Hz), 3.31-3.30 (m, 1H), 3.24-3.20 (m, 2H), 3.16 (t, 2H, J = 7.2 Hz), 3.01-3.00 (m, 2H), 2.05-1.97 (m, 3H), 1.88-1.85 (m, 2H), 0.78-0.74 (m, 4H).





#	Structure	LCMS (M+1)	NMR
245		472	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.55-4.52 (m, 1H), 3.93-3.91 (m, 3H), 3.70 (br. s., 2H), 3.53-3.47 (m, 4H), 3.15-3.09 (m, 1H), 2.78-2.73 (m, 1H), 2.61-2.50 (m, 1H), 2.04-2.00 (m, 4H), 1.82-1.76 (m, 2H), 1.63-1.59 (m, 1H), 1.47-1.43 (m, 1H), 0.77-0.74 (m, 4H).
246		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.72-3.68 (m, 2H), 3.55-3.50 (m, 2H), 3.47-3.42 (m, 2H), 2.89-2.85 (m, 2H), 2.73-2.67 (m, 1H), 2.46-2.42 (m, 1H), 2.21 (t, 2H, J = 3.2 Hz), 2.05-1.98 (m, 1H), 1.77-1.74 (m, 2H), 1.67-1.55 (m, 2H), 0.98 (d, 6H, J = 6.8 Hz), 0.80-0.71 (m, 4H).
247		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 3.93-3.92 (m, 2H), 3.71-3.70 (m, 2H), 3.54-3.50 (m, 2H), 3.49-3.43 (m, 2H), 2.81-2.79 (m, 2H), 2.74-2.10 (m, 1H), 2.67-2.63 (m, 1H), 2.19-2.14 (m, 2H), 2.04-2.00 (m, 1H), 1.81-1.80 (m, 1H), 1.71-1.70 (m, 1H), 1.59-1.50 (m, 1H), 1.45-1.30 (m, 1H), 0.98-0.92 (m, 6H), 0.79-0.74 (m, 4H).
248		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 3.93-3.92 (m, 2H), 3.71-3.70 (m, 2H), 3.54-3.50 (m, 2H), 3.49-3.43 (m, 2H), 2.81-2.79 (m, 2H), 2.74-2.10 (m, 1H), 2.67-2.63 (m, 1H), 2.19-2.14 (m, 2H), 2.04-2.00 (m, 1H), 1.81-1.80 (m, 1H), 1.71-1.70 (m, 1H), 1.59-1.50 (m, 1H), 1.45-1.30 (m, 1H), 0.98-0.92 (m, 6H), 0.79-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
249		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.70-3.64 (m, 2H), 3.60-3.54 (m, 2H), 3.45-3.40 (m, 4H), 3.40-3.38 (m, 1H), 3.30-3.28 (m, 1H), 2.98 (t, 1H, J = 8.0 Hz), 2.75-2.71 (m, 1H), 2.70-2.67 (m, 1H), 2.50-2.45 (m, 1H), 2.44-2.38 (m, 1H), 2.25-2.20 (m, 3H), 2.19-2.11 (m, 2H), 1.90-1.92 (m, 1H), 1.74-1.77 (m, 2H), 1.05 (t, 6H, J = 6.4 Hz).
250		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.69-3.66 (m, 2H), 3.57-3.54 (m, 2H), 3.46-3.42 (m, 4H), 3.40-3.33 (m, 2H), 3.10-3.00 (m, 2H), 2.58-2.50 (m, 2H), 2.19 (q, 2H, J = 7.2 Hz), 2.20-2.11 (m, 4H), 1.98-1.86 (m, 1H), 1.83-1.73 (m, 4H), 1.72-1.69 (m, 1H), 1.08 (t, 3H, J = 7.2 Hz).
251		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (s, 1H), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 7.02 (s, 1H), 5.97 (d, 1H, J = 5.2 Hz), 3.86-3.82 (m, 1H), 3.80-3.71 (m, 2H), 3.68-3.61 (m, 2H), 3.47-3.43 (m, 2H), 3.31-3.26 (m, 3H), 2.93-2.89 (m, 1H), 2.68-2.61 (m, 2H), 2.47-2.40 (m, 2H), 2.24-2.20 (m, 1H), 1.78-1.72 (m, 2H), 1.19 (s, 3H), 1.05 (t, 3H, J = 6.8 Hz), 0.97 (s, 3H), 0.96-0.93 (m, 1H), 0.68-0.65 (m, 1H).
252		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.31 (s, 1H), 8.14 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 6.94 (d, 1H, J = 1.6 Hz), 5.95 (d, 1H, J = 5.6 Hz), 4.65-4.64 (m, 1H), 4.20-4.16 (m, 1H), 3.94-3.87 (m, 2H), 3.75-3.63 (m, 1H), 3.25-3.23 (m, 2H), 3.01-2.98 (m, 2H), 2.54-2.49 (m, 1H), 2.38 (q, 2H, J = 7.2 Hz), 2.02-1.96 (m, 3H), 1.77-1.70 (m, 2H), 1.70-1.64 (m, 2H), 1.41-1.23 (m, 2H), 1.02 (t, 2H, J = 7.2 Hz), 0.81-0.71 (m, 4H).

#	Structure	LCMS (M+1)	NMR
253		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (s, 1H), 7.87 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 7.2 Hz), 7.33 (d, 2H, J = 7.2 Hz), 6.95 (s, 1H), 5.94 (d, 1H, J = 5.2 Hz), 4.68-4.62 (m, 1H), 4.22-4.14 (m, 1H), 3.97-3.84 (m, 2H), 3.74-3.50 (m, 2H), 3.30-3.00 (m, 4H), 2.47-2.40 (m, 2H), 2.35-2.25 (m, 1H), 2.05-1.95 (m, 1H), 1.90-1.83 (m, 1H), 1.45-1.10 (m, 3H), 1.00 (m, 6H), 0.82-0.70 (m, 4H).
254		472	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.70 (d, 2H, J = 8.0 Hz), 7.30 (d, 2H, J = 8.0 Hz), 6.94 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.6 Hz), 4.65-4.64 (m, 1H), 4.19-4.16 (m, 1H), 3.95-3.92 (m, 2H), 3.30-3.24 (m, 3H), 2.99-2.97 (m, 1H), 2.72-2.71 (m, 2H), 2.49-2.39 (m, 3H), 2.23-2.22 (m, 1H), 2.00-1.97 (m, 1H), 1.78-1.75 (m, 1H), 1.43-1.38 (m, 3H), 1.07-1.03 (m, 6H), 0.75-0.74 (m, 4H).
255		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.93 (d, 1H, J = 5.6 Hz), 7.87 (s, 1H), 7.41 (d, 1H, J = 7.2 Hz), 7.20 (s, 1H), 7.17 (d, 1H, J = 7.2 Hz), 6.77 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.00-3.80 (m, 2H), 3.75-3.60 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.35-3.30 (m, 2H), 3.20-2.99 (m, 2H), 2.80-2.60 (m, 2H), 2.43 (s, 3H), 2.33-2.18 (m, 1H), 2.06-1.94 (m, 1H), 1.89-1.70 (m, 1H), 1.21-0.92 (m, 6H), 0.83-0.71 (m, 4H).
256		472	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.94 (d, 1H, J = 5.6 Hz), 7.89 (d, 1H, J = 1.6 Hz), 7.48 (d, 1H, J = 8.0 Hz), 7.28 (s, 1H), 7.23 (d, 1H, J = 8.0 Hz), 6.78 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.75-3.70 (m, 2H), 3.60-3.53 (m, 2H), 3.50-3.40 (m, 2H), 3.35-3.30 (m, 4H), 3.30-3.00 (m, 1H), 2.45 (s, 3H), 2.45-2.25 (m, 2H), 2.20-1.91 (m, 2H), 1.30-0.14 (m, 6H), 0.83-0.71 (m, 4H).






#	Structure	LCMS (M+1)	NMR
257		473	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 7.8 Hz, 2H), 7.30 (t, J = 7.8 Hz, 2H), 7.02 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.92 (s, 1H), 3.70 (s, 2H), 3.60 – 3.43 (m, 6H), 2.91 (d, J = 11.4 Hz, 1H), 2.74 (q, J = 6.7 Hz, 1H), 2.20 – 2.06 (m, 1H), 2.01 (tt, J = 7.9, 3.8 Hz, 1H), 1.67 (dt, J = 25.8, 13.4 Hz, 3H), 1.47 (t, J = 11.7 Hz, 1H), 0.93 (d, J = 7.1 Hz, 4H), 0.83 – 0.66 (m, 8H).
258		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (s, 1H), 7.92 (d, 1H, J = 5.2 Hz), 7.86 (d, 2H, J = 7.6 Hz), 7.46 (d, 2H, J = 7.6 Hz), 7.11 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 4.10-3.90 (m, 2H), 3.80-3.57 (m, 2H), 3.70-3.57 (m, 2H), 3.55-3.48 (m, 2H), 3.40-3.35 (m, 2H), 3.25-3.20 (m, 1H), 3.10-3.00 (m, 1H), 2.95-2.90 (m, 1H), 2.81-2.74 (m, 1H), 2.20-2.08 (m, 2H), 2.05-1.97 (m, 2H), 1.93-1.78 (m, 1H), 1.74-1.62 (m, 1H), 1.50-1.43 (m, 1H), 1.40-1.28 (m, 1H).
259		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.81 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 3.91 (br. s, 2H), 3.70-3.65 (m, 2H), 3.55-3.50 (m, 4H), 3.48-3.45 (m, 2H), 3.14 (s, 2H), 2.76 (t, 2H, J = 5.2 Hz), 2.44 (q, 2H, J = 7.6 Hz), 1.99 (quintet, 1H, J = 5.2 Hz), 1.04 (t, 3H, J = 7.6 Hz), 0.77-0.72 (m, 4H).
260		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.08 (d, 1H, J = 1.2 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.67 (d, 2H, J = 8.8 Hz), 7.01 (d, 2H, J = 8.8 Hz), 6.95 (d, 1H, J = 1.2 Hz), 5.95 (d, 1H, J = 5.2 Hz), 4.40-4.30 (m, 4H), 4.00-3.80 (m, 2H), 3.78-3.62 (m, 2H), 3.58-3.50 (m, 2H), 3.48-3.42 (m, 2H), 3.26-3.20 (m, 2H), 3.05-2.95 (m, 2H), 2.80-2.75 (m, 2H), 2.10-1.90 (m, 1H), 0.80-0.60 (m, 4H).
261		473	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.88 (s, 1H), 7.83 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.98 (d, 2H, J = 8.4 Hz), 6.63 (s, 1H), 5.83 (d, 1H, J = 5.2 Hz), 3.94-3.89 (m, 4H), 3.55-3.47 (m, 4H), 3.31-3.28 (m, 4H), 2.80-2.74 (m, 5H), 1.82-1.75 (m, 1H), 1.15 (d, 6H, J = 5.2 Hz), 1.06-1.03 (m, 2H), 0.85-0.81 (m, 2H).

#	Structure	LCMS (M+1)	NMR
262		473	¹ H-NMR (500 MHz, DMSO-d ₆) δ ppm 8.06 (s, 1H), 7.86 (d, 1H, J = 5.5 Hz), 7.66-7.64 (m, 2H), 6.96-6.92 (m, 3H), 5.94 (d, 1H, J = 5.0 Hz), 3.68-3.66 (m, 2H), 3.56-3.54 (m, 2H), 3.45-3.39 (m, 6H), 3.32-3.30 (m, 2H), 3.18-3.14 (m, 4H), 2.43-2.34 (m, 2H), 2.24-2.11 (m, 5H), 1.94-1.89 (m, 1H), 1.79-1.75 (m, 1H), 1.04 (t, 3H, J = 7.0 Hz).
263		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 1.2 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.94 (s, 1H), 5.93 (d, 1H, J = 5.2 Hz), 4.53-4.10 (m, 3H), 3.72-3.39 (m, 6H), 3.29-2.96 (m, 6H), 2.36 (q, 2H, J = 7.2 Hz), 2.07-1.97 (m, 1H), 1.08-0.98 (m, 3H), 1.03 (t, 3H, J = 7.2 Hz), 0.79-0.74 (m, 4H).
264		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.64 (d, 2H, J = 8.4 Hz), 6.94 (d, 2H, J = 8.4 Hz), 6.93 (s, 1H), 5.95 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.70-3.69 (m, 2H), 3.51-3.45 (m, 6H), 2.88-2.80 (m, 1H), 2.80-2.72 (m, 2H), 2.47 (q, 2H, J = 7.2 Hz), 2.34-2.32 (m, 2H), 2.02-2.01 (m, 1H), 1.05 (d, 3H, J = 5.2 Hz), 0.98 (t, 3H, J = 7.2 Hz), 0.77-0.74 (m, 4H).
265		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.2 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.63 (d, 2H, J = 8.8 Hz), 6.94 (d, 2H, J = 8.8 Hz), 6.93 (s, 1H), 5.94 (d, 1H, J = 6.0 Hz), 3.91-3.90 (m, 2H), 3.70-3.69 (m, 2H), 3.49-3.45 (m, 6H), 2.88-2.76 (m, 3H), 2.49-2.47 (m, 2H), 2.33-2.28 (m, 2H), 2.01-1.99 (m, 1H), 1.05 (d, 3H, J = 4.8 Hz), 0.98 (t, 3H, J = 7.2 Hz), 0.77-0.73 (m, 4H).
266		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.47 (d, 1H, J = 1.6 Hz), 8.27 (d, 1H, J = 1.2 Hz), 7.94-7.90 (m, 2H), 7.67 (d, 1H, J = 6.8 Hz), 7.18 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.97-3.94 (m, 2H), 3.73-3.69 (m, 2H), 3.60-3.55 (m, 2H), 3.54-3.49 (m, 2H), 3.34-3.28 (m, 2H), 3.01-2.90 (m, 2H), 2.88-2.81 (m, 2H), 2.06-2.00 (m, 2H), 1.99-1.91 (m, 2H), 1.23-1.15 (m, 7H), 0.79-0.73 (m, 4H).

#	Structure	LCMS (M+1)	NMR
267		473	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.97 (s, 1H), 8.27 (s, 1H), 8.11 (dd, 1H, J = 8.0, 2.0 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.29 (d, 1H, J = 8.0 Hz), 7.14 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.69-3.68 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 2.97-2.81 (m, 2H), 2.75-2.61 (m, 2H), 2.33-2.08 (m, 2H), 2.06-2.02 (m, 1H), 1.90-1.80 (m, 2H), 1.79-1.61 (m, 2H), 1.08-1.00 (m, 6H), 0.77-0.74 (m, 4H).
268		474	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.86 (d, 1H, J = 1.6 Hz), 7.82 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.65 (d, 1H, J = 1.6 Hz), 5.82 (d, 1H, J = 5.2 Hz), 4.38 (quintet, 1H, J = 3.6 Hz), 3.99-3.90 (m, 2H), 3.90-3.75 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 2.83-2.72 (m, 2H), 2.50 (q, 2H, J = 7.2 Hz), 2.46-2.34 (m, 2H), 2.12-2.05 (m, 2H), 1.94-1.87 (m, 2H), 1.79-1.74 (m, 1H), 1.14 (t, 3H, J = 7.2 Hz), 1.06-1.01 (m, 2H), 0.84-0.78 (m, 2H).
269		474	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 2.0 Hz), 7.83 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.98 (d, 2H, J = 8.8 Hz), 6.65 (d, 1H, J = 2.0 Hz), 5.83 (d, 1H, J = 5.2 Hz), 4.46 (heptet, 1H, J = 4.4 Hz), 3.97-3.90 (m, 2H), 3.90-3.85 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.38 (m, 2H), 2.88-2.78 (m, 1H), 2.52 (q, 2H, J = 7.2 Hz), 2.22-2.08 (m, 4H), 1.89-1.82 (m, 1H), 1.81-1.75 (m, 1H), 1.71-1.64 (m, 1H), 1.55-1.43 (m, 1H), 1.12 (t, 3H, J = 7.2 Hz), 1.07-1.01 (m, 2H), 0.85-0.78 (m, 2H).
270		474	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.29 (s, 1H), 8.10 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.8 Hz), 6.98 (d, 1H, J = 1.6 Hz), 6.95 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.92-3.89 (m, 4H), 3.71-3.70 (m, 2H), 3.54-3.50 (m, 2H), 3.48-3.43 (m, 2H), 2.76-2.73 (m, 1H), 2.70-2.65 (m, 1H), 2.65-2.62 (m, 2H), 2.60 (q, 2H, J = 7.2 Hz), 2.51-2.49 (m, 1H), 2.02-2.00 (m, 1H), 1.99-1.95 (m, 1H), 1.59-1.56 (m, 1H), 1.06 (t, 3H, J = 7.2 Hz), 0.78-0.74 (m, 4H).





#	Structure	LCMS (M+1)	NMR
271		474	¹ H-NMR (400 MHz, MeOD) δ ppm 7.98 (d, 1H, J = 1.6 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.56 (d, 2H, J = 8.4 Hz), 6.94 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 4.04-3.98 (m, 2H), 3.85-3.78 (m, 2H), 3.80-3.75 (m, 1H), 3.72-3.65 (m, 2H), 3.65-3.58 (m, 2H), 3.55-3.40 (m, 4H), 2.62-2.47 (m, 1H), 2.40-2.26 (m, 1H), 2.06-1.94 (m, 1H), 1.45-1.35 (m, 6H), 0.95-0.83 (m, 4H).
272		474	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (s, 1H), 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 4.8 Hz), 7.64 (d, 2H, J = 8.4 Hz), 6.95 (d, 2H, J = 8.4 Hz), 6.91 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.2 Hz), 3.94 (t, 4H, J = 7.6 Hz), 3.45-3.38 (m, 10H), 3.17-3.15 (m, 6H), 2.38 (q, 2H, J = 7.2 Hz), 2.17 (quintet, 2H, J = 7.6 Hz), 1.04 (t, 3H, J = 7.2 Hz).
273		475	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.11 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.75-7.73 (m, 2H), 6.99-6.96 (m, 3H), 5.97 (d, 1H, J = 5.6 Hz), 4.11 (d, 1H, J = 5.6 Hz), 3.92 (br. s., 2H), 3.70 (br. s., 3H), 3.52-3.45 (m, 4H), 2.95 (br. s., 4H), 2.74 (t, 2H, J = 5.6 Hz), 2.61 (br. s., 4H), 2.06-1.99 (m, 1H), 0.78-0.75 (m, 4H).
274		475	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 4.8 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.94 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 4.44 (t, 1H, J = 5.2 Hz), 3.93-3.91 (m, 2H), 3.71-3.69 (m, 2H), 3.56-3.45 (m, 6H), 3.15 (t, 4H, J = 4.4 Hz), 2.56 (t, 4H, J = 4.4 Hz), 2.44 (t, 2H, J = 6.4 Hz), 2.04-2.00 (m, 1H), 0.78-0.72 (m, 4H).
275		475	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.12 (s, 1H), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.24 (d, 2H, J = 8.0 Hz), 6.98 (s, 1H), 6.59 (t, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.54-3.48 (m, 4H), 3.45-3.40 (m, 4H), 3.09-3.05 (m, 2H), 2.89-2.85 (m, 2H), 2.73-2.68 (m, 1H), 2.45-2.40 (m, 1H), 2.23-2.17 (m, 2H), 1.78-1.72 (m, 2H), 1.63-1.57 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz), 0.93 (d, 6H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
276		475	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.60 (t, 1H, J = 5.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.53-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.10-3.05 (m, 2H), 2.82-2.75 (m, 2H), 2.74-2.64 (m, 2H), 2.18-2.15 (m, 2H), 1.83-1.81 (m, 1H), 1.72-1.71 (m, 1H), 1.56-1.50 (m, 1H), 1.50-1.45 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.99-0.97 (m, 6H).
277		475	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.2 Hz), 6.62 (t, 1H, J = 5.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.52-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.10-3.05 (m, 2H), 2.82-2.75 (m, 2H), 2.73-2.67 (m, 2H), 2.19-2.16 (m, 2H), 1.83-1.81 (m, 1H), 1.75-1.72 (m, 1H), 1.54-1.50 (m, 1H), 1.49-1.44 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.99-0.97 (m, 6H).
278		476	¹ H-NMR (400 MHz, DMSO-d6) δ ppm 8.17 (s, 1H), 7.89 (d, 1H, J = 4.8 Hz), 7.75 (d, 2H, J = 7.6 Hz), 7.32 (d, 2H, J = 7.2 Hz), 7.04 (s, 1H), 5.96 (d, 1H, J = 4.8 Hz), 4.78-4.62 (m, 1H), 3.92 (br. s., 2H), 3.70 (br. s., 2H), 3.54 (br. s., 2H), 3.47 (br. s., 2H), 2.89-2.87 (m, 1H), 2.66-2.64 (m, 1H), 2.45-2.42 (m, 2H), 2.00-1.93 (m, 3H), 1.78-1.65 (m, 2H), 1.22 (s, 1H), 1.03 (t, 3H, J = 6.4 Hz), 0.76-0.74 (m, 4H).
279		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.89 (d, 1H, J = 4.8 Hz), 7.75 (d, 2H, J = 7.6 Hz), 7.32 (d, 2H, J = 7.2 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 4.8 Hz), 4.80-4.62 (m, 1H), 3.92 (br. s., 2H), 3.70 (br. s., 2H), 3.54 (br. s., 2H), 3.47 (br. s., 2H), 2.89-2.87 (m, 1H), 2.70-2.60 (m, 1H), 2.45-2.42 (m, 2H), 2.03-1.94 (m, 3H), 1.81-1.68 (m, 2H), 1.28 (s, 1H), 1.03 (t, 3H, J = 7.2 Hz), 0.77-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
280		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.89 (d, 1H, J = 4.8 Hz), 7.75 (d, 2H, J = 7.6 Hz), 7.32 (d, 2H, J = 7.2 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 4.8 Hz), 4.80-4.62 (m, 1H), 3.92 (s, 2H), 3.70 (s, 2H), 3.54 (s, 2H), 3.47 (s, 2H), 2.89-2.87 (m, 1H), 2.70-2.60 (m, 1H), 2.45-2.42 (m, 2H), 2.03-1.94 (m, 3H), 1.81-1.68 (m, 2H), 1.28 (s, 1H), 1.03 (t, 3H, J = 7.2 Hz), 0.77-0.74 (m, 4H).
281		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.90 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 7.04 (s, 1H), 5.99 (d, 1H, J = 5.2 Hz), 5.05-4.88 (m, 1H), 3.92-3.89 (m, 2H), 3.74-3.73 (m, 2H), 3.58-3.48 (m, 6H), 3.00-2.97 (m, 2H), 2.35 (q, 2H, J = 6.8 Hz), 2.22-2.20 (m, 1H), 1.99-1.96 (m, 2H), 1.74-1.65 (m, 4H), 1.03 (t, 3H, J = 6.8 Hz).
282		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.31 (d, 2H, J = 8.4 Hz), 7.04 (s, 1H), 5.96 (d, 1H, J = 5.2 Hz), 5.26-5.00 (m, 1H), 4.00-3.86 (m, 2H), 3.76-3.64 (m, 2H), 3.54-3.50 (m, 2H), 3.45-3.40 (m, 2H), 3.32-3.26 (m, 2H), 3.16-3.00 (m, 1H), 2.86-2.70 (m, 1H), 2.46-2.36 (m, 1H), 2.34-2.26 (m, 1H), 2.06-1.96 (m, 1H), 1.10-1.00 (m, 6H), 0.80-0.70 (m, 4H).
283		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.71-7.62 (m, 2H), 7.49-7.40 (m, 1H), 7.11 (m, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.98-3.88 (m, 2H), 3.76-3.66 (m, 2H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 3.43-3.33 (m, 2H), 3.32-3.28 (m, 1H), 3.00-2.67 (m, 3H), 2.30-2.23 (m, 1H), 2.06-1.98 (m, 1H), 1.96-1.90 (m, 1H), 1.21-1.10 (m, 6H), 0.80-0.74 (m, 4H).
284		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.68-7.64 (m, 1H), 7.61 (d, 1H, J = 8.0 Hz), 7.41 (t, 1H, J = 8.0 Hz), 7.10 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.95-3.91 (m, 2H), 3.73-3.65 (m, 2H), 3.60-3.50 (m, 2H), 3.49-3.41 (m, 2H), 3.32-3.27 (m, 1H), 3.10-2.92 (m, 1H), 2.82-2.74 (m, 2H), 2.64-2.57 (m, 1H), 2.28-2.19 (m, 1H), 2.04-1.98 (m, 1H), 1.86-1.78 (m, 1H), 1.12-1.04 (m, 6H), 0.79-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
285		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.28-8.27 (m, 2H), 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.90-4.70 (m, 1H), 3.70-3.52 (m, 4H), 3.50-3.35 (m, 4H), 3.10-2.90 (m, 2H), 2.44 (q, 2H, J = 7.2 Hz), 2.15-2.00 (m, 2H), 1.82-1.70 (m, 4H), 1.70-1.60 (m, 1H), 1.30-1.12 (m, 6H), 1.04 (t, 3H, J = 7.2 Hz).
286		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.12 (s, 1H), 7.87 (d, 1H, J = 5.6 Hz), 7.69 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 6.91 (s, 1H), 5.94 (d, 1H, J = 5.6 Hz), 4.31-4.30 (m, 1H), 4.09-4.07 (m, 2H), 3.89-3.85 (m, 2H), 3.85-3.83 (m, 1H), 3.22-3.21 (m, 2H), 3.03-3.00 (m, 1H), 3.00-2.97 (m, 2H), 3.50-2.45 (m, 1H), 2.35 (q, 2H, J = 7.2 Hz), 1.99-1.94 (m, 2H), 1.73-1.68 (m, 2H), 1.68-1.63 (m, 2H), 1.29-1.27 (d, 3H, J = 6.4 Hz), 1.20 (t, 3H, J = 7.2 Hz), 1.02 (t, 3H, J = 7.2 Hz).
287		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 6.97 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.60-3.59 (m, 4H), 3.46-3.43 (m, 4H), 2.80-2.75 (m, 2H), 2.72-2.67 (m, 2H), 2.18-2.08 (m, 2H), 1.82-1.78 (m, 1H), 1.73-1.68 (m, 1H), 1.58-1.51 (m, 1H), 1.47-1.40 (m, 1H), 1.21 (t, 3H, J = 7.2 Hz), 1.11-1.10 (m, 6H).
288		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 6.97 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.60-3.59 (m, 4H), 3.46-3.43 (m, 4H), 2.80-2.75 (m, 2H), 2.72-2.67 (m, 2H), 2.18-2.08 (m, 2H), 1.82-1.78 (m, 1H), 1.73-1.68 (m, 1H), 1.58-1.51 (m, 1H), 1.47-1.40 (m, 1H), 1.21 (t, 3H, J = 7.2 Hz), 1.15-1.00 (m, 6H).

#	Structure	LCMS (M+1)	NMR
289		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 4.8 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.94 (d, 2H, J = 8.8 Hz), 6.91 (d, 1H, J = 1.6 Hz), 6.60 (t, 1H, J = 5.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.52-3.51 (m, 4H), 3.43-3.41 (m, 4H), 3.15-3.14 (m, 4H), 3.10-3.07 (m, 2H), 2.68-2.66 (m, 1H), 2.59-2.58 (m, 4H), 1.03 (t, 3H, J = 7.2 Hz), 1.00 (d, 2H, J = 6.8 Hz).
290		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.44 (d, 1H, J = 2.0 Hz), 8.23 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.84 (d, 1H, J = 8.4 Hz), 7.67 (dd, 1H, J = 8.4, 2.0 Hz), 7.13 (d, 1H, J = 1.2 Hz), 6.60 (t, 1H, J = 5.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.53-3.50 (m, 4H), 3.45-3.42 (m, 4H), 3.33-3.31 (m, 1H), 3.11-3.04 (m, 2H), 2.90-2.87 (m, 2H), 2.73-2.69 (m, 1H), 2.24-2.19 (m, 2H), 1.78-1.75 (m, 2H), 1.70-1.63 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz), 0.98 (d, 6H, J = 6.8 Hz).
291		477	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.95 (d, 2H, J = 8.4 Hz), 6.94 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.05-4.80 (m, 1H), 4.00-3.80 (m, 2H), 3.78-3.65 (m, 2H), 3.60-3.50 (m, 1H), 3.49-3.35 (m, 3H), 3.30-3.21 (m, 4H), 3.20-3.05 (m, 4H), 2.36 (q, 2H, J = 7.2 Hz), 2.30-2.10 (m, 1H), 1.60-1.40 (m, 1H), 1.10-1.00 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz).
292		478	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.2 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.95 (d, 2H, J = 8.4 Hz), 6.95-6.93 (m, 1H), 5.96 (d, 1H, J = 5.2 Hz), 4.95-4.70 (m, 1H), 4.00-3.85 (m, 2H), 3.70-3.65 (m, 2H), 3.60-3.40 (m, 4H), 3.30-3.25 (m, 4H), 3.18-3.10 (m, 4H), 2.70-2.60 (m, 1H), 2.36 (q, 2H, J = 7.2 Hz), 1.50-1.30 (m, 1H), 1.25-1.10 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
298		483	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.90 (m, 2H), 3.75-3.65 (m, 2H), 3.60-3.55 (m, 2H), 3.54-3.50 (m, 2H), 3.30-3.20 (m, 1H), 3.05-2.97 (m, 1H), 2.95-2.90 (m, 1H), 2.80-2.60 (m, 2H), 2.45-2.35 (m, 2H), 2.25-2.15 (m, 1H), 2.13-2.05 (m, 1H), 1.80-1.70 (m, 1H), 1.50-1.40 (m, 1H), 1.38-1.30 (m, 1H), 1.10-1.00 (m, 6H).
299		483	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.96-3.94 (m, 2H), 3.71-3.70 (m, 2H), 3.58-3.55 (m, 2H), 3.49-3.47 (m, 2H), 3.29-3.25 (m, 1H), 3.02-2.96 (m, 1H), 2.95-2.90 (m, 1H), 2.73-2.69 (m, 2H), 2.50-2.49 (m, 2H), 2.45-3.39 (m, 1H), 2.24-2.19 (m, 1H), 2.18-2.08 (m, 1H), 1.79-1.74 (m, 1H), 1.49-1.46 (m, 1H), 1.36-1.33 (m, 1H), 1.06 (t, 3H, J = 6.4 Hz).
300		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.81 (d, 2H, J = 8.4 Hz), 7.38 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.22 (t, 2H, J = 8.8 Hz), 4.11-4.08 (m, 2H), 3.80-3.73 (m, 4H), 3.55-3.30 (s, 9H), 3.25-3.20 (m, 2H), 3.10-3.04 (m, 1H), 2.40-2.42 (m, 1H), 2.10-2.00 (br., 1H), 1.27 (t, 3H, J = 7.2 Hz).
301		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.08 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.67 (d, 2H, J = 8.4 Hz), 6.99-6.95 (m, 3H), 5.99 (d, 1H, J = 5.6 Hz), 3.96-3.88 (m, 2H), 3.75-3.66 (m, 1H), 3.65-3.52 (m, 2H), 3.50-3.45 (m, 1H), 3.41-3.36 (m, 1H), 3.23-3.07 (m, 4H), 2.68-2.62 (m, 2H), 2.58-2.54 (m, 1H), 2.22-2.15 (m, 1H), 1.54-1.50 (m, 1H), 1.39-1.34 (m, 1H), 1.26-1.22 (m, 1H), 1.12-1.01 (m, 3H).






#	Structure	LCMS (M+1)	NMR
302		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.09 (s, 1H), 7.88 (d, 1H, J = 5.2 Hz), 7.68 (d, 2H, J = 8.4 Hz), 6.98 (d, 2H, J = 8.4 Hz), 6.97 (s, 1H), 5.98 (d, 1H, J = 5.2 Hz), 4.00-3.91 (m, 2H), 3.75-3.66 (m, 2H), 3.61-3.51 (m, 3H), 3.50-3.43 (m, 2H), 3.21-3.08 (m, 4H), 2.96-2.91 (m, 1H), 2.13-2.08 (m, 1H), 1.50-1.44 (m, 1H), 1.38-1.32 (m, 1H), 1.27-1.21 (m, 1H), 1.17-1.02 (m, 3H).
303		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.70 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 6.97 (d, 1H, J = 1.2 Hz), 5.94 (d, 1H, J = 5.6 Hz), 4.21 (t, 2H, J = 8.4 Hz), 4.08 (dd, 2H, J = 7.2, 6.0 Hz), 3.78-3.71 (m, 1H), 3.46-3.45 (m, 8H), 3.37-3.24 (m, 2H), 2.91 (t, 1H, J = 8.4 Hz), 2.71-2.59 (m, 2H), 2.49-2.40 (m, 2H), 2.27-2.18 (m, 1H), 1.79-1.71 (m, 1H), 1.05 (t, 3H, J = 7.2 Hz).
304		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 6.98 (d, 1H, J = 1.2 Hz), 5.94 (d, 1H, J = 5.6 Hz), 4.21 (t, 2H, J = 8.4 Hz), 4.08 (dd, 2H, J = 7.2, 6.0 Hz), 3.78-3.71 (m, 1H), 3.46-3.45 (m, 8H), 3.32-3.26 (m, 2H), 2.96 (t, 1H, J = 8.4 Hz), 2.75-2.64 (m, 2H), 2.50-2.44 (m, 2H), 2.28-2.19 (m, 1H), 1.81-1.72 (m, 1H), 1.06 (t, 3H, J = 7.2 Hz).
305		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.2 Hz), 3.92-3.91 (m, 2H), 3.70-3.69 (m, 2H), 3.53-3.50 (m, 2H), 3.49-3.45 (m, 2H), 3.35-3.30 (m, 1H), 2.80-2.78 (m, 2H), 2.69-2.66 (m, 2H), 2.06-1.95 (m, 3H), 1.79-1.68 (m, 6H), 1.62-1.58 (m, 2H), 1.58-1.55 (m, 1H), 0.77-0.74 (m, 4H).





#	Structure	LCMS (M+1)	NMR
306		485	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.08 (d, J = 1.7 Hz, 1H), 7.74 – 7.62 (m, 3H), 7.32 – 7.18 (m, 3H), 5.60 (d, J = 5.9 Hz, 1H), 4.83 (s, 1H), 4.40 (s, 1H), 4.22 (d, J = 11.6 Hz, 2H), 3.98 (d, J = 11.5 Hz, 1H), 3.85 (d, J = 11.3 Hz, 1H), 2.87 (d, J = 10.9 Hz, 2H), 2.69 (q, J = 7.2 Hz, 2H), 2.48 – 2.38 (m, 1H), 2.20 (t, J = 11.3 Hz, 2H), 1.74 (q, J = 7.1, 4.5 Hz, 3H), 1.69 – 1.48 (m, 3H), 0.98 (d, J = 6.6 Hz, 6H), 0.91 – 0.72 (m, 2H), 0.71 (s, 1H), 0.51 (s, 1H).
307		485	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.30 (br. s., 1H), 8.06 (s, 1H), 7.87 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.0 Hz), 6.95 (d, 2H, J = 8.0 Hz), 6.93 (s, 1H), 5.96 (d, 1H, J = 5.2 Hz), 3.95-3.90 (br., 2H), 3.71-3.45 (m, 8H), 2.79-2.66 (m, 2H), 2.38-2.33 (m, 1H), 2.27-2.20 (m, 1H), 2.03-1.92 (m, 2H), 1.74-1.45 (m, 4H), 1.31-1.08 (m, 4H), 0.77-0.74 (m, 4H).
308		485	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 8.03 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.8 Hz), 6.00 (d, 1H, J = 1.6 Hz), 6.45 (d, 2H, J = 8.8 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.96-3.89 (m, 6H), 3.72-3.68 (m, 2H), 3.45-3.45 (m, 9H), 2.04-2.00 (m, 1H), 0.91 (d, 6H, J = 6.0 Hz), 0.78-0.74 (m, 4H).
309		486	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.11 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.4 Hz), 6.99 (d, 2H, J = 8.4 Hz), 6.94 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.48-4.41 (m, 1H), 4.22 (d, 2H, J = 8.4 Hz), 4.09 (dd, 2H, J = 8.4, 6.4 Hz), 3.80-3.72 (m, 1H), 3.47-3.45 (m, 8H), 3.21-3.18 (m, 1H), 2.92-2.86 (m, 1H), 2.82-2.67 (m, 2H), 2.01-1.96 (m, 1H), 1.80-1.72 (m, 1H), 1.68-1.60 (m, 1H), 1.57-1.43 (m, 2H).
310		486	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.16 (s, 1H), 7.90 (d, 2H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.03 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 4.55 (t, 2H, J = 6.4 Hz), 4.45 (t, 2H, J = 6.4 Hz), 3.93 (br. s., 2H), 3.71 (br. s., 2H), 3.54-3.47 (m, 4H), 3.42-3.35 (m, 2H), 2.82-2.79 (m, 2H), 2.06-2.00 (m, 1H), 1.86-1.64 (m, 6H), 0.78-0.75 (m, 4H).





#	Structure	LCMS (M+1)	NMR
311		486	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.73-3.71 (m, 2H), 3.64-3.62 (m, 2H), 3.48-3.46 (m, 4H), 3.30-3.25 (m, 1H), 2.96-2.92 (m, 1H), 2.73-2.65 (m, 2H), 2.46-2.42 (m, 1H), 2.25-2.19 (m, 1H), 2.09 (s, 3H), 2.03-1.92 (m, 1H), 1.80-1.72 (m, 1H), 1.49-1.46 (m, 2H), 1.34-1.31 (m, 2H), 1.06 (t, 3H, J = 7.2 Hz).
312		486	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.65-3.66 (m, 2H), 3.55-3.52 (m, 2H), 3.46-3.43 (m, 4H), 3.42-3.39 (m, 1H), 2.85-2.75 (m, 2H), 2.73-2.65 (m, 2H), 2.22-2.17 (m, 2H), 2.16-2.09 (m, 4H), 1.95-1.87 (m, 1H), 1.85-1.78 (m, 1H), 1.77-1.69 (m, 2H), 1.60-1.50 (m, 1H), 1.50-1.39 (m, 1H), 0.95-0.98 (m, 6H).
313		486	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.23 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.70-3.68 (m, 2H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 2.82-2.78 (m, 2H), 2.50-2.45 (m, 2H), 2.40-2.35 (m, 1H), 2.22-2.17 (m, 1H), 2.03-1.98 (m, 1H), 1.78-1.72 (m, 2H), 1.67-1.60 (m, 2H), 1.57-1.49 (m, 1H), 1.30-1.22 (m, 1H), 0.92 (d, 3H, J = 6.8 Hz), 0.86 (t, 3H, J = 7.6 Hz), 0.78-0.71 (m, 4H).
314		487	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.88 (d, 1H, J = 1.6 Hz), 7.83 (d, 1H, J = 5.2 Hz), 7.57 (d, 2H, J = 8.8 Hz), 6.98 (d, 2H, J = 8.8 Hz), 6.66 (d, 1H, J = 0.8 Hz), 5.83 (d, 1H, J = 5.2 Hz), 4.73-4.66 (m, 4H), 3.95-3.88 (m, 4H), 3.59-3.46 (m, 5H), 3.29-3.27 (m, 4H), 2.54-2.52 (m, 4H), 1.81-1.76 (m, 1H), 1.07-1.03 (m, 2H), 0.85-0.81 (m, 2H).

#	Structure	LCMS (M+1)	NMR
315		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 8.07 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 4.8 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.96 (d, 2H, J = 8.8 Hz), 6.94 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 3.95-3.90 (br., 2H), 3.79-3.65 (m, 5H), 3.57-3.46 (m, 6H), 3.20-3.15 (m, 1H), 2.84-2.67 (m, 3H), 2.33-2.20 (m, 4H), 2.04-2.00 (m, 1H), 0.79-0.74 (m, 4H).
316		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.63 (d, 2H, J = 8.8 Hz), 6.94-6.91 (m, 3H), 5.95 (d, 1H, J = 5.6 Hz), 3.95-3.85 (m, 2H), 3.75-3.65 (m, 2H), 3.60-3.40 (m, 4H), 3.14-3.10 (m, 2H), 2.88 (s, 2H), 2.63-2.59 (m, 2H), 2.39 (q, 2H, J = 7.2 Hz), 2.05-1.95 (m, 1H), 1.05 (s, 6H), 0.99 (t, 3H, J = 7.2 Hz), 0.80-0.70 (m, 4H).
317		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.46-3.45 (m, 4H), 3.41-3.40 (m, 4H), 3.34-3.31 (m, 4H), 3.01-2.98 (m, 2H), 2.36 (q, 2H, J = 7.2 Hz), 2.01-1.95 (m, 2H), 1.80-1.72 (m, 8H), 1.70-1.64 (m, 1H), 1.02 (t, 3H, J = 7.2 Hz).
318		487	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.08 (d, 1H, J = 1.0 Hz), 7.86 (d, 1H, J = 5.5 Hz), 7.63 (d, 2H, J = 8.5 Hz), 6.96 (d, 1H, J = 8.5 Hz), 6.87 (d, 1H, J = 2.0 Hz), 5.95 (d, 1H, J = 5.5 Hz), 4.66-4.65 (m, 1H), 4.21-4.16 (m, 1H), 3.95-3.93 (m, 1H), 3.93-3.90 (m, 1H), 3.22-3.19 (m, 2H), 3.19-3.09 (m, 4H), 2.70-2.65 (m, 1H), 2.60-2.54 (m, 4H), 2.01-1.96 (m, 1H), 1.43-1.35 (m, 2H), 1.30-1.21 (m, 2H), 1.06-0.96 (m, 6H), 0.82-0.71 (m, 4H).
319		488	¹ H NMR (400 MHz, Methanol-d ₄) δ ppm 7.90 (d, J = 1.8 Hz, 1H), 7.72 (d, J = 5.4 Hz, 1H), 7.67 – 7.59 (m, 2H), 7.30 – 7.20 (m, 2H), 6.82 (d, J = 1.8 Hz, 1H), 5.72 (d, J = 5.5 Hz, 1H), 4.73 (d, J = 33.7 Hz, 2H), 3.97 (d, J = 11.7 Hz, 2H), 3.44 (d, J = 11.8 Hz, 2H), 3.15 – 3.00 (m, 4H), 2.95 (q, J = 7.0 Hz, 1H), 2.79 (p, J = 6.5 Hz, 1H), 2.54 (tt, J = 12.1, 4.3 Hz, 1H), 2.36 (td, J = 11.8, 2.8 Hz, 3H), 1.90 – 1.67 (m, 5H), 1.12 (dd, J = 6.7, 3.8 Hz, 7H), 1.02 (t, J = 7.2 Hz, 2H).





#	Structure	LCMS (M+1)	NMR
320		488	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 6.0 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.2 Hz), 5.10 (d, 1H, J = 6.0 Hz), 4.20-4.00 (m, 1H), 3.70-3.60 (m, 2H), 3.58-3.50 (m, 2H), 3.48-3.35 (m, 4H), 3.30-3.15 (m, 2H), 3.05-2.95 (m, 1H), 2.80-2.60 (m, 2H), 2.47-2.30 (m, 4H), 2.25-2.15 (m, 1H), 2.10-2.00 (m, 2H), 1.80-1.70 (m, 1H), 1.10-1.00 (m, 6H).
321		488	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.92 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.59 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.0 Hz), 6.68 (d, 1H, J = 1.6 Hz), 5.84 (d, 1H, J = 5.2 Hz), 4.54-4.48 (m, 1H), 3.92-3.83 (m, 2H), 3.65-3.57 (m, 2H), 3.50-3.42 (m, 4H), 3.41-3.37 (m, 1H), 3.35-3.25 (m, 2H), 3.11-3.00 (m, 1H), 2.81-2.72 (m, 1H), 2.69-2.63 (m, 2H), 2.59-2.51 (m, 2H), 2.40-2.32 (m, 1H), 2.28-2.21 (m, 2H), 1.99-1.94 (m, 2H), 1.20-1.16 (dd, 6H, J = 6.4, 1.2 Hz).
322		488	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.36 (s, 1H), 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.86 (quintet, 1H, J = 7.2 Hz), 3.60-3.52 (m, 4H), 3.50-3.40 (m, 4H), 3.10-2.90 (m, 2H), 2.48-2.42 (m, 1H), 2.36 (q, 2H, J = 7.2 Hz), 2.30-2.20 (m, 2H), 2.10-1.90 (m, 4H), 1.80-1.62 (m, 4H), 1.60-1.50 (m, 2H), 1.02 (t, 3H, J = 7.2 Hz).
323		488	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26-8.17 (m, 1H), 7.92-7.88 (m, 1H), 7.77-7.72 (m, 2H), 7.57 (d, 2H, J = 7.6 Hz), 7.05 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.56-4.44 (m, 1H), 4.00-3.80 (m, 2H), 3.80-3.60 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.31-3.00 (m, 1H), 2.90-2.75 (m, 1H), 2.70-2.51 (m, 1H), 2.45-2.38 (m, 1H), 2.05-1.98 (m, 2H), 1.90-1.79 (m, 2H), 1.58-1.50 (m, 1H), 1.50-1.41 (m, 1H), 1.03-0.98 (m, 6H), 0.79-0.73 (m, 4H).

#	Structure	LCMS (M+1)	NMR
324		488	¹ H-NMR (400 MHz, MeOD) δ ppm 7.89 (s, 1H), 7.75 (d, 1H, J = 5.6 Hz), 7.66 (d, 1H, J = 8.8 Hz), 7.44 (d, 1H, J = 8.8 Hz), 6.84 (s, 1H), 5.90 (d, 1H, J = 5.6 Hz), 3.93 (m, 2H), 3.75 (m, 2H), 3.53-3.33 (m, 9H), 2.28-2.19 (m, 2H), 1.99-1.94 (m, 2H), 1.91 (m, 1H), 1.33 (d, 6H, J = 6.4 Hz), 0.85-0.83 (m, 2H), 0.78-0.74 (m, 2H).
325		488	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.0 Hz), 7.56 (d, 2H, J = 8.0 Hz), 7.05 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.53-4.48 (m, 1H), 4.00-3.80 (m, 2H), 3.80-3.65 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.33-3.25 (m, 1H), 2.85-2.51 (m, 2H), 2.50-2.30 (m, 2H), 2.05-1.95 (m, 1H), 1.90-1.70 (m, 2H), 1.62-1.55 (m, 1H), 1.55-1.48 (m, 1H), 1.15-0.85 (m, 6H), 0.79-0.70 (m, 4H).
326		488	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (s, 1H), 7.91 (d, 1H, J = 5.6 Hz), 7.89-7.65 (m, 2H), 7.57 (d, 2H, J = 8.0 Hz), 7.06 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 2H), 3.80-3.65 (m, 2H), 3.60-3.50 (m, 2H), 3.50-3.40 (m, 2H), 3.33-3.25 (m, 1H), 2.50-2.10 (m, 4H), 2.05-1.40 (m, 5H), 1.30-0.80 (m, 6H), 0.79-0.70 (m, 4H).
327		488	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.65-7.63 (m, 2H), 6.96-6.94 (m, 2H), 6.91 (d, 1H, J = 1.2 Hz), 5.95 (d, 1H, J = 5.6 Hz), 3.44-3.40 (m, 4H), 3.39-3.35 (m, 8H), 3.16 (br. s., 4H), 2.56 (br. s., 4H), 2.49 (br. s., 2H), 1.76 (br. s., 4H), 1.05 (d, 3H, J = 7.2 Hz).
328		488	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (s, 1H), 8.02 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.61 (d, 2H, J = 8.4 Hz), 6.88 (s, 1H), 6.61 (t, 1H, J = 5.6 Hz), 6.44 (d, 2H, J = 8.4 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.87 (s, 4H), 3.55-3.50 (m, 4H), 3.48-3.45 (m, 4H), 3.32 (s, 4H), 3.15-3.04 (m, 2H), 2.34-2.32 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz), 0.87 (d, 6H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
329		489	
330		489	
331		489	
332		489	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.85 (d, 1H, J = 1.6 Hz), 7.82 (d, 1H, J = 5.2 Hz), 7.51 (d, 2H, J = 8.4 Hz), 6.62 (d, 1H, J = 1.6 Hz), 6.51 (d, 1H, J = 8.4 Hz), 5.83 (d, 1H, J = 5.2 Hz), 4.93-4.70 (m, 1H), 4.13-4.05 (m, 1H), 4.03-3.95 (m, 1H), 3.93-3.84 (m, 1H), 3.80-3.71 (m, 1H), 3.68-3.60 (m, 1H), 3.58-3.50 (m, 1H), 3.47-3.41 (m, 1H), 3.38 (s, 1H), 3.38-3.32 (m, 1H), 2.46 (q, 2H, J = 7.2 Hz), 1.97-1.95 (m, 1H), 1.95-1.90 (m, 1H), 1.14-1.06 (m, 1H), 0.98 (t, 3H, J = 7.2 Hz).

#	Structure	LCMS (M+1)	NMR
333		489	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.56 (s, 1H), 7.86 (d, 1H, J = 1.6 Hz), 7.83 (d, 1H, J = 5.2 Hz), 7.52 (d, 2H, J = 8.4 Hz), 6.63 (d, 1H, J = 1.6 Hz), 6.51 (d, 2H, J = 8.4 Hz), 5.84 (d, 1H, J = 5.2 Hz), 4.93-4.70 (m, 1H), 4.13-4.05 (m, 1H), 4.04-4.01 (m, 4H), 4.00-3.95 (m, 1H), 3.94-3.83 (m, 1H), 3.80-3.75 (m, 1H), 3.72-3.70 (m, 4H), 3.68-3.60 (m, 1H), 3.59-3.50 (m, 1H), 3.48-3.45 (m, 1H), 3.42-3.39 (m, 1H), 2.72 (q, 2H, J = 7.2 Hz), 1.98-1.92 (m, 1H), 1.92-1.87 (m, 1H), 1.15-1.05 (m, 4H).
334		489	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.88 (d, 1H, J = 2.0 Hz), 7.82 (d, 1H, J = 5.2 Hz), 7.55 (d, 2H, J = 8.8 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.66 (d, 1H, J = 2.0 Hz), 5.83 (d, 1H, J = 6.0 Hz), 3.94-3.86 (m, 4H), 3.57 (t, 2H, J = 5.6 Hz), 3.58-3.46 (m, 4H), 3.38 (s, 3H), 3.27 (t, 4H, J = 5.2 Hz), 2.70 (t, 4H, J = 5.2 Hz), 2.67 (t, 2H, J = 5.6 Hz), 1.80-1.76 (m, 1H), 1.06-1.03 (m, 2H), 0.84-0.81 (m, 2H).
335		489	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 1.6 Hz), 7.82 (d, 1H, J = 5.2 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.98 (d, 2H, J = 8.8 Hz), 6.63 (d, 1H, J = 1.6 Hz), 5.82 (d, 1H, J = 5.2 Hz), 4.24-4.21 (m, 1H), 3.88-3.84 (m, 2H), 3.66-3.62 (m, 2H), 3.45-3.40 (m, 4H), 3.37-3.30 (m, 4H), 2.82-2.78 (m, 1H), 2.78-2.73 (m, 4H), 2.65-2.57 (m, 4H), 2.26-2.23 (m, 2H), 1.26-1.20 (m, 3H).
336		489	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 2.0 Hz), 7.82 (d, 1H, J = 5.6 Hz), 7.56 (d, 2H, J = 8.8 Hz), 6.98 (d, 2H, J = 8.8 Hz), 6.63 (d, 1H, J = 2.0 Hz), 5.82 (d, 1H, J = 5.6 Hz), 4.51 (quintet, 1H, J = 6.4 Hz), 3.88-3.85 (m, 2H), 3.62-3.58 (m, 2H), 3.45-3.42 (m, 4H), 3.30-3.25 (m, 5H), 2.69-2.62 (m, 6H), 2.50 (d, 2H, J = 7.2 Hz), 2.27-2.19 (m, 2H), 1.16 (t, 3H, J = 7.2 Hz).

#	Structure	LCMS (M+1)	NMR
337		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 7.0 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.6 Hz), 5.09 (d, 1H, J = 5.6 Hz), 4.14-4.09 (m, 1H), 3.69-3.66 (m, 2H), 3.55-3.52 (m, 2H), 3.46-3.41 (m, 4H), 3.26-3.18 (m, 1H), 2.99-2.96 (m, 2H), 2.43-2.37 (m, 3H), 2.34 (q, 2H, J = 7.2 Hz), 2.10-2.02 (m, 2H), 1.97-1.92 (m, 2H), 1.77-1.70 (m, 2H), 1.68-1.61 (m, 2H), 1.02 (t, 3H, J = 7.2 Hz).
338		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (br s, 1H), 8.03 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.61 (d, 2H, J = 8.4 Hz), 6.88 (d, 1H, J = 1.6 Hz), 6.44 (d, 2H, J = 8.4 Hz), 5.96 (d, 2H, J = 5.6 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.87 (s, 4H), 3.63-3.57 (m, 4H), 3.47-3.41 (m, 4H), 3.28 (s, 4H), 2.30-2.22 (m, 1H), 1.22 (t, 3H, J = 7.2 Hz), 0.86 (d, 6H, J = 6.4 Hz).
339		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (s, 1H), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 6.91 (s, 1H), 6.52 (t, 1H, J = 5.2 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.32-4.28 (m, 1H), 3.88-3.80 (m, 3H), 3.32-3.31 (m, 1H), 3.29-3.24 (m, 1H), 3.18-3.14 (m, 1H), 3.12-3.06 (m, 2H), 3.02-2.96 (m, 1H), 2.92-2.86 (m, 2H), 2.74-2.68 (m, 1H), 2.46-2.42 (m, 1H), 2.25-2.20 (m, 2H), 1.78-1.72 (m, 2H), 1.66-1.56 (m, 2H), 1.24 (d, 3H, J = 6.0 Hz), 1.04 (t, 3H, J = 7.2 Hz), 1.00 (d, 6H, J = 6.4 Hz).
340		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (s, 1H), 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 6.32 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.82-3.75 (m, 1H), 3.54-3.50 (m, 4H), 3.46-3.41 (m, 4H), 3.03-2.98 (m, 2H), 2.92-2.86 (m, 1H), 2.52-2.50 (m, 1H), 2.44-2.36 (m, 2H), 1.85-1.75 (m, 2H), 1.72-1.68 (m, 2H), 1.10-1.05 (m, 12H).

#	Structure	LCMS (M+1)	NMR
341		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (s, 1H), 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 6.62 (t, 1H, J = 5.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.54-3.50 (m, 4H), 3.46-3.41 (m, 4H), 3.03-2.94 (m, 4H), 2.88-2.83 (m, 1H), 2.52-2.50 (m, 1H), 2.40-2.32 (m, 2H), 1.84-1.77 (m, 2H), 1.75-1.66 (m, 2H), 1.43 (q, 2H, J = 7.2 Hz), 1.05 (d, 6H, J = 6.8 Hz), 0.85 (t, 3H, J = 7.2 Hz).
342		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (s, 1H), 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.50-3.45 (m, 4H), 3.36-3.30 (m, 4H), 3.17 (q, 2H, J = 7.2 Hz), 3.05-3.02 (m, 2H), 2.95-2.92 (m, 1H), 2.79 (s, 3H), 2.59-2.56 (m, 1H), 2.50-2.44 (m, 2H), 1.86-1.80 (m, 2H), 1.78-1.70 (m, 2H), 1.12-1.05 (m, 9H).
343		490	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.65-4.50 (m, 1H), 4.40-4.30 (m, 1H), 3.95-3.85 (m, 1H), 3.79-3.75 (m, 2H), 3.70-3.65 (m, 2H), 3.60-3.55 (m, 2H), 3.52-3.45 (m, 2H), 3.42-3.38 (m, 2H), 3.00-2.85 (m, 1H), 2.80-2.65 (m, 2H), 2.60-2.52 (m, 2H), 1.90-1.80 (m, 2H), 1.70-1.60 (m, 2H), 1.50-1.30 (m, 2H), 1.25-1.10 (m, 2H).
344		490	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.05 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.91 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.2 Hz), 5.61 (d, 1H, J = 6.0 Hz), 4.43-4.69 (m, 1H), 4.10 (t, 2H, J = 8.8 Hz), 3.70 (dd, 2H, J = 8.8, 4.8 Hz), 2.45-2.41 (m, 8H), 3.35-3.30 (m, 4H), 3.17-3.15 (m, 4H), 2.37 (q, 2H, J = 7.2 Hz), 1.04 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
345		490	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.65 (s, 1H), 7.62-7.59 (m, 1H), 7.34-7.30 (m, 1H), 7.09 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.71-3.69 (m, 2H), 3.54-3.50 (m, 2H), 3.50-3.46 (m, 2H), 2.90-2.87 (m, 2H), 2.73-2.70 (m, 2H), 2.25-2.21 (m, 2H), 2.20-2.19 (m, 1H), 2.00-1.70 (m, 4H), 0.99 (d, 6H, J = 6.4 Hz), 0.77-0.73 (m, 4H).
346		490	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 7.8 Hz, 2H), 7.24 (d, J = 8.2 Hz, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.81 (p, J = 6.2 Hz, 1H), 3.58 (q, J = 5.6, 5.1 Hz, 4H), 3.44 (t, J = 5.2 Hz, 4H), 2.91 (s, 2H), 2.23 (s, 1H), 1.73 (d, J = 52.3 Hz, 5H), 1.21 (d, J = 6.2 Hz, 6H), 1.03 (bs, 8H).
347		491	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.7 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.1 Hz, 2H), 7.24 (d, J = 8.2 Hz, 2H), 6.97 (d, J = 1.9 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 5.90 (s, 1H), 3.45 (ddd, J = 28.3, 7.6, 4.0 Hz, 9H), 2.88 (d, J = 10.8 Hz, 2H), 2.80 – 2.57 (m, 1H), 2.21 (t, J = 11.1 Hz, 2H), 1.76 (d, J = 12.4 Hz, 2H), 1.70 – 1.49 (m, 2H), 1.27 (s, 9H), 0.99 (d, J = 6.5 Hz, 6H).
348		490	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.4 Hz), 6.93 (d, 2H, J = 8.4 Hz), 6.90 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.80-3.70 (m, 1H), 3.54-3.48 (m, 4H), 3.44-3.36 (m, 4H), 3.20-3.10 (m, 4H), 2.72-2.62 (m, 1H), 2.60-2.54 (m, 4H), 1.07 (d, 6H, J = 6.4 Hz), 1.00 (d, 6H, J = 6.4 Hz).
349		491	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.89 (d, 1H, J = 1.6 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.57 (d, 2H, J = 8.4 Hz), 6.98 (d, 2H, J = 8.4 Hz), 6.66 (d, 1H, J = 1.6 Hz), 5.85 (d, 1H, J = 8.4 Hz), 4.95-4.75 (m, 1H), 3.96-3.92 (m, 2H), 3.90-3.84 (m, 2H), 3.90-3.84 (m, 2H), 3.58-3.54 (m, 2H), 2.49-3.43 (m, 2H), 3.41-3.35 (m, 4H), 2.98-2.90 (m, 1H), 2.90-2.83 (m, 4H), 2.33-2.20 (m, 1H), 1.47-1.40 (m, 2H), 1.22 (d, 6H, J = 5.6 Hz).






#	Structure	LCMS (M+1)	NMR
350		491	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.95 (d, 1H, J = 1.6 Hz), 6.94 (d, 2H, J = 8.8 Hz), 5.97 (d, 1H, J = 5.2 Hz), 5.06-4.85 (m, 1H), 3.97-3.85 (m, 2H), 3.74-7.71 (m, 2H), 3.60-3.43 (m, 4H), 3.15-3.12 (m, 4H), 2.66 (heptet, 1H, J = 6.4 Hz), 2.51-2.49 (m, 4H), 2.24-2.19 (m, 1H), 1.60-1.50 (m, 1H), 1.08-1.03 (m, 1H), 1.01 (d, 6H, J = 6.4 Hz).
351		491	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.90 (d, 1H, J = 6.0 Hz), 7.77 (d, 2H, J = 8.0 Hz), 7.56 (d, 2H, J = 8.0 Hz), 7.02 (s, 1H), 6.61 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.55-3.50 (m, 4H), 3.48-3.43 (m, 4H), 3.43-3.33 (m, 2H), 3.33-3.25 (m, 2H), 3.20-3.05 (m, 2H), 1.91-1.74(m, 2H), 1.70-1.45 (m, 2H), 1.14-0.95 (m, 9H).
352		491	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.88 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.48 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.74 (s, 1H), 3.52-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.13-3.05 (m, 2H), 2.74-2.70 (m, 1H), 2.60-2.57 (m, 4H), 1.95-1.85 (m, 2H), 1.63-1.60 (m, 2H), 1.04 (t, 3H, J = 7.2 Hz), 1.00 (d, 6H, J = 6.4 Hz).
353		491	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.63 (t, 1H, J = 5.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.65 (t, 1H, J = 5.6 Hz), 3.55-3.52 (m, 4H), 3.48-3.40 (m, 6H), 3.12 (q, 2H, J = 5.6 Hz), 2.90-2.86 (m, 2H), 2.74-2.68 (m, 1H), 2.50-2.45 (m, 1H), 2.25-2.18 (m, 2H), 1.80-1.74 (m, 2H), 1.65-1.60 (m, 2H), 1.00 (d, 6H, J = 6.4 Hz).
354		493	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.65 (s, 1H), 7.65-7.60 (m, 1H), 7.33-7.30 (m, 1H), 7.06 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.2 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.51-3.50 (m, 4H), 3.43-3.40 (m, 4H), 3.07-3.04 (m, 2H), 2.91-2.89 (m, 2H), 2.73-2.71 (m, 2H), 2.30-2.20 (m, 2H), 1.73-1.65 (m, 4H), 1.21-1.20 (m, 9H).

#	Structure	LCMS (M+1)	NMR
355		494	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 6.0 Hz), 7.76 (d, 2H, J = 8.8 Hz), 7.30 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.2 Hz), 5.97 (d, 1H, J = 4.2 Hz), 3.92 (br. s., 2H), 3.70 (br. s., 2H), 3.54 (br. s., 2H), 3.47 (br. s., 2H), 3.16-2.98 (m, 3H), 2.47-2.43 (m, 2H), 2.36-2.26 (m, 1H), 2.16-2.07 (m, 2H), 2.05-1.98 (m, 1H), 1.82-1.78 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz), 0.77-0.74 (m, 4H).
356		494	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (s, 1H), 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.31 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 6.12-5.84 (m, 1H), 5.99 (d, 1H, J = 5.2 Hz), 3.79-3.78 (m, 2H), 3.55-3.45 (m, 4H), 3.37-3.31 (m, 3H), 3.03-3.02 (m, 1H), 2.77-2.75 (m, 2H), 2.57-2.53 (m, 4H), 2.25-2.17 (m, 1H), 1.84-1.74 (m, 1H), 1.10-1.07 (m, 6H).
357		494	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.65-7.59 (m, 2H), 7.33-7.30 (m, 1H), 7.06 (d, 1H, J = 1.2 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.61-3.59 (m, 4H), 3.49-3.46 (m, 4H), 2.89-2.88 (m, 2H), 2.72-2.67 (m, 2H), 2.24-2.20 (m, 2H), 1.73-1.65 (m, 4H), 1.21 (t, 3H, J = 7.2 Hz), 0.98 (d, 6H, J = 6.4 Hz).
358		495	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 8.13 (s, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 7.8 Hz, 2H), 7.31 – 7.16 (m, 2H), 6.99 (s, 1H), 5.98 (d, J = 5.5 Hz, 1H), 4.77 – 4.64 (m, 1H), 4.63 – 4.52 (m, 1H), 4.37 – 4.27 (m, 1H), 4.25 (t, J = 4.0 Hz, 1H), 3.62 (s, 4H), 3.47 (t, J = 5.1 Hz, 4H), 2.90 (s, 2H), 2.70 (d, J = 29.0 Hz, 1H), 2.24 (s, 2H), 1.88 – 1.54 (m, 5H), 1.00 (d, J = 6.6 Hz, 6H).
359		495	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.07 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.66 (d, 2H, J = 8.4 Hz), 6.96-6.94 (m, 3H), 6.20-6.15 (m, 1H), 5.95 (d, 1H, J = 5.6 Hz), 3.92-3.90 (m, 2H), 3.75-3.65 (m, 2H), 3.51-3.44 (m, 4H), 3.15 (t, 4H, J = 4.8 Hz), 2.83-2.74 (m, 2H), 2.68-2.66 (m, 4H), 2.02-1.98 (m, 1H), 0.78-0.72 (m, 4H).





#	Structure	LCMS (M+1)	NMR
360		495	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.27 (s, 1H), 7.90 (d, 1H, J = 4.8 Hz), 7.57 (d, 2H, J = 10.8 Hz), 7.14 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 3.92 (br. s., 2H), 3.70 (br. s., 2H), 3.53 (br. s., 2H), 3.47 (br. s., 2H), 3.13 (s, 4H), 2.50 (br. s., 4H), 2.38 (q, 2H, J = 7.2 Hz), 2.02-2.00 (m, 1H), 1.02 (t, 3H, J = 7.2 Hz), 0.78-0.75 (m, 4H).
361		495	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 2.0 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.94-6.93 (m, 1H), 5.97 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 1H), 3.80-3.60 (m, 3H), 3.58-3.40 (m, 4H), 3.30-3.24 (m, 4H), 3.23-3.20 (m, 1H), 3.18-3.10 (m, 4H), 2.37 (q, 2H, J = 7.2 Hz), 2.00-1.90 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
362		495	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.94 (d, 1H, J = 5.6 Hz), 7.90 (d, 1H, J = 1.2 Hz), 6.83 (s, 1H), 7.76 (d, 2H, J = 12.8 Hz), 6.01 (d, 1H, J = 5.6 Hz), 3.93-3.91 (m, 2H), 3.69-3.68 (m, 2H), 3.55-3.54 (m, 2H), 3.51-3.50 (m, 2H), 3.25-3.22 (m, 4H), 2.48-2.47 (m, 4H), 2.37 (q, 2H, J = 7.2 Hz), 2.04-2.00 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.78-0.72 (m, 4H).
363		497	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.00 (s, 1H), 7.88 (d, 1H, J = 5.6 Hz), 7.75 (d, 1H, J = 8.8 Hz), 7.28 (t, 1H, J = 7.4 Hz), 6.96 (s, 1H), 6.89 (d, 1H, J = 8.4 Hz), 6.76 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 3.91-3.90 (m, 2H), 3.69-3.67 (m, 2H), 3.52-3.46 (m, 4H), 3.24-3.20 (m, 4H), 2.79-2.74 (m, 4H), 2.01-1.98 (m, 1H), 0.84-0.73 (m, 4H).
364		497	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.77-3.68 (m, 2H), 3.68-3.62 (m, 1H), 3.57-3.50 (m, 2H), 3.48-3.40 (m, 4H), 3.30-3.20 (m, 2H), 3.05-2.95 (m, 1H), 2.80-2.68 (m, 2H), 2.65-2.55 (m, 2H), 2.50-2.42 (m, 3H), 2.40-2.35 (m, 1H), 2.25-2.15 (m, 1H), 1.80-1.70 (m, 1H), 1.10-1.00 (m, 6H).





#	Structure	LCMS (M+1)	NMR
365		497	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.97-3.93 (m, 2H), 3.72-3.70 (m, 2H), 3.60-3.50 (m, 2H), 3.49-3.44 (m, 2H), 2.96-2.92 (m, 3H), 2.81-2.77 (m, 1H), 2.52-2.50 (m, 1H), 2.34-2.26 (m, 2H), 2.15-2.05 (m, 1H), 1.82-1.75 (m, 2H), 1.71-1.61 (m, 2H), 1.50-1.44 (m, 1H), 1.38-1.34 (m, 1H), 1.03 (d, 6H, J = 6.8 Hz).
366		497	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.4 Hz), 7.30 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.73-3.67 (m, 2H), 3.67-3.64 (m, 1H), 3.54-3.50 (m, 2H), 3.45-3.39 (m, 4H), 3.27-3.25 (m, 2H), 3.01-2.96 (m, 1H), 2.75-2.71 (m, 2H), 2.66-2.63 (m, 2H), 2.61-2.53 (m, 2H), 2.50-2.46 (m, 1H), 2.45-2.41 (m, 1H), 2.22-2.19 (m, 1H), 1.79-1.73 (m, 1H), 1.07-1.04 (m, 6H).
367		498	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.87 (d, 1H, J = 2.0 Hz), 7.82 (d, 1H, J = 5.2 Hz), 7.55 (d, 2H, J = 8.8 Hz), 6.97 (d, 2H, J = 8.8 Hz), 6.62 (d, 1H, J = 2.0 Hz), 5.82 (d, 1H, J = 5.2 Hz), 3.87-3.83 (m, 2H), 3.62-3.58 (m, 1H), 3.57-3.54 (m, 2H), 3.43-3.41 (m, 4H), 3.29-3.24 (m, 4H), 3.22-3.14 (m, 1H), 2.82-2.74 (m, 2H), 2.66-2.62 (m, 4H), 2.61-2.55 (m, 2H), 2.50 (q, 2H, J = 7.2 Hz), 1.15 (t, 3H, J = 7.2 Hz).
368		498	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 2.0 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.0 Hz), 6.98 (d, 1H, J = 1.2 Hz), 5.95 (d, 1H, J = 5.2 Hz), 4.21 (t, 2H, J = 8.8 Hz), 4.08 (dd, 2H, J = 8.0, 6.4 Hz), 3.80-3.70 (m, 1H), 3.50-3.40 (m, 8H), 3.30-3.20 (m, 1H), 3.10-2.90 (m, 1H), 2.80-2.60 (m, 2H), 2.45-2.35 (m, 2H), 2.30-2.10 (m, 1H), 1.80-1.70 (m, 1H), 1.05 (t, 6H, J = 6.4 Hz).





#	Structure	LCMS (M+1)	NMR
369		498	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.92 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.59 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.0 Hz), 6.67 (d, 1H, J = 1.6 Hz), 5.84 (d, 1H, J = 5.2 Hz), 4.33-4.22 (m, 4H), 3.62-3.56 (m, 4H), 3.51-3.47 (m, 4H), 3.50-3.40 (m, 2H), 3.33-3.26 (m, 1H), 3.10-3.00 (m, 1H), 2.79-2.69 (m, 1H), 2.60-2.49 (m, 2H), 2.41-2.30 (m, 1H), 2.01-1.90 (m, 1H), 1.18 (d, 6H, J = 4.4 Hz).
370		498	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 2.0 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 6.98 (d, 1H, J = 1.2 Hz), 5.94 (d, 1H, J = 5.6 Hz), 4.21 (t, 2H, J = 8.4 Hz), 4.08 (dd, 2H, J = 8.4, 6.0 Hz), 3.77-3.75 (m, 1H), 3.46-3.45 (m, 8H), 2.99-2.95 (m, 2H), 2.49-2.44 (m, 1H), 2.34 (q, 2H, J = 7.2 Hz), 1.97-1.92 (m, 2H), 1.76-1.70 (m, 2H), 1.68-1.63 (m, 2H), 1.01 (t, 3H, J = 7.2 Hz).
371		498	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.30 (s, 1H), 8.08 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.67 (d, 2H, J = 8.4 Hz), 6.96 (d, 2H, J = 8.4 Hz), 6.94 (d, 1H, J = 1.6 Hz), 5.95 (d, 1H, J = 5.6 Hz), 3.75-3.67 (m, 2H), 3.67-3.60 (m, 2H), 3.50-3.42 (m, 2H), 3.42-3.35 (m, 2H), 3.22-3.14 (m, 4H), 3.06-2.94 (m, 4H), 2.76-2.64 (m, 2H), 2.10-2.00 (m, 2H), 1.78-1.68 (m, 2H), 1.66-1.50 (m, 2H), 1.48-1.34 (m, 2H).
372		498	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.30 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.92-3.86 (m, 2H), 3.72-3.68 (m, 2H), 3.54-3.49 (m, 2H), 3.48-3.43 (m, 2H), 3.31-3.27 (m, 1H), 2.94-2.91 (m, 1H), 2.68-2.62 (m, 2H), 2.46-2.42 (m, 2H), 2.26-2.24 (m, 1H), 2.16-2.10 (m, 1H), 1.98-1.92 (m, 2H), 1.80-1.66 (m, 6H), 1.28-1.22 (m, 2H), 1.06 (t, 3H, J = 7.2 Hz), 0.99-0.95 (m, 1H), 0.65-0.60 (m, 1H).






#	Structure	LCMS (M+1)	NMR
373		499	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.09 (s, 1H), 7.87 (d, 1H, J = 5.6 Hz), 7.67 (d, 2H, J = 8.4 Hz), 6.99 (d, 2H, J = 8.4 Hz), 6.95 (s, 1H), 5.96 (d, 1H, J = 5.6 Hz), 3.92 (br. s., 2H), 3.69-3.63 (m, 3H), 3.51-3.45 (m, 6H), 3.04-3.01 (m, 1H), 2.94-2.89 (m, 1H), 2.86-2.79 (m, 1H), 2.76-2.66 (m, 2H), 2.06-2.00 (m, 1H), 0.78-0.71 (m, 4H).
374		499	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.34 (s, 1H), 8.07 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.8 Hz), 7.01 (d, 2H, J = 8.8 Hz), 6.95 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.2 Hz), 4.71-4.68 (m, 1H), 3.92 (br. s., 2H), 3.71 (br. s., 2H), 3.53-3.35 (m, 4H), 3.28-3.09 (m, 3H), 3.03-2.96 (m, 2H), 2.74-2.67 (m, 1H).
375		499	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.27 (s, 1H), 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.50-4.30 (m, 1H), 3.95-3.85 (m, 1H), 3.79-3.75 (m, 2H), 3.70-3.65 (m, 2H), 3.60-3.55 (m, 2H), 3.52-3.45 (m, 2H), 3.42-3.38 (m, 2H), 3.00-2.90 (m, 1H), 2.80-2.71 (m, 2H), 2.70-2.65 (m, 2H), 2.10-2.00 (m, 2H), 1.80-1.68 (m, 2H), 1.65-1.50 (m, 2H), 1.48-1.30 (m, 2H).
376		499	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 8.06 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.96 (d, 2H, J = 8.8 Hz), 6.91 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.6 Hz), 4.22 (t, 2H, J = 8.8 Hz), 4.09 (dd, 2H, J = 8.0, 6.0 Hz), 3.78-3.73 (m, 1H), 2.47-2.46 (m, 4H), 2.45-2.44 (m, 4H), 3.18-3.16 (m, 4H), 2.55-2.53 (m, 4H), 2.40 (q, 2H, J = 7.2 Hz), 1.05 (t, 3H, J = 7.2 Hz).
377		500	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.30 (s, 1H), 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.07 (t, 1H, J = 5.6 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.58-3.52 (m, 4H), 3.48-3.42 (m, 4H), 3.33-3.24 (m, 2H), 3.03-2.98 (m, 2H), 2.90-2.85 (m, 1H), 2.69-2.64 (m, 2H), 2.55-2.50 (m, 1H), 2.43-2.39 (m, 2H), 1.84-1.75 (m, 2H), 1.74-1.68 (m, 2H), 1.06 (d, 6H, J = 6.4 Hz).





#	Structure	LCMS (M+1)	NMR
378		501	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.2 Hz), 5.97 (d, 1H, J = 5.2 Hz), 5.30-5.00 (m, 1H), 3.80-3.70 (m, 2H), 3.70-3.60 (m, 2H), 3.54-3.45 (m, 2H), 3.42-3.33 (m, 4H), 3.18-3.10 (m, 1H), 3.08-3.02 (m, 1H), 2.80-2.60 (m, 3H), 2.10-1.98 (m, 2H), 1.80-1.66 (m, 2H), 1.62-1.50 (m, 2H), 1.46-1.30 (m, 2H).
379		501	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 6.70 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.45-3.44 (m, 4H), 3.35-3.33 (m, 4H), 3.16-3.13 (m, 4H), 2.99-2.96 (m, 2H), 2.48-2.42 (m, 1H), 2.34 (q, 2H, J = 7.2 Hz), 1.98-1.92 (m, 2H), 1.76-1.71 (m, 2H), 1.70-1.60 (m, 2H), 1.60-1.50 (m, 2H), 1.50-1.40 (m, 4H), 1.02 (t, 3H, J = 7.2 Hz).
380		502	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.1 Hz, 2H), 7.24 (d, J = 8.0 Hz, 2H), 6.97 (d, J = 1.9 Hz, 1H), 6.74 (d, J = 7.6 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.12 (h, J = 8.3 Hz, 1H), 3.52 (dd, J = 6.9, 3.3 Hz, 4H), 3.42 (dd, J = 6.7, 3.5 Hz, 4H), 2.88 (d, J = 10.6 Hz, 2H), 2.71 (p, J = 6.6 Hz, 1H), 2.27 – 2.17 (m, 2H), 2.12 (qt, J = 7.7, 2.6 Hz, 2H), 1.93 (pd, J = 9.1, 2.7 Hz, 2H), 1.76 (d, J = 12.3 Hz, 2H), 1.69 – 1.47 (m, 4H), 0.99 (d, J = 6.5 Hz, 6H).
381		502	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.00 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 5.11 (d, 1H, J = 6.0 Hz), 4.15-4.08 (m, 1H), 3.72-3.65 (m, 2H), 3.60-3.50 (m, 2H), 3.48-3.40 (m, 4H), 3.10-2.99 (m, 1H), 3.85-3.78 (m, 2H), 3.78-3.69 (m, 2H), 2.44-2.39 (m, 2H), 2.20-2.10 (m, 2H), 2.09-2.03 (m, 2H), 1.85-1.79 (m, 1H), 1.77-1.70 (m, 1H), 1.60-1.42 (m, 2H), 0.97 (dd, 6H, J = 6.4, 3.2 Hz).





#	Structure	LCMS (M+1)	NMR
382		502	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (s, 1H), 7.88 (d, 1H, J = 4.4 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 6.93 (s, 1H), 5.97-5.93 (m, 1H), 5.10 (d, 1H, J = 2.0 Hz), 4.70-4.30 (m, 1H), 4.12-4.08 (m, 1H), 3.89-3.86 (m, 2H), 3.60-3.50 (m, 1H), 3.21-3.05 (m, 2H), 3.05-3.00 (m, 1H), 3.00-2.96 (m, 2H), 2.49-2.48 (m, 1H), 2.37-2.35 (m, 4H), 2.10-2.00 (m, 2H), 2.00-1.94 (m, 2H), 1.78-1.75 (m, 2H), 1.70-1.67 (m, 2H), 1.37-1.32 (m, 1H), 1.27-1.21 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
383		503	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 7.8 Hz, 2H), 7.25 (d, J = 7.9 Hz, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.95 (d, J = 5.5 Hz, 1H), 4.97 (s, 1H), 3.63 (dt, J = 27.9, 4.8 Hz, 4H), 3.43 (q, J = 5.9, 4.1 Hz, 4H), 2.98 (d, J = 11.0 Hz, 2H), 2.89 (q, J = 8.8 Hz, 1H), 2.34 (q, J = 7.2 Hz, 2H), 2.19 (t, J = 10.1 Hz, 2H), 2.08 (td, J = 8.5, 2.6 Hz, 2H), 2.03 – 1.88 (m, 2H), 1.75 (d, J = 11.3 Hz, 2H), 1.65 (qd, J = 12.3, 3.5 Hz, 2H), 1.28 (s, 3H), 1.02 (t, J = 7.2 Hz, 3H).
384		503	
385		502	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.00 (d, 1H, J = 1.2 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.10 (d, 1H, J = 6.0 Hz), 4.13-4.11 (m, 1H), 3.70-3.69 (m, 2H), 3.55-3.54 (m, 2H), 3.45-3.43 (m, 4H), 3.24-3.22 (m, 1H), 2.82-2.77 (m, 2H), 2.76-2.67 (m, 2H), 2.44-2.38 (m, 2H), 2.20-2.14 (m, 2H), 2.10-2.03 (m, 2H), 1.81-1.79 (m, 1H), 1.72-1.70 (m, 1H), 1.59-1.50 (m, 1H), 1.48-1.40 (m, 1H), 0.99-0.96 (m, 6H).

#	Structure	LCMS (M+1)	NMR
386		502	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (s, 1H), 8.15 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.89 (d, 2H, J = 6.8 Hz), 3.65-3.59 (m, 4H), 3.50-3.40 (m, 4H), 2.97-2.2.93 (m, 2H), 2.82-2.78 (m, 1H), 2.52-2.50 (m, 1H), 2.35-2.28 (m, 2H), 1.82-1.77 (m, 2H), 1.70-1.64 (m, 2H), 1.14-1.11 (m, 1H), 1.03 (d, 6H, J = 6.8 Hz), 0.54-0.51 (m, 2H), 0.30-0.27 (m, 2H).
387		502	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.11 (d, 1H, J = 5.6 Hz), 4.14-4.11 (m, 1H), 3.69-3.68 (m, 2H), 3.55-3.54 (m, 2H), 3.46-3.44 (m, 4H), 3.22-3.21 (m, 1H), 2.81-2.75 (m, 2H), 2.74-2.68 (m, 2H), 2.43-2.38 (m, 2H), 2.19-2.10 (m, 2H), 2.08-2.01 (m, 2H), 1.72-1.71 (m, 1H), 1.70-1.69 (m, 1H), 1.60-1.40 (m, 2H), 0.98-0.96 (m, 6H).
388		502	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (s, 1H), 8.19 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.80 (d, 2H, J = 8.4 Hz), 7.48 (d, 2H, J = 8.4 Hz), 7.06 (s, 1H), 5.98 (d, 1H, J = 5.2 Hz), 3.94-3.91 (m, 2H), 3.74-3.70 (m, 2H), 3.60-3.52 (m, 2H), 3.50-3.45 (m, 2H), 2.92 (s, 3H), 2.92-2.81 (m, 3H), 2.60-2.50 (m, 1H), 2.03-2.01 (m, 1H), 1.85-1.80 (m, 2H), 1.80-1.75 (m, 1H), 1.45-1.40 (m, 1H), 1.04 (d, 6H, J = 6.4 Hz), 0.78-0.74 (m, 4H).
389		502	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.19 (s, 1H), 7.91 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.06 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.01-3.91 (m, 2H), 3.76-3.69 (m, 2H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 2.89 (s, 3H), 2.71-2.69 (m, 1H), 2.61-2.59 (m, 2H), 2.47-2.45 (m, 2H), 2.03-1.96 (m, 2H), 1.88-1.81 (m, 2H), 1.00 (d, 6H, J = 6.0 Hz), 0.78-0.75 (m, 4H).





#	Structure	LCMS (M+1)	NMR
394		503	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.12 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.24 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.25 (d, 1H, J = 8.0 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.68-3.55 (m, 1H), 3.54-3.48 (m, 4H), 3.46-3.38 (m, 4H), 2.92-2.82 (m, 2H), 2.74-2.64 (m, 1H), 2.46-2.40 (m, 1H), 2.26-2.14 (m, 2H), 1.80-1.70 (m, 2H), 1.68-1.54 (m, 2H), 1.50-1.30 (m, 2H), 1.04 (d, 3H, J = 6.4 Hz), 0.98 (d, 6H, J = 6.4 Hz), 0.83 (t, 3H, J = 7.2 Hz).
395		504	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.7 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.1 Hz, 2H), 7.24 (d, J = 8.2 Hz, 2H), 6.97 (d, J = 1.9 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 5.90 (s, 1H), 3.45 (ddd, J = 28.3, 7.6, 4.0 Hz, 9H), 2.88 (d, J = 10.8 Hz, 2H), 2.80 – 2.57 (m, 1H), 2.21 (t, J = 11.1 Hz, 2H), 1.76 (d, J = 12.4 Hz, 2H), 1.70 – 1.49 (m, 2H), 1.27 (s, 9H), 0.99 (d, J = 6.5 Hz, 6H).
396		504	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21 (s, 1H), 8.12 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.76 - 7.67 (m, 2H), 7.28 - 7.19 (m, 2H), 6.98 (d, J = 1.9 Hz, 1H), 6.56 (t, J = 5.5 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.56 - 3.47 (m, 4H), 3.42 (dd, J = 6.8, 3.4 Hz, 4H), 3.09 - 3.00 (m, 2H), 2.95 (d, J = 11.0 Hz, 2H), 2.81 (p, J = 6.5 Hz, 1H), 2.32 (dd, J = 12.6, 10.1 Hz, 2H), 1.79 (d, J = 12.4 Hz, 2H), 1.74 - 1.59 (m, 2H), 1.46 - 1.34 (m, 2H), 1.27 (h, J = 7.2 Hz, 2H), 1.02 (d, J = 6.6 Hz, 5H), 0.87 (t, J = 7.3 Hz, 3H).
397		504	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (br. s., 1H), 8.06 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.65 (d, 2H, J = 8.4 Hz), 6.95 (d, 2H, J = 8.4 Hz), 6.91 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.2 Hz), 4.15-4.12 (m, 3H), 3.78-3.75 (m, 2H), 3.46-3.44 (m, 8H), 3.20 (s, 3H), 3.16-3.15 (m, 4H), 2.55-2.54 (m, 4H), 2.41 (q, 2H, J = 7.2 Hz), 1.05 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
398		505	¹ H NMR (400 MHz, DMSO-d ₆) δ 9.63 (s, 1H), 8.16 (d, J = 1.7 Hz, 1H), 7.90 (d, J = 5.4 Hz, 1H), 7.78 (d, J = 7.9 Hz, 2H), 7.25 (d, J = 8.1 Hz, 2H), 7.01 (d, J = 1.8 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 3.83 (d, J = 6.5 Hz, 2H), 3.61 (br. s, 4H), 3.52 - 3.41 (m, 7H), 3.15 - 3.01 (m, 2H), 2.87 (p, J = 9.4, 8.5 Hz, 1H), 2.03 (t, J = 9.3 Hz, 4H), 1.89 (hept, J = 6.8 Hz, 1H), 1.29 (d, J = 6.6 Hz, 6H), 0.91 (d, J = 6.7 Hz, 6H)
399		505	¹ H-NMR (400 MHz, MeOD) δ ppm 7.98 (d, 1H, J = 1.2 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.74 (d, 1H, J = 8.4 Hz), 7.45 (d, 1H, J = 8.4 Hz), 6.92 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 5.2 Hz), 3.67-3.64 (m, 4H), 3.56-3.54 (m, 4H), 3.24 (q, 2H, J = 7.2 Hz), 3.00 (s, 3H), 2.87-2.82 (m, 3H), 2.77-2.71 (m, 2H), 2.20-2.16 (m, 2H), 2.08-2.02 (m, 2H), 1.17 (d, 6H, J = 5.4 Hz), 1.15 (t, 3H, J = 7.2 Hz).
400		505	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.48 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.2 Hz), 6.62 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.53-3.51 (m, 4H), 3.45-3.43 (m, 4H), 3.33-3.30 (m, 1H), 3.20-3.10 (m, 2H), 2.92 (s, 3H), 2.80-2.74 (m, 2H), 2.72-2.60 (m, 1H), 2.50-2.40 (m, 2H), 1.83-1.71 (m, 3H), 1.06-1.01 (m, 9H).
401		507	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.1 Hz, 2H), 7.24 (d, J = 8.2 Hz, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.21 - 4.10 (m, 2H), 3.68 - 3.57 (m, 4H), 3.57 - 3.50 (m, 2H), 3.50 - 3.41 (m, 4H), 3.28 (s, 3H), 2.87 (s, 2H), 2.68 (d, J = 17.6 Hz, 1H), 2.21 (s, 2H), 1.75 (s, 2H), 1.62 (d, J = 13.0 Hz, 2H), 0.99 (d, J = 6.4 Hz, 6H).
402		509	¹ H-NMR (500 MHz, DMSO-d ₆) δ ppm 8.07 (s, 1H), 7.88 (d, 1H, J = 5.5 Hz), 7.65 (d, 2H, J = 8.0 Hz), 6.97-6.93 (m, 3H), 5.96 (d, 1H, J = 5.5 Hz), 3.72-3.71 (m, 2H), 3.63-3.62 (m, 2H), 3.47-3.46 (m, 4H), 3.40-3.39 (m, 4H), 3.17-3.16 (m, 4H), 2.86-2.80 (m, 4H), 2.39-2.37 (m, 2H), 1.24 (s, 1H), 1.04 (t, 3H, J = 7.5 Hz).






#	Structure	LCMS (M+1)	NMR
407		511	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.33 (br.s., 1H), 8.15 (s, 1H), 7.90 (d, 1H, J = 4.8 Hz), 7.73 (d, 2H, J = 7.6 Hz), 7.27 (d, 2H, J = 7.6 Hz), 6.96 (s, 1H), 5.97 (s, 1H), 4.66-4.62 (m, 1H), 4.22-4.16 (m, 1H), 3.98-3.88 (m, 2H), 3.42-3.16 (m, 2H), 3.08-2.98 (m, 2H), 2.96-2.84 (m, 2H), 2.62-2.52 (m, 1H), 2.46-2.40 (m, 2H), 2.12-2.06 (m, 1H), 1.88-1.72 (m, 4H), 1.50-1.40 (m, 3H), 1.32-1.24 (m, 3H), 1.08 (d, 6H, J = 5.2 Hz).
408		511	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 6.99 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.69-3.67 (m, 2H), 3.64-3.60 (m, 1H), 3.54-3.53 (m, 2H), 3.45-3.43 (m, 4H), 3.32-3.30 (m, 1H), 2.80-2.78 (m, 2H), 2.72-2.71 (m, 2H), 2.63-2.61 (m, 2H), 2.52-2.51 (m, 1H), 2.46-2.45 (m, 1H), 2.19-2.11 (m, 2H), 1.83-1.80 (m, 1H), 1.74-1.71 (m, 1H), 1.58-1.50 (m, 1H), 1.49-1.42 (m, 1H), 0.98-0.96 (m, 6H).
409		511	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.00 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.75-3.69 (m, 2H), 3.70-3.67 (m, 1H), 3.54-3.53 (m, 2H), 3.45-3.33 (m, 4H), 3.31-3.28 (m, 1H), 2.81-2.79 (m, 2H), 2.74-2.70 (m, 2H), 2.66-2.63 (m, 2H), 2.50-2.46 (m, 2H), 2.19-2.09 (m, 2H), 1.82-1.81 (m, 1H), 1.72-1.71 (m, 1H), 1.56-1.50 (m, 1H), 1.50-1.42 (m, 1H), 0.98-0.96 (m, 6H).
410		512	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.92 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.60 (d, 2H, J = 8.0 Hz), 7.33 (d, 2H, J = 8.0 Hz), 6.68 (d, 1H, J = 1.6 Hz), 5.87 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 4H), 3.47-3.45 (m, 4H), 3.44-3.43 (m, 1H), 3.18-3.17 (m, 1H), 2.96-2.94 (m, 1H), 2.68-2.67 (m, 2H), 2.53-2.52 (m, 2H), 2.37-2.36 (m, 1H), 1.96-1.94 (m, 1H), 1.39-1.37 (m, 2H), 1.26-1.25 (m, 2H), 1.21-1.16 (m, 3H).





#	Structure	LCMS (M+1)	NMR
411		512	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (s, 1H), 8.15 (s, 1H), 7.89 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 6.99 (s, 1H), 5.95 (d, 1H, J = 5.2 Hz), 4.24-4.19(m, 2H), 4.11-4.01(m, 2H), 3.77-3.74 (m, 1H), 2.89-2.84(m, 4H), 2.41-2.27 (m, 2H), 1.85-1.41(m, 4H), 1.03-1.00 (m, 6H).
412		512	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (br. s., 1H), 8.15 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.70 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 6.90 (d, 1H, J = 1.6 Hz), 5.95 (d, 1H, J = 5.6 Hz), 4.23-4.20 (m, 2H), 4.19-4.16 (m, 1H), 4.10-4.06 (m, 2H), 3.86-3.83 (m, 2H), 3.75-3.74 (m, 1H), 3.64-3.63 (m, 1H), 3.35-3.34 (m, 2H), 3.16-3.03 (m, 2H), 3.01-2.99 (m, 1H), 2.39 (q, 2H, J = 7.2 Hz), 2.01-1.98 (m, 2H), 1.79-1.74 (m, 2H), 1.72-1.65 (m, 2H), 1.31 (d, 3H, J = 6.4 Hz), 1.04 (t, 3H, J = 7.2 Hz).
413		512	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (s, 1H), 7.86 (d, 1H, J = 5.6 Hz), 7.64 (d, 2H, J = 8.4 Hz), 6.93 (d, 2H, J = 8.4 Hz), 6.92-6.90 (m, 1H), 5.94 (d, 1H, J = 5.6 Hz), 3.70-3.60 (m, 3H), 3.55-3.48 (m, 2H), 3.45-3.38 (m, 4H), 3.30-3.22 (m, 1H), 3.17-3.08 (m, 4H), 2.70-2.60 (m, 1H), 2.58-2.54 (m, 6H), 2.48-2.42 (m, 2H), 1.00 (d, 6H, J = 6.4 Hz).
414		512	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.90-3.60 (m, 4H), 3.55-3.45 (m, 4H), 2.91-2.86 (m, 2H), 2.71 (heptet, 1H, J = 6.8 Hz), 2.50-2.46 (m, 1H), 2.26-2.20 (m, 2H), 1.80-1.75 (m, 2H), 1.69-1.56 (m, 2H), 1.24-1.20 (m, 1H), 1.00 (d, 6H, J = 6.8 Hz), 0.78-0.74 (m, 2H), 0.53-0.50 (m, 2H), 0.48-0.43 (m, 2H), 0.15-0.11 (m, 2H).






#	Structure	LCMS (M+1)	NMR
415		515	¹ H NMR (400 MHz, Chloroform-d) δ 7.92 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.3 Hz, 1H), 7.59 (d, J = 7.9 Hz, 2H), 7.28 (d, J = 8.0 Hz, 3H), 6.69 (d, J = 1.9 Hz, 1H), 5.85 (d, J = 5.4 Hz, 1H), 3.86 (d, J = 26.3 Hz, 5H), 3.49 (d, J = 31.7 Hz, 5H), 3.15 (s, 3H), 2.56 (td, J = 15.0, 5.8 Hz, 3H), 2.34 (ddd, J = 33.9, 16.5, 9.0 Hz, 1H), 2.17 – 2.06 (m, 7H), 1.89 (bs, 3H), 1.24 (d, J = 5.2 Hz, 3H), 1.16 (t, J = 7.1 Hz, 4H).
416		516	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (s, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.74 (d, J = 7.8 Hz, 2H), 7.28 (dd, J = 24.1, 7.9 Hz, 3H), 6.98 (s, 1H), 6.36 (d, J = 7.0 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 5.75 (s, 1H), 3.93 (p, J = 7.1 Hz, 1H), 3.60 – 3.40 (m, 10H), 2.11 – 1.72 (m, 2H), 1.63 (q, J = 6.6, 6.2 Hz, 2H), 1.44 (ddt, J = 27.1, 11.9, 5.6 Hz, 4H), 1.33 – 0.90 (m, 12H).
417		516	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.29 (br. s., 1H), 8.06 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.65 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.90 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.2 Hz), 4.66-4.42 (m, 4H), 4.11-4.03 (m, 4H), 3.49-3.45 (m, 4H), 3.44-3.40 (m, 4H), 3.17-3.11 (m, 4H), 2.53-2.50 (m, 4H), 2.38 (q, 2H, J = 7.2 Hz), 1.04 (t, 3H, J = 7.2 Hz).
418		516	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.2 Hz), 4.58 (d, 1H, J = 4.0 Hz), 3.80-3.70 (m, 2H), 3.70-3.60 (m, 2H), 3.55-3.47 (m, 2H), 3.47-3.40 (m, 2H), 3.39-3.35 (m, 1H), 3.30-3.20 (m, 1H), 3.05-2.95 (m, 1H), 2.80-2.60 (m, 2H), 2.58-2.51 (m, 1H), 2.45-2.30 (m, 2H), 2.28-2.10 (m, 1H), 1.90-1.81 (m, 2H), 1.80-1.75 (m, 1H), 1.70-1.60 (m, 2H), 1.50-1.30 (m, 2H), 1.25-1.10 (m, 2H), 1.08-1.00 (m, 6H).





#	Structure	LCMS (M+1)	NMR
419		516	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.93 (d, 1H, J = 2.0 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.61 (d, 2H, J = 8.0 Hz), 7.33 (d, 2H, J = 8.0 Hz), 6.69 (d, 1H, J = 1.6 Hz), 5.85 (d, 1H, J = 5.6 Hz), 3.92-3.82 (m, 2H), 3.80-3.70 (m, 2H), 3.69-3.65 (m, 1H), 3.56-3.45 (m, 3H), 3.44-3.41 (m, 2H), 3.40-3.38 (m, 1H), 3.24-3.12 (m, 1H), 2.99-2.87 (m, 1H), 2.80-2.66 (m, 2H), 2.53-2.36 (m, 2H), 2.14-2.00 (m, 4H), 1.90-1.78 (m, 2H), 1.74-1.71 (m, 1H), 1.39-1.34 (m, 2H), 1.27 (d, 6H, J = 6.4 Hz).
420		517	¹ H NMR (400 MHz, Chloroform-d) δ 7.91 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.3 Hz, 1H), 7.58 (d, J = 8.1 Hz, 2H), 7.28 (d, J = 8.2 Hz, 2H), 6.67 (d, J = 1.8 Hz, 1H), 5.83 (d, J = 5.3 Hz, 1H), 4.12 (q, J = 7.1 Hz, 2H), 3.87 (d, J = 5.6 Hz, 2H), 3.66 (s, 2H), 3.45 (s, 2H), 3.03 (d, J = 11.2 Hz, 2H), 2.98 – 2.86 (m, 1H), 2.78 (p, J = 6.5 Hz, 1H), 2.59 – 2.44 (m, 2H), 2.44 – 2.35 (m, 4H), 2.27 (t, J = 11.0 Hz, 2H), 1.94 – 1.77 (m, 4H), 1.25 (t, J = 7.1 Hz, 3H), 1.10 (d, J = 6.6 Hz, 6H).
421		517	¹ H NMR (400 MHz, Chloroform-d) δ 7.91 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.3 Hz, 1H), 7.61 – 7.49 (m, 2H), 7.29 (d, J = 8.2 Hz, 2H), 6.67 (d, J = 1.9 Hz, 1H), 5.83 (d, J = 5.4 Hz, 1H), 4.11 (q, J = 7.1 Hz, 1H), 3.87 (t, J = 5.1 Hz, 2H), 3.73 – 3.62 (m, 3H), 3.51 – 3.36 (m, 2H), 3.18 (s, 2H), 2.93 (p, J = 8.1 Hz, 1H), 2.58 (s, 2H), 2.52 – 2.29 (m, 5H), 1.94 (d, J = 12.9 Hz, 2H), 1.41 (s, 3H), 1.24 (dt, J = 10.8, 6.8 Hz, 10H).
422		517	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 6.0 Hz), 7.64 (d, 2H, J = 8.8 Hz), 6.95 (d, 2H, J = 8.8 Hz), 6.93 (d, 1H, J = 1.6 Hz), 5.94 (d, 1H, J = 5.6 Hz), 4.57 (d, 1H, J = 4.4 Hz), 3.73-3.70 (m, 2H), 3.70-3.64 (m, 2H), 3.48-3.45 (m, 2H), 3.44-3.38 (m, 2H), 3.17-3.13 (m, 4H), 2.53-2.58 (m, 4H), 1.86-1.83 (m, 2H), 1.70-1.66 (m, 2H), 1.56-1.53 (m, 1H), 1.43-1.16 (m, 5H), 1.06-1.04 (m, 3H), 0.93 (t, 3H, J = 7.2 Hz).

#	Structure	LCMS (M+1)	NMR
423		517	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.2 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.72 (d, 1H, J = 3.6 Hz), 3.65-3.60 (m, 1H), 3.52-3.49 (m, 2H), 3.49-3.43 (m, 4H), 3.40-3.35 (m, 4H), 3.40-3.24 (m, 2H), 3.08-2.98 (m, 2H), 2.93-2.87 (m, 2H), 2.41 (q, 2H, J = 7.2 Hz), 2.07-2.01 (m, 2H), 1.78-1.66 (m, 6H), 1.33 (q, 2H, J = 10.4 Hz), 1.03 (t, 3H, J = 7.2 Hz).
424		518	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21 (s, 1H), 8.12 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.76 - 7.65 (m, 2H), 7.29 - 7.20 (m, 2H), 6.98 (d, J = 1.9 Hz, 1H), 6.57 (t, J = 5.5 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.51 (dd, J = 6.9, 3.3 Hz, 4H), 3.42 (dd, J = 6.7, 3.5 Hz, 4H), 3.03 (q, J = 6.6 Hz, 2H), 2.94 (d, J = 11.0 Hz, 2H), 2.79 (p, J = 6.6 Hz, 1H), 2.30 (dd, J = 12.6, 10.0 Hz, 2H), 1.78 (d, J = 12.5 Hz, 2H), 1.66 (tt, J = 12.8, 6.4 Hz, 2H), 1.42 (p, J = 7.3 Hz, 2H), 1.26 (dtd, J = 14.3, 8.3, 7.8, 2.9 Hz, 4H), 1.02 (d, J = 6.5 Hz, 6H), 0.86 (t, J = 6.9 Hz, 3H).
425		518	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 8.08 (d, 1H, J = 2.0 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.67 (d, 2H, J = 8.8 Hz), 6.99 (d, 2H, J = 8.8 Hz), 6.92 (d, 1H, J = 2.0 Hz), 5.94 (d, 1H, J = 5.2 Hz), 3.97 (t, 3H, J = 8.0 Hz), 3.68-3.64 (m, 4H), 3.47-3.46 (m, 2H), 3.45-3.43 (m, 5H), 3.29-3.27 (m, 2H), 3.26 (s, 3H), 2.93-2.91 (m, 4H), 2.79-2.76 (m, 4H), 1.15 (t, 3H, J = 7.2 Hz).
426		524	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.03 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.61 (d, 2H, J = 8.4 Hz), 6.89 (d, 1H, J = 1.6 Hz), 6.45 (d, 2H, J = 8.4 Hz), 5.95 (d, 2H, J = 5.6 Hz), 3.90-3.86 (m, 4H), 3.72-3.68 (m, 2H), 3.68-3.65 (m, 1H), 3.54-3.51 (m, 2H), 3.42-3.41 (m, 4H), 3.30-3.24 (m, 4H), 2.65-2.60 (m, 2H), 2.52-2.50 (m, 1H), 2.50-2.48 (m, 2H), 0.90-0.85 (m, 6H).

#	Structure	LCMS (M+1)	NMR
427		525	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 7.98 (s, 1H), 7.88 (d, 1H, J = 3.6 Hz), 7.73 (d, 1H, J = 8.4 Hz), 7.26 (t, 1H, J = 74.4 Hz), 6.95 (s, 1H), 6.88 (d, 1H, J = 6.8 Hz), 6.73 (s, 1H), 5.96 (d, 1H, J = 4.4 Hz), 3.91-3.90 (m, 2H), 3.69-3.67 (m, 2H), 3.53-3.45 (m, 4H), 3.23-3.19 (m, 4H), 2.55-2.51 (m, 4H), 2.36 (q, 2H, J = 7.2 Hz), 2.01-2.00 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz), 0.76-0.74 (m, 4H).
428		526	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.4 Hz), 7.25 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.45-3.44 (m, 4H), 3.39-3.37 (m, 4H), 3.35-3.36 (m, 3H), 3.15-3.04 (m, 6H), 2.58-2.54 (m, 1H), 2.17-2.10 (m, 2H), 1.90-1.86 (m, 2H), 1.82-1.77 (m, 2H), 1.74-1.66 (m, 4H), 1.06 (t, 3H, J = 7.2 Hz).
429		526	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.06 (s, 1H), 7.86 (d, 1H, J = 5.2 Hz), 7.64 (d, 2H, J = 8.4 Hz), 6.94 (d, 2H, J = 8.4 Hz), 6.93 (s, 1H), 5.94 (d, 1H, J = 5.2 Hz), 3.75-3.70 (m, 2H), 3.69-3.65 (m, 2H), 3.50-3.45 (m, 2H), 3.44-3.40 (m, 2H), 3.15-3.13 (m, 4H), 2.73-2.67 (m, 2H), 2.50-2.49 (m, 4H), 2.37 (q, 2H, J = 6.8 Hz), 2.05-2.02 (m, 2H), 1.73-1.70 (m, 2H), 1.63-1.53 (m, 2H), 1.44-1.36 (m, 2H), 1.03 (t, 3H, J = 6.8 Hz).
430		531	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.77 – 7.64 (m, 2H), 7.24 (d, J = 8.1 Hz, 2H), 7.00 (d, J = 1.9 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 4.75 (q, J = 9.1 Hz, 2H), 3.65 (s, 4H), 3.58 – 3.39 (m, 4H), 2.88 (d, J = 10.9 Hz, 3H), 2.70 (p, J = 6.7 Hz, 1H), 2.21 (t, J = 11.1 Hz, 2H), 1.76 (d, J = 12.4 Hz, 2H), 1.62 (tt, J = 12.6, 6.3 Hz, 2H), 0.99 (d, J = 6.5 Hz, 6H).
431		531	¹ H NMR (400 MHz, Chloroform-d) δ 7.92 (d, J = 1.6 Hz, 1H), 7.84 (d, J = 5.3 Hz, 1H), 7.59 (d, J = 7.9 Hz, 2H), 7.28 (d, J = 8.0 Hz, 2H), 6.69 (d, J = 1.3 Hz, 1H), 5.84 (d, J = 5.4 Hz, 1H), 3.87 (s, 2H), 3.75 (s, 2H), 3.47 (d, J = 25.7 Hz, 5H), 3.15 (s, 2H), 2.54 (d, J = 9.7 Hz, 5H), 2.08 (s, 1H), 1.91 (q, J = 20.9, 17.1 Hz, 3H), 1.81 – 1.72 (m, 1H), 1.66 (bs, 6H), 1.58 – 1.39 (m, 1H), 1.31 (s, 2H), 1.24 (d, J = 3.8 Hz, 4H), 1.16 (s, 1H).





#	Structure	LCMS (M+1)	NMR
432		531	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.44 (d, 1H, J = 2.0 Hz), 8.24 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.85 (d, 1H, J = 8.0 Hz), 7.67 (dd, 1H, J = 8.4, 2.0 Hz), 7.16 (d, 1H, J = 2.0 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.58 (d, 1H, J = 4.0 Hz), 3.76-3.72 (m, 2H), 3.68-3.64 (m, 2H), 3.53-3.48 (m, 2H), 3.45-3.40 (m, 2H), 2.93-2.88 (m, 2H), 2.76-2.69 (m, 1H), 2.57-2.54 (m, 1H), 2.25-2.20 (m, 2H), 1.86-1.81 (m, 2H), 1.80-1.72 (m, 2H), 1.70-1.60 (m, 5H), 1.44-1.34 (m, 2H), 1.25-1.14 (m, 2H), 0.99 (d, 6H, J = 6.8 Hz).
433		536	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.43 (d, 1H, J = 2.4 Hz), 8.43 (s, 1H), 8.03 (d, 1H, J = 1.2 Hz), 7.83 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.4 Hz), 7.61-7.58 (m, 1H), 7.36 (dd, 1H, J = 8.4, 5.2 Hz), 6.93 (d, 2H, J = 8.4 Hz), 6.88 (d, 1H, J = 1.2 Hz), 5.90 (d, 1H, J = 5.6 Hz), 3.69-3.67 (m, 4H), 3.42-3.30 (m, 8H), 3.16-3.13 (m, 4H), 2.37 (q, 2H, J = 6.4 Hz), 1.42-1.40 (m, 2H), 1.31-1.28 (m, 2H), 1.04 (t, 3H, J = 7.2 Hz).
434		539	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.76-3.72 (m, 2H), 3.72-3.68 (m, 2H), 3.51-3.47 (m, 2H), 3.45-3.40 (m, 2H), 2.79-2.75 (m, 2H), 2.72-2.69 (m, 4H), 2.18-2.15 (m, 2H), 2.04-2.03 (m, 2H), 1.82-1.81 (m, 1H), 1.75-1.72 (m, 3H), 1.61-1.57 (m, 3H), 1.43-1.40 (m, 3H), 0.99-0.97 (m, 6H).
435		539	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.77-3.72 (m, 2H), 3.70-3.65 (m, 2H), 3.50-3.45 (m, 2H), 3.45-3.40 (m, 2H), 2.82-2.79 (m, 2H), 2.75-2.67 (m, 4H), 2.18-2.15 (m, 2H), 2.07-2.06 (m, 2H), 1.84-1.83 (m, 1H), 1.80-1.75 (m, 3H), 1.60-1.54 (m, 3H), 1.46-1.40 (m, 3H), 0.99-0.96 (m, 6H).

#	Structure	LCMS (M+1)	NMR
436		540	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (s, 0.3H), 8.09 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.68 (d, 2H, J = 8.4 Hz), 7.00 (d, 2H, J = 8.4 Hz), 6.94 (d, 1H, J = 1.2 Hz), 6.54 (s, 0.4H), 5.96 (d, 1H, J = 5.6 Hz), 3.80-3.70 (m, 2H), 3.70-3.60 (m, 2H), 3.56-3.48 (m, 2H), 3.45-3.38 (m, 2H), 3.35-3.20 (m, 8H), 2.80-2.60 (m, 2H), 2.10-2.00 (m, 2H), 1.78-1.68 (m, 2H), 1.66-1.50 (m, 2H), 1.48-1.34 (m, 2H), 1.24-1.10 (m, 6H).
437		540	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.45 (s, 1H), 8.24 (s, 1H), 7.91 (d, 1H, J = 6.0 Hz), 7.85 (d, 1H, J = 8.0 Hz), 7.68 (d, 1H, J = 8.0 Hz), 7.16 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.76-3.73 (m, 2H), 3.69-3.63 (m, 2H), 3.53-3.48 (m, 2H), 3.46-3.42 (m, 2H), 2.90-2.87 (m, 2H), 2.74-2.68 (m, 3H), 2.57-2.49 (m, 1H), 2.24-2.19 (m, 2H), 2.05-2.02 (m, 2H), 1.78-1.70 (m, 4H), 1.70-1.57 (m, 4H), 1.45-1.39 (m, 2H), 0.98 (d, 6H, J = 6.8 Hz).
438		541	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 1.6 Hz), 7.83 (d, 1H, J = 5.6 Hz), 7.63 (d, 2H, J = 9.2 Hz), 7.35 (dd, 1H, J = 4.8, 2.4 Hz), 6.93 (d, 2H, J = 8.4 Hz), 6.96-6.91 (m, 3H), 5.91 (d, 1H, J = 5.6 Hz), 3.73-3.55 (m, 4H), 3.41-3.18 (m, 8H), 3.15-3.13 (m, 4H), 2.36 (q, 2H, J = 7.2 Hz), 1.46-1.43 (m, 2H), 1.21-1.18 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
439		541	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 2.0 Hz), 7.84 (d, 1H, J = 5.6 Hz), 7.63 (d, 2H, J = 8.8 Hz), 7.50 (dd, 1H, J = 5.2, 2.8 Hz), 7.23 (dd, 1H, J = 2.8, 1.2 Hz), 6.94-6.90 (m, 4H), 5.91 (d, 1H, J = 5.6 Hz), 3.72-3.69 (m, 4H), 3.42-3.30 (m, 8H), 3.16-3.13 (m, 4H), 2.37 (q, 2H, J = 7.2 Hz), 1.35-1.32 (m, 2H), 1.17-1.15 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
440		542	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 1.6 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.69 (d, 1H, J = 3.2 Hz), 7.64 (d, 2H, J = 8.8 Hz), 7.61 (d, 1H, J = 3.2 Hz), 6.93 (d, 2H, J = 8.8 Hz), 6.92 (d, 1H, J = 1.6 Hz), 5.93 (d, 1H, J = 5.2 Hz), 3.80-3.60 (m, 4H), 3.50-3.36 (m, 4H), 3.30-3.28 (m, 4H), 3.20-3.10 (m, 4H), 2.36 (q, 2H, J = 7.2 Hz), 1.60-1.55 (m, 2H), 1.50-1.40 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
441		543	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (br s, 1H), 8.04 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.4 Hz), 6.90 (d, 1H, J = 1.6 Hz), 6.45 (d, 2H, J = 8.4 Hz), 5.95 (d, 2H, J = 5.6 Hz), 3.88 (s, 4H), 3.75-3.71 (m, 2H), 3.69-3.64 (m, 2H), 3.52-3.48 (m, 2H), 3.48-3.45 (m, 4H), 3.45-3.33 (m, 3H), 2.56-2.52 (m, 1H), 2.50-2.41 (m, 1H), 1.88-1.82 (m, 2H), 1.72-1.65 (m, 2H), 1.43-1.33 (m, 2H), 1.26-1.18 (m, 2H), 0.87 (d, 6H, J = 6.4 Hz).
442		552	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.34 (s, 1H), 8.03 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.61 (d, 2H, J = 8.8 Hz), 6.90 (d, 2H, J = 1.6 Hz), 6.44 (d, 1H, J = 8.8 Hz), 5.95 (d, 1H, J = 5.6 Hz), 3.86 (s, 4H), 3.78-3.70 (m, 2H), 3.69-3.62 (m, 2H), 3.50-3.45 (m, 2H), 3.44-3.38 (m, 2H), 3.25 (s, 4H), 2.76-2.66 (m, 2H), 2.26-2.20 (m, 1H), 2.10-2.01 (m, 2H), 1.75-1.71 (m, 2H), 1.62-1.55 (m, 2H), 1.46-1.40 (m, 2H), 1.10-1.07 (m, 6H).
443		553	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.03 (d, 1H, J = 1.2 Hz), 7.83 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.8 Hz), 7.37 (dd, 1H, J = 14.4, 8.0 Hz), 7.07-7.02 (m, 2H), 6.99-6.95 (m, 1H), 6.92 (d, 2H, J = 8.8 Hz), 6.89 (d, 1H, J = 1.2 Hz), 5.90 (d, 1H, J = 5.2 Hz), 3.73-3.55 (m, 4H), 3.41-3.18 (m, 8H), 3.14 (t, 4H, J = 4.4 Hz), 2.36 (q, 2H, J = 7.2 Hz), 1.41-1.37 (m, 2H), 1.27-1.23 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
444		554	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (s, 1H), 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.0 Hz), 7.24 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.97-3.89 (m, 1H), 3.85-3.70 (m, 2H), 3.70-3.66 (m, 1H), 3.36-3.26 (m, 4H), 3.06-2.98 (m, 1H), 2.79-2.71 (m, 2H), 2.61-2.54 (m, 2H), 2.42-2.32 (m, 3H), 2.29-2.17 (m, 1H), 1.84-1.74 (m, 1H), 1.51-1.44 (m, 1H), 1.28-1.21 (m, 1H), 1.08 (t, 3H, J = 7.2 Hz).






#	Structure	LCMS (M+1)	NMR
445		557	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.93 (d, J = 1.8 Hz, 1H), 7.82 (d, J = 5.4 Hz, 1H), 7.68 – 7.59 (m, 2H), 7.31 – 7.21 (m, 2H), 6.86 (d, J = 1.8 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 3.86 – 3.75 (m, 2H), 3.65 (dd, J = 6.9, 3.5 Hz, 2H), 3.56 – 3.40 (m, 4H), 3.24 – 3.06 (m, 3H), 2.80 – 2.67 (m, 2H), 2.54 (dh, J = 14.0, 7.6 Hz, 4H), 2.14 (td, J = 11.8, 2.9 Hz, 2H), 1.91 – 1.75 (m, 5H), 1.15 (t, J = 7.3 Hz, 3H).
446		565	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.76 – 7.62 (m, 2H), 7.57 (dd, J = 7.6, 1.6 Hz, 2H), 7.36 (t, J = 7.6 Hz, 2H), 7.25 (dd, J = 7.7, 5.4 Hz, 3H), 6.98 (d, J = 1.9 Hz, 1H), 5.95 (d, J = 5.5 Hz, 1H), 5.63 (s, 1H), 3.77 – 3.63 (m, 2H), 3.58 (d, J = 5.8 Hz, 2H), 3.44 (q, J = 5.4 Hz, 4H), 2.99 (dd, J = 17.5, 9.4 Hz, 3H), 2.59 (ddt, J = 11.7, 9.7, 6.1 Hz, 4H), 2.34 (q, J = 7.1 Hz, 2H), 2.02 – 1.87 (m, 2H), 1.81 – 1.55 (m, 4H), 1.02 (t, J = 7.1 Hz, 3H).
447		569	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.03 (d, 1H, J = 1.6Hz), 7.83 (d, 1H, J = 5.2 Hz), 7.62 (d, 2H, J = 8.8 Hz), 7.38-7.34 (m, 1H), 7.28 (d, 1H, J = 8.4 Hz), 7.18 (s, 1H), 7.17 (d, 1H, J = 8.0 Hz), 6.92 (d, 2H, J = 8.8 Hz), 6.89 (d, 1H, J = 1.6 Hz), 5.90 (d, 1H, J = 5.6 Hz), 3.73-3.55 (m, 4H), 3.41-3.18 (m, 8H), 3.14 (t, 4H, J = 4.8 Hz), 2.36 (q, 2H, J = 7.2 Hz), 1.41-1.37 (m, 2H), 1.27-1.24 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz).
448		479	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.68 (s, 1H), 8.31 (s, 1H), 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.61-3.60 (m, 4H), 3.56-3.54 (m, 4H), 3.47-3.46 (m, 4H), 2.73-2.67 (m, 1H), 2.57-2.54 (m, 4H), 1.22 (t, 3H, J = 7.2 Hz), 1.00 (d, 6H, J = 6.8 Hz).
449		402	

#	Structure	LCMS (M+1)	NMR
450		417	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 9.07 (d, 1H, J = 2.0 Hz), 8.38 (d, 1H, J = 1.2 Hz), 8.29 (dd, 1H, J = 8.4, 2.0 Hz), 7.94 (d, 2H, J = 5.6 Hz), 7.54 (d, 1H, J = 8.4 Hz), 7.21 (d, 1H, J = 1.2 Hz), 6.01 (d, 1H, J = 5.6 Hz), 4.66-4.62 (m, 1H), 3.94-3.91 (m, 2H), 3.72-3.70 (m, 2H), 3.57-3.54 (m, 2H), 3.53-3.49 (m, 2H), 3.30-3.19 (m, 2H), 2.40-2.34 (m, 1H), 1.99-1.85 (m, 4H), 0.79-0.75 (m, 4H).
451		417	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 9.06 (s, 1H), 8.36 (s, 1H), 8.26 (dd, 1H, J1 = 8.4 Hz, J2 = 2.0 Hz), 7.94 (d, 1H, J = 5.6 Hz), 7.53 (d, 1H, J = 8.4 Hz), 7.21 (s, 1H), 5.99 (d, 1H, J = 5.6 Hz), 4.62-4.58 (m, 1H), 4.02-3.98 (m, 2H), 3.95-3.91 (m, 2H), 3.72-3.50 (m, 4H), 3.30-3.16 (m, 2H), 2.35-2.30 (m, 1H), 2.08-2.04 (m, 1H), 1.99-1.87 (m, 3H), 0.79-0.73 (m, 4H).
452		417	¹ H NMR (400 MHz, DMSO-d6) δ 8.52 (d, J = 2.2 Hz, 1H), 8.25 (d, J = 1.8 Hz, 1H), 7.92 (d, J = 5.4 Hz, 1H), 7.86 (d, J = 8.1 Hz, 1H), 7.75 (dd, J = 8.2, 2.3 Hz, 1H), 7.17 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.06 (t, J = 7.6 Hz, 1H), 3.93 (s, 2H), 3.70 (s, 2H), 3.52 (d, J = 32.7 Hz, 4H), 3.01 (ddd, J = 9.9, 7.6, 5.3 Hz, 1H), 2.89 (ddd, J = 9.9, 8.2, 6.6 Hz, 1H), 2.13 (dtd, J = 12.3, 7.7, 4.9 Hz, 1H), 2.02 (tt, J = 7.7, 4.9 Hz, 1H), 1.85 - 1.69 (m, 2H), 1.58 - 1.44 (m, 1H), 0.76 (tt, J = 7.9, 2.9 Hz, 4H).
453		417	¹ H NMR (400 MHz, DMSO-d6) δ 8.48 (d, J = 2.2 Hz, 1H), 8.21 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.82 (d, J = 8.1 Hz, 1H), 7.71 (dd, J = 8.2, 2.3 Hz, 1H), 7.14 (d, J = 1.8 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 4.01 (t, J = 7.6 Hz, 1H), 3.89 (s, 2H), 3.67 (s, 2H), 3.53 (s, 2H), 3.45 (s, 2H), 2.98 (ddd, J = 9.9, 7.5, 5.4 Hz, 1H), 2.85 (ddd, J = 9.9, 8.2, 6.6 Hz, 1H), 2.09 (dtd, J = 12.1, 7.6, 4.8 Hz, 1H), 2.02 - 1.93 (m, 1H), 1.80 - 1.66 (m, 2H), 1.53 - 1.42 (m, 1H), 0.72 (tt, J = 7.9, 2.9 Hz, 4H).
454		417	






#	Structure	LCMS (M+1)	NMR
455		420	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 8.2 Hz, 2H), 7.40 (d, J = 8.0 Hz, 2H), 6.93 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.4 Hz, 1H), 4.03 (s, 2H), 3.85 (s, 2H), 3.63 (s, 2H), 3.51 (d, J = 25.3 Hz, 4H), 2.80 (s, 2H), 2.60 (d, J = 9.9 Hz, 2H), 2.02 (tt, J = 7.8, 4.7 Hz, 1H), 1.51 (d, J = 6.6 Hz, 3H), 0.92 (dt, J = 4.9, 2.8 Hz, 2H), 0.86 (ddt, J = 7.5, 4.6, 2.6 Hz, 2H).
456		420	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 9.04 (s, 1H), 8.34 (s, 1H), 8.25 (d, 1H, J = 8.0 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.52 (d, 1H, J = 8.0 Hz), 7.17 (s, 1H), 6.65-6.62 (m, 1H), 6.00 (d, 1H, J = 5.6 Hz), 4.56-4.54 (m, 1H), 3.53-3.51 (m, 4H), 3.45-3.44 (m, 4H), 3.25-3.22 (m, 1H), 3.21-3.16 (m, 1H), 3.08 (quintet, 2H, J = 7.2 Hz), 2.32-2.30 (m, 1H), 1.91-1.86 (m, 3H), 1.03 (t, 3H, J = 7.2 Hz).
457		420	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 9.05 (d, 1H, J = 8.4 Hz), 8.36 (s, 1H), 8.26 (dd, 1H, J ₁ = 8.0 Hz, J ₂ = 2.0 Hz), 7.93 (d, 1H, J = 5.2 Hz), 7.53 (d, 1H, J = 8.0 Hz), 7.17 (d, 1H, J = 1.2 Hz), 6.64 (t, 1H, J = 5.2 Hz), 6.01 (d, 1H, J = 5.6 Hz), 4.58-4.54 (m, 1H), 3.54-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.23-3.22 (m, 1H), 3.20-3.11 (m, 1H), 3.09-3.06 (m, 2H), 2.34-2.28 (m, 1H), 1.94-1.85 (m, 3H), 1.04 (t, 3H, J = 7.2 Hz).
458		428	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.99 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.5 Hz, 1H), 7.80 – 7.67 (m, 2H), 7.58 – 7.46 (m, 2H), 6.94 (d, J = 1.8 Hz, 1H), 6.20 (tt, J = 3.5, 1.7 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 4.01 (s, 2H), 3.93 – 3.75 (m, 4H), 3.61 (s, 3H), 3.53 (d, J = 5.3 Hz, 3H), 3.48 (t, J = 6.1 Hz, 2H), 2.84 (tq, J = 5.8, 1.9 Hz, 2H), 2.01 (tt, J = 7.8, 4.7 Hz, 1H), 0.99 – 0.90 (m, 2H), 0.86 (ddt, J = 7.5, 4.6, 2.5 Hz, 2H).

#	Structure	LCMS (M+1)	NMR
459		430	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.76 - 7.66 (m, 2H), 7.54 - 7.42 (m, 2H), 7.01 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.91 (qd, J = 5.9, 2.8 Hz, 2H), 3.70 (s, 2H), 3.50 (d, J = 29.4 Hz, 4H), 3.00 (ddd, J = 10.6, 8.3, 5.9 Hz, 1H), 2.76 (ddd, J = 10.3, 8.5, 4.7 Hz, 1H), 2.06 - 1.95 (m, 2H), 1.82 - 1.69 (m, 2H), 1.58 - 1.46 (m, 1H), 1.35 (s, 3H), 0.75 (tt, J = 7.9, 2.9 Hz, 4H).
460		430	
461		431	¹ H NMR (400 MHz, MeOD) δ ppm 8.55-8.48 (m, 1H), 7.86 (d, 1H, J = 2.0 Hz), 7.82 (d, 1H, J = 5.6 Hz), 7.60 (d, 2H, J = 8.8 Hz), 6.82 (d, 1H, J = 2.0 Hz), 6.70 (d, 2H, J = 8.8 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.07-4.01 (m, 2H), 3.98-3.95 (m, 1H), 3.89-3.83 (m, 2H), 3.65-3.58 (m, 4H), 3.56-3.50 (m, 2H), 3.42-3.36 (m, 2H), 2.49-2.40 (m, 1H), 2.14-2.07 (m, 1H), 2.05-2.00 (m, 1H), 0.97-0.90 (m, 2H), 0.89-0.86 (m, 2H).
462		431	¹ H-NMR (400 MHz, MeOD) δ ppm 8.51 (s, 1H), 7.82 (d, 1H, J = 1.6 Hz), 7.78 (d, 1H, J = 5.2 Hz), 7.57 (d, 2H, J = 8.8 Hz), 6.78 (d, 1H, J = 1.2 Hz), 6.67 (d, 2H, J = 8.8 Hz), 5.94 (d, 1H, J = 5.6 Hz), 3.98-3.95 (m, 3H), 3.82-3.80 (m, 2H), 3.60-3.54 (m, 4H), 3.48-3.46 (m, 2H), 3.40-3.36 (m, 2H), 2.42-2.40 (m, 1H), 2.13-2.11 (m, 1H), 2.00-1.98 (m, 1H), 0.90-0.88 (m, 2H), 0.84-0.80 (m, 2H).
463		431.2	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.63 (dd, J = 2.1, 1.0 Hz, 1H), 8.21 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.83 - 7.76 (m, 2H), 7.13 (d, J = 1.8 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 3.90 (br.s, 2H), 3.67 (br.s, 2H), 3.52 (br.s, 2H), 3.44 (br.s, 2H), 2.99 (dt, J = 10.8, 7.1 Hz, 1H), 2.71 (ddd, J = 13.5, 9.2, 4.7 Hz, 1H), 1.98 (ddd, J = 10.8, 8.2, 5.5 Hz, 2H), 1.75 (tt, J = 8.7, 6.2 Hz, 2H), 1.57 - 1.43 (m, 1H), 1.35 (s, 3H), 0.72 (ddd, J = 10.9, 5.0, 2.4 Hz, 4H).

#	Structure	LCMS (M+1)	NMR
464		432	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.94 (d, J = 1.8 Hz, 1H), 7.81 (d, J = 5.4 Hz, 1H), 7.73 – 7.65 (m, 2H), 7.39 (d, J = 8.0 Hz, 2H), 6.89 (d, J = 1.8 Hz, 1H), 5.92 (d, J = 5.4 Hz, 1H), 3.96 (d, J = 5.4 Hz, 2H), 3.84 (ddt, J = 16.0, 8.6, 4.2 Hz, 5H), 3.68 – 3.53 (m, 3H), 3.53 – 3.38 (m, 3H), 3.31 (p, J = 1.6 Hz, 1H), 3.11 – 2.91 (m, 2H), 1.98 (ddd, J = 9.3, 6.6, 4.0 Hz, 1H), 0.92 (ddd, J = 5.9, 4.7, 2.5 Hz, 2H), 0.85 (tdd, J = 7.4, 5.4, 1.9 Hz, 2H).
465		434	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.48 (d, J = 2.2 Hz, 1H), 8.20 (d, J = 1.7 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.82 (d, J = 8.2 Hz, 1H), 7.70 (dd, J = 8.2, 2.3 Hz, 1H), 7.10 (d, J = 1.9 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.95 (d, J = 5.5 Hz, 1H), 4.01 (t, J = 7.6 Hz, 1H), 3.75 (dt, J = 13.5, 6.7 Hz, 1H), 3.48 (dd, J = 6.8, 3.2 Hz, 4H), 3.44 – 3.36 (m, 4H), 2.97 (ddd, J = 9.9, 7.6, 5.3 Hz, 1H), 2.85 (ddd, J = 9.9, 8.2, 6.7 Hz, 1H), 2.09 (dtd, J = 12.4, 7.7, 5.0 Hz, 1H), 1.81 – 1.64 (m, 2H), 1.52 – 1.42 (m, 1H), 1.04 (d, J = 6.6 Hz, 6H).
466		434	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.48 (d, J = 2.2 Hz, 1H), 8.20 (d, J = 1.7 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.82 (d, J = 8.2 Hz, 1H), 7.70 (dd, J = 8.2, 2.3 Hz, 1H), 7.10 (d, J = 1.9 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.95 (d, J = 5.5 Hz, 1H), 4.01 (t, J = 7.6 Hz, 1H), 3.75 (dt, J = 13.5, 6.7 Hz, 1H), 3.48 (dd, J = 6.8, 3.2 Hz, 4H), 3.44 – 3.36 (m, 4H), 2.97 (ddd, J = 9.9, 7.6, 5.3 Hz, 1H), 2.85 (ddd, J = 9.9, 8.2, 6.7 Hz, 1H), 2.09 (dtd, J = 12.4, 7.7, 5.0 Hz, 1H), 1.81 – 1.64 (m, 2H), 1.52 – 1.42 (m, 1H), 1.04 (d, J = 6.6 Hz, 6H).
467		434	





#	Structure	LCMS (M+1)	NMR
468		436	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (dd, J = 5.9, 2.5 Hz, 3H), 7.71 – 7.55 (m, 2H), 4.55 (dd, J = 10.8, 3.6 Hz, 1H), 4.21 (q, J = 7.1 Hz, 2H), 4.18 – 4.09 (m, 2H), 4.09 – 3.91 (m, 2H), 3.82 – 3.69 (m, 4H), 3.63 – 3.53 (m, 4H), 3.49 – 3.37 (m, 2H), 3.30 (p, J = 1.7 Hz, 1H), 2.65 (p, J = 1.9 Hz, 1H), 1.33 (t, J = 7.1 Hz, 3H).
469		437	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (s, 1H), 7.91 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.80 (s, 1H), 6.79 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 8.0 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.11-4.10 (m, 1H), 3.81-3.76 (m, 1H), 3.51-3.50 (m, 4H), 3.41-3.39 (m, 4H), 3.21-3.07 (m, 1H), 2.95-2.90 (m, 1H), 2.75-2.70 (m, 1H), 2.46-2.43 (m, 2H), 2.15-2.11 (m, 1H), 1.90 1.87 (m, 1H), 1.74 1.70 (m, 1H), 1.53 1.50 (m, 1H), 1.08 (d, 6H, J = 6.4 Hz).
470		437	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (s, 1H), 7.91 (s, 1H), 7.85 (d, 1H, J = 5.6 Hz), 7.80 (s, 1H), 6.78 (s, 1H), 6.33 (d, 1H, J = 8.0 Hz), 5.96 (d, 1H, J = 5.6 Hz), 4.14-4.08 (m, 1H), 3.80-3.74 (m, 1H), 3.48-3.58 (m, 6H), 3.22-3.16 (m, 4H), 2.90-2.86 (m, 1H), 2.78-2.74 (m, 1H), 2.46-2.43 (m, 1H), 2.13-2.10 (m, 1H), 1.89-1.85 (m, 1H), 1.73-1.69 (m, 1H), 1.52-1.48 (m, 1H), 1.08 (d, 6H, J = 6.8 Hz).
471		444	
472		444	¹ H NMR (500 MHz, Methanol-d ₄) δ 8.02 (d, J = 1.8 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.80 – 7.70 (m, 2H), 7.56 (d, J = 8.0 Hz, 1H), 6.98 (d, J = 2.0 Hz, 1H), 6.02 (d, J = 5.4 Hz, 1H), 4.91 (s, 2H), 4.04 (s, 2H), 3.86 (s, 2H), 3.60 (d, J = 31.4 Hz, 5H), 3.22 (t, J = 7.4 Hz, 6H), 2.22 (s, 1H), 2.03 (s, 1H), 1.44 (t, J = 7.3 Hz, 6H), 0.93 (d, J = 4.1 Hz, 2H), 0.88 (d, J = 7.8 Hz, 1H).





#	Structure	LCMS (M+1)	NMR
473		446	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.49 (d, 2H, J = 8.4 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.80-3.60 (m, 2H), 3.60-3.55(m, 2H), 3.50-3.46 (m, 4H), 3.35-3.29 (m, 1H), 2.54 2.53 (m, 1H), 2.02-1.99 (m, 1H), 1.37 (s, 6H), 0.84 (d, 6H, J = 6.4 Hz), 0.76-0.72 (m, 4H).
474		428	1H NMR (400 MHz, DMSO-d6) δ 8.16 (s, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.79 (d, J = 8.0 Hz, 2H), 7.37 (d, J = 8.0 Hz, 2H), 7.06 - 6.98 (m, 1H), 5.94 (d, J = 5.5 Hz, 1H), 3.88 (br.s, 2H), 3.66 (br.s, 2H), 3.47 (br.d, J = 27.3 Hz, 4H), 2.90 - 2.72 (m, 4H), 2.05 - 1.72 (m, 5H), 0.72 (dd, J = 9.3, 6.3 Hz, 4H).
475		449	1H NMR (400 MHz, DMSO-d6) δ 8.48 (d, J = 2.1 Hz, 1H), 8.21 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.82 (d, J = 8.1 Hz, 1H), 7.71 (dd, J = 8.2, 2.3 Hz, 1H), 7.11 (d, J = 1.8 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 5.33 - 5.25 (m, 1H), 4.75 (t, J = 6.9 Hz, 2H), 4.49 (dd, J = 7.5, 5.1 Hz, 2H), 4.02 (t, J = 7.6 Hz, 1H), 3.66 (br.s, 2H), 3.56 (br.s, 2H), 3.50 - 3.41 (m, 4H), 2.98 (ddd, J = 9.9, 7.6, 5.4 Hz, 1H), 2.85 (ddd, J = 9.9, 8.2, 6.7 Hz, 1H), 2.10 (dtd, J = 12.4, 7.7, 5.0 Hz, 1H), 1.81 - 1.64 (m, 2H), 1.54 - 1.41 (m, 1H).
476		449	1H NMR (400 MHz, DMSO-d6) δ 8.48 (d, J = 2.1 Hz, 1H), 8.21 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.82 (d, J = 8.1 Hz, 1H), 7.71 (dd, J = 8.2, 2.3 Hz, 1H), 7.11 (d, J = 1.8 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 5.33 - 5.25 (m, 1H), 4.75 (t, J = 6.9 Hz, 2H), 4.49 (dd, J = 7.5, 5.1 Hz, 2H), 4.02 (t, J = 7.6 Hz, 1H), 3.66 (br.s, 2H), 3.56 (br.s, 2H), 3.50 - 3.41 (m, 4H), 2.98 (ddd, J = 9.9, 7.6, 5.4 Hz, 1H), 2.85 (ddd, J = 9.9, 8.2, 6.7 Hz, 1H), 2.10 (dtd, J = 12.4, 7.7, 5.0 Hz, 1H), 1.81 - 1.64 (m, 2H), 1.54 - 1.41 (m, 1H).

#	Structure	LCMS (M+1)	NMR
477		449	
478		449	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.43 (s, 1H), 8.00 (d, J = 1.8 Hz, 1H), 7.88 – 7.74 (m, 3H), 7.56 – 7.40 (m, 2H), 6.93 (d, J = 1.9 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.37 (dd, J = 10.7, 3.5 Hz, 1H), 4.08 (dd, J = 12.5, 3.3 Hz, 2H), 3.93 (p, J = 6.6 Hz, 1H), 3.90 – 3.77 (m, 2H), 3.66 – 3.58 (m, 4H), 3.58 – 3.46 (m, 4H), 3.39 – 3.29 (m, 4H), 1.24 – 1.07 (m, 6H).
479		452	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.5 Hz), 7.91 (d, 1H, J = 5.5 Hz), 7.82 (d, 2H, J = 8.5 Hz), 7.43 (d, 2H, J = 8.5 Hz), 7.04 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 6.0 Hz), 4.09 (q, 2H, J = 7.0 Hz), 3.65-3.55 (m, 4H), 3.50-3.38 (m, 4H), 2.98-2.85 (m, 3H), 2.70-2.55 (m, 1H), 2.45-2.35 (m, 1H), 2.23-2.05 (m, 1H), 1.99-1.93 (m, 1H), 1.80-1.70 (m, 1H), 1.60-1.50 (m, 1H), 1.22 (t, 3H, J = 7.0 Hz).
480		452	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 6.8 Hz), 3.65-3.50 (m, 4H), 3.50-3.38 (m, 4H), 2.98-2.80 (m, 3H), 2.70-2.55 (m, 1H), 2.45-2.15 (m, 2H), 2.03-1.95 (m, 1H), 1.85-1.70 (m, 1H), 1.65-1.48 (m, 1H), 1.22 (t, 3H, J = 6.8 Hz).
481		455	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21 (d, J = 1.7 Hz, 1H), 7.98 – 7.84 (m, 3H), 7.57 – 7.44 (m, 2H), 7.07 (d, J = 1.9 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 3.92 (s, 2H), 3.71 (s, 2H), 3.51 (d, J = 25.6 Hz, 4H), 3.27 – 3.07 (m, 3H), 2.90 (t, J = 12.8 Hz, 2H), 2.13 (d, J = 13.2 Hz, 2H), 2.08 – 1.90 (m, 4H), 0.89 – 0.60 (m, 4H).

#	Structure	LCMS (M+1)	NMR
482		456	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.77 - 7.67 (m, 2H), 7.34 - 7.24 (m, 2H), 7.01 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.92 (br.s, 2H), 3.69 (br.s, 2H), 3.50 (br.d, J = 28.4 Hz, 4H), 2.94 - 2.78 (m, 6H), 2.02 (tt, J = 7.6, 4.9 Hz, 1H), 1.74 - 1.59 (m, 6H), 0.75 (tt, J = 7.9, 2.9 Hz, 4H).
483		459	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.46 (s, 1H), 8.02 (d, J = 1.8 Hz, 1H), 7.89 - 7.81 (m, 3H), 7.63 - 7.52 (m, 2H), 6.94 (d, J = 1.9 Hz, 1H), 6.02 (d, J = 5.4 Hz, 1H), 5.49 (d, J = 5.3 Hz, 1H), 4.18 (q, J = 7.1 Hz, 2H), 3.72 (d, J = 5.8 Hz, 4H), 3.59 (d, J = 13.5 Hz, 2H), 3.52 (t, J = 5.1 Hz, 4H), 3.38 (dd, J = 13.1, 2.9 Hz, 2H), 2.44 (d, J = 14.5 Hz, 2H), 2.39 - 2.23 (m, 2H), 1.30 (t, J = 7.1 Hz, 3H).
484		459	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.94 (s, 1H), 8.29 (s, 1H), 8.18 (s, 1H), 8.17 (d, 1H, J = 8.0 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.54 (d, 1H, J = 8.0 Hz), 7.15 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 4.02-3.98 (m, 1H), 3.94-3.91 (m, 2H), 3.72-3.70 (m, 2H), 3.55-3.51 (m, 2H), 3.50-3.45 (m, 2H), 3.16-3.14 (m, 1H), 2.86-2.83 (m, 1H), 2.75-2.69 (m, 1H), 2.23-2.18 (m, 1H), 2.04-2.01 (m, 1H), 1.85-1.79 (m, 2H), 1.72-1.69 (m, 1H), 0.99-0.97 (m, 6H), 0.78-0.74 (m, 4H).
485		459	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.48 (d, 1H, J = 1.6 Hz), 8.26 (d, 1H, J = 1.2 Hz), 7.93 (d, 1H, J = 5.2 Hz), 7.86 (d, 1H, J = 8.4 Hz), 7.72 (q, 1H, J1 = 8.4 Hz, J2 = 2.4 Hz), 7.14 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 4.01-3.92 (m, 2H), 3.73-3.70 (m, 2H), 3.57-3.36 (m, 4H), 3.30-3.28 (m, 1H), 3.02-2.96 (m, 1H), 2.76-2.70 (m, 2H), 2.50-2.48 (m, 1H), 2.44-2.38 (m, 1H), 2.29-2.20 (m, 1H), 2.06-1.99 (m, 1H), 1.81-1.74 (m, 1H), 1.08-1.04 (t, 6H, J = 6.8 Hz), 0.79-0.75 (m, 4H).

#	Structure	LCMS (M+1)	NMR
486		459	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.51 (d, 1H, J = 2.0 Hz), 8.13 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.65 (d, 1H, J = 2.4 Hz), 7.60 (d, 1H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.82 (d, 1H, J = 5.2 Hz), 3.95-3.90 (m, 2H), 3.89-3.85 (m, 2H), 3.62-3.57 (m, 2H), 3.51-3.47 (m, 2H), 3.48-3.42 (m, 1H), 3.35-3.31 (m, 1H), 3.11-3.05 (m, 1H), 2.93-2.85 (m, 1H), 2.71-2.65 (m, 2H), 2.45-2.38 (m, 1H), 2.01-1.89 (m, 1H), 1.79-1.74 (m, 1H), 1.55-1.45 (m, 1H), 1.22-1.19 (m, 6H), 0.84-0.78 (m, 4H).
487		459	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.90 (d, 1H, J = 2.0 Hz), 8.26 (d, 1H, J = 1.6 Hz), 8.13 (dd, 1H, J ₁ = 8.0 Hz, J ₂ = 2.0 Hz), 7.93 (d, 1H, J = 5.2 Hz), 7.52 (d, 1H, J = 8.0 Hz), 7.13 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.93-3.92 (m, 2H), 3.85-3.81 (m, 1H), 3.71-3.70 (m, 2H), 3.56-3.49 (m, 4H), 3.10-3.05 (m, 1H), 2.75-2.69 (m, 1H), 2.62-2.59 (m, 1H), 2.20-2.11 (m, 1H), 2.06-2.00 (m, 1H), 1.83-1.72 (m, 2H), 1.68-1.61 (m, 1H), 0.94 (d, 6H, J = 6.4 Hz), 0.79-0.75 (m, 4H).
488		460	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.00 (dd, J = 5.1, 1.8 Hz, 1H), 7.86 (dd, J = 5.5, 3.8 Hz, 1H), 7.82 – 7.74 (m, 2H), 7.51 – 7.40 (m, 2H), 6.96 (t, J = 2.4 Hz, 1H), 6.02 (d, J = 5.5 Hz, 1H), 4.06 (d, J = 12.5 Hz, 2H), 3.86 (s, 2H), 3.64 (s, 2H), 3.56 (s, 2H), 3.27 – 3.11 (m, 4H), 3.01 (d, J = 11.9 Hz, 3H), 2.27 (d, J = 14.3 Hz, 2H), 2.13 – 1.96 (m, 1H), 0.99 – 0.81 (m, 3H).
489		460	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 9.03-9.00 (m, 1H), 8.46-8.35 (m, 1H), 8.25 (d, 1H, J = 1.6 Hz), 7.93 (d, 1H, J = 5.2 Hz), 7.91 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.09 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 3.94-3.93 (m, 2H), 3.74-3.73 (m, 2H), 3.57-3.53 (m, 2H), 3.52-3.49 (m, 2H), 3.45-3.42 (m, 1H), 3.24-3.20 (m, 1H), 3.18-3.14 (m, 1H), 3.00 (s, 3H), 2.95-2.92 (m, 1H), 2.23-2.22 (m, 1H), 2.11-2.09 (m, 2H), 2.00-1.83 (m, 2H), 0.79-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
490		460	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.34 (s, 1H), 8.22 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.84 (d, 2H, J = 8.4 Hz), 7.42 (d, 2H, J = 8.4 Hz), 7.08 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.93-3.92 (m, 2H), 3.72-3.71 (m, 2H), 3.58-3.53 (m, 2H), 3.51-3.45 (m, 2H), 3.17-3.14 (m, 1H), 3.06-3.035 (m, 1H), 2.98 (s, 3H), 2.91-2.88 (m, 1H), 2.70-2.69 (m, 1H), 2.17-2.16 (m, 1H), 2.05-2.01 (m, 2H), 1.75-1.74 (m, 1H), 1.64-1.60 (m, 1H), 0.79-0.73 (m, 4H).
491		461	¹ H NMR (400 MHz, Methanol-d ₄) δ ppm 8.53 (d, J = 1.9 Hz, 1H), 8.19 (t, J = 2.3 Hz, 1H), 7.92 – 7.73 (m, 3H), 7.17 (t, J = 2.1 Hz, 1H), 5.96 (dd, J = 5.6, 1.9 Hz, 1H), 4.03 (d, J = 24.3 Hz, 2H), 3.83 (s, 2H), 3.62 (d, J = 6.6 Hz, 2H), 3.54 – 3.53 (m, 3H), 3.30 (p, J = 1.6 Hz, 2H), 3.09 (tt, J = 12.4, 2.3 Hz, 2H), 3.01 – 2.89 (m, 2H), 2.17 – 2.05 (m, 2H), 2.06 – 1.88 (m, 3H), 0.97 – 0.89 (m, 2H), 0.88 (d, J = 2.6 Hz, 2H).
492		461	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.91 (d, 1H, J = 5.5 Hz), 7.78 (d, 2H, J = 8.0 Hz), 7.39 (d, 2H, J = 8.0 Hz), 7.00 (s, 1H), 6.33 (d, 2H, J = 7.5 Hz), 5.99 (d, 1H, J = 5.5 Hz), 3.89-3.80 (m, 1H), 3.79-3.65 (m, 8H), 3.45-3.41 (m, 1H), 3.29-3.19 (m, 2H), 3.13-3.11 (m, 2H), 2.51-2.15 (m, 2H), 1.80-1.79 (m, 2H), 1.23 (s, 3H), 1.08 (d, 6H, J = 6.5 Hz).
493		462	¹ H NMR (400 MHz, Methanol-d ₄) δ ppm 7.98 (d, J = 1.8 Hz, 1H), 7.85 (dd, J = 5.5, 2.3 Hz, 1H), 7.77 – 7.66 (m, 2H), 7.45 – 7.33 (m, 2H), 6.94 (d, J = 1.9 Hz, 1H), 6.00 (dd, J = 5.6, 2.2 Hz, 1H), 5.49 (d, J = 2.1 Hz, 1H), 4.04 (s, 2H), 3.85 (s, 2H), 3.63 (s, 2H), 3.54 (s, 2H), 3.42 (s, 1H), 2.75 (s, 2H), 2.53 (s, 2H), 2.03 (tt, J = 7.9, 4.8 Hz, 1H), 1.84 (d, J = 6.9 Hz, 5H), 1.49 (dd, J = 6.7, 2.2 Hz, 3H), 1.00 – 0.86 (m, 2H), 0.86 (q, J = 2.8 Hz, 1H).






#	Structure	LCMS (M+1)	NMR
494		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.92 (d, 1H, J = 2.0 Hz), 8.27 (s, 1H), 8.27(s, 1H), 8.16 (dd, 2H, J = 8.4, 2.0 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.53 (d, 1H, J = 8.4 Hz), 7.11 (s, 1H), 6.62 (t, 1H, J = 5.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.97-3.93 (m, 1H), 3.53-3.51 (m, 4H), 3.45-3.44 (m, 4H), 3.16-3.12 (m, 1H), 3.07 (quintet, 2H, J = 6.8 Hz), 2.84-2.77 (m, 1H), 2.71-2.65 (m, 1H), 2.24-2.15 (m, 1H), 1.86-1.75 (m, 2H), 1.72-1.65 (m, 1H), 1.03 (t, 3H, J = 6.8 Hz), 0.98-0.96 (m, 6H).
495		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.47 (d, 1H, J = 1.6 Hz), 8.23 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.85 (d, 1H, J = 8.0 Hz), 7.71 (d, 1H, J = 2.4 Hz), 7.69 (d, 1H, J = 2.0 Hz), 7.13 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.52-3.51 (m, 4H), 3.48-3.44 (m, 4H), 3.31-3.25 (m, 1H), 3.11-3.04 (m, 2H), 2.98-2.94 (m, 1H), 2.74-2.68 (m, 2H), 2.49-2.46 (m, 1H), 2.41-2.36 (m, 1H), 2.27-2.18 (m, 1H), 1.77-1.70 (m, 1H), 1.06-1.01 (m, 9H).
496		462	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.51 (d, 1H, J = 1.6 Hz), 8.12 (d, 1H, J = 1.6 Hz), 7.84 (d, 1H, J = 5.2 Hz), 7.64 (dd, 1H, J = 8.0, 2.0 Hz), 7.58 (d, 1H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.80 (d, 1H, J = 5.6 Hz), 4.49 (t, 1H, J = 7.2 Hz), 3.64-3.60 (m, 4H), 3.54-3.50 (m, 4H), 3.41-3.36 (m, 1H), 3.36-3.28 (m, 2H), 3.23-3.18 (m, 1H), 2.97-2.94 (m, 1H), 2.77-2.75 (m, 1H), 2.55-2.21 (m, 1H), 2.51-2.49 (m, 1H), 2.41-2.36 (m, 1H), 1.91-1.88 (m, 1H), 1.18 (t, 3H, J = 7.2 Hz), 1.16-1.12 (m, 6H).
497		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.90 (d, 1H, J = 1.6 Hz), 8.25 (d, 1H, J = 1.6 Hz), 8.13 (dd, 1H, J1 = 8.4 Hz, J2 = 2.4 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.52 (d, 1H, J = 8.4 Hz), 7.10 (d, 1H, J = 1.2 Hz), 6.62 (t, 1H, J = 5.2 Hz), 6.00 (d, 1H, J = 5.2 Hz), 3.83 (m, 1H), 3.54-3.52 (m, 4H), 3.45-3.43 (m, 4H), 3.12-3.05 (m, 3H), 2.75-2.69 (m, 1H), 2.62-2.58 (m, 1H), 2.20-2.11 (m, 1H), 1.83-1.73 (m, 2H), 1.68-1.61 (m, 1H), 1.04 (t, 3H, J = 7.2 Hz), 0.94 (d, 6H, J = 6.0 Hz).






#	Structure	LCMS (M+1)	NMR
498		464	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.53 (s, 1H), 7.99 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.82 – 7.73 (m, 2H), 7.53 – 7.37 (m, 2H), 6.91 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.18 (q, J = 7.1 Hz, 2H), 3.72 (t, J = 4.9 Hz, 4H), 3.50 (dd, J = 6.4, 3.8 Hz, 4H), 3.43 – 3.24 (m, 6H), 2.35 (dd, J = 15.2, 2.6 Hz, 2H), 2.14 (ddd, J = 14.5, 11.0, 6.5 Hz, 2H), 1.29 (t, J = 7.1 Hz, 3H).
499		465	
500		470	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.45 (d, 1H, J = 2.0 Hz), 8.26 (d, 1H, J = 1.0 Hz), 7.93 (d, 1H, J = 5.5 Hz), 7.86 (d, 1H, J = 8.0 Hz), 7.67 (dd, 1H, J = 8.0, 2.5 Hz), 7.15 (d, 1H, J = 1.5 Hz), 5.98 (d, 1H, J = 6.0 Hz), 3.74-3.69 (m, 2H), 3.69-3.62 (m, 1H), 3.60-3.53 (m, 2H), 3.52-3.44 (m, 4H), 3.30-3.20 (m, 4H), 3.04-2.92 (m, 2H), 2.70-2.56 (m, 4H), 1.94-1.86 (m, 1H), 1.72-1.67 (m, 1H), 1.67-1.61 (m, 1H), 1.56-1.44 (m, 1H).
501		472	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.52 (s, 1H), 8.01 (d, J = 1.8 Hz, 1H), 7.87 (d, J = 5.5 Hz, 1H), 7.85 – 7.77 (m, 2H), 7.66 – 7.54 (m, 2H), 6.97 (d, J = 1.9 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 4.08 – 3.94 (m, 3H), 3.85 (s, 2H), 3.59 (d, J = 29.0 Hz, 4H), 3.54 – 3.36 (m, 3H), 2.53 (d, J = 4.0 Hz, 2H), 2.03 (tt, J = 7.9, 4.7 Hz, 1H), 1.99 – 1.85 (m, 2H), 1.77 (q, J = 10.4 Hz, 1H), 0.93 (dt, J = 5.3, 2.8 Hz, 2H), 0.87 (ddt, J = 7.6, 4.7, 2.4 Hz, 2H).
502		472	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.47 (s, 1H), 8.02 (d, J = 1.8 Hz, 1H), 7.92 – 7.77 (m, 3H), 7.66 – 7.44 (m, 2H), 6.94 (d, J = 1.9 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 3.93 (dt, J = 13.1, 6.6 Hz, 1H), 3.65 (dd, J = 6.8, 3.5 Hz, 4H), 3.59 – 3.48 (m, 7H), 2.43 (d, J = 14.3 Hz, 2H), 2.30 (td, J = 13.7, 3.9 Hz, 2H), 1.18 (d, J = 6.6 Hz, 6H).






#	Structure	LCMS (M+1)	NMR
503		472	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (s, 1H), 7.90 (d, 1H, J = 6.0 Hz), 7.70-7.55 (m, 2H), 7.35 (d, 1H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.80 (m, 2H), 3.80-3.60 (m, 2H), 3.60-3.40 (m, 5H), 3.32-2.54 (m, 5H), 2.35 (s, 3H), 2.30-2.19 (m, 1H), 2.06-1.94 (m, 1H), 1.85-1.65 (m, 1H), 1.21-0.95 (m, 6H), 0.83-0.65 (m, 4H).
504		472	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.70-7.55 (m, 2H), 7.35 (d, 1H, J = 8.8 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.80 (m, 2H), 3.80-3.60 (m, 2H), 3.60-3.40 (m, 5H), 3.32-2.54 (m, 5H), 2.35 (s, 3H), 2.30-2.19 (m, 1H), 2.06-1.94 (m, 1H), 1.85-1.70 (m, 1H), 1.21-0.95 (m, 6H), 0.83-0.65 (m, 4H).
505		472	¹ H-NMR (400 MHz, MeOD) δ ppm 7.98 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.37 (d, 2H, J = 8.0 Hz), 6.94 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.2 Hz), 4.08-4.02 (m, 2H), 3.89-3.81 (m, 2H), 3.79-3.38 (m, 2H), 3.67-3.52 (m, 6H), 3.31-3.20 (m, 2H), 2.64-2.56 (m, 1H), 2.08-2.00 (m, 1H).
506		472	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (s, 1H), 7.88 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.4 Hz), 7.01 (s, 1H), 5.96 (s, 1H), 3.96-3.87 (m, 2H), 3.74-3.66 (m, 2H), 3.58-3.42 (m, 4H), 3.31-3.20 (m, 2H), 3.18-3.10 (m, 1H), 3.06-2.96 (m, 1H), 2.92-2.82 (m, 1H), 2.08-1.92 (m, 2H), 1.82-1.76 (m, 1H), 1.08-1.02 (m, 6H), 0.95 (d, 3H, J = 6.8 Hz), 0.79-0.72 (m, 4H).
507		472	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.32 (d, 2H, J = 8.4 Hz), 7.02 (d, 1H, J = 5.2 Hz), 5.97 (d, 1H, J = 5.2 Hz), 3.92-3.91 (m, 2H), 3.70-3.60 (m, 2H), 3.53-3.52 (m, 2H), 3.47-3.46 (m, 2H), 3.16-3.15 (m, 3H), 2.81-2.80 (m, 1H), 2.49-2.34 (m, 1H), 2.02-2.01 (m, 1H), 1.46-1.45 (m, 1H), 1.22-1.21 (m, 1H), 1.12-1.11 (m, 6H), 0.96(s, 3H), 0.78-0.73 (m, 4H).






#	Structure	LCMS (M+1)	NMR
508		472	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (s, 1H), 7.90 (d, 1H, J = 5.2 Hz), 7.79-7.78 (m, 2H), 7.35-7.34 (m, 2H), 7.05-7.04 (m, 1H), 5.97 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.70-3.69 (m, 4H), 3.54-3.47 (m, 6H), 2.02-2.01 (m, 2H), 1.34-1.22 (m, 6H), 1.08-1.03 (m, 5H), 0.78-0.74 (m, 4H).
509		472	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.2 Hz, 2H), 7.50 (d, J = 8.4 Hz, 2H), 7.02 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.99 - 3.85 (m, 2H), 3.70 (s, 2H), 3.54 (s, 2H), 3.47 (s, 2H), 3.14 - 3.04 (m, 1H), 2.82 - 2.67 (m, 2H), 2.06 - 1.96 (m, 1H), 1.77 (d, J = 4.7 Hz, 4H), 1.38 (s, 3H), 0.99 (d, J = 6.3 Hz, 3H), 0.82 (d, J = 6.5 Hz, 3H), 0.79 - 0.70 (m, 4H).
510		472	
511		473	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.21 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.83 (d, 2H, J = 8.0 Hz), 7.43 (d, 2H, J = 8.0 Hz), 7.08 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.04-3.90 (m, 2H), 3.80-3.60 (m, 2H), 3.60-3.40 (m, 5H), 3.05-2.85 (m, 3H), 2.70-2.57 (m, 1H), 2.15-1.65 (m, 5H), 1.60-1.50 (m, 1H), 1.50-1.40 (m, 1H), 1.40-1.30 (m, 1H).
512		473	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.23 (s, 1H), 7.92 (d, 1H, J = 5.2 Hz), 7.86 (d, 2H, J = 8.0 Hz), 7.45 (d, 2H, J = 8.4 Hz), 7.10 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 4.10-3.87 (m, 2H), 3.80-3.65 (m, 2H), 3.65-3.38 (m, 5H), 3.20-2.60 (m, 5H), 2.35-1.70 (m, 4H), 1.70-1.55 (m, 1H), 1.55-1.40 (m, 1H), 1.40-1.30 (m, 1H).
513		473	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.22 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.84 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.09 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 6.0 Hz), 4.04-3.90 (m, 2H), 3.80-3.65 (m, 2H), 3.65-3.40 (m, 5H), 3.25-2.80 (m, 4H), 2.75-2.60 (m, 1H), 2.30-1.90 (m, 3H), 1.90-1.50 (m, 2H), 1.50-1.40 (m, 1H), 1.40-1.30 (m, 1H).





#	Structure	LCMS (M+1)	NMR
514		473	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.22 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.84 (d, 2H, J = 8.0 Hz), 7.44 (d, 2H, J = 8.4 Hz), 7.09 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.04-3.90 (m, 2H), 3.80-3.65 (m, 2H), 3.65-3.40 (m, 5H), 3.25-2.80 (m, 4H), 2.75-2.60 (m, 1H), 2.30-1.90 (m, 3H), 1.90-1.50 (m, 2H), 1.50-1.40 (m, 1H), 1.40-1.30 (m, 1H).
515		473	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (d, J = 1.9 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 8.2 Hz, 2H), 7.37 (d, J = 7.9 Hz, 2H), 6.94 (d, J = 1.9 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 5.42 (t, J = 9.0 Hz, 1H), 4.04 (s, 2H), 3.85 (s, 2H), 3.75 (dt, J = 18.2, 8.7 Hz, 2H), 3.66 (ddd, J = 11.0, 6.6, 4.2 Hz, 3H), 3.54 (s, 2H), 3.46 (ddd, J = 8.9, 6.5, 4.4 Hz, 1H), 3.23 (td, J = 8.7, 6.6 Hz, 1H), 3.06 (t, J = 9.0 Hz, 1H), 2.07 – 1.98 (m, 1H), 0.93 (dt, J = 5.3, 2.8 Hz, 2H), 0.87 (dt, J = 8.1, 2.9 Hz, 2H).
516		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.90 (d, 1H, J = 5.2 Hz), 7.76 (d, 2H, J = 7.6 Hz), 7.41 (d, 2H, J = 7.6 Hz), 7.04 (s, 1H), 5.98 (d, 1H, J = 5.2 Hz), 3.94-3.92 (m, 2H), 3.71-3.69 (m, 3H), 3.56-3.50 (m, 2H), 3.50-3.44 (m, 2H), 2.98-2.94 (m, 1H), 2.78-2.72 (m, 1H), 2.71-2.62 (m, 3H), 2.23-2.21 (m, 1H), 2.04-1.98 (m, 2H), 0.97 (d, 6H, J = 6.0 Hz), 0.78-0.74 (m, 4H).
517		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 7.6 Hz), 7.42 (d, 2H, J = 7.6 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.94-3.92 (m, 2H), 3.71-3.69 (m, 3H), 3.56-3.40 (m, 5H), 3.01-2.98 (m, 1H), 2.84-2.64 (m, 4H), 2.28-2.22 (m, 1H), 2.10-1.98 (m, 2H), 0.98 (d, 6H, J = 6.0 Hz), 0.80-0.72 (m, 4H).
518		473	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 7.6 Hz), 7.42 (d, 2H, J = 7.6 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.94-3.92 (m, 2H), 3.71-3.69 (m, 3H), 3.56-3.40 (m, 5H), 3.01-2.98 (m, 1H), 2.84-2.64 (m, 4H), 2.28-2.22 (m, 1H), 2.10-1.98 (m, 2H), 0.98 (d, 6H, J = 6.0 Hz), 0.80-0.72 (m, 4H).

#	Structure	LCMS (M+1)	NMR
519		473	
520		474	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.94 (d, J = 1.8 Hz, 1H), 7.82 (d, J = 5.5 Hz, 1H), 7.73 – 7.60 (m, 2H), 7.45 – 7.33 (m, 2H), 6.90 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.00 (s, 2H), 3.91 (d, J = 10.7 Hz, 1H), 3.82 (s, 2H), 3.70 (td, J = 11.3, 2.4 Hz, 1H), 3.66 – 3.56 (m, 4H), 3.51 (s, 2H), 3.47 – 3.36 (m, 1H), 2.88 (p, J = 6.7 Hz, 1H), 2.83 – 2.72 (m, 1H), 2.55 (td, J = 11.7, 3.3 Hz, 1H), 1.99 (tt, J = 7.9, 4.8 Hz, 1H), 1.02 (d, J = 6.9 Hz, 3H), 0.90 (dt, J = 4.8, 2.8 Hz, 2H), 0.84 (dd, J = 7.2, 2.6 Hz, 6H)
521		474	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.21 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.84 (d, 2H, J = 8.0 Hz), 8.47 (d, 2H, J = 8.0 Hz), 7.07 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 2H), 3.75-3.70 (m, 2H), 3.58-3.52 (m, 4H), 3.50-3.45 (m, 2H), 3.23-3.21 (m, 2H), 2.95 (s, 3H), 2.35-2.31 (m, 1H), 2.06-2.01 (m, 1H), 0.90 (d, 6H, J = 8.4 Hz), 0.81-0.72 (m, 4H).
522		474	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.16 (d, J = 1.8 Hz, 1H), 7.90 (d, J = 5.4 Hz, 1H), 7.82 - 7.69 (m, 2H), 7.54 - 7.43 (m, 2H), 7.03 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 5.14 (s, 1H), 3.92 (s, 2H), 3.70 (s, 2H), 3.50 (d, J = 29.4 Hz, 4H), 2.94 - 2.71 (m, 4H), 2.18 - 1.93 (m, 3H), 1.04 (t, J = 6.4 Hz, 7H), 0.76 (tt, J = 7.9, 2.9 Hz, 4H).
523		474	

#	Structure	LCMS (M+1)	NMR
524		474	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.5 Hz, 1H), 7.79 – 7.66 (m, 2H), 7.50 – 7.41 (m, 2H), 6.93 (d, J = 1.8 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 4.02 (s, 2H), 3.84 (s, 2H), 3.62 (s, 2H), 3.53 (s, 2H), 3.20 – 3.04 (m, 4H), 3.04 – 2.89 (m, 2H), 2.17 – 2.04 (m, 2H), 2.04 – 1.85 (m, 3H), 1.14 (t, J = 7.0 Hz, 3H), 0.92 (dt, J = 4.7, 2.8 Hz, 2H), 0.85 (ddt, J = 7.5, 4.6, 2.6 Hz, 2H).
525		474	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 9.18 (s, 2H), 8.36 (d, 1H, J = 1.6 Hz), 7.94 (d, 1H, J = 5.6 Hz), 7.24 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.92-3.91 (m, 2H), 3.70-3.69 (m, 2H), 3.55-3.49 (m, 2H), 3.32-3.21 (m, 2H), 2.88-2.85 (m, 2H), 2.76-2.70 (m, 2H), 2.26-2.21 (m, 2H), 2.03-2.02 (m, 1H), 1.94-1.92 (m, 2H), 1.78-1.75 (m, 2H), 0.98 (d, 6H, J = 6.4 Hz), 0.77-0.74 (m, 4H).
526		475	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.72 – 8.64 (m, 1H), 8.26 (d, J = 1.8 Hz, 1H), 7.92 (d, J = 5.4 Hz, 1H), 7.85 (dd, J = 5.6, 1.6 Hz, 2H), 7.18 (d, J = 1.8 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 5.36 (s, 1H), 3.96 – 3.89 (m, 2H), 3.70 (s, 2H), 3.52 (d, J = 31.6 Hz, 4H), 2.92 (d, J = 9.8 Hz, 1H), 2.87 – 2.73 (m, 3H), 2.13 (dt, J = 12.9, 8.0 Hz, 1H), 2.03 (dtd, J = 12.7, 7.8, 7.3, 4.8 Hz, 2H), 1.42 – 1.35 (m, 1H), 1.11 (d, J = 15.0 Hz, 6H), 0.76 (tt, J = 7.9, 2.9 Hz, 4H).
527		475	
528		475	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.84 (s, 2H), 8.18 (d, 1H, J = 1.6 Hz), 8.17 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.06 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.93-3.90 (m, 2H), 3.78-3.72 (m, 4H), 3.72-3.69 (m, 2H), 3.55-3.45 (m, 4H), 2.77-2.72 (m, 1H), 2.57-2.50 (m, 4H), 2.04-2.00 (m, 1H), 1.01 (d, 6H, J = 6.4 Hz), 0.79-0.74 (m, 4H).






#	Structure	LCMS (M+1)	NMR
529		475	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.68 (d, 1H, J = 0.8 Hz), 8.31 (s, 1H), 8.17 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.08 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.93-3.90 (m, 2H), 3.71-3.70 (m, 2H), 3.56-3.54 (m, 6H), 3.33-3.31 (m, 2H), 2.73-2.67 (m, 1H), 2.57-2.54 (m, 4H), 2.05-1.99 (m, 1H), 1.00 (d, 6H, J = 6.4 Hz), 0.78-0.74 (m, 4H).
530		475	¹ H-NMR (400 MHz, DMSO-d6) δ ppm 8.14 (d, 1H, J = 0.8 Hz), 7.89 (d, 1H, J = 4.8 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.63-6.60 (m, 1H), 5.98 (d, 1H, J = 5.6 Hz), 3.54-3.50 (m, 4H), 3.43-3.40 (m, 4H), 3.29-3.25 (m, 1H), 3.04-2.97 (m, 3H), 2.75-2.69 (m, 2H), 2.47-2.39 (m, 2H), 2.24-2.20 (m, 1H), 1.78-1.73 (m, 1H), 1.46-1.41 (m, 2H), 1.09-1.04 (m, 6H), 0.85 (t, 3H, J = 7.6 Hz).
531		475	¹ H-NMR (400 MHz, DMSO-d6) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 6.0 Hz), 7.72 (d, 2H, J = 7.6 Hz), 7.29 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.81-3.76 (m, 1H), 3.53-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.29-3.25 (m, 1H), 3.01-2.97 (m, 1H), 2.75-2.68 (m, 2H), 2.47-2.37 (m, 2H), 2.24-2.19 (m, 1H), 1.78-1.73 (m, 1H), 1.09-1.04 (m, 6H).
532		475	¹ H-NMR (400 MHz, DMSO-d6) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 2.0 Hz), 6.63-6.60 (m, 1H), 5.98 (d, 1H, J = 5.6 Hz), 3.53-3.52 (m, 4H), 3.44-3.43 (m, 4H), 3.29-3.25 (m, 1H), 3.04-2.97 (m, 3H), 2.73-2.69 (m, 2H), 2.47-2.39 (m, 2H), 2.23-2.20 (m, 1H), 1.78-1.71 (m, 1H), 1.46-1.41 (m, 2H), 1.07-1.04 (m, 6H), 0.85 (t, 3H, J = 7.6 Hz).
533		475	¹ H-NMR (400 MHz, DMSO-d6) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.29 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.2 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.81-3.76 (m, 1H), 3.52-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.29-3.25 (m, 1H), 3.02-2.98 (m, 1H), 2.75-2.69 (m, 2H), 2.47-2.37 (m, 2H), 2.24-2.19 (m, 1H), 1.78-1.75 (m, 1H), 1.09-1.04 (m, 6H).

#	Structure	LCMS (M+1)	NMR
534		476	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.92 (d, J = 1.8 Hz, 1H), 7.81 (d, J = 5.4 Hz, 1H), 7.72 – 7.58 (m, 2H), 7.31 – 7.18 (m, 2H), 6.85 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.4 Hz, 1H), 4.17 (q, J = 7.1 Hz, 2H), 3.70 (t, J = 5.0 Hz, 4H), 3.56 – 3.37 (m, 4H), 3.03 (dt, J = 12.2, 3.1 Hz, 2H), 2.77 (hept, J = 6.6 Hz, 1H), 2.53 (tt, J = 11.9, 4.1 Hz, 1H), 2.34 (td, J = 11.8, 2.8 Hz, 2H), 1.95 – 1.84 (m, 2H), 1.83 – 1.69 (m, 2H), 1.28 (t, J = 7.1 Hz, 3H), 1.12 (d, J = 6.6 Hz, 6H).
535		476	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.73 – 7.67 (m, 2H), 7.52 – 7.43 (m, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.08 (q, J = 7.1 Hz, 2H), 3.59 (s, 4H), 3.49 – 3.40 (m, 4H), 3.09 (dd, J = 9.4, 4.2 Hz, 1H), 2.82 – 2.65 (m, 2H), 1.78 (dd, J = 7.2, 4.4 Hz, 4H), 1.38 (s, 3H), 1.21 (t, J = 7.1 Hz, 3H), 0.99 (d, J = 6.4 Hz, 3H), 0.82 (d, J = 6.5 Hz, 3H).
536		476	
537		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.35 (d, 2H, J = 8.0 Hz), 7.04 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.77 (d, 1H, J = 8.4 Hz), 3.53-3.51 (m, 4H), 3.45-3.43 (m, 4H), 3.12-3.06 (m, 2H), 3.02-2.98 (m, 1H), 2.85-2.79 (m, 1H), 2.74-2.66 (m, 3H), 2.25 (t, 1H, J = 11.2 Hz), 2.09 (t, 1H, J = 10.4 Hz), 1.04 (t, 3H, J = 7.2 Hz), 0.98 (d, 6H, J = 6.4 Hz).
538		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.49 (s, 1H), 8.25 (s, 1H), 7.92 (d, 1H, J = 5.6 Hz), 7.86 (d, 1H, J = 8.4 Hz), 7.71 (d, 1H, J = 8.4 Hz), 7.14 (s, 1H), 6.61 (t, 1H, J = 5.2 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.55-3.50 (m, 4H), 3.47-3.42 (m, 4H), 3.12-3.04 (m, 2H), 2.85-2.71 (m, 4H), 2.25-2.17 (m, 2H), 1.85-1.81 (m, 1H), 1.75-1.71 (m, 1H), 1.59-1.45 (m, 2H), 1.04 (t, 3H, J = 7.2 Hz), 1.04-0.97 (m, 6H).

#	Structure	LCMS (M+1)	NMR
539		476	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.48 (d, 1H, J = 1.6 Hz), 8.24 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.85 (d, 1H, J = 8.0 Hz), 7.71-7.68 (m, 1H), 7.14 (d, 1H, J = 1.2 Hz), 6.60 (t, 1H, J = 5.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.53-3.51 (m, 4H), 3.45-3.44 (m, 4H), 3.12-3.05 (m, 2H), 2.81-2.72 (m, 4H), 2.24-2.14 (m, 2H), 1.84-1.81 (m, 1H), 1.76-1.72 (m, 1H), 1.60-1.43 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz), 0.99-0.96 (m, 6H).
540		477	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.62 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.79 (s, 1H), 3.82-3.66 (m, 1H), 3.60-3.47 (m, 6H), 3.46-3.38 (m, 4H), 3.24-3.18 (m, 2H), 2.44-2.34 (m, 1H), 1.07 (d, 6H, J = 6.8 Hz), 0.90 (d, 6H, J = 6.0 Hz).
541		477	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.46 (s, 1H), 8.00 (d, J = 1.8 Hz, 1H), 7.86 (d, J = 5.5 Hz, 1H), 7.82 – 7.69 (m, 2H), 7.54 – 7.36 (m, 2H), 6.93 (d, J = 1.9 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 3.93 (p, J = 6.6 Hz, 1H), 3.65 (dd, J = 6.6, 3.6 Hz, 4H), 3.54 (dd, J = 6.7, 3.5 Hz, 4H), 3.42 – 3.34 (m, 4H), 3.04 (s, 3H), 2.38 (d, J = 14.5 Hz, 2H), 2.13 (dd, J = 26.5, 5.7 Hz, 1H), 1.23 – 1.12 (m, 7H).
542		477	¹ H NMR (400 MHz, 4d-MeOD) δ ppm 8.55 (m, 1H), 8.04 (dd, 1H, J = 7.6, 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.52 (d, 2H, J = 8.4 Hz), 6.95 (d, 1H, J = 2.0 Hz), 6.01 (d, 1H, J = 5.6 Hz), 3.96-3.95 (m, 1H), 3.68-3.65 (m, 4H), 3.56-3.55 (m, 4H), 3.44-3.42 (m, 2H), 3.26-3.25 (m, 1H), 3.12 (s, 3H), 3.04-3.00 (m, 2H), 2.25-2.19 (m, 3H), 1.21-1.17 (m, 6H).

#	Structure	LCMS (M+1)	NMR
543		477	¹ H NMR (400 MHz, 6d-DMSO) δ ppm 8.32 (s, 1H), 8.19 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.41 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 6.0 Hz), 6.32 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.80-3.75 (m, 1H), 3.55-3.50 (m, 4H), 3.50-3.45 (m, 4H), 3.18-3.10 (m, 1H), 2.98 (s, 3H), 2.93-2.92 (m, 1H), 2.82-2.80 (m, 1H), 2.70-2.66 (m, 1H), 2.19-2.18 (m, 1H), 2.03-1.98 (m, 1H), 1.73-1.68 (m, 1H), 1.60-1.55 (m, 1H), 1.09-1.06 (m, 6H).
544		477	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.48 (d, 1H, J = 2.0 Hz), 8.24 (d, 1H, J = 2.0 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.85 (d, 1H, J = 8.4 Hz), 7.71 (dd 1H, J = 8.4, 2.4 Hz), 7.14 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.62-3.59 (m, 4H), 3.48-3.46 (m, 4H), 2.82-2.72 (m, 4H), 2.25-2.14 (m, 2H), 1.84-1.80 (m, 1H), 1.75-1.72 (m, 1H), 1.60-1.45 (m, 2H), 1.22 (t, 3H, J = 7.2 Hz), 0.99-0.96 (m, 6H).
545		477	¹ H-NMR (500 MHz, CDCl ₃) δ ppm 8.53 (s, 1H), 8.14 (d, 1H, J = 1.5 Hz), 7.87 (d, 1H, J = 5.0 Hz), 7.63-7.59 (m, 2H), 7.00 (d, 1H, J = 1.5 Hz), 5.84 (d, 1H, J = 5.5 Hz), 4.21 (q, 2H, J = 7.2 Hz), 3.75-3.70 (m, 4H), 3.52-3.49 (m, 1H), 3.50-3.45 (m, 4H), 3.33-3.20 (m, 3H), 2.60-2.53 (m, 2H), 2.36-2.31 (m, 1H), 2.14-2.10 (m, 1H), 2.01-1.95 (m, 1H), 1.70-1.64 (m, 1H), 1.35-1.28 (m, 9H).
546		477	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.33 (d, 1H, J = 2.8 Hz), 8.06 (d, 1H, J = 1.6 Hz), 7.83 (d, 1H, J = 5.6 Hz), 7.54 (d, 1H, J = 8.4 Hz), 7.26-7.22 (m, 1H), 6.96 (d, 1H, J = 1.6 Hz), 5.80 (d, 1H, J = 5.6 Hz), 4.52-4.48 (m, 1H), 3.65-3.59 (m, 4H), 3.55-3.50 (m, 4H), 3.34 (q, 2H, J = 7.2 Hz), 3.28-3.24 (m, 4H), 2.77-2.70 (m, 5H), 1.19 (t, 3H, J = 7.2 Hz), 1.11 (d, 6H, J = 6.4 Hz).
547		477	¹ H NMR (400 MHz, 4d-MeOD) δ ppm 9.08 (s, 2H), 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.06 (d, 1H, J = 0.8 Hz), 6.02 (d, 1H, J = 5.6 Hz), 3.66-3.65 (m, 4H), 3.65-3.57 (m, 4H), 3.25 (q, 2H, J = 7.2 Hz), 3.11-3.09 (m, 2H), 2.93-2.91 (m, 1H), 2.86-2.83 (m, 1H), 2.48-2.43 (m, 2H), 2.07-2.01 (m, 4H), 1.17-1.13 (m, 9H).

#	Structure	LCMS (M+1)	NMR
548		477	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.42 (d, 1H, J = 1.6 Hz), 8.31 (s, 1H), 8.16 (d, 1H, J = 8.8 Hz), 7.96 (d, 1H, J = 6.0 Hz), 7.65 (d, 1H, J = 8.8 Hz), 7.30 (d, 1H, J = 1.2 Hz), 6.66-6.61 (m, 1H), 6.02 (d, 1H, J = 6.0 Hz), 3.56-3.51 (m, 4H), 3.50-3.45 (m, 4H), 3.10-3.05 (m, 4H), 3.00-2.94 (m, 2H), 2.58-2.50 (m, 2H), 2.00-1.90 (m, 4H), 1.10 (d, 6H, J = 6.4 Hz), 1.04 (t, 3H, J = 7.2 Hz).
549		446	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.04 (t, J = 2.1 Hz, 1H), 7.95 - 7.84 (m, 2H), 7.28 - 7.13 (m, 2H), 7.04 (d, J = 1.8 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 3.89 (br.s, 2H), 3.66 (br.s, 2H), 3.48 (br.d, J = 28.4 Hz, 4H), 2.88 (s, 3H), 2.84 - 2.72 (m, 4H), 1.98 (td, J = 7.7, 3.8 Hz, 1H), 1.89 (d, J = 13.6 Hz, 2H), 1.76 (td, J = 13.5, 12.6, 4.8 Hz, 2H), 0.79 - 0.65 (m, 4H).
550		478	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.15 (d, J = 1.8 Hz, 1H), 7.90 (d, J = 5.4 Hz, 1H), 7.79 - 7.68 (m, 2H), 7.54 - 7.41 (m, 2H), 7.00 (d, J = 1.9 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 5.14 (s, 1H), 4.08 (q, J = 7.1 Hz, 2H), 3.63 - 3.56 (m, 4H), 3.45 (t, J = 5.1 Hz, 4H), 2.90 (d, J = 9.7 Hz, 1H), 2.85 - 2.70 (m, 3H), 2.10 (dt, J = 12.9, 8.0 Hz, 1H), 1.99 (dt, J = 12.5, 5.6 Hz, 1H), 1.21 (t, J = 7.1 Hz, 3H), 1.04 (t, J = 6.4 Hz, 6H).
551		478	
552		478	¹ H NMR (400 MHz, 4d-MeOD) δ ppm 9.11 (s, 2H), 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.07 (d, 1H, J = 2.0 Hz), 6.05 (d, 1H, J = 5.2 Hz), 4.19 (q, 2H, J = 7.2 Hz), 3.74-3.73 (m, 4H), 3.55-3.52 (m, 4H), 3.28-3.27 (m, 2H), 3.14-3.06 (m, 2H), 2.76-2.75 (m, 2H), 2.22-2.10 (m, 4H), 1.31 (t, 3H, J = 7.2 Hz), 1.25 (d, 6H, J = 6.4 Hz).





#	Structure	LCMS (M+1)	NMR
553		478	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.84 (s, 2H), 8.17 (d, 1H, J = 1.6 Hz), 8.16 (s, 1H), 7.89 (d, 1H, J = 5.6 Hz), 7.03 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.78-3.72 (m, 4H), 3.53-3.48 (m, 4H), 3.46-3.40 (m, 4H), 3.08 (q, 2H, J = 7.2 Hz), 2.80-2.70 (m, 1H), 2.56-2.52 (m, 4H), 1.10-1.00 (m, 9H).
554		478	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.68 (s, 1H), 8.31 (s, 1H), 8.16 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.05 (d, 1H, J = 1.2 Hz), 6.60 (t, 1H, J = 5.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.56-3.51 (m, 8H), 3.45-3.43 (m, 4H), 3.11-3.05 (m, 2H), 2.73-2.67 (m, 1H), 2.57-2.54 (m, 4H), 1.05-1.00 (m, 9H).
555		479	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.84 (s, 2H), 8.19 (s, 1H), 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.80-3.73 (m, 4H), 3.61-3.58 (m, 4H), 3.46-3.42 (m, 4H), 2.79-2.72 (m, 1H), 2.59-2.55 (m, 4H), 1.22 (t, 3H, J = 7.2 Hz), 1.02 (d, 6H, J = 6.4 Hz).
556		479	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.91 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.81 (s, 1H), 6.79 (d, 1H, J = 1.6 Hz), 6.32 (d, 1H, J = 7.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.20-4.19 (m, 1H), 3.81-3.76 (m, 1H), 3.51-3.50 (m, 4H), 3.41-3.39 (m, 4H), 3.07-3.04 (m, 1H), 2.80-2.74 (m, 2H), 2.41-2.35 (m, 1H), 2.20-2.17 (m, 1H), 2.09-2.07 (m, 1H), 1.79-1.76 (m, 2H), 1.58-1.55 (m, 1H), 1.23-1.09 (m, 6H), 0.99-0.97 (m, 6H).
557		479	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.92 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.81 (s, 1H), 6.80 (d, 1H, J = 1.6 Hz), 6.33 (d, 1H, J = 7.2 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.20-4.18 (m, 1H), 3.81-3.75 (m, 1H), 3.52-3.50 (m, 4H), 3.41-3.39 (m, 4H), 3.06-3.04 (m, 1H), 2.81-2.74 (m, 2H), 2.40-2.35 (m, 1H), 2.18 (t, 1H, J = 11.2 Hz), 2.10-2.06 (m, 1H), 1.82-1.72 (m, 2H), 1.62-1.56 (m, 1H), 1.08 (d, 6H, J = 6.8 Hz), 0.10-0.96 (m, 6H).






#	Structure	LCMS (M+1)	NMR
558		480	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (d, 1H, J = 2.0 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 6.8 Hz), 4.53 (dd, 2H, J = 7.2, 1.2 Hz), 3.72-3.68 (m, 2H), 3.64-3.60 (m, 2H), 3.55-3.40 (m, 4H), 2.99-2.90 (m, 3H), 2.70-2.55 (m, 1H), 2.25-2.15 (m, 1H), 2.10-1.98 (m, 1H), 1.92-1.66 (m, 1H), 1.60-1.50 (m, 1H).
559		480	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (d, 1H, J = 2.0 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.43 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 6.8 Hz), 4.53 (dd, 2H, J = 7.2, 1.2 Hz), 3.72-3.68 (m, 2H), 3.64-3.60 (m, 2H), 3.55-3.40 (m, 4H), 2.99-2.90 (m, 3H), 2.70-2.55 (m, 1H), 2.25-2.15 (m, 1H), 2.10-1.98 (m, 1H), 1.92-1.66 (m, 1H), 1.60-1.47 (m, 1H).
560		486	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (s, 1H), 7.89 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.28-7.25 (m, 2H), 7.01 (s, 1H), 5.97 (d, 1H, J = 5.2 Hz), 4.00-3.82 (m, 2H), 3.80-3.60 (m, 2H), 3.58-3.40 (m, 4H), 2.80-2.60 (m, 3H), 2.47-2.44 (m, 1H), 2.38-2.24 (m, 1H), 2.20-1.96 (m, 2H), 1.86-1.76 (m, 1H), 1.75-1.66 (m, 1H), 1.62-1.38 (m, 3H), 1.32-1.16 (m, 1H), 0.90 (dd, 3H, J = 6.4, 2.8 Hz), 0.85 (t, 3H, J = 7.6 Hz), 0.80-0.70 (m, 4H).
561		486	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.15 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 4.00-3.82 (m, 2H), 3.80-3.60 (m, 2H), 3.58-3.40 (m, 4H), 3.00-2.60 (m, 3H), 2.46-2.40 (m, 1H), 2.38-2.15 (m, 2H), 2.10-1.94 (m, 1H), 1.86-1.70 (m, 2H), 1.66-1.40 (m, 3H), 1.36-1.24 (m, 1H), 1.00-0.90 (m, 3H), 0.86 (t, 3H, J = 7.2 Hz), 0.80-0.70 (m, 4H).

#	Structure	LCMS (M+1)	NMR
562		486	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.2 Hz), 4.00-3.82 (m, 2H), 3.80-3.62 (m, 2H), 3.58-3.40 (m, 4H), 2.80-2.65 (m, 3H), 2.47-2.40 (m, 1H), 2.37-2.25 (m, 1H), 2.16-2.06 (m, 1H), 2.05-1.94 (m, 1H), 1.86-1.76 (m, 1H), 1.75-1.66 (m, 1H), 1.60-1.32 (m, 3H), 1.30-1.24 (m, 1H), 0.90 (d, 3H, J = 6.8 Hz), 0.85 (t, 3H, J = 7.2 Hz), 0.80-0.70 (m, 4H).
563		486	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.94-3.91 (m, 2H), 3.71-3.70 (m, 2H), 3.54-3.46 (m, 4H), 2.84-2.81 (m, 2H), 2.46-2.30 (m, 2H), 2.22-2.19 (m, 1H), 2.03-2.01 (m, 1H), 1.75-1.20 (m, 7H), 0.95-0.71 (m, 10H).
564		486	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.2 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.2 Hz), 5.97 (d, 1H, J = 5.2 Hz), 3.94-3.91 (m, 2H), 3.71-3.70 (m, 2H), 3.54-3.46 (m, 4H), 2.84-2.81 (m, 2H), 2.46-2.30 (m, 2H), 2.03-2.01 (m, 1H), 1.75-1.20 (m, 8H), 0.95-0.71 (m, 10H).
565		486	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 7.97 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.42 (d, 2H, J = 8.4 Hz), 6.93 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.08-4.02 (m, 2H), 3.90-3.84 (m, 2H), 3.68-3.62 (m, 2H), 3.58-3.54 (m, 2H), 2.88-2.78 (m, 3H), 2.70-2.60 (m, 2H), 2.36-2.25 (m, 2H), 2.08-2.00 (m, 1H), 1.94-1.84 (m, 2H), 1.34-1.22 (m, 3H), 1.12 (d, 6H, J = 6.8 Hz), 0.97-0.84 (m, 4H).
566		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.35 (d, 2H, J = 8.0 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.94-3.92 (m, 2H), 3.72-3.70 (m, 2H), 3.56-3.46 (m, 4H), 2.97-2.93 (m, 2H), 2.77-2.73 (m, 1H), 2.65-2.61 (m, 2H), 2.36-2.32 (m, 1H), 2.24-2.20 (m, 1H), 2.10 (t, 1H, J = 10.4 Hz), 2.04-2.00 (m, 1H), 1.96 (s, 3H), 0.96 (d, 6H, J = 6.8 Hz), 0.80-0.74 (m, 4H).






#	Structure	LCMS (M+1)	NMR
567		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.35 (d, 2H, J = 8.0 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.94-3.92 (m, 2H), 3.72-3.70 (m, 2H), 3.56-3.46 (m, 4H), 2.97-2.94 (m, 1H), 2.90-2.86 (m, 1H), 2.77-2.73 (m, 1H), 2.65-2.61 (m, 2H), 2.36-2.32 (m, 1H), 2.24-2.20 (m, 1H), 2.10 (t, 1H, J = 10.4 Hz), 2.04-2.00 (m, 1H), 1.96 (s, 3H), 0.96 (d, 6H, J = 6.8 Hz), 0.80-0.74 (m, 4H).
568		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.34 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.94-3.92 (m, 2H), 3.72-3.70 (m, 2H), 3.50-3.45 (m, 2H), 3.37-3.32 (m, 2H), 2.97-2.92 (m, 1H), 2.91-2.86 (m, 1H), 2.77-2.71 (m, 1H), 2.65-2.61 (m, 2H), 2.36-2.32 (m, 1H), 2.24-2.20 (m, 1H), 2.13-2.08 (m, 1H), 2.04-2.00 (m, 1H), 1.96 (s, 3H), 0.96 (d, 6H, J = 6.8 Hz), 0.80-0.74 (m, 4H).
569		487	¹ H NMR (400 MHz, CDCl ₃) δ ppm 7.92 (d, 1H, J = 1.6 Hz), 7.83 (d, 1H, J = 5.2 Hz), 7.63 (d, 2H, J = 8.4 Hz), 7.54 (d, 2H, J = 8.4 Hz), 6.71 (d, 1H, J = 1.6 Hz), 5.82 (d, 1H, J = 5.2 Hz), 3.94-3.90 (m, 2H), 3.90-3.85 (m, 2H), 3.57-3.51 (m, 2H), 3.48-3.43 (m, 2H), 3.21-3.15 (m, 1H), 2.95-2.87 (m, 1H), 2.85-2.77 (m, 1H), 2.74 (heptet, 1H, J = 6.4 Hz), 2.55-2.45 (m, 2H), 2.45-2.38 (m, 1H), 1.79-1.74 (m, 1H), 1.39 (s, 3H), 1.10-1.07 (m, 3H), 1.06-1.01 (m, 5H), 0.84-0.79 (m, 2H).
570		487	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 2.0 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.50 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 2.0 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.96-3.90 (m, 2H), 3.73-3.69 (m, 2H), 3.59-3.54 (m, 2H), 3.53-3.47 (m, 2H), 3.10-3.05 (m, 1H), 2.72-2.67 (m, 1H), 2.64-2.56 (m, 2H), 2.41-2.37 (m, 1H), 2.33-2.28 (m, 2H), 2.26-2.22 (m, 1H), 2.06-1.99 (m, 1H), 1.21 (s, 3H), 1.00-0.95 (m, 6H), 0.78-0.74 (m, 4H).






#	Structure	LCMS (M+1)	NMR
571		488	
572		488	
573		488	
574		488	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.47 (s, 1H), 7.98 (d, J = 1.8 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.81 – 7.63 (m, 2H), 7.41 – 7.24 (m, 2H), 6.90 (d, J = 1.9 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.87 (s, 2H), 4.26 (p, J = 7.8 Hz, 1H), 3.94 – 3.83 (m, 2H), 3.63 – 3.52 (m, 3H), 3.33 (p, J = 1.7 Hz, 4H), 3.21 (td, J = 12.6, 3.0 Hz, 2H), 3.00 – 2.84 (m, 0H), 2.67 (s, 2H), 2.23 – 2.12 (m, 2H), 2.12 – 1.97 (m, 2H), 1.42 (d, J = 6.6 Hz, 6H).
575		488	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.78 (d, 1H, J = 2.0 Hz), 8.14 (d, 1H, J = 1.2 Hz), 7.98-7.92 (m, 1H), 7.85 (d, 1H, J = 6.4 Hz), 7.64 (d, 1H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.82 (d, 1H, J = 5.2 Hz), 3.97-3.92 (m, 2H), 3.90-3.85 (m, 2H), 3.62-3.57 (m, 2H), 3.52-3.46 (m, 2H), 3.30-3.24 (m, 1H), 3.01-2.93 (m, 1H), 2.81-2.75 (m, 2H), 2.65-2.56 (m, 1H), 2.52-2.45 (m, 3H), 1.80-1.75 (m, 1H), 1.46 (s, 3H), 1.48 (s, 3H), 1.11-1.01 (m, 8H), 0.85-0.79 (m, 2H).





#	Structure	LCMS (M+1)	NMR
576		488	¹ H-NMR (500 MHz, CDCl ₃) δ ppm 8.81 (d, 1H, J = 1.5 Hz), 8.15 (d, 1H, J = 2.0 Hz), 7.99 (d, 1H, J = 7.5 Hz), 7.87 (d, 1H, J = 5.0 Hz), 7.66 (d, 1H, J = 8.5 Hz), 7.07 (s, 1H), 5.83 (d, 1H, J = 5.0 Hz), 3.98-3.91 (m, 2H), 3.90-3.86 (m, 2H), 3.76-3.80 (m, 1H), 3.68-3.60 (m, 2H), 3.52-3.46 (m, 2H), 3.28-3.25 (m, 1H), 3.08-3.02 (m, 1H), 2.90-2.80 (m, 2H), 2.70-2.62 (m, 2H), 2.62-2.55 (m, 1H), 1.81-1.77 (m, 1H), 2.24-2.21 (m, 1H), 1.54 (s, 3H), 1.14-1.08 (m, 6H), 1.06-1.04 (m, 2H), 0.90-0.82 (m, 2H).
577		489	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.88 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 6.98 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.85-3.70 (m, 1H), 3.60-3.48 (m, 4H), 3.46-3.38 (m, 4H), 2.84-2.75 (m, 2H), 2.73-2.60 (m, 2H), 2.22-2.08 (m, 2H), 1.86-1.77 (m, 1H), 1.76-1.68 (m, 1H), 1.60-1.50 (m, 1H), 1.50-1.38 (m, 1H), 1.07 (d, 6H, J = 6.4 Hz), 1.00-0.90 (m, 6H).
578		489	¹ H-NMR (500 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.5 Hz), 7.89 (d, 1H, J = 5.0 Hz), 7.72 (d, 2H, J = 8.5 Hz), 7.27 (d, 2H, J = 8.5 Hz), 6.99 (s, 1H), 6.31 (d, 2H, J = 7.5 Hz), 5.98 (d, 1H, J = 5.5 Hz), 3.79-3.78 (m, 1H), 3.52-3.51 (m, 4H), 3.44-3.43 (m, 4H), 2.81-2.79 (m, 2H), 2.74-2.70 (m, 2H), 2.19-2.14 (m, 2H), 1.83-1.81 (m, 1H), 1.74-1.71 (m, 1H), 1.56-1.54 (m, 1H), 1.46-1.44 (m, 1H), 1.09-1.08 (m, 6H), 0.98-0.96 (m, 6H).
579		489	¹ H-NMR (500 MHz, DMSO-d ₆) δ ppm 8.14 (s, 1H), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 6.99 (s, 1H), 6.63-6.60 (m, 1H), 5.98 (d, 1H, J = 5.6 Hz), 3.54-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.03-2.99 (m, 2H), 2.81-2.79 (m, 2H), 2.73-2.67 (m, 2H), 2.19-2.14 (m, 2H), 1.84-1.81 (m, 1H), 1.75-1.74 (m, 1H), 1.56-1.41 (m, 4H), 0.98-0.96 (m, 6H), 0.85 (t, 3H, J = 7.6 Hz).

#	Structure	LCMS (M+1)	NMR
580		489	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.13 (d, 1H, J = 1.2 Hz), 7.88 (d, 1H, J = 5.2 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.0 Hz), 6.98 (d, 1H, J = 1.2 Hz), 6.61 (t, 1H, J = 5.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.60-3.38 (m, 8H), 3.05-2.95 (m, 2H), 2.85-2.60 (m, 4H), 2.22-2.05 (m, 2H), 1.86-1.68 (m, 2H), 1.60-1.34 (m, 4H), 1.00-0.90 (m, 6H), 0.84 (t, 3H, J = 7.2 Hz).
581		489	
582		489	
583		490	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.16 (d, J = 1.8 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.78 (d, J = 8.1 Hz, 2H), 7.39 (d, J = 8.2 Hz, 2H), 7.03 (d, J = 1.9 Hz, 1H), 5.93 (d, J = 5.5 Hz, 1H), 3.88 (br.s, 2H), 3.66 br.(s, 2H), 3.47 (br.d, J = 27.5 Hz, 4H), 2.70 (d, J = 11.3 Hz, 3H), 2.13 - 1.80 (m, 5H), 0.98 (d, J = 6.5 Hz, 6H), 0.80 - 0.64 (m, 4H).
584		490	¹ H NMR (500 MHz, Methanol-d ₄) δ 8.01 (t, J = 2.2 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.76 (t, J = 8.1 Hz, 1H), 7.13 (dd, J = 8.0, 1.7 Hz, 1H), 7.07 (dd, J = 12.9, 1.7 Hz, 1H), 6.98 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.4 Hz, 1H), 4.04 (s, 2H), 3.86 (s, 2H), 3.65 (s, 2H), 3.55 (s, 2H), 3.09 (d, J = 11.5 Hz, 2H), 2.85 (s, 1H), 2.62 (t, J = 12.4 Hz, 1H), 2.42 (s, 2H), 2.03 (tt, J = 8.2, 4.7 Hz, 1H), 1.93 (d, J = 13.0 Hz, 2H), 1.86 - 1.72 (m, 2H), 1.15 (d, J = 6.6 Hz, 7H), 0.93 (dt, J = 5.6, 3.0 Hz, 2H), 0.87 (dt, J = 8.0, 3.1 Hz, 2H).

#	Structure	LCMS (M+1)	NMR
585		490	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.34 (s, 1H), 7.15 (d, 1H, J = 1.5 Hz), 7.90 (d, 1H, J = 5.5 Hz), 7.74 (d, 2H, J = 8.5 Hz), 7.37 (d, 2H, J = 8.5 Hz), 6.99 (d, 1H, J = 2.0 Hz), 5.98 (d, 1H, J = 5.5 Hz), 4.09 (q, 2H, J = 7.0 Hz), 3.70-3.41 (m, 8H), 2.68-2.65 (m, 1H), 2.56-2.54 (m, 2H), 2.49-2.45 (m, 2H), 2.05-2.03 (m, 2H), 1.74-1.71 (m, 2H), 1.22 (t, 3H, J = 7.0 Hz), 1.18 (s, 3H), 0.95 (d, 6H, J = 6.5 Hz).
586		490	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.26 (s, 1H), 8.14 (d, 1H, J = 1.5 Hz), 7.89 (d, 1H, J = 5.0 Hz), 7.70 (d, 2H, J = 8.0 Hz), 7.28 (d, 2H, J = 8.0 Hz), 6.91 (d, 1H, J = 1.5 Hz), 5.95 (d, 1H, J = 5.5 Hz), 4.29-4.24 (m, 1H), 4.11-4.08 (m, 2H), 3.90-3.86 (m, 3H), 3.41-3.40 (m, 1H), 3.23-3.20 (m, 1H), 3.05-3.04 (m, 1H), 2.85-2.83 (m, 2H), 2.80-2.73 (m, 2H), 2.50-2.20 (m, 2H), 1.75-1.74 (m, 1H), 1.73-1.72 (m, 1H), 1.54-1.53 (m, 1H), 1.52-1.51 (m, 1H), 1.29 (d, 3H, J = 6.5 Hz), 1.22(t, 3H, J = 7.0 Hz), 1.01-0.99 (m, 6H).
587		490	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.35 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 6.62 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.2Hz), 3.53-3.51 (m, 4H), 3.45-3.43 (m, 4H), 3.12-3.06 (m, 2H), 2.98-2.94 (m, 1H), 2.90-2.88 (m, 1H), 2.77-2.75 (m, 1H), 2.66-2.60 (m, 2H), 2.36 (t, 1H, J = 10.8 Hz), 2.22 (t, 1H, J = 9.2 Hz), 2.11 (t, 1H, J = 10.4 Hz), 1.96 (s, 3H), 1.03 (t, 3H, J = 7.6 Hz), 0.98 (d, 6H, J = 6.0 Hz).
588		490	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.47 (d, 1H, J = 2.0 Hz), 8.24 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.85 (d, 1H, J = 7.6 Hz), 7.69 (dd, 1H, J = 8.0, 2.0 Hz), 7.13 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.84-3.74 (m, 1H), 3.53-3.52 (m, 4H), 3.46-3.44 (m, 4H), 2.82-2.72 (m, 4H), 2.25-2.14 (m, 2H), 1.84-1.72 (m, 2H), 1.60-1.45 (m, 2H), 1.08 (d, 6H, J = 6.8 Hz), 0.99-0.96 (m, 6H).






#	Structure	LCMS (M+1)	NMR
589		490	¹ H-NMR (500 MHz, CDCl ₃) δ ppm 8.52 (s, 1H), 8.13 (d, 1H, J = 1.5 Hz), 7.86 (d, 1H, J = 5.5 Hz), 7.61-7.57 (m, 2H), 7.02 (d, 1H, J = 1.5 Hz), 5.81 (d, 1H, J = 5.0 Hz), 4.31 (d, 1H, J = 7.0 Hz), 4.05-3.98 (m, 1H), 3.64-3.60 (m, 4H), 3.58-3.52 (m, 4H), 3.33-3.29 (m, 1H), 3.25-3.10 (m, 2H), 2.48-2.42 (m, 2H), 2.10-2.04 (m, 2H), 2.00-1.91 (m, 2H), 1.60-1.54 (m, 1H), 1.30-1.20 (m, 6H), 1.20 (d, 6H, J = 6.5 Hz).
590		491	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.74 (d, J = 7.8 Hz, 2H), 7.25 (d, J = 8.1 Hz, 2H), 7.00 (d, J = 1.9 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 4.56 – 4.42 (m, 2H), 3.63 – 3.51 (m, 4H), 3.44 (t, J = 4.9 Hz, 4H), 3.17 (s, 3H), 1.83 (s, 3H), 1.08 (s, 9H).
591		491	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.16 (s, 1H), 7.90 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.0 Hz), 7.56 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 6.63-6.59 (m, 1H), 5.99 (d, 1H, J = 5.6 Hz), 4.56-4.43 (s, 1H), 3.55-3.38 (m, 8H), 3.15-3.00 (m, 2H), 2.90-2.52 (m, 3H), 2.49-2.20 (m, 2H), 1.91-1.70(m, 2H), 1.70-1.40 (m, 2H), 1.10-0.85 (m, 9H).
592		491	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.17 (s, 1H), 7.90 (d, 1H, J = 5.2 Hz), 7.89-7.70 (m, 2H), 7.57 (d, 2H, J = 8.4 Hz), 7.02 (s, 1H), 6.63-6.59 (m, 1H), 5.99 (d, 1H, J = 5.2 Hz), 4.63-4.38 (s, 1H), 3.55-3.38 (m, 8H), 3.15-3.00 (m, 2H), 2.90-2.52 (m, 3H), 2.49-2.20 (m, 2H), 2.00-1.40(m, 4H), 1.40-0.75 (m, 9H).
593		491	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.19 (s, 1H), 7.92 (d, 1H, J = 5.6 Hz), 7.83 (d, 2H, J = 8.0 Hz), 8.46 (d, 2H, J = 8.0 Hz), 7.04 (s, 1H), 6.32 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.82-3.76 (m, 1H), 3.55-3.50 (m, 6H), 3.48-3.43 (m, 4H), 3.22-3.18 (m, 2H), 2.95 (s, 3H), 2.36-2.31 (m, 1H), 1.09 (d, 6H, J = 6.4 Hz), 0.90 (d, 6H, J = 6.0 Hz).

#	Structure	LCMS (M+1)	NMR
594		491	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.94 (d, J = 1.8 Hz, 1H), 7.87 – 7.77 (m, 1H), 7.75 – 7.64 (m, 2H), 7.48 – 7.35 (m, 2H), 6.88 (d, J = 1.7 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 4.02 – 3.82 (m, 2H), 3.77 – 3.59 (m, 7H), 3.55 – 3.38 (m, 6H), 2.91 (p, J = 6.7 Hz, 1H), 2.80 (dt, J = 11.7, 2.0 Hz, 1H), 2.58 (td, J = 11.7, 3.3 Hz, 1H), 1.17 (d, J = 6.6 Hz, 6H), 1.04 (d, J = 6.9 Hz, 3H), 0.87 (d, J = 6.5 Hz, 3H).
595		491	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.11 (d, J = 1.7 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.70 (d, J = 8.1 Hz, 2H), 7.45 (d, J = 8.1 Hz, 2H), 6.96 (d, J = 1.8 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 5.12 (s, 1H), 3.74 (h, J = 6.7 Hz, 1H), 3.48 (dd, J = 6.8, 3.3 Hz, 4H), 3.38 (dd, J = 6.8, 3.4 Hz, 4H), 2.94 - 2.66 (m, 4H), 2.11 - 2.03 (m, 1H), 2.01 - 1.90 (m, 1H), 1.06 - 0.97 (m, 12H).
596		491	
597		491	¹ H NMR (400 MHz, CDCl ₃) δ ppm 7.91 (d, 1H, J = 1.2 Hz), 7.83 (d, 1H, J = 5.2 Hz), 7.62 (d, 2H, J = 8.4 Hz), 7.54 (d, 2H, J = 8.4 Hz), 6.68 (d, 1H, J = 1.2 Hz), 5.82 (d, 1H, J = 5.2 Hz), 4.19 (q, 2H, J = 7.2 Hz), 3.72-3.69 (m, 4H), 3.45-3.41 (m, 4H), 3.14-3.10 (m, 1H), 2.92-2.85 (m, 1H), 2.83-2.74 (m, 1H), 2.72 (heptet, 1H, J = 6.4 Hz), 2.53-2.46 (m, 2H), 2.45-2.38 (m, 1H), 1.39 (s, 3H), 1.29 (t, 3H, J = 7.2 Hz), 1.08 (d, 3H, J = 6.4 Hz), 1.03 (d, 3H, J = 6.4 Hz).
598		491	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.90 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.50 (d, 2H, J = 8.0 Hz), 7.00 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 4.90 (q, 2H, J = 7.2 Hz), 3.62-3.60 (m, 4H), 3.47-3.45 (m, 4H), 3.12-3.04 (m, 1H), 2.72-2.66 (m, 1H), 2.64-2.56 (m, 2H), 2.41-2.37 (m, 1H), 2.33-2.22 (m, 3H), 1.24-1.20 (m, 6H), 1.00-0.95 (m, 6H).






#	Structure	LCMS (M+1)	NMR
599		492	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.98 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.78 – 7.70 (m, 2H), 7.48 – 7.40 (m, 2H), 6.91 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.2, 5.1 Hz, 1H), 4.90 (ddd, J = 7.3, 6.2, 1.0 Hz, 2H), 4.65 (ddd, J = 7.6, 5.1, 0.9 Hz, 2H), 3.75 (d, J = 35.1 Hz, 5H), 3.53 (dd, J = 6.6, 3.9 Hz, 4H), 3.19 – 3.05 (m, 2H), 3.00 (s, 3H), 2.99 – 2.90 (m, 2H), 2.18 – 2.05 (m, 2H), 1.96 (ddd, J = 13.8, 12.2, 4.4 Hz, 2H).
600		492	¹ H NMR (500 MHz, 4d-MeOD) δ ppm 8.57 (s, 1H), 8.03 (d, 1H, J = 2.0 Hz), 7.88 (d, 1H, J = 5.5 Hz), 7.82 (d, 2H, J = 8.5 Hz), 7.50 (d, 2H, J = 8.5 Hz), 6.95 (d, 1H, J = 1.5 Hz), 6.04 (d, 1H, J = 5.5 Hz), 5.43 (quintet, 1H, J = 5.5 Hz), 4.93 (t, 2H, J = 7.0 Hz), 4.67 (dd, 2H, J = 8.0, 5.5 Hz), 3.83-3.82 (m, 2H), 3.73-3.72 (m, 2H), 3.56-3.54 (m, 4H), 3.47-3.46 (m, 1H), 3.44-3.43 (m, 1H), 3.11 (s, 3H), 3.08-3.06 (m, 1H), 2.99-2.98 (m, 1H), 2.41-2.40 (m, 1H), 2.20-2.19 (m, 1H), 2.16-2.12 (m, 1H), 1.94-1.93 (m, 1H).
601		492	¹ H NMR (500 MHz, 4d-MeOD) δ ppm 8.57 (s, 1H), 8.03 (d, 1H, J = 2.0 Hz), 7.88 (d, 1H, J = 5.5 Hz), 7.82 (d, 2H, J = 8.5 Hz), 7.50 (d, 2H, J = 8.5 Hz), 6.95 (d, 1H, J = 1.5 Hz), 6.04 (d, 1H, J = 5.5 Hz), 5.43 (quintet, 1H, J = 5.5 Hz), 4.93 (t, 2H, J = 7.0 Hz), 4.67 (dd, 2H, J = 8.0, 5.5 Hz), 3.83-3.82 (m, 2H), 3.73-3.72 (m, 2H), 3.56-3.54 (m, 4H), 3.47-3.46 (m, 1H), 3.44-3.43 (m, 1H), 3.11 (s, 3H), 3.08-3.06 (m, 1H), 2.99-2.98 (m, 1H), 2.41-2.40 (m, 1H), 2.20-2.19 (m, 1H), 2.16-2.12 (m, 1H), 1.94-1.93 (m, 1H).
602		492	

#	Structure	LCMS (M+1)	NMR
603		492	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.14 (d, J = 1.8 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.77 - 7.71 (m, 2H), 7.38 - 7.31 (m, 2H), 6.98 (d, J = 1.8 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 4.04 (q, J = 7.1 Hz, 2H), 3.58 - 3.51 (m, 4H), 3.45 - 3.38 (m, 4H), 2.95 - 2.86 (m, 5H), 2.76 - 2.63 (m, 2H), 2.39 (p, J = 6.3 Hz, 1H), 2.16 (ddd, J = 12.3, 7.2, 4.7 Hz, 1H), 2.04 (dt, J = 13.1, 7.7 Hz, 1H), 1.17 (t, J = 7.1 Hz, 3H), 0.99 (d, J = 6.2 Hz, 6H).
604		492	
605		492	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.76 (s, 1H), 8.25 (s, 1H), 7.95-7.91 (m, 2H), 7.88-7.85 (m, 1H), 7.15 (s, 1H), 6.61-6.59 (m, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.76 (s, 1H), 3.53-3.52 (m, 4H), 3.45-3.44 (m, 4H), 3.11-3.05 (m, 2H), 2.81-2.75 (m, 1H), 2.69-2.66 (m, 1H), 2.59-2.56 (m, 1H), 2.46-2.37 (m, 2H), 1.90-1.84 (m, 1H), 1.77-1.74 (m, 1H), 1.61-1.57 (m, 1H), 1.45-1.42 (m, 1H), 1.03 (t, 3H, J = 7.2 Hz), 1.0-0.97 (m, 6H).
606		492	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.76 (d, 1H, J = 1.6 Hz), 8.28-8.25 (m, 1H), 7.94-7.90 (m, 2H), 7.87-7.85 (m, 1H), 7.15 (d, 1H, J = 1.2 Hz), 6.61-6.59 (m, 1H), 5.98 (d, 1H, J = 5.6 Hz), 4.76 (s, 1H), 3.53-3.52 (m, 4H), 3.45-3.44 (m, 4H), 3.11-3.05 (m, 2H), 2.81-2.74 (m, 1H), 2.68-2.66 (m, 1H), 2.58-2.56 (m, 1H), 2.46-2.36 (m, 2H), 1.90-1.84 (m, 1H), 1.79-1.74 (m, 1H), 1.60-1.57 (m, 1H), 1.45-1.43 (m, 1H), 1.03 (t, 3H, J = 7.6 Hz), 1.0-0.97 (m, 6H).
607		492	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.68 (d, J = 2.1 Hz, 1H), 8.24 (d, J = 1.7 Hz, 1H), 7.91 (d, J = 5.4 Hz, 1H), 7.90 - 7.80 (m, 2H), 7.14 (d, J = 1.8 Hz, 1H), 6.30 (d, J = 7.6 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 5.36 (s, 1H), 3.78 (h, J = 6.7 Hz, 1H), 3.57 - 3.48 (m, 4H), 3.48 - 3.40 (m, 4H), 2.93 (d, J = 9.7 Hz, 1H), 2.88 - 2.72 (m, 3H), 2.14 (dt, J = 15.1, 7.9 Hz, 1H), 2.09 - 1.98 (m, 1H), 1.10 - 1.00 (m, 12H).





#	Structure	LCMS (M+1)	NMR
608		492	
609		493	1H NMR (400 MHz, Methanol-d4) δ 8.54 (dd, J = 2.2, 1.1 Hz, 1H), 8.20 (d, J = 1.9 Hz, 1H), 7.94 – 7.78 (m, 3H), 7.24 – 7.06 (m, 1H), 6.01 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.3, 5.1 Hz, 1H), 4.90 (ddd, J = 7.3, 6.3, 1.0 Hz, 2H), 4.66 (ddd, J = 7.5, 5.1, 0.9 Hz, 2H), 3.89 – 3.64 (m, 5H), 3.55 (dd, J = 6.5, 3.8 Hz, 4H), 3.30 (p, J = 1.6 Hz, 4H), 3.17 – 3.06 (m, 2H), 3.01 – 2.93 (m, 1H), 2.14 (dd, J = 14.5, 2.4 Hz, 2H), 2.04 – 1.89 (m, 2H).
610		494	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.49 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 6.8 Hz), 3.65-3.55 (m, 4H), 3.55-3.40 (m, 4H), 2.82-2.72 (m, 3H), 2.70-2.60 (m, 1H), 2.40-2.25 (m, 1H), 2.05-1.90 (m, 2H), 1.80-1.70 (m, 1H), 1.60-1.50 (m, 1H), 1.22 (t, 3H, J = 6.8 Hz), 1.05-0.92 (m, 6H).
611		494	1H-NMR (400 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.82 (d, 2H, J = 8.4 Hz), 7.49 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 6.8 Hz), 3.65-3.55 (m, 4H), 3.55-3.40 (m, 4H), 2.82-2.72 (m, 3H), 2.70-2.60 (m, 1H), 2.40-2.25 (m, 1H), 2.05-1.90 (m, 2H), 1.80-1.70 (m, 1H), 1.60-1.50 (m, 1H), 1.22 (t, 3H, J = 6.8 Hz), 1.05-0.92 (m, 6H).





#	Structure	LCMS (M+1)	NMR
612		494	¹ H NMR (500 MHz, Methanol-d ₄) δ 8.00 (t, J = 2.2 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.75 (t, J = 8.1 Hz, 1H), 7.12 (dd, J = 8.0, 1.7 Hz, 1H), 7.06 (dd, J = 12.9, 1.7 Hz, 1H), 6.94 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 4.18 (q, J = 7.1 Hz, 2H), 3.72 (d, J = 5.9 Hz, 4H), 3.51 (dd, J = 6.3, 3.9 Hz, 4H), 3.08 (d, J = 11.5 Hz, 2H), 2.91 – 2.77 (m, 1H), 2.61 (dd, J = 13.9, 10.0 Hz, 1H), 2.42 (t, J = 11.8 Hz, 2H), 1.91 (t, J = 17.0 Hz, 2H), 1.86 – 1.67 (m, 2H), 1.30 (t, J = 7.1 Hz, 2H), 1.14 (t, J = 6.6 Hz, 7H).
613		495	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.72 (s, 1H), 8.31 (s, 1H), 7.97-7.91 (m, 3H), 7.20 (s, 1H), 6.02-6.0 (m, 1H), 4.11-4.08 (m, 2H), 3.66-3.58 (m, 4H), 3.52-3.49 (m, 4H), 2.85-2.80 (m, 2H), 2.70-2.65 (m, 2H), 2.43-2.35 (m, 1H), 2.13-1.99 (m, 1H), 1.97-1.89 (m, 1H), 1.85-1.75 (m, 1H), 1.60-1.51 (m, 1H), 1.23 (t, 3H, J = 7.0 Hz), 1.02-0.99 (m, 6H).
614		495	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.73 (s, 1H), 8.31 (s, 1H), 7.96-7.90 (m, 3H), 7.20 (s, 1H), 6.0 (d, 1H, J = 5.6 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.66-3.59 (m, 4H), 3.53-3.46 (m, 4H), 2.85-2.80 (m, 2H), 2.70-2.65 (m, 2H), 2.43-2.31 (m, 1H), 2.07-1.93 (m, 2H), 1.85-1.71 (m, 1H), 1.61-1.50 (m, 1H), 1.23 (t, 3H, J = 7.2 Hz), 1.01-0.98 (m, 6H).
615		497	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.99 (d, J = 1.8 Hz, 1H), 7.85 (t, J = 5.2 Hz, 1H), 7.83 – 7.74 (m, 2H), 7.59 – 7.48 (m, 2H), 6.94 (d, J = 1.9 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.04 (s, 2H), 3.86 (s, 2H), 3.65 (s, 2H), 3.55 (s, 2H), 3.07 (d, J = 12.2 Hz, 2H), 2.86 (p, J = 6.6 Hz, 1H), 2.75 – 2.62 (m, 2H), 2.26 – 2.07 (m, 4H), 2.00 (tt, J = 7.9, 4.7 Hz, 1H), 1.16 (d, J = 6.6 Hz, 6H), 0.94 (dt, J = 4.9, 3.0 Hz, 2H), 0.92 – 0.81 (m, 2H).
616		499	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.64 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.80 (s, 1H), 3.76-3.69 (m, 2H), 3.65-3.62 (m, 1H), 3.60-3.50 (m, 4H), 3.49-3.40 (m, 4H), 3.32-3.27 (m, 1H), 3.26-3.20 (m, 2H), 2.70-2.58 (m, 2H), 2.48-2.30 (m, 3H), 0.92 (d, 6H, J = 6.0 Hz).

#	Structure	LCMS (M+1)	NMR
617		500	¹ H-NMR (500 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 7.08-7.05 (m, 1H), 7.00 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.55-3.51 (m, 4H), 3.45-3.43 (m, 4H), 3.30-3.27 (m, 2H), 2.82-2.80 (m, 2H), 2.74-2.65 (m, 4H), 2.18-2.16 (m, 2H), 1.84-1.81 (m, 1H), 1.75-1.74 (m, 1H), 1.46-1.41 (m, 2H), 0.99-0.97 (m, 6H).
618		500	¹ H-NMR (500 MHz, DMSO-d ₆) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 7.07-7.04 (m, 1H), 7.00 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.55-3.54 (m, 4H), 3.46-3.45 (m, 4H), 3.30-3.27 (m, 2H), 2.81-2.79 (m, 2H), 2.74-2.65 (m, 4H), 2.19-2.14 (m, 2H), 1.83-1.80 (m, 1H), 1.75-1.71 (m, 1H), 1.56-1.45 (m, 2H), 0.99-0.96 (m, 6H).
619		501	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.17(s, 1H), 7.92(d, 1H, J = 5.5 Hz), 7.75 (d, 2H, J = 7.0 Hz), 7.58(d, 2H, J = 7.0 Hz), 7.03(s, 1H), 5.99 (d, 1H, J = 5.0 Hz), 3.94-3.93 (m, 2H), 3.72-3.71 (m, 2H), 3.56-3.52 (m, 2H), 3.48-3.43(m, 2H), 2.65-2.64 (m, 1H), 2.63-2.60 (m, 3H), 2.39-2.37 (m, 2H), 2.07-2.03 (m, 2H), 1.92 (s, 3H), 1.39 (s, 3H), 0.94-0.91 (m, 6H), 0.79-0.76 (m, 4H).
620		501	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.95 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 6.4 Hz), 7.68-7.60 (m, 4H), 6.73 (d, 1H, J = 2.0 Hz), 5.85 (d, 1H, J = 5.2 Hz), 4.00-3.95 (m, 2H), 3.93-3.87 (m, 2H), 3.60-3.55 (m, 2H), 3.55-3.50 (m, 2H), 3.52-3.48 (m, 2H), 2.81-2.71 (m, 2H), 2.70-2.60 (m, 1H), 2.47-2.45 (m, 1H), 2.25-2.22 (m, 1H), 2.02 (s, 3H), 1.48 (s, 3H), 1.16-1.14 (m, 1H), 1.07-1.04 (m, 2H), 1.01-0.94 (m, 6H), 0.85-0.82 (m, 2H).





#	Structure	LCMS (M+1)	NMR
621		502	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 7.9 Hz, 2H), 7.24 (d, J = 8.1 Hz, 2H), 6.98 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.4 Hz, 1H), 4.87 (p, J = 7.4 Hz, 1H), 3.59 (s, 4H), 3.44 (t, J = 5.1 Hz, 4H), 2.89 (d, J = 10.8 Hz, 3H), 2.69 (d, J = 21.9 Hz, 1H), 2.26 (dtd, J = 12.2, 8.6, 7.7, 3.6 Hz, 4H), 2.01 (ddt, J = 16.1, 10.9, 5.4 Hz, 2H), 1.76 (d, J = 13.2 Hz, 3H), 1.70 – 1.50 (m, 2H), 0.99 (d, J = 6.5 Hz, 7H).
622		503	
623		503	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.04 (d, 1H, J = 1.2 Hz), 7.87 (d, 1H, J = 5.6 Hz), 7.62 (d, 2H, J = 8.4 Hz), 6.91 (d, 1H, J = 1.6 Hz), 6.45 (d, 2H, J = 8.4 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.93-4.75 (s, 1H), 3.94-3.92 (m, 2H), 3.85 (s, 4H), 3.70-3.66 (m, 2H), 3.54-3.42 (m, 4H), 3.21 (m, 4H), 2.70-2.60 (m, 1H), 2.22-2.16 (m, 1H), 1.48-1.38 (m, 1H), 1.24-1.14 (m, 1H), 0.85 (d, 6H, J = 6.0 Hz).
624		503	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 8.00-7.90 (m, 2H), 7.80-7.75 (m, 1H), 7.21 (d, 1H, J = 1.6 Hz), 5.99 (d, J = 5.6 Hz), 4.00-3.85 (m, 2H), 3.80-3.65 (m, 2H), 3.63-3.40 (m, 4H), 2.92 (s, 3H), 2.75-2.65 (m, 1H), 2.65-2.55 (m, 2H), 2.50-2.40 (m, 2H), 2.10-1.95 (m, 3H), 1.95-1.80 (m, 2H), 1.00 (d, 6H, J = 6.8 Hz), 0.83-0.70 (m, 4H).
625		503	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.67 (s, 1H), 8.29 (s, 1H), 7.94-7.90 (m, 2H), 7.86-7.84 (m, 1H), 7.20 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 3.92-3.94 (m, 2H), 3.70-3.72 (m, 2H), 3.57-3.49 (m, 4H), 2.95-2.94 (m, 1H), 2.93 (s, 3H), 2.81-2.78 (m, 1H), 2.66-2.63 (m, 1H), 2.49-2.45 (m, 2H), 2.03-2.01 (m, 1H), 1.89-1.87 (m, 1H), 1.74-1.72 (m, 2H), 1.34-1.32 (m, 1H), 1.0-1.01 (m, 6H), 0.78-0.74 (m, 4H).

#	Structure	LCMS (M+1)	NMR
626		503	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.67 (s, 1H), 8.29 (s, 1H), 7.94-7.90 (m, 2H), 7.86-7.84 (m, 1H), 7.20 (s, 1H), 5.98 (d, 1H, J = 5.2 Hz), 3.92-3.94 (m, 2H), 3.70-3.72 (m, 2H), 3.57-3.49 (m, 4H), 2.95-2.94 (m, 1H), 2.93 (s, 3H), 2.81-2.78 (m, 1H), 2.65-2.63 (m, 1H), 2.49-2.45 (m, 2H), 2.01-2.0 (m, 1H), 1.95-1.89 (m, 1H), 1.76-1.72 (m, 2H), 1.32-1.29 (m, 1H), 1.0 (d, 6H, J = 2.0 Hz), 0.78-0.74 (m, 4H).
627		503	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.94 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.70 – 7.60 (m, 2H), 7.32 – 7.21 (m, 2H), 6.88 (d, J = 1.9 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.67 – 4.56 (m, 2H), 3.68 (dd, J = 6.6, 3.6 Hz, 4H), 3.54 (dd, J = 6.6, 3.7 Hz, 5H), 3.08 (d, J = 11.3 Hz, 2H), 2.83 (p, J = 6.6 Hz, 1H), 2.57 (ddt, J = 12.0, 8.0, 4.1 Hz, 1H), 2.41 (dd, J = 13.0, 10.3 Hz, 2H), 1.95 – 1.71 (m, 5H), 1.15 (d, J = 6.6 Hz, 6H).
628		503	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.35 (s, 1H), 8.15 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 6.34 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.81-3.78 (m, 1H), 3.55-3.50 (m, 4H), 3.45-3.40 (m, 4H), 2.76-2.73 (m, 1H), 2.65-2.58 (m, 2H), 2.51-2.50 (m, 2H), 2.06-2.03 (m, 2H), 1.75-1.72 (m, 2H), 1.18 (s, 3H), 1.09 (d, 6H, J = 6.4 Hz), 0.97 (d, 6H, J = 6.4 Hz).
629		504	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.76 – 7.64 (m, 2H), 7.24 (d, J = 8.2 Hz, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 5.40 – 5.24 (m, 1H), 4.78 (t, J = 6.9 Hz, 2H), 4.52 (dd, J = 7.6, 5.1 Hz, 2H), 3.63 (d, J = 36.0 Hz, 4H), 3.47 (t, J = 5.0 Hz, 4H), 2.88 (d, J = 10.8 Hz, 2H), 2.77 – 2.62 (m, 1H), 2.21 (t, J = 11.4 Hz, 2H), 1.76 (d, J = 12.4 Hz, 2H), 1.69 – 1.53 (m, 2H), 0.99 (d, J = 6.5 Hz, 6H).





#	Structure	LCMS (M+1)	NMR
630		504	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 6.4 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.28 (d, 2H, J = 8.0 Hz), 7.00 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 6.4 Hz), 5.34-5.31 (m, 1H), 4.79 (t, 2H, J = 7.6 Hz), 4.53 (dd, 2H, J = 7.6, 5.6 Hz), 3.71-3.65 (m, 2H), 3.65-3.59 (m, 2H), 3.50-3.47 (m, 4H), 2.82-2.78 (m, 2H), 2.75-2.69 (m, 2H), 2.20-2.14 (m, 2H), 1.80-1.75 (m, 1H), 1.75-1.70 (m, 1H), 1.60-1.50 (m, 1H), 1.50-1.40 (m, 1H), 0.99-0.96 (m, 6H).
631		504	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.33 (quintet, 1H, J = 5.2 Hz), 4.78 (t, 2H, J = 7.2 Hz), 4.55-4.45 (dd, 2H, J = 7.2, 5.2 Hz), 3.67-3.62 (m, 2H), 3.59-3.53 (m, 2H), 3.50-3.60 (m, 4H), 2.85-2.75 (m, 2H), 2.74-2.63 (m, 2H), 2.20-2.05 (m, 2H), 1.86-1.77 (m, 1H), 1.76-1.66 (m, 1H), 1.60-1.36 (m, 2H), 1.00-0.90 (m, 6H).
632		504	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.76 – 7.67 (m, 2H), 7.24 (d, J = 8.1 Hz, 2H), 6.98 (d, J = 1.9 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 3.54 (d, J = 5.6 Hz, 4H), 3.43 (dd, J = 6.7, 3.7 Hz, 4H), 2.88 (d, J = 10.9 Hz, 3H), 2.78 – 2.63 (m, 1H), 2.21 (t, J = 11.3 Hz, 2H), 1.76 (d, J = 12.5 Hz, 2H), 1.70 – 1.51 (m, 2H), 1.43 (s, 9H), 0.99 (d, J = 6.6 Hz, 7H).
633		504	¹ H NMR (400 MHz, CDCl ₃) δ ppm 7.90 (d, 1H, J = 1.6 Hz), 7.80 (d, 1H, J = 5.2 Hz), 7.61 (d, 2H, J = 8.4 Hz), 7.52 (d, 2H, J = 8.4 Hz), 6.68 (d, 1H, J = 1.6 Hz), 5.78 (d, 1H, J = 5.2 Hz), 4.39-4.36 (m, 1H), 4.02-3.97 (m, 1H), 3.61-3.56 (m, 4H), 3.50-3.46 (m, 4H), 3.11-3.06 (m, 1H), 2.92-2.85 (m, 1H), 2.83-2.74 (m, 1H), 2.72 (heptet, 1H, J = 6.4 Hz), 2.53-2.45 (m, 2H), 2.45-2.38 (m, 1H), 1.37 (s, 3H), 1.17 (d, 6H, J = 6.8 Hz), 1.07 (d, 3H, J = 6.4 Hz), 1.02 (d, 3H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
634		504	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.49 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 8.0 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.83-3.74 (m, 1H), 3.53-3.51 (m, 4H), 3.44-3.42 (m, 4H), 3.13-3.04 (m, 1H), 2.73-2.66 (m, 1H), 2.63-2.55 (m, 2H), 2.42-2.37 (m, 1H), 2.33-2.28 (m, 2H), 2.26-2.22 (m, 1H), 1.21 (s, 3H), 1.07 (d, 6H, J = 6.4 Hz), 1.00-0.95 (m, 6H).
635		505	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.49 (d, 1H, J = 2.0 Hz), 8.26 (d, 1H, J = 1.6 Hz), 7.94 (d, 1H, J = 5.6 Hz), 7.86 (d, 1H, J = 8.4 Hz), 7.71 (dd, 1H, J = 8.4, 2.0 Hz), 7.16 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.35 (quintet, 1H, J = 5.6 Hz), 4.80 (t, 2H, J = 7.2 Hz), 4.54 (dd, 2H, J = 7.2, 5.6 Hz), 3.73-3.68 (m, 2H), 3.62-3.55 (m, 2H), 3.53-3.45 (m, 4H), 2.85-2.70 (m, 4H), 2.30-2.10 (m, 2H), 1.90-1.80 (m, 1H), 1.78-1.70 (m, 1H), 1.60-1.40 (m, 2H), 1.05-0.90 (m, 6H).
636		505	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.49 (d, 1H, J = 2.0 Hz), 8.26 (d, 1H, J = 1.6 Hz), 7.94 (d, 1H, J = 5.6 Hz), 7.86 (d, 1H, J = 8.4 Hz), 7.71 (dd, 1H, J = 8.4, 2.0 Hz), 7.16 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.35 (quintet, 1H, J = 5.6 Hz), 4.80 (t, 2H, J = 7.2 Hz), 4.54 (dd, 2H, J = 7.2, 5.6 Hz), 3.73-3.68 (m, 2H), 3.62-3.55 (m, 2H), 3.53-3.45 (m, 4H), 2.85-2.70 (m, 4H), 2.30-2.10 (m, 2H), 1.90-1.80 (m, 1H), 1.78-1.70 (m, 1H), 1.60-1.40 (m, 2H), 1.05-0.90 (m, 6H).
637		505	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.78 - 7.69 (m, 2H), 7.40 - 7.29 (m, 2H), 6.98 (d, J = 1.8 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.94 (d, J = 5.5 Hz, 1H), 3.74 (h, J = 6.7 Hz, 1H), 3.53 - 3.43 (m, 4H), 3.43 - 3.32 (m, 4H), 2.89 (d, J = 13.7 Hz, 5H), 2.76 - 2.61 (m, 2H), 2.45 - 2.37 (m, 1H), 2.16 (ddd, J = 12.3, 7.2, 4.7 Hz, 1H), 2.04 (dt, J = 13.1, 7.7 Hz, 1H), 1.06 - 0.97 (m, 12H).





#	Structure	LCMS (M+1)	NMR
638		505	
639		505	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (s, 1H), 7.90 (d, 1H, J = 5.5 Hz), 7.81 (d, 2H, J = 6.0 Hz), 7.48 (d, 2H, J = 8.5 Hz), 7.03 (s, 1H), 6.63-6.50 (m, 1H), 5.99 (d, 1H, J = 5.5 Hz), 3.53-3.51 (m, 4H), 3.45-3.44 (m, 4H), 3.10-3.07 (m, 3H), 2.94(s, 3H), 2.90-2.52 (m, 3H), 2.49-2.20 (m, 2H), 1.90 -1.40 (m, 4H), 1.06-1.02 (m, 6H), 1.04 (t, 3H, J = 7.0 Hz).
640		505	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22-8.17 (m, 1H), 7.91 (d, 2H, J = 5.2 Hz), 7.80-7.79 (m, 1H), 7.48 (d, 2H, J = 8.4 Hz), 7.03 (s, 1H), 6.63-6.50 (m, 1H), 5.99 (d, 1H, J = 5.2 Hz), 3.53-3.51 (m, 4H), 3.45-3.44 (m, 4H), 3.10-3.07 (m, 3H), 2.94(s, 3H), 2.90-2.52 (m, 3H), 2.49-2.20 (m, 2H), 1.90 -1.40 (m, 2H), 1.30-1.23(m, 2H) 1.06-1.02 (m, 9H).
641		505	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.16 (s, 1H), 7.92 (d, 1H, J = 5.0 Hz), 7.75 (d, 2H, J = 8.0 Hz), 7.57(d, 2H, J = 8.0 Hz), 7.00(s, 1H), 6.00 (d, 1H, J = 5.5 Hz), 4.10 (q, 2H, J = 7.0 Hz), 3.61-3.60 (m, 4H), 3.48-3.47 (m, 4H), 2.65-2.64 (m, 1H), 2.61-2.60 (m, 3H), 2.39-2.37 (m, 2H), 2.09-2.08 (m, 1H), 1.91 (s, 3H), 1.38 (s, 3H), 1.23(t, 3H, J = 7.0 Hz), 0.94-0.90 (m, 6H).
642		505	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.94 (d, 1H, J = 1.6 Hz), 7.85 (d, 1H, J = 5.6 Hz), 7.68-7.60 (m, 4H), 6.70 (d, 1H, J = 1.6 Hz), 5.85 (d, 1H, J = 5.6 Hz), 4.21 (q, 2H, J = 7.2 Hz), 3.80-3.70 (m, 4H), 3.56-3.52 (m, 1H), 3.48-3.42 (m, 4H), 3.42-3.39 (m, 1H), 2.80-2.76 (m, 2H), 2.67-2.62 (m, 1H), 2.48-2.44 (m, 1H), 2.24-2.21 (m, 1H), 2.01 (s, 3H), 1.47 (s, 3H), 1.31 (t, 3H, J = 7.2 Hz), 1.05-0.95 (m, 6H).






#	Structure	LCMS (M+1)	NMR
651		511	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 9.24 (br. s., 1H), 8.07 (d, 1H, J = 2.0 Hz), 7.95 (d, 1H, J = 5.5 Hz), 7.73 (d, 1H, J = 8.5 Hz), 7.39 (d, 1H, J = 1.5 Hz), 7.27 (dd, 1H, J = 8.5, 2.0 Hz), 6.91 (d, 1H, J = 2.0 Hz), 6.02 (d, 1H, J = 5.5 Hz), 4.08 (q, 2H, J = 7.5 Hz), 3.64-3.58 (m, 4H), 3.57-3.52 (m, 1H), 3.52-3.45 (m, 6H), 3.13-3.07 (m, 2H), 2.95-2.90 (m, 1H), 2.07-2.01 (m, 2H), 1.97-1.89 (m, 2H), 1.30 (d, 6H, J = 6.5 Hz), 1.22 (t, 3H, J = 7.5 Hz).
652		512	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.93 (dd, J = 6.3, 1.8 Hz, 1H), 7.83 (dd, J = 5.4, 1.8 Hz, 1H), 7.73 – 7.61 (m, 2H), 7.33 – 7.20 (m, 2H), 6.87 (dd, J = 6.8, 1.8 Hz, 1H), 6.29 – 5.88 (m, 2H), 4.34 (td, J = 14.4, 3.7 Hz, 2H), 3.74 (s, 5H), 3.51 (t, J = 5.2 Hz, 5H), 3.22 – 2.95 (m, 2H), 2.70 – 2.50 (m, 2H), 2.01 – 1.77 (m, 3H), 1.31 (dd, J = 85.1, 6.5 Hz, 6H).
653		512	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.49 (d, 1H, J = 2.0 Hz), 8.26 (d, 1H, J = 1.5 Hz), 7.93 (d, 1H, J = 5.5 Hz), 7.86 (d, 1H, J = 8.5 Hz), 7.71 (dd, 1H, J = 8.5, 2.0 Hz), 7.16 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 5.0 Hz), 3.74-3.69 (m, 2H), 3.69-3.62 (m, 1H), 3.60-3.53 (m, 2H), 3.52-3.44 (m, 4H), 3.33-3.24 (m, 3H), 2.84-2.80 (m, 2H), 2.80-2.76 (m, 2H), 2.68-2.60 (m, 2H), 2.26-2.20 (m, 1H), 2.20-2.14 (m, 1H), 1.88-1.80 (m, 1H), 1.78-1.70 (m, 1H), 1.60-1.55 (m, 1H), 1.55-1.47 (m, 1H), 1.02-0.90 (m, 6H).
654		512	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.49 (d, 1H, J = 1.6 Hz), 8.26 (d, 1H, J = 1.6 Hz), 7.93 (d, 1H, J = 5.6 Hz), 7.86 (d, 1H, J = 8.0 Hz), 7.72 (dd, 1H, J = 8.0, 2.0 Hz), 7.16 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.74-3.69 (m, 2H), 3.69-3.62 (m, 1H), 3.60-3.53 (m, 2H), 3.52-3.44 (m, 4H), 3.33-3.24 (m, 3H), 2.90-2.70 (m, 4H), 2.68-2.56 (m, 2H), 2.40-2.10 (m, 2H), 1.90-1.85 (m, 1H), 1.82-1.75 (m, 1H), 1.65-1.40 (m, 2H), 1.10-0.90 (m, 6H).

#	Structure	LCMS (M+1)	NMR
655		513	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (s, 1H), 8.22 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.84 (d, 2H, J = 8.4 Hz), 8.47 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.71-3.68 (m, 2H), 3.67-3.63 (m, 3H), 3.56-3.52 (m, 2H), 3.48-3.45 (m, 4H), 3.35-3.31 (m, 2H), 3.30-3.25 (m, 2H), 2.96 (s, 3H), 2.65-2.59 (m, 2H), 2.53-2.45 (m, 2H), 0.93 (d, 6H, J = 6.4 Hz).
656		516	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.78 (d, 2H, J = 8.0 Hz), 7.51 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.94-3.93 (m, 2H), 3.72-3.71 (m, 2H), 3.55-3.48 (m, 4H), 3.19-3.16 (m, 1H), 3.05-3.00 (m, 1H), 2.88-2.80 (m, 2H), 2.69-2.64 (m, 1H), 2.43-2.33 (m, 2H), 2.05-2.00 (m, 1H), 1.82-1.69 (m, 3H), 1.41-1.29 (m, 1H), 1.05-1.02 (m, 9H), 0.79-0.74 (m, 4H).
657		516	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.78 (d, 2H, J = 8.4 Hz), 7.51 (d, 2H, J = 8.8 Hz), 7.06 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.94-3.92 (m, 2H), 3.71-3.70 (m, 2H), 3.54-3.48 (m, 4H), 3.32-3.31 (m, 1H), 3.23-3.19 (m, 1H), 3.03-2.99 (m, 1H), 2.87-2.77 (m, 2H), 2.68-2.65 (m, 1H), 2.42-2.41 (m, 1H), 2.05-2.01 (m, 1H), 1.81-1.73 (m, 3H), 1.37-1.35 (m, 1H), 1.05-0.99 (m, 9H), 0.79-0.73 (m, 4H).
658		516	¹ H NMR (400 MHz, 4d-MeOD) δ ppm 8.00 (d, 1H, J = 2.0 Hz), 7.86 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.47 (d, 2H, J = 8.0 Hz), 6.95 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 4.05-4.04 (m, 2H), 3.87-3.86 (m, 2H), 3.64-3.63 (m, 2H), 3.56-3.55 (m, 2H), 3.15 (q, 2H, J = 6.8 Hz), 3.01-2.85 (m, 5H), 2.24-2.20 (m, 2H), 2.11-2.03 (m, 3H), 1.22 (d, 6H, J = 6.4 Hz), 1.15 (t, 3H, J = 6.8 Hz), 0.95-0.93 (m, 2H), 0.89-0.86 (m, 2H).

#	Structure	LCMS (M+1)	NMR
659		516	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.23 (br .s, 1H), 8.18 (d, 1H, J = 1.5 Hz), 7.90 (d, 1H, J = 5.0 Hz), 7.78 (d, 2H, J = 8.5 Hz), 7.40 (d, 2H, J = 8.5 Hz), 6.98 (d, 1H, J = 1.0 Hz), 5.96 (d, 1H, J = 5.5 Hz), 4.63-4.62 (m, 1H), 4.20-4.19 (m, 1H), 3.93-3.92 (m, 2H), 2.89 (s, 3H), 2.75-2.74 (m, 1H), 2.63-2.62 (m, 2H), 2.56-2.55 (m, 2H), 2.01-1.98 (m, 4H), 1.90-1.87 (m, 2H), 1.41-1.40 (m, 2H), 1.25-1.21 (m, 3H), 1.04-1.01 (m, 6H), 0.78-0.76 (m, 4H).
660		517	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.6 Hz), 8.31 (d, 1H, J = 1.6 Hz), 7.95 (s, 1H), 7.93 (d, 1H, J = 1.6 Hz), 7.80 (dd, 1H, J = 8.4, 2.4 Hz), 7.22 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.56 (t, 2H, J = 6.4 Hz), 4.45 (t, 2H, J = 6.4 Hz), 4.00-3.80 (m, 2H), 3.70-3.55 (m, 2H), 3.60-3.55 (m, 2H), 3.54-3.48 (m, 2H), 3.47-3.40 (m, 1H), 2.93 (s, 3H), 2.60-2.52 (m, 2H), 2.22-2.10 (m, 2H), 2.08-1.88 (m, 5H), 0.82-0.70 (m, 4H).
661		517	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (s, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.74 (d, J = 8.1 Hz, 2H), 7.24 (d, J = 8.0 Hz, 2H), 6.99 (s, 1H), 6.77 (s, 1H), 5.97 (d, J = 5.4 Hz, 1H), 4.59 (dd, J = 7.8, 6.0 Hz, 2H), 4.29 (t, J = 6.0 Hz, 2H), 3.51 (s, 4H), 3.44 (s, 4H), 3.25 – 2.94 (m, 2H), 1.95 – 1.63 (m, 4H), 1.11 (s, 6H).
662		517	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.5 Hz), 8.30 (d, 1H, J = 1.5 Hz), 7.94 (d, 1H, J = 5.5 Hz), 7.92 (s, 1H), 7.79 (dd, 1H, J = 8.5, 2.5 Hz), 7.21 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 3.96-3.94 (m, 2H), 3.72-3.71 (m, 2H), 3.60-3.55 (m, 2H), 3.55-3.50 (m, 2H), 3.08 (q, 2H, J = 7.0 Hz), 2.70-2.64 (m, 1H), 2.62-2.58 (m, 2H), 2.59-2.54 (m, 2H), 2.04-2.00 (m, 3H), 1.99-1.90 (m, 2H), 1.09 (t, 3H, J = 7.0 Hz), 1.01 (d, 6H, J = 5.5 Hz), 0.78-0.72 (m, 4H).





#	Structure	LCMS (M+1)	NMR
663		517	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 7.9 Hz, 2H), 7.24 (d, J = 8.1 Hz, 2H), 6.98 (d, J = 1.8 Hz, 1H), 6.20 (d, J = 8.3 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.58 – 3.47 (m, 5H), 3.42 (t, J = 5.0 Hz, 4H), 2.88 (d, J = 10.8 Hz, 2H), 2.79 – 2.62 (m, 1H), 2.21 (t, J = 11.4 Hz, 2H), 1.76 (d, J = 12.2 Hz, 2H), 1.69 – 1.51 (m, 3H), 1.00 (t, J = 6.4 Hz, 9H), 0.84 (dd, J = 6.7, 4.1 Hz, 6H).
664		517	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.77 – 7.66 (m, 2H), 7.24 (d, J = 8.1 Hz, 2H), 6.99 (d, J = 1.8 Hz, 1H), 6.49 (t, J = 6.2 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.62 – 3.47 (m, 4H), 3.43 (dd, J = 6.7, 3.6 Hz, 4H), 2.89 (dd, J = 14.5, 8.5 Hz, 4H), 2.70 (p, J = 6.6 Hz, 1H), 2.52 – 2.35 (m, 1H), 2.20 (td, J = 11.5, 2.3 Hz, 2H), 1.83 – 1.70 (m, 2H), 1.61 (qd, J = 12.2, 3.7 Hz, 2H), 0.98 (d, J = 6.6 Hz, 6H), 0.83 (s, 8H).
665		518	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.13 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.75 – 7.68 (m, 2H), 7.28 – 7.20 (m, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.65 (dd, J = 8.0, 6.1 Hz, 2H), 4.39 – 4.35 (m, 2H), 4.24 (d, J = 6.3 Hz, 2H), 3.61 (s, 4H), 3.46 (dd, J = 6.7, 3.7 Hz, 4H), 3.29 – 3.21 (m, 1H), 2.88 (d, J = 10.9 Hz, 2H), 2.74 – 2.65 (m, 1H), 2.21 (t, J = 11.3 Hz, 2H), 1.76 (d, J = 12.3 Hz, 2H), 1.69 – 1.54 (m, 2H), 0.99 (d, J = 6.5 Hz, 6H).
666		518	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.74 – 7.68 (m, 2H), 7.26 – 7.19 (m, 2H), 6.97 (d, J = 1.9 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 5.17 (ddt, J = 6.5, 4.3, 1.9 Hz, 1H), 3.82 – 3.68 (m, 4H), 3.58 (d, J = 5.5 Hz, 4H), 3.45 (dd, J = 6.6, 3.7 Hz, 4H), 2.91 (d, J = 11.1 Hz, 2H), 2.75 (p, J = 6.5 Hz, 1H), 2.30 – 2.21 (m, 2H), 2.12 (dtd, J = 13.6, 8.3, 6.3 Hz, 1H), 1.92 (dt, J = 11.9, 5.3 Hz, 1H), 1.77 (d, J = 12.1 Hz, 2H), 1.63 (qd, J = 12.2, 3.7 Hz, 2H), 1.00 (d, J = 6.5 Hz, 6H).

#	Structure	LCMS (M+1)	NMR
667		518	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.12 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.75 - 7.67 (m, 2H), 7.26 - 7.20 (m, 2H), 6.98 (d, J = 1.8 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 5.17 (ddt, J = 6.5, 4.3, 1.9 Hz, 1H), 3.83 - 3.68 (m, 4H), 3.58 (d, J = 5.5 Hz, 4H), 3.45 (d, J = 5.7 Hz, 4H), 2.96 (d, J = 11.2 Hz, 2H), 2.83 (p, J = 6.6 Hz, 1H), 2.40 - 2.27 (m, 2H), 2.12 (dt, J = 13.6, 8.3, 6.3 Hz, 1H), 1.97 - 1.88 (m, 1H), 1.84 - 1.75 (m, 2H), 1.73 - 1.60 (m, 2H), 1.03 (d, J = 6.6 Hz, 6H).
668		518	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.91 (d, 1H, J = 5.0 Hz), 7.73 (d, 2H, J = 8.5 Hz), 7.37 (d, 2H, J = 8.5 Hz), 7.00 (s, 1H), 6.00 (d, 1H, J = 5.0 Hz), 4.43-4.40 (m, 1H), 3.88-3.84 (m, 1H), 3.62-3.64 (m, 4H), 3.46-3.48 (m, 4H), 3.32-3.30 (m, 4H), 2.81-2.79 (m, 1H), 2.66-2.64 (m, 2H), 2.46-2.38 (m, 2H), 2.08-1.98 (m, 2H), 1.76-1.68 (m, 2H), 1.17 (s, 3H), 0.94 (d, 1H, J = 6.0 Hz).
669		518	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.25 (s, 1H), 8.14 (d, 1H, J = 2.0 Hz), 7.89 (d, 1H, J = 5.5 Hz), 7.71 (d, 2H, J = 8.0 Hz), 7.26 (d, 2H, J = 8.0 Hz), 6.93 (d, 1H, J = 1.5 Hz), 5.97 (d, 1H, J = 5.0 Hz), 5.36-5.33 (m, 1H), 4.79 (t, 2H, J = 7.0 Hz), 4.53-4.51 (m, 2H), 3.93-3.91 (m, 2H), 3.25-3.21 (m, 2H), 3.07-3.03 (m, 1H), 2.94-2.92 (m, 2H), 2.78-2.77 (m, 1H), 2.28-2.27 (m, 2H), 2.00-1.99 (m, 1H), 1.79-1.77 (m, 2H), 1.67-1.64 (m, 2H), 1.34-1.32 (m, 2H), 1.25-1.20 (m, 3H), 1.06-1.01 (m, 6H).
670		518	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.16 (d, J = 1.7 Hz, 1H), 7.90 (d, J = 5.4 Hz, 1H), 7.78 (d, J = 8.0 Hz, 2H), 7.25 (d, J = 8.0 Hz, 2H), 7.01 (d, J = 1.9 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 3.63 (s, 5H), 3.48 (d, J = 5.6 Hz, 6H), 2.01 (s, 6H), 1.28 (s, 8H), 0.93 (s, 10H)





#	Structure	LCMS (M+1)	NMR
671		518	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.15 (s, 1H), 7.90 (d, 1H, J = 5.5 Hz), 7.74 (d, 2H, J = 8.0 Hz), 7.57 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.0 Hz), 6.32 (d, 1H, J = 7.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 3.79-3.78 (m, 1H), 3.54-3.52 (m, 4H), 3.45-3.43 (m, 4H), 2.75-2.74 (m, 1H), 2.62-2.61 (m, 3H), 2.37-2.36 (m, 2H), 2.09-2.08 (m, 1H), 1.92 (s, 3H), 1.39 (s, 3H), 1.08 (d, 6H, J = 6.5 Hz), 0.95-0.90 (m, 6H).
672		518	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 7.94 (d, 1H, J = 1.6 Hz), 7.84 (d, 1H, J = 6.4 Hz), 7.68-7.60 (m, 4H), 6.72 (d, 1H, J = 1.6 Hz), 5.83 (d, 1H, J = 6.4 Hz), 4.28 (d, 1H, J = 7.2 Hz), 4.10-4.00 (m, 1H), 3.65-3.58 (m, 4H), 3.56-3.46 (m, 4H), 3.45-3.40 (m, 1H), 2.80-2.75 (m, 2H), 2.74-2.70 (m, 1H), 2.50-2.45 (m, 1H), 2.25-2.21 (m, 1H), 2.02 (s, 3H), 1.48 (s, 3H), 1.20 (d, 6H, J = 6.4 Hz), 1.05-0.95 (m, 6H).
673		519	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.51 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.6 Hz), 6.62 (t, 1H, J = 5.2 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.53-3.51 (m, 4H), 3.45-3.43 (m, 4H), 3.23-3.18 (m, 1H), 3.12-3.07 (m, 2H), 3.01-2.99 (m, 1H), 2.89-2.77 (m, 2H), 2.67-2.65 (m, 1H), 2.45-2.40 (m, 2H), 1.81-1.70 (m, 3H), 1.40-1.30 (m, 1H), 1.05-0.99 (m, 12H).
674		519	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.17 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.77 (d, 2H, J = 8.4 Hz), 7.51 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.2 Hz), 6.62 (s, 1H), 5.98 (d, 1H, J = 5.6 Hz), 3.53-3.52 (m, 4H), 3.45-3.43 (m, 4H), 3.34-3.33 (m, 1H), 3.23-3.21 (m, 1H), 3.12-3.00 (m, 3H), 2.87-2.77 (m, 2H), 2.68-2.65 (m, 1H), 2.41-2.40 (m, 1H), 1.81-1.73 (m, 3H), 1.37-1.35 (m, 1H), 1.06-0.99 (m, 12H).
675		519	¹ H NMR (400 MHz, 4d-MeOD) δ ppm 7.99 (d, 1H, J = 1.6 Hz), 7.86 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.48 (d, 2H, J = 8.4 Hz), 6.92 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.2 Hz), 3.67-3.65 (m, 4H), 3.56-3.54 (m, 4H), 3.24 (q, 2H, J = 7.2 Hz), 3.17 (d, 2H, J = 6.8 Hz), 3.11-3.10 (m, 2H), 3.07-3.06 (m, 2H), 2.32-2.28 (m, 2H), 2.12-2.11 (m, 2H), 1.29 (d, 6H, J = 6.4 Hz), 1.18-1.14 (m, 7H).

#	Structure	LCMS (M+1)	NMR
676		519	
677		519	
678		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.28 (d, 2H, J = 8.4 Hz), 7.0 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.80 (t, 2H, J = 7.2 Hz), 4.54 (dd, 2H, J = 7.2, 5.6 Hz), 4.54-4.49 (m, 1H), 3.69-3.65 (m, 2H), 3.65-3.60 (m, 3H), 3.51-3.49 (m, 4H), 2.89-2.75 (m, 3H), 2.65-2.59 (m, 1H), 2.33-2.24 (m, 2H), 1.94-1.92 (m, 1H), 1.53-1.50 (m, 1H), 0.98 (s, 6H).
679		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (s, 1H), 8.16 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.73 (d, 2H, J = 8.0 Hz), 7.28 (d, 2H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.80 (t, 2H, J = 6.8 Hz), 4.54 (dd, 2H, J = 7.2, 5.2 Hz), 3.79-3.70 (m, 5H), 3.57-3.50 (m, 4H), 2.90-2.78 (m, 4H), 2.69-2.60 (m, 1H), 2.41-2.32 (m, 2H), 1.95-1.92 (m, 1H), 1.61-1.49 (m, 1H), 1.01-0.99 (m, 6H).
680		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.68-7.62 (m, 2H), 7.37-7.33 (m, 1H), 7.13 (d, 1H, J = 1.6 Hz), 5.96 (d, 1H, J = 5.2 Hz), 3.93-3.91 (m, 2H), 3.67-3.66 (m, 2H), 3.56-3.47 (m, 4H), 2.96 (s, 3H), 2.67-2.57 (m, 3H), 2.49-2.45 (m, 2H), 2.13-1.95 (m, 5H), 0.98 (d, 6H, J = 6.8 Hz), 0.78-0.73 (m, 4H).

#	Structure	LCMS (M+1)	NMR
681		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.00-4.70 (m, 1H), 4.00-3.90 (m, 2H), 3.75-3.65 (m, 2H), 3.61-3.49 (m, 2H), 3.37-3.40 (m, 2H), 2.88 (s, 3H), 2.72-2.65 (m, 2H), 2.60-2.54 (m, 2H), 2.48-2.42 (m, 2H), 2.00-1.92 (m, 2H), 1.90-1.76 (m, 2H), 1.50-1.36 (m, 1H), 1.24-1.10 (m, 1H), 0.99 (d, 6H, J = 6.4 Hz).
682		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.27 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.68-7.62 (m, 2H), 7.36 (t, 1H, J = 8.4 Hz), 7.14 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.94-3.92 (m, 2H), 3.72-3.70 (m, 2H), 3.56-3.51 (m, 2H), 3.50-3.45 (m, 2H), 2.97 (s, 3H), 2.69-2.65 (m, 1H), 2.60-2.56 (m, 2H), 2.46-2.44 (m, 2H), 2.14-2.10 (m, 2H), 2.04-2.00 (m, 1H), 1.98-1.92 (m, 2H), 1.00 (d, 6H, J = 6.8 Hz), 0.78-0.74 (m, 4H).
683		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.65-7.63 (m, 2H), 7.51 (t, 1H, J = 8.4 Hz), 7.14 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.93-3.80 (m, 2H), 3.72-3.70 (m, 2H), 3.65-3.54 (m, 2H), 3.55-3.46 (m, 2H), 3.03-3.00 (m, 1H), 2.98 (s, 3H), 2.80-2.76 (m, 1H), 2.68-2.64 (m, 1H), 2.50-2.42 (m, 2H), 2.07-2.00 (m, 2H), 1.76-1.71 (m, 2H), 1.39-1.36 (m, 1H), 1.01-0.99 (m, 6H), 0.79-0.73 (m, 4H).
684		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.65-7.63 (m, 2H), 7.52 (t, 1H, J = 8.4 Hz), 7.14 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.93-3.80 (m, 2H), 3.72-3.70 (m, 2H), 3.65-3.54 (m, 2H), 3.55-3.46 (m, 2H), 3.30 (s, 3H), 2.80-2.76 (m, 1H), 2.68-2.64 (m, 1H), 2.50-2.42 (m, 3H), 2.07-2.00 (m, 2H), 1.76-1.71 (m, 2H), 1.39-1.36 (m, 1H), 1.01-0.99 (m, 6H), 0.79-0.73 (m, 4H).





#	Structure	LCMS (M+1)	NMR
685		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.29 (s, 1H), 8.19 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.80 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 4.82 (heptet, 1H, J = 6.4 Hz), 3.60-3.59 (m, 4H), 3.46-3.45 (m, 4H), 2.90 (s, 3H), 2.84 (heptet, 1H, J = 6.8 Hz), 2.74-2.71 (m, 2H), 2.64-2.58 (m, 2H), 2.04-1.97 (m, 2H), 1.96-1.91 (m, 2H), 1.22 (d, 6H, J = 6.4 Hz), 1.05 (d, 6H, J = 6.8 Hz).
686		520	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.66 (d, 1H, J = 1.6 Hz), 8.27 (d, 1H, J = 1.6 Hz), 7.93-7.90 (m, 2H), 7.86-7.84 (m, 1H), 7.16 (d, 1H, J = 1.6 Hz), 6.30 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.82-3.73 (m, 1H), 3.53-3.52 (m, 4H), 3.45-3.44 (m, 4H), 2.98-2.88 (m, 1H), 2.93 (s, 3H), 2.83-2.78 (m, 1H), 2.66-2.63 (m, 1H), 2.46-2.40 (m, 2H), 1.93-1.87 (m, 1H), 1.75-1.68 (m, 2H), 1.37-1.33 (m, 1H), 1.08 (d, 6H, J = 6.8 Hz), 1.02-0.99 (m, 6H).
687		520	¹ H-NMR (400 MHz, DMSO-d ₆) δ ppm 8.66 (d, 1H, J = 1.6 Hz), 8.27 (d, 1H, J = 1.6 Hz), 7.93-7.90 (m, 2H), 7.86-7.84 (m, 1H), 7.16 (d, 1H, J = 2.0 Hz), 6.30 (d, 1H, J = 7.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.82-3.74 (m, 1H), 3.53-3.52 (m, 4H), 3.45-3.44 (m, 4H), 2.98-2.89 (m, 1H), 2.93 (s, 3H), 2.83-2.75 (m, 1H), 2.66-2.63 (m, 1H), 2.46-2.43 (m, 2H), 1.93-1.84 (m, 1H), 1.78-1.65 (m, 2H), 1.38-1.29 (m, 1H), 1.08 (d, 6H, J = 6.4 Hz), 1.02-0.99 (m, 6H).
688		520	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.2 Hz), 7.94-7.90 (m, 2H), 7.77 (dd, 1H, J = 8.0, 2.0 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.30 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 3.82-3.75 (m, 1H), 3.55-3.50 (m, 4H), 3.47-3.42 (m, 4H), 2.92 (s, 3H), 2.92 (s, 3H), 2.77-2.70 (m, 1H), 2.60-2.50 (m, 1H), 2.49-2.40 (m, 1H), 2.10-1.95 (m, 2H), 1.95-1.80 (m, 2H), 1.08 (d, 6H, J = 6.4 Hz), 1.00 (d, 6H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
689		521	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.46 (d, 1H, J = 5.2 Hz), 8.19 (s, 1H), 7.98 (d, 1H, J = 5.2 Hz), 7.29 (t, 1H, J = 5.2 Hz), 7.21 (s, 1H), 6.02 (d, 1H, J = 5.6 Hz), 3.96-3.94 (m, 2H), 3.72-3.71 (m, 2H), 3.62-3.60 (m, 2H), 3.51-3.50 (m, 2H), 3.04 (s, 3H), 2.74-2.66 (m, 3H), 2.55-2.50 (m, 1H), 2.18-2.14 (m, 2H), 2.07-2.01 (m, 2H), 1.02 (d, 6H, J = 6.0 Hz), 0.78-0.74 (m, 4H).
690		522	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.5 Hz), 7.92 (d, 1H, J = 5.5 Hz), 7.82 (d, 2H, J = 8.5 Hz), 7.50 (d, 2H, J = 8.5 Hz), 7.05 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 5.5 Hz), 5.34 (quintet, 1H, J = 5.5 Hz), 4.79 (t, 2H, J = 7.5 Hz), 4.53 (dd, 2H, J = 7.5, 5.5 Hz), 3.78-3.74 (m, 2H), 3.62-3.55 (m, 2H), 3.55-3.40 (m, 4H), 2.83-2.75 (m, 2H), 2.75-2.70 (m, 1H), -2.68-2.58 (m, 1H), 2.38-2.27 (m, 1H), 2.08-2.00 (m, 1H), 1.98-1.90 (m, 1H), 1.87-1.75 (m, 1H), 1.61-1.51 (m, 1H), 1.03-0.90 (m, 6H).
691		522	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.20 (d, 1H, J = 1.5 Hz), 7.92 (d, 1H, J = 5.5 Hz), 7.82 (d, 2H, J = 8.5 Hz), 7.50 (d, 2H, J = 8.5 Hz), 7.05 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 5.5 Hz), 5.34 (quintet, 1H, J = 5.5 Hz), 4.79 (t, 2H, J = 7.5 Hz), 4.53 (dd, 2H, J = 7.5, 5.5 Hz), 3.78-3.74 (m, 2H), 3.62-3.55 (m, 2H), 3.55-3.40 (m, 4H), 2.83-2.75 (m, 2H), 2.75-2.70 (m, 1H), -2.68-2.58 (m, 1H), 2.38-2.27 (m, 1H), 2.08-2.00 (m, 1H), 1.98-1.90 (m, 1H), 1.87-1.75 (m, 1H), 1.61-1.51 (m, 1H), 1.03-0.90 (m, 6H).
692		522	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.5 Hz), 7.92 (d, 1H, J = 5.5 Hz), 7.66 (d, 1H, J = 11.5 Hz), 7.61 (d, 1H, J = 7.5 Hz), 7.38 (t, 1H, J = 6.5 Hz), 7.09 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 5.35 (quintet, 1H, J = 5.5 Hz), 4.80 (t, 2H, J = 7.5 Hz), 4.54 (dd, 2H, J = 7.5, 5.5 Hz), 3.70-3.60 (m, 2H), 3.60-3.55 (m, 2H), 3.50-3.48 (m, 4H), 3.10-2.98 (m, 1H), 2.82-2.81 (m, 2H), 2.21-2.19 (m, 2H), 1.79-1.76 (m, 2H), 1.56-1.55 (m, 2H), 1.00-0.99 (m, 6H).





#	Structure	LCMS (M+1)	NMR
693		522	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.5 Hz), 7.92 (d, 1H, J = 5.5 Hz), 7.66 (d, 1H, J = 11.5 Hz), 7.61 (d, 1H, J = 7.5 Hz), 7.38 (t, 1H, J = 6.5 Hz), 7.09 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 5.35 (quintet, 1H, J = 5.5 Hz), 4.80 (t, 2H, J = 7.5 Hz), 4.54 (dd, 2H, J = 7.5, 5.5 Hz), 3.70-3.60 (m, 2H), 3.60-3.55 (m, 2H), 3.50-3.48 (m, 4H), 3.10-2.98 (m, 1H), 2.82-2.81 (m, 2H), 2.21-2.19 (m, 2H), 1.79-1.76 (m, 2H), 1.56-1.55 (m, 2H), 1.00-0.99 (m, 6H).
694		552	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 8.00 (dd, J = 2.8, 1.7 Hz, 1H), 7.90 (d, J = 5.4 Hz, 1H), 7.81 (t, J = 8.4 Hz, 1H), 7.17 - 7.05 (m, 2H), 6.98 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 5.29 (tt, J = 6.3, 5.1 Hz, 1H), 4.81 - 4.68 (m, 2H), 4.48 (dd, J = 7.6, 5.1 Hz, 2H), 3.59 (d, J = 40.5 Hz, 4H), 3.45 (dd, J = 6.7, 3.7 Hz, 4H), 2.84 (d, J = 10.9 Hz, 2H), 2.72 - 2.60 (m, 1H), 2.22 - 2.10 (m, 2H), 1.74 (d, J = 12.4 Hz, 2H), 1.67 - 1.49 (m, 2H), 0.95 (d, J = 6.6 Hz, 6H).
695		523	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.73 (s, 1H), 8.31 (s, 1H), 7.96-7.92 (m, 3H), 7.21 (s, 1H), 6.06-6.01 (m, 1H), 5.37-5.32 (m, 1H), 4.81-4.78 (m, 2H), 4.55-4.53 (m, 2H), 3.74-3.69 (m, 2H), 3.68-3.62 (m, 2H), 3.52-3.49 (m, 4H), 2.90-2.85 (m, 2H), 2.69-2.64 (m, 2H), 2.44-2.36 (m, 1H), 2.12-1.98 (m, 1H), 1.96-1.93 (m, 1H), 1.80-1.75 (m, 1H), 1.61-1.53 (m, 1H), 1.01-0.98 (m, 6H).
696		523	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.72 (s, 1H), 8.31 (d, 1H, J = 1.6 Hz), 7.97-7.91 (m, 3H), 7.21 (s, 1H), 6.02 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.80 (d, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.75-3.70 (m, 2H), 3.65-3.60 (m, 2H), 3.56-3.46 (m, 4H), 2.85-2.79 (m, 2H), 2.75-2.68 (m, 2H), 2.41-2.33 (m, 1H), 2.15-2.09 (m, 1H), 2.01-1.73 (m, 2H), 1.60-1.53 (m, 1H), 1.02-0.97 (m, 6H).





#	Structure	LCMS (M+1)	NMR
697		523	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.68-7.62 (m, 2H), 7.37-7.32 (m, 1H), 7.10 (s, 1H), 6.59 (t, 1H, J = 5.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.52-3.51 (m, 4H), 3.44-3.43 (m, 4H), 3.10-3.04 (m, 2H), 2.96 (s, 3H), 2.67-2.57 (m, 3H), 2.49-2.45 (m, 2H), 2.13-2.09 (m, 2H), 2.01-1.92 (m, 2H), 1.03 (t, 3H, J = 7.2 Hz), 0.99 (d, 6H, J = 6.8 Hz).
698		523	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.65-7.63 (m, 2H), 7.52 (t, 1H, J = 8.4 Hz), 7.11 (d, 1H, J = 1.6 Hz), 6.62 (t, 1H, J = 5.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.56-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.10-3.08 (m, 2H), 3.05-3.00 (m, 1H), 2.99 (s, 3H), 2.80-2.77 (m, 1H), 2.77-2.68 (m, 1H), 2.50-2.42 (m, 2H), 2.08-2.07 (m, 1H), 1.76-1.70 (m, 2H), 1.37-1.36 (m, 1H), 1.06-1.00 (m, 9H).
699		523	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.65-7.63 (m, 2H), 7.52 (t, 1H, J = 8.4 Hz), 7.11 (d, 1H, J = 1.6 Hz), 6.61 (t, 1H, J = 5.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.56-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.10-3.08 (m, 2H), 3.05-3.00 (m, 1H), 2.99 (s, 3H), 2.80-2.77 (m, 1H), 2.77-2.68 (m, 1H), 2.50-2.42 (m, 2H), 2.08-2.07 (m, 1H), 1.76-1.70 (m, 2H), 1.37-1.36 (m, 1H), 1.06-1.00 (m, 9H).
700		524	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 9.24 (br. s., 1H), 8.07 (d, 1H, J = 1.0 Hz), 7.95 (d, 1H, J = 5.5 Hz), 7.73 (d, 1H, J = 7.5 Hz), 7.39 (s, 1H), 7.27 (d, 1H, J = 7.0 Hz), 6.91 (d, 1H, J = 1.5 Hz), 6.31 (d, 1H, J = 7.0 Hz), 6.02 (d, 1H, J = 6.0 Hz), 3.80-3.76 (m, 1H), 3.55-3.51 (m, 6H), 3.51-3.49 (m, 1H), 3.49-3.42 (m, 4H), 3.13-3.07 (m, 2H), 2.95-2.90 (m, 1H), 2.11-2.08 (m, 2H), 1.97-1.90 (m, 2H), 1.29 (d, 6H, J = 6.5 Hz), 1.07 (d, 6H, J = 6.5 Hz).

#	Structure	LCMS (M+1)	NMR
701		524	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.03 (t, J = 2.2 Hz, 1H), 7.93 - 7.84 (m, 2H), 7.25 - 7.16 (m, 2H), 7.00 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.04 (q, J = 7.1 Hz, 2H), 3.55 (d, J = 5.8 Hz, 4H), 3.43 (dd, J = 6.7, 3.7 Hz, 4H), 2.87 (s, 3H), 2.60 (dd, J = 32.7, 8.3 Hz, 3H), 1.93 (d, J = 13.0 Hz, 2H), 1.81 (t, J = 11.6 Hz, 2H), 1.17 (t, J = 7.1 Hz, 3H), 0.95 (d, J = 6.5 Hz, 6H).
702		526	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.34 (s, 1H), 8.07 (d, 1H, J = 2.0 Hz), 7.77 (d, 1H, J = 5.2 Hz), 7.73-7.68 (m, 2H), 7.04 (d, 1H, J = 1.6 Hz), 5.87 (d, 1H, J = 5.6 Hz), 3.77-3.69 (m, 4H), 3.48-3.42 (m, 4H), 3.39-3.33 (m, 1H), 3.31-3.14 (m, 1H), 2.94-2.88 (m, 1H), 2.79-2.73 (m, 1H), 2.71-2.64 (m, 1H), 2.57-2.46 (m, 3H), 2.31-2.23 (m, 1H), 2.09-2.04 (m, 2H), 1.88-1.81 (m, 1H), 1.79-1.70 (m, 2H), 1.62-1.52 (m, 2H), 1.48-1.39 (m, 2H), 1.08 (m, 6H).
703		526	¹ H-NMR (400 MHz, CDCl ₃) δ ppm 8.52 (d, 1H, J = 2.0 Hz), 8.13 (d, 1H, J = 1.6 Hz), 7.87 (d, 1H, J = 5.2 Hz), 7.64 (dd, 1H, J = 8.0, 2.0 Hz), 7.59 (d, 1H, J = 8.0 Hz), 7.01 (d, 1H, J = 1.6 Hz), 5.83 (d, 1H, J = 5.6 Hz), 3.88-3.83 (m, 2H), 3.77-3.72 (m, 2H), 3.57-3.52 (m, 2H), 3.47-3.42 (m, 2H), 3.41-3.37 (m, 1H), 3.23-3.18 (m, 1H), 2.98-2.93 (m, 1H), 2.78-2.73 (m, 1H), 2.60-2.47 (m, 4H), 2.45-2.38 (m, 1H), 2.26-2.23 (m, 2H), 1.91-1.85 (m, 3H), 1.71-1.65 (m, 2H), 1.64-1.59 (m, 2H), 1.17-1.13 (m, 6H).
704		527	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.19 (s, 1H), 8.13 (d, J = 1.7 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 8.0 Hz, 2H), 7.25 (d, J = 8.1 Hz, 2H), 6.98 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.86 (p, J = 7.3 Hz, 1H), 3.58 (d, J = 17.1 Hz, 5H), 3.46 (t, J = 5.2 Hz, 4H), 3.06 (t, J = 8.8 Hz, 1H), 2.93 (d, J = 11.2 Hz, 2H), 2.83 - 2.70 (m, 2H), 2.59 (d, J = 7.0 Hz, 1H), 2.42 - 2.22 (m, 3H), 1.78 (d, J = 12.0 Hz, 2H), 1.73 - 1.57 (m, 2H), 0.99 (d, J = 8.3 Hz, 4H).





#	Structure	LCMS (M+1)	NMR
705		527	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.19 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 4.00-3.90 (m, 2H), 3.76-3.68 (m, 2H), 3.64-3.52 (m, 2H), 3.51-3.40 (m, 2H), 3.00-2.90 (m, 1H), 2.88 (s, 3H), 2.70 (heptet, 1H, J = 6.4 Hz), 2.62-2.54 (m, 2H), 2.48-2.42 (m, 2H), 2.15-2.05 (m, 1H), 2.00-1.92 (m, 2H), 1.90-1.78 (m, 2H), 1.50-1.40 (m, 1H), 1.38-1.30 (m, 1H), 0.99 (d, 6H, J = 6.4 Hz).
706		527	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 2.0 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.78 (d, 2H, J = 8.4 Hz), 7.63 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.80 (s, 1H), 3.80-3.70 (m, 2H), 3.68-3.60 (m, 2H), 3.56-3.47 (m, 4H), 3.45-3.40 (m, 2H), 3.26-3.16 (m, 2H), 2.80-2.60 (m, 2H), 2.44-2.34 (m, 1H), 2.10-1.94 (m, 2H), 1.80-1.68 (m, 2H), 1.66-1.50 (m, 2H), 1.48-1.32 (m, 2H), 0.90 (d, 6H, J = 6.0 Hz).
707		527	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.56 (d, 2H, J = 8.4 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.51 (s, 1H), 3.71-3.64 (m, 3H), 3.56-3.54 (m, 2H), 3.46-3.45 (m, 4H), 3.31-3.24 (m, 2H), 2.81-2.76 (m, 1H), 2.67-2.59 (m, 4H), 2.46-2.43 (m, 2H), 2.34-2.30 (m, 1H), 1.84-1.80 (m, 2H), 1.56-1.51 (m, 2H), 0.99 (t, 6H, J = 5.6 Hz).
708		529	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 8.19 (s, 1H), 8.13 (d, J = 1.7 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 8.1 Hz, 2H), 7.26 (dd, J = 9.8, 7.1 Hz, 3H), 7.00 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 3.86 (qd, J = 9.8, 6.3 Hz, 2H), 3.59-3.53 (m, 4H), 3.45 (dd, J = 6.6, 3.6 Hz, 4H), 2.97 (d, J = 11.3 Hz, 2H), 2.83 (p, J = 6.6 Hz, 1H), 2.42-2.26 (m, 2H), 1.88-1.74 (m, 2H), 1.67 (qd, J = 12.3, 3.7 Hz, 2H), 1.03 (d, J = 6.6 Hz, 6H).

#	Structure	LCMS (M+1)	NMR
709		532	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21 (s, 1H), 8.13 (d, J = 1.6 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 8.0 Hz, 2H), 7.24 (d, J = 8.1 Hz, 2H), 6.99 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.79 (tt, J = 8.1, 4.0 Hz, 1H), 3.79 (dt, J = 10.4, 4.5 Hz, 2H), 3.61 (s, 4H), 3.48 (ddd, J = 11.4, 8.4, 3.4 Hz, 6H), 2.96 (d, J = 11.0 Hz, 2H), 2.83 (p, J = 6.6 Hz, 1H), 2.34 (dd, J = 12.5, 10.0 Hz, 2H), 1.88 (dd, J = 12.8, 4.6 Hz, 2H), 1.83 – 1.75 (m, 2H), 1.68 (tt, J = 12.5, 6.3 Hz, 2H), 1.56 (dtd, J = 12.6, 8.5, 4.0 Hz, 2H), 1.03 (d, J = 6.6 Hz, 6H).
710		533	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.57 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.96-7.88 (m, 2H), 7.76 (dd, 1H, J = 8.4, 2.4 Hz), 7.20 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.90 (m, 2H), 3.80-3.65 (m, 2H), 3.61-3.55 (m, 2H), 3.53-3.47 (m, 2H), 3.46-3.40 (m, 1H), 3.26-3.22 (m, 1H), 3.24 (s, 3H), 2.91 (s, 3H), 2.80-2.70 (m, 1H), 2.68-2.56 (m, 4H), 2.10-1.94 (m, 3H), 1.92-1.80 (m, 2H), 0.98 (d, 3H, J = 6.4 Hz), 0.80-0.70 (m, 4H).
711		533	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.30 (d, 1H, J = 1.6 Hz), 8.00-7.90 (m, 2H), 7.78 (dd, 1H, J = 8.4, 2.0 Hz), 7.21 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.00-3.90 (m, 2H), 3.80-3.65 (m, 2H), 3.65-3.59 (m, 2H), 3.55-3.47 (m, 1H), 3.47-3.40 (m, 1H), 3.28-3.24 (m, 1H), 3.27 (s, 3H), 2.92 (s, 3H), 2.80-2.70 (m, 1H), 2.68-2.55 (m, 4H), 2.10-1.95 (m, 3H), 1.94-1.80 (m, 2H), 0.98 (d, 3H, J = 6.4 Hz), 0.82-0.70 (m, 4H).
712		533	¹ H NMR (400 MHz, CD ₃ OD) δ ppm 7.98 (d, 1H, J = 2.0 Hz), 7.85(d, 1H, J = 5.6 Hz), 7.74 (d, 2H, J = 8.0 Hz), 7.47 (d, 2H, J = 8.0 Hz), 6.92 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.95-3.90 (m, 1H), 3.67-3.65 (m, 4H), 3.55-3.53 (m, 4H), 3.17-3.16 (q, 2H, J = 7.2 Hz), 2.95-2.94 (m, 3H), 2.93-2.90 (m, 2H), 2.17-2.16 (m, 2H), 2.09-2.05 (m, 2H), 1.21-1.18 (m, 12H), 1.17-1.13 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
713		534	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.30 (s, 1H), 8.20 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.81 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 5.36-5.33 (m, 1H), 4.79 (t, 2H, J = 6.4 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.60-3.59 (m, 4H), 3.50-3.49 (m, 4H), 2.93-2.91 (m, 1H), 2.90 (s, 3H), 2.82-2.79 (m, 2H), 2.70-2.65 (m, 2H), 2.07-1.98 (m, 4H), 1.08 (d, 6H, J = 6.4 Hz).
714		534	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.96-7.90 (m, 2H), 7.78 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.60-4.51 (m, 2H), 4.50-4.40 (m, 2H), 3.84-3.72 (m, 1H), 3.56-3.50 (m, 4H), 3.48-3.40 (m, 4H), 2.91 (s, 3H), 2.60-2.46 (m, 3H), 2.30-2.10 (m, 2H), 2.08-1.88 (m, 4H), 1.08 (d, 6H, J = 6.4 Hz).
715		534	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.5 Hz), 8.27 (d, 1H, J = 2.0 Hz), 7.93 (d, 1H, J = 5.5 Hz), 7.91 (s, 1H), 7.78 (dd, 1H, J = 8.5, 2.0 Hz), 7.17 (d, 1H, J = 1.5 Hz), 6.31 (d, 1H, J = 7.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 3.81-3.76 (m, 1H), 3.53-3.50 (m, 4H), 3.46-3.45 (m, 4H), 3.08 (q, 2H, J = 7.0 Hz), 2.70-2.62 (m, 1H), 2.60-2.54 (m, 2H), 2.54-2.50 (m, 2H), 2.01-1.98 (m, 2H), 1.91-1.87 (m, 2H), 1.09-1.04 (m, 9H), 1.01-0.99 (m, 6H).
716		534	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.69 (s, 1H), 8.27 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.90-7.88 (m, 2H), 7.16 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.80-3.75 (m, 1H), 3.53-3.52 (m, 4H), 3.45-3.44 (m, 4H), 3.26-3.22 (m, 1H), 3.03-2.98 (m, 2H), 2.82-2.78 (m, 1H), 2.62-2.60 (m, 1H), 2.45-2.41 (m, 2H), 1.94-1.85 (m, 1H), 1.74-1.66 (m, 2H), 1.36-1.26 (m, 1H), 1.07 (d, 6H, J = 6.4 Hz), 1.04-0.99 (m, 9H).





#	Structure	LCMS (M+1)	NMR
717		534	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.68 (s, 1H), 8.26 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.90-7.88 (m, 2H), 7.16 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 3.81-3.75 (m, 1H), 3.53-3.52 (m, 4H), 3.45-3.44 (m, 4H), 3.26-3.22 (m, 1H), 3.03-3.0 (m, 2H), 2.81-2.78 (m, 1H), 2.62-2.59 (m, 1H), 2.45-2.40 (m, 2H), 1.94-1.85 (m, 1H), 1.78-1.63 (m, 2H), 1.35-1.24 (m, 1H), 1.07 (d, 6H, J = 6.4 Hz), 1.04-0.99 (m, 9H).
718		535	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.68 (d, 1H, J = 1.6 Hz), 8.30 (d, 1H, J = 1.2 Hz), 8.20 (s, 1H), 7.94 (d, 1H, J = 5.6 Hz), 7.91 (s, 1H), 7.85 (dd, 1H, J = 8.0, 2.4 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 5.35-5.30 (m, 1H), 4.78 (t, 2H, J = 7.2 Hz), 4.52 (dd, 2H, J = 7.6, 5.6 Hz), 3.72-3.66 (m, 2H), 3.65-3.59 (m, 2H), 3.51-3.50 (m, 4H), 2.98-2.95 (m, 1H), 2.94 (s, 3H), 2.90-2.84 (m, 1H), 2.76-2.73 (m, 1H), 2.59-2.53 (m, 1H), 1.95-1.90 (m, 1H), 1.83-1.81 (m, 1H), 1.74-1.71 (m, 1H), 1.45-1.33 (m, 1H), 1.03 (d, 6H, J = 6.4 Hz).
719		535	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.66 (d, 1H, J = 1.6 Hz), 8.29 (d, 1H, J = 1.6 Hz), 8.21-8.17 (m, 1H), 7.94 (d, 1H, J = 5.6 Hz), 7.91 (s, 1H), 7.85 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.0 (d, 1H, J = 5.2 Hz), 5.34-5.30 (m, 1H), 4.78 (t, 2H, J = 7.2 Hz), 4.52 (dd, 2H, J = 7.6, 5.6 Hz), 3.72-3.66 (m, 2H), 3.65-3.59 (m, 2H), 3.51-3.50 (m, 4H), 2.98-2.95 (m, 1H), 2.94 (s, 3H), 2.90-2.84 (m, 1H), 2.78-2.75 (m, 1H), 2.58-2.53 (m, 1H), 1.96-1.90 (m, 1H), 1.84-1.72 (m, 2H), 1.46-1.36 (m, 1H), 1.04 (d, 6H, J = 6.4 Hz).
720		537	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.25 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.68-7.62 (m, 2H), 7.37-7.32 (m, 1H), 7.10 (s, 1H), 6.32 (d, 1H, J = 7.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 3.80-3.77 (m, 1H), 3.60-3.57 (m, 4H), 3.52-3.51 (m, 4H), 2.96 (s, 3H), 2.67-2.57 (m, 3H), 2.49-2.45 (m, 2H), 2.13-2.09 (m, 2H), 2.01-1.97 (m, 2H), 1.07 (d, 6H, J = 6.4 Hz), 0.98 (d, 6H, J = 6.8 Hz).





#	Structure	LCMS (M+1)	NMR
721		537	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.68-7.62 (m, 2H), 7.36 (t, 1H, J = 8.4 Hz), 7.11 (d, 1H, J = 1.6 Hz), 6.33 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.82-3.76 (m, 1H), 3.54-3.50 (m, 4H), 3.44-3.40 (m, 4H), 2.96 (s, 3H), 2.70-2.67 (m, 1H), 2.60-2.56 (m, 2H), 2.48-2.46 (m, 2H), 2.14-2.10 (m, 2H), 2.00-1.92 (m, 2H), 1.09 (d, 6H, J = 6.8 Hz), 1.00 (d, 6H, J = 6.8 Hz).
722		537	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (s, 1H), 7.91 (d, 1H, J = 5.2 Hz), 7.65-7.63 (m, 2H), 7.52 (t, 1H, J = 8.4 Hz), 7.12 (s, 1H), 6.33 (d, 1H, J = 7.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.81-3.76 (m, 1H), 3.56-3.52 (m, 4H), 3.46-3.44 (m, 4H), 2.99 (s, 3H), 2.80-2.70 (m, 3H), 2.52-2.49 (m, 2H), 2.08-2.07 (m, 1H), 1.76-1.70 (m, 2H), 1.38-1.36 (m, 1H), 1.09 (d, 6H, J = 6.8 Hz), 1.10-0.90 (m, 6H).
723		537	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.26 (d, 1H, J = 1.6 Hz), 8.21 (s, 1H), 7.91 (d, 1H, J = 5.2 Hz), 7.68-7.65 (m, 2H), 7.64 (s, 1H), 7.54-7.50 (m, 1H), 7.11 (s, 1H), 6.33 (d, 1H, J = 7.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.81-3.76 (m, 1H), 3.56-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.02-2.97 (m, 1H), 2.99 (s, 3H), 2.90-2.80 (m, 1H), 2.79-2.70 (m, 1H), 2.61-2.55 (m, 1H), 2.08-2.07 (m, 1H), 1.89-1.79 (m, 1H), 1.79-1.70 (m, 1H), 1.38-1.36 (m, 1H), 1.09 (d, 6H, J = 6.8 Hz), 1.10-0.90 (m, 6H).
724		537	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.88-8.85 (m, 1H), 8.33 (d, 1H, J = 1.6 Hz), 7.93 (d, 1H, J = 5.6 Hz), 7.65-7.63 (m, 2H), 7.43 (t, 1H, J = 8.4 Hz), 7.15 (d, 1H, J = 1.2 Hz), 6.45-6.41 (m, 1H), 6.00 (d, 1H, J = 5.6 Hz), 3.81-3.80 (m, 1H), 3.78-3.77 (m, 1H), 3.56-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.30-3.27 (m, 2H), 3.25-3.06 (m, 2H), 3.06 (s, 3H), 2.50-2.49 (m, 1H), 2.39-2.36 (m, 1H), 2.13-2.12 (m, 1H), 1.98-1.91 (m, 1H), 1.29-1.26 (m, 6H), 1.09 (d, 6H, J = 6.4 Hz).





#	Structure	LCMS (M+1)	NMR
725		538	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.46 (d, 1H, J = 4.8 Hz), 8.18 (s, 1H), 7.98 (d, 1H, J = 5.6 Hz), 7.29 (t, 1H, J = 5.6 Hz), 7.17 (s, 1H), 6.29 (d, 1H, J = 7.6 Hz), 6.03 (d, 1H, J = 5.6 Hz), 3.80-3.76 (m, 1H), 3.54-3.52 (m, 4H), 3.48-3.47 (m, 4H), 3.04 (s, 3H), 2.74-2.72 (m, 2H), 2.58-2.50 (m, 2H), 2.20-2.16 (m, 2H), 2.10-2.04 (m, 2H), 1.09 (d, 6H, J = 6.8 Hz), 1.05 (d, 6H, J = 6.0 Hz).
726		539	¹ H-NMR (500 MHz, 4d-MeOD) δ ppm 8.00 (d, 1H, J = 1.5 Hz), 7.90 (d, 1H, J = 5.5 Hz), 7.63 (d, 1H, J = 8.0 Hz), 7.41 (d, 1H, J = 1.0 Hz), 7.27 (dd, 1H, J = 8.0, 1.0 Hz), 6.86 (d, 1H, J = 1.0 Hz), 6.03 (d, 1H, J = 5.0 Hz), 5.43-5.41 (m, 1H), 4.92-4.90 (m, 2H), 4.68-4.66 (m, 2H), 3.80-3.75 (m, 2H), 3.75-3.70 (m, 2H), 3.56-3.52 (m, 4H), 3.37-3.32 (m, 2H), 3.25-3.18 (m, 1H), 2.84-2.76 (m, 3H), 2.09-2.05 (m, 2H), 1.96-1.88 (m, 2H), 1.30 (d, 6H, J = 6.5 Hz).
727		540	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.84 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.09 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.00-3.82 (m, 2H), 3.82-3.65 (m, 2H), 3.58-3.54 (m, 2H), 3.52-3.48 (m, 2H), 2.80-2.65 (m, 2H), 2.65-2.50 (m, 3H), 2.08-1.90 (m, 5H), 0.85 (d, 6H, J = 6.4 Hz), 0.84-0.74 (m, 4H).
728		541	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.78 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 3.74-3.68 (m, 2H), 3.68-3.62 (m, 1H), 3.58-3.50 (m, 2H), 3.48-3.40 (m, 4H), 3.30-3.20 (m, 1H), 2.88 (s, 3H), 2.74-2.70 (m, 1H), 2.68-2.56 (m, 4H), 2.54-2.50 (m, 2H), 2.47-2.40 (m, 2H), 2.03-1.92 (m, 2H), 1.90-1.78 (m, 2H), 0.99 (d, 6H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
729		541	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (s, 1H), 7.92 (d, 1H, J = 5.2 Hz), 7.85 (d, 2H, J = 7.2 Hz), 8.46 (d, 2H, J = 7.2 Hz), 7.07 (s, 1H), 5.99 (d, 1H, J = 5.2 Hz), 3.78-3.74 (m, 2H), 3.70-3.65 (m, 2H), 3.52-3.43 (m, 1H), 3.52-3.48 (m, 2H), 3.47-3.44 (m, 2H), 3.30-3.21 (m, 2H), 2.96 (s, 3H), 2.75-2.70 (m, 2H), 2.35-2.30 (m, 2H), 2.10-2.01 (m, 2H), 1.75-1.71 (m, 2H), 1.62-1.57 (m, 2H), 1.43-1.39 (m, 2H), 1.01-0.80 (m, 6H).
730		541	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.0 Hz), 7.90 (d, 1H, J = 5.5 Hz), 7.77 (d, 2H, J = 8.0 Hz), 7.48 (d, 2H, J = 8.0 Hz), 7.03 (d, 1H, J = 1.5 Hz), 5.98 (d, 1H, J = 5.5 Hz), 3.71-3.68 (m, 2H), 3.68-3.65 (m, 1H), 3.55-3.53 (m, 2H), 3.47-3.43 (m, 4H), 3.30-3.25 (m, 2H), 2.91 (s, 3H), 2.81-2.77 (m, 2H), 2.71-2.68 (m, 1H), 2.65-2.60 (m, 2H), 2.54-2.51 (m, 2H), 2.39-2.36 (m, 1H), 1.80-1.78 (m, 2H), 1.72-1.69 (m, 1H), 1.40-1.35 (m, 1H), 1.01 (t, 6H, J = 6.0 Hz).
731		542	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.5 Hz), 8.29 (d, 1H, J = 1.5 Hz), 7.94-7.91 (m, 2H), 7.77 (dd, 1H, J = 8.5, 2.0 Hz), 7.19 (d, 1H, J = 1.5 Hz), 5.98 (d, 1H, J = 5.0 Hz), 3.71-3.66 (m, 2H), 3.65-3.63 (m, 1H), 3.56-3.55 (m, 2H), 3.48-3.46 (m, 4H), 3.30-3.25 (m, 1H), 2.92 (s, 3H), 2.72-2.69 (m, 1H), 2.63-2.60 (m, 4H), 2.52-2.47 (m, 4H), 2.02-2.00 (m, 2H), 1.92-1.86 (m, 2H), 1.00 (d, 6H, J = 6.5 Hz).
732		545	
733		549	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.5 Hz), 7.95-7.93 (m, 2H), 7.79 (dd, 1H, J = 8.5, 2.5 Hz), 7.19 (d, 1H, J = 1.5 Hz), 6.01 (d, 1H, J = 5.5 Hz), 5.34 (quintet, 1H, J = 6.0 Hz), 4.79 (t, 2H, J = 7.0 Hz), 4.56-4.52 (m, 4H), 4.45 (t, 2H, J = 6.0 Hz), 3.72-3.68 (m, 2H), 3.63-3.59 (m, 2H), 3.52-3.50 (m, 4H), 3.45-3.43 (m, 1H), 2.92 (s, 3H), 2.54-2.52 (m, 2H), 2.19-2.15 (m, 2H), 2.04-1.98 (m, 2H), 1.95-1.91 (m, 2H).






#	Structure	LCMS (M+1)	NMR
734		549	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.5 Hz), 8.29 (d, 1H, J = 2.0 Hz), 7.95-7.92 (m, 2H), 7.79 (dd, 1H, J = 8.5, 2.0 Hz), 7.19 (d, 1H, J = 2.0 Hz), 6.01 (d, 1H, J = 5.5 Hz), 5.35 (quintet, 1H, J = 5.5 Hz), 4.80 (t, 2H, J = 6.5 Hz), 4.54 (dd, 2H, J = 7.5, 5.5 Hz), 3.71-3.70 (m, 2H), 3.67-3.60 (m, 2H), 3.52-3.51 (m, 4H), 3.33-3.30 (m, 1H), 3.09 (q, 2H, J = 6.5 Hz), 2.70-2.55 (m, 2H), 2.55-2.50 (m, 2H), 2.02-1.98 (m, 2H), 1.09 (t, 3H, J = 6.5 Hz), 1.01 (d, 6H, J = 6.5 Hz).
735		550	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.57 (d, 1H, J = 1.6 Hz), 8.27 (d, 1H, J = 1.2 Hz), 7.96-7.88 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.0 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.30 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.80-3.70 (m, 1H), 3.60-3.50 (m, 4H), 3.48-3.42 (m, 4H), 3.44-3.41 (m, 1H), 3.26-3.20 (m, 1H), 3.25 (s, 3H), 2.91 (s, 3H), 2.80-2.70 (m, 1H), 2.67-2.55 (m, 4H), 2.05-1.95 (m, 2H), 1.92-1.80 (m, 2H), 1.08 (d, 6H, J = 6.8 Hz), 0.98 (d, 3H, J = 7.2 Hz).
736		550	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 8.00-7.90 (m, 2H), 7.77 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.32 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 3.90-3.70 (m, 1H), 3.60-3.50 (m, 4H), 3.47-3.42 (m, 4H), 3.43-3.40 (m, 1H), 3.30-3.20 (m, 1H), 3.25 (s, 3H), 2.92 (s, 3H), 2.80-2.72 (m, 1H), 2.70-2.58 (m, 4H), 2.40-1.96 (m, 2H), 1.94-1.80 (m, 2H), 1.09 (d, 6H, J = 6.8 Hz), 0.99 (d, 3H, J = 6.8 Hz).
737		552	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.04 (t, J = 2.2 Hz, 1H), 7.97 - 7.81 (m, 2H), 7.26 - 7.16 (m, 2H), 7.01 (d, J = 1.8 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 5.33 - 5.23 (m, 1H), 4.74 (t, J = 6.9 Hz, 2H), 4.48 (dd, J = 7.6, 5.1 Hz, 2H), 3.65 (br.s, 2H), 3.55 (br.s, 2H), 3.45 (dd, J = 6.7, 3.7 Hz, 4H), 3.29 (s, 5H), 2.87 (s, 3H), 2.72 - 2.50 (m, 3H), 1.93 (d, J = 13.0 Hz, 2H), 1.82 (d, J = 13.1 Hz, 2H), 0.95 (d, J = 6.5 Hz, 6H).

#	Structure	LCMS (M+1)	NMR
738		557	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.22 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.84 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 6.32 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 3.85-3.75 (m, 1H), 3.55-3.50 (m, 4H), 3.47-3.42 (m, 4H), 2.80-2.65 (m, 2H), 2.65-2.50 (m, 3H), 2.08-1.90 (m, 4H), 1.08 (d, 6H, J = 6.4 Hz), 0.85 (d, 6H, J = 6.4 Hz).
739		557	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.04 (s, 1H), 5.97 (d, 1H, J = 5.6 Hz), 4.23-4.22 (m, 1H), 3.69-3.65 (m, 2H), 3.65-3.63 (m, 1H), 3.55-3.54 (m, 2H), 3.52-3.50 (m, 1H), 3.49-3.45 (m, 4H), 3.30-3.26 (m, 3H), 2.89 (s, 3H), 2.74-2.66 (m, 2H), 2.65-2.50 (m, 6H), 1.98-1.93 (m, 2H), 1.92-1.88 (m, 2H), 0.95 (d, 3H, J = 6.4 Hz).
740		557	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.5 Hz), 7.91 (d, 1H, J = 5.5 Hz), 7.79 (d, 2H, J = 8.5 Hz), 7.39 (d, 2H, J = 8.5 Hz), 7.04 (d, 1H, J = 1.5 Hz), 5.97 (d, 1H, J = 5.5 Hz), 4.30-4.20 (m, 1H), 3.69-3.65 (m, 2H), 3.65-3.61 (m, 1H), 3.55-3.54 (m, 2H), 3.51-3.48 (m, 1H), 3.49-3.45 (m, 4H), 3.30-3.26 (m, 3H), 2.89 (s, 3H), 2.69-2.65 (m, 2H), 2.65-2.50 (m, 6H), 1.98-1.95 (m, 2H), 1.90-1.88 (m, 2H), 0.95 (d, 3H, J = 6.5 Hz).
741		558	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.5 Hz), 8.30 (d, 1H, J = 1.5 Hz), 7.95-7.91 (m, 2H), 7.80-7.78 (m, 1H), 7.19 (d, 1H, J = 1.5 Hz), 5.99 (d, 1H, J = 5.0 Hz), 4.23-4.21 (m, 1H), 3.74-3.69 (m, 2H), 3.70-3.65 (m, 1H), 3.56-3.55 (m, 2H), 3.49-3.47 (m, 5H), 3.31-3.29 (m, 3H), 2.93 (s, 3H), 2.65-2.64 (m, 1H), 2.64-2.60 (m, 6H), 2.51-2.50 (m, 1H), 2.05-2.01 (m, 2H), 1.99-1.98 (m, 2H), 0.96-0.95 (d, 3H, J = 6.5 Hz).

#	Structure	LCMS (M+1)	NMR
742		558	¹ H NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.30 (d, 1H, J = 2.0 Hz), 7.96-7.93 (m, 2H), 7.79 (dd, 1H, J = 8.0, 2.0 Hz), 7.19 (d, 1H, J = 1.5 Hz), 5.99 (d, 1H, J = 5.5 Hz), 3.73-3.68 (m, 2H), 3.69-3.65 (m, 1H), 3.67-3.56 (m, 2H), 3.55-3.49 (m, 4H), 3.31-3.28 (m, 4H), 2.94 (s, 3H), 2.95-2.85 (m, 2H), 2.66-2.65 (m, 2H), 2.56-2.53 (m, 2H), 2.49-2.48 (m, 2H), 2.18-1.99 (m, 4H), 1.04-1.03 (m, 3H).
743		565	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.57 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.95-7.90 (m, 2H), 7.77 (dd, 1H, J = 8.0 Hz, 2.0 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.2 Hz), 5.39-5.28 (m, 1H), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.80-3.75 (m, 2H), 3.72-3.67 (m, 2H), 3.63-3.58 (m, 4H), 3.53-3.49 (m, 1H), 3.32-3.20 (m, 1H), 3.26 (s, 3H), 2.91 (s, 3H), 2.80-3.75 (m, 1H), 2.70-2.56 (m, 4H), 2.05-1.98 (m, 2H), 1.95-1.79 (m, 2H), 0.98 (d, 3H, J = 6.8 Hz).
744		565	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.57 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.95-7.90 (m, 2H), 7.77 (dd, 1H, J = 8.0 Hz, 2.0 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.2 Hz), 5.39-5.28 (m, 1H), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.80-3.75 (m, 2H), 3.72-3.67 (m, 2H), 3.63-3.58 (m, 4H), 3.53-3.49 (m, 1H), 3.32-3.20 (m, 1H), 3.26 (s, 3H), 2.91 (s, 3H), 2.80-3.75 (m, 1H), 2.70-2.56 (m, 4H), 2.05-1.98 (m, 2H), 1.95-1.79 (m, 2H), 0.98 (d, 3H, J = 6.8 Hz).
745		571	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.5 Hz), 7.90 (d, 1H, J = 5.5 Hz), 7.78 (d, 2H, J = 8.5 Hz), 7.38 (d, 2H, J = 8.5 Hz), 7.03 (d, 1H, J = 1.0 Hz), 5.96 (d, 1H, J = 5.5 Hz), 3.70-3.65 (m, 2H), 3.65-3.58 (m, 1H), 3.55-3.51 (m, 2H), 3.46-3.41 (m, 6H), 3.30-3.22 (m, 2H), 3.24 (s, 3H), 2.87 (s, 3H), 2.77-2.73 (m, 1H), 2.65-2.58 (m, 6H), 2.50-2.47 (m, 1H), 1.97-1.94 (m, 2H), 1.86-1.82 (m, 2H), 1.00 (d, 3H, J = 6.0 Hz).

#	Structure	LCMS (M+1)	NMR
746		571	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.5 Hz), 7.90 (d, 1H, J = 5.5 Hz), 7.78 (d, 2H, J = 8.5 Hz), 7.38 (d, 2H, J = 8.5 Hz), 7.03 (d, 1H, J = 1.0 Hz), 5.96 (d, 1H, J = 5.5 Hz), 3.70-3.65 (m, 2H), 3.65-3.58 (m, 1H), 3.55-3.51 (m, 2H), 3.46-3.41 (m, 6H), 3.30-3.22 (m, 2H), 3.24 (s, 3H), 2.87 (s, 3H), 2.77-2.73 (m, 1H), 2.65-2.58 (m, 6H), 2.50-2.47 (m, 1H), 1.97-1.94 (m, 2H), 1.86-1.82 (m, 2H), 1.00 (d, 3H, J = 6.0 Hz).
747		572	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.6 Hz), 7.93 (d, 1H, J = 5.6 Hz), 7.84 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 2.0 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 8.0, 5.6 Hz), 3.80-3.75 (m, 2H), 3.70-3.65 (m, 2H), 3.55-3.45 (m, 4H), 2.78-2.70 (m, 2H), 2.65-2.50 (m, 3H), 2.08-1.80 (m, 4H), 0.85 (d, 6H, J = 6.4 Hz).
748		572	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.57 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 2.0 Hz), 7.95-7.90 (m, 2H), 7.77 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.73-3.67 (m, 2H), 3.58-3.53 (m, 2H), 3.49-3.40 (m, 4H), 3.40-3.35 (m, 2H), 3.32-3.20 (m, 4H), 3.25 (s, 3H), 2.92 (s, 3H), 2.81-2.75 (m, 2H), 2.60-2.52 (m, 4H), 2.50-2.47 (m, 1H), 2.03-1.95 (m, 2H), 1.90-1.84 (m, 2H), 0.99 (d, 3H, J = 6.8 Hz).
749		572	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.57 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 2.0 Hz), 7.95-7.90 (m, 2H), 7.77 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.73-3.67 (m, 2H), 3.58-3.53 (m, 2H), 3.49-3.40 (m, 4H), 3.40-3.35 (m, 2H), 3.32-3.20 (m, 4H), 3.25 (s, 3H), 2.92 (s, 3H), 2.81-2.75 (m, 2H), 2.60-2.52 (m, 4H), 2.50-2.47 (m, 1H), 2.03-1.95 (m, 2H), 1.90-1.84 (m, 2H), 0.99 (d, 3H, J = 6.8 Hz).

#	Structure	LCMS (M+1)	NMR
750		586	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.5 Hz), 8.28 (d, 1H, J = 1.0 Hz), 7.93-7.92 (m, 2H), 7.78 (dd, 1H, J = 7.0, 2.0 Hz), 7.18 (s, 1H), 6.69 (d, 1H, J = 76.0 Hz), 6.31 (d, 1H, J = 7.5 Hz), 6.00 (d, 1H, J = 5.0 Hz), 3.94-3.89 (m, 1H), 3.84-3.75 (m, 1H), 3.55-3.52 (m, 4H), 3.48-3.45 (m, 4H), 2.93 (s, 3H), 2.87-2.81 (m, 1H), 2.70-2.61 (m, 4H), 2.06-2.00 (m, 2H), 1.95-1.86 (m, 2H), 1.09 (d, 6H, J = 6.5 Hz), 1.02 (d, 3H, J = 6.5 Hz).
751		407	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.54 (s, 1H), 7.98 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.5 Hz, 1H), 7.81 – 7.70 (m, 2H), 7.55 – 7.38 (m, 2H), 6.94 (d, J = 1.8 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 4.86 (d, J = 3.6 Hz, 1H), 4.03 (s, 2H), 3.85 (s, 2H), 3.62 (s, 2H), 3.59 – 3.47 (m, 3H), 3.16 – 3.07 (m, 1H), 3.00 (dd, J = 12.8, 9.3 Hz, 1H), 2.09 – 1.96 (m, 1H), 0.92 (dt, J = 4.8, 2.8 Hz, 2H), 0.86 (ddt, J = 7.5, 4.6, 2.5 Hz, 2H).
752		407	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.95 (d, J = 1.8 Hz, 1H), 7.83 (d, J = 5.4 Hz, 1H), 7.79 – 7.66 (m, 2H), 7.44 – 7.35 (m, 2H), 6.92 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 5.48 (s, 1H), 4.02 (s, 2H), 3.97 (dd, J = 8.0, 4.7 Hz, 1H), 3.84 (s, 2H), 3.71 (dd, J = 10.9, 4.7 Hz, 1H), 3.63 (s, 2H), 3.60 – 3.55 (m, 1H), 3.54 (d, J = 8.0 Hz, 2H), 2.01 (tt, J = 8.0, 4.7 Hz, 1H), 0.92 (dt, J = 4.8, 2.8 Hz, 2H), 0.85 (ddt, J = 7.4, 4.6, 2.5 Hz, 2H).
753		407	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.96 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.5 Hz, 1H), 7.77 – 7.63 (m, 2H), 7.48 – 7.35 (m, 2H), 6.92 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.03 (s, 2H), 3.98 (dd, J = 8.0, 4.7 Hz, 1H), 3.85 (s, 2H), 3.72 (dd, J = 10.9, 4.8 Hz, 1H), 3.63 (s, 2H), 3.58 (dd, J = 10.8, 8.0 Hz, 1H), 3.54 (s, 2H), 2.02 (tt, J = 7.9, 4.7 Hz, 1H), 0.92 (dt, J = 4.8, 2.8 Hz, 2H), 0.86 (ddt, J = 7.5, 4.6, 2.5 Hz, 2H).
754		411	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.05 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.84 – 7.69 (m, 2H), 7.51 – 7.33 (m, 2H), 6.96 (d, J = 1.8 Hz, 1H), 6.04 (d, J = 5.4 Hz, 1H), 4.19 (q, J = 7.1 Hz, 2H), 4.02 (dd, J = 7.9, 4.7 Hz, 1H), 3.80 – 3.68 (m, 5H), 3.63 – 3.48 (m, 5H), 2.63 (p, J = 1.9 Hz, 3H), 1.31 (t, J = 7.1 Hz, 3H).





#	Structure	LCMS (M+1)	NMR
755		419	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.18 (d, J = 1.8 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.86 – 7.77 (m, 2H), 7.63 – 7.53 (m, 2H), 7.05 (d, J = 1.8 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 4.72 (d, J = 6.1 Hz, 2H), 4.65 (d, J = 6.0 Hz, 2H), 3.91 (s, 2H), 3.70 (s, 2H), 3.51 (d, J = 27.0 Hz, 5H), 2.01 (tt, J = 7.7, 4.9 Hz, 1H), 0.81 – 0.65 (m, 4H).
756		421	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.74 – 7.66 (m, 2H), 7.42 – 7.35 (m, 2H), 7.36 – 7.21 (m, 1H), 6.93 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.04 (d, J = 9.6 Hz, 2H), 3.91 – 3.85 (m, 0H), 3.83 – 3.72 (m, 7H), 3.69 – 3.61 (m, 6H), 3.55 (d, J = 6.9 Hz, 2H), 2.77 (t, J = 5.6 Hz, 2H), 2.72 (dt, J = 11.3, 5.6 Hz, 5H), 2.02 (tt, J = 7.8, 4.7 Hz, 1H), 0.92 (dt, J = 4.8, 2.8 Hz, 2H), 0.86 (ddt, J = 7.5, 4.6, 2.6 Hz, 2H).
757		421	
758		421	
759		425	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.02 (dd, J = 2.8, 1.8 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.81 (t, J = 8.2 Hz, 1H), 7.30 – 7.15 (m, 2H), 7.04 – 6.95 (m, 1H), 6.00 (d, J = 5.5 Hz, 1H), 4.10 – 3.94 (m, 3H), 3.85 (s, 2H), 3.72 (dd, J = 10.9, 4.9 Hz, 1H), 3.68 – 3.48 (m, 5H), 2.02 (tt, J = 7.9, 4.8 Hz, 1H), 0.92 (dt, J = 4.9, 2.8 Hz, 2H), 0.86 (ddt, J = 7.5, 4.7, 2.5 Hz, 2H).





#	Structure	LCMS (M+1)	NMR
760		433	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.96 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.73 – 7.63 (m, 2H), 7.41 – 7.32 (m, 2H), 6.93 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.70 (t, J = 6.8 Hz, 2H), 4.44 (t, J = 6.4 Hz, 2H), 4.08 – 3.98 (m, 3H), 3.85 (s, 2H), 3.71 (s, 2H), 3.63 (s, 2H), 3.54 (s, 2H), 2.02 (tt, J = 7.9, 4.8 Hz, 1H), 0.92 (dt, J = 4.8, 2.8 Hz, 2H), 0.86 (ddt, J = 7.5, 4.6, 2.5 Hz, 2H).
761		435	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.96 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.73 – 7.65 (m, 2H), 7.42 – 7.36 (m, 2H), 6.92 (d, J = 1.8 Hz, 1H), 5.98 (d, J = 5.5 Hz, 1H), 4.02 (s, 2H), 3.84 (s, 2H), 3.79 (s, 1H), 3.62 (s, 2H), 3.59 – 3.47 (m, 2H), 3.34 (s, 6H), 2.01 (tt, J = 7.9, 4.7 Hz, 1H), 1.20 (s, 3H), 1.12 (s, 3H), 0.96 – 0.90 (m, 2H), 0.86 (tdd, J = 7.5, 5.3, 2.4 Hz, 2H).
762		439	¹ H NMR (400 MHz, Chloroform-d) δ 7.95 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.5 Hz, 1H), 7.75 – 7.67 (m, 2H), 7.43 – 7.34 (m, 2H), 6.89 (d, J = 1.9 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.2, 5.1 Hz, 1H), 4.89 (ddd, J = 7.4, 6.2, 1.0 Hz, 2H), 4.65 (ddd, J = 7.5, 5.2, 0.9 Hz, 2H), 3.97 (dd, J = 8.0, 4.7 Hz, 1H), 3.74 – 3.68 (m, 3H), 3.57 (dd, J = 10.8, 8.0 Hz, 1H), 3.53 (dd, J = 6.5, 3.9 Hz, 4H).
763		439	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.96 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.77 – 7.62 (m, 2H), 7.47 – 7.33 (m, 2H), 6.89 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.3, 5.1 Hz, 1H), 4.90 (ddd, J = 7.4, 6.2, 1.0 Hz, 2H), 4.66 (ddd, J = 7.5, 5.1, 0.9 Hz, 2H), 3.99 (dd, J = 7.9, 4.7 Hz, 1H), 3.80 (s, 3H), 3.77 – 3.64 (m, 2H), 3.59 (dd, J = 10.9, 8.0 Hz, 1H), 3.54 – 3.44 (m, 4H).
764		439	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.95 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.5 Hz, 1H), 7.76 – 7.59 (m, 2H), 7.49 – 7.32 (m, 2H), 6.89 (d, J = 1.8 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 4.17 (q, J = 7.1 Hz, 2H), 3.79 (s, 1H), 3.72 (d, J = 6.1 Hz, 4H), 3.57 – 3.41 (m, 4H), 1.29 (t, J = 7.1 Hz, 3H), 1.20 (s, 3H), 1.12 (s, 3H).






#	Structure	LCMS (M+1)	NMR
765		440	
766		443	1H NMR (500 MHz, Methanol-d4) δ 7.98 (dd, J = 12.3, 1.8 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.69 (dd, J = 16.3, 7.9 Hz, 2H), 7.25 (t, J = 7.2 Hz, 2H), 6.94 (dd, J = 10.9, 1.8 Hz, 1H), 6.42 (s, 0H), 6.02 (d, J = 5.4 Hz, 1H), 5.57 (s, 1H), 4.07 (d, J = 18.4 Hz, 2H), 3.87 (s, 2H), 3.65 (s, 2H), 3.57 (s, 2H), 3.48 (s, 1H), 3.38 (s, 2H), 3.05 (dt, J = 18.3, 6.0 Hz, 2H), 2.94 (t, J = 5.8 Hz, 1H), 2.61 (t, J = 5.9 Hz, 1H), 2.46 (t, J = 5.9 Hz, 1H), 2.15 (s, 1H), 2.04 (tt, J = 8.2, 5.0 Hz, 1H), 0.95 (dt, J = 5.5, 3.1 Hz, 2H), 0.88 (dq, J = 10.4, 4.0, 3.6 Hz, 2H).
767		443	1H NMR (400 MHz, Methanol-d4) δ 7.94 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.5 Hz, 1H), 7.66 (d, J = 8.3 Hz, 2H), 7.24 (d, J = 8.0 Hz, 2H), 6.90 (d, J = 1.9 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 4.25 (s, 3H), 4.03 (s, 3H), 3.84 (s, 2H), 3.58 (d, J = 26.6 Hz, 5H), 2.75 – 2.66 (m, 2H), 2.42 (td, J = 9.7, 2.9 Hz, 2H), 2.06 – 1.99 (m, 1H), 1.30 (d, J = 14.4 Hz, 1H), 0.99 – 0.81 (m, 4H).
768		447	1H NMR (400 MHz, DMSO-d6) δ 8.14 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.78 – 7.68 (m, 2H), 7.56 – 7.47 (m, 2H), 7.02 (d, J = 1.9 Hz, 1H), 5.96 (d, J = 5.5 Hz, 1H), 3.91 (s, 2H), 3.86 (td, J = 11.2, 2.3 Hz, 2H), 3.69 (s, 2H), 3.62 (dt, J = 11.1, 3.8 Hz, 2H), 3.50 (d, J = 25.8 Hz, 4H), 2.07 – 1.91 (m, 3H), 1.53 (d, J = 13.1 Hz, 2H), 0.84 – 0.67 (m, 4H).
769		447	





#	Structure	LCMS (M+1)	NMR
770		449	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.54 (s, 2H), 7.98 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.75 (d, J = 8.0 Hz, 2H), 7.43 (d, J = 8.0 Hz, 2H), 6.94 (d, J = 1.8 Hz, 1H), 6.00 (d, J = 5.5 Hz, 1H), 4.55 (s, 1H), 4.03 (s, 3H), 3.82 (d, J = 23.8 Hz, 2H), 3.79 – 3.67 (m, 1H), 3.63 (s, 2H), 3.53 (d, J = 18.9 Hz, 2H), 2.05 – 1.94 (m, 1H), 1.28 (s, 1H), 1.16 (dd, J = 16.5, 9.4 Hz, 6H), 0.96 – 0.81 (m, 5H).
771		453	¹ H NMR (500 MHz, Methanol-d ₄) δ 8.45 (s, 2H), 8.11 – 7.96 (m, 1H), 7.87 (dd, J = 17.6, 6.7 Hz, 3H), 7.51 (d, J = 8.2 Hz, 2H), 6.96 (d, J = 1.9 Hz, 1H), 6.05 (d, J = 5.5 Hz, 1H), 5.44 (p, J = 5.6 Hz, 1H), 4.93 (t, J = 6.9 Hz, 2H), 4.68 (dd, J = 7.6, 5.1 Hz, 2H), 3.96 – 3.66 (m, 6H), 3.56 (t, J = 5.1 Hz, 4H), 3.01 (s, 2H), 2.88 (s, 1H), 1.74 (s, 3H).
772		561	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.55 (s, 0H), 8.14 (d, J = 2.0 Hz, 1H), 7.89 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.0 Hz, 2H), 7.33 (d, J = 8.2 Hz, 2H), 7.01 (d, J = 1.9 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.13 (s, 2H), 3.92 (s, 2H), 3.70 (s, 2H), 3.57 (d, J = 25.1 Hz, 3H), 3.46 (s, 1H), 3.15 (s, 4H), 2.79 (d, J = 10.7 Hz, 1H), 2.07 – 1.98 (m, 1H), 1.94 – 1.80 (m, 2H), 1.63 (d, J = 10.9 Hz, 1H), 1.51 (s, 1H), 1.22 (s, 1H), 0.75 (t, J = 7.9, 2.9 Hz, 4H).
773		462	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.16 (s, 1H), 7.91 (d, 1H, J = 5.5 Hz), 7.74 (d, 2H, J = 8.0 Hz), 7.25 (d, 2H, J = 8.0 Hz), 7.01 (s, 1H), 5.99 (d, 1H, J = 5.5 Hz), 5.34 (quintet, 1H, d, 1H, J = 5.5 Hz), 4.80 (t, 2H, d, 1H, J = 7.0 Hz), 4.53 (dd, 2H, d, 1H, J = 7.0, 5.5 Hz), 3.72-3.67 (m, 2H), 3.63-3.58 (m, 2H), 3.55-3.45 (m, 4H), 3.20-3.15 (m, 1H), 3.06-3.02 (m, 2H), 2.64-2.59 (m, 2H), 1.75-1.70 (m, 2H), 1.57-1.52 (m, 2H).
774		462	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.2 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.24 (d, 2H, J = 8.4 Hz), 7.00 (s, 1H), 6.00 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.69 (m, 2H), 3.59 (m, 2H), 3.49-3.48 (m, 4H), 2.96-2.93 (m, 2H), 2.60-2.52 (m, 2H), 2.47 (m, 1H), 1.89-1.87 (m, 1H), 1.70-1.65 (m, 1H), 1.62-1.58 (m, 1H), 1.55-1.40 (m, 2H).





#	Structure	LCMS (M+1)	NMR
775		463	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.27 (s, 1H), 8.08 (d, 1H, J = 1.5 Hz), 7.88 (d, 1H, J = 5.5 Hz), 7.67 (d, 2H, J = 9.0 Hz), 6.97 (d, 2H, J = 9.0 Hz), 6.93 (d, 1H, J = 1.5 Hz), 5.98 (d, 1H, J = 5.5 Hz), 5.33 (quintet, 1H, J = 5.0 Hz), 4.79 (t, 2H, J = 7.0 Hz), 4.53 (dd, 2H, J = 7.5, 5.0 Hz), 3.70-3.68 (m, 2H), 3.65-3.64 (m, 2H), 3.48-3.46 (m, 4H), 3.18-3.16 (m, 4H), 2.99-2.97 (m, 4H).
776		464	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.28 (d, 1H, J = 2.4 Hz), 8.15 (d, 1H, J = 1.2 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.70 (d, 1H, J = 8.8 Hz), 7.73 (dd, 1H, J = 8.8, 2.4 Hz), 7.06 (d, 1H, J = 1.2 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.33 (quintet, 1H, J = 5.6 Hz), 4.78 (t, 2H, J = 7.2 Hz), 4.52 (dd, 2H, J = 7.2, 5.6 Hz), 3.70-3.65 (m, 2H), 3.65-3.58 (m, 2H), 3.51-3.45 (m, 4H), 3.25-3.15 (m, 4H), 3.05-2.93 (m, 4H).
777		464	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.62-3.60 (m, 4H), 3.47-3.46 (m, 4H), 2.96-2.87 (m, 5H), 2.70-2.67 (m, 1H), 2.49-2.46 (m, 1H), 2.11-2.07 (m, 1H), 1.98-1.92 (m, 1H), 1.69-1.59 (m, 1H), 1.46-1.42 (m, 1H), 1.22 (t, 3H, J = 7.2 Hz).
778		465	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.61 (s, 1H), 8.32 (d, 1H, J = 1.5 Hz), 8.29 (s, 1H), 7.99 (d, 1H, J = 8.0 Hz), 7.95 (d, 1H, J = 5.0 Hz), 7.82 (d, 1H, J = 7.5 Hz), 7.20 (d, 1H, J = 1.5 Hz), 6.01 (d, 1H, J = 5.5 Hz), 4.10 (q, 2H, J = 7.0 Hz), 3.70-3.58 (m, 4H), 3.54-3.46 (m, 4H), 3.38-3.25 (m, 1H), 3.15-3.05 (m, 2H), 3.02 (s, 3H), 2.90-2.76 (m, 1H), 2.26-2.18 (m, 1H), 2.14-2.04 (m, 1H), 1.90-1.78 (m, 1H), 1.76-1.66 (m, 1H), 1.23 (t, 3H, J = 7.0 Hz).
779		467	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.96 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.73 – 7.62 (m, 2H), 7.43 – 7.36 (m, 2H), 6.90 (d, J = 1.9 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.2, 5.1 Hz, 1H), 4.90 (ddd, J = 7.4, 6.2, 1.0 Hz, 2H), 4.66 (ddd, J = 7.5, 5.1, 0.9 Hz, 2H), 3.76 (d, J = 26.0 Hz, 1H), 3.58 – 3.47 (m, 4H), 1.16 (d, J = 29.4 Hz, 6H).





#	Structure	LCMS (M+1)	NMR
780		467	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.72 (d, J = 8.1 Hz, 2H), 7.38 (d, J = 8.2 Hz, 2H), 6.90 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.4 Hz, 1H), 5.46 – 5.35 (m, 1H), 4.90 (dd, J = 7.5, 6.5 Hz, 2H), 4.66 (dd, J = 7.8, 5.2 Hz, 2H), 3.88 – 3.61 (m, 8H), 3.61 – 3.46 (m, 5H), 2.58 (q, J = 7.0 Hz, 2H), 1.29 (s, 2H), 1.13 (t, J = 7.2 Hz, 3H).
781		475	
782		478	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.78 (d, 2H, J = 8.4 Hz), 7.41 (d, 2H, J = 8.4 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.62-3.61 (m, 4H), 3.47-3.46 (m, 4H), 3.19-3.15 (m, 1H), 3.08-3.04 (m, 1H), 2.90-2.87 (m, 2H), 2.68-2.65 (m, 1H), 2.51-2.50 (m, 1H), 2.05-2.04 (m, 1H), 1.97-1.96 (m, 1H), 1.62-1.60 (m, 1H), 1.41-1.39 (m, 1H), 1.22 (t, 3H, J = 7.2 Hz), 1.09 (t, 3H, J = 6.4 Hz).
783		478	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.40 (d, 2H, J = 8.4 Hz), 7.02 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.62-3.55 (m, 4H), 3.20-3.45 (m, 4H), 3.20-3.15 (m, 1H), 3.08-3.03 (m, 1H), 2.95-2.88 (m, 2H), 2.73-2.68 (m, 1H), 2.54-2.50 (m, 1H), 2.08-2.04 (m, 1H), 2.00-1.95 (m, 1H), 1.69-1.64 (m, 1H), 1.48-1.43 (m, 1H), 1.22 (t, 3H, J = 7.2 Hz), 1.11 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
784		478	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (s, 1H), 8.29 (d, 1H, J = 1.6 Hz), 8.27 (s, 1H), 7.97 (d, 1H, J = 8.0 Hz), 7.93 (d, 1H, J = 5.2 Hz), 7.80 (d, 1H, J = 7.6 Hz), 7.18 (d, 1H, J = 1.2 Hz), 6.32 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.75-3.60 (m, 1H), 3.67-3.48 (m, 4H), 3.47-3.40 (m, 4H), 3.30-3.20 (m, 1H), 3.12-3.02 (m, 2H), 3.00 (s, 3H), 2.84-2.66 (m, 1H), 2.24-2.14 (m, 1H), 2.12-2.00 (m, 1H), 1.86-1.72 (m, 1H), 1.70-1.60 (m, 1H), 1.07 (d, 6H, J = 6.4 Hz).
785		478	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.56 (d, 1H, J = 1.6 Hz), 8.27 (d, 1H, J = 1.6 Hz), 7.96-7.90 (m, 2H), 7.75 (dd, 1H, J = 8.0, 2.0 Hz), 7.16 (d, 1H, J = 1.6 Hz), 6.30 (d, 1H, J = 7.2 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.80-3.70 (m, 1H), 3.60-3.50 (m, 4H), 3.48-3.40 (m, 4H), 2.91 (s, 3H), 2.90-2.80 (m, 2H), 2.78-2.70 (m, 2H), 2.00-1.88 (m, 2H), 1.86-1.74 (m, 2H), 1.08 (d, 6H, J = 6.4 Hz).
786		478	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.5 Hz), 8.29 (d, 1H, J = 1.0 Hz), 7.96-7.93 (m, 2H), 7.80 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.5 Hz), 6.31 (d, 1H, J = 7.0 Hz), 6.00 (d, 1H, J = 5.5 Hz), 3.81-3.75 (m, 1H), 3.54-3.53 (m, 4H), 3.46-3.45 (m, 4H), 3.17-3.14 (m, 1H), 3.03-3.01 (m, 1H), 3.01 (s, 3H), 3.00-2.92 (m, 1H), 2.70-2.64 (m, 1H), 2.18 (m, 1H), 2.07-2.02 (m, 1H), 1.75-1.72 (m, 1H), 1.60-1.58 (m, 1H), 1.08 (d, 6H, J = 6.5 Hz).
787		479	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.4 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.95-7.90 (m, 2H), 7.79 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.09 (q, 2H, J = 7.2 Hz), 3.64-3.60 (m, 4H), 3.50-3.45 (m, 4H), 3.22-3.18 (m, 1H), 3.11-3.07 (m, 1H), 2.92-2.84 (m, 2H), 2.78-2.74 (m, 1H), 2.51-2.50 (m, 1H), 2.05-1.99 (m, 2H), 1.54-1.53 (m, 1H), 1.44-1.43 (m, 1H), 1.22 (t, 3H, J = 7.2 Hz), 1.12 (t, 3H, J = 7.2 Hz).





#	Structure	LCMS (M+1)	NMR
788		479	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.31 (s, 1H), 8.29 (d, 1H, J = 2.0 Hz), 7.97-7.91 (m, 2H), 7.79 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 1.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 4.08 (q, 2H, J = 7.2 Hz), 3.68-3.56 (m, 4H), 3.52-3.42 (m, 4H), 3.26-3.12 (m, 2H), 3.10-2.98 (m, 2H), 2.97-2.90 (m, 1H), 2.76-2.64 (m, 1H), 2.20-2.00 (m, 2H), 1.84-1.68 (m, 1H), 1.66-1.54 (m, 1H), 1.21 (t, 3H, J = 6.8 Hz), 1.13 (t, 3H, J = 6.8 Hz).
789		480	¹ H-NMR (500 MHz, 4d-MeOD) δ ppm 7.97 (d, 1H, J = 1.5 Hz), 7.86 (d, 1H, J = 5.0 Hz), 7.71 (d, 2H, J = 8.5 Hz), 7.32 (d, 2H, J = 8.5 Hz), 6.90 (d, 1H, J = 1.5 Hz), 6.02 (d, 1H, J = 5.5 Hz), 4.85-4.81 (m, 1H), 4.26-4.23 (m, 1H), 4.16-4.13 (m, 1H), 3.91-3.86 (m, 1H), 3.78-3.71 (m, 4H), 3.61-3.59 (m, 1H), 3.55-3.51 (m, 6H), 3.14 (td, 2H, J = 12.5, 2.5 Hz), 2.93 (tt, 1H, J = 12.5, 3.5 Hz), 2.12-2.09 (m, 2H), 1.98-1.89 (m, 2H).
790		480	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.38 (s, 1H), 8.16 (s, 1H), 7.91 (d, 1H, J = 5.0 Hz), 7.75 (d, 2H, J = 8.0 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.01 (s, 1H), 5.99 (d, 1H, J = 5.5 Hz), 4.62 (quintet, 1H, J = 5.5 Hz), 3.60-3.55 (m, 12H), 3.20-3.14 (m, 2H), 2.81-2.28 (m, 3H), 2.74-2.71 (m, 1H), 1.91-1.89 (m, 1H), 1.380-1.78 (m, 1H), 1.69-1.64 (m, 2H).
791		481	
792		481	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.96 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.71 (d, J = 8.2 Hz, 2H), 7.38 (d, J = 8.2 Hz, 2H), 6.90 (d, J = 1.9 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.3, 5.1 Hz, 1H), 4.90 (ddd, J = 7.3, 6.2, 0.9 Hz, 2H), 4.70-4.59 (m, 2H), 3.92 (t, J = 6.3 Hz, 1H), 3.87-3.59 (m, 7H), 3.53 (t, J = 5.2 Hz, 4H), 2.81-2.68 (m, 1H), 1.07 (dd, J = 15.0, 6.3 Hz, 6H).





#	Structure	LCMS (M+1)	NMR
793		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.61 (d, 1H, J = 1.6 Hz), 8.36 (s, 1H), 8.31 (d, 1H, J = 1.6 Hz), 7.97-7.93 (m, 2H), 7.82-7.79 (m, 1H), 7.19 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 6.4 Hz), 3.64-3.60 (m, 4H), 3.52-3.44 (m, 4H), 3.23-3.19 (m, 2H), 3.14-3.04 (m, 2H), 3.00-2.95 (m, 1H), 2.77-2.70 (m, 1H), 2.20-2.02 (m, 2H), 1.82-1.61 (m, 2H), 1.14 (t, 3H, J = 7.2 Hz).
794		484	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.61 (d, 1H, J = 1.6 Hz), 8.36 (s, 1H), 8.31 (d, 1H, J = 1.6 Hz), 7.97-7.93 (m, 2H), 7.82-7.79 (m, 1H), 7.19 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 6.4 Hz), 3.64-3.60 (m, 4H), 3.52-3.44 (m, 4H), 3.24-3.19 (m, 2H), 3.12-3.03 (m, 2H), 3.00-2.94 (m, 1H), 2.77-2.70 (m, 1H), 2.20-2.02 (m, 2H), 1.82-1.60 (m, 2H), 1.14 (t, 3H, J = 7.2 Hz).
795		489	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.93 (d, J = 1.8 Hz, 1H), 7.83 (d, J = 5.5 Hz, 1H), 7.73 – 7.56 (m, 2H), 7.30 – 7.19 (m, 2H), 6.85 (d, J = 1.8 Hz, 1H), 5.99 (d, J = 5.5 Hz, 1H), 4.17 – 4.00 (m, 1H), 3.69 (s, 4H), 3.48 (d, J = 7.0 Hz, 4H), 3.19 – 2.99 (m, 3H), 2.82 (p, J = 6.6 Hz, 1H), 2.68 – 2.47 (m, 1H), 2.40 (t, J = 11.6 Hz, 2H), 1.98 – 1.67 (m, 4H), 1.14 (d, J = 6.6 Hz, 6H), 0.77 – 0.64 (m, 4H).
796		465	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 1.5 Hz), 8.32 (s, 1H), 8.30 (d, 1H, J = 1.5 Hz), 7.96-7.94 (m, 2H), 7.79 (dd, 1H, J = 8.5, 2.0 Hz), 7.18 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 4.09 (q, 2H, J = 6.5 Hz), 3.62-3.61 (m, 4H), 3.49-3.48 (m, 4H), 3.16-3.13 (m, 1H), 3.00 (s, 3H), 2.95-2.92 (m, 2H), 2.70-2.65 (m, 1H), 2.18-2.15 (m, 1H), 2.07-2.02 (m, 1H), 1.75-1.73 (m, 1H), 1.60-1.57 (m, 1H), 1.22 (t, 3H, J = 6.5 Hz).





#	Structure	LCMS (M+1)	NMR
797		492	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.61 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.97-7.92 (m, 2H), 7.81 (dd, 1H, J = 8.4, 2.4 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.33 (d, 1H, J = 7.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 3.80-3.78 (m, 1H), 3.54-3.53 (m, 4H), 3.46-3.45 (m, 4H), 3.25-3.18 (m, 2H), 3.17-3.07 (m, 2H), 3.05-2.97 (m, 2H), 2.70-2.63 (m, 1H), 2.10-2.01 (m, 2H), 1.76-1.75 (m, 1H), 1.74-1.73 (m, 1H), 1.14 (t, 3H, J = 6.8 Hz), 1.08 (d, 6H, J = 6.8 Hz).
798		492	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 1.6 Hz), 8.30-8.25 (m, 2H), 8.00-7.90 (m, 2H), 7.79 (dd, 1H, J = 8.4, 2.0 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.30 (d, 1H, J = 8.4 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.84-3.70 (m, 1H), 3.56-3.50 (m, 4H), 3.48-3.40 (m, 4H), 3.26-3.16 (m, 1H), 3.14-3.04 (m, 2H), 3.02-2.86 (m, 2H), 2.72-2.60 (m, 1H), 2.18-1.94 (m, 2H), 1.80-1.64 (m, 1H), 1.62-1.50 (m, 1H), 1.13 (t, 3H, J = 6.8 Hz), 1.07 (d, 6H, J = 6.8 Hz).
799		493	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.21 (s, 1H), 7.93 (d, 1H, J = 5.2 Hz), 7.83 (d, 2H, J = 8.4 Hz), 7.21 (d, 2H, J = 8.4 Hz), 7.06 (s, 1H), 6.01 (d, 1H, J = 5.6 Hz), 5.35 (quintet, 1H, J = 5.2 Hz), 4.80 (t, 2H, J = 6.8 Hz), 4.55-4.52 (m, 2H), 3.75-3.70 (m, 4H), 3.70-3.65 (m, 2H), 3.65-3.60 (m, 2H), 3.50-3.49 (m, 4H), 2.91 (s, 3H), 1.99-1.94 (m, 4H).
800		493	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.61 (d, 1H, J = 1.5 Hz), 8.32 (d, 1H, J = 2.0 Hz), 7.99-7.95 (m, 2H), 7.81 (d, 1H, J = 9.5 Hz), 7.20 (d, 1H, J = 1.5 Hz), 6.02 (d, 1H, J = 4.5 Hz), 5.34 (quintet, 1H, J = 5.5 Hz), 4.80 (t, 2H, J = 7.5 Hz), 4.53 (dd, 2H, J = 7.5, 5.5 Hz), 3.71-3.65 (m, 2H), 3.65-3.60 (m, 2H), 3.53-3.51 (m, 4H), 3.34-3.31 (m, 1H), 3.13-3.06 (m, 2H), 3.01 (s, 3H), 2.82-2.77 (m, 1H), 2.24-2.21 (m, 1H), 2.11-2.06 (m, 1H), 1.83-1.80 (m, 1H), 1.72-1.69 (m, 1H).





#	Structure	LCMS (M+1)	NMR
801		493	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.2 Hz), 7.94 (d, 1H, J = 5.6 Hz), 7.92 (d, 1H, J = 8.8 Hz), 7.77 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 5.6 Hz), 5.40-5.25 (m, 1H), 4.78 (t, 2H, J = 7.2 Hz), 4.55-4.50 (m, 2H), 3.80-3.72 (m, 2H), 3.72-3.65 (m, 2H), 3.54-3.46 (m, 4H), 2.97 (s, 3H), 2.96-2.91 (m, 1H), 2.90-2.82 (m, 1H), 2.80-2.72 (m, 1H), 2.56-2.51 (m, 1H), 2.15-2.05 (m, 1H), 2.05-1.95 (m, 1H), 1.70-1.55 (m, 1H), 1.50-1.35 (m, 1H).
802		494	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.59 – 8.47 (m, 1H), 8.20 (d, J = 1.8 Hz, 1H), 8.00 – 7.77 (m, 3H), 7.16 (d, J = 1.8 Hz, 1H), 6.02 (d, J = 5.5 Hz, 1H), 4.18 (q, J = 7.1 Hz, 2H), 3.73 (d, J = 5.4 Hz, 4H), 3.54 (dd, J = 6.6, 3.9 Hz, 4H), 3.28 – 3.14 (m, 1H), 3.00 (s, 3H), 2.94 – 2.82 (m, 1H), 2.25 – 2.05 (m, 2H), 1.91 (td, J = 13.1, 4.3 Hz, 1H), 1.66 (d, J = 14.2 Hz, 1H), 1.40 (s, 3H), 1.29 (t, J = 7.1 Hz, 4H), 1.14 (s, 3H).
803		494	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.37 – 8.28 (m, 1H), 7.95 (dd, J = 3.6, 1.8 Hz, 1H), 7.85 (dd, J = 5.4, 3.6 Hz, 1H), 7.76 – 7.62 (m, 2H), 7.51 – 7.33 (m, 1H), 7.28 (d, J = 7.9 Hz, 2H), 6.90 (dd, J = 3.9, 1.8 Hz, 1H), 6.88 – 6.68 (m, 1H), 6.10 – 5.98 (m, 1H), 3.97 (s, 1H), 3.80 (d, J = 18.0 Hz, 1H), 3.62 (s, 4H), 3.14 – 3.04 (m, 3H), 2.88 (d, J = 15.9 Hz, 1H), 2.60 (t, J = 12.2 Hz, 1H), 2.43 (d, J = 12.3 Hz, 2H), 2.00 – 1.73 (m, 5H), 1.28 (s, 1H), 1.16 (d, J = 6.5 Hz, 6H).
804		495	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.97 (d, J = 1.8 Hz, 1H), 7.84 (d, J = 5.4 Hz, 1H), 7.76 – 7.64 (m, 2H), 7.59 – 7.47 (m, 2H), 6.90 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.5 Hz, 1H), 5.41 (tt, J = 6.2, 5.1 Hz, 1H), 4.90 (ddd, J = 7.4, 6.3, 1.0 Hz, 2H), 4.66 (ddd, J = 7.5, 5.1, 1.0 Hz, 2H), 3.76 (d, J = 36.2 Hz, 5H), 3.61 (d, J = 13.2 Hz, 1H), 3.57 – 3.48 (m, 4H), 2.82 – 2.73 (m, 1H), 1.57 (s, 3H), 1.03 (d, J = 6.3 Hz, 3H), 0.91 (d, J = 6.4 Hz, 3H).





#	Structure	LCMS (M+1)	NMR
805		495	
806		495	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.30-8.28 (m, 2H), 7.92 (d, 1H, J = 5.6 Hz), 7.72 (s, 1H), 7.69-7.67 (m, 1H), 7.39 (t, 1H, J = 7.2 Hz), 7.12 (d, 1H, J = 2.0 Hz), 6.32 (d, 1H, J = 7.5 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.80-3.76 (m, 1H), 3.54-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.26-3.22 (m, 1H), 3.05 (s, 3H), 3.05-2.96 (m, 2H), 2.68-2.64 (m, 1H), 2.24-2.20 (m, 1H), 2.09-2.04 (m, 1H), 1.77-1.75 (m, 1H), 1.59-1.56 (m, 1H), 1.09-1.03 (d, 6H, J = 6.8 Hz).
807		495	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.30-8.28 (m, 2H), 7.92 (d, 1H, J = 5.5 Hz), 7.72 (s, 1H), 7.68 (d, 1H, J = 7.0 Hz), 7.38 (t, 1H, J = 7.5 Hz), 7.12 (d, 1H, J = 1.5 Hz), 6.32 (d, 1H, J = 7.5 Hz), 5.99 (d, 1H, J = 5.5 Hz), 3.80-3.76 (m, 1H), 3.54-3.52 (m, 4H), 3.46-3.44 (m, 4H), 3.26-3.22 (m, 2H), 3.05 (s, 3H), 3.05-2.96 (m, 2H), 2.68-2.64 (m, 1H), 2.24-2.20 (m, 1H), 2.09-2.04 (m, 1H), 1.77-1.75 (m, 1H), 1.59-1.56 (m, 1H), 1.09-1.03 (d, 6H, J = 6.5 Hz).
808		498	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (d, 1H, J = 1.6 Hz), 7.92 (d, 1H, J = 5.6 Hz), 7.85 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.09 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.00-3.82 (m, 2H), 3.82-3.65 (m, 2H), 3.57-3.52 (m, 2H), 3.52-3.47 (m, 2H), 2.87-2.80 (m, 2H), 2.50-2.25 (m, 4H), 2.10-1.95 (m, 1H), 1.95-1.75 (m, 2H), 0.82-0.74 (m, 4H).





#	Structure	LCMS (M+1)	NMR
817		506	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.54 (t, J = 1.6 Hz, 1H), 8.20 (d, J = 1.8 Hz, 1H), 7.88 (dd, J = 3.5, 1.8 Hz, 3H), 7.17 (d, J = 1.8 Hz, 1H), 6.00 (d, J = 5.6 Hz, 1H), 3.93 (hept, J = 6.4 Hz, 1H), 3.65 (dd, J = 6.7, 3.5 Hz, 4H), 3.60 – 3.48 (m, 5H), 3.01 (s, 4H), 2.31 – 2.11 (m, 2H), 2.06 – 1.87 (m, 1H), 1.73 (d, J = 14.4 Hz, 1H), 1.47 (s, 3H), 1.22 (s, 3H), 1.17 (d, J = 6.6 Hz, 6H).
818		507	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 2.0 Hz), 8.35-8.25 (m, 2H), 8.00-7.90 (m, 2H), 7.80 (dd, 1H, J = 8.4, 2.4 Hz), 7.19 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 5.33 (quintet, 1H, J = 5.6 Hz), 4.78 (t, 2H, J = 7.2 Hz), 4.56-4.50 (m, 2H), 3.70-3.65 (m, 2H), 3.65-3.59 (m, 2H), 3.54-3.46 (m, 4H), 3.26-3.14 (m, 2H), 3.12-23.00 (m, 2H), 2.98-2.90 (m, 1H), 2.78-2.64 (m, 1H), 2.20-2.11 (m, 1H), 2.10-2.00 (m, 1H), 1.84-1.70 (m, 1H), 1.66-1.56 (m, 1H), 1.13 (t, 3H, J = 7.2 Hz).
819		507	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.6 Hz), 8.30 (d, 1H, J = 2.0 Hz), 7.96-7.91 (m, 2H), 7.80 (dd, 1H, J = 8.4, 2.4 Hz), 7.19 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.70-3.65 (m, 2H), 3.65-3.60 (m, 2H), 3.52-3.45 (m, 4H), 3.23-3.18 (m, 1H), 3.11-3.05 (m, 2H), 2.98-2.95 (m, 1H), 2.89-2.86 (m, 1H), 2.66-2.60 (m, 1H), 2.14-2.10 (m, 2H), 1.73-1.70 (m, 1H), 1.55-1.52 (m, 1H), 1.13 (t, 3H, J = 7.2 Hz).
820		507	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.5 Hz), 8.22 (s, 1H), 7.96-7.91 (m, 2H), 7.78 (dd, 1H, J = 2.0 Hz, 8.0 Hz), 7.18 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 6.0 Hz), 4.09 (q, 2H, J = 6.5 Hz), 3.70-3.55 (m, 4H), 3.55-3.40 (m, 4H), 2.93 (s, 3H), 2.90-2.80 (m, 1H), 2.80-2.70 (m, 2H), 2.70-2.55 (m, 2H), 2.15-2.03 (m, 2H), 2.03-1.90 (m, 2H), 1.22 (t, 3H, J = 6.5 Hz), 1.05 (d, 6H, J = 6.0 Hz).

#	Structure	LCMS (M+1)	NMR
821		508	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.5 Hz), 8.30 (d, 1H, J = 1.5 Hz), 7.96 (s, 1H), 7.95-7.93 (m, 1H), 7.80 (dd, 1H, J = 8.5, 2.0 Hz), 7.19 (d, 1H, J = 1.0 Hz), 6.01 (d, 1H, J = 5.5 Hz), 5.33 (quintet, 1H, J = 5.5 Hz), 4.79 (t, 2H, J = 7.0 Hz), 4.53 (dd, 2H, J = 7.0, 5.5 Hz), 3.77-3.74 (m, 2H), 3.74-3.70 (m, 4H), 3.62-3.58 (m, 2H), 3.54-3.50 (m, 2H), 3.09 (q, 2H, J = 7.0 Hz), 2.02-1.97 (m, 4H), 1.10 (t, 3H, J = 7.0 Hz).
822		512	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.39 – 8.24 (m, 2H), 8.03 (d, J = 1.8 Hz, 1H), 7.90 (d, J = 5.4 Hz, 1H), 7.79 – 7.67 (m, 2H), 7.52 – 7.41 (m, 2H), 7.41 – 7.23 (m, 2H), 6.96 (d, J = 1.8 Hz, 1H), 6.08 (d, J = 5.5 Hz, 1H), 3.96 (d, J = 16.6 Hz, 2H), 3.87 (d, J = 27.2 Hz, 2H), 3.70 – 3.57 (m, 5H), 3.57 – 3.37 (m, 4H), 3.15 – 2.97 (m, 2H), 2.95 – 2.75 (m, 1H), 2.13 (d, J = 13.9 Hz, 2H), 2.08 – 1.96 (m, 2H), 1.93 (s, 3H), 1.40 – 1.31 (m, 8H), 1.23 – 1.12 (m, 3H).
823		515	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.86 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 6.32 (d, 1H, J = 7.2 Hz), 6.00 (d, 1H, J = 5.6 Hz), 3.85-3.70 (m, 1H), 3.55-3.50 (m, 4H), 3.50-3.45 (m, 4H), 2.91-2.87 (m, 2H), 2.50-2.25 (m, 4H), 1.95-1.80 (m, 2H), 1.08 (d, 6H, J = 7.2 Hz).
824		517	¹ H NMR (500 MHz, Methanol-d ₄) δ 8.01 (d, J = 1.8 Hz, 1H), 7.88 (d, J = 5.4 Hz, 1H), 7.79 (d, J = 8.0 Hz, 2H), 7.48 (d, J = 8.1 Hz, 2H), 6.96 (d, J = 1.9 Hz, 1H), 6.03 (d, J = 5.4 Hz, 1H), 5.51 (s, 0H), 4.06 (s, 2H), 3.87 (s, 2H), 3.74 (q, J = 6.6 Hz, 4H), 3.68 (d, J = 24.2 Hz, 2H), 3.58 (s, 2H), 3.23 (q, J = 7.6 Hz, 5H), 2.88 (s, 0H), 2.50 (s, 2H), 2.28 (s, 2H), 2.05 (dd, J = 8.9, 4.3 Hz, 1H), 1.25 (d, J = 6.5 Hz, 6H), 0.95 (q, J = 3.3 Hz, 2H), 0.91 – 0.86 (m, 2H).





#	Structure	LCMS (M+1)	NMR
825		518	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.19-5.14 (m, 1H), 3.83-3.76 (m, 2H), 3.76-3.68 (m, 2H), 3.65-3.55 (m, 4H), 3.50-3.40 (m, 4H), 2.80-2.75 (m, 2H), 2.72-2.65 (m, 2H), 2.20-2.05 (m, 3H), 2.00-1.88 (m, 1H), 1.86-1.76 (m, 1H), 1.75-1.68 (m, 1H), 1.59-1.49 (m, 1H), 1.49-1.40 (m, 1H), 1.00-0.94 (m, 6H).
826		518	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.89 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.26 (d, 2H, J = 8.4 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.97 (d, 1H, J = 5.6 Hz), 5.19-5.14 (m, 1H), 3.83-3.76 (m, 2H), 3.76-3.68 (m, 2H), 3.65-3.55 (m, 4H), 3.50-3.40 (m, 4H), 2.80-2.75 (m, 2H), 2.72-2.65 (m, 2H), 2.20-2.05 (m, 3H), 2.00-1.88 (m, 1H), 1.86-1.76 (m, 1H), 1.75-1.68 (m, 1H), 1.59-1.49 (m, 1H), 1.49-1.40 (m, 1H), 1.00-0.94 (m, 6H).
827		518	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.13 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 6.98 (d, 1H, J = 2.0 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.65 (d, 2H, J = 7.2 Hz), 4.41 (d, 2H, J = 7.2 Hz), 3.65-3.60 (m, 2H), 3.60-3.54 (m, 2H), 3.52-3.42 (m, 4H), 2.83-2.74 (m, 2H), 2.72-2.66 (m, 2H), 2.22-2.08 (m, 2H), 1.86-1.68 (m, 2H), 1.66 (s, 3H), 1.60-1.36 (m, 2H), 1.00-0.94 (m, 6H).
828		518	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.0 Hz), 7.90 (d, 1H, J = 5.5 Hz), 7.72 (d, 2H, J = 8.5 Hz), 7.27 (d, 2H, J = 8.0 Hz), 7.00-6.99 (m, 1H), 5.99 (d, 1H, J = 5.5 Hz), 5.40-5.30 (m, 0.25H), 5.05-4.95 (m, 0.25H), 4.95-4.85 (m, 0.75H), 4.74-4.61 (m, 1H), 4.60-4.55 (m, 0.75H), 4.45-4.35 (m, 1H), 3.65-3.60 (m, 2H), 3.60-3.55 (m, 2H), 3.52-3.45 (m, 4H), 2.90-2.80 (m, 2H), 2.80-2.70 (m, 2H), 2.25-2.10 (m, 2H), 1.90-1.80 (m, 1H), 1.80-1.70 (m, 1H), 1.6-1.50 (m, 1H), 1.50-1.40 (m, 1H), 1.38 (d, 2.25H, J = 6.5 Hz), 1.30 (d, 0.75H, J = 6.5 Hz), 1.05-0.90 (m, 6H).

#	Structure	LCMS (M+1)	NMR
829		519	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (s, 1H), 7.89 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 7.6 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.03 (s, 1H), 6.31 (s, 1H), 5.98 (d, 1H, J = 5.2 Hz), 3.81-3.76 (m, 1H), 3.53-3.50 (m, 4H), 3.45-3.42 (m, 4H), 2.89 (s, 3H), 2.71-2.67 (m, 1H), 2.60-2.58 (m, 2H), 2.48-2.46 (m, 2H), 1.99-1.96 (m, 2H), 1.87-1.81 (m, 2H), 1.08 (d, 6H, J = 6.4 Hz), 0.99 (d, 6H, J = 6.8 Hz).
830		521	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.32 (s, 1H), 8.19 (d, J = 1.8 Hz, 1H), 7.91 (d, J = 5.4 Hz, 1H), 7.80 (dd, J = 8.0, 5.9 Hz, 2H), 7.36 (d, J = 8.5 Hz, 2H), 7.03 (d, J = 1.9 Hz, 1H), 5.99 (d, J = 5.6 Hz, 1H), 4.88 – 4.76 (m, 1H), 3.90 – 3.74 (m, 2H), 3.60 – 3.53 (m, 2H), 3.46 (t, J = 5.2 Hz, 5H), 3.11 (d, J = 12.6 Hz, 2H), 2.94 (s, 0H), 2.79 (t, J = 11.9 Hz, 2H), 1.99 (t, J = 12.3 Hz, 2H).
831		520	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.30 (s, 1H), 8.18 (s, 1H), 7.91 (d, 1H, J = 5.0 Hz), 7.89 (d, 2H, J = 8.5 Hz), 7.40 (d, 2H, J = 8.5 Hz), 7.03 (s, 1H), 5.98 (d, 1H, J = 5.0 Hz), 4.09 (q, 2H, J = 7.0 Hz), 4.65-4.55 (m, 4H), 4.51-4.41 (m, 4H), 3.06 (q, 2H, J = 6.5 Hz), 2.82-2.77 (m, 1H), 2.72-2.68 (m, 2H), 2.63-2.56 (m, 2H), 2.03-1.98 (m, 2H), 1.93-1.88 (m, 2H), 1.22 (t, 3H, J = 6.5 Hz), 1.07 (t, 3H, J = 7.0 Hz), 1.03 (d, 6H, J = 7.0 Hz).
832		521	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.98 (d, J = 1.8 Hz, 1H), 7.86 (d, J = 5.4 Hz, 1H), 7.76 – 7.62 (m, 2H), 7.34 – 7.22 (m, 2H), 6.91 (d, J = 1.8 Hz, 1H), 6.01 (d, J = 5.4 Hz, 1H), 5.14 – 5.01 (m, 2H), 4.66 – 4.54 (m, 3H), 3.78 (s, 4H), 3.60 – 3.47 (m, 4H), 3.34 (s, 4H), 3.10 – 2.99 (m, 2H), 2.80 (h, J = 6.5 Hz, 1H), 2.62 (p, J = 1.9 Hz, 1H), 2.56 (dq, J = 11.9, 4.0 Hz, 1H), 2.38 (td, J = 11.8, 2.7 Hz, 2H), 1.96 – 1.68 (m, 4H), 1.14 (d, J = 6.6 Hz, 6H).

#	Structure	LCMS (M+1)	NMR
833		522	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.54 (t, J = 1.5 Hz, 1H), 8.20 (d, J = 1.8 Hz, 1H), 8.01-7.79 (m, 3H), 7.16 (d, J = 1.8 Hz, 1H), 6.02 (d, J = 5.5 Hz, 1H), 5.42 (tt, J = 6.2, 5.1 Hz, 1H), 4.90 (ddd, J = 7.3, 6.2, 0.9 Hz, 2H), 4.66 (ddd, J = 7.5, 5.1, 0.9 Hz, 2H), 3.77 (d, J = 35.0 Hz, 5H), 3.56 (t, J = 5.2 Hz, 4H), 3.48 (q, J = 7.0 Hz, 1H), 3.26–3.07 (m, 1H), 3.00 (s, 3H), 2.87 (dt, J = 13.2, 3.6 Hz, 1H), 2.22–2.00 (m, 3H), 1.96–1.78 (m, 1H), 1.65 (d, J = 14.2 Hz, 1H), 1.40 (s, 3H), 1.20–1.11 (m, 4H).
834		521	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.96-7.90 (m, 2H), 7.78 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 2.0 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.60-4.50 (m, 2H), 4.48-4.40 (m, 2H), 4.08 (q, 2H, J = 7.2 Hz), 3.66-3.56 (m, 4H), 3.52-3.38 (m, 5H), 2.91 (s, 3H), 2.60-2.52 (m, 2H), 2.20-2.10 (m, 2H), 2.05-1.85 (m, 4H), 1.21 (t, 3H, J = 6.8 Hz).
835		521	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.5 Hz), 8.28 (d, 1H, J = 2.0 Hz), 7.94-7.91 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.0 Hz), 7.17 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 4.09 (q, 2H, J = 7.0 Hz), 3.62-3.61 (m, 4H), 3.49-3.48 (m, 4H), 3.08 (q, 2H, J = 7.0 Hz), 2.69-2.68 (m, 1H), 2.62-2.60 (m, 2H), 2.53-2.51 (m, 2H), 2.01-1.98 (m, 2H), 1.91-1.89 (m, 2H), 1.22 (d, 3H, J = 7.0 Hz), 1.08 (t, 3H, J = 7.0 Hz), 1.00-0.99 (m, 6H).
836		522	¹ H-NMR (500 MHz, 4d-MeOD) δ ppm 7.95 (d, 1H, J = 1.5 Hz), 7.83 (d, 1H, J = 5.0 Hz), 7.65 (d, 2H, J = 8.5 Hz), 7.27 (d, 2H, J = 8.5 Hz), 6.86 (d, 1H, J = 1.5 Hz), 5.98 (d, 1H, J = 5.5 Hz), 4.85-4.81 (m, 0.5H), 4.26-4.13 (m, 1H), 3.90-3.86 (m, 0.5H), 3.78-3.71 (m, 6H), 3.61-3.60 (m, 1H), 3.52-3.49 (m, 4H), 3.06-3.03 (m, 2H), 2.81-2.76 (m, 1H), 2.57-2.52 (m, 1H), 2.38-2.32 (m, 2H), 1.89-1.87 (m, 2H), 1.85-1.79 (m, 2H), 1.13 (d, 6H, J = 6.5 Hz)





#	Structure	LCMS (M+1)	NMR
837		522	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.27 (s, 1H), 8.16 (d, 1H, J = 2.4 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.74 (d, 2H, J = 8.4 Hz), 7.29 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.2 Hz), 4.62 (quintet, 1H, J = 5.2 Hz), 3.57-3.47 (m, 12H, mixed with H ₂ O peak), 2.99-2.95 (m, 2H), 2.95-2.91 (m, 1H), 2.81-2.80 (m, 1H), 2.43-2.36 (m, 2H), 1.58-1.77 (m, 2H), 1.69-1.60 (m, 1H), 1.53-1.47 (m, 1H), 1.07-1.04 (m, 6H).
838		524	
839		527	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.6 Hz), 7.99 (d, 1H, J = 5.6 Hz), 7.92 (d, 1H, J = 8.4 Hz), 7.82 (d, 1H, J = 1.6 Hz), 7.75 (dd, 1H, J = 8.4, 2.0 Hz), 7.13 (d, 1H, J = 1.6 Hz), 6.03 (d, 1H, J = 5.6 Hz), 3.93-3.91 (m, 2H), 3.72-3.70 (m, 2H), 3.59-3.58 (m, 2H), 3.53-3.51 (m, 2H), 2.91 (s, 3H), 2.72-2.68 (m, 1H), 2.63-2.59 (m, 2H), 2.48-2.44 (m, 2H), 2.04-1.99 (m, 3H), 1.91-1.85 (m, 2H), 1.01 (d, 6H, J = 6.8 Hz), 0.79-0.73 (m, 4H).
840		529	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.15 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.72 (d, 2H, J = 8.4 Hz), 7.27 (d, 2H, J = 8.4 Hz), 7.00 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.01 (d, 2H, J = 8.8 Hz), 4.82 (d, 2H, J = 8.8 Hz), 3.73-3.68 (m, 2H), 3.66-3.60 (m, 2H), 3.55-3.45 (m, 4H), 2.85-2.75 (m, 2H), 2.74-2.64 (m, 2H), 2.22-2.08 (m, 2H), 1.86-1.77 (m, 1H), 1.76-1.66 (m, 1H), 1.60-1.40 (m, 2H), 1.00-0.90 (m, 6H).





#	Structure	LCMS (M+1)	NMR
841		530	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.93 (d, J = 1.7 Hz, 1H), 7.82 (d, J = 5.5 Hz, 1H), 7.71 – 7.58 (m, 2H), 7.35 – 7.21 (m, 2H), 6.87 (d, J = 1.8 Hz, 1H), 5.97 (d, J = 5.5 Hz, 1H), 4.79 (s, 4H), 4.22 (s, 4H), 3.63 – 3.56 (m, 4H), 3.52 (dd, J = 6.7, 3.5 Hz, 4H), 3.16 – 3.07 (m, 2H), 2.89 (d, J = 10.8 Hz, 1H), 2.60 (t, J = 12.0 Hz, 1H), 2.46 (t, J = 11.6 Hz, 2H), 1.93 (d, J = 13.0 Hz, 2H), 1.88 – 1.74 (m, 2H), 1.16 (d, J = 6.6 Hz, 6H).
842		530	¹ H NMR (400 MHz, Methanol-d ₄) δ 7.93 (d, J = 1.8 Hz, 1H), 7.83 (dd, J = 5.4, 2.9 Hz, 1H), 7.65 (d, J = 8.0 Hz, 2H), 7.27 (d, J = 8.2 Hz, 2H), 6.87 (dd, J = 3.8, 1.8 Hz, 1H), 5.98 (dd, J = 10.4, 5.5 Hz, 1H), 5.46 (d, J = 6.7 Hz, 1H), 4.63 (d, J = 6.8 Hz, 1H), 4.10 (t, J = 7.3 Hz, 1H), 3.68 – 3.60 (m, 1H), 3.60 – 3.45 (m, 4H), 3.19 – 3.05 (m, 2H), 2.64 – 2.40 (m, 4H), 1.99 – 1.70 (m, 4H), 1.17 (d, J = 6.5 Hz, 6H).
843		530	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.6 Hz), 7.93 (d, 1H, J = 5.2 Hz), 7.85 (d, 2H, J = 8.4 Hz), 7.53 (d, 2H, J = 8.4 Hz), 7.06 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.2 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.72-3.67 (m, 2H), 3.62-3.56 (m, 2H), 3.52-3.47 (m, 4H), 2.87-2.80 (m, 2H), 2.50-2.25 (m, 4H), 1.95-1.80 (m, 2H).
844		533	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (s, 0.26 H), 8.18 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 4.2 Hz), 7.80 (d, 2H, J = 8.4 Hz), 7.40 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.2 Hz), 6.32 (d, 1H, J = 4.0 Hz), 5.98 (d, 1H, J = 4.2 Hz), 4.56-4.53 (m, 2H), 4.46-4.43 (m, 2H), 3.80-3.76 (m, 1H), 3.53-3.51 (m, 4H), 3.47-3.43 (m, 4H), 3.43-3.33 (m, 1H), 2.89 (s, 3H), 2.53-2.49 (m, 2H), 2.18-2.13 (m, 2H), 2.01-1.92 (m, 4H), 1.08 (d, 6H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
845		535	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 2.0 Hz), 8.30 (d, 1H, J = 1.6 Hz), 7.96-7.93 (m, 1H), 7.93 (d, 1H, J = 5.6 Hz), 7.81 (dd, 1H, J = 8.4, 2.4 Hz), 7.19 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.56 (t, 2H, J = 6.4 Hz), 4.45 (t, 2H, J = 6.4 Hz), 4.10 (q, 2H, J = 7.2 Hz), 3.68-3.58 (m, 4H), 3.52-3.43 (m, 4H), 3.43-3.40 (m, 1H), 3.08 (q, 2H, J = 6.8 Hz), 2.60-2.52 (m, 2H), 2.24-2.12 (m, 2H), 2.06-1.86 (m, 4H), 1.23 (t, 3H, J = 7.2 Hz), 1.08 (t, 3H, J = 6.8 Hz).
846		535	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.31-8.26 (m, 2H), 7.96-7.91 (m, 2H), 7.78 (dd, 1H, J = 2.0 Hz, 8.0 Hz), 7.18 (d, 1H, J = 1.5 Hz), 6.00 (d, 1H, J = 5.5 Hz), 5.40-5.30 (m, 1H), 4.82-4.75 (m, 2H), 4.58-4.50 (m, 2H), 3.80-3.55 (m, 4H), 3.55-3.45 (m, 4H), 2.93 (s, 3H), 2.85-2.75 (m, 1H), 2.75-2.65 (m, 2H), 2.65-2.55 (m, 2H), 2.12-1.90 (m, 4H), 1.04 (d, 6H, J = 6.5 Hz).
847		536	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.13 (s, 1H), 7.90 (d, 1H, J = 5.5 Hz), 7.72 (d, 2H, J = 8.0 Hz), 7.24 (d, 2H, J = 8.0 Hz), 6.99 (s, 1H), 5.99 (d, 1H, J = 6.0 Hz), 4.78-4.46 (m, 1H), 3.75-3.65 (m, 4H), 3.65-3.58 (m, 2H), 3.51-3.42 (m, 4H), 3.08-3.00 (m, 2H), 2.93-2.90 (m, 1H), 2.56-2.51 (m, 1H), 2.49-2.42 (m, 2H), 1.83-1.79 (m, 2H), 1.79-1.72 (m, 2H), 1.08 (d, 6H, J = 7.0 Hz).
848		538	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.5 Hz), 7.91 (d, 1H, J = 5.5 Hz), 7.75 (d, 2H, J = 8.5 Hz), 7.34 (d, 2H, J = 8.5 Hz), 7.05 (d, 1H, J = 1.5 Hz), 6.49 (t, 1H, J = 76.5 Hz), 5.98 (d, 1H, J = 5.5 Hz), 4.35-4.29 (m, 1H), 3.93-3.91 (m, 2H), 3.72-3.70 (m, 2H), 3.54-3.50 (m, 2H), 3.50-3.46 (m, 2H), 2.88-2.80 (m, 2H), 2.80-2.72 (m, 2H), 2.31 (t, 2H, J = 11.0 Hz), 2.12-2.10 (m, 1H), 2.04-2.00 (m, 1H), 1.68-1.63 (m, 1H), 0.98-0.95 (m, 6H), 0.79-0.74 (m, 4H).





#	Structure	LCMS (M+1)	NMR
849		538	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 2.0 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.34 (d, 2H, J = 8.4 Hz), 7.04 (d, 1H, J = 1.6 Hz), 6.49 (t, 1H, J = 76.4 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.35-4.29 (m, 1H), 3.93-3.91 (m, 2H), 3.72-3.70 (m, 2H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 2.86-2.79 (m, 2H), 2.79-2.72 (m, 2H), 2.31 (t, 2H, J = 11.2 Hz), 2.12-2.10 (m, 1H), 2.04-2.00 (m, 1H), 1.68-1.63 (m, 1H), 0.98-0.94 (m, 6H), 0.80-0.74 (m, 4H).
850		538	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 1.6 Hz), 8.29 (d, 1H, J = 1.2 Hz), 8.14 (s, 1H), 7.96-7.92 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.2 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.60-4.43 (m, 2H), 3.81-3.77 (m, 1H), 3.55-3.51 (m, 4H), 3.50-3.45 (m, 4H), 3.08-3.00 (m, 1H), 2.93 (s, 3H), 2.80-2.72 (m, 4H), 2.11-2.00 (m, 2H), 2.00-1.92 (m, 2H), 1.15-1.05 (m, 9H).
851		542	¹ H NMR (400 MHz, Methanol-d ₄) δ 8.54 (s, 3H), 7.96 (d, J = 1.8 Hz, 1H), 7.85 (d, J = 5.4 Hz, 1H), 7.70 (d, J = 8.2 Hz, 2H), 7.56 – 7.43 (m, 1H), 7.30 (d, J = 8.1 Hz, 2H), 7.22 – 7.07 (m, 3H), 6.91 (d, J = 1.8 Hz, 1H), 6.02 (d, J = 5.6 Hz, 1H), 3.90 – 3.75 (m, 4H), 3.68 – 3.58 (m, 4H), 3.50 (d, J = 13.0 Hz, 4H), 3.18 – 3.02 (m, 2H), 2.88 (s, 1H), 2.15 (d, J = 14.4 Hz, 3H), 2.00 (t, J = 13.1 Hz, 2H), 1.37 (d, J = 6.6 Hz, 6H).
852		544	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.23 (d, 1H, J = 2.0 Hz), 7.99 (d, 1H, J = 5.2 Hz), 7.92 (d, 1H, J = 8.8 Hz), 7.82 (d, 1H, J = 1.6 Hz), 7.75 (dd, 1H, J = 8.8, 2.0 Hz), 7.12 (d, 1H, J = 1.6 Hz), 6.34 (d, 1H, J = 7.6 Hz), 6.05 (d, 1H, J = 5.6 Hz), 3.81-3.75 (m, 1H), 3.52-3.50 (m, 4H), 3.46-3.44 (m, 4H), 2.91 (s, 3H), 2.72-2.68 (m, 1H), 2.63-2.59 (m, 2H), 2.48-2.44 (m, 2H), 2.01-1.97 (m, 2H), 1.91-1.85 (m, 2H), 1.08 (d, 6H, J = 6.8 Hz), 1.00 (d, 6H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
853		545	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.14 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.71 (d, 2H, J = 7.8 Hz), 7.27 (d, 2H, J = 8.0 Hz), 6.99 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 5.11-5.05 (m, 1H), 4.43 (dd, 1H, J = 9.6, 7.2 Hz), 4.14 (dd, 1H, J = 10.4, 7.2 Hz), 4.09 (dd, 1H, J = 9.6, 4.0 Hz), 3.78 (dd, 1H, J = 10.4, 4.0 Hz), 3.72-3.65 (m, 2H), 3.65-3.58 (m, 2H), 3.51-3.45 (m, 4H), 2.83-2.76 (m, 2H), 2.73-2.65 (m, 2H), 2.22-2.08 (m, 2H), 1.84-1.77 (m, 1H), 1.77 (s, 3H), 1.76-1.71 (m, 1H), 1.60-1.50 (m, 1H), 1.49-1.39 (m, 1H), 1.00-0.92 (m, 6H).
854		546	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.5 Hz), 8.21 (s, 1H), 7.95-7.92 (m, 2H), 7.79 (dd, 1H, J = 8.5, 2.5 Hz), 7.19 (d, 1H, J = 1.5 Hz), 6.01 (d, 1H, J = 5.0 Hz), 4.24 (t, 2H, J = 6.0 Hz), 3.68-3.64 (m, 4H), 3.53-3.48 (m, 4H), 3.09 (q, 2H, J = 7.0 Hz), 2.92 (t, 2H, J = 6.0 Hz), 2.83-2.82 (m, 1H), 2.74-2.72 (m, 2H), 2.64-2.60 (m, 2H), 2.06-2.04 (m, 2H), 1.98-1.95 (m, 2H), 1.08 (t, 3H, J = 7.0 Hz), 1.05 (d, 6H, J = 7.0 Hz).
855		547	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 1.6 Hz), 8.29 (d, 1H, J = 1.6 Hz), 8.17 (s, 1H), 7.95-7.90 (m, 2H), 7.80-7.76 (m, 1H), 7.18 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.36-5.31 (m, 1H), 4.82-4.76 (m, 2H), 4.55-4.51 (m, 2H), 3.80-3.55 (m, 4H), 3.51-3.48 (m, 4H), 3.10 (q, 2H, J = 7.2 Hz), 2.79-2.75 (m, 2H), 2.61-2.55 (m, 2H), 2.00-1.96 (m, 2H), 1.88-1.84 (m, 2H), 1.67-1.65 (m, 1H), 1.10 (t, 3H, J = 7.2 Hz), 0.44-0.41 (m, 2H), 0.31-0.29 (m, 2H).
856		547	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.19 (s, 1H), 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.78 (d, 2H, J = 8.4 Hz), 7.40 (d, 2H, J = 8.0 Hz), 7.02 (d, 1H, J = 1.2 Hz), 6.31 (d, 1H, J = 7.6 Hz), 5.97 (d, 1H, J = 5.2 Hz), 4.60-4.50 (m, 2H), 4.48-4.40 (m, 2H), 3.80-3.70 (m, 1H), 3.60-3.48 (m, 4H), 3.46-3.36 (m, 5H), 3.03 (q, 2H, J = 6.8 Hz), 2.60-2.50 (m, 2H), 2.22-2.10 (m, 2H), 2.00-1.80 (m, 4H), 1.12-1.00 (m, 9H).





#	Structure	LCMS (M+1)	NMR
857		548	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.30 (s, 1H), 8.18 (d, 1H, J = 1.5 Hz), 7.91 (d, 1H, J = 5.5 Hz), 7.80 (d, 2H, J = 8.5 Hz), 7.40 (d, 2H, J = 8.5 Hz), 7.04 (d, 1H, J = 1.5 Hz), 5.99 (d, 1H, J = 5.5 Hz), 5.34 (quintet, 1H, J = 5.5 Hz), 4.79 (t, 2H, J = 7.0 Hz), 4.53 (dd, 2H, d, 2H, J = 7.0, 5.5 Hz), 3.80-3.65 (m, 2H), 3.65-3.50 (m, 2H), 3.50-3.40 (m, 4H), 3.06 (q, 2H, J = 6.5 Hz), 2.90-2.80 (m, 1H), 2.80-2.70 (m, 2H), 2.68-2.61 (m, 2H), 2.03-2.00 (m, 2H), 1.98-1.92 (m, 2H), 1.09-1.05 (m, 9H).
858		548	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.27 (d, 1H, J = 1.2 Hz), 8.16 (s, 1H), 7.92 (d, 1H, J = 5.6 Hz), 7.91-7.88 (m, 1H), 7.78 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.30 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 4.54 (t, 2H, J = 6.4 Hz), 4.43 (t, 2H, J = 6.4 Hz), 3.84-3.72 (m, 1H), 3.56-3.49 (m, 4H), 3.48-3.43 (m, 4H), 3.44-3.40 (m, 1H), 3.06 (q, 2H, J = 6.8 Hz), 2.56-2.51 (m, 2H), 2.22-2.12 (m, 2H), 2.06-1.86 (m, 4H), 1.08 (d, 6H, J = 6.4 Hz), 1.06 (t, 3H, J = 6.8 Hz).
859		552	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.76 (d, 2H, J = 8.4 Hz), 7.40 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 6.54 (t, 1H, J = 75.6 Hz), 5.99 (d, 1H, J = 5.6 Hz), 3.92 (m, 2H), 3.83 (s, 2H), 3.71 (m, 2H), 3.57-3.50 (m, 2H), 3.50-3.43 (m, 2H), 2.60-2.57 (m, 3H), 2.30-2.25 (m, 2H), 2.15-2.11 (m, 2H), 2.14-1.99 (m, 1H), 1.88-1.83 (m, 2H), 0.90 (d, 6H, J = 6.8 Hz), 0.79-0.77 (m, 4H).
860		552	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.94-7.90 (m, 2H), 7.79 (dd, 1H, J = 8.4, 2.4 Hz), 7.17 (d, 1H, J = 2.0 Hz), 6.31 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.56-4.32 (m, 2H), 3.82-3.76 (m, 1H), 3.55-3.50 (m, 4H), 3.48-3.43 (m, 4H), 3.08 (q, 2H, J = 7.2 Hz), 2.93-2.84 (m, 1H), 2.71-2.63 (m, 4H), 2.03-1.95 (m, 2H), 1.95-1.85 (m, 2H), 1.09-1.06 (m, 9H), 1.18 (d, 3H, J = 5.6 Hz).





#	Structure	LCMS (M+1)	NMR
861		552	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.6 Hz), 8.28 (d, 1H, J = 1.6 Hz), 8.17 (s, 1H), 7.95-7.91 (m, 2H), 7.81-7.77 (m, 1H), 7.18 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.95-4.78 (m, 1H), 3.82-3.76 (m, 1H), 3.55-3.51 (m, 4H), 3.48-3.45 (m, 4H), 3.07 (q, 2H, J = 7.2 Hz), 2.80-2.65 (m, 2H), 2.64-2.50 (m, 2H), 2.50-2.45 (m, 2H), 2.08-1.93 (m, 4H), 1.27 (dd, 2H, J = 23.6 Hz, J = 6.4 Hz), 1.10-1.06 (m, 9H).
862		552	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.6 Hz), 8.28 (d, 1H, J = 1.6 Hz), 8.16 (s, 1H), 7.95-7.91 (m, 2H), 7.81-7.77 (m, 1H), 7.18 (d, 1H, J = 1.6 Hz), 6.31 (d, 1H, J = 7.6 Hz), 6.00 (d, 1H, J = 5.6 Hz), 4.97-4.78 (m, 1H), 3.82-3.76 (m, 1H), 3.55-3.51 (m, 4H), 3.48-3.45 (m, 4H), 3.08 (q, 2H, J = 7.2 Hz), 2.80-2.65 (m, 2H), 2.64-2.50 (m, 2H), 2.50-2.43 (m, 2H), 1.99-1.93 (m, 4H), 1.27 (dd, 2H, J = 23.6 Hz, J = 6.4 Hz), 1.10-1.06 (m, 9H).
863		553	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.30 (d, 1H, J = 2.0 Hz), 8.16 (s, 1H), 7.96-7.92 (m, 2H), 7.78 (dd, 1H, J = 8.4, 2.0 Hz), 7.19 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.2 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.80 (t, 2H, J = 7.2 Hz), 4.55 (dd, 2H, J = 7.2, 5.2 Hz), 4.60-4.41 (m, 2H), 3.73-3.68 (m, 2H), 3.68-3.60 (m, 2H), 3.55-3.50 (m, 4H), 3.10-2.95 (m, 1H), 2.93 (s, 3H), 2.85-2.70 (m, 4H), 2.10-2.00 (m, 2H), 2.00-1.92 (m, 2H), 1.06 (d, 3H, J = 6.4 Hz).
864		553	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.28 (s, 1H), 7.94-7.91 (m, 2H), 7.80-7.77 (m, 1H), 7.18 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 5.01-4.92 (m, 1H), 4.61-4.38 (m, 2H), 3.63-3.62 (m, 4H), 3.50-3.49 (m, 4H), 3.08 (q, 2H, J = 6.8 Hz), 2.70-2.54 (m, 4H), 2.44-2.33 (m, 1H), 2.01-1.98 (m, 2H), 1.91-1.85 (m, 2H), 1.22 (d, 3H, J = 6.4 Hz), 1.08 (t, 3H, J = 6.8 Hz), 1.00 (d, 3H, J = 6.8 Hz).





#	Structure	LCMS (M+1)	NMR
865		553	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.4 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.95-7.91 (m, 2H), 7.80-7.77 (m, 1H), 7.18 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.00-4.81 (m, 1H), 4.28-4.06 (m, 2H), 3.65-3.64 (m, 4H), 3.52-3.50 (m, 4H), 3.08 (q, 2H, J = 6.8 Hz), 2.73-2.61 (m, 5H), 2.03-1.91 (m, 4H), 1.31 (dd, 3H, J = 24.0, 6.8 Hz), 1.07 (t, 3H, J = 6.8 Hz), 0.99 (d, 6H, J = 6.4 Hz).
866		553	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.28 (s, 1H), 7.94-7.91 (m, 2H), 7.80-7.77 (m, 1H), 7.18 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 5.01-4.92 (m, 1H), 4.61-4.38 (m, 2H), 3.63-3.62 (m, 4H), 3.50-3.49 (m, 4H), 3.08 (q, 2H, J = 6.8 Hz), 2.70-2.54 (m, 4H), 2.44-2.33 (m, 1H), 2.01-1.98 (m, 2H), 1.91-1.85 (m, 2H), 1.22 (d, 3H, J = 6.4 Hz), 1.08 (t, 3H, J = 6.8 Hz), 1.00 (d, 3H, J = 6.8 Hz).
867		553	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.4 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.95-7.91 (m, 2H), 7.80-7.77 (m, 1H), 7.18 (d, 1H, J = 1.2 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.00-4.81 (m, 1H), 4.28-4.06 (m, 2H), 3.65-3.64 (m, 4H), 3.52-3.50 (m, 4H), 3.08 (q, 2H, J = 6.8 Hz), 2.73-2.61 (m, 5H), 2.03-1.91 (m, 4H), 1.31 (dd, 3H, J = 24.0, 6.8 Hz), 1.07 (t, 3H, J = 6.8 Hz), 0.99 (d, 6H, J = 6.4 Hz).
868		555	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 2.0 Hz), 7.90 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 6.49 (t, 1H, J = 75.2 Hz), 6.32 (s, 1H), 5.99 (d, 1H, J = 5.6 Hz), 4.35-4.29 (m, 1H), 3.82-3.74 (m, 1H), 3.53-3.51 (m, 4H), 3.46-3.43 (m, 4H), 2.89-2.80 (m, 2H), 2.80-2.74 (m, 2H), 2.31 (t, 2H, J = 10.8 Hz), 2.12-2.10 (m, 1H), 1.68-1.64 (m, 1H), 1.09 (d, 6H, J = 6.4 Hz), 0.98-0.94 (m, 6H).





#	Structure	LCMS (M+1)	NMR
869		555	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.16 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.33 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.6 Hz), 6.49 (t, 1H, J = 76.4 Hz), 6.32-6.30 (m, 1H), 5.98 (d, 1H, J = 5.2 Hz), 4.32 (td, 1H, J = 10.4, 4.4 Hz), 3.82-3.74 (m, 1H), 3.53-3.51 (m, 4H), 3.46-3.43 (m, 4H), 2.89-2.80 (m, 2H), 2.80-2.74 (m, 2H), 2.31 (t, 2H, J = 10.8 Hz), 2.12-2.10 (m, 1H), 1.68-1.64 (m, 1H), 1.09 (d, 6H, J = 6.4 Hz), 0.98-0.94 (m, 6H).
870		556	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.96-7.90 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.4 Hz), 7.18 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.58-4.50 (m, 2H), 4.48-4.40 (m, 2H), 3.72-3.60 (m, 3H), 3.58-3.52 (m, 2H), 3.50-3.38 (m, 5H), 3.30-3.20 (m, 1H), 2.91 (s, 3H), 2.68-2.57 (m, 2H), 2.56-2.50 (m, 4H), 2.20-2.10 (m, 2H), 2.05-1.85 (m, 4H).
871		556	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.30 (d, 1H, J = 2.0 Hz), 8.17 (s, 1H), 7.96-7.91 (m, 2H), 7.80 (dd, 1H, J = 2.5 Hz, 8.0 Hz), 7.19 (d, 1H, J = 1.5 Hz), 5.99 (d, 1H, J = 5.5 Hz), 3.75-3.60 (m, 3H), 3.60-3.52 (m, 2H), 3.52-3.40 (m, 4H), 3.35-3.20 (m, 3H), 3.15-3.05 (m, 2H), 3.00-2.88 (m, 1H), 2.88-2.75 (m, 2H), 2.75-2.55 (m, 4H), 2.15-1.90 (m, 4H), 1.15-1.00 (m, 9H).
872		557	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.0 Hz), 7.95-7.91 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.0 Hz), 7.19 (d, 1H, J = 1.0 Hz), 6.28 (tt, 1H, J = 54.5, 3.5 Hz), 6.01 (d, 1H, J = 5.0 Hz), 4.35 (td, 2H, J = 15.5, 3.5 Hz), 3.66-3.65 (m, 4H), 3.52-3.49 (m, 4H), 3.08 (q, 2H, J = 6.5 Hz), 2.70-2.67 (m, 1H), 2.62-2.60 (m, 2H), 2.53-2.51 (m, 2H), 2.01-1.98 (m, 2H), 1.91-1.87 (m, 2H), 1.08 (l, 3H, J = 6.5 Hz), 1.00 (d, 6H, J = 6.5 Hz).




#	Structure	LCMS (M+1)	NMR
873		559	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.24 (d, 1H, J = 1.6 Hz), 8.00 (d, 1H, J = 5.2 Hz), 7.92 (d, 1H, J = 8.4 Hz), 7.82 (d, 1H, J = 1.6 Hz), 7.75 (dd, 1H, J = 8.4, 2.0 Hz), 7.12 (d, 1H, J = 2.0 Hz), 6.07 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.69-3.67 (m, 2H), 3.59-3.57 (m, 2H), 3.53-3.51 (m, 4H), 2.91 (s, 3H), 2.72-2.68 (m, 1H), 2.63-2.59 (m, 2H), 2.48-2.44 (m, 2H), 2.01-1.97 (m, 2H), 1.91-1.85 (m, 2H), 1.01 (d, 6H, J = 6.4 Hz).
874		563	¹ H NMR (500 MHz, Methanol-d ₄) δ 8.51 (s, 1H), 8.00 (d, J = 1.8 Hz, 1H), 7.87 (d, J = 5.4 Hz, 1H), 7.78 (d, J = 8.0 Hz, 2H), 7.46 (d, J = 8.1 Hz, 2H), 6.92 (d, J = 1.9 Hz, 1H), 6.03 (d, J = 5.5 Hz, 1H), 4.99 – 4.92 (m, 1H), 3.88 – 3.63 (m, 8H), 3.55 (s, 5H), 3.44 (d, J = 15.6 Hz, 1H), 3.12 (d, J = 53.8 Hz, 2H), 2.97 – 2.67 (m, 2H), 2.23 (s, 2H), 1.35 (d, J = 6.6 Hz, 6H).
875		563	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.94 (d, 1H, J = 5.2 Hz), 7.93-7.90 (m, 1H), 7.79 (dd, 1H, J = 8.4, 2.4 Hz), 7.18 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.33 (quintet, 1H, J = 5.2 Hz), 4.78 (t, 2H, J = 7.2 Hz), 4.60-4.50 (m, 4H), 4.44 (t, 2H, J = 5.6 Hz), 3.73-3.66 (m, 2H), 3.66-3.58 (m, 2H), 3.54-3.47 (m, 4H), 3.46-3.36 (m, 1H), 3.06 (q, 2H, J = 6.8 Hz), 2.58-2.51 (m, 2H), 2.24-2.10 (m, 2H), 2.06-1.86 (m, 4H), 1.07 (t, 3H, J = 6.8 Hz).
876		563	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.5 Hz), 7.94-7.91 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.5 Hz), 7.18 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 6.0 Hz), 5.19-5.18 (m, 1H), 3.82-3.79 (m, 2H), 3.76-3.71 (m, 2H), 3.62-3.60 (m, 4H), 3.50-3.48 (m, 4H), 3.08 (q, 2H, J = 7.0 Hz), 2.70-2.68 (m, 1H), 2.62-2.60 (m, 2H), 2.54-2.51 (m, 2H), 2.16-2.12 (m, 1H), 2.01-1.92 (m, 3H), 1.92-1.86 (m, 2H), 1.08(t, 3H, J = 7.0 Hz), 1.00 (d, 6H, J = 7.0 Hz).

#	Structure	LCMS (M+1)	NMR
877		563	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.5 Hz), 7.94-7.91 (m, 2H), 7.78 (dd, 1H, J = 8.0, 2.5 Hz), 7.18 (d, 1H, J = 2.0 Hz), 6.00 (d, 1H, J = 6.0 Hz), 5.19-5.18 (m, 1H), 3.82-3.79 (m, 2H), 3.76-3.71 (m, 2H), 3.62-3.60 (m, 4H), 3.50-3.48 (m, 4H), 3.08 (q, 2H, J = 7.0 Hz), 2.70-2.68 (m, 1H), 2.62-2.60 (m, 2H), 2.54-2.51 (m, 2H), 2.16-2.12 (m, 1H), 2.01-1.92 (m, 3H), 1.92-1.86 (m, 2H), 1.08(t, 3H, J = 7.0 Hz), 1.00 (d, 6H, J = 7.0 Hz).
878		563	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (s, 1H), 8.29 (s, 1H), 7.94 (d, 1H, J = 5.2 Hz), 7.93-7.91 (m, 1H), 7.79 (d, 1H, J = 8.0 Hz), 7.18 (s, 1H), 6.00 (d, 1H, J = 5.2 Hz), 4.66 (d, 2H, J = 7.2 Hz), 4.42 (d, 2H, J = 7.2 Hz), 3.66-3.63 (m, 2H), 3.63-3.58 (m, 2H), 3.51-3.48 (m, 4H), 3.08 (q, 2H, J = 7.2 Hz), 2.80-2.70 (m, 2H), 2.70-2.60 (m, 2H), 2.02-1.99 (m, 2H), 1.92-1.90 (m, 2H), 1.67 (s, 3H), 1.08(t, 3H, J = 7.2 Hz), 1.01-0.99 (m, 6H).
879		563	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.61 (s, 1H), 8.31(d, 1H, J = 1.0 Hz), 7.98-7.94 (m, 2H), 7.81-7.78 (m, 1H), 7.21-7.18 (m, 1H), 6.02-6.00 (m, 1H), 5.40-5.30 (m, 0.25H), 5.05-4.95 (m, 0.25H), 4.95-4.85 (m, 0.75H), 4.80-4.70 (m, 1H), 4.65-4.58 (m, 0.75H), 4.45-4.30 (m, 1H), 3.75-3.40 (br., 2H), 3.70-3.62 (m, 2H), 3.62-3.55 (m, 2H), 3.55-3.48 (m, 4H), 3.30-3.25 (m, 1H), 3.11 (q, 2H, J = 7.0 Hz), 2.45-2.10 (m, 4H), 1.38 (d, 6H, J = 7.0 Hz), 1.30 (d, 3H, J = 6.0 Hz), 1.30-1.20 (m, 2H), 1.11 (t, 3H, J = 7.0 Hz).
880		567	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 1.6 Hz), 8.29 (d, 1H, J = 1.6 Hz), 7.95-7.92 (m, 2H), 7.80 (dd, 1H, J = 8.0, 2.4 Hz), 7.18 (d, 1H, J = 1.6 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.2 Hz), 4.97-4.90 (m, 1H), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 7.2, 5.2 Hz), 3.73-3.65 (m, 2H), 3.65-3.59 (m, 2H), 3.52-3.50 (m, 4H), 3.33-3.31 (m, 2H), 3.08 (q, 2H, J = 6.8 Hz), 2.75-2.66 (m, 2H), 2.50-2.49 (m, 2H), 1.91-1.75 (m, 4H), 1.27 (dd, 3H, J = 24.0, 6.4 Hz), 1.08 (t, 3H, J = 6.8 Hz).

#	Structure	LCMS (M+1)	NMR
881		567	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 1.6 Hz), 8.30 (d, 1H, J = 1.6 Hz), 8.22 (s, 1H), 7.96-7.91 (m, 2H), 7.79 (dd, 1H, J = 8.4, 2.4 Hz), 7.19 (d, 1H, J = 1.2 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.54 (dd, 2H, J = 7.2, 5.6 Hz), 4.50-4.33 (m, 2H), 3.75-3.65 (m, 2H), 3.65-3.55 (m, 2H), 3.55-3.48 (m, 4H), 3.08 (q, 2H, J = 7.2 Hz), 2.95-2.85 (m, 1H), 2.73-2.65 (m, 4H), 2.05-1.95 (m, 2H), 1.95-1.85 (m, 2H), 1.08 (t, 3H, J = 7.2 Hz), 1.26 (d, 3H, J = 6.8 Hz).
882		567	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 2.0 Hz), 8.30 (d, 1H, J = 1.6 Hz), 8.18 (s, 1H), 7.96-7.91 (m, 2H), 7.79 (dd, 1H, J = 8.0, 2.4 Hz), 7.19 (d, 1H, J = 1.2 Hz), 6.01 (d, 1H, J = 5.6 Hz), 5.33 (quintet, 1H, J = 5.2 Hz), 4.95-4.77 (m, 1H), 4.80 (t, 2H, J = 7.2 Hz), 4.54 (dd, 2H, J = 7.2, 5.2 Hz), 3.72-3.65 (m, 2H), 3.65-3.57 (m, 2H), 3.57-3.48 (m, 4H), 3.07 (q, 2H, J = 7.2 Hz), 2.80-2.71 (m, 2H), 2.70-2.50 (m, 2H), 2.50-2.48 (m, 2H), 2.02-1.90 (m, 4H), 1.27 (dd, 3H, J = 23.6 Hz, 6.4 Hz), 1.08 (t, 3H, J = 7.2 Hz).
883		568	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 6.69 (d, 1H, J = 76.4 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 3H), 3.82-3.65 (m, 3H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 2.89 (s, 3H), 2.88-2.82 (m, 1H), 2.75-2.53 (m, 4H), 2.08-1.95 (m, 3H), 1.92-1.85 (m, 2H), 1.01 (d, 3H, J = 6.8 Hz), 0.82-.072 (m, 4H).
884		568	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.05 (d, 1H, J = 1.6 Hz), 6.69 (d, 1H, J = 76.4 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.00-3.85 (m, 3H), 3.82-3.65 (m, 3H), 3.55-3.50 (m, 2H), 3.50-3.45 (m, 2H), 2.89 (s, 3H), 2.88-2.82 (m, 1H), 2.75-2.53 (m, 4H), 2.08-1.95 (m, 3H), 1.92-1.85 (m, 2H), 1.01 (d, 3H, J = 6.8 Hz), 0.82-.072 (m, 4H).

#	Structure	LCMS (M+1)	NMR
889		570	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.6 Hz), 7.91 (d, 1H, J = 5.2 Hz), 7.75 (d, 2H, J = 8.4 Hz), 7.34 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 6.49 (t, 1H, J = 76.4 Hz), 5.99 (d, 1H, J = 5.6 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 7.2 Hz), 4.53 (dd, 2H, J = 8.0, 5.6 Hz), 4.35-4.29 (m, 1H), 3.70-3.65 (m, 2H), 3.65-3.59 (m, 2H), 3.50-3.48 (m, 4H), 2.86-2.79 (m, 2H), 2.79-2.72 (m, 2H), 2.31 (t, 2H, J = 11.2 Hz), 2.12-2.10 (m, 1H), 1.68-1.62 (m, 1H), 0.98-0.94 (m, 6H).
890		570	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.60 (d, 1H, J = 1.6 Hz), 8.30 (d, 1H, J = 1.6 Hz), 8.17 (s, 1H), 7.95-7.91 (m, 2H), 7.81-7.78 (m, 1H), 7.19 (d, 1H, J = 1.6 Hz), 5.98 (d, 1H, J = 5.6 Hz), 4.55 (t, 2H, J = 6.4 Hz), 4.44 (t, 2H, J = 6.4 Hz), 3.71-3.65 (m, 3H), 3.57-3.54 (m, 2H), 3.49-3.46 (m, 4H), 3.45-3.42 (m, 1H), 3.31-3.26 (m, 1H), 3.07 (q, 2H, J = 7.2 Hz), 2.67-2.59 (m, 2H), 2.54-2.51 (m, 2H), 2.50-2.48 (m, 2H), 2.21-2.15 (m, 2H), 2.03-1.90 (m, 4H), 1.08 (t, 3H, J = 7.2 Hz).
891		584	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.18 (s, 1H), 7.92 (d, 1H, J = 5.5 Hz), 7.76 (d, 2H, J = 8.0 Hz), 7.40 (d, 2H, J = 8.0 Hz), 7.02 (s, 1H), 6.53 (t, 1H, J = 76.0 Hz), 6.00 (d, 1H, J = 5.5 Hz), 5.33 (quintet, 1H, J = 5.0 Hz), 4.79 (t, 2H, J = 7.0 Hz), 4.53 (dd, 2H, J = 7.0, 5.0 Hz), 3.83 (s, 2H), 3.75-3.70 (m, 2H), 3.70-3.65 (m, 2H), 3.52-3.47 (m, 4H), 2.60-2.56 (m, 3H), 2.32-2.28 (m, 2H), 2.15-2.13 (m, 2H), 1.90-1.87 (m, 2H), 0.91 (d, 6H, J = 7.0 Hz).
892		585	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 6.68 (t, 1H, J = 76.4 Hz), 6.30 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.91-3.87 (m, 1H), 3.80-3.73 (m, 2H), 3.53-3.48 (m, 4H), 3.46-3.41 (m, 4H), 2.89 (s, 3H), 2.88-2.83 (m, 1H), 2.75-2.53 (m, 4H), 2.03-1.93 (m, 2H), 1.90-1.82 (m, 2H), 1.06 (d, 6H, J = 8.4 Hz), 1.01 (d, 3H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
893		585	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.17 (d, 1H, J = 1.6 Hz), 7.90 (d, 1H, J = 5.6 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.01 (d, 1H, J = 1.2 Hz), 6.68 (t, 1H, J = 76.4 Hz), 6.30 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.91-3.87 (m, 1H), 3.80-3.73 (m, 2H), 3.53-3.48 (m, 4H), 3.46-3.41 (m, 4H), 2.89 (s, 3H), 2.88-2.83 (m, 1H), 2.75-2.53 (m, 4H), 2.03-1.93 (m, 2H), 1.90-1.82 (m, 2H), 1.06 (d, 6H, J = 8.4 Hz), 1.01 (d, 3H, J = 6.4 Hz).
894		586	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 1.6 Hz), 8.28 (d, 1H, J = 1.6 Hz), 7.93-7.91 (m, 2H), 7.77 (dd, 1H, J = 8.4, 2.0 Hz), 7.17 (d, 1H, J = 1.6 Hz), 6.68 (t, 1H, J = 76.4 Hz), 6.30 (d, 1H, J = 7.6 Hz), 5.99 (d, 1H, J = 5.2 Hz), 3.93-3.89 (m, 1H), 3.80-3.77 (m, 1H), 3.77-3.72 (m, 1H), 3.54-3.53 (m, 4H), 3.47-3.46 (m, 4H), 2.92 (s, 3H), 2.89-2.86 (m, 1H), 2.67-2.62 (m, 1H), 2.61-2.56 (m, 3H), 2.04-2.01 (m, 2H), 1.89-1.86 (m, 2H), 1.08 (d, 6H, J = 6.8 Hz), 1.02 (d, 3H, J = 6.4 Hz).
895		590	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.59 (d, 1H, J = 2.0 Hz), 8.28 (d, 1H, J = 1.5 Hz), 7.94 (d, 1H, J = 5.5 Hz), 7.91 (d, 1H, J = 8.0 Hz), 7.78 (dd, 1H, J = 8.0, 2.0 Hz), 7.18 (d, 1H, J = 1.5 Hz), 6.01 (d, 1H, J = 5.5 Hz), 5.10-5.08 (m, 1H), 4.43 (dd, 1H, J = 9.5, 7.0 Hz), 4.14 (dd, 1H, J = 10.5, 7.0 Hz), 4.09 (dd, 1H, J = 9.5, 4.0 Hz), 3.78 (dd, 1H, J = 10.5, 4.0 Hz), 3.71-3.65 (m, 2H), 3.65-3.59 (m, 2H), 3.52-3.51 (m, 4H), 3.08 (q, 2H, J = 7.0 Hz), 2.70-2.68 (m, 1H), 2.62-2.60 (m, 2H), 2.54-2.51 (m, 2H), 2.01-1.98 (m, 2H), 1.91-1.89 (m, 2H), 1.78 (s, 3H), 1.08 (t, 3H, J = 7.0 Hz), 1.00 (d, 6H, J = 7.0 Hz).
896		600	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 6.69 (d, 1H, J = 76.4 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 6.8 Hz), 4.53 (dd, 2H, J = 6.8, 5.6 Hz), 3.92-3.87 (m, 1H), 3.77-3.72 (m, 1H), 3.75-3.69 (m, 2H), 3.69-3.66 (m, 2H), 3.62-3.58 (m, 4H), 2.89 (s, 3H), 2.88-2.83 (m, 1H), 2.75-2.55 (m, 4H), 2.03-1.96 (m, 2H), 1.90-1.81 (m, 2H), 1.01 (d, 3H, J = 6.4 Hz).

#	Structure	LCMS (M+1)	NMR
897		600	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.18 (d, 1H, J = 1.2 Hz), 7.92 (d, 1H, J = 5.2 Hz), 7.79 (d, 2H, J = 8.4 Hz), 7.39 (d, 2H, J = 8.4 Hz), 7.03 (d, 1H, J = 1.6 Hz), 6.69 (d, 1H, J = 76.4 Hz), 6.00 (d, 1H, J = 5.2 Hz), 5.34 (quintet, 1H, J = 5.6 Hz), 4.79 (t, 2H, J = 6.8 Hz), 4.53 (dd, 2H, J = 6.8, 5.6 Hz), 3.92-3.87 (m, 1H), 3.77-3.72 (m, 1H), 3.75-3.69 (m, 2H), 3.69-3.66 (m, 2H), 3.62-3.58 (m, 4H), 2.89 (s, 3H), 2.88-2.83 (m, 1H), 2.75-2.55 (m, 4H), 2.03-1.96 (m, 2H), 1.90-1.81 (m, 2H), 1.01 (d, 3H, J = 6.4 Hz).
898		601	¹ H-NMR (400 MHz, 6d-DMSO) δ ppm 8.58 (s, 1H), 8.29 (s, 1H), 8.27 (s, 0.4H, HCOOH), 7.94 (d, 1H, J = 5.6 Hz), 7.92 (d, 1H, J = 4.4 Hz), 7.78 (d, 1H, J = 4.4 Hz), 7.19 (s, 1H), 6.69 (t, 1H, J = 76.4 Hz), 6.01 (d, 1H, J = 5.2 Hz), 5.35-5.31 (m, 1H), 4.81-4.77 (m, 2H), 4.55-4.51 (m, 2H), 3.93-3.89 (m, 1H), 3.78-3.68 (m, 8H), 2.92 (s, 3H), 2.89-2.84 (m, 2H), 2.70-2.58 (m, 4H), 2.03-2.00 (m, 2H), 1.89-1.78 (m, 2H), 1.01 (d, 3H, J = 6.4 Hz).
899		601	¹ H-NMR (500 MHz, 6d-DMSO) δ ppm 8.58 (d, 1H, J = 2.0 Hz), 8.29 (d, 1H, J = 1.5 Hz), 7.95-7.92 (m, 2H), 7.78 (dd, 1H, J = 8.5, 2.5 Hz), 7.19 (d, 1H, J = 1.5 Hz), 6.69 (t, 1H, J = 76.0 Hz), 6.01 (d, 1H, J = 5.5 Hz), 5.34 (quintet, 1H, J = 5.5 Hz), 4.79 (t, 2H, J = 7.5 Hz), 4.53 (dd, 2H, J = 7.5, 5.5 Hz), 3.91-3.89 (m, 1H), 3.76-3.74 (m, 1H), 3.75-3.70 (m, 2H), 3.62-3.57 (m, 2H), 3.51-3.49 (m, 4H), 2.92 (s, 3H), 2.87-2.86 (m, 1H), 2.70-2.67 (m, 1H), 2.67-2.61 (m, 3H), 2.03-2.01 (m, 2H), 1.90-1.89 (m, 2H), 1.02 (d, 3H, J = 6.5 Hz).

[0084] In another aspect, the present invention features the compounds of the invention as described herein for use in a method of treating or ameliorating fibrodysplasia ossificans progressiva in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound described herein (e.g., a compound in Table 1) or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof.

[0085] In another aspect, the present invention features the compounds of the invention as described herein for use in a method of treating or ameliorating diffuse intrinsic pontine glioma in a subject, comprising administering to the subject a pharmaceutically effective amount of a compound described herein (e.g., a compound in Table 1) or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof.

[0086] Pharmaceutically acceptable salts of these compounds are also contemplated for the uses described herein.

[0087] "Pharmaceutically acceptable salt" refers to any salt of a compound of the disclosure which retains its biological properties and which is not toxic or otherwise undesirable for pharmaceutical use. Pharmaceutically acceptable salts may be

derived from a variety of organic and inorganic counter-ions. Such salts include one or more of: (1) acid addition salts formed with organic or inorganic acids such as hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, sulfamic, acetic, trifluoroacetic, trichloroacetic, propionic, hexanoic, cyclopentylpropionic, glycolic, glutaric, pyruvic, lactic, malonic, succinic, sorbic, ascorbic, malic, maleic, fumaric, tartaric, citric, benzoic, 3-(4-hydroxybenzoyl)benzoic, picric, cinnamic, mandelic, phthalic, lauric, methanesulfonic, ethanesulfonic, 1,2-ethane-disulfonic, 2-hydroxyethanesulfonic, benzenesulfonic, 4-chlorobenzenesulfonic, 2-naphthalenesulfonic, 4-toluenesulfonic, camphoric, camphorsulfonic, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic, glucoheptonic, 3-phenylpropionic, trimethylacetic, tert-butylacetic, lauryl sulfuric, gluconic, benzoic, glutamic, hydroxynaphthoic, salicylic, stearic, cyclohexylsulfamic, quinic, muconic acid and the like acids; or (2) salts formed when an acidic proton present in the parent compound either (a) is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion or an aluminum ion, or alkali metal or alkaline earth metal hydroxides, such as sodium, potassium, calcium, magnesium, aluminum, lithium, zinc, and barium hydroxide, ammonia or (b) coordinates with an organic base, such as aliphatic, alicyclic, or aromatic organic amines, such as ammonia, methylamine, dimethylamine, diethylamine, picoline, ethanolamine, diethanolamine, triethanolamine, ethylenediamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylene-diamine, chlorprocaine, diethanolamine, procaine, N-benzylphenethylamine, N-methylglucamine piperazine, tris(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, and the like. Pharmaceutically acceptable salts further include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium and the like, and when the compound contains a basic functionality, salts of non-toxic organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, besylate, acetate, maleate, oxalate and the like. A pharmaceutically acceptable salt according to the disclosure includes at least one salt, and also may be mixtures of more than one salt.

Pharmaceutical Compositions

[0088] Pharmaceutical compositions of the invention and disclosure comprise one or more compounds of the invention and one or more pharmaceutically acceptable carrier(s). The term "pharmaceutically acceptable carrier" refers to a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting any subject composition or component thereof. Each carrier must be "acceptable" in the sense of being compatible with the subject composition and its components and not injurious to the patient. Some examples of materials which may serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate;

(13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

5 **[0089]** The compositions of the disclosure may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally, or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, 10 intralesional and intracranial injection or infusion techniques. In an embodiment, the compositions of the disclosure are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this disclosure may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile 15 injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a 20 solvent or suspending medium.

[0090] For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically- 25 acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such 30 as Tween, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[0091] The pharmaceutically acceptable compositions of this disclosure may be 35 orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions, or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous 40 suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring, or coloring agents may also be added.

[0092] Alternatively, the pharmaceutically acceptable compositions of this disclosure 45 may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the

rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0093] The pharmaceutically acceptable compositions of this disclosure may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs. Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[0094] For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this disclosure include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0095] The pharmaceutically acceptable compositions of this disclosure may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[0096] The amount of the compounds of the present disclosure that may be combined with the carrier to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration, and other factors determined by the person administering the single dosage form.

Dosages

[0097] Toxicity and therapeutic efficacy of compounds of the disclosure, including pharmaceutically acceptable salts and deuterated variants, can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The LD₅₀ is the dose lethal to 50% of the population. The ED₅₀ is the dose therapeutically effective in 50% of the population. The dose ratio between toxic and therapeutic effects (LD₅₀/ ED₅₀) is the therapeutic index. Compounds that exhibit large therapeutic indexes are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0098] Data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds may lie within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (*i.e.*, the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0099] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including but not limited to the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present disclosure in the composition will also depend upon the particular compound in the composition.

Treatment

[0100] Mutations in ALK2 cause the kinase to be inappropriately active and are associated with various diseases. The disclosure provides compounds that inhibit a mutant ALK2 gene, *e.g.*, a mutant ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification. In another aspect, the disclosure provides compounds that inhibit both wild type (WT) ALK2 protein and mutant forms of ALK2 protein. For the purposes of this disclosure, sequence information for ALK2 is found on the National Center for Biological Information (NCBI) webpage (<https://www.ncbi.nlm.nih.gov/>) under ACVR1 activin A receptor type 1 [Homo sapiens (human)]; Entrez Gene ID (NCBI): 90. It is also known as: FOP; ALK2; SKR1; TSRI; ACTRI; ACVR1A; ACVRLK2 said sequence information is incorporated herein.

[0101] In an embodiment, the disclosure provides the compounds for use in a method of inhibiting aberrant ALK2 activity in a subject comprising the step of administering to the subject in need thereof a pharmaceutically effective amount of at least one compound or pharmaceutical composition described herein. In an embodiment, the aberrant ALK2 activity is caused by a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification selected from one or more of L196P, PF197-8L, R202I, R206H, Q207E, R258S, R258G, G328A, G328V, G328W, G328E, G328R, G356D, and R375P. In an embodiment, the ALK2 enzyme has the amino acid modification R206H.

[0102] Because of their activity against ALK2, the compounds described herein can be used to treat a patient with a condition associated with aberrant ALK2 activity. In

an embodiment, the condition associated with aberrant ALK2 activity is fibrodysplasia ossificans progressiva. FOP diagnosis is based on the presence of congenital malformations of the great toes (hallux valgus) and the formation of fibrous nodules in soft tissues. The nodules may or may not transform into heterotopic bone. These soft tissue lesions are often first noted in the head, neck back. -97% of FOP patients have the same c.617G>A; R206H mutation in the ACVR1 (Alk2) gene. There is a genetic test available through the University of Pennsylvania (Kaplan et al, Pediatrics 2008, 121(5): e1295-e1300).

[0103] Other common congenital anomalies include malformations of the thumbs, short broad femoral necks, tibial osteochondromas and fused facet joints of the cervical spine. The fused facet joints in the neck often cause toddlers to scoot on their buttocks rather than crawl. FOP is commonly misdiagnosed (~80%; cancer or fibromatosis) and patients are frequently subjected to inappropriate diagnostic procedures such as biopsies that exacerbate disease and cause permanent disability.

[0104] In an embodiment, the present invention provides the compounds of the invention as described herein for use in a method of treating or ameliorating fibrodysplasia ossificans progressiva in a subject, comprising administering to the subject in need thereof a pharmaceutically effective amount of a compound or pharmaceutical composition described herein.

[0105] In an embodiment, the condition associated with aberrant ALK2 activity is fibrodysplasia ossificans progressiva (FOP) and the subject has a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification selected from one or more of L196P, PF197-8L, R202I, R206H, Q207E, R258S, R258G, G328A, G328W, G328E, G328R, G356D, and R375P. In one aspect of this embodiment, the ALK2 enzyme has the amino acid modification R206H.

[0106] The present disclosure includes methods of identifying and/or diagnosing patients for treatment with one or more of the compounds or pharmaceutical compositions described herein. In an embodiment, the disclosure provides a method of detecting a condition associated with aberrant ALK2 activity e.g., FOB in a subject, wherein the method includes a. obtaining a sample e.g., plasma from the subject e.g., a human patient; and b. detecting whether one or more mutations in an ALK2 gene as described herein are present in the sample. In another embodiment, the disclosure provides a method of diagnosing a condition associated with aberrant ALK2 activity in a subject, said method comprising: a. obtaining a sample from the subject; b. detecting whether one or more mutations in an ALK2 gene as described herein are present in the sample using a detection method described herein; and c. diagnosing the subject with the condition when the presence of the one or more mutations is detected. Methods for detecting a mutation include but are not limited to hybridization-based methods, amplification-based methods, microarray analysis, flow cytometry analysis, DNA sequencing, next-generation sequencing (NGS), primer extension, PCR, in situ hybridization, dot blot, and Southern blot. In an embodiment, the present disclosure provides a method of diagnosing and treating a

condition associated with aberrant ALK2 activity in a subject, said method comprising a. obtaining a sample from a subject; b. detecting whether one or more mutations in an ALK2 gene as described herein are present in the sample; diagnosing the subject with the condition when the one or more mutations in the sample are detected; and d. administering an effective amount of one or more of the compounds or a pharmaceutical composition described herein to the diagnosed patient. In an embodiment, the disclosure provides a method of treating a condition associated with aberrant ALK2 activity in a subject, said method comprising a. determining if, having determined if, or receiving information that the subject has one or more mutations in an ALK2 gene as described herein; b. identifying the subject as responsive to one or more compounds or a pharmaceutical composition described herein; and c. administering an effective amount of the one or more compounds or pharmaceutical compositions to the subject.

[0107] In an embodiment, the condition associated with aberrant ALK2 activity is a brain tumor, e.g., glial tumor. In an embodiment, the glial tumor is diffuse intrinsic pontine glioma (DIPG). In an embodiment, the invention provides the compounds of the invention as described herein for use in a method of treating or ameliorating diffuse intrinsic pontine glioma in a subject, comprising administering to the subject in need thereof a pharmaceutically effective amount of a compound or pharmaceutical composition described herein.

[0108] In an embodiment, the condition associated with aberrant ALK2 activity is diffuse intrinsic pontine glioma and the subject has a mutation in an ALK2 gene that results in the expression of an ALK2 enzyme having an amino acid modification selected from one or more of R206H, G328V, G328W, G328E, and G356D. In one aspect of this embodiment, the ALK2 enzyme has the amino acid modification R206H.

[0109] Other conditions associated with aberrant ALK2 activity are anemia associated with inflammation, cancer or chronic disease or trauma- or surgery-induced heterotopic ossification.

[0110] A compound of the invention is suitable for co-administration (either as part of a combination dosage form or as a separate dosage form administered prior to, sequentially with, or after administration) with a second therapeutic agent useful in treating the disease to be treated e.g., FOP, for example a steroid (e.g., prednisone) or other anti-allergenic agents such as omalizumab.

[0111] A compound of the invention is suitable for co-administration with a RAR- γ agonist or an antibody against activating for treating the disease to be treated e.g., FOP, such as the RAR- γ agonist palovarotene and the antibody against activin REGN2477.

[0112] A compound of the invention is suitable for co-administration with therapies that target mast cells useful in treating FOP, such as a mast cell inhibitor including, but not limited to a KIT inhibitor. Examples of a mast cell inhibitor to be co-administered are cromolyn sodium (or sodium cromoglicate); brentuximab


(ADCETRIS®); ibrutinib (IMBRUVICA®); omalizumab (XOLAIR®); anti-leukotriene agents (e.g., montelukast (SINGULAIR®) or zileuton (ZYFLO® or ZYFLO CR®)); and KIT inhibitors (e.g., imatinib (GLEEVEC®), midostaurin (PKC412A), masitinib (MASIVET® or KINAVET®), BLU-285, DCC-2618, PLX9486).

5 Synthesis

[0113] The Schemes below are meant to provide general guidance in connection with preparing the compounds of the disclosure. One skilled in the art would understand that the preparations shown in the Schemes can be modified or optimized using general knowledge of organic chemistry to prepare various compounds of the disclosure.

Synthetic Protocol 1

[0114]




Tf = trifluoromethyl sulfonyl; LG^2 = leaving group; PG = protecting group; R^C is an optional capping group; $R^{2'} = R^2$ or a precursor to R^2 if capped with R^C ; Z = a Pd-mediated cross coupling partner for a halide, e.g., a boronic ester/acid, a tin group or a zinc group.

[0115] The pyrrolopyridazine **1** having a leaving group (LG^2) can be coupled with a functionalized piperazine **2** via a substitution reaction to provide an intermediate **3** with a new carbon-nitrogen bond formed. The functionalized piperazine **2** can be formed by reaction with such groups as carboxylic acids/acid chlorides, chloroformates and isocyanates (or activated carbamates, etc.) to form amides, carbamates and ureas, respectively, via well-established reaction protocols; followed by deprotection (if necessary). The resulting pyrrolopyridazine **3** can be coupled to intermediate **4** via a palladium-mediated coupling reaction, e.g., Suzuki, Stille, or Negishi coupling, to provide an intermediate (**5**) with a new carbon-carbon

bond formed. (LG^2 can be, e.g., Cl, Br, or I. Z can be boronate, boronate ester, or trialkyltin. R^2 can be, for example, Br, OH, N-linked alkyl or cycloalkylamine, or C-linked alkyl or cycloalkylamine. The resulting intermediate **5** can be further functionalized (following protecting group removal, if necessary) by capping reactions including alkylation, reductive amination with carbonyl containing compounds, acylation, ether formation via Mitsunobu coupling, amination using the Buchwald reaction, or alkyl/cycloalkylamine formation achieved through vinylboronic acid addition followed by hydrogenation. Synthesis of exemplary compounds prepared using Synthetic Protocol 1 are disclosed in certain of the Examples below.

10 Synthetic Protocol 2

[0116]




[0117] The pyrrolopyridazine **1** can be coupled with a functionalized piperazine **2** via a substitution reaction to provide an intermediate with a new carbon-nitrogen bond formed. The resulting intermediate can be converted to a boronate ester by palladium mediated coupling with bis-pinacolato diboron to give intermediate **3'**. In some instances, the R^1 group can be replaced with a different R^1 group e.g., replacement nitrophenyloxy with oxetanyloxy. The resulting pyrrolopyridazine boronate can be coupled to an aryl halide (**4'**) via a palladium-mediated coupling reaction, e.g., Suzuki coupling, to provide an intermediate with a new carbon-carbon bond formed. (LG^2 can be Cl, Br, I, OTf; R^2 can be, for example, Br, OH, N-linked alkyl or cycloalkylamine, or C-linked alkyl or cycloalkylamine.) This resulting di-substituted pyrrolopyridazine, intermediate **5**, can be further functionalized (following protecting group removal, if necessary) by capping reactions including alkylation, reductive amination with carbonyl containing compounds, acylation, ether formation via Mitsunobu coupling between a phenol and alcohols, amination using the Buchwald reaction or alkyl/cycloalkylamine formation achieved through vinylboronic acid addition followed by hydrogenation via well-established reaction protocols.

Synthesis of exemplary compounds prepared using Synthetic Protocol 2 are disclosed in certain of the Examples below.


Synthetic Protocol 3

5


[0118]




- [0119]** The pyrrolopyridazine **7** can be coupled with substituted aryl boronic acids **4** ($R^{2'}$ can be, for example, Br, OH, N-linked alkyl or cycloalkylamine, or C-linked alkyl or cycloalkylamine) to form a new carbon-carbon bond. The resulting intermediate **8** can be further functionalized (following protecting group removal, if necessary) by capping reactions including alkylation, reductive amination with carbonyl containing compounds, acylation, ether formation via Mitsunobu coupling, or amination using the Buchwald reaction. Removal of the BOC group in intermediate **9** can be followed by capping reactions of the resulting free NH using activated carboxylic acids, chloroformates, carbamoyl chlorides/isocyanates to give amides, carbamates, or ureas respectively via well-established reaction protocols. Synthesis of exemplary compounds prepared using Synthetic Protocol 3 are disclosed in certain of the Examples below.

Synthetic Protocol 4**[0120]**


- 5 **[0121]** The pyrrolopyridazine boronate ester **10** can be coupled with aryl halides 4' to provide intermediate **8**. (LG² can be, e.g., Cl, Br, or I. R^{2'} can be, for example, Br, OH, N-linked alkyl or cycloalkylamine, or C-linked alkyl or cycloalkylamine).
- 10 Intermediate **8** can be further functionalized (following protecting group removal, if necessary) by capping reactions including alkylation, reductive amination with carbonyl containing compounds, acylation, ether formation via Mitsunobu coupling, or amination using the Buchwald reaction to provide intermediate **9**. Removal of the BOC group in intermediate **9** can be followed by capping reactions of the resulting free NH using activated carboxylic acids, chloroformates, carbamoyl chlorides/isocyanates to give amides, carbamates, or ureas respectively via well-
- 15 established reaction protocols. Synthesis of exemplary compounds prepared using Synthetic Protocol 4 are disclosed in certain of the Examples below.

Synthetic Protocol 5**[0122]**

- 5 **[0123]** The pyrrolopyridazine bromide **7** can be converted to the boronate ester **10**, via a Pd-mediate reaction. Removal of the BOC group in intermediate **10** can be followed by capping reactions of the resulting free NH using activated carboxylic acids, chloroformates, carbamoyl chlorides/isocyanates to give amides, carbamates, or ureas respectively via well-established reaction protocols.
- 10 The pyrrolopyridazine boronate ester **3'** can be coupled with aryl halides **4'** to provide intermediate **5**. (LG² can be, e.g., Cl, Br, or I. R^{2'} can be, for example, Br, OH, N-linked alkyl or cycloalkylamine, or C-linked alky or cycloalkylamine). Intermediate **5** can be further functionalized (following protecting group removal, if necessary) by capping reactions including alkylation, reductive amination with carbonyl containing compounds, acylation, ether formation via Mitsunobu coupling, or amination using the Buchwald reaction to provide **6**. Synthesis of exemplary compounds prepared using Synthetic Protocol 5 are disclosed in certain of the Examples below.
- 15
- 20 **[0124]** In yet another embodiment, the invention provides an intermediate for synthesizing compounds of the disclosure. The intermediate is 6-bromopyrrolo[1,2-b]pyridazin-4-ol:




- In another embodiment, the invention provides a process for synthesizing said intermediate. The method of synthesizing 6-bromopyrrolo[1,2-b]pyridazin-4-ol comprises the step of combining a compound of Formula C-1:
- 25



(C-1),

with a compound of Formula D-1:



(D-1),

wherein:

- 5 R^{21} is selected from chloro, bromo, and iodo;
 R^{22} is a leaving group; and
 R^{23} is an electron withdrawing group.
- 10 **[0125]** In some embodiments, R^{21} is bromo.
- [0126]** In some embodiments, R^{22} is selected from $-N(R^{24})(R^{25})$ and $-OR^{24}$, wherein each of R^{24} and R^{25} is an independently selected C_1 - C_4 alkyl. In more specific aspects of these embodiments, R^{22} is $-N(CH_3)_2$.
- 15 **[0127]** In some embodiments, R^{23} is selected from methylcarbonyl, t-butylcarbonyl, 4-nitrophenylcarbonyl, 4-cyanophenylcarbonyl, 4-trifluoromethylphenylcarbonyl, 4-fluorophenylcarbonyl, 4-trifluoromethylcarbonylphenylcarbonyl, 4-ethoxycarbonylphenylcarbonyl, 4-trifluoromethylsulfonylphenylcarbonyl, 2,4,6-trimethylphenylcarbonyl, 2,4,6-trimethyl-3,5-dinitrophenylcarbonyl, 2-trifluoromethyl-20 4-nitrophenyl, 2,4-dinitrophenyl and diphenylphosphinyl. In more specific aspects of these embodiments, R^{23} is 4-nitrophenylcarbonyl.
- [0128]** In the method of synthesizing 6-bromopyrrolo[1,2-b]pyridazin-4-ol from C-1 and D-1, the starting materials are dissolved in a polar solvent. The choice of polar solvent can be made from any known in the art. More specifically, the polar solvent is selected from N-methyl-2-pyrrolidine ("NMP"), N,N-dimethylacetamide ("DMAC"), dimethylformamide ("DMF"), tetrahydrofuran ("THF"), methyl-tetrahydrofuran ("MeTHF"), dimethyl sulfoxide ("DMSO"), and cyclopentylmethyl ether ("CPME").
25 Even more specifically, the polar solvent is NMP or DMAC.
- [0129]** In the first step of the synthesis method C-1 dissolved in the polar solvent. This is done at the lowest temperature possible which allows dissolution. Dissolved C-1 is then treated with a base, typically 1.15-1.5 equivalents thereof, optionally
35 maintained under a N_2 atmosphere. The choice of base can be made from any known in the art. More specifically, the base is selected from $KOC(CH_3)_3$, $NaOC(CH_3)_3$, $LiOC(CH_3)_3$, $LiC(CH_3)_3$, $Li(CH_2)_3CH_3$, $LiN(C_3H_7)_2$, $NaOCH_3$, $NaOCH_2CH_3$, $KOCH_3$, $LiOCH_3$, $LiOCH_2CH_3$, and $KOCH_2CH_3$. Even more specifically, the base is $KOC(CH_3)_3$.
- 40 **[0130]** Treatment of C-1 with a base is performed at a temperature of between about 15 to $30^\circ C$ for 0.5-2 hours with stirring. The base-treated C-1 solution is then optionally cooled to -8 to $-5^\circ C$ before adding reagent D-1.

[0131] **D-1** is also dissolved in a polar solvent at a temperature of between -5 to 30°C optionally under a N₂ atmosphere and then slowly added to base-treated **C-1**. The resulting mixture is stirred for 1-2 hrs until at least 90% of **C-1** has disappeared as determined by LCMS or IPC.

[0132] At that point a protonating agent is added at acid pH and at a temperature of between about -5 to 10°C. The choice of protonating agent can be made from any known in the art. More specifically, the protonating agent is selected from NH₄Cl, NaHCO₃, KHCO₃, LiHCO₃, acetic acid, HCl, HBr, and H₂SO₄. Even more specifically, the protonating agent is NH₄Cl. In some specific aspects, the pH of the reaction with the protonating agent is adjusted to between about 1 and 5, more specifically between about 2 and 4 with an acidifying agent. In certain specific embodiments, the acidifying agent is HCl. The protonation reaction is allowed to proceed for between 0.5-2 hrs at 0-10°C.

[0133] The resulting mixture is then optionally filtered before extracting the insoluble material with an extraction agent. If filtered, the filter cake is extracted multiple times with the extraction agent and filtered after each extraction. The original filtrate is then combined with all of the extraction filtrates and the resulting solution is allowed to separate into an organic and an aqueous phase. The aqueous phase is then extracted several more times with the extraction agent and all organic phases are combined. If the mixture resulting from the protonation reaction is not filtered, it is extracted multiple times with the extraction agent, with all organic phases resulting from the extractions being pooled.

[0134] The choice of extraction agent may be made from any agent known in the art that is capable of extracting material from an aqueous phase into an organic phase. More specifically, the extraction agent is selected from methyl tert-butyl ether ("MTBE"), MeTHF, dichloromethane ("DCM"), CPME, diethyl ether, ethyl acetate, toluene, and isopropyl acetate. Even more specifically, the extraction agent is MTBE.

[0135] The pooled organic layers resulting from the extractions are optionally washed with saturated NaCl, dried over anhydrous Na₂SO₄, filtered if any insoluble material remains, and the soluble material then concentrated to dryness. This process typically results in at least 90% pure 6-bromopyrrolo[1,2-b]pyridazin-4-ol as determined by LCMS, HPLC or quantitative ¹H-NMR.

EXAMPLES

[0136] Compounds of the invention and disclosure, including salts and N -oxides thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes, such as those in the Schemes below. The reactions for preparing compounds of the disclosure can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, e.g., temperatures which can range from the

solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.

[0137] Preparation of compounds of the disclosure can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Wuts and Greene, *Protective Groups in Organic Synthesis*, 4th ed., John Wiley & Sons: New Jersey, (2006), which is incorporated herein by reference in its entirety.

[0138] Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance (NMR) spectroscopy (e.g., ^1H or ^{13}C), infrared (IR) spectroscopy, spectrophotometry (e.g., UV-visible), mass spectrometry (MS), or by chromatographic methods such as high performance liquid chromatography (HPLC) or thin layer chromatography (TLC). Analytical instruments and methods for compound characterization include the following:

LC-MS: Unless otherwise indicated, all liquid chromatography-mass spectrometry (LC-MS) data (sample analyzed for purity and identity) were obtained with an Agilent model-1260 LC system using an Agilent model 6120 or model 1956 mass spectrometer utilizing ES-API ionization fitted with an Agilent Poroshel 120 (EC-C18, 2.7 μm particle size, 3.0 $\times$ 50mm dimensions) reverse-phase column at 22.4 degrees Celsius. The mobile phase consisted of a mixture of solvent 0.1% formic acid in water and 0.1% formic acid in acetonitrile. A constant gradient from 95% aqueous/5% organic to 5% aqueous/95% organic mobile phase over the course of 4 minutes was utilized. The flow rate was constant at 1mL/min.

[0139] Alternatively, LC-MS data was obtained using the following columns and mobile phases. Basic Mobile Phase: A: water (10 mM NH_4HCO_3) B: ACN; Gradient: 5% B increase to 95% B within 1.2 min, 95% B for 1.3 min, back to 5% B within 0.01 min; Flow Rate: 2 mL/min; Column: XBridge, 3.5 μm , 50*4.6 mm; Oven Temperature: 50 $^\circ\text{C}$. Acidic Mobile Phase: A: water (0.01% TFA) B: CAN (0.01% TFA); Gradient: 5% B increase to 95% B within 1.2 min, 95% B for 1.3 min, back to 5% B within 0.01 min; Flow Rate: 2 mL/min; Column: Sunfire, 3.5 μm , 50*4.6 mm; Oven Temperature: 50 $^\circ\text{C}$.

[0140] Alternatively, HPLC data was obtained using the following columns and mobile phases. Basic Mobile Phase: A: water (10 mM NH_4HCO_3) B: ACN; Gradient: 5% B increase to 95% B within 1.2 min, 95% B for 1.3 min, back to 5% B within 0.01 min; Flow Rate: 2 mL/min; Column: XBridge, 3.5 μm , 50*4.6 mm; Oven Temperature: 50 $^\circ\text{C}$. Acidic Mobile Phase: A: water (0.01% TFA) B: ACN (0.01% TFA); Gradient: 0 min 5% B, 3 min 5% B, 10 min 95% B, 15 min 95% B; Flow Rate: 1.2 mL/min; Column: Eclipse XDB-C18, 4.6*150 mm, 5 μm ; Oven Temperature: 40 $^\circ\text{C}$.

[0141] Prep LC-MS: Preparative HPLC was performed on a Shimadzu Discovery

VP® Preparative system fitted with a Luna 5u C18(2) 100A, AXIA packed, 250 × 21.2 mm reverse-phase column at 22.4 degrees Celsius. The mobile phase consisted of a mixture of solvent 0.1% formic acid in water and 0.1% formic acid in acetonitrile. A constant gradient from 95% aqueous/5% organic to 5% aqueous/95% organic mobile phase over the course of 25 minutes was utilized. The flow rate was constant at 20 mL/min. Reactions carried out in a microwave were done so in a Biotage Initiator microwave unit.

[0142] Chiral HPLC: Preparative HPLC to resolve chiral mixtures was performed on one of the following systems. For SFC using either a SFC-80 or SFC-200 (Thar, Waters) instrument, we used an AD-H 20x250mm, 5µm Diacel column run at 35°C using a mobile phase gradient of CO₂/Methanol (0.1%NH₄OH)= 40:60 - 90:10 at a flow rate of 80-180 g/min and detection at 214-360 nm. For HPLC using a Gilson-281 instrument, we used an AD-H 20x250mm, 10µm Diacel column run at 40°C using a mobile phase gradient of Hexane(0.1%DEA): EtOH(0.1%DEA)=0:100 - 100:0.

[0143] Silica gel chromatography: Silica gel chromatography was performed on either a Teledyne Isco COMBIFLASH® Rf unit or a BIOTAGE® Isolera Four unit.

[0144] Proton NMR: Unless otherwise indicated, all ¹H NMR spectra were obtained with a Varian 400MHz Unity Inova 400 MHz NMR instrument (acquisition time = 3.5 seconds with a 1 second delay; 16 to 64 scans), a Bruker, AVANCE III 500MHz UltraShield-Plus digital NMR spectrometer, or a Bruker, AVANCE III 400MHz UltraShield-Plus digital NMR spectrometer.. Where characterized, all protons were reported in DMSO-*d*₆ solvent as parts-per million (ppm) with respect to residual DMSO (2.50 ppm).


EXAMPLES

[0145] The below Schemes are meant to provide general guidance in connection with preparing the compounds of the invention. One skilled in the art would understand that the preparations shown in the Schemes can be modified or optimized using general knowledge of organic chemistry to prepare various compounds of the invention and disclosure.

Example 1. Synthesis of 6-bromopyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate

Step 1: Synthesis of 1-(4-bromo-1H-pyrrol-2-yl)ethanone (Intermediate B)


[0146]



- 5 **[0147]** Amberlyst 15 (0.09 g/g-bulk-LR) was added to a solution of commercially available 1-(1H-pyrrol-2-yl)ethanone (70 g; 1.00 equiv; 641.45 mmoles) in tetrahydrofuran (10 mL/g-bulk-LR; 9.71 moles; 700.00 mL; 700.00 g) at room temperature (RT) (around 25 °C). Next, 1-bromopyrrolidine-2,5-dione (1 equiv (molar); 641.45 mmoles; 114.17 g) was added in portions at -30 to -20°C and stirred for approximately 1 h until LCMS indicated that the reaction was complete. The reaction mixture was then filtered and the filtrate quenched with saturated Na₂SO₃ aqueous (350 mL) and extracted with DCM (700 mL×2). The organic layer was concentrated and then diluted with MTBE (700mL). The organic layers were combined and then washed with sat. NaHCO₃ (350mL × 2) and concentrated on a rotavapor under reduced pressure to give 1-(4-bromo-1H-pyrrol-2-yl)ethanone (Intermediate B; **91 g**; 0.75 equiv; 483.98 mmoles; 91.00 g; 75.45% yield) as a white solid. LCMS: 100% purity.

Step 2: Synthesis of (E)-1-(4-bromo-1H-pyrrol-2-yl)-3-(dimethylamino)prop-2-en-1-one (Intermediate C)


20 [0148]



- 25 **[0149]** 1-(4-bromo-1H-pyrrol-2-yl)ethanone (50 g; 1.00 equiv; 265.92 mmoles; 50.00 g) was added to 1,1-dimethoxy-N,N-dimethylmethanamine (5 mL/g-pure-LR; 2.10 moles; 250.00 mL; 250.00 g;) at room temperature (around 25 °C) and then the reaction mixture was heated at 70-80 °C for 12 h when LCMS indicated that the reaction was complete resulting in a suspension. The reaction mixture was filtered and the cake washed with PE (300mL). The wet cake was dried in air for 16 h to give (E)-1-(4-bromo-1H-pyrrol-2-yl)-3-(dimethylamino)prop-2-en-1-one (35 g; 0.54 equiv; 143.97 mmoles; 35.00 g; 54.14% yield) as a yellow solid. LCMS: >95%.

Step 3: Synthesis of 6-bromopyrrolo [1,2-b]pyridazin-4-ol (Intermediate E)


[0150]



[0151] (E)-1-(4-bromo-1H-pyrrol-2-yl)-3-(dimethylamino)prop-2-en-1-one (**C**; 20 g; 1.00 equiv; 82.27 mmol; 20.00 g) was taken up in 1-methylpyrrolidin-2-one (30 mL/g-bulk-LR; 6.05 moles; 600.00 mL; 600.00 g) to form a solution. Potassium 2-methylpropan-2-olate (1.5 equiv (molar); 123.40 mmol; 13.85 g) was then added in portions. The solution temperature was kept at 10 to 25 °C and the solution was then stirred at 15 to 25°C for 0.5 h. Commercially available O-(4-nitrobenzoyl)hydroxylamine (**D**; 1.5 equiv (molar); 123.40 mmol; 22.48 g) was then added into the reaction mixture maintaining the temperature at 20 to 30°C and then stirred at 30 °C for 2 h until LCMS indicated that the starting material was gone. Saturated aqueous ammonium chloride (200 mL) was added dropwise into the reaction mixture cooled in an ice bath (0 °C), diluted with water (200 mL), and the pH was adjusted to between 3 and 4 with hydrochloric acid (1 M). The resulting solution was extracted with MTBE (3×150 mL), the combined organic layers were dried over anhydrous sodium sulfate, and then filtered and concentrated to dryness. The residue was purified by silica gel chromatography (petroleum ether: ethyl acetate =10:1-5:1) to give 6-bromopyrrolo [1,2-b]pyridazin-4-ol (**E**; 16 g, 90% yield; Purity:91.6%) and isolated as a yellow solid, which is used for next step without further purifications. LCMS: 91.6%.

Step 3 Alternate: Synthesis of 6-bromopyrrolo [1,2-b]pyridazin-4-ol (Intermediate E)

[0152]




[0153] A synthesis of Intermediate E using the same reagents, but different conditions, is described in this Step 3 Alternate.

[0154] Two hundred kilograms of DMAc was added into a reactor to which was then quickly added. potassium tert-butoxide (1.15equiv) under protection of N₂. The mixture was stirred until the reagents were dissolved. ((E)-1-(4-bromo-1H-pyrrol-2-yl)-3-(dimethylamino)prop-2-en-1-one)(**C**; 20.5kg, 84.32mol, 1 equiv) was added and was kept stirring at 20-30°C for 1-2 hours. The reaction mixture was then cooled to -8 to -5°C. O-(4-nitrobenzoyl)hydroxylamine (**D**; 16.1kg, 88.54mol, 1.05equiv) was then dissolved into DMAc(100kg) in a separate container and the solution maintained at -5 - 0°C, then slowly the resulting solution of **D** was added to the reaction mixture. During addition, the temperature of solution **D** was maintained at -5 - 0°C and kept under N₂ protection. The addition of **D** was completed after about 4 hrs. The resulting mixture was continually stirred at -5 - 0°C for an additional 1-2 hr, until less than 8% of starting material **C** was present as determined by IPC. Saturated NH₄Cl (150kg) was added at -5 - 10°C and the pH was adjusted to 2-2.5 with hydrochloric acid also maintained at -5~10°C. The mixture was continually stirred for an additional 1-2 hrs at 0 - 10°C. The resulting mixture was then filtered, and the filter cake washed twice with MTBE (100kg x 2). The filtrates were

combined, and the aqueous layer separated from the organic layer. The aqueous layer was then extracted with MTBE 4-5 times and all of the organic phases were combined. The organic phases were then washed with sat. NaCl (40kg x 3). The organic phases were dried over anhydrous Na₂SO₄, filtered and the filtrate was concentrated at 35-45°C under vacuum until the solution volume was ~50L (this concentration process should be completed within 3hrs). The resulting concentrated solution was separated into several smaller batches and each batch transferred to a rotary evaporator for further and faster concentration to give wet solid (this process should be completed within 2hrs). The resulting wet solids were combined and then DCM (40kg) was added to slurry wash the solid at 10-15°C for 0.5h. The slurry was then filtered and dried to give 7.45kg of **E** (HPLC: 98.51%, RRT=~1.4 impurity is 1.28%, QHNMR : 96.72%, assay by external standard method is 94.5%, yield is 41.4%).

Step 4: Synthesis of 6-bromopyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate (Intermediate F)

[0155]




[0156] 6-bromopyrrolo[1,2-b]pyridazin-4-ol (**E**; 10 g; 1.00 equiv; 46.94 mmol; 10.00 g), dichloromethane (15 mL/g-bulk-LR; 2.34 moles; 150.00 mL; 198.75 g) and triethylamine (1.18 equiv (molar); 55.39 mmol; 7.68 mL; 5.61 g) were combined in a 250 mL reactor. Trifluoromethanesulfonic anhydride (1.15 equiv (molar); 1.15 equiv; 53.98 mmol; 9.08 mL; 15.23 g) was added dropwise and the temperature was kept between 0-20°C. The reaction mixture was warmed to 25 °C and stirred for an additional 2 h until LC-MS showed the reaction was completed. The mixture was then diluted with DCM (160 mL) and washed with sat. NaHCO₃ solution (2 x 80 mL). The organic phases were combined and dried over Na₂SO₄, filtered and concentrated under reduced pressure. MTBE (80mL) and PE (80mL) were added to dilute the crude product with stirring. Any solid that precipitated out at the bottom was removed by filtration. The filtrate was washed with sat. NaHCO₃ (40 mL x 2) and water (40 mL) and sat. NaCl (40mL) and then concentrated to give a crude product. Further purification was achieved with silica chromatography (PE/MTBE=100/0 to 50/1) to give 6-bromopyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate (**F**; 9 g; 0.56 equiv; 26.08 mmol; 9.00 g; 55.56% yield; [Actual]) as a dark green liquid. LC-MS: 345 (M+H)⁺, 98% purity (214 nm).

Alternate Synthesis of Intermediate E


[0157] An alternate synthesis of Intermediate E was carried out as follows

Step 1: 6-bromo-4-hydroxy-pyrrolo[1,2-b]pyridazine-3-carbonitrile

[0158]

- 5 **[0159]** To a solution of methyl 1-amino-4-bromo-pyrrole-2-carboxylate (4.0 g, 18 mmol, 1.0 eq) and 3,3-dimethoxypropanenitrile (12.6 g, 109 mmol, 6.0 eq) was added TsOH (629 mg, 4 mmol, 0.2 eq). The reaction mixture was stirred at 80 °C for 6 h. Then, DBU (16.7 g, 109 mmol, 6.0 eq) was added into the reaction mixture and stirred for another 10 h at 80 °C. TLC showed the reaction was complete. The
- 10 mixture was diluted with water (5mL) and extracted with EA (10 mLx2). The combined organic phases were washed with brine, dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by silica gel column chromatography (DCM:MeOH=10:1) to give the title product (3.7 g, 15 mmol, 85% yield) as a yellow solid.

15 **Step 2: 6-bromo-4-hydroxypyrrolo[1,2-b]pyridazine-3-carboxamide**

[0160]

- 20 **[0161]** To the solution of 6-bromo-4-hydroxy-pyrrolo[1,2-b]pyridazine-3-carbonitrile (2.0 g, 8.4 mmol, 1.0 eq) in EtOH (20 mL) was added a solution of NaOH (16.0 g, 400 mmol) in H₂O (50 mL). The reaction mixture was stirred for 48 h at 100 °C until TLC (petroleum ether (PE): ethyl acetate (EA) =0:1) indicated that most of the starting material was consumed. The reaction mixture was concentrated to remove the EtOH. The pH of the resulting aqueous solution was adjusted to 5-6 and then
- 25 extracted with ethyl acetate. The combined organic layer was dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by silica gel column chromatography (petroleum ether: ethyl acetate = 5:1-1:1) to afford the title product (1.1 g, 4 mmol, 51% yield) as a yellow solid.

Step 3: 6-bromopyrrolo[1,2-b]pyridazin-4-ol (Intermediate E)

30 **[0162]**


- [0163]** To a solution of 6-bromo-4-hydroxypyrrolo[1,2-b]pyridazine-3-carboxamide

(1.0 g, 4 mmol) in concentrated HCl (aq., 30 mL) was added dioxane (2 mL) and EtOH (2 mL). The reaction mixture was stirred for 48 h at 100 °C. TLC (petroleum ether: ethyl acetate = 0:1) indicated that most of the starting material was consumed and the reaction mixture was concentrated to remove organic solvents. The pH of the resulting aqueous solution was adjusted to 4-6 and then extracted (twice) with ethyl acetate. The combined organic layer was dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by silica gel column chromatography (petroleum ether: ethyl acetate = 10:1-5:1) to afford crude Intermediate E (150 mg, contains two byproducts (halogen-exchange and de-halogen)) as a yellow solid.

Example 2. Synthesis of cyclopropyl(4-(6-(4-(piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 127).

Step 1: Synthesis of (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone.


[0164]



[0165] A mixture of 6-bromopyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate (30 g, 86.9 mmol), cyclopropyl(piperazin-1-yl)methanone (16.0 g, 104 mmol), and triethylamine (13.1 g, 130 mmol) in NMP (300 mL) was stirred at 100 °C for 30 min. The reaction mixture was cooled and diluted with EA. The organic layer was washed with water and brine, concentrated and purified by silica gel column to give the title product (26.0 g, yield 86%) as a yellow solid. MS (ES+) $C_{15}H_{17}BrN_4O$ requires: 348, found: 349 $[M+H]^+$.

Step 2: Synthesis of cyclopropyl(4-(6-(4-(piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 127).

[0166]




[0167] A mixture of (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (3.0 g, 8.59 mmol), 1-(4-(4,4,5,5-tetramethyl-1,3,2-

dioxaborolan-2-yl)phenyl)piperazine hydrochloride (3.10 g, 12.8 mmol), K_2CO_3 (4.73 g, 34.3 mmol) and $Pd(dppf)Cl_2 \cdot CH_2Cl_2$ (700 mg, 859 μ mol) in 1,4-dioxane/water (30 mL/5 mL) was degassed with N_2 , and then stirred at 100 °C for 16 h under N_2 . The mixture was cooled to RT and concentrated. The residue was purified by silica gel column to give the title product (1.95 g, yield 52.8%) as a white solid. MS (ES+) $C_{25}H_{30}N_6O$ requires: 430, found: 431 $[M+H]^+$.

Example 3. Synthesis of cyclopropyl(4-(6-(4-(4-(2-hydroxyethyl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 274)

10 [0168]




[0169] A mixture of cyclopropyl(4-(6-(4-(piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (100 mg, 232 μ mol), 2-bromoethanol (57.9 mg, 464 μ mol) and potassium carbonate (32 mg, 0.232 mmol) was stirred at 70 °C overnight (-12 h). The reaction mixture was cooled and concentrated. The residue was purified by Prep-HPLC to afford the title compound as a white solid (10.5 mg, yield 9.5%). MS (ES+) $C_{27}H_{34}N_6O_2$, requires: 474, found: 475 $[M+H]^+$.

20 **Example 4. Synthesis of cyclopropyl(4-(6-(4-(4-(oxetan-3-yl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 314).**

Step 1: Synthesis of tert-butyl 4-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo(1,2-b]pyridazin-6-yl)phenyl)piperazine-1-carboxylate


25 [0170]



[0171] A mixture of (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (65 mg, 0.19 mmol), tert-butyl 4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine-1-carboxylate (86 mg, 0.22 mmol), K_2CO_3 (51 mg, 0.37 mmol) and $Pd(dppf)Cl_2$ (14 mg, 0.019 mmol) in dioxane/water (10/1) was irradiated in the microwave at 100 °C for 1 h. Concentrated and purified by flash column (PE/EA = 2/1 to 1/10) to give the title product (66 mg, yield 65.4%). MS (ES+) $C_{30}H_{38}N_6O_3$, requires: 530, found 531 $[M+H]^+$.

Step 2: Synthesis of cyclopropyl(4-(6-(4-(piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone

[0172]




5

[0173] To a solution of tert-butyl 4-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)phenyl)piperazine-1-carboxylate (66 mg, 0.12 mmol) in DCM was added TFA (TFA/DCM, 10:1). The reaction mixture was stirred at RT for 1 h. Saturated NaHCO₃ solution was added to the mixture to bring the pH to 8-9, and then the mixture was extracted with DCM. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated to give the title product (55 mg, crude). MS (ES⁺) C₂₅H₃₀N₆O, requires: 430, found 431 [M+H]⁺.

10

Step 3: Synthesis of cyclopropyl(4-(6-(4-(4-(oxetan-3-yl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone

[0174]



15

[0175] To a solution of cyclopropyl(4-(6-(4-(piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (55 mg, crude) in 1,2-dichloroethane was added oxetan-3-one (92 mg, 1.27 mmol), followed by addition of NaBH(OAc)₃ (269 mg, 1.27 mmol). The reaction mixture was stirred at RT overnight. Concentration and purification by Prep-HPLC gave the title product (3.7 mg, yield 6.3%). MS (ES⁺) C₂₈H₃₄N₆O₂ requires: 486, found 487 [M+H]⁺.


20

Example 5. Synthesis of cyclopropyl(4-(6-(4-(2-(piperazin-1-yl)ethoxy)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 273) and (4-(6-(4-(2-(4-ethylpiperazin-1-yl)ethoxy)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 390).

25


Step 1: Synthesis of tert-butyl 4-(2-(4-bromophenoxy)ethyl)piperazine-1-carboxylate

30

[0176]


- 5 **[0177]** A mixture of tert-butyl 4-(2-hydroxyethyl)piperazine-1-carboxylate (400 mg, 1.73 mmol), MsCl (596 mg, 5.19 mmol) and triethylamine (524 mg, 5.19 mmol) in DCM (25 mL) was stirred at RT for 2 h. The solution was diluted with DCM, and washed with sat. NaHCO₃ and brine. The organic layer was concentrated. The residue was dissolved in DMF (15 mL), followed by addition of 4-bromophenol (451 mg, 2.61 mmol) and Cs₂CO₃ (1.70 g, 5.22 mmol) at RT. The resultant mixture was stirred at 100 °C for 18 h. After that, the solution was diluted with EA and washed with brine. The organic layer was concentrated and purified by silica gel chromatography to get the title compound (320 mg, yield 48%) as a colorless oil. MS (ES+) C₁₇H₂₅BrN₂O₃ requires: 384, 386 found: 385, 387 [M+H]⁺.

- 15 **Step 2: Synthesis of tert-butyl 4-(2-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)ethyl)piperazine-1-carboxylate**

[0178]

- 20 **[0179]** A mixture of tert-butyl 4-(2-(4-bromophenoxy)ethyl)piperazine-1-carboxylate (320 mg, 830 μmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (271 mg, 1.07 mmol), Pd(dppf)Cl₂ (60.6 mg, 83.0 μmol) and KOAc (325 mg, 3.32 mmol) in dioxane/water (10 mL) was purged with N₂, and stirred at 100 °C for 18 h under N₂. After that, the solution was cooled and concentrated. The residue was purified by silica gel chromatography (EA/PE = 1/3) to get the title compound (300 mg, yield 84%) as a brown oil. MS (ES+) C₂₃H₂₂BN₂O₅ requires: 432, found: 433 [M+H]⁺.

Step 3: Synthesis of tert-butyl 4-(2-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)phenoxy)ethyl)piperazine-1-carboxylate


[0180]

- 30 **[0181]** A mixture of (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-

yl)(cyclopropyl)methanone (200 mg, 572 μmol), tert-butyl 4-(2-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)ethyl)piperazine-1-carboxylate (271 mg, 629 μmol), Pd(dppf)Cl₂ (41.8 mg, 57.2 μmol), and K₂CO₃ (314 mg, 2.28 mmol) in dioxane (5 mL) and water (1 mL) was purged with N₂, and then stirred at 100 °C for 18 h under N₂. After that, the solution was cooled and concentrated. The residue was purified by silica gel chromatography to get the title compound (160 mg, yield 49%) as a yellow oil. MS (ES+) C₃₂H₄₂N₆O₄ requires: 574, found: 575 [M+H]⁺.

Step 4: Synthesis of cyclopropyl(4-(6-(4-(2-(piperazin-1-yl)ethoxy)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 273)


[0182]



[0183] A mixture of tert-butyl 4-(2-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)phenoxy)ethyl)piperazine-1-carboxylate (80 mg, 139 μmol) in HCl/dioxane (4 N, 1 mL) was stirred at RT for 2 h. After that, the solution was concentrated to afford the title compound (80 mg, crude) as a yellow solid. MS (ES+) C₂₇H₃₄N₆O₂ requires: 474 found: 475 [M+H]⁺.

Step 5: Synthesis of cyclopropyl(4-(6-(4-(2-(4-ethylpiperazin-1-yl)ethoxy)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 390)

[0184]




[0185] A mixture of tert-butyl 4-(2-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)phenoxy)ethyl)piperazine-1-carboxylate (40 mg, 84.2 μmol), CH₃CHO (11.0 mg, 252 μmol), NaBH₃CN (7.93 mg, 126 μmol) and AcOH (5.05 mg, 84.2 μmol) in DCM (5 mL) and MeOH (2 mL) was stirred at RT for 2 h. After that, the solution was concentrated and purified by Prep-HPLC to give the title compound (12 mg, 28%) as a yellow solid. MS (ES+) C₂₉H₃₈N₆O₂ requires: 502 found: 503 [M+H]⁺.

Example 6. Synthesis of cyclopropyl(4-(6-(1-(5,5-difluoropiperidin-3-yl)-1H-pyrazol-4-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 194)

Step 1: Synthesis of tert-butyl 5-oxo-5,6-dihydropyridine-1(2H)-carboxylate


[0186]



- 10 [0187] To a solution of 1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-3-ol (300 mg, 1.50 mmol) in DCM (25 mL) was added Dess-Martin Oxidant (1.27 g, 3.00 mmol). The reaction solution was stirred at RT for 12 hours and then filtered. The filtrate was washed with saturated aqueous Na₂CO₃ (50 mL) and concentrated in *vacuo*. The residue was purified by silica gel column chromatography (PE/EtOAc (v/v) = 2/1) to give the title compound as a colorless oil (280mg, yield 94%). MS (ES⁺) C₁₀H₁₅NO₃ requires: 197, found: 142 [M+H-56]⁺.
- 15

Step 2: Synthesis of (4-(6-(1H-pyrazol-4-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone


[0188]



- 20 [0189] A mixture of tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylate (335 mg, 1.14 mmol), (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (400 mg, 1.14 mmol), Na₂CO₃ (362 mg, 3.42 mmol) and Pd(t-Bu₃P)₂ (116 mg, 0.228 mmol) in dioxane/water (v/v = 3:1, 10 mL) was degassed with N₂ three times, and then stirred at 85 °C for 12 hours. The reaction mixture was cooled and evaporated in *vacuo*. The residue was purified by flash column (PE:EA = 3:1 to 1:3) to give the title compound as a yellow solid (340 mg, yield 88%). MS (ES⁺) C₁₈H₂₀N₆O requires: 336, found: 337 [M+H]⁺.
- 25

- 30 **Step 3: Synthesis of tert-butyl 3-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)-1H-pyrazol-1-yl)-5-oxopiperidine-1-carboxylate**


[0190]



[0191] A mixture of tert-butyl 5-oxo-5,6-dihydropyridine-1(2H)-carboxylate (100 mg, 0.51 mmol) and (4-(6-(1H-pyrazol-4-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (180 mg, 0.54 mmol) in MeCN (5 mL) in a flask was evaporated in *vacuo* at 50 °C. The residue was diluted with 5 mL of MeCN, and then evaporated to dryness. The dilution/evaporation was repeated three times. Purification by flash column (PE/EA to EA) gave the title compound as an off-white solid (140 mg, yield 49%). MS (ES+) C₂₈H₃₅N₇O₄ requires: 533, found: 534 [M+H]⁺.


Step 4: Synthesis of tert-butyl 5-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)-1H-pyrazol-1-yl)-3,3-difluoropiperidine-1-carboxylate

[0192]



[0193] To a solution of tert-butyl 3-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)-1H-pyrazol-1-yl)-5-oxopiperidine-1-carboxylate (60 mg, 0.1124 mmol) in DCM (4 mL) was added DAST (180 mg, 1.12 mmol) at 0 °C. The reaction mixture was then stirred at 0 °C for 10 minutes. Quenched by water and extracted with DCM. The organic layers were evaporated and purified by flash column (PE/EA = 1:4 to 4:1) to give the title compound as a yellow solid (12 mg, yield 19%). MS (ES+) C₂₈H₃₅F₂N₇O₃ requires: 555, found: 556 [M+H]⁺.


Step 5: Synthesis of cyclopropyl(4-(6-(1-(5,5-difluoropiperidin-3-yl)-1H-pyrazol-4-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone

[0194]

[0195] To a solution of tert-butyl 5-(4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)-1H-pyrazol-1-yl)-3,3-difluoropiperidine-1-carboxylate (12 mg, 0.02159 mmol) in DCM (2.0 mL) was added TFA (1.0 mL). The reaction solution was stirred at 25 °C for 1 hour. The reaction solution was concentrated in vacuo. The residue was dissolved in DCM and neutralized by sat. aqueous NaHCO₃. The organic layer was washed with brine, dried and concentrated to give the title compound (8.5 mg, yield 86%) as a yellow solid. MS (ES⁺) C₂₃H₂₇F₂N₇O requires: 455, found 456 [M+H]⁺.

Example 7. Synthesis of (S)-cyclopropyl(4-(6-(4-(morpholin-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 130) and (R)-cyclopropyl(4-(6-(4-(morpholin-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 131).


Step 1: Synthesis of cyclopropyl(4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone

[0196]

[0197] A mixture of (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (11.6 g, 33.2 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (16.8 g, 66.4 mmol), Pd(dppf)Cl₂ (3.63 g, 4.97 mmol) and KOAc (9.76 g, 99.6 mmol) in CH₃CN (300 mL) was purged with N₂ and then stirred at 65 °C for 24 hrs under N₂. The reaction mixture was concentrated and purified by flash chromatography (PE/EA = 10:1 to 2:1) to afford the title compound (11.2 g, 80% yield) as a yellow solid. MS (ES⁺) C₂₁H₂₉BN₄O₃ requires: 396, found: 397 [M+H]⁺.

Step 2: Synthesis of (S)-cyclopropyl(4-(6-(4-(morpholin-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 130) and (R)-cyclopropyl(4-(6-(4-(morpholin-2-yl)phenyl)pyrrolo(1,2-b)pyridazin-4-yl)piperazin-1-yl)methanone (Compound 131)


5 **[0198]**



[0199] A mixture of cyclopropyl(4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (300 mg, 757 μ mol), 2-(4-bromophenyl)morpholine (183 mg, 757 μ mol), Pd(dppf)Cl₂-CH₂Cl₂ (61.7 mg, 75.7 μ mol) and K₂CO₃ (208 mg, 1.51 mmol) in dioxane/water (3 mL/ 0.5 mL) was purged with N₂, and then stirred at 100 °C for 16 hrs under N₂. The mixture was concentrated and purified by flash column chromatography (DCM/MeOH = 10:1) to give a yellow oil (300 mg, crude). Chiral separation was performed to afford the title compounds using a CE-3 column. Mobile Phase: Hexane/EtOH/DEA =30/70/0.1; flow rate: 50 mL/min; 0.4 ml injection; 25 minute run time; sample solution: 3.2g in 30mL MeOH; elution measured at 214 and 254 nm using a Gilson-281. Compound 130: (56.3 mg, 17% yield) as a yellow solid MS (ES+) C₂₅H₂₉N₅O₂ requires: 431, found: 432 [M+H]⁺. Compound 131: (32.6 mg, 10% yield) as a yellow solid. MS (ES+) C₂₅H₂₉N₅O₂ requires: 431, found: 432 [M+H]⁺.

Example 8. Synthesis of cyclopropyl(4-(6-(4-(4-ethylmorpholin-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 214)

[0200]




[0201] A mixture of cyclopropyl(4-(6-(4-(morpholin-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (100 mg, 231 μ mol) and acetaldehyde (40.7 mg, 924 μ mol) in DCM/MeOH/CH₃COOH (2 mL/2 mL/0.5 mL) was stirred at 25 °C for 30 min, followed by addition of NaBH₃CN (72.2 mg, 1.15 mmol). The mixture was stirred at 25 °C for 2 hours. The mixture was purified by Prep-HPLC to afford the title compound (8.4 mg, yield 8%) as a yellow solid. MS (ES+) C₂₇H₃₃N₅O₂ requires: 459, found: 460 [M+H]⁺.

Example 9. Synthesis of cyclopropyl(4-(6-(4-(2-(trifluoromethyl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 374)

Step 1: Synthesis of 4-benzyl-1-(4-bromophenyl)-2-(trifluoromethyl)piperazine


[0202]



[0203] A mixture of 1-benzyl-3-(trifluoromethyl)piperazine (200 mg, 818 μ mol), 1-bromo-4-iodobenzene (461 mg, 1.63 mmol), bis(tri-*t*-butylphosphine)palladium (83.3 mg, 163 μ mol) and cesium carbonate (798 mg, 2.45 mmol) in toluene (4 mL) was purged with N₂ and stirred at 80 °C overnight. TLC and LCMS showed completed reaction. The mixture was cooled to RT and concentrated. The residue was purified by silica gel column (PE/EA = 10/1) to give the title compound (40 mg, yield 12%) as a yellow solid. MS (ES+) C₁₈H₁₈BrF₃N₂ requires: 398, found: 399 [M+H]⁺.

Step 2: Synthesis of (4-(6-(4-(4-benzyl-2-(trifluoromethyl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone

[0204]




[0205] A mixture of 4-benzyl-1-(4-bromophenyl)-2-(trifluoromethyl)piperazine (60 mg, 150 μ mol), cyclopropyl(4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (118 mg, 300 μ mol), Pd(dppf)Cl₂ (21.9 mg, 30.0 μ mol) and potassium carbonate (62.1 mg, 450 μ mol) in dioxane/water (4 mL) was purged with N₂ and stirred at 100 °C for 2 h. TLC and LCMS showed completed reaction. The mixture was cooled to RT and concentrated.

The residue was purified by silica gel column (PE/EA = 2/1) to give the title compound (80 mg, crude) as a yellow solid. MS (ES+) $C_{33}H_{35}F_3N_6O$ requires: 588, found: 589 $[M+H]^+$.

Step 3: Synthesis of cyclopropyl(4-(6-(4-(2-(trifluoromethyl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 374)

[0206]




[0207] A mixture of (4-(6-(4-(4-benzyl-2-(trifluoromethyl)piperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (70 mg, 118 μ mol) and Pd/C (14 mg, 20% wt) in i-PrOH (4 mL) was stirred at 45 °C overnight under hydrogen (balloon). The mixture was filtered and concentrated. The residue was purified by Prep-HPLC to give the title compound (1.6 mg, yield 3%) as a white solid. MS (ES+) $C_{26}H_{29}F_3N_6O$ requires: 498, found: 499 $[M+H]^+$.

Example 10. Synthesis of N-ethyl 4-(6-(4-(1-ethylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 230).

Step 1: Synthesis of 6-(4-(1-ethylpiperidin-4-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine

[0208]




[0209] A mixture of 6-bromo-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine hydrochloride (1.2 g, 3.77 mmol), 1-ethyl-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine (1.78 g, 5.65 mmol), K_2CO_3 (1.57 g, 11.3 mmol) and $Pd(t-Bu_3P)_2$ in dioxane/water (20 mL/5 mL) was degassed with nitrogen three times. The mixture was then heated to 70 °C and the temperature maintained overnight. The reaction mixture was cooled to RT and concentrated to give a residue, which was

purified by silica gel chromatography to afford the title compound (1.2 g, yield 82%) as a yellow oil. MS (ES+) $C_{24}H_{31}N_5$ requires: 389, found: 390 $[M+H]^+$.

Step 2: Synthesis of ethyl 4-(6-(4-(1-ethylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 230).

5 [0210]




10 [0211] To a solution 6-(4-(1-ethylpiperidin-4-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine (50 mg, 128 μ mol) and DIPEA (82 mg, 640 μ mol) in DCM (10 mL) was added ethyl ethyl chloroformate (41.5 mg, 384 μ mol). The reaction mixture was stirred at RT for 3 h. The mixture was concentrated under reduced pressure. The residue was purified by Prep-HPLC to afford the title compound (9 mg, yield 15%). MS (ES+) $C_{27}H_{35}N_5O_2$ requires: 461, found: 462 $[M+H]^+$.

15 **Example 11. Synthesis of N-ethyl-4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxamide (Compound 275)**

Step 1: Synthesis of tert-butyl-4-(6-bromo-pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate


[0212]



20


25 [0213] A mixture of 6-bromopyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate (200 g, 579.7 mmol), tert-butyl piperazine-1-carboxylate (113.2 g, 609 mmol) and triethylamine (176 g, 1740 mmol) in NMP (180 mL) was stirred at 100 °C for 1 hr. Monitored by LC-MS, the reaction was completed. The mixture was diluted with EtOAc, washed with water and brine, and dried over sodium sulfate. The organic layer was concentrated and purified by flash chromatography (silica gel, 10-40% EtOAc in PE) to afford the title compound (220.0 g, crude). MS (ES+) $C_{16}H_{21}BrN_4O_2$ require: 380, 382, found: 381, 383 $[M+H]^+$.

30 **Step 2: Synthesis of tert-butyl-4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate**

[0214]

[0215] A mixture of tert-butyl 4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (220 g crude, 0.58 mol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (294 g, 1.158 mol), Pd(dppf)Cl₂-CH₂Cl₂ (42.4 g, 0.058 mol) and KOAc (170.2 g, 1.737 mol) in CH₃CN (500 mL) was purged with N₂ and stirred at 65 °C for 24 hrs under N₂. Monitored by LC-MS, the reaction was completed. The mixture was cooled to RT, concentrated and purified by flash column chromatography (Petroleum ether/ethyl acetate = 10: 1 to 2: 1) to afford the title compound (189.0 g, 76%) as a white solid. MS (ES+) C₂₂H₃₃BN₄O₄ requires: 428, found: 429 [M+H]⁺.


Step 3: Synthesis of tert-butyl tert-butyl 4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0216]

[0217] To a solution of 4-(4-bromophenyl)-1-isopropylpiperidine (300 mg, 1.06 mmol) and tert-butyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (550 mg, 1.59 mmol) in dioxane/water (30 mL/10 mL) was added Pd(t-Bu₃P)₂ (108.3 mg, 0.212 mmol) and Na₂CO₃ (337.1 mg, 3.18 mmol) at RT under nitrogen. The resulting mixture was heated at 75 °C for 3 hours; LCMS showed that the reaction was completed. Water (50 mL) was added and extracted with EtOAc (50 mLx2). The organic layer was washed with brine (50 mL), dried over Na₂SO₄, and concentrated to dryness to afford crude product, which was purified with flash column (DCM/MeOH, 10: 1) to give the title compound (346 mg, yield 65%) as a white powder. MS (ES+) C₃₀H₄₁N₅O₂, requires: 503, found: 504 [M+H]⁺.

Step 4: Synthesis of 6-(4-(1-isopropylpiperidin-4-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine hydrochloride


[0218]



[0219] To a solution of tert-butyl 4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (346 mg, 0.69 mmol) in dioxane (10 mL) was added 6 M HCl/dioxane (3 mL) at RT. The resulting mixture was stirred for 6 hours; LCMS showed that the reaction was completed. The mixture was then evaporated to dryness to afford the title product (282 mg, 93% yield) as a white solid. MS (ES+) C₂₅H₃₃N₅ requires: 403, found: 404 [M+H]⁺.

Step 5: Synthesis of N-ethyl-4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxamide (Compound 275)


[0220]



[0221] To a solution of 6-(4-(1-isopropylpiperidin-4-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine hydrochloride (282 mg, 0.64 mmol) in DCM (10 mL) was added triethylamine (0.5 mL) and isocyanatoethane (0.1 mL) at 0 °C. The resulting mixture was stirred at RT for 30 min; LCMS indicated that the reaction was completed. Extracted with EtOAc (50 mLx2) after water (50 mL) was added. The organic layer was washed with brine (50 mL), dried over Na₂SO₄, and concentrated to dryness to afford crude product which was purified by Prep-HPLC to give the title compound (89.9 mg, yield 30%) as a white powder. MS (ES+) C₂₇H₃₇N₇O, requires: 475, found: 476 [M+H]⁺.


Example 12. Synthesis of cyclopropyl(4-(6-(4-(1-isopropylazetidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 150).

Step 1: Synthesis of tert-butyl 4-(6-(4-(1-isopropylazetidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0222]

- 5 **[0223]** To a solution of tert-butyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (680 mg, 1.59 mmol) in dioxane/water (15 mL) was added 3-(4-bromophenyl)-1-isopropylazetidine (480 mg, 1.9 mmol), Pd(t-Bu₃P)₂ (81 mg, 0.16 mmol) and Na₂CO₃ (337 mg, 3.18 mmol) at room temperature under nitrogen. The resulting mixture was stirred at 70 °C for 3 h
- 10 under nitrogen. Water was added and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated to dryness to afford crude compound, which was purified by flash chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (430 mg, 57% yield) as a light yellow powder. MS (ES⁺) C₂₈H₃₇N₅O₂ requires: 475, found 476 [M+H]⁺.


- 15 **Step 2: Synthesis of 6-(4-(1-isopropylazetidin-3-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine**

[0224]

- 20 **[0225]** To a solution of tert-butyl 4-(6-(4-(1-isopropylazetidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (430 mg, 0.91 mmol) in dioxane (10 mL) was added HCl/dioxane (3 mL, 4.0 M) at RT. The resulting mixture was stirred for 5 hours. LC-MS showed that the reaction was completed. The mixture was evaporated to dryness to afford the title compound (360 mg, 99% yield) as a white powder. MS
- 25 (ES⁺) C₂₃H₂₉N₅ requires: 375, found 376 [M+H]⁺.

Step 3: Synthesis of cyclopropyl(4-(6-(4-(1-isopropylazetidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone

[0226]




- 5 **[0227]** To a solution of 6-(4-(1-isopropylazetidin-3-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine (120 mg, 0.32 mmol) in DMF (10 mL) was added cyclopropanecarboxylic acid (33 mg, 0.38 mmol) and triethylamine (80 mg, 0.8 mmol) at RT, followed by addition of HATU (144 mg, 0.38 mmol). The resulting mixture was stirred at RT for 2 h, diluted with water and extracted with EtOAc. The
- 10 combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated to dryness to afford crude product, which was purified by flash chromatography on silica gel eluting with DCM/MeOH (5 : 1) to give the title compound (101 mg, 72% yield) as a white powder. MS (ES⁺) C₂₇H₃₃N₅O requires: 443, found 444 [M+H]⁺.

15 **Example 13. Synthesis of ethyl 4-(6-(5-(4-isopropylpiperazin-1-yl)pyrazin-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 448)**

Step 1: Synthesis of tert-butyl 4-(5-bromopyrazin-2-yl)piperazine-1-carboxylate


20 [0228]



- 25 **[0229]** A mixture of 2,5-dibromopyrazine (2 g, 8.40 mmol), tert-butyl piperazine-1-carboxylate (1.86 g, 10.0 mmol) and DIPEA (1.62 g, 12.6 mmol) in NMP (40 mL) was stirred at 110 °C for 2 h. Diluted with ethyl acetate, washed with water and brine, dried over Na₂SO₄, and concentrated to give the title product (2.8 g, yield 97%). MS (ES⁺) C₁₃H₁₉BrN₄O₂ requires: 342, found: 343 [M+H]⁺.

Step 2: Synthesis of 2-bromo-5-(piperazin-1-yl)pyrazine


[0230]



- 5 **[0231]** To a solution of tert-butyl 4-(5-bromopyrazin-2-yl)piperazine-1-carboxylate (2.8 g, 8.15 mmol) in DCM (20 mL) was added TFA (5 mL). The mixture was stirred at 20 °C for 1 h. Concentrated and diluted with ethyl acetate. Adjusted pH to 7-8 with sat. NaHCO₃ solution. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated to give the title product (1.98 g, crude). MS (ES+) C₈H₁₁BrN₄ requires: 242, found: 243 [M+H]⁺.

10 **Step 3: Synthesis of 2-bromo-5-(4-isopropylpiperazin-1-yl)pyrazine**


[0232]



- 15 **[0233]** A mixture of 2-bromo-5-(piperazin-1-yl)pyrazine (458 mg, 1.88 mmol), 2-iodopropane (479 mg, 2.82 mmol) and K₂CO₃ (518 mg, 3.76 mmol) in MeCN (10 mL) was stirred at 60 °C overnight. Concentrated and purified by silica gel column (MeOH/EA = 1:10) to give the title product (388 mg, yield 72%). MS (ES+) C₁₁H₁₇BrN₄ requires: 284, found: 285[M+H]⁺.

20 **Step 4: Synthesis of tert-butyl 4-(6-(5-(4-isopropylpiperazin-1-yl)pyrazin-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate**

[0234]




- [0235]** A mixture of 2-bromo-5-(4-isopropylpiperazin-1-yl)pyrazine (350 mg, 1.22

mmol), tert-butyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (625 mg, 1.46 mmol), Na₂CO₃ (193 mg, 1.83 mmol) and Pd(t-Bu₃P)₂ (62.3 mg, 122 μmol) in dioxane/water (20 mL) was purged with N₂, and then stirred at 90 °C overnight under nitrogen. Concentrated and purified by silica gel column (MeOH/EA = 1:10) to give the title product (388 mg, yield 63%). MS (ES+) C₂₇H₃₈N₈O₂ requires: 506, found: 507 [M+H]⁺.

Step 5: Synthesis of 6-(5-(4-isopropylpiperazin-1-yl)pyrazin-2-yl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine hydrochloride


[0236]



[0237] To a solution of tert-butyl 4-(6-(5-(4-isopropylpiperazin-1-yl)pyrazin-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (388 mg, 765 μmol) in DCM (4 mL) was added HCl/dioxane (4 mL, 4 M). The mixture was stirred at 20 °C overnight and then concentrated to give the title product (338 mg, crude). MS (ES+) C₂₂H₃₁ClN₈ requires: 406, found: 407 [M+H]⁺.

Step 6: Synthesis of ethyl 4-(6-(5-(4-isopropylpiperazin-1-yl)pyrazin-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 448).

[0238]




[0239] To a solution of 6-(5-(4-isopropylpiperazin-1-yl)pyrazin-2-yl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine hydrochloride (100 mg, 225 μmol) and DIPEA (87.0 mg, 675 μmol) in DCM (10 mL) was added ethyl chloroformate (48.8 mg, 450 μmol). The mixture was stirred at 20 °C for 1 h and then concentrated and purified by Prep-HPLC to give the title product (30.0 mg, yield 28%). MS (ES+) C₂₅H₃₄N₈O₂ requires: 478, found: 479 [M+H]⁺.

Example 14. Synthesis of 1-(4-(6-(4-(1-ethylpyrrolidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carbonyl)azetidine-3-carbonitrile (Compound 300)

Step 1: Synthesis of tert-butyl 4-(6-(4-(1-ethylpyrrolidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0240]



5


[0241] A solution of 3-(4-bromophenyl)-1-ethylpyrrolidine (200 mg, 786 μmol) in dioxane/water (5 mL) was added tert-butyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (336 mg, 786 μmol), $\text{Pd}(\text{t-Bu}_3\text{P})_2$ (40.1 mg, 78.6 μmol) and Cs_2CO_3 (763 mg, 2.35 mmol) was degassed with nitrogen, and then heated at 60 $^\circ\text{C}$ under MW for 1 h. The reaction mixture was cooled to RT and concentrated. The residue was purified by flash chromatography to afford the title compound (300 mg, yield 80%). MS (ES+) $\text{C}_{28}\text{H}_{37}\text{N}_5\text{O}_2$, requires: 475, found: 476 $[\text{M}+\text{H}]^+$.

10

Step 2: Synthesis of 6-(4-(1-ethylpyrrolidin-3-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine

15

[0242]




[0243] To a solution of tert-butyl 4-(6-(4-(1-ethylpyrrolidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (300 mg, 630 μmol) in DCM (8 mL) was added TFA (5 mL). The reaction mixture was stirred at 20 $^\circ\text{C}$ for 18 h. The mixture was concentrated to give the title compound (230 mg, yield 97%). MS (ES+) $\text{C}_{28}\text{H}_{34}\text{N}_6\text{O}_2$, requires: 375, found: 376 $[\text{M}+\text{H}]^+$.

20

Step 3: Synthesis of 1-(4-(6-(4-(1-ethylpyrrolidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carbonyl)azetidine-3-carbonitrile

25


[0244]



[0245] To a solution of 6-(4-(1-ethylpyrrolidin-3-yl)phenyl)-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine (200 mg, 532 μmol) in DMF (5 mL) was added 4-nitrophenyl 3-cyanoazetidine-1-carboxylate (131 mg, 532 μmol) and Na_2CO_3 (168 mg, 1.59 mmol). The reaction mixture was stirred at 20 °C for 18 hrs. Water was added and extracted with DCM/MeOH(10/1). The mixture was concentrated to give a residue, which was purified by Prep-HPLC to give the title compound (130 mg, yield 51%). MS (ES+) $\text{C}_{28}\text{H}_{33}\text{N}_7\text{O}$, requires: 483, found: 484[M+H]⁺.


Example 15. Synthesis of (4-(6-(4-(2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (Compound 260).

Step 1: Synthesis of 2-oxa-5,8-diazaspiro[3.5]nonane



[0247] To a mixture of tert-butyl 2-oxa-5,8-diazaspiro[3.5]nonane-8-carboxylate (100 mg, 438 μmol) in DCM (3 mL) was added TFA ((1 mL). The reaction mixture was stirred at 25 °C for 16 h. NH_3 (7 N in MeOH) was added to adjust pH to 8-9. The reaction mixture was concentrated in vacuo to afford the title compound (56 mg, crude) as a white solid MS (ES+) $\text{C}_6\text{H}_{12}\text{N}_2\text{O}$ requires: 128, found 129 [M+H]⁺.

Step 2: Synthesis of (4-(6-(4-(2-oxa-5,8-diazaspiro[3.5]nonan-8-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (Compound 260)




[0249] A mixture of 2-oxa-5,8-diazaspiro[3.5]nonane (30 mg, 234 μmol), (4-(6-(4-bromophenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (129 mg, 304 μmol), $\text{Pd}[(t\text{-Bu})_3\text{P}]_2$ (11.9 mg, 23.4 μmol) and Na_2CO_3 (49.6 mg, 468 μmol) in toluene (3 mL) was degassed with nitrogen and stirred at 100 °C for 16 h under N_2 . LCMS showed the reaction was completed. Concentrated in vacuo, the

residue was purified by Prep-HPLC to afford the title compound (7.2 mg, yield 6%) as a yellow solid MS (ES+) $C_{27}H_{32}N_6O_2$ requires: 472, found 473 $[M+H]^+$.

Example 16. Synthesis of 2-(6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)hexahydropyrrolo[1,2-a]pyrazin-6(7H)-one (Compound 157)

5 Step 1: Synthesis of 1-(4-bromophenyl)-4-ethylpiperazine

[0250]




10

[0251] A mixture of 1-(4-bromophenyl)piperazine (40 g, 165 mmol), bromoethane (17.9 g, 165 mmol) and triethylamine (16.6 g, 165 mmol) in THF (500 ml) was stirred at 60 °C overnight. The reaction was cooled to RT, diluted with EA and washed with water. The organic layer was concentrated to give the title compound (50 g, crude).

15 MS (ES+) $C_{12}H_{17}BrN_2$ requires: 268, found 269 $[M+H]^+$.

Step 2: Synthesis of 1-ethyl-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine

[0252]




20

[0253] A mixture of 1-(4-bromophenyl)-4-ethylpiperazine (50 g, 185 mmol), KOAc (54.3 g, 554 mmol) and Pd(dppf)Cl₂ (13.5 g, 18.5 mmol) in dioxane (500 mL) was purged with N₂ and stirred at 60 °C overnight. The reaction mixture was cooled, concentrated and purified by silica gel column (PE/EA = 5/1 to EA) to give the title compound (40 g, 69.5%). MS (ES+) $C_{18}H_{29}BN_2O_2$ requires: 316, found 317 $[M+H]^+$.

25

Step 3: Synthesis of 6-bromo-4-((2-(trimethylsilyl)ethoxy)methoxy)pyrrolo[1,2-b]pyridazine

[0254]




30

[0255] To a mixture of 6-bromopyrrolo[1,2-b]pyridazin-4-ol (22.0 g, 103 mmol) and triethylamine (31.2 g, 309 mmol) in THF (100 mL) was added (2-(chloromethoxy)ethyl)trimethylsilane (20.5 g, 123 mmol) dropwise at 0 °C. The reaction mixture was stirred at 20 °C for 1 h. The mixture was concentrated, diluted

with EA and washed with water. The organic layer was concentrated and purified by flash column chromatography (PE/EA = 5:1) to afford the title compound (31g, yield 87%) as a yellow oil MS (ES+) $C_{13}H_{19}BrN_2O_2Si$ requires: 342, found 343 $[M+H]^+$.

Step 4: Synthesis of 6-(4-(4-ethylpiperazin-1-yl)phenyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)pyrrolo[1,2-b]pyridazine


[0256]



[0257] A mixture of 6-bromo-4-((2-(trimethylsilyl)ethoxy)methoxy)pyrrolo[1,2-b]pyridazine (5 g, 14.5 mmol), 1-ethyl-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine (5.94 g, 18.8 mmol), K_2CO_3 (6.0 g, 43.5 mmol) and $Pd[(t-Bu)_3P]_2$ (370 mg, 725 μ mol) in dioxane/water (30 mL, 4/1) was purged with N_2 and then stirred at 70 °C for 4 h under N_2 . The mixture was purified by flash column chromatography (PE/EA = 10:1 to 1:2) to afford the title compound (6 g, yield 91%) as a grey solid. MS (ES+) $C_{25}H_{36}N_4O_2Si$ requires: 452, found 453 $[M+H]^+$.

Step 5: Synthesis of 6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-ol


[0258]



[0259] To a mixture of 6-(4-(4-ethylpiperazin-1-yl)phenyl)-4-((2-(trimethylsilyl)ethoxy)methoxy)pyrrolo[1,2-b]pyridazine (8.0 g, 17.6 mmol) in dioxane (20 mL) was added HCl (4 N in dioxane, 40 mL). The mixture was stirred at 25 °C for 1 h. The mixture was concentrated in *vacuo* to afford the title compound (4.77 g, crude) as a yellow solid. MS (ES+) $C_{19}H_{22}N_4O$ requires: 322, found: 323 $[M+H]^+$.

Step 6: Synthesis of 6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo(1,2-b)pyridazin-4-yl trifluoromethanesulfonate

[0260]




[0261] To a solution of 6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-ol (3.0 g, 9.30 mmol) in DCM (30 mL) at 0 °C was added triethylamine (2.82 g, 27.9 mmol), followed by addition of 1,1,1-trifluoro-N-phenyl-N-

((trifluoromethyl)sulfonyl)methanesulfonamide (3.96 g, 11.1 mmol). The reaction mixture was stirred at 0 °C for 2 h. The mixture was diluted with DCM and washed with brine. The organic layer was evaporated and purified by flash column (PE:EA = 5:1) to afford the title compound (1.9 g, yield 45%) as a yellow solid MS (ES+)

5 C₂₀H₂₁F₃N₄O₃S requires: 454, found 455 [M+H]⁺.

Step 7: Synthesis of 2-(6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo(1,2-b)pyridazin-4-yl)hexahydropyrrolo[1,2-a]pyrazin-6(7H)-one

[0262]



10

[0263] A mixture of hexahydropyrrolo[1,2-a]pyrazin-6(2H)-one (30 mg, 214 μmol), 6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate (116 mg, 256 μmol) and triethylamine (64.9 mg, 642 μmol) in NMP (2 mL) was stirred at 100 °C for 1 h. The mixture was concentrated and purified by Prep-HPLC to afford the title compound (4.0 mg, yield 4%) as a white solid MS (ES+) C₂₆H₃₂N₆O requires: 444, found 445 [M+H]⁺.

15

Example 17. Synthesis of cyclopropyl(2-(difluoromethyl)-4-(6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 403)

20 **Step 1: Synthesis of 2-(difluoromethyl)piperazine**

[0264]




25

[0265] To a solution of tert-butyl 3-(difluoromethyl)piperazine-1-carboxylate (30 mg, 0.127 mmol) in dioxane (2.0 mL) was added HCl/dioxane (4 N, 1.0 mL). The reaction solution was stirred at 25 °C for 1 h. The reaction mixture was concentrated. The residue was used in next step directly without any further purification.

Step 2: Synthesis of 4-(3-(difluoromethyl)piperazin-1-yl)-6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazine


[0266]



[0267] A mixture of 6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl trifluoromethanesulfonate (60 mg, 0.13 mmol) and 2-(difluoromethyl)piperazine (17 mg, 0.13 mmol) and triethylamine (40 mg, 0.40 mmol) in NMP (3 mL) was stirred at 100 °C for 2 h. The reaction solution was used in next step directly without any further purification. MS (ES+) $C_{24}H_{30}F_2N_6$ requires: 440, found 441 $[M+H]^+$.

Step 3: Synthesis of cyclopropyl(2-(difluoromethyl)-4-(6-(4-(4-ethylpiperazin-1-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 403)


[0268]




[0269] To above reaction mixture was added triethylamine (40 mg, 0.40 mmol) and cyclopropanecarbonyl chloride (27 mg, 0.26 mmol). The mixture was stirred at RT for 2 h. The reaction mixture was concentrated and purified by Prep-HPLC to give the title product as a yellow solid (3.8 mg, yield 7.6%). MS (ES+) $C_{28}H_{34}F_2N_6O$ requires: 508, found 509 $[M+H]^+$.

Example 18. Synthesis of cyclopropyl(4-(6-(4-(4,5-dihydro-1H-imidazol-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 111)

[0270]




Step 1: Synthesis of 4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo(1,2-b)pyridazin-6-yl)benzaldehyde

[0271]

[0272] A mixture of cyclopropyl(4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (250 mg, 0.63 mmol), 4-bromobenzaldehyde (120 mg, 0.65 mmol), sodium carbonate (200 mg, 1.89 mmol) and Pd(t-Bu₃P)₂ (65 mg, 0.13 mmol) in dioxane/water (15 mL) was purged with N₂, and then stirred at 75 °C for 3 h. Cooled to RT and diluted with EtOAc. The organic layer was washed with water and brine, concentrated and purified by flash column (PE/EtOAc = 10/1 to 5/1) to give the title product (180 mg, crude). MS (ES+) C₂₂H₂₂N₄O₂ requires: 374, found 375 [M+H]⁺.


Step 2: Synthesis of cyclopropyl(4-(6-(4-(4,5-dihydro-1H-imidazol-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 111)

[0273]

[0274] To a solution of 4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)benzaldehyde (180 mg, 0.48 mmol) in t-BuOH (20 mL) was added potassium carbonate (200 mg, 1.44 mmol) and iodine (122 mg, 0.48 mmol). The reaction mixture was heated at 70 degrees for 2 h. Quenched by sat. aq. Na₂S₂O₃ (20 mL) and diluted with EtOAc. The organic layer was washed with water and brine, concentrated and purified by column (DCM/MeOH = 20/1 to 5/1) to give the title product (98.99 mg, 49.7 %). MS (ES+) C₂₄H₂₆N₆O requires: 414, found 415 [M+H]⁺.


Example 19. Synthesis of cyclopropyl(4-(6-(4-(4,5-dihydro-1H-imidazol-2-ylamino)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 121)

Step 1: Synthesis of (4-(6-(4-aminophenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone

[0275]


- 5 **[0276]** A mixture of (4-(6-bromopyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (250 mg, 0.72 mmol), 4-aminophenylboronic acid (150 mg, 1.08 mmol), K_2CO_3 (300 mg, 2.16 mmol) and $Pd(dppf)Cl_2$ (100 mg) in 1,4-dioxane (20 mL) and water (4 mL) was purged with N_2 , and then stirred at 100 °C for 6 h. Cooled to RT and diluted with DCM. The organic layer was washed with water and
- 10 brine, concentrated and purified by flash column (DCM/MeOH = 100/1 to 20/1) to give the title product (130 mg, yield: 50%). MS (ES+) $C_{21}H_{23}N_5O$ requires: 361, found 362 $[M+H]^+$.

Step 2: Synthesis of dimethyl 4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)phenylcarbonimidodithioate

15 **[0277]**

- [0278]** To a solution of (4-(6-(4-aminophenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)(cyclopropyl)methanone (100 mg, 0.28 mmol) in DMF (6 mL) was added CS_2 (1 mL). The reaction mixture was added 6 ml of NaOH in water (1 N) and stirred at RT
- 20 for 30 min. The reaction mixture was added methyl iodide (0.3 ml) and stirred overnight at RT. The mixture was extracted with DCM. The combined organic layer was washed with brine, dried over Na_2SO_4 and concentrated to give the title product (120 mg, crude). MS (ES+) $C_{24}H_{27}N_5OS_2$ requires: 465, found 466 $[M+H]^+$.

- 25 **Step 3: Synthesis of cyclopropyl(4-(6-(4-(4,5-dihydro-1H-imidazol-2-ylamino)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazin-1-yl)methanone (Compound 121)**


[0279]

[0280] A mixture of dimethyl 4-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)pyrrolo[1,2-b]pyridazin-6-yl)phenylcarbonimidodithioate (120 mg, crude) and ethane-1,2-diamine (300 mg) in DMF (5 mL) was heated to 120 °C for 10 h. The reaction mixture was concentrated and purification by Pre-HPLC to give the title product as a white solid (48.2 mg, yield 40.2%). MS (ES+) $C_{24}H_{27}N_7O$ requires: 429, found 430 $[M+H]^+$.

Example 20. Synthesis of ethyl 4-(6-(2-fluoro-4-(1-isopropyl-4-methoxypiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 701)

Step 1: Synthesis of 4-(piperazin-1-yl)-6-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)pyrrolo[1,2-b]pyridazine


[0281]



[0282] A mixture of tert-butyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (100 g, 0.234 mol) in DCM (200 mL) was added HCl in dioxane (584 mL, 2.34 mol) at 15 °C for 16 hrs. The reaction mixture was cooled to RT and concentrated to afford the title product (100 g, crude) as a yellow solid. MS (ES+) $C_{17}H_{25}BN_4O_2$ requires: 328, found: 329 $[M+H]^+$.

Step 2: Synthesis of ethyl 4-(6-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0283]




[0284] 4-(piperazin-1-yl)-6-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)pyrrolo[1,2-b]pyridazine, 2TFA (917mg, 1.45mmole) was dissolved in 7mL dichloromethane and Hunig's base (1.0mmole, 5.8mmole) was added followed by

ethyl chloroformate (167 μ L, 1.7 mmole) and the reaction mixture allowed to stir overnight at room temperature. The reaction mixture was evaporated exhaustively and then the residue was preloaded onto silica gel and subjected to flash chromatography using a gradient of 0 to 100% ethyl acetate/hexane. Pure fractions were combined and evaporated to give 403 mg (58%) of the title compound as a pale yellow foam.

Step 3: Synthesis of ethyl 4-(6-(2-fluoro-4-(1-isopropyl-4-methoxypiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 701)

[0285]




[0286] Ethyl 4-(6-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (106 mg, 0.22 mmole), 4-(4-bromo-3-fluorophenyl)-1-isopropyl-4-methoxypiperidine (70 mg, 0.21 mmole), palladium (II) acetate (1.0 mg, 2 mol%), SPhos (3.5 mg, 4 mol%), and potassium carbonate (88 mg, 0.64 mmole) were combined in a vial and purged with nitrogen. Acetonitrile (0.8 mL) and water (0.4 mL) were added and then the reaction was heated to 100 degrees in an oil bath for 3 hours. The reaction mixture was diluted with ethyl acetate and then filtered through celite and transferred to a separatory funnel. The organic layer was washed with brine and dried over sodium sulfate. Filtration and evaporation gave the crude product, which was subjected to flash chromatography using a gradient of 0 to 10% methanol/dichloromethane containing 1% ammonium hydroxide. Pure fractions were combined and evaporated to give 96 mg (87%) of the title compound as a tan-colored foam.

Example 21. Synthesis of 3-hydroxy-2-(4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxyloxy)propanoic acid (Compound 847)

Step 1. Synthesis of benzyl 2,3-dihydroxypropanoate


[0287]



[0288] To a solution of 2,3-dihydroxypropanoic acid (10.0 g, 18.8 mmol, 20% aq) in DMF (20 mL) was added (bromomethyl)benzene (8.0 g, 47.1 mmol) and K₂CO₃ (6.5 g, 47.1 mmol). The resulting mixture was stirred at room temperature for 48 h. LC-MS showed that the reaction was completed. The solvent was evaporated. The residue was dissolved in EtOAc and washed with brine. The organic layer was dried over Na₂SO₄, filtered and evaporated. The crude product was purified by flash chromatography on silica gel eluting with EtOAc/petroleum ether (3: 1) to give the title compound (1.7 g, 46% yield) as a colorless oil.

Step 2. Synthesis of benzyl 3-(tert-butyldiphenylsilyloxy)-2-hydroxypropanoate


[0289]



[0290] To a solution of benzyl 2,3-dihydroxypropanoate (1.7 g, 8.6 mmol) in DMF (10 mL) was added tert-butyldichlorodiphenylsilane (2.6 g, 9.5 mmol) and imidazole (1.2 g, 17.2 mmol). The resulting mixture was stirred at room temperature for 16 hours; LC-MS showed that the reaction was completed. Water was added and extracted with DCM. The organic was washed with brine, dried over Na₂SO₄ and concentrated to afford crude product which was purified by flash chromatography on silica gel eluting with PE/EA (3: 1) to give the title product (510 mg, 14% yield) as a colorless oil.

Step 3. Synthesis of 1-(benzyloxy)-3-(tert-butyldiphenylsilyloxy)-1-oxopropan-2-yl 4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0291]




[0292] To a solution of benzyl 3-(tert-butyldiphenylsilyloxy)-2-hydroxypropanoate (510 mg, 1.2 mmol) in DCM (10 mL) was added CDI (194 mg, 1.2 mmol). The solution was stirred at room temperature for 6 h.

[0293] In another flask, 6-(4-(1-isopropylpiperidin-4-yl)phenyl)-4-(piperazin-1-yl)pyrrolo [1,2-b]pyridazine (322 mg, 0.8 mmol) was dissolved in DCM (5 mL) and TEA (244 mg, 2.4 mmol) was added thereto. The above solution of benzyl 3-(tert-butyl-diphenylsilyloxy)-2-hydroxypropanoate and CDI was added the solution. The mixture was stirred at room temperature for 16 hours; LC-MS showed that the reaction was completed. Water was added and extracted with EtOAc. The organic was washed with brine, dried over Na₂SO₄ and concentrated to afford crude product, which was purified by silica gel chromatography eluting with PE/EA (1: 1) to give the title compound (310 mg, 45% yield) as a light yellow solid.

Step 4. Synthesis of 1-(benzyloxy)-3-hydroxy-1-oxopropan-2-yl 4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate


[0294]



[0295] To a solution of 1-(benzyloxy)-3-(tert-butyl-diphenylsilyloxy)-1-oxopropan-2-yl 4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (250 mg, 0.29 mmol) in THF (5 mL) was added TBAF (2.0 mL, 1.0 M). The solution was heated at 70 °C for 6 hours. The reaction mixture was concentrated and subjected to silica gel eluting with DCM/MeOH (10: 1) to give the title compound (80 mg, 44% yield) as a light yellow solid.

Step 5. Synthesis of 3-hydroxy-2-(4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxyloxy)propanoic acid

[0296]




[0297] To a solution of 1-(benzyloxy)-3-hydroxy-1-oxopropan-2-yl 4-(6-(4-(1-isopropylpiperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

(80 mg, 0.12 mmol) in MeOH (5 mL) was added Pd/C (50 mg, 10% wet). The suspension was hydrogenated with H₂ balloon at room temperature for 16 hours. Filtered off and the filtrate was evaporated to afford crude product which was purified by Prep-HPLC to give the title compound (5.3 mg, 8% yield) as a light white solid.

5 **Example 22. Synthesis of 1,3-dihydroxypropan-2-yl 4-(6-(4-(piperidin-4-yl)phenyl) pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 789)**

10 ***Step 1. Synthesis of 2-phenyl-1,3-dioxan-5-yl 4-(6-(4-(1-(tert-butoxycarbonyl)piperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate***


[0298]



15 **[0299]** To a solution of 2-phenyl-1,3-dioxan-5-ol (327 mg, 1.82 mmol) in DCM (5 mL) was added CDI (295 mg, 1.82 mmol). The resulting mixture was stirred at room temperature for 6 h. In another flask, tert-butyl 4-(4-(4-(piperazin-1-yl)pyrrolo- [1,2-b]pyridazin-6-yl)phenyl)piperidine-1-carboxylate (420 mg, 0.91 mmol) was dissolved in DCM (5 mL) and TEA (278 mg, 2.73 mmol) was added thereto. The above solution of 2-phenyl-1,3-dioxan-5-ol and CDI was added into the solution. The reaction mixture was stirred at room temperature for 16 hours where LC-MS showed that the reaction was completed. Water was added and extracted with EtOAc. The organic layers were washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by PE/EA (2: 1) to give the title compound (340 mg, 56%) as a yellow powder.

25 ***Step 2. Synthesis of 1,3-dihydroxypropan-2-yl 4-(6-(4-(1-(tert-butoxycarbonyl)piperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate***


[0300]



- [0301] To a solution of 2-phenyl-1,3-dioxan-5-yl 4-(6-(4-(1-(tert-butoxycarbonyl)piperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (310 mg, 0.46 mmol) in MeOH (10 mL) was added Pd(OH)₂/C (100 mg). The suspension was hydrogenated with hydrogen balloon at room temperature for 16 hours. LC-MS showed that the reaction was completed. The suspension was then filtered and the solvent was removed under reduced pressure to give the title product (210 mg, 78% yield) as a yellow solid.

Step 3. Synthesis of 1,3-dihydroxypropan-2-yl 4-(6-(4-(piperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0302]




- [0303] To a solution of 1,3-dihydroxypropan-2-yl 4-(6-(4-(1-(tert-butoxycarbonyl)piperidin-4-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (210 mg, 0.36 mmol) in DCM (5 mL) was added TFA (2.5 mL) at 0 °C. The resulting mixture was stirred at room temperature for 3 hours. LC-MS indicated that the reaction was completed. The resulting mixture was evaporated to dryness to afford crude product which was purified by Prep-HPLC to give the title compound as a white powder.

[0304] Example 23. Synthesis of (S)-oxetan-3-yl 4-(6-(4-(1-isopropylpiperidin-3-yl)phenyl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (Compound 631):

- Step 1: Synthesis of 4-(piperazin-1-yl)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazine hydrochloride**


[0305]



- 5 **[0306]** To a solution of tert-butyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (7 g, 16.3 mmol) in dioxane (200 mL) was added HCl/dioxane (4 M, 16 mL). The reaction mixture was stirred at 20 °C overnight. The resulting mixture was concentrated to give the title compound as a yellow solid (9 g, crude). MS (ES+) $C_{17}H_{26}BN_4O_2$ requires: 328, found: 329
- 10 $[M+H]^+$.

Step 2: Synthesis of 4-nitrophenyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate


[0307]



- 15 To a solution of 4-(piperazin-1-yl)-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazine hydrochloride (7 g, 19.1 mmol) in dichloromethane (DCM) (100 mL) was added triethylamine (TEA) (11.6 g, 115 mmol) and 4-nitrophenyl carbonochloridate (4.62 g, 23 mmol). The mixture was stirred at room temperature (RT) for 4 h. The solution was concentrated to give a residue, which was purified by
- 20 flash chromatography to afford the title compound (7 g, 85%). MS (ES+) $C_{24}H_{28}BN_5O_6$ requires: 493, found: 494 $[M+H]^+$.

Step 3: Synthesis of oxetan-3-yl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate

[0308]



[0309] To a solution of oxetan-3-ol (1.56 g, 21.1 mmol) in tetrahydrofuran (THF) (200 mL) at 0 °C was added 60% NaH (2.26 g, 56.4 mmol). The mixture was stirred at 25 °C for 1 h, followed by 4-nitrophenyl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (7 g, 14.1 mmol). The reaction mixture was stirred at 25 °C for another 5 h. The reaction was quenched with NH₄Cl solution and extracted with ethyl acetate.


Combined, dried and concentrated to give the title compound (3.5 g, 71%).

MS

(ES+) C₂₁H₂₉BN₄O₅ requires: 428, found: 429[M+H]⁺.

Step 4: Synthesis of (S)-oxetan-3-yl 4-(6-(4-(1-isopropylpiperidin-3-yl)phenyl)pyrrolo(1,2-b)pyridazin-4-yl)piperazine-1-carboxylate

[0310]




[0311] A solution of oxetan-3-yl 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[1,2-b]pyridazin-4-yl)piperazine-1-carboxylate (3 g, 7.00 mmol) in dioxane/water (20 mL, 10/1) was added (S)-3-(4-bromophenyl)-1-isopropylpiperidine (1.77 g, 6.30 mmol), Pd(dppf)Cl₂ (511 mg, 700 μmol) and K₂CO₃ (2.89 g, 21.0 mmol) was degassed with nitrogen and heated at 90 °C for 5 h. The reaction mixture was cooled to RT and concentrated to give a residue, which was purified by flash chromatography to afford the title compound (1.5 g, 43%)

Example 24. Building block synthesis

A. Synthesis of 1-cyclopentyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole

[0312]




[0313] A mixture of bromocyclopentane (1.0 g, 6.71 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (1.32 g, 6.84 mmol) and Cs₂CO₃ (6.54 g, 20.1 mmol) in CH₃CN (50 mL) was refluxed overnight. The mixture was cooled, concentrated and purified by flash column chromatography (PE:EA = 5:1) to give the title compound as a white solid (1.5 g, yield 85.7 %). MS (ES⁺)

C₁₄H₂₃BN₂O₂ requires: 262, found 263 [M+H]⁺.

B. Synthesis of 1-(2,2-difluoroethyl)-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine

[0314] Step 1: Synthesis of 2,2-difluoroacetaldehyde




5

[0315] To a solution of 2,2-difluoroethanol (2.0 g, 24.3 mmol) in DMF (20 mL) was added Dess-Martin periodinane (25.7 g, 60.7 mmol) in portions at 0 °C. The solution was stirred at RT for 6 h. After that, the solution was used directly for the next reaction.

10 Step 2: Synthesis of 1-(4-bromophenyl)-4-(2,2-difluoroethyl)piperazine


[0316]



[0317] Above solution (24.3 mmol, assumed) in DMF was diluted with DCM (30 mL) and CH₃OH (30 mL). 1-(4-Bromophenyl)piperazine (6.0 g, 24.9 mmol) and acetic acid (1.49 g, 24.9 mmol) was added, followed by addition of sodium cyanoborohydride (2.34 g, 37.3 mmol) at 0 °C. The solution was stirred at RT for 12 h. After that, the solution was cooled to 0 °C, quenched with NaHCO₃ (aq) and brine, and diluted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by flash column (PE/EtOAc = 6/1) to get the title compound as a yellow solid (400 mg, 5%). MS (ES+) C₁₂H₁₅BrF₂N₂ requires: 304, found 305 [M+H]⁺.

Step 3: Synthesis of 1-(2,2-difluoroethyl)-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine

25 [0318]




[0319] A mixture of 1-(4-bromophenyl)-4-(2,2-difluoroethyl)piperazine (400 mg, 1.31 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (497 mg, 1.96 mmol), potassium acetate (385 mg, 3.93 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (191 mg, 262 mmol) in dioxane (20 mL) was purged with N₂, and then stirred at 80 °C for 16 h. After that, the mixture was cooled, concentrated and purified by flash column (PE/EtOAc = 5/1) to get the title compound as a white solid (350 mg, 76%). MS (ES+) C₁₈H₂₇BF₂N₂O₂ requires: 352, found 353 [M+H]⁺.

35

C. Synthesis of 1-(4-bromophenyl)-2-(difluoromethyl)piperazine

Step 1: Synthesis of tert-butyl 4-benzyl-3-(hydroxymethyl)piperazine-1-carboxylate


[0320]



[0321] A mixture of tert-butyl 3-(hydroxymethyl)piperazine-1-carboxylate (5 g, 23.1 mmol), (bromomethyl)benzene (4.73 g, 27.7 mmol) and triethylamine (4.67 g, 46.2 mmol) in acetonitrile (40 mL) was stirred at 80 °C overnight. The reaction solution was cooled, concentrated and purified by silica gel chromatograph (EA:PE = 1:5) to give the title product (6.0 g, yield: 85%) as colorless oil. MS (ES+) $C_{17}H_{26}N_2O_3$ requires: 306, found: 307 $[M+H]^+$.

Step 2: Synthesis of tert-butyl 4-benzyl-3-(hydroxymethyl)piperazine-1-carboxylate


[0322]



[0323] A solution of oxalyl chloride (1.98 g, 15.6 mmol) in DCM (10 mL) was added DMSO (1.52 g, 19.5 mmol) in DCM (10 mL) at -78 °C. The mixture was stirred for 15 min, followed by addition of tert-butyl 4-benzyl-3-(hydroxymethyl)piperazine-1-carboxylate (4.00 g, 13.0 mmol). The resulting mixture was stirred at RT for 2 h, followed by addition of Et₃N. LC-MS showed full conversion. The reaction mixture was diluted with DCM, washed with water and brine, and dried over sodium sulfate. The organic layer was concentrated to afford the title compound (4.00 g, crude) as a yellow oil, which was used in the next step without further purification. MS (ES+) $C_{17}H_{24}N_2O_3$ requires: 304, found: 305 $[M+H]^+$.

Step 3: Synthesis of tert-butyl 4-benzyl-3-(difluoromethyl)piperazine-1-carboxylate


[0324]



[0325] A mixture of tert-butyl 4-benzyl-3-formylpiperazine-1-carboxylate (4.00 g, 13.1 mmol) in DCM (10 mL) was added diethylaminosulfurtrifluoride (3.45 mL, 26.2 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 2 h. LC-MS showed full conversion. The reaction solution was poured into ice water and extracted with DCM. The organic layer was washed with water and brine, dried over sodium sulfate, concentrated and purified by silica gel chromatograph (EA:PE = 1:5) to give the title product (1.40 g, yield 33%) as a light yellow oil. MS (ES+) $C_{17}H_{24}F_2N_2O_2$ requires: 326, found: 327 $[M+H]^+$.

Step 4: Synthesis of tert-butyl 3-(difluoromethyl)piperazine-1-carboxylate


[0326]



[0327] A mixture of tert-butyl 3-(difluoromethyl)piperazine-1-carboxylate (1.20 g, 3.67 mmol) in methanol (10 mL) was added Pd/C (388 mg). The suspension was stirred at RT overnight under H_2 (balloon). LC-MS showed full conversion. The reaction mixture was filtered through a pad of Celite. The filtrate was concentrated to give the title product (800 mg, crude) as a light yellow oil, which was used in the next step without further purification.

Step 5: Synthesis of tert-butyl 4-(4-bromophenyl)-3-(difluoromethyl)piperazine-1-carboxylate

[0328]




[0329] A mixture of tert-butyl 3-(difluoromethyl)piperazine-1-carboxylate (250 mg, 1.05 mmol), 1-bromo-4-iodobenzene (1.48 g, 5.25 mmol), bis(tri-*t*-butylphosphine)palladium (268 mg, 525 μ mol) and sodium tert-butoxide (201 mg, 2.10 mmol) in toluene (20 mL) was purged with N_2 , and then stirred at 60 °C for 15 h. LCMS showed full conversion. The reaction mixture was concentrated. The

residue was purified by silica gel chromatograph (EA:PE = 1: 10 to EA:PE = 1:1) to give the title product (310 mg, yield 75%) as a colorless oil. MS (ES+) $C_{16}H_{21}BrF_2N_2O_2$ requires: 390, found: 391, $[M+H]^+$.

Step 6: Synthesis of 1-(4-bromophenyl)-2-(difluoromethyl)piperazine

5 [0330]




10 [0331] A mixture of tert-butyl 4-(4-bromophenyl)-3-(difluoromethyl)piperazine-1-carboxylate (150 mg, 383 μ mol) in HCl/dioxane (4 M, 2 mL) was stirred at RT for 1.5 h. LCMS showed full conversion. The reaction mixture was concentrated and purified by Prep-HPLC to give the title product (80 mg, yield 72%) as colorless oil. MS (ES+) $C_{11}H_{13}BrF_2N_2$ requires: 290, found: 291 $[M+H]^+$.

D. Synthesis of tert-butyl 3-(4-bromophenyl)-3-fluoropiperidine-1-carboxylate


15 **Step 1: Synthesis of tert-butyl 3-(4-bromophenyl)-3-hydroxypiperidine-1-carboxylate**

[0332]



20 [0333] A solution of 1-bromo-4-iodobenzene (1.0 g, 3.53 mmol) in THF (20 mL) was cooled to -78°C , followed by addition of n-BuLi (2.5 N in hexane, 1.4 mL, 3.53 mmol). After 15 minutes, tert-butyl 3-oxopiperidine-1-carboxylate (703 mg, 3.53 mmol) in THF (5 mL) was added slowly. The reaction solution was stirred for 2 h at -78°C , allowed to warm up to 0°C , and quenched using saturated aqueous NH_4Cl and extracted with EA. The organic layer was concentrated and purified by flash column
25 (silica gel, PE:EA = 3:1) to give the title product (0.8 g, yield 64%) as a colorless oil. MS (ES+) $C_{16}H_{22}BrNO_3$ requires: 355, found 356 $[M+H]^+$.


Step 2: Synthesis of tert-butyl 3-(4-bromophenyl)-3-fluoropiperidine-1-carboxylate

[0334]

[0335] To a solution of tert-butyl 3-(4-bromophenyl)-3-hydroxypiperidine-1-carboxylate (3.0 g, 8.42 mmol) in DCM (30 mL) was added Dast (2.0 g, 12.6 mmol) at 0 °C. The reaction mixture was stirred at RT for 5 h. The mixture was diluted with DCM, washed with aq. NaHCO₃ and brine, evaporated and purified by flash column (silica gel, PE:EA = 5:1) to give the title product (1.5 g, yield 50%) as a colorless oil. MS (ES+) C₁₆H₂₁BrFNO₂ requires: 357, found 358 [M+H]⁺.

E. Synthesis of tert-butyl 3-(4-bromophenyl)-4-fluoropyrrolidine-1-carboxylate

Step 1: Synthesis of tert-butyl 3-(4-bromophenyl)-2,5-dihydro-1H-pyrrole-1-carboxylate

[0336]

[0337] A mixture of tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxylate (1.0 g, 3.38 mmol), 1-bromo-4-iodobenzene (1.91 g, 6.76 mmol), Pd(dppf)Cl₂ (247 mg, 338 μmol) and K₂CO₃ (932 mg, 6.76 mmol) in dioxane/water (5 mL, 4/1) was purged with N₂ and stirred at 80 °C for 16 h under N₂. The mixture was cooled and concentrated in *vacuo*. The residue was purified by flash column chromatography (PE/EA = 10:1) to afford the title compound (800 mg, yield 73%) as a white solid MS (ES+) C₁₅H₁₈BrNO₂ requires: 323, found 324 [M+H]⁺.

Step 2: Synthesis of tert-butyl 3-(4-bromophenyl)-4-hydroxypyrrolidine-1-carboxylate


[0338]

[0339] A solution of BH₃ (1 N in THF, 21.5 mL, 21.5 mmol) was added to a stirring solution of tert-butyl 3-(4-bromophenyl)-2,5-dihydro-1H-pyrrole-1-carboxylate (1.4 g, 4.31 mmol) in THF (20 mL) at 0 °C. After stirred at RT for 4 h, the mixture was cooled to 0 °C, followed by addition of NaOH aqueous (4 N, 6.45 mL, 25.8 mmol). After 10 min, H₂O₂ (2.92 g, 25.8 mmol) was added. The resulting mixture was allowed to warm up to RT and stirred for 90 min. Monitored by LC-MS. Quenched by

water and extracted with ethyl acetate. The organic layer was concentrated and purified by flash column chromatography (PE/EA = 10:1 to 1:1) to afford the title compound (1.3 g, yield 88%) as a light oil MS (ES+) $C_{15}H_{20}BrNO_3$ requires: 341, found 342 $[M+H]^+$.

5 **Step 3: Synthesis of tert-butyl 3-(4-bromophenyl)-4-fluoropyrrolidine-1-carboxylate**

[0340]




- 10 **[0341]** To a mixture of tert-butyl 3-(4-bromophenyl)-4-hydroxypyrrolidine-1-carboxylate (500 mg, 1.46 mmol) in DCM (10 mL) was added dropwise DAST (1.17 g, 7.29 mmol) at -78°C . The mixture was stirred at 25°C for 2 h. Monitored by LC-MS. Diluted with DCM and quenched with saturated NaHCO_3 solution. The organic layer was separated and concentrated in *vacuo* to afford the title compound (400
- 15 mg, crude) as a yellow oil. MS (ES+) $C_{15}H_{19}BrFNO_2$ requires: 343, found: 344 $[M+H]^+$.

F. Synthesis of 4-(4-bromophenyl)-4-methoxypiperidine


Step 1: Synthesis of tert-butyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate

20 **[0342]**



- [0343]** To a solution of 1-bromo-4-iodobenzene (12 g, 42.6 mmol) in THF (60 mL) at -78°C was added BuLi (20 mL, 2.4 M in hexane) dropwise. The solution was stirred at -78°C for 2 h. A solution of tert-butyl 4-oxopiperidine-1-carboxylate (8.5 g, 42.4 mmol) in THF (20 mL) was added dropwise. The resulting solution was stirred at -78°C for 1 h. The reaction was quenched carefully by addition of water and extracted with EtOAc. The organic layer was washed with water and brine, concentrated and purified by flash column (PE/EtOAc = 10/1 to 3/1) to give the title product (12.6 g,
- 25 yield 83.4%). MS (ES+) $C_{16}H_{22}BrNO_3$ requires: 355, found 282 $[M-73]^+$.
- 30


Step 2: Synthesis of tert-butyl 4-(4-bromophenyl)-4-methoxypiperidine-1-carboxylate

[0344]

- 5 **[0345]** To a solution of tert-butyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate (178 mg, 499 μ mol) in dry DMF (2 mL) was added NaH (26 mg, 0.646 mmol) at 0 °C. The mixture was stirred at 0 °C for 1 h. Iodomethane (106 mg, 0.746 mmol) was added. The mixture was stirred at RT overnight. The reaction was quenched carefully by addition of water, extracted with EtOAc, washed with water and brine, concentrated and purified by flash column (PE/EtOAc = 10/1 to 5/1) to give the title product (160 mg, yield 86.5%). MS (ES+) $C_{17}H_{24}BrNO_3$ requires: 369, found 282 $[M+H-88]^+$.
- 10


Step 3: Synthesis of 4-(4-bromophenyl)-4-methoxypiperidine**[0346]**

- 15 **[0347]** To a solution of tert-butyl 4-(4-bromophenyl)-4-methoxypiperidine-1-carboxylate (740 mg, 1.99 mmol) in dioxane (4 mL) was added HCl (4 M in dioxane, 3 mL). The resulting solution was stirred at RT overnight. The solvent was removed in *vacuo* to afford the title product (690 mg, crude). MS (ES+) $C_{12}H_{16}BrNO$ requires: 269, found 270 $[M+H]^+$.
- 20

Step 4: Synthesis of 4-(4-bromophenyl)-4-methoxypiperidine**[0348]**


- 25 **[0349]** To a solution of 4-(4-bromophenyl)-4-methoxypiperidine (1 g, 3.70 mmol) in CH_3CN (50 mL) was added K_2CO_3 (1.53 g, 11.1 mmol) and 2-bromopropane (2.27 g, 18.5 mmol). The resulting mixture was stirred at 80 °C for 5 h. The solvent was removed in *vacuo*. The residue was purified by flash column (PE/EtOAc = 10/1 to 5/1) to afford the title product (1.06 g, yield 91.8%). MS (ES+) $C_{15}H_{22}BrNO$ requires: 311, found 312 $[M+H]^+$.
- 30

G. Synthesis of 1-isopropyl-4-methoxy-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine**Step 1: Synthesis of tert-butyl 4-methoxy-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine-1-carboxylate**

[0350]


[0351] A mixture of tert-butyl 4-(4-bromophenyl)-4-methoxypiperidine-1-carboxylate (200 mg, 540 μ mol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (178 mg, 702 μ mol), Pd(dppf)Cl₂ (39.5 mg, 54.0 μ mol) and K₂CO₃ (105 mg, 1.08 mmol) in dioxane (10 mL) was stirred at 65 °C for 4 h. The reaction mixture was concentrated and purified by silica gel chromatography (PE:EA = 10:1 to 5:1) to give the title product (158 mg, yield: 70%) as a yellow solid. MS (ES+) C₂₃H₃₆BNO₅ requires: 417, found 418 [M+H]⁺.

Step 2: Synthesis of 4-methoxy-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine

[0352]


[0353] A mixture of tert-butyl 4-methoxy-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine-1-carboxylate (6 g, 14.3 mmol) and HCl/dioxane (30 mL) in DCM (30 mL) was stirred at RT for 1 h. The reaction mixture was concentrated to give the title product (4.6 g, crude) as a light yellow solid, which was used for the next step without purification. MS (ES+) C₁₈H₂₈BNO₃ requires: 317, found 318 [M+H]⁺.

Step 3: Synthesis of 1-isopropyl-4-methoxy-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine


[0354]

[0355] A mixture of 4-methoxy-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperidine (4.5 g, 14.1 mmol), 2-iodopropane (3.58 g, 21.1 mmol) and K₂CO₃ (5.84 g, 42.3 mmol) in acetonitrile (50 mL) was stirred at 85 °C for 2 h. The reaction mixture was concentrated and purified by silica gel chromatography (DCM:MeOH = 15:1) to give the title product (4.8 g, yield: 95%) as a light yellow solid. MS (ES+) C₂₁H₃₄BNO₃ requires: 359, found 360 [M+H]⁺.

H. Synthesis of tert-butyl 4-(3-(difluoromethoxy)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine-1-carboxylate


Step 1: Synthesis of 1,4-dibromo-2-(difluoromethoxy)benzene**[0356]**

[0357] To a solution of 2,5-dibromophenol (1.5 g, 6.0 mmol) in DMF/water (30 mL/10 mL) was added sodium 2-chloro-2,2-difluoroacetate (2.3 g, 15.0 mmol) and Cs₂CO₃ (3.9 g, 12.0 mmol) at RT. The resulting mixture was stirred at 100 °C for 3 h. Quenched with water and extracted with EtOAc. The combined organic layer was washed with brine, dried over Na₂SO₄, and concentrated to dryness. The residue was purified by flash chromatography on silica gel eluting with PE/EtOAc (5:1) to give the title compound (1.2 g, yield 67%) as a light yellow powder. MS (ES+) C₇H₄Br₂F₂O requires: 300, found 301 [M+H]⁺ (weak ion mass).

Step 2: Synthesis of tert-butyl 4-(4-bromo-3-(difluoromethoxy)phenyl)piperazine-1-carboxylate**[0358]**


[0359] To a solution of 1,4-dibromo-2-(difluoromethoxy)benzene (500 mg, 1.66 mmol) and tert-butyl piperazine-1-carboxylate (309 mg, 1.66 mmol) in DMF (20 mL) was added Pd₂(dba)₃ (155 mg, 0.17 mmol), XantPhos (98 mg, 0.17 mmol) and Cs₂CO₃ (1.6 g, 5.0 mmol) at RT under nitrogen. The resulting mixture was stirred at 90 °C for 1 h under Microwave. LC-MS showed that the reaction was completed. Quenched with water and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated to dryness. The residue was purified by flash chromatography on silica gel eluting with PE/EtOAc (3:1 to 1: 1) to give the title compound (370 mg, yield 55%) as white solid. MS (ES+) C₁₆H₂₁BrF₂N₂O₃ requires: 406, found 351 [M+H-56]⁺.

Step 3: Synthesis of tert-butyl 4-(3-(difluoromethoxy)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)piperazine-1-carboxylate

[0360]

[0361] To a solution of tert-butyl 4-(4-bromo-3-(difluoromethoxy)phenyl)piperazine-1-carboxylate (320 mg, 0.78 mmol) and 4,4',4'',5,5',5'',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (309 mg, 1.66 mmol) in dioxane (10 mL) was added Pd(dppf)Cl₂ (63 mg, 0.078 mmol) and KOAc (229 mg, 2.34 mmol) at RT under nitrogen. The resulting mixture was stirred at 90 °C for 6 h under nitrogen; LC-MS showed that the reaction was completed. Quenched with water and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated to dryness to afford crude product, which was purified by flash chromatography on silica gel eluting with PE/ EtOAc (10:1 to 3: 1) to give the title compound (260 mg, yield 73%) as white solid. MS (ES+) C₂₂H₃₃BF₂N₂O₅ requires: 454, found 399 [M+H-56]⁺.

I. Synthesis of 2-(4-chlorophenyl)-1-isopropyl-2-methylpyrrolidine

[0362]

Step 1: Synthesis of tert-butyl 2-(4-chlorophenyl)-2-methylpyrrolidine-1-carboxylate

[0363] 5-(4-chlorophenyl)-3,4-dihydro-2H-pyrrole (2.00g, 11.1mmole) was dissolved in 33mL dry THF and cooled to -78°C. Boron trifluoride diethyletherate (2.8mL, 22.3mmole) was added dropwise and stirred at -78°C for 40 minutes and then 0.5M methyllithium in ether (13.9mL, 22.3mmole) was added dropwise and then allowed to warm slowly to room temperature overnight. The reaction mixture was quenched with water and diluted with ethyl acetate. The reaction mixture was acidified with 1M HCl and then transferred to a separatory funnel. The aqueous layer was then made basic to pH-12-13 with 6M NaOH and then extracted with ethyl acetate (x2) and combined organics washed with brine and dried over sodium sulfate. Filter and evaporate to give the crude product, which consisted of desired product and the starting material. The two were difficult to separate by chromatography so were converted to the Boc protected product to facilitate separation: The crude product was dissolved in 30mL dichloromethane and then Boc anhydride (1.74g, 8.0mmole) was added as a solution in 5mL dichloromethane, followed by DMAP (100mg, 0.8mmole) and the mixture stirred at room temperature overnight. The reaction


mixture was evaporated and then subjected to flash chromatography (0 to 40% ethyl acetate/hexane, collecting all fractions due to low UV activity). Pure fractions combined and evaporated to give 920mg (28%) of the title compound as a colorless oil that crystallized on standing.

5 **Step 2: Synthesis of 2-(4-chlorophenyl)-1-isopropyl-2-methylpyrrolidine**

[0364] tert-Butyl 2-(4-chlorophenyl)-2-methylpyrrolidine-1-carboxylate (915mg, 3.1mmole) was dissolved in 12mL dichloromethane and trifluoroacetic acid (3.6mmole, 46.4mmole) was added and stirred at room temperature for 2 hours. The reaction mixture was evaporated exhaustively and then the residue partitioned between
 10 dichloromethane and 1M NaOH. The organic layer was dried over sodium sulfate, filtered and evaporated to give 610mg (100%) of 2-(4-chlorophenyl)-2-methylpyrrolidine as a viscous orange oil. This material was then dissolved in 10mL acetonitrile in a heavy-walled pressure vessel and potassium carbonate (646mg, 4.7mmole) was added followed by 2-iodopropane (374uL, 3.7mmole). The reaction
 15 mixture was heated to 90°C for 3 days and then diluted with ethyl acetate. The reaction was filtered through celite and evaporated. The crude product was subjected to flash chromatography using an Isco amine column, gradient of 0 to 30% ethyl acetate/hexane. Pure fractions were combined and evaporated to give 509mg (69%) of the desired product as a pale yellow oil.

20 **J. Synthesis of benzyl 4-(4-bromophenyl)-4-fluoropiperidine-1-carboxylate**

[0365]



Step 1: Synthesis of benzyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate.

[0366] Benzyl 4-oxopiperidine-1-carboxylate (937mg, 4.0mmole) and 1,4-dibromobenzene (790mg, 3.4mmole) were dissolved in 15mL THF and cooled to -78°C. nBuLi (1.47mL of 2.5M solution in hexanes, 3.7mmole) was added dropwise and the reaction mixture was allowed to warm to room temperature over several hours. The reaction was quenched with saturated ammonium chloride solution and
 30 diluted with ethyl acetate and transferred to a separatory funnel. The organic layer was washed with brine and dried over sodium sulfate. Filtration and evaporation gave the crude product, which was subjected to flash chromatography using a gradient of 0 to 35% ethyl acetate/hexane, collecting all fractions and monitoring by ELSD. Clean fractions combined and evaporated to give 732mg (56%) of the desired
 35 product as a colorless oil.


Step 2: Synthesis of benzyl 4-(4-bromophenyl)-4-fluoropiperidine-1-carboxylate

[0367] Benzyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate (731 mg, 1.87 mmole) was dissolved in 8 mL dichloromethane and cooled to -78 °C. DAST (272 μ L, 2.06 mmole) was added dropwise and then allowed to warm slowly to room temperature over a couple hours. The reaction was quenched with saturated sodium bicarbonate solution and diluted with dichloromethane and transferred to a separatory funnel. The organic layer was dried over sodium sulfate, filtered and evaporated to give the crude product. Purification by flash chromatography using a gradient of 0 to 35% ethyl acetate/hexane, collecting all fractions and monitoring by ELSD. Clean fractions combined and evaporated to give 236 mg (32%) of the desired product as a colorless oil.

K. Synthesis of tert-butyl 3-(6-chloropyridin-3-yl)-3-methylpiperazine-1-carboxylate

Step 1: Synthesis of ethyl 2-(6-chloropyridin-3-yl)propanoate


[0368]



[0369] A solution of lithium bis(trimethylsilyl)amide in THF (1 M, 14 mL, 14.0 mmol) was added to a solution of ethyl 2-(6-chloropyridin-3-yl)acetate (2.5 g, 12.5 mmol) in THF at -78 °C under nitrogen. After stirred at -78 °C for 2 h, methyl iodide (1.94 g, 13.7 mmol) was added. The mixture was stirred at 20 °C for another 8 h. Worked up, concentrated and purified with silica gel column chromatography, eluting with 0%~10% EA/PE, to give the title compound (1.5 g, 83% purity in LCMS, yield 46%) as yellow oil. MS (ES+) $C_{10}H_{12}ClNO_2$ requires: 213, 215, found: 214, 216 $[M+H]^+$.


Step 2: Synthesis of ethyl 2-bromo-2-(6-chloropyridin-3-yl)propanoate

[0370]




[0371] A solution of ethyl 2-(6-chloropyridin-3-yl)propanoate (1.5 g, 5.80 mmol), N-bromosuccinimide (1.23 g, 6.95 mmol) and (E)-azobis(isobutyronitrile) (95 mg, 0.58 mmol) in perchloromethane (50 mL) was stirred at 80 °C for 48 h under nitrogen. The mixture was cooled and concentrated in *vacuo*. The residue was purified with silica gel column chromatography, eluting with 0%-10% EA/PE, to give the title compound (1.9 g, 75% purity in LCMS, yield 84%) as a yellow oil. MS (ES+) $C_{10}H_{11}BrClNO_2$ requires: 291, 293, found: 292, 294 $[M+H]^+$.

Step 3: Synthesis of 3-(6-chloropyridin-3-yl)-3-methylpiperazin-2-one

[0372]

[0373] A solution of ethyl 2-bromo-2-(6-chloropyridin-3-yl)propanoate (1.2 g, 4.10 mmol) in ethane-1,2-diamine (5 mL) was stirred at 25 °C for 18 h. Diluted with DCM and washed with brine. Concentrated, the residue was purified with silica gel column chromatography, eluting with 100% EA, to give the title compound (600 mg, yield 63%) as a yellow solid. MS (ES+) $C_{10}H_{12}ClN_3O$ requires: 225, 227, found: 226, 228 $[M+H]^+$.


Step 4: Synthesis of tert-butyl 3-(6-chloropyridin-3-yl)-3-methylpiperazine-1-carboxylate

[0374]

[0375] A solution of borane in tetrahydrofuran (22 mL, 1 M, 22.0 mmol) was added dropwise to a solution of 3-(6-chloropyridin-3-yl)-3-methylpiperazin-2-one (500 mg, 2.21 mmol) in tetrahydrofuran (10 mL), and then stirred at 80 °C for 18 h under nitrogen. Cooled, quenched with MeOH and refluxed with HCl/dioxane solution. The mixture was basified with 1 M NaOH solution to pH = 10~12, and then di-tert-butyl dicarbonate (1.23 g, 5.64 mmol) was added. Stirred at 20 °C for 18 h. Diluted with EA and washed with water. Concentrated, the residue was purified with silica gel column chromatography, eluting with 0%~10% MeOH/EA, to give the title compound (200 mg, yield 34%) as a white solid. MS (ES+) $C_{15}H_{22}ClN_3O_2$ requires: 311, 313, found: 312, 314 $[M+H]^+$.

L. Synthesis of 2-chloro-3-fluoro-5-(1-isopropyl-4-methoxypiperidin-4-yl)pyridine


Step 1: Synthesis of tert-butyl 4-(6-chloro-5-fluoropyridin-3-yl)-4-hydroxypiperidine-1-carboxylate

[0376]

[0377] To a solution of 5-bromo-2-chloro-3-fluoropyridine (2 g, 9.50 mmol) in THF (60 mL) at -78 °C was added nBuLi (4 mL, 2.4 M in hexane) dropwise. The solution was stirred at -78 °C for 2 h. Then a solution of tert-butyl 4-oxopiperidine-1-carboxylate (1.89 g, 9.50 mmol) in THF (10 mL) was added dropwise. The resulting solution was stirred at -78 °C for 2 h. The reaction was quenched carefully by addition of water and extracted with EtOAc. The organic layer was washed with water and brine, concentrated and purified by silica gel column chromatography (petroleum ether/ethyl acetate = 10/1 to 2/1) to afford the title product (2 g, 64%) as a white solid. MS (ES+) C₁₅H₂₀ClFN₂O₃ requires: 330, found: 331 [M+H]⁺.

Step 2: Synthesis of tert-butyl 4-(6-chloro-5-fluoropyridin-3-yl)-4-methoxypiperidine-1-carboxylate


[0378]



[0379] To a solution of tert-butyl 4-(6-chloro-5-fluoropyridin-3-yl)-4-hydroxypiperidine-1-carboxylate (2 g, 6.04 mmol) in dry DMF (20 mL) was added NaH (60%) (313 mg, 7.85 mmol) at 0 °C. The mixture was stirred at 0 °C for 1 h. Then iodomethane (1.28 g, 9.06 mmol) was added. The mixture was stirred at RT overnight. After that, the mixture was slowly poured into ice water (200 mL) and stirred for 1 h. The solid was collected by filtration and dried to afford the title compound (1.5 g, 72%) as a white solid. MS (ES+) C₁₆H₂₂ClFN₂O₃ requires: 344, found: 345 [M+H]⁺.

Step 3: Synthesis of 2-chloro-3-fluoro-5-(4-methoxypiperidin-4-yl)pyridine

[0380]



[0381] To a solution of tert-butyl 4-(6-chloro-5-fluoropyridin-3-yl)-4-methoxypiperidine-1-carboxylate (1.5 g, 4.35 mmol) in dioxane (10 mL) was added HCl/dioxane (4 N, 10 mL). The mixture was stirred at RT for 12 h. After that, the solution was concentrated. The residue was used in the next step without further purification. MS (ES+) C₁₁H₁₄ClFN₂O requires: 244, found: 245 [M+H]⁺.

Step 4: Synthesis of 2-chloro-3-fluoro-5-(1-isopropyl-4-methoxypiperidin-4-yl)pyridine

[0382]




[0383] A mixture of 2-chloro-3-fluoro-5-(4-methoxypiperidin-4-yl)pyridine (1 g, 4.08 mmol), 2-iodopropane (693 mg, 4.08 mmol) and triethylamine (1.23 g, 12.2 mmol) in CH₃CN (15 mL) was stirred at 80 °C overnight. After that, the solution was concentrated and purified by silica gel column chromatography (DCM/ MeOH=10/1) to afford the title compound (800 mg, 68%) as a yellow solid. MS (ES+) C₁₄H₂₀ClFN₂O requires: 286, found: 287 [M+H]⁺.

10 M. Synthesis of (R)-3-(4-bromophenyl)-1-isopropylpiperidine and (S)-3-(4-bromophenyl)-1-isopropylpiperidine

Step 1: Chiral separation of 3-(4-bromophenyl)piperidine hydrochloride


[0384]



[0385] Racemic 3-(4-bromophenyl)piperidine hydrochloride was separated into single enantiomers using the following appropriately scaled chiral HPLC conditions: Column: CD-PH 250×4.6mm I.D., Sum Mobile phase: A: water with 0.1% TFA B: acetonitrile with 0.1% TFA A / B = 70/30 Flow rate: 1.0mL/min Wavelength: 220nm.

20 Step 2: Synthesis of (S)-3-(4-bromophenyl)-1-isopropylpiperidine

[0386]




[0387] A mixture of (S)-3-(4-bromophenyl)piperidine hydrochloride (3.1 g, 11.2 mmol), 2-bromopropane (2.75 g, 22.4 mmol) and K₂CO₃ (4.62 g, 33.5 mmol) in CH₃CN (20 mL) was stirred at 70 °C for 16 hrs. The mixture was diluted with EtOAc and washed with brine. The organic layer was concentrated in *vacuo* to afford the title compound (2.9 g, yield 91%) as a yellow oil. MS (ES+) C₁₄H₂₀BrN requires: 281, found: 282 [M+H]⁺.

30 N. Synthesis of 5-(4-bromo-2-fluorophenyl)-1-isopropyl-1,2,3,6-tetrahydropyridine

Step 1: Synthesis of tert-butyl 3-(4-bromo-2-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate


5 [0388]



10 [0389] A mixture of tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (4.3 g, 13.9 mmol), 4-bromo-2-fluoro-1-iodobenzene (6.25 g, 20.8 mmol), [1,1'-
Bis(diphenylphosphino)ferrocene]dichloropalladium(II) (2.54 g, 2.78 mmol) and potassium carbonate (5.75 g, 41.7 mmol) in dioxane/H₂O (L/10 mL) was degassed with Nitrogen for three times and then heated at 70 °C for 3 h. The reaction mixture was cooled to room temperature and concentrated to get crude product, which was
15 purified by silica gel chromatography (petroleum ether/ethyl acetate = 4/1) to afford the title compound (3.7 g, yield 75%) as a brown oil. MS (ES+) C₁₆H₁₉BrFNO₂ requires: 355, found: 300 [M-56+H]⁺.

Step 2: Synthesis of 5-(4-bromo-2-fluorophenyl)-1,2,3,6-tetrahydropyridine HCl salt

20 [0390]




25 [0391] To a mixture of tert-butyl 3-(4-bromo-2-fluorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (3.7 g, 10.3 mmol) in dioxane (20 ml) was added dioxane/HCl (4 M, 20 mL). The reaction mixture was stirred at room temperature for 3 hours. Concentrated under reduced pressure to afford the title compound (2.6 g, yield 86%) as a brown solid, which was directly used into the next step without further purification. MS (ES+) C₁₁H₁₁BrFN requires: 255, found: 256 [M+H]⁺.

Step 3: Synthesis of 5-(4-bromo-2-fluorophenyl)-1-isopropyl-1,2,3,6-tetrahydropyridine

30

[0392]




[0393] To a solution of 5-(4-bromo-2-fluorophenyl)-1,2,3,6-tetrahydropyridine HCl salt (150 mg, 585 μ mol) and potassium carbonate (241 mg, 1.75 mmol) in acetonitrile (10 ml) was added 2-iodopropane (496 mg, 2.92 mmol). The solution was stirred at 60 °C for 6 hours. Cooled to room temperature and extracted with EtOAc after water was added. The organic phase was washed with water and brine, dried over Na₂SO₄, and concentrated to dryness to afford the title compound as a yellow solid which was used without further purification (140 mg, crude). MS (ES+) C₁₄H₁₇BrFN requires: 297, found: 298 [M+H]⁺.

10 O. Synthesis of 3-(4-bromophenyl)-1-isopropyl-3-methylpiperazine

Step 1: Synthesis of methyl 2-(4-bromophenyl)acetate

[0394]



[0395] A mixture of 2-(4-bromophenyl)acetic acid (10 g, 46.5 mmol) in MeOH (100 mL, with 2 mL of conc. H₂SO₄) was refluxed for 4 h. The mixture was concentrated and diluted with EtOAc. The organic was washed with water and brine, dried and concentrated to give the title product (10.6 g, yield 100%). MS (ES+) C₉H₉BrO₂ requires: 229 found 230 [M+H]⁺.

20 Step 2: Synthesis of methyl 2-(4-bromophenyl)propanoate

[0396]



[0397] To a solution of methyl 2-(4-bromophenyl)acetate (1 g, 4.36 mmol) in dry THF (20 mL) was added LiHMDS (5.23 mL, 1 M) dropwise at -78 °C under N₂. The mixture was stirred at 0 °C for 2 h. After MeI (1.23 g, 8.72 mmol) was added at -78 °C, the mixture was stirred at 20 °C for 16 h. Quenched with NH₄Cl sat. and extracted with EtOAc. The organic was concentrated and purified by silica gel column (PE/EtOAc = 1/0 to 100/1) to give the title product (642 mg, yield 61%). MS (ES+) C₁₀H₁₁BrO₂ requires: 243, found 244 [M+H]⁺.

Step 3: Synthesis of methyl 2-bromo-2-(4-bromophenyl)propanoate


[0398]



[0399] A mixture of methyl 2-(4-bromophenyl)propanoate (30 g, 123 mmol), AIBN (2.01 g, 12.3 mmol) and NBS (32.9 g, 184 mmol) in CCl_4 (300 mL) was refluxed for 16 h. Cooled to RT and filtered. After concentrated, the residue was used into the next step without further purification.

Step 4: Synthesis of 3-(4-bromophenyl)-3-methylpiperazin-2-one


[0400]



[0401] To a solution of methyl 2-bromo-2-(4-bromophenyl)propanoate (39 g, 121 mmol) in EtOH (200 mL) was added ethane-1,2-diamine (14.5 g, 242 mmol). The mixture was stirred at RT for 16 h. Filtered off solid, concentrated and purified by silica gel column (EtOAc) to give the title product (12 g, yield 37%). MS (ES+) $\text{C}_{11}\text{H}_{13}\text{BrN}_2\text{O}$ requires: 269, found 270 $[\text{M}+\text{H}]^+$.


Step 5: Synthesis of tert-butyl 3-(4-bromophenyl)-3-methylpiperazine-1-carboxylate

[0402]




[0403] A solution of 3-(4-bromophenyl)-3-methylpiperazin-2-one (12 g, 44.5 mmol) and BH_3 (250 mL, 1 M in THF) was stirred at 80 °C overnight. Cooled to RT and quenched with MeOH. After concentrated, the residue was dissolved in MeOH (100 mL), followed by addition of HCl (250 mL, aq, 1 M). The mixture was stirred at 80 °C for 30 min. Cooled to RT. NaOH (12 g, 0.3 mol) was added, followed by addition of Boc_2O (11.6 g, 53.3 mmol). The mixture was stirred at RT overnight. Extracted with EtOAc. The organic was concentrated and purified by silica gel column DCM/MeOH (50/1) to give the title product (10 g, yield 63%). MS (ES+) $\text{C}_{16}\text{H}_{23}\text{BrN}_2\text{O}_2$ requires: 355, found 356 $[\text{M}+\text{H}]^+$.


[0404] The racemic product (10 g) was separated by chiral-HPLC to give two single enantiomers: P1 (4.63 g) and P2 (4.46 g). Chiral conditions: Co-Solvent: MeOH(0.2%Methanol Ammonia); Column: OZ-H 100*4.6mm 5um; Column Temperature: 39.9.

Step 6: Synthesis of 2-(4-bromophenyl)-2-methylpiperazine**[0405]**

- 5 **[0406]** To a mixture of tert-butyl 3-(4-bromophenyl)-3-methylpiperazine-1-carboxylate (1.8 g, 5.06 mmol) in DCM/MeOH (10 mL/10 mL) was added HCl/dioxane (14 mL, 4 M). The mixture was stirred at RT for 1 h, concentrated, and the residue was dissolved in NH₃. MeOH (8 mL, 7 mL). DCM (20 mL) was added. Filtered off solid and then concentrated to give the title product (1.2 g, yield 98%).
- 10 MS (ES⁺) C₁₆H₂₃BrN₂O₂ requires: 255, found 256 [M+H]⁺.

Step 7: Synthesis of 3-(4-bromophenyl)-1-isopropyl-3-methylpiperazine**[0407]**

- 15 **[0408]** A mixture of 2-(4-bromophenyl)-2-methylpiperazine (1.3 g, 5.09 mmol), 2-iodopropane (994 mg, 5.85 mmol) and DIPEA (3.28 g, 25.4 mmol) in THF (50 mL) was stirred at 60 °C for 16 h. Concentrated and purified by silica gel column (EtOAc) to give the title product (1.3 g, yield 86%). MS (ES⁺) C₁₄H₂₁BrN₂ requires: 297, found 298 [M+H]⁺.
- 20 **P. Synthesis of 6'-bromo-1-isopropyl-1,2,5,6-tetrahydro-3,3'-bipyridine**


Step 1: Synthesis of tert-butyl 6'-bromo-5,6-dihydro-[3,3'-bipyridine]-1(2H)-carboxylate**[0409]**

- 25 **[0410]** A mixture of 2-bromo-5-iodopyridine (3.66 g, 12.9 mmol), tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (4 g, 12.9 mmol), Pd(dppf)Cl₂ (944 mg, 1.29 mmol) and K₂CO₃ (2.66 g, 19.3 mmol) in dioxane/water (10/1, 20 mL) was irradiated under microwave at 120 °C for 2 h.
- 30 Concentrated and purified by silica gel column PE/EtOAc (10/1) to give title product (2.12 g, yield 48%). MS (ES⁺) C₁₅H₁₉BrN₂O₂ requires: 339 found 340 [M+H]⁺.

Step 2: Synthesis of 6'-bromo-1,2,5,6-tetrahydro-3,3'-bipyridine

[0411]


- 5 **[0412]** To a solution of tert-butyl 6'-bromo-5,6-dihydro-[3,3'-bipyridine]-1(2H)-carboxylate (2.12 g, 6.24 mmol) in DCM (20 mL) was added TFA (4 mL) at RT, and then stirred for 1 h. The mixture was diluted with water, adjusted pH to 7-8 with NaHCO₃ sat., and extracted with DCM. Dried and concentrated to give the title product (1.49 g, yield 100%). MS (ES+) C₁₀H₁₁BrN₂ requires: 239, found 240 [M+H]⁺.

Step 3: Synthesis of 6'-bromo-1-isopropyl-1,2,5,6-tetrahydro-3,3'-bipyridine10 **[0413]**

- 15 **[0414]** A mixture of 6'-bromo-1,2,5,6-tetrahydro-3,3'-bipyridine (1.49 g, 6.23 mmol), DIPEA (2.39 g, 18.6 mmol) and 2-iodopropane (3.16 g, 18.6 mmol) in MeCN (50 mL) was stirred at 60 °C for 16 h. Concentrated and purified by silica gel column DCM/MeOH (20/1) to give title product (1.75 g, yield 100%). MS (ES+) C₁₃H₁₇BrN₂ requires: 281 found 282 [M+H]⁺.


Q. Synthesis of 2-bromo-5-(4-ethoxy-1-isopropylpiperidin-4-yl)pyridine

- 20 **Step 1: Synthesis of tert-butyl 4-(6-bromopyridin-3-yl)-4-hydroxypiperidine-1-carboxylate**


[0415]

- 25 **[0416]** To a solution of 2-bromo-5-iodopyridine (18.5 g, 65.2 mmol) in 150 mL of THF was added n-BuLi (26 mL, 2.5 M in hexane) dropwise under N₂ at -78 °C. The resulting solution was stirred at -78 °C for 2 h. A solution of tert-butyl 4-oxopiperidine-1-carboxylate (10 g, 50.2 mmol) in 30 mL of THF was added dropwise. The mixture was stirred at -78 °C for another 1 h. The reaction was quenched by addition of sat. NH₄Cl solution and extracted with EtOAc. The organic phase was
30 dried over Na₂SO₄ and concentrated. The residue was purified by silica gel chromatography to afford the title product (12.6 g, 70% yield) as a white solid.


Step 2: Synthesis of tert-butyl 4-(6-bromopyridin-3-yl)-4-ethoxypiperidine-1-carboxylate

[0417]

[0418] To a solution of tert-butyl 4-(6-bromopyridin-3-yl)-4-hydroxypiperidine-1-carboxylate (2.5 g, 6.99 mmol) in 100 mL of THF was added NaH (60%, 415 mg, 10.4 mmol) at 0 °C. The mixture was stirred at 0 °C for 30 min. Then bromoethane (1.13 g, 10.4 mmol) was added. The resulting mixture was stirred at rt for 4 h. The reaction was quenched by addition of water carefully and extracted with EtOAc. The organic phase was washed with brine and water, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography to afford the title compound (2.4 g, 89% yield) as a white solid.


Step 3: Synthesis of 2-bromo-5-(4-ethoxypiperidin-4-yl)pyridine**[0419]**

[0420] To a solution of tert-butyl 4-(6-bromopyridin-3-yl)-4-ethoxypiperidine-1-carboxylate (2.4 g, 6.22 mmol) in dioxane (80 mL) was added HCl/dioxane (15 mL, 4.0 M) at 25 °C, and the resulting mixture was stirred at 25 °C for 2 h. LC-MS showed that the reaction was completed. Evaporated to dryness to afford the title compound (1.5 g) as HCl salt, which was used for the next step without further purification.

Step 4: Synthesis of 2-bromo-5-(4-ethoxy-1-isopropylpiperidin-4-yl)pyridine**[0421]**


[0422] A mixture of 2-bromo-5-(4-ethoxypiperidin-4-yl)pyridine (1.5 g, 5.25 mmol) and potassium carbonate (2.18 g, 15.7 mmol) in MeCN (100 mL) was added iodomethane (2.66 g, 15.7 mmol) and then stirred at 60 °C for 5 h. Concentrated, the residue was diluted with DCM/water (200 mL/200 mL) and extracted with DCM. The combined organic layers were dried over Na₂SO₄ and concentrated to give the title product (1.4 g), which was used for the next step without further purification.

R. Synthesis of 2-bromo-5-(1-(1-(difluoromethoxy)propan-2-yl)-4-methoxypiperidin-4-yl)pyridine**Step 1: Synthesis of 2-(4-(6-bromopyridin-3-yl)-4-methoxypiperidin-1-yl)propan-1-ol**

[0423]


[0424] To a solution of 2-bromo-5-(4-methoxypiperidin-4-yl)pyridine (10 g, 36.8 mmol) and 1-hydroxypropan-2-one (54.5 g, 736 mmol) in DCM (100 mL) and CH₃OH (100 mL) was added HOAc (1.0 mL) at 0 °C, and then sodium cyanoborohydride (11.5 g, 184 mmol) was added. The mixture was stirred at 35 °C for 16 h. After that, the mixture was quenched by addition of NaHCO₃ solution and extracted with DCM. The organic layers were concentrated and purified by silica gel column (DCM/CH₃OH = 10/1) to get the title compound (8.0 g, yield 66%) as a colorless oil. MS (ES⁺) C₁₄H₂₁BrN₂O₂ requires: 328, 330, found 329, 331 [M+H]⁺.

Step 2: Chiral separation of 2-(4-(6-bromopyridin-3-yl)-4-methoxypiperidin-1-yl)propan-1-ol

[0425]

[0426] The two enantiomers were separated on an AY-H column (250*4.6mm, 5um; Mobile Phase: n-Hexane (0.1% DEA): EtOH(0.1% DEA) = 70: 30; Temperature: 40 °C; Flow: 1.0 mL/min) and the fractions were measured at wavelengths 214nm and 254nm on a SHIMADZU Instrument.

Step 3: Synthesis of 2-bromo-5-(1-(1-(difluoromethoxy)propan-2-yl)-4-methoxypiperidin-4-yl)pyridine

[0427]

[0428] To a mixture of 2-(4-(6-bromopyridin-3-yl)-4-methoxypiperidin-1-yl)propan-1-ol (100 mg, 303 μmol) and CuI (86 mg, 0.45 mmol) in 4 mL of CH₃CN was added a solution of 2,2-difluoro-2-(fluorosulfonyl)acetic acid (268 mg, 1.51 mmol) dropwise in 2 mL of CH₃CN at 45 °C under N₂. The resulting mixture was stirred at 45 °C for 2 hrs. The solvent was removed in *vacuo* and the residue was purified by silica gel column (EtOAc/CH₃OH = 20/1) to afford the title product (40 mg, yield 35%) as a yellow oil. MS (ES⁺) C₁₅H₂₁BrF₂N₂O₂ requires: 378, 380, found 379, 381 [M+H]⁺.

S. Synthesis of 2-bromo-5-(4-methoxy-1-(1-methoxypropan-2-yl)piperidin-4-yl)pyridine

Step 1: Synthesis of (S)-2-bromo-5-(4-methoxy-1-(1-methoxypropan-2-yl)piperidin-4-yl)pyridine

5 [0429]



10 [0430] A mixture of 2-bromo-5-(4-methoxypiperidin-4-yl)pyridine (800 mg, 2.95 mmol), (R)-1-methoxypropan-2-yl methanesulfonate (992 mg, 5.90 mmol) and K_2CO_3 (1.22 g, 8.85 mmol) in CH_3CN (30 mL) was heated to 70 °C for 72 h. Concentrated and passed a column (silica gel, DCM:MeOH = 20: 1) to give the title product (0.5 g, 50%) as a yellow oil. MS (ES+) $C_{15}H_{23}BrN_2O_2$ requires: 342, found 343 $[M+H]^+$.

Step 2: Synthesis of (R)-2-bromo-5-(4-methoxy-1-(1-methoxypropan-2-yl)piperidin-4-yl)pyridine

15

[0431]




20 [0432] A mixture of 2-bromo-5-(4-methoxypiperidin-4-yl)pyridine hydrochloride (200 mg, 0.65 mmol), (S)-1-methoxypropan-2-yl methanesulfonate (218 mg, 1.30 mmol) and K_2CO_3 (269 mg, 1.95 mmol) in CH_3CN (5 mL) was heated to 70 °C for 72 h. Concentrated and passed a column (silica gel, DCM:MeOH = 20:1) to give the title product (100 mg, 45%) as a yellow oil. MS (ES+) $C_{15}H_{23}BrN_2O_2$ requires: 342, found 343 $[M+H]^+$.

25 **T. Synthesis of (3S,4S)-3-(4-bromophenyl)-1-isopropylpiperidin-4-ol**

Step 1: Synthesis of tert-butyl 3-(4-bromophenyl)-5,6-dihydropyridine-1(2H)-carboxylate

[0433]




30

35 [0434] A mixture of tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (10 g, 32.3 mmol), 1-bromo-4-iodobenzene (9.13 g, 32.3 mmol), [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II) (2.35 g, 3.22 mmol) and sodium carbonate (10.2 g, 96.8 mmol) in dioxane (80 mL) and water (20 mL) was purged with N_2 and stirred at 75 °C for 2 h. After that, the solution was cooled to RT and purified by silica gel column chromatography (petroleum

ether/ethyl acetate = 10/1 to 5/1) to afford the title compound (8 g, 73%) as a colorless oil. MS (ES+) $C_{16}H_{20}BrNO_2$ requires: 337, 339, found: 282, 284 [M-55]⁺.

Step 2: Synthesis of (3*r*,4*r*)-tert-butyl 3-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate


5 [0435]



[0436] To a cooled solution of borane-methyl sulfide complex (26.5 mL, 26.5 mmol) in anhydrous tetrahydrofuran (80 mL) under an atmosphere of nitrogen was added a solution of tert-butyl 3-(4-bromophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (7.5 g, 22.1 mmol) in tetrahydrofuran (20 mL). The resulting mixture was stirred at room temperature for 17 hours, then cooled in an ice bath, and sodium hydroxide (12.65 mL of a 2 N solution, 24.25 mmol) added in a dropwise manner, followed by hydrogen peroxide (9.2 mL of a 30% solution). The resulting mixture was stirred at RT for 3-5 hours, then poured into water (100 mL) and extracted with ethyl acetate (3 × 150 mL). The combined organic layers were washed with water (150 mL), saturated $NaHCO_3$ (150 mL) and saturated $NaCl$ (150 mL), dried over Na_2SO_4 , filtered, concentrated and purified by silica gel column chromatography (petroleum ether/ethyl acetate = 5/1 to 1/1) to afford the title compound (*trans*-, 3.5 g, 44%) as a white solid. MS (ES+) $C_{16}H_{20}BrNO_2$ requires: 355, 357, found: 300, 302 [M-55]⁺.


Step 3: Separation of (3*S*,4*S*)-tert-butyl 3-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate and (3*R*,4*R*)-tert-butyl 3-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate

[0437]




[0438] The enantiomers were separated on a S,S-Whelk-O1 column (4.6*100*5μm; Co-Solvent: MeOH (0.2% Methanol Ammonia); Column Temperature: 40°C; CO₂ Flow Rate:3.6.

Step 4: Synthesis of (3*S*,4*S*)-3-(4-bromophenyl)piperidin-4-ol

[0439]

- 5 **[0440]** To a solution of (3S,4S)-tert-butyl 3-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate (1.3 g, 3.64 mmol) in dioxane (20 mL) was added HCl/dioxane (4 mol/L, 10 mL). The mixture was stirred at RT for 12 h. After that, the solution was concentrated, and the residue was used in the next step without further purification. MS (ES+) $C_{11}H_{14}BrNO$ requires: 255, 257, found: 256, 258 $[M+H]^+$.


Step 5: Synthesis of (3S,4S)-3-(4-bromophenyl)-1-isopropylpiperidin-4-ol

10 **[0441]**

- 15 **[0442]** A mixture of (3S,4S)-3-(4-bromophenyl)piperidin-4-ol (900 mg, 3.51 mmol), 2-iodopropane (1.19 g, 7.02 mmol) and triethylamine (1.06 g, 10.5 mmol) in CH_3CN (30 mL) was stirred at 70 °C overnight. After that, the solution was concentrated and purified by silica gel column chromatography (petroleum ether/ethyl acetate/ NH_3 .MeOH=10/10/1) to afford the title compound (900 mg, 87%) as a white solid. MS (ES+) $C_{14}H_{20}BrNO$ requires: 297, 299, found: 298, 300 $[M+H]^+$.

20 **U. Synthesis of 4-(4-bromophenyl)-4-methoxy-1-(1-methoxypropan-2-yl)piperidine**

Step 1: Synthesis of tert-butyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate

[0443]


25

- 30 **[0444]** To a solution of 1,4-dibromobenzene (25.9 g, 110 mmol) in dry THF (250 mL) was added n-BuLi (2.5 M, 48.0 mL, 120.0 mmol) dropwise at -78 °C. The mixture was stirred at -78 °C for 1 h before tert-butyl 4-oxopiperidine-1-carboxylate (20 g, 100 mmol) in dry THF (100 mL) was added dropwise at -78 °C. The reaction mixture was stirred at -78 °C for another 1 h and warmed to RT slowly. LC-MS showed the reaction was completed. The reaction mixture was quenched with saturated NH_4Cl and extracted with EA (200 mLx3). The organic layer was washed with water and brine, dried over Na_2SO_4 , filtrated, concentrated and purified by silica gel

chromatograph (PE:EA = 5:1) to give the title product (22 g, yield: 62%) as a white solid. MS (ES+) $C_{16}H_{22}BrNO_3$ requires: 355, 357 found 356, 358 $[M+H]^+$.

Step 2: Synthesis of tert-butyl 4-(4-bromophenyl)-4-methoxypiperidine-1-carboxylate


5 [0445]



10 [0446] Sodium hydride (60%, 782 mg, 32.6 mmol) was added to a solution of tert-butyl 4-(4-bromophenyl)-4-hydroxypiperidine-1-carboxylate (7.8 g, 21.8 mmol) in DMF. This mixture was stirred at 20 °C for 30 min under nitrogen. Iodomethane (32.6mmol, 4.62 g) was then added. Stirred at 20 °C for 2 h. Dissolved in water and extracted with EA. EA phase was dried and purified by silica gel chromatograph, eluting with 1/4 EA/PE to give the title product as a yellow oil (8.0 g, ~80% in LCMS, yield 79%). MS (ES+) $C_{17}H_{24}BrNO_3$ requires: 369, 371, found: 282, 284 $[M+H]^+$.

15 **Step 3: Synthesis of 4-(4-bromophenyl)-4-methoxypiperidine HCl salt**


[0447]



20 [0448] A solution of 4 M HCl in dioxane (25 mL, 100.0 mmol) was added to a stirred solution of tert-butyl 4-(4-bromophenyl)-4-methoxypiperidine-1-carboxylate (3.7 g, 10.0 mmol) in MeOH (20 mL) at 25 °C, and stirred for 3 h under N_2 . After concentrated, the crude product (2.5 g, yield 89%, yellow solid) was used into the next reaction directly without further purification. MS (ES+) $C_{12}H_{16}BrNO$ requires: 269, 271, found: 270, 272 $[M+H]^+$.


25 **Step 4a: Synthesis of (R)-1-methoxypropan-2-yl methanesulfonate**

[0449]




30 [0450] To a mixture of (R)-1-methoxypropan-2-ol (2.0 g, 22.1 mmol) and triethylamine (6.70 g, 66.3 mmol) in DCM (10 mL) was added methanesulfonyl chloride (3.79 g, 33.1 mmol) dropwise at 0 °C. The mixture was stirred at rt for 16 hrs. The mixture was quenched with sat. $NaHCO_3$ solution, diluted with DCM, washed with water and dried over Na_2SO_4 anhydrous. The organic layer was concentrated in *vacuo* to afford the title compound (3.0 g, yield 80%) as a yellow oil

35 MS (ES+) $C_5H_{12}O_4S$ requires: 168, found: 169 $[M+H]^+$.

Step 4b: Synthesis of (S)-1-methoxypropan-2-yl methanesulfonate**[0451]**


- 5 **[0452]** To a mixture of (S)-1-methoxypropan-2-ol (1.5 g, 16.6 mmol) and triethylamine (5.03 g, 49.8 mmol) was added methanesulfonyl chloride (1.92 mL, 24.9 mmol) at 0 °C. The mixture was stirred at rt for 16 hrs. The mixture was diluted with DCM, washed with sat. NaHCO₃ solution and dried over Na₂SO₄ anhydrous. The organic layer was concentrated in *vacuo* to afford the title compound (2.3 g, yield 82%) as a yellow oil MS (ES⁺) C₅H₁₂O₄S requires: 168, found: 169 [M+H]⁺.
- 10

Step 5a: Synthesis of (S)-4-(4-bromophenyl)-4-methoxy-1-(1-methoxypropan-2-yl)piperidine**[0453]**

- 15 **[0454]** A solution of 4-(4-bromophenyl)-4-methoxypiperidine (250 mg, 0.93 mmol), potassium carbonate (127 mg, 0.93 mmol), potassium iodide (153 mg, 0.93 mmol) and (R)-1-methoxypropan-2-yl methanesulfonate (186 mg, 1.11 mmol) in DMF (5 mL) was stirred at 60 °C for 5 h under N₂. Dissolved in EA, and washed with water and brine. After dried and concentrated, the residue was purified by silica gel chromatograph, eluting with 1/10 MeOH/EA to give desired product (100 mg, 65% purity in LCMS, yield 21%) as a brown paste. MS (ES⁺) C₁₆H₂₄BrNO₂ requires: 341, found: 342, 344 [M+H]⁺.
- 20

Step 5b: Synthesis of (R)-4-(4-bromophenyl)-4-methoxy-1-(1-methoxypropan-2-yl)piperidine

25

[0455]


- 30 **[0456]** A solution of 4-(4-bromophenyl)-4-methoxypiperidine (250 mg, 0.93 mmol), potassium carbonate (127 mg, 0.93 mmol), potassium iodide (153 mg, 0.93 mmol) and (S)-1-methoxypropan-2-yl methanesulfonate (155 mg, 0.93 mmol) in DMF (5 mL) was stirred at 60 °C for 5 h under N₂. Dissolved in EA, and washed with water and brine. After dried and concentrated, the residue was purified by silica gel chromatograph, eluting with 1/10 MeOH/EA, to give the title compound (106 mg,

85% purity in LCMS, yield 34%) as a yellow paste. MS (ES+) $C_{16}H_{24}BrNO_2$ requires: 341, 343, found: 342, 344 $[M+H]^+$.

V. Synthesis of (S)-5-(1-isopropylpiperidin-3-yl)pyridin-2-yl trifluoromethanesulfonate

5 Step 1: Synthesis of tert-butyl 3-(6-(benzyloxy)pyridin-3-yl)-5,6-dihydropyridine-1(2H)-carboxylate


[0457]



15 [0458] A mixture of tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (5 g, 16.1 mmol), 2-(benzyloxy)-5-bromopyridine (6.36 g, 24.1 mmol), Pd(dppf)Cl₂ (818 mg, 1.12 mmol) and K₂CO₃ (4.44 g, 32.2 mmol) in dioxane/water (40 mL/5 mL) was purged with N₂ for three times and stirred at 100 °C for 16 hrs. The mixture was concentrated and purified by flash column chromatography (PE/EtOAc = 10:1) to afford the title compound (5.0 g, yield 84%) as a yellow oil. MS (ES+) $C_{22}H_{26}N_2O_3$ requires: 366, found: 367 $[M+H]^+$.


20 Step 2: Synthesis of (S)-tert-butyl 3-(6-hydroxypyridin-3-yl)piperidine-1-carboxylate (P1) and (R)-tert-butyl 3-(6-hydroxypyridin-3-yl)piperidine-1-carboxylate (P2)

[0459]




25 [0460] A mixture of tert-butyl 6'-(benzyloxy)-5,6-dihydro-[3,3'-bipyridine]-1(2H)-carboxylate (2.5 g, 6.82 mmol) and Pd/C (1.44 g) in EtOAc (10 mL) was stirred at RT for 16 hrs under H₂ (H₂ balloon). The mixture was filtered through a pad of Celite and purified by Prep-HPLC to afford racemic compound (1.1 g) as a white solid, which was separated by chiral HPLC (Chiral condition: Co-Solvent: MeOH(0.2% Methanol Ammonia); Column: OZ-H 100*4.6mm 5um; Column Temperature: 36.8; CO₂ Flow Rate:3; Co-Solvent Flow Rate:1) to afford the title compound **P1** (500 mg, yield 26%) as a white solid. MS (ES+) $C_{15}H_{22}N_2O_3$ requires: 278, found: 279 $[M+H]^+$ and **P2** (500 mg, yield 26%) as a white solid. MS (ES+) $C_{15}H_{22}N_2O_3$ requires: 278, found: 279 $[M+H]^+$.

35 Step 3: Synthesis of (S)-tert-butyl 3-(6-(trifluoromethylsulfonyloxy)pyridin-3-yl)piperidine-1-carboxylate

[0461]**[0462]** To a mixture of (S)-tert-butyl 3-(6-hydroxypyridin-3-yl)piperidine-1-


- 5 carboxylate (500 mg, 1.79 mmol) and pyridine (431 μ L, 5.37 mmol) in DCM (5 mL) was added trifluoromethanesulfonyl anhydride (450 μ L, 2.68 mmol) dropwise at 0 °C. The mixture was stirred at 0 °C for 30 min, diluted with DCM, washed with ice water and dried over Na₂SO₄ anhydrous. The organic layer was concentrated in *vacuo* to afford the title compound (700 mg, yield 95%) as a yellow oil. MS (ES+)
- 10 C₁₆H₂₁F₃N₂O₅S requires: 410, found: 411 [M+H]⁺.

Step 4: Synthesis of (S)-5-(piperidin-3-yl)pyridin-2-yl trifluoromethanesulfonate hydrochloride**[0463]**

15

[0464] To a mixture of (S)-tert-butyl 3-(6-(((trifluoromethyl)sulfonyl)oxy)pyridin-3-yl)piperidine-1-carboxylate (400 mg, 974 μ mol) in DCM (2 mL) was added HCl in dioxane (1.21 mL, 4.84 mmol). The mixture was stirred at RT for 4 hrs. The mixture was concentrated in *vacuo* to afford the title compound (330 mg, crude) as a yellow solid. MS (ES+) C₁₁H₁₄ClF₃N₂O₃S requires: 310, found: 311 [M+H]⁺.

20

Step 5: Synthesis of (S)-5-(1-isopropylpiperidin-3-yl)pyridin-2-yl trifluoromethanesulfonate**[0465]**

25


[0466] A mixture of (S)-5-(piperidin-3-yl)pyridin-2-yl trifluoromethanesulfonate hydrochloride (330 mg, 951 μ mol), 2-iodopropane (322 mg, 1.90 mmol) and triethylamine (288 mg, 2.85 mmol) in ACN (5 mL) was stirred at 60 °C for 16 hrs. The mixture was concentrated and purified by flash column chromatography (DCM/MeOH = 10:1) to afford the title compound (250 mg, yield 74%) as a yellow oil. MS (ES+) C₁₄H₁₉F₃N₂O₃S requires: 352, found: 353 [M+H]⁺.

30

W. Synthesis of 4-(4-bromo-3-chlorophenyl)-1-isopropyl-1,2,3,6-tetrahydropyridine:

Step 1: Synthesis of tert-butyl 4-(4-bromo-3-chlorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate


5 [0467]



10 [0468] To a solution of 1-bromo-2-chloro-4-iodobenzene (4.9, 15.4 mmol) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (4.3 g, 13.9 mmol) in dioxane/H₂O (50 mL/20 mL) was added Pd(dppf)Cl₂ (626 mg, 0.77 mmol) and K₂CO₃ (6.4 g, 46.2 mmol) at room temperature under nitrogen. The resulting mixture was stirred at 70 °C for 2 hours. Cooled to room temperature and extracted with EtOAc after additional water was added. The organic was washed with brine, dried over Na₂SO₄ and concentrated to dryness to afford the crude product which was purified by flash chromatography on silica gel eluting with PE/EA (10: 1) to afford the title product (3.6 g, yield 70%) as a yellow solid. MS (ES⁺) C₁₆H₁₉BrClNO₂ requires: 371, found 372 [M+H]⁺.

Step 2: Synthesis of 4-(4-bromo-3-chlorophenyl)-1,2,3,6-tetrahydropyridine TFA salt


20 [0469]



25 [0470] To a solution of tert-butyl 4-(4-bromo-3-chlorophenyl)-5,6-dihydropyridine-1(2H)-carboxylate (3.6 g, 9.7 mmol) in DCM (20 mL) was added TFA (5 mL) dropwise at room temperature. The resulting mixture was stirred at room temperature for 3 hours; LC-MS showed that the reaction was completed. The solution was evaporated to dryness to afford the crude product which was used for the next step without further purification (3.4 g, TFA salt). MS (ES⁺) C₁₁H₁₁BrClN requires: 271, found 272 [M+H]⁺.

30 **Step 3: Synthesis of 4-(4-bromo-3-chlorophenyl)-1-isopropyl-1,2,3,6-tetrahydropyridine**

[0471]



5 **[0472]** To a solution of 4-(4-bromo-3-chlorophenyl)-1,2,3,6-tetrahydropyridine TFA salt (3.4 g, 9.2 mmol) in acetonitrile (15 mL) was added TEA (2.8 g, 27.6 mmol), followed by addition of isopropyl iodide (4.7 g, 27.6 mmol). The solution was heated at 70 °C for 6 hours; LC-MS showed that the reaction was completed. The crude product was purified by flash chromatography on silica gel eluting with PE/EA (1:1 to 1:3) to give the title product (2.4 g, yield 83%) as viscous oil. MS (ES+) C₁₄H₁₇BrClN requires: 313, found 314 [M+H]⁺.


10

X. Synthesis of 4-(4-bromo-3-fluorophenyl)-1-isopropyl-4-methoxypiperidine

Step 1: Synthesis of *tert*-butyl 4-(4-bromo-3-fluorophenyl)-4-hydroxypiperidine-1-carboxylate

15


[0473]




20 **[0474]** To a solution of 1-bromo-2-fluoro-4-iodobenzene (15.0 g, 50.1 mmol) in THF (150 mL) at -78 °C was added n-BuLi (20 mL, 2.4 M in hexane) dropwise. The solution was stirred at -78 °C for 2 h. Then, a solution of *tert*-butyl 4-oxopiperidine-1-carboxylate (10 g, 50.1 mmol) in THF (20 mL) was added dropwise. After that, the resulting solution was stirred at -78 °C for 2 h. The reaction was quenched carefully by addition of water and extracted with EtOAc. The organic layer was washed with water and brine, concentrated and purified by silica gel column chromatography (petroleum ether/ethyl acetate = 10/1 to 2/1) to afford the title product (3 g, 16%) as a white solid. MS (ES+) C₁₆H₂₁BrFNO₃ requires: 373, 375, found: 300, 302 [M-73]⁺.

25


Step 2: Synthesis of *(tert*-butyl 4-(4-bromo-3-fluorophenyl)-4-methoxypiperidine-1-carboxylate

[0475]

[0476] To a solution of tert-butyl 4-(4-bromo-3-fluorophenyl)-4-hydroxypiperidine-1-carboxylate (3 g, 8.01 mmol) in dry DMF (50 mL) was added NaH (60%) (416 mg, 10.4 mmol) at 0 °C. The mixture was stirred at 0 °C for 1 h. Then, iodomethane (1.7 g, 12.0 mmol) was added. The mixture was stirred at RT overnight. After that, the mixture was slowly poured into ice water (200 mL) and stirred for 1 h. The solid was collected *via* filtration and purified by silica gel column chromatography (petroleum ether/ethyl acetate = 10/1 to 4/1) to afford the title compound (2.5 g, 80%) as a white solid. MS (ES+) C₁₇H₂₃BrFNO₃ requires: 387, 389, found: 300, 302 [M-87]⁺.

Step 3: Synthesis of 4-(4-bromo-3-fluorophenyl)-4-methoxypiperidine**[0477]**

[0478] To a solution of tert-butyl 4-(4-bromo-3-fluorophenyl)-4-methoxypiperidine-1-carboxylate (2.5 g, 6.43 mmol) in dioxane (20 mL) was added HCl/dioxane (4 N, 20 mL). The mixture was stirred at RT for 12 h. After that, the solution was concentrated and used in next step without further purification. MS (ES+) C₁₂H₁₅BrFNO requires: 287, 289, found: 288, 290 [M+H]⁺.


Step 4: Synthesis of 4-(4-bromo-3-fluorophenyl)-1-isopropyl-4-methoxypiperidine**[0479]**

[0480] A mixture of 4-(4-bromo-3-fluorophenyl)-4-methoxypiperidine (1.6 g, 5.55 mmol), 2-iodopropane (1.88 g, 11.1 mmol) and triethylamine (1.67 g, 16.6 mmol) in CH₃CN (30 mL) was stirred at 80 °C overnight. After that, the solution was concentrated and purified by silica gel column chromatography (DCM/ MeOH=10/1) to afford the title compound (1.5 g, 82%) as a yellow solid. MS (ES+) C₁₅H₂₁BrFNO requires: 329, 331, found: 330, 332 [M+H]⁺.

Y. Synthesis of tert-butyl 4-methyl-4-(4-(trifluoromethylsulfonyloxy)phenyl)piperidine-1-carboxylate

Step 1: Synthesis of tert-butyl 4-(4-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate


5 [0481]



10 [0482] To a solution of 1-bromo-4-methoxybenzene (6.0 g, 32.0 mmol) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (10.0 g, 32.0 mmol) in dioxane/H₂O (60 mL/30 mL) was added Pd(t-Bu₃P)₂ (163 mg, 0.32 mmol) and Na₂CO₃ (6.8 g, 64 mmol) at room temperature under nitrogen. The resulting mixture was stirred at 70 °C for 2 hours. LC-MS showed that the reaction was completed. Cooled to room temperature and extracted with EtOAc after additional water was added. The organic was washed with brine, dried over Na₂SO₄ and concentrated to dryness to afford the crude product which was purified by flash chromatography on silica gel eluting with PE/EA (10: 1) to afford the title product (9.3 g, yield 97%) as a white solid. MS (ES⁺) C₁₇H₂₃NO₃ requires: 289, found 234 [M+H-56]⁺.

20 **Step 2: Synthesis of tert-butyl 4-(4-methoxyphenyl)-4-methyl-3,4-dihydropyridine-1(2H)-carboxylate**

[0483]




25 [0484] To a solution of tert-butyl 4-(4-methoxyphenyl)-5,6-dihydropyridine-1(2H)-carboxylate (9.3 g, 32.0 mmol) in THF (50 mL) was added n-BuLi (25.6 mL, 2.5 M in hexane) at -15 °C under nitrogen. The blood-red solution formed as the end of dropwise addition of n-BuLi. The reaction mixture was stirred at this temperature for 15 minutes. Me₂SO₄ (8.1 g, 64.0 mmol) was added dropwise to the above solution at -15 °C. The resulting mixture was stirred at room temperature for 30 minutes. NH₄OH (40 mL, 2.0 M) was added and the resulting mixture was extracted with EtOAc after water was added. The combined organics were washed with water and brine, dried over Na₂SO₄, and concentrated to dryness to afford the crude product

which was purified by flash chromatography on silica gel eluting with PE/EA (5: 1) to give the title product (1.5 g, yield 15%) MS (ES+) $C_{18}H_{25}NO_3$ requires: 303, found 248 $[M+H-56]^+$.

Step 3: Synthesis of 4-(4-methyl-1,2,3,4-tetrahydropyridin-4-yl)phenol


5 [0485]



10 [0486] To a solution of tert-butyl 4-(4-methoxyphenyl)-4-methyl-3,4-dihydropyridine-1(2H)-carboxylate (750 mg, 2.5 mmol) in anhydrous DCM (15 mL) was added BBr_3 (2 mL, 17% in DCM) at 0 °C. The resulting mixture was stirred at 0 °C for 6 hours; LC-MS showed that reaction was completed. The reaction mixture was used for the next step without work-up and purification.

Step 4: Synthesis of tert-butyl 4-(4-(tert-butoxycarbonyloxy)phenyl)-4-methyl-3,4-dihydropyridine-1(2H)-carboxylate

15 [0487]




20 [0488] To a solution of 4-(4-methyl-1,2,3,4-tetrahydropyridin-4-yl)phenol from last step was added TEA (7 mL) at 0 °C to adjust pH value to 9.0. Boc_2O (1.6 g, 7.5 mmol) and catalytic DMAP (30 mg, 0.25 mmol) was added. The resulting mixture was stirred at room temperature for 16 hours; LC-MS showed that the reaction was completed. Water was added and extracted with EtOAc (x2). The combined organics were washed with brine, dried over Na_2SO_4 , and concentrated to dryness to afford the crude product which was purified by flash chromatography on silica gel eluting with PE/EA (20: 1 to 10: 1) to afford the title product (470 mg, yield 51%) MS (ES+) $C_{22}H_{31}NO_5$ requires: 389, found 390 $[M+H]^+$.

25

Step 5: Synthesis of tert-butyl 4-(4-(tert-butoxycarbonyloxy)phenyl)-4-methylpiperidine-1-carboxylate


[0489]



[0490] To a solution of tert-butyl 4-(4-(tert-butoxycarbonyloxy)phenyl)-4-methyl-3,4-dihydropyridine-1(2H)-carboxylate (470 mg, 1.2 mmol) in MeOH (10 mL) was added Pd/C (100 mg, 10% wet) at room temperature. The resulting mixture was stirred under hydrogen balloon at room temperature for 16 hours; LC-MS showed that the reaction was completed. Filtered off, the filtration was evaporated to dryness to afford the title product (410 mg, yield 87%) MS (ES+) C₂₂H₃₃NO₅ requires: 391, found 392 [M+H]⁺.


Step 6: Synthesis of tert-butyl 4-(4-hydroxyphenyl)-4-methylpiperidine-1-carboxylate

[0491]



[0492] To a solution of tert-butyl 4-(4-(tert-butoxycarbonyloxy)phenyl)-4-methylpiperidine-1-carboxylate (410 mg, 1.0 mmol) in MeOH (5.0 mL) was added K₂CO₃ (691 mg, 5.0 mmol) at room temperature. The resulting mixture was stirred at room temperature at for 1 hour. LC-MS showed that the reaction was completed. The mixture was then extracted with EtOAc after water was added. The organic was washed with brine, dried over Na₂OS₄, and concentrated to dryness to afford the title product (310 mg, quantitative) which was used for the next step without further purification MS (ES+) C₁₇H₂₅NO₃ requires: 291, found 236 [M+H-56]⁺.


Step 7: Synthesis of tert-butyl 4-methyl-4-(4-(trifluoromethylsulfonyloxy)phenyl)piperidine-1-carboxylate

[0493]

[0494] To a solution of tert-butyl 4-(4-hydroxyphenyl)-4-methylpiperidine-1-carboxylate (310 mg, 1.1 mmol) in DCM (15.0 mL) was added TEA (333 mg, 3.3 mmol) at room temperature. Tf₂O (372 mg, 1.3 mmol) in DCM (3 mL) was added at 0 °C dropwise during 5 minutes. The resulting mixture was stirred at 0 °C for 4 hours; LC-MS showed that the reaction was completed. The mixture was then extracted with EtOAc after water was added. The organic was washed with brine, dried over Na₂SO₄ and concentrated to dryness to afford the title product, which was used for the next step without further purification (270 mg, 8%) MS (ES⁺) C₁₈H₂₄F₃NO₅S requires: 423, found 424 [M+H]⁺.


Z. Synthesis of 2-bromo-5-(4-methoxy-1-(oxetan-3-yl)piperidin-4-yl)pyridine:

Step 1: Synthesis of tert-butyl 4-(6-bromopyridin-3-yl)-4-hydroxypiperidine-1-carboxylate

[0495]

[0496] To a solution of 2-bromo-5-iodopyridine (11.3 g, 40 mmol) in 120 mL of THF was added nBuLi (17.6 mL, 2.5M in hexane) dropwise under N₂ at -78 °C. The resulting solution was stirred at -78 °C for 2 h. Then a solution of tert-butyl 4-oxopiperidine-1-carboxylate (7.96 g, 40.0 mmol) in 20 mL of THF was added dropwise. The mixture was stirred at -78 °C for another 1 h. The reaction was quenched by addition of sat. NH₄Cl solution. After extracted with EtOAc, the organic phase was dried over Na₂SO₄ and concentrated. The residue was purified by silica gel column (PE/EtOAc = 10:1) to afford the title product (8 g, yield 56%) as a yellow solid. MS (ES⁺) C₁₅H₂₁BrN₂O₃ requires: 356, 358, found 357, 359 [M+H]⁺.


Step 2: Synthesis of tert-butyl 4-(6-bromopyridin-3-yl)-4-methoxypiperidine-1-carboxylate

[0497]

[0498] To a solution of tert-butyl 4-(6-bromopyridin-3-yl)-4-hydroxypiperidine-1-carboxylate (6.5 g, 18.1 mmol) in 20 mL of DMF was added NaH (60%, 3.62 g, 90.5 mmol) at 0 °C. The mixture was stirred at 0 °C for 30 min. Then MeI (7.7 g, 54.3 mmol) was added. The resulting mixture was stirred at RT overnight. The reaction was quenched by addition of water carefully, and extracted with EtOAc. The organic phase was washed with brine, dried over Na₂SO₄ and concentrated in *vacuo* to afford crude product which was used in the next step without purification. MS (ES+) C₁₆H₂₃BrN₂O₃ requires: 370, 372, found 315, 317 [M+H-56]⁺.

Step 3: Synthesis of 2-bromo-5-(4-methoxypiperidin-4-yl)pyridine hydrochloride


[0499]



[0500] To a solution of tert-butyl 4-(6-bromopyridin-3-yl)-4-methoxypiperidine-1-carboxylate (740 mg, 2 mmol) in 5 mL of dioxane was added HCl (4 mL, 4M in dioxane). The resulting mixture was stirred at RT for 4 h. The solvent was removed in *vacuo* to afford crude product which was used in the next step without purification. MS (ES+) C₁₁H₁₅BrN₂O requires: 270, 272, found 271, 273 [M+H]⁺.

Step 4: Synthesis of 2-bromo-5-(4-methoxy-1-(oxetan-3-yl)piperidin-4-yl)pyridine


[0501]




[0502] To a solution of 2-bromo-5-(4-methoxypiperidin-4-yl)pyridine hydrochloride (400 mg, 1.30 mmol) and oxetan-3-one (468 mg, 6.50 mmol) in 10 mL of DCM and 10 mL of MeOH was added NaCNBH₃ (410 mg, 6.5 mmol) and drops of AcOH. The resulting solution was stirred at RT overnight. The reaction was quenched by addition of water and extracted with EtOAc. The organic phase was dried over Na₂SO₄ and concentrated in *vacuo* to afford crude product which was used in the next step without purification. MS (ES+) C₁₄H₁₉BrN₂O₂ requires: 326, 328, found 327, 329 [M+H]⁺.

AA. Synthesis of (S)-tert-butyl 3-(6-bromopyridin-3-yl)-3-ethoxypiperidine-1-carboxylate and (R)-tert-butyl 3-(6-bromopyridin-3-yl)-3-ethoxypiperidine-1-carboxylate


Step 1: Synthesis of tert-butyl 3-(6-bromopyridin-3-yl)-3-hydroxypiperidine-1-carboxylate

[0503]

[0504] To a solution of 2-bromo-5-iodopyridine (15 g, 52.8 mmol) in dry THF (300 mL) at -78 °C was added n-BuLi (2.5 M, 21 mL), and stirred at -78 °C for 1 hrs, followed by addition of tert-butyl 3-oxopiperidine-1-carboxylate (10.5 g, 52.8 mmol) in THF. The reaction mixture was stirred at -78 °C~60 °C for 2 hrs, quenched by NH₄Cl/H₂O and extracted by EA. The organic was dried to give a residue, which was purified by flash chromatography to afford the title compound (8.4 g, 45%). MS (ES+) C₁₅H₂₁BrN₂O₃ requires: 356, found: 357 [M+H]⁺.

Step 2: Synthesis of 6-bromo-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine**[0505]**


[0506] To a solution of tert-butyl 3-(6-bromopyridin-3-yl)-3-hydroxypiperidine-1-carboxylate (6.5 g, 18.1 mmol) in DMF (150 mL) at 0 °C was added 60% NaH (3.00 g, 125 mmol), and stirred at 25 °C for 2 hrs, followed by addition of bromoethane (7.88 g, 72.4 mmol). The mixture was stirred at 25 °C for another 5 hrs. Quenched by water and extracted with EA. Concentrated and purified by flash chromatography to give the title compound (5.7 g, 97%). MS (ES+) C₁₇H₂₅BrN₂O₃ requires: 385, found: 386[M+H]⁺.

Step 3: Chiral separation of 6-bromo-4-(piperazin-1-yl)pyrrolo[1,2-b]pyridazine**[0507]**


absolute configuration of P1 and P2 unknown.

[0508] The title compound (5.7g) was separated by Chiral-HPLC (Chiral conditions: Co-Solvent: PA (0.1%DEA); Column: Cellulose-SC 4.6*100mm 5um; Column Temperature:39.2; CO₂ Flow Rate: 3.6; Co-Solvent Flow Rate: 0.4) to give P1 (1.8 g) and P2 (1.8 g).


BB. Synthesis of 3-(trimethylsilyloxy)oxetane-3-carbonitrile

[0509]

[0510] To a mixture of oxetan-3-one (500 mg, 6.93 mmol) and triethylamine (1.39 g, 13.8 mmol) in DCM (5 mL) was added trimethylsilyl cyanide (1.71 g, 17.3 mmol). The mixture was stirred at RT for 16 h, then diluted with DCM, washed with saturated Na₂CO₃ solution and dried with Na₂SO₄ anhydrous. The organic layer was separated and concentrated in *vacuo* to afford the title compound (800 mg, yield 67%) as a brown oil.

CC. Synthesis of 3-cyanooxetan-3-yl 1H-imidazole-1-carboxylate**[0511]**

[0512] To a mixture of 3-((trimethylsilyl)oxy)oxetane-3-carbonitrile (50 mg, 291 μmol) in DCM (2 mL) was added CDI (51.8 mg, 320 μmol). The mixture was stirred at rt for 4 hrs, which was added directly into the next reaction solution. MS (ES⁺) C₈H₇N₃O₃ requires: 193, found: 194 [M+H]⁺.

DD. Synthesis of (3S,4S)-3-(4-bromophenyl)-4-(difluoromethoxy)-1-isopropylpiperidine**[0513]**

[0514] To a mixture of (3S,4S)-3-(4-bromophenyl)-1-isopropylpiperidin-4-ol (900 mg, 3.01 mmol) and CuI (571 mg, 3.01 mmol) in dry CH₃CN (10 mL) was dropwise added a solution of 2,2-difluoro-2-(fluorosulfonyl)acetic acid (2.67 g, 15.00 mmol) in dry CH₃CN (5 mL) at 45 °C. After that, the solution was stirred at 45 °C for 2 hrs. The reaction was quenched with water (1.0 mL) and extracted. The combined organic layers were washed with water, dried over Na₂SO₄, filtered and concentrated. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate /NH₃.MeOH = 10/10/1) to afford the title compound (500 mg, 48%) as a yellow oil. MS (ES⁺) C₁₅H₂₀BrF₂NO requires: 347, 349, found: 348, 350 [M+H]⁺.

Example 25. Synthesis of other Exemplary Compounds.

[0515] Table 1 indicates which Synthetic Protocol 1, 2, 3, or 4 described in the disclosure was used to synthesize various compounds described herein. Blank values indicate that a synthetic scheme other than one of Synthetic Protocols 1-4 was used and that the synthetic scheme for such compound is set forth in the Examples.

Table 1. Synthetic Protocols used for Exemplary Compounds

Compound	Synthetic Scheme
100	2
101	2
102	2
103	1
104	1
105	2
106	2
107	2
108	2
109	2
110	1
111	Ex. 18
112	1
113	2
114	1
115	4
116	2
117	1
118	1
119	2
120	2
121	Ex. 19
122	2
123	2
124	2
125	2
126	2

Compound	Synthetic Scheme
127	1
128	2
129	2
130	1
131	1
132	4
133	2
134	2
135	1
136	1
137	2
138	1
139	2
140	2
141	4
142	4
143	2
144	2
145	2
146	2
147	2
148	1
149	1
150	4
151	2
152	1
153	1
154	1
155	2
156	2
157	Ex. 16
158	2
159	2

Compound	Synthetic Scheme
160	2
161	1
162	1
163	1
164	2
165	2
166	2
167	2
168	2
169	2
170	1
171	1
172	2
173	1
174	2
175	1
176	1
177	2
178	2
179	2
180	2
181	2
182	4
183	2
184	1
185	1
186	1
187	2
188	2
189	2
190	2
191	2
192	2

Compound	Synthetic Scheme
193	2
194	1
195	4
196	2
197	4
198	2
199	1
200	2
201	1
202	2
203	1
204	1
205	3
206	2
207	2
208	4
209	2
210	2
211	2
212	1
213	1
214	2
215	3
216	2
217	2
218	2
219	2
220	2
221	1
222	2
223	3
224	4
225	4

Compound	Synthetic Scheme
226	3
227	3
228	2
229	1
230	3
231	2
232	4
233	2
234	3
235	2
236	2
237	4
238	2
239	2
240	4
241	1
242	2
243	2
244	3
245	2
246	2
247	2
248	2
249	2
250	3
251	4
252	2
253	2
254	2
255	2
256	2
257	2
258	2

Compound	Synthetic Scheme
259	1
260	Ex. 15
261	1
262	1
263	1
264	1
265	2
266	2
267	2
268	2
269	2
270	2
271	2
272	1
273	1
274	1
275	4
276	2
277	4
278	1
279	1
280	1
281	3
282	2
283	2
284	2
285	2
286	4
287	2
288	2
289	2
290	2
291	3

Compound	Synthetic Scheme
292	1
293	3
294	3
295	2
296	1
297	2
298	4
299	4
300	4
301	3
302	3
303	1
304	1
305	2
306	1
307	2
308	1
309	1
310	1
311	4
312	2
313	2
314	1
315	2
316	2
317	4
318	1
319	1
320	2
321	1
322	2
323	2
324	4

Compound	Synthetic Scheme
325	2
326	2
327	1
328	4
329	1
330	1
331	1
332	3
333	2
334	1
335	1
336	3
337	4
338	4
339	2
340	4
341	4
342	4
343	2
344	1
345	2
346	4
347	4
348	4
349	2
350	1
351	2
352	2
353	4
354	2
355	1
356	4
357	2

Compound	Synthetic Scheme
358	4
359	1
360	2
361	3
362	2
363	1
364	4
365	4
366	4
367	1
368	3
369	4
370	4
371	2
372	4
373	2
374	2
375	2
376	1
377	4
378	2
379	4
380	4
381	4
382	4
383	4
384	4
385	4
386	4
387	4
388	2
389	2
390	1

Compound	Synthetic Scheme
391	2
392	4
393	4
394	4
395	4
396	4
397	2
398	4
399	2
400	2
401	4
402	1
403	Ex. 17
404	4
405	4
406	4
407	2
408	4
409	4
410	4
411	4
412	4
413	4
414	4
415	4
416	4
417	2
418	2
419	1
420	4
421	4
422	3
423	4

Compound	Synthetic Scheme
424	4
425	2
426	4
427	1
428	4
429	3
430	4
431	4
432	4
433	3
434	4
435	4
436	2
437	2
438	1
439	3
440	3
441	4
442	4
443	1
444	4
445	4
446	4
447	3
448	4
449	2
450	2
451	2
452	5
453	2
454	2
455	2
456	2

Compound	Synthetic Scheme
457	2
458	2
459	1
460	1
461	2
462	2
463	2
464	2
465	5
466	5
467	5
468	2
469	2
470	2
471	2
472	2
473	2
474	2
475	2
476	5
477	5
478	2
479	2
480	2
481	2
482	2
483	2
484	2
485	2
486	2
487	2
488	2
489	2

Compound	Synthetic Scheme
490	2
491	2
492	2
493	2
494	2
495	2
496	2
497	2
498	2
499	5
500	2
501	2
502	2
503	2
504	2
505	2
506	2
507	2
508	2
509	4
510	4
511	4
512	2
513	2
514	4
515	2
516	2
517	2
518	2
519	2
520	2
521	2
522	1

Compound	Synthetic Scheme
523	1
524	2
525	4
526	4
527	4
528	2
529	4
530	4
531	2
532	4
533	2
534	4
535	4
536	4
537	2
538	2
539	2
540	2
541	2
542	2
543	2
544	2
545	2
546	2
547	4
548	2
549	2
550	4
551	4
552	4
553	2
554	4
555	2

Compound	Synthetic Scheme
556	2
557	2
558	2
559	2
560	2
561	2
562	2
563	2
564	2
565	2
566	2
567	2
568	2
569	2
570	2
571	5
572	4
573	4
574	4
575	2
576	2
577	2
578	2
579	4
580	4
581	5
582	5
583	2
584	5
585	2
586	2
587	2
588	2

Compound	Synthetic Scheme
589	2
590	4
591	2
592	2
593	2
594	2
595	4
596	4
597	2
598	2
599	2
600	2
601	2
602	5
603	4
604	4
605	2
606	2
607	4
608	4
609	2
610	2
611	2
612	5
613	2
614	2
615	2
616	2
617	4
618	4
619	2
620	2
621	4

Compound	Synthetic Scheme
622	2
623	4
624	2
625	2
626	2
627	4
628	2
629	4
630	2
631	2
632	4
633	2
634	2
635	2
636	2
637	4
638	4
639	2
640	2
641	2
642	2
643	2
644	2
645	2
646	2
647	2
648	5
649	5
650	5
651	2
652	4
653	2
654	2

Compound	Synthetic Scheme
655	2
656	2
657	2
658	2
659	2
660	2
661	4
662	2
663	4
664	4
665	3
666	4
667	4
668	2
669	2
670	4
671	2
672	2
673	2
674	2
675	2
676	4
677	4
678	2
679	2
680	2
681	4
682	2
683	2
684	2
685	2
686	2
687	2

Compound	Synthetic Scheme
688	2
689	2
690	2
691	2
692	2
693	2
694	5
695	2
696	2
697	2
698	2
699	2
700	2
701	5
702	4
703	4
704	4
705	4
706	2
707	2
708	4
709	4
710	2
711	2
712	2
713	2
714	2
715	2
716	2
717	2
718	2
719	2
720	2

Compound	Synthetic Scheme
721	2
722	2
723	2
724	2
725	2
726	2
727	2
728	4
729	2
730	2
731	2
732	2
733	2
734	2
735	2
736	2
737	5
738	2
739	2
740	2
741	2
742	4
743	2
744	2
745	2
746	2
747	2
748	2
749	2
750	2
751	2
752	2
753	2

Compound	Synthetic Scheme
754	2
755	2
756	2
757	2
758	2
759	2
760	2
761	2
762	2
763	2
764	2
765	2
766	2
767	2
768	2
769	2
770	2
771	2
772	2
773	2
774	2
775	2
776	2
777	2
778	2
779	2
780	2
781	2
782	2
783	2
784	2
785	2
786	2

Compound	Synthetic Scheme
787	2
788	2
789	3
790	3
791	2
792	2
793	3
794	3
795	3
796	2
797	2
798	2
799	2
800	2
801	2
802	2
803	3
804	2
805	2
806	2
807	2
808	2
809	3
810	3
811	2
812	3
813	2
814	3
815	2
816	2
817	2
818	2
819	2

Compound	Synthetic Scheme
820	2
821	2
822	3
823	2
824	2
825	3
826	3
827	3
828	3
829	3
830	2
831	2
832	3
833	2
834	2
835	2
836	3
837	3
838	3
839	2
840	3
841	3
842	3
843	2
844	2
845	2
846	2
847	3
848	2
849	2
850	2
851	3
852	2

Compound	Synthetic Scheme
853	3
854	3
855	2
856	2
857	2
858	2
859	2
860	2
861	2
862	2
863	2
864	3
865	3
866	3
867	3
868	2
869	2
870	3
871	3
872	3
873	2
874	2
875	2
876	3
877	3
878	3
879	3
880	2
881	2
882	2
883	2
884	2
885	2

Compound	Synthetic Scheme
886	2
887	2
888	2
889	2
890	3
891	2
892	2
893	2
894	2
895	2
896	2
897	2
898	2
899	2

Example 26. ALK Binding Assays

[0516] Binding assays were conducted using the LANTHASCREEN® technology (ThermoFisher Scientific). LANTHASCREEN® is a competitive binding, displacement assay where the assumed steady state occupancy of the binding site is measured using a time-resolved, fluorescence energy transfer (TR-FRET) readout between a fluorescent tracer and Europium (Eu)-tagged antibody specific for the kinase or expression tag (e.g. GST) of interest. Displacement of the tracer by a compound of the disclosure reduces, and is directly proportional to, the TR-FRET between the tracer and the antibody. Tracer was used at a concentration equal to or near to its K_d for the kinase. The Eu-tagged antibody was typically used in excess to ensure sampling of all competent protein capable of binding the tracer.

[0517] For these assays a mutant ALK2 was used that was N-terminally tagged with GST (ALK2 R206H Carma Bioscience (09-148) or ALK2 R206H ThermoFisher (PV6232)); a Eu-tagged anti-GST antibody (ThermoFisher) and Kinase Tracer 178 (ThermoFisher). In all cases, the kinase (2-5 nM) was mixed with Eu-tagged antibody (10 nM) mix and tracer (50 nM) were incubated with test compound titrations prepared in 100% DMSO (1% DMSO final) for 30 minutes at room temperature. All reagents and compounds were dissolved in Kinase Buffer A (ThermoFisher) to achieve the final concentration. The plates were read on a PerkinElmer EnVision® multilabel plate reader or a BioTek Synergy Neo plate reader, and the assay signal was represented as a ratio of the TR-FRET emission (λ_{ex} 330 nm, λ_{em} 662 nm and λ_{em} 665 nm). This readout was normalized to 0% and

100% inhibited control wells, plotted against inhibitor concentration, and fitted to a 4-parameter log dose response curve.

- 5 **[0518]** The results of this assay are shown in Table 2 in the column labelled "Binding Assay" wherein "A" represents an IC₅₀ of less than or equal to 10 nM; "B" represents an IC₅₀ of greater than 10 nM and less than or equal to 50 nM; and "C" represents an IC₅₀ of greater than 50 nM. Blank values in the table indicate that the particular compound was not tested in this assay.

Example 27. Cell-based ALK2-R206H Cell Activity Assay

10 **A. Cell line HEK293-ALK2-R206H**

- 15 **[0519]** A HEK293 (ATCC, Cat# CRL1573) based stable cell line expressing human ALK2 R206H cDNA (synthesized by GeneScript, Piscataway, NJ) and a FLAG tag at the C-terminus was generated by lentivirus transduction and subsequent blasticidin (Life Technologies, Cat#-A1113902) selection at 10 µg/ml for >2 wks. This cell line was named HEK293-ALK2-R206H.

B. Measurement of Smad1 -Ser463/Ser465 Phosphorylation by AlphaLISA

- 20 **[0520]** HEK293-ALK2-R206H cells were grown, harvested and then resuspended in serum-free, phenol red-free DMEM high glucose media (Life Technologies, Cat#-31053). The media also contained 50 units/ml penicillin and 50 µg/ml streptomycin (Life Technologies, Cat#-15070-063). HEK293-ALK2-R206H cells were then plated in white opaque 384-wells microplates (2 × 10⁴/well) (OptiPlate-384, PerkinElmer, Waltham, MA, Cat#6007299) overnight (>16h) at 37° C, 5% CO₂ for use in the assay.

- 25 **[0521]** Test compounds were first diluted to 4 mM or 0.4 mM and then serially diluted 3-fold into 10 different concentrations using DMSO. Each concentration of compound was further diluted 40-fold with phenol red-free DMEM (Life Technologies, Cat#-31053). Two µl of the diluted compounds were then dispensed into the HEK293-ALK2-R206H cell-containing wells of the microplates in duplicates. In this way, each compound was tested in 10 doses (3-fold serial dilution with the top concentration being 10 µM or 1 µM). Liquid handling was achieved using a Bravo Automated Liquid Handling Platform (Agilent Technologies). DMSO without compound was used as a negative control. The positive control was 1 µM
35 LDN193189, a known bone morphogenetic protein (BMP inhibitor).

- 40 **[0522]** After 2-3 hours of incubation with test compound or control, the cells were lysed and signal was developed using ALPHASCREEN® SUREFIRE® SMAD1 (p-Ser463/465) cellular kinase assay kit (PerkinElmer, Cat# TGRSM1S10K) following the manufacturer's recommended protocol. The microplates were read using Perkin Elmer ENVISION® plate reader (emission 520 - 620 nM). The signal reflected the level of phospho-Ser/463/465-Smad1 in the lysate. The raw data were plotted using the DMSO negative and LDN193189 positive controls as the 0% and 100%

inhibition, respectively. The 10-point dose response curve was used to calculate the IC₅₀ values.

[0523] The results of this assay are shown in Table 2 in the column labelled "Cell Assay" wherein "A" represents an IC₅₀ of less than or equal to 100 nM; "B" an IC₅₀ of greater than 100 nM and less than or equal to 500 nM; "C" an IC₅₀ of greater than 500 nM. Blank values in the table indicate that the particular compound was not tested in this assay.

[0524] In Table 2, the following designations are used:

For "Binding Assay" data: ≤ 10 nM = A; ≥ 10 -50 nM = B; > 50 nM = C; and a blank value in the table indicates that the particular compound was not tested in this assay.

For "Cell Line" data: ≤ 100 nM = A; ≥ 100 -500 nM = B; > 500 nM = C; and a blank value in the table indicates that the particular compound was not tested in this assay.

Table 2. ALK2 Inhibitory Activity of Exemplary Compounds of the Invention

Compound	Binding Assay	Cell Line
100	A	B
101	C	
102	A	B
103	B	C
104	A	B
105	B	
106	C	
107	A	A
108	B	
109	A	A
110	B	B
111	A	B
112	A	B
113	A	A
114	A	C
115	A	B
116	A	A
117	A	B
118	A	B
119	C	
120	A	B

Compound	Binding Assay	Cell Line
121	A	
122	A	A
123	A	B
124	A	A
125	A	A
126	B	
127	A	A
128	B	B
129	A	A
130	A	B
131	A	B
132	A	A
133	A	A
134	A	B
135	B	B
136	B	
137	A	A
138	B	
139	A	
140	A	A
141	B	
142	A	B
143	A	A
144	A	B
145	A	A
146	A	B
147	A	A
148	A	A
149	A	A
150	A	A
151	C	
152	A	A
153	A	B

Compound	Binding Assay	Cell Line
154	A	B
155	A	B
156	A	A
157	C	
158	A	B
159	A	A
160	A	B
161	A	A
162	A	A
163	A	A
164	A	B
165	B	
166	A	
167	A	A
168	B	
169	A	B
170	B	B
171	B	
172	A	B
173	B	C
174	A	B
175	B	
176	B	
177	A	A
178	A	A
179	A	A
180	A	A
181	A	A
182	A	A
183	B	
184	A	A
185	A	A
186	A	B

Compound	Binding Assay	Cell Line
187	A	A
188	A	A
189	A	A
190	A	
191	A	A
192	A	A
193	A	A
194	B	B
195	A	A
196	A	B
197	B	B
198	A	
199	A	A
200	A	A
201	A	A
202	A	A
203	A	A
204	A	A
205	B	B
206	A	
207	A	A
208	A	A
209	A	A
210	A	A
211	A	A
212	A	B
213	A	A
214	B	B
215	B	
216	A	A
217	A	B
218	B	
219	A	A

Compound	Binding Assay	Cell Line
220	A	
221	A	B
222	A	A
223	B	B
224	A	A
225	A	A
226	A	A
227	A	A
228	A	A
229	A	A
230	A	A
231	A	A
232	A	B
233	B	B
234	A	A
235	A	B
236	A	B
237	A	A
238	A	B
239	A	
240	A	A
241	A	A
242	A	B
243	A	A
244	A	
245	B	B
246	A	A
247	A	A
248	A	A
249	A	A
250	A	A
251	A	A
252	A	B

Compound	Binding Assay	Cell Line
253	A	B
254	A	A
255	A	B
256	A	B
257	A	A
258	A	A
259	B	C
260	A	B
261	A	A
262	A	B
263	A	B
264	A	B
265	A	A
266	A	A
267	A	A
268	B	B
269	A	A
270	A	A
271	A	B
272	B	B
273	B	
274	A	B
275	A	A
276	A	A
277	A	A
278	A	A
279	B	B
280	B	B
281	A	A
282	A	A
283	A	B
284	A	A
285	A	A

Compound	Binding Assay	Cell Line
286	A	B
287	A	A
288	A	A
289	A	A
290	A	A
291	B	
292	A	A
293	A	A
294	B	
295	A	A
296	A	B
297	A	A
298	A	A
299	A	A
300	A	A
301	A	B
302	A	A
303	A	A
304	A	A
305	A	A
306	B	
307	A	B
308	A	A
309	A	B
310	A	B
311	C	
312	A	A
313	A	A
314	B	C
315	B	B
316	A	B
317	A	B
318	A	B

Compound	Binding Assay	Cell Line
319	C	
320	A	A
321	A	A
322	A	A
323	A	A
324	A	A
325	A	A
326	A	A
327	B	B
328	A	B
329	A	B
330	A	B
331	B	
332	A	A
333	A	A
334	B	B
335	C	C
336	A	A
337	A	A
338	A	B
339	A	A
340	A	A
341	A	A
342	B	
343	A	B
344	B	B
345	A	A
348	A	A
349	A	A
350	A	A
351	A	A
352	A	B
353	A	B

Compound	Binding Assay	Cell Line
354	A	A
355	B	C
356	B	
357	A	A
359	A	C
360	A	B
361	A	B
362	B	
363	A	B
364	A	A
365	A	A
366	A	A
367	A	A
368	A	A
369	A	A
370	A	A
371	A	A
372	A	
373	B	C
374	B	
375	A	A
376	A	A
377	A	B
378	A	
379	A	B
380	A	A
381	A	A
382	A	A
383	A	A
384	A	A
385	A	A
386	A	B
387	A	A

Compound	Binding Assay	Cell Line
388	A	A
390	B	B
391	A	A
392	B	
393	A	B
397	B	
399	A	A
400	A	A
401	A	B
402	A	B
403	C	
404	A	B
405	A	
406	A	A
407	A	A
408	A	A
409	A	A
410	C	
411	A	B
412	A	A
413	A	A
414	B	
415	A	A
417	B	
418	A	A
419	A	A
420	A	A
421	A	A
422	A	A
423	A	B
425	C	
426	A	A
427	A	B

Compound	Binding Assay	Cell Line
428	A	B
429	A	A
430	A	A
431	A	B
432	A	A
433	C	
434	A	A
435	A	A
436	A	A
437	A	A
438	B	
439	B	
440	C	
441	A	A
442	A	A
443	B	
444	B	
445	A	A
446	A	
447	B	C
448	A	B
449	A	
450	A	B
451	A	A
452	A	A
453	A	A
454	A	B
455	A	A
456	A	B
457	A	B
458	A	
459	A	A
460	A	A

Compound	Binding Assay	Cell Line
461	A	
462	A	B
463	A	A
464	A	A
465	A	B
466	A	A
467	A	B
468	A	A
469	A	A
470	A	A
471	C	
472	A	
473	A	A
474	A	B
475	A	A
476	A	A
477	A	A
478	A	A
479	A	A
480	A	A
481	A	A
482	A	A
483	A	A
484	A	A
485	A	A
486	A	A
487	A	A
488	A	A
489	A	A
490	A	A
491	A	A
492	A	A
493	A	A

Compound	Binding Assay	Cell Line
494	A	A
495	A	A
496	A	A
497	A	A
498	A	A
499	A	B
500	A	B
501	A	B
502	A	A
503	A	B
504	A	B
505	A	A
506	B	
507	A	A
508	A	B
509	A	A
510	A	A
511	A	A
512	A	A
513	A	A
514	A	A
515	A	A
516	A	A
517	A	A
518	A	A
519	A	A
520	A	A
521	A	A
522	A	A
523	A	A
524	A	A
525	A	
526	A	A

Compound	Binding Assay	Cell Line
527	A	A
528	B	
529	A	
530	A	A
531	A	A
532	A	A
533	A	A
534	A	A
535	A	
536	A	
537	A	A
538	A	A
539	A	A
540	A	A
541	A	A
542	A	A
543	A	A
544	A	A
545	A	A
546	A	A
547	A	
548	A	C
549	A	A
550	A	A
551	A	A
552	A	
553	B	
554	A	B
555	A	
556	A	A
557	A	A
558	A	A
559	A	A

Compound	Binding Assay	Cell Line
560	A	A
561	A	A
562	A	A
563	A	A
564	A	A
565	A	A
566	A	A
567	A	A
568	A	A
569	A	
570	A	A
571	A	
572	A	A
573	A	A
574	A	
575	A	B
576	A	A
577	A	A
578	A	A
579	A	A
580	A	A
581	A	A
582	A	A
583	A	A
584	A	
585	A	A
586	A	B
587	A	A
588	A	A
589	A	A
590	A	
591	A	A
592	A	A

Compound	Binding Assay	Cell Line
593	A	A
594	A	A
595	A	A
596	A	B
597	A	A
598	A	A
599	A	A
600	A	A
601	A	A
602	A	
603	A	A
604	A	A
605	A	A
606	A	A
607	A	
608	A	B
609	A	A
610	A	A
611	A	A
612	A	
613	A	A
614	A	A
615	A	A
616	A	A
617	A	B
618	A	A
619	A	A
620	A	A
621	A	B
622	A	C
623	A	B
624	A	A
625	A	A

Compound	Binding Assay	Cell Line
626	A	A
627	A	A
628	A	A
629	A	A
630	A	A
631	A	A
632	A	B
633	A	A
634	A	A
635	A	A
636	A	A
637	A	A
638	A	A
639	A	
640	A	
641	A	A
642	A	A
643	A	A
644	A	A
645	A	A
646	A	A
647	A	A
648	A	A
649	A	A
650	A	A
651	A	A
652	A	A
653	A	A
654	A	A
655	A	A
656	A	A
657	A	A
658	A	A

Compound	Binding Assay	Cell Line
659	A	A
660	A	A
661	A	B
662	A	A
663	A	A
664	A	B
665	A	A
666	A	A
667	A	A
668	A	B
669	A	A
670	A	
671	A	A
672	A	A
673	A	A
674	A	A
675	A	A
676	A	A
677	A	A
678	A	A
679	A	A
680	A	A
681	A	A
682	A	A
683	A	A
684	A	A
685	A	A
686	A	A
687	A	A
688	A	A
689	A	B
690	A	A
691	A	A

Compound	Binding Assay	Cell Line
692	A	A
693	A	A
694	A	A
695	A	A
696	A	A
697	A	A
698	A	A
699	A	A
700	A	
701	A	A
702	A	A
703	A	A
704	A	A
705	A	A
706	A	A
707	A	
708	A	A
709	A	B
710	A	A
711	A	A
712	A	A
713	A	A
714	A	A
715	A	A
716	A	A
717	A	
718	A	A
719	A	A
720	A	A
721	A	A
722	A	A
723	A	A
724	A	A

Compound	Binding Assay	Cell Line
725	A	A
726	A	B
727	A	A
728	A	A
729	A	A
730	A	
731	A	A
732	A	B
733	A	A
734	A	A
735	A	A
736	A	A
737	A	A
738	A	A
739	A	A
740	A	A
741	A	A
742	A	
743	A	A
744	A	A
745	A	A
746	A	A
747	A	A
748	A	A
749	A	A
750	A	A
751	A	
752	A	B
753	A	A
754	A	A
755	A	B
756	A	
757	A	

Compound	Binding Assay	Cell Line
758	A	
759	A	
760	A	B
761	A	A
762	A	B
763	A	A
764	A	A
765	A	
766	A	B
767	A	
768	A	A
769	A	
770	A	A
771	A	A
772	A	
773	A	B
774	A	A
775	A	A
776	A	A
777	A	A
778	A	A
779	A	A
780	A	A
781	B	
782	A	A
783	A	A
784	A	A
785	A	B
786	A	A
787	A	A
788	A	A
789	A	C
790	A	C

Compound	Binding Assay	Cell Line
791	A	A
792	A	A
793	A	A
794	A	A
795	A	A
796	A	A
797	A	A
798	A	A
799	A	A
800	A	A
801	A	A
802	A	A
803	A	
804	A	A
805	A	A
806	A	
807	A	
808	A	A
809	A	A
810	A	A
811	A	A
812	A	B
813	A	A
814	A	A
815	A	A
816	A	A
817	A	A
818	A	A
819	A	A
820	A	A
821	A	A
822	B	
823	A	A

Compound	Binding Assay	Cell Line
824	A	A
825	A	B
826	A	A
827	A	B
828	A	B
829	A	A
830	A	B
831	A	A
832	A	A
833	A	A
834	A	A
835	A	A
836	A	B
837	A	B
838	A	B
839	A	A
840	A	B
841	B	
842	B	
843	A	A
844	A	A
845	A	A
846	A	A
847	A	C
848	A	A
849	A	A
850	A	A
851	A	B
852	A	A
853	A	B
854	A	A
855	A	A
856	A	A


Compound	Binding Assay	Cell Line
857	A	A
858	A	A
859	A	A
860	A	A
861	A	A
862	A	
863	A	A
864	A	A
865	A	A
866	A	
867	A	A
868	A	A
869	A	A
870	A	A
871	A	A
872	A	A
873	A	A
874	A	B
875	A	A
876	A	A
877	A	A
878	A	A
879	A	A
880	A	A
881	A	
882	A	A
883	A	A
884	A	A
885	A	A
886	A	A
887	A	A
888	A	A
889	A	A

Compound	Binding Assay	Cell Line
890	A	A
891	A	A
892	A	A
893	A	A
894	A	A
895	A	B
896	A	A
897	A	A
898	A	A
899	A	A

Equivalents

[0525] In the claims articles such as "a," "an," and "the" may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The disclosure includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The disclosure includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[0526] Furthermore, the disclosure encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims are introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, e.g., in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the disclosure, or aspects of the disclosure, is/are referred to as comprising particular elements and/or features, certain embodiments of the disclosure or aspects of the disclosure consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth *in haec verba* herein. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the disclosure, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

Patentkrav**1. Forbindelse med formelen (I):**

(I),

eller et farmaceutisk acceptabelt salt deraf, hvor:

ring A er phenyl eller heteroaryl, hvor ring A eventuelt er substitueret med 1, 2 eller 3 substituent, der uafhængigt af hinanden er valgt blandt halogen, =O, cyano, $-\text{OR}^c$, $-\text{NR}^d\text{R}^e$, $-\text{S}(\text{O})_k\text{R}^c$, $-\text{NR}^c\text{S}(\text{O})_2\text{R}^c$, $-\text{S}(\text{O})_2\text{NR}^d\text{R}^e$, $-\text{C}(=\text{O})\text{OR}^c$, $-\text{OC}(=\text{O})\text{OR}^c$, $-\text{OC}(=\text{O})\text{R}^c$, $-\text{OC}(=\text{S})\text{OR}^c$, $-\text{C}(=\text{S})\text{OR}^c$, $-\text{O}(\text{C}=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{C}(=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{S})\text{R}^c$, $-\text{NR}^c(\text{C}=\text{O})\text{OR}^c$, $-\text{O}(\text{C}=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{S})\text{OR}^c$, $-\text{O}(\text{C}=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{S})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{R}^c$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-halogenalkyl}$, $\text{C}_1\text{-C}_6\text{-heteroalkyl}$, carbocyclyl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-carbocyclyl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-carbocyclyl}$, heterocyclyl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-heterocyclyl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-heterocyclyl}$, aryl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-aryl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-aryl}$, heteroaryl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-heteroaryl}$ og $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-heteroaryl}$, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclen, heterocyclen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c , $-\text{NOa}$, cyano, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{NR}^d\text{R}^e$, $-\text{S}(\text{O})_k\text{R}^c$, $-\text{C}(=\text{O})\text{OR}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{O})\text{R}^c$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-halogenalkyl}$ eller $\text{C}_1\text{-C}_6\text{-heteroalkyl}$ ud over R^2 ;

R^1 er valgt blandt $\text{NH}(\text{C}_1\text{-C}_6\text{-alkyl})$, $\text{N}(\text{C}_1\text{-C}_6\text{-alkyl})_2$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $-\text{O}-\text{C}_1\text{-C}_6\text{-alkyl}$, $-\text{C}(\text{O})-\text{C}_1\text{-C}_4\text{-alkyl}$, carbocyclyl, heterocyclyl, $-\text{O}-(\text{C}_0\text{-C}_4\text{-alkylen})\text{-carbocyclyl}$, $-\text{O}-(\text{C}_0\text{-C}_4\text{-alkylen})\text{-heterocyclyl}$, $-\text{NH}-(\text{C}_0\text{-C}_4\text{-alkylen})\text{-carbocyclyl}$, $-\text{NH-aryl}$, $-\text{NH-O}-(\text{C}_1\text{-C}_4\text{-alkyl})$, $-\text{S-heterocyclyl}$, $-\text{S}-(\text{C}_0\text{-C}_3\text{-alkylen})\text{-(O-holdig heterocyclyl)}$ og $-\text{NH}-(\text{C}_0\text{-C}_4\text{-alkylen})\text{-heterocyclyl}$, hvor hver alkyl-, alkylen-, carbocyclyl-, aryl- og heterocyclyl del af R^1 eventuelt er substitueret med 1, 2, 3 eller 4 substituent, der uafhængigt af hinanden er valgt blandt deuterium, halogen, =O, cyano, $-\text{OR}^c$, -

NR^dR^e, -S(O)_kR^c, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -OC(=O)OR^c, -OC(=O)R^c, -OC(=S)OR^c, -C(=S)OR^c, -O(C=S)R^c, -C(=O)NR^dR^e, -NR^cC(=O)R^c, -C(=S)NR^dR^e, -NR^cC(=S)R^c, -NR^c(C=O)OR^c, -O(C=O)NR^dR^e, -NR^c(C=S)OR^c, -O(C=S)NR^dR^e, -NR^c(C=O)NR^dR^e, -NR^c(C=S)NR^dR^e, -C(=S)R^c, -C(=O)R^c, C₁-C₆-alkyl, C₁-C₆-halogenalkyl,

C₁-C₆-heteroalkyl, carbocyclyl, (C₁-C₆-alkylen)-carbocyclyl, (C₁-C₆-heteroalkylen)-carbocyclyl, heterocyclyl, (C₁-C₆-alkylen)-heterocyclyl, (C₁-C₆-heteroalkylen)-heterocyclyl, aryl, (C₁-C₆-alkylen)-aryl, (C₁-C₆-heteroalkylen)-aryl, heteroaryl, (C₁-C₆-alkylen)-heteroaryl og (C₁-C₆-heteroalkylen)-heteroaryl, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclen, heterocyclen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c, -NO₂, cyano, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆-alkyl, C₁-C₆-halogenalkyl eller C₁-C₆-heteroalkyl; eller

R¹ sammen med ét R³ danner en mættet ring, der er kondenseret til piperazinringen i formel (I), og hvor den ring, der dannes af R¹ og R³, eventuelt er substitueret med 1, 2 eller 3 substituent, der uafhængigt af hinanden er valgt blandt halogen, =O, cyano, -OR^c, -NR^dR^e, -S(O)_kR^c, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -OC(=O)OR^c, -OC(=O)R^c, -OC(=S)OR^c, -C(=S)OR^c, -O(C=S)R^c, -C(=O)NR^dR^e, -NR^cC(=O)R^c, -C(=S)NR^dR^e, -NR^cC(=S)R^c, -NR^c(C=O)OR^c, -O(C=O)NR^dR^e, -NR^c(C=S)OR^c, -O(C=S)NR^dR^e, -NR^c(C=O)NR^dR^e, -NR^c(C=S)NR^dR^e, -C(=S)R^c, -C(=O)R^c, C₁-C₆-alkyl, C₁-C₆-halogenalkyl, C₁-C₆-heteroalkyl, carbocyclyl,

(C₁-C₆-alkylen)-carbocyclyl, (C₁-C₆-heteroalkylen)-carbocyclyl, heterocyclyl, (C₁-C₆-alkylen)-heterocyclyl, (C₁-C₆-heteroalkylen)-heterocyclyl, aryl, (C₁-C₆-alkylen)-aryl, (C₁-C₆-heteroalkylen)-aryl, heteroaryl, (C₁-C₆-alkylen)-heteroaryl og (C₁-C₆-heteroalkylen)-heteroaryl, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclen, heterocyclen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c, -NO₂, cyano, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆-alkyl, C₁-C₆-halogenalkyl eller C₁-C₆-heteroalkyl;

R^2 er valgt blandt halogen, C_1 - C_6 -alkyl, heterocyclyl, cycloalkyl, $-NH-(C_0-C_4\text{-alkylen})\text{-heterocyclyl}$, $-(C_1-C_4\text{-alkylen})\text{-heterocyclyl}$, $-(C_1-C_4\text{-alkylen})\text{-NH-heterocyclyl}$ og $-O-(C_0-C_4\text{-alkylen})\text{-heterocyclyl}$, hvor en hvilken som helst heterocyclyl-, cycloalkyl-, alkyl- eller alkylendel af R^2 eventuelt er substitueret med 1, 2, 3 eller 4 substituent, der uafhængigt af hinanden er valgt blandt halogen, $=O$, cyano, $-OR^c$, $-NR^dR^e$, $-S(O)_kR^c$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$, $-OC(=O)R^c$, $-OC(=S)OR^c$, $-C(=S)OR^c$, $-O(C=S)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, $-NR^cC(=S)R^c$, $-NR^c(C=O)OR^c$, $-O(C=O)NR^dR^e$, $-NR^c(C=S)OR^c$, $-O(C=S)NR^dR^e$, $-NR^c(C=O)NR^dR^e$, $-NR^c(C=S)NR^dR^e$, $-C(=S)R^c$, $-C(=O)R^c$, C_1 - C_6 -alkyl, C_1 - C_6 -halogenalkyl, C_1 - C_6 -heteroalkyl, carbocyclyl, $(C_1-C_6\text{-alkylen})\text{-carbocyclyl}$, $(C_1-C_6\text{-heteroalkylen})\text{-carbocyclyl}$, heterocyclyl, $(C_1-C_6\text{-alkylen})\text{-heterocyclyl}$, $(C_1-C_6\text{-heteroalkylen})\text{-heterocyclyl}$, aryl, $(C_1-C_6\text{-alkylen})\text{-aryl}$, $(C_1-C_6\text{-heteroalkylen})\text{-aryl}$, heteroaryl, $(C_1-C_6\text{-alkylen})\text{-heteroaryl}$ og $(C_1-C_6\text{-heteroalkylen})\text{-heteroaryl}$, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclilen, heterocyclilen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c , $-NO_a$, cyano, $-NR^cC(=O)R^c$, $-NR^dR^e$, $-S(O)_kR^c$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$, $-C(=O)R^c$, C_1 - C_6 -alkyl, C_1 - C_6 -halogenalkyl eller C_1 - C_6 -heteroalkyl; eller R^2 sammen med et hvilket som helst ringatom i ring A danner en cycloalkyl- eller mættet heterocyclylring, der er kondenseret til ring A, og hvor den ring, der dannes af R^2 og ringatomet i ring A, eventuelt er substitueret med 1, 2 eller 3 substituent, der uafhængigt af hinanden er valgt blandt halogen, $=O$, cyano, $-OR^c$, $-NR^dR^e$, $-S(O)_kR^c$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$, $-OC(=O)R^c$, $-OC(=S)OR^c$, $-C(=S)OR^c$, $-O(C=S)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, $-NR^cC(=S)R^c$, $-NR^c(C=O)OR^c$, $-O(C=O)NR^dR^e$, $-NR^c(C=S)OR^c$, $-O(C=S)NR^dR^e$, $-NR^c(C=O)NR^dR^e$, $-NR^c(C=S)NR^dR^e$, $-C(=S)R^c$, $-C(=O)R^c$, C_1 - C_6 -alkyl, C_1 - C_6 -halogenalkyl, C_1 - C_6 -heteroalkyl, carbocyclyl, $(C_1-C_6\text{-alkylen})\text{-carbocyclyl}$, $(C_1-C_6\text{-heteroalkylen})\text{-carbocyclyl}$, heterocyclyl, $(C_1-C_6\text{-alkylen})\text{-heterocyclyl}$, $(C_1-C_6\text{-heteroalkylen})\text{-heterocyclyl}$, aryl, $(C_1-C_6\text{-alkylen})\text{-aryl}$, $(C_1-C_6\text{-heteroalkylen})\text{-aryl}$, heteroaryl, $(C_1-C_6\text{-alkylen})\text{-heteroaryl}$ og $(C_1-C_6\text{-heteroalkylen})\text{-heteroaryl}$, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclilen,

heterocyclylen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c , $-\text{NO}_2$, cyano, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{NR}^d\text{R}^e$, $-\text{S}(\text{O})_k\text{R}^c$, $-\text{C}(=\text{O})\text{OR}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{O})\text{R}^c$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-halogenalkyl}$ eller $\text{C}_1\text{-C}_6\text{-heteroalkyl}$; eller

R^2 sammen med et hvilket som helst mættet ringatom i ring A danner en cycloalkyl- eller mættet heterocyclylring, der er kondenseret, spirokondenseret eller broforbundet til ring A, og hvor den ring, der dannes af R^2 og ringatomet i ring A, eventuelt er substitueret med 1, 2 eller 3 substituenten, der uafhængigt af hinanden er valgt blandt halogen, $=\text{O}$, cyano, $-\text{OR}^c$, $-\text{NR}^d\text{R}^e$, $-\text{S}(\text{O})_k\text{R}^c$, $-\text{NR}^c\text{S}(\text{O})_2\text{R}^c$, $-\text{S}(\text{O})_2\text{NR}^d\text{R}^e$, $-\text{C}(=\text{O})\text{OR}^c$, $-\text{OC}(=\text{O})\text{OR}^c$, $-\text{OC}(=\text{O})\text{R}^c$, $-\text{OC}(=\text{S})\text{OR}^c$, $-\text{C}(=\text{S})\text{OR}^c$, $-\text{O}(\text{C}=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{C}(=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{S})\text{R}^c$, $-\text{NR}^c(\text{C}=\text{O})\text{OR}^c$, $-\text{O}(\text{C}=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{S})\text{OR}^c$, $-\text{O}(\text{C}=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{S})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{R}^c$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-halogenalkyl}$, $\text{C}_1\text{-C}_6\text{-heteroalkyl}$, carbocyclyl,

$(\text{C}_1\text{-C}_6\text{-alkylen})\text{-carbocyclyl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-carbocyclyl}$, heterocyclyl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-heterocyclyl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-heterocyclyl}$, aryl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-aryl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-aryl}$, heteroaryl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-heteroaryl}$ og $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-heteroaryl}$, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclylen, heterocyclylen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c , $-\text{NO}_2$, cyano, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{NR}^d\text{R}^e$, $-\text{S}(\text{O})_k\text{R}^c$, $-\text{C}(=\text{O})\text{OR}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{O})\text{R}^c$, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-halogenalkyl}$ eller $\text{C}_1\text{-C}_6\text{-heteroalkyl}$;

hvert R^3 uafhængigt er valgt blandt $\text{C}_1\text{-C}_4\text{-alkyl}$ og $\text{C}_1\text{-C}_4\text{-halogenalkyl}$;

hvert R^c er valgt blandt hydrogen, hydroxy, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-heteroalkyl}$, carbocyclyl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-carbocyclyl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-carbocyclyl}$, heterocyclyl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-heterocyclyl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-heterocyclyl}$, aryl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-aryl}$, $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-aryl}$, heteroaryl, $(\text{C}_1\text{-C}_6\text{-alkylen})\text{-heteroaryl}$ eller $(\text{C}_1\text{-C}_6\text{-heteroalkylen})\text{-heteroaryl}$, der hver især eventuelt er substitueret med én eller flere af halogen, hydroxy, $\text{C}_1\text{-C}_6\text{-alkyl}$, $\text{C}_1\text{-C}_6\text{-halogenalkyl}$, $\text{C}_1\text{-C}_6\text{-heteroalkyl}$, carbocyclyl, heterocyclyl, aryl eller heteroaryl;


hvert R^d og R^e uafhængigt af hinanden er valgt blandt hydrogen, C_1 - C_6 -alkyl eller C_1 - C_6 -heteroalkyl;

hvert k uafhængigt er 0, 1 eller 2; og


n er 0, 1, 2 eller 3.

2. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor n er 0; eller n er 1, og R^3 er valgt blandt methyl, ethyl og $-CHF_2$.

3. Forbindelse ifølge krav 1 eller krav 2 eller et farmaceutisk acceptabelt salt deraf, hvor ring A er valgt blandt:



og




hvor:

"1" angiver en del af ring A, der er bundet til en pyrrolo[1,2-b]pyridazindel;

"2" angiver en del af ring A, der er bundet til R^2 ; og

ring A eventuelt er substitueret med 1, 2 eller 3 substituenten, der uafhængigt af hinanden er valgt blandt halogen, $=O$, cyano, $-OR^c$, $-NR^dR^e$, $-S(O)_kR^c$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$, $-OC(=O)R^c$, $-OC(=S)OR^c$, $-C(=S)OR^c$, $-O(C=S)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, $-NR^cC(=S)R^c$, $-NR^c(C=O)OR^c$, $-O(C=O)NR^dR^e$, $-NR^c(C=S)OR^c$, $-O(C=S)NR^dR^e$, $-NR^c(C=O)NR^dR^e$, $-NR^c(C=S)NR^dR^e$, $-C(=S)R^c$, $-C(=O)R^c$, C_1 - C_6 -alkyl, C_1 - C_6 -halogenalkyl,

C₁-C₆-heteroalkyl, carbocyclyl, (C₁-C₆-alkylen)-carbocyclyl, (C₁-C₆-heteroalkylen)-carbocyclyl, heterocyclyl, (C₁-C₆-alkylen)-heterocyclyl, (C₁-C₆-heteroalkylen)-heterocyclyl, aryl, (C₁-C₆-alkylen)-aryl, (C₁-C₆-heteroalkylen)-aryl, heteroaryl, (C₁-C₆-alkylen)-heteroaryl og (C₁-C₆-heteroalkylen)-heteroaryl, hvor hver af alkylen, alkylenen, heteroalkylen, heteroalkylenen, carbocyclen, heterocyclen, arylen og heteroarylen eventuelt er substitueret med én eller flere af halogen, OR^c, -NO₂, cyano, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆-alkyl, C₁-C₆-halogenalkyl eller C₁-C₆-heteroalkyl ud over R², eventuelt hvor ring A er valgt blandt



og/eller

hvor ring A eventuelt er substitueret med 1 eller 2 substituent ud over R², hvor hver eventuelle substituent uafhængigt er valgt blandt cyano, halogen, methyl og -OCHF₂.

4. Forbindelse ifølge et hvilket som helst af kravene 1-3 eller et farmaceutisk acceptabelt salt deraf, hvor R¹ er valgt blandt -C(O)-(C₁-C₃-alkyl), C₁-C₃-alkyl, -O-(C₁-C₅-alkyl), -NH(C₁-C₅-alkyl), -N(C₁-C₄-alkyl)₂, -NH-(C₃-C₆-cycloalkyl), C₃-C₆-cycloalkyl, -O-(C₃-C₆-cycloalkyl), -O-(C₁-C₃-alkylen)-(C₃-C₆-cycloalkyl), -(C₁-C₃-alkylen)-(C₃-C₆-cycloalkyl), -O-(C₀-C₃-alkylen)-(O-holdig heterocyclyl), -NH-(C₀-C₃-alkylen)-(O-holdig heterocyclyl), en O-holdig heterocyclyl, en N-holdig heterocyclyl, -O-(C₀-C₃-alkylen)-(N-holdig heterocyclyl), -S-(C₀-C₃-alkylen)-(O-holdig heterocyclyl), -NH-O-(C₁-C₃-alkyl) og -NH-phenyl, hvor en hvilken som helst alkyl-, cycloalkyl-, phenyl- eller heterocyclyl-del af R¹ eventuelt er substitueret med 1, 2 eller 3 substituent, der uafhængigt af hinanden er valgt blandt deuterium, halogen, cyano, acetyl, C₁-C₄-alkyl, C₁-C₄-halogenalkyl, -O-C₁-C₄-alkyl, -C₁-C₄-alkylen-O-C₁-C₄-alkyl, heteroaryl, phenyl, cycloalkyl, -COOH og hydroxy, hvor hver af heteroarylen, phenylen og cycloalkylen eventuelt er substitueret med én eller flere af halogen, OR^c, -NO₂, cyano, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, C₁-C₆-alkyl, C₁-C₆-halogenalkyl eller C₁-C₆-heteroalkyl; eller R¹

sammen med et hvilket som helst ringatom i piperazindelen med formlen (I) danner en carbocycl- eller heterocyclring, der er kondenseret til piperazindelen,

og/eller

hvor R¹ er valgt blandt 1-(3-chlorphenyl)cyclopropyl, 1-(3-fluorphenyl)cyclopropyl, 1-acetylcyclopropyl, 1-cyclopropylcyclopropyl, 1-difluormethylcyclopropyl, 1-fluorcyclopropyl, 1-methylpropylaminol-pyridin-3-ylcyclopropyl, 1-thiazol-2-ylcyclopropyl, 1-thien-2-ylcyclopropyl, 1-trifluormethylcyclopropyl, 2-(4-chlorphenyl)cyclopropyl, 2,2,2-trifluorethoxy, 2,2-difluorcyclopropyl, 2,2-dimethylcyclopropyl, 2-cyanocyclopropyl, 2-cyanoethyl, 2-cyanoethylamino-2-cyclobutylcyclopropyl, 2-fluorcyclopropyl, 2-fluorethoxy, 2-hydroxyethylamino, 2-methoxyethoxy, 2-methylcyclopropyl, 2-oxa-6-azaspiro[3.3]heptan-6-yl, 3,3-difluorcyclobutyl, 3-cyanoazetidin-1-yl, 3-cyanocyclobutyl, 3-fluorcyclobutyl, 3-hydroxy-3-methylcyclobutyl, 3-hydroxy-3-trifluormethylcyclobutyl, 3-hydroxyazetidin-1-yl, 3-hydroxycyclobutyl, 3-methoxyazetidin-1-yl, 3-methoxymethylazetidin-1-yl, 3-phenyl-3-hydroxycyclobutyl, 4-cyanocyclohexyl, 4-cyanopiperidin-1-yl, 4-hydroxy-4-methylcyclohexyl, 4-hydroxycyclohexyl, 4-hydroxypiperidin-1-yl, 4-methylcyclohexyl, acetyl, azetidin-1-yl, cyclobutoxy, cyclobutyl, cyclobutylamino, cyclopentylamino, cyclopropyl, cyclopropylmethyl, diethylamino, ethoxy, ethyl, ethylamino, isobutoxy, isopropoxy, isopropyl, isopropylamino, methoxymethyl, N-ethyl-N-methylamino, pentylamino, piperidin-1-yl, propylamino, propyloxypyrrolidin-1-yl og t-butylamino, t-butoxy, 2,2-dimethylpropoxy, 2,2-difluorethoxy, N-(2,2-dimethylpropyl)amino, N-(1,2-dimethylpropyl)amino, 2,2,2-trifluorethylamino, N-(methoxymethyl)amino, oxetan-3-yloxy, oxetan-3-yl, oxetan-3-ylamino, oxetan-3-ylmethoxy, N-(oxetan-3-ylmethyl)amino, tetrahydrofuran-3-yloxy, tetrahydropyran-4-yloxy, 3-cyanocyclobutoxy, 1,3-dihydroxypropan-2-yloxy, 1-acetylazetidin-3-yloxy, 1-hydroxy-2-hydroxycarbonylethan-2-yloxy, 1-methyl-2-fluorethoxy, 2,2-difluorethylamino, 2-cyanoethan-1-yloxy, 2-fluorethylamino, 2-fluorphenylamino, 2-fluorpropoxy, 2-methyloxetan-3-yloxy, 3-cyano-oxetan-3-yloxy, 3-deutero-oxetan-3-yloxy, 6-oxa-1-azaspiro[3.3]heptan-1-yl, cyclopropoxy, ethoxyamino, oxetan-3-ylthio, perdeuteroethoxy, phenylamino, tetrahydrofuran-2-yloxy og tetrahydrofuran-3-yl, eller

R¹ sammen med et ringatom i piperazindelen danner en 6-oxohexahydropyrrolo[1,2-a]pyrazin-2-yl eller 2-ethyl-3-oxohexahydroimidazo[1,5-a]pyrazin-7-yl.

5. Forbindelse ifølge et hvilket som helst af kravene 1-4 eller et farmaceutisk acceptabelt salt deraf, hvor R^2 er valgt blandt halogen, cycloalkyl, heterocyclyl, $-O-(C_0-C_4\text{-alkylen})-$ (heterocyclyl), $-(C_1-C_3\text{-alkylen})\text{-heterocyclyl}$, $-(C_1-C_3\text{-alkylen})\text{-NH}-(C_1-C_3\text{-alkyl})$, $-(\text{hydroxy-substitueret } C_1-C_3\text{-alkylen})\text{-NH}-(C_1-C_3\text{-alkyl})$, $C_1-C_4\text{-alkyl}$, substitueret med både hydroxy og én eller flere af amino, $C_1-C_4\text{-alkylamino}$ eller $\text{di-}C_1-C_4\text{-alkylamino}$, $\text{cyanosubstitueret } C_1-C_4\text{-alkyl}$, hydroxy, $-S(O)_2-C_1-C_4\text{-alkyl}$ og $-(\text{aminosubstitueret } C_1-C_3\text{-alkylen})\text{-heterocyclyl}$, eller R^2 sammen med et ringatom i ring A danner en heterocyclyl eller en carbocyclyl, der er kondenseret til ring A, hvor en hvilken som helst heterocyclyl eller carbocyclyl eventuelt er substitueret med 1, 2 eller 3 substituent, der uafhængigt af hinanden er valgt blandt halogen, cyano, hydroxy, $-\text{NH}_2$, $-\text{NH}(C_1-C_4\text{-alkyl})$, $-\text{NH-C(O)-O}-(C_1-C_4\text{-alkyl})$, $=O$, $-C(O)-C_1-C_4\text{-alkyl}$, $-C_1-C_4\text{-alkyl}$, $\text{deutereret } C_1-C_4\text{-alkyl}$, $-C_1-C_4\text{-halogenalkyl}$, $\text{hydroxysubstitueret } -C_1-C_4\text{-alkyl}$, $-O-C_1-C_4\text{-alkyl}$, $-O-C_1-C_4\text{-halogenalkyl}$, $-(C_1-C_4\text{-alkylen})\text{-O}-(C_1-C_4\text{-alkyl})$, $-(\text{aminosubstitueret } C_1-C_4\text{-alkylen})\text{-O}-(C_1-C_4\text{-alkyl})$, $-(C_1-C_4\text{-alkylen})\text{-O}-(C_1-C_4\text{-halogenalkyl})$, $-C(O)\text{-O-}C_1-C_4\text{-alkyl}$, $-\text{COOH}$, $C_3-C_6\text{-cycloalkyl}$, heterocyclyl og $-\text{NH-heterocyclyl}$, hvor hver heterocyclyl eventuelt er substitueret med én eller flere af halogen, OR^c , $-\text{NO}_2$, cyano, $-\text{NR}^c\text{C(=O)OR}^c$, $-\text{NR}^d\text{R}^e$, $-S(O)_k\text{R}^c$, $-C(=O)\text{OR}^c$, $-C(=O)\text{NR}^d\text{R}^e$, $-C(=O)\text{R}^c$, $C_1-C_6\text{-alkyl}$, $C_1-C_6\text{-halogenalkyl}$ eller $C_1-C_6\text{-heteroalkyl}$,
og/eller

hvor R^2 er valgt blandt 1-(1-hydroxypropan-2-yl)-4-methoxypiperidin-4-yl, 1-(3-difluormethoxy)propan-2-yl-4-methoxypiperidin-4-yl, 1-(3-methoxy)propan-2-yl-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)piperidin-4-yl, 1-(propan-2-yl)piperidin-3-yl, 1-(propan-2-yl)piperidin-4-yl, 1-(pyrrolidin-1-yl)ethan-1-yl, 1,2,3,6-tetrahydropyridin-4-yl, 1,4-diazabicyclo[4.2.0]octan-4-yl, 1-acetylpiperidin-4-yl, 1-cyclobutylpiperidin-3-yl, 1-ethyl-3,3-difluorpiperidin-4-yl, 1-ethyl-3-fluorpiperidin-4-yl, 1-ethyl-3-hydroxyazetidin-3-yl, 1-ethyl-4-fluorpyrrolidin-3-yl, 1-ethylazetidin-3-yl, 1-ethylazetidin-3-yloxy, 1-ethylpiperidin-3-yl, 1-ethylpiperidin-3-yloxy, 1-ethylpiperidin-4-yl, 1-ethylpiperidin-4-yloxy, 1-ethylpyrrolidin-3-yl, 1-ethylpyrrolidin-3-ylmethoxy, 1-ethylpyrrolidin-3-yloxy, 1H-pyrrolidin-2-yl, 1-hydroxy-2-aminoprop-2-yl, 1-isopropyl-1,2,3,6-tetrahydropyridin-4-yl, 1-isopropyl-2-methylpyrrolidin-2-yl, 1-isopropyl-3,4-dimethylpiperazin-3-yl, 1-isopropyl-3-ethoxypiperidin-3-yl, 1-


isopropyl-3-fluoropiperidin-3-yl, 1-isopropyl-3-hydroxyazetidin-3-yl, 1-isopropyl-3-hydroxypiperidin-3-yl, 1-isopropyl-3-hydroxypyrrolidin-3-yl, 1-isopropyl-3-methoxyazetidin-3-yl, 1-isopropyl-3-methoxypiperidin-3-yl, 1-isopropyl-3-methoxypyrrolidin-3-yl, 1-isopropyl-3-methylpiperazin-3-yl, 1-isopropyl-4-cyanopiperidin-4-yl, 1-isopropyl-4-ethoxypiperidin-4-yl, 1-isopropyl-4-fluoropiperidin-4-yl, 1-isopropyl-4-fluorpyrrolidin-3-yl, 1-isopropyl-4-hydroxypiperidin-3-yl, 1-isopropyl-4-hydroxypiperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-isopropyl-4-methylpiperazin-3-yl, 1-isopropyl-4-methylpiperidin-4-yl, 1-isopropyl-4-trifluormethylpiperidin-4-yl, 1-isopropyl-5-methylpyrrolidin-3-yl, 1-isopropylazetidin-2-ylmethoxy, 1-isopropylazetidin-3-yl, 1-isopropylazetidin-3-ylmethoxy, 1-isopropylazetidin-3-yloxy, 1-isopropylpiperazin-3-yl, 1-isopropylpiperazin-4-yl, 1-isopropylpiperidin-2-yl, 1-isopropylpiperidin-3-yl, 1-isopropylpiperidin-4-yl, 1-isopropylpyrrolidin-2-yl, 1-isopropylpyrrolidin-3-yl, 1-methyl-1-cyanoethyl, 1-sec-butylpiperidin-4-yl, 1-t-butoxycarbonyl-4-aminopiperidin-4-yl, 2-(isopropylamino)-3-hydroxypropan-2-yl, 2-(isopropylamino)-propan-2-yl, 2,3,5,6-tetrahydroimidazo[2,1-b]thiazol-6-yl, 2,6-diazaspiro[3.3]heptan-2-yl, 2,6-diazaspiro[3.4]octan-2-yl, 2,6-diazaspiro[3.4]octan-6-yl, 2,7-diazaspiro[4.4]nonan-2-yl, 2-difluormethylpiperazin-1-yl, 2-isopropyl-2,6-diazaspiro[3.3]heptan-6-yl, 2-methyl-1H-pyrrolidin-2-yl, 2-oxa-5,8-diazaspiro[3.5]nonan-8-yl, 2-oxo-4-ethylpiperazin-1-yl, 2-oxopiperazin-1-yl, 2-trifluormethylpiperazin-1-yl, 3,3-difluoropiperidin-4-yl, 3,3-dimethyl-4-ethylpiperazin-1-yl, 3-aminopyrrolidin-1-yl, 3-fluoropiperidin-3-yl, 3-fluoropiperidin-4-yl, 3-hydroxyazetidin-3-yl, 3-hydroxyquinuclidin-3-yl, 3-methyl-4-ethylpiperazin-1-yl, 3-methylpiperazin-1-yl, 3-trifluormethylpiperazin-1-yl, 4-(1,1,2,2-pentadeuteroethyl)piperazin-1-yl, 4-(2,2-difluorethyl)piperazin-1-yl, 4-(2-hydroxyethyl)piperazin-1-yl, 4-(2-methoxyethyl)piperazin-1-yl, 4-(methoxycarbonylamino)piperidin-4-yl, 4-(oxetan-3-yl)piperazin-1-yl, 4,5-dihydro-1H-imidazol-2-yl, 4,5-dihydro-1H-imidazol-2-ylamino, 4-aminopiperidin-1-yl, 4-cyanopiperidin-4-yl, 4-ethoxypiperidin-4-yl, 4-ethylmorpholin-2-yl, 4-ethylpiperazin-1-yl, 4-ethylpiperazin-1-ylethoxy, 4-fluoropiperidin-4-yl, 4-fluorpyrrolidin-3-yl, 4-hydroxy-tetrahydro-2H-pyran-4-yl, 4-isopropylmorpholin-3-

yl, 4-isopropylpiperazin-1-yl, 4-methoxypiperidin-4-yl, 4-methylpiperazin-1-yl, 4-methylpiperidin-4-yl, 5,5-difluorpiperidin-3-yl, 5-ethyl-2,5-diazabicyclo[2.2.1]heptan-2-yl, 6-ethyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-isopropyl-2,6-diazaspiro[3.3]heptan-2-yl, 6-methylmorpholin-2-yl, azetidin-2-ylmethoxy, azetidin-3-yl, brom, cyclopentyl, hexahydropyrazino[2,1-c][1,4]oxazin-8(1H)-yl, hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl, morpholin-2-yl, morpholin-3-yl, octahydro-2H-pyrido[1,2-a]pyrazin-2-yl, piperazin-1-yl, piperazin-1-ylethoxy, piperdin-4-yl, piperidin-2-yl, piperidin-3-yl, piperidin-3-yloxy, piperidin-4-yloxy, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrrolidin-3-ylmethoxy, pyrrolidin-3-yloxy, quinuclidin-4-yl, tetrahydro-2H-pyran-4-yl, 1-(1-fluorpropan-2-yl)-4-ethoxypiperidin-4-yl, 1-(1-fluorpropan-2-yl)-4-methoxypiperidin-4-yl, 1-(2-fluorpropyl)-4-ethoxypiperidin-4-yl, 1-(2-fluorpropyl)-4-ethoxypiperidin-4-yl, 1-(3-(difluormethoxy)propan-2-yl)-4-methoxypiperidin-4-yl, 1-(oxetan-3-yl)-4-ethoxypiperidin-4-yl, 1-(tetrahydrofuran-2-yl)-1-aminomethyl, 1-amino-2-hydroxy-2-methylpropyl, 1-amino-2-methoxyethyl, 1-azabicyclo[2.2.1]heptan-4-yl, 1-cyclopropyl-4-ethoxypiperidin-4-yl, 1-diethylamino-2-hydroxyethyl, 1-ethylamino-2-hydroxyethyl, 1-isopropyl-4-difluormethoxypiperidin-3-yl, 1-isopropyl-4-difluormethoxypiperidin-4-yl, 1-isopropyl-4-hydroxymethylpiperidin-4-yl, 1-isopropyl-4-methoxycarbonylpiperidin-4-yl, 1-isopropyl-4-(methoxymethyl)piperidin-4-yl, 1-isopropyl-4-methoxypiperidin-4-yl, 1-methyl-1-isopropylamino-2-hydroxyethyl, 2,2,5,5-tetramethyl-4-hydroxypiperidin-4-yl, 2,2-dimethyl-4-methoxypiperidin-4-yl, 2-amino-1-hydroxyethyl, 2-amino-3-hydroxypropyl, 2-azaspiro[3.3]heptan-6-yl, 2-hydroxy-1-aminoethyl, 2-hydroxy-1-isopropylaminoethyl, 2-hydroxyethylaminomethyl, 3-amino-oxetan-3-yl, 3-ethoxypiperidin-3-yl, 3-methoxypiperidin-3-yl, 4-amino-tetrahydropyran-4-yl, 4-ethoxytetrahydropyran-4-yl, 4-hydroxycarbonylpiperidin-4-yl, 4-hydroxymethylpiperidin-4-yl, 4-methoxycarbonylpiperidin-4-yl, 4-methoxytetrahydropyran-4-yl, 4-trifluormethylpiperidin-4-yl, ethylsulfonyl og oxetan-3-ylaminomethyl, eller

R² sammen med ring A danner 3'H-spiro[azetidin-3,1'-isobenzofuran]-5'-yl, 6-isopropyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 4,5,6,7-

tetrahydrothiazolo[4,5-c]pyridin-2-yl, 4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 5,6,7,8-tetrahydroimidazo[1,5-a]pyrazin-3-yl, 7-methyl-5,6,7,8-tetrahydroimidazo[1,5-a]pyrazin-3-yl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridin-2-yl, 5-isopropyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridin-2-yl, 1-amino-2,3-dihydro-1H-inden-5-yl eller 1-(isopropylamino)-2,3-dihydro-1H-inden-5-yl.

6. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor forbindelsen er en forbindelse med formlen (II):



hvor:

X er C(R¹³) eller N;

R¹¹ er valgt blandt -NH-(C₃-C₄-cycloalkyl); -NH-C₁-C₃-alkyl; -O-C₃-C₄-cycloalkyl; -O-C₁-C₃-alkyl, eventuelt substitueret med én eller flere substituer, der er valgt blandt fluor, hydroxy, cyano og deuterium; og -O-(O-holdig heterocykel);

R¹² er valgt blandt:

piperidin-3-yl, eventuelt 3-substitueret med C₁-C₃-alkoxy, fluor, C₁-C₃-alkyl eller cyano; og


piperidin-4-yl, eventuelt 4-substitueret med C₁-C₃-alkoxy, fluor, C₁-C₃-alkyl, cyano, hvor R¹² endvidere eventuelt er 1-substitueret med C₁-C₅-alkyl, eventuelt substitueret med én eller flere hydroxy og/eller én eller flere -NH₂;

R¹³ er valgt blandt hydrogen, cyano og fluor; og

R¹⁴ er fluor.

7. Forbindelse ifølge krav 6 eller et farmaceutisk acceptabelt salt deraf, hvor forbindelsen er:

(A) en forbindelse med formlen (IIa):



(IIa),

eller et farmaceutisk acceptabelt salt deraf, hvor:

X er C(R¹³) eller N;

R¹¹ er valgt blandt: -NH-(C₃-C₄-cycloalkyl); -NH-C₁-C₃-alkyl; -O-C₃-C₄-cycloalkyl; -O-(C₁-C₃-alkyl), eventuelt substitueret med én eller flere substituent, der er valgt blandt fluor, hydroxy, cyano og deuterium; og -O-(O-holdig heterocykel);


R¹³ er valgt blandt hydrogen, cyano og fluor;

R¹⁴ er fluor;

R¹⁵ er valgt blandt hydrogen, C₁-C₃-alkoxy, fluor, C₁-C₃-alkyl og cyano; og

R¹⁶ er hydrogen eller C₁-C₅-alkyl, eventuelt substitueret med én eller flere hydroxy og/eller én eller flere -NH₂; eller

(B) en forbindelse med formlen (IIb):



(IIb),

eller et farmaceutisk acceptabelt salt deraf, hvor:

X er C(R¹³) eller N;

R¹¹ er valgt blandt: -NH-(C₃-C₄-cycloalkyl); -NH-C₁-C₃-alkyl; -O-C₃-C₄-cycloalkyl; -O-(C₁-C₃-alkyl), eventuelt substitueret med én eller flere substituent, der er valgt blandt fluor, hydroxy, cyano og deuterium; og -O-(O-holdig heterocykel);


R¹³ er valgt blandt hydrogen, cyano og fluor;

R¹⁴ er fluor;

R^{15} er valgt blandt hydrogen, C_1 - C_3 -alkoxy, fluor, C_1 - C_3 -alkyl og cyano; og

R^{16} er hydrogen eller C_1 - C_5 -alkyl, eventuelt substitueret med én eller flere hydroxy og/eller én eller flere $-NH_2$,


eventuelt hvor forbindelsen med formlen IIb enten er en forbindelse med formlen (IIb-1):



(IIb-1),

eller et farmaceutisk acceptabelt salt deraf,

eller en forbindelse med formlen (IIb-2):



(IIb-2),

eller et farmaceutisk acceptabelt salt deraf.


8. Forbindelse ifølge krav 6 eller krav 7 eller et farmaceutisk acceptabelt salt deraf, hvor:







R^{14} er fraværende; eller







R^{13} er hydrogen; eller







R^{11} er valgt blandt $-NH-C_1-C_3$ -alkyl; $-O-C_1-C_3$ -alkyl, eventuelt substitueret med én eller flere substituenten, der er valgt blandt fluor, hydroxy, cyano og deuterium; oxetan-3-yloxy og tetrahydrofuran-3-yloxy, eventuelt hvor R^{11} er valgt blandt $-OCH_2CH_3$, $-NHCH(CH_3)_2$, oxetan-3-yloxy og tetrahydrofuran-3-yloxy.







9. Forbindelse ifølge krav 1, hvor forbindelsen er valgt blandt en hvilken som helst af følgende forbindelser:







#	Struktur
100	 The chemical structure of compound 100 consists of a central pyridine ring. At the 2-position of the pyridine ring, there is a piperazine ring. The nitrogen at the 1-position of the piperazine ring is substituted with a cyclopropyl carbonyl group (a cyclopropyl ring attached to a carbonyl group, which is in turn attached to the nitrogen). The nitrogen at the 4-position of the piperazine ring is substituted with a 2,6-diazapyridine ring system. The 2-position of this diazapyridine ring is further substituted with a 1,4-dihydropyridine ring, which has a nitrogen atom at the 4-position.







#	Struktur
101	
102	
103	
104	
105	
106	






#	Struktur
107	
108	
109	
110	
111	
112	

#	Struktur
113	
114	
115	
116	
117	
118	







#	Struktur
119	
120	
121	
122	
123	
124	







#	Struktur
125	
126	
127	
128	
129	
130	






#	Struktur
131	
132	
133	
134	
135	
136	







#	Struktur
137	
138	
139	
140	
141	






#	Struktur
142	 <chem>CCNC(=O)N1CCN(C1)c2nc3cc(ccn3n2)-c4ccc(cc4)[C@H]5CCNCC5</chem>
143	 <chem>CCOC(=O)N1CCN(C1)c2nc3cc(ccn3n2)-c4ccc(cc4)N5CCNCC5</chem>
144	 <chem>CCOC(=O)N1CCN(C1)c2nc3cc(ccn3n2)-c4ccc(cc4)[C@H]5CCNCC5</chem>
145	 <chem>CCOC(=O)N1CCN(C1)c2nc3cc(ccn3n2)-c4ccc(cc4)[C@@H]5CNC(F)C5</chem>
146	 <chem>C1CC1C(=O)N1CCN(C1)c2nc3cc(ccn3n2)-c4ccc(cc4)N5CC6CCNCC6C5</chem>






#	Struktur
147	
148	
149	
150	
151	
152	






#	Struktur
153	
154	
155	
156	
157	
158	

#	Struktur
159	
160	
161	
162	
163	






#	Struktur
164	
165	
166	
167	
168	
169	

#	Struktur
170	
171	
172	
173	
174	






#	Struktur
175	
176	
177	
178	
179	






#	Struktur
180	
181	
182	
183	
184	






#	Struktur
185	
186	
187	
188	
189	






#	Struktur
190	
191	
192	
193	
194	






#	Struktur
195	 <chem>CN1CCN(CC1C(=O)C2CC2C#N)C3=CC=C4C(=CC=C4C5=CC=CC=C5N6C=CC=CC=C6N)N7CCOCC7</chem>
196	 <chem>CN1CCN(CC1C(=O)C2CC2)C3=CC=C4C(=CC=C4C5=CC=CC=C5N6C=CC=CC=C6N)N7CCN8CCNCC87</chem>
197	 <chem>CN1CCN(CC1C(=O)C2CC2)C3=CC=C4C(=CC=C4C5=CC=CC=C5N6C=CC=CC=C6N)N7CCN8CCNCC87</chem>
198	 <chem>CN1CCN(CC1C(=O)C2CC2)C3=CC=C4C(=CC=C4C5=CC=CC=C5N6C=CC=CC=C6N)N7CCN8CCNCC87</chem>
199	 <chem>CN1CCN(CC1C(=O)CC#N)C3=CC=C4C(=CC=C4C5=CC=CC=C5N6C=CC=CC=C6N)N7CCOCC7</chem>
200	 <chem>CN1CCN(CC1C(=O)C2CC2)C3=CC=C4C(=CC=C4C5=CC=CC=C5N6C=CC=CC=C6N)N7CCNCC7</chem>






#	Struktur
201	
202	
203	
204	
205	






#	Struktur
206	
207	
208	
209	
210	






#	Struktur
211	
212	
213	
214	
215	







#	Struktur
216	
217	
218	
219	
220	






#	Struktur
221	
222	
223	
224	
225	






#	Struktur
226	 <chem>CCN1CC[C@H](C1)c2ccc(cc2)-c3cc4c(cc3)nn(C4)C5CN(CCN5)C(=O)C6CC6F</chem>
227	 <chem>CCN1CC[C@H](C1)c2ccc(cc2)-c3cc4c(cc3)nn(C4)C5CN(CCN5)C(=O)C6CC6F</chem>
228	 <chem>CCN1CC[C@H](C1)c2ccc(cc2)-c3cc4c(cc3)nn(C4)C5CN(CCN5)C(=O)C6CC6</chem>
229	 <chem>CCN1CC[C@H](C1)c2ccc(cc2)-c3cc4c(cc3)nn(C4)C5CN(CCN5)C(=O)C6CC6</chem>
230	 <chem>CCN1CC[C@H](C1)c2ccc(cc2)-c3cc4c(cc3)nn(C4)C5CN(CCN5)C(=O)OCC</chem>







#	Struktur
231	
232	
233	
234	
235	






#	Struktur
236	
237	
238	
239	
240	






#	Struktur
241	
242	
243	
244	
245	
246	






#	Struktur
247	
248	
249	
250	
251	






#	Struktur
252	
253	
254	
255	
256	






#	Struktur
257	
258	
259	
260	
261	
262	






#	Struktur
263	
264	
265	
266	
267	






#	Struktur
268	
269	
270	
271	
272	






#	Struktur
273	
274	
275	
276	
277	






#	Struktur
278	
279	
280	
281	
282	






#	Struktur
283	
284	
285	
286	
287	






#	Struktur
288	 <chem>CCOC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccc(cc4)[C@H]5CCN(C)CC5</chem>
289	 <chem>CCNC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccc(cc4)N5CCN(C)CC5</chem>
290	 <chem>CCNC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccncc4[C@H]5CCN(C)CC5</chem>
291	 <chem>CC1CCN(C1)c2cnc3cc(ccn32)c4ccc(cc4)N5CCN(CC)CC5C(=O)[C@H]6CC6F</chem>
292	 <chem>CC1CCN(C1)c2cnc3cc(ccn32)c4ccc(cc4)N5CCN(CC)CC5C(=O)[C@H]6CC6F</chem>






#	Struktur
293	
294	
295	
296	
297	






#	Struktur
298	
299	
300	
301	
302	





#	Struktur
303	
304	
305	
306	
307	






#	Struktur
308	
309	
310	
311	
312	






#	Struktur
313	 <chem>CC1CCN(C1)Cc2ccc(cc2)-c3cc4c(cc3n5ccccc45)nc6ccccc6N7CCCN(C7)C(=O)C8CC8</chem>
314	 <chem>C1CCN(C1)Cc2ccc(cc2)-c3cc4c(cc3n5ccccc45)nc6ccccc6N7CCCN(C7)C(=O)C8CC8</chem>
315	 <chem>C1CCN(C1)Cc2ccc(cc2)-c3cc4c(cc3n5ccccc45)nc6ccccc6N7CCCN(C7)C(=O)C8CC8</chem>
316	 <chem>CC1CCN(C1)Cc2ccc(cc2)-c3cc4c(cc3n5ccccc45)nc6ccccc6N7CCCN(C7)C(=O)C8CC8</chem>
317	 <chem>CC1CCN(C1)Cc2ccc(cc2)-c3cc4c(cc3n5ccccc45)nc6ccccc6N7CCCN(C7)C(=O)C8CC8</chem>






#	Struktur
318	
319	
320	
321	
322	






#	Struktur
323	
324	
325	
326	
327	

#	Struktur
328	 <chem>CCN(CC)C(=O)N1CCN(CC1)c2nc3cc(ccc3n2)-c4ccc(cc4)N5CCN(CC5)C6C(C)C</chem>
329	 <chem>CCN(CC)C(=O)N1CCN(CC1)c2nc3cc(ccc3n2)-c4ccc(cc4)N5CCN(CC5)C6C(C)C</chem>
330	 <chem>CCN(CC)C(=O)N1CCN(CC1)c2nc3cc(ccc3n2)-c4ccc(cc4)N5CCN(CC5)C6C(C)C</chem>
331	 <chem>CCN(CC)C(=O)N1CCN(CC1)c2nc3cc(ccc3n2)-c4ccc(cc4)N5CCN(CC5)C6C(C)C</chem>






#	Struktur
332	
333	
334	
335	
336	






#	Struktur
337	
338	
339	
340	
341	






#	Struktur
342	
343	
344	
345	
346	






#	Struktur
347	
348	
349	
350	
351	






#	Struktur
357	
358	
359	
360	
361	






#	Struktur
362	
363	
364	
365	
366	






#	Struktur
367	
368	
369	
370	
371	

#	Struktur
372	
373	
374	
375	
376	

#	Struktur
377	
378	
379	
380	
381	

#	Struktur
382	
383	
384	
385	
386	






#	Struktur
387	
388	
389	
390	
391	

#	Struktur
397	
398	
399	
400	
401	






#	Struktur
402	<chem>CCN1CCN(CC1)c2ccc(cc2)c3cc4c(cc3)ncn4C(=O)C5(CCC5)F</chem>
403	<chem>CCN1CCN(CC1)c2ccc(cc2)c3cc4c(cc3)ncn4C(=O)C5(CCC5)F</chem>
404	<chem>CCN1CCN(CC1)c2ccc(cc2)c3cc4c(cc3)ncn4C(=O)C5(CCC5)F</chem>
405	<chem>CCN1CCN(CC1)c2ccc(cc2)c3cc4c(cc3)ncn4C(=O)C5(CCC5)F</chem>
406	<chem>CCN1CCN(CC1)c2ccc(cc2)c3cc4c(cc3)ncn4C(=O)C5(CCC5)F</chem>






#	Struktur
407	<chem>CC1(C)CCN(C1)C2=CC=C(C=C2)C3=CC=CC=C3C4=CC=CC=C4C5=CN=C6C(=C5)N(C6)C(=O)N7C8CC8C#N7</chem>
408	<chem>CC1(C)CCN(C1)C2=CC=C(C=C2)C3=CC=CC=C3C4=CC=CC=C4C5=CN=C6C(=C5)N(C6)C(=O)N7C8CCC8C#N7</chem>
409	<chem>CC1(C)CCN(C1)C2=CC=C(C=C2)C3=CC=CC=C3C4=CC=CC=C4C5=CN=C6C(=C5)N(C6)C(=O)N7C8CCC8C#N7</chem>
410	<chem>CCN1CCN(C1)C2=CC=C(C=C2)C3=CC=CC=C3C4=CC=CC=C4C5=CN=C6C(=C5)N(C6)C(=O)N7C8CC8C(F)(F)F7</chem>
411	<chem>CC1(C)CCN(C1)C2=CC=C(C=C2)C3=CC=CC=C3C4=CC=CC=C4C5=CN=C6C(=C5)N(C6)C(=O)N7C8CCN8C#N7</chem>






#	Struktur
412	 <chem>CCN1CCc2ccc(cc2-c3cc4nc5ccccc5n4n3)N6CCN(C(=O)N7CC[C@H](C#N)C7)CC6</chem>
413	 <chem>CC(C)N1CCc2ccc(cc2-c3cc4nc5ccccc5n4n3)N6CCN(C(=O)N7CC[C@H](C#N)C7)CC6</chem>
414	 <chem>CC(C)N1CCc2ccc(cc2-c3cc4nc5ccccc5n4n3)N6CCN(C(=O)N7CC8CC8N7)CC6</chem>
415	 <chem>CCN1CCc2ccc(cc2-c3cc4nc5ccccc5n4n3)N6CCN(C(=O)N7CCCCC7)CC6</chem>
416	 <chem>CC(C)N1CCc2ccc(cc2-c3cc4nc5ccccc5n4n3)N6CCN(C(=O)N7CCCC7)CC6</chem>






#	Struktur
417	
418	
419	
420	
421	






#	Struktur
422	 <chem>CCN1CCN(CC1)c2ccc(cc2)-c3cc4c(cc3)nn5ccc(cc45)C(=O)[C@H]6CCCC(O)C6</chem>
423	 <chem>CCN1CCN(CC1)c2ccc(cc2)-c3cc4c(cc3)nn5ccc(cc45)C(=O)N6CCCC(O)C6</chem>
424	 <chem>CCCCCNC(=O)N1CCN(CC1)c2cc3c(cc2)nn4ccc(cc34)C5=CC=C(C=C5)C6CCCC(C)N6</chem>
425	 <chem>CCN1CCN(CC1)c2ccc(cc2)-c3cc4c(cc3)nn5ccc(cc45)C(=O)N6CCOC7CCOC7N6</chem>
426	 <chem>CC1CN2CCN2C1c3ccc(cc3)-c4cc5c(cc4)nn6ccc(cc56)C7NCCN7C(=O)[C@H]8CC(C#N)C8</chem>






#	Struktur
427	
428	
429	
430	
431	







#	Struktur
432	
433	
434	
435	
436	






#	Struktur
437	
438	
439	
440	
441	






#	Struktur
442	
443	
444	
445	
446	






#	Struktur
447	
448	
449	
450	
451	






#	Struktur
452	
453	
454	
455	
456	

#	Struktur
457	
458	
459	
460	
461	
462	






#	Struktur
463	
464	
465	
466	
467	






#	Struktur
468	
469	
470	
471	
472	






#	Struktur
473	
474	
475	
476	
477	







#	Struktur
478	
479	
480	
481	
482	







#	Struktur
483	 <chem>CCOC(=O)N1CCN(CC1)N2C=CN3C=CC=C(C3=N2)C4=CC=C(C=C4)C5(C#N)C6CCNCC6</chem>
484	 <chem>CC1(C)CCN(C1)C2=CC=CC=C2N3C=CC=C(C3=N)C4=CC=CC=C4N5C=CC=C(C5=N)C6=CC=CC=C6N7CCN(CC7)C(=O)O</chem>
485	 <chem>CC1(C)CCN(C1)C2=CC=CC=C2N3C=CC=C(C3=N)C4=CC=CC=C4N5C=CC=C(C5=N)C6=CC=CC=C6N7CCN(CC7)C(=O)O</chem>
486	 <chem>CC1(C)CCN(C1)C2=CC=CC=C2N3C=CC=C(C3=N)C4=CC=CC=C4N5C=CC=C(C5=N)C6=CC=CC=C6N7CCN(CC7)C(=O)O</chem>
487	 <chem>CC1(C)CCN(C1)C2=CC=CC=C2N3C=CC=C(C3=N)C4=CC=CC=C4N5C=CC=C(C5=N)C6=CC=CC=C6N7CCN(CC7)C(=O)O</chem>






#	Struktur
488	 <chem>CO[C@H]1CCNCC1c2ccc(cc2-c3ccc4nc5cccnc5n4)C(=O)N6CCNCC6C7CC7</chem>
489	 <chem>CO[C@@H]1CCNCC1c2ccc(cc2-c3ccc4nc5cccnc5n4)C(=O)N6CCNCC6C7CC7</chem>
490	 <chem>CO[C@H]1CCNCC1c2ccc(cc2-c3ccc4nc5cccnc5n4)C(=O)N6CCNCC6C7CC7</chem>
491	 <chem>CO[C@H]1CCNCC1c2ccncc2-c3ccc4nc5cccnc5n4C(=O)N6CCNCC6C7CC7</chem>
492	 <chem>CC(C)C(=O)N1CCNCC1c2ccc3cc4ccccc4n3c2C(=O)N5CCNCC5C6CC6</chem>






#	Struktur
493	
494	
495	
496	
497	






#	Struktur
498	
499	
500	
501	
502	






#	Struktur
503	
504	
505	
506	
507	
508	






#	Struktur
509	
510	
511	
512	
513	
514	






#	Struktur
515	
516	
517	
518	
519	






#	Struktur
520	
521	
522	
523	
524	






#	Struktur
525	
526	
527	
528	
529	






#	Struktur
530	
531	
532	
533	
534	






#	Struktur
535	
536	
537	
538	
539	






#	Struktur
540	
541	
542	
543	
544	






#	Struktur
545	
546	
547	
548	
549	





#	Struktur
550	 <chem>CCOC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccc(cc4)[C@H]5CCN(C)C5</chem>
551	 <chem>CCOC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccc(cc4)[C@@H]5CCN(C)C5</chem>
552	 <chem>CCOC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccnnc4c5ccccc5N(C)C</chem>
553	 <chem>CCNC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccnnc4c5ccccc5N(C)C</chem>
554	 <chem>CCNC(=O)N1CCN(C1)c2cnc3cc(ccn32)c4ccnnc4c5ccccc5N(C)C</chem>






#	Struktur
555	
556	
557	
558	
559	






#	Struktur
560	
561	
562	
563	
564	

#	Struktur
565	
566	
567	
568	
569	






#	Struktur
570	
571	
572	
573	
574	






#	Struktur
575	
576	
577	
578	






#	Struktur
579	
580	
581	
582	
583	






#	Struktur
584	
585	
586	
587	
588	





#	Struktur
589	
590	
591	
592	
593	






#	Struktur
594	
595	
596	
597	
598	






#	Struktur
599	
600	
601	
602	
603	






#	Struktur
604	
605	
606	
607	
608	






#	Struktur
609	
610	
611	
612	
613	






#	Struktur
614	
615	
616	
617	






#	Struktur
618	
619	
620	
621	
622	






#	Struktur
623	
624	
625	
626	
627	






#	Struktur
628	
629	
630	
631	
632	






#	Struktur
633	
634	
635	
636	
637	






#	Struktur
638	
639	
640	
641	
642	






#	Struktur
643	
644	
645	
646	
647	






#	Struktur
648	
649	
650	
651	
652	





#	Struktur
653	
654	
655	
656	
657	

#	Struktur
658	
659	
660	
661	
662	





#	Struktur
663	
664	
665	
666	
667	





#	Struktur
668	
669	
670	
671	
672	






#	Struktur
673	
674	
675	
676	
677	






#	Struktur
678	
679	
680	
681	





#	Struktur
682	
683	
684	
685	
686	






#	Struktur
687	
688	
689	
690	






#	Struktur
691	
692	
693	
694	






#	Struktur
695	
696	
697	
698	
699	






#	Struktur
700	
701	
702	
703	
704	






#	Struktur
705	
706	
707	
708	





#	Struktur
709	
710	
711	
712	
713	





#	Struktur
714	
715	
716	
717	
718	





#	Struktur
719	
720	
721	
722	
723	







#	Struktur
724	
725	
726	
727	
728	

#	Struktur
729	
730	
731	
732	
733	






#	Struktur
734	
735	
736	
737	






#	Struktur
742	
743	
744	
745	






#	Struktur
746	
747	
748	
749	






#	Struktur
755	
756	
757	
758	
759	
760	





#	Struktur
761	 <chem>CC(C)(O)C(N)c1ccc(cc1)-c2nc3ccncc3nc2N4CCN(CC4)C(=O)C5CC5</chem>
762	 <chem>NC[C@H](O)c1ccc(cc1)-c2nc3ccncc3nc2N4CCN(CC4)C(=O)O[C@H]5CCOC5</chem>
763	 <chem>NC[C@@H](O)c1ccc(cc1)-c2nc3ccncc3nc2N4CCN(CC4)C(=O)O[C@H]5CCOC5</chem>
764	 <chem>CCOC(=O)N1CCN(CC1)c2nc3ccncc3nc2-c4ccc(cc4)C(C)(O)C(N)</chem>
765	 <chem>CCS(=O)(=O)c1ccc(cc1)-c2nc3ccncc3nc2N4CCN(CC4)C(=O)C5CC5</chem>






#	Struktur
766	
767	
768	
769	
770	





#	Struktur
771	
772	
773	
774	
775	






#	Struktur
781	
782	
783	
784	
785	






#	Struktur
786	
787	
788	
789	
790	






#	Struktur
791	
792	
793	
794	






#	Struktur
795	
796	
797	
798	
799	





#	Struktur
800	
801	
802	
803	





#	Struktur
804	
805	
806	
807	
808	






#	Struktur
809	
810	
811	
812	
813	





#	Struktur
814	
815	
816	
817	
818	






#	Struktur
819	
820	
821	
822	
823	






#	Struktur
824	
825	
826	
827	





#	Struktur
828	
829	
830	
831	





#	Struktur
832	
833	
834	
835	
836	





#	Struktur
837	 <chem>CC(C)N1CCN(C1)C2=CC=C(C=C2)C3=CC4=CC=CC=C4C5=CC6=CC=CC=C6N5C7CCN(C7)C(=O)OCCO</chem>
838	 <chem>CC(C)N1CCN(C1)C2=CC=C(C=C2)C3=CC4=CC=CC=C4C5=CC6=CC=CC=C6N5C7CCN(C7)C(=O)Nc8ccccc8</chem>
839	 <chem>CC(C)N1CCN(C1)C2=CC=C(C=C2)C3=CC(=CC=C3C#N)C(OC)=C4C=CC=CC=C4C5=CC6=CC=CC=C6N5C7CCN(C7)C(=O)OC8CC8</chem>
840	 <chem>CC(C)N1CCN(C1)C2=CC=C(C=C2)C3=CC4=CC=CC=C4C5=CC6=CC=CC=C6N5C7CCN(C7)C(=O)OC8CC8C#N</chem>

#	Struktur
841	
842	
843	
844	
845	





#	Struktur
846	
847	
848	
849	
850	





#	Struktur
851	 <chem>CC1(C)CCN(CC1)c2ccc(cc2)-c3cc4nncn4c3N5CCN(CC5)C(=O)Nc6cccc(F)c6</chem>
852	 <chem>CC1(C)CCN(CC1)c2ccc(OC)c(C#N)c2-c3cc4nncn4c3N5CCN(CC5)C(=O)Nc6cc(C)cc(C)c6</chem>
853	 <chem>CC1(C)CCN(CC1)c2ccc(cc2)-c3cc4nncn4c3N5CCN(CC5)C(=O)Oc6ccc(cc6)C(=O)N(C)C7=CC=CC=C7C8=CC=CC=C8C9=CC=CC=C9C10=CC=CC=C10</chem>
854	 <chem>CC1(C)CCN(CC1)c2ccc(OC)c(C#N)c2-c3cc4nncn4c3N5CCN(CC5)C(=O)Oc6ccc(cc6)C(=O)N(C)C7=CC=CC=C7C8=CC=CC=C8C9=CC=CC=C9C10=CC=CC=C10</chem>






#	Struktur
855	
856	
857	
858	





#	Struktur
864	
865	
866	
867	





#	Struktur
868	 <chem>CC(C)NC(=O)N1CCN(CC1)c2cnc3cc(ccn32)-c4ccc(cc4)[C@H]5CCN(CC5)C(F)(F)F</chem>
869	 <chem>CC(C)NC(=O)N1CCN(CC1)c2cnc3cc(ccn32)-c4ccc(cc4)[C@@H]5CCN(CC5)C(F)(F)F</chem>
870	 <chem>CC(C)NC(=O)N1CCN(CC1)c2cnc3cc(ccn32)-c4ccc(cc4)[C@H]5CCN(CC5)C(F)(F)F</chem>
871	 <chem>CC(C)NC(=O)N1CCN(CC1)c2cnc3cc(ccn32)-c4ccc(cc4)[C@H]5CCN(CC5)C(F)(F)F</chem>

#	Struktur
872	
873	
874	
875	

#	Struktur
876	
877	
878	
879	

#	Struktur
880	
881	
882	
883	
884	


#	Struktur
885	
886	
887	
888	

#	Struktur
889	
890	
891	
892	

11. Farmaceutisk sammensætning, der omfatter mindst én forbindelse ifølge et hvilket som helst af kravene 1 til 10 eller et farmaceutisk acceptabelt salt deraf og en farmaceutisk acceptabel bærer.

12. Forbindelse, der er 6-halogen-pyrrolo[1,2-b]pyridazin-4-ol, hvor halogen er valgt blandt Cl, I og Br, særligt 6-brompyrrolo[1,2-b]pyridazin-4-ol, idet forbindelsen eventuelt har en renhed på mere end 90 %.

13. Fremgangsmåde til syntetisering en forbindelse ifølge krav 12, hvilken fremgangsmåde omfatter det trin, at en forbindelse med formlen C-1:



kombineres med en forbindelse med formlen D-1:



hvor:

R^{21} er chlor, brom eller iod og fortrinsvis er brom;

R^{22} er en fraspaltelig enhed, der fortrinsvis er valgt blandt $-N(R^{24})(R^{25})$ og $-OR^{24}$, hvor hvert af R^{24} og R^{25} uafhængigt er C_1 - C_4 -alkyl og mere fortrinsvis er $-N(CH_3)_2$; og

R^{23} er en elektrontiltrækkende gruppe, der fortrinsvis er valgt blandt methylcarbonyl, t-butylcarbonyl, 4-nitrophenylcarbonyl, 4-cyanophenylcarbonyl, 4-trifluormethylphenylcarbonyl, 4-fluorphenylcarbonyl, 4-trifluormethylcarbonylphenylcarbonyl, 4-ethoxycarbonylphenylcarbonyl, 4-trifluormethylsulfonylphenylcarbonyl, 2,4,6-trimethylphenylcarbonyl, 2,4,6-trimethyl-3,5-dinitrophenylcarbonyl, 2-trifluormethyl-4-nitrophenyl, 2,4-dinitrophenyl og diphenylphosphinyl og mere fortrinsvis er 4-nitrophenylcarbonyl.

14. Fremgangsmåde ifølge krav 13, hvor forbindelsen med formlen C-1 og forbindelsen med formlen D-1 opløses i et polært opløsningsmiddel, hvor det polære opløsningsmiddel fortrinsvis er valgt blandt N-methyl-2-pyrrolidin ("NMP"), N,N-dimethylacetamid ("DMAC"), dimethylformamid ("DMF"), tetrahydrofuran ("THF"), methyl-tetrahydrofuran ("MeTHF"), dimethylsulfoxid ("DMSO") og cyclopentylmethylether ("CPME") og mere fortrinsvis er valgt blandt NMP og DMAC.

15. Fremgangsmåde ifølge krav 13, hvor forbindelsen med formelen C-1 behandles med en base, hvor den fortrinsvis er valgt blandt $\text{KOC}(\text{CH}_3)_3$, $\text{NaOC}(\text{CH}_3)_3$, $\text{LiOC}(\text{CH}_3)_3$, $\text{LiC}(\text{CH}_3)_3$, $\text{Li}(\text{CH}_2)_3\text{CH}_3$, $\text{LiN}(\text{C}_3\text{H}_7)_2$, NaOCH_3 , $\text{NaOCH}_2\text{CH}_3$, KOCH_3 , LiOCH_3 , $\text{LiOCH}_2\text{CH}_3$ og KOCH_2CH_3 og mere fortrinsvis er $\text{KOC}(\text{CH}_3)_3$.

16. Fremgangsmåde ifølge krav 13, hvor fremgangsmåden yderligere omfatter tilsætning af et protoneringsmiddel, hvor protoneringsmidlet fortrinsvis er valgt blandt NH_4Cl , NaHCO_3 , KHCO_3 , LiHCO_3 , eddikesyre, HCl , HBr og H_2SO_4 og mere fortrinsvis er NH_4Cl .

17. Forbindelse til anvendelse i en fremgangsmåde til behandling eller bedring af fibrodysplasia ossificans progressiva hos et individ, hvilken fremgangsmåde omfatter det trin, at der til individet, som har behov derfor, administreres en farmaceutisk effektiv mængde af mindst én forbindelse ifølge et hvilket som helst af kravene 1 til 10 eller et farmaceutisk acceptabelt salt deraf eller en farmaceutisk sammensætning ifølge krav 11, eventuelt hvor individet har en mutation i et ALK2-gen, der resulterer i ekspression af et ALK2-enzym med en aminosyremodifikation, der er valgt blandt én eller flere af L196P, PF197-8L, R202I, R206H, Q207E, R258S, R258G, G328A, G328W, G328E, G328R, G356D og R375P, for eksempel hvor ALK2-enzymet har aminosyremodifikationen R206H.

18. Forbindelse til anvendelse i en fremgangsmåde til behandling eller bedring af diffust intrinsisk pontint gliom hos et individ, hvilken fremgangsmåde omfatter det trin, at der til individet, som har behov derfor, administreres en farmaceutisk effektiv mængde af mindst én forbindelse ifølge et hvilket som helst af kravene 1 til 10 eller et farmaceutisk acceptabelt salt deraf eller en farmaceutisk sammensætning ifølge krav 11, eventuelt hvor individet har en mutation i et ALK2-gen, der resulterer i ekspression af et ALK2-enzym med en aminosyremodifikation, der er valgt blandt én eller flere af R206H, G328V, G328W, G328E og G356D, for eksempel hvor ALK2-enzymet har aminosyremodifikationen R206H.