



US 20180147226A1

(19) **United States**

(12) **Patent Application Publication**
FILLMORE et al.

(10) **Pub. No.: US 2018/0147226 A1**

(43) **Pub. Date: May 31, 2018**

(54) **METHODS AND ASSAYS FOR
COMBINATION TREATMENT OF CANCER**

(60) Provisional application No. 61/735,303, filed on Dec.
10, 2012.

(71) Applicant: **CHILDREN'S MEDICAL CENTER
CORPORATION**, Boston, MA (US)

Publication Classification

(72) Inventors: **Christine M. FILLMORE**, Waltham,
MA (US); **Carla F. KIM**, Boston, MA
(US)

(51) **Int. Cl.**
A61K 31/706 (2006.01)
A61K 31/7048 (2006.01)
C12Q 1/6886 (2006.01)
A61K 45/06 (2006.01)

(73) Assignee: **CHILDREN'S MEDICAL CENTER
CORPORATION**, Boston, MA (US)

(52) **U.S. Cl.**
CPC *A61K 31/706* (2013.01); *A61K 31/7048*
(2013.01); *C12Q 2600/156* (2013.01); *C12Q*
2600/106 (2013.01); *A61K 45/06* (2013.01);
C12Q 1/6886 (2013.01)

(21) Appl. No.: **15/865,647**

(22) Filed: **Jan. 9, 2018**

Related U.S. Application Data

(62) Division of application No. 14/650,473, filed on Jun.
8, 2015, now Pat. No. 9,895,390, filed as application
No. PCT/US2013/069557 on Nov. 12, 2013.

(57) **ABSTRACT**

The technology described herein is directed to the treatment
of cancer, e.g. methods and assays relating to selecting and
administering a chemotherapy with or without an EZH2
inhibitor.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A




FIG. 3B








FIG. 4A

**FIG. 4B**






FIG. 5A



FIG. 5B




FIG. 5C




FIG. 6A

FIG. 6B




FIG. 6C




FIG. 6D




FIG. 7A




FIG. 7B






FIG. 7C






FIG. 8C




FIG. 8D




FIG. 9A






FIG. 9B



H157-SENSITIZED LINE

FIG. 10A



H460-PROTECTED LINE

FIG. 10B



FIG. 10C




FIG. 10D




FIG. 10E




FIG. 10F




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D




FIG. 12A




FIG. 12B




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17A

FIG. 17B




FIG. 18A




FIG. 18B




FIG. 18C






FIG. 18D






FIG. 19A




FIG. 19B




FIG. 19C




FIG. 19D




FIG. 20B




FIG. 20A




FIG. 21A




FIG. 21B




FIG. 21C




FIG. 21D




FIG. 21E




FIG. 22




FIG. 23A








FIG. 23B








FIG. 23G




FIG. 24A




FIG. 24B




FIG. 24C




FIG. 25A




FIG. 25B





FIG. 25C




FIG. 26A




FIG. 26B




FIG. 26C




FIG. 27A




FIG. 27B




FIG. 28A




FIG. 28B




FIG. 28C

Cell Line	Gender	Age	Ethnicity	Subtype	Stage	Smoke (pack/yr)	Biopsy	PIK3CA	Kras	Nras	p53	SMARCA4	EGFR	CDKN2A	LKB1	B-Raf
HCC2450	M	52	Caucasian	SCC	IIIB	unknown	primary	H1047R	N/A	N/A	deletion	N/A	N/A	N/A	N/A	WT
PC9	M	45	East Asian	AdC	N/A	unknown	N/A	WT	WT	WT	N/A	N/A	E746 A750del	G67V	WT	WT
H522	M	60	Caucasian	AdC	II	60	primary	WT	WT	WT	P191*	P279*	WT	WT	WT	WT
A549	M	58	Caucasian	AdC	N/A	unknown	primary	WT	G12S	WT	WT	Q729*	WT	deletion	Q37*	WT
Calu1	M	47	Caucasian	EpiC	III	unknown	pleura	N/A	G12C	WT	WT	WT	WT	WT	WT	WT
H2126	M	65	Caucasian	LC	IIIB	non-smoker	pleura	WT	WT	WT	E62*	W764R	WT	deletion	deletion	WT
H157	M	59	Caucasian	SCC	IIIB	50	metastasis	WT	G12R	WT	E298*	T58*	WT	E69*	deletion	WT
H1975	F	N/A	N/A	AdC	N/A	non-smoker	primary	G118D	WT	WT	R273H	WT	T790M, L858R	E69*	Q37* het	WT
HCC15	M	47	African- American	SCC	N/A	unknown	primary	WT	WT	mut	mut	M272*	WT	mut/hypem	deletion	WT
H2030	M	N/A	N/A	AdC	N/A	unknown	metastasis	WT	G12C	WT	G262V	deletion	WT	WT	E317*	WT
H1650	M	27	Caucasian	AdC (BAC)	IIIB	10	pleura	WT	WT	WT	deletion	WT	E746X	deletion	WT	WT
HCC4006	M	N/A	Caucasian	AdC	N/A	non-smoker	N/A	WT	WT	WT	N/A	N/A	L747 E749del	N/A	WT	WT
HCC927	F	39	Caucasian	AdC	N/A	4	N/A	WT	WT	WT	N/A	WT	E746 A750del	N/A	WT	WT
H1299	M	43	Caucasian	LC	IIIA	50	lymph node	WT	WT	Q61K	deletion	Y560*	WT	hypermeth	WT	WT
H322	M	52	Caucasian	AdC (BAC)	IV	60	lymph node	WT	WT	WT	N/A	N/A	WT	deletion	WT	WT
HCC95	M	64	Caucasian	SCC	IV	unknown	pleura	Amplified	WT	WT	WT	WT	WT	mut/hypem	WT	WT
H2009	F	68	Caucasian	AdC	IV	30	lymph node	E711D	G12A	WT	R273L	WT	WT	WT	WT	WT
Sw1573	F	44	Caucasian	AdC	N/A	unknown	primary	K111E	G12C	WT	WT	WT	WT	deletion	WT	N/A
H2122	F	46	Caucasian	AdC	IV	30	pleura	WT	G12C	WT	Q16L, C176F	WT	WT	deletion	P281*	WT
H520	M	N/A	N/A	SCC	N/A	non-smoker	primary	WT	WT	WT	W146*	WT	WT	G45*	WT	WT
Calu6	F	61	Caucasian	AdC (Ana)	N/A	unknown	primary	N/A	Q61K	WT	R196*	WT	WT	WT	WT	WT
Calu3	M	25	Caucasian	AdC	N/A	unknown	pleura	Amplified	WT	WT	M237I	WT	WT	hypermeth	WT	WT
H460	M	N/A	N/A	LC	N/A	non-smoker	primary	E545K	Q61H	WT	mut	WT	WT	deletion	Q37*	WT
H3255	F	47	Caucasian	AdC	IIIB	non-smoker	N/A	Amplified	G12C	WT	N/A	N/A	L858R	N/A	N/A	WT
H23	M	51	African- American	AdC	N/A	40	primary	WT	G12C	WT	M246I	K1533N	WT	hypermeth	W322*	WT
H441	M	33	N/A	AdC (pap)	IIIA	non-smoker	pleura	WT	G12V	WT	R158L	WT	WT	hypermeth	WT	WT

FIG. 29

METHODS AND ASSAYS FOR COMBINATION TREATMENT OF CANCER

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application is divisional of co-pending application Ser. No. 14/650,473 filed Jun. 8, 2015 which is a 35 U.S.C. § 371 National Phase Entry Application of International Application No. PCT/US2013/069557 filed Nov. 12, 2013, which designates the U.S., and which claims benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 61/735,303 filed Dec. 10, 2012, the contents of which are incorporated herein by reference in their entireties.

GOVERNMENT SUPPORT

[0002] This invention was made with Government support under grants Nos. RO1 HL090136 and U01 HL100402 awarded by the National Institutes of Health and the National Heart, Lung and Blood Institute. The government has certain rights in the invention.

SEQUENCE LISTING

[0003] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Nov. 6, 2013, is named 701039-073361-PCT_SL.txt and is 77,118 bytes in size.

TECHNICAL FIELD

[0004] The technology described herein relates to the treatment of cancer.

BACKGROUND

[0005] The gene expression of a cell can be regulated by epigenetic mechanisms, i.e. by altering the physical state of the chromosomes themselves. One such epigenetic mechanism is based upon the modification of histones. Depending upon the modification, an area of the genome can be made more or less accessible to transcriptional machinery, essentially up or down regulating gene expression in that area, leading to changes in cell behavior and characteristics.

[0006] Such epigenetic mechanisms, such as histone modifications, are thought to be crucial for the survival of cancer cells, particularly metastatic cells and those which are resistant to standard therapies (Baylin, 2011, Crea et al., 2011a, Min et al., 2010, Iliopoulos et al., 2010). Therefore, combining epigenetic therapies with chemotherapy and radiation treatments may allow for more complete treatment responses.

[0007] Polycomb Repressive Complexes (PRCs) are key regulators of, e.g. histone modifications in cancer cells (Simon and Lange, 2008, Lee et al., 2006, Ben-Porath et al., 2008). PRC2 often contains EZH2, a methyltransferase that tri-methylates histone H3 at lysine 27 (H3K27me3). Accordingly, EZH2 inhibition has been identified as a possible therapeutic strategy for the treatment of cancer.

SUMMARY

[0008] As described herein, the inventors have discovered that cancer cells can exhibit one of two phenotypes in response to EZH2 inhibition. In the first phenotype, the cell is rendered more sensitive to treatment with a chemothera-

peutic agent (i.e. an agent other than the agent which inhibits EZH2), while in the second phenotype, the cell becomes more protected from treatment with the chemotherapeutic agent. Further, the inventors have identified methods and assays for determining which phenotype a cell will exhibit, permitting subjects who will benefit from the combination therapy to be identified and treated; and permitting practitioners to avoid administering a treatment that would be harmful to a subject with the resistant phenotype.

[0009] In some aspects, described herein is a method of treating cancer, the method comprising: administering or prescribing a chemotherapeutic agent and an EZH2 inhibitor to a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell.

[0010] In some aspects, described herein, is a method of identifying a subject who is a candidate for treatment of cancer with a combination therapy comprising an EZH2 inhibitor and a chemotherapeutic agent, the method comprising: detecting a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject; wherein if the sensitizing mutation is detected, the subject is identified as a candidate for cancer treatment with a combination therapy comprising a chemotherapeutic agent and an EZH2 inhibitor; and wherein if the sensitizing mutation is not detected, the subject is identified as a candidate for cancer treatment which does not comprise administering an EZH2 inhibitor.

[0011] In some aspects, described herein is a method of classifying a tumor cell as an EZH2 combination treatment responsive tumor, the method comprising: detecting a sensitizing mutation of BRG1, EGFR, or B-RAF in a sample comprising the tumor cell; wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF classifies the tumor as a EZH2 combination treatment responsive tumor and the absence of a sensitizing mutation of BRG1, EGFR, or B-RAF classifies the tumor as a EZH2 combination treatment non-responsive tumor.

[0012] In some embodiments of the foregoing aspects, the EZH2 inhibitor is selected from the group consisting of: an inhibitory nucleic acid; DZNep; and S-adenosyl-L-homocysteine. In some embodiments of the foregoing aspects, the chemotherapeutic agent is selected from the group consisting of a topoisomerase inhibitor; a topoisomerase I inhibitor; and a topoisomerase II inhibitor. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of: camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of: doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a PARP inhibitor. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a CDK1 inhibitor. In some embodiments of the foregoing aspects, the chemotherapeutic agent is an EGFR inhibitor. In some embodiments of the foregoing aspects, the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of: a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*. In some embodiments of the foregoing aspects, the sensi-

tizing mutation of EGFR comprises a mutation selected from the group consisting of: a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R. In some embodiments of the foregoing aspects, the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of: G496A and L597V. In some embodiments of the foregoing aspects, the presence of the mutation is determined using an assay selected from the group consisting of: hybridization; sequencing; exome capture; PCR; and high-throughput sequencing. In some embodiments of the foregoing aspects, the mutation is present in the genomic DNA of the tumor cell. In some embodiments of the foregoing aspects, the mutation is present in the mRNA transcripts of the tumor cell. In some embodiments of the foregoing aspects, the cancer is selected from the group consisting of: lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers. In some embodiments of the foregoing aspects, the method further comprises the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.

[0013] In some aspects, described herein is an assay comprising: subjecting a tumor cell sample from a subject to at least one analysis to detect the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF; wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with an EZH2 inhibitor and a chemotherapeutic agent. In some embodiments, the presence of the mutation is determined using an assay selected from the group consisting of: hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.

[0014] In some aspects, described herein is an assay for selecting a treatment regimen for a subject with cancer, comprising: subjecting a nucleotide molecule derived from a biological sample of a subject, who is determined to suffer from or have a risk for cancer, to at least one genotyping analysis adapted to determine the presence of a sensitizing mutation in one or more of B-RAF, EGFR, and BRG1; wherein if at least one sensitizing mutation is determined to be present, a treatment regimen comprising a combination of an EZH2 inhibitor and a chemotherapeutic agent is administered. In some embodiments, wherein if no sensitizing mutations are determined to be present, a treatment regimen comprising a combination of an EZH2 inhibitor and a chemotherapeutic agent is not administered. In some embodiments, the biological sample comprises a tumor cell.

[0015] In some aspects, described herein is an assay comprising: contacting a tumor cell sample obtained from a human subject having cancer with a nucleic acid probe to detect the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF; and detecting the presence or intensity of a signal which indicates the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF; wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with an EZH2 inhibitor and a chemotherapeutic agent.

[0016] In some embodiments of the foregoing aspects, a detectable signal is generated by the probe when a sensitizing mutation is present. In some embodiments of the foregoing aspects, the probe is detectably labeled. In some

embodiments of the foregoing aspects, the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of: a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*. In some embodiments of the foregoing aspects, the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of: a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R. In some embodiments of the foregoing aspects, the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of: G496A and L597V. In some embodiments of the foregoing aspects, the mutation is present in the genomic DNA of the tumor cell. In some embodiments of the foregoing aspects, the mutation is present in the mRNA transcripts of the tumor cell. In some embodiments of the foregoing aspects, the assay further comprises the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.

[0017] In some aspects, described herein is a method of determining whether a subject is likely to respond to a combination treatment for cancer, the method comprising: detecting the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject; wherein the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF indicates the subject has an increased likelihood of responding to a treatment to cancer; wherein the treatment for cancer comprises administration of an EZH2 inhibitor and a chemotherapeutic agent.

[0018] In some embodiments of the foregoing aspects, the EZH2 inhibitor is selected from the group consisting of: an inhibitory nucleic acid; DZNep; S-adenosyl-L-homocysteine. In some embodiments of the foregoing aspects, the chemotherapeutic agent is selected from the group consisting of: a topoisomerase inhibitor; a topoisomerase I inhibitor; a topoisomerase II inhibitor. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of: camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of: doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a PARP inhibitor. In some embodiments of the foregoing aspects, the chemotherapeutic agent is a CDK1 inhibitor. In some embodiments of the foregoing aspects, the chemotherapeutic agent is an EGFR inhibitor. In some embodiments of the foregoing aspects, the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of: a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*. In some embodiments of the foregoing aspects, the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of: a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R. In some embodiments of the foregoing aspects, the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of: G496A and L597V. In some embodiments of the foregoing aspects, the

presence of the mutation is determined using an assay selected from the group consisting of: hybridization; sequencing; exome capture; PCR; and high-throughput sequencing. In some embodiments of the foregoing aspects, the mutation is present in the genomic DNA of the tumor cell. In some embodiments of the foregoing aspects, the mutation is present in the mRNA transcripts of the tumor cell. In some embodiments of the foregoing aspects, the cancer is selected from the group consisting of: lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.

[0019] In some aspects, described herein is a computer system for determining if a subject will be responsive to a cancer treatment, the system comprising: a determination module configured to detect the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from a subject; a storage module configured to store output data from the determination module; a comparison module adapted to compare the data stored on the storage module with a reference level, and to provide a retrieved content, and a display module for displaying whether a BRG1, EGFR, or B-RAF sensitizing mutation was detected in the tumor cell sample; wherein the cancer treatment comprises the administration of an EZH2 inhibitor and a chemotherapeutic agent. In some embodiments, the determining module measures the intensity of a detectable signal from a RT-PCR assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample. In some embodiments, the determining module measures the intensity of a detectable signal from a sequencing assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample. In some embodiments, the determining module measures the intensity of a detectable signal from a hybridization assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample. In some embodiments, if the computing module determines that the a sensitizing mutation of BRG1, EGFR, or B-RAF is present in the sample obtained from the subject, the display module displays a signal indicating that a sensitizing mutation has been detected. In some embodiments, the signal indicates that the subject has an increased likelihood of responding to treatment with an EZH2 inhibitor and chemotherapeutic agent. In some embodiments, the EZH2 inhibitor is selected from the group consisting of: an inhibitory nucleic acid; DZNep; S-adenosyl-L-homocysteine. In some embodiments, the chemotherapeutic agent is selected from the group consisting of: a topoisomerase inhibitor; a topoisomerase I inhibitor; a topoisomerase II inhibitor. In some embodiments, the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of: camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D. In some embodiments, the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of: doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331. In some embodiments, the chemotherapeutic agent is a PARP inhibitor. In some embodiments, the chemotherapeutic agent is a CDK1 inhibitor. In some embodiments, the chemotherapeutic agent is an EGFR inhibitor. In some embodiments, the sensitizing mutation of BRG1 com-

prises a mutation selected from the group consisting of: a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*. In some embodiments, the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of: a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R. In some embodiments, the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of: G496A and L597V. In some embodiments, the presence of the mutation is determined using an assay selected from the group consisting of: hybridization; sequencing; exome capture; PCR; and high-throughput sequencing. In some embodiments, the mutation is present in the genomic DNA of the tumor cell. In some embodiments, the mutation is present in the mRNA transcripts of the tumor cell. In some embodiments, the cancer is selected from the group consisting of: lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.

[0020] In some aspects, described herein is a kit comprising: a nucleic acid probe which is specific for a sensitizing mutation of BRG1, EGFR, or B-RAF. In some embodiments, the probe is detectably labeled. In some aspects, described herein is a kit comprising: a nucleic acid probe which can amplify a nucleic acid sequence comprising a sensitizing mutation of BRG1, EGFR, or B-RAF. In some embodiments, the kit further comprises a reagent for producing a detectable signal.

[0021] In some aspects, described herein is the use of an EZH2 inhibitor in combination with a chemotherapeutic agent in treatment of a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell obtained from the subject. In some embodiments, the EZH2 inhibitor is selected from the group consisting of: an inhibitory nucleic acid; DZNep; and S-adenosyl-L-homocysteine. In some embodiments, the chemotherapeutic agent is selected from the group consisting of: a topoisomerase inhibitor; a topoisomerase I inhibitor; and a topoisomerase II inhibitor. In some embodiments, the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of: camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D. In some embodiments, the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of: doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331. In some embodiments, the chemotherapeutic agent is a PARP inhibitor. In some embodiments, the chemotherapeutic agent is a CDK1 inhibitor. In some embodiments, the chemotherapeutic agent is an EGFR inhibitor. In some embodiments, the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of: a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*. In some embodiments, the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of: a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R. In some embodiments, the sensitizing mutation of B-RAF comprises a mutation selected from the group con-

sisting of: G496A and L597V. In some embodiments, the presence of the mutation is determined using an assay selected from the group consisting of: hybridization; sequencing; exome capture; PCR; and high-throughput sequencing. In some embodiments, the mutation is present in the genomic DNA of the tumor cell. In some embodiments, the mutation is present in the mRNA transcripts of the tumor cell. In some embodiments, the cancer is selected from the group consisting of: lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers. In some embodiments, the use further comprises the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] FIGS. 1A-1C demonstrate that EZH2 knock-down has differential effects on chemotherapy sensitivities. FIG. 1A depicts images of film. Cell lines were infected with pLK0.1 lentivirus encoding either shGFP or shEZH2 and selected with puromycin for 1 week. Western Blot for EZH2 and H3K27me3 was performed on whole cell extracts from indicated lines for EZH2 and its catalytic mark H3K27me3, β -actin is shown as loading control. FIG. 1B depicts a graph of growth rates of matched shGFP and shEZH2 cell lines assayed by Cell Titer Glo at days 3, 5, 7, and 10 post plating at equal densities. Average luminescence \pm s.e.m. for 3 replicate wells is shown based on when each shGFP cell line reached logarithmic growth, * indicates $p<0.03$. FIG. 1C depicts a graph of cell survival of shGFP and shEZH2 cell lines plated and treated with etoposide for 4 days. Average change \pm s.e.m. in % survival between shGFP and shEZH2 lines is graphed for 1 μ M and 10 μ M etoposide, $n=3$ technical replicates, ** indicates $p<0.0001$.

[0023] FIGS. 2A-2C demonstrate that EZH2 inhibition by DZNep has differential effects on chemotherapy sensitivities. FIG. 2A depicts a Western Blot for EZH2 and H3K27me3 performed on whole cell extracts 4 days after indicated treatments, β -actin shown as a loading control. FIG. 2B depicts a graph of the growth rates of matched shGFP and shGFP cells treated with 1 μ M DZNep and assayed by Cell Titer Glo at days 3, 5, 7, and 10 post plating at equal densities. Average luminescence \pm s.e.m. for 3 replicate wells is shown based on when each shGFP cell line reached logarithmic growth, * indicates $p<0.03$. FIG. 2C depicts a graph of the response to Etoposide. Cell lines were plated at equal densities and treated a range of etoposide doses with or without 1 μ M DZNep. On the 4th day, cell viability was measured using Cell Titer Glo and the percent survival at each dose of etoposide was calculated. Dose response curves were generated for the no DZNep and DZNep treated cells using the GraphPad prism software and IC50s were extrapolated. The fold change \pm s.e.m. of etoposide IC50 in response to DZNep is plotted, $n=4$ biological replicates, * indicates $p<0.05$. Cell lines with mutations in BRG1, EGFR and B-Raf are indicated.

[0024] FIGS. 3A-3B demonstrate that in vitro sensitivities to DZNep/etoposide predict in vivo responses. Either the H157 (FIG. 3A) or H23 (FIG. 3B) cell lines were injected into the flanks of Nude mice and tumors were allowed to form. On day 12, mice were randomly segregated into cohorts that received either placebo or etoposide for five

consecutive days. For H157 mice, a subset of the placebo and etoposide cohorts also received 2 mg/kg/day DZNep twice in the first week (dosing scheme A), * indicates $p=0.03$ untreated vs. DZNep, ** indicates $p=0.0002$ untreated vs. dual, and *** indicates $p=0.015$ for etoposide vs. dual. For the H23 mice, the DZNep dose administered was 1 mg/kg/day twice per week for two weeks (dosing scheme B), * indicates $p=0.001$ untreated vs. DZNep, ** indicates $p=0.01$ dual vs. etoposide, and *** indicates $p<0.0001$ for dual vs. DZNep. Tumor growth was measured by caliper every other day until the tumor burden for the control mice was the maximum allowed by CHB IACUC. Results for at least 8 xenografts per condition were averaged and graphed \pm s.e.m. Each cell line with the alternative dosing strategy is shown in FIGS. 9A-9B.

[0025] FIGS. 4A-4E demonstrate that EZH2 inhibition in presence of chemotherapy differentially affects cell cycle. FIG. 4A depicts a graph of Gene Ontology term enrichment for the EZH2 co-expressed gene signature, of 64 enriched GO terms with $p<0.001$, 12 selected gene sets are graphed. All 64 GO terms are described in Table 3. FIG. 4B depicts a graph of survival of patients whose tumor samples had either high expression or low expression of the EZH2 co-expression gene signature. FIG. 4C depicts a graph of 7-AAD cell cycle flow cytometry performed on cell lines were plated cultured with or without 10 μ M etoposide or 1 μ M DZNep for 4 days. The average change in % of each cell cycle stage in response to DZNep is plotted, * indicates $p=0.0001$. FIG. 4D depicts a graph of the same analysis in (FIG. 4C) performed with shGFP and shEZH2 cells, ** indicates $p=0.0006$. For all these studies the average of 3 biological replicates \pm s.e.m. is shown. FIG. 4E depicts a graph of quantitative RT-PCR for change in expression of p57, Brg1 and EGFR in response to etoposide and etoposide/DZNep treatment. Results from two biological replicates of HCC2450, H157, H522 and HCC15 (sensitized) and H441, H460, H23, and Calu6 (protected) lines were averaged and graphed \pm s.e.m., * indicates $p=0.03$ for Brg1, ** indicates $p=0.005$ for EGFR, and *** indicates $p=0.01$ for p57.

[0026] FIGS. 5A-5C demonstrate that Brg1 and EGFR mutations are negatively correlated and predict sensitized phenotypes. FIG. 5A depicts a graph of EGFR and BRG1 expression in tumors. Tumors from the Director's Challenge with a normalized expression of more than 1400 for either EGFR or Brg1 were plotted and correlation was assessed, Spearman's correlation coefficient -0.619 , $p=0.0002$. FIG. 5B depicts a graph of average probe intensity of EGFR probe (201983_s_at) on the U133A Affymetrix array in primary squamous cell carcinoma samples from TCGA with various EGFR and Brg1 mutational statuses, * indicates $p=0.003$, ** indicates $p<0.0001$. FIG. 5C depicts a graph of average probe intensity of EGFR probe (201983_s_at) on the U133A Affymetrix array for cell lines with various EGFR and Brg1 mutational statuses, *** indicates $p=0.02$.

[0027] FIGS. 6A-6D demonstrate that Brg1 and EGFR genetically interact to control the sensitized phenotype. FIGS. 6A and 6B depict graphs of HCC15 (FIG. 6A) and H460 (FIG. 6B) cells as analyzed by 7AAD flow cytometry to assess changes in cell cycle status in response to DZNep, $n=2$ technical replicates, * indicates $p=0.002$, ** indicates $p=0.02$. FIGS. 6C and 6D depict graphs of quantitative RT-PCR for p57, Brg1 and EGFR expression in the various indicated treated transduced cell lines * indicates $p=x$, ** indicates $p=y$, *** indicates $p=z$.

[0028] FIGS. 7A-7C depict the formation of tumorspheres. FIG. 7A depicts images of western blots of shGFP and shEZH2 cell line lysates for H3K27me3, β -actin is shown as loading control. FIG. 7B depicts a graph of primary tumorsphere formation, visible spheres per 5,000 cells plated were counted at day 10. FIG. 7C depicts a graph of secondary non-adherent tumorsphere formation quantified 10 days after plating single dissociated cells from primary tumorspheres on super-low attachment plates. Average sphere number from 3 replicate wells is plotted.

[0029] FIGS. 8A-8D depict the effects of EZH2 combination treatment. FIG. 8A depicts a waterfall plot of cell line growth in 1 μ M DZNep for 4 days compared to untreated cells. FIG. 8B depicts a graph of the average growth of all sensitized and all protected lines grown in DZNep for 4 days compared to untreated cells. FIG. 8C depicts a graph of fold change in doxorubicin IC50 in response to DZNep, n=2 biological replicates. FIG. 8D depicts a graph of fold change in cis-platinum IC50 in response to DZNep, n=3 biological replicates.

[0030] FIGS. 9A-9B depict alternative Dosing Strategies for DZNep and etoposide for H157 and H23 xenografts. FIG. 9A depicts H157 xenografts with dosing scheme B, * indicates p=0.21 DZNep vs untreated and ** indicates p=0.009 DZNep vs dual. FIG. 9B depicts H23 xenografts with dosing scheme A, * indicates p=0.18 DZNep vs dual, ** indicates p=0.011 DZNep vs untreated.

[0031] FIGS. 10A-10F depict the effects of EZH2 inhibition and chemotherapy combination treatment. FIGS. 10A-10B depict representative 7AAD flow cytometry histograms analyzed with ModFit LT software. Both the 'sensitized' H157 line (FIG. 10A) and the 'protected' H460 lines (FIG. 10B) are shown. FIG. 10C depicts a graph of RT qPCR for EZH2 expression in indicated cell lines after 4 days of indicated treatments. FIG. 10D depicts a graph of RT qPCR for p27 and p21 expression. FIG. 10E depicts a graph of the results of quantitative RT-PCR for p27 and p21 in H522, H157, HCC15 (sensitized) and H460, H23, H441 (protected) lines in indicated treatments. FIG. 10F depicts a graph of the results of quantitative RT-PCR for p57 in H522, H157, HCC15, HCC2450, A549 (sensitized) and H460, H23, H441, Sw1573, and Calu6 (protected) lines. Average changes relative to untreated for the groups of lines are plotted.

[0032] FIGS. 11A-11D demonstrate gene expression in different cell lines/genotypes. FIG. 11A depicts a Venn diagram of differential gene expression over-lap between cell lines of various genotypes. FIG. 11B depicts a graph of averaged EZH2 probe (203358_s_at) intensity within each cell line genotype indicated. FIG. 11C depicts a graph of RT-qPCR of EZH2 and Suz12 gene expression in the indicated cell lines graphed relative to the average expression among the lines. A trend line for EZH2 expression was added to show no significant difference between sensitized (left side) and protected (right side) cell lines. FIG. 11D depicts graphs of additional Brg1 and EGFR modulation cell line RT-qPCR, the protected cell line H2122 and the sensitized cell line H1299 are shown.

[0033] FIGS. 12A-12B depict etoposide dose response curves. FIG. 12A depicts etoposide dose response curves for HCC15 cells corresponding to data graphed in FIG. 6A. FIG. 12B depicts etoposide dose response curves for H460 cells corresponding to data graphed in FIG. 6B.

[0034] FIG. 13 depicts a graphical representation of the effect of EZH2 inhibition combination treatment on responsive (sensitized) and nonresponsive cells (protected).

[0035] FIG. 14 is a diagram of an exemplary embodiment of a system for performing an assay for determining the presence of a sensitizing mutation in sample obtained from a subject.

[0036] FIG. 15 is a diagram of an embodiment of a comparison module as described herein.

[0037] FIG. 16 is a diagram of an exemplary embodiment of an operating system and instructions for a computing system as described herein.

[0038] FIGS. 17A-17B depict graphs of the results of quantitative RT-PCR for Brg1 and EGFR expression in the various indicated treated transduced cell lines.

[0039] FIGS. 18A-18F demonstrate that EZH2 expression is clinically relevant and influences etoposide sensitivities. FIG. 18A depicts graphs of survival of lung adenocarcinoma patients in the Director's Challenge dataset. Samples were hierarchically clustered using the primary tumor generated EZH2 co-expression signature (Table 2) into two risk groups. The Kaplan-Meier curves for the groups are shown for all patients (n=416), only Stage 1 patients (n=142) or only patients with moderately differentiated tumors (n=74). FIG. 18B depicts a graph of expression of EZH2 mRNA in human NSCLC cell lines compared to lung cells isolated from 4 normal patient samples (see methods). FIG. 18C depicts western blot results. Cell lines were infected with pLK0.1 lentivirus encoding either shGFP or shEZH2 and selected with puromycin. Western Blot for EZH2 and H3K27me3 was performed on whole cell extracts from indicated lines for EZH2 and its catalytic mark H3K27me3, total Histone H3 is shown as loading control. CR indicates a coding region targeting hairpin. FIG. 18D depicts graphs of shGFP and shEZH2 cell lines plated and treated with etoposide for 4 days. Cell growth was measured using CellTiter-Glo and dose response curves were constructed using GraphPad Prism software. Representative H157 and H460 curves are shown. FIG. 18E depicts a graph of average fold change in etoposide IC₅₀±s.e.m. is graphed, n=3-4 biological replicates, * indicates p<0.04, ** p<0.002. FIG. 18F depicts a graph of rescue of 3'UTR hairpin by over-expression of EZH2 cDNA. Also see FIG. 24A.

[0040] FIGS. 19A-19D demonstrate that chemical EZH2 inhibition sensitizes BRG1 or EGFR mutants to TopoII inhibitors. FIG. 19A depicts the results of a Western Blot. Western Blot for EZH2 and H3K27me3 was performed on whole cell extracts 4 days after administration of 1 μ M DZNep or vehicle, total Histone H3 is shown as a loading control. FIG. 19B depicts graphs of cell viability. Cell lines were plated at equal densities and treated a range of etoposide doses with or without 1 μ M DZNep. On the 4th day, cell viability was measured using CellTiter-Glo and the percent survival at each dose of etoposide was calculated. Dose response curves were generated for the vehicle- and DZNep-treated cells using the GraphPad prism software and IC₅₀s were extrapolated. The fold change±s.e.m. of etoposide IC₅₀ in response to DZNep is plotted, n=4 biological replicates, * indicates p<0.01. Cell lines with mutations in BRG1 or EGFR are indicated. FIG. 19C depicts a Western Blot for EZH2 and H3K27me3 performed on whole cell extracts from cultures treated with 10 μ M GSK126 or vehicle, total Histone H3 is shown as a loading control. FIG. 19D depicts a graph of dose response curves generated as described in

FIG. 19B using 10 μ M GSK126 in the place of 1 μ M DZNep, * indicates $p<0.03$, ** $p<0.01$. Also see FIGS. 25A-25C.

[0041] FIGS. 20A-20B demonstrate that in vitro sensitivities to DZNep/etoposide predict in vivo responses. FIG. 20A depicts a waterfall plot depicting tumor growth of EGFR-driven tumors after 2 weeks and 4 weeks of treatment with vehicle, etoposide, DZNep, and etoposide/DZNep. The y-axis indicates % tumor growth vs. day 0. Each bar represents an individual mouse. Statistical analyses were performed on the 4 week log2 transformed data, $p=0.008$ dual vs. DZNep and $p=0.004$ dual vs. etoposide. FIG. 20B depicts a waterfall plot depicting tumor growth of Kras/p53 tumors after 2 weeks and 4 weeks of treatment with vehicle and etoposide/DZNep.

[0042] FIGS. 21A-21E demonstrate that dual etoposide/EZH2 inhibition differentially affects cell cycle, apoptosis and anaphase bridging. FIG. 21A depicts graphs of the % S phase of the indicated cells. 7-AAD cell cycle flow cytometry was performed on cell lines cultured with or without 5 μ M etoposide or 1 μ M DZNep for 4 days. The average % S phase \pm s.e.m. of each culture is plotted, $n=3-4$ biological replicates, * indicates $p<0.05$ ** $p<0.009$. FIG. 21B depicts graphs of Annexin V+/7AAD- cells quantified by flow cytometry from cultures 4 days after indicated treatments. $n=3-4$ biological replicates, * indicates $p<0.03$ dual vs. etoposide, ** $p=0.001$ DZNep vs. vehicle. FIG. 21C depicts representative images of nuclei undergoing a normal anaphase and of nuclei that scored positively for the presence of anaphase bridges. FIG. 21D depicts a graph of the quantification of the percentage of anaphase cells that showed bridges in untreated or DZNep treated cell lines. $n=4$ biological replicates of 18-25 anaphase nuclei, * indicates $p<0.05$, ** $p<0.03$. FIG. 21E depicts a graph of real time RT-PCR for BRG1 levels in indicated cell lines in response to 4 days of 1 μ M DZNep, * indicates $p<0.02$, ** $p<0.0005$.

[0043] FIG. 22 demonstrates that BRG1 and EGFR are genetically antagonistic. Chromatin immuno-precipitation for GFP (control), BRG1 or the FLAG-tag on the over-expressed BRG1 in the HCC15 shGFP (control) and BRG1 over-expressing cell lines. Real time PCR was performed with primers for indicated genomic sites, $n=3$ biological replicates, * indicates $p<0.03$.

[0044] FIGS. 23A-23G demonstrate that BRG1 and EGFR control the sensitized phenotype. FIGS. 23A and 23B depict graphs of etoposide dose response curves performed in the presence or absence of DZNep in the indicated HCC15 and H460 stably transduced cell lines. The fold change \pm range of etoposide IC_{50} in response to DZNep is plotted, $n=3$ biological replicates, * indicates $p<0.04$, ** $p<0.0001$. FIGS. 23C and 23D depict graphs for the indicated HCC15 and H460 stably transduced etoposide treated cell lines, where 7AAD flow cytometry was used to assess changes in S phase percentage in response to DZNep, $n=3$ biological replicates, * indicates $p=0.02$, ** $p<0.001$. FIGS. 23E and 23F depict graphs of the quantification of the percentage of anaphase cells that showed bridges in untreated or DZNep treated cell lines, $n=3$ biological replicates of 11-25 anaphase nuclei, * indicates $p<0.04$, ** $p<0.02$. FIG. 23G depicts a model of the interaction of EZH2 with BRG1 and EGFR.

[0045] FIGS. 24A-24C demonstrate the effects of EZH2 inhibition. FIG. 24A depicts the results of a Western blot. Western Blot was performed on whole cell extracts from indicated lines for EZH2 and its catalytic mark H3K27me3,

total Histone H3 is shown as loading control. CR indicates a coding region targeting hairpin. FIG. 24B depicts a graph of growth. The indicated stably transduced cell lines were plated at equal density. On the 4th day of culture, cell viability was measured using CellTiter-Glo and the percent survival relative to the shGFP line was calculated, $n=3-4$ biological replicates, * indicates $p<0.03$, ** $p<0.002$; y-axis begins at 60% to more clearly show the differences between cell lines. FIG. 24C depicts a graph of Real time RT-PCR for EZH2 mRNA in the indicated cell lines after plating at equal density and treating for 4 days with indicated treatments. Each cell line is normalized to its shGFP control.

[0046] FIGS. 25A-25C demonstrate the effect of combination treatment using DZNep. FIG. 25A is a Waterfall plot of cell line growth in 1 μ M DZNep for 4 days compared to untreated cells, $n=3-5$ biological replicates, * indicates $p<0.04$, ** $p<0.007$. FIG. 25B is a graph of average etoposide IC_{50} for two classes of cell lines, sensitized to etoposide by EZH2 inhibition, and protected from etoposide by EZH2 inhibition, in the presence or absence of DZNep, * indicates $p=0.032$. FIG. 25C is a pair of representative etoposide dose response curves performed in the presence of 5 μ M cisplatin, the sensitized line PC9 (4.2-fold decrease, $p<0.0001$) and the protected line H23 (3.4-fold increase, $p<0.0001$) are shown.

[0047] FIGS. 26A-26C demonstrate the effect of EZH2 on cell cycle. FIG. 26A depicts a graph of average change in % of S phase between DZNep treated cultures and vehicle treated is plotted, * indicates $p=0.0001$. FIG. 26B depicts a graph of the average change in % of S phase in shEZH2 cultures and shGFP cultures is plotted, * indicates $p=0.0006$. FIG. 26C depicts graphs of 7-AAD cell cycle flow cytometry on cultures corresponding to the matched controls for each experiment shown in FIGS. 3A. The average % S phase \pm s.e.m. of each culture is plotted, $n=3-4$ biological replicates.

[0048] FIG. 27A depicts a graph of averaged EZH2 probe (203358_s_at) intensity within each NSCLC sample of the genotype indicated. FIG. 27B depicts a graph of gefitinib dose response curves performed on the indicated cell lines after 4 days of treatment. At 1 μ M gefitinib, the relative survival of the 3 BRG1 mutant cell lines was compared to that of the 3 BRG1 wild-type (WT) cell lines, $p=0.01$. The EGFR mutant cell line PC9 is shown for comparison.

[0049] FIG. 28A-28C demonstrates the interaction of EZH2 and BRG1 and EGFR. FIG. 28A depicts graphs of real time RT-PCR for BRG1 and EGFR expression in the various indicated treated transduced cell lines. FIG. 28B depicts graphs of cell survival of the indicated stably transduced cell lines were plated at equal density. On the 4th day of culture, cell viability was measured using CellTiter-Glo and the percent survival relative to the shGFP line was calculated, $n=3$ biological replicates, * indicates $p<0.05$, ** $p<0.008$. FIG. 28C depicts graphs of sub-G1 fractions of the indicated 4-day cultures as assessed during 7AAD cell cycle flow cytometry analysis. Critically, for these assays the supernatant of each culture was retained and combined with the trypsinized adherent cells to accurately reflect the total amount of apoptosis/necrosis in each culture, $n=3$ biological replicates, * indicates $p=0.03$.

[0050] FIG. 29 depicts a table of the mutational status of the cell lines used in the Examples.

DETAILED DESCRIPTION

[0051] One route of improving cancer treatment is to seek to combine therapeutics, which target different molecules or pathways within a cancerous cell. Occasionally, such combination therapies can produce additive or even synergistic effects. As described in the Examples herein, the inventors were examining combination therapies involving EZH2 inhibition and found that while in some cases a combination therapy performs better than administration of the chemotherapeutic alone, in other cases, the combination of an inhibitor of EZH2 and a chemotherapeutic actually increased the cancer's resistance to the chemotherapeutic, worsening the outcome. As further described herein, the inventors have determined the molecular and genetic basis for these divergent resistant/susceptible phenotypes.

[0052] The inventors' discoveries regarding the effect of EZH2 inhibition on different populations of cancer cells as regards their responsiveness to chemotherapy permits the identification and treatment of subjects who can benefit from the described EZH2 inhibition combination therapies. Conversely, patients who should not receive such combination therapies can also be identified, permitting practitioners to avoid administering therapies that would be ineffective or actually worsen a cancer. Described herein are methods of treatment, assays, and systems related to these discoveries.

[0053] As described in the Examples herein, the inventors have determined that sensitizing mutations in BRG1, EGFR, or B-RAF result in an "EZH2 inhibitor sensitizing" phenotype, i.e. inhibiting EZH2 in such cells will sensitize them to the action of a chemotherapeutic. When cells that do not comprise such sensitizing mutations are contacted with an EZH2 inhibitor, they will either experience no change in their response to a chemotherapeutic or become protected from the chemotherapeutic. Accordingly, in one aspect, the technology described herein relates to a method of treating cancer, the method comprising: administering a chemotherapeutic agent and an EZH2 inhibitor to a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell. In some embodiments, the subject can be determined to have more than one sensitizing mutation, e.g. a sensitizing mutation in BRG1 and a sensitizing mutation in EGFR, or two sensitizing mutations in BRG1. In some embodiments, the technology described herein relates to a method of treating cancer, the method comprising: administering a chemotherapeutic agent to a subject determined not to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell, and wherein the subject is not administered an EZH2 inhibitor.

[0054] As used herein, "brahma-related gene 1" or "BRG1" (e.g. NCBI Gene ID: 6597) refers to a member of the SWI/SNF family of proteins which is similar to the brahma protein of *Drosophila*. Members of this family have helicase and ATPase activities and are thought to regulate transcription of certain genes by altering the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent chromatin remodeling complex SNF/SWI, which is required for transcriptional activation of genes normally repressed by chromatin. In addition, this protein can bind BRCA1, as well as regulate the expression of the tumorigenic protein CD44. Multiple isoforms of BRG1 are known. The sequences of BRG1 isoforms, genes, and transcripts are known in a variety of species, e.g. human BRG1 isoform 1 mRNA (e.g. SEQ ID NO: 004; NCBI Ref

Seq: NM_001128844) and amino acid (e.g. SEQ ID NO: 003; NCBI Ref Seq: NP_001122316) sequences.

[0055] As used herein, "epidermal growth factor receptor" or "EGFR" (e.g. NCBI Gene ID: 1956) refers to a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Multiple isoforms of EGFR are known. The sequences of EGFR isoforms, genes, and transcripts are known in a variety of species, e.g. human EGFR isoform 1 mRNA (e.g. SEQ ID NO: 006; NCBI Ref Seq: NM_005228) and amino acid (e.g. SEQ ID NO: 005; NCBI Ref Seq: NP_005219) sequences.

[0056] As used herein, "v-raf murine sarcoma viral oncogene homolog B1" or "B-RAF" (e.g. NCBI Gene ID: 6635) refers to a member of the Raf kinase family of serine/threonine-specific protein kinases. This protein plays a role in regulating the MAP kinase/ERKs signaling pathway. The sequences of B-RAF genes, transcripts, and polypeptides are known in a variety of species, e.g. human B-RAF mRNA (e.g. SEQ ID NO: 008; NCBI Ref Seq: NM_004333) and polypeptide (e.g. SEQ ID NO: 009; NCBI Ref Seq: NP_004324) sequences.

[0057] As referred to herein, a "sensitizing mutation" is a mutation which causes, at least in part, or is found associated with, a phenotype in which a cell carrying the mutation, when contacted with an agent that causes EZH2 inhibition will thereafter be more sensitive to the action of a chemotherapeutic agent, i.e. the growth of the cell will be reduced after being contacted with a chemotherapeutic agent or be more likely to die after being contacted with a chemotherapeutic agent. Described herein are sensitizing mutations in BRG1, EGFR, and B-RAF. Exemplary sensitizing mutations of BRG1 include, but are not limited to a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; and/or mutations corresponding to P270*; Q729*; W764R; T58*; and/or M272* relative to SEQ ID NO: 003. In some embodiments, the sensitizing mutation of BRG1 can be a mutation corresponding to P270*; Q729*; W764R; T58*; and/or M272* relative to SEQ ID NO: 003.

[0058] Exemplary sensitizing mutations of EGFR include, but are not limited to, a mutation which increases the expression level of EGFR; and/or a mutation corresponding to E746_A750del; E746_A749del; T790M; and/or L858R relative to SEQ ID NO: 005. In some embodiments, the sensitizing mutation of EGFR can be E746_A750del. In some embodiments, the sensitizing mutation of EGFR can be a mutation which increases the expression level of EGFR; and/or a mutation corresponding to E746_A750del; E746_A749del; T790M relative to SEQ ID NO: 005; and/or L858R. In some embodiments, the sensitizing mutation of EGFR can be a mutation corresponding to E746_A750del; E746_A749del; and/or T790M relative to SEQ ID NO: 005. In some embodiments, the sensitizing mutation of EGFR can be a mutation which deletes exon 19 of EGFR, e.g. a deletion of any or all of the nucleotides corresponding to nucleotides 2530-2715 of SEQ ID NO: 6.

[0059] Exemplary sensitizing mutations of B-RAF include, but are not limited to mutations corresponding to G496A and/or L597V relative to SEQ ID NO: 007.

[0060] In some embodiments, a cell can comprise a single sensitizing mutation. In some embodiments, a cell can

comprise two or more sensitizing mutations e.g. two sensitizing mutations, three sensitizing mutations, four sensitizing mutations, or more sensitizing mutations. Multiple sensitizing mutations can be present in the same gene, e.g. two sensitizing mutations in EGFR or present in two or more genes, e.g. sensitizing mutations in B-RAF and EGFR. Exemplary examples of combinations of sensitizing mutations are shown in Table 6.

TABLE A

	EZH2a	EZH2b	EZH2c	EZH2d	EZH2e
EZH2a	X	X	X	X	X
EZH2b	X	X	X	X	X
EZH2c	X	X	X	X	X

TABLE 6

Exemplary combinations of sensitizing mutations												
Additional possible mutations (not mutually exclusive)												
First sensitizing mutation		EGFR				B-RAF		BRG1				
		E746_A750 del	E746_A749dc②	T790M	L858R	G496A	L597V	P270*	Q729*	W764R	T58*	M272*
EGFR	E746_A750 del	X	X	X	X	X	X	X	X	X	X	X
	E746_A749del	X	X	X	X	X	X	X	X	X	X	X
	T790M	X	X	X	X	X	X	X	X	X	X	X
	L858R	X	X	X	X	X	X	X	X	X	X	X
B-RAF	G496A	X	X	X	X	X	X	X	X	X	X	X
	L597V	X	X	X	X	X	X	X	X	X	X	X
BRG1	P270*	X	X	X	X	X	X	X	X	X	X	X
	Q729*	X	X	X	X	X	X	X	X	X	X	X
	W764R	X	X	X	X	X	X	X	X	X	X	X
	T58*	X	X	X	X	X	X	X	X	X	X	X
	M272*	X	X	X	X	X	X	X	X	X	X	X

② indicates text missing or illegible when filed

[0061] As used herein, “enhancer of zeste homolog 2” or “EZH2” (e.g. NCBI Gene ID: 2146) refers to Polycomb-group enzyme which acts as a transcriptional silencer by adding 3 methyl groups to Lysine 27 of histone 3. This methylation leads to condensation of chromosomes. Multiple isoforms of EZH2 are known to exist, e.g. five different isoforms are known in humans. The sequences of EZH2 isoforms, genes, and transcripts are known in a variety of species, e.g. human (NCBI Gene ID NO: 2146: isoform c (protein: SEQ ID NO: 10 (NCBI Ref Seq: NP_001190176); mRNA: SEQ ID NO: 9 (NCBI Ref Seq: NM_001203247); or isoform a (protein: SEQ ID NO: 1 (NCBI Ref Seq: NP_004447); mRNA: SEQ ID NO: 2 (NCBI Ref Seq: NM_004456)). In humans, five isoforms of EZH2, isoforms, namely EZH2a; EZH2b; EZH2c; EZH2d; EZH2e, are known to exist, with isoform c being the form most prevalently expressed. In some aspects of all the embodiments of the invention, the term “EZH2” refers to any of the five isoforms of EZH2. In some aspects of all the embodiments of the invention the EZH2 is EZH2a. In some aspects of all the embodiments of the invention the EZH2 is EZH2b. In some aspects of all the embodiments of the invention the EZH2 is EZH2c. In some aspects of all the embodiments of the invention the EZH2 is EZH2d. In some aspects of all the embodiments of the invention the EZH2 is EZH2e.

[0062] The inhibition in the methods of the invention can relate to one specific isoform or a combination thereof Table A sets forth the different combinations of the isoforms that are contemplated to be inhibited in the methods of the invention. For example, in some aspects one can inhibit all or any combination of the EZH2 isoforms.

TABLE A-continued

	EZH2a	EZH2b	EZH2c	EZH2d	EZH2e
EZH2d	X	X	X	X	X
EZH2e	X	X	X	X	X

[0063] In some embodiments, the methods described herein relate to the treatment of cancer, wherein the treatment comprises administering to the human subject an EZH2 inhibitor if the human subject has been determined to be sensitive to EZH2 inhibitors. As used herein, the term “inhibitor of EZH2” or “EZH2 inhibitor” refers to an agent, which reduces the expression and/or activity of EZH2. The reduction in expression and/or activity can be, for example, by at least about 10%, e.g. by 10% or more, 20% or more, 30% or more, 50% or more, 75% or more, 90% or more, 95% or more, 98% or more, or 99% or more. The reduction of the expression and/or activity of EZH2 is not intended to encompass complete inhibition or total reduction of EZH2 expression and/or activity.


[0064] In some aspects of all the embodiments, the reduction of EZH2 expression and/or activity can be a “complete reduction” or “complete inhibition”, i.e. a 100% reduction of the expression and/or activity of EZH2. In some aspects of all the EZH2 inhibitor can be an agent that inhibits the expression level, e.g. mRNA and/or polypeptide expression product level of EZH2. In some aspects of all the embodiments, an EZH2 inhibitor can be an agent that inhibits the activity of EZH2.

[0065] EZH2 is known to interact with SUZ12, ATRX, HDAC1, Histone deacetylase 2, VAV1, EED (4 possible isoforms), RBBP4, RBBP7, PHF1, MTF2, PHF19, JARID1A, JARID1B, DNMT1, DNMT3A, DNMT3B,

HDAC1, HDAC2. EZH2 polypeptide can be located within a multiple polypeptide complex, e.g. PRC2, PRC3, and/or PRC4 (for further discussion of the PRC complexes, see, e.g. Kuzmichev et al. *Molecular Cell* 2004 14:183-193; Simon and Lange. *Mutation Research* 2008 647:21-9; Simon and Kingston. *Mol Cell Biol* 2009 10:697-708; and Sauvageau and Sauvageau. *Cell Stem Cell* 2010 7:299-313; each of which is incorporated by reference herein in its entirety). An inhibitor of EZH2 can inhibit the EZH2 function or reduce the EZH2 amount which is present in any of these complexes.

[0066] In some instances, the function of EZH2 can be replicated by EZH1 (see, e.g. Ho and Crabtree. *Cell Stem Cell* 2008 3:577-8; which is incorporated by reference herein in its entirety). Accordingly, in some aspects of all the embodiments, an EZH2 inhibitor inhibits the EZH2-like activity of EZH1.

[0067] The term “agent” refers generally to any entity which is normally not present or not present at the levels being administered to a cell, tissue or subject. An agent can be selected from a group comprising: polynucleotides; polypeptides; small molecules; antibodies; or functional fragments thereof. Exemplary, but non-limiting examples of EZH2 inhibitors can an inhibitory nucleic acid; DZNep (CAS 102052-95-9); S-adenosyl-L-homocysteine. EZH2 inhibitors are commercially available, e.g. DZNep (Cat No: 13828, Cayman Chemical; Ann Arbor, Mich.) Additional examples of EZH2 inhibitors and methods of producing them are described in in U.S. Patent Publications 2012/0071418 (see, e.g. compound 75), which is incorporated by reference herein in its entirety; 2011/0251216 which is incorporated by reference herein in its entirety; 2009/0012031 which is incorporated by reference herein in its entirety; isoliquiritigenin; inhibitors as described in International Patent Publications WO/05/034845 which is incorporated by reference herein in its entirety; WO/12/068589 (see, e.g. paragraphs [0037]-[105], particularly Table 1), which is incorporated by reference herein in its entirety; WO/12/075080 (see, e.g. page 2, line 20—page 4 line 33) which is incorporated by reference herein in its entirety; WO/11/140324 (see, e.g. page 2, line 20—page 4, line 29; pages 37-79) which is incorporated by reference herein in its entirety; WO/12/005805 (see, e.g. page 2, line 20—page 3, line 21; the structures shown in Examples 1-125), which is incorporated by reference herein in its entirety; WO/11/140325 (see, e.g. page 3, line 1—page 5, line 6; the structures shown in Examples 1-131), which is incorporated by reference herein in its entirety; and including nucleic acid inhibitors as described in US Patent Publications 2005/0159382, which is incorporated by reference herein in its entirety; and 2011/0286990 which is incorporated by reference herein in its entirety. EZH2 inhibitors and methods of screening for EZH2 inhibitors are also described, e.g. in Diaz et al. *J Biomol Screen* 2012; which is incorporated by reference herein in its entirety. In some embodiments, an EZH2 inhibitor can be GSK126, e.g. an agent with the structure of formula I.



[0068] In some embodiments, the technology described herein relates to the treatment of cancer, comprising at least in part, administering a drug that inhibits EZH2 in combination with one or more chemotherapeutic agent. As used herein the term “chemotherapeutic agent” refers to any chemical or biological agent with therapeutic usefulness in the treatment of diseases characterized by abnormal cell growth. Such diseases include tumors, neoplasms and cancer as well as diseases characterized by hyperplastic growth. These agents can function to inhibit a cellular activity upon which the cancer cell depends for continued proliferation. In some aspect of all the embodiments, a chemotherapeutic agent is a cell cycle inhibitor or a cell division inhibitor. Categories of chemotherapeutic agents that are useful in the methods of the invention include alkylating/alkaloid agents, antimetabolites, hormones or hormone analogs, and miscellaneous antineoplastic drugs. Most of these agents are directly or indirectly toxic to cancer cells. In one embodiment, a chemotherapeutic agent is a radioactive molecule. One of skill in the art can readily identify a chemotherapeutic agent of use (e.g. see Slapak and Kufe, *Principles of Cancer Therapy*, Chapter 86 in Harrison’s *Principles of Internal Medicine*, 14th edition; Perry et al., *Chemotherapy*, Ch. 17 in Abeloff, *Clinical Oncology* 2nd ed. 2000 Churchill Livingstone, Inc; Baltzer L, Berkery R (eds): *Oncology Pocket Guide to Chemotherapy*, 2nd ed. St. Louis, Mosby-Year Book, 1995; Fischer D S, Knobf M F, Durivage H J (eds): *The Cancer Chemotherapy Handbook*, 4th ed. St. Louis, Mosby-Year Book, 1993). In some embodiments, the chemotherapeutic agent can be a cytotoxic chemotherapeutic. The term “cytotoxic agent” as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g. At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32 and radioactive isotopes of Lu), chemotherapeutic agents, and toxins, such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof. Exemplary chemotherapeutic agents can include, but are not limited to, a topoisomerase inhibitor; a topoisomerase I inhibitor; a topoisomerase II inhibitor; a PARP inhibitor; a CDK1 inhibitor; and an EGFR inhibitor.

[0069] As used herein, a “topoisomerase inhibitor” refers to an agent that can reduce the level and/or activity of a topoisomerase, e.g. topoisomerase I and/or topoisomerase

II. A topoisomerase inhibitor can be specific for either topoisomerase I or topoisomerase II; or inhibit both topoisomerase I and II. Topoisomerase I inhibitors are known in the art and include, by way of non-limiting example, camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D. Topoisomerase II inhibitors are known in the art and include, by way of non-limiting example, doxorubicin (e.g. ADRIAMYCIN™); etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331.

[0070] In some embodiments, a chemotherapeutic agent can be a targeted therapeutic, i.e. an agent that prevents proliferation of only a subset of cells, e.g. cells that have high levels of EGFR signaling. Exemplary examples of targeted therapies include, but are not limited to, a PARP inhibitor; a CDK1 inhibitor; and an EGFR inhibitor.

[0071] As used herein, a “PARP inhibitor” refers to an agent that can reduce the level and/or activity of poly ADP ribose polymerase (PARP). PARP inhibitors are known in the art and include, by way of non-limiting example, olaparib, veliparib, iniparib, rucaparib, AG14361, INO-1001, A-966492, BMN-673, CEP 9722; 3-aminobenzamide; and MK-4827. PARP inhibitors are commercially available, e.g. olaparib (Cat No. S1060 SelleckBio; Houston, Tex.).

[0072] As used herein, a “CDK1 inhibitor” refers to an agent that can reduce the level and/or activity of cyclin-dependent kinase 1 (CDK1). CDK1 inhibitors are known in the art and include, by way of non-limiting example, PD 0332991, flavopiridol, roscovitine, SNS-032, AT7519, JNJ-7706621, PHA-793887, dinaciclib, BMS-265246, PHA-848125, PHA-767491, SCH 900776, R547, A-674563, AZD5438, BS-181 HCl, RO-3306, CGP754514A, and NU6102. CDK1 inhibitors are commercially available, e.g. flavopiridol (Cat No. S1230 SelleckBio; Houston, Tex.).

[0073] As used herein, an “EGFR inhibitor” refers to an agent that can reduce the level and/or activity of epidermal growth factor receptor (EGFR). EGFR inhibitors are known in the art and include, by way of non-limiting example, erlotinib, gefitinib, lapatinib, afatinib, CI-1033, neratinib, CP-724714, TAK-285, AST-1306, ARRY334543, Arry-380, AG-1478, dacomitinib, desmethyl erlotinib, OSI-420, AZD8931, AEE788, pelitinib, CUDC-101, WS8040, WZ4002, WZ3146, AG-490, XL647, PD153035 HCl, BMS-599626, AG 825, AG 494, AG 555, BIBU 1361, BIBX 1382, DAPH, JNJ 28871063, lavendustin A, hypericin, etc. EGFR inhibitors are commercially available, e.g. erlotinib (Cat No. S1023 SelleckBio; Houston, Tex.).

[0074] In some embodiments, the technology described herein relates to the treatment of cancer, at least in part, by administering an EZH2 inhibitor in combination with one or more chemotherapeutic agents. In some embodiments, an EZH2 inhibitor and a second chemotherapeutic agent can be administered. In some embodiments, an EZH2 inhibitor and two or more chemotherapeutic agents can be administered, e.g. two chemotherapeutic agents can be administered, three chemotherapeutic agents can be administered, or four chemotherapeutic agents can be administered in addition to an EZH2 inhibitor. In some embodiments, multiple agents belonging to the same class of agents can be administered, e.g. two EGFR inhibitors, two EZH2 inhibitors, or two CDK1 inhibitors can be administered. In some embodiments, multiple agents belonging to different classes of agents can be administered, e.g. an EZH2 inhibitor, a CDK1

inhibitor, and a topoisomerase inhibitor can be administered. In some embodiments of any of the aspects described herein, a subject can be administered an EZH2 inhibitor, a CDK1 inhibitor, and a further chemotherapeutic agent. In some embodiments, the further chemotherapeutic agent can include, but is not limited to, a topoisomerase inhibitor; a topoisomerase I inhibitor; a topoisomerase II inhibitor; a PARP inhibitor; and an EGFR inhibitor.

[0075] In one aspect, provided herein is a method of identifying a subject who is a candidate for treatment of cancer, the method comprising: detecting a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject; wherein the treatment comprises the administration of a chemotherapeutic agent and an EZH2 inhibitor. In one aspect, provided herein is a method of identifying a subject who is a candidate for treatment of cancer, the method comprising: detecting the absence of a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject; wherein the treatment comprises the administration of a chemotherapeutic agent and does not comprise the administration of an EZH2 inhibitor. In one aspect, provided herein is a method of determining whether a subject is likely to respond to a treatment for cancer, the method comprising: detecting the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject; wherein the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF indicates the subject has an increased likelihood of responding to a treatment to cancer; wherein the treatment for cancer comprises administration of an EZH2 inhibitor and a chemotherapeutic agent. In one aspect, provided herein is a method of determining whether a subject is likely to respond to a treatment for cancer, the method comprising: detecting the absence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject; wherein the absence of a sensitizing mutation in BRG1, EGFR, or B-RAF indicates the subject has an increased likelihood of responding to a treatment to cancer; wherein the treatment for cancer comprises administration of a chemotherapeutic agent and does not comprise the administration of an EZH2 inhibitor.

[0076] In one aspect, provided herein is an assay comprising: subjecting a tumor cell sample from a subject to at least one analysis to detect the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF; wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with an EZH2 inhibitor and a chemotherapeutic agent. In some embodiments, the presence of the mutation can be determined using an assay selected from the group consisting of: hybridization; sequencing; exome capture; PCR; high-throughput sequencing; allele-specific probe hybridization; allele-specific primer extension, allele-specific amplification; 5' nuclease digestion; molecular beacon assay; oligonucleotide ligation assay; size analysis; single-stranded conformation polymorphism; real-time quantitative PCR, and any combinations thereof. In some embodiments, the absence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with a chemotherapeutic agent in the absence of an EZH2 inhibitor.

[0077] In some embodiments, the presence and/or absence of a sensitizing mutation, e.g. in BRG1, EGFR, and/or B-RAF can be detected by determining the sequence of a

genomic locus and/or an mRNA transcript. Such molecules can be isolated, derived, or amplified from a biological sample, such as a tumor sample. Nucleic acid (e.g. DNA) and ribonucleic acid (RNA) molecules can be isolated from a particular biological sample using any of a number of procedures, which are well-known in the art, the particular isolation procedure chosen being appropriate for the particular biological sample. For example, freeze-thaw and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from solid materials; and proteinase K extraction can be used to obtain nucleic acid from blood (Roiff, A et al. PCR: Clinical Diagnostics and Research, Springer (1994)).

[0078] In some embodiments, the nucleic acid sequence of a target gene (e.g. BRG1, EGFR, and/or B-RAF) in a sample obtained from a subject can be determined and compared to a reference sequence to determine if a sensitizing mutation is present in the subject. In some embodiments, the sequence of the target gene can be determined by sequence the target gene (e.g. the genomic sequence and/or the mRNA transcript thereof). Methods of sequencing a nucleic acid sequence are well known in the art. Briefly, a sample obtained from a subject can be contacted with one or more primers which specifically hybridize to a single-strand nucleic acid sequence flanking the target gene sequence and a complementary strand is synthesized. In some next-generation technologies, an adaptor (double or single-stranded) is ligated to nucleic acid molecules in the sample and synthesis proceeds from the adaptor or adaptor compatible primers. In some third-generation technologies, the sequence can be determined, e.g. by determining the location and pattern of the hybridization of probes, or measuring one or more characteristics of a single molecule as it passes through a sensor (e.g. the modulation of an electrical field as a nucleic acid molecule passes through a nanopore). Exemplary methods of sequencing include, but are not limited to, Sanger sequencing, dideoxy chain termination, high-throughput sequencing, next generation sequencing, 454 sequencing, SOLiD sequencing, polony sequencing, Illumina sequencing, Ion Torrent sequencing, sequencing by hybridization, nanopore sequencing, Helioscope sequencing, single molecule real time sequencing, RNAP sequencing, and the like. Methods and protocols for performing these sequencing methods are known in the art, see, e.g. "Next Generation Genome Sequencing" Ed. Michal Janitz, Wiley-VCH; "High-Throughput Next Generation Sequencing" Eds. Kwon and Ricke, Humanna Press, 2011; and Sambrook et al., Molecular Cloning: A Laboratory Manual (3 ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA (2001); which are incorporated by reference herein in their entireties.

[0079] In some embodiments, sequencing can comprise exome sequencing (i.e. targeted exome capture). Exome sequencing comprises enriching for an exome(s) of interest and then sequencing the nucleic acids comprised by the enriched sample. Sequencing can be according to any method known in the art, e.g. those described above herein. Methods of enrichment can include, e.g. PCR, molecular inversion probes, hybrid capture, and in solution capture. Exome capture methodologies are well known in the art, see, e.g. Sulonen et al. Genome Biology 2011 12:R94; and Teer and Mullikin. Hum Mol Genet 2010 19:R2; which are incorporated by reference herein in their entireties. Kits for performing exome capture are available commercially, e.g.

the TRUSEQ™ Exome Enrichment Kit (Cat. No. FC-121-1008; Illumina, San Diego, Calif.). Exome capture methods can also readily be adapted by one of skill in the art to enrich specific exomes of interest.

[0080] In some embodiments, the presence of a sensitizing mutation can be determined using a probe that is specific for the sensitizing mutation. In some embodiments, the probe can be detectably labeled. In some embodiments, a detectable signal can be generated by the probe when a sensitizing mutation is present.

[0081] In some embodiments, the probe specific for the sensitizing mutation can be a probe in a hybridization assay, i.e. the probe can specifically hybridize to a nucleic acid comprising a sensitizing mutation (as opposed to a wild-type nucleic acid sequence) and the hybridization can be detected, e.g. by having the probe and/or the target nucleic acid be detectably labeled. Hybridization assays are well known in the art and include, e.g. northern blots and Southern blots.

[0082] In some embodiments, the probe specific for the sensitizing mutation can be a probe in a PCR assay, i.e. a primer. In general, the PCR procedure describes a method of gene amplification which is comprised of (i) sequence-specific hybridization of primers to specific genes within a nucleic acid sample or library, (ii) subsequent amplification involving multiple rounds of annealing, elongation, and denaturation using a thermostable DNA polymerase, and optionally, (iii) screening the PCR products for a band or product of the correct size. The primers used are oligonucleotides of sufficient length and appropriate sequence to provide initiation of polymerization, i.e. each primer is specifically designed to be complementary to a strand of the genomic locus to be amplified. In an alternative embodiment, the presence of a sensitizing mutation in an mRNA transcript can be determined by reverse-transcription (RT) PCR and by quantitative RT-PCR (QRT-PCR) or real-time PCR methods. Methods of RT-PCR and QRT-PCR are well known in the art. In some embodiments, the PCR product can be labeled, e.g. the primers can comprise a detectable label, or a label can be incorporated and/or bound to the PCR product, e.g. EtBr detection methods. Other non-limiting detection methods can include the detection of a product by mass spectroscopy or MALDI-TOF.

[0083] The term "label" refers to a composition capable of producing a detectable signal indicative of the presence of an antibody reagent (e.g. a bound antibody reagent). Suitable labels include radioisotopes, nucleotide chromophores, enzymes, substrates, fluorescent molecules, chemiluminescent moieties, magnetic particles, bioluminescent moieties, and the like. As such, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means.

[0084] The nucleic acid sequences of BRG1, B-RAF, and EGFR have been assigned NCBI accession numbers for different species such as human, mouse and rat. In particular, the NCBI accession numbers for the nucleic acid sequences of the human expression products are included herein (SEQ ID NOs: 004, 006, and 008). Accordingly, a skilled artisan can design appropriate primers based on the known sequence for determining the mRNA level of the respective gene.

[0085] A sensitizing mutation will typically be present in the genomic DNA of a tumor (e.g. cancerous) cell. Accordingly, the mutation can be detected in either or both of the

genomic DNA or the mRNA transcripts of a tumor cell. In some embodiments, the sensitizing mutation can occur within a DNA and/or RNA sequence that is translated. Accordingly, in some embodiments, the sensitizing mutation can be detected in the polypeptide of a tumor cell.

[0086] Detection of polypeptides comprising a sensitizing mutation can be according to any method known in the art (e.g. mass spectroscopy, flow cytometry, and/or immunological-based methods). Immunological methods to detect polypeptides comprising a sensitizing mutation in accordance with the present technology include, but are not limited to antibody techniques such as immunohistochemistry, immunocytochemistry, flow cytometry, fluorescent-activated cell sorting (FACS), immunoblotting, radioimmunoassays, western blotting, immunoprecipitation, enzyme-linked immunosorbent assays (ELISA), and derivative techniques that make use of antibody reagents as described herein.

[0087] Immunochemical methods require the use of an antibody reagent specific for the target molecule (e.g. the antigen or in the embodiments described herein, a polypeptide or fragment thereof comprising a sensitizing mutation). In some embodiments, an antibody reagent for determining the presence of a sensitizing mutation in a sample can be an antibody reagent specific for a polypeptide comprising a sensitizing mutation, e.g. a sensitizing mutant of BRG1, EGFR, and/or B-RAF as described herein.

[0088] In some embodiments, the assays, methods, and/or systems described herein can comprise: an anti-sensitizing mutation antibody reagent, i.e. an antibody reagent specific for a polypeptide comprising a sensitizing mutation. In some embodiments, the antibody reagent can be detectably labeled. In some embodiments, the antibody reagent can be attached to a solid support (e.g. bound to a solid support). In some embodiments, the solid support can comprise a particle (including, but not limited to an agarose or latex bead or particle or a magnetic particle), a bead, a nanoparticle, a polymer, a substrate, a slide, a coverslip, a plate, a dish, a well, a membrane, and/or a grating. The solid support can include many different materials including, but not limited to, polymers, plastics, resins, polysaccharides, silicon or silica based materials, carbon, metals, inorganic glasses, and membranes.

[0089] In one embodiment, an assay, method, and/or system as described herein can comprise an ELISA. In an exemplary embodiment, a first antibody reagent can be immobilized on a solid support (usually a polystyrene micro titer plate). The solid support can be contacted with a sample obtained from a subject, and the antibody reagent will bind ("capture") antigens for which it is specific (e.g. a polypeptide comprising a sensitizing mutation). The solid support can then be contacted with a second labeled antibody reagent (e.g. a detection antibody reagent). The detection antibody reagent can, e.g. comprise a detectable signal, be covalently linked to an enzyme, or can itself be detected by a secondary antibody, which is linked to an enzyme through bio-conjugation. The presence of a signal indicates that both the first antibody reagent immobilized on the support and the second "detection" antibody reagent have bound to an antigen, i.e. the presence of a signal indicated the presence of polypeptide comprising a sensitizing mutation. Between each step the plate is typically washed with a mild detergent solution to remove any proteins or antibodies that are not specifically bound. After the final wash step the plate is

developed by adding an enzymatic substrate to produce a visible signal, which indicates the presence of a sensitizing mutation in the sample. Older ELISAs utilize chromogenic substrates, though newer assays employ fluorogenic substrates with much higher sensitivity. There are other different forms of ELISA, which are well known to those skilled in the art. The standard techniques known in the art for ELISA are described in "Methods in Immunodiagnosis", 2nd Edition, Rose and Bigazzi, eds. John Wiley & Sons, 1980; Campbell et al., "Methods and Immunology", W. A. Benjamin, Inc., 1964; and Oellerich, M. 1984, J. Clin. Chem. Clin. Biochem. 22:895-904. These references are hereby incorporated by reference in their entirety.

[0090] In one embodiment, the assays, systems, and methods described herein can comprise a lateral flow immunoassay test (LFIA), also known as the immunochromatographic assay, or strip test to measure or determine the presence of a polypeptide comprising a sensitizing mutation. LFIA is a simple device intended to detect the presence (or absence) of a target in a sample. There are currently many LFIA tests are used for medical diagnostics either for home testing, point of care testing, or laboratory use. LFIA tests are a form of immunoassay in which the test sample flows along a solid substrate via capillary action. After the sample is applied to the test it encounters a colored antibody reagent, which mixes with the sample, and if bound to a portion of the sample, transits the substrate encountering lines or zones which have been pretreated with a second antibody reagent. Depending upon the presence or absence of the target in the sample the colored antibody reagent can become bound at the test line or zone. LFIA is essentially immunoassays adapted to operate along a single axis to suit the test strip format or a dipstick format. Strip tests are extremely versatile and can be easily modified by one skilled in the art for detecting an enormous range of antigens from fluid samples such as urine, blood, tumor cell lysates etc. Strip tests are also known as dip stick test, the name bearing from the literal action of "dipping" the test strip into a fluid sample to be tested. LFIA strip test are easy to use, require minimum training and can easily be included as components of point-of-care test (POCT) diagnostics to be used on site in the field. LFIA tests can be operated as either competitive or sandwich assays. Sandwich LFIA is similar to sandwich ELISA. The sample first encounters colored particles, which are labeled with antibody reagents specific for a target. The test line will also contain antibody reagents. The test line will show as a colored band in positive samples. In some embodiments, the lateral flow immunoassay can be a double antibody sandwich assay, a competitive assay, a quantitative assay or variations thereof. There are a number of variations on lateral flow technology. It is also possible to apply multiple capture zones to create a multiplex test.

[0091] A typical test strip consists of the following components: (1) sample application area comprising an absorbent pad (i.e. the matrix or material) onto which the test sample is applied; (2) conjugate or reagent pad- this contains antibody reagent(s) specific to the target which can be conjugated to colored particles (usually colloidal gold particles, or latex microspheres); (3) test results area comprising a reaction membrane - typically a hydrophobic nitrocellulose or cellulose acetate membrane onto which antibody reagents are immobilized in a line across the membrane as a capture zone or test line (a control zone may also be present, containing antibodies specific for the antibody

reagents conjugated to the particles or microspheres); and (4) optional wick or waste reservoir—a further absorbent pad designed to draw the sample across the reaction membrane by capillary action and collect it. The components of the strip are usually fixed to an inert backing material and may be presented in a simple dipstick format or within a plastic casing with a sample port and reaction window showing the capture and control zones. While not strictly necessary, most tests will incorporate a second line, which contains an antibody that picks up free latex/gold in order to confirm the test has operated correctly.

[0092] The use of “dip sticks” or LFIA test strips and other solid supports have been described in the art in the context of an immunoassay for a number of antigen biomarkers. U.S. Pat. Nos. 4,943,522; 6,485,982; 6,187,598; 5,770,460; 5,622,871; 6,565,808, U.S. patent applications Ser. No. 10/278,676; U.S. Ser. No. 09/579,673 and U.S. Ser. No. 10/717,082, which are incorporated herein by reference in their entirety, are non-limiting examples of such lateral flow test devices. Three U.S. patents (U.S. Pat. No. 4,444,880, issued to H. Tom; U.S. Pat. No. 4,305,924, issued to R. N. Piasio; and U.S. Pat. No. 4,135,884, issued to J. T. Shen) describe the use of “dip stick” technology to detect soluble antigens via immunochemical assays. The apparatuses and methods of these three patents broadly describe a first component fixed to a solid surface on a “dip stick” which is exposed to a solution containing a soluble antigen that binds to the component fixed upon the “dip stick,” prior to detection of the component-antigen complex upon the stick. It is within the skill of one in the art to modify the teaching of these “dip stick” technology for the detection of a sensitizing mutation.

[0093] Immunochemistry is a family of techniques based on the use of a specific antibody, wherein antibodies are used to specifically target molecules inside or on the surface of cells. In some embodiments, immunohistochemistry (“IHC”) and immunocytochemistry (“ICC”) techniques can be used to detect the presence of a sensitizing mutation. IHC is the application of immunochemistry to tissue sections, whereas ICC is the application of immunochemistry to cells or tissue imprints after they have undergone specific cytological preparations such as, for example, liquid-based preparations. In some instances, signal amplification may be integrated into the particular protocol, wherein a secondary antibody, that includes a label, follows the application of an antibody reagent specific for a polypeptide comprising a sensitizing mutation. Typically, for immunohistochemistry, tissue obtained from a subject and fixed by a suitable fixing agent such as alcohol, acetone, and paraformaldehyde, is sectioned and reacted with an antibody. Conventional methods for immunohistochemistry are described in Buchwalow and Bocker (Eds) “Immunohistochemistry: Basics and Methods” Springer (2010); Lin and Prichard “Handbook of Practical Immunohistochemistry” Springer (2011); which are incorporated by reference herein in their entireties. In some embodiments, immunocytochemistry may be utilized where, in general, tissue or cells are obtained from a subject are fixed by a suitable fixing agent such as alcohol, acetone, and paraformaldehyde, to which is reacted an antibody. Methods of immunocytological staining of human samples is known to those of skill in the art and described, for example, in Burry. “Immunocytochemistry: A Practical Guide for Biomedical Research” Springer (2009); which is incorporated by reference herein in its entirety.

[0094] In some embodiments, one or more of the antibody reagents described herein can comprise a detectable label and/or comprise the ability to generate a detectable signal (e.g. by catalyzing reaction converting a compound to a detectable product). Detectable labels can comprise, for example, a light-absorbing dye, a fluorescent dye or a radioactive label. Detectable labels, methods of detecting them, and methods of incorporating them into an antibody reagent are well known in the art.

[0095] In some embodiments, detectable labels can include labels that can be detected by spectroscopic, photochemical, biochemical, immunochemical, electromagnetic, radiochemical, or chemical means, such as fluorescence, chemifluorescence, or chemiluminescence, or any other appropriate means. The detectable labels used in the methods described herein can be primary labels (where the label comprises a moiety that is directly detectable or that produces a directly detectable moiety) or secondary labels (where the detectable label binds to another moiety to produce a detectable signal, e.g., as is common in immunological labeling using secondary and tertiary antibodies). The detectable label can be linked by covalent or non-covalent means to the antibody reagent. Alternatively, a detectable label can be linked such as by directly labeling a molecule that achieves binding to the antibody reagent via a ligand-receptor binding pair arrangement or other such specific recognition molecules. Detectable labels can include, but are not limited to radioisotopes, bioluminescent compounds, chromophores, antibodies, chemiluminescent compounds, fluorescent compounds, metal chelates, and enzymes.

[0096] In other embodiments, the detection antibody is label with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wavelength, its presence can then be detected due to fluorescence. In some embodiments, a detectable label can be a fluorescent dye molecule, or fluorophore including, but not limited to fluorescein, phycoerythrin, phycocyanin, o-phthaldehyde, fluorescamine, Cy3TM, Cy5TM, allophycocyanine, Texas Red, peridinin chlorophyll, cyanine, tandem conjugates such as phycoerythrin-Cy5TM, green fluorescent protein, rhodamine, fluorescein isothiocyanate (FITC) and Oregon GreenTM, rhodamine and derivatives (e.g., Texas red and tetra-rhodamine isothiocyanate (TRITC)), biotin, phycoerythrin, AMCA, CyDyesTM, 6-carboxyfluorescein (commonly known by the abbreviations FAM and F), 6-carboxy-2',4',7',4',7'-hexachlorofluorescein (HEX), 6-carboxy-4',5'-dichloro-2',7'-dimethoxyfluorescein (JOE or J), N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA or T), 6-carboxy-X-rhodamine (ROX or R), 5-carboxyrhodamine-6G (R6G5 or G5), 6-carboxyrhodamine-6G (R6G6 or G6), and rhodamine 110; cyanine dyes, e.g. Cy3, Cy5 and Cy7 dyes; coumarins, e.g. umbelliferone; benzimide dyes, e.g. Hoechst 33258; phenanthridine dyes, e.g. Texas Red; ethidium dyes; acridine dyes; carbarole dyes; phenoxazine dyes; porphyrin dyes; polymethine dyes, e.g. cyanine dyes such as Cy3, Cy5, etc.; BODIPY dyes and quinoline dyes. In some embodiments, a detectable label can be a radiolabel including, but not limited to ³H, ¹²⁵I, ³⁵S, ¹⁴C, ³²P, and ³³P. In some embodiments, a detectable label can be an enzyme including, but not limited to horseradish peroxidase and alkaline phosphatase. An enzymatic label can produce, for example, a chemiluminescent signal, a color signal, or a fluorescent signal. Enzymes contemplated for use to detectably label an

antibody reagent include, but are not limited to, malate dehydrogenase, *staphylococcal* nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. In some embodiments, a detectable label is a chemiluminescent label, including, but not limited to lucigenin, luminol, luciferin, isoluminol, therratic acridinium ester, imidazole, acridinium salt and oxalate ester. In some embodiments, a detectable label can be a spectral colorimetric label including, but not limited to colloidal gold or colored glass or plastic (e.g., polystyrene, polypropylene, and latex) beads

[0097] In some embodiments, antibodies can also be labeled with a detectable tag, such as c-Myc, HA, VSV-G, HSV, FLAG, V5, HIS, or biotin. Other detection systems can also be used, for example, a biotin-streptavidin system. In this system, the antibodies immunoreactive (i.e. specific for) with the biomarker of interest is biotinylated. Quantity of biotinylated antibody bound to the biomarker is determined using a streptavidin-peroxidase conjugate and a chromagenic substrate. Such streptavidin peroxidase detection kits are commercially available, e.g. from DAKO; Carpinteria, Calif. An antibody reagent can also be detectably labeled using fluorescence emitting metals such as ¹⁵²Eu, or others of the lanthanide series. These metals can be attached to the antibody reagent using such metal chelating groups as diethylenetriaminepentaacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

[0098] In some embodiments of any of the aspects described herein, the sequence, expression level, and/or mutational status of more than one gene can be determined simultaneously (e.g. a multiplex assay) or in parallel. In some embodiments, the sequence, expression level, and/or mutational status of no more than 20 other genes is determined. In some embodiments, the sequence, expression level, and/or mutational status of no more than 10 other genes is determined.

[0099] In some embodiments, the sequence and/or mutational status of BRG1, EGFR, and B-RAF can be determined simultaneously (e.g. a multiplex assay). Where the sequences of sensitizing mutations and wildtype sequences are known as described herein (both in nucleic acid and polypeptide sequences), methods of creating multiplex assays for both nucleic acid or polypeptide targets are known in the art (e.g. multiplex assays can include, but are not limited to multiplexed RTPCR assays, multiplex nucleic acid hybridization assays, and multiplex immunochemical assays). By way of non-limiting example, primers for detecting the level of, e.g. BRG1 and EGFR expression in a quantitative RTPCR assay can be used to detect the presence of 1) a sensitizing mutation which increases the expression level of EGFR and 2) a sensitizing mutation which decreases the expression of BRG1 in one reaction. Non-limiting examples of such primers include those described in the Examples herein, i.e. BRG1 F AGCGATGACGTCTCTGAGGT (SEQ ID NO: 11); BRG1 R GTACAGGGACAC-CAGCCACT (SEQ ID NO: 12); EGFR F TAACAAGCT-CACGCAGTTGG (SEQ ID NO: 13); and EGFR R GTTGAGGGCAATGAGGACAT (SEQ ID NO: 14).

[0100] By way of non-limiting example, probes for detecting the level of, e.g. BRG1 and EGFR expression in a

microarray assay can be used to detect the presence of 1) a sensitizing mutation which increases the expression level of EGFR and 2) a sensitizing mutation which decreases the expression of BRG1 in one reaction. As but one example, the Human Genome U133A 2.0 Array (Cat No. 900471, Affymetrix, Cleveland, Ohio), or the probes comprised therein can be used to detect the level of, e.g. BRG1 and EGFR expression in a single assay, as described in the Examples herein.

[0101] By way of further non-limiting example, antibodies specific for polypeptides comprising a sensitizing mutation as described herein can be produced and used to detect the presence or absence of multiple sensitizing mutations in a single assay.

[0102] The term "sample" or "test sample" as used herein denotes a sample taken or isolated from a biological organism, e.g., a tumor sample from a subject. Exemplary biological samples include, but are not limited to, a biofluid sample; serum; plasma; urine; saliva; a tumor sample; a tumor biopsy and/or tissue sample etc. The term also includes a mixture of the above-mentioned samples. The term "test sample" also includes untreated or pretreated (or pre-processed) biological samples. In some embodiments, a test sample can comprise cells from subject. In some embodiments, a test sample can be a tumor cell test sample, e.g. the sample can comprise cancerous cells, cells from a tumor, and/or a tumor biopsy.

[0103] The test sample can be obtained by removing a sample of cells from a subject, but can also be accomplished by using previously isolated cells (e.g. isolated at a prior timepoint and isolated by the same or another person). In addition, the test sample can be freshly collected or a previously collected sample.

[0104] In some embodiments, the test sample can be an untreated test sample. As used herein, the phrase "untreated test sample" refers to a test sample that has not had any prior sample pre-treatment except for dilution and/or suspension in a solution. Exemplary methods for treating a test sample include, but are not limited to, centrifugation, filtration, sonication, homogenization, heating, freezing and thawing, and combinations thereof. In some embodiments, the test sample can be a frozen test sample, e.g., a frozen tissue. The frozen sample can be thawed before employing methods, assays and systems described herein. After thawing, a frozen sample can be centrifuged before being subjected to methods, assays and systems described herein. In some embodiments, the test sample is a clarified test sample, for example, by centrifugation and collection of a supernatant comprising the clarified test sample. In some embodiments, a test sample can be a pre-processed test sample, for example, supernatant or filtrate resulting from a treatment selected from the group consisting of centrifugation, filtration, thawing, purification, and any combinations thereof. In some embodiments, the test sample can be treated with a chemical and/or biological reagent. Chemical and/or biological reagents can be employed to protect and/or maintain the stability of the sample, including biomolecules (e.g., nucleic acid and protein) therein, during processing. One exemplary reagent is a protease inhibitor, which is generally used to protect or maintain the stability of protein during processing. The skilled artisan is well aware of methods and processes appropriate for pre-processing of biological samples required for determination of the presence of a sensitizing mutation as described herein.

[0105] In some embodiments, the methods, assays, and systems described herein can further comprise a step of obtaining a test sample from a subject. In some embodiments, the subject can be a human subject.

[0106] In one aspect, provided herein is an assay comprising: contacting a tumor cell sample obtained from a subject having cancer with a nucleic acid probe to detect the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF; and detecting the presence or intensity of a signal which indicates the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF; wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with an EZH2 inhibitor and a chemotherapeutic agent. In some embodiments, the absence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with a chemotherapeutic agent, wherein the treatment does not comprise, or in other words, specifically excludes administering an EZH2 inhibitor to the subject if the subject has been determined to be unresponsive to EZH2 inhibitors using the assays or methods of the invention.

[0107] In any of the aspects and embodiments described herein, the subject can be human. In some embodiments, the subject can be a subject having, at risk of having, or in need of treatment for cancer. In some embodiments, the cancer can be selected from the group consisting of lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers. In some embodiments, the cancer can be lung cancer. In some embodiments, the cancer can be non-small cell lung cancer. In some embodiments, the cancer is not lymphoma.

[0108] In one aspect, provided herein is a method of classifying a tumor as an EZH2 combination treatment responsive tumor, the method comprising: detecting a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell sample; wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the tumor is a EZH2 combination treatment responsive tumor. As used herein, the term “EZH2 combination treatment responsive tumor” refers to a tumor comprising cells that, after being contacted with an effective dose of an EZH2 inhibitor will be more sensitive to at least one chemotherapeutic agent than the same cell would be in the absence of being contacted with the EZH2 inhibitor.

[0109] In any embodiment of the methods and assays described herein, the method or assay can further comprise the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF. In some embodiments, the report can be generated by a non-human machine, e.g. a computer system as described elsewhere herein.

[0110] In some embodiments of various aspects described herein, the combination of an EZH2 inhibitor and a chemotherapeutic agent is indicated for the treatment of cancer (e.g. lung cancer) in patients with one or more of the following genetic variants as determined by an FDA approved diagnostic test: a variant corresponding to P270*; Q729*; W764R; T58*; and/or M272* of SEQ ID NO: 003; a variant corresponding to E746_A750del; E746_A749del; T790M; and/or L858R of SEQ ID NO: 003; and/or a variant

corresponding to G496A and/or L597V of SEQ ID NO: 003 or any combination of the foregoing variants.

[0111] In some embodiments, the methods described herein relate to treating a subject having or diagnosed as having cancer with an EZH2 inhibitor and a chemotherapeutic agent. Subjects having cancer can be identified by a physician using current methods of diagnosing cancer. Symptoms and/or complications of cancer which characterize these conditions and aid in diagnosis are well known in the art and include but are not limited to, growth of a tumor, impaired function of the organ or tissue harboring cancer cells, etc. Tests that may aid in a diagnosis of, e.g. cancer include, but are not limited to, tissue biopsies and histological examination. A family history of cancer, or exposure to risk factors for cancer (e.g. tobacco products, radiation, etc.) can also aid in determining if a subject is likely to have cancer or in making a diagnosis of cancer.

[0112] The compositions and methods described herein can be administered to a subject having or diagnosed as having cancer. In some embodiments, the methods described herein comprise administering an effective amount of compositions described herein, e.g. an EZH2 inhibitor and chemotherapeutic agent to a subject in order to alleviate a symptom of a cancer. As used herein, “alleviating a symptom of a cancer” is ameliorating any condition or symptom associated with the cancer. As compared with an equivalent untreated control, such reduction is by at least 5%, 10%, 20%, 40%, 50%, 60%, 80%, 90%, 95%, 99% or more as measured by any standard technique. A variety of means for administering the compositions described herein to subjects are known to those of skill in the art. Such methods can include, but are not limited to oral, parenteral, intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), pulmonary, cutaneous, topical, injection, or intratumoral administration. Administration can be local or systemic.

[0113] The term “effective amount” as used herein refers to the amount of an EZH2 inhibitor and/or chemotherapeutic agent needed to alleviate at least one or more symptom of the disease or disorder, and relates to a sufficient amount of pharmacological composition to provide the desired effect. The term “therapeutically effective amount” therefore refers to an amount of an EZH2 inhibitor and/or chemotherapeutic agent that is sufficient to effect a particular anti-cancer effect when administered to a typical subject. An effective amount as used herein, in various contexts, would also include an amount sufficient to delay the development of a symptom of the disease, alter the course of a symptom disease (for example but not limited to, slowing the progression of a symptom of the disease), or reverse a symptom of the disease. Thus, it is not generally practicable to specify an exact “effective amount”. However, for any given case, an appropriate “effective amount” can be determined by one of ordinary skill in the art using only routine experimentation.

[0114] Effective amounts, toxicity, and therapeutic efficacy can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dosage can vary depending upon the dosage form employed and the route of administration utilized. The dose ratio between toxic and therapeutic effects is the therapeutic index and can be expressed as the ratio LD50/ED50. Compositions and methods that exhibit large therapeutic indices are preferred. A therapeutically

effective dose can be estimated initially from cell culture assays. Also, a dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of an EZH2 inhibitor and/or chemotherapeutic agent, which achieves a half-maximal inhibition of symptoms) as determined in cell culture, or in an appropriate animal model. Levels in plasma can be measured, for example, by high performance liquid chromatography. The effects of any particular dosage can be monitored by a suitable bioassay, e.g., assay for cell death, among others. The dosage can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment.

[0115] In some embodiments, the technology described herein relates to a pharmaceutical composition comprising an EZH2 inhibitor and/or chemotherapeutic agent as described herein, and optionally a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers and diluents include saline, aqueous buffer solutions, solvents and/or dispersion media. The use of such carriers and diluents is well known in the art. Some non-limiting examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, methylcellulose, ethyl cellulose, microcrystalline cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) lubricating agents, such as magnesium stearate, sodium lauryl sulfate and talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol (PEG); (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; (22) bulking agents, such as polypeptides and amino acids (23) serum component, such as serum albumin, HDL and LDL; (22) C₂-C₁₂ alcohols, such as ethanol; and (23) other non-toxic compatible substances employed in pharmaceutical formulations. Wetting agents, coloring agents, release agents, coating agents, sweetening agents, flavoring agents, perfuming agents, preservative and antioxidants can also be present in the formulation. The terms such as "excipient", "carrier", "pharmaceutically acceptable carrier" or the like are used interchangeably herein. In some embodiments, the carrier inhibits the degradation of the active agent, e.g. an EZH2 inhibitor and/or chemotherapeutic agent as described herein.

[0116] In some embodiments, the pharmaceutical composition comprising an EZH2 inhibitor and/or chemotherapeutic agent as described herein can be a parenteral dose form. Since administration of parenteral dosage forms typically bypasses the patient's natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions. In addition, controlled-release parent-

eral dosage forms can be prepared for administration of a patient, including, but not limited to, administration DUROS®-type dosage forms, and dose-dumping.

[0117] Suitable vehicles that can be used to provide parenteral dosage forms of an EZH2 inhibitor and/or chemotherapeutic agent as disclosed within are well known to those skilled in the art. Examples include, without limitation: sterile water; water for injection USP; saline solution; glucose solution; aqueous vehicles such as but not limited to, sodium chloride injection, Ringer's injection, dextrose injection, dextrose and sodium chloride injection, and lactated Ringer's injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and propylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate. Compounds that alter or modify the solubility of a pharmaceutically acceptable salt of an EZH2 inhibitor and/or chemotherapeutic agent as disclosed herein can also be incorporated into the parenteral dosage forms of the disclosure, including conventional and controlled-release parenteral dosage forms.

[0118] Pharmaceutical compositions comprising an EZH2 inhibitor and/or chemotherapeutic agent can also be formulated to be suitable for oral administration, for example as discrete dosage forms, such as, but not limited to, tablets (including without limitation scored or coated tablets), pills, caplets, capsules, chewable tablets, powder packets, cachets, troches, wafers, aerosol sprays, or liquids, such as but not limited to, syrups, elixirs, solutions or suspensions in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil emulsion. Such compositions contain a predetermined amount of the pharmaceutically acceptable salt of the disclosed compounds, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, Remington: The Science and Practice of Pharmacy, 21st Ed., Lippincott, Williams, and Wilkins, Philadelphia Pa. (2005).

[0119] Conventional dosage forms generally provide rapid or immediate drug release from the formulation. Depending on the pharmacology and pharmacokinetics of the drug, use of conventional dosage forms can lead to wide fluctuations in the concentrations of the drug in a patient's blood and other tissues. These fluctuations can impact a number of parameters, such as dose frequency, onset of action, duration of efficacy, maintenance of therapeutic blood levels, toxicity, side effects, and the like. Advantageously, controlled-release formulations can be used to control a drug's onset of action, duration of action, plasma levels within the therapeutic window, and peak blood levels. In particular, controlled- or extended-release dosage forms or formulations can be used to ensure that the maximum effectiveness of a drug is achieved while minimizing potential adverse effects and safety concerns, which can occur both from under-dosing a drug (i.e., going below the minimum therapeutic levels) as well as exceeding the toxicity level for the drug. In some embodiments, the an EZH2 inhibitor and/or chemotherapeutic agent can be administered in a sustained release formulation.

[0120] Controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled release counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug

substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include: 1) extended activity of the drug; 2) reduced dosage frequency; 3) increased patient compliance; 4) usage of less total drug; 5) reduction in local or systemic side effects; 6) minimization of drug accumulation; 7) reduction in blood level fluctuations; 8) improvement in efficacy of treatment; 9) reduction of potentiation or loss of drug activity; and 10) improvement in speed of control of diseases or conditions. Kim, Cherng-ju, *Controlled Release Dosage Form Design*, 2 (Technomic Publishing, Lancaster, Pa.: 2000).

[0121] Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, ionic strength, osmotic pressure, temperature, enzymes, water, and other physiological conditions or compounds.

[0122] A variety of known controlled- or extended-release dosage forms, formulations, and devices can be adapted for use with the salts and compositions of the disclosure. Examples include, but are not limited to, those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; 4,008,719; 5,674,533; 5,059,595; 5,591,767; 5,120,548; 5,073,543; 5,639,476; 5,354,556; 5,733,566; and 6,365,185 B1; each of which is incorporated herein by reference. These dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydroxypropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems (such as OROS® (Alza Corporation, Mountain View, Calif. USA)), or a combination thereof to provide the desired release profile in varying proportions.

[0123] The methods described herein can further comprise administering a second agent and/or treatment to the subject, e.g., as part of a combinatorial therapy. Non-limiting examples of a second agent and/or treatment can include radiation therapy, surgery, gemcitabine, cisplatin, paclitaxel, carboplatin, bortezomib, AMG479, vorinostat, rituximab, temozolomide, rapamycin, ABT-737, PI-103; alkylating agents such as thiopeta and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolmelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride,

melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammaII and calicheamicin omegaII (see, e.g., Agnew, Chem. Intl. Ed. Engl., 33: 183-186 (1994))); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, caminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-di-azo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitiostane, testosterone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguanzone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, Oreg.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiopeta; taxoids, e.g., TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE® Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Ill.), and TAXOTERE® doxetaxel (Rhône-Poulenc Rorer, Antony, France); chlorambucil; GEMZAR® gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin, oxaliplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE™ vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (Camptosar, CPT-11) (including the treatment regimen of irinotecan with 5-FU and leucovorin); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; combretastatin; leucovorin (LV); oxaliplatin, including the oxaliplatin treatment regimen (FOLFOX); lapatinib (Tykerb™); inhibitors of PKC-alpha, Raf, H-Ras,

EGFR (e.g., erlotinib (Tarceva®)) and VEGF-A that reduce cell proliferation and pharmaceutically acceptable salts, acids or derivatives of any of the above. In addition, the methods of treatment can further include the use of radiation or radiation therapy. In some embodiments, cisplatin can be administered in combination with a topoisomerase inhibitor as described above herein, e.g. cisplatin can be administered in combination with etoposide. Further, the methods of treatment can further include the use of surgical treatments.

[0124] In some embodiments, an EZH2 inhibitor can be administered in any combination with, e.g. concurrently or sequentially, with a chemotherapeutic agent. In some embodiments, an EZH2 inhibitor and a chemotherapeutic agent can be administered in a single composition. In some embodiments, an EZH2 inhibitor and a chemotherapeutic agent can be administered in separate compositions. In embodiments where multiple compositions are administered (e.g. one composition comprising an EZH2 inhibitor and a second composition comprising a chemotherapeutic agent), the compositions can be administered at varying times and/or for varying durations.

[0125] In certain embodiments, an effective dose of a composition comprising an EZH2 inhibitor and/or chemotherapeutic agent as described herein can be administered to a patient once. In certain embodiments, an effective dose of a composition comprising an EZH2 inhibitor and/or chemotherapeutic agent can be administered to a patient repeatedly. For systemic administration, subjects can be administered a therapeutic amount of a composition comprising an EZH2 inhibitor and/or chemotherapeutic agent, such as, e.g. 0.1 mg/kg, 0.5 mg/kg, 1.0 mg/kg, 2.0 mg/kg, 2.5 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 40 mg/kg, 50 mg/kg, or more. A composition comprising an EZH2 inhibitor and/or chemotherapeutic agent can be administered over a period of time, such as over a 5 minute, 10 minute, 15 minute, 20 minute, or 25 minute period. The administration can be repeated, for example, on a regular basis, such as hourly for 3 hours, 6 hours, 12 hours or longer or such as biweekly (i.e., every two weeks) for one month, two months, three months, four months or longer.

[0126] In some embodiments, after an initial treatment regimen, the treatments can be administered on a less frequent basis. For example, after treatment biweekly for three months, treatment can be repeated once per month, for six months or a year or longer. Treatment according to the methods described herein can reduce levels of a marker or symptom of a condition, e.g. tumor size by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80% or at least 90% or more.

[0127] In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered daily. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered daily for at least 3 days, e.g. 3 days, 4 days, 5 days, 6 days, 7 days, or longer. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered daily for 2 weeks or less, e.g. 14 days, 13 days, 12 days, 11 days, 10 days, 9 days, 8 days, or 7 days or less. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered daily for between 1 days and 14 days. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered daily for between 3 days and 10 days. In some embodiments, the EZH2 inhibitor and chemotherapeutic

agent can be administered daily for between 3 days and 10 days. In some embodiments, the EZH2 inhibitor, e.g. DZNep can be administered at a dose of about 0.1 mg/kg/day to 20 mg/kg/day. In some embodiments, the EZH2 inhibitor, e.g. DZNep can be administered at a dose of about 1 mg/kg/day to 10 mg/kg/day. In some embodiments, the EZH2 inhibitor, e.g. DZNep, can be administered at a dose of about 2 mg/kg/day.

[0128] In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered weekly. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered twice per week or more frequently, e.g. twice per week, three times per week, four times per week, five times per week, six times per week, or daily.

[0129] In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered for at least 3 days, e.g. 3 days, 4 days, 5 days, 6 days, 7 days, or longer. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered for 2 weeks or less, e.g. 14 days, 13 days, 12 days, 11 days, 10 days, 9 days, 8 days, or 7 days or less. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered for between 1 days and 14 days. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered for between 3 days and 10 days. In some embodiments, the EZH2 inhibitor and chemotherapeutic agent can be administered for between 3 days and 10 days.

[0130] In some embodiments, the EZH2 inhibitor, e.g. DZNep can be administered at a dose of about 0.1 mg/kg/day to 20 mg/kg/day. In some embodiments, the EZH2 inhibitor, e.g. DZNep can be administered at a dose of about 1 mg/kg/day to 10 mg/kg/day. In some embodiments, the EZH2 inhibitor, e.g. DZNep, can be administered at a dose of about 2 mg/kg/day.

[0131] The dosage of a composition as described herein can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment. With respect to duration and frequency of treatment, it is typical for skilled clinicians to monitor subjects in order to determine when the treatment is providing therapeutic benefit, and to determine whether to increase or decrease dosage, increase or decrease administration frequency, discontinue treatment, resume treatment, or make other alterations to the treatment regimen. The dosing schedule can vary from once a week to daily depending on a number of clinical factors, such as the subject's sensitivity to an EZH2 inhibitor and/or chemotherapeutic agent. The desired dose or amount of activation can be administered at one time or divided into subdoses, e.g., 2-4 subdoses and administered over a period of time, e.g., at appropriate intervals through the day or other appropriate schedule. In some embodiments, administration can be chronic, e.g., one or more doses and/or treatments daily over a period of weeks or months. Examples of dosing and/or treatment schedules are administration daily, twice daily, three times daily or four or more times daily over a period of 1 week, 2 weeks, 3 weeks, 4 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, or 6 months, or more.

[0132] The dosage ranges for the administration of an EZH2 inhibitor and/or chemotherapeutic agent, according to the methods described herein depend upon, for example, the form of an EZH2 inhibitor and/or chemotherapeutic agent, its potency, and the extent to which symptoms, markers, or

indicators of a condition described herein are desired to be reduced, for example the percentage reduction desired for tumor size and/or growth. The dosage should not be so large as to cause adverse side effects. Generally, the dosage will vary with the age, condition, and sex of the patient and can be determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication.

[0133] The efficacy of an EZH2 inhibitor and/or chemotherapeutic agent in, e.g. the treatment of a condition described herein, or to induce a response as described herein (e.g. a decrease in tumor size and/or growth rate) can be determined by the skilled clinician. However, a treatment is considered “effective treatment,” as the term is used herein, if any one or all of the signs or symptoms of a condition described herein are altered in a beneficial manner, other clinically accepted symptoms are improved, or even ameliorated, or a desired response is induced e.g., by at least 10% following treatment according to the methods described herein. Efficacy can be assessed, for example, by measuring a marker, indicator, symptom, and/or the incidence of a condition treated according to the methods described herein or any other measurable parameter appropriate, e.g. tumor growth rate. Efficacy can also be measured by a failure of an individual to worsen as assessed by hospitalization, or need for medical interventions (i.e., progression of the disease is halted). Methods of measuring these indicators are known to those of skill in the art and/or are described herein. Treatment includes any treatment of a disease in an individual or an animal (some non-limiting examples include a human or an animal) and includes: (1) inhibiting the disease, e.g., preventing a worsening of symptoms (e.g. pain or impaired organ function); or (2) relieving the disease, e.g., causing regression of symptoms. An effective amount for the treatment of a disease means that amount which, when administered to a subject in need thereof, is sufficient to result in effective treatment as that term is defined herein, for that disease. Efficacy of an agent can be determined by assessing physical indicators of a condition or desired response, (e.g. tumor size, metastasis, and/or mortality). It is well within the ability of one skilled in the art to monitor efficacy of administration and/or treatment by measuring any one of such parameters, or any combination of parameters. Efficacy can be assessed in animal models of a condition described herein, for example treatment of a mouse model of cancer. When using an experimental animal model, efficacy of treatment is evidenced when a statistically significant change in a marker is observed, e.g. tumor size and/or mortality.

[0134] In vitro and animal model assays are provided herein which allow the assessment of a given dose of an EZH2 inhibitor and/or chemotherapeutic agent. By way of non-limiting example, the effects of a dose of an EZH2 inhibitor and/or chemotherapeutic agent can be assessed by a cytotoxicity assay. A non-limiting example of a protocol for such an assay is as follows: Cell lines are dissociated, counted and plated at 5000 cells per well in flat bottom opaque tissue culture treated 96 well plates (CytoOne). The following day, an EZH2 inhibitor and one or more chemotherapeutic agents can be added to each well. After 4 days, Cell Titer Glo™ (Promega) can be added and luminescence read on a BioTec plate reader to determine relative cell number in each well.

[0135] The efficacy of a given dosage combination can also be assessed in an animal model, e.g. a mouse model of xenotransplanted lung cancer cells. For example, lung cancer cell lines can be injected subcutaneously with 1×10⁶ cells in four spots on flanks of 8- to 16-week-old Foxnlnu (nude) mice (Harlan). An EZH2 inhibitor and one or more chemotherapeutic agents can be administered from day 12 to day 25 post injections. Tumor growth can be measured every other day by caliper.

[0136] In some embodiments of the assays and/or methods described herein, the assay/method comprises or consists essentially of a system for determining (e.g. transforming and measuring) the presence of a sensitizing mutation in BRG1, EGFR, and/or B-RAF as described herein. If the comparison system, which can be a computer implemented system, indicates that one or more sensitizing mutations are present, the subject from which the sample is collected can be identified as, e.g. having an increased likelihood of responding to a treatment comprising the administration of an EZH2 inhibitor and a chemotherapeutic agent. In some embodiments, the sequence of BRG1, EGFR, and/or B-RAF or an expression product thereof can be compared to a reference sequence, e.g. SEQ ID NOS: 003-008. In some embodiments, the absence of a sensitizing mutation in BRG1, EGFR, and/or B-RAF indicates the subject as, e.g. having an increased likelihood of responding to a treatment that does not comprise the administration of an EZH2 inhibitor.

[0137] In one embodiment, provided herein is a system comprising: (a) at least one memory containing at least one computer program adapted to control the operation of the computer system to implement a method that includes (i) a determination module configured to identify and detect the presence of a sensitizing mutation in BRG1, EGFR, and/or B-RAF in a test sample obtained from a subject; (ii) a storage module configured to store output data from the determination module; (iii) a computing module adapted to identify from the output data whether a sensitizing mutation in BRG1, EGFR, and/or B-RAF is present in the test sample obtained from the subject, and (iv) a display module for displaying whether a sensitizing mutation in BRG1, EGFR, and/or B-RAF is present in the test sample, optionally, as compared to a reference sequence (see FIG. 14).

[0138] Embodiments can be described through functional modules, which are defined by computer executable instructions recorded on computer readable media and which cause a computer to perform method steps when executed. The modules are segregated by function for the sake of clarity. However, it should be understood that the modules/systems need not correspond to discreet blocks of code and the described functions can be carried out by the execution of various code portions stored on various media and executed at various times. Furthermore, it should be appreciated that the modules can perform other functions, thus the modules are not limited to having any particular functions or set of functions.

[0139] The computer readable storage media can be any available tangible media that can be accessed by a computer. Computer readable storage media includes volatile and nonvolatile, removable and non-removable tangible media implemented in any method or technology for storage of information such as computer readable instructions, data structures, program modules or other data. Computer readable storage media includes, but is not limited to, RAM

(random access memory), ROM (read only memory), EPROM (erasable programmable read only memory), EEPROM (electrically erasable programmable read only memory), flash memory or other memory technology, CD-ROM (compact disc read only memory), DVDs (digital versatile disks) or other optical storage media, magnetic cassettes, magnetic tape, magnetic disk storage or other magnetic storage media, other types of volatile and non-volatile memory, and any other tangible medium which can be used to store the desired information and which can be accessed by a computer including any suitable combination of the foregoing. Computer-readable storage medium do not include a signal.

[0140] Computer-readable data embodied on one or more computer-readable media may define instructions, for example, as part of one or more programs that, as a result of being executed by a computer, instruct the computer to perform one or more of the functions described herein, and/or various embodiments, variations and combinations thereof. Such instructions may be written in any of a plurality of programming languages, for example, Java, J#, Visual Basic, C, C#, C++, Fortran, Pascal, Eiffel, Basic, COBOL assembly language, and the like, or any of a variety of combinations thereof. The computer-readable media on which such instructions are embodied may reside on one or more of the components of either of a system, or a computer readable storage medium described herein, may be distributed across one or more of such components.

[0141] The computer-readable media may be transportable such that the instructions stored thereon can be loaded onto any computer resource to implement the aspects of the technology discussed herein. In addition, it should be appreciated that the instructions stored on the computer-readable medium, described above, are not limited to instructions embodied as part of an application program running on a host computer. Rather, the instructions may be embodied as any type of computer code (e.g., software or microcode) that can be employed to program a computer to implement aspects of the technology described herein. The computer executable instructions may be written in a suitable computer language or combination of several languages. Basic computational biology methods are known to those of ordinary skill in the art and are described in, for example, Setubal and Meidanis et al., *Introduction to Computational Biology Methods* (PWS Publishing Company, Boston, 1997); Salzberg, Searles, Kasif, (Ed.), *Computational Methods in Molecular Biology*, (Elsevier, Amsterdam, 1998); Rashidi and Buehler, *Bioinformatics Basics: Application in Biological Science and Medicine* (CRC Press, London, 2000) and Ouellette and Bzevanis *Bioinformatics: A Practical Guide for Analysis of Gene and Proteins* (Wiley & Sons, Inc., 2nd ed., 2001).

[0142] The functional modules of certain embodiments can include at minimum a determination module, a storage module, a computing module, and a display module. The functional modules can be executed on one, or multiple, computers, or by using one, or multiple, computer networks. The determination module has computer executable instructions to provide e.g., levels of expression products etc. in computer readable form.

[0143] The determination module can comprise any system for detecting a signal elicited from the genomic sequence and/or expression products of BRG1, EGFR, and/or B-RAF in a biological sample. In some embodiments,

such systems can include an instrument, e.g., the HiSeq 2500 (Illumina; San Diego, Calif.) for sequencing, e.g. nucleic acids isolated by exome capture (e.g. with the TruSeq Exome Enrichment Kit (Illumina; San Diego, Calif.). In another embodiment, the determination module can comprise multiple units for different functions, such as amplification and hybridization. In some embodiments, the determination module can be further configured to identify and detect the presence of at least one additional gene, allele, and/or gene expression product.

[0144] The information determined in the determination system can be read by the storage module. As used herein the “storage module” is intended to include any suitable computing or processing apparatus or other device configured or adapted for storing data or information. Examples of electronic apparatus suitable for use with the technology described herein include stand-alone computing apparatus, data telecommunications networks, including local area networks (LAN), wide area networks (WAN), Internet, Intranet, and Extranet, and local and distributed computer processing systems. Storage modules also include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage media, magnetic tape, optical storage media such as CD-ROM, DVD, electronic storage media such as RAM, ROM, EPROM, EEPROM and the like, general hard disks and hybrids of these categories such as magnetic/optical storage media. The storage module is adapted or configured for having recorded thereon, for example, sample name, allelic variants (e.g. sensitizing mutations or wild-type sequences), and frequency of each allelic variant. Such information may be provided in digital form that can be transmitted and read electronically, e.g., via the Internet, on diskette, via USB (universal serial bus) or via any other suitable mode of communication.

[0145] As used herein, “stored” refers to a process for encoding information on the storage module. Those skilled in the art can readily adopt any of the presently known methods for recording information on known media to generate manufactures comprising sequence information.

[0146] In one embodiment of any of the systems described herein, the storage module stores the output data from the determination module. In additional embodiments, the storage module stores the reference information such as the sequence of BRG1, EGFR, and/or B-RAF in cells which are not EZH2 combination treatment responsive, wild-type sequences, and/or sequences comprising sensitizing mutations.

[0147] The “computing module” can use a variety of available software programs and formats for computing the presence and/or absence of a sensitizing mutation in BRG1, EGFR, and/or B-RAF, e.g. aligning two or more sequences and determining if a sensitizing mutation is present in a test sequence as compared to a reference and/or wild-type sequence. Such algorithms are well established in the art. A skilled artisan is readily able to determine the appropriate algorithms based on the size and quality of the sample and type of data. The data analysis can be implemented in the computing module. In one embodiment, the computing module further comprises a comparison module, which compares the sequence of BRG1, EGFR, and/or B-RAF in the test sample obtained from a subject as described herein with the reference sequence of those genes and/or their expression products (FIG. 15). By way of example, when the sequence of BRG1 in the test sample obtained from a subject

is measured, a comparison module can compare or match the output data, e.g. with the reference sequence of BRG1 in a reference sample. In certain embodiments, the reference sequence can have been pre-stored in the storage module. During the comparison or matching process, the comparison module can determine whether the sequence in the test sample obtained from a subject differs from the reference sequence in such a way as to indicate a sensitizing mutation. Alternatively, the comparison module can determine whether the sequence in the test sample obtained from a subject matches a sequence known to comprise a sensitizing mutation. In various embodiments, the comparison module can be configured using existing commercially-available or freely-available software for comparison purpose, and may be optimized for particular data comparisons that are conducted.

[0148] The computing and/or comparison module, or any other module, can include an operating system (e.g., UNIX) on which runs a relational database management system, a World Wide Web application, and a World Wide Web server. World Wide Web application includes the executable code necessary for generation of database language statements (e.g., Structured Query Language (SQL) statements). Generally, the executables will include embedded SQL statements. In addition, the World Wide Web application may include a configuration file, which contains pointers and addresses to the various software entities that comprise the server as well as the various external and internal databases which must be accessed to service user requests. The Configuration file also directs requests for server resources to the appropriate hardware as may be necessary should the server be distributed over two or more separate computers. In one embodiment, the World Wide Web server supports a TCP/IP protocol. Local networks such as this are sometimes referred to as "Intranets." An advantage of such Intranets is that they allow easy communication with public domain databases residing on the World Wide Web (e.g., the GenBank or Swiss Pro World Wide Web site). Thus, in a particular preferred embodiment, users can directly access data (via Hypertext links for example) residing on Internet databases using a HTML interface provided by Web browsers and Web servers (FIG. 16).

[0149] The computing and/or comparison module provides a computer readable comparison result that can be processed in computer readable form by predefined criteria, or criteria defined by a user, to provide content based in part on the comparison result that may be stored and output as requested by a user using an output module, e.g., a display module.

[0150] In some embodiments, described herein is a computer system for determining if a subject will be responsive to a cancer treatment, the system comprising: a determination module configured to detect the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from a subject; a storage module configured to store output data from the determination module; a comparison module adapted to compare the data stored on the storage module with a reference level, and to provide a retrieved content, and a display module for displaying whether a BRG1, EGFR, or B-RAF sensitizing mutation was detected in the tumor cell sample; wherein the cancer treatment comprises the administration of an EZH2 inhibitor and a chemotherapeutic agent. In some embodiments, the determining module can measure the intensity of a detect-

able signal from a RT-PCR assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample. In some embodiments, the determining module can measure the intensity of a detectable signal from a sequencing assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample. In some embodiments, the determining module can measure the intensity of a detectable signal from a hybridization assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample.

[0151] In some embodiments, if the computing module determines that a sensitizing mutation of BRG1, EGFR, or B-RAF is present in the sample obtained from the subject, the display module can display a signal indicating that a sensitizing mutation has been detected. In some embodiments, the signal can indicate that the subject has an increased likelihood of responding to treatment with an EZH2 inhibitor and chemotherapeutic agent. In some embodiments, the content displayed on the display module can be a numerical value indicating one of these risks or probabilities. In such embodiments, the probability can be expressed in percentages or a fraction. For example, higher percentage or a fraction closer to 1 indicates a higher likelihood of a subject being responsive. In some embodiments, the content displayed on the display module can be single word or phrases to qualitatively indicate a risk or probability. For example, a word "unlikely" can be used to indicate a lower likelihood for being responsive to an EZH2 inhibitor combination therapy as described herein, while "likely" can be used to indicate a high likelihood for being responsive to an EZH2 inhibitor combination therapy as described herein.

[0152] In one embodiment, the content based on the computing and/or comparison result is displayed on a computer monitor. In one embodiment, the content based on the computing and/or comparison result is displayed through printable media. The display module can be any suitable device configured to receive from a computer and display computer readable information to a user. Non-limiting examples include, for example, general-purpose computers such as those based on Intel PENTIUM-type processor, Motorola PowerPC, Sun UltraSPARC, Hewlett-Packard PA-RISC processors, any of a variety of processors available from Advanced Micro Devices (AMD) of Sunnyvale, Calif., or any other type of processor, visual display devices such as flat panel displays, cathode ray tubes and the like, as well as computer printers of various types.

[0153] In one embodiment, a World Wide Web browser is used for providing a user interface for display of the content based on the computing/comparison result. It should be understood that other modules can be adapted to have a web browser interface. Through the Web browser, a user can construct requests for retrieving data from the computing/comparison module. Thus, the user will typically point and click to user interface elements such as buttons, pull down menus, scroll bars and the like conventionally employed in graphical user interfaces.

[0154] Systems and computer readable media described herein are merely illustrative embodiments of the technology relating to determining the presence of a sensitizing mutation, and therefore are not intended to limit the scope of the invention. Variations of the systems and computer readable media described herein are possible and are intended to fall within the scope of the invention.

[0155] The modules of the machine, or those used in the computer readable medium, may assume numerous configurations. For example, function may be provided on a single machine or distributed over multiple machines.

[0156] Provided herein are kits and devices for practicing the assays and methods described herein. In some embodiments, described herein is a device for determining the presence of a sensitizing mutation in BRG1, EGFR, and/or B-RAF in a test sample from a subject comprising a nucleic acid probe which is specific for a sensitizing mutation of BRG1, EGFR, or B-RAF. In some embodiments, described herein is a device for the presence of a sensitizing mutation in BRG1, EGFR, and/or B-RAF in a test sample from a subject comprising a nucleic acid probe which can amplify a nucleic acid sequence comprising a sensitizing mutation of BRG1, EGFR, or B-RAF. In some embodiments, the probe can be detectably labeled. In some embodiments, the kit can further comprise a reagent for producing a detectable signal.

[0157] In some embodiments, the kit can comprise further reagents, e.g. a restriction enzyme, a universal adaptor to be conjugated to a nucleotide molecule, a primer complementary to the universal adaptor, a wash agent, free nucleotide bases, a polymerase, or any combinations thereof.

[0158] In some embodiments, the kit can comprise at least one protein-binding moiety (e.g. an antibody reagent) specific for a protein comprising a sensitizing mutation of BRG1, EGFR, or B-RAF. In some embodiments, the kit can further comprise a solid substrate support affixed with the at least one protein-based binding moiety. In some embodiments, the solid substrate support is a microtiter plate for ELISA. In some embodiments, the solid substrate support is a dipstick. In some embodiments, the solid substrate support comprises a magnetic bead. In some embodiments, the kit comprises at least one further reagent, e.g. a reagent for detecting a detectable label.

[0159] In some embodiments, the kit or device can comprise a reference, e.g. a reference sample or reference signal. In some embodiments, the reference can comprise a wild-type sequence and/or wild-type expression product of one or more of BRG1, EGFR, and/or B-RAF.

[0160] In some embodiments, the kit or device can comprise a multiplex assay or the reagents for performing a multiplex assay, e.g. multiple sets of probes and/or primers or multiple antibody reagents such that the presence or absence of multiple sensitizing mutations can be detected in a single reaction mixture. In some embodiments, the kit or device can comprise an assay or the reagents for performing an assay for multiple sensitizing mutations in parallel, e.g. multiple sets of probes and/or primers or multiple antibody reagents such that the presence or absence of multiple sensitizing mutations can be detected in parallel.

[0161] For convenience, the meaning of some terms and phrases used in the specification, examples, and appended claims, are provided below. Unless stated otherwise, or implicit from context, the following terms and phrases include the meanings provided below. The definitions are provided to aid in describing particular embodiments, and are not intended to limit the claimed invention, because the scope of the invention is limited only by the claims. Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. If there is an apparent discrepancy between the

usage of a term in the art and its definition provided herein, the definition provided within the specification shall prevail.

[0162] For convenience, certain terms employed herein, in the specification, examples and appended claims are collected here.

[0163] The terms “decrease”, “reduced”, “reduction”, or “inhibit” are all used herein to mean a decrease by a statistically significant amount. In some embodiments, the terms “reduced”, “reduction”, “decrease”, or “inhibit” can mean a decrease by at least 10% as compared to a reference level, for example a decrease by at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or more or any decrease of at least 10% as compared to a reference level. These terms are not intended to cover complete reduction or inhibition.

[0164] In some aspects of all the embodiments, inhibition or reduction can be complete inhibition or complete reduction, i.e., a 100% decrease, i.e. absent or non-detectable level as compared to a reference level. In the context of a marker or symptom, a “decrease” is a statistically significant decrease in such level. The decrease can be, for example, at least 10%, at least 20%, at least 30%, at least 40% or more, and is preferably down to a level accepted as within the range of normal for an individual without such disorder.

[0165] The terms “increased”, “increase”, “enhance”, or “activate” are all used herein to mean an increase by a statistically significant amount. In some embodiments, the terms “increased”, “increase”, “enhance”, or “activate” can mean an increase of at least 10% as compared to a reference level, for example an increase of at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% increase or any increase between 10-100% as compared to a reference level, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold increase, or any increase between 2-fold and 10-fold or greater as compared to a reference level. In the context of a marker or symptom, a “increase” is a statistically significant increase in such level.

[0166] As used herein, a “subject” means a human or animal. Usually the animal is a vertebrate such as a primate, rodent, domestic animal or game animal. Primates include chimpanzees, cynomolgous monkeys, spider monkeys, and macaques, e.g., Rhesus. Rodents include mice, rats, woodchucks, ferrets, rabbits and hamsters. Domestic and game animals include cows, horses, pigs, deer, bison, buffalo, feline species, e.g., domestic cat, canine species, e.g., dog, fox, wolf, avian species, e.g., chicken, emu, ostrich, and fish, e.g., trout, catfish and salmon. In some embodiments, the subject is a mammal, e.g., a primate, e.g., a human. The terms, “individual,” “patient” and “subject” are used interchangeably herein.

[0167] Preferably, the subject is a mammal. The mammal can be a human, non-human primate, mouse, rat, dog, cat, horse, or cow, but is not limited to these examples. Mammals other than humans can be advantageously used as subjects that represent animal models of cancer. A subject can be male or female.

[0168] A subject can be one who has been previously diagnosed with or identified as suffering from or having a condition in need of treatment (e.g. cancer) or one or more complications related to such a condition, and optionally,

have already undergone treatment for cancer or the one or more complications related to cancer. Alternatively, a subject can also be one who has not been previously diagnosed as having cancer or one or more complications related to cancer. For example, a subject can be one who exhibits one or more risk factors for cancer or one or more complications related to cancer or a subject who does not exhibit risk factors.

[0169] A “subject in need” of treatment for a particular condition can be a subject having that condition, diagnosed as having that condition, or at risk of developing that condition.

[0170] A “cancer” or “tumor” as used herein refers to an uncontrolled growth of cells which interferes with the normal functioning of the bodily organs and systems. A subject that has a cancer or a tumor is a subject having objectively measurable cancer cells present in the subject's body. Included in this definition are benign and malignant cancers, as well as dormant tumors or micrometastases. Cancers which migrate from their original location and seed vital organs can eventually lead to the death of the subject through the functional deterioration of the affected organs. In some embodiments, a cancer cell can be a cell obtained from a tumor. By “metastasis” is meant the spread of cancer from its primary site to other places in the body. Cancer cells can break away from a primary tumor, penetrate into lymphatic and blood vessels, circulate through the bloodstream, and grow in a distant focus (metastasize) in normal tissues elsewhere in the body. Metastasis can be local or distant. Metastasis is a sequential process, contingent on tumor cells breaking off from the primary tumor, traveling through the bloodstream, and stopping at a distant site. At the new site, the cells establish a blood supply and can grow to form a life-threatening mass. Both stimulatory and inhibitory molecular pathways within the tumor cell regulate this behavior, and interactions between the tumor cell and host cells in the distant site are also significant. Metastases are most often detected through the sole or combined use of magnetic resonance imaging (MRI) scans, computed tomography (CT) scans, blood and platelet counts, liver function studies, chest X-rays and bone scans in addition to the monitoring of specific symptoms.

[0171] Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include, but are not limited to, basal cell carcinoma, biliary tract cancer; bladder cancer; bone cancer; brain and CNS cancer; breast cancer; cancer of the peritoneum; cervical cancer; choriocarcinoma; colon and rectum cancer; connective tissue cancer; cancer of the digestive system; endometrial cancer; esophageal cancer; eye cancer; cancer of the head and neck; gastric cancer (including gastrointestinal cancer); glioblastoma (GBM); hepatic carcinoma; hepatoma; intra-epithelial neoplasm; kidney or renal cancer; larynx cancer; leukemia; liver cancer; lung cancer (e.g., small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, and squamous carcinoma of the lung); lymphoma including Hodgkin's and non-Hodgkin's lymphoma; melanoma; myeloma; neuroblastoma; oral cavity cancer (e.g., lip, tongue, mouth, and pharynx); ovarian cancer; pancreatic cancer; prostate cancer; retinoblastoma; rhabdomyosarcoma; rectal cancer; cancer of the respiratory system; salivary gland carcinoma; sarcoma; skin cancer; squamous cell cancer; stomach cancer; testicular cancer; thyroid cancer;

uterine or endometrial cancer; cancer of the urinary system; vulval cancer; as well as other carcinomas and sarcomas; as well as B-cell lymphoma (including low grade/follicular non-Hodgkin's lymphoma (NHL); small lymphocytic (SL) NHL; intermediate grade/follicular NHL; intermediate grade diffuse NHL; high grade immunoblastic NHL; high grade lymphoblastic NHL; high grade small non-cleaved cell NHL; bulky disease NHL; mantle cell lymphoma; AIDS-related lymphoma; and Waldenstrom's Macroglobulinemia); chronic lymphocytic leukemia (CLL); acute lymphoblastic leukemia (ALL); Hairy cell leukemia; chronic myeloblastic leukemia; and post-transplant lymphoproliferative disorder (PTLD), as well as abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome.

[0172] The terms “compound” and “agent” refer to any entity which is normally not present or not present at the levels being administered and/or provided to a cell, tissue or subject. An agent can be selected from a group comprising: chemicals; small organic or inorganic molecules; signaling molecules; nucleic acid sequences; nucleic acid analogues; proteins; peptides; enzymes; aptamers; peptidomimetic, peptide derivative, peptide analogs, antibodies; intrabodies; biological macromolecules, extracts made from biological materials such as bacteria, plants, fungi, or animal cells or tissues; naturally occurring or synthetic compositions or functional fragments thereof. In some embodiments, the agent is any chemical, entity or moiety, including without limitation synthetic and naturally-occurring non-proteinaceous entities. In certain embodiments the agent is a small molecule having a chemical moiety. For example, chemical moieties include unsubstituted or substituted alkyl, aromatic, or heterocyclyl moieties including macrolides, leptomycins and related natural products or analogues thereof. Agents can be known to have a desired activity and/or property, or can be selected from a library of diverse compounds.

[0173] As used herein, the term “small molecule” refers to a chemical agent which can include, but is not limited to, a peptide, a peptidomimetic, an amino acid, an amino acid analog, a polynucleotide, a polynucleotide analog, an aptamer, a nucleotide, a nucleotide analog, an organic or inorganic compound (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

[0174] As used herein, the terms “protein” and “polypeptide” are used interchangeably herein to designate a series of amino acid residues, connected to each other by peptide bonds between the alpha-amino and carboxy groups of adjacent residues. The terms “protein”, and “polypeptide” refer to a polymer of amino acids, including modified amino acids (e.g., phosphorylated, glycosylated, etc.) and amino acid analogs, regardless of its size or function. “Protein” and “polypeptide” are often used in reference to relatively large polypeptides, whereas the term “peptide” is often used in reference to small polypeptides, but usage of these terms in the art overlaps. The terms “protein” and

“polypeptide” are used interchangeably herein when referring to a gene product and fragments thereof. Thus, exemplary polypeptides or proteins include gene products, naturally occurring proteins, homologs, orthologs, paralogs, fragments and other equivalents, variants, fragments, and analogs of the foregoing.

[0175] As used herein, the term “nucleic acid” or “nucleic acid sequence” refers to any molecule, preferably a polymeric molecule, incorporating units of ribonucleic acid, deoxyribonucleic acid or an analog thereof. The nucleic acid can be either single-stranded or double-stranded. A single-stranded nucleic acid can be one strand nucleic acid of a denatured double-stranded DNA. Alternatively, it can be a single-stranded nucleic acid not derived from any double-stranded DNA. In one aspect, the nucleic acid can be DNA. In another aspect, the nucleic acid can be RNA. Suitable nucleic acid molecules are DNA, including genomic DNA or cDNA. Other suitable nucleic acid molecules are RNA, including mRNA.

[0176] As used herein, “specific” when used in the context of an oligonucleotide primer or specific for a target nucleic acid refers to a level of complementarity between the primer and the target such that there exists an annealing temperature at which the primer will anneal to and mediate amplification of the target nucleic acid and will not anneal to or mediate amplification of non-target sequences present in a sample.

[0177] As used herein, the terms “treat,” “treatment,” “treating,” or “amelioration” refer to therapeutic treatments, wherein the object is to alleviate, ameliorate, inhibit, slow down or stop the progression or severity of a condition associated with a disease or disorder, e.g. cancer. The terms are not intended to encompass a complete cure. The term “treating” includes reducing or alleviating at least one adverse effect or symptom of a condition, disease or disorder associated with cancer.

[0178] In some aspects, a complete cure can be achieved, and in those situations, the method is considered a method for curing cancer.

[0179] Treatment is generally “effective” if one or more symptoms or clinical markers associated with the disease are objectively or subjectively reduced. Alternatively, treatment is “effective” if the progression of a disease is reduced or halted. That is, “treatment” includes not just the improvement of symptoms or markers, but also a cessation of some symptoms, or at least slowing of, progress or worsening of symptoms compared to what would be expected in the absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and/or decreased mortality. The term “treatment” of a disease also includes providing at least some relief from the symptoms or side-effects of the disease (including palliative treatment).

[0180] As used herein, the term “pharmaceutical composition” refers to the active agent in combination with a pharmaceutically acceptable carrier e.g. a carrier commonly used in the pharmaceutical industry. The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic

response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0181] As used herein, the term “administering,” refers to the placement of a compound as disclosed herein into a subject by a method or route which results in at least partial delivery of the agent at a desired site. Pharmaceutical compositions comprising the compounds disclosed herein can be administered by any appropriate route which results in an effective treatment in the subject.

[0182] The term “statistically significant” or “significantly” refers to statistical significance and generally means a two standard deviation (2SD) or greater difference.

[0183] Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients or reaction conditions used herein should be understood as modified in all instances by the term “about.” The term “about” when used in connection with percentages can mean $\pm 1\%$.

[0184] As used herein the term “comprising” or “comprises” is used in reference to compositions, methods, and respective component(s) thereof, that are essential to the method or composition, yet open to the inclusion of unspecified elements, whether essential or not.

[0185] The term “consisting of” refers to compositions, methods, and respective components thereof as described herein, which are exclusive of any element not recited in that description of the embodiment.

[0186] As used herein the term “consisting essentially of” refers to those elements required for a given embodiment. The term permits the presence of elements that do not materially affect the basic and novel or functional characteristic(s) of that embodiment.

[0187] The singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. Similarly, the word “or” is intended to include “and” unless the context clearly indicates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below. The abbreviation, “e.g.” is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation “e.g.” is synonymous with the term “for example.”

[0188] Definitions of common terms in cell biology and molecular biology can be found in “The Merck Manual of Diagnosis and Therapy”, 19th Edition, published by Merck Research Laboratories, 2006 (ISBN 0-911910-19-0); Robert S. Porter et al. (eds.), The Encyclopedia of Molecular Biology, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); Benjamin Lewin, Genes X, published by Jones & Bartlett Publishing, 2009 (ISBN-10: 0763766321); Kendrew et al. (eds.), Molecular Biology and Biotechnology: a Comprehensive Desk Reference, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8) and Current Protocols in Protein Sciences 2009, Wiley Intersciences, Coligan et al., eds.

[0189] Unless otherwise stated, the present invention was performed using standard procedures, as described, for example in Sambrook et al., Molecular Cloning: A Laboratory Manual (3 ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA (2001); Davis et al., Basic Methods in Molecular Biology, Elsevier Science Publishing, Inc., New York, USA (1995); Current Protocols in Cell Biology (CPCB) (Juan S. Bonifacino et. al. ed., John Wiley

and Sons, Inc.), and Culture of Animal Cells: A Manual of Basic Technique by R. Ian Freshney, Publisher: Wiley-Liss; 5th edition (2005), Animal Cell Culture Methods (Methods in Cell Biology, Vol. 57, Jennie P. Mather and David Barnes editors, Academic Press, 1st edition, 1998) which are all incorporated by reference herein in their entireties.

[0190] Other terms are defined herein within the description of the various aspects of the invention.

[0191] All patents and other publications; including literature references, issued patents, published patent applications, and co-pending patent applications; cited throughout this application are expressly incorporated herein by reference for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with the technology described herein. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents.

[0192] The description of embodiments of the disclosure is not intended to be exhaustive or to limit the disclosure to the precise form disclosed. While specific embodiments of, and examples for, the disclosure are described herein for illustrative purposes, various equivalent modifications are possible within the scope of the disclosure, as those skilled in the relevant art will recognize. For example, while method steps or functions are presented in a given order, alternative embodiments may perform functions in a different order, or functions may be performed substantially concurrently. The teachings of the disclosure provided herein can be applied to other procedures or methods as appropriate. The various embodiments described herein can be combined to provide further embodiments. Aspects of the disclosure can be modified, if necessary, to employ the compositions, functions and concepts of the above references and application to provide yet further embodiments of the disclosure. These and other changes can be made to the disclosure in light of the detailed description. All such modifications are intended to be included within the scope of the appended claims.

[0193] Specific elements of any of the foregoing embodiments can be combined or substituted for elements in other embodiments. Furthermore, while advantages associated with certain embodiments of the disclosure have been described in the context of these embodiments, other embodiments may also exhibit such advantages, and not all embodiments need necessarily exhibit such advantages to fall within the scope of the disclosure.

[0194] The technology described herein is further illustrated by the following examples which in no way should be construed as being further limiting.

[0195] Some embodiments of the technology described herein can be defined according to any of the following numbered paragraphs:

- [0196]** 1. A method of treating cancer, the method comprising:
- [0197]** administering or prescribing a chemotherapeutic agent and an EZH2 inhibitor to a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell.
- [0198]** 2. A method of identifying a subject who is a candidate for treatment of cancer with a combination therapy comprising an EZH2 inhibitor and a chemotherapeutic agent, the method comprising:
- [0199]** detecting a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject;
- [0200]** wherein if the sensitizing mutation is detected, the subject is identified as a candidate for cancer treatment with a combination therapy comprising a chemotherapeutic agent and an EZH2 inhibitor; and
- [0201]** wherein if the sensitizing mutation is not detected, the subject is identified as a candidate for cancer treatment which does not comprise administering an EZH2 inhibitor.
- [0202]** 3. A method of classifying a tumor cell as an EZH2 combination treatment responsive tumor, the method comprising:
- [0203]** detecting a sensitizing mutation of BRG1, EGFR, or B-RAF in a sample comprising the tumor cell;
- [0204]** wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF classifies the tumor as a EZH2 combination treatment responsive tumor and the absence of a sensitizing mutation of BRG1, EGFR, or B-RAF classifies the tumor as a EZH2 combination treatment non-responsive tumor.
- [0205]** 4. The method of any of paragraphs 1-3, wherein the EZH2 inhibitor is selected from the group consisting of:
- [0206]** an inhibitory nucleic acid; DZNep; and S-adenosyl-L-homocysteine; and GSK126.
- [0207]** 5. The method of any of paragraphs 1-4, wherein the chemotherapeutic agent is selected from the group consisting of:
- [0208]** a topoisomerase inhibitor; a topoisomerase I inhibitor; and a topoisomerase II inhibitor.
- [0209]** 6. The method of paragraph 5, wherein the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of:
- [0210]** camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D.
- [0211]** 7. The method of paragraph 5, wherein the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of:
- [0212]** doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331.
- [0213]** 8. The method of any of paragraphs 5-7, wherein the subject is further administered cisplatin.
- [0214]** 9. The method of any of paragraphs 1-4, wherein the chemotherapeutic agent is a PARP inhibitor.
- [0215]** 10. The method of any of paragraphs 1-4, wherein the chemotherapeutic agent is a CDK1 inhibitor.
- [0216]** 11. The method of any of paragraphs 1-4, wherein the chemotherapeutic agent is an EGFR inhibitor.
- [0217]** 12. The method of any of paragraphs 1-11, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:

- [0218] a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*.
- [0219] 13. The method of any of paragraphs 1-12, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:
- [0220] a mutation which increased the expression level of EGFR; an exon 19 deletion of EGFR; E746_A750del; E746_A749del; T790M; and L858R.
- [0221] 14. The method of any of paragraphs 1-13, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:
- [0222] G496A and L597V.
- [0223] 15. The method of any of paragraphs 1-14, wherein the presence of the mutation is determined using an assay selected from the group consisting of:
- [0224] hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.
- [0225] 16. The method of any of paragraphs 1-15, wherein the mutation is present in the genomic DNA of the tumor cell.
- [0226] 17. The method of any of paragraphs 1-15, wherein the mutation is present in the mRNA transcripts of the tumor cell.
- [0227] 18. The method of any of paragraphs 1-17, wherein the cancer is selected from the group consisting of:
- [0228] lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.
- [0229] 19. The method of any of paragraphs 1-18, further comprising the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.
- [0230] 20. An assay comprising:
- [0231] subjecting a tumor cell sample from a subject to at least one analysis to detect the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF;
- [0232] wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with an EZH2 inhibitor and a chemotherapeutic agent.
- [0233] 21. The assay of paragraph 20, wherein the presence of the mutation is determined using an assay selected from the group consisting of:
- [0234] hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.
- [0235] 22. An assay for selecting a treatment regimen for a subject with cancer, comprising:
- [0236] subjecting a nucleotide molecule derived from a biological sample of a subject, who is determined to suffer from or have a risk for cancer, to at least one genotyping analysis adapted to determine the presence of a sensitizing mutation in one or more of B-RAF, EGFR, and BRG1;
- [0237] wherein if at least one sensitizing mutation is determined to be present, a treatment regimen comprising a combination of an EZH2 inhibitor and a chemotherapeutic agent is administered.
- [0238] 23. The assay of paragraph 22, wherein if no sensitizing mutations are determined to be present, a treatment regimen comprising a combination of an EZH2 inhibitor and a chemotherapeutic agent is not administered.
- [0239] 24. The assay of any of paragraphs 20-23, wherein the biological sample comprises a tumor cell.
- [0240] 25. An assay comprising:
- [0241] contacting a tumor cell sample obtained from a human subject having cancer with a nucleic acid probe to detect the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF; and
- [0242] detecting the presence or intensity of a signal which indicates the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF;
- [0243] wherein the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF indicates the subject has a cancer which will respond to treatment with an EZH2 inhibitor and a chemotherapeutic agent.
- [0244] 26. The assay of any of paragraphs 20-25, wherein a detectable signal is generated by the probe when a sensitizing mutation is present.
- [0245] 27. The assay of any of paragraphs 20-26, wherein the probe is detectably labeled.
- [0246] 28. The assay of any of paragraphs 20-27, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:
- [0247] a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*.
- [0248] 29. The assay of any of paragraphs 20-28, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:
- [0249] a mutation which increased the expression level of EGFR; an exon 19 deletion of EGFR; E746_A750del; E746_A749del; T790M; and L858R.
- [0250] 30. The assay of any of paragraphs 20-29, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:
- [0251] G496A and L597V.
- [0252] 31. The assay of any of paragraphs 20-30, wherein the mutation is present in the genomic DNA of the tumor cell.
- [0253] 32. The assay of any of paragraphs 20-31, wherein the mutation is present in the mRNA transcripts of the tumor cell.
- [0254] 33. The assay of any of paragraphs 20-32, further comprising the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.
- [0255] 34. A method of determining whether a subject is likely to respond to a combination treatment for cancer, the method comprising:
- [0256] detecting the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from the subject;
- [0257] wherein the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF indicates the subject has an increased likelihood of responding to a treatment to cancer;
- [0258] wherein the treatment for cancer comprises administration of an EZH2 inhibitor and a chemotherapeutic agent.
- [0259] 35. The method of paragraph 34, wherein the EZH2 inhibitor is selected from the group consisting of:

- [0260] an inhibitory nucleic acid; DZNep; S-adenosyl-L-homocysteine; and GSK126.
- [0261] 36. The method of any of paragraphs 34-35, wherein the chemotherapeutic agent is selected from the group consisting of:
- [0262] a topoisomerase inhibitor; a topoisomerase I inhibitor; a topoisomerase II inhibitor
- [0263] 37. The method of paragraph 36, wherein the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of:
- [0264] camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D.
- [0265] 38. The method of paragraph 36, wherein the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of:
- [0266] doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331.
- [0267] 39. The method of any of paragraphs 36-38, wherein the subject is further administered cisplatin.
- [0268] 40. The method of any of paragraphs 34-35, wherein the chemotherapeutic agent is a PARP inhibitor.
- [0269] 41. The method of any of paragraphs 34-35, wherein the chemotherapeutic agent is a CDK1 inhibitor.
- [0270] 42. The method of any of paragraphs 34-35, wherein the chemotherapeutic agent is an EGFR inhibitor.
- [0271] 43. The method of any of paragraphs 34-42, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:
- [0272] a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*.
- [0273] 44. The method of any of paragraphs 34-43, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:
- [0274] a mutation which increased the expression level of EGFR; an exon 19 deletion of EGFR; E746_A750del; E746_A749del; T790M; and L858R.
- [0275] 45. The method of any of paragraphs 34-44, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:
- [0276] G496A and L597V.
- [0277] 46. The method of any of paragraphs 34-45, wherein the presence of the mutation is determined using an assay selected from the group consisting of:
- [0278] hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.
- [0279] 47. The method of any of paragraphs 34-46, wherein the mutation is present in the genomic DNA of the tumor cell.
- [0280] 48. The method of any of paragraphs 34-47, wherein the mutation is present in the mRNA transcripts of the tumor cell.
- [0281] 49. The method of any of paragraphs 34-48, wherein the cancer is selected from the group consisting of:
- [0282] lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.
- [0283] 50. A computer system for determining if a subject will be responsive to a cancer treatment, the system comprising:
- [0284] a determination module configured to detect the presence of a sensitizing mutation in BRG1, EGFR, or B-RAF in a tumor cell sample obtained from a subject;
- [0285] a storage module configured to store output data from the determination module;
- [0286] a comparison module adapted to compare the data stored on the storage module with a reference level, and to provide a retrieved content, and
- [0287] a display module for displaying whether a BRG1, EGFR, or B-RAF sensitizing mutation was detected in the tumor cell sample;
- [0288] wherein the cancer treatment comprises the administration of an EZH2 inhibitor and a chemotherapeutic agent.
- [0289] 51. The system of paragraph 50, wherein the determining module measures the intensity of a detectable signal from a RT-PCR assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample.
- [0290] 52. The system of paragraph 50, wherein the determining module measures the intensity of a detectable signal from a sequencing assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample.
- [0291] 53. The system of paragraph 50, wherein the determining module measures the intensity of a detectable signal from a hybridization assay indicating the presence of a sensitizing mutation of BRG1, EGFR, or B-RAF tumor cell sample.
- [0292] 54. The system of any of paragraphs 50-53, wherein if the computing module determines that the a sensitizing mutation of BRG1, EGFR, or B-RAF is present in the sample obtained from the subject, the display module displays a signal indicating that a sensitizing mutation has been detected.
- [0293] 55. The system of any of paragraphs 50-54, wherein the signal indicates that the subject has an increased likelihood of responding to treatment with an EZH2 inhibitor and chemotherapeutic agent.
- [0294] 56. The system of any of paragraphs 50-55, wherein the EZH2 inhibitor is selected from the group consisting of:
- [0295] an inhibitory nucleic acid; DZNep; S-adenosyl-L-homocysteine; and GSK126.
- [0296] 57. The system of any of paragraphs 50-56, wherein the chemotherapeutic agent is selected from the group consisting of:
- [0297] a topoisomerase inhibitor; a topoisomerase I inhibitor; a topoisomerase II inhibitor
- [0298] 58. The system of paragraph 55, wherein the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of:
- [0299] camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and
- [0300] indimitecan; and lamellarin D.
- [0301] 59. The system of paragraph 55, wherein the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of:
- [0302] doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331.

- [0303] 60. The system of any of paragraphs 57-59, wherein the subject is further administered cisplatin.
- [0304] 61. The system of any of paragraphs 50-54, wherein the chemotherapeutic agent is a PARP inhibitor.
- [0305] 62. The system of any of paragraphs 50-54, wherein the chemotherapeutic agent is a CDK1 inhibitor.
- [0306] 63. The system of any of paragraphs 50-54, wherein the chemotherapeutic agent is an EGFR inhibitor.
- [0307] 64. The system of any of paragraphs 50-63, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:
- [0308] a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*.
- [0309] 65. The system of any of paragraphs 50-64, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:
- [0310] a mutation which increased the expression level of EGFR; an exon 19 deletion of EGFR; E746_A750del; E746_A749del; T790M; and L858R.
- [0311] 66. The system of any of paragraphs 50-65, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:
- [0312] G496A and L597V.
- [0313] 67. The system of any of paragraphs 50-66, wherein the presence of the mutation is determined using an assay selected from the group consisting of:
- [0314] hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.
- [0315] 68. The system of any of paragraphs 50-67, wherein the mutation is present in the genomic DNA of the tumor cell.
- [0316] 69. The system of any of paragraphs 50-68, wherein the mutation is present in the mRNA transcripts of the tumor cell.
- [0317] 70. The system of any of paragraphs 50-69, wherein the cancer is selected from the group consisting of:
- [0318] lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.
- [0319] 71. A kit comprising:
- [0320] a nucleic acid probe which is specific for a sensitizing mutation of BRG1, EGFR, or B-RAF.
- [0321] 72. The kit of paragraph 71, wherein the probe is detectably labeled.
- [0322] 73. A kit comprising:
- [0323] a nucleic acid probe which can amplify a nucleic acid sequence comprising a sensitizing mutation of BRG1, EGFR, or B-RAF.
- [0324] 74. The kit of any of paragraphs 71-73, further comprising a reagent for producing a detectable signal.
- [0325] 75. The use of an EZH2 inhibitor in combination with a chemotherapeutic agent in treatment of a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell obtained from the subject.
- [0326] 76. The use of paragraph 75, wherein the EZH2 inhibitor is selected from the group consisting of:
- [0327] an inhibitory nucleic acid; DZNep; and S-adenosyl-L-homocysteine; and GSK126.
- [0328] 77. The use of any of paragraphs 75-76, wherein the chemotherapeutic agent is selected from the group consisting of:
- [0329] a topoisomerase inhibitor; a topoisomerase I inhibitor; and a topoisomerase II inhibitor.
- [0330] 78. The use of paragraph 77, wherein the chemotherapeutic agent is a topoisomerase I inhibitor selected from the group consisting of:
- [0331] camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and
- [0332] indimitecan; and lamellarin D.
- [0333] 79. The use of paragraph 77, wherein the chemotherapeutic agent is a topoisomerase II inhibitor selected from the group consisting of:
- [0334] doxorubicin; etoposide; amsacrine; teniposide; ICRF-193; genistein; daunorubicin; mitoxantrone; ellipticines; aurintricarboxylic acid; and HU-331.
- [0335] 80. The use of any of paragraphs 77-79, wherein the subject is further administered cisplatin.
- [0336] 81. The use of any of paragraphs 75-76, wherein the chemotherapeutic agent is a PARP inhibitor.
- [0337] 82. The use of any of paragraphs 75-76, wherein the chemotherapeutic agent is a CDK1 inhibitor.
- [0338] 83. The use of any of paragraphs 75-76, wherein the chemotherapeutic agent is an EGFR inhibitor.
- [0339] 84. The use of any of paragraphs 75-83, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:
- [0340] a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272*.
- [0341] 85. The use of any of paragraphs 75-84, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:
- [0342] a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R.
- [0343] 86. The use of any of paragraphs 75-85, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:
- [0344] G496A and L597V. p0
87. The use of any of paragraphs 75-86, wherein the presence of the mutation is determined using an assay selected from the group consisting of:
- [0345] hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.
- [0346] 88. The use of any of paragraphs 75-87, wherein the mutation is present in the genomic DNA of the tumor cell.
- [0347] 89. The use of any of paragraphs 75-88, wherein the mutation is present in the mRNA transcripts of the tumor cell.
- [0348] 90. The use of any of paragraphs 75-89, wherein the cancer is selected from the group consisting of:
- [0349] lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.
- [0350] 91. The use of any of paragraphs 75-90, further comprising the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.

EXAMPLES

Example 1

[0351] Drugs targeting epigenetic enzymes such as EZH2 may help to combat many cancers, including non-small cell lung cancer (NSCLC). As described herein, the effects of EZH2 inhibition on lung cancer growth and chemotherapy response both were examined both in vitro and in vivo. EZH2 inhibition via small hairpin or drug DZNep slowed growth of the majority of cell lines, and sensitized half of the lines to the chemotherapy etoposide. Unexpectedly, the remaining lines (termed “protected lines”) showed an increase in etoposide IC50 in response to EZH2 inhibition. Cell lines sensitized to etoposide entered S phase in response to EZH2 inhibition, while protected lines showed cell cycle arrest and up-regulation of the cell cycle inhibitor p57. Sensitized lines were highly enriched for Brg1 and EGFR mutations, and modulation of Brg1 or EGFR levels changed the sensitized and protected phenotypes. Genes co-expressed with EZH2 in primary human lung cancer samples were highly enriched for cell cycle regulators and strongly predicted poor survival. These data further knowledge of the effects of EZH2 inhibition on cancer cells, and permit methods, assays, and systems to predict whether patients will benefit from dual chemotherapy and EZH2 inhibitor therapies.

[0352] Understanding which tumors will respond to epigenetic therapies will be essential for this new class of drugs to benefit patients. It is demonstrated herein, for the first time, that EZH2 inhibition changes chemotherapy sensitivities of lung cancer cells dependent on genetic interactions of the Brg1-containing SWI/SNF complex and EGFR signaling. These findings suggest that patients with lung tumors bearing, e.g., EGFR or Brg1 mutations will benefit from EZH2 inhibition treatment combined with chemotherapy, while patients with wild-type EGFR and Brg1 will respond well to EZH2 inhibition as a single therapy. EZH2 inhibition differentially affects lung cancer chemotherapy sensitivities; Brg1 and EGFR mutations are negatively correlated and predict sensitization; Brg1 and EGFR interact genetically to control EZH2 inhibition response; and an EZH2 expression gene signature significantly predicts patient survival.

[0353] Lung cancer is the leading cause of cancer related deaths worldwide, and its poor prognosis is attributable to metastasis and therapy resistance (Herbst et al., 2008). Epigenetic mechanisms, such as histone modifications, are thought to be crucial for the survival of metastatic and therapy resistant cells (Baylin, 2011, Crea et al., 2011a, Min et al., 2010, Iliopoulos et al., 2010). Therefore, combining epigenetic therapies with chemotherapy and radiation treatments may allow for more complete treatment responses. Polycomb Repressive Complexes (PRCs) are key regulators of chromatin states in stem cells and cancer (Simon and Lange, 2008, Lee et al., 2006, Ben-Porath et al., 2008). Binding of PRCs to regions of chromatin, and histone modifications mediated by these PRCs, can affect higher order chromatin structure, rendering areas in the genome inaccessible to transcriptional machinery (Simon and Kingston, 2009).

[0354] PRC2 often contains EZH2, a methyltransferase that tri-methylates histone H3 at lysine 27 (H3K27me3). H3K27me3 moieties are known to be docking sites for the

Bmi1 containing complex PRC1, which can subsequently ubiquitylate histone H2A (Spamann and van Lohuizen, 2006, Bracken et al., 2009).

[0355] Many lines of evidence suggest that EZH2 inhibition may help to eradicate aggressive lung tumor cells, including, by way of non-limiting example, non-small cell lung cancer cells. Previous work by the inventors demonstrated that in the absence of Bmi1, progression of lung adenocarcinomas driven by oncogenic Kras was impaired (Dovey et al., 2007). Further, expression of EZH2 is highly correlated with poor prognosis in all stages of both adenocarcinoma and squamous cell carcinoma, the two major subtypes of NSCLC (Kikuchi et al., 2010, Takawa et al., 2011, Huqun et al., 2011). Of note, studies also found that Bmi1 expression has no predictive value (Kikuchi et al., 2010, Breuer et al., 2004), suggesting that EZH2 is more important for human lung cancers than is Bmi1. Because EZH2 is an enzyme, it is a particularly attractive target for development of small molecule inhibitors (Simon and Lange, 2008). Furthermore, EZH2 and other PRC components home to sites of DNA damage after ionizing radiation (Chou et al., 2010). Therefore, without wishing to be bound by theory, incorporation of PRC complex inhibitors may lead to more complete responses of tumors to standard therapies by preventing proper DNA repair (Crea et al., 2011, Iliopoulos et al., 2010). In support of this concept, EZH2 inhibition was shown to increase apoptosis of one lung cancer cell line when combined with chemotherapy in vitro (Wu et al., 2011).

[0356] Lung cancer is one of the most genetically complex diseases in humans. The most common genetic lesions in lung cancer include inactivating mutations in p53 (50-70% of NSCLC), deletion and mutation of LKB1 (20-35% of NSCLC), and activating mutations in K-ras (10-30% of NSCLC, Herbst et al., 2008, TCGA, 2012, Imielinski et al., 2012). Also frequently observed are activating mutations and amplifications in EGFR (10-40% of NSCLC), which cause increased Raf-MEK-MAPK signaling (Herbst et al., 2008, Rodriguez-Nieto et al., 2011, TCGA, 2012, Imielinski et al., 2012). The Raf-MEK-MAPK pathway can also be activated in lung tumors through activating mutations in B-Raf (3% of NSCLC, Herbst et al., 2008, Ji et al., 2007, TCGA, 2012, Imielinski et al., 2012). Recently, Brg1, the ATPase in the SWI/SNF chromatin-remodeling complex, was identified as a gene commonly deleted or truncated in lung tumors (2-10% of NSCLC, Rodriguez-Nieto et al., 2011, TCGA, 2012, Imielinski et al., 2012). Because of this genetic complexity, testing EZH2 inhibitors must be performed by carefully considering the mutations driving each tumor tested. As the inventors have previously demonstrated, laboratories studies found that consideration of lung tumor genotype is essential to understanding the complex biological processes which allow tumor cells to survive and propagate (Curtis et al., 2010).

[0357] In order to validate the use of EZH2 inhibition to treat NSCLC, the effects of EZH2 inhibition on lung tumor progression and chemotherapy responses were examined, using a large panel of human lung cancer cell lines grown both in vivo and in vitro. It was found that mutations in EGFR, B-Raf and Brg1 significantly predict whether a cell line will be sensitized or protected from chemotherapy by EZH2 inhibition. These data link, for the first time, EZH2 inhibition phenotypes to common mutations in NSCLC.

Results

[0358] EZH2 knock-down has differential effects on chemotherapy sensitivities. In order to test EZH2 inhibition as a therapy for NSCLC, we obtained ~30 human NSCLC lines that vary in genotype, stage and subtype (Table SI) and used a lentiviral small hairpin targeting EZH2 to stably knock-down EZH2 expression. By Western Blot, EZH2 protein and the PRC2 catalytic mark, H3K27me3, was diminished in every cell line transduced with this hairpin when compared to matched shGFP transduced cells (FIG. 1A, FIG. 7A). Growth rates in standard 2D adherent cultures were attenuated in 9 out of 12 cell lines in response to small-hairpin mediated EZH2 ablation over a period of 10 days (FIG. 1B). The shEZH2 Calu6, H441 and H157 lines were the most affected cell lines with 4.5-fold, 3.9-fold and 2.1-fold decreases in proliferation respectively when compared to their shGFP control lines. HCC95 shEZH2 was the outlier, which proliferated 3.2-fold more than the matched shGFP control line. To assess self-renewal capacity of the shEZH2 lines, efficiency of primary and secondary 3D sphere colonies was quantified (FIGS. 7B-7C). All of the shEZH2 cell lines had diminished secondary sphere forming capacity relative to their shGFP controls.

[0359] To test if EZH2 depleted cells are more sensitive to standard therapeutics such as chemotherapy, cell viability assays were performed with the commonly used chemotherapy etoposide. Both shGFP or shEZH2 cell lines were treated for 4 days with several doses of etoposide, and cell viability was measured. Each cell line was normalized to the untreated control thereby accounting for the decrease in proliferation of the shEZH2 lines, and change in percent survival between the shGFP and shEZH2 lines for both 1 μ M and 10 μ M etoposide was determined (FIG. 1C). Unexpectedly, though some cell lines were more sensitive to etoposide when EZH2 was knocked down, others were more viable. For example the H522 shEZH2 was an average of 19% less viable in 1 μ M and 10 μ M etoposide than H522 shGFP, while the H23 shEZH2 was an average of 31% more viable than its shGFP control line. These two classes of cell lines are referred to herein as those that are sensitized to etoposide by EZH2 inhibition (i.e. EZH2 inhibitor combination treatment responsive) and those that are protected.

[0360] EZH2 inhibition by DZNep has differential effects on chemotherapy sensitivities. Next, to learn if these results could be reproduced by chemical inhibition of EZH2, a small molecule known to target EZH2 termed DZNep was obtained (Tan et al., 2007, Puppe et al., 2009, Crea et al., 2011a, Crea et al., 2011b, Wu et al., 2011, Kikuchi et al., 2012). Western Blot confirmed that a 1 μ M dose of DZNep during a 4-day treatment effectively reduced EZH2 protein abundance and the H3K27me3 catalytic mark (FIG. 2A). At the 4-day time point, 1 μ M DZNep caused an average of 20% reduction in growth of the cell lines (FIG. 8A). This dose of DZNep also lead to reduced growth rates of 11 out of 12 cell lines assessed over a 10 day time-course (FIG. 2B). Similarly to the results with shEZH2, Calu6, H441 and H157 all showed 2.1-fold decreases in proliferation in response to DZNep. HCC15 was the most affected cell line with a 3.2-fold decrease in response to DZNep, while HCC95 was again the outlier that increased proliferation in response to the EZH2 inhibitor. Together these data indicate that an EZH2 inhibitor may be a viable therapeutic strategy to slow the growth of the majority of NSCLCs.

[0361] The synergy of DZNep with etoposide was assessed by performing etoposide dose response curves in the presence and absence of DZNep. After 4 days, cell viability was measured and the etoposide IC50s for each cell line with and without DZNep were compared. 16 of the 29 cell lines, including H157, HCC15 and PC9, were more sensitive to etoposide, while the other lines, such as H23, H460 and H441, were less sensitive to the chemotherapy in the presence of DZNep (FIG. 2C). Importantly, when the etoposide IC50 values were averaged for the two classes, no significant difference between the sensitized and protected cell lines was observed, suggesting that sensitization was not unique to chemo-resistant cell lines as seen in previous studies (FIG. 25B, Ougolkov et al., 2008).

[0362] Etoposide acts by binding within the Topoisomerase II (TopoII) complex and preventing re-ligation of the double strand break required to unwind DNA during S phase (Baldwin and Osheroff, 2005). To learn if the described results were specific to TopoII inhibition, two other common chemotherapies were used: doxorubicin, which also targets the TopoII complex, and cis-platinum, which causes double strand breaks through pyrimidine dimers. Cell lines segregated into the same protected and sensitized classes with doxorubicin combined with DZNep, but the pattern was not observed when cis-platinum and DZNep were combined (FIGS. 8C, 8D).

[0363] In vitro sensitivities to DZNep/Etoposide predict in vivo responses. To determine if the protected and sensitized phenotypes could be observed in vivo, cells were transplanted subcutaneously, tumors allowed to initiate, and mice treated with etoposide and DZNep. 12 days after cell injection, tumor bearing mice were randomized into groups that received either etoposide (20 mg/kg/day) or placebo (corn oil) daily for five days. The cohorts were further subdivided into groups that also received DZNep, either 2 mg/kg/day twice during the first week (dosing scheme A), or 1 mg/kg/day twice during the first two weeks (dosing scheme B).

[0364] For the sensitized cell line H157, dosing scheme A yielded the most significant results. DZNep alone was effective at reducing tumor burden (FIG. 3A, $p=0.03$ untreated vs. DZNep), and dual therapy prevented tumors from forming in 7/8 xenografts, proving more efficacious than etoposide alone ($p=0.0002$ untreated vs. dual, $p=0.015$ etoposide vs. dual). Dosing scheme B was not as effective for the H157 line. In particular, treatment with DZNep during the second week appeared to have the unwanted effect of causing tumors to grow larger (FIG. 9A, $p=0.21$ untreated vs. DZNep, $p=0.49$ etoposide vs. dual). This result indicates that for sensitized lines, higher doses of DZNep earlier in tumor development are more effective than lower doses administered over a longer time period.

[0365] In contrast, for the protected cell line H23, dosing scheme B was most effective. H23 xenografts were very sensitive to DZNep as a single therapy (FIG. 3B, $p=0.001$ untreated vs. DZNep), but those that received dual therapy grew significantly larger than those treated with either DZNep or etoposide alone ($p<0.0001$ dual vs. DZNep, $p=0.01$ dual vs. etoposide). For this line, treatment with DZNep during the second week was required for the long-term cytostatic effect of DZNep alone. Using dosing scheme A, the difference between untreated xenografts and those that received only DZNep was less significant (FIG. 9B, $p=0.01$ untreated vs. DZNep). Together these data demonstrate that sensitized and protected phenotypes can be

observed in xenografts, and that dosing schemes for both types of tumors can effectively reduce tumor burden.

[0366] EZH2 inhibition in presence of chemotherapy differentially affects cell cycle. It was next sought to learn the mechanism through which the protected cells were surviving chemotherapy in the presence of EZH2 inhibition. Using OncoPrint, an EZH2 expression gene signature was generated by annotating the top 20 genes co-expressed with EZH2 in 9 studies consisting of over 1,000 primary non-small cell lung cancer samples. Of the 180 probes, 64 were redundant, leading to a list of 116 genes highly co-expressed with EZH2 (Table 2). By gene ontology, these 116 genes are highly enriched for cell cycle, DNA synthesis and DNA repair (Table 3, FIG. 4A, $p < 0.001$). Whether this EZH2 expression signature had predictive power for cancer progression was queried using the Director's Challenge dataset of 410 human lung adenocarcinomas (Director's Challenge, 2008). When patients were clustered into 2 groups, those whose tumors had high expression of the EZH2 expression signature, and those whose tumors had low expression, a significant difference was observed in survival between the two groups (FIG. 4B, $p < 0.00001$). Patients with EZH2 expression high tumors survived a median of 3 years post tumor biopsy, while patients with EZH2 expression low tumors survived a median of 6 years.

[0367] Due to the significant enrichment of cell cycle regulatory genes in the EZH2 expression signature, it was hypothesized that EZH2 inhibition caused changes in cell cycle status that rendered cells more or less sensitive to chemotherapy. It was further hypothesized that protected lines undergo a cell cycle arrest in response to EZH2 inhibition, thus sparing them from etoposide-induced DNA damage, while sensitized lines are unable to arrest and instead continue to cycle in the presence of etoposide. To test this hypothesis, various cell lines were treated with 10 μ M etoposide (IC50), 1 μ M DZNep, or both for 4 days, and cells were fixed and stained with 7AAD to visualize ploidy by flow cytometry. The protected cell lines had an average of 10% fewer cells in S phase when treated with etoposide and DZNep compared to treatment with etoposide alone (FIG. 4C, FIG. 10A-10B S phase $p = 0.0001$). These protected cell lines also appeared to be collecting in the G1 and G2/M phases of the cell cycle, consistent with cell cycle arrest. In contrast, etoposide-treated sensitized cell lines had an average of 20% more cells in S phase in response to DZNep. Similar results were seen when comparing shGFP to shEZH2 lines treated with etoposide, and it was confirmed that EZH2 remains knocked down during the 4 day drug treatment (FIG. 4D, FIG. 10A-10C, S phase $p = 0.0006$). These data indicate that EZH2 inhibition has differential effects on cell cycle regulation, which in turn mediates differential survival in the presence of TopoII inhibitors.

[0368] To learn how cell cycle arrest is mediated in protected cell lines, RNA was isolated from the treated cell lines (4 sensitized lines: H522, HCC2450, H157 and HCC15, 4 protected lines: H460, H441, H23 and Calu6) and quantitative RT-PCR performed for cell cycle inhibitors. Several non-mutated cell cycle regulators are up-regulated in response to dual EZH2 inhibition and etoposide treatment, including p21, while others, such as p27, were unchanged (FIG. 10D). Of note, p57 is down-regulated in sensitized lines in response to etop/DZNep, and up-regulated in protected lines, following the pattern predicted by the cell cycle data (FIG. 4E, $p = 0.01$).

[0369] Brg1 and EGFR mutations are negatively correlated and predict sensitized phenotypes. By using mutation annotation available for the NSCLC lines, it was discovered that 14 of the 16 sensitized cell lines harbored mutations in Brg1 (also known as SMARCA4), B-Raf or EGFR, while 10 of the 13 protected cell lines were WT for the three genes (Table 1, Fisher's exact test $p = 0.001$). Of note, H23 and H1819 lines are protected from etoposide by DZNep, yet have Brg1 mutations. However, these mutations are both late in the coding region and are predicted to produce a protein with an intact ATPase (Medina et al., 2008). Expression changes were assessed in EGFR and Brg1 in response to therapy using the RNA isolated for cell cycle inhibitor analysis. In protected cell lines, Brg1 expression is significantly up-regulated and EGFR expression is significantly down-regulated in response to etoposide/DZNep, while remaining unchanged in sensitized lines (FIG. 18A, $p = 0.03$ Brg1, $p = 0.005$ EGFR).

[0370] In addition to correlating with etoposide/DZNep sensitivities, Brg1 and EGFR/B-Raf mutations appeared to be negatively correlated. In our panel of 29 cell lines, 31% have a Brg1 mutation and 27.5% have an EGFR or B-Raf mutation. With these mutation frequencies, 8.5% of cell lines are expected to have both Brg1 and EGFR/B-Raf mutations. However, no NSCLC cell line in this panel or elsewhere was observed that harbored both mutations (Table 1, Table 4, Fisher's exact test $p = 0.001$). To examine the pattern of Brg1 and EGFR mutations in primary human lung cancers, lung adenocarcinoma whole exome sequencing from The Cancer Genome Atlas (TCGA) and a recent publication were queried (Imielinski et al., 2012). Of the 424 adenocarcinoma samples, 86 (20.3%) had exon 19 deletion or missense mutation in EGFR or missense mutations in B-Raf, while 46 (10.8%) had missense, nonsense, deletion or insertion mutations in Brg1. With these allele frequencies, one would expect 2% of the tumor samples to have both Brg1 and EGFR/B-Raf mutations, though only 3 (0.7%) were observed (Fisher's exact test, $p = 0.006$, B-Raf G466E/SMARCA4 G594A, EGFR R574L/SMARCA4 A1249S, EGFR H1129Y/SMARCA4_homozygous focal deletion, all from Imielinski dataset). If the cell lines are included with the primary tumor data, the anti-correlation of EGFR/B-Raf mutations and SMARCA4 mutations becomes very significant (Fisher's exact test, $p < 0.0001$). In squamous cell carcinoma (TCGA, 2012), a similar trend was observed.

[0371] The negative correlation of Brg1 and EGFR mutations suggests that these pathways interact genetically and that these mutations may be functionally redundant. In the Director's Challenge dataset of primary human lung adenocarcinomas, samples that showed high expression of EGFR and/or Brg1 were analyzed for a correlation between these two genes. A strong negative correlation was found, which supports the theory that loss of Brg1 may be permissive for high expression of EGFR (Spearman's correlation = -0.619, $p = 0.0002$, FIG. 5A). Likewise in the primary squamous cell carcinoma data-set (TCGA, 2012), tumors with mutations in Brg1 had significantly higher expression of EGFR than Brg1 WT/EGFR WT tumors (FIG. 5B, $p = 0.003$). To explore the relationship between these mutations in the cell lines used in the experiments described above, Affymetrix arrays were used to compare gene expression signatures in the various genotypes. Overlap between differentially expressed genes in EGFR mutant lines vs. EGFR/Brg1 WT lines, and Brg1 mutant lines vs. EGFR/Brg1 WT lines was queried (FIG.

11A). Interestingly, it was found that EGFR was the first shared up-regulated gene, over-expressed in 4.5-fold in Brg1 mutant cell lines compared to EGFR/Brg1 WT lines and 4.7-fold in EGFR mutant cell lines compared to EGFR/Brg1 WT lines (FIG. 5C, Table 5, $p=0.014$). Immuno-fluorescence for EGFR confirmed that Brg1 mutant cell lines have high expression of cell surface bound EGFR compared to Brg1 WT cell lines, which have very little detectable EGFR staining (data not shown). Importantly, it was also confirmed that EZH2 levels are not different in the various genotypes (FIG. 11B). This observation was in agreement with RT PCR analysis for EZH2 and Suz12 in the cell lines showing that PRC2 gene levels were not different between sensitized and protected lines (FIG. 11C).

[0372] To further explore the genetic relationship between these two pathways, EGFR and Brg1 expression were manipulated via knock-down and over-expression viruses in protected and sensitized cell lines. In the Brg1 mutant cell line HCC15, which expressed high levels of WT EGFR, EGFR knock-down or re-introduction of Brg1 both decreased the amount of EGFR transcript (FIG. 18A, making bioreps for p values currently). In contrast, in the H460 line, which is WT for Brg1 and has little detectable EGFR expression, knock-down of Brg1 led to up-regulation of EGFR ($p=0.009$) and over-expression of WT EGFR led to a decrease in Brg1 transcript (FIG. 18B, $p=0.002$). Similar results were observed in several cell lines (FIG. 11D, making bioreps for p values currently).

[0373] Brg1 and EGFR genetically interact to control the sensitized phenotype. To learn if Brg1 and EGFR mutations are functional mediators, or merely predictors, of DZNep/etoposide response, we performed etoposide dose response curves on the Brg1 and EGFR knock-down and over-expression cell lines. For the HCC15 control cells, DZNep decreased the etoposide IC50 2-fold consistent with its sensitized phenotype ($p<0.0001$). However, both the EGFR knock-down and the Brg1 over-expressing HCC15 lines showed an increase in etoposide IC50, suggesting that the phenotype of the line was converted from sensitized to protected through loss of EGFR or re-introduction of Brg1 ($p=0.004$ and $p<0.0001$ respectively) (FIG. 23A, FIG. 12A). For the protected cell line H460, DZNep raised the etoposide IC50 2-fold in the control cells ($p=0.0005$), yet had no effect on etoposide IC50 when EGFR was over-expressed ($p=0.78$) or Brg1 was knocked-down ($p=0.2$) (FIG. 6B, FIG. 12B). These data demonstrate that the sensitized and protected phenotypes are partially reliant upon the levels of Brg1 and EGFR within the cells.

[0374] To learn if the changes in etoposide sensitivity of the Brg1- and EGFR-modulated DZNep-treated cell lines were attributable to cell cycle changes, 7AAD cell cycle analysis was performed. Etoposide treated HCC15 Brg1 over-expression cell lines showed a dramatically smaller accumulation in S phase in response to DZNep than the control shGFP cells (FIG. 6A). The HCC15 shEGFR line, which was significantly less sensitive to etoposide in the presence of DZNep, showed a decrease in S phase instead of the increase normally observed with this cell line. Add p57 qPCR data for HCC15 treated lines here (FIG. 6B). Consistent with the increase in etoposide sensitivity, the etoposide treated H460 shBrg1 and EGFR over-expression cell lines were enriched for S phase (FIG. 6C). This was in contrast to the control H460 cell line, which again showed accumulation in G1 and G2/M phases and a loss of S phase

in response to the dual therapy. Consistent with previous results, shGFP H460 cells showed a significant increase in p57 levels when treated with etoposide and DZNep (FIG. 6D). However, the levels of p57 were much higher in untreated H460 shBrg1 and EGFR oe cells, and the addition of etoposide and DZNep had no effect on p57 levels. Together these data demonstrate that EZH2 inhibition in the presence of etoposide altered cell cycle dynamics, and that this effect was dependent on the level of Brg1 and EGFR expression in the cells.

Discussion

[0375] Understanding the molecular determinants of epigenetic therapy responses will be essential for this new class of drugs to benefit cancer patients. It is demonstrated herein that EZH2 inhibition had differential effects on lung cancer chemotherapy sensitivities both in vitro and in vivo. Nearly half of the cell lines tested were sensitized to etoposide by EZH2 inhibition, while the other half were protected from the chemotherapy by the epigenetic therapy. These sensitized and protected phenotypes were mediated by differential cell cycle regulation and correlated with regulation of p57 expression. Within the panel of lung cancer cell lines, Brg1 and EGFR mutations were mutually exclusive and significantly predicted chemotherapy sensitization by EZH2 inhibition. Using primary tumor data and genetic manipulation of Brg1 and EGFR, it was demonstrated that these pathways genetically interact, and control response to chemotherapy and EZH2 inhibition. Changing the level of EGFR or Brg1 in a cell line can switch a line from sensitized to protected. These data link, for the first time, common mutations found in non-small cell lung cancer to responses to epigenetic therapy, and provide a strategy to predict which patients will benefit from dual EZH2 inhibitor and chemotherapy treatments or epigenetic therapy alone.

[0376] The differential effects of EZH2 inhibition on chemotherapy in vitro predicted response of the cell lines to therapies in vivo. Using two dosing schemes for DZNep, growth of xenografts of both sensitized and protected lines could be effectively prevented. For protected lines, a two week time course of DZNep alone was most effective, while for sensitized line, a one week time-course of DZNep and etoposide together significantly reduced tumor growth. Without wishing to be bound by theory, the observation that a second week of DZNep treatment increased H157 tumor burden suggests that in vivo, DZNep may cause etoposide challenged H157 cells to enter S phase just as they do in vitro. Of important clinical note, the combination of chemotherapy and DZNep for the protected line caused tumors to grow faster than tumors treated with either drug alone. These data reveal that planning dosing schemes to maximize cell cycle entry when chemotherapy is present is likely essential for this phenomenon to benefit patients. Furthermore, these data suggest that while EGFR and Brg1 mutation carriers may benefit from dual chemotherapy/EZH2 inhibitor strategies, the combination may be ineffective, or even dangerous, for patients with EGFR/Brg1 WT tumors, who can benefit from treatment with an EZH2 inhibitor alone.

[0377] In addition to EGFR and Brg1 mutations predicting chemo-sensitization by EZH2 inhibition, Brg1 inactivating mutations and EGFR activating mutations were significantly anti-correlated. By knocking down and over-expressing Brg1 and EGFR, a genetic interaction between the two

pathways became evident. Genetic antagonism between the Brg1 containing SWI/SNF complex and EZH2 has been demonstrated for lymphomas (Wilson et al., 2010), and genetic interaction between Brg1 and EGFR has been shown in *Drosophila* (Herr et al., 2010). However, to the inventors' knowledge, this is the first reported data to implicate cooperation of these three pathways in lung cancer. Together these data suggest that one main effect of Brg1 inactivation is de-repression of EGFR signaling. Because B-Raf mutations also predicted chemo-sensitization via EZH2 inhibition, it is likely that activated MAPK signaling is involved in the sensitized phenotype. However, more than half of the protected cell lines harbor mutations in K-Ras, suggesting that expression of mutant K-Ras causes effects unique to activation of the EGFR-Raf-MAPK pathway in respect to the chemo-sensitization phenotype. Cells with high levels of EGFR react to EZH2 inhibition differently than cells that have lower levels of EGFR. The data described herein indicate that the cell cycle inhibitor p57 is a mediator in the lung cancer cell therapy protection phenotype. Whereas p57 mutations have not been reported in lung cancer, previous reports have characterized p57 as a target of PRC-mediated repression (Yang et al., 2009, Guo et al., 2011, Zacharek et al., 2011). Thus, de-repression of direct targets of Polycomb, such as p57, can be critical in regulating lung cancer response to targeted therapies.

[0378] These data indicate several possible combination therapies that may also be useful to combat lung cancers. In Brg1 and EGFR mutant cell lines, EZH2 inhibition caused sensitization to the TopoII inhibitors etoposide and doxorubicin. Double strand breaks (DSBs) caused by these agents cause formation of BRCA1 foci and can be repaired by homologous recombination if the sister chromatid can be found, or non-homologous end-joining if the homologous template is not available. BRCA1 mutant breast cancers express more EZH2, and are therefore intrinsically more sensitive to EZH2 inhibition (Puppe et al., 2009). PARP inhibitors, which lead to double strand breaks by preventing single strand break repair, are synthetically lethal with CDK1 inhibition, in part because BRCA1 requires CDK1 phosphorylation to function (Johnson et al., 2011). Because another substrate for CDK1 is EZH2 (Wei et al., 2010, Chen et al., 2010), the data indicate that inhibition of EZH2 can also contribute to the synthetic lethality of PARP and CDK1 inhibitors. Furthermore, because of the reliance of BRCA1 mutant tumors on EZH2, tumors treated with CDK1 inhibitors may be exquisitely sensitive to drugs such as DZNep. Therefore, a combination of all three therapies (DSB initiation, CDK1 inhibition and EZH2 inhibition) can be effective.

[0379] Therapeutic strategies targeting epigenetic factors have risen as promising new candidates for lung cancer and other difficult to treat cancers. Whereas many studies suggest that epigenetic therapy may offer a treatment avenue for broader patient groups than drugs targeting specific oncogenic changes, the work described herein indicates that selection of patient subsets will be critical in designing clinical trials and treatment strategies for drugs inspired by epigenetics. The initial cell culture approach examining dual treatment responses has revealed important, predictable in vivo drug sensitivities driven by lung cancer oncogenotype. Interestingly, the histological subtype of NSCLC was not a major factor in determining response to EZH2 inhibition.

Experimental Procedures

[0380] Cell Lines and small hairpins. Cell lines used are listed in Table 1, Table 4, and FIG. 29. All cell lines were maintained in RPMI 1640 media with 10% fetal bovine serum, 4 mM L-glutamine and penicillin/streptomycin at 37° C., 5% CO₂. The pLKO.1 EZH2 shRNA construct clone TRCN0000040076 was purchased from SIGMA and the shGFP plasmid 12273 is available on Addgene. Both shBrg1 and the matched empty vector were provided by the Smale lab (Ramirez et al., 2006), the Brg1 over-expression plasmid 19148 was purchased through Addgene, and the shEGFR and EGFR WT overexpression constructs were provided by the Janne lab (Engelman et al., 2006). Small hairpin sequences:

```
GFP: (SEQ ID NO: 15)
GCCC (GCAAGCTGACCCCTGAAGTTCAT) TCAAGAG
(ATGAACTTCAGGGTCAGCTTGC) TTTT

EZH2: (SEQ ID NO: 16)
CCGG (CGGAAATCTTAAACCAAGAAT) CTCGAG
(ATTCTTGGTTTAAGATTTCG) TTTT

EGFR: (SEQ ID NO: 17)
CCGG (GCTGAGAATGTGGAATACCTA) CTCGAG
(TAGGTATTCACATTCTCAGC) TTTT

BRG1: (SEQ ID NO: 18)
TTTG (TGGATAAGCAGCACAAGATT) TCAAGAG
(AATCTTCTGCTGCTTCTCCA) TTTT.
```

[0381] Drugs. Etoposide, cis-platinum, and doxorubicin (SIGMA) were diluted to a stock of 100 mM in DMSO for all cell culture experiments. DZNep was diluted in DMSO to a stock of 10 mM. All stocks were diluted in DMSO to 1000× concentration prior to addition into media at 2× concentration and final dilution onto plated cells 1:1.

[0382] Cytotox Assays. Cell lines were dissociated, counted and plated at 5000 cells per well in flat bottom opaque tissue culture treated 96 well plates (CytoOne). Edge wells were filled with PBS. The following day, 2× drug diluted in media was added to each well such that the well then contained 100 ul media with 1× drug concentration. After 4 days, Cell Titer Glo™ (Promega) was added and luminescence was read on a BioTec plate reader to determine relative cell number in each well. Data were averaged for triplicate or quadruplicate technical replicates, normalized to the untreated wells and input into GraphPad Prism software to extrapolate IC₅₀.

[0383] Cell Cycle Flow Cytometry. Cell lines were plated at 1.5×10⁶ cells per 10 cm plate and treated with drug for 4 days. Cells were then dissociated, fixed with 100% ice cold Ethanol for at least 2 hours, incubated for 30 minutes with 1 mg/mL RNase A and resuspended in 20 µg/mL 7AAD. 30,000 events were collected on the BD FacsCalibur™ and analyzed with the ModFit LT™ software. In cases where etoposide treated cells had such abnormal ploidy that ModFit LT™ was unable to correctly model the cell cycle, the data was analyzed manually on FlowJo™ software by

setting the gates on the etoposide sample and copying them to the etoposide+DZNep and etoposide shEZH2 samples.

[0384] Quantitative RT PCR. RNA from treated cell lines was extracted from using Trizol. Equal amounts of RNA were incubated with DNase I before cDNA was made using the SuperScript III™ kit (Invitrogen). Relative gene expression was assayed with Sybr green on the StepOnePlus™ Real-Time PCR System (Applied Biosystems). Relative expression was calculated by Gene of Interest (Ctreference-Ctexperimental)-CYPA(Ctreference-Ctexperimental) and graphed on the log2 scale. For all experiments, the reference sample was a matched untreated or control transduced cell line. Primer sequences:

CYPA F	TCATCTGCACTGCCAAGACTG	(SEQ ID NO: 19)
CYPA R	CATGCCTTCTTTCACTTTGCC	(SEQ ID NO: 20)
EZH2 F	AGGAGTTTGCTGCTGCTCTC	(SEQ ID NO: 21)
EZH2 R	CCGAGAATTGCTTTCAGAGG	(SEQ ID NO: 22)
BRG1 F	AGCGATGACGTCTCTGAGGT	(SEQ ID NO: 11)
BRG1 R	GTACAGGGACACCAGCCACT	(SEQ ID NO: 12)
EGFR F	TAACAAGCTCACGCAGTTGG	(SEQ ID NO: 13)
EGFR R	GTTGAGGGCAATGAGGACAT	(SEQ ID NO: 14)
SUZ12 F	TACGGCTCTATTGCCAAAC	(SEQ ID NO: 23)
SUZ12 R	TGCTTCAGTTTGTTCCTTG	(SEQ ID NO: 24)
P57 F	GGCCTCTGATCTCCGATTTTC	(SEQ ID NO: 25)
P57 R	TGGGCTCTAAATTGGCTCAC	(SEQ ID NO: 26)
P21 F	ATGAAATTCACCCCTTTCC	(SEQ ID NO: 27)
P21 R	CCCTAGGCTGTGCTCACTTC	(SEQ ID NO: 28)
P27 F	CCGGCTAACTCTGAGGACAC	(SEQ ID NO: 29)
P27 R	CGAGCTGTTTACGTTTGACG	(SEQ ID NO: 30)

[0385] Mice. Cell lines were dissociated into single cells, counted and resuspended at 1×10⁶ cells per 2504 of 1:1 media/matrigel (BD). 8- to 16-week-old Foxnlnu (nude) mice (Harlan) were injected subcutaneously with 1×10⁶ cells in four spots on flanks. Chemo and DZNep were administered from day 12 to day 17 post injections, Etoposide: 20 mg/kg/day i.p. in corn oil once per day for 5 consecutive days, and DZNep 2 mg/kg/day i.p. in corn oil twice per week for 1 week (dosing scheme A) OR 1 mg/kg/day i.p. in corn oil twice per week for 2 weeks (dosing scheme B). Tumor growth was measured every other day by caliper.

[0386] Generation of the EZH2 Co-Expressed Gene Signature. Oncomine was used to query the top 20 genes co-expressed with EZH2 in all datasets containing human non-small cell lung cancer samples and co-expression data. The 9 datasets correspond to Beer et al., 2008, Director's Challenge, 2008, Garber et al., 2001, Gordon et al., 2002, Landi et al., 2008, Rohrbeck et al., 2008, Su et al., 2001, Yu et al., 2008, and Kim Lung unpublished. Of the 180 probes, 64 were redundant, leading to a list of 116 genes highly co-expressed with EZH2. Because data sets on Oncomine were from various microarray platforms, the gene list was then used to generate a probe list for the 116 genes corre-

sponding to probes on the U133A Affymetrix array using the batch query function on the NetAffx website (available on the world wide web at

[www\(dot\)Affymetrix\(dot\)com\(backslash\)analysis\(backslash\)index\(dot\)affx](http://www(dot)Affymetrix(dot)com(backslash)analysis(backslash)index(dot)affx)).

[0387] Kaplan Meier Analysis. Raw gene expression data from human lung adenocarcinoma samples were obtained (Director's Challenge). Probe intensities from the Affymetrix U133A platform used in these studies were normalized and modeled using dChip software (Li C and Wong W H. PNAS 2001 Vol. 98, 31-36; which is incorporated by reference herein in its entirety). Kaplan-Meier survival analyses were implemented after the samples were hierarchically clustered using centroid linkage into two risk groups using the EZH2 co-expressed gene signature. Differences of the survival risk between the two risk groups were assessed using the Mantel-Haenszel log rank test. The larger area between the two risk groups and its associated smaller p value from the Mantel-Haenszel log rank test implicate a better classification model.

[0388] Cell Line Micro-Array. Array quality was assessed using the R/Bioconductor package (see, e.g. Gentleman R. C. et al. Genome Biol. 2004 5(10): R80; which is incorporated by reference herein in its entirety). Raw CEL files from U133A Affymetrix arrays were processed using the robust multiarray average (RMA) algorithm (Irizarry et al., 2003). To identify genes correlating with the phenotypic groups, limma (Smyth, 2004) was used to fit a statistical linear model to the data and then tested for differential gene expression in the three groups, WT: H460, H441, H2122, H2009, Calu6, HCC95, EGFR mutant: H1650, HCC827, HCC4006, H1975, H3255, PC9, Brg1 mutant: A549, H1299, H157, H2126, H522, HCC15. Results were adjusted for multiple testing using the Benjamini and Hochberg (BH) method (Benjamini and Hochberg, 1995), and significance was determined using a False-Discovery-Rate cutoff of less than 5%.

[0389] Statistical Analysis. Except where indicated, Student's t-test was used to compare measurements between 2 conditions. Unless noted otherwise, pooled data is represented by the mean and standard error. p-values less than 0.05 were considered significant.

[0390] Western blot. For Western blotting, 25 µg of protein extract per sample was denatured with heat and reducing agents, separated on a 4-12% acrylimide gel and transferred to nitrocellulose. Antibodies used for Western blotting were: EZH2 (Cell Signalling), β-actin (AbCAM) and H3K27me3 (Upstate).

REFERENCES

- [0391]** Each of the following references is incorporated by reference herein in its entirety.
- [0392]** 1. Avan, A., Crea, F, Paolicchi, E, Funel, N., Galvani, E. Marquez, V. E., Honeywell, R. J., Danesi, R., Peters, G. J., and Giovannetti, E. (2012). Molecular mechanisms involved in the synergistic interaction of the EZH2 inhibitor 3-deazaneplanocin A (DZNep) with gemcitabine in pancreatic cancer cells. Mol. Cancer Ther. Epub.
- [0393]** 2. Baldwin, E. L. and Osherooff, N. (2005). Etoposide, topoisomerase II and cancer. Curr. Med. Chem. Anticancer Agents 5, 363-372.
- [0394]** 3. Baylin, S. B. (2011). Resistance, epigenetics and the cancer ecosystem. Nat. Med. 17, 288-289.

- [0395] 4. Beer, D. G., S. L. Kardia, S. L., C. C. Huang, C. C., T. J. Giordano, T. J., A. M. Levin, A. M., D. E. Misek, D. E., L. Lin, L., G. Chen, G., T. G. Gharib, T. G., D. G. Thomas, D. G. et al. (2002). Gene-expression profiles predict survival of patients with lung adenocarcinoma. *Nat. Med.* 8, 816-824.
- [0396] 5. Ben-Porath, I., Thomson, M. W., Carey, V. J., Ge, R., Bell, G. W., Regev, A., and Weinberg, R. A. (2008). An embryonic stem cell-like gene expression signature in poorly differentiated aggressive human tumors. *Nat. Gen.* 40, 499-507.
- [0397] 6. Bracken, A. P. and Helin, K. 2009. Polycomb group proteins: navigators of lineage pathways led astray in cancer. *Nat. Rev. Cancer* 9, 7730-784.
- [0398] 7. Breuer, R. H. J., Snijders, P. J. F., Smit, E. F., Sutedja, T. G., Sewalt, R. G. A. B., Otte, A. P., van Kemenade, F. J., Postmus, P. E., Meijer, C. J. L. M., and Raaphorst, F. M. (2004). Increased expression of the *Ezh2* polycomb group gene in *Bmi1*-positive neoplastic cells during bronchial carcinogenesis. *Neoplasia* 6, 736-743.
- [0399] 8. Chen, S., Bohrer, L. R., Rai, A. N., Pan, Y., Gan, L., Zhou, X., Bagchi, A., Simon, J. A. and Huang, H. (2010). Cyclin-dependent kinases regulate epigenetic gene silencing through phosphorylation of *EZH2*. *Nat. Cell Biol.* 12, 1108-1114.
- [0400] 9. Chou, D. M., Adamson, B., Dephoure, N. E., Tan, X., Nottke, A. C., Hurov, K. E., Gygi, S. P., Colaiacovo, M. P., and Elledge, S. J. (2010). A chromatin localization screen reveals poly (ADP ribose)-regulated recruitment of the repressive Polycomb and NuRD complexes to sites of DNA damage. *Proc. Natl. Acad. Sci. USA* 107, 18475-18480.
- [0401] 10. Crea, F., Paolicchi, E., Marquez, V. E. and Danesi, R. (2011a). Polycomb genes and cancer: Time for clinical application? *Crit. Rev. Oncol./Hematol.* Epub.
- [0402] 11. Crea, F., Hurt, E. M., Mathews, L. A., Cabarcas, S. M., Sun, L., Marquez, V. E., Danesi, R., and Farrar, W. L. (2011b). Pharmacologic disruption of Polycomb Repressive Complex 2 inhibits tumorigenicity and tumor progression in prostate cancer. *Mol. Cancer* 10, Epub.
- [0403] 12. Curtis, S. J., Sinkevicius, K. W., Li, D., Lau, A. N., Roach, R. R., Zamponi, R., Woollenden, A. E., Kirsch, D. G., Wong, K. K., and Kim, C. F. (2010). Primary tumor genotype is an important determinant in identification of lung cancer propagating cells. *Cell Stem Cell* 7, 127-133.
- [0404] 13. Director's Challenge Consortium for the Molecular Classification of Lung Adenocarcinoma: Shedden, K., Taylor, J. M. G., Enkemann, S. A., Tsao, M., Yeatman, T. J., Gerald, W. L., Eschrich, S., Jurisica, I., Giordano, T. J., Misek, D. E. et al. (2008). Gene expression-based survival prediction in lung adenocarcinoma: a multi-site, blinded validation study. *Nat. Med.* 14, 822-827.
- [0405] 14. Dovey, J. S., Zacharek, S. J., Kim, C. F., and Lees, J. A. (2008). *Bmi1* is critical for lung tumorigenesis and bronchioalveolar stem cell expansion. *Proc. Natl. Acad. Sci. USA* 105, 11857-11862.
- [0406] 15. Engelman, J. A., Mukohara, T., Zejnullahu, K., Lifshits, E., Borras, A. M., Gale, C. M., Naumov, G. N., Yeap, B. Y., Jarrell, E., Sun, J. et al. (2006). Allelic dilution obscures detection of a biologically significant resistance mutation in *EGFR*-amplified lung cancer. *J. Clin. Invest.* 116, 2695-26706.
- [0407] 16. Garber, M. E., Troyanskaya, O. G., Schluens, K., Petersen, S., Thaessler, Z., Pacyna-Gengelbach, M., van de Rijn, M., Rosen, G. D., Perou, C. M., Whyte, R. I. et al. (2001). Diversity of gene expression in adenocarcinoma of the lung. *Proc. Natl. Acad. Sci. U S A* 98, 13784-13789.
- [0408] 17. Gordon G. J., Jensen, R. V., Hsiao, L. L., Gullans, S. R., Blumenstock, J. E., Ramaswamy, S., Richard, W. D., Sugarbaker, D. J. and Bueno, R. (2002). Translation of microarray data into clinically relevant cancer diagnostic tests using gene expression ratios in lung cancer and mesothelioma. *Cancer Res.* 62, 4963-4967.
- [0409] 18. Hassan, K. A., Chen, G., Kalemkerian, G. P., Wicha, M. S., and Beer, D. G. (2009). An embryonic stem cell-like signature identifies poorly differentiated lung adenocarcinoma but not squamous cell carcinoma. *Clin. Cancer Res.* 15, 6386-6390.
- [0410] 19. Herbst, R. S., and Heymach J. V. (2008). Lung cancer. *N. Engl. J. Med.* 359, 1367-1380.
- [0411] 20. Herr, A., McKenzie, L., Suryadinata, R., Sadowski, M., Parsons, L. M., Sarcevic, B., and Richardson, H. E. (2010). Geminin and Brahma act antagonistically to regulate *EGFR*-Ras-MAPK signaling in *Drosophila*. *Dev. Biol.* 344, 36-51.
- [0412] 21. Huqun, Ishikawa, R., Zhang, J., Miyazawa, H., Goto, Y., Shimizu, Y., Hagiwara, K., and Koyama, N. (2012) Enhancer of Zeste Homolog 2 is a novel prognostic biomarker in nonsmall cell lung cancer. *Cancer* 118, 1599-1606.
- [0413] 22. Iliopoulos, D., Lindahl-Alten, M., Polyarchou, C., Hirsch, H. A., Tschlis, P. N., and Struhl, K. (2010). Loss of miR-200 inhibition of *Suz12* leads to Polycomb-mediated repression required for the formation and maintenance of cancer stem cells. *Mol. Cell* 39, 761-772.
- [0414] 23. Imielinski, M., Berger, A. H., Hammerman, P. S., Hernandez, B., Pugh, T. J., Hodis, E., Cho, J., Suh, J., Capelletti, M., Sivachenko, A., et al. (2012). Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. *Cell* 150, 1107-1120.
- [0415] 24. Ji, H., Wang, Z., Perera, S. A., Li, D., Liang, M. C., Zaghlul, S., McNamara, K., Chen, L., Albert, M., Sun, Y. et al. (2007). Mutations in *BRAF* and *KRAS* converge on activation of the mitogen-activated protein kinase pathway in lung cancer mouse models. *Cancer Res.* 67, 4933-4939.
- [0416] 25. Johnson, N., Li, Y., Walton, Z. E., Cheng, K. A., Li, D., Rodig, S., Moreau, L. A., Unitt, C., Bronson, R. T., Thomas, H. D., Newell, D. R. et al. (2011). Compromised *CDK1* activity sensitizes *BRCA*-proficient cancers to *PARP* inhibition. *Nat. Med.* 17, 875-883.
- [0417] 26. Kikuchi, J., Kinoshita, I., Shimizu, Y., Kikuchi, E., Konishi, J., Oizumi, S., Kaga, K., Matsuno, Y., Nishimura, M., and Dosaka-Akita, H. (2010). Distinctive expression of the Polycomb group proteins *Bmi1* polycomb ring finger oncogene and enhancer of zeste homolog 2 in nonsmall cell lung cancers and their clinical and clinicopathologic significance. *Cancer* 116, 3015-3024.
- [0418] 27. Kikuchi, J., Takashina, T., Kinoshita, I., Kikuchi, E., Shimizu, Y., Sakakibara-Konishi, J., Oizumi, S., Marquez, V. E., Nishimura, M., and Dosaka-Akita, H. (2012). Epigenetic therapy with 3-deazaneplanocin A, an

- inhibitor of the histone methyltransferase EZH2, inhibits growth of non-small cell lung cancer cells. *Lung Cancer Epub*.
- [0419] 28. Landi, M. T., Dracheva, T., Rotunno, M., Figueroa, J. D., Liu, H., Dasgupta, A., Mann, F. E., Fukuoka, J., Hames, M., Bergen, A. W. et al. (2008). Gene expression signature of cigarette smoking and its role in lung adenocarcinoma development and survival. *PLoS One* 3, e1651.
- [0420] 29. Lee, T. I., Jenner, R. G., Boyer, L. A., Guenther, M. G., Levine, S. S., Kumar, R. M., Chevalier, B., Johnstone, S. E., Cole, M. F., Isono, K. et al. (2006). Control of developmental regulators by polycomb in human embryonic stem cells. *Cell* 125, 301-313.
- [0421] 30. Medina, P. P., Romero, O. A., Kohno, T., Montuenga, L. M., Pio, R., Yokota, J., and Sanchez-Cespedes, M. (2008). Frequent BRG1/SMARCA4-inactivating mutations in human lung cancer cell lines. *Hum. Mutat.* 29, 617-622.
- [0422] 31. Min, J., Zaslavsky, A., Fedele, G., McLaughlin, S. K., Reczek, E. E., De Raedt, T., Guney, I., Strohlic, D. E., Macconail, L. E., Beroukhi, R. et al. (2010). An oncogene-tumor suppressor cascade drives metastatic prostate cancer by coordinately activating Ras and nuclear factor-kappaB. *Nat. Med.* 16, 286-294.
- [0423] 32. Ougolkov, A. V., Bilim, V. N., and Billadeau, D. D. (2008). Regulation of pancreatic tumor cell proliferation and chemoresistance by the histone methyltransferase Enhancer of Zeste Homologue 2. *Clin. Cancer Res.* 14, 6790-6796.
- [0424] 33. Puppe, J., Drost, R., Liu, X., Joosse, S. A., Evers, B., Cornelissen-Stijger, P., Nederlof, P., Qiang, Y., Jonkers, J., van Lohuizen, M., and Pietersen, A. M. (2009). BRCA1-deficient mammary tumor cells are dependent on EZH2 expression and sensitive to Polycomb Repressive Complex 2-inhibitor 3-deazaneplanocin A. *Breast Cancer Res.* 11, R36.
- [0425] 34. Ramirez-Carrozzi, V. R., Nazarian, A. A., Li, C. C., Gore, S. L., Sridharan, R., Imbalzano, A. N. and Smale, S. T. (2006). Selective and antagonistic functions of SWI/SNF and Mi-2beta nucleosome remodeling complexes during an inflammatory response. *Genes Dev.* 20, 282-296.
- [0426] 35. Rodriguez-Nieto, S. A., Canada, A., Pros, E., Pinto, A. I., Torres-Lanzas, J., Lopez-Rios, F., Sanchez-Verde, L., Pisano, D. G. and Sanchez-Cespedes, M. (2011). Massive parallel DNA pyrosequencing analysis of the tumor suppressor BRG1/SMARCA4 in lung primary tumors. *Hum. Mutat.* 32, E1999-2017.
- [0427] 36. Rohrbeck, A., Neukirchen, J., Roskopf, M., Pardillos, G. G., Geddert, H., Schwalen, A., Gabbert, H. E., von Haeseler, A., Pitschke, G., Schott, M. et al. (2008). Gene expression profiling for molecular distinction and characterization of laser captured primary lung cancers. *J. Transl. Med.* 6, e69.
- [0428] 37. Sauvageau, M. and Sauvageau, G. (2010). Polycomb group proteins: Multi-faceted regulators of somatic stem cells and cancer. *Cell Stem Cell* 7, 299-313.
- [0429] 38. Simon, J. A., and Lange, C. A. (2008). Roles of EZH2 histone methyltransferase in cancer epigenetics. *Mutat. Res.* 647, 21-29.
- [0430] 39. Simon, J. A. and Kingston, R. E. (2009). Mechanisms of Polycomb gene silencing: knowns and unknowns. *Nat. Rev. Mol. Cell Biol.* 10, 697-708.
- [0431] 40. Smyth, G. K. (2004). Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* 3.
- [0432] 41. Spamann, A., and van Lohuizen, M. (2006). Polycomb silencers control cell fate, development and cancer. *Nat. Rev. Genet.* 6, 846-856.
- [0433] 42. Su, A. I., Welsh, J. B., Sapinoso, L. M., Kern, S. G., Dimitrov, P., Lapp, H., Schultz, P. G., Powell, S. M., Moskaluk, C. A., Frierson, H. F., and Hampton, G. M. (2001). Molecular classification of human carcinomas by use of gene expression signatures. *Cancer Res.* 61, 7388-7393.
- [0434] 43. Takawa, M., Masuda, K., Kunizaki, M., Daigo, Y., Takagi, K., Iwai, Y., Cho, H., Toyokawa, G., Yamne, Y., Maejima, K. et al. (2011). Validation of the histone methyltransferase EZH2 as a therapeutic target for various types of human cancer and as a prognostic marker. *Cancer Sci.* 102, 1298-1305.
- [0435] 44. Tan, J., Yang, X., Zhuang, L., Jiang, X., Chen, W., Lee, P. L., Karuturi, R. K., Tan, P. B., Liu, E. T. and Yu, Q. (2007). Pharmacologic disruption of Polycomb-repressive complex 2-mediated gene repression selectively induces apoptosis in cancer cells. *Genes Dev.* 21, 1050-1063.
- [0436] 45. The Cancer Genome Atlas Research Network. (2012). Comprehensive genomic characterization of squamous cell lung cancers. *Nature Epub*.
- [0437] 46. Wei, Y., Chen, Y. H., Li, L. Y., Lang, J., Yeh, S. P., Shi, B., Yang, C. C., Yang, J. Y., Lin, C. Y., Lai, C. C., and Hung, M. C. (2010). CDK1-dependent phosphorylation of EZH2 suppresses methylation of H3K27 and promotes osteogenic differentiation of human mesenchymal stem cells. *Nat. Cell Biol.* 13, 87-94.
- [0438] 47. Wilson, B. G., Wang, X., Shen, X., McKenna, E. S., Lemieux, M. E., Cho, Y. J., Koellhoffer, E. C., Pomeroy, S. L., Orkin, S. H., and Roberts, C. W. (2010). Epigenetic antagonism between polycomb and SWI/SNF complexes during oncogenic transformation. *Cancer Cell.* 18, 316-328.
- [0439] 48. Wu, Z., Lee, S. T., Qiao, Y., Li, Z., Lee, P. L., Lee, Y. J., Jiang, X., Tan, J., Aau, M., Lim, C. Z., and Yu, Q. (2011). Polycomb protein EZH2 regulates cancer cell fate decision in response to DNA damage repair. *Cell Death Differ.* 18, 1771-1779.
- [0440] 49. Yamamoto, H., Shigematsu, H., Nomura, M., Lockwood, W. W., Sato, M., Okumura, N., Soh, J., Suzuki, M., and Wistuba, L. I., Fong, K. M. et al. (2008). PIK3CA mutations and copy number gains in human lung cancers. *Cancer Res.* 68, 6913-6921.
- [0441] 50. Yu K., Ganesan, K., Tan, L. K., Laban, M., Wu, J., Zhao, X. D., Li, H., Leung, C. H., Zhu, Y., Wei, C. L. et al. (2008). A precisely regulated gene expression cassette potently modulates metastasis and survival in multiple solid cancers. *PLoS Genet.* 4, e1000129.

TABLE 1

B-Raf, EGFR, and BRG1 Annotation			
Cell Line	B-Raf	EGFR	BRG1
HCC2450	WT	WT	n/a
PC9	WT	E746_A750del	n/a
H1395	G469A	WT	WT

TABLE 1-continued

B-Raf, EGFR, and BRG1 Annotation			
Cell Line	B-Raf	EGFR	BRG1
H522	WT	WT	P270*
A549	WT	WT	Q729*
Calu1	WT	WT	WT
H2126	WT	WT	W764R
H157	WT	WT	T58*
H1975	WT	T290M, L858R	WT
HCC15	WT	WT	M272*
H2030	WT	WT	Null
H1650	WT	E746X	WT
HCC4006	WT	L747_A749del	WT
H2087	L597V	WT	WT
HCC827	WT	E746_A750del	WT
H1299	WT	WT	Y560*
H322	WT	WT	WT
HCC95	WT	WT	WT
H2009	WT	WT	WT
Sw1573	WT	WT	WT
H2122	WT	WT	WT
H1819	WT	WT	L1085*
H520	WT	WT	WT
Calu6	WT	WT	WT
Calu3	WT	WT	WT
H460	WT	WT	WT
H3255	WT	L858R	WT
H23	WT	WT	K1533N
H441	WT	WT	WT

TABLE 2

EZH2 Co-Expression Gene Signature. Top 116 genes co-expressed with EZH2 across 9 Oncomine studies (Beer et al., 2008, Director's Challenge, 2008, Garber et al., 2001, Gordon et al., 2002, Landi et al., 2008, Rohrbeck et al., 2008, Su et al., 2001, Vu et al., 2008, and Kim Lung unpublished)	
Gene Symbol	Description
ALS2CR4	amyotrophic lateral sclerosis 2 (juvenile) chromosome region, candidate 4
ATIC	5-aminoimidazole-4-carboxamide ribonucleotide formyltransferase/IMP cyclohydrolase
ATP2A2	ATPase, Ca++ transporting, cardiac muscle, slow twitch 2
AURKA	aurora kinase A
BIRC5	baculoviral LAP repeat-containing 5
BOP1	block of proliferation 1
BUB1	budding uninhibited by benzimidazoles 1 homolog (yeast)
BUB1B	budding uninhibited by benzimidazoles 1 homolog beta (yeast)
C13orf34	chromosome 13 open reading frame 34
C1orf112	chromosome 1 open reading frame 112
CCNA2	cyclin A2
CCNB1	cyclin B1
CCNB2	cyclin B2
CDC20	cell division cycle 20 homolog (<i>S. cerevisiae</i>)
CDC25A	cell division cycle 25 homolog A (<i>S. pombe</i>)
CDC6	cell division cycle 6 homolog (<i>S. cerevisiae</i>)
CDC7	cell division cycle 7 homolog (<i>S. cerevisiae</i>)
CDCA3	cell division cycle associated 3
CDCA8	cell division cycle associated 8
CDK1	cyclin-dependent kinase 1
CDKN3	cyclin-dependent kinase inhibitor 3
CENPE	centromere protein E, 312 kDa
CENPF	centromere protein F, 350/400 kDa (mitosin)
CENPM	centromere protein M
CENPW	centromere protein W
CHEK1	CHK1 checkpoint homolog (<i>S. pombe</i>)
CKAP2	cytoskeleton associated protein 2
CKS1B	CDC28 protein kinase regulatory subunit 1B

TABLE 2-continued

EZH2 Co-Expression Gene Signature. Top 116 genes co-expressed with EZH2 across 9 Oncomine studies (Beer et al., 2008, Director's Challenge, 2008, Garber et al., 2001, Gordon et al., 2002, Landi et al., 2008, Rohrbeck et al., 2008, Su et al., 2001, Vu et al., 2008, and Kim Lung unpublished)	
Gene Symbol	Description
CKS2	CDC28 protein kinase regulatory subunit 2
CSE1L	CSE1 chromosome segregation 1-like (yeast)
CSTF1	cleavage stimulation factor, 3' pre-RNA, subunit 1, 50 kDa
CTPS	CTP synthase
DBF4	DBF4 homolog (<i>S. cerevisiae</i>)
DBN1	drebrin 1
DEPDC1	DEP domain containing 1
DLGAP5	discs, large (<i>Drosophila</i>) homolog-associated protein 5
DTL	denticless homolog (<i>Drosophila</i>)
ECT2	Epithelial cell transforming sequence 2 oncogene
ESCO2	establishment of cohesion 1 homolog 2 (<i>S. cerevisiae</i>)
ESPL1	extra spindle pole bodies homolog 1 (<i>S. cerevisiae</i>)
EXO1	exonuclease 1
EZH2	enhancer of zeste homolog 2 (<i>Drosophila</i>)
FANCI	Fanconi anemia, complementation group I
FBXO5	F-box protein 5
FEN1	flap structure-specific endonuclease 1
FOXM1	forkhead box M1
GIN51	GIN5 complex subunit 1 (Psf1 homolog)
GIN52	GIN5 complex subunit 2 (Psf2 homolog)
GMNN	geminin, DNA replication inhibitor
GOT1	glutamic-oxaloacetic transaminase 1, soluble (aspartate aminotransferase 1)
GPD2	glycerol-3-phosphate dehydrogenase 2 (mitochondrial)
H2AFX	H2A histone family, member X
H2AFZ	H2A histone family, member Z
HMGA1	high mobility group AT-hook 1
HMGB2	high-mobility group box 2
HMMR	hyaluronan-mediated motility receptor (RHAMM)
HN1	hematological and neurological expressed 1
KIAA0101	KIAA0101
KIF11	kinesin family member 11
KIF14	kinesin family member 14
KIF15	kinesin family member 15
KIF18A	kinesin family member 18A
KIF18B	kinesin family member 18B
KIF20A	kinesin family member 20A
KIF20B	kinesin family member 20B
KIF23	kinesin family member 23
KIF2C	kinesin family member 2C
KIFC1	kinesin family member C1
KNTC1	kinetochore associated 1
KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)
LMNB1	lamin B1
LMNB2	lamin B2
MAD2L1	MAD2 mitotic arrest deficient-like 1 (yeast)
MCM10	minichromosome maintenance complex component 10
MCM7	minichromosome maintenance complex component 7
MKI67	antigen identified by monoclonal antibody Ki-67
MLF1IP	MLF1 interacting protein
MTHFD2	methylenetetrahydrofolate dehydrogenase (NADP+ dependent) 2,
NCAPG	non-SMC condensin I complex, subunit G
NCAPG2	non-SMC condensin II complex, subunit G2
NCAPH	non-SMC condensin I complex, subunit H
NDC80	NDC80 homolog, kinetochore complex component (<i>S. cerevisiae</i>)
NUP205	nucleoporin 205 kDa
OLA1	Obg-like ATPase 1
PAICS	phosphoribosylaminoimidazole carboxylase
PBK	PDZ binding kinase
PCNA	proliferating cell nuclear antigen
PFN2	profilin 2
PLK4	polo-like kinase 4
POLA1	polymerase (DNA directed), alpha 1, catalytic subunit
POLE2	polymerase (DNA directed), epsilon 2 (p59 subunit)
PPAT	Phosphoribosyl pyrophosphate amidotransferase
PRIM1	primase, DNA, polypeptide 1 (49 kDa)

TABLE 2-continued

EZH2 Co-Expression Gene Signature. Top 116 genes co-expressed with EZH2 across 9 Oncomine studies (Beer et al., 2008, Director's Challenge, 2008, Garber et al., 2001, Gordon et al., 2002, Landi et al., 2008, Rohrbeck et al., 2008, Su et al., 2001, Vu et al., 2008, and Kim Lung unpublished)	
Gene Symbol	Description
PSMD14	proteasome (prosome, macropain) 26S subunit, non-ATPase, 14
PTTG1	pituitary tumor-transforming 1
RFC4	replication factor C (activator 1) 4, 37 kDa
SLC7A5	solute carrier family 7 (cationic amino acid transporter, y+ system), member 5
SMC2	structural maintenance of chromosomes 2
SMC4	structural maintenance of chromosomes 4
SPAG5	sperm associated antigen 5
SRM	spermidine synthase
STIL	SCL/TAL1 interrupting locus
STMN1	stathmin 1
TACC3	transforming, acidic coiled-coil containing protein 3

TABLE 2-continued

EZH2 Co-Expression Gene Signature. Top 116 genes co-expressed with EZH2 across 9 Oncomine studies (Beer et al., 2008, Director's Challenge, 2008, Garber et al., 2001, Gordon et al., 2002, Landi et al., 2008, Rohrbeck et al., 2008, Su et al., 2001, Vu et al., 2008, and Kim Lung unpublished)	
Gene Symbol	Description
TFRC	Transferrin receptor (p90, CD71)
TMEM106C	transmembrane protein 106C
TOP2A	Topoisomerase (DNA) II alpha 170 kDa
TOPBP1	topoisomerase (DNA) II binding protein 1
TPX2	TPX2, microtubule-associated, homolog (<i>Xenopus laevis</i>)
TRIP13	thyroid hormone receptor interactor 13
TROAP	trophinin associated protein (tastin)
TYMS	thymidylate synthetase
UBE2C	ubiquitin-conjugating enzyme E2C
UBE2S	ubiquitin-conjugating enzyme E2S
UNG	uracil-DNA glycosylase
WDR12	WD repeat domain 12

TABLE 3

Gene Function enrichment analysis of EZH2 expressed gene signature						
C1: number of genes in a cluster or list that have this annotation term						
C2: number of annotated genes in this cluster or list						
C3: number of all genes on array that have this annotation term						
C4: number of all annotated genes on array						
P-value: binomial approximated p-value for hypergeometric distribution						
***Gene Ontology**				P-		
C1	C2	C3	C4	value	Term Name	
81	112	5956	12364	0	protein binding	
73	112	3967	12364	0	nucleus	
58	112	3873	12364	0.000005	cytoplasm	
49	112	439	12364	0	cell cycle	
38	112	243	12364	0	cell division	
36	112	186	12364	0	mitosis	
36	112	1621	12364	0	nucleotide binding	
35	112	1250	12364	0	ATP binding	
30	112	436	12364	0	nucleoplasm	
24	112	667	12364	0	cytoskeleton	
20	112	144	12364	0	DNA replication	
18	112	294	12364	0	cell proliferation	
18	112	103	12364	0	spindle	
16	112	203	12364	0	microtubule	
15	112	224	12364	0	DNA repair	
14	112	155	12364	0	chromosome	
12	112	46	12364	0	phosphoinositide-mediated signaling	
11	112	53	12364	0	condensed chromosome	
11	112	71	12364	0	kinetochore	
11	112	54	12364	0	microtubule motor activity	
11	112	65	12364	0	microtubule-based movement	
11	112	109	12364	0	motor activity	
10	112	236	12364	0.000062	response to DNA damage stimulus	
10	112	47	12364	0	spindle pole	
8	112	59	12364	0	anaphase-promoting complex-dependent proteasomal	
8	112	41	12364	0	chromosome segregation	
7	112	47	12364	0	chromosome, centromeric region	
7	112	135	12364	0.000247	microtubule organizing center	
7	112	14	12364	0	spindle organization	
6	112	12	12364	0	chromosome condensation	

TABLE 3-continued

Gene Function enrichment analysis of EZH2 expressed gene signature					
6	112	58	12364	0.000017	negative regulation of mitotic ubiquitin-protein
					②
6	112	61	12364	0.000022	positive regulation of mitotic ubiquitin-protein
					②
6	112	38	12364	0.000002	regulation of cyclin-dependent protein kinase
					②
5	112	49	12364	0.000092	microtubule cytoskeleton
5	112	14	12364	0	mitotic sister chromatid segregation
5	112	25	12364	0.000004	spindle microtubule
4	112	19	12364	0.00003	chromosome organization
4	112	23	12364	0.000063	condensed chromosome
4	112	5	12364	0	condensin complex
4	112	36	12364	0.000347	DNA metabolic process
4	112	44	12364	0.000734	G1/S transition of mitotic cell cycle
4	112	8	12364	0.000001	M phase of mitotic cell cycle
4	112	19	12364	0.00003	midbody
4	112	12	12364	0.000005	mitotic cell cycle checkpoint
4	112	11	12364	0.000004	mitotic cell cycle spindle assembly checkpoint
4	112	12	12364	0.000005	mitotic chromosome condensation
4	112	26	12364	0.000101	nuclear chromosome
3	112	6	12364	0.000025	'de novo' IMP biosynthetic process
3	112	20	12364	0.000846	anaphase-promoting complex
3	112	14	12364	0.000302	DNA unwinding involved in replication
3	112	20	12364	0.000846	G2/M transition of mitotic cell cycle
3	112	13	12364	0.000243	kinesin complex
3	112	15	12364	0.000369	male germ cell nucleus
3	112	10	12364	0.000113	mitotic cell cycle
3	112	15	12364	0.000369	mitotic spindle organization
3	112	17	12364	0.00053	nucleotide-excision repair, DNA gap filling
3	112	13	12364	0.000243	oocyte maturation
3	112	8	12364	0.000059	outer kinetochore of condensed chromosome
3	112	15	12364	0.000369	purine nucleotide biosynthetic process
3	112	15	12364	0.000369	regulation of mitosis
3	112	11	12364	0.000149	replication fork
2	112	5	12364	0.000987	lamin filament
2	112	5	12364	0.000987	purine base biosynthetic process
2	112	3	12364	0.00036	regulation of chromosome segregation

64 reported significant, 0 expected false positive
(209 terms assessed for enrichment at p-value threshold 0.001)

② indicates text missing or illegible when filed

TABLE 4

B-RAF, EGFR, and BRG1 mutations in additional NSCLC cell lines.		
Cell Line	EGFR	BRG1
HCC2279	E746_A750d ②	WT
HCC2935	E746_A750d ②	WT

TABLE 4-continued

B-RAF, EGFR, and BRG1 mutations in additional NSCLC cell lines.		
Cell Line	EGFR	BRG1
PC14	E746_A750d ②	WT
A427	WT	Null
H661	WT	L1161fs*3

TABLE 4-continued

B-RAF, EGFR, and BRG1 mutations in additional NSCLC cell lines.		
Cell Line	EGFR	BRG1
H1703	WT	E668_Q758del
DMS-114	WT	E1310*
RERF-LC-MS	WT	A1245fs*13
H1573	WT	E1399*
H1581	WT	E1310*
H1693	WT	L1085fs*32
H838	WT	Null
H1819	WT	L1085fs*6
H596	WT	WT
H1648	WT	WT
H1437	WT	WT
H1755	WT	WT
H2087	WT	WT
H1395	WT	WT

② indicates text missing or illegible when filed

TABLE 5

Shared differentially expressed genes in EGFR and BRG1 mutants ("DEAD (Asp-Glu-Ala-Asp) box" disclosed as SEQ ID NO: 31)					
Probe	Symbol	Description	p- value	log2 fold change for Brg1mut - WT	log2 fold change for EGFRmut - WT
201983_s_at	EGFR	epidermal growth factor receptor	0.014	2.178	2.232
210986_s_at	TPM1	tropomyosin 1 (alpha)	0.007	1.525	1.743
218368_s_at	TNFRSF12A	tumor necrosis factor receptor, 12A	0.004	1.172	1.173
205130_at	MOK	MOK protein kinase	0.013	0.928	0.891
217785_s_at	YKT6	YKT6 v-SNARE homolog (<i>S. cerevisiae</i>)	0.01	0.913	1.195
219088_s_at	ZNF576	zinc finger protein 576	7E-04	0.907	1.008
202408_s_at	PRPF31	PRP31 pre-mRNA processing factor 31	0.001	0.755	0.974
218261_at	AP1M2	adaptor-related protein complex 1, mu	0.013	0.748	0.937
202578_s_at	DDX19A	DEAD (Asp-Glu-Ala-Asp) box 19A	0.002	0.716	0.531
201144_s_at	EIF2S1	eukaryotic translation initiation factor 2	0.012	0.69	0.79
216289_s_at	XRCC3	X-ray repair complementing defective	0.003	0.574	0.572
217784_at	YKT6	YKT6 v-SNARE homolog (<i>S. cerevisiae</i>)	0.002	0.531	0.81
1773_at	FNTB	farnesyltransferase, CAAX box, beta	0.003	0.514	0.408
218408_at	TMM10	membrane 10 homolog (yeast)	0.004	0.475	0.727
221769_at	SPSB3	SOCS box containing 3	0.008	0.471	0.614
202927_at	PIN1	peptidylprolyl cis/NIMA-interacting 1	0.007	0.456	0.451
211289_x_at	CDK11	cyclin dependent kinase 11	0.006	0.438	0.373
218206_x_at	SCAND1	SCAN domain containing 1	0.005	0.396	0.543
214244_s_at	ATP6V0E1	ATPase V0 subunit e1	0.005	0.384	0.481
221907_at	TRMT61A	tRNA methyltransferase	0.006	0.378	0.522
221650_s_at	MED18	mediator complex subunit 18	0.008	0.251	0.29

Example 2

EZH2 Inhibition Sensitizes BRG1 and EGFR
Mutant Lung Tumors to TopoII Inhibitors

[0442] Lung cancer is a complex disease for which epigenetic therapies may improve outcomes. It is demonstrated herein that EZH2 inhibition had differential effects on topoisomerase II (TopoII) inhibitor response of non-small cell lung cancers in vitro and in vivo. EGFR and BRG1 mutations could be used as genetic biomarkers to predict enhanced sensitivity to TopoII inhibitors in response to EZH2 inhibition. BRG1 loss-of-function lung cancer cell lines responded to EZH2 inhibition with increased S phase, anaphase bridging, apoptosis, and TopoII inhibitor sensitivity. EGFR mutant lung cancer cells were also sensitive to EZH2/TopoII inhibition, due to genetic antagonism between EGFR and BRG1. In stark contrast, EGFR/BRG1 wild-type

tumors upregulated BRG1 in response to EZH2 inhibition and ultimately became more resistant to TopoII inhibition. These findings suggest a genetic network involving EZH2, BRG1, EGFR and TopoII that, under proper circumstances, can be exploited to cause lung cancer cell death. [0443] Non-small cell lung cancer (NSCLC) is the leading cause of cancer-related death worldwide¹. This year, NSCLC will claim the lives of approximately 160,000 individuals in the United States alone². The current standard of care for NSCLC is combined modality therapy consisting of radiation treatment and chemotherapy^{3,4}. A combination of cisplatin and etoposide (VP-16) are often used to treat NSCLC⁵⁻⁷. Cisplatin induces apoptosis through formation of DNA adducts, while etoposide acts by binding within the Topoisomerase II (TopoII) complex and preventing re-ligation of the double strand break required to decatenate DNA; during mitosis, TopoII dysfunction can be visualized by anaphase bridges⁸⁻¹⁰. Despite some advances in timing and dosing strategies, combination chemotherapies effectively control disease in only ~15% of NSCLC patients^{3,4}.

[0444] In order to achieve more durable treatment response, alternative drug targets, including epigenetic enzymes, are under consideration for therapeutic intervention¹¹. Among potential epigenetic targets is the methyltransferase EZH2, which in the context of the Polycomb Repressive Complex 2 (PRC2), is well known to tri-methylate Histone H3 at lysine 27 (H3K27me3) to elicit gene silencing¹². EZH2 also has non-histone substrates, including STAT3 and GATA4, and acts a transcriptional co-activator in concert with androgen receptor¹³⁻¹⁵. Furthermore, EZH2 has been implicated in both gene transcription control and also DNA replication and repair, and appears to be a critical regulator of both stem cell and cancer phenotypes^{12,16,17}. The chemical 3-Deazaneplanocin A (DZNep) is a prototypical drug that decreases EZH2 levels and causes growth suppression in a variety of tumor cell lines, including lung cancer cell lines¹⁸⁻²³; however, DZNep may also target other

methyltransferases²⁴. More recently, several small molecule inhibitors that specifically target EZH2 methyltransferase activity including GSK126 were described²⁵⁻²⁷.

[0445] Despite the genetic complexity of lung cancer, epigenetic therapy, and specifically EZH2 inhibition, has not been tested with regard to the predominant onco-genotypes identified in patient populations. The most common genetic lesions in NSCLCs include inactivating mutations in TP53 (50-70%), deletion and mutation of LKB1 (20-35%), and activating mutations in KRAS (10-30%)²⁸⁻³⁰. Activating mutations and amplifications in EGFR (10-40%), which cause increased RAF-MEK-MAPK signaling, are also frequently observed. Recently, BRG1 (aka SMARCA4), an ATPase in the SWI/SNF chromatin-remodeling complex, was identified as a gene commonly deleted or truncated in NSCLCs (2-12%)²⁹⁻³¹. Several of these common mutations in lung cancer define patient subsets that respond to therapeutics targeting the oncogenic changes, such as EGFR inhibitors for tumors harboring activating EGFR mutations³²⁻³⁴.

[0446] To explore the potential of EZH2 inhibition as a treatment for lung cancer, EZH2 was targeted, as described herein, via small hairpin RNA, DZNep, or GSK126 in combination with chemotherapy. EZH2-inhibited lung cancer cells segregated into 2 distinct classes when challenged with chemotherapy: those with BRG1-inactivating mutations or EGFR-activating mutations were sensitized to the TopoII inhibitor etoposide, while EZH2 inhibition conferred less sensitivity ("protection") to the same chemotherapy for cells that were wild-type for BRG1/EGFR. Protected lines showed cell cycle arrest, up-regulation of BRG1 and a decrease in incidence of anaphase bridging. In contrast, sensitized lines showed increases in S phase, apoptosis and anaphase bridging in response to EZH2 inhibition and etoposide dual therapy. The sensitized phenotypes were observed in response to dual DZNep and etoposide treatment in mice bearing BRG1-mutant xenografts and mice harboring EGFR-driven endogenous lung tumors, while the protected phenotype was observed in mice bearing BRG1/EGFR wild-type xenografts or BRG1/EGFR wild-type endogenous lung tumors. EGFR and BRG1 mutations were mutually exclusive in lung cancer, and BRG1 directly binds to the EGFR promoter region to down-regulate EGFR expression. These studies define a method to predict which lung cancer patients may benefit from epigenetic therapy alone or combined with chemotherapeutics.

Results

[0447] EZH2 expression is clinically relevant and influences etoposide sensitivities. It was first sought to determine whether expression of EZH2 in primary tumors supported the hypothesis that EZH2 is an important target for NSCLC. Using Oncomine³⁵, an EZH2 co-expression gene signature was generated by annotating the top 20 genes co-expressed with EZH2 in 9 studies consisting of over 1,000 primary NSCLC samples³⁶⁻⁴³. Of 180 probes (top 20 probes from each of the 9 studies), 64 were redundant, leading to a list of 116 genes highly co-expressed with EZH2 (Table 2). It was determined whether this EZH2 co-expression signature had predictive power for cancer progression using the Director's Challenge dataset of 410 human lung adenocarcinomas⁴¹. When patients were clustered into 2 groups, those whose tumors had high expression of the EZH2 co-expression signature, and those whose tumors had low expression, a

significant difference in survival was observed between the two groups (FIG. 18A, $p < 0.00001$). This survival difference was significant even when only the Stage 1 lung cancer patients or only tumors of moderate differentiation were considered ($p = 0.003$ and $p = 0.00002$, respectively), suggesting that this EZH2 co-expression signature is an independent, robust method to further stratify patients into risk groups. Gene ontology analysis revealed that the 116 gene EZH2 co-expression signature was highly enriched for cell cycle, DNA synthesis and DNA repair (Table 3), $p < 0.001$). One of the genes highly co-expressed with EZH2 in primary tumors was Topoisomerase 2A (TOP2A), a subunit of the TopoII complex targeted by etoposide⁹.

[0448] Next, in order to begin testing EZH2 inhibition as a therapy for NSCLC, 26 human NSCLC lines that vary in genotype, stage and subtype^{44,45} were obtained (FIG. 29). It was confirmed that several of these lines express high levels of EZH2 compared to cells isolated from normal lung tissue, consistent with previous reports (FIG. 18B). To stably knock-down EZH2 expression, several lines were infected with one of two different small hairpin constructs targeting EZH2. By Western Blot, EZH2 protein and the PRC2 catalytic mark, H3K27me3, were decreased in every cell line transduced with these hairpins compared to matched shGFP transduced cells, though the knock-down and loss of H3K27me3 were both more complete with the coding region (CR) hairpin than the 3' UTR-targeting hairpin (FIGS. 18C and 24A). At the 4-day time-point, shEZH2 CR knock-down cell lines showed 5-20% reduction in growth compared to matched shGFP cultures (FIG. 24B).

[0449] To test if EZH2 knock-down cells were more sensitive to standard chemotherapy, cell viability assays were performed with the commonly used chemotherapy etoposide. shGFP and shEZH2 cell lines were treated for 4 days with a range of doses of etoposide, and cell viability was measured. Dose response curves were performed in biological triplicates, and the etoposide IC_{50} s were extrapolated for each shGFP and shEZH2 pair of lines (FIG. 18D). The fold changes in etoposide IC_{50} were graphed (FIG. 18E). Some cell lines, termed 'sensitized', were more sensitive to etoposide, with a lower etoposide IC_{50} , when EZH2 was knocked down, whereas other cell lines, termed 'protected' were more viable, with a higher etoposide IC_{50} , as shEZH2 lines. EZH2 mRNA levels were assessed after the 4 day etoposide treatment to ensure that the knock-down remained, and results demonstrate that the sensitized and protected phenotypes were not due to differential degree of EZH2 knock-down nor to significant differential expression of EZH2 in response to drug treatment (FIG. 24C).

[0450] To clearly demonstrate that EZH2 knock-down was the primary reason for differential etoposide response, virus was used to introduce ectopic EZH2 cDNA into both the A549 (sensitized) and Sw1573 (protected) shEZH2 3'UTR expressing lines. In these rescue experiments, cultures infected with the EZH2 cDNA construct and sorted for infection showed levels of EZH2 similar to those observed in the shGFP control cell lines (FIG. 24A). Etoposide dose response curves were performed with these cell lines, using sorted shGFP and empty vector-infected/sorted shEZH2 3'UTR cell lines as controls. Both the A549 and Sw1573 empty-vector expressing shEZH2 3'UTR cell lines showed the same fold changes in etoposide response relative to shGFP controls as non-infected cells from our previous experiments (compare FIG. 18F to 18E). Importantly, there

was no statistical difference in etoposide response between shGFP and EZH2 cDNA-expressing shEZH2 3'UTR cells. These results indicated that EZH2 levels determine how NSCLC cell lines respond to etoposide.

[0451] Chemical EZH2 inhibition sensitizes BRG1 and EGFR mutant cells to TopoII inhibitors. Next, to learn if etoposide sensitivities could be changed by another method of EZH2 inhibition, chemical reduction of EZH2 with the S-adenosylhomocystein hydrolase inhibitor DZNep was tested. DZNep reduces pools of methyl groups required for PRC2 function, and leads to proteosomal degradation of PRC2 components including EZH2^{18,20}. Western Blot confirmed that 4 days of 1 μ M DZNep effectively reduced EZH2 protein abundance and the H3K27me3 catalytic mark (FIG. 19A). 11 of 26 cell lines showed a significant reduction in growth at the 4-day time-point in response to 1 μ M DZNep (FIG. 25A). Next, etoposide dose response curves were performed in the presence or absence of DZNep at the same 4-day time-point. 14 of the 26 cell lines, including H157, A549 and PC9 ($p=0.001$, 0.009 and <0.0001 respectively), were more sensitive to etoposide in the presence of DZNep, while the other lines, such as H23, H460 and H441 ($p=0.0003$, 0.001 and 0.0002 respectively), were less sensitive to the chemotherapy in the presence of DZNep (FIGS. 25B and 8C). The classification of a cell line as protected or sensitized was the same as was observed for shEZH2. There was no significant difference in the etoposide IC₅₀ values of the sensitized and protected cell lines in the absence of DZNep; the etoposide IC₅₀ difference between the groups was only evident in the presence of DZNep ($p=0.43$ for no DZNep, $p=0.032$ for DZNep, FIG. 25B). This indicates that sensitization was not unique to chemotherapy resistant cell lines as seen in previous studies^{46,47}.

[0452] To learn if the changes in chemotherapy sensitivity that EZH2 inhibition caused were specific to TopoII-targeted chemotherapies, doxorubicin, which also targets the TopoII complex, was tested⁹. Cell lines segregated into the same protected and sensitized classes with doxorubicin combined with DZNep (FIG. 8C). Etoposide is almost always administered with cisplatin to NSCLC patients³. When cisplatin (5 μ M, ~IC₃₀) was administered in combination with etoposide/DZNep treatment, the changes in etoposide IC₅₀ in response to DZNep were still evident, showing that the effect of DZNep on etoposide sensitivity is intact in this more clinically relevant situation (FIG. 25C). Furthermore, the pattern of sensitized and protected cells was not observed when cisplatin and DZNep were combined, suggesting that cisplatin response is not mechanistically linked to the EZH2/TopoII inhibition phenotype (FIG. 8D). Lastly, it was sought to validate our results using a novel compound that specifically targets EZH2 methyltransferase activity, GSK126. Cell lines were treated in culture for 7 days with 10 μ M GSK126, and observed a marked decrease in H3K27me3 levels while EZH2 levels remained unchanged as expected (FIG. 19C). Etoposide dose response curves showed that GSK126 treated cell lines fell into the same TopoII inhibitor-sensitized and -protected classes as observed with DZNep or shEZH2 (FIG. 19D).

[0453] To uncover biomarkers that could potentially be used to identify the protected and sensitized phenotypes, the mutational annotation available for the NSCLC lines was examined. 12 of the 14 sensitized cell lines harbored inactivating mutations in BRG1 or activating mutations in EGFR, while 10 of the 12 protected cell lines were wild-type

for the two genes (FIG. 29, Fisher's exact test $p=0.001$). H23 is a line protected from etoposide by DZNep, yet has a late coding region BRG1 mutation predicted to produce a protein with an intact ATPase domain⁴⁴, supporting the lack of sensitization in the EZH2/TopoII inhibitor assay. No correlation of TP53, KRAS, NRAS, LKB1 or PIK3CA mutations with the protected and sensitized phenotypes was observed (FIG. 29). However, for both KRAS and PIK3CA, results suggest that protected lines may be enriched for these mutations, though neither reached significance with the current cohort of cell lines ($p=0.054$ for KRAS, $p=0.051$ for PIK3CA, Fisher's Exact Test). Together these results indicate that EZH2 inhibition sensitizes some NSCLC cell lines to TopoII inhibition, while causing TopoII inhibitor resistance in others, and that these phenotypes correlated with mutations in BRG1 and EGFR.

[0454] In vitro sensitivities to DZNep/etoposide predict in vivo responses. To determine if the protected and sensitized phenotypes could be observed in vivo, xenograft-bearing mice were treated with etoposide and DZNep. 12 days after cell injection, when small but visible tumors had developed (data not shown), mice were randomized into groups that received etoposide, DZNep, or both. For the cell line H157, which was sensitized to etoposide by DZNep in vitro, DZNep alone was weakly effective at reducing tumor size compared to vehicle treated tumors (FIG. 3A and data not shown, vehicle vs. DZNep $p=0.08$). However, dual etoposide and DZNep therapy prevented tumors from forming in 4/6 mice, proving more efficacious than etoposide or DZNep alone (etoposide vs. dual $p=0.003$, DZNep vs. dual $p=0.03$). Similarly, the protected phenotype of the H23 cell line was recapitulated in vivo in response to dual etoposide and DZNep therapy. H23 xenografts were sensitive to DZNep as a single therapy (FIG. 3B, data not shown, vehicle vs. DZNep $p=0.001$), but those that received dual therapy grew significantly larger than those treated with either DZNep or etoposide alone (dual vs. DZNep $p=0.005$, dual vs. etoposide $p=0.0005$). Together these data demonstrate that the sensitized and protected phenotypes can be observed in vivo.

[0455] Next, mouse models of lung cancer representing predicted sensitized (EGFR-T790M/L858R transgenic; EGFR hereafter⁴⁸) and protected (Kras-G12D/p53-null; Kras/p53 hereafter⁴⁹) tumor types were treated to test our ability to predict response to DZNep and etoposide. These models represent a predicted 'protected' cancer in the Kras/p53 model, which is wild-type for Brg1 and Egfr, and a predicted 'sensitized' cancer in the EGFR mutant model, which is wild-type for Brg1 (confirmed by unpublished RNA-seq). Mice with radiographically documented lung masses reflecting active lung cancer were used for each experimental and control condition at the start of treatment (week 0). Chemotherapy, EZH2 inhibitor, or combination therapy was then administered to cohorts of mice for 4 weeks. MRI was performed at 2 weeks and 4 weeks post treatment start date (data not shown). Marked tumor regression in the EGFR mouse model was observed in response to 4 weeks of dual etoposide/DZNep treatment, while mice in the other treatment arms showed continued tumor growth (data not shown and FIG. 20A, $p=0.008$ dual vs. DZNep, $p=0.004$ dual vs. etoposide). This is in striking contrast to the protected Kras/p53 model, in which tumors proceeded to grow during the 4 weeks despite dual treatment (FIG. 20B, $p=0.4$). Drug efficacy in both models was confirmed by

histology and immuno-histochemistry for EZH2, which was dramatically reduced in both the EGFR and Kras/p53 tumors in response to dual therapy (data not shown). In addition pEGFR is greatly reduced by dual treatment, fitting with the mechanism we demonstrate in the next figures. These data indicate that tumors mutant for BRG1 (H157 xenografts) or relying upon activation of EGFR (EGFR mice), are sensitized to TopoII inhibition by DZNep in vivo, suggesting a new possible therapeutic avenue for these two genotypes of NSCLC.

[0456] Dual etoposide/EZH2 inhibition differentially affects cell cycle, apoptosis and anaphase bridging. Next, the hypothesis that EZH2 inhibition caused changes in cell cycle status, rendering cells more or less sensitive to chemotherapy was tested. It was reasoned that protected lines may undergo a cell cycle arrest in response to EZH2 inhibition, thus sparing them from etoposide-induced DNA damage, while sensitized lines may be unable to arrest and instead continue to cycle in the presence of etoposide. shGFP, shEZH2, and shGFP cells treated with 1 μ M DZNep were treated for 4 days with vehicle or 5 μ M etoposide (\sim IC₅₀) and stained with 7AAD to assess cell cycle profiles by flow cytometry. Protected etoposide-treated lines had an average of 14% fewer cells in S phase when in response to DZNep or shEZH2 compared to treatment with etoposide alone; in contrast, sensitized etoposide-treated lines had an average of 14% more cells in S phase in response to DZNep or shEZH2 (FIGS. 21A, 26A, and 26B). Importantly, the S phase content of the sensitized and protected lines was not significantly changed in response to EZH2 inhibition alone, suggesting that cell cycle changes are specific to dual treatment (FIG. 26C). To test the hypothesis that the increase in S phase in sensitized etoposide-treated lines translated into higher levels of apoptosis, Annexin V/7AAD flow cytometry was performed. Cultures treated for 4 days with 1 μ M DZNep, 5 μ M etoposide, or both were collected and stained with Annexin V and 7AAD, and the percentage of apoptotic cells (Annexin V+/7AAD-) cells was assessed by flow cytometry (FIG. 21B). While the protected lines H2009 and H23 showed no difference in apoptotic levels in etoposide compared to dual treated cultures, the sensitized lines H157 and PC9 had significantly higher apoptotic fractions in dual treated cultures than in cultures treated with etoposide as a single agent ($p=0.017$ and $p=0.026$, respectively). Together these data indicate that sensitized cell lines respond to combined EZH2/TopoII inhibition with altered cell cycle responses that predispose cells to increased etoposide-induced apoptosis.

[0457] To further address the mechanism through which EZH2 inhibition changes sensitivity specifically to TopoII inhibitors, the recent report of a physical interaction between BRG1 and TopoII that allows for increased TopoII function¹⁰ was considered. Given that BRG1 and EZH2 are known to be genetically antagonistic⁵⁰, it was hypothesized that protected cell lines had up-regulation of BRG1 in response to EZH2 inhibition and therefore had increased TopoII function. In order to test this hypothesis, anaphase structures in DZNep and vehicle-treated cultures were imaged, quantifying the numbers of anaphase bridges that are indicative of a failure of TopoII to decatenate DNA prior to mitosis (FIG. 21C). BRG1 mutant cell lines had lower baseline levels of anaphase bridging than Brg1 wild-type cell lines. However, when challenged with DZNep, BRG1 mutant cell lines showed increased anaphase bridging, while

BRG1 wild-type lines showed a significant decrease (FIG. 21D). To learn if the decrease in anaphase bridging of the BRG1 wild-type cell lines corresponded to an increase in BRG1 levels, real time RT-PCR for BRG1 transcript was performed (FIG. 21E). In 5 different cell lines, 4 of which are wild-type for EGFR (Sw1573, Calu6, H23 and H460) and one of which is EGFR mutant (PC9), BRG1 transcript was reproducibly increased by DZNep treatment. Together these data indicate that increased BRG1 is an effect of EZH2 inhibition, which leads to a decrease in anaphase bridging and protection from TopoII inhibition. In contrast, the increased anaphase bridging after EZH2 inhibition in BRG1 mutants likely explains their increased sensitivity to TopoII inhibition and subsequent apoptosis.

[0458] BRG1 and EGFR are genetically antagonistic. While the observed anaphase bridging of BRG1 mutants in response to EZH2 inhibition explained their increased sensitivity to TopoII inhibition, it was still unclear why EGFR mutants also fell into this chemo-sensitized category. In the panel of cell lines, 30% have a BRG1 mutation and 23% have an EGFR mutation. With these mutation frequencies, \sim 7% of cell lines are expected to have both BRG1 and EGFR mutations. However, no NSCLC cell line in this panel or elsewhere that harbored both mutations was observed (FIG. 29, Table 4, Fisher's exact test $p=0.005$). To examine the pattern of BRG1 and EGFR mutations in primary human lung cancers, lung adenocarcinoma whole exome sequencing from The Cancer Genome Atlas (TCGA) and a recent publication were queried²⁹. Of the 412 adenocarcinoma samples, 56 (14%) had exon 19 deletion or missense mutation in EGFR, while 41 (10%) had missense, nonsense, deletion or insertion mutations in BRG1. With these allele frequencies, 1.4% of the tumor samples were expected to have both BRG1 and EGFR mutations, though only 2 (0.5%) were observed (Fisher's exact test, $p=0.03$), and both contain EGFR mutations that may be passenger/non-functional mutations (E574L, H1129Y). With compilation of the cell line and primary tumor data, the anti-correlation of EGFR mutations and BRG1 mutations was more significant (Fisher's exact test, $p=0.002$).

[0459] The negative correlation of BRG1 and EGFR mutations suggests that these pathways interact genetically and that these mutations may be functionally redundant. Supporting this idea, in the Director's Challenge dataset of primary human lung adenocarcinomas⁴¹, a strong negative correlation was found between EGFR and BRG1 expression, supporting the theory that loss of BRG1 may be permissive for high expression of EGFR (FIG. 5A, Spearman's correlation $=-0.619$, $p=0.0002$). Likewise in a primary NSCLC dataset³⁰, tumors with mutations in BRG1 had significantly higher expression of EGFR than EGFR/BRG1 wild-type tumors, while there was no difference in EZH2 levels among the genotypes (FIGS. 5B and 27A, $p=0.002$). To explore the relationship between these mutations in our cell lines, available gene array data was used to compare gene expression signatures in the various genotypes. Overlap between differentially expressed genes in EGFR mutant lines vs. EGFR/BRG1 wild-type lines, and BRG1 mutant lines vs. EGFR/BRG1 wild-type lines was queried (FIG. 11A). Interestingly, it was found that EGFR was the top shared up-regulated gene, over-expressed 3.1-fold in BRG1 mutant cell lines and 3.25-fold in EGFR mutant cell lines, both compared to EGFR/BRG1 wild-type lines (FIG. 5C, Table 5, $p=0.014$). Importantly, we also confirmed with

these arrays that baseline EZH2 transcript levels do not differ among the various genotypes (FIG. 11B, $p=0.5$). Immuno-fluorescence for EGFR confirmed that BRG1 mutant cell lines have high expression of EGFR compared to BRG1 wild-type cell lines, which have very little detectable EGFR staining (data not shown). Although gefitinib, and other EGFR specific inhibitors, are thought to cause death preferentially of cells with activating mutations in EGFR³², we assessed whether there was any detectable difference in gefitinib sensitivities of the BRG1-mutant/EGFR^{high} cells vs. the BRG1-wild-type/EGFR^{low} cells. It was observed that the IC₅₀ for gefitinib among the cultures is identical, but the growth inhibition at sub-IC₅₀ concentrations of gefitinib was significantly different between BRG1-mutant and BRG1-wild-type cell lines (FIG. 27B).

[0460] To further explore the genetic relationship between these two pathways, EGFR and BRG1 expression were manipulated via knock-down and over-expression in NSCLC cell lines. In the BRG1 mutant cell line HCC15, which expressed high levels of wild-type EGFR, re-introduction of BRG1 significantly decreased the amount of EGFR transcript, while EGFR knock-down led to an increase in the truncated mutant Brg1 mRNA (FIG. 18A, $p=0.005$). These changes were confirmed by immuno-fluorescence (data not shown). In contrast, in the H460 line, which is wild-type for BRG1 and has little detectable EGFR expression, knock-down of BRG1 led to up-regulation of EGFR ($p=0.009$) and over-expression of wild-type EGFR led to a decrease in BRG1 transcript (FIG. 18B, $p=0.002$). These results were observed in several cell lines (FIG. 11D). The ENCODE database⁵¹ shows a chromatin immuno-precipitation (ChIP) peak for BRG1 in the regulatory element upstream of the EGFR transcriptional start site in HeLa-S3 cells. Therefore, it was hypothesized that the mechanism through which BRG1 over-expression was down-regulating EGFR was through direct binding of the SWI/SNF complex to the EGFR regulatory element. To validate this hypothesis, ChIP assays were performed in the HCC15 shGFP (control) and HCC15 BRG1 over-expressing cell lines (FIG. 22). It was found that BRG1, immuno-precipitated with an antibody recognizing the protein itself or the FLAG-tag, was significantly associated with the EGFR regulatory element ($p=0.02$ for both IPs). Together these data indicate that BRG1 loss-of-function mutations and EGFR gain-of-function mutations are mutually exclusive in NSCLC, and that BRG1 and EGFR are genetically antagonistic due to binding of BRG1 to an EGFR regulatory element that leads to down-regulation of EGFR transcript.

BrG1 and EGFR Control the Sensitized Phenotype

[0461] To learn what impact modulation of EGFR and BRG1 levels had on the protected and sensitized phenotypes, etoposide dose response curves were performed on the BRG1 and EGFR knock-down and over-expression cell lines. For the HCC15 control cell line, DZNep decreased the etoposide IC₅₀ 2-fold, consistent with its sensitized phenotype ($p<0.0001$). However, BRG1 over-expressing HCC15 lines showed an increase in etoposide IC₅₀, suggesting that the phenotype of the line was converted from sensitized to protected through re-introduction of BRG1 (FIG. 23A, $p=0.037$). In the HCC15 shEGFR line, the line was no longer sensitized to etoposide by DZNep, but was not protected either, consistent with the fact that this line still lacks functional BRG1 protein. For the protected cell line, DZNep

raised the etoposide IC₅₀ 2-fold in the H460 control cells ($p=0.028$), yet had no effect on etoposide IC₅₀ when EGFR was over-expressed ($p=0.35$) or BRG1 was knocked-down (FIG. 6b, $p=0.5$). We confirmed that 4 days of etoposide, DZNep or dual treatment did not change the over-expression and knock-down of EGFR and BRG1 by RT-qPCR (FIG. 28B). These data demonstrate that the sensitized and protected phenotypes are partially reliant upon the levels of BRG1 and EGFR within the cells.

[0462] Finally, it examined whether changes in etoposide sensitivity of the BRG1- and EGFR-modulated DZNep-treated cell lines were attributable to cell cycle, apoptotic and anaphase bridge changes. While shBRG1 in the BRG1 wild-type line H460 led to decreased proliferation, BRG1 over-expression in the mutant line HCC15 led to increased proliferation (FIG. 28B). Despite their increased proliferation in the absence of etoposide, etoposide-treated HCC15 BRG1 over-expressing cell lines did not accumulate in S phase in response to DZNep as the control shGFP cells did (FIG. 23C, $p=0.02$). HCC15 shEGFR showed higher percent S phase in response to etoposide alone than do shGFP control cells, but have no additional increase in S phase in response to DZNep/etoposide dual treatment. Consistent with results from other sensitized lines, the HCC15 control cells showed an increase in sub-G1/apoptotic cells in response to dual therapy, but this change was not evident in either shEGFR or BRG1 over-expression cultures (FIG. 28C). Likewise, the anaphase bridging phenotype clearly demonstrated an increase in TopoII dysfunction in response to DZNep in the HCC15 control line, but not in the shEGFR or BRG1 over-expressing lines (FIG. 23D). The protected H460 control cell line showed a depletion of S phase in response to the dual therapy; however, the etoposide treated H460 shBRG1 and EGFR over-expressing cell lines were enriched for S phase (FIG. 23E, $p=0.02$ for shBrg1, $p<0.001$ for EGFR over-expression). While none of these cultures showed differences in sub-G1/apoptotic cells (FIG. 28C), the decrease in anaphase bridges normally observed in H460 cells was reversed by BRG1 knock-down or EGFR over-expression (FIG. 23F, $p<0.04$). Together these data demonstrate that EZH2/TopoII inhibition altered cell cycle dynamics, apoptosis and anaphase bridging, and that these effects were mediated by the levels of BRG1 and EGFR expression in the cells.

Discussion

[0463] Targeting of epigenetic enzymes such as EZH2 could be a novel therapeutic approach for lung cancer patients; yet, clear demonstration of how EZH2 inhibitors interact with current therapies and specific genotypes of NSCLCs is lacking. It is demonstrated that susceptibility to combined TopoII and EZH2 inhibition is predictable by NSCLC genotype. BRG1 loss-of-function mutants represent on class of TopoII/EZH2-inhibitor 'sensitized' cells that cannot up-regulate Brg1 in response to EZH2 and therefore have impaired TopoII function that rendered them sensitive to etoposide. Another, unexpected class of sensitized tumors is EGFR mutants, which respond to EZH2 inhibition with BRG1 up-regulation and apoptosis, the later likely due to direct negative regulation of EGFR by BRG1⁵² (FIG. 23G). Endogenous mouse lung cancer models were used to verify the link between tumor genotype and EZH2 inhibitor effects on chemotherapy response. The differential effects of EZH2 inhibition on chemotherapy in vitro allowed the prediction

of responses to therapies *in vivo*, suggesting that dual EZH2/TopoII inhibitor therapy will be an option for patients with either EGFR or BRG1 mutant NSCLCs.

[0464] The data described herein indicate that patients with lung cancers bearing EGFR or BRG1 mutations will benefit from dual EZH2/TopoII inhibitor strategies. This work also indicates that this combination may be ineffective or even detrimental to patients with EGFR/BRG1 WT tumors due to the reduced efficacy of TopoII inhibitors such as etoposide. The mouse model that was used to validate the sensitized phenotype harbors lung tumors driven by EGFR T790M and L858R mutations⁴⁸. The T790M mutation represents the canonical gefitinib/erlotinib “gatekeeper” resistance mutation, suggesting that dual EZH2/TopoII inhibition could be an option for patients that developed resistance to first line EGFR TKI strategies³⁴. Because we showed BRG1 could bind at a site upstream of the EGFR start codon, it is possible that EZH2 inhibitors, through BRG1 up-regulation will negatively regulate both point mutant and exon 19 deletion forms of EGFR. Furthermore, BRG1 mutations have only recently been identified in lung cancers and there are currently no specific treatments available for this subset of lung cancer patients; EZH2/TopoII inhibitor combination could offer the first therapeutic intervention for BRG1 mutant lung cancers. No correlation of TP53, KRAS, NRAS, LKB1 or PIK3CA mutations with the protected and sensitized phenotypes was observed, yet it remains possible that other predominant NSCLC oncogenotypes will exhibit differential responses to EZH2/TopoII inhibitor combinations.

[0465] It was found that in NSCLC patient datasets, BRG1 inactivating mutations and EGFR activating mutations were significantly anti-correlated. This finding is in agreement with two recent publications showing that EGFR mutant tumors in Asian populations, where EGFR mutations occur at a much higher frequency than in US patients, also appear to preferentially lack BRG1 mutations^{53,54}. It was found that BRG1 mutant tumors and cell lines had higher total EGFR levels, and it is not yet known if this translates into higher active EGFR (pEGFR) in BRG1 mutant tumors⁵³. In support of the hypothesis that BRG1 mutants have more EGFR signaling, BRG1 mutant cell lines were slightly more sensitive to gefitinib than BRG1 wild-type cell lines (Fillmore and Kim data not shown). Some NSCLC patients with wild-type EGFR clinically benefit from gefitinib and erlotinib therapy by stabilizing disease and preventing further progression^{55,56}, and it will be interesting to observe whether these patients were BRG1 mutation carriers. It was also found that re-expression of BRG1 protein in a BRG1 mutant cell line led to down-regulation of EGFR, and that BRG1 was significantly associated with an upstream regulatory element of EGFR in these cells. One theory of how BRG1, which is often thought to activate genes through displacement of nucleosomes from key promoter elements⁵⁷, can elicit gene silencing is through blocking the binding of transcriptional co-activators such as p300⁵⁸. Also, recent work suggests that on a genome-wide scale, BRG1 deletion causes derepression of more than 1,000 genes in MEFs, again supporting that the SWI/SNF complex participates in transcriptional silencing of target genes⁵⁹.

[0466] In agreement with previous work, it was found herein that expression of an EZH2-expression signature was highly correlated with poor prognosis in all stages and differentiation states of lung adenocarcinoma. On this list of

EZH2 co-expressed genes was TOP2A and numerous genes involved in the decatenation checkpoint, suggesting at the minimum a genetic interaction between these pathways. It is demonstrated herein that in BRG1 and EGFR mutant cell lines, EZH2 inhibition caused sensitization to the TopoII inhibitors etoposide and doxorubicin. Unlike Brg1-deficient MEFs, BRG1 mutant cell lung cancer cell lines did not show dysfunction in TopoII function, as assessed by anaphase bridging, until EZH2 was inhibited. Recently, research has uncovered roles of EZH2 outside of histone methylation^{13, 14}, and it is possible that genetic interaction with TopoII may be another non-histone mechanism for EZH2 action.

[0467] Therapeutic strategies targeting epigenetic factors have risen as promising new candidates for lung cancer and other difficult-to-treat cancers. Rationales for epigenetic therapy approaches include the possibility that targeting universally required chromatin enzymes will benefit a broader patient population than drugs targeting specific oncogenic changes.

Methods

[0468] Cell Lines and small hairpins. Cell lines used are listed in FIG. 29. All cell lines were maintained in RPMI 1640 media with 10% fetal bovine serum, 4 mM L-glutamine and penicillin/streptomycin at 37° C., 5% CO₂. The pLK0.1 EZH2 shRNA construct clones TRCN0000040076 and TRCN0000040073 were purchased from SIGMA and the shGFP plasmid 12273 is available on Addgene. Both shBRG1 and the matched empty vector were provided by the Smale lab⁶⁰ and are available on Addgene, the Brg1 over-expression plasmid 19148 was purchased through Addgene, and the shEGFR and EGFR WT over-expression constructs were provided by the Janne lab⁶¹. The EZH2 over-expression construct was derived by cloning human EZH2 cDNA into pLenti7.3N5-DEST (Invitrogen). Lentivirus was packaged in 293T cells using established protocols⁶², and retrovirus was packaged in PlatE cells again using established protocols⁶³. Cell lines were infected with viral-containing supernatant containing 6 µg/mL polybrene (SIGMA) for a period of 10-18 hours. Infected cultures were selected with 1 µg/mL puromycin (all sh constructs and EGFR oe, SIGMA), 200 µg/mL hygromycin (BRG1 oe, Invitrogen), or by flow cytometry for GFP (EZH2 oe) 5 days post infection.

Small Hairpin Sequences:

[0469]

```
GFP: (SEQ ID NO: 15)
GCCC(GCAAGCTGACCCTGAAGTTCAT) TCAAGAG

(ATGAACCTCAGGGTCAGCTTGC) TTTT

EZH2 coding region: (SEQ ID NO: 16)
CCGG(CGGAAATCTTAAACCAAGAAT) CTCGAG

(ATTCTTGGTTTAAGATTCCG) TTTT

EZH2 3' UTR: (SEQ ID NO: 32)
CCGG(TATTGCCTTCTCACCAGCTGC) CTCGAG

(GCAGCTGGTGAGAAGCAATA) TTTT
```

-continued

EGFR:
(SEQ ID NO: 17)
CCGG (GCTGAGAATGTGGAATACCTA) CTCGAG
(TAGGTATTCACATTCTCAGC) TTTT
BRG1:
(SEQ ID NO: 18)
TTTG (TGGATAAGCAGCACAGATT) TCAAGAG
(AATCTTCTGCTGCTTCTCCA) TTTT

[0470] Drugs. Etoposide, cisplatin, and doxorubicin (SIGMA) were diluted to a stock of 100 mM in DMSO for all cell culture experiments. DZNep was diluted in DMSO to a stock of 10 mM. GSK126 was purchased from Xcess Bio as a 10 mM stock in DMSO. All stocks were diluted in DMSO to 1000× concentration prior to addition into media at 2× concentration and final dilution onto plated cells 1:1.

[0471] Cytotox Assays. Cell lines were dissociated, counted and plated at 5000 cells per well in flat bottom opaque tissue culture treated 96 well plates (CytoOne). Edge wells were filled with PBS. The following day, 2× drug diluted in media was added to each well such that the well then contained 100 μ l media with 1× drug concentration at the following doses: etoposide; 0, 0.1, 1, 3, 5, 7, 10, 50, 100, 500 μ M, cisplatin; 0, 0.1, 1, 5, 10, 30, 70, 100, 500 μ M, or doxorubicin; 0, 0.01, 0.1, 0.5, 1, 3, 5, 7, 10, 50 μ M. After 4 days, CELLTITER-GLO™ (Promega) was added and luminescence was read on a BioTec plate reader to determine relative cell number in each well. Data were averaged for triplicate or quadruplicate technical replicates and normalized to the untreated wells. Results from three or four independent biological replicate experiments were input into GRAPHPAD PRISM™ software to extrapolate IC₅₀ and s.e.m. of IC₅₀ for a given cell line.

[0472] Flow Cytometry. For 7AAD-cell cycle analysis, cell lines were plated at 1.5×10^6 cells per 10 cm plate and treated with drug for 4 days. Cells were then dissociated, fixed with 100% ice cold Ethanol for at least 2 hours, incubated for 30 minutes with 1mg/mL DNase-free RNase A (Thermo) and resuspended in 20 μ g/mL 7-Aminoactinomycin D (7AAD; Invitrogen). 30,000 events were collected on the BD FORTESSA™ and analyzed with the MODFIT LT™ software. Results from three to four independent biological replicate experiments were averaged for each cell line.

[0473] For Annexin V/7AAD apoptosis analysis, cell lines were plated at 1×10^5 cells per well of 6 well plate and treated with drug for 4 days. Supernatant was retained and added to trypsinized suspensions of adherent cells. Cells were stained with Annexin V-FITC (BD Biosciences) according to manufacturer's instructions, and resuspended with 1 μ g/mL 7AAD prior to analysis on BD FORTESSA™.

[0474] Quantitative RT PCR RNA from treated cell lines was extracted using Absolutely RNA™ kits (Agilent) and cDNA was made using the SuperScript III™ kit (Invitrogen). Relative gene expression was assayed with Sybr green on the StepOnePlus™ Real-Time PCR System (Applied Biosystems). Relative expression was calculated by Gene of Interest($C_{t_{reference}} - C_{t_{experimental}}$)-CYP A($C_{t_{reference}} - C_{t_{experimental}}$) and graphed on the log₂ scale. For all experiments, the reference sample was a matched untreated or control transduced cell line.

[0475] For normal human lung samples, tissue adjacent to lung tumors was provided and dissociated with Dispase (BD Biosciences) for 45 minutes at 37° C. then titrated through a 5 mL pipet. Tissue was then incubated in DMEM, 10% fetal bovine serum, 4 mM L-glutamine and penicillin/streptomycin at 37° C., 5% CO₂ on tissue culture treated plates overnight to allow adherence of fibroblasts. The following day, supernatant, which included visible alveolar and bronchiolar structures, was collected, pelleted, and RNA was isolated as described. 4 patient samples were assayed and averaged for FIG. 1b.

Primer Sequences:

[0476]

CYP A:
F TCATCTGCACTGCCAAGACTG (SEQ ID NO: 19)
R CATGCCTTCTTCACTTTGCC (SEQ ID NO: 20)
EZH2:
F AGGAGTTTGCTGCTGCTCTC (SEQ ID NO: 21)
R CCGAGAATTTGCTTCAGAGG (SEQ ID NO: 22)
BRG1:
F AGCGATGACGTCTCTGAGGT (SEQ ID NO: 11)
R GTACAGGGACACCGCCACT (SEQ ID NO: 12)
EGFR:
F TAACAAGCTCAGCAGTTGG (SEQ ID NO: 13)
R GTTGAGGGCAATGAGGACAT (SEQ ID NO: 14)

[0477] Xenograft Experiments. Cell lines were dissociated into single cells, counted and resuspended at 1×10^6 cells per 2504 of 1:1 media/matrigel (BD). 8- to 16-week-old Foxn-lu (nude) mice (Harlan) were injected subcutaneously with 1×10^6 cells in 2-4 spots on flanks. Etoposide and DZNep were administered from day 12 to day 17 post injections; etoposide: 20 mg/kg/day i.p. in corn oil once per day for 5 consecutive days, and DZNep 2 mg/kg/day i.p. in corn oil twice per week for 1 week, or 1 mg/kg/day i.p. in corn oil twice per week for 2 weeks. Tumor growth was measured every other day by caliper. All mouse experiments were approved by the CHB Animal Care and Use Committee and by the Dana-Farber Cancer Institute Institutional Animal Care and Use Committee, both accredited by AAALAC, and were performed in accordance with relevant institutional and national guidelines and regulations.

[0478] Generation of the EZH2 Co-Expressed Gene Signature. ONCOMINE™³⁵ was used to query the top 20 genes co-expressed with EZH2 in all datasets containing human non-small cell lung cancer samples and co-expression data³⁶⁻⁴³. 20 was the number of probes chosen to examine from each study in order to yield a list between 100-200 genes, which allowed for robust hierarchal clustering of samples similar to that in previous studies³⁶. Of the 180 probes, 64 were redundant, leading to a list of 116 genes highly co-expressed with EZH2 (Table 2). Because data sets on Oncomine were from various microarray platforms, the gene list was then used to generate a probe list for the 116 genes corresponding to probes on the U133A Affymetrix array using the batch query function on the NetAffx™ website (available on the World Wide Web at <http://www.affymetrix.com/analysis/index.affx>).

[0479] Kaplan-Meier Analysis Raw gene expression data from human lung adenocarcinoma samples were obtained.

Probe intensities from the Affymetrix U133A™ platform used in these studies were normalized and modeled using dChip™ software (available on the World Wide Web at <http://biosun1.harvard.edu/complab/dchip>). Kaplan-Meier survival analyses were implemented after the samples were hierarchically clustered using centroid linkage into two risk groups using the EZH2 co-expressed gene signature. Differences of the survival risk between the two risk groups were assessed using the Mantel-Haenszel log rank test. The large area between the two risk groups and its associated small p value from the Mantel-Haenszel log rank test implicate a robust survival classification model.

[0480] Cell Line Micro-Array All array data is publically available on Gene Expression Omnibus (available on the World Wide Web at <http://www.ncbi.nlm.nih.gov/geo/>). Array quality was assessed using the R/Bioconductor™ package (available on the World Wide Web at <http://www.bioconductor.org/>). Raw CEL files from U133A Affymetrix arrays were processed using the robust multiarray average (RMA) algorithm⁶⁴. To identify genes correlating with the phenotypic groups, we used limma⁶⁵ to fit a statistical linear model to the data and then tested for differential gene expression in the three groups, WT: H460, H441, H2122, H2009, Calu6, HCC95, EGFR mutant: H1650, HCC827, HCC4006, H1975, H3255, PC9, Brg1 mutant: A549, H1299, H157, H2126, H522, HCC15. Results were adjusted for multiple testing using the Benjamini and Hochberg (BH) method⁶⁶, and significance was determined using a False-Discovery-Rate cutoff of less than 5%.

[0481] Statistical Analysis. Except where indicated, a 2-tailed Student's t-test with equal variance was used to compare measurements between 2 conditions. Unless noted otherwise, pooled data is represented by the mean and standard error. Except where indicated p-values less than 0.05 were considered significant.

[0482] Western blot Whole cell extracts were made in RIPA buffer (0.5% Deoxycholate, 1% IGEPAL-CA630, 0.1% sodium dodecyl sulfate, 150 mM NaCl, 50 mM Tris-8.1), lysates were cleared by centrifugation, and protein concentrations were quantified with the Pierce BCA Protein Assay Kit (Thermo). For Western blotting, 25 µg of protein extract per sample was denatured with heat and reducing agents, separated on a 4-12% acrylimide gel (BioRad) and transferred to nitrocellulose (GE Healthcare). Antibodies used for Western blotting were: EZH2 (clone D2C9; Cell Signaling; 1:200), Histone H3 (polyclonal; AbCAM ab1791; 1:2000) and H3K27me3 (polyclonal; Millipore 07-449; 1:1000) all incubated overnight at 4° C. Secondary antibody, anti-rabbit-HRP (Santa Cruz sc-2313; 1:5,000), was incubated for 1 hour at room temperature. After washing, chemiluminescence was visualized with Western Lightning Plus-ECL (PerkinElmer) and exposure onto KODAK BioMax XAR™ film.

[0483] Anaphase Bridge Analysis. To quantify anaphase bridges, cells were grown on 4-well cultures slides (Lab Tek II). Adherent cells were fixed with 4% paraformaldehyde for 20 min, washed and stained with Vectashield™ with 4',6-diamidino-2-phenylindole (DAPI; Vector Labs). Images were taken of each anaphase structure, and the number of anaphases with bridges over the total number of anaphases (between 11 and 34 total anaphases per well of a 4-well chamber slide) was recorded for each of three or more

independent biological replicate experiments. Imaging was performed with a Nikon 90i™ camera and NIS-Elements™ software and processed with NIS-Elements and Adobe Photoshop™.

[0484] Immuno-fluorescence Cells were fixed in 4% paraformaldehyde and permeabilized with 10% Normal Donkey Serum (NDS; Jackson ImmunoResearch), 0.25% Triton-X (SIGMA), both in PBS. Primary antibodies, Brg1 (clone G-7, Santa Cruz), EGFR (polyclonal; Cell Signaling 2232), were incubated over-night at 1:100 dilution in PBS, 10% NDS. Slides were washed 3× and secondary antibodies, anti-mouse-AlexaFluor594 and anti-rabbit-AlexaFluor488 (Invitrogen) were incubated at 1:500 for 1 hour. After washing, cover slips were mounted with Vectashield™ with 4',6-diamidino-2-phenylindole (DAPI; Vector Labs). Imaging was performed with a Nikon 90i™ camera and NIS-Elements™ software and processed with NIS-Elements™ and Adobe Photoshop™.

[0485] Treatment and MRI of Endogenous Mouse Models Doxycycline inducible EGFR-TL (L858R/T790M) transgenic mice⁴⁸, and Lox-Stop-Lox-Kras^{G12D};p53^{fl/fl} (Kras/p53) mice^{49,67} were housed in a pathogen-free environment at the Harvard School of Public Health and were handled in strict accordance with Good Animal Practice as defined by the Office of Laboratory Animal Welfare. All animal work was done with Dana-Farber Cancer Institute IACUC approval. Cohorts of EGFR TL/CCSP-rtTA were put on a doxycycline diet at 6 weeks of age to induce the expression of mutant EGFR, while Kras/p53 mice received intranasal adeno-Cre between 6 and 8 weeks of age. Mice were evaluated by Magnetic resonance imaging (MRI) 12 to 16 weeks after doxycycline diet or adeno-Cre infection to document and quantify the lung cancer burden before being assigned to various treatment study cohorts. Treated mice in all cohorts have the similar initial tumor burden. Tumor bearing mice were treated either with vehicle (corn oil), etoposide 10 mg/kg i.p. 3× per week for 4 weeks; DZNep 4 mg/kg i.p. 2× per week for 4 weeks, or both etoposide and DZNep. The mice were imaged by MRI biweekly to determine the reduction in tumor volume during the respective treatments as described previously³⁴. The tumor burden volume and quantification were reconstructed on 3D slicer software (<http://www.slicer.org>). Immunohistochemistry was performed as described with anti-EZH2 (clone D2C9; Cell Signaling) or anti-pEGFR (Y1068; clone D7A5; Cell Signaling) and developed using Vectastain Elite ABC kit™ (Vector Labs). Imaging was performed with a Nikon 90i™ camera and NIS-Elements™ software and processed with NIS-Elements™ and Adobe Photoshop™.

[0486] Chromatin Immunoprecipitation (ChIP) 5×10⁶ cells were fixed in 1% formaldehyde for 10 minutes prior to addition of glycine to a concentration of 1 mM. Cells were pelleted, washed and resuspended in ChIP sonication buffer (1% Triton X-100, 0.1% Deoxycholate, 50 mM Tris 8.1, 150 mM NaCl, 5 mM EDTA) containing protease and phosphatase inhibitors (Roche). Samples were sonicated for a total of 3 minutes in 30 second cycles with 1 minute breaks. Sonicated samples were centrifuged for 15 minutes to clear the lysates, and resulting whole cell extracts were used for pull-downs. Antibodies directed against GFP (Ab-1; Neomarkers), Brg1 (G-7; Santa Cruz) and Flag (M2; SIGMA) were incubated with equal proportions of whole cell extracts at 1:30 dilution overnight, rotating at 4° C. 1:1 Protein A and Protein G agarose beads (GE Healthcare) were added and

incubated for 2 hours at 4° C. Beads were then pelleted and washed with high salt wash buffer (1% Triton X-100, 0.1% Deoxycholate, 50 mM Tris-8.1, 500 mM NaCl, 5 mM EDTA), followed by LiCl immune complex buffer (250 mM LiCl, 0.5% IGEPAL-CA630, 0.5% Deoxycholate, 10 mM Tris-8.1, 1 mM EDTA), and TE (10 mM Tris-8.1, 1 mM EDTA) prior to suspension in Elution buffer (1% SDS, 0.1 M NaHCO₂, 0.01 mg/ml salmon sperm DNA (GE Healthcare)). Crosslinks were reversed at 65° C. overnight, beads were pelleted, and resulting supernatant was incubated with 0.4 mg/mL Proteinase K (SIGMA) for 2 hours at 37° C. DNA from each sample was purified using Qiagen PCR purification columns following the manufacturer's instructions. Samples were resuspended in 1004 10 mM Tris-8.1 and 24 were used for each Sybr green™ PCR reaction (Applied Biosystems). Enrichment was calculated by 2^Δ average(Ctreference-Ctexprimento) for each genomic region of interest. Reference samples were the GFP, both Brg1 and FLAG samples were experimental. Statistical analyses were performed on log2 transformed data for 3 independent biological replicates.

Primer Sequences:

[0487]

β-Actin:		
F	TCGAGCCATAAAAGGCAACT	(SEQ ID NO: 33)
R	TCTCCCTCCTCTCTTCCTC	(SEQ ID NO: 34)
EGFR regulatory element:		
F	CCTTGATAGATTGGGCTGAG	(SEQ ID NO: 35)
R	AGTTTGGGGGTGAAGAAAG	(SEQ ID NO: 36)
30kb upstream of regulatory element:		
F	GGCTGAGACAGAGGAACAC	(SEQ ID NO: 37)
R	CCATCTCAGCCTCCAAGTA	(SEQ ID NO: 38)

REFERENCES

[0488] 1. Jemal, A. et al. Global cancer statistics. *CA: A Cancer Journal for Clinicians* 61, 69-90 (2011).

[0489] 2. Siegel, R., Naishadham, D. & Jemal, A. Cancer statistics, 2013. *CA: A Cancer Journal for Clinicians* 63, 11-30 (2013).

[0490] 3. Langer, C. J. et al. Patterns of care survey (PCS) in lung cancer: how well does current U.S. practice with chemotherapy in the non-metastatic setting follow the literature? *Lung Cancer* 48, 93-102 (2005).

[0491] 4. Zornosa, C. et al. First-Line Systemic Therapy Practice Patterns and Concordance With NCCN Guidelines for Patients Diagnosed With Metastatic NSCLC Treated at NCCN Institutions. *Journal of the National Comprehensive Cancer Network* 10, 847-856 (2012).

[0492] 5. Movsas, B. et al. Randomized Phase II Trial of Cisplatin, Etoposide, and Radiation Followed by Gemcitabine Alone or by Combined Gemcitabine and Docetaxel in Stage III A/B Unresectable Non-small Cell Lung Cancer. *Journal of Thoracic Oncology* 5, 673-679 10.1097/JTO.0b013e3181d60e8f (2010).

[0493] 6. Jalal, S. I. et al. Updated survival and outcomes for older adults with inoperable stage III non-small-cell lung cancer treated with cisplatin, etoposide, and concurrent chest radiation with or without consolidation doc-

etaxel: analysis of a phase III trial from the Hoosier Oncology Group (HOG) and US Oncology. *Annals of Oncology* 23, 1730-1738 (2013).

[0494] 7. Wang, L. et al. Randomized phase II study of concurrent cisplatin/etoposide or paclitaxel/carboplatin and thoracic radiotherapy in patients with stage III non-small cell lung cancer. *Lung Cancer* 77, 89-96 (2012).

[0495] 8. Siddik, Z. H. Cisplatin: mode of cytotoxic action and molecular basis of resistance. *Oncogene* 22, 7265-7279 (2003).

[0496] 9. Deweese, J. E. & Osheroff, N. The DNA cleavage reaction of topoisomerase II: wolf in sheep's clothing. *Nucleic Acids Research* 37, 738-748 (2009).

[0497] 10. Dykhuizen, E. C. et al. BAF complexes facilitate decatenation of DNA by topoisomerase II. *Nature advance online publication* (2013).

[0498] 11. Baylin, S. B. & Jones, P. A. A decade of exploring the cancer epigenome—biological and translational implications. *Nat Rev Cancer* 11, 726-734 (2011).

[0499] 12. Simon, J. A. & Lange, C. A. Roles of the EZH2 histone methyltransferase in cancer epigenetics. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 647, 21-29 (2008).

[0500] 13. Kim, E. et al. Phosphorylation of EZH2 Activates STAT3 Signaling via STAT3 Methylation and Promotes Tumorigenicity of Glioblastoma Stem-like Cells. *Cancer Cell* 23, 839-852 (2013).

[0501] 14. He, A. et al. PRC2 directly methylates GATA4 and represses its transcriptional activity. *Genes & Development* 26, 37-42 (2012).

[0502] 15. Xu, K. et al. EZH2 Oncogenic Activity in Castration-Resistant Prostate Cancer Cells Is Polycomb-Independent. *Science* 338, 1465-1469 (2012).

[0503] 16. Chou, D. M. et al. A chromatin localization screen reveals poly (ADP ribose)-regulated recruitment of the repressive polycomb and NuRD complexes to sites of DNA damage. *Proceedings of the National Academy of Sciences* 107, 18475-18480 (2010).

[0504] 17. Sauvageau, M. & Sauvageau, G. Polycomb Group Proteins: Multi-Faceted Regulators of Somatic Stem Cells and Cancer. *Cell Stem Cell* 7, 299-313 (2010).

[0505] 18. Tan, J. et al. Pharmacologic disruption of Polycomb-repressive complex 2-mediated gene repression selectively induces apoptosis in cancer cells. *Genes & Development* 21, 1050-1063 (2007).

[0506] 19. Avan, A. et al. Molecular Mechanisms Involved in the Synergistic Interaction of the EZH2 Inhibitor 3-Deazaneplanocin A with Gemcitabine in Pancreatic Cancer Cells. *Molecular Cancer Therapeutics* 11, 1735-1746 (2012).

[0507] 20. Choudhury, S. R. et al. (–)-Epigallocatechin-3-gallate and DZNep reduce polycomb protein level via a proteasome-dependent mechanism in skin cancer cells. *Carcinogenesis* 32, 1525-1532 (2011).

[0508] 21. Crea, F. et al. Pharmacologic disruption of Polycomb Repressive Complex 2 inhibits tumorigenicity and tumor progression in prostate cancer. *Molecular Cancer* 10, 40 (2011).

[0509] 22. Puppe, J. et al. BRCA1-deficient mammary tumor cells are dependent on EZH2 expression and sensitive to Polycomb Repressive Complex 2-inhibitor 3-deazaneplanocin A. *Breast Cancer Research* 11, R63 (2009).

- [0510] 23. Kikuchi, J. et al. Epigenetic therapy with 3-deazaneplanocin A, an inhibitor of the histone methyltransferase EZH2, inhibits growth of non-small cell lung cancer cells. *Lung Cancer* 78, 138-143 (2012).
- [0511] 24. Miranda, T. B. et al. DZNep is a global histone methylation inhibitor that reactivates developmental genes not silenced by DNA methylation. *Molecular Cancer Therapeutics* 8, 1579-1588 (2009).
- [0512] 25. Qi, W. et al. Selective inhibition of Ezh2 by a small molecule inhibitor blocks tumor cells proliferation. *Proceedings of the National Academy of Sciences* 109, 21360-21365 (2012).
- [0513] 26. Knutson, S. K. et al. Durable tumor regression in genetically altered malignant rhabdoid tumors by inhibition of methyltransferase EZH2. *Proceedings of the National Academy of Sciences* 110, 7922-7927 (2012).
- [0514] 27. McCabe, M. T. et al. EZH2 inhibition as a therapeutic strategy for lymphoma with EZH2-activating mutations. *Nature* 492, 108-112 (2012).
- [0515] 28. Herbst, R. S., Heymach, J. V. & Lippman, S. M. Lung Cancer. *New England Journal of Medicine* 359, 1367-1380 (2008).
- [0516] 29. Imielinski, M. et al. Mapping the Hallmarks of Lung Adenocarcinoma with Massively Parallel Sequencing. *Cell* 150, 1107-1120 (2012).
- [0517] 30. TCGA. Comprehensive genomic characterization of squamous cell lung cancers. *Nature* 489, 519-525 (2012).
- [0518] 31. Rodriguez-Nieto, S. et al. Massive parallel DNA pyrosequencing analysis of the tumor suppressor BRG1/SMARCA4 in lung primary tumors. *Human Mutation* 32, E1999-E2017 (2011).
- [0519] 32. Lynch, T. J. et al. Activating Mutations in the Epidermal Growth Factor Receptor Underlying Responsiveness of Non-Small-Cell Lung Cancer to Gefitinib. *New England Journal of Medicine* 350, 2129-2139 (2004).
- [0520] 33. Paez, J. G. et al. EGFR Mutations in Lung Cancer: Correlation with Clinical Response to Gefitinib Therapy. *Science* 304, 1497-1500 (2004).
- [0521] 34. Ji, H. et al. The impact of human EGFR kinase domain mutations on lung tumorigenesis and in vivo sensitivity to EGFR-targeted therapies. *Cancer Cell* 9, 485-495 (2006).
- [0522] 35. Rhodes, D. R. et al. Oncomine 3.0: genes, pathways, and networks in a collection of 18,000 cancer gene expression profiles. *Neoplasia* 9(2007).
- [0523] 36. Beer, D. G. et al. Gene-expression profiles predict survival of patients with lung adenocarcinoma. *Nat Med* 8, 816-824 (2002).
- [0524] 37. Garber, M. E. et al. Diversity of gene expression in adenocarcinoma of the lung. *Proceedings of the National Academy of Sciences* 98, 13784-13789 (2001).
- [0525] 38. Gordon, G. J. et al. Translation of Microarray Data into Clinically Relevant Cancer Diagnostic Tests Using Gene Expression Ratios in Lung Cancer and Mesothelioma. *Cancer Research* 62, 4963-4967 (2002).
- [0526] 39. Landi, M. T. et al. Gene Expression Signature of Cigarette Smoking and Its Role in Lung Adenocarcinoma Development and Survival. *PLoS ONE* 3, e1651 (2008).
- [0527] 40. Rohrbeck, A. et al. Gene expression profiling for molecular distinction and characterization of laser captured primary lung cancers. *Journal of Translational Medicine* 6, 69 (2008).
- [0528] 41. Shedden, K. et al. Gene expression-based survival prediction in lung adenocarcinoma: a multi-site, blinded validation study. *Nat Med* 14, 822-827 (2008).
- [0529] 42. Su, A. I. et al. Molecular Classification of Human Carcinomas by Use of Gene Expression Signatures. *Cancer Research* 61, 7388-7393 (2001).
- [0530] 43. Yu, K. et al. A Precisely Regulated Gene Expression Cassette Potently Modulates Metastasis and Survival in Multiple Solid Cancers. *PLoS Genet* 4, e1000129 (2008).
- [0531] 44. Medina, P. P. et al. Frequent BRG1/SMARCA4-inactivating mutations in human lung cancer cell lines. *Human Mutation* 29, 617-622 (2008).
- [0532] 45. Yamamoto, H. et al. PIK3CA Mutations and Copy Number Gains in Human Lung Cancers. *Cancer Research* 68, 6913-6921 (2008).
- [0533] 46. Ougolkov, A. V., Bilim, V. N. & Billadeau, D. D. Regulation of Pancreatic Tumor Cell Proliferation and Chemoresistance by the Histone Methyltransferase Enhancer of Zeste Homologue 2. *Clinical Cancer Research* 14, 6790-6796 (2008).
- [0534] 47. Lv, Y. et al. The expression and significance of the enhancer of zest homolog 2 in lung adenocarcinoma. *Oncology Reports* 28, 147-154 (2013).
- [0535] 48. Li, D. et al. Bronchial and Peripheral Murine Lung Carcinomas Induced by T790M-L858R Mutant EGFR Respond to HKI-272 and Rapamycin Combination Therapy. *Cancer Cell* 12, 81-93 (2007).
- [0536] 49. Jackson, E. L. et al. The Differential Effects of Mutant p53 Alleles on Advanced Murine Lung Cancer. *Cancer Research* 65, 10280-10288 (2005).
- [0537] 50. Wilson, B. G. et al. Epigenetic Antagonism between Polycomb and SWI/SNF Complexes during Oncogenic Transformation. *Cancer Cell* 18, 316-328 (2010).
- [0538] 51. ENCODE. An integrated encyclopedia of DNA elements in the human genome. *Nature* 489, 57-74 (2013).
- [0539] 52. Rothenberg, S. M. et al. Modeling oncogene addiction using RNA interference. *Proceedings of the National Academy of Sciences* 105, 12480-12484 (2008).
- [0540] 53. Matsubara, D. et al. Lung cancer with loss of BRG1/BRM, shows epithelial mesenchymal transition phenotype and distinct histologic and genetic features. *Cancer Science* 104, 266-273 (2013).
- [0541] 54. Oike, T. et al. A Synthetic Lethality-Based Strategy to Treat Cancers Harboring a Genetic Deficiency in the Chromatin Remodeling Factor BRG1. *Cancer Research* 73, 5508-5518 (2013).
- [0542] 55. Engelman, J. A. & Janne, P. A. Factors Predicting Response to EGFR Tyrosine Kinase Inhibitors. *Semin Respir Crit Care Med* 26, 314-322 (2005).
- [0543] 56. Giaccone, G. Epidermal Growth Factor Receptor Inhibitors in the Treatment of Non-Small-Cell Lung Cancer. *Journal of Clinical Oncology* 23, 3235-3242 (2005).
- [0544] 57. Simon, J. A. & Kingston, R. E. Mechanisms of Polycomb gene silencing: knowns and unknowns. *Nat Rev Mol Cell Biol* 10, 697-708 (2009).

- [0547] 58. Dai, Y., Ngo, D., Jacob, J., Forman, L. W. & Faller, D. V. Prohibitin and the SWI/SNF ATPase subunit BRG1 are required for effective androgen antagonist-mediated transcriptional repression of androgen receptor-regulated genes. *Carcinogenesis* 29, 1725-1733 (2008).
- [0548] 59. Tolstorukov, M. Y. et al. Swi/Snf chromatin remodeling/tumor suppressor complex establishes nucleosome occupancy at target promoters. *Proceedings of the National Academy of Sciences* 110, 10165-10170 (2013).
- [0549] 60. Ramirez-Carrozzi, V. R. et al. Selective and antagonistic functions of SWI/SNF and Mi-2 nucleosome remodeling complexes during an inflammatory response. *Genes & Development* 20, 282-296 (2006).
- [0550] 61. Engelman, J. A. et al. Allelic dilution obscures detection of a biologically significant resistance mutation in EGFR-amplified lung cancer. *The Journal of Clinical Investigation* 116, 2695-2706 (2006).
- [0551] 62. Fillmore, C. M. et al. Estrogen expands breast cancer stem-like cells through paracrine FGF/Tbx3 signaling. *Proceedings of the National Academy of Sciences* 107, 21737-21742 (2010).
- [0552] 63. Zacharek, S. J. et al. Lung Stem Cell Self-Renewal Relies on BMII-Dependent Control of Expression at Imprinted Loci. *Cell Stem Cell* 9, 272-281 (2010).
- [0553] 64. Irizarry, R. A. et al. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4, 249-264 (2003).
- [0554] 65. Smyth, G. K., Michaud, J. I. & Scott, H. S. Use of within-array replicate spots for assessing differential expression in microarray experiments. *Bioinformatics* 21, 2067-2075 (2005).
- [0555] 66. Hochberg, Y. & Benjamini, Y. More powerful procedures for multiple hypothesis testing. *Statistical Medicine*, 811-818 (1990).
- [0556] 67. Curtis, S. J. et al. Primary Tumor Genotype Is an Important Determinant in Identification of Lung Cancer Propagating Cells. *Cell Stem Cell* 7, 127-133 (2010).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 38

<210> SEQ ID NO 1

<211> LENGTH: 751

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
1          5          10          15

Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
          20          25          30

Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
          35          40          45

Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
          50          55          60

Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
65          70          75          80

Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
          85          90          95

Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
          100         105         110

Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
          115         120         125

His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
          130         135         140

Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp
          145         150         155         160

Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn
          165         170         175

Ala Leu Gly Gln Tyr Asn Asp Asp Asp Asp Asp Asp Gly Asp Asp
          180         185         190

Pro Glu Glu Arg Glu Glu Lys Gln Lys Asp Leu Glu Asp His Arg Asp
          195         200         205

Asp Lys Glu Ser Arg Pro Pro Arg Lys Phe Pro Ser Asp Lys Ile Phe
          210         215         220

```

-continued

Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu	Leu	225	230	235	240
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala	Leu	245	250	255	
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser	Val	260	265	270	
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg	Arg	275	280	285	
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Arg	Lys	Cys	Asn	Tyr	Ser	Phe	290	295	300	
His	Ala	Thr	Pro	Asn	Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	305	310	315	320
Asp	Asn	Lys	Pro	Cys	Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	325	330	335	
Lys	Glu	Phe	Ala	Ala	Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	340	345	350	
Lys	Arg	Pro	Gly	Gly	Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	355	360	365	
Arg	Pro	Ser	Thr	Pro	Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	370	375	380	
Ser	Asp	Arg	Glu	Ala	Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	385	390	395	400
Glu	Glu	Glu	Glu	Lys	Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	405	410	415	
Ser	Arg	Cys	Gln	Thr	Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	420	425	430	
Glu	Asn	Val	Glu	Trp	Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	435	440	445	
Ile	Gly	Thr	Tyr	Tyr	Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	450	455	460	
Thr	Lys	Thr	Cys	Arg	Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	465	470	475	480
Ile	Ile	Ala	Pro	Ala	Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	485	490	495	
Lys	Lys	Arg	Lys	His	Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	500	505	510	
Leu	Lys	Lys	Asp	Gly	Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	515	520	525	
Asp	His	Pro	Arg	Gln	Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	530	535	540	
Gln	Asn	Phe	Cys	Glu	Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	545	550	555	560
Arg	Phe	Pro	Gly	Cys	Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	565	570	575	
Pro	Cys	Tyr	Leu	Ala	Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	580	585	590	
Cys	Gly	Ala	Ala	Asp	His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	595	600	605	
Cys	Ser	Ile	Gln	Arg	Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	610	615	620	

-continued

Asp	Val	Ala	Gly	Trp	Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn
625					630					635					640
Glu	Phe	Ile	Ser	Glu	Tyr	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala
			645						650					655	
Asp	Arg	Arg	Gly	Lys	Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe
			660					665					670		
Asn	Leu	Asn	Asn	Asp	Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys
	675						680					685			
Ile	Arg	Phe	Ala	Asn	His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val
	690					695					700				
Met	Met	Val	Asn	Gly	Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala
705					710					715					720
Ile	Gln	Thr	Gly	Glu	Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala
			725						730					735	
Asp	Ala	Leu	Lys	Tyr	Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro	
		740						745					750		

<210> SEQ ID NO 2

<211> LENGTH: 2723

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

ggcggcgctt gattgggctg ggggggccaa ataaaagcga tggcgattgg gctgccgcgt      60
ttggcgctcg gtcgggtcgc gtccgacacc cgggtgggact cagaaggcag tggagccccg      120
gcggcggcgg cggcggcgcg cgggggcgac gcgcgggaac aacgcgagtc ggcgcgcggg      180
acgaagaata atcatgggcc agactgggaa gaaatctgag aagggaccag ttgttgggcg      240
gaagcgtgta aaatcagagt acatgcgact gagacagctc aagaggttca gacgagctga      300
tgaagtaaag agtatgttta gttccaatcg tcagaaaatt ttggaaagaa cggaatctt      360
aaaccaagaa tggaacagc gaaggataca gcctgtgcac atcctgactt ctgtgagctc      420
attgcgcggg actagggagt gttcggtgac cagtgacttg gattttccaa cacaagtcac      480
cccattaaag actctgaatg cagttgcttc agtaccata atgtattctt ggtctcccct      540
acagcagaat tttatggtgg aagatgaaac tgttttacat aacattcctt atatgggaga      600
tgaagtttta gatcaggatg gtactttcat tgaagaacta ataaaaaatt atgatgggaa      660
agtacacggg gatagagaat gtgggtttat aaatgatgaa attttgtgag agttggtgaa      720
tgcccttggt caatataatg atgatgacga tgatgatgat ggagacgac ctgaagaaag      780
agaagaaaag cagaaagatc tggaggatca ccgagatgat aaagaaagcc gccacctcgc      840
gaaatttcct tctgataaaa tttttgaagc catttcctca atgtttccag ataagggcac      900
agcagaagaa ctaaggaaa aatataaaga actcaccgaa cagcagctcc caggcgccact      960
tcctcctgaa tgtaccccca acatagatgg accaaatgct aaatctgttc agagagagca     1020
aagcttacac tcctttcata cgtttttctg taggcgatgt tttaaatag actgcttcct     1080
acatcgtaag tgcaattatt cttttcatgc aacacccaac acttataagc ggaagaacac     1140
agaaacagct ctagacaaca aaccttggtg accacagtgt taccagcatt tggagggagc     1200
aaaggagttt gctgctgctc tcaccgctga gcggataaag accccaccaa aacgtccagc     1260
agggcgcgag agaggacggc ttccaataaa cagtagcagg ccagcacccc ccaccattaa     1320

```


-continued

tgtgctggaa tcaaaggata cagacagtga tagggaagca gggactgaaa cggggggaga	1380
gaacaatgat aaagaagaag aagagaagaa agatgaaact tcgagctcct ctgaagcaaa	1440
ttctcggtgt caaacaccaa taaagatgaa gccaaatatt gaacctcctg agaatgtgga	1500
gtggagtggg gctgaagcct caatgtttag agtcctcatt ggcacttact atgacaattt	1560
ctgtgccatt gctaggttaa ttgggaccaa aacatgtaga caggtgtatg agtttagagt	1620
caaagaatct agcatcatag ctccagctcc cgctgaggat gtggatactc ctccaaggaa	1680
aaagaagagg aaacaccggt tgtgggctgc aactgcaga aagatacagc tgaaaaagga	1740
cggctcctct aaccatgttt acaactatca accctgtgat catccacggc agccttgtga	1800
cagttcgtgc ccttgtgtga tagcacaaaa tttttgtgaa aagttttgtc aatgtagttc	1860
agagtgtcaa aaccgctttc cggtatgccc ctgcaaagca cagtgaaca ccaagcagtg	1920
cccgtgttac ctggctgtcc gagagtgtga ccttgacctc tgtcttactt gtggagccgc	1980
tgaccattgg gacagtaaaa atgtgtcctg caagaactgc agtattcagc ggggctccaa	2040
aaagcatcta ttgctggcac catctgacgt ggcaggctgg gggattttta tcaaagatcc	2100
tgtgcagaaa aatgaattca tctcagaata ctgtggagag attatttctc aagatgaagc	2160
tgacagaaga gggaaagtgt atgataaata catgtgcagc tttctgttca acttgaacaa	2220
tgattttgtg gtggatgcaa cccgcaaggg taacaaaatt cgttttgcaa atcattcggg	2280
aatccaaac tgctatgcaa aagttatgat ggttaacggt gatcacagga taggtatttt	2340
tgccaagaga gccatccaga ctggcgaaga gctgtttttt gattacagat acagccaggc	2400
tgatgccctg aagtatgtcg gcatcgaaag agaaatggaa atcccttgac atctgctacc	2460
tcctcccccc tcctctgaaa cagctgcctt agcttcagga acctcgagta ctgtgggcaa	2520
tttagaaaaa gaacatgcag tttgaaatc tgaatttgca aagtactgta agaataattt	2580
atagtaatga gtttaaaaat caacttttta ttgccttctc accagctgca aagtgttttg	2640
taccagttaa tttttgcaat aatgcagtat ggtacatttt tcaactttga ataaagaata	2700
cttgaacttg tccttgttga atc	2723

<210> SEQ ID NO 3

<211> LENGTH: 1647

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Met	Ser	Thr	Pro	Asp	Pro	Pro	Leu	Gly	Gly	Thr	Pro	Arg	Pro	Gly	Pro
1				5				10					15		
Ser	Pro	Gly	Pro	Gly	Pro	Ser	Pro	Gly	Ala	Met	Leu	Gly	Pro	Ser	Pro
	20						25					30			
Gly	Pro	Ser	Pro	Gly	Ser	Ala	His	Ser	Met	Met	Gly	Pro	Ser	Pro	Gly
	35					40					45				
Pro	Pro	Ser	Ala	Gly	His	Pro	Ile	Pro	Thr	Gln	Gly	Pro	Gly	Gly	Tyr
	50					55				60					
Pro	Gln	Asp	Asn	Met	His	Gln	Met	His	Lys	Pro	Met	Glu	Ser	Met	His
65					70				75					80	
Glu	Lys	Gly	Met	Ser	Asp	Asp	Pro	Arg	Tyr	Asn	Gln	Met	Lys	Gly	Met
		85						90					95		
Gly	Met	Arg	Ser	Gly	Gly	His	Ala	Gly	Met	Gly	Pro	Pro	Pro	Ser	Pro
	100						105						110		

-continued

Met	Asp	Gln	His	Ser	Gln	Gly	Tyr	Pro	Ser	Pro	Leu	Gly	Gly	Ser	Glu
	115						120					125			
His	Ala	Ser	Ser	Pro	Val	Pro	Ala	Ser	Gly	Pro	Ser	Ser	Gly	Pro	Gln
	130					135					140				
Met	Ser	Ser	Gly	Pro	Gly	Gly	Ala	Pro	Leu	Asp	Gly	Ala	Asp	Pro	Gln
145					150					155					160
Ala	Leu	Gly	Gln	Gln	Asn	Arg	Gly	Pro	Thr	Pro	Phe	Asn	Gln	Asn	Gln
					165				170					175	
Leu	His	Gln	Leu	Arg	Ala	Gln	Ile	Met	Ala	Tyr	Lys	Met	Leu	Ala	Arg
			180					185					190		
Gly	Gln	Pro	Leu	Pro	Asp	His	Leu	Gln	Met	Ala	Val	Gln	Gly	Lys	Arg
		195					200					205			
Pro	Met	Pro	Gly	Met	Gln	Gln	Gln	Met	Pro	Thr	Leu	Pro	Pro	Pro	Ser
	210					215					220				
Val	Ser	Ala	Thr	Gly	Pro	Gly	Pro	Gly	Pro	Gly	Pro	Gly	Pro	Gly	Pro
225					230					235					240
Gly	Pro	Gly	Pro	Ala	Pro	Pro	Asn	Tyr	Ser	Arg	Pro	His	Gly	Met	Gly
				245					250					255	
Gly	Pro	Asn	Met	Pro	Pro	Pro	Gly	Pro	Ser	Gly	Val	Pro	Pro	Gly	Met
		260						265					270		
Pro	Gly	Gln	Pro	Pro	Gly	Gly	Pro	Pro	Lys	Pro	Trp	Pro	Glu	Gly	Pro
		275					280					285			
Met	Ala	Asn	Ala	Ala	Ala	Pro	Thr	Ser	Thr	Pro	Gln	Lys	Leu	Ile	Pro
	290					295					300				
Pro	Gln	Pro	Thr	Gly	Arg	Pro	Ser	Pro	Ala	Pro	Pro	Ala	Val	Pro	Pro
305					310					315					320
Ala	Ala	Ser	Pro	Val	Met	Pro	Pro	Gln	Thr	Gln	Ser	Pro	Gly	Gln	Pro
				325					330					335	
Ala	Gln	Pro	Ala	Pro	Met	Val	Pro	Leu	His	Gln	Lys	Gln	Ser	Arg	Ile
		340						345					350		
Thr	Pro	Ile	Gln	Lys	Pro	Arg	Gly	Leu	Asp	Pro	Val	Glu	Ile	Leu	Gln
		355					360					365			
Glu	Arg	Glu	Tyr	Arg	Leu	Gln	Ala	Arg	Ile	Ala	His	Arg	Ile	Gln	Glu
	370					375				380					
Leu	Glu	Asn	Leu	Pro	Gly	Ser	Leu	Ala	Gly	Asp	Leu	Arg	Thr	Lys	Ala
385				390						395					400
Thr	Ile	Glu	Leu	Lys	Ala	Leu	Arg	Leu	Leu	Asn	Phe	Gln	Arg	Gln	Leu
			405					410						415	
Arg	Gln	Glu	Val	Val	Val	Cys	Met	Arg	Arg	Asp	Thr	Ala	Leu	Glu	Thr
		420						425					430		
Ala	Leu	Asn	Ala	Lys	Ala	Tyr	Lys	Arg	Ser	Lys	Arg	Gln	Ser	Leu	Arg
		435					440					445			
Glu	Ala	Arg	Ile	Thr	Glu	Lys	Leu	Glu	Lys	Gln	Gln	Lys	Ile	Glu	Gln
	450					455					460				
Glu	Arg	Lys	Arg	Arg	Gln	Lys	His	Gln	Glu	Tyr	Leu	Asn	Ser	Ile	Leu
465					470					475					480
Gln	His	Ala	Lys	Asp	Phe	Lys	Glu	Tyr	His	Arg	Ser	Val	Thr	Gly	Lys
			485						490					495	
Ile	Gln	Lys	Leu	Thr	Lys	Ala	Val	Ala	Thr	Tyr	His	Ala	Asn	Thr	Glu
		500						505						510	

-continued

Arg	Glu	Gln	Lys	Lys	Glu	Asn	Glu	Arg	Ile	Glu	Lys	Glu	Arg	Met	Arg
	515						520					525			
Arg	Leu	Met	Ala	Glu	Asp	Glu	Glu	Gly	Tyr	Arg	Lys	Leu	Ile	Asp	Gln
	530					535					540				
Lys	Lys	Asp	Lys	Arg	Leu	Ala	Tyr	Leu	Leu	Gln	Gln	Thr	Asp	Glu	Tyr
545					550					555					560
Val	Ala	Asn	Leu	Thr	Glu	Leu	Val	Arg	Gln	His	Lys	Ala	Ala	Gln	Val
			565						570					575	
Ala	Lys	Glu	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Ala	Glu	Asn	Ala
			580					585					590		
Glu	Gly	Gln	Thr	Pro	Ala	Ile	Gly	Pro	Asp	Gly	Glu	Pro	Leu	Asp	Glu
		595					600					605			
Thr	Ser	Gln	Met	Ser	Asp	Leu	Pro	Val	Lys	Val	Ile	His	Val	Glu	Ser
	610					615					620				
Gly	Lys	Ile	Leu	Thr	Gly	Thr	Asp	Ala	Pro	Lys	Ala	Gly	Gln	Leu	Glu
625					630					635					640
Ala	Trp	Leu	Glu	Met	Asn	Pro	Gly	Tyr	Glu	Val	Ala	Pro	Arg	Ser	Asp
				645					650					655	
Ser	Glu	Glu	Ser	Gly	Ser	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu	Glu
			660					665					670		
Gln	Pro	Gln	Ala	Ala	Gln	Pro	Pro	Thr	Leu	Pro	Val	Glu	Glu	Lys	Lys
		675					680					685			
Lys	Ile	Pro	Asp	Pro	Asp	Ser	Asp	Asp	Val	Ser	Glu	Val	Asp	Ala	Arg
	690					695					700				
His	Ile	Ile	Glu	Asn	Ala	Lys	Gln	Asp	Val	Asp	Asp	Glu	Tyr	Gly	Val
705				710						715					720
Ser	Gln	Ala	Leu	Ala	Arg	Gly	Leu	Gln	Ser	Tyr	Tyr	Ala	Val	Ala	His
				725					730					735	
Ala	Val	Thr	Glu	Arg	Val	Asp	Lys	Gln	Ser	Ala	Leu	Met	Val	Asn	Gly
			740					745					750		
Val	Leu	Lys	Gln	Tyr	Gln	Ile	Lys	Gly	Leu	Glu	Trp	Leu	Val	Ser	Leu
		755					760					765			
Tyr	Asn	Asn	Asn	Leu	Asn	Gly	Ile	Leu	Ala	Asp	Glu	Met	Gly	Leu	Gly
	770					775					780				
Lys	Thr	Ile	Gln	Thr	Ile	Ala	Leu	Ile	Thr	Tyr	Leu	Met	Glu	His	Lys
785					790					795					800
Arg	Ile	Asn	Gly	Pro	Phe	Leu	Ile	Ile	Val	Pro	Leu	Ser	Thr	Leu	Ser
				805					810					815	
Asn	Trp	Ala	Tyr	Glu	Phe	Asp	Lys	Trp	Ala	Pro	Ser	Val	Val	Lys	Val
			820					825					830		
Ser	Tyr	Lys	Gly	Ser	Pro	Ala	Ala	Arg	Arg	Ala	Phe	Val	Pro	Gln	Leu
		835					840					845			
Arg	Ser	Gly	Lys	Phe	Asn	Val	Leu	Leu	Thr	Thr	Tyr	Glu	Tyr	Ile	Ile
	850					855					860				
Lys	Asp	Lys	His	Ile	Leu	Ala	Lys	Ile	Arg	Trp	Lys	Tyr	Met	Ile	Val
865					870					875					880
Asp	Glu	Gly	His	Arg	Met	Lys	Asn	His	His	Cys	Lys	Leu	Thr	Gln	Val
			885					890						895	
Leu	Asn	Thr	His	Tyr	Val	Ala	Pro	Arg	Arg	Leu	Leu	Leu	Thr	Gly	Thr
			900					905						910	
Pro	Leu	Gln	Asn	Lys	Leu	Pro	Glu	Leu	Trp	Ala	Leu	Leu	Asn	Phe	Leu

-continued

915					920					925					
Leu 930	Pro	Thr	Ile	Phe	Lys	Ser 935	Cys	Ser	Thr	Phe	Glu 940	Gln	Trp	Phe	Asn
Ala 945	Pro	Phe	Ala	Met	Thr 950	Gly	Glu	Lys	Val	Asp 955	Leu	Asn	Glu	Glu	Glu 960
Thr	Ile	Leu	Ile	Ile 965	Arg	Arg	Leu	His	Lys 970	Val	Leu	Arg	Pro	Phe 975	Leu
Leu	Arg	Arg	Leu	Lys 980	Lys	Glu	Val 985	Glu	Ala	Gln	Leu	Pro 990	Glu	Lys	Val
Glu	Tyr 995	Val	Ile	Lys	Cys	Asp 1000	Met	Ser	Ala	Leu	Gln	Arg 1005	Val	Leu	Tyr
Arg 1010	His	Met	Gln	Ala	Lys	Gly 1015	Val	Leu	Leu	Thr	Asp 1020	Gly	Ser	Glu	
Lys 1025	Asp	Lys	Lys	Gly	Lys	Gly 1030	Gly	Thr	Lys	Thr	Leu 1035	Met	Asn	Thr	
Ile 1040	Met	Gln	Leu	Arg	Lys	Ile 1045	Cys	Asn	His	Pro	Tyr 1050	Met	Phe	Gln	
His 1055	Ile	Glu	Glu	Ser	Phe	Ser 1060	Glu	His	Leu	Gly	Phe 1065	Thr	Gly	Gly	
Ile 1070	Val	Gln	Gly	Leu	Asp	Leu 1075	Tyr	Arg	Ala	Ser	Gly 1080	Lys	Phe	Glu	
Leu 1085	Leu	Asp	Arg	Ile	Leu	Pro 1090	Lys	Leu	Arg	Ala	Thr 1095	Asn	His	Lys	
Val 1100	Leu	Leu	Phe	Cys	Gln	Met 1105	Thr	Ser	Leu	Met	Thr 1110	Ile	Met	Glu	
Asp 1115	Tyr	Phe	Ala	Tyr	Arg	Gly 1120	Phe	Lys	Tyr	Leu	Arg 1125	Leu	Asp	Gly	
Thr 1130	Thr	Lys	Ala	Glu	Asp	Arg 1135	Gly	Met	Leu	Leu	Lys 1140	Thr	Phe	Asn	
Glu 1145	Pro	Gly	Ser	Glu	Tyr	Phe 1150	Ile	Phe	Leu	Leu	Ser 1155	Thr	Arg	Ala	
Gly 1160	Gly	Leu	Gly	Leu	Asn	Leu 1165	Gln	Ser	Ala	Asp	Thr 1170	Val	Ile	Ile	
Phe 1175	Asp	Ser	Asp	Trp	Asn	Pro 1180	His	Gln	Asp	Leu	Gln 1185	Ala	Gln	Asp	
Arg 1190	Ala	His	Arg	Ile	Gly	Gln 1195	Gln	Asn	Glu	Val	Arg 1200	Val	Leu	Arg	
Leu 1205	Cys	Thr	Val	Asn	Ser	Val 1210	Glu	Glu	Lys	Ile	Leu 1215	Ala	Ala	Ala	
Lys 1220	Tyr	Lys	Leu	Asn	Val	Asp 1225	Gln	Lys	Val	Ile	Gln 1230	Ala	Gly	Met	
Phe 1235	Asp	Gln	Lys	Ser	Ser	Ser 1240	His	Glu	Arg	Arg	Ala 1245	Phe	Leu	Gln	
Ala 1250	Ile	Leu	Glu	His	Glu	Glu 1255	Gln	Asp	Glu	Ser	Arg 1260	His	Cys	Ser	
Thr 1265	Gly	Ser	Gly	Ser	Ala	Ser 1270	Phe	Ala	His	Thr	Ala 1275	Pro	Pro	Pro	
Ala 1280	Gly	Val	Asn	Pro	Asp	Leu 1285	Glu	Glu	Pro	Pro	Leu 1290	Lys	Glu	Glu	
Asp 1295	Glu	Val	Pro	Asp	Asp	Glu 1300	Thr	Val	Asn	Gln	Met 1305	Ile	Ala	Arg	

-continued

His	Glu	Glu	Glu	Phe	Asp	Leu	Phe	Met	Arg	Met	Asp	Leu	Asp	Arg
1310						1315					1320			
Arg	Arg	Glu	Glu	Ala	Arg	Asn	Pro	Lys	Arg	Lys	Pro	Arg	Leu	Met
1325						1330					1335			
Glu	Glu	Asp	Glu	Leu	Pro	Ser	Trp	Ile	Ile	Lys	Asp	Asp	Ala	Glu
1340						1345					1350			
Val	Glu	Arg	Leu	Thr	Cys	Glu	Glu	Glu	Glu	Glu	Lys	Met	Phe	Gly
1355						1360					1365			
Arg	Gly	Ser	Arg	His	Arg	Lys	Glu	Val	Asp	Tyr	Ser	Asp	Ser	Leu
1370						1375					1380			
Thr	Glu	Lys	Gln	Trp	Leu	Lys	Ala	Ile	Glu	Glu	Gly	Thr	Leu	Glu
1385						1390					1395			
Glu	Ile	Glu	Glu	Glu	Val	Arg	Gln	Lys	Lys	Ser	Ser	Arg	Lys	Arg
1400						1405					1410			
Lys	Arg	Asp	Ser	Asp	Ala	Gly	Ser	Ser	Thr	Pro	Thr	Thr	Ser	Thr
1415						1420					1425			
Arg	Ser	Arg	Asp	Lys	Asp	Asp	Glu	Ser	Lys	Lys	Gln	Lys	Lys	Arg
1430						1435					1440			
Gly	Arg	Pro	Pro	Ala	Glu	Lys	Leu	Ser	Pro	Asn	Pro	Pro	Asn	Leu
1445						1450					1455			
Thr	Lys	Lys	Met	Lys	Lys	Ile	Val	Asp	Ala	Val	Ile	Lys	Tyr	Lys
1460						1465					1470			
Asp	Ser	Ser	Ser	Gly	Arg	Gln	Leu	Ser	Glu	Val	Phe	Ile	Gln	Leu
1475						1480					1485			
Pro	Ser	Arg	Lys	Glu	Leu	Pro	Glu	Tyr	Tyr	Glu	Leu	Ile	Arg	Lys
1490						1495					1500			
Pro	Val	Asp	Phe	Lys	Lys	Ile	Lys	Glu	Arg	Ile	Arg	Asn	His	Lys
1505						1510					1515			
Tyr	Arg	Ser	Leu	Asn	Asp	Leu	Glu	Lys	Asp	Val	Met	Leu	Leu	Cys
1520						1525					1530			
Gln	Asn	Ala	Gln	Thr	Phe	Asn	Leu	Glu	Gly	Ser	Leu	Ile	Tyr	Glu
1535						1540					1545			
Asp	Ser	Ile	Val	Leu	Gln	Ser	Val	Phe	Thr	Ser	Val	Arg	Gln	Lys
1550						1555					1560			
Ile	Glu	Lys	Glu	Asp	Asp	Ser	Glu	Gly	Glu	Glu	Ser	Glu	Glu	Glu
1565						1570					1575			
Glu	Glu	Gly	Glu	Glu	Glu	Gly	Ser	Glu	Ser	Glu	Ser	Arg	Ser	Val
1580						1585					1590			
Lys	Val	Lys	Ile	Lys	Leu	Gly	Arg	Lys	Glu	Lys	Ala	Gln	Asp	Arg
1595						1600					1605			
Leu	Lys	Gly	Gly	Arg	Arg	Arg	Pro	Ser	Arg	Gly	Ser	Arg	Ala	Lys
1610						1615					1620			
Pro	Val	Val	Ser	Asp	Asp	Asp	Ser	Glu	Glu	Glu	Gln	Glu	Glu	Asp
1625						1630					1635			
Arg	Ser	Gly	Ser	Gly	Ser	Glu	Glu	Asp						
1640						1645								

<210> SEQ ID NO 4

<211> LENGTH: 5779

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 4

ggagaggcgc cgcggtgct gagggggagg ggagcgcgcg agcgcgcgcg cagcgggggc	60
gcggttggcg cgcgtgtgtg tgaagggggg gcggtggcgc aggcggggcg gcgcgcgcgc	120
gaggcttccc ctggtttggc ggcggcggcg gcttctttgt ttcgtgaaga gaagcgagac	180
gccattctg ccccgggccc cgcgcggagg ggcgggggag gcgcgggaa gtcgacggcg	240
cggcgggctc ctgcgtctcg cctttttgcc caggctagag tgcagtgggt cggtcatggt	300
tactgcagc ctcaacctcc tggactcagc aggaggccac tgtctgcagc tcccgtaag	360
atgtccactc cagaccacc cctgggcgga actcctcggc caggtccttc cccggggcct	420
ggccttccc ctggagccat gctgggcctt agcccggtc cctcgccggg ctcgcccac	480
agcatgatgg gggccagccc agggccgccc tcagcaggac accccatccc caccagggg	540
cctggagggt accctcagga caacatgcac cagatgcaca agcccatgga gtccatgcat	600
gagaagggca tgtcggaaga cccgcgctac aaccagatga aaggaatggg gatgcggta	660
gggggccatg ctgggatggg gcccccgccc agcccatgg accagcactc ccaaggttac	720
ccctcgcccc tgggtggctc tgagcatgcc tctagtccag ttccagccag tggcccgctc	780
tcggggcccc agatgtcttc cgggccagga ggtgccccgc tggatggtgc tgaccccag	840
gccttggggc agcagaaccg gggcccaacc ccatttaacc agaaccagct gcaccagctc	900
agagctcaga tcattggcta caagatgtg gccagggggc agccctccc cgaccacctg	960
cagatggcgg tgcagggcaa gcggccgatg cccgggatgc agcagcagat gccaacgcta	1020
cctccacctc cgggtgtcgc aacaggaccc ggcctggcc ctggccctgg ccccggccg	1080
ggtcccgccc cggcacctcc aaattacagc aggcctcatg gtatgggagg gcccaacatg	1140
cctccccag gacctcggg cgtgcccccc gggatgccag gccagcctcc tggagggcct	1200
cccaagccct ggcctgaagg acccatggcg aatgctgtg ccccaacgag caccctcag	1260
aagctgatcc ccccgagcc aacgggccc ccttccccgc cgccccctgc cgtcccaacc	1320
gccgcctcgc cgtgatgcc accgcagacc cagtccccgc ggcagccggc ccagcccgcg	1380
cccatggtgc cactgcacca gaagcagagc cgcacacccc ccacccagaa gccgcggggc	1440
ctcgacctg tggagatcct gcaggagcgc gagtacaggc tgcaggctcg catcgcacac	1500
cgaattcagg aacttgaaaa ccttccccgg tccctggccc gggatttgcg aaccaaagcg	1560
accattgagc tcaaggccct caggctgtg aacttcaga ggcagctgcg ccaggagggtg	1620
gtggtgtgca tgcggaggga cacagcgtg gagacagccc tcaatgctaa ggcctacaag	1680
cgcagcaagc gccagtcct gcgcgaggcc cgcacactg agaagctgga gaagcagcag	1740
aagatcgagc aggagcgcaa gcgccggcag aagcaccagg aatacctcaa tagcattctc	1800
cagcatgcca aggatttcaa ggaatatcac agatccgta caggcaaaat ccagaagctg	1860
accaaggcag tggccacgta ccatgccaac acggagcggg agcagaagaa agagaacgag	1920
cggatcgaga aggagcgcac gcggaggctc atggctgaag atgaggagg gtaccgcaag	1980
ctcatcgacc agaagaagga caagcgctg gcctacctct tgcagcagac agacgagtac	2040
gtggctaacc tcacggagct ggtgcggcag cacaaggctg cccaggctgc caaggagaaa	2100
aagaagaaaa agaaaaagaa gaaggcagaa aatgcagaag gacagacgcc tgccattggg	2160
cggatggcg agcctctgga cgagaccagc cagatgagcg acctcccggt gaaggatgat	2220

-continued

cacgtggaga	gtgggaagat	cctcacaggc	acagatgccc	ccaaagccgg	gcagctggag	2280
gcctggctcg	agatgaaccc	ggggtatgaa	gtagctccga	ggtctgatag	tgaagaaagt	2340
ggctcagaag	aagaggaaga	ggaggaggag	gaagagcagc	cgcaggcagc	acagcctccc	2400
accctgcccc	tggaggagaa	gaagaagatt	ccagatccag	acagcgatga	cgtctctgag	2460
gtggacgcgc	ggcacatcat	tgagaatgcc	aagcaagatg	tcgatgatga	atatggcgtg	2520
tcccaggccc	ttgcacgtgg	cctgcagtcc	tactatgccg	tggcccattg	tgtaactgag	2580
agagtggaca	agcagtcagc	gcttatggtc	aatggtgtcc	tcaaacagta	ccagatcaaa	2640
ggtttgaggt	ggctgggtgc	cctgtacaac	aacaacctga	acggcaccct	ggcgcacgag	2700
atgggcctgg	ggaagaccat	ccagaccatc	gcgctcatca	cgtacctcat	ggagcacaaa	2760
cgcacatcat	ggcccttctc	catcatcgtg	cctctctcaa	cgtgttccaa	ctgggcgtac	2820
gagtttgaca	agtggtggcc	ctccgtggtg	aaggtgtctt	acaagggatc	cccagcagca	2880
agacgggcct	ttgtccccc	gctccggagt	gggaagtcca	acgtcttctg	gacgacgtac	2940
gagtacatca	tcaaagacaa	gcacatcttc	gccaaagatc	ggtggaagta	catgattgtg	3000
gacgaaggtc	accgatgaa	gaaccaccac	tgcaagctga	cgcagggtgt	caacacgcac	3060
tatgtggcac	cccgcgcct	gctgctgacg	ggcacaccgc	tgcaagaaca	gcttcccagag	3120
ctctgggcgc	tgctcaactt	cctgctgccc	accatcttca	agagctgcag	caccttcgag	3180
cagtgggtta	acgcaccctt	tgccatgacc	ggggaaaagg	tggacctgaa	tgaggaggaa	3240
accattctca	tcattccggc	tctccacaaa	gtgctgcggc	ccttcttctg	ccgacgactc	3300
aagaaggaa	tcgaggccca	gttgcctgaa	aaggtggagt	acgtcatcaa	gtgcgacatg	3360
tctgcgctgc	agcagtgct	ctaccgccac	atgcaggcca	agggcgtgct	gctgactgat	3420
ggctccgaga	aggacaagaa	gggcaaaggc	ggcaccaaga	ccctgatgaa	caccatcatg	3480
cagctgcgga	agatctgcaa	ccacccttac	atgttccagc	acatcgagga	gtccttttcc	3540
gagcacttgg	ggttcactgg	cggcattgtc	caagggtggc	acctgtaccg	agcctcgggt	3600
aaatttgagc	ttcttgatag	aattcttccc	aaactccgag	caaccaacca	caaagtgtgt	3660
ctgttctgcc	aatgacctc	cctcatgacc	atcatggaag	attactttgc	gtatcgcggc	3720
tttaaatacc	tcaggcttga	tgaaccacag	aaggcggagg	accggggcat	gctgctgaaa	3780
accttcaacg	agcccggtc	tgagtacttc	atcttctctg	tcagcaccgc	ggctgggggg	3840
ctcggcctga	acctccagtc	ggcagacact	gtgatcattt	ttgacagcga	ctggaatcct	3900
caccaggacc	tgcaagcgca	ggaccgagcc	caccgcatcg	ggcagcagaa	cagggtgcgt	3960
gtgctccgcc	tctgcaccgt	caacagcgtg	gaggagaaga	tcctagctgc	agccaagtac	4020
aagctcaacg	tggaccagaa	ggtgatccag	gccggcatgt	tcgaccagaa	gtcctccagc	4080
catgagcggc	gcgccttctc	gcaggccatc	ctggagcagc	aggagcagga	tgagagcaga	4140
cactgcagca	cgggcagcgg	cagtgccagc	ttcgcaccac	ctgcccctcc	gccagcgggc	4200
gtcaaccccg	acttgaggga	gccacctcta	aaggagggaag	acgagggtgc	cgacgacgag	4260
accgtcaacc	agatgatcgc	ccggcacgag	gaggagtgtt	atctgttcat	gcgatggac	4320
ctggaccgca	ggcgcgagga	ggcccgcac	cccaagcggg	agccgcgcct	catggaggag	4380
gacgagctcc	cctcgtggat	catcaaggac	gacgcggagg	tggagcggct	gacctgtgag	4440
gaggaggagg	agaagatgtt	cggccgtggc	tcccgcacc	gcaaggagggt	ggactacagc	4500

-continued

```

gactcactga cggagaagca gtggctcaag gccatcgagg agggcacgct ggaggagatc 4560
gaagaggagg tccggcagaa gaaatcatca cggaagcgca agcgagacag cgacgccggc 4620
tcttccaccc cgaccaccag caccgcagc cgcgacaagg acgacgagag caagaagcag 4680
aagaagcgcg ggcggccgccc tgccgagaaa ctctccccta acccacccaa cctcaccaag 4740
aagatgaaga agattgtgga tgccgtgatc aagtacaagg acagcagcag tggacgtcag 4800
ctcagcgagg tcttcatcca gctgccctcg cgaaaggagc tgcccagta ctacgagctc 4860
atccgcaagc ccgtggactt caagaagata aaggagcgca ttcgcaacca caagtaccgc 4920
agcctcaacg acctagagaa ggacgtcatg ctctgtgtcc agaacgcaca gacctcaac 4980
ctggagggct ccctgatcta tgaagactcc atcgtcttgc agtcggtctt caccagcgtg 5040
cggcagaaaa tcgagaagga ggatgacagt gaaggcgagg agagtgagga ggaggaagag 5100
ggcgaggagg aaggctccga atccgaatct cggtcctgca aagtgaagat caagcttggc 5160
cggaaggaga aggcacagga ccggctgaag ggcggccggc ggcggccgag ccgaggggtcc 5220
cgagccaagc cggtcgtgag tgacgatgac agtgaggagg aacaagagga ggaccgctca 5280
ggaagtggca gcgaagaaga ctgagccccg acattccagt ctcgaccccc agcccctcgt 5340
tccagagctg agatggcata ggccttagca gtaacgggta gcagcagatg tagtttcaga 5400
cttgagtagt aactgtataa acaaaagaat ctcccatatt tatacagcag agaagctgta 5460
ggactgtttg tgactggccc tgcctctggc tcagtagcat ctgtaacagc attaactgtc 5520
ttaaagagag agagagagaa ttccgaattg gggaacacac gatacctgtt tttcttttcc 5580
gttgctggca gtactgttgc gccgcagttt ggagtcactg tagttaagtg tggatgcatg 5640
tgcgtcaccc tccactcctc ctactgtatt ttattggaca ggtcagactc gccggggggc 5700
cggcgagggt atgtcagtg cactggatgt caaacagtaa taaattaaac caacaacaaa 5760
acgcacagcc aaaaaaaaaa 5779

```

<210> SEQ ID NO 5

<211> LENGTH: 1210

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

Met Arg Pro Ser Gly Thr Ala Gly Ala Ala Leu Leu Ala Leu Ala
1          5          10          15

Ala Leu Cys Pro Ala Ser Arg Ala Leu Glu Glu Lys Lys Val Cys Gln
20        25        30

Gly Thr Ser Asn Lys Leu Thr Gln Leu Gly Thr Phe Glu Asp His Phe
35        40        45

Leu Ser Leu Gln Arg Met Phe Asn Asn Cys Glu Val Val Leu Gly Asn
50        55        60

Leu Glu Ile Thr Tyr Val Gln Arg Asn Tyr Asp Leu Ser Phe Leu Lys
65        70        75        80

Thr Ile Gln Glu Val Ala Gly Tyr Val Leu Ile Ala Leu Asn Thr Val
85        90        95

Glu Arg Ile Pro Leu Glu Asn Leu Gln Ile Ile Arg Gly Asn Met Tyr
100       105       110

Tyr Glu Asn Ser Tyr Ala Leu Ala Val Leu Ser Asn Tyr Asp Ala Asn
115       120       125

```


-continued

Lys	Thr	Gly	Leu	Lys	Glu	Leu	Pro	Met	Arg	Asn	Leu	Gln	Glu	Ile	Leu
130						135					140				
His	Gly	Ala	Val	Arg	Phe	Ser	Asn	Asn	Pro	Ala	Leu	Cys	Asn	Val	Glu
145					150					155					160
Ser	Ile	Gln	Trp	Arg	Asp	Ile	Val	Ser	Ser	Asp	Phe	Leu	Ser	Asn	Met
				165					170					175	
Ser	Met	Asp	Phe	Gln	Asn	His	Leu	Gly	Ser	Cys	Gln	Lys	Cys	Asp	Pro
		180						185					190		
Ser	Cys	Pro	Asn	Gly	Ser	Cys	Trp	Gly	Ala	Gly	Glu	Glu	Asn	Cys	Gln
		195					200					205			
Lys	Leu	Thr	Lys	Ile	Ile	Cys	Ala	Gln	Gln	Cys	Ser	Gly	Arg	Cys	Arg
210						215					220				
Gly	Lys	Ser	Pro	Ser	Asp	Cys	Cys	His	Asn	Gln	Cys	Ala	Ala	Gly	Cys
225					230					235					240
Thr	Gly	Pro	Arg	Glu	Ser	Asp	Cys	Leu	Val	Cys	Arg	Lys	Phe	Arg	Asp
				245					250					255	
Glu	Ala	Thr	Cys	Lys	Asp	Thr	Cys	Pro	Pro	Leu	Met	Leu	Tyr	Asn	Pro
			260					265					270		
Thr	Thr	Tyr	Gln	Met	Asp	Val	Asn	Pro	Glu	Gly	Lys	Tyr	Ser	Phe	Gly
		275					280					285			
Ala	Thr	Cys	Val	Lys	Lys	Cys	Pro	Arg	Asn	Tyr	Val	Val	Thr	Asp	His
290						295					300				
Gly	Ser	Cys	Val	Arg	Ala	Cys	Gly	Ala	Asp	Ser	Tyr	Glu	Met	Glu	Glu
305					310					315					320
Asp	Gly	Val	Arg	Lys	Cys	Lys	Lys	Cys	Glu	Gly	Pro	Cys	Arg	Lys	Val
				325					330					335	
Cys	Asn	Gly	Ile	Gly	Ile	Gly	Glu	Phe	Lys	Asp	Ser	Leu	Ser	Ile	Asn
			340					345					350		
Ala	Thr	Asn	Ile	Lys	His	Phe	Lys	Asn	Cys	Thr	Ser	Ile	Ser	Gly	Asp
		355					360					365			
Leu	His	Ile	Leu	Pro	Val	Ala	Phe	Arg	Gly	Asp	Ser	Phe	Thr	His	Thr
370						375					380				
Pro	Pro	Leu	Asp	Pro	Gln	Glu	Leu	Asp	Ile	Leu	Lys	Thr	Val	Lys	Glu
385					390					395					400
Ile	Thr	Gly	Phe	Leu	Leu	Ile	Gln	Ala	Trp	Pro	Glu	Asn	Arg	Thr	Asp
				405					410					415	
Leu	His	Ala	Phe	Glu	Asn	Leu	Glu	Ile	Ile	Arg	Gly	Arg	Thr	Lys	Gln
			420					425					430		
His	Gly	Gln	Phe	Ser	Leu	Ala	Val	Val	Ser	Leu	Asn	Ile	Thr	Ser	Leu
			435					440				445			
Gly	Leu	Arg	Ser	Leu	Lys	Glu	Ile	Ser	Asp	Gly	Asp	Val	Ile	Ile	Ser
450						455					460				
Gly	Asn	Lys	Asn	Leu	Cys	Tyr	Ala	Asn	Thr	Ile	Asn	Trp	Lys	Lys	Leu
465					470					475					480
Phe	Gly	Thr	Ser	Gly	Gln	Lys	Thr	Lys	Ile	Ile	Ser	Asn	Arg	Gly	Glu
				485					490					495	
Asn	Ser	Cys	Lys	Ala	Thr	Gly	Gln	Val	Cys	His	Ala	Leu	Cys	Ser	Pro
			500					505					510		
Glu	Gly	Cys	Trp	Gly	Pro	Glu	Pro	Arg	Asp	Cys	Val	Ser	Cys	Arg	Asn
		515						520				525			
Val	Ser	Arg	Gly	Arg	Glu	Cys	Val	Asp	Lys	Cys	Asn	Leu	Leu	Glu	Gly

530					535					540						
Glu 545	Pro	Arg	Glu	Phe	Val 550	Glu	Asn	Ser	Glu	Cys 555	Ile	Gln	Cys	His	Pro 560	
Glu	Cys	Leu	Pro	Gln 565	Ala	Met	Asn	Ile	Thr 570	Cys	Thr	Gly	Arg	Gly	Pro 575	
Asp	Asn	Cys	Ile 580	Gln	Cys	Ala	His	Tyr 585	Ile	Asp	Gly	Pro	His 590	Cys	Val	
Lys	Thr	Cys 595	Pro	Ala	Gly	Val	Met 600	Gly	Glu	Asn	Asn	Thr 605	Leu	Val	Trp	
Lys	Tyr 610	Ala	Asp	Ala	Gly	His 615	Val	Cys	His	Leu	Cys 620	His	Pro	Asn	Cys	
Thr 625	Tyr	Gly	Cys	Thr	Gly 630	Pro	Gly	Leu	Glu	Gly 635	Cys	Pro	Thr	Asn	Gly 640	
Pro	Lys	Ile	Pro	Ser 645	Ile	Ala	Thr	Gly	Met 650	Val	Gly	Ala	Leu	Leu 655	Leu	
Leu	Leu	Val	Val 660	Ala	Leu	Gly	Ile	Gly 665	Leu	Phe	Met	Arg 670	Arg	Arg	His	
Ile	Val	Arg 675	Lys	Arg	Thr	Leu	Arg 680	Arg	Leu	Leu	Gln	Glu 685	Arg	Glu	Leu	
Val	Glu 690	Pro	Leu	Thr	Pro	Ser 695	Gly	Glu	Ala	Pro	Asn 700	Gln	Ala	Leu	Leu	
Arg 705	Ile	Leu	Lys	Glu	Thr 710	Glu	Phe	Lys	Lys	Ile 715	Lys	Val	Leu	Gly	Ser 720	
Gly	Ala	Phe	Gly	Thr 725	Val	Tyr	Lys	Gly	Leu 730	Trp	Ile	Pro	Glu	Gly 735	Glu	
Lys	Val	Lys 740	Ile	Pro	Val	Ala	Ile	Lys 745	Glu	Leu	Arg	Glu	Ala 750	Thr	Ser	
Pro	Lys 755	Ala	Asn	Lys	Glu	Ile	Leu 760	Asp	Glu	Ala	Tyr	Val 765	Met	Ala	Ser	
Val	Asp 770	Asn	Pro	His	Val	Cys 775	Arg	Leu	Leu	Gly 780	Ile	Cys	Leu	Thr	Ser	
Thr 785	Val	Gln	Leu	Ile	Thr 790	Gln	Leu	Met	Pro	Phe 795	Gly	Cys	Leu	Leu	Asp 800	
Tyr	Val	Arg	Glu	His 805	Lys	Asp	Asn	Ile	Gly 810	Ser	Gln	Tyr	Leu	Leu 815	Asn	
Trp	Cys	Val	Gln 820	Ile	Ala	Lys	Gly	Met 825	Asn	Tyr	Leu	Glu	Asp 830	Arg	Arg	
Leu	Val	His 835	Arg	Asp	Leu	Ala	Ala 840	Arg	Asn	Val	Leu	Val 845	Lys	Thr	Pro	
Gln	His 850	Val	Lys	Ile	Thr	Asp 855	Phe	Gly	Leu	Ala	Lys 860	Leu	Leu	Gly	Ala	
Glu 865	Glu	Lys	Glu	Tyr	His 870	Ala	Glu	Gly	Gly	Lys 875	Val	Pro	Ile	Lys	Trp 880	
Met	Ala	Leu	Glu	Ser 885	Ile	Leu	His	Arg	Ile	Tyr	Thr	His	Gln	Ser	Asp 895	
Val	Trp	Ser	Tyr 900	Gly	Val	Thr	Val	Trp 905	Glu	Leu	Met	Thr	Phe	Gly	Ser	
Lys	Pro	Tyr 915	Asp	Gly	Ile	Pro	Ala 920	Ser	Glu	Ile	Ser	Ser 925	Ile	Leu	Glu	
Lys	Gly 930	Glu	Arg	Leu	Pro	Gln 935	Pro	Pro	Ile	Cys	Thr	Ile 940	Asp	Val	Tyr	

-continued

Met Ile Met Val Lys Cys Trp Met Ile Asp Ala Asp Ser Arg Pro Lys
 945 950 955 960

Phe Arg Glu Leu Ile Ile Glu Phe Ser Lys Met Ala Arg Asp Pro Gln
 965 970 975

Arg Tyr Leu Val Ile Gln Gly Asp Glu Arg Met His Leu Pro Ser Pro
 980 985 990

Thr Asp Ser Asn Phe Tyr Arg Ala Leu Met Asp Glu Glu Asp Met Asp
 995 1000 1005

Asp Val Val Asp Ala Asp Glu Tyr Leu Ile Pro Gln Gln Gly Phe
 1010 1015 1020

Phe Ser Ser Pro Ser Thr Ser Arg Thr Pro Leu Leu Ser Ser Leu
 1025 1030 1035

Ser Ala Thr Ser Asn Asn Ser Thr Val Ala Cys Ile Asp Arg Asn
 1040 1045 1050

Gly Leu Gln Ser Cys Pro Ile Lys Glu Asp Ser Phe Leu Gln Arg
 1055 1060 1065

Tyr Ser Ser Asp Pro Thr Gly Ala Leu Thr Glu Asp Ser Ile Asp
 1070 1075 1080

Asp Thr Phe Leu Pro Val Pro Glu Tyr Ile Asn Gln Ser Val Pro
 1085 1090 1095

Lys Arg Pro Ala Gly Ser Val Gln Asn Pro Val Tyr His Asn Gln
 1100 1105 1110

Pro Leu Asn Pro Ala Pro Ser Arg Asp Pro His Tyr Gln Asp Pro
 1115 1120 1125

His Ser Thr Ala Val Gly Asn Pro Glu Tyr Leu Asn Thr Val Gln
 1130 1135 1140

Pro Thr Cys Val Asn Ser Thr Phe Asp Ser Pro Ala His Trp Ala
 1145 1150 1155

Gln Lys Gly Ser His Gln Ile Ser Leu Asp Asn Pro Asp Tyr Gln
 1160 1165 1170

Gln Asp Phe Phe Pro Lys Glu Ala Lys Pro Asn Gly Ile Phe Lys
 1175 1180 1185

Gly Ser Thr Ala Glu Asn Ala Glu Tyr Leu Arg Val Ala Pro Gln
 1190 1195 1200

Ser Ser Glu Phe Ile Gly Ala
 1205 1210

<210> SEQ ID NO 6

<211> LENGTH: 5616

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

```

ccccggcgca gcgcgccgc agcagcctcc gcccccgca cgggtgtgagc gcccgacgcg      60
gccgaggcgg ccggagtccc gagctagccc cgccggccgc cgccgccag accggacgac      120
aggccacctc gtcggcgctc gcccgagtcc ccgcctcgcc gccaacgcca caaccaccgc      180
gcacggcccc ctgactccgt ccagtattga tcgggagagc cggagcgagc tcttcgggga      240
gcagcgatgc gacctccgg gacggccggg gcagcgctcc tggcgctgct ggetgcgctc      300
tgcccgcgca gtcgggctct ggaggaaaag aaagtttgcc aaggcacgag taacaagctc      360
acgcagttgg gcacttttga agatcatttt ctcagcctcc agaggatgtt caataactgt      420

```

-continued

gagggtggtcc	ttgggaat	ttt	ggaaattacc	tatgtgcaga	ggaattatga	tctttccttc	480
ttaaagacca	tccaggaggt	ggctgggtat	gtcctcattg	ccctcaacac	agtgagagca		540
attcctttgg	aaaacctgca	gatcatcaga	ggaaatatgt	actacgaaaa	ttcctatgcc		600
ttagcagtct	tatctaacta	tgatgcaa	aaaaccggac	tgaaggagct	gcccattgaga		660
aatttacagg	aaatcctgca	tggcgccgtg	cggttcagca	acaacctgc	cctgtgcaac		720
gtggagagca	tccagtggcg	ggacatagtc	agcagtgact	ttctcagcaa	catgtcgatg		780
gacttccaga	accacctggg	cagctgccaa	aagtgtgac	caagctgtcc	caatgggagc		840
tgctgggggtg	caggagagga	gaactgccag	aaactgacca	aaatcatctg	tgcccagcag		900
tgctccgggc	gctgccgtgg	caagtcctcc	agtgactgct	gccacaacca	gtgtgctgca		960
ggctgcacag	gcccccgga	gagcgactgc	ctgggtctgcc	gcaaattccg	agacgaagcc		1020
acgtgcaagg	acacctgcc	cccactcatg	ctctacaacc	ccaccacgta	ccagatggat		1080
gtgaaccccg	agggcaata	cagctttggt	gccacctgcg	tgaagaagtg	tccccgta		1140
tatgtggtga	cagatcacgg	ctcgtgcgtc	cgagcctgtg	gggcccagac	ctatgagatg		1200
gaggaagacg	gcgtccgcaa	gtgtaagaag	tgccaagggc	cttgccgcaa	agtgtgtaac		1260
ggaataggta	ttggtgaatt	taaagactca	ctctccataa	atgctacgaa	tattaaacac		1320
ttcaaaaact	gcacctccat	cagtggcgat	ctccacatcc	tgccggtggc	atttaggggt		1380
gactccttca	cacatactcc	tcctctggat	ccacaggaac	tggatattct	gaaaaccgta		1440
aaggaaatca	cagggttttt	gctgattcag	gcttgccctg	aaaacaggac	ggacctccat		1500
gcctttgaga	acctagaaat	catacgccgc	aggaccaagc	aacatggtca	gtttttctct		1560
gcagtcgtca	gcctgaacat	aacatccttg	ggattacgct	ccctcaagga	gataagtgat		1620
ggagatgtga	taatttcagg	aaacaaaaat	ttgtgctatg	caaatacaat	aaactggaaa		1680
aaactgtttg	ggacctccgg	tcagaaaaacc	aaaattataa	gcaacagagg	tgaaaacagc		1740
tgcaaggcca	caggccaggt	ctgccatgcc	ttgtgctccc	ccgagggtcg	ctggggcccg		1800
gagccacagg	actgcgtctc	ttgccggaat	gtcagccgag	gcagggaatg	cgtggacaag		1860
tgcaaccttc	tggagggtga	gccaaggag	tttgtggaga	actctgagtg	catacagtgc		1920
caccagagtg	gcctgcctca	ggccatgaac	atcacctgca	caggacgggg	accagacaac		1980
tgtatccagt	gtgcccacta	cattgacggc	ccccactgcg	tcaagacctg	cccggcagga		2040
gtcatggggag	aaaacaacac	cctgggtctgg	aagtaacgag	acgcccggcca	tgtgtgccac		2100
ctgtgccatc	caaaactgcac	ctacggatgc	actgggcccag	gtcttgaagg	ctgtccaacg		2160
aatgggccta	agatcccgtc	catcgccact	gggatgggtg	gggcccctct	ctgtgctgctg		2220
gtggtggccc	tggggatcgg	cctcttcatg	cgaaggcgcc	acatcgttcg	gaagcgcacg		2280
ctgcggaggc	tgctgcagga	gaggagagctt	gtggagcctc	ttacaccag	tggagaagct		2340
cccaaccaag	ctctcttgag	gatcttgaag	gaaactgaat	tcaaaaagat	caaagtgtctg		2400
ggctccgggtg	cgttcggcac	ggtgtataag	ggactctgga	tcccagaagg	tgagaaaagt		2460
aaaattcccg	tcgtatcaaa	ggaattaaga	gaagcaacat	ctccgaaagc	caacaaggaa		2520
atcctcgatg	aagcctacgt	gatggccagc	gtggacaacc	cccagctgtg	ccgctgctg		2580
ggcatctgcc	tcacctccac	cgtgcagctc	atcacgcagc	tcattgccctt	cggtgcctc		2640
ctggactatg	tccgggaaca	caaagacaat	attggctccc	agtacctgct	caactggtgt		2700

-continued

gtgcagatcg	caaagggcat	gaactacttg	gaggaccgtc	gcttggtgca	ccgcgacctg	2760
gcagccagga	acgtactggt	gaaaacaccg	cagcatgtca	agatcacaga	ttttgggctg	2820
gccaactgc	tgggtgcgga	agagaagaa	taccatgcag	aaggaggcaa	agtgcctatc	2880
aagtggatgg	cattggaatc	aattttacac	agaatctata	cccaccagag	tgatgtctgg	2940
agctacgggg	tgaccgtttg	ggagttgatg	acctttggat	ccaagccata	tgacggaatc	3000
cctgccagcg	agatctcctc	catcctggag	aaaggagaac	gcctccctca	gccacccata	3060
tgtaccatcg	atgtctacat	gatcatggtc	aagtgtctga	tgatagacgc	agatagtctc	3120
ccaaagtcc	gtgagttgat	catcgaatc	tccaaaatgg	cccagagacc	ccagcgctac	3180
cttgtcattc	agggggatga	aagaatgcat	ttgccaagtc	ctacagactc	caactctctac	3240
cgtgccctga	tggatgaaga	agacatggac	gacgtggtgg	atgccgacga	gtacctcatc	3300
ccacagcagg	gcttcttcag	cagccccctc	acgtcacgga	ctccccctct	gagctctctg	3360
agtgaacca	gcaacaatc	caccgtggct	tgcatgtgata	gaaatgggct	gcaaagctgt	3420
cccatcaagg	aagacagctt	cttgacagga	tacagctcag	acccacacag	cgcttgact	3480
gaggacagca	tagacgacac	cttcctccca	gtgcctgaat	acataaacca	gtccgttccc	3540
aaaaggcccg	ctggctctgt	gcagaatcct	gtctatcaca	atcagcctct	gaaccccgcg	3600
cccagcagag	accacacta	ccaggacccc	cacagcactg	cagtgggcaa	ccccgagtat	3660
ctcaacactg	tccagcccac	ctgtgtcaac	agcacattcg	acagccctgc	ccactggggc	3720
cagaaaggca	gccaccaaat	tagcctggac	aacctgact	accagcagga	cttctttccc	3780
aaggaagcca	agccaaatgg	catctttaag	ggctccacag	ctgaaaatgc	agaataccta	3840
agggtcgcgc	cacaaagcag	tgaatttatt	ggagcatgac	cacggaggat	agtatgagcc	3900
ctaaaaatcc	agactctttc	gatacccagg	accaagccac	agcaggctct	ccatcccaac	3960
agccatgccc	gcattagctc	ttagacccac	agactggttt	tgcaacgttt	acaccgacta	4020
gccaggaagt	acttccacct	cgggcacatt	ttgggaagtt	gcattccttt	gtcttcaaac	4080
tgtgaagcat	ttacagaaac	gcattccagca	agaatattgt	ccctttgagc	agaaatttat	4140
ctttcaaaga	ggatatattg	aaaaaaaaaa	aaagtatatg	tgaggatttt	tattgattgg	4200
ggatcttgga	gtttttcatt	gtcgctattg	atttttactt	caatgggctc	ttccaacaag	4260
gaagaagctt	gctggtagca	cttgctaccc	tgagttcatc	caggcccaac	tgtgagcaag	4320
gagcacaagc	cacaagtctt	ccagaggatg	cttgattcca	gtggttctgc	ttcaaggctt	4380
ccactgcaaa	acactaaaga	tccaagaagg	ccttcatggc	cccagcaggc	cgatcggtta	4440
ctgtatcaag	tcatggcagg	tacagtagga	taagccactc	tgtcccttcc	tgggcaaaga	4500
agaaacggag	gggatggaat	tcttccttag	acttactttt	gtaaaaatgt	ccccacggta	4560
cttactcccc	actgatggac	cagtggtttc	cagtcatgag	cgtagactg	acttgtttgt	4620
cttccattcc	attgttttga	aactcagtat	gctgcccctg	tcttgctgtc	atgaaatcag	4680
caagagagga	tgacacatca	aataataact	cggattccag	cccacattgg	attcatcagc	4740
atttggacca	atagcccaca	gctgagaatg	tggaaatcct	aaggatagca	ccgcttttgt	4800
tctcgcaaaa	acgtatctcc	taatttgagg	ctcagatgaa	atgcatcagg	tcctttgggg	4860
catagatcag	aagactacaa	aatgaagct	gctctgaaat	ctccttttagc	catcacccca	4920
accccccaaa	attagtttgt	gttacttatg	gaagatagtt	ttctcctttt	acttcacttc	4980

-continued

```

aaaagctttt tactcaaaga gtatatgttc cctccaggtc agctgcccc aaacccctc 5040
cttacgcttt gtcacacaaa aagtgtctct gccttgagtc atctattcaa gcacttacag 5100
ctctggccac aacagggcat ttacagggtg cgaatgacag tagcattatg agtagtggtg 5160
aattcaggta gtaaatatga aactagggtt tgaaattgat aatgctttca caacatttgc 5220
agatgtttta gaaggaaaaa agttccttcc taaaataatt tctctacaat tggaagattg 5280
gaagattcag ctagttagga gccacacctt ttctctaata tgtgtgtgcc ctgtaacctg 5340
actggttaac agcagtcctt tgtaaacagt gttttaaact ctctagtca atatccacc 5400
catccaattt atcaaggaag aaatggttca gaaaatattt tcagcctaca gttatgttca 5460
gtcacacaca catacaaaat gttccttttg cttttaaagt aatttttgac tcccagatca 5520
gtcagagccc ctacagcatt gttaagaaag tatttgattt ttgtctcaat gaaaataaaa 5580
ctatattcat ttccactcta aaaaaaaaaa aaaaaa 5616

```

<210> SEQ ID NO 7

<211> LENGTH: 766

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

Met Ala Ala Leu Ser Gly Gly Gly Gly Gly Ala Glu Pro Gly Gln
1           5           10          15
Ala Leu Phe Asn Gly Asp Met Glu Pro Glu Ala Gly Ala Gly Ala Gly
20          25          30
Ala Ala Ala Ser Ser Ala Ala Asp Pro Ala Ile Pro Glu Glu Val Trp
35          40          45
Asn Ile Lys Gln Met Ile Lys Leu Thr Gln Glu His Ile Glu Ala Leu
50          55          60
Leu Asp Lys Phe Gly Gly Glu His Asn Pro Pro Ser Ile Tyr Leu Glu
65          70          75          80
Ala Tyr Glu Glu Tyr Thr Ser Lys Leu Asp Ala Leu Gln Gln Arg Glu
85          90          95
Gln Gln Leu Leu Glu Ser Leu Gly Asn Gly Thr Asp Phe Ser Val Ser
100         105         110
Ser Ser Ala Ser Met Asp Thr Val Thr Ser Ser Ser Ser Ser Ser Leu
115        120        125
Ser Val Leu Pro Ser Ser Leu Ser Val Phe Gln Asn Pro Thr Asp Val
130        135        140
Ala Arg Ser Asn Pro Lys Ser Pro Gln Lys Pro Ile Val Arg Val Phe
145        150        155        160
Leu Pro Asn Lys Gln Arg Thr Val Val Pro Ala Arg Cys Gly Val Thr
165        170        175
Val Arg Asp Ser Leu Lys Lys Ala Leu Met Met Arg Gly Leu Ile Pro
180        185        190
Glu Cys Cys Ala Val Tyr Arg Ile Gln Asp Gly Glu Lys Lys Pro Ile
195        200        205
Gly Trp Asp Thr Asp Ile Ser Trp Leu Thr Gly Glu Glu Leu His Val
210        215        220
Glu Val Leu Glu Asn Val Pro Leu Thr Thr His Asn Phe Val Arg Lys
225        230        235        240

```

-continued

Thr	Phe	Phe	Thr	Leu	Ala	Phe	Cys	Asp	Phe	Cys	Arg	Lys	Leu	Leu	Phe
				245					250				255		
Gln	Gly	Phe	Arg	Cys	Gln	Thr	Cys	Gly	Tyr	Lys	Phe	His	Gln	Arg	Cys
		260						265					270		
Ser	Thr	Glu	Val	Pro	Leu	Met	Cys	Val	Asn	Tyr	Asp	Gln	Leu	Asp	Leu
		275					280					285			
Leu	Phe	Val	Ser	Lys	Phe	Phe	Glu	His	His	Pro	Ile	Pro	Gln	Glu	Glu
	290				295						300				
Ala	Ser	Leu	Ala	Glu	Thr	Ala	Leu	Thr	Ser	Gly	Ser	Ser	Pro	Ser	Ala
305					310					315					320
Pro	Ala	Ser	Asp	Ser	Ile	Gly	Pro	Gln	Ile	Leu	Thr	Ser	Pro	Ser	Pro
			325						330					335	
Ser	Lys	Ser	Ile	Pro	Ile	Pro	Gln	Pro	Phe	Arg	Pro	Ala	Asp	Glu	Asp
		340						345					350		
His	Arg	Asn	Gln	Phe	Gly	Gln	Arg	Asp	Arg	Ser	Ser	Ser	Ala	Pro	Asn
		355					360					365			
Val	His	Ile	Asn	Thr	Ile	Glu	Pro	Val	Asn	Ile	Asp	Asp	Leu	Ile	Arg
	370					375					380				
Asp	Gln	Gly	Phe	Arg	Gly	Asp	Gly	Gly	Ser	Thr	Thr	Gly	Leu	Ser	Ala
385					390					395					400
Thr	Pro	Pro	Ala	Ser	Leu	Pro	Gly	Ser	Leu	Thr	Asn	Val	Lys	Ala	Leu
			405						410					415	
Gln	Lys	Ser	Pro	Gly	Pro	Gln	Arg	Glu	Arg	Lys	Ser	Ser	Ser	Ser	Ser
			420					425					430		
Glu	Asp	Arg	Asn	Arg	Met	Lys	Thr	Leu	Gly	Arg	Arg	Asp	Ser	Ser	Asp
		435					440					445			
Asp	Trp	Glu	Ile	Pro	Asp	Gly	Gln	Ile	Thr	Val	Gly	Gln	Arg	Ile	Gly
450					455						460				
Ser	Gly	Ser	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Lys	Trp	His	Gly	Asp	Val
465				470						475					480
Ala	Val	Lys	Met	Leu	Asn	Val	Thr	Ala	Pro	Thr	Pro	Gln	Gln	Leu	Gln
			485					490						495	
Ala	Phe	Lys	Asn	Glu	Val	Gly	Val	Leu	Arg	Lys	Thr	Arg	His	Val	Asn
		500						505					510		
Ile	Leu	Leu	Phe	Met	Gly	Tyr	Ser	Thr	Lys	Pro	Gln	Leu	Ala	Ile	Val
	515						520					525			
Thr	Gln	Trp	Cys	Glu	Gly	Ser	Ser	Leu	Tyr	His	His	Leu	His	Ile	Ile
	530					535					540				
Glu	Thr	Lys	Phe	Glu	Met	Ile	Lys	Leu	Ile	Asp	Ile	Ala	Arg	Gln	Thr
545					550					555					560
Ala	Gln	Gly	Met	Asp	Tyr	Leu	His	Ala	Lys	Ser	Ile	Ile	His	Arg	Asp
			565					570						575	
Leu	Lys	Ser	Asn	Asn	Ile	Phe	Leu	His	Glu	Asp	Leu	Thr	Val	Lys	Ile
			580					585					590		
Gly	Asp	Phe	Gly	Leu	Ala	Thr	Val	Lys	Ser	Arg	Trp	Ser	Gly	Ser	His
		595					600					605			
Gln	Phe	Glu	Gln	Leu	Ser	Gly	Ser	Ile	Leu	Trp	Met	Ala	Pro	Glu	Val
	610					615					620				
Ile	Arg	Met	Gln	Asp	Lys	Asn	Pro	Tyr	Ser	Phe	Gln	Ser	Asp	Val	Tyr
625					630					635					640
Ala	Phe	Gly	Ile	Val	Leu	Tyr	Glu	Leu	Met	Thr	Gly	Gln	Leu	Pro	Tyr

-continued

645					650					655					
Ser	Asn	Ile	Asn	Asn	Arg	Asp	Gln	Ile	Ile	Phe	Met	Val	Gly	Arg	Gly
	660						665						670		
Tyr	Leu	Ser	Pro	Asp	Leu	Ser	Lys	Val	Arg	Ser	Asn	Cys	Pro	Lys	Ala
	675						680					685			
Met	Lys	Arg	Leu	Met	Ala	Glu	Cys	Leu	Lys	Lys	Lys	Arg	Asp	Glu	Arg
	690					695					700				
Pro	Leu	Phe	Pro	Gln	Ile	Leu	Ala	Ser	Ile	Glu	Leu	Leu	Ala	Arg	Ser
	705				710					715					720
Leu	Pro	Lys	Ile	His	Arg	Ser	Ala	Ser	Glu	Pro	Ser	Leu	Asn	Arg	Ala
			725							730				735	
Gly	Phe	Gln	Thr	Glu	Asp	Phe	Ser	Leu	Tyr	Ala	Cys	Ala	Ser	Pro	Lys
			740						745				750		
Thr	Pro	Ile	Gln	Ala	Gly	Gly	Tyr	Gly	Ala	Phe	Pro	Val	His		
		755					760					765			

<210> SEQ ID NO 8

<211> LENGTH: 2949

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```

cgctccctt cccctcccc gcccgacagc gcccgctcgg gcccggtc tcggttataa    60
gatggcggcg ctgagcgggtg gcggtggtgg cggcgcgagg ccgggccagg ctctgttcaa    120
cggggacatg gagcccgagg ccggcgccgg gcccgggccc gcggcctctt cggctgcgga    180
ccctgccatt ccggaggagg tgtggaatat caaacaaatg attaagtga cacaggaaca    240
tatagaggcc ctattggaca aatttggtgg ggagcataat ccaccatcaa tatatctgga    300
ggcctatgaa gaatacacca gcaagctaga tgcaactcaa caaagagaac aacagttatt    360
ggaatctctg gggaaacggaa ctgatttttc tgtttctagc tctgcatcaa tggataaccgt    420
tacatcttct tcctcttcta gcctttcagt gctacettca tctctttcag tttttcaaaa    480
tcccacagat gtggcacgga gcaaccccaa gtcaccacaa aaacctatcg ttagagtctt    540
cctgcccac aaacagagga cagtgggtacc tgcaagggtg ggagttacag tccgagacag    600
tctaaagaaa gcactgatga tgagaggtct aatcccagag tgctgtgctg ttacagaat    660
tcaggatgga gagaagaac caattggtg ggacactgat atttctggc ttactggaga    720
agaattgcat gtggaagtgt tggagaatgt tccacttaca acacacaact ttgtacgaaa    780
aacgtttttc accttagcat tttgtgactt ttgtcgaaag ctgcttttcc agggtttccg    840
ctgtcaaaca tgtggttata aatttcacca gcgttgtagt acagaagttc cactgatgtg    900
tgtaattat gaccaacttg atttctgtt tgtctccaag ttctttgaac accacccaat    960
accacaggaa gaggcgtcct tagcagagac tgccctaaca tctggatcat ccccttcgcg    1020
acccgcctcg gactctattg ggcccaaat tctcaccagt ccgtctcctt caaaatccat    1080
tccaattcca cagcccttcc gaccagcaga tgaagatcat cgaaatcaat ttgggcaacg    1140
agaccgatcc tcatcagctc ccaatgtgca tataaacaca atagaacctg tcaatattga    1200
tgacttgatt agagaccaag gatttcgtgg tgatggagga tcaaccacag gtttgtctgc    1260
tccccccct gcctcattac ctggctcact aactaacgtg aaagccttac agaaatctcc    1320
aggacctcag cgagaaagga agtcatcttc atcctcagaa gacaggaatc gaatgaaaac    1380

```


-continued

acttggtaga	cgggactoga	gtgatgattg	ggagattcct	gatgggcaga	ttacagtggg	1440
acaaagaatt	ggatctggat	catttggaac	agtctacaag	ggaaagtggc	atggatgatgt	1500
ggcagtga	atgttgatg	tgacagcacc	tacacctcag	cagttacaag	ccttcaaaaa	1560
tgaagtagga	gtactcagga	aaacacgaca	tgtgaatata	ctactcttca	tgggctattc	1620
cacaaagcca	caactggcta	ttgttaccca	gtggtgtgag	ggctccagct	tgtatcacca	1680
tctccatata	attgagacca	aatttgagat	gatcaaactt	atagatatgt	cacgacagac	1740
tgcacagggc	atggattact	tacacgcca	gtcaatcata	cacagagacc	tcaagagtaa	1800
taatatat	cttcatgaag	acctcacagt	aaaaataggt	gattttgggt	tagctacagt	1860
gaaatctcga	tggagtgggt	cccatcagtt	tgaacagttg	tctggatcca	ttttgtggat	1920
ggcaccagaa	gtcatcagaa	tgaagataa	aatccatac	agctttcagt	cagatgtata	1980
tgcatttga	attgttctgt	atgaattgat	gactggacag	ttaccttatt	caaacatcaa	2040
caacagggac	cagataat	ttatggtggg	acgaggatag	ctgtctccag	atctcagtaa	2100
ggtacggagt	aactgtccaa	aagccatgaa	gagattaatg	gcagagtgc	tcaaaaagaa	2160
aagagatgag	agaccactct	ttccccaaat	tctcgctct	attgagctgc	tggcccgctc	2220
attgccaaaa	attcacccga	gtgcatcaga	acctccttg	aatcgggctg	gtttccaaac	2280
agaggatttt	agtctatatg	cttgtgcttc	tccaaaaaca	cccatccagg	cagggggata	2340
tgggtgcgtt	cctgtccact	gaaacaaatg	agtgagagag	ttcaggagag	tagcaacaaa	2400
aggaaaaata	atgaacatat	gtttgcttat	atgttaaatt	gaataaaata	ctctcttttt	2460
ttttaaggtg	aaccaaagaa	cacttgtgtg	gttaaagact	agatataatt	tttcccaaaa	2520
ctaaaattta	tacttaacat	tggattttta	acatccaagg	gttaaaatac	atagacattg	2580
ctaaaatttg	gcagagcctc	ttctagaggc	ttactttct	gttcggggtt	tgtatcattc	2640
acttggttat	tttaagtagt	aaacttcagt	ttctcatgca	actttgtgtg	ccagctatca	2700
catgtccact	agggactcca	gaagaagacc	ctacctatgc	ctgtgtttgc	aggtgagaag	2760
ttggcagtcg	gttagcctgg	gttagataag	gcaaactgaa	cagatcta	ttagggaagtc	2820
agtagaattt	aataattcta	ttattattct	taataatttt	tctataacta	tttcttttta	2880
taacaatttg	gaaaatgtgg	atgtctttta	tttccttgaa	gcaataaact	aagtttcttt	2940
ttataaaaa						2949

<210> SEQ ID NO 9

<211> LENGTH: 2708

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

ggcggcgctt	gattgggctg	ggggggccaa	ataaaagcga	tggcgattgg	gctgccgcgt	60
ttggcgctcg	gtccggctgc	gtccgacacc	cgggtgggact	cagaaggcag	tggagccccg	120
gcggcgccgg	cggcgccgcg	cgggggcgac	gcgcgggaac	aacgcgagtc	ggcgcgccgg	180
acgaagaata	atcatgggcc	agactgggaa	gaaatctgag	aagggaaccag	ttgtgtggcg	240
gaagcgtgta	aatcagagtc	acatgcgact	gagacagctc	aagaggttca	gacgagctga	300
tgaagtaaag	agtatgttta	gttccaatcg	tcagaaaatt	ttggaaagaa	cggaaatctt	360
aaaccaagaa	tggaaacagc	gaaggatata	gcctgtgcac	atcctgactt	ctgtgagctc	420

-continued

attgcgcggg	actagggagt	gttcggtgac	cagtgacttg	gattttccaa	cacaagtcac	480
cccattaaag	actctgaatg	cagttgcttc	agtaccata	atgtattctt	ggtctcccct	540
acagcagaat	tttatggtgg	aagatgaaac	tgttttacac	aacattcctt	atatgggaga	600
tgaagtttta	gatcaggatg	gtactttcat	tgaagaacta	ataaaaaatt	atgatgggaa	660
agtacacggg	gatagagaat	gtgggtttat	aatgatgaa	atttttgttg	agttggtgaa	720
tgcccttggt	caatataatg	atgatgacga	tgatgatgat	ggagacgac	ctgaagaaag	780
agaagaaaag	cagaaagatc	tggaggatca	cagagatgat	aaagaaagcc	gcccacctcg	840
gaaatttcct	tctgataaaa	tttttgaagc	catttctca	atgtttccag	ataagggcac	900
agcagaagaa	ctaagggaaa	aatataaaga	actcaccgaa	cagcagctcc	caggcgccact	960
tcctcctgaa	tgtaccccca	acatagatgg	accaaagtct	aaatctgttc	agagagagca	1020
aagcttacac	tcctttcata	cgcttttctg	taggcgatgt	tttaaatatg	actgcttcct	1080
acatcctttt	catgcaacac	ccaacactta	taagcggaag	aacacagaaa	cagctctaga	1140
caacaaacct	tgtggaccac	agtgttacca	gcatttgag	ggagcaaagg	agtttgctgc	1200
tgctctcacc	gctgagcgga	taaagacccc	accaaaacgt	ccaggaggcc	gcagaagagg	1260
acggcttccc	aataacagta	gcaggcccag	cacccccacc	attaatgtgc	tggaatcaaa	1320
ggatacacag	agtgatagg	aagcaggac	tgaacgggg	ggagagaaca	atgataaaga	1380
agaagaagag	aagaaagatg	aaacttcgag	ctcctctgaa	gcaaattctc	ggtgtcaaac	1440
accaataaag	atgaagccaa	atattgaacc	tcctgagaat	gtggagtgga	gtggtgctga	1500
agcctcaatg	tttagagtcc	tcattggcac	ttactatgac	aatttctgtg	ccattgctag	1560
gttaattggg	acaaaacat	gtagacaggt	gtatgagttt	agagtcaaag	aatctagcat	1620
catagctcca	gctcccgtg	aggatgtgga	tactcctcca	aggaaaaaga	agaggaaaca	1680
cgggttggtg	gctgcacact	gcagaaagat	acagctgaaa	aaggacggct	cctctaacca	1740
tgtttacaac	tatcaacct	gtgatcatcc	acggcagcct	tgtgacagtt	cgtgcccttg	1800
tgtgatagca	caaaattttt	gtgaaaagtt	tgtcaatgt	agttcagagt	gtcaaaaccg	1860
ctttccggga	tgccgctgca	aagcacagt	caacaccaag	cagtgcccg	gctacctggc	1920
tgtccgagag	tgtgaccctg	acctctgtct	tacttggtga	gccgctgacc	attgggacag	1980
taaaaatgtg	tcctgcaaga	actgcagtat	tcagcggggc	tccaaaaagc	atctattgct	2040
ggcaccatct	gacgtggcag	gctgggggat	ttttatcaaa	gatcctgtgc	agaaaaatga	2100
attcatctca	gaatactgtg	gagagattat	ttctcaagat	gaagctgaca	gaagagggaa	2160
agtgtatgat	aaatacatgt	gcagctttct	gttcaacttg	aacaatgatt	ttgtggtgga	2220
tgcaaccgcg	aagggtaaca	aaattcggtt	tgcaaatcat	tcggtaaatc	caaaactgcta	2280
tgcaaaagtt	atgatggtta	acggtgatca	caggataggt	atttttgcca	agagagccat	2340
ccagactggc	gaagagctgt	tttttgatta	cagatcacag	caggctgatg	ccctgaagta	2400
tgtcggcatc	gaaagagaaa	tggaaatccc	ttgacatctg	ctacctctc	ccccctctc	2460
tgaacacagct	gccttagctt	caggaaacct	gagtactgtg	ggcaatttag	aaaaagaaca	2520
tgcagtttga	aattctgaat	ttgcaaagta	ctgtaagaat	aatttatagt	aatgagttaa	2580
aaaatcaact	ttttattgcc	ttctcaccag	ctgcaaagtg	ttttgtacca	gtgaattttt	2640
gcaataatgc	agtatggtac	atttttcaac	tttgaataaa	gaatacttga	acttgcctct	2700

-continued

gttgaatc

2708

<210> SEQ ID NO 10

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
 1 5 10 15
 Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
 20 25 30
 Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
 35 40 45
 Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
 50 55 60
 Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
 65 70 75 80
 Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
 85 90 95
 Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
 100 105 110
 Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
 115 120 125
 His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
 130 135 140
 Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp
 145 150 155 160
 Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn
 165 170 175
 Ala Leu Gly Gln Tyr Asn Asp Asp Asp Asp Asp Asp Gly Asp Asp
 180 185 190
 Pro Glu Glu Arg Glu Glu Lys Gln Lys Asp Leu Glu Asp His Arg Asp
 195 200 205
 Asp Lys Glu Ser Arg Pro Pro Arg Lys Phe Pro Ser Asp Lys Ile Phe
 210 215 220
 Glu Ala Ile Ser Ser Met Phe Pro Asp Lys Gly Thr Ala Glu Glu Leu
 225 230 235 240
 Lys Glu Lys Tyr Lys Glu Leu Thr Glu Gln Gln Leu Pro Gly Ala Leu
 245 250 255
 Pro Pro Glu Cys Thr Pro Asn Ile Asp Gly Pro Asn Ala Lys Ser Val
 260 265 270
 Gln Arg Glu Gln Ser Leu His Ser Phe His Thr Leu Phe Cys Arg Arg
 275 280 285
 Cys Phe Lys Tyr Asp Cys Phe Leu His Pro Phe His Ala Thr Pro Asn
 290 295 300
 Thr Tyr Lys Arg Lys Asn Thr Glu Thr Ala Leu Asp Asn Lys Pro Cys
 305 310 315 320
 Gly Pro Gln Cys Tyr Gln His Leu Glu Gly Ala Lys Glu Phe Ala Ala
 325 330 335
 Ala Leu Thr Ala Glu Arg Ile Lys Thr Pro Pro Lys Arg Pro Gly Gly
 340 345 350

Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr	Pro
		355					360					365			
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu	Ala
	370					375					380				
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Glu	Lys
385					390					395					400
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln	Thr
				405					410					415	
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu	Trp
			420					425					430		
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr	Tyr
		435					440					445			
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys	Arg
	450					455					460				
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro	Ala
465					470					475					480
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys	His
				485					490					495	
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp	Gly
			500					505					510		
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg	Gln
		515					520					525			
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys	Glu
	530					535					540				
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly	Cys
545					550					555					560
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala
			565						570					575	
Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp
			580					585					590		
His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg
		595					600					605			
Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp
	610					615					620				
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625					630					635					640
Tyr	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
			645						650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Arg	Phe	Ala	Asn
		675					680					685			
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
	690					695					700				
Asp	His	Arg	Ile	Gly</											

-continued

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 11
agcgatgacg tctctgaggt 20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 12
gtacagggac accagccact 20

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 13
taacaagctc acgcagttgg 20

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 14
gttgaggcca atgaggacat 20

<210> SEQ ID NO 15
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 15
gcccgaagc tgaccctgaa gttcattcaa gagatgaact tcagggtcag cttgctttt 59

<210> SEQ ID NO 16
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 16

-continued

ccggcggaaa tcttaacca agaattcga gattcttgg ttaagatttc cgttttt 57

<210> SEQ ID NO 17
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 17

ccgggctgag aatgtggaat acctactga gtaggtattc cacattctca gcttttt 57

<210> SEQ ID NO 18
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 18

tttgtgata agcagcacia gatttcaaga gaattctctg ctgcttctcc attttt 56

<210> SEQ ID NO 19
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 19

tcatctgcac tgccaagact g 21

<210> SEQ ID NO 20
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 20

catgccttct ttcactttgc c 21

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 21

aggagtttgc tgctgctctc 20

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 22

ccgagaattt gcttcagagg 20

<210> SEQ ID NO 23

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 23

tacggctcct attgcaaac 20

<210> SEQ ID NO 24

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 24

tgcttcagtt tgttgcttg 20

<210> SEQ ID NO 25

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 25

ggcctctgat ctccgatttc 20

<210> SEQ ID NO 26

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 26

tgggctctaa attggctcac 20

<210> SEQ ID NO 27

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 27

atgaaattca cccctttcc 20

<210> SEQ ID NO 28

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

primer

<400> SEQUENCE: 28

ccctaggctg tgctcacttc 20

<210> SEQ ID NO 29

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 29

ccggctaact ctgaggacac 20

<210> SEQ ID NO 30

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 30

cgagctgttt acgtttgacg 20

<210> SEQ ID NO 31

<211> LENGTH: 4

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: Description of Unknown: "DEAD" box motif peptide

<400> SEQUENCE: 31

Asp Glu Ala Asp
1

<210> SEQ ID NO 32

<211> LENGTH: 57

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 32

ccggtattgc cttctcacca gctgcctcga ggcagctggg gagaaggcaa tattttt 57

<210> SEQ ID NO 33

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 33

tcgagccata aaaggcaact 20

<210> SEQ ID NO 34

<211> LENGTH: 20

<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 34

tctccctcct cctcttcctc                                20

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 35

ccttgtagat tggggctgag                                20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 36

agtttggggg tggaagaaag                                20

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 37

ggctgagaca gaggaacac                                20

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 38

ccatctcagc ctccaagta                                20

```

What is claimed herein is:

1. A method of treating cancer, the method comprising: administering or prescribing a topoisomerase I inhibitor and an EZH2 inhibitor to a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell.
2. The method of claim 1, wherein the EZH2 inhibitor is selected from the group consisting of: an inhibitory nucleic acid; DZNep; and S-adenosyl-L-homocysteine.
3. The method of claim 1, wherein the topoisomerase I inhibitor is selected from the group consisting of:

camptothecins; topotecan; irinotecan; indenoisoquinolines; indotecan; and indimitecan; and lamellarin D.

4. The method of claim 1, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:

a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272* wherein the * denotes truncation.

5. The method of claim 1, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:

a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R.

6. The method of claim 1, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:

G496A and L597V.

7. The method of claim 1, wherein the presence of the mutation is determined using an assay selected from the group consisting of:

hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.

8. The method of claim 1, wherein the mutation is present in the genomic DNA or in the mRNA transcripts of the tumor cell.

9. The method of claim 1, wherein the cancer is selected from the group consisting of:

lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.

10. The method of claim 1, further comprising the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.

11. A method of treating cancer, the method comprising: administering or prescribing a EGFR inhibitor and an EZH2 inhibitor to a subject determined to have a sensitizing mutation of BRG1, EGFR, or B-RAF in a tumor cell.

12. The method of claim 11, wherein the EZH2 inhibitor is selected from the group consisting of:

an inhibitory nucleic acid; DZNep; and S-adenosyl-L-homocysteine.

13. The method of claim 11, wherein the EGFR inhibitor is selected from the group consisting of:

erlotinib; gefitinib; lapatinib; afatinib; CI-1033; neratinib; CP-724714; TAK-285; AST-1306; ARRY334543; Arry-380; AG-1478; dacomitinib; desmethyl erlotinib;

OSI-420; AZD8931; AEE788; pelitinib; CUDC-101; WS8040; WZ4002; WZ3146; AG-490; XL647; PD153035 HC1; BMS-599626; AG 825; AG 494; AG 555; BIBU 1361; BIBX 1382; DAPH; JNJ 28871063; lavendustin A; and hypericin.

14. The method of claim 11, wherein the sensitizing mutation of BRG1 comprises a mutation selected from the group consisting of:

a mutation which inactivates the ATPase activity of BRG1; a mutation which decreases the expression of BRG1; P270*; Q729*; W764R; T58*; and M272* wherein the * denotes truncation.

15. The method of claim 11, wherein the sensitizing mutation of EGFR comprises a mutation selected from the group consisting of:

a mutation which increased the expression level of EGFR; E746_A750del; E746_A749del; T790M; and L858R.

16. The method of claim 11, wherein the sensitizing mutation of B-RAF comprises a mutation selected from the group consisting of:

G496A and L597V.

17. The method of claim 11, wherein the presence of the mutation is determined using an assay selected from the group consisting of:

hybridization; sequencing; exome capture; PCR; and high-throughput sequencing.

18. The method of claim 11, wherein the mutation is present in the genomic DNA or mRNA transcripts of the tumor cell.

19. The method of claim 11, wherein the cancer is selected from the group consisting of:

lung cancer; non-small cell lung cancer; ovarian cancer; EGFR-expressing ovarian cancer; B-Raf V600E melanomas; breast cancer; colon cancer; EGFR-mutated cancers; EGFR-mutated breast cancers; and EGFR-mutated colon cancers.

20. The method of claim 11, further comprising the step of generating a report based on the detection of a sensitizing mutation in BRG1, EGFR, or B-RAF by a non-human machine.

* * * * *