



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/08/16
(87) Date publication PCT/PCT Publication Date: 2020/02/20
(85) Entrée phase nationale/National Entry: 2021/02/04
(86) N° demande PCT/PCT Application No.: US 2019/046726
(87) N° publication PCT/PCT Publication No.: 2020/037164
(30) Priorité/Priority: 2018/08/17 (US62/765,090)

(51) Cl.Int./Int.Cl. *C07D 213/75* (2006.01),
A61K 31/44 (2006.01), *A61K 31/496* (2006.01),
A61P 9/04 (2006.01), *C07C 309/04* (2006.01),
C07C 309/08 (2006.01), *C07C 309/35* (2006.01),
C07C 55/06 (2006.01), *C07C 55/08* (2006.01),
C07C 55/10 (2006.01), *C07C 57/145* (2006.01),
C07C 57/15 (2006.01), ...

(71) Demandeurs/Applicants:
AMGEN INC., US;
CYTOKINETICS, INC., US

(72) Inventeurs/Inventors:
MORRISON, HENRY, US;
CUI, SHENG, US;

(54) Titre : SEL ET FORMES CRISTALLINES D'OMECAMTIV MECARBIL

(54) Title: SALTS AND CRYSTAL FORMS OF OMECAMTIV MECARBIL

(57) Abrégé/Abstract:

Provided herein are free base crystalline forms, crystalline salts, and an amorphous salts of omeamtiv mecarbیل.

(51) Cl.Int./Int.Cl. (suite/continued) *C07C 59/255* (2006.01), *C07C 65/10* (2006.01), *C07D 213/80* (2006.01)

(72) **Inventeurs(suite)/Inventors(continued)**: QUASDORF, KYLE, US; YANEZ, EVELYN, US; QUAN, BIN PETER, US; KELLY, RON C., US; CAILLE, SEBASTIEN, US; XIAO, LINGYUN, US; PEARSON, DAVID, GB; RAEBURN, JACLYN, GB; GOLIGHTLY, AILSA, GB; MORGAN, BRADLEY, US; PETERSON, MATTHEW, US; EBERLIN, ALEX, US; OBLEZOV, ALEXANDER, US

(74) **Agent**: SMART & BIGGAR LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

20 February 2020 (20.02.2020)



(10) International Publication Number

WO 2020/037164 A1

(51) International Patent Classification:

A61P 9/00 (2006.01) C07C 57/145 (2006.01)
 C07D 213/75 (2006.01) C07C 57/15 (2006.01)
 A61K 31/496 (2006.01) C07C 59/255 (2006.01)
 C07C 55/06 (2006.01) C07C 309/04 (2006.01)
 C07C 55/08 (2006.01) C07C 309/08 (2006.01)
 C07C 55/10 (2006.01) C07C 309/35 (2006.01)

(21) International Application Number:

PCT/US2019/046726

(22) International Filing Date:

16 August 2019 (16.08.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/765,090 17 August 2018 (17.08.2018) US

(71) Applicants: AMGEN INC. [US/US]; Law Department-patent Operations, One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). CYTOKINETICS, INC. [US/US]; 280 E. Grand Ave., South San Francisco, CA 94080 (US).

(72) Inventors: MORRISON, Henry; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). CUI, Sheng; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). QUASDORF, Kyle; One Amgen Drive Drive, Thousand Oaks, CA 91320-1799 (US). YANEZ, Evelyn; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). QUAN, Bin, Peter; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). KELLY, Ron, C.; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). CAILLE, Sebastien; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). XIAO, Lingyun; One Amgen Center Drive, Thousand Oakd, CA 91320-1799 (US). PEARSON, David; 19 Edinburgh Road, Meadowburn, Berwickshire TD2 6TW (GB). RAEBURN, Jacylyn; 1 The Fleming Building, Edinburgh Technopole, Milton Bridge EH26 0BE (GB). GOLIGHTLY, Ailsa; 26 Sanderson's Grove, Tranent EH33 1JY (GB). MORGAN, Bradley; 5948 Monzal Avenue, Oakland, CA 94611 (US). PETERSON, Matthew; 4939 Sandshore Court, San Diego, CA 92130 (US). EBERLIN, Alex; 280 East Grand Avenue, South San Francisco, CA 94080 (US). OBLEZOV, Alexander; 280 East Grand Avenue, South San Francisco, CA 94080 (US).

(74) Agent: DANEK, Shelley, C.; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, 6300 Willis Tower, Chicago, IL 60606-6357 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: SALTS AND CRYSTAL FORMS OF OMECAMTIV MECARBIL

(57) Abstract: Provided herein are free base crystalline forms, crystalline salts, and an amorphous salts of omecamtiv mecarbil.



WO 2020/037164 A1

SALTS AND CRYSTAL FORMS OF OMECAMPV MECARBIL**TECHNICAL FIELD OF INVENTION**


[0001] The present disclosure relates to salts and crystal forms of omecamtiv mecarbil, pharmaceutical compositions thereof and methods of using the same.

BACKGROUND

[0002] The cardiac sarcomere is the basic unit of muscle contraction in the heart. The cardiac sarcomere is a highly ordered cytoskeletal structure composed of cardiac muscle myosin, actin and a set of regulatory proteins. The discovery and development of small molecule cardiac muscle myosin activators would lead to promising treatments for acute and chronic heart failure and dilated cardiomyopathy (DCM) and conditions associated with left and/or right ventricular systolic dysfunction or systolic reserve. Cardiac muscle myosin is the cytoskeletal motor protein in the cardiac muscle cell. It is directly responsible for converting chemical energy into the mechanical force, resulting in cardiac muscle contraction.

[0003] Current positive inotropic agents, such as beta-adrenergic receptor agonists or inhibitors of phosphodiesterase activity, increase the concentration of intracellular calcium, thereby increasing cardiac sarcomere contractility. However, the increase in calcium levels increase the velocity of cardiac muscle contraction and shortens systolic ejection time, which has been linked to potentially life-threatening side effects. In contrast, cardiac muscle myosin activators work by a mechanism that directly stimulates the activity of the cardiac muscle myosin motor protein, without increasing the intracellular calcium concentration. They accelerate the rate-limiting step of the myosin enzymatic cycle and shift it in favor of the force-producing state. Rather than increasing the velocity of cardiac contraction, this mechanism instead lengthens the systolic ejection time, which results in increased cardiac muscle contractility and cardiac output in a potentially more oxygen-efficient manner.

[0004] Omecamtiv mecarbil is a first in class direct activator of cardiac myosin, the motor protein that causes cardiac contraction. It is being evaluated as a potential treatment of heart failure in both intravenous and oral formulations with the goal of establishing a new continuum of care for patients in both the in-hospital and outpatient settings. Omecamtiv



mecarbil has a structure of

(see, e.g., US

Patent No. 7,507,735), and has alternatively been referred to as methyl 4-(2-fluoro-3-(3-(6-methylpyridin-3-yl)ureido)benzyl)piperazine-1-carboxylate, AMG 423, and CK 1827452.

[0005] There is a need for various new salt and crystalline forms of omecamtiv mecarbil with different chemical and physical stabilities, and formulations and uses of the same.

SUMMARY

[0006] Provided herein are salts and crystal forms of omecamtiv mecarbil, including free base crystalline forms, crystalline salts and amorphous salt forms of omecamtiv mecarbil. In some embodiments, provided herein is the omecamtiv mecarbil free base crystalline form III. In some embodiments, provided herein is the omecamtiv mecarbil free base crystalline form IV. In some embodiments, provided herein is the omecamtiv mecarbil free base crystalline form V. In some embodiments, provided herein is the omecamtiv mecarbil free base crystalline form VI. In some embodiments, provided herein is the omecamtiv mecarbil free base crystalline form VII. In some embodiments, provided herein is the omecamtiv mecarbil amorphous hydrochloride salt. In some embodiments, provided herein is the omecamtiv mecarbil ethane sulfonate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil bis-fumarate crystalline salt form A. In some embodiments, provided herein is the omecamtiv mecarbil bis-fumarate crystalline salt form B. In some embodiments, provided herein is the omecamtiv mecarbil bis-fumarate crystalline salt form C. In some embodiments, provided herein is the omecamtiv mecarbil mono-fumarate crystalline salt form D. In some embodiments, provided herein is the omecamtiv mecarbil bis-maleate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil bis-malonate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil mesylate crystalline salt form A. In some embodiments, provided herein is the omecamtiv mecarbil bis-mesylate crystalline salt form B. In some embodiments, provided herein is the omecamtiv mecarbil bis-naphthalate-2-sulfonate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil mono-napadisylate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil nicotinate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil oxalate crystalline salt form A. In some embodiments, provided herein is the omecamtiv mecarbil oxalate crystalline salt form B. In some embodiments, provided herein is the omecamtiv mecarbil salicylate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil hemi-succinate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil bis-sulfate crystalline salt form A. In some embodiments, provided herein is the omecamtiv mecarbil bis-sulfate crystalline salt form B. In some embodiments, provided herein is the omecamtiv mecarbil bis-sulfate crystalline salt form C. In some embodiments, provided herein is the omecamtiv mecarbil sulfate crystalline salt form D. In some embodiments, provided herein is the omecamtiv mecarbil 2-hydroxyethane sulfonate crystalline salt. In some embodiments, provided herein is the omecamtiv mecarbil bis-tartrate crystalline salt form A. In some

embodiments, provided herein is the omecamtiv mecarbil bis-tartrate crystalline salt form B. In some embodiments, provided herein is the omecamtiv mecarbil bis-tartrate crystalline salt form C. In some embodiments, provided herein is the omecamtiv mecarbil mono-tartrate crystalline salt form D.

[0007] Also provided are pharmaceutical compositions comprising a salt or crystal form of omecamtiv mecarbil as disclosed herein and a pharmaceutically acceptable excipient.

[0008] Further provided are methods of treating heart failure in a subject in need thereof comprising administering to the subject a salt or crystal form of omecamtiv mecarbil as disclosed herein in an amount effective to treat heart failure.

BRIEF DESCRIPTION OF FIGURES

[0009] Figure 1 depicts an X-ray powder diffraction ("XRPD") pattern of the free base crystalline form III.

[0010] Figure 2 depicts a differential scanning calorimetry ("DSC") thermograph of the free crystalline form III.

[0011] Figure 3 depicts a thermogravimetric analysis ("TGA") trace of the free base crystalline form III.

[0012] Figure 4 depicts an XRPD pattern of the free base crystalline form IV.

[0013] Figure 5 depicts a DSC thermograph of the free base crystalline form IV.

[0014] Figure 6 depicts a TGA trace of the free base crystalline form IV.

[0015] Figure 7 depicts an XRPD pattern of the free base crystalline form V.

[0016] Figure 8 depicts a DSC thermograph of the free base crystalline form V.

[0017] Figure 9 depicts a TGA trace of the free base crystalline form V.

[0018] Figure 10 depicts an XRPD pattern of the free base crystalline form VI.

[0019] Figure 11 depicts a DSC thermograph of the free base crystalline form VI.

[0020] Figure 12 depicts an XRPD pattern of the free base crystalline form VII.

[0021] Figure 13 depicts an XRPD pattern overlay of the free base crystalline forms I-VII.

[0022] Figure 14 depicts an XRPD pattern of the amorphous hydrochloride salt.

[0023] Figure 15 depicts a DSC thermograph of the amorphous hydrochloride salt indicating a T_g of ~149.16°C.

[0024] Figure 16 depicts a TGA trace of the amorphous hydrochloride salt indicating a weight loss of ~7.9% from ~26°C to 160°C.

[0025] Figure 17 depicts a moisture sorption profile of the amorphous hydrochloride salt indicating a weight gain of ~11% by 50% RH then weight loss due to crystallization to Form A up to 95% RH.

[0026] Figure 18 depicts an XRPD pattern of the ethane sulfonate crystalline salt.

[0027] Figure 19 depicts a thermal gravimetric/differential thermal analysis ("TG/DTA") thermograph of the ethane sulfonate crystalline salt.

[0028] Figure 20 depicts an XRPD pattern of the bis-fumarate crystalline salt form A.

[0029] Figure 21 depicts a TGA trace of the bis-fumarate crystalline salt form A.

[0030] Figure 22 depicts an XRPD pattern of the bis-fumarate crystalline salt form B.

[0031] Figure 23 depicts a TGA trace of the bis-fumarate crystalline salt form B.

[0032] Figure 24 depicts an XRPD pattern of the bis-fumarate crystalline salt form C.

[0033] Figure 25 depicts a TGA trace of the bis-fumarate crystalline salt form C.

[0034] Figure 26 depicts an XRPD pattern of the mono-fumarate crystalline salt form D.

[0035] Figure 27 depicts a DSC thermograph of the mono-fumarate crystalline salt form D.

[0036] Figure 28 depicts a TGA trace of the mono-fumarate crystalline salt form D.

[0037] Figure 29 depicts an XRPD pattern overlay of the fumarate crystalline salts A-D.

[0038] Figure 30 depicts an XRPD pattern of the bis-maleate crystalline salt.

[0039] Figure 31 depicts a DSC thermograph of the bis-maleate crystalline salt.

[0040] Figure 32 depicts a TGA trace of the bis-maleate crystalline salt.

[0041] Figure 33 depicts an XRPD pattern of the bis-malonate crystalline salt.

[0042] Figure 34 depicts a TGA trace of the bis-malonate crystalline salt.

[0043] Figure 35 depicts an XRPD pattern of the mesylate crystalline salt form A.

[0044] Figure 36 depicts a TGA trace of the mesylate crystalline salt form A.

[0045] Figure 37 depicts an XRPD of the bis-mesylate crystalline salt form B.

[0046] Figure 38 depicts a thermal gravimetric/differential thermal analysis ("TG/DTA") of the bis-mesylate crystalline salt form B.

[0047] Figure 39 depicts an XRPD pattern overlay of the mesylate crystalline salt form A and B.

- [0048] Figure 40 depicts an XRPD pattern of the bis-naphthalate-2-sulfonate crystalline salt.
- [0049] Figure 41 depicts a TG/DTA of the bis-naphthalate-2-sulfonate crystalline salt.
- [0050] Figure 42 depicts an XRPD pattern of the mono-napadisylate crystalline salt.
- [0051] Figure 43 depicts a DSC thermograph of the mono-napadisylate crystalline salt.
- [0052] Figure 44 depicts a TGA trace of the mono-napadisylate crystalline salt.
- [0053] Figure 45 depicts an XRPD pattern of the nicotinate crystalline salt.
- [0054] Figure 46 depicts a TG/DTA of the nicotinate crystalline salt.
- [0055] Figure 47 depicts an XRPD pattern of the oxalate crystalline salt form A.
- [0056] Figure 48 depicts a DSC thermograph of the oxalate crystalline salt form A.
- [0057] Figure 49 depicts a TGA trace of the oxalate crystalline salt form A.
- [0058] Figure 50 depicts an XRPD pattern of the oxalate crystalline salt form B.
- [0059] Figure 51 depicts a TGA trace of the oxalate crystalline salt form B.
- [0060] Figure 52 depicts an XRPD overlay of the oxalate crystalline salt forms A and B.
- [0061] Figure 53 depicts an XRPD pattern of the salicylate crystalline salt.
- [0062] Figure 54 depicts a TG/DTA of the salicylate crystalline salt.
- [0063] Figure 55 depicts XRPD patterns of the hemi-succinate crystalline salts overlaid.
- [0064] Figure 56 depicts a DSC thermograph of the hemi-succinate crystalline salt.
- [0065] Figure 57 depicts an XRPD pattern of the bis-sulfate crystalline salt form A.
- [0066] Figure 58 depicts an XRPD pattern of the bis-sulfate crystalline salt form B.
- [0067] Figure 59 depicts a TGA trace of the bis-sulfate crystalline salt form B.
- [0068] Figure 60 depicts an XRPD pattern of the bis-sulfate crystalline salt form C.
- [0069] Figure 61 depicts a TGA trace of the bis-sulfate crystalline salt form C.
- [0070] Figure 62 depicts an XRPD pattern of the sulfate crystalline salt form D.
- [0071] Figure 63 depicts a TG/DTA of the sulfate crystalline salt form D.
- [0072] Figure 64 depicts an XRPD pattern overlay of the sulfate crystalline salt forms A-D.
- [0073] Figure 65 depicts an XRPD pattern of the 2-hydroxyethane sulfonate crystalline salt.

- [0074] Figure 66 depicts a TG/DTA of the 2-hydroxyethane sulfonate crystalline salt.
- [0075] Figure 67 depicts an XRPD pattern of the bis-tartrate crystalline salt form A.
- [0076] Figure 68 depicts a TGA trace of the bis-tartrate crystalline salt form A.
- [0077] Figure 69 depicts an XRPD pattern of the bis-tartrate crystalline salt form B.
- [0078] Figure 70 depicts a TGA trace of the bis-tartrate crystalline salt form B.
- [0079] Figure 71 depicts an XRPD pattern of the bis-tartrate crystalline salt form C.
- [0080] Figure 72 depicts a TGA trace of the bis-tartrate crystalline salt form C.
- [0081] Figure 73 depicts an XRPD pattern of the mono-tartrate crystalline salt form D.
- [0082] Figure 74 depicts a TGA trace of the mono-tartrate crystalline salt form D.
- [0083] Figure 75 depicts an XRPD pattern overlay of the tartrate crystalline salt form A-D.

DETAILED DESCRIPTION

[0084] The present disclosure provides salts and crystal forms of omecamtiv mecarbil.

[0085] Embodiments of the free base crystalline forms, crystalline salts and amorphous salt of omecamtiv mecarbil can be characterized by one or more of the parameters described in further detail below.

Free base crystalline forms of omecamtiv mecarbil

[0086] Provided herein are free base crystalline forms of omecamtiv mecarbil. In some embodiments, the free base crystalline forms of omecamtiv mecarbil can be nonionic forms of omecamtiv mecarbil. In some embodiments, the free base crystalline forms III-VII of omecamtiv mecarbil can be anhydrous. The free base crystalline forms I and II of omecamtiv mecarbil depicted in Figure 13 are prepared and discussed in detail in Morrison et. al, *Organic Process Research & Development*, 2015, 19, 1842-1848.

[0087] *Free base crystalline form III*

[0088] Free base crystalline form III of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 9.50, 19.06, and $23.01 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form III optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 14.27, 15.25, 16.10, 17.78, and $23.87 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form III optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 7.91, 20.65, 28.11, 31.01, 31.95, and $32.34 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form III optionally can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 1 set forth

in the Examples. In some embodiments, free base crystalline form III has an X-ray powder diffraction pattern substantially as shown in Figure 1, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. It is well known in the field of XRPD that while relative peak heights in spectra are dependent on a number of factors, such as sample preparation and instrument geometry, peak positions are relatively insensitive to experimental details.

[0089] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for free base crystalline form III. The DSC curve indicates an endothermic transition at about $186^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, free base crystalline form III can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 175°C to about 190°C . For example, in some embodiments free base crystalline form III is characterized by DSC, as shown in Figure 2.

[0090] Free base crystalline form III also can be characterized by thermogravimetric analysis (TGA). Thus, free base crystalline form III can be characterized by a weight loss in a range of about 0% to about 1% with an onset temperature in a range of about 25°C to about 100°C . For example, free base crystalline form III can be characterized by a weight loss of about 0%, up to about 150°C . In some embodiments, free base crystalline form III has a thermogravimetric analysis substantially as depicted in Figure 3, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0091] *Free base crystalline form IV*

[0092] Free base crystalline form IV of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 5.18, 10.35, 14.84, 15.54, 18.10, and $19.92 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form IV optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 14.21, 20.62, 20.77, 22.86, 24.05, 24.36, 27.81, and $29.42 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form IV optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 7.42, 7.70, 21.70, 22.40, 23.09, 25.20, 25.72, 27.40, 28.18, 28.63, 28.98, and $30.51 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form IV can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 2 set forth in the Examples. In some embodiments, free base crystalline form IV has an X-ray powder diffraction pattern substantially as shown in Figure 4, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0093] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for free base crystalline form IV. The DSC curve indicates an endothermic transition at about $185^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, free base crystalline form IV can be characterized by a DSC thermograph having a decomposition endotherm with an

onset in a range of about 175°C to about 190°C. For example, in some embodiments free base crystalline form IV is characterized by DSC, as shown in Figure 5.

[0094] Free base crystalline form IV also can be characterized by thermogravimetric analysis (TGA). Thus, free base crystalline form IV can be characterized by a weight loss in a range of about 0% to about 1% with an onset temperature in a range of about 25°C to about 100°C. For example, free base crystalline form IV can be characterized by a weight loss of about 0%, up to about 150°C. In some embodiments, free base crystalline form IV has a thermogravimetric analysis substantially as depicted in Figure 6, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0095] *Free base crystalline form V*

[0096] Free base crystalline form V of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 7.38, 8.56, 9.14, and $18.28 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form V optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 8.93, 10.03, 10.73, 11.71, 13.69, 15.08, 16.85, 17.85, 18.86, 20.05, 20.72, 21.74, 23.56, 24.03, 26.23, and $27.62 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form V optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 5.40, 16.04, 22.83, 25.45, 26.23, 27.62, 28.58, 29.85, 32.10, and $33.37 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Free base crystalline form V can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 3 set forth in the Examples. In some embodiments, free base crystalline form V has an X-ray powder diffraction pattern substantially as shown in Figure 7, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0097] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for Form V. The DSC curve indicates an endothermic transition at about $185^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, free base crystalline form V can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 175°C to about 190°C. For example, in some embodiments free base crystalline form V is characterized by DSC, as shown in Figure 8.

[0098] Free base crystalline form V also can be characterized by thermogravimetric analysis (TGA). Thus, free base crystalline form V can be characterized by a weight loss in a range of about 2% to about 6% with an onset temperature in a range of about 25°C to about 100°C. For example, free base crystalline form V can be characterized by a weight loss of about 4.2%, up to about 150°C. In some embodiments, free base crystalline form V has a thermogravimetric analysis substantially as depicted in Figure 9, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0099] *Free base crystalline form VI*

[0100] Free base crystalline form VI of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 9.07, 16.67, 18.18, 19.70, 20.89, and $21.28 \pm 0.2^\circ$ 2θ using Cu K α radiation. Free base crystalline form VI optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 15.88, 17.81, 18.80, 23.72, 24.26, 26.80, 27.59, and $29.82 \pm 0.2^\circ$ 2θ using Cu K α radiation. Free base crystalline form VI optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 14.28, 20.23, 26.19, and $28.90 \pm 0.2^\circ$ 2θ using Cu K α radiation. Free base crystalline form VI can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 4 set forth in the Examples. In some embodiments, free base crystalline form VI has an X-ray powder diffraction pattern substantially as shown in Figure 10, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0101] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for Form VI. The DSC curve indicates an endothermic transition at about $185^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, free base crystalline form VI can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 175°C to about 190°C . For example, in some embodiments free base crystalline form VI is characterized by DSC, as shown in Figure 11.

[0102] *Free base crystalline form VII*

[0103] Free base crystalline form VII of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 8.40, 8.71, 13.08, 15.66, and $19.61 \pm 0.2^\circ$ 2θ using Cu K α radiation. Free base crystalline form VII optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 4.37, 16.83, 18.92, 20.32, 20.49, 22.26, 24.21, and $25.41 \pm 0.2^\circ$ 2θ using Cu K α radiation. Free base crystalline form VII optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 7.84, 10.81, 21.61, 23.22, 23.46, 27.58, 29.53, 30.13, and $31.32 \pm 0.2^\circ$ 2θ using Cu K α radiation. Free base crystalline form VII can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 5 set forth in the Examples. In some embodiments, the free base crystalline form VII has an X-ray powder diffraction pattern substantially as shown in Figure 12, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

Crystalline salts of omecamtiv mecarbil

[0104] Provided herein are various crystalline salts of omecamtiv mecarbil. Specifically provided are crystalline salts of omecamtiv mecarbil, wherein the salt is an ethane sulfonate

salt, a fumarate salt, a maleate salt, a malonate salt, a mesylate salt, a naphthalate-2-sulfonate salt, a napadisylate salt, a nicotinate salt, an oxalate salt, a salicylate salt, a succinate salt, a sulfate salt, a hydroxyethane sulfonate salt, or a tartrate salt.

[0105] *Ethane sulfonate crystalline salt*

[0106] The ethane sulfonate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 8.61, 16.14, 16.76, 16.97, 20.73, 20.96, 25.95, and $26.30 \pm 0.2^\circ$ 2θ using Cu K α radiation. The ethane sulfonate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 17.23, 18.35, 19.20, 20.27, 23.73, 25.24, and $27.09 \pm 0.2^\circ$ 2θ using Cu K α radiation. The ethane sulfonate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 7 set forth in the Examples. In some embodiments, the ethane sulfonate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 18, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In some embodiments, ethane sulfonate crystalline salt has an TG/DTA substantially as shown in Figure 19.

[0107] *Bis-fumarate crystalline salt form A*

[0108] Bis-fumarate crystalline salt form A of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 5.64, 15.76, 22.03, and $23.87 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-fumarate crystalline salt form A optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 16.80, 21.55, 21.87, 23.61, 23.87, 26.01, and $27.20 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-fumarate crystalline salt form A optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.80, 12.04, 15.76, 16.40, 17.94, 18.32, 19.87, 20.61, 22.88, 27.86, 32.73, and $36.54 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-fumarate crystalline salt form A can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 8 set forth in the Examples. In some embodiments, the bis-fumarate crystalline salt form A has an X-ray powder diffraction pattern substantially as shown in Figure 20, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0109] Bis-fumarate crystalline salt form A also can be characterized by thermogravimetric analysis (TGA). Thus, bis-fumarate crystalline salt form A can be characterized by a weight loss in a range of about 6% to about 10% with an onset temperature in a range of about 25°C to about 100°C. For example, bis-fumarate crystalline salt form A can be characterized by a weight loss of about 7.9%, up to about 150°C. In some embodiments, bis-fumarate crystalline salt form A has a thermogravimetric analysis substantially as

depicted in Figure 21, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^{\circ}\text{C}$.

[0110] *Bis-fumarate crystalline salt form B*

[0111] Bis-fumarate crystalline salt form B of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 5.68, 6.11, 13.13, 18.08, and $22.47 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-fumarate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 9.69, 11.43, 12.92, 15.95, 20.81, 22.95, 26.04, 27.01, and $28.43 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-fumarate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 19.52, 24.53, 31.37, 32.32, 34.89, 35.89, and $37.16 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-fumarate crystalline salt form B optionally can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 9 set forth in the Examples. In some embodiments, the bis-fumarate crystalline salt form B has an X-ray powder diffraction pattern substantially as shown in Figure 22, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^{\circ}$.

[0112] Bis-fumarate crystalline salt form B also can be characterized by thermogravimetric analysis (TGA). Thus, bis-fumarate crystalline salt form B can be characterized by a weight loss in a range of about 4% to about 8% with an onset temperature in a range of about 25°C to about 100°C . For example, bis-fumarate crystalline salt form B can be characterized by a weight loss of about 5.6%, up to about 150°C . In some embodiments, bis-fumarate crystalline salt form B has a thermogravimetric analysis substantially as depicted in Figure 23, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^{\circ}\text{C}$.

[0113] *Bis-fumarate crystalline salt form C*

[0114] Bis-fumarate crystalline salt form C of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 5.88, 18.79, 25.41, and $26.86 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-fumarate crystalline salt form C optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 12.74, 13.56, 17.15, 17.63, 20.29, 21.47, 21.77, 22.21, 22.92, 23.58, 24.15, 25.41, 26.78, and $27.83 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-fumarate crystalline salt form C can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 10 set forth in the Examples. In some embodiments, the bis-fumarate crystalline salt form C has an X-ray powder diffraction pattern substantially as shown in

Figure 24, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0115] Bis-fumarate crystalline salt form C also can be characterized by thermogravimetric analysis (TGA). Thus, bis-fumarate crystalline salt form C can be characterized by a weight loss in a range of about 6% to about 10% with an onset temperature in a range of about 25°C to about 100°C . For example, bis-fumarate crystalline salt form C can be characterized by a weight loss of about 8.4%, up to about 150°C . In some embodiments, bis-fumarate crystalline salt form C has a thermogravimetric analysis substantially as depicted in Figure 25, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$. This weight loss was determined to be water via Karl Fischer (KF) analysis. KF analysis shows that the water content of bis-fumarate crystalline salt form C can be about 8.4%, corresponding to a trihydrate.

[0116] *Mono-fumarate crystalline salt form D*

[0117] Mono-fumarate crystalline salt form D of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 8.01, 15.20, and $20.02 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Mono-fumarate crystalline salt form D optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 12.11, 12.67, 14.46, 16.01, 16.57, 17.04, 17.63, 20.51, 21.75, 22.86, 24.25, 24.97, 25.84, 26.17, 27.10, 27.97, and $29.21 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Mono-fumarate crystalline salt form D can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 11 set forth in the Examples. In some embodiments, mono-fumarate crystalline salt form D has an X-ray powder diffraction pattern substantially as shown in Figure 26, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0118] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for mono-fumarate crystalline salt form D. The DSC curve indicates an endothermic transition at about $125^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, mono-fumarate crystalline salt form D can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 110°C to about 130°C . For example, in some embodiments mono-fumarate crystalline salt form D is characterized by DSC, as shown in Figure 27.

[0119] Mono-fumarate crystalline salt form D also can be characterized by thermogravimetric analysis (TGA). Thus, mono-fumarate crystalline salt form D can be characterized by a weight loss in a range of about 5% to about 9% with an onset temperature in a range of about 25°C to about 100°C . For example, mono-fumarate crystalline salt form D can be characterized by a weight loss of about 6.7%, up to about

150°C. In some embodiments, mono-fumarate crystalline salt form D has a thermogravimetric analysis substantially as depicted in Figure 28, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0120] *Bis-maleate crystalline salt*

[0121] The bis-maleate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 9.97, 15.31, 16.04, and $26.96 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The bis-maleate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.56, 13.25, 15.53, 16.38, 17.44, 17.70, 18.17, 19.00, 20.13, 21.47, 22.31, 22.44, 24.38, 24.64, 25.66, 26.66, and $27.83 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The bis-maleate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at 4.99, 14.83, 17.10, 22.02, 28.55, 30.76, 32.01, 34.39, and $34.51 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The bis-maleate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 13 set forth in the Examples. In some embodiments, the bis-maleate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 30, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0122] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for the bis-maleate crystalline salt. The DSC curve indicates an endothermic transition at about $190^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, the bis-maleate crystalline salt can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 160°C to about 210°C . For example, in some embodiments the bis-maleate crystalline salt is characterized by DSC, as shown in Figure 31.

[0123] The bis-maleate crystalline salt also can be characterized by thermogravimetric analysis (TGA). Thus, the bis-maleate crystalline salt can be characterized by a weight loss in a range of about 0% to about 1% with an onset temperature in a range of about 25°C to about 150°C . For example, the bis-maleate crystalline salt can be characterized by a weight loss of about 0%, up to about 150°C . In some embodiments, the bis-maleate crystalline salt has a thermogravimetric analysis substantially as depicted in Figure 32, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0124] *Bis-malonate crystalline salt*

[0125] The bis-malonate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 4.74, 11.37, 14.25, 15.13, 18.29, 20.14, 23.87, 27.78, and $28.01 \pm 0.2^\circ 2\theta$ using Cu

K α radiation. The bis-malonate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 9.30, 13.73, 16.45, 16.83, 18.08, 18.88, 19.54, 20.77, 21.21, 23.32, 24.67, 26.51, 27.59, and $28.90 \pm 0.2^\circ$ 2θ using Cu K α radiation. The bis-malonate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 15.69, 25.72, 30.18, 33.70, $34.19 \pm 0.2^\circ$ 2θ using Cu K α radiation. The bis-malonate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 14 set forth in the Examples. In some embodiments, the bis-malonate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 33, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0126] The bis-malonate crystalline salt also can be characterized by thermogravimetric analysis (TGA). Thus, the bis-malonate crystalline salt can be characterized by a weight loss in a range of about 0% to about 1% with an onset temperature in a range of about 25°C to about 140°C . For example, the bis-malonate crystalline salt can be characterized by a weight loss of about 0%, up to about 140°C . In some embodiments, the bis-malonate crystalline salt has a thermogravimetric analysis substantially as depicted in Figure 34, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0127] *Mesylate crystalline salt form A*

[0128] Mesylate crystalline salt form A of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 4.02, 4.87, 15.21, 15.86, 20.53, and $24.39 \pm 0.2^\circ$ 2θ using Cu K α radiation. Mesylate crystalline salt form A optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 7.79, 11.61, 16.51, 17.57, 18.42, 19.26, 21.55, 23.17, 25.51, 26.38, and $27.63 \pm 0.2^\circ$ 2θ using Cu K α radiation. Mesylate crystalline salt form A can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 15 set forth in the Examples. In some embodiments, mesylate crystalline salt form A has an X-ray powder diffraction pattern substantially as shown in Figure 35, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0129] Mesylate crystalline salt form A also can be characterized by thermogravimetric analysis (TGA). Thus, mesylate crystalline salt form A can be characterized by a weight loss in a range of about 0% to about 2% with an onset temperature in a range of about 25°C to about 175°C . For example, mesylate crystalline salt form A can be characterized by a weight loss of about 1.0%, up to about 200°C . In some embodiments, mesylate crystalline salt form A has a thermogravimetric analysis substantially as depicted in Figure 36, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0130] *Bis-mesylate crystalline salt form B*

[0131] Bis-mesylate crystalline salt form B of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 8.30, 8.94, 9.59, 12.15, 14.37, 19.82, 20.29, 22.04, and $25.02 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-mesylate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 11.66, 16.18, 16.64, 16.81, 17.07, 17.19, 17.41, 17.76, 19.24, 20.66, 21.62, 22.39, 23.95, 24.60, 25.59, 25.89, 27.14, 27.35, 27.41, and $29.45 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-mesylate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.78, 11.15, 14.93, 15.36, 15.57, 23.54, 26.14, 26.49, 27.89, 28.86, 29.89, 31.11, 32.47, 33.10, 33.51, $34.56 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-mesylate crystalline salt form B can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 16 set forth in the Examples. In some embodiments, bis-mesylate crystalline salt form B has an X-ray powder diffraction pattern substantially as shown in Figure 37, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In some embodiments, bis-mesylate crystalline salt form B has a TG/DTA substantially as shown in Figure 38.

[0132] *Bis-naphthalate-2-sulfonate crystalline salt*

[0133] The bis-naphthalate-2-sulfonate crystalline salt can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 4.49, 18.20, 18.62, 21.38, 21.52, and $26.11 \pm 0.2^\circ$ 2θ using Cu K α radiation. The bis-naphthalate-2-sulfonate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 6.25, 6.65, 13.44, 14.39, 14.92, 16.28, 18.90, 19.53, 20.82, 22.02, 22.43, 22.80, 24.40, 25.16, 27.01, 29.67, 31.63, and $33.42 \pm 0.2^\circ$ 2θ using Cu K α radiation. The bis-naphthalate-2-sulfonate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 18 set forth in the Examples. In some embodiments, the bis-naphthalate-2-sulfonate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 40, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In some embodiments, bis-naphthalate-2-sulfonate crystalline salt has a TG/DTA substantially as shown in Figure 41.

[0134] *Mono-napadisylate crystalline salt*

[0135] The mono-napadisylate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 12.27, 15.75, 16.5, 17.83, 19.94, 21.83, and $22.87 \pm 0.2^\circ$ 2θ using Cu K α radiation. The mono-napadisylate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 13.41, 14.57, 15.14,

18.82, 23.49, 24.34, and $25.26 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The mono-napadisylate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 19 set forth in the Examples. In some embodiments, the mono-napadisylate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 42, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0136] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for the mono-napadisylate crystalline salt. The DSC curve indicates an endothermic transition at about $100^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, the mono-napadisylate crystalline salt can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 80°C to about 115°C . For example, in some embodiments the mono-napadisylate crystalline salt is characterized by DSC, as shown in Figure 43.

[0137] The mono-napadisylate crystalline salt also can be characterized by thermogravimetric analysis (TGA). Thus, the mono-napadisylate crystalline salt can be characterized by a weight loss in a range of about 4% to about 8% with an onset temperature in a range of about 20°C to about 100°C . For example, the mono-napadisylate crystalline salt can be characterized by a weight loss of about 5.8%, up to about 100°C . In some embodiments, the mono-napadisylate crystalline salt has a thermogravimetric analysis substantially as depicted in Figure 44, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0138] *Nicotinate crystalline salt*

[0139] The nicotinate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 3.69, 8.55, 9.13, 16.70, 16.84, 18.30, 19.99, 20.76, 23.43, 24.83, and $25.95 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The nicotinate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 7.36, 10.01, 12.43, 14.74, 15.50, 17.62, 18.58, 19.59, 20.34, 21.32, 22.03, 22.91, 23.87, 24.92, 25.40, 26.85, 26.94, 27.32, 28.01, and $28.94 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The nicotinate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 20 set forth in the Examples. In some embodiments, the nicotinate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 45, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In some embodiments, the nicotinate crystalline salt form D has an TG/DTA substantially as shown in Figure 46.

[0140] *Oxalate crystalline salt form A*

[0141] Oxalate crystalline salt form A of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 6.48, 13.01, and $23.82 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Oxalate crystalline salt form A optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.36, 11.85, 14.79, 15.35, 17.11, 19.23, 19.91, 21.48, 22.07, 22.75, 25.70, 28.55, and $30.71 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Oxalate crystalline salt form A can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 21 set forth in the Examples. In some embodiments, oxalate crystalline salt form A has an X-ray powder diffraction pattern substantially as shown in Figure 47, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0142] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for oxalate crystalline salt form A. The DSC curve indicates an endothermic transition at about $209^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, oxalate crystalline salt form A can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 190°C to about 230°C . For example, in some embodiments oxalate crystalline salt form A is characterized by DSC, as shown in Figure 48.

[0143] Oxalate crystalline salt form A also can be characterized by thermogravimetric analysis (TGA). Thus, oxalate crystalline salt form A can be characterized by a weight loss in a range of about 0.5% to about 4.5% with an onset temperature in a range of about 25°C to about 100°C . For example, oxalate crystalline salt form A can be characterized by a weight loss of about 2.5%, up to about 150°C . In some embodiments, oxalate crystalline salt form A has a thermogravimetric analysis substantially as depicted in Figure 49, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0144] *Oxalate crystalline salt form B*

[0145] Oxalate crystalline salt form B of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 7.38, 13.30, and $16.54 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Oxalate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 17.11, 17.95, 18.45, 21.25, 22.63, 24.82, and $25.77 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Oxalate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 14.76, 24.35, 28.61, 29.58, 30.49, 31.76, 34.46, and $37.35 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Oxalate crystalline salt form B can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 22 set forth in the Examples. In some embodiments, oxalate crystalline salt form B has an X-ray powder diffraction pattern substantially as shown in Figure 50, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0146] Oxalate crystalline salt form B also can be characterized by thermogravimetric analysis (TGA). Thus, oxalate crystalline salt form B can be characterized by a weight loss in a range of about 0% to about 1% with an onset temperature in a range of about 25°C to about 100°C. For example, oxalate crystalline salt form B can be characterized by a weight loss of about 0%, up to about 150°C. In some embodiments, oxalate crystalline salt form B has a thermogravimetric analysis substantially as depicted in Figure 51, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0147] *Salicylate crystalline salt*

[0148] The salicylate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 8.36, 16.75, 17.56, 23.58, and $28.21 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The salicylate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.08, 11.30, 13.69, 17.77, 17.86, 18.67, 19.11, 20.22, 21.07, 25.23, and $27.40 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The salicylate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 9.78, 12.00, 13.80, 15.51, 19.27, 19.62, 20.02, 20.79, 22.19, 22.39, 22.75, 22.92, 24.99, 25.59, 26.79, 29.94, and $34.07 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The salicylate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 24 set forth in the Examples. In some embodiments, the salicylate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 53, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In some embodiments, the salicylate crystalline salt has an TG/DTA substantially as shown in Figure 54.

[0149] *Hemi-succinate crystalline salt*

[0150] The hemi-succinate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 6.32, 18.87, 19.32, 20.5, 21.24, 21.89, 23.49, 24.23, and $26.71 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The hemi-succinate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 12.93, 15.08, 16.97, 25.36, 27.39, and $28.32 \pm 0.2^\circ 2\theta$ using Cu K α radiation. The hemi-succinate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 25 set forth in the Examples. In some embodiments, the hemi-succinate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 55, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0151] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for the hemi-succinate crystalline salt. The DSC curve indicates an

endothermic transition at about $171^{\circ}\text{C} \pm 3^{\circ}\text{C}$. Thus, in some embodiments, the hemi-succinate crystalline salt can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 155°C to about 190°C . For example, in some embodiments the hemi-succinate crystalline salt is characterized by DSC, as shown in Figure 56.

[0152] *Bis-sulfate crystalline salt form A*

[0153] Bis-sulfate crystalline salt form A of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 5.39, 7.55, 14.35, 19.26, and $20.22 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-sulfate crystalline salt form A optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 16.17, 16.71, 16.92, 17.07, 18.60, 20.83, 21.38, 22.27, 22.77, 23.14, 23.42, 23.76, 24.32, 25.11, 25.74, 26.46, 27.71, 28.15, and $29.92 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-sulfate crystalline salt form A can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 26 set forth in the Examples. In some embodiments, bis-sulfate crystalline salt form A has an X-ray powder diffraction pattern substantially as shown in Figure 57, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^{\circ}$.

[0154] *Bis-sulfate crystalline salt form B*

[0155] Bis-sulfate crystalline salt form B of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 11.72 and $20.48 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-sulfate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 12.17, 12.93, 17.79, 18.39, 18.76, 19.84, 23.60, 25.13, 25.63, and $30.12 \pm 0.2^{\circ} 2\theta$ using Cu K α radiation. Bis-sulfate crystalline salt form B can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 27 set forth in the Examples. In some embodiments, bis-sulfate crystalline salt form B has an X-ray powder diffraction pattern substantially as shown in Figure 58, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^{\circ}$.

[0156] Bis-sulfate crystalline salt form B also can be characterized by thermogravimetric analysis (TGA). Thus, bis-sulfate crystalline salt form B can be characterized by a weight loss in a range of about 10% to about 14% with an onset temperature in a range of about 25°C to about 100°C . For example, bis-sulfate crystalline salt form B can be characterized by a weight loss of about 12.2%, up to about 150°C . In some embodiments, bis-sulfate crystalline salt form B has a thermogravimetric analysis substantially as depicted in Figure 59, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^{\circ}\text{C}$. This weight loss was determined to be water via Karl Fischer (KF) analysis. KF analysis

shows that the water content of bis-sulfate crystalline salt form B can be about 12%, corresponding to a pentahydrate.

[0157] *Bis-sulfate crystalline salt form C*

[0158] Bis-sulfate crystalline salt form C of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 10.98, 11.49, 18.04, and $19.60 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-sulfate crystalline salt form C optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.39, 10.72, 12.52, 12.99, 17.11, 17.43, 20.94, 24.76, 25.25, 25.87, and $26.51 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-sulfate crystalline salt form C can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 28 set forth in the Examples. In some embodiments, bis-sulfate crystalline salt form C has an X-ray powder diffraction pattern substantially as shown in Figure 60, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0159] Bis-sulfate crystalline salt form C also can be characterized by thermogravimetric analysis (TGA). Thus, bis-sulfate crystalline salt form C can be characterized by a weight loss in a range of about 7% to about 11% with an onset temperature in a range of about 25°C to about 60°C . For example, bis-sulfate crystalline salt form C can be characterized by a weight loss of about 9.0%, up to about 150°C . In some embodiments, bis-sulfate crystalline salt form C has a thermogravimetric analysis substantially as depicted in Figure 61, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0160] *Sulfate crystalline salt form D*

[0161] Sulfate crystalline salt form D of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 7.32, 8.02, and $20.44 \pm 0.2^\circ$ 2θ using Cu K α radiation. Sulfate crystalline salt form D optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 13.57, 14.54, 16.29, 16.41, 16.91, 17.36, 18.70, 21.02, 21.77, 22.37, 22.90, 23.72, 24.28, 25.14, 25.88, 26.58, 27.25, 28.10, and $29.43 \pm 0.2^\circ$ 2θ using Cu K α radiation. Sulfate crystalline salt form D can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 29 set forth in the Examples. In some embodiments, sulfate crystalline salt form D has an X-ray powder diffraction pattern substantially as shown in Figure 62, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In embodiments, sulfate crystalline salt form D has a TG/DTA substantially as shown in Figure 63.

[0162] *2-Hydroxyethane sulfonate crystalline salt*

[0163] The 2-hydroxyethane sulfonate crystalline salt of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 9.95, 17.85, 19.93, 20.07, 20.46, 25.06, and $26.20 \pm 0.2^\circ$ 2θ using Cu K α radiation. The 2-hydroxyethane sulfonate crystalline salt optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 6.26, 6.69, 14.99, 16.37, 19.61, 20.95, 29.98, 32.16, and $34.39 \pm 0.2^\circ$ 2θ using Cu K α radiation. The 2-hydroxyethane sulfonate crystalline salt can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 31 set forth in the Examples. In some embodiments, the 2-hydroxyethane sulfonate crystalline salt has an X-ray powder diffraction pattern substantially as shown in Figure 65, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$. In some embodiments, the 2-hydroxyethane sulfonate crystalline salt has a TG/DTA substantially as shown in Figure 66.

[0164] *Bis-tartrate crystalline salt form A*

[0165] Bis-tartrate crystalline salt form A of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 4.20, 7.49, 8.22, 11.88, 16.42, and $21.19 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-tartrate crystalline salt form A optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 4.77, 7.67, 8.43, 9.49, 13.05, 13.26, 14.98, 15.14, 17.34, 17.47, 18.02, 18.23, 18.72, 19.20, 22.50, 24.53, 25.67, 26.30, and $28.14 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-tartrate crystalline salt form A can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 32 set forth in the Examples. In some embodiments, bis-tartrate crystalline salt form A has an X-ray powder diffraction pattern substantially as shown in Figure 67, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0166] Bis-tartrate crystalline salt form A also can be characterized by thermogravimetric analysis (TGA). Thus, bis-tartrate crystalline salt form A can be characterized by a weight loss in a range of about 1% to about 5% with an onset temperature in a range of about 25°C to about 120°C . For example, bis-tartrate crystalline salt form A can be characterized by a weight loss of about 3.2%, up to about 150°C . In some embodiments, bis-tartrate crystalline salt form A has a thermogravimetric analysis substantially as depicted in Figure 68, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0167] *Bis-tartrate crystalline salt form B*

[0168] Bis-tartrate crystalline salt form B of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 3.77, 5.69, and $10.07 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-tartrate crystalline salt form B optionally can be further characterized by an X-ray powder diffraction pattern having

additional peaks at about 4.72, 6.95, 9.34, 11.18, 12.63, 15.18, 17.69, 22.35, and $25.46 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-tartrate crystalline salt form B can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 33 set forth in the Examples. In some embodiments, bis-tartrate crystalline salt form B has an X-ray powder diffraction pattern substantially as shown in Figure 69, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0169] Bis-tartrate crystalline salt form B also can be characterized by thermogravimetric analysis (TGA). Thus, bis-tartrate crystalline salt form B can be characterized by a weight loss in a range of about 3% to about 7% with an onset temperature in a range of about 20°C to about 100°C . For example, bis-tartrate crystalline salt form B can be characterized by a weight loss of about 5.4%, up to about 150°C . In some embodiments, bis-tartrate crystalline salt form B has a thermogravimetric analysis substantially as depicted in Figure 70, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0170] *Bis-tartrate crystalline salt form C*

[0171] Bis-tartrate crystalline salt form C of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks at about 3.57, 6.23, and $15.84 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-tartrate crystalline salt form C optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 3.86, 4.78, 7.04, 9.36, 13.08, 13.96, 16.88, 17.60, 18.20, 18.73, 20.40, 22.58, 25.44, 26.06, and $28.61 \pm 0.2^\circ$ 2θ using Cu K α radiation. Bis-tartrate crystalline salt form C can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 34 set forth in the Examples. In some embodiments, bis-tartrate crystalline salt form C has an X-ray powder diffraction pattern substantially as shown in Figure 71, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0172] Bis-tartrate crystalline salt form C also can be characterized by thermogravimetric analysis (TGA). Thus, bis-tartrate crystalline salt form C can be characterized by a weight loss in a range of about 12% to about 17% with an onset temperature in a range of about 20°C to about 100°C . For example, bis-tartrate crystalline salt form C can be characterized by a weight loss of about 14.6%, up to about 150°C . In some embodiments, bis-tartrate crystalline salt form C has a thermogravimetric analysis substantially as depicted in Figure 72, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

[0173] *Mono-tartrate crystalline salt form D*

[0174] Mono-tartrate crystalline salt form D of omecamtiv mecarbil can be characterized by an X-ray powder diffraction pattern, obtained as set forth in the Examples, having peaks

at about 9.77 and $15.40 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Mono-tartrate crystalline salt form D optionally can be further characterized by an X-ray powder diffraction pattern having additional peaks at about 10.87 , 13.79 , 17.36 , 17.74 , 18.58 , 18.87 , 21.78 , 25.43 , and $26.24 \pm 0.2^\circ 2\theta$ using Cu K α radiation. Mono-tartrate crystalline salt form D can be characterized by an X-ray powder diffraction pattern having peaks shown in Table 35 set forth in the Examples. In some embodiments, mono-tartrate crystalline salt form D has an X-ray powder diffraction pattern substantially as shown in Figure 73, wherein by “substantially” is meant that the reported peaks can vary by about $\pm 0.2^\circ$.

[0175] Mono-tartrate crystalline salt form D also can be characterized by thermogravimetric analysis (TGA). Thus, mono-tartrate crystalline salt form D can be characterized by a weight loss in a range of about 4% to about 8% with an onset temperature in a range of about 20°C to about 75°C . For example, mono-tartrate crystalline salt form D can be characterized by a weight loss of about 6.4%, up to about 150°C . In some embodiments, mono-tartrate crystalline salt form D has a thermogravimetric analysis substantially as depicted in Figure 74, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$. This weight loss was determined to be water via Karl Fischer (KF) analysis. KF analysis shows that the water content of mono-tartrate crystalline salt form D can be about 6.9%, corresponding to a dihydrate.

[0176] *Amorphous hydrochloride salt*

[0177] Also provided herein is an amorphous hydrochloride salt of omecamtiv mecarbil, and confirmation of its amorphous nature is provided by its X-ray powder diffraction pattern as shown in Figure 14.

[0178] Differential scanning calorimetry (DSC) thermographs were obtained, as set forth in the Examples, for the amorphous hydrochloride salt. The DSC curve indicates an endothermic transition at about $171^\circ\text{C} \pm 3^\circ\text{C}$. Thus, in some embodiments, the amorphous hydrochloride salt can be characterized by a DSC thermograph having a decomposition endotherm with an onset in a range of about 155°C to about 190°C . For example, in some embodiments the amorphous hydrochloride salt is characterized by DSC, as shown in Figure 15.

[0179] The amorphous hydrochloride salt also can be characterized by thermogravimetric analysis (TGA). Thus, the amorphous hydrochloride salt can be characterized by a weight loss in a range of about 6% to about 10% with an onset temperature in a range of about 20°C to about 60°C . For example, the amorphous hydrochloride salt can be characterized by a weight loss of about 7.9%, up to about 150°C . In some embodiments, the amorphous hydrochloride salt has a thermogravimetric analysis substantially as depicted in Figure 16, wherein by “substantially” is meant that the reported TGA features can vary by about $\pm 5^\circ\text{C}$.

In some embodiments, the amorphous hydrochloride salt has a moisture sorption profile substantially as depicted in Figure 17.

Pharmaceutical compositions

[0180] Also provided herein are pharmaceutical compositions comprising a salt or crystalline form of omecamtiv mecarbil described herein; and a pharmaceutically acceptable excipient.

[0181] The phrase “pharmaceutically acceptable” is employed herein to refer to those ligands, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. The compositions described herein can be formulated for any form of administration. In various cases, the composition is for oral administration. In various cases, the composition is in tablet form.

[0182] In some embodiments, the pharmaceutical compositions can include a pharmaceutically acceptable carrier. The phrase “pharmaceutically acceptable carrier” as used herein means a pharmaceutically acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material. As used herein the language “pharmaceutically acceptable carrier” includes buffers, sterile water for injection, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose, and sucrose; (2) starches, such as corn starch, potato starch, and substituted or unsubstituted β -cyclodextrin; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil, and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol, and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations. In certain embodiments, pharmaceutical compositions provided herein are non-pyrogenic, i.e., do not induce significant temperature elevations when administered to a patient.

[0183] Wetting agents, emulsifiers, and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring, and perfuming agents, preservatives and antioxidants can also be present in the compositions as excipients.

[0184] Examples of pharmaceutically acceptable antioxidants as excipient include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite, and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

[0185] A pharmaceutical composition may also contain adjuvants such as preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include tonicity-adjusting agents, such as sugars and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

[0186] In some cases, in order to prolong the effect of one or more compounds provided herein, it is desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. For example, delayed absorption of a parenterally administered compound can be accomplished by dissolving or suspending the compound in an oil vehicle.

[0187] The composition should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, and sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate and gelatin.

[0188] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are

prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the methods of preparation are freeze-drying (lyophilization), which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0189] Injectable depot forms can be made by forming microencapsule or nanoencapsule matrices of a compound provided herein in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes, microemulsions or nanoemulsions, which are compatible with body tissue.

[0190] In one embodiment, the therapeutic crystalline salts are prepared with carriers that will protect the therapeutic compounds against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Such formulations can be prepared using standard techniques, or obtained commercially, e.g., from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to selected cells with monoclonal antibodies to cellular antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811, which is incorporated herein by reference in its entirety.

[0191] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[0192] Controlled Release Compositions

[0193] In various cases, the pharmaceutical formulations described herein are capable of releasing omecamtiv mecarbil evenly at a pace controlled by the diffusion of omecamtiv mecarbil through a gel layer formed by the hydration of the control release agents in the tablets. In some embodiments, in conjunction with other above or below embodiments, the present modified release matrix tablets demonstrate a minimal pH-dependent release in-vitro. In some embodiments, in conjunction with other above or below embodiments, complete release of omecamtiv mecarbil is achieved in both pH 2 and 6.8 dissolution medium within 24 hours, possibly resulting in less inter- and intra-subject variability and food effect. It is found that the present modified release matrix tablet dosage form is superior to

the former immediate release dosage form in minimizing the plasma peak-trough ratio. As a result, the present modified release matrix tablets reduce plasma concentration fluctuation, leading to reduced side effects, and improved safety and efficacy. It is also expected that the present modified release matrix tablets will improve patient compliance by reducing the dosing frequency. Additionally, the present modified release matrix tablets are physicochemically stable—resulting in no physical attribute, assay, impurity, or dissolution profile changes after storage at 40 °C/75%RH for 6 months.

[0194] Provided are pharmaceutical formulations comprising the salt or crystalline omecamtiv mecarbil as disclosed herein; a control release agent; a pH modifying agent; a filler; and a lubricant.

[0195] As used herein, the term “control release agents” refer to agents that facilitate the release of the active ingredient from the present composition in a controlled fashion. In some embodiments, in conjunction with other above or below embodiments, the control release agents form a gel upon hydration. Control release agents include pullulan, dextrin, sodium and calcium acid, polyacrylic acid, polymethacrylic acid, polymethylvinylether co-maleic anhydride, polyvinylpyrrolidone, polyethylene oxide, polyethylene glycol, hydroxypropylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose, hydroxymethyl methacrylate, sodium carboxymethylcellulose, calcium carboxymethylcellulose, methylcellulose, maltodextrin, xanthan gum, tragacanth gum, agar, gellan gum, kayara gum, alginic acids, pectins, pre-gelatinized starch, polyvinyl alcohol, carboxymethylethylcellulose, cellulose acetate phthalate, cellulose acetate succinate, methylcellulose phthate, hydroxymethylethylcellulosephthate, hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate succinate, polyvinyl alcohol phthalate, polyvinyl butylate phthalate, polyvinyl acetal phthalate, a copolymer of vinyl acetate/maleic anhydride, a copolymer of styrene/maleic acid monoester, a copolymer of methyl acrylate/methacrylic acid, a copolymer of styrene/acrylic acid, a copolymer of methyl acrylate/methacrylic acid/octyl acrylate, a copolymer of methacrylic acid/methyl methacrylate, benzylaminomethylcellulose, diethylaminomethylcellulose, piperidylethylhydroxyethylcellulose, cellulose acetate dimethylaminoacetate, a copolymer of vinyl diethylamine/vinyl acetate, a copolymer of vinyl benzylamine/vinyl acetate, polyvinyl acetaldimethylamino acetate, a copolymer of vinylpiperidylacetoacetal/vinyl acetate, polydiethylaminomethylstyrene, a copolymer of methyl methacrylate/butyl methacrylate/dimethylaminoethyl methacrylate and polydimethylaminoethylmethacrylate, a copolymer of 2-methyl-5-vinylpyridine/methylmethacrylate/methacrylic acid, a copolymer of 2-methyl-5-vinylpyridine/methyl acrylate/methacrylic acid, a copolymer of 2-vinyl-5-ethylpyridine/methacrylic acid/methyl acrylate, a copolymer of 2-vinylpyridine/methacrylic

acid/acrylonitrile, carboxymethylpiperidyl starch, carboxy-methylbenzylaminocellulose, a copolymer of N-vinylglycine/styrene, chitosan, poly(vinyl alcohol), maleic anhydride copolymer, poly (vinyl pyrrolidone), starch and starch-based polymers, poly (2-ethyl-2-oxazoline), poly(ethyleneimine), polyurethane hydrogels, welan gum, rhamosan gum, polyvinyl acetates, ethylcellulose, eudragit RL, RS, NE 30D, Kollicoat EMM 30D, or combinations thereof.

[0196] In some embodiments, in conjunction with other above or below embodiments, the control release agent is a polymer.

[0197] In some embodiments, in conjunction with other above or below embodiments, the control release agent is selected from pullulan, dextrin, sodium and calcium alginate, polyacrylic acid, polymethacrylic acid, polymethylvinylether co-maleic anhydride, polyvinylpyrrolidone, polyethylene oxide, polyethylene glycol, hydroxypropylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose, hydroxymethyl methacrylate, sodium carboxymethylcellulose, calcium carboxymethylcellulose, methylcellulose, maltodextrin, xanthan gum, tragacanth gum, agar, gellan gum, kayara gum, alginic acids, pectins, pregelatinized starch, polyvinyl alcohol, carboxymethylcellulose, cellulose acetate phthalate, cellulose acetate succinate, methylcellulose phthalate, hydroxymethylcellulosephthalate, hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate succinate, polyvinyl alcohol phthalate, polyvinyl butyrate phthalate, polyvinyl acetate phthalate, a copolymer of vinyl acetate/maleic anhydride, a copolymer of styrene/maleic acid monoester, a copolymer of methyl acrylate/methacrylic acid, a copolymer of styrene/acrylic acid, a copolymer of methyl acrylate/methacrylic acid/octyl acrylate, a copolymer of methacrylic acid/methyl methacrylate, benzylaminomethylcellulose, diethylaminomethylcellulose, piperidylethylhydroxyethylcellulose, cellulose acetate dimethylaminoacetate, a copolymer of vinyl diethylamine/vinyl acetate, a copolymer of vinyl benzylamine/vinyl acetate, polyvinyl acetaldimethylamino acetate, a copolymer of vinylpiperidylacetoacetal/vinyl acetate, polydiethylaminomethylstyrene, a copolymer of methyl methacrylate/butyl methacrylate/dimethylaminoethyl methacrylate and polydimethylaminoethyl methacrylate, a copolymer of 2-methyl-5-vinylpyridine/methylmethacrylate/methacrylic acid, a copolymer of 2-methyl-5-vinylpyridine/methyl acrylate/methacrylic acid, a copolymer of 2-vinyl-5-ethylpyridine/methacrylic acid/methyl acrylate, a copolymer of 2-vinylpyridine/methacrylic acid/acrylonitrile, carboxymethylpiperidyl starch, carboxy-methylbenzylaminocellulose, a copolymer of N-vinylglycine/styrene, chitosan, poly(vinyl alcohol), maleic anhydride copolymer, poly (vinyl pyrrolidone), starch and starch-based polymers, poly (2-ethyl-2-oxazoline), poly(ethyleneimine), polyurethane hydrogels, welan gum, rhamosan gum,

polyvinyl acetates, ethylcellulose, eudragit RL, RS, NE 30D, and Kollicoat EMM 30D, or any combination thereof.

[0198] As used herein, the term “pH modifying agent” refers to an agent capable of modulating the pH to a desired range. In some embodiments, in conjunction with other above or below embodiments, the pH modifying agent is an acidifying agent. In some embodiments, in conjunction with other above or below embodiments, the pH modifying agent is present in an amount sufficient to lower the pH. pH modifying agents include maleic acid, citric acid, tartaric acid, pamoic acid, fumaric acid, salicylic acid, 2,6-diaminohexanoic acid, camphorsulfonic acid, glycerophosphoric acid, 2-hydroxyethanesulfonic acid, isethionic acid, succinic acid, carbonic acid, p-toluenesulfonic acid, aspartic acid, 8-chloro-theophylline, benzenesulfonic acid, malic acid, orotic acid, oxalic acid, benzoic acid, 2-naphthalenesulfonic acid, stearic acid, adipic acid, p-amino-salicylic acid, 5-aminosalicylic acid, ascorbic acid, sulfuric acid, cyclamic acid, sodium lauryl sulfate, glucoheptonic acid, glucuronic acid, glycine, sulfuric acid, mandelic acid, 1,5-naphthalenedisulfonic acid, nicotinic acid, oleic acid, 2-oxoglutaric acid, pyridoxal 5-phosphate, undecanoic acid, p-acetamidobenzoic acid, o-acetamido-benzoic acid, m-acetamidobenzoic acid, N-acetyl-L-aspartic acid, camphoric acid, dehydrocholic acid, malonic acid, edetic acid, ethylenediaminetetraacetic acid, ethylsulfuric acid, hydroxyphenylbenzoylbenzoic acid, glutamic acid, glycyrrhizic acid, 4-hexylresorcinol, hippuric acid, p-phenolsulfonic acid, 4-hydroxybenzoic acid, 3-hydroxybenzoic acid, 3-hydroxy-2-naphthoic acid, 1-hydroxy-2-naphthoic acid, lactobionic acid, 3'-adenylic acid, 5'-adenylic acid, mucic acid, galactaric acid, pantothenic acid, pectic acid, polygalacturonic acid, 5-sulfosalicylic acid, 1,2,3,6-tetrahydro-1,3-dimethyl-2,6-dioxopurine-7-propanesulfonic acid, terephthalic acid, 1-hydroxy-2-naphthoic acid, and combinations thereof. In some embodiments, in conjunction with other above or below embodiments, pH modifying agents include, for example, maleic acid, citric acid, malic acid, fumaric acid, sulfuric acid, tartaric acid, lactic acid, salicylic acid, aspartic acid, aminosalicylic acid, malonic acid, glutamic acid, and combinations thereof. In some embodiments, in conjunction with other above or below embodiments, the pH modifying agent comprises fumaric acid, tartaric acid, glutamic acid, or a combination thereof.

[0199] In some embodiments, in conjunction with other above or below embodiments, pH modifying agent includes maleic acid, citric acid, tartaric acid, pamoic acid, fumaric acid, salicylic acid, 2,6-diaminohexanoic acid, camphorsulfonic acid, glycerophosphoric acid, 2-hydroxyethanesulfonic acid, isethionic acid, succinic acid, carbonic acid, p-toluenesulfonic acid, aspartic acid, 8-chlorotheophylline, benzenesulfonic acid, malic acid, orotic acid, oxalic acid, benzoic acid, 2-naphthalenesulfonic acid, stearic acid, adipic acid, p-amino-salicylic acid, 5-aminosalicylic acid, ascorbic acid, sulfuric acid, cyclamic acid, sodium lauryl

sulfate, glucoheptonic acid, glucuronic acid, glycine, sulfuric acid, mandelic acid, 1,5-naphthalenedisulfonic acid, nicotinic acid, oleic acid, 2-oxoglutaric acid, pyridoxal 5-phosphate, undecanoic acid, p-acetamidobenzoic acid, o-acetamidobenzoic acid, m-acetamidobenzoic acid, N-acetyl-L-aspartic acid, camphoric acid, dehydrocholic acid, malonic acid, edetic acid, ethylenediaminetetraacetic acid, ethylsulfuric acid, hydroxyphenylbenzoylbenzoic acid, glutamic acid, glycyrrhizic acid, 4-hexylresorcinol, hippuric acid, p-phenolsulfonic acid, 4-hydroxybenzoic acid, 3-hydroxybenzoic acid, 3-hydroxy-2-naphthoic acid, 1-hydroxy-2-naphthoic acid, lactobionic acid, 3'-adenylic acid, 5'-adenylic acid, mucic acid, galactaric acid, pantothenic acid, pectic acid, polygalacturonic acid, 5-sulfosalicylic acid, 1,2,3,6-tetrahydro-1,3-dimethyl-2,6-dioxopurine-7-propanesulfonic acid, terephthalic acid, 1-hydroxy-2-naphthoic acid, and combinations thereof.

[0200] In some embodiments, in conjunction with other above or below embodiments, the pH modifying agent is selected from maleic acid, citric acid, malic acid, fumaric acid, sulfuric acid, tartaric acid, lactic acid, salicylic acid, aspartic acid, aminosalicic acid, malonic acid, glutamic acid, and any combination thereof.

[0201] In some embodiments, in conjunction with other above or below embodiments, fumaric acid is used as the pH modifying agent as it is less hygroscopic and more compatible with omecamtiv mecarbil dihydrochloride hydrate than citric acid, resulting in less or no active form transformation and no changes in tablet appearance when stored at about 40 °C/75%RH for about 6 months, leading to improved final product quality. Additionally, fumaric acid is more acidic (by about 2-fold) than citric acid. Therefore, it is more efficient, i.e., about 1:1 weight ratio to active instead of about 2:1, to use fumaric acid to modulate the microenvironmental pH to enhance omecamtiv mecarbil release at neutral environment. Fumaric acid also has a very slow dissolution rate. As a result, fumaric acid will stay in the tablet longer and maintain the low micro-environmental pH better, resulting in more complete release of omecamtiv mecarbil within about 24 hours.

[0202] As used herein, the term “fillers” refers to one or more substances that can be added to components of a pharmaceutical composition to increase bulk weight of the material to be formulated, e.g. tableted, in order to achieve the desired weight. Fillers include but are not limited to starches, lactose, mannitol (such as Pearlitol™ SD 200), cellulose derivatives, calcium phosphate, sugar and the like.

[0203] Different grades of lactose include, but are not limited, to lactose monohydrate, lactose DT (direct tableting), lactose anhydrous, Flowlac™ (available from Meggle products), Pharmatose™ (available from DMV) and others. Different grades of starches include, but are not limited to, maize starch, potato starch, rice starch, wheat starch, pregelatinized starch

(commercially available as PCS PC10 from Signet Chemical Corporation) and Starch 1500, Starch 1500 LM grade (low moisture content grade) from Colorcon, fully pregelatinized starch (commercially available as National 78-1551 from Essex Grain Products) and others. Different cellulose compounds that can be used include crystalline cellulose and powdered cellulose. Examples of crystalline cellulose products include but are not limited to CEOLUS™ KG801, Avicel™ PH 101, PH102, PH301, PH302 and PH-F20, microcrystalline cellulose 114, and microcrystalline cellulose 112. Other useful fillers include, but are not limited to, carmellose, sugar alcohols such as mannitol, sorbitol and xylitol, calcium carbonate, magnesium carbonate, dibasic calcium phosphate, and tribasic calcium phosphate.

[0204] In some embodiments, in conjunction with other above or below embodiments, the filler is selected from starch, lactose, mannitol (such as Pearlitol™ SD 200), cellulose derivatives, calcium phosphate, and a sugar.

[0205] In some embodiments, in conjunction with other above or below embodiments, the filler is lactose anhydrous or lactose monohydrate. In some embodiments, in conjunction with other above or below embodiments, the filler is lactose DT, Flowlac™, or Pharmatose™.

[0206] In some embodiments, in conjunction with other above or below embodiments, the filler is maize starch, potato starch, rice starch, wheat starch, pregelatinized starch (such as Starch 1500 or Starch 1500 LM grade (low moisture content grade)), or fully pregelatinized starch.

[0207] In some embodiments, in conjunction with other above or below embodiments, the filler is microcrystalline cellulose, such as CEOLUS™ KG801, Avicel™ PH 101, PH102, PH301, PH302 and PH-F20, microcrystalline cellulose 114, or microcrystalline cellulose 112.

[0208] In In some embodiments, in conjunction with other above or below embodiments, the filler is carmellose, mannitol, sorbitol, xylitol, calcium carbonate, magnesium carbonate, dibasic calcium phosphate, or tribasic calcium phosphate.

[0209] As used herein, the term “lubricants” refers to one or more substances that can be added to components of the present compositions to reduce sticking by a solid formulation to the equipment used for production of a unit dose form. Lubricants include stearic acid, hydrogenated vegetable oils, hydrogenated soybean oil and hydrogenated soybean oil & castor wax, stearyl alcohol, leucine, polyethylene glycol, magnesium stearate, glycerylmonostearate, stearic acid, glycerylbehenate, polyethylene glycol, ethylene oxide polymers, sodium lauryl sulfate, magnesium lauryl sulfate, sodium oleate, sodium stearyl fumarate, DL-leucine, colloidal silica, and mixtures thereof.

[0210] In some embodiments, in conjunction with other above or below embodiments, the lubricant is stearic acid, hydrogenated vegetable oil, hydrogenated soybean oil, hydrogenated soybean oil, castor wax, stearyl alcohol, leucine, polyethylene glycol, magnesium stearate, glycerylmonostearate, stearic acid, glyceryl behenate, polyethylene glycol, ethylene oxide polymers, sodium lauryl sulfate, magnesium lauryl sulfate, sodium oleate, sodium stearyl fumarate, DL-leucine, colloidal silica, or any mixture thereof.

[0211] Swellable Core Formulations

[0212] Provided herein are pharmaceutical formulations comprising: a drug layer comprising omecamtiv mecarbil as a salt or crystal form as disclosed herein; a sweller layer; and a semi-permeable membrane coating having at least one delivery port.

[0213] Further provided are formulations comprising

a drug layer comprising:

omecamtiv mecarbil as a salt or crystal form as disclosed herein;

a drug layer polymer; and

a lubricant;

a sweller layer comprising:

a sweller layer polymer;

an osmotic agent;

a diluent; and

a lubricant; and

a semi-permeable membrane coating having at least one delivery port comprising:

an insoluble polymer; and

a pore forming polymer.

[0214] In some embodiments, in conjunction with other above or below embodiments, the pharmaceutical formulation comprises:

a drug layer comprising: 10-20 (w/w%) omecamtiv mecarbil salt or crystal form; 40-60 (w/w%) polyethylene oxide; and 0-2% (w/w%) lubricant;

a sweller layer comprising: 12-30 (w/w%) polyethylene oxide; 2-10 (w/w%) an osmotic agent; 1-8 (w/w%) microcrystalline cellulose; and 0.1-2 (w/w%) lubricant; and

a semi-permeable membrane having at least one delivery port comprising: 5-15 (w/w%) cellulose acetate; 0.3-5 (w/w%) polyethylene glycol.

[0215] In some embodiments, in conjunction with other above or below embodiments, the pharmaceutical formulation comprises:

a drug layer comprising: 14-17 (w/w%) omecamtiv mecarbil salt or crystal form; 48-55 (w/w%) polyethylene oxide; and 0.1-0.5% (w/w%) lubricant;

a sweller layer comprising: 18-25 (w/w%) polyethylene oxide; 4-9 (w/w%) an osmotic agent; 3-6 (w/w%) microcrystalline cellulose; and 0.1-0.5 (w/w%) lubricant; and

a semi-permeable membrane having at least one delivery port comprising: 8-10 (w/w%) cellulose acetate; 0.5-3 (w/w%) polyethylene glycol.

[0216] In some embodiments, in conjunction with other above or below embodiments, the pharmaceutical formulation comprises:

a drug layer comprising: 10-20 (w/w%) omecamtiv mecarbil salt or crystal form; 40-60 (w/w%) polyethylene oxide; and 0-2% (w/w%) magnesium stearate;

a sweller layer comprising: 12-30 (w/w%) polyethylene oxide; 2-10 (w/w%) sodium chloride; 1-8 (w/w%) microcrystalline cellulose; and 0.1-2 (w/w%) magnesium stearate; and

a semi-permeable membrane having at least one delivery port comprising: 5-15 (w/w%) cellulose acetate; 0.3-5 (w/w%) polyethylene glycol 3350.

[0217] In some embodiments, in conjunction with other above or below embodiments, the pharmaceutical formulation comprises:

a drug layer comprising: 15-16 (w/w%) omecamtiv mecarbil salt or crystal form; 50-52 (w/w%) PolyOxTM WSR N-80; and 0.1-0.5% (w/w%) magnesium stearate;

a sweller layer comprising: 20-23 (w/w%) PolyOxTM WSR Coagulant; 4-9 (w/w%) sodium chloride; 3-6 (w/w%) Avicel PH 200; and 0.1-0.5 (w/w%) lubricant; and

a semi-permeable membrane having at least one delivery port comprising: 8-10 (w/w%) cellulose acetate; 0.5-3 (w/w%) polyethylene glycol 3350.

[0218] In some embodiments, in conjunction with other above or below embodiments, the pharmaceutical formulation comprises:

a drug layer comprising: 15-16 (w/w%) omecamtiv mecarbil salt or crystal form; 50-52 (w/w%) PolyOxTM WSR N-80; and 0.1-0.5% (w/w%) magnesium stearate;

a sweller layer comprising: 20-23 (w/w%) PolyOxTM WSR Coagulant; 4-9 (w/w%) sodium chloride; 3-6 (w/w%) Avicel PH 200; and 0.1-0.5 (w/w%) lubricant; and

a semi-permeable membrane having at least one delivery port comprising: 8-9 (w/w%) cellulose acetate; 2-3 (w/w%) polyethylene glycol 3350.

[0219] In some embodiments, in conjunction with other above or below embodiments, the pharmaceutical formulation comprises:

a drug layer comprising: 15-16 (w/w%) omecamtiv mecarbil salt or crystal form; 50-52 (w/w%) PolyOxTM WSR N-80; and 0.1-0.5% (w/w%) magnesium stearate;

a sweller layer comprising: 20-23 (w/w%) PolyOxTM WSR Coagulant; 4-9 (w/w%) sodium chloride; 3-6 (w/w%) Avicel PH 200; and 0.1-0.5 (w/w%) lubricant; and

a semi-permeable membrane having at least one delivery port comprising: 9-10

(w/w%) cellulose acetate; 0.5-2 (w/w%) polyethylene glycol 3350.

Methods of Use

[0220] The omecamtiv mecarbil salt or crystal forms disclosed herein, or the pharmaceutical compositions described herein, may be used in the treatment or prevention of heart failure, including but not limited to: acute (or decompensated) congestive heart failure, and chronic congestive heart failure; particularly diseases associated with systolic heart dysfunction.

[0221] Also provided herein are methods of treating or preventing heart failure in a subject in need thereof comprising administering to the subject one or more of the omecamtiv mecarbil salt or crystal forms disclosed herein, or one or more of the pharmaceutical compositions described herein in an amount effective to treat or prevent heart failure. Further provided are methods for the use of the disclosed salts and crystal forms of omecamtiv mecarbil, or compositions thereof, for the treatment or prevention of heart failure, including but not limited to: acute (or decompensated) congestive heart failure, and chronic congestive heart failure; particularly diseases associated with systolic heart dysfunction.

[0222] Also provided herein is the use of the omecamtiv mecarbil salt or crystal forms disclosed herein, or the pharmaceutical compositions described herein, in the manufacture of a medicament for the treatment or prevention of heart failure. In some embodiments, the present disclosure provides use of the omecamtiv mecarbil salt or crystal forms disclosed herein, or the pharmaceutical compositions described herein, in the manufacture of a medicament for the treatment of acute (or decompensated) congestive heart failure, and chronic congestive heart failure; particularly diseases associated with systolic heart dysfunction.

[0223] "Treatment" or "treating" includes one or more of : a) inhibiting a disease or disorder; b) slowing or arresting the development of clinical symptoms of a disease or disorder; and/or c) relieving a disease or disorder, that is, causing the regression of clinical symptoms. The term covers both complete and partial reduction of the condition or disorder, and complete or partial reduction of clinical symptoms of a disease or disorder. Thus, the omecamtiv mecarbil salt or crystal forms described herein, or the pharmaceutical compositions described herein may prevent an existing disease or disorder from worsening, assist in the management of the disease or disorder, or reduce or eliminate the disease or disorder. "Prevention," that is, causing the clinical symptoms of the disease or disorder not to develop, includes the prophylactic administration of a pharmaceutical formulation described herein to a subject (i.e., an animal, preferably a mammal, most preferably a human) believed to be in need of preventative treatment, such as, for example, chronic heart failure.

EXAMPLES

Methods

X-ray Powder Diffraction

[0224] Procedure A: XRPD analysis was carried out on a PANalytical X'pert pro, scanning the samples between 3 and 35° 2 θ . The material was gently ground to release any agglomerates and loaded onto a multi-well plate with Kapton or Mylar polymer film to support the sample. The multi-well plate was then placed into the diffractometer and analyzed using Cu K radiation ($\alpha_1\lambda = 1.54060 \text{ \AA}$; $\alpha_2 = 1.54443 \text{ \AA}$; $\beta = 1.39225 \text{ \AA}$; $\alpha_1 : \alpha_2$ ratio = 0.5) running in transmission mode (step size 0.0130° 2 θ) using 40 kV / 40 mA generator settings.

[0225] Procedure B: XRPD analysis was carried out on a PANalytical X'pert pro, scanning the samples between 3 and 40° 2 θ . The material was loaded onto a zero background sample holder then placed into the diffractometer on a spinning stage at 1 rotation per second and analyzed using Cu K radiation ($\alpha_1\lambda = 1.54060 \text{ \AA}$; $\alpha_2 = 1.54443 \text{ \AA}$; $\beta = 1.39225 \text{ \AA}$; $\alpha_1 : \alpha_2$ ratio = 0.5) running in transmission mode (step size 0.0167° 2 θ) using 45 kV / 40 mA generator settings.

[0226] Procedure C: X-Ray Powder Diffraction patterns were collected on a Bruker AXS C2 GADDS diffractometer using Cu K α radiation (40 kV, 40 mA), automated XYZ stage, laser video microscope for auto-sample positioning and a HiStar 2-dimensional area detector. X-ray optics consists of a single GObel multilayer mirror coupled with a pinhole collimator of 0.3 mm. The beam divergence, i.e. the effective size of the X-ray beam on the sample, was approximately 4 mm. A θ - θ continuous scan mode was employed with a sample — detector distance of 20 cm which gives an effective 2 θ range of 3.2°-29.7°. Typically the sample would be exposed to the X-ray beam for 120 seconds. Ambient conditions. Samples run under ambient conditions were prepared as flat plate specimens using powder as received with and without grinding. Approximately 1-2 mg of the sample was lightly pressed on a glass slide to obtain a flat surface.

Differential Scanning Calorimetry (DSC)

[0227] Procedure A: Thermal properties were characterized using a DSC Q1000 or DSC Q100 model, TA Instruments, differential scanning calorimetry and a Q500, TA Instruments, thermogravimetric analyzer. Data analysis was performed utilizing Universal Analysis 2000, TA Instruments. Heating rates of 1, 10 and 100 °C/min were used over a variety of temperature ranges for differential scanning calorimetry and thermogravimetric analysis. Samples ranging from 1-5 mg were prepared in crimped, hermetic or open aluminum pans for DSC analysis.

[0228] Procedure B: DSC data were collected on a TA Instruments Q2000 equipped with a 50 position autosampler. The instrument was calibrated for energy and temperature calibration using certified indium. Typically 0.5-3 mg of each sample, in a pin-holed aluminum pan, was heated at 10 °C/min from 25 °C to 350 °C. A nitrogen purge at 50 ml/min was maintained over the sample. The instrument control software was Thermal Advantage v4.6.6 and the data were analyzed using Universal Analysis v4.3A.

Thermal Gravimetric/Differential Thermal Analysis (TG/DTA)

[0229] Approximately 5 mg of material was weighed into an open aluminum pan and loaded into a simultaneous thermogravimetric/differential thermal analyzer (TG/DTA) and held at room temperature. The sample was then heated at a rate of 10 °C/min from 20 °C to 300 °C during which time the change in sample weight was recorded along with any differential thermal events (DTA). Nitrogen was used as the purge gas at a flow rate of 300 cm³/min.

Thermal Gravimetric Analysis

[0230] Procedure A: Thermograms were collected on a TA Instruments Q500 thermogravimetric analyzer. Samples were loaded onto a platinum pan, 1-10 mg, and heated at 10 °C/min from ambient to 300 °C.

[0231] Procedure B: TGA data were collected on a TA Instruments Q500 TGA, equipped with a 16 position auto-sampler. The instrument was temperature calibrated using certified Alumel. Typically 5-30 mg of each sample was loaded onto a pre-tared platinum crucible and aluminum DSC pan, and was heated at 10 °C/min from ambient temperature to 350 °C. A nitrogen purge at 60 ml/min was maintained over the sample. The instrument control software was Thermal Advantage v4.6.6 and the data were analyzed using Universal Analysis v4.3A.

Moisture Sorption

[0232] Moisture balance was collected using a dynamic vapor sorption (DVS) analyzer. Relative humidity (RH) was set to 0, 5, 15, 25, 35, 45, 55, 65, 75, 85 and 95% RH for 2 sorption/desorption cycles at 25 °C. Equilibration criteria was set at 0.001% weight. Approximately 10 mg of sample was used.

Solubility

[0233] An excess of solid was added to water, pH 1.0 or pH 4.5 buffer to produce a suspension and dispersed for at least 24 at room temperature. Suspensions were filtered.

Filtrate was analyzed by HPLC-UV and compared against a standard curve to determine the solution concentration of the crystal form.

Experimental Section

[0234] Free base crystalline form III: Procedure A - prepared by adding 429 mg of omecamtiv mecarbil to 20 mL of 2-propanol and 20 mL of water then heating to 50 °C to dissolve then precipitate with 200 mL of water.

[0235] Procedure B - prepared during a solubility screen from 7 different solvents (2-BuOAc slurry, Cumene slurry, Isopropyl acetate slurry, MTBE slurry, heptane slurry, tBuOAc slurry or toluene slurry). 10 mg of free base was placed in a vial and 50 μ L aliquots for the first 300 μ L, 100 μ L thereafter (up to 1 mL), of the solvents were added to the vial. Between each addition the mixture was checked for dissolution and if no dissolution was apparent the mixture was heated to ca. 50 °C and checked again. This procedure was continued until dissolution was observed or until 100 volumes of solvent had been added. If no dissolution occurred the solid was filtered and an XRPD collected. If dissolution occurred, the cap was removed to allow evaporation of the solvent and an XRPD of the remaining solid was collected. Free base crystalline form III was then reproduced from a slurry in cumene, isopropyl acetate, MTBE or tBuOAc by adding 60 mg of lyophilized free base to the solvent and gently heating to about 40 °C.

[0236] The omecamtiv mecarbil free base crystalline form III was characterized by an XRPD pattern comprising peaks in Table 1.

[0237] Table 1

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
7.91	0.07	11.17	533.03	2.67
9.50	0.08	9.31	3776.76	18.91
14.27	0.10	6.21	15559.92	77.89
15.25	0.08	5.81	1095.46	5.48
16.10	0.10	5.51	1128.65	5.65
17.78	0.10	4.99	19975.52	100.00
19.06	0.10	4.66	2511.63	12.57
19.64	0.20	4.52	143.81	0.72
20.65	0.13	4.30	421.10	2.11
21.24	0.15	4.18	293.88	1.47
23.01	0.10	3.87	1285.85	6.44
23.87	0.10	3.73	1434.94	7.18
24.83	0.13	3.59	201.27	1.01
25.59	0.10	3.48	226.37	1.13
26.23	0.20	3.40	166.45	0.83

27.23	0.20	3.28	89.66	0.45
28.11	0.15	3.17	805.44	4.03
28.80	0.13	3.10	362.99	1.82
31.01	0.17	2.88	701.54	3.51
31.95	0.12	2.80	844.68	4.23
32.34	0.20	2.77	605.43	3.03
33.67	0.13	2.66	248.53	1.24
34.69	0.13	2.59	293.11	1.47
37.26	0.13	2.41	245.07	1.23
38.73	0.23	2.33	152.34	0.76

[0238] Free base crystalline form IV: was prepared by precipitation of 46 mg of omecamtiv mecarbil from 2 mL THF with 2 mL n-butyl ether.

[0239] The omecamtiv mecarbil free base crystalline form IV was characterized by an XRPD pattern comprising peaks in Table 2.

[0240] Table 2

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
5.18	0.07	17.04	12067.81	100.00
7.42	0.10	11.91	322.11	2.67
7.70	0.08	11.48	416.99	3.46
10.35	0.07	8.55	814.51	6.75
10.81	0.10	8.19	170.45	1.41
14.21	0.08	6.23	789.68	6.54
14.84	0.07	5.97	1276.45	10.58
15.54	0.08	5.70	2686.48	22.26
17.10	0.10	5.18	122.93	1.02
18.10	0.10	4.90	10344.17	85.72
19.92	0.08	4.46	939.94	7.79
20.62	0.07	4.31	802.71	6.65
20.77	0.07	4.28	931.21	7.72
21.70	0.08	4.10	481.64	3.99
22.40	0.12	3.97	418.41	3.47
22.86	0.10	3.89	673.49	5.58
23.09	0.08	3.85	571.53	4.74
24.05	0.08	3.70	774.58	6.42
24.36	0.12	3.65	752.02	6.23
25.20	0.10	3.53	582.91	4.83
25.72	0.07	3.46	378.21	3.13
26.00	0.10	3.43	138.03	1.14
26.68	0.27	3.34	93.44	0.77
27.40	0.10	3.26	400.49	3.32

27.81	0.12	3.21	724.88	6.01
28.18	0.10	3.17	278.71	2.31
28.63	0.08	3.12	399.78	3.31
28.98	0.08	3.08	480.22	3.98
29.42	0.10	3.04	644.30	5.34
30.51	0.13	2.93	247.42	2.05
32.80	0.13	2.73	137.46	1.14
33.98	0.20	2.64	155.02	1.28
35.34	0.20	2.54	147.52	1.22
36.38	0.20	2.47	110.92	0.92
37.12	0.20	2.42	126.42	1.05

[0241] Free base crystalline form V: Procedure A - was prepared by adding 50 mg of omecamtiv mecarbil to 2 mL of THF at 60°C, filtering, then crash cooling by placing sample into an acetone/dry ice bath.

[0242] Procedure B - Free base crystalline form V was prepared during a solubility screen from 1,4-dioxane slurry. 10 mg of free base was placed in a Vial and 50 μ L aliquots for the first 300 μ L, 100 μ L thereafter (up to 1 mL), of the solvent was added to the vial. Between each addition the mixture was checked for dissolution and if no dissolution was apparent the mixture was heated to ca. 50 °C and checked again. This procedure was continued until dissolution was observed or until 100 volumes of solvent had been added. If no dissolution occurred the solid was filtered and an XRPD collected. If dissolution occurred, the cap was removed to allow evaporation of the solvent and an XRPD of the remaining solid was collected.

[0243] The omecamtiv mecarbil free base crystalline form V was characterized by an XRPD pattern comprising peaks in Table 3.

[0244] Table 3

Pos. [°2 θ .]	FWHM [°2 θ .]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
5.40	0.10	16.37	213.79	7.88
7.38	0.07	11.98	470.25	17.33
8.56	0.07	10.33	430.26	15.86
8.93	0.08	9.91	563.71	20.78
9.14	0.07	9.67	392.66	14.47
10.03	0.08	8.82	375.08	13.82
10.73	0.13	8.25	332.70	12.26
11.71	0.12	7.56	840.50	30.98
13.69	0.17	6.47	335.68	12.37
15.08	0.18	5.88	653.88	24.10
16.04	0.33	5.52	105.31	3.88

16.85	0.08	5.26	920.36	33.92
17.85	0.13	4.97	2713.30	100.00
18.28	0.10	4.85	347.67	12.81
18.86	0.12	4.71	1897.24	69.92
20.05	0.20	4.43	1456.16	53.67
20.72	0.07	4.29	659.49	24.31
21.74	0.17	4.09	354.74	13.07
22.83	0.27	3.90	189.17	6.97
23.56	0.10	3.78	1152.90	42.49
24.03	0.20	3.70	645.49	23.79
25.45	0.20	3.50	217.43	8.01
26.23	0.10	3.40	366.63	13.51
27.62	0.23	3.23	654.84	24.13
28.58	0.23	3.12	253.96	9.36
29.85	0.27	2.99	254.72	9.39
32.10	0.20	2.79	233.50	8.61
33.37	0.33	2.69	165.44	6.10
35.47	0.27	2.53	105.15	3.88
37.02	0.40	2.43	122.14	4.50

[0245] Free base crystalline form VI: was prepared by heating free base crystalline form V to 150°C.

[0246] The omecamtiv mecarbil free base crystalline form VI was characterized by an XRPD pattern comprising peaks in Table 4.

[0247] Table 4

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
9.07	0.08	9.75	895.54	31.71
14.28	0.13	6.20	147.01	5.20
15.19	0.20	5.83	42.70	1.51
15.88	0.12	5.58	539.11	19.09
16.67	0.08	5.32	796.74	28.21
17.81	0.08	4.98	1094.93	38.77
18.18	0.17	4.88	2824.51	100.00
18.80	0.12	4.72	850.75	30.12
19.70	0.10	4.51	1045.62	37.02
20.23	0.13	4.39	185.58	6.57
20.89	0.10	4.25	1310.32	46.39
21.28	0.12	4.18	925.51	32.77
23.72	0.08	3.75	691.60	24.49
24.26	0.08	3.67	784.97	27.79
26.19	0.20	3.40	146.84	5.20

26.80	0.17	3.33	235.58	8.34
27.59	0.13	3.23	347.69	12.31
28.90	0.33	3.09	163.33	5.78
29.82	0.12	3.00	368.54	13.05

[0248] Free base crystalline form VII: was prepared during a solubility screen from water slurry. 10 mg of free base was placed in a vial and 50 μ L aliquots for the first 300 μ L, 100 μ L thereafter (up to 1 mL), of water was added to the vial. Between each addition the mixture was checked for dissolution and if no dissolution was apparent the mixture was heated to ca. 50 °C and checked again. This procedure was continued until dissolution was observed or until 100 volumes of solvent had been added. The solid was filtered and an XRPD collected.

[0249] The omecamtiv mecarbil free base crystalline form VII was characterized by an XRPD pattern comprising peaks in Table 5.

[0250] Table 5

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
4.37	0.08	20.24	4137.34	100.00
7.84	0.13	11.27	89.35	2.16
8.40	0.06	10.52	285.34	6.90
8.71	0.08	10.15	172.30	4.16
10.81	0.15	8.19	74.73	1.81
13.08	0.08	6.77	387.47	9.37
15.66	0.08	5.66	481.77	11.64
16.83	0.10	5.27	1596.66	38.59
18.92	0.10	4.69	1248.28	30.17
19.61	0.09	4.53	1696.18	41.00
20.32	0.05	4.37	584.96	14.14
20.49	0.10	4.33	961.99	23.25
21.61	0.08	4.11	319.30	7.72
22.26	0.10	3.99	1149.06	27.77
23.22	0.08	3.83	265.13	6.41
23.46	0.10	3.79	299.05	7.23
24.21	0.13	3.68	1205.74	29.14
25.41	0.05	3.51	1016.23	24.56
27.58	0.13	3.23	318.93	7.71
29.53	0.15	3.02	279.27	6.75
30.13	0.15	2.97	221.71	5.36
31.32	0.61	2.86	95.23	2.30

[0251] The XRPD peaks unique to each of the free base crystalline forms III-VII disclosed herein are shown in Table 6.

[0252] Table 6

Free Base Crystalline Form	Peaks Unique to Each Form (KA1 °)					
Form III	9.50	19.06	23.01			
Form IV	5.18	10.35	14.84	15.54	18.10	19.92
Form V	7.38	8.56	9.14	18.28		
Form VI	9.07	16.67	18.18	19.70	20.89	21.28
Form VII	8.40	8.71	13.08	15.66	19.61	

[0253] **Amorphous hydrochloride salt:** An amorphous bis-hydrochloride salt was prepared by dissolving 0.505g of bis-hydrochloride monohydrate salt Form A in 20 mL of water, flash frozen in liquid nitrogen and lyophilized. Chloride analysis gave a result of 14.6% Cl which is in agreement with a bis-hydrochloride. Thermal analysis indicates approximately a 7.9% weight loss during heating due to loss of water and a Tg around 149 °C. Vapor sorption studies show that the amorphous form is hygroscopic and converts to the crystalline bis-hydrochloride monohydrate Form A.

[0254] **Ethane Sulfonate crystalline salt:** was prepared in a primary salt screen. A 2 mL aliquot of acetone was added to ~40 mg of free base. 1.05 equivalents of ethane sulfonate as a 1M solution in THF were added and the sample was temperature cycled for 3-5 days.

[0255] The omecamtiv mecarbil ethane sulfonate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 7.

[0256] Table 7

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
8.61	0.10	10.27	1864.81	100.00
13.61	0.15	6.51	141.09	7.57
16.14	0.10	5.49	1152.97	61.83
16.76	0.13	5.29	825.27	44.26
16.97	0.08	5.22	914.05	49.02
17.23	0.13	5.15	1148.35	61.58
18.35	0.15	4.84	579.44	31.07
19.20	0.20	4.62	710.28	38.09
20.27	0.20	4.38	781.89	41.93
20.73	0.08	4.28	1275.35	68.39

20.96	0.10	4.24	1275.82	68.42
23.73	0.20	3.75	494.48	26.52
25.24	0.15	3.53	262.30	14.07
25.95	0.15	3.43	442.47	23.73
26.30	0.10	3.39	512.57	27.49
27.09	0.20	3.29	204.40	10.96
29.01	0.41	3.08	74.97	4.02

[0257] Bis-fumarate crystalline salt form A: was prepared by dissolving 1 equivalent of free base (4.104 g) and 2.1 equivalents of fumaric acid (as 84 mL in a 95% EtOH/water solution) in 20 mL of 90% THF in water then distilling under house vacuum at 50°C. An additional 92 mL of water and mono-fumarate Form A seed were added to induce precipitation.

[0258] The omecamtiv mecarbil bis-fumarate crystalline salt form A was characterized by an XRPD pattern comprising peaks in Table 8.

[0259] Table 8

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
5.64	0.18	15.67	1058.97	74.12
8.83	0.27	10.01	92.99	6.51
10.80	0.17	8.20	144.21	10.09
12.04	0.20	7.35	271.90	19.03
15.76	0.17	5.62	249.66	17.47
16.40	0.10	5.41	417.23	29.20
16.80	0.13	5.28	1428.74	100.00
17.94	0.20	4.94	167.95	11.76
18.32	0.13	4.84	247.68	17.34
19.38	0.17	4.58	130.62	9.14
19.87	0.13	4.47	159.85	11.19
20.61	0.27	4.31	198.11	13.87
21.55	0.13	4.12	445.40	31.17
21.87	0.13	4.06	589.24	41.24
22.03	0.13	4.04	583.89	40.87
22.88	0.13	3.89	300.01	21.00
23.61	0.10	3.77	1008.70	70.60
23.87	0.13	3.73	1042.62	72.97
25.07	0.27	3.55	109.91	7.69
26.01	0.18	3.43	795.76	55.70
27.20	0.27	3.28	900.61	63.04
27.86	0.23	3.20	317.54	22.23
29.55	0.27	3.02	61.88	4.33
31.04	0.20	2.88	103.03	7.21

32.73	0.27	2.74	145.12	10.16
35.00	0.20	2.56	80.10	5.61
36.01	0.27	2.49	62.22	4.35
36.54	0.23	2.46	138.74	9.71

[0260] Bis-fumarate crystalline salt form B: was prepared during DVS cycle of bis-fumarate Form A.

[0261] The omecamtiv mecarbil bis-fumarate crystalline salt form B was characterized by an XRPD pattern comprising peaks in Table 9.

[0262] Table 9

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
5.68	0.20	15.57	557.86	53.41
6.11	0.15	14.47	941.86	90.18
9.69	0.07	9.13	213.36	20.43
11.43	0.17	7.74	195.35	18.70
12.92	0.10	6.85	428.18	41.00
13.13	0.08	6.74	739.41	70.80
14.34	0.40	6.18	52.05	4.98
15.95	0.20	5.56	656.27	62.84
16.83	0.23	5.27	142.59	13.65
17.22	0.20	5.15	178.09	17.05
18.08	0.20	4.91	494.57	47.35
19.05	0.20	4.66	135.39	12.96
19.52	0.33	4.55	99.22	9.50
20.81	0.20	4.27	362.18	34.68
22.47	0.15	3.96	547.41	52.41
22.95	0.27	3.87	553.11	52.96
24.53	0.20	3.63	147.89	14.16
26.04	0.20	3.42	434.11	41.57
27.01	0.13	3.30	1044.39	100.00
28.43	0.13	3.14	313.18	29.99
31.37	0.27	2.85	59.28	5.68
32.32	0.23	2.77	49.90	4.78
34.89	0.40	2.57	65.58	6.28
35.89	0.23	2.50	54.73	5.24
37.16	0.20	2.42	68.51	6.56

[0263] Bis-fumarate crystalline salt form C: was prepared from slurry of bis-fumarate Forms A and B in 2 mL water at RT. Water solubility was determined to be 13.7 mg/mL (pH 3.1).

[0264] The omecamtiv mecarbil bis-fumarate crystalline salt form C was characterized by an XRPD pattern comprising peaks in Table 10.

[0265] Table 10

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
5.88	0.17	15.03	5167.95	100.00
10.38	0.54	8.53	75.90	1.47
10.70	0.10	8.27	162.04	3.14
11.56	0.12	7.65	325.84	6.30
12.74	0.10	6.95	1330.99	25.75
13.56	0.12	6.53	529.65	10.25
15.33	0.13	5.78	181.25	3.51
16.85	0.10	5.26	336.53	6.51
17.15	0.12	5.17	1259.92	24.38
17.63	0.10	5.03	864.50	16.73
18.79	0.12	4.72	557.14	10.78
19.52	0.12	4.55	229.88	4.45
20.29	0.12	4.38	906.12	17.53
20.60	0.12	4.31	510.74	9.88
20.86	0.10	4.26	460.43	8.91
21.47	0.07	4.14	668.33	12.93
21.77	0.08	4.08	697.87	13.50
22.21	0.13	4.00	865.69	16.75
22.92	0.13	3.88	864.48	16.73
23.58	0.13	3.77	792.02	15.33
24.15	0.07	3.68	600.60	11.62
24.55	0.13	3.63	373.42	7.23
25.41	0.17	3.51	1064.69	20.60
26.37	0.12	3.38	254.11	4.92
26.78	0.12	3.33	1601.63	30.99
26.86	0.06	3.33	1640.39	31.74
27.31	0.12	3.26	450.94	8.73
27.83	0.16	3.20	618.62	11.97
28.97	0.06	3.08	340.41	6.59
29.93	0.24	2.98	283.61	5.49
30.37	0.24	2.94	102.99	1.99
31.97	0.20	2.80	88.93	1.72
32.37	0.29	2.76	67.32	1.30
33.10	0.16	2.70	128.77	2.49
34.02	0.33	2.63	173.14	3.35
35.76	0.29	2.51	160.08	3.10
38.14	0.24	2.36	76.60	1.48
38.88	0.33	2.31	187.08	3.62

39.55	0.20	2.28	104.38	2.02
-------	------	------	--------	------

[0266] Mono-fumarate crystalline salt form D: was first prepared during a high throughput screen by adding 0.3 mL of a 0.05M solution of free base in methanol and 0.3 mL of a fumaric acid 0.05M solution in methanol to a glass plate then solvent was evaporated then 0.4 mL of solvent (water, 0.001M HCl aq, acetone, acetonitrile or hexanes) was added and heated to 50 °C for 4 hours then evaporated.

[0267] The omecamtiv mecarbil mono-fumarate crystalline salt form D was characterized by an XRPD pattern comprising peaks in Table 11.

[0268] Table 11

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
8.01	0.09	11.04	102.83	59.94
12.11	0.11	7.31	72.99	42.55
12.67	0.13	6.99	50.43	29.40
14.46	0.13	6.12	83.35	48.59
15.20	0.19	5.83	55.38	32.28
16.01	0.19	5.54	50.34	29.34
16.57	0.11	5.35	76.69	44.71
17.04	0.14	5.20	32.05	18.69
17.63	0.16	5.03	37.88	22.08
18.48	0.16	4.80	16.35	9.53
20.02	0.11	4.43	34.71	20.23
20.51	0.22	4.33	31.27	18.23
21.75	0.16	4.09	37.60	21.92
22.86	0.19	3.89	18.19	10.60
24.25	0.13	3.67	171.54	100.00
24.97	0.19	3.57	82.03	47.82
25.84	0.13	3.45	41.01	23.91
26.17	0.16	3.41	74.19	43.25
27.10	0.13	3.29	37.05	21.60
27.97	0.19	3.19	21.24	12.38
28.61	0.19	3.12	11.11	6.48
29.21	0.16	3.06	37.43	21.82
30.69	0.25	2.91	10.74	6.26
34.70	0.50	2.59	7.97	4.64
37.41	0.19	2.40	11.12	6.48
38.48	0.38	2.34	6.46	3.77

[0269] The XRPD peaks unique to each of the fumarate crystalline salts forms A-D disclosed herein are shown in Table 12.

[0270] Table 12

Fumarate Form	Peaks Unique to Each Form (°2Th.)					
Form A	5.64	15.76	22.03	23.87		
Form B	5.68	6.11	13.13	18.08	22.47	
Form C	5.88	18.79	25.41	26.86		
Form D	8.01	15.20	20.02			

[0271] **Bis-Maleate crystalline salt form A:** Procedure A: was first prepared during a high throughput screen by adding 0.2 mL of a 0.124M solution of free base in methanol and 0.2 mL of a maleic acid 0.25M solution in methanol to a glass plate then solvent was evaporated then 0.2 mL of solvent (water, 0.001M HCl aq or acetone) was added and heated to 50°C for 4 hours then evaporated. Salt was scaled up by adding 100 mg free base to an 8 mL vial and 5 mL methanol and gently heating to dissolve. Maleic acid (2 mL of a 0.25M solution in acetone) was added at room temperature. Precipitate was isolated by filtration.

[0272] Procedure B - bis-maleate crystalline salt form A was prepared by dissolving 3.011g of free base (1 eq) and 1.828g of maleic acid (solution in 8 mL MeOH; 2.1 eq) in 45 mL methanol at 60°C then cooled to precipitate. Water solubility was determined to be 3.8 mg/mL (pH 3.7).

[0273] Procedure C - bis-maleate crystalline salt form A was also prepared in a primary salt screen. A 2 mL aliquot of 2-propanol, THF, acetonitrile, isopropyl acetate or acetone was added to ~40 mg of free base. 1.05 equivalents of maleic acid as a 1M solution in THF were added and the sample was temperature cycled for 3-5 days.

[0274] The omecamtiv mecarbil bis-maleate crystalline salt form A was characterized by an XRPD pattern comprising peaks in Table 13.

[0275] Table 13

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
3.32	0.82	26.60	267.10	3.31
4.99	0.05	17.71	482.89	5.98
6.60	0.08	13.38	216.56	2.68
6.95	0.41	12.72	56.27	0.70
7.25	0.08	12.19	198.55	2.46
9.17	0.15	9.64	73.48	0.91
9.97	0.08	8.87	7989.17	98.97
10.56	0.08	8.38	1761.58	21.82

12.96	0.09	6.83	311.81	3.86
13.25	0.09	6.68	1227.24	15.20
14.54	0.12	6.09	356.32	4.41
14.83	0.08	5.98	743.84	9.21
15.31	0.10	5.79	8072.29	100.00
15.53	0.09	5.71	2487.57	30.82
16.04	0.10	5.52	6086.12	75.40
16.38	0.10	5.41	2253.00	27.91
17.10	0.08	5.19	675.31	8.37
17.44	0.12	5.08	3071.51	38.05
17.70	0.13	5.01	2667.01	33.04
18.17	0.10	4.88	932.80	11.56
18.45	0.10	4.81	273.07	3.38
19.00	0.13	4.67	2099.18	26.00
19.33	0.08	4.59	392.68	4.86
20.13	0.14	4.41	1667.83	20.66
21.47	0.12	4.14	2989.90	37.04
21.83	0.08	4.07	1329.81	16.47
22.02	0.06	4.04	747.10	9.26
22.31	0.10	3.98	822.83	10.19
22.44	0.09	3.96	1010.70	12.52
23.02	0.10	3.86	347.42	4.30
23.15	0.10	3.84	372.20	4.61
23.58	0.13	3.77	194.06	2.40
24.38	0.06	3.65	1003.21	12.43
24.64	0.09	3.61	876.08	10.85
25.66	0.15	3.47	924.66	11.45
26.66	0.13	3.34	1042.71	12.92
26.96	0.13	3.31	4976.38	61.65
27.83	0.13	3.21	845.81	10.48
28.10	0.10	3.18	293.57	3.64
28.55	0.15	3.13	766.92	9.50
29.31	0.08	3.05	151.67	1.88
30.17	0.20	2.96	186.07	2.31
30.76	0.23	2.91	458.79	5.68
31.67	0.13	2.83	362.78	4.49
32.01	0.13	2.80	453.36	5.62
32.25	0.13	2.78	561.46	6.96
32.67	0.18	2.74	427.72	5.30
33.11	0.20	2.71	342.20	4.24
33.65	0.13	2.66	375.70	4.65
34.17	0.10	2.62	370.55	4.59
34.39	0.10	2.61	643.90	7.98
34.51	0.15	2.60	639.10	7.92

[0276] Bis-malonate crystalline salt: was prepared by dissolving 200.7 mg of free base (1 eq) and 525 μ L of malonic acid in methanol (2.1 eq) in 5 mL methanol at 50°C then 0.5 mL IPAc was added to precipitate then heat cycled twice 40°C/RT. Water solubility was determined to be greater than 62 mg/mL (pH 3.69).

[0277] The omecamtiv mecarbil bis-malonate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 14.

[0278] Table 14

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
4.74	0.09	18.64	734.38	65.86
9.30	0.13	9.51	262.93	23.58
11.37	0.13	7.78	1115.14	100.00
13.73	0.13	6.45	154.92	13.89
14.25	0.11	6.22	902.82	80.96
15.13	0.16	5.86	567.42	50.88
15.69	0.11	5.65	94.13	8.44
16.45	0.13	5.39	438.97	39.36
16.83	0.13	5.27	165.27	14.82
18.08	0.13	4.91	552.17	49.52
18.29	0.11	4.85	855.08	76.68
18.88	0.16	4.70	287.13	25.75
19.54	0.13	4.54	232.73	20.87
20.14	0.08	4.41	819.11	73.45
20.77	0.11	4.28	300.42	26.94
21.21	0.25	4.19	192.92	17.30
23.32	0.16	3.81	218.40	19.58
23.87	0.13	3.73	766.34	68.72
24.67	0.13	3.61	193.04	17.31
25.72	0.22	3.46	82.90	7.43
26.51	0.22	3.36	211.46	18.96
27.59	0.13	3.23	520.24	46.65
27.78	0.08	3.21	805.51	72.23
28.01	0.20	3.19	630.55	56.54
28.90	0.28	3.09	273.00	24.48
29.68	0.22	3.01	33.94	3.04
30.18	0.22	2.96	51.90	4.65
33.70	0.22	2.66	96.86	8.69
34.19	0.19	2.62	54.63	4.90
35.52	0.22	2.53	39.53	3.54
36.82	0.19	2.44	42.45	3.81
37.62	0.63	2.39	36.76	3.30

[0279] Mesylate crystalline salt form A: was prepared by dissolving 200.7 mg of free base (1 eq) and 68.1 μ L of methanesulfonic acid (2.1eq) in 5 mL methanol at 50°C, 2 mL IPAc and 5 mL acetone were added to precipitate. Water solubility was determined to be greater than 72 mg/mL (pH 1.39).

[0280] The omecamtiv mecarbil mesylate crystalline salt form A was characterized by an XRPD pattern comprising peaks in Table 15.

[0281] Table 15

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
4.02	0.38	21.96	492.38	100.00
4.87	0.25	18.15	472.41	95.94
7.79	0.19	11.35	87.26	17.72
11.61	0.25	7.62	110.78	22.50
15.21	0.25	5.82	120.46	24.46
15.86	0.25	5.59	76.35	15.51
16.51	0.19	5.37	103.12	20.94
17.57	0.28	5.05	160.72	32.64
18.42	0.31	4.82	126.44	25.68
19.26	0.38	4.61	218.37	44.35
20.53	0.19	4.33	98.62	20.03
21.55	0.13	4.12	184.64	37.50
23.17	0.50	3.84	115.61	23.48
24.39	0.28	3.65	207.28	42.10
25.51	0.16	3.49	126.17	25.63
26.38	0.25	3.38	65.52	13.31
27.63	0.25	3.23	83.43	16.94
30.85	0.38	2.90	29.23	5.94

[0282] Bis-mesylate crystalline salt form B: was prepared in a primary salt screen. A 2 mL aliquot of 2-propanol was added to ~40 mg of free base. 1.05 equivalents of methanesulfonic acid as a 1M solution in THF were added and the sample was temperature cycled for 3-5 days then tert-butyl methyl ether was added as an antisolvent. Solubility in pH 1 and 4.5 buffers was determined to be greater than 23 mg/mL.

[0283] The omecamtiv mecarbil bis-mesylate crystalline salt form B was characterized by an XRPD pattern comprising peaks in Table 16.

[0284] Table 16

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
--------------	--------------	---------------	--------------	---------------

8.30	0.08	10.65	4139.68	77.69
8.94	0.06	9.89	358.87	6.73
9.59	0.08	9.22	222.95	4.18
10.78	0.08	8.21	360.97	6.77
11.15	0.05	7.94	127.57	2.39
11.66	0.09	7.59	724.49	13.60
12.15	0.09	7.28	3000.51	56.31
14.37	0.12	6.17	521.87	9.79
14.93	0.10	5.93	402.24	7.55
15.36	0.08	5.77	258.87	4.86
15.57	0.08	5.69	436.88	8.20
16.18	0.10	5.48	560.02	10.51
16.64	0.08	5.33	1864.30	34.99
16.81	0.08	5.27	744.70	13.98
17.07	0.06	5.19	2428.76	45.58
17.19	0.09	5.16	3783.32	71.00
17.41	0.10	5.09	2291.40	43.00
17.76	0.12	4.99	1630.15	30.59
19.24	0.09	4.61	757.57	14.22
19.82	0.10	4.48	1081.40	20.29
20.29	0.12	4.38	2158.58	40.51
20.66	0.14	4.30	1716.49	32.21
21.62	0.13	4.11	1236.28	23.20
22.04	0.13	4.03	739.53	13.88
22.39	0.15	3.97	5328.66	100.00
23.54	0.18	3.78	176.98	3.32
23.95	0.06	3.72	593.98	11.15
24.60	0.12	3.62	1334.05	25.04
25.02	0.13	3.56	2954.89	55.45
25.59	0.17	3.48	1527.25	28.66
25.89	0.12	3.44	581.20	10.91
26.14	0.18	3.41	329.04	6.17
26.49	0.13	3.36	171.39	3.22
27.14	0.12	3.29	637.23	11.96
27.35	0.08	3.26	1259.20	23.63
27.41	0.05	3.26	1229.76	23.08
27.89	0.19	3.20	181.37	3.40
28.86	0.09	3.09	126.53	2.37
29.45	0.16	3.03	536.88	10.08
29.89	0.25	2.99	142.78	2.68
31.11	0.31	2.87	220.88	4.15
32.47	0.25	2.76	129.04	2.42
33.10	0.19	2.70	166.58	3.13
33.51	0.25	2.67	417.91	7.84

34.56	0.16	2.59	235.62	4.42
-------	------	------	--------	------

[0285] The XRPD peaks unique to each of the mesylate crystalline salts forms A and B disclosed herein are shown in Table 17.

[0286] Table 17

Mesylate Form	Peaks Unique to Each Form (°2Th.)								
Form A	4.02	4.87	15.21	15.86	20.53	24.39			
Form B	8.30	8.94	9.59	12.15	14.37	19.82	20.29	22.04	25.02

[0287] Bis-naphthalate-2-sulfonate crystalline salt: was prepared. For the primary (small scale) screen a 2 mL aliquot of 2-propanol, THF, acetonitrile, isopropyl acetate, acetone or toluene was added to 40 mg of free base, 1.05 equivalents of naphthalate-2-sulfonate sodium salt and 1.0 eq of 1M hydrochloric acid were added then temperature cycled for 3-5 days. The salt was analyzed by XRPD. For the secondary screen (scale-up) 1.05 eq of naphthalene-2-sulfonate and 2M hydrochloric acid in THF were added to 700 mg of free base in 7 mL of 2-propanol then temperature cycled (RT to 40 C) for 3 days, filtered, dried in vacuum oven at ambient temperature. The salt was analyzed by XRPD, IR, HPLC, ¹H NMR, PLM, TG/DTA, DSC, DVS, VH-XRPD, stability studies, thermodynamic solubility studies, disproportionation studies and hydration studies. Solubility in pH 1 and 4.5 buffers was determined to be greater than 10 mg/mL.

[0288] The omecamtiv mecarbil bis-naphthalate-2-sulfonate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 18.

[0289] Table 18

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
4.49	0.06	19.70	5776.56	100.00
6.25	0.08	14.14	572.59	9.91
6.65	0.20	13.29	561.77	9.73
8.95	0.10	9.88	138.06	2.39
9.72	0.15	9.10	54.78	0.95
13.44	0.08	6.59	367.63	6.36
14.39	0.10	6.16	341.11	5.91
14.92	0.12	5.94	904.43	15.66
15.51	0.20	5.71	276.65	4.79
16.28	0.08	5.44	918.98	15.91
17.02	0.20	5.21	280.34	4.85
18.20	0.10	4.87	1575.43	27.27

18.62	0.13	4.77	3005.63	52.03
18.90	0.12	4.69	792.38	13.72
19.53	0.15	4.54	806.67	13.96
20.82	0.26	4.27	664.39	11.50
21.38	0.06	4.16	1580.84	27.37
21.52	0.13	4.13	1906.58	33.01
22.02	0.18	4.04	410.81	7.11
22.43	0.15	3.96	364.89	6.32
22.80	0.23	3.90	479.37	8.30
24.40	0.10	3.65	608.59	10.54
25.16	0.09	3.54	424.12	7.34
26.11	0.10	3.41	2521.03	43.64
27.01	0.13	3.30	388.79	6.73
27.77	0.15	3.21	210.15	3.64
29.67	0.15	3.01	314.29	5.44
30.21	0.26	2.96	233.20	4.04
30.78	0.20	2.90	266.96	4.62
31.63	0.36	2.83	418.96	7.25
33.42	0.31	2.68	268.71	4.65
34.00	0.20	2.64	207.67	3.60

[0290] Mono-napadisylate crystalline salt: was prepared during a high throughput screen by adding 0.2 mL of a 0.124M solution of free base in methanol and 0.2 mL of a 1,5-naphthalene-disulfonic acid 0.25M solution in methanol to a glass plate then solvent was evaporated then 0.2 mL of solvent (acetonitrile or 0.001M HCl aq) was added and heated to 50°C for 4 hours then evaporated. Salt was scaled up by adding 100 mg free base to an 8 mL Vial and 3 mL methanol and gently heating to dissolve. 1,5-naphthalene-disulfonic acid (2 mL of a 0.25 M solution) was added at room temperature. Solids were collected after evaporation and filtration. Water solubility was determined to be 0.3 mg/mL (pH 2.35).

[0291] The omecamtiv mecarbil mono-napadisylate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 19.

[0292] Table 19

Angle 2-Theta°	d value Angstrom	Intensity Count (Height)	Rel. Int. [%]
6.671	13.251	82.2	6.4
7.052	12.535	171	13.2
10.837	8.1642	360	27.9
12.266	7.2159	516	39.9
13.409	6.6034	442	34.2
14.572	6.0789	460	35.6

15.144	5.8503	356	27.6
15.75	5.6266	753	58.3
16.466	5.3837	788	61
17.831	4.9745	1292	100
18.824	4.7141	469	36.3
19.938	4.4533	955	73.9
21.828	4.0716	829	64.2
22.868	3.8888	605	46.9
23.487	3.7877	477	36.9
24.339	3.6571	307	23.8
25.263	3.5253	393	30.5

[0293] Nicotinate crystalline salt: was prepared in a primary salt screen. A 2 mL aliquot of THF was added to ~40 mg of free base. 1.05 equivalents of nicotinic acid were added and the sample was temperature cycled for 3-5 days then evaporated.

[0294] The omecamtiv mecarbil nicotinate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 20.

[0295] Table 20

Pos, [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
3.69	0.05	23.96	4182.86	73.17
7.36	0.06	12.01	1796.70	31.43
8.55	0.08	10.34	3192.46	55.85
9.13	0.08	9.69	3381.54	59.16
10.01	0.08	8.84	1830.40	32.02
11.70	0.10	7.56	135.02	2.36
12.43	0.09	7.12	759.00	13.28
13.83	0.08	6.40	329.03	5.76
14.74	0.10	6.01	648.74	11.35
15.04	0.15	5.89	392.48	6.87
15.50	0.12	5.72	2090.66	36.57
16.70	0.09	5.31	3962.66	69.32
16.84	0.09	5.27	5013.53	87.70
17.62	0.10	5.03	2110.97	36.93
17.87	0.12	4.96	510.00	8.92
18.30	0.12	4.85	3434.06	60.07
18.58	0.13	4.78	1509.52	26.41
18.85	0.10	4.71	353.23	6.18
19.59	0.13	4.53	2097.05	36.69
19.99	0.15	4.44	5716.36	100.00
20.34	0.17	4.37	2282.98	39.94
20.76	0.15	4.28	2868.05	50.17

21.32	0.15	4.17	2273.73	39.78
22.03	0.14	4.03	1235.06	21.61
22.91	0.13	3.88	1620.93	28.36
23.43	0.14	3.80	3577.00	62.57
23.87	0.12	3.73	895.16	15.66
24.83	0.20	3.58	3276.98	57.33
24.92	0.05	3.58	2632.73	46.06
25.40	0.16	3.50	1715.21	30.01
25.95	0.23	3.43	2961.06	51.80
26.85	0.11	3.32	1483.29	25.95
26.94	0.09	3.31	1729.54	30.26
27.32	0.14	3.26	2177.84	38.10
28.01	0.19	3.18	1526.39	26.70
28.94	0.09	3.08	887.28	15.52
29.34	0.22	3.04	154.50	2.70
29.93	0.19	2.98	217.00	3.80
31.00	0.22	2.88	319.34	5.59
31.44	0.12	2.84	286.19	5.01
31.78	0.16	2.81	214.18	3.75
32.10	0.22	2.79	385.58	6.75
32.59	0.19	2.75	320.31	5.60
32.94	0.12	2.72	259.73	4.54
33.23	0.22	2.69	207.23	3.63
34.14	0.28	2.62	300.22	5.25
34.65	0.22	2.59	283.29	4.96

[0296] Oxalate crystalline salt form A: Procedure A - was prepared a high throughput screen by adding 0.3 mL of a 0.05M solution of free base in methanol and 0.3 mL of a oxalic acid 0.05M solution in methanol to a glass plate then solvent was evaporated then 0.4 mL of solvent (water, 0.001M HCl aq, acetone, acetonitrile or hexanes) was added and heated to 50°C for 4 hours then evaporated.

[0297] Procedure B - was prepared by dissolving 47 mg of free base in ~10 mL of methanol, then 145 µL of a 102 mg/mL solution of oxalic acid was added and sample was placed in a N₂ box. Solids were then slurried in water/ethanol at RT.

[0298] The omecamtiv mecarbil oxalate crystalline salt form A was characterized by an XRPD pattern comprising peaks in Table 21.

[0299] Table 21

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
6.48	0.16	13.64	163.25	62.74

10.36	0.22	8.54	17.19	6.61
11.85	0.14	7.47	24.67	9.48
13.01	0.16	6.81	86.11	33.10
14.79	0.11	5.99	75.91	29.17
15.35	0.13	5.77	45.24	17.39
17.11	0.14	5.18	260.20	100.00
18.24	0.50	4.86	7.90	3.04
19.23	0.13	4.62	28.45	10.94
19.91	0.31	4.46	40.62	15.61
21.48	0.19	4.14	55.28	21.25
22.07	0.19	4.03	25.22	9.69
22.75	0.19	3.91	42.95	16.50
23.82	0.08	3.74	258.28	99.26
24.88	0.38	3.58	9.55	3.67
25.70	0.16	3.47	30.99	11.91
28.55	0.44	3.13	31.95	12.28
29.86	0.19	2.99	16.81	6.46
30.71	0.19	2.91	31.18	11.98
33.32	0.19	2.69	16.02	6.16

[0300] Oxalate crystalline salt form B: was prepared during vapor sorption analysis of oxalate crystalline salt form A.

[0301] The omecamtiv mecarbil oxalate crystalline salt form B was characterized by an XRPD pattern comprising peaks in Table 22.

[0302] Table 22

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
7.38	0.11	11.97	219.37	77.59
13.30	0.14	6.66	127.50	45.10
14.76	0.19	6.00	21.36	7.55
16.54	0.13	5.36	64.20	22.71
17.11	0.13	5.18	245.68	86.90
17.95	0.09	4.94	166.30	58.82
18.45	0.11	4.81	91.52	32.37
21.25	0.09	4.18	106.18	37.56
22.63	0.13	3.93	114.73	40.58
24.35	0.19	3.66	25.92	9.17
24.82	0.19	3.59	73.08	25.85
25.77	0.16	3.46	282.73	100.00
28.61	0.25	3.12	25.78	9.12
29.58	0.13	3.02	25.44	9.00
30.49	0.38	2.93	17.89	6.33
31.76	0.25	2.82	24.02	8.50

34.46	0.50	2.60	17.34	6.13
36.26	0.63	2.48	8.50	3.00
37.35	0.19	2.41	14.20	5.02

[0303] The XRPD peaks unique to each of the oxalate crystalline salt forms A and B disclosed herein are shown in Table 23.

[0304] Table 23

Oxalate Form	Peaks Unique to Each Form (°2Th.)		
Form A	6.48	13.01	23.82
Form B	7.38	13.30	16.54

[0305] Salicylate crystalline salt: was prepared in a primary salt screen. A 2 mL aliquot of 2-propanol or toluene was added to ~40 mg of free base. 1.05 equivalents of salicylic acid as a 1M solution in THF were added and the sample was temperature cycled for 3-5 days then the 2-propanol was evaporated.

[0306] The omecamtiv mecarbil salicylate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 24.

[0307] Table 24

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
8.36	0.06	10.58	5425.26	100.00
9.78	0.06	9.05	634.48	11.69
10.08	0.06	8.78	1112.62	20.51
10.37	0.05	8.53	164.07	3.02
11.30	0.06	7.83	1739.85	32.07
11.82	0.05	7.48	271.53	5.00
12.00	0.05	7.38	412.51	7.60
13.69	0.06	6.47	843.47	15.55
13.80	0.04	6.42	717.86	13.23
14.25	0.15	6.21	74.68	1.38
15.51	0.09	5.71	588.40	10.85
16.75	0.09	5.29	3941.54	72.65
17.56	0.06	5.05	2277.70	41.98
17.77	0.05	4.99	1649.96	30.41
17.86	0.08	4.97	2140.62	39.46
18.67	0.09	4.75	1989.60	36.67
19.11	0.08	4.64	1216.29	22.42
19.27	0.06	4.61	812.90	14.98
19.62	0.08	4.52	795.44	14.66

20.02	0.08	4.43	709.58	13.08
20.22	0.13	4.39	1916.47	35.33
20.79	0.08	4.27	721.20	13.29
21.07	0.13	4.22	1935.15	35.67
21.78	0.09	4.08	313.94	5.79
22.19	0.05	4.01	532.25	9.81
22.39	0.10	3.97	665.30	12.26
22.75	0.08	3.91	677.94	12.50
22.92	0.06	3.88	673.25	12.41
23.58	0.13	3.77	3464.58	63.86
24.10	0.10	3.69	257.32	4.74
24.99	0.10	3.56	457.77	8.44
25.23	0.09	3.53	1162.00	21.42
25.59	0.09	3.48	506.81	9.34
26.79	0.09	3.33	620.95	11.45
27.40	0.17	3.26	1709.43	31.51
27.78	0.15	3.21	231.91	4.27
28.21	0.18	3.16	2479.45	45.70
28.76	0.10	3.10	278.54	5.13
29.32	0.13	3.05	236.55	4.36
29.65	0.15	3.01	309.68	5.71
29.94	0.06	2.98	491.39	9.06
30.50	0.13	2.93	198.15	3.65
31.34	0.31	2.85	86.35	1.59
32.15	0.20	2.78	138.16	2.55
32.74	0.10	2.74	175.34	3.23
34.07	0.06	2.63	656.95	12.11

[0308] Hemi-succinate crystalline salt: as prepared during a high throughput screen by adding 0.3 mL of a 0.05M solution of free base in methanol and 0.3 mL of a succinic acid 0.05M solution in methanol to a glass plate then solvent was evaporated then 0.4 mL of solvent (water, 0.001M HCl aq, acetone, acetonitrile or hexanes) was added and heated to 50°C for 4 hours then evaporated. Water solubility was determined to be 7.4 mg/mL (pH 4.7).

[0309] The omecamtiv mecarbil hemi-succinate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 25.

[0310] Table 25

Angle 2-Theta°	d value Angstrom	Intensity Count (Height)	Rel. Int. [%]
5.488	16.102	191	11.5

6.319	13.988	504	30.5
7.596	11.638	237	14.3
12.932	6.8459	437	26.4
14.092	6.2847	245	14.8
15.079	5.8756	361	21.8
16.971	5.2244	307	18.6
18.768	4.728	1654	100
19.318	4.5947	1074	64.9
20.495	4.3334	1229	74.3
21.244	4.1822	790	47.8
21.889	4.0604	849	51.4
23.492	3.7869	1651	99.8
24.228	3.6736	531	32.1
25.364	3.5116	488	29.5
26.671	3.3424	749	45.3
27.389	3.2563	636	38.5
28.316	3.1518	635	38.4

[0311] Bis-sulfate crystalline salt form A: Procedure A - prepared during a high throughput screen by adding 0.2 mL of a 0.124M solution of free base in methanol and 0.2 mL of a sulfuric acid 0.25M solution in methanol to a glass plate then solvent was evaporated then 0.2 mL of solvent (THF or 0.001M HCl aq) was added and heated to 50°C for 4 hours then evaporated. Salt was scaled up by adding 100 mg free base to an 8 mL vial and 4 mL methanol and gently heating to dissolve. Sulphuric acid (2 mL of a 0.25 M solution) was added at room temperature. Solids were collected after evaporation to dryness.

[0312] Procedure B - prepared by dissolving 200.7 mg free base (1eq) and 1.05 mL of 1M sulfuric acid (2.1eq) in 5 mL methanol at 50°C then cooled to precipitate.

[0313] Procedure C - formed when Form D was exposed to 40C/75% RH storage conditions for 3 days. Water solubility was determined to be 16 mg/mL (pH 1.5).

[0314] The omecamtiv mecarbil bis-sulfate crystalline salt form A was characterized by an XRPD pattern comprising peaks in Table 26.

[0315] Table 26

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
5.39	0.05	16.40	803.64	14.66
7.55	0.06	11.71	915.17	16.70
8.33	0.05	10.62	247.72	4.52
10.13	0.08	8.73	62.00	1.13
14.35	0.09	6.17	3211.66	58.60

14.63	0.08	6.05	340.15	6.21
15.12	0.09	5.86	414.53	7.56
15.64	0.09	5.66	517.83	9.45
16.00	0.10	5.54	503.75	9.19
16.17	0.09	5.48	833.42	15.21
16.39	0.08	5.41	379.73	6.93
16.71	0.08	5.31	565.39	10.32
16.92	0.06	5.24	1727.49	31.52
17.07	0.08	5.19	1764.97	32.20
17.68	0.09	5.02	464.46	8.47
18.33	0.09	4.84	344.30	6.28
18.60	0.08	4.77	648.56	11.83
19.26	0.12	4.61	5480.54	100.00
19.75	0.10	4.50	350.13	6.39
20.22	0.08	4.39	1216.75	22.20
20.83	0.10	4.26	1272.51	23.22
21.03	0.08	4.23	507.32	9.26
21.38	0.09	4.16	1278.02	23.32
22.27	0.12	3.99	1717.52	31.34
22.77	0.09	3.90	602.87	11.00
23.14	0.13	3.84	1861.14	33.96
23.42	0.08	3.80	727.12	13.27
23.76	0.12	3.75	1376.36	25.11
24.32	0.12	3.66	1360.34	24.82
25.11	0.13	3.55	1608.13	29.34
25.48	0.41	3.50	169.13	3.09
25.74	0.13	3.46	2483.24	45.31
26.30	0.08	3.39	445.14	8.12
26.46	0.06	3.37	589.33	10.75
27.71	0.06	3.22	1509.03	27.53
28.15	0.08	3.17	1156.37	21.10
28.90	0.08	3.09	415.94	7.59
29.24	0.09	3.05	516.64	9.43
29.92	0.06	2.99	927.24	16.92
30.38	0.13	2.94	210.94	3.85
30.65	0.08	2.92	266.96	4.87
31.57	0.13	2.83	212.99	3.89
32.14	0.09	2.79	339.08	6.19
33.10	0.18	2.71	172.88	3.15
33.78	0.20	2.65	181.24	3.31
34.22	0.15	2.62	131.00	2.39

[0316] Bis-sulfate crystalline salt form B: was prepared by dissolving 150 mg free base in 20 mL of acetone and adding 22 μ L of 17.6M sulfuric acid then sonicating. Isolated solids were slurried in water at RT. Water solubility was determined to be 9 mg/mL (pH 3.5).

[0317] The omecamtiv mecarbil bis-sulfate crystalline salt form B was characterized by an XRPD pattern comprising peaks in Table 27.

[0318] Table 27

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
10.67	0.19	8.29	12.89	6.12
11.72	0.19	7.55	160.31	76.09
12.17	0.19	7.27	80.11	38.02
12.93	0.13	6.85	43.21	20.51
14.57	0.22	6.08	17.85	8.47
17.79	0.13	4.99	30.92	14.67
18.39	0.11	4.82	82.79	39.29
18.76	0.09	4.73	32.73	15.54
19.84	0.22	4.47	101.51	48.18
20.48	0.11	4.34	124.27	58.98
21.34	0.25	4.16	11.41	5.41
21.90	0.25	4.06	10.23	4.86
23.60	0.17	3.77	210.68	100.00
24.23	0.13	3.67	35.79	16.99
25.13	0.16	3.54	44.10	20.93
25.63	0.19	3.48	98.90	46.94
29.30	0.31	3.05	29.96	14.22
30.12	0.13	2.97	47.50	22.55
30.98	0.19	2.89	15.83	7.51
31.48	0.19	2.84	18.11	8.60
31.94	0.19	2.80	13.68	6.49
36.02	0.25	2.49	10.23	4.86
36.85	0.50	2.44	7.64	3.63
38.16	0.25	2.36	11.73	5.57

[0319] Bis-sulfate crystalline salt form C: was prepared by heating bis-sulfate Form B on TGA.

[0320] The omecamtiv mecarbil bis-sulfate crystalline salt form C was characterized by an XRPD pattern comprising peaks in Table 28.

[0321] Table 28

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
8.20	0.19	10.79	30.31	4.27

10.39	0.13	8.51	179.40	25.28
10.72	0.09	8.25	226.72	31.94
10.98	0.13	8.06	252.05	35.51
11.49	0.22	7.70	136.57	19.24
12.17	0.19	7.27	132.34	18.65
12.52	0.16	7.07	159.11	22.42
12.99	0.22	6.82	238.57	33.61
13.56	0.22	6.53	122.29	17.23
15.98	0.25	5.55	37.67	5.31
16.74	0.19	5.30	135.94	19.15
17.11	0.16	5.18	296.18	41.73
17.43	0.22	5.09	332.88	46.90
18.04	0.31	4.92	107.59	15.16
19.60	0.14	4.53	249.29	35.12
20.94	0.22	4.24	215.86	30.41
21.53	0.22	4.13	109.98	15.50
22.47	0.31	3.96	100.45	14.15
23.09	0.28	3.85	136.51	19.23
23.98	0.31	3.71	58.62	8.26
24.76	0.19	3.60	230.19	32.43
25.25	0.31	3.53	709.74	100.00
25.87	0.16	3.44	263.23	37.09
26.51	0.19	3.36	179.86	25.34
27.63	0.76	3.23	91.65	12.91
29.33	0.50	3.04	30.11	4.24
32.14	0.38	2.79	44.36	6.25
33.91	0.38	2.64	17.82	2.51
35.27	0.38	2.55	15.44	2.18
39.25	0.25	2.30	22.20	3.13

[0322] Sulfate crystalline salt form D: was prepared in a primary salt screen. To prepare Form D a 2 mL aliquot of acetone was added to ~40 mg of free base. 1.05 equivalents of sulfuric acid as a 1M solution in THF were added and the sample was temperature cycled for 3-5 days.

[0323] The omecamtiv mecarbil sulfate crystalline salt form D was characterized by an XRPD pattern comprising peaks in Table 29.

[0324] Table 29

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
7.32	0.08	12.08	844.93	59.30
8.02	0.05	11.03	224.63	15.77
11.67	0.10	7.58	138.68	9.73

12.20	0.10	7.25	169.56	11.90
12.55	0.10	7.05	121.39	8.52
13.57	0.08	6.52	1067.44	74.92
14.54	0.10	6.09	805.82	56.56
16.29	0.08	5.44	758.59	53.24
16.41	0.08	5.40	705.84	49.54
16.91	0.15	5.24	520.14	36.51
17.36	0.20	5.11	373.69	26.23
18.70	0.06	4.74	783.62	55.00
20.44	0.15	4.34	1424.79	100.00
21.02	0.15	4.23	587.18	41.21
21.77	0.15	4.08	625.52	43.90
22.37	0.15	3.97	403.56	28.32
22.90	0.15	3.88	1136.86	79.79
23.72	0.15	3.75	749.74	52.62
24.28	0.08	3.67	771.24	54.13
25.14	0.13	3.54	490.08	34.40
25.88	0.10	3.44	399.16	28.02
26.58	0.09	3.35	808.90	56.77
27.25	0.15	3.27	783.97	55.02
28.10	0.15	3.18	200.02	14.04
29.43	0.18	3.04	218.32	15.32
30.45	0.41	2.94	99.72	7.00
33.15	0.31	2.70	89.81	6.30
33.88	0.15	2.65	121.18	8.51

[0325] The XRPD peaks unique to each of the sulfate crystalline salt forms A-D disclosed herein are shown in Table 30.

[0326] Table 30

Sulfate Form	Peaks Unique to Each Form (°2Th.)				
Form A	5.39	7.55	14.35	19.26	20.22
Form B	11.72	20.48			
Form C	10.98	11.49	18.04	19.60	
Form D	7.32	8.02	20.44		

[0327] **2-Hydroxyethane sulfonate crystalline salt:** was prepared in a primary salt screen. A 2 mL aliquot of THF, acetonitrile or isopropyl acetate was added to ~40 mg of free base. 1.05 equivalents of 2-hydroxyethane sulfonic acid as a solid and 1 equivalent of 1M hydrochloric acid were added and the sample was temperature cycled for 3-5 days.

[0328] The omecamtiv mecarbil 2-hydroxyethane sulfonate crystalline salt was characterized by an XRPD pattern comprising peaks in Table 31.

[0329] Table 31

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
6.26	0.08	14.13	219.45	9.02
6.69	0.18	13.21	374.53	15.39
9.95	0.06	8.89	2434.20	100.00
14.43	0.15	6.14	98.15	4.03
14.99	0.23	5.91	360.35	14.80
15.51	0.15	5.71	182.59	7.50
16.37	0.09	5.41	438.44	18.01
17.06	0.26	5.20	148.91	6.12
17.85	0.08	4.97	916.01	37.63
19.61	0.15	4.53	272.20	11.18
19.93	0.05	4.46	695.74	28.58
20.07	0.06	4.42	590.93	24.28
20.46	0.12	4.34	487.31	20.02
20.95	0.18	4.24	261.07	10.73
22.06	0.18	4.03	158.42	6.51
22.89	0.26	3.88	158.38	6.51
23.96	0.15	3.71	128.47	5.28
24.41	0.13	3.65	155.88	6.40
25.06	0.15	3.55	488.47	20.07
26.20	0.18	3.40	632.72	25.99
26.98	0.20	3.30	133.39	5.48
27.92	0.08	3.20	162.41	6.67
28.43	0.15	3.14	116.74	4.80
29.98	0.04	2.98	438.04	18.00
32.16	0.04	2.78	443.24	18.21
33.38	0.41	2.68	86.61	3.56
34.39	0.06	2.61	387.48	15.92

[0330] Bis-tartrate crystalline salt form A: was prepared by dissolving 200.7 mg (1 eq) of free base and 525 μ L (30.02 g in 100 mL methanol, 2.1 eq) in 20 mL methanol and heat cycle 40°C/RT twice. Water solubility was determined to be >53 mg/mL (pH 3.28).

[0331] The omecamtiv mecarbil bis-tartrate crystalline salt form A was characterized by an XRPD pattern comprising peaks in Table 32.

[0332] Table 32

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
4.20	0.19	21.02	968.44	59.18

4.77	0.19	18.52	1062.98	64.96
7.49	0.16	11.80	715.72	43.74
7.67	0.09	11.52	629.63	38.48
8.22	0.19	10.75	793.41	48.48
8.43	0.09	10.49	699.24	42.73
9.49	0.19	9.32	377.23	23.05
11.18	0.25	7.91	193.33	11.81
11.88	0.31	7.45	208.48	12.74
13.05	0.22	6.79	590.85	36.11
13.26	0.13	6.68	608.71	37.20
14.98	0.11	5.92	1399.17	85.50
15.14	0.22	5.85	1490.06	91.05
16.42	0.35	5.40	437.01	26.70
17.34	0.08	5.11	983.61	60.11
17.47	0.14	5.08	1247.90	76.26
18.02	0.19	4.92	1121.36	68.52
18.23	0.16	4.87	911.65	55.71
18.72	0.13	4.74	441.84	27.00
19.20	0.16	4.62	370.82	22.66
21.19	0.11	4.19	1636.46	100.00
22.50	0.19	3.95	667.89	40.81
23.86	0.16	3.73	212.59	12.99
24.53	0.41	3.63	339.28	20.73
25.67	0.38	3.47	528.93	32.32
26.30	0.31	3.39	451.92	27.62
28.14	0.25	3.17	388.50	23.74
28.44	0.22	3.14	285.60	17.45
29.87	0.38	2.99	258.33	15.79
31.32	0.31	2.86	101.08	6.18
32.48	0.25	2.76	128.85	7.87
33.69	0.25	2.66	103.91	6.35
34.42	0.25	2.61	158.44	9.68
35.36	0.31	2.54	97.83	5.98
35.92	0.19	2.50	84.54	5.17
36.60	0.31	2.46	82.62	5.05
37.41	0.25	2.40	136.23	8.32
37.88	0.25	2.38	96.44	5.89

[0333] Bis-tartrate crystalline salt form B: was prepared by dissolving 1.004g of omecamtiv mecarbil and 0.788g L-tartaric acid (2.1 eq) in 50 mL of MeOH at 50°C then cooling to precipitate.

[0334] The omecamtiv mecarbil bis-tartrate crystalline salt form B was characterized by an XRPD pattern comprising peaks in Table 33.

[0335] Table 33

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
3.77	0.17	23.43	971.26	100.00
4.72	0.27	18.74	635.71	65.45
5.69	0.27	15.52	516.33	53.16
6.95	0.27	12.73	358.34	36.89
9.34	0.27	9.46	200.35	20.63
10.07	0.23	8.78	434.70	44.76
11.18	0.54	7.92	167.11	17.21
12.63	0.20	7.01	158.00	16.27
15.18	0.20	5.84	190.67	19.63
17.69	0.17	5.01	282.08	29.04
22.35	0.20	3.98	177.49	18.27
25.46	0.40	3.50	62.01	6.38

[0336] **Bis-tartrate crystalline salt form C:** was prepared by slurry of bis-tartrate Form B in water.

[0337] The omecamtiv mecarbil bis-tartrate crystalline salt form C was characterized by an XRPD pattern comprising peaks in Table 34.

[0338] Table 34

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
3.57	0.13	24.73	920.53	89.99
3.86	0.15	22.88	1022.92	100.00
4.78	0.40	18.49	513.98	50.25
6.23	0.20	14.19	427.74	41.82
7.04	0.13	12.56	270.90	26.48
9.36	0.17	9.45	637.85	62.36
13.08	0.10	6.77	549.91	53.76
13.96	0.17	6.34	239.78	23.44
15.84	0.17	5.59	348.37	34.06
16.88	0.20	5.25	121.74	11.90
17.60	0.27	5.04	146.72	14.34
18.20	0.20	4.87	147.06	14.38
18.73	0.23	4.74	245.18	23.97
20.40	0.20	4.35	268.32	26.23
22.58	0.27	3.94	120.04	11.73
25.44	0.40	3.50	192.03	18.77
26.06	0.47	3.42	213.11	20.83

28.61	0.40	3.12	106.58	10.42
-------	------	------	--------	-------

[0339] Mono-tartrate crystalline salt form D: was prepared by mixing 1.004g of omecamtiv mecarbil and 0.375g of L-tartaric acid (1 eq) in 10 mL of 5% H₂O in THF at 50°C then cooling to precipitate.

[0340] The omecamtiv mecarbil mono-tartrate crystalline salt form D was characterized by an XRPD pattern comprising peaks in Table 35.

[0341] Table 35

Pos. [°2Th.]	FWHM [°2Th.]	d-spacing [Å]	Height [cts]	Rel. Int. [%]
6.94	0.12	12.74	377.02	15.46
9.77	0.17	9.05	478.47	19.62
10.87	0.17	8.14	2097.98	86.02
12.74	0.13	6.95	204.81	8.40
13.04	0.13	6.79	144.17	5.91
13.79	0.15	6.42	871.83	35.75
14.54	0.20	6.09	418.22	17.15
14.86	0.13	5.96	315.83	12.95
15.40	0.18	5.75	2345.40	96.17
17.36	0.18	5.11	1127.77	46.24
17.74	0.17	5.00	512.38	21.01
18.58	0.12	4.78	598.49	24.54
18.87	0.17	4.70	693.22	28.42
19.25	0.17	4.61	322.02	13.20
20.71	0.13	4.29	293.45	12.03
21.78	0.17	4.08	1734.64	71.13
23.11	0.20	3.85	97.51	4.00
23.58	0.13	3.77	214.55	8.80
24.69	0.12	3.61	423.81	17.38
25.43	0.17	3.50	2438.82	100.00
26.24	0.20	3.40	529.42	21.71
26.50	0.13	3.36	437.75	17.95
26.99	0.27	3.30	259.96	10.66
28.58	0.27	3.12	73.70	3.02
29.43	0.27	3.03	100.12	4.11
30.40	0.13	2.94	161.38	6.62
32.74	0.27	2.74	127.69	5.24
34.76	0.17	2.58	210.18	8.62
35.46	0.17	2.53	118.22	4.85
36.31	0.20	2.47	191.26	7.84
37.01	0.13	2.43	138.71	5.69
37.64	0.27	2.39	131.07	5.37

[0342] The XRPD peaks unique to each of the tartrate crystalline salt forms A-D disclosed herein are shown in Table 36.

[0343] Table 36

Tartrate Form	Peaks Unique to Each Form (°2Th.)					
Form A	4.20	7.49	8.22	11.88	16.42	21.19
Form B	3.77	5.69	10.07			
Form C	3.57	6.23	15.84			
Form D	9.77	15.40				

What is Claimed:

1. An omecamtiv mecarbil free base crystalline form III, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 9.50, 19.06, and $23.01 \pm 0.2^\circ$ 2θ using Cu K α radiation.
2. The omecamtiv mecarbil of claim 1, further characterized by XRPD pattern peaks at 14.27, 15.25, 16.10, 17.78, and $23.87 \pm 0.2^\circ$ 2θ using Cu K α radiation.
3. The omecamtiv mecarbil of claim 2, further characterized by XRPD pattern peaks at 7.91, 20.65, 28.11, 31.01, 31.95, and $32.34 \pm 0.2^\circ$ 2θ using Cu K α radiation.
4. The omecamtiv mecarbil of any one of claims 1 to 3, having an XRPD pattern substantially as shown in Figure 1.
5. The omecamtiv mecarbil of any one of claims 1 to 4, having an endothermic transition at 175°C to 190°C, as measured by differential scanning calorimetry.
6. The omecamtiv mecarbil of claim 5, wherein the endothermic transition is at $186^\circ\text{C} \pm 3^\circ\text{C}$.
7. The omecamtiv mecarbil of any one of claims 1 to 6, having a thermogravimetric analysis (TGA) substantially as shown in Figure 3.
8. An omecamtiv mecarbil free base crystalline form IV, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 5.18, 10.35, 14.84, 15.54, 18.10, and $19.92 \pm 0.2^\circ$ 2θ using Cu K α radiation.
9. The omecamtiv mecarbil of claim 8, further characterized by XRPD pattern peaks at 14.21, 20.62, 20.77, 22.86, 24.05, 24.36, 27.81, and $29.42 \pm 0.2^\circ$ 2θ using Cu K α radiation.
10. The omecamtiv mecarbil of claim 9, further characterized by XRPD pattern peaks at 7.42, 7.70, 21.70, 22.40, 23.09, 25.20, 25.72, 27.40, 28.18, 28.63, 28.98, and $30.51 \pm 0.2^\circ$ 2θ using Cu K α radiation.
11. The omecamtiv mecarbil of any one of claims 8 to 10, having an XRPD pattern substantially as shown in Figure 4.
12. The omecamtiv mecarbil of any one of claims 8 to 11, having an endothermic transition at 175°C to about 190°C, as measured by differential scanning calorimetry.
13. The omecamtiv mecarbil of claim 12, wherein the endothermic transition is at $185^\circ\text{C} \pm 3^\circ\text{C}$.
14. The omecamtiv mecarbil of any one of claims 8 to 13, having a thermogravimetric analysis (TGA) as shown in Figure 6.
15. An omecamtiv mecarbil free base crystalline form V, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 7.38, 8.56, 9.14, and $18.28 \pm 0.2^\circ$ 2θ using Cu K α radiation.
16. The omecamtiv mecarbil of claim 15, further characterized by XRPD pattern peaks at 8.93, 10.03, 10.73, 11.71, 13.69, 15.08, 16.85, 17.85, 18.86, 20.05, 20.72, 21.74, 23.56, 24.03, 26.23, and $27.62 \pm 0.2^\circ$ 2θ using Cu K α radiation.

17. The omecamtiv mecarbil of claim 16, further characterized by XRPD pattern peaks at 5.40, 16.04, 22.83, 25.45, 26.23, 27.62, 28.58, 29.85, 32.10, and $33.37 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
18. The omecamtiv mecarbil of any one of claims 15 to 17, having an XRPD pattern substantially as shown in Figure 7.
19. The omecamtiv mecarbil of any one of claims 15 to 18, having an endothermic transition at 175°C to 190°C, as measured by differential scanning calorimetry.
20. The omecamtiv mecarbil of claim 19, wherein the endothermic transition is at 185°C $\pm 3^\circ\text{C}$.
21. The omecamtiv mecarbil of any one of claims 15 to 20, having a thermogravimetric analysis (TGA) as shown in Figure 9.
22. An omecamtiv mecarbil free base crystalline form VI, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 9.07, 16.67, 18.18, 19.70, 20.89, and $21.28 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
23. The omecamtiv mecarbil of claim 22, further characterized by XRPD pattern peaks at 15.88, 17.81, 18.80, 23.72, 24.26, 26.80, 27.59, and $29.82 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
24. The omecamtiv mecarbil of claim 23, further characterized by XRPD pattern peaks at 14.28, 20.23, 26.19, and $28.90 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
25. The omecamtiv mecarbil of any one of claims 22 to 24, having an XRPD pattern substantially as shown in Figure 10.
26. The omecamtiv mecarbil of any one of claims 22 to 25, having an endothermic transition at 175°C to 190°C, as measured by differential scanning calorimetry.
27. The omecamtiv mecarbil of claim 26, wherein the endothermic transition is at 185°C $\pm 3^\circ\text{C}$.
28. An omecamtiv mecarbil free base crystalline form VII, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 8.40, 8.71, 13.08, 15.66, and $19.61 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
29. The omecamtiv mecarbil of claim 28, further characterized by XRPD pattern peaks at 4.37, 16.83, 18.92, 20.32, 20.49, 22.26, 24.21, and $25.41 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
30. The omecamtiv mecarbil of claim 29, further characterized by XRPD pattern peaks at 7.84, 10.81, 21.61, 23.22, 23.46, 27.58, 29.53, 30.13, and $31.32 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
31. The omecamtiv mecarbil of any one of claims 28 to 30, having an XRPD pattern substantially as shown in Figure 12.
32. An omecamtiv mecarbil ethane sulfonate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 8.61, 16.14, 16.76, 16.97, 20.73, 20.96, 25.95, and $26.30 \pm 0.2^\circ 2\theta$ using Cu K α radiation.

33. The omecamtiv mecarbil of claim 32, further characterized by XRPD pattern peaks at 17.23, 18.35, 19.20, 20.27, 23.73, 25.24, and $27.09 \pm 0.2^\circ$ 2θ using Cu K α radiation.
34. The omecamtiv mecarbil of claim 32 or 33, having an XRPD pattern substantially as shown in Figure 18.
35. An omecamtiv mecarbil bis-fumarate crystalline salt, form A, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 5.64, 15.76, 22.03, and $23.87 \pm 0.2^\circ$ 2θ using Cu K α radiation.
36. The omecamtiv mecarbil of claim 35, further characterized by XRPD pattern peaks at 16.80, 21.55, 21.87, 23.61, 23.87, 26.01, and $27.20 \pm 0.2^\circ$ 2θ using Cu K α radiation.
37. The omecamtiv mecarbil of claim 36, further characterized by XRPD pattern peaks at 10.80, 12.04, 15.76, 16.40, 17.94, 18.32, 19.87, 20.61, 22.88, 27.86, 32.73, and $36.54 \pm 0.2^\circ$ 2θ using Cu K α radiation.
38. The omecamtiv mecarbil of any one of claims 35 to 37, having an XRPD pattern substantially as shown in Figure 20.
39. The omecamtiv mecarbil of any one of claims 35 to 38, having a thermogravimetric analysis (TGA) as shown in Figure 21.
40. An omecamtiv mecarbil bis-fumarate crystalline salt, form B, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 5.68, 6.11, 13.13, 18.08, and $22.47 \pm 0.2^\circ$ 2θ using Cu K α radiation.
41. The omecamtiv mecarbil of claim 40, further characterized by XRPD pattern peaks at 9.69, 11.43, 12.92, 15.95, 20.81, 22.95, 26.04, 27.01, and $28.43 \pm 0.2^\circ$ 2θ using Cu K α radiation.
42. The omecamtiv mecarbil of claim 41, further characterized by XRPD pattern peaks at 19.52, 24.53, 31.37, 32.32, 34.89, 35.89, and $37.16 \pm 0.2^\circ$ 2θ using Cu K α radiation.
43. The omecamtiv mecarbil of any one of claims 40 to 42, having an XRPD pattern substantially as shown in Figure 22.
44. The omecamtiv mecarbil of any one of claims 40 to 43, having a thermogravimetric analysis (TGA) as shown in Figure 23.
45. An omecamtiv mecarbil bis-fumarate crystalline salt, form C, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 5.88, 18.79, 25.41, and $26.86 \pm 0.2^\circ$ 2θ using Cu K α radiation.
46. The omecamtiv mecarbil of claim 45, further characterized by XRPD pattern peaks at 12.74, 13.56, 17.15, 17.63, 20.29, 21.47, 21.77, 22.21, 22.92, 23.58, 24.15, 25.41, 26.78, and $27.83 \pm 0.2^\circ$ 2θ using Cu K α radiation.
47. The omecamtiv mecarbil of claim 45 or 46, having an XRPD pattern substantially as shown in Figure 24.
48. The omecamtiv mecarbil of any one of claims 45 to 47, having a thermogravimetric analysis (TGA) as shown in Figure 25.

49. An omecamtiv mecarbil mono-fumarate crystalline salt, form D, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 8.01, 15.20, and 20.02 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
50. The omecamtiv mecarbil of claim 49, further characterized by XRPD pattern peaks at 12.11, 12.67, 14.46, 16.01, 16.57, 17.04, 17.63, 20.51, 21.75, 22.86, 24.25, 24.97, 25.84, 26.17, 27.10, 27.97, and 29.21 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
51. The omecamtiv mecarbil of claim 49 or 50, having an XRPD pattern substantially as shown in Figure 26.
52. The omecamtiv mecarbil of any one of claims 49 to 51, having an endothermic transition at 110°C to 130°C, as measured by differential scanning calorimetry.
53. The omecamtiv mecarbil of claim 52, wherein the endothermic transition is at 125°C $\pm 3^\circ\text{C}$.
54. The omecamtiv mecarbil of any one of claims 49 to 53, having a thermogravimetric analysis (TGA) as shown in Figure 28.
55. An omecamtiv mecarbil bis-maleate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 9.97, 15.31, 16.04, and 26.96 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
56. The omecamtiv mecarbil of claim 55, further characterized by XRPD pattern peaks at 10.56, 13.25, 15.53, 16.38, 17.44, 17.70, 18.17, 19.00, 20.13, 21.47, 22.31, 22.44, 24.38, 24.64, 25.66, 26.66, and 27.83 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
57. The omecamtiv mecarbil of claim 56, further characterized by XRPD pattern peaks at 4.99, 14.83, 17.10, 22.02, 28.55, 30.76, 32.01, 34.39, and 34.51 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
58. The omecamtiv mecarbil of any one of claims 55 to 57, having an XRPD pattern substantially as shown in Figure 30.
59. The omecamtiv mecarbil of any one of claims 55 to 58, having an endothermic transition at 160°C to 210°C, as measured by differential scanning calorimetry.
60. The omecamtiv mecarbil of claim 59, wherein the endothermic transition is at 190°C $\pm 3^\circ\text{C}$.
61. The omecamtiv mecarbil of any one of claims 55 to 60, having a thermogravimetric analysis (TGA) as shown in Figure 32.
62. An omecamtiv mecarbil bis-malonate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 4.74, 11.37, 14.25, 15.13, 18.29, 20.14, 23.87, 27.78, and 28.01 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
63. The omecamtiv mecarbil of claim 62, further characterized by XRPD pattern peaks at 9.30, 13.73, 16.45, 16.83, 18.08, 18.88, 19.54, 20.77, 21.21, 23.32, 24.67, 26.51, 27.59, and 28.90 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.
64. The omecamtiv mecarbil of claim 63, further characterized by XRPD pattern peaks at 15.69, 25.72, 30.18, 33.70, 34.19 $\pm 0.2^\circ 2\theta$ using Cu K α radiation.

65. The omecamtiv mecarbil of any one of claims 62 to 64, having an XRPD pattern substantially as shown in Figure 33.
66. The omecamtiv mecarbil of any one of claims 62 to 65, having a thermogravimetric analysis (TGA) as shown in Figure 34.
67. An omecamtiv mecarbil mesylate crystalline salt form A, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 4.02, 4.87, 15.21, 15.86, 20.53, and $24.39 \pm 0.2^\circ$ 2θ using Cu K α radiation.
68. The omecamtiv mecarbil of claim 68, further characterized by XRPD pattern peaks at 7.79, 11.61, 16.51, 17.57, 18.42, 19.26, 21.55, 23.17, 25.51, 26.38, and $27.63 \pm 0.2^\circ$ 2θ using Cu K α radiation.
69. The omecamtiv mecarbil of claim 67 or 68, having an XRPD pattern substantially as shown in Figure 35.
70. The omecamtiv mecarbil of any one of claims 67 to 69, having a thermogravimetric analysis (TGA) as shown in Figure 36.
71. An omecamtiv mecarbil bis-mesylate crystalline salt form B, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 8.30, 8.94, 9.59, 12.15, 14.37, 19.82, 20.29, 22.04, and $25.02 \pm 0.2^\circ$ 2θ using Cu K α radiation.
72. The omecamtiv mecarbil of claim 71, further characterized by XRPD pattern peaks at 11.66, 16.18, 16.64, 16.81, 17.07, 17.19, 17.41, 17.76, 19.24, 20.66, 21.62, 22.39, 23.95, 24.60, 25.59, 25.89, 27.14, 27.35, 27.41, and $29.45 \pm 0.2^\circ$ 2θ using Cu K α radiation.
73. The omecamtiv mecarbil of claim 72, further characterized by XRPD pattern peaks at 10.78, 11.15, 14.93, 15.36, 15.57, 23.54, 26.14, 26.49, 27.89, 28.86, 29.89, 31.11, 32.47, 33.10, 33.51, $34.56 \pm 0.2^\circ$ 2θ using Cu K α radiation.
74. The omecamtiv mecarbil of any one of claims 71 to 73, having an XRPD pattern substantially as shown in Figure 37.
75. An omecamtiv mecarbil bis-naphthalate-2-sulfonate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 4.49, 18.20, 18.62, 21.38, 21.52, and $26.11 \pm 0.2^\circ$ 2θ using Cu K α radiation.
76. The omecamtiv mecarbil of claim 75, further characterized by XRPD pattern peaks at 6.25, 6.65, 13.44, 14.39, 14.92, 16.28, 18.90, 19.53, 20.82, 22.02, 22.43, 22.80, 24.40, 25.16, 27.01, 29.67, 31.63, and $33.42 \pm 0.2^\circ$ 2θ using Cu K α radiation.
77. The omecamtiv mecarbil of claim 75 or 76, having an XRPD pattern substantially as shown in Figure 40.
78. An omecamtiv mecarbil mono-napadisylate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 12.27, 15.75, 16.5, 17.83, 19.94, 21.83, and $22.87 \pm 0.2^\circ$ 2θ using Cu K α radiation.
79. The omecamtiv mecarbil of claim 78, further comprising XRPD pattern peaks at 10.84, 13.41, 14.57, 15.14, 18.82, 23.49, 24.34, and $25.26 \pm 0.2^\circ$ 2θ using Cu K α radiation.

80. The omecamtiv mecarbil of claim 78 or 79, having an XRPD pattern substantially as shown in Figure 42.
81. The omecamtiv mecarbil of any one of claims 78 to 80, having a differential scanning calorimetry (DSC) as shown in Figure 43.
82. The omecamtiv mecarbil of any one of claims 78 to 81, having a thermogravimetric analysis (TGA) as shown in Figure 44.
83. An omecamtiv mecarbil nicotinate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 3.69, 8.55, 9.13, 16.70, 16.84, 18.30, 19.99, 20.76, 23.43, 24.83, and $25.95 \pm 0.2^\circ$ 2θ using Cu K α radiation.
84. The omecamtiv mecarbil of claim 83, further characterized by XRPD pattern peaks at 7.36, 10.01, 12.43, 14.74, 15.50, 17.62, 18.58, 19.59, 20.34, 21.32, 22.03, 22.91, 23.87, 24.92, 25.40, 26.85, 26.94, 27.32, 28.01, and $28.94 \pm 0.2^\circ$ 2θ using Cu K α radiation.
85. The omecamtiv mecarbil of claim 83 or 84, having an XRPD pattern substantially as shown in Figure 45.
86. An omecamtiv mecarbil oxalate crystalline salt, form A, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 6.48, 13.01, and $23.82 \pm 0.2^\circ$ 2θ using Cu K α radiation.
87. The omecamtiv mecarbil of claim 86, further characterized by XRPD pattern peaks at 10.36, 11.85, 14.79, 15.35, 17.11, 19.23, 19.91, 21.48, 22.07, 22.75, 25.70, 28.55, and $30.71 \pm 0.2^\circ$ 2θ using Cu K α radiation.
88. The omecamtiv mecarbil of claim 86 or 87, having an XRPD pattern substantially as shown in Figure 47.
89. The omecamtiv mecarbil of any one of claims 86 to 88, having an endothermic transition at 190°C to 230°C, as measured by differential scanning calorimetry.
90. The omecamtiv mecarbil of claim 89, wherein the endothermic transition is at $209^\circ\text{C} \pm 3^\circ\text{C}$.
91. The omecamtiv mecarbil of any one of claims 86 to 87, having a thermogravimetric analysis (TGA) as shown in Figure 49.
92. An omecamtiv mecarbil oxalate crystalline salt, form B, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 7.38, 13.30, and $16.54 \pm 0.2^\circ$ 2θ using Cu K α radiation.
93. The omecamtiv mecarbil of claim 92, further characterized by XRPD pattern peaks at 17.11, 17.95, 18.45, 21.25, 22.63, 24.82, and $25.77 \pm 0.2^\circ$ 2θ using Cu K α radiation.
94. The omecamtiv mecarbil of claim 93, further characterized by XRPD pattern peaks at 14.76, 24.35, 28.61, 29.58, 30.49, 31.76, 34.46, and $37.35 \pm 0.2^\circ$ 2θ using Cu K α radiation.
95. The omecamtiv mecarbil of any one of claims 92 to 94, having an XRPD pattern substantially as shown in Figure 50.

96. The omecamtiv mecarbil of any one of claims 92 to 95, having a thermogravimetric analysis (TGA) as shown in Figure 51.
97. An omecamtiv mecarbil salicylate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 8.36, 16.75, 17.56, 23.58, and 28.21 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
98. The omecamtiv mecarbil of claim 97, further characterized by XRPD pattern peaks at 10.08, 11.30, 13.69, 17.77, 17.86, 18.67, 19.11, 20.22, 21.07, 25.23, and 27.40 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
99. The omecamtiv mecarbil of claim 98, further characterized by XRPD pattern peaks at 9.78, 12.00, 13.80, 15.51, 19.27, 19.62, 20.02, 20.79, 22.19, 22.39, 22.75, 22.92, 24.99, 25.59, 26.79, 29.94, and 34.07 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
100. The omecamtiv mecarbil of any one of claims 97 to 99, having an XRPD pattern substantially as shown in Figure 53.
101. An omecamtiv mecarbil hemi-succinate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 6.32, 18.87, 19.32, 20.5, 21.24, 21.89, 23.49, 24.23, and 26.71 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
102. The omecamtiv mecarbil of claim 101, further comprising XRPD pattern peaks at 12.93, 15.08, 16.97, 25.36, 27.39, and 28.32 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
103. The omecamtiv mecarbil of claim 101 or 102, having an XRPD pattern substantially as shown in Figure 55.
104. The omecamtiv mecarbil of any one of claims 101 to 103, having an endothermic transition at 155°C to 190°C, as measured by differential scanning calorimetry.
105. The omecamtiv mecarbil of claim 104, wherein the endothermic transition is at 171°C $\pm 3^\circ$ C.
106. An omecamtiv mecarbil bis-sulfate crystalline salt, form A, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 5.39, 7.55, 14.35, 19.26, and 20.22 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
107. The omecamtiv mecarbil of claim 106, further characterized by XRPD pattern peaks at 16.17, 16.71, 16.92, 17.07, 18.60, 20.83, 21.38, 22.27, 22.77, 23.14, 23.42, 23.76, 24.32, 25.11, 25.74, 26.46, 27.71, 28.15, and 29.92 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
108. The omecamtiv mecarbil of claim 106 or 107, having an XRPD pattern substantially as shown in Figure 57.
109. An omecamtiv mecarbil bis-sulfate crystalline salt, form B, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 11.72 and 20.48 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.
110. The omecamtiv mecarbil of claim 109, further characterized by XRPD pattern peaks at 12.17, 12.93, 17.79, 18.39, 18.76, 19.84, 23.60, 25.13, 25.63, and 30.12 $\pm 0.2^\circ$ 2 θ using Cu K α radiation.

111. The omecamtiv mecarbil of claim 109 or 111, having an XRPD pattern substantially as shown in Figure 58.
112. The omecamtiv mecarbil of any one of claims 109 to 111, having a thermogravimetric analysis (TGA) as shown in Figure 59.
113. An omecamtiv mecarbil bis-sulfate crystalline salt, form C, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 10.98, 11.49, 18.04, and $19.60 \pm 0.2^\circ$ 2θ using Cu K α radiation.
114. The omecamtiv mecarbil of claim 113, further characterized by XRPD pattern peaks at 10.39, 10.72, 12.52, 12.99, 17.11, 17.43, 20.94, 24.76, 25.25, 25.87, and $26.51 \pm 0.2^\circ$ 2θ using Cu K α radiation.
115. The omecamtiv mecarbil of claim 113 or 114, having an XRPD pattern substantially as shown in Figure 60.
116. The omecamtiv mecarbil of any one of claims 113 to 115, having a thermogravimetric analysis (TGA) as shown in Figure 61.
117. An omecamtiv mecarbil sulfate crystalline salt, form D, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 7.32, 8.02, and $20.44 \pm 0.2^\circ$ 2θ using Cu K α radiation.
118. The omecamtiv mecarbil of claim 117, further characterized by XRPD pattern peaks at 13.57, 14.54, 16.29, 16.41, 16.91, 17.36, 18.70, 21.02, 21.77, 22.37, 22.90, 23.72, 24.28, 25.14, 25.88, 26.58, 27.25, 28.10, and $29.43 \pm 0.2^\circ$ 2θ using Cu K α radiation.
119. The omecamtiv mecarbil of claim 117 or 119, having an XRPD pattern substantially as shown in Figure 62.
120. An omecamtiv mecarbil 2-hydroxyethane sulfonate crystalline salt, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 9.95, 17.85, 19.93, 20.07, 20.46, 25.06, and $26.20 \pm 0.2^\circ$ 2θ using Cu K α radiation.
121. The omecamtiv mecarbil of claim 120, further characterized by XRPD pattern peaks at 6.26, 6.69, 14.99, 16.37, 19.61, 20.95, 29.98, 32.16, and $34.39 \pm 0.2^\circ$ 2θ using Cu K α radiation.
122. The omecamtiv mecarbil of claim 120 or 122, having an XRPD pattern substantially as shown in Figure 65.
123. An omecamtiv mecarbil bis-tartrate crystalline salt, form A, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 4.20, 7.49, 8.22, 11.88, 16.42, and $21.19 \pm 0.2^\circ$ 2θ using Cu K α radiation.
124. The omecamtiv mecarbil of claim 123, further characterized by XRPD pattern peaks at 4.77, 7.67, 8.43, 9.49, 13.05, 13.26, 14.98, 15.14, 17.34, 17.47, 18.02, 18.23, 18.72, 19.20, 22.50, 24.53, 25.67, 26.30, and $28.14 \pm 0.2^\circ$ 2θ using Cu K α radiation.
125. The omecamtiv mecarbil of claim 123 or 124, having an XRPD pattern substantially as shown in Figure 67.

126. The omecamtiv mecarbil of any one of claims 123 to 125, having a thermogravimetric analysis (TGA) as shown in Figure 68.
127. An omecamtiv mecarbil bis-tartrate crystalline salt, form B, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 3.77, 5.69, and $10.07 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
128. The omecamtiv mecarbil of claim 127, further characterized by XRPD pattern peaks at 4.72, 6.95, 9.34, 11.18, 12.63, 15.18, 17.69, 22.35, and $25.46 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
129. The omecamtiv mecarbil of claim 127 or 128, having an XRPD pattern substantially as shown in Figure 69.
130. The omecamtiv mecarbil of any one of claims 127 to 129, having a thermogravimetric analysis (TGA) as shown in Figure 70.
131. An omecamtiv mecarbil bis-tartrate crystalline salt, form C, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 3.57, 6.23, and $15.84 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
132. The omecamtiv mecarbil of claim 131, further characterized by XRPD pattern peaks at 3.86, 4.78, 7.04, 9.36, 13.08, 13.96, 16.88, 17.60, 18.20, 18.73, 20.40, 22.58, 25.44, 26.06, and $28.61 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
133. The omecamtiv mecarbil of claim 131 or 132, having an XRPD pattern substantially as shown in Figure 71.
134. The omecamtiv mecarbil of any one of claims 131 to 133, having a thermogravimetric analysis (TGA) as substantially shown in Figure 72.
135. An omecamtiv mecarbil mono-tartrate crystalline salt, form D, characterized by an X-ray powder diffraction (XRPD) pattern comprising peaks at 9.77 and $15.40 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
136. The omecamtiv mecarbil of claim 135, further characterized by XRPD pattern peaks at 10.87, 13.79, 17.36, 17.74, 18.58, 18.87, 21.78, 25.43, and $26.24 \pm 0.2^\circ 2\theta$ using Cu K α radiation.
137. The omecamtiv mecarbil of claim 135 or 136, having an XRPD pattern substantially as shown in Figure 73.
138. The omecamtiv mecarbil of any one of claims 135 to 137, having a thermogravimetric analysis (TGA) as shown in Figure 74.
139. An omecamtiv mecarbil amorphous hydrochloride salt.
140. The omecamtiv mecarbil of claim 139, having a differential scanning calorimetry (DSC) transition as substantially shown in Figure 15.
141. The omecamtiv mecarbil of claim 139 or 140, having a TGA as substantially shown in Figure 16.
142. The omecamtiv mecarbil of any one of claims 139 to 141, having a moisture sorption profile as substantially shown in Figure 17.

143. A pharmaceutical composition comprising the omecamtiv mecarbil of any one of claims 1 to 142 and a pharmaceutically acceptable excipient.
144. A method of treating heart failure in a subject in need thereof comprising administering to the subject the omecamtiv mecarbil of any one of claims 1 to 142 or the composition of claim 143 in an amount effective to treat heart failure.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28




Figure 29




Figure 30




Figure 31




Figure 32




Figure 33




Figure 34




Figure 35




Figure 36




Figure 37




Figure 38




Figure 39




Figure 40




Figure 41




Figure 42




Figure 43




Figure 44




Figure 45




Figure 46




Figure 47




Figure 48




Figure 49




Figure 50




Figure 51




Figure 52




Figure 53






Figure 54





Temperature (°C)

Universal V4.DC TA Instrument

Figure 56




Figure 57




Figure 58




Figure 59




Figure 60




Figure 61




Figure 62




Figure 63




Figure 64

65/75




Figure 65




Figure 66




Figure 67




Figure 68




Figure 69




Figure 70




Figure 71




Figure 72




Figure 73




Figure 74




Figure 75