

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number  
**WO 2004/005516 A1**

- (51) International Patent Classification<sup>7</sup>: C12N 15/54, 15/60, 15/82, 5/10, A01H 5/00
- (21) International Application Number: PCT/EP2003/007085
- (22) International Filing Date: 3 July 2003 (03.07.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 02015247.6 9 July 2002 (09.07.2002) EP
- (71) Applicant (for all designated States except US): **BASF PLANT SCIENCE GMBH** [DE/DE]; 67056 Ludwigshafen (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ANDERSSON, Mariette** [SE/SE]; Pilegatan 20, S-22229 Lund (SE). **TRIFONOVA, Adelina** [SE/SE]; Nordanväg 10a, 2v, S-24438 Kävlinge (SE). **HOFVANDER, Per** [SE/SE]; Doppingränd 8, S-23941 Falsterbo (SE).
- (74) Agent: **PREBLER, Uwe**; BASF Aktiengesellschaft, 67056 Ludwigshafen (DE).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
  - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 2004/005516 A1

(54) Title: USE OF AHAS MUTANT GENES AS SELECTION MARKER IN POTATO TRANSFORMATION

(57) Abstract: The present invention relates to the use of mutated AHAS genes conferring resistance to herbicides and resulting in an highly efficient selection system for the production of transgenic potato lines. The invention provides a key advantage by minimizing the escapes to almost zero of all shoots regenerated without interfering with the high number of shoots regenerated per explant.

Use of AHAS mutant genes as selection marker in potato transformation

5 The present invention relates to an improved selection system, using a mutated acetohydroxy acid synthase (AHAS) gene, for production of transgenic potato lines. Genetic insertion of mutated AHAS genes can give plants tolerance to a number of herbicide  
10 compounds. This invention yields a higher transformation efficiency and a reduced escape rate than previously described selection systems for potato plants.

The invention relates to a method for the generation of stably transformed fertile plants of the genus *Solanum*, which comprises  
15 the following steps:

- (a) growing the plant to be transformed, and obtaining the suitable explant,
- 20 (b) transferring DNA sequences into plant cells,
- (c) selecting transformed plant cells and
- (d) regenerating transgenic fertile plants.

25 Moreover, the method comprises the possibility of transferring homologous or heterologous DNA sequences into the plant to be transformed or explants thereof, for example by *Agrobacterium tumefaciens*.

30 Moreover, the method comprises the possibility of using leaves as explants.

Acetohydroxyacid synthase (EC 4.1.3.18; AHAS; acetolactate synthase) is an enzyme catalysing, in two parallel pathways, the  
35 first step of the synthesis of the branched-chain aminoacids valin, leucin and isoleucin. In the valin and leucin biosynthesis, AHAS is catalysing the production of acetolactate by condensation of two pyruvate molecules. While in the isoleucin  
40 biosynthesis AHAS condensates one pyruvate molecule with one 2-oxobutyrate molecule to form acetohydroxybutyrate (Umbarger, 1975). Sequence comparison of the AHAS gene in higher plants shows high conservation in at least 10 regions. These regions are probably of great importance for the AHAS function. In tobacco  
45 two unlinked genes named SuRA and SuRB code for the AHAS enzyme

catalytic subunit. This is not the case in *Arabidopsis thaliana* where only one gene is coding for the enzyme.

AHAS is the target enzyme of several classes of herbicides including sulphanylureas (Ray, 1984), imidazolinones (Shaner et al, 1984), triazolopyrimidines (Subrimanian et al, 1989) and pyrimidinyl oxybenzoat (Hawkes et al, 1989). The herbicides prevent branched amino acids to be synthesised by the plant and may lead to plant death. However, mutations in the AHAS genes can result in higher tolerance to these herbicides (US 5,013,659) because of reduced affinity between enzyme and the herbicide.

In plants mutations conferring resistance to herbicides occur as a result of exposure to the compound repeatedly as it was reported for *Arabidopsis thaliana*. Once the mutated genes are isolated they can be used to genetically engineer plants for improved tolerance to the herbicides. For example mutations in the *Arabidopsis thaliana* AHAS gene have been produced and successfully confer resistance to imidazolinones as described in WO 00/26390, US 5,767,366 and US 6,225,105. Mutations in a corn AHAS gene confer imidazolinone resistance to monocot plants as described in EP 0 525 384.

The mutated AHAS genes can be used for production of herbicide resistant plants, yielding field resistance to a specific herbicide or can be used as a selection marker for genetic engineering of plants.

When genetic material is introduced into a population of plant cells, only a minor part of the cells are successfully transformed. For the production of novel genetically modified plants a selection system is transformed together with the trait genes providing the transformed cells with a selective growth advantage. This construct allows to select transformed from non-transformed cells by adding a compound favoring the regeneration of transformed shoots.

A number of selection systems have been developed; most of them are based on the use of herbicide or antibiotic resistance genes. For many plant species, several of the used selectable marker genes yield a low transformation efficiency and many non-transgenic escapes. Thus more efficient selection systems are required.

The use of antibiotic and herbicide resistance is called negative selection due to that the non-transformed cells are greatly retarded in growth or even killed. Selectable marker genes which

can be used are for example the bialaphos resistance gene (bar) and the kanamycin or G418 resistance gene (NPTII). By growing the tissue on media containing for example kanamycin, in theory only the transformed cells will divide and take part in morphogenesis.

- 5 The NPTII gene encodes a neomycin phosphotransferase, which reduces the inhibitory action of kanamycin, neomycin, G418 and paromomycin by a phosphorylation reaction.

In contrast to negative selection research has also been done on  
10 positive selection, based on giving the transformed cells a metabolic advantage while the non-transformed cells are starving with a concomitant slow reduction in viability. An example is the use of the mannose or xylose based selection system as described e.g. in US 5,767,378.

15 *Solanum tuberosum* is one of the major target crops for genetic engineering. Main traits for potato are tuber quality, nutritional composition, starch quality, starch yield, insect and virus resistance. One trait with high market potential is the  
20 production of high amylopectin starch. The high amylopectin starch has an improved performance in the adhesive and paper industry compared to ordinary starch. In the paper production it will be used as a binder and for coating with printing quality better than latex used today.

25 *Solanum tuberosum* is a tetraploid plant with a high level of genetic heterozygosity. Conventional breeding of potato is therefore complicated because of segregation of important characteristics. Genetic engineering has during the last decade been an  
30 alternative to conventional breeding when it comes to improving potato varieties. A single trait or a combination of traits are more efficiently introduced by transformation than by using conventional breeding. Due to the fact that genetically modified potato plants can efficiently be vegetatively propagated the  
35 trait is fixed in the commercial line without the normal subsequent breeding.

*Agrobacterium tumefaciens* mediated transformation has successfully been achieved in potato since 1986. Almost exclusively the  
40 nptII gene has been used as a selection marker in potato.

The highest documented efficiency using nptII as selection marker in potato transformation is described by Libiakova, G. et al. in 2001. They have produced 120 leaf discs. Of those 81% produced  
45 one or more shoots. 90% of the shoots were confirmed transformants and showed resistance to kanamycin. This is equivalent to a

regeneration efficiency of 81%, an escape rate of 10% and yields a transformation efficiency of 73%.

It is an object of the present invention to provide an efficient  
5 method for transformation of potato plants with a low escape rate and a high transformation efficiency.

Leaf segments are the explants of choice used in Agrobacterium mediated potato transformation. Transgenic plants are regenerated  
10 via direct or indirect organogenesis under selective conditions. Cytokinines as growth hormones are mainly involved in provoking cell division and adventitious bud formation in in vitro potato leaf segments. When Imazamox - (RS)-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-methoxymethylnicotinic acid, an  
15 imidazolinone type herbicide - is applied in low concentrations it was surprisingly found that it possesses a cytokinin-like effect in tissue culture, increasing plant cell growth. At higher concentrations the substance is acting as an herbicide. The invention shows that Imazamox can efficiently be used as a  
20 selective agent in the recovery of transgenic potato plants and as a cytokinin-like substance it is supporting regeneration during transformation. This phenomenon classifies Imazamox as a unique selection agent that is combining both lethality for non-transgenic cells and improved growth for transgenic cells.  
25

The present invention is describing a new selection system for efficient recovery of transgenic plants. The invention is also describing an innovative selection system that is acting as  
restriction for untransformed cells and support for transformed  
30 ones. Finally the invention is presenting an efficient system for recovery of transgenic potato plants and yields only a very small escape rate.

The present invention relates to the use of a mutated AHAS gene  
35 conferring imidazoline type herbicide resistance resulting in a highly efficient selection system for production of transgenic potato lines.

The invention provides a key advantage by minimizing the escapes  
40 to almost zero of all shoots regenerated without interfering with the high number of shoots regenerated per explant.

Another advantage is that the method does not comprise a gene conferring resistance to an antibiotic.  
45

For example a mutated gene coding for a functional AHAS enzyme with tolerance to the imadazolinone type herbicides as described in Sathasiavan, K. et al, 1991 is used, see SEQ-ID No. 1. The mutated AHAS gene is used for insertion as a selection marker  
5 together with any other homologous or heterologous gene into the potato genome for production of transgenic potato plants. The AHAS enzyme naturally exists in higher plants and the use of a modified plant AHAS gene as selection marker conferring resistance to AHAS inhibiting herbicides does not result in the  
10 addition of a new biosynthetic function.

Furthermore

- 15 (i) the mutated AHAS gene can be of eukaryotic or prokaryotic origin;
- (ii) the mutated AHAS gene can be regulated by any promoter functional in plants;
- (iii) any plant transformation method for insertion of the gene constructs can be used;
- 20 (iiii) any mutations in the AHAS gene resulting in tolerance to an AHAS inhibiting herbicide can be used;
- (iiiii) any potato variety can be transformed.

A mutated AHAS gene with mutation S653N as described in  
25 SEQ-ID No. 1 and US 5,767,366 was cloned in a vector with the reporter gene beta-glucuronidase (GUS) (Jefferson, R.A. et al., 1987 ) and transformed to potato cells, see example 1 to 4. The GUS activity is tested in the transgenic lines produced and results in a transformation frequency of 93 to 100% of all  
30 transgenic shoots analysed. The transformation method produces a very high amount of shoots per explants, 3 to 5 independent transgenic shoots dependent on the variety used. With a calculation on transformation efficiency per explants the number can be as high as 500%. This is the highest transformation fre-  
35 quency of potato published. The mutated AHAS gene has also been used in combination with a *gbss* antisense gene for production of high amylopectin potato lines, see example 12. The AHAS selection system yielded in this case the same high trans-  
40 formation efficiency, which establishes the use of the AHAS gene as a selection marker also for commercially important traits.

The highest efficiency of *nptII* as a selection marker is 74% (see example 10) of totally analyzed shoots yielding a transformation efficiency of 370% at the maximum. Another selection system that  
45 can be used for potato transformation is the Mannose or xylose based selection system. Haldrup et al. (1998) have used the xylose isomerase gene in potato transformation, which yielded a

transformation efficiency of 29% based on number of explants producing GUS positive shoots. High percentage of shoots positively transformed has been published for sugar beet and tobacco when using a very high mannose concentration, however the high concentration of the selective agent inhibit the number of shoots per explant distinctively (Joersbo et al., 1998).

The process furthermore comprises the possibility that compounds with herbicidal activity inhibiting the AHAS synthase can be employed for the selection of transgenic potato plants by using different combinations of growth regulators as for example auxins, cytokinines and/or auxin or cytokinines conjugates for the regeneration into intact transgenic plants.

The method is suitable for transforming the species *Solanum tuberosum*.

The invention also relates to transgenic fertile *Solanum* plants themselves which have been generated by the method described and to their transgenic seeds.

The method according to the invention allows for the first time to transform plant cells of the genus *Solanum* with a transformation efficiency of more than 93% to be regenerated into transgenic plants as shown in example 9.

The method for the generation of stably transformed *Solanum* plants is composed of the following steps: (a) growing the plant to be transformed and obtaining the suitable explant, (b) transferring DNA sequences into plant cells, (c) selecting transformed plant cells with compounds which inhibit in non transformed plant cells the AHAS synthase and (d) regenerating these transformed plant cells into fertile transgenic plants.

A variety of methods is currently available for the transformation. The most frequently employed method for transforming dicotyledonous plants is the *Agrobacterium-tumefaciens*-mediated gene transfer. This method exploits the natural ability of the soil bacterium to integrate genetic material into the plant genome. Other suitable methods are for example protoplast transformation by polyethylene-glycol-induced DNA uptake, electroporation, sonication or microinjection, the transformation of intact cells or tissue by micro- or macroinjection into tissue or embryos, tissue electroporation, incubation of dry embryos in DNA-containing solution, vacuum infiltration of seeds and the biolistic gene transfer.

The use of *Agrobacterium tumefaciens* for the transformation of plants using tissue culture explants has been described by Horsch et al., Science 228(1985), 1229-1231; Fraley et al., Proc. Natl. Acad. Sci. USA 80(1983), 4803-4807 and Bevans et al., Nature 5 304(1983), 184-187. Many strains of *Agrobacterium tumefaciens* are capable of transferring genetic material into *Solanum* species such as for example the strains EHA 101, EHA105, LBA4404 and C58C1 with various disarmed Ti-plasmids.

10 The agrobacterial strain employed for the transformation of potatoes contains in addition to its disarmed Ti-plasmid a binary plasmid with the T-DNA to be transferred which contains a gene for selecting the transformed cells and the gene to be transferred. Both genes must be equipped with transcriptional and 15 translational initiation and termination signals. The binary plasmid can be transferred into the agrobacterial strain for example by electroporation or other transformation methods, see Mozo & Hooykaas, Plant Mol. Biol. 16(1991), 917-918. Coculture of the plant explants with the agrobacterial strain takes two to 20 three days.

Foreign genes can be expressed in a constitutive, inducible (by biotic and abiotic factors), tissue-specific or development-specific way. A relatively constitutive expression is achieved for 25 example in plants by the cauliflower mosaic virus 35S promoter. This expression has been described by Shewmaker et al., Virology 140, 281-288 (1985) and Gardner et al., Plant Mol. Biol. 6, 221-228(1986).

30 Both the direct and the indirect gene transfer are suitable transformation methods. The *Agrobacterium*-mediated transformation using a wide range of starting explants such as for example cotyledons, leaves, hypocotyls, shoots, roots, callus, mature and immature seeds or floral tissue can successfully be employed. The 35 gene transfer can be effected both by simple cocultivation/incubation or wetting with the agrobacterial strain and by a supporting vacuum infiltration of the explants with the bacterial culture in question. The use of feeder cultures as aids may be advantageous in the process. Each agrobacterial strain which con- 40 tains a Ti- or Ri-plasmid with the genetic information required for the transfer is suitable as vector for the transformation. Suitable agrobacterial strains are for example EHA101[pEHA101], EHA105[pEHA105], LBA4404[pAL4404], C58C1[pMP90] and C58C1[pGV2260]. Viral vectors also seem to be suitable for the 45 transformation of *Solanum*. Other methods for transferring genetic material into *Solanum* are for example the polyethylene glycol (PEG)-mediated protoplast transformation, the electroporation,

- the sonication or microinjection and the transformation of intact cells or tissue by micro- or macroinjection into tissue or embryos, tissue electroporation, incubation of dry embryos in DNA-containing solution, the vacuum infiltration of seeds, see also Eds. Galun, E. and Breiman, A. In: Transgenic Plants, Imperial College Press, 1997. The use of the particle gun for bombarding a wide range of explants such as for example leaves and callus, is also possible.
- 10 All gene transfer vectors which contain the border sequences necessary for transferring the T-DNA (left and right border, LB and RB respectively), carry one or more selectable marker or reporter genes under the control of suitable promoters and terminators and/or contain further useful genes or target genes, also under the control of suitable transcriptional and translational regulatory units, can be employed for the *Agrobacteria*-mediated transformation. The transfer of DNA sequences into the *Solanum* plant to be transformed or explants thereof is preferably done by *Agrobacterium-tumefaciens*-mediated gene transfer.
- 20 The regeneration medium according to the invention furthermore comprises growth regulators such as auxins and/or auxin conjugates, and cytokinins and/or cytokinin conjugates. Suitable auxins are both natural and synthetic auxins. The natural auxin is indoleacetic acid (IAA), examples of synthetic auxins are 3-indolebutyric acid (IBA), 1-naphthylacetic acid (NAA) and the herbicide 2,4-dichlorophenoxyacetic acid (2,4-D). Other herbicides which act as auxins are also feasible as growth regulators. Auxin conjugates are compounds of auxin (IAA) with aspartic acid, glucose, myoinositol and others. In the case of the cytokinins, again, natural cytokinins such as, for example, zeatin and synthetic components, such as 6-benzylaminopurine (BAP) and 6-furfurylaminopurine (kinetin) may also be employed. Zeatin riboside is frequently employed as cytokinin conjugate. The auxins and the cytokinins can be employed in each case as individual components, but also as auxin or cytokinin mixtures. The concentration of the individual growth regulators is 0.05 to 10 mg/l, preferably 0.1 to 5 mg/l.
- 40 The transformation method according to the invention can be applied, inter alia, to Solanaceae species as *Solanum tuberosum* (potato), *Lycopersicon esculentum* (tomato) and pepper.
- The mutant AHAS genes of the present invention confer resistance to imidazolinone herbicides. Types of herbicides to which resistance is conferred are described for example in US Patent Nos.: 4,188,487; 4,201,565; 4,221,586; 4,297,128; 4,554,013; 4,608,079;

4,638,068; 4,747,301; 4,650,514; 4,698,092; 4,701,208; 4,709,036;  
4,752,323; 4,772,311 and 4,798,619.

Mutant AHAS genes conferring resistance to AHAS inhibiting  
5 herbicides are described in US 5,013,659, US 5,141,870 and  
US 5,378,824.

Other mutant AHAS genes of the present invention could also  
confer resistance to sulfonylurea herbicides. Types of mutants  
10 which confer sulfonylurea resistance are described for example  
in US 5,853,973 and US 5,928,937.

Mutant AHAS genes conferring resistance to imidazolinone type  
herbicides are described in WO 00/26390 and US 5,767,366.

15 Furthermore Duggleby, R.G. and Pang, S.S. in Journal of Bio-  
chemistry and Molecular Biology 33(1), 1-36 (2000) describe muta-  
tions of the AHAS genes which could be used in the invention for  
conferring herbicide resistance to transgenic potato plants.

20 In WO 00/26390 additional genomic and cDNA sequences coding for  
an eukaryotic AHAS small subunit protein are disclosed. The DNA  
sequences and vectors are used to transform plants to produce  
transgenic plants which possess elevated levels of tolerance or  
25 resistance to herbicides such as imidazolinones.

It will be understood by those working in the field that the  
nucleic acid sequence depicted in SEQ-ID No. 1 is not the only  
sequence which can be used to confer imidazolinone-specific  
30 resistance. Also contemplated are those nucleic acid sequences  
which encode an identical protein but which, because of the  
degeneracy of the genetic code, possess a different nucleotide  
sequence. The invention also encompasses genes encoding AHAS  
sequences in which the above-mentioned mutation is present, but  
35 which also encode one or more silent amino acid changes in posi-  
tions of the molecule not relevant for resistance to herbicides  
or to the catalytic function. Also contemplated are gene  
sequences from other imidazolinone resistant monocot or dicot  
plants which have a mutation in the corresponding region of the  
40 sequence.

For example, alterations in the gene sequence which results in  
the production of a chemically equivalent amino acid at a given  
site are contemplated; thus, a codon for the amino acid alanine,  
45 a hydrophobic amino acid, can readily be substituted by a codon  
encoding another hydrophobic residue, such as glycine, or may  
be substituted with a more hydrophobic residue such as valine,

leucine or isoleucine. Similarly, changes which result in a substitution of a negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be  
5 expected to produced a biologically equivalent product.

The invention also encompasses chimaeric genes, in which the substituted portion of the corn or Arabidopsis AHAS gene is recombined with unaltered portions of the normal AHAS gene, in  
10 which the substituted portion of the corn or Arabidopsis AHAS gene is recombined with unaltered portions of the normal AHAS gene from other species. Thus, throughout the specification and claims, wherever the term "herbicide resistant AHAS gene" is used, it is intended to cover each of these alternate embodiments  
15 as well as the sequence of SEQ-ID No. 1.

For expression of the mutated AHAS gene conferring herbicide resistance in potato the following promoters can be used:

20 the tuber specific gbss-promoter from potato described in WO 92/11376, and the patatin promoter described in Rocha-Sosa et al., 1989 EMBO J. 8:23-29;

the light inducible promoter: cytosolic FBPase from potato  
25 described in WO 98/18940;

the octopine promoters (US 5,428,147); the triple OCS enhanced vATPase c1 promoter from Beta vulgaris (Plant Mol Biol (1999) 39: 463-475);

30 Constitutive promoters: for reference see Benfey et al., EMBO J. 8 (1989), 2195-2202; the 35S promoter (Franck et al., Cell 21, (1980), 285-294) or enhanced versions; the 19S promoter, see US 5,352,605 and WO 84/02913;

35 the RUBISCO small subunit SSU promoter: see US 4,962,028;

plastid specific promoter: see e.g. the RNA polymerase promoter  
40 in WO 95/16783 and WO 97/06250 or the clpP-promoter in WO 99/46394;

the AHAS promoter as described in US 6,025,541;

45

Other promoters for the expression of genes in the leaf, in the callus, in specific tissues as e.g. in the tubers or other parts of the potato plant could also be used.

- 5 The invention can especially be carried out by using the AHAS promoter, the AHAS resistance gene S653N as described in US 5,767,366 and the AHAS terminator from *Arabidopsis thaliana* cut from pAC321 as described in example 3.
- 10 The *Arabidopsis* AHAS gene (S653N) used for transformation and selection contains most of the common restriction sites for cloning such as HindIII, BamHI, EcoRI, SstI, BglII and EcoRV. Alteration of the molecular composition of the AHAS gene can be performed to eliminate restriction sites without altering the amino
- 15 acid sequence of the resulting protein. Special care must be taken not to alter the codon usage profile from the one found in *Arabidopsis*, which could lead to reduced translation efficiency.

- The designed gene can be made synthetically and accommodates
- 20 suitable restriction sites at 5' and 3' ends for cloning.

Comparative studies have to be performed in potato in order to confirm that the synthetically made gene does not differ from the original mutated AHAS gene in conferring herbicide resistance.

- 25 For selection of transgenic potato plants chemical compounds inhibiting the AHAS enzyme can be used. Useful compounds are the imidazoline type herbicides. Especially useful compounds are selected from the group consisting of imazethapyr (Pursuit™), imazamox (Raptor™), imazamethabenz (Assert™), imazapyr (Aresenal™),
- 30 imazapic (Cadre™) and imazaquinon (Scepter™).

- For selection of transgenic plants chemical compounds as described in the review article by Duggleby, R.G. and Pang, S.S.
- 35 in *Journal of Biochemistry and Molecular Biology* 33(1), 1-36 (2000) can be used.

- The mutated AHAS gene as described in SEQ-ID No. 1 has been established as a selection marker in five different potato
- 40 varieties, see table 2 and 3, and can thus be used for transformation of potato varieties in general. Two different selection pressures, using 0.3 and 0.5  $\mu\text{M}$  Imazamox, have been tested, see table 2. For selection on 0.3  $\mu\text{M}$  Imazamox the transformation efficiency is ranging from 7 to 76 % depending on the variety
- 45 used. When increasing the selection to 0.5  $\mu\text{M}$  Imazamox the transformation rate is increasing to 93 to 100% respectively. The high transformation efficiency at 0.5  $\mu\text{M}$  Imazamox is independent from

the variety used. The fifth variety Seresta has as can be seen in table 3 and 4 a transformation efficiency of 98 to 100% at 0.5  $\mu$ M Imazamox.

- 5 The invention furthermore relates to a plant expression vector comprising SEQ-ID No. 1 wherein the heterologous DNA sequence encodes a peptide, protein, antisense-, sense-RNA, viral RNA or ribozyme.
- 10 The invention furthermore relates to a plant expression vector comprising SEQ-ID No. 1 wherein the heterologous DNA sequence contains information that causes changes in the carbohydrate concentration and the carbohydrate composition of regenerated potato plants.
- 15 The invention furthermore relates to a plant expression vector comprising SEQ-ID No. 1 wherein the heterologous DNA sequence contains information that causes the increased production of amylose type starches by using gene constructs as described in
- 20 WO 92/11375, WO 92/14827, WO 96/34968 and WO 97/20040.

- The invention furthermore relates to the use of a DNA sequence SEQ-ID No. 1 or a DNA sequence comprising a nucleotide sequence which hybridizes to a complementary strand of the nucleotide SEQ-
- 25 ID No. 1 or a DNA sequence comprising a nucleotide sequence which is degenerated to the nucleotide sequence SEQ-ID No. 1 or a DNA sequence being a derivative, analogue or fragment of a nucleotide sequence encoding a protein possessing AHA synthase activity and conferring resistance to AHA synthase inhibitors.
- 30 The experiment shown in example 12 with the high amylopectin trait establishes the use of the AHAS resistance gene conferring resistance to herbicides inhibiting the AHAS enzyme as a selection marker also for commercially important traits. The trans-
- 35 formation efficiency is 98 and 100% respectively. Analysis of the amylopectin/amylose ratio shows that in the transgenic shoots selected the amylopectin content is higher than in non transgenic control shoots. This is proof that by using the invention as described commercially relevant transgenic potato plants can be
- 40 generated.

- The invention now having been generally described will be more readily understood by reference to the following examples, which are included for the purpose of illustration only, and are not
- 45 intended to limit scope of the present invention.

## Example 1

Test for transgenicity - Detection of glucuronidase expression

- 5 Microtubers of the regenerated in vitro potato plants were subjected to a qualitative glucuronidase (GUS) enzyme detection by infiltrating cut microtubers for 1 minute in vacuo with the GUS substrate 5-bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronic acid (X-GlcA) in a 100 mM sodium phosphate buffer, pH 7.0, which contained
- 10 10 mM EDTA, 0.1% Triton X100 and 10 mM DTT, and subsequently incubating them for approximately 15 hours at 37°C. Microtubers showed an intense blue coloration, which proves that the reporter gene is expressed in *Solanum*.

## 15 Example 2

Investigation of selection pressure

- Potato leaf segments from in vitro propagated plants of four
- 20 different potato varieties Prevalent, Producent, Karnico and Desiree were tested for natural tolerance on different concentrations of the imidazolinone herbicide Imazamox. It is important to choose a selection concentration at which the potato leaf tissue is viable long enough to regenerate shoots but have a high
- 25 enough concentration to prevent the regeneration of untransformed shoots. Doses 0.1, 0.5, 1, 5, 10, 20, 30 and 50  $\mu$ M Imazamox were tested.

- Fully expended potato leaves are diagonally cut in 2 pieces
- 30 and precultivated on MC-plates for 2 to 3 days at 23 to 24°C, see table 1.

- The leaf tissues are transferred on MS300 medium for additional 2 days and cultivated under modest light at 23 to 24°C simulating
- 35 a co-cultivation step. Subsequently the leaf segments are moved to MS400 plates containing 400mg/l Claforan. Claforan is added in order to suppress growth of *Agrobacterium tumefaciens*. Therefore the regeneration of shoots on different concentrations of Imazamox need to be monitored in presence of Claforan. Selection
- 40 is taking place within 4 to 5 days when the explants are moved to MS400 medium supplemented with 400mg/l Claforan and Imazamox at above concentrations.

- The explants are transferred to fresh selection medium every
- 45 fortnight.

5	MC plates  MS300 plates with 1.5–2 ml liquid MS100 medium and covered with one sterile filter paper	MS300  X MS–medium 2mg/l NAA (naphtyl acetic acid) 1mg/l BAP (6–benzyl amino pyridine) 3% (w/v) sucrose pH 5.2
10	MS 10  4.4 g/l MS–medium (Murashige and Skoog) 1% (w/v) sucrose pH 5.8	MS400  4.4 g/l MS–medium 2mg/l zeatine 0.01mg/l NAA (naphtyl acetic acid) 0.1 mg/l GA3 (giberellic acid) 10% (w/v) sucrose 400 mg/l claforan Imazamox or kanamycin pH5.8
15	MS 30  4.4 g/l MS–medium 3% (w/v) sucrose pH 5.8	Microtuber medium  4.4 g/l MS–medium 2.5 mg/l kinetin 0.5 mg/l ABA (abscisic acid) 8% sucrose 200 mg/l claforan
20	MS100  4.4 mg/l MS–medium 30g/l sucrose 0.5 mg/l thiamin–HCl 0.5 mg/l pyridoxin–HCl 1 mg/l nicotinacid 0.5 mg/l kinetin 29.8 mg/l ferrous sulfate hepta hydrate 1 mg/l 2,4–Dichlorophenoxyacetic acid 2 g/l caseinhydrolysate pH 5.2	

30

Table 1: Media used for potato transformation.

Example 3

35 Construction of binary vectors pAHASGUS, pAP1 and pAP2

For all constructs the same AHAS mutant gene with mutation S653N originating from *Arabidopsis thaliana* was used (Sathasivan, K. et al., 1991).

40

pAHASGUS

AHAS gene with mutation S653N as described in SEQ ID No. 1, originating from *Arabidopsis thaliana*, was used together with the  
45 nos promoter ( Herrera, L. et al., 1983 ) and the OCS terminator (Wesley S. V. et al., 2001) in the binary vector pGPTVkan (Becker, D. et al., 1992 ). The pBIN19 based pGPTVkan was

digested with ApaI and BamHI discarding the *nptII* gene. The remaining 12600 bp contained from right border the nos terminator, the *uidA* gene (also known as GUS), the cloning cassette, the nos promoter and the pAg7 terminator. The AHAS gene with  
5 mutation S653N and the OCS terminator was ligated to the digested pGPTVkan and named pABA1, see figure 1.

For expression of the GUS gene a tuber specific gbss promoter 988bp (WO 92/11376) was cloned at the SmaI site of pABA1.

10

The resulting pAHASGUS construct was used for *Agrobacterium tumefaciens* transformation of potato, where regenerated plantlets were analysed for GUS expression to determine the transformation efficiency.

15

pAP1

AHAS gene with mutation S653N, driven by the *Arabidopsis thaliana* AHAS promoter as described in US 5,750,866, was used as a selec-  
20 tion gene for co-transformation with a granule-bound starch synthase (gbss) antisense gene (EP-A 0 563 189).

A 293 bp nos terminator was cut out from pHaxw0 and ligated to pBluescript with EcoRI. The 2952 bp gbss promoter and gbss gene  
25 in antisense direction (EP-A 0 563 189) was cut from pHaxw0 and ligated in front of the nos terminator with HindIII resulting in pMJ2. The gbss complex was cut out from pMJ2 with XbaI-XhoI and ligated to pSUN1 (WO 02/00900) cut with XbaI-SalI resulting in the construct named pMJ3.

30

pAC321 was cut with XbaI which yielded a fragment of 5717 bp containing AHAS promoter, AHAS gene and AHAS terminator all originating from *Arabidopsis thaliana* as described in SEQ-ID No. 1.

35 SEQ-ID No.1 contains the following elements:

1-2483 At AHAS promoter  
2484-4496 At AHAS gene  
4497-5717 At AHAS terminator

40

The pAC321 fragment was ligated to pMJ3 opened with XbaI.

The resulting pAP1 (figure 1) construct was used for *Agrobacterium tumefaciens* transformation of potato, which were analysed for  
45 high amylopectin starch quality.

pAP2

AHAS gene with mutation S653N, driven by the nos promoter (US 6,174,724), was used as a selection marker for co-transformation with a gbss antisense gene (WO 92/11376).

A nos promoter of 600 bp was cut out from pGPTVKan (Becker et al., 1992) with HindIII, blunted and cut with BglII. The fragment was ligated to pBluescript (Stratagene) cut with SpeI, blunted and cut with BamHI. The construct was named pPnos2.

A 2986 bp fragment of pACGH101 containing the AHAS gene with mutation S653N as described in SEQ ID No. 1 and OCS terminator was cut out with PstI and PvuII. The AHAS gene originates from *Arabidopsis thaliana* see pAP1. The pACGH101 fragment was ligated to pPnos2 cut with PstI and EcoRV and named pPnoas.

The 3406 bp AHAS complex was cut out from pPnoas with XbaI-PvuII and ligated to pMJ3 (see pAP1) cut open with XbaI.

The resulting pAP2 (figure 2) construct was used for *Agrobacterium tumefaciens* transformation of potato, which were analysed for high amylopectin starch quality.

**25** Example 4

## Transformation method

Fully expanded leaves from in vitro propagated potato plants are diagonally cut in 2 pieces and precultivated on MC-plates for 2 to 3 days at 23 to 24°C.

*Agrobacterium tumefaciens* containing pAHASGUS is grown in YEB medium containing 25 µg/ml kanamycin and 100 µg rifampicin and grown over night on constant shaking (200 rpm) at 28°C.

The *Agrobacterium* culture is prepared for infection by dilution 1:20 with MS10 medium. The leaf explants are infected for 8-10 min in the bacterial solution and afterwards drained on filter paper for 5 to 20 seconds. The leaf segments are placed on the MS300 plates for 2 days co-cultivation under modest light at 23 to 24°C. At the end of co-cultivation the leaf segments are moved to M400 plates containing 400 mg/l Claforan to suppress bacterial growth. After 4 to 5 days the explants are moved to selection medium MS400 supplemented with 400 mg/l Claforan and Imazamox at the previously detected concentrations of 0.3 or 0.5 µM.

## 17

Leaf segments are transferred to fresh MS 400 selection medium every fortnight. The regenerated putative transgenic shoots are collected and cultivated on MS30 plates with 200mg/l Claforan aiming at shoot elongation.

5

When the shoots are 3 to 5 cm long, 1 to 2 cm are cut off and grown on microtuber medium in the dark at 25°C. After 2 to 5 weeks microtubers are produced. Putative transgenic plants are analysed for GUS expression in microtubers to determine the transformation

10 efficiency.

## Example 5

GUS expression analysis for determination of the transformation  
15 efficiency

A thin slice from each microtuber was placed in a microtiter plate. The slices were incubated with 1 mM X-gluc in 50 mM NaH<sub>2</sub>PO<sub>4</sub> pH 7.2 at 37°C for approximately 1 hour. A very distinct blue  
20 colour appeared on tuber slices if a successful integration and expression of the GUS gene was achieved.

## Example 6

25 Transformation efficiency determination of pAP1 and pAP2 by PCR analysis

Transgenic shoots were distinguished from non-transgenic shoots - for example escapes - by using PCR. DNA was extracted according  
30 to DNeasy 96 Plant protocol (Qiagen, Germany). In a 96 well microtiter plate, 10 to 15 mg leaf tissue was added to each well together with a 5 mm steel ball each well then representing one individual shoot. The plates were frozen in N<sub>2</sub>(l) before homogenisation. The homogenisation was done at 30 Hz in a Mixer-  
35 mill 300 for 1 min. The DNA was at the end of the extraction protocol eluted in 75 ml H<sub>2</sub>O.

Specific primers were used to amplify a fragment of the mutant S653N AHAS gene. Successful integration of the mutant AHAS gene  
40 results in a fragment of 509 bp upon PCR amplification using the specific primers.

Forward primer: AHAS1\_frw:           AACACAACATCTTCTTCGATC  
Reverse primer: AHAS1\_rev:         TAACGAGATTTGTAGCTCCG

45

The PCR reactions were done with the extracted DNA setup and run as follows:

Reaction:

<b>5</b>		
	10x PCR Mix	2,0 µl
	Primer frw (25 µM)	0,4 µl
	Primer rev (25 µM)	0,4 µl
	dNTPs (10 mM)	0,4 µl
<b>10</b>	RedTAQ (Sigma)	1,0 µl
	Templat (~20 ng/µl)	4,0 µl
	H <sub>2</sub> O	11,8 µl

PCR program:

<b>15</b>		
	94°C	30 s
	59°C	30 s x 29 cycles
	72°C	30 s
	72°C	7 min
<b>20</b>	8°C	Hold

Negative and positive control was included in all runs. The reactions were analysed on 1.5% agarose gels.

#### **25** Example 7

Starch quality analysis of amylopectin/amylose ratio

The screening for an altered amylopectin/amylose ratio was per-  
**30** formed on transgenic lines selected for high quality integration of the intended genetic insert. The screening was done by staining with iodine (Lugol's solution: (6.7 g/l KI + 3.3 g/l I<sub>2</sub>) and (glycerol); ratio 1:1). Iodine stains starch containing amylose blue and starch exclusively containing amylopectin red-brown. A  
**35** microtuber was crushed and a few drops of Lugol's solution were added. The starch amylopectin quality was analysed under the microscope according to the staining colour of the starch.

Example 8

**40**

Leaf tissues were tested for the use of the AHAS resistance gene contained in SEQ ID No. 1 and for the selection of transgenic potato plant cells as described above. Four different potato cultivars Prevalent, Producent, Karnico and Desiree were cultivated  
**45** on different concentrations of Imazamox, an imidazolinones type herbicides. It is essential to use the proper concentration for selection to yield an acceptable number of shoots per explant

without high escape rate. The lethality of various Imazamox concentrations on regeneration was evaluated.

Lethal dose for regeneration was detected within 0.1 to 1  $\mu\text{M}$  ranges of concentrations. Slight variation in lethality was detected between varieties. No regeneration occurs when 0.5  $\mu\text{M}$  Imazamox was included in the medium. The concentrations 0.3  $\mu\text{M}$  and 0.5  $\mu\text{M}$  are selected for transformation experiments providing a lethality of 80 and 100% in regeneration.

10

Example 9

Transformation of four different potato varieties (Prevalent, Producent, Kuras and Desiree) was performed with pAHASGUS as described above. For selection a concentration of 0.3 or 0.5  $\mu\text{M}$  Imazamox was used. Each shoot was analysed according to GUS activity and the number of GUS positive shoots in total is listed in table 2. A distinct blue colour appears on the microtuber slices if a successful integration of the GUS gene was achieved. From each explant 3 to 5 independent shoots were regenerated depending on the variety used. In all experiments the efficiency of positive shoots per explants is stated and the figures in table 2 have to be multiplied 3 to 5 times, yielding a transformation efficiency of up to 500%. Compared with selection systems used previously in potato plants the big advantage of the invention described is that together with a very high efficiency the number of shoots per explant is very high. For all four varieties transformed an extremely high transformation efficiency of up to 93% to 100% of all shoots regenerated was achieved at an Imazamox concentration of 0.5  $\mu\text{M}$ . The experiment using 0.3  $\mu\text{M}$  of Imazamox for selection shows how important the Imazamox concentration is for an efficient transformation rate. This can be seen from the results for variety Kuras in which a transformation efficiency of 7% at 0.3  $\mu\text{M}$  Imazamox is increased to 93% at the 0.5  $\mu\text{M}$  level.

40

45

Variety	Construct	Selection Imazamox ( $\mu\text{M}$ )	Selection Imazamox (mg/l)	Number analysed	Number of positive	% positive
Desiree	pAHASGUS	0.3 $\mu\text{M}$	0.0915mg/l	19	12	63
5 Prevalent	pAHASGUS	0.3 $\mu\text{M}$	0.0915mg/l	26	16	62
5 Producent	pAHASGUS	0.3 $\mu\text{M}$	0.0915mg/l	25	19	76
5 Kuras	pAHASGUS	0.3 $\mu\text{M}$	0.0915mg/l	58	4	7
10 Prevalent	pAHASGUS	0.5 $\mu\text{M}$	0.1525mg/l	23	23	100
10 Producent	pAHASGUS	0.5 $\mu\text{M}$	0.1525mg/l	40	38	95
10 Kuras	pAHASGUS	0.5 $\mu\text{M}$	0.1525mg/l	33	32	97
10 Desiree	pAHASGUS	0.5 $\mu\text{M}$	0.1525mg/l	29	27	93

Table 2: Transformation efficiency of four different varieties with 0.3 and 0.5  $\mu\text{M}$  Imazamox used as the selective agent.

15

#### Example 10

In a similar experiment compared to example 9 the *nptII* gene was used as selection marker being controlled by the same promoter nos as in pAHASGUS. The shoots are selected on 50 mg/l kanamycin, which is a standard kanamycin concentration used for potato transformation as described in Ooms, G. et al., 1987 and Tavazza, R. et al., 1988. The transformation frequency was much lower compared to when using a mutated AHAS gene e.g. the S653N gene as selection marker. The highest transformation efficiency based on analysis of the same GUS gene driven by the same GBSS promoter resulted in 74% transformed shoots regenerated.

#### Example 11

30

The successful use of a mutated AHAS gene as a selection marker has been demonstrated using the promoter from the wild-type *Arabidopsis thaliana* AHAS gene (pAP1 construct) as well as a recombinant nos promoter (pAP2 construct) in the potato variety Seresta, see table 3. This shows that any promoter with sufficient expression can be used in connection with the mutated AHAS gene and thus the invention could be used with different promoters as regulatory elements for the mutated AHAS gene successfully yielding selection of transgenic shoots on selective medium. The AHAS gene can be driven by different promoters such as e.g. the AHAS-, nos-, 35S-, Rubisco- or vATPase - promoter.

45

Variety	Construct	Promoter	Number shoots analysed	Number shoots positive	% positive
Seresta	pAP1	A.t AHAS	50	49	98
Seresta	pAP2	nos	120	117	98

Table 3: Transformation efficiency of the mutated AHAS gene driven by two different promoters: *Arabidopsis thaliana* (A.t.) AHAS promoter and the nos promoter.

#### Example 12

The mutated AHAS gene conferring resistance to Imazamox was used as a selection marker for the transformation of a commercially important trait to *Solanum tuberosum* (variety Seresta) resulting in potato lines producing amylopectine type starch. The starch component amylose is synthesized by granule-bound starch synthase (GBSS). Inhibition of the gene coding for GBSS directs starch production completely to amylopectin. An antisense gene fragment inhibiting the expression of gbss as described in EP-A 0 563 189 was co-transformed with the mutated AHAS gene conferring resistance to Imazamox.

The gbss promoter was used to drive the gbss antisense gene fragment. The Nos promoter or the AHAS promoter was used to drive the AHAS selection gene and resulted in a total transformation efficiency per shoots analysed of 98 to 100%, see table 4.

For restriction map of the T-DNA of the pAP1 and pAP2 constructs used, see figure 1 and 2. There were 3 to 5 independent shoots produced per explant yielding a transformation efficiency of 280 to 480% or up to 500% per explant. There were no differences in transformation efficiency between the different promoters and *Solanum tuberosum* varieties used.

Experiment	Variety	Construct	Number shoots analysed	Number shoots positive	% positive	high amylopectin content
1	Seresta	pAP2	120	117	98	+
2	Seresta	pAP1	50	49	98	+
3	Seresta	pAP2	30	30	100	+
4	Seresta	pAP2	3	3	100	+

Table 4: pAP1 and pAP2 transgenic lines analysed for transformation efficiency and high amylopectin quality.

Starch quality analysis was done with microtubers of the positive shoots with high quality integration of the intended genetic insert. The starch was mixed with iodine and the shoots exclusively containing amylopectin stains red-brown compared to the starch containing amylose that stains blue.

## References

- Hawkes, T.R., in Prospects for Amino Acid Biosynthesis Inhibitors in crop protection and Pharmaceutical Chemistry, pp131-138,  
5 Britttish Crop Protection Council Monograph No 42, Surrey, U.K. (1989)
- Herrera-Estrella, L., Depicker, a., Van Montagu, M., Schell, J.,  
Expression of chimaeric genes transferred into plant cells using  
10 Ti-plasmid-derived vector, Nature 303:209-213 (1983)
- Becker, D., Kemper, E., Schell, J., Masterson, R., New plant  
binary vectors with selectable markers located proximately to the  
left T-DNA border, Plant Molecular Biology 20:1195-1197, (1992)  
15
- Jefferson, R.A., Kavanagh, T.A., Bevan, M.W., GUS fusions: b-glu-  
curonidase as a sensitive and versatile gene fusion marker in  
higher plants, EMBO Journal 6: 3901-3907 (1987)
- 20 Joersbo, M., Donaldson, I., Kreiberg, J., Guldager Petersen, S.,  
Brunstedt, J., and Okkels, F.T., Analysis of mannose selection  
used for transformation of sugar beet, Molecular Breeding  
4:111-117 (1998)
- 25 Libiakova, G., Jorgensen, B., Palmgren, G., Ulskov, P.,  
Johansen, E., Efficacy of an intron containing kanamycin resist-  
ance gene as a selectable marker in plant transformation, Plant  
Cell Rep 20:610-615(2001)
- 30 Ooms, G., Burell, M.M., Karp, A., Bevan, M., Hille, J., Genetic  
transformation in two potato cultivars with T-DNA from disarmed  
*Agrobacterium*,  
Theoretical and Applied Genetics (1987) 73:744-750,
- 35 Ray, T.B., Plant Physiology (1984), 75, 827-831
- Sathasivan, K., Haughn, G.W., Murai, N., Plant Physiology  
97(1991), 1044-1050
- 40 Shaner, D.L., Andersson, P.C., Stidham, M.A., Plant Physiology 76  
(1984), 545-546
- Subrimanian, M.V., Gerwick, B.C., in Biocatalysis in Agricultural  
Biotechnology, pp277-288, ACS Symposium Series No389, American  
45 Chemical Society, Washington DC (1989)

Tavazza, R., Tavazza, M., Ordas, R.J., Ancora, G., Benvenuto, E., Genetic transformation of potato (*Solanum tuberosum*): An efficient method to obtain transgenic plants, *Plant Science*, 59 (1988), 175-181

5

Umbarger, H.E., in *Synthesis of Amino Acids and Proteins*, (1975) pp1-56, MTP International Review of Science, Butterworth, London.

Wesley S.V., Helliwell C.A., Smith N.A., Wang M.B., Rouse D.T.,  
10 Liu Q., Gooding P.S., Singh S.P., Abbott D., Stoutjesdijk P.A.,  
Robinson S.P., Gleave A.P., Green A.G., Waterhouse P.M.,  
Construct design for efficient, effective and high-throughput  
gene silencing in plants, *Plant J* 2001 Sep;27(6):581-90

15 Haldrup A, Petersen, GP, Okkels FT (1998) The xylose isomerase  
gene from *Thermoanaerobacterium thermosulfurogenes* allows effec-  
tive selection of transgenic plant cells using D-xylose as the  
selection agent. *Plant Molecular Biology* 37:287-296

20

25

30

35

40

45

## Claims

1. A method for transformation of potato plants by transforming  
5 potato plant cells with an expression vector comprising
- a) regulatory sequences of a promoter active in plants;
  - 10 b) operably linked thereto a DNA sequence encoding a protein with the biological activity of an AHA synthase resistant to inhibitors of potato plant wildtype AHA synthase; and
  - 15 c) operably linked thereto regulatory sequences which serve as transcription termination and/or polyadenylation signals in plants,
- selecting for AHA synthase inhibitor resistant cells and  
regenerating them to transgenic plants.
- 20 2. A method for transformation according to claim 1, wherein the expression vector comprises a DNA sequence according to SEQ-ID No. 1.
3. A method for transformation according to claim 1, wherein the  
25 DNA sequence encoding a protein with the biological activity of an AHA synthase resistant to inhibitors of potato plant wildtype AHA synthase is selected from the group consisting of
- 30 a) DNA sequence comprising a nucleotide according to SEQ-ID NO. 1;
  - b) DNA sequence comprising a nucleotide sequence which hybridizes to a complementary strand of the nucleotide  
35 sequence a)
  - c) DNA sequence comprising a nucleotide sequence which is degenerate to the nucleotide sequence of a) and
  - 40 d) DNA sequence being a derivative, analogue or fragment of a nucleotide sequence of a), b) or c) and encoding a protein possessing AHA synthase activity and conferring resistance to AHA synthase inhibitors.

45

Zeichn. + Sequ.

4. A method for transformation according to any of claims 1 to 3, wherein the AHA synthase promoter from *Arabidopsis thaliana* or the nos promoter is used.
- 5 5. A method for transformation according to any of claims 1 to 4, wherein the AHA synthase terminator from *Arabidopsis thaliana* or the OCS terminator is used.
6. A method for transformation according to any of claims 1 to 5, wherein for selection a imidazolinone type herbicide is used.  
10
7. A method for transformation according to claim 6, wherein for selection (RS)-2-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-5-methoxymethylnicotinic acid is used.  
15
8. A plant expression vector according to claim 1 or 2 additionally comprising a heterologous DNA sequence.
9. A plant expression vector according to claim 8, wherein the heterologous DNA sequence encodes a peptide, protein, antisense-, sense-RNA, viral RNA or ribozyme.  
20
10. A plant expression vector according to claim 9, wherein the heterologous DNA sequence contains information that causes changes in the carbohydrate concentration and the carbohydrate composition of regenerated potato plants.  
25
11. A plant expression vector according to claim 10, wherein the heterologous DNA sequence contains information that causes the increased production of amylopectin type starches.  
30
12. A plant expression vector according to claim 10, wherein the heterologous DNA sequence contains information that causes the increased production of amylose type starches.  
35
13. A transgenic potato plant cell produced by the method for transformation according to any of claims 1 to 7 and containing a plant expression vector according to any of claims 8 to 12.  
40
14. A transgenic potato plant produced by the method of transformation according to any of claims 1 to 7, wherein the regenerated plant exhibits an elevated resistance to imidazolinone type herbicides.  
45

15. Harvest product of the transgenic potato plant according to any of claims 13 and 14 comprising a DNA sequence SEQ ID No. 1 or a DNA sequence according to claim 3.
- 5 16. Harvest product according to claim 15 wherein the harvest product is a tuber.
17. Propagation material of transgenic potato plants comprising a DNA sequence SEQ ID No. 1 or a DNA sequence according to claim 3.
- 10
18. Use of a DNA sequence SEQ ID No. 1 according to claim 2 or a DNA sequence according to claim 3, or a plant expression vector according to of any of claims 8 to 12 in potato plant cells, potato tissue cultures, potato plants and/or potato plant breeding.
- 15
- 20
- 25
- 30
- 35
- 40
- 45




Fig. 1. T-DNA of pAP1 construct




Fig. 2. T-DNA of pAP2 construct

## SEQUENCE LISTING

<110> BASF Plant Science GmbH

<120> Use of AHAS mutant genes as selection marker in potato transformation

<130> NAE387-02

<140> 0093-53700

<141> 2002-07-08

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 5717

<212> DNA

<213> *Arabidopsis thaliana*

<220>

<221> CDS

<222> (2484)..(4493)

<400> 1

tctagattat gtatcccaca ctttcattaa caatataatc gcatataaat gaaaaatcgt 60  
ttccaggata atatcttgat gaaatctcat attattgttc gtactcggat tgatggtgaa 120  
ggcttgaagc gcttcaaatt atagaccaga ttatttaagt ttttcttttg tttactccat 180  
atcaatttga tccattatac tacctaagaa aatttaggta acatagaatt atttattggt 240  
atagtaaaaa aaaggaaaac cacaaaaata atctactttt acgtatatac tattttcatg 300  
acataagtaa ttaagttgta caactttttt ttaatgaaaa gagagagtaa atttatcatg 360  
ttcatgtgta gttacctcgt gaataaccga cggttatata gacgcctaac atgaattggt 420  
cagttgaaga cagttcaaaa catgtgtttc actctaaaat cctcaacaaa aaaaaagtgt 480  
taaaatttgt aaacctcttt caagcaaaaa aagaaaaagt gttagaatcc caagattcct 540  
tcataatccg gaatcttggc tgaaaacgta taaaagagat tgacgtagta acaaggagtc 600  
ttggtagtct tccatgcttt ttatcctttt ttgtcatgga accatgattt ggttaccatt 660  
tattatgtaa ccgaaatctt cattgtaata atgaatattt aaatcttagt caaaaaaaaa 720  
caaaaaaaaa caaggagtct tgtcttcggt ctcaaatttc agagctcttg cacttttcaa 780  
gagttttact ttgatgagtg agacatttgt ctttttagtg tttattttct aaacttaaaa 840  
tagtagcatc aacatcactc aattataatt ctttaagatgt ttagtagaaaa tattttatag 900  
atggaaagta atcgatatta agacaaataa gaaaccaaac cggactttgt gttcagaccg 960

aatcaaactct gaattggaga aattatggtg gaggcgaaag tcaacggaac taaagtataa 1020  
aaccaaatgt caaaaataaa acccaatttt catccttaaa cgaacctgct gaaaccctaa 1080  
tttcgattac caattccgat ctaaaaagaa gtcattggaag ccattgattc cgcaatcgat 1140  
cctctcagag atttcgctaa gagcagtgtt cgtctcgtcc agcgctgtca caaacccgat 1200  
cgcaagggtta acgccttttc tcaaaaaaat ctcaatttccg atttttgatc thtagattag 1260  
ggttttctga aattttgata tcatttgtaa ttgaattggt taccagaatt cacgaaagta 1320  
gctgtgcgta cggcgattgg atttgtggtg atgggattcg ttggattctt cgtgaagctc 1380  
gttttcatcc caatcaaca catcatcggt ggatcttctt agttagtagt tttctttacg 1440  
aggtaattga tctcgcatta tatacttaca ttttggttat gttacttgac atatagtcac 1500  
tgattcaata gttctgtaa ttcttttaa gatcattttg actagaccac attcttggtt 1560  
cattcctcaa taatttgtaa tcatattggt ggatatagaa gtagattggt tatagatcag 1620  
atagtggag actttaggat gaatttcagc tagttttttt ttttggtta ttgtctcaaa 1680  
agattagtgc tttgctgtct ccattgcttc tgctatcgac acgcttctgt ctcttctgtat 1740  
ctttattata tctattcgtc ccatgagttt tgtttggtct gtattcgttc gctctggtgt 1800  
catggatgga gtctctgttc catgtttctg taatgcatgt tgggttggtt catgcaagaa 1860  
atgctgagat aacactcat ttgtgaaagt ttctaaactc tgaatcgcgc tacaggcaat 1920  
gctccgagga gtaggaggag aagaacgaa caaacgacat taccagccct ttgaggaagc 1980  
tcttagtttt gttattggtt ttgtagccaa attctccatt ctattccat tttcacttat 2040  
ctcttggtcc ttatagacct tataagtttt ttattcatgt atacaaatta tattgtcatc 2100  
aagaagtatc tttaaaactc aaatctcaaa tcaccaggac tatgtttttg tccaattcgt 2160  
ggaaccaact tgcagcttgt atccattctc ttaaccaata aaaaaagaaa gaaagatcaa 2220  
tttgataaat ttctcagcca caaattctac atttaggttt tagcatatcg aaggctcaat 2280  
cacaataca atagatagac tagagattcc agcgtcacgt gagttttatc tataaataaa 2340  
ggaccaaaaa tcaaatcccg agggcatttt cgtaatccaa cataaaacc ttaaacttca 2400  
agtctcattt ttaaacaaat catgttcaca agtctcttct tcttctctgt ttctctatct 2460  
cttgctcatc tttctctga acc atg gcg gcg gca aca aca aca aca aca aca 2513  
Met Ala Ala Ala Thr Thr Thr Thr Thr Thr  
1 5 10  
tct tct tcg atc tcc ttc tcc acc aaa cca tct cct tcc tcc tcc aaa 2561  
Ser Ser Ser Ile Ser Phe Ser Thr Lys Pro Ser Pro Ser Ser Ser Lys



ttg gtt gat gtt cct aaa gat att caa caa cag ctt gcg att cct aat	3281
Leu Val Asp Val Pro Lys Asp Ile Gln Gln Gln Leu Ala Ile Pro Asn	
255 260 265	
tgg gaa cag gct atg aga tta cct ggt tat atg tct agg atg cct aaa	3329
Trp Glu Gln Ala Met Arg Leu Pro Gly Tyr Met Ser Arg Met Pro Lys	
270 275 280	
cct ccg gaa gat tct cat ttg gag cag att gtt agg ttg att tct gag	3377
Pro Pro Glu Asp Ser His Leu Glu Gln Ile Val Arg Leu Ile Ser Glu	
285 290 295	
tct aag aag cct gtg ttg tat gtt ggt ggt ggt tgt ttg aat tct agc	3425
Ser Lys Lys Pro Val Leu Tyr Val Gly Gly Gly Cys Leu Asn Ser Ser	
300 305 310	
gat gaa ttg ggt agg ttt gtt gag ctt acg ggg atc cct gtt gcg agt	3473
Asp Glu Leu Gly Arg Phe Val Glu Leu Thr Gly Ile Pro Val Ala Ser	
315 320 325 330	
acg ttg atg ggg ctg gga tct tat cct tgt gat gat gag ttg tcg tta	3521
Thr Leu Met Gly Leu Gly Ser Tyr Pro Cys Asp Asp Glu Leu Ser Leu	
335 340 345	
cat atg ctt gga atg cat ggg act gtg tat gca aat tac gct gtg gag	3569
His Met Leu Gly Met His Gly Thr Val Tyr Ala Asn Tyr Ala Val Glu	
350 355 360	
cat agt gat ttg ttg ttg gcg ttt ggg gta agg ttt gat gat cgt gtc	3617
His Ser Asp Leu Leu Leu Ala Phe Gly Val Arg Phe Asp Asp Arg Val	
365 370 375	
acg ggt aag ctt gag gct ttt gct agt agg gct aag att gtt cat att	3665
Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala Lys Ile Val His Ile	
380 385 390	
gat att gac tcg gct gag att ggg aag aat aag act cct cat gtg tct	3713
Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys Thr Pro His Val Ser	
395 400 405 410	
gtg tgt ggt gat gtt aag ctg gct ttg caa ggg atg aat aag gtt ctt	3761
Val Cys Gly Asp Val Lys Leu Ala Leu Gln Gly Met Asn Lys Val Leu	
415 420 425	
gag aac cga gcg gag gag ctt aag ctt gat ttt gga gtt tgg agg aat	3809
Glu Asn Arg Ala Glu Glu Leu Lys Leu Asp Phe Gly Val Trp Arg Asn	
430 435 440	
gag ttg aac gta cag aaa cag aag ttt ccg ttg agc ttt aag acg ttt	3857
Glu Leu Asn Val Gln Lys Gln Lys Phe Pro Leu Ser Phe Lys Thr Phe	
445 450 455	
ggg gaa gct att cct cca cag tat gcg att aag gtc ctt gat gag ttg	3905
Gly Glu Ala Ile Pro Pro Gln Tyr Ala Ile Lys Val Leu Asp Glu Leu	
460 465 470	

act gat gga aaa gcc ata ata agt act ggt gtc ggg caa cat caa atg 3953  
 Thr Asp Gly Lys Ala Ile Ile Ser Thr Gly Val Gly Gln His Gln Met  
 475 480 485 490

tgg gcg gcg cag ttc tac aat tac aag aaa cca agg cag tgg cta tca 4001  
 Trp Ala Ala Gln Phe Tyr Asn Tyr Lys Lys Pro Arg Gln Trp Leu Ser  
 495 500 505

tca gga ggc ctt gga gct atg gga ttt gga ctt cct gct gcg att gga 4049  
 Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu Pro Ala Ala Ile Gly  
 510 515 520

gcg tct gtt gct aac cct gat gcg ata gtt gtg gat att gac gga gat 4097  
 Ala Ser Val Ala Asn Pro Asp Ala Ile Val Val Asp Ile Asp Gly Asp  
 525 530 535

gga agc ttt ata atg aat gtg caa gag cta gcc act att cgt gta gag 4145  
 Gly Ser Phe Ile Met Asn Val Gln Glu Leu Ala Thr Ile Arg Val Glu  
 540 545 550

aat ctt cca gtg aag gta ctt tta tta aac aac cag cat ctt ggc atg 4193  
 Asn Leu Pro Val Lys Val Leu Leu Leu Asn Asn Gln His Leu Gly Met  
 555 560 565 570

gtt atg caa tgg gaa gat cgg ttc tac aaa gct aac cga gct cac aca 4241  
 Val Met Gln Trp Glu Asp Arg Phe Tyr Lys Ala Asn Arg Ala His Thr  
 575 580 585

ttt ctc ggg gat ccg gct cag gag gac gag ata ttc ccg aac atg ttg 4289  
 Phe Leu Gly Asp Pro Ala Gln Glu Asp Glu Ile Phe Pro Asn Met Leu  
 590 595 600

ctg ttt gca gca gct tgc ggg att cca gcg gcg agg gtg aca aag aaa 4337  
 Leu Phe Ala Ala Ala Cys Gly Ile Pro Ala Ala Arg Val Thr Lys Lys  
 605 610 615

gca gat ctc cga gaa gct att cag aca atg ctg gat aca cca gga cct 4385  
 Ala Asp Leu Arg Glu Ala Ile Gln Thr Met Leu Asp Thr Pro Gly Pro  
 620 625 630

tac ctg ttg gat gtg att tgt ccg cac caa gaa cat gtg ttg ccg atg 4433  
 Tyr Leu Leu Asp Val Ile Cys Pro His Gln Glu His Val Leu Pro Met  
 635 640 645 650

atc ccg aat ggt ggc act ttc aac gat gtc ata acg gaa gga gat ggc 4481  
 Ile Pro Asn Gly Gly Thr Phe Asn Asp Val Ile Thr Glu Gly Asp Gly  
 655 660 665

cgg att aaa tac tgagagatga aaccggtgat tatcagaacc ttttatggtc 4533  
 Arg Ile Lys Tyr  
 670

tttgtatgca tatggtaaaa aaacttagtt tgcaatttcc tgtttgtttt ggtaatttga 4593

gtttctttta gttgttgatc tgctgtcttt ttggtttacg tcagactact actgctgttg 4653

ttgtttggtt tcctttcttt cattttataa ataaataatc cggttcggtt tactccttgt 4713

gactggctca gtttggttat tgcgaaatgc gaatggtaaa ttgagtaatt gaaattcggt 4773  
 attagggttc taagctgttt taacagtcac tgggtaata tctctcgaat cttgcatgga 4833  
 aaatgctctt accattgggt ttttaattgaa atgtgctcat atgggccgtg gtttccaaat 4893  
 taaataaaac tacgatgtca tcgagaagta aaatcaactg tgtccacatt atcagttttg 4953  
 tgtatacgat gaaatagggt aattcaaaat cttagcttgat atgccttttg gttcatttta 5013  
 accttctgta aacatttttt cagattttga acaagtaaat ccaaaaaaaaa aaaaaaaaaa 5073  
 tctcaactca aactaaatt attttaatgt ataaaagatg cttaaaacat ttggcttaaa 5133  
 agaaagaagc taaaaacata gagaactcct gtaaattgaa gtatgaaaat atactgaatt 5193  
 gggattata tgaatttttc tgatttagga ttcacatgat ccaaaaagga aatccagaag 5253  
 cactaatcag acattggaag taggaatatt tcaaaaagtt tttttttttt aagtaagtga 5313  
 caaaagcttt taaaaaatag aaaagaaact agtattaaag ttgtaaattt aataaacaaa 5373  
 agaaattttt tatatttttt catttctttt tccagcatga ggttatgatg gcaggatgtg 5433  
 gatttcattt ttttctttt gatagccttt taattgatct attataattg acgaaaaaat 5493  
 attagttaat tatagatata ttttaggtag tattagcaat ttacacttcc aaaagactat 5553  
 gtaagttgta aatatgatgc gttgatctct tcatcattca atggttagtc aaaaaataa 5613  
 aagcttaact agtaactaa agtagtcaaa aattgtactt tagtttaaaa tattacatga 5673  
 ataatccaaa acgacattta tgtgaaacaa aaacaatc taga 5717

<210> 2

<211> 670

<212> PRT

<213> Arabidopsis thaliana

<400> 2

Met Ala Ala Ala Thr Thr Thr Thr Thr Thr Ser Ser Ser Ile Ser Phe  
 1 5 10 15  
 Ser Thr Lys Pro Ser Pro Ser Ser Ser Lys Ser Pro Leu Pro Ile Ser  
 20 25 30  
 Arg Phe Ser Leu Pro Phe Ser Leu Asn Pro Asn Lys Ser Ser Ser Ser  
 35 40 45  
 Ser Arg Arg Arg Gly Ile Lys Ser Ser Ser Pro Ser Ser Ile Ser Ala  
 50 55 60  
 Val Leu Asn Thr Thr Thr Asn Val Thr Thr Thr Pro Ser Pro Thr Lys  
 65 70 75 80

Pro Thr Lys Pro Glu Thr Phe Ile Ser Arg Phe Ala Pro Asp Gln Pro  
 85 90 95

Arg Lys Gly Ala Asp Ile Leu Val Glu Ala Leu Glu Arg Gln Gly Val  
 100 105 110

Glu Thr Val Phe Ala Tyr Pro Gly Gly Ala Ser Met Glu Ile His Gln  
 115 120 125

Ala Leu Thr Arg Ser Ser Ser Ile Arg Asn Val Leu Pro Arg His Glu  
 130 135 140

Gln Gly Gly Val Phe Ala Ala Glu Gly Tyr Ala Arg Ser Ser Gly Lys  
 145 150 155 160

Pro Gly Ile Cys Ile Ala Thr Ser Gly Pro Gly Ala Thr Asn Leu Val  
 165 170 175

Ser Gly Leu Ala Asp Ala Leu Leu Asp Ser Val Pro Leu Val Ala Ile  
 180 185 190

Thr Gly Gln Val Pro Arg Arg Met Ile Gly Thr Asp Ala Phe Gln Glu  
 195 200 205

Thr Pro Ile Val Glu Val Thr Arg Ser Ile Thr Lys His Asn Tyr Leu  
 210 215 220

Val Met Asp Val Glu Asp Ile Pro Arg Ile Ile Glu Glu Ala Phe Phe  
 225 230 235 240

Leu Ala Thr Ser Gly Arg Pro Gly Pro Val Leu Val Asp Val Pro Lys  
 245 250 255

Asp Ile Gln Gln Gln Leu Ala Ile Pro Asn Trp Glu Gln Ala Met Arg  
 260 265 270

Leu Pro Gly Tyr Met Ser Arg Met Pro Lys Pro Pro Glu Asp Ser His  
 275 280 285

Leu Glu Gln Ile Val Arg Leu Ile Ser Glu Ser Lys Lys Pro Val Leu  
 290 295 300

Tyr Val Gly Gly Gly Cys Leu Asn Ser Ser Asp Glu Leu Gly Arg Phe  
 305 310 315 320

Val Glu Leu Thr Gly Ile Pro Val Ala Ser Thr Leu Met Gly Leu Gly  
 325 330 335

Ser Tyr Pro Cys Asp Asp Glu Leu Ser Leu His Met Leu Gly Met His  
 340 345 350

Gly Thr Val Tyr Ala Asn Tyr Ala Val Glu His Ser Asp Leu Leu Leu  
 355 360 365

Ala Phe Gly Val Arg Phe Asp Asp Arg Val Thr Gly Lys Leu Glu Ala  
 370 375 380

Phe Ala Ser Arg Ala Lys Ile Val His Ile Asp Ile Asp Ser Ala Glu  
 385 390 395 400

Ile Gly Lys Asn Lys Thr Pro His Val Ser Val Cys Gly Asp Val Lys  
 405 410 415

Leu Ala Leu Gln Gly Met Asn Lys Val Leu Glu Asn Arg Ala Glu Glu  
 420 425 430

Leu Lys Leu Asp Phe Gly Val Trp Arg Asn Glu Leu Asn Val Gln Lys  
 435 440 445

Gln Lys Phe Pro Leu Ser Phe Lys Thr Phe Gly Glu Ala Ile Pro Pro  
 450 455 460

Gln Tyr Ala Ile Lys Val Leu Asp Glu Leu Thr Asp Gly Lys Ala Ile  
 465 470 475 480

Ile Ser Thr Gly Val Gly Gln His Gln Met Trp Ala Ala Gln Phe Tyr  
 485 490 495

Asn Tyr Lys Lys Pro Arg Gln Trp Leu Ser Ser Gly Gly Leu Gly Ala  
 500 505 510

Met Gly Phe Gly Leu Pro Ala Ala Ile Gly Ala Ser Val Ala Asn Pro  
 515 520 525

Asp Ala Ile Val Val Asp Ile Asp Gly Asp Gly Ser Phe Ile Met Asn  
 530 535 540

Val Gln Glu Leu Ala Thr Ile Arg Val Glu Asn Leu Pro Val Lys Val  
 545 550 555 560

Leu Leu Leu Asn Asn Gln His Leu Gly Met Val Met Gln Trp Glu Asp  
 565 570 575

Arg Phe Tyr Lys Ala Asn Arg Ala His Thr Phe Leu Gly Asp Pro Ala  
 580 585 590

Gln Glu Asp Glu Ile Phe Pro Asn Met Leu Leu Phe Ala Ala Ala Cys  
 595 600 605

Gly Ile Pro Ala Ala Arg Val Thr Lys Lys Ala Asp Leu Arg Glu Ala  
 610 615 620

Ile Gln Thr Met Leu Asp Thr Pro Gly Pro Tyr Leu Leu Asp Val Ile  
 625 630 635 640

Cys Pro His Gln Glu His Val Leu Pro Met Ile Pro Asn Gly Gly Thr  
 645 650 655

Phe Asn Asp Val Ile Thr Glu Gly Asp Gly Arg Ile Lys Tyr  
 660 665 670

INTERNATIONAL SEARCH REPORT

PCT/EP 03/07085

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/60 C12N15/82 C12N5/10 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	STOROZHENKO S V ET AL: "Isolation of the transgenic potato plants expressing the mutant gene coding for acetolactate synthase from Arabidopsis thaliana." BIOPOLIMERY I KLETKA, vol. 10, no. 1, 1994, pages 67-71, XP002220128 ISSN: 0233-7657 abstract	1-18
Y	WO 01 82685 A (PENG JIANYING ;HIRAYAMA LYNNE (US); AMERICAN CYANAMID CO (US); LOC) 8 November 2001 (2001-11-08) the whole document	1-18
	-/--	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

3 November 2003

Date of mailing of the international search report

12/11/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2260 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

## INTERNATIONAL SEARCH REPORT

PCT/EP 03/07085

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 767 366 A (MURAI NORIMOTO ET AL) 16 June 1998 (1998-06-16) cited in the application the whole document ---	
A	WO 96 33270 A (AMERICAN CYANAMID CO ;KAKEFU DA GENICHI (US); OTT KARL HEINZ (US);) 24 October 1996 (1996-10-24) figure 15 ---	
A	LIBIAKOVA G ET AL: "Efficacy of an intron-containing kanamycin resistance gene as a selectable marker in plant transformation." PLANT CELL REPORTS, vol. 20, no. 7, October 2001 (2001-10), pages 610-615, XP002220129 ISSN: 0721-7714 cited in the application -----	
A	WO 92 11376 A (AMYLOGENE HB) 9 July 1992 (1992-07-09) the whole document ---	10,11
A	WO 92 11375 A (AMYLOGENE HB) 9 July 1992 (1992-07-09) the whole document -----	10,12

## INTERNATIONAL SEARCH REPORT

PCT/EP 03/07085

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0182685	A	08-11-2001	AU 5920601	A 12-11-2001
			BR 0110410	A 01-07-2003
			CA 2407396	A1 08-11-2001
			EP 1284597	A1 26-02-2003
			WO 0182685	A1 08-11-2001
			US 2003167538	A1 04-09-2003
			US 5767366	A
WO 9633270	A	24-10-1996	US 5853973	A 29-12-1998
			US 5928937	A 27-07-1999
			AU 5575896	A 07-11-1996
			BR 9604993	A 30-11-1999
			CA 2218526	A1 24-10-1996
			CZ 9703317	A3 17-06-1998
			EP 0821729	A1 04-02-1998
			HU 9900852	A2 28-07-1999
			JP 11504213	T 20-04-1999
			NO 974803	A 19-12-1997
			NZ 307012	A 28-01-2000
			NZ 500281	A 28-02-2000
			NZ 502491	A 28-04-2000
			PL 322899	A1 02-03-1998
			WO 9633270	A1 24-10-1996
			US 6576455	B1 10-06-2003
			US 2003180929	A1 25-09-2003
WO 9211376	A	09-07-1992	SE 467358	B 06-07-1992
			AT 243752	T 15-07-2003
			AU 9114891	A 22-07-1992
			CA 2098171	A1 21-06-1992
			DE 69133285	D1 31-07-2003
			DK 563189	T3 13-10-2003
			EP 0563189	A1 06-10-1993
			EP 0788735	A1 13-08-1997
			EP 0921191	A1 09-06-1999
			FI 932804	A 17-06-1993
			HU 66754	A2 28-12-1994
			JP 6507064	T 11-08-1994
			JP 2003034702	A 07-02-2003
			KR 210352	B1 15-07-1999
			NO 932227	A 11-08-1993
			PL 169848	B1 30-09-1996
			WO 9211376	A1 09-07-1992
			US 5824798	A 20-10-1998
			WO 9211375	A
AU 9109791	A 22-07-1992			
EP 0563201	A1 06-10-1993			
PL 169859	B1 30-09-1996			
SE 9004095	A 01-06-1992			
WO 9211375	A1 09-07-1992			
US 5856467	A 05-01-1999			