



(86) Date de dépôt PCT/PCT Filing Date: 2009/10/16
 (87) Date publication PCT/PCT Publication Date: 2010/04/22
 (45) Date de délivrance/Issue Date: 2016/02/23
 (85) Entrée phase nationale/National Entry: 2011/04/15
 (86) N° demande PCT/PCT Application No.: JP 2009/067935
 (87) N° publication PCT/PCT Publication No.: 2010/044469
 (30) Priorité/Priority: 2008/10/17 (JP2008-268106)

(51) Cl.Int./Int.Cl. *C12P 23/00* (2006.01),
C12N 1/20 (2006.01), *C12P 5/02* (2006.01),
C12Q 1/68 (2006.01)
 (72) Inventeurs/Inventors:
 HIRASAWA KAZUAKI, JP;
 TSUBOKURA AKIRA, JP;
 SATOU HIROSHI, JP;
 YATA TETSUHISA, JP
 (73) Propriétaire/Owner:
 JX NIPPON OIL & ENERGY CORPORATION, JP
 (74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PROCEDE DE FERMENTATION DE CAROTENOIDE
 (54) Title: CAROTENOID FERMENTATION METHOD

(57) **Abrégé/Abstract:**

The present invention provides a method for producing a carotenoid, which comprises culturing a carotenoid-producing bacterium in an amino acid-supplemented medium, and collecting the carotenoid from the resulting cultured product, wherein the amino acid is at least one selected from the group consisting of glutamic acid, aspartic acid, glutamine, asparagine, alanine, glycine, serine, threonine, arginine, tyrosine, proline, phenylalanine and leucine, and salts thereof.

ABSTRACT

The present invention provides a method for producing a carotenoid, which comprises culturing a carotenoid-producing bacterium in an amino acid-supplemented medium, and collecting the carotenoid from the resulting cultured product, wherein the amino acid is at least one selected from the group consisting of glutamic acid, aspartic acid, glutamine, asparagine, alanine, glycine, serine, threonine, arginine, tyrosine, proline, phenylalanine and leucine, and salts thereof.

SPECIFICATION

CAROTENOID FERMENTATION METHOD

TECHNICAL FIELD

[0001]

The present invention relates to a method for microbiological production of carotenoids. More specifically, the present invention relates to a method for microbial fermentation-based production of carotenoids such as astaxanthin, canthaxanthin, zeaxanthin, β -cryptoxanthin, lycopene, β -carotene, phoenicoxanthin, adonixanthin, echinenone, asteroidenone and 3-hydroxyechinenone.

BACKGROUND ART

[0002]

Carotenoids are natural pigments useful as feed additives, food additives, pharmaceutical ingredients and so on. Carotenoids include astaxanthin, canthaxanthin, zeaxanthin, β -cryptoxanthin, lycopene, β -carotene, phoenicoxanthin, adonixanthin, echinenone, asteroidenone and 3-hydroxyechinenone. Among them, astaxanthin is useful as a body color improver for cultured fish including salmon, trout and red sea bream and/or useful as a feed additive such as a color improver for poultry egg yolk. Astaxanthin is also industrially valuable as a safe and natural food additive or health food material. As in the case of astaxanthin, adonixanthin and phoenicoxanthin are also expected for use as feed additives, food additives, pharmaceutical ingredients or the like once their industrial production has been established. In addition, β -carotene is used as a feed additive, a food additive, a pharmaceutical ingredient or the like, canthaxanthin is used as a feed additive, a food additive, a cosmetic ingredient or the like, and zeaxanthin is used as a food additive, a feed additive or the like. Further, lycopene, echinenone, β -cryptoxanthin, 3-hydroxyechinenone, asteroidenone and others are also expected for use as feed additives, food materials or the like. For production of these carotenoids, chemical synthesis, extraction from natural products, microbial production or other techniques are known.

[0003]

For chemical synthesis of astaxanthin, conversion from β -carotene (Non-patent Document 1: Pure Appl. Chem., 57, 741, 1985) and synthesis from C15 phosphonium salt (Non-patent Document 2: Helv. Chim. Acta, 64, 2436, 1981) are known. Astaxanthin produced by these chemical synthesis techniques is commercially available as a feed additive. Astaxanthin can also be extracted from fish (e.g., red sea bream, salmon) and crustaceans (e.g., shrimp, crab, krill) because astaxanthin is found in these

organisms.

For microbial production of astaxanthin, there are reports of culture in green algae *Haematococcus pluvialis* (Patent Document 1: JP 2007-97584 A), fermentation in red yeast *Phaffia rhodozyma* (Patent Document 2: JP H11-69969 A), and fermentation in bacteria belonging to the genus *Paracoccus* (hereinafter also referred to as “*Paracoccus* sp.”). Examples of astaxanthin-producing bacteria belonging to the genus *Paracoccus* include strains E-396 and A-581-1 (Patent Document 3: JP H7-79796 A and Non-patent Document 3: *International Journal of Systematic Bacteriology* (1999), 49, 277-282). Other astaxanthin-producing bacteria belonging to the genus *Paracoccus* include *Paracoccus marcusii* strain MH1 (Patent Document 4: JP 2001-512030 A), *Paracoccus haeundaensis* strain BC74171 (Non-patent Document 4: *International Journal of Systematic and Evolutionary Microbiology* (2004), 54, 1699-1702), *Paracoccus* sp. strain N-81106 (Patent Document 5: JP 2007-244205 A), *Paracoccus zeaxanthinifaciens* (Non-patent Document 5: *International Journal of Systematic and Evolutionary Microbiology* (2003), 53, 231-238) and *Paracoccus* sp. strain PC-1 (Patent Document 6: WO 2005/118812), etc.

[0004]

However, the above techniques for carotenoid production have some problems. For example, chemical synthesis would make an unfavorable impression on consumers in terms of safety. Likewise, extraction from natural products requires high production costs. Moreover, production in green algae or yeast not only provides low productivity, but also makes difficult carotenoid extraction because of their rigid cell walls.

[0005]

In contrast, bacteria belonging to the genus *Paracoccus* are advantageous, e.g., in having high growth rate and in achieving high productivity and easy extraction of carotenoids, and some reports have been issued for methods of their culture. JP 2007-143492 A (Patent Document 7) discloses a method in which an iron salt is added during culture, while JP 2008-167665 A (Patent Document 8) discloses a method in which the carbon source concentration is limited. However, these methods are not practical for commercial or industrial purposes because an expensive yeast extract is used in a large amount as a material of the medium.

Related Art Documents

Patent Documents

[0006]

[Patent Document 1] JP 2007-97584 A

[Patent Document 2] JP H11-69969 A

[Patent Document 3] JP H7-79796 A

[Patent Document 4] JP 2001-512030 A

[Patent Document 5] JP 2007-244205 A

[Patent Document 6] WO2005/118812

[Patent Document 7] JP 2007-143492 A

[Patent Document 8] JP 2008-167665 A

Non-patent Documents

[0007]

[Non-patent Document 1] Pure Appl. Chem., 57, 741, 1985

[Non-patent Document 2] Helv. Chim. Acta, 64, 2436, 1981

[Non-patent Document 3] International Journal of Systematic Bacteriology (1999), 49, 277-282

[Non-patent Document 4] International Journal of Systematic and Evolutionary Microbiology (2004), 54, 1699-1702

[Non-patent Document 5] International Journal of Systematic and Evolutionary Microbiology (2003), 53, 231-238

GENERAL DESCRIPTIONS OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0008]

The present invention has been made under these circumstances and aims to provide a method for microbiological production of carotenoids in high yields and at low costs.

MEANS FOR SOLVING THE PROBLEMS

[0009]

As a result of various efforts made to solve the problems stated above, the inventors of the present invention have found that it is possible to improve the productivity of carotenoids when a medium commonly used for bacterial culture is further supplemented with an amino acid or a salt thereof (e.g., sodium glutamate) in the culture of carotenoid-producing bacteria. This finding led to the completion of the present invention.

[0010]

Namely, the present invention relates to a method for producing a carotenoid, which comprises culturing a carotenoid-producing bacterium in an amino acid-supplemented medium, and collecting the carotenoid from the resulting cultured product,

wherein the amino acid is at least one selected from the group consisting of glutamic acid, aspartic acid, glutamine, asparagine, alanine, glycine, serine, threonine, arginine, tyrosine, proline, phenylalanine and leucine, and salts thereof.

In the above method, a preferred amino acid is glutamic acid or a glutamate salt.

30179-209

The supplemented amino acid concentration may be, for example, 1 mmol/L to 200 mmol/L. As used herein, the term “supplemented amino acid concentration” is intended to mean the amino acid concentration achieved in the medium in which the amino acid is supplemented (i.e., the concentration of supplemented amino acid(s) in the medium).

5 The carotenoid may be, for example, at least one selected from the group consisting of astaxanthin, canthaxanthin, zeaxanthin, β -cryptoxanthin, lycopene, carotene, phoenicoxanthin, adonixanthin, echinenone, asteroidenone and 3-hydroxyechinenone.

In the above method, bacteria preferred for use are those belonging to the genus *Paracoccus*. Such bacteria may also be those having a homology of 95% or more in the
10 base sequence of DNA corresponding to 16S ribosomal RNA with respect to the base sequence represented by SEQ ID NO: 1. In particular, strain E-396 (FERM BP-4283) or A-581-1 (FERM BP-4671) or a mutant thereof is preferred for use.

[0010A]

The present invention as claimed relates to:

15 - a method for producing astaxanthin, which comprises culturing an astaxanthin-producing bacterium of the genus *Paracoccus* in a culture medium supplemented with glutamic acid or a glutamate salt, and collecting the astaxanthin from the resulting culture; wherein the bacterium is a mutant that produces astaxanthin at a higher concentration in the culture medium than its parent strain.

30179-209

EFFECT OF THE INVENTION

[0011]

The present invention aims to provide more efficient production of carotenoids at high concentrations. The present invention also aims to provide microbiological production
5 of carotenoids at low costs.

MODES FOR CARRYING OUT THE INVENTION

[0012]

The present invention will be further described in more detail below. The scope of the present invention is not limited to the following description, and may be carried
10 out in appropriate modifications other than the following illustrative embodiments without departing from the invention as defined in the claims.

The specification of Japanese Patent Application No. 2008-268106, is the document upon which the present application claims the benefit of priority.

[0013]

15 The present invention relates to a method for producing a carotenoid by culturing a carotenoid-producing bacterium. This method is characterized by that the medium is supplemented with a specific amino acid(s). The method of the present invention allows more efficient and low-cost production of carotenoids at high concentrations.

[0014]

Although bacteria used in the present invention are not limited in any way as long as they are carotenoid-producing bacteria, preferred for use are bacteria belonging to the genus *Paracoccus*. Among such bacteria belonging to the genus *Paracoccus*, preferred for use are *Paracoccus carotinifaciens*, *Paracoccus marcusii*, *Paracoccus haeundaensis* and *Paracoccus zeaxanthinifaciens*, especially *Paracoccus carotinifaciens*. Specific exemplary strains of bacteria belonging to the genus *Paracoccus* include *Paracoccus carotinifaciens* strain E-396 and *Paracoccus* sp. strain A-581-1 (FERM BP-4671), which are also preferred for use in the present invention.

[0015]

Other carotenoid-producing bacteria preferred for use are those having high homology in the base sequence of DNA corresponding to 16S ribosomal RNA with respect to the base sequence of the strain E-396 represented by SEQ ID NO: 1. As used herein, the phrase "having high homology" is, for example, intended to mean that there is a homology of preferably 95% or more, more preferably 96% or more, even more preferably 97% or more, particularly preferably 98% or more, most preferably 99% or more between the base sequence represented by SEQ ID NO: 1 and the corresponding base sequences of bacteria to be compared.

The base sequence of DNA corresponding to 16S ribosomal RNA refers to a base sequence having T (thymine) instead of U (uracil) in the base sequence of 16S ribosomal RNA.

Microbial classification based on homology of the base sequence of this 16S ribosomal RNA has become mainstream in recent years. Conventional microbial classifications are based on microbiological properties of microorganisms, such as motility, auxotrophy, sugar assimilation property, etc., and may cause misclassification of microorganisms in some cases when a spontaneous mutation has induced a phenotypic change or the like. In contrast, the base sequence of 16S ribosomal RNA is very stable inheritantly; and hence classification based on the homology of this sequence ensures remarkably improved confidence in classification results when compared to conventional classifications.

[0016]

The base sequence of 16S ribosomal RNA in *Paracoccus carotinifaciens* strain E-396 has a homology of 99.7%, 99.7%, 99.6%, 99.4%, 95.7% and 95.4% with the base sequences of 16S ribosomal RNA in other carotenoid-producing bacteria, i.e., *Paracoccus marcusii* strain DSM 11574, *Paracoccus* sp. strain N-81106, *Paracoccus haeundaensis* strain BC 74171, *Paracoccus* sp. strain A-581-1, *Paracoccus zeaxanthinifaciens* strain ATCC 21588 and *Paracoccus* sp. strain PC-1, respectively, which indicates that they are taxonomically very closely related strains. Thus, these strains appear to constitute a group of carotenoid-producing bacteria. For this reason, these strains are preferred for use in the present invention and allow efficient production

of carotenoids.

[0017]

In the present invention, it is also possible to use mutant strains with improved productivity of carotenoids. Examples of improved mutant strains include those having high ability to produce astaxanthin (JP 2001-95500 A), those selectively producing canthaxanthin at high levels (JP 2003-304875 A), those selectively producing zeaxanthin and β -cryptoxanthin at high levels (JP 2005-87097 A), and those selectively producing lycopene (JP 2005-87100 A).

[0018]

Such mutant strains with improved productivity of carotenoids can be obtained by mutagenesis and screening. Any technique may be used for mutagenesis as long as it induces a mutation(s). For example, it is possible to use chemical techniques using a mutagen such as N-methyl-N'-nitro-N-nitrosoguanidine (NTG) or ethylmethane sulfonate (EMS), physical techniques such as ultraviolet irradiation or X-ray irradiation, or biological techniques using gene recombination or transposons, etc. Although bacteria to be mutated are not limited in any way, they are preferably carotenoid-producing bacteria. Alternatively, such mutant strains may be generated as a result of spontaneous mutation.

Any technique may be used for screening of mutant strains, including selection of a desired mutant strain on the basis of colony color on agar medium, as well as selection of a desired mutant strain by carotenoid pigment analysis using absorbance, high performance liquid chromatography, thin-layer chromatography or the like from among mutant strains cultured in test tubes, flasks, fermentation tanks, etc.

Such mutagenesis and screening steps may be performed once, or alternatively, may be repeated twice or more, for example, such that a mutant strain obtained by mutagenesis and screening is subjected to further mutagenesis and screening to obtain a mutant strain with improved productivity.

[0019]

The strain E-396 listed as an example of carotenoid-producing bacteria used in the present invention has been internationally deposited to the International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology, as shown below.

International Deposition Authority: International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology (former National Institute of Bioscience and Human-Technology, Agency of Science and Technology, Ministry of International Trade and Industry), Chuoh 6, Higashi 1-1-1, Tsukuba-shi, Ibaraki-ken, 305-8566

Identification No.:	E-396
Deposition No.:	FERM BP-4283

Date of original deposition: April 27, 1993

[0020]

Likewise, the strain A-581-1 listed as another example of carotenoid-producing bacteria used in the present invention has been internationally deposited to the above depositary, as shown below.

Identification No.: A-581-1
Deposition No.: FERM BP-4671
Date of original deposition: May 20, 1994

[0021]

In the present invention, when cultured in a medium supplemented with a specific amino acid(s), the above carotenoid-producing bacteria can produce larger amounts of carotenoids at higher concentrations than in a medium not supplemented with the amino acid(s).

Carotenoids produced by the method of the present invention are not limited in any way. Examples include astaxanthin, canthaxanthin, zeaxanthin, β -cryptoxanthin, lycopene, β -carotene, phoenicoxanthin, adonixanthin, echinenone, asteroidenone or 3-hydroxyechinenone. Preferred is astaxanthin, canthaxanthin, zeaxanthin or β -cryptoxanthin, and more preferred is astaxanthin, zeaxanthin or β -cryptoxanthin. These carotenoids may be produced, either alone or in combination, in the present invention.

[0022]

The method for culturing the above bacteria in the present invention will be described below.

For culture in the present invention, any medium for carotenoid production may be used, without limitation, as long as it is an amino acid-supplemented medium containing a specific amino acid(s) and allows carotenoid-producing bacteria to grow and produce carotenoids. Preferred for use is a medium containing a carbon source, a nitrogen source, inorganic salts and optionally vitamins, etc. Namely, in the present invention, an amino acid(s) may be added to any medium (e.g., standard medium for carotenoid production) in which carotenoid-producing bacteria can grow and produce carotenoids.

[0023]

Examples of a carbon source include sugars such as glucose, sucrose, lactose, fructose, trehalose, mannose, mannitol and maltose; organic acids such as acetic acid, fumaric acid, citric acid, propionic acid, malic acid, malonic acid and pyruvic acid; alcohols such as ethanol, propanol, butanol, pentanol, hexanol, isobutanol and glycerol; as well as fats and oils such as soybean oil, rice bran oil, olive oil, corn oil, sesame oil

and linseed oil. Among them, glucose or sucrose is preferred for use. One or more of these carbon sources may be used. The amount to be added to the medium before culture (initial medium) will vary depending on the type of carbon source and may be adjusted as appropriate. It is usually 1 to 100 g, preferably 2 to 50 g per liter of the medium. Moreover, such carbon sources are not only added to the initial medium, but also may preferably be supplied sequentially or continuously during culture.

[0024]

Examples of an inorganic nitrogen source include ammonium salts (e.g., ammonium nitrate, ammonium sulfate, ammonium chloride, ammonium phosphate), nitrate salts (e.g., potassium nitrate), ammonia and urea, which may be used either alone or in combination. The amount to be added will vary depending on the type of nitrogen source and may be adjusted as appropriate. It is usually 0.1 g to 20 g, preferably 0.2 to 10 g per liter of the medium.

Examples of an organic nitrogen source include corn steep liquor (including filtered corn steep liquor), Pharmamedia, soybean meal, soybean powder, peanut meal, distiller's solubles and dry yeast, which may be used either alone or in combination. The concentration to be added will vary depending on the type of nitrogen source and may be adjusted as appropriate. It is usually 0 to 80 g/L, preferably 0 to 30 g/L.

Such inorganic and organic nitrogen sources are normally added to the initial medium, and may also preferably be supplied sequentially or continuously during culture.

[0025]

Examples of inorganic salts include phosphate salts (e.g., potassium dihydrogen phosphate, dipotassium hydrogen phosphate, disodium hydrogen phosphate), magnesium salts (e.g., magnesium sulfate, magnesium chloride), iron salts (e.g., iron sulfate, iron chloride), calcium salts (e.g., calcium chloride, calcium carbonate), sodium salts (e.g., sodium carbonate, sodium chloride), manganese salts (e.g., manganese sulfate), cobalt salts (e.g., cobalt chloride), copper salts (e.g., copper sulfate), zinc salts (e.g., zinc sulfate), molybdenum salts (e.g., sodium molybdate), nickel salts (e.g., nickel sulfate), selenium salts (e.g., sodium selenate), boric acid and potassium iodide, which may be used either alone or in combination. The amount to be added will vary depending on the type of inorganic salt and may be adjusted as appropriate. It is usually 0.0001 to 15 g per liter of the medium. A preferred concentration is 0.02 to 15 g/L for phosphate salts, magnesium salts, calcium salts, sodium salts and iron salts, and 0.1 to 15 mg/L for manganese salts, cobalt salts, copper salts, zinc salts, molybdenum salts, nickel salts, selenium salts, boric acid, potassium iodide and so on. Such inorganic salts are normally added to the initial medium, and may also be supplied sequentially or continuously during culture.

[0026]

Examples of vitamins available for use include cyanocobalamin, riboflavin,

pantothenic acid, pyridoxine, thiamine, ascorbic acid, folic acid, niacin, p-aminobenzoic acid, biotin, inositol, choline and the like. The ratio to be added will vary depending on the type of vitamin and may be adjusted as appropriate. It is usually 0.001 to 1000 mg, preferably 0.01 to 100 mg per liter of the medium. Such vitamins are normally added to the initial medium, and may also be supplied sequentially or continuously during culture.

[0027]

The present invention is characterized by that carotenoid-producing bacteria are cultured in an amino acid-supplemented medium for carotenoid production supplemented with an amino acid(s). When cultured in such an amino acid-supplemented medium for carotenoid production, carotenoid-producing bacteria can produce larger amounts of carotenoids at higher concentrations than in a medium not supplemented with the amino acid(s).

[0028]

Amino acids used in the present invention are pure products (single products) purified to some extent, i.e., isolated products, but not amino acids contained in naturally occurring mixtures of complex composition (e.g., casamino acid, yeast extract, peptone). Naturally occurring mixtures may contain not only effective amino acids, but also unwanted or inhibitory components, and may further have lot-to-lot variation in their composition. Moreover, naturally occurring mixtures such as casamino acid, yeast extract and peptone are expensive and thus less valuable for industrial use.

Purified amino acids preferably have a purity of 90% or more, more preferably 95% or more, even more preferably 98% or more, particularly preferably 99% or more. It should be noted that amino acids used in the present invention may further comprise other non-amino acid components to the extent not to inhibit the growth of carotenoid-producing bacteria or to the extent not to inhibit carotenoid production by carotenoid-producing bacteria. Although amino acids used for this purpose are preferably, for example, pure amino acids free from other components including impurities, unpurified amino acids (e.g., amino acids with a purity of less than 90%) may also be used as long as they do not inhibit carotenoid production.

[0029]

As amino acids added to the medium for carotenoid production, glutamic acid, aspartic acid, glutamine, asparagine, alanine, glycine, serine, threonine, arginine, tyrosine, proline, phenylalanine and leucine or salts thereof are preferably used. These amino acids are preferably in the L-form, but may also be a mixture of L- and D-forms. More preferred are glutamic acid, aspartic acid, glutamine and asparagine or salts thereof, and even more preferred are glutamic acid and aspartic acid or salts thereof. Among them, glutamic acid or a salt thereof is preferred because of its high effect on carotenoid production. Sodium L-glutamate or a hydrate thereof is inexpensive and thus particularly preferred for use.

Examples of salts with acids include inorganic acid salts such as hydrochloride salt, hydrobromide salt, sulfate salt and phosphate salt, as well as organic acid salts such as salts with formic acid, acetic acid or lactic acid. Likewise, examples of salts with bases include alkali metal salts (e.g., sodium salt, potassium salt), alkaline earth metal salts (e.g., calcium salt, magnesium salt), organic base salts (e.g., salts with trimethylamine, triethylamine or pyridine), and ammonium salt.

Amino acids added to the medium for carotenoid production are at least one or more of the above amino acids. Although a single amino acid may be used, it is also possible to add two or more amino acids.

Among essential amino acids, cysteine, lysine, isoleucine and methionine have an inhibitory effect on carotenoid production. In the present invention, it is therefore preferred that amino acids except for these amino acids are added to the medium.

[0030]

Although amino acids are normally added to the initial medium, they may be added intermittently or continuously during culture. Alternatively, amino acids may be added to the initial medium and further added intermittently or continuously during culture.

In the method of the present invention, there is no lower limit for the supplemented amino acid concentration (i.e., the concentration of supplemented amino acids in the medium), but it is preferably 1 mmol/L or more, more preferably 3 mmol/L or more, even more preferably 5 mmol/L or more, particularly preferably 10 mmol/L, and most preferably 15 mmol/L or more. Likewise, there is no upper limit for the supplemented amino acid concentration, but it is preferably 200 mmol/L or less, more preferably 150 mmol/L or less, even more preferably 100 mmol/L or less, even still more preferably 80 mmol/L or less, particularly preferably 60 mmol/L or less, and most preferably 50 mmol/L or less. Thus, in the present invention, the supplemented amino acid concentration is 1 mmol/L to 200 mmol/L, by way of example.

[0031]

In the present invention, a defoaming agent may preferably be used to suppress bubbling of the culture solution. Such a defoaming agent may be of any type, without limitation, as long as it has the effect of suppressing bubble formation or breaking the bubbles formed and has low inhibitory effect on the producing bacteria. Examples include alcohol-based defoaming agents, polyether-based defoaming agents, ester-based defoaming agents, fatty acid-based defoaming agents, silicon-based defoaming agents, sulfonate-based defoaming agents and so on. The amount to be added will vary depending on the type of defoaming agent and may be adjusted as appropriate. It is usually 0.01 g to 10 g per liter of the medium.

Such a defoaming agent is normally added to the initial medium before sterilization, and may further be added continuously or intermittently during culture. For addition during culture, such a defoaming agent may be automatically added upon

detection of bubbles with a sensor, or may be added at given time intervals with a programmed timer, or may be added in response to the growth rate in admixture with, e.g., a carbon source, a nitrogen source or a pH adjustor to be fed, by way of example. The defoaming agent added to the initial medium and the defoaming agent added to the culture solution during culture may be either of the same or different type, depending on the intended effect.

[0032]

In the present invention, the amino acid-supplemented medium which is supplemented with an amino acid(s) is adjusted to have an initial pH of 2 to 12, preferably 6 to 9, more preferably 6.5 to 8.0. It is desirable to also maintain the pH within the above range during culture. Examples of a pH adjustor include aqueous sodium hydroxide, aqueous potassium hydroxide, aqueous sodium carbonate, aqueous ammonia, ammonia gas, aqueous sulfuric acid or mixtures thereof.

[0033]

In the present invention, the amino acid-supplemented medium is sterilized before use and used for bacterial culture. Sterilization may be accomplished as appropriate by those skilled in the art. For example, the medium in an appropriate vessel may be sterilized by heating in an autoclave. Alternatively, the medium may be sterilized by filtration through a sterile filter.

[0034]

In the present invention, carotenoid-producing bacteria are inoculated into the amino acid-supplemented medium thus prepared and cultured under given conditions. Inoculation may be accomplished as follows: strains are grown as appropriate by seed culture in such as test tubes, flasks or fermentation tanks and the resulting cultured products are each added to the amino acid-supplemented medium for carotenoid production. Any medium, without limitation, either with or without specific amino acids, may be used for seed culture as long as it ensures good growth of carotenoid-producing bacteria.

Culture is performed in an appropriate culture vessel. Such a culture vessel may be selected as appropriate depending on the culture volume and is exemplified by test tubes, flasks, fermentation tanks and so on.

The culture temperature is set to 15°C to 80°C, preferably 20°C to 35°C, more preferably 25°C to 32°C, and culture is performed under aerobic conditions for usually 1 to 20 days, preferably 2 to 12 days, more preferably 3 to 9 days. Culture under aerobic conditions includes, for example, shaking culture or aeration agitation culture, during which the dissolved oxygen concentration is preferably controlled within a certain range. Control of the dissolved oxygen concentration may be accomplished, for example, by varying the number of agitation rotations, the aeration volume, the internal pressure, etc. The dissolved oxygen concentration is controlled to preferably 0.3 to 10 ppm, more preferably 0.5 to 7 ppm, and even more preferably 1 to 5 ppm.

[0035]

In the present invention, carotenoids in the cultured product obtained by culturing carotenoid-producing bacterial cells or carotenoids collected from the cultured product after some purification procedure can be quantified by high performance liquid chromatography.

[0036]

Carotenoid-producing bacterial cells can be cultured as described above and carotenoids can be collected from the resulting cultured product.

Such a cultured product may be, for example, a culture solution, a culture supernatant, a concentrated microbial cell solution, wet microbial cells, dry microbial cells, a microbial cell lysate, etc. A culture supernatant may be prepared from a culture solution by centrifugation or filtration to remove microbial cells from the culture solution. A concentrated microbial cell solution may be obtained from a culture solution upon concentration by centrifugation or membrane filtration. Wet microbial cells may be obtained from a culture solution by centrifugation or filtration. Dry microbial cells may be obtained from wet microbial cells or a concentrated microbial cell solution upon drying in a standard manner. The carotenoid-containing dry microbial cells thus obtained may be used directly as feed additives.

In the present invention, any technique may be used, without limitation, to collect carotenoids from the above cultured product as long as it allows stable and efficient collection of carotenoids. Such a technique may be selected as appropriate from extraction and purification techniques known to those skilled in the art.

[0037]

Prior to carotenoid extraction from the cultured product, the cultured product may be subjected to one or more treatments selected from chemical treatments with alkaline reagents, surfactants or the like, biochemical treatments with lytic enzymes, lipolytic enzymes, proteolytic enzymes or the like, or physical treatments such as ultrasonication or homogenization.

[0038]

For example, when carotenoids are extracted from the cultured product, any solvent may be used, without limitation, for extraction and washing, including lower alcohols (e.g., methanol, ethanol, isopropanol), acetone, tetrahydrofuran, methylethyl ketone, methylisobutyl ketone, dichloromethane, chloroform, dimethylformamide, dimethyl sulfoxide and so on.

To minimize oxidation of carotenoids during the extraction step, the cultured product may be treated in an inert gas atmosphere such as nitrogen gas. Moreover, an antioxidant used in pharmaceutical preparations or food products may also be selected and added to the extraction solvent. Alternatively, these treatments may be used in combination.

In addition, to minimize light-induced degradation of carotenoids, the cultured

30179-209

product may be treated under light shielding conditions.

[0039]

The extract thus obtained may be used directly as a carotenoid fraction or may further be purified before use.

[0040]

Any technique may be used, without limitation, to separate bacterial cells or others from the extract after the extraction step. Examples include membrane filtration, centrifugation, decantation, etc.

[0041]

To obtain carotenoid precipitates from the extract, techniques commonly used for this purpose include concentration under heating and/or reduced pressure, as well as crystallization. Alternatively, carotenoid pigments may also be separated, without being concentrated, by precipitation at low temperature or by precipitation with acid and/or alkaline agents or with various salts.

For industrial purposes, crystallization is desired.

[0042]

The resulting carotenoid precipitates may optionally be suspended and stirred using a small volume of a solvent (e.g., lower alcohol) for washing purposes.

Washing procedures are not limited in any way, and practically preferred procedures include those in which the precipitates are collected by filtration after being suspended and stirred, or those in which a solution is passed from above the precipitates.

[0043]

The cultured product, extract or purified product obtained as described above may be used as a carotenoid fraction, either alone or in admixture at any ratio.

EXAMPLES

[0044]

The present invention will be further described in more detail by way of the following illustrative examples. The scope of the present invention is not limited to the following examples.

In the examples, carotenoids were quantified by high performance liquid chromatography (HPLC) as follows.

Two columns of Wakosil-II 5 SIL-100 ($\phi 4.6 \times 250$ mm) (Wako Pure Chemical Industries, Ltd., Japan) were connected to each other for use as a column. Elution was performed by flowing an n-hexane-tetrahydrofuran-methanol mixed solution (40:20:1), which was a mobile phase, at a flow rate of 1.0 mL/minute at a constant temperature around room temperature. The measurement was performed as follows. Samples were each dissolved in tetrahydrofuran and then diluted 100-fold with the mobile phase. Each dilution was injected in a volume of 20 μ L. The column eluate was detected at a

wavelength of 470 nm. As a reference for quantification, astaxanthin (SIGMA, Cat. No. A9335) was used. The astaxanthin concentration of the reference solution was determined using the following equation after measuring the absorbance (A) of the reference solution at 477 nm and the area percentage % (B) of the astaxanthin peak at the time of HPLC analysis under the above conditions.

$$\text{Astaxanthin concentration (mg/L)} = A/2150 \times B \times 100$$

[0045]

[Example 1]

A medium of the following composition (sucrose 30 g/L, corn steep liquor 30 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 0.3 g/L, pH 7.2) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm and sterilized in an autoclave at 121°C for 15 minutes to prepare a test tube medium for seed culture.

Next, a medium of the following composition (glucose 30 g/L, filtered corn steep liquor 5 g/L, ammonium sulfate 1.5 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 0.6 g/L, ester-based defoaming agent 0.2 g/L) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm (21 tubes in total).

The mediums in these test tubes were supplemented with 20 amino acids, i.e., glycine, alanine, valine, leucine, isoleucine, serine, threonine, aspartic acid, glutamic acid, asparagine, glutamine, lysine, arginine, cysteine, methionine, phenylalanine, tyrosine, tryptophan, histidine and proline, respectively, at 1.0 g/L. One of the tubes was supplemented with no amino acid as a control. Finally, the mediums in the tubes were adjusted to pH 7.1 with aqueous sodium hydroxide or aqueous sulfuric acid and then sterilized in an autoclave at 121°C for 20 minutes.

[0046]

Paracoccus carotinifaciens strain E-396 (FERM BP-4283) was inoculated into the test tube medium for seed culture and then cultured with shaking at 28°C for 2 days at 300 rpm. The resulting culture solution was then inoculated in 0.1 ml volumes into the 21 test tube media, respectively, and cultured with shaking at 28°C for 4 days at 300 rpm.

[0047]

Each culture solution was measured for its carotenoid concentration by HPLC and microbial cell growth was determined by OD610 (absorbance at 610 nm). As shown in Table 1, glutamic acid, aspartic acid, glutamine, asparagine, alanine, glycine, serine, threonine, arginine, tyrosine, proline, phenylalanine and leucine were found to

have a stimulatory effect on the production of carotenoid pigments. In contrast, cysteine, lysine, isoleucine and methionine were found to have a clear inhibitory effect on carotenoid production.

[0048]

[Table 1]

Supplimented component	OD610	Astaxanthin mg/L	Total carotenoid mg/L
Glycine	18	6.2	19.0
Alanine	13	6.2	20.3
Valine	10	3.0	9.2
Leucine	11	4.7	10.1
Isoleucine	6	1.3	4.5
Serine	13	6.0	16.4
Threonine	8	5.2	12.5
Aspartic Acid	18	9.4	27.0
Glutamic Acid	17	10.7	28.2
Asparagine	14	8.1	24.2
Glutamine	17	8.3	25.8
Lysine	11	1.1	4.9
Arginine	13	5.1	11.7
Cysteine	4	0.7	3.2
Methionine	7	1.4	5.0
Phenylalanine	10	4.9	10.6
Tyrosine	12	5.0	11.3
Tryptophan	10	3.0	7.0
Histidine	8	3.3	8.1
Proline	13	5.0	10.9
None	9	2.6	5.9

[0049]

[Example 2]

A medium of the following composition (glucose 20 g/L, filtered corn steep liquor 5 g/L, potassium dihydrogenphosphate 0.54 g/L, dipotassium hydrogenphosphate 12-hydrate 2.78 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 3.0 g/L, alcohol-based defoaming agent 0.2 g/L, pH 7.5) was dispensed in 100 ml volumes into 500 mL cotton-plugged Erlenmeyer flasks and sterilized in an autoclave at 121°C for 15 minutes to prepare a flask medium for seed culture in 8 flasks.

Next, a medium of the following composition (glucose 40 g/L, corn steep liquor 30 g/L, ammonium sulfate 0.5 g/L, potassium dihydrogenphosphate 2.25 g/L, disodium hydrogenphosphate 12-hydrate 5.7 g/L, calcium chloride dihydrate 0.1 g/L, magnesium sulfate 7-hydrate 0.5 g/L, iron sulfate 7-hydrate 5 g/L, alcohol-based

defoaming agent 0.5 g/L) was dispensed in 2.0 L volumes into 5 L fermentation tanks (8 tanks in total). To these tanks, monosodium L-glutamate monohydrate was added at 0, 1, 5, 15, 30, 50, 100 and 200 mmol/L, respectively, followed by sterilization in an autoclave at 121°C for 30 minutes.

[0050]

A loopful of *Paracoccus carotinifaciens* strain E-396 (FERM BP-4283) was inoculated into the flask medium for seed culture and then cultured with rotary shaking at 29°C for 2 days at 100 rpm. The resulting culture solution was then inoculated in 80 mL volumes into the individual fermentation tanks, followed by aerobic culture at 29°C at an aeration volume of 1 vvm for 100 hours. To maintain a pH of 7.2 during culture, the pH was continuously controlled with 15% aqueous ammonia. Glucose was added in an amount of 30 g on day 1 and day 2 of culture to prevent glucose depletion. In addition, the minimum number of agitation rotations was set to 200 rpm, and the number of agitation rotations was varied such that the dissolved oxygen concentration in the culture solution was maintained at 2 to 4 ppm. Bubbling was detected with a bubble sensor and suppressed by automatic addition of an alcohol-based defoaming agent.

[0051]

At the completion of culture, each culture solution was measured for its carotenoid concentration by HPLC. The results obtained are as shown in Table 2. All the samples containing glutamic acid at concentrations from 1 to 200 mmol/L were found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid.

[0052]

[Table 2]

Glutamic acid concentration mmol/L	0	1	5	15	30	50	100	200
	Produced carotenoid concentration mg/L							
β -Carotene	3.5	4.3	6.2	5.8	5.9	4.9	5.5	4.6
Echinenone	1.3	1.6	1.9	2.1	2.2	2.3	2.0	1.7
3-Hydroxyechinenone	0.2	0.2	0.3	0.3	0.3	0.4	0.3	0.3
Canthaxanthin	1.8	2.0	3.7	3.0	3.1	2.0	2.8	2.4
Phoenicoxanthin	4.4	5.1	7.8	7.2	7.4	6.4	7.0	5.9
β -Cryptoxanthin	0.04	0.05	0.07	0.07	0.08	0.07	0.06	0.05
Astaxanthin	12.9	18.3	27.3	30.8	30.3	30.9	28.6	23.5
Asteroidenone	0.4	0.4	0.4	0.6	0.6	0.9	0.6	0.5
Adonixanthin	5.9	7.8	7.8	9.6	9.3	10.8	9.2	7.6
Zeaxanthin	0.12	0.17	0.18	0.19	0.20	0.21	0.18	0.15
Total Carotenoid	30.6	39.9	55.7	59.7	59.4	58.9	56.2	46.7

[0053]

[Example 3]

Paracoccus carotinifaciens strain E-396 was mutated with N-methyl-N'-nitro-N-nitrosoguanidine to select colonies with darker red color. The selected strains were analyzed for carotenoids in their culture solutions to select a mutant strain Y-1071 with improved astaxanthin productivity.

[0054]

A medium of the following composition (sucrose 30 g/L, Pharmamedia 30 g/L, potassium dihydrogenphosphate 0.8 g/L, dipotassium hydrogenphosphate 4.2 g/L, calcium chloride dihydrate 1 g/L, magnesium sulfate 7-hydrate 12 g/L, iron sulfate 7-hydrate 1 g/L, pH 7.2) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm and sterilized in an autoclave at 121°C for 15 minutes to prepare a test tube medium for seed culture.

Next, a medium of the following composition (sucrose 30 g/L, Pharmamedia 20 g/L, ammonium sulfate 1.5 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 0.1 g/L, magnesium sulfate 7-hydrate 4.5 g/L, iron sulfate 7-hydrate 5 g/L, biotin 1 mg/L, silicon-based defoaming agent 1 g/L) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm (2 tubes in total). One of the tubes was supplemented with monosodium L-glutamate monohydrate to 30 mmol/L, while the other was supplemented with nothing as a control. Finally, the mediums in the tubes were adjusted to pH 7.1 with aqueous sodium hydroxide and then sterilized in an autoclave at 121°C for 20 minutes.

[0055]

The Paracoccus sp. strain Y-1071 selected above was inoculated into the test tube medium for seed culture and then cultured with shaking at 28°C for 2 days at 300 spm. The resulting culture solution was then inoculated in 0.1 ml volumes into the two test tube media and cultured with shaking at 28°C for 4 days at 300 spm.

Each culture solution was measured for its carotenoid concentration by HPLC. The results obtained are as shown in Table 3.

In the mutant Paracoccus sp. strain Y-1071, the sample containing glutamic acid was also found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid.

[0056]

[Table 3]

Glutamic acid concentration		0 mmol/L	30 mmol/L
β-Carotene	mg/L	6.9	13.1
Echinenone	mg/L	3.4	6.5
3-Hydroxyechinenone	mg/L	0.4	0.7
Canthaxanthin	mg/L	5.5	10.0

Phoenicoxanthin	mg/L	13.6	24.8
β -Cryptoxanthin	mg/L	0.04	0.10
Astaxanthin	mg/L	46.4	94.8
Asteroidenone	mg/L	0.9	1.5
Adonixanthin	mg/L	11.3	18.4
Zeaxanthin	mg/L	0.17	0.28
Total Carotenoid	mg/L	88.7	170.2

[0057]

[Example 4]

A medium of the following composition (glucose 20 g/L, dry yeast 5 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 0.1 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 3 g/L, pH 7.2) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm and sterilized in an autoclave at 121°C for 15 minutes to prepare a test tube medium for seed culture.

[0058]

Next, a medium of the following composition (glucose 40 g/L, ammonium sulfate 1.5 g/L, potassium dihydrogenphosphate 0.54 g/L, dipotassium hydrogenphosphate 2.78 g/L, calcium chloride dihydrate 1 g/L, sodium chloride 3 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 5 g/L, zinc sulfate 7-hydrate 2 mg/L, cobalt chloride 6-hydrate 2 mg/L, copper sulfate 5-hydrate 1 mg/L, manganese sulfate 5-hydrate 4 mg/L, sodium molybdate dihydrate 2 mg/L, nickel sulfate 6-hydrate 1 mg/L, sodium selenate 0.5 mg/L, boric acid 5 mg/L, potassium iodide 1 mg/L, cyanocobalamin 1 mg/L, riboflavin 10 mg/L, calcium pantothenate 15 mg/L, pyridoxine hydrochloride salt 20 mg/L, thiamine hydrochloride salt 30 mg/L, ascorbic acid 30 mg/L, folic acid 1 mg/L, niacin 15 mg/L, p-aminobenzoic acid 10 mg/L, biotin 0.1 mg/L, myo-inositol 50 mg/L, choline 10 mg/L, polyether-based defoaming agent 0.2 g/L) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm (4 tubes in total).

For use in this medium, glucose, inorganic salts, trace metals and vitamins were prepared separately. Glucose, inorganic salts and trace metals were sterilized by heating at 121°C for 15 minutes, while vitamins were sterilized by filtration. Then, these 4 solutions were mixed together.

Further, one of the test tubes was supplemented with a heat-sterilized aqueous solution of monosodium L-glutamate monohydrate to 6 g/L (32 mmol/L), another tube was supplemented with a heat-sterilized aqueous yeast extract to 6 g/L, another tube was supplemented with the same yeast extract to 12 g/L, and the other tube was supplemented with nothing. Finally, the mediums in the tubes were aseptically supplemented with 12% aqueous ammonia to give pH 7.2.

[0059]

The mutant *Paracoccus* sp. strain Y-1071 selected in Example 3 was inoculated into the test tube medium for seed culture and then cultured with shaking at 30°C for 2 days at 300 spm. The resulting culture solution was then inoculated in 0.1 ml volumes into the 4 test tube media and cultured with shaking at 30°C for 3 days at 300 spm.

Each culture solution was measured for its carotenoid concentration by HPLC. As shown in Table 4, the sample containing glutamic acid was found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid. Neither of the samples containing yeast extract was found to have a significant improving effect comparable to that of the sample containing glutamic acid.

[0060]

[Table 4]

	Produced carotenoid concentration mg/L			
	Control	Glutamic acid 6 g/L	Yeast extract 6 g/L	Yeast extract 12 g/L
β -Carotene	1.1	2.0	1.5	1.4
Echinenone	0.5	0.9	0.7	0.5
3-Hydroxyechinenone	0.1	0.2	0.2	0.1
Canthaxanthin	0.9	1.8	1.4	1.2
Phoenicoxanthin	1.9	4.1	2.6	2.4
β -Cryptoxanthin	0.01	0.02	0.01	0.01
Astaxanthin	5.3	13.7	7.7	6.1
Asteroidenone	0.1	0.2	0.1	0.1
Adonixanthin	1.4	3.3	1.7	1.6
Zeaxanthin	0.03	0.06	0.04	0.04
Total Carotenoid	11.1	26.3	16.0	13.5

[0061]

[Example 5]

A medium of the following composition (sucrose 20 g/L, filtered corn steep liquor 5 g/L, potassium dihydrogenphosphate 0.54 g/L, dipotassium hydrogenphosphate 12-hydrate 2.78 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 3.0 g/L, alcohol-based defoaming agent 0.2 g/L, pH 7.5) was dispensed in 100 ml volumes into 500 mL cotton-plugged Erlenmeyer flasks and sterilized in an autoclave at 121°C for 15 minutes to prepare a flask medium for seed culture in 2 flasks.

Next, a medium of the following composition (glucose 40 g/L, corn steep liquor 30 g/L, ammonium sulfate 0.5 g/L, potassium dihydrogenphosphate 2.25 g/L, disodium hydrogenphosphate 12-hydrate 5.7 g/L, calcium chloride dihydrate 0.1 g/L, magnesium sulfate 7-hydrate 0.5 g/L, iron sulfate 7-hydrate 5 g/L, alcohol-based defoaming agent 0.5 g/L) was dispensed in 2.0 L volumes into 5 L fermentation tanks (2

tanks in total). One of the fermentation tanks was supplemented with monosodium L-glutamate monohydrate to 15 mmol/L, while the other was supplemented with nothing as a control. These fermentation tanks were sterilized in an autoclave at 121°C for 30 minutes.

[0062]

A loopful of *Paracoccus* sp. strain A-581-1 (FERM BP-4671) was inoculated into the flask medium for seed culture and then cultured with rotary shaking at 27°C for 2 days at 150 rpm. The resulting culture solution was then inoculated in 90 mL volumes into the individual fermentation tanks, followed by aerobic culture at 27°C at an aeration volume of 1 vvm for 100 hours. To maintain a pH of 7.1 during culture, the pH was continuously controlled with 20% aqueous sodium hydroxide. Glucose was added in an amount of 30 g on day 1 and day 2 of culture to prevent glucose depletion. At 22 and 29 hours of culture, monosodium L-glutamate monohydrate and ammonium sulfate were added in amounts of 5 g and 3 g, respectively, per liter of the initial medium. The minimum number of agitation rotations was set to 100 rpm, and the number of agitation rotations was varied such that the dissolved oxygen concentration in the culture solution was maintained at 2 to 4 ppm. An alcohol-based defoaming agent was added in an amount of 0.1 g per hour to prevent bubble formation.

[0063]

At the completion of culture, each culture solution was measured for its carotenoid concentration by HPLC. The results obtained are as shown in Table 5. The sample containing glutamic acid was found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid.

[0064]

[Table 5]

Glutamic acid concentration		0 mmol/L	15 mmol/L
β -Carotene	mg/L	0.92	1.76
Echinenone	mg/L	0.30	0.51
3-Hydroxyechinenone	mg/L	0.03	0.05
Canthaxanthin	mg/L	0.56	1.22
Phoenicoxanthin	mg/L	1.13	2.01
β -Cryptoxanthin	mg/L	0.00	0.01
Astaxanthin	mg/L	3.13	5.97
Asteroidenone	mg/L	0.01	0.02
Adonixanthin	mg/L	1.55	2.10
Zeaxanthin	mg/L	0.02	0.03
Total Carotenoid	mg/L	7.65	13.68

[0065]

[Example 6]

Paracoccus sp. strain A-581-1 (FERM BP-4671) was mutated by ultraviolet irradiation to select colonies with darker red color. The selected strains were analyzed for carotenoids in their culture solutions to select a mutant strain K-185 with improved astaxanthin productivity.

[0066]

A medium of the following composition (sucrose 30 g/L, corn steep liquor 30 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 0.3 g/L, pH 7.2) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm and sterilized in an autoclave at 121°C for 15 minutes to prepare a test tube medium for seed culture.

Next, a medium of the following composition (glucose 30 g/L, soybean meal 20 g/L, ammonium sulfate 1.5 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 0.6 g/L, ester-based defoaming agent 0.2 g/L) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm (2 tubes in total). One of the test tubes was supplemented with monosodium L-glutamate monohydrate to 30 mmol/L, while the other was supplemented with nothing as a control. Finally, the mediums in the tubes were adjusted to pH 7.1 with aqueous ammonia and then sterilized in an autoclave at 121°C for 20 minutes.

[0067]

The Paracoccus sp. strain K-185 was inoculated into the test tube medium for seed culture and then cultured with shaking at 28°C for 2 days at 300 rpm. The resulting culture solution was then inoculated in 0.1 ml volumes into the two test tube mediums and cultured with shaking at 28°C for 3 days at 300 rpm. Each culture solution was measured for its carotenoid concentration by HPLC. The results obtained are as shown in Table 6.

In the mutant Paracoccus sp. strain K-185, the sample containing glutamic acid was also found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid.

[0068]

[Table 6]

Glutamic acid concentration		0 mmol/L	30 mmol/L
β-Carotene	mg/L	1.3	2.1
Echinenone	mg/L	0.4	0.6
3-Hydroxyechinenone	mg/L	0.1	0.1
Canthaxanthin	mg/L	0.9	1.3
Phoenicoxanthin	mg/L	1.3	2.0
	mg/L	0.01	0.02

β -Cryptoxanthin	mg/L	5.6	9.2
Astaxanthin	mg/L	0.1	0.1
Asteroidenone	mg/L	1.2	1.6
Adonixanthin	mg/L	0.02	0.03
Zeaxanthin	mg/L	10.9	17.1
Total Carotenoid			

[0069]

[Example 7]

The strain E-396 (FERM BP-4283) was mutated with N-methyl-N'-nitro-N-nitrosoguanidine to select mutant strain colonies with red-purple color. Carotenoid compounds in their culture solutions were further analyzed by high performance liquid chromatography to select a strain L-25 specifically producing lycopene.

[0070]

A medium of the following composition (sucrose 30 g/L, corn steep liquor 30 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 0.3 g/L, pH 7.2) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm and sterilized in an autoclave at 121°C for 15 minutes to prepare a test tube medium for seed culture.

Next, a medium of the following composition (glucose 30 g/L, filtered corn steep liquor 5 g/L, ammonium sulfate 1.5 g/L, potassium dihydrogenphosphate 1.5 g/L, disodium hydrogenphosphate 12-hydrate 3.8 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 0.6 g/L, ester-based defoaming agent 0.2 g/L) was dispensed in 8 ml volumes into cotton-plugged test tubes having an inner diameter of 18 mm (2 tubes in total). One of the test tubes was supplemented with monosodium L-glutamate monohydrate to 30 mmol/L, while the other was supplemented with nothing as a control. Finally, the mediums in the tubes were adjusted to pH 7.1 with aqueous ammonia and then sterilized in an autoclave at 121°C for 20 minutes.

[0071]

The *Paracoccus* sp. strain L-25 selected above was inoculated into the test tube medium for seed culture and then cultured with shaking at 28°C for 2 days at 300 rpm. The resulting culture solution was then inoculated in 0.1 ml volumes into the two test tube mediums and cultured with shaking at 28°C for 3 days at 300 rpm. Each culture solution was measured for its carotenoid concentration by HPLC. The results obtained are as shown in Table 7.

In the mutant *Paracoccus* sp. strain L-25, the sample containing glutamic acid was also found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid.

[0072]

[Table 7]

Glutamic acid concentration		0 mmol/L	30 mmol/L
Lycopene	mg/L	12.3	19.6
Astaxanthin	mg/L	0.2	0.3
Adonixanthin	mg/L	0.2	0.3
Total Carotenoid	mg/L	12.7	20.2

[0073]

[Example 8]

A medium of the following composition (sucrose 20 g/L, filtered corn steep liquor 5 g/L, potassium dihydrogenphosphate 0.54 g/L, dipotassium hydrogenphosphate 12-hydrate 2.78 g/L, calcium chloride dihydrate 5.0 g/L, magnesium sulfate 7-hydrate 0.7 g/L, iron sulfate 7-hydrate 3.0 g/L, fatty acid-based defoaming agent 0.2 g/L, pH 7.5) was dispensed in 100 ml volumes into 500 mL cotton-plugged Erlenmeyer flasks and sterilized in an autoclave at 121°C for 15 minutes to prepare a flask medium for seed culture in 2 flasks.

Next, a medium of the following composition (sucrose 40 g/L, corn steep liquor 30 g/L, ammonium sulfate 0.5 g/L, potassium dihydrogenphosphate 2.25 g/L, disodium hydrogenphosphate 12-hydrate 5.7 g/L, calcium chloride dihydrate 0.1 g/L, magnesium sulfate 7-hydrate 0.5 g/L, iron sulfate 7-hydrate 5 g/L, fatty acid-based defoaming agent 0.5 g/L) was dispensed in 2.0 L volumes into 5 L fermentation tanks (2 tanks in total). One of the fermentation tanks was supplemented with monosodium L-glutamate monohydrate to 50 mmol/L, while the other was supplemented with nothing as a control. These fermentation tanks were sterilized in an autoclave at 121°C for 30 minutes.

[0074]

A loopful of the mutant *Paracoccus* sp. strain Y-1071 selected in Example 3 was inoculated into the flask medium for seed culture and then cultured with rotary shaking at 28°C for 2 days at 150 rpm. The resulting culture solution was then inoculated in 80 mL volumes into the individual fermentation tanks, followed by aerobic culture at 28°C at an aeration volume of 1 vvm for 120 hours. To maintain a pH of 7.2 during culture, the pH was continuously controlled with 15% aqueous ammonia. Glucose was added in an amount of 30 g on day 1, day 2 and day 3 of culture to prevent glucose depletion. The minimum number of agitation rotations was set to 100 rpm, and the number of agitation rotations was varied such that the dissolved oxygen concentration in the culture solution was maintained at 2 to 3 ppm. Bubbling was detected with a bubble sensor and suppressed by automatic addition of a fatty acid-based defoaming agent.

At the completion of culture, each culture solution was measured for its

carotenoid concentration by HPLC. The results obtained are as shown in Table 8. The sample containing glutamic acid was found to show a higher produced carotenoid concentration when compared to the sample without glutamic acid.

[0075]

[Table 8]

Glutamic acid concentration		0 mmol/L	50 mmol/L
β -Carotene	mg/L	57	93
Echinenone	mg/L	32	62
3-Hydroxyechinenone	mg/L	5	10
Canthaxanthin	mg/L	58	101
Phoenicoxanthin	mg/L	131	217
β -Cryptoxanthin	mg/L	0.3	0.5
Astaxanthin	mg/L	376	805
Asteroidenone	mg/L	5	9
Adonixanthin	mg/L	70	158
Zeaxanthin	mg/L	2	3
Total Carotenoid	mg/L	737	1458

Sequence Listing Free Text

[0076]

SEQ ID NO: 1: Explanation on unknown organism (E-396)

n = a, c, g or t (Location: 1350)

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 30179-209 Seq 11-04-11 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110> Nippon Oil Corporation

<120> Fermentation Method of Carotenoid

<130> PCT09-0034

<150> JP 2008-268106

<151> 2008-10-17

<160> 1

<170> PatentIn version 3.4

<210> 1

<211> 1452

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: E-396

<220>

<221> misc_feature

<222> (1350)..(1350)

<223> n is a, c, g, or t

<400> 1

agtttgatcc	tggtcagaa	cgaacgctgg	cggcaggctt	aacacatgca	agtcgagcga	60
gaccttcggg	tctagcggcg	gacgggtgag	taacgcgtgg	gaacgtgccc	ttctctacgg	120
aatagccccg	ggaaactggg	agtaataccg	tatacgccct	ttgggggaaa	gatttatcgg	180
agaaggatcg	gcccgcgttg	gattaggtag	ttggtggggg	aatggcccac	caagccgacg	240
atccatagct	ggtttgagag	gatgatcagc	cacactggga	ctgagacacg	gcccagactc	300
ctacgggagg	cagcagtggg	gaatcttaga	caatgggggc	aaccctgatc	tagccatgcc	360
gcgtgagtga	tgaaggcctt	agggttgtaa	agctctttca	gctgggaaga	taatgacggt	420
accagcagaa	gaagccccgg	ctaactccgt	gccagcagcc	gcggtataac	ggagggggct	480
agcgttggtc	ggaattactg	ggcgtaaagc	gcacgtaggc	ggactggaaa	gtcagaggtg	540
aaatcccagg	gctcaacctt	ggaactgcct	ttgaaactat	cagtctggag	ttcgagagag	600
gtgagtggaa	ttccgagtgt	agaggtgaaa	ttcgtagata	ttcggaggaa	caccagtggc	660
gaaggcggct	cactggctcg	atactgacgc	tgaggtgcga	aagcgtgggg	agcaaacagg	720

attagatacc	ctggtagtcc	acgccgtaaa	cgatgaatgc	cagacgtcgg	caagcatgct	780
tgtcgggtgc	acacctaacg	gattaagcat	tccgcctggg	gagtacggtc	gcaagattaa	840
aactcaaagg	aattgacggg	ggccccacaca	agcgggtggag	catgtggttt	aattcgaagc	900
aacgcgcaga	accttaccaa	cccttgacat	ggcaggaccg	ctggagagat	tcagctttct	960
cgtaagagac	ctgcacacag	gtgctgcatg	gctgtcgtca	gctcgtgctg	tgagatgttc	1020
ggttaagtcc	ggcaacgagc	gcaaccacg	tccctagtgtg	ccagcaattc	agttgggaac	1080
tctatggaaa	ctgccgatga	taagtcggag	gaaggtgtgg	atgacgtcaa	gtcctcatgg	1140
gccttacggg	ttgggctaca	cacgtgctac	aatgggtggg	acagtgggtt	aatccccaaa	1200
agccatctca	gttcggattg	tcctctgcaa	ctcgagggca	tgaagttgga	atcgctagta	1260
atcgcggaac	agcatgccgc	ggtgaatacg	ttcccggggc	ttgtacacac	cgcccgtcac	1320
accatgggag	ttggttctac	ccgacgacgn	tgcgctaacc	ttcggggggc	aggcggccac	1380
ggtaggatca	gcgactgggg	tgaagtcgta	acaaggtagc	cgtaggggaa	cctgcggctg	1440
gatcacctcc	tt					1452

30179-209

CLAIMS:

1. A method for producing astaxanthin, which comprises culturing an astaxanthin-producing bacterium of the genus *Paracoccus* in a culture medium supplemented with glutamic acid or a glutamate salt, and collecting the astaxanthin from the resulting
5 culture; wherein the bacterium is a mutant that produces astaxanthin at a higher concentration in the culture medium than its parent strain.
2. The method according to claim 1, wherein the bacterium is a mutant of strain E-396 (FERM BP-4283) or A-581-1 (FERM BP-4671) that produces astaxanthin at a higher
10 concentration in the culture medium than either strain E-396 (FERM BP-4283) or A-581-1 (FERM BP-4671).
3. The method according to claim 1 or 2, wherein the concentration of the glutamic acid or glutamate salt in the medium is 1 mmol/L to 200 mmol/L.
4. The method according to claim 1 or 2, wherein the concentration of the glutamic acid or glutamate salt in the medium is 10 mmol/L to 100 mmol/L.
- 15 5. The method according to any one of claims 1 to 4, wherein the bacterium produces at least one further carotenoid selected from the group consisting of canthaxanthin, β -cryptoxanthin, phoenicoxanthin, adonixanthin, echinenone, asteroidenone and 3-hydroxyechinenone at a higher concentration in the culture medium than its parent strain.
- 20 6. The method according to any one of claims 1 to 5, wherein the bacterium has 95% or more sequence identity in the base sequence of DNA corresponding to 16S ribosomal RNA with respect to the nucleotide sequence represented by SEQ ID NO: 1.