



US00PP28850P3

(12) **United States Plant Patent**  
**Koning**

(10) **Patent No.:** **US PP28,850 P3**

(45) **Date of Patent:** **Jan. 2, 2018**

(54) **LAVANDULA PLANT NAMED ‘ANOUK DELUXE 173’**

(50) Latin Name: *Lavandula stoechas*  
Varietal Denomination: **Anouk Deluxe 173**

(71) Applicant: **Lammert Koning**, Nuis (NL)

(72) Inventor: **Lammert Koning**, Nuis (NL)

(73) Assignee: **L. Koning Beheer B.V.**, Nuis (NL)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/999,140**

(22) Filed: **Mar. 31, 2016**

(65) **Prior Publication Data**

US 2017/0290250 P1 Oct. 5, 2017

(51) **Int. Cl.**  
**A01H 5/02** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./445**

(58) **Field of Classification Search**

USPC ..... Plt./445  
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

PLUTO Plant Variety Database Mar. 2, 2017. p. 1.\*

\* cited by examiner

*Primary Examiner* — Annette Para

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Lavandula* plant named ‘Anouk Deluxe 173’, characterized by its upright to slightly outward plant habit; freely branching growth habit; strong durable plants; freely flowering habit; dark violet-colored flowers with purple violet-colored terminal flower bracts positioned on strong and erect peduncles; long flowering period and strong re-flowering habit; and good garden performance.

**2 Drawing Sheets**

**1**

Botanical designation: *Lavandula stoechas*.  
Cultivar denomination: ‘ANOUK DELUXE 173’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct *Lavandula* plant, botanically known as *Lavandula stoechas*, commonly referred to as Lavender and hereinafter referred to by the name ‘Anouk Deluxe 173’.

The new *Lavandula* plant is a product of a planned breeding program conducted by the Inventor in Nuis, The Netherlands. The objective of the breeding program was to develop new unique *Lavandula* plants with good performance and attractive plant form and flower coloration.

The new *Lavandula* plant originated from an open-pollination in May, 2011 of *Lavandula stoechas* ‘Anouk’, disclosed in U.S. Plant Pat. No. 16,685, as the female, or seed, parent with an unknown selection of *Lavandula stoechas*, as the male, or pollen, parent. The new *Lavandula* plant was discovered and selected by the Inventor as a single flowering plant within the progeny of the stated open-pollination in a controlled environment in Nuis, The Netherlands in May, 2012.

Asexual reproduction of the new *Lavandula* plant by terminal cuttings in a controlled greenhouse environment in Boskoop, The Netherlands since August, 2012 has shown that the unique features of this new *Lavandula* plant are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Lavandula* have not been observed under all possible combinations of environmental conditions

**2**

and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity, without, however, any variance in genotype.

5 The following traits have been repeatedly observed and are determined to be the unique characteristics of the new *Lavandula* plant. These characteristics in combination distinguish ‘Anouk Deluxe 173’ as a new and distinct *Lavandula* plant:

1. Upright to slightly outward plant habit.
2. Freely branching growth habit.
3. Strong durable plants.
4. Freely flowering habit.
5. Dark violet-colored flowers with purple violet-colored terminal flower bracts positioned on strong and erect peduncles.
6. Long flowering period and strong re-flowering habit.
7. Good garden performance.

10 Plants of the new *Lavandula* differ primarily from plants of the female parent, ‘Anouk’, in the following characteristics:

1. Plants of the new *Lavandula* and ‘Anouk’ differ in terminal flower bract color as plants of ‘Anouk’ have light purple-colored terminal flower bracts.
2. Plants of the new *Lavandula* have a stronger re-flowering habit than plants of ‘Anouk’.

15 Plants of the new *Lavandula* can be compared to plants of the *Lavandula stoechas* ‘Sanne’, disclosed in U.S. Plant Pat. No. 20,632. In side-by-side comparisons, plants of the new *Lavandula* differ primarily from plants of ‘Sanne’ in the following characteristics:

1. Plants of the new *Lavandula* are more upright than plants of 'Sanne'.
2. Plants of the new *Lavandula* and 'Sanne' differ in terminal flower bract color as plants of 'Sanne' have purple-colored terminal flower bracts.

Plants of the new *Lavandula* can also be compared to plants of the *Lavandula stoechas* 'Toscane', disclosed in U.S. Plant Pat. No. 18,046. In side-by-side comparisons, plants of the new *Lavandula* differ primarily from plants of 'Toscane' in the following characteristics:

1. Plants of the new *Lavandula* and 'Toscane' differ in terminal flower bract color as plants of 'Toscane' have light purple-colored terminal flower bracts.
2. Plants of the new *Lavandula* have a stronger re-flowering habit than plants of 'Toscane'.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Lavandula* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Lavandula* plant.

The photograph on the first sheet is a side perspective view of a typical flowering plant of 'Anouk Deluxe 173' grown in a container.

The photograph on the second sheet is a close-up view of a typical flowering plant of 'Anouk Deluxe 173'.

#### DETAILED BOTANICAL DESCRIPTION

Plants used in the aforementioned photographs and following detailed description were grown in 12-cm containers during the spring and early summer in an outdoor nursery in Boskoop, The Netherlands and under cultural practices typical of commercial *Lavandula* production. Plants were ten months old when the photographs and description were taken. During the production of the plants, day temperatures ranged from 10° C. to 30° C. and night temperatures ranged from 5° C. to 20° C. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Lavandula stoechas* 'Anouk Deluxe 173'.

#### Parentage:

*Female, or seed, parent.*—*Lavandula stoechas* 'Anouk', disclosed in U.S. Plant Pat. No. 16,685.

*Male, or pollen, parent.*—Unknown selection of *Lavandula stoechas*, not patented.

#### Propagation:

*Type.*—Terminal vegetative cuttings.

*Time to initiate roots, summer and winter.*—About ten days at temperatures about 18° C.

*Time to produce a rooted young plant, summer.*—About 60 days at temperatures about 20° C.

*Time to produce a rooted young plant, winter.*—About 70 days at temperatures about 20° C.

*Root description.*—Fine, fibrous; initially white in color and becoming more brown with development; actual color of the roots is dependent on substrate composition, water quality, fertilizer type and formulation.

*Rooting habit.*—Freely branching; dense.

#### Plant description:

*Plant and growth habit.*—Herbaceous perennial; upright to slightly outward plant habit; freely branching habit; moderately vigorous growth habit; flowers arranged in verticillasters on crowded spikes with showy terminal flower bracts.

*Plant height, soil level to top of floral plane.*—About 28.2 cm.

*Plant width.*—About 29.8 cm.

*Lateral branch description.*—Quantity per plant: About 58 primary lateral branches developing per plant with numerous secondary branches. Length: About 10.4 cm. Diameter: About 2.5 mm. Internode length: About 1.6 mm. Strength: Strong. Aspect: Upright to slightly outward. Texture and luster: Moderately tomentose; matte. Color: Close to 145A to 145B; pubescence, close to 157D.

*Leaf description.*—Arrangement: Opposite, simple; sessile. Length: About 3.3 cm. Width: About 6 mm. Shape: Narrowly oblanceolate to lanceolate. Apex: Acute. Base: Cuneate. Margin: Entire; revolute. Texture and luster, upper and lower surfaces: Densely tomentose; matte. Fragrance: Strongly aromatic, pungent. Venation pattern: Pinnate. Color: Developing leaves, upper surface: Close to 137B. Developing leaves, lower surface: Close to 137C. Fully expanded leaves, upper surface: Close to between 137B and 189A; venation, close to 144B; pubescence, close to 192D. Fully expanded leaves, lower surface: Close to 138A; venation, close to 148B; pubescence, close to 192D.

#### Flower description:

*Flower type, arrangement and habit.*—Small single salverform flowers arranged in compact verticillasters on crowded terminal spikes; freely flowering, about 120 flowers developing per inflorescence and more than 7,000 flowers developing per plant; flowers with two-lobed upper lip and three-lobed lower lip; flowers face mostly outward to slightly upright on the spike; inflorescences with showy terminal flower bracts.

*Natural flowering season.*—Long flowering period; continuous from late spring into late summer in The Netherlands; strong re-flowering habit.

*Flower longevity on the plant.*—Individual flowers last about ten days on the plant; flowers not persistent.

*Fragrance.*—None detected.

*Flower buds.*—Length: About 6 mm. Diameter: About 2 mm. Shape: Narrowly oblong. Texture and luster: Moderately tomentose; matte. Color: Close to 145C; towards the apex, close to N187A.

*Inflorescence size.*—Height: About 5.3 cm. Diameter, at terminal flower bracts: About 2.1 cm. Diameter, below flower bracts: About 1.5 cm.

*Flower size.*—Diameter: About 3.5 mm. Depth (height): About 8.5 mm.

*Petals.*—Quantity and arrangement: Upper lip, two-lobed and lower lip, three-lobed. Length, upper lip: About 8.5 mm. Length, lower lip: About 7 mm. Width, upper lip: About 2.5 mm. Width, lower lip: About 4 mm. Shape: Roughly spatulate; lower 70% fused into a tube. Apex: Obtuse, rounded. Margin: Entire. Texture and luster, upper (inner) and lower (outer) surfaces: Smooth, glabrous, velvety; matte. Color: When opening and fully opened, upper (in-

ner) surface: Upper lip, slightly darker than 86A; lower lip, close to 86B, towards the base, close to NN155D; color does not fade with development. When opening and fully opened, lower (outer) surface: Close to 86A to 86B; towards the base, close to NN155D; color does not fade with development.

*Terminal flower bracts*.—Quantity and arrangement: About four positioned upright at inflorescence apex. Length: About 2.4 cm. Width: About 8 mm. Shape: Narrowly obovate. Apex: Obtuse. Base: Narrowly cuneate. Margin: Entire; undulate. Texture and luster, upper and lower surfaces: Moderately tomentose (minute hairs); matte. Color, upper surface: Close to N81B; venation, close to N81B; pubescence, close to N187D. Color, lower surface: Close to N81B; venation, close to N187A; pubescence, close to N187D.

*Basal flower bracts*.—Quantity and arrangement: Each group of five flowers is subtended by a single basal flower bract. Length: About 8 mm. Width: About 7 mm. Shape: Broadly rhomboidal. Apex: Aristate. Base: Cuneate. Margin: Entire. Texture and luster, upper and lower surfaces: Moderately tomentose; matte. Color, upper surface: Close to 147D; towards the margins, close to N77B; pubescence, close to NN155C. Color, lower surface: Close to 148C; venation, close to N187A; pubescence, close to NN155C.

*Sepals*.—Quantity and arrangement: Five, fused into a campanulate tube. Length: About 4 mm. Width: About 0.75 mm. Shape: Lanceolate. Apex: Acute. Margin: Entire. Texture and luster, upper (inner) and lower (outer) surfaces: Densely pubescent; matte. Color: When opening, upper (inner) surface: Close to 147C tinged with close to 195A; pubescence, close to NN155C. When opening, lower (outer) surface: Close to 145C; towards the apex, close to

N187A; pubescence, close to NN155C. Fully opened, upper (inner) surface: Close to 147C tinged with close to 195A; pubescence, close to NN155C. Fully opened, lower (outer) surface: Close to 145C; towards the apex, tinged with close to N187B; pubescence, close to NN155C.

*Peduncles*.—Length: About 6 cm. Diameter: About 2 mm. Aspect: Mostly upright. Strength: Strong. Texture and luster: Densely tomentose; matte. Color: Close to 143A and 145C; towards the apex, close to 141A and N77C to N77D; pubescence, close to 157D.

*Reproductive organs*.—Stamens: Quantity per flower: Four. Filament length: About 1 mm. Filament color: Close to 86D. Anther shape: Reniform. Anther length: About 0.5 mm. Anther color: Close to 148D. Pollen amount: Moderate. Pollen color: Close to 13B. Pistils: Quantity per flower: One. Pistil length: About 3.5 mm. Stigma shape: Club-shaped. Stigma color: Close to N92A. Style length: About 3 mm. Style color: Close to 85D. Ovary color: Close to 143C.

*Seeds and fruits*.—Seed and fruit production has not been observed on plants of the new *Lavandula*.

Disease & pest resistance: Plants of the new *Lavandula* have not been noted to be resistant to pathogens and pests common to *Lavandula* plants.

Garden performance: Plants of the new *Lavandula* have exhibited good tolerance to rain and wind and have been observed to tolerate high temperatures about 40° C. and to be hardy to USDA Hardiness Zone 7.

It is claimed:

1. A new and distinct *Lavandula* plant named 'Anouk Deluxe 173' as illustrated and described.

\* \* \* \* \*



