



(86) **Date de dépôt PCT/PCT Filing Date:** 2007/02/15
(87) **Date publication PCT/PCT Publication Date:** 2007/08/23
(45) **Date de délivrance/Issue Date:** 2014/04/08
(85) **Entrée phase nationale/National Entry:** 2008/08/07
(86) **N° demande PCT/PCT Application No.:** US 2007/062260
(87) **N° publication PCT/PCT Publication No.:** 2007/095628
(30) **Priorité/Priority:** 2006/02/15 (US60/773,513)


(51) **Cl.Int./Int.Cl.** **C07D 471/04** (2006.01),
A61K 31/438 (2006.01), **A61P 35/00** (2006.01),
C07D 491/04 (2006.01), **C07D 495/04** (2006.01)

(72) **Inventeurs/Inventors:**
PENNING, THOMAS D., US;
ZHU, GUI-DONG, US;
GANDHI, VIRAJKUMAR B., US;
GONG, JIANCHUN, US;
GIRANDA, VINCENT L., US

(73) **Propriétaire/Owner:**
ABBVIE INC., US

(74) **Agent:** TORYS LLP

(54) **Titre : PYRAZOLOQUINOLONES EN TANT QU'INHIBITEURS EFFICACES DE PARP**
(54) **Title: PYRAZOLOQUINOLONES ARE POTENT PARP INHIBITORS**



Formula (I)

(57) **Abrégé/Abstract:**

Compounds of Formula (I) inhibit the PARP enzyme and are useful for treating a disease or a disorder associated with PARP. Also disclosed are pharmaceutical compositions comprising compounds of Formula (I), methods of treatment comprising compounds of Formula (I), and methods of inhibiting the PARP enzyme comprising compounds of Formula (I).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 August 2007 (23.08.2007)

PCT

(10) International Publication Number
WO 2007/095628 A1

(51) International Patent Classification:

C07D 471/04 (2006.01) **C07D 491/04** (2006.01)
A61K 31/438 (2006.01) **C07D 495/04** (2006.01)
A61P 35/00 (2006.01)

(74) Agents: **CORBIN, Johanna, M.** et al.; Dept. 377 Bldg
AP6A-1, 100 Abbott Park Road, Abbott Park, Illinois
60064-6008 (US).

(21) International Application Number:

PCT/US2007/062260

(22) International Filing Date:

15 February 2007 (15.02.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/773,513 15 February 2006 (15.02.2006) US

(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; Dept. 377 Bldg AP6A-1, 100 Abbott Park Road, Abbott Park, Illinois 60064-6008 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PENNING, Thomas, D.** [US/US]; 321 Highview Avenue, Elmhurst, Illinois 60126 (US). **ZHU, Gui-dong** [US/US]; 1395 Almaden Lane, Gurnee, Illinois 60031 (US). **GANDHI, Virajkumar, B.** [IN/US]; 5335 Mahogany Ct., Gurnee, Illinois 60031 (US). **GONG, Jianchun** [CN/US]; 130 Ferndale Road, Deerfield, Illinois 60015 (US). **GIRANDA, Vincent, L.** [US/US]; 272 South Fork Drive, Gurnee, Illinois 60031 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRAZOLOQUINOLONES ARE POTENT PARP INHIBITORS



(I)

(57) Abstract: Compounds of Formula (I) inhibit the PARP enzyme and are useful for treating a disease or a disorder associated with PARP. Also disclosed are pharmaceutical compositions comprising compounds of Formula (I), methods of treatment comprising compounds of Formula (I), and methods of inhibiting the PARP enzyme comprising compounds of Formula (I).

WO 2007/095628 A1

PYRAZOLOQUINOLONES ARE POTENT PARP INHIBITORS

This application claims priority from U.S. Provisional Patent Application Serial No. 60/773,513, filed February 15, 2006 .

5

Technical Field

The present invention relates to pyrazoloquinolones, their preparation, and their use as inhibitors of the enzyme poly(ADP-ribose)polymerase for the preparation of drugs.


10 Background

Poly(ADP-ribose)polymerase (PARP) or poly(ADP-ribose)synthase (PARS) has an essential role in facilitating DNA repair, controlling RNA transcription, mediating cell death, and regulating immune response. These actions make PARP inhibitors targets for a broad spectrum of disorders. PARP inhibitors have demonstrated efficacy in numerous models of disease, particularly in models of ischemia reperfusion injury, inflammatory disease, degenerative diseases, protection from adverse effects of cytotoxic compounds, and the potentiation of cytotoxic cancer therapy. PARP has also been indicated in retroviral infection and thus inhibitors may have use in antiretroviral therapy. PARP inhibitors have been efficacious in preventing ischemia reperfusion injury in models of myocardial infarction, stroke, other neural trauma, organ transplantation, as well as reperfusion of the eye, kidney, gut and skeletal muscle. Inhibitors have been efficacious in inflammatory diseases such as arthritis, gout, inflammatory bowel disease, CNS inflammation such as MS and allergic encephalitis, sepsis, septic shock, hemorrhagic shock, pulmonary fibrosis, and uveitis. PARP inhibitors have also shown benefit in several models of degenerative disease including diabetes (as well as complications) and Parkinsons disease. PARP inhibitors can ameliorate the liver toxicity following acetaminophen overdose, cardiac and kidney toxicities from doxorubicin and platinum based antineoplastic agents, as well as skin damage secondary to sulfur mustards. In various cancer models, PARP inhibitors have been shown to potentiate radiation and chemotherapy by increasing cell death of cancer cells, limiting tumor growth, decreasing metastasis, and prolonging the survival of tumor-bearing animals.

15
20
25
30

Summary of the Invention

In one embodiment, the present invention provides compounds of Formula (I)



Formula (I)

and therapeutically acceptable salts, prodrugs and salts of prodrugs thereof, wherein

R¹ and R² together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl or heterocycle, the cycloalkyl or heterocycle may be unsubstituted or substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C1-C6 alkyl, alkynyl, aryl, arylalkoxycarbonyl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydroxy, hydroxyalkyl, nitro, NR^AR^B, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, and (NR^AR^B)sulfonyl;

R³ is selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C1-C6 alkyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydrogen, hydroxy, hydroxyalkyl, nitro, NR^AR^B, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, and (NR^AR^B)sulfonyl;

X is NR⁴;

R⁴ is selected from the group consisting of C1-C6 alkyl, alkoxycarbonylalkyl, arylalkyl, carboxyalkyl, cyanoalkyl, cycloalkyl, cycloalkylalkyl, hydrogen, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, -alkyl-O-C(O)-(NR^AR^B), formylalkyl, heteroarylalkyl, heterocyclealkyl, hydroxyalkyl, alkoxyalkyl, and haloalkyl; and

R^A and R^B are independently selected from the group consisting of hydrogen, alkyl, aryl, arylalkyl, arylalkylcarbonyl, arylalkoxy, arylalkoxycarbonyl, cycloalkyl, cycloalkylalkyl, and heterocycle;


if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is heterocycle then the heterocycle may not be substituted with methyl or methylcarbonyl;

and if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is a cycloalkyl then the cycloalkyl must be substituted; and

wherein the aryl, cycloalkyl, heterocycle and heterocyclealkyl represented by R^3 and R^4 either themselves or as part of another moiety, are independently unsubstituted or substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, C1-C6-alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, oxo, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, $(NR^A R^B)$ sulfonyl, alkylsulfonyl, heterocycle, heterocyclealkyl, heteroaryl, and heteroarylalkyl, wherein the substituent moieties are themselves further unsubstituted.

Detailed Description of the Invention

In another embodiment, the present invention provides compounds of Formula (I)



(I),

and therapeutically acceptable salts, prodrugs and salts of prodrugs thereof, wherein

R^1 and R^2 together with the atoms to which they are attached form a 5, 6, or 7 membered cycloalkyl or heterocycle, the cycloalkyl or heterocycle may be unsubstituted or substituted with a substituent independently selected from the group consisting of heterocycle and $NR^A R^B$;

R^3 is selected from the group consisting of C1-C6 alkyl, aryl, cycloalkyl, heterocycle, heteroaryl, hydrogen, and $(NR^A R^B)$ alkyl;

X is NR^4 ;

R^4 is selected from the group consisting of C1-C6 alkyl, -alkyl-O-C(O)- $(NR^A R^B)$, alkoxycarbonylalkyl, arylalkyl, carboxyalkyl, cyanoalkyl, hydrogen, haloalkyl, heteroarylalkyl, heterocyclealkyl, hydroxyalkyl, formylalkyl, and $(NR^A R^B)$ alkyl;


R^A and R^B are independently selected from the group consisting of hydrogen, C1-C6 alkyl, arylalkoxycarbonyl, cycloalkyl, cycloalkylalkyl, and heterocycle;

if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is heterocycle then the heterocycle may not be substituted with methyl or methylcarbonyl;

and if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is a cycloalkyl then the cycloalkyl must be substituted; and

wherein the aryl, cycloalkyl, heterocycle and heterocyclealkyl represented by R^3 and R^4 either themselves or as part of another moiety, are independently unsubstituted or substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, C1-C6-alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, oxo, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, $(NR^A R^B)$ sulfonyl, alkylsulfonyl, heterocycle, heterocyclealkyl, heteroaryl, and heteroarylalkyl, wherein the substituent moieties are themselves further unsubstituted.

In another embodiment, the present invention provides compounds of Formula (I)



Formula (I)

5 and therapeutically acceptable salts, prodrugs and salts of prodrugs thereof, wherein

R¹ and R² together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl or heterocycle, the cycloalkyl or heterocycle may be unsubstituted or substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C1-C6 alkyl, alkynyl, aryl, 10 arylalkoxycarbonyl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydroxy, hydroxyalkyl, nitro, NR^AR^B, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, and (NR^AR^B)sulfonyl;

R³ is selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, 15 alkoxycarbonylalkyl, C2-C6 alkyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydrogen, hydroxy, hydroxyalkyl, nitro, NR^AR^B, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, and (NR^AR^B)sulfonyl;

20 X is NR⁴;

R⁴ is selected from the group consisting of C2-C6 alkyl, arylalkyl, cycloalkyl, cycloalkylalkyl, hydrogen, (NR^AR^B)alkyl, hydroxyalkyl, alkoxyalkyl, and haloalkyl; and R^A and R^B are independently selected from the group consisting of hydrogen, alkyl, aryl, arylalkyl, arylalkylcarbonyl, arylalkoxy, arylalkoxycarbonyl, cycloalkyl, cycloalkylalkyl, and 25 heterocycle; and

wherein the aryl, cycloalkyl, heterocycle and heterocyclealkyl represented by R^3 and R^4 either themselves or as part of another moiety, are independently unsubstituted or substituted with 1, 2, or 3 substituents independently selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, C1-C6-alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, oxo, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, $(NR^A R^B)$ sulfonyl, alkylsulfonyl, heterocycle, heterocyclealkyl, heteroaryl, and heteroarylalkyl, wherein the substituent moieties are themselves further unsubstituted.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered heterocycle.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein X is NR^4 .

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl..

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered heterocycle.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl; and R^3 is selected from the group consisting of C1-C6 alkyl, aryl, cycloalkyl, heterocycle, heteroaryl, hydrogen, and $(NR^A R^B)$ alkyl.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl; and R^3 is selected from the group consisting of aryl, cycloalkyl, heterocycle, and heteroaryl.

5 In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl; X is NR^4 ; R^4 is C1-C6 alkyl; and R^3 is selected from the group consisting of aryl, cycloalkyl, heterocycle, and heteroaryl.

10 In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^3 is selected from the group consisting of aryl, cycloalkyl, heterocycle, and heteroaryl.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^4 is selected from the group consisting of C1-C6 alkyl, -alkyl-O-C(O)-(NR^AR^B), alkoxy carbonylalkyl, arylalkyl, carboxyalkyl, cyanoalkyl, 15 hydrogen, haloalkyl, heteroarylalkyl, heterocyclealkyl, hydroxyalkyl, formylalkyl, and (NR^AR^B)alkyl.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 6 membered cycloalkyl.

20 In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^3 is a heterocycle.

In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^3 is pyrrolidinyl.

25 In another embodiment, the present invention provides compounds of Formula (I) or a therapeutically acceptable salt thereof wherein R^1 and R^2 together with the atoms to which they are attached form a 6 membered cycloalkyl; and R^3 is pyrrolidinyl.

In another embodiment, the present invention provides a pharmaceutical composition comprising a compound of Formula (I), or a therapeutically acceptable salt thereof, in combination with a therapeutically acceptable carrier.

In another embodiment, the present invention provides a method of inhibiting PARP in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

5 In another embodiment, the present invention provides a method of treating cancer in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

10 In another embodiment, the present invention provides a method for decreasing tumor volume in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

15 In another embodiment, the present invention provides a method of treating leukemia, colon cancer, glioblastomas, lymphomas, melanomas, carcinomas of the breast, or cervical carcinomas in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

20 In another embodiment, the present invention provides a method of potentiation of cytotoxic cancer therapy in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

25 In another embodiment, the present invention provides a method of potentiation of radiation therapy in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

30 In another embodiment, the present invention provides a method of treating ischemia reperfusion injury associated with, but not limited to, myocardial infarction, stroke, other neural trauma, and organ transplantation, in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of reperfusion including, but not limited to, reperfusion of the eye, kidney, gut and skeletal muscle, in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating inflammatory diseases including, but not limited to, arthritis, gout, inflammatory bowel disease, CNS inflammation, multiple sclerosis, allergic encephalitis, sepsis, septic shock, hemorrhagic shock, pulmonary fibrosis, and uveitis in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating immunological diseases or disorders such as rheumatoid arthritis and septic shock in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating degenerative disease including, but not limited to, diabetes and Parkinsons disease, in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating hypoglycemia in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating retroviral infection in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating liver toxicity following acetaminophen overdose in a mammal in recognized need of such treatment

comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating cardiac and kidney toxicities from doxorubicin and platinum based antineoplastic agents in a mammal in
5 recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a method of treating skin damage secondary to sulfur mustards in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I)
10 or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for inhibiting the PARP enzyme in a mammal in recognized need of such treatment.

In another embodiment, the present invention provides a use of a compound of Formula
15 (I), or a therapeutically acceptable salt thereof, to prepare a medicament for inhibiting tumor growth in a mammal in recognized need of such treatment.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating cancer in a mammal in recognized need of such treatment.

20 In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating leukemia, colon cancer, glioblastomas, lymphomas, melanomas, carcinomas of the breast, or cervical carcinomas in a mammal in a mammal in recognized need of such treatment.

In another embodiment, the present invention provides a use of a compound of Formula
25 (I), or a therapeutically acceptable salt thereof, to prepare a medicament for potentiation of cytotoxic cancer therapy in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula
30 (I), or a therapeutically acceptable salt thereof, to prepare a medicament for potentiation of

radiation in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating ischemia reperfusion injury associated with, but not limited to, myocardial infarction, stroke, other neural trauma, and organ transplantation, in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating reperfusion including, but not limited to, reperfusion of the eye, kidney, gut and skeletal muscle, in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating inflammatory diseases including, but not limited to, arthritis, gout, inflammatory bowel disease, CNS inflammation, multiple sclerosis, allergic encephalitis, sepsis, septic shock, hemorrhagic shock, pulmonary fibrosis, and uveitis in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating immunological diseases or disorders such as rheumatoid arthritis and septic shock in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating degenerative disease including, but not limited to, diabetes and Parkinsons disease, in a mammal in recognized

need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating
5 hypoglycemia in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating retroviral
10 infection in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating liver toxicity
15 following acetaminophen overdose in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating cardiac and
20 kidney toxicities from doxorubicin and platinum based antineoplastic agents in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

In another embodiment, the present invention provides a use of a compound of Formula (I), or a therapeutically acceptable salt thereof, to prepare a medicament for treating skin damage
25 secondary to sulfur mustards in a mammal in recognized need of such treatment comprising administering to the mammal a therapeutically acceptable amount of a compound of Formula (I) or a therapeutically acceptable salt thereof.

Definitions

As used throughout this specification and the appended claims, the following terms have the following meanings:

The term "alkenyl" as used herein, means a straight or branched chain hydrocarbon containing from 2 to 10 carbons and containing at least one carbon-carbon double bond formed by the removal of two hydrogens. Representative examples of alkenyl include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl.

The term "alkoxy" as used herein, means an alkyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy, tert-butoxy, pentyloxy, and hexyloxy.

The term "alkoxyalkyl" as used herein, means at least one alkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkoxyalkyl include, but are not limited to, tert-butoxymethyl, 2-ethoxyethyl, 2-methoxyethyl, and methoxymethyl.

The term "alkoxycarbonyl" as used herein, means an alkoxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkoxycarbonyl include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, and tert-butoxycarbonyl.

The term "alkoxycarbonylalkyl" as used herein, means an alkoxycarbonyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

The term "alkyl" as used herein, means a straight or branched chain hydrocarbon containing from 1 to 10 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, n-heptyl, n-octyl, n-nonyl, and n-decyl.

The term "C1-C6 alkyl" as used herein, means a straight or branched chain hydrocarbon containing from 1 to 6 carbon atoms.

The term "C₁-alkyl" means methyl.

The term "C₂-alkyl" means ethyl.

The term "C₃-alkyl" means prop-1-yl and prop-2-yl (isopropyl).

The term "C₄-alkyl" means but-1-yl, but-2-yl, 2-methylprop-1-yl, and 2-methylprop-2-yl (tert-butyl).

5 The term "C₅-alkyl" means 2,2-dimethylprop-1-yl (neo-pentyl), 2-methylbut-1-yl, 2-methylbut-2-yl, 3-methylbut-1-yl, 3-methylbut-2-yl, pent-1-yl, pent-2-yl, and pent-3-yl.

10 The term "C₆-alkyl" means 2,2-dimethylbut-1-yl, 2,3-dimethylbut-1-yl, 2,3-dimethylbut-2-yl, 3,3-dimethylbut-1-yl, 3,3-dimethylbut-2-yl, 2-ethylbut-1-yl, hex-1-yl, hex-2-yl, hex-3-yl, 2-methylpent-1-yl, 2-methylpent-2-yl, 2-methylpent-3-yl, 3-methylpent-1-yl, 3-methylpent-2-yl, 3-methylpent-3-yl, 4-methylpent-1-yl, and 4-methylpent-2-yl.

The term "alkylcarbonyl" as used herein, means an alkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkylcarbonyl include, but are not limited to, acetyl, 1-oxopropyl, 2,2-dimethyl-1-oxopropyl, 1-oxobutyl, and 1-oxopentyl.

15 The term "alkylcarbonyloxy" as used herein, means an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkylcarbonyloxy include, but are not limited to, acetyloxy, ethylcarbonyloxy, and tert-butylcarbonyloxy.

20 The term "-alkyl-O-C(O)-(NR^AR^B)" as used herein, means a (NR^AR^B)carbonyloxy group, as defined herein, appended to the parent molecule through an alkyl group.

The term "alkylsulfonyl" as used herein, means an alkyl group appended to the parent molecular moiety through a sulfonyl.

25 The term "alkylthio" as used herein, means an alkyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom. Representative examples of alkylthio include, but are not limited, methylthio, ethylthio, tert-butylthio, and hexylthio.

The term "alkylthioalkyl" as used herein, means an alkylthio group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkylthioalkyl include, but are not limited, methylthiomethyl and 2-(ethylthio)ethyl.

The term "alkynyl" as used herein, means a straight or branched chain hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond. Representative examples of alkynyl include, but are not limited, to acetylenyl, 1-propynyl, 2-propynyl, 3-butynyl, 2-pentynyl, and 1-butynyl.

5 The term "aryl," as used herein, means a phenyl group or a naphthyl group.

The aryl groups are unsubstituted or substituted with one, two, three, four, or five substituents independently selected from the group consisting of alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, carboxy, cyano, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro,
10 $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, and tetrazolyl.

The term "arylalkyl" as used herein, means an aryl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, 1-methyl-3-phenylpropyl, and 2-naphth-2-ylethyl.

15 The term "arylalkoxy" as used herein, means an aryl group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of arylalkoxy include, but are not limited to, 2-phenylethoxy, 3-naphth-2-ylpropoxy, and 5-phenylpentyloxy.

The term "arylalkoxycarbonyl" as used herein, means an arylalkoxy group, as defined
20 herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylalkoxycarbonyl include, but are not limited to, benzyloxycarbonyl and naphth-2-ylmethoxycarbonyl.

The term "carbonyl" as used herein, means a $-C(O)-$ group.

The term "carboxy" as used herein, means a $-CO_2H$ group.

25 The term "carboxyalkyl" as used herein, means a carboxy group, as defined herein, attached to the parent molecular moiety through an alkyl group.

The term "cyano" as used herein, means a $-CN$ group.

The term "cyanoalkyl" as used herein, means a cyano group appended to the parent molecular moiety through an alkyl group.

The term "cycloalkyl" as used herein, means a saturated cyclic hydrocarbon group containing from 3 to 8 carbons, examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

5 The cycloalkyl groups are unsubstituted or substituted with 1, 2, 3, or 4 substituents selected from alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, carboxy, cyano, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, oxo, $-NR^A R^B$, and $(NR^A R^B)$ carbonyl.

The term "cycloalkylalkyl" as used herein, means a cycloalkyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

10 Representative examples of cycloalkylalkyl include, but are not limited to, cyclopropylmethyl, 2-cyclobutylethyl, cyclopentylmethyl, cyclohexylmethyl, and 4-cycloheptylbutyl.

The term "formyl" as used herein, means a $-C(O)H$ group.

The term "formylalkyl" as used herein, means a formyl group, as defined herein, attached to the parent molecular moiety through an alkyl group.

15 The term "halo" or "halogen" as used herein, means $-Cl$, $-Br$, $-I$ or $-F$.

The term "haloalkoxy" as used herein, means at least one halogen, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of haloalkoxy include, but are not limited to, chloromethoxy, 2-fluoroethoxy, trifluoromethoxy, and pentafluoroethoxy.

20 The term "haloalkyl" as used herein, means at least one halogen, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of haloalkyl include, but are not limited to, chloromethyl, 2-fluoroethyl, trifluoromethyl, pentafluoroethyl, and 2-chloro-3-fluoropentyl.

25 The term "heteroaryl," as used herein, means a monocyclic heteroaryl ring or a bicyclic heteroaryl ring. The monocyclic heteroaryl ring is a 5 or 6 membered ring. The 5 membered ring has two double bonds and contains one, two, three or four heteroatoms independently selected from the group consisting of N, O, and S. The 6 membered ring has three double bonds and contains one, two, three or four heteroatoms independently selected from the group consisting of N, O, and S. The bicyclic heteroaryl ring consists of the 5 or 6 membered
30 heteroaryl ring fused to a phenyl group or the 5 or 6 membered heteroaryl ring is fused to another

5 or 6 membered heteroaryl ring. Nitrogen heteroatoms contained within the heteroaryl may be optionally oxidized to the N-oxide. The heteroaryl is connected to the parent molecular moiety through any carbon atom contained within the heteroaryl while maintaining proper valence.

Representative examples of heteroaryl include, but are not limited to, benzothienyl,

5 benzoxadiazolyl, cinnolinyl, furopyridinyl, furyl, imidazolyl, indazolyl, indolyl, isoxazolyl, isoquinolinyl, isothiazolyl, naphthyridinyl, oxadiazolyl, oxazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrazolyl, pyrrolyl, pyridinium N-oxide, quinolinyl, tetrazolyl, thiadiazolyl, thiazolyl, thienopyridinyl, thienyl, triazolyl, and triazinyl.

The heteroaryl groups are unsubstituted or substituted with 1, 2, 3, or 4 substituents
10 independently selected from alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, carboxy, cyano, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, $-NR^A R^B$, and $(NR^A R^B)$ carbonyl.

The term "heteroarylalkyl" as used herein, means a heteroaryl, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

15 Representative examples of heteroarylalkyl include, but are not limited to, pyridinymethyl.

The term "heterocycle" or "heterocyclic" as used herein, means a monocyclic or bicyclic heterocyclic ring. The monocyclic heterocyclic ring consists of a 3, 4, 5, 6, 7, or 8 membered ring containing at least one heteroatom independently selected from O, N, and S. The 3 or 4 membered ring contains 1 heteroatom selected from the group consisting of O, N and S. The 5
20 membered ring contains zero or one double bond and one, two or three heteroatoms selected from the group consisting of O, N and S. The 6 or 7 membered ring contains zero, one or two double bonds and one, two or three heteroatoms selected from the group consisting of O, N and S. The bicyclic heterocyclic ring consists of a monocyclic heterocyclic ring fused to a cycloalkyl group or the monocyclic heterocyclic ring fused to a phenyl group or the monocyclic
25 heterocyclic ring fused to another monocyclic heterocyclic ring. The heterocycle is connected to the parent molecular moiety through any carbon or nitrogen atom contained within the heterocycle while maintaining proper valence. Representative examples of heterocycle include, but are not limited to, azetidiny, azepanyl, aziridinyl, diazepanyl, 1,3-dioxanyl, 1,3-dioxolanyl, 1,3-dithiolanyl, 1,3-dithianyl, imidazolinyl, imidazolidinyl, isothiazolinyl, isothiazolidinyl,
30 isoxazolinyl, isoxazolidinyl, morpholinyl, oxadiazolinyl, oxadiazolidinyl, oxazolinyl,

oxazolidinyl, piperazinyl, piperidinyl, pyrazolinyl, pyrazolidinyl, pyrrolinyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothienyl, thiadiazolinyl, thiadiazolidinyl, thiazolinyl, thiazolidinyl, thiomorpholinyl, 1,1-dioxidothiomorpholinyl (thiomorpholine sulfone), thiopyranyl, and trithianyl.

5 The heterocycles are unsubstituted or substituted with 1, 2, or 3 substituents independently selected from alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, 10 sulfonyl, alkylsulfonyl, tetrahydropyranyl, and heteroarylalkyl wherein the heteroaryl is pyridinyl.

 The term "heterocyclealkyl" as used herein, means a heterocycle, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

 The term "heterocyclecarbonyl" as used herein, means a heterocycle group, as defined 15 herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclecarbonyl include, but are not limited to, 1,3-benzodioxol-4-ylcarbonyl, pyridin-3-ylcarbonyl, pyrimidin-2-ylcarbonyl, tetrahydrofuran-2-ylcarbonyl, tetrahydrofuran-3-ylcarbonyl, tetrahydro-2H-pyran-2-ylcarbonyl, tetrahydro-2H-pyran-4-ylcarbonyl, tetrahydrothien-2-ylcarbonyl and tetrahydrothien-3-ylcarbonyl.

20 The term "heterocyclecarbonylalkyl" as used herein, means a heterocyclecarbonyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

 The term "heterocyclesulfonyl" as used herein, means a heterocycle group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein.

25 The term "hydroxy" as used herein, means an -OH group.

 The term "hydroxyalkyl" as used herein, means at least one hydroxy group, as defined herein, is appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of hydroxyalkyl include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2,3-dihydroxypentyl, and 2-ethyl-4-hydroxyheptyl.

30 The term "mercapto" as used herein, means a -SH group.

The term "nitro" as used herein, means a -NO_2 group.

The term "nonaromatic" as used herein, means that a 4 membered nonaromatic ring contains zero double bonds, a 5 membered nonaromatic ring contains zero or one double bond, a 6, 7, or 8 membered nonaromatic ring contains zero, one, or two double bonds.

5 The term " $\text{NR}^{\text{A}}\text{R}^{\text{B}}$ " as used herein, means two groups, R^{A} and R^{B} , which are appended to the parent molecular moiety through a nitrogen atom. R^{A} and R^{B} are each independently hydrogen, alkyl, and alkylcarbonyl. Representative examples of $\text{NR}^{\text{A}}\text{R}^{\text{B}}$ include, but are not limited to, amino, methylamino, acetylamino, and acetylmethylamino.

10 The term " $(\text{NR}^{\text{A}}\text{R}^{\text{B}})\text{carbonyl}$ " as used herein, means a $\text{NR}^{\text{A}}\text{R}^{\text{B}}$ group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of $(\text{NR}^{\text{A}}\text{R}^{\text{B}})\text{carbonyl}$ include, but are not limited to, aminocarbonyl, (methylamino)carbonyl, (dimethylamino)carbonyl, and (ethylmethylamino)carbonyl.

The term " $(\text{NR}^{\text{A}}\text{R}^{\text{B}})\text{carbonyloxy}$ " as used herein, means a $(\text{NR}^{\text{A}}\text{R}^{\text{B}})\text{carbonyl}$ group as defined herein appended to the parent molecular moiety through an oxygen.

15 The term " $\text{NR}_{\text{C}}\text{R}_{\text{D}}$ " as used herein, means two groups, R_{C} and R_{D} , which are appended to the parent molecular moiety through a nitrogen atom. R_{C} and R_{D} are each independently hydrogen, alkyl, and alkylcarbonyl. Representative examples of $\text{NR}_{\text{C}}\text{R}_{\text{D}}$ include, but are not limited to, amino, methylamino, acetylamino, and acetylmethylamino.

20 The term " $(\text{NR}_{\text{C}}\text{R}_{\text{D}})\text{carbonyl}$ " as used herein, means a $\text{NR}_{\text{C}}\text{R}_{\text{D}}$ group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of $(\text{NR}_{\text{C}}\text{R}_{\text{D}})\text{carbonyl}$ include, but are not limited to, aminocarbonyl, (methylamino)carbonyl, (dimethylamino)carbonyl, and (ethylmethylamino)carbonyl.

25 The term " $(\text{NR}_{\text{C}}\text{R}_{\text{D}})\text{carbonylalkyl}$ " as used herein, means a $(\text{NR}_{\text{C}}\text{R}_{\text{D}})\text{carbonyl}$ group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

The term " $(\text{NR}_{\text{C}}\text{R}_{\text{D}})\text{sulfonyl}$ " as used herein, means a $\text{NR}_{\text{C}}\text{R}_{\text{D}}$ group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of $(\text{NR}_{\text{C}}\text{R}_{\text{D}})\text{sulfonyl}$ include, but are not limited to, aminosulfonyl, (methylamino)sulfonyl, (dimethylamino)sulfonyl, and (ethylmethylamino)sulfonyl.

The term "NR^AR^B" as used herein, means two groups, R^A and R^B, which are appended to the parent molecular moiety through a nitrogen atom. R^A and R^B are each independently hydrogen, alkyl, cycloalkyl, and alkylcarbonyl. Representative examples of NR^AR^B include, but are not limited to, amino, methylamino, acetylamino, and acetylmethylamino.

5 The term "(NR^AR^B)carbonyl" as used herein, means a NR^AR^B group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of (NR^AR^B)carbonyl include, but are not limited to, aminocarbonyl, (methylamino)carbonyl, (dimethylamino)carbonyl, and (ethylmethylamino)carbonyl.

The term "oxo" as used herein, means a =O moiety.

10 The term "sulfinyl" as used herein, means a -SO- group.

The term "sulfonyl" as used herein, means a -SO₂- group.

Compounds of the present invention can exist as stereoisomers, wherein asymmetric or chiral centers are present. Stereoisomers are designated (R) or (S) depending on the configuration of substituents around the chiral carbon atom. The terms (R) and (S) used herein
15 are configurations as defined in IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem., (1976), 45: 13-30. The present invention contemplates various stereoisomers and mixtures thereof and are specifically included within the scope of this invention. Stereoisomers include enantiomers, diastereomers, and mixtures of enantiomers or diastereomers. Individual stereoisomers of compounds of the
20 present invention may be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the
25 optically pure product from the auxiliary or (2) direct separation of the mixture of optical enantiomers on chiral chromatographic columns.

Compounds of the present invention were named by ACD/ChemSketch version 5.06 (developed by Advanced Chemistry Development, Inc., Toronto, ON, Canada) or were given names which appeared to be consistent with ACD nomenclature.

Determination of Biological ActivityInhibition of PARP

5 Nicotinamide[2,5',8-3H]adenine dinucleotide and strepavidin SPA beads were purchased from Amersham Biosciences (UK) Recombinant Human Poly(ADP-Ribose) Polymerase (PARP) purified from E.coli and 6-Biotin-17-NAD⁺, were purchase from Trevigen, Gaithersburg, MD. NAD⁺, Histone, aminobenzamide, 3-amino benzamide and Calf Thymus DNA (dcDNA) were purchased from Sigma, St. Louis, MO. Stem loop oligonucleotide containing MCAT sequence
 10 was obtained from Qiagen. The oligos were dissolved to 1mM in annealing buffer containing 10mM Tris HCl pH 7.5, 1mM EDTA, and 50mM NaCl, incubated for 5min at 95°C, and followed by annealing at 45°C for 45 minutes. Histone H1 (95% electrophoretically pure) was purchased from Roche, Indianapolis, IN. Biotinylated histone H1 was prepared by treating the protein with Sulfo-NHS-LC-Biotin from Pierce Rockford, IL. The biotinylation reaction was
 15 conducted by slowly and intermittently adding 3 equivalents of 10mM Sulfo-NHS-LC-Biotin to 100μM Histone H1 in phosphate-buffered saline, pH 7.5, at 4°C with gentle vortexing over 1min followed by subsequent 4°C incubation for 1hr. Streptavidin coated (FlashPlate Plus) microplates were purchased from Perkin Elmer, Boston, MA.

PARP1 assay was conducted in PARP assay buffer containing 50 mM Tris pH 8.0, 1mM
 20 DTT, 4 mM MgCl₂. PARP reactions contained 1.5 μM [³H]-NAD⁺ (1.6uCi/mmol), 200 nM biotinylated histone H1, 200 nM slDNA, and 1nM PARP enzyme. Auto reactions utilizing SPA bead-based detection were carried out in 100μl volumes in white 96 well plates. Reactions were initiated by adding 50 μl of 2X NAD⁺ substrate mixture to 50μl of 2X enzyme mixture containing PARP and DNA. These reactions were terminated by the addition of 150 μl of 1.5
 25 mM benzamide (~1000-fold over its IC₅₀). 170 μl of the stopped reaction mixtures were transferred to streptavidin Flash Plates, incubated for 1hr, and counted using a TopCount microplate scintillation counter. The K_i data was determined from inhibition curves at various substrate concentrations and are shown in Table 1 for compounds of the present invention

30

Table 1

IC₅₀ Values for PARP Inhibitors of the Invention (nM)

46	14.2	7.1	158	4.5	21.5	6.3	13	450	81
11	62	59	19.6	1.6	11.6	22.5	43	29	1.7
9	29	16.8	64	3.5	3.1	5.7	13.3	800	180
52	10	34	55	2.6	472	4.1	6.3	3	10
11	2.8	1.2	69	3.3	20.7	2.4	15.3	>8300	3.6
24.9	6	91	3.1	4.6	47	6.5	15.7	16.2	4.5
6.1	1.1	1.6	2.9	3.6	6.1	23	3.7	18.6	5
3.5	116	4.3	8.2	4.9	5.5	5.5	.7	56	395
2.7	800	9.6	16.4	5.3	9.5	8.5	31	226	20.2
10.3	866	5.8	11.4	3.5	4.5	17	45	35	12.2
5.3	2.4	5.4	3700	3.2	5.6	13	44	11.9	2.8
9.8	11.3	2.1	6.5	4.6	34	33	52	1.1	2.0
70	4.3	3.4	18	6.5	17	211	4	20	5.6
231	2.9	57	37	20	19	256	6	10	2
109	1.8	35	2	32	26	315	7	23	2.5
456	4	52	4	42	242	24	6	37	3.7
590	2.7	90	4	5	14	15	8	31	3
640	2	135	5	122	14	2.3	5	41	7.2
2.7	4.7	129	10	65	18	2.9	6	27	16

8	54	131	251	1	53	134	86	69	167
20	49	228	233	23	135	65	181	25	183
18	59	3	12	30	27	217	26	65	239
5	49	.4	3	5	40	245	17	88	9
3	39	14	25	12	60	72	61	27	155
307	163								

Cellular PARP assay:

C41 cells were treated with a compound of the present invention for 30 minutes in 96 well plate. PARP was then activated by damaging DNA with 1 mM H₂O₂ for 10 minutes. The cells were then washed with ice-cold PBS once and fixed with pre-chilled methanol:acetone (7:3) at -20°C for 10 minutes. After air-drying, the plates were rehydrated with PBS and blocked 5% non-fat dry milk in PBS-tween (0.05%) (blocking solution) for 30 minutes at room temperature. The cells were incubated with anti-PAR antibody 10H (1:50) in Blocking solution at 37°C for 60 minutes followed by washing with PBS-Tween20 5 times, and incubation with goat anti-mouse fluorescein 5(6)-isothiocyanate-coupled antibody (1:50) and 1 µg/ml 4',6-diamidino-2-phenylindole (DAPI) in blocking solution at 37°C for 60 minutes. After washing with PBS-Tween20 5 times, the analysis was performed using an fmax Fluorescence Microplate Reader (Molecular Devices, Sunnyvale, CA), set at the excitation wavelength of 490 nm and emission wavelength of 528 nm fluorescein 5(6)-isothiocyanate (FITC) or the excitation wavelength of 355 nm and emission wavelength of 460 nm (DAPI). The PARP activity (FITC signal) was normalized with cell numbers (DAPI).

The cellular assay measures the formation of poly ADP-ribose by PARP within cells and demonstrates that compounds of the present invention penetrate cell membranes and inhibit PARP in intact cells. The EC_{50s} for representative compounds of the present invention are provided in Table 2.

Table 2

EC₅₀ Values for PARP Inhibitors of the Invention (nM)

2.3	1.5	45	1	18	2.6	7	0.3	11.9	35
0.9	46	1	14	115	<0.3	>1000	0.5	12.1	4.8
11	>1000	1.3	1.2	0.6	17.5	463	0.2	3.7	4.5
0.3	1.2	.6	2.3	7	.5	1.9	.6	1.2	.5
1.4	.8	.6	1	1.7	.3	.9	.5	1.6	.4
.3	.3	.7	17	2.1	.7	2.7	.5	1.9	>300
.7	1.8	3.2	.4	2.3	.5	3.6	.5	5.7	7.6

.5	.8	3.2	.7	40	1.1	1.5	.6	1	.5
.5	.3	5.6	.3	4.1	3.4	1.8	.5	4	.5
.6	.6	2.4	.7	3.5	1.7	.9	.6	3.3	1.5
>300	>300	.7	6.2	>300	10.6	8.7	>300	1.5	3.1
.5	2.4	1.8	14	>300	>1000	14	>300	63	7.9
.8	5.6	4.5	15	16	259	112	52	8.2	8.9
>300	57	>300	>300	191	>300	61			

As PARP inhibitors, the compounds of the present invention have numerous therapeutic applications related to, ischemia reperfusion injury, inflammatory diseases, degenerative diseases, protection from adverse effects of cytotoxic compounds, and potentiation of cytotoxic cancer therapy. In particular, compounds of the present invention potentiate radiation and chemotherapy by increasing cell death of cancer cells, limiting tumor growth, decreasing metastasis, and prolonging the survival of tumor-bearing mammals. Compounds of Formula (I) can treat leukemia, colon cancer, glioblastomas, lymphomas, melanomas, carcinomas of the breast, and cervical carcinomas.

Other therapeutic applications include, but are not limited to, retroviral infection, arthritis, gout, inflammatory bowel disease, CNS inflammation, multiple sclerosis, allergic encephalitis, sepsis, septic shock, hemorrhagic shock, pulmonary fibrosis, uveitis, diabetes, Parkinsons disease, myocardial infarction, stroke, other neural trauma, organ transplantation, reperfusion of the eye, reperfusion of the kidney, reperfusion of the gut, reperfusion of skeletal muscle, liver toxicity following acetaminophen overdose, cardiac and kidney toxicities from doxorubicin and platinum based antineoplastic agents, and skin damage secondary to sulfur mustards. (G. Chen et al. Cancer Chemo. Pharmacol. 22 (1988), 303; C. Thiernemann et al., Proc. Natl. Acad. Sci. USA 94 (1997), 679-683 D. Weltin et al. Int. J. Immunopharmacol. 17 (1995), 265- 271; H. Kröger et al. Inflammation 20 (1996), 203-215; W. Ehrlich et al. Rheumatol. Int. 15 (1995), 171-172; C. Szabo et al., Proc. Natl. Acad. Sci. USA 95 (1998), 3867-3872; S. Cuzzocrea et al. Eur. J. Pharmacol. 342 (1998), 67-76; V. Burkhart et al., Nature Medicine (1999), 5314-19).

When used in the above or other treatments, a therapeutically effective amount of one of the compounds of the present invention can be employed as a zwitterion or as a pharmaceutically acceptable salt. By a "therapeutically effective amount" of the compound of the invention is meant a sufficient amount of the compound to treat or prevent a disease or disorder ameliorated by a PARP inhibitor at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

By "pharmaceutically acceptable salt" is meant those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well-known in the art. The salts can be prepared in situ during the final isolation and purification of the compounds of the present invention or separately by reacting the free base of a compound of the present invention with a suitable acid. Representative acids include, but are not limited to acetic, citric, aspartic, benzoic, benzenesulfonic, butyric, fumaric, hydrochloric, hydrobromic, hydroiodic, lactic, maleic, methanesulfonic, pantoic, pectinic, pivalic, propionic, succinic, tartaric, phosphic, glutamic, and p-toluenesulfonic. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides;

arylalkyl halides like benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

A compound of the present invention may be administered as a pharmaceutical composition containing a compound of the present invention in combination with one or more pharmaceutically acceptable excipients. A pharmaceutically acceptable carrier or excipient refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The compositions can be administered parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), rectally, or buccally. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions for parenteral injection comprise pharmaceutically-acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions can also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

Compounds of the present invention may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are


dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 *et seq.*

Total daily dose of the compositions of the invention to be administered to a human or other mammal host in single or divided doses may be in amounts, for example, from 0.0001 to 300 mg/kg body weight daily and more usually 1 to 300 mg/kg body weight. The dose, from 0.0001 to 300 mg/kg body, may be given twice a day.

Abbreviations which have been used in the descriptions of the examples that follow are: DBU for 1,8-diazabicyclo[5.4.0]undec-7-ene; DMF for N,N-dimethylformamide; DMSO for dimethylsulfoxide; Et₂O for diethyl ether; EtOAc for ethyl acetate; EtOH for ethanol; HPLC for high pressure liquid chromatography; LDA for lithium diisopropylamide; MeOH for methanol; psi for pounds per square inch; TFA for trifluoroacetic acid; THF for tetrahydrofuran, and TMS for trimethylsilane.

Compounds having formula I may be made by synthetic chemical processes, examples of which are shown herein below. It is meant to be understood that the order of the steps in the processes may be varied, that reagents, solvents and reaction conditions may be substituted for those specifically mentioned, and that vulnerable moieties may be protected and deprotected, as necessary.

Scheme 1




Compounds of formula (4) wherein X and R³ are as defined in formula (I) and ring A is cycloalkyl or heterocycle, unsubstituted or substituted as defined in formula (I), can be prepared from amines of formula (1) as shown in Scheme 1.

Amines of formula (1) can be reacted with unsubstituted or substituted cyclic ketones of formula (2) in the presence of acid such as, but not limited to, acetic acid, to provide alkenes of formula (3). The reaction is generally conducted at a temperature from about room temperature to about 70 °C.


5 Compounds of formula (3) when treated with isocyanates such as, but not limited to, ethyl isocyanates, in pyridine or 4-methylpyridine, at reflux, provides compounds of formula (4).

Scheme 2



10 Aminopyrazoles (7) wherein R^3 and R^4 are as defined in formula (I) can be prepared from the reaction of hydrazines of formula (5) with an α -cyanoketone of formula (6), in a solvent such as ethanol and the like, at the reflux temperature of the solvent employed.

Scheme 3




15 Compounds of formula (8) wherein X and R^3 are as defined in formula (I), Y is $NR^A R^B$, alkoxy or haloalkoxy, and ring A is cycloalkyl or heterocycle, unsubstituted or substituted as defined in formula (I), and R^A and R^B are as defined in formula (I), can be prepared by (a) allylic halogenation, such as chlorination or bromination of ring A, and (b) displacement of the halide with an appropriate nucleophile of formula Y-H.

20 Allylic halogenation of compounds of formula (4) can be accomplished by treatment of (4) with a halogenating agent such as, but not limited to, bromosuccinimide or chlorosuccinimide, in the presence of catalytic amount of radical initiator such as, but not limited to, 2,2'-azobis(2-methylpropionitrile) or m-chloroperbenzoic acid (MCPBA), at elevated (for example 40 – 70 °C) or ambient temperature.

Displacement of the halide with a nucleophile can be achieved in the presence of a base such as, but not limited to, triethylamine or diisopropylethylamine, at a temperature from about room temperature to about 100 °C.

Scheme 4




Compounds of formula (11) wherein X is as defined in formula (I), Q is alkylene, ring A is cycloalkyl or heterocycle, unsubstituted or substituted as defined in formula (I), ring B is aryl, heteroaryl, heterocycle or cycloalkyl, each of which is independently unsubstituted or substituted as described in formula (I), and R¹⁰¹ and R¹⁰² are independently selected from the group consisting of alkyl, cycloalkyl and heterocycle, can be prepared from primary amines of formula (9) as shown in Scheme 4.

Primary amines of formula (9) can be monoalkylated by compounds of formula R¹⁰¹-Z, wherein Z is halides, triflates or mesylates, in the presence of a base such as, but not limited to, triethylamine or diisopropylethylamine, at a temperature from about room temperature to about 100 °C. The second alkylation can be achieved with compounds of formula R¹⁰²-Z using similar reaction conditions to afford compounds of formula (11). The alkylation can be conducted sequentially or in one pot.

Alternatively, compounds of formula (10) can be obtained from primary amines of formula (9) via reductive amination conditions wherein (9) is treated with a ketones or aldehydes, in the presence of an acid such as, but not limited to, acetic acid, and a reducing agent such as, but not limited to, sodium cyanoborohydride. Compounds of formula (11) wherein R^A and R^B are independently alkyl, cycloalkyl or heterocycle, can be prepared from (10) using similar reaction conditions.

Scheme 5



Compounds of formula (14) wherein X is as defined in formula (I), Q is alkylene, ring A is cycloalkyl or heterocycle, unsubstituted or substituted as defined in formula (I), and R^{101} and R^{102} are independently alkyl, cycloalkyl or heterocycle, can be prepared from primary amines of formula (12) as shown in Scheme 5, using reaction conditions as described in Scheme 4.

The following Examples are intended as an illustration of and not a limitation upon the scope of the invention as defined in the appended claims. The compounds of this invention can be prepared by a variety of synthetic routes.

EXAMPLE 1

1,3-dimethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as described in Winters, G; Sala, A; De Paoli, A.; Ferri, V., Easy Synthesis of New Ring-Fused Pyridones from Heteroaromatic β -Vinylamines, *Synthesis* 1984, 1052-1054.

EXAMPLE 2

9-cyclobutylamino-1,3-dimethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 2A

9-bromo-1,3-dimethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

A solution of EXAMPLE 1 (966 mg, 4.45 mmol) in a mixture of chloroform (30 mL) and N,N'-dimethylformamide (2 mL) was treated with N-bromosuccinimide (791 mg, 4.45 mmol) and catalytic 2,2'-azobis(2-methylpropionitrile) (5 mg) at 65°C for 18 hours. The mixture was cooled, concentrated and the residue partitioned between ethyl acetate and brine. The organic phase was washed with water, concentrated and purified by flash chromatography on silica gel

eluting with 50% ethyl acetate in hexanes to provide 1.14 g (87%) of the title compound. MS (DCI/NH₃) m/z 297 (M+H)⁺.

EXAMPLE 2B

5 9-cyclobutylamino-1,3-dimethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one
Step A

To a solution of EXAMPLE 2A (40 mg, 0.14 mmol) in 1:1 tetrahydrofuran/N,N'-dimethylformamide (4 mL) was added cyclobutylamine (20 mg, 0.28 mmol) and diisopropylethylamine (49 μ L, 0.28 mmol) and the mixture heated at 50°C for 18 hours. The
10 mixture was cooled, concentrated, and the residue purified by flash chromatography on silica gel using 70% ethyl acetate in hexanes to give the title compound.

Step B

Purification of the free base from Step A by HPLC (Zorbax, C-18, 250x2.54 column, Mobile phase A: 0.1% trifluoroacetic acid in water; B: 0.1% trifluoroacetic acid in acetonitrile;
15 0-100% gradient) provided 12 mg (30%) of the title compound as a trifluoroacetate salt. ¹H NMR (CD₃OD) δ 1.85-1.97 (m, 3H), 1.98-2.12 (m, 2H), 2.18-2.39 (m, 4H), 2.40-2.53 (m, 2H), 2.56 (s, 3H), 2.76-2.90 (m, 1H), 3.84 (s, 3H), 4.04-4.19 (m, 1H), 4.79 (d, J=3.0 Hz, 1H), 13.70 (br s, 1H).

20

EXAMPLE 3

1,3-dimethyl-9-morpholin-4-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in Step A of EXAMPLE 2B, substituting morpholine for cyclobutylamine (21% yield). ¹H NMR (CDCl₃)
 δ 1.60 (s, 1H), 1.65-1.79 (m, 2H), 1.84-1.96 (m, 1H), 2.04-2.16 (m, 1H), 2.50-2.57 (m, 2H),
25 2.57-2.64 (m, 6H), 3.54-3.62 (m, 2H), 3.62-3.71 (m, 2H), 3.99 (s, 3H), 4.08 (t, J=5.7 Hz, 1H), 13.70 (br s, 1H).

EXAMPLE 4

1,3-dimethyl-9-piperidin-1-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in Step A of EXAMPLE 2B, substituting piperidine for cyclobutylamine (72% yield). ¹H NMR (DMSO-*d*₆) δ 1.26-1.39 (m, 4H), 1.42-1.50 (m, 2H), 1.54-1.65 (m, 2H), 1.71-1.82 (m, 1H), 1.95-2.04 (m, 1H), 2.32-2.39 (m, 2H), 2.39-2.44 (m, 2H), 2.47 (s, 3H), 3.22-3.28 (m, 2H), 3.73 (s, 3H), 3.99-4.07 (m, 1H), 13.76 (br s, 1H).

EXAMPLE 5

1,3-dimethyl-9-(4-methylpiperazin-1-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared according to the procedure as described in Step A of EXAMPLE 2B, substituting 1-methylpiperazine for cyclobutylamine (37% yield). ¹H NMR (C₅D₅N) δ 1.52-1.66 (m, 2H), 1.74-1.83 (m, 1H), 1.96-2.05 (m, 1H), 2.27-2.36 (m, 2H), 2.74 (s, 3H), 2.74 (s, 3H), 2.76-2.79 (m, 1H), 3.01-3.12 (m, 3H), 3.13-3.24 (m, 2H), 3.29-3.36 (m, 1H), 3.60-3.68 (m, 1H), 3.98 (s, 3H), 4.22 (t, *J*=5.7 Hz, 1H), 14.18 (br s, 1H).

EXAMPLE 6

1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one

A mixture of 8.5 g of pyridine and 10 mL of concentrated hydrochloric acid was heated to 230°C, added to EXAMPLE 1 (217 mg, 1.0 mmol) and the mixture heated at 230°C for 2 hours. After cooling, water was added and the suspension neutralized with 10% sodium hydroxide. Dichloromethane (50 mL) was added, the mixture was filtered and the solid collected, washed with dichloromethane and water and dried. Purification by HPLC (Zorbax, C-18, 250x2.54 column, Mobile phase A: 0.1% trifluoroacetic acid in water; B: 0.1% trifluoroacetic acid in acetonitrile; 0-100% gradient) provided 173 mg (73%) of the title compound as the trifluoroacetate salt. ¹H NMR (DMSO-*d*₆) δ 1.63-1.73 (m, 4H), 2.29-2.34 (m, 2H), 2.46 (s, 3H), 2.75-2.80 (m, 2H), 11.28 (br s, 1H).

EXAMPLE 7

1-methyl-3-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one

EXAMPLE 7A

4-cyclohex-1-enyl-5-methyl-2-phenyl-2H-pyrazol-3-ylamine

A solution of 5-amino-3-methyl-1-phenylpyrazole (8.66 g, 50 mmol) in acetic acid (70 mL) was treated with cyclohexanone (10.6 mL, 100 mmol) at ambient temperature for 40 hours. The volatiles were removed and the residue diluted with water and 10% sodium hydroxide added until the pH = 12. The mixture was stirred with 10 mL of hexane for 30 minutes and the solid collected by filtration, washed with water and hexane and dried to give 9.43 g (74%) of the title compound. MS (DCI): m/z 254 (M+H)⁺.

EXAMPLE 7B

1-methyl-3-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

To a solution of EXAMPLE 7A (1.26 g, 5 mmol) in pyridine (10 mL) was added ethyl isocyanate (1.18 mL, 15 mmol). The solution was heated at reflux for 7 hours, cooled, concentrated and the residue triturated with methanol. The solid was collected by filtration, washed with methanol and dried to give 493 mg (35%) of the title compound. ¹H NMR (DMSO-d₆) δ 1.72-1.83 (m, 4H), 2.49-2.55 (m, 2H), 2.57 (s, 3H), 3.05 (t, J=5.2 Hz, 2H), 7.24 (t, J=7.4 Hz, 1H), 7.43-7.53 (m, 2H), 8.19 (d, J=7.7 Hz, 2H), 11.27 (br s, 1H).

EXAMPLE 8

3-(2-dimethylaminoethyl)-1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 8A

4-cyclohex-1-enyl-2-(2-dimethylaminoethyl)-5-methyl-2H-pyrazol-3-ylamine

A solution of 2-(2-dimethylaminoethyl)-5-methyl-2H-pyrazol-3-ylamine (0.998 g, 5.93 mmol) in acetic acid (10 mL) was treated with cyclohexanone (1.23 mL, 11.86 mmol) at ambient temperature for two days. The volatiles were removed and the residue partitioned between dilute sodium hydroxide and ethyl acetate. The organic phase was washed with water, concentrated and the residue purified by flash chromatography on silica gel using 10% methanol in dichloromethane to give a solid (1.10 g, 75%). MS (DCI): m/z 249 (M+H)⁺.

EXAMPLE 8B

3-(2-dimethylaminoethyl)-1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

To a solution of EXAMPLE 8A (1.05 g, 4.22 mmol) in pyridine (10 mL) was added ethyl isocyanate (1.0 mL, 12.68 mmol) and the solution heated at reflux overnight. The mixture was cooled, stirred with 15 mL ethyl acetate at ambient temperature for 5 minutes and the solid collected by filtration, washed with ethyl acetate and dried to give 793 mg of the title compound. The mother liquor was concentrated and the residue recrystallized from ethyl acetate to give an additional 300 mg of the title compound (94% overall yield). ¹H NMR (DMSO-d₆) δ 1.64-1.75 (m, 4H), 2.16 (s, 6H), 2.34-2.39 (m, 5H), 2.59 (t, *J*=6.4 Hz, 2H), 2.83-2.87 (m, 2H), 4.18 (t, *J*=6.6 Hz, 2H).

10

EXAMPLE 9

3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure for EXAMPLE 8, substituting 1-methyl-1H-pyrazol-5-ylamine for 2-(2-dimethylaminoethyl)-5-methyl-2H-pyrazol-3-ylamine. ¹H NMR (DMSO-d₆): δ 1.67-1.77 (m, 4H), 2.38 (br d, *J*=1.8 Hz, 2H), 2.72 (br d, *J*=1.8 Hz, 2 H), 3.82 (s, 3H), 7.68 (s, 1H), 11.90 (br s, 1H).

15

EXAMPLE 10

1-methyl-6,7,8,9-tetrahydro-4H-isoxazolo[5,4-c]isoquinolin-5-one

20

EXAMPLE 10A

4-Cyclohex-1-enyl-3-methyl-isoxazol-5-ylamine

The title compound was prepared according to the procedure for EXAMPLE 7A, substituting 5-amino-3-methylisoxazole for 5-amino-3-methyl-1-phenylpyrazole (72% yield).

MS (DCI/NH₃) *m/z* 179 (M+H)⁺.

25

EXAMPLE 10B

1-methyl-6,7,8,9-tetrahydro-4H-isoxazolo[5,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 10A for EXAMPLE 7A (67% yield). ¹H NMR (DMSO-*d*₆) δ 1.69-1.79 (m, 4H), 2.46 (t, *J*=4.6 Hz, 2H), 2.49 (s, 3H), 2.95 (t, *J*=4.8 Hz, 2H), 12.16 (br s, 1H).

5

EXAMPLE 11

1-phenyl-6,7,8,9-tetrahydro-4H-isoxazolo[5,4-*c*]isoquinolin-5-one

EXAMPLE 11A

10

4-cyclohex-1-enyl-3-phenylisoxazol-5-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 5-amino-3-phenylisoxazole for 5-amino-3-methyl-1-phenylpyrazole (50% yield). MS (DCI/NH₃) *m/z* 241 (M+H)⁺.

15

EXAMPLE 11B

1-phenyl-6,7,8,9-tetrahydro-4H-isoxazolo[5,4-*c*]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 11A for EXAMPLE 7A (3% yield). ¹H NMR (C₅D₅N) δ 1.43-1.52 (m, 2H), 1.65-1.75 (m, 2H), 2.55 (t, *J*=6.3 Hz, 2H), 2.81 (t, *J*=6.4 Hz, 2H), 7.50-7.57 (m, 3H), 7.80-7.87 (m, 2H), 14.15 (br s, 1H).

20

EXAMPLE 12

3-methyl-1-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one

25

EXAMPLE 12A

4-cyclohex-1-enyl-2-methyl-5-phenyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 5-amino-1-methyl-3-phenylpyrazole for 5-amino-3-methyl-1-phenylpyrazole (36% yield). MS (DCI/NH₃) *m/z* 254 (M+H)⁺.

30

EXAMPLE 12B

3-methyl-1-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 12A for EXAMPLE 7A (66% yield). ¹H NMR (C₅D₅N) δ 1.44-1.56 (m, 2H), 1.66-1.76 (m, 2H), 2.68 (t, *J*=5.7 Hz, 2H), 2.87 (t, *J*=5.8 Hz, 2H), 4.06 (s, 3H), 7.43-7.49 (m, 1H), 7.51-7.56 (m, 2H), 7.56-7.58 (m, 1H), 7.89 (t, *J*=1.7 Hz, 1H), 14.15 (br s, 1H).

EXAMPLE 13

3-benzyl-1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 13A

2-benzyl-4-cyclohex-1-enyl-5-methyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 2-benzyl-5-methyl-2H-pyrazol-3-ylamine for 5-amino-3-methyl-1-phenylpyrazole (45% yield). MS (DCI/NH₃) *m/z* 268 (M+H)⁺.

EXAMPLE 13B

3-benzyl-1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 13A for EXAMPLE 7A (55% yield). ¹H NMR (C₅D₅N) δ 1.64-1.78 (m, 4H), 2.57 (s, 3H), 2.80-2.91 (m, 4H), 5.67 (s, 2H), 7.21-7.24 (m, 1H), 7.26-7.32 (m, 2H), 7.44-7.49 (m, 2H), 14.15 (br s, 1H).

EXAMPLE 14

3-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 14A

4-cyclohex-1-enyl-2-phenyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure for EXAMPLE 7A, substituting 5-amino-1-phenylpyrazole for 5-amino-3-methyl-1-phenylpyrazole (18% yield). MS (DCI/NH₃) m/z 240 (M+H)⁺.

5

EXAMPLE 14B

3-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 14A for EXAMPLE 7A (23% yield). ¹H NMR (C₅D₅N) δ 1.66-1.80 (m, 4H), 2.79 (t, J=6.1 Hz, 2H), 2.85 (t, J=6.1 Hz, 2H), 7.25-7.30 (m, 1H), 7.41-7.50 (m, 10 2H), 8.19 (s, 1H), 8.54 (dd, J=8.6, 1.2 Hz, 2H), 14.15 (br s, 1H).

EXAMPLE 15

3-methyl-1-thiophen-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

15

EXAMPLE 15A

4-cyclohex-1-enyl-2-methyl-5-thiophen-2-yl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 5-amino-1-methyl-3-(2-thienyl)pyrazole for 5-amino-3-methyl-1-phenylpyrazole (30% yield). MS (DCI/NH₃) m/z 260 (M+H)⁺.

20

EXAMPLE 15B

3-methyl-1-thiophen-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 15A for EXAMPLE 7A (73% yield). ¹H NMR (C₅D₅N) δ 1.54-1.60 (m, 2H), 1.69-1.77 (m, 2H), 2.87 (t, J=6.4 Hz, 4H), 4.02 (s, 3H), 7.24 (dd, J=5.2, 3.7 Hz, 1H), 7.54 (dd, J=3.4, 0.9 Hz, 1H), 7.57 (d, J=0.9 Hz, 1H), 14.15 (br s, 1H).

30

EXAMPLE 16

1-isopropyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 16A

4-cyclohex-1-enyl-5-isopropyl-2-methyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 3-isopropyl-1-methyl-1H-pyrazol-5-amine for 5-amino-3-methyl-1-phenylpyrazole (13% yield). MS (DCI/NH₃) m/z 220 (M+H)⁺.

EXAMPLE 16B

1-isopropyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 16A for EXAMPLE 7A (44% yield). ¹H NMR (C₅D₅N) δ 1.51 (d, J=6.7 Hz, 6H), 1.68-1.81 (m, 4H), 2.87 (m, 2H), 2.93 (m, 2H), 3.32-3.47 (m, 1H), 3.95 (s, 3H), 13.45 (br, s 1H).

EXAMPLE 17

1-cyclopropyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 17A

4-cyclohex-1-enyl-5-cyclopropyl-2-methyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 3-cyclopropyl-1-methyl-1H-pyrazol-5-amine for 5-amino-3-methyl-1-phenylpyrazole (50% yield). MS (DCI/NH₃) m/z 218 (M+H)⁺.

EXAMPLE 17B

1-cyclopropyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 17A for EXAMPLE 7A (83% yield). ¹H NMR (C₅D₅N) δ 0.92-1.01 (m, 2H), 1.17-1.28 (m, 2H), 1.69-1.82 (m, 4H), 2.16-2.27 (m, 1H), 2.86 (t, J=4.4 Hz, 2H), 3.06 (t, J=4.4 Hz, 2H), 3.91 (s, 3H), 13.46 (br s, 1H).

30

EXAMPLE 19

1-cyclopropyl-3-methyl-4,6,8,9-tetrahydro-3H-7-oxa-2,3,4-triaza-cyclopenta[a]naphthalen-5-one

EXAMPLE 19A

5-cyclopropyl-4-(3,6-dihydro-2H-pyran-4-yl)-2-methyl-2H-pyrazol-3-ylamine

5 The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 3-cyclopropyl-1-methyl-1H-pyrazol-5-amine for 5-amino-3-methyl-1-phenylpyrazole and substituting tetrahydro-4H-pyran-4-one for cyclohexanone (67% yield). MS (DCI/NH₃) m/z 220 (M+H)⁺.

10

EXAMPLE 19B

1-cyclopropyl-3-methyl-4,6,8,9-tetrahydro-3H-7-oxa-2,3,4-triaza-cyclopenta[a]naphthalen-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 19A for EXAMPLE 7A (20% yield). ¹H NMR (DMSO-*d*₆) δ 0.77-0.83 (m, 2H), 0.83-0.90 (m, 2H), 2.04-2.18 (m, 1H), 2.96-3.11 (m, 2H), 3.72 (s, 3H), 3.87 (t, 15 *J*=5.6 Hz, 2H), 4.41 (s, 2H), 11.99 (br s, 1H)

EXAMPLE 20

1-(3-chlorophenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one

20

EXAMPLE 20A

5-(3-chlorophenyl)-4-cyclohex-1-enyl-2-methyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 5-(3-chlorophenyl)-2-methyl-2H-pyrazole-3-ylamine for 5-amino-3-methyl-1-phenylpyrazole (14% yield). MS (DCI/NH₃) m/z 288 (M+H)⁺.

25

EXAMPLE 20B

1-(3-chlorophenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 20A for EXAMPLE 7A (3% yield). ¹H NMR (DMSO-*d*₆) δ 1.52-

1.62 (m, 2H), 1.64-1.74 (m, 2H), 2.37-2.48 (m, 4H), 3.88 (s, 3H), 7.47-7.51 (m, 3H), 7.53-7.58 (m, 1H), 12.04 (br s, 1H).

EXAMPLE 21

5 3-methyl-1-thiophen-2-yl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one

EXAMPLE 21A

4-cyclopent-1-enyl-2-methyl-5-thiophen-2-yl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE
10 7A, substituting 5-amino-1-methyl-3-(2-thienyl)pyrazole for 5-amino-3-methyl-1-phenylpyrazole and substituting cyclopentanone for cyclohexanone (20% yield). MS (DCI/NH₃) m/z 246 (M+H)⁺.

EXAMPLE 21B

15 3-methyl-1-thiophen-2-yl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one

The title compound was prepared according to the procedure as described in EXAMPLE
7B, substituting EXAMPLE 21A for EXAMPLE 7A (70% yield). ¹H NMR (DMSO-*d*₆) δ 2.07 (dt, *J*=15.0, 7.5 Hz, 2H), 2.71 (t, *J*=7.4 Hz, 2H), 3.16 (t, *J*=7.4 Hz, 2H), 3.89 (s, 3H), 7.15 (dd, *J*=4.6, 4.0 Hz, 1H), 7.45 (d, *J*=3.7 Hz, 1H), 7.56 (d, *J*=5.2 Hz, 1H), 11.94 (br s, 1H).

20

EXAMPLE 22

1-cyclopropyl-3-methyl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one

EXAMPLE 22A

25 4-cyclopent-1-enyl-5-cyclopropyl-2-methyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE
7A, substituting 3-cyclopropyl-1-methyl-1H-pyrazol-5-amine for 5-amino-3-methyl-1-phenylpyrazole and substituting cyclopentanone for cyclohexanone (30% yield). MS (DCI/NH₃) m/z 204 (M+H)⁺.

30

EXAMPLE 22B

1-cyclopropyl-3-methyl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 22A for EXAMPLE 7A (60% yield). ¹H NMR (DMSO-*d*₆) δ 0.74-0.83 (m, 2H), 0.88 (dd, *J*=8.2, 2.4 Hz, 2H), 1.97-2.12 (m, 3H), 2.65 (t, *J*=6.9 Hz, 2H), 3.12 (t, *J*=7.3 Hz, 2H), 3.72 (s, 3H), 11.78 (br s, 1H).

EXAMPLE 23

1-cyclopropyl-3-methyl-4,6,7,8,9,10-hexahydro-3H-2,3,4-triazacyclohepta[e]inden-5-one

EXAMPLE 23A

4-Cyclohept-1-enyl-5-cyclopropyl-2-methyl-2H-pyrazol-3-ylamine

The title compound was prepared according to the procedure as described in EXAMPLE 7A, substituting 3-cyclopropyl-1-methyl-1H-pyrazol-5-amine for 5-amino-3-methyl-1-phenylpyrazole and substituting cycloheptanone for cyclohexanone (6% yield). MS (DCI/NH₃) *m/z* 232 (M+H)⁺.

EXAMPLE 23B

1-cyclopropyl-3-methyl-4,6,7,8,9,10-hexahydro-3H-2,3,4-triazacyclohepta[e]inden-5-one

The title compound was prepared according to the procedure as described in EXAMPLE 7B, substituting EXAMPLE 23A for EXAMPLE 7A (30% yield). ¹H NMR (DMSO-*d*₆) δ 0.76-0.80 (m, 2H), 0.82-0.89 (m, 2H), 1.41-1.48 (m, 2H), 1.59-1.66 (m, 2H), 1.83 (ddd, *J*=11.2, 6.0, 5.7 Hz, 2H), 2.02-2.10 (m, 1H), 2.74-2.83 (m, 2H), 3.14-3.22 (m, 2H), 3.72 (s, 3H), 11.70 (br s, 1H).

EXAMPLE 24

benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-1-yl)-pyrrolidine-1-carboxylate

EXAMPLE 24A

N-carbobenzyloxypyrrolidine-3-carboxylic acid methyl ester

To a solution of N-carbobenzyloxypyrrolidine-3-carboxylic acid (10.15 g, 40.72 mmol) and iodomethane (5.08 mL, 81.44 mmol) in N,N'-dimethylformamide (80 mL) was added powdered potassium hydrogen carbonate (8.15 g, 81.44 mmol) and the reaction mixture stirred at ambient temperature for 5 hours. The mixture was partitioned between ethyl acetate and brine and the organic phase washed with water and concentrated. The residue was purified by flash chromatography on silica gel using 5-50% ethyl acetate in hexane to give the title compound (10.59 g, 99%). MS(DCI): m/z 264 (M+1)⁺.

EXAMPLE 24B

benzyl 3-(2-cyanoacetyl)-pyrrolidine-1-carboxylate

To a solution of acetonitrile (708 μ L, 13.5 mmol) in tetrahydrofuran (30 mL) was added n-butyl lithium (1.6 M solution in hexane, 7.75 mL, 12.4 mmol) at -78°C. The solution was stirred at -78 °C for 10 minutes and a solution of EXAMPLE 24A (3.0 g, 11.34 mmol) in 10 mL of tetrahydrofuran was added. The mixture was stirred at -78°C for 1 hour and was warmed up to -20°C over 3 hours. Water was added and the mixture acidified to pH 3 with 2N hydrochloric acid. The mixture was partitioned between ethyl acetate and brine and the organic phase concentrated. The residue was purified by flash chromatography on silica gel using 30-80% ethyl acetate in hexane to give the title compound (1.91 g, 61%). MS(DCI): m/z 273 (M+1)⁺.

EXAMPLE 24C

benzyl 3-(5-amino-1-methyl-1H-pyrazol-3-yl)-pyrrolidine-1-carboxylate

A solution of EXAMPLE 24B (1.9 g, 7 mmol) and methylhydrazine (367 μ L, 7 mmol) in ethanol (15 mL) was heated at reflux for 4 hours. After cooling, the mixture was concentrated and the residue purified by flash chromatography on silica gel using 0-15% methanol in dichloromethane to give the title compound (1.78 g, 85%). MS(DCI): m/z 301 (M+1)⁺.

EXAMPLE 24D

benzyl 3-(5-amino-4-cyclohex-1-enyl-1-methyl-1H-pyrazol-3-yl)pyrrolidine-1-carboxylate

A solution of EXAMPLE 24C (1.78 g, 5.92 mmol) and cyclohexanone (1.23 mL, 11.85 mmol) in acetic acid (10 mL) was stirred at ambient temperature overnight and at 55°C for 1 day. The acetic acid was removed and the residue partitioned between sodium bicarbonate solution and ethyl acetate. The organic phase was washed with water, concentrated and the residue
5 purified by flash chromatography on silica gel using 70-90% ethyl acetate in hexane to provide the title compound as a light yellow solid (1.19 g, 53%). MS (DCI): m/z 381 ($M+H$)⁺.

EXAMPLE 24E

benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-1-yl)-pyrrolidine-1-
10 carboxylate

To a stirring solution of EXAMPLE 24D (0.967 g, 2.54 mmol) in pyridine (10 mL) was added ethyl isocyanate (0.6 mL, 7.62 mmol) and the solution heated at reflux for 24 hours. After cooling, the mixture was concentrated and the residue recrystallized from methanol. The white
15 solid was collected by filtration, washed with methanol and dried to give the title compound (794 mg, 77%). ¹H NMR (DMSO-d₆): δ 1.66-1.77 (m, 4H), 2.01-2.17 (m, 1H), 2.18-2.27 (m, 1H), 2.37-2.43 (m, 2H), 2.86-2.92 (m, 2H), 3.33-3.46 (m, 1H), 3.49-3.64 (m, 2H), 3.68-3.76 (m, 1H), 3.76 (s, 3H), 3.77-3.85 (m, 1H), 5.08 (s, 2H), 7.29-7.35 (m, 1H), 7.34-7.39 (m, 5H).

EXAMPLE 25

20 3-methyl-1-pyrrolidin-3-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

A solution of EXAMPLE 24E (766 mg, 1.88 mmol) in a mixture of dichloromethane (30 mL) and methanol (30 mL) was treated with 10% palladium on carbon (150 mg) under hydrogen at ambient temperature for 6 hours. Solid material was filtered off and the filtrate concentrated to provide the title compound (564 mg, 100%). ¹H NMR (DMSO-d₆): δ 1.67-1.79 (m, 4H),
25 2.01-2.11 (m, 1H), 2.22-2.32 (m, 1H), 2.38-2.43 (m, 2H), 2.85-2.91 (m, 2H), 3.16-3.26 (m, 2H), 3.30 (dd, $J=11.0, 8.0$ Hz, 1H), 3.46 (dd, $J=11.2, 7.5$ Hz, 1H), 3.78 (s, 3H), 3.81-3.88 (m, 1H).

EXAMPLE 26

3-methyl-1-(1-methylpyrrolidin-3-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

To a solution of EXAMPLE 25 (86 mg, 0.316 mmol) in methanol (6 mL) was added formaldehyde (37 wt% in water, 118 μ L, 1.58 mmol) at ambient temperature and the solution stirred for 1 hour. Sodium cyanoborohydride (99 mg, 1.58 mmol) was added and the mixture stirred overnight and concentrated. The residue was purified by HPLC (Zorbax, C-18, 250x2.54 column, Mobile phase A: 0.1% trifluoroacetic acid in water; B: 0.1% trifluoroacetic acid in acetonitrile; 0-100% gradient) to provide the title compound as the trifluoroacetate salt (110 mg, 79%). ^1H NMR (CD_3OD): δ 1.77-1.87 (m, 4H), 2.24-2.37 (m, 1H), 2.49 (t, $J=6.0$ Hz, 2H), 2.66 (dd, $J=13.6, 6.0$ Hz, 0.5H), 2.90-2.96 (m, 2.5H), 3.02 (s, 1H), 3.08 (s, 1H), 3.24-3.31 (m, 1H), 3.44 (dd, $J=11.4, 7.8$ Hz, 0.5H), 3.62 (dd, $J=11.3, 7.6$ Hz, 0.5H), 3.72-3.83 (m, 0.5H), 3.84 (s, 3H), 3.84 (m, 0.5H), 3.97-4.03 (m, 0.5H), 4.04-4.14 (m, 1H), 4.19-4.26 (m, 0.5H).

EXAMPLE 27

1-(1-isopropylpyrrolidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLE 26, substituting acetone for formaldehyde (77% yield). ^1H NMR (CD_3OD): δ 1.39-1.46 (m, 6H), 1.77-1.88 (m, 4H), 2.23-2.32 (m, 1H), 2.44-2.53 (m, 2.5H), 2.54-2.62 (m, 0.5H), 2.87-3.00 (m, 2H), 3.33-3.42 (m, 1H), 3.51-3.56 (m, 0.5H), 3.58-3.66 (m, 1H), 3.66-3.72 (m, 1H), 3.74-3.79 (m, 0.5H), 3.84 (s, 3H), 3.92 (dd, $J=11.6, 7.3$ Hz, 0.5H), 3.97 (dd, $J=11.4, 5.0$ Hz, 0.5H), 4.03-4.11 (m, 0.5H), 4.14-4.21 (m, 0.5H).

EXAMPLE 28

3-methyl-1-(1-propylpyrrolidin-3-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLE 26, substituting propionaldehyde for formaldehyde (67% yield). ^1H NMR (CD_3OD): δ 1.06 (t, $J=7.5$ Hz, 3H), 1.75-1.89 (m, 6H), 2.21-2.35 (m, 1H), 2.43-2.54 (m, 2.5H), 2.58-2.66 (m, 0.5H), 2.89-2.99 (m, 2H), 3.19-3.26 (m, 1H), 3.29-3.35 (m, 1H), 3.48 (dd, $J=11.5, 8.1$ Hz, 0.5H), 3.64 (dd, $J=11.0, 7.4$ Hz, 0.5H), 3.71-3.82 (m, 1H), 3.84 (s, 3H), 3.98 (dd, $J=11.4, 7.67$ Hz, 0.5H), 4.04-4.11 (m, 1H), 4.17-4.23 (m, 0.5H).

EXAMPLE 29

1-(1-cyclopropylmethylpyrrolidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLE 26, substituting cyclopropanecarboxyaldehyde for formaldehyde (66%
5 yield). ¹H NMR (CD₃OD): δ 0.45-0.50 (m, 2H), 0.74-0.79 (m, 2H), 1.14-1.23 (m, 1H), 1.77-1.88 (m, 4H), 2.23-2.35 (m, 1H), 2.47-2.54 (m, 2.5H), 2.59-2.67 (m, 0.5H), 2.88-3.00 (m, 2H), 3.18 (dd, J=7.2, 2.0 Hz, 1H), 3.26 (dd, J=7.3, 1.2 Hz, 1H), 3.29-3.40 (m, 1H), 3.56 (dd, J=11.6, 8.2 Hz, 0.5H), 3.69 (dd, J=11.6, 7.6 Hz, 0.5H), 3.76-3.84 (m, 1H), 3.84 (s, 1.5H), 3.85 (s, 1.5H), 3.97-4.04 (m, 0.5H), 4.08-4.16 (m, 1H), 4.18-4.24 (m, 0.5H).

10

EXAMPLE 30

1-(1-cyclobutylpyrrolidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLE 26, substituting cyclobutanone for formaldehyde (69% yield). ¹H
15 NMR (CD₃OD): δ 1.77-1.88 (m, 4H), 1.89-2.00 (m, 2H), 2.21-2.33 (m, 3H), 2.34-2.43 (m, 2H), 2.49 (t, J=5.8 Hz, 2.5H), 2.54-2.63 (m, 0.5H), 2.87-3.00 (m, 2H), 3.19-3.27 (m, 1H), 3.42 (dd, J=11.6, 8.2 Hz, 0.5H), 3.53 (dd, J=11.6, 7.6 Hz, 0.5H), 3.62-3.74 (m, 1H), 3.84 (s, 3H), 3.88-3.93 (m, 1H), 3.95-4.01 (m, 0.5H), 4.02-4.06 (m, 0.5H), 4.07-4.13 (m, 0.5H), 4.16-4.22 (m, 0.5H).

20

EXAMPLE 31

(S)-3-methyl-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLES 24B-E and EXAMPLE 25, substituting N-carbobenzyloxy-L-proline
25 methyl ester for EXAMPLE 24A. ¹H NMR (CD₃OD): δ 1.78-1.91 (m, 4H), 2.13-2.20 (m, 1H), 2.22-2.31 (m, 2H), 2.47-2.57 (m, 3H), 2.87-2.99 (m, 2H), 3.39-3.48 (m, 1H), 3.51-3.61 (m, 1H), 3.88 (s, 3H), 4.98-5.41 (m, 1H).

EXAMPLE 32

(S)-3-Methyl-1-(1-methylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

A solution of EXAMPLE 31 (100 mg, 0.37 mmol), triethylamine (52 μ L, 0.37 mmol) and formaldehyde (37% in water, 31 μ L, 1.11 mmol) in 1:1 methanol/N,N'-dimethylformamide
5 (6 mL) was stirred at ambient temperature for 2 hours. Sodium cyanoborohydride (70 mg, 1.11 mmol) and zinc chloride (50 mg, 0.37 mmol) were added and the mixture stirred at 50°C for 18 hours. The mixture was concentrated and the residue purified by HPLC (Zorbax C-8, 0.1% trifluoroacetic acid/acetonitrile/water) to provide the trifluoroacetate salt of the title compound. Treatment of a methanol solution of the trifluoroacetate salt with a solution of hydrochloric acid
10 in ether gave the title compound as the hydrochloride salt (13 mg, 12%). ¹H NMR (CD₃OD): δ 1.79-1.91 (m, 4H), 2.10-2.24 (m, 2H), 2.26-2.36 (m, 1H), 2.49-2.55 (m, 2H), 2.69-2.78 (m, 1H), 2.80-2.88 (m, 1H), 2.93-3.01 (m, 1H), 2.96 (s, 3H), 3.32-3.39 (m, 1H), 3.81-3.87 (m, 1H), 3.91 (s, 3H), 4.97 (t, J =8.4 Hz, 1H).

15 EXAMPLE 33

(S)-3-methyl-1-(1-propyl-pyrrolidin-2-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The title compound as the hydrochloride salt was prepared according to the procedure for EXAMPLE 32, substituting propionaldehyde for formaldehyde (10% yield). ¹H NMR
20 (CD₃OD): δ 0.97 (t, J =7.3 Hz, 3H), 1.64-1.77 (m, 2H), 1.80-1.90 (m, 4H), 2.06-2.14 (m, 1H), 2.15-2.23 (m, 1H), 2.23-2.32 (m, 1H), 2.48-2.56 (m, 2H), 2.66-2.74 (m, 1H), 2.79-2.87 (m, 1H), 2.91-3.00 (m, 1H), 3.07-3.16 (m, 1H), 3.22-3.29 (m, 2H), 3.85-3.90 (m, 1H), 3.91 (s, 3H), 5.01 (t, J =8.4 Hz, 1H).

25 EXAMPLE 34

(S)-1-(1-isopropylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The hydrochloride salt of the title compound was prepared according to the procedure as described in EXAMPLE 32, substituting acetone for formaldehyde (42% yield). ¹H NMR
30 (CD₃OD): δ 1.33 (dd, J =6.7, 3.0 Hz, 6H), 1.79-1.92 (m, 4H), 2.03-2.12 (m, 1H), 2.15-2.28 (m,

2H), 2.52 (t, $J=6.1$ Hz, 2H), 2.63-2.72 (m, 1H), 2.81-2.91 (m, 1H), 2.92-3.01 (m, 1H), 3.35-3.43 (m, 1H), 3.69-3.77 (m, 2H), 3.92 (s, 3H), 5.10 (t, $J=7.8$ Hz, 1H).

EXAMPLE 35 (A-936655.3)

5 (S)-1-(1-cyclopropylmethylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The hydrochloride salt of the title compound was prepared according to the procedure as described in EXAMPLE 32, substituting cyclopropane carboxaldehyde for formaldehyde (36% yield). ^1H NMR (CD_3OD): δ 0.35-0.42 (m, 2H), 0.65-0.71 (m, 2H), 1.06-1.13 (m, 1H), 1.78-
10 1.92 (m, 4H), 2.05-2.13 (m, 1H), 2.18-2.25 (m, 1H), 2.27-2.33 (m, 1H), 2.48-2.55 (m, 2H), 2.67-2.76 (m, 1H), 2.79-2.88 (m, 1H), 2.91-3.00 (m, 1H), 3.08-3.19 (m, 2H), 3.36-3.42 (m, 1H), 3.91 (s, 3H), 3.96-4.01 (m, 1H), 5.02 (t, $J=8.5$ Hz, 1H).

EXAMPLE 36

15 (S)-1-(1-cyclobutylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The hydrochloride salt of the title compound was prepared according to the procedure as described in EXAMPLE 32, substituting cyclobutanone for formaldehyde (27% yield). ^1H NMR (CD_3OD): δ 1.71-1.83 (m, 4H), 1.85-1.93 (m, 3H), 1.94-2.02 (m, 1H), 2.17-2.26 (m, 3H), 2.26-
20 2.31 (m, 1H), 2.32-2.39 (m, 1H), 2.49-2.56 (m, 2H), 2.62-2.71 (m, 1H), 2.83-2.91 (m, 1H), 2.91-3.00 (m, 1H), 3.23-3.29 (m, 1H), 3.73-3.80 (m, 1H), 3.93 (s, 3H), 3.94-4.00 (m, 1H), 4.93 (t, $J=7.6$ Hz, 1H).

EXAMPLE 37

25 (S)-1-(1-isobutylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The hydrochloride salt of the title compound was prepared according to the procedure as described in EXAMPLE 32, substituting isobutyraldehyde for formaldehyde (29% yield). ^1H NMR (CD_3OD): δ 1.09 (d, $J=6.7$ Hz, 6H), 1.80-1.90 (m, 4H), 2.07-2.12 (m, 1H), 2.14-2.19 (m,
30 1H), 2.23-2.30 (m, 1H), 2.51 (s, 2H), 2.66-2.74 (m, 1H), 2.79-2.88 (m, 1H), 2.91-3.00 (m, 1H),

3.07 (t, $J=6.0$ Hz, 2H), 3.11-3.16 (m, 1H), 3.32-3.36 (m, 1H), 3.91 (s, 3H), 3.93-3.97 (m, 1H), 5.00-5.06 (m, 1H).

EXAMPLE 38

5 3-methyl-1-methylaminomethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 38A

methyl-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-ylmethyl)carbamic acid benzyl ester

10 The title compound was prepared according to the procedure as described in EXAMPLE 24B-E, substituting methyl (benzyloxycarbonylmethylamino)acetate for EXAMPLE 24A. MS (DCI/ NH_3) m/z 381 ($\text{M}+\text{H}$)⁺.

EXAMPLE 38B

15 3-methyl-1-methylaminomethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

A solution of EXAMPLE 38A (160 mg) in trifluoroacetic acid (10 mL) was heated at 60°C for 18 hours. The mixture was cooled, diluted with acetonitrile and concentrated. The residue was purified by HPLC (Zorbax C-8, 0.1% trifluoroacetic acid/acetonitrile/water) and the trifluoroacetate salt dissolved in methanol and treated with a solution of hydrochloric acid in ether to provide the title compound as the hydrochloride salt (82 mg, 79%). ¹H NMR (DMSO- d_6): δ 1.67-1.79 (m, 4H), 2.38-2.45 (m, 2H), 2.66 (t, $J=4.3$ Hz, 3H), 2.84-2.95 (m, 2H), 3.86 (s, 3H), 4.40-4.44 (m, 2H), 9.06 (br s, 1H), 12.17 (br s, 1H).

EXAMPLE 39

25 benzyl [4-(3-methyl-5-oxo-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-1-yl)benzyl]carbamate

The title compound was prepared according to the procedure for Example 24B-E, substituting methyl 4-(benzyloxycarbonylaminomethyl) benzoate for Example 24A. ¹H NMR (DMSO- d_6) δ 1.48-1.59 (m, 2H), 1.62-1.74 (m, 2H), 2.36-2.49 (m, 4H), 3.87 (s, 3H), 4.28 (d,

$J=6.1$ Hz, 2H), 5.07 (s, 2H), 7.28-7.34 (m, 3H), 7.35-7.41 (m, 3H), 7.46 (d, $J=8.3$ Hz, 2H), 7.86 (t, $J=6.0$ Hz, 1H).

EXAMPLE 40

5 1-(4-aminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

A solution of Example 39 (90 mg, 0.2 mmol) in trifluoroacetic acid (10 mL) was stirred at 50°C for 4 hours. After cooling, the mixture was concentrated and the residue purified by HPLC (Zorbax, C-18, 250x2.54 column, Mobile phase A 0.1% trifluoroacetic acid in water, B 0.1% trifluoroacetic acid in acetonitrile, 0-100% gradient) to provide 70 mg of the title
10 compound as the trifluoroacetate salt. ^1H NMR (CD_3OD) δ 1.53-1.63 (m, 2H), 1.71-1.80 (m, 2H), 2.42-2.54 (m, 4H), 3.92 (s, 3H), 4.21 (s, 2H), 7.52-7.65 (m, 4H).

Example 41

15 1-(4-dimethylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting Example 40 for Example 25. ^1H NMR (CD_3OD) δ 1.58-1.65 (m, 2H), 1.73-1.81 (m, 2H), 2.49 (q, $J=6.4$ Hz, 4H), 2.91 (s, 6H), 3.93 (s, 3H), 4.40 (s, 2H), 7.58-7.68 (m, 4H).

20

Example 42

1-[4-(isopropylamino-methyl)-phenyl]-3-methyl-3,4,6,7,8,9-hexahydropyrazolo [3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as
25 described in Example 26, substituting Example 40 for Example 25, and substituting acetone for formaldehyde. ^1H NMR (CD_3OD) δ 1.43 (d, $J=6.8$ Hz, 6H), 1.52-1.65 (m, 2H), 1.72-1.82 (m, 2H), 2.41-2.54 (m, 4H), 3.44-3.55 (m, 1H), 3.93 (s, 3H), 4.29 (s, 2H), 7.56-7.67 (m, 4H).

Example 43

30 1-(4-cyclohexylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-

5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting Example 40 for Example 25 and substituting cyclohexanone for formaldehyde. ¹H NMR (CD₃OD) δ 1.20-1.32 (m, 2H), 1.32-1.50 (m, 3H), 1.54-1.64 (m, 2H), 1.70-1.82 (m, 3H), 1.92 (dd, *J*=9.0, 2.9 Hz, 2H), 2.16-2.27 (m, 2H), 2.39-2.55 (m, 4H), 3.10-3.25 (m, 1H), 3.92 (s, 3H), 4.30 (s, 2H), 7.56-7.66 (m, 4H).

Example 441-(4-cyclopentylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure for Example 26, substituting Example 40 for Example 25 and substituting cyclopentanone for formaldehyde. ¹H NMR (CD₃OD) δ 1.50-1.63 (m, 2H), 1.64-1.80 (m, 6H), 1.81-1.94 (m, 2H), 2.12-2.32 (m, 2H), 2.33-2.61 (m, 4H), 3.52-3.77 (m, 1H), 3.92 (s, 3H), 4.28 (s, 2H), 7.43-7.71 (m, 4H).

EXAMPLE 45

1,3-dimethyl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one

The title compound was prepared as described in Winters, G; Sala, A; De Paoli, A.; Ferri, V. *Synthesis* **1984**, 1052-1054.

EXAMPLE 46

1-(4-cycloheptylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLE 26, substituting EXAMPLE 40 for EXAMPLE 25 and substituting cycloheptanone for formaldehyde. ¹H NMR (CD₃OD) δ 1.52-1.59 (m, 4H), 1.61-1.68 (m, 4H), 1.71-1.78 (m, 4H), 1.78-1.90 (m, 2H), 2.10-2.28 (m, 2H), 2.37-2.59 (m, 4H), 3.26-3.43 (m, 1H), 3.92 (s, 3H), 4.30 (s, 2H), 7.45-7.69 (m, 4H).

EXAMPLE 47

1-[4-(isobutylaminomethyl)phenyl]-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting EXAMPLE 40 for EXAMPLE 25 and substituting 2-methylpropionaldehyde for formaldehyde. ¹H NMR (CD₃OD) δ 1.05 (d, *J*=6.7 Hz, 6H), 1.50-1.67 (m, 2H), 1.67-1.85 (m, 2H), 1.95-2.16 (m, 1H), 2.35-2.60 (m, 4H), 2.94 (d, *J*=7.0 Hz, 2H), 3.93 (s, 3H), 4.29 (s, 2H), 7.44-7.76 (m, 4H).

EXAMPLE 48

10 1-4-(cyclobutylaminomethyl)phenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting EXAMPLE 40 for EXAMPLE 25 and substituting cyclobutanone for formaldehyde. ¹H NMR (CD₃OD) δ 1.42-1.66 (m, 2H), 1.69-1.83 (m, 2H), 15 1.86-2.02 (m, 2H), 2.17-2.30 (m, 2H), 2.28-2.42 (m, 2H), 2.41-2.58 (m, 4H), 3.81-3.90 (m, 1H), 3.92 (s, 3H), 4.17 (s, 2H), 7.34-7.77 (m, 4H).

EXAMPLE 49

20 1-{4-[(cyclopropylmethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting EXAMPLE 40 for EXAMPLE 25 and substituting cyclopropyl aldehyde for formaldehyde. ¹H NMR (CD₃OD) δ 0.26-0.53 (m, 2H), 0.59-0.81 (m, 2H), 1.02-1.24 (m, 1H), 1.49-1.65 (m, 2H), 1.68-1.85 (m, 2H), 2.34-2.59 (m, 4H), 3.00 (d, *J*=7.3 25 Hz, 2H), 3.93 (s, 3H), 4.30 (s, 2H), 7.50-7.71 (m, 4H).

EXAMPLE 50

1-{4-[(biscyclopropylmethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was isolated as a side product from the synthesis of EXAMPLE 49. ¹H NMR (CD₃OD) δ 0.20-0.49 (m, 4H), 0.80 (d, *J*=7.9 Hz, 4H), 1.01-1.26 (m, 2H), 1.52-1.69 (m, 2H), 1.70-1.90 (m, 2H), 2.37-2.61 (m, 4H), 2.88-3.19 (m, 2H), 3.19-3.36 (m, 2H), 3.93 (s, 3H), 4.58 (s, 2H), 7.48-7.81 (m, 4H).

5

EXAMPLE 51

3-methyl-1-(4-propylaminomethylphenyl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting EXAMPLE 40 for EXAMPLE 25 and substituting propionaldehyde for formaldehyde. ¹H NMR (CD₃OD) δ 1.04 (t, *J*=7.5 Hz, 3H), 1.48-1.65 (m, 2H), 1.66-1.86 (m, 4H), 2.37-2.60 (m, 4H), 2.92-3.16 (m, 2H), 3.92 (s, 3H), 4.28 (s, 2H), 7.39-7.81 (m, 4H).

15

EXAMPLE 52

1-dimethylaminomethyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

Using the procedure as described in EXAMPLE 26, substituting EXAMPLE 38 for EXAMPLE 25 provided the trifluoroacetate salt of the title compound. Treatment of the methanol solution of the trifluoroacetate salt with a solution of hydrochloric acid in ether gave the title compound as the hydrochloride salt (30% yield). ¹H NMR (CD₃OD) δ 1.78-1.92 (m, 4H), 2.51 (t, *J*=6.1 Hz, 2H), 2.92 (t, *J*=6.0 Hz, 2H), 3.02 (s, 6H), 3.92 (s, 3H), 4.63 (s, 2H).

EXAMPLE 53

1-[(isopropylmethylamino)methyl]-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

25

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in Example 26, substituting EXAMPLE 38 for EXAMPLE 25 and substituting acetone for formaldehyde. Treatment of the methanol solution of the trifluoroacetate salt with a solution of hydrochloric acid in ether gave the title compound as the hydrochloride salt (39% yield). ¹H NMR (CD₃OD) δ 1.45 (dd, *J*=6.7, 1.8 Hz, 6H), 1.78-1.93 (m, 4H), 2.51 (t, *J*=6.1 Hz, 2H), 2.91

30

(s, 3H), 2.93-2.98 (m, 2H), 3.80-3.88 (m, 1H), 3.92 (s, 3H), 4.43 (d, $J=14.7$ Hz, 1H), 4.74 (d, $J=14.7$ Hz, 1H).

EXAMPLE 54

5 (S)-3-methyl-1-piperidin-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedures as described in EXAMPLES 24 and 25, substituting (R)-(+)-1-(carboxybenzyl)-2-piperidine carboxylic acid for 1-N-carboxybenzylpyrrolidine-3-carboxylic acid, followed by purification by HPLC (Zorbax, C-18, 250x2.54 column, Mobile phase A: 0.1% trifluoroacetic acid in water; B: 0.1% trifluoroacetic acid in acetonitrile; 0-100% gradient). ^1H NMR (CD_3OD) δ 1.76-1.83 (m, 4H), 1.85 (d, $J=6.1$ Hz, 2H), 1.87-1.90 (m, 1H), 1.93-2.03 (m, 2H), 2.28 (d, $J=13.4$ Hz, 1H), 2.44-2.57 (m, 2H), 2.78-2.88 (m, 1H), 2.94-3.05 (m, 1H), 3.16-3.27 (m, 1H), 3.47 (d, $J=12.5$ Hz, 1H), 3.89 (s, 3H), 4.65 (d, $J=8.8$ Hz, 1H).

15

EXAMPLE 55

3-methyl-1-{4-[(2-methyltetrahydrofuran-3-ylamino)methyl]phenyl}-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure as described in EXAMPLE 26, substituting EXAMPLE 40 for EXAMPLE 25 and substituting 2-methyltetrahydrofuran-3-one for formaldehyde. ^1H NMR (CD_3OD) δ 1.32 (d, $J=6.4$ Hz, 3H), 1.52-1.63 (m, 2 H), 1.69-1.84 (m, 2H), 2.10-2.24 (m, 1H), 2.40-2.57 (m, 4H), 3.55-3.68 (m, 1H), 3.93 (s, 3H), 3.94-4.01 (m, 2H), 4.00-4.14 (m, 1H), 4.18-4.27 (m, 1H), 4.34 (s, 2H), 7.47-7.75 (m, 4H).

25

EXAMPLE 56

1-(3-aminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 56A

methyl 3-[(4-methoxybenzylamino)methyl]benzoate

To a solution of 4-methoxybenzylamine (5.6 g, 41.2 mmol) in methanol (100 mL) was added methyl 3-formylbenzoate (5.2 g, 31.7 mmol) at ambient temperature. The solution was stirred for 30 minutes and sodium cyanoborohydride (3.0 g) was added. The mixture was stirred at ambient temperature overnight and concentrated to about 20 mL. Water (100 mL) was added and the mixture acidified with 2N hydrochloric acid to a pH of 3. The mixture was partitioned between ethyl acetate and brine and the organic phase separated and concentrated. The residue was purified by flash chromatography on silica gel using 10% methanol in ethyl acetate to give 6.0 g (67%) of the title compound. MS (DCI): m/z 286 (M+H)⁺.

EXAMPLE 56B

methyl 3-{[benzyloxycarbonyl-(4-methoxybenzyl)amino]methyl}benzoate

To a mixture of EXAMPLE 56A (6 g, 21 mmol), potassium carbonate (5 g, 36 mmol), dioxane (150 mL) and water (50 mL) was added benzyl chloroformate (3.4 mL, 23 mmol) at ambient temperature and the mixture stirred overnight. Catalytic piperazine was added and the mixture stirred for another 30 minutes and concentrated. The residue was partitioned between dilute hydrochloric acid and ethyl acetate and the organic phase was washed with water and concentrated. The residue was purified by flash chromatography on silica gel using 20% ethyl acetate in hexane to give 8.2 g (93%) of the title compound. MS (DCI): m/z 420 (M+H)⁺.

EXAMPLE 56C

benzyl [3-(2-cyanoacetyl)benzyl]-(4-methoxybenzyl)carbamate

A solution of acetonitrile (1.46, 28 mmol) in tetrahydrofuran (180 mL) was treated with *n*-butyl lithium (1.6 M solution in hexane, 14 mL, 22.2 mmol) at -78°C for 10 minutes. A solution of EXAMPLE 56B (7.8 g, 18.6 mmol) in 40 mL of tetrahydrofuran was added and the mixture stirred at -78°C for 1 hour and at ambient temperature for 1 hour. After quenching with water, the mixture was acidified with 2N hydrochloric acid to a pH of 3. The mixture was partitioned between ethyl acetate and brine and the organic phase washed with water and concentrated. The residue was purified by flash chromatography on silica gel using 30% ethyl acetate in hexane to give 7.7g (96%) of the title compound. MS (DCI): m/z 429 (M+H)⁺.

EXAMPLE 56D

benzyl [3-(5-amino-1-methyl-1H-pyrazol-3-yl)benzyl]-(4-methoxybenzyl)carbamate

To a suspension of EXAMPLE 56C (7.7 g, 18 mmol) in ethanol (60 mL) was added methylhydrazine (1.4 mL, 27 mmol) at ambient temperature and the solution heated to reflux for 4 hours. After cooling, the mixture was concentrated and the residue purified by flash chromatography on silica gel using ethyl acetate to give 4.5 g (56%) of the title compound. MS (DCI): m/z 456 (M+H)⁺.

EXAMPLE 56E

benzyl [3-(5-amino-4-cyclohex-1-enyl-1-methyl-1H-pyrazol-3-yl)benzyl]-(4-methoxybenzyl)carbamate

A solution of EXAMPLE 56D (4.5 g, 10 mmol) in acetic acid (100 mL) was treated with cyclohexanone (2 mL, 20 mmol) at ambient temperature for 4 days. The mixture was concentrated and the residue partitioned between dilute sodium hydroxide and ethyl acetate. The organic layer was washed with water and concentrated and the residue purified by flash chromatography on silica gel using ethyl acetate to give 3.8 g (71%) of the title compound. MS (DCI): m/z 537 (M+H)⁺.

EXAMPLE 56F

benzyl (4-methoxybenzyl)-[3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-1-yl)benzyl]carbamate

To a solution of EXAMPLE 56E (3.8 g, 7.1 mmol) in pyridine (100 mL) was added ethyl isocyanate (1.6 mL, 21 mmol) and the solution stirred at ambient temperature for 24 hours. The mixture was heated at reflux overnight, cooled, concentrated and the residue purified by flash chromatography on silica gel using ethyl acetate to give 3.4 g (85%) of the title compound. MS (DCI): m/z 563 (M+H)⁺.

EXAMPLE 56G

1-(3-aminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

A solution of EXAMPLE 56F (3.4 g) in trifluoroacetic acid (20 mL) was stirred at 55°C for 6 hours. The mixture was concentrated and the residue purified by HPLC (Zorbax, C-18, 250x2.54 column, Mobile phase A: 0.1% trifluoroacetic acid in water; B: 0.1% trifluoroacetic acid in acetonitrile; 0-100% gradient) to give the title compound (1.3 g, 50%) as the trifluoroacetate salt. ¹H NMR (CD₃OD) δ 2.28-2.40 (m, 2H), 2.44-2.57 (m, 2H), 3.08-3.37 (m, 4H), 4.69 (s, 3H), 4.93 (s, 2H), 8.26-8.37 (m, 3H), 8.43 (s, 1H).

Example 57

1-{3-[(dimethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 57A

benzyl [3-(3-methyl-5-oxo-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-1-yl)benzyl]carbamate

The title compound was prepared according to the procedure for Example 24B-E, substituting methyl 3-(benzyloxycarbonylaminomethyl) benzoate for Example 24A.

EXAMPLE 57B

1-(3-aminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared as described in Example 40, substituting Example 57A for Example 39.

Example 57C

1-{3-[(dimethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure for Example 26, substituting Example 57B for Example 25. MS (DCI): m/z 337 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.58 - 1.64 (m, 2 H), 1.74 - 1.79 (m, 2 H), 2.46 - 2.53 (m, 4 H), 2.90 (s, 6 H), 3.93 (s, 3 H), 4.40 (s, 2 H), 7.60 - 7.62 (m, 2 H), 7.65 - 7.69 (m, 2 H); Anal. Calcd for C₂₀H₂₄N₄O•1.5 TFA: C, 54.44; H, 5.06; N, 11.04. Found: C, 54.14; H, 4.75; N, 11.01.

Example 58

3-methyl-1-[(2S)-1-propylpiperidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The hydrochloride salt of the title compound was prepared according to procedure as describe in Example 26, substituting Example 54 for Example 25, and substituting propionaldehyde for formaldehyde. The trifluoroacetate salt was treated with 1.0 M solution of HCl in ether to yield the HCl salt. MS (DCI/NH₃) m/z 329 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD): δ 0.86 (t, J=7.36 Hz, 3H), 1.60 - 1.71 (m, 1H), 1.73 - 1.79 (m, 2H), 1.79 - 1.84 (m, J=8.59 Hz, 2H), 1.84 - 1.90 (m, 2H), 1.90 - 1.97 (m, 2H), 1.98 - 2.08 (m, 2H), 2.23 - 2.33 (m, 1H), 2.49 - 2.58 (m, 2H), 2.76 - 2.88 (m, 1H), 2.93 - 3.07 (m, 3H), 3.18 - 3.28 (m, 1H), 3.76 (d, J=11.97 Hz, 1H), 3.92 (s, 3H), 4.66 (dd, J=11.66, 2.76 Hz, 1H). Anal. Calcd for C₁₉H₂₈N₄O•1.9 HCl: C, 57.38; H, 7.58; N, 14.09. Found: C, 57.08; H, 7.97; N, 13.79.

Example 59

15 1-azetidin-3-yl-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedures for Examples 24A-24E and Example 25, substituting N-carbobenzyloxazetidine-3-carboxylic acid for N-carbobenzyloxypyrrolidine-3-carboxylic acid. MS (DCI): m/z 259 (M+H)⁺; ¹H NMR (400 MHz, DMSO-d₆): δ 1.66 - 1.75 (m, 4 H), 2.37 - 2.41 (m, 2 H), 2.70 (br s, 2 H), 3.84 (s, 3 H), 4.22 - 4.31 (m, 4 H), 4.42 (q, J=8.08 Hz, 1 H), 8.94 (br s, 1 H), 11.92 (br s, 1 H); Anal. Calcd for C₁₄H₁₈N₄O•1.32 TFA: C, 48.88; H, 4.76; N, 13.70. Found: C, 48.86; H, 4.53; N, 13.80.

Example 60

25 1-(1-cyclobutylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure for Example 26, substituting Example 59 for Example 25, and cyclobutanone for formaldehyde. MS (DCI): m/z 313 (M+H)⁺; ¹H NMR (500 MHz, Pyridine-d₅): δ 1.51 - 1.77 (m, 6 H), 2.07 - 2.14 (m, 2 H), 2.30 - 2.37 (m, 2 H), 2.70 (t, J=5.95 Hz, 2 H), 2.81 (t, J=5.80 Hz, 2 H), 3.91 - 3.96 (m, 1 H), 4.01 (s, 3 H), 4.37 (t, J=7.93 Hz, 2 H), 4.61 - 4.66 (m, 1 H), 4.71 (t, J=8.85 Hz, 2 H);

Anal. Calcd for C₁₈H₂₄N₄O•1.6 TFA: C, 51.46; H, 5.21; N, 11.32. Found: C, 51.46; H, 5.32; N, 11.28.

Example 61

5 1-[4-(aminomethyl)benzyl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to the procedure for Examples 24 B-24E and Example 25, substituting methyl

4-((benzyloxycarbonylamino)methyl)phenyl)acetate for Example 24A. MS (DCI): m/z 323

10 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.61 - 1.72 (m, 4 H), 2.42 (t, *J*=5.37 Hz, 2 H), 2.65 (t, *J*=5.98 Hz, 2 H), 3.86 (s, 3 H), 4.07 (s, 2 H), 4.24 (s, 2 H), 7.23 (d, *J*=8.59 Hz, 2 H), 7.36 (d, *J*=8.29 Hz, 2 H).

Example 62

15 3-methyl-1-[(2R)-1-(2-phenylethyl)pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The hydrochloride salt of the title compound was prepared according to procedure for Example 26 substituting phenylacetaldehyde for formaldehyde, and substituting Example 102 for Example 25. The trifluoroacetate salt was treated with 1.0 M solution of HCl in ether to yield

20 the HCl salt. MS (DCI/NH₃) m/z 377 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.75 - 1.89 (m, 4H), 2.03 - 2.12 (m, 1H), 2.14 - 2.22 (m, 1H), 2.24 - 2.33 (m, 1H), 2.46 - 2.55 (m, 2H), 2.66 - 2.74 (m, 1H), 2.77 - 2.87 (m, 2H), 2.93 - 3.03 (m, 1H), 3.04 - 3.12 (m, 1H), 3.34 - 3.42 (m, 1H), 3.46 - 3.54 (m, 1H), 3.59 - 3.68 (m, 1H), 3.91 (s, 3H), 3.92 - 3.97 (m, 1H), 5.04 (t, *J*=8.09 Hz, 1H), 7.17 - 7.20 (m, 2H), 7.21 - 7.23 (m, 1H), 7.25 - 7.29 (m, 2H).

25

Example 63

3-methyl-1-piperidin-3-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedures for Examples 24A-24E and Example 25 substituting 1-benzyl piperidine-1,3-dicarboxylate for 1-benzyl pyrrolidine-1,3-

30 dicarboxylate. The product as TFA salt was treated with 1.0 M solution of HCl in ether to yield the HCl salt. MS (DCI/NH₃) m/z 287 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.65 - 1.78 (m, 5H), 1.80 - 1.88 (m, 2H), 1.98 - 2.07 (m, 1H), 2.36 - 2.46 (m, 2H), 2.81 - 2.97 (m, 3H), 2.99 - 3.11 (m, 1H), 3.27 (d, *J*=12.21 Hz, 1H), 3.37 (d, *J*=11.90 Hz, 1H), 3.52 - 3.59 (m, 1H), 3.79 (s,

3H). Anal. Calcd for $C_{16}H_{22}N_4O \cdot 2.4 HCl$: C, 51.40; H, 6.58; N, 14.99. Found: C, 51.69; H, 6.46; N, 14.64.

Example 64

5 3-methyl-1-piperidin-4-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as trifluoroacetate salt according to procedure for Examples 24A-24E and Example 25 substituting 1-benzyl piperidine-1,4-dicarboxylate for 1-benzyl pyrrolidine-1,3-dicarboxylate. MS (DCI/ NH_3) m/z 287 ($M+H$)⁺; ¹H NMR (500 MHz, CD_3OD): δ 1.67 - 1.79 (m, 4H), 1.85 - 1.95 (m, 2H), 1.97 - 2.05 (m, 2H), 2.35 - 2.46 (m, 2H),
10 2.85 - 2.94 (m, 2H), 3.06 (q, $J=11.60$ Hz, 2H), 3.27 - 3.34 (m, 3H), 3.77 (s, 3H); Anal. Calcd for $C_{16}H_{22}N_4O \cdot 1.0 TFA$: C, 54.00; H, 5.79; N, 13.99. Found: C, 53.98; H, 5.61; N, 13.87.

Example 65

1-cyclopropyl-3,7-dimethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared according to procedure for Examples 7A-7B substituting 4-methylcyclohexanone for cyclohexanone. Yield: 47%. MS (DCI/ NH_3) m/z 258 ($M+H$)⁺; ¹H NMR (500 MHz, $DMSO-d_6$): δ 0.72 - 0.76 (m, 1H), 0.82 - 0.87 (m, 3H), 1.04 (d, $J=6.41$ Hz, 3H), 1.26 - 1.38 (m, 1H), 1.67 - 1.78 (m, 1H), 1.82 - 1.91 (m, 2H), 2.07 - 2.17 (m, 1H), 2.62 (dd, $J=16.48, 3.97$ Hz, 1H), 2.88 - 3.01 (m, 1H), 3.16 (d, $J=20.14$ Hz, 1H), 3.70 (s, 3H), 11.86 (br s, 1H). Anal. Calcd for $C_{15}H_{19}N_3O$: C, 70.01; H, 7.44; N, 16.33. Found: C, 70.00; H, 7.56; N, 16.36.

Example 66

1-cyclopropyl-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]-2,7-naphthyridin-5-one

25

Example 66A

benzyl 1-cyclopropyl-3-methyl-5-oxo-4,5,8,9-tetrahydro-3H-pyrazolo[3,4-c][2,7]naphthyridine-7(6H)-carboxylate

The title compound was prepared according to procedure for Examples 7A-7B substituting benzyl-4-oxo-1-piperidine carboxylate for cyclohexanone. MS (DCI/ NH_3) m/z 379 ($M+H$)⁺.

30

Example 66B

1-cyclopropyl-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]-2,7-naphthyridin-5-one

5 A solution of Example 66A (180 mg, 0.48 mmol) in trifluoroacetic acid (5 mL) was heated at 50 °C for 16 h. After cooling, acetonitrile was added and the mixture concentrated on a rotary evaporator. The residue was separated by HPLC (Zorbax C-18, 0.1% TFA/CH₃CN/H₂O) to provide the title product as trifluoroacetate salt. This product was treated with 1.0 M solution of HCl in ether to yield the HCl salt. Yield: 92 mg (60%). MS (DCI/NH₃) m/z 245 (M+H)⁺; ¹H NMR (500 MHz, DMSO-*d*₆): δ 0.77 - 0.85 (m, 2H), 0.86 - 0.92 (m, 2H), 2.04 - 2.17 (m, 1H), 3.24 - 3.32 (m, 2H), 3.33 - 3.41 (m, 2H), 3.74 (s, 3H), 3.84 - 3.96 (m, 2H), 5.27 (br s, 1H), 9.64 (br s, 1H). Anal. Calcd for C₁₃H₁₆N₄O•2.1 HCl: C, 48.91; H, 5.69; N, 17.46. Found: C, 48.91; H, 5.88; N, 17.11.

Example 67

3-ethyl-1-(1-methylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as TFA salt according to procedure for Example 69A-69F, substituting (±)-1-benzyl 2-methyl pyrrolidine-1,2-dicarboxylate for (R)-1-benzyl 2-methyl pyrrolidine-1,2-dicarboxylate and ethylhydrazine for (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt. MS (DCI): m/z 301 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.42 (t, *J*=7.36 Hz, 2 H), 1.75 - 1.91 (m, 4 H), 2.02 - 2.26 (m, 2 H), 2.23 - 2.37 (m, 1 H), 2.52 (t, *J*=5.68 Hz, 2 H), 2.59 - 2.77 (m, 20 1 H), 2.77 - 2.89 (m, 1 H), 2.88 - 3.03 (m, 1 H), 2.96 (s, 3 H), 3.26 - 3.41 (m, 1 H), 3.75 - 3.93 (m, 1 H), 4.30 (q, *J*=7.36 Hz, 2 H), 4.96 (t, *J*=8.29 Hz, 1 H).

Example 68

25 1-{4-[(dimethylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The trifluoroacetate salt of the title compound was prepared according to procedure for Example 26 substituting Example 61 for Example 25. MS (DCI/NH₃) m/z 351 (M+H)⁺; ¹H NMR (500 MHz, DMSO-*d*₆): δ 1.61 (m, 4 H), 2.34 - 2.37 (m, 2 H), 2.61 - 2.65 (m, 2 H), 2.70 (s, 3 H), 2.71 (s, 3 H), 3.80 (s, 3 H), 4.19 (s, 2 H), 4.22 (d, *J*=5.19 Hz, 2 H), 7.25 (d, *J*=8.24 Hz, 2 30 H), 7.39 (d, *J*=7.93 Hz, 2 H), 9.61 (s, 1 H). Anal. Calcd for C₂₁H₂₆N₄O•2 TFA: C, 51.90; H, 4.88; N, 9.68. Found: C, 51.91; H, 4.20; N, 9.42.

Example 69

1-[(2R)-1-methylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

EXAMPLE 69A

5 (R)-benzyl 2-(2-cyanoacetyl)pyrrolidine-1-carboxylate

The title compound was prepared according to procedure for Example 24B, substituting (R)-1-benzyl 2-methyl pyrrolidine-1,2-dicarboxylate for EXAMPLE 24A. MS(DCI): m/z 273 (M+1)⁺.

10 Example 69B

(R)-2-[5-Amino-1-(2,2,2-trifluoro-ethyl)-1H-pyrazol-3-yl]-pyrrolidine-1-carboxylic acid benzyl ester

To a solution of Example 69A (3.0 g, 11 mmol) in ethanol (10 mL) was added (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt (2.3 g, 11 mmol) and triethylamine (1.84 mL, 13.2 mmol) at
15 rt. The solution was heated under reflux for 4 h. After cooling, the reaction mixture was concentrated and the residue was separated by flash chromatography to give Example 69B. Yield: 3.5 g (88%). MS (DCI): m/z 369 (M+H)⁺.

Example 69C

20 (R)-2-[5-Amino-4-cyclohex-1-enyl-1-(2,2,2-trifluoro-ethyl)-1H-pyrazol-3-yl]-pyrrolidine-1-carboxylic acid benzyl ester

The title compound was prepared according to the procedure for Example 24D, substituting Example 69B for Example 24C. Yield: 50%. MS (DCI): m/z 449 (M+H)⁺.

25 Example 69D

benzyl (2R)-2-[5-oxo-3-(2,2,2-trifluoroethyl)-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl]pyrrolidine-1-carboxylate

The title compound was prepared according to the procedure for 24E, substituting Example 69D for Example 24D. Yield: 33%. MS (DCI): m/z 475 (M+H)⁺.

30

Example 69E

1-[(2*R*)-pyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one

A solution of Example 69D (0.55 g, 1.16 mmol) in TFA (10 ml) was heated at 50 °C for 6 h. Volatile were removed and the residual was purified by HPLC (C-18, 0-100% gradient CH₃CN with 0.1% TFA in water with 0.1% TFA) to provide Example 69E. MS (DCI): *m/z* 341 (M+H)⁺.

Example 69F

1-[(2*R*)-1-methylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for Example 26, substituting Example 69E for 25. MS (DCI): *m/z* 355 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD) δ 1.74 - 1.98 (m, 5 H), 2.04 - 2.26 (m, 2 H), 2.26 - 2.39 (m, 1 H), 2.53 - 2.67 (m, 2 H), 2.71 - 2.81 (m, 1 H), 2.82 - 2.95 (m, 1 H), 2.98 (s, 2 H), 3.00 - 3.09 (m, 1 H), 3.30 - 3.43 (m, 1 H), 3.67 - 3.94 (m, 1 H), 4.95 - 5.21 (m, 3 H).

Example 70

3-methyl-1-(1-methylpiperidin-3-yl)-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one
The title compound was prepared according to procedure for Example 26 substituting Example 63 for Example 25. The product was treated with 1.0 M solution of HCl in ether to yield the HCl salt. Yield: 31%. MS (DCI/NH₃) *m/z* 301 (M+H)⁺; ¹H NMR (500 MHz, pyridine-*d*₅): δ 1.65 - 1.75 (m, 4H), 1.77 - 1.84 (m, 1H), 1.85 - 1.93 (m, 1H), 2.24 - 2.32 (m, 1H), 2.58 - 2.68 (m, 1H), 2.78 - 2.83 (m, 3H), 2.84 (s, 3H), 3.08 - 3.15 (m, 1H), 3.29 - 3.42 (m, 2H), 3.47 - 3.57 (m, 1H), 3.82 - 3.91 (m, 1H), 3.97 (s, 3H), 4.38 - 4.47 (m, 1H), 5.40 (br s, 1H). Anal. Calcd for C₁₇H₂₄N₄O•2.1 HCl: C, 54.17; H, 6.98; N, 14.86. Found: C, 54.37; H, 6.95; N, 14.56.

Example 71

2-{1-[(2*R*)-1-methylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3*H*-pyrazolo[3,4-*c*]isoquinolin-3-yl}ethyl ethylcarbamate

The title compound was prepared according to the procedure for Example 69A-69F, substituting 2-hydrazino-ethanol for (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt. MS (DCI): *m/z* 388 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD) δ 0.94 (t, *J*=7.17 Hz, 3 H), 1.62 - 1.96 (m, 4 H), 2.06

- 2.24 (m, 2 H), 2.25 - 2.35 (m, 1 H), 2.51 (s, 3 H), 2.69 - 2.75 (m, 1 H), 2.77 - 2.86 (m, 1 H), 2.89 - 3.09 (m, 4 H), 3.32 - 3.43 (m, 2 H), 3.62 - 3.92 (m, 1 H), 4.31 - 4.43 (m, 1 H), 4.42 - 4.63 (m, 3 H), 4.96 (t, $J=8.39$ Hz, 1 H).

5

Example 72

3-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanamide

Example 72A

10 benzyl (2R)-2-[3-(2-cyanoethyl)-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl]pyrrolidine-1-carboxylate

The title compound was prepared according to the procedure for Example 69D, substituting 3-hydrazino-propionitrile for (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt. MS (DCI): m/z 446 (M+H)⁺;

15

Example 72B

3-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanamide

The title compound was prepared as TFA salt according to the procedure for Example 20 69E, substituting Example 72A for Example 69D. MS (DCI): m/z 330 (M+H)⁺. ¹H NMR (500 MHz, CD₃OD) δ 1.72 - 1.94 (m, 4 H), 2.07 - 2.32 (m, 3 H), 2.42 - 2.58 (m, 3 H), 2.79 (t, $J=6.41$ Hz, 2 H), 2.83 - 3.00 (m, 2 H), 3.37 - 3.50 (m, 1 H), 3.48 - 3.65 (m, 1 H), 4.42 - 4.55 (m, 2 H), 5.11 - 5.22 (m, 1 H).

25

Example 73

3-methyl-1-(1-methylpiperidin-4-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 26 substituting Example 64 for Example 25. The product was treated with 1.0 M solution of HCl in ether to yield the HCl salt. Yield: 59%. MS (DCI/NH₃) m/z 301 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): 30 δ 1.77 - 1.88 (m, 5H), 2.07 - 2.18 (m, 2H), 2.19 - 2.25 (m, 2H), 2.50 (t, $J=5.95$ Hz, 2H), 2.92 (s, 3H), 2.96 - 3.02 (m, 2H), 3.15 - 3.24 (m, 2H), 3.57 - 3.65 (m, 2H), 3.83 (s, 3H). Anal. Calcd for C₁₇H₂₄N₄O•2.1 HCl: C, 54.18; H, 6.98; N, 14.86. Found: C, 54.11; H, 7.09; N, 14.70.

Example 74

3-{1-[(2R)-1-methylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-
c]isoquinolin-3-yl}propanenitrile

5

Example 74A

3-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-
c]isoquinolin-3-yl}propanenitrile

10 The title compound was prepared according to the procedure for Example 69E,
substituting Example 72A for Example 69D. MS (DCI): m/z 326 (M+H)⁺.

Example 74B

3-{1-[(2R)-1-methylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-
c]isoquinolin-3-yl}propanenitrile

15 The title compound was prepared according to the procedure for Example 26, substituting
Example 74A for Example 25. MS (DCI): m/z 326 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD) δ
1.86 (d, J=5.52 Hz, 3 H), 2.04 - 2.42 (m, 4 H), 2.56 (s, 3 H), 2.74 (s, 1 H), 2.89 (s, 1 H), 2.99 (s,
3 H), 3.07 (t, J=6.29 Hz, 2 H), 3.31 - 3.47 (m, 1 H), 3.60 - 3.97 (m, 1 H), 4.32 - 4.74 (m, 2 H),
5.01 (t, J=7.98 Hz, 1 H).

20

Example 75

3-[2-(dimethylamino)benzyl]-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-
c]isoquinolin-5-one

25 The title compound was prepared according to the procedures for Examples 69B-69E,
substituting (2-hydrazino-phenyl)-dimethyl-amine for (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt.
MS (DCI): m/z 392 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD) δ 1.54 - 2.00 (m, 4 H), 2.07 - 2.32
(m, 3 H), 2.44 - 2.67 (m, 3 H), 2.95 (t, J=5.98 Hz, 2 H), 3.01 (s, 6 H), 3.35 - 3.47 (m, 1 H), 3.46 -
3.62 (m, 1 H), 5.18 (t, J=7.36 Hz, 1 H), 5.59 (s, 2 H), 7.04 - 7.21 (m, 2 H), 7.34 - 7.44 (m, 1 H),
7.47 - 7.50 (m, 1 H).

30

Example 76

1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5

Example 76A

3-(3-hydroxybenzyl)-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to the procedure for Example 69E, substituting 3-hydroxybenzylhydrazine for (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt. MS

10 (DCI): m/z 365 (M+H)⁺;

Example 76B

1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15

The title compound was prepared as TFA salt according to the procedure for Example 26, substituting Example 76A for Example 25, and cyclobutanone for formaldehyde. MS (DCI):

m/z 419 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD) δ 1.55 - 1.77 (m, 3 H), 1.76 - 2.01 (m, 5 H),

2.08 - 2.38 (m, 5 H), 2.54 (t, J=5.68 Hz, 2 H), 2.58 - 2.75 (m, 1 H), 2.77 - 3.03 (m, 2 H), 3.12 -

3.28 (m, 1 H), 3.63 - 3.77 (m, 1 H), 3.80 - 4.01 (m, 1 H), 4.93 (t, J=7.67 Hz, 1 H), 5.29 - 5.43 (m,

20 1 H), 5.44 - 5.61 (m, 1 H), 6.56 (s, 1 H), 6.59 - 6.78 (m, 2 H), 7.11 (t, J=7.82 Hz, 1 H).

Example 77

3-[3-(dimethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25

Example 77A

benzyl (2R)-2-[5-oxo-3-(3-oxopropyl)-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl]pyrrolidine-1-carboxylate

A solution of Example 72A (2.0 g, 4.5 mmol) in acetic acid (10 ml) was treated with
30 Raney nickel (200 mg) at rt. The reaction mixture was stirred under hydrogen for 6 h. Solid

material was filtered off and the filtrate was concentrated. The residue was separated by flash chromatography to give Example 77A. Yield: 0.5 g (25%). MS (DCI): m/z 449 ($M+H$)⁺.

Example 77B

5 benzyl (2*R*)-2-{3-[3-(dimethylamino)propyl]-5-oxo-4,5,6,7,8,9-hexahydro-3*H*-pyrazolo[3,4-
c]isoquinolin-1-yl}pyrrolidine-1-carboxylate

The title compound was prepared according to the procedure for Example 26, substituting Example 77A for formaldehyde, and dimethyl amine for Example 25. MS (DCI): m/z 4 480 (M+H)⁺.

Example 77C

3-[3-(dimethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for Example 69E, substituting Example 77B for Example 69D. MS (DCI): m/z 344 ($M+H$)⁺; ¹H NMR (500 MHz, CD₃OD) δ 1.73 - 1.94 (m, 5 H), 2.07 - 2.36 (m, 6 H), 2.44 - 2.67 (m, 3 H), 2.87 (s, 6 H), 2.91 - 3.07 (m, 1 H), 3.06 - 3.24 (m, 2 H), 3.39 - 3.51 (m, 1 H), 3.52 - 3.62 (m, 1 H), 5.21 (t, $J=7.17$ Hz, 2 H).

Example 78

1-cyclobutyl-3-[3-(ethylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

Example 78A

25 3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3*H*-pyrazolo[3,4-*c*]isoquinolin-3-yl)propanenitrile

The title compound was prepared according to the procedure for Example 85B-85D, substituting 3-hydrazino-propionitrile for Example 85A. MS (DCI): m/z 299 (M+H)⁺;

Example 78B

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3*H*-pyrazolo[3,4-*c*]isoquinolin-3-yl)propanal

The title compound was prepared according to the procedure for Example 77A, substituting Example 78A for Example 72A. MS (DCI): m/z 4300 ($M+H$)⁺.

5

Example 78C

1-cyclobutyl-3-[3-(ethylamino)propyl]-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for Example 26, substituting Example 78B for formaldehyde and ethylamine for Example 25. MS (DCI): m/z 329 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.15 (t, $J=7.32$ Hz, 3 H), 1.56 - 1.80 (m, 3 H), 1.77 - 1.90 (m, 1 H), 1.92 - 2.08 (m, 4 H), 2.22 - 2.32 (m, 4 H), 2.33 - 2.41 (m, 2 H), 2.79 (s, 2 H), 2.83 - 3.07 (m, 4 H), 3.54 - 3.91 (m, 2 H), 4.25 (t, $J=6.71$ Hz, 1 H), 8.41 (s, 2 H).

10

Example 79

15 1-cyclobutyl-3-(3-pyrrolidin-1-ylpropyl)-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for Example 26, substituting Example 78B for formaldehyde and pyrrolidine for Example 25. MS (DCI): m/z 355 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.60 - 1.77 (m, 4 H), 1.77 - 1.89 (m, 2 H), 1.92 - 2.03 (m, 3 H), 2.05 - 2.16 (m, 2 H), 2.16 - 2.40 (m, 6 H), 2.79 (s, 2 H), 2.86 - 3.03 (m, 2 H), 3.04 - 3.21 (m, 2 H), 3.55 (dd, $J=10.37, 5.19$ Hz, 3 H), 3.67 - 3.95 (m, 2 H), 4.23 (t, $J=6.71$ Hz, 1 H), 9.64 (s, 1 H).

20

Example 80

25 1-cyclobutyl-3-[3-(dimethylamino)propyl]-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for Example 26, substituting Example 78B for formaldehyde and dimethyl amine for Example 25. MS (DCI): m/z 329 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.52 - 1.78 (m, 3 H), 1.78 - 1.88 (m, 1 H),

1.91 - 2.06 (m, 1 H), 2.05 - 2.18 (m, 2 H), 2.21 - 2.35 (m, 4 H), 2.37 (s, 2 H), 2.72 (d, $J=4.88$ Hz, 6 H), 2.79 (s, 2 H), 2.90 - 3.12 (m, 2 H), 3.71 - 3.91 (m, 2 H), 4.24 (t, $J=6.87$ Hz, 2 H), 10.63 (s, 1 H).

5

Example 81

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)propanoic acid

A suspension of Example 78A (100 mg, 0.34 mmol) in 6N HCl (20 ml) was heated at 100 °C for 20 h. After cooling to room temperature the formed solid was collected by filtration, washed with water and MeOH to give Example 81 (100 mg, 90%). MS (DCI): m/z 318 (M+H)⁺;

10 ¹H NMR (500 MHz, DMSO- d_6) δ 1.60 - 1.77 (m, 5 H), 1.76 - 1.91 (m, 1 H), 1.90 - 2.08 (m, 1 H), 2.16 - 2.31 (m, 4 H), 2.30 - 2.43 (m, 2 H), 2.65 - 2.86 (m, 5 H), 3.77 (t, $J=8.31$ Hz, 1 H), 4.36 (t, $J=6.95$ Hz, 2 H), 12.00 (s, 2 H).

Example 82

15 3-isopropyl-1-(1-methylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to procedure for Example 69A-69F, substituting ($\pm$)-1-benzyl 2-methyl pyrrolidine-1,2-dicarboxylate for (R)-1-benzyl 2-methyl pyrrolidine-1,2-dicarboxylate and isopropylhydrazine for (2,2,2-Trifluoro-ethyl)-hydrazine HCl salt. MS (DCI): m/z 315 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.51 (dd, $J=6.71, 3.66$ Hz, 6 H), 1.71 - 1.95 (m, 3 H), 2.05 - 2.25 (m, 2 H), 2.24 - 2.37 (m, 1 H), 2.41 - 2.56 (m, 2 H), 2.69 - 2.78 (m, 1 H), 2.77 - 2.89 (m, 1 H), 2.90 - 3.03 (m, 1 H), 2.96 (s, 3 H), 3.29 - 3.44 (m, 2 H), 3.77 - 3.96 (m, 1 H), 4.73 - 4.91 (m, 2 H).

25

Example 83

4-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzonitrile

The title compound was prepared according to procedure for Example 85A-85D substituting 4-cyanobenzaldehyde for pyridine-3-aldehyde. MS (DCI/NH₃) m/z 359 (M+H)⁺; ¹H NMR (500 MHz, CDCl₃): δ 1.56 - 1.59 (m, 1 H), 1.76 - 1.84 (m, 2 H), 1.91 - 1.98 (m, 1 H),

30

2.03 – 2.10 (m, 1 H), 2.32 – 2.39 (m, 2 H), 2.41 – 2.50 (m, 5 H), 2.86 (t, $J=5.34$ Hz, 2 H), 3.77 – 3.82 (m, 1 H), 5.61 (s, 2 H), 7.51 (d, $J=8.8$ Hz, 2 H), 7.56 (d, $J=8.8$ Hz, 2 H), 14.20 (br s, 1H).

Example 84

5 1-cyclobutyl-3-{4-[(isopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

Example 84A

10 4-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzaldehyde

The title compound was prepared according to procedure for Example 77A, substituting Example 83 for Example 72A. MS (DCI): m/z 362 ($M+H$)⁺.

Example 84B

15 1-cyclobutyl-3-{4-[(isopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to procedure for Example 26, substituting Example 84A for formaldehyde and isopropyl amine for Example 25. MS (DCI): m/z 405 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO-d₆): δ 1.35 (d, $J=6.71$ Hz, 6 H), 1.55 – 1.82 (m, 3 H), 1.90 (s, 1 H), 1.95 – 2.15 (m, 1 H), 2.17 – 2.51 (m, 6 H), 2.50 – 2.67 (m, 2 H), 2.68 – 2.88 (m, 1 H), 3.27 (d, $J=5.80$ Hz, 1 H), 3.71 – 3.85 (m, 1 H), 4.04 (s, 2 H), 5.46 (s, 2 H), 7.23 (d, $J=8.24$ Hz, 2 H), 7.43 (d, $J=8.24$ Hz, 2 H), 9.54 (s, 2 H).

Example 85

25 1-cyclobutyl-3-(pyridin-3-ylmethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

Example 85A

Pyridin-3-ylmethyl-hydrazine

To a solution of pyridine-3-aldehyde (5 g, 46.7 mmol) in methanol (100 mL) was added hydrazine (1.05 mL, 46.7 mmol). The reaction mixture was stirred at rt for 1 h. After purging

with nitrogen, 10% Pd/C (200 mg) was added. The reaction mixture was purged with hydrogen and stirred under hydrogen at rt overnight. Solid material was removed, and the filtrate was concentrated to give a crude Example 85A. This material was used without further purification. Yield: 5.2 g (91%). MS (DCI): m/z 124 (M+H)⁺.

5

Example 85B

5-Cyclobutyl-2-pyridin-3-ylmethyl-2H-pyrazol-3-ylamine

To a solution of 3-cyclobutyl-3-oxo-propionitrile (3.3 g, 32 mmol) in ethanol (100 mL) was added Example 85A (3.9 g, 32 mmol) at rt. The solution was heated under reflux for 4 h.

10 After cooling, the reaction mixture was concentrated and the residue was separated by flash chromatography (silica gel, 0-15% gradient MeOH in CH₂Cl₂) to give Example 85B (6.8 g). Yield: 93%. MS (DCI): m/z 229 (M+H)⁺.

Example 85C

15 5-Cyclobutyl-4-cyclohex-1-enyl-2-pyridin-3-ylmethyl-2H-pyrazol-3-ylamine

To a solution of Example 85B (4g, 17.5 mmol) in acetic acid (150 mL) was added cyclohexanone (3 mL). The solution was stirred at 50 °C for 24 h. After cooling, the reaction mixture was concentrated. The residual oil was partitioned between sodium bicarbonate solution and ethyl acetate. The organic phase was washed with water and concentrated. The residue was
20 purified by flash column chromatography to give Example 85C. Yield: 3.2 g (59 %). MS (DCI): m/z 309 (M+H)⁺.

Example 85D

1-cyclobutyl-3-(pyridin-3-ylmethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 To a solution of Example 85C (4 g, 18 mmol) in anhydrous pyridine (50 mL) was added ethyl isocyanate (3 mL) under nitrogen. The solution was heated at 50 °C for 6 h and under reflux overnight. The volatile were removed and the residual solid was purified by flash chromatography to give Example 85D. Yield: 4.2g (70%). MS (DCI): m/z 335 (M+H)⁺, ¹H NMR (300 MHz, CDCl₃) δ 1.58 (s, 1 H), 1.72 - 1.86 (m, 5 H), 1.87 - 2.17 (m, 2 H), 2.27 - 2.61

(m, 6 H), 2.84 (s, 2 H), 3.62 - 3.90 (m, 1 H), 7.20 (dd, $J=7.80$, 4.75 Hz, 1 H), 7.78 (d, $J=7.80$ Hz, 1 H), 8.48 (d, $J=3.05$ Hz, 1 H), 8.77 (s, 1 H).

Example 86

5 1-cyclobutyl-3-(piperidin-3-ylmethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

Example 85D was hydrogenated in the presence of 10% palladium on carbon overnight to provide Example 86 (1.8 g). Yield: 56%. MS (DCI): m/z 341 ($M+H$)⁺. ¹H NMR (500 MHz, DMSO-d₆) δ 0.92 - 1.28 (m, 1 H), 1.49 - 1.62 (m, 1 H), 1.62 - 1.76 (m, 7 H), 1.74 - 1.91 (m, 1
10 H), 1.91 - 2.08 (m, 1 H), 2.12 - 2.44 (m, 6 H), 2.57 - 2.89 (m, 4 H), 3.01 (d, $J=11.60$ Hz, 1 H), 3.10 - 3.29 (m, 1 H), 3.71 - 3.84 (m, 1 H), 3.98 - 4.22 (m, 2 H).

Example 87

15 1-cyclobutyl-3-{3-[(dimethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

Example 87A

3-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzonitrile

20 The title compound was prepared according to procedure for Example 85B-85D, substituting 3-hydrazinomethyl-benzonitrile for Example 85A. MS (DCI): m/z 359 ($M+H$)⁺.

Example 87B

25 3-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzaldehyde

The title compound was prepared according to procedure for Example 77A, substituting Example 87A for Example 72A. MS (DCI): m/z 362 ($M+H$)⁺.

Example 87C

1-cyclobutyl-3-{3-[(dimethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to procedure for Example 26, substituting Example 87B for formaldehyde and dimethylamine for Example 25. MS (DCI): m/z 391 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.72 - 1.85 (m, 5 H), 1.84 - 2.00 (m, 1 H), 1.96 - 2.18 (m, 1 H), 2.28 - 2.41 (m, 4 H), 2.48 (t, *J*=5.34 Hz, 2 H), 2.81 (s, 6 H), 2.91 (t, *J*=5.19 Hz, 2 H), 3.88 (t, *J*=8.54 Hz, 1 H), 4.27 (s, 2 H), 5.49 (s, 2 H), 7.11 - 7.31 (m, 1 H), 7.38 - 7.43 (m, 1 H), 7.46 (t, *J*=7.48 Hz, 1 H).

10

Example 88

1-cyclobutyl-3-[3-(hydroxymethyl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

A solution of Example 87B (50 mg, 0.14 mmol) in methylene chloride (10 ml) was treated with sodium borohydride (20 mg, 0.58 mmol) at rt for 1 h. Volatiles were removed and the residual solid was purified by HPLC (C-18, 0-100% gradient CH₃CN with 0.1% TFA in water with 0.1% TFA) to give Example 88 as TFA salt. Yield: 36 mg (72%). MS (DCI): m/z 364 (M+H)⁺; ¹H NMR (500 MHz, DMSO-d₆): δ 1.71 (d, *J*=4.27 Hz, 5 H), 1.74 - 1.88 (m, 1 H), 1.87 - 2.08 (m, 1 H), 2.16 - 2.33 (m, 4 H), 2.39 (s, 1 H), 2.43 - 2.56 (m, 2 H), 2.82 (s, 2 H), 3.64 - 3.90 (m, 1 H), 3.99 (s, 2 H), 5.40 - 5.45 (m, 2 H), 7.11 (d, *J*=7.02 Hz, 1 H), 7.25 (s, 1 H), 7.29 - 7.53 (m, 2 H), 8.17 (s, 2 H).

20

Example 89

1-cyclobutyl-3-[(1-methylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for Example 26, substituting Example 86 for 25. MS (DCI): m/z 355 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD) δ 1.15 - 1.41 (m, 1 H), 1.59 - 1.84 (m, 6 H), 1.85 - 1.96 (m, 1 H), 1.96 - 2.20 (m, 2 H), 2.24 - 2.59 (m, 7 H), 2.80 - 2.85 (m, 3 H), 2.86 - 2.98 (m, 4 H), 3.31 - 3.41 (m, 1 H), 3.48 (d, *J*=12.58 Hz, 1 H), 3.77 - 3.95 (m, 1 H), 4.08 - 4.29 (m, 2 H).

30

Example 90

ethyl (1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)acetate

The title compound was prepared according to procedure for Example 85B-85D, substituting hydrazino-acetic acid ethyl ester for Example 85A. MS (DCI): m/z 330 ($M+H$)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.27 (t, $J=7.06$ Hz, 3 H), 1.70 - 1.85 (m, 4 H), 1.85 - 1.97 (m, 1 H), 1.96 - 2.20 (m, 1 H), 2.27 - 2.45 (m, 4 H), 2.48 (t, $J=4.91$ Hz, 2 H), 2.90 (t, $J=4.91$ Hz, 2 H), 3.69 - 3.94 (m, 1 H), 4.22 (q, $J=7.06$ Hz, 2 H), 5.02 (s, 2 H).

Example 91

1-cyclobutyl-3-(2-hydroxyethyl)-3,4,6,7,8,9-hexahydro-5H-pyrrolo[2,3-c]isoquinolin-5-one

A solution of Example 90 (100 mg, 0.3 mmol) in methylene chloride (10 ml) was treated with sodium borohydride (20 mg, 0.58 mmol) at rt for 1 h. Volatiles were removed, and the residual solid was washed with methanol and dried to give Example 91 (40 mg, 47%). MS (DCI): m/z 288 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO- d_6): δ 1.65 - 1.76 (m, 4 H), 1.76 - 1.88 (m, 1 H), 1.90 - 2.05 (m, 1 H), 2.17 - 2.30 (m, 3 H), 2.32 - 2.43 (m, 2 H), 2.78 (s, 2 H), 3.70 (d, $J=4.58$ Hz, 2 H), 3.73 - 3.88 (m, 1 H), 4.15 - 4.26 (m, 2 H), 4.62 - 4.85 (m, 1 H), 11.67 (s, 2 H).

Example 92

(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)acetic acid

A solution of Example 90 (100 mg, 0.3 mmol) in a mixture of THF (5 ml) and methanol (5 ml) was treated with a 1N solution of lithium hydroxide (2 mL) at rt for 10 h. The mixture was neutralized with 1N HCl to a pH of 7. The formed solid was collected by filtration, washed with methanol and dried to give Example 92 (30 mg, 32%). MS (DCI): m/z 302 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO- d_6): δ 1.65 - 1.76 (m, 5 H), 1.75 - 1.88 (m, 1 H), 1.87 - 2.06 (m, 1 H), 2.18 - 2.34 (m, 4 H), 2.33 - 2.45 (m, 1 H), 2.80 (s, 2 H), 3.70 - 3.87 (m, 1 H), 4.93 (s, 2 H).

Example 93

1-cyclobutyl-3-[2-(1H-tetrazol-5-yl)ethyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

To a suspension of Example 78A (150 mg, 0.5 mmol) and trimethylsilyl azide (582 mg, 5.0 mmol) in toluene (20 ml) was added dibutyltin oxide (28 mg, 0.34 mmol). The reaction mixture was heated at 100 °C for 3 days. Volatiles were removed, and the residual solid was

washed with methanol and dried to give Example 93 (100 mg, 60%). MS (DCI): m/z 340 ($M+H$)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.70 - 1.85 (m, 4 H), 1.82 - 1.95 (m, 1 H), 1.94 - 2.13 (m, 1 H), 2.24 - 2.38 (m, 4 H), 2.48 (t, $J=5.03$ Hz, 2 H), 2.72 - 2.90 (m, 2 H), 3.48 (q, $J=6.92$ Hz, 2 H), 3.69 - 3.89 (m, 1 H), 4.63 (t, $J=6.87$ Hz, 2 H).

5

Example 94

1-cyclobutyl-3-{[1-(methanesulfonyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

To a solution of Example 86 (170 mg, 0.5 mmol) in pyridine (5 mL) was added
10 methanesulfonyl chloride (68 mg, 0.6 mmol). The reaction mixture was stirred at rt for 1 h and concentrated. The residual solid was washed with MeOH, and dried to give Example 94. Yield: 102 mg (49%). MS (DCI): m/z 419 ($M+H$)⁺; ¹H NMR (500 MHz, DMSO- d_6) δ 0.86 - 1.24 (m, 1 H), 1.26 - 1.57 (m, 1 H), 1.64 (s, 1 H), 1.67 - 1.82 (m, 5 H), 1.81 - 1.93 (m, 1 H), 1.92 - 2.07 (m, 2 H), 2.15 (s, 1 H), 2.22 - 2.37 (m, 5 H), 2.35 - 2.48 (m, 2 H), 2.67 - 2.80 (m, 1 H), 2.80 (s, 3
15 H), 2.81 - 2.87 (m, 3 H), 3.46 (d, $J=3.36$ Hz, 2 H), 3.81 (d, $J=8.24$ Hz, 1 H).

Example 95

1-cyclobutyl-3-[3-(1H-tetrazol-5-yl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

Example 95A

3-((1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl)benzonitrile

The title compound was prepared according to procedure for Example 85A-85D
25 substituting 3-cyanobenzaldehyde for pyridine-3-aldehyde. MS (DCI/NH₃) m/z 359 ($M+H$)⁺.

Example 95B

1-cyclobutyl-3-[3-(1H-tetrazol-5-yl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

30

To a suspension of Example 95A (150 mg) and trimethylsilyl azide (582 mg) in toluene (20 ml) was added dibutyltin oxide (28 mg). The mixture was heated at 100 °C for 3 days. Solid material was collected by filtration, washed with methanol and dried to give Example 95. Yield: 100 mg. MS (DCI/NH₃) m/z 402 ($M+H$)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.81 (s, 2 H), 1.92 (m, 1 H), 2.02 - 2.11 (m, 1 H), 2.35 - 2.44 (m, 4 H), 2.51 (m, 2 H), 2.91 (br s, 2 H), 3.84 - 3.90 (m, 1 H), 5.37 (br s, 1 H), 5.51 (s, 2 H), 5.51 (s, 2 H), 7.30 (m, 2 H), 7.97 (m, 2 H),
35

Example 96

7,9-dimethyl-1,2,3,4,6,7-hexahydro-5*H*-pyrazolo[3,4-*h*]-1,6-naphthyridin-5-one

5

Example 96A

5-chloro-N-(4-cyano-1,3-dimethyl-1*H*-pyrazol-5-yl)pentanamide

To a solution of 5-amino-1,3-dimethyl-1*H*-pyrazole-4-carbonitrile (1.59 g, 11.67 mmol) in anhydrous pyridine (15 mL) was slowly added 5-chlorovaleryl chloride (1.80 mL, 14.0 mmol) at 0 °C. After addition, the reaction mixture was allowed to warm up to rt and stir at rt overnight.

10 The reaction mixture was concentrated on a rotavapor and the residue was partitioned between ethyl acetate and brine. The organic phase was washed with brine and concentrated. The residue was separated by flash chromatography (silica gel, 50 – 80% gradient EtOAc in hexane) to provide Example 96A. Yield: 2.42 g (81%). MS (DCI): m/z 255 ($M+H$)⁺.

15

Example 96B

7,9-dimethyl-1,2,3,4,6,7-hexahydro-5*H*-pyrazolo[3,4-*h*]-1,6-naphthyridin-5-one

To a stirring solution of 2,2,6,6-tetramethylpiperidine (1.66 mL, 9.8 mmol) in anhydrous THF (20 mL) was added *n*-butyllithium (2.5 M solution in hexane, 3.92 mL, 9.8 mmol) at – 20 °C. The solution was stirred at –20 °C for 30 min, and a solution of Example 96A (1.0 g, 3.92 mmol) in 10 mL of anhydrous THF was added dropwise. The resulting dark red mixture was stirred at the same temperature for 30 min. The reaction was quenched with water, and the mixture was concentrated. The residue was partitioned between ethyl acetate and brine. The formed solid material in the bi-phase mixture was collected by filtration, washed with EtOAc and water. The obtained crude product was recrystallized from methanol to give Example 96.

25 Yield: 343 mg (40%). MS (DCI): m/z 219 ($M+H$)⁺; ¹H NMR (300 MHz, DMSO-*d*₆): δ 1.71 - 1.79 (m, 2 H), 2.33 - 2.37 (m, 5 H), 3.20 - 3.25 (m, 2 H), 3.65 (s, 3 H), 6.04 (br s, 1 H), 11.06 (br s, 1 H).

Example 97

30 1-{{cyclohexyl(methyl)amino}methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5*H*-pyrazolo[3,4-*c*]isoquinolin-5-one

The title compound was prepared according to procedure for Example 53 substituting cyclohexanone for acetone. Yield: 34%. MS (DCI/NH₃) m/z 329 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.23 - 1.32 (m, 1H), 1.38 - 1.48 (m, 2H), 1.58 - 1.69 (m, 2H), 1.71 - 1.77 (m, 1H), 1.79 - 1.84 (m, 2H), 1.85 - 1.90 (m, 2H), 1.94 - 2.01 (m, 2H), 2.13 - 2.22 (m, 2H), 2.50 (t, *J*=5.95 Hz, 2H), 2.92 (s, 3H), 2.95 - 3.03 (m, 2H), 3.43 - 3.52 (m, 1H), 3.92 (s, 3H), 4.42 (d, *J*=14.65 Hz, 1H), 4.80 (d, *J*=14.65 Hz, 1H); Anal. Calcd for C₁₉H₂₈N₄O•1.6 HCl: C, 59.00; H, 7.71; N, 14.48. Found: C, 59.07; H, 7.36; N, 14.20.

Example 98

10 3-methyl-1-[1-(2-phenylethyl)piperidin-4-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 73 substituting phenylacetaldehyde for formaldehyde. Yield: 34%. MS (DCI/NH₃) m/z 391 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.78 - 1.89 (m, 5H), 2.11 - 2.20 (m, 2H), 2.22 - 2.29 (m, 2H), 2.46 - 2.54 (m, 2H), 2.98 (q, *J*=5.59 Hz, 2H), 3.08 - 3.14 (m, 2H), 3.18 - 3.26 (m, 2H), 3.36 - 3.42 (m, 2H), 3.77 (d, *J*=12.51 Hz, 2H), 3.83 (s, 3H), 7.27 - 7.30 (m, 1H), 7.31 - 7.33 (m, 1H), 7.33 - 7.36 (m, 2H), 7.36 - 7.38 (m, 1H); Anal. Calcd for C₂₄H₃₀N₄O•2.1 HCl: C, 61.71; H, 6.93; N, 11.99. Found: C, 61.89; H, 6.75; N, 11.81.

Example 99

20 3-methyl-1-[(2S)-1-methylpiperidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 26 substituting Example 54 for Example 25. The TFA salt was treated with 1.0 M soln of HCl in ether to yield the HCl salt. Yield: 54 mg, 58%. MS (DCI/NH₃) m/z 301 (M+H)⁺; ¹H NMR (400 MHz, CD₃OD): δ 1.72 - 1.85 (m, 3H), 1.86 - 1.95 (m, 3H), 1.96 - 2.08 (m, 3H), 2.22 - 2.32 (m, 2H), 2.49 - 2.57 (m, 2H), 2.73 - 2.76 (m, 3H), 2.78 - 2.87 (m, 1H), 2.98 - 3.09 (m, 1H), 3.55 - 3.66 (m, 1H), 3.92 (s, 3H), 4.60 (d, *J*=11.66 Hz, 1H).

Example 100

30 3-methyl-1-{[methyl(propyl)amino]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 53 substituting propionaldehyde for acetone. Yield: 50%. MS (DCI/NH₃) m/z 259 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.04 (t, *J*=7.48 Hz, 3H), 1.78 - 1.92 (m, 6H), 2.50 (t, *J*=6.10 Hz, 2H), 2.88 - 2.96 (m, 2H), 2.98 (s, 3H), 3.15 - 3.26 (m, 1H), 3.33 - 3.43 (m, 1H), 3.92 (s, 3H), 4.54 (d, *J*=14.95 Hz, 1H), 4.73 (d, *J*=14.65 Hz, 1H).

Example 101

1-{[(cyclopropylmethyl)(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 53 substituting cyclopropanecarboxaldehyde for acetone. Yield: 49%. MS (DCI/NH₃) m/z 301 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 0.47 - 0.52 (m, 2H), 0.78 - 0.84 (m, 2H), 1.21 - 1.29 (m, 1H), 1.80 - 1.90 (m, 4H), 2.51 (t, J=5.95 Hz, 2H), 2.95 (t, J=5.80 Hz, 2H), 3.04 (s, 3H), 3.17 (dd, J=13.27, 7.48 Hz, 1H), 3.32 - 3.37 (m, 1H), 3.92 (s, 3H), 4.53 (d, J=14.65 Hz, 1H), 4.84 (d, J=14.65 Hz, 1H).

Example 102

3-methyl-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Examples 24B-24E and Example 25 substituting (R)-1-benzyl 2-methyl pyrrolidine-1,2-dicarboxylate for Example 24A. MS (DCI/NH₃) m/z 273 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.79 - 1.91 (m, 4H), 2.16 - 2.30 (m, 3H), 2.46 - 2.59 (m, 3H), 2.88 - 3.01 (m, 2H), 3.39 - 3.49 (m, 1H), 3.53 - 3.60 (m, 1H), 3.89 (s, 3H), 5.18 (t, J=6.90 Hz, 1H).

Example 103

3-methyl-1-[(2R)-1-methylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 26 substituting Example 102 for Example 25. The TFA salt was treated with 1.0 M solution of HCl in ether to yield the HCl salt. Yield: 33 mg (56%). MS (DCI/NH₃) m/z 287 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 1.79 - 1.91 (m, 4H), 2.10 - 2.24 (m, 2H), 2.26 - 2.34 (m, 1H), 2.49 - 2.55 (m, 2H), 2.70 - 2.76 (m, 1H), 2.80 - 2.88 (m, 1H), 2.93 - 2.96 (m, 1H), 2.96 (s, 3H), 3.33 - 3.39 (m, 1H), 3.81 - 3.87 (m, 1H), 3.91 (s, 3H), 4.98 (t, J=8.39 Hz, 1H).

Example 104

1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared according to procedure for Example 26, substituting Example 102 for Example 25, and cyclopropanecarboxaldehyde for formaldehyde. The TFA salt was treated with 1.0 M solution of HCl in ether to yield the HCl salt. Yield: 41 mg (56%). MS (DCI/NH₃) m/z 327 (M+H)⁺; ¹H NMR (500 MHz, CD₃OD): δ 0.33 - 0.44 (m, 2H), 0.63 - 0.71 (m, 2H), 1.05 - 1.16 (m, 1H), 1.77 - 1.91 (m, 4H), 2.04 - 2.13 (m, 1H), 2.19 - 2.26 (m, 1H), 2.26 - 2.35 (m, 1H), 2.47 - 2.56 (m, 2H), 2.67 - 2.76 (m, 1H), 2.78 - 2.88 (m, 1H), 2.90 - 3.01 (m, 1H), 3.09 - 3.19 (m, 2H), 3.35 - 3.43 (m, 1H), 3.91 (s, 3H), 3.96 - 4.03 (m, 1H), 5.04 (t, J=8.54 Hz, 1H).

Example 105

3-methyl-1-(1-propylpiperidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared according to procedure for Example 26 substituting Example 63 for Example 25, and substituting propionaldehyde for formaldehyde. The TFA salt was treated with 1.0 M solution of HCl in ether to yield the HCl salt. Yield: 16 mg (12%). MS (DCI/NH₃) m/z 329 (M+H)⁺; ¹H NMR (500 MHz, C₅D₅N): δ 0.78 (t, *J*=7.32 Hz, 3H), 1.67 - 1.72 (m, 2H), 1.73 - 1.77 (m, 2H), 1.79 - 1.84 (m, 1H), 1.87 - 1.93 (m, 1H), 1.94 - 2.03 (m, 2H), 2.32 (d, *J*=13.43 Hz, 1H), 2.65 - 2.78 (m, 2H), 2.84 (t, *J*=5.80 Hz, 2H), 2.94 - 3.04 (m, 2H), 3.10 - 3.18 (m, 1H), 3.27 - 3.42 (m, 2H), 3.59 (d, *J*=7.63 Hz, 1H), 3.92 (d, *J*=10.98 Hz, 1H), 3.99 (s, 10 3H), 4.43 - 4.60 (m, 1H), 5.22 (br s, 1H)

Example 106

1-(3-{{(cyclopropylmethyl)amino}methyl}phenyl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 363 (M+H)⁺.

Example 107

20 1-(3-{{[bis(cyclopropylmethyl)amino]methyl}phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 417 (M+H)⁺.

Example 108

25 1-{3-[(cyclobutylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 363 (M+H)⁺.

30

Example 109

1-{3-[(dicyclobutylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples.

MS (DCI): m/z 417 (M+H)⁺.

Example 110

5 1-{3-[(cyclopentylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples.

MS (DCI): m/z 377 (M+H)⁺.

Example 111

10 3-methyl-1-{3-[(propylamino)methyl]phenyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 351 (M+H)⁺.

15

Example 112

1-{3-[(cycloheptylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 405 (M+H)⁺.

Example 113

1-{3-[(cyclohexylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 391 (M+H)⁺.

Example 114

1-{3-[(isopropylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 351 (M+H)⁺.

5

Example 115

1-{3-[(isobutylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 365 (M+H)⁺.

Example 116

benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)azetidine-1-carboxylate

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 393 (M+H)⁺.

Example 117

3-methyl-1-(1-methylazetidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 273 (M+H)⁺.

Example 118

3-methyl-1-(1-propylazetidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 301 (M+H)⁺.

Example 119

1-(1-isopropylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 301 (M+H)⁺.

5

Example 120

1-(1-cyclopentylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 327 (M+H)⁺.

Example 121

1-(1-cyclohexylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 341 (M+H)⁺.

Example 122

1-(1-cycloheptylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 355 (M+H)⁺.

Example 123

25 1-[1-(cyclopropylmethyl)azetidin-3-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 313 (M+H)⁺.

Example 124

1-(1-isobutylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 315 (M+H)⁺.

5

Example 125

benzyl 2-(3-ethyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)pyrrolidine-1-carboxylate

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 421 (M+H)⁺.

10

Example 126

3-ethyl-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 287 (M+H)⁺.

15

Example 127

benzyl 4-[(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)methyl]benzylcarbamate

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 457 (M+H)⁺.

20

Example 128

benzyl 2-[5-oxo-3-(2,2,2-trifluoroethyl)-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl]pyrrolidine-1-carboxylate

25

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 475 (M+H)⁺.

Example 129

1-(1-cyclobutylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 341 ($M+H$)⁺.

5

Example 130

1-[1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 341 ($M+H$)⁺.

Example 131

1-(1-cyclopentylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 355 ($M+H$)⁺.

Example 132

1-(1-cyclohexylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 369 ($M+H$)⁺.

Example 133

25 1-(1-cycloheptylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 383 ($M+H$)⁺.

Example 134

3-ethyl-1-(1-isopropylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 329 (M+H)⁺.

Example 135

3-ethyl-1-[1-(1-ethylpropyl)pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 357 (M+H)⁺.

Example 136

15 1-{4-[(cyclobutylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 376 (M+H)⁺.

Example 137

20 1-{4-[(dicyclobutylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 431 (M+H)⁺.

25 Example 138

1-{4-[(cyclopentylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 391 (M+H)⁺.

Example 139

1-{4-[(cyclohexylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 405 (M+H)⁺.

Example 140

10 1-[(2R)-pyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 341 (M+H)⁺.

Example 141

15 1-[(2R)-1-isopropylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 383 (M+H)⁺.

Example 142

20 1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 395 (M+H)⁺.

Example 143

25 1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 395 (M+H)⁺.

Example 144

5 1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 409 (M+H)⁺.

Example 145

10 1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 423 (M+H)⁺.

15

Example 146

1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 437 (M+H)⁺.

Example 147

2- {5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl} ethyl ethylcarbamate

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 374 (M+H)⁺.

Example 148

2-{1-[(2R)-1-isopropylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 415 (M+H)⁺.

5

Example 149

2-{5-oxo-1-[(2R)-1-propylpyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 415 (M+H)⁺.

Example 150

2-{1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 428 (M+H)⁺.

Example 151

20 2-{1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 428 (M+H)⁺.

Example 152

25 2-{1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 442 (M+H)⁺.

Example 153

2-{1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 456 (M+H)⁺.

Example 154

2-{1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 470 (M+H)⁺.

Example 155

3-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

15

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 312 (M+H)⁺.

Example 156

3-{1-[(2R)-1-isopropylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 354 (M+H)⁺.

Example 157

3-{1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

25

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 366 (M+H)⁺.

Example 158

3-{1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 366 (M+H)⁺.

Example 159

10 3-{1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 380 (M+H)⁺.

Example 160

15 3-{1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 394 (M+H)⁺.

20 Example 161

3-{1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 408 (M+H)⁺.

25

Example 162

3-(3-hydroxybenzyl)-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 365 (M+H)⁺.

Example 163

5 3-(3-hydroxypropyl)-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 451 (M+H)⁺.

10 Example 164

3-[3-(cyclopentylamino)propyl]-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 384 (M+H)⁺.

15

Example 165

3-(3-hydroxybenzyl)-1-[(2R)-1-methylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 379 (M+H)⁺.

Example 166

3-(3-hydroxybenzyl)-1-[(2R)-1-isopropylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 407 (M+H)⁺.

Example 167

1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 419 (M+H)⁺.

5

Example 168

1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 433 (M+H)⁺.

Example 169

1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 447 (M+H)⁺.

Example 170

20 1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 461 (M+H)⁺.

Example 171

25 3-[3-(dimethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 344 (M+H)⁺.

Example 172

3-[3-(methyamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 330 (M+H)⁺.

Example 173

3-[3-(ethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 344 (M+H)⁺.

Example 174

15 3-[3-(isopropylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 358 (M+H)⁺.

Example 175

20 3-[3-(cyclopropylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 356 (M+H)⁺.

25 Example 176

3-[3-(cyclobutylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 370 (M+H)⁺.

Example 177

3-[3-(cyclohexylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 398 (M+H)⁺.

Example 178

1-pyrrolidin-2-yl-3-(3-pyrrolidin-1-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 370 (M+H)⁺.

Example 179

3-(3-piperidin-1-ylpropyl)-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 384 (M+H)⁺.

Example 180

3-(3-morpholin-4-ylpropyl)-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 386 (M+H)⁺.

25

Example 181

1-cyclobutyl-3-[3-(methylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 315 (M+H)⁺.

Example 182

5 1-cyclobutyl-3-[3-(isopropylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 343 (M+H)⁺.

Example 183

10 1-cyclobutyl-3-[3-(cyclopropylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 341 (M+H)⁺.

15

Example 184

1-cyclobutyl-3-[3-(cyclobutylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related
20 examples. MS (DCI): m/z 355 (M+H)⁺.

Example 185

1-cyclobutyl-3-[3-(cyclopentylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 369 (M+H)⁺.

Example 186

1-cyclobutyl-3-[3-(cyclohexylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 382 (M+H)⁺.

5

Example 187

1-cyclobutyl-3-(3-piperidin-1-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 369 (M+H)⁺.

10

Example 188

1-cyclobutyl-3-(3-morpholin-4-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 371 (M+H)⁺.

15

Example 189

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)propanenitrile

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 297 (M+H)⁺.

20

Example 190

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)propanal

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 300 (M+H)⁺.

25

Example 191

1-cyclobutyl-3-(3-hydroxypropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 302 (M+H)⁺.

Example 192

5 1-cyclobutyl-3-[4-(dimethylamino)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 377 (M+H)⁺.

Example 193

10 3-isopropyl-1-(1-isopropylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 343 (M+H)⁺.

15

Example 194

1-[1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 355 (M+H)⁺.

Example 195

1-(1-cyclobutylpyrrolidin-2-yl)-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 355 (M+H)⁺.

Example 196

1-(1-cyclopentylpyrrolidin-2-yl)-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 369 (M+H)⁺.

5

Example 197

1-(1-cyclohexylpyrrolidin-2-yl)-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 383 (M+H)⁺.

Example 198

3-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzonitrile

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 359 (M+H)⁺.

Example 199

1-cyclobutyl-3-{4-[(methylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 377 (M+H)⁺.

Example 200

1-cyclobutyl-3-{4-[(ethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 391 (M+H)⁺.

Example 201

1-cyclobutyl-3-{4-[(dimethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 391 (M+H)⁺.

Example 202

1-cyclobutyl-3-{4-[(cyclopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 403 (M+H)⁺.

Example 203

1-cyclobutyl-3-{4-[(cyclobutylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 417 (M+H)⁺.

Example 204

1-cyclobutyl-3-{4-[(cyclohexylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 445 (M+H)⁺.

Example 205

1-cyclobutyl-3-[4-(hydroxymethyl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 364 (M+H)⁺.

Example 206

1-cyclobutyl-3-{4-[(cyclopentylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 431 (M+H)⁺.

Example 207

1-cyclobutyl-3-{3-[(methylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 377 (M+H)⁺.

Example 208

1-cyclobutyl-3-{3-[(ethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 391 (M+H)⁺.

Example 209

1-cyclobutyl-3-{3-[(cyclopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 403 (M+H)⁺.

25

Example 210

1-cyclobutyl-3-{3-[(cyclobutylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 417 (M+H)⁺.

Example 211

5 1-cyclobutyl-3- {[1-(cyclopropylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 395 (M+H)⁺.

Example 212

10 1-cyclobutyl-3- [(1-isopropylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 383 (M+H)⁺.

15

Example 213

1-cyclobutyl-3- [(1-isobutylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 397 (M+H)⁺.

20

Example 214

1-cyclobutyl-3- [(1-cyclobutylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 395 (M+H)⁺.

Example 215

1-cyclobutyl-3-[(1-cyclopentylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 409 (M+H)⁺.

5

Example 216

1-cyclobutyl-3-[(1-cyclohexylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 423 (M+H)⁺.

Example 217

1-cyclobutyl-3-[(1-cycloheptylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 437 (M+H)⁺.

Example 218

20 1-cyclobutyl-3-[(1-tetrahydro-2H-pyran-4-ylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 425 (M+H)⁺.

Example 219

25 1-cyclobutyl-3-{[1-(pyridin-2-ylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 432 (M+H)⁺.

Example 220

1-cyclobutyl-3- {[1-(pyridin-3-ylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

5 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 432 (M+H)⁺.

Example 221

1-cyclobutyl-3- {[1-(pyridin-4-ylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

10 The title compound was prepared as TFA salt according to the procedure for related examples. MS (DCI): m/z 432 (M+H)⁺.

Example 222

3-isopropyl-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 MS (DCI): m/z 301 (M+H)⁺.

Example 223

benzyl methyl[(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)methyl]carbamate

20 The title compound was prepared according to procedure for related example. MS (DCI): m/z 381 (M+H)⁺.

Example 224

3-methyl-1-[(2S)-piperidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25 The title compound was prepared as TFA salt according to procedure for related example. MS (DCI): m/z 287 (M+H)⁺.

Example 225

1- {[cyclobutyl(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.

MS (DCI): m/z 301 (M+H)⁺.

5

Example 226

1- {[cyclopentyl(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.

10 MS (DCI): m/z 315 (M+H)⁺.

Example 227

1-[(2S)-1-(cyclopropylmethyl)piperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15 The title compound was prepared as HCl salt according to procedure for related example.

MS (DCI): m/z 341 (M+H)⁺.

Example 228

1-[(2S)-1-cyclobutylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

20

The title compound was prepared as HCl salt according to procedure for related example.

MS (DCI): m/z 341 (M+H)⁺.

Example 229

1-[(2S)-1-isobutylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

25

The title compound was prepared as HCl salt according to procedure for related example.

MS (DCI): m/z 343 (M+H)⁺.

Example 230

1-[(2S)-1-cyclopentylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.

5 MS (DCI): m/z 355 (M+H)⁺.

Example 231

benzyl (2R)-2-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)pyrrolidine-1-carboxylate

10 The title compound was prepared according to procedure for related example. MS (DCI): m/z 407 (M+H)⁺.

Example 232

3-methyl-1-[(2R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

15

The title compound was prepared as HCl salt according to procedure for related example. MS (DCI): m/z 357 (M+H)⁺.

Example 233

20 benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)piperidine-1-carboxylate

The title compound was prepared according to procedure for related example. MS (DCI): m/z 421 (M+H)⁺.

25

Example 234

benzyl 1-cyclopropyl-3-methyl-5-oxo-3,4,5,6,8,9-hexahydro-7H-pyrazolo[3,4-c]-2,7-naphthyridine-7-carboxylate

The title compound was prepared according to procedure for related example. MS (DCI): m/z 379 (M+H)⁺.

Example 235

benzyl 4-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)piperidine-
1-carboxylate

The title compound was prepared according to procedure for related example. MS (DCI):
m/z 421 (M+H)⁺.

Example 236

1-(1-isopropylpiperidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-
one

The title compound was prepared as HCl salt according to procedure for related example. MS
(DCI): m/z 329 (M+H)⁺.

Example 237

1-[1-(cyclopropylmethyl)piperidin-3-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-
c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 341 (M+H)⁺.

Example 238

1-(1-cyclobutylpiperidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-
one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 341 (M+H)⁺.

Example 239

1-(1-cyclopentylpiperidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-
5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 355 (M+H)⁺.

Example 240

5 3-methyl-1-[1-(2-phenylethyl)piperidin-3-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 391 (M+H)⁺.

Example 241

10 3-methyl-1-(1-tetrahydro-2H-pyran-4-yl)piperidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 371 (M+H)⁺.

15

Example 242

1-[1-(cyclopentylmethyl)piperidin-3-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
20 MS (DCI): m/z 369 (M+H)⁺.

Example 243

3-methyl-1-(1-propylpiperidin-4-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
25 MS (DCI): m/z 329 (M+H)⁺.

Example 244

1-(1-isopropylpiperidin-4-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 329 (M+H)⁺.

Example 245

5 1-[1-(cyclopropylmethyl)piperidin-4-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 341 (M+H)⁺.

Example 246

10 1-(1-cyclobutylpiperidin-4-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

The title compound was prepared as HCl salt according to procedure for related example.
MS (DCI): m/z 341 (M+H)⁺.

15

Example 247

1-cyclopropyl-3,6-dimethyl-4,6,7,8-tetrahydrocyclopenta[d]pyrazolo[3,4-b]pyridin-5(3H)-one
compound with 1-cyclopropyl-3,7-dimethyl-4,6,7,8-tetrahydrocyclopenta[d]pyrazolo[3,4-b]pyridin-5(3H)-one (1:1)
20

The title compound was prepared according to procedure for related example. MS (DCI):
 m/z 244 (M+H)⁺.

Example 248

25 1-[(2S)-1-isopropylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one


Example 249

1-(1-cyclobutylpyrrolidin-3-yl)-3,4-dimethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one

MS (DCI): m/z 341 (M+H)⁺.

What is claimed:

1. A compound of Formula (I)



Formula (I)

or a therapeutically acceptable salt thereof, wherein

R¹ and R² together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl or heterocycle, the cycloalkyl or heterocycle may be unsubstituted or substituted with 1, 2, or 3 substituents that are independently alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C1-C6 alkyl, alkynyl, aryl, arylalkoxycarbonyl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydroxy, hydroxyalkyl, nitro, NR^AR^B, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, or (NR^AR^B)sulfonyl;

R³ is alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C1-C6 alkyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydrogen, hydroxy, hydroxyalkyl, nitro, NR^AR^B, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, or (NR^AR^B)sulfonyl;

X is NR⁴;


R⁴ is C1-C6 alkyl, alkoxycarbonylalkyl, arylalkyl, carboxyalkyl, cyanoalkyl, cycloalkyl, cycloalkylalkyl, hydrogen, (NR^AR^B)alkyl, (NR^AR^B)carbonyl, (NR^AR^B)carbonylalkyl, -alkyl-O-C(O)-(NR^AR^B), formylalkyl, heteroarylalkyl, heterocyclealkyl, hydroxyalkyl, alkoxyalkyl, or haloalkyl; and

R^A and R^B are independently hydrogen, C1-C6 alkyl, aryl, arylalkyl, arylalkylcarbonyl, arylalkoxy, arylalkoxycarbonyl, cycloalkyl, cycloalkylalkyl, or heterocycle;

if R³ is C1-alkyl and R⁴ is C1-alkyl and the ring formed by R¹ and R² is heterocycle then the heterocycle may not be substituted with methyl or methylcarbonyl;

and if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is a cycloalkyl then the cycloalkyl must be substituted; and wherein the aryl, cycloalkyl, heterocycle and heterocyclealkyl represented by R^3 and R^4 either themselves or as part of another moiety, are independently unsubstituted or substituted with 1, 2, or 3 substituents that are independently alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, C1-C6-alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, oxo, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, $(NR^A R^B)$ sulfonyl, alkylsulfonyl, heterocycle, heterocyclealkyl, heteroaryl, or heteroarylalkyl, wherein the substituent moieties are themselves further unsubstituted.

2. A compound of Formula (I)



(I),

or a therapeutically acceptable salt thereof, wherein

R^1 and R^2 together with the atoms to which they are attached form a 5, 6, or 7 membered cycloalkyl or heterocycle, the cycloalkyl or heterocycle may be unsubstituted or substituted with a substituent that is independently heterocycle or $NR^A R^B$;

R^3 is C1-C6 alkyl, aryl, cycloalkyl, heterocycle, heteroaryl, hydrogen, or $(NR^A R^B)$ alkyl;

X is NR^4 ;

R^4 is C1-C6 alkyl, -alkyl-O-C(O)- $(NR^A R^B)$, alkoxycarbonylalkyl, arylalkyl, carboxyalkyl, cyanoalkyl, hydrogen, haloalkyl, heteroarylalkyl, heterocyclealkyl, hydroxyalkyl, formylalkyl, or $(NR^A R^B)$ alkyl;


R^A and R^B are independently hydrogen, C1-C6 alkyl, arylalkoxycarbonyl, cycloalkyl, cycloalkylalkyl, or heterocycle;

if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is heterocycle then the heterocycle may not be substituted with methyl or methylcarbonyl; and

if R^3 is C1-alkyl and R^4 is C1-alkyl and the ring formed by R^1 and R^2 is a cycloalkyl then the cycloalkyl must be substituted; and

wherein the aryl, cycloalkyl, heterocycle and heterocyclealkyl represented by R^3 and R^4 either themselves or as part of another moiety, are independently unsubstituted or substituted with 1, 2, or 3 substituents that are independently alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, C1-C6-alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, oxo, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, $(NR^A R^B)$ sulfonyl, alkylsulfonyl, heterocycle, heterocyclealkyl, heteroaryl, or heteroarylalkyl, wherein the substituent moieties are themselves further unsubstituted.

3. A compound of Formula (I)



Formula (I)

or a therapeutically acceptable salt thereof, wherein

R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl or heterocycle, the cycloalkyl or heterocycle may be unsubstituted or substituted with 1, 2, or 3 substituents that are independently alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C1-C6 alkyl, alkynyl, aryl, arylalkoxycarbonyl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydroxy, hydroxyalkyl, nitro, $NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, or $(NR^A R^B)$ sulfonyl;

R^3 is alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, C2-C6 alkyl, alkynyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, cyano, haloalkoxy, haloalkyl, halogen, heterocycle, heterocyclealkyl, heterocyclecarbonyl, heterocyclecarbonylalkyl, heterocyclesulfonyl, heteroaryl, heteroarylalkyl, hydrogen, hydroxy, hydroxyalkyl, nitro, $NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, or $(NR^A R^B)$ sulfonyl;

X is NR^4 ;

R^4 is C2-C6 alkyl, arylalkyl, cycloalkyl, cycloalkylalkyl, hydrogen, $(NR^A R^B)$ alkyl, hydroxyalkyl, alkoxyalkyl, or haloalkyl; and

R^A and R^B are independently hydrogen, alkyl, aryl, arylalkyl, arylalkylcarbonyl, arylalkoxy, arylalkoxycarbonyl, cycloalkyl, cycloalkylalkyl, or heterocycle; and

wherein the aryl, cycloalkyl, heterocycle and heterocyclealkyl represented by R^3 and R^4 either themselves or as part of another moiety, are independently unsubstituted or substituted with 1, 2, or 3 substituents that are independently alkenyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, C1-C6-alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylthio, alkylthioalkyl, alkynyl, aryl, arylalkyl, arylalkoxy, arylalkoxycarbonyl, carboxy, cyano, cycloalkyl, cycloalkylalkyl, formyl, haloalkoxy, haloalkyl, halogen, hydroxy, hydroxyalkyl, mercapto, nitro, oxo, $-NR^A R^B$, $(NR^A R^B)$ alkyl, $(NR^A R^B)$ carbonyl, $(NR^A R^B)$ carbonylalkyl, $(NR^A R^B)$ sulfonyl, alkylsulfonyl, heterocycle, heterocyclealkyl, heteroaryl, or heteroarylalkyl, wherein the substituent moieties are themselves further unsubstituted.

4. A compound according to claim 1 wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl.

5. A compound according to claim 1 wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered heterocycle.

6. A compound according to claim 1 wherein R^1 and R^2 together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl; and

R^3 is C1-C6 alkyl, aryl, cycloalkyl, heterocycle, heteroaryl, hydrogen, or $(NR^A R^B)$ alkyl.

7. A compound according to claim 1 wherein R^3 is aryl, cycloalkyl, heterocycle, or heteroaryl.

8. A compound according to claim 1 wherein R¹ and R² together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl; and
R³ is aryl, cycloalkyl, heterocycle, or heteroaryl.
9. A compound according to claim 1 wherein R¹ and R² together with the atoms to which they are attached form a 5, 6, 7, or 8 membered cycloalkyl;
R⁴ is C1-C6 alkyl; and
R³ is aryl, cycloalkyl, heterocycle, or heteroaryl.
10. A compound according to claim 1 wherein R⁴ is C1-C6 alkyl, -alkyl-O-C(O)-(NR^AR^B), alkoxycarbonylalkyl, arylalkyl, carboxyalkyl, cyanoalkyl, hydrogen, haloalkyl, heteroarylalkyl, heterocyclealkyl, hydroxyalkyl, formylalkyl, or (NR^AR^B)alkyl.
11. A compound according to claim 1 that is
9-cyclobutylamino-1,3-dimethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1,3-dimethyl-9-morpholin-4-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1,3-dimethyl-9-piperidin-1-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1,3-dimethyl-9-(4-methylpiperazin-1-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
3-(2-dimethylaminoethyl)-1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
3-methyl-1-phenyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
3-benzyl-1-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
3-methyl-1-thiophen-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1-isopropyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1-cyclopropyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;
1-cyclopropyl-3-methyl-4,6,8,9-tetrahydro-3H-7-oxa-2,3,4-triazacyclopenta[a]naphthalen-5-one;
1-(3-chlorophenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-thiophen-2-yl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one;
 1-cyclopropyl-3-methyl-4,6,7,8-tetrahydro-3H-2,3,4-triaza-*as*-indacen-5-one;
 1-cyclopropyl-3-methyl-4,6,7,8,9,10-hexahydro-3H-2,3,4-triazacyclohepta[e]inden-5-one;
 benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-1-yl)-pyrrolidine-1-carboxylate;
 3-methyl-1-pyrrolidin-3-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 3-methyl-1-(1-methylpyrrolidin-3-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 1-(1-isopropylpyrrolidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 3-methyl-1-(1-propylpyrrolidin-3-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 1-(1-cyclopropylmethylpyrrolidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 1-(1-cyclobutylpyrrolidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-3-methyl-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-3-Methyl-1-(1-methylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-3-methyl-1-(1-propyl-pyrrolidin-2-yl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-1-(1-isopropylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-1-(1-cyclopropylmethylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-1-(1-cyclobutylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 (S)-1-(1-isobutylpyrrolidin-2-yl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;
 3-methyl-1-methylaminomethyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-*c*]isoquinolin-5-one;

benzyl [4-(3-methyl-5-oxo-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-1-yl)benzyl]carbamate;

1-(4-aminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-(4-dimethylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-[4-(isopropylamino-methyl)-phenyl]-3-methyl-3,4,6,7,8,9-hexahydropyrazolo [3,4-c]isoquinolin-5-one;

1-(4-cyclohexylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-(4-cyclopentylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-(4-cycloheptylaminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-[4-(isobutylaminomethyl)phenyl]-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-4-(cyclobutylaminomethyl)phenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(cyclopropylmethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(biscyclopropylmethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-(4-propylaminomethylphenyl)-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-dimethylaminomethyl-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-[(isopropylmethylamino)methyl]-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

(S)-3-methyl-1-piperidin-2-yl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-{4-[(2-methyltetrahydrofuran-3-ylamino)methyl]phenyl}-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-(3-aminomethylphenyl)-3-methyl-3,4,6,7,8,9-hexahydropyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(dimethylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[(2S)-1-propylpiperidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-azetidin-3-yl-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclobutylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[4-(aminomethyl)benzyl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[(2R)-1-(2-phenylethyl)pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-piperidin-3-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-piperidin-4-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclopropyl-3,7-dimethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclopropyl-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]-2,7-naphthyridin-5-one;

3-ethyl-1-(1-methylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(dimethylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-methylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-(1-methylpiperidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

2-{1-[(2R)-1-methylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

3-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanamide;

3-methyl-1-(1-methylpiperidin-4-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-{1-[(2R)-1-methylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-[2-(dimethylamino)benzyl]-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(dimethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(ethylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-(3-pyrrolidin-1-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(dimethylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)propanoic acid;

3-isopropyl-1-(1-methylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

4-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzonitrile;

1-cyclobutyl-3-{4-[(isopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-(pyridin-3-ylmethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-(piperidin-3-ylmethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{3-[(dimethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(hydroxymethyl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-methylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

ethyl (1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)acetate;

1-cyclobutyl-3-(2-hydroxyethyl)-3,4,6,7,8,9-hexahydro-5H-pyrrolo[2,3-c]isoquinolin-5-one;

(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)acetic acid;

1-cyclobutyl-3-[2-(1H-tetrazol-5-yl)ethyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{[1-(methylsulfonyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(1H-tetrazol-5-yl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

7,9-dimethyl-1,2,3,4,6,7-hexahydro-5H-pyrazolo[3,4-*h*]-1,6-naphthyridin-5-one;

1-{[cyclohexyl(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[1-(2-phenylethyl)piperidin-4-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[(2S)-1-methylpiperidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-{[methyl(propyl)amino]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{[(cyclopropylmethyl)(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[(2R)-1-methylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-(1-propylpiperidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(3-{[(cyclopropylmethyl)amino]methyl}phenyl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(3-{[bis(cyclopropylmethyl)amino]methyl}phenyl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(cyclobutylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(dicyclobutylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(cyclopentylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-{3-[(propylamino)methyl]phenyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(cycloheptylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(cyclohexylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(isopropylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{3-[(isobutylamino)methyl]phenyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)azetidine-1-carboxylate;

3-methyl-1-(1-methylazetidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-(1-propylazetidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-isopropylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclopentylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclohexylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cycloheptylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[1-(cyclopropylmethyl)azetidin-3-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-isobutylazetidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

benzyl 2-(3-ethyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)pyrrolidine-1-carboxylate;

3-ethyl-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

benzyl 4-[(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)methyl]benzylcarbamate;

benzyl 2-[5-oxo-3-(2,2,2-trifluoroethyl)-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl]pyrrolidine-1-carboxylate;

1-(1-cyclobutylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclopentylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclohexylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cycloheptylpyrrolidin-2-yl)-3-ethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-ethyl-1-(1-isopropylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-ethyl-1-[1-(1-ethylpropyl)pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(cyclobutylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(dicyclobutylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(cyclopentylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{4-[(cyclohexylamino)methyl]benzyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-pyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-isopropylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-3-(2,2,2-trifluoroethyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

2-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{1-[(2R)-1-isopropylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{5-oxo-1-[(2R)-1-propylpyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

2-{1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}ethyl ethylcarbamate;

3-{5-oxo-1-[(2R)-pyrrolidin-2-yl]-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-{1-[(2R)-1-isopropylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-{1-[(2R)-1-cyclobutylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-{1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-{1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-{1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-{1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl}propanenitrile;

3-(3-hydroxybenzyl)-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(3-hydroxypropyl)-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(cyclopentylamino)propyl]-1-[(2R)-pyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(3-hydroxybenzyl)-1-[(2R)-1-methylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(3-hydroxybenzyl)-1-[(2R)-1-isopropylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cyclopentylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cyclohexylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2R)-1-cycloheptylpyrrolidin-2-yl]-3-(3-hydroxybenzyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(dimethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(methylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(ethylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(isopropylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(cyclopropylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(cyclobutylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[3-(cyclohexylamino)propyl]-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-pyrrolidin-2-yl-3-(3-pyrrolidin-1-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(3-piperidin-1-ylpropyl)-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(3-morpholin-4-ylpropyl)-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(methylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(isopropylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(cyclopropylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(cyclobutylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(cyclopentylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[3-(cyclohexylamino)propyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-(3-piperidin-1-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-(3-morpholin-4-ylpropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)propanenitrile;

3-(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)propanal;

1-cyclobutyl-3-(3-hydroxypropyl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[4-(dimethylamino)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-isopropyl-1-(1-isopropylpyrrolidin-2-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[1-(cyclopropylmethyl)pyrrolidin-2-yl]-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclobutylpyrrolidin-2-yl)-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclopentylpyrrolidin-2-yl)-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclohexylpyrrolidin-2-yl)-3-isopropyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-[(1-cyclobutyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-3-yl)methyl]benzonitrile;

1-cyclobutyl-3-{4-[(methylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{4-[(ethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{4-[(dimethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{4-[(cyclopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{4-[(cyclobutylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{4-[(cyclohexylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[4-(hydroxymethyl)benzyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{4-[(cyclopentylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{3-[(methylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{3-[(ethylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{3-[(cyclopropylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{3-[(cyclobutylamino)methyl]benzyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{[1-(cyclopropylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-isopropylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-isobutylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-cyclobutylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-cyclopentylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-cyclohexylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-cycloheptylpiperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-[(1-tetrahydro-2H-pyran-4-yl)piperidin-3-yl)methyl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{[1-(pyridin-2-ylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{[1-(pyridin-3-ylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclobutyl-3-{[1-(pyridin-4-ylmethyl)piperidin-3-yl]methyl}-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-isopropyl-1-pyrrolidin-2-yl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

benzyl methyl[(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)methyl]carbamate;

3-methyl-1-[(2S)-piperidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{[cyclobutyl(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-{[cyclopentyl(methyl)amino]methyl}-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2S)-1-(cyclopropylmethyl)piperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2S)-1-cyclobutylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2S)-1-isobutylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[(2S)-1-cyclopentylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

benzyl (2R)-2-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)pyrrolidine-1-carboxylate;

3-methyl-1-[(2R)-1-tetrahydro-2H-pyran-4-ylpyrrolidin-2-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

benzyl 3-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)piperidine-1-carboxylate;

benzyl 1-cyclopropyl-3-methyl-5-oxo-3,4,5,6,8,9-hexahydro-7H-pyrazolo[3,4-c]-2,7-naphthyridine-7-carboxylate;

benzyl 4-(3-methyl-5-oxo-4,5,6,7,8,9-hexahydro-3H-pyrazolo[3,4-c]isoquinolin-1-yl)piperidine-1-carboxylate;

1-(1-isopropylpiperidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[1-(cyclopropylmethyl)piperidin-3-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclobutylpiperidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclopentylpiperidin-3-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-[1-(2-phenylethyl)piperidin-3-yl]-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-(1-tetrahydro-2H-pyran-4-ylpiperidin-3-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[1-(cyclopentylmethyl)piperidin-3-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

3-methyl-1-(1-propylpiperidin-4-yl)-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-isopropylpiperidin-4-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-[1-(cyclopropylmethyl)piperidin-4-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-(1-cyclobutylpiperidin-4-yl)-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

1-cyclopropyl-3,6-dimethyl-4,6,7,8-tetrahydrocyclopenta[d]pyrazolo[3,4-b]pyridin-5(3H)-one compound with 1-cyclopropyl-3,7-dimethyl-4,6,7,8-tetrahydrocyclopenta[d]pyrazolo[3,4-b]pyridin-5(3H)-one (1:1);

1-[(2S)-1-isopropylpiperidin-2-yl]-3-methyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one; or

1-(1-cyclobutylpyrrolidin-3-yl)-3,4-dimethyl-3,4,6,7,8,9-hexahydro-5H-pyrazolo[3,4-c]isoquinolin-5-one;

or a pharmaceutically acceptable salt thereof.

12. A pharmaceutical composition comprising a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof, in combination with a therapeutically acceptable carrier.


13. Use of a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof for treating inflammation in a mammal in recognized need of such treatment.

14. Use of a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof for the manufacture of a medicament for treating inflammation in a mammal in recognized need of such treatment.

15. Use of a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof for treating sepsis in a mammal in recognized need of such treatment.

16. Use of a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof for the manufacture of a medicament for treating sepsis in a mammal in recognized need of such treatment.

17. Use of a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof for treating cancer in a mammal in recognized need of such treatment.
18. Use of a compound of Formula (I) according to claim 1 or a therapeutically acceptable salt thereof for the manufacture of a medicament for treating cancer in a mammal in recognized need of such treatment.



Formula (I)