



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2095926 C 2002/07/16

(11)(21) **2 095 926**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1992/09/24
(87) Date publication PCT/PCT Publication Date: 1993/04/01
(45) Date de délivrance/Issue Date: 2002/07/16
(85) Entrée phase nationale/National Entry: 1993/05/10
(86) N° demande PCT/PCT Application No.: GB 1992/001759
(87) N° publication PCT/PCT Publication No.: 1993/006231
(30) Priorité/Priority: 1991/09/26 (9120467.7) GB

(51) Cl.Int.⁵/Int.Cl.⁵ C12N 15/13, A61K 47/48, A61K 39/395,
C07K 16/18

(72) Inventeurs/Inventors:
Adair, John Robert, GB;
Owens, Raymond John, GB;
Baker, Terence Seward, GB;
Lyons, Alan Howard, GB;
Hamann, Philip R., US;
Menendez, Ana T., US;
Hinman, Lois M., US

(73) Propriétaire/Owner:
CELLTECH THERAPEUTICS LIMITED, GB

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : ANTICORPS HUMANISES CONTRE LES GLOBULES DE MATIERE GRASSE DU LAIT
(54) Title: ANTI-HUMAN MILK FAT GLOBULE HUMANISED ANTIBODIES

(57) **Abrégé/Abstract:**

Humanised antibody molecules (HAMs) are described having specificity for human milk fat globate and having an antigen binding site wherein at least one of the complementarity determining regions (CDRs) of the variable domains is derived from the mouse monoclonal antibody CTMO1 and the remaining immunoglobulin-derived parts of the HAM are derived from a human immunoglobulin. The HAMs maybe chimeric humanised antibodies or CDR-grafted humanised antibodies and are preferably produced by recombinant DNA techniques. The HAMs may be conjugated to an effector or reporter molecule, particularly methyltrithio anti-tumour agents, and are useful for in vivo diagnosis and therapy.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12P 21/08, C12N 15/13 A61K 47/48, 39/395	A1	(11) International Publication Number: WO 93/06231 (43) International Publication Date: 1 April 1993 (01.04.93)
(21) International Application Number: PCT/GB92/01759 (22) International Filing Date: 24 September 1992 (24.09.92) (30) Priority data: 9120467.7 26 September 1991 (26.09.91) GB (71) Applicant: CELLTECH LIMITED [GB/GB]; 216 Bath Road, Slough, Berkshire SL1 4EN (GB). (72) Inventors: ADAIR, John, Robert ; 23 George Road, Stokenchurch, High Wycombe, Bucks. HP14 3RB (GB). OWENS, Raymond, John ; 23 Hamilton Avenue, Henley-on-Thames, Oxon. RG9 1SH (GB). BAKER, Terence, Seward ; The Mages, 4 Garson Lane, Wraysbury, Staines, Middlesex TW19 5JF (GB). LYONS, Alan, Howard ; 15 Beaumont Close, Woodlands Park, Maidenhead, Berkshire SL6 3XN (GB). HAMANN, Philip, R. ; 9 Alice Street, Garnerville, NY 10923 (US). MENENDEZ, Ana, T. ; 2 Country Club Lane, Monsey, New York, NY 10951 (US). HINMAN, Lois, M. ; 126 Harwood Avenue, North Tarrytown, NY 10591 (US).		(74) Agent: MERCER, Christopher, Paul; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB). (81) Designated States: AU, CA, CS, FI, HU, JP, KR, NO. Published <i>With international search report.</i>
(54) Title: ANTI-HUMAN MILK FAT GLOBULE HUMANISED ANTIBODIES (57) Abstract Humanised antibody molecules (HAMs) are described having specificity for human milk fat globule and having an antigen binding site wherein at least one of the complementarity determining regions (CDRs) of the variable domains is derived from the mouse monoclonal antibody CTMO1 and the remaining immunoglobulin-derived parts of the HAM are derived from a human immunoglobulin. The HAMs may be chimeric humanised antibodies or CDR-grafted humanised antibodies and are preferably produced by recombinant DNA techniques. The HAMs may be conjugated to an effector or reporter molecule, particularly methyltrithio anti-tumour agents, and are useful for <i>in vivo</i> diagnosis and therapy.		

2095926

ANTI-HUMAN MILK FAT GLOBULE HUMANISED ANTIBODIES

FIELD OF THE INVENTION

The present invention relates to humanised antibody molecules (HAMs) having specificity for human milk fat globule (HMFG) and to processes for their production using recombinant DNA technology.

BACKGROUND TO THE INVENTION

The term "humanised antibody molecule" (HAM) is used to describe a molecule having an antigen binding site derived from an immunoglobulin from a non-human species, the remaining immunoglobulin-derived parts of the molecule being derived from a human immunoglobulin. The antigen binding site may comprise: either a complete variable domain from the non-human immunoglobulin fused onto one or more human constant domains; or one or more of the complementarity determining regions (CDRs) grafted onto appropriate human framework regions in the variable domain. The abbreviation "MAb" is used to indicate a monoclonal antibody.

In the description, reference is made to publications by number. These numbers are placed in square brackets []. The publications are listed in numerical order at the end of the description.

Natural immunoglobulins have been known for many years, as have the various fragments thereof, such as the Fab, Fab' (Fab')₂ and Fc fragments, which can be derived by enzymatic cleavage. Natural immunoglobulins comprise a generally Y-shaped molecule having an antigen-binding site towards the outer end of each arm. The remainder of the structure, and particularly the stem of the Y, mediates the effector functions associated with immunoglobulins.

Natural immunoglobulins have been used in assay, diagnosis and, to a more limited extent, therapy. However, such uses, especially in therapy, have been hindered by the polyclonal nature of natural

immunoglobulins. A significant step towards the realisation of the potential of immunoglobulins as therapeutic agents was the discovery of procedures for the production of monoclonal antibodies of defined specificity [1]. However, most MAbs are produced by hybridomas which are fusions of rodent spleen cells with rodent myeloma cells. The resultant MAbs are therefore essentially rodent proteins. There are few reports of the production of human MAbs.

Since most available MAbs are of rodent origin, they are naturally antigenic in humans and thus can give rise to an undesirable immune response termed the HAMA (Human Anti-Mouse Antibody) response. Therefore, the use of rodent MAbs as therapeutic agents in humans is inherently limited by the fact that the human subject will mount an immunological response to the MAb and will either remove it entirely or at least reduce its effectiveness.

Therefore proposals have been made for making non-human MAbs less antigenic in humans. Such techniques can be generically termed "humanisation" techniques. These techniques generally involve the use of recombinant DNA technology to manipulate DNA sequences encoding the polypeptide chains of the antibody molecule.

Early methods for humanising MAbs related to production of chimeric antibodies in which an antigen binding site comprising the complete variable domains of one antibody are fused to constant domains derived from a second antibody. Methods for carrying out such chimerisation procedures are described in EP-A-0 120 694 (Celltech Limited), European Patent Application EP-A-0 125 023 City of Hope and published on November 14, 1984 (Genentech Inc.), EP-A-0 171 496 (Res. Dev. Corp. Japan), European Patent Application EP-A-0 173 494 published on February 19, 1986 (Stanford University) and European Patent Application EP-A-0 194 276 published on September 17, 1986 (Celltech Limited).

EP-A-0 194 276 discloses a process for preparing an antibody molecule having the variable domains from a mouse MAb and the constant domains from a human immunoglobulin. It also describes the production of an antibody molecule comprising the variable domains of a mouse MAb, the CH1 and CL domains of a human immunoglobulin and a non-immunoglobulin-derived protein in place of the Fc portion of the human immunoglobulin.

Subsequently, a number of further patent applications have been published relating to chimeric antibodies, including tumour specific chimeric antibodies. Among these applications are WC-A-87/02671 published on May 7, 1987 (Int. Gen. Eng. Inc.), European Patent Application EP-A-0 256 654 published on February 24, 1988 (Centocor), European Patent Application EP-A-0 266 663 published on May 11, 1988 (Int. Gen. Eng. Inc. & Oncogen), PCT Application WO-A-89/00999 published on February 9, 1989 (Int. Gen. Eng. Inc.) and European Patent Application EP-A-0 332 424 published on September 13, 1989 (Hybritech Inc.).

Such humanised chimeric antibodies, however, still contain a significant proportion of non-human amino acid sequence, i.e. the complete variable domains. Thus, such humanised antibodies may elicit some HAMA response, particularly if administered over a prolonged period [2].

In an alternative approach, described in European Patent Application EP-A-0 239 400 published on September 30, 1987 (Winter), the complementarity determining regions (CDRs) of a mouse MAb have been grafted onto the framework regions of the variable domains of a human immunoglobulin by site directed mutagenesis using long oligonucleotides. Such CDR-grafted humanised antibodies are less likely to give rise to a HAMA response than humanised chimeric antibodies in view of the lower proportion of non-human amino acid sequence which they contain. There are three CDRs (CDR1, CDR2 and CDR3) in each of the heavy and light chain variable domains.

The earliest work on CDR-grafted humanised MAbs was carried out on a MAb recognising the synthetic antigen NP or NIP. However, subsequently, examples in which a mouse MAb recognising lysozyme and a rat MAb recognising an antigen on human T cells respectively were humanised have been described [3,4]. The preparation of the CDR-grafted antibody to the antigen on human T cells is also described in PCT Application WO-A-89/07452 published on August 16, 1989 (Medical Research Council). Recently the preparation of a humanised CDR-grafted antibody that binds to the interleukin 2 receptor has been described [5]. Further examples of humanised CDR-grated antibodies having specificity for anti-viral [6,7], anti-tumour [8] and anti-T cell [9 and European Patent Application EP-A-0 403 156 published on December 19, 1990 (Genzyme Corp.)] antigens have been described more recently.

Our copending PCT Patent Specification No. WO-A-91/09967 published on July 11, 1991 relates to the CDR grafting of antibodies in general.

It has been widely suggested that immunoglobulins, and in particular MAbs, could potentially be very useful in the diagnosis and treatment of cancer [10,11]. There has therefore been much activity in trying to produce immunoglobulins or MAbs directed against tumour-specific antigens. So far, over one hundred MAbs directed against a variety of human carcinomas have been used in various aspects of tumour diagnosis or treatment [12].

There have been a number of papers published concerning the production of chimeric monoclonal antibodies recognizing cell surface antigens. For instance, genetically engineered murine/human chimeric antibodies which retain specificity for tumour-associated antigens have been described [13 and PCT Application WO-A-89/01783 published on March 9, 1989

(Celltech)]. Also, a recombinant murine/human chimeric monoclonal antibody specific for common acute lymphocytic leukaemia antigen has been described [14].

SUMMARY OF THE INVENTION

We have now prepared humanized antibodies to human milk fat globule (HMFG) derived from the anti-HMFG mouse MAb CTMO1 [15].

According to the present invention, there is provided a humanised antibody molecule (HAM) having specificity for human milk fat globule (HMFG) and having an antigen binding site wherein at least one of the complementarity determining regions (CDRs) of the variable domain is derived from the mouse monoclonal antibody CTMO1 (CTMO1 MAb) and the remaining immunoglobulin-derived parts of the HAM are derived from a human immunoglobulin or an analogue thereof.

In accordance with one aspect of the present invention there is provided an antibody molecule having specificity for human milk fat globule (HMFG) comprising a composite heavy chain and a complementary light chain wherein: the variable domain of said composite heavy chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof; amino acid residues 2, 26 to 35, 37, 50 to 65, 71, 73, 95 to 105, and 107 (according to the Kabat numbering system) at least in said heavy chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 1); and the remaining immunoglobulin derived parts of the heavy chain are derived from a human immunoglobulin or an analogue thereof.

In accordance with another aspect of the present invention there is provided a process for producing the aforementioned antibody molecules.

In accordance with yet another aspect of the present invention there is provided a conjugate molecule comprising the aforementioned antibody molecule conjugated to an effector molecule or a reporter molecule.

In accordance with still yet another aspect of the present invention there is provided a pharmaceutical composition comprising an antibody molecule of the invention or a conjugate molecule of the invention.

In accordance with still yet another aspect of the present invention there is provided a diagnostic aid comprising an antibody molecule of the invention or a conjugate molecule of the invention.

In accordance with still yet another aspect of the present invention there is provided a DNA sequence encoding an antibody molecule of the invention.

The HAM may comprise a chimeric humanized antibody or a CDR-grafted humanised antibody. When the HAM comprises a CDR-grafted humanized antibody, each heavy or light chain variable domain may comprise only one or two CTMO1-derived CDRs. Preferably, however, all three heavy and light chain CDRs are derived from CTMO1.

The CTMO1 MAb is a mouse MAb of the type IgG1-kappa raised against the membrane-associated antigen of HMFG and has been extensively studied [15]. The CTMO1 MAb has been shown to recognize breast, ovarian and non-small cell lung cancers. It has been shown to internalise rapidly into target cells. Conjugates of CTMO1 and calichaemicin display highly specific cytotoxicity against appropriate cell lines, (see United States Patent No. 5053394 published on October 1, 1991 (American Cyanamid Co.)).

High levels of the antigen recognised by the CTMO1 MAb have been detected circulating in the blood of patients suffering from breast cancer. This may have a deleterious effect on pharmacokinetics and tumour localisation *in vivo*. However, circulating antigen levels in the blood of patients suffering from ovarian cancer are lower than those in breast cancer patients. It is therefore believed that the HAM of the present invention will be of particular use in the treatment of ovarian cancer.

It is believed that the CTMO1 MAb recognises the polymorphic epithelial mucin (PEM) of HMFG. Thus, preferably, the present invention provides a HAM which recognises the PEM of HMFG.

Surprisingly, it has been found that humanising the CTMO1 MAb does not substantially adversely affect its binding activity or internalisation, and can create, particularly by CDR grafting, a HAM which has better binding and internalisation characteristics than the murine antibody (see Table 1 hereinafter). This produces a HAM which is of use in both therapy and diagnosis of certain human carcinomas, for example carcinomas of ovary, breast, uterus and lung.

Preferably, the HAM of the present invention is produced by recombinant DNA technology.

The HAM of the present invention may comprise: a complete antibody molecule, having full length heavy and light chains; a fragment thereof,

2095926

such as an Fab, Fab', (Fab')₂ or Fv fragment; a single chain antibody fragment, e.g. a single chain Fv; a light chain or heavy chain monomer or dimer; or a fragment or analogue of any of these or any other molecule with the same specificity as the CTMO1 MAb.

The remaining non-CTMO1 immunoglobulin-derived parts of the HAM may be derived from a suitable human immunoglobulin. For instance, when the HAM is a CDR-grafted HAM, appropriate variable region framework sequences may be used having regard to the class or type of the CTMO1 donor antibody from which the antigen binding regions are derived. Preferably, the type of human framework used is of the same or similar class or type as the donor antibody (CTMO1 is IgG1-kappa). Advantageously, the framework is chosen to maximise or optimise homology with the donor antibody sequence, particularly at positions spatially close to or adjacent the CDRs. Examples of human frameworks which may be used to construct CDR-grafted HAMs are LAY, POM, TUR, TEI, KOL, NEWM, REI and EU [16]. KOL and NEWM are suitable for heavy chain construction. REI is suitable for light chain construction. EU is particularly suitable for both heavy chain and light chain construction. Preferably, the EU framework is used as the human framework for both heavy and light chain variable domains in view of its high level of homology with the CTMO1 MAb.

The light or heavy chain variable regions of the HAM may be fused to human light or heavy chain constant domains as appropriate, (the term "heavy chain constant domains" as used herein are to be understood to include hinge regions unless specified otherwise). The human constant domains of the HAM, where present, may be selected having regard to the proposed function of the antibody, in particular the effector functions which may be required. For example, the heavy chain constant domains fused to the heavy chain variable region may be human IgA, IgG or IgM domains. Preferably human IgG domains are used. IgG1 and IgG3 isotype domains may be used when the HAM is intended for therapeutic purposes and antibody effector functions are required. Alternatively, IgG2 and IgG4 isotype domains may be used when the HAM is intended for purposes for which antibody effector functions are not required, e.g. for imaging, diagnostic

or cytotoxic targeting purposes. Light chain human constant domains which may be fused to the light chain variable region include human Lambda or, especially, human Kappa chains.

Analogues of human constant domains may alternatively be advantageously used. These include those constant domains containing one or more additional amino acids than the corresponding human domain, or those constant domains wherein one or more existing amino acids of the corresponding human domain has been deleted or altered. Such domains may be obtained, for example, by oligonucleotide directed mutagenesis. In the present invention, a particularly useful analogue of a heavy chain constant domain is an IgG4 constant domain in which a serine residue at position 241 of the corresponding naturally occurring human domain has been altered to a proline residue.

The remainder of the HAM need not comprise only protein sequences from human immunoglobulins. For instance, a gene may be constructed in which a DNA sequence encoding part of a human immunoglobulin chain is fused to a DNA sequence encoding the amino acid sequences of a polypeptide effector or reporter molecule.

According to a second aspect of the present invention, there is provided a process for producing the HAM of the first aspect of the invention, which process comprises:

- (a) producing in an expression vector an operon having a DNA sequence which encodes an antibody heavy or light chain comprising a variable domain wherein at least one of the CDRs of the variable domain is derived from the CTMO1 MAb and the remaining immunoglobulin-derived parts of the antibody chain are derived from a human immunoglobulin;
- (b) producing in an expression vector an operon having a DNA sequence which encodes a complementary antibody light or heavy chain comprising a variable domain wherein at least one of the CDRs of the variable domain is derived from the

2095926

CTMO1 MAb and the remaining immunoglobulin-derived parts of the antibody chain are derived from a human immunoglobulin;

- (c) transfecting a host cell with both operons; and
- (d) culturing the transfected cell line to produce the HAM.

The cell line may be transfected with two vectors, the first vector containing the operon encoding the light chain-derived polypeptide and the second vector containing the operon encoding the heavy chain-derived polypeptide. Preferably, the vectors are identical except in so far as the coding sequences and selectable markers are concerned so as to ensure as far as possible that each polypeptide chain is equally expressed.

Alternatively, a single vector may be used, the vector including the operons encoding both light chain- and heavy chain-derived polypeptides.

In further aspects, the invention also includes DNA sequences coding for the heavy and light chains of the HAM of the present invention, cloning and expression vectors containing these DNA sequences, host cells transformed with these DNA sequences and processes for producing the heavy or light chains and antibody molecules comprising expressing these DNA sequences in a transformed host cell.

The general methods by which the vectors may be constructed, transfection methods and culture methods are well known per se [17, 18].

The DNA sequences which encode the CTMO1 heavy and light chain variable domain amino acid sequences (and the corresponding deduced amino acid sequences) are given hereinafter in the sequence listing as Sequence ID No. 1 and Sequence ID No. 2 respectively.

DNA coding for human immunoglobulin sequences may be obtained in any appropriate way. For example, amino acid sequences of preferred human acceptor frameworks, such as LAY, POM, KOL, REI, EU, TUR, TEI and NEWM, are widely available to workers in the art. Corresponding DNA sequences which code for these amino acid sequences may be inferred or deduced by reverse application of the genetic code. Similarly, the amino acid sequences of human constant region domains are well known and DNA sequences which code for them may be readily deduced.

The standard techniques of molecular biology may be used to prepare DNA sequences coding for CDR-grafted products. Desired DNA sequences may be synthesised completely or in part using oligonucleotide synthesis techniques. Site-directed mutagenesis and polymerase chain reaction (PCR) techniques may be used as appropriate. For example, oligonucleotide directed synthesis [19] may be used. Also, oligonucleotide directed mutagenesis of a pre-existing variable domain region [3, 4] may be used. Enzymatic filling-in of gapped oligonucleotides using T4 DNA polymerase [5] may be used.

Any suitable host cell/vector system may be used for expression of the DNA sequences coding for the chimeric or CDR-grafted heavy and light chains. Bacterial, e.g. *E. coli*, and other microbial systems may be used, in particular for expression of antibody fragments, e.g. Fv, Fab and Fab' fragments and single chain antibody fragments, e.g. single chain Fvs. Eucaryotic, e.g. mammalian host cell, expression systems may be used for production of larger chimeric or CDR-grafted antibody products, including complete antibody molecules. Suitable mammalian host cells include CHO cells and myeloma or hybridoma cell lines, for example NSO cells.

In a further aspect of the invention we provide a conjugate molecule comprising a HAM conjugated to an effector or reporter molecule. Thus for example the HAM of the present invention may have attached to it an effector molecule such as a cytotoxic or cytostatic agent, or a reporter group, for example an atom or molecule such as a radionuclide, or complexed radionuclide capable of being detected

while inside the human body. For instance, the HAM may have an organic group, such as a macrocycle, capable of binding a metal atom, or a toxin, such as ricin, or an anti-tumour agent as hereinafter defined, attached to it by a covalent bridging structure. Alternatively, the procedures of recombinant DNA technology may be used to produce a HAM in which the Fc fragment, CH2 or CH3 domain of a complete molecule has been replaced by or has attached thereto by peptide linkage a functional non-immunoglobulin protein, such as an enzyme or toxin molecule.

A particularly useful conjugate molecule according to this aspect of the invention is a HAM conjugated to a methyltrithio anti-tumour agent. Particular methyltrithio anti-tumour agents include the disulphide analogues of the α_1 , α_2 , α_3 , α_4 , β_1 , β_2 , γ_1 , δ_1 and pseudoaglycone components of the LL-E33288 complex and derivatives thereof, as well as the disulphide analogues of BBM-1675, FR-900405, FR-900406, PD 114759, PD 115028, CL-1577A, CL-1577B, CL-1577D, CL-1577E and CL 1724 antitumour antibiotics and derivatives thereof.

The family of antibacterial and antitumor agents, known collectively as the LL-E33288 complex are described and claimed in U.S. Pat. No. 4,970,198 published on November 13, 1990 (American Cyanamid Co.) and are used to prepare the disulphide antitumour agents which are some of the starting materials for the conjugate molecules of the invention.

U.S. Patent 4,970,198 describes the LL-E33288 complex, the components thereof, namely, LL-E33288 α_1^{Br} , LL-E33288 α_1^{I} , LL-E33288 α_2^{Br} , LL-E33288 α_2^{I} , LL-E33288 α_3^{Br} , LL-E33288 α_3^{I} , LL-E33288 α_4^{Br} , LL-E33288 β_1^{Br} , LL-E33288 β_1^{I} , LL-E33288 β_2^{Br} , LL-E33288 β_2^{I} , LL-E33288 δ_1^{Br} , LL-E33288 δ_1^{I} , and methods for their production by aerobic fermentation utilizing a new strain of *Micromonospora echinospora ssp calichensis* or natural or derived mutants thereof. U.S. Pat. No. 4,970,198 also discloses

proposed structures for some of the above-named components.

Additional members of the LL-E33288 complex (the calicheamicins) are described and claimed in U.S. Pat. No. 4,939,244 published on July 3, 1990 (American Cyanamid Co.) and are likewise useful for preparing the conjugate molecules of the invention. This patent also describes the LL-E33288 bromo- and iodo-pseudoaglycones of the series, which have been prepared by chemical means. The patent also describes dihydro derivatives accessible from all the above-named antitumor antibiotics through sodium borohydride reduction of the ketone at C₁₁ to a hydroxyl group.

Still other members of the LL-E33288 family of antitumour antibiotics are described and claimed in U.S. Patent 5,079,233 published on January 7, 1992 (American Cyanamid Co.), and also are useful for preparing additional conjugate molecules of the invention. This patent describes N-acyl derivatives of several members of the LL-E33288 complex which have been prepared by chemical means.

Other antibiotics are useful to prepare conjugate molecules of the invention, namely:

- 1) Esperamicin BBM-1675, [M. Konishi, et. al., J. Antibiotics, 38, 1605 (1985); M. Konishi, et. al., U.K. Patent Specification 2,141,425A, and U.S. Pat. No. 4,675,187 published June 23, 1987 (M. Konishi et al.)].
- 2) Antitumour antibiotics, FR-900405 and FR-900406. [M. Iwami, et. al., J. Antibiotics, 38, 835 (1985), S. Kiyoto, et. al., J. Antibiotics, 38 340 (1985)].
- 3) PD 114759 and PD 115028, [R.H. Bunge, et. al., J. Antibiotics, 37, 1566 (1984) U.S. Pat. No. 4,554,162 published November 19, 1985 (C.A. Young et al.) D.W.

Fry et. al., Investigational New Drugs, 4,3 (1986)].

- 4) Antibiotic complex CL-1577A, CL-1566B produced by *Streptomyces* asp. ATCC 39363. U.S. Pat. No. 4,539,203 published September 3, 1985 (Alex J. Brankiewicz et al.).
- 5) CL-1577D and CL-1577E Antibiotic antitumor compounds, U.S. Pat. No. 4,539,203.
- 6) CL-1724 Antibiotic compounds, U.S. Pat. No. 4,554,162.
- 7) Antitumour antibiotics BBM-1675-A3 and BBM-1675-A4, obtained by fermentation of actinomadura verrucosospora strains H964-92 (ATCC 39334) or AB27Y (ATCC 39638). U.S. Pat. No. 4,675,187.
- 8) N-acetyl-esperamicin A_1 , A_2 and $A_{1\beta}$ derivatives with antimicrobial and antitumor activities. European Patent Specification 289,030.

All of the information regarding the LL-E33288 family of antitumor antibiotics, BBM-1675, FR-900405, FR-900406, PD 114759, PD 115028, CL-1577A, CL-1577B, CL-1577D, CL-1577E and CL-1724 contained in the above citations is of interest to the present invention.

The α_1 , α_2 , α_3 , α_4 , β_1 , β_2 , γ_1 , δ_1 , and pseudoaglycone components of the LL-E33288 complex their dihydro and N-acyl counterparts, as well as the BBM-1675, FR-900405, FR-900406, PD 114759, PD 115028, CL-1577A, CL-1577B, CL-1577D, CL-1577E and CL-1724 antibiotics and their N-acyl counterparts, each contain a methyltrithio group in their structure. The methyltrithio moiety of the above-named antibiotics is subject to displacement by a variety of thiol-containing organic molecules resulting in the formation of




a new class of anticancer and antibacterial agents as described in U.S. Patent Nos. 5,606,040 and 5,770,710. Displacement of the methyltrithio unit of the antitumour antibiotics as depicted in Scheme I, below, can be used to introduce a spacer (Sp), the judicious choice of which enables the introduction of a HAM of the invention (hereinafter Hu:CT-M-01) into the compounds of the above-named patents and applications to form a conjugate molecule according to the invention.

Scheme 1




With reference to Scheme I $\text{CH}_3\text{-SSS-W}$ is the antitumour antibiotic, Sp is a straight or branched-chain divalent or trivalent ($\text{C}_1\text{-C}_{18}$) radical,


divalent or trivalent aryl or heteroaryl radical, divalent or trivalent (C₃-C₁₈) cycloalkyl or heterocycloalkyl radical, divalent or trivalent aryl- or heteroaryl-alkyl (C₁-C₁₈) radical, divalent or trivalent cycloalkyl- or heterocycloalkyl-alkyl (C₁-C₁₈) radical, or divalent or trivalent (C₂-C₁₈) unsaturated alkyl radical, wherein if Sp is a trivalent radical, it can be additionally substituted by amino, alkylamino, arylamino, heteroarylamino, carboxyl, lower alkoxy, hydroxy, thiol or lower alkylthio groups; Q is, or can be subsequently converted to, halogen, amino, alkylamino, carboxyl, carboxaldehyde, hydroxy, thiol, α-haloacetyloxy, lower alkyldicarboxyl, -CONHNH₂, -NHCONHNH₂, -NHCSNHNH₂, -ONH₂, -CON₃,



As long as the product from Scheme I contains at least one functional group which can be converted to, or is directly reactive with Hu:CT-M-01, targeted forms of the antitumor antibiotics of the above-named patents and applications can be generated, as shown in Scheme II below:

Scheme II

wherein Q, Sp, and W are as hereinbefore defined, Hu:CT-M-01 is a HAM, its fragments, or an analogue thereof; Y is a side-chain amino, carboxy, or thiol group of a protein, an aldehyde derived from carbohydrate residues, or an amidoalkylthio group; n is an integer of from 1 to 100; Z is formed from covalent reaction of the groups Q and Y directly or after subsequent reduction and Z is -CONH-, -CONHN=CH-, -CONHNHCH₂-, -NH-CONHN=CH-, -NHCONHNHCH₂-, -NHCSNHN=CH-, -NHCSNHNHCH₂-, -ON=CH-, -NH-, -NHCH₂-, -N=CH-, -CO₂-, -NHCH₂CO₂-, -SS-,




and *m* is 0.1 to 15.

As an example, and with reference to Scheme II, above, the 3-mercaptopropionic acid derivative of E-33288γ₁^I (Q=CO₂H, Sp=-CH₂CH₂-), when converted to its activated hydroxysuccinimide form (Q=CO₂Su, Sp=-CH₂CH₂-) can be used to react with some of the ε-amino groups of lysine residues (e.g., Y=-NH₂ wherein n=50-100 from available lysine residues), of Hu:CT-M-O1 at a pH between 7.0 and 9.5 in aqueous buffered solutions at temperatures between 4°C to 40°C to produce conjugate molecules of the invention with the antibiotics attached at random sites along the protein backbone (Z=-NHCO-, Sp=-CH₂CH₂-, *m*=1-10). Only a fraction of the available lysine residues are substituted in this manner, since high loading is generally

not considered compatible with preserving the antibody immunoreactivity. The same randomly-substituted immunoconjugates can also be prepared from the 3-mercaptopropionic acid derivative using other carboxyl group activating agents such as a variety of carbodiimides, or the corresponding acyl azide. Alternatively, a 3-mercaptopropionyl hydrazide derivative of E-33288_{y1}¹ ($Q = H_2NNHCO-$, $Sp = -CH_2CH_2-$), when reacted with a periodate-oxidized antibody ($Y = -CHO$, $N = 1-15$) as described in U.S. Pat. No. 4,671,958 published on June 9, 1987 (Cytogen Corp.) at a pH between 4 and 7, in a buffered aqueous solution at a temperature of between 4°C and 40°C, reacts only at the aldehyde functionality (derived from cleavage of vic-diols of carbohydrate residues situated on the Fc portion of the antibodies) to generate Hu:CT-M-01 conjugates containing the drug substituted at specific sites along the backbone of the protein ($Z = -CH=NNHCO-$, $Sp = -CH_2CH_2-$, $m = 0.5-10$). Other aldehyde-reactive groups as part of the drug construct are within our invention to generate the products of Scheme II. Such functional groups are preferably, though not limited to, those which react with aldehydes under acidic aqueous conditions. The reactivity of protein lysines under basic conditions is sufficiently great such that their amines compete with the products of Scheme II for available aldehydes of the monoclonal antibody. Alternative aldehyde-reactive groups are for example, the semicarbazide, the thiosemicarbazide, and the O-substituted hydroxylamine functionalities.

Assembly of conjugate molecules of the invention is not restricted to the sequence outlined in Scheme II. The Hu:CT-M-01 antibody can be first modified to contain a thiol group, which is then reacted with the antitumour antibiotics useful in the invention in accordance with Scheme III below:

Scheme III




wherein Hu:CT-M-O1, Y, Q, Sp, W, *n*, and *m* are as hereinbefore defined, and P is hydrogen or 2-(pyridylthio), with the proviso that when Y is a thiol derived from a backbone amino acid residue of Hu:CT-M-O1, Z-Sp taken together is a covalent bond.

As an example, and with references to Scheme III, above, the Hu:CT-M-01 monoclonal antibody can be reacted with 3-(2-dithiopyridyl)propionic acid hydroxysuccinimide ester to modify the protein through lysine residues ($Y=NH_2$, $n=50-100$, $Q=-CO_2Su$, $Sp=-CH_2CH_2-$, $P=2$ -pyridylthio). Following reduction with, for example, dithiothreitol, an intermediate is generated ($Z=-NHCO-$, $Sp=-CH_2CH_2-$, $P=H$, $m=1-15$) which can be reacted with the antitumour antibiotics to generate the subject immunoconjugates. Similarly, 2-iminothiolane can be reacted with Hu:CT-M-01 to introduce thiol groups onto the surface of the protein directly, without requiring a reduction step ($Z=-NHCO-$, $Sp=-(CH_2)_3-$, $P=H$, $m=1-15$), and this intermediate can be reacted with the CH_3 -SSS-W as before. Alternatively, sulfhydryl groups inherent within the structure of Hu:CT-M-01 in dimeric form as cystine residues can be used to participate in the reaction of Scheme III directly. Such sulfhydryls are traditionally exposed by a combination of enzymatic digestion and reduction of native monoclonal antibodies (Hu:CT-M-01=Fab' fragment, $Z-Sp=Bond$, $Y=SH$).


A preferred embodiment of the present invention is a protein-drug conjugate of the formula:

2095926




prepared from the class of antitumour antibiotics designated LL-E33288 (CH₃-SSS-W) comprising:

displacing the dithiomethyl moiety with a compound of formula Q-Sp-SH, wherein Sp is straight or branched-chain divalent or trivalent (C₂-C₁₀) radicals or divalent or trivalent (C₂-C₅) arylalkyl or heteroarylalkyl radicals, wherein if Sp is a trivalent radical, it can be additionally substituted by amino, heteroarylamino, hydroxy, or thiol groups; and Q is carboxyl, lower alkylidicarboxyl anhydride, -CO₂Su, -CONHNH₂, or




to produce an intermediate of general formula Q-Sp-SS-W, wherein Q, Sp, and W are as hereinbefore defined, and

reacting Q-Sp-SS-W with a molecule of the formula Hu:CT-M-O1-(Y)_n wherein Y is a side-chain amino group on the antibody, or an aldehyde generated by oxidation of the carbohydrate groups of the antibody, and *n* is an integer of from 1 to 100, to produce a compound of the formula:



wherein Y, Sp, W, and *n* are as hereinbefore defined, and Z is formed from covalent reaction of the groups Q and Y directly or after subsequent reduction, and Z is -CONH-, -CONHN=CH-, -CONHNHCH₂-, or



and m is 0.1 to 15.

The present invention also includes therapeutic and diagnostic compositions containing the HAM of the invention, particularly a conjugate molecule comprising a HAM conjugated to an effector or reporter molecule and uses of such compositions in therapy and diagnosis. Such therapeutic and diagnostic compositions typically comprise a HAM according to the invention together with a pharmaceutically acceptable excipient, diluent or carrier, e.g. for *in vivo* use.

Therapeutic and diagnostic uses typically comprise administering a pharmaceutically effective amount of a HAM according to the invention to a human subject. The exact dose to be administered will vary according to the intended use of the HAM and on the age and condition of the patient but may be typically varied from about 0.1mg to 1000mg, for example from about 1mg to 500 mg. The HAM may be administered as a single dose, or in a continuous manner over a period of time. Doses may be repeated as appropriate. The HAM may be formulated in accordance with conventional practice for administration by any suitable route, and may generally be in a liquid form [e.g. a solution of the antibody in a sterile physiologically acceptable buffer] for administration by for example an intravenous, intraperitoneal or intramuscular route.

In the HAM of the first aspect of the invention and the process of the second aspect of the invention, the heavy and light chain variable domains of the HAM may comprise either the entire variable domains of the CTMO1 MAb or may comprise framework regions of a human variable domain having grafted thereon one, two or all three of the CDRs of the CTMO1 MAb. Thus, the HAM may comprise a chimeric humanised antibody or a CDR-grafted humanised antibody.

When the HAM is a CDR-grafted humanised antibody. in addition to the CDRs, specific variable region framework residues may be altered to correspond to non-human, i.e. the CTMO1 mouse, residues. Preferably, the CDR-grafted humanised antibodies of the present invention include CDR-grafted humanised antibodies as defined in our International Patent Specification WO-A-91/09967.

Preferably, the CDRs of the light chain correspond to the Kabat CTMO1 MAb CDRs at CDR1 (residues 24-34) and CDR2 (residues 50-56) and to the structural loop residues (residues 91-96) or Kabat CTMO1 MAb CDR residues (residues 89-97) in CDR3. (The residue designations given above and elsewhere in the present application are numbered according to the Kabat numbering system [16]). In addition, the light chain may have mouse CTMO1 residues at one or more of residues 1, 2, 3, 36, 37, 45, 48, 49, 60, 63, 70, 84, 85, 87 and 108. In preferred embodiments, when the human framework used is EU, the light chain comprises Kabat CTMO1 MAb CDRs at all of CDR1, CDR2 and CDR3 and preferably additional CTMO1 residues at positions 3, 36, 37, 45, 48, 63 and 108, or especially additional CTMO1 residues at positions 3, 36, 63 and 108 only.

Preferably, the CDRs of the heavy chain correspond to the Kabat CTMO1 MAb CDRs at all of CDR1 (26 to 35), CDR2 (50 to 65) and CDR3 (94 to 100). In addition, the heavy chain may have mouse CTMO1 residues at one or more of residues, 2, 6, 23, 37, 48, 49, 67, 69, 73, 76, 78, 80, 88, 91 and 94. In particularly preferred embodiments, when the human framework used is EU, the heavy chain framework comprises additional CTMO1 MAb residues at positions 2, 37, 71 and 73, and especially in addition at positions 48, 67 and 69.

In addition, EU has a particularly idiosyncratic J region between residues 103 to 113 and it may be useful to include the murine amino acids, a consensus human J region or a suitable combination of both at residues 103 to 108 inclusive. When the EU framework is used,

preferably heavy chain residues 94, 103, 104, 105 and 107 are murine residues, since in the case of these residues, the murine sequence is more frequently found in human VH sequences than the EU residues.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention is now described, by way of example only, with reference to the accompanying drawings, in which:

- Figure 1 is a schematic diagram of plasmid pRR62;
- Figure 2 is a schematic diagram of plasmid pAL41;
- Figure 3 is a schematic diagram of plasmid pMRR017;
- Figure 4 is a schematic diagram of plasmid pHMC34;
- Figure 5 is a schematic diagram of plasmid pMRR011;
- Figure 6 is a schematic diagram of plasmid pHMC32;
- Figure 7 is a schematic diagram of plasmid pMRR022;
- Figure 8 is a schematic diagram of plasmid pMRR014;
- Figure 9 is a schematic diagram of plasmid pHMC33;
- Figure 10 is a schematic diagram of plasmid pMRR001;
- Figure 11 is a schematic diagram of plasmid pHMC35;
- Figure 12 is a schematic diagram of plasmid pHMC38;
- Figure 13 is a schematic diagram of plasmid pHMC40;
- Figure 14 is a schematic diagram of plasmid pHMC41;
- Figure 15 is a schematic diagram of plasmid pHMC42;
- Figure 16 shows the alignment of oligonucleotides H1 to H8 in the formation of the gH1 coding sequence;
- Figure 17 is a schematic diagram of plasmid pAL51;
- Figure 18 is a schematic diagram of plasmid pAL52;
- Figure 19 is a schematic diagram of plasmid pMRR010;
- Figure 20 is a schematic diagram of plasmid pAL47;
- Figure 21 is a schematic diagram of plasmid pAL48;
- Figure 22 is a graph of a direct binding ELISA on transiently expressed chimeric antibodies;
- Figure 23 is a graph of a direct binding ELISA on transiently expressed CDR-grafted antibodies; and
- Figure 24 is a graph of a competition EIA on transiently expressed chimeric and CDR-grafted antibodies.

Figure 25 is a graph comparing the effects on tumour size obtained by treating nude mice implanted with a human ovarian xenograft tumour with a humanised CDR grafted CTMO1 and a murine CTMO1 antibody each conjugated to the hydroxysuccinimide derivative of 4-mercapto-4-methyl-pentanoic acid disulphide of N-acetyl calicheamicin γ_1^1 .

DESCRIPTION OF SPECIFIC EMBODIMENTS

The following description of certain embodiments of the invention is provided by way of example only and is not to be regarded as placing any limitation on the scope of the protection claimed.

MOLECULAR CLONING AND CONSTRUCTION OF THE CTMO1 CHIMERIC HEAVY CHAIN

The heavy chain variable domain of CTMO1 was cloned using the polymerase chain reaction. This enabled the construction of the chimeric version in a single step as described below.

Polyadenylated RNA was isolated from the CTMO1 hybridoma cell line using the guanidinium isothiocyanate/lithium chloride method [17]. Double stranded cDNA was synthesised and used as a template for PCR amplification of the VH gene. A set of twenty four 5' forward primers were synthesised to complement a sequence within the murine leader sequence of VH domains [16] and to introduce a BstEII restriction site. A set of twelve 3' reverse primers was synthesised to complement the framework 4 region of VH [20] and included an ApaI restriction site.

The sequence of the basic 5' primer is given in the Sequence Listing as ID No. 3. The set of twenty four primers was based on this primer as follows. In one group of twelve primers, residue 27 remained as C. In three subgroups of four primers, residue 25 either remained as G or is altered to C or T. In each subgroup, the four primers differed at residue 28, which was A, C, G or T. In the subgroups where residue 25 is C or T, the sixth amino acid is His.

In the second group of twelve primers, residue 27 is changed to G. In three subgroups of four primers, residue 25 either remains as G or is altered to C or T. In each subgroup, the four primers differed at residue 28, which was A, C, G or T. Where residue 25 is C or G, amino acid 6 is Gln and where residue 25 is T, amino acid residue G is His. Where residue 28 is T or C, amino acid residue 7 is Cys. Where residue 28 is G, amino acid residue 7 is Trp.

The sequence of the basic 3' PCR primer is given in the Sequence Listing as ID No. 4. The set of twelve primers was based on this primer as follows. Residue 5 could remain as G or could be altered to A or T. Residue 11 either remains as A or is altered to G. Residue 12 either remains as A or is altered to C.

PCR amplification of CTMO1 VH was carried out using the following conditions:

10 pmoles each primer; 20 ng cDNA; 0.5 U Taq polymerase; 94°C 1 min; 50 °C 2 min; 72°C 3 min; for 40 cycles.

The PCR amplified VH fragment was restricted with BstEII and Apal and ligated to an adaptor to reconstruct the leader sequence and add a 5' HindIII restriction enzyme site. The sequence of the adaptor used is given in the Sequence Listing as Sequence ID No. 5 and codes in part for the leader amino acid sequence of the VH domain of the murine anti TAG-72 monoclonal antibody B72.3 (WO-A-89/01783).

The adapted fragment was then cloned into the HindIII/Apal sites of the vector pE1004 to give plasmid pRR62 shown in Figure 1. Plasmid pRR62 consists of an SV40 origin of replication followed by the hCMV-MIE promoter/enhancer region. The promoter/enhancer controls a nucleotide sequence encoding a chimeric heavy chain comprising the CTMO1 heavy chain variable domain fused to human g4 constant domains. Downstream of the coding sequence is a poly A site and gpt gene.

The heavy chain variable region of several independent clones of pRR62 were sequenced. The DNA sequence and deduced amino sequence for CTMO1 VH are given in Sequence ID No. 1.

MOLECULAR CLONING AND CONSTRUCTION OF THE CTMO1 CHIMERIC LIGHT CHAIN

Polyadenylated RNA was isolated from the CTMO1 hybridoma cell line using the guanidinium isothiocyanate/lithium chloride method [17]. Double stranded cDNA was synthesised [21] and a cDNA library was constructed in plasmid pSP64 [22] using EcoRI linkers. A screening probe was synthesised, complementary to mouse immunoglobulin light chain constant region by PCR amplification. The light chain probe was a 318 bp PCR fragment encoding the mouse kappa light chain constant region [23].

The probe was radio-labelled ($g^{32}P$) ATP by random hexanucleotide priming and was used to screen the cDNA library.

The clone which encoded the complete leader, variable and constant domains of light chain was isolated and designated as pRB63.

A fragment of pRB63, which encodes the variable domain of the light chain was recovered by PCR amplification. The PCR primers introduced a BstBI and SphI restriction sites at the 5' and 3' ends of the VL region respectively to enable subsequent cloning of the fragment.

The PCR amplified fragment of plasmid pRB63 was restricted with BstBI/SphI and ligated between the BstBI/SphI sites of plasmid pMRR010 to produce plasmid pAL41, which is shown in Figure 2. Plasmid pAL41 consists of a glutamine synthetase cDNA having downstream of it the hCMV-MIE promoter/enhancer region. The promoter/enhancer region controls a nucleotide sequence encoding a chimeric light chain comprising the CTMO1 light chain variable domain fused to a human CK constant domain. Downstream of the coding sequence is a poly A site.

Nucleotide sequence analysis was carried out according to the chain termination procedure [24]. The VH coding sequence insert in pRR62 and the VL coding sequence insert in pAL41 were fully sequenced. The DNA and predicted amino acid sequences for the unprocessed variable domains of the CTMO1 heavy and light chains are shown in the Sequence Listing appended to the end of the description as Sequence ID No. 1 and No. 2 respectively.

Sequence No. 1 shows the sequence coding for the VH domain and the predicted amino acid sequence. The leader sequence for the heavy chain runs from residue 1 to residue 19 as shown in Sequence No. 1. Sequence No. 2 shows the sequence coding for the VL domain together with the predicted amino acid sequence. The leader sequence for the light chain runs from residue 1 to residue 20 as shown in Sequence No. 2. Examination of the derived amino acid sequences revealed considerable homology with other characterised immunoglobulin genes. The CTMO1 MAb was confirmed to be an IgG1-kappa antibody.

PREPARATION OF CHIMERIC ANTIBODY PRODUCTS

CHIMERIC LIGHT CHAIN VECTOR

A ClaI-EcoRI fragment of pAL41 carrying the hCMV promoter and chimeric light chain was cloned into plasmid pMRR017 which is shown in Figure 3. Plasmid pMRR017 has a GS mini gene (PCT Application WO-A-87/04462 published on July 30, 1987 (Celltech Ltd.)), hCMV-MIE promoter/enhancer region, a polylinker sequence and a poly A site. This produced plasmid pHMC34, which is shown in Figure 4. In plasmid pHMC34, the chimeric light chain gene is under the control of the hCMV-MIE promoter/enhancer sequence.

CHIMERIC HEAVY CHAIN VECTORS

IgG1 CONSTRUCT

A HindIII-ApaI fragment containing the sequence encoding the VH domain was excised from plasmid pRR62 (Figure 1). This fragment was inserted between the HindIII and ApaI sites of plasmid pMRR011. Plasmid pMRR011 is shown in Figure 5 and comprises an hCMV-MIE

promoter/enhancer region, an SV40 polyadenylation sequence, a gpt gene and a sequence encoding a human IgG1 heavy chain lacking a variable domain. The plasmid thus produced, pHMC32, is shown in Figure 6 and has a chimeric heavy chain coding sequence under the control of the hCMV-MIE promoter/enhancer. The chimeric heavy chain has the VH domain from the CTM01 MAb fused to human IgG1 constant domains.

IgG2 CONSTRUCT

The HindIII-ApaI fragment of pRR62 (Figure 1) was inserted between the HindIII and ApaI sites of a plasmid containing an hCMV-MIE promoter, a polylinker site and a nucleotide coding sequence which encodes the three constant domains of a human IgG2 antibody. This yielded plasmid pMRR022 which encodes a chimeric heavy chain having the CTM01 variable domain linked to the human IgG2 constant domains.

IgG4 CONSTRUCT

The HindIII-ApaI fragment of pRR62 (Figure 1) was inserted between the HindIII and ApaI sites of plasmid pMRR014 to produce plasmid pHMC33. Plasmids pMRR014 and pHMC33 are shown in Figure 8 and 9 respectively. Plasmid pMRR014 has an hCMV-MIE promoter, a polylinker site and a nucleotide coding sequence which encodes the three constant domains of a human IgG4 antibody. Plasmid pHMC33 is identical to plasmid pHMC32 except that the coding sequence encodes a chimeric heavy chain having the CTM01 variable domain and human IgG4 constant domains in place of the human IgG1 constant domains.

ALTERED IgG4 CONSTRUCT

The HindIII-ApaI fragment was reisolated from plasmid pHMC33. Plasmid pMRR001 shown in Figure 10 was digested with HindIII and ApaI. The large fragment was isolated and ligated to the HindIII-ApaI fragment of pHMC33 to produce plasmid pHMC35, shown in Figure 11. Plasmid pHMC35 is almost identical to plasmid pHMC32 except that the coding sequence encodes a chimeric heavy chain having the

CTM01 variable domain and altered human IgG4 (hereinafter referred to as IgG4P) constant domains in place of the human IgG1 constant domains.

The alteration in the constant domains comprises a change of a serine residue in the hinge region at position 241 to a proline residue. This change advantageously abolished the formation of an 80 KD half antibody which otherwise occasionally is formed with IgG4 constant domains.

CHIMERIC HEAVY AND LIGHT CHAIN VECTORS

Vectors were constructed having operons coding for both heavy and light chains within the same vector.

A NotI-SalI fragment carrying the hCMV-MIE promoter/enhancer, the chimeric light chain encoding sequence and the SV40 poly A site together with the GS mini gene was excised from plasmid pHMC34 (Figure 4). A NotI-HindIII fragment carrying the hCMV-MIE promoter/enhancer was excised from plasmid pHMC35 (Figure 11). A HindIII-SalI fragment carrying the altered IgG4 heavy chain coding sequence and SV40 poly A site was excised from plasmid pHMC35 (Figure 11). These three fragments were ligated together to produce plasmid pHMC38, which is shown in Figure 12, and codes for expression of chimeric light chain together with the altered IgG4 chimeric heavy chain.

Plasmids pHMC32, pMRR022 and pHMC33 were digested with HindIII and EcoRI and the fragments containing the chimeric heavy chain encoding sequences were isolated. The isolated fragments were each ligated with the large HindIII-SalI fragment of pHMC38 (Figure 12) and an EcoRI-SalI fragment comprising the SV40 poly A region. The ligations produced plasmids pHMC40, pHMC41 and pHMC42 (shown in Figures 13 to 15 respectively). pHMC40 encodes a heavy chain having IgG1 constant domains. pHMC41 encodes IgG2 constant domains and pHMC42 encodes IgG4 constant domains.

PREPARATION OF CDR-GRAFTED ANTIBODY PRODUCTS

It was decided to use the EU human antibody framework [16] for carrying out the CDR-grafting. The strategy followed for CDR-grafting was as set out in our International Patent Specification No. WO-A-91/09967.

Two CDR-grafted heavy chains were designed. In the first, gH1, all three CDRs [as defined by Kabat, ref. 16] were changed to murine residues. In addition, residues 2, 37, 71, 73, 94, 103, 104, 105 and 107, which are outside the Kabat CDRs, were also changed to murine residues. In the second, gH2, in addition to those murine residues in gH1, residues 48, 67 and 69 were changed to murine residues with a view to improving packing of the VH domain.

Two CDR-grafted light chains were also designed. In the first, gL1, all three CDRs [as defined by Kabat, ref. 16] were changed to murine residues. In addition residues 3, 36, 63 and 108, which are outside the Kabat CDRs, were changed to murine residues. In the second, gL2, in addition to those murine residues in gL1, residues 37, 45 and 48 were changed to murine residues with a view to improving packing.

A nucleotide sequence coding for the gH1 variable domain was produced by oligonucleotide assembly using oligonucleotides H1 to H8. The sequences for these oligonucleotides are given in the Sequence Listing at the end of the description under Sequence ID Nos. 6 to 13. The way in which these oligonucleotides are assembled to produce the gH1 coding sequence is shown in Figure 16. The amino acid sequence coded for by this gH1 sequence is shown in the sequence listing under Sequence ID No. 14.

A nucleotide sequence coding for the gH2 variable domain was also produced by oligonucleotide assembly using oligonucleotides H1, H2, H3A, H4, H5, H6A, H7 and H8. Oligonucleotide H3A differs from oligonucleotide H3 (Sequence ID No. 8) in that residues 55 to 57 have been changed from GTG to GCA and residues 61 to 63 have been changed from ATT to CTG. Oligonucleotide H6A differs from oligonucleotide H6 (Sequence ID No. 11) in that residues 70 to 72

have been changed from TAC to TAA. Thus, the gH2 variable domain encodes the same sequence as is shown under Sequence ID No. 14, except that at residue 67, MET has been changed to ILE; at residue 87, VAL has been changed to ALA; and at residue 89, ILE has been changed to LEU.

A nucleotide sequence coding for the gL1 variable domain was produced by oligonucleotide assembly using oligonucleotides L1 to L8. The sequences for these oligonucleotides are given in the Sequence Listing at the end of the description under Sequence ID Nos. 15 to 22. The way in which these nucleotides are assembled is similar to that shown in Figure 16 for the gH1 coding sequence (except that L is substituted for H). The amino acid sequence coded for by the assembled gL1 variable domain coding sequence is shown in the Sequence Listing under Sequence ID No. 20.

A nucleotide sequence coding for the gL2 variable domain was produced by oligonucleotide assembly using oligonucleotides L1, L2A, L3A and L4 to L8. Oligonucleotide L2A differs from oligonucleotide L2 (Sequence ID No. 16) in that residues 28 to 30 have been changed from CAG to GTA. Oligonucleotide L3A differs from oligonucleotide L3 (Sequence ID No. 17) in that residues 25 - 27 have been changed from CAG to CTC, residues 49 - 52 have been changed from AAG to CAG and residues 59 - 61 have been changed from CAT to ATC. Thus, the gL2 variable domain encodes the same sequence as is shown under Sequence ID NO. 23, except that: at residue 23, Gln has been changed to Val; at residue 62, Gln has been changed to Leu; at residue 60, Lys has been changed to Gln; and at residue 73, Met has been changed to Ile.

For gene assembly 1 pmol of H2 - H7 or L2 - L7 was mixed with 10 pmol of H1 and H8 or L1 and L8 in a 100 µl reaction with 5U Taq polymerase. A PCR reaction was done using 30 cycles (95°C, 1 min. 50°C 1 min; 72°C 1 min). The resulting fragments were cut with HindIII and ApaI for VL with Bstb1 and SphI for VH.

The nucleotide sequences coding for gH1 and gH2 were cloned as HindIII-ApaI fragments into plasmid pMRR014 (Figure 8) to produce plasmids pAL51 and pAL52 (Figure 17 and 18 respectively).

The nucleotide sequences coding for gL1 and gL2 were cloned as HindIII-ApaI fragments into plasmid pMRR010 (Figure 19) to produce plasmids pAL47 and pAL48 (Figures 20 and 21 respectively).

TRANSIENT EXPRESSION OF CHIMERIC/CHIMERIC OR CDR-GRAFTED/CHIMERIC ANTIBODIES

The following plasmids:

pHMC38, pHMC40, pHMC41 and pHMC42

and the following pairs of plasmids:

pAL47, pHMC33; pAL48, pHMC33; pAL51, pAL41; pAL52, pAL41; and pAL48, pAL41;

were each transfected or cotransfected into CHO-L761h cells for transient expression.

Assembly ELISA assays on culture supernatants resulting from the single transfected cells showed that they contained assembled antibody.

The assembly ELISA assay for quantifying antibody yields used microwell plates coated with a goat F(ab')₂ anti-human IgG₁Fc. Following incubation with transfected culture supernatants, bound chimeric or CDR-grafted antibody was revealed with a horseradish peroxidase (HRP)-conjugated murine anti-human IgK antibody using tetramethyl benzidine (TMB) as the substrate. Concentrations of chimeric or CDR-grafted whole antibody in the samples were interpolated from a calibration curve generated from serial dilutions of purified chimeric B72.3 γ 4 antibody [25].

BINDING ACTIVITY OF TRANSIENTLY EXPRESSED CHIMERIC OR CDR-GRAFTED ANTIBODIES

Direct binding ELISA assays for determining the binding activity of the transiently expressed antibodies were carried out as follows.

2095926

An affinity column was prepared by attaching the CTM01 MAb to a suitable chromatographic medium in conventional manner. In a first method, pooled human urine samples were applied directly to the affinity column. In a second method, human milk was subjected to low speed centrifugation to separate the cream from skimmed milk. The skimmed milk was then subjected to high speed centrifugation to produce an aqueous and a lipid component. The aqueous component was applied to the affinity column.

Once the affinity column was loaded, by either of the two methods, column fractions were eluted at high and low pHs, neutralised and assayed for reactivity with the CTM01 MAb. Fractions showing reactivity were pooled and dialysed. The pooled fractions contained the polymorphic epithelial mucin (PEM) recognised by the CTM01 MAb.

Microwell plates were coated with PEM obtained as described above. The microwells were then incubated with serial dilutions of culture supernatants. Binding of chimeric or CDR-grafted antibody was revealed and quantified by use of an HRP-conjugated murine anti-human IgK antibody.

The results of direct binding ELISA assays on the supernatants from singly transfected cells are shown in Figure 22. These assays confirm that all the supernatants contained antibodies capable of binding to PEM. No significant differences in binding activity were observed.

The direct binding ELISA assays on the supernatants from doubly transfected cells confirmed that the supernatants contained antibodies capable of binding to PEM and that the chimeric/chimeric antibody bound better than any of the CDR-grafted/chimeric antibodies.

A competition binding assay was carried out using polystyrene beads coated with PEM obtained as described above. CTM01 MAb was radiolabelled with ¹²⁵I and was used to compete with the antibody produced by the pHMC40 (IgG1) transfected cells. The potency of the chimeric antibody was 84-102% that of the CTM01 MAb.

2095926

TRANSIENT EXPRESSION OF CDR-GRAFTED/CDR-GRAFTED ANTIBODIES

The following pairs of plasmids:

pAL47, pAL51; pAL47, pAL52; pAL48, pAL51; and pAL48, pAL52;

were cotransfected into CHO-L761 cells.

Direct binding assays were carried out on the culture supernatants produced by the doubly transfected cell lines.

The results of these assays are shown in Figure 23, together with some results for chimeric/CDR-grafted antibodies.

From all the direct binding assays referred to above, it can be determined that the order of binding activity of the various antibodies produced by transient expression is as follows:

$$cLcH3 > gL1ch = gL1gH2 > cLgH2 = gL2H2 = gL1gH1 = gL2cH > gL2gH1.$$

(wherein: cL = chimeric light chain;

ch = chimeric heavy chain

gL1 = CDR-grafted light chain with lowest number of amino acid changes;

gL2 = CDR-grafted light chain with highest number of amino acid changes;

gH1 = CDR-grafted heavy chain with lowest number of amino acid changes;

gH2 = CDR-grafted heavy chain with highest number of amino acid changes).

The more active variants (cLcH, gL1cH, gL1gH2 and gL2gH2) together with the CTM01 MAb were tested in a competition enzyme immunoassay (EIA). Microwell plates were coated with PEM obtained as described above. The CTM01 MAb was biotinylated and was used to compete with the four variants referred to above. Bound biotinylated

CTMO1 MAb was revealed and quantified using a streptavidin-HRP conjugate and TMB.

The results of the competition EIA are shown in Figure 24, which shows the same ranking of binding activity as set out above, except that the gL1cH combination shows greater activity than the cLcH combination.

It can thus be seen that chimeric, chimeric/CDR-grafted and CDR-grafted antibodies which recognise the same antigen as the CTMO1 MAb have successfully been produced.

IN VITRO CELL BINDING AND INTERNALISATION OF CDR-GRAFTED CTMO1 ANTIBODIES

Stable NSO cell lines expressing gL1gH2IgG2 CTMO1 (hereinafter hu1: CTMO1) and gL1gH2IgG4P CTMO1 (hereinafter hu:CTMO1) antibody variants were made by transfecting into NSO cells by electroporation double gene expression plasmids assembled by ligating the large (7.8 kbp) Not1/BamH1 fragment of pAL47 to the 2.4 kbp Not1/Apa1 fragment from pAL52 and either a 1.9Kbp BamH1/Apa1 (partial) fragment carrying the IgG2 constant domains or a 2kbp Apa1/BamH1 fragment carrying the IgG4P constant domains as appropriate.

Antibody, purified from the supernatant of each cultured cell line by protein-A SEPHAROSE (trade-mark), chromatography was radiolabelled (^{125}I) and incubated using a conventional continuous exposure method with either MX-1 or MCF-7 breast carcinoma cells. Radiolabelled murine CTMO1 was used in all tests as a comparison. All antibodies were incubated at $2\mu\text{g}/\text{million}$ cells. The total binding of antibodies to the cells and the peak net uptake of the antibodies by the cells was determined. The results are shown in Table 1 below. With both cell lines each CDR grafted antibody exhibited better binding and internalisation than the murine form.

TABLE 1

Antibody	Cell Line	Total Binding, 0° (molecules/cell)	Peak Net Uptake (molecules/cell)
hu1:CTMO1	MCF-7	650,000	150,000
hu:CTMO1	MCF-7	450,000	90,000
Murine CTMO1	MCF-7		300,000
			70,000
hu1:CTMO1	MX-1	1,200,000	150,000
hu:CTMO1	MX-1	1,100,000	150,000
Murine CTMO1	MX-1	800,000	80,000

IN VIVO ANTI-TUMOUR ACTIVITY OF A CONJUGATE OF
hu:CTMO1 AND AN ANTI-TUMOUR ANTIBIOTIC

hu:CTMO1 was conjugated to the hydroxysuccinimide derivative of 4-mercapto-4-methyl-pentanoic acid disulphide of N-acetyl calicheamicin γ_1^I as follows:

SYNTHESIS OF THE 4-MERCAPTO-4-METHYL-PENTANOIC ACID
DISULPHIDE DERIVATIVE OF N-ACETYL CALICHEAMICIN γ_1^I

To N-acetyl calicheamicin γ_1^I [US Patent No. 5079233] at a concentration of 2 mg/mL in acetonitrile at -15°C was added 5 molar equivalents of 4-mercapto-4-methyl-pentanoic acid and 6 molar equivalents of triethylamine. After 24 hours at -15°C the reaction was checked by C₁₈-HPLC. [If the reaction is incomplete, additional amounts of 4-mercapto-4-methyl-pentanoic acid and triethylamine are added]. Upon completion of the reaction the volatile organics were evaporated under reduced pressure and the crude product was chromatographed on BIO-SIL A (trade-mark) using a gradient of 1 to 5% methanol in chloroform. Pure fractions as assessed by tlc were pooled and evaporated to a glass. The ¹H-NMR of the product was similar to N-acetyl calicheamicin γ_1^I , but was missing the absorbance for -SSSMe and exhibits absorbances for the methylpentanoic acid moiety as expected. FAB-MS gave m/z = 1478(M + H) and 1500 (M + Na).

SYNTHESIS OF THE HYDROXYSUCCINIMIDE DERIVATIVE OF 4-MERCAPTO-4-METHYL-PENTANOIC ACID DISULPHIDE OF N-ACETYL CALICHEAMICIN γ_1

To the 4-mercapto-4-methyl-pentanoic acid disulphide derivative of N-acetyl calicheamicin γ_1 described above at a concentration of 5 mg/mL in acetonitrile at ambient temperature was added 3 molar equivalents of N-hydroxysuccinimide and 5 molar equivalents of 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride. After 1 hour the reaction was checked by C₁₈-HPLC. [If the reaction is incomplete, then additional 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride is added]. Upon completion of the reaction the volatile organics were evaporated under reduced pressure and the crude product was chromatographed on BIO-SIL A using a gradient of 0 to 5% methanol in chloroform. Pure fractions as assessed by tlc were pooled and stripped to a glass. The ¹H-NMR was similar to that of the product described above, but with absorbances present for succinimide, as expected. FAB-MS gave m/z = 1575 (M + H) and 1597 (M + Na).

SYNTHESIS OF hu:CTM01 CONJUGATE USING THE HYDROXY-SUCCINIMIDE DERIVATIVE OF 4-MERCAPTO-4-METHYL-PENTANOIC ACID DISULPHIDE OF N-ACETYL CALICHEAMICIN γ_1

To hu:CTM01 in phosphate buffer at a pH of about 7.4 was added 2-6 molar equivalents of the hydroxysuccinimide derivative of 4-mercapto-4-methyl-pentanoic acid disulphide of n-acetyl calicheamicin γ_1 , described above, in dimethylformamide (DMF) such that the final concentration of DMF was 10-15%. After completion of the reaction (2-24 hours) the low-molecular-weight organic material was removed by passing through a desalting column using pH 7.4 phosphate buffer. The product was further purified by chromatography on a gel exclusion column and concentrated to give a monomeric product with an average loading of 1-3 molecules of calicheamicin derivative per molecule of antibody.

2095926

IN VIVO TEST FOR ANTITUMOUR ACTIVITY

The human ovarian xenograft tumour, OvCar3, implanted subcutaneously in nude mice was used as a test system to study the efficacy of the hu:CTMO1 conjugate in vivo. A murine CTMO1 conjugate containing the same calicheamicin was also tested for comparison. Tumours were implanted subcutaneously into athymic mice and test samples were inoculated intraperitoneally (IP) at several dose levels on a q4 day x 3 schedule, starting 2-3 days after tumour implantation with 6 mice per group and 10 in each control group. Tumour mass was determined by measuring the tumour diameter once weekly during 42 days post tumour implantation. Significant antitumour activity was defined as a sustained 58% inhibition of mean tumour mass compared with untreated controls in groups with greater than 65% survivors. At both the 1 and 3 μ g doses of drug equivalents the hu:CTMO1 conjugate showed significant inhibition of tumour growth (Figure 25). No deaths were noted in the 42 days observation period in any test group. In all test groups, n=6, in the control group n=10, error bars = $\pm$ Standard Error Mean for each data point.

REFERENCES

1. Kohler & Milstein, Nature 265, 495-497, 1975
2. Begent et al, Br. J. Cancer, 62, 487, 1990
3. Verhoeyen et al, Science, 293, 1534-1536, 1988
4. Riechmann et al, Nature, 332, 323, 324, 1988
5. Queen et al, Proc. Natl. Acad. Sci., USA, 86, 10029-10033, 1989 and WO-A-90/078861
6. Tempest et al, Biotechnology, 9, 266-271, 1991
7. Co et al, Proc. Natl. Acad. Sci., USA, 88, 2869-2873, 1991
8. Verhoeyen et al, 1991 in Epenetos, A.A., (ed.), - "Monoclonal Antibodies: Applications in Clinical Oncology"
9. Gorman et al, Proc. Natl. Acad. Sci., USA, 88, 4181-4885, 1991
10. Ehrlich, P., Collected Studies on Immunity, 2, John Wiley & Sons, New York, 1986
11. Levy & Miller, Ann. Rev. Med., 34, 107-116, 198-(?)

2095926

12. Schlom & Weeks, Important Advances in Oncology, 170-192, Wippincott, Philadelphia, 1985
13. Sahagan et al, J. Immunol., 137, 3, 1066-1074, 1986
14. Nishimura et al, Cancer Res., 47, 999-1005, 1987
15. Aboud-Pirak et al, Cancer Res., 48, 3188-3196, 1988
16. Kabat et al, Sequences of Proteins of Immunological Interest, US Department of Health and Human Services, NIH, USA, 1987 and Wu, T.T. and Kabat, E.A. J. Exp. Med., 132, 211-250, 1970
17. Maniatis et al, Molecular Cloning, Cold Spring Harbour, New York, 1982
18. Primrose and Old, Principles of Gene Manipulation, Blackwell, Oxford, 1980
19. Jones et al, Nature, 54, 75-82, 1986
20. Oriandi et al, Proc. Natl. Acad. Sci., USA, 86, 3833-3837, 1989
21. Gubler and Hoffman, Gene, 25, 263-269, 1983
22. Melton et al, Nuc. Acids. Res., 12, 7035-7056, 1984
23. Max et al, J. Biol. Chem., 256, 5116-5120, 1981
24. Sanger et al, PNAS, 74, 5463-5467, 1977
25. Colcher et al, Proc. Natl. Acad. Sci., USA, 86, 3833-3837, 1989.

2095926

SEQUENCE LISTING

SEQUENCE ID NO: 1.

5 SEQUENCE TYPE: Nucleotide with deduced protein sequence.

SEQUENCE LENGTH: 416 bases.

STRANDEDNESS: Single.

TOPOLOGY: Linear.

MOLECULE TYPE: cDNA.

10 ORIGINAL SOURCE ORGANISM: Murine.

IMMEDIATE EXPERIMENTAL SOURCE

NAME OF CELL LINE: Hybridoma CTM01.

PROPERTIES: Coding sequence for variable domain of heavy chain of the CTM01 monoclonal antibody.

15 FEATURES: Leader sequence from residues 1 to 19.

ATG GAA TGG AGC TGG GTC TTT CTC TTC TTC CTG TCG GTA ACC ACA GGT
Met Glu Trp Ser Trp Val Phe Leu Phe Phe Leu Ser Val Thr Thr Gly

20 GTC CAT TGC CAG ATC CAG CTG CAG CAG TCT GGA CCT GAG CTG GTG AAG
Val His Cys Gln Ile Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys

CCT GGG GCT TCA GTG AAG ATA TCC TGC AAG GCT TCT GGC TAC ACC TTC
Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe

25

ACT GAC TAC TAT ATA AAC TGG ATG AAG CAG AAG CCT GGA CAG GGA CTT 192
Thr Asp Tyr Tyr Ile Asn Trp Met Lys Gln Lys Pro Gly Gln Gly Leu 64

GAG TGG ATT GGA TGG ATT GAT CCT GGA AGC GGT AAT ACT AAG TAC AAT 240
5 Glu Trp Ile Gly Trp Ile Asp Pro Gly Ser Gly Asn Thr Lys Tyr Asn 80

GAG AAG TTC AAG GGC AAG GCC ACA TTG ACT GTA GAC ACA TCC TCC AGC 288
Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser Ser Ser 96

10 ACA GCC TAC ATG CAG CTC AGC AGC CTG ACA TCT GAG GAC ACT GCT GTC 336
Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val 112

TAT TTC TGT GCA AGA GAG AAA ACG ACC TAT TAC TAT GCT ATG GAC TAC 384
Tyr Phe Cys Ala Arg Glu Lys Ile Thr Tyr Tyr Tyr Ala Met Asp Tyr 128

15 TGG GGT CAA GGA ACC TCA GTC ACT GTC TCC GC 416
Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ala 139

2095926

SEQUENCE ID NO: 2.
SEQUENCE TYPE: Nucleotide with deduced protein sequence.
SEQUENCE LENGTH: 399 bases.
5 STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: cDNA.
ORIGINAL SOURCE ORGANISM: Murine.
IMMEDIATE EXPERIMENTAL SOURCE
10 NAME OF CELLINE: Hybridoma CTM01.
PROPERTIES: Coding sequence for variable domain of light chain of the CTM01 monoclonal antibody
FEATURES: Leader sequence from residues 1 to 20.

ATG	AGG	TGC	CTA	GCT	GAG	TTC	CTG	GGG	CTG	CTT	GTG	CTC	TGG	ATC	CCT	48	
15	Met	Arg	Cys	Leu	Ala	Gly	Phe	Leu	Gly	Leu	Val	Leu	Trp	Ile	Pro	16	
GGA	CCC	ATT	GGG	GAT	ATT	GTG	ATG	ACT	CAG	GCT	GCA	CCC	TCT	GTT	CCT	96	
Gly	Ala	Ile	Gly	Asp	Ile	Val	Met	Thr	Gln	Ala	Ala	Pro	Ser	Val	Pro	32	
20	GTC	ACT	CCT	GGA	GAG	TCA	TTA	TCC	ATT	TCC	TGC	AGG	TCT	AGT	AAG	AGT	144
	Val	Thr	Pro	Gly	Gly	Ser	Leu	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Lys	Ser	48
CTC	CTT	CAT	AGT	AAT	GGC	GAC	ACT	TTC	TTG	TAT	TGG	TTC	CTG	CAG	AGG	192	
Leu	Leu	His	Ser	Asn	Gly	Asp	Thr	Phe	Leu	Tyr	Trp	Phe	Leu	Gln	Arg	64	

5	TCC	GGA	GTC	CCA	GAC	AGG	TTC	AGT	GGC	AGT	GGG	TCA	GGA	ACT	GCT	TTC	288
	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Ala	Phe	96

10

CTG	GAG	CTG	AAA	CGG	399	
15	Leu	Gly	Leu	Lys	Arg	133

SEQUENCE ID NO. 3
 SEQUENCE TYPE: Nucleotide sequence with corresponding amino acid sequence
 SEQUENCE LENGTH: 28 bases
 5 STRANDEDNESS: Single
 TOPOLOGY: Linear
 MOLECULAR TYPE: Synthesised DNA
 PROPERTIES: 5' forward primers for PCR amplification of murine VH domains
 FEATURES: Complementary to leader sequence of murine VH domains with introduced BstEII
 10 restriction site at residues 7 to 13

GGTGGCG GTA ACC ACA GGT GTC CAG TCA 28
 Val Thr Thr Gly Val Gln Ser 7

15 SEQUENCE ID NO. 4
 SEQUENCE TYPE: Nucleotide sequence
 SEQUENCE LENGTH: 36 bases
 STRANDEDNESS: Single
 TOPOLOGY: Linear
 20 MOLECULE TYPE: Synthesised DNA
 PROPERTIES: 3' reverse primer for PCR amplification of murine VH domains
 FEATURES: Complementary to framework 4 region of murine VH including an ApaI restriction
 site at residues 25 to 30

25 AGTGGCAGAG AAGTCGGAGT TGCTCCCGG GTAGAC 36

SEQUENCE ID NO. 5
SEQUENCE TYPE: Nucleotide sequences with corresponding amino acid sequence
SEQUENCE LENGTH: 50 bases (sense strand) and 51 bases (anti-sense strand)
STRANDEDNESS: Double
5 TOPOLOGY: Linear
MOLECULE TYPE: Synthesised DNA
PROPERTIES: Leader sequence adaptor for murine VH domain
FEATURES: Comprises a HindIII-BstEII fragment in part coding for the leader sequence of the
VH domain of murine monoclonal antibody B72.3

10

AGCTTGCCGC CACC ATG GAA TGG AGC TGG GTC TTT CTC TTC TTC CTG TCG 50
ACGGCG GTGG TAC CTT ACC TCG ACC CAG AAA GAG AAG AAG GAC AGC CATTG 51
Met Glu Trp Ser Trp Val Phe Leu Phe Leu Ser 12

SEQUENCE ID NO: 6
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 21.
5 STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide
PROPERTIES: Used for assembly of CDR-grafted heavy chain
FEATURES: HindIII site at residues 7-12

10

GCGCGCAAGC TTGCCGCCAC C

21

SEQUENCE ID NO: 7.
SEQUENCE TYPE: Nucleotide
15 SEQUENCE LENGTH: 96.
STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted heavy chain.

20

TCTCAGATTC AGCTGGTGCA GTCTGGAGCA GAGGTGAAGA AGCCTGGATC
TTCTGTGAAG GTGTCTTGTA AGGCATCTGG ATACACCTTC ACCGAC

50

96

2095926

SEQUENCE ID NO: 8.
 SEQUENCE TYPE: Nucleotide.
 SEQUENCE LENGTH: 96.
 5 STRANDEDNESS: Single.
 TOPOLOGY: Linear.
 MOLECULE TYPE: Synthetic oligonucleotide.
 PROPERTIES: Used for assembly of CDR-grafted heavy chain.

10 TGGATTGACC CTGGATCTGG AAATACAAAG TACAATGAGA AGTTCAAGGG 50
 AAGAGTGACA ATTACAGTGG ACACATCCAC GAATACCGCC TACATG 96

SEQUENCE ID NO: 9.
 SEQUENCE TYPE: Nucleotide.
 15 SEQUENCE LENGTH: 89.
 STRANDEDNESS: Single.
 TOPOLOGY: Linear.
 MOLECULE TYPE: Synthetic oligonucleotide.
 PROPERTIES: Used for assembly of CDR-grafted heavy chain.
 20 FEATURES: ApaI site at residues 78-83

GAGAAGACCA CCTACTACTA CGCAATGGAC TACTGGGGAC AGGGAACACT 50
 GGTGACAGTG TCTTCTGCCT CAACGAAGGG CCCGCGCGC 89

SEQUENCE ID NO: 10.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 96.
5 STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted heavy chain.

10 CTGCACCAGC TGAATCTGAG AATGGACTCC TGTAGTTACT GACAGGAAGA 50
AGAGAAAGAC CCAGCTCCAT TCCATGGTGG CGGCAAGCTT GCGCGC 96

SEQUENCE ID NO: 11.
SEQUENCE TYPE: Nucleotide.
15 SEQUENCE LENGTH: 96.
STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted heavy chain.

20
TCCAGATCCA GGGTCAATCC ATCCCATCCA CTCGAGTCCC TGTCAGGTG 50
CCTGTCTCAT CCAATTAATG TAGTAGTCGG TGAAGGTGTA TCCAGA 96

SEQUENCE ID NO: 12.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 93.
STRANDEDNESS: Single.
5 TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted heavy chain.

GTAGTAGTAG GTGGTCTTCT CTCTTGCACA GAAGTAGAAT GCTGTGTCCT 50
10 CAGATCTCAG AGAAGACAGC TCCATGTAGG CCGTATTCGT GGA 93

SEQUENCE ID NO: 13.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 21.
15 STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted heavy chain.
FEATURES: ApaI site at residues 7 - 12.

20 GCGCGGGGC CCTTCGTTGA G 21

2095926

SEQUENCE ID NO: 14.

SEQUENCE TYPE: Amino acid.

SEQUENCE LENGTH: 139.

5 MOLECULE TYPE: Immunoglobulin heavy chain variable domain.

FEATURES: CDRs at residues 45 - 54, 69 - 85 and 118 - 128

Met	Glu	Trp	Ser	Trp	Val	Phe	Leu	Phe	Phe	Leu	Ser	Val	Thr	Thr	Gly	16
Val	His	Ser	Gln	Ile	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	32
10	Pro	Gly	Ser	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	48
	Thr	Asp	Tyr	Tyr	Ile	Asn	Trp	Met	Arg	Gln	Ala	Pro	Gly	Gln	Gly	64
	Glu	Trp	Met	Gly	Trp	Ile	Asp	Pro	Gly	Ser	Gly	Asn	Thr	Lys	Tyr	80
	Glu	Lys	Phe	Lys	Gly	Arg	Val	Thr	Ile	Thr	Val	Asp	Thr	Ser	Thr	96
	Thr	Ala	Tyr	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	112
15	Tyr	Phe	Cys	Ala	Arg	Glu	Lys	Thr	Thr	Tyr	Tyr	Tyr	Ala	Met	Asp	128
	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					139

SEQUENCE ID NO: 15.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 21.
STRANDEDNESS: Single.
5 TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted light chain.
FEATURES: BstBI site at residues 7 to 11

10 GGACTGTTCG AAGCCGCCAC C 21

SEQUENCE ID NO: 16.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 81.
15 STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted light chain.

20 TGGCTTACAG ATGCCAGATG CGATATCCAG ATGACTCAGA GTCCAAGTAC 50
TCTCAGTGCC AGTGTAGGTG ATAGGGTCAC C 81

SEQUENCE ID NO: 17.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 90.
STRANDEDNESS: Single.
5 TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted light chain.

GGTGACACCT TCCTCTATTG GTTCCAGCAG AAACCAGGTA AAGCCCCAAA 50
10 GCTCCTCATG TATAGGATGA GTAACTCGC CAGTGGTGTA 90

SEQUENCE ID NO: 18.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 99.
15 STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted light chain

20 CAGCCAGATG ATTCGCCAC TTATTATTGT ATGCAGCATC TCGAATATCC 50
ATTCACTTTC GGTGAGGGTA CTAAAGTAGA AGTAAACGT ACGGGCCGG 99

2095926

SEQUENCE ID NO: 19.
 SEQUENCE TYPE: Nucleotide.
 SEQUENCE LENGTH: 81.
 STRANDEDNESS: Single.
 5 TOPOLOGY: Linear.
 MOLECULE TYPE: Synthetic oligonucleotide.
 PROPERTIES: Used for assembly of CDR-grafted light chain.
 FEATURES: BstBI site at residues 70 to 75.

10 GCATCTGGCA TCTGTAAGCC ACAGCAGCAG GAGTCCGAGG ACTGGGTGG 50
 GGACAGACAT GGTGGCGGCT TCGAACAGTC C 81

SEQUENCE ID NO: 20.
 SEQUENCE TYPE: Nucleotide.
 15 SEQUENCE LENGTH: 81.
 STRANDEDNESS: Single.
 TOPOLOGY: Linear.
 MOLECULE TYPE: Synthetic oligonucleotide.
 PROPERTIES: Used for assembly of CDR-grafted light chain.
 20 CCAATAGAGG AAGGTGTCAC CGTTACTATG GAGGAGACTT TTACTACTCC 50
 TACAAGTGAT GGTGACCCCTA TCACCTACAC T 81

2095926

SEQUENCE ID NO: 21.
SEQUENCE TYPE: Nucleotide.
SEQUENCE LENGTH: 102.
STRANDEDNESS: Single.
5 TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted light chain.

AGTGGCGAAA TCATCTGGCT GGAGACTACT GATAGTGAGA GTGAAC TCAG
10 TACCACTACC ACTACCACTG AATCTAGATG GTACACCACT GCCGAGGTTA
CT
50
100
102

SEQUENCE ID NO: 22.
SEQUENCE TYPE: Nucleotide.
15 SEQUENCE LENGTH: 21.
STRANDEDNESS: Single.
TOPOLOGY: Linear.
MOLECULE TYPE: Synthetic oligonucleotide.
PROPERTIES: Used for assembly of CDR-grafted light chain.
20 FEATURES: SplI site at residues 7 - 12.
CCGGCCCGTA CGTTTACTT C
21

SEQUENCE ID NO: 23.

SEQUENCE TYPE: Aminoacid.

SEQUENCE LENGTH: 133.

5 MOLECULE TYPE: Immunoglobulin light chain variable domain.

FEATURES: CDRs at residues 44 - 59, 75 - 81 and 114 - 122

[illegible]

2095926

Claims:

1. An antibody molecule having specificity for human milk fat globule (HMFG) comprising a composite heavy chain and a complementary light chain wherein:
 - the variable domain of said composite heavy chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;
 - amino acid residues 2, 26 to 35, 37, 50 to 65, 71, 73, 95 to 105, and 107 (according to the Kabat numbering system) at least in said heavy chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 1); and
 - the remaining immunoglobulin derived parts of the heavy chain are derived from a human immunoglobulin or an analogue thereof.
2. An antibody molecule having specificity for human milk fat globule (HMFG) comprising a composite heavy chain and a complementary light chain wherein:
 - the variable domain of said composite heavy chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;
 - amino acid residues 2, 26 to 35, 37, 48, 50 to 65, 67, 69, 71, 73, 95 to 105, and 107 (according to the Kabat numbering system) at least in said heavy chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 1); and
 - the remaining immunoglobulin derived parts of the heavy chain are derived from a human immunoglobulin or an analogue thereof.
3. The antibody molecule of claim 1 or claim 2, wherein additionally in said composite heavy chain at least one of residues 6, 23, 49, 76, 78, 80, 88 and 91 are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO:1).
4. The antibody molecule of any one of claims 1 to 3, wherein the complementary light chain is a composite light chain wherein:

the variable domain of said composite light chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;

amino acid residues 3, 24 to 34, 36, 50 to 56, 63, 91 to 96 and 108 (according to the Kabat numbering system) at least in said light chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2); and

the remaining immunoglobulin derived parts of the light chain are derived from a human immunoglobulin or an analogue thereof.

5. The antibody molecule of any one of claims 1 to 3, wherein the complementary light chain is a composite light chain wherein:

the variable domain of said composite light chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;

amino acid residues 3, 24 to 34, 36, 37, 45, 48, 50 to 56, 63, 91 to 96 and 108 (according to the Kabat numbering system) at least in said light chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2); and

the remaining immunoglobulin derived parts of the light chain are derived from a human immunoglobulin or an analogue thereof.

6. An antibody molecule having specificity for human milk fat globule (HMFG) comprising a composite light chain and a complementary heavy chain wherein:

the variable domain of said composite light chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;

amino acid residues 3, 24 to 34, 36, 50 to 56, 63, 91 to 96 and 108 (according to the Kabat numbering system) at least in said light chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2); and


the remaining immunoglobulin derived parts of the light chain are derived from a human immunoglobulin or an analogue thereof.



7. An antibody molecule having specificity for human milk fat globule (HMFG) comprising a composite light chain and a complementary heavy chain wherein:
 - the variable domain of said composite light chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;
 - amino acid residues 3, 24 to 34, 36, 37, 45, 48, 50 to 56, 63, 91 to 96 and 108 (according to the Kabat numbering system) at least in said light chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2); and
 - the remaining immunoglobulin derived parts of the light chain are derived from a human immunoglobulin or an analogue thereof.
8. The antibody molecule of any one of claims 4 to 7, wherein additionally in said composite light chain, residues 89, 90 and 97 are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2).
9. The antibody molecule of any one of claims 4 to 8 wherein additionally in said composite light chain, at least one of residues 1, 2, 49, 60, 70, 84, 85 and 87 are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2).
10. The antibody molecule of any one of claims 1 to 9, wherein the human framework residues in the composite light chain variable domain and/or composite heavy chain variable domain are derived from the human LAY, POM, TUR, TEI, KOL, NEWM, REI OR EU variable domain sequences.
11. The antibody molecule of claim 10, wherein the human framework residues are derived from the human EU sequence.
12. The antibody molecule of any one of claims 1 to 11, which is a complete immunoglobulin.

13. The antibody molecule of claim 12, wherein the constant region of the heavy chain is of human IgG class.
14. The antibody molecule of claim 13, wherein the
5 constant region of the heavy chain is of human IgG4 subclass.
15. The antibody molecule of claim 13, wherein the constant region of the heavy chain is of human IgG4 subclass
10 with a proline residue at position 241.
16. The antibody molecule of any one of claims 1 to 11, which is a Fab, Fab' or F(ab')₂ fragment.
- 15 17. The antibody molecule of any one of claims 12 to 16, wherein the constant domain of the light chain is of the human kappa class.
18. The antibody molecule of any one of claims 1 to 11,
20 which is an Fv or single chain Fv fragment.
19. The antibody molecule of any one of claims 1 to 18, which is produced by recombinant DNA technology.
- 25 20. A process for producing the antibody molecule of any one of claims 1 to 19, which process comprises:
- (a) producing in an expression vector an operon having a DNA sequence which encodes an antibody molecule heavy chain as defined in any one of claims 1 to 19;
 - 30 (b) producing in an expression vector an operon having a DNA sequence which encodes an antibody molecule light chain as defined in any one of claims 1 to 19;
 - (c) transfecting a host cell with the or each vector; and
 - 35 (d) culturing the transfected cell line to produce the antibody molecule.


21. The process of claim 20, for the production of an antibody fragment according to claim 16 or claim 18, in which the host cell is a bacterial host cell.
22. The process of claim 20, in which the host cell is a mammalian host cell.
23. A conjugate molecule comprising the antibody molecule of any one of claims 1 to 19 conjugated to an effector molecule or to a reporter molecule.
24. A conjugate molecule according to claim 23, wherein the effector molecule is a methyltrithio anti-tumour agent.
25. A conjugate molecule according to claim 24, wherein the methyltrithio anti-tumour agent is a disulphide analogue of the α_1 , α_2 , α_3 , α_4 , β_1 , β_2 , γ_1 , δ_1 and pseudoaglycone components of the LL-E33288 complex and derivatives thereof, BBM-1675, FR-900405, FR-900406, PD 114759, PD 115028, CL-1577A, CL-1577B, CL-1577D, CL-1577E and CL 1724 anti-tumour antibiotics and derivatives thereof.
26. A conjugate molecule according to claim 25, of the formula



prepared from a compound of formula $\text{CH}_3\text{-SSS-W}$ wherein $\text{CH}_3\text{-SSS-W}$ is an anti-tumour antibiotic designated as LL-E33288 $\alpha_1^{\text{Br}}, \alpha_1^{\text{I}}, \alpha_2^{\text{Br}}, \alpha_3^{\text{Br}}, \alpha_3^{\text{I}}, \alpha_4^{\text{Br}}, \beta_1^{\text{Br}}, \beta_1^{\text{I}}, \beta_2^{\text{Br}}, \beta_2^{\text{I}}, \gamma_1^{\text{Br}}, \gamma_1^{\text{I}}, \delta_1^{\text{I}}$, the iodo or bromo pseudoaglycone, their dihydro or N-acyl counterparts, BBM-1675, FR-900405, FR-900406, PD 114759, PD 115028, CL-1577A, CL-1577B, CL-1577D, CL-1577E, CL 1724 or their N-acetyl counterparts comprising:


reacting $\text{CH}_3\text{-SSS-W}$ with a compound of general formula Q-Sp-SH , wherein Sp is a straight or branched -chain divalent or trivalent ($\text{C}_1\text{-C}_{18}$) radical, divalent or trivalent aryl or heteroaryl radical,

divalent or trivalent (C_3 - C_{18}) cycloalkyl or heterocycloalkyl radical, divalent or trivalent aryl- or heteroaryl-alkyl (C_1 - C_{18}) radical, divalent or trivalent cycloalkyl- or heterocycloalkyl-alkyl (C_1 - C_{18}) radical or divalent or trivalent (C_2 - C_{18}) unsaturated alkyl radical, wherein if Sp is a trivalent radical, it can be additionally substituted by amino, alkylamino, arylamino, heteroarylamino, carboxyl, lower alkoxy, hydroxy, thiol, or lower alkylthio groups; and Q is, or can be subsequently converted to, halogen, amino, alkylamino, carboxyl, carboxaldehyde, hydroxy, thiol, α -haloacetyloxy, lower alkyldi-carboxyl, $-\text{CONHNH}_2$, $-\text{NHCONHNH}_2$, $-\text{NHCSNHNH}_2$, $-\text{ONH}_2$, $-\text{CON}_3$,




to produce an intermediate formula of the formula Q-Sp-SS-W, wherein Q, Sp, and W are as hereinbefore defined, reacting Q-Sp-SS-W with a molecule of the formula Hu:CT-M-O1-(Y) n wherein Hu:CT-M-O1 is humanized antibody molecule as defined in Claim 1, Claim 2, Claim 6 or Claim 7 and Y is a side-chain amino, carboxyl, or thiol group of a protein, an aldehyde derived from glycoprotein carbohydrate residues, or an amidoalkylthio group; and n is an integer of from 1 to 100, to produce a compound of the formula:

2095926




wherein Y, Sp, W, an n are as hereinbefore defined, and Z is formed from covalent reaction of the groups Q and Y directly or after subsequent reduction, and Z is -CONH-, -CONHN=CH-, -CONHNHCH₂-, -NHCSNHN=CH-, -NHCH₂-, -N=CH-, -CO₂-, -NHCH₂CO₂-, -SS-,



and m is 0.1 to 15.


27. A conjugate according to Claim 26 of the formula



prepared from the class of antitumor antibiotics designated LL-E33288 (CH₃-SSS-W) comprising:


displacing the dithiomethyl moiety with a compound of formula Q-Sp-SH, wherein Sp is straight or branched-chain divalent or trivalent (C₂-C₁₀) radicals or divalent or trivalent aryl- or heteroarylalkyl (C₂-C₅) radicals, wherein if Sp is a trivalent radical, it can be additionally substituted by amino, heteroarylamino, hydroxy, or thiol groups; and Q is, or can be subsequently converted to, carboxyl, lower alkyldicarboxyl anhydride, -CONHNH₂, or

2095926




to produce an intermediate of general formula Q-Sp-SS-W, wherein Q, Sp, and W are as hereinbefore defined,

reacting Q-Sp-SS-W with a molecule of the formula Hu:CT-M-O1-(Y)_n wherein Y is a side-chain amino group on the antibody, or an aldehyde generated by oxidation of the carbohydrate groups of the antibody, and *n* is an integer of from 1 to 100, to produce a compound of the formula:




wherein Y, Sp, W, and *n* are as hereinbefore defined, and Z is formed from covalent reaction of the groups Q and Y directly or after subsequent reduction, and Z is -CONH-, -CONHN=CH-, -CONHNHCH₂-, or



and *m* is 0.1 to 15.

28. A conjugate according to Claim 27 wherein CH₃-SSS-W is the antitumour antibiotic designated LL-E33288γ¹.
29. A conjugate according to Claim 27 wherein CH₃-SSS-W is the antitumour antibiotic designated LL-E33288α₂¹.
30. A conjugate according to Claim 27 wherein CH₃-SSS-W is the antitumour antibiotic designated LL-E33288α₃¹.

31. A conjugate according to Claim 27 wherein CH₃-SSS-W is the antitumour antibiotic designated N-acetyl LL-E33288γ¹.
32. A conjugate according to Claim 27 wherein CH₃-SSS-W is the antitumour antibiotic designated iodo LL-E33288 pseudoaglycone.
33. A conjugate according to Claim 27 wherein Q is the hydroxysuccinimide ester of a carboxyl group, Sp is -CH₂CH₂-, Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
34. A conjugate according to Claim 27 wherein Q is the hydroxysuccinimide ester of a carboxyl group, Sp is -CH₂CH(CH₃)-, Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
35. A conjugate according to Claim 27 wherein Q is the 4-nitrophenyl ester of a carboxyl group, Sp is -CH₂CH₂-, Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
36. A conjugate according to Claim 27 wherein Q is the hydroxysuccinimide ester of a carboxyl group, Sp is -CH₂C(CH₃)₂-, Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
37. A conjugate according to Claim 27 wherein Q is the hydroxysuccinimide ester of a carboxyl group, Sp is




Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.

38. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is -CH₂CH₂-, Y is -CHO, Z is -CONHN=CH-, and *m* is 0.1 to 10.
39. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is -CH₂CH₂-, Y is -CHO, Z is -CONHNHCH₂-, and *m* is 0.1 to 10.


2095926

40. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is -CH₂CH(CH₃)-, Y is -CHO, Z is -CONHN=CH-, and *m* is 0.1 to 10.
41. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is -CH₂CH(CH₃)-, Y is -CHO, Z is -CONHNHCH₂-, and *m* is 0.1 to 10.
42. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is -CH₂C(CH₃)₂-, Y is -CHO, Z is -CONHN=CH-, and *m* is 0.1 to 10.
43. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is -CH₂C(CH₃)₂-, Y is -CHO, Z is -CONHNHCH₂-, and *m* is 0.1 to 10.
44. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is




Y is -CHO, Z is -CONHN=CH-, and *m* is 0.1 to 10.

45. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is




Y is -CHO, Z is -CONHNHCH₂-, and *m* is 0.1 to 10.

46. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is




Y is -CHO, Z is -CONHN=CH-, and *m* is 0.1 to 10.

47. A conjugate according to Claim 27 wherein Q is -CONHNH₂, Sp is



Y is -CHO, Z is -CONHNHCH₂-, and *m* is 0.1 to 10.

48. A conjugate according to Claim 38 wherein CH₃-SSS-W is LL-E33288γ¹, Q is -CONHNH₂, Sp is -CH₂CH₂-, Y is -CHO, Z is -CONHN=CH-, and *m* is 0.1 to 10.
49. A conjugate according to Claim 39 wherein CH₃-SSS-W is LL-E33288α₃¹, Q is -CONHNH₂, Sp is -CH₂CH₂-, Y is -CHO, Z is -CONHNHCH₂-, and *m* is 0.1 to 10.
50. A conjugate according to Claim 33 wherein CH₃-SSS-W is N-acetyl LL-E33288γ¹, Q is hydroxysuccinimidocarbonyl, Sp is -CH₂CH₂-, Y = -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
51. A conjugate according to Claim 34 wherein CH₃-SSS-W is N-acetyl LL-E33288γ¹, Q is hydroxysuccinimidocarbonyl, Sp is -CH₂CH(CH₃)-, Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
52. A conjugate according to Claim 36 wherein CH₃-SSS-W is N-acetyl LL-E33288γ¹, Y, Q is hydroxysuccinimidocarbonyl, Sp is -CH₂C(CH₃)₂-, Y is -NH₂, Z is -CONH-, and *m* is 0.5 to 15.
53. A conjugate according to Claim 44 wherein CH₃-SSS-W is N-acetyl LL-E33288γ¹, Q is -CONHNH₂, Sp is



Y is -CHO, Z is -CONNH=CH-, and *m* is 0.1 to 10.

54. A pharmaceutical composition comprising a humanized antibody molecule according to any one of claims 1 to 19, together with a pharmaceutically acceptable excipient, diluent or carrier.
55. A pharmaceutical composition comprising a conjugate molecule according to any one of claims 23 to 53, together with a pharmaceutically acceptable excipient, diluent or carrier.
56. A pharmaceutical composition according to claim 54 or 55, for the treatment or prevention of cancer.
57. A diagnostic aid comprising an antibody molecule according to any one of claims 1 to 19.
58. A diagnostic aid according to claim 57, wherein the antibody molecule is conjugated to an effector or reporter molecule.
59. A diagnostic aid comprising a conjugate molecule according to any one of claims 23 to 53.
60. Use of an antibody molecule as claimed in any one of claims 1 to 19, for the treatment of carcinoma in a human patient.
61. Use of a conjugate molecule as claimed in any one of claims 23 to 53, for the treatment of carcinoma in a human patient.
62. Use of an antibody molecule as claimed in any one of claims 1 to 19, for the manufacture of a medicament for the treatment of carcinoma in a human patient.
63. Use of a conjugate molecule as claimed in any one of claims 23 to 53, for the manufacture of a medicament for the treatment of carcinoma in a human patient.
64. A DNA sequence which encodes the composite heavy chain of an antibody molecule having specificity for human milk fat globule

(HMFG), wherein:

the variable domain of said composite heavy chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;

amino acid residues 2, 26 to 35, 37, 50 to 65, 71, 73, 95 to 105 and 107 (according to the Kabat numbering system) at least in said heavy chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 1); and

the remaining immunoglobulin derived parts of the heavy chain are derived from a human immunoglobulin or an analogue thereof.

65. A DNA sequence which encodes the composite heavy chain of an antibody molecule having specificity for human milk fat globule (HMFG), wherein:

the variable domain of said composite heavy chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;

amino acid residues 2, 26 to 35, 37, 48, 50 to 65, 67, 69, 71, 73, 95 to 105 and 107 (according to the Kabat numbering system) at least in said heavy chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 1); and

the remaining immunoglobulin derived parts of the heavy chain are derived from a human immunoglobulin or an analogue thereof.

66. The DNA sequence of claim 64 or 65, wherein additionally in said composite heavy chain at least one of residues 6, 23, 49, 76, 78, 80, 88 and 91 are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 1).

67. A DNA sequence which encodes the composite light chain of an antibody molecule having specificity for human milk fat globule (HMFG), wherein:

the variable domain of said composite light chain comprises predominantly framework region residues from a human

immunoglobulin or an analogue thereof;

amino acid residues 3, 24 to 34, 36, 50 to 56, 63, 91 to 96 and 108 (according to the Kabat numbering system) at least in said light chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2); and

the remaining immunoglobulin derived parts of the light chain are derived from a human immunoglobulin or an analogue thereof.

68. A DNA sequence which encodes the composite light chain of an antibody molecule having specificity for human milk fat globule (HMFG), wherein:

the variable domain of said composite light chain comprises predominantly framework region residues from a human immunoglobulin or an analogue thereof;

amino acid residues 3, 24 to 34, 36, 37, 45, 48, 50 to 56, 63, 91 to 96 and 108 (according to the Kabat numbering system) at least in said light chain variable domain are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2); and

the remaining immunoglobulin derived parts of the light chain are derived from a human immunoglobulin or an analogue thereof.



69. The DNA sequence of claim 67 or 68, wherein additionally in said composite light chain, residues 89, 90 and 97 are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2).

70. The DNA sequence of any one of claims 67 to 69, wherein additionally in said composite light chain, at least one of the residues 1, 2, 49, 60, 70, 84, 85 and 87 are derived from the corresponding residues in the monoclonal antibody CTMO1 (as shown in SEQ ID NO: 2).

71. The DNA sequence of any one of claims 64 to 70, wherein the human framework residues in the composite light chain variable domain or composite heavy chain variable domain are derived from the human

LAY, POM, TUR, TEI, KOL, NEWM, REI or EU variable domain sequences.

72. The DNA sequence of any one of claims 64 to 66 or claim 71, wherein the constant region of the heavy chain is of human IgG class.
73. The DNA sequence of claim 72, wherein the constant region of the heavy chain is of human IgG4 subclass.
74. The DNA sequence of claim 72, wherein the constant region of the heavy chain is of human IgG4 subclass with a proline residue at position 241.
75. The DNA sequence of any one of claims 67 to 71, wherein the constant domain of the light chain is of the human kappa class.
76. The DNA sequence of any one of claims 64 to 66 or 71 to 74, which encodes the heavy chain of a Fab, Fab', F(ab')₂, or Fv fragment.
77. The DNA sequence of any one of claims 67 to 71 or 75, which encodes the light chain of a Fab, Fab', F(ab')₂, or Fv fragment.
78. The DNA sequence of any one of claims 64 to 75, which encodes a single chain Fv fragment.
79. A cloning or expression vector containing DNA sequence(s) according to any one of claims 64 to 78.
80. A host cell transformed with (i) DNA sequence(s) according to any one of claims 64 to 78 or (ii) a cloning or expression vector according to claim 79.



SUBSTITUTE SHEET

2095926

2/14




FIG. 3






FIG. 4



4/14




FIG. 7




FIG. 8




FIG. 9






FIG. 10



SUBSTITUTE SHEET

2095926

7/14

FIG. 13





FIG. 12



2095926

8/14

FIG. 15





FIG. 14



2095926

9/14




FIG. 16

FIG. 17





FIG. 18



SUBSTITUTE SHEET

2095926

10/14

FIG. 21





FIG. 20



2095926

11/14




FIG. 22

2095926

12/14





FIG. 23

2095926

13/14

FIG. 24



14 / 14 2095926




FIG. 25