



**(12) PATENT ABRIDGMENT (11) Document No. AU-B-73907/91**  
**(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 647464**

(54) Title  
**FUOPYRIMIDINE DERIVATIVES**

International Patent Classification(s)

(51)<sup>5</sup> C07D 491/048 A01N 043/90 A01N 055/00 C07D 307/68  
 C07D 307/84 C07D 307/85 C07D 405/12 C07D 409/12  
 C07D 409/14 C07D 413/12 C07D 417/12 C07F 007/08

(21) Application No. : 73907/91

(22) Application Date : 28.03.91

(30) Priority Data

(31) Number (32) Date (33) Country  
 502364 30.03.90 US UNITED STATES OF AMERICA

(43) Publication Date : 31.10.91

(44) Publication Date of Accepted Application : 24.03.94

(71) Applicant(s)  
**DOWELANCO**

(72) Inventor(s)  
**RONNIE GERALD EDIE; RONALD ERVIN HACKLER; ERIKS VIKTORS KRUMKAINS**


(74) Attorney or Agent  
**PHILLIPS ORMONDE & FITZPATRICK , 367 Collins Street, MELBOURNE VIC 3000**

(56) Prior Art Documents  
 US 4199584  
 US 3632763  
 EP 196524

(57) Claim

1. A compound of formula (1)

wherein



$R^1$  is H,  $C_1-C_4$  alkyl, or phenyl optionally substituted with halo,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$  alkoxy, halo  $(C_1-C_4)$  alkyl, or halo  $(C_1-C_4)$  alkoxy;

$R^2$  is H,  $C_1-C_4$  alkyl, nitro, halo; or

$R^1$  and  $R^2$  combine to form  $-(CH_2)_4-$ ;

.../2

(11) AU-B-73907/91  
(10) 647464

-2-

$R^3$  is H,  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkyl, phenyl, or substituted phenyl;

-XYZ together form 1H-1,2,4-triazol-1-yl; or

X is O, S,  $NR^4$ , or  $CR^5R^6$ , where  $R^4$  is H,  $(C_1-C_4)$  alkyl, or  $(C_1-C_4)$  acyl, and  $R^5$  and  $R^6$  are independently H,  $(C_1-C_4)$  acyl,  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  alkenyl or -alkynyl, CN, or OH, or  $R^5$  and  $R^6$  combine to form a carbocyclic ring containing four to six carbon atoms; and

~~YZ together form a  $C_5-C_{12}$  saturated or unsaturated hydrocarbon chain, straight chain or branched, optionally including a hetero atom selected from O,  $NR^4$ , S, SO,  $SO_2$ , or  $SiR^7R^8$ , where  $R^7$  and  $R^8$  are independently~~


$(C_1-C_4)$  alkyl,  $(C_3-C_4)$  branched alkyl, phenyl, or substituted phenyl, and optionally substituted with  $(C_1-C_4)$  alkyl,  $(C_2-C_4)$  alkenyl or -alkynyl, branched  $(C_3-C_7)$  alkyl,  $(C_3-C_7)$  cycloalkyl or -cycloalkenyl, halo, halo  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkoxy, hydroxy, or  $(C_1-C_4)$  acyl; ~~or~~

Y is a bond or an alkylene chain one to six carbon atoms long, ~~optionally including a carbocyclic ring, and optionally including a hetero atom selected from O,  $NR^4$ , S, SO,  $SO_2$ , or  $SiR^7R^8$ , where  $R^4$ ,  $R^7$ , and  $R^8$  are as defined above, and optionally substituted with~~  $(C_1-C_4)$  alkyl,  $(C_2-C_4)$  alkenyl or -alkynyl, branched  $(C_3-C_7)$  alkyl,  $(C_3-C_7)$  cycloalkyl or -cycloalkenyl, halo, halo  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkoxy, hydroxy, or  $(C_1-C_4)$  acyl; and

Z is

(a)  $(C_3-C_8)$  cycloalkyl or cycloalkenyl, optionally substituted with  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  alkoxy, halo  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkoxy, halo, hydroxy, or  $(C_1-C_4)$  acyl;

(b) a phenyl group of the formula (2)



where

R<sup>9</sup> to R<sup>13</sup> are independently

H,

halo,

I, \_\_\_\_\_

(C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or -cycloalkenyl,

phenoxy, or

substituted phenoxy,

phenylthio, or substituted phenylthio,

phenyl, or substituted phenyl,

NO<sub>2</sub>,

acetoxy,

OH,

CN,

SiR<sup>7</sup>R<sup>8</sup>R<sup>14</sup> or OSiR<sup>7</sup>R<sup>8</sup>R<sup>14</sup>, where R<sup>7</sup>, and R<sup>8</sup> are as defined above and R<sup>14</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>) branched alkyl, phenyl, or substituted phenyl,

NR<sup>15</sup>R<sup>16</sup>, where R<sup>15</sup> and R<sup>16</sup> are independently H, (C<sub>1</sub>-C<sub>4</sub>) alkyl, or (C<sub>1</sub>-C<sub>4</sub>) acyl,

S(O)R<sup>17</sup>, or SO<sub>2</sub>R<sup>17</sup>, where R<sup>17</sup> is (C<sub>1</sub>-C<sub>10</sub>) alkyl, phenyl, or substituted phenyl;

a C<sub>1</sub>-C<sub>12</sub> saturated or unsaturated hydrocarbon chain, straight chain or branched optionally including a hetero atom selected from O, S, SO, SO<sub>2</sub>, NR<sup>4</sup>, or SiR<sup>7</sup>R<sup>8</sup>, where R<sup>4</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above, and optionally substituted with halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy, (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or cycloalkenyl, (C<sub>1</sub>-C<sub>4</sub>) acyl, phenoxy, substituted phenoxy, phenyl, substituted phenyl, phenylthio, or substituted phenylthio;

(C<sub>1</sub>-C<sub>7</sub>) alkoxy optionally substituted with halo, phenyl, substituted phenyl, (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or cycloalkenyl, phenoxy, or substituted phenoxy;


(11) AU-B-73907/91  
(10) 647464

-4-

(C<sub>1</sub>-C<sub>7</sub>) alkylthio optionally substituted with halo, phenyl, substituted phenyl, (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or cycloalkenyl, phenoxy or substituted phenoxy; or


R<sup>10</sup> and R<sup>11</sup> combine to form -O-CF<sub>2</sub>-O- or -O-(CH<sub>2</sub>)<sub>2</sub>-O-;

(c) a furyl group of formula (3)



where R<sup>18</sup> is H, halo, halomethyl, CN, NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>) branched alkyl, phenyl, (C<sub>1</sub>-C<sub>4</sub>) alkoxy;

(d) a thienyl group of the formula (4)



where R<sup>19</sup> is R<sup>18</sup> as defined in paragraph (c) or thienyl;

(e) a group of formula (5) or (6)



where R<sup>18</sup> is as defined in paragraph (c), J is N or CH, and G is O, NR<sup>20</sup>, or S, provided that if J is not N then G is NR<sup>20</sup>, where R<sup>20</sup> is H, (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>1</sub>-C<sub>4</sub>) acyl, phenylsulfonyl, or substituted phenylsulfonyl;

(f) a group selected from

optionally substituted naphthyl, dihydronaphthyl, tetrahydronaphthyl, and decahydronaphthyl;

optionally substituted pyridyl;

optionally substituted indolyl;

1,3-benzodioxolyl;

2,6-dimethyl-4-morpholinyl; and

1-adamantyl;

or an acid addition salt of a compound of formula (1);  
provided that:

if the compound of formula (1) is a furo[2,3-d]-pyrimidin-4-amine, then Y is an alkylene chain two to six

carbon atoms long, optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, hydroxy, or acetyl, and Z is one of the groups defined in paragraphs (a) through (f).

14. A composition for use as a fungicide, insecticide or miticide which includes a compound according to any one of claims 1 to 11 in combination with a phytologically-acceptable carrier.



## FUROPYRIMIDINE DERIVATIVES

### Field of the Invention

5           This invention provides new compounds that have excellent plant fungicide activity. Some of the compounds have also demonstrated insecticidal and miticidal activity. The invention also provides compositions and combination products that contain a compound of the  
10 invention as active ingredient, as well as providing fungicidal, miticidal, and insecticidal methods.

          There is an acute need for new fungicides, insecticides, and miticides, because target pathogens are rapidly developing resistance to currently used  
15 pesticides. Widespread failure of N-substituted azole fungicides to control barley mildew was observed in 1988, and has been attributed to the development of resistance. At least 50 species of fungi have developed resistance to the benzimidazole fungicides. The field performance of  
20 DMI (demethylation inhibitor) fungicides, which are now widely relied on to protect cereal crops from powdery mildew has declined since they were introduced in the 1970's. Even recently introduced fungicides, like the acylalanines, which initially exhibited excellent control  
25 of potato late blight and grape downy mildew in the field, have become less effective because of widespread resistance. Similarly, mites and insects are developing resistance to the miticides and insecticides in current use. Resistance to insecticides in arthropods is  
30 widespread, with at least 400 species resistant to one or more insecticides. The

35

39


-1A-

GD


development of resistance to some of the older insecticides, such as DDT, the carbamates, and the organophosphates, is well known. But resistance has even developed to some of the newer pyrethroid insecticides and miticides. Therefore a need exists for new fungicides, insecticides, and miticides.

Summary of the Invention

This invention provides compounds of the formula (1):



wherein



$R^1$  is H,  $(C_1-C_4)$  alkyl, or phenyl optionally substituted with halo,  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  alkoxy, halo  $(C_1-C_4)$  alkyl, or halo  $(C_1-C_4)$  alkoxy;

$R^2$  is H,  $C_1-C_4$  alkyl, nitro, halo; or

$R^1$  and  $R^2$  combine to form  $-(CH_2)_4-$ ;

$R^3$  is H,  $C_1-C_4$  alkyl, halo  $(C_1-C_4)$  alkyl, phenyl, or substituted phenyl

-XYZ combine to form 1H-1,2,4-triazol-1-yl;

or

X is O, S, NR<sup>4</sup>, or CR<sup>5</sup>R<sup>6</sup>, where R<sup>4</sup> is H, (C<sub>1</sub>-C<sub>4</sub>) alkyl, or (C<sub>1</sub>-C<sub>4</sub>) acyl, and R<sup>5</sup> and R<sup>6</sup> are independently H, (C<sub>1</sub>-C<sub>4</sub>) acyl, alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, CN, or OH, or R<sup>5</sup> and R<sup>6</sup> combine to form a carbocyclic ring containing four to six carbon atoms; and

-YZ combine to form a C<sub>5</sub>-C<sub>12</sub> saturated or unsaturated hydrocarbon chain, straight chain or branched; optionally including a hetero atom selected from O, NR<sup>4</sup>, S, SO, SO<sub>2</sub>, or SiR<sup>7</sup>R<sup>8</sup>, where R<sup>4</sup> is as defined above and R<sup>7</sup> and R<sup>8</sup> are independently (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>) branched alkyl, phenyl, or substituted phenyl, and optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy, or (C<sub>1</sub>-C<sub>4</sub>) acyl; or

Y is a bond or a bivalent hydrocarbon radical one to six carbon atoms long, optionally including a carbocyclic ring, and optionally including a hetero atom selected from O, NR<sup>4</sup>, S, SO, SO<sub>2</sub>, or SiR<sup>7</sup>R<sup>8</sup>, where R<sup>4</sup>, R<sup>7</sup>, and R<sup>8</sup> are as defined above, and optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy, or (C<sub>1</sub>-C<sub>4</sub>) acyl; and

Z is

(a) (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or cycloalkenyl, optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>1</sub>-C<sub>4</sub>) alkoxy, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, halo, hydroxy, or (C<sub>1</sub>-C<sub>4</sub>) acyl;


35

39



(b) a phenyl group of the formula (2)

5



where

10

R<sup>9</sup> to R<sup>13</sup> are independently

H,

halo,

I,

(C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or -cycloalkenyl,

15

phenoxy, or

substituted phenoxy,

phenylthio, or substituted phenylthio,

phenyl, or substituted phenyl,

NO<sub>2</sub>,

20

acetoxy,

OH,

CN,

SiR<sup>7</sup>R<sup>8</sup>R<sup>14</sup> or OSiR<sup>7</sup>R<sup>8</sup>R<sup>14</sup>, where R<sup>7</sup>, and

25

R<sup>8</sup> are as defined above and R<sup>14</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>) branched alkyl, phenyl, or substituted phenyl.

NR<sup>15</sup>R<sup>16</sup>, where R<sup>15</sup> and R<sup>16</sup> are independently

H, (C<sub>1</sub>-C<sub>4</sub>) alkyl, or (C<sub>1</sub>-C<sub>4</sub>) acyl,

S(O)R<sup>17</sup>, or SO<sub>2</sub>R<sup>17</sup>, where R<sup>17</sup> is

30

(C<sub>1</sub>-C<sub>10</sub>) alkyl, phenyl, or substituted phenyl;

a C<sub>1</sub>-C<sub>12</sub> saturated or unsaturated hydrocarbon chain, straight chain or branched optionally including a hetero atom selected from O, S, SO, SO<sub>2</sub>, NR<sup>4</sup> or SiR<sup>7</sup>R<sup>8</sup>,

35

39


where  $R^4$ ,  $R^7$  and  $R^8$  are as defined above, and optionally substituted with halo, halo ( $C_1-C_4$ ) alkoxy, hydroxy, ( $C_3-C_8$ ) cycloalkyl or cycloalkenyl, ( $C_1-C_4$ ) acyl, phenoxy, substituted phenoxy, phenyl, substituted phenyl, phenylthio, or substituted phenylthio;

( $C_1-C_7$ ) alkoxy optionally substituted with halo, phenyl, substituted phenyl, ( $C_3-C_8$ ) cycloalkyl or cycloalkenyl, phenoxy, or substituted phenoxy;

( $C_1-C_7$ ) alkylthio optionally substituted with halo, phenyl, substituted phenyl, ( $C_3-C_8$ ) cycloalkyl or cycloalkenyl, phenoxy or substituted phenoxy; or


$R^{10}$  and  $R^{11}$  combine to form  $-O-CF_2-O-$  or  $-O-(CH_2)_2-O-$ ;

(c) a furyl group of formula (3)





where  $R^{18}$  is H, halo, halomethyl, CN,  $NO_2$ , ( $C_1-C_4$ ) alkyl, ( $C_3-C_4$ ) branched alkyl, phenyl, ( $C_1-C_4$ ) alkoxy;

(d) a thienyl group of the formula (4)



where  $R^{19}$  is  $R^{18}$  as defined in paragraph (c) or thienyl;

(e) a group of formula (5) or (6)



39

-5-



where  $R^{18}$  is as defined in paragraph (c), J is N or CH,  
and G is O,  $NR^{20}$ , or S, provided that if J is not N then  
G is  $NR^{20}$ , where  $R^{20}$  is H,  $(C_1-C_4)$  alkyl,  
( $C_1-C_4$ ) acyl, phenylsulfonyl, or substituted  
5 phenylsulfonyl;


(f) a group selected from  
optionally substituted naphthyl,  
dihydronaphthyl, tetrahydronaphthyl, and decahydronaphthyl;  
optionally substituted pyridyl;  
10 optionally substituted indolyl;  
1,3-benzodioxolyl;  
2,6-dimethyl-4-morpholinyl; and  
1-adamantyl;

or an acid addition salt of a compound of formula (1);  
15 provided that:

if the compound of formula (1) is a furo[2,3-d]-  
pyrimidin-4-amine, then Y is an alkylene chain two to six  
carbon atoms long, optionally including a carbocyclic  
ring, and optionally including a hetero atom selected from  
20 O,  $NR^4$ , S, SO,  $SO_2$ , or  $SiR^7R^8$ , where  $R^4$ ,  $R^7$ ,  
and  $R^8$  are as defined above, and optionally substituted  
with  $(C_1-C_4)$  alkyl,  $(C_2-C_4)$  alkenyl or -alkenyl,  
branched  $(C_3-C_7)$  alkyl,  $(C_3-C_7)$  cycloalkyl or  
-cycloalkenyl, halo, hydroxy, or acetyl, and Z is one of  
25 the groups defined in paragraphs (a) through (f).

Compounds of formula (1) are useful as fungicides,  
insecticides, and miticides.

The invention also provides novel compounds of the  
formula (7)




35

39

GD

where



and R<sup>3</sup> are as defined in


5 formula (1);

Y is an alkylene chain two to six carbon atoms long, optionally including a carbocyclic ring, and optionally including a hetero atom selected from O, NR<sup>4</sup>, S, SO, SO<sub>2</sub>, or SiR<sup>7</sup>R<sup>8</sup>, where R<sup>4</sup>, R<sup>7</sup>, and R<sup>8</sup> are as defined


10 above, and optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy, or (C<sub>1</sub>-C<sub>4</sub>) acyl; and

15 Z is one of the groups defined in paragraphs (a) through (f). Compounds of formula (7) are useful as fungicides, in some cases as insecticides and miticides, and also as intermediates in preparation of compounds of formula (1).

20 The invention also provides novel compounds of the formula (8)



where



and R<sup>3</sup> are as defined in

formula (1);

Y is an alkylene chain two to six carbon atoms long, optionally including a carbocyclic ring, and optionally including a hetero atom selected from O, NR<sup>4</sup>, S, SO, SO<sub>2</sub>, or SiR<sup>7</sup>R<sup>8</sup>, where R<sup>4</sup>, R<sup>7</sup>, and R<sup>8</sup> are as defined

35 above, and optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl,



(C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkenyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, hydroxy, or acetyl; and

5 Z is one of the groups defined in paragraphs (a) through (f). Compounds of formula (8) are useful as intermediates in preparation of compounds of formula (1) and (7).

The fungicide combinations of the invention, <sup>include</sup> ~~comprise~~ at least 1% by weight of a compound of formula (1) or (7) in combination with a second plant fungicide.

10 The fungicide compositions of the invention, <sup>include</sup> ~~comprise~~ a disease inhibiting and phytologically acceptable amount of compound of formula (1) or (7) in combination with a phytologically-acceptable carrier. Such compositions may optionally contain additional active ingredients, such as an additional fungicidal, miticidal, or insecticidal ingredient.

15 The fungicidal method of the invention, <sup>includes</sup> ~~comprises~~ applying to the locus of a plant pathogen a disease inhibiting and phytologically acceptable amount of a compound of formula (1) or (7).

20 The insecticide and miticide combinations of the invention, <sup>include</sup> ~~comprise~~ at least 1% by weight of a compound of formula (1) or (7) in combination with a second insecticide or miticide.

25 The insecticide and miticide compositions of the invention, <sup>include</sup> ~~comprise~~ an insect- or mite-inactivating amount of a compound of formula (1) or (7) in combination with a carrier. Such compositions may optionally contain additional active ingredients, such as an additional fungicidal, miticidal, or insecticidal ingredient.

30

35

-8-



The insecticidal or miticidal method of the invention, <sup>includes</sup> ~~comprises~~ applying to a locus of an insect or mite an insect- or mite-inactivating amount of a compound of formula (1) or (7), or of a combination described above.

5

### Detailed Description of the Invention

Throughout this document, all temperatures are given in degrees Celsius, and all percentages are weight percentages unless otherwise stated.

The term "halo" refers to a F, Cl, or Br atom.

The term "(C<sub>1</sub>-C<sub>7</sub>) alkoxy" refers to straight or branched chain alkoxy groups.

The term "(C<sub>1</sub>-C<sub>7</sub>) alkylthio" refers to straight and branched chain alkylthio groups.

The term "halo (C<sub>1</sub>-C<sub>7</sub>) alkyl" refers to a (C<sub>1</sub>-C<sub>7</sub>) alkyl group, straight chain or branched, substituted with one or more halo atoms.

The term "halo (C<sub>1</sub>-C<sub>7</sub>) alkoxy" refers to a (C<sub>1</sub>-C<sub>7</sub>) alkoxy group substituted with one or more halo groups.

The term "halo (C<sub>1</sub>-C<sub>4</sub>) alkylthio" refers to a (C<sub>1</sub>-C<sub>4</sub>) alkylthio group, straight chain or branched, substituted with one or more halo atoms.

The term "substituted phenyl" refers to phenyl substituted with up to three groups selected from halo, I, (C<sub>1</sub>-C<sub>10</sub>) alkyl, branched (C<sub>3</sub>-C<sub>6</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, hydroxy (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>1</sub>-C<sub>4</sub>) alkoxy, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, phenoxy, substituted phenoxy, phenyl, substituted phenyl, NO<sub>2</sub>, OH, CN, (C<sub>1</sub>-C<sub>4</sub>) alkanoyloxy, or benzyloxy.

35

39

-9-



The terms "substituted naphthyl", "substituted pyridyl" and "substituted indolyl" refer to these ring systems substituted with halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, CN, NO<sub>2</sub>, (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>) branched alkyl, phenyl, (C<sub>1</sub>-C<sub>4</sub>) alkoxy, or halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy.

The term "substituted phenoxy" refers to phenoxy substituted with up to three groups selected from halo, I, (C<sub>1</sub>-C<sub>10</sub>) alkyl, branched (C<sub>3</sub>-C<sub>6</sub>) alkyl, halo (C<sub>1</sub>-C<sub>7</sub>) alkyl, hydroxy (C<sub>1</sub>-C<sub>7</sub>) alkyl, (C<sub>1</sub>-C<sub>7</sub>) alkoxy, halo (C<sub>1</sub>-C<sub>7</sub>) alkoxy, phenoxy, substituted phenoxy, phenyl, substituted phenyl, NO<sub>2</sub>, OH, CN, (C<sub>1</sub>-C<sub>4</sub>) alkanoyloxy, or benzyloxy.

The term "carbocyclic ring" refers to a saturated or unsaturated carbocyclic ring containing three to seven carbon atoms.

The terms "substituted phenylthio" and "substituted phenyl sulfonyl" refer to such groups substituted with up to three groups selected from halo, I, (C<sub>1</sub>-C<sub>10</sub>) alkyl, branched (C<sub>3</sub>-C<sub>6</sub>) alkyl, halo (C<sub>1</sub>-C<sub>7</sub>) alkyl, hydroxy (C<sub>1</sub>-C<sub>7</sub>) alkyl, (C<sub>1</sub>-C<sub>7</sub>) alkoxy, halo (C<sub>1</sub>-C<sub>7</sub>) alkoxy, phenoxy, substituted phenoxy, phenyl, substituted phenyl, NO<sub>2</sub>, OH, CN, (C<sub>1</sub>-C<sub>4</sub>) alkanoyloxy, or benzyloxy.

The term "unsaturated hydrocarbon chain" refers to a hydrocarbon chain containing one or two sites of unsaturation.

The term "HPLC" refers to a high pressure liquid chromatography.

The term "bivalent hydrocarbon radical" refers to bivalent radicals derived from normal alkanes by removal of hydrogen atoms from each of the two terminal carbon atoms of the chain, e.g. methylene, ethylene, trimethylene, tetramethylene, etc.


35

39

Compounds

Compounds of formula (1) include furo(2,3-d)-  
pyrimidines of formula (1a):


5



10

furo(3,2-d)pyrimidines of formula (1b)


15



20

and furo(3,4-d)pyrimidines of formula (1c)

25



30

While all of the compounds of the invention are  
useful fungicides, certain classes are preferred for  
reasons of greater efficacy or ease of synthesis, viz:

- a) furo[2,3-d]pyrimidines of formula (1a);
- b) 5-methylfuro[2,3-d]pyrimidines of formula (1a);
- c) 6-methylfuro[2,3-d]pyrimidines of formula (1a);

35

39

GD

d) 5,6-dimethylfuro[2,3-d]pyrimidines of formula (1a);

e) compounds of formula (1) wherein Y is an alkylene chain two to six carbon atoms long;

5 f) compounds of formula (1) wherein Y is  $-(CH_2)_2-$ ;

g) compounds of formula (1) wherein Z is phenyl;

h) compounds of formula (1) wherein Z is a substituted phenyl group of formula (2);

10 i) compounds of formula (1) wherein Z is a phenyl group of formula (2) in which one of the substituents  $R^9$  to  $R^{13}$  is a haloalkoxy group;

j) compounds of formula (1) wherein Z is 2-naphthyl;

15 k) compounds of formula (1) wherein Z is 3-( $C_1-C_4$ )alkoxyphenyl;

l) compounds of formula (1) wherein Z is 3-methoxyphenyl;

20 m) compounds of formula (1) wherein Z is 4-( $C_1-C_4$ )alkoxyphenyl;

n) compounds of formula (1) wherein Z is 4-methoxyphenyl;

o) compounds of formula (1) wherein Z is 3-(halo( $C_1-C_4$ )alkyl)phenyl;

25 p) compounds of formula (1) wherein Z is 3-(Trifluoromethyl)phenyl;

q) compounds of formula (1) wherein Z is 4-(halo( $C_1-C_4$ )alkyl)phenyl;

30

35

39

r) compounds of formula (1) wherein Z is 4-(trifluoromethyl)phenyl;

s) compounds of formula (1) wherein Z is 4-(1,1,2,2-tetrafluoroethoxy)phenyl;

5 t) compounds of formula (1) wherein Z is phenyl substituted with a branched (C<sub>3</sub>-C<sub>7</sub>)alkyl group;

u) compounds of formula (1) wherein Z is 4-(t-butyl)phenyl;

10 v) compounds of formula (1) wherein Z is phenyl substituted with a (C<sub>1</sub>-C<sub>4</sub>)alkyl group;

w) compounds of formula (1) wherein Z is 4-(n-butyl)phenyl;

x) compounds of formula (1) wherein Z is phenyl substituted with a halo atom;

15 y) compounds of formula (1) wherein Z is 4-chlorophenyl;

z) compounds of formula (1) wherein Z is 4-fluorophenyl.

20


Synthesis

The compounds of this invention are made using well known chemical procedures. The required starting materials are commercially available, or they are readily  
25 synthesized using standard procedures.

Synthesis of Compounds Wherein X is NR<sup>4</sup>

Compounds of formula (1) wherein X is NH are represented by the following formula (1d):


5



10

wherein each of the variable substituents are as defined above for formula (1). Compounds of formula (1d) wherein R<sup>3</sup> is H can be prepared by the procedure shown in the following Scheme 1:

15



20

25

Scheme 1 represents an application of the general procedure that is described in Edward C. Taylor and Alexander McKillop, The Chemistry of Cyclic Enaminonitriles and o-Aminonitriles (Interscience 1970).

5 Application of this procedure to preparation of furo[2,3-d]pyrimidines is described in Chemical Abstracts 80:96102f (1974), in U.S. 4,199,584 to Cox et al., and in Chemica Scripta 26, 337-342 (1986).

In step 1 of Scheme 1 an o-aminonitrile (10) is  
10 condensed with an orthoester of the formula  $\text{HC}(\text{OC}_2\text{H}_5)_3$  or  $\text{HC}(\text{OCH}_3)_3$ . The reaction is optionally carried out in the presence of acetic anhydride.

In step 2 of Scheme 1, the imino ether (9) produced  
15 in step 1 is reacted with a primary amine of the formula  $\text{NH}_2\text{-Y-Z}$ , where Y and Z are as defined in formula (1). The preferred solvent is ethyl alcohol. If step 2 is carried out at room temperature, the novel o-cyano-amidine of formula (8) can be isolated. At elevated temperatures cyclization of the o-cyanoamidine (step 3) proceeds,  
20 delivering the imine of formula (7).

In step 4, Dimroth rearrangement of the imine of  
formula (7) provides the N-substituted-4-amine of formula (1d). Rearrangement is favored by basic conditions or elevated temperatures.

25 o-Aminonitriles of formula (10), used as starting materials in Scheme 1, are prepared using known procedures, e.g., those described in Acta Chemica Scandinavica B29 224-232 (1975) (3-amino-2-cyanofurans); U.S. Patent No. 4,240,820 (2-amino-3-cyanofurans).


30

35

39

Compounds of formula (1d) wherein R<sup>3</sup> is other than H can be prepared by the general procedure described in Chemica Scripta 25, 227-229 (1985), as illustrated in Scheme 2:

5




In step 1 of Scheme 2, the o-(acylamino)nitrile of formula (21) is reacted with phosphorus pentoxide, a tertiary amine such as triethylamine or N,N-dimethylcyclo-hexylamine (DMCA), and an amine hydrochloride of the formula Z-Y-NH<sub>3</sub>.HCl to give the amidine intermediate of formula (8), which can cyclize, as in Scheme 1 to give

39


GD

the imino compound of formula (7). Upon heating, particularly in the presence of base, the imino compounds rearrange to provide the N-substituted-4-amino compounds of formula (1d) (step 3).

5 Compounds of formula (1) wherein X is NH can also be prepared by the process disclosed in Chemica Scripta 26, 337-342 (1986), illustrated in Scheme 3:



30 Compounds of formula (1) wherein X is  $\text{NR}^4$  and  $\text{R}^4$  is H or  $(\text{C}_1\text{-C}_4)$  alkyl may also be prepared by aminating a suitably substituted intermediate of the formula (23)



where L is a leaving group, such as F, Cl, Br, I, NO<sub>2</sub>,  
1,2,4-triazol-1-yl, O-SiMe<sub>3</sub>, arylthio, alkylthio,  
alkylsulfonyl, arylsulfonyl, alkoxy, or arylsulfinyl. The  
general procedure is described in, e.g. Bull. Soc. Chim.  
5 Fr. 12, 4344 (1969); Diss. Abstr. Int. B32 (12), 6915  
(1972); and Ger. Offen 1,817,146; U.S. 4,199,584 to Cox,  
and U.S. 4,007,187 to Fauran et al. A chloride of formula  
(23) wherein L is Cl is allowed to react with an  
appropriate amine at a wide variety of temperatures  
10 (20-180°), preferably in the presence of an acid  
acceptor, such as triethylamine. The reaction may be  
carried out neat, or in a non-reactive organic solvent.  
Compounds where R<sup>4</sup> is acyl are prepared from amines  
where R<sup>4</sup> is H, which were allowed to react with an  
15 acylating agent such as acetyl chloride or acetic  
anhydride.

Preparation of 4-chloro compounds of formula (23) is  
illustrated in U.S. 3,632,763 and U.S. 3,577,420.

#### 20 Synthesis of Compounds Wherein X is O

The compounds of formula (1) wherein X is O are made  
by condensing a compound of formula (23) with an alcohol  
or phenol of the formula (24):



where

Y and Z are as previously defined.


The reaction is preferably carried out in the presence of a strong base, such as sodium hydride, in a non-reactive organic solvent, such as DMF, at a temperature in the range of 0 to 25°C.

5 Compounds wherein X is S are made by the same procedure using corresponding mercaptans of formula HS-Y-Z.

Synthesis of Compounds Wherein X is CH<sub>2</sub>

10 The compounds of formula (1) wherein X is CH<sub>2</sub> can be made using the process described in the J. Heterocyclic Chemistry, Vol. 14, 1081-1083 (1977).

This procedure involves hydrolysis and decarboxylation of 5-substituted-5-(4-fuopyrimidyl)-barbituric acids of the formula



25 wherein the variable substituents are as defined for formula (1). The 5-substituted-5-(4-fuopyrimidyl)-barbituric acid is dissolved in a solution of sodium hydroxide and water and refluxed. The solution is then made slightly acidic and again refluxed.

30 Compounds of formula (1) wherein X is CR<sup>5</sup>R<sup>6</sup> may also be prepared by reacting a compound of formula (23) where L is H with a Grignard reagent of the formula Z-Y-CR<sup>5</sup>R<sup>6</sup>-MgX' or a lithio reagent of the formula Z-Y-

35


39

CR<sup>5</sup>R<sup>6</sup>-Li, where X' is halo, to provide a 3,4-dihydrofuroprymidine which is then oxidized to provide a compound of the invention. Typical reaction conditions are those described in Armarego and Smith, J. Chem. Soc., page 5360 (1965).


The acid addition salts and N-oxides of compounds of formula (1) are obtained in the usual way.

Accordingly, the invention also provides a process for preparing a compound of formula (1) which comprises

(a) condensing a compound of formula (23)




wherein R is as previously defined, and L is a leaving group, with an alcohol of the formula (8):



wherein Y and Z are as previously defined to produce a compound of formula (1) wherein X is O; or

(b) condensing a compound of formula (23) as defined above with an amine of the formula (9)




wherein  $R^4$  is H or  $(C_1-C_4)$  alkyl, and Y and Z are as previously defined, to provide a compound of formula (1) where X is  $NR^4$ ; or

5 (c) acylating a compound of formula (1) wherein X is NH to provide a compound of formula (1) wherein X is  $NR^4$  and  $R^4$  is acyl; or


(d) reacting a compound of formula (23) wherein L is H with a Grignard reagent of formula  $Z-Y-CR^5R^6-MgX'$ , where  $X'$  is halo and Z and Y are as defined above, or with a lithio reagent of formula  $Z-Y-CR^5R^6-Li$ , to provide a 3,4-dihydro furopyrimidine, which is then oxidized to produce a compound of formula (1) where X is  $CR^5R^6$ ; or

15 (e) hydrolyzing and decarboxylating a 5-substituted-5-(4-(furopyrimidinyl)barbituric acid of the formula



wherein R,  $R^3$ ,  $R^5$ ,  $R^6$ , Y, and Z are as previously defined to provide a compound of formula (1) wherein X is  $CR^5R^6$ ; or

25 (f) heating a compound of the formula (7)




35

39

optionally in the presence of base to provide a compound of the formula (1d)

5



10

Examples 1-89

15

The following examples are compounds actually prepared by the above described general procedures. The melting point is given for each compound. In addition, although the data has not been included, each compound was fully characterized by one or more of NMR, IR, mass spectra, and combustion analysis. Specific illustrative preparations for the compounds of representative compounds follow the tabular listing.

25

30

35

39

GD

<u>EXAMPLE NUMBER</u>	<u>COMPOUND</u>	<u>M.P.</u>
1	N-[2-(4-methoxyphenyl)ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	150-151°C
5	2 5-methyl-N-[2-(4-methoxyphenyl)-ethyl]furo[2,3-d]pyrimidin-4-amine	154°C
	3 5,6-dimethyl-N-[2-(4-ethoxyphenyl)-ethyl]furo[2,3-d]pyrimidin-4-amine	151°C
10	4 N-[2-(4-methoxyphenyl)ethyl]-6-methylfuro[2,3-d]pyrimidin-4-amine	186°C
	5 6-methyl-4-(4-fluorophenoxy)furo[2,3-d]pyrimidine	105° dec.
15	6 6-methyl-N-[2-(4-chlorophenyl)ethyl]-furo[2,3-d]pyrimidin-4-amine	180°C
20	7 4-[2-(1,1'-biphenyl)-4-ylethoxy]6-methylfuro[2,3-d]pyrimidine	124-125°C
	8 6-methyl-N-[2-[3-(trifluoromethyl)-phenyl]ethyl]furo[2,3-d]pyrimidin-4-amine	119°C
25	9 6-methyl-N-[2-(4-ethoxyphenyl)ethyl]-furo[2,3-d]pyrimidin-4-amine	164°C
	10 N-[2-(4-chlorophenyl)ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	145°C
30	11 5-methyl-N-[2-(4-chlorophenyl)ethyl]-furo[2,3-d]pyrimidin-4-amine	140°C
	12 5,6-dimethyl-N-(2-phenylethyl)-furo[2,3-d]pyrimidin-4-amine	135°C
35	13 5,6-dimethyl-4-[3-(3-pyridinyl)-propoxy]furo[2,3-d]pyrimidine	75°C
39		

<u>EXAMPLE NUMBER</u>	<u>COMPOUND</u>	<u>M.P.</u>
14	5,6-dimethyl-N-[2-[[2-chloro-6-fluorophenyl)methyl]thio]ethyl]-furo[2,3-d]pyrimidin-4-amine	147°C
5		
15	N-[2-(4-methoxyphenyl)ethyl]furo[2,3-d]pyrimidin-4-amine	135-136°C
16	5-methyl-N-[2-[[2-chloro-6-fluorophenyl)methyl]thio]ethyl]-furo[2,3-d]pyrimidin-4-amine	167°C
10		
17	5,6-dimethyl-4-(1-azabicyclo[2.2.2]oct-3-yloxy)furo[2,3-d]pyrimidine	105°C
15		
18	N-[2-(decahydro-2-naphthyl)ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	138°C
19	5-methyl-N-(2-phenylethyl)furo[2,3-d]pyrimidin-4-amine	132-133°C
20		
20	N-[2-[4-(t-butyl)phenyl]ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	148-150°C
21	5-methyl-N-[2-(2-naphthyl)ethyl]-furo[2,3-d]pyrimidin-4-amine	141-142°C
22	5,6-dimethyl-N-[2-(2-naphthyl)ethyl]-furo[2,3-d]pyrimidin-4-amine	125°C
30		
23	5,6-dimethyl-N-[2-[3-(trifluoromethyl)phenyl]ethyl]furo[2,3-d]pyrimidin-4-amine	139-140°C
24	5,6-dimethyl-N-[2-(4-methylphenyl)ethyl]furo[2,3-d]pyrimidin-4-amine	140°C
35		
25	5,6-dimethyl-N-[2-(3,4-dimethoxyphenyl)ethyl]furo[2,3-d]pyrimidin-4-amine	171°C

39

GD

	<u>EXAMPLE NUMBER</u>	<u>COMPOUND</u>	<u>M.P.</u>
	26	N-[2-(4-chlorophenyl)-3-(4-fluorophenyl)propyl]- 5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	147°C
5	27	N-[2-(3,4-dimethoxyphenyl)ethyl]-5- methylfuro[2,3-d]pyrimidin-4-amine	164°C
	28	5,6-dimethyl-4-(2-methoxyphenoxy)- furo[2,3-d]pyrimidine	121-122°C
10	29	5,6-dimethyl-4-(2-methylphenoxy)- furo[2,3-d]pyrimidine	78-79°C
	30	N-[2-(4-methoxyphenyl)ethyl]-5,6,7,8-tetra- hydrobenzofuro[2,3-d]pyrimidin-4-amine	135°C
15	31	6-methyl-N-[2-(2-naphthalenyl)ethyl]- furo[2,3-d]pyrimidin-4-amine	127°C
20	32	5,6-dimethyl-4-[4-(trifluoromethoxy)- phenoxy]furo[2,3-d]pyrimidine	105-106°C
	33	5-methyl-N-[2-(3-methoxyphenyl)- ethyl]furo[2,3-d]pyrimidin-4-amine	116°C
25	34	4-(4-fluorophenoxy)-5,6,7,8-tetra- hydrobenzofuro[2,3-d]pyrimidine	96-97°C
	35	5-methyl-N-[2-(2-methoxyphenyl)- ethyl]furo[2,3-d]pyrimidin-4-amine	122°C
30	36	4-[(5,6-dimethylfuro[2,3-d]pyrimidin- 4-yl)oxy]benzotrile	N/A
	37	2-[(5,6-dimethylfuro[2,3-d]pyrimidin- 4-yl)oxy]benzotrile	225°C
35	38	6-methyl-4-[2-(1-adamantyl)ethoxy]- furo[2,3-d]pyrimidine	99-100°C
39			

	<u>EXAMPLE NUMBER</u>	<u>COMPOUND</u>	<u>M.P.</u>
	39	6-methyl-4-[2-[4-(phenylmethoxy)-phenyl]ethoxy]furo[2,3-d]pyrimidine	90°C
5	40	6-methyl-4-[2-(4-methoxyphenyl)-ethoxy]furo[2,3-d]pyrimidine	75°C
	41	5,6-dimethyl-4-[2-[4-[4-(trifluoromethyl)-phenoxy]phenyl]ethoxy]furo[2,3-d]pyrimidine	oil
10	42	6-methyl-4-[2-[4-(trifluoromethyl)-phenyl]ethoxy]furo[2,3-d]pyrimidine	73°C
	43	6-methyl-4-[2-[4-(t-butoxy)phenyl]-ethoxy]furo[2,3-d]pyrimidine	75°C
15	44	4-[2-(4-butoxyphenyl)ethoxy]-6-methylfuro[2,3-d]pyrimidine	60-61°C
20	45	5,6-dimethyl-4-[2-[4-(n-butoxy)-phenyl]ethoxy]furo[2,3-d]pyrimidine	60°C
	46	4-[2-[4-(1,1-dimethylethyl)phenyl]-ethoxy]-6-methylfuro[2,3-d]pyrimidine	N/A
25	47	N-[2-[4-(1,1,2,2-tetrafluoroethoxy)phenyl]-ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	111°C
	48	5-methyl-N-[2-(4-methylphenyl)ethyl]-furo[2,3-d]pyrimidin-4-amine	139°C
30	49	5,6-dimethyl-4-(4-fluorophenoxy)furo[2,3-d]pyrimidine	126°C
	50	4-[2-[4-(t-butyl)phenyl]ethoxy]-5,6-dimethylfuro[2,3-d]pyrimidine	107-108°C
35	51	4-[2-[4-biphenyl]ethoxy]-5,6-dimethylfuro[2,3-d]pyrimidine	83°C
39			

	<u>EXAMPLE NUMBER</u>	<u>COMPOUND</u>	<u>M.P.</u>
	52	4-[2-(4-methoxyphenyl)ethoxy]-5,6-dimethylfuro[2,3-d]pyrimidine	69-70°C
5	53	4-[2-(4-chlorophenyl)ethoxy]-5,6-dimethylfuro[2,3-d]pyrimidine	84°C
	54	5,6-dimethyl-4-[2-[4-(trifluoromethyl)-phenyl]ethoxy]furo[2,3-d]pyrimidine	74-75°C
10	55	5,6-dimethyl-4-[2-(2-naphthyl)-ethoxy]furo[2,3-d]pyrimidine	85°C
	56	5,6-dimethyl-4-[(4-fluorophenyl)-thio]furo[2,3-d]pyrimidine	115°C
15	57	Cis-5,6-dimethyl-4-[2-(2,6-dimethyl-4-morpholinyl)ethoxy]furo[2,3-d]pyrimidine	71-74°C
20	58	Trans-5,6-dimethyl-4-[2-(2,6-dimethyl-4-morpholinyl)ethoxy]furo[2,3-d]pyrimidine	164-165°C
	59	4-(2-chloro-4-fluorophenoxy)-5,6-dimethylfuro[2,3-d]pyrimidine	164-165°C
25	60	5,6-dimethyl-4-[(2-methyloxolan-5-yl)methoxy]furo[2,3-d]pyrimidine	49-50°C
	61	5,6-dimethyl-4-[2-(2-thienyl)-ethoxy]furo[2,3-d]pyrimidine	72°C
30	62	5,6-dimethyl-4-(4-methylphenoxy)-furo[2,3-d]pyrimidine	103°C
35	63	5,6-dimethyl-4-[3-(trifluoromethyl)phenoxy]furo[2,3-d]pyrimidine	125°C
	64	5,6-dimethyl-4-phenoxyfuro[2,3-d]pyrimidine	74°C
39			

	<u>EXAMPLE NUMBER</u>	<u>COMPOUND</u>	<u>M.P.</u>
	65	5,6-dimethyl-4-(4-methoxyphenoxy)-furo[2,3-d]pyrimidine	125-126°C
5	66	4-(2-fluorophenoxy)-5,6-dimethyl-furo[2,3-d]pyrimidine	87°C
	67	4-[2-[4-(2,2-dimethylethyl)phenyl]-2-methoxy]5,6-dimethylfuro[2,3-d]pyrimidine	oil
10	68	5,6-dimethyl-N-[2-[4-(trifluoromethoxy)phenyl]ethyl]furo[2,3-d]pyrimidin-4-amine	N/A
	69	5,6-dimethyl-4-[2-[4-(trifluoromethoxy)phenyl]ethoxy]furo[2,3-d]pyrimidine	78°C
15	70	5,6-dimethyl-4-[2-[3,4-difluoromethylene-dioxy]phenyl]ethoxy]furo[2,3-d]pyrimidine	95°C
	71	5,6-dimethyl-N-[3-[4-(n-hexyloxy)phenyl]-2-(1-naphthyl)propyl]furo[2,3-d]pyrimidin-4-amine	105°C
	72	5,6-dimethyl-N-[2-(1-cyclohexenyl)-ethyl]furo[2,3-d]pyrimidin-4-amine	124°C
25	73	5,6-dimethyl-N-[2-[4-(2,2,2-trifluoroethoxy)phenyl]ethyl]furo[2,3-d]pyrimidin-4-amine	155-156°C
	74	5,6-dimethyl-4-[2-[4-(1-ethyl-1-methylbutoxy)phenyl]ethoxy]furo[2,3-d]pyrimidine	oil
30	75	5,6-dimethyl-4-[2-[benzo[b]-1,4-dioxan-6-yl]ethoxy]furo[2,3-d]pyrimidine	146°C
35	76	5,6-dimethyl-4-[2-[2(trifluoromethyl)phenyl]ethoxy]furo[2,3-d]pyrimidine	84°C
39			

EXAMPLE NUMBER	COMPOUND	M.P.
5	77 5,6-dimethyl-N-[1,1-dimethyl-2-phenylethyl]furo[2,3-d]pyrimidin-4-amine	59°C
	78 5,6-dimethyl-4-[4-(2,2,2-trifluoroethoxy)phenyl]ethoxy]furo[2,3-d]pyrimidine	72-73°C
10	82 N-[2-(4-phenoxyphenyl)ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	104°C
	83 5,6-dimethyl-N-[2-[4-(trifluoromethyl)phenyl]ethyl]-furo[2,3-d]pyrimidin-4-amine	146°C
15	84 5,6-dimethyl-4-[2-(4-fluorophenoxy)ethoxy]furo[2,3-d]pyrimidine	98-99°C
	85 5,6-dimethyl-N-[2-[4-(1-methylethyl)phenyl]ethyl]furo[2,3-d]pyrimidin-4-amine	122-124°C
20	86 5,6-dimethyl-4-[2-[4-(trimethylsilyl)phenyl]ethoxy]-furo[2,3-d]pyrimidine	80°C
	87 5,6-dimethyl-4-[2-[3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]ethyl]furo[2,3-d]pyrimidine	85°C

25

The following detailed examples illustrate the procedures used to prepare the compounds of formula (1).

Example 2

30

5-Methyl-N-[2-(4-methoxyphenyl)ethyl]furo[2,3-d]pyrimidin-4-amine

A mixture of 4 g of 2-amino-3-cyano-4-methylfuran, 3 mL of triethylorthoformate and 0.5 mL of acetic anhydride was heated at 135° for three hours. To the

hot solution was added 4.5 g of 2-(4-methoxyphenyl)ethylamine, 5 mL of acetic anhydride, and 3.1 g of sodium acetate. The temperature of the mixture was kept at 135°C for three additional hours, then the mixture was cooled and combined with a mixture of water and ether. The ether layer was separated, and product was extracted from the remaining aqueous layer with large volumes of ether. The ether extracts were combined, dried over magnesium sulfate, filtered, and concentrated. The residue was recrystallized from acetone/ether. Yield 2.6 g. M.P. 184°C

5

10

15

20

25

30

Example 8

6-Methyl-N-[2-[3-(trifluoromethyl)phenyl]ethyl]-  
furo[2,3-d]pyrimidin-4-amine

5 A mixture of 1.9 g of 2-[3-(trifluoromethyl)phenyl]-  
ethylamine and 1.6 g of 4-chloro-6-methylfuro[2,3-d]-  
pyrimidine was heated for ten minutes at 230°C. The  
mixture was then cooled and the product extracted in ethyl  
acetate/water. The organic phase was separated, washed  
with water, dried over magnesium sulfate, filtered, and  
10 concentrated. The residue was chromatographed on a silica  
column using toluene and acetone as eluent.  
Yield 0.97 g. M.P. 119°C.

Example 11

15 5-Methyl-N-[2-(4-chlorophenyl)ethyl]furo[2,3-d]-  
pyrimidin-4-amine

A mixture of 4.0 g of 2-amino-3-cyano-4-methylfuran,  
3 cc of triethylorthoformate, and .05 cc of acetic  
anhydride was heated to 135°C for three hours. Then 5 g  
20 of 2-(4-chlorophenyl)ethylamine, 5 cc of glacial acetic  
acid, and 3.2 g of sodium acetate were added to the  
reaction mixture. The temperature of the mixture was  
maintained at 135°C for three additional hours, after  
which the mixture was cooled, treated with water and  
25 extracted with dichloromethane. The organic phase was  
then concentrated and washed with water. The solids were  
extracted from the organic phase with large volumes of  
ether, which were combined and concentrated to provide  
1.07 g of product. M.P. 140°C.

30

35

39

Example 15

N-[2-(4-methoxyphenyl)ethyl]furo[2,3-d]-  
pyrimidin-4-amine

5 a. A mixture of 5.8 g of malononitrile, 4.8 mL of  
triethylamine, and 3.8 g of glycoaldehyde was refluxed for  
10 minutes in 100 cc of toluene. Then the mixture was  
cooled and the toluene layer decanted. The dark residue  
was extracted with fresh toluene several additional  
times. The extracts were combined, washed with water,  
10 dried over magnesium sulfate, and concentrated under  
vacuum to produce 2-amino-3-cyanofuran as an oil.

b. The 2-amino-3-cyanofuran produced in the  
preceding step was combined with 4.2 mL of triethylortho-  
formate, and 0.5 mL of acetic anhydride, and the mixture  
15 was heated to 130°C for two hours. Then was added 1.8 g  
of sodium acetate and 3 g of 2-(4-methoxyphenyl)ethyl-  
amine, followed by 3 mL of acetic acid. The resulting  
mixture was heated for two hours at 130-138°C, then cooled  
and diluted with water. The product was extracted with  
20 large volumes of ether. The ether extracts were combined,  
washed with water, dried over magnesium sulfate, and  
concentrated. The residue was recrystallized from ether  
and petroleum ether. Yield 0.65 g. M.P. 135-136°C.

25

Example 43

4-[2-[4-(t-butoxy)phenyl]ethoxy-6-methyl-  
furo[2,3-d]pyrimidine

To a dispersion of 0.3 g of 60% sodium hydride in  
mineral oil in 25 mL of DMF was added 1.13 g of 2-[4-(t-  
30 butoxy)phenyl]ethanol, and the mixture was agitated

35

39

for two hours. Then 0.9 g of 4-chloro-6-methylfuro[2,3-d]pyrimidine was added, and the mixture was agitated at room temperature overnight, after which the mixture was poured on ice. Gray solids precipitated and were collected by filtration and dried. Yield 0.76 g. M.P. 75°C.

Example 44

4-[2-[4-(n-butoxy)phenyl]ethoxy]-6-methylfuro[2,3-d]pyrimidine

To a dispersion of 0.52 g of sodium hydride in 35 mL of DMF was added 2 g of 2-[4-(n-butoxy)phenyl]ethanol at room temperature to provide the sodium salt of the alcohol. To this mixture was added 6.5 g of 4-chloro-6-methylfuro[2,3-d]pyrimidine, and the resulting mixture was agitated at room temperature overnight, then poured on ice. Pale yellow solid precipitated and were isolated by filtration, dried, and recrystallized first from ether, and then from a mixture of ether and petroleum ether. Yield 1.2 g. M.P. 60-61°C.

Example 52

4-[2-(4-methoxyphenyl)ethoxy]-5,6-dimethylfuro[2,3-d]pyrimidine

To a dispersion of 0.48 g of 60% sodium hydride in mineral oil in 50 mL of DMF was added 1.5 g of 2-(4-methoxyphenyl)ethanol to produce the sodium salt of the alcohol. To this mixture was added 1.2 g of 4-chloro-5,6-dimethylfuro[2,3-d]pyrimidine in small portions at room temperature and the dispersion was allowed to react overnight. The reaction mixture was concentrated under

vacuum and the residue dispersed in water. The product was extracted into large volumes of ether, which were combined, dried over magnesium sulfate, filtered, and concentrated. The residue was recrystallized from ether and petroleum ether. M.P. 69-70°C.

5

The following compounds of formula (7) were prepared and isolated.

EXAMPLE NUMBER	COMPOUND	M.P.	
10	79	5-methyl-3-[2-(2-naphthyl)ethyl]-furo[2,3-d]pyrimidin-4(3H)-imine	145°C
	80	3-[2-(4-methoxyphenyl)ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4(3H)-imine	125°C
15	81	3-[2-[3-(trifluoromethyl)phenyl]ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4(3H)-imine	148-149°C
	88	3-[2-[4-(1,1,2,2-tetrafluoroethoxy)phenyl]ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4(3H)-imine	133-134°C
20	89	3-[2-(4-phenoxyphenyl)ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4(3H)-imine	130-132°C

The following detailed preparation illustrates the general procedure used to prepare compounds of formula (7) (see also Schemes 1 and 2, supra).

Example 80

30 3-[2-(4-methoxyphenyl)ethyl]-5,6-dimethyl-furo[2,3-d]pyrimidin-4(3H)-imine

2-Amino-4,5-dimethyl-3-cyanofuran and trimethylorthoformate were refluxed for one hour. The solution was then concentrated under vacuum. To the solution was added 3 g of 4-methoxy-phenethylamine dissolved in 10 mL

of benzene. After setting overnight at room temperature, the mixture was concentrated under vacuum. The residue was dissolved in acetone/ether and cooled. A yellow precipitate formed, which was isolated, digested in ether and air dried. The solids were again digested in acetone/ether, and filtered. M.P. 125°C.

#### Fungicide Utility

The compounds of the present invention have been found to control fungi, particularly plant pathogens. When employed in the treatment of plant fungal diseases, the compounds are applied to the plants in a disease inhibiting and phytologically acceptable amount. The term "disease inhibiting and phytologically acceptable amount," as used herein, refers to an amount of a compound of the invention which kills or inhibits the plant disease for which control is desired, but is not significantly toxic to the plant. This amount will generally be from about 1 to 1000 ppm, with 10 to 500 ppm being preferred. The exact concentration of compound required varies with the fungal disease to be controlled, the type formulation employed, the method of application, the particular plant species, climate conditions and the like. A suitable application rate is typically in the range from .25 to 4 lb/A. The compounds of the invention may also be used to protect stored grain and other non-plant loci from fungal infestation.

Greenhouse Tests

5 The following experiments were performed in the laboratory to determine the fungicidal efficacy of the compounds of the invention.

Test 1

10 This screen was used to evaluate the efficacy of the present compounds against a variety of different organisms that cause plant diseases.

15 The test compounds were formulated for application by dissolving 50 mg of the compound into 1.25 ml of solvent. The solvent was prepared by mixing 50 ml of "Tween 20" (polyoxyethylene (20) sorbitan monolaurate emulsifier) with 475 ml of acetone and 475 ml of ethanol. The solvent/compound solution was diluted to 125 ml with deionized water. The resulting formulation contains 400 ppm test chemical. Lower concentrations were obtained by serial dilution with the solvent-surfactant mixture.

20

The formulated test compounds were applied by foliar spray. The following plant pathogens and their corresponding plants were employed.

	<u>Pathogen</u>	<u>Designation in Following Table</u>	<u>Host</u>
5	<u>Erysiphe graminis tritici</u> (powdery mildew)	POWD MDEW	wheat
	<u>Pyricularia oryzae</u> (rice blast)	RICE BLAS	rice
10	<u>Puccinia recondita tritici</u> (leaf rust)	LEAF RUST	wheat
	<u>Botrytis cinerea</u> (gray mold)	GRAY MOLD	grape berries
15	<u>Pseudoperonospora cubensis</u> (downy mildew)	DOWN MDEW	squash

20                    The formulated technical compounds were sprayed on all foliar surfaces of the host plants (or cut berry) to past run-off. Single pots of each host plant were placed on raised, revolving pedestals in a fume hood. Test solutions were sprayed on all foliar surfaces. All treatments were allowed to dry and the plants were inoculated with the appropriate pathogens within 2-4 hours.

25                    Table 1 presents the activity of typical compounds of the present invention when evaluated in this experiment. The effectiveness of test compounds in controlling disease was rated using the following scale.

- 30
- 0     =     not tested against specific organism
  - =     0-19% control at 400 ppm
  - +     =     20-89% control at 400 ppm
  - 35    ++    =     90-100% control at 400 ppm
  - +++ =     90-100% control at 100 ppm

PLANT PATHOLOGY SCREEN

EXAMPLE NUMBER	POWD MDEW	RICE BLAST	LEAF RUST	GRAY MOLD	DOWN MDEW	
5	1	-	++	+	-	+
	2	+	++	+++	-	+++
	3	-	+	+	-	+
10	4	-	+	++	-	-
	5	+	-	++	-	-
	6	-	-	-	-	-
	7	-	+	-	-	+
	8	+	+	+++	-	+++
15	9	-	-	++	-	+
	10	-	++	+	-	+
	11	+	++	+++	-	++
	12	-	+	++	-	++
	13	-	-	-	-	-
20	14	-	-	-	-	-
	15	+	+	++	-	+++
	16	-	-	-	-	-
	17	-	-	-	-	-
	18	-	-	-	+	-
25	19	+	-	+++	-	+++
	20	-	+	+++	-	+++
	21	++	-	+++	-	+++
	22	-	-	-	-	-
	23	-	-	+	-	-

PLANT PATHOLOGY SCREEN

EXAMPLE NUMBER	POWD MDEW	RICE BLAST	LEAF RUST	GRAY MOLD	DOWN MDEW	
5	24	-	+	++	-	-
	25	-	-	+	-	-
	26	-	-	-	-	-
10	27	-	-	++	-	++
	28	-	-	-	-	-
	29	+	+	+	-	+
	30	-	-	+	-	+
	31	+	+	+++	-	+++
15	32	-	-	-	0	-
	33	++	+	+++	0	+++
	34	-	-	+	0	-
	35	-	-	++	0	+
	36	-	-	-	0	-
20	37	-	-	-	0	+
	38	-	-	-	-	-
	39	-	-	++	-	-
	40	+	-	++	-	++
	41	++	+	+++	-	+++
25	42	+	+	++	-	++
	43	++	+	++	-	+++
	44	-	-	++	-	+
	45	+	-	+++	-	+
	46	-	-	-	-	-

PLANT PATHOLOGY SCREEN

	EXAMPLE NUMBER	POWD MDEW	RICE BLAST	LEAF RUST	GRAY MOLD	DOWN MDEW
5	47	+	+++	+++	-	+++
	48	++	++	+++	-	++
	49	++	-	++	-	-
10	50	+	+	+	-	+
	51	-	-	-	-	++
	52	++	++	+++	-	+++
	53	+	+	++	-	-
	54	++	++	++	-	-
15	55	-	-	+	-	-
	56	+	+	+	-	-
	57	-	-	-	-	+
	58	-	-	+	-	+
	59	++	-	+	-	-
20	60	-	-	-	-	++
	61	++	+	+	-	++
	62	++	-	-	-	-
	63	-	-	-	-	+
	64	++	-	+	-	++
25	65	+	-	-	-	-
	66	++	-	+	-	++
	67	+	++	+	+	+
	68	+	++	+	-	++
	69	++	+	+	-	-
30	70	-	-	-	-	-

PLANT PATHOLOGY SCREEN

	<u>EXAMPLE NUMBER</u>	<u>POWD MDEW</u>	<u>RICE BLAST</u>	<u>LEAF RUST</u>	<u>GRAY MOLD</u>	<u>DOWN MDEW</u>
5	71	-	-	-	-	-
	72	+	-	++	-	++
	73	-	-	+	-	+
	74	-	-	-	-	+
10	75	-	-	-	-	+
	76	++	-	-	-	+
	77	++	++	++	-	++
	79	-	+	+	-	+++
15	80	-	-	-	-	0
	81	+	++	++	-	++
	82	+	+	+++	-	+++
	83	+	+	++	0	++
20	84	-	-	-	0	-
	85	+	+	+	0	+++
	86	++	-	-	0	-
	87	++	+	+++	0	0
	88	-	-	-	0	+
25	89	-	-	-	0	0

Combinations

30 Fungal disease pathogens are known to develop resistance to fungicides. When strains resistant to a fungicide do develop, it becomes necessary to apply larger and larger amounts of the fungicide to obtain desired results. To retard the development of resistance to new fungicides, it is desirable to apply the new fungicides in combination with other fungicides.

Use of a combination product also permits the product's spectrum of activity to be adjusted.

5


10

15

20

25

30



Accordingly, another aspect of the invention is a fungicidal combination comprising at least 1% by weight of a compound of formula (1) in combination with a second fungicide.

5 Contemplated classes of fungicides from which the second fungicide may be selected include:

1) N-substituted azoles, for example propiconazole, triademefon, flusilazol, diniconazole, ethyltrianol, myclobutanil, and prochloraz;

10 2) pyrimidines, such as fenarimol and nuarimol;

3) morpholines, such as fenpropimorph and tridemorph;

4) piperazines, such as triforine; and

5) pyridines, such as pyrifenox.

15 Fungicides in these five classes all function by inhibiting sterol biosynthesis. Additional classes of contemplated fungicides, which have other mechanisms of action include:

6) dithiocarbamates, such as maneb and mancozeb;

7) phthalimides, such as captafol;

8) isophthalonitrites, such as chlorothalonil;

9) dicarboximides, such as iprodione;

10) benzimidazoles, such as benomyl and carbendazim;

25 11) 2-aminopyrimidines, such as ethirimol;

12) carboxamides, such as carboxin; and

13) dinitrophenols, such as dinocap.

The fungicide combinations of the invention contain at least 1%, ordinarily 20 to 80%, and more typically 30 cally 50 to 75% by weight of a compound of formula (1).

Insecticide and Miticide Utility

The compounds of the invention are also useful for the control of insects and mites. Therefore, the present invention also is directed to a method for inhibiting an insect or mite which comprises applying to a locus of the insect or mite an insect- or mite-inhibiting amount of a compound of formula (1).

The compounds of the invention show activity against a number of insects and mites. More specifically, the compounds show activity against melon aphid, which is a member of the insect order Homoptera. Other members of the Homoptera include leafhoppers, planthoppers, pear psylla, apple sucker, scale insects, whiteflies, spittle bugs as well as numerous other host specific aphid species. Activity has also been observed against greenhouse thrips, which are members of the order Thysanoptera. The compounds also show activity against Southern armyworm, which is a member of the insect order Lepidoptera. Other typical members of this order are codling moth, cutworm, clothes moth, Indianmeal moth, leaf rollers, corn earworm, European corn borer, cabbage worm, cabbage looper, cotton bollworm, bagworm, eastern tent caterpillar, sod webworm, and fall armyworm.

Representative mite species with which it is contemplated that the present invention can be practiced include those listed below.

<u>FAMILY</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	
<b>ACARIDAE</b>			
5	<u>Aleurobius farinae</u>		
	<u>Rhizoglyphus echinopus</u>	Bulb mite	
	<u>Rhizoglyphus elongatus</u>		
	<u>Rhizoglyphus rhizophagus</u>		
	<u>Rhizoglyphus sagittatae</u>		
	<u>Rhizoglyphus tarsalis</u>		
10			
<b>ERIOPHYIDAE</b>			
6	<u>Abacarus farinae</u>	Grain rust mite	
	<u>Aceria brachytarsus</u>		
15	<u>Acalitus essigi</u>	Redberry mite	
	<u>Aceria ficus</u>		
	<u>Aceria fraaxinivorus</u>		
20	<u>Aceria granati</u>		
	<u>Aceria parapopuli</u>		
	<u>Eriophyes sheldoni</u>	Citrus bud mite	
	<u>Aceria tulipae</u>		
	<u>Aculus carnutus</u>	Peach silver mite	
	<u>Aculus schlechtendali</u>	Apple rust mite	
	<u>Colomerus vitis</u>	Grape erineum mite	
	<u>Eriophyes convolvens</u>		
	25	<u>Eriophyes insidiosus</u>	
		<u>Eriophyes malifoliae</u>	
<u>Eriophyes padi</u>			
30	<u>Eriophyes pruni</u>		
	<u>Epitrimerus pyri</u>	Pear leaf blister mite	
	<u>Eriophyes ramosus</u>		

<u>FAMILY</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>
5	<u>Eriophyes sheldoni</u>	Citrus bud mite
	<u>Eriophyes ribis</u>	
	<u>Phyllocoptes gracilis</u>	Dryberry mite
	<u>Phyllocoptruta oleivora</u>	Citrus rust mite
	<u>Phytoptus ribis</u>	
10	<u>Trisetacus pini</u>	
	<u>Vasates amygdalina</u>	
	<u>Vasates eurynotus</u>	
	<u>Vasates quadripedes</u>	Maple bladder gall mite
	<u>Vasates schlechtendali</u>	
EUPODIDAE		
15	<u>Penthaleus major</u>	Winter grain mite
	<u>Linopodes spp.</u>	
NALEPELLIDAE		
20	<u>Phyllocoptella avellanae</u>	Filbert bud mite
	PENTHALEIDAE	
	<u>Halotydeus destructor</u>	
PYEMOTIDAE		
25	<u>Pyemotes tritici</u>	Straw itch mite
	<u>Siteroptes cerealium</u>	
TARSONEMIDAE		
30	<u>Polyphagotarsonemus latus</u>	Broad mite
	<u>Steneotarsonemus pallidus</u>	Cyclamen mite

FAMILYSCIENTIFIC NAMECOMMON NAME

## TENUIPALPIDAE

5

Brevipalpus californicusBrevipalpus obovatus

Privet mite

Brevipalpus lewisi

Citrus flat mite

Dolichotetranychus floridanusPineapple flase spider  
mite

10

Tenuipalpes granatiTenuipalpes pacificus

## TETRANYCHIDAE

15

Bryobia arboreaBryobia practiosa

Clover mite

Bryobia rubrioculus

Brown mite

Eotetranychus coryliEotetranychus hicoriae

Pecan deaf scorch mite

Eotetranychus lewisiEotetranychus sexmaculatus

Sixspotted spider mite

20

Eotetranychus willamettiEotetranychus banksi

Texas citrus mite

Oligonychus ilicis

Southern red mite

Oligonychus pratensis

Banks grass mite

Oligonychus ununguis

Spruce spider mite

25

Panonychus citri

Citrus red mite

Panonychus ulmi

European red mite

Paratetranychus modestusParatetranychus pratensisParatetranychus viridis

30

Petrobia latens

Brown wheat mite

FAMILY

SCIENTIFIC NAME

COMMON NAME

5

10

Schizotetranychus celarius

Bamboo spider mite

Schizotetranychus pratensis

Tetranychus canadensis

Fourspotted spider mite

Tetranychus cinnabarinus

Carmine spider mite

Tetranychus mcdanieli

McDaniel spider mite

Tetranychus pacificus

Pacific spider mite

Tetranychus schoenei

Schoene spider mite

Tetranychus urticae


Twospotted spider mite

Tetranychus turkestanii

Strawberry spider mite

Tetranychus desertorum

Desert spider mite



The compounds are useful for reducing populations of insects and mites, and are used in a method of inhibiting an insect or mite population which comprises applying to a locus of the insect or arachnid an effective insect- or mite-inactivating amount of a compound of formula (1). The "locus" of insects or mites is a term used herein to refer to the environment in which the insects or mites live or where their eggs are present, including the air surrounding them, the food they eat, or objects which they contact. For example, plant-ingesting insects or mites can be controlled by applying the active compound to plant parts, which the insects or mites eat, particularly the foliage. It is contemplated that the compounds might also be useful to protect textiles, paper, stored grain, or seeds by applying an active compound to such substance. The term "inhibiting an insect or mite" refers to a decrease in the numbers of living insects or mites; or a decrease in the number of viable insect or mite eggs. The extent of reduction accomplished by a compound depends, of course, upon the application rate of the compound, the particular compound used, and the target insect or mite species. At least an insect-inactivating or mite-inactivating amount should be used. The terms "insect-inactivating amount" and "mite-inactivating amount" are used to describe the amount, which is sufficient to cause a measurable reduction in the treated insect or mite population. Generally an amount in the range from about 1 to about 1000 ppm active compound is used.

In a preferred embodiment, the present invention is directed to a method for inhibiting a mite which comprises applying to a plant an effective mite-inactivating amount of a compound of formula (1) in accordance with the present invention.

MITE/INSECT SCREEN

The compounds of Examples were tested for miticidal and insecticidal activity in the following mite/insect screen.

Each test compound was formulated by dissolving the compound in acetone/alcohol (50:50) mixture containing 23 g of "TOXIMUL R" (sulfonate/nonionic emulsifier blend) and 13 g of "TOXIMUL S" (sulfonate/nonionic emulsifier blend) per liter. These mixtures were then diluted with water to give the indicated concentrations.

Twospotted spider mites (Tetranychus urticae Koch) and melon aphids (Aphis gossypii Glover) were introduced on squash cotyledons and allowed to establish on both leaf surfaces. Other plants in the same treatment pot were left uninfested. The leaves were then sprayed with 5 ml of test solution using a DeVilbiss atomizing sprayer at 10 psi. Both surfaces of the leaves were covered until runoff, and then allowed to dry for one hour. Two uninfested leaves were then excised and placed into a Petri dish containing larval southern armyworm (Spodopetra eridania Cramer).

Activity on Southern corn rootworm (Diabrotica undecimpunctata howardi Barber) was evaluated by adding two ml of tap water, a presoaked corn seed, and 15 g of dry sandy soil to a one ounce plastic container. The soil was treated with 1 mL of test solution containing a pre-determined concentration of test compound. After six to 12 hours of drying, five 2-3 instar corn rootworm larvae were added to the individual cups, which were then capped and held at 23°C.

5  
10

After standard exposure periods, percent mortality and phytotoxicity were evaluated. Results for the compounds found to be active are reported in Table 3. The remaining compounds showed no activity. The following abbreviations are used in Table 3:

15

CRW refers to corn rootworm  
SAW refers to Southern armyworm  
SM refers to twospotted spider mites  
MA refers to melon aphids.

20

MITE & INSECT SCREEN

		CRW	CRW	SAW	SAW	SM	MA
	EXAMPLE	RATE	RESULTS	SM & MA	RESULTS	RESULTS	RESULTS
	NUMEER	PPM	%	RATE	%	%	%
				PPM			
5	1	12.00	0	200	0	0	80
10		24.00	100	400	0	0	0
	2	12.00	0	200	0	0	0
		24.00	0	400	0	0	80
15	3	12.00	0	200	30	0	0
		24.00	0	400	0	0	0
	4	24.00	0	400	0	0	0
20	5	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	6	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
25	7	12.00	0	200	0	30	100
		24.00	0	400	0	0	0
30	8	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	9	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
35	10	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	11	12.00	0	200	0	30	90
40		24.00	0	400	60	0	0
	12	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
45	13	12.00	0	200	0	0	0
		24.00	100	400	0	0	0

MITE & INSECT SCREEN

		CRW	CRW	SAW	SAW	SM	MA
	EXAMPLE	RATE	RESULTS	SM & MA	RESULTS	RESULTS	RESULTS
	NUMBER	PPM	%	RATE	%	%	%
				PPM			
5							
	14	12.00	0	200	0	0	0
10		24.00	0	400	0	0	0
	15	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
15	16	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	17	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
20	18	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	19	12.00	0	200	0	0	0
25		24.00	0	400	0	0	90
	20	12.00	0	200	0	C	0
		24.00	0	400	0	10C	0
		0	400	0	0	C	
30	21	12.00	100	200	0	8C	100
		24.00	100	400	0	6C	80
	22	12.00	0	200	0	C	0
35		24.00	0	400	0	C	0
	23	12.00	100	200	0	C	0
		24.00	0	400	0	C	0
40	24	12.00	0	200	0	C	0
		24.00	0	400	0	C	0

MITE & INSECT SCREEN

		CRW	CRW	SAW	SAW	SM	MA
	EXAMPLE	RATE	RESULTS	SM & MA	RESULTS	RESULTS	RESULTS
	NUMBER	PPM	%	RATE	%	%	%
				PPM			
5							
	25	12.00	0	200	0	0	0
10		24.00	0	400	0	0	0
	26	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
15	27	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	28	12.00	0	200	0	90	30
		24.00	0	400	0	0	0
20	29	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	30	12.00	0	200	0	0	0
25		24.00	0	400	90	0	0
	31	12.00	0	200	100	20	90
		24.00	0	400	80	100	0
30	32	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	33	12.00	0	200	0	80	40
		24.00	0	400	0	0	0
35	34	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	35	12.00	0	200	0	0	0
40		24.00	0	400	0	0	30
	36	12.00	0	200	0	0	0
		24.00	0	400	0	0	0

MITE & INSECT SCREEN

		CRW	CRW	SAW	SAW	SM	MA
	EXAMPLE	RATE	RESULTS	SM & MA	RESULTS	RESULTS	RESULTS
	NUMBER	PPM	%	RATE	%	%	%
				PPM			
5							
	37	12.00	0	200	0	90	0
10		24.00	0	400	0	0	0
	38	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
15	39	12.00	0	200	0	0	0
		24.00	0	400	0	0	10
	40	12.00	0	200	0	0	0
		24.00	0	400	0	20	80
20	41	12.00	0	200	0	0	0
		24.00	60	400	0	100	100
	42	12.00	0	200	0	0	100
25		24.00	0	400	0	100	100
	43	12.00	0	200	0	90	90
		24.00	0	400	0	80	80
30	44	24.00	0	400	0	0	0
	45	12.00	0	200	0	90	90
		24.00	0	400	0	100	90
35	46	12.00	0	200	0	80	90
		24.00	0	400	0	80	60
	47	12.00	100	200	100	90	100
		24.00	80	400	0	100	100
40	48	12.00	60	200	0	0	0
		24.00	100	400	0	0	80

MITE & INSECT SCREEN

		CRW	CRW	SAW	SAW	SM	MA
	EXAMPLE	RATE	RESULTS	SM & MA	RESULTS	RESULTS	RESULTS
	NUMBER	PPM	%	RATE	%	%	%
				PPM			
5							
	49	12.00	0	200	80	0	0
10		24.00	0	400	0	0	0
	50	12.00	0	200	0	0	0
		24.00	0	400	0	100	100
15	51	12.00	0	200	0	100	90
		24.00	0	400	0	80	80
	52	12.00	0	200	0	0	80
		24.00	0	400	0	0	100
20	53	12.00	0	200	0	0	0
		24.00	0	400	0	30	60
25	54	12.00	0	200	0	100	100
		24.00	0	400	40	60	40
	55	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
30	56	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	57	12.00	0	200	0	0	0
			0	200	0	0	0
35		24.00	0	400	0	0	0
			0	400	0	0	0
	58	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
40							
	59	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
45	60	12.00	0	200	0	0	50
		24.00	0	400	0	80	60

MITE & INSECT SCREEN

		CRW	CRW	SAW	SAW	SM	MA
	EXAMPLE	RATE	RESULTS	SM & MA	RESULTS	RESULTS	RESULTS
	NUMBER	PPM	%	RATE	%	%	%
				PPM			
5							
	61	12.00	0	200	0	90	90
10		24.00	0	400	0	0	0
	62	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
15	63	12.00	0	200	0	0	0
		24.00	0	400	0	50	80
	64	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
20	65	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
25	66	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	67	12.00	0	200	0	100	100
		24.00	0	400	0	100	100
30	68	24.00	0	400	100	30	90
	69	24.00	0	400	0	90	90
	70	24.00	0	400	0	0	0
35	71	24.00	0	400	0	0	0
	72	24.00	0	400	60	0	100
40	73	12.00		200			
		24.00	0	400	100	0	90
	74	24.00	0	400	0	100	90

MITE & INSECT SCREEN

5	COMPOUND	CRW RATE PPM	CRW RESULTS %	SAW SM & MA RATE PPM	SAW RESULTS %	SM RESULTS %	MA RESULTS %
	75	24.00	0	400	40	0	0
	76	24.00	60	400	0	0	0
10	77	12.00	0	200	0	50	100
		24.00	0	400	0	90	80
	78	12.00	0	200	0	20	90
		24.00	0	400	100	0	100
	79	24.00	0	400	0	0	0
15	80	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	81	24.00	0	400	0	0	0
	82	12.00	100	200	40	100	100
		24.00	100	400	0	100	60
20	83	12.00	0	200	0	0	0
		24.00	0	400	100	0	0
	84	12.00	0	200	0	0	0
		24.00	0	400	0	0	0
	85	12.00	0	200	0	100	100
		24.00	0	400	0	20	0
25	86	12.00	0	200	0	90	60
		24.00	0	400	0	100	100
	87	12.00	0	200	0	0	0
		24.00	0	400	80	0	0
30	88	12.00	0	200	0	90	60
		24.00	0	400	0	0	0

## Compositions

The compounds of this invention are applied in the form of compositions which are important embodiments of the invention, and which comprise a compound of this invention and a phytologically-acceptable inert carrier. The compositions are either concentrated formulations which are dispersed in water for application, or are dust or granular formulations which are applied without further treatment. The compositions are prepared according to procedures and formulae which are conventional in the agricultural chemical art, but which are novel and important because of the presence therein of the compounds of this invention. Some description of the formulation of the compositions will be given, however, to assure that agricultural chemists can readily prepare any desired composition.

The dispersions in which the compounds are applied are most often aqueous suspensions or emulsions prepared from concentrated formulations of the compounds. Such water-soluble, water-suspendable or emulsifiable formulations are either solids usually known as wettable powders, or liquids usually known as emulsifiable concentrates or aqueous suspensions. Wettable powders, which may be compacted to form water dispersible granules, comprise an intimate mixture of the active compound, an inert carrier and surfactants. The concentration of the active compound is usually from about 10% to about 90% by weight. The inert carrier is usually chosen from among the attapulgite clays, the montmorillonite clays, the diatomaceous

earths, or the purified silicates. Effective surfactants, comprising from about 0.5% to about 10% of the wettable powder, are found among the sulfonated lignins, the condensed naphthalenesulfonates, the naphthalene-sulfonates, the alkylbenzenesulfonates, the alkyl sulfates, and nonionic surfactants such as ethylene oxide adducts of alkyl phenols.

Emulsifiable concentrates of the compounds comprise a convenient concentration of a compound, such as from about 50 to about 500 grams per liter of liquid, equivalent to about 10% to about 50%, dissolved in an inert carrier which is either water miscible solvent or a mixture of water-immiscible organic solvent and emulsifiers. Useful organic solvents include aromatics, especially the xylenes, and the petroleum fractions, especially the high-boiling naphthalenic and olefinic portions of petroleum such as heavy aromatic naphtha. Other organic solvents may also be used, such as the terpenic solvents including rosin derivatives, aliphatic ketones such as cyclohexanone, and complex alcohols such as 2-ethoxyethanol. Suitable emulsifiers for emulsifiable concentrates are chosen from conventional nonionic surfactants, such as those discussed above.

Aqueous suspensions comprise suspensions of water-insoluble compounds of this invention, dispersed in an aqueous vehicle at a concentration in the range from about 5% to about 50% by weight. Suspensions are prepared by finely grinding the compound, and vigorously mixing it into a vehicle comprised of water and surfactants chosen from the same types discussed above. Inert

35

39

ingredients, such as inorganic salts and synthetic or natural gums, may also be added, to increase the density and viscosity of the aqueous vehicle. It is often most effective to grind and mix the compound at the same time  
5 by preparing the aqueous mixture, and homogenizing it in an implement such as a sand mill, ball mill, or piston-type homogenizer.

The compounds may also be applied as granular compositions, which are particularly useful for  
10 applications to the soil. Granular compositions usually contain from about 0.5% to about 10% by weight of the compound, dispersed in an inert carrier which consists entirely or in large part of clay or a similar inexpensive substance. Such compositions are usually prepared by  
15 dissolving the compound in a suitable solvent, and applying it to a granular carrier which has been pre-formed to the appropriate particle size, in the range of from about 0.5 to 3 mm. Such compositions may also be formulated by making a dough or paste of the carrier and  
20 compound, and crushing and drying to obtain the desired granular particle size.

Dusts containing the compounds are prepared simply by intimately mixing the compound in powdered form with a suitable dusty agricultural carrier, such as kaolin clay,  
25 ground volcanic rock and the like. Dusts can suitably contain from about 1% to about 10% of the compound.

It is equally practical, when desirable for any reason, to apply the compound in the form of a solution in an appropriate organic solvent, usually a bland petroleum  
30 oil, such as the spray oils, which are widely used in agricultural chemistry.

35

39

Insecticides and miticides are generally applied in the form of a dispersion of the active ingredient in a liquid carrier. It is conventional to refer to application rates in terms of the concentration of active ingredient in the carrier. The most widely used carrier is water.

The compounds of the invention can also be applied in the form of an aerosol composition. In such compositions the active compound is dissolved or dispersed in an inert carrier, which is a pressure-generating propellant mixture. The aerosol composition is packaged in a container from which the mixture is dispensed through an atomizing valve. Propellant mixtures comprise either low-boiling halocarbons, which may be mixed with organic solvents, or aqueous suspensions pressurized with inert gases or gaseous hydrocarbons.

The actual amount of compound to be applied to loci of insects and mites is not critical and can readily be determined by those skilled in the art in view of the examples above. In general, concentrations of from 10 ppm to 5000 ppm of compound are expected to provide good control. With many of the compounds, concentrations of from 100 to 1500 ppm will suffice. For field crops, such as soybeans and cotton, a suitable application rate for the compounds is about 0.5 to 1.5 lb/A, typically applied in 50 gal/A of spray formulation containing 1200 to 3600 ppm of compound. For citrus crops, a suitable application rate is from about 100 to 1500 gal/A spray formulation, which is a rate of 100 to 1000 ppm.

The locus to which a compound is applied can be any locus inhabited by an insect or arachnid, for example, vegetable crops, fruit and nut trees, grape vines, and ornamental plants. Inasmuch as many mite species are specific to a particular host, the foregoing list of mite species provides exemplification of the wide range of settings in which the present compounds can be used.

Because of the unique ability of mite eggs to resist toxicant action, repeated applications may be desirable to control newly emerged larvae, as is true of other known acaricides.

The following formulations of compounds of the invention are typical of compositions useful in the practice of the present invention.

15

20

25

30

35

39

A. 0.75 Emulsifiable Concentrate

	5-methyl-N-[2-(4-methoxyphenyl)ethyl]- 5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	9.38%
5	"TOXIMUL D" (nonionic/anionic surfactant blend)	2.50%
	"TOXIMUL H" (nonionic/anionic surfactant blend)	2.50%
10	"EXXON 200" (naphthalenic solvent)	85.62%

15 B. 1.5 Emulsifiable Concentrate

	5-methyl-N-[2-(2-naphthyl)ethyl]furo- [2,3-d]pyrimidin-4-amine	18.50%
20	"TOXIMUL D"	2.50%
	"TOXIMUL H"	2.50%
	"EXXON 200"	76.50%

25

C. 0.75 Emulsifiable Concentrate

	N-[2-[4-(1,1,2,2-tetrafluoroethoxy)phenyl]- ethyl]-5,6-dimethylfuro[2,3-d]pyrimidin-4-amine	9.38%
30	"TOXIMUL D"	2.50%
	"TOXIMUL H"	2.50%
35	"EXXON 200"	85.62%

39

D. 1.0 Emulsifiable Concentrate

5	5,6-dimethyl-4-[2-[4-[4-(trifluoromethyl)-phenoxy]phenyl]ethoxy]furo[2,3-d]pyrimidine	12.50%
	N-methylpyrrolidone	25.00%
	"TOXIMUL D"	2.50%
10	"TOXIMUL H"	2.50%
	"EXXON 200"	57.50%

E. 1.0 Aqueous Suspension

15	5,6-dimethyl-4-[2-[4-(n-butoxy)phenyl]-ethoxy]furo[2,3-d]pyrimidine	12.00%
	"PLURONIC P-103"	1.50%
20	(block copolymer of propylene oxide and ethylene oxide, surfactant)	
	"PROXEL GXL"	.05%
	(biocide/preservative)	
25	"AF-100"	.20%
	(silicon based antifoam agent)	
	"REAX 88B"	1.00%
30	(lignosulfonate dispersing agent)	
	propylene glycol	10.00%
	veegum	.75%
35	xanthan	.25%
	water	74.25%

39

GD

F. 1.0 Aqueous Suspension

	5-methyl-N-[2-(4-methoxyphenyl)ethyl]- furo[2,3-d]pyrimidin-4-amine	12.50%
5	"MAKON 10" (10 moles ethyleneoxide nonylphenol surfactant)	1.00%
	"ZEOSYL 200" (silica)	1.00%
10	"AF-100"	0.20%
	"AGRIWET FR" (surfactant)	3.00%
15	2% xanthan hydrate	10.00%
	water	72.30%

G. 1.0 Aqueous Suspension

20	5-methyl-N-[2-(4-chlorophenyl)ethyl]- furo[2,3-d]pyrimidin-4-amine	12.50%
	"MAKON 10"	1.50%
25	"ZEOSYL 200" (silica)	1.00%
	"AF-100"	0.20%
30	"POLYFON H" (lignosulfonate dispersing agent)	0.20%
	2% xanthan hydrate	10.00%
35	water	74.60%

H. Wettable Powder

	5,6-dimethyl-4-[2-[4-[4-(trifluoro-	25.80%
5	methyl)phenoxy]phenyl]ethoxy]furo[2,3-d]- pyrimidine	
	"POLYFON H"	3.50%
	"SELLOGEN HR"	5.00%
10	"STEPANOL ME DRY"	1.00%
	gum arabic	0.50%
15	"HISIL 233"	2.50%
	Barden clay	61.70%

I. 1.0 Aqueous Suspension

20	5,6-dimethyl-4-[2-[4-[4-(trifluoro-	12.40%
	methyl)phenoxy]phenyl]ethoxy]furo[2,3-d]- pyrimidine	
25	"TERGITOL 158-7"	5.00%
	"ZEOSYL 200"	1.00%
	"AF-100"	0.20%
30	"POLYFON H"	0.50%
	2% xanthan solution	10.00%
35	tap water	70.90%

J. 1.0 Emulsifiable Concentrate

	5-methyl-N-[2-(2-naphthyl)ethyl]- furo[2,3-d]pyrimidin-4-amine	12.40%
5	"TOXIMUL D"	2.50%
	"TOXIMUL H"	2.50%
10	"EXXON 200"	82.60%

K. Wettable Powder

	5,6-dimethyl-4-[2-[4-(n-butoxy)phenyl]- ethoxy]furo[2,3-d]pyrimidine	25.80%
15	"SELLOGEN HR"	5.00%
	"POLYFON H"	4.00%
20	"STEPANOL ME DRY"	2.00%
	"HISIL 233"	3.00%
25	Barden clay	60.20%

L. 0.5 Emulsifiable Concentrate

	5-methyl-N-[2,(2-naphthyl)ethyl]furo- [2,3-d]pyrimidin-4-amine	6.19%
30	"TOXIMUL H"	3.60%
	"TOXIMUL D"	0.40%
35	"EXXON 200"	89.81%

39

GD

M. Wettable Powder

	5-methyl-N-[2-(4-methoxyphenyl)ethyl]- furo[2,3-d]pyrimidin-4-amine	25.80%
5	"SELLOGEN HR"	5.00%
	"POLYFON H"	4.00%
10	"STEPANOL ME DRY"	2.00%
	"HISIL 233"	3.00%
15	Barden Clay	60.20%

N. 1.0 Aqueous Suspension

	N-[2-[4-(1,1,2,2-tetrafluoroethoxy)- phenyl]ethyl]-5,6-dimethylfuro[2,3-d]- pyrimidin-4-amine	12.40%
20	"TERGITOL 158-7"	5.00%
	"ZEOSYL 200"	1.00%
25	"POLYFON H"	0.50%
	"AF-100"	0.20%
30	xanthan solution (2%)	10.00%
	tap water	70.90%

35

39


GD

The claims defining the invention are as follows:


1. A compound of formula (1)

5

10 wherein



15



$R^1$  is H,  $C_1-C_4$  alkyl, or phenyl optionally substituted with halo,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$  alkoxy, halo  $(C_1-C_4)$  alkyl, or halo  $(C_1-C_4)$  alkoxy;  
 $R^2$  is H,  $C_1-C_4$  alkyl, nitro, halo; or  
 $R^1$  and  $R^2$  combine to form  $-(CH_2)_4-$ ;  
 $R^3$  is H,  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkyl, phenyl, or substituted phenyl;

$-XYZ$  together form 1H-1,2,4-triazol-1-yl; or  
X is O, S,  $NR^4$ , or  $CR^5R^6$ , where  $R^4$  is H,  $(C_1-C_4)$  alkyl, or  $(C_1-C_4)$  acyl, and  $R^5$  and  $R^6$  are independently H,  $(C_1-C_4)$  acyl,  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  alkenyl or -alkynyl, CN, or OH, or  $R^5$  and  $R^6$  combine to form a carbocyclic ring containing four to six carbon atoms; and

~~YZ together form a  $C_5-C_{12}$  saturated or unsaturated hydrocarbon chain, straight chain or branched; optionally including a hetero atom selected from O,  $NR^4$ , S, SO,  $SO_2$ , or  $SiR^7R^8$ , where  $R^7$  and  $R^8$  are independently~~

39

-70-




(C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>) branched alkyl, phenyl, or substituted phenyl, and optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, halo  
 5 (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy, or (C<sub>1</sub>-C<sub>4</sub>) acyl; ~~or~~

Y is a bond or an alkylene chain one to six carbon atoms long, ~~optionally including a carbocyclic ring, and optionally including a hetero atom selected~~  
 10 ~~from O, NR<sup>4</sup>, S, SO, SO<sub>2</sub>, or SiR<sup>7</sup>R<sup>8</sup>, where R<sup>4</sup>, R<sup>7</sup>, and R<sup>8</sup> are as defined above, and optionally substituted with~~  
 (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched (C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or -cycloalkenyl, halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy, or  
 15 (C<sub>1</sub>-C<sub>4</sub>) acyl; and

Z is

(a) (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or cycloalkenyl, optionally substituted with (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>1</sub>-C<sub>4</sub>) alkoxy, halo (C<sub>1</sub>-C<sub>4</sub>) alkyl, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, halo, hydroxy, or  
 20 (C<sub>1</sub>-C<sub>4</sub>) acyl;

(b) a phenyl group of the formula (2)



25

where

R<sup>9</sup> to R<sup>13</sup> are independently

H,

halo,

I,




- (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or -cycloalkenyl,  
 phenoxy, or  
 substituted phenoxy,  
 phenylthio, or substituted phenylthio,  
 5 phenyl, or substituted phenyl,  
 NO<sub>2</sub>,  
 acetoxy,  
 OH,  
 CN,  
 10 SiR<sup>7</sup>R<sup>8</sup>R<sup>14</sup> or OSiR<sup>7</sup>R<sup>8</sup>R<sup>14</sup>, where R<sup>7</sup>, and R<sup>8</sup> are  
 as defined above and R<sup>14</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl, (C<sub>3</sub>-C<sub>4</sub>)  
 branched alkyl, phenyl, or substituted phenyl,  
 NR<sup>15</sup>R<sup>16</sup>, where R<sup>15</sup> and R<sup>16</sup> are independently H,  
 (C<sub>1</sub>-C<sub>4</sub>) alkyl, or (C<sub>1</sub>-C<sub>4</sub>) acyl,  
 15 S(O)R<sup>17</sup>, or SO<sub>2</sub>R<sup>17</sup>, where R<sup>17</sup> is (C<sub>1</sub>-C<sub>10</sub>) alkyl,  
 phenyl, or substituted phenyl;  
 a C<sub>1</sub>-C<sub>12</sub> saturated or unsaturated hydrocarbon  
 chain, straight chain or branched optionally including a  
 hetero atom selected from O, S, SO, SO<sub>2</sub>, NR<sup>4</sup>, or SiR<sup>7</sup>R<sup>8</sup>,  
 20 where R<sup>4</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above, and optionally  
 substituted with halo, halo (C<sub>1</sub>-C<sub>4</sub>) alkoxy, hydroxy,  
 (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or cycloalkenyl, (C<sub>1</sub>-C<sub>4</sub>) acyl,  
 phenoxy, substituted phenoxy, phenyl, substituted  
 phenyl, phenylthio, or substituted phenylthio;  
 25 (C<sub>1</sub>-C<sub>7</sub>) alkoxy optionally substituted with  
 halo, phenyl, substituted phenyl, (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or  
 cycloalkenyl, phenoxy, or substituted phenoxy;  
 (C<sub>1</sub>-C<sub>7</sub>) alkylthio optionally substituted with  
 halo, phenyl, substituted phenyl, (C<sub>3</sub>-C<sub>8</sub>) cycloalkyl or  
 30 cycloalkenyl, phenoxy or substituted phenoxy; or

$R^{10}$  and  $R^{11}$  combine to form  $-O-CF_2-O-$  or  $-O-(CH_2)_2-O-$ ;

(c) a furyl group of formula (3)


5



where  $R^{18}$  is H, halo, halomethyl, CN,  $NO_2$ ,  $(C_1-C_4)$  alkyl,  $(C_3-C_4)$  branched alkyl, phenyl,  $(C_1-C_4)$  alkoxy;

(d) a thienyl group of the formula (4)

10



where  $R^{19}$  is  $R^{18}$  as defined in paragraph (c) or thienyl;

(e) a group of formula (5) or (6)

15



where  $R^{18}$  is as defined in paragraph (c), J is N or CH, and G is O,  $NR^{20}$ , or S, provided that if J is not N then G is  $NR^{20}$ , where  $R^{20}$  is H,  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  acyl, phenylsulfonyl, or substituted phenylsulfonyl;

(f) a group selected from

optionally substituted naphthyl, dihydro-  
25 naphthyl, tetrahydronaphthyl, and decahydronaphthyl;

optionally substituted pyridyl;

optionally substituted indolyl;

1,3-benzodioxolyl;

2,6-dimethyl-4-morpholinyl; and

30

1-adamantyl;

or an acid addition salt of a compound of formula (1);  
provided that:

5 if the compound of formula (1) is a furo[2,3-d]-  
pyrimidin-4-amine, then Y is an alkylene chain two to six  
carbon atoms long, optionally substituted with (C<sub>1</sub>-C<sub>4</sub>)  
alkyl, (C<sub>2</sub>-C<sub>4</sub>) alkenyl or -alkynyl, branched  
(C<sub>3</sub>-C<sub>7</sub>) alkyl, (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl or  
10 -cycloalkenyl, halo, hydroxy, or acetyl, and Z is one of  
the groups defined in paragraphs (a) through (f).

2. A compound according to claim 1 that is a  
furo-[2,3-d]pyrimidine.

15 3. A compound according to either claim 1 or 2  
wherein one of R<sup>1</sup> and R<sup>2</sup> is methyl and the other is H  
or methyl.

20 4. A compound according to any one of claims 1  
to 3 wherein Y is -CH<sub>2</sub>CH<sub>2</sub>-.

5. A compound according to any one of claims 1  
to 4 wherein Z is a phenyl group of formula (2).

25 6. A compound according to any one of claims 1  
to 5 wherein X is NH.

30 7. A compound according to any one of claims 1  
to 6 wherein X is O.

35



8. A compound according to claim 5 wherein Z is a phenyl group having at least one haloalkoxy substituent.

5 9. A compound according to either claim 5 or 8 wherein Z is a phenyl group having at least one haloalkyl substituent.

10 10. A compound according to any one of claims 5, 8 or 9 wherein Z is a phenyl group having a phenoxy or substituted phenoxy substituent.

15

20

25


30  
35

40




11. A compound of formula (7)

5



10 where



15

$R^1$  is H,  $C_1-C_4$  alkyl, or phenyl optionally substituted with halo,  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  alkoxy, halo  $(C_1-C_4)$  alkyl, or halo  $(C_1-C_4)$  alkoxy;

$R^2$  is H,  $C_1-C_4$  alkyl, nitro, halo; or

$R^1$  and  $R^2$  combine to form  $-(CH_2)_4-$ ;

$R^3$  is H,  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkyl, phenyl, or substituted phenyl;


Y is an alkylene chain two to six carbon atoms long, optionally substituted with  $(C_1-C_4)$  alkyl,  $(C_2-C_4)$  alkenyl or -alkynyl, branched  $(C_3-C_7)$  alkyl,  $(C_3-C_7)$  cycloalkyl or -cycloalkenyl, halo, halo  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkoxy, hydroxy, or  $(C_1-C_4)$  acyl; and Z is

(a)  $(C_3-C_8)$  cycloalkyl or cycloalkenyl, optionally substituted with  $(C_1-C_4)$  alkyl,  $(C_1-C_4)$  alkoxy, halo  $(C_1-C_4)$  alkyl, halo  $(C_1-C_4)$  alkoxy, halo, hydroxy, or  $(C_1-C_4)$  acyl;

40



(b) a phenyl group of the formula (2)



where

$R^9$  to  $R^{13}$  are independently

H,

halo,

I,

( $C_3$ - $C_8$ ) cycloalkyl or -cycloalkenyl,

phenoxy, or

substituted phenoxy,

phenylthio, or substituted phenylthio,

phenyl, or substituted phenyl,

$NO_2$ ,

acetoxy,

OH,

CN,

$SiR^7R^8R^{14}$  or  $OSiR^7R^8R^{14}$ , where  $R^7$ , and  $R^8$  are as defined above and  $R^{14}$  is ( $C_1$ - $C_4$ ) alkyl, ( $C_1$ - $C_4$ ) branched alkyl, phenyl, or substituted phenyl,

$NR^{15}R^{16}$ , where  $R^{15}$  and  $R^{16}$  are independently H, ( $C_1$ - $C_4$ ) alkyl, or ( $C_1$ - $C_4$ ) acyl,

$S(O)R^{17}$ , or  $SO_2R^{17}$ , where  $R^{17}$  is ( $C_1$ - $C_{10}$ ) alkyl, phenyl, or substituted phenyl;

a  $C_1$ - $C_{12}$  saturated or unsaturated hydrocarbon chain, straight chain or branched optionally including a hetero atom selected from O, S, SO,  $SO_2$ ,  $NR^4$ , or  $SiR^7R^8$ ,

where  $R^4$ ,  $R^7$  and  $R^8$  are as defined above, and optionally substituted with halo, halo ( $C_1-C_4$ ) alkoxy, hydroxy, ( $C_3-C_8$ ) cycloalkyl or cycloalkenyl, ( $C_1-C_4$ ) acyl, phenoxy, substituted phenoxy, phenyl, substituted phenyl, phenylthio, or substituted phenylthio;


5 (C<sub>1</sub>-C<sub>7</sub>) alkoxy optionally substituted with halo, phenyl, substituted phenyl, ( $C_3-C_8$ ) cycloalkyl or cycloalkenyl, phenoxy, or substituted phenoxy;

10 (C<sub>1</sub>-C<sub>7</sub>) alkylthio optionally substituted with halo, phenyl, substituted phenyl, ( $C_3-C_8$ ) cycloalkyl or cycloalkenyl, phenoxy or substituted phenoxy; or

$R^{10}$  and  $R^{11}$  combine to form  $-O-CF_2-O-$  or  $-O-(CH_2)_2-O-$ ;

(c) a furyl group of formula (3)


15



where  $R^{18}$  is H, halo, halomethyl, CN,  $NO_2$ , ( $C_1-C_4$ ) alkyl, ( $C_3-C_4$ ) branched alkyl, phenyl, ( $C_1-C_4$ ) alkoxy;

20



(d) a thienyl group of the formula (4)



where  $R^{19}$  is  $R^{18}$  as defined in paragraph (c) or thienyl;

25

(e) a group of formula (5) or (6)



where R<sup>18</sup> is as defined in paragraph (c), J is N or CH,  
and G is O, NR<sup>20</sup>, or S, provided that if J is not N then  
G is NR<sup>20</sup>, where R<sup>20</sup> is H, (C<sub>1</sub>-C<sub>4</sub>) alkyl,  
(C<sub>1</sub>-C<sub>4</sub>) acyl, phenylsulfonyl, or substituted  
5 phenylsulfonyl;

(f) a group selected from optionally  
substituted naphthyl, dihydronaphthyl, tetrahydronaphthyl,  
and decahydronaphthyl;

optionally substituted pyridyl;  
10 optionally substituted indolyl;  
1,3-benzodioxolyl;  
2,6-dimethyl-4-morpholinyl; and  
1-adamantyl.

15 12. A fungicidal method which includes applying  
to the locus of a plant pathogen a  
phytologically-acceptable amount of a compound according  
to any one of claims 1 to 10.

20 13. A fungicidal method which includes applying  
to the locus of a plant pathogen a  
phytologically-acceptable amount of a compound according  
to claim 11.

25 14. A composition for use as a fungicide,  
insecticide or miticide which includes a compound  
according to any one of claims 1 to 11 in combination with  
a phytologically-acceptable carrier.

30 15. An insecticidal or miticidal method which  
includes applying to a locus of an insect or mite an  
insect- or mite-inactivating amount of a compound  
according to any one of claims 1 to 11.

35 16. A compound of formula (1) or formula (7)  
substantially as described herein with reference to any  
one of the examples.

17. A method of preparation of a compound of



formula (1) or formula (7) substantially as described herein with reference to any one of the examples.

18. A composition for use as a fungicide,  
5 insecticide or miticide substantially as described herein with reference to any one of the examples.

19. A fungicidal, insecticidal or miticidal  
10 method substantially as described herein with reference to any one of the examples.

DATED: 9 December, 1993

PHILLIPS ORMONDE & FITZPATRICK

Patent Attorneys For:

15 DOWELANCO

*David B Fitzpatrick*

20

25

30

35

(6102h)

