



HU000025872T2

(19) **HU**(11) Lajstromszám: **E 025 872**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 10 161291**
- (22) A bejelentés napja: **2003. 06. 18.**
- (96) Az európai bejelentés bejelentési száma:
EP 20030161291
- (97) Az európai bejelentés közzétételi adatai:
EP 2213730 A1 **2010. 08. 04.**
- (97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2213730 B1 **2015. 12. 02.**
- (51) Int. Cl.: **C12N 5/10** (2006.01)
A61K 48/00 (2006.01)
C12N 15/09 (2006.01)

- | | |
|--|--|
| (30) Elsőbbségi adatok:
2002177648 2002. 06. 18. JP
2002237974 2002. 08. 19. JP | (73) Jogosult(ak):
Eisai R&D Management Co., Ltd., Tokyo
112-8088 (JP) |
| (72) Feltaláló(k):
Ito, Masashi, Tsukaba-shi, Ibaraki 300-2635 (JP)
SAITO, Yasushi, Chiba-shi, Chiba 260-0853 (JP) | (74) Képviselő:
Danubia Szabadalmi és Jogi Iroda Kft.,
Budapest |

- (54) **Primer, tenyésztett adipociták génterápiára**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 213 730 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
02.12.2015 Bulletin 2015/49

(51) Int Cl.:
C12N 5/10^(2006.01) C12N 15/09^(2006.01)
A61K 48/00^(2006.01)

(21) Application number: **10161291.9**

(22) Date of filing: **18.06.2003**

(54) **Primary cultured adipocytes for gene therapy**

Primäre kultivierte Adipozyten für die Gentherapie

Adipocytes primaires cultivés pour thérapie génique

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PT RO SE SI SK TR
Designated Extension States:
AL LT LV MK

(30) Priority: **18.06.2002 JP 2002177648**
19.08.2002 JP 2002237974

(43) Date of publication of application:
04.08.2010 Bulletin 2010/31

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
03736235.7 / 1 541 674

(73) Proprietor: **Eisai R&D Management Co., Ltd.**
Tokyo 112-8088 (JP)

(72) Inventors:
• **Ito, Masashi**
Tsukaba-shi,
Ibaraki 300-2635 (JP)
• **Saito, Yasushi**
Chiba-shi, Chiba 260-0853 (JP)

(74) Representative: **Vossius & Partner**
Patentanwälte Rechtsanwälte mbB
Siebertstrasse 3
81675 München (DE)

(56) References cited:
WO-A-00/31267

- **NAGAMATSU<A> S ET AL: "Adenovirus-mediated preproinsulin gene transfer into adipose tissues ameliorates hyperglycemia in obese diabetic KKA<y> mice" FEBS LETTERS, ELSEVIER, AMSTERDAM, NL LNKD- DOI: 10.1016/S0014-5793(01)03146-5, vol. 509, no. 1, 30 November 2001 (2001-11-30), pages 106-110, XP004329153 ISSN: 0014-5793**
- **RAPER S E ET AL: "Cell transplantation in liver-directed gene therapy" CELL TRANSPLANTATION, ELSEVIER SCIENCE, US, vol. 2, no. 5, 1 January 1993 (1993-01-01) , pages 381-400, XP002970482 ISSN: 0963-6897**
- **CHRISTENSEN R ET AL: "Cutaneous gene therapy - an update" HISTOCHEMISTRY AND CELL BIOLOGY, SPRINGER, BERLIN, DE, vol. 115, no. 1, 1 January 2001 (2001-01-01), pages 73-82, XP002970483 ISSN: 0948-6143**
- **HERTZEL A V ET AL: "Adenovirus-mediated gene transfer in primary murine adipocytes" JOURNAL OF LIPID RESEARCH, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, INC, US, vol. 41, no. 7, 1 January 2000 (2000-01-01), pages 1082-1086, XP002970484 ISSN: 0022-2275**
- **SIMONSON G D ET AL: "SYNTHESIS AND PROCESSING OF GENETICALLY MODIFIED HUMAN PROINSULIN BY RAT MYOBLAST PRIMARY CULTURES" HUMAN GENE THERAPY, MARY ANN LIEBERT, NEW YORK ,NY, US, vol. 7, no. 1, 1996, pages 71-78, XP000900066 ISSN: 1043-0342**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 213 730 B1

- SHIBASAKI M ET AL: "Alterations of insulin sensitivity by the implantation of 3T3-L1 cells in nude mice. A role for TNF-alpha?"
DIABETOLOGIA. APR 2002, vol. 45, no. 4, April 2002 (2002-04), pages 518-526, XP002403318
ISSN: 0012-186X
- ITO M ET AL: "Implantation of primary cultured adipocytes that secrete insulin modifies blood glucose levels in diabetic mice."
DIABETOLOGIA. AUG 2005, vol. 48, no. 8, August 2005 (2005-08), pages 1614-1620, XP002403319
ISSN: 0012-186X

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

DescriptionTechnical Field

5 **[0001]** The present invention relates to primary cultured adipocytes for gene therapy, to which a foreign gene(s) has been transferred.

Background Art

10 **[0002]** Current gene therapies (Toyooka et al., Folia Pharmacol. Jpn. 116:158-162, 2000) can be classified into two groups: (1) methods of transferring therapeutic genes into patients by directly administering viral vectors, naked plasmids, or such that encode the gene (*in vivo*), and (2) methods of temporarily removing cells from patients, transferring a gene to these cells, and then returning these cells to the patient (*ex vivo*).

15 **[0003]** In the *in vivo* methods, major problems remain to be solved, such as transfer efficiency, continuous expression, and selective gene transfer to target cells. *Ex vivo* methods, on the other hand, can potentially overcome these problems. The majority of examples of *ex vivo* methods have been performed using blood-system cells (peripheral lymphocytes and bone marrow cells), since their collection and transplantation is relatively easy and the burden on patients is reduced (Tani et al., Saishin Igaku, 56:258-267, 2001). With regards to cells other than blood-system cells, methods that transfer genes to hepatocytes and then return these cells to the patient have been carried out (Raper, S.E. et al., Cell Transplant 2(5):381-400, 1993), but most of these methods focus on the recovery, maintenance, and enhancement of the function of the transfected cells themselves.

20 **[0004]** WO00/31267-A1 relates to the introduction of insulin gene into muscle cells by retrovirus *in vitro*, and re-implantation of the cells into the body, as a treatment for diabetes.

25 Disclosure of the Invention

[0005] While searching for cells suitable for *ex vivo* gene therapy, the present inventors developed the idea of using primary cultured adipocytes. The use of adipocytes has the following advantages:

30 (1) there are many reports of humoral factors secreted from adipocytes, and adipocytes comprise the functions of hormone production and can act as secretory organs (Bradley R.D., et al., Recent Prog. Horm. Res., 2001, 56, 329-358);

(2) adipocytes can be easily collected since they also exist subcutaneously, and techniques relating to their extirpation are being developed in the fields of plastic and cosmetic surgery; furthermore, even when adipocytes are grafted to subcutaneous tissue, which allows easy implantation, these cells are not heterotropic since they originally belonged to this region;

(3) since isolated primary cultured adipocytes actively proliferate, even *in vitro*, they are appropriate for procedures such as gene transfer;

40 (4) since adipocytes are likely to stay in a limited area after implantation, the grafted cells can be extirpated after implantation if so desired (specifically, when wanting to eliminate the gene expression);

(5) since adipocytes themselves produce angiogenic factors (Mick, G.J., et al., Endocrinology 2002, 143(3):948-53), a high level of engraftment can be expected after implantation;

(6) adipocyte extirpation or implantation has a small impact on the human body because the weight of this organ changes greatly in adults; and

45 (7) adipocytes are widely recognized as superfluous and obstructive, and consent for their collection may be obtained easily.

[0006] Although investigations with similar objectives are currently underway using keratinocytes (J. Gene. Med. 2001 Jan-Feb, 3(1):21-31; Histochem. Cell Biol. 2001 Jan, 115(1):73-82), removing the biological barrier of the skin in the process of isolating the primary culture is problematic considering infection risk. Patient pain during extirpation and implantation is predicted to be severe, and re-extirpation (4, mentioned above) to eliminate expression is not easy. Furthermore, when using keratinocytes or skin, which can only be grafted two-dimensionally, the amount of the graft can only be increased by increasing the graft surface area. Therefore, adipocytes, which allow three-dimensional transplantation, are considered more useful.

55 **[0007]** The present inventors designed methods for efficiently transferring genes into primary cultured adipocytes. They also confirmed that the transferred genes are functioning after implantation, and found that adipocytes can be effectively utilized in gene therapy. Furthermore, adipocytes that stably express the transferred foreign gene *in vivo* for a long period of time can be obtained by the methods of this invention. The implanted mature adipocytes can continue

to express foreign genes for one year or longer. Furthermore, if expression of the foreign gene becomes unnecessary after adipocyte implantation, expression can be stopped by removing the graft.

[0008] The present invention relates to the following items:

5 1. A pharmaceutical composition comprising a primary cultured preadipocyte, wherein the preadipocyte stably maintains a foreign gene encoding a protein that is secreted outside of the cell and wherein the gene has been inserted into a retroviral vector and has been transferred to the cell by the retroviral vector, wherein the protein is lecithin cholesterol acyltransferase (LCAT).

10 2. A primary cultured preadipocyte, wherein the preadipocyte stably maintains a foreign gene encoding a protein that is secreted outside of the cell, and wherein the gene has been inserted into a retroviral vector and has been transferred to the cell by the retroviral vector, for use in gene therapy, wherein the protein is LCAT.

15 3. The pharmaceutical composition of item 1, or the preadipocyte for use according to item 2, wherein the preadipocyte has the ability to significantly express the protein *in vivo* for at least 20 days.

4. The pharmaceutical composition of item 1 or 3, or the preadipocyte for use according to item 2 or 3, wherein the preadipocyte is used to release the protein into the blood flow.

20 **[0009]** An *in vitro* method of producing a preadipocyte for use in gene therapy, wherein the method comprises the steps of:

- (1) primary culturing a preadipocyte; and
- (2) transferring, and then stably holding a foreign gene encoding a protein that is secreted outside of the cell,

25 wherein the protein is LCAT.

[0010] The method of item 5, wherein the foreign gene is transferred by a retroviral vector.

[0011] The preadipocyte for use according to item 2 which has been produced by the method of item 5 or 6.

30 **[0012]** An implant composition, wherein the composition comprises a primary cultured preadipocyte, which stably holds a foreign gene encoding a protein that is secreted outside of the cell, and wherein the gene has been inserted into a retroviral vector and has been transferred to the cell by the retroviral vector, and a pharmaceutically acceptable carrier, for use in gene therapy, wherein the protein is LCAT.

[0013] The implant composition for use according to item 8, or the pharmaceutical composition of any one of items 1, 3 and 4, which further comprises an extracellular matrix component.

35 **[0014]** The implant composition for use according to item 8 or 9, or the pharmaceutical composition of any one of items 1, 3, 4 and 9, which further comprises an angiogenesis factor.

11. A non-human animal, the body of which has been implanted with the primary cultured preadipocyte of item 2.

40 **[0015]** Hereinafter, the mode for carrying out this invention will be described.

[0016] First, the present invention provides primary cultured adipocytes for gene therapy, where the adipocytes stably maintain a foreign gene(s) encoding a protein(s) that is secreted to the cell exterior.

45 **[0017]** Herein, a foreign gene refers to a lecithin cholesterol acyltransferase (LCAT) gene transferred into primary cultured adipocytes from the outside. Furthermore, primary cultured cells refer to non-established cells that are cultured from tissues-removed from a living body. Adipocytes refer to mature adipocytes and cells comprising the ability to differentiate into adipose tissue, such as preadipocytes. More specifically, unless the adipocytes are particularly said to be "mature" adipocytes, they also include preadipocytes. Mature adipocytes are spherical cells that store fat, and contain lipid droplets. Fat stored in mature adipocytes can be identified using oil red O staining. Mature adipocytes generally secrete leptin in response to insulin. Preadipocytes normally exist as stromal cells that have not yet differentiated into mature adipocytes. Preadipocytes can be isolated by treating adipose tissue with collagenase, or can be isolated as a result of the division of mature adipocytes, using the ceiling culture method described below (Sugihara, et al. Nippon Rinsho 1995, 53, 115-120; Sugihara, H., et al. J. Lipid Res. 1987, 28, 1038-1045; Zhang H.H., et al. J. Endocrinol. 2000, 164, 119-128). Although the existence of adipocyte-specific surface antigens has not been confirmed, high levels of CD36 expression and such have been found in mature adipocytes (Abumrad N.A., et al. J. Biol. Chem. 1993 Aug 25, 268(24):17665-8). Therefore, extremely pure adipocytes may be collected by using such molecules as markers. By inducing differentiation as described below, preadipocytes can differentiate into mature adipocytes within a few days to few weeks (Hauner H., et al., J. Clin. Invest. 84, 1663-1670, 1989; Marko, et al., Endocrinology 136, 4582-4588, 1994). Primary cultured adipocytes can be isolated from a desired tissue, for example, subcutaneous adipose tissue or visceral

adipose tissue such as tissue surrounding the epididymis or mesenteric tissue.

[0018] The phrase "for gene therapy" refers to using the *in vivo* expression of a protein(s) encoded by a foreign gene(s) in anticipation of a therapeutic effect. Furthermore, cells for gene therapy refer to cells carrying a foreign gene(s), in which the cells are used for administering the foreign gene into a body by *ex vivo* administration, and the cells comprise the ability to express the protein in that body. *Ex vivo* administration refers to removing adipose tissues or adipocytes from an individual, performing gene transfer *in vitro*, and then implanting the cells to the same or a different individual.

[0019] Cells for gene therapy preferably refer to cells used for treating disorders, which are cells that are implanted so that a specific protein is produced. Preferably, treatment by a specific protein includes replacement therapy, which uses a protein whose physical or functional deficiency or absence causes a disorder.

[0020] The specific protein is LCAT that shows activity in the bloodstream, or is supplied to a target tissue via the bloodstream, and functions at the cell surface of that tissue. A continuous supply of the specific protein is also preferably required for a certain period of time (for example, for a few days to a few weeks or more). Factors and disorders for which protein replacement therapy is already being carried out, or is predicted to be effective, may all become targets.

[0021] Hereinafter, representative targets are listed according to their classification, but their use is not to be understood as being limited to these examples, and the use of similar factors for similar purposes is included within the scope of this invention.

[0022] Replacement therapy includes supplementation against disorders that develop or are exacerbated by a lack or reduced function of a hormone, supplementation against disorders due to a congenital genetic defect, and supplementation of a factor for pathological improvement:

[0023] Furthermore, the adipocytes of the present invention are not limited to those used for so-called "therapy", but include cells used for *in vivo* expression of a desired secretory protein. For example, the methods of this invention enable production of model animals by a *posteriori* expression of a particular protein. Using these methods, disease model animals with a *posteriori* expression of pathogenesis or aggravative factors can be produced, and these animals can be used to screen drugs. Furthermore, by expressing pathologic improvement factors, these methods can be utilized as proof of working hypotheses for novel drug discoveries in which a given factor improves a pathologic condition. The animals that are used include desired non-human animals, and preferably non-human mammals (including rodents and primates).

[0024] The primary cultured adipocytes for the gene therapies of this invention stably maintain a LCAT gene(s) that encodes a LCAT protein(s) that is secreted outside of the cell. The phrase "stably maintains" means that the foreign gene is passed on to daughter cells during cell division, and more specifically, this phrase refers to the incorporation of the foreign gene into a cell chromosome. The adipocytes for gene therapy of this invention comprise a LCAT gene(s), stably transferred by a chromosome-incorporating viral vector. The LCAT gene is transferred by a retroviral vector.

[0025] The retroviral vector is stably integrated into a cell chromosome and comprises the ability to express a transferred gene for a long period. The vector's transfer efficiency and continuation of expression of the transferred gene depends on the cell type. For example, a gene transferred by a retroviral vector can show continued expression while the cells are growing, but expression may stop when cell growth stops (Lund, A.H., et al., J. Biomed. Sci. 1996, 3:365-378; Niwa, O. et al., 1983, Cell, 32:1105-1113). Foreign gene expression is often observed to be suppressed, particularly after introducing the gene into a body by *in vivo* or *ex vivo* methods. Such suppression of expression is said to involve *de novo* methylation of the promoter or coding sequence of the transferred gene (Jahner, D. and Jaenisch, R., Nature 315:594-597, 1985; Challita, P.-M. and Kohn, D.B., Proc. Natl. Acad. Sci. USA 91:2567-2571, 1994; Hoeben, R.C. et al., J. Virol. 65:904-912, 1991). Furthermore, deacetylation of histone is involved in silencing the transferred gene (Chen, W.Y. et al., Proc. Natl. Acad. Sci. USA 97:377-382, 2000; Chen, W. Y. et al., Proc. Natl. Acad. Sci. USA 94:5798-5803, 1997). However, when the present inventors transferred a foreign gene into primary cultured adipocytes using a retroviral vector, surprisingly, expression of the transferred gene was found to persist extremely stably, both *in vitro* and *in vivo*. Expression of transferred genes is stable in adipocytes before differentiation and also in mature adipocytes. Expression of the transferred gene was confirmed to persist for the entire duration of the experiment for *in vitro* cultures (80 days or more), and for the entire duration of the experiment when implanted into the body (360 days or more). Therefore, primary cultured adipocytes, to which a LCAT gene (s) has been stably transferred, can be used as implants that stably express a gene(s) for a long period.

[0026] The adipocytes for gene therapy of this invention comprise the ability to significantly express a LCAT protein(s) encoded by a LCAT gene(s) for at least 20 days or more *in vitro*, or more preferably *in vivo*. The phrase "significantly express" means, for example, expression is detected at a statistically significant level compared to when the foreign gene is not transferred (for example, with a significance level of 5% or a higher significance). More preferably, the adipocytes of the present invention, when transplanted into a body, comprise the ability to significantly express a LCAT protein(s) encoded by a LCAT gene(s) in the body for at least 30 days or more, preferably 40 days or more, more preferably 50 days or more, even more preferably 60 days or more, still more preferably 80 days or more, yet even more preferably 100 days or more, yet even more preferably 150 days or more, yet even more preferably 200 days or more, yet even more preferably 250 days or more, yet even more preferably 300 days or more, and yet even more preferably

350 days or more.

[0027] The adipocytes for gene therapy of this invention are particularly useful as cells for releasing LCAT proteins, that are encoded by LCAT genes carried by the cells, into the blood flow. The LCAT proteins released into the blood flow are desired secretory proteins that demonstrate activity in the blood stream or at the surface of cells of target tissues.

[0028] The present invention also relates to methods of producing adipocytes for use in gene therapy, where the methods comprise the steps of:

- (1) primary culturing pre-adipocytes, and
- (2) transferring cells with a LCAT gene(s) that encodes a LCAT protein(s) that is secreted to the cell exterior, preferably using a retroviral vector or an adeno-associated viral vector, so that the gene is stably maintained.

The present invention also relates to the adipocytes for gene therapy produced by this method. "Stably maintained" means transfer of a foreign gene(s) such that it is passed on to daughter cells when the cell divides, and more specifically, it refers to integration of the foreign gene into the chromosome of the cells. Southern blotting or PCR using genomic DNA can molecular biologically demonstrate that the foreign gene has achieved stable expression by integrating into a chromosome. Furthermore, to concentrate the stably transfected cells, for example, a method using fluorescence activated cell sorting (FACS), which concentrates cells by recognizing the GFP coexpressed by the cells along with the target gene, may be used.

1. Methods of collecting primary cultured adipocytes

[0029] Primary cultured adipocytes can be collected by methods described in the report by Sugihara *et al.* (Sugihara, H. *et al.*, *Differentiation*, 31:42-49, 1986). More specifically, adipose tissue, and preferably the implant recipient's own subcutaneous adipose tissue or visceral adipose tissue, such as tissue surrounding the epididymis or mesenteric tissue, is extirpated under sterile conditions, and for example, after washing with PBS, is morcellated using a pair of scissors or a surgical knife. This morcellated tissue is digested by shaking at 37°C in a medium comprising an appropriate amount of collagenase, preferably 1 to 3 mg/mL, for an appropriate length of time, preferably for 20 to 60 minutes, and then separated into a precipitated residue and floating layer by centrifugation.

[0030] The floating layer is preferably further washed once or twice by centrifugation, and is then added to a culture flask filled with medium. Bubbles are removed, and the flask is left to stand in a CO₂ incubator for culturing, such that the conventional culture surface is a ceiling (ceiling culture). After culturing for an appropriate period, preferably ten to 14 days, cells adhered to the ceiling surface are collected by trypsin treatment. These cells are subsequently subcultured in a conventional culturing system.

[0031] Primary cultured adipocytes may be stored by freezing before or after gene transfer. This procedure allows multiple use of adipocytes after a single collection.

2. Gene transfer to adipocytes

[0032] Gene transfer can be performed using gene transfer reagents (Fugene 6, Roche; Lipofectamin, Invitrogen; Cellfect transfection kit (calcium phosphate method), Amersham; etc.), electroporation methods (Chen, H. *et al.*, *J. Biol. Chem.* 1997, 272(12), 8026-31), or viral vectors (Kay, M.A., *et al.*, *Nat. Med.* 2001, 7, 33-40). Transfer is preferably performed using viral vectors, and more preferably using retroviral vectors (*e.g.*, Arai, T. *et al.*, *J. Virol.*, 1998, 72, pp1115-21).

[0033] When gene transfer is performed using a plasmid, the plasmid is transfected into adipocytes, and those adipocytes stably maintaining the transferred foreign gene can be selected. Such adipocytes can be selected by, for example, equipping the plasmid encoding the foreign gene with a drug-resistance gene, or by performing the transfection together with a plasmid carrying a drug-resistance gene, and then selecting the transfected cells using this drug. Otherwise, the cells can be obtained by cloning the transfected cells by limiting dilution techniques. Furthermore, when gene transfer is performed using a plasmid, a method of transiently expressing a phage-derived integrase can be used to increase the efficiency of chromosomal insertion (*Mol. Cell Biol.* 2001 Jun, 21(12):3926-34).

[0034] In the present invention, the LCAT gene is transferred into adipocytes using a retroviral vector. Retroviruses refer to viruses that belong to the Retroviridae family, and include oncoviruses, foamy viruses (Russell, D.W. and Miller, A.D., *J. Virol.* 1996, 70:217-222; Wu, M. *et al.*, *J. Virol.* 1999, 73:4498-4501), and lentiviruses (for example, HIV-1 (Naldini, L. *et al.*, *Science* 1996, 272:263-267; Poeschla, E. *et al.*, *Proc. Natl. Acad. Sci. USA* 1996, 93:11395-11399; Srinivasakumar, N. *et al.*, *J. Virol.* 1997, 71:5841-5848; Zufferey, R., *et al.* *Nat. Biotechnol.* 1997, 15:871-875; Kim, V.N., *et al.*, *J. Virol.* 1998, 72:811-816) and feline immunodeficiency virus (Johnston, J.C. *et al.*, *J. Virol.* 1999, 73:4991-5000; Johnston, J. and Power, C., *J. Virol.* 1999, 73:2491-2498; Poeschla, E.M. *et al.*, *Nat. Med.* 1998, 4:354-357). A preferable retroviral vector for use in this invention is a Moloney murine leukemia virus (MoMLV) vector (T. M. Shinnick, R. A. Lerner

and J. G. Sutcliffe, Nature 293, 543-548, 1981).

[0035] The retroviruses may be self-inactivating (SIN) vectors. A SIN vector can be prepared by deleting a portion of the 3' LTR during viral packaging (Yu S.F. et al. (1986) Proc. Natl. Acad. Sci. USA 83:3194; Yee, J. K. et al., 1987, Proc. Natl. Acad. Sci. USA 84:5197-5201; Zufferey, R. et al., 1998, J. Virology, 72, 9873-9880). The foreign gene in the retrovirus can be transcribed by LTR, or it may be expressed from another promoter inside the vector. For example, a constitutive expression promoter such as CMV promoter, EF-1 α promoter, or CAG promoter, or a desired inducible promoter may be used. Furthermore, a chimeric promoter, in which a portion of LTR is substituted with another promoter, may be used.

[0036] To transfer genes using retroviruses, specifically, a plasmid carrying a gene to be transferred, such as pBabe CL-SEAP-IRES-GFP, is gene-transferred to packaging cells, such as 293-EBNA cells (Invitrogen), using a gene transfer reagent and such. This is then cultured for an appropriate period of time, preferably one to three days, and the produced recombinant viruses in the supernatant are collected. These viruses are then infected into the adipocytes to be transfected.

[0037] The retroviral vectors preferably comprise an envelope protein with broad tropism, so that they can infect a wide range of mammalian adipocytes, including those of humans. For example, amphotropic envelope protein may be used (for example 4070A) (Accession K02729; Sorge, J. et al., Mol. Cell. Biol. 4 (9), 1730-1737 (1984)). In the present invention the retrovirus is preferably pseudotyped (Emi, T. Friedmann and J. K. Yee, J. Virol., 65 (3), 1202-1207 (1991); Yee, J.-K. et al. (1994) Methods Cell Biol. 43 43:99-112; Burns, J. C. et al. (1993) Proc. Natl. Acad. Sci. USA 90 90:8033-8037) by vesicular stomatitis virus G protein (VSV-G) (Rose, J.K. and Gallione, C.J., J. Virol. 39 (2), 519-528 (1981)). Pseudotyping by VSV-G enables highly efficient transfer -of genes into adipocytes. VSV-G pseudotyped vector can be produced by expressing VSV-G in packaging cells. More specifically, for example, packaging cells that can inducibly express VSV-G may be used favorably (for example, Arai T. et al., J. Virol., 1998: 72, pp1115-21).

[0038] The titer of the produced viruses can be determined by infecting cells with virus solutions that have been stepwise diluted, and counting the number of colonies of infected cells (for details, see Ausubel *et al.*) (Ausubel, F.M. et al. Eds. (1995) Current Protocols in Molecular Biology. (John Wiley & Sons, NY)). Alternatively, the titer can be determined by the method of Byun *et al.* (Byun, J. et al. (1996) Gene Ther. 33333:1018-1020), Tafuro *et al.* (Tafuro, S. et al. (1996) Gene Ther. 33333:679-684), Miyao *et al.* (Miyao, Y. et al. (1995) Cell Struct. Funct. 20 20:177-183), Claudio *et al.* (Claudio, P. P. et al. (2001) Anal. Biochem. 291: 96-101), or Cashion *et al.* (Cashion, L. M. et al. (1999) Biotechniques 26 26: 924-930).

[0039] Primary cultured adipocytes can be introduced with viral vectors by contacting the vectors to the cells. For example, primary cultured adipocytes are incubated in a culture solution comprising viral vectors. Adipocytes are preferably infected in the form of preadipocytes. Infection efficiency can be increased by adding 0.5 to 8 μ g/mL or so of polybrene. Multiplicity of infection (MOI) is not particularly limited, but can be appropriately adjusted within the range of 0.1 to 100. Gene transferred cells can be selected using a marker gene, for example. However, if infection is carried out at an MOI of approximately 2 or more, or preferably approximately 3, 4, 5, or more, the gene can be transferred to most cells, even without selection. The gene-transferred adipocytes can be used for implantation without further treatment, or in certain cases, they can be converted to mature adipocytes by culturing in a medium comprising 3-isobutyl-1-methylxanthine (IBMX), dexamethasone, and insulin. In such cases, since IBMX and dexamethasone are used mainly to activate the adipocyte peroxisome proliferator-activated receptor- γ (PPAR- γ), drugs that directly activate this receptor (for example the thiazolidine derivatives, pioglitazone/Takeda Pharmaceutical Company Limited and rosiglitazone/GlaxoSmithKline) may be added at the same time.

[0040] The primary cultured adipocytes of this invention, which carry a desired therapeutic LCAT gene, can be implanted into the body of an immunologically matched recipient, thus enabling gene therapy by *in vivo* expression of the secretory protein encoded by the therapeutic gene. The primary cultured adipocytes to be implanted are preferably cells from the same host as the recipient. The gene therapy methods in which the primary cultured adipocytes of this invention are implanted can be applied by expressing a desired LCAT secretory protein in a body, in anticipation of that protein's effects. For example, a disorder can be treated or prevented by implanting the adipocytes of this invention, which maintain a foreign LCAT gene(s) encoding a LCAT protein (s) comprising a therapeutic or preventive effect against the disorder. Furthermore, the present invention relates to methods of releasing proteins into the blood flow, where the methods comprise the step of administering the primary cultured adipocytes of this invention into a body. Using these methods, the LCAT protein encoded by a foreign LCAT gene can be significantly secreted into the blood flow for at least 20 days or more, preferably 30 days or more, more preferably 40 days or more, even more preferably 50 days or more, still more preferably 60 days or more, yet even more preferably 80 days or more, yet even more preferably 100 days or more, yet even more preferably 150 days or more, yet even more preferably 200 days or more, yet even more preferably 250 days or more, yet even more preferably 300 days or more, and yet even more preferably 350 days or more. The foreign LCAT gene expressed in a body can be detected and/or quantified, for example by immunoassays such as EIA. Removal of the transplanted cells can stop the expression of the administered foreign LCAT gene at any time. In certain cases, by transferring an inducible suicide gene (*e.g.*, HSV-tk) to the graft cells, the graft cells can be eliminated by administering ganciclovir, for example.

[0041] The present invention also provides implant compositions for use in gene therapy, where the compositions comprise primary cultured adipocytes that stably hold a LCAT gene(s) that encodes a LCAT protein(s) secreted to the cell exterior, and pharmaceutically acceptable carriers. Examples of the carriers are physiological saline, phosphate buffer, culture solutions, serums, and body fluids. These may also be combined with a solid or gel support that becomes a scaffold for cells.

[0042] The implant compositions of the present invention preferably comprise an extracellular matrix (ECM) component. An extracellular matrix component refers to a component such as a protein or mucopolysaccharide comprised in an insoluble network or fibrous structure accumulated between cells. They may be isolated from organisms or artificially reconstructed. ECM components preferably used in this invention are collagen, fibronectin, vitronectin, laminin, heparan sulfate, proteoglycan, glycosaminoglycan, chondroitin sulfate, hyaluronate, dermatan sulfate, keratin sulfate, elastin, or combinations of two or more of the above. Preferably, these ECM components are formed into a gel and then mixed with adipocytes. ECM gels used in this invention are not particularly limited, as long as at least one or more of the above-mentioned components are comprised, but preferably comprise at least type IV collagen, laminin, and heparan sulfate. Such ECMs include a substrate extracted from Engelbreth-Holm-Swarm mouse tumor (Matrigel®) (Becton Dickinson Labware) (US Patent No. 4,829,000). The structure of the compositions comprising the ECM component and adipocytes used in the present invention is not particularly limited, and may be, for example, a gel or paste network structure, a fibrous structure, flat (disc) structure, honeycomb structure, and sponge-like structure. ECM components can be gelled according to conventional methods. For example, gelation can be performed by incubating an aqueous solution comprising approximately 0.3 to 0.5% collagen at 37°C for ten to 30 minutes. Otherwise, ECM components can be gelled using a gelation agent.

[0043] Furthermore, the implant compositions of the present invention preferably comprise an angiogenesis factor. The implant compositions of this invention that comprise an angiogenesis factor cause blood vessels to form around them after implantation, and can secrete a foreign protein into the blood flow with higher efficiency. The angiogenesis factors are not particularly limited, as long as they are factors that may induce angiogenesis *in vivo*, and examples are vascular endothelial cell growth factor (VEGF), basic fibroblast growth factor (bFGF), acidic fibroblast growth factor (aFGF), platelet-derived growth factor, transforming growth factor- β (TGF- β), osteonectin, angiopoietin, and hepatocyte growth factor (HGF). The most preferred example is bFGF. bFGFs, which are also called FGF2, are not only fibroblast growth factors, but also comprise the activity of promoting the growth of various cells such as vascular endothelial cells, cartilage, osteoblasts, and epidermal cells (Abraham et al., EMBO J., 5, 2523-2528, 1986; Prats et al., Proc. Natl. Acad. Sci. USA, 86, 1836-1840, 1989). The bFGFs used in the present invention are not only natural proteins, but may also be produced by genetic engineering by recombinant DNA technology, and modified forms thereof. Examples of bFGFs are those described in WO87/01728, WO89/04832, WO86/07595, WO87/03885, European Patent Application Publication Nos. 237966, 281822, 326907, 394951, and 493737. Alternatively, another expression vector that transiently expresses an angiogenesis factor may be introduced into the adipocytes (see WO97/49827). The main objective of angiogenesis factors used in this manner is to form blood vessels around the transplanted cells, so that the foreign protein can be efficiently secreted into the blood flow from the adipocytes of this invention. Therefore, when using a vector encoding a vascular inducing factor to express that vascular inducing factor from adipocytes, the use of a transient expression vector (more specifically, a vector that is not incorporated into the chromosome) is preferred. When the adipocytes express a vascular inducing factor for a long period, excess amounts of blood vessels form around the implanted adipocytes, which may cause systemic side effects. Therefore, it is preferable that the foreign gene encoding an angiogenesis factor is not stably transferred to the primary cultured adipocytes of this invention.

3. Implantation of adipocytes

[0044] Gene transferred adipocytes are prepared at an appropriate cell concentration, preferably 0.2×10^7 to 2×10^7 cells/mL, or 0.2×10^6 to 5×10^6 cells/mL when transfected with a retrovirus. They are infused as is into the subcutaneous tissue or adipose tissue, preferably subcutaneous tissue, or by mixing with an effective media, preferably a solution comprising an extracellular matrix such as collagen. Injection into adipose tissue can be performed by making an incision and exposing the adipose tissue. Cells that have terminally differentiated into mature adipocytes will not proliferate after transplantation, and will express the foreign gene for a long period at a constant level. The expression level of a foreign gene in a body that receives an implant is proportional to the number of implanted cells. Therefore, when performing an implantation, a desired expression level can be maintained for a long period in a body receiving an implant by adjusting the amount of adipocytes that are implanted to align with a pre-measured *in vitro* foreign gene expression level.

Brief Description of the Drawings

[0045]

Fig. 1 is a set of microphotographs of primary cultured adipocytes isolated from the subcutaneous fat of three-week old ICR mice. (A) shows adipocytes that adhered to the ceiling-side culture surface after 14 days of ceiling culture, (B) shows primary cultured adipocytes grown in a normal culture, (C) shows mature adipocytes that have stored lipid droplets due to differentiation induction, and (D) shows an oil red O-stained image of differentiation-induced cells.

Fig. 2 shows the plasma alkaline phosphatase (AP) activity obtained by implanting ICR nude mice with primary cultured adipocytes (derived from subcutaneous fat of ICR mice) that are transiently transfected with AP-expressing plasmid pcDNA3.1-SEAPmh.

Fig. 3 shows a comparison of gene transfer efficiency when retroviral vector MLV(VSV)/pBabeCL(PLAP)IP is transduced to primary cultured adipocytes derived from various adipose tissues.

Fig. 4 is a set of microphotographs showing images of the differentiation induction of primary cultured adipocytes transduced with MLV(VSV)/pBabeCL(GFP)IP. (A) and (B) respectively show a light-microphotograph, and a GFP fluorescence photograph of the same visual field.

Fig. 5 shows the duration of AP expression in subcultures of primary cultured adipocytes transduced with an AP-expressing viral vector. (A) shows the result of transferring SEAP gene (MLV(VSV)/pBabeCL(SEAPmh)I2G) or PLAP gene (MLV(VSV)/pBabeCL(PLAP)IP) to cells derived from C57BL/6 mice subcutaneous fat. (B) shows the result of transferring PLAP gene (MLV(VSV)/pBabeCL(PLAP)IP) or GFP gene (MLV(VSV)/pBabeCL(GFP)IP) into adipocytes derived from ICR mice.

Fig. 6 is a set of photographs and a graph showing the change in expression in differentiation-induced gene-transferred adipocytes. (A) shows a GFP light microscope image of primary cultured adipocytes under non-differentiation-inducing conditions, where the adipocytes transfected with MLV(VSV)/pBabeCL(GFP)IP are derived from ICR subcutaneous fat. (B) shows a similar GFP microscope image taken under differentiation-inducing conditions. (C) shows AP production by MLV(VSV)/pBabeCL(PLAP)IP-transfected primary cultured adipocytes (derived from ICR subcutaneous fat) under non-differentiation-inducing conditions (non-differentiation) and differentiation-inducing conditions (differentiation).

Fig. 7 shows (pro)insulin production by plasmid transfection into primary cultured adipocytes.

Fig. 8 shows the stable expression of AP in primary cultured adipocytes (derived from C57BL/6 mice subcutaneous fat) transfected with AP-expressing AAV.

Fig. 9 shows insulin expression at the time of differentiation induction in primary cultured adipocytes transfected with s1s2B10 insulin-expressing retroviral vector. (A) shows the results using an EIA produced by Morinaga and (B) shows the results using an EIA produced by IBL.

Fig. 10 shows the expression of GLP-1(7-37) in primary cultured adipocytes transfected with GLP-1(7-37)-expressing retroviral vector. Measurements were made in triplicate, and their average values and standard deviations are shown.

Fig. 11 shows the effect of the presence or absence of pre-implantation stimulation of differentiation induction on *in vivo* AP expression in the implantation of AP-expressing primary cultured adipocytes.

Fig. 12 is a set of graphs and a photograph. (A) shows the change in plasma AP activity when AP-expressing primary cultured adipocytes are implanted in the presence of differentiation stimulation using a basic FGF-supplemented Matrigel. (B) shows the loss of plasma AP activity on extirpation of the implanted Matrigel (individual A). (C) shows a GFP light microscope image of the Matrigel extirpated from the control group which received GFP-transfected cells. For the PLAP-implanted group shown in (A), the values shown are the group average and standard deviation of values measured for each individual up to the 32nd day. The remaining values are average values.

Fig. 13 shows the results of the long-term examination of AP activity in the blood of mice receiving an implant by the method of Fig. 12(A), and by a variety of other methods.

Fig. 14 shows the results of performing an extirpation test similar to that of Fig. 12(B) in the late stage of transplantation.

Fig. 15 shows the dependence of blood AP activity on the number of implanted cells when implanting AP-expressing adipocytes. The values indicated are the group average and standard deviation of the measurements of each individual.

Fig. 16 shows the effect of implanting s1s2B10 insulin-expressing adipocytes to STZ-induced diabetic mice. (A) shows the effect on fasting plasma glucose level, and (B) shows the effect on body weight. The values indicated are the group average and standard deviation of the measurements of each individual.

Best Mode for Carrying out the Invention

[0046] The present invention will be described in detail below with reference to Examples, but it is not to be construed as being limited thereto. All references cited herein are incorporated into this description.

[Example 1] Primary culture of murine adipocytes

[Methods]

5 **[0047]** Three-week old male ICR mice or four- to five-week old male C57BL/6 mice (both from Charles River) were anesthetized with diethyl ether, and sacrificed by collection of whole blood from the heart. Next, inguinal subcutaneous fat, or fat surrounding the epididymis, and mesenteric adipose tissue were individually extirpated under sterile conditions. The extirpated tissues were washed with PBS, and then morcellated using a pair of scissors or a surgical knife. This morcellated tissue was digested with shaking at 37°C for 20 to 60 minutes in normal medium (DMEM-high glucose/SIGMA, 10% FCS) comprising 1 mg/mL of collagenase (S1 fraction/Nitta gelatin), and then separated into precipitate and suspended layer by centrifugation (300 g, five minutes).

10 **[0048]** The floating layer was further centrifuged once or twice to remove the collagenase by dilution, and then added to a T-25 flask (IWAKI) filled with medium. Bubbles were removed, and this was cultured under a 5% CO₂ atmosphere in a CO₂ incubator at 37°C so the conventional culture surface was upside (ceiling culture). Ten to 14 days after culturing, the cells adhering to the ceiling surface were collected by trypsin treatment and transferred to a normal culturing system. Subculturing was then performed at a ratio of 1:3 to 1:10.

15 **[0049]** To induce differentiation, the medium of cells cultured to confluency in a 6-well plate was transferred to an induction medium (normal medium supplemented with 0.5 mM IBMX, 0.25 μM dexamethasone, and 10 μg/mL insulin). This stimulation was continued for 48 hours. Next, the cells were differentiated in a maturation medium (normal medium supplemented with 10 μg/mL insulin). The maturation medium was exchanged every three days.

20 **[0050]** Oil red O staining solution was prepared by mixing a stock solution, prepared by mixing 0.3 g of oil red O in 100 mL isopropanol (99%), with distilled water in a 3:2 ratio at the time of use. The cells were washed with PBS and then fixed with 10% neutral formalin solution (WAKO). After washing again with PBS, the cells were stained with oil red O staining solution at room temperature for ten minutes. The cells were washed with PBS again, and then examined by microscope.

[Results]

30 **[0051]** Fig. 1 is a set of microphotographs of primary cultured adipocytes isolated from the subcutaneous fat of three-week old ICR mice. After 14 days of ceiling culture, adhesion of adipocytes carrying lipid droplets was observed on the ceiling-side culture surface (A). When these cells were transferred to a normal culturing system, they showed fibroblast-like growth, as shown in (B). However, when differentiation was induced by IBMX, dexamethasone, and insulin, the cells again differentiated into mature adipocytes that carry lipid droplets (C). Stored fat was stained red with oil red O staining (D). Cells isolated by this method were shown to be primary cultured adipocytes comprising the ability to differentiate.

35 [Example 2] Transient transfer of thermostable secretory alkaline phosphatase (AP) gene into primary cultured adipocytes, and implantation of transfected adipocytes into mice

40 **[0052]** As a model system for gene expression, AP gene, more specifically, SEAP gene (Clontech) or PLAP gene (Goto, M. et al. Mol. Pharmacol. vol.49 860-873 (1996)) was transferred to primary cultured adipocytes, and changes in AP activity were examined. (Both AP gene products are thermostable and can be easily distinguished from endogenous alkaline phosphatases by thermal treatment.)

[Methods]

45 (1) Production of primary cultured adipocytes transiently transfected with the SEAP gene

50 **[0053]** AP-expressing plasmid (pcDNA3.1-SEAPmh) was constructed by inserting the SEAP sequence, obtained by double digestion of pSEAP2-basic vector (Clontech) with restriction enzymes HindIII-XbaI, into the *HindIII-XbaI* site of pcDNA3.1Myc-HisA (Invitrogen), which is a vector for expression in mammalian cells.

55 **[0054]** For every gene transfer to a 10-cm dish, 500 μL of FCS-free DMEM medium and 15 μL of Fugene 6 reagent (Roche) were mixed, then 5 μg of pcDNA3.1-SEAPmh was added. This mixture was left to stand at room temperature for 15 minutes. This mixture was added to primary cultured cells (derived from ICR subcutaneous fat) cultured to 70 to 80% confluency in a 10-cm dish. This was then cultured for 24 hours in a CO₂ incubator.

(2) Implanting mice with alkaline phosphatase gene-transferred primary cultured adipocytes

[0055] Gene-transferred cells were collected by trypsin treatment, and washed twice with PBS by centrifugation. The

cells were then suspended in PBS at 1×10^7 cells/mL. The animals (ICR nude mice, five-weeks old at the time of operation) were anesthetized by intraperitoneal administration of 50 mg/kg of sodium pentobarbital (Nembutal; Dainippon Pharmaceutical). After disinfecting the area to be operated with dilute Hibitane solution (Sumitomo Pharmaceuticals), a 3 mm to 5 mm or so incision was made to the skin near the base of the right hind leg, and the inguinal subcutaneous fat was exposed. 0.55 mL of the prepared cell suspension solution (5.5×10^6 cells/head) was loaded into a 1-mL syringe, and this was injected into the subcutaneous fat using a 22 G injection needle. As a control, PBS was injected to the same site. To compare this to the protein supplementation method, 1 μ g of purified AP (Roche) was dissolved in PBS under sterile conditions, and this was injected in a similar manner. The incised skin was sutured and the operated site was disinfecting with surgical Isodine (Meiji Seika).

[0056] Blood was collected using a heparin-coated capillary (Dramond) from the postorbital venous plexus before implantation (day 0) and after implantation over time. Plasma was obtained from the whole blood by centrifugation at 2000 g for 15 minutes. AP activity in this plasma was measured using an assay kit (SEAP reporter gene assay kit, Roche) by following the attached instructions.

[Results]

[0057] Fig. 2 shows the plasma AP activity achieved by implanting mice with primary cultured cells, which have been transiently transfected with alkaline phosphatase (AP)-expressing plasmid pcDNA3.1-SEAPmh. For purposes of comparison, mice were administered with 1 μ g of purified AP protein (Roche) by injection. Seven days after administration the blood AP activity in these mice decreased to the level of the control. On the other hand, blood AP activity in mice that received an implant of cells holding transiently transferred genes was confirmed to peak on the fourth day after implantation, and the duration of expression was 14 days. The duration of *in vivo* expression by implanting cells carrying transiently transferred gene was short, and the concentration in the blood was found to vary greatly, although it was maintained longer than by injecting protein.

[Example 3] Production, by using a viral vector, of adipocytes that stably express AP

[Methods]

(1) Construction of AP- and control GFP-expression vectors

[0058] The PLAP gene was excised from pTK-PLAP using *HindIII* and *BglII*, as described in the literature (Goto, M. et al. Mol. Pharmacol. vol. 49, 860-873 (1996)). The SEAP gene was obtained by double digestion of pcDNA 3.1-SEAPmh with *HindIII/PmeI*. The GFP gene was excised from pEGFP-N2 using *NotI-NcoI*.

[0059] The plasmid, pBabeCLX12G, used for viral vector production, was produced based on pBabePuro (Morgenstern, J.P. et al. Nucleic Acids Res. vol.18, 3587-3596 (1990)), by excising its SV40 promoter and neomycin resistance genes using *SalI-ClaI*, and blunting those ends with Klenow fragments, then replacing these with the internal ribosome re-entry site (IRES) of encephalomyocarditis virus (EMCV), which was excised from pIRES2-EGFP by *HincII-HincII*, and the green fluorescent protein (GFP); then replacing the portion from the long terminal repeat (LTR) to the foreign gene insertion site (multicloning site) (*SspI-BamHI*) with a sequence corresponding (*SspI-BamHI*) of pCLXSN (IMGENEX). Furthermore, pBabeCLXIP, in which the IRES-GFP portion of pBabeCLX12G had been replaced with IRES-puromycin resistance gene, was also used.

[0060] Each of the DNA fragments of the above-mentioned PLAP, SEAP, and GFP were blunted with Klenow fragments, then inserted into pBabeCLXIP or pBabeCLX12G vector cleaved with *HpaI*, yielding pBabeCL(PLAP)IP, pBabeCL(SEAPmh)12G, and pBabeCL(GFP)IP, respectively.

(2) Production of viral vectors

[0061] Each gene transfer to a 10-cm dish was performed as follows: 30 μ L of plasmid transfection reagent TransIT (MIRUS) was mixed into 500 μ L of FCS-free DMEM medium, and left to stand at room temperature for five minutes (mixed DMEM/TransIT solution). In a separate tube, 3.3 μ g of a vector encoding VSV-G (pCALG, modified according to Arai, T. et al., J. Virol., 1998, 72, pp1115-21), 3.3 μ g of a vector encoding Gag-Pol (pCLAmpho/RetroMax system (IMGENEX)), and 3.3 μ g of a vector comprising a packaging signal and the transferred gene (pBabeCL(PLAP)IP, pBabeCL(SEAPmh)12G, or pBabeCL(GFP)IP), were mixed, totaling 9.9 μ g (plasmid solution). The plasmid solution was added to the mixed DMEM/TransIT solution, thoroughly mixed, and then left to stand at room temperature for 15 minutes. This was then added to 293-EBNA cells (Invitrogen), cultured overnight from 2×10^6 cells/10-cm dish on the previous day.

[0062] Medium was exchanged eight hours after addition, and the culture supernatant was collected after culturing for another two days. The collected culture supernatant was centrifuged (300 g, five minutes) or filtered through a 0.45

EP 2 213 730 B1

μ m syringe filter (Millipore) to remove contaminants, and this supernatant was used as the virus solution (MLV(VSV)/pBabeCL(PLAP)IP, MLV(VSV)/pBabeCL(SEAPmh)I2G, and MLV(VSV)/pBabeCL(GFP)IP, respectively). Some of the virus solution was concentrated by ultracentrifugation (19,500 rpm, 100 minutes) and then used.

5 (3) Gene transfer to and culturing of primary cultured adipocytes

[0063] Adipocytes to be used for gene transfer (derived from subcutaneous fat, fat surrounding the epididymis, and mesenteric fat of ICR mice, and the subcutaneous fat of C57BL/6 mice) were prepared in 6-well or 96-well plates so that they were 50 to 80% confluent by the day before transfection. The medium was discarded, and equal amounts of
10 4 μ g/mL Polybrene (SIGMA) solution and virus solution were added to the cells to transduce the viral vector. Eight hours after transduction, the medium was changed to a normal medium, and further culturing and subculturing were performed. The AP activity of a portion of the cells was measured by collecting the 24-hour culture supernatant on day four after transfection (Fig. 3).

[0064] Subculturing was performed according to the method of Example 1 on a 10-cm-dish scale. Cells were cultured
15 for four to seven days, and medium was exchanged on reaching confluence. AP activity was measured in the culture supernatant 17 hours later. These cells were continuously subcultured and by appropriately performing similar manipulations, maintenance of expression was examined (Figs. 5 and 6). AP activity was not measured every time subculturing was performed.

[0065] Differentiation was induced in 6-well plates according to the method of Example 1. However, treatment was
20 performed for three days with induction medium, which was replaced with maturation medium every three days thereafter. The AP activity of the culture supernatant was measured using the culture supernatant obtained every three days, and the x-axes in the figures show the day on which the supernatant was collected. For the GFP-transfected cells, microphotographs were taken under appropriate GFP light (Figs. 4 and 6). Non-differentiation-inducing conditions refer to conditions in which culturing is continued in a normal medium instead of an induction medium or mature medium.

25 [Results]

[0066] Fig. 3 is a comparison of the gene transfer efficiency for each kind of tissue-derived cell when using retroviral
30 vectors. AP activity was confirmed in the culture supernatant of all cells when gene transfer was performed on the primary cultured adipocytes isolated from each of the adipose tissues existing in the inguinal subcutaneous tissue, area around the epididymis, and mesentery of ICR mice. This showed that retroviral vectors can transfer genes regardless of the site of cell origin.

[0067] Fig. 4 is a set of microphotographs showing images of the differentiation induction of cells transduced with a
35 GFP-expressing retroviral vector. Differentiation induction was initiated 13 days after gene transfer, and the photographs were taken three weeks later. GFP fluorescence was observed in cells containing lipid droplets, which showed that the viral vector can transfer genes into preadipocytes that possess the ability to differentiate, and that gene transfer by the vector does not affect their ability to differentiate.

[0068] Fig. 5 shows the continuity of expression in the subcultures of primary cultured adipocytes transfected with an
40 AP-expressing viral vector. AP activity was measured in culture supernatant taken 17 hours after cells reached confluency in a 10-cm dish. Continuous AP production was confirmed over the 87 days for which primary cultured adipocytes derived from C57BL/6 mice subcutaneous fat were examined (A), and over the 63 days for which primary cultured adipocytes derived from ICR mice subcutaneous fat were examined (B). These results showed that transduction of the viral vector to primary cultured adipocytes can produce stable expression cells that maintain the foreign genes in the daughter cells produced after division.

[0069] Fig. 6 is a set of photographs and a graph showing changes of expression in differentiation-induced gene-
45 transferred adipocytes. GFP-expressing adipocytes derived from ICR subcutaneous fat showed strong GFP expression under both normal culture conditions (A), and differentiation-inducing conditions (B). Furthermore, AP-expressing adipocytes derived from ICR subcutaneous fat showed continuous expression of AP under both non-differentiation-inducing conditions (non-differentiation) and differentiation-inducing conditions (differentiation) (C). The primary cultured adipocytes that were gene transferred by the viral vectors were found to stably express genes at any phase, not only under
50 the proliferation conditions described in Fig. 5, but also under non-differentiation-inducing conditions, or more specifically under non-proliferative conditions or mature conditions.

[Example 4] Production, by using a plasmid vector, of adipocytes that stably express insulin

55 **[0070]** Methods of gene transfer include methods that use plasmid vectors.

[Methods]

(1) Isolation and modification of the human insulin gene

5 **[0071]** PCR was performed on a human pancreas-derived cDNA library (Stratagene), using the primers shown in Table 1 (Insulin Fw and Rv). A human insulin gene fragment was obtained. The nucleotide sequence of this obtained 354-bp fragment was determined, and the fragment was subcloned into pCR2.1TOPO vector (Invitrogen) as native insulin.

Table 1
Primer sequences used for PCR

Primer	Nucleotide sequence (5'-)
Insulin Fw	CATAAGCTTACCA TGG CCCTGTGGATGCGC (SEQ ID NO: 1)
Insulin Rv	CATTCTAGACT AG TTGCAGTAGTTCTCCAG (SEQ ID NO: 2)
15 site1	CTTCTACACACCCAG <u>GACCAAG</u> CGGGAGGCAGAGGAC (SEQ ID NO: 3)
site2	CCCTGGAGGG <u>ATCCCGG</u> CAGAAGCGTGG (SEQ ID NO: 4)
B10	CACCTGTGCGG <u>ATCCG</u> ACCTGGTGAAGC (SEQ ID NO: 5)
sPL-GLP-1Fw	TTCCACCATGCTGCTGCTGCTGCTGCTGCTGCTGGGCCTGAGGCTACAGCTCT- 20 -CCCTGGGCCATGCTGAAGGGACCTTTACCAGTG (SEQ ID NO: 6)
sPL-GLP-1Rv	AATTATCCTCGGCCTTTACCAGCCAAGCAATGAACTCCTTGGCAGCTTG- 25 -GCCTTCCAAATAAGAACTTACATCACTGGTAAAGGTCCTTCAGC (SEQ ID NO: 7)
GLP-5'	TTCCACCA TG CTGCTGCTGC (SEQ ID NO: 8)
GLP-3'	AATTA T CCTCGGCCTTTACCAG (SEQ ID NO: 9)

(The bold letters denote the initiation codon in Fw, and the antisense of the stop codon in Rv. The underline indicates mutated portions.)

30 **[0072]** Next, in order to express mature insulin in the adipocytes, genetic modification was performed based on literature (JBC, 1994, 269(8), 6241-). More specifically, primers of both directions were individually synthesized to contain mutations at each of the junction sites between the human insulin B chain and the C peptide (site1) between the same C peptide and A chain (site2), and the 10th histidine residue of the B chain (B10) (Table 1). The mutants were obtained using a Quikchange mutagenesis kit (Stratagene). Performing this reaction on site1 and site2 yielded the sls2 mutant. Performing the reaction on site1, site2, and B10 yielded sls2B10 mutant insulin. After confirming the nucleotide sequence of the obtained modified human insulin gene, the gene was incorporated into pcDNA3.1 vector, and then used for gene transfer.

40 (2) Gene transfer into primary cultured adipocytes

[0073] After mixing 500 μ L of FCS-free DMEM medium and 15 μ L of Fugene 6 reagent (Roche), 5 μ g of transfection plasmid was added, and then this was left to stand at room temperature for 15 minutes. The mixed solution was added to primary cultured adipocytes (derived from adipose tissue around the C57BL/6 mice epididymis), which had been cultured to 70 to 80% confluency in a 10-cm dish. This was cultured for 24 hours in a CO₂ incubator. Four days after gene transfer, the cells were subcultured in a T225 flask, and cultured overnight. The medium was then exchanged for a medium comprising 0.2 mgU/mL of G418 (SIGMA), and culture was continued for three weeks, whereupon gene-transferred cells were selected. The obtained G418-resistant cells were plated onto a 10-cm dish, and the amount of insulin in the culture supernatant was measured using an ultrasensitive insulin EIA kit (Morinaga). This EIA kit detects both proinsulin, which has not yet been processed, and mature insulin.

[Results]

55 **[0074]** Fig. 7 shows (pro)insulin production by plasmid transfection into primary cultured adipocytes. Each of pcDNA3.1Myc-His vectors individually incorporating the native human insulin gene (native) and the site1/site2/B10-modified form (s1s2B10), or an empty vector (mock), was transfected into adipocytes derived from the adipose tissue surrounding the C57BL/6 mice epididymis. Human (pro)insulin was detected in the culture supernatant of resistant cells obtained by G418 selection. This showed that stable gene transfer to primary cultured adipocytes is also possible using

a plasmid vector.

[Example 5] Production, using an adeno-associated virus, of adipocytes that stably express AP

5 **[0075]** Methods of gene transfer include methods that use adeno-associated viruses (AAV).

[Methods]

10 **[0076]** The study was carried out using AAV Helper-Free System (Stratagene). The PLAP fragment of Example 2 (a fragment excised by using *Hind*III and *Bgl*II) was inserted into the same restriction enzyme site of the pAAV-MCS vector, yielding pAAV-PLAP.

15 **[0077]** AAV vector production was carried out as follows: 1.75 mL of OPTI-MEM (Invitrogen) was mixed with 220 μ L of the plasmid transfection reagent Fugene, then 25 μ g each of pAAV-PLAP, pAAV-RC, and pHelper were mixed in, and these were left to stand at room temperature for 15 minutes (Fugene/plasmid solutions). Meanwhile, 293-EBNA cells grown to 60 to 70% confluency in a 15-cm dish were prepared. The culture solution was changed to FCS-free DMEM, then Fugene/plasmid solution was instilled evenly, and this was cultured for two to three hours. FCS was then added to a final concentration of 10%, and this was cultured for two more days. The cells were collected by trypsin treatment and centrifugation, and then suspended in 50 mM Tris-HCl and 150 mM NaCl solution so that the final volume was 3 mL. Cells were disrupted by performing three cycles of dry ice-ethanol/37°C freeze-thawing on this suspension solution. Furthermore, after degrading the host genomic DNA using Benzonase (SIGMA), the virus solution was produced by centrifugation at 9,000 rpm for 30 minutes, followed by filtration of the supernatant.

20 **[0078]** Primary cultured adipocytes (derived from C57BL/6 mice subcutaneous fat) were plated onto a 12-well plate at 1×10^4 cells/well the day before gene transfer, and were cultured. They were then treated for six hours in a medium containing 40 mM of Hydroxyurea and 1 mM of butyric acid (both from SIGMA). After removing this medium, 0.5 mL/well of the virus solution, diluted to 1/100 with FCS-free DMEM, was added. After culturing for one hour, FCS-containing medium was added to a final concentration of 10%, and this was cultured overnight. Thereafter, normal medium exchanges were performed, and subculturing was performed on the 24th day.

30 **[0079]** Medium was exchanged on the first, seventh, and 25th day of transfer, and the culture supernatant collected two days after each exchange were used for the AP assays. 10 μ L of the supernatant, which was diluted as necessary, was warmed at 65°C for 20 minutes, then 50 μ L of assay buffer (16 mM NaHCO₃, 12 mM Na₂CO₃, 0.8 mM MgSO₄), and 50 μ L of luminescent stain reagent (CDP-Star Ready to Use with Sapphire II, TROPIX), were mixed, reacted in the dark for 30 minutes, and then measured with a luminometer.

[Results]

35 **[0080]** Fig. 8 shows stable expression of AP in primary cultured adipocytes (derived from C57BL/6 mice subcutaneous fat) transfected with AP-expressing AAV. AP activity was detected in the culture supernatant over the entire examination period. This showed that stable gene transfer to primary cultured adipocytes can be accomplished using an AAV vector.

40 [Example 6] Construction of a human insulin-expressing retroviral vector, and transduction thereof into adipocytes

[Methods]

45 **[0081]** The modified human insulin gene constructed in Example 4 (s1s2B10Ins) was inserted into pBabeCLX12G vector following the method of Example 3 (pBabeCL(s1s2B10Ins)I2G). This plasmid along with a VSV-G-encoding vector (pVPack-VSV-G/Stratagene), and Gag-Pol-encoding vector (modified from pVPack-gp/Stratagene) were introduced into 293-EBNA cells according to the method of Example 3, thus producing the modified insulin-expressing retroviral vector (MLV(VSV)/pBabeCL(s1s2B10Ins)I2G). The culture supernatant (approximately 200 mL) of 293-EBNA cells from twenty-two 10-cm dishes was collected, insoluble material was removed by centrifugation/filtration treatment, and then the concentrated virus solution was yielded by ultracentrifugation (19,500 rpm, 100 minutes). This was transferred to primary cultured adipocytes (derived from C57BL/6 subcutaneous fat), which had been plated onto a 6-well plate on the previous day.

50 **[0082]** The gene-transferred cells were re-plated onto a 6-well plate, and differentiation was induced according to the method of Example 1. Culture supernatants were each collected for three days, from three days before induction to the day of induction initiation (pre-induction), and for three days from the 14th to 17th day of induction (post-induction). The amount of insulin was assayed by the same method as in Example 4. Furthermore, to confirm that processing occurred at the desired sites, and that mature insulin was produced, measurements were made using insulin EIA kit (IBL), which only recognizes mature insulin. The culture supernatant of non-gene-transferred cells, which were simultaneously sub-

jected to differentiation induction, was used as a control.

[Results]

5 **[0083]** Fig. 9 shows insulin expression at the time of differentiation induction in primary cultured adipocytes transduced with s1s2B10 insulin-expressing retroviral vector. (A) shows the results of using EIA produced by Morinaga, and (B) shows the results of using EIA produced by IBL. These results show that insulin is stably secreted both before and after differentiation induction, and that transfer of mutant insulin gene may cause the production of mature insulin from adipocytes.

10

[Example 7] Construction of a retroviral vector that expresses human glucagon-like peptide-1 (GLP-1), and transduction thereof into adipocytes

15 **[0084]** GLP-1 is a peptide that is produced from small intestinal L-cells during food intake, and comprises the effect of stimulating insulin secretion by acting on pancreatic β -cells. GLP-1 is also known to have a variety of other antidiabetic and antiobesity effects such as a regeneration effect on pancreatic β -cells, an appetite-suppressing effect, and an inhibitory effect on gastric emptying (Meier, J.J. et al. Eur. J. Pharmacol. 2002, 12; 440(2-3):269-79; Drucker, D.J. Gastroenterology 2002; 122 (2) :531-544). A peptide comprising positions 7 to 37 of the amino acid sequence of GLP-1 (or up to position 36 in the amide form), is formed by tissue-specific processing of the polypeptide produced from the preproglucagon gene, and is known to comprise the main pharmacological activity (Drucker, D.J. et al. Proc. Natl. Acad. Sci. USA. 1987 May; 84(10):3434-3438; Kreymann, B. et al. Lancet. 1987, 5; 2(8571):1300-1304; Mojsov, S. et al. J. Clin. Invest. 1987 Feb; 79(2):616-619). The following examination was carried out in order to produce this factor from adipocytes.

25 [Methods]

30 **[0085]** A nucleotide sequence with a total of 156 base pairs was designed, comprising a sequence (SEQ ID NO: 10 shows the coding sequence) in which human GLP-1 (7-37) and a stop codon are linked to the signal peptide (17 amino acids) of the PLAP gene used in Example 3. Nucleotides were synthesized so that a 22mer overlap was comprised at the center (sPL-GLP-1Fw and sPL-GLP-1Rv in Table 1). These were annealed and a double strand was formed using Pfu polymerase (Stratagene). The target fragment was then amplified by PCR using 5'-end and 3'-end primers (GLP-5' and GLP-3' in Table 1). This fragment was subcloned into pCR2.1 vector, then excised using restriction enzymes, and subsequently inserted into pBabeCLX12G vector, as in Example 3 (pBabeGL(sPL-GLP1)12G). This was transfected into 293-EBNA cells by a method similar to that of Example 6, producing a GLP-1-expressing retroviral vector (MLV(VSV)/pBabeCL(sPL-GLP-1)12G). Approximately 90 mL of the culture supernatant of 293-EBNA cells from nine 10-cm dishes was collected. Insoluble material was removed by centrifugation/filtration treatment, and the supernatant was then ultracentrifuged (19,500 rpm, 100 minutes) to yield a concentrated virus solution. This was transduced into primary cultured adipocytes (derived from C57BL/6 subcutaneous fat) that had been plated onto a 6-well plate the previous day. The transfected adipocytes were again plated onto a 12-well plate, and differentiation induction was carried out according to the method of Example 1. "Non-induced" refers to a condition in which culture was continued in a normal medium instead of in an induction medium or mature medium. Seven days later, the medium was exchanged to FCS-free DMEM medium comprising 1 mM Valine-pyrrolidine (GLP-1 degradation enzyme inhibitor; synthesized at Eisai). The culture supernatant was collected 18 hours later, and the amount of active GLP-1(7-37) was measured using ELISA (LINCO).

45

[Results]

50 **[0086]** Fig. 10 shows the level of expression in primary cultured adipocytes transfected with GLP-1(7-37)-expressing retroviral vector. Expression of active form GLP-1(7-37) was observed in the culture supernatant of both non-differentiation-induced and differentiation-induced adipocytes. This showed that even when a factor is produced as the prepropeptide and then cut out by processing, this method allows production of only that factor from adipocytes.

[Example 8] Implanting mice with cells that stably express AP (Test 1)

55 [Methods]

[0087] After culturing the AP-expressing adipocytes (transduced with MLV(VSV)/pBabeCL(PLAP)IP; derived from C57BL/6 subcutaneous fat) produced by the method of Example 3 to confluency, the cells were collected by trypsin

EP 2 213 730 B1

treatment, washed with PBS, and suspended at 5×10^7 cells/mL in ice-cold Matrigel (Becton Dickinson). Implantation was performed by injecting this to the dorsal subcutaneous area (Sc) of C57BL/6 mice (eight weeks old at the time of operation; Charles River) at a dose of 0.2 mL per mouse (1×10^6 cells/head) (without differentiation induction). On the other hand, the same cells were cultured to confluency, then cultured for three days in the inducing medium of Example 1, and then implanted by similar methods (with differentiation induction). Blood was collected over time by the method indicated in Example 2, and AP activity in the plasma was measured.

[Results]

[0088] Fig. 11 shows the change in plasma AP activity in mice implanted with AP-expressing primary cultured adipocytes. Individuals that received an implant of cells subjected to differentiation-inducing stimulation for three days before implantation ("with differentiation induction") showed less fluctuation in the continued expression than individuals that received an implant of cells that were not induced. However, both methods of implantation showed continued expression over the entire 50 or so days of examination. This shows that the post-transplantation survival rate of cells may be improved by providing differentiation-inducing stimulation.

[Example 9] Implanting mice with cells that stably express AP (Test 2)

[Methods]

(1) Implantation

[0089] The AP-expressing adipocytes (transduced with MLV(VSV)/pBabeCL(PLAP)IP; derived from ICR subcutaneous fat) produced in Example 3 were cultured to confluency. The cells were collected by trypsin treatment, washed with PBS, and suspended at 5×10^7 cells/mL in an ice-cold Matrigel (Becton Dickinson) to which $1 \mu\text{g/mL}$ of bFGF (Genzyme Techne) was added. Implantation was performed by injecting this at a dose of 0.2 mL per mouse (1×10^6 cells/head) to each site (dorsal subcutaneous area (Sc), inguinal subcutaneous fat (fat), and intraperitoneal region (ip)), of the ICR nude mice (six weeks old at the time of operation, Charles River). As a control, GFP-expressing-adipocytes were treated similarly and implanted into the subcutaneous tissue.

[0090] Some of the AP-expressing cells were cultured for three days by the induction medium of Example 1, and then collected and implanted in the same manner (Dif). After using the induction medium, some of these cells were cultured for four days in a maturation medium, and then collected and implanted in the same manner (Mat).

[0091] Furthermore, some of the AP-expressing cells were plated onto an 8-well-Labteck chamber (Nunc) under the same conditions used for implantation ($1 \times 10^6/0.2$ mL bFGF-added Matrigel), and the cells were solidified by heating at 37°C . Implantation was accomplished by inserting this solidified gel into the mouse subcutaneous area. Herein, cells cultured in a normal medium after solidification were referred to as pre-fixed (pf)/gr, and cells cultured in a differentiation-inducing medium were referred to as pf/dif. Implantation was carried out after seven days of culturing.

[0092] AP activity in the plasma was measured over time, before implantation (day 0) and after implantation, according to the method of Example 2.

(2) Extirpation

[0093] In the group implanted after differentiation induction (Dif/Sc), the implanted cell masses were extirpated, along with the Matrigel, from individuals A and B, five and 43 weeks after implantation, respectively. Extirpation was performed on the control sample in the fifth week since implantation. Each individual was intraperitoneally administered with 50 mg/kg of Nembutal, as anesthesia. Their skin was then incised and a visually confirmed implanted Matrigel section was extirpated. The site of the surgery was sutured and disinfected with Iodine (Meiji). The animals were then raised in the same manner, and blood was collected over time.

[Results]

[0094] Fig. 12 (A) shows the result of examining the change in plasma AP activity over 50 days, when AP-expressing primary cultured adipocytes were implanted using basic FGF-added Matrigel, in the presence of differentiation stimulation (Dif/Sc group). Change in blood AP activity was stable for 49 days over an about 5-fold range. This showed that bFGF addition-at the time of implantation can further improve the post-implantation engraftment rate. (B) shows the disappearance of plasma AP activity due to the extirpation of the implanted Matrigel (individual A) over the same period. AP activity in the extirpated individuals was significantly decreased compared to the average value for the PLAP transduced group. This showed that blood AP is derived from the implanted cells, and that graft extirpation can quickly eliminate gene

expression. At this time, extirpation was also performed on a portion of the control group, which was implanted with GFP-transfected cells. GFP-positive cells were found in the extirpated Matrigel, and many of them displayed a vacuole image (C) similar to that shown in Fig. 6(B). This showed that primary cultured adipocytes implanted by this method may be engrafted as mature adipose tissue *in vivo*.

5 **[0095]** Fig. 13 shows the result of a long-term examination of blood AP activity in the implanted mice of Fig. 12 (A), and in mice receiving an implant by a variety of other methods. In the group implanted with PLAP-transfected cells, a clear increase of blood AP activity was confirmed for all implantation sites and implantation methods. Blood AP activity was maintained for a long period, and in particular, stable AP expression was observed for one year during the Dif/Sc group testing period (the group described in Fig. 12 (A)). Continuous AP production was also confirmed for the other
10 implantation methods, all during the examination period (316 days for the ip group, 54 days for the fat group, 225 days for the Sc group, 317 days for the Mat/Sc group, and 314 days for the two pre-fix groups). The peak of activity observed within one week of implantation was highest in the ip group. The highest values were then in the order of Sc>fat>Dif/Sc≈pf-dif>pf-gr≈Mat/Sc. The range of variation after implantation was observed as a ratio between the activity after 13 weeks and the peak activity, which can be compared in all groups. Variance was smallest, approximately three-fold, in the two
15 pre-fix groups, approximately five-fold in the ip, Dif/Sc, and Mat/Sc groups, and approximately ten-fold in the Sc and fat groups. The peak value immediately after implantation, and the range of variation after implantation differed for each implantation method. Any of these methods can thus be used according to the characteristics of the gene product used, the pathologic characteristics, and the simplicity of the technique. This showed that implantation of primary cultured adipocytes, to which genes were stably introduced *ex vivo*, can be performed by a variety of methods, and that long-term stable *in vivo* gene expression is possible after implantation.

[0096] Fig. 14 shows the result of performing an extirpation experiment, similar to that described in Fig. 12(B), in the later stage of implantation. Blood AP activity after extirpation was confirmed to quickly disappear, not only in individuals in which extirpation was performed in the early stages of implantation (individual A), but also in individuals in which extirpation was performed in the later stages of implantation (individual B). This showed that adipocytes implanted by this method are localized at the implanted site for a long period after implantation, and their extirpation, when appropriate, can eliminate the gene expression regardless of the timing.
25

[Example 10] Transplanting mice with cells that stably express AP

30 (Test 3)

[0097] The following examinations were carried out to confirm 'dose' dependence on the number of implanted cells

[Methods]

35

[0098] The AP-expressing adipocytes produced in Example 3 (transfected with MLV(VSV)/pBabeCL(PLAP)IP; derived from ICR subcutaneous fat) were cultured to confluency. The cells were cultured for three days in the induction medium indicated in Example 1, and then collected by trypsin treatment. After washing with PBS, the cells were suspended at 5×10^7 cells/mL into Matrigel. A five-fold stepwise dilution was carried out on the AP cell suspension solution using Matrigel, and 1×10^7 cells/mL and 2×10^6 cells/mL solutions were respectively prepared. bFGF was added to these
40 solutions at a final concentration of $1 \mu\text{g/mL}$, and they were then implanted to the dorsal subcutaneous area of ICR nude mice at a dose of 0.2 mL per mouse (high dose: 1×10^6 cells/head; medium dose: 2×10^5 cells/head; low dose: 4×10^4 cells/head). As a control, GFP-expressing adipocytes were similarly treated, and were implanted into the subcutaneous tissue under the same conditions as for high-dose conditions (1×10^5 cells/head).

45

[Results]

[0099] Fig. 15 shows the dependence of blood AP activity on the number of implanted cells when implanting AP-expressing adipocytes. Dose-dependent blood AP activity was observed on changing the number of implanted cells, and this was not influenced by duration. More specifically, the medium or low dose groups did not show a peak at the early stage of implantation, which was observed in the high dose group, and the range of fluctuation was narrower. This showed that *in vivo* expression level can be easily adjusted using the number of implanted cells, and that by adjusting the optimal number of cells, the post-implantation blood concentration (expression level) can be stabilized.
50

55

[Example 11] Hypoglycemic effect on diabetes model mice due to implantation of insulin-expressing adipocytes

[Methods]

- 5 **[0100]** Diabetic mice were produced by intravenously administering eight-week old male C57BL/6 mice with 10 mL/kg of 170 mg/kg streptozotocin (STZ, SIGMA). Fasting blood glucose (FBG) levels were measured individually at one and two weeks after STZ administration, and individuals with an FBG of 300 mg/dl or more were determined to have diabetes. The blood sugar level was measured by performing a perchlorate treatment immediately after collection of whole blood, and then using Glucose Test-II (WAKO).
- 10 **[0101]** The MLV(VSV)/pBabeCL(s1s2B10Ins)I2G-transfected adipocytes produced in Example 6 were subjected to differentiation induction stimulation using the same method as in Example 10, and then suspended at 5×10^7 cells/mL in Matrigel to which 1 μ g/mL of bFGF had been added. This suspension solution was implanted in the dorsal subcutaneous area of each diabetic mouse, at 0.2 mL per site, to a total of four sites (four $\times 10^6$ /head). For the control group, non-gene-transferred adipocytes were implanted by the same method. Implantation was performed 19 days after STZ treatment, and thereafter, FBG level was measured over time. Statistical analysis was carried out by comparison with the control group (unpaired t test).
- 15

[Results]

- 20 **[0102]** Fig. 16 shows the effect of implanting s1s2B10 insulin-expressing adipocytes in to STZ-induced diabetic mice. Non-gene-transferred cells were implanted as a control. The blood glucose level of the group implanted with insulin-expressing cells tended to decrease from the seventh day of implantation, and a significant hypoglycemic effect was indicated on the 13th and 21st day of implantation (A). The body weight 20 days after implantation was significantly higher in the group implanted with insulin-expressing cells than in the control group, and weight loss due to diabetes was therefore improved (B). The results of examination using AP suggest that this hypoglycemic effect will be maintained for a long period. Therefore, the foreign gene product produced from the implanted primary cultured adipocytes was shown to be able to contribute to the modification of the pathology of the recipient, indicating that this method may be able to treat diabetes.
- 25

30 Industrial Applicability

[0103] The present invention established methods of *ex vivo* transfer of a foreign gene into primary cultured adipocytes suitable for gene therapy, and established primary cultured adipocytes that stably maintain a foreign gene.

35 SEQUENCE LISTING

[0104]

- 40 <110> EISAI CO., LTD.
- <120> PRIMARY CULTURED ADIPOCYTES FOR GENE THERAPY
- <130> K2989 EP/1 S3
- 45 <150> JP 2002-177648
<151> 2002-06-18
- <150> JP 2002-237974
<151> 2002-08-19
- 50 <160> 11
- <170> PatentIn version 3.1
- 55 <210> 1
<211> 30
<212> DNA
<213> Artificial

EP 2 213 730 B1

<220>

<223> an artificially synthesized oligonucleotide for amplifying human insulin gene

<400> 1

5 cataagctta ccatggccct gtggatgcg 30

<210> 2

<211> 30

<212> DNA

10 <213> Artificial

<220>

<223> an artificially synthesized oligonucleotide for amplifying human insulin gene

<400> 2

15 cattctagac tagttgcagt agttctccag 30

<210> 3

<211> 37

20 <212> DNA

<213> Artificial

<220>

<223> an artificially synthesized oligonucleotide for mutagenesis

25

<400> 3

ctctacaca cccaggacca agcgggaggc agaggac 37

<210> 4

30 <211> 28

<212> DNA

<213> Artificial

<220>

35 <223> an artificially synthesized oligonucleotide for mutagenesis

<400> 4

ccctggaggg atcccgacag aagcgtgg 28

40 <210> 5

<211> 29

<212> DNA

<213> Artificial

45 <220>

<223> an artificially synthesized oligonucleotide for mutagenesis

<400> 5

50 cacctgtgcg gatccgacct ggtgaagc 29

<210> 6

<211> 83

<212> DNA

55 <213> Artificial

<220>

<223> an artificially synthesized oligonucleotide for constructing a gene encoding a secretory human GLP-1

EP 2 213 730 B1

atg ctg ctg ctg ctg ctg ctg ctg ggc ctg agg cta cag ctc tcc ctg 48
 Met Leu Leu Leu Leu Leu Leu Leu Leu Gly Leu Arg Leu Gln Leu Ser Leu
 1 5 10 15

5 ggc cat gct gaa ggg acc ttt acc agt gat gta agt tct tat ttg gaa 96
 Gly His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu
 20 25 30

10 ggc caa gct gcc aag gag ttc att gct tgg ctg gtg aaa ggc cga gga 144
 Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
 35 40 45

<210> 11
 <211> 48
 15 <212> PRT
 <213> Artificial

<220>
 <223> a secretory human GLP-1
 20

<400> 11

Met Leu Leu Leu Leu Leu Leu Leu Gly Leu Arg Leu Gln Leu Ser Leu
 1 5 10 15

25 Gly His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu
 20 25 30

30 Gly Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
 35 40 45

Claims

- 35 1. A pharmaceutical composition comprising a primary cultured preadipocyte, wherein the preadipocyte stably maintains a foreign gene encoding a protein that is secreted outside of the cell and wherein the gene has been inserted into a retroviral vector and has been transferred to the cell by the retroviral vector, wherein the protein is lecithin cholesterol acyltransferase (LCAT).
- 40 2. A primary cultured preadipocyte, wherein the preadipocyte stably maintains a foreign gene encoding a protein that is secreted outside of the cell, and wherein the gene has been inserted into a retroviral vector and has been transferred to the cell by the retroviral vector, for use in gene therapy, wherein the protein is LCAT.
- 45 3. The pharmaceutical composition of claim 1, or the preadipocyte for use according to claim 2, wherein the preadipocyte has the ability to significantly express the protein *in vivo* for at least 20 days.
4. The pharmaceutical composition of claim 1 or 3, or the preadipocyte for use according to claim 2 or 3, wherein the preadipocyte is used to release the protein into the blood flow.
- 50 5. An *in vitro* method of producing a preadipocyte for use in gene therapy, wherein the method comprises the steps of:
- (1) primary culturing a preadipocyte; and
 (2) transferring, and then stably holding a foreign gene encoding a protein that is secreted outside of the cell,
- 55 wherein the protein is LCAT.
6. The method of claim 5, wherein the foreign gene is transferred by a retroviral vector.

EP 2 213 730 B1

7. The preadipocyte for use according to claim 2 which has been produced by the method of claim 5 or 6.
8. An implant composition, wherein the composition comprises a primary cultured preadipocyte, which stably holds a foreign gene encoding a protein that is secreted outside of the cell, and wherein the gene has been inserted into a retroviral vector and has been transferred to the cell by the retroviral vector, and a pharmaceutically acceptable carrier, for use in gene therapy, wherein the protein is LCAT.
9. The implant composition for use according to claim 8, or the pharmaceutical composition of any one of claims 1, 3 and 4, which further comprises an extracellular matrix component.
10. The implant composition for use according to claim 8 or 9, or the pharmaceutical composition of any one of claims 1, 3, 4 and 9, which further comprises an angiogenesis factor.
11. A non-human animal, the body of which has been implanted with the primary cultured preadipocyte of claim 2.

Patentansprüche

1. Pharmazeutische Zusammensetzung, die einen primär kultivierten Präadipozyten umfasst, wobei der Präadipozyt ein Fremdgen stabil erhält, das ein Protein codiert, das aus der Zelle sezerniert wird, und wobei das Gen in einen retroviralen Vektor inseriert und durch den retroviralen Vektor in die Zelle transferiert wurde, wobei das Protein Lecithin-Cholesterin-Acyltransferase (LCAT) ist.
2. Primär kultivierter Präadipozyt zur Verwendung in der Gentherapie, wobei der Präadipozyt ein Fremdgen stabil erhält, das ein Protein codiert, das aus der Zelle sezerniert wird, und wobei das Gen in einen retroviralen Vektor inseriert und durch den retroviralen Vektor in die Zelle transferiert wurde, wobei das Protein LCAT ist.
3. Arzneimittel nach Anspruch 1 oder Präadipozyt zur Verwendung nach Anspruch 2, wobei der Präadipozyt die Fähigkeit hat, das Protein *in vivo* für mindestens 20 Tage signifikant zu exprimieren.
4. Arzneimittel nach Anspruch 1 oder 3 oder Präadipozyt zur Verwendung nach Anspruch 2 oder 3, wobei der Präadipozyt verwendet wird, um das Protein in den Blutfluss abzugeben.
5. *In vitro*-Verfahren zur Herstellung eines Präadipozyten zur Verwendung in der Gentherapie, wobei das Verfahren die Schritte umfasst:
- (1) Primärkultivierung eines Präadipozyten; und
(2) Transfer und anschließend stabiles Erhalten eines Fremdgens, das ein Protein codiert, das aus der Zelle sezerniert wird,
- wobei das Protein LCAT ist.
6. Verfahren nach Anspruch 5, wobei das Fremdgen durch einen retroviralen Vektor transferiert wird.
7. Präadipozyt zur Verwendung nach Anspruch 2, der durch das Verfahren nach Anspruch 5 oder 6 hergestellt wurde.
8. Implantatzusammensetzung zur Verwendung in der Gentherapie, wobei die Zusammensetzung einen primär kultivierten Präadipozyten umfasst, der ein Fremdgen stabil erhält, das ein Protein codiert, das aus der Zelle sezerniert wird, und wobei das Gen in einen retroviralen Vektor inseriert und durch den retroviralen Vektor in die Zelle transferiert wurde, und einen pharmazeutisch verträglichen Träger, wobei das Protein LCAT ist.
9. Implantatzusammensetzung zur Verwendung nach Anspruch 8 oder Arzneimittel nach einem der Ansprüche 1, 3 und 4, die/das des Weiteren einen extrazellulären Matrixbestandteil umfasst.
10. Implantatzusammensetzung zur Verwendung nach Anspruch 8 oder 9 oder Arzneimittel nach einem der Ansprüche 1, 3, 4 und 9, die/das des Weiteren einen Angiogenesefaktor umfasst.
11. Nicht-humanes Tier, in dessen Körper der primär kultivierte Präadipozyt nach Anspruch 2 implantiert wurde.

Revendications


- 5
1. Composition pharmaceutique comprenant un pré-adipocyte issu d'une culture primaire, dans laquelle le pré-adipocyte maintient de manière stable un gène étranger codant pour une protéine qui est sécrétée à l'extérieur de la cellule et dans laquelle le gène a été inséré dans un vecteur rétroviral et a été transféré dans la cellule par le vecteur rétroviral, la protéine étant la lécithine cholestérol acyltransférase (LCAT).
- 10
2. Pré-adipocyte issu d'une culture primaire, dans lequel le pré-adipocyte maintient de manière stable un gène étranger codant pour une protéine qui est sécrétée à l'extérieur de la cellule et dans lequel le gène a été inséré dans un vecteur rétroviral et a été transféré dans la cellule par le vecteur rétroviral, pour une utilisation en thérapie génique, la protéine étant LCAT.
- 15
3. Composition pharmaceutique selon la revendication 1, ou pré-adipocyte pour une utilisation selon la revendication 2, dans lequel le pré-adipocyte a la capacité d'exprimer de manière significative la protéine *in vivo* pendant au moins 20 jours.
- 20
4. Composition pharmaceutique selon la revendication 1 ou 3, ou pré-adipocyte pour une utilisation selon la revendication 2 ou 3, dans lequel le pré-adipocyte est utilisé pour libérer la protéine dans la circulation sanguine.
- 25
5. Méthode *in vitro* de production d'un pré-adipocyte pour une utilisation en thérapie génique, dans laquelle la méthode comprend les étapes consistant à :
- (1) mettre un pré-adipocyte en culture primaire ; et
- (2) transférer, et puis maintenir de manière stable un gène étranger codant pour une protéine qui est sécrétée à l'extérieur de la cellule,
- la protéine étant LCAT.
- 30
6. Méthode selon la revendication 5, dans laquelle le gène étranger est transféré par un vecteur rétroviral.
- 35
7. Pré-adipocyte pour une utilisation selon la revendication 2 qui a été produit par la méthode selon la revendication 5 ou 6.
- 40
8. Composition d'implant, dans laquelle la composition comprend un pré-adipocyte issu d'une culture primaire, qui maintient de manière stable un gène étranger codant pour une protéine qui est sécrétée à l'extérieur de la cellule, et dans laquelle le gène a été inséré dans un vecteur rétroviral et a été transféré dans la cellule par le vecteur rétroviral, et un véhicule pharmaceutiquement acceptable, pour une utilisation en thérapie génique, la protéine étant LCAT.
- 45
9. Composition d'implant pour une utilisation selon la revendication 8, ou composition pharmaceutique selon l'une quelconque des revendications 1, 3 et 4, qui comprend en outre un composant de matrice extracellulaire.
- 50
10. Composition d'implant pour une utilisation selon la revendication 8 ou 9, ou composition pharmaceutique selon l'une quelconque des revendications 1, 3, 4 et 9, qui comprend en outre un facteur d'angiogenèse.
- 55
11. Animal non-humain, dans le corps duquel a été implanté un pré-adipocyte issu d'une culture primaire selon la revendication 2.

FIG. 1

(A)



(B)



(C)



(D)




FIG. 2




FIG. 3





FIG. 4

(A)



(B)




FIG. 5





FIG. 6





FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11






FIG. 12

(A)



(B)



(C)




FIG. 13




FIG. 14




FIG. 15






FIG. 16



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 0031267 A1 [0004]
- US 4829000 A [0042]
- WO 8701728 A [0043]
- WO 8904832 A [0043]
- WO 8607595 A [0043]
- WO 8703885 A [0043]
- EP 237966 A [0043]
- EP 281822 A [0043]
- EP 326907 A [0043]
- EP 394951 A [0043]
- EP 493737 A [0043]
- WO 9749827 A [0043]
- JP 2002177648 A [0104]
- JP 2002237974 A [0104]

Non-patent literature cited in the description

- TOYOOKA et al. *Folia Pharmacol. Jpn.*, 2000, vol. 116, 158-162 [0002]
- TANI et al. *Saishin Igaku*, 2001, vol. 56, 258-267 [0003]
- RAPER, S.E. et al. *Cell Transplant*, 1993, vol. 2 (5), 381-400 [0003]
- BRADLEY R.D. et al. *Recent Prog. Horm. Res.*, 2001, vol. 56, 329-358 [0005]
- MICK, G.J. et al. *Endocrinology*, 2002, vol. 143 (3), 948-53 [0005]
- *J. Gene. Med.*, January 2001, vol. 3 (1), 21-31 [0006]
- *Histochem. Cell Biol.*, January 2001, vol. 115 (1), 73-82 [0006]
- SUGIHARA et al. *Nippon Rinsho*, 1995, vol. 53, 115-120 [0017]
- SUGIHARA, H. et al. *J. Lipid Res.*, 1987, vol. 28, 1038-1045 [0017]
- ZHANGH.H. et al. *J. Endocrinol.*, 2000, vol. 164, 119-128 [0017]
- ABUMRAD N.A. et al. *J. Biol. Chem.*, 25 August 1993, vol. 268 (24), 17665-8 [0017]
- HAUNER H. et al. *J. Clin. Invest.*, 1989, vol. 84, 1663-1670 [0017]
- MARKO et al. *Endocrinology*, 1994, vol. 136, 4582-4588 [0017]
- LUND, A.H. et al. *J. Biomed. Sci.*, 1996, vol. 3, 365-378 [0025]
- NIWA, O. et al. *Cell*, 1983, vol. 32, 1105-1113 [0025]
- JAHNER, D. ; JAENISCH, R. *Nature*, 1985, vol. 315, 594-597 [0025]
- CHALLITA, P.-M. ; KOHN, D.B. *Proc. Natl. Acad. Sci. USA*, 1994, vol. 91, 2567-2571 [0025]
- HOEBEN, R.C. et al. *J. Virol.*, 1991, vol. 65, 904-912 [0025]
- CHEN, W.Y. et al. *Proc. Natl. Acad. Sci. USA*, 2000, vol. 97, 377-382 [0025]
- CHEN, W. Y. et al. *Proc. Natl. Acad. Sci. USA*, 1997, vol. 94, 5798-5803 [0025]
- SUGIHARA, H. et al. *Differentiation*, 1986, vol. 31, 42-49 [0029]
- CHEN, H. et al. *J. Biol. Chem.*, 1997, vol. 272 (12), 8026-31 [0032]
- KAY, M.A. et al. *Nat. Med.*, 2001, vol. 7, 33-40 [0032]
- ARAI, T. et al. *J. Virol.*, 1998, vol. 72, 1115-21 [0032]
- *Mol. Cell Biol.*, June 2001, vol. 21 (12), 3926-34 [0033]
- RUSSELL, D.W. ; MILLER, A.D. *J. Virol.*, 1996, vol. 70, 217-222 [0034]
- WU, M. et al. *J. Virol.*, 1999, vol. 73, 4498-4501 [0034]
- NALDINI, L. et al. *Science*, 1996, vol. 272, 263-267 [0034]
- POESCHLA, E et al. *Proc. Natl. Acad. Sci. USA*, 1996, vol. 93, 11395-11399 [0034]
- SRINIVASAKUMAR, N. et al. *J. Virol.*, 1997, vol. 71, 5841-5848 [0034]
- ZUFFEREY, R. et al. *Nat. Biotechnol.*, 1997, vol. 15, 871-875 [0034]
- KIM, V.N. et al. *J. Virol.*, 1998, vol. 72, 811-816 [0034]
- JOHNSTON, J.C. et al. *J. Virol.*, 1999, vol. 73, 4991-5000 [0034]
- JOHNSTON, J. ; POWER, C. *J. Virol.*, 1999, vol. 73, 2491-2498 [0034]
- POESCHLA, E.M. et al. *Nat. Med.*, 1998, vol. 4, 354-357 [0034]
- T. M. SHINNICK ; R. A. LERNER ; J. G. SUTCLIFFE. *Nature*, 1981, vol. 293, 543-548 [0034]
- YU S.F. et al. *Proc. Natl. Acad. Sci. USA*, 1986, vol. 83, 3194 [0035]
- YEE, J. K. et al. *Proc. Natl. Acad. Sci. USA*, 1987, vol. 84, 5197-5201 [0035]
- ZUFFEREY, R. et al. *J. Virology*, 1998, vol. 72, 9873-9880 [0035]
- SORGE, J. et al. *Mol. Cell. Biol.*, 1984, vol. 4 (9), 1730-1737 [0037]

EP 2 213 730 B1

- **EMI, T. FRIEDMANN ; J. K. YEE.** *J. Virol.*, 1991, vol. 65 (3), 1202-1207 [0037]
- **YEE, J.-K. et al.** *Methods Cell Biol.*, 1994, vol. 43 (43), 99-112 [0037]
- **BURNS, J. C. et al.** *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90 (90), 8033-8037 [0037]
- **ROSE, J.K. ; GALLIONE, C.J.** *J. Virol.*, 1981, vol. 39 (2), 519-528 [0037]
- **ARAI T. et al.** *J. Virol.*, 1998, vol. 72, 1115-21 [0037]
- *Current Protocols in Molecular Biology.* John Wiley & Sons, 1995 [0038]
- **BYUN, J. et al.** *Gene Ther.*, 1996, vol. 33333, 1018-1020 [0038]
- **TAFURO, S. et al.** *Gene Ther.*, 1996, vol. 33333, 679-684 [0038]
- **MIYAO, Y. et al.** *Cell Struct. Funct.*, 1995, vol. 20 (20), 177-183 [0038]
- **CLAUDIO, P. P. et al.** *Anal. Biochem.*, 2001, vol. 291, 96-101 [0038]
- **CASHION, L. M. et al.** *Biotechniques*, 1999, vol. 26 (26), 924-930 [0038]
- **ABRAHAM et al.** *EMBO J.*, 1986, vol. 5, 2523-2528 [0043]
- **PRATS et al.** *Proc. Natl. Acad. Sci. USA*, 1989, vol. 86, 1836-1840 [0043]
- **GOTO, M. et al.** *Mol. Pharmacol.*, 1996, vol. 49, 860-873 [0052] [0058]
- **MORGENSTERN, J.P et al.** *Nucleic Acids Res.*, 1990, vol. 18, 3587-3596 [0059]
- *JBC*, 1994, vol. 269 (8), 6241 [0072]
- **MEIER, J.J. et al.** *Eur. J. Pharmacol.*, 2002, vol. 440 (2-3), 269-79 [0084]
- **DRUCKER, D.J.** *Gastroenterology*, 2002, vol. 122 (2), 531-544 [0084]
- **DRUCKER, D.J. et al.** *Proc. Natl. Acad. Sci. USA.*, 1987, vol. 84 (10), 3434-3438 [0084]
- **KREYMANN, B. et al.** *Lancet*, 1987, vol. 2 (8571), 1300-1304 [0084]
- **MOJSOV, S. et al.** *J. Clin. Invest.*, February 1987, vol. 79 (2), 616-619 [0084]

PRIMER, TENYÉSZTETT ADIPOCITÁK GÉNTERÁPIÁRA

Szabadalmi igénypontok

1. Gyógyászati készítmény, amely primer, tenyésztett pre-adipocitát tartalmaz, ahol a pre-adipocita stabilan tartalmaz a sejten kívülre szekretálódó fehérjét kódoló, idegen gént, és ahol a gén retrovirális vektorba inszertált és a retrovirális vektorral a sejthez juttatott, és ahol a fehérje lecitin-koleszterin-aciltranszferáz (LCAT).

2. Primer, tenyésztett pre-adipocita, ahol a pre-adipocita stabilan tartalmaz a sejten kívülre szekretálódó fehérjét kódoló, idegen gént, és ahol a gén retrovirális vektorba inszertált és a retrovirális vektorral a sejthez juttatott, génterápiában történő alkalmazásra, ahol a fehérje LCAT.

3. Az 1. igénypont szerinti gyógyászati készítmény vagy a 2. igénypont szerinti pre-adipocita az ott meghatározott alkalmazásra, ahol a pre-adipocita képes szignifikáns módon expresszálni a fehérjét *in vivo*, legalább 20 napon keresztül.

4. Az 1. vagy 3. igénypont szerinti gyógyászati készítmény vagy a 2. vagy 3. igénypont szerinti pre-adipocita az ott meghatározott alkalmazásra, ahol a pre-adipocita a fehérjét a véráramba szabadítja fel.

5. *In vitro* eljárás génterápiában történő alkalmazásra szolgáló pre-adipocita előállítására, amely magában foglalja a következő lépéseket:

(1) pre-adipocita primer tenyésztése és

(2) a sejten kívülre szekretált fehérjét kódoló idegen gén transzfere és stabil megtartása, ahol a fehérje LCAT.

6. Az 5. igénypont szerinti eljárás, ahol az idegen gén retrovirális vektorral transzferált.

7. A 2. igénypont szerinti pre-adipocita az ott meghatározott alkalmazásra, amely az 5. vagy 6. igénypont szerinti eljárással előállított.

8. Implantátum-készítmény, amely tartalmaz primer, tenyésztett pre-adipocitát, ahol a pre-adipocita stabilan tartalmaz a sejten kívülre szekretálódó fehérjét kódoló, idegen gént, és ahol a gén retrovirális vektorba inszertált és a retrovirális vektorral a sejthez juttatott, és gyógyászatiilag elfogadható hordozót, génterápiában történő alkalmazásra, ahol a fehérje LCAT.

9. A 8. igénypont szerinti implantátum-készítmény az ott meghatározott alkalmazásra vagy az 1., 3. vagy 4. igénypontok bármelyike szerinti gyógyászati készítmény, amely extracellulárismátrix-komponenst is tartalmaz.

10. A 8. vagy 9. igénypont szerinti implantátum-készítmény az ott meghatározott alkalmazásra vagy az 1., 3., 4. vagy 9. igénypontok bármelyike szerinti gyógyászati készítmény, amely angiogenezis faktort is tartalmaz.

11. Embertől különböző állat, amelynek testébe a 2. igénypont szerinti primer, tenyésztett pre-adipocita van beültetve.