

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 September 2010 (30.09.2010)

PCT

(10) International Publication Number
WO 2010/111273 A1

(51) International Patent Classification:

A61K 39/085 (2006.01) **C07K 14/00** (2006.01)
A61K 39/38 (2006.01)

(21) International Application Number:

PCT/US2010/028326

(22) International Filing Date:

23 March 2010 (23.03.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/210,772 23 March 2009 (23.03.2009) US

(71) Applicant (for all designated States except US): **EPI-TOPIX, LLC** [US/US]; 3735 County Road 5, Willmar, MN 56201 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HERRON-OLSON, Lisa, L.** [US/US]; 5001 Columbus Avenue, Minneapolis, MN 55417 (US). **CATRON, Drew, M.** [US/US]; 535 Elaine Avenue, Shoreview, MN 55126 (US).

(74) Agent: **GRAM, Christopher, D.**; Muetting, Raach & Gebhardt, P.A., P.O. Box 581336, Minneapolis, MN 55458-1336 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: POLYPEPTIDES AND IMMUNIZING COMPOSITIONS CONTAINING GRAM POSITIVE POLYPEPTIDES AND METHODS OF USE

(57) Abstract: The present invention provides isolated polypeptides isolatable from a *Staphylococcus* spp. Also provided by the present invention are compositions that include one or more of the polypeptides, and methods for making and methods for using the polypeptides.



WO 2010/111273 A1

POLYPEPTIDES AND IMMUNIZING COMPOSITIONS CONTAINING
GRAM POSITIVE POLYPEPTIDES AND METHODS OF USE

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority to U.S. Provisional Patent Application Serial
No. 61/210,772, filed March 23, 2009.

BACKGROUND

Gram-positive bacteria are a remarkably diverse group of organisms that
cause a variety of diseases in both humans and animals. Some of the pathogens
recognized as important in human and/or animal health include bacteria belonging to
the families of Corynebacteriaceae, Enterococcaceae, Micrococcaceae,
Mycobacteriaceae, Nocardaceae, and Peptococcaceae, which include such bacterial
species as *Actinomyces* spp., *Bifidobacterium* spp., *Corynebacterium* spp.,
Enterococcus spp., *Erysipelothrix* spp., *Eubacterium* spp., *Kytococcus* spp.,
Lactobacillus spp., *Micrococcus* spp., *Mobiluncus* spp., *Mycobacteria* spp.,
Peptostreptococcus spp., *Propionibacterium* spp., and *Staphylococcus* spp. These
pathogens cause a multitude of clinical manifestations in many different animal
species. The treatment for such infections has historically been antibiotics that attack
the common structures and functions of gram-positive organisms. However, many of
the more ubiquitous gram-positive organisms have developed resistance to several
classes of antibiotics, making treatment of infections difficult. The widespread use of
antibiotics in the treatment of bacterial diseases in both humans and food production
animals is likely a major contributing factor in the proliferation of antibiotic-resistant
strains of many species of gram-positive organisms. Therefore, there is a great need
to find different treatments that prevent or eliminate infections by gram-positive
organisms in animals as well as humans.

Staphylococcal infections in agricultural animals

In the agricultural industry a number of important diseases are caused by gram-positive organisms. Examples of clinical conditions caused by gram positive bacterial infections include, mastitis, septicemia, pneumonia, osteomyelitis, meningoencephalitis, lymphangitis, dermatitis, genital tract infections, metritis, perinatal disease, pituitary abscesses, arthritis, bursitis, orchitis, cystitis and pyelonephritis, caseous lymphadenitis, tuberculosis, ulcerative lymphangitis, erysipelas, laminitis, tyzzer's disease, tetanus, botulism, enteritis, malignant edema, braxy, bacillary hemoglobinuria, enterotoxemia. *Staphylococcus* spp., in particular, are capable of infecting many different species of agricultural animals and can cause enormous economic losses. For example, the United States dairy industry is estimated to lose approximately \$185 per cow annually due to mastitis, a disease often caused by *Staphylococcus aureus*. Since there are 9.5 million head of milking cows in the U.S., the annual cost of mastitis is approximately \$1.8 billion. This is approximately 10% of the total value of farm milk sales, and about two-thirds of this loss is due to reduced milk production in sub-clinically infected cows. Other losses are due to discarded abnormal milk and milk withheld from cows treated with antibiotic, costs of early replacement of affected cows, reduced sale value of culled cows, costs of drugs and veterinary services, and increased labor costs. In addition to its prevalence within the bovine dairy industry, mastitis caused by gram-positive cocci is also common among goats and sheep. Additional animal diseases caused by *S. aureus* include botryomycosis in horses, purulent synovitis and osteomyelitis in poultry, snuffles in rabbits, abortions in swine, and tick pyemia in lambs. Other species of staphylococci are major skin pathogens of canine (*S. intermedius*) and swine (*S. hyicus*). In poultry species, staphylococcal pathogens cause endocarditis and septicemia.

Staphylococcal infections in humans

Staphylococcus spp. are also human pathogens causing a wide variety of infections. The species *Staphylococcus aureus*, a common colonizer of human mucosa and skin, is an opportunistic pathogen that can cause diverse human infections. For example, *S. aureus* is the causative agent of several skin infections, including impetigo, furunculosis, cellulitis, and scalded skin syndrome, as well as

potentially fatal post-surgical wound infections. In addition, the exposure of immunocompromised individuals to *S. aureus* in hospital settings has resulted in organ infections such as pneumonia, urinary tract infections, osteomyelitis, arthritis, bacteremia, and endocarditis. *S. aureus* is also the causative agent of toxinoses, most notably toxic shock syndrome and food poisoning. Food poisoning caused by the staphylococcal enterotoxin B is the most common cause of food-borne illness, surpassing even salmonellosis, campylobacteriosis and listeriosis. Other species of staphylococci also cause human disease; *S. epidermidis*, *S. haemolyticus* and *S. hominis* commonly infect implanted medical devices and *S. saprophyticus* is associated with urinary tract infections in women.

Virulence mechanisms of staphylococci

Staphylococci infect a variety of host tissues and evade the immune system through the production of several types of secreted proteins, surface expressed virulence factors and metabolic systems designed for survival amidst the limited resources and active defenses associated with the host environment. Colonization is the necessary first step in establishing infection; numerous factors including capsule, lipoteichoic acid, and teichoic acid are common structural components contributing to colonization. In addition, surface proteins such as staphylococcal fibronectin-binding protein and bone-sialoprotein binding proteins specifically bind host tissue components. Toxins are commonly produced among staphylococcal pathogens and are highly damaging; several human diseases, including food poisoning, toxic shock syndrome and exfoliative skin conditions, are the direct result of extracellular secreted toxin proteins. A single isolate may encode genes for 20-30 different secreted toxins. Some of the secreted protein products are superantigens that can bind nonspecifically to the MHC class II molecule of an antigen-presenting cell and, simultaneously, to the T-cell receptor of a T cell. The binding induces T cell signaling and leads to the release of high levels of proinflammatory factors, ultimately inducing host damage due to the overwhelming immune response. Another class of virulence factors expressed on the surface disguise the bacteria from the host immune system. For example, the *S. aureus* surface-expressed Protein A inhibits opsonization and phagocytosis by binding of the Fc component of host antibody. Numerous proteases, hemolysins (alpha, beta, gamma and delta),

nucleases, lipases, hyaluronidase, and collagenase also aid bacteria in extracting nutrients from surrounding cells and protecting them against host defenses.

Antibiotic resistance among staphylococci

5 The CDC estimates that each year nearly 2 million people in the United States acquire a nosocomial infection, resulting in 90,000 deaths annually. Of these fatal infections, 70% are caused by antibiotic-resistant bacteria. The increase in antibiotic-resistance among microbial species is particularly pronounced in skin and mucosal colonizers such as *S. aureus*. For example, the vast majority of *S. aureus* isolated
10 from hospital settings are resistant to penicillin, and 50% are also resistant to the semisynthetic penicillins, such as methicillin, nafcillin, and oxacillin. These isolates, referred to as MRSA (methicillin resistant *S. aureus*) were first seen in the 1970s, and are now firmly established in hospital settings. Recently there have been several cases of MRSA infections in the community, where the infected individuals had no
15 previous exposure to hospitals or healthcare workers. This alarming trend is intensified by the isolation of MRSA isolates that are less susceptible to vancomycin, a glycopeptide used to treat MRSA. Very few strains have been shown to be truly resistant to vancomycin according to the CDC's definition of vancomycin resistance, but several MRSA strains have been characterized as consisting of subpopulations
20 with reduced susceptibility to vancomycin, or VISA (vancomycin intermediate *S. aureus*). Since the isolation of vancomycin resistant and vancomycin intermediate strains is a relatively new development, there is little data concerning their prevalence in hospitals and/or the community. Occasionally, VRSA (vancomycin resistant *S. aureus*) with full resistance to vancomycin and carrying a resistance plasmid likely
25 acquired from *Enterococcus spp.* have also been recovered from humans.

Strategies for the prevention and treatment of *Staphylococcus* infections

 The emergence of numerous gram-positive pathogens that are resistant to multiple antibiotics has fueled research efforts aimed at developing preventative
30 vaccines to protect against disease. Vaccines are designed to be administered to patients in order to elicit a long-term memory response from the immune system, so that if the pathogen is encountered at a future time, the immune system can more quickly and efficiently clear the pathogen. To date, a broadly-protective vaccine

against gram-positive pathogens associated with a number of severe human diseases, particularly those disease associated with staphylococcal infections, is not available. Vaccine development approaches for the prevention of staphylococcal infections include those reporting the use of microbial surface components recognizing

5 adhesion matrix molecules [MSCRAMMS (Nilsson et al. 1998. J Clin Invest 101:2640-9; Menzies et al. 2002. J Infect Dis 185:937-43; Fattom et al. 2004. Vaccine 22:880-7], surface polysaccharides (McKenney et al. 2000; McKenzie et al. 1999. Science 284:1523-7; Maira-Litran et al. 2002. Infect Immun 70:4433-40; Maira-Litran et al. 2004. Vaccine 22:872-9; Maira-Litran et al. 2005. Infect Immun

10 73:6752-62) and mutated exoproteins (Lowell et al. 1996. Infect Immun 64:4686-93; Stiles et al. 2001. Infect Immun 69:2031-6; Gampfer et al. 2002. Vaccine 20:3675-84), as antigens in subunit vaccine compositions, as well as one live avirulent strain (Reinoso et al. 2002. Can J Vet Res 66:285-8) and several DNA vaccine approaches (Ohwada et al. 1999. J Antimicrob Chemother 44:767-74); Brouillette et al. 2002.

15 Vaccine 20:2348-57; Senna et al. 2003. Vaccine 21:2661-6). Although many of these compositions have shown some degree of protection, they have achieved little cross-protection against diverse staphylococcal strains and have additionally failed to elicit substantial immune responses in immunocompromised patients, an important at-risk population for nosocomial infections.

20 The most severe staphylococcal diseases are those mediated by the aforementioned supernantigenic pyrogenic exotoxins (SPEs) that nonspecifically stimulate T-cells independent of antigen presentation. Such diseases include toxic shock syndrome, exfoliative skin disease, and possibly Kawasaki syndrome. For these SPE-mediated diseases, immunotherapeutic agents that boost the immune

25 system during an active infection are often more effective than vaccines, which are typically administered prior to infection. The overwhelming nature of the immune response to SPE necessitates rapid reduction in toxin activity as the first objective in therapy. To date, toxin neutralization in *S. aureus*-mediated disease has been most effectively accomplished by the administration of intravenous human

30 immunoglobulin (IVIG), a purified, concentrated human antibody preparation from several thousand human donors (Takei et al. 1993. J Clin Invest 91:602-7; Stohl and Elliot. 1996. Clin Immunol Immunopathol 79:122-33). The widespread distribution of *S. aureus*, which colonizes approximately 30% of healthy human adults, coincides

with high exposure rates for the majority of the population, so the level of anti-staphylococcal anti-toxin antibodies in IVIG is often sufficient to neutralize toxin long enough to stabilize the immune response until the bacterial load is reduced with antibiotics (Schlievert, 2001. J Allergy Clin Immunol 108(4 Suppl):S107-110).

5 IVIG preparations from multiple manufacturers have been shown to neutralize toxin in proliferation assays with human peripheral blood mononuclear cells, inhibit toxin-induced human T cell-driven B cell differentiation *in vitro* (Stohl and Elliot. 1996. Clin Immunol Immunopathol 79:122-33; Stohl and Elliott. 1995. J Immunol 155:1838-50; Stohl et al. 1994. J Immunol 153:117-27) and reduce IL-4 and IL-2

10 secretion in PBMCs stimulated with staphylococcal enterotoxin B (Takei et al. 1993. J Clin Invest 91:602-7; Darenberg et al. 2004. Clin Infect Dis 38:836-42). IVIG therapy, with its proven ability to neutralize SPE, is now a recommended therapy for Kawasaki syndrome and is gaining favor as a treatment method for staphylococcal toxic shock syndrome (Schlievert 2001. J Allergy Clin Immunol 108(4 Suppl):S107-

15 110). Use of IVIG as an immunoprotective wound lavage during surgery has also been investigated in mice (Poelstra et al. 2000. Tissue Eng 6(4):401-411). Although standard IVIG has utility for limiting the advance of some staphylococcal SPE-mediated disease, the safety, efficacy and consistency of human IVIG preparations generated from thousands of unselected human donors remains controversial (Baker

20 et al. 1992. N Engl J Med 327:213-9; Miller et al. 2001. J Allergy Clin Immunol 108:S91-4; Sacher, 2001. J Allergy Clin Immunol 108:S139-46; Darenberg et al. 2004. Clin Infect Dis 38:836-42). Furthermore, the benefit of IVIG in *preventing* some staphylococcal infections is doubtful (Baker et al. 1992. N Engl J Med 327:213-9; Hill, H. R. 2000. J Pediatr 137:595-7; Darenberg et al. 2004. Clin Infect

25 Dis 38:836-42). In order to increase the effectiveness of IVIG in treating staphylococcal infections in certain at-risk populations, a plasma-derived, donor-selected, polyclonal anti-staphylococcal human IgG with high titers of antibody directed toward the staphylococcal MSCRAMMS clumping factor A (ClfA) and fibrinogen-binding protein G (SdrG) was created and tested with success in very low

30 birthweight infants to prevent staphylococcal sepsis (Vernachio et al. 2003. Antimicrob Agents Chemother 47:3400-6; Bloom et al. 2005. Pediatr Infect Dis J 24:858-866; Capparelli et al. 2005. Antimicrob Agents Chemother 49:4121-7). A specific humanized monoclonal antibody toward the *S. aureus* MSCRAMM

Clumping factor A, is also being developed. The antibody was selected from a pool of thousands of murine anti-ClfA antibodies for its ability to bind ClfA in a manner that abrogates *S. aureus* binding to human fibronectin and was subsequently humanized by mutating specific targeted residues to mimic the homologous human germline subgroup antibody (Hall et al. 2003. Infect Immun 71:6864-70; Domanski et al. 2005. Infect Immun 73:5229-32). The specific antibody is being designed for use in conjunction with antibiotics for the treatment of severe life-threatening *S. aureus* infection, although animal studies also demonstrated a prophylactic protective effect.

SUMMARY

In one aspect, the present invention provides compositions including two or more isolated polypeptides. An isolated polypeptide in the composition have a molecular weight as determined by electrophoresis on a sodium dodecyl sulfate-polyacrylamide gel of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa. For instance, a composition may include isolated proteins of 88 kDa and 55 kDa. In some aspects the composition may include isolated polypeptides having molecular weights of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, and 33 kDa. The polypeptides are isolatable from a *Staphylococcus aureus* when incubated in media including an iron chelator and not isolatable when grown in the media without the iron chelator. The composition protects an animal, such as a mouse or cow or human, against challenge with an *S. aureus* strain such as, for instance, ATCC strain 19636. The composition may further include a pharmaceutically acceptable carrier, and may further include one or more isolated polypeptides having a molecular weight of 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, or 40 kDa and isolatable from a *S. aureus* when grown in the media without the iron chelator. In some aspects the polypeptides of the composition may be isolated from *S. aureus* ATCC strain 19636.

In some embodiments, each polypeptide of the composition has a mass fingerprint of at least 80% similarity to a mass fingerprint of a polypeptide of the same molecular weight polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, wherein the polypeptide is isolatable from a *Staphylococcus aureus*

when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator. For instance, the isolated polypeptide with a molecular weight of 88 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 88 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and the isolated polypeptide with a molecular weight of 55 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 55 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636.

In another aspect, the present invention provides compositions that include an isolated polypeptide having at least 80% sequence similarity to an amino acid sequence selected from SEQ ID NO:408 and SEQ ID NO:397. The composition may further include at least one second polypeptide, wherein the second polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator. In some cases, the second polypeptide can include an amino acid sequence having at least 80% similarity to an amino acid sequence selected from SEQ ID NO:353, SEQ ID NO:364, SEQ ID NO:375, SEQ ID NO:386, and SEQ ID NO:419. In other cases, the second polypeptide can have a molecular weight as determined by electrophoresis on a sodium dodecyl sulfate-polyacrylamide gel of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa. The composition may further include one or more isolated polypeptides isolatable from a *S. aureus* when grown in media without the iron chelator and has a molecular weight of 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, 40 kDa.

The present invention also provides methods for using the compositions. In one aspect the method is for treating in infection in a subject, and includes administering an effective amount of a composition of the present invention to a subject having or at risk of having an infection caused by a *Staphylococcus* spp. In another aspect, the method is for treating a symptom in a subject, and it includes administering an effective amount of a composition of the present invention to a subject having an infection caused by a *Staphylococcus* spp. The subject may be a mammal, such as a human, horse, or cow. The *Staphylococcus* spp. may be *S. aureus*.

The present invention further provides methods for using antibody, for instance, polyclonal antibody, that specifically binds polypeptides of the present

invention. In one aspect, the method is for treating an infection in a subject, and includes administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp., wherein the composition includes antibody that specifically binds at least one, and in some cases more than one, isolated polypeptide of the present invention. In another aspect, the method is for treating a symptom in a subject, and includes administering an effective amount of a composition to a subject having an infection caused by a *Staphylococcus* spp., wherein the composition includes antibody that specifically binds at least one, and in some cases more than one, isolated polypeptide of the present invention. The subject may be a mammal, such as a human, horse, or cow. The *Staphylococcus* spp. may be *S. aureus*.

Also provided by the present invention are methods for decreasing colonization in a subject. In one aspect, the method includes administering an effective amount of a composition of the present invention to a subject colonized by a *Staphylococcus* spp. In another aspect, the method includes administering an effective amount of a composition to a subject colonized by *Staphylococcus* spp., wherein the composition includes antibody that specifically binds at least one, and in some cases more than one, isolated polypeptide of the present invention.

The present invention provides a kit for detecting antibody that specifically binds a polypeptide. The kit includes, in separate containers, an isolated polypeptide of the present invention, and a reagent that detects an antibody that specifically binds the polypeptide.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. The electrophoretic profile of the proteins of different strains *Staphylococcus aureus* derived from different species grown with and without iron (lanes marked Fe⁺⁺ and DP, respectively).

Figure 2. The difference in mortality between vaccinated and non-vaccinated mice after homologous and heterologous challenge with *Staphylococcus aureus*.

Figure 3. Kaplan-Meier survival curve showing percent survival after vaccination and homologous challenge with *S. aureus* ATCC 19636.

Figure 4. Kaplan-Meier survival curve showing percent survival after vaccination and heterologous challenge with *S. aureus* ATCC 19636.

Figure 5. Kaplan-Meier survival curve showing percent survival after passive immunization and homologous challenge with *S. aureus* ATCC 19636.

5 Figure 6. Kaplan-Meier survival curve showing percent survival after passive immunization and heterologous challenge with *S. aureus* strain 1477.

Figure 7. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* ATCC 19636. (SEQ ID NO:353).

Figure 8. The amino acid sequence of a metal-regulated polypeptide obtained
10 from *S. aureus* RF122. (SEQ ID NO:354).

Figure 9. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:355).

Figure 10. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:356).

15 Figure 11. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:357).

Figure 12. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:358).

Figure 13. The amino acid sequence of a metal-regulated polypeptide
20 obtained from *S. aureus* JH9. (SEQ ID NO:359).

Figure 14. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:360).

Figure 15. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:361).

25 Figure 16. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:362).

Figure 17. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:363).

Figure 18. The amino acid sequence of a metal-regulated polypeptide
30 obtained from *S. aureus* ATCC19636. (SEQ ID NO:364).

Figure 19. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:365).

Figure 20. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:366).

Figure 21. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:367).

5 Figure 22. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:368).

Figure 23. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:369).

10 Figure 24. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:370).

Figure 25. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:371).

Figure 26. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:372).

15 Figure 27. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:373).

Figure 28. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:374).

20 Figure 29. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:375).

Figure 30. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:376).

Figure 31. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:377).

25 Figure 32. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:378).

Figure 33. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:379).

30 Figure 34. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:380).

Figure 35. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:381).

Figure 36. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:382).

Figure 37. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:383).

5 Figure 38. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:384).

Figure 39. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:385).

10 Figure 40. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:386).

Figure 41. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:387).

Figure 42. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:388).

15 Figure 43. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:389).

Figure 44. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:390).

20 Figure 45. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:391).

Figure 46. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:392).

Figure 47. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:393).

25 Figure 48. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:394).

Figure 49. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:395).

30 Figure 50. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:396).

Figure 51. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:397).

Figure 52. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:398).

Figure 53. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:399).

5 Figure 54. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:400).

Figure 55. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:401).

10 Figure 56. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:402).

Figure 57. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:403).

Figure 58. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:404).

15 Figure 59. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:405).

Figure 60. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:406).

20 Figure 61. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:407).

Figure 62. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:408).

Figure 63. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:409).

25 Figure 64. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:410).

Figure 65. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:411).

30 Figure 66. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:412).

Figure 67. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:413).

Figure 68. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:414).

Figure 69. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:415).

5 Figure 70. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:416).

Figure 71. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:417).

10 Figure 72. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:418).

Figure 73. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:419).

Figure 74. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:420).

15 Figure 75. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:421).

Figure 76. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:422).

20 Figure 77. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:423).

Figure 78. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:424).

Figure 79. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:425).

25 Figure 80. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:426).

Figure 81. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:427).

30 Figure 82. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:428).

Figure 83. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:429).

Figure 84. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC 19636. (SEQ ID NO:430).

Figure 85. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:431).

5 Figure 86. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:432).

Figure 87. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:433).

10 Figure 88. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:434).

Figure 89. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:435).

Figure 90. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:436).

15 Figure 91. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:437).

Figure 92. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:438).

20 Figure 93. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:439).

Figure 94. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:440).

Figure 95. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:441).

25 Figure 96. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:442).

Figure 97. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:443).

30 Figure 98. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:444).

Figure 99. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:445).

Figure 100. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:446).

Figure 101. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:447).

5 Figure 102. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:448).

Figure 103. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:449).

10 Figure 104. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:450).

Figure 105. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:451).

Figure 106. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:452).

15 Figure 107. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:453).

Figure 108. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:454).

20 Figure 109. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:455).

Figure 110. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:456).

Figure 111. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:457).

25 Figure 112. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:458).

Figure 113. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:459).

30 Figure 114. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:460).

Figure 115. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:461).

Figure 116. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:462).

Figure 117. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:463).

5 Figure 118. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:464).

Figure 119. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:465).

10 Figure 120. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:466).

Figure 121. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:467).

Figure 122. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:468).

15 Figure 123. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:469).

Figure 124. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:470).

20 Figure 125. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:471).

Figure 126. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:472).

Figure 127. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:473).

25 Figure 128. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:474).

Figure 129. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:475).

30 Figure 130. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:476).

Figure 131. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:477).

Figure 132. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:478).

Figure 133. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:479).

5 Figure 134. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:480).

Figure 135. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:481).

10 Figure 136. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:482).

Figure 137. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:483).

Figure 138. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:484).

15 Figure 139. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:485).

Figure 140. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:486).

20 Figure 141. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:487).

Figure 142. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:488).

Figure 143. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:489).

25 Figure 144. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:490).

Figure 145. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:491).

30 Figure 146. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:492).

Figure 147. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:493).

Figure 148. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:494).

Figure 149. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:495).

5 Figure 150. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* ATCC19636. (SEQ ID NO:496).

Figure 151. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:497).

10 Figure 152. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:498).

Figure 153. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:499).

Figure 154. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:500).

15 Figure 155. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:501).

Figure 156. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:502).

20 Figure 157. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:503).

Figure 158. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:504).

Figure 159. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:505).

25 Figure 160. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:506).

Figure 161. Kaplan-Meier survival curves showing percent survival after passive immunization and homologous challenge with *S. aureus* ATCC 25904. *A*, intravenous challenge after vaccination with rMntC; *B*, intraperitoneal challenge after 2x vaccination with SIRP extract, 2x vaccination with rSIRP7, or 3x vaccination with rSIRP7; *C*, intravenous challenge after vaccination with rSIRP7.

30 Figure 162. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:543).

Figure 163. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:544).

Figure 164. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:545).

5 Figure 165. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:546).

Figure 166. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:547).

10 Figure 167. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:548).

Figure 168. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:549).

Figure 169. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:550).

15 Figure 170. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:551).

Figure 171. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:552).

20 Figure 172. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:553).

Figure 173. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:554).

Figure 174. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:555).

25 Figure 175. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:556).

Figure 176. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:557).

30 Figure 177. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:558).

Figure 178. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:559).

Figure 179. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:560).

Figure 180. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:561).

5 Figure 181. The amino acid sequence of a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:562).

Figure 182. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:563).

10 Figure 183. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:564).

Figure 184. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:565).

Figure 185. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:566).

15 Figure 186. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:567).

Figure 187. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:568).

20 Figure 188. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:569).

Figure 189. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:570).

Figure 190. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:571).

25 Figure 191. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:572).

Figure 192. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* RF122. (SEQ ID NO:573).

30 Figure 193. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Mu50. (SEQ ID NO:574).

Figure 194. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MRSA252. (SEQ ID NO:575).

Figure 195. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MW2. (SEQ ID NO:576).

Figure 196. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* Newman. (SEQ ID NO:577).

5 Figure 197. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* JH9. (SEQ ID NO:578).

Figure 198. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* USA300. (SEQ ID NO:579).

10 Figure 199. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* COL. (SEQ ID NO:580).

Figure 200. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* NCTC8325. (SEQ ID NO:581).

Figure 201. A nucleic acid sequence encoding a metal-regulated polypeptide obtained from *S. aureus* MSSA476. (SEQ ID NO:582).

15 Figure 202. A Western blot showing binding of mouse convalescent sera to recombinantly-produced metal-regulated polypeptides.

Figure 203. A Western blot showing binding of sera from healthy humans to recombinantly-produced metal-regulated polypeptides.

20 Figure 204. A Western blot showing binding of sera from convalescent humans to recombinantly-produced metal-regulated polypeptides.

Figure 205. Flow cytometry data showing *S. aureus* DU5875 surface expression of metal-regulated polypeptides.

Figure 206. Cytokine induction following vaccination with rSIRP7 and restimulation with either SIRPextract or rSIRP7.

25

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS OF THE INVENTION

30 The present invention provides polypeptides and compositions including polypeptides. As used herein, "polypeptide" refers to a polymer of amino acids linked by peptide bonds. Thus, for example, the terms peptide, oligopeptide, protein, and enzyme are included within the definition of polypeptide. This term also includes post-expression modifications of the polypeptide, such as glycosylations,

acetylations, phosphorylations, and the like. The term polypeptide does not connote a specific length of a polymer of amino acids. A polypeptide may be isolatable directly from a natural source, or can be prepared with the aid of recombinant, enzymatic, or chemical techniques. In the case of a polypeptide that is naturally
5 occurring, such a polypeptide is typically isolated.

An “isolated” polypeptide is one that has been removed from its natural environment. For instance, an isolated polypeptide is a polypeptide that has been removed from the cytoplasm or from the membrane of a cell, and many of the polypeptides, nucleic acids, and other cellular material of its natural environment are
10 no longer present.

A polypeptide characterized as “isolatable” from a particular source is a polypeptide that, under appropriate conditions, is produced by the identified source, although the polypeptide may be obtained from alternate sources using, for example, recombinant, chemical, or enzymatic techniques well known to those skilled in the art.
15 Thus, characterizing a polypeptide as “isolatable” from a particular source does not imply any specific source from which the polypeptide must be obtained or any particular conditions or processes under which the polypeptide must be obtained.

A “purified” polypeptide is one that is at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which
20 they are naturally associated. Polypeptides that are produced outside the organism in which they naturally occur, e.g., through chemical or recombinant means, are considered to be isolated and purified by definition, since they were never present in a natural environment.

As used herein, a “polypeptide fragment” refers to a portion of a polypeptide
25 that results from digestion of a polypeptide with a protease.

Unless otherwise specified, “a,” “an,” “the,” and “at least one” are used interchangeably and mean one or more than one. The terms “comprises” and variations thereof do not have a limiting meaning where these terms appear in the description and claims.

30 A polypeptide of the present invention may be characterized by molecular weight, mass fingerprint, amino acid sequence, nucleic acid that encodes the polypeptide, immunological activity, or any combination of two or more such characteristics. The molecular weight of a polypeptide, typically expressed in

kilodaltons (kDa), can be determined using routine methods including, for instance, gel filtration, gel electrophoresis including sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis (PAGE), capillary electrophoresis, mass spectrometry, liquid chromatography (including HPLC), and calculating the
5 molecular weight from an observed or predicted amino acid sequence. Unless indicated otherwise, molecular weight refers to molecular weight as determined by resolving a polypeptide using an SDS polyacrylamide gel having a stacking gel of about 4% and a resolving gel of about 10% under reducing and denaturing conditions.

10 As used herein, a “mass fingerprint” refers to a population of polypeptide fragments obtained from a polypeptide after digestion with a protease. Typically, the polypeptide fragments resulting from a digestion are analyzed using a mass spectrometric method. Each polypeptide fragment is characterized by a mass, or by a mass (m) to charge (z) ratio, which is referred to as an “m/z ratio” or an “m/z value.”
15 Methods for generating a mass fingerprint of a polypeptide are routine. An example of such a method is disclosed in Example 13.

A polypeptide of the present invention may be a metal-regulated polypeptide. As used herein, a “metal-regulated polypeptide” is a polypeptide that is expressed by a microbe at a greater level when the microbe is grown in low metal conditions
20 compared to growth of the same microbe in high metal conditions. Low metal and high metal conditions are described herein. For instance, one class of metal-regulated polypeptide produced by *Staphylococcus* spp. is not expressed at detectable levels during growth of the microbe in high metal conditions but is expressed at detectable levels during growth in low metal conditions.

25 Examples of metal-regulated polypeptides isolatable from *S. aureus* after growth in low iron conditions have molecular weights of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, and 33 kDa. Examples of metal-regulated polypeptides isolatable from *S. aureus* after growth in low zinc or low copper conditions have molecular weights of 115 kDa, 88 kDa, 80 kDa, 71 kDa, 69 kDa, 35 kDa, 30 kDa,
30 29, kDa, and 27 kDa.

Additional examples of metal-regulated polypeptides include recombinantly-produced versions of polypeptides described herein. A recombinantly-produced polypeptide may include the entire amino acid sequence translatable from an mRNA

transcript. Alternatively, a recombinantly-produced metal-regulated polypeptide can include a fragment or portion of the entire translatable amino acid sequence. For example, a recombinantly-produced metal-regulated polypeptide may lack a cleavable sequence at either terminal of the polypeptide—e.g., a cleavable signal
5 sequence at the amino terminal of the polypeptide.

Thus, a metal-regulated polypeptide can be a polypeptide that includes the amino acid sequence depicted in, for example, SEQ ID NO:353, SEQ ID NO:364, SEQ ID NO:375, SEQ ID NO:386, SEQ ID NO:397, SEQ ID NO:408, and SEQ ID NO:419.

10 The present invention also includes polypeptides that are not metal-regulated. Such polypeptides are expressed in the presence of a metal ion such as, for example, in the presence of ferric chloride, and also expressed when grown in low iron conditions. Examples of such polypeptides isolatable from *S. aureus* have molecular weights of 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41
15 kDa, and 40 kDa.

Whether a polypeptide is a metal-regulated polypeptide or not can be determined by methods useful for comparing the presence of polypeptides, including, for example, gel filtration, gel electrophoresis including sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), capillary electrophoresis, mass
20 spectrometry, and liquid chromatography including HPLC. Separate cultures of a microbe are grown under high metal conditions and under low metal conditions, polypeptides of the present invention are isolated as described herein, and the polypeptides present in each culture are resolved and compared. Typically, an equal amount of polypeptides from each culture is used. Preferably, the polypeptides are
25 resolved using an SDS polyacrylamide gel having a stacking gel of about 4% and a resolving gel of about 10% under reducing and denaturing conditions. For instance, 30 micrograms (μg) of total polypeptide from each culture may be used and loaded into wells of a gel. After running the gel and staining the polypeptides with Coomassie Brilliant Blue, the two lanes can be compared. When determining whether
30 a polypeptide is or is not expressed at a detectable level, 30 μg of total polypeptide from a culture is resolved on an SDS-PAGE gel and stained with Coomassie Brilliant Blue using methods known in the art. A polypeptide that can be visualized by eye is

considered to be expressed at a detectable level, while a polypeptide that cannot be visualized by eye is considered to not be expressed at a detectable level.

Alternatively, whether a polypeptide is metal-regulated or not can be determined using microarray-based gene expression analysis. Separate cultures of a
5 microbe are grown under high metal conditions and under low metal conditions, RNA is extracted from cells of each culture, and differences in RNA expression in cells grown in high metal conditions versus RNA expression in cells grown in low metal conditions are detected and compared. For example, labeled cDNA can be prepared from 8-10 µg of bacterial RNA using established protocols. The labeled
10 cDNA can be applied to a microarray of the *S. aureus* genome. Such microarrays are commercially available and gene expression using such arrays is routine.

Polypeptides of the present invention may have immunological activity. “Immunological activity” refers to the ability of a polypeptide to elicit an immunological response in an animal. An immunological response to a polypeptide
15 is the development in an animal of a cellular and/or antibody-mediated immune response to the polypeptide. Usually, an immunological response includes but is not limited to one or more of the following effects: the production of antibodies, B cells, helper T cells, suppressor T cells, and/or cytotoxic T cells, directed to an epitope or epitopes of the polypeptide. “Epitope” refers to the site on an antigen to which
20 specific B cells and/or T cells respond so that antibody is produced. The immunological activity may be protective. “Protective immunological activity” refers to the ability of a polypeptide to elicit an immunological response in an animal that prevents or inhibits infection by *Staphylococcus* spp., for instance, *S. aureus*. Whether a polypeptide has protective immunological activity can be determined by
25 methods known in the art such as, for example, methods described in Examples 5, 9, or 12. For example, a polypeptide of the present invention, or combination of polypeptides of the present invention, protects a rodent such as a mouse against challenge with a *Staphylococcus* spp. A polypeptide of the present invention may have seroactive activity. “Seroactive activity” refers to the ability of a candidate
30 polypeptide to react with antibody present in convalescent serum from an animal infected with a *Staphylococcus* spp., for instance, *S. aureus*. In some aspects, the convalescent serum may be from an animal infected with the ATCC isolate 19636, strain SAAV1, strain 2176, or strain 1477. Polypeptides of the present invention may

have immunoregulatory activity. “Immunoregulatory activity” refers to the ability of a polypeptide to act in a nonspecific manner to enhance an immune response to a particular antigen. Methods for determining whether a polypeptide has immunoregulatory activity are known in the art.

- 5 A polypeptide of the present invention may have the characteristics of a polypeptide expressed by a reference microbe—i.e., a reference polypeptide. The characteristics can include, for example, molecular weight, mass fingerprint, amino acid sequence, or any combination thereof. The reference microbe can be a gram positive, preferably a member of the family Micrococcaceae, preferably,
- 10 *Staphylococcus* spp., more preferably, *Staphylococcus aureus*. Preferred examples of strains are detailed in Table 1.

Table 1. Bacterial strains.

<u>Bacterial cell</u>	<u>Laboratory designation</u>
<i>S. aureus</i>	ATCC isolate 19636
<i>S. aureus</i>	strain SAAV1
<i>S. aureus</i>	strain 1477
<i>S. aureus</i>	strain 2176

- 15 When the reference microbe is *S. aureus* ATCC isolate 19636, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has a molecular weight of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa 35 kDa, or 33 kDa, and has a mass fingerprint that is similar to the mass fingerprint of a metal-regulated polypeptide expressed by a reference microbe and having a molecular weight of 88
- 20 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa 35 kDa, or 33 kDa, respectively. Preferably, such polypeptides are metal-regulated. For instance, a candidate polypeptide can be a polypeptide of the present invention if it has a molecular weight of 88 kDa and has a mass fingerprint similar to the mass fingerprint of an 88 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* ATCC isolate 19636.

- 25 Alternatively, when the reference microbe is *S. aureus* ATCC isolate 19636, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:353, SEQ ID NO:364, SEQ ID NO:375, SEQ ID NO:386, SEQ ID NO:397, SEQ ID NO:408, or SEQ ID NO:419.

Alternatively, when the reference microbe is *S. aureus* RF122, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:354, SEQ ID NO:365, SEQ ID NO:376, SEQ
5 ID NO:387, SEQ ID NO:398, SEQ ID NO:409, or SEQ ID NO:420.

Alternatively, when the reference microbe is *S. aureus* Mu50, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:355, SEQ ID NO:366, SEQ ID NO:377, SEQ
10 ID NO:388, SEQ ID NO:399, SEQ ID NO:410, or SEQ ID NO:421.

Alternatively, when the reference microbe is *S. aureus* MRSA252, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:356, SEQ ID NO:367, SEQ ID
15 NO:378, SEQ ID NO:389, SEQ ID NO:400, SEQ ID NO:411, or SEQ ID NO:422.

Alternatively, when the reference microbe is *S. aureus* MW2, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:357, SEQ ID NO:368, SEQ ID NO:379, SEQ
20 ID NO:390, SEQ ID NO:401, SEQ ID NO:412, or SEQ ID NO:423.

Alternatively, when the reference microbe is *S. aureus* Newman, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:358, SEQ ID NO:369, SEQ ID NO:380, SEQ
25 ID NO:391, SEQ ID NO:402, SEQ ID NO:413, or SEQ ID NO:424.

Alternatively, when the reference microbe is *S. aureus* JH9, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:359, SEQ ID NO:370, SEQ ID NO:381, SEQ
30 ID NO:392, SEQ ID NO:403, SEQ ID NO:414, or SEQ ID NO:425.

Alternatively, when the reference microbe is *S. aureus* USA300, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the

amino acid sequence of SEQ ID NO:360, SEQ ID NO:371, SEQ ID NO:382, SEQ ID NO:393, SEQ ID NO:404, SEQ ID NO:415, or SEQ ID NO:426.

Alternatively, when the reference microbe is *S. aureus* COL, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:361, SEQ ID NO:372, SEQ ID NO:383, SEQ ID NO:394, SEQ ID NO:405, SEQ ID NO:416, or SEQ ID NO:427.

Alternatively, when the reference microbe is *S. aureus* NCTC 8325, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:362, SEQ ID NO:373, SEQ ID NO:384, SEQ ID NO:395, SEQ ID NO:406, SEQ ID NO:417, or SEQ ID NO:428.

Alternatively, when the reference microbe is *S. aureus* MSSA476, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has an amino acid sequence that is structurally similar, as described in detail below, to the amino acid sequence of SEQ ID NO:363, SEQ ID NO:374, SEQ ID NO:385, SEQ ID NO:396, SEQ ID NO:407, SEQ ID NO:418, or SEQ ID NO:429.

When the reference microbe is *S. aureus* isolate SAAV1, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has a molecular weight of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa, and has a mass fingerprint that is similar to the mass fingerprint of a polypeptide expressed by a reference microbe and having a molecular weight of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa, respectively. Preferably, such polypeptides are metal-regulated. For instance, a candidate polypeptide can be a polypeptide of the present invention if it has a molecular weight of 88 kDa and has a mass fingerprint similar to the mass fingerprint of an 88 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* isolate SAAV1.

When the reference microbe is *S. aureus* strain 2176, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has a molecular weight of 88 kDa, 80 kDa, 65 kDa, 55 kDa, 37 kDa, 36 kDa, 35 kDa, 33 kDa, or 32 kDa, and has a mass fingerprint that is similar to the mass fingerprint of a polypeptide expressed by a reference microbe and having a molecular weight of 88 kDa, 80 kDa, 65 kDa, 55 kDa, 37 kDa, 36 kDa, 35 kDa, 33 kDa, or 32 kDa,

respectively. Preferably, such polypeptides are metal-regulated. For instance, a candidate polypeptide can be a polypeptide of the present invention if it has a molecular weight of 88 kDa and has a mass fingerprint similar to the mass fingerprint of an 88 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* isolate 2176.

When the reference microbe is *S. aureus* strain 1477, a candidate polypeptide can be considered to be a polypeptide of the present invention if it has a molecular weight of 88 kDa, 80 kDa, 65 kDa, 55 kDa, 37 kDa, 36 kDa, 35 kDa, 33 kDa, or 32 kDa, and has a mass fingerprint that is similar to the mass fingerprint of a polypeptide expressed by a reference microbe and having a molecular weight of 88 kDa, 80 kDa, 65 kDa, 55 kDa, 37 kDa, 36 kDa, 35 kDa, 33 kDa, or 32 kDa, respectively. Preferably, such polypeptides are metal-regulated. For instance, a candidate polypeptide can be a polypeptide of the present invention if it has a molecular weight of 88 kDa and has a mass fingerprint similar to the mass fingerprint of an 88 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* isolate 1477.

As used herein, a polypeptide may be “structurally similar” to a reference polypeptide if the amino acid sequence of the polypeptide possesses a specified amount of sequence similarity and/or sequence identity compared to the reference polypeptide. A polypeptide also may be “structurally similar” to a reference polypeptide if the polypeptide exhibits a mass fingerprint possessing a specified amount of identity compared to a comparable mass fingerprint of the reference polypeptide. Thus, a polypeptide may be “structurally similar” to a reference polypeptide if, compared to the reference polypeptide, it possesses a sufficient level of amino acid sequence identity, amino acid sequence similarity, mass fingerprint similarity, or any combination thereof.

Polypeptide sequence similarity and polypeptide sequence identity

Structural similarity of two polypeptides can be determined by aligning the residues of the two polypeptides (for example, a candidate polypeptide and any appropriate reference polypeptide described herein) to optimize the number of identical amino acids along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the number of

identical amino acids, although the amino acids in each sequence must nonetheless remain in their proper order. A reference polypeptide may be a polypeptide described herein or any known metal-regulated polypeptide, as appropriate. A candidate polypeptide is the polypeptide being compared to the reference
5 polypeptide. A candidate polypeptide can be isolated, for example, from a microbe, or can be produced using recombinant techniques, or chemically or enzymatically synthesized.

Unless modified as otherwise described herein, a pair-wise comparison analysis of amino acid sequences can be carried out using the BESTFIT algorithm in
10 the GCG package (version 10.2, Madison WI). Alternatively, polypeptides may be compared using the Blastp program of the BLAST 2 search algorithm, as described by Tatiana et al., (*FEMS Microbiol Lett*, 174, 247-250 (1999)), and available on the National Center for Biotechnology Information (NCBI) website. The default values for all BLAST 2 search parameters may be used, including matrix = BLOSUM62;
15 open gap penalty = 11, extension gap penalty = 1, gap x_dropoff = 50, expect = 10, wordsize = 3, and filter on.

In the comparison of two amino acid sequences, structural similarity may be referred to by percent "identity" or may be referred to by percent "similarity." "Identity" refers to the presence of identical amino acids. "Similarity" refers to the
20 presence of not only identical amino acids but also the presence of conservative substitutions. A conservative substitution for an amino acid in a polypeptide of the invention may be selected from other members of the class to which the amino acid belongs. For example, it is well-known in the art of protein biochemistry that an amino acid belonging to a grouping of amino acids having a particular size or
25 characteristic (such as charge, hydrophobicity and hydrophilicity) can be substituted for another amino acid without altering the activity of a protein, particularly in regions of the protein that are not directly associated with biological activity. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and tyrosine. Polar neutral amino acids
30 include glycine, serine, threonine, cysteine, tyrosine, asparagine and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Conservative substitutions include, for example, Lys for Arg and vice versa to

maintain a positive charge; Glu for Asp and vice versa to maintain a negative charge; Ser for Thr so that a free -OH is maintained; and Gln for Asn to maintain a free -NH₂. Likewise, biologically active analogs of a polypeptide containing deletions or additions of one or more contiguous or noncontiguous amino acids that do not
5 eliminate a functional activity—such as, for example, immunological activity—of the polypeptide are also contemplated.

Thus, as used herein, reference to a polypeptide of the present invention and/or reference to the amino acid sequence of one or more SEQ ID NOs can include a polypeptide with at least 50%, at least 55%, at least 60%, at least 65%, at least
10 70%, at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino acid sequence similarity to the reference amino acid sequence.

Alternatively, as used herein, reference to a polypeptide of the present
15 invention and/or reference to the amino acid sequence of one or more SEQ ID NOs can include a polypeptide with at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino
20 acid sequence identity to the reference amino acid sequence.

Consequently, a polypeptide of the present invention can include certain variants including, for example, homologous polypeptides that originate—biologically and/or recombinantly—from microbial species or strains other than the microbial species or strain from which the polypeptide was originally isolated and/or
25 identified.

For example, a polypeptide of the invention can include a polypeptide commonly known as formate acetyltransferase (PflB). One embodiment of this polypeptide is reflected in SEQ ID NO:353. Variant embodiments are reflected in SEQ ID NO:354, SEQ ID NO:355, SEQ ID NO:356, SEQ ID NO:357, SEQ ID
30 NO:358, SEQ ID NO:359, SEQ ID NO:360, SEQ ID NO:361, SEQ ID NO:362, and SEQ ID NO:363.

As another example, a polypeptide of the invention can include a polypeptide commonly known as oligopeptide permease, peptide-binding protein (Opp1A). One

embodiment of this polypeptide is reflected in SEQ ID NO:364. Variant embodiments are reflected in SEQ ID NO:365, SEQ ID NO:366, SEQ ID NO:367, SEQ ID NO:368, SEQ ID NO:369, SEQ ID NO:370, SEQ ID NO:371, SEQ ID NO:372, SEQ ID NO:373, and SEQ ID NO:374.

5 As another example, a polypeptide of the invention can include a polypeptide commonly known as siderophore compound ABC transporter binding protein (SirA). One embodiment of this polypeptide is reflected in SEQ ID NO:375. Variant embodiments are reflected in SEQ ID NO:376, SEQ ID NO:377, SEQ ID NO:378, SEQ ID NO:379, SEQ ID NO:380, SEQ ID NO:381, SEQ ID NO:382, SEQ ID
10 NO:383, SEQ ID NO:384, and SEQ ID NO:385.

 As another example, a polypeptide of the invention can include a polypeptide sometimes referred to herein as SYN2. One embodiment of this polypeptide is reflected in SEQ ID NO:386. Variant embodiments are reflected in SEQ ID NO:387, SEQ ID NO:388, SEQ ID NO:389, SEQ ID NO:390, SEQ ID NO:391, SEQ ID
15 NO:392, SEQ ID NO:393, SEQ ID NO:394, SEQ ID NO:395, and SEQ ID NO:396.

 As another example, a polypeptide of the invention can include a polypeptide commonly known as ferric hydroxamate-binding lipoprotein (FhuD). One embodiment of this polypeptide is reflected in SEQ ID NO:397. Variant embodiments are reflected in SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400,
20 SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, and SEQ ID NO:407.

 As another example, a polypeptide of the invention can include a polypeptide sometimes referred to herein as SYN1. One embodiment of this polypeptide is reflected in SEQ ID NO:408. Variant embodiments are reflected in SEQ ID NO:409,
25 SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413, SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, and SEQ ID NO:418.

 As another example, a polypeptide of the invention can include a polypeptide commonly known as manganese transport system membrane protein (MntC). One embodiment of this polypeptide is reflected in SEQ ID NO:419. Variant
30 embodiments are reflected in SEQ ID NO:420, SEQ ID NO:421, SEQ ID NO:422, SEQ ID NO:423, SEQ ID NO:424, SEQ ID NO:425, SEQ ID NO:426, SEQ ID NO:427, SEQ ID NO:428, and SEQ ID NO:429.

As another example, a polypeptide of the invention can include a polypeptide commonly known as ferrichrome ABC transporter lipoprotein (SstD). Embodiments of this polypeptide are reflected in SEQ ID NO:543, SEQ ID NO:544, SEQ ID NO:545, SEQ ID NO:546, SEQ ID NO:547, SEQ ID NO:548, SEQ ID NO:549, SEQ ID NO:550, SEQ ID NO:551, and SEQ ID NO:552.

As another example, a polypeptide of the invention can include a polypeptide commonly known as iron compound ABC transporter (FhuD2). Embodiments of this polypeptide are reflected in SEQ ID NO:553, SEQ ID NO:554, SEQ ID NO:555, SEQ ID NO:556, SEQ ID NO:557, SEQ ID NO:558, SEQ ID NO:559, SEQ ID NO:560, SEQ ID NO:561, and SEQ ID NO:562.

A polypeptide of the present invention also can be designed to provide one or more additional sequences such as, for example, the addition of coding sequences for added C-terminal and/or N-terminal amino acids that may facilitate purification by trapping on columns or use of antibodies. Such tags include, for example, histidine-rich tags that allow purification of polypeptides on nickel columns. Such gene modification techniques and suitable additional sequences are well known in the molecular biology arts.

A polypeptide of the present invention also may be designed so that certain amino acids at the C-terminal and/or N-terminal are deleted. For example, one difference between the amino acid sequences of SEQ ID NO:364 and SEQ ID NO:365 is that SEQ ID NO:365 possesses an N-terminal 29 amino acid addition that is not present in the amino acid sequence of the reference polypeptide of SEQ ID NO:364. Similar exemplary N-terminal additions, typically varying from about 20 amino acids to about 35 amino acids, are apparent when one compares, for example, the amino acid sequence of reference peptide SEQ ID NO:353, SEQ ID NO:364, SEQ ID NO:375, SEQ ID NO:386, SEQ ID NO:397, SEQ ID NO:408, or SEQ ID NO:419 with certain variant embodiments of the respective reference polypeptide. Other amino acids additions and/or deletions, at either the N-terminal or the C-terminal, are possible.

A "modification" of a polypeptide of the present invention includes polypeptides (or analogs thereof such as, e.g., fragments thereof) that are chemically or enzymatically derivatized at one or more constituent amino acid. Such modifications can include, for example, side chain modifications, backbone

modifications, and N- and C- terminal modifications such as, for example, acetylation, hydroxylation, methylation, amidation, and the attachment of carbohydrate or lipid moieties, cofactors, and the like, and combinations thereof. Modified polypeptides of the invention may retain the biological activity—such as, for example, immunological activity—of the unmodified polypeptide or may exhibit a reduced or increased biological activity.

The polypeptides of the present invention (including biologically active analogs thereof and modifications thereof) include native (naturally occurring), recombinant, and chemically or enzymatically synthesized polypeptides. For example, a polypeptide of the present invention may be prepared by isolating the polypeptide from a natural source or may be prepared recombinantly by well known methods including, for example, preparation as fusion proteins in bacteria or other host cells.

The polypeptides expressed by a reference microbe can be obtained by growth of the reference microbe under low metal conditions and the subsequent isolation of a polypeptide by the processes disclosed herein. Alternatively, polypeptides expressed by a reference microbe can be obtained by identifying genes expressed at higher levels when the microbe is grown in low metal conditions—i.e., metal-regulated genes. The metal-regulated genes can be cloned and expressed, and the expressed metal-regulated polypeptides may be identified by the processes described herein. A candidate polypeptide can be isolatable from a microbe or identified from a microbe, preferably a gram positive microbe, more preferably, a member of the family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *Staphylococcus aureus*. Other gram positive microbes from which polypeptides can be isolated and/or identified include *Corynebacterium* spp., *Enterococcus* spp., *Erysipelothrix* spp., *Kytococcus* spp., and *Micrococcus* spp., *Mycobacterium* spp., and *Erysipelothrix* spp. A candidate polypeptide may also be produced using enzymatic or chemical techniques.

30

Mass fingerprint similarity

A candidate polypeptide may be evaluated by mass spectrometric analysis to determine whether the candidate polypeptide has a mass fingerprint similar to one of

the polypeptides expressed by a reference microbe and referred to above by molecular weight. Typically, the candidate polypeptide can be isolated, for instance by resolving the candidate polypeptide by gel electrophoresis and excising the portion of the gel containing the candidate polypeptide. Any gel electrophoresis method that separates polypeptides based on differing characteristics can be used, including 1 dimensional or 2 dimensional gel electrophoresis, as well as liquid chromatographic separation based on, for instance, hydrophobicity, pI, or size. The candidate polypeptide can be fragmented, for instance by digestion with a protease. Preferably, the protease can cleave the peptide bond on the carboxy-terminal side of the amino acid lysine and the amino acid arginine, except when the amino acid following the lysine or the arginine is a proline. An example of such a protease is trypsin. Methods for digesting a polypeptide with trypsin are routine and known in the art. An example of such a method is disclosed in Example 13.

Methods for the mass spectrometric analysis of polypeptides are routine and known in the art and include, but are not limited to, matrix assisted laser desorption/ionization time of flight mass spectroscopy (MALDI-TOF MS). Typically, a mixture containing the polypeptide fragments obtained from a candidate polypeptide is mixed with a matrix that functions to transform the laser energy to the sample and produce ionized, preferably monoisotopic, polypeptide fragments. Examples of matrices that can be used include, for instance, sinapinic acid or cyano-4-hydroxycinnamic acid. An example of a method for the analysis of polypeptides by MALDI-TOF MS is described in Example 13. The ionized polypeptide fragments are separated according to their m/z ratio, and detected to yield a spectrum of m/z ratio versus intensity. The spectrum includes m/z values that represent the polypeptide fragments derived from the candidate polypeptide. For any given polypeptide, the amount of each polypeptide fragment resulting from a trypsin digestion should be equimolar. However, it is known that trypsin digestion is not always 100% efficient, for instance, some sites are more efficiently cleaved. Thus, when MALDI-TOF MS is used to determine m/z values, the intensity of each m/z value is typically not identical. Generally, a spectrum has a background level of noise present across most of the x-axis (i.e., the axis having the values of the m/z ratios). This background level of noise varies depending on the running conditions and the machine used, and is easily identified by visual inspection of the spectrum.

An m/z value is generally considered to represent a polypeptide fragment when the intensity is at least 2 times greater, at least 3 times greater, or at least 4 times greater than the background level of noise. The spectrum usually includes other m/z values that are artifacts resulting from, for instance, incomplete digestion, over digestion,
5 other polypeptides that may be present in the mixture, or the protease used to digest the polypeptide including m/z values resulting from autolysis of the protease. This method of digesting a polypeptide with a protease is recognized in the art as resulting in a mass fingerprint of great specificity that can be used to accurately characterize the polypeptide and distinguish it from other polypeptides.

10 In this aspect of the invention, when a candidate polypeptide is analyzed by mass spectroscopy, preferably both the candidate polypeptide and the polypeptide from the reference microbe are prepared and analyzed together, thereby decreasing any potential artifacts resulting from differences in sample handling and running conditions. Preferably, all reagents used to prepare and analyze the two polypeptides
15 are the same. For instance, the polypeptide from the reference microbe and the candidate polypeptide are isolated under substantially the same conditions, fragmented under substantially the same conditions, and analyzed by MALDI-TOF MS on the same machine under substantially the same conditions. A candidate polypeptide may be considered to be "structurally similar" to a reference polypeptide
20 if it exhibits a mass fingerprint possessing at least 80%, at least 90%, at least 95%, or substantially all of the m/z values present in the spectrum of the reference microbe polypeptide and above the background level of noise are also present in the spectrum of the candidate polypeptide. (See, e.g., United States Patent Application Publication No. 2006/0233824 A1).

25 In another aspect, a polypeptide can be considered to be a polypeptide of the present invention if it has a molecular weight of a reference polypeptide described in Table 2, 3, 4, or 5 and has a mass fingerprint that includes a subpopulation including at least a specified percentage of the polypeptide fragments of the reference polypeptide as listed in Table 2, 3, 4, or 5. For instance, a polypeptide of the present
30 invention includes a polypeptide of 88 kDa and a mass fingerprint that includes a specified percentage of polypeptide fragments having masses of HVDVR (SEQ ID NO: 1), YSYER (SEQ ID NO: 2), IIGDYRR (SEQ ID NO: 3), IFTDYRK (SEQ ID NO: 4), ELKELGQK (SEQ ID NO: 5), YAQVKPIR (SEQ ID NO: 6), QMQFFGAR

(SEQ ID NO: 7), SMQPFGGIR (SEQ ID NO: 8), VSGYAVNFIK (SEQ ID NO: 9),
 NHATAWQGFK (SEQ ID NO: 10), LWEQVMQLSK (SEQ ID NO: 11),
 SLGKEPEDQNR (SEQ ID NO: 12), DGISNTFSIVPK (SEQ ID NO: 13),
 AGVITGLPDAYGR (SEQ ID NO: 14), TSTFLDIYAER (SEQ ID NO: 15),
 5 SMQPFGGIRMAK (SEQ ID NO: 16), THNQGVFDAYSIR (SEQ ID NO: 17),
 KAGVITGLPDAYGR (SEQ ID NO: 18), TLLYAINGGKDEK (SEQ ID NO: 19),
 IEMALHDTEIVR (SEQ ID NO: 20), AGEPFAPGANPMHGR (SEQ ID NO: 21),
 VALYGVDFLMEEK (SEQ ID NO: 22), KTHNQGVFDAYSIR (SEQ ID NO: 23),
 YGFDLSRPAENFK (SEQ ID NO: 24), TSSIQYENDDIMR (SEQ ID NO: 25),
 10 KAGEPFAPGANPMHGR (SEQ ID NO: 26), RVALYGVDFLMEEK (SEQ ID NO:
 27), LWEQVMQLSKEER (SEQ ID NO: 28), MLETNKNHATAWQGFK (SEQ ID
 NO: 29), MHDFNTMSTEMSEDVIR (SEQ ID NO: 30),
 YGNNDDRVDDIAVDLVER (SEQ ID NO: 31), ETLIDAMEHPPEEYPQLTIR
 (SEQ ID NO: 32), YAQVKPIRNEEGLVVDFFEIEGDFPK (SEQ ID NO: 33).

15 The mass fingerprint of a candidate polypeptide can be determined by a mass
 spectrometric method, for instance by MALDI-TOF MS. The mass fingerprint of a
 candidate polypeptide will generally have additional polypeptide fragments and,
 therefore, can have additional m/z values other than those listed for a polypeptide in
 Table 2, 3, 4, or 5. When the candidate polypeptide is being compared to a
 20 polypeptide in Table 2, 3, 4, or 5, the candidate polypeptide can be isolatable from a
 microbe, preferably a gram positive microbe, more preferably, a member of the
 family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably,
Staphylococcus aureus. Other gram positive microbes include *Corynebacterium*
 spp., *Enterococcus* spp., *Erysipelothrix* spp., *Kytococcus* spp., *Listeria* spp.,
 25 *Micrococcus* spp., and *Mycobacterium* spp., and *Erysipelothrix* spp.

A candidate polypeptide can be obtained by growth of a microbe under low
 metal conditions and the subsequent isolation of a polypeptide by the processes
 described herein. Alternatively, a candidate polypeptide can be obtained by
 recombinant expression of a polynucleotide that encodes the candidate polypeptide.

30 It is well known in the art that modifications of amino acids can be
 accidentally introduced during sample handling, such as oxidation, and formation of
 carbamidomethyl derivatives. Further, these types of modifications alter the m/z
 value of a polypeptide fragment. For instance, if a polypeptide fragment contains a

methionine that is oxidized, the m/z value will be increased by 16 relative to the same fragment that does not contain the oxidized methionine. Accordingly, those polypeptide fragments in Tables 2, 3, 4, or 5 having the notation "oxidation (M)" have an m/z value that is increased by 16 relative to the same fragment that does not contain the oxidized methionine. It is understood that the polypeptide fragments of Table 2, 3, 4, or 5 can be modified during sample handling.

Polynucleotide sequence similarity and polynucleotide sequence identity

Polypeptides of the invention also may be identified in terms the polynucleotide that encodes the polypeptide. Thus, the invention includes polynucleotides that encode a polypeptide of the invention or hybridize, under standard hybridization conditions, to a polynucleotide that encodes a polypeptide of the invention, and the complements of such polynucleotide sequences.

As used herein, reference to a polynucleotide of the present invention and/or reference to the nucleic acid sequence of one or more SEQ ID NOs can include polynucleotides having a sequence identity of at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to an identified reference polynucleotide sequence.

Table 2. Characteristics of polypeptides obtained from *S. aureus* ATCC isolate 19636.

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P23	88	625.4	HVDVR	1
		717.3	YSYER	2
		892.5	IIGDYRR	3
		942.5	IETDYRK	4
		944.5	ELKELGQK	5
		974.6	YQVVKPIR	6
		984.5	QMQLFGAR	7
		992.5	SMQPFGGIR	8
		1097.6	VSGYAVNFIK	9
		1159.5	NHATAWQGFK	10
		1261.7	LWEQVMQLSK	11
		1272.7	SLGKEPEDQNR	12
		1277.7	DGISTFSIVPK	13
		1289.7	AGVITGLPDAYGR	14
		1315.7	TSTFLDIYAEK	15
		1322.7	SMQPFGGIRMAK	16
		1394.7	THNQGVFDAYS	17
		1417.8	KAGVITGLPDAYGR	18
		1421.8	TLLYAINGGKDEK	19
		1426.8	IEMALHDTIVR	20
		1508.8	AGEPFAPGANPMHGR	21
		1513.9	VALYGVDFLMEEK	22
		1522.8	KTHNQGVFDAYS	23
		1543.9	YGFDLSRPAENFK	24

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1571.8	TSSIQYENDDIMR	25
		1636.9	KAGEPFAPGANPMHGR	26
		1670.0	RVALYGVDFLMEEK	27
		1676.0	LWEQVMQLSKEER	28
		1876.2	MLETNKNHATAWQGFK	29
		2043.1	MHDFNTMSTMSSEVIR	30
		2078.2	YGNNDDDRVDIAVDLVER	31
		2285.5	ETLIDAMEHPPEYPQLTIR	32
		2892.9	YAVKPIRNEEGLVVDFEIEGDFPK	33
P25	55	783.6	LHSWLK	34
		911.7	KLHSWLK	35
		937.6	TYTFHLR	36
		996.6	KFDGTGPFK	37
		1025.6	QAIGHMVMNR	38
		1063.6	KWDVSEDGK	39
		1185.6	IYNSIDDAFK	40
		1277.6	NLEMAMYDYK	41
		1324.7	ENKQLTYTTVK	42
		1346.7	AESLLDEAGWKK	43
		1381.8	TVRQAIGHMVMNR	44
		1394.8	TYTFHLRDDVK	45

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1400.7	KGETNFAFTDDR	46
		1419.7	FHDGTPFDADAVK	47
		1422.8	NVTDINFDMPTR	48
		1428.8	DKIYNSIDDAFK	49
		1483.8	EQAELYLQAEFKK	50
		1509.8	VMPAGETAFLSMKK	51
		1547.9	FHDGTPFDADAVKK	52
		1550.9	NVTDINFDMPTRK	53
		1559.9	LNINGETSDKIAER	54
		1788.1	EILDGQEKPATQLFAK	55
		1930.1	GSSSQKEQAEYLQAEFK	56
		1946.0	DESADFNKNDQYWGEK	57
		2100.4	IAKEILDGQEKPATQLFAK	58
		2239.3	VSFTQSQYELPFNEMQYK	59
		2493.5	EAYQPALAEELAMPRPYVFSVK + Oxidation (M)	60
		2900.6	DIGDMNPHVYGGSMSAESMIYEPLVR + 2 Oxidation (M)	61
		2916.6	DIGDMNPHVYGGSMSAESMIYEPLVR + 3 Oxidation (M)	62

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P26	38	993.6	IVYVGADEK	63
		996.7	QALNNPVLK	64
		1237.7	ETVKIENNYK	65
		1272.7	ENPDVILAMDR	66
		1502.0	LAATKPEVIFISGR	67
		1507.9	NAVVDYGALDVMK	68
		1523.9	ALPNFLESFKDDK	69
		1559.9	LWYFAAGSTTTTK	70
		1716.0	FGGLVYDTLGFNAVDK	71
		1737.0	IVYVGADEKNLIGSMK	72
		1844.1	FGGLVYDTLGFNAVDKK	73
		1929.1	GRFGGLVYDTLGFNAVDK	74
		1998.2	TVMYLLVNEGELSTFGPK	75
		2234.4	EVNFDKIAATKPEVIFISGR	76
		3143.8	VSNSNHGQNVSNQYVKNENPDVILAMDR	77
P27	37	699.5	FEYIK	78
		729.4	DAWPLK	79
		792.5	ASVVNFR	80
		852.4	VYDQLSK	81
		987.5	HAMGTTEIK	82
		1008.5	LIDDLYEK	83
		1020.5	YKDAWPLK	84
		1074.5	EKEAEDLLK	85
		1083.6	LKPDILVASK	86
		1169.5	FEYIKNDLK	87
		1182.5	KTESEWTSSK	88

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1184.5	YDDKVAAFQK	89
		1223.5	NEKVYDQLSK	90
		1278.6	IAPTVSTDTIVFK	91
		1497.6	TESEWTSSKEWK	92
		1502.7	DAWPLKASVVNFR	93
		1558.8	QVDNGKDIQLTSK	94
		1605.8	LIDDLYEKLNIEK	95
		1623.8	IVGQEPAPNLEEISK	96
		1712.8	ESIPLMNADHIFVVK	97
		1800.9	IYAGGYAGEILNDLGFK	98
		1957.0	IYAGGYAGEILNDLGFKR	99
		2252.0	NNQVSDDLDEITWNLGGYK	100
		3383.9	RVVTLYQGATDVAVSLGVKPVGAVESWTQKPK	101
P28	36	646.4	DVWAR	102
		725.5	IIKPVR	103
		1068.4	IGDYTSVGTR	104
		1185.5	KQPNLEEISK	105
		1327.6	LKPDLIADSSR	106
		1343.6	VDIVDRDVWAR	107
		2080.9	GPYLQDTEHLADLNPER	108
		2438.1	AGLLAHPNYSYVGQFLNELGFK	109
		2789.4	IVVLEYSEADALAALDVKPVGIADDDGK	110

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P29	35	760.5	AGWAEVK	111
		1012.6	TVDPKDPK	112
		1107.6	KDWEETAK	113
		1204.7	VAPTVDYDK	114
		1238.6	YLEQEMLGK	115
		1244.6	LYTYGDNWGR	116
		1259.7	IADVPTYAGGLK	117
		1281.7	GGEVLYQAFGLK	118
		1516.8	AGWAEVKQEEIEK	119
		1683.9	LGANIVAVNQVDQSK	120
		1877.1	EKPDLIIVYSTDKDIK	121
		1884.0	AIGQDATVSLDEFDKK	122
		2227.1	VDAGTYWYNDPYTLDFMR	123
		2781.4	YAGDYIVSTSEKPTPGYESTNMWK	124
		834.5	QAIEFVK	125
		864.5	YIAQLEK	126
		946.5	QGTPEQMR	127
		962.5	QAIEFVKK	128
P30	33	976.5	DKFNDIPK	129
		1054.5	AMITSEGAFK	130
		1202.5	SNITVHGSMK	131
		1268.6	HLLVETSVDDKK	132

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1443.6	DIFGEVYTDSIGK	133
		1450.7	TIQQTIFDNDKK	134
		1454.7	VVTINSILYDMAK	135
		1571.7	KDIFGEVYTDSIGK	136
		1593.7	QDPHAWLSLDNGIK	137
		1818.9	DVKPIYLNAGEEGNKDK	138
		1836.9	DKQDPHAWLSLDNGIK	139
		1911.9	QYGITPGYIWEINTEK	140
		2582.3	LTDAADVILYNGLNLETGNGWFEK	141
		2710.2	KLTDADVILYNGLNLETGNGWFEK	142
		2942.4	NVGGDNVDIHSIVPGQDPHEYEYKPK	143

1. Molecular weight as determined by SDS-PAGE.

2. The m/z value of a polypeptide fragment can be converted to mass by subtracting 1 from the m/z value. Each mass includes a range of plus or minus 300 parts per million (ppm), or plus or minus 1 Da.

Table 3. Characteristics of polypeptides obtained from *S. aureus* isolate SAAV1.

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P33A	55	783.4	LHSWLK	144
		911.5	KLHSWLK	145
		937.5	TYTFHLR	146

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) [†]	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		996.5	KFDGTGPFK	147
		1025.5	QAIGHMVR	148
		1039.4	NDQYWGEK	149
		1178.5	GTDSDKDSLK	150
		1185.5	IYNSDDAFK	151
		1222.6	DKYTVLNLK	152
		1229.5	ISTLDNVKVK	153
		1346.6	AESLLDEAGWKK	154
		1355.5	EQAEYLQAEFK	155
		1381.6	VMPAGETAFLSMK	156
		1400.5	KGETNFAFTDDR	157
		1419.6	FHDGTPFDADAVK	158
		1422.6	NVTDINFDMPTR	159
		1483.6	EQAEYLQAEFKK	160
		1547.7	FHDGTPFDADAVKK	161
		1550.6	NVTDINFDMPTRK	162
		1559.7	LNINGETSDKIAER	163
		1787.9	EILDGQEKPATQLFAK	164
		1945.8	DESADFNKNDQYWGEK	165
		2239.0	VSFTQSQYELPFNEMQYK	166
		2354.1	QIDDEGIFIPISHGSMTVVAPK	167
P33B	55	2868.1	DIGDMNPHVYGGMSAESMIYEPLVR	168
		895.4	FPYAANGR	169
		904.5	ALLHASHR	170
		1045.5	EEGLAIKASK	171
		1384.5	GEAYFVDNNSLR	172

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1435.7	TIEADYVLVTVGR	173
		1669.8	RPNTDELGLEELGVK	174
		1841.0	NAILATGSRPIEPNFK	175
		2179.2	TSISNIYAIGDIVPGLPLAHK	176
		2546.2	FVEAQHSENGLVIAESVSLNFQK	177
		2587.3	VVGDFPIETDTIVIGAGPGGYVAAIR	178
P35	37	699.4	FEYIK	179
		729.4	DAWPLK	180
		792.4	ASVVNFR	181
		852.4	VYDQLSK	182
		1008.4	LIDDLYEK	183
		1020.4	YKDAWPLK	184
		1074.4	EKEAEDLLK	185
		1083.5	LKPDLLIVASK	186
		1169.5	FEYIKNDLK	187
		1182.4	KTESEWTSSK	188
		1184.4	YDDKVAAAFQK	189
		1278.5	IAPTVSTDTVEK	190
		1558.7	QVDNGKDIQLTSK	191
		1623.7	IVGQEPAPNLEEISK	192
		1712.7	ESIPLMNADHIFVVK	193
		1800.7	IYAGGYAGEILNDLGFK	194
		1956.8	IYAGGYAGEILNDLGFKR	195
		2251.9	NNQVSDDLDEITWNLGGYK	196
		3227.5	VVTLYQGATDVAVSLGVKPYGAVESWTQKPK	197
P38	33	864.5	YIAQLEK	198

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		946.4	QGTPEQMR	199
		976.5	DKFNIDPK	200
		1054.5	AMITSEGAFK	201
		1146.5	ENDIPKEQR	202
		1268.6	HLLVETSVDDK	203
		1322.5	TIQQTFIDNDK	204
		1443.6	DIFGEVYTDISIGK	205
		1450.6	TIQQTFIDNDKK	206
		1454.6	VVTTNSILYDMAK	207
		1593.7	QDPHAWLSLDNGIK	208
		1818.9	DVKPIYLNAGEEGNKDK	209
		1836.8	DKQDPHAWLSLDNGIK	210
		1911.9	QYGITPGYIWEINTEK	211
		2942.4	NVGDDNVDIHSIVPGQDPHEVEVKPK	212

1. Molecular weight as determined by SDS-PAGE.

2. The m/z value of a polypeptide fragment can be converted to mass by subtracting 1 from the m/z value. Each mass includes a range of plus or minus 300 parts per million (ppm) or plus or minus 1 Da.

Table 4. Characteristics of polypeptides obtained from *S. aureus* isolate 2176.

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P478	88	736.35	IIGDYR	213
		814.49	IFIDYR	214
		942.42	IFTDYRK	4
		945.36	TGNTPDGRK	215
		974.40	YAOVKPIR	6
		984.27	QMFFGAR	7
		992.41	SMQFFGIR	8
		1087.31	EQQLDVISR	216
		1097.31	VSGYAVNFIK	9
		1159.37	NHATAWQGFK	10
		1261.37	LWEQVMQLSK	11
		1289.46	AGVITGLPDAYGR	14
		1315.42	TSTFLDIYAER	15
		1322.39	LREELSEQYR	217
		1394.37	THNQGVFDAYS	17
		1417.52	KAGVITGLPDAYGR	18
		1426.36	IEMALHDTIVR	20
		1487.39	NHATAWQGFKNR	218
		1508.42	AGEPFAPGANPMHGR	21
		1513.52	VALYGVDFLMEEK	22
		1543.43	YGFDLSPAEFIK	24
		1571.50	TSSIQYENDDIMR	25
		1636.56	KAGEPFAPGANPMHGR	26
		1859.80	DLETIVGVQTEKPFKR	219

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1876.77	TMATGIAGLSVAADSLSAIK	220
		2042.57	MHDENTMSTEMSESVIR	30
		2077.68	YGNNDRRVDDIAVDLVER	31
		2158.88	AGVITSEVQEIDHFIMK	221
		2284.90	ETLIDAMEHPPEYPQLTIR	32
		2575.08	FLHSLDNLGPAPEPNLTVLWSVR	222
		2628.01	SGAQVGNFEGINSEVLEYDEVFK	223
		2756.06	SGAQVGNFEGINSEVLEYDEVFKK	224
		3262.33	VASTITSHDAGYLDKDLLETIVGVQTEKPEK	225
P479	80	625.27	HVDVR	1
		736.26	IIGDYR	226
		814.22	IFTDYR	227
		942.27	IFTDYRK	4
		974.26	YQVKPIR	6
		984.18	QMFFGAR	7
		992.23	SMQPFGGIR	8
		1087.16	EQQLDVISR	228
		1097.24	VSGYAVNFIK	9
		1159.12	NHATAWQGFK	10
		1243.14	VDDIAVDLVER	229
		1261.22	LWEQVMQLSK	11
		1272.24	SLGKEPEDQNR	12
		1277.18	DGISTNFTSIVPK	13
		1289.21	AGVITGLPDAYGR	14
		1315.19	TSTFLDIYAER	15
		1322.21	LREELSEQYR	230

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1394.16	THNQGVFDAYS	17
		1417.32	KAGVITGLPDAYGR	18
		1426.23	IEMALHDTIVR	20
		1487.19	NHATAWQGFKNR	231
		1508.25	AGEPFAPGANPMHGR	21
		1513.21	VALYGVDFLMEEK	22
		1522.25	KTHNQGVFDAYS	23
		1543.26	YGFDSLSPAENFK	24
		1571.23	TSSIQYENDDIMR	25
		1636.29	KAGEPFAPGANPMHGR	26
		1703.43	DLETIVGVQTEKPFK	232
		1751.45	EAVQWLYLAYLAALK	233
		1859.53	DLETIVGVQTEKPFKR	234
		1876.50	TMATGIAGLSVAADSLSAIK	235
		1936.37	NEEGLVVDFFIEGDFPK	236
		2042.43	MHDFNTMSTMSSESVIR	30
		2077.45	YGNNDRRVDDIAVDLVER	31
		2158.57	AGVITSEVQEIDHFIMK	237
		2284.61	ETLIDAMEHPPEYPLTIR	32
		2574.77	FLHSLDNLGPAPEPNLTVLWSVR	238
		2627.61	SGAQVGNFEGINSEVLEYDEVFK	239
		2755.70	SGAQVGNFEGINSEVLEYDEVFK	240
		2907.65	EFIQLNYTLYEGNDSFLAGPTEATSK	241
		3261.91	VASTTSHDAGYLDKDLTIVGVQTEKPFK	242
		3421.02	TPDYNELFSGDPTWVTESIGGVGIDGRPLVTK	243
P480	65	625.35	HVDVR	1

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		717.38	YSYER	2
		733.42	LPDNFK	244
		736.44	IIGDYR	245
		814.33	IFTDYR	246
		853.31	YGNDDRR	247
		942.33	IFTDYRK	4
		944.39	ELKELGQK	5
		974.52	YQVVKPIR	6
		984.36	QMFFGAR	7
		992.44	SMQFFGGIR	8
		1049.44	TLLYAINGGK	248
		1087.43	EQQLDVISR	249
		1097.51	VSGYAVNFIK	9
		1159.52	NHATAWQGFK	10
		1289.53	AGVITGLPDAYGR	14
		1315.51	TSTFLDIYAER	15
		1322.46	LREELSEQYR	250
		1394.50	THNQGVFDAYS	17
		1417.65	KAGVITGLPDAYGR	18
		1442.56	IEMALHDTIVR + Oxidation (M)	251
		1467.60	VSGYAVNFIKLTR	252
		1522.61	KTHNQGVFDAYS	23
		1524.55	AGEPFAPGANPMHGR + Oxidation (M)	253
		1529.64	VALYGVDFLMEEEK + Oxidation (M)	254
		1543.62	YGFDSLSPAEINFK	24
		1652.68	KAGEPFAPGANPMHGR + Oxidation (M)	255

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1671.76	TSTFLDIYAERDLK	256
		1766.76	VDDIAVDLVERFMK + Oxidation (M)	257
		1876.86	TMATGIAGLSVAADSLSAIK	258
		2077.93	YGNNDDRVDDIAVDLVER	31
		2225.07	DSEHTMSVLTTISNVVYGKK + Oxidation (M)	259
		2575.33	FLHSLDNLGPAPEPNI.TVLWSVR	260
		2628.25	SGAQVGNPNEGINSVLEYDEVFK	261
		2748.36	NLTSMILDGYAMQCGHHLNINVFNR	262
		2756.63	SGAQVGNPNEGINSVLEYDEVFK	263
		3001.02	DEKSGAQVGNPNEGINSVLEYDEVFK	264
		3420.75	TPDYNELFSGDPTWVTESIGGVGIDGRPLVTK	265
P481	55	634.33	AKSNSK	266
		883.24	TFYPEAR	267
		1014.24	QFWGHLVK	268
		1131.17	WIPLMMKGR	269
		1207.21	VINEEFESK	270
		1324.10	NEDWQLYTAK	271
		1360.28	TLLFGPFANVGPK	272
		1386.31	LDRPAIESSNER	273
		1565.30	IDEGTDVNFGLTR	274
		1584.34	EFINPLPHISYVR	275
		1699.29	EIEPDWNIHVYER	276
		1744.36	EPPGTPPMTVPFLDTR	277
		2046.52	QVTDYVFIGAGGGAIPLLQK	278
		2189.43	TFYPEARNEDWQLYTAK	279

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P482	37	2806.58	HLGGFPISGQFLACTNPQVIEQHDAK	280
		699.28	FEYIK	281
		729.26	DAWPLK	282
		792.33	ASVVNFR	283
		852.28	VYDQLSK	284
		1008.30	LIDDLYEK	285
		1020.31	YKDAWPLK	286
		1083.43	LKPDLIIVASK	287
		1278.36	IAPTVSTDTVFK	288
		1623.44	IVGQEPAPNLEEISK	289
		1712.62	ESIPLMNADHIFVVK	290
		1800.61	IYAGGYAGEILNDLGFK	291
		1956.77	IYAGGYAGEILNDLGFKR	292
		2251.77	NNQVSDDLDEITWNLGGYK	293
P483	36	3227.44	VVTLYQGATDVAVSLGVKPVGAVESWTQKPK	294
		646.50	DVWAR	295
		672.41	KLNAVK	296
		716.41	VDIVDR	297
		725.61	IIKPVR	298
		842.50	IAPTLILK	299
		850.47	QNINSFK	300
		1068.50	IGDYTSVGTR	301
		1075.42	MIIMTDHAK + Oxidation (M)	302
		1185.53	KQPNLEEISK	303
		1327.59	LKPDLIADSSR	304
		1343.58	VDIVDRDVWAR	305

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1592.76	LKPDLIADSSRHK	306
		2081.00	GPYLQDTEHLADLNPER	307
		2438.24	AGLLAHPNYSYVGQFLNELGFK	308
		2789.48	IVVLEYSFADALAAALDVKPVGIADDGK	309
		2917.60	IVVLEYSFADALAAALDVKPVGIADDGKK	310
P484	35	857.38	AAADLAGR	311
		1022.23	NIEADTGMR + Oxidation (M)	312
		1056.32	VVDANIAAQR	313
		1075.36	ADIDLPER	314
		1285.44	LVGGAGEETIAR	315
		1435.44	AMAVATEQEMKAR	316
		1632.50	HHEVLEPNPNISK	317
		1813.65	VVEAESEVPLAMAEALR	318
		1887.67	VIETFIAGVAMNGIEVK	319
		2299.85	AGLALTTNQLESHYLAGGNVDR	320
		2806.95	TVLSKGLDSGTAFELSIDIADVDSK	321
		3337.42	AGLALTTNQLESHYLAGGNVDRVVDANIAAQR	322
P485	33	625.28	ADYEK	323
		864.28	YIAQLEK	324
		946.23	QGTPEQMR	325
		1045.26	ALEQAGKSLK	326
		1268.35	HLLVETSVDDKK	327
		1443.34	DIFGEVYTDGK	328
		1450.40	TIQQTFIDNDKK	329
		1454.37	VVTNSILYDMAK	330
		1571.45	KDIFGEVYTDGK	331

Polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1576.44	DVKPIYLNKEEGNK	332
		1593.47	QDPHAWLSLDNGIK	333
		1819.59	DVKPIYLNKEEGNKDK	334
		1836.62	DKQDPHAWLSLDNGIK	335
		1911.66	QYGITPGYIWEINTEK	336
		2172.83	VIAVSKDVKPIYLNKEEGNK	337
		2582.00	LTDADVILYNGLNLETGNGWFEEK	338
		2942.26	NVGGDNVDIHSIVPVGGDPHEYEVKPK	339
P486	32	625.42	ADYEK	340
		864.41	YIAQLEK	341
		1268.48	HLLVETSVDDKK	342
		1443.49	DIFGEVYTDISIGK	343
		1450.53	TIQQTIFIDNDKK	344
		1454.61	VVTNSILYDMAK	345
		1576.64	DVKPIYLNKEEGNK	346
		1593.57	QDPHAWLSLDNGIK	347
		1818.77	DVKPIYLNKEEGNKDK	348
		1836.78	DKQDPHAWLSLDNGIK	349
		1911.81	QYGITPGYIWEINTEK	350
		2582.18	LTDADVILYNGLNLETGNGWFEEK	351
		2942.32	NVGGDNVDIHSIVPVGGDPHEYEVKPK	352

1. Molecular weight as determined by SDS-PAGE.

2. The m/z value of a polypeptide fragment can be converted to mass by subtracting 1 from the m/z value. Each mass includes a range of plus or minus 400 parts per million (ppm) or 1 Dalton.

Table 5. Characteristics of polypeptides obtained from *S. aureus* bovine isolate 1477.

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
P487	88	717.39	YSYER	2
		736.52	IIGDYR	583
		814.46	IFTDYR	584
		942.46	IFTDYRK	4
		974.54	YQVKPIR	6
		984.41	QMFFGAR	7
		992.40	SMQFFGIR	8
		1087.49	EQQLDVISR	585
		1097.50	VSGYAVNFIK	9
		1159.39	NHATAWQGFK	10
		1261.45	LWEQVMQLSK	11
		1272.50	SLGKEPEDQNR	12
		1277.50	DGISTFIVPK	13
		1289.54	AGVITGLPDAYGR	14
		1315.54	TSTFLDIYAEK	15
		1322.53	LREELSEQYR	586
		1394.50	THNQGVFDAYS	17
		1417.62	KAGVITGLPDAYGR	18
		1426.65	IEMALHDTIVR	20
		1508.59	AGEPFAPGANPMHGR	21
		1522.61	KTHNQGVFDAYS	23
		1543.68	YGFDSLSPAENFK	24
		1877.74	TMATGIAGLSVAADSLSAIK	587
		2077.86	YGNNDDRVDDIAVDLVER	31

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		2159.08	AGVITSEVQEIIDHFIMK	588
		2285.07	ETLIDAMEHPPEYPQLTIR	32
		2575.32	FLHSLDNLGPAPEPNTVLWSVR	589
		2628.24	SGAQVGNFEGINSEVLEYDEVFK	590
		2756.41	SGAQVGNFEGINSEVLEYDEVFKK	591
		3262.68	VASTITSHDAGYLDKDLETIVGVQTEKPFK	592
P488	80	625.49	HVDVR	1
		814.54	IFTDYR	593
		942.66	IFTDYRK	4
		974.69	YQVVKPIR	6
		984.59	QMQLFFGAR	7
		992.55	SMQPFGGIR	8
		1159.64	NHATAWQGFK	10
		1261.63	LWEQVMQLSK	11
		1272.74	SLGKEPEDQNR	12
		1277.69	DGISTFTSIVPK	13
		1289.76	AGVITGLPDAYGR	14
		1315.73	TSTFLDIYAER	15
		1322.72	SMQPFGGIRMAK	16
		1394.73	THNQGVFDAYS	17
		1417.86	KAGVITGLPDAYGR	18
		1422.76	TLLYAINGGKDEK	19
		1426.80	IEMALHDTIVR	20
		1508.82	AGEPFAPGANPMHGR	21
		1513.80	VALYGVDFLMEEK	22
		1543.82	YGFDSLSPAENFK	24

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1571.82	TSSIQYENDDIMR	25
		1703.99	DLETIVGVQTEKPFK	594
		1860.23	DLETIVGVQTEKPFKR	595
		1877.07	TMATGIAGLSVAADSLSAIK	596
		1937.09	NEEGLVDFEIEGDFPK	597
		2078.13	YGNNDLDRVDDIAVDLVER	31
		2575.56	FLHSLDNLGPAPEPNLTVLWSVR	598
		2628.30	SGAQVGPNFEGINSEVLEYDEVFK	599
		2908.63	EFIQLNYTLYEGNDSFLAGPTEATSK	600
P489	65	733.67	IVKFAR	601
		944.71	ELKELGQK	5
		974.79	YAVVKPIR	6
		984.69	QMFFGAR	7
		1049.83	TLLYAINGGK	602
		1087.78	EQQLDVISR	603
		1097.79	VSGYAVNFIK	9
		1243.80	VDDIAVDLVER	604
		1272.82	SLGKEPEDQNR	12
		1289.87	AGVITGLPDAYGR	14
		1299.92	LPDNFKTYCAK	605
		1315.83	TSTFLDIYAER	15
		1322.84	SMQPFGGIRMAK	16
		1390.93	DQKGALSSLSVAK	606
		1394.84	THNQGVFDAYS	17
		1577.94	VASTITSHDAGYLDK	607
		1637.09	KAGEFFAPGANPMHGR	26

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1704.16	DLETIVGVQTEKPFK	608
		2030.42	MSIKTSSIQYENDDIMR	608
		2078.34	YGNNDDRVDLIAVDLVER	31
		2284.60	ETLIDAMEHPPEYPQLTIR	32
		2575.77	FLHSLDNLGPAPEPNLTVLWVSR	610
		2628.64	SGAQVGPNFEGINSEVLEYDEVFK	611
P490	55	883.81	TFYPEAR	612
		1014.87	QFWGHLVK	613
		1131.97	WPLMMKGR	614
		1207.99	VINEEFEISK	615
		1231.97	YSFDQVIMTK	616
		1325.02	NEDWQLYTAGK	617
		1361.17	TLFGPFANVGPK	618
		1362.14	GRDNPFGIMAAASK + Oxidation (M)	619
		1387.14	LDRPAIESSNER	620
		1481.24	NEDWQLYTAGKR	621
		1566.28	IDEGTDVNFGEIIR	622
		1585.34	EFINPLPHISYVR	623
		1700.36	EIEPDWNIHVIYR	624
		1761.49	EPPGTPPMTVPRLDTR + Oxidation (M)	625
		2047.67	QVTDYVFIGAGGGAIPLLQK	626
		2208.82	VYGKEPPGTPPMTVPRLDTR + Oxidation (M)	627
		2865.21	HLGGFPISGQFLACTNPQVIEQHDAK	628
P492	36	857.57	AAAILAGR	629
		1056.59	VVDANIAAQR	630
		1075.61	ADILPFER	631

polypeptide designation	Approximate molecular weight in kilodaltons (kDa) ¹	m/z value of polypeptide fragments resulting from trypsin digest ²	Predicted amino acid sequence of the polypeptide fragment	SEQ ID NO:
		1285.74	LVGGAGEETILAR	632
		1632.95	HHTEVLEPNPNISK	633
		1814.09	VVEAESEVPLAMAEALR	634
		2284.45	AAIDLAGRDVLEAVQMSVNPk + Oxidation (M)	635
		2300.40	AGLALTNQLESHYLAGGNVDR	636
		2807.80	TVLSKGLDSGTAFEILSIDIADVDSK	637
P493	35	762.46	FVFHGR	638
		964.39	DGFNNIER	639
		1363.56	GHVYNGISGGQFK	640
		1443.56	YTPTSILYFNPK	641
		1450.64	QLAEDLQKHLGAK	642
		1819.88	NHSEYVTDMLRIGR + Oxidation (M)	643
		1875.84	DLPPMEQVFDTLDLK	644
		1941.00	IRPEDMHIMANIFLPK + Oxidation (M)	645
		2081.10	RIRPEDMHIMANIFLPK	646
		2283.30	ISHLVLTRTGLYIIDSQLLK	647
P495	32			

1. Molecular weight as determined by SDS-PAGE.

2. The m/z value of a polypeptide fragment can be converted to mass by subtracting 1 from the m/z value. Each mass includes a range of plus or minus 430 parts per million (ppm) or 1 Dalton.

In this context, “sequence identity” refers to the identity between two polynucleotide sequences. Sequence identity is generally determined by aligning the bases of the two polynucleotides (for example, aligning the nucleotide sequence of the candidate sequence and a nucleotide sequence that includes, for example, the nucleotide sequence of SEQ ID NO:474 or SEQ ID NO:485) to optimize the number of identical nucleotides along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the number of shared nucleotides, although the nucleotides in each sequence must nonetheless remain in their proper order. A candidate sequence is the sequence being compared to a known sequence—e.g., a nucleotide sequence that includes the nucleotide sequence of, for example, SEQ ID NO:474 or SEQ ID NO:485. For example, two polynucleotide sequences can be compared using the Blastn program of the BLAST 2 search algorithm, as described by Tatiana et al., *FEMS Microbiol Lett.*, 1999;174: 247-250, and available on the world wide web at ncbi.nlm.nih.gov/BLAST/. The default values for all BLAST 2 search parameters may be used, including reward for match = 1, penalty for mismatch = -2, open gap penalty = 5, extension gap penalty = 2, gap x_dropoff = 50, expect = 10, wordsize = 11, and filter on.

For example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide commonly known as formate acetyltransferase (PflB). One embodiment of such a polynucleotide is reflected in SEQ ID NO:430. Variant embodiments are reflected in SEQ ID NO:431, SEQ ID NO:432, SEQ ID NO:433, SEQ ID NO:434, SEQ ID NO:435, SEQ ID NO:436, SEQ ID NO:437, SEQ ID NO:438, SEQ ID NO:439, and SEQ ID NO:440.

As another example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide commonly known as oligopeptide permease, peptide-binding protein (Opp1A). One embodiment of such a polynucleotide is reflected in SEQ ID NO:441. Variant embodiments are reflected in SEQ ID NO:442, SEQ ID NO:443, SEQ ID NO:444, SEQ ID NO:445, SEQ ID NO:446, SEQ ID NO:447, SEQ ID NO:448, SEQ ID NO:449, SEQ ID NO:450, and SEQ ID NO:451.

As another example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide commonly known as siderophore compound ABC transporter binding protein (SirA). One embodiment of such a polynucleotide is reflected in SEQ ID NO:452. Variant embodiments are
5 reflected in SEQ ID NO:453, SEQ ID NO:454, SEQ ID NO:455, SEQ ID NO:456, SEQ ID NO:457, SEQ ID NO:458, SEQ ID NO:459, SEQ ID NO:460, SEQ ID NO:461, and SEQ ID NO:462.

As another example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide referred to herein as SYN2. One
10 embodiment of such a polynucleotide is reflected in SEQ ID NO:463. Variant embodiments are reflected in SEQ ID NO:464, SEQ ID NO:465, SEQ ID NO:466, SEQ ID NO:467, SEQ ID NO:468, SEQ ID NO:469, SEQ ID NO:470, SEQ ID NO:471, SEQ ID NO:472, and SEQ ID NO:473.

As another example, a polynucleotide of the invention can include a
15 polynucleotide that encodes a polypeptide commonly known as FhuD. One embodiment of such a polynucleotide is reflected in SEQ ID NO:474. Variant embodiments are reflected in SEQ ID NO:475, SEQ ID NO:476, SEQ ID NO:477, SEQ ID NO:478, SEQ ID NO:479, SEQ ID NO:480, SEQ ID NO:481, SEQ ID NO:482, SEQ ID NO:483, and SEQ ID NO:484.

As another example, a polynucleotide of the invention can include a
20 polynucleotide that encodes a polypeptide referred to herein as SYN1. One embodiment of such a polynucleotide is reflected in SEQ ID NO:485. Variant embodiments are reflected in SEQ ID NO:486, SEQ ID NO:487, SEQ ID NO:488, SEQ ID NO:489, SEQ ID NO:490, SEQ ID NO:491, SEQ ID NO:492,
25 SEQ ID NO:493, SEQ ID NO:494, and SEQ ID NO:495.

As another example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide commonly known as MntC. One embodiment of such a polynucleotide is reflected in SEQ ID NO:496. Variant
30 embodiments are reflected in SEQ ID NO:497, SEQ ID NO:498, SEQ ID NO:499, SEQ ID NO:500, SEQ ID NO:501, SEQ ID NO:502, SEQ ID NO:503, SEQ ID NO:504, SEQ ID NO:505, and SEQ ID NO:506.

As another example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide commonly known as ferrichrome

ABC transporter lipoprotein (SstD). Embodiments of such a polynucleotide are reflected in SEQ ID NO:563, SEQ ID NO:564, SEQ ID NO:565, SEQ ID NO:566, SEQ ID NO:567, SEQ ID NO:568, SEQ ID NO:569, SEQ ID NO:570, SEQ ID NO:571, and SEQ ID NO:572.

5 As another example, a polynucleotide of the invention can include a polynucleotide that encodes a polypeptide commonly known as iron compound ABC transporter (FhuD2). Embodiments of such a polynucleotide are reflected in SEQ ID NO:573, SEQ ID NO:574, SEQ ID NO:575, SEQ ID NO:576, SEQ ID NO:577, SEQ ID NO:578, SEQ ID NO:579, SEQ ID NO:580, SEQ ID
10 NO:581, and SEQ ID NO:582.

Also provided by the present invention are whole cell preparations of a microbe, where the microbe expresses one or more of the polypeptides of the present invention. The cells present in a whole cell preparation are preferably inactivated such that the cells cannot replicate, but the immunological activity
15 of the polypeptides of the present invention expressed by the microbe is maintained. Typically, the cells are killed by exposure to agents such as glutaraldehyde, formalin, or formaldehyde.

Compositions

20 A composition of the present invention may include at least one isolated polypeptide described herein, or a number of polypeptides that is an integer greater than one (e.g., at least two, at least three, at least four). For example, a composition can include an isolated polypeptide that includes the amino acid sequence of SEQ ID NO:408 and/or an isolated polypeptide that includes the
25 amino acid sequence of SEQ ID NO:397. Unless a specific level of sequence similarity and/or identity is expressly indicated herein (e.g., at least 80% sequence similarity, at least 90% sequence identity, etc.), reference to the amino acid sequence of an identified SEQ ID NO includes variants having the levels of sequence similarity and/or the levels of sequence identity described herein in the
30 section headed "*Polypeptide sequence similarity and polypeptide sequence identity.*"

In some embodiments, the composition can include one or more additional isolated polypeptides. In some embodiments, the additional isolated

polypeptide or polypeptides may include one or more metal-regulated polypeptides. Thus, a composition can include at least one isolated metal-regulated polypeptide that includes an amino acid sequence depicted in one or more of SEQ ID NO:353 through SEQ ID NO:429 and/or one or more of SEQ ID NO:543 through SEQ ID NO:562. In addition or in the alternative, a composition can include at least one isolated metal-regulated polypeptide having a molecular weight of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa. In addition or in the alternative, a composition can include at least one metal-regulated isolated polypeptide that includes an amino acid sequence encoded by a polynucleotide that includes a nucleotide sequence depicted in one or more of SEQ ID NO:430 through SEQ ID NO:506 and/or one or more of SEQ ID NO:563 through SEQ ID NO:582.

In one embodiment, the composition includes a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:397 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID NO:407).

In another embodiment, the composition includes a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:408 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413, SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, or SEQ ID NO:418).

In another embodiment, the composition includes a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:419 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in SEQ ID NO:420, SEQ ID NO:421, SEQ ID NO:422, SEQ ID NO:423, SEQ ID NO:424, SEQ ID NO:425, SEQ ID NO:426, SEQ ID NO:427, SEQ ID NO:428, or SEQ ID NO:429).

In another embodiment, the composition includes a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:375 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in

SEQ ID NO:376, SEQ ID NO:377, SEQ ID NO:378, SEQ ID NO:379, SEQ ID NO:380, SEQ ID NO:381, SEQ ID NO:382, SEQ ID NO:383, SEQ ID NO:384, or SEQ ID NO:385).

In another embodiment, the composition includes a polypeptide that
5 includes the amino acid sequence depicted in SEQ ID NO:386 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in SEQ ID NO:387, SEQ ID NO:388, SEQ ID NO:389, SEQ ID NO:390, SEQ ID NO:391, SEQ ID NO:392, SEQ ID NO:393, SEQ ID NO:394, SEQ ID NO:395, or SEQ ID NO:396).

10 In another embodiment, the composition includes a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:364 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in SEQ ID NO:365, SEQ ID NO:366, SEQ ID NO:367, SEQ ID NO:368, SEQ ID NO:369, SEQ ID NO:370, SEQ ID NO:371, SEQ ID NO:372, SEQ ID NO:373,
15 or SEQ ID NO:374).

In another embodiment, the composition includes a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:353 (or a variant thereof such as, for example, any one of the amino acid sequences depicted in SEQ ID NO:354, SEQ ID NO:355, SEQ ID NO:356, SEQ ID NO:357, SEQ ID
20 NO:358, SEQ ID NO:359, SEQ ID NO:360, SEQ ID NO:361, SEQ ID NO:362, or SEQ ID NO:363).

In another embodiment, the composition includes a polypeptide that includes the amino acid sequence, or a variant thereof, of any one of the amino acid sequences depicted in SEQ ID NO:543, SEQ ID NO:544, SEQ ID NO:545,
25 SEQ ID NO:546, SEQ ID NO:547, SEQ ID NO:548, SEQ ID NO:549, SEQ ID NO:550, SEQ ID NO:551, or through SEQ ID NO:552.

In another embodiment, the composition includes a polypeptide that includes the amino acid sequence, or variant thereof, of any one of the amino acid sequences depicted in SEQ ID NO:553, SEQ ID NO:554, SEQ ID NO:555,
30 SEQ ID NO:556, SEQ ID NO:557, SEQ ID NO:558, SEQ ID NO:559, SEQ ID NO:560, SEQ ID NO:561, or SEQ ID NO:562.

In some embodiments, the composition can include a combination of two or more polypeptides selected from the following: a SYN1 polypeptide, a

MntC polypeptide, a FhuD polypeptide, a SYN2 polypeptide, a SirA polypeptide, an Opp1A polypeptide, and a Pflb polypeptide.

- Thus, the composition can include at least one or any combination that includes at least two, at least three, at least four, at least five, at least six, or all
- 5 seven of: a SYN1 polypeptide, a MntC polypeptide, a FhuD polypeptide, a SYN2 polypeptide, a SirA polypeptide, an Opp1A polypeptide, and a Pflb polypeptide. Exemplary compositions that include combinations of polypeptides are identified in Table 6.

<u>Composition</u>	<u>SYN1</u>	<u>MntC</u>	<u>FhuD</u>	<u>SYN2</u>	<u>SirA</u>	<u>Opp1A</u>	<u>Pflb</u>
2 pPeptides*							
1	X	X					
2	X		X				
3	X			X			
4	X				X		
5	X					X	
6	X						X
7		X	X				
8		X		X			
9		X			X		
10		X				X	
11		X					X
12			X	X			
13			X		X		
14			X			X	
15			X				X
16				X	X		
17				X		X	
18				X			X
19					X	X	
20					X		X
21						X	X
3 pPeptides*							
22	X	X	X				

23	X	X		X			
24	X	X			X		
25	X	X				X	
26	X	X					X
27	X		X	X			
28	X		X		X		
29	X		X			X	
30	X		X				X
31	X			X	X		
32	X			X		X	
33	X			X			X
34	X				X	X	
35	X				X		X
36	X					X	X
37		X	X	X			
38		X	X		X		
39		X	X			X	
40		X	X				X
41		X		X	X		
42		X		X		X	
43		X		X			X
44		X			X	X	
45		X			X		X
46		X				X	X
47			X	X	X		
48			X	X		X	
49			X	X			X
50			X		X	X	
51			X		X		X
52			X			X	X
53				X	X	X	
54				X	X		X
55				X		X	X
56					X	X	X
4 pPeptides*	X	X	X	X			

57	X	X	X		X		
58	X	X	X			X	
59	X	X	X				X
60	X	X		X	X		
61	X	X		X		X	
62	X	X		X			X
63	X	X			X		
64	X	X			X	X	
65	X	X			X		X
67	X	X				X	X
68	X		X	X	X		
69	X		X	X		X	
70	X		X	X			X
71	X		X		X	X	
72	X		X		X		X
73	X		X			X	X
74	X			X	X	X	
75	X			X	X		X
76	X				X	X	X
77		X	X	X	X		
78		X	X	X		X	
79		X	X	X			X
80		X	X		X	X	
81		X	X		X		X
82		X	X			X	X
83		X		X	X	X	
84		X		X	X		X
85		X		X		X	X
86		X			X	X	X
87			X	X	X	X	
88			X	X	X		X
89			X	X		X	X
90			X		X	X	X
91				X	X	X	X
5 pPeptides*							

92	X	X	X	X	X		
93	X	X	X	X		X	
94	X	X	X	X			X
95	X	X	X		X	X	
96	X	X	X		X		X
97	X	X	X			X	X
98	X	X		X	X	X	
99	X	X		X	X		X
100	X	X		X		X	X
101	X	X			X	X	X
102	X		X	X	X	X	
103	X		X	X	X		X
104	X		X	X		X	X
105	X		X		X	X	X
106	X			X	X	X	X
107		X	X	X	X	X	
108		X	X	X	X		X
109		X	X	X		X	X
110		X	X		X	X	X
111		X		X	X	X	X
112			X	X	X	X	X
6 pPeptides*							
113	X	X	X	X	X	X	
114	X	X	X	X	X		X
115	X	X	X	X		X	X
116	X	X	X		X	X	X
117	X	X		X	X	X	X
118	X		X	X	X	X	X
119		X	X	X	X	X	X

* pPeptides = polypeptides

"X" identifies polypeptides included in a particular composition.

- Throughout this description, a SYN1 polypeptide may be characterized
- 5 by one or more of the following: an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:408, SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413, SEQ ID

NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, or SEQ ID NO:418, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:485, SEQ ID NO:486, SEQ ID NO:487, SEQ ID NO:488, SEQ ID NO:489, SEQ ID NO:490, SEQ ID NO:491, 5 SEQ ID NO:492, SEQ ID NO:493, SEQ ID NO:494, SEQ ID NO:495, and/or a calculated molecular weight of about 33.1 kDa.

Throughout this description, an MntC polypeptide may be characterized by one or more of the following: possessing a molecular weight as determined by SDS-PAGE of 33kDa, a mass fingerprint at least 80% similar to the mass 10 fingerprint of an 33 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* ATCC isolate 19636, an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:419, SEQ ID NO:420, SEQ ID NO:421, SEQ ID NO:422, SEQ ID NO:423, SEQ ID NO:424, SEQ ID NO:425, SEQ ID NO:426, SEQ ID NO:427, SEQ ID NO:428, or SEQ ID 15 NO:429, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:496, SEQ ID NO:497, SEQ ID NO:498, SEQ ID NO:499, SEQ ID NO:500, SEQ ID NO:501, SEQ ID NO:502, SEQ ID NO:503, SEQ ID NO:504, SEQ ID NO:505, or SEQ ID NO:506, and/or a calculated molecular weight of about 34.6 kDa.

20 Throughout this description, a FhuD polypeptide may be characterized by one or more of the following: an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:397, SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID 25 NO:407, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:474, SEQ ID NO:475, SEQ ID NO:476, SEQ ID NO:477, SEQ ID NO:478, SEQ ID NO:479, SEQ ID NO:480, SEQ ID NO:481, SEQ ID NO:482, SEQ ID NO:483, or SEQ ID NO:484, and/or a calculated molecular weight of about 35.4 kDa.

30 Throughout this description, a SYN2 polypeptide may be characterized by one or more of the following: a molecular weight as determined by SDS-PAGE of 36 kDa, a mass fingerprint at least 80% similar to the mass fingerprint of an 36 kDa metal-regulated polypeptide produced by the reference strain *S.*

aureus ATCC isolate 19636, an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:386, SEQ ID NO:387, SEQ ID NO:388, SEQ ID NO:389, SEQ ID NO:390, SEQ ID NO:391, SEQ ID NO:392, SEQ ID NO:393, SEQ ID NO:394, SEQ ID NO:395, or SEQ ID NO:396, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:463, SEQ ID NO:464, SEQ ID NO:465, SEQ ID NO:466, SEQ ID NO:467, SEQ ID NO:468, SEQ ID NO:469, SEQ ID NO:470, SEQ ID NO:471, SEQ ID NO:472, or SEQ ID NO:473, and/or a calculated molecular weight of about 36.5 kDa.

Throughout this description, a SirA polypeptide may be characterized by one or more of the following: a molecular weight as determined by SDS-PAGE of 37 kDa, a mass fingerprint at least 80% similar to the mass fingerprint of an 37 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* ATCC isolate 19636, an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:375, SEQ ID NO:376, SEQ ID NO:377, SEQ ID NO:378, SEQ ID NO:379, SEQ ID NO:380, SEQ ID NO:381, SEQ ID NO:382, SEQ ID NO:383, SEQ ID NO:384, or SEQ ID NO:385, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:452, SEQ ID NO:453, SEQ ID NO:454, SEQ ID NO:455, SEQ ID NO:456, SEQ ID NO:457, SEQ ID NO:458, SEQ ID NO:459, SEQ ID NO:460, SEQ ID NO:461, or SEQ ID NO:462, and/or a calculated molecular weight of about 36.6 kDa.

Throughout this description, a Opp1A polypeptide may be characterized by one or more of the following: a molecular weight as determined by SDS-PAGE of 55 kDa, a mass fingerprint at least 80% similar to the mass fingerprint of an 55 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* ATCC isolate 19636, an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:364, SEQ ID NO:365, SEQ ID NO:366, SEQ ID NO:367, SEQ ID NO:368, SEQ ID NO:369, SEQ ID NO:370, SEQ ID NO:371, SEQ ID NO:372, SEQ ID NO:373, or SEQ ID NO:374, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:441, SEQ ID NO:442, SEQ ID NO:443, SEQ ID NO:444, SEQ ID NO:445, SEQ ID NO:446, SEQ ID NO:447,

SEQ ID NO:448, SEQ ID NO:449, SEQ ID NO:450, or SEQ ID NO:451,
and/or a calculated molecular weight of about 59.9 kDa.

Throughout this description, a PflB polypeptide may be characterized by one or more of the following: a molecular weight as determined by SDS-PAGE
5 of 88 kDa, a mass fingerprint at least 80% similar to the mass fingerprint of an 88 kDa metal-regulated polypeptide produced by the reference strain *S. aureus* ATCC isolate 19636, an amino acid sequence that includes the amino acid sequence depicted in any one of SEQ ID NO:353, SEQ ID NO:354, SEQ ID NO:355, SEQ ID NO:356, SEQ ID NO:357, SEQ ID NO:358, SEQ ID NO:359,
10 SEQ ID NO:360, SEQ ID NO:361, SEQ ID NO:362, or SEQ ID NO:363, being encoded by a polynucleotide that includes the nucleic acid sequence depicted in any one of SEQ ID NO:430, SEQ ID NO:431, SEQ ID NO:432, SEQ ID NO:433, SEQ ID NO:434, SEQ ID NO:435, SEQ ID NO:436, SEQ ID NO:437, SEQ ID NO:438, SEQ ID NO:439, or SEQ ID NO:440, and/or a calculated
15 molecular weight of about 84.7 kDa.

In another particular embodiment, the composition can include a combination of polypeptides such as, for example, a MntC polypeptide, a FluD polypeptide, a SirA polypeptide, and a SYN2 polypeptide (Composition 77 of Table 6), each polypeptide being characterized as described immediately above.

20 In some embodiments, a composition can include one or more polypeptides that are produced recombinantly. For example, a composition can include a recombinantly-produced Pflb polypeptide such as, for example, a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:353, although a recombinantly-produced Pflb polypeptide may be characterized in
25 any manner in which a Pflb polypeptide may be characterized, as described above, in addition to being recombinantly-produced. Such a composition can include one or more recombinantly-produced polypeptides, one or more polypeptides isolated from *S. aureus*, or any combination thereof.

As another example, a composition can include a recombinantly-
30 produced Opp1A polypeptide such as, for example, a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:364, although a recombinantly-produced Opp1A polypeptide may be characterized in any manner in which an Opp1A polypeptide may be characterized, as described

above, in addition to being recombinantly-produced. Such a composition can further include one or more recombinantly-produced polypeptides, one or more polypeptides isolated from *S. aureus*, or any combination thereof.

As another example, a composition can include a recombinantly-produced SirA polypeptide such as, for example, a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:375, although a recombinantly-produced SirA polypeptide may be characterized in any manner in which a SirA polypeptide may be characterized, as described above, in addition to being recombinantly-produced. Such a composition can further include one or more recombinantly-produced polypeptides, one or more polypeptides isolated from *S. aureus*, or any combination thereof.

As another example, a composition can include a recombinantly-produced SYN2 polypeptide such as, for example, a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:386, although a recombinantly-produced SYN2 polypeptide may be characterized in any manner in which a SYN2 polypeptide may be characterized, as described above, in addition to being recombinantly-produced. Such a composition can further include one or more recombinantly-produced polypeptides, one or more polypeptides isolated from *S. aureus*, or any combination thereof.

As another example, a composition can include a recombinantly-produced FhuD polypeptide such as, for example, a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:397, although a recombinantly-produced FhuD polypeptide may be characterized in any manner in which a FhuD polypeptide may be characterized, as described above, in addition to being recombinantly-produced. Such a composition can further include one or more recombinantly-produced polypeptides, one or more polypeptides isolated from *S. aureus*, or any combination thereof.

As another example, a composition can include a recombinantly-produced SYN1 polypeptide such as, for example, a polypeptide that includes the amino acid sequence depicted in SEQ ID NO:408, although a recombinantly-produced SYN1 polypeptide may be characterized in any manner in which a SYN1 polypeptide may be characterized, as described above, in addition to being recombinantly-produced. Such a composition can further

include one or more recombinantly-produced polypeptides, one or more polypeptides isolated from *S. aureus*, or any combination thereof.

As another example, a composition can include a recombinantly-produced MntC polypeptide such as, for example, a polypeptide that includes
5 the amino acid sequence depicted in SEQ ID NO:419, although a recombinantly-produced MntC polypeptide may be characterized in any manner in which a MntC polypeptide may be characterized, as described above, in addition to being recombinantly-produced. Such a composition can further include one or more recombinantly-produced polypeptides, one or more
10 polypeptides isolated from *S. aureus*, or any combination thereof.

In some embodiments, a recombinantly-produced polypeptide can represent an immunologically active fragment of the native version of the polypeptide. An immunologically active fragment may include amino acid additions to the amino terminal and/or carboxy terminal of the core (e.g., the
15 amino acid sequence of SEQ ID NO: 353, SEQ ID NO: 364, SEQ ID NO: 375, SEQ ID NO: 386, SEQ ID NO: 397, SEQ ID NO: 408, or SEQ ID NO: 419) of the immunologically active fragment. In certain embodiments, any addition to the amino terminal of the core of the immunologically active fragment can include one or more amino acid additions, deletions, substitutions (collectively,
20 “modifications”), or any combination of modification compared to longer version—e.g., wild-type or other native form—of the polypeptide. Thus, for example, in embodiments in which the immunologically active fragment includes SEQ ID NO:397, an addition to the amino terminal of SEQ ID NO:397 can include at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least
25 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, or at least 26 modifications in the amino terminal addition compared to, for example, amino acids 1-26 of SEQ ID NO:399. As another example, in embodiments in which the immunologically
30 active fragment includes SEQ ID NO:408, an addition to the amino terminal of SEQ ID NO:408 can include at least 1, at least 2, at least 3, at least 4, or at least 5 modifications in the amino terminal addition compared to, for example, amino acids 1-26 of SEQ ID NO:415.

When comparing the amino acid sequence similarity and/or amino acid sequence identity of a reference polypeptide and a candidate polypeptide of a different length (e.g., an immunologically active fragment of the reference polypeptide) the similarity and/or identity may be computed over the full length
5 of the longer polypeptide, counting each amino acid residue of greater length in the longer polypeptide contributing as a mismatch.

In some embodiments, therefore, an isolated polypeptide of the invention can include a polypeptide having at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%
10 sequence similarity and/or identity to the amino acid sequence of SEQ ID NO:397, with the proviso that if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-26 of SEQ ID NO:399.

15 In other embodiments, an isolated polypeptide of the invention can include a polypeptide having at least 98% or at least 99% sequence similarity and/or identity to the amino acid sequence of SEQ ID NO:408, with the proviso that if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional
20 amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-5 of SEQ ID NO:415.

As noted above in the section headed *Polypeptide sequence similarity and polypeptide sequence identity*, a polypeptide identified by reference to the amino acid sequence of a particular SEQ ID NO can include a polypeptide with
25 at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% amino acid sequence similarity to the reference amino acid sequence (e.g., the amino acid sequence
30 provided in a specified SEQ ID NO:) and/or a polypeptide with at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at

least 97%, at least 98%, or at least 99% amino acid sequence identity to the reference amino acid sequence (e.g., the amino acid sequence provided in a specified SEQ ID NO:).

5 A recombinantly-produced polypeptide may be expressed from a vector that permits expression of the polypeptide when the vector is introduced into an appropriate host cell. A host cell may be constructed to produce one or more recombinantly-produced polypeptides of the invention and, therefore, can include one more vectors that include at least one polynucleotide that encodes a polypeptide of the invention. Thus, each vector can include one or more
10 polynucleotides of the invention—i.e., a polynucleotide that encodes a polypeptide of the invention.

Certain compositions such as, for example, those including recombinantly-produced polypeptides, can include a maximum number of polypeptides. In some embodiments, the maximum number of polypeptides can
15 refer to the maximum total number of polypeptides. Certain compositions can include, for example, no more than 50 polypeptides such as, for example, no more than 40 polypeptides, no more than 30 polypeptides, no more than 25 polypeptides, no more than 20 polypeptides, no more than 15 polypeptides, no more than 10 polypeptides, no more than eight polypeptides, no more than
20 seven polypeptides, no more than six polypeptides, no more than five polypeptides, no more than four polypeptides, no more than three polypeptides, no more than two polypeptides, or no more than one polypeptide. In other embodiments, a maximum number of recombinantly-produced polypeptides may be specified in a similar manner. In still other embodiments, a maximum
25 number of nonrecombinantly-produced polypeptides may be specified in a similar manner.

A composition can include polypeptides isolatable from one microbe, or can be isolatable from a combination of two or more microbes. For instance, a composition can include polypeptides isolatable from two or more
30 *Staphylococcus* spp., or from a *Staphylococcus* spp. and a different microbe that is not a member of the genus *Staphylococcus*. The present invention also provides compositions including a whole cell preparation, where the whole cell expresses one or more of the polypeptides of the present invention. For

instance, the whole cell can be a *Staphylococcus* spp. In some aspects, a composition can include whole preparations from two, three, four, five, or six strains.

Optionally, a polypeptide of the present invention can be covalently
5 bound or conjugated to a carrier polypeptide to improve the immunological properties of the polypeptide. Useful carrier polypeptides are known in the art. The chemical coupling of polypeptides of the present invention can be carried out using known and routine methods. For instance, various homobifunctional and/or heterobifunctional cross-linker reagents such as bis(sulfosuccinimidyl) suberate, bis(diazobenzidine), dimethyl adipimide, dimethyl pimelimide,
10 dimethyl superimide, disuccinimidyl suberate, glutaraldehyde, *m*-maleimidobenzoyl-*N*-hydroxysuccinimide, sulfo-*m*-maleimidobenzoyl-*N*-hydroxysuccinimide, sulfosuccinimidyl 4-(*N*-maleimidomethyl) cycloheane-1-carboxylate, sulfosuccinimidyl 4-(*p*-maleimido-phenyl) butyrate and (1-ethyl-3-
15 (dimethyl-aminopropyl) carbodiimide can be used (see, for instance, Harlow and Lane, Antibodies, A Laboratory Manual, generally and Chapter 5, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, NY (1988)).

The compositions of the present invention optionally further include a pharmaceutically acceptable carrier. "Pharmaceutically acceptable" refers to a
20 diluent, carrier, excipient, salt, etc, that is compatible with the other ingredients of the composition, and not deleterious to the recipient thereof. Typically, the composition includes a pharmaceutically acceptable carrier when the composition is used as described herein. The compositions of the present invention may be formulated in pharmaceutical preparations in a variety of
25 forms adapted to the chosen route of administration, including routes suitable for stimulating an immune response to an antigen. Thus, a composition of the present invention can be administered via known routes including, for example, oral; parenteral including intradermal, transcutaneous and subcutaneous; intramuscular, intravenous, intraperitoneal, etc. and topically, such as,
30 intranasal, intrapulmonary, intramammary, intravaginal, intrauterine, intradermal, transcutaneous and rectally, etc. It is foreseen that a composition can be administered to a mucosal surface, such as by administration to the nasal or respiratory mucosa (e.g., via a spray or aerosol), in order to stimulate

mucosal immunity, such as production of secretory IgA antibodies, throughout the animal's body.

A composition of the present invention can also be administered via a sustained or delayed release implant. Implants suitable for use according to the invention are known and include, for example, those disclosed in Emery and Straub (WO 01/37810 (2001)), and Emery *et al.*, (WO 96/01620 (1996)). Implants can be produced at sizes small enough to be administered by aerosol or spray. Implants also can include nanospheres and microspheres.

A composition of the present invention may be administered in an amount sufficient to treat certain conditions as described herein. The amount of polypeptides or whole cells present in a composition of the present invention can vary. For instance, the dosage of polypeptides can be between 0.01 micrograms (μg) and 300 mg, typically between 0.1 mg and 10 mg. When the composition is a whole cell preparation, the cells can be present at a concentration of, for instance, 10^2 bacteria/ml, 10^3 bacteria/ml, 10^4 bacteria/ml, 10^5 bacteria/ml, 10^6 bacteria/ml, 10^7 bacteria/ml, 10^8 bacteria/ml, or 10^9 bacteria/ml. For an injectable composition (e.g. subcutaneous, intramuscular, etc.) the polypeptides may be present in the composition in an amount such that the total volume of the composition administered is 0.5 ml to 5.0 ml, typically 1.0 to 2.0 ml. When the composition is a whole cell preparation, the cells are preferably present in the composition in an amount that the total volume of the composition administered is 0.5 ml to 5.0 ml, typically 1.0 to 2.0 ml. The amount administered will vary depending on various factors including, but not limited to, the specific polypeptides chosen, the weight, physical condition and age of the animal, and the route of administration. Thus, the absolute weight of the polypeptide included in a given unit dosage form can vary widely, and depends upon factors such as the species, age, weight and physical condition of the animal, as well as the method of administration. Such factors can be determined by one of skill in the art. Other examples of dosages suitable for the invention are disclosed in Emery *et al.*, (U.S. Patent 6,027,736).

The formulations may be conveniently presented in unit dosage form and may be prepared by methods well known in the art of pharmacy. Methods of preparing a composition with a pharmaceutically acceptable carrier include

the step of bringing the active compound (e.g., a polypeptide or whole cell of the present invention) into association with a carrier that constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active compound into association with a liquid
5 carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product into the desired formulations.

A composition including a pharmaceutically acceptable carrier can also include an adjuvant. An "adjuvant" refers to an agent that can act in a nonspecific manner to enhance an immune response to a particular antigen, thus
10 potentially reducing the quantity of antigen necessary in any given immunizing composition, and/or the frequency of injection necessary in order to generate an adequate immune response to the antigen of interest. Adjuvants may include, for example, IL-1, IL-2, emulsifiers, muramyl dipeptides, dimethyl dioctadecyl ammonium bromide (DDA), avridine, aluminum hydroxide, oils, saponins,
15 alpha-tocopherol, polysaccharides, emulsified paraffins (including, for instance, those available from under the tradename EMULSIGEN from MVP Laboratories, Ralston, Nebraska), ISA-70, RIBI and other substances known in the art. It is expected that polypeptides of the present invention will have immunoregulatory activity and that such polypeptides may be used as adjuvants
20 that directly act as T and/or B cell activators or act on specific cell types that enhance the synthesis of various cytokines or activate intracellular signaling pathways. Such polypeptides are expected to augment the immune response to increase the protective index of the existing composition.

In another embodiment, a composition of the invention including a
25 pharmaceutically acceptable carrier can include a biological response modifier, such as, for example, IL-2, IL-4 and/or IL-6, TNF, IFN-alpha, IFN-gamma, and other cytokines that effect immune cells. An immunizing composition can also include other components known in the art such as an antibiotic, a preservative, an anti-oxidant, or a chelating agent.

30

Methods of Making

The present invention also provides methods for obtaining the polypeptides described herein. The polypeptides and whole cells of the present invention may be isolatable from a member of the family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *Staphylococcus aureus*. Other gram positive microbes from which polypeptides can be isolated include *Corynebacterium* spp., *Erysipelothrix* spp., *Mycobacterium* spp., and *Erysipelothrix* spp. Microbes useful for obtaining polypeptides of the present invention and making whole cell preparations are commercially available from a depository such as American Type Culture Collection (ATCC). In addition, such microbes are readily obtainable by techniques routine and known to the art. The microbes may be derived from an infected animal as a field isolate, and used to obtain polypeptides and/or whole cell preparations of the present invention, or stored for future use, for example, in a frozen repository at -20°C to -95°C, or -40°C to -50°C, in bacteriological media containing 20% glycerol, and other like media.

When a polypeptide of the present invention is to be obtained from a microbe, the microbe can be incubated under low metal conditions. As used herein, the phrase "low metal conditions" refers to an environment, typically bacteriological media, which contains amounts of a free metal that cause a microbe to express metal-regulated polypeptides at a detectable level. As used herein, the phrase "high metal conditions" refers to an environment that contains amounts of a free metal that cause a microbe to either not express one or more of the metal-regulated polypeptides described herein at a detectable level, or to express such a polypeptide at a decreased level compared to expression of the metal-regulated polypeptide under low metal conditions. In some cases, "high metal conditions" can include a metal-rich natural environment and/or culture in a metal-rich medium without a metal chelator. In contrast, in some cases, "low metal conditions" can include culture in a medium that includes a metal chelator, as described in more detail below. Metals are those present in the periodic table under Groups 1 through 17 (IUPAC notation; also referred to as Groups I-A, II-A, III-B, IV-B, V-B, VI-B, VII-B, VIII, I-B, II-B, III-A, IV-A, V-A, VI-A, and VII-A, respectively, under CAS notation).

Preferably, metals are those in Groups 2 through 12, more preferably, Groups 3-12. Even more preferably, the metal is iron, zinc, copper, magnesium, nickel, cobalt, manganese, molybdenum, or selenium, most preferably, iron.

Low metal conditions are generally the result of the addition of a metal
5 chelating compound to a bacteriological medium, the use of a bacteriological medium that contains low amounts of a metal, or the combination thereof. High metal conditions are generally present when a chelator is not present in the medium, a metal is added to the medium, or the combination thereof. Examples of metal chelators include natural and synthetic compounds. Examples of
10 natural compounds include plant phenolic compounds, such as flavenoids. Examples of flavenoids include the copper chelators catechin and naringenin, and the iron chelators myricetin and quercetin. Examples of synthetic copper chelators include, for instance, tetrathiomolybdate, and examples of synthetic zinc chelators include, for instance, N,N,N',N' -Tetrakis (2-pyridylmethyl)-
15 ethylene diamine. Examples of synthetic iron chelators include 2,2'-dipyridyl (also referred to in the art as α,α' -bipyridyl), 8-hydroxyquinoline, ethylenediamine-di-O-hydroxyphenylacetic acid (EDDHA), desferrioxamine methanesulphonate (desferol), transferrin, lactoferrin, ovotransferrin, biological siderophores, such as, the catecholates and hydroxamates, and citrate. An
20 example of a general divalent cation chelator is CHELEX resin. Preferably, 2,2'-dipyridyl is used for the chelation of iron. Typically, 2,2'-dipyridyl is added to the media at a concentration of at least 300 micrograms/milliliter ($\mu\text{g/ml}$), at least 600 $\mu\text{g/ml}$, or at least 900 $\mu\text{g/ml}$. High levels of 2,2'-dipyridyl can be 1200 $\mu\text{g/ml}$, 1500 $\mu\text{g/ml}$, or 1800 $\mu\text{g/ml}$.

25 The *S. aureus* genome encodes three Fur homologs: Fur, PerR, and Zur. While the Zur and PerR proteins appear to be primarily involved in regulating zinc homeostasis and peroxide stress genes, respectively, the Fur protein has been demonstrated to regulate several iron-siderophore uptake systems in response to iron limitation. The Fur protein also plays a role in
30 oxidative stress resistance and virulence. It is expected that a gram positive organism, preferably, an *S. aureus*, with a mutation in a *fur* gene will result in the constitutive expression of many, if not all, of the metal-regulated polypeptides of the present invention. The production of a *fur* mutation in a

gram positive, preferably, an *S. aureus*, can be produced using routine methods including, for instance, transposon, chemical, or site-directed mutagenesis useful for generating gene knock-out mutations in gram positive bacteria.

5 The medium used to incubate the microbe and the volume of media used to incubate the microbe can vary. When a microbe is being evaluated for the ability to produce one or more of the polypeptides described herein, the microbe can be grown in a suitable volume, for instance, 10 milliliters to 1 liter of medium. When a microbe is being grown to obtain polypeptides for use in, for instance, administration to animals, the microbe may be grown in a fermentor to
10 allow the isolation of larger amounts of polypeptides. Methods for growing microbes in a fermentor are routine and known to the art. The conditions used for growing a microbe preferably include a metal chelator, more preferably an iron chelator, for instance 2,2'-dipyridyl, a pH of between 6.5 and 7.5, preferably between 6.9 and 7.1, and a temperature of 37°C.

15 In some aspects of the invention, a microbe may be harvested after growth. Harvesting includes concentrating the microbe into a smaller volume and suspending in a media different than the growth media. Methods for concentrating a microbe are routine and known in the art, and include, for example, filtration or centrifugation. Typically, the concentrated microbe is
20 suspended in an appropriate buffer. An example of a buffer that can be used contains Tris-base (7.3 grams /liter), at a pH of 8.5. Optionally, the final buffer also minimizes proteolytic degradation. This can be accomplished by having the final buffer at a pH of greater than 8.0, preferably, at least 8.5, and/or including one or more proteinase inhibitors (e.g., phenylmethanesulfonyl
25 fluoride). Optionally and preferably, the concentrated microbe is frozen at -20°C or below until disrupted.

When the microbe is to be used as a whole cell preparation, the harvested cells may be processed using routine and known methods to inactivate the cells. Alternatively, when a microbe is to be used to prepare polypeptides of
30 the present invention, the microbe may be disrupted using chemical, physical, or mechanical methods routine and known to the art, including, for example, boiling, french press, sonication, digestion of peptidoglycan (for instance, by digestion with lysozyme), or homogenization. An example of a suitable device

useful for homogenization is a model C500-B AVESTIN homogenizer, (Avestin Inc, Ottawa Canada). As used herein, "disruption" refers to the breaking up of the cell. Disruption of a microbe can be measured by methods that are routine and known to the art, including, for instance, changes in optical density. Typically, a microbe is subjected to disruption until the percent transmittance is increased by 20% when a 1:100 dilution is measured. When physical or mechanical methods are used, the temperature during disruption is typically kept low, preferably at 4°C, to further minimize proteolytic degradation. When chemical methods are used the temperature may be increased to optimize for the cell disruption. A combination of chemical, physical, and mechanical methods may also be used to solubilize the cell wall of microbe. As used herein, the term "solubilize" refers to dissolving cellular materials (e.g., polypeptides, nucleic acids, carbohydrates) into the aqueous phase of the buffer in which the microbe was disrupted, and the formation of aggregates of insoluble cellular materials. Without intending to be limited by theory, the conditions for solubilization are believed to result in the aggregation of polypeptides of the present invention into insoluble aggregates that are large enough to allow easy isolation by, for instance, centrifugation.

The insoluble aggregates that include one or more of the polypeptides of the present invention may be isolated by methods that are routine and known to the art. Preferably, the insoluble aggregates are isolated by centrifugation. Typically, centrifugation of polypeptides, such as membrane polypeptides, can be accomplished by centrifugal forces of 100,000 x g. The use of such centrifugal forces requires the use of ultracentrifuges, and scale-up to process large volumes of sample is often difficult and not economical with these types of centrifuges. The methods described herein provide for the production of insoluble aggregates large enough to allow the use of continuous flow centrifuges, for instance T-1 Sharples (Alfa Laval Separations, Warminster, PA), which can be used with a flow rate of 250 ml/minute at 17 psi at a centrifugal force of 46,000 x g to 60,000 x g. Other large scale centrifuges can be used, such as the tubular bowl, chamber, and disc configurations. Such centrifuges are routinely used and known in the art, and are commercially available from such manufactures as Pennwalt, Westfalia and alpha-Laval.

The final harvested proteins are washed and/or dialyzed against an appropriate buffer using methods known in the art, for instance diafiltration, precipitation, hydrophobic chromatography, ion-exchange chromatography, or affinity chromatography, or ultra filtration and washing the polypeptides, for instance, in alcohol, by diafiltration. After isolation, the polypeptides
5 suspended in buffer and stored at low temperature, for instance, -20°C or below.

In those aspects of the present invention where a whole cell preparation is to be made, after growth a microbe can be killed with the addition of an agent such as glutaraldehyde, formalin, or formaldehyde, at a concentration sufficient
10 to inactivate the cells in the culture. For instance, formalin can be added at a concentration of 0.3% (vol:vol). After a period of time sufficient to inactivate the cells, the cells can be harvested by, for instance, diafiltration and/or centrifugation, and washed.

In other aspects, an isolated polypeptide of the invention may be prepared recombinantly. When prepared recombinantly, a polynucleotide encoding the polypeptide may be identified and cloned into an appropriate expression host as described below in Example 14. The recombinant expression host may be grown in an appropriate medium, disrupted, and the polypeptides isolated as described above.

20

Methods of Use

An aspect of the present invention is further directed to methods of using the compositions of the present invention. The methods include administering to an animal an effective amount of a composition of the present invention. The
25 animal can be, for instance, avian (including, for instance, chickens or turkeys), bovine (including, for instance, cattle), caprine (including, for instance, goats), ovine (including, for instance, sheep), porcine (including, for instance, swine), bison (including, for instance, buffalo), equine (including, for instance, horses), a companion animal (including, for instance, dogs or cats), members of the
30 family Cervidae (including, for instance, deer, elk, moose, caribou and reindeer), or human.

In some aspects, the methods may further include additional administrations (e.g., one or more booster administrations) of the composition to

the animal to enhance or stimulate a secondary immune response. A booster can be administered at a time after the first administration, for instance, one to eight weeks, preferably two to four weeks, after the first administration of the composition. Subsequent boosters can be administered one, two, three, four, or
5 more times annually. Without intending to be limited by theory, it is expected that in some aspects of the present invention annual boosters will not be necessary, as an animal will be challenged in the field by exposure to microbes expressing polypeptides present in the compositions having epitopes that are identical to or structurally related to epitopes present on polypeptides of the
10 composition administered to the animal.

In one aspect, the invention is directed to methods for making antibodies, for instance by inducing the production of antibody in an animal, or by recombinant techniques. The antibody produced includes antibody that specifically binds at least one polypeptide present in the composition. In this
15 aspect of the invention, an "effective amount" is an amount effective to result in the production of antibody in the animal. Methods for determining whether an animal has produced antibodies that specifically bind polypeptides present in a composition of the present invention can be determined as described herein. The present invention further includes antibody that specifically bind to a
20 polypeptide of the present invention, and compositions including such antibodies.

The method may be used to produce antibody that specifically binds polypeptides expressed by a microbe other than the microbe from which the polypeptides of the composition were isolated. As used herein, an antibody that
25 can "specifically bind" a polypeptide is an antibody that interacts with the epitope of the antigen that induced the synthesis of the antibody, or interacts with a structurally related epitope. At least some of the polypeptides present in the compositions of the present invention typically include epitopes that are conserved in the polypeptides of different species and different genera of
30 microbes. Accordingly, antibody produced using a composition derived from one microbe is expected to bind to polypeptides expressed by other microbes and provide broad spectrum protection against gram positive organisms. Examples of gram positive microbes to which the antibody may specifically

bind are Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *Staphylococcus aureus*; members of the family Streptococcaceae, preferably, *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*, *Streptococcus uberis*, *Streptococcus bovis*, *Streptococcus equi*, or *Streptococcus dysgalactiae*; and *Bacillus* spp., *Clostridium* spp., *Corynebacterium* spp.,
5 *Enterococcus* spp., *Erysipelothrix* spp., *Listeria* spp., *Micrococcus* spp., and *Mycobacterium* spp., *Kytococcus* spp., and *Erysipelothrix* spp. Therefore, antibody produced using a composition of polypeptides of the invention may be used to identify and characterize polypeptides of the invention independent of
10 the origin, source, and/or manner of obtaining the polypeptide.

The present invention is also directed to the use of such antibody to target a microbe expressing a polypeptide of the present invention or a polypeptide having an epitope structurally related to an epitope present on a polypeptide of the present invention. A compound can be covalently bound to
15 an antibody, where the compound can be, for instance, a toxin. Likewise, such compounds can be covalently bound to a bacterial siderophore to target the microbe. The chemical coupling or conjugation of an antibody of the present invention, or a portion thereof (such as an Fab fragment), can be carried out using known and routine methods.

20 In one aspect the invention is also directed to treating an infection in an animal, including a human, caused by a gram positive microbe, preferably by a member of the family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *S. aureus*; members of the family Streptococcaceae, preferably, *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*,
25 *Streptococcus uberis*, *Streptococcus bovis*, *Streptococcus equi*, or *Streptococcus dysgalactiae*; *Bacillus* spp., *Clostridium* spp., *Corynebacterium* spp., *Enterococcus* spp., *Erysipelothrix* spp., *Kytococcus* spp., *Listeria* spp., *Micrococcus* spp., *Mycobacterium* spp., and *Erysipelothrix* spp. As used herein, the term "infection" refers to the presence of a gram positive microbe in an
30 animal's body, which may or may not be clinically apparent. An animal with an infection by a member of the genus *Staphylococcus* that is not clinically apparent is often referred to as an asymptomatic carrier.

Treating an infection can be prophylactic or, alternatively, can be initiated after the animal is infected by the microbe. Treatment that is prophylactic—e.g., initiated before a subject is infected by a microbe or while any infection remains subclinical—is referred to herein as treatment of a subject
5 that is “at risk” of infection. As used herein, the term “at risk” refers to an animal that may or may not actually possess the described risk. Thus, typically, an animal “at risk” of infection by a microbe is an animal present in an area where animals have been identified as infected by the microbe and/or is likely to be exposed to the microbe even if the animal has not yet manifested any
10 detectable indication of infection by the microbe and regardless of whether the animal may harbor a subclinical amount of the microbe. Accordingly, administration of a composition can be performed before, during, or after the animal has first contact with the microbe. Treatment initiated after the animal’s first contact with the microbe may result in decreasing the severity of symptoms
15 and/or clinical signs of infection by the microbe, completely removing the microbe, and/or decreasing the likelihood of experiencing a clinically evident infection compared to an animal to which the composition is not administered. The method includes administering an effective amount of the composition of the present invention to an animal having, or at risk of having, an infection
20 caused by a gram positive microbe, and determining whether the number of microbes causing the infection has decreased. In this aspect of the invention, an “effective amount” is an amount effective to reduce the number of the specified microbes in an animal or reduce the likelihood that the animal experiences a clinically-evident infection compared to an animal to which the composition is
25 not administered. Methods for determining whether an infection is caused by a gram positive microbe are routine and known in the art, as are methods for determining whether the infection has decreased.

In another aspect, the present invention is directed to methods for treating one or more symptoms or clinical signs of certain conditions in an
30 animal that may be caused by infection by a gram positive microbe, preferably by a member of the family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *S. aureus*; members of the family Streptococcaceae, preferably, *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Streptococcus*

agalactiae, *Streptococcus uberis*, *Streptococcus bovis*, *Streptococcus equi*, or *Streptococcus dysgalactiae*; *Bacillus* spp., *Clostridium* spp., *Corynebacterium* spp., *Enterococcus* spp., *Erysipelothrix* spp., *Kytococcus* spp., *Listeria* spp., *Micrococcus* spp., *Mycobacterium* spp., and *Erysipelothrix* spp. The method

5 includes administering an effective amount of a composition of the present invention to an animal having or at risk of having a condition, or exhibiting symptoms and/or clinical signs of a condition, and determining whether at least one symptom and/or clinical sign of the condition is changed, preferably, reduced. Examples of conditions and/or clinical signs caused by microbial

10 infections include, for instance, mastitis, septicemia, pneumonia, meningoencephalitis, lymphangitis, dermatitis, genital tract infections, strangles, metritis, perinatal disease, pituitary abscesses, arthritis, bursitis, orchitis, cystitis and pyelonephritis, caseous lymphadenitis, tuberculosis, ulcerative lymphangitis, listeriosis, erysipelas, laminitis, anthrax, tyzzer's disease, tetanus,

15 botulism, enteritis, malignant edema, braxy, bacillary hemoglobinuria, enterotoxemia, necrotic skin lesions, and nosocomial infections. Examples of conditions caused by *S. aureus* also include, for instance, botryomycosis in horses, purulent synovitis and osteomyelitis in poultry, abortions in swine, and tick pyemia in lambs. Examples of conditions caused by *Streptococcus* spp.

20 also include, for instance, sore throat, scarlet fever, impetigo, ulcerative endocarditis, rheumatic fever and post streptococcal glomerulonephritis cervicitis in humans, cervicitis in equine and swine, and meningitis and jowl abscesses in swine.

Treatment of symptoms and/or clinical signs associated with these

25 conditions can be prophylactic or, alternatively, can be initiated after the development of a condition described herein. As used herein, the term "symptom" refers to subjective evidence of disease or condition experienced by the patient and caused by infection by a microbe. As used herein, the term "clinical sign" or, simply, "sign" refers to objective evidence of disease or

30 condition caused by infection by a microbe. Symptoms and/or clinical signs associated with conditions referred to herein and the evaluations of such symptoms are routine and known in the art. Treatment that is prophylactic, for instance, initiated before a subject manifests symptoms or signs of a condition

caused by a microbe, is referred to herein as treatment of a subject that is “at risk” of developing the condition. Thus, typically, an animal “at risk” of developing a condition is an animal present in an area where animals having the condition have been diagnosed and/or is likely to be exposed to a microbe causing the condition even if the animal has not yet manifested symptoms or signs of any condition caused by the microbe. Accordingly, administration of a composition can be performed before, during, or after the occurrence of the conditions described herein. Treatment initiated after the development of a condition may result in decreasing the severity of the symptoms of one of the conditions, or completely removing the symptoms. In this aspect of the invention, an “effective amount” is an amount effective to prevent the manifestation of symptoms of a disease, decrease the severity of the symptoms of a disease, and/or completely remove the symptoms. The successful treatment of a gram positive microbial infection in an animal is disclosed in Example 5, which demonstrates the protection against disease caused by *S. aureus* in mouse models by administering a composition of the present invention. These mouse models are a commonly accepted model for the study of human disease caused by these microbes. The successful treatment of a gram positive microbial infection in an animal is also disclosed in Examples 10 to 12, which demonstrate that administering a composition of the present invention provides protection against disease caused by *S. aureus* in cows.

The present invention also provides methods for decreasing colonization by gram positive microbes, for instance blocking the attachment sites of gram positive microbe, including tissues of the skeletal system (for instance, bones, cartilage, tendons and ligaments), muscular system, (for instance, skeletal and smooth muscles), circulatory system (for instance, heart, blood vessels, capillaries and blood), nervous system (for instance, brain, spinal cord, and peripheral nerves), respiratory system (for instance, nose, trachea lungs, bronchi, bronchioles, alveoli), digestive system (for instance, mouth, salivary glands oesophagus liver stomach large and small intestine), excretory system (for instance, kidneys, ureters, bladder and urethra), endocrine system (for instance, hypothalamus, pituitary, thyroid, pancreas and adrenal glands), reproductive system (for instance, ovaries, oviduct, uterus, vagina, mammary

glands, testes, and seminal vesicles), lymphatic/immune systems (for instance, lymph, lymph nodes and vessels, mononuclear or white blood cells, such as macrophages, neutrophils, monocytes, eosinophils, basophils, and lymphocytes, including T cells and B cells), and specific cell lineages (for instance, precursor
5 cells, epithelial cells, stem cells), and the like. Preferably, the gram positive microbe is a member of the family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *S. aureus*; a member of the family Streptococcaceae, preferably, *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*, *Streptococcus uberis*, *Streptococcus bovis*, *Streptococcus equi*, or
10 *Streptococcus dysgalactiae*; *Bacillus* spp., *Clostridium* spp., *Corynebacterium* spp., *Enterococcus* spp., *Erysipelothrix* spp., *Kytococcus* spp., *Listeria* spp., *Micrococcus* spp., *Mycobacterium* spp., and *Erysipelothrix* spp.

Decreasing colonization in an animal may be performed prophylactically or, alternatively, can be initiated after the animal is colonized by the microbe.
15 Treatment that is prophylactic—e.g., initiated before a subject is colonized by a microbe or while any colonization remains undetected—is referred to herein as treatment of a subject that is “at risk” of colonization by the microbe. Thus, typically, an animal “at risk” of colonization by a microbe is an animal present in an area where animals have been identified as colonized by the microbe
20 and/or is likely to be exposed to the microbe even if the animal has not yet manifested any detectable indication of colonization by the microbe and regardless of whether the animal may harbor a subcolonization number of the microbe. Accordingly, administration of a composition can be performed before, during, or after the animal has first contact with the microbe. Treatment
25 initiated after the animal’s first contact with the microbe may result in decreasing the extent of colonization by the microbe, completely removing the microbe, and/or decreasing the likelihood that the animal becomes colonized by the microbe compared to an animal to which the composition is not administered. Thus, the method includes administering an effective amount of a
30 composition of the present invention to an animal colonized by, or at risk of being colonized by, a gram positive microbe. In this aspect of the invention, an “effective amount” is an amount sufficient to decrease colonization of the animal by the microbe, where decreasing colonization refers to one or more of:

decreasing the extent of colonization by the microbe, completely removing the microbe, and/or decreasing the likelihood that the animal becomes colonized by the microbe compared to an animal to which the composition is not administered. Methods for evaluating the colonization of an animal by a
5 microbe are routine and known in the art. For instance, colonization of an animal's intestinal tract by a microbe can be determined by measuring the presence of the microbe in the animal's feces. It is expected that decreasing the colonization of an animal by a microbe will reduce transmission of the microbe to humans.

10 A composition of the invention can be used to provide for active or passive immunization against bacterial infection. Generally, the composition can be administered to an animal to provide active immunization. However, the composition can also be used to induce production of immune products, such as antibodies, which can be collected from the producing animal and administered
15 to another animal to provide passive immunity. Immune components, such as antibodies, can be collected to prepare compositions (preferably containing antibody) from serum, plasma, blood, colostrum, etc. for passive immunization therapies. Antibody compositions including monoclonal antibodies and/or anti-idiotypes can also be prepared using known methods. Chimeric antibodies
20 include human-derived constant regions of both heavy and light chains and murine-derived variable regions that are antigen-specific (Morrison et al., Proc. Natl. Acad. Sci. USA, 1984, 81(21):6851-5; LoBuglio et al., Proc. Natl. Acad. Sci. USA, 1989, 86(11):4220-4; Boulianne et al., Nature, 1984, 312(5995):643-6.). Humanized antibodies substitute the murine constant and framework (FR)
25 (of the variable region) with the human counterparts (Jones et al., Nature, 1986, 321(6069):522-5; Riechmann et al., Nature, 1988, 332(6162):323-7; Verhoeyen et al., Science, 1988, 239(4847):1534-6; Queen et al., Proc. Natl. Acad. Sci. USA, 1989, 86(24):10029-33; Daugherty et al., Nucleic Acids Res., 1991, 19(9): 2471-6.). Alternatively, certain mouse strains can be used that have been
30 genetically engineered to produce antibodies that are almost completely of human origin; following immunization the B cells of these mice are harvested and immortalized for the production of human monoclonal antibodies (Bruggeman and Taussig, Curr. Opin. Biotechnol., 1997, 8(4):455-8; Lonberg

and Huszar, Int. Rev. Immunol., 1995;13(1):65-93; Lonberg et al., Nature, 1994, 368:856-9; Taylor et al., Nucleic Acids Res., 1992, 20:6287-95.). Passive antibody compositions and fragments thereof, e.g., scFv, Fab, F(ab')₂ or Fv or other modified forms thereof, may be administered to a recipient in the form of serum, plasma, blood, colostrum, and the like. However, the antibodies may also be isolated from serum, plasma, blood, colostrum, and the like, using known methods for later use in a concentrated or reconstituted form such as, for instance, lavage solutions, impregnated dressings and/or topical agents and the like. Passive immunization preparations may be particularly advantageous for the treatment of acute systemic illness, or passive immunization of young animals that failed to receive adequate levels of passive immunity through maternal colostrum. Antibodies useful for passive immunization may also be useful to conjugate to various drugs or antibiotics that could be directly targeted to bacteria expressing during a systemic or localized infection a polypeptide of the present invention or a polypeptide having an epitope structurally related to an epitope present on a polypeptide of the present invention.

Animal models, in particular mouse models, are available for experimentally evaluating the compositions of the present invention. These mouse models are commonly accepted models for the study of human disease caused by members of the genus *Staphylococcus*, and *S. aureus* in particular. In those cases where a member of the genus *Staphylococcus* causes disease in an animal, for instance a cow, the natural host can be used to experimentally evaluate the compositions of the present invention.

However, protection in a mouse model is not the only way to assess whether a composition can confer protection to an animal against infection by a *Staphylococcus* spp. The adaptive immune response consists of two primary divisions: the humoral (antibody) response and the cellular (T cell) response. Following infection by a bacterial pathogen, dendritic cells at the infection site encounter microbial antigens and produce signaling molecules such as, for example, surface receptors and cytokines in response to conserved molecular patterns associated with the specific bacterium. These signals are shaped by the nature of the pathogen and ideally lead to the appropriate antibody and T cell responses that protect the host from disease. While some bacterial diseases are

controlled primarily through antibody functions, others require T cell responses or both antibody and T cell responses for protection. The goal of vaccine biology is to identify the immune responses that provide protection and then design a vaccine to reproduce one or more of these responses in humans.

5 Antibodies can have many different functions in the conferring protection against infection such as, for example, complement fixation, opsonization, neutralization, and/or agglutination. Moreover, some subclasses of antibodies are better than others at specific functions; for example, for complement fixation the following hierarchy exists for human IgG subclasses:

10 IgG3>IgG1>IgG2>IgG4).

 Antibody immunological functions can be studied in a variety of ways. For instance, Western blots are used to identify antigen-specific binding based on size of separated proteins, while the standard enzyme-linked immunosorbent assay (ELISA) is used to produce quantitative information about antibody titers
15 within serum. Antibody surface binding studies are used to determine whether antibody in serum are able to recognize antigens on the surface of intact bacteria, an important indicator of whether the antibodies have the potential to work *in vivo*. Thus, one skilled in the art recognizes that antibody binding assays such as a Western blot, ELISA (e.g., using human antisera), and/or surface binding
20 correlate positively with the specifically-bound antigens providing immunological activity against infection by *Staphylococcus* spp. (Vytvytska *et al.* 2002, *Proteomics* 2:580-590; Kuklin *et al.* 2006, *Infect. Immun.* 74(4):2215-2223; Dryla *et al.* 2005, *Clin. Diag. Lab. Immunol.* 12(3):387-398). However, one skilled in the art further recognizes that a lack of antibody binding in an
25 assay such as, for example, a Western blot, ELISA, or surface binding assay does not mean that the assayed antigen fails to provide immunological activity against infection by *Staphylococcus* spp. (Kim HK *et al.* IsdA and IsdB antibodies protect mice against *Staphylococcus aureus* abscess formation and lethal challenge. *Vaccine* (2010), doi:10.1016/j.vaccine.2010.02.097).

30 Figure 202 shows that convalescent mouse serum binds to at least recombinantly-produced MntC (SEQ ID NO:419), recombinantly-produced SYN2 (SEQ ID NO:386), recombinantly-produced SirA (SEQ ID NO:375), and recombinantly-produced Opp1A (SEQ ID NO:364), indicating that each of the

bound recombinantly-produced polypeptides can induce immunological activity against infection by *Staphylococcus* spp.

Figure 204 shows that convalescent human serum binds to recombinantly-produced PflB (SEQ ID NO:353), recombinantly-produced
 5 Opp1A (SEQ ID NO:364), recombinantly-produced SirA (SEQ ID NO:375), recombinantly-produced SYN2 (SEQ ID NO:386), recombinantly-produced FhuD (SEQ ID NO:397), recombinantly-produced SYN1 (SEQ ID NO:408), and recombinantly-produced MntC (SEQ ID NO:419), indicating that each of the recombinantly-produced polypeptides can induce immunological activity against
 10 infection by *Staphylococcus* spp.

Figure 205 shows that antibody raised against recombinantly-produced FhuD (SEQ ID NO:397), recombinantly-produced Opp1A (SEQ ID NO:364), and recombinantly-produced PflB (SEQ ID NO:353) binds to the surface of *Staphylococcus* spp. cells, indicating that each of the polypeptide targets of the
 15 cell-binding antibody can induce immunological activity against infection by *Staphylococcus* spp.

Techniques such as opsonophagocytosis assays (OPA), in which antibody and complement-bound bacteria are combined with human or mouse phagocytes to determine levels of bacterial killing, are useful for studying
 20 antibody function. Positive OPA results correlate with vaccine-induced protection in a mouse model. (Stranger-Jones *et al.* 2006, *Proc. Natl. Acad. Sci.* 103(45):16942-16947). A similar oxidative burst assay can be used to assess the level of reactive oxygen species (ROS) by fresh human or mouse neutrophils following interaction with antibody and complement-bound bacteria.

25 In some cases, one can determine that a candidate polypeptide possesses cell-mediated immunological activity and, therefore, the candidate polypeptide may exhibit immunological activity in the absence of inducing the production of antibodies. (Spellberg *et al.* 2008, *Infect. Immun.* 76(10):4575-4580). Cytotoxic or CD8 T cells primarily kill infected cells directly through various effector
 30 mechanisms, while helper CD4 T cells function to provide important signaling in the way of cytokines. These T cell classes can be further subdivided based on the cytokines they produce, and different subclasses are effective against different bacterial pathogens. T cells are often studied by assessing their

- phenotypes with flow cytometry, where antibodies are used to visualize the levels of specific surface markers that enable classification of the T cells as, for example, a recently activated CD4⁺ T cell, a memory CD8⁺ T cell, etc. In addition, cytokines and other products of T cells can be studied by isolating the
- 5 T cells from lymphoid tissue and restimulating them with cognate antigen. Following antigen stimulation the T cells produce cytokines that may be visualized by, for example, intracellular cytokine staining coupled with flow cytometry, or collecting the cell supernatants and using Luminex bead technology to measure 15-25 cytokines simultaneously.
- 10 Figure 206 shows that a composition (rSIRP7) that includes recombinantly-produced PflB (SEQ ID NO:353), recombinantly-produced Opp1A (SEQ ID NO:364), recombinantly-produced SirA (SEQ ID NO:375), recombinantly-produced SYN2 (SEQ ID NO:386), recombinantly-produced FhuD (SEQ ID NO:397), recombinantly-produced SYN1 (SEQ ID NO:408), and
- 15 recombinantly-produced MntC (SEQ ID NO:419) induces a cytokine profile similar to the cytokine profile induced by the SIRP extract demonstrated to provide immunological activity against infection by *Staphylococcus* spp. The rSIRP7 composition induced the production of, for example, IL-2, IL-6, IL-17, IFN- γ , MIP-2, and GM-CSF.
- 20 Thus, in addition to mouse models, those of ordinary skill in the art recognize that immunological activity commensurate with the methods described herein may correlate with any one or more of the following: Western blot data showing that serum from animals exposed to a *Staphylococcus* spp. contains antibody that specifically binds to a candidate polypeptide, cell surface
- 25 binding assays demonstrating that antibody that specifically binds to a candidate polypeptide specifically binds to a *Staphylococcus* spp., opsonophagocytosis data, and cytokine induction.
- Another aspect of the present invention provides methods for detecting antibody that specifically binds polypeptides of the present invention. These
- 30 methods are useful in, for instance, detecting whether an animal has antibody that specifically binds polypeptides of the present invention, and diagnosing whether an animal may have a condition caused by a microbe expressing polypeptides described herein, or expressing polypeptides that share epitopes

with the polypeptides described herein. Such diagnostic systems may be in kit form. The methods include contacting an antibody with a preparation that includes a polypeptide of the present invention to result in a mixture. The antibody may be present in a biological sample, for instance, blood, milk, or
5 colostrum. The method further includes incubating the mixture under conditions to allow the antibody to specifically bind the polypeptide to form a polypeptide:antibody complex. As used herein, the term "polypeptide:antibody complex" refers to the complex that results when an antibody specifically binds to a polypeptide. The preparation that includes the polypeptides of the present
10 invention may also include reagents, for instance a buffer, that provide conditions appropriate for the formation of the polypeptide:antibody complex. The polypeptide:antibody complex is then detected. The detection of antibodies is known in the art and can include, for instance, immunofluorescence or peroxidase. The methods for detecting the presence of antibodies that
15 specifically bind to polypeptides of the present invention can be used in various formats that have been used to detect antibody, including radioimmunoassay and enzyme-linked immunosorbent assay.

The present invention also provides a kit for detecting antibody that specifically binds polypeptides of the present invention. The antibody detected
20 may be obtained from an animal suspected to have an infection caused by a gram positive microbe, more preferably, a member of the family Micrococcaceae, preferably, *Staphylococcus* spp., more preferably, *S. aureus*; *Streptococcus* spp., *Bacillus* spp., *Clostridium* spp., *Corynebacterium* spp., *Enterococcus* spp., *Erysipelothrix* spp., *Kytococcus* spp., *Listeria* spp.,
25 *Micrococcus* spp., *Mycobacterium* spp., and *Erysipelothrix* spp.

The kit includes at least one of the polypeptides of the present invention (e.g., one, at least two, at least three, etc.), in a suitable packaging material in an amount sufficient for at least one assay. Optionally, other reagents such as buffers and solutions needed to practice the invention are also included. For
30 instance, a kit may also include a reagent to permit detection of an antibody that specifically binds to a polypeptide of the present invention, such as a detectably labeled secondary antibody designed to specifically bind to an antibody obtained from an animal. Instructions for use of the packaged polypeptides are

also typically included. As used herein, the phrase “packaging material” refers to one or more physical structures used to house the contents of the kit. The packaging material is constructed by well known methods, generally to provide a sterile, contaminant-free environment. The packaging material may have a label which indicates that the polypeptides can be used for detecting antibody that specifically binds polypeptides of the present invention. In addition, the packaging material contains instructions indicating how the materials within the kit are employed to detect the antibody. As used herein, the term “package” refers to a container such as glass, plastic, paper, foil, and the like, capable of holding within fixed limits the polypeptides, and other reagents, for instance a secondary antibody. Thus, for example, a package can be a microtiter plate well to which microgram quantities of polypeptides have been affixed. A package can also contain a secondary antibody. “Instructions for use” typically include a tangible expression describing the reagent concentration or at least one assay method parameter, such as the relative amounts of reagent and sample to be admixed, maintenance time periods for reagent/sample admixtures, temperature, buffer conditions, and the like.

The present invention is illustrated by the following examples. It is to be understood that the particular examples, materials, amounts, and procedures are to be interpreted broadly in accordance with the scope and spirit of the invention as set forth herein.

EXAMPLES

Example 1

Preparation of iron regulated proteins

Laboratory Scale

Compositions derived from different strains of *Staphylococcus aureus* including novel proteins expressed under iron-restriction and/or other degrees of metal ion chelation were evaluated for efficacy against a virulent challenge in

mice. The efficacy of the composition was evaluated by collecting data on the following parameters (1) the efficacy of each composition to provide homologous and heterologous protection against a live virulent challenge in mice, (2) the efficacy of each composition to reduce necrotic skin lesions, and
5 (3) the efficacy of compositions derived from *Staphylococcus* grown in replete and deplete iron conditions to provide protection.

The *Staphylococcus aureus* strains evaluated in this study originated from three animal species; avian, human and bovine. The avian isolate SAAV1 was a field isolate originating from a flock of diseased turkeys having a high
10 degree of osteomyelitis and synovitis. The bovine isolates (strain 1477 and strain 2176) were isolated from two different commercial dairy herds having a high incidence of clinical mastitis. The human isolate was obtained from the ATCC (strain 19636), and originated from a patient having clinical osteomyelitis.

15 Master seed stocks of each isolate were prepared by inoculating the appropriate isolate into 200 ml of Tryptic Soy Broth (TSB, Difco Laboratories, Detroit, MI) containing 300 μ M 2,2-dipyridyl (Sigma-Aldrich St. Louis, MO). The culture was grown while stirring at 200 rpm for 6 hours at 37°C, and collected by centrifugation at 10,000 x g. The bacterial pellet was re-suspended
20 into 100 ml TSB broth containing 20% glycerol, and sterilely dispensed into 2 ml cryogenic vials (1 ml per vial) and stored at -90°C until use.

Each master seed stock was expanded into a working seed. One vial of each master seed isolate was inoculated into 200 ml of Tryptic Soy Broth (TSB, Difco Laboratories, Detroit, MI) containing 1000 μ M 2,2-dipyridyl (Sigma-
25 Aldrich St. Louis, MO). The culture was grown while stirring at 200 rpm for 6 hours at 37°C, and collected by centrifugation at 10,000 x g. The bacterial pellet was resuspended into 100 ml TSB broth containing 20% glycerol, and sterilely dispensed into 2 ml cryogenic vials (1 ml per vial) and stored at -90°C until use. The working seed was used for the production of compositions
30 enriched with iron-regulated membrane proteins, including iron-regulated membrane proteins.

All strains were adapted to grow in highly iron-depleted media (i.e., media containing very low levels of free iron). This was accomplished by sub-

culturing the bacteria in TSB containing increasing concentrations of 2,2-dipyridyl (from 300 to 1600 μ M).

Proteins were prepared from bacteria as follows. The bacteria were grown from frozen working seed stocks by subculturing into 25 ml of iron-deplete media (containing 1000 μ M 2,2'-dipyridyl) and iron-replete media, then incubated at 37°C while shaking at 400 rpm. Following 12 hours of incubation, 5 ml of each culture was transferred into 500 ml of iron-deplete or iron-replete media pre-incubated at 37°C. Cultures were incubated for 8 hours at 37°C while shaking at 100 rpm, then cells were pelleted by centrifugation at 10,000 x g for 20 minutes. Bacterial pellets were resuspended in 100 ml of sterile physiological saline and centrifuged at 10,000 x g for 10 minutes. Pellets were then resuspended in 45 ml of Tris-buffered saline, pH 7.2 (TBS; 25 mM Tris, 150 mM NaCl) and the resulting bacterial suspensions were dispensed as 9-ml aliquots into 5 individual tubes. One milliliter of TBS containing 50 units of lysostaphin (Sigma, St. Louis, MO) was added to each tube to give a final volume of 5 units/ml. Following incubation at 37°C for 30 minutes while shaking at 200 rpm, 1 ml of TBS containing 0.1 mg of lysozyme (Sigma) was added to each tube. The bacterial suspensions were then incubated for an additional 45 minutes while shaking at 200 rpm. Next, suspensions were centrifuged at 3050 x g for 12 minutes at 4°C to pellet large cellular debris. The supernatants were collected by aspiration without disturbing the pellet. The supernatant was then centrifuged at 39,000 x g for 2.5 hours. The resulting pellets containing the proteins were resuspended into 200 μ L Tris buffer, pH 7.2, without saline. The protein solution for each isolate were combined for a total volume of 1 ml and stored at -90°C.

The protein-enriched extracts derived from *S. aureus* were size-fractionated on SDS-PAGE gels using a 4% stacking gel and 10% resolving gel. Samples for electrophoresis were prepared by combining 10 μ L of sample with 30 μ L of SDS reducing sample buffer (62.5mM Tris-HCL pH 6.8, 20% glycerol, 2% SDS, 5% β -mercaptoethanol) and boiled for 4 minutes. Samples were electrophoresed at 18 mA constant current for 5 hours at 4°C using a Protein II xi cell power supply (BioRad Laboratories, Richmond, CA, model 1000/500). The molecular weight of each individual protein as visually seen in the SDS-

PAGE gel was estimated using a GS-800 densitometer (BioRad) using a broad range molecular weight marker as a reference standard (BioRad).

The SDS-PAGE patterns of the proteins from each isolate when grown in the presence of 1600 μ M dipyrindyl showed a very different protein expression pattern compared to the same strain when grown in the presence of 300 μ M dipyrindyl. For instance, when grown in 300 μ M dipyrindyl isolate SAAV1 resulted in metal-regulated proteins of 90 kDa, 84 kDa, 72 kDa, 66 kDa, 36 kDa, 32 kDa, and 22 kDa, while growth in 1600 μ M dipyrindyl resulted in metal-regulated proteins of 87.73 kDa, 54.53 kDa, 38.42 kDa, 37.37 kDa, 35.70 kDa, 34.91 kDa, and 33.0 kDa. Likewise, when grown in 300 μ M dipyrindyl isolate 19636 resulted in proteins of 42 kDa and 36 kDa, while growth in 1600 μ M dipyrindyl resulted in metal-regulated proteins of 87.73 kDa, 54.53 kDa, 38.42 kDa, 37.37 kDa, 35.70 kDa, 34.91 kDa, and 33.0 kDa. All conditions, including growth in iron-replete media, resulted in the expression of the following proteins that were presumably not metal-regulated: 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa 43kDa 41 kDa, and 40 kDa.

Furthermore, growth of the different strains of *S. aureus* in 1600 μ M dipyrindyl resulted in similar protein expression patterns. The compositions enriched in iron-regulated membrane proteins from the avian isolate (SAAV1) included proteins with molecular weights of 87.73 kDa, 54.53 kDa, 38.42 kDa, 37.37 kDa, 35.70 kDa, 34.91kDa, and 33.0 kDa. The molecular weights of the proteins from the ATCC isolate 19636 were essentially identical to those from the avian isolate. Both bovine isolates, when grown with 1600 μ M 2,2-dipyrindyl, expressed similar banding profiles as the avian and ATCC isolates for the majority of the proteins (87.73 kDa, 54.53 kDa, 37.7 kDa, 35.70 kDa, 34.91 kDa, and 33.0 kDa). However, neither of the bovine isolates produced the 38.42 kDa protein seen with the avian and ATCC isolates, and the bovine isolates expressed three proteins (80.46 kDa, 65.08 kDa, and 31.83 kDa) not observed with the avian and ATCC strains (see Figure 1 and Table 7). All conditions resulted in the expression of the following proteins that were not metal-regulated: 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, and 40 kDa.

Table 7. Molecular weights of metal-regulated polypeptides obtained from *Staphylococcus aureus* isolates.

Avian SAAV1	Human 19636	Bovine 1477	Bovine 2176
87.73	87.73	87.73	87.73
-	-	80.46	80.46
-	-	65.08	65.08
54.53	54.53	54.53	54.53
38.42	38.42	-	-
37.37	37.37	37.37	37.37
35.70	35.70	35.70	35.70
34.91	34.91	34.91	34.91
33.0	33.0	33.0	33.0
		31.83	31.83

5

Interestingly, there was no difference in the protein profiles as examined by SDS-PAGE between the clarified supernatant and the bacterial pellet after treating the bacteria with lysostaphin/lysozyme. Both the extracted bacterial pellet and the supernatant had exactly the same protein profiles as examined by SDS-PAGE. This same observation was also seen when disrupting the bacterial cells using an AVESTIN homogenizer at 30,000 psi. The resultant bacterial pellet, after slow speed centrifugation was identical in its protein profile as compared to the clarified supernatant after high speed centrifugation at 30,000 x g for 2.0 hours at 4°C.

15

Example 2

Preparation of the immunizing compositions derived from *Staphylococcus aureus*

20

The proteins from the human isolate ATCC 19636 and the bovine isolate 1477, grown in iron-deplete conditions and prepared as described in Example 1, were used to formulate two vaccine compositions. The proteins from the ATCC isolate had molecular weights of 87.73 kDa, 54.53 kDa, 38.42 kDa, 37.37 kDa, 35.70 kDa, 34.91 kDa, and 33.0 kDa, while the bovine isolate expressed

25

proteins having molecular weights 87.73 kDa, 80.46 kDa, 65.08 kDa, 54.53 kDa, 37.37 kDa, 35.70 kDa, 34.91 kDa, 33.0 kDa, and 31.83. Each composition also contained the following proteins that were not metal-regulated: 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, and 40 kDa. Stock vaccines were prepared from the two strains by emulsifying each aqueous protein suspension (500 µg total protein/ml) into a commercial adjuvant (EMULSIGEN, MVP Laboratories, Ralston, Nebraska) using an IKA Ultra Turrax T-50 homogenizing vessel (IKA, Cincinnati, OH) to give a final dose of 50 µg total protein in a 0.1 ml injectable volume with an adjuvant concentration of 22.5% vol/vol. As a control vaccination, a protein composition was prepared from the bovine isolate 1477 grown under iron-replete conditions (TSB supplemented with 300 µM ferric chloride) as described in Example 1. A placebo vaccine was prepared by substituting physiological saline for the aqueous protein suspension in the above protocol.

15

Example 3

Mouse vaccination

Seventy (N=70) female CF-1 mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams were equally distributed into 7 groups (10 mice/group). Mice were housed in polycarbonate mouse cages (Ancore Corporation, Bellmore, NY). A single cage was used for each treatment group and food and water was supplied *ad libitum* to all mice. All mice were vaccinated intraperitoneally with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

- Group-1: Placebo-Vaccinated
- Group-2: Vaccinated with ATCC 19636 proteins expressed under iron-restriction.
- Group-3: Placebo-Vaccinated
- Group-4: Vaccinated with Bovine 1477 proteins expressed under iron-restriction.

Group-5: Vaccinated with Bovine 1477 proteins expressed under iron-restriction.

Group-6: Vaccinated with ATCC 19636 proteins expressed under iron-restriction.

- 5 Group-7: Bovine 1477 FeCl₃ -Vaccinated, where "Bovine 1477 FeCl₃" refers to proteins obtained from Bovine 1477 grown in TSB supplemented with 300 µM ferric chloride.

10

Example 4

Preparation of challenge organism

- The previously described *Staphylococcus aureus* strains ATCC 19636 and strain 1477 were used as challenge organisms. Briefly, the isolates from frozen stocks (previously described) were streaked onto blood agar plates and incubated at 37°C for 18 hours. A single colony of each isolate was subcultured into 50 ml Tryptic Soy Broth (Difco) containing 1600 µM 2,2' dipyridyl. The cultures were incubated at 37°C for 6 hours while rotating at 200 rpm, then centrifuged at 10,000 x g for 10 minutes at 4°C to pellet the bacteria. The bacterial pellets were washed twice by centrifugation in TBS at 4°C. The final pellets were resuspended in TBS to an optical density of 42 % Transmittance (T) at 562 nm in a volume of approximately 25 ml of TBS and used for challenge. Just prior to challenge, 1 ml of these bacterial suspensions was serially diluted and plated on agar to enumerate the number of colony-forming units (CFU) per mouse dose.

30

Example 5

Challenge

Fourteen days after the second vaccination, mice in all groups (1-7) were subcutaneously challenged in the back of the neck with 0.1 ml of the appropriate organism. The seven groups of mice were challenged as follows:

Group-1 (Placebo-Vaccinated): Challenged with ATCC 19636

Group-2 (Vaccinated with ATCC 19636 proteins expressed under iron-restriction): Challenged with ATCC 19636

5 Group-3 (Placebo-Vaccinated): Challenged with Bovine 1477

Group-4 (Vaccinated Bovine 1477 proteins expressed under iron-restriction): Challenged with Bovine 1477

Group-5 (Vaccinated Bovine 1477 proteins expressed under iron-restriction): Challenged with ATCC 19636

10 Group-6 (Vaccinated ATCC 19636 proteins expressed under iron-restriction): Challenged with Bovine 1477

Group-7 (Bovine 1477 FeCl₃ -Vaccinated): Challenged with Bovine 1477

15 As determined by the enumeration protocol described in Example 4, the concentration of *S. aureus* 19636 used for challenge was 1.35×10^8 CFU per mouse dose, and the concentration of *S. aureus* 1477 used for challenge was 1.65×10^8 colony CFU per mouse dose. Morbidity, mortality and gross pathology were recorded daily for 7 days after challenge.

20 When comparing the mice challenged with the ATCC 19636 isolate, 70% of the placebo-vaccinated Group 1 mice died within 7 days of challenge (Table 8 and Fig. 2). This demonstrated that strain 19636 caused a high rate of mortality in mice at the dose level administered. In contrast to the mice in Group 1, only 10% of the mice in Group 2 died within 7 days post-challenge.

25 These results illustrated that the mice challenged with strain 19636 were significantly protected by vaccination with the 19636 composition ($p = 0.020$, Fischer's Exact test). Furthermore, a Kaplan-Meier analysis of the time-to-death data indicated that the vaccine afforded significant ($p = 0.0042$, logrank test) protection against homologous challenge (Fig. 3). In addition, only 20% of
30 the mice in Group 5 died within 7 days of challenge, indicating that the bovine 1477 composition offered significant protection against challenge with the ATCC 19636 strain ($p = 0.015$ logrank test for mortality). When the data was analyzed by a Kaplan- Meier survival curve and logrank test (Fig. 4), the

protection against mortality was determined to be significant ($p = 0.015$ logrank test for mortality), indicating that the vaccine composition derived from strain 1477 provided heterologous protection against challenge with strain 19636..

5

Table 8. Mortality of Vaccinated and Non-Vaccinated Mice Following Challenge with *Staphylococcus aureus* (human ATCC isolate 19636 and bovine isolate 1477).

Groups	# Mice	# Dead	Percent mortality (%)
Group-1* (Placebo, ATCC 19636 Chlg)	10	7/10	70
Group-2* (ATCC 19636, Homologous Chlg)	10	1/10	10
Group-3* (Placebo, Bovine 1477 Chlg)	10	2/10	20
Group-4* (Bovine 1477, Homologous Chlg)	10	1/10	10
Group-5* (Bovine 1477, Heterologous Chlg)	10	2/10	20
Group-6* (ATCC 19636, Heterologous Chlg)	10	0/10	0
Group-7* (Bovine 1477 FeCl ₃ , Bovine 1477 Chlg)	10	2/10	20

*Group-1, (Placebo-Vaccinated/Challenged with ATCC 19636)

10 *Group-2 (Vaccinated with ATCC 19636 proteins expressed under iron-restriction /Challenged with ATCC 19636)

*Group-3 (Placebo-Vaccinated/Challenged with Bovine 1477)

*Group-4 (Vaccinated with Bovine 1477 proteins expressed under iron-restriction /Challenged with Bovine 1477)

15 *Group-5 (Vaccinated with Bovine 1477 proteins expressed under iron-restriction /Challenged with ATCC 19636)

*Group-6 (Vaccinated with ATCC 19636 proteins expressed under iron-restriction /Challenged with Bovine 1477)

*Group-7 (Bovine 1477 FeCl₃ -Vaccinated/Challenged with Bovine 1477)

20

When comparing the mice challenged with the bovine 1477 isolate, only 20% of the mice in the placebo-vaccinated group (Group 3) died within 7 days of challenge. However, challenge with the bovine 1477 isolate elicited the development of necrotic skin lesions on 6 (75%) of the surviving mice of Group 3. These lesions were measured and the average size of the lesions on the surviving mice was 18.5 mm (Table 9). In contrast, 20% of the Group 4 mice died within 7 days of challenge, but only three (38%) of the surviving mice developed lesions (average diameter, 2.7 mm). These results indicate that the

25

bovine 1477 composition offered significant homologous protection against development of lesions in the mice challenged with the bovine strain 1477 ($p = 0.009$, Student's *t*-test). In addition, vaccination with the ATCC 19636 composition protected against challenge with strain 1477, since no mice died in Group 6 and only three (30%) of the mice developed skin lesions (average diameter, 3.7 mm). Taken together, the reduced mortality and/or lesion development in the mice in Groups 5 and 6 demonstrate the significant cross-protective nature of the compositions derived from strains 19636 and 1477 ($p = 0.012$, Student's *t*-test based on lesion size). In demonstration of the efficacy of the composition as compared to the non-iron regulated proteins, 20% of the mice in Group 7 died and 4 of the survivors developed skin lesions (average diameter, 15.8 mm). The mice of Group 7 demonstrated some degree of protection by vaccination with the proteins of the 1477 isolate since fewer mice developed lesions compared to the placebo-vaccinated Group 3. However, the skin lesions observed on the mice in group 7 were more frequent and of a larger diameter than the lesions on the mice of Group 4, indicating that, relative to proteins isolated from cells grown under iron-replete conditions, the proteins isolated from bacteria grown under iron restriction offered superior protection against an identical challenge.

The cross-protective nature of the proteins observed in the mouse challenge study is supported by the similar molecular weights of the proteins from the *S. aureus* strains described in Example 1 (Figure 1). Although there were noticeable differences in the SDS-PAGE profile of the proteins from the bovine-derived isolates, specifically the absence of a 38.4 kDa protein and the presence of 3 additional proteins, the proteins from both strains 1477 and ATCC 19636 elicited heterologous protection. These results indicate that the similar proteins between strains 19636 and 1477 are likely responsible for the cross-protection observed in Groups 5 and 6. By contrast, the protein profiles from strain 1477 grown under iron-deplete and iron-replete conditions are observably different. Those proteins isolated under iron-depleted conditions are more protective when compared to the proteins isolated under iron-replete conditions, demonstrated by the reduction in lesion development among the mice of Group 4 compared to the mice of Group 7.

Table 9. The Induction of Necrotic Lesions in Mice Seven Days Post-Challenge with *Staphylococcus aureus* (ATCC Isolate 19636 and/or Bovine Isolate 1477)

Lesion diameter (millimeter) per mouse						
Group-1	Group-2	Group-3	Group-4	Group-5	Group-6	Group-7
No lesion	No lesion	26	5	5	5	25
No lesion	No lesion	25	2	No lesion	5	25
No lesion	No lesion	24	1	No lesion	1	10
Dead	No lesion	24	No lesion	No lesion	No lesion	3
Dead	No lesion	7	No lesion	No lesion	No lesion	No lesion
Dead	No lesion	5	No lesion	No lesion	No lesion	No lesion
Dead	No lesion	No lesion	No lesion	No lesion	No lesion	No lesion
Dead	No lesion	No lesion	No lesion	No lesion	No lesion	No lesion
Dead	No lesion	Dead	No lesion	Dead	No lesion	Dead
Dead	Dead	Dead	Dead	Dead	No lesion	Dead
Average lesion diameter (mm) among surviving mice						
0	0	18.5	2.7	5	3.7	15.8

*Group-1, (Placebo -Vaccinated/Challenged ATCC 19636)

*Group-2 (Vaccinated with ATCC 19636 proteins expressed under iron-restriction /Challenged ATCC 19636)

*Group-3 (Placebo -Vaccinated/Challenged Bovine 1477)

*Group-4 (Vaccinated with Bovine 1477 proteins expressed under iron-restriction /Challenged Bovine 1477)

*Group-5 (Vaccinated with Bovine 1477 proteins expressed under iron-restriction /Challenged ATCC 19636)

*Group-6 (Vaccinated with ATCC 19636 proteins expressed under iron-restriction /Challenged Bovine 1477)

*Group-7 (Bovine 1477 FeCl₃ Vaccinated/Challenged Bovine 1477)

Example 6

In mammals, it has been shown that the response to tissue injury or bacterial infection results in an acute inflammatory response. This response
5 increases capillary permeability and phagocytic infiltration resulting in the clinical signs recognized as inflammation; swelling, fever, pain and redness; if left uncontrolled, this may lead to death. The activation of humoral factors and the release of cytokines mediate systemic events collectively known as the acute phase protein response which results in a cascade of physiological and
10 biochemical events. The duration of this response is directly related to the severity of the injury and magnitude of the systemic infection. It has been well-documented that during bacterial sepsis, major surgery, burns and other bodily trauma there is an alteration in the concentration of a number of metal ions in serum such as, iron, copper, and zinc. For instance, during the acute phase of an
15 infection there is a decrease in plasma levels of iron and zinc and an increase in copper. The alteration of these trace metal ions in serum may directly affect the severity or progression of any bacterial infection.

In this study we examined the expression of proteins of *Staphylococcus aureus* under various conditions of metal ion restriction in order to mimic the
20 expression of novel proteins that may be expressed during systemic invasion. The *Staphylococcus aureus* strains evaluated in this study originated from clinical samples of three different species of animal; avian (strain SAAV1), human (strain 19636), and bovine (strains 1477 and 2176). Briefly, cultures of each isolate were prepared from master seed stocks in 200 ml of Tryptic Soy
25 Broth (TSB). Each culture was grown while stirring at 200 rpm for 6 hours at 37°C. Ten ml of each culture were transferred into 500 ml of deplete TSB containing one of four metal ion chelators; 2, 2-dipyridyl (Dp), 2-pyridylmethyl-ethylene diamine (TPEN), catechin, and naringenin (all obtained from Sigma, St. Louis, MO). In addition each culture was also grown in cation-
30 replete media containing ferric chloride, zinc chloride and/or copper chloride prepared at 300 µM concentrations. The metal ion chelators were used at the following concentration; 2,2-dipyridyl (800 µM), catechin and naringenin were used at 300 µM, and 2-pyridylmethyl-ethylene diamine was used at a

concentration of 100 μ M. Cultures were grown with each chelator for 8 hours, at which point the culture was subcultured a second time for an additional 12 hours. Each culture was subcultured for three consecutive passes at 12-hour intervals. At the end of the third pass, each culture was harvested by
5 centrifugation at 10,000 x g for 20 minutes. Each culture was washed twice by centrifugation at 10,000 x g and resuspended in 20 ml Tris-buffered saline, pH 7.2 at 4°C.

Each bacterial pellet was resuspended in 45 ml of Tris-buffered saline, pH 7.2 (25 mM Tris and 150 mM NaCl) and the resulting bacterial suspensions
10 were dispensed as 9-ml aliquots into 5 individual tubes, twenty tubes total. One milliliter of TBS containing 50 units of lysostaphin (Sigma, St. Louis, MO) was added to each tube to give a final concentration of 5 units/ml. Following incubation at 37°C for 30 minutes while shaking at 200 rpm, 1 ml of TBS containing 0.1 mg of lysozyme (Sigma) was added to each tube. The bacterial
15 suspensions were then incubated for an additional 45 minutes while shaking at 200 rpm. Next, suspensions were centrifuged at 3050 x g for 12 minutes at 4°C to pellet large cellular debris. The supernatants were collected by aspiration without disturbing the pellet. The supernatant was then centrifuged at 39,000 x g for 2.5 hours. The resulting pellets, enriched for metal-regulated membrane
20 proteins, were resuspended in 200 μ L Tris buffer, pH 7.2. The protein solutions for each isolate were combined for a total volume of 1 ml and stored at -90°C.

The proteins obtained from the SAAV1, 19636, 1477 and 2176 *S. aureus* isolates grown under iron, zinc and copper deplete conditions included metal-regulated polypeptides.

25 Cell extracts, derived from each isolate were size-fractionated on SDS-PAGE gels using a 4% stacking gel and 10% resolving gel. Samples for electrophoresis were prepared by combining 10 μ l of sample with 30 μ l of SDS reducing sample buffer (62.5mM Tris-HCL pH 6.8, 20% glycerol, 2% SDS, 5% beta-mercaptoethanol) boiled for 4 minutes. Samples were electrophoresed at
30 18 mA of constant current for 5 hours at 4°C using a Protein II xi cell power supply (BioRad Laboratories, Richmond, CA, model 1000/500).

The SDS-PAGE patterns of the proteins grown under zinc and/or copper chelation showed unique banding patterns in all isolates that were different

when compared to the same isolates grown under iron-restriction in the presence of 2,2'-diipyridyl. For example, when the 19636 isolate was grown under iron-restriction or in the presence of the chelator 2,2'-diipyridyl, unique iron-regulated proteins were expressed at the 87.73 kDa, 54.53 kDa, 38.42 kDa, 5 37.37 kDa, 35.70 kDa, 34.91 kDa and 33.0 kDa regions. These proteins were downregulated when the isolate was grown in the presence of ferric chloride. However, when the same isolate was grown in the presence of the zinc and or copper chelator, a novel subsets of proteins was expressed relative to the proteins expressed under iron-restriction; the new proteins having molecular 10 weights of approximately 115 kDa, 88 kDa, 80 kDa, 71 kDa, 69 kDa, 35 kDa, 30 kDa, 29, kDa and 27 kDa. In addition, an 87.73 kDa protein was expressed under conditions of iron restriction or copper-restriction but not when cultures were zinc-restricted. The proteins expressed under iron-restriction appeared to be downregulated when growth was under either zinc-restriction and/or copper- 15 restriction, but not completely shut off as seen when the isolate was grown in ferric chloride.

It appears that there are novel proteins expressed when the organism is grown under copper-restriction and/or zinc-restriction that are not expressed when the same isolate is grown under iron-restricted conditions. Since 20 transitional metals are used by organisms to build enzymes that catalyze various biochemical reactions, the metal ions may play a vital role in microbial survival during a systemic infection. It is perhaps for this reason that during sepsis there is a transient decrease in the availability of these transitional metals, making them unavailable for growth of the organism. These novel proteins could very 25 well enhance the protective efficacy of the existing composition grown under iron-restriction because they may also be expressed by the bacteria under the metal ion restriction experienced during systemic invasion.

30

Example 7

Compositions of the present invention can also be produced under large scale commercial conditions.

5 *Fermentation*

A cryogenic vial of the working seed (2 ml at 10^9 CFU/ml) as described in Example 1 was used to inoculate 500 ml of Tryptic Soy Broth (TSB) without dextrose (Difco) pre-warmed to 37°C containing 0.125 g/l 2,2-dipyridyl (Sigma), 2.7 grams BiTek yeast extract (Difco) and glycerol (3% vol/vol). The culture was incubated at 37°C for 12 hours while stirring at 200 rpm at which time it was used to inoculate 2 liters of the above media and allowed to grow for an additional 4 hours at 37°C. This culture was used to inoculate a 20-liter VIRTIS bench-top fermentor, (Virtis, Gardiner, NY) charged with 13 liters of the above-described media. The pH was held constant between 6.9 and 7.1 by automatic titration with 50% NaOH and 10% HCL. The stirring speed was adjusted at 400 rev/minute, and the culture aerated with 11 liters air/minute at 37°C. Foaming was controlled automatically by the addition of 11 ml defoamer (Mazu DF 204 Chem/Serv, Minneapolis, MN). The culture was allowed to grow continuously at these conditions for 4 hours at which time was sterilely pumped into a 150-liter fermentor (W. B. Moore, Easton, PA). The fermentor was charged with 120 liters tryptic soy broth without dextrose (3,600.0 grams), BiTek yeast extract (600 grams), glycerol (3,600 ml), 2,2-dipyridyl (3.0 grams) and Mazu DF 204 defoamer (60 ml). The parameters of the fermentation were as follows: dissolved oxygen (DO) was maintained at 30% +/- 10% by increasing agitation to 220 rev/minute sparged with 60 liters of air/minute and 10 pounds per square inch (psi) back pressure. The pH was held constant between 6.9 and 7.1 by automatic titration with 50% NaOH and 10% HCL and the temperature maintained at 37°C. At hour 4.5 (OD_{540} 8-9) of the fermentation the culture was transferred to a 1,500 liter New Brunswick Scientific fermentor IF-15000 charged with 1200 liters tryptic soy broth without dextrose (36,000 grams), BiTek yeast extract (6,000 grams), glycerol (36,000 ml), 2,2-dipyridyl (30.0 grams) and Mazu DF 204 defoamer (600 ml). The parameters of the fermentation were as follows: dissolved oxygen (DO) was

maintained at 60% +/- 10% with supplemental oxygen by increasing agitation to 300 rev/minute sparged with 300 to 1100 liters of air/minute and 5 pounds per square inch (psi) back pressure. As fermentation progressed supplemental oxygen was added from 0-90 liters/minute to assist in the control of dissolved
5 oxygen. The pH was held constant between 6.9 and 7.4 by automatic titration with 50% NaOH and 10% HCL and the temperature was maintained at 37°C.

At approximately 5 hours post inoculation of the large fermentor the culture was supplemented with additional nutrients by feeding 70 liters of media containing 18,000 grams TSB without dextrose, 3,000 grams yeast extract 30.0
10 grams 2,2-dipyridyl and 18,000 ml of glycerol. The rate of feed was adjusted to approximately 28 liters/hour while increasing agitation. At the end of the feed the fermentation was allowed to continue for an additional 4 hours at which point the fermentation was terminated by lowering the temperature of the fermentor to 18°C (OD₅₄₀ 35-40 at a 1:100 dilution).

15

Harvest

The bacterial fermentation was concentrated and washed using a Pall Filtron Tangential Flow Maxiset-25 (Pall Filtron Corporation, Northboro, MA) equipped with three 30ft² Alpha 300-K open channel filters, catalog No.
20 AS300C5, (Pall Filtron) connected to a Waukesha Model U-60 feed pump (Waukesha Cherry-Burrell, Delevan, WI) The original culture volume of 1250 liters was reduced to 50 liters (2.5 liters/minute) using a filter inlet pressure of 30 psi and a retentate pressure of 5-6 psi. The bacterial retentate was adjusted back up to 150 liters using Tris-buffered Saline pH 8.5 and then concentrated
25 again to 50 liters to help remove any contaminating exogenous proteins, such as exoproteins to include secreted toxins and proteases. The elevated pH of the tris-buffered saline helps prevent much of the proteolytic degradation that can occur during storage of the whole cell suspension. Protease inhibitors may be used instead of, or in addition to, an elevated pH. The retentate was mixed
30 thoroughly while in the 200-liter tank using a bottom mount magnetically driven mixer. The retentate was sterilely dispensed (3.5 liters) into sterile 4 liter Nalgene containers No. 2122 and placed into a -20°C freezer for storage as a breaking point in the manufacture, or could be further processed. The pellet

mass was calculated by centrifuging 30 ml samples of the fermented culture and final harvest. Briefly, pre-weighted 50 ml Nalgene conical tubes were centrifuged at 39,000 x g for 90 minutes in a Beckman J2-21 centrifuge using a JA-21 rotor (Beckman Instruments, Palo Alto CA). At the end of the run, the supernate was poured off and the tubes were weighed again. The pellet mass was calculated for each stage. The fermentation process yielded a wet pellet mass of approximately 60 kilograms.

Disruption

Eighty kilograms of bacterial cell slurry in Tris-buffered Saline pH 8.5 was aseptically transferred into a steam in place 1000 liter jacketed process tank (Lee, Model 259LU) with a top mounted mixer (Eastern, Model TME-1/2, EMI Incorporated, Clinton, CT) containing 900 liters TBS pH 8.5. The bulk bacterial suspension was chilled to 4°C with continuous mixing for 18 hours at 200 rpm at which time was disrupted by homogenization. Briefly, the 1000 liter tank containing the bacterial suspension was connected to a model C-500-B AVESTIN homogenizer, (Avestin Inc, Ottawa Canada). A second 1000 liter jacketed process tank (empty) was connected to the homogenizer such that the fluid in the process tank could be passed through the homogenizer, into the empty tank and back again, allowing for multiple homogenizing passes while still maintaining a closed system. The temperature during homogenization was kept at 4°C. At the start of the first pass, fluid was circulated at 70 psi via a Waukesha model 10DO pump (Waukesha) through the homogenizer (500 gallons/hour), while the homogenizer pressure was adjusted to 30,000 psi. Prior to the first pass, two pre-homogenizing samples were withdrawn from the homogenizer to establish a baseline for determining the degree of disruption and monitoring of pH. The degree of disruption was monitored by transmittance (% T at 540 nm at 1:100 dilution) compared to the non-homogenized sample. The number of passes through the homogenizer was standardized to give a final percent transmittance between 78-91%T at a 1:100 dilution preferably between 86-91%. After homogenization, the tank was removed from the homogenizer and put onto a chiller loop at 4°C and mixed at 240 rpm.

Protein Harvest

The disrupted bacterial suspension containing the iron-regulated proteins as illustrated in Figure 1 were collected by centrifugation using T-1 Sharples, (Alfa Laval Separations, Warminster, PA). Briefly, the 1000 liter jacketed
5 process tank containing the disrupted bacterial homogenate was fed into 12 Sharples with a feed rate of 250 ml/minute at 17 psi at a centrifugal force of 60,000 x g. The effluent was collected into a second 1000 liter jacketed process tank through a closed sterile loop allowing for multiple passes through the centrifuges while maintaining a closed system. The temperature during
10 centrifugation was kept at 4°C. The homogenate was passed 8 times across the centrifuges. Approximately 50 % of the protein was collected after the second pass, at which point, the homogenate fluid was concentrated to 1/3 of its original volume, which shortened the process time for the next 6 passes. The homogenate tank was aseptically disconnected from the centrifuges and
15 connected to a Millipore Pellicon Tangential Flow Filter assembly (Millipore Corporation, Bedford, MA), equipped with a 25ft² screen-channel series Alpha 30K Centrasette filter (Pall Filtron) connected to a Waukesha Model U30 feed pump for concentration. After concentration, centrifugation was continued until the process was completed. Protein was collected after each pass. The protein
20 was collected, resuspended and dispensed in 50 liters Tris-buffered saline pH 8.5 containing 0.15% formulin (Sigma) as preservative.

Diafiltration

The protein suspension was washed by diafiltration at 4°C to remove any
25 exogenous proteins (proteases, toxins, cytoplasmic and metabolic enzymes etc). Briefly, the 50 liters of protein was sterilely transferred into a 200 liter process tank containing 150 liters sterile Tris-buffer saline, pH 8.5 equipped with a bottom mount Dayton mixer, Model 2Z846 (Dayton Electric, Chicago, IL) rotating at 125 rev/minute. The process tank was sterilely connected to a
30 Millipore Pellicon Tangential Flow Filter assembly (Millipore Corporation), equipped with a 25ft² screen-channel series Alpha 30K Centrasette filter (Pall Filtron) connected to a Waukesha Model U30 feed pump. The 200 liter protein solution was concentrated by filtration to a target volume 50 liters at which

point 150 liters of sterile saline was added. The protein suspension was then concentrated to approximately 50 liters. The protein concentrate was stored in a 50 liter jacketed process tank equipped with a top mounted mixer and stored at 4°C.

- 5 It is interesting to note that the composition derived from the large scale process using homogenization as a means of disruption generated identical banding profiles as examined by SDS-PAGE as compared to the smaller scale process described in Example 1. These results show that lysostaphin could be replaced as the bacterial lysis agent using the AVESTIN homogenizer C500-B.
- 10 This discovery allows for the low cost generation of large volumes of iron-regulated proteins from staphylococci.

Example 8

- 15 Hyper-immunization of mice and preparation of polyclonal antibody

- Passive immunization with purified antibody isolated from mice
- 20 vaccinated with proteins derived from *S. aureus* strain ATCC 19636 grown under iron-limiting conditions was protective against a homologous and heterologous *S. aureus* challenge. Fifteen adult CD1 mice were vaccinated as described in Example 3 with the protein composition derived from *S. aureus* strain ATCC 19636 grown under iron-deplete conditions as described in
- 25 Examples 1 and 2. Mice were vaccinated intraperitoneally 3 times at 7 day intervals with 50 µg of protein composition at each vaccination. Seven days after the third immunization, mice were bled completely by cardiac puncture. Serum was pooled and antibody purified using standard ammonium sulfate precipitation. Exogenous serum proteins were removed first prior to antibody
- 30 precipitation by adding 0.5 volumes of saturated ammonium sulfate pH 7.2. The solution was stirred at 100 rpm for 24 hours at 4°C. The solution was again centrifuged at 3000 x g for 30 minutes. The supernatant was collected and precipitated again by adding enough saturated ammonium sulfate to bring the final concentration to 55 % saturation. The solution was stirred at 100 rpm for
- 35 24 hours at 4°C. The precipitate was centrifuged at 3000 x g for 30 minutes.

The final pellet from each sample was resuspended into 2 ml PBS pH 7.2. The precipitated antibodies were then dialyzed using a 50,000 molecular cut off dialysis tubing (Pierce, Rockford IL) for 30 hours against three 1 liter changes of phosphate-buffered saline to remove ammonium sulfate. The first two liter changes were preserved with .02 % sodium azide. The final 1 liter buffer change contained no preservative. The dialysate was collected and centrifuged again to remove any remaining debris at 3000 x g for 30 minutes. The antibody solution was stored at 4°C for less than 48 hours prior to use. Each sample was plated on blood agar to verify sterility prior to infusion.

10

Example 9

Passive immunization and challenge

15 In order to evaluate the protective effect of infused antibody raised against *S. aureus* proteins expressed during iron-limitation, two groups of 15 mice each were infused intraperitoneally with either the purified antibody preparation (Group 1) or physiological saline (Group 2) in a 200 µL infusion. An additional two groups of 15 mice each were infused subcutaneously with either the purified antibody preparation (Group 3) or physiological saline (Group 4). After 60 minutes, the 2 groups of 15 mice receiving an intraperitoneal infusion were challenged intraperitoneally with 1.3×10^8 cfu of *S. aureus* strain 19636. Similarly, the 2 groups of 15 mice receiving a subcutaneous infusion were challenged subcutaneously with 1.3×10^8 cfu of *S. aureus* strain 1477 to test for cross-protection against challenge by a different *S. aureus* strain. Mortality and/or lesion size was recorded for 5 days and the livers of all mice were removed post-mortem, homogenized and plated to determine the number of *S. aureus* present as a measure of systemic infection. The Kaplan-Meier survival curves (Figures 5 and 6) show the protective effect afforded by the infusion of antibodies from mice vaccinated with the *S. aureus* proteins expressed during iron restriction. Although the difference between the infused and control groups for the ATCC 19636-challenge groups was not significant ($p = 0.076$, log-rank test), the liver of the single mouse that died

20
25
30

within the antibody-infused group at Day 1 was cultured on blood agar to determine the absence and/or presence of the challenge organism (*S. aureus*). The culture derived from this mouse was negative for *Staphylococcus* and showed no growth on the blood agar plate or culture medium. In contrast, the
5 livers of the mice that died within the placebo group, were all positive for the presence of *Staphylococcus*, in fact, pure cultures were obtained on each blood agar plate derived from the livers of these mice. While the liver data do not preclude the possibility that the mouse that died within the antibody-infused group died of *S. aureus* infection, the infection was not systemic, as it was in the
10 placebo group, and the mouse may have died for other reasons. Censoring of this antibody-infused mouse death results in a significant difference between antibody-infused and placebo treatments ($p = 0.015$, log-rank test). The data for the cross-challenge, where mice were infused with antibody generated after vaccination with ATCC 19636-derived proteins and challenged by *S. aureus*
15 strain 1477, also showed a protective trend. Between 7 and 14 days post challenge, all mice in the infused and non-infused groups began to develop necrotic skin lesions. However, gross examination of mice clearly revealed a visible delay in the formation of an observable lesion as well as the severity of the lesion between the groups. Infused mice developed lesions more slowly as
20 compared to non-infused control mice which developed lesion faster than infused mice and at a greater degree of severity. The infused mice healed faster than non-infused mice. This was clearly evident between 21 and 35 days post challenge. Gross examination of mice at 35 days post challenge showed that non-infused mice were severely disfigured and revealed a greater degree of
25 scarring. In fact, many of these mice lost normal posture, in that they appeared twisted in appearance, in contrast to infused mice that did not develop nearly the extensive scar tissue and/or disfigurement as illustrated by the twisted appearance that the non-infused mice developed. Overall, these data suggest that interperitoneal infusion of antibodies raised against *S. aureus* iron-induced
30 proteins can both protect against and limit the severity of *S. aureus* infection.

Example 10

Evaluation of a vaccine composition derived from *Staphylococcus aureus* in a chronically infected dairy herd

5 A commercial Dairy herd having a history of chronically high somatic cell counts attributable to *Staphylococcus aureus* was chosen for the evaluation of a vaccine composition as described in Example 1. The criterion for establishing vaccine efficacy of this experimental study was: 1) decreased incidence of clinical mastitis caused by *Staphylococcus aureus* among
10 vaccinates compared to non-vaccinated controls, 2) improvement (i.e., a decrease) in somatic cell count among vaccinates compared to controls and 3) decrease in culture positive isolation rates of *S. aureus* between vaccinated and non-vaccinated controls. Blood will be taken at the time of the first vaccination (day 0) and again at 3 and 6 weeks post initial immunization. Injection site
15 reactions or systemic reactions following vaccinations were monitored throughout the study. In addition, bulk tank milk samples were cultured and quantitatively enumerated to determine if there was a decrease in the number of CFU of *Staphylococcus aureus* cultured after vaccination.

 Three of the *Staphylococcus* isolates derived from the chronically
20 infected lactating cows within the herd were grown under conditions of iron-restriction and non-iron restricted conditions as described in Example 1. The three isolates were designated TTX101, TTX102, and TTX103. Extracted samples were examined by SDS-PAGE to compare banding profiles between isolates. Identical banding profiles were observed among isolates examined; the
25 compositions made from each isolate included proteins having molecular weights of 87.73 kDa, 80.46 kDa, 65.08 kDa, 54.53 kDa, 37.37 kDa, 35.70 kDa, 34.91 kDa, 33.0 kDa and 31.83 kDa. These proteins are the same molecular weights as previously described in Table 7. In addition, when comparing the isolates identical banding profiles were seen with those proteins that were
30 expressed in all conditions that were not regulated by iron: 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, and 40 kDa. These results were consistent with previous observations. One isolate designated as

TTX101 was chosen as the isolate to manufacture a composition to be used in this study.

5

Example 11

Vaccine preparation of *Staphylococcus aureus* (TTX101)

A composition was prepared as described in Example 1 using the isolate TTX101. The composition included proteins expressed under iron deplete
10 conditions having molecular weights of 87.73 kDa, 80.46 kDa, 65.08 kDa, 54.53 kDa, 37.37 kDa, 35.70 kDa, 34.91 kDa, 33.0 kDa, and 31.83 kDa as well as non-metal-regulated proteins having molecular weights of 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa 43kDa 41 kDa, and 40 kDa. The immunizing composition derived from strain TTX101 was used to prepare the
15 experimental vaccine by emulsifying the extracted protein suspension (400 µg total protein per milliliter) into a commercial adjuvant (EMULSIGEN, MVP Laboratories, Ralston NE) using an IKA Ultra Turrax T-50 homogenizing vessel (IKA, Cincinnati, OH) to give a final dose of 800 µg total protein in a 2.0 ml injectable volume with an adjuvant concentration of 22.5 % vol/vol. The
20 vaccine was administered subcutaneously 2 times at 21 day intervals.

Example 12

Experimental design and herd vaccination

25

Eighteen days before the first vaccination all lactating cows enrolled in the study (N=80) were tested for *S. aureus* by standardized aerobic bacteriological culture methods by culturing individual milk samples derived from each lactating cow. In addition, the Somatic Cell Counts (SCC) were
30 enumerated by the Dairy Herd Improvement Association using standard methods. Fourteen of the 80 cows were clinically diagnosed with mastitis and were culture positive for *S. aureus*. The remaining cows (N=66) tested negative for *S. aureus*. The eighty cows were equally divided into two groups designated

as group-1, vaccinated (N=40) and group-2, non-vaccinated (N=40). The fourteen clinically diagnosed Staphylococcus positive cows were equally distributed between both groups so that each study group contained 7 cows with clinical mastitis. The average SCC between groups prior to the first vaccination
5 was 203,219 in the non-vaccinated controls compared to 240,443 in vaccinates (not statistically different $p = 0.7$).

Eighteen days after the first sampling all cows in group 1 were vaccinated subcutaneously in the upper right shoulder with 2 ml of vaccine as described in Example 11. Ten days after the first vaccination milk samples
10 were taken at this time period by the DHIA for the enumeration of somatic cells from each individual cow. Milk samples were not bacteriologically tested at this time period for determining the presence of Staphylococcus. The difference in the SCC between groups at this time period was 125,241 (vaccinates) compared to 196,297 (controls). This was a 36% difference in the number of
15 somatic cells between vaccinates as compared to non-vaccinated controls. The difference in the SCC between the controls and vaccinates at this sampling period was not statistically different ($p = 0.5$). The lack of statistical difference in the SCC between groups at both sampling periods was due to the large variation in individual SCC between cows. The injection site of each
20 vaccinated cow was also examined at this same time period. None of the cows examined showed any adverse tissue reaction at the site of injection by physical examination. In addition, there was no measurable loss in milk production due to vaccination.

Twenty one days after the first vaccination all cows in group-1
25 (vaccinates) were given their second vaccination or booster. During the time period between first and second vaccination, cows in both groups (vaccinates and controls) developed teat damage due to a dramatic drop in the environmental temperature resulting in the formation of lesions at the end of the teat, resulting in the development of infected teats and potentially increasing the
30 isolation of Staphylococcus during sampling, which was observed at the third sampling period. Twenty three days after the second vaccination milk samples were taken by the DHIA for the enumeration of Somatic Cells from each individual cow. Milk samples were also bacteriologically tested for the

presence of *Staphylococcus aureus*. There was a dramatic increase in isolation rate of *S. aureus* at this time period in the cows that tested negative at the first sampling period. In the non-vaccinated controls 42.9% of these cows now tested positive for *S. aureus*, in contrast to the vaccinates, which only showed
5 and increase of 35.5 %. This was a 7.4 % difference between vaccinates as compared to the non vaccinated controls. It's difficult to say that the improvement in the isolation rate of *S. aureus* in the vaccinated group was due to the effect of the vaccine alone. One cannot overlook the difficulty in obtaining clean milk samples from cows that had teat damage which could
10 increase the potential contamination of the milk by *S. aureus* when obtaining the sample. Nevertheless, there was a significant difference in the average SCC between vaccinates compared to controls. The average SCC of the vaccinated group was 222,679 compared to 404,278 somatic cells as measured in the control group. This was a 44.9 % difference between vaccinates when
15 compared to the non vaccinated controls. It's interesting to speculate that the difference seen in the SCC between these groups also coincides with the difference in the isolation rate of *S. aureus* between groups. However, due to the large variation in SCC between individual animals and the small sample size of the experimental trial in the number of animals the difference was not
20 statistically different ($p = 0.28$).

At this same time period the injection site of each vaccinated cow was examined for any adverse tissue reaction that may have been caused by the vaccine composition. None of the cows examined showed any adverse reaction at the site of injection by physical examination. The vaccine compositions
25 appeared to be highly tissue compatible and caused no measurable loss in milk production after each vaccination.

Monitoring of the cows is continued by measuring SCC and milk samples for the presence or absence of *Staphylococcus aureus*. Some of the cows of each group are vaccinated a third time at 42 days after the second
30 vaccination. There appears to be a difference favoring the use of the vaccine composition for decreasing somatic cell counts and controlling infection caused by *Staphylococcus aureus*. Further monitoring includes serology based on antibody titers to the vaccine composition, changes in milk production in

vaccinated cows due the improvement in health, and decreased SCC of vaccinated animals compared to non-vaccinated cohorts. In addition, other experiments are conducted to investigate the protective index of the vaccine based on dose response following challenge with a virulent *S. aureus*.

5

Example 13

Since the molecular weights of the proteins among the different *S.*
10 *aureus* strains have been demonstrated to be similar and since heterologous protection was observed in the mouse challenge study, we sought to determine if the proteins sharing similar molecular weights in Figure 1 were similar proteins. The technique chosen to characterize the proteins was matrix-assisted laser desorption/ionization mass spectrometry (MALDI-MS). A portion of the
15 composition was resolved using SDS-PAGE as described in Example 1, and the gel was stained with Coomassie Brilliant blue to visualize the proteins.

Materials and Methods

Excision and washing. The gel was washed for 10 minutes with water
20 twice. Each protein band of interest was excised by cutting as close to the protein band as possible to reduce the amount of gel present in the sample.

Each gel slice was cut into 1x1 mm cubes and placed in 1.5 ml tube. The gel pieces were washed with water for 15 minutes. All the solvent volumes used in the wash steps were approximately equal to twice the volume of the gel
25 slice. The gel slice was next washed with water/acetonitrile (1:1) for 15 minutes. When the proteins had been stained with silver, the water/acetonitrile mixture was removed, the gel pieces dried in a SPEEDVAC vacuum concentrator/dryer (ThermoSavant, Holbrook, NY) and then reduced and alkylated as described below. When the gel pieces were not silver-stained, the
30 water/acetonitrile mixture was removed, and acetonitrile was added to cover until the gel pieces turned a sticky white, at which time the acetonitrile was removed. The gel pieces were rehydrated in 100 mM NH_4HCO_3 , and after 5 minutes, a volume of acetonitrile equal to twice the volume of the gel pieces

was added. This was incubated for 15 minutes, the liquid removed, and the gel pieces dried in a SPEEDVAC.

Reduction & alkylation. The dried gel pieces were rehydrated in 10mM DTT and 100 mM NH_4HCO_3 , and incubated for 45 minutes at 56°C. After
5 allowing the tubes to cool to room temperature, the liquid was removed and the same volume of a mixture of 55mM iodoacetamide and 100 mM NH_4HCO_3 was immediately added. This was incubated for 30 minutes at room temperature in the dark. The liquid was removed, acetonitrile was added to cover until the gel pieces turned a sticky white, at which time the acetonitrile was removed. The
10 gel pieces were rehydrated in 100 mM NH_4HCO_3 , and after 5 minutes, a volume of acetonitrile equal to twice the volume of the gel pieces was added. This was incubated for 15 minutes, the liquid removed, and the gel pieces dried in a Speed vac. If the gel was stained with coomassie blue, and residual coomassie still remained, the wash with 100 mM NH_4HCO_3 /acetonitrile was repeated.

In-gel digestion. Gel pieces were completely dried down in a Speed
15 Vac. The pieces were rehydrated in digestion buffer (50 mM NH_4HCO_3 , 5 mM CaCl_2 , 12.5 nanograms per microliter (ng/ μl) trypsin) at 4°C. Enough buffer was added to cover the gel pieces, and more was added as needed. The gel pieces were incubated on ice for 45 minutes, and the supernatant removed and
20 replaced with 5-2 μl of same buffer without trypsin. This was incubated at 37°C overnight in an air incubator.

Extraction of peptides. A sufficient volume of 25 mM NH_4HCO_3 was added to cover gel pieces, and incubated for 15 minutes (typically in a bath sonicator). The same volume of acetonitrile was added and incubated for 15
25 minutes (in a bath sonicator if possible), and the supernatant was recovered. The extraction was repeated twice, using 5% formic acid instead of NH_4HCO_3 . A sufficient volume of 5% formic acid was added to cover gel pieces, and incubated for 15 minutes (typically in a bath sonicator). The same volume of acetonitrile was added and incubated for 15 minutes (typically in a bath
30 sonicator), and the supernatant was recovered. The extracts were pooled, and 10 mM DTT was added to a final concentration of 1 mM DTT. The sample was dried in a SPEEDVAC vacuum concentrator/dryer to a final volume of approximately 5 μl .

Desalting of peptides. The samples were desalted using a ZIPTIP pipette tips (C18, Millipore, Billerica, MA) as suggested by the manufacturer. Briefly, a sample was reconstituted in reconstitution solution (5:95 acetonitrile:H₂O, 0.1% - 0.5% trifluoroacetic acid), centrifuged, and the pH
5 checked to verify that it was less than 3. A ZIPTIP was hydrated by aspirating 10 µl of solution 1 (50:50 acetonitrile:H₂O, 0.1% trifluoroacetic acid) and discarding the aspirated aliquots. This was followed by aspirating 10 µl of solution 2 (0.1% trifluoroacetic acid in deionized H₂O) and discarding the aspirated aliquots. The sample was loaded into the tip by aspirating 10 µl of the
10 sample slowly into the tip, expelling it into the sample tube, and repeating this 5 to 6 times. Ten microliters of solution 2 was aspirated into the tip, the solution discarded by expelling, and this process was repeated 5-7 times to wash. The peptides were eluted by aspirating 2.5 µl of ice cold solution 3 (60:40, acetonitrile:H₂O, 0.1% trifluoroacetic acid), expelling, and then re-aspirating
15 the same aliquot in and out of the tip 3 times. After the solution has been expelled from the tip, the tube is capped and stored on ice.

Mass spectrometric peptide mapping. The peptides were suspended in 10 µl to 30 µl of 5% formic acid, and analyzed by MALDI-TOF MS (Bruker Daltonics Inc., Billerica, MA). The mass spectrum of the peptide fragments
20 was determined as suggested by the manufacturer. Briefly, a sample containing the peptides resulting from a tryptic digest were mixed with matrix cyano-4-hydroxycinnamic acid, transferred to a target, and allowed to dry. The dried sample was placed in the mass spectrometer, irradiated, and the time of flight of each ion detected and used to determine a peptide mass fingerprint for each
25 protein present in the composition. Known polypeptides were used to standardize the machine.

Data analysis. The experimentally observed masses for the peptides in each mass spectrum were compared to the expected masses of proteins using the Peptide Mass Fingerprint search method of the Mascot search engine (Matrix
30 Science Ltd., London, UK, and www.matrixscience.com, see Perkins et al., Electrophoresis 20, 3551-3567 (1999)). The search parameters included: database, MSDB or NCBI nr; taxonomy, bacteria (eubacteria) or Firmicutes (gram-positive bacteria); type of search, peptide mass fingerprint; enzyme,

trypsin; fixed modifications, carbamidomethyl (C) or none; variable
 modifications, oxidation (M), carbamidomethyl (C), the combination, or none;
 mass values, monoisotopic; protein mass, unrestricted; peptide mass tolerance,
 between ± 150 ppm and ± 430 ppm, or ± 1 Da; peptide charge state, Mr; max
 5 missed cleavages, 0 or 1; number of queries, 20.

Results

The result of this search was a mass fingerprint for each protein present
 in the composition is shown in Tables 2, 3, 4, and 5.

10

Example 14

Identification of iron-regulated protein families using microarray-based gene
 expression analysis of *S. aureus* grown under low iron conditions

15

For microarray analysis, bacteria were cultured in chemically defined
 media (CDM) made from individual stock solutions (Table 10).

Table 10: **Chemically defined medium (CDM) for *Staphylococcus aureus***

20

	[Final]	Stock composition	Add to 1 L
	g/L	g/500 ml	50 ml
	Salts (20X)		
	K ₂ HPO ₄	7	70
25	KH ₂ PO ₄	2	20
	Na ₃ citrate	1.47	14.7
	(NH ₄) ₂ SO ₄	1	10
	Carbohydrate (40X)	g/L	g/500 ml
30	Glucose	5	100
	Vitamins (1000X)	mg/L	mg/100 ml
	Thiamine	1	100
35	Nicotinic acid	0.5	50
	Biotin	0.005	dilution*
	Calcium pantothenate	0.25	25

	Nucleotides (100X)	mg/L	mg/100 ml*	10 ml
	<i>*Dissolve in 100 ml 2N HCl</i>			
	Adenine	5	50	
	Guanine	5	50	
5	Cytosine	5	50	
	Uracil	5	50	
	Thymine	10	200	
	Micronutrients (1000X)	μM	mg/100 ml stockAB*	1 ml
10	<i>*Make stocks A and B, then add 1 ml of each to 98 ml ddH2O to make final stock</i>			
	(A)			
	CaCl ₂	0.5	735	
	H ₃ BO ₃	0.5	309	
15	CoCl ₂	0.05	118	
	(NH ₄) ₆ Mo ₇ O ₂₄	0.005	62	
	(B)			
	CuSO ₄	0.1	125	
	MnSO ₄	0.1	169	
20	ZnSO ₄	0.05	144	
	Individual			
	MgSO ₄	100		
	FeSO ₄ or other	10-50		
	Amino acids (200X)	mg/L	g/100 ml	5 ml
25	<i>*Autoclave unless otherwise noted</i>			
	<i>Refrigerate</i>			
	Aspartic acid (0.1 M HCl)	90	1.8	
30	Proline	80	1.6	
	Alanine	60	1.2	
	Histidine	20	0.4	
	Valine	80	1.6	
	Arginine	50	1.0	
35	Serine	30	0.6	
	Methionine	3	0.06	
	Isoleucine	30	0.6	
	<i>Dark refrigerated filtered</i>			
40	Tryptophan	10	0.2	
	Tyrosine (0.5 M NaOH)	50	1.0	
	<i>Room temperature</i>			
	Glutamic acid	100	2.0	
45	Leucine	90	1.8	
	Phenylalanine	40	0.8	
	Glycine	50	1.0	
	Threonine	30	0.6	
	Lysine	50	1.0	

Fresh daily

Cysteine

20

0.4

- 5 **METHOD OF MAKING.** For making iron-deplete media, combine all stock solutions except cations and micronutrients and bring to proper volume in volumetric flask using MilliQ purified water and leaving enough void volume to accommodate cation additions. Add 15 g CHELEX resin per 1 L media and stir at room temperature for at least 2 hours. Filter solution into sulfuric acid (10%)
 10 treated glass bottle using 2 μ m bottle-top filter. Add filtered cation stock solutions and store at 4°C in the dark for up to 2 weeks.

- S. aureus* strains RF122 (isolated from bovine mastitis) and MSA553 (isolated
 15 from human toxic shock syndrome) were used. Both isolates were streaked on tryptic soy broth agar directly from secondary-passage freezer stocks prior to use in experiments. CDM contained final citrate concentrations approximately analogous to that in bovine milk (5 mM). For iron-free CDM, the following components were combined (0.998 L total volume) and added to 15 g CHELEX
 20 resin (BioRad Laboratories, Hercules, CA), then stirred for 1.5 hours at room temperature: salt, glucose, amino acids, vitamins, and nucleotides. The deferrated base media was then filtered using 2 μ m bottle-top filters (Nalgene Nunc International, Rochester NY) into sulfuric acid-treated bottles, after which micronutrients and 100 μ M MgCl₂ (both solutions prepared in acid-treated
 25 glassware with MilliQ water) were added. CDM was stored at 4°C in the dark until use.

- A single bacterial colony was inoculated into 3 ml iron-deplete CDM in a 25 ml acid-treated glass culture tube and shaken overnight at 250 rpm in a 37°C incubator. One milliliter of the subculture was then used to inoculate 500
 30 ml of CDM in a 2500 ml Erlenmeyer flask the following day. Cultures were incubated at 37°C with shaking at 250 rpm. Iron-deplete CDM cultures took approximately twice as long as CDM + 50 μ M FeSO₄ to reach an OD of 1.0 (18 hrs versus 36 hours). At mid-log phase (OD = 0.600), 4x100 ml culture aliquots were distributed into 500 ml Erlenmeyer flasks and allowed to shake in
 35 the incubator for 10 minutes prior to the addition of experimental iron solutions. To one flask, 300 μ l bovine lactoferrin (50 mg/ml, Sigma-Aldrich, St. Louis,

MO) were added for a final concentration of 150 µg/ml. To another flask, 50 µl ferric citrate (100 mM) were added. The remaining two control flasks received no supplements. At 5, 30, 60 and 120 minutes, 7.5 ml of culture were collected and added to 5 ml guanidine thiocyanate solution containing β-mercaptoethanol and 0.5% sodium lauryl sarcosine. Solutions were mixed thoroughly to halt transcription and centrifuged at 4,000 x g for 8 minutes at 8°C; supernatant was poured off and cells were frozen in 250 µl Trizol (Invitrogen) using an ethanol/dry ice bath, then stored at -80°C until RNA extraction.

10 For RNA extraction, cell pellets were thawed on ice and 750 µl Trizol (Invitrogen, Carlsbad CA) were added. Cells were resuspended by vortex and the slurry was transferred to a 2 ml screw cap tube containing 0.1 mm silica-zirconium beads, then beat 3 x 2 minutes in a BeadBeater (Biospec Products, Inc., Bartlesville, OK) with ice incubation between repetitions. Slurries were
15 incubated an additional 20 minutes at room temperature, followed by centrifugation to pellet beads and cellular components. 400 µl chloroform were added and mixed by inversion, incubated for 10 minutes at room temperature and tubes were centrifuged 8 minutes at 12,000 X g at 8°C. The aqueous layer was removed and the RNA precipitated with 400 µl isopropanol followed by
20 washing with 1 ml 70% ethanol. Clear RNA pellets were air-dried briefly and resuspended in 100 µl RNase-free H₂O. DNA was digested using standard DNase kit (Qiagen, Valencia CA) followed by cleanup according to manufacturer's recommendations on RNeasy columns (Qiagen). Finally, a Turbo DNA-free kit (Ambion, Austin TX) was used to ensure elimination of
25 DNA from the preparation. RNA was measured on a spectrophotometer and run on an Agilent Bioanalyzer (Agilent, Palo Alto CA) to verify quality and quantity prior to the generation of cDNA for microarray hybridization.

Microarray analysis was carried out according to established protocols. The array, featuring 3841 70mer oligonucleotides (Illumina, San Diego CA) representing open reading frames (ORFs) from nine sequenced *S. aureus*
30 genomes including RF122 and MSA553, was spotted in triplicate on Gaps II aminosaline coated slides (Corning, Acton MA) using a BioRobotics Microgrid II Array Spotter (BioRobotics, Cambridge UK). Slides were rehydrated, UV

cross-linked and stored under dessication. Immediately prior to hybridization, slides were incubated for 1 hour at 42°C in prehybridization buffer consisting of 25 ml formamide, 12.5 ml 20X SSC, 12 ml dH₂O, 500 µL 10% SDS and 0.5 g BSA. Slides were rinsed with 2 L MilliQ water and dried by centrifugation. To
5 prepare samples, 8-10 µg total bacterial RNA were incubated with 20 µg random hexamers at 70°C for 10 minutes, followed by reverse transcription with amino-allyl incorporation using Superscript II (Invitrogen) and amino-allyl coupled dUTP (Sigma). Labeled cDNA was neutralized, purified, dried and resuspended with Cy3 or Cy5 fluorescent dyes (Amersham Biosciences Corp.,
10 Piscataway NJ); coupling proceeded for 2 hours. Fluorescently labeled cDNA samples (12 µl each) were washed using a Qiagen PCR purification kit, combined and added to 9.8 µl formamide, 6.8 µL 20X SSC, 3.4 µl salmon sperm DNA (10 mg/ml, Invitrogen) and 1 µl 10% SDS. Samples were incubated for 2 minutes at 99.9°C in a thermal cycler and allowed to cool prior
15 to array application. Probes were then applied to array, covered with a glass coverslip and incubated overnight in a 42°C waterbath. Slides were washed thoroughly in diluted SSC buffers after 12-16 hours of incubation and scanned using an Axon 4100B Scanner and Axon GenePix Software (Axon Instruments, Union City CA). Raw intensity data were exported to GeneSpring (Agilent
20 Technologies, [Silicon Genetics], Palo Alto CA) for normalization and filtering. Spots were globally normalized, filtered based on minimum (>1500) raw intensity values and the triplicates were averaged. Each experiment was run twice and a single slide was run for each using a dye-swap between matching timepoints. Thus, at least 6 dye-swapped datapoints were generated for each
25 gene at each timepoint, representing at least 2 biological replicates. Data were further analyzed by hierarchical clustering (Euclidian distance, average linkage, UPGMA) and K-means clustering (uncentered correlation based measured distance) using EPCLUST (Jaak Vilo, EBI) and SpotFire (SpotFire, Somerville, MA). Significance Analysis for Microarrays (SAM, (157)) was used on
30 median-centered log ratios using the one-class model across all timepoints to determine if expression of the gene differed significantly from zero. Stringent delta values were used so that the percentage of false positives was estimated to

be zero. A summary of operons showing similar up or downregulation by SAM analysis is shown in Table 11, supporting the ability of the arrays to detect biological responses.

- 5 Table 11. Operonic clusters with coordinated transcriptional responses identified using microarray analysis of gene expression of *S. aureus*

<u>Operon ID</u>	<u>Function</u>	<u>Response</u>	<u>No. of coordinately expressed probes</u>	
			<u>Contiguous unregulated genes</u>	<u>Functionally related but not contiguous</u>
Sir	cation transport	upregulated in low iron	3	0
Fhu	cation transport	upregulated in low iron	3	1
Opp	oligopeptide transport	upregulated in low iron	9	N/A
Mnt	cation transport	upregulated in low iron	3	1
Pfl (Formate acetyltransferase)	fermentation	upregulated in presence of lactoferrin	2	4

- 10 For standard cloning of proteins, the appropriate genes were amplified from DNA extracted from *S. aureus* (strain ATCC19636) by standard polymerase chain reaction. Primers were designed to incorporate StuI and KpnI restriction endonuclease sites and are shown below.

- 15 Table 12. Cloning Primers:

Gene (primer)	Primer Sequence	SEQ ID NO
Pflb (5' to 3')	GCAGGCCTTTAGAAACAAATAAAAAATCATG	507
Pflb (3' to 5')	TATGGTACCTTACATACTTTCATGGAATGTACG	508
Opp1A (5' to 3')	GCAGGCCTAAAAAGAAAACAAGCAATTAA	509
Opp1A (3' to 5')	TATGGTACCTTATTTATACTGCATTTCATTGAA	510
SirA (5' to 3')	GCAGGCCTTCATCTGATAGCA AAGATAAGG	511
SirA (3' to 5')	TATGGTACCTTATTTTGATTGTTTTCAATAATT	512
SYN2 (5' to 3')	GCAGGCCTAAAGAATCATCAACTAAA	513
SYN2 (3' to 5')	TATGGTACCCTTTTGTCTTTTTTTGA	514

FhuD (5' to 3')	GCAGGCCTACTGAAGAGAAAACTGAAATGA	515
FhuD (3' to 5')	TATGGTACCTTATTTTGCTTTTCTGCAATTT	516
SYN1 (5' to 3')	GCAGGCCTGGTAGCGACGATAATGGCTCGT	517
SYN1 (3' to 5')	TATGGTACCTTATTTTCTATAAATTGCATCTCT	518
MntC (5' to 3')	GCAGGCCTAGTGATAAGTCAAATGGCAAATA	519
MntC (3' to 5')	TATGGTACCTTATTTTCATGCTTCCGTGTACAG	520
SstD (5' to 3')	GCAGGCCTTCAGAACTAAAGGTTCTAAAGAT	521
SstD (3' to 5')	TATGGTACCTTATTTTACAACCTTTTCAAGTT	522
FhuD2 (5' to 3')	GCAGGCCTACTAAATCTTATAAAATGGACGAT	523
FhuD2 (3' to 5')	TATGGTACCTTATTTTGCAGCTTAATTAATT	524

DNA extracted from *S. aureus* ATCC19636 was used as the template. DNA amplicons were verified by gel electrophoresis and the amplified bands of DNA were excised, purified, digested and ligated into cut pQE30-Xa vector, transformed into competent XL-1 *E. coli* and screened for ampicillin resistance. Resistant clones were screened for plasmid inserts using colony PCR.

10

Example 15

Screening of immunoreactivity of protective protein candidates

To evaluate the antibody reactivity of the proteins identified from MALDI-TOF analysis (Example 13) and/or microarray and genomic analysis (Example 14), a two-part screen was used to evaluate individually expressed staphylococcal proteins. The rapid first screen used transcriptionally active PCR fragments to survey antibody binding to small amounts of candidate protein expressed using a cell-free *E. coli* lysate. The second screen used standard PCR-based cloning, expression and purification of proteins in *E. coli* using a commercial vector (pQE30Xa, Qiagen, Valencia CA) in order to validate positive candidates from the first screen. The second screen also generated master seed stocks of *E. coli* host cells containing the expression

20

vector corresponding to each immunoreactive protein candidate for production and purification of sufficient protein for vaccination and experimentation.

A high-throughput method for generating individual SIRP antigens was used to test several candidate genes encoding *S. aureus* proteins involved in metal metabolism. This method generates a transcriptionally active PCR (TAP) amplicon using a 2-step PCR reaction with primers that add a promoter, terminator and C-terminal His₆ tag. The resulting transcriptionally active amplicons were used as a template for protein production in a cell-free *in vitro* transcription/translation reaction consisting of *E. coli* cell lysate, amino acids and buffers. The two-step PCR reaction required a first set of primers specific to the gene of interest that also included a linker sequence matching the second set of primers. Each primer set for the first step of PCR was designed to exclude membrane-processing signal sequences to prevent integration into the cell membrane and are shown below.

Table 13. TAP Primers:

Gene (primer)	Primer Sequence	SEQ ID NO
Pflb (5' to 3')	AGAAGGAGATATACCATGTTAGAAACAAAT	525
Pflb (3' to 5')	TTAATGATGATGATGATGATGCATACTTTCATG	526
Opp1A (5' to 3')	AGAAGGAGATATACCATGAGAAAATACT	527
Opp1A (3' to 5')	TTAATGATGATGATGATGATGTTTATACTGCAT	528
SirA (5' to 3')	ATAAGGAGATATACCATGAATAAAGTAATT	529
SirA (3' to 5')	TTAATGATGATGATGATGATGTTTTGATTGTTT	530
SYN2 (5' to 3')	AGAAGGAGGATATACCATGAGAGGTCTAAAACTTTT	531
SYN2 (3' to 5')	TTAATGATGATGATGATGATGCTTTTGTCTTTTTTTGA	532
FhuD (5' to 3')	AGAAGGAGGATATACCATGAATAGGAATATCGTTAAA	533
FhuD (3' to 5')	TTAATGATGATGATGATGATGTTTTGCTTTTCTGCAAT	534
SYN1 (5' to 3')	AGAAGGAGGATATACCATGAAGAAATCGTTAATTGCT	535
SYN1 (3' to 5')	TTAATGATGATGATGATGATGTTTTCTATAAATTGCATC	536
MntC (5' to 3')	AGAAGGAGATATACCAAAAAATTAGTA	537
MntC (3' to 5')	TTAATGATGATGATGATGATGTTTCATGCTTCC	538
SstD (5' to 3')	AGAAGGAGATATACCATGAAGAAAACAGTC	539

SstID (3' to 5')	TTAATGATGATGATGATGATGTTTTACAACCTT	540
FhuD2 (5' to 3')	AGAAGGAGATATACCATGAAAAAACTATTA	541
FhuD2 (3' to 5')	TTAATGATGATGATGATGATGTTTTGCAGCTT	542

A standard 50 μ l PCR reaction was performed using (1 unit High Fidelity Taq DNA polymerase (Invitrogen), 0.2 μ M primers, 2 mM dNTP (each), 2 mM final Mg⁺⁺, and approximately 5 ng of starting DNA template, buffered to 60 mM TrisSO₄ (pH 8.9) and 18 mM ammonium sulfate). The PCR cycling protocol included 1 minute of initial denaturation at 94°C, followed by 30 cycles as follows: Denature/94°C/30s; Anneal/55°C/30s; Extend/68°C, 90s. Identical primers with the appropriate overlap were utilized for the second step PCR reaction and were supplied by the manufacturer (Genlantis). The resulting DNA PCR product was purified to eliminate residual primer, salt and DNA fragments and used as a template for a second reaction with a standard set of primers to add the promoter and terminator sequences using similar conditions. The DNA template was then purified and added to an *E. coli* cell-free Rapid Translation System RTS 100 reaction mix (Roche) containing 12 μ l of *E. coli* lysate, 12 μ l of amino acids, 10 μ l of reaction mix, 1 μ l of added methionine, 5 μ l of reconstitution buffer and 10 μ l of purified DNA template from the two-step PCR reaction. Following incubation for 5 hours at 30°C, one microliter of each protein sample (approximately 0.5 μ g/ μ l total protein) was applied to polyvinylidene fluoride (PVDF) membrane after methanol saturation. The blot was blocked overnight with 5% NFDM/TTBS, incubated with iron-restricted protein, enhanced (IRPE) hyperimmunized mouse sera diluted 1:500 or anti-His₆ antibody (1:500), washed, incubated with secondary goat-anti mouse alkaline phosphatase (AP) conjugate (1:3000), washed and developed chromogenically (Bio-Rad AP color development kit) for 20 minutes. Lysates containing seroreactive polypeptides were identified.

Clones produced as described in Example 14 were grown to mid-log phase, induced with 1 mM IPTG and grown for four hours. Cells were pelleted, washed, and lysed in boiling SDS-PAGE loading buffer. Crude lysates were separated by SDS-PAGE and Coomassie stained. A second set of separated

proteins was transferred to PVDF membrane and immunblotted with IRPE vaccine hyperimmunized mouse sera diluted 1:500 in 1% NFDM/TTBS. The blot was washed, incubated with alkaline phosphatase (AP)-conjugated goat anti-mouse secondary antibody, washed, and developed with chromogenic substrate.

Example 16

Preparation of immunizing compositions from recombinantly-produced polypeptides

In order to isolate recombinant *S. aureus* polypeptides for formulating a vaccine, the *E. coli* clones described in Example 14 were grown at 37°C (225 rpm) to mid-log phase ($OD_{600} = 0.4-0.6$) in 1L of Trypic Soy Broth and then induced for 4 hours with 1 mM isopropyl β -D-1-thiogalactopyranoside (IPTG). Cultures were pelleted for 10 minutes at 4°C in a Sorvall centrifuge (4000 x g) and frozen at -80°C prior to undergoing the purification procedure. Bacterial pellets were then processed by two different methods, depending on the solubility of the over-expressed *S. aureus* polypeptide.

For soluble polypeptides (e.g., MntC, FhuD, SYN2, SirA, or SYN1), bacterial pellets were resuspended in 25 ml of BUGBUSTER reagent (Novagen) and subjected to 15 minutes of sonication on ice using a Branson sonifier fitted with a microtip (65% duty cycle, 5 output). Insoluble material was removed by 10 minutes of centrifugation (4000 x g). The soluble supernatant was filtered (0.2 μ m) and subjected to metal affinity chromatography (Ni-NTA His-Bind, Novagen) according to the instructions provided by the manufacturer.

For insoluble polypeptides (e.g., PflB, or Opp1A), bacterial pellets were resuspended in 25 ml of BUGBUSTER reagent (Novagen), placed on a rocker platform for 10 minutes, and then subjected to centrifugation (15,000 x g for 12 minutes). The resulting pellet was resuspended in 10 ml of BUGBUSTER plus 20 ml of diluted BUGBUSTER (1:10 in PBS) and subjected to centrifugation (5000 x g for 12 minutes). The pellet was then resuspended in 20 ml of diluted BUGBUSTER and subjected to a final step of centrifugation (15,000 x g for 12

minutes). The final pellet was resuspended in 10 ml of Buffer A (0.1 M NaH_2PO_4 , 0.01 Tris-HCl, 8M Urea, pH 8.0) and incubated for 10 minutes on a rocking platform at room temperature. The samples were then subjected to centrifugation (12,000 x g for 20 minutes), and the resulting supernatant was separated by metal affinity chromatography (Ni-NTA His-Bind, Novagen) according to the instructions provided by the manufacturer, but with the following modifications. After charging the column, 10 ml of Buffer A was used to equilibrate the resin. Following the binding of polypeptide, the column was washed with 15 ml of Buffer B (0.1 M NaH_2PO_4 , 0.01 Tris-HCl, 8M Urea, pH 6.0), and eluted using 15 ml of Buffer C (0.1 M NaH_2PO_4 , 0.01 Tris-HCl, 8M Urea, pH 4.5).

Isolated recombinant polypeptides were eluted from the columns in a volume of 15 ml and placed in 20kDa cutoff dialysis cassettes (Pierce) for dialysis against 2L of phosphate-buffered saline (PBS). Following three buffer changes over the course of 30 hours, the polypeptides were removed, filtered (0.2 μm), and concentrated to 2-3 ml of volume using 20kDa cutoff Centricon devices (Millipore). The concentrations of the purified polypeptides were determined using the standard BCA method (Pierce).

10 μg of each polypeptide was combined, and the volume adjusted to 100 μl with PBS, to form an immunizing composition.

Example 17

Mouse vaccination

25

IV Challenge (Study A)

Fifty (N=50) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams were distributed into 3 groups (10-20 mice/group). Mice were housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water were supplied *ad libitum* to all mice. All vaccines were formulated with 50% IFA as an adjuvant. Mice were vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

Group 1: Placebo, vaccinated with ovalbumin (70 µg/100 µl) (Placebo, 20 mice).

Group 2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (70 µg/100 µl) (SIRP Extract, 20 mice).

5 Group 3: Vaccinated with recombinant polypeptide MntC (10 µg/100 µl) (rMntC, 10 mice).

IP Challenge (Study B)

Forty (N=40) female BALB/C mice obtained from Harlan Breeding
10 Laboratories (Indianapolis, IN) weighing 16-22 grams were equally distributed into 4 groups (10 mice/group). Mice were housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water were supplied *ad libitum* to all mice. All vaccines were formulated with 50% IFA as an adjuvant. Mice were
15 vaccinated subcutaneously with 0.1 ml of the appropriate composition at 14 day intervals using either two vaccinations (Groups 1-3) or three vaccinations (Group 4) as follows:

Group 1: Placebo, vaccinated twice with ovalbumin (70 µg/100 µl) (Placebo).

Group 2: Vaccinated twice with ATCC 25904 proteins expressed under
20 iron-restriction (70 µg/100 µl) (SIRP Extract).

Group 3: Vaccinated twice with recombinant polypeptides PflB, Opp1A, SirA, SYN2, FhuD, SYN1, and MntC (each 10 µg/100 µl, total protein 70 µg/100 µl) (rSIRP7 (2x)).

Group 4: Vaccinated three times with recombinant polypeptides PflB,
25 Opp1A, SirA, SYN2, FhuD, SYN1, and MntC (each 10 µg/100 µl, total protein 70 µg/100 µl) (rSIRP7 (3x)).

IC Challenge (Study C)

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding
30 Laboratories (Indianapolis, IN) weighing 16-22 grams were equally distributed into 3 groups (10 mice/group). Mice were housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water were supplied *ad libitum* to all

mice. Mice were vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

Group 1: Placebo, vaccinated with ovalbumin (70 µg/100 µl) (Placebo).

Group 2: Vaccinated with ATCC 25904 proteins expressed under iron-
5 restriction (70 µg/100 µl) (SIRP Extract).

Group 3: Vaccinated with recombinant polypeptides PflB, Opp1A, SirA, SYN2, FhuD, SYN1, and MntC (each 10 µg/100 µl, total protein 70 µg/100 µl) (rSIRP7).

10

Example 18

Preparation of challenge organism

IV Challenge (Study A)

15 *Staphylococcus aureus* strain ATCC 25904 was used as a challenge organism. Briefly, a 1 µl loop of bacteria from a frozen glycerol stock grown in standard TSB (no iron restriction) was used to inoculate a 20 ml culture of TSB and incubated at 37°C for 18 hours. 2.5 ml of this culture were passaged into 500 ml of fresh TSB. The culture was incubated at 37°C for approximately
20 two hours while rotating at 250 rpm until an optical density (OD₆₀₀) of 0.4 (Absorbance) was reached (mid-log phase), then the cells were centrifuged at 10,000 x g for 10 minutes at 4°C to pellet the bacteria. The bacterial pellet was washed by centrifugation in PBS at 4°C. The final pellet was resuspended in 20 ml of PBS. The final challenge dose was prepared by adding an aliquot of this
25 concentrated bacterial culture to PBS to generate a solution with an OD₆₀₀ of 4.0 (A), corresponding to approximately 6.67×10^8 CFU/ml. Just prior to challenge, 1 ml of these bacterial suspensions was serially diluted and plated on agar to enumerate the number of colony-forming units (CFU) per mouse dose.

30 IP Challenge (Study B)

Staphylococcus aureus strain ATCC 25904 was used as a challenge organism. Briefly, a 1 µl loop of bacteria from a frozen glycerol stock grown in standard TSB (no iron restriction) was used to inoculate a 20 ml culture of

TSB and incubated at 37°C for 18 hours. 2.5 ml of this culture were passaged into 500 ml of fresh TSB. The culture was incubated at 37°C for approximately two hours while rotating at 250 rpm until an optical density (OD₆₀₀) of 0.4 (Absorbance) was reached (mid-log phase), then the cells were centrifuged at 5 10,000 x g for 10 minutes at 4°C to pellet the bacteria. The bacterial pellet was washed by centrifugation in PBS at 4°C. The final pellet was resuspended in 20 ml of PBS. The final challenge dose was prepared by adding an aliquot of this concentrated bacterial culture to PBS to generate a solution with an OD₆₀₀ of 6.0 (A), corresponding to approximately 3.33×10^9 CFU/ml. Just prior to 10 challenge, 1 ml of these bacterial suspensions was serially diluted and plated on agar to enumerate the number of colony-forming units (CFU) per mouse dose.

IV Challenge (Study C)

Staphylococcus aureus strain ATCC 25904 was used as a challenge 15 organism. Briefly, a 1 µl loop of bacteria from a frozen glycerol stock grown in standard TSB (no iron restriction) was used to inoculate a 20 ml culture of TSB and incubated at 37°C for 18 hours. 2.5 ml of this culture were passaged into 500 ml of fresh TSB. The culture was incubated at 37°C for approximately two hours while rotating at 250 rpm until an optical density (OD₆₀₀) of 0.4 20 (Absorbance) was reached (mid-log phase), then the cells were centrifuged at 10,000 x g for 10 minutes at 4°C to pellet the bacteria. The bacterial pellet was washed by centrifugation in PBS at 4°C. The final pellet was resuspended in 20 ml of PBS. The final challenge dose was prepared by adding an aliquot of this concentrated bacterial culture to PBS to generate a solution with an OD₆₀₀ of 4.0 25 (A), corresponding to approximately 6.67×10^8 CFU/ml. Just prior to challenge, 1 ml of these bacterial suspensions was serially diluted and plated on agar to enumerate the number of colony-forming units (CFU) per mouse dose.

30

Example 19
Challenge

IV Challenge (Study A)

5 Fourteen days after the second vaccination, mice in all groups (1-3) were intravenously challenged in the lateral tail vein with 0.3 ml of the appropriate organism. The three groups of mice were challenged identically with 2×10^8 CFU of *S. aureus* strain ATCC 25904 per mouse. Mortality was recorded daily for 10 days after challenge.

10 When comparing the mice challenged with the ATCC 25904 isolate, 80% of the placebo-vaccinated Group 1 mice died within 10 days of challenge (Table 14). This demonstrated that strain ATCC 25904 caused a high rate of mortality in mice at the dose level administered. In contrast to the mice in Group 1, only 25% of the mice in Group 2 (vaccinated with proteins extracted
15 from strain ATCC 25904 following growth in iron-depleted conditions, SIRP Extract) died within 10 days post-challenge. These results illustrated that the mice challenged with strain ATCC 25904 were significantly protected by vaccination with the protein composition derived from iron-depleted ATCC 25904 ($p = 0.0006$, logrank test for mortality). In addition, only 50% of the
20 mice in Group 3 (vaccinated with recombinant MntC polypeptides, rMntC) died within 10 days of challenge, indicating that recombinant proteins offered protection against challenge with the ATCC 25904 strain ($p = 0.100$, logrank test for mortality).

25 IP Challenge (Study B)

 Fourteen days after the second vaccination, mice in all groups (1-4) were intraperitoneally challenged with 0.5 ml of the appropriate organism. The three groups of mice were challenged identically with 1×10^9 CFU of *S. aureus* strain ATCC 25904 per mouse. Mortality was recorded daily for 10 days after
30 challenge.

 When comparing the mice challenged with the ATCC 25904 isolate, 60% of the placebo-vaccinated Group 1 mice died within 10 days of challenge (Table 14). This demonstrated that strain ATCC 25904 caused a moderate rate

of mortality in mice at the dose level administered. In contrast to the mice in Group 1, only 30% of the mice in Group 2 (vaccinated with proteins extracted from strain ATCC 25904 following growth in iron-depleted conditions, SIRP Extract) died within 10 days post-challenge. These results illustrated that the mice vaccinated with the protein composition derived from iron-depleted ATCC 25904 died at half the rate of placebo-vaccinated mice when challenged with strain ATCC 25904 ($p = 0.143$, logrank test for mortality). Mice vaccinated with the combination of seven recombinant proteins showed a higher level of protection relative to placebo. Only 20% of the mice in Group 3 (vaccinated 2x with recombinant polypeptides, rSIRP7 2x) died within 10 days of challenge and only 10% of the mice in Group 4 (vaccinated 3x with recombinant polypeptides, rSIRP7 3x) died within 10 days of challenge, indicating that three vaccinations with recombinant proteins offered significant protection against challenge with the ATCC 25904 strain ($p = 0.040$, logrank test for mortality).

IV Challenge (Study C)

Fourteen days after the second vaccination, mice in all groups (1-3) were intravenously challenged in the lateral tail vein with 0.3 ml of the appropriate organism. The three groups of mice were challenged identically with 2×10^8 CFU of *S. aureus* strain ATCC 25904. Mortality was recorded daily for 10 days after challenge.

When comparing the mice challenged with the ATCC 25904 isolate, 90% of the placebo-vaccinated Group 1 mice died within 10 days of challenge (Table 14). This demonstrated that strain ATCC 25904 caused a high rate of mortality in mice at the dose level administered. In contrast to the mice in Group 1, only 40% of the mice in Group 2 (vaccinated with proteins extracted from strain ATCC 25904 following growth in iron-depleted conditions, SIRP Extract) died within 10 days post-challenge. These results illustrated that the mice challenged with strain ATCC 25904 were significantly protected by vaccination with the protein composition derived from iron-depleted ATCC 25904 ($p = 0.0164$, logrank test for mortality). In addition, only 40% of the mice in Group 3 (vaccinated with recombinant polypeptides, rSIRP7) died within 10 days of challenge, indicating that recombinant proteins offered

significant protection against challenge with the ATCC 25904 strain ($p = 0.0255$, logrank test for mortality).

Results are shown in Table 14 and Fig. 161.

- 5 Table 14. Mortality of Vaccinated and Non-Vaccinated Mice Following Challenge with *Staphylococcus aureus* ATCC isolate 25904.

Groups	# Mice	# Dead	Percent mortality (%)
IV Challenge (Study A)			
Group 1 (Placebo)	20	16/20	80
Group 2 (SIRP Extract)	10	5/20	25
Group 3 (rMntC)	10	5/10	50
IP Challenge (Study B)			
Group 1 (Placebo)	10	6/10	60
Group 2 (SIRP Extract)	10	3/10	30
Group 3 (rSIRP (2x))	10	2/10	20
Group 4 (rSIRP (3x))	10	1/10	10
IV Challenge (Study C)			
Group 1 (Placebo)	10	9/10	90
Group 2 (SIRP Extract)	10	4/10	40
Group 3 (rMntC)	10	4/10	40

10

Example 20

Passive immunization using recombinantly-prepared polypeptides

A polyclonal antibody composition is prepared as described in Example 8, except that the mice are vaccinated with a composition of recombinant

15 polypeptides prepared as described in Example 16.

The resulting antibody composition is used to passively immunize mice as described in Example 9. The immunized mice are challenged as described in Example 9.

Immunized mice will show decreased mortality compared to
5 unvaccinated and placebo vaccinated mice.

Example 21

Large scale fermentation and isolation of recombinantly-produced polypeptides

10

A master seed stock of the recombinant *E. coli* of Example 14 can be prepared by growing the organism in 2000 ml of sterile RM medium (20 g casamino acids, 60 g Na₂HPO₄, 30 g KH₂PO₄, 5 g NaCl 10 g NH₄Cl per liter and 100 ug/ml ampicillin) for 8 hours at 37°C. The bacteria can be harvested by
15 centrifugation at 10,000 x g for 30 minutes. The culture can be washed twice by centrifugation (10,000 x g) and the final bacterial pellet resuspended in 500 ml of sterile RM medium containing 20% sterile glycerol. One milliliter of the culture will be transferred to a 2 ml cryovial and stored at -85°C.

A cryovial (1 ml) of the recombinant master seed stock can be used to
20 inoculate a 100 ml culture flask containing the medium described above, with the exception of having 2 g casamino acids and 0.5 % glucose ("modified RM medium"). The culture can be incubated at 37°C for seven hours, at which time it can be inoculated into 2 liters of modified RM medium and allowed to grow for an additional four hours at 37°C. This culture can be used to inoculate a 30-
25 liter New Brunswick BIOFLOW 4 bench-top fermenter charged with 20 liters of modified RM medium except the final concentration of casamino acids will be 20 g/liter ("fermentation RM medium"). The pH of the fermentation medium can be held maintained between 6.9 and 7.2 by automatic titration with 30% NaOH and 10% HCl. The fermentation culture can be stirred at 350
30 rev/minute, and the culture can be aerated with 11 liters/minute at 37°C. Foaming can be controlled automatically by the addition of 0.4 % silicone defoamer (Antifoam-B, J. T. Baker, NJ). The culture can be allowed to grow continuously at these conditions for four hours (OD₆₀₀=4.0-6.0), at which time

the culture can be pumped into a 150-liter fermenter (W. B. Moore, Easton PN), charged with 110 liters of fermentation RM medium and 0.2 % defoamer. The parameters of the fermentation can be as follows: 650 rpm, 60% DO, 50 slpm air, 10 psi back pressure, 37°C, and pH held at 7.2 with NaOH. After reaching
5 late exponential phase growth (approximately six hours, $OD_{600}=15.0$), recombinant proteins can be induced by adding of 150 ug/ml IPTG. The fermentation can be allowed to grow for an additional three hours at which point the fermentation can be terminated by lowering the temperature of the fermentor to 18°C (OD_{600} 20-25 at a 1:100 dilution).

10 After fermentation, the recombinantly-produced polypeptides can be harvested by conventional means. The cells are disrupted (e.g., by homogenization) to release the recombinantly-produced polypeptides, some of which are soluble and some of which are insoluble.

Soluble polypeptides can be concentrated by tangential flow filtration,
15 detergent solubilized, and harvested by metal affinity chromatography. The collected soluble polypeptides can be isolated as described for the isolation of soluble polypeptides in Example 16.

Insoluble proteins can be harvested by high speed centrifugation. The pellet of insoluble polypeptides can be collected and then isolated as described
20 for the isolation of insoluble polypeptides in Example 16.

Example 22

25 Mouse vaccination

Recombinantly-produced SirA, SYN2, FhuD, and MntC polypeptides are prepared and isolated as described in Example 16.

30 Thirty (N=30) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed into three groups (10 mice/group). Mice are housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all

mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (40 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-
5 restriction (40 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptides SirA, SYN2, FhuD, and MntC (each 10 µg/100 µl, total protein 40 µg/100 µl).

Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

10

Example 23

Mouse vaccination

15 Recombinantly-produced PflB polypeptide is prepared and isolated as described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed into three groups (10 mice/group). Mice are housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all
20 mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-
25 restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide PflB (each 10 µg/100 µl).

Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

30

Example 24
Mouse vaccination

Recombinantly-produced Opp1A polypeptide is prepared and isolated as
5 described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding
Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed
into three groups (10 mice/group). Mice are housed in polycarbonate mouse
cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all
10 mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate
composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-
restriction (10 µg/100 µl).

15 Group-3: Vaccinated with recombinant polypeptide Opp1A (each 10
µg/100 µl).

Mice are challenged as described in Example 19. Mice in Group 3 will
exhibit reduced mortality compared to mice in Group 1.

20

Example 25
Mouse vaccination

Recombinantly-produced SirA polypeptide is prepared and isolated as
25 described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding
Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed
into three groups (10 mice/group). Mice are housed in polycarbonate mouse
cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all
30 mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate
composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide SirA (each 10 µg/100 µl).

- 5 Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

Example 26

10 Mouse vaccination

Recombinantly-produced SYN2 polypeptide is prepared and isolated as described in Example 16.

- 15 Thirty (N=30) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed into three groups (10 mice/group). Mice are housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

- 20 Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide SYN2 (each 10 µg/100 µl).

- 25 Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

Example 27

30 Mouse vaccination

Recombinantly-produced FhuD polypeptide is prepared and isolated as described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed into three groups (10 mice/group). Mice are housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide FhuD (each 10 µg/100 µl).

Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

Example 28

Mouse vaccination

Recombinantly-produced SYN1 polypeptide is prepared and isolated as described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed into three groups (10 mice/group). Mice are housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide SYN1 (each 10 µg/100 µl).

Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

Example 29

Mouse vaccination

Recombinantly-produced MntC polypeptide is prepared and isolated as
5 described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding
Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed
into three groups (10 mice/group). Mice are housed in polycarbonate mouse
cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all
10 mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate
composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-
restriction (10 µg/100 µl).

15 Group-3: Vaccinated with recombinant polypeptide MntC (each 10
µg/100 µl).

Mice are challenged as described in Example 19. Mice in Group 3 will
exhibit reduced mortality compared to mice in Group 1.

20

Example 30

Mouse vaccination

Recombinantly-produced SstD polypeptide is prepared and isolated as
25 described in Example 16.

Thirty (N=30) female BALB/C mice obtained from Harlan Breeding
Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed
into three groups (10 mice/group). Mice are housed in polycarbonate mouse
cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all
30 mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate
composition two times at 14 day intervals as follows:

Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide SstD (each 10 µg/100 µl).

5 Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

Example 31

10 Mouse vaccination

Recombinantly-produced FhuD2 polypeptide is prepared and isolated as described in Example 16.

15 Thirty (N=30) female BALB/C mice obtained from Harlan Breeding Laboratories (Indianapolis, IN) weighing 16-22 grams are equally distributed into three groups (10 mice/group). Mice are housed in polycarbonate mouse cages (N = 5 mice per cage). Food and water are supplied *ad libitum* to all mice. Mice are vaccinated subcutaneously with 0.1 ml of the appropriate composition two times at 14 day intervals as follows:

20 Group-1: Placebo, vaccinated with ovalbumin (10 µg/100 µl)

Group-2: Vaccinated with ATCC 25904 proteins expressed under iron-restriction (10 µg/100 µl).

Group-3: Vaccinated with recombinant polypeptide FhuD2 (each 10 µg/100 µl).

25 Mice are challenged as described in Example 19. Mice in Group 3 will exhibit reduced mortality compared to mice in Group 1.

Example 32

30 Recombinant polypeptides are specifically bound by sera generated by *S. aureus* infection

S. aureus were prepared as challenge organisms as described in Example

4 and used to challenge mice as described in Examples 5 or 19. Sera were obtained from mice either prior to challenge or from mice which had received placebo vaccine, had been challenged with *S. aureus* Newman and recovered (convalescent). Blood was collected from challenged and unchallenged mice
5 and sera were obtained by centrifugation. Sera were used to evaluate the reactivity of the individual recombinantly-produced polypeptides using an immunoblot method as described in Example 15.

Sera collected from mice prior to challenge did not react with any of the recombinant polypeptides.

10 Antibody in the sera produced as a result of *S. aureus* challenge reacted with the recombinantly-produced polypeptides as shown in Figure 202. Antibody raised against polypeptides expressed during infection by *S. aureus* recognizes recombinantly-produced variants of the polypeptides. Thus, recombinant polypeptides are immunological substitutes for—e.g.,
15 immunological fragments of—immunological polypeptides expressed by the challenge *S. aureus*.

Example 33

20 Antibody against recombinantly-produced polypeptides and iron-regulated membrane polypeptides extracted directly from *S. aureus* cells crossreact

Recombinant iron-regulated polypeptides are expressed and purified as described in Example 16. Iron-regulated membrane polypeptides are extracted
25 directly from *S. aureus* cells grown in low-iron conditions as described in Example 1. The recombinant polypeptides are formulated into a vaccine as described in Example 16 and used to vaccinate mice as described in Example 17. The *S. aureus*-extracted polypeptides are formulated into vaccines as described in Example 2 and used to vaccinate mice as described in Example 3.
30 For the recombinant polypeptide vaccines, a single recombinant polypeptide will be formulated into the vaccine. Antisera from vaccinated mice are collected and used for immunoblots as described in Example 15.

The antisera from animals vaccinated with recombinantly-produced iron-regulated polypeptides are further used to evaluate reactivity with extracted iron-regulated polypeptides following the separation of the extracted iron-regulated polypeptides by SDS-PAGE. Antisera from animals vaccinated with recombinant polypeptides will bind to appropriate SDS-PAGE-separated extracted polypeptides where antibody epitopes are conserved between the extracted polypeptides and the recombinantly-produced polypeptides.

The antisera from animals vaccinated with extracted iron-regulated polypeptides are further used to evaluate reactivity with individual recombinantly-produced iron-regulated polypeptides following the separation of the recombinant iron-regulated polypeptides by SDS-PAGE. Antisera from animals vaccinated with extracted polypeptides will bind to appropriate SDS-PAGE-separated recombinant polypeptides where antibody epitopes are conserved between the recombinantly-produced polypeptides and the extracted polypeptides.

Example 34

Western blot analysis of recombinant *S. aureus* proteins

Recombinant *S. aureus* proteins were prepared as described in Example 16, then subjected to sodium dodecyl sulfate-10% polyacrylamide gel electrophoresis and transferred to nitrocellulose membranes (BioRad, Hercules, CA) for Western blot analysis. Individual blots were reacted with serum from healthy (no reported Staphylococcus infection) or convalescent (Methicillin Resistant *S. aureus*) human donors.

As a control to identify each recombinant Histidine-tagged protein, each blot was co-incubated with anti-Histidine antibodies (Rockland Immunochemicals, Inc., Gilbertsville, PA and Qiagen GmbH, Hilden, Germany) to identify the recombinant proteins using two-color analysis. All primary antisera were used at a 1:1000 dilution and incubated for 1 hour on a rocker. Following several washes with TBS+0.05% Tween to remove unbound antibody, the membranes were subsequently incubated with a 800CW dye-

conjugated human secondary antibody (Rockland Immunochemicals, Inc.,
Gilbertsville, PA), 680 dye-conjugated mouse secondary antibody (Li-cor
Biosciences, Lincoln, NE) or a 800CW dye-conjugated rabbit secondary
antibody (Li-Cor Biosciences) at the dilution recommended by the manufacturer,
5 for 1 hour, in the dark. The membranes were washed an additional three to five
times with TBS+0.05% Tween with the last wash in TBS-only. Fluorescent
signals (680 and 800) were detected using the Odyssey Infrared Imaging System
(Li-Cor Biosciences). Results are shown In Figure 203 (healthy) and Figure 204
(convalescent).

10

Example 35

S. aureus DU5875 surface expression of metal-regulated polypeptides

15 *S. aureus* strain DU5875 (cap-, spa-) was grown in iron-replete (TSB+
0.3 mM ferric chloride) or iron-deplete (TSB+1mM dipyrldyl) conditions to an
OD₆₀₀ of approximately 0.6.

In panels A-C of Figure 205, *S. aureus* strain DU5875 was grown in iron-
deplete conditions to an OD₆₀₀ of approximately 0.75 and frozen. Bacteria were
20 thawed, washed in PBS and resuspended in PBS+ 1% Pig IgG+1% BSA as a
blocking step. Mouse antiserum raised against FhuD, Opp1A or PflB was used
at a dilution of 1:100 to stain approximately 2×10^6 bacteria. Preimmune mouse
serum was used as a negative control. Bacteria were then washed in blocking
buffer and incubated with an AF633-conjugated goat anti-mouse secondary
25 antibody, and analyzed on a flow cytometer. Bacteria incubated with
preimmune mouse serum, at a 1:100, were used as a negative control.

In panel D of Figure 205, *S. aureus* strain DU5875 was grown in iron-
deplete conditions to an OD₆₀₀ of approximately 2.0 and frozen. Bacteria were
thawed, washed in PBS and resuspended in PBS+ 0.2% Pig IgG+1% BSA as a
30 blocking step. Mouse antiserum raised against rSIRP7 was used at a dilution of
1:50 to stain approximately 5×10^7 bacteria, with preimmune mouse serum used
as a negative control. Bacteria were then washed in blocking buffer and

incubated with an AF633-conjugated goat anti-mouse secondary antibody, and analyzed on a flow cytometer.

The results of this assay indicate that murine antibodies raised against the SIRP proteins bind to *S. aureus* cells. Cells grown under iron-deplete conditions
5 bind more antibody than cells grown under iron-replete conditions, providing further evidence that FhuD, Opp1A, and PflB are expressed at higher levels under low-iron conditions and are antigens that can induce immunological activity against *Staphylococcus* spp. The increase in median fluorescence intensity (MFI) demonstrates the relative increase in fluorescence when anti-
10 SIRP antibodies bind to *S. aureus* cells compared to the MFI of preimmune mouse serum. Results are shown in Figure 205.

Example 36

15 A Luminex assay was used to evaluate cytokine expression by splenocytes from mice immunized with the combination (rSIRP7) of recombinant SIRP components PflB (SEQ ID NO:353), Opp1A (SEQ ID NO:364), SirA (SEQ ID NO:375), SYN2 (SEQ ID NO:386), FhuD (SEQ ID NO:397), SYN1 (SEQ ID NO:408, and MntC (SEQ ID NO:419) or placebo,
20 then restimulated with SIRP Extract (SIRPE) or rSIRP7. Several cytokines were upregulated upon restimulation, and the cytokine profiles induced by SIRPE and rSIRP7 restimulation were similar. The overall cytokine profile in response to rSIRP7 or SIRPE restimulation resembled that expected from a
25 Th1/Th17-type immune response, and demonstrates that vaccination with the recombinant SIRP components induces a cellular immune response that can be measured based on cytokine expression.

Methods. Mice were vaccinated two times, 14 days apart, with 70 µg total protein (OVA, SIRP Extract, or rSIRP7) formulated with 50% IFA. CD4⁺
30 T cells were purified from splenocyte suspensions by negative selection using a CD4⁺ T cell isolation kit and LD columns (Miltenyi Biotec, Inc., Auburn, CA). Briefly, biotinylated antibodies were used to label all cells except CD4⁺ T cells, and then streptavidin-conjugated magnetic beads were used to remove these cells from the mixture with a magnetic column, leaving highly purified CD4⁺ T cells.

The resulting CD4⁺ T cells were found to be greater than 95% pure based on CD3 and CD4 expression. Naïve splenocytes were treated with Mitomycin-C to generate mitotically inactive antigen presenting cells. 4 x 10⁵ APC added to 5 x 10⁵ CD4⁺ T cells plus stimulation antigen, followed by 42 hours of incubation.

- 5 Supernatants were analyzed by Luminex using the standard assay parameters. Results are shown in Figure 206.

10

Example 37

S. aureus extracts from several strains (Newman, Reynolds) were prepared using the method described in Example 1. *S. aureus* cells were grown in either iron restricted media containing 1000 µM 2,2-dipyridyl (Sigma-Aldrich St. Louis, MO) or iron replete media containing 300 µM FeCl₃ (Sigma-Aldrich St. Louis, MO).

Proteins within *S. aureus* membrane extracts from iron-replete and iron-deplete cultures were identified and quantified using ITRAQ and LCQ mass spectrometry. Amine-modified labeling of membrane extracts for iron deficient and iron replete *Staphylococcus aureus* Newman strain were performed with ITRAQ-8plex reagents (Applied Biosystems, Inc., Foster City, CA) using 40.0 µg of membrane extract (reagents 113 versus 115) according to the manufacturer's 8Plex protocol. Cation exchange chromatography was applied using an MCX column (Waters Corp., Milford, MA) and the peptides separated using an ULTIMATE 3000 NANO LC system (Dionex Corp. Bannockburn, IL) coupled to ESI mode using a QSTAR XL mass spectrometer (Applied Biosystems, Inc., Foster City, CA).

The ratio of metal-regulated polypeptides produced by cells grown in iron-restricted media compared to iron-replete media was measured. The ratio is a relative measure of protein expression and does not provide data indicating an absolute amount of protein present in the extract. Results are shown in Table 15.

35

Table 15.

Protein	Identified in extract	Fold increase in low iron
MntC	Yes	22
SYN1	Yes	Not determined
FhuD	No	Not determined
SYN2	Yes	23
SirA	Yes	36
Opp1A	Yes	Not determined
PflB	Yes	Not determined
FhuD2	Yes	6
SstD	Yes	14

Example 38

5

An oxidative burst assay can be used to measure the production of reactive oxygen species by neutrophils, an indication of an inflammatory response. To obtain neutrophils from fresh blood, red blood cells from fresh human blood are lysed by the addition of lysis buffer (150 mM NH₄Cl, 10 mM KHCO₃, 1 mM disodium EDTA, pH 7.4) at a 1:10 dilution, incubated for 10 minutes at room temperature and centrifuged for 10 minutes at 430 x g. The supernatant is removed by tube inversion and pellets are washed twice with PBS, then resuspended in 5 ml of RPMI-Hepes 5% FCS + Glutamine (complete RPMI) and enumerated using a MULTISIZER (Beckman Coulter, Inc., Brea, CA) after a 1/500 dilution of cell suspension in ISOTON (20 µl of cells in 10 ml ISOTON, Beckman Coulter, Inc., Brea, CA).

For the preparation of bacteria, *S. aureus* strain Lowenstein is seeded in TSB medium and grown for 20 hours at 37°C in 50 ml of medium. From this culture, 5 ml are pelleted for 10 minutes at 4000 rpm at 4°C. The pellet is then washed with 50 ml of PBS and repelleted by centrifugation for 10 minutes at 4000 rpm at 4°C. The wash step is repeated and the bacterial pellet is resuspended in 5 ml of PBS. Bacteria are adjusted to a theoretical density of

1.10^9 CFU/ml and cell dilutions are plated on agar and incubated for exact enumeration the following day.

To deplement the sera, all sera are incubated for 30 minutes at 56°C. Mixtures of sera and cells are performed in polypropylene sterile DW plates in a final volume of 500 µl per well. Into each well, the following reagents are added (as shown in Table 16), in order: culture medium (RPMI-Hepes, glutamine, 5% FCS), live bacteria at the appropriate concentration, the sera at the appropriate dilution, the complement, hPMNs at the appropriate concentration and, finally, the DHR molecule (Life Technologies, Inc., Carlsbad, CA) as the marker of oxidative burst. Plates are incubated for 25 minutes at 37°C with shaking. The reaction is stopped by incubating the plates for five minutes on ice.

Table 16.

Reagent	Identification	Working Conc.	Vol./ well	Final concentration (in 500 µl)
Live bacteria	TSB, 20h, 37°C 10^9 CFU/ml	$1.25.10^8$ CFU/ml	200 µl	5.10^7 CFU/ml
Whole blood leukocytes	From 2 different donors after the red blood cell lysis	$2.5.10^6$ cells/ml	100 µl	$0.5.10^6$ cells/ml
DHR	Life Technologies, Inc. Cat. No. D632 (10 mg/ml)	100 µg/ml	50 µl	10 µg/ml
Baby rabbit complement	Produced in-house	1/10	50 µl	1/100 (1%)
Sera	Serum from mouse immunized with adjuvant alone	1/10	50µl	1/100
	Anti-whole cell control	1/100	50µl	1/1000
	Anti- <i>S. aureus</i> protein	1/10 and 1/100	50µl	1/100 and 1/1000

For flow cytometric analysis, the human PMNs are first identified according to their size and granularity and then verified by surface expression of specific markers (CD35, CD16, GR1, etc.), then oxidative burst marker. The data are given in terms of percent of activated hPMNs able to induce oxidative burst in comparison with the negative control group. (Didier, 2003 ; Ploppa, 2008).

10

Example 39

Opsonophagocytic Assay

15

An opsonophagocytic assay (OPA) has been developed to estimate the functional phagocytic activity of serum for *Staphylococcus aureus* by measuring the complement dependent opsonic activity of the serum. The OPA is summarized on Table 17. Two strains of *S. aureus* are used in the assay. Strain DU5875, which does not produce capsule or protein A, is used to best control the assay. Strain LST4 Lowenstein, which does express capsule and protein A, is used as a wild-type strain in the assay. The number of bacteria used in the assay is dependent on the source of the effector white blood cells and ranges in concentration from 1×10^5 cfu/ml to 5×10^7 cfu/ml. White blood cells from healthy human volunteers or from a human promyelocytic leukemia cell line, HL60, are used as phagocytic effector cells. As Table 17 indicates, the number of effector cells used in the assay is dependent on the source of the cells. Baby rabbit serum is used as a source of complement and is diluted from 1% to 10% depending on the lot of serum in order to maximize the functional complement activity of the rabbit serum while minimizing the toxicity. Sera undergoing testing for opsonic activity, pre- or post-immunization, are decompemented at 56°C for 30 minutes and are tested in the assay at a dilution of 1:20 to 1:2,000,000. Phagocytosis is determined by viable counts (cfu) of *S. aureus*. Test serum which demonstrates a significant loss of cfu in combination with active complement when compared to preimmune serum, is considered opsonic. The assay data are analyzed by the Student's t-Test using one-tailed

30

distribution with unequal variance. (Kim 2010; Stranger-Jones 2006; Dryla 2005).

Table 17.

Reagent	Final Concentration
Live Bacteria	approx. 1×10^5 cfu/ml when using HL-60 effector cells approx. 5×10^7 cfu/ml when using healthy human WBCs
Effector cells	HL-60 human cell line chemically induced to differentiate with N,N- dimethylformamide diluted to 10×10^6 /ml or human white blood cells from healthy volunteers diluted to 1×10^7 /ml
Complement	Baby rabbit serum diluted from 1% to 10% according to Lot of serum
Test Serum	Serum obtained after the last immunization, decompemented and serially diluted from 1:20 to 1:2,000,000

5

Example 40

Immune mechanism studies (*in vivo*)

10 In another example, the immune mechanism by which vaccine proteins confer protection to mice will be assessed. These experiments can include two types: (1) using gene knockout mice in vaccine-challenge experiments to determine whether specific immune components are necessary for protection; and (2) adoptive transfer experiments, where immune cells from immunized
15 donor mice are transferred into naïve recipients prior to bacterial challenge, in order to test whether the transferred components are sufficient to confer protection.

For examples involving vaccine challenge experiments in gene knockout mice, the mice can be purchased from commercial vendors and can include
20 several well characterized strains such as, for example, B cell knockouts (μ MT), T cell knockouts (TCR α), and a variety of cytokine knockouts, such as IFN- γ , IL-1 α , TNF α , IL-17, etc.). The mice can be immunized as described in Example 3, and then challenge with *S. aureus*, as described in Example 5. The use of wild

type mice with the same genetic background (Balb/c) can provide suitable controls to measure the effect of the knockout on vaccine-mediated protection against *S. aureus*. For example, if the vaccinated B cell knockout mice die more rapidly, or in larger numbers relative to the vaccinated control mice in response to bacterial challenge, it can be concluded that B cells (or their products) are important for the vaccine-mediated protection against *S. aureus*. These strategies are standard practice in the field for measuring contributions of various immune components to vaccine protection (Spellberg, 2008; Lin, 2009).

For examples involving adoptive transfer of immune components, wild type Balb/c donor mice can be immunized as described in Example 3 in order to generate tissue for adoptive transfer. These mice can then be euthanized 2-4 weeks after the second immunization and the blood collected via cardiac puncture and secondary lymphoid tissue (lymph nodes and spleen) collected. Lymph nodes collected can include: axillary, brachial, mesenteric, inguinal, superficial cervical, deep cervical, and lumbar. Serum can be isolated from the blood using standard methodology, such as centrifugation-based serum separators, and then transferred back into a separate set of recipient mice via intravenous or intraperitoneal injection. The use of 1-3 donor mice per recipient is a reasonable ratio for serum transfer, and can occur in volumes up to 0.5 ml. In addition, T cells from the donor lymphoid tissue can be purified using antibodies and magnetic bead enrichment technology (Miltenyi Biotec) that is standard practice in the field. It is typical to achieve 95-99% cell purity using these methods, as assessed by staining the cell surface proteins with antibodies to specific lineage markers and examining the cells by flow cytometry. The cell populations of interest (e.g., CD4⁺ T cells, CD8⁺ T cells, etc.) can be transferred back into recipient animals via intravenous injection (between 2,000,000 and 5,000,000 T cells per recipient). Recipient mice can receive T cells (or subsets of T cells), immune serum, or both.

As a negative control, placebo-immunized animals can also be used for serum and T cell isolation followed by transfer back into naïve recipients. As a positive control, a group of recipients that gets immunized with the standard protective vaccine can be included in order to provide a baseline assessment of protective efficacy upon bacterial challenge.

Once the recipient mice have received various transferred cells or sera, they can be challenged with *S. aureus* as described in Example 5. Based on the percentage of death and the rate of death, the relative contribution of various immune components to vaccine protection can be assessed. For instance, if recipients that have been administered T cells from vaccinated donors are protected against challenge at the same rate as the positive control, it can be concluded that T cells are sufficient for vaccine-mediated protection. This experimental strategy is standard practice in the field for assessing the immunological mechanisms of vaccines (Spellberg, 2008; Lin, 2009).

Example 41

Inhibition of iron uptake assay

For assessing whether antibodies directed against SIRP components inhibit cell growth by blocking iron uptake, an iron uptake/transport assay can be run using bacterial cells that are pre-incubated with anti-SIRP antisera. *S. aureus* cells from any strain are grown overnight in chelated or iron-rich media. Cells are subcultured in the same media and grown to mid to late log phase, pelleted and incubated with anti-SIRP antisera or control antisera in PBS for up to one hour. Cells were then harvested by filtration using 0.45 μ M filters, then resuspended in Chelex-100 treated minimal medium to eliminate environmental iron. Cells are shaken briefly. Meanwhile, an iron source (for example, ferrichrome) is mixed with radiolabeled $^{55}\text{FeCl}_2$ (or another radiolabeled iron molecule) with nitrilotriacetic acid and allowed to incubate for several minutes.

To commence iron uptake, a small aliquot (for example, 10 μ l) of the radiolabeled iron mixture was added to cells (in a 1 ml volume) in a 10 ml Fe-free culture tube. The tube is incubated with periodic vortexing and sampling of aliquots filtered onto membrane filters and washed with LiCl. Following drying, membranes are counted in scintillation fluid to quantify the iron uptake. Cells pre-incubated with anti-SIRP antisera should be slower to take up iron than cells pre-incubated with control antisera (Sebulsky, 2000; Goel, 2001).

Example 42

High-yield protein purification protocol

- 5 In some instances, recombinant rSIRPs were purified using a high-yield method. This method is optimized for higher yields and higher purity of the polypeptide. The method is performed by resuspending the bacterial pellet containing the recombinant-produced polypeptide in 20 mM Tris pH 9; 300 mM NaCl complemented with lysozyme (100 µg/ml final) and MgCl₂ (1 mM final).
- 10 The sample is then incubated with gentle rocking for 15 minutes at 4°C. After 15 minutes, 1U/ml of benzonase is added and the sample is incubated for one hour at room temperature. The soluble lysate obtained after centrifugation (20,000 x g, 20 minutes at 4°C) should be filtered with a 0.45 µm filter and added to an equilibrated 5 ml gravity column packed with nickel-His binding
- 15 resin (Novagen 69670-4, EMD Chemicals, Inc., Gibbstown, NJ). The purification of the N-His₆-protein is performed following manufacturer's instructions. The purification of the N-His₆-protein is finalized using a HiLoad 26/60 Superdex 75 prep grade size-exclusion column (GE Healthcare Biosciences, Piscataway, NJ). The column is equilibrated with 20 mM Tris pH 9;
- 20 300 mM NaCl using a BioCAD FPLC (Applied Biosystems Inc., Foster City, CA) and 15 ml of the protein sample is loaded onto the HiLoad column at a flow-rate of 1.5ml/min. The sample is eluted following 3.5 hours of run time at a flow-rate 1.8 ml/min using 20 mM Tris pH 9; 300 mM NaCl as the mobile phase.
- 25 The polypeptide is quantified using a modified version of the BCA (Thermo Fisher Scientific, Inc., Rockford, IL) procedure where 15% sodium dodecyl sulphate (SDS), 8M urea, and 2.5% 3-([3-Cholamidopropyl]dimethylammonio)-2-hydroxy-1-propanesulfonate (CHAPS) is used to ensure complete solubility of all protein within the sample. The assay
- 30 also consists of 5% SDS added to the BCA working reagent to maintain solubility of the protein during the BCA reactive phase. The reading and analysis of the BCA is performed according to the product literature.

For densitometric analysis, 3.0 µg of final product antigen was quantified for purity using 10% SDS PAGE, stained with coomassie and imaged using an ODYSSEY scanner (LiCor Biosciences, Lincoln, NE). The stained gel was scanned and the areas of the major components were determined relative to the
5 total area. Residual Endotoxin was removed using an Endotrap blue one/endosafe kit (Hyglos GmbH, Regensburg, Germany). The final batch is stored with a concentration range of 1.0 mg/ml to 4.0 mg/ml in PBS storage buffer at less than -70°C in appropriate aliquots.

10 The complete disclosure of all patents, patent applications, and publications, and electronically available material (including, for instance, nucleotide sequence submissions in, e.g., GenBank and RefSeq, and amino acid sequence submissions in, e.g., SwissProt, PIR, PRF, PDB, and translations from annotated coding regions in GenBank and RefSeq) cited herein are incorporated
15 by reference. In the event that any inconsistency exists between the disclosure of the present application and the disclosure(s) of any document incorporated herein by reference, the disclosure of the present application shall govern. The foregoing detailed description and examples have been given for clarity of understanding only. No unnecessary limitations are to be understood therefrom.
20 The invention is not limited to the exact details shown and described, for variations obvious to one skilled in the art will be included within the invention defined by the claims.

Unless otherwise indicated, all numbers expressing quantities of components, molecular weights, and so forth used in the specification and
25 claims are to be understood as being modified in all instances by the term "about." Accordingly, unless otherwise indicated to the contrary, the numerical parameters set forth in the specification and claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt
30 to limit the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. All numerical values, however, inherently contain a range necessarily resulting from the standard deviation found in their respective testing measurements.

5

All headings are for the convenience of the reader and should not be used to limit the meaning of the text that follows the heading, unless so specified.

What is claimed is:

1. A composition comprising:
5 an isolated polypeptide having at least 92% sequence similarity to the amino acid sequence of SEQ ID NO:397, with the proviso that if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared
10 to amino acids 1-26 of SEQ ID NO:399.
2. A composition comprising:
an isolated polypeptide having at least 98% sequence similarity to the amino acid sequence of SEQ ID NO:408, with the proviso that if the
15 isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-5 of SEQ ID NO:415.
- 20 3. The composition of claim 1 or claim 2 wherein the composition protects an animal against challenge with a *S. aureus* selected from ATCC 19636 or ATCC 25904.
4. The composition of claim 1 or claim 2 further comprising a
25 pharmaceutically acceptable carrier.
5. The composition of claim 1 wherein the polypeptide comprises an amino acid sequence having at least 95% sequence similarity to the amino acid sequence of SEQ ID NO:397.
30
6. The composition of claim 1 wherein the polypeptide comprises an amino acid sequence having at least 92% sequence identity to the amino acid sequence of SEQ ID NO:397.

7. The composition of claim 6 wherein the polypeptide comprises an amino acid sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO:397.
- 5 8. The composition of claim 2 wherein the polypeptide comprises an amino acid sequence having at least 98% sequence identity to the amino acid sequence of SEQ ID NO:397.
- 10 9. The composition of claim 2 wherein the polypeptide comprises an amino acid sequence having at least 98% sequence identity to the amino acid sequence of SEQ ID NO:408.
- 15 10. The composition of claim 1 or claim 2 further comprising at least one second polypeptide, wherein the second polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator.
- 20 11. The composition of claim 10 wherein the second polypeptide comprises an amino acid sequence having at least 85% similarity to the amino acid sequence of any one of: SEQ ID NO:353, SEQ ID NO:354, SEQ ID NO:355, SEQ ID NO:356, SEQ ID NO:357, SEQ ID NO:358, SEQ ID NO:359, SEQ ID NO:360, SEQ ID NO:361, SEQ ID NO:362, SEQ ID NO:363, SEQ ID NO:364, SEQ ID NO:365, SEQ ID NO:366, SEQ ID NO:367, SEQ ID NO:368, SEQ ID NO:369, SEQ ID NO:370, SEQ ID NO:371, SEQ ID NO:372, SEQ ID NO:373, SEQ ID NO:374, SEQ ID NO:375, SEQ ID NO:376, SEQ ID NO:377, SEQ ID NO:378, SEQ ID NO:379, SEQ ID NO:380, SEQ ID NO:381, SEQ ID NO:382, SEQ ID NO:383, SEQ ID NO:384, SEQ ID NO:385, SEQ ID NO:386, SEQ ID NO:387, SEQ ID NO:388, SEQ ID NO:389, SEQ ID NO:390, SEQ ID NO:391, SEQ ID NO:392, SEQ ID NO:393, SEQ ID NO:394, SEQ ID NO:395, SEQ ID NO:396, SEQ ID NO:419, SEQ ID NO:420,
- 25 30

SEQ ID NO:421, SEQ ID NO:422, SEQ ID NO:423, SEQ ID NO:424,
SEQ ID NO:425, SEQ ID NO:426, SEQ ID NO:427, SEQ ID NO:428, or
SEQ ID NO:429.

5 12. The composition of claim 1 comprising further comprising at
least one second polypeptide, wherein the second polypeptide is
isolatable from a *Staphylococcus aureus* when incubated in media
comprising an iron chelator, is not isolatable when grown in the media
without the iron chelator, and comprises an amino acid sequence having
10 at least 85% similarity to the amino acid sequence of any one of: SEQ ID
NO:408, SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID
NO:412, SEQ ID NO:413, SEQ ID NO:414, SEQ ID NO:415, SEQ ID
NO:416, SEQ ID NO:417, or SEQ ID NO:418.

15 13. The composition of claim 12 wherein the second polypeptide
comprises an amino acid sequence having at least 85% identity to the
amino acid sequence of any one of: SEQ ID NO:408, SEQ ID NO:409,
SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413,
SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, or
20 SEQ ID NO:418.

14. The composition of claim 2 further comprising at least one
second polypeptide, wherein the second polypeptide is isolatable from a
Staphylococcus aureus when incubated in media comprising an iron
25 chelator, is not isolatable when grown in the media without the iron
chelator, and comprises an amino acid sequence having at least 85%
similarity to the amino acid sequence of any one of: SEQ ID NO:397,
SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401,
SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405,
30 SEQ ID NO:406, or SEQ ID NO:407.

15. The composition of claim 14 wherein the second polypeptide
comprises an amino acid sequence having at least 85% identity to the

amino acid sequence of any one of: SEQ ID NO:397, SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID NO:407.

5

16. The composition of claim 1 or claim 2 further comprising:

an isolated polypeptide is isolatable from a *S. aureus* when grown in media without the iron chelator and has a molecular weight of 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41

10 kDa, 40 kDa; or

a combination of such isolated polypeptides.

17. A method for treating an infection in a subject, the method comprising:

15 administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:

an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397.

20

18. A method for treating a symptom in a subject, the method comprising:

administering an effective amount of a composition to a subject having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:

25 an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397.

19. A method for treating an infection in a subject, the method comprising:

30 administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:

an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397.

20. A method for treating an infection in a subject, the method
5 comprising:

administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:

an isolated polypeptide having at least 80% sequence similarity to
10 the amino acid sequence of SEQ ID NO:408.

21. A method for treating a symptom in a subject, the method comprising:
administering an effective amount of a composition to a subject
having an infection caused by a *Staphylococcus* spp., wherein the
15 composition comprises:

an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:408.

22. A method for treating an infection in a subject, the method
20 comprising:

administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:

an isolated polypeptide having at least 80% sequence similarity to
25 the amino acid sequence of SEQ ID NO:408.

23. The method of any one of claims 17-22 wherein the composition further comprises at least one second polypeptide, wherein the second polypeptide is isolatable from a *Staphylococcus aureus* when incubated
30 in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator.

24. The method of claim 23 wherein the second polypeptide has a molecular weight as determined by electrophoresis on a sodium dodecyl sulfate-polyacrylamide gel of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa.

5

25. The method of any one of claims 17-22 wherein the composition further comprises at least one second polypeptide comprising an amino acid sequence having at least 85% similarity to the amino acid sequence of any one of: SEQ ID NO:353, SEQ ID NO:354, SEQ ID NO:355, SEQ
10 ID NO:356, SEQ ID NO:357, SEQ ID NO:358, SEQ ID NO:359, SEQ ID NO:360, SEQ ID NO:361, SEQ ID NO:362, SEQ ID NO:363, SEQ ID NO:364, SEQ ID NO:365, SEQ ID NO:366, SEQ ID NO:367, SEQ ID NO:368, SEQ ID NO:369, SEQ ID NO:370, SEQ ID NO:371, SEQ ID NO:372, SEQ ID NO:373, SEQ ID NO:374, SEQ ID NO:375, SEQ
15 ID NO:376, SEQ ID NO:377, SEQ ID NO:378, SEQ ID NO:379, SEQ ID NO:380, SEQ ID NO:381, SEQ ID NO:382, SEQ ID NO:383, SEQ ID NO:384, SEQ ID NO:385, SEQ ID NO:386, SEQ ID NO:387, SEQ ID NO:388, SEQ ID NO:389, SEQ ID NO:390, SEQ ID NO:391, SEQ ID NO:392, SEQ ID NO:393, SEQ ID NO:394, SEQ ID NO:395, SEQ ID
20 NO:396, SEQ ID NO:419, SEQ ID NO:420, SEQ ID NO:421, SEQ ID NO:422, SEQ ID NO:423, SEQ ID NO:424, SEQ ID NO:425, SEQ ID NO:426, SEQ ID NO:427, SEQ ID NO:428, or SEQ ID NO:429.

26. The method of any one of claims 17-19 wherein the composition
25 further comprises at least one second polypeptide, wherein the second polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator, is not isolatable when grown in the media without the iron chelator, and comprises an amino acid sequence having at least 85% similarity to the amino acid sequence of any one of:
30 SEQ ID NO:408, SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413, SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, or SEQ ID NO:418.

27. The method of claim 26 wherein the second polypeptide comprises an amino acid sequence having at least 85% identity to the amino acid sequence of any one of: SEQ ID NO:408, SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413, 5 SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, or SEQ ID NO:418.

28. The method of any one of claims 20-22 wherein the composition further comprises at least one second polypeptide, wherein the second 10 polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator, is not isolatable when grown in the media without the iron chelator, and comprises an amino acid sequence having at least 85% similarity to the amino acid sequence of any one of: SEQ ID NO:397, SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, 15 SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID NO:407.

29. The method of claim 29 wherein the second polypeptide comprises an amino acid sequence having at least 85% identity to the 20 amino acid sequence of any one of: SEQ ID NO:397, SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID NO:407.

25 30. The method of any one of claims 17-22 wherein the composition further comprises:

an isolated polypeptide that is isolatable from a *S. aureus* when grown in media without an iron chelator and has a molecular weight as determined by electrophoresis on a sodium dodecyl sulfate- 30 polyacrylamide gel of 150 kDa, 132 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, 40 kDa; or
a combination of such isolated polypeptides.

31. The method of any one of claims 17-22 wherein the subject is a mammal.
32. The method of claim 31 wherein the mammal is a human.
- 5 33. The method of any one of claims 17-22 wherein the *Staphylococcus* spp. is *S. aureus*.
- 10 34. A method for treating an infection in a subject, the method comprising:
administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:
antibody that specifically binds an isolated polypeptide having at
15 least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397.
- 20 35. A method for treating a symptom in a subject comprising:
administering an effective amount of a composition to a subject having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:
antibody that specifically binds an isolated polypeptide having at
least 80% sequence similarity to the amino acid sequence selected of
SEQ ID NO:397.
- 25 36. A method for decreasing colonization in a subject, the method comprising:
administering an effective amount of a composition to a subject colonized by a *Staphylococcus* spp., wherein the composition comprises:
30 an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397.

37. A method for treating an infection in a subject, the method comprising:
administering an effective amount of a composition to a subject having or at risk of having an infection caused by a *Staphylococcus* spp.,
5 wherein the composition comprises:
antibody that specifically binds an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:408.
- 10 38. A method for treating a symptom in a subject comprising:
administering an effective amount of a composition to a subject having an infection caused by a *Staphylococcus* spp., wherein the composition comprises:
antibody that specifically binds an isolated polypeptide having at
15 least 80% sequence similarity to the amino acid sequence of SEQ ID NO:408.
39. A method for decreasing colonization in a subject, the method comprising:
20 administering an effective amount of a composition to a subject colonized by a *Staphylococcus* spp., wherein the composition comprises:
an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:408.
- 25 40. The method of any one of claims 34-39 wherein the composition further comprises antibody that specifically binds to at least one second polypeptide, wherein the second polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron
30 chelator.
41. The method of claim 40 wherein the second polypeptide has a molecular weight as determined by electrophoresis on a sodium dodecyl

sulfate-polyacrylamide gel of 88 kDa, 55 kDa, 38 kDa, 37 kDa, 36 kDa, 35 kDa, or 33 kDa.

42. The method of claim 40 wherein the second polypeptide
5 comprises an amino acid sequence having at least 85% similarity to the
amino acid sequence of any one of: SEQ ID NO:353, SEQ ID NO:354,
SEQ ID NO:355, SEQ ID NO:356, SEQ ID NO:357, SEQ ID NO:358,
SEQ ID NO:359, SEQ ID NO:360, SEQ ID NO:361, SEQ ID NO:362,
SEQ ID NO:363, SEQ ID NO:364, SEQ ID NO:365, SEQ ID NO:366,
10 SEQ ID NO:367, SEQ ID NO:368, SEQ ID NO:369, SEQ ID NO:370,
SEQ ID NO:371, SEQ ID NO:372, SEQ ID NO:373, SEQ ID NO:374,
SEQ ID NO:375, SEQ ID NO:376, SEQ ID NO:377, SEQ ID NO:378,
SEQ ID NO:379, SEQ ID NO:380, SEQ ID NO:381, SEQ ID NO:382,
SEQ ID NO:383, SEQ ID NO:384, SEQ ID NO:385, SEQ ID NO:386,
15 SEQ ID NO:387, SEQ ID NO:388, SEQ ID NO:389, SEQ ID NO:390,
SEQ ID NO:391, SEQ ID NO:392, SEQ ID NO:393, SEQ ID NO:394,
SEQ ID NO:395, SEQ ID NO:396, SEQ ID NO:419, SEQ ID NO:420,
SEQ ID NO:421, SEQ ID NO:422, SEQ ID NO:423, SEQ ID NO:424,
SEQ ID NO:425, SEQ ID NO:426, SEQ ID NO:427, SEQ ID NO:428, or
20 SEQ ID NO:429.

43. The method of any one of claims 34-36 wherein the composition
further comprises antibody that specifically binds to at least one second
polypeptide, wherein the second polypeptide is isolatable from a
25 *Staphylococcus aureus* when incubated in media comprising an iron
chelator, is not isolatable when grown in the media without the iron
chelator, and comprises an amino acid sequence having at least 85%
similarity to the amino acid sequence of any one of: SEQ ID NO:408,
SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412,
30 SEQ ID NO:413, SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416,
SEQ ID NO:417, or SEQ ID NO:418.

44. The method of claim 43 wherein the second polypeptide comprises an amino acid sequence having at least 85% identity to the amino acid sequence of any one of: SEQ ID NO:408, SEQ ID NO:409, SEQ ID NO:410, SEQ ID NO:411, SEQ ID NO:412, SEQ ID NO:413, 5 SEQ ID NO:414, SEQ ID NO:415, SEQ ID NO:416, SEQ ID NO:417, or SEQ ID NO:418.

45. The method of any one of claims 37-39 wherein the composition further comprises antibody that specifically binds to at least one second 10 polypeptide, wherein the second polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator, is not isolatable when grown in the media without the iron chelator, and comprises an amino acid sequence having at least 85% similarity to the amino acid sequence of any one of: SEQ ID NO:397, 15 SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID NO:407.

46. The method of claim 45 wherein the second polypeptide 20 comprises an amino acid sequence having at least 85% similarity to the amino acid sequence of any one of: SEQ ID NO:397, SEQ ID NO:398, SEQ ID NO:399, SEQ ID NO:400, SEQ ID NO:401, SEQ ID NO:402, SEQ ID NO:403, SEQ ID NO:404, SEQ ID NO:405, SEQ ID NO:406, or SEQ ID NO:407.

25

47. The method of any one of claims 34-39 wherein the composition further comprises:

antibody that specifically binds to a polypeptide that is isolatable from a *S. aureus* when grown in media without an iron chelator and has a 30 molecular weight as determined by electrophoresis on a sodium dodecyl sulfate-polyacrylamide gel of 150 kDa, 292 kDa, 120 kDa, 75 kDa, 58 kDa, 50 kDa, 44 kDa, 43kDa, 41 kDa, 40 kDa; or a combination of such isolated polypeptides.

48. The method of any one of claims 34-39 wherein the subject is a mammal.
- 5 49. The method of claim 48 wherein the mammal is a human.
50. The method of any one of claims 34-39 wherein the *Staphylococcus* spp. is *S. aureus*.
- 10 51. The method of any one of claims 34-39 wherein the antibody is polyclonal antibody.
52. The method of any one of claims 34-39 wherein the antibody is polyclonal antibody.
- 15 53. The method of any one of claims 34-39 wherein the antibody is monoclonal antibody.
54. A kit for detecting antibody that specifically binds a polypeptide,
20 comprising in separate containers:
an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:408 or SEQ ID NO:397; and
a reagent that detects an antibody that specifically binds the polypeptide.
- 25 55. The method of claim 24 or claim 41,
wherein the isolated polypeptide with a molecular weight of 88 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of an 88 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and wherein the polypeptide is isolatable from a
30 *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator;

wherein the isolated polypeptide with a molecular weight of 55 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 55 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and wherein the polypeptide is isolatable from a *Staphylococcus aureus* when
5 incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator;

wherein the isolated polypeptide with a molecular weight of 38 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 38 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and
10 wherein the polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator;

wherein the isolated polypeptide with a molecular weight of 37 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 37 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and
15 wherein the polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator;

wherein the isolated polypeptide with a molecular weight of 36 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 36 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and wherein the polypeptide is isolatable from a *Staphylococcus aureus* when
20 incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator;


wherein the isolated polypeptide with a molecular weight of 35 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 35 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and wherein the polypeptide is isolatable from a *Staphylococcus aureus* when
25 incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator; and
30

wherein the isolated polypeptide with a molecular weight of 33 kDa has a mass fingerprint of at least 80% similarity to a mass fingerprint of a 33 kDa polypeptide expressed by *Staphylococcus aureus* ATCC strain 19636, and

wherein the polypeptide is isolatable from a *Staphylococcus aureus* when incubated in media comprising an iron chelator and not isolatable when grown in the media without the iron chelator.

- 5 56. A composition comprising;
 antibody that specifically binds to an isolated polypeptide having
 at least 92% sequence similarity to the amino acid sequence of SEQ ID
 NO:397, with the proviso that if the isolated polypeptide includes one or
10 more additional amino acids at the amino terminal, the one or more
 additional amino acids include at least one amino acid deletion or at least
 one amino acid substitution compared to amino acids 1-26 of SEQ ID
 NO:399.
57. A composition comprising:
15 antibody that specifically binds to an isolated polypeptide having
 at least 98% sequence similarity to the amino acid sequence of SEQ ID
 NO:408, with the proviso that if the isolated polypeptide includes one or
 more additional amino acids at the amino terminal, the one or more
 additional amino acids include at least one amino acid deletion or at least
20 one amino acid substitution compared to amino acids 1-5 of SEQ ID
 NO:415.


Figure 1



* Fe^{++} The bacteria grown in the presence of ferric chloride (Fe^{++})

*Dp The bacteria grown in the presence of an iron chelator (2,2 dipyridyl (Dp))

Figure 2



3 / 112

Figure 3

Kaplan-Meier survival curve showing percent survival after vaccination and homologous challenge with *S. aureus* ATCC19636




Figure 4

Kaplan-Meier survival curve showing percent survival after vaccination and heterologous challenge with *S. aureus* ATCC19636




Figure 5

Kaplan-Meier survival curve showing percent survival after passive immunization and homologous challenge with *S. aureus* ATCC19636




Figure 6

Kaplan-Meier survival curve showing percent survival after passive immunization and heterologous challenge with *S. aureus* 1477




Figure 7

LETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSKL
WEQVMQLSKEERERGGMWDMDTKVASITITSHDAGYLDKDLETIVGVQTEKPFK
RSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNCR
KAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIRL
REELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGAA
MSLGRTSTYLDIYAERDLKAGVITESEVQEIIDHFIMKLRIKVFARTPDYNELFSGD
PTWVTVSIGGVGIDGRPLVTKNFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKTY
CAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAVTIGKQMQFFGARANIAKTLL
YAINGGKDEKSGAQVGNFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLNVI
HYMHDKYSYERIEALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEEG
LVVD FEIEGDFPKYGNND DRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVL TIT
SNVVYVKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSSVAKIPYDCCKD
GISNTFSIVPKSLGKEPEDQNRNLTSM LDGYAMQCGHHLNINVFNR ETLIDAMEHP
EEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:353)

Figure 8

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASITITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTVSIGGVGIDGRPLVTKNFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQFFGARANLAKT
LLYAINGGKDEKSGAQVGNFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVD FEIEGDFPKYGNND DRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVL TI
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSM LDGYAMQCGHHLNINVFNR ETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:354)

Figure 9

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNSFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQUFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEMALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFEIEGDFPKYGNNDDEVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:355)

Figure 10

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNSFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQUFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEMALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFEIEGDFPKYGNNDDEVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:356)

Figure 11

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITeseVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNSFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEMALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFEIEGDFPKYGNNDDRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSVAKIPYDCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:357)

Figure 12

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITeseVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNSFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEMALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFEIEGDFPKYGNNDDRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSVAKIPYDCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:358)

Figure 13

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDSLSPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIVKFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNSFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEMALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFFEIEGDFPKYGNNDDEVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:359)

Figure 14

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDSLSPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIVKFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNSFRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEMALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFFEIEGDFPKYGNNDDEVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:360)

Figure 15

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSANTIGKQMQLFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFFEIEGDFPKYGNNDNRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLTII
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:361)

Figure 16

MLETNKNHATAWQGFKNGRWNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSRPAENFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITESEVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSANTIGKQMQLFFGARANLAKT
LLYAINGGKDEKSGAQVGNPFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVDFFEIEGDFPKYGNNDNRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLTII
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:362)

10 / 112

Figure 17

MLETNKNHATAWQGFKNRWRNRHVDVREFIQLNYTLYEGNDSFLAGPTEATSK
LWEQVMQLSKEERERGGMWDMDTKVASTITSHDAGYLDKDLETIVGVQTEKPF
KRSMQPFGGIRMAKAACEAYGYELDEETEKIFTDYRKTHNQGVFDAYSREMLNC
RKAGVITGLPDAYGRGRIIGDYRRVALYGVDFLMEEKMHDFNTMSTEMSEDVIR
LREELSEQYRALKELKELGQKYGFDLSPAEKFKEAVQWLYLAYLAAIKEQNGA
AMSLGRTSTFLDIYAERDLKAGVITSEVQEIIDHFIMKLRIKVFARTPDYNELFSG
DPTWVTESIGGVGIDGRPLVTKNRFLHSLDNLGPAPEPNLTVLWSVRLPDNFKT
YCAKMSIKTSSIQYENDDIMRESYGDDYGIACCVSAMTIGKQMQQFFGARANLAKT
LLYAINGGKDEKSGAQVGPNFEGINSEVLEYDEVFKKFDQMMDWLAGVYINSLN
VIHYMHDKYSYERIEALHDTEIVRTMATGIAGLSVAADSLSAIKYAQVKPIRNEE
GLVVD FEIEGDFPKYGNND DRVDDIAVDLVERFMTKLRSHKTYRDSEHTMSVLT
TSNVVYGKKTGNTPDGRKAGEPFAPGANPMHGRDQKGALSSLSSVAKIPYDCCK
DGISNTFSIVPKSLGKEPEDQNRNLTSMLDGYAMQCGHHLNINVFNRETLIDAME
HPEEYPQLTIRVSGYAVNFIKLTREQQLDVISRTFHESM

(SEQ ID NO:363)

Figure 18

KKENKQLTYTTVKDIGDMNPHVYGGSMSAESMIYEPLVRNTKDGKIPLLAKKWG
VSEDGKTYTFHLRDDVKFHDGTPFDADAVKKNIDAVQENKKLHSLWKISTLIDN
VKVKDKYTVELNLKEAYQPALAE LAMP RP YV FVSPKDFKNGTTKDG VKKFDGT
GPFKLGEHKKDESADFNKNDQYWGEKSKLNKVQAKVMPAGETAFLSMKKGETN
FAFTDDGGTDSLDDSLKQLKDTGDYQVKRSQPMNTKMLVVNSGKKDNAVSDK
TVRQAIGHMVNRDKIAKEILDGQEKPATQLFAKNVTDINFDMPTRKYDLKKAESL
LDEAGWKKGKDSVVRQKDGNLEMAMYDYDKGSSSQKEQAEYLQAEFKKMGIK
LNINGETSDKIAERRTSGDYDLMFNQTWGLLYDPQSTIAAFKAKNGMKVQHQA
RIKIKYTTALMTHLKS KTVKSVQTLIKTFNKLMMKVSLSLFHTVVQLLRKIKKY
HSHNHSTNYHSMKCSIN

(SEQ ID NO:364)

Figure 19

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVY
GSM SAE SMIYEPLVRNTKDGKIPLLAKKWVSEDGKTYTFHLRDDVKFHDGTPF
DADAVKKNIDAVQENKKLHSLWKISTLIDNVKVKDKYTVELNLKEAYQPALAE

11 / 112

AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFFKLGEHKKDESADFNKNDQYWG
 EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSDLKQLKDTG
 DYQVKRSQPMNTKMLVVNSGKKDSAVSDKTVRQAIGHMVNRDKIAKEILDGQE
 KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGKNL
 EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
 NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
 YKNILKQVNDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
 (SEQ ID NO:365)

Figure 20

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVYG
 GSMSAESMIYEPLVRNTKDGKIPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTF
 DADAVKKNIDAVQQNKKLHSLWKISTLIDNVKVKDKYTVELNLKEAYQPALAE
 AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFFKLGEHKKDESADFNKNDQYWG
 EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSDLKQLKDTG
 DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
 KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGKNL
 EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
 NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
 YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
 (SEQ ID NO:366)

Figure 21

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVYG
 GSMSAESMIYEPLVRNTKDGKIPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTF
 DADAVKKNIDAVQENKKLHSLWKISTLIDNVKVKDKYTVELNLKEAYQPALAE
 AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFFKLGEHKKDESADFNKNDQYWG
 EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSDLKQLKDTG
 DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
 KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGKNL
 EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
 NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
 YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVAFTQSQYELPFNEMQYK
 (SEQ ID NO:367)

Figure 22

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDGIKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTPF
DADAVKKNIDAVQQNKKLHSLWKISTLIDNVKVKDKYTVELNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSDLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGNL
EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSGA
YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:368)

Figure 23

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDGIKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTPF
DADAVKKNIDAVQENKKLHSLWKISTLIDNVKVKDKYTVELNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSDLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGNL
EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:369)

Figure 24

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDGIKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTTF
DADAVKKNIDAVQQNKKLHSLWKISTLIDNVKVKDKYTVELNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSDLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGNL

EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:370)

Figure 25

MRKLTKMSAMLLASGLILTGC GGNGGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDG IKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTPF
DADAVKKNIDAVQENKKLHSWLKISTLIDNVKVKDKYTVELNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDG VKKFDGTGPFKLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSL DKDSLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGKNL
EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:371)

Figure 26

MRKLTKMSAMLLTSG LILTGC GGNGGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDG IKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTPF
DADAVKKNIDAVQENKKLHSWLKISTLIDNVKVKDKYTVELNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDG VKKFDGTGPFKLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSL DKDSLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGKNL
EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDA
YKNILKQIDDEGIFIPISHGSMTVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:372)

Figure 27

MRKLTKMSAMLLASGLILTGC GGNGGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDG IKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTPF

14 / 112

DADAVKKNIDAVQENKKLHSLWKISTLIDNVKVKDKYTVLNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGNL
EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKEKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSDAY
KNILKQIDDEGIFIPISHGSM TVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:373)

Figure 28

MRKLTKMSAMLLASGLILTGCGGNKGLEEKKENKQLTYTTVKDIGDMNPHVYG
GSMSAESMIYEPLVRNTKDGIKPLLAKKWDVSEDGKTYTFHLRDDVKFHDGTPF
DADAVKKNIDAVQQNKKLHSLWKISTLIDNVKVKDKYTVLNLKEAYQPALAE
AMPRPYVVFVSPKDFKNGTTKDGVKKFDGTGPFLGEHKKDESADFNKNDQYWG
EKSKLNKVQAKVMPAGETAFLSMKKGETNFAFTDDRGTDSLKDSLKQLKDTG
DYQVKRSQPMNTKMLVVNSGKKDNAVSDKTVRQAIGHMVNRDKIAKEILDGQE
KPATQLFAKNVTDINFDMPTRKYDLKKAESLLDEAGWKKGKDS DVRQKDGNL
EMAMYYDKGSSSQKEQAEYLQAEFKKMGIKLNINGETSDKIAERRTSGDYDLMF
NQTWGLLYDPQSTIAAFKAKNGYESATSGIENKDKIYNSIDDAFKIQNGKERSGA
YKNILKQIDDEGIFIPISHGSM TVVAPKDLEKVSFTQSQYELPFNEMQYK
(SEQ ID NO:374)

Figure 29

SSDSKDKETTSIKHAMGTTEIKGKPKRVVTLYQGATDVA VSLGVKPVGAVESWT
QKPKFEYIKNDLKDTKIVGQEPAPNLEEISKLKPD LIVASKVRNEKVYDQLSKIAPT
VSTD TVFKFKDTTKLMGKALGKEKEAEDLLKKYDDKVA AFQKDAKAKYKDAW
PLKASVVNFRADHTRIYAGGYAGEILNDLGFKRNKDLQKQVDNGKDIIQLTSKESI
PLMNADHIFVVKSDPNAKDAALVKKTESEWTSSKEWKNLDAVKNNQVSDDLDEI
TWNLAGGYKSSLKLIDDLYEKLNIEKQSK
(SEQ ID NO:375)

Figure 30

MNKVIKMLVVTLAFLLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNL
EEISKLKPD LIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVA AFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK

(SEQ ID NO:376)

Figure 31

MNKVIKMLVVTLAFLLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNL
EEISKLKPD LIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVA AFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK

(SEQ ID NO:377)

Figure 32

MNKVIKMLVVTLAFLLVLAGCSGNSNKQSSDSKDKETTSIKHAMGTTEIKGKPKR
VVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNLE
EISKLKPD LIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVA AFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK

(SEQ ID NO:378)

Figure 33

MNKVIKMLVVTLAFLLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNL
EEISKLKPD LIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE

16 / 112

AEDLLKKYDDKVAAFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK
(SEQ ID NO:379)

Figure 34

MNKVIKMLVVTLAFLVLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNL
EEISKLKPDIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVAAFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK
(SEQ ID NO:380)

Figure 35

MNKVIKMLVVTLAFLVLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNL
EEISKLKPDIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVAAFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK
(SEQ ID NO:381)

Figure 36

MNKVIKMLVVTLAFLVLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKDTKIVGQEPAPNL
EEISKLKPDIVASKVRNEKVYDQLSKIAPT VSTD TVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVAAFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKLNIE
KQSK (SEQ ID NO:382)

17 / 112

Figure 37

MNKVIKMLVVTLAFLVLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKD TKIVGQEPAPNL
EEISKLPDLIVASKVRNEKVYDQLSKIAPT VSTDTVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVA AFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKL NIE
KQSK

(SEQ ID NO:383)

Figure 38

MNKVIKMLVVTLAFLVLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKD TKIVGQEPAPNL
EEISKLPDLIVASKVRNEKVYDQLSKIAPT VSTDTVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVA AFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKL NIE
KQSK

(SEQ ID NO:384)

Figure 39

MNKVIKMLVVTLAFLVLVLAGCSGNSNKQSSDNKDKETTSIKHAMGTTEIKGKPK
RVVTLYQGATDVAVSLGVKPVGAVESWTQKPKFEYIKNDLKD TKIVGQEPAPNL
EEISKLPDLIVASKVRNEKVYDQLSKIAPT VSTDTVFKFKD TTKLMGKALGKEKE
AEDLLKKYDDKVA AFQKDAKAKYKDAWPLKASVVNFRADHTRIYAGGYAGEIL
NDLGFKRNKDLQKQVDNGKDIIQLTSKESIPLMNADHIFVVKSDPNAKDAALVKK
TESEWTSSKEWKNLDAVKNNQVSDDLDEITWNLAGGYKSSLKLIDDLYEKL NIE
KQSK

(SEQ ID NO:385)

Figure 40

KESSTKDTISVKDENGTVKVPKDAKRIVVLEYSFADALAALDVKPVGIADDGKKK
RIIKPVREKIGDYTSVGTRKQPNLEEISKLPDLIIADSSRHKGINKELNKIAP TSLK
SFDGDYKQNINSFKTIAKALNKEKEGEKRLAEHDKLIK KYKDEIKFDRNQKV LPA

18 / 112

VVAKAGLLAHPNYSYVGQFLNELGFKNALSDDVTKGLSKYLKGPYLQLDTEHLA
DLNPERMIIMTDNAKKDSAEFKKLQEDPTWKKLNAVKNRVDIVDRDVWARSR
GLISSEEMAKELVELSKKEQK

(SEQ ID NO:386)

Figure 41

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKS YKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVT KIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLD FMRKDLKEKLIKAAK

(SEQ ID NO:387)

Figure 42

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKS YKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVT KIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLD FMRKDLKEKLIKAAK

(SEQ ID NO:388)

Figure 43

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKS YKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVT KIGDGDIEKVAKEKPDII
YSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
TTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQPE
QQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLD FMRKDLKEKLIKAAK

(SEQ ID NO:389)

19 / 112

Figure 44

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKWDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:390)

Figure 45

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKWDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:391)

Figure 46

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKWDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:392)

Figure 47

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKWDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:393)

Figure 48

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:394)

Figure 49

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:395)

Figure 50

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:396)

Figure 51

TEEKTEMTTIKDELGTEKIKKNPKRVVVLEYSFADYLAALDMKPVGIADDGSSKN
ITKSVRDKIGAYESVGSRSQPNMEVISKLPDLIADVSRHKKIKSELSKIAPTIMLV
SGTGDYNANIEAFKTVAKAVGREKEGEKRLEKHKILAEIRKKIEQSTLKSALG
ISRAGMFINNEDTFMGQFLIKMGIQPEVTKDKTAHVGERKGGPYIYLNNEELANIN
PKVMILATDGKTDKNRTKFIDPAVWWSLKA VKDNKVYDVDRNKWLKSRGIIASE
SMAEDLEKIAEKAK
(SEQ ID NO:397)

Figure 52

MEVISKLKPDLLIADVNRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAV
GKEKEGEKRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIK
MGIQPEVTKDKTMHVGERKGGPYTYLNNEELANINPKVMILATDGKTDKNRTKFI
DPAVWWSLKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK
(SEQ ID NO:398)

Figure 53

MNRNIVKLVVFMILLVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRVVV
LEYSFADYLAALDMKPVGIADDGSTKNITKSVRDKIGAYESVGSRPQPNMEVISK
LKPDLLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEG
EKRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIKMGIQPE
VTKDKTTHVGERKGGPYTYLNNEELANINPKVMILATDGKTDKNRTKFIIDPAVW
KSLKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK
(SEQ ID NO:399)

Figure 54

MNRNIVKLVVFMILLVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRIVVL
EYSFADYLAALDMKPVGIADDGSTKNITKSVRDKIGAYESVGSRPQPNMEVISKL
KPDLLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEGE
KRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIKMGIQPEV
TKDKTAHVGERKGGPYTYLNNEELANINPKVMILATNGKTDKNRTKFIIDPAVWKS
LKAVKDNKVYDVDRNKWLQSRGIMASESMAEDLEKIAEKAK
(SEQ ID NO:400)

Figure 55

MNRNIVKLVVFMILLVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRVVV
LEYSFADYLAALDMKPVGIADDGSSKNITKSVRDKIGAYESVGSRPQPNMEVISKL
KPDLLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEGE
KRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLLKMGIQPEV
TKDKTTHVGERKGGPYTYLNNEELANINPKVMILATDGKTDKNRTKFIIDPAVWKS
LKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK
(SEQ ID NO:401)

Figure 56

MNRNIVKL VVFMLIL VVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRVVV
LEYSFADYLAALDMKPVGIADDGSTKNITKSVRDKIGAYESVGSRPQPNMEVISK
LKPDLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEG
EKRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIKMGIQPE
VTKDKTTHVGERKGGPYIYLNNEELANINPKVMILATDGKTDKNRTKFIDPAVW
KSLKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK
(SEQ ID NO:402)

Figure 57

MNRNIVKL VVFMLIL VVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRVVV
LEYSFADYLAALDMKPVGIADDGSTKNITKSVRDKIGAYESVGSRPQPNMEVISK
LKPDLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEG
EKRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIKMGIQPE
VTKDKTTHVGERKGGPYIYLNNEELANINPKVMILATDGKTDKNRTKFIDPAVW
KSLKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK
(SEQ ID NO:403)

Figure 58

MTTIKDELGTEKIKKNPKRVVVLEYSFADYLAALDMKPVGIADDGSTKNITKSVR
DKIGAYESVGSRPQPNMEVISKLKPDLIADVSRHKKIKSELSKIAPTIMLVSGTGD
YNANIEAFKTVAKAVGKEKEGEKREKLEKHDKILAEIRKKIEQSTLKSFAFGISRAG
MFINNEDTFMGQFLIKMGIQPEVTKDKTTHVGERKGGPYIYLNNEELANINPKVMI
LATDGKTDKNRTKFIDPAVWKSLKAVKDNKVYDVDRNKWLKSRGIIASESMAED
LEKIAEKAK
(SEQ ID NO:404)

Figure 59

MNRNIVKL VVFMLIL VVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRVVV
LEYSFADYLAALDMKPVGIADDGSTKNITKSVRDKIGAYESVGSRPQPNMEVISK
LKPDLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEG
EKRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIKMGIQPE
VTKDKTTHVGERKGGPYIYLNNEELANINPKVMILATDGKTDKNRTKFIDPAVW
KSLKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK
(SEQ ID NO:405)

Figure 60

MTTIKDELGTEKIKKNPKRVVVLEYSFADYLAALDMKPVGIADDGSTKNITKSVR
DKIGAYESVGSRPQPNMEVISKLPDLIIADVSRHKKIKSELSKIAPTIMLVSGTGD
YNANIEAFKTVAKAVGKEKEGEKRLEKHDKILAEIRKKIEQSTLKSFAFGISRAG
MFINNEDTFMGQFLIKMGIQPEVTKDKTTHVGERKGGPYIYLNNEELANINPKVMI
LATDGKTDKNRTKFIDPAVWWSLKAVKDNKVYDVDRNKWLKSRGIIASESMAED
LEKIAEKAK

(SEQ ID NO:406)

Figure 61

MNRNIVKLVVFMILVAVAGCGQKDTEEKTEMTTIKDELGTEKIKKNPKRVVV
LEYSFADYLAALDMKPVGIADDGSSKNITKSVRDKIGAYESVGSRPQPNMEVISKL
KPDLLIADVSRHKKIKSELSKIAPTIMLVSGTGDYNANIEAFKTVAKAVGKEKEGE
KRLEKHDKILAEIRKKIEQSTLKSFAFGISRAGMFINNEDTFMGQFLIKMGIQPEV
TKDKTTHVGERKGGPYIYLNNEELANINPKVMI
LATDGKTDKNRTKFIDPAVWWSLKAVKDNKVYDVDRNKWLKSRGIIASESMAEDLEKIAEKAK

(SEQ ID NO:407)

Figure 62

GSDDNGSSKSPYHRIVSLMPSNTEILYELGLGKYIVGVSTVDDYPKDVKKGKKQF
DALNLNKEELLKAKPDLILAHESQKATANKVLSSLEKQGIKVYVYKDAQSIDETY
NTFKQIGKLTHHDKQAEQLVEETKDNIDKVIDSIPAHKKSKVFIEVSSKPEIYTAG
KHTFFNDMLEKLEAQNVSIDINGWNPVTKESIHKKNPDILISTEAKTRSDYMDIHK
RGGFNKINAVKNTRIEVVNGDEVSRPGPRIDEGLKELRDAIYRK

(SEQ ID NO:408)

Figure 63

MKKSLIAFILIFMLVLSGCGMKDNDKQGSDDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKKGKKQFDALNLNKEELLKAKPDLILAHESQKATA
NKVLSSLEKQGIKVYVYKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDNIDK
VIDSIPAHKKSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVSIDINGWNPVT
KESIHKKNPDILISTEAKTRSDYMDIHKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK

(SEQ ID NO:409)

Figure 64

MKKS LIAFILIFMLVLSGCGMKDNDKQGSDDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKEGKKQFDALNLNKEELLKAKPDLILAHESQKATA
NKVLSSLEKQGKIKVVYVKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDNID
KVIDSIPAHHKSSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVS DINGWNPV
TKESIHKKNPDILISTEAKTRSDYMDIHKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK
(SEQ ID NO:410)

Figure 65

MKKS LIAFILIFMLVLSGCGMKDNDKQGSDDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKKGKKQFDALNLNKEELLKAKPDLILAHESQKATA
NKVLSSLEKQGKIKVVYVKDAQSIDETYNTFKQIGKLTHEKQAEQLVEETKDNID
KVIDSIPAHHKSSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVS DINGWNPV
TKESIHKKNPDILISTEAKTRSDYMDIHKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK
(SEQ ID NO:411)

Figure 66

MKKS LIAFILIFMLVLSGCGMKDNDKQGSNDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKEGKKQFDALNLNKEELLKEKPDLILAHESQKATA
NKVLSSLEKQGKIKVVYVKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDNID
KVIDSIPAHHKSSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVS DINGWNPV
TKESIHKKNPDILISTEAKTRSDYMDIHKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK
(SEQ ID NO:412)

Figure 67

MKKS LIAFILIFMLVLSGCGMKDNDKQGSNDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKKGKKQFDALNLNKEELLKAKPDLILAHESQKATA
NKVLSSLEKQGKIKVVYVKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDNID
KVIDSIPAHHKSSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVS DINGWNPV
TKESIHKKNPDILISTEAKTRSDYMDIHKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK
(SEQ ID NO:413)

25 / 112

Figure 68

MKKSIIAFILIFMLVLSGCGMKDNDKQGSDDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKEGKKQFDALNLNKEELLKAKPDLILAHESQKATA
NKVLSSLEKQGIKVYVVKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDND
KVIDSIPAHKKSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVSINGWNPV
TKESIHKKNPDILISTEAKTRSDYMDIHKKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK

(SEQ ID NO:414)

Figure 69

MKDNDKQGSNDNGSSKSPYHRIVSLMPSNTEILYELGLGKYIVGVSTVDDYPKDV
KKGKKQFDALNLNKEELLKAKPDLILAHESQKATANKVLSSLEKQGIKVYVVKD
AQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDNDKVIDSIPAHKKSKVFIEVSS
KPEIYTAGKHTFFNDMLEKLEAQNVSINGWNPVTKESIHKKNPDILISTEAKTRS
DYMDIHKKRGGFNKINAVKNTRIEVVNGDEVSRPGPRIDEGLKELRDAIYRK

(SEQ ID NO:415)

Figure 70

MKKSIIAFILIFMLVLSGCGMKDNDKQGSNDNGSSKSPYHRIVSLMPSNTEILYEL
GLGKYIVGVSTVDDYPKDVKKGKKQFDALNLNKEELLKAKPDLILAHESQKATA
NKVLSSLEKQGIKVYVVKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDND
KVIDSIPAHKKSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVSINGWNPV
TKESIHKKNPDILISTEAKTRSDYMDIHKKRGGFNKINAVKNTRIEVVNGDEVSRPGP
RIDEGLKELRDAIYRK

(SEQ ID NO:416)

Figure 71

MKDNDKQGSNDNGSSKSPYHRIVSLMPSNTEILYELGLGKYIVGVSTVDDYPKDV
KKGKKQFDALNLNKEELLKAKPDLILAHESQKATANKVLSSLEKQGIKVYVVKD
AQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDNDKVIDSIPAHKKSKVFIEVSS
KPEIYTAGKHTFFNDMLEKLEAQNVSINGWNPVTKESIHKKNPDILISTEAKTRS
DYMDIHKKRGGFNKINAVKNTRIEVVNGDEVSRPGPRIDEGLKELRDAIYRK

(SEQ ID NO:417)

Figure 72

MKKSILAFILIFMLVLSGCGMKDNDKQGSNDNGSSKSPYHRIVSLMPNTEILYEL
GLGKYIVGVSTVDDYPKDVKEGKKQFDALNLNKEELLKEKPDLILAHESQKATA
NKVLSSLEKQGKVVVYVKDAQSIDETYNTFKQIGKLTHHDKQAEQLVEETKDND
KVIDSIPAHKKSKVFIEVSSKPEIYTAGKHTFFNDMLEKLEAQNVSNDINGWNPV
TKESIKKNPDLISTEAKTRSDYMDIKKRGGFNKINAVKNTRIEVVNGDEVSRPGP
R IDEGLKELRDAIYRK

(SEQ ID NO:418)

Figure 73

SDKSNGKLVVTTNSILYDMAKNVGGDNVDIHSIVPVGQDPHEYEVKPKDIKKLT
DADVILYNGLNLETGNGWFEKALEQAGKSLKDKKVIKAVSKDVKPIYLNKEEGNK
DKQDPHAWLSLDNGIKYVKTIQQTFIDNDKKHKADYEKQGNKYIAQLEKLNND
KDKFNDIPKEQRAMITSEGAFKYFSKQYGITPGYIWEINTEKQGTPEQMRQAIEFV
KKHKLKHLLVETSVDDKAMESLSEETKKDIFGEVYTD SIGKEGTKGDSYYKMMK
SNIETVHGSMK

(SEQ ID NO:419)

Figure 74

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKKLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIKAVSKDVKPIYLNKEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNNDKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLLVETSVDDKAMESLSEETKKD
IFGEVYTD SIGKEGTKGDSYYKMMKSNIETVHGSMK

(SEQ ID NO:420)

Figure 75

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKKLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIKAVSKDVKPIYLNKEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNNDKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLLVETSVDDKAMESLSEETKKD
IFGEVYTD SIGKEGTKGDSYYKMMKSNIETVHGSMK

(SEQ ID NO:421)

Figure 76

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK
(SEQ ID NO:422)

Figure 77

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK
(SEQ ID NO:423)

Figure 78

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK
(SEQ ID NO:424)

Figure 79

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK
(SEQ ID NO:425)

Figure 80

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK

(SEQ ID NO:426)

Figure 81

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK

(SEQ ID NO:427)

Figure 82

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSK
QYGITPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEE
TKKDIFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK

(SEQ ID NO:428)

Figure 83

MKKLVPLLLALLLLVAACGTGGKQSSDKSNGKLVVTTNSILYDMAKNVGGDN
VDIHSIVPVGQDPHEYEVKPKDIKCLTDADVILYNGLNLETGNGWFEKALEQAGK
SLKDKKVIAVSKDVKPIYLN GEEGNKDKQDPHAWLSLDNGIKYVKTIQQTFIDND
KKHKADYEKQGNKYIAQLEKLNND SKDKFNDIPKEQRAMITSEGAFKYFSKQYGI
TPGYIWEINTEKQGTPEQMRQAIEFVKKHKLKHLVETSVDKKAMESLSEETKKD
IFGEVYTDSIGKEGTKGDSYYKMMKSNIETVHGSMK

(SEQ ID NO:429)

Figure 84

TTAGAAACAAATAAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATGGAA
GATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAAACTACACTCTT
TATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTAAACT
TTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGGCATG
TGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTGGTTA
TTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAAACTGAAAAGCCATTCA
AACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTGTGAA
GCTTACGGTTACGAATTAGACGAAGAAACTGAAAAAATCTTTACAGATTATCG
TAAACACATAACCAAGGTGTATTTCGATGCATATTCTAGAGAAATGTTGAACT
GCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGGACGT
ATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAATGGAA
GAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGATGTAA
TTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATTAAAA
GAACTTGGGCAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAACCTCAA
AGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGAACAAA
ACGGTGCAGCAATGAGTTTAGGTTCGTACATCAACATACTTAGATATCTATGCT
GAACGTGACCTTAAAGCAGGTGTTATTACTGAAAGCGAAGTTCAAGAAATTAT
TGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACCTGATTA
CAATGAATTATTCTCTGGAGACCCAACTTGGGTAAGTGTATCTATCGGTGGTG
TAGGTATTGACGGACGTCCACTTGTTACGAAAAACTCATTCCGTTTCTTACACT
CATTAGATAACTTAGGTCCAGCACCAGAACCAAACTTAACAGTATTATGGTCA
GTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTATTA AAC
AAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATGGCGATG
ACTATGGTATCGCATGTTGTGTATCAGCGGTGACAATTGGTAAACAAATGCAA
TTCTTCGGTGCACGTGCGAACATAGCTAAAACATTACTTTACGCTATCAATGG
TGGTAAAGATGAAAAGTCTGGTGCACAAGTTGGTCCAACTTCGAAGGTATTA
ACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAAATGAT
GGATTGGCTAGCAGGTGTTTACATTA ACTCATTAAATGTTATTCACTACATGCA
CGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGAAATTG
TACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACTCATT
TCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGTCTTGT
AGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGACGACC
GTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTTATGACTAAATTACGT

30 / 112

AGTCATAAAACATATCGTGATTGAGAACATACAATGAGTGTATTAACAATTAC
TTCAAACGTTGTATACGTTAAGAAAACCTGGTAACACACCAGACGGACGTAAA
GCTGGCGAACCATTTGCACCAGGTGCAAACCCAATGCATGGCCGTGACCAAA
AAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTGCTGTA
AAGATGGTATTTCAAATACATTTCAGTATCGTACCAAAAATCATTAGGTAAAGAA
CCAGAAGATCAAAACCGTAACTTAACTAGTATGTTAGATGGTTACGCAATGCA
ATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAATAGATG
CAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTGGTTAC
GCTGTAACTTCATTAAATTAACACGTGAACAACAGTTAGATGTAATTTCTCGT
ACATTCCATGAAAGTATGTAA
(SEQ ID NO:430)

Figure 85

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGTGTTATTACTGAAAGCGAAGTTCAAGAA
ATTATTGACCACTTCATCATGAAATTACGTATTGTAAATTTGCTCGTACACCT
GATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACTGAATCTATCGG
TGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCTT
ACACTCATTAGATAACTTAGGTCCAGCACCAGAACCAAACCTTAACAGTATTAT

GGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTATT
AAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATGG
CGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAAA
TGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATCA
ATGGTGGTAAAGATGAAAAATCTGGCGCACAAGTTGGTCCAAACTTCGAAGG
TATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAAA
TGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTACA
TGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGAA
ATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACTC
ATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGTC
TTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGAC
GACCGTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTTATGACTAAATT
ACGTAGTCATAAAACATATCGTGATTTCAGAACATACAATGAGTGTATTAACAA
TTACTTCAAACGTTGTATACGGTAAGAAAACCTGGTAACACACCAGACGGACGT
AAAGCTGGCGAACCATTTGCACCAGGTGCAAACCCAATGCATGGCCCGTGACC
AAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTGCT
GTAAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTAAA
GAACCAGAAGATCAAAACCGTAACTTAACTAGTATGTTAGATGGTTACGCAAT
GCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAATAG
ATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTGGT
TACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAATTTCT
CGTACATTCCATGAAAGTATGTAA
(SEQ ID NO:431)

Figure 86

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG

AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAAGCTTGGACAAAAATATGGTTTTCGATTTAAGCCGTCCAGCAGAAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAAACCTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGCGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCAAT
CAATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAA
GGTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCA
AATGATGGATTGGCTAGCAGGTGTTTACATTAACCTATTAAATGTTATTCTACTA
CATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAG
AAATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGAC
TCATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGG
TCTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATG
ACGACCGTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTCATGACTAAA
TTACGTAGTCATAAAACATATCGTGATTGAGAACATACAATGAGTGTATTAAC
AATTACTTCAAACGTTGTATACGGTAAGAAAACTGGTAACACACCAGACGGA
CGTAAAGCTGGCGAACCATTTGCTCCAGGTGCAAACCCAATGCATGGCCGTGA
CCAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTG
CTGTAAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTA
AAGAACCAGAAGATCAAAACCGTAACCTTAAGTATGTTAGATGGTTACGC
AATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAA
TAGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCT

GGTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAAT
TTCTCGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:432)

Figure 87

ATGTTAGAAACAAATAAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCAATTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGGCGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGTGTTATTACTGAAAGCGAAGTTCAAGAA
ATTATTGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACCT
GATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACTGAATCTATCGG
TGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCTT
ACACTCATTAGATAACTTAGGTCCAGCACCAGAACCAAACCTTAACAGTATTAT
GGTCAGTACGTTTACCAGACAACTTCAAAACATACTGTGCAAAAATGAGTATT
AAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATGG
CGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAAA
TGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATCA
ATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAAGG
TATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAAA
TGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTACA
TGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGAA

ATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACTC
ATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGTC
TTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGAC
GACCGTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTTATGACTAAATT
ACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACAA
TACTTCAAACGTTGTATACGGTAAGAAAACCTGGTAACACACCAGACGGACGT
AAAGCTGGCGAACCATTTGCACCAGGTGCAAACCCAATGCATGGCCGTGACC
AAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATTCCTTACGATTGCT
GTAAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTAAA
GAACCAGAAGATCAAAACCGTAACTTAACTAGTATGTTAGATGGTTACGCAAT
GCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAATAG
ATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTGGT
TACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAATTTCT
CGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:433)

Figure 88

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTCCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAACACATAACCAAGGTGTATTCGATGCATATTCCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAACTTGGACAAAAATATGGTTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA

35 / 112

AATTATTGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACCTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAAACCTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATC
AATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAAG
GTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAA
ATGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTAC
ATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGA
AATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACT
CATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGT
CTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGA
CGACCGTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTTATGACTAAAT
TACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACA
ATTACTTCAAACGTTGTATACGGTAAGAAAACCTGGTAACACACCAGACGGAC
GTAAAGCTGGCGAACCATTTGCACCAGGTGCAAACCCAATGCATGGCCGTGA
CCAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTG
CTGTAAAGATGGTATTTCAAATACATTCAGTATCGTACCAAAAATCATTAGGTA
AAGAACCAGAAGATCAAAACCGTAACTTAAGTATGTTAGATGGTTACGC
AATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAA
TAGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCT
GGTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAAT
TTCTCGTACATTCCATGA AAGTATGTAA

(SEQ ID NO:434)

Figure 89

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG

GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAACTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATC
AATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAAG
GTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAA
ATGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTAC
ATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGA
AATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACT
CATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGT
CTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGA
CGACCGTGTAGATGATATTGCAGTTGATTTAGTAGAACGCTTCATGACTAAAT
TACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACA
ATTACTTCAAACGTTGTATACGGTAAGAAAACTGGTAACACACCAGACGGAC
GTAAAGCTGGCGAACCATTTGCTCCAGGTGCAAACCCAATGCATGGCCGTGAC
CAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTGC
TGTAAGATGGTATTTCAAATACATTCAGTATCGTACCAAAATCATTAGGTAA
AGAACCAGAAGATCAAAACCGTAACTTAAGTAGTATGTTAGATGGTTACGCA

37 / 112

ATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAAT
AGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTG
GTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAATT
TCTCGTACATTCCATGAAAGTATGTAA
(SEQ ID NO:435)

Figure 90

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCTGATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACCTTGGGTAACTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAAACCTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGCGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCAAT
CAATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAA
GGTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCA

38 / 112

AATGATGGATTGGCTAGCAGGTGTTTACATTAAC TCATTAATGTTATTCACTA
CATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAG
AAATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGAC
TCATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGG
TCTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATG
ACGACCGTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTCATGACTAAA
TTACGTAGTCATAAAACATATCGTGATT CAGAACATACAATGAGTGTATTAAC
AATTACTTCAAACGTTGTATACGGTAAGAAA ACTGGTAACACACCAGACGGA
CGTAAAGCTGGCGAACCATTTGCTCCAGGTGCAAACCCAATGCATGGCCGTGA
CCAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTG
CTGTAAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTA
AAGAACCAGAAGATCAAAACCGTA ACTTAAGTAGTATGTTAGATGGTTACGC
AATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAA
TAGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCT
GGTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAAT
TTCTCGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:436)

Figure 91

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTCCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGA ACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA

39 / 112

ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACCTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAAACCTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATC
AATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAAG
GTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAA
ATGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTAC
ATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGA
AATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACT
CATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGT
CTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGA
CGACCGTGTAGATGATATTGCAGTTGATTTAGTAGAACGCTTCATGACTAAAT
TACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACA
ATTACTTCAAACGTTGTATACGGTAAGAAAACCTGGTAACACACCAGACGGAC
GTAAAGCTGGCGAACCATTTGCTCCAGGTGCAAACCCAATGCATGGCCGTGAC
CAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTGC
TGTAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTAA
AGAACCAGAAGATCAAAACCGTAACTTAACCTAGTATGTTAGATGGTTACGCA
ATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAAT
AGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTG
GTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAATT
TCTCGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:437)

Figure 92

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA

AAC TTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACTTGGGTAACCTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACC AAACTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATC
AATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTTCGAAG
GTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAA
ATGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTAC
ATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGA
AATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACT
CATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGT
CTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGA
CGACCGTGTAGATGATATTGCAGTTGATTTAGTAGAACGCTTCATGACTAAAT
TACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACA
ATTACTTCAAACGTTGTATACGGTAAGAAAACTGGTAACACACCAGACGGAC
GTAAAGCTGGCGAACCATTTGCTCCAGGTGCAAACCCAATGCATGGCCGTGAC
CAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTGC

41 / 112

TGTAAAGATGGTATTTCAAATACATTTCAGTATCGTACCAAAATCATTAGGTAA
AGAACCAGAAGATCAAAACCGTAACTTAAGTAGTATGTTAGATGGTTACGCA
ATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAAT
AGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTG
GTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAATT
TCTCGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:438)

Figure 93

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT
AAAAGAAGCTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAAGTTGGGTAACTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAAACTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATC

42 / 112

AATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAAACTTCGAAG
GTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAA
ATGATGGATTGGCTAGCAGGTGTTTACATTAAC TCATTAAATGTTATTCACTAC
ATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGA
AATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACT
CATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGT
CTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGA
CGACCGTGTAGATGATATTGCAGTTGATTTAGTAGAACGCTTCATGACTAAAT
TACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACA
ATTACTTCAAACGTTGTATACGGTAAGAAAAC TGGTAACACACCAGACGGAC
GTAAAGCTGGCGAACCATTTGCTCCAGGTGCAAACCCAATGCATGGCCGTGAC
CAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTGC
TGTAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTAA
AGAACCAGAAGATCAAACCGTAAC TTAAC TAGTATGTTAGATGGTTACGCA
ATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAAT
AGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCTG
GTTACGCTGTTAAC TTCATTAAATTAACACGTGAACAACAATTAGATGTAATT
TCTCGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:439)

Figure 94

ATGTTAGAAACAAATAAAAATCATGCAACAGCTTGGCAAGGATTTAAAAATG
GAAGATGGAACAGACACGTAGATGTAAGAGAGTTTATCCAATTAACTACAC
TCTTTATGAAGGTAATGATTCATTTTTAGCAGGACCAACAGAAGCAACTTCTA
AACTTTGGGAACAAGTAATGCAGTTATCGAAAGAAGAACGTGAACGTGGCGG
CATGTGGGATATGGACACGAAAGTAGCTTCAACAATCACATCTCATGATGCTG
GTTATTTAGACAAAGATTTAGAAACAATTGTAGGTGTACAACTGAAAAGCCA
TTCAAACGTTCAATGCAACCATTTCGGTGGTATTCGTATGGCGAAAGCAGCTTG
TGAAGCTTACGGTTACGAATTAGACGAAGAACTGAAAAAATCTTTACAGATT
ATCGTAAACACATAACCAAGGTGTATTCGATGCATATTCTAGAGAAATGTTG
AACTGCCGTAAAGCAGGTGTAATCACTGGTTTACCTGATGCATACGGACGTGG
ACGTATTATCGGTGACTATCGTCGTGTAGCTTTATATGGTGTAGATTTCTTAAT
GGAAGAAAAAATGCACGACTTCAACACGATGTCTACAGAAATGTCAGAAGAT
GTAATTCGTTTACGTGAAGAATTATCAGAACAATATCGTGCATTAAAAGAATT

AAAAGAACTTGGACAAAAATATGGTTTCGATTTAAGCCGTCCAGCAGAAAAC
TTCAAAGAAGCAGTTCAATGGTTATACTTAGCATACCTTGCTGCAATTAAAGA
ACAAAACGGTGCAGCAATGAGTTTAGGTCGTACATCAACATTCTTAGATATCT
ATGCTGAACGTGACCTTAAAGCAGGCGTTATTACTGAAAGCGAAGTTCAAGA
AATTATTGACCACTTCATCATGAAATTACGTATTGTAAATTTGCTCGTACACC
TGATTACAATGAATTATTCTCTGGAGACCCAACCTTGGGTAACCTGAATCTATCG
GTGGTGTAGGTATTGACGGACGTCCACTTGTTACGAAAACTCATTCCGTTTCT
TACACTCATTAGATAACTTAGGTCCAGCTCCAGAACCAAACCTTAACAGTATTA
TGGTCAGTACGTTTACCTGACAACTTCAAAACATACTGTGCAAAAATGAGTAT
TAAAACAAGTTCTATCCAATATGAAAATGATGACATTATGCGTGAAAGCTATG
GCGATGACTATGGTATCGCATGTTGTGTATCAGCGATGACAATTGGTAAACAA
ATGCAATTCCTTCGGTGCACGTGCGAACTTAGCTAAAACATTACTTTACGCTATC
AATGGTGGTAAAGATGAAAAATCTGGTGCACAAGTTGGTCCAACTTCGAAG
GTATTAACAGCGAAGTATTAGAATATGACGAAGTATTCAAGAAATTTGATCAA
ATGATGGATTGGCTAGCAGGTGTTTACATTAACCTCATTAAATGTTATTCACTAC
ATGCACGATAAATACAGCTATGAACGTATTGAAATGGCATTACATGATACAGA
AATTGTACGTACAATGGCAACAGGTATCGCTGGTTTATCAGTAGCAGCTGACT
CATTATCTGCAATTAAATATGCACAAGTTAAACCAATTCGTAACGAAGAAGGT
CTTGTAGTAGACTTTGAAATCGAAGGCGACTTCCCTAAATACGGTAACAATGA
CGACCGTGTAGATGATATCGCAGTTGATTTAGTAGAACGCTTTATGACTAAAT
TACGTAGTCATAAAACATATCGTGATTCAGAACATACAATGAGTGTATTAACA
ATTACTTCAAACGTTGTATACGGTAAGAAAACCTGGTAACACACCAGACGGAC
GTAAAGCTGGCGAACCATTTGCACCAGGTGCAAACCCAATGCATGGCCGTGA
CCAAAAAGGTGCATTATCTTCATTAAGTTCTGTAGCTAAGATCCCTTACGATTG
CTGTAAAGATGGTATTTCAAATACATTCAGTATCGTACCAAATCATTAGGTA
AAGAACCAGAAGATCAAAACCGTAACTTAACCTAGTATGTTAGATGGTTACGC
AATGCAATGTGGTCACCACTTAAATATTAACGTATTTAACCGTGAAACATTAA
TAGATGCAATGGAACATCCAGAAGAATATCCACAGTTAACAATCCGTGTATCT
GGTTACGCTGTAACTTCATTAAATTAACACGTGAACAACAATTAGATGTAAT
TTCTCGTACATTCCATGAAAGTATGTAA

(SEQ ID NO:440)

Figure 95

AAAAAAGAAAACAAGCAATTAACGTATACGACGGTTAAAGATATCGGGGATA
TGAACCCGCATGTTTACGGTGGATCGATGTCTGCTGAAAGTATGATATACGAG
CCGCTTGTACGTAACACGAAAGACGGGATTAAGCCTTTACTAGCTAAAAAATG
GGGTGTGTCTGAAGATGGGAAGACATACACGTTCCATTTGAGAGATGACGTTA
AATTCCATGATGGTACGCCATTTGATGCTGACGCAGTTAAGAAAAATATTGAC
GCAGTTCAAGAAAACAAAAAATTGCATTCTTGGTTAAAGATTTCAACATTAAT
TGACAATGTTAAAGTTAAAGATAAGTACACGGTTGAATTGAATTTGAAAGAA
GCATATCAACCTGCATTGGCTGAATTAGCGATGCCTCGTCCATATGTATTTGTG
TCTCCAAAAGACTTTAAAAACGGCACAACAAAAGATGGCGTTAAAAAGTTCTG
ATGGTACTGGTCCGTTTAAATTAGGTGAACACAAAAAAGATGAGTCTGCAGAC
TTTAACAAAAATGATCAATACTGGGGCGAAAAGTCTAACTTAACAAAGTAC
AAGCAAAAGTAATGCCTGCTGGTGAAACAGCATTCTATCAATGAAAAAAGG
TGAAACGAACTTTGCCTTCACAGATGATGGAGGTACAGATAGCTTAGACAAA
GACTCTTTAAACAATTGAAAGATACTGGTGACTATCAAGTTAAGCGTAGTCA
ACCTATGAATACGAAAATGTTAGTTGTCAATTCTGGTAAAAAAGATAACGCTG
TCAGTGACAAAACAGTCAGACAAGCGATTGGTCATATGGTAAACAGAGATAA
AATTGCCAAAGAAATTTTAGATGGTCAAGAAAAACCAGCAACTCAATTATTTG
CGAAAAATGTAACAGACATTAATTTGATATGCCAACACGTAAGTATGACCTT
AAAAAAGCAGAATCATTATTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACA
GCGATGTTCGTCAAAAAGATGGTAAAAACCTTGAAATGGCAATGTACTATGAC
AAAGGTTCTTCAAGTCAAAAAGAACAAGCAGAATACTTACAAGCAGAATTTA
AGAAAATGGGTATTAAGTTAAACATCAATGGCGAAACATCAGATAAAATTGC
TGAACGTCGTACTTCTGGTGATTATGACTTAATGTTCAACCAAACCTTGGGGATT
ATTGTACGATCCACAAAGTACTATTGCAGCATTTAAAGCGAAAAATGGTATGA
AAGTGCAACATCAGGCATTGAGAATAAAGATAAAATATACAACAGCATTGAT
GACGCATTTAAAATCCAAAACGGTAAAGAGCGTTCAGACGCTTATAAAAACA
TTTTGAAACAAGTTGATGATGAAGGTATCTTTATCCCTGTTTCACACGGTAGTA
TGACAGTTGTTGCGCCGAAAGATTTAGAAAAAGTATCATTCACACAATCACAG
TACGAATTACCATTCAATGAAATGCAGTATAAATAA

(SEQ ID NO:441)

Figure 96

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGGGATATGAACCCGCATGTTTACGG
TGGATCGATGTCTGCTGAAAGTATGATATACGAGCCACTTGTACGTAACACGA
AAGATGGGATTAAGCCTTTATTAGCTAAAAAATGGGATGTTTCTGAAGATGGG
AAGACATATACATTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACGCC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGACGCGGTTCAAGAAAACAAA
AAATTGCATTCTTGGTTAAAGATTTCAACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGCACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTAA
ATTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAA
TACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTG
CTGGTGAAACAGCATTCCCTATCAATGAAAAAAGGTGAAACGAACTTTGCCTTC
ACAGATGATAGAGGTACAGATAGCTTAGACAAAGACTCTTTAAAACAATTGA
AAGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAAT
GTTAGTTGTCAATTCTGGTAAAAAAGATAGCGCTGTCAGTGACAAAACAGTCA
GACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTT
AGATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGAC
ATTAATTTTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
ATTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAA
GATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCA
AAAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAAATGGGTATTAAG
TTAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGG
TGATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAA
GTACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGC
ATTGAGAATAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCC
AAAACGGTAAAGAGCGTTCAGACGCTTATAAAAAACATTTTGAAACAAGTTAA
TGATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCGCC
GAAAGATTTAGAAAAAGTATCATTACACAATCACAGTATGAATTACCATTCA
ATGAAATGCAGTATAAATAA

(SEQ ID NO:442)

Figure 97

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCAATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGTATTAAGCCTTTACTAGCTAAAAAATGGGATGTGTCTGAAGATGGG
AAGACTTATACATTCCATTTAAGAGATGATGTGAAATTCCATGATGGAACAAC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGATGCGGTTCAACAAAATAAAA
AATTGCATTCTTGGTTAAAGATTTCAACATTAATTGACAATGTTAAAGTTAAA
GATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGGC
TGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTTCCTATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACGGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTCAGTGACAAAACAGTCAGA
CAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTAG
ATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGACAT
TAATTTTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATTAT
TAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGTGATGTTTCGTCAAAAAGA
TGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAAA
AAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGTT
AAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGTG
ATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAAGT
ACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCAT
TGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCCAA
AACGGTAAAGAGCGTTCAGACGCTTATAAAAACATTTTGAAACAAATTGATG
ATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCGCCGA
AAGATTTAGAAAAAGTATCATTACACAATCACAGTATGAATTACCATTCAAT
GAAATGCAGTATAAATAA

(SEQ ID NO:443)

Figure 98

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGGGATATGAATCCGCATGTTTACGG
TGGATCGATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGGATTAAGCCTTTACTAGCTAAAAAATGGGATGTGTCAGAAGATGG
GAAGACATACACGTTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACGC
CATTTGATGCTGACGCAGTTAAGAAAAATATTGACGCAGTTCAAGAAAACAA
AAAATTGCATTCTTGGTTAAAGATTTCAACATTAATTGACAATGTAAAGTTA
AAGATAAGTACACGGTTGAATTAAATTTGAAAGAAGCATATCAACCTGCATTG
GCTGAATTAGCGATGCCTCGTCCATATGTATTTGTATCTCCAAAAGACTTTAAA
AACGGCACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTA
AATTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAATAAAAAATGATCA
ATACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCT
GCTGGTGAAACAGCATTCCATCAATGAAAAAAGGTGAAACAACTTTGCCTT
CACAGACGATAGAGGTACAGATAGCTTAGACAAAGACTCTTTAAAACAATTG
AAAGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAA
TGTTAGTGGTCAATTCTGGTAAAAAAGATAACGCTGTCAGTGACAAAACAGTC
AGACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTT
TAGATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGA
CATTAAATTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCAT
TATTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAA
AGATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTC
AAAAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAA
GTTAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTG
GTGATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAA
AGTACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAG
GCATTGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAT
CCAAAACGGTAAAGAGCGTTCAGACGCTTATAAAAACATTTTGAAACAAATT
GATGATGAAGGTATTTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCG
CCGAAAGATTTAGAAAAAGTAGCATTACACAATCACAGTATGAATTACCATT
CAATGAAATGCAGTATAAATAA

(SEQ ID NO:444)

Figure 99

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCGATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGAATTAAGCCTTTACTAGCTAAAAAATGGGATGTGTCTGAAGATGG
GAAGACATATACATTCCATTTGAGAGATGACGTAAATTCCATGATGGTACAC
CATTTGATGCTGACGCAGTTAAGAAAAATATTGATGCGGTTCAACAAAATAAA
AAATTGCATTCTTGGTTAAAGATTTCAACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACATAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTCCATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACGGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTGAGTGACAAAACAGTCAG
ACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTA
GATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGATA
TTAATTTTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
TTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAAG
ATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAA
AAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGT
TAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGT
GATTATGACTTAATGTTCAACCAAACTTGGGGATTATTGTACGATCCACAAAG
TACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCA
TTGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCCA
AAACGGTAAAGAGCGTTCAGGCGCTTATAAAAACATTTTGAAACAAATTGAT
GATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCGCCG
AAAGATTTAGAAAAAGTATCATTCACACAATCACAGTATGAATTACCATTCAA
TGAAATGCAGTATAAATAA

(SEQ ID NO:445)

Figure 100

ATGAGAAAATACTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCAATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGTATTAAGCCTTTACTAGCTAAAAAGTGGGATGTGTCTGAAGATGGG
AAGACATACACGTTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACGCC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGACGCAGTTCAAGAAAACAAA
AAATTGCATTCTTGTTAAAGATTTTCGACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTCCATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACAGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTGAGTGACAAAACAGTCAG
ACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTA
GATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGACA
TTAATTTGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
TTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAAG
ATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAA
AAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGT
TAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGT
GATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAAG
TACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCA
TTGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCCA
AAACGGTAAAGAGCGTTCAGACGCTTATAAAAACATTTTGAAACAAATTGAT
GATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCACCA
AAAGATTTAGAAAAAGTATCATTCACACAATCACAGTATGAATTACCATTCAA
TGAAATGCAGTATAAATAA

(SEQ ID NO:446)

Figure 101

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAAGAAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCAATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGTATTAAGCCTTTACTAGCTAAAAAATGGGATGTGTCTGAAGATGGG
AAGACTTATACATTCCATTTAAGAGATGATGTGAAATTCCATGATGGAACAAC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGATGCGGTTCAACAAAATAAAA
AATTGCATTCTTGGTTAAAGATTTCAACATTAATTGACAATGTTAAAGTTAAA
GATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGGC
TGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAAGTCTAAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTTCCTATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACGGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTCAGTGACAAAACAGTCAGA
CAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTAG
ATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGACAT
TAATTTTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATTAT
TAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGTGATGTTTCGTCAAAAAGA
TGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAAA
AAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAAATGGGTATTAAGTT
AAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGTG
ATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAAGT
ACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCAT
TGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCCAA
AACGGTAAAGAGCGTTCAGACGCTTATAAAAAACATTTTGAAACAAATTGATG
ATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCGCCGA
AAGATTTAGAAAAAGTATCATTCACACAATCACAGTATGAATTACCATTCAAT
GAAATGCAGTATAAATAA

(SEQ ID NO:447)

Figure 102

ATGAGAAAATACTAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCAATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGTATTAAGCCTTTACTAGCTAAAAAGTGGGATGTGTCTGAAGATGGG
AAGACATACACGTTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACGCC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGACGCAGTTCAAGAAAACAAA
AAATTGCATTCTTGTTAAAGATTTTCGACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTCCATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACAGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTGAGTGACAAAACAGTCAG
ACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTA
GATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGACA
TTAATTTGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
TTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAAG
ATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAA
AAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGT
TAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGT
GATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAAG
TACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCA
TTGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAATCCA
AAACGGTAAAGAGCGTTCAGACGCTTATAAAAACATTTTGAAACAAATTGAT
GATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCACCA
AAAGATTTAGAAAAAGTATCATTCACACAATCACAGTATGAATTACCATTCAA
TGAAATGCAGTATAAATAA

(SEQ ID NO:448)

Figure 103

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTACATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCAATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGTATTAAGCCTTTACTAGCTAAAAAGTGGGATGTGTCTGAAGATGGG
AAGACATACACGTTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACGCC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGACGCAGTTCAAGAAAACAAA
AAATTGCATTCTTGTTAAAGATTTTCGACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTCCATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACAGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTGAGTGACAAAACAGTCAG
ACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTA
GATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGACA
TTAATTTTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
TTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAAG
ATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAA
AAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGT
TAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGT
GATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAAG
TACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCA
TTGAGAACAAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCCA
AAACGGTAAAGAGCGTTCAGACGCTTATAAAAACATTTTGAAACAAATTGAT
GATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCACCA
AAAGATTTAGAAAAAGTATCATTCACACAATCACAGTATGAATTACCATTCAA
TGAAATGCAGTATAAATAA

(SEQ ID NO:449)

Figure 104

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCAATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGTATTAAGCCTTTACTAGCTAAAAAGTGGGATGTGTCTGAAGATGGG
AAGACATACACGTTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACGCC
ATTTGATGCTGACGCAGTTAAGAAAAATATTGACGCAGTTCAAGAAAACAAA
AAATTGCATTCTTGGTTAAAGATTTTCGACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACACAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAGTCTAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTCCATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACAGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTGAGTGACAAAACAGTCAG
ACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTA
GATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGACA
TTAATTTGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
TTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAAG
ATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAA
AAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGT
TAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGT
GATTATGACTTAATGTTCAACCAAACTTGGGGATTATTGTACGATCCACAAAG
TACTATTGCAGCATTTAAAGAGAAAAATGGTTATGAAAGTGCAACATCAGGC
ATTGAGAACAAGATAAAATATACAACAGCATTGATGACGCATTTAAAATCC
AAAACGGTAAAGAGCGTTCAGACGCTTATAAAAAACATTTTGAAACAAATTGA
TGATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCACC
AAAAGATTTAGAAAAAGTATCATTACACAATCACAGTATGAATTACCATTCA
ATGAAATGCAGTATAAATAA

(SEQ ID NO:450)

Figure 105

ATGAGAAAACCTAACTAAAATGAGTGCAATGTTACTTGCATCAGGGCTAATTTT
AACTGGTTGTGGCGGTAATAAAGGTTTAGAGGAGAAAAAAGAAAACAAGCAA
TTAACGTATACGACGGTTAAAGATATCGGTGATATGAATCCGCATGTTTACGG
TGGATCGATGTCTGCTGAAAGTATGATATACGAGCCGCTTGTACGTAACACGA
AAGATGGAATTAAGCCTTTACTAGCTAAAAAATGGGATGTGTCTGAAGATGG
GAAGACATATACATTCCATTTGAGAGATGACGTTAAATTCCATGATGGTACAC
CATTTGATGCTGACGCAGTTAAGAAAAATATTGATGCGGTTCAACAAAATAAA
AAATTGCATTCTTGGTTAAAGATTTCAACATTAATTGACAATGTTAAAGTTAA
AGATAAGTACACGGTTGAATTGAATTTGAAAGAAGCATATCAACCTGCATTGG
CTGAATTAGCGATGCCTCGTCCATATGTATTTGTGTCTCCAAAAGACTTTAAAA
ACGGTACAACAAAAGATGGCGTTAAAAAGTTTCGATGGTACTGGTCCATTTAAA
TTAGGTGAACATAAAAAAGATGAGTCTGCAGACTTTAACAAAAATGATCAAT
ACTGGGGCGAAAAAGTCTAAACTTAACAAAGTACAAGCAAAAGTAATGCCTGC
TGGTGAAACAGCATTTCCTATCAATGAAAAAAGGTGAAACGAACTTTGCCTTCA
CAGATGATAGAGGTACGGATAGCTTAGACAAAGACTCTTTAAAACAATTGAA
AGATACAGGTGACTATCAAGTTAAGCGTAGTCAACCTATGAATACGAAAATGT
TAGTTGTCAATTCTGGTAAAAAAGATAACGCTGTGAGTGACAAAACAGTCAG
ACAAGCGATTGGTCATATGGTAAACAGAGATAAAATTGCCAAAGAAATTTTA
GATGGTCAAGAAAAACCAGCAACTCAATTATTTGCGAAAAATGTAACAGATA
TTAATTTTCGATATGCCAACACGTAAGTATGACCTTAAAAAAGCAGAATCATT
TTAGATGAAGCTGGTTGGAAGAAAGGTAAAGACAGCGATGTTTCGTCAAAAAG
ATGGTAAAAACCTTGAAATGGCAATGTACTATGACAAAGGTTCTTCAAGTCAA
AAAGAACAAGCAGAATACTTACAAGCAGAATTTAAGAAAATGGGTATTAAGT
TAAACATCAATGGCGAAACATCAGATAAAATTGCTGAACGTCGTACTTCTGGT
GATTATGACTTAATGTTCAACCAAACCTTGGGGATTATTGTACGATCCACAAAG
TACTATTGCAGCATTTAAAGCGAAAAATGGTTATGAAAGTGCAACATCAGGCA
TTGAGAACAAAGATAAAAATATACAACAGCATTGATGACGCATTTAAAATCCA
AAACGGTAAAGAGCGTTCAGGCGCTTATAAAAACATTTTGAAACAAATTGAT
GATGAAGGTATCTTTATCCCTATTTACACGGTAGTATGACAGTTGTTGCGCCG
AAAGATTTAGAAAAAGTATCATTCACACAATCACAGTATGAATTACCATTCAA
TGAAATGCAGTATAAATAA

(SEQ ID NO:451)

55 / 112

Figure 106

TCATCTGATAGCAAAGATAAGGAAACAACCTTCAATTAAACATGCAATGGGTA
CAACTGAAATTAAAGGGAAACCAAAGCGTGTTGTTACGCTATATCAAGGTGCC
ACTGACGTCGCTGTATCTTTAGGTGTTAAACCTGTAGGTGCTGTAGAATCATG
GACACAAAAACCGAAATTCGAATACATAAAAAATGATTTAAAAGATACTAAG
ATTGTAGGTCAAGAACCTGCACCTAACTTAGAGGAAATCTCTAAATTAAAACC
GGACTTAATTGTCGCGTCAAAAGTTAGAAATGAAAAAGTTTACGATCAATTAT
CTAAAATCGCACCAACAGTTTCTACTGATACAGTTTCAAATTCAAAGATACA
ACTAAGTTAATGGGGAAAGCTTTAGGGAAAGAAAAAGAAGCTGAAGATTTAC
TTAAAAAGTACGATGATAAAGTAGCTGCATTCCAAAAAGATGCAAAAGCAAA
GTATAAAGATGCATGGCCATTGAAAGCTTCAGTTGTTAACTTCCGTGCTGATC
ATACAAGAATTTATGCTGGTGGATATGCTGGTGAAATCTTAAATGATTTAGGA
TTCAAACGTAATAAAGACTTACAAAAACAAGTTGATAATGGTAAAGATATTAT
CCAATTACATCTAAAGAAAGTATTCCATTAATGAACGCTGATCATATTTTTGT
AGTAAATCAGATCCAAATGCGAAAGATGCTGCATTAGTTAAAAAGACTGAA
AGCGAATGGACTTCAAGTAAAGAGTGGAAAAATTTAGACGCAGTTAAAAACA
ACCAAGTATCTGATGATTTAGATGAAATCACTTGGAAGTTAGCTGGCGGATAT
AAATCATCATTAAACTTATTGACGATTTATATGAAAAGTTAAATATTGAAAA
ACAATCAAAATAA

(SEQ ID NO:452)

Figure 107

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCG
CCTAACTTAGAGGAAATCTCTAAATTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTCAAATTCAAAGATACAATAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG

56 / 112

GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAATTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGA AAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGA ACTTAGCTGGCGGATATAAATCATCATTAAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA
(SEQ ID NO:453)

Figure 108

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAACAATCATCTGATAACAAAGATAAGG
AAACA ACTTCAATTAAACATGCAATGGGTACA ACTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACA ACTAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAATTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGA AAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGA ACTTAGCTGGCGGATATAAATCTTCATTAAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA
(SEQ ID NO:454)

Figure 109

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAGCAATCATCTGATAGCAAAGATAAGG

57 / 112

AAACAACCTTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGCGTTGTTACACTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAAGAAATCTCTAAATTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAACCTAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAACCTTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGA AAAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATT
AGATGAAATCACTTGGAACCTTAGCTGGCGGATATAAATCATCATTAAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA
(SEQ ID NO:455)

Figure 110

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAACCTAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAACCTTACATCCAAAGAAA

58 / 112

GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAATGGAAAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGAACCTAGCTGGCGGATATAAATCATCATTAAAATTGA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA
(SEQ ID NO:456)

Figure 111

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTA AAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAATAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATAACAAGATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAACCTTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGA AAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGAACCTAGCTGGCGGATATAAATCTTCATTAAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA
(SEQ ID NO:457)

Figure 112

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG

59 / 112

GTGTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAATAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAATTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGA AAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGA ACTTAGCTGGCGGATATAAATCTTCATTAAAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA

(SEQ ID NO:458)

Figure 113

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACA ACTTCAATTAAACATGCAATGGGTACA ACTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAATAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAATTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA

60 / 112

AAGAGTGGAAAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGAACCTTAGCTGGCGGATATAAATCTTCATTA AAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA

(SEQ ID NO:459)

Figure 114

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAACCTAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAACCTTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGAAAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGAACCTTAGCTGGCGGATATAAATCTTCATTA AAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA

(SEQ ID NO:460)

Figure 115

ATGAATAAAGTAATTAAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTTCAATTAAACATGCAATGGGTACAACCTGAAATTAAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA

61 / 112

CCTAACTTAGAGGAAATCTCTAAATTTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAACCTAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAACCTACATCTAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAGTGGAATAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT
AGATGAAATCACTTGGAACCTAGCTGGCGGATATAAATCTTCATTAAACTTA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAATAA
(SEQ ID NO:461)

Figure 116

ATGAATAAAGTAATTTAAATGCTTGTTGTTACGCTTGCTTTCCTACTTGTTTTA
GCAGGATGTAGTGGGAATTCAAATAAACAATCATCTGATAACAAAGATAAGG
AAACAACCTCAATTAAACATGCAATGGGTACAACCTGAAATTAAGGGAAACC
AAAGCGTGTTGTTACGCTATATCAAGGTGCCACTGACGTCGCTGTATCTTTAG
GTGTTAAACCTGTAGGTGCTGTAGAATCATGGACACAAAAACCGAAATTCGA
ATACATAAAAAATGATTTAAAAGATACTAAGATTGTAGGTCAAGAACCTGCA
CCTAACTTAGAGGAAATCTCTAAATTTAAAACCGGACTTAATTGTCGCGTCAAA
AGTTAGAAATGAAAAAGTTTACGATCAATTATCTAAAATCGCACCAACAGTTT
CTACTGATACAGTTTTCAAATTCAAAGATACAACCTAAGTTAATGGGGAAAGCT
TTAGGGAAAGAAAAAGAAGCTGAAGATTTACTTAAAAAGTACGATGATAAAG
TAGCTGCATTCCAAAAAGATGCAAAAGCAAAGTATAAAGATGCATGGCCATT
GAAAGCTTCAGTTGTAACTTCCGTGCTGATCATACAAGAATTTATGCTGGTG
GATATGCTGGTGAAATCTTAAATGATTTAGGATTCAAACGTAATAAAGACTTA
CAAAAACAAGTTGATAATGGTAAAGATATTATCCAACCTACATCCAAAGAAA
GCATTCCATTAATGAACGCTGATCATATTTTTGTAGTAAAATCAGATCCAAAT
GCGAAAGATGCTGCATTAGTTAAAAAGACTGAAAGCGAATGGACTTCAAGTA
AAGAATGGAAAAATTTAGACGCAGTTAAAAACAACCAAGTATCTGATGATTT

62 / 112

AGATGAAATCACTTGGAAGTTAGCTGGCGGATATAAATCATCATTAAAATTGA
TTGACGATTTATATGAAAAGTTAAATATTGAAAAACAATCAAAATAA
(SEQ ID NO:462)

Figure 117

AAAGAATCATCAACTAAAGATACTATTTTCGGTAAAAGATGAAAATGGTACAG
TAAAAGTACCTAAAGATGCAAAACGTATCGTTGTATTAGAGTACTCATTTGCA
GATGCATTAGCAGCATTAGACGTTAAACCAGTTGGTATTGCTGATGATGGTAA
GAAAAAACGTATCATTAACCAGTTAGAGAAAAAATTGGGGATTATACTTCTG
TAGGTACACGTAAACAGCCAACTTAGAGGAAATTAGTAAATTAAAACCGGA
TTAATTATCGCTGATAGCAGTAGACATAAAGGTATTAATAAAGAATTAAACA
AAATTGCACCAACATTATCATTAAGAGTTTTGATGGAGACTACAAACAAAAC
ATTAATTCGTTCAAAACAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCG
AAAAGCGTCTTGCTGAACATGATAAATTAATCAAAAAGTATAAAGATGAAAT
TAAGTTTGATAGAAATCAAAAAGTGCTTCCAGCAGTTGTTGCTAAAGCTGGTT
TATTAGCACATCCAACTATTCATATGTTGGACAATTTTTAAACGAACTTGGAT
TTAAAAATGCATTAAGTGATGATGTAACAAAAGGTTTAAGTAAATACTTGAAA
GGACCTTACTTACAATTAGATACTGAACATTTAGCTGACTTAAATCCTGAACG
CATGATTATTATGACAGATAATGCTAAAAAAGATTCTGCTGAATTCAAGAAGT
TACAAGAAGATCCAACTTGGA AAAAGTTGAACGCAGTTAAAAATAATCGCGT
GGATATTGTTGACCGTGATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGA
AGAAATGGCTAAAGAACTTGTTGAATTATCAAAAAAAGAACAAAAG
(SEQ ID NO:463)

Figure 118

ATGAAGTTATACATATTAAGGAGTGGAACGATGAGAGGTCTAAAACTTTTAG
TATATTGGGATTAATAGTTGCCTTATTTTTAGTTGCAGCTTGTGGTAATACGGA
TAATTCAAGTAAAAAAGAATCATCAACTAAAGATACTATTTTCGGTAAAAGATG
AAAATGGCACAGTAAAAGTACCTAAAGATGCAAAACGTATCGTTGTATTAGA
GTACTCATTTGCAGATGCATTAGCAGCATTAGACGTTAAACCAGTTGGTATTG
CTGATGATGGTAAGAAAAAACGTATCATTAACCAGTTAGAGAAAAAATTGG
GGATTATACTTCTGTAGGTACACGTAAACAGCCAACTTAGAGGAAATTAGTA
AATTAAAACCGGATTTAATTATCGCTGATAGCAGTAGACATAAAGGTATTAAT
AAAGAATTAAACAAAATTGCACCAACATTATCATTAAGAGTTTTGATGGAGA

63 / 112

CTACAAACAAAACATTAATTCGTTCAAAACAATTGCTAAAGCTTTAAATAAAG
AAAAAGAAGGCGAAAAGCGTCTTGCTGAACATGATAAATTAATCAAAAAGTA
TAAAGATGAAATTAAGTTTGATAGAAATCAAAAAGTGCTTCCAGCAGTTGTTG
CTAAAGCTGGTTTATTAGCACATCCAAACTATTCATATGTTGGACAATTTTTAA
ACGAACTTGGATTTAAAAATGCATTAAGTGATGATGTAACAAAAGGTTTAAGT
AAATACTTGAAAGGACCTTACTTACAATTAGATACTGAACATTTAGCTGACTT
AAATCCTGAACGCATGATTATTATGACAGATAATGCTAAAAAAGATTCTGCTG
AATTCAAGAAGTTACAAGAAGATCCAACCTTGAAAAAGTTGAATGCAGTTAA
AAATAATCGCGTGGATATTGTTGACCGTGATGTTTGGGCAAGATCTCGTGGCT
TAATTTCTTCTGAAGAAATGGCTAAAGAACTTGTTGAATTATCAAAAAAAGAA
CAAAAGTAA
(SEQ ID NO:464)

Figure 119

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCGAGATGCATTAGCAGCATT
AGACGTTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAGGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAATATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC
AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAACTAGGATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACA
GATCATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAA
CATGGAAAAAGTTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA
(SEQ ID NO:465)

Figure 120

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTATTTTAA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATTAA
AACCAGTTAGAGAAAAAATTCGGGATTATACTTCTGTAGGTACACGTAAACAG
CCAAACTTAGAGGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATAG
CAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATTA
TCATTAAAGAGTTTTGATGGTGACTACAAGCAAAATATCGATGCTTTCAAGAC
AATTGCGAAAGCTTTAGATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGAA
CATGATAAATTAATCAATAAGTATAAAGATGAAATTAATTTGATAGAAATCA
AAAAGTGCTTCCAGCAGTTGTTGCGAAAGCTGGTTTATTAGCACATCCAACT
ATTCGTATGTTGGACAATTTTTAAACGAACTTGGATTAAAAATGCATTAAGT
GATGATGTAACAAAAGGTTTAAGTAAATACTTGAAAGGACCTTACTTACAATT
AGATACTGAACATTTAGCTGACTTAAATCCTGAACGCATGATTATTATGACAG
ATAATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAAC
ATGGAAAAAATTGAATGCAGTTAAAAATAATCGTGTTGATATTGTTGACCGTG
ATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGAA
CTTGTTGAATTATCAAAAAAAGAACAAAAGTAA

(SEQ ID NO:466)

Figure 121

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTAAATCCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATTAA
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAAGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAACATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA

65 / 112

GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC
AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAACTAGGATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACT
GATCACGCTAAAAAAGATTCTGCTGAATTCAAGAAATTACAAGAAGATGCAA
CATGGAAAAAATTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA
(SEQ ID NO:467)

Figure 122

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAAGAAATTAGTAAATTAAAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAATATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC
AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAACTAGGATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACA
GATCATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAA
CATGGAAAAAGTTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA
(SEQ ID NO:468)

Figure 123

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAGGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAATATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC
AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTAAACGAAGTATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACA
GATCATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAA
CATGGAAAAAGTTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA

(SEQ ID NO:469)

Figure 124

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAAGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAATATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC

67 / 112

AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAACTAGGATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACA
GATCATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAA
CATGGAAAAAGTTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA

(SEQ ID NO:470)

Figure 125

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAAGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAATATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAATTTGATAGAAATC
AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAACTAGGATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACA
GATCATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAA
CATGGAAAAAGTTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA

(SEQ ID NO:471)

Figure 126

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTAAACCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAAGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAATATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC
AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAAGTAGGATTTAAAAATGCATTAAG
TGACGATGTAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACA
GATCATGCTAAAAAAGATTCTGCTGAATTCAAGAAGTTACAAGAAGATGCAA
CATGGAAAAAGTTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA

(SEQ ID NO:472)

Figure 127

ATGAGAGGTCTAAAACTTTTAGTATATTGGGATTAATAGTTGCCTTACTTTTA
GTTGCAGCTTGTGGTAATACGGATAATTCAAGTAAAAAAGAATCATCAACTAA
AGATACTATTTTCGGTAAAAGATGAAAATGGTACAGTAAAAGTACCTAAAGAT
GCAAAACGTATCGTTGTATTAGAGTACTCATTTCAGATGCATTAGCAGCATT
AGACGTAAATCCAGTTGGTATTGCTGATGATGGTAAGAAAAAACGTATCATT
AACCAGTTAGAGAAAAAATTGGGGATTATACTTCTGTAGGTACACGTAAACA
GCCAACTTAGAAGAAATTAGTAAATTAACCGGATTTAATTATCGCTGATA
GCAGTAGACATAAAGGTATTAATAAAGAATTAAACAAAATTGCACCAACATT
ATCATTAAAGAGTTTTGATGGAGACTACAAACAAAACATTAATTCGTTCAAAA
CAATTGCTAAAGCTTTAAATAAAGAAAAAGAAGGCGAAAAACGTCTTGCTGA
GCATGATAAATTAATCAATAAGTATAAAGATGAAATTAAATTTGATAGAAATC

69 / 112

AAAAAGTGCTTCCAGCAGTAGTTGCTAAAGCTGGTTTATTAGCACATCCAAAC
TATTCATATGTTGGACAATTTTTAAACGAACTAGGATTTAAAAATGCATTAAG
TGACGATGTAAACAAAAGGTTTAAAGTAAATATTTGAAAGGACCTTACTTACAAT
TAGACACTGAACATTTAGCTGATTTAAATCCAGAGCGTATGATCATTATGACT
GATCACGCTAAAAAAGATTCTGCTGAATTCAAGAAATTACAAGAAGATGCAA
CATGGAAAAAATTGAATGCAGTTAAAAATAATCGCGTGGATATTGTTGACCGT
GATGTTTGGGCAAGATCTCGTGGCTTAATTTCTTCTGAAGAAATGGCTAAAGA
ACTTGTTGAATTATCAAAAAAAGAACAAAAGTAA

(SEQ ID NO:473)

Figure 128

ACTGAAGAGAAAACCTGAAATGACGACAATAAAAGATGAATTAGGAACTGAAA
AAATTAAGAAAAATCCTAAACGTGTTGTTGTATTAGAATATAGTTTTGCTGAT
TATTTAGCAGCATTAGATATGAAACCTGTTGGTATTGCAGATGATGGCAGCAG
TAAAAATATAACAAAGTCAGTAAGAGATAAGATTGGGGCATATGAATCGGTT
GGATCTAGATCGCAACCGAATATGGAAGTGATAAGTAAATTAACCGGATT
TGATCATTGCAGATGTCAGCAGACATAAGAAAAATCAAATCAGAATTGAGCAA
AATTGCGCCGACAATTATGTTAGTCAGTGGTACGGGAGATTACAATGCAAATA
TTGAAGCATTTAACAGTCGCTAAAGCAGTTGGCAGAGAGAAAGAAGGCCGA
AAAGCGTCTGGAAAAGCATGATAAAATATTAGCGGAGATTAGAAAGAAAATT
GAACAGAGTACGTTAATAATCTGCATTTGCACTTGGTATCTCAAGAGCAGGTAT
GTTTATTAATAATGAAGATACATTTATGGGACAATTTTAAATTAATAATGGGTA
TTCAACCTGAAGTCACAAAAGACAAAACCTGCGCATGTGGGTGAACGCAAGGG
TGGCCCTTATATTTATTTAAATAATGAAGAGCTTGCCAATATCAATCCAAAAG
TTATGATTTTAGCCACTGACGGAAAAACGGACAAAAATAGAACGAAATTCATT
GATCCTGCAGTTTGGAATCATTAAAAGCTGTGAAAGATAACAAAGTTTATGA
CGTTGACCGAAATAAGTGGTTGAAATCAAGGGGTATTATCGCAAGTGAAAGT
ATGGCAGAAGATTTAGAAAAAATTGCAGAAAAAGCAAAATAA

(SEQ ID NO:474)

Figure 129

ATGGAAGTGATAAGTAAATTAACCGGATTTGATCATTGCAGATGTAAACAG
ACATAAGAAAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGTTA
GTTAGCGGTACGGGAGATTATAATGCAAATATTGAAGCATTTAACAGTTGC

70 / 112

TAAAGCAGTAGGCAAAGAGAAAGAAGGCGAGAAACGTCTGGAAAAGCATGA
TAAAATATTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCT
GCATTTGCATTTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATAC
ATTTATGGGACAATTCTTAATTAATAATGGGTATTCAACCTGAAGTCACAAAAG
ACAAAACCTATGCATGTTGGTGAACGCAAGGGTGGTCCTTATATTTATTTAAAT
AATGAAGAACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGG
AAAAACGGACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCA
TTAAAAGCTGTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTT
GAAATCAAGGGGTATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAA
ATTGCAGAAAAAGCAAAATAA

(SEQ ID NO:475)

Figure 130

GTGAATAGGAATATCGTTAAATTAGTTGTGTTTATGCTAATCTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAACCTGAAATGACGACAA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTAAACGTGTTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGCAGCACTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAACCGGATTTGATCATTGCAGATGTTAGCAGACATAAG
AAAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGTTAGTTAGCGG
TACGGGAGATTATAATGCAAATATTGAAGCATTTAACAGTCGCTAAAGCA
GTAGGCAAAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCATGATAAAATA
TTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTTGC
ATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTATGG
GACAATTCTTAATTAATAATGGGTATTCAACCTGAAGTCACAAAAGACAAAAC
ACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTTAAATAATGAAGA
ACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGGAAAAACGG
ACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAAGCT
GTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTTGAAATCAA
GGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAAATTGCAGA
AAAAGCAAAATAA

(SEQ ID NO:476)

Figure 131

GTGAATAGGAATATCGTTAAATTAGTTGTGTTTATGCTAATCTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAAGTAAATGACGACGA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCGAAACGTATTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGTAGTACTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAAAACCGGATTTGATTATTGCAGATGTCAGCAGACATAAG
AAAATCAAATCAGAATTGAGCAAAATTGCGCCGACAATCATGTTAGTTAGCG
GTACGGGAGATTACAATGCAAATATTGAAGCATTTAAAACAGTCGCTAAAGC
AGTTGGCAAAGAGAGAAAGAAGGCGAAAAGCGTCTGGAAAAGCATAATAAAAT
ATTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTT
GCATTTGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTAT
GGGACAATTCTTAATTAATAATGGGTATTCAACCTGAAGTCACAAAAGACAAA
ACTGCGCATGTTGGTGAACGCAAGGGTGGCCCTTATATTTATTTAAATAATGA
AGAGCTTGCCAATATCAATCCAAAAGTTATGATTTTAGCTACGAATGGAAAAA
CAGATAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAATA
GCTGTGAAAGATAATAAAGTATATGATGTTGATCGAAATAAGTGGTTGCAATC
AAGAGGTATTATGGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAAATTGCA
GAAAAAGCAAAATAA

(SEQ ID NO:477)

Figure 132

GTGAATAGGAATATCGTTAAACTAGTTGTGTTTATGCTAATCTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAAGTAAATGACGACAA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTAAACGTGTTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGCAGCAGTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGCAGACATAAG
AAAATCAAATCAGAATTGAGCAAAATTGCGCCGACAATTATGTTAGTCAGTGG
TACGGGAGATTACAATGCAAATATTGAAGCATTTAAAACAGTCGCTAAAGCA
GTTGGCAAAGAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCATGATAAAATAT
TAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTTGC

72 / 112

ATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTATGG
GACAATTCTTACTTAAAATGGGTATTCAACCTGAAGTCACAAAAGACAAAAC
ACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTTAAATAATGAAGA
ACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGGGAAAACGG
ACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAAAAGCT
GTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTTGAAATCAA
GGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAAATTGCAGA
AAAAGCAAAATAA

(SEQ ID NO:478)

Figure 133

GTGAATAGGAATATCGTTAAACTAGTTGTGTTTCATGCTAATTTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAACCTGAAATGACGACAA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTAAACGTGTTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGCAGCACTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGCAGACATAAG
AAAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGTTAGTTAGCGG
TACGGGAGATTATAATGCAAATATTGAAGCATTTAAAACAGTCGCTAAAGCA
GTAGGCAAAGAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCATGATAAAATA
TTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTTGC
ATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTATGG
GACAATTCTTAATTAAAATGGGTATTCAACCTGAAGTCACAAAAGACAAAAC
ACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTTAAATAATGAAGA
ACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGGAAAAACGG
ACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAAAAGCT
GTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTTGAAATCAA
GGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAAATTGCAGA
AAAAGCAAAATAA

(SEQ ID NO:479)

Figure 134

GTGAATAGGAATATCGTTAAATTAGTTGTGTTTATGCTAATCTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAAC TGAAATGACGACAA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTAAACGTGTTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGCAGCACTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGCAGACATAAG
AAAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGTTAGTTAGCGG
TACGGGAGATTATAATGCAAATATTGAAGCATT TAAAACAGTCGCTAAAGCA
GTAGGCAAAGAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCATGATAAAATA
TTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTTGC
ATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTATGG
GACAATTCTTAATTAATAATGGGTATTCAACCTGAAGTCACAAAAGACAAAAC T
ACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTTAAATAATGAAGA
ACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGGAAAAACGG
ACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAAGCT
GTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTTGAAATCAA
GGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAAATTGCAGA
AAAAGCAAAATAA

(SEQ ID NO:480)

Figure 135

ATGACGACAATAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTA
AACGTGTTGTTGTATTAGAATATAGTTTTTGCTGATTATTTAGCAGCATTAGATA
TGAAACCTGTTGGTATTGCAGATGATGGCAGCACTAAAAATATAACAAAGTCA
GTAAGAGATAAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGA
ATATGGAAGTGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGC
AGACATAAGAAAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGT
TAGTTAGCGGTACGGGAGATTATAATGCAAATATTGAAGCATT TAAAACAGTC
GCTAAAGCAGTAGGCAAAGAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCAT
GATAAAATATTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAAT
CTGCATTTGCATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGAT
ACATTTATGGGACAATTCTTAATTAATAATGGGTATTCAACCTGAAGTCACAAA

74 / 112

AGACAAAACACTACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTAA
ATAATGAAGAACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGAC
GGAAAAACGGACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAAT
CATTAAAAGCTGTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTG
GTTGAAATCAAGGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTAGAA
AAAATTGCAGAAAAAGCAAAATAA

(SEQ ID NO:481)

Figure 136

GTGAATAGGAATATCGTTAACTAGTTGTGTTTCATGCTAATTTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAAGTAAATGACGACAA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTAAACGTGTTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGCAGCACTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGCAGACATAAG
AAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGTTAGTTAGCGG
TACGGGAGATTATAATGCAAATATTGAAGCATTTAAAACAGTCGCTAAAGCA
GTAGGCAAAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCATGATAAAATA
TTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTTGC
ATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTATGG
GACAATTCTTAATTAATAATGGGTATTCAACCTGAAGTCACAAAAGACAAAAGT
ACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTTAAATAATGAAGA
ACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGGAAAAACGG
ACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAAAAGCT
GTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTTGAAATCAA
GGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTAGAAAAAATTGCAGA
AAAAGCAAAATAA

(SEQ ID NO:482)

Figure 137

ATGACGACAATAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTA
AACGTGTTGTTGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATA
TGAAACCTGTTGGTATTGCAGATGATGGCAGCACTAAAAATATAACAAAGTCA

75 / 112

GTAAGAGATAAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGA
ATATGGAAGTGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGC
AGACATAAGAAAAATCAAATCAGAATTGAGCAAAATTGCTCCGACAATCATGT
TAGTTAGCGGTACGGGAGATTATAATGCAAATATTGAAGCATTTAAAACAGTC
GCTAAAGCAGTAGGCAAAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCAT
GATAAAATATTAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAAT
CTGCATTTGCATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGAT
ACATTTATGGGACAATTCTTAATTAAAATGGGTATTCAACCTGAAGTCACAAA
AGACAAAACCTACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTAA
ATAATGAAGAACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGAC
GGAAAAACGGACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAAT
CATTAAAAGCTGTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTG
GTTGAAATCAAGGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAA
AAAATTGCAGAAAAAGCAAAATAA

(SEQ ID NO:483)

Figure 138

GTGAATAGGAATATCGTTAACTAGTTGTGTTTATGCTAATCTTAGTTGTAGCA
GTAGCGGGTTGTGGTCAAAAAGATACTGAAGAGAAAACCTGAAATGACGACAA
TAAAAGATGAATTAGGAACTGAAAAAATTAAGAAAAATCCTAAACGTGTTGT
TGTATTAGAATATAGTTTTGCTGATTATTTAGCAGCATTAGATATGAAACCTGT
TGGTATTGCAGATGATGGCAGCAGTAAAAATATAACAAAGTCAGTAAGAGAT
AAGATTGGGGCATATGAATCGGTTGGATCTAGACCGCAACCGAATATGGAAG
TGATAAGTAAATTAAAACCGGATTTGATCATTGCAGATGTTAGCAGACATAAG
AAAATCAAATCAGAATTGAGCAAAATTGCGCCGACAATTATGTTAGTCAGTGG
TACGGGAGATTACAATGCAAATATTGAAGCATTTAAAACAGTCGCTAAAGCA
GTTGGCAAAGAGAAAGAAGGCGAGAAGCGTCTGGAAAAGCATGATAAAATAT
TAGCGGAGATTAGAAAGAAAATTGAACAGAGTACGTTAAAATCTGCATTTGC
ATTCGGTATCTCAAGAGCAGGTATGTTTATTAATAATGAAGATACATTTATGG
GACAATTCTTACTTAAAATGGGTATTCAACCTGAAGTCACAAAAGACAAAACCT
ACGCATGTTGGTGAACGCAAGGGTGGTCCTTATATATATTTAAATAATGAAGA
ACTTGCCAATATCAATCCAAAAGTTATGATTTTAGCCACTGACGGGAAAACGG
ACAAAAATAGAACGAAATTCATTGATCCTGCAGTTTGGAAATCATTAAAAGCT
GTGAAAGATAACAAAGTTTATGACGTTGACCGAAATAAGTGGTTGAAATCAA

76 / 112

GGGGGATTATCGCAAGTGAAAGTATGGCAGAAGATTTAGAAAAAATTGCAGA
AAAAGCAAAATAA

(SEQ ID NO:484)

Figure 139

GGTAGCGACGATAATGGCTCGTCTAAATCGCCGTACCATAGAATTGTTTCGTT
AATGCCTAGTAATACTGAAATTTTATATGAATTAGGATTAGGTAAATACATAG
TTGGTGTTTCAACGGTTGATGATTATCCAAAAGATGTGAAAAAGGGTAAGAAA
CAATTTGATGCTTTGAATCTAAATAAAGAGGAACTTTTAAAGGCAAAGCCAGA
TCTAATTCTTGCGCATGAGTCGCAAAAGGCAACTGCAAATAAAGTATTGTCAT
CATTAGAGAAACAAGGCATCAAAGTAGTGTATGTTAAAGATGCACAATCAAT
TGATGAAACTTACAACACATTTAAGCAAATTGGGAAATTAACGCATCATGATA
AGCAGGCTGAACAACTTGTTGAGGAACTAAAGATAATATCGATAAAGTCAT
AGATTCAATTCCTGCTCATCATAAAAAATCAAAAGTATTTATTGAGGTTTCATC
AAAGCCTGAAATATATACAGCAGGGAAGCATAACATTTTCAATGATATGTTAG
AAAAATTAGAAGCCCAAATGTTTATAGTGACATTAATGGTTGGAACCCTGTA
ACGAAGGAAAGTATTATTA AAAAAGAACCCAGATATATTAATTTGACGGAAG
CTAAGACAAGATCAGATTATATGGATATCATCAAAAAAAGAGGTGGATTCAA
TAAAATTAATGCTGTCAAGAATACACGTATTGAAGTTGTAAATGGTGATGAAG
TGTCAAGACCAGGTCCACGTATTGATGAAGGATTAAAAGAATTAAGAGATGC
AATTTATAGAAAATAA

(SEQ ID NO:485)

Figure 140

GTGAAGAAATCGTTAATTGCTTTTATTTTGATTTTATGCTTGTTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCGATGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT
CCAAAAGATGTGAAAAAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAA
AAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAG
TAGTGTATATTAAGATGCACAATCAATTGATGAACTTACAACACATTTAAG
CAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAGG
AAACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAA

77 / 112

AAATCAAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGG
GAAGCATACATTTTTCAATGATATGTTAGAAAAATTAGAAGCCCCAAAATGTTT
ATAGTGACATTAATGGTTGGAACCCTGTAACGAAGGAAAGTATTATTA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTAGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTG
ATGAAGGATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:486)

Figure 141

GTGAAGAAATCGTTAATTGCTTTTATTTTGATTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCGATGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT
CCAAAAGATGTGAAAGAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAA
AAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAG
TAGTGTATGTTAAAGATGCACAATCAATTGATGAAACATACAACACATTTAAG
CAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAGG
AAACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAA
AAATCAAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGG
GAAGCATACATTTTTTAATGATATGTTAGAAAAATTAGAAGCCCCAAAATGTTT
ATAGTGACATTAATGGTTGGAACCCTGTAACGAAGGAAAGTATTATTA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGGGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTG
ATGAAGGATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:487)

Figure 142

GTGAAGAAATCGTTAATTGCTTTTATTTTGATTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCGACGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT

78 / 112

CCAAAAGATGTGAAAAAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAA
AAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAG
TAGTGTATGTTAAAGATGCACAATCAATTGATGAAACTTACAACACATTTAAG
CAAATTGGGAAATTAACGCATCATGAAAAGCAGGCTGAACAACTTGTTGAGG
AAACTAAGGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAA
AAATCAAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGG
GAAGCATACATTTTTCAATGATATGTTAGAAAAATTAGAAGCCCAAATGTTT
ATAGTGACATTAATGGTTGGAACCCTGTAACGAAGGAAAGTATTATTAATAA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCCGGTCCACGTATTG
ATGAAGGATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:488)

Figure 143

GTGAAGAAATCGTTAATTGCTTTTTATTTTGATTTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCAATGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT
CCAAAAGATGTGAAAGAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGAAAAGCCAGATCTAATTCTTGCGCATGAGTCGCA
AAAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAA
GTAGTGTATGTTAAAGATGCACAATCAATTGATGAAACATACAACACATTTAA
GCAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAG
GAAACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAA
AAAATCAAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAG
GAAAGCATACATTTTTCAATGATATGTTAGAAAAATTAGAAGCCCAAATGTT
TATAGTGACATTAATGGTTGGAACCCTGTAACGAAGGAAAGTATTATTAATAA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCCGGTCCACGTATTG
ATGAAGGATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:489)

Figure 144

GTGAAGAAATCGTTAATTGCTTTTATTTTGATTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCAATGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT
CCAAAAGATGTGAAAAAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAA
AAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAG
TAGTGTATGTTAAAGATGCACAATCAATTGATGAACTTACAACACATTTAAG
CAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAGG
AACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAA
AAATCAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGG
GAAGCATACATTTTTTAATGATATGTTAGAAAAATTAGAAGCCCAAAATGTGT
ATAGTGACATTAATGGTTGGAACCCTGTAAACGAAGGAAAGTATTATTA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTG
ATGAAGGATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA

(SEQ ID NO:490)

Figure 145

GTGAAGAAATCGTTAATTGCTTTTATTTTGATTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCGATGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT
CCAAAAGATGTGAAAGAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAA
AAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAG
TAGTGTATGTTAAAGATGCACAATCAATTGATGAAACATACAACACATTTAAG
CAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAGG
AACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAA
AAATCAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGG
GAAGCATACATTTTTTAATGATATGTTAGAAAAATTAGAAGCCCAAAATGTTT

80 / 112

ATAGTGACATTAATGGTTGGAACCCCTGTAACGAAGGAAAGTATTATTA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGGGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTG
ATGAAGGATTAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:491)

Figure 146

ATGAAAGATAATGATAAACAAGGTAGCAATGATAATGGCTCGTCTAAATCGC
CGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTATATGAAT
TAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTATCCAAAA
GATGTGAAAAAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATAAAGAGG
AACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAAAAGGCA
ACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAGTAGTGTA
TGTTAAAGATGCACAATCAATTGATGAACTTACAACACATTTAAGCAAATTG
GGAAATTAACGCATCATGATAAGCAGGCTGAACAACCTGTTGAGGAACTAA
AGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCAAAAAATCAA
AAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGGGAAGCAT
ACATTTTTTAATGATATGTTAGAAAAATTAGAAGCCCAAATGTGTATAGTGA
CATTAAATGGTTGGAACCCCTGTAACGAAGGAAAGTATTATTAAGAACCCA
GATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGGATATCAT
CAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATACACGTATT
GAAGTTGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTGATGAAG
GATTAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:492)

Figure 147

GTGAAGAAATCGTTAATTGCTTTTATTTTGATTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCAATGATAATGGCTCGTCTA
AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTAT
CCAAAAGATGTGAAAAAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAA
AAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAG

81 / 112

TAGTGTATGTTAAAGATGCACAATCAATTGATGAACTTACAACACATTTAAG
CAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAGG
AAACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAA
AAATCAAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGG
GAAGCATACATTTTTTAATGATATGTTAGAAAAATTAGAAGCCCCAAAATGTGT
ATAGTGACATTAATGGTTGGAACCCTGTAACGAAGGAAAGTATTATTAATAA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTG
ATGAAGGATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:493)

Figure 148

ATGAAAGATAATGATAAACAAGGTAGCAATGATAATGGCTCGTCTAAATCGC
CGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTATATGAAT
TAGGATTAGGTAAATACATAGTTGGTGTTTCAACGGTTGATGATTATCCAAAA
GATGTGAAAAAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATAAAGAGG
AACTTTTAAAGGCCAAAGCCAGATCTAATTCTTGCGCATGAGTCGCAAAAGGCA
ACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAAGTAGTGTA
TGTTAAAGATGCACAATCAATTGATGAACTTACAACACATTTAAGCAAATTG
GGAAATTAACGCATCATGATAAGCAGGCTGAACAACTTGTTGAGGAAACTAA
AGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAAAAAATCAA
AAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAGGGAAGCAT
ACATTTTTTAATGATATGTTAGAAAAATTAGAAGCCCCAAAATGTGTATAGTGA
CATTAAATGGTTGGAACCCTGTAACGAAGGAAAGTATTATTAATAAAGAACCCA
GATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGGATATCAT
CAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATACACGTATT
GAAGTTGTAAATGGTGATGAAGTATCAAGACCAGGTCCACGTATTGATGAAG
GATTAAAAGAATTAAGAGATGCAATTTATAGAAAATAA
(SEQ ID NO:494)

Figure 149

GTGAAGAAATCGTTAATTGCTTTTTATTTTGATTTTATGCTTGTCTGAGTGGC
TGTGGTATGAAAGATAATGATAAACAAGGTAGCAATGATAATGGCTCGTCTA

82 / 112

AATCGCCGTACCATAGAATTGTTTCGTTAATGCCTAGTAATACTGAAATTTTAT
ATGAATTAGGATTAGGTAAATACATAGTTGGTGTTC AACGGTTGATGATTAT
CCAAAAGATGTGAAAGAGGGTAAGAAACAATTTGATGCTTTGAATCTAAATA
AAGAGGAACTTTTAAAGGAAAAGCCAGATCTAATTCTTGCGCATGAGTCGCA
AAAGGCAACTGCTAATAAAGTATTGTCATCATTAGAGAAACAAGGCATCAAA
GTAGTGTATGTTAAAGATGCACAATCAATTGATGAAACATACAACACATTTAA
GCAAATTGGGAAATTAACGCATCATGATAAGCAGGCTGAACAACCTTGTTGAG
GAACTAAAGATAATATCGATAAAGTCATAGATTCAATTCCTGCTCATCATAA
AAAATCAAAAGTATTTATTGAGGTTTCATCAAAGCCTGAAATATATACAGCAG
GAAAGCATACATTTTTCAATGATATGTTAGAAAAATTAGAAGCCCCAAAATGTT
TATAGTGACATTAATGGTTGGAACCCTGTAAACGAAGGAAAGTATTATTA AAAA
GAACCCAGATATATTAATTTTCGACGGAAGCTAAGACAAGATCAGATTATATGG
ATATCATCAAAAAAAGAGGTGGATTCAATAAAATTAATGCTGTCAAGAATAC
ACGTATTGAAGTTGTAAATGGTGATGAAGTATCAAGACCCGGTCCACGTATTG
ATGAAGGATTA AAAAGAATTAAGAGATGCAATTTATAGAAAATAA

(SEQ ID NO:495)

Figure 150

AGTGATAAGTCAAATGGCAA ACTAAAAGTAGTAACGACGAATTCAATTTTATA
TGATATGGCTAAAAATGTTGGTGGAGACAACGTCGATATTCATAGTATTGTAC
CTGTTGGTCAAGATCCTCATGAATATGAAGTTAAACCTAAAGATATTA AAAAAG
TTAACTGACGCTGACGTTATTTTATACAACGGATTAAATTTAGAGACTGGTAA
CGGTTGGTTTGAAAAAGCCTTAGAACAGGCTGGTAAATCATTAAAAGATAAA
AAAGTTATCGCAGTATCAAAAGATGTAAACCTATCTATTTAAACGGTGAAGA
AGGCAACAAAGATAAACAAGATCCACACGCATGGTTAAGTTTAGATAACGGT
ATTAAATACGTAAAAACAATTCAACAAACATTTATCGATAACGACAAAAAAC
ATAAAGCAGATTATGAAAAGCAAGGTAACAAATACATTGCTCAATTGGAAAA
ATTAAATAATGACAGTAAAGACAAATTTAATGACATTCCAAAAGAACAACGT
GCCATGATTACAAGTGAAGGTGCCTTCAAGTACTTCTCAAAACAATACGGTAT
TACACCAGGTTATATTTGGGAAATTAACACTGAAAAACAAGGTACACCAGAA
CAAATGAGACAAGCTATTGAGTTTGTTAAAAAGCACAAATTA AAACACTTATT
AGTAGAAACAAGTGTTGATAAGAAAGCAATGGAAAGTTTATCTGAAGAAACG
AAGAAAGATATCTTTGGTGAAGGTGTACACAGATTCAATCGGTAAAGAAGGCA

83 / 112

CTAAAGGTGACTCTTACTACAAAATGATGAAATCAAATATTGAAACTGTACAC
GGAAGCATGAAATAA

(SEQ ID NO:496)

Figure 151

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTTAAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTTAAAAGATAAAAAAGTTATCGCAGTATCAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCAGAACAATGAGACAAGCTATTGAGTTTGTTAAAAAGCA
CAAATTTAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA

(SEQ ID NO:497)

Figure 152

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTTAAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTTAAAAGATAAAAAAGTTATCGCAGTATCAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT

84 / 112

GGTTAAGTTTAGATAATGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTTAAAAAGCA
CAAATTAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA
(SEQ ID NO:498)

Figure 153

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTTTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAACATAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAGATAAAAAAGTTATCGCAGTATCAAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAACGACAGTAAAGACAAATTTAATGA
CATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACT
TCTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAA
AAACAAGGTACACCAGAACAAATGAGACAAGCTATTGAGTTTGTTAAAAAGC
ACAAATTAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGA
AAGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATT
CAATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATC
AAATATTGAAACTGTACACGGAAGCATGAAATAA
(SEQ ID NO:499)

Figure 154

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTTAAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAAGATAAAAAAGTTATCGCAGTATCAAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTAAAAAGCA
CAAATTTAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA

(SEQ ID NO:500)

Figure 155

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTTAAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAAGATAAAAAAGTTATCGCAGTATCAAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT

86 / 112

CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTTAAAAAGCA
CAAATTAAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA

(SEQ ID NO:501)

Figure 156

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAGATAAAAAAGTTATCGCAGTATCAAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAATGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTTAAAAAGCA
CAAATTAAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA

(SEQ ID NO:502)

Figure 157

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT

87 / 112

AAACCTAAAGATATTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAAGATAAAAAAGTTATCGCAGTATCAAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTTAAAAAGCA
CAAATTA AACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAACTGTACACGGAAGCATGAAATAA
(SEQ ID NO:503)

Figure 158

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTA AAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAAGATAAAAAAGTTATCGCAGTATCAAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTTAAAAAGCA
CAAATTA AACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC

88 / 112

AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA
(SEQ ID NO:504)

Figure 159

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTTAAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAAGATAAAAAAGTTATCGCAGTATCAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT
GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAGTAAAGACAA
ATTTAATGACATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCT
TCAAGTACTTCTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATT
AACACTGAAAAACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTG
TTAAAAAGCACAAATTTAAACACTTATTAGTAGAAACAAGTGTGATAAGAA
AGCAATGGAAAGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTG
TACACAGATTCAATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAAT
GATGAAATCAAATATTGAAACTGTACACGGAAGCATGAAATAA
(SEQ ID NO:505)

Figure 160

ATGAAAAAATTAGTACCTTTATTATTAGCCTTATTACTTCTAGTTGCTGCATGT
GGTACTGGTGGTAAACAAAGCAGTGATAAGTCAAATGGCAAATTTAAAAGTAG
TAACGACGAATTCAATTTTATATGATATGGCTAAAAATGTTGGTGGAGACAAC
GTCGATATTCATAGTATTGTACCTGTTGGTCAAGATCCTCATGAATATGAAGTT
AAACCTAAAGATATTTAAAAAGTTAACTGACGCTGACGTTATTTTATACAACGG
ATTAAATTTAGAGACTGGTAACGGTTGGTTTGAAAAAGCCTTAGAACAGGCTG
GTAAATCATTAAAAGATAAAAAAGTTATCGCAGTATCAAAGATGTTAAACCT
ATCTATTTAAACGGTGAAGAAGGCAACAAAGATAAACAAGATCCACACGCAT

89 / 112

GGTTAAGTTTAGATAACGGTATTAAATACGTAAAAACAATTCAACAAACATTT
ATCGATAACGACAAAAACATAAAGCAGATTATGAAAAGCAAGGTAACAAAT
ACATTGCTCAATTGGAAAAATTAAATAATGACAGTAAAGACAAATTTAATGAC
ATTCCAAAAGAACAACGTGCCATGATTACAAGTGAAGGTGCCTTCAAGTACTT
CTCAAAACAATACGGTATTACACCAGGTTATATTTGGGAAATTAACACTGAAA
AACAAGGTACACCTGAACAAATGAGACAAGCTATTGAGTTTGTAAAAAGCA
CAAATTAAACACTTATTAGTAGAAACAAGTGTTGATAAGAAAGCAATGGAA
AGTTTATCTGAAGAAACGAAGAAAGATATCTTTGGTGAAGTGTACACAGATTC
AATCGGTAAAGAAGGCACTAAAGGTGACTCTTACTACAAAATGATGAAATCA
AATATTGAAACTGTACACGGAAGCATGAAATAA

(SEQ ID NO:506)

Figure 161 A




Figure 161 B





Figure 161 C



91 / 112

Figure 162

MKKTVLVLYLVLAVMFLLAACGNNSDKEQSKSETKGSKETVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPNFLESFKDD
KYTNIGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKAKIVYVGADE
KNLIGSMKQNTENIGKIYDKEVKAKELNKDLDNKIASMKDKTKNFNKTVMYLLV
NEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN EYVNKENPDVIL
AMDRGQAVSGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYFAAGSTTTTI
KQIEELDKVVK

(SEQ ID NO:543)

Figure 163

MKKTVLVLYLVLAVMFLLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN EY
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK

(SEQ ID NO:544)

Figure 164

MKKTVLVLYLVVAVMFLLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLISSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKKF
NKSVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKNVSN SNHGQNVSN EYI
NKENPDVILAMDRGQAVSGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYF
AAGSTTTTIKQIDELEKVVK

(SEQ ID NO:545)

Figure 165

MKKTVLVLYLVLAVMFLLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA

92 / 112

KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK
(SEQ ID NO:546)

Figure 166

MKKTVLYLVLA VMFLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK
(SEQ ID NO:547)

Figure 167

MKKTVLYLVLA VMFLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK
(SEQ ID NO:548)

Figure 168

MKKTVLYLVLA VMFLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKAIKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK
(SEQ ID NO:549)

Figure 169

MKKTVLYLVLA VMFLLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN EY
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKA IKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK

(SEQ ID NO:550)

Figure 170

MKKTVLYLVLA VMFLLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN EY
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKA IKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK

(SEQ ID NO:551)

Figure 171

MKKTVLYLVLA VMFLLAACGNNSDKEQSKSETKGSKDTVKIENNYKMRGEKKD
GSDAKKVKETVEVPKNPKNAVVL DYGALDVMKEMGLSDKVKALPKGEGGKSLP
NFLESFKDDKYTNVGNLKEVNFDKIAATKPEVIFISGRTANQKNLDEFKKAAPKA
KIVYVGADEKNLIGSMKQNTENIGKIYDKEDKAKELNKDLDNKIASMKDKTKNF
NKTVMYLLVNEGELSTFGPKGRFGGLVYDTLGFNAVDKKVSNSNHGQNVSN EY
VNKENPDVILAMDRGQAISGKSTAKQALNNPVLKNVKA IKEDKVYNLDPKLWYF
AAGSTTTTIKQIEELDKVVK

(SEQ ID NO:552)

Figure 172

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTV DIPKDPKRIAVVA
PTYAGGLKKL GANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPD LII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKDWE

94 / 112

ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKA

(SEQ ID NO:553)

Figure 173

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKA

(SEQ ID NO:554)

Figure 174

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDIEKVAKEKPDII
YSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWEE
TTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQPE
QQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKA

(SEQ ID NO:555)

Figure 175

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNLKATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKA

(SEQ ID NO:556)

Figure 176

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTIKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:557)

Figure 177

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTIKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:558)

Figure 178

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTIKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:559)

Figure 179

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTIKIGDGDVEKVAKEKPDLII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKAWKKDWE
ETTAKDGKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:560)

Figure 180

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKD GKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:561)

Figure 181

MKKLLLPLIIMLLVLAACGNQGEKNNKAETKSYKMDDGKTVDIPKDPKRIAVVA
PTYAGGLKKLGANIVAVNQVDQSKVLKDKFKGVTKIGDGDVEKVAKEKPDII
VYSTDKDIKKYQKVAPT VVVDYNKHKYLEQQEMLGKIVGKEDKVKA WKKDWE
ETTAKD GKEIKKAIGQDATVSLFDEFDKKLYTYGDNWGRGGEVLYQAFGLKMQP
EQQKLTAKAGWAEVKQEEIEKYAGDYIVSTSEGKPTPGYESTNMWKNL KATKEG
HIVKVDAGTYWYNDPYTLDFMRKDLKEKLIKAAK
(SEQ ID NO:562)

Figure 182

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAAACTAAAGGTTCA
AAAGAAACGGTTAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAGAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGACAAAGTAAAAGCATTACCGAATTTCTTAGAATCATTTAAAGAT
GATAAATATACAAACATTGGTAATTTAAAAGAAGTGAATTTTGATAAAATTGC
TGCGACGAAACCCGAAGTAATCTTTATCTCTGGACGTACAGCTAATCAAAAGA
ATTTAGATGAATTCAAAAAAGCTGCACCTAAAGCGAAAATTGTTTATGTTGGT
GCAGATGAAAAGAACTTAATTGGTTCAATGAAACAAAACACTGAAAATATCG
GTAAAATCTACGACAAAGAAGTCAAAGCTAAAGAGTTAAATAAAGATTTAGA
TAATAAAATTGCTTCAATGAAAGATAAAACGAAAACTTCAATAAAACTGTTA
TGTATTTACTAGTTAACGAAGGTGAATTATCAACATTTGGACCTAAAGGTCGT
TTTGGTGGATTAGTTTACGATACATTAGGATTCAATGCAGTTGATAAAAAAGT
AAGTAATAGTAATCATGGACAAAATGTTTCTAACGAATATGTAAATAAAGAA

97 / 112

AATCCAGATGTTATTTTAGCGATGGATAGAGGTCAAGCGGTAAGTGGTAAATC
AACTGCGAAACAAGCATTAAATAATCCTGTATTAAAAAATGTTAAAGCAATTA
AAGAAGACAAAGTTTATAATTTAGATCCTAAATTATGGTACTTTGCAGCTGGA
TCAACTACAACACTACAATTAAACAAATTGAGGAACTTGATAAAGTTGTAAAATA
A

(SEQ ID NO:563)

Figure 183

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAAGTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGTAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAACTTCAATAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACACTACAATTAAACAAA
TTGAGGAACTTGATAAAGTTGTAAAATAA

(SEQ ID NO:564)

Figure 184

ATGAAGAAAACAGTCTTATATTTAGTAGTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAACAACACTATAAAATGCGTGGCGAGAAAAAAG

98 / 112

ATGGTAGCGATGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAGAACCC
TAAAAATGCAGTTGTATTAGATTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGACAAAGTAAAAGCATTACCTAAAGGTGAAGGCGGCAAGTCATT
ACCGAATTTCTTAGAGTCATTTAAAGATGACAAGTATACAAATGTTGGTAACT
TGAAAGAAGTTAATTTTCGATAAAATTGCAGCAACAAAACCGGAAGTAATCTTT
ATCTCTGGCCGTACAGCAAATCAAAAGAATTTAGATGAATTTAAAAAAGCTGC
ACCAAAAGCAAAAATTGTTTACGTTGGTGCAGATGAAAAGAATTTAATTAGTT
CAATGAAACAAAACACTGAAAATATTGGTAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAATAAGAAGTTCAACAAATCAGTTATGTATTTATTAGTTAATGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTCGGTGGATTAGTTTACGATACATT
AGGTTTCAACGCGGTTGATAAAAACGTAAGCAATAGTAACCACGGACAAAAT
GTCTCTAACGAATACATTAATAAAGAAAATCCAGATGTTATTTTAGCAATGGA
TAGAGGACAGGCTGTAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTTTATAATTTAGA
CCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACATAAATAAATAA
TTGATGAACCTTGAAAAAGTTGTAAAATAA

(SEQ ID NO:565)

Figure 185

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAACTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGTAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAAACTTCAATAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT

99 / 112

AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACATAATTAACAAA
TTGAGGAACCTTGATAAAGTTGTAAAATAA

(SEQ ID NO:566)

Figure 186

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAACTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGAAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAACTTCAATAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACATAATTAACAAA
TTGAGGAACCTTGATAAAGTTGTAAAATAA

(SEQ ID NO:567)

Figure 187

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT

100 / 112

AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAACTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGTAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAAACTTCAATAAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACATAATTAACAAA
TTGAGGAACTTGATAAAGTTGTAAAATAA

(SEQ ID NO:568)

Figure 188

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAACTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGAAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAAACTTCAATAAAACTGTTATGTATTTACTAGTTAACGAAGGTGA

101 / 112

ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACCTACAATTAAACAAA
TTGAGGAACTTGATAAAGTTGTAAAATAA

(SEQ ID NO:569)

Figure 189

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAAGCTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGAAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAAACTTCAATAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACCTACAATTAAACAAA
TTGAGGAACTTGATAAAGTTGTAAAATAA

(SEQ ID NO:570)

102 / 112

Figure 190

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAACTTAATTGGTT
CAATGAAACAAAACACTGAAAATATCGGAAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAAACTTCAATAAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACACTACAATTAAACAAA
TTGAGGAACTTGATAAAGTTGTAAAATAA

(SEQ ID NO:571)

Figure 191

ATGAAGAAAACAGTCTTATATTTAGTATTAGCAGTAATGTTTTTATTAGCGGC
ATGCGGTAACAATTCTGATAAAGAACAATCAAAATCAGAACTAAAGGTTCT
AAAGATACAGTAAAAATTGAAAATAACTATAAAATGCGTGGCGAGAAAAAAG
ATGGTAGTGACGCTAAAAAAGTTAAAGAACTGTTGAAGTACCAAAAAATCC
TAAAAATGCAGTTGTGTTAGACTATGGCGCATTAGATGTAATGAAAGAAATGG
GCTTATCAGATAAAGTAAAAGCATTACCTAAAGGGGAAGGCGGTAAGTCATT
ACCGAATTTCTTAGAATCATTTAAAGATGATAAATATACAAACGTTGGTAATT
TAAAAGAAGTGAATTTTGATAAAATTGCTGCGACGAAACCCGAAGTAATCTTT
ATCTCTGGACGTACAGCTAATCAAAAGAATTTAGATGAATTCAAAAAAGCTGC
ACCTAAAGCGAAAATTGTTTATGTTGGTGCAGATGAAAAGAACTTAATTGGTT

103 / 112

CAATGAAACAAAACACTGAAAATATCGGTAAAATTTACGATAAAGAAGATAA
AGCTAAAGAATTAAATAAAGATTTAGATAACAAAATTGCTTCAATGAAAGAT
AAAACGAAAAACTTCAATAAAACTGTTATGTATTTACTAGTTAACGAAGGTGA
ATTATCAACATTTGGACCTAAAGGTCGTTTTTGGTGGATTAGTTTACGATACATT
AGGATTCAATGCAGTTGATAAAAAAGTAAGTAATAGCAATCATGGACAAAAT
GTTTCTAACGAATATGTTAATAAAGAAAATCCAGATGTTATTTTAGCGATGGA
TAGAGGTCAAGCGATAAGTGGTAAATCAACTGCGAAACAAGCATTAAATAAT
CCTGTATTAAAAAATGTTAAAGCAATTAAAGAAGACAAAGTATATAATTTAGA
TCCTAAATTATGGTACTTTGCAGCTGGATCAACTACAACATAAATAACAAA
TTGAGGAACTTGATAAAGTTGTAAAATAA
(SEQ ID NO:572)

Figure 192

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAGATAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAGGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGACATTAAAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACACTGCTAAAGACGGTAAAGAAATTAAGCAATTGG
ACAAGATGCAACAGTGTCGTTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTACTATATCAAGCATTGTTTGA
AAAATGCAACCAGAACAACAAAAGTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATATGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAGAAAAATTAA
TTAAAGCTGCAAAATAA
(SEQ ID NO:573)

Figure 193

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAAAGATAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTAAAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTGGTTG
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATATGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:574)

Figure 194

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAAACATTGTAGCTG
TAAATCAACAAGTTGATCAAAGCAAAGTGTTAAAAGATAAATTTAAAGGTGTT
ACTAAAATTGGTGATGGTGATATAGAAAAAGTTGCTAAAGAAAAGCCAGATT
TAATTATTGTATACTCTACTGACAAAGACATTAAAAAGTATCAAAAAGTAGCA
CCAACAGTAGTTGTTGACTATAATAAGCATAAATACTTAGAACAACAAGAAAT
GTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAAGCTTGGAAGAAAGAT
TGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGGAC
AAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTTAC
GGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTGTTAAA

105 / 112

AATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAGTG
AAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAGTG
AAGGTAAACCTACACCAGGATACGAATCAACAAACATGTGGAAGAATTTGAA
AGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTACA
ACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAATT
AAAGCTGCAAAATAA

(SEQ ID NO:575)

Figure 195

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAGATAAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTAATAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTTGGTTTA
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATATGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTA
TTAAAGCTGCAAAATAA

(SEQ ID NO:576)

Figure 196

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG

106 / 112

TAAATCAACAAGTCGATCAAAGCAAAGTATTA AAAAGATAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTA AAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTTG
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATACGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:577)

Figure 197

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTA AAAAGATAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTA AAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAAGCCTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTTG
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATATGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA

107 / 112

CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:578)

Figure 198

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAGATAAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTAAGAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTTGGTTTG
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATACGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:579)

Figure 199

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAGATAAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTAAGAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA

108 / 112

ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTGTTG
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATACGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:580)

Figure 200

ATGAAAAAACTATTATTACCATTAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAGGATAAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTAATAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTGTTG
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATACGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:581)

Figure 201

ATGAAAAAACTATTATTACCATTAAATAATTATGTTATTAGTGTTAGCTGCGTGT
GGGAACCAAGGTGAAAAAAATAACAAAGCTGAAACTAAATCTTATAAAATGG
ACGATGGCAAAACGGTAGATATTCGAAAGACCCTAAACGCATTGCAGTAGT
TGCGCCAACATATGCTGGTGGACTTAAAAAATTAGGTGCAAACATTGTAGCTG
TAAATCAACAAGTCGATCAAAGCAAAGTATTAAAAGATAAATTTAAAGGTGT
TACAAAAATTGGTGATGGCGATGTAGAAAAAGTTGCTAAAGAAAAGCCAGAT
TTAATTATTGTATACTCTACTGACAAAGATATTAAAAAATATCAAAAAGTAGC
ACCAACAGTAGTTGTTGACTATAATAAGCATAAATATTTAGAACAACAAGAA
ATGTTAGGGAAAATTGTTGGTAAAGAAGATAAAAGTAAAAGCTTGGAAGAAAG
ATTGGGAAGAAACAACCTGCTAAAGACGGTAAAGAAATTAAAAAAGCAATTGG
ACAAGATGCAACAGTGTCATTGTTTGATGAATTTGATAAAAAATTATACACTT
ACGGCGATAACTGGGGTCGTGGTGGAGAAGTATTATATCAAGCATTGTTTA
AAAATGCAACCAGAACAACAAAAGTTAACTGCAAAAGCAGGTTGGGCTGAAG
TGAAACAAGAAGAAATTGAAAAATATGCTGGTGATTACATTGTGAGTACAAG
TGAAGGTAAACCTACACCAGGATATGAATCAACAAACATGTGGAAGAATTTG
AAAGCTACTAAAGAAGGACATATTGTTAAAGTTGATGCTGGTACATACTGGTA
CAACGATCCTTATACATTAGATTTTCATGCGTAAAGATTTAAAAGAAAAATTAA
TTAAAGCTGCAAAATAA

(SEQ ID NO:582)

Figure 202

Western Blot: Convalescent Mouse Serum




Figure 203

Western Blot: Healthy Human Serum




Figure 204





Figure 205






Figure 206

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/28326

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 39/085; A61K 39/38; C07K 14/00 (2010.01)

USPC - 424/243.1; 424/184.1; 530/350

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC: 424/243.1; 424/184.1; 530/350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC: 424/243.1; 424/184.1; 530/350; 424/190.1; 435/6 (text search)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Electronic data bases searched: PubWEST (EPAB, JPAB, UPST, PGPB); Google Scholar: Gram positive bacteria, Staphylococcus spp. or aureus, media, iron chelator, prevention or treatment Staphylococcal infections, ATCC 19636, ATCC 25904. GenCore 6.3 SEQ ID NOs: 353, 397, 399

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ----- Y	US 2006/0115490 A1 (MASIGNANI et al.) 1 June 2006 (01.06.2006) para [0006], [0024]-[0025], [0148], [0153], [0159], [0209]-[0210], claim 1, SEQ ID NOs: 2544, 4190	1, 4-7, 17-19, 25, 31-33, 54 ----- 3, 10-11, 16, 23-24, 30
Y	US 2008/0200650 A1 (EMERY et al.) 21 August 2008 (21.08.2008) para [0009]; SEQ ID NO: 418	3, 10-11, 16, 23-24, 30

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 August 2010 (23.08.2010)

Date of mailing of the international search report

30 AUG 2010

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/28326

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 55
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I+: claims 1, 3-8, 10-13, 16-19, 23-33, 54, drawn to a composition comprising: an isolated polypeptide having at least 92% sequence similarity to the amino acid sequence of SEQ ID NO:397, wherein if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-26 of SEQ ID NO:399, and a method of using said composition. The first invention encompasses a composition comprising a second peptide of SEQ ID NO: 353. Due to the number of sequences in this application, additional inventions of Group I+ will be defined as necessary depending on Applicant's ultimate payment of additional fees. Additional sequences will be searched if applicant pays for each additional sequence or shows that the sequences share a special technical feature, i.e. a common structure or feature that defines a contribution over the prior art. Note that each additional sequence to be searched must be specified by the Applicant in the response to this invitation and must either (1) have an additional invention fee paid or -----continued on Separate Sheet-----

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Claims 1, 3-7, 10-11, 16-19, 23-25, 30-33, 54, restricted to SEQ ID NO:353

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/28326

***** Supplemental Box *****

Continuation of Box III (Lack of Unity of Invention):

(2) have a showing that the sequences share a common structure or feature that defines a contribution over the prior art. The exact claims searched will depend on the specifically elected SEQ ID NO(s). Please NOTE that SEQ ID NOs: 354-363 are identical with respect to each other; SEQ ID NOs: 366, 368-369, 371-374 are identical with respect to each other; SEQ ID NOs: 376-377 and 379-385 are identical with respect to each other; SEQ ID NOs: 387-388 and 390-396 are identical with respect to each other; SEQ ID NOs: 420-427 and 429 are also identical with respect to each other.

[NOTE: Claims 8, 12-13 and 26-29 were excluded from group I, because they are drawn to a non-elected subject matter.]

Group II, claims 2-4, 9-11, 14-16, 20-33, 37-42, 45-54, 57, drawn to a composition comprising an isolated polypeptide having at least 98% sequence similarity to the amino acid sequence of SEQ ID NO:408, with the proviso that if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-5 of SEQ ID NO:415, and a method of using said composition, and an antibody against said polypeptide.

Group III, claims 34-36, 40-44, 47-53, 56, drawn to a composition comprising an antibody that specifically binds an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397, and a method of using said composition.

The inventions listed as Groups I+ through III do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

A technical feature of the inventions listed as Groups I+ through III is a specific polypeptide of the specific amino acid sequence recited therein. The inventions do not share a special technical feature, because 1) no significant structural similarities can readily be ascertained among the amino acid sequences SEQ ID NO:397 and SEQ ID NO: 408; 2) US 2006/0115490 A1 to Massignani; et al. discloses a *Staphylococcus aureus* polypeptide comprising a polypeptide (Massignani; et al., SEQ ID NO 4190) having 100% identity to the claimed SEQ ID NO:397 and, while said peptide includes amino acids 1-26 of SEQ ID NO:399, it also includes one amino acid substitution comparing with amino acids 1-26 of SEQ ID NO:399, specifically Methionine in the position 1 of SEQ ID NO:399 corresponds Valine 7 of SEQ ID NO 4190 of Massignani; et al. Without a shared special technical feature, the inventions lack unity with one another.

The inventions of Group I+ do not include the inventive concept of a composition comprising an antibody that specifically binds an isolated polypeptide having at least 80% sequence similarity to the amino acid sequence of SEQ ID NO:397, as required by Groups III.

The inventions of Group I+ and III share the technical feature of a polypeptide having at least 92% sequence similarity to the amino acid sequence of SEQ ID NO:397, with the proviso that if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-26 of SEQ ID NO:399, and a method of using said composition. However, this shared technical feature does not represent a contribution over prior art. Specifically, Massignani; et al. discloses said polypeptide as set forth above (Massignani; et al., SEQ ID NO 4190). As said polypeptide was known at the time of the invention, this cannot be considered a special technical feature that would otherwise unify the groups.

The inventions of Group I+ share the technical feature of a composition comprising a polypeptide having at least 92% sequence similarity to the amino acid sequence of SEQ ID NO:397, with the proviso that if the isolated polypeptide includes one or more additional amino acids at the amino terminal, the one or more additional amino acids include at least one amino acid deletion or at least one amino acid substitution compared to amino acids 1-26 of SEQ ID NO:399, and a method of using said composition. However, this shared technical feature does not represent a contribution over prior art. Specifically, Massignani; et al. discloses said polypeptide (Massignani; et al., SEQ ID NO 4190), as set forth in the immediately preceding paragraph, and further teaches a composition comprising said polypeptide (claims 11-16). As said composition would have been obvious to one of ordinary skill in the art at the time of the invention, this cannot be considered a special technical feature that would otherwise unify the groups.

Another technical feature of the inventions listed as Group I+ is a second polypeptide of the specific amino acid sequence recited therein. The inventions do not share a special technical feature, because 1) no significant structural similarities can readily be ascertained among the amino acid sequences (NOTE: SEQ ID NOs: 354-363 are the same; SEQ ID NOs: 366, 368-369, 371-374 are the same; SEQ ID NOs: 376-377 and 379-385 are the same; SEQ ID NOs: 387-388 and 390-396 are the same; SEQ ID NOs: 420-427 and 429 are also the same; 2) US 2008/0200650 A1 to EMERY, et al. discloses, in the context of polypeptides and immunizing compositions containing gram positive polypeptides and methods of use (title), discloses the claimed polypeptide of SEQ ID 354 (SEQ ID NO 418). Without a shared special technical feature, the inventions lack unity with one another.

Groups I+ through III therefore lack unity under PCT Rule 13 because they do not share a same or corresponding special technical feature.