




(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/01/31
(87) Date publication PCT/PCT Publication Date: 2019/08/15
(85) Entrée phase nationale/National Entry: 2020/07/31
(86) N° demande PCT/PCT Application No.: KR 2019/001404
(87) N° publication PCT/PCT Publication No.: 2019/156439
(30) Priorité/Priority: 2018/02/07 (KR10-2018-0015170)

(51) Cl.Int./Int.Cl. *C07D 231/38* (2006.01),
A61K 31/415 (2006.01), *A61P 35/00* (2006.01)
(71) Demandeurs/Applicants:
KOREA RESEARCH INSTITUTE OF CHEMICAL
TECHNOLOGY, KR;
INDUSTRY-ACADEMIC COOPERATION FOUNDATION,
YONSEI UNIVERSITY, KR
(72) Inventeurs/Inventors:
CHANG, SUNG YOUN, KR;
LEE, HYUK, KR;
KIM, KI YOUNG, KR;
KIM, BUM TAE, KR;
KIM, SUNG SOO, KR; ...

(54) Titre : COMPOSES DESTINES A L'INHIBITION DE TNIK ET UTILISATIONS MEDICALES ASSOCIEES
(54) Title: COMPOUNDS FOR INHIBITING TNIK AND MEDICAL USES THEREOF



(57) **Abrégé/Abstract:**

The present disclosure provides the compound having inhibitory property against TNIK having a specific chemical structure or its pharmaceutically acceptable salt. The present disclosure also provides a composition comprising the compound or its pharmaceutically acceptable salt. The present disclosure also provides a medical use of the compound, its salt or the composition comprising the compound or its pharmaceutically acceptable salt for treating or preventing cancer. The present disclosure also provides a method of treatment or prevention of cancer comprising administering the compound, its salt or the composition comprising the compound or its salt to a subject in need of such treatment or prevention.

(72) **Inventeurs(suite)/Inventors(continued):** KIM, SEONG HWAN, KR; LIM, HWAN JUNG, KR; HEO, JUNG NYOUNG, KR;
SHIN, SANG JOON, KR; PARK, SANG YOUN, KR

(74) **Agent:** INTEGRAL IP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

15 August 2019 (15.08.2019)



(10) International Publication Number

WO 2019/156439 A1

(51) International Patent Classification:

C07D 231/38 (2006.01) *A61P 35/00* (2006.01)*A61K 31/415* (2006.01)

(21) International Application Number:

PCT/KR2019/001404

(22) International Filing Date:

31 January 2019 (31.01.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10-2018-0015170 07 February 2018 (07.02.2018) KR

(71) Applicants: KOREA RESEARCH INSTITUTE

OF CHEMICAL TECHNOLOGY [KR/KR]; 141,

Gajeong-ro, Yuseong-gu, Daejeon 34114 (KR).

INDUSTRY-ACADEMIC COOPERATION FOUN-

DATION, YONSEI UNIVERSITY [KR/KR]; 50, Yon-


sei-ro, Seodaemun-gu, Seoul 03722 (KR).

(72) Inventors: **CHANG, Sung Youn**; 102-902, 63, Gajeong-ro, Yuseong-gu, Daejeon 34119 (KR). **LEE, Hyuk**; 203-1604, 448, Expo-ro, Yuseong-gu, Daejeon 34049 (KR). **KIM, Ki Young**; 203-502, 127, Dunsannam-ro, Seogu, Daejeon 35249 (KR). **KIM, Bum Tae**; 102-1502, 92, Gyeryong-ro, Yuseong-gu, Daejeon 34179 (KR). **KIM, Sung Soo**; 1-803, 212, Samseong-ro, Gangnam-gu, Seoul 06284 (KR). **KIM, Seong Hwan**; 101-801, 65, Gajeong-ro, Yuseong-gu, Daejeon 34120 (KR). **LIM, Hwan Jung**; 306-1601, 448, Expo-ro, Yuseong-gu, Daejeon 34049 (KR). **HEO, Jung Nyoung**; 103, 32-10, Yudeung-ro 669beon-gil, Seo-gu, Daejeon 35252 (KR). **SHIN, Sang Joon**; 101-2007, 23, Sinnmok-ro, Yangcheon-gu, Seoul 08017 (KR). **PARK, Sang Youn**; 204-401, 47, Sangdo-ro 55-gil, Dongjak-gu, Seoul 06978 (KR).

(74) Agent: **PHIL & ONZI INT INT'L PATENT & LAW FIRM**; 3F., 36, Seochojungang-ro, Seocho-gu, Seoul 06643 (KR).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,

(54) Title: COMPOUNDS FOR INHIBITING TNK AND MEDICAL USES THEREOF



(57) Abstract: The present disclosure provides the compound having inhibitory property against TNK having a specific chemical structure or its pharmaceutically acceptable salt. The present disclosure also provides a composition comprising the compound or its pharmaceutically acceptable salt. The present disclosure also provides a medical use of the compound, its salt or the composition comprising the compound or its pharmaceutically acceptable salt for treating or preventing cancer. The present disclosure also provides a method of treatment or prevention of cancer comprising administering the compound, its salt or the composition comprising the compound or its salt to a subject in need of such treatment or prevention.

[Continued on next page]



WO 2019/156439 A1

WO 2019/156439 A1

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
 CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
 DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
 HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
 KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
 OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
 SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

Description

Title of Invention: COMPOUNDS FOR INHIBITING TNIK AND MEDICAL USES THEREOF

Technical Field

- [1] This application claims priority to Patent Application No. 10-2018-0015170 filed in the Republic of Korea on February 7, 2018, the entire contents of which are incorporated herein by reference.
- [2] The present disclosure relates to a group of compounds having an activity inhibiting Traf2- and Nck-interacting kinase (TNIK). The present disclosure also relates to pharmaceutical compositions comprising the compound. The present disclosure relates to methods useful for treating specific diseases, including cancer or tumor, using the compound. That is, the present disclosure relates to medical-uses of those compounds according to the present disclosure for treating or preventing cancer or tumor.

Background Art

- [3] It is well known that inhibitors of Traf2- and Nck-interacting kinase (TNIK) are useful for treating cancer (e.g., U.S. Patent Application Publication no. 2010/0216795). More concretely, it is known that TNIK is It is well known that inhibitors of Traf2- and Nck-interacting kinase (TNIK) are useful for treating cancer (e.g., U.S. Patent Application Publication no. 2010/0216795). More concretely, it is known that TNIK is hyperactive in colorectal cancer, breast cancer, brain tumor, gastric cancer, liver cancer, ovarian cancer and so on (J.S. Boehm et al., Cell 129, 1065-1079, 2007). Specifically, TNIK plays an important role in the growth of colorectal cancer, and TNIK was reported to be a target that is able to control aberrant Wnt signaling in colorectal cancer (*Cancer Res* 2010;70:5024-5033). TNIK gene was over-expressed in 7% of gastric cancer patients' tissue samples, and TNIK was reported to be a new target for treating gastric cancer (Oncogenesis 2014, 3, e89). In addition, TNIK was reported to be associated with proliferation and differentiation of leukemia stem cells in chronic myelogenous leukemia (The journal of clinical investigation 2012 122 624). In addition to these cancers or tumors, hepatocellular carcinoma, desmoid tumor, medulloblastoma (pediatric brain tumor), Wilms tumor (pediatric kidney cancer), thyroid tumor and so no is related to aberrant Wnt signaling, and thus medicine for these diseases can be developed based on TNIK inhibition.
- [4] Therefore, drugs inhibiting TNIK block the pathway of TNIK signaling, thereby inhibiting proliferation, survival, and angiogenesis of cancers. Drugs inhibiting TNIK thus are expected to be useful as medicine for treating cancer (See WO2010-100431 and WO2009-030890).

- [5] Meanwhile, TNIK inhibition was reported to be useful in treating chronic obstructive pulmonary disease (COPD), lupus nephritis, diabetic nephropathy, focal segmental glomerulosclerosis, renal fibrosis, Pulmonary fibrosis, scleroderma and so on.
- [6] Thus, TNIK inhibitors are expected to be useful in treating or preventing various diseases including inflammatory diseases as well as cancers.

Disclosure of Invention

Technical Problem

- [7] Thus one object of the present disclosure is to provide a compound having activity inhibiting TNIK (Traf2- and Nck-interacting kinase), pharmaceutical compositions comprising the compound as an effective agent, and medical-uses thereof for treating or preventing cancers.
- [8] Another object of the present disclosure is to provide a method for treating or ameliorating cancer comprising administering to a subject in need of treatment, amelioration or prevention of cancer a compound inhibiting TNIK activity according to the present disclosure.
- [9] Yet another object of the present disclosure is to provide a compound exhibiting synergic effect with other anticancer drugs by co-administration, pharmaceutical compositions comprising the compound as an effective agent, and medical uses thereof for treating or preventing cancer.
- [10] Yet another object of the present disclosure is to provide a method for treating or ameliorating cancer comprising administering to a subject in need of treatment, amelioration or prevention of cancer a compound inhibiting TNIK activity and anti-cancer drug having other mechanism simultaneously or sequentially.


Solution to Problem

[11] Summary

- [12] To achieve the object, in one embodiment, there is provided a compound of Chemical Formula 1:

[13] [Chemical Formula 1]

[14]



- [15] or a pharmaceutically acceptable salt thereof,
- [16] in the Chemical Formula 1

- [17] Y is N or CH,
- [18] Z is N or C-V,
- [19] A is H, halogen, -OH, -CO₂-C₁₋₆ alkyl, -CO₂H, -CN, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -OR¹, -NH₂, -NHR², -substituted or unsubstituted piperazine, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,
- [20] B is H, -C₁₋₆ haloalkyl, C₁₋₆ alkyl, halogen, or C₁₋₆ alkoxy,
- [21] V is H, -CH₂OH, halogen, -CO₂H, -CO₂-C₁₋₆ alkyl, -OH, -NH₂, phenoxy, or -NHCO-C₁₋₆ alkyl,
- [22] X is H or F,
- [23] W is substituted or unsubstituted, aromatic ring, heteroaryl, or fused heteroaryl,
- [24] wherein,
- [25] R¹ is C₁₋₆ alkyl, benzyl, C₁₋₆ haloalkyl, or phenyl,
- [26] R² is C₁₋₆ alkyl, C₁₋₆ haloalkyl, -CH₂CH₂-morpholin, or phenyl,
- [27] R³ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, or substituted or unsubstituted phenyl,
- [28] R⁴ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.
- [29] In another embodiment, there is provided a pharmaceutical composition comprising a compound of Chemical Formula 1 or a pharmaceutically acceptable salt thereof, and a pharmaceutically-acceptable carrier or additive.
- [30] In various embodiments, the pharmaceutical composition further comprises one or more additional pharmaceutically active agents, preferably, irinotecan or pharmaceutically acceptable salt thereof.
- [31] In yet another embodiment, there is provided a method for treating a condition comprising administering to a subject a therapeutically effective amount of a compound of Chemical Formula 1 or a pharmaceutically acceptable salt thereof, wherein the condition to be treated includes, but is not limited to, cancer such as colorectal cancer, breast cancer, brain tumor, gastric cancer, liver cancer, ovarian cancer, lung cancer, gastrointestinal cancer, leukemia, and melanoma, neoplasia, or tumor. The compound of Chemical Formula 1 or a pharmaceutically acceptable salt thereof according to the present disclosure is also useful in preventing metastasis and recurrence of tumor by targeting cancer stem cells. That is, there is provided medical-uses of the compound of Chemical Formula 1 or a pharmaceutically acceptable salt thereof for treating or preventing the diseases mentioned above.
- [32] In various embodiments, the method comprises administering combination agents of the compound of the present disclosure or its salt and other pharmaceutically active compounds (preferably, irinotecan or pharmaceutically acceptable salt thereof). That is, there is provided a medical-use of combination medicine comprising the compound of the present disclosure or its pharmaceutically acceptable salt and other active agent

(preferably, irinotecan or pharmaceutically acceptable salt thereof) for treating or preventing the diseases above.

[33] The compounds, the pharmaceutical composition, and their medical use above are more fully described in the detailed description that follows.

[34]

[35] **Detailed Description**

[36] The following description is merely exemplary in nature and is not intended to limit the present disclosure, application, or uses.

[37] **Definitions**

[38] The generic terms used in the present disclosure are herein defined for clarity.

[39] This specification uses the terms "substituent", "radical", "group", "moiety", and "fragment" interchangeably.

[40] As used herein, the term "patient" means an animal, preferably a mammal such as a non-primate (e.g., cow, horse, sheep, pig, chicken, turkey, quail, cat, dog, mouse, rat, rabbit or guinea pig) or a primate (e.g., monkey and human), most preferably a human.

[41] As used herein, the term "alkyl" means a saturated straight chain or branched non-cyclic hydrocarbon, unless the context clearly dictates otherwise, having from 1 to 10 carbon atoms. "lower alkyl" means alkyl having from 1 to 4 carbon atoms. Representative saturated straight chain alkyls include -methyl, -ethyl, -n-propyl, -n-butyl, -n-pentyl, -n-hexyl, -n-heptyl, -n-octyl, -n-nonyl and -n-decyl; while saturated branched alkyls include -isopropyl, -sec-butyl, -isobutyl, -tert-butyl, -isopentyl, 2-methylbutyl, 3-methylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 2,3-dimethylbutyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylpentyl, 2,2-dimethylhexyl, 3,3-dimethylpentyl, 3,3-dimethylhexyl, 4,4-dimethylhexyl, 2-ethylpentyl, 3-ethylpentyl, 2-ethylhexyl, 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-ethylpentyl, 2-methyl-4-ethylpentyl, 2-methyl-2-ethylhexyl, 2-methyl-3-ethylhexyl, 2-methyl-4-ethylhexyl, 2,2-diethylpentyl, 3,3-diethylhexyl, 2,2-diethylhexyl, 3,3-diethylhexyl and the like.

[42] As used herein, if the term "C₁₋₆" is used, it means the number of carbon atoms is from 1 to 6. For example, C₁₋₆ alkyl means an alkyl which carbon number is any integer of from 1 to 6.

[43] The terms "halogen" and "halo" mean fluorine, chlorine, bromine or iodine.

[44] As used herein, the term "haloalkyl", "haloalkoxy", "haloalkenyl" or "haloalkynyl" means an alkyl, alkoxy, alkenyl or alkynyl group, respectively, wherein one or more hydrogen atoms are substituted with halogen atoms. For example, the haloalkyl includes -CF₃, -CHF₂, -CH₂F, -CBr₃, -CHBr₂, -CH₂Br, -CCl₃, -CHCl₂, -CH₂Cl, -CI₃, -CHI₂, -CH₂I, -CH₂CF₃, -CH₂CHF₂, -CH₂CH₂F, -CH₂CBr₃, -CH₂CHBr

- $_{2}$, $-\text{CH}_2-\text{CH}_2\text{Br}$, $-\text{CH}_2-\text{CCl}_3$, $-\text{CH}_2-\text{CHCl}_2$, $-\text{CH}_2-\text{CH}_2\text{Cl}$, $-\text{CH}_2-\text{Cl}_3$, $-\text{CH}_2-\text{CHI}_2$, $-\text{CH}_2-\text{CH}_2\text{I}$, and the like, wherein alkyl and halogen are as described above.
- [45] The term "alkanoyl" or "acyl" means an $-\text{C}(\text{O})\text{alkyl}$ group, wherein alkyl is defined above, including $-\text{C}(\text{O})\text{CH}_3$, $-\text{C}(\text{O})\text{CH}_2\text{CH}_3$, $-\text{C}(\text{O})(\text{CH}_2)_2\text{CH}_3$, $-\text{C}(\text{O})(\text{CH}_2)_3\text{CH}_3$, $-\text{C}(\text{O})(\text{CH}_2)_4\text{CH}_3$, $-\text{C}(\text{O})(\text{CH}_2)_5\text{CH}_3$, and the like.
- [46] The term "alkanoyloxy" or "acyloxy" means an $-\text{OC}(\text{O})\text{alkyl}$ group, wherein alkyl is defined above, including $-\text{OC}(\text{O})\text{CH}_3$, $-\text{OC}(\text{O})\text{CH}_2\text{CH}_3$, $-\text{OC}(\text{O})(\text{CH}_2)_2\text{CH}_3$, $-\text{OC}(\text{O})(\text{CH}_2)_3\text{CH}_3$, $-\text{OC}(\text{O})(\text{CH}_2)_4\text{CH}_3$, $-\text{OC}(\text{O})(\text{CH}_2)_5\text{CH}_3$, and the like.
- [47] The term "alkoxy" means an $-\text{O}(\text{alkyl})$ group, wherein alkyl is defined above, including $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{O}(\text{CH}_2)_2\text{CH}_3$, $-\text{O}(\text{CH}_2)_3\text{CH}_3$, $-\text{O}(\text{CH}_2)_4\text{CH}_3$, $-\text{O}(\text{CH}_2)_5\text{CH}_3$, and the like.
- [48] The term "lower alkoxy" means $-\text{O}(\text{lower alkyl})$, wherein lower alkyl is as described above.
- [49] The term "aryl" means a carbocyclic aromatic group containing from 5 to 10 ring atoms. Representative examples include, but are not limited to, phenyl, tolyl, xylyl, naphthyl, tetrahydronaphthyl, anthracenyl, fluorenyl, indenyl, and azulenyl. A carbocyclic aromatic group can be unsubstituted or optionally substituted.
- [50] The term "aryloxy" is $\text{RO}-$, where R is aryl as defined above. "arylthio" is $\text{RS}-$, where R is aryl as defined above.
- [51] The term "cycloalkyl" means a monocyclic or polycyclic saturated ring having carbon and hydrogen atoms and having no carbon-carbon multiple bonds. Examples of cycloalkyl groups include, but are not limited to, $(\text{C}_3-\text{C}_7)\text{cycloalkyl}$ groups, including cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. A cycloalkyl group can be unsubstituted or optionally substituted. In one embodiment, the cycloalkyl group is a monocyclic ring or bicyclic ring.
- [52] The term "mono-alkylamino" means $-\text{NH}(\text{alkyl})$, wherein alkyl is defined above, such as $-\text{NHCH}_3$, $-\text{NHCH}_2\text{CH}_3$, $-\text{NH}(\text{CH}_2)_2\text{CH}_3$, $-\text{NH}(\text{CH}_2)_3\text{CH}_3$, $-\text{NH}(\text{CH}_2)_4\text{CH}_3$, $-\text{NH}(\text{CH}_2)_5\text{CH}_3$, and the like.
- [53] The term "di-alkylamino" means $-\text{N}(\text{alkyl})(\text{alkyl})$, wherein each alkyl is independently an alkyl group as defined above, including $-\text{N}(\text{CH}_3)_2$, $-\text{N}(\text{CH}_2\text{CH}_3)_2$, $-\text{N}((\text{CH}_2)_2\text{CH}_3)_2$, $-\text{N}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$, and the like.
- [54] The term "alkylamino" means mono-alkylamino or di-alkylamino as defined above.
- [55] The term "carboxyl" and "carboxy" mean $-\text{COOH}$.
- [56] The term "aminoalkyl" means $-(\text{alkyl})-\text{NH}_2$, wherein alkyl is defined above, including $-\text{CH}_2-\text{NH}_2$, $-(\text{CH}_2)_2-\text{NH}_2$, $-(\text{CH}_2)_3-\text{NH}_2$, $-(\text{CH}_2)_4-\text{NH}_2$, $-(\text{CH}_2)_5-\text{NH}_2$ and the like.
- [57] The term "mono-alkylaminoalkyl" means $-(\text{alkyl})-\text{NH}(\text{alkyl})$, wherein each alkyl is independently an alkyl group defined above, including $-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{NHCH}_2$

CH₃, -CH₂-NH(CH₂)₂CH₃, -CH₂-NH(CH₂)₃CH₃, -CH₂-NH(CH₂)₄CH₃, -CH₂-NH(CH₂)₅CH₃, -(CH₂)₂-NH-CH₃, and the like.

- [58] The term "heteroaryl" means an aromatic heterocycle ring of 5- to 10 members and having at least one heteroatom selected from nitrogen, oxygen and sulfur, and containing at least 1 carbon atom, including both mono- and bicyclic ring systems. Representative heteroaryls are triazolyl, tetrazolyl, oxadiazolyl, pyridyl, furyl, benzofuranyl, thiophenyl, benzothiophenyl, quinolinyl, pyrrolyl, indolyl, oxazolyl, benzoxazolyl, imidazolyl, benzimidazolyl, thiazolyl, benzothiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, cinnolinyl, phthalazinyl, quinazolinyl, pyrimidyl, oxetanyl, azepinyl, piperazinyl, morpholinyl, dioxanyl, thietanyl and oxazolyl.
- [59] The term "heterocycle" means a 5- to 7-membered monocyclic, or 7- to 10-membered bicyclic, heterocyclic ring which is either saturated, unsaturated, and which contains from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulfur, and wherein the nitrogen and sulfur heteroatoms can be optionally oxidized, and the nitrogen heteroatom can be optionally quaternized, including bicyclic rings in which any of the above heterocycles are fused to a benzene ring. The heterocycle can be attached via any heteroatom or carbon atom. Heterocycles include heteroaryls as defined above. Representative heterocycles include morpholinyl, pyrrolidinyl, pyrrolidinyl, piperidinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranlyl, tetrahydropyranyl, tetrahydropyridinyl, tetrahydroprimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, and tetrahydrothiopyranyl.
- [60] The term "heterocycle fused to phenyl" means a heterocycle, wherein heterocycle is defined as above, that is attached to a phenyl ring at two adjacent carbon atoms of the phenyl ring.
- [61] The term "hydroxyalkyl" means alkyl, wherein alkyl is as defined above, having one or more hydrogen atoms replaced with hydroxy, including -CH₂OH, -CH₂CH₂OH, -(CH₂)₃CH₂OH, -(CH₂)₄CH₂OH, -(CH₂)₅CH₂OH, -CH(OH)-CH₃, -CH₂CH(OH)CH₃, and the like.
- [62] The term "sulfonyl" means -SO₂.
- [63] The term "sulfonylalkyl" means -SO₂-(alkyl), wherein alkyl is defined above, including -SO₂-CH₃, -SO₂-CH₂CH₃, -SO₂-(CH₂)₂CH₃, -SO₂-(CH₂)₃CH₃, -SO₂-(CH₂)₄CH₃, -SO₂-(CH₂)₅CH₃, and the like.
- [64] The term "sulfinylalkyl" means -SO-(alkyl), wherein alkyl is defined above, including -SO-CH₃, -SO-CH₂CH₃, -SO-(CH₂)₂CH₃, -SO-(CH₂)₃CH₃, -SO-(CH₂)₄CH₃, -SO-(CH₂)₅CH₃, and the like.
- [65] The term "thioalkyl" means -S-(alkyl), wherein alkyl is defined above, including -

S-CH₃, -S-CH₂CH₃, -S-(CH₂)₂CH₃, -S-(CH₂)₃CH₃, -S-(CH₂)₄CH₃, -S-(CH₂)₅CH₃, and the like.

[66] As used herein, the term "substituted" means any of the above groups (i.e., alkyl, aryl, heteroaryl, heterocycle or cycloalkyl) wherein at least one hydrogen atom of the moiety being substituted is replaced with a substituent. In one embodiment, each carbon atom of the group being substituted is substituted with no more than two substituents. In another embodiment, each carbon atom of the group being substituted is substituted with no more than one substituent. In the case of a keto substituent, two hydrogen atoms are replaced with an oxygen which is attached to the carbon via a double bond. Unless specifically defined, substituents include halogen, hydroxyl, (lower) alkyl, haloalkyl, mono- or di-alkylamino, aryl, heterocycle, -NO₂, -NR_aR_b, -NR_aC(=O)R_b, -NR_aC(=O)NR_aR_b, -NR_aC(=O)OR_b, -NR_aSO₂R_b, -OR_a, -CN, -C(=O)R_a, -C(=O)OR_a, -C(=O)NR_aR_b, -OC(=O)R_a, -OC(=O)OR_a, -OC(=O)NR_aR_b, -NR_aSO₂R_b, -PO₃R_a, -PO(OR_a)(OR_b), -SO₂R_a, -S(O)R_a, -SO(N)R_a (e.g., sulfoximine), -(R_a)S=NR_b (e.g., sulfilimine) and -SR_a, wherein R_a and R_b are the same or different and independently hydrogen, halogen, amino, alkyl, haloalkyl, aryl or heterocycle, or wherein R_a and R_b taken together with the nitrogen atom to which they are attached form a heterocycle. R_a and R_b may be in the plural based on atoms which those are attached to.

[67] As used herein, the term "pharmaceutically acceptable salt(s)" refers to a salt prepared from active compounds according to the present disclosure with relatively non-toxic acids or bases with active compounds, depending on the particular substituents of those compounds. When the compounds have a relatively acidic group, base-added salts can be obtained by contacting the neutral compounds with a sufficient amount of the desired base and a pure or inert solvent. Suitable pharmaceutically acceptable base addition salts include, but are not limited to sodium, potassium, calcium, aluminum, organic amino, magnesium salts and the like. When the compounds have a relatively basic group, acid-added salts can be obtained by contacting the neutral compounds with a sufficient amount of the desired acid and pure or inert solvent. Suitable pharmaceutically acceptable acid addition salts include salts derived from non-toxic organic acids including, but are not limited to, acetic acid, propionic acid, isobutyl acid, oxalic acid, maleic acid, malonic acid, benzoic acid, succinic acid, suberic acid, fumaric acid, mandelic acid, phthalic acid, benzenesulfonic acid, p-tolylsulfonic acid, citric acid, tartaric acid, methanesulfonic acid, and the like, and non-toxic inorganic acids including, but are not limited to, hydrochloric acid, hydrobromic acid, nitric acid, carbonic acid, monohydrogencarbonic acid, phosphoric acid, monohydrogenphosphoric acid, dihydrogenphosphoric acid, sulfuric acid, monohydrogen-sulfuric acid, hydrogen iodide, phosphorous acid and the like. Also it includes a salt of

amino acid such as arginate or its analogues, and it also includes analogues of organic acid such as glucuronic or galacturonic acid. Some specific compounds of this disclosure have both basic and acidic functionality for the conversion of compounds with a basic or acidic portion (addition) salts. Other examples of salts are disclosed in well-known literature on the art, for example, *Remington's Pharmaceutical Sciences*, 18th eds., Mack Publishing, Easton PA (1990) or *Remington : The Science and Practice of Pharmacy*, 19th eds, are disclosed in Mack Publishing, Easton PA (1995).

- [68] As used herein, the term "effective amount" includes that amount of a compound of this disclosure sufficient to destroy, modify, control or remove a primary, regional or metastatic cancer cell or tissue; delay or minimize the spread of cancer; or provide a therapeutic benefit in the treatment or management of cancer, a neoplastic disorder, or tumor. An "effective amount" also includes the amount of a compound of this disclosure sufficient to result in cancer or neoplastic cell death. An "effective amount" also includes the amount of a compound of this disclosure sufficient to inhibit or decrease TNIK activity either *in vitro* or *in vivo*.
- [69] As used herein, the term "prophylactically effective amount" refers to the amount of a compound sufficient to prevent the recurrence or spread of cancer or the occurrence of cancer in a patient, including but not limited to those predisposed to cancer or previously exposed to a carcinogen.
- [70] As used herein, the term "neoplastic" means an abnormal growth of a cell or tissue (e.g., a tumor) which may be benign or cancerous.
- [71] As used herein, the term "prevention" includes the prevention of the recurrence, spread or onset of cancer in a patient.
- [72] As used herein, the term "treatment" includes the eradication, removal, modification, or control of primary, regional, or metastatic cancer tissue; and the minimizing or delay of the spread of cancer.
- [73] As used herein, the phrase "Compound(s) of this/the Disclosure" includes any compound(s) of Chemical Formula 1, as well as clathrates, hydrates, solvates, or polymorphs thereof. And, even if the term "Compound(s) of the Disclosure" does not mention its pharmaceutically acceptable salt, the term includes salts thereof. In one embodiment, the compounds of this disclosure include stereo-chemically pure compounds, e.g., those substantially free (e.g., greater than 85% ee, greater than 90% ee, greater than 95% ee, greater than 97% ee, or greater than 99% ee) of other stereoisomers. That is, if the compounds of Chemical Formula 1 according to the present disclosure or salts thereof are tautomeric isomers and/or stereoisomers (e.g., geometrical isomers and conformational isomers), such isolated isomers and their mixtures also are included in the scope of this disclosure. If the compounds of the present disclosure or salts thereof have an asymmetric carbon in their structures, their

active optical isomers and their racemic mixtures also are included in the scope of this disclosure.

- [74] As used herein, the term "polymorph" refers to solid crystalline forms of a compound of this disclosure or complex thereof. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (e.g., to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which can affect bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both (e.g., tablets of one polymorph are more susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it.
- [75] As used herein, the term "solvate" means a compound or its salt according to this disclosure that further includes a stoichiometric or non-stoichiometric amount of a solvent bound by non-covalent intermolecular forces. Preferred solvents are volatile, non-toxic, and/or acceptable for administration to humans in trace amounts.
- [76] As used herein, the term "hydrate" means a compound or its salt according to this disclosure that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.
- [77] As used herein, the term "clathrate" means a compound or its salt in the form of a crystal lattice that contains spaces (e.g., channels) that have a guest molecule (e.g., a solvent or water) trapped within.
- [78] If any compound (prodrug) produces the compound or its salt of this disclosure after degrading *in vivo*, such compound is included in this disclosure. As used herein and unless otherwise indicated, the term "prodrug" means a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (*in vitro* or *in vivo*) to provide an active compound, particularly a compound of this disclosure. Examples of prodrugs include, but are not limited to, metabolites of a compound that include biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Preferably, prodrugs of compounds with carboxyl functional groups are the lower alkyl esters of the carboxylic acid. The carboxylate esters are conveniently formed by esterifying any of the carboxylic acid moieties present on the

molecule. Prodrugs can typically be prepared using well-known methods, such as those described by *Burger's Medicinal Chemistry and Drug Discovery* 6th ed. (Donald J. Abraham ed., 2001, Wiley) and *Design and Application of Prodrugs* (H. Bundgaard ed., 1985, Harwood Academic Publishers GmHh).

[79] As used herein, the term "purified" means that when isolated, the isolate is greater than 90% pure, in one embodiment greater than 95% pure, in another embodiment greater than 99% pure and in another embodiment greater than 99.9% pure.

[80] The term "hydrido" means a single -H atom (H) and may be used interchangeably with the symbol "H" or the term "hydrogen".

[81] If a substituent is described as being "optionally substituted," the substituent may be either (1) not substituted or (2) substituted. If a substitutable position is not substituted, the default substituent is a hydrido radical.

[82] As used herein, the singular forms "a" and "an" may include plural reference unless the context clearly dictates otherwise.

[83] The term "pharmaceutically-acceptable" means suitable for use in pharmaceutical preparations, generally considered as safe for such use, officially approved by a regulatory agency of a national or state government for such use, or being listed in South Korea or the U. S. Pharmacopoeia or other generally recognized pharmacopoeia for use in animals, and more particularly in humans.


[84]

[85] **Compounds of the present disclosure**

[86] There is provided a compound of Chemical Formula 1:

[87] [Chemical Formula 1]

[88]



[89] or a pharmaceutically acceptable salt thereof,

[90] in the Chemical Formula 1

[91] Y is N or CH,


[92] Z is N or C-V,

[93] A is H, halogen, -OH, -CO₂-C₁₋₆ alkyl, -CO₂H, -CN, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -OR¹, -NH₂, -NHR², -substituted or unsubstituted piperazine, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,


[94] B is H, -C₁₋₆ haloalkyl, C₁₋₆ alkyl, halogen, or C₁₋₆ alkoxy,

- [95] V is H, -CH₂OH, halogen, -CO₂H, -CO₂-C₁₋₆ alkyl, -OH, -NH₂, phenoxy, or -NHCO-C₁₋₆ alkyl,
- [96] X is H or F,
- [97] W is substituted or unsubstituted, aromatic ring, heteroaryl, or fused heteroaryl,
- [98] wherein,
- [99] R¹ is C₁₋₆ alkyl, benzyl, C₁₋₆ haloalkyl, or phenyl,
- [100] R² is C₁₋₆ alkyl, C₁₋₆ haloalkyl, -CH₂CH₂-morpholin, or phenyl,
- [101] R³ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, or substituted or unsubstituted phenyl,
- [102] R⁴ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.
- [103]
- [104] In another embodiment, in the Chemical Formula 1,
- [105] Y is N or CH,
- [106] Z is N or C-V,
- [107] A is H, halogen, -OH, -CO₂-C₁₋₃ alkyl, -CO₂H, -CN, -C₁₋₃ alkyl, -OR¹, -NH₂, -NHR², -substituted or unsubstituted piperazine, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,
- [108] B is H, -C₁₋₃ haloalkyl, C₁₋₃ alkyl, halogen, or C₁₋₃ alkoxy,
- [109] V is H, -CH₂OH, halogen, -CO₂H, -CO₂-C₁₋₃ alkyl, -OH, -NH₂, phenoxy, or -NHCO-C₁₋₃ alkyl,
- [110] X is H or F,
- [111] W is substituted or unsubstituted, phenyl, pyridyl, thiophene, thiazole, pyrrole, benzothienophene, indole, oxazole, pyrazole, imidazole, pyrimidine, benzopyrazole, benzothiazole, benzoxazole, benzoimidazole, or benzothiophene,
- [112] wherein,
- [113] R¹ is benzyl, C₁₋₃ haloalkyl, or phenyl,
- [114] R² is CF₃, C₁₋₃ alkyl, -CH₂CH₂-morpholin, or phenyl,
- [115] R³ is C₁₋₃ alkyl, or substituted or unsubstituted phenyl,
- [116] R⁴ is C₁₋₃ alkyl, CF₃, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.
- [117]
- [118] Preferably, in yet another embodiment, in the Chemical Formula 1,
- [119] Y is N or CH,
- [120] Z is C-V,
- [121] A is -OH, -CO₂-C₁₋₂ alkyl, -methyl, -OR¹, -NH₂, -NHR², -substituted or unsubstituted piperazine, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,
- [122] B is H, -C₁₋₃ haloalkyl, C₁₋₃ alkyl, halogen, or C₁₋₃ alkoxy,


- [123] V is H, -CH₂OH, F, -CO₂H, -CO₂-C₁₋₂ alkyl, -OH, -NH₂, phenoxy, or -NHCOCH₃,
- [124] X is H or F,
- [125] W is substituted or unsubstituted, phenyl, pyridyl, thiophene, thiazole, pyrrole, benzothiophene, or indole,
- [126] wherein,
- [127] R¹ is benzyl, CF₃, or phenyl,
- [128] R² is CF₃, C₁₋₃ alkyl, -CH₂CH₂-morpholin, or phenyl,
- [129] R³ is C₁₋₃ alkyl, or substituted or unsubstituted phenyl,
- [130] R⁴ is C₁₋₃ alkyl, CF₃, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.
- [131]
- [132] More preferably, in yet another embodiment, in the Chemical Formula 1,
- [133] Y is CH,
- [134] Z is C-V,
- [135] A is -OH, -NHR², -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,
- [136] B is H, -C₁₋₃ haloalkyl, C₁₋₃ alkyl, halogen, or C₁₋₃ alkoxy,
- [137] V is H, -CH₂OH, F, -OH, or -NHCOCH₃,
- [138] X is H or F,
- [139] W is
- [140]




[141]



- [142] wherein,
- [143] R² is CF₃, C₁₋₃ alkyl, -CH₂CH₂-morpholin, or phenyl,
- [144] R³ is C₁₋₃ alkyl, or substituted or unsubstituted phenyl,
- [145] R⁴ is C₁₋₃ alkyl, CF₃, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.
- [146]
- [147] Much more preferably, in yet another embodiment, in the Chemical Formula 1,
- [148] Y is CH,
- [149] Z is C-V,
- [150] A is -OH, -NHR², -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,
- [151] B is H, -C₁₋₃ haloalkyl, C₁₋₃ alkyl, halogen, or C₁₋₃ alkoxy,
- [152] V is H, F, -OH, or -NHCOCH₃,
- [153] X is H or F,
- [154] W is
- [155]



[156]



[157] wherein,

[158] R² is CF₃, or C₁₋₃ alkyl,

[159] R³ is C₁₋₃ alkyl,

[160] R⁴ is C₁₋₃ alkyl, CF₃, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.

[161]

[162] Much more preferably, in yet another embodiment, in the Chemical Formula 1,

[163] Y is CH,

[164] Z is C-V,

[165] A is -OH, -NHCF₃, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,


[166] B is C₁₋₃ alkyl, or halogen,

[167] V is H, or F,

[168] X is H or F,

[169] W is

[170]



[171] wherein,

[172] R^3 is C_{1-3} alkyl,

[173] R^4 is C_{1-3} alkyl, CF_3 , $-CH_2CH_2Cl$, $-CH_2CH_2NMe_2$, $-CH_2NMe_2$, or $-CH_2CH_2$ - morpholin.







[174]

[175] The inventors had synthesized and evaluated lots of compounds to find out compounds having good TNIK inhibition activity and high selectivity against TNIK, thereby having good inhibition effect against cancer cells and low side effects about normal cells. Finally, the compounds of the present disclosure are identified to be suitable for the object of the present disclosure.

[176] Non-limiting examples of the compounds of the present disclosure include the compounds of Table 1 below and pharmaceutically acceptable salts thereof.











[177]

[Table 1]










Compound No.	Structure	Name
1		N-(4-(benzyloxy)phenyl)-5-(4-methoxyphenyl)-1H-pyrazol-3-amine
2		4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)phenol
3		4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenol
4		4-((5-(o-tolyl)-1H-pyrazol-3-yl)amino)phenol
5		4-((5-(2,4-dimethylphenyl)-1H-pyrazol-3-yl)amino)phenol
6		N-(4-chloro-2-methylphenyl)-5-(4-methoxyphenyl)-1H-pyrazol-3-amine

[178]

7		N-(4-chloro-2-methylphenyl)-5-(2,4-dimethylphenyl)-1H-pyrazol-3-amine
8		5-(4-chlorophenyl)-N-(4-fluoro-2-methylphenyl)-1H-pyrazol-3-amine
9		N-(4-fluoro-2-methylphenyl)-5-(4-methoxyphenyl)-1H-pyrazol-3-amine
10		N-(4-fluoro-2-methylphenyl)-5-(o-tolyl)-1H-pyrazol-3-amine
11		ethyl 4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)benzoate
12		4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
13		N1-(5-(4-methoxyphenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine

14		N-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
15		4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)benzoic acid
16		N1-(5-(4-methoxyphenyl)-1H-pyrazol-3-yl)benzene-1,3-diamine
17		ethyl 3-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)benzoate
18		N-(3-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
19		3-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenol
20		3-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)benzoic acid
21		N1-(5-(4-nitrophenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine
22		N1-(5-(4-aminophenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine
23		3-methyl-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol

[180]

24		4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
25		4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol
26		5-(4-methoxyphenyl)-N-(3-phenoxyphenyl)-1H-pyrazol-3-amine
27		4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-methoxyphenol
28		4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-isopropoxyphenol
29		N1-(5-(4-chlorophenyl)-1H-pyrazol-3-yl)-N4-propylbenzene-1,4-diamine
30		N1-(5-(4-chlorophenyl)-1H-pyrazol-3-yl)-N4-phenylbenzene-1,4-diamine
31		3-methyl-4-((5-(4-phenoxyphenyl)-1H-pyrazol-3-yl)amino)phenol
32		4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol

33		4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzonitrile
34		4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methoxyphenol
35		4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-isopropoxyphenol
36		N1-(5-(4-aminophenyl)-1H-pyrazol-3-yl)-N4-propylbenzene-1,4-diamine
37		N1-(5-(4-aminophenyl)-1H-pyrazol-3-yl)-N4-phenylbenzene-1,4-diamine
38		4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)phenol
39		4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
40		4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
41		4-((5-(3-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol









42		4-((5-(4-fluorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
43		4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol
44		4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol
45		4-((5-([1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
46		4-((5-(4'-bromo-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
47		3-methyl-4-((5-(pyridin-3-yl)-1H-pyrazol-3-yl)amino)phenol
48		N-(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)phenyl)-4-fluorobenzenesulfonamide
49		3-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
50		N-(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide









51		N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
52		4-fluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)benzenesulfonamide
53		N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
54		N-(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
55		N-(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
56		4-((5-(4-aminophenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenol
57		N-(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-4-fluorobenzenesulfonamide
58		N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide

59		N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
60		4-((5-(4-chlorophenyl)-4-fluoro-1H-pyrazol-3-yl)amino)phenol
61		4-(4-fluoro-3-((4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)phenol
62		4-((4-fluoro-5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
63		4-(3-((4-bromo-2-methylphenyl)amino)-4-fluoro-1H-pyrazol-5-yl)phenol
64		4-(3-((4-bromo-2-(trifluoromethyl)phenyl)amino)-4-fluoro-1H-pyrazol-5-yl)phenol
65		4-(4-fluoro-3-((3-fluoro-4-methylphenyl)amino)-1H-pyrazol-5-yl)phenol
66		4-((5-(6-chloropyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol

[185]

67		3-fluoro-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol
68		2-fluoro-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol
69		4-fluoro-N-(4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenyl)benzenesulfonamide
70		N-(3-methyl-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
71		4-((4-fluoro-5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
72		4-fluoro-N-(3-methyl-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenyl)benzenesulfonamide
73		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
74		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
75		4-((5-([1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)phenol

76		4-((5-(4'-bromo-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)phenol
77		N-(4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
78		N-(4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
79		4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)phenol
80		4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
81		3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
82		4-(3-((4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)-2-methylphenol
83		N-(4-((5-(4-hydroxy-3-methylphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide

84		N-(3-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)acetamide
85		4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)-2-methylphenol
86		N-(4-((5-(4-hydroxy-3-methylphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
87		4-((5-(3-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
88		2-fluoro-4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol
89		4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol
90		methyl 4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzoate
91		4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzoic acid

92		2-fluoro-4-(3-((4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)phenol
93		4-((5-(6-chloropyridin-3-yl)-1H-pyrazol-3-yl)amino)phenol
94		2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
95		4-((5-(6-(benzyloxy)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
96		(3-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-4-methylphenyl)methanol
97		4-((5-(6-(benzyloxy)pyridin-3-yl)-1H-pyrazol-3-yl)amino)phenol
98		4-((5-(5-methylthiophen-2-yl)-1H-pyrazol-3-yl)amino)phenol
99		4-((5-(thiophen-2-yl)-1H-pyrazol-3-yl)amino)phenol









100		3-methyl-4-((5-(5-methylthiophen-2-yl)-1H-pyrazol-3-yl)amino)phenol
101		3-methyl-4-((5-(thiophen-2-yl)-1H-pyrazol-3-yl)amino)phenol
102		N-(3-methyl-4-((5-(5-methylthiophen-2-yl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
103		4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
104		4-((5-(thiazol-2-yl)-1H-pyrazol-3-yl)amino)phenol Molecular Weight: 258.299
105		3-methyl-4-((5-(thiazol-2-yl)-1H-pyrazol-3-yl)amino)phenol
106		N-(4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)phenyl)acetamide









[190]









107		N-(4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
108		4-(3-((5-(hydroxymethyl)-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol
109		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
110		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
111		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
112		4-((5-(1-methyl-1H-pyrrol-3-yl)-1H-pyrazol-3-yl)amino)phenol
113		3-methyl-4-((5-(1-methyl-1H-pyrrol-3-yl)-1H-pyrazol-3-yl)amino)phenol
114		4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol

115		4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
116		4-((5-(benzo[b]thiophen-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
117		4-((5-(benzo[b]thiophen-3-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
118		4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
119		4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
120		3-methyl-4-((5-(thiazol-2-yl)-1H-pyrazol-3-yl)amino)phenol
121		3-ethyl-4-((5-(thiazol-2-yl)-1H-pyrazol-3-yl)amino)phenol
122		tert-butyl 3-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)-1H-indol-1-carboxylate
123		tert-butyl 3-(3-((2-ethyl-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)-1H-indol-1-carboxylate

124		4-(3-((3-hydroxycyclohexyl)amino)-1H-pyrazol-5-yl)phenol
125		N-(5-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-4-methylpyridin-2-yl)acetamide
126		3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
127		N-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
128		N-(4-((5-(benzo[b]thiophen-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
129		4-((5-(6-bromopyridin-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
130		4-((5-(6-bromopyridin-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
131		3-methyl-4-((5-(pyridin-2-yl)-1H-pyrazol-3-yl)amino)phenol
132		3-ethyl-4-((5-(pyridin-2-yl)-1H-pyrazol-3-yl)amino)phenol









133		4-((5-(5-bromopyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
134		4-((5-(4-bromopyridin-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
135		N-(3-methyl-4-((5-(thiazol-2-yl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
136		N-(4-((5-(6-bromopyridin-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
137		1-(4-(4-(3-((2-ethyl-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)phenyl)piperazin-1-yl)ethan-1-one
138		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
139		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
140		4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
141		4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol

142		N-(3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
143		N-(5-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-4-methylpyridin-2-yl)acetamide
144		N-(3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
145		N-(2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenyl)acetamide
146		3-ethyl-4-((5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)phenol
147		N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)butyramide
148		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
149		N-(4-((5-(4-hydroxy-3,5-dimethylphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide

150		4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)-2,6-dimethylphenol
151		2,2,2-trifluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
152		4-(3-((4-amino-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol
153		N-(4-((5-(4-((tert-butyl)dimethylsilyl)oxy)-3,5-dimethylphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
154		N-(3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
155		N-(2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenyl)methanesulfonamide
156		4-((5-(6-chloropyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
157		ethyl 4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl carbamate

158		N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
159		1-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
160		4-(3-((2-ethyl-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)-2-fluorophenol
161		1-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
162		2-fluoro-4-(3-((2-fluoro-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)phenol
163		4-(3-((6-fluoropyridin-3-yl)amino)-1H-pyrazol-5-yl)phenol
164		4-(3-((2-methyl-4-((2-morpholinoethyl)amino)phenyl)amino)-1H-pyrazol-5-yl)phenol
165		4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylbenzonitrile










[197]

166		1-(4-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)piperazin-1-yl)ethan-1-one
167		5-(4-(benzyloxy)phenyl)-N-(2-methyl-4-(piperazin-1-yl)phenyl)-1H-pyrazol-3-amine
168		4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3,5-dimethylphenol
169		4-(3-((2-methyl-4-(piperazin-1-yl)phenyl)amino)-1H-pyrazol-5-yl)phenol
170		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
171		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
172		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol
173		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol










174		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenyl)acetamide
175		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol
176		4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenol
177		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenyl)acetamide
178		N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenyl)acetamide
179		N-(2-fluoro-4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenyl)acetamide
180		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-chloropropanamide
181		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide
182		2-(dimethylamino)-N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
183		N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide









184		3-chloro-N-(5-(4-fluorophenyl)-1H-pyrazol-3-yl)pyridin-2-amine
185		3-(dimethylamino)-N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)propanamide
186		4-(3-((3-chloropyridin-2-yl)amino)-1H-pyrazol-5-yl)-2-fluorophenol
187		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
188		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol
189		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol
190		N-(4-((5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
191		N-(4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
192		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol









193		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenol
194		3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenol
195		N-(3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenyl)acetamide
196		N-(3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
197		N-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
198		N-(4-((5-(4-cyanophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
199		N-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
200		N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide

201		N-(3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
202		3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenol
203		N-(3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
204		N-(4-((5-(4-fluorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
205		1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)pyrrolidin-2-one
206		1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)piperidin-2-one
207		1-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
208		methyl 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl carbamate
209		ethyl 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl carbamate









[202]

210		N-(3-methyl-4-((5-(4-(2-oxopyrrolidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
211		N-(3-methyl-4-((5-(4-(2-oxopiperidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
212		4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
213		N-(4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
214		N-(4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
215		1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)azetidin-2-one
216		4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
217		N-(4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
218		N-(4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide








219		N-(3-chloro-4-((5-(4-(2-oxopyrrolidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
220		4-((5-(4-(dimethylamino)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenol
221		N-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
222		1-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
223		ethyl 4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl carbamate
224		methyl 4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl carbamate
225		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
226		N-(3-chloro-4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide

227		4-((5-(3-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
228		N-(4-((5-(3-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
229		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)isobutyramide
230		1-(4-(4-fluoro-3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)pyrrolidin-2-one
231		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
232		methyl (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
233		ethyl (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
234		4-((4-fluoro-5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol








[205]

235		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
236		1-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
237		3-ethyl-4-((4-fluoro-5-(4-(1H-pyrazol-3-yl)amino)phenyl)-1H-pyrazol-3-yl)phenol
238		3-methyl-4-((5-(4-(4-methyl-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
239		N-(3-methyl-4-((5-(4-(4-methyl-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
240		3-methyl-4-((5-(3-(pyridin-3-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
241		3-methyl-4-((5-(4-(pyridin-3-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
242		4-((5-(4-(4-fluoro-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol

[206]

243		N-(4-((5-(4-(4-fluoro-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
244		4-((4-fluoro-5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
245		3-ethyl-4-((5-(3-iodophenyl)-1H-pyrazol-3-yl)amino)phenol
246		4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
247		N-(4-(benzyloxy)phenyl)-5-(4-chlorophenyl)-1H-pyrazol-3-amine
248		N-(4-(benzyloxy)phenyl)-5-(o-tolyl)-1H-pyrazol-3-amine
249		N-(4-(benzyloxy)phenyl)-5-(2,4-dimethylphenyl)-1H-pyrazol-3-amine

[207]

250		N-(4-chloro-2-methylphenyl)-5-(4-chlorophenyl)-1H-pyrazol-3-amine
251		N-(4-chloro-2-methylphenyl)-5-(o-tolyl)-1H-pyrazol-3-amine
252		5-(2,4-dimethylphenyl)-N-(4-fluorophenyl)-1H-pyrazol-3-amine
253		N-(4-fluorophenyl)-5-(o-tolyl)-1H-pyrazol-3-amine
254		5-(4-methoxyphenyl)-N-(4-phenoxyphenyl)-1H-pyrazol-3-amine
255		4-(4-fluoro-5-((4-(trifluoromethoxy)phenyl)amino)-1H-pyrazol-3-yl)phenol
256		4-fluoro-N-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)benzenesulfonamide









[208]

257		4-(5-((4-bromophenyl)amino)-4-fluoro-1H-pyrazol-3-yl)phenol
258		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-2-hydroxyacetamide
259		2-((4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)amino)propane-1,3-diol
260		3-((4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)amino)propan-1-ol
261		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-2-(4-methylpiperazin-1-yl)acetamide
262		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-2-(4-acetylpiperazin-1-yl)acetamide
263		N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)isobutyramide
264		N1-(5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)-2-methyl-N4-(trifluoromethyl)benzene-1,4-diamine
265		4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol

[209]









266		4-(3-((2-methyl-4-((trifluoromethyl)amino)phenyl)amino)-1H-pyrazol-5-yl)phenol
267		4-((5-(4-((trifluoromethyl)amino)phenyl)-1H-pyrazol-3-yl)amino)phenol
268		2-methyl-N4-(trifluoromethyl)-N1-(5-(4-((trifluoromethyl)amino)phenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine
269		3-ethyl-4-((5-(4-((trifluoromethyl)amino)phenyl)-1H-pyrazol-3-yl)amino)phenol
270		2-fluoro-4-(3-((2-methyl-4-((trifluoromethyl)amino)phenyl)amino)-1H-pyrazol-5-yl)phenol
271		N1-(5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)-2-methyl-N4-(trifluoromethyl)benzene-1,4-diamine
272		3-methyl-4-((5-(3-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol




[210]

273		4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
274		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol
275		2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenol
276		3-methyl-4-((5-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
277		4-((5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
278		3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
279		4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
280		1-(4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea

281		methyl 4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenylcarbamate
282		4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol
283		1-(4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
284		methyl 4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenylcarbamate
285		1-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
286		(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
287		4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2,5-difluorophenol
288		1-methyl-3-(3-methyl-4-((5-(4-(methylsulfonyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)urea

[212]

289		methyl (3-methyl-4-((5-(4-(methylsulfonyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
290		1-methyl-3-(3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)urea
291		methyl (3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
292		1-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
293		methyl (4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
294		methyl (3-methyl-4-((5-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
295		3-methyl-4-((5-(4-(trifluoromethyl)phenyl)-1H-pyrazol-3-yl)amino)phenol
296		methyl (3-methyl-4-((5-(4-(trifluoromethyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate

297		methyl (4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
298		methyl (4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
299		methyl (4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate

[214]

[215] Preferably, the compound of the present disclosure is chosen in Table 2 below.

[216]

[Table 2]

Compo und No.	Name
12	4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
24	4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
29	N1-(5-(4-chlorophenyl)-1H-pyrazol-3-yl)-N4-propylbenzene-1,4-diamine
33	4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzonitrile
34	4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methoxyphenol
39	4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
40	4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
49	3-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
58	N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
59	N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
62	4-((5-(4-fluoro-5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
73	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
74	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
77	N-(4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
80	4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
81	3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
85	4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)-2-methylphenol
86	N-(4-((5-(4-hydroxy-3-methylphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
88	2-fluoro-4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol
89	4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol
94	2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
109	N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
110	N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-m

	ethylphenyl)methanesulfonamide
111	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
114	4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
115	4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
117	4-((5-(benzo[b]thiophen-3-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
119	4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
126	3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol
133	4-((5-(5-bromopyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
138	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
139	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
140	4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
141	4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
142	N-(3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
144	N-(3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
146	3-ethyl-4-((5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)phenol
147	N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)butyramide
151	2,2,2-trifluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
152	4-(3-((4-amino-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol
154	N-(3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
155	N-(2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenyl)methanesulfonamide
156	4-((5-(6-chloropyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
157	ethyl (4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
158	N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)

	l)methanesulfonamide
160	4-(3-((2-ethyl-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)-2-fluorophenol
161	1-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
162	2-fluoro-4-(3-((2-fluoro-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)phenol
170	N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
171	N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
172	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol
173	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol
174	N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenyl)acetamide
175	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol
176	4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenol
178	N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenyl)acetamide
180	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-chloropropanamide
181	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide
182	2-(dimethylamino)-N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
183	N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide
185	3-(dimethylamino)-N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)propanamide
187	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
188	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol
189	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol

190	N-(4-((5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
191	N-(4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
192	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol
193	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenol
194	3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenol
195	N-(3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenyl)acetamide
196	N-(3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
197	N-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
198	N-(4-((5-(4-cyanophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
199	N-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
200	N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
201	N-(3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
202	3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenol
203	N-(3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide
204	N-(4-((5-(4-fluorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
205	1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)pyrrolidin-2-one
206	1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)piperidin-2-one
207	1-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
208	methyl (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)

	carbamate
209	ethyl (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl) carbamate
210	N-(3-methyl-4-((5-(4-(2-oxopyrrolidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino) phenyl)acetamide
211	N-(3-methyl-4-((5-(4-(2-oxopiperidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)p henyl)acetamide
212	4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
213	N-(4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl) methanesulfonamide
214	N-(4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)a cetamide
215	1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)azetidin -2-one
216	4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphe nol
217	N-(4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methyl phenyl)acetamide
218	N-(4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methyl phenyl)methanesulfonamide
219	N-(3-chloro-4-((5-(4-(2-oxopyrrolidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)p henyl)acetamide
220	4-((5-(4-(dimethylamino)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methyl phenol
221	N-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)m ethanesulfonamide
222	1-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3 -methylurea
223	ethyl (4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carb amate
224	methyl (4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carb

	amate
225	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
226	N-(3-chloro-4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide
227	4-((5-(3-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
228	N-(4-((5-(3-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
229	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)isobutyramide
230	1-(4-(4-fluoro-3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)pyrrolidin-2-one
231	N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
232	methyl (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
233	ethyl (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
234	4-((4-fluoro-5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
236	1-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
237	3-ethyl-4-((4-fluoro-5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)phenol
240	3-methyl-4-((5-(3-(pyridin-3-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
241	3-methyl-4-((5-(4-(pyridin-3-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
242	4-((5-(4-(4-fluoro-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
243	N-(4-((5-(4-(4-fluoro-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide
244	4-((4-fluoro-5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
245	3-ethyl-4-((5-(3-iodophenyl)-1H-pyrazol-3-yl)amino)phenol
246	4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-1H-pyrazol-3-yl)amino)-3-

	methylphenol
265	4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
272	3-methyl-4-((5-(3-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
273	4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol
274	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol
275	2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenol
276	3-methyl-4-((5-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
277	4-((5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol
278	3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
279	4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol
280	1-(4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
281	methyl (4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
282	4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol
283	1-(4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
284	methyl (4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
285	1-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
286	(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
287	4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2,5-difluorophenol

288	1-methyl-3-(3-methyl-4-((5-(4-(methylsulfonyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)urea
289	methyl (3-methyl-4-((5-(4-(methylsulfonyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
290	1-methyl-3-(3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)urea
291	methyl (3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
292	1-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea
293	methyl (4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
294	methyl (3-methyl-4-((5-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
295	3-methyl-4-((5-(4-(trifluoromethyl)phenyl)-1H-pyrazol-3-yl)amino)phenol
296	methyl (3-methyl-4-((5-(4-(trifluoromethyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate
297	methyl (4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
298	methyl (4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
299	methyl (4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate

[217]

[218]

In yet another embodiment, there is provided a pharmaceutical composition comprising a therapeutically effective amount of a compound of Chemical Formula 1 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[219]

In yet another embodiment, there is provided a pharmaceutical composition comprising a therapeutically effective amount of a compound of Chemical Formula 1

or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable carrier, and a therapeutically effective amount of an active pharmaceutical ingredient, which is not the compound of the present disclosure, selected from the group consisting of cytostatic drugs, angiogenesis inhibitors, kinase inhibitors, cytokine blockers and inhibitors of cell adhesion molecule inhibitors.

[220] In another embodiment, there is provided a method for treating a disease or condition comprising administering to a subject in need thereof a therapeutically effective amount of a compound of Chemical Formula 1 or pharmaceutically acceptable salt thereof, wherein the disease or condition is selected from the group consisting of cancers, neoplasia, or tumors including colorectal cancer, breast cancer, brain tumor, gastric cancer, liver cancer, ovarian cancer, lung cancer, gastrointestinal cancer, leukemia, or melanoma. In another embodiment, the subject is a human. In another embodiment, the disease or condition is colorectal cancer.

[221] That is, there is provided a medical use of Chemical Formula 1 or pharmaceutically acceptable salt thereof, wherein Chemical Formula 1 or pharmaceutically acceptable salt thereof is used as an effective agent. In one embodiment, the medical-use is for treatment or prevention of the disease or condition described above.

[222]

[223] **Medical uses and Methods of treatment of the compounds according to the present disclosure**

[224] The present disclosure further provides methods for treating a disease or condition in a subject having or susceptible to having such a disease or condition, by administering to the subject a therapeutically-effective amount of one or more compounds as described above. In one embodiment, the treatment is preventative treatment. In another embodiment, the treatment is palliative treatment. In another embodiment, the treatment is restorative treatment.

[225] **1. Diseases or conditions**

[226] The compound of the present disclosure for inhibiting TNIK activity is useful for treatment or prevention of various conditions (for example, anti-tumor). The compound can be used for inhibiting or hindering TNIK activity, and for treating a tumor or cancer, or for preventing aggravation of such disease. Thus, the present disclosure provides a method for inhibiting or hindering TNIK activity in a cell, wherein the cell is contacted with an effective amount of a compound of the present disclosure. In one embodiment, such cell is present in a subject (for example, cancer patients). In another embodiment, there is provided a medical use for treating cancer or preventing proliferation of tumor in a subject, using the compound according to the present disclosure. The method of the present disclosure comprises administering to a subject in need of treatment or prevention a pharmaceutical composition containing a

therapeutically or prophylactically effective amount of TNiK inhibitor.

[227] In one embodiment, there is provided a method for inhibiting or blocking TNiK activity in a tumor or cancer cell. For example, the present disclosure is used for inhibiting TNiK activity in a cell such as colorectal cancer cell, breast cancer cell, brain tumor cell, gastric cancer cell, liver cancer cell, ovarian cancer cell, lung cancer cell, gastrointestinal cancer cell, leukemia cell, or melanoma cell. In this method, the present disclosure provides a method for inhibiting the growth or proliferation of cells, particularly tumor or cancer cells, in a subject. In this method, tumor cells are present *in vivo*. The compound of the present disclosure can be administered to the subject as a form of the pharmaceutical composition described herein.

[228] In another embodiment, there is provided a method for treating or preventing a cancer or tumor in a subject. The cancer includes, but is not limited to, colorectal cancer, breast cancer, brain tumor, gastric cancer, liver cancer, ovarian cancer, lung cancer, gastrointestinal cancer, leukemia, or melanoma. The method comprises administering to a subject in need of treatment an enough amount of the compound, that is, a therapeutically amount of the compound of the present disclosure.

[229] **2. Subjects**

[230] Suitable subjects to be treated according to the present disclosure include mammalian subjects. Mammals according to the present disclosure include, but are not limited to, human, canine, feline, bovine, caprine, equine, ovine, porcine, rodents, lagomorphs, primates, and the like, and encompass mammals *in utero*. Subjects may be of either gender and at any stage of development.

[231] In one embodiment, the suitable subject to be treated according to the present disclosure is human.

[232] **3. Administration and dosing**

[233] The compounds of the present disclosure are generally administered in a therapeutically effective amount.

[234] The compounds of the present disclosure can be administered by any suitable route in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. An effective dosage is typically in the range of about 0.001 to about 100 mg per kg body weight per day, preferably about 0.01 to about 50 mg/kg/day, in single or divided doses. Depending on age, species and disease or condition being treated, dosage levels below the lower limit of this range may be suitable. In other cases, still larger doses may be used without harmful side effects. Larger doses may also be divided into several smaller doses, for administration throughout the day. Methods for determining suitable doses are well known in the art to which the present disclosure pertains. For example, Remington: The Science and Practice of Pharmacy, Mack Publishing Co., 20th ed., 2000 can be used.

[235]

[236] **Pharmaceutical Compositions, Dosage Forms and Administration Routes**

[237] For the treatment of the diseases or conditions referred to above, the compounds described herein or pharmaceutically acceptable salts thereof can be administered as follows:

[238] **Oral administration**

[239] The compounds of the present disclosure may be administered orally, including by swallowing, so that the compound enters the gastrointestinal tract, or absorbed into the blood stream directly from the mouth (e.g., buccal or sublingual administration).

[240] Suitable compositions for oral administration include solid, liquid, gel or powder formulations, and have a dosage form such as tablet, lozenge, capsule, granule or powder.

[241] Compositions for oral administration may be formulated as immediate or modified release, including delayed or sustained release, optionally with enteric coating.

[242] Liquid formulations can include solutions, syrups and suspensions, which can be used in soft or hard capsules. Such formulations may include a pharmaceutically acceptable carrier, for example, water, ethanol, polyethylene glycol, cellulose, or an oil. The formulation may also include one or more emulsifying agents and/or suspending agents.

[243] In a tablet dosage form the amount of drug present may be from about 0.05% to about 95% by weight, more typically from about 2% to about 50% by weight of the dosage form. In addition, tablets may contain a disintegrant, comprising from about 0.5% to about 35% by weight, more typically from about 2% to about 25% of the dosage form. Examples of disintegrants include, but are not limited to, lactose, starch, sodium starch glycolate, crospovidone, croscarmellose sodium, maltodextrin, or mixtures thereof.

[244] Suitable lubricants, for use in a tablet, may be present in amounts from about 0.1% to about 5% by weight, and include, but are not limited to, talc, silicon dioxide, stearic acid, calcium, zinc or magnesium stearate, sodium stearyl fumarate and the like.

[245] Suitable binders, for use in a tablet, include, but are not limited to, gelatin, polyethylene glycol, sugars, gums, starch, polyvinyl pyrrolidone, hydroxypropyl cellulose, hydroxypropylmethyl cellulose and the like. Suitable diluents, for use in a tablet, include, but are not limited to, mannitol, xylitol, lactose, dextrose, sucrose, sorbitol, microcrystalline cellulose and starch.

[246] Suitable solubilizers, for use in a tablet, may be present in amounts from about 0.1% to about 3% by weight, and include, but are not limited to, polysorbates, sodium lauryl sulfate, sodium dodecyl sulfate, propylene carbonate, diethyleneglycol monoethyl ether, dimethyl isosorbide, polyethylene glycol (natural or hydrogenated) castor oil, HCORTM (Nikkol), oleyl ester, GelucireTM, caprylic/caprylic acid mono/diglyceride,

sorbitan fatty acid esters, and Solutol HS TM.

[247] **Parenteral Administration**

[248] Compounds of the present disclosure may be administered directly into the blood stream, muscle, or internal organs. Suitable means for parenteral administration include intravenous, intra-muscular, subcutaneous intraarterial, intraperitoneal, intrathecal, intracranial, and the like. Suitable devices for parenteral administration include injectors (including needle and needle-free injectors) and infusion methods.

[249] Compositions for parenteral administration may be formulated as immediate or modified release, including delayed or sustained release.

[250] Most parenteral formulations are aqueous solutions containing excipients, including salts, buffering agents and isotonic agents.

[251] Parenteral formulations may also be prepared in a dehydrated form (e.g., by lyophilization) or as sterile non-aqueous solutions. These formulations can be used with a suitable vehicle, such as sterile water. Solubility-enhancing agents may also be used in preparation of parenteral solutions.

[252] **Topical Administration**

[253] Compounds of the present disclosure may be administered topically to the skin or transdermally. Formulations for this topical administration can include lotions, solutions, creams, gels, hydrogels, ointments, foams, implants, patches and the like. Pharmaceutically acceptable carriers for topical administration formulations can include water, alcohol, mineral oil, glycerin, polyethylene glycol and the like. Topical administration can also be performed by electroporation, iontophoresis, phonophoresis and the like.

[254] Compositions for topical administration may be formulated as immediate or modified release, including delayed or sustained release.

[255] **References for preparing pharmaceutical compositions**

[256] Methods for preparing pharmaceutical compositions for treating or preventing a disease or condition are well known in the art to which the present disclosure pertains. For example, based on *Handbook of Pharmaceutical Excipients* (7th ed.), *Remington: The Science and Practice of Pharmacy* (20th ed.), *Encyclopedia of Pharmaceutical Technology* (3rd ed.), or *Sustained and Controlled Release Drug Delivery Systems* (1978), pharmaceutically acceptable excipients, carriers, additives and so on can be selected and then mixed with the compounds of the present disclosure for making the pharmaceutical compositions.

[257]

[258] **Combinations and Combination Therapy**

[259] The compounds of the present disclosure can be used, alone or in combination with other pharmaceutically active compounds, to treat conditions such as those previously

described above. The compound(s) of the present invention and other pharmaceutically active compound(s) can be administered simultaneously (either in the same dosage form or in separate dosage forms) or sequentially. Accordingly, in one embodiment, the present disclosure comprises methods for treating a condition by administering to the subject a therapeutically-effective amount of one or more compounds of the present disclosure and one or more additional pharmaceutically active compounds.

[260] In another embodiment, there is provided a pharmaceutical composition comprising one or more compounds of the present disclosure, one or more additional pharmaceutically active compounds, and a pharmaceutically acceptable carrier

[261] In another embodiment, the one or more additional pharmaceutically active compounds is an anti-cancer drug. For example, the anti-cancer drug is EGFR kinase inhibitors, MEK inhibitors, VEGFR inhibitors, anti-VEGFR2 antibodies, KDR antibodies, AKT inhibitors, PDK-1 inhibitors, PI3K inhibitors, c-kit/Kdr tyrosine kinase inhibitors, Bcr-Abl tyrosine kinase inhibitors, VEGFR2 inhibitors, PDGFR-beta inhibitors, KIT inhibitors, Flt3 tyrosine kinase inhibitors, PDGF receptor family inhibitors, Flt3 tyrosine kinase inhibitors, RET tyrosine kinase receptor family inhibitor, VEGF-3 receptor antagonists, Raf protein kinase family inhibitor, angiogenesis inhibitors, Erb2 inhibitors, mTOR inhibitors, IGF-1R antibodies, NFkB inhibitors, proteasome inhibitors, chemotherapy agents, or glucose reduction agents.

[262] In one embodiment, an active agent used in the combination medicine and/or combination therapy with the compound(s) of the present disclosure is an anti-cancer drug. That is, the compound(s) of the present invention may be administered concurrently or sequentially to a subject who is taking one or more anti-cancer drug. Such anti-cancer drug includes, but is not limited to, alkylating agents such as nitrogen mustard, chlorambucil, cytoxan, ifosfamide, melphalan, thiopeta and busulfan; antimetabolites such as methotrexate, 5-fluorouracil, cytosine arabinoside (ara-C), 5-azacitidine, 6-mercaptopurine, 6-thioguanine, and fludarabine phosphat; antitumor antibiotics such as doxorubicin, adriamycin, daunorubicin, dactinomycin, bleomycin, mitomycin C, plicamycin, idarubicin, and mitoxantrone; vinca alkaloids and epipodophyllotoxins such as vincristine, vinblastine, vindesine, etoposide, and teniposide; nitrosoureas such as carmustine, lomustine, semustine and streptozotocin; synthetic agents such as dacarbazine, hexamethyl melamine, hydroxyurea, mitotane procabazine, cisplatin, cisplatinum and carboplatin; corticosteroids (cortisone acetate, hydrocortisone, prednisone, prednisolone, methyl prednisolone and dexamethasone), estrogens (diethylstilbestrol, estradiol, esterified estrogens, conjugated estrogens, chlorotrianisene), progestones (medroxyprogesterone acetate, hydroxy progesterone caproate, megestrol acetate), anti-estrogens (tamoxifen), aromatase inhibitors (aminoglutethimide), androgens (testosterone propionate), methyl testosterone, flu-

oxymesterone, testolactone), anti-androgens (flutamide), LHRH analogues (leuprolide acetate), and endocrines for prostate cancer (ketoconazole).

- [263] In one embodiment, an active agent used in the combination medicine and/or combination therapy with the compound(s) of the present disclosure is a drug for colorectal cancer. In another embodiment, the drug for colorectal cancer is based on regimens FOLFOX or FOLFIRI including 5-FU, leucovorin, oxaliplatin, irinotecan or their combinations. In a conventional standard method of treatment, the combination therapy is used together with cetuximab and/or bevacizumab. When the compound of the present disclosure is used with other anti-colon drugs, the compound of the present disclosure may be locally administered by injection to treat non-invasive colon cancer.
- [264] Therapeutically effective dosages vary when the drugs are used in treatment combinations. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient. In any case, the multiple therapeutic agents (one of which is a TNIK inhibitor as described herein) are administered in any order, or even simultaneously. If simultaneously, the multiple therapeutic agents are optionally provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills).
- [265] In some embodiments, one of the therapeutic agents is given in multiple doses, or both are given as multiple doses. If not simultaneous, the timing between the multiple doses optionally varies from more than zero weeks to less than twelve weeks.
- [266] In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents, the use of multiple therapeutic combinations are also envisioned. It is understood that the dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, is optionally modified in accordance with a variety of factors. These factors include the disorder from which the subject suffers, as well as the age, weight, sex, diet, and medical condition of the subject. Thus, the dosage regimen actually employed varies widely, in some embodiments, and therefore deviates from the dosage regimens set forth herein.
- [267] The pharmaceutical agents which make up the combination therapy disclosed herein are optionally a combined dosage form or in separate dosage forms intended for substantially simultaneous administration. The pharmaceutical agents that make up the combination therapy are optionally also administered sequentially, with either agent being administered by a regimen calling for two-step administration. The two-step administration regimen optionally calls for sequential administration of the active agents or spaced-apart administration of the separate active agents. The time period between the multiple administration steps ranges from, a few minutes to several hours, depending upon the properties of each pharmaceutical agent, such as potency, solubility, bioavailability, plasma half-life and kinetic profile of the pharmaceutical

agent. Circadian variation of the target molecule concentration is optionally used to determine the optimal dose interval.

Advantageous Effects of Invention

[268] The present disclosure provides a compound having various pharmacological effects by inhibiting TNIK activity, a pharmaceutical composition having the compound as an effective agent, a medical use, particularly for treating cancers, of the compound, and a method of treatment or prevention comprising administering the compound to a subject in need of such treatment or prevention. The compounds of the present disclosure and pharmaceutically acceptable salts thereof have good safety and high selectivity for TNIK, and thus exhibit superior property as a drug.

Brief Description of Drawings

[269] Figures 1, 2 and 3 are graphs showing the results of the mouse xenograft test performed with the compounds according to some examples of the present disclosure.

Mode for the Invention

[270] Hereinafter, the present disclosure is described in considerable detail with examples to help those skilled in the art understand the present disclosure. However, the following examples are offered by way of illustration and are not intended to limit the scope of the invention. It is apparent that various changes may be made without departing from the spirit and scope of the invention or sacrificing all of its material advantages.

[271]

Preparation of Compounds of the Present Disclosure

[272] Reagents and solvents used below were purchased from Aldrich Chemical Co. (Milwaukee, Wisconsin, USA). ¹H-NMR spectra were evaluated with Bruker Avance 300MHz, Bruker Avance III HD 300MHz, Bruker Avance 500MHz NMR spectrometer and so on.


[274] Below, the illustrating synthetic examples of some compounds of the present disclosure are described, and other compounds can be prepared by the similar method to one described below with different starting or reacting materials.

[275]

[276] Synthesis Example 1: Preparation of Compound 2

[277] Step 1. 1-(4-chlorophenyl)-3,3-bis(methylthio)prop-2-en-1-one

[278]




[279] 4-chloroacetophenone (5.4 mL, 66.6 mmol) and benzene (220 mL) were mixed at ambient temperature in a 500 mL round flask. After cooled using an ice bath, NaH (5.32 g, 133.0 mmol, 2.0 eq.) was added slowly and stirred for 5 minutes. Carbon disulfide (6.0 mL, 99.9 mmol, 1.5 eq.) was added slowly and stirred at ambient temperature for 5 minutes, and then iodomethane (12.4 mL, 199.8 mmol, 3.0 eq.) was added slowly and stirred at ambient temperature for 5 minutes. After that, *N,N*-dimethylacetamide (13 mL, 139.9 mmol, 2.1 eq.) was added slowly and stirred at ambient temperature for 3 hours. After the reaction ended, H₂O was added for quenching, and the reactant was extracted with EtOAc. The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, it was filtered with Et₂O to provide the compound as a yellow solid (8.1 g) (yield: 47%).

[280] Step 2.

(*Z*)-3-((4-(benzyloxy)phenyl)amino)-1-(4-chlorophenyl)-3-(methylthio)prop-2-en-1-one


[281]



[282] 1-(4-chlorophenyl)-3,3-bis(methylthio)prop-2-en-1-one (1.94 g, 7.5 mmol) prepared in Step 1 and toluene (40 mL) were stirred in a 100 mL round flask. Then BF₃·OEt₂ (200 μL, 1.5 mmol, 0.2 eq.) was added, and 4-benzyloxyaniline (2.65 g, 11.25 mmol, 1.5 eq.) was added. After that, the reaction mixture was reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor and the remaining reactant was filtered with MeOH to provide the compound as a yellow solid (1.83 g) (yield: 60%).

[283] Step 3. *N*-(4-(benzyloxy)phenyl)-5-(4-chlorophenyl)-1H-pyrazol-3-amine


[284]



[285] (Z)-3-((4-(benzyloxy)phenyl)amino)-1-(4-chlorophenyl)-3-(methylthio)prop-2-en-1-one (410 mg, 1.0 mmol) prepared in Step 2, *t*-BuOH (7 mL) and AcOH (75 μ L) were mixed in a 50 mL round flask. Then Hydrazine hydrate (75 μ L, 1.5 mmol, 1.5 eq.) was added and reacted under reflux for 3 hours. After the reaction ended, the solvent was removed with a rotavapor and the reaction was filtered with MeOH to provide the compound as a yellow solid (336 mg) (yield: 89%).

[286] Step 4. 4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)phenol

[287]



[288] N-(4-(benzyloxy)phenyl)-5-(4-chlorophenyl)-1H-pyrazol-3-amine (94 mg, 0.25 mmol) prepared in Step 3 and dichloromethane (10 mL) were stirred at -78 $^{\circ}$ C for 15 minutes in a 50 mL round flask. BBr₃ (1M in DCM) (1.2 mL) was added slowly and stirred at ambient temperature for 30 minutes. After the reaction ended, it was cooled using an ice bath and MeOH was added for quenching. The solvent was removed with a rotavapor. The reactant was extracted with H₂O (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound as a white solid (50 mg) (yield: 70%).

[289] ¹H-NMR (300 MHz, DMSO-*d*₆) δ 12.31 (s, 1H), 8.67 (s, 1H), 7.97 (s, 1H), 7.73 (d, 2H, *J*=9.0 Hz), 7.47 (d, 2H, *J*=6.0 Hz), 7.12 (s, 1H), 6.62 (d, 2H, *J*=6.0 Hz), 6.17 (s, 1H).


[290]

[291] Synthesis Example 2: Preparation of

4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-isopropoxyphenol (Compound **28**)


[292] Step 1. 3-isopropoxy-4-nitrophenol

[293]




[294] THF (30 mL) and NaH (1.2 g, 30.0 mmol, 3.0 eq.) were stirred in a 100 mL round flask. *i*-PrOH (2.3 mL, 30.0 mmol, 3.0 eq.) was added and then stirred at ambient temperature for 10 minutes. After that, 3-fluoro-4-nitrophenol (1.57 g, 10.0 mmol) was added and reacted under reflux for 12 hours. After the reaction ended, it was cooled at ambient temperature and H₂O was added for quenching. The reactant was extracted with 1N HCl (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was dried with MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound as a dark brown solid (1.87 g) (yield: 95%).

[295] Step 2. 4-amino-3-isopropoxyphenol



[297] 3-isopropoxy-4-nitrophenol (1.87 g, 9.48 mmol) prepared in Step 1 and EtOAc/THF (10/10 mL) were stirred at ambient temperature in a 100 mL round flask. Pd/C (370 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter, and silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (836 mg) (yield: 53%).

[298] Step 3. 4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-isopropoxyphenol



[300] 1-(4-chlorophenyl)-3,3-bis(methylthio)prop-2-en-1-one (259 mg, 1.0 mmol) prepared in Step 1 for preparation of Compound **2** and Toluene (10 mL) were stirred in a vial for microwave reaction. $\text{BF}_3 \cdot \text{OEt}_2$ (25 μL , 0.2 mmol, 0.2 eq.) and 4-amino-3-isopropoxyphenol (251 mg, 1.5 mmol, 1.5 eq.) prepared in Step 2 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and extracted with H_2O (50ml) and EtOAc (50 mL x 3). And, the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (10 mL), AcOH (75 μL) and Hydrazine hydrate (75 μL , 1.5 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and extracted with H_2O (50ml) and EtOAc (50 mL x 3). Then, the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound as a dark brown solid (2 steps, 96 mg) (yield: 28%).


[301] ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.37 (s, 1H), 8.80 (s, 1H), 7.73 (d, $J = 8.1$ Hz, 2H), 7.47 (d, $J = 8.0$ Hz, 2H), 6.77 (s, 1H), 6.43 (d, $J = 2.7$ Hz, 1H), 6.35 - 6.23 (m, 2H), 4.51 (p, $J = 6.0$ Hz, 1H), 1.30 (d, $J = 6.0$ Hz, 6H).

[302]

[303] Synthesis Example 3: Preparation of 3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol (Compound **81**)

[304] Step 1. 3-ethyl-4-nitrophenol


[305]



[306] EtOH/conc. HCl (45 mL/45 mL) and 3-ethylphenol (74 mmol, 11 mL) were added in a 250 mL RBF. The mixture was stirred in an ice bath for 30 minutes. NaNO_2 (1.5 eq, 7.8 g) then was added slowly and stirred to provide a solid. The reactant was stirred for 30 minutes and then water (100 mL) was added to end the reaction. The produced solid was filtered to provide 3-ethyl-4-nitrophenol (10.38g) as a brown solid (yield: 84%).

[307] Step 2. 4-amino-3-ethylphenol

[308]




[309] 3-ethyl-4-nitrophenol (62 mmol, 10.4 g) was dissolved in EA/MeOH (90/10 mL) and Pd/C (1.0 g) was added. The mixture was reacted with a hydrogenator for 4 hours. After the reaction ended, Pd/C was removed using a celite filter and the solvent was removed under reduced pressure. The solid was filtered with EA to provide 4-amino-3-ethylphenol (7.99 g) (yield: 94%).

[310] Step 3.

(Z)-1-(4-(benzyloxy)phenyl)-3-((2-ethyl-4-hydroxyphenyl)amino)-3-(methylthio)prop-2-en-1-one


[311]



[312] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (496 mg, 1.5 mmol) and toluene (15 mL) were stirred in a 100 mL round flask. $\text{BF}_3 \cdot \text{OEt}_2$ (37 μL , 0.3 mmol, 0.2 eq.) was added and then 4-amino-3-ethylphenol 3-aminophenol (308 mg, 2.25 mmol, 1.5 eq.) prepared in Step 2 was added. The mixture was reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and the reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3). The organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (298 mg) (yield: 47%).

[313] Step 4. 4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol


[314]



[315] (Z)-1-(4-(benzyloxy)phenyl)-3-((2-ethyl-4-hydroxyphenyl)amino)-3-(methylthio)prop-2-en-1-one (250 mg, 0.6 mmol), *t*-BuOH (8 mL), and AcOH (46 μL) were stirred in a 50 mL round flask. Hydrazine hydrate (46 μL , 0.75 mmol, 1.5 eq.) was added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor and the reactant was filtered with MeOH to provide the compound (227 mg) (yield: 99%).

[316] Step 5. 3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol

[317]



[318] 4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol (300 mg, 0.78 mmol) prepared in Step 4 and DCM (8 mL) were stirred at -78 °C for 15 minutes in a 50 mL round flask. BBr₃ (1M in DCM) (3.9 mL) was added slowly and the mixture was stirred at ambient temperature for 3 hours. After the reaction ended, the reactant was cooled using an ice bath and MeOH was used for quenching, and the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (44 mg) (yield: 20%).


[319] ¹H NMR (500 MHz, DMSO) δ 12.01 - 11.82 (m, 1H), 9.57 (s, 1H), 8.78 (s, 1H), 7.48 (d, *J* = 8.6 Hz, 2H), 7.25 (s, 1H), 6.89 (s, 1H), 6.78 (d, *J* = 8.6 Hz, 2H), 6.58 (d, *J* = 2.7 Hz, 1H), 6.51 (dd, *J* = 8.6, 2.8 Hz, 1H), 5.87 (s, 1H), 2.56 (q, *J* = 7.5 Hz, 2H), 1.13 (t, *J* = 7.5 Hz, 3H).

[320]

[321] Synthesis Example 4: Preparation of
N1-(5-(4-chlorophenyl)-1H-pyrazol-3-yl)-N4-phenylbenzene-1,4-diamine (Compound 30)

[322] Step 1. tert-butyl (4-nitrophenyl)(phenyl)carbamate


[323]



[324] 4-nitrodiphenylamine (1.07 g, 5.0 mmol) and DCM (30 mL) were stirred in a 100 mL round flask. Et₃N (1.4 mL, 10.0 mmol, 2.0 eq.) and (Boc)₂O (1.64 g, 7.5 mmol, 1.5 eq.) were added and the mixture was reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and extracted with H₂O (50ml) and DCM (50 mL x 3). The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. Next step was performed without purification.

[325] Step 2. tert-butyl (4-aminophenyl)(phenyl)carbamate

[326]




[327]

Tert-butyl (4-nitrophenyl)(phenyl)carbamate of tert-butyl (4-nitrophenyl)(phenyl)carbamate (2.53 g, 8.0 mmol) prepared in Step 1 and THF(60 mL) were stirred in a 500 mL round flask. MeOH (6 mL) and H₂O (90 mL) were added, and then NaS₂O₄ (12.54 g, 72.0 mmol, 9.0 eq.) and NaHCO₃ (6.05 g, 72.0 mmol, 9.0 eq.) were added slowly, and the mixture was stirred for 5 minutes. After the reaction ended, the reactant was extracted several times with EtOAc and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 597 mg) (yield: 42%).

[328]

Step 3. N1-(5-(4-chlorophenyl)-1H-pyrazol-3-yl)-N4-phenylbenzene-1,4-diamine

[329]



[330]

1-(4-chlorophenyl)-3,3-bis(methylthio)prop-2-en-1-one (259 mg, 1.0 mmol) and Toluene (10 mL) were stirred in a vial for microwave reaction. BF₃.OEt₂ (25 μL, 0.2 mmol, 0.2 eq.) and tert-butyl (4-aminophenyl)(phenyl)carbamate (426 mg, 1.5 mmol, 1.5 eq.) prepared in Step 2 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and the reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3). The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. *t*-BuOH(10 mL), AcOH (75 μL) and Hydrazine hydrate (75 μL, 1.5 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and the organic

solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 154 mg) (yield: 43%).


[331] ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.43 (s, 1H), 8.28 (s, 1H), 7.80 - 7.67 (m, 3H), 7.51 (s, 2H), 7.29 (s, 1H), 7.13 (t, $J = 7.6$ Hz, 2H), 6.99 (d, $J = 8.2$ Hz, 2H), 6.88 (d, $J = 8.0$ Hz, 2H), 6.66 (t, $J = 7.3$ Hz, 1H), 6.26 (s, 1H).

[332]

[333] Synthesis Example 5: Preparation of 4-fluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)benzenesulfonamide (Compound **52**)

[334] Step 1. 4-fluoro-N-(4-nitrophenyl)benzenesulfonamide


[335]



[336] 4-nitroaniline (1.41 g, 10.0 mmol) and DCM (30 mL) were stirred in a 100 mL round flask. Pyridine (2.4 mL, 30.0 mmol, 3.0 eq.) was added and cooled with an ice bath. 4-fluorobenzenesulfonyl chloride (1.98 g, 10.0 mmol, 1.0 eq.) was added slowly and the mixture was stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. The produced solid was filtered with Ether to provide the compound (2.9 g) (yield: 98%).

[337] Step 2. N-(4-aminophenyl)-4-fluorobenzenesulfonamide

[338]




[339] 4-fluoro-N-(4-nitrophenyl)benzenesulfonamide (2.9 g, 9.8 mmol) prepared in Step 1 and EtOAc/MeOH (30/30 mL) were stirred at ambient temperature in a 250 mL round flask. Pd/C (580 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2.09 g) (yield: 80%).

[340] Step 3. N-

4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)phenyl)-4-fluorobenzenesulfonamide

mide

[341]




[342] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (496 mg, 1.5 mmol) and Toluene (15 mL) were stirred in a vial for microwave reaction. $\text{BF}_3 \cdot \text{OEt}_2$ (37 μL , 0.3 mmol, 0.2 eq.) and N-(4-aminophenyl)-4-fluorobenzenesulfonamide (599 mg, 2.25 mmol, 1.5 eq.) prepared in Step 2 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor H_2O (50ml) and the reactant was extracted with EtOAc (50 mL x 3). Then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (30 mL), AcOH (113 μL) and Hydrazine hydrate (113 μL , 2.25 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and the reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3). Then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 476mg) (yield: 62%).

[343] Step 4.

4-fluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)benzenesulfonamide

[344]



[345] N-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)phenyl)-4-fluorobenzenesulfonamide (257 mg, 0.5 mmol) prepared in Step 3 and EtOAc/MeOH (8/8 mL) were stirred at ambient temperature in a round flask. Pd/C (51 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was

removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (168 mg) (yield: 79%).


[346] ^1H NMR (300 MHz, DMSO- d_6) δ 12.17 (s, 1H), 9.66 (s, 2H), 8.37 (s, 1H), 7.76 - 7.65 (m, 2H), 7.51 (d, J = 8.3 Hz, 2H), 7.37 (t, J = 8.7 Hz, 2H), 7.19 (d, J = 8.4 Hz, 2H), 6.89 - 6.75 (m, 4H), 6.02 (s, 1H).

[347]

[348] Synthesis Example 6: Preparation of N-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide (Compound **58**)

[349] Step 1. N-(3-methyl-4-nitrophenyl)methanesulfonamide


[350]



[351] 3-methyl-4-nitroaniline (1.57 g, 10.0 mmol) and DCM (30 mL) were stirred in a 100 mL round flask. Pyridine (2.4 mL, 30.0 mmol, 3.0 eq.) was added and the mixture was cooled using an ice bath. Then methanesulfonyl chloride (774 μL , 10.0 mmol, 1.0 eq.) was added slowly and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a rotavapor and the reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3). Then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. The produced solid was filtered with ether to provide the compound (1.98 g) (yield: 86%).

[352] Step 2. N-(4-amino-3-methylphenyl)methanesulfonamide

[353]




[354] N-(3-methyl-4-nitrophenyl)methanesulfonamide (1.98 g, 8.6 mmol) prepared in Step 1 and EtOAc/MeOH (40/40 mL) were stirred at ambient temperature in a 250 mL round flask. Pd/C (400 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter, and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (1.36 g) (yield: 79%).

[355] Step 3. N-

(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide


[356]



[357] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (496 mg, 1.5 mmol) and Toluene (15 mL) were stirred in a MV vial. $\text{BF}_3 \cdot \text{OEt}_2$ (37 μL , 0.3 mmol, 0.2 eq.) and N-(4-amino-3-methylphenyl)methanesulfonamide (451 mg, 2.25 mmol, 1.5 eq.) prepared in Step 2 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor and the reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3). Then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (30 mL), AcOH (113 μL) and Hydrazine hydrate (113 μL , 2.25 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 543 mg) (yield: 81%).

[358] Step 4. N-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide

[359]



[360] N-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesu

lfonamide (224 mg, 1.0 mmol) prepared in Step 3 and EtOAc/MeOH/THF (10/10/10 mL) were stirred at ambient temperature in a round flask. Pd/C (90 mg, 20 wt%) was added and stirred at ambient temperature for 48 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (24 mg) (yield: 13%).


[361] ^1H NMR (300 MHz, DMSO- d_6) δ 12.21 (s, 1H), 9.63 (s, 1H), 9.17 (s, 1H), 7.68 (s, 1H), 7.51 (d, J = 8.5 Hz, 2H), 7.29 (s, 1H), 6.94 (d, J = 8.9 Hz, 2H), 6.81 (d, J = 8.5 Hz, 2H), 6.15 (s, 1H), 2.86 (s, 3H), 2.22 (s, 3H).

[362]

[363] Synthesis Example 7: Preparation of N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide (Compound **59**)

[364] Step 1. N-(3-methyl-4-nitrophenyl)acetamide


[365]



[366] 3-methyl-4-nitroaniline (3.14 g, 20.0 mmol) and pyridine (20 mL) were stirred in a 100 mL round flask. The mixture was cooled in an ice bath, and acetic anhydride (1.9 mL, 20.0 mmol, 2.0 eq.) was added slowly and stirred at ambient temperature for 24 hours. After the reaction ended, the reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. Next step was performed without purification.

[367] Step 2. N-(4-amino-3-methylphenyl)acetamide


[368]



56% (2 steps)


[369] N-(3-methyl-4-nitrophenyl)acetamide (4.78 g, 24.6 mmol) and EtOAc/MeOH (100/100 mL) were stirred at ambient temperature in a round flask. Pd/C (1.9 g, 40 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 1.85 g) (yield: 56%).

- [370] Step 3. N-
 (4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
 [371]



- [372] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (496 mg, 1.5 mmol) and toluene (15 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (37 μL , 0.3 mmol, 0.2 eq.) and N-(4-amino-3-methylphenyl)acetamide (451 mg, 2.25 mmol, 1.5 eq.) were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. Then the reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (30 mL), AcOH (113 μL) and Hydrazine hydrate (113 μL , 2.25 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. Next step was performed without purification.

- [373] Step 4. N-
 (4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide
 [374]



- [375] N-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide (610 mg, 1.48 mmol) prepared in Step 3 and EtOAc/MeOH (10/10 mL) were stirred at ambient temperature in a round flask. Pd/C (120 mg, 20 wt%) was added and stirred at ambient temperature for 48 hours. After the reaction ended, Pd/C was removed with a

celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (3 steps, 24 mg) (yield: 34%).


[376] ^1H NMR (300 MHz, DMSO- d_6) δ 12.14 (s, 1H), 9.62 (s, 1H), 7.51 (d, J = 8.2 Hz, 2H), 7.30 (s, 1H), 7.27 - 7.15 (m, 2H), 6.80 (d, J = 8.2 Hz, 2H), 6.10 (s, 1H), 2.20 (s, 3H), 1.98 (s, 3H).

[377]

[378] Synthesis Example 8: Preparation of
2,2,2-trifluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide (Compound **151**)

[379] Step 1. 2,2,2-trifluoro-N-(3-methyl-4-nitrophenyl)acetamide


[380]



[381] 3-methyl-4-nitroaniline (1.0 g, 6.6 mmol) and DCM (55 mL) were stirred at -78 °C in a 100 mL round flask. Tridluoroacetic anhydride (1.4 mL, 9.9 mmol, 1.5 eq.) was added slowly and then trimethylamine (3.4 mL, 18.48 mmol, 2.8 eq.) was added and stirred for 3 hours. After the reaction ended, the reactant was cooled in an ice bath, and 1N HCl was added and extracted with DCM several times. Then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (1.5 g) (yield: 92%).

[382] Step 2. N-(4-amino-3-methylphenyl)-2,2,2-trifluoroacetamide

[383]




[384] 2,2,2-trifluoro-N-(3-methyl-4-nitrophenyl)acetamide (1.5 g, 6.0 mmol) prepared in Step 1 and EtOAc/MeOH (20/15 mL) were stirred at ambient temperature in a round flask. Pd/C (300 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (1.19 g) (yield: 91%).

[385] Step 3.

(Z)-N-(4-((3-(4-(benzyloxy)phenyl)-1-(methylthio)-3-oxoprop-1-en-1-yl)amino)-3-methylphenyl)-2,2,2-trifluoroacetamide


[386]



[387] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (350 mg, 1.06 mmol) and THF (4 mL) were stirred in a round flask. $\text{BF}_3 \cdot \text{OEt}_2$ (26 μL , 0.22 mmol, 0.2 eq.) was added and 2,2,2-trifluoro-N-(3-methylphenyl)acetamide (347 mg, 1.59 mmol, 1.5 eq.) prepared in Step 2 was added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (228 mg) (yield: 43%).

[388] Step 4. N-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-2,2,2-trifluoroacetamide


[389]



[390] (Z)-N-(4-((3-(4-(benzyloxy)phenyl)-1-(methylthio)-3-oxoprop-1-en-1-yl)amino)-3-methylphenyl)-2,2,2-trifluoroacetamide (220 mg, 0.45 mmol) prepared in Step 3, *t*-BuOH (2 mL) and AcOH (33 μL) were stirred in a round flask. Hydrazine hydrate (33 μL , 0.68 mmol, 1.5 eq.) was added and reacted under reflux for 20 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was filtered with MeOH to provide the compound (152 mg) (yield: 73%).

[391] Step 5.
2,2,2-trifluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide

[392]



[393] N-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-2,2,2-trifluoroacetamide (150 mg, 0.32 mmol) prepared in Step 4 and EtOAc/MeOH (3/3 mL) were stirred at ambient temperature in a round flask. Pd/C (30 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (yield: 49%).


[394] ¹H NMR (300 MHz, DMSO) δ 12.24 (s, 1H), 10.94 (s, 1H), 9.63 (s, 1H), 7.71 (s, 1H), 7.52 (d, J = 8.2 Hz, 2H), 7.37-7.31 (m, 3H), 6.81 (d, J = 8.1 Hz, 2H), 6.18 (s, 1H), 2.24 (s, 3H).

[395]

[396] Synthesis Example 9: Preparation of
1-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea (Compound **159**)

[397] Step 1. 1-(4-amino-3-methylphenyl)-3-methylurea

[398]




[399] Triphosgene (5.94 g, 20.0 mmol, 1.0 eq.) and THF were stirred at 0 °C in a 250 mL round flask. 3-methyl-4-nitroaniline (3.14 g, 20.0 mmol) in THF was added slowly and stirred at ambient temperature for 1 hour. The reactant was cooled in an ice bath, and methylamine (40% in MeOH, 1.5 mL, 1.0 eq.) was added slowly and stirred at ambient temperature for 24 hours. After the reaction ended, the reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. EtOAc/MeOH (100/100 mL) was added and then Pd/C (1.7 g, 20 wt%) was added, and the mixture was stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2.11 g) (yield: 59%).

[400] Step 2.

1-(4-((5-(4-((tert-butyldimethylsilyl)oxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea

[401]




[402] 1-(4-((tert-butyldimethylsilyl)oxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (709 mg, 2.0 mmol) and Toluene (15 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (49 μL , 0.3 mmol, 0.2 eq.) and 1-(4-amino-3-methylphenyl)-3-methylurea (538 mg, 3.0 mmol, 1.5 eq.) prepared in Step 1 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (10 mL), AcOH (113 μL) and Hydrazine hydrate (150 μL , 3.0 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. Then reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 775 mg) (yield: 86%).

[403] Step 3.

1-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea

[404]



[405] The Compound (774 mg, 1.71 mmol, 1.0 eq.) of Step 2 and THF were stirred at 0 °C in a 50 mL round flask. TBAF (2.24 g, 8.55 mmol) was added slowly and stirred at ambient temperature for 5 minutes. After the reaction ended, the reactant was extracted

with H₂O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (69 mg) (yield: 10%).


[406] ¹H NMR (300 MHz, Chloroform-*d*) δ 7.85 (d, *J* = 7.9 Hz, 2H), 7.35 (s, 1H), 7.23 (s, 1H), 7.12 (d, *J* = 8.4 Hz, 1H), 6.88 (d, *J* = 7.9 Hz, 2H), 6.76 (s, 1H), 6.13 (s, 1H), 3.78 (s, 3H), 2.21 (s, 3H).

[407]

[408] Synthesis Example 10: Preparation of
4-(3-((2-methyl-4-((2-morpholinoethyl)amino)phenyl)amino)-1H-pyrazol-5-yl)phenol
(Compound **164**)

[409] Step 1. 3-methyl-N1-(2-morpholinoethyl)benzene-1,4-diamine

[410]




[411] 5-fluoro-2-nitrotoluene (1.2 mL, 10.0 mmol) and THF were stirred in a 100 mL round flask. Trimethylamine (2.8 mL, 20.0 mmol, 2.0 eq.) and 4-(2-aminoethyl)morpholine (3.9 mL, 30.0 mmol, 3.0 eq.) were added and reacted under reflux for 48 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and DCM (50 mL x 3), and then the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. The produced compound (3.9 g) and EtOAc/MeOH/THF (50/70/70 mL) were added in a 250 mL round flask and then Pd/C (800 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2.14 g) (yield: 91%).

[412] Step 2.

N1-(5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)-2-methyl-N4-(2-morpholinoethyl)benzene-1,4-diamine

[413]




[414] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (660 mg, 2.0 mmol) and Toluene (10 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (49 μL , 0.3 mmol, 0.2 eq.) and the aniline (706 mg, 3.0 mmol, 1.5 eq.) prepared in Step 1 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (10 mL), AcOH (150 μL) and Hydrazine hydrate (150 μL , 3.0 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 460 mg) (yield: 48%).

[415] Step 3.

4-(3-((2-methyl-4-((2-morpholinoethyl)amino)phenyl)amino)-1H-pyrazol-5-yl)phenol

[416]



[417] N1-(5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)-2-methyl-N4-(2-morpholinoethyl)benzene-1,4-diamine (450 mg, 0.93 mmol) prepared in Step 2 and EtOAc/MeOH/THF (5/10/10 mL) were stirred at ambient temperature in a round flask. Pd/C (90 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (140 mg) (yield: 68%).

[418] ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.87 (s, 1H), 9.57 (s, 1H), 7.47 (d, J = 8.6 Hz,


2H), 7.19 (d, $J = 8.5$ Hz, 1H), 6.80 (d, $J = 3.1$ Hz, 2H), 6.76 (s, 1H), 6.44 (d, $J = 2.4$ Hz, 1H), 6.38 (dd, $J = 8.6, 2.6$ Hz, 1H), 5.82 (s, 1H), 4.85 (s, 1H), 3.67 - 3.52 (m, 4H), 3.08 (t, $J = 6.8$ Hz, 2H), 2.47 (s, 2H), 2.45 - 2.36 (m, 4H), 2.14 (s, 3H).

[419]

[420] Synthesis Example 11: Preparation of
1-(4-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)piperazin-1-yl)ethan-1-one (Compound **166**)

[421] Step 1. 1-(4-(4-amino-3-methylphenyl)piperazin-1-yl)ethan-1-one

[422]




[423] 5-fluoro-2-nitrotoluene (1.2 mL, 10.0 mmol) and THF (20 mL) were stirred in a 100 mL round flask. Et₃N (2.8 mL, 20.0 mmol, 2.0 eq.) and 1-acetylpiperazine (3.7 mL, 30.0 mmol, 3.0 eq.) were added and reacted under reflux for 48 hours. After the reaction ended, the reactant was cooled at ambient temperature, and then the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. EtOAc/MeOH/THF (50/50/50 mL) was added to the produced compound and the mixture was stirred at ambient temperature. Pd/C (600 mg, 20 wt%) was added and stirred at ambient temperature for 48 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (1.99 g) (yield: 85%).

[424] Step 2.

1-(4-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)piperazin-1-yl)ethan-1-one

[425]




[426] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (660 mg, 2.0 mmol) and Toluene (10 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (49 μL , 0.3 mmol, 0.2 eq.) and 1-(4-(4-amino-3-methylphenyl)piperazin-1-yl)ethan-1-one (700 mg, 3.0 mmol, 1.5 eq.) prepared in Step 1 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. EtOH (10 mL) and Hydrazine hydrate (150 μL , 3.0 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 277 mg) (yield: 29%).

[427] Step 3.

1-(4-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)piperazin-1-yl)ethan-1-one

[428]



[429] 1-(4-(4-((5-(4-(benzyloxy)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)piperazin-1-yl)ethan-1-one (270 mg, 0.56 mmol) prepared in Step 2 and EtOAc/MeOH/THF (5/10/10 mL) were stirred at ambient temperature in a round flask. Pd/C (60 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (153 mg) (yield: 70%).

[430] ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.06 (s, 1H), 9.65 (s, 1H), 7.49 (d, *J* = 8.7 Hz, 2H), 7.04 (s, 1H), 6.79 (d, *J* = 8.9 Hz, 3H), 6.70 (dd, *J* = 8.7, 2.8 Hz, 1H), 6.01 (s, 1H), 3.56 (q, *J* = 4.3 Hz, 4H), 2.97 (dt, *J* = 19.4, 5.1 Hz, 4H), 2.20 (s, 3H), 2.03 (s, 3H).

[431]


[432] Synthesis Example 12: Preparation of

4-(3-((2-methyl-4-(piperazin-1-yl)phenyl)amino)-1H-pyrazol-5-yl)phenol (Compound **169**)

[433] Step 1.

5-(4-(benzyloxy)phenyl)-N-(2-methyl-4-(piperazin-1-yl)phenyl)-1H-pyrazol-3-amine


[434]



- [435] 1-(4-(benzyloxy)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (660 mg, 2.0 mmol) and Toluene (10 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (49 μL , 0.3 mmol, 0.2 eq.) and 1-(4-(4-amino-3-methylphenyl)piperazin-1-yl)ethan-1-one (700 mg, 3.0 mmol, 1.5 eq.) prepared in Step 1 of Compound **166** synthesis were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (10 mL), AcOH (150 μL) and Hydrazine hydrate (150 μL , 3.0 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 377 mg) (yield: 43%).

- [436] Step 2. 4-(3-((2-methyl-4-(piperazin-1-yl)phenyl)amino)-1H-pyrazol-5-yl)phenol

[437]



- [438] 5-(4-(benzyloxy)phenyl)-N-(2-methyl-4-(piperazin-1-yl)phenyl)-1H-pyrazol-3-amine (241 mg, 0.55 mmol) prepared in Step 1 and EtOAc/MeOH/THF (10/10/10 mL) were

stirred at ambient temperature in a round flask. Pd/C (50 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite filter and the solvent was removed with a rotavapor. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (92 mg) (yield: 48%).


[439] ^1H NMR (300 MHz, DMSO- d_6) δ 12.08 (s, 1H), 9.63 (s, 1H), 8.82 (s, 2H), 7.49 (d, J = 8.6 Hz, 2H), 7.09 (s, 1H), 6.80 (d, J = 8.6 Hz, 3H), 6.73 (dd, J = 8.8, 2.9 Hz, 1H), 6.03 (s, 1H), 3.33 (s, 4H), 3.17 (d, J = 4.2 Hz, 4H), 2.21 (s, 3H).

[440]

[441] Synthesis Example 13: Preparation of
N1-(5-(4-nitrophenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine (Compound **21**)

[442] Step 1. 3,3-bis(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one

[443]




[444] 4-nitroacetophenone (2.4 g, 14.5 mmol) and benzene (73 mL) were stirred at ambient temperature in a round flask. The mixture was cooled in an ice bath, and then NaH (1.16 g, 29.0 mmol, 2.0 eq.) was added slowly and stirred at ambient temperature for 5 minutes. Carbon disulfide (1.3 mL, 21.8 mmol, 1.5 eq.) was added slowly and stirred at ambient temperature for 5 minutes. Iodomethane (2.7 mL, 43.5 mmol, 3.0 eq.) was then added slowly and stirred at ambient temperature for 5 minutes. *N,N*-dimethylacetamide (2.8 mL, 30.5 mmol, 2.1 eq.) was added slowly and stirred at ambient temperature for 3 hours. After the reaction ended, H₂O was added for quenching, and the reactant was extracted with EtOAc and then the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. Then the reactant was filtered with Et₂O to provide the compound (2.2 g) (yield: 56%).

[445] Step 2.

(*Z*)-3-((4-aminophenyl)amino)-3-(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one

[446]




[447] 3,3-bis(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one (269 mg, 1.0 mmol) prepared

in Step 1 and toluene (4 mL) were stirred in a round flask. $\text{BF}_3 \cdot \text{OEt}_2$ (25 μL , 0.2 mmol, 0.2 eq.) was added and 1,4-phenylenediamine (162 mg, 1.5 mmol, 1.5 eq.) was added, and then reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with MeOH to provide the compound (290 mg) (yield: 88%).

[448] Step 3. N1-(5-(4-nitrophenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine

[449]




[450] (Z)-3-((4-aminophenyl)amino)-3-(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one (290 mg, 0.88 mmol) prepared in Step 2, *t*-BuOH (10 mL) and AcOH (45 μL) were stirred in a round flask. Hydrazine hydrate (45 μL , 1.3 mmol, 1.5 eq.) was added and reacted under reflux for 20 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with MeOH to provide the compound (250 mg) (yield: 85%).

[451] $^1\text{H-NMR}$ (300 MHz, DMSO-d_6) δ 12.58 (s, 1H), 8.23-8.21 (m, 2H), 7.97-7.64 (m, 2H) 7.83 (s, 1H), 7.02 (s, 1H), 6.50-6.47 (m, 2H), 6.27 (s, 1H)

[452]

[453] Synthesis Example 14: Preparation of
N1-(5-(4-aminophenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine (Compound **22**)

[454]



[455] N1-(5-(4-nitrophenyl)-1H-pyrazol-3-yl)benzene-1,4-diamine (207 mg, 0.7 mmol) and THF/MeOH (5/5 mL) were stirred at ambient temperature in a 50 mL round flask. Pd/C (20 mg, 10 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, Pd/C was removed with a celite pad and the solvent was removed with a rotavapor. After that, silica gel column chromatography (DCM : MeOH) was performed to provide the compound (120 mg) (yield: 65%).


[456] $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 7.33 (d, 2H, $J=8.3$ Hz), 7.03 (d, 2H, $J=8.2$ Hz), 6.72-6.65 (m, 4H), 6.02 (s, 1H).

[457]

[458] Synthesis Example 15: Preparation of
4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methoxyphenol (Compound **34**)

[459] Step 1. 3-methoxy-4-nitrophenol


[460]



[461] MeOH (10 mL) and Na (368 mg, 2.0 eq.) were stirred in a 100 mL round flask. 3-fluoro-4-nitrophenol (1.26 g, 8.0 mmol) was added and reacted under reflux for 12 hours. After the reaction ended, the reactant was cooled at ambient temperature, and then the solvent was removed with a rotavapor. The reactant was extracted with 1N HCl (50ml) and DCM (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo* to provide the compound (1.23 g) (yield: 91%).

[462] Step 2. 4-amino-3-methoxyphenol


[463]



[464] 3-methoxy-4-nitrophenol (1.23 g, 7.2 mmol) prepared in Step 1 and EtOAc/THF (10/10 mL) were stirred at ambient temperature in a 100 mL round flask. Pd/C (240 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (410 mg) (yield: 41%).

[465] Step 3. 3-methoxy-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol


[466]



[467] 3,3-bis(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one (404 mg, 1.5 mmol) and Toluene (15 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (38 μL , 0.3 mmol, 0.2 eq.) and 4-amino-3-methoxyphenol (313 mg, 2.25 mmol, 1.5 eq.) prepared in Step 2 were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (10 mL), AcOH (113 μL) and Hydrazine hydrate (113 μL , 2.25 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50 mL) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 51 mg) (yield: 10%).

[468] Step 4. 4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methoxyphenol

[469]



[470] 3-methoxy-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol (50 mg, 0.15 mmol) prepared in Step 3 and MeOH/EtOAc (3/3 mL) were stirred at ambient temperature in a round flask. Pd/C (10 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (13 mg) (yield: 30%).


[471] ^1H NMR (300 MHz, Methanol- d_4) δ 7.40 (d, J = 8.2 Hz, 2H), 7.26 (d, J = 8.5 Hz, 1H), 6.73 (d, J = 8.2 Hz, 2H), 6.47 (d, J = 2.6 Hz, 1H), 6.35 (dd, J = 8.5, 2.6 Hz, 1H), 6.05 (s, 1H), 3.84 (s, 3H).

[472]

[473] Synthesis Example 16: Preparation of

4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol (Compound **24**)

[474]



[475] 3-methyl-4-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenol (88 mg, 0.28 mmol) and MeOH/THF/EtOAc (2/2/2 mL) were stirred at ambient temperature in a round


flask. Pd/C (9 mg, 10 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound as a brown solid (15 mg) (yield: 20%).

[476] $^1\text{H-NMR}$ (300 MHz, CD_3OD) δ 7.37 (d, 2H, $J=8.5$ Hz), 7.10 (d, 1H, $J=8.5$ Hz), 6.71 (d, 2H, $J=8.5$ Hz), 6.64-6.63 (m, 1H), 6.57 (dd, 1H, $J=8.6$, 2.8 Hz), 5.81 (s, 1H), 2.20 (s, 3H).

[477]

[478] Synthesis Example 17: Preparation of 4-fluoro-N-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)benzenesulfonamide (Compound **256**)

[479]




[480] Compound **24** (112 mg, 0.4 mmol) and DCM (3 mL) were stirred in a 100 mL round flask. Pyridine (97 μL , 1.2 mmol, 3.0 eq.) was added and the mixture was cooled in an ice bath. After that, 4-fluorobenzenesulfonyl chloride (79 mg, 0.4 mmol, 1.0 eq.) was added slowly and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and DCM (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound as a brown solid (80 mg) (yield: 46%).

[481] $^1\text{H NMR}$ (300 MHz, Chloroform- d) δ 8.08 (dd, $J = 8.6$, 5.0 Hz, 2H), 7.75 (dd, $J = 8.7$, 5.1 Hz, 2H), 7.59 (d, $J = 8.4$ Hz, 2H), 7.21 (t, $J = 8.5$ Hz, 2H), 7.12 - 7.00 (m, 4H), 6.75 (s, 1H), 6.72 - 6.62 (m, 2H), 5.48 (s, 1H), 4.98 (s, 1H), 2.21 (s, 3H).

[482]

[483] Synthesis Example 18: Preparation of 4-((4-fluoro-5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol (Compound **71**)

[484]




[485] (Z)-3-((4-hydroxy-2-methylphenyl)amino)-3-(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one (344 mg, 1.0 mmol) and CH_3CN (7 mL) were stirred in a 25 mL round flask. The mixture was cooled in an ice bath, and Selectfluor (373 mg, 1.0 mmol, 1.0 eq.) was slowly added and stirred for 30 minutes at 0 °C. After the reaction ended, H_2O (50 mL) was added and extracted with DCM (50 mL x 3). After that, the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *i*-PrOH (5 mL) and Hydrazine hydrate (50 μL , 1.0 mmol, 1.0 eq.) were added and stirred at 80 °C for 2 hours. After the reaction ended, the reactant was cooled at ambient temperature and the solvent was removed with a rotavapor. After that, the reactant was filtered with MeOH to provide the compound (2 steps, 158 mg) (yield: 48%).

[486] ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.81 (s, 1H), 8.35 (d, $J = 8.5$ Hz, 2H), 7.97 (d, $J = 8.5$ Hz, 2H), 7.04 (s, 1H), 6.73 (s, 1H), 6.56 (s, 1H), 6.47 (d, $J = 8.6$ Hz, 1H), 2.16 (s, 3H).

[487]

[488] Synthesis Example 19: Preparation of 4-((5-(4-aminophenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenol (Compound 56)

[489]



[490] Compound **71** (150 mg, 0.46 mmol) and EtOAc/MeOH/THF (5/5/5 mL) were stirred at ambient temperature in a round flask. Pd/C (30 mg, 20 wt%) was added and stirred

at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (58 mg) (yield: 42%).


[491] ^1H NMR (300 MHz, DMSO- d_6) δ 12.01 (s, 1H), 8.69 (s, 1H), 7.35 (d, J = 9.1 Hz, 2H), 6.86 (s, 1H), 6.68 (s, 1H), 6.63 (d, J = 8.2 Hz, 2H), 6.53 (s, 1H), 6.44 (d, J = 8.2 Hz, 1H), 5.36 (s, 2H), 2.16 (s, 3H).

[492]

[493] Synthesis Example 20: Preparation of N-(3-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)acetamide (Compound **84**)

[494] Step 1. tert-butyl acetyl(3-acetylphenyl)carbamate


[495]



[496] N-(3-acetylphenyl)acetamide (10 mmol, 1.77 g), (Boc) $_2$ O (1.5 eq, 3.2 g) and ACN (35 mL) were added in a 100 ml Round-bottom flask. Triethyl amine (2.0 eq, 2.8 mL) and DMAP (0.2 eq, 24 mg) were added to the mixture and stirred at 78 °C for 3 days. After the reaction ended, the solvent was removed under reduced pressure. The reactant was then extracted with H $_2$ O/EA three times, and the organic solvent layer was dried over MgSO $_4$ and concentrated *in vacuo*. After that, silica gel column chromatography was performed to provide the compound, tert-butyl acetyl(3-acetylphenyl)carbamate (1.63 g) (yield: 59%).

[497] Step 2. tert-butyl acetyl(3-(3,3-bis(methylthio)acryloyl)phenyl)carbamate

[498]




[499] Tert-butyl acetyl(3-acetylphenyl)carbamate (1.5g, 5.4 mmol), benzene (18 mL), and NaH (432 mg, 2.0 eq) were added in a 100 mL RBF at 0 °C and stirred at ambient temperature. After 5 minutes, carbon disulfide (0.67 mL, 1.5 eq) was added and stirred at

ambient temperature. After 5 minutes, iodomethane (1.03 mL, 3.0 eq) was added and stirred at ambient temperature. After 5 minutes, N,N-dimethylacetamide (1.05 mL, 2.1 eq) was added and stirred at ambient temperature for 3 hours. After the reaction ended, H₂O (60 mL) was added for quenching, and the reactant was extracted with EtOAc (100 mL x 3) and dried over MgSO₄ and concentrated *in vacuo*. The solvent was removed and silica gel column chromatography was performed to provide tert-butyl acetyl(3-(3,3-bis(methylthio)acryloyl)phenyl)carbamate (255 mg) (yield: 12%).

[500] Step 3. N-

(3-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)acetamide

[501]



[502] S.M. (0.65 mmol) and aniline derivative (1.5 eq) were dissolved in toluene (7 mL) in a 20 mL vial. Boron trifluoride diethyl etherate (0.2 eq) was added to the solution. The sealed vial was stirred at 120 °C for 6 hours. After the reaction ended, the solvent was removed under reduced pressure and the reactant was extracted with H₂O/EA three times. The organic solvent layer was dried over MgSO₄. The solvent was concentrated *in vacuo*. t-BuOH (5 mL) was added to the obtained intermediate, and hydrazine (1.5 eq) and acetic acid (1.5 eq) was added and stirred at 95 °C for 16 hours. After the reaction ended, the solvent was removed under reduced pressure and the reactant was extracted with H₂O/EA three times. The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. Then silica gel column chromatography was performed to provide N-(3-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)acetamide (11 mg) (yield: 5%).

[503] ¹H NMR (300 MHz, CDCl₃) δ 7.70 (s, 1H), 7.54 (dd, *J* = 5.5, 3.2 Hz, 1H), 7.14 (d, *J* = 8.6 Hz, 2H), 6.74 - 6.61 (m, 3H), 5.99 (s, 1H), 2.23 (s, 3H), 2.17 (s, 3H).


[504]

[505] Synthesis Example 21: Preparation of methyl

4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzoate (Compound **90**)

[506] Step 1. methyl 4-(3,3-bis(methylthio)acryloyl)benzoate

[507]




[508] 4-methylesteracetophenone (2.0 g, 11.2 mmol) and benzene (37 mL) were stirred at ambient temperature in a round flask. The mixture was cooled in an ice bath, and NaH (538 mg, 22.4 mmol, 2.0 eq.) was added slowly and stirred at ambient temperature for 5 minutes. Carbon disulfide (1.0 mL, 16.8 mmol, 1.5 eq.) was added slowly to the reactant and stirred at ambient temperature for 5 minutes. After that, iodomethane (2.1 mL, 33.7 mmol, 3.0 eq.) was added slowly to the reactant and stirred at ambient temperature for 5 minutes. *N, N*-dimethylacetamide (2.2 mL, 23.6 mmol, 2.1 eq.) was added slowly to the reactant and stirred at ambient temperature for 3 hours. After the reaction ended, H₂O was added for quenching. The reactant was extracted with EtOAc and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, the reactant was filtered with Et₂O to provide the compound (yield: 72%).

[509] Step 2. methyl

(*Z*)-4-(3-((4-hydroxy-2-methylphenyl)amino)-3-(methylthio)acryloyl)benzoate


[510]



[511] Methyl 4-(3,3-bis(methylthio)acryloyl)benzoate (1.7 g, 6.0 mmol) prepared in Step 1 and toluene (20 mL) were stirred at ambient temperature in a 250 mL round flask. Boron trifluoride (170 mg, 1.2 mmol, 0.2 eq) and 4-amino-3-methylphenol (1.1 g, 9.0 mmol, 1.5 eq) were added and stirred at 120 °C for 24 hours. After the reaction ended, toluene was removed with a rotavapor, and the reactant was extracted with *sat.* NaHCO₃ (30ml) and ethyl acetate (30 mL x 2). The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. Silica gel column chromatography (EtOAc : Hex = 1 : 9) was performed to provide the reactant. The reactant was sonicated in DCM and filtered to the compound (yield: 79 %).

[512] Step 3. methyl 4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzoate

[513]



[514] Methyl

(Z)-4-(3-((4-hydroxy-2-methylphenyl)amino)-3-(methylthio)acryloyl)benzoate (1.5 g, 4.3 mmol) prepared in Step 2, *t*-BuOH (13 mL), acetic acid (1.0 mL), and hydrazine hydrate (206 mg, 6.44 mmol, 1.5 eq.) were stirred at 95 °C for 18 hours in a 250 mL round flask. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (30 mL) and DCM (30 mL x 3), and was dried over MgSO₄ and concentrated *in vacuo*. Silica gel column chromatography (EtOAc : Hex = 1 : 4) was performed to provide the reactant. The reactant was sonicated in DCM and filtered to the compound (yield: 49 %).


[515] ¹H NMR (300 MHz, CDCl₃) δ 8.11 (s, 1H), 8.06 (d, *J* = 8.2 Hz, 2H), 7.98 (d, *J* = 8.2 Hz, 1H), 7.68 - 7.60 (m, 2H), 7.19 (d, *J* = 8.5 Hz, 1H), 6.71 (d, *J* = 2.9 Hz, 1H), 6.66 (dd, *J* = 8.5, 2.8 Hz, 1H), 6.13 (s, 1H), 5.53 (s, 1H), 3.93 (s, 3H), 2.23 (s, 3H)

[516]

[517] Synthesis Example 22: Preparation of

4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)benzoic acid (Compound **91**)

[518]



[519] Compound **90** (1.95 mmol, 630 mg) was stirred and dissolved in the solvent (H₂O:THF:EtOH = 2.0:1.0:2.0) in a 50 ml round bottom flask. Lithium hydroxide (9.74 mmol, 5.0 eq, 408.8 mg) was added and stirred at room temperature for 4 hours. After the reaction ended, solvent was removed for concentration. 30 ml of H₂O was added to the reactant, and 6 N HCl was added dropwise to adjust the pH to 1.0. The reactant was extracted with ethyl acetate (30ml x 3). The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo* to provide the compound as a white solid (yield: 13 %).

[520] ¹H NMR (300 MHz, DMSO) δ 12.67 (s, 1H), 8.81 (s, 1H), 8.01 (d, *J* = 12.4 Hz, 1H),

7.95 (d, $J = 7.9$ Hz, 2H), 7.81 (d, $J = 8.6$ Hz, 2H), 7.17 (d, $J = 8.3$ Hz, 1H), 7.10 (s, 1H), 6.58 (s, 1H), 6.52 (d, $J = 8.1$ Hz, 1H), 6.12 (s, 1H), 2.16 (s, 3H)

[521]


[522]

Synthesis Example 23: Preparation of
3-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-4-methylphenyl)methanol
(Compound **96**)

[523]

Step 1. (4-methyl-3-((5-(4-nitrophenyl)-1H-pyrazol-3-yl)amino)phenyl)methanol

[524]




[525]

3,3-bis(methylthio)-1-(4-nitrophenyl)prop-2-en-1-one (808 mg, 3.0 mmol) and Toluene (20 mL) were added in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (74 μL , 0.6 mmol, 0.2 eq.) and 3-amino-4-methylbenzylalcohol (636 mg, 4.5 mmol, 1.5 eq.) were added and reacted at 120 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. *t*-BuOH (25 mL), AcOH (150 μL) and Hydrazine hydrate (150 μL , 4.5 mmol, 1.5 eq.) were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and then the organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. Silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 200 mg) (yield: 21%).

[526]

Step 2. 3-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-4-methylphenyl)methanol

[527]



[528]

Compound **96** (200 mg, 0.68 mmol) and EtOAc/MeOH/THF (5/5/5 mL) were stirred

at ambient temperature in a round flask. Pd/C(40 mg, 20 wt%) was added and stirred at ambient temperature for 24 hours. After the reaction ended, the solvent was removed with a celite filter. Silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (63 mg) (yield: 31%).


[529] ¹H NMR (300 MHz, DMSO- *d*₆) δ 7.60 (s, 1H), 7.35 (d, *J* = 8.1 Hz, 2H), 7.16 (s, 1H), 6.99 (d, *J* = 7.3 Hz, 1H), 6.61 (t, *J* = 9.6 Hz, 3H), 6.08 (s, 1H), 5.28 (s, 2H), 4.98 (s, 1H), 4.37 (s, 2H), 2.20 (s, 3H).

[530]

[531] Synthesis Example 24: Preparation of N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-chloro propanamide (Compound **180**)

[532] Step 1. 1-(4-(1H-pyrazol-1-yl)phenyl)ethan-1-one


[533]



[534] 4'-iodoacetophenone (1.23 g, 5.0 mmol), pyrazole (0.68 g, 10.0 mmol, 2.0 eq), Cu(OAc)₂ (91 mg, 0.5 mmol, 0.1 eq.), Cs₂CO₃ (3.3 mg, 10.0 mmol, 2.0 eq.) and DMF (5 mL) were stirred at 110 °C in the sealed vial. The same reaction was repeated four times. After the reaction ended, all the reactant was combined and filtered with a celite pad. The filtrate was concentrated with a rotavapor and extracted with H₂O (50ml) and EtOAc (50 mL x 3). The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed and filtered with Et₂O and Hex to provide the compound (2.91 g) (yield: 78%).

[535] Step 2. 1-(4-(1H-pyrazol-1-yl)phenyl)-3,3-bis(methylthio)prop-2-en-1-one

[536]




[537] 1-(4-(1H-pyrazol-1-yl)phenyl)ethan-1-one (1.26 g, 6.8 mmol) and THF (25 mL) were stirred at ambient temperature in a 100 mL round flask. The mixture was cooled in an ice bath, and NaH (544 mg, 13.6 mmol, 2.0 eq.) was added slowly and stirred at ambient temperature for 5 minutes. CS₂ (620 μL, 10.2 mmol, 1.5 eq.) was added slowly and stirred at ambient temperature for 5 minutes. MeI (1.3 mL, 20.4 mmol, 3.0 eq.) was added slowly and stirred at 50 °C for 12 hours. After the reaction ended, H₂O

was added for quenching, and the reactant was extracted with EtOAc. The organic solvent layer was dried over MgSO_4 and concentrated *in vacuo*. After that, Et_2O was added to the reactant and sonicated and filtered to provide the compound (1.3 g) (yield: 66%).

[538] Step 3.

(Z)-1-(4-(1H-pyrazol-1-yl)phenyl)-3-((2-methyl-4-nitrophenyl)amino)-3-(methylthio)prop-2-en-1-one

[539]




[540] 1-(4-(1H-pyrazol-1-yl)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (871 mg, 3.0 mmol) prepared in Step 2 and toluene (15 mL) were stirred in a round flask. $\text{BF}_3 \cdot \text{OEt}_2$ (74 μL , 0.6 mmol, 0.2 eq.) was added and then 2-methyl-4-nitroaniline (685 mg, 4.5 mmol, 1.5 eq.) was added to the mixture. The mixture was stirred and reacted under reflux for 48 hours. After the reaction ended, the solvent was removed with a rotavapor and the reactant was filtered with MeOH to provide the compound (1.22 g) (yield: 77%).

[541] Step 4.

5-(4-(1H-pyrazol-1-yl)phenyl)-N-(2-methyl-4-nitrophenyl)-1H-pyrazol-3-amine

[542]




[543] (Z)-1-(4-(1H-pyrazol-1-yl)phenyl)-3-((2-methyl-4-nitrophenyl)amino)-3-(methylthio)prop-2-en-1-one (1.22 g, 3.1 mmol) prepared in Step 3, *t*-BuOH (30 mL) and AcOH (233 μL) were stirred in a round flask. Hydrazine hydrate (233 μL , 4.65 mmol, 1.5 eq.) was added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor and the reactant was filtered with MeOH to provide the compound (1.02 g) (yield: 91%).

[544] Step 5. tert-butyl

5-(4-(1H-pyrazol-1-yl)phenyl)-3-((tert-butoxycarbonyl)(2-methyl-4-nitrophenyl)amino)-1H-pyrazole-1-carboxylate

[545]




- [546] 5-(4-(1H-pyrazol-1-yl)phenyl)-N-(2-methyl-4-nitrophenyl)-1H-pyrazol-3-amine (1.51 g, 4.2 mmol) prepared in Step 4 and THF were stirred in a 250 mL round flask. DMAP (103 mg, 0.84 mmol, 0.2 eq.), Et₃N (1.8 mL, 12.6 mmol, 3.0 eq.), and (Boc)₂O (2.75 g, 12.6 mmol, 3.0 eq.) were added and stirred for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (1.54 g) (yield: 99%).

- [547] Step 6. tert-butyl

5-(4-(1H-pyrazol-1-yl)phenyl)-3-((4-amino-2-methylphenyl)(tert-butoxycarbonyl)amino)-1H-pyrazole-1-carboxylate

- [548]




- [549] tert-butyl

5-(4-(1H-pyrazol-1-yl)phenyl)-3-(((tert-butoxycarbonyl)(2-methyl-4-nitrophenyl)amino)-1H-pyrazole-1-carboxylate (1.54 g, 2.75 mmol) prepared in Step 5 and MeOH/THF (10/10 mL) were stirred at ambient temperature in a round flask. Pd/C(300 mg, 20 wt%) was added and stirred at ambient temperature for 48 hours. After the reaction ended, the solvent was removed with a celite filter. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (1.16 g) (yield: 79%).

- [550] Step 7. tert-butyl

5-(4-(1H-pyrazol-1-yl)phenyl)-3-(((tert-butoxycarbonyl)(4-(3-chloropropanamido)-2-methylphenyl)amino)-1H-pyrazole-1-carboxylate

- [551]




- [552] tert-butyl

5-(4-(1H-pyrazol-1-yl)phenyl)-3-((4-amino-2-methylphenyl)(tert-butoxycarbonyl)amino)-1H-pyrazole-1-carboxylate prepared in Step 6 and THF(15 mL) were stirred in a 100 mL round flask. 3-chloropropionyl chloride (600 μ L, 6.21 mmol, 3.0 eq.) was added and reacted at 50 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. Next step was performed without purification.

[553] Step 8. N-

(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-chloro propanamide

[554]



[555] tert-butyl

5-(4-(1H-pyrazol-1-yl)phenyl)-3-((tert-butoxycarbonyl)(4-(3-chloropropanamido)-2-methylphenyl)amino)-1H-pyrazole-1-carboxylate prepared in Step 7 and THF(10 mL) were stirred in a 100 mL round flask. Morpholine (450 μ L, 5.1 mmol, 3.0 eq.) was added and reacted at 50 °C for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50ml) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide both tert-butyl (5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)(4-(3-chloropropanamido)-2-methylphenyl)carbamate (360 mg) (yield: 41%) and N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-chloropropanamide (16 mg) (yield: 2%).

[556] ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.47 (s, 1H), 9.79 (s, 1H), 8.55 (d, *J* = 2.5 Hz, 1H), 7.96 - 7.74 (m, 5H), 7.60 (s, 1H), 7.39 - 7.24 (m, 3H), 6.57 (t, *J* = 2.1 Hz, 1H), 6.31 (s, 1H), 3.87 (t, *J* = 6.3 Hz, 2H), 2.77 (t, *J* = 6.3 Hz, 2H), 2.23 (s, 3H).

[557]


[558] Synthesis Example 25: Preparation of N-

(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide (Compound **181**)

[559] Step 1. tert-butyl

(5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)(2-methyl-4-(3-morpholinopropanamido)phenyl)carbamate

[560]




[561] tert-butyl

(5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)(4-(3-chloropropanamido)-2-methylphenyl)carbamate (360 mg, 0.69 mmol) obtained as a by-product in Step 8 of the preparation of Compound **180** and THF (10 mL) were stirred in a 100 mL round flask. Morpholine (180 μ L, 2.07 mmol, 3.0 eq.) was added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H₂O (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (DCM : MeOH) was performed to provide the compound (310 mg) (yield: 79%).

[562] Step 2. N-

(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide

[563]



[564] tert-butyl

(5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)(2-methyl-4-(3-morpholinopropanamido)phenyl)carbamate (300 mg, 0.52 mmol) prepared in Step 1 and DCM (5 mL) were stirred in a 100 mL round flask. The mixture was cooled in an ice bath, and TFA (1.2 mL, 15.6 mmol, 30.0 eq.) was added and stirred at room temperature for 30 hours. After the reaction ended, the reactant was extracted with H₂O (50 mL) and EtOAc (50 mL x 3), and the organic solvent layer was washed with sat. NaHCO₃. The reactant was dried over MgSO₄ and concentrated *in vacuo*. After that, the reactant was filtered with DCM to provide the compound (81 mg) (yield: 33%).

[565] ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.92 (s, 1H), 9.62 (s, 1H), 8.55 (d, *J* = 2.5 Hz, 1H), 7.91 (d, *J* = 8.8 Hz, 2H), 7.84 (d, *J* = 8.9 Hz, 2H), 7.77 (d, *J* = 1.7 Hz, 1H), 7.50 (d, *J* = 8.7 Hz, 1H), 7.35 (d, *J* = 2.4 Hz, 2H), 7.29 (dd, *J* = 8.6, 2.5 Hz, 1H), 6.60 - 6.54 (m, 1H), 6.32 (s, 1H), 4.00 (d, *J* = 12.8 Hz, 12H), 2.23 (s, 3H).


[566]

[567] Synthesis Example 26: Preparation of N-

(4-((5-(4-fluorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide

(Compound **204**)

[568]



[569] XPhos Pd G1 (5 mol%, 0.0113 mmol, 5.4 mg), XPhos (7 mol%, 0.0158 mmol, 11.6 mg), NaO-tBu (2.1 eq, 0.475 mmol, 45 mg), pyrazole amine (1.0 eq, 0.226 mol, 40 mg), N-(4-bromo-3-methylphenyl)methanesulfonamide (1.2 eq, 0.271 mmol, 71 mg), and degassed t-BuOH (4 mL) were added in a microwave vial. The vial was irradiated at 150 °C for 4 hours by microwave. The vial was cooled to room temperature and the solvent was removed under reduced pressure. The reactant was diluted with EA, and extracted with distilled water and NH₄Cl solution. The organic solvent layer was dried over Na₂SO₄ and filtered. The reactant was concentrated *in vacuo*, and separated by column chromatography to provide the compound as a brown liquid (11 mg) (yield: 13%).


[570]

[571] Synthesis Example 27: Preparation of 4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol (Compound **246**)

[572] Step 1.

1-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-3,3-bis(methylthio)prop-2-en-1-one

[573]




[574] 1-(3-iodophenyl)-3,3-bis(methylthio)prop-2-en-1-one (700 mg, 2.0 mmol), 2-fluoro-5-methoxyphenylboronic acid (340 mg, 2.0 mmol, 1.0 eq.), Pd(PPh₃)₂Cl₂ (264 mg, 0.6 mmol, 30 mol%), and K₂CO₃ (553 mg, 4.0 mmol, 2.0 eq.) were added to a mixture of H₂O (2 mL) and 1,4-dioxane (4 mL) in a round flask and stirred at 90 °C for 24 hours. After the reaction ended, the solvent was removed with a celite filter. The reactant was extracted with H₂O (30 mL) and ethyl acetate (30 mL x 3), and then dried over MgSO₄ and concentrated *in vacuo*. Then, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (297 mg) (yield: 43%).

[575] Step 2.

(Z)-1-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-3-((4-hydroxy-2-methylphenyl)amino)-3-(methylthio)prop-2-en-1-one


[576]



[577] 1-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-3,3-bis(methylthio)prop-2-en-1-one (297 mg, 0.88 mmol) prepared in Step 1 and toluene (9 mL) were stirred in a round flask. $\text{BF}_3 \cdot \text{OEt}_2$ (21 μL , 0.17 mmol, 0.2 eq.) was added, and then 4-amino-3-methylphenol (157 mg, 1.28 mmol, 1.5 eq.) was added to the mixture. The mixture was reacted under reflux for 24 hours. After the reaction ended, the reactant was extracted with H_2O (30 mL) and ethyl acetate (30 mL x 3), and dried over MgSO_4 and concentrated *in vacuo*. Then, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (133 mg) (yield: 37%).

[578] Step 3. 4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol

[579]



[580] (Z)-1-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-3-((4-hydroxy-2-methylphenyl)amino)-3-(methylthio)prop-2-en-1-one (133 mg, 0.31 mmol) prepared in Step 2, *t*-BuOH (3 mL) and AcOH (23 μL) were stirred in a round flask. Hydrazine hydrate (23 μL , 0.46 mmol, 1.5 eq.) was added and the mixture was reacted under reflux for 24 hours. After the reaction ended, the reactant was extracted with H_2O (30 mL) and ethyl acetate (30 mL x 3), and dried over MgSO_4 and concentrated *in vacuo*. Then, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (96 mg) (yield: 79%).

[581] ^1H NMR (300 MHz, Chloroform-*d*) δ 7.74 (q, J = 1.7 Hz, 1H), 7.58 (dt, J = 7.6, 1.5 Hz, 1H), 7.54 - 7.47 (m, 1H), 7.42 (t, J = 7.6 Hz, 1H), 7.13 (d, J = 8.4 Hz, 1H), 7.06 (dd, J = 9.9, 8.9 Hz, 1H), 6.94 (dd, J = 6.3, 3.1 Hz, 1H), 6.83 (dt, J = 8.9, 3.5 Hz, 1H),


6.67 - 6.58 (m, 2H), 6.07 (s, 1H), 3.80 (s, 3H), 2.13 (s, 3H).

[582]

[583] Synthesis Example 28: Preparation of
4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol
(Compound **274**)

[584] Step 1. 2-fluoro-5-methyl-4-nitrophenol


[585]



[586] 2-fluoro-5-methylphenol (4.3 mL, 39.0 mmol), AcOH (12 mL), and H₂SO₄ (1.7 mL) were stirred in a 25 mL round flask. The mixture was cooled in an ice bath. NaNO₂ (2.7 g, 39.0 mmol, 1.0 eq.) in H₂O (7.0 mL) was added slowly to the reactant and then stirred at room temperature for 1 hour. The prepared orange solid was filtered and put in a 25 mL round flask to which H₂O (19.0 mL) and HNO₃ (4.0 mL) were added. The mixture was stirred at 40 °C for 2 hours (until the solid became pale yellow). After the reaction ended, the prepared solid was filtered with H₂O to provide a beige solid (3.13 g) (yield: 47%).

[587] Step 2. 4-amino-2-fluoro-5-methylphenol

[588]




[589] 2-fluoro-5-methyl-4-nitrophenol (3.13 g) prepared in Step 1, THF/H₂O (30/30 mL), Fe (4.09 g, 73.2 mmol, 4.0 eq.), and NH₄Cl (3.92g, 73.2 mol, 4.0 eq.) were added in a 250 mL round flask and reacted under reflux for 4 hours. After the reaction ended, a celite filtration was performed and the reactant was extracted with H₂O and EtOAc. The organic solvent layer was dried over MgSO₄ and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed. The reactant was sonicated in Et₂O and filtered to provide a beige solid (984 mg) (yield: 38%).

[590] Step 3.

4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol

[591]



[592] 1-(4-(1H-pyrazol-1-yl)phenyl)-3,3-bis(methylthio)prop-2-en-1-one (436 mg, 1.5 mmol) and Toluene (15 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (37 μL , 0.3 mmol, 0.2 eq.) and 4-amino-2-fluoro-5-methylphenol (254 mg, 1.8 mmol, 1.2 eq.) prepared in Step 2 were added hereto and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor, and the reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3). The organic layer was dried over MgSO_4 and concentrated *in vacuo*. EtOH (15 mL) and Hydrazine hydrate (112 μL , 2.25 mmol, 1.5 eq.) were added hereto and the mixture was reacted under reflux. After the reaction ended, the solvent was removed with a rotavapor, and the reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3). The organic layer was dried over MgSO_4 and and concentrated *in vacuo*. After that, silica gel column chromatography (EtOAc : Hex) was performed to provide a white solid (2 steps, 65 mg) (yield: 12%).

[593] ^1H NMR (300 MHz, DMSO-d_6) δ 12.47 (s, 1H), 10.43 (s, 1H), 8.56 (d, $J = 2.5$ Hz, 1H), 8.42 (s, 1H), 7.92 (d, $J = 8.5$ Hz, 2H), 7.84 (d, $J = 8.5$ Hz, 2H), 7.77 (d, $J = 1.7$ Hz, 1H), 7.14 (s, 1H), 6.90 (d, $J = 6.9$ Hz, 1H), 6.74 (d, $J = 8.4$ Hz, 1H), 6.57 (t, $J = 2.1$ Hz, 1H), 6.24 (s, 1H), 4.49 (s, 2H).


[594]

[595] Synthesis Example 29: Preparation of 4-((5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol (Compound **277**)

[596] Step 1. tert-butyl

3-((tert-butoxycarbonyl)(4-hydroxy-2-methylphenyl)amino)-5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazole-1-carboxylate

[597]




[598] tert-butyl

5-(4-bromophenyl)-3-((tert-butoxycarbonyl)(4-hydroxy-2-methylphenyl)amino)-1H-pyrazole-1-carboxylate (100 mg, 0.18 mmol), 3,5-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (61 mg, 0.27 mmol, 1.5 eq.), Pd(PPh₃)₂Cl₂ (16 mg, 0.036 mmol, 20 mol%), and Na₂CO₃ (38 mg, 0.36 mmol, 2.0 eq.) were added to a mixture of EtOH (1 mL), H₂O (1 mL) and DMF (2 mL) in a round flask, and then stirred at 80 °C for 19 hours. After the reaction ended, the solvent was removed with a celite filter. The reactant was extracted with H₂O (30 mL) and ethyl acetate (30 mL x 3), and dried over MgSO₄ and concentrated *in vacuo*. Then, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (46 mg) (yield: 50%).

[599] Step 2.

4-((5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol

[600]



[601] tert-butyl

3-((tert-butoxycarbonyl)(4-hydroxy-2-methylphenyl)amino)-5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazole-1-carboxylate (46 mg, 0.08 mmol) prepared in Step 1 and DCM (1 mL) were stirred at room temperature in a round flask. The mixture was cooled in an ice bath. TFA (61 µL, 0.8 mmol, 10.0 eq.) was added to the mixture and stirred at room temperature for 5 hours. After the reaction ended, the reactant was extracted with H₂O (30 mL) and ethyl acetate (30 mL x 3). The organic layer was washed with sat. NaHCO₃ and then dried over MgSO₄ and concentrated *in vacuo*. Then, silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (17 mg) (yield: 59%).

[602] ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.85 (s, 1H), 7.70 - 7.54 (m, 4H), 7.05 (s, 1H),


6.58 (s, 1H), 6.52 (d, $J = 8.6$ Hz, 1H), 6.07 (s, 1H), 2.15 (s, 3H), 1.24 (s, 1H), 1.07 (s, 1H).

[603]

[604] Synthesis Example 30: Preparation of methyl
(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate
(Compound **286**)

[605] Step 1. methyl (3-methyl-4-nitrophenyl)carbamate


[606]



[607] 3-methyl-4-nitroaniline (15.2 g, 100.0 mmol) and DCM (500 mL) were stirred in a 1000 mL round flask. Pyridine (24.0 mL, 300.0 mmol, 3.0 eq.) was added to the mixture in an ice bath, and methyl chloroformate (9.3 mL, 120.0 mmol, 1.2 eq.) was added slowly and stirred at room temperature for 5 hours. After the reaction ended, H₂O was added and the reactant was extracted with EtOAc. The organic layer was dried over MgSO₄ and concentrated *in vacuo*. Then, the reactant was sonicated in Et₂O to provide the compound (20.0 g) (yield: 95%).

[608] Step 2. methyl (4-amino-3-methylphenyl)carbamate

[609]




[610] methyl (3-methyl-4-nitrophenyl)carbamate (20.0 g, 95.0 mmol) and THF/MeOH (250/250 mL), Pd/C (4 g, 20 wt%) were added in a hydrogenator flask, and reacted with a hydrogenator for 24 hours. After the reaction ended, the solvent was removed with a celite filter, and silica gel column chromatography (EtOAc : Hex) was performed to provide the brown oil (8.6 g) (yield: 50%).

[611] Step 3.

(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate

[612]



[613] 1-(4-methoxyphenyl)-3,3-bis(methylthio)prop-2-en-1-one (509 mg, 2.0 mmol) and Toluene (15 mL) were stirred in a MW vial. $\text{BF}_3 \cdot \text{OEt}_2$ (49 μL , 0.4 mmol, 0.2 eq.) and methyl (4-amino-3-methylphenyl)carbamate (433 mg, 2.4 mmol, 1.2 eq.) prepared in Step 2 were added and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3). The organic layer was dried over MgSO_4 and concentrated *in vacuo*. EtOH (15 mL) and Hydrazine hydrate (150 μL , 3.0 mmol, 1.5 eq.) were added to the reactant and reacted under reflux for 24 hours. After the reaction ended, the solvent was removed with a rotavapor. The reactant was extracted with H_2O (50ml) and EtOAc (50 mL x 3), and the organic layer was dried over MgSO_4 and concentrated *in vacuo*. Silica gel column chromatography (EtOAc : Hex) was performed to provide the compound (2 steps, 258 mg) (yield: 37%).

[614] ^1H NMR (300 MHz, DMSO-d_6) δ 12.22 (s, 1H), 8.92 (s, 1H), 7.62 (d, $J = 8.6$ Hz, 3H), 7.51 - 7.32 (m, 5H), 7.19 (s, 1H), 7.07 (d, $J = 8.4$ Hz, 2H), 6.69 (d, $J = 9.9$ Hz, 1H), 6.12 (s, 1H), 5.15 (s, 2H), 2.14 (s, 3H).

[615]

[616] The compounds of the present disclosure were synthesized using the method described above by adjusting starting materials and/or intermediates. Results of ^1H NMR tests are written in Table 3 below.

[617]

[Table 3]

Compound No.	¹ H NMR
1	¹ H-NMR (300 MHz, DMSO-d ₆) δ 7.48 (d, 2H, J=8.3 Hz), 7.33-7.30 (m, 2H), 7.19 (d, 2H, J=8.4 Hz), 7.07-6.86 (m, 6H), 6.68 (d, 2H, J=8.3 Hz), 6.17 (s, 1H), 5.98 (s, NH), 3.85 (s, 3H).
2	¹ H-NMR (300 MHz, DMSO-d ₆) δ 12.31 (s, 1H), 8.67 (s, 1H), 7.97 (s, 1H), 7.73 (d, 2H, J=9.0 Hz), 7.47 (d, 2H, J=6.0 Hz), 7.12 (s, 1H), 6.62 (d, 2H, J=6.0 Hz), 6.17 (s, 1H).
3	¹ H-NMR (300 MHz, DMSO-d ₆) δ 7.69 (d, 2H, J=9.0 Hz), 7.11 (d, 2H, J=9.0 Hz), 7.02 (d, 2H, J=9.0 Hz), 6.68 (d, 2H, J=9.0 Hz), 6.15 (s, 1H), 3.79 (s, 3H).
4	¹ H-NMR (300 MHz, CDCl ₃) δ 7.41-6.84 (m, 4H), 7.11 (d, 2H, J=8.9 Hz), 6.85 (d, 2H, J=8.7 Hz), 5.95 (s, 1H), 2.41 (s, 3H).
5	¹ H-NMR (300 MHz, CDCl ₃) δ 7.50 (d, 2H, J=8.2 Hz), 7.41 (d, 2H, J=8.5 Hz), 7.35 (d, 1H, J=8.5 Hz), 7.16-7.05 (m, 2H), 6.24 (s, 1H), 5.74 (s, NH), 2.26 (s, 3H).
6	¹ H-NMR (300 MHz, CDCl ₃) δ 7.49-7.47 (m, 2H), 7.40 (d, 1H, J=8.4 Hz), 7.13-7.09 (m, 2H), 6.96 (d, 2H, 8.6 Hz), 6.18 (s, 1H), 5.76 (s, NH), 3.85 (s, 3H), 2.25 (s, 3H).
7	¹ H-NMR (300 MHz, CDCl ₃) δ 7.42 (d, 1H, 8.5 Hz), 7.29 (s, 1H), 7.13-7.06 (m, 4H), 6.10 (s, 1H), 5.83 (s, NH), 2.40 (s, 3H), 2.36 (s, 3H), 2.26 (s, 3H).
8	¹ H-NMR (300 MHz, CDCl ₃) δ 7.51-7.48 (m, 2H), 7.41-7.26 (m, 3H), 6.93-6.84 (m, 2H), 6.15 (s, 1H), 5.30 (s, 1H), 2.27 (s, 3H).
9	¹ H-NMR (300 MHz, CDCl ₃) δ 7.50-7.46 (m, 2H), 7.38-7.27 (m, 1H), 6.94-6.85 (m, 4H), 6.11 (s, 1H), 5.64 (s, 1H), 3.84 (s, 3H), 2.25 (s, 3H).
10	¹ H-NMR (300 MHz, CDCl ₃) δ 7.40-7.35 (m, 2H), 7.32-7.21 (m, 3H), 6.91-6.80 (m, 2H), 6.05 (s, 1H), 5.73 (s, NH), 2.42 (s, 3H), 2.26 (s, 3H).
11	¹ H-NMR (300 MHz, DMSO-d ₆) δ 9.11 (s, 1H), 7.79 (d, 2H J=8.9 Hz), 7.66 (d, 2H, J=8.7 Hz), 7.38 (d, 2H, J=8.5 Hz), 7.00 (d, 2H, J=8.8 Hz), 6.23 (s, 1H), 4.23 (q, 2H, J=7.0 Hz), 3.78 (s, 3H), 1.28 (t, J=7.1 Hz).
12	¹ H-NMR (300 MHz, CDCl ₃) δ 7.48 (d, 2H, J=8.7 Hz), 7.21 (d, 1H, J=8.5 Hz), 6.92 (d, 2H, J=8.9 Hz), 6.69-6.61 (m, 1H), 6.00 (s, 1H), 3.83 (s, 3H), 2.21 (s, 3H).

13	¹ H-NMR (300 MHz, CDCl ₃) δ 6.77 (d, 2H, J=8.9 Hz), 6.20 (d, 2H, J=8.2 Hz), 6.15 (d, 2H, J=8.9 Hz), 5.90(d, 2H, J=8.3 Hz), 5.24 (s, 1H), 3.00 (s, 3H).
14	¹ H-NMR (300 MHz, DMSO-d ₆) δ 9.63 (s, 1H), 8.27 (s, 1H), 7.64 (d, 2H, J=8.3 Hz), 7.37 (d, 2H, 8.5 Hz), 7.28-7.25 (m, 2H), 6.99 (d, 2H, J=8.4 Hz), 6.11 (s, 1H), 3.79 (s, 3), 1.99 (s, 3H).
15	¹ H-NMR (300 MHz, DMSO-d ₆) δ 12.50 (s, 1H), 9.08 (s, 1H), 7.78 (d, 2H J=8.8 Hz), 7.65 (d, 2H, J=8.2 Hz), 7.39 (d, 2H, J=8.5 Hz), 6.99 (d, 2H, J=8.3 Hz), 6.21 (s, 1H), 3.75 (s, 3H).
16	¹ H-NMR (300 MHz, CDCl ₃) δ 7.51-7.48 (m, 2H), 7.08-7.02 (m, 1H), 6.97-6.94 (m, 2H), 6.57-6.56 (m, 1H), 6.53-6.50 (m, 1H), 6.26-6.22 (2H), 3.85 (s, 3H).
17	¹ H-NMR (300 MHz, CDCl ₃) δ 7.81 (s, 1H), 7.57-7.54 (m, 1H), 7.49 (d, 2H, J=8.7 Hz), 7.44-7.42 (m, 1H), 7.35-7.30 (m, 1H),), 6.96 (d, 2H, J= 8.7 Hz), 6.21 (s, 1H), 6.14 (s, 1H), 4.37 (q, 2H, J=7.1 Hz), 3.85 (s, 3H), 1.39 (t, 3H, J=7.1Hz)
18	¹ H-NMR (300 MHz, CDCl ₃) δ 12.14 (s, NH), 9.75 (s, 1H), 8.38 (s, 1H), 7.63 (d, 1H, J=8.4 Hz), 7.50 (s, 1H), 7.08-6.91 (m, 4H), 6.15 (s, 1H), 3.78 (s, 3H), 2.00 (s, 3H).
19	¹ H-NMR (300 MHz, CDCl ₃) δ 7.46 (d, J=8.5 Hz), 7.12 (t, 1H, J=7.5 Hz), 6.95 (d, 2H, J=8.6 Hz), 6.84-6.83 (m, 1H), 6.65-6.62 (m, 1H), 6.39-6.36 (m, 2H), 6.20 (s, 1H), 6.03-6.02 (m, 1H) 4.12 (q, 2H, J=7.2 Hz), 3.85 (s, 3H), 1.26 (t, 3H, J=7.1Hz)
20	¹ H-NMR (300 MHz, DMSO-d ₆) δ 8.68 (s, 1H), 8.06 (s, 1H), 7.65 (d, 2H, J=8.4 Hz), 7.53-7.50 (m, 1H), 7.31-7.27 (m, 2H), 7.00 (d, 2H, J=8.4 H), 6.14 (s, 1H), 3.78 (s, 3H).
21	¹ H-NMR (300 MHz, DMSO-d ₆) δ) 12.58 (s, 1H), 8.23-8.21 (m, 2H), 7.97-7.64 (m, 2H) 7.83 (s, 1H), 7.02 (s, 1H), 6.50-6.47 (m, 2H), 6.27 (s, 1H)
22	¹ H-NMR (300 MHz, CDCl ₃) δ 7.33 (d, 2H, J=8.3 Hz), 7.03 (d, 2H, J=8.2 Hz), 6.72-6.65 (m, 4H), 6.02 (s, 1H).
23	¹ H-NMR (300 MHz, CD ₃ OD) δ 8.21 (d, 2H, J=8.9 Hz), 7.86 (d, 2H, J=8.9 Hz), 7.07 (d, 2H, J=8.5 Hz), 6.66 (s, 1H), 6.60 (dd, 1H, 8.6, 2.8 Hz), 6.04 (s, 1H), 2.20 (s, 3H).
24	¹ H-NMR (300 MHz, CD ₃ OD) δ 7.37 (d, 2H, J=8.5 Hz), 7.10 (d, 1H, J=8.5 Hz), 6.71 (d, 2H, J=8.5 Hz), 6.64-6.63 (m, 1H), 6.57 (dd, 1H, J=8.6, 2.8 Hz), 5.81 (s, 1H), 2.20 (s, 3H).

25	¹ H-NMR (300 MHz, CDCl ₃) δ 8.27 (d, 2H, J=8.6 Hz), 7.77 (d, 2H, J=8.6 Hz), 7.04 (d, 2H, J=8.5 Hz), 6.81 (d, 2H, J=8.5 Hz).
26	¹ H-NMR (300 MHz, DMSO-d ₆) δ 12.32 (s, NH), 8.60 (s, 1H), 7.62 (d, 2H, J=8.7 Hz), 7.40-7.33 (m, 3H), 7.20-6.95 (m, 9H), 6.36-6.29 (m, 2H), 6.17-6.09 (m, 2H), 5.21 (s, NH), 3.78 (s, 3H).
27	¹ H NMR (300 MHz, DMSO- d ₆) δ 12.27 (s, 1H), 8.87 (s, 1H), 7.71 (d, J = 8.3 Hz, 2H), 7.48 (d, J = 8.2 Hz, 2H), 7.04 (s, 1H), 6.43 (s, 1H), 6.29 (d, J = 8.4 Hz, 2H), 3.79 (s, 3H).
28	¹ H NMR (300 MHz, DMSO- d ₆) δ 12.37 (s, 1H), 8.80 (s, 1H), 7.73 (d, J = 8.1 Hz, 2H), 7.47 (d, J = 8.0 Hz, 2H), 6.77 (s, 1H), 6.43 (d, J = 2.7 Hz, 1H), 6.35 - 6.23 (m, 2H), 4.51 (p, J = 6.0 Hz, 1H), 1.30 (d, J = 6.0 Hz, 6H).
29	¹ H NMR (300 MHz, Methanol- d ₄) δ 7.65 (d, J = 8.1 Hz, 2H), 7.40 (d, J = 8.3 Hz, 2H), 7.02 (s, 2H), 6.67 (d, J = 8.2 Hz, 2H), 6.12 (s, 1H), 3.01 (t, J = 7.5 Hz, 2H), 1.62 (q, J = 7.2 Hz, 2H), 0.99 (t, J = 7.4 Hz, 3H).
30	¹ H NMR (300 MHz, DMSO- d ₆) δ 12.43 (s, 1H), 8.28 (s, 1H), 7.80 - 7.67 (m, 3H), 7.51 (s, 2H), 7.29 (s, 1H), 7.13 (t, J = 7.6 Hz, 2H), 6.99 (d, J = 8.2 Hz, 2H), 6.88 (d, J = 8.0 Hz, 2H), 6.66 (t, J = 7.3 Hz, 1H), 6.26 (s, 1H).
31	¹ H NMR (300 MHz, Chloroform- d) δ 7.50 (d, J = 8.7 Hz, 2H), 7.34 (t, J = 7.9 Hz, 2H), 7.12 (dt, J = 7.4, 3.1 Hz, 2H), 6.99 (dd, J = 13.0, 8.2 Hz, 4H), 6.64 - 6.52 (m, 3H), 5.98 (s, 1H), 2.13 (s, 3H).
32	¹ H NMR (300 MHz, DMSO- d ₆) δ 12.23 (s, 1H), 8.81 (s, 1H), 7.65 (d, J = 8.4 Hz, 2H), 7.58 (d, J = 8.3 Hz, 2H), 7.08 (s, 1H), 6.57 (d, J = 2.9 Hz, 1H), 6.51 (dd, J = 8.6, 2.7 Hz, 1H), 6.05 (s, 1H), 2.14 (s, 3H).
33	¹ H NMR (300 MHz, DMSO- d ₆) δ 12.52 (s, 1H), 12.18 (s, 1H), 8.84 (s, 1H), 7.87 (d, J = 7.3 Hz, 4H), 7.17 (s, 1H), 6.61 - 6.47 (m, 2H), 6.17 (s, 1H), 2.15 (s, 3H).
34	¹ H NMR (300 MHz, Methanol- d ₄) δ 7.40 (d, J = 8.2 Hz, 2H), 7.26 (d, J = 8.5 Hz, 1H), 6.73 (d, J = 8.2 Hz, 2H), 6.47 (d, J = 2.6 Hz, 1H), 6.35 (dd, J = 8.5, 2.6 Hz, 1H), 6.05 (s, 1H), 3.84 (s, 3H).
35	¹ H NMR (300 MHz, Methanol- d ₄) δ 7.41 (d, J = 8.6 Hz, 2H), 7.24 (d, J = 8.6 Hz, 1H), 6.74 (d, J = 8.6 Hz, 2H), 6.48 (d, J = 2.6 Hz, 1H), 6.36 (dd, J = 8.5, 2.6 Hz, 1H), 6.09 (s, 1H), 4.55 (p, J = 6.1 Hz, 1H), 1.36 (d, J = 6.0 Hz, 6H).
36	¹ H NMR (300 MHz, DMSO- d ₆) δ 11.79 (s, 1H), 7.67 (s, 1H), 7.33 (d, J = 8.2 Hz, 2H), 7.09 (d, J = 8.4 Hz, 2H), 6.57 (d, J = 8.3 Hz, 2H), 6.47 (d, J = 8.4 Hz, 2H), 5.85 (s, 1H), 5.26 (s, 2H), 4.88 (s, 1H), 2.90 (t, J = 7.0 Hz, 2H), 1.54 (q, J

	= 7.2 Hz, 2H), 0.93 (t, $J = 7.4$ Hz, 3H).
37	^1H NMR (300 MHz, Methanol- d_4) δ 7.42 (d, $J = 8.6$ Hz, 2H), 7.21 - 7.07 (m, 5H), 7.02 (d, $J = 8.8$ Hz, 2H), 6.92 (d, $J = 7.8$ Hz, 2H), 6.77 - 6.71 (m, 2H), 6.06 (s, 1H).
38	^1H NMR (300 MHz, DMSO- d_6) δ 11.85 (s, 1H), 8.63 (s, 1H), 7.86 (s, 1H), 7.34 (d, $J = 8.4$ Hz, 2H), 7.14 (d, $J = 8.3$ Hz, 2H), 6.59 (t, $J = 8.9$ Hz, 4H), 5.88 (s, 1H), 5.27 (s, 2H).
39	^1H NMR (500 MHz, DMSO- d_6) δ 11.98 (s, 1H), 9.58 (s, 1H), 7.49 (d, $J = 8.5$ Hz, 2H), 6.90 (s, 1H), 6.79 (d, $J = 8.5$ Hz, 2H), 6.56 (d, $J = 2.8$ Hz, 1H), 6.50 (dd, $J = 8.5, 2.9$ Hz, 1H), 5.91 (s, 1H), 2.15 (s, 3H).
40	^1H NMR (500 MHz, DMSO- d_6) δ 12.02 (s, 1H), 9.62 (s, 1H), 8.63 (s, 1H), 7.90 (s, 1H), 7.51 (d, $J = 8.3$ Hz, 2H), 7.17 (s, 1H), 6.81 (d, $J = 8.3$ Hz, 2H), 6.65 - 6.59 (m, 2H), 5.98 (s, 1H).
41	^1H NMR (300 MHz, DMSO- d_6) δ 12.05 (s, 1H), 8.75 (s, 1H), 7.31 (s, 1H), 7.01 (dd, $J = 17.9, 10.2$ Hz, 2H), 6.86 - 6.77 (m, 2H), 6.55 (s, 1H), 6.53 - 6.46 (m, 2H), 5.87 (s, 1H), 5.11 (s, 2H), 2.14 (s, 3H).
42	^1H NMR (300 MHz, DMSO- d_6) δ 12.14 (s, 1H), 8.79 (s, 1H), 7.78 - 7.67 (m, 2H), 7.24 (t, $J = 8.7$ Hz, 3H), 7.03 (s, 1H), 6.57 (d, $J = 2.8$ Hz, 1H), 6.51 (dd, $J = 8.6, 2.7$ Hz, 1H), 6.03 (s, 1H), 2.15 (s, 3H).
43	^1H NMR (300 MHz, DMSO- d_6) δ 11.89 (s, 1H), 9.14 (s, 1H), 7.74 (s, 1H), 7.52 (s, 1H), 7.32 (d, $J = 8.4$ Hz, 2H), 6.62 - 6.52 (m, 3H), 6.49 (dd, $J = 8.9, 2.7$ Hz, 1H), 5.92 (s, 1H), 5.28 (s, 2H).
44	^1H NMR (300 MHz, DMSO- d_6) δ 11.96 (s, 1H), 8.90 (s, 1H), 8.18 (s, 1H), 7.35 (d, $J = 8.2$ Hz, 3H), 6.85 (d, $J = 9.3$ Hz, 1H), 6.77 (t, $J = 9.2$ Hz, 1H), 6.58 (d, $J = 8.3$ Hz, 2H), 5.89 (s, 1H), 5.30 (s, 2H), 3.35 (s, 3H), 2.08 (s, 3H).
45	^1H NMR (300 MHz, DMSO- d_6) δ 12.29 (s, 1H), 8.78 (s, 1H), 7.78 (d, $J = 7.9$ Hz, 2H), 7.71 (d, $J = 8.1$ Hz, 4H), 7.47 (t, $J = 7.5$ Hz, 2H), 7.38 (d, $J = 7.1$ Hz, 1H), 7.05 (s, 1H), 6.57 (s, 1H), 6.52 (d, $J = 7.3$ Hz, 1H), 6.11 (s, 1H), 2.17 (s, 3H).
46	^1H NMR (300 MHz, DMSO- d_6) δ 12.31 (s, 1H), 8.79 (s, 1H), 7.79 (d, $J = 8.2$ Hz, 2H), 7.73 (s, 2H), 7.67 (t, $J = 10.0$ Hz, 5H), 7.29 (s, 1H), 7.05 (s, 1H), 6.57 (s, 1H), 6.52 (dd, $J = 8.8, 2.4$ Hz, 1H), 6.11 (s, 1H), 2.16 (s, 3H).
47	^1H NMR (300 MHz, DMSO- d_6) δ 12.27 (s, 1H), 8.92 (s, 1H), 8.82 (s, 1H), 8.48 (d, $J = 3.7$ Hz, 1H), 8.06 (d, $J = 8.1$ Hz, 1H), 7.42 (t, $J = 6.5$ Hz, 1H), 7.13 (s, 1H), 6.57 (s, 1H), 6.52 (d, $J = 8.8$ Hz, 1H), 6.13 (s, 1H), 2.15 (s, 3H).

48	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.01 (s, 1H), 9.69 (s, 1H), 8.33 (s, 1H), 7.76 - 7.65 (m, 2H), 7.37 (t, <i>J</i> = 9.2 Hz, 4H), 7.19 (d, <i>J</i> = 8.4 Hz, 2H), 6.83 (d, <i>J</i> = 8.4 Hz, 2H), 6.58 (d, <i>J</i> = 8.2 Hz, 2H), 5.93 (s, 1H), 5.29 (s, 2H).
49	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.04 (s, 1H), 9.62 (s, 1H), 9.17 (s, 1H), 7.71 (s, 1H), 7.58 (s, 1H), 7.48 (d, <i>J</i> = 8.2 Hz, 2H), 6.80 (d, <i>J</i> = 8.3 Hz, 2H), 6.57 (d, <i>J</i> = 13.1 Hz, 1H), 6.50 (dd, <i>J</i> = 9.1, 2.0 Hz, 1H), 6.00 (s, 1H).
50	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.03 (s, 1H), 9.12 (s, 1H), 8.37 (s, 1H), 7.36 (d, <i>J</i> = 8.0 Hz, 2H), 7.31 (d, <i>J</i> = 7.8 Hz, 2H), 7.04 (d, <i>J</i> = 8.9 Hz, 2H), 6.58 (d, <i>J</i> = 8.9 Hz, 2H), 5.97 (s, 1H), 5.30 (s, 2H), 2.86 (s, 3H).
51	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.19 (s, 1H), 9.64 (s, 1H), 9.13 (s, 1H), 8.42 (s, 1H), 7.52 (d, <i>J</i> = 8.3 Hz, 2H), 7.30 (d, <i>J</i> = 8.4 Hz, 2H), 7.05 (d, <i>J</i> = 8.5 Hz, 2H), 6.81 (d, <i>J</i> = 8.3 Hz, 2H), 6.06 (s, 1H), 2.86 (s, 3H).
52	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.17 (s, 1H), 9.66 (s, 2H), 8.37 (s, 1H), 7.76 - 7.65 (m, 2H), 7.51 (d, <i>J</i> = 8.3 Hz, 2H), 7.37 (t, <i>J</i> = 8.7 Hz, 2H), 7.19 (d, <i>J</i> = 8.4 Hz, 2H), 6.89 - 6.75 (m, 4H), 6.02 (s, 1H).
53	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.14 (s, 1H), 9.64 (s, 2H), 8.27 (s, 1H), 7.52 (d, <i>J</i> = 8.3 Hz, 2H), 7.35 (s, 2H), 7.26 (s, 2H), 6.81 (d, <i>J</i> = 7.4 Hz, 2H), 6.04 (s, 1H), 2.09 (s, 3H).
54	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 11.98 (s, 1H), 9.63 (s, 1H), 8.23 (s, 1H), 7.36 (d, <i>J</i> = 8.4 Hz, 4H), 7.25 (d, <i>J</i> = 8.3 Hz, 2H), 6.58 (d, <i>J</i> = 8.2 Hz, 2H), 5.95 (s, 1H), 5.29 (s, 2H), 1.98 (s, 3H).
55	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.05 (s, 1H), 9.13 (s, 1H), 7.62 (s, 1H), 7.35 (d, <i>J</i> = 8.3 Hz, 2H), 7.23 (s, 1H), 6.93 (d, <i>J</i> = 8.8 Hz, 2H), 6.58 (d, <i>J</i> = 8.4 Hz, 2H), 6.06 (s, 1H), 5.29 (s, 2H), 2.86 (s, 3H), 2.21 (s, 3H).
56	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.01 (s, 1H), 8.69 (s, 1H), 7.35 (d, <i>J</i> = 9.1 Hz, 2H), 6.86 (s, 1H), 6.68 (s, 1H), 6.63 (d, <i>J</i> = 8.2 Hz, 2H), 6.53 (s, 1H), 6.44 (d, <i>J</i> = 8.2 Hz, 1H), 5.36 (s, 2H), 2.16 (s, 3H).
57	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.03 (s, 1H), 9.71 (s, 1H), 7.73 (dd, <i>J</i> = 8.6, 5.1 Hz, 2H), 7.50 (s, 1H), 7.43 - 7.29 (m, 4H), 7.17 (s, 1H), 6.76 (s, 1H), 6.74 - 6.68 (m, 1H), 6.58 (d, <i>J</i> = 8.2 Hz, 2H), 6.02 (s, 1H), 5.28 (s, 2H), 2.12 (s, 3H).
58	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.21 (s, 1H), 9.63 (s, 1H), 9.17 (s, 1H), 7.68 (s, 1H), 7.51 (d, <i>J</i> = 8.5 Hz, 2H), 7.29 (s, 1H), 6.94 (d, <i>J</i> = 8.9 Hz, 2H), 6.81 (d, <i>J</i> = 8.5 Hz, 2H), 6.15 (s, 1H), 2.86 (s, 3H), 2.22 (s, 3H).
59	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.14 (s, 1H), 9.62 (s, 1H), 7.51 (d, <i>J</i> = 8.2

	Hz, 2H), 7.30 (s, 1H), 7.27 - 7.15 (m, 2H), 6.80 (d, $J = 8.2$ Hz, 2H), 6.10 (s, 1H), 2.20 (s, 3H), 1.98 (s, 3H).
66	^1H NMR (300 MHz, DMSO- d_6) δ 12.13 (s, 1H), 8.85 (s, 1H), 8.74 (d, $J = 2.5$ Hz, 1H), 8.13 (dd, $J = 8.3, 2.5$ Hz, 1H), 7.53 (s, 1H), 7.18 (s, 1H), 6.57 (s, 1H), 6.52 (d, $J = 9.0$ Hz, 1H), 6.11 (s, 1H), 2.14 (s, 3H).
67	^1H NMR (300 MHz, Acetone- d_6) δ 8.24 (d, $J = 8.7$ Hz, 2H), 7.99 (d, $J = 8.7$ Hz, 2H), 7.61 (t, $J = 9.6$ Hz, 1H), 7.19 (s, 1H), 6.68 - 6.54 (m, 2H), 6.42 (s, 1H).
68	^1H NMR (300 MHz, Methanol- d_4) δ 8.22 (d, $J = 8.9$ Hz, 2H), 7.86 (d, $J = 9.0$ Hz, 2H), 7.03 (d, $J = 12.2$ Hz, 1H), 6.85 - 6.75 (m, 2H), 6.31 (s, 1H).
69	^1H NMR (300 MHz, DMSO- d_6) δ 12.79 (s, 1H), 9.66 (s, 1H), 8.54 (s, 1H), 8.28 (d, $J = 8.5$ Hz, 2H), 8.01 (d, $J = 8.5$ Hz, 2H), 7.72 (dd, $J = 8.6, 5.2$ Hz, 2H), 7.38 (t, $J = 8.7$ Hz, 2H), 7.14 (s, 2H), 6.88 (d, $J = 8.4$ Hz, 2H), 6.47 (s, 1H).
70	^1H NMR (300 MHz, DMSO- d_6) δ 12.74 (s, 1H), 9.22 (s, 1H), 8.28 (d, $J = 8.5$ Hz, 2H), 8.01 (d, $J = 8.5$ Hz, 2H), 7.53 (s, 1H), 6.97 (d, $J = 11.9$ Hz, 2H), 6.50 (s, 1H), 2.88 (s, 3H), 2.23 (s, 3H).
71	^1H NMR (300 MHz, DMSO- d_6) δ 8.81 (s, 1H), 8.35 (d, $J = 8.5$ Hz, 2H), 7.97 (d, $J = 8.5$ Hz, 2H), 7.04 (s, 1H), 6.73 (s, 1H), 6.56 (s, 1H), 6.47 (d, $J = 8.6$ Hz, 1H), 2.16 (s, 3H).
72	^1H NMR (300 MHz, DMSO- d_6) δ 12.74 (s, 1H), 9.84 (s, 1H), 8.27 (d, $J = 8.5$ Hz, 2H), 7.99 (d, $J = 8.4$ Hz, 2H), 7.81 - 7.69 (m, 2H), 7.46 (s, 1H), 7.38 (t, $J = 8.8$ Hz, 2H), 6.82 (s, 1H), 6.76 (d, $J = 8.1$ Hz, 1H), 6.48 (s, 1H), 2.14 (s, 3H).
73	^1H NMR (500 MHz, DMSO- d_6) δ 12.34 (s, 1H), 8.76 (s, 1H), 8.32 (s, 1H), 7.82 (d, $J = 11.3$ Hz, 4H), 7.72 (s, 2H), 7.13 (s, 2H), 6.58 (s, 2H), 6.53 (s, 1H), 6.15 (s, 1H), 2.17 (d, $J = 2.9$ Hz, 4H).
74	^1H NMR (500 MHz, DMSO- d_6) δ 12.38 (s, 1H), 8.70 (s, 1H), 8.33 (d, $J = 1.2$ Hz, 1H), 8.00 (s, 1H), 7.86 (d, $J = 8.5$ Hz, 2H), 7.81 (t, $J = 1.3$ Hz, 1H), 7.73 (d, $J = 8.3$ Hz, 2H), 7.13 (t, $J = 1.2$ Hz, 1H), 6.68 - 6.62 (m, 2H), 6.24 (s, 1H).
75	^1H NMR (500 MHz, DMSO- d_6) δ 12.36 (s, 1H), 8.66 (s, 1H), 8.01 (s, 1H), 7.81 (d, $J = 8.0$ Hz, 2H), 7.79 - 7.71 (m, 4H), 7.49 (t, $J = 7.7$ Hz, 2H), 7.39 (t, $J = 7.3$ Hz, 1H), 7.20 (d, $J = 8.1$ Hz, 2H), 6.65 (d, $J = 8.4$ Hz, 2H), 6.21 (s, 1H).
76	^1H NMR (500 MHz, DMSO- d_6) δ 12.37 (s, 1H), 8.67 (s, 1H), 8.01 (s, 1H), 7.82 (d, $J = 8.0$ Hz, 2H), 7.76 (d, $J = 8.3$ Hz, 2H), 7.72 - 7.65 (m, 4H), 7.20 (s, 1H), 6.65 (d, $J = 8.4$ Hz, 2H), 6.22 (s, 1H).

80	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 11.93 (s, 1H), 8.71 (s, 1H), 7.54 - 7.46 (m, 2H), 7.33 (s, 1H), 6.87 (s, 1H), 6.77 - 6.71 (m, 2H), 6.56 (d, <i>J</i> = 2.9 Hz, 1H), 6.50 (dd, <i>J</i> = 8.5, 2.9 Hz, 1H), 5.90 (s, 1H), 2.93 (s, 6H).
81	¹ H NMR (500 MHz, DMSO) δ 12.01 - 11.82 (m, 1H), 9.57 (s, 1H), 8.78 (s, 1H), 7.48 (d, <i>J</i> = 8.6 Hz, 2H), 7.25 (s, 1H), 6.89 (s, 1H), 6.78 (d, <i>J</i> = 8.6 Hz, 2H), 6.58 (d, <i>J</i> = 2.7 Hz, 1H), 6.51 (dd, <i>J</i> = 8.6, 2.8 Hz, 1H), 5.87 (s, 1H), 2.56 (q, <i>J</i> = 7.5 Hz, 2H), 1.13 (t, <i>J</i> = 7.5 Hz, 3H).
82	¹ H NMR (300 MHz, DMSO) δ 12.06 - 11.76 (m, 1H), 9.46 (s, 1H), 8.62 (s, 1H), 7.86 (s, 1H), 7.38 (s, 1H), 7.30 (dd, <i>J</i> = 8.3, 2.4 Hz, 1H), 7.11 (d, <i>J</i> = 8.6 Hz, 2H), 6.77 (d, <i>J</i> = 8.4 Hz, 1H), 6.59 (d, <i>J</i> = 8.6 Hz, 2H), 5.93 (s, 1H), 2.13 (s, 3H).
83	¹ H NMR (300 MHz, CD ₃ OD) δ 7.44 - 7.26 (m, 4H), 7.14 (d, <i>J</i> = 8.5 Hz, 2H), 6.77 (d, <i>J</i> = 8.4 Hz, 1H), 6.10 (s, 1H), 2.22 (s, 3H), 2.08 (s, 3H).
84	¹ H NMR (300 MHz, CDCl ₃) δ 7.70 (s, 1H), 7.54 (dd, <i>J</i> = 5.5, 3.2 Hz, 1H), 7.14 (d, <i>J</i> = 8.6 Hz, 2H), 6.74 - 6.61 (m, 3H), 5.99 (s, 1H), 2.23 (s, 3H), 2.17 (s, 3H).
85	¹ H NMR (300 MHz, DMSO) δ 11.87 (s, 1H), 9.41 (s, 1H), 8.67 (s, 1H), 7.33 (s, 1H), 7.32 (t, <i>J</i> = 9.2 Hz, 2H), 6.83 (s, 1H), 6.73 (d, <i>J</i> = 8.3 Hz, 1H), 6.54-6.39 (m, 1H), 5.83 (s, 1H), 2.09 (s, 6H).
86	¹ H NMR (500 MHz, DMSO) δ 12.13 - 12.04 (m, 1H), 9.63 (s, 1H), 9.50 (s, 1H), 7.51 (s, 1H), 7.43 (d, <i>J</i> = 2.2 Hz, 1H), 7.34 (dd, <i>J</i> = 8.3, 2.3 Hz, 1H), 7.31 (d, <i>J</i> = 2.4 Hz, 1H), 7.24 (dd, <i>J</i> = 8.7, 2.4 Hz, 1H), 7.17 (s, 1H), 6.81 (d, <i>J</i> = 8.3 Hz, 1H), 6.09 (s, 1H), 2.21 (s, 3H), 2.16 (s, 3H), 1.99 (s, 3H).
87	¹ H NMR (300 MHz, CD ₃ OD) δ 7.20 (t, <i>J</i> = 7.8 Hz, 1H), 7.11 (d, <i>J</i> = 7.9 Hz, 2H), 7.07 (s, 1H), 6.75 (d, <i>J</i> = 8.5 Hz, 1H), 6.66 (s, 1H), 6.60 (dd, <i>J</i> = 8.5, 2.7 Hz, 1H), 5.92 (s, 3H), 2.21 (s, 3H)
88	¹ H NMR (300 MHz, CD ₃ OD) δ 7.35 (d, <i>J</i> = 12.3 Hz, 1H), 7.26 (d, <i>J</i> = 8.4 Hz, 1H), 7.09 (d, <i>J</i> = 8.4 Hz, 1H), 6.91 (t, <i>J</i> = 8.7 Hz, 1H), 6.64 (s, 1H), 6.58 (dd, <i>J</i> = 8.4, 2.8 Hz, 1H), 5.85 (d, <i>J</i> = 1.4 Hz, 1H), 2.19 (s, 3H)
89	¹ H NMR (300 MHz, CD ₃ OD) δ 7.49 (d, <i>J</i> = 8.4 Hz, 1H), 7.37 (d, <i>J</i> = 8.8 Hz, 1H), 6.99 (s, 1H), 6.94 (d, <i>J</i> = 8.9 Hz, 1H), 6.82 (d, <i>J</i> = 8.4 Hz, 2H), 6.07 (s, 1H)
90	¹ H NMR (300 MHz, CDCl ₃) δ 8.11 (s, 1H), 8.06 (d, <i>J</i> = 8.2 Hz, 2H), 7.98 (d, <i>J</i> = 8.2 Hz, 1H), 7.68 - 7.60 (m, 2H), 7.19 (d, <i>J</i> = 8.5 Hz, 1H), 6.71 (d, <i>J</i> = 2.9 Hz, 1H), 6.66 (dd, <i>J</i> = 8.5, 2.8 Hz, 1H), 6.13 (s, 1H), 5.53 (s, 1H), 3.93 (s, 3H),

	2.23 (s, 3H)
91	¹ H NMR (300 MHz, DMSO) δ 12.67 (s, 1H), 8.81 (s, 1H), 8.01 (d, J = 12.4 Hz, 1H), 7.95 (d, J = 7.9 Hz, 2H), 7.81 (d, J = 8.6 Hz, 2H), 7.17 (d, J = 8.3 Hz, 1H), 7.10 (s, 1H), 6.58 (s, 1H), 6.52 (d, J = 8.1 Hz, 1H), 6.12 (s, 1H), 2.16 (s, 3H)
92	¹ H NMR (300 MHz, CD ₃ OD) δ 7.38 (dd, J = 12.2, 2.1 Hz, 1H), 7.29 (d, J = 8.2 Hz, 1H), 7.03 (d, J = 8.8 Hz, 2H), 6.94 (t, J = 8.7 Hz, 1H), 6.70 (d, J = 8.8 Hz, 2H), 6.05 (s, 3H)
93	¹ H NMR (300 MHz, DMSO- d_6) δ 12.49 (s, 1H), 8.79 (d, J = 2.5 Hz, 1H), 8.70 (s, 1H), 8.22 - 8.11 (m, 1H), 8.06 (s, 1H), 7.58 (s, 1H), 7.15 (s, 1H), 6.91 (s, 1H), 6.64 (d, J = 8.4 Hz, 2H), 6.31 (s, 1H).
94	¹ H NMR (300 MHz, DMSO- d_6) δ 12.11 (s, 1H), 9.63 (s, 1H), 8.91 (s, 1H), 8.21 (s, 1H), 7.51 (d, J = 8.7 Hz, 2H), 6.87 - 6.74 (m, 4H), 5.98 (s, 1H).
95	¹ H NMR (300 MHz, DMSO- d_6) δ 11.91 (s, 1H), 8.83 (s, 1H), 8.19 (s, 1H), 7.81 (d, J = 9.5 Hz, 1H), 7.31 (dt, J = 8.6, 5.6 Hz, 5H), 7.05 (s, 1H), 6.57 (d, J = 2.8 Hz, 1H), 6.50 (td, J = 6.9, 6.4, 3.4 Hz, 2H), 5.85 (s, 1H), 5.13 (s, 2H), 2.13 (s, 3H).
96	¹ H NMR (300 MHz, DMSO- d_6) δ 7.60 (s, 1H), 7.35 (d, J = 8.1 Hz, 2H), 7.16 (s, 1H), 6.99 (d, J = 7.3 Hz, 1H), 6.61 (t, J = 9.6 Hz, 3H), 6.08 (s, 1H), 5.28 (s, 2H), 4.98 (s, 1H), 4.37 (s, 2H), 2.20 (s, 3H).
97	¹ H NMR (300 MHz, DMSO- d_6) δ 12.06 (s, 1H), 8.72 (s, 1H), 8.24 (s, 1H), 7.95 (s, 1H), 7.82 (d, J = 9.5 Hz, 1H), 7.41 - 7.23 (m, 5H), 7.05 (s, 1H), 6.62 (d, J = 8.3 Hz, 2H), 6.51 (d, J = 9.6 Hz, 1H), 6.02 (s, 1H), 5.14 (s, 2H).
98	¹ H NMR (300 MHz, DMSO) δ 12.11 (s, 1H), 8.72 (s, 1H), 7.94 (s, 1H), 7.15 (d, J = 3.6 Hz, 1H), 7.04 (q, J = 9.1, 8.6 Hz, 2H), 6.77 (s, 1H), 6.63 (d, J = 8.7 Hz, 2H), 5.89 (s, 1H), 2.44 (s, 3H)
99	¹ H NMR (300 MHz, DMSO) δ 9.92 (s, 1H), 8.74 (s, 1H), 7.96 (s, 1H), 7.47 (s, 1H), 7.36 (s, 1H), 7.08 (d, J = 5.3 Hz, 2H), 6.63 (d, J = 8.6 Hz, 2H), 6.43 (q, J = 8.5 Hz, 1H), 5.97 (s, 1H)
100	¹ H NMR (300 MHz, DMSO) δ 8.81 (s, 1H), 7.14 (d, J = 8.5 Hz, 1H), 7.11-7.02 (m, 2H), 6.73 (s, 1H), 6.56 (s, 1H), 6.51 (dd, J = 8.5, 2.8 Hz, 1H), 6.43-6.37 (m, 1H), 5.77 (s, 1H), 2.43 (s, 3H), 2.13 (s, 3H)
101	¹ H NMR (300 MHz, DMSO) δ 12.04 (s, 1H), 8.84 (s, 1H), 7.44 (d, J = 4.8 Hz, 1H), 7.32 (d, J = 3.6 Hz, 1H), 7.16-7.01 (m, 3H), 6.57 (s, 1H), 6.51 (d, J = 8.4 Hz, 1H), 5.83 (s, 1H), 2.14 (s, 3H)

102	¹ H NMR (300 MHz, DMSO) δ 9.65 (s, 1H), 7.37-7.20 (m, 3H), 7.15 (s, 1H), 6.77 (s, 2H), 6.42 (s, 1H), 6.00 (d, J = 2.5 Hz, 1H), 2.44 (s, 2H), 2.19 (s, 2H), 1.99 (s, 3H)
103	¹ H NMR (300 MHz, DMSO) δ 11.85 (s, 1H), 8.91 (s, 1H), 7.15 (s, 2H), 7.05 (s, 1H), 6.93 (s, 1H), 6.58 (s, 1H), 6.52 (dd, J = 9.0, 2.5 Hz, 1H), 5.81 (s, 1H), 2.13 (s, 3H)
104	¹ H NMR (300 MHz, DMSO) δ 12.42 (s, 1H), 8.84 (s, 1H), 8.06 (s, 1H), 7.83 (s, 1H), 7.67 (s, 1H), 7.11 (d, J = 8.0 Hz, 1H), 6.86-6.76 (m, 1H), 6.66 (d, J = 8.2 Hz, 2H), 6.16 (s, 1H)
105	¹ H NMR (300 MHz, DMSO) δ 12.18 (s, 1H), 8.94 (s, 1H), 7.80 (s, 1H), 7.63 (s, 1H), 7.29 (s, 1H), 6.97 (s, 1H), 6.60 (s, 1H), 6.54 (d, J = 8.7 Hz, 1H), 5.93 (s, 1H), 2.14 (s, 3H)
109	¹ H NMR (300 MHz, DMSO- d_6) δ 12.53 (s, 1H), 8.32 (s, 1H), 7.86 (d, J = 8.4 Hz, 2H), 7.80 (s, 1H), 7.73 (d, J = 8.3 Hz, 2H), 7.38 (s, 1H), 7.12 (s, 1H), 6.97 (d, J = 8.7 Hz, 2H), 6.38 (s, 1H), 2.88 (s, 3H), 2.24 (s, 3H).
110	¹ H NMR (300 MHz, DMSO- d_6) δ 8.34 (s, 1H), 7.90 - 7.76 (m, 5H), 7.23 (s, 1H), 7.14 (s, 1H), 6.98 (d, J = 2.1 Hz, 1H), 6.92 (dd, J = 8.3, 2.5 Hz, 1H), 2.87 (s, 3H), 2.24 (s, 3H).
111	¹ H NMR (300 MHz, DMSO- d_6) δ 8.84 (s, 1H), 8.30 (s, 1H), 7.81 (d, J = 8.6 Hz, 2H), 7.79 (s, 1H), 7.68 (d, J = 8.5 Hz, 2H), 7.18 (d, J = 8.4 Hz, 1H), 7.11 (s, 1H), 7.03 (s, 1H), 6.59 (d, J = 2.8 Hz, 1H), 6.53 (dd, J = 8.5, 2.9 Hz, 1H), 6.06 (s, 1H), 2.62 - 2.53 (m, 2H), 1.13 (t, J = 7.5 Hz, 3H).
112	¹ H NMR (300 MHz, DMSO) δ 11.7 (s, 1H), 8.61 (s, 1H), 7.80 (s, 1H), 7.13 (s, 2H), 7.04 (s, 1H), 6.71 (s, 1H), 6.60 (s, 1H), 6.29 (s, 1H), 5.73 (s, 1H), 3.62 (s, 3H)
115	¹ H NMR (500 MHz, DMSO- d_6) δ 11.91 (s, 1H), 9.01 (s, 1H), 7.92 (s, 1H), 7.77 (s, 1H), 7.62 (s, 1H), 7.34 (s, 3H), 6.97 (s, 1H), 6.63 (s, 1H), 6.57 (d, J = 8.5 Hz, 1H), 5.94 (s, 1H), 2.59 - 2.53 (m, 2H), 1.14 (t, J = 7.5 Hz, 3H).
117	¹ H NMR (500 MHz, DMSO- d_6) δ 8.87 (s, 1H), 8.04 (d, J = 7.9 Hz, 1H), 7.94 (s, 1H), 7.45 (dt, J = 22.4, 7.3 Hz, 2H), 7.11 (s, 1H), 6.61 (d, J = 2.8 Hz, 1H), 6.55 (dd, J = 8.5, 2.9 Hz, 1H), 6.03 (s, 1H), 2.58 (q, J = 7.4 Hz, 2H), 1.15 (t, J = 7.5 Hz, 3H).
119	¹ H NMR (500 MHz, DMSO- d_6) δ 11.92 (s, 1H), 8.96 (s, 1H), 7.17 (t, J = 3.7 Hz, 1H), 7.06 (d, J = 3.9 Hz, 1H), 7.04 - 6.98 (m, 1H), 6.61 (d, J = 2.9 Hz, 1H), 6.55 (dd, J = 8.5, 2.8 Hz, 1H), 5.78 (s, 1H), 2.56 - 2.51 (m, 2H), 1.14 - 1.08 (m,

	3H).
121	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 12.10 (s, 1H), 9.04 (s, 1H), 7.78 (d, <i>J</i> = 3.2 Hz, 1H), 7.60 (d, <i>J</i> = 3.3 Hz, 1H), 7.32 (s, 1H), 6.93 (d, <i>J</i> = 8.5 Hz, 1H), 6.64 (s, 1H), 6.58 (d, <i>J</i> = 8.4 Hz, 1H), 5.83 (s, 1H), 2.55 (d, <i>J</i> = 7.1 Hz, 2H), 1.12 (t, <i>J</i> = 7.5 Hz, 3H).
123	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 12.18 (s, 1H), 8.81 (s, 1H), 8.13 (s, 2H), 7.37 (d, <i>J</i> = 32.0 Hz, 3H), 7.20 - 6.92 (m, 2H), 6.57 (d, <i>J</i> = 29.6 Hz, 3H), 2.62 - 2.54 (m, 2H), 1.66 (s, H), 1.15 (t, <i>J</i> = 7.5 Hz, 3H).
125	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.98 (s, 1H), 10.56 (s, 1H), 8.81 (s, 1H), 8.16 (d, <i>J</i> = 10.8 Hz, 3H), 8.04 (d, <i>J</i> = 10.5 Hz, 2H), 7.82 (d, <i>J</i> = 8.7 Hz, 2H), 7.39 (s, 1H), 6.08 (s, 1H), 2.26 (s, 3H), 2.10 (s, 3H).
126	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 12.17 (s, 1H), 9.63 (s, 1H), 9.20 (s, 1H), 7.70 (s, 1H), 7.51 (d, <i>J</i> = 8.6 Hz, 2H), 7.19 (s, 1H), 6.87 - 6.76 (m, 3H), 6.68 (dd, <i>J</i> = 8.9, 2.8 Hz, 1H), 6.14 (s, 1H).
127	¹ H NMR (300 MHz, CDCl ₃) δ 7.84 - 7.74 (m, 2H), 7.60 (s, 1H), 7.38 - 7.28 (m, 3H), 7.18 - 7.14 (m, 2H), 6.23 (s, 1H), 5.80 (s, 1H), 2.20 (s, 3H), 2.17 (s, 3H).
128	¹ H NMR (300 MHz, CDCl ₃) δ 8.07 (dd, <i>J</i> = 9.0, 2.2 Hz, 1H), 7.91 (dd, <i>J</i> = 6.2, 2.2 Hz, 1H), 7.58 (s, 1H), 7.44 - 7.38 (m, 3H), 7.34 (d, <i>J</i> = 2.1 Hz, 1H), 7.20 - 7.13 (m, 2H), 6.29 (s, 1H), 5.85 (s, 1H), 2.26 (s, 3H), 2.16 (s, 3H).
132	¹ H NMR (300 MHz, DMSO) δ 8.91 (d, <i>J</i> = 1.7 Hz, 2H), 8.48 (d, <i>J</i> = 3.8 Hz, 1H), 8.05 (d, <i>J</i> = 8.2 Hz, 1H), 7.41 (s, 1H), 7.12 (s, 2H), 6.59 (d, <i>J</i> = 2.5 Hz, 1H), 6.53 (d, <i>J</i> = 8.6 Hz, 1H), 6.09 (s, 1H), 2.55 (q, <i>J</i> = 7.5 Hz, 2H), 1.12 (t, <i>J</i> = 7.5 Hz, 3H).
133	¹ H NMR (300 MHz, DMSO) δ 8.93 (s, 1H), 8.87 (s, 1H), 8.60 (s, 1H), 8.35 (s, 1H), 7.18 (s, 1H), 6.59 (d, <i>J</i> = 2.2 Hz, 1H), 6.53 (dd, <i>J</i> = 8.7, 2.5 Hz, 1H), 6.21 (s, 1H), 2.15 (s, 3H).
135	¹ H NMR (300 MHz, DMSO) δ 12.54 (s, 1H), 9.26 (s, 1H), 7.85 - 7.60 (m, 3H), 7.01 - 6.97 (m, 2H), 6.27 (s, 1H), 2.90 (s, 3H), 2.23 (s, 3H).
137	¹ H NMR (300 MHz, DMSO) δ 9.60 (s, 1H), 7.95 (d, <i>J</i> = 5.4 Hz, 2H), 7.04 (d, <i>J</i> = 5.4 Hz, 2H), 6.85 (d, <i>J</i> = 5.0 Hz, 1H), 6.73 (d, <i>J</i> = 1.4 Hz, 1H), 6.66 (dd, <i>J</i> = 5.0, 1.4 Hz, 1H), 6.39 (s, 1H), 3.59 (t, <i>J</i> = 2.9 Hz, 4H), 3.45 (d, <i>J</i> = 2.9 Hz, 2H), 3.38 (d, <i>J</i> = 3.0 Hz, 2H), 2.41 - 2.35 (m, 2H), 2.05(s, 3H), 1.12 (t, <i>J</i> = 4.5 Hz, 3H).
138	¹ H NMR (300 MHz, DMSO) δ 12.15 (s, 1H), 8.84 (s, 1H), 8.54 (d, <i>J</i> = 2.5 Hz,

	1H), 7.89 - 7.75 (m, 5H), 7.19 (s, 1H), 7.02 (s, 1H), 6.59 - 6.52 (m, 3H), 6.04 (s, 1H), 2.56 (q, $J = 7.4$ Hz, 2H), 1.12 (t, $J = 7.4$ Hz, 3H).
140	¹ H NMR (300 MHz, MeOD) δ 7.80 (d, $J = 7.8$ Hz, 1H), 7.59 (s, 1H), 7.43 (d, $J = 7.6$ Hz, 1H), 7.23 - 7.08 (m, 3H), 6.68 (d, $J = 2.7$ Hz, 1H), 6.62 (dd, $J = 8.5$, 2.7 Hz, 1H), 5.99 (s, 1H), 2.27 (s, 3H).
141	¹ H NMR (300 MHz, DMSO) δ 7.77 (d, $J = 7.3$ Hz, 1H), 7.57 (s, 1H), 7.41 (d, $J = 7.7$ Hz, 1H), 7.20 - 7.10 (m, 3H), 6.69 (d, $J = 2.7$ Hz, 1H), 6.61 (dd, $J = 8.5$, 2.9 Hz, 1H), 5.94 (s, 1H), 2.65 (q, $J = 7.5$ Hz, 2H), 1.23 (t, $J = 7.5$ Hz, 3H).
142	¹ H NMR (300 MHz, DMSO) δ 12.31 (s, 1H), 9.83 (s, 1H), 9.66 (s, 1H), 7.89 (d, $J = 7.5$ Hz, 1H), 7.76 (s, 1H), 7.52-7.48 (m, 3H), 7.28-7.25 (m, 1H), 6.82 (d, $J = 8.3$ Hz, 2H), 6.25 (s, 1H), 2.00 (s, 3H).
143	¹ H NMR (300 MHz, DMSO- d_6) δ 12.24 (s, 1H), 10.18 (s, 1H), 9.69 (s, 1H), 8.70 (s, 1H), 7.84 (s, 1H), 7.52 (s, 1H), 7.48 (d, $J = 3.9$ Hz, 2H), 6.81 (d, $J = 8.3$ Hz, 2H), 6.11 (s, 1H), 2.25 (s, 3H), 2.03 (s, 3H).
146	¹ H NMR (300 MHz, MeOD) δ 7.75 (d, $J = 8.4$ Hz, 2H), 7.45 (d, $J = 8.4$ Hz, 2H), 7.11 (d, $J = 8.5$ Hz, 1H), 6.70 (d, $J = 2.7$ Hz, 1H), 6.62 (dd, $J = 8.5$, 2.8 Hz, 1H), 5.93 (s, 1H), 2.63 (q, $J = 7.5$ Hz, 2H), 1.21 (t, $J = 7.5$ Hz, 3H).
147	¹ H NMR (500 MHz, DMSO- d_6) δ 12.12 (s, 1H), 9.61 (s, 1H), 9.56 (s, 1H), 7.55 - 7.48 (m, 2H), 7.34 (d, $J = 2.4$ Hz, 1H), 7.26 (dd, $J = 8.7$, 2.5 Hz, 1H), 7.19 (s, 1H), 6.82 - 6.78 (m, 2H), 6.10 (s, 1H), 2.24 (d, $J = 7.2$ Hz, 2H), 2.21 (d, $J = 1.9$ Hz, 3H), 1.60 (h, $J = 7.4$ Hz, 2H), 0.91 (t, $J = 7.4$ Hz, 3H).
148	¹ H NMR (300 MHz, DMSO) δ 9.99 (s, 1H), 8.40 (s, 1H), 8.22 (s, 1H), 7.99 (d, $J = 8.5$ Hz, 2H), 7.88 (s, 1H), 7.83 (d, $J = 8.0$ Hz, 2H), 7.55 (s, 1H), 7.46 (s, 1H), 7.15 (s, 1H), 7.00 (s, 1H), 6.74 (s, 1H), 2.09 (s, 3H), 2.06 (s, 3H).
149	¹ H NMR (300 MHz, DMSO) δ 12.21 (s, 1H), 9.61 (s, 1H), 7.30 (s, 1H), 7.26-7.22 (m, 5H), 7.13 (s, 1H), 6.07 (s, 1H), 2.16 (s, 6H), 1.98 (s, 3H).
150	¹ H NMR (300 MHz, DMSO) δ 11.91 (s, 1H), 8.71 (s, 1H), 8.41 (s, 1H), 7.28-7.23 (m, 2H), 6.85 (s, 1H), 6.55-6.48 (m, 2H), 5.87 (s, 1H), 2.18 (s, 6H), 2.14 (s, 3H).
151	¹ H NMR (300 MHz, DMSO) δ 12.24 (s, 1H), 10.94 (s, 1H), 9.63 (s, 1H), 7.71 (s, 1H), 7.52 (d, $J = 8.2$ Hz, 2H), 7.37-7.31 (m, 3H), 6.81 (d, $J = 8.1$ Hz, 2H), 6.18 (s, 1H), 2.24 (s, 3H).
152	¹ H NMR (300 MHz, DMSO) δ 11.84 (s, 1H), 9.54 (s, 1H), 7.46 (d, $J = 8.7$ Hz, 2H), 7.09 (s, 1H), 6.76 (d, $J = 8.3$ Hz, 2H), 6.40 (s, 1H), 6.35-6.32 (m, 1H), 5.78 (s, 1H), 4.55 (s, 2H), 2.09 (s, 3H).

153	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.12 (s, 1H), 9.63 (s, 1H), 7.45 (s, 1H), 7.34-7.30 (m, 3 H), 7.25-7.19 (m, 2H), 6.13 (s, 1H), 2.20 (s, 9H), 1.98 (s, 3H), 1.00 (s, 9H), 0.20 (s, 6H).
156	¹ H NMR (300 MHz, DMSO) δ 12.25 (s, 1H), 8.85 (s, 1H), 8.75 (d, <i>J</i> = 2.4 Hz, 1H), 8.14 (dd, <i>J</i> = 8.4, 2.4 Hz, 1H), 7.54 (d, <i>J</i> = 8.2 Hz, 1H), 7.17 (s, 1H), 7.10 (s, 1H), 6.58 (d, <i>J</i> = 2.5 Hz, 1H), 6.52 (dd, <i>J</i> = 8.5, 2.7 Hz, 1H), 6.12 (s, 1H), 2.14 (s, 3H).
157	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.11 (s, 1H), 9.62 (s, 1H), 9.21 (s, 1H), 7.50 (d, <i>J</i> = 8.1 Hz, 2H), 7.15 (d, <i>J</i> = 7.9 Hz, 3H), 6.80 (d, <i>J</i> = 8.0 Hz, 2H), 6.07 (s, 1H), 4.08 (q, <i>J</i> = 6.6 Hz, 2H), 2.19 (s, 3H), 1.23 (t, <i>J</i> = 6.5 Hz, 3H).
158	¹ H NMR (300 MHz, MeOD) δ 7.39 (dd, <i>J</i> = 12.3, 1.9 Hz, 1H), 7.31 (d, <i>J</i> = 8.5 Hz, 2H), 7.04 (s, 1H), 7.00 (dd, <i>J</i> = 8.5, 2.4 Hz, 1H), 6.94 (t, <i>J</i> = 8.7 Hz, 1H), 6.14 (s, 1H), 2.87 (s, 3H), 2.26 (s, 3H).
159	¹ H NMR (300 MHz, Chloroform- <i>d</i>) δ 7.85 (d, <i>J</i> = 7.9 Hz, 2H), 7.35 (s, 1H), 7.23 (s, 1H), 7.12 (d, <i>J</i> = 8.4 Hz, 1H), 6.88 (d, <i>J</i> = 7.9 Hz, 2H), 6.76 (s, 1H), 6.13 (s, 1H), 3.78 (s, 3H), 2.21 (s, 3H).
160	¹ H NMR (300 MHz, MeOD) δ 7.38 (dd, <i>J</i> = 12.3, 2.1 Hz, 1H), 7.29 (d, <i>J</i> = 8.3 Hz, 1H), 7.12 (d, <i>J</i> = 8.5 Hz, 1H), 6.94 (t, <i>J</i> = 8.7 Hz, 1H), 6.70 (d, <i>J</i> = 2.8 Hz, 1H), 6.62 (dd, <i>J</i> = 8.5, 2.9 Hz, 1H), 5.85 (s, 1H), 2.63 (q, <i>J</i> = 7.5 Hz, 2H), 1.21 (t, <i>J</i> = 7.5 Hz, 3H).
161	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.18 (s, 1H), 10.04 (s, 1H), 9.26 (s, 1H), 7.49 (d, <i>J</i> = 12.5 Hz, 1H), 7.34 (d, <i>J</i> = 8.2 Hz, 1H), 7.22 - 7.09 (m, 3H), 6.97 (t, <i>J</i> = 8.7 Hz, 1H), 6.14 (s, 1H), 3.62 (s, 3H), 2.19 (s, 3H).
162	¹ H NMR (300 MHz, DMSO) δ 9.18 (s, 1H), 7.71 (s, 1H), 7.58 (s, 1H), 7.49-7.44 (m, 1H), 7.30 (d, <i>J</i> = 8.6 Hz, 1H), 6.97 (t, <i>J</i> = 8.7 Hz, 1H), 6.59-6.51 (m, 2H), 6.06 (s, 1H).
163	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.28 (s, 1H), 9.69 (s, 1H), 8.70 (s, 1H), 8.26 (s, 1H), 7.96 (t, <i>J</i> = 8.3 Hz, 1H), 7.53 (d, <i>J</i> = 8.6 Hz, 2H), 7.01 (dd, <i>J</i> = 8.8, 3.3 Hz, 1H), 6.82 (d, <i>J</i> = 8.5 Hz, 2H), 6.06 (s, 1H).
164	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 11.87 (s, 1H), 9.57 (s, 1H), 7.47 (d, <i>J</i> = 8.6 Hz, 2H), 7.19 (d, <i>J</i> = 8.5 Hz, 1H), 6.80 (d, <i>J</i> = 3.1 Hz, 2H), 6.76 (s, 1H), 6.44 (d, <i>J</i> = 2.4 Hz, 1H), 6.38 (dd, <i>J</i> = 8.6, 2.6 Hz, 1H), 5.82 (s, 1H), 4.85 (s, 1H), 3.67 - 3.52 (m, 4H), 3.08 (t, <i>J</i> = 6.8 Hz, 2H), 2.47 (s, 2H), 2.45 - 2.36 (m, 4H), 2.14 (s, 3H).
165	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.53 (s, 1H), 9.73 (s, 1H), 7.99 (s, 1H),

	7.83 (s, 1H), 7.53 (d, $J = 8.6$ Hz, 2H), 7.46 (d, $J = 6.8$ Hz, 2H), 6.82 (d, $J = 8.6$ Hz, 2H), 6.31 (s, 1H), 2.26 (s, 3H).
166	¹ H NMR (300 MHz, DMSO- d_6) δ 12.06 (s, 1H), 9.65 (s, 1H), 7.49 (d, $J = 8.7$ Hz, 2H), 7.04 (s, 1H), 6.79 (d, $J = 8.9$ Hz, 3H), 6.70 (dd, $J = 8.7, 2.8$ Hz, 1H), 6.01 (s, 1H), 3.56 (q, $J = 4.3$ Hz, 4H), 2.97 (dt, $J = 19.4, 5.1$ Hz, 4H), 2.20 (s, 3H), 2.03 (s, 3H).
167	¹ H NMR (300 MHz, DMSO- d_6) δ 7.61 (d, $J = 8.7$ Hz, 2H), 7.48 - 7.33 (m, 6H), 7.05 (d, $J = 9.0$ Hz, 3H), 6.73 (d, $J = 2.6$ Hz, 1H), 6.66 (dd, $J = 8.7, 2.8$ Hz, 1H), 6.04 (s, 1H), 5.14 (s, 2H), 2.97 - 2.87 (m, 4H), 2.87 - 2.78 (m, 4H), 2.20 (s, 3H).
168	¹ H NMR (300 MHz, DMSO- d_6) δ 11.64 (s, 1H), 9.50 (s, 1H), 8.96 (s, 1H), 7.41 (d, $J = 8.6$ Hz, 2H), 6.77 - 6.69 (m, 2H), 6.67 - 6.57 (m, 1H), 6.45 (s, 2H), 5.33 (s, 1H), 2.10 (s, 6H).
169	¹ H NMR (300 MHz, DMSO- d_6) δ 12.08 (s, 1H), 9.63 (s, 1H), 8.82 (s, 2H), 7.49 (d, $J = 8.6$ Hz, 2H), 7.09 (s, 1H), 6.80 (d, $J = 8.6$ Hz, 3H), 6.73 (dd, $J = 8.8, 2.9$ Hz, 1H), 6.03 (s, 1H), 3.33 (s, 4H), 3.17 (d, $J = 4.2$ Hz, 4H), 2.21 (s, 3H).
170	¹ H NMR (300 MHz, DMSO) δ 12.48 (s, 1H), 9.66 (s, 1H), 8.33 (d, 2H), 7.87 (d, $J = 8.6$ Hz, 2H), 7.81 (s, 1H), 7.74 (d, $J = 7.7$ Hz, 2H), 7.39 (d, $J = 8.8$ Hz, 1H), 7.28 (s, 1H), 7.13 (s, 1H), 6.31 (s, 1H), 1.99 (s, 3H).
171	¹ H NMR (300 MHz, DMSO) δ 8.51 (s, 1H), 8.33 (s, 1H), 7.87 (d, $J = 8.7$ Hz, 2H), 7.81 (s, 1H), 7.74 (d, $J = 8.7$ Hz, 2H), 7.30 (d, $J = 7.7$ Hz, 1H), 7.13 (s, 1H), 7.08 (d, $J = 8.8$ Hz, 2H), 6.33 (s, 1H), 2.87 (s, 3H).
172	¹ H NMR (300 MHz, DMSO) δ 12.48 (s, 1H), 9.20 (s, 1H), 8.32 (s, 1H), 7.81 - 7.68 (m, 7H), 7.13 (s, 1H), 6.62 - 6.51 (m, 2H), 6.23 (s, 1H).
173	¹ H NMR (300 MHz, DMSO) δ 8.92 (s, 1H), 8.32 (s, 1H), 7.84 (d, $J = 8.7$ Hz, 2H), 7.80 (s, 1H), 7.72 (d, $J = 8.6$ Hz, 2H), 7.48 (s, 1H), 7.12 (s, 1H), 6.98 - 6.96 (m, 2H), 6.30 (s, 1H).
174	¹ H NMR (300 MHz, DMSO) δ 12.66 (s, 1H), 9.87 (s, 1H), 8.93 (s, 1H), 8.33 (s, 1H), 7.82 - 7.76 (m, 6H), 7.61 (s, 1H), 7.30 (d, $J = 9.0$ Hz, 1H), 7.13 (s, 1H), 6.49 (s, 1H), 2.01 (s, 3H).
175	¹ H NMR (300 MHz, DMSO) δ 8.40 (s, 1H), 7.98 (d, $J = 8.6$ Hz, 2H), 7.87 (s, 1H), 7.82 (d, $J = 8.6$ Hz, 2H), 7.15 (s, 1H), 7.01 (t, $J = 9.2$ Hz, 1H), 6.92 (d, $J = 14$ Hz, 1H), 6.81 (d, $J = 9.4$ Hz, 1H), 6.73 (s, 1H).
176	¹ H NMR (300 MHz, DMSO) δ 12.50 (s, 1H), 9.21 (s, 1H), 8.32 (s, 1H), 7.81 -

	7.75 (m, 5H), 7.25 (s, 1H), 7.13 (s, 1H), 6.81 (s, 1H), 6.69 (d, $J = 8.8$ Hz, 1H), 6.35 (s, 1H).
177	^1H NMR (300 MHz, DMSO) δ 12.61 (s, 1H), 9.70 (s, 1H), 8.33 (s, 1H), 7.84 - 7.76 (m, 5H), 7.31 (s, 1H), 7.13 (s, 1H), 7.06 (s, 1H), 6.98 (d, $J = 8.3$ Hz, 1H), 6.42 (s, 1H), 2.59 (q, $J = 7.4$ Hz, 2H), 1.97 (s, 3H), 1.15 (t, $J = 7.4$ Hz, 3H).
178	^1H NMR (300 MHz, DMSO) δ 9.78 (s, 1H), 8.41 (s, 1H), 8.00 (d, $J = 8.5$ Hz, 2H), 7.88 - 7.81 (m, 4H), 7.15 (s, 1H), 6.90 (d, $J = 11.5$ Hz, 1H), 6.75 (s, 1H), 2.11 (s, 3H), 2.07 (s, 3H).
180	^1H NMR (300 MHz, DMSO- d_6) δ 12.47 (s, 1H), 9.79 (s, 1H), 8.55 (d, $J = 2.5$ Hz, 1H), 7.96 - 7.74 (m, 5H), 7.60 (s, 1H), 7.39 - 7.24 (m, 3H), 6.57 (t, $J = 2.1$ Hz, 1H), 6.31 (s, 1H), 3.87 (t, $J = 6.3$ Hz, 2H), 2.77 (t, $J = 6.3$ Hz, 2H), 2.23 (s, 3H).
181	^1H NMR (300 MHz, DMSO- d_6) δ 9.92 (s, 1H), 9.62 (s, 1H), 8.55 (d, $J = 2.5$ Hz, 1H), 7.91 (d, $J = 8.8$ Hz, 2H), 7.84 (d, $J = 8.9$ Hz, 2H), 7.77 (d, $J = 1.7$ Hz, 1H), 7.50 (d, $J = 8.7$ Hz, 1H), 7.35 (d, $J = 2.4$ Hz, 2H), 7.29 (dd, $J = 8.6, 2.5$ Hz, 1H), 6.60 - 6.54 (m, 1H), 6.32 (s, 1H), 4.00 (d, $J = 12.8$ Hz, 12H), 2.23 (s, 3H).
182	^1H NMR (300 MHz, Methanol- d_4) δ 7.40 (dd, $J = 12.4, 1.7$ Hz, 1H), 7.37-7.18 (m, 4H), 6.95 (t, $J = 8.7$ Hz, 1H), 6.13 (s, 1H), 3.12 (s, 2H), 2.38 (s, 6H), 2.28 (s, 3H).
183	^1H NMR (300 MHz, Methanol- d_4) δ 7.40 (dd, $J = 12.2, 2.0$ Hz, 1H), 7.36-7.22 (m, 4H), 6.95 (t, $J = 8.7$ Hz, 1H), 6.12 (s, 1H), 3.73 (t, $J = 4.7$ Hz, 4H), 2.78 (t, $J = 7.0$ Hz, 2H), 2.64-2.50 (m, 6H), 2.27 (s, 3H).
184	^1H NMR (300 MHz, DMSO- d_6) δ 12.54 (s, 1H), 9.25 (s, 1H), 8.24-8.04 (m, 1H), 7.79 (dd, $J = 8.6, 5.4$ Hz, 3H), 7.38-7.13 (m, 2H), 7.03-6.45 (m, 2H).
185	^1H NMR (300 MHz, Methanol- d_4) δ 7.40 (dd, $J = 12.3, 2.1$ Hz, 1H), 7.36-7.24 (m, 4H), 6.96 (t, $J = 8.7$ Hz, 1H), 6.13 (s, 1H), 3.20 (t, $J = 6.5$ Hz, 2H), 2.76 (t, $J = 6.7$ Hz, 2H), 2.71 (s, 6H), 2.27 (s, 3H).
187	^1H NMR (300 MHz, DMSO- d_6) δ 12.51 (s, 1H), 9.18 (s, 1H), 8.56 (d, $J = 2.3$ Hz, 1H), 7.91 (d, $J = 8.7$ Hz, 2H), 7.84 (d, $J = 8.4$ Hz, 2H), 7.77 (s, 1H), 7.40 (s, 1H), 6.98-6.94 (m, 2H), 6.61 - 6.53 (m, 1H), 6.36 (s, 1H), 2.88 (s, 3H), 2.23 (s, 3H).
188	^1H NMR (300 MHz, DMSO- d_6) δ 12.47 (s, 1H), 9.53 (s, 1H), 8.54 (d, $J = 2.5$ Hz, 1H), 7.90 (d, $J = 8.6$ Hz, 2H), 7.82 (d, $J = 8.6$ Hz, 2H), 7.76 (d, $J = 1.7$ Hz, 1H), 7.53 (s, 1H), 6.97 (d, $J = 6.3$ Hz, 3H), 6.56 (t, $J = 2.2$ Hz, 1H), 6.27 (s,

	1H), 5.75 (s, 1H).
189	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.33 (s, 1H), 9.20 (s, 1H), 8.54 (d, <i>J</i> = 2.5 Hz, 1H), 7.90 (d, <i>J</i> = 8.5 Hz, 2H), 7.80 (d, <i>J</i> = 8.6 Hz, 2H), 7.76 (d, <i>J</i> = 1.7 Hz, 1H), 7.65 (s, 1H), 6.63 - 6.55 (m, 2H), 6.52 (dd, <i>J</i> = 8.8, 2.8 Hz, 1H), 6.20 (s, 1H).
190	¹ H NMR (300 MHz, DMSO) δ 12.52 (s, 1H), 9.17 (s, 1H), 7.79 (d, <i>J</i> = 7.6 Hz, 2H), 7.62 (s, 1H), 7.53 (d, <i>J</i> = 7.3 Hz, 2H), 7.37 (s, 1H), 6.97 - 6.94 (m, 2H), 6.32 (s, 1H), 2.88 (s, 3H), 2.22 (s, 3H).
191	¹ H NMR (300 MHz, DMSO) δ 11.38 (s, 1H), 7.83 (d, <i>J</i> = 8.0 Hz, 1H), 7.73 (d, <i>J</i> = 2.5 Hz, 1H), 7.45 (d, <i>J</i> = 7.9 Hz, 1H), 7.29 (s, 1H), 7.21 - 7.08 (m, 2H), 6.96 - 6.94 (m, 2H), 6.25 (s, 1H), 5.71 (s, 1H), 2.87 (s, 3H), 2.24 (s, 3H).
192	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.33 (s, 1H), 9.20 (s, 1H), 8.54 (d, <i>J</i> = 2.5 Hz, 1H), 7.90 (d, <i>J</i> = 8.5 Hz, 2H), 7.80 (d, <i>J</i> = 8.6 Hz, 2H), 7.76 (d, <i>J</i> = 1.7 Hz, 1H), 7.65 (s, 1H), 6.63 - 6.55 (m, 2H), 6.52 (dd, <i>J</i> = 8.8, 2.8 Hz, 1H), 6.20 (s, 1H).
193	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.38 (s, 1H), 9.26 (s, 1H), 8.55 (d, <i>J</i> = 2.5 Hz, 1H), 7.91 (d, <i>J</i> = 8.6 Hz, 2H), 7.82 (d, <i>J</i> = 8.6 Hz, 2H), 7.77 (d, <i>J</i> = 1.6 Hz, 1H), 7.62 (d, <i>J</i> = 8.9 Hz, 1H), 7.32 (s, 1H), 6.82 (d, <i>J</i> = 2.7 Hz, 1H), 6.69 (dd, <i>J</i> = 8.9, 2.8 Hz, 1H), 6.56 (t, <i>J</i> = 2.1 Hz, 1H), 6.33 (s, 1H).
194	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.15 (s, 1H), 8.74 (s, 1H), 7.68 (d, <i>J</i> = 7.0 Hz, 2H), 7.40 (t, <i>J</i> = 7.4 Hz, 2H), 7.29 (t, <i>J</i> = 7.3 Hz, 1H), 6.98 (s, 1H), 6.57 (d, <i>J</i> = 2.8 Hz, 1H), 6.51 (dd, <i>J</i> = 8.5, 2.9 Hz, 1H), 6.05 (s, 1H), 2.16 (s, 3H).
195	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.41 (s, 1H), 9.63 (s, 1H), 7.70 (d, <i>J</i> = 7.5 Hz, 2H), 7.58 (s, 1H), 7.43 (t, <i>J</i> = 7.2 Hz, 2H), 7.33 (d, <i>J</i> = 7.5 Hz, 2H), 7.25 (d, <i>J</i> = 8.0 Hz, 2H), 6.26 (s, 1H), 2.21 (s, 3H), 1.99 (s, 3H).
196	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.48 (s, 1H), 9.17 (s, 1H), 7.71 (d, <i>J</i> = 7.6 Hz, 2H), 7.49 - 7.29 (m, 4H), 6.95 (d, <i>J</i> = 9.7 Hz, 2H), 6.31 (s, 1H), 2.87 (s, 3H), 2.23 (s, 3H).
197	¹ H NMR (300 MHz, DMSO-d ₆) δ 12.31 (s, 1H), 9.15 (s, 1H), 7.64 (d, <i>J</i> = 8.4 Hz, 3H), 7.30 (s, 1H), 7.00 (d, <i>J</i> = 8.5 Hz, 2H), 6.95 (d, <i>J</i> = 8.8 Hz, 2H), 6.21 (s, 1H), 3.79 (s, 3H), 2.87 (s, 3H), 2.22 (s, 3H).
198	¹ H NMR (300 MHz, DMSO) δ 12.76 (s, 1H), 9.19 (s, 1H), 7.91 (s, 4H), 7.54 (m, 2H), 6.98 - 6.95 (m, 2H), 6.47 (s, 1H), 2.88 (s, 3H), 2.23 (s, 3H).
199	¹ H NMR (300 MHz, DMSO) δ 12.55 (s, 1H), 9.19 (s, 1H), 7.75 - 7.62 (m, 5H), 7.40 (s, 1H), 6.98 - 6.95 (m, 2H), 6.34 (s, 1H), 2.88 (s, 3H), 2.23 (s, 3H).

200	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.29 (s, 1H), 10.03 (s, 1H), 9.16 (s, 1H), 7.61 (s, 1H), 7.51 (dd, <i>J</i> = 12.5, 2.1 Hz, 1H), 7.40 - 7.27 (m, 2H), 7.05 - 6.89 (m, 3H), 6.22 (s, 1H), 2.87 (s, 3H), 2.22 (s, 3H).
201	¹ H NMR (300 MHz, DMSO- <i>d</i> 6) δ 12.19 (s, 1H), 9.62 (s, 1H), 7.56 (d, <i>J</i> = 8.4 Hz, 3H), 7.30 (s, 1H), 7.24 (d, <i>J</i> = 8.9 Hz, 1H), 7.18 (s, 1H), 6.99 (d, <i>J</i> = 8.5 Hz, 2H), 6.13 (s, 1H), 4.10 (q, <i>J</i> = 5.3 Hz, 4H), 3.74 (t, <i>J</i> = 4.8 Hz, 4H), 2.20 (s, 3H), 1.98 (s, 3H).
202	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.02 (s, 1H), 8.70 (s, 1H), 7.53 (d, <i>J</i> = 8.3 Hz, 2H), 7.34 (s, 1H), 6.97 (d, <i>J</i> = 8.5 Hz, 2H), 6.89 (s, 1H), 6.55 (d, <i>J</i> = 2.9 Hz, 1H), 6.50 (d, <i>J</i> = 8.6 Hz, 1H), 5.94 (s, 1H), 3.74 (dd, <i>J</i> = 5.9, 3.6 Hz, 4H), 3.14 (t, <i>J</i> = 4.9 Hz, 4H), 2.15 (s, 3H).
203	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.25 (s, 1H), 7.65 (s, 1H), 7.57 (d, <i>J</i> = 8.2 Hz, 2H), 7.28 (s, 1H), 7.04 - 6.89 (m, 4H), 6.18 (s, 1H), 3.74 (t, <i>J</i> = 4.2 Hz, 4H), 3.23 - 3.10 (m, 4H), 2.87 (s, 3H), 2.22 (s, 3H).
204	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.46 (s, 1H), 9.16 (s, 1H), 7.85-7.70 (m, 2H), 7.65 (s, 1H), 7.36 (s, 1H), 7.34-7.18 (m, 2H), 7.01-6.90 (m, 2H), 6.30 (s, 1H), 2.87 (s, 3H), 2.23 (s, 3H).
205	(300 MHz, DMSO- <i>d</i> ₆) δ 12.19 (s, 1H), 8.76 (s, 1H), 7.69 (s, 4H), 7.30 (s, 1H), 6.99 (s, 1H), 6.57 (s, 1H), 6.52 (d, <i>J</i> = 8.8 Hz, 1H), 6.03 (s, 1H), 3.86 (t, <i>J</i> = 7.0 Hz, 2H), 2.16 (s, 3H), 2.08 (p, <i>J</i> = 7.6 Hz, 3H)
206	(300 MHz, DMSO- <i>d</i> ₆)δ 12.22 (s, 1H) 8.94 (s, 2H), 8.78 (s, 1H), 7.67 (d, <i>J</i> = 8.3 Hz, 2H), 7.31 (d, <i>J</i> = 8.2 Hz, 3H), 7.02 (s, 1H), 6.72 - 6.62 (m, 1H), 6.57 (d, <i>J</i> = 2.8 Hz, 1H), 6.52 (dd, <i>J</i> = 8.4, 2.8 Hz, 1H), 6.05 (s, 1H), 4.14 (s, 4H), 3.62 (t, <i>J</i> = 5.6 Hz, 2H), 2.40 (t, <i>J</i> = 6.2 Hz, 2H), 2.16 (s, 3H), 2.12 - 2.04 (m, 1H), 1.91 - 1.79 (m, 5H), 1.75 (s, 6H), 1.56 (d, <i>J</i> = 14.3 Hz, 1H), 1.24 (s, 1H), 0.97 - 0.72 (m, 1H).
207	(300 MHz, DMSO- <i>d</i> 6) δ 12.45 (s, 1H), 9.28 (s, 1H), 8.55 (s, 1H), 7.91 (d, <i>J</i> = 7.4 Hz, 2H), 7.83 (d, <i>J</i> = 8.6 Hz, 2H), 7.77 (d, <i>J</i> = 1.3 Hz, 1H), 7.57 (s, 1H), 7.24 (s, 1H), 7.16 (d, <i>J</i> = 7.8 Hz, 2H), 6.57 (t, <i>J</i> = 2.0 Hz, 1H), 6.29 (s, 1H), 3.63 (s, 3H), 2.21 (s, 3H).
208	(300 MHz, DMSO- <i>d</i> 6) δ 12.46 (s, 1H), 9.29 (s, 1H), 8.56 (s, 1H), 7.92 (d, <i>J</i> = 7.9 Hz, 2H), 7.84 (d, <i>J</i> = 8.5 Hz, 2H), 7.78 (d, <i>J</i> = 1.7 Hz, 1H), 7.57 (s, 1H), 7.26 (s, 1H), 7.17 (d, <i>J</i> = 9.8 Hz, 2H), 6.58 (t, <i>J</i> = 2.1 Hz, 1H), 6.30 (s, 1H), 3.64 (s, 3H), 2.22 (s, 3H).
209	(300 MHz, DMSO- <i>d</i> 6) δ 12.44 (s, 1H), 9.26 (s, 1H), 8.55 (d, <i>J</i> = 2.5 Hz, 1H), 7.90 (d, <i>J</i> = 8.6 Hz, 2H), 7.83 (d, <i>J</i> = 8.5 Hz, 2H), 7.77 (d, <i>J</i> = 1.7 Hz, 1H),

	7.56 (s, 1H), 7.25 (s, 1H), 7.17 (d, $J = 12.5$ Hz, 2H), 6.57 (t, $J = 2.1$ Hz, 1H), 6.28 (s, 1H), 4.09 (q, $J = 7.1$ Hz, 2H), 2.21 (s, 3H), 1.23 (t, $J = 7.1$ Hz, 3H).
210	(300 MHz, DMSO- d_6) δ 12.37 (s, 1H), 9.72 - 9.56 (m, 1H), 7.72 (s, 4H), 7.58 (s, 1H), 7.32 (s, 1H), 7.29 - 7.20 (m, 2H), 6.24 (s, 1H), 3.90 - 3.82 (m, 2H), 2.22 (s, 3H), 2.13 - 2.04 (m, 9H), 1.99 (d, $J = 2.0$ Hz, 3H).
211	(300 MHz, DMSO- d_6) δ 12.40 (s, 1H), 9.64 (s, 1H), 7.70 (d, $J = 7.9$ Hz, 2H), 7.59 (s, 1H), 7.39 - 7.21 (m, 5H), 6.25 (s, 1H), 4.11 (q, $J = 5.2$ Hz, 2H), 3.63 (d, $J = 6.1$ Hz, 2H), 3.18 (d, $J = 5.2$ Hz, 4H), 2.41 (t, $J = 6.1$ Hz, 3H), 2.22 (s, 3H), 2.00 (s, 3H), 1.87 (d, $J = 5.4$ Hz, 4H).
212	(500 MHz, Methanol- d_4) δ 8.37 (d, $J = 2.3$ Hz, 1H), 7.86 (dd, $J = 8.7, 2.5$ Hz, 1H), 7.05 (d, $J = 8.5$ Hz, 1H), 6.75 (d, $J = 8.6$ Hz, 1H), 6.62 (d, $J = 2.8$ Hz, 1H), 6.56 (dd, $J = 8.5, 2.9$ Hz, 1H), 5.85 (s, 1H), 3.86 (s, 3H).
213	(300 MHz, DMSO- d_6) δ 12.45 (s, 1H), 9.17 (s, 1H), 8.54 (d, $J = 2.4$ Hz, 1H), 8.02 (d, $J = 8.6$ Hz, 1H), 7.64 (s, 1H), 7.38 (s, 1H), 7.00 - 6.86 (m, 3H), 6.30 (s, 1H), 3.88 (s, 3H), 2.87 (s, 3H), 2.22 (s, 3H).
214	(300 MHz, DMSO- d_6) δ 12.39 (s, 1H), 9.64 (s, 1H), 8.53 (d, $J = 2.4$ Hz, 1H), 8.02 (d, $J = 8.7$ Hz, 1H), 7.57 (s, 1H), 7.38 - 7.21 (m, 3H), 6.90 (d, $J = 7.6$ Hz, 1H), 6.24 (s, 1H), 3.88 (s, 3H), 2.21 (s, 3H), 1.99 (s, 3H).
215	(300 MHz, DMSO- d_6) δ 12.08 (s, 1H), 9.19 (s, 1H), 9.07 (s, 1H), 8.77 (s, 1H), 7.72 - 7.65 (m, 2H), 7.37 (d, $J = 8.6$ Hz, 2H), 7.22 (s, 1H), 7.05 - 6.97 (m, 2H), 6.58 (dd, $J = 5.8, 2.8$ Hz, 2H), 6.52 (dt, $J = 8.0, 3.7$ Hz, 2H), 6.00 (s, 1H), 4.10 (d, $J = 5.4$ Hz, 1H), 3.66 (t, $J = 4.5$ Hz, 2H), 3.18 (d, $J = 4.4$ Hz, 3H), 3.09 (t, $J = 4.5$ Hz, 2H), 2.16 (s, 3H), 2.08 (s, 2H), 1.99 (s, 2H).
216	(300 MHz, DMSO- d_6) δ 12.26 (s, 1H), 8.85 (s, 1H), 8.81 (d, $J = 1.7$ Hz, 1H), 8.64 (d, $J = 2.2$ Hz, 1H), 8.28 (dd, $J = 8.6, 2.2$ Hz, 1H), 7.95 (d, $J = 8.6$ Hz, 1H), 7.85 (s, 1H), 7.16 (s, 2H), 6.60 (d, $J = 2.1$ Hz, 2H), 6.54 (dd, $J = 8.6, 2.5$ Hz, 1H), 6.15 (s, 1H), 2.17 (s, 3H).
217	(300 MHz, DMSO- d_6) δ 12.60 (s, 1H), 9.66 (s, 1H), 8.84 (s, 1H), 8.64 (d, $J = 2.6$ Hz, 1H), 8.30 (d, $J = 9.8$ Hz, 1H), 7.97 (d, $J = 8.4$ Hz, 1H), 7.85 (s, 1H), 7.39 (s, 1H), 7.34 (s, 1H), 7.27 (d, $J = 8.2$ Hz, 1H), 6.64 - 6.56 (m, 1H), 6.38 (s, 1H), 2.22 (s, 3H), 1.99 (s, 3H).
218	(300 MHz, DMSO- d_6) δ 12.60 (s, 1H), 9.20 (s, 1H), 8.85 (s, 1H), 8.64 (d, $J = 2.6$ Hz, 1H), 8.31 (d, $J = 8.5$ Hz, 1H), 7.98 (d, $J = 8.5$ Hz, 1H), 7.85 (s, 1H), 7.47 (s, 1H), 6.98-6.95 (m, 2H), 6.64 - 6.56 (m, 1H), 6.44 (s, 1H), 2.88 (s, 3H), 2.23 (s, 3H).

219	(500 MHz, DMSO- d_6) δ 9.97 (s, 3H), 7.74 (d, J = 8.4 Hz, 4H), 7.63 (s, 4H), 7.29 (d, J = 8.6 Hz, 4H), 6.41 (s, 3H), 5.70 (s, 3H), 2.10 (s, 3H), 2.04 (s, 1H).
220	(500 MHz, DMSO-d δ) δ 12.10 (s, 1H), 8.69 (s, 1H), 7.51 (d, J = 8.3 Hz, 2H), 6.89 (s, 1H), 6.80 (d, J = 8.5 Hz, 2H), 6.69 (s, 1H), 6.54 (d, J = 2.7 Hz, 1H), 6.45 (dd, J = 8.5, 2.8 Hz, 1H), 2.94 (s, 6H), 2.16 (s, 3H).
221	(300 MHz, DMSO-d δ) δ 12.50 (s, 1H), 9.23 (s, 1H), 7.95 (d, J = 7.6 Hz, 1H), 7.81 (d, J = 6.9 Hz, 1H), 7.70 (s, 1H), 7.50 (s, 1H), 7.41 - 7.29 (m, 2H), 6.98 (d, J = 9.2 Hz, 2H), 6.29 (s, 1H), 2.88 (s, 3H), 2.22 (s, 3H).
222	(300 MHz, DMSO-d δ) δ 12.17 (s, 1H), 9.35 (s, 1H), 8.03 - 7.64 (m, 3H), 7.56 - 6.90 (m, 6H), 6.21 (s, 1H), 3.64 (s, 3H), 2.21 (s, 3H).
223	(300 MHz, DMSO-d δ) δ 12.63 (d, J = 7.0 Hz, 1H), 12.21 (s, 1H), 9.32 (s, 1H), 7.95 (s, 1H), 7.80 (s, 1H), 7.69 (s, 1H), 7.43 (d, J = 42.1 Hz, 3H), 7.28 - 7.14 (m, 2H), 7.00 (s, 1H), 6.21 (s, 1H), 4.15 - 4.07 (m, 2H), 2.21 (s, 3H), 1.24 (t, J = 7.1 Hz, 3H).
224	(300 MHz, DMSO-d δ) δ 12.63 (s, 1H), 12.18 (s, 1H), 9.34 (s, 1H), 7.94 (d, J = 7.4 Hz, 1H), 7.80 (d, J = 7.2 Hz, 1H), 7.69 (s, 1H), 7.34 (h, J = 6.7, 6.2 Hz, 3H), 7.24 - 7.17 (m, 2H), 6.22 (s, 1H), 3.34 (s, 3H), 2.21 (s, 3H).
225	(300 MHz, DMSO-d δ) δ 9.94 (s, 1H), 8.70 (d, J = 2.1 Hz, 1H), 8.24 (d, J = 8.7 Hz, 2H), 8.06 (d, J = 8.8 Hz, 2H), 7.85 (s, 1H), 7.15 (d, J = 6.6 Hz, 2H), 7.07 (d, J = 8.8 Hz, 1H), 6.67 (s, 1H), 6.63 (s, 1H), 3.07 (s, 3H), 2.71 (s, 3H), 2.15 (s, 3H).
226	(300 MHz, DMSO- d_6) δ 12.42 (s, 1H), 9.86 (s, 1H), 7.90 (s, 1H), 7.77 (d, J = 2.4 Hz, 1H), 7.65 (d, J = 8.4 Hz, 2H), 7.54 (s, 1H), 7.28 (dd, J = 8.9, 2.5 Hz, 1H), 7.02 (d, J = 8.6 Hz, 2H), 6.33 (s, 1H), 3.80 (s, 3H), 2.01 (s, 3H).
227	(300 MHz, DMSO-d δ) δ 12.32 (s, 1H), 8.80 (s, 1H), 8.56 (d, J = 2.5 Hz, 1H), 8.13 (s, 1H), 7.77 (q, J = 3.5 Hz, 2H), 7.63 (d, J = 7.7 Hz, 1H), 7.51 (t, J = 7.9 Hz, 1H), 7.22 (s, 1H), 7.05 (s, 1H), 6.61 - 6.56 (m, 2H), 6.53 (dd, J = 8.5, 2.8 Hz, 1H), 6.16 (s, 1H), 2.16 (s, 3H).
228	^1H NMR (300 MHz, DMSO-d δ) δ 12.58 (s, 1H), 9.20 (s, 1H), 8.57 (d, J = 2.5 Hz, 1H), 8.18 (s, 1H), 7.78 (d, J = 1.5 Hz, 2H), 7.66 (d, J = 7.3 Hz, 1H), 7.55 (t, J = 7.9 Hz, 1H), 7.41 (s, 1H), 6.98 (d, J = 9.0 Hz, 2H), 6.62 - 6.54 (m, 1H), 6.45 (s, 1H), 2.88 (s, 3H), 2.24 (s, 3H).
229	(300 MHz, DMSO- d_6) δ 9.55 (s, 1H), 8.56 (d, J = 2.6 Hz, 1H), 7.91 (d, J = 8.7 Hz, 1H), 7.84 (d, J = 8.6 Hz, 2H), 7.77 (s, 1H), 7.40 - 7.26 (m, 4H), 6.58 (t, J = 2.2 Hz, 1H), 6.30 (s, 1H), 2.22 (s, 3H), 1.09 (d, J = 6.8 Hz, 6H).

230	(500 MHz, DMSO- d_6) δ 11.54 (s, 1H), 9.29 (s, 1H), 7.89 - 7.76 (m, 4H), 6.85 (d, J = 8.5 Hz, 1H), 6.68 (d, J = 2.7 Hz, 1H), 6.62 (dd, J = 8.4, 2.8 Hz, 1H), 4.11 (q, J = 5.3 Hz, 1H), 3.89 (t, J = 7.1 Hz, 2H), 3.18 (d, J = 5.2 Hz, 2H), 2.56 (d, J = 8.0 Hz, 3H), 2.14 (s, 3H), 2.09 (q, J = 7.6 Hz, 2H), 1.24 (s, 1H).
231	(300 MHz, DMSO- d_6) δ 8.56 (d, J = 2.5 Hz, 1H), 7.99 (d, J = 8.5 Hz, 2H), 7.92 - 7.75 (m, 3H), 7.34 (s, 1H), 7.20 (qd, J = 13.8, 12.6, 7.3 Hz, 2H), 6.89 (s, 1H), 6.63 - 6.55 (m, 1H), 3.34 (d, J = 7.0 Hz, 1H), 2.31 (s, 1H), 2.24 (s, 3H), 1.99 (d, J = 1.1 Hz, 4H), 1.18 (t, J = 7.1 Hz, 1H).
232	(300 MHz, DMSO- d_6) δ 12.71 (s, 1H), 9.28 (s, 1H), 8.57 (d, J = 2.5 Hz, 1H), 7.99 (d, J = 8.4 Hz, 2H), 7.84 (d, J = 8.3 Hz, 2H), 7.79 (d, J = 1.7 Hz, 1H), 7.18 (s, 1H), 7.11 (d, J = 6.6 Hz, 2H), 6.59 (t, J = 2.1 Hz, 1H), 3.63 (s, 3H), 2.23 (s, 3H).
233	(300 MHz, DMSO- d_6) δ 12.70 (s, 1H), 9.24 (s, 1H), 8.57 (d, J = 2.5 Hz, 1H), 7.99 (d, J = 8.4 Hz, 2H), 7.84 (d, J = 8.4 Hz, 2H), 7.79 (d, J = 1.7 Hz, 1H), 7.19 (s, 1H), 7.11 (d, J = 7.9 Hz, 2H), 6.59 (t, J = 2.1 Hz, 1H), 4.09 (q, J = 7.1 Hz, 2H), 2.22 (s, 3H), 1.23 (t, J = 7.1 Hz, 3H).
234	(300 MHz, Methanol- d_4) δ 7.70 - 7.60 (m, 2H), 7.05 - 6.98 (m, 2H), 6.94 (d, J = 8.5 Hz, 1H), 6.65 (d, J = 2.8 Hz, 1H), 6.56 (dd, J = 8.6, 2.9 Hz, 1H), 3.84 (s, 3H), 2.25 (s, 3H).
235	(300 MHz, DMSO- d_6) δ 12.82 (s, 1H), 9.18 (s, 1H), 8.57 (d, J = 2.6 Hz, 1H), 7.99 (d, J = 8.5 Hz, 2H), 7.84 (d, J = 8.5 Hz, 2H), 7.78 (d, J = 1.7 Hz, 1H), 7.24 (s, 1H), 6.98 (s, 1H), 6.92 (d, J = 7.4 Hz, 1H), 6.62 - 6.54 (m, 1H), 2.87 (s, 3H), 2.24 (s, 3H).
236	(300 MHz, DMSO- d_6) δ 12.75 (s, 1H), 9.29 (s, 1H), 8.58 (s, 1H), 7.99 (d, J = 8.4 Hz, 2H), 7.90 - 7.75 (m, 3H), 7.16 (d, J = 11.0 Hz, 3H), 6.84 (s, 1H), 6.64 - 6.53 (m, 1H), 3.63 (s, 3H), 2.23 (s, 3H).
237	^1H NMR (300 MHz, DMSO- d_6) δ 12.44 (s, 1H), 8.80 (s, 1H), 7.86 (d, J = 8.0 Hz, 2H), 7.51 (s, 2H), 6.86 (s, 1H), 6.57 (d, J = 2.6 Hz, 1H), 6.47 (dd, J = 8.5, 2.8 Hz, 1H), 2.66 - 2.54 (m, 2H), 1.17 - 1.09 (m, 3H).
238	(300 MHz, DMSO- d_6) δ 13.01 (s, 1H), 9.58 (s, 1H), 8.39 (s, 1H), 8.07 (d, J = 8.7 Hz, 2H), 7.87 (d, J = 8.7 Hz, 2H), 7.63 (s, 1H), 7.07 (d, J = 8.5 Hz, 1H), 6.72 (d, J = 2.7 Hz, 1H), 6.64 (dd, J = 8.5, 2.7 Hz, 1H), 5.99 (s, 1H), 2.14 (d, J = 9.1 Hz, 5H).
239	^1H NMR (500 MHz, DMSO- d_6) δ 10.07 (s, 1H), 8.48 (s, 1H), 8.22 (d, J = 8.9 Hz, 2H), 7.99 (d, J = 8.7 Hz, 2H), 7.68 (s, 1H), 7.58 (s, 1H), 7.51 (d, J = 8.6 Hz, 1H), 7.06 - 7.00 (m, 1H), 6.65 (s, 1H), 2.70 (s, 3H), 2.14 (s, 5H), 2.07 (s,

	3H).
240	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.83 (s, 1H), 7.92 (t, <i>J</i> = 1.8 Hz, 1H), 7.70 (d, <i>J</i> = 7.7 Hz, 1H), 7.48 (d, <i>J</i> = 8.2 Hz, 1H), 7.36 (t, <i>J</i> = 7.8 Hz, 1H), 7.10 (s, 2H), 6.58 (d, <i>J</i> = 2.8 Hz, 1H), 6.52 (dd, <i>J</i> = 8.5, 2.9 Hz, 1H), 6.11 (s, 1H), 2.15 (s, 3H).
241	(300 MHz, DMSO- <i>d</i> ₆) δ 11.16 (s, 1H), 9.33 (s, 1H), 8.97 (d, <i>J</i> = 2.4 Hz, 1H), 8.58 (dd, <i>J</i> = 4.7, 1.6 Hz, 1H), 8.16 (s, 1H), 8.06 (d, <i>J</i> = 8.3 Hz, 2H), 7.80 (d, <i>J</i> = 8.2 Hz, 2H), 7.55 - 7.49 (m, 1H), 6.97 (d, <i>J</i> = 2.2 Hz, 1H), 6.58 (d, <i>J</i> = 2.2 Hz, 1H), 6.35 (s, 1H), 2.40 (s, 3H).
242	¹ H NMR (500 MHz, Acetone- <i>d</i> ₆) δ 8.45 (d, <i>J</i> = 4.6 Hz, 1H), 7.86 (q, <i>J</i> = 8.9 Hz, 4H), 7.70 (d, <i>J</i> = 4.4 Hz, 1H), 7.34 (s, 1H), 6.69 (d, <i>J</i> = 2.9 Hz, 1H), 6.63 (dd, <i>J</i> = 8.6, 3.0 Hz, 1H), 6.54 (s, 1H), 6.14 (s, 1H), 2.24 (s, 3H).
243	¹ H NMR (300 MHz, Acetone- <i>d</i> ₆) δ 11.76 (s, 1H), 8.46 (d, <i>J</i> = 4.6 Hz, 1H), 8.12 (s, 1H), 7.89 (s, 4H), 7.72 (d, <i>J</i> = 4.3 Hz, 1H), 7.13 (dd, <i>J</i> = 11.9, 3.3 Hz, 2H), 6.89 (s, 1H), 6.41 (s, 1H), 2.32 (s, 3H).
244	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 11.53 (s, 1H), 8.73 (s, 1H), 7.65 (s, 1H), 7.46 (d, <i>J</i> = 8.2 Hz, 1H), 7.14 (dt, <i>J</i> = 23.9, 7.2 Hz, 2H), 6.77 (s, 1H), 6.55 (s, 1H), 6.47 (d, <i>J</i> = 9.1 Hz, 1H), 2.19 (s, 2H).
245	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 8.07 (s, 1H), 7.77 - 7.59 (m, 2H), 7.20 (s, 2H), 7.02 (s, 1H), 6.59 (s, 1H), 6.53 (d, <i>J</i> = 8.7 Hz, 1H), 6.05 (s, 1H), 1.99 (s, 2H), 1.12 (t, <i>J</i> = 7.5 Hz, 3H).
246	¹ H NMR (300 MHz, Chloroform- <i>d</i>) δ 7.74 (q, <i>J</i> = 1.7 Hz, 1H), 7.58 (dt, <i>J</i> = 7.6, 1.5 Hz, 1H), 7.54 - 7.47 (m, 1H), 7.42 (t, <i>J</i> = 7.6 Hz, 1H), 7.13 (d, <i>J</i> = 8.4 Hz, 1H), 7.06 (dd, <i>J</i> = 9.9, 8.9 Hz, 1H), 6.94 (dd, <i>J</i> = 6.3, 3.1 Hz, 1H), 6.83 (dt, <i>J</i> = 8.9, 3.5 Hz, 1H), 6.67 - 6.58 (m, 2H), 6.07 (s, 1H), 3.80 (s, 3H), 2.13 (s, 3H).
256	¹ H NMR (300 MHz, Chloroform- <i>d</i>) δ 8.08 (dd, <i>J</i> = 8.6, 5.0 Hz, 2H), 7.75 (dd, <i>J</i> = 8.7, 5.1 Hz, 2H), 7.59 (d, <i>J</i> = 8.4 Hz, 2H), 7.21 (t, <i>J</i> = 8.5 Hz, 2H), 7.12 - 7.00 (m, 4H), 6.75 (s, 1H), 6.72 - 6.62 (m, 2H), 5.48 (s, 1H), 4.98 (s, 1H), 2.21 (s, 3H).
265	¹ H NMR (300 MHz, Methanol- <i>d</i> ₄) δ 8.01 (s, 2H), 7.66 (d, <i>J</i> = 3.1 Hz, 4H), 7.15 (d, <i>J</i> = 8.5 Hz, 1H), 6.67 (s, 1H), 6.62 (dd, <i>J</i> = 8.6, 2.8 Hz, 1H), 5.99 (s, 1H), 2.25 (s, 3H).
272	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 8.78 (s, 1H), 8.19 (s, 1H), 7.92 (s, 1H), 7.89 (s, 1H), 7.49 (d, <i>J</i> = 6.7 Hz, 2H), 7.39 (d, <i>J</i> = 7.7 Hz, 1H), 7.01 (s, 1H), 6.57 (s,

	1H), 6.53 (d, $J = 8.5$ Hz, 1H), 6.14 (s, 1H), 3.88 (s, 3H), 2.17 (s, 3H).
273	¹ H NMR (500 MHz, Acetone- d_6) δ 7.85 (d, $J = 8.4$ Hz, 2H), 7.63 (dd, $J = 8.3$, 1.8 Hz, 2H), 7.38 (d, $J = 8.5$ Hz, 1H), 7.16 (dd, $J = 10.4$, 8.9 Hz, 1H), 7.06 (dd, $J = 6.4$, 3.2 Hz, 1H), 6.93 (dt, $J = 8.9$, 3.5 Hz, 1H), 6.70 (d, $J = 2.7$ Hz, 1H), 6.64 (dd, $J = 8.6$, 2.8 Hz, 1H), 6.19 (s, 1H), 3.84 (s, 3H), 2.25 (s, 3H).
274	¹ H NMR (300 MHz, DMSO- d_6) δ 12.47 (s, 1H), 10.43 (s, 1H), 8.56 (d, $J = 2.5$ Hz, 1H), 8.42 (s, 1H), 7.92 (d, $J = 8.5$ Hz, 2H), 7.84 (d, $J = 8.5$ Hz, 2H), 7.77 (d, $J = 1.7$ Hz, 1H), 7.14 (s, 1H), 6.90 (d, $J = 6.9$ Hz, 1H), 6.74 (d, $J = 8.4$ Hz, 1H), 6.57 (t, $J = 2.1$ Hz, 1H), 6.24 (s, 1H), 4.49 (s, 2H).
275	¹ H NMR (300 MHz, DMSO- d_6) δ 12.17 (s, 1H), 10.41 (s, 1H), 9.66 (s, 1H), 8.33 (s, 1H), 7.52 (d, $J = 8.4$ Hz, 2H), 7.14 (s, 1H), 6.88 (d, $J = 8.7$ Hz, 1H), 6.81 (d, $J = 8.5$ Hz, 2H), 6.72 (d, $J = 8.5$ Hz, 1H), 6.01 (s, 1H), 4.49 (s, 2H).
276	¹ H NMR (500 MHz, DMSO- d_6) δ 12.21 (s, 1H), 8.79 (s, 1H), 8.17 (s, 1H), 7.90 (s, 1H), 7.67 (d, $J = 8.1$ Hz, 2H), 7.60 (d, $J = 8.1$ Hz, 2H), 7.29 (s, 1H), 7.00 (s, 1H), 6.57 (d, $J = 2.8$ Hz, 1H), 6.52 (dd, $J = 8.6$, 2.9 Hz, 1H), 6.06 (s, 1H), 3.87 (s, 3H), 2.16 (s, 3H).
277	¹ H NMR (500 MHz, DMSO- d_6) δ 8.85 (s, 1H), 7.70 - 7.54 (m, 4H), 7.05 (s, 1H), 6.58 (s, 1H), 6.52 (d, $J = 8.6$ Hz, 1H), 6.07 (s, 1H), 2.15 (s, 3H), 1.24 (s, 1H), 1.07 (s, 1H).
278	¹ H NMR (300 MHz, Acetone- d_6) δ 7.75 - 7.68 (m, 2H), 7.61 - 7.54 (m, 2H), 7.30 (d, $J = 8.5$ Hz, 1H), 6.71 (d, $J = 2.8$ Hz, 1H), 6.68 - 6.58 (m, 2H), 6.12 (s, 1H), 2.24 (s, 3H).
279	¹ H NMR (300 MHz, Acetone- d_6) δ 7.72 (d, $J = 8.4$ Hz, 2H), 7.57 (d, $J = 8.4$ Hz, 2H), 7.28 (d, $J = 8.5$ Hz, 1H), 6.74 (d, $J = 2.8$ Hz, 1H), 6.66 (dd, $J = 8.5$, 2.9 Hz, 1H), 6.59 (s, 1H), 6.08 (s, 1H), 2.65 (q, $J = 7.5$ Hz, 2H), 1.19 (t, $J = 7.5$ Hz, 3H).
280	¹ H NMR (300 MHz, Acetone- d_6) δ 11.67 (s, 1H), 8.37 (s, 1H), 7.77 - 7.71 (m, 2H), 7.64 - 7.57 (m, 2H), 7.53 (d, $J = 8.6$ Hz, 1H), 7.38 - 7.34 (m, 1H), 7.33 - 7.27 (m, 1H), 6.78 (s, 1H), 6.31 (s, 1H), 3.69 (s, 3H), 2.29 (s, 3H).
281	¹ H NMR (300 MHz, Acetone- d_6) δ 8.39 (s, 1H), 7.79 - 7.70 (m, 2H), 7.65 - 7.58 (m, 2H), 7.53 (d, $J = 8.6$ Hz, 1H), 7.36 (d, $J = 2.5$ Hz, 1H), 7.30 (dd, $J = 8.8$, 2.6 Hz, 1H), 6.80 (s, 1H), 6.31 (s, 1H), 3.69 (s, 3H), 2.29 (s, 3H).
282	¹ H NMR (300 MHz, DMSO- d_6) δ 12.45 (s, 1H), 9.01 (s, 1H), 7.65 (d, $J = 5.3$ Hz, 5H), 7.29 (s, 1H), 6.71 (d, $J = 9.9$ Hz, 1H), 6.24 (s, 1H), 2.14 (s, 3H).
283	¹ H NMR (300 MHz, Chloroform- d) δ 7.73 - 7.64 (m, 2H), 7.62 - 7.53 (m, 2H),

	7.52 - 7.43 (m, 1H), 7.35 (d, $J = 8.6$ Hz, 1H), 7.20 (s, 1H), 7.10 (t, $J = 9.3$ Hz, 2H), 6.94 (dd, $J = 6.4, 3.2$ Hz, 1H), 6.86 (dt, $J = 8.8, 3.5$ Hz, 1H), 6.65 (s, 1H), 6.28 (s, 1H), 5.80 (s, 1H), 3.85 (s, 3H), 3.79 (s, 3H), 2.20 (d, $J = 5.8$ Hz, 3H).
284	^1H NMR (300 MHz, Chloroform- d) δ 7.66 (d, $J = 8.2$ Hz, 2H), 7.51 (dd, $J = 8.4, 1.6$ Hz, 2H), 7.32 (d, $J = 8.6$ Hz, 1H), 7.15 (s, 1H), 7.13 - 7.04 (m, 2H), 6.91 (dd, $J = 6.3, 3.1$ Hz, 1H), 6.85 (dt, $J = 8.9, 3.5$ Hz, 1H), 6.75 (s, 1H), 6.26 (s, 1H), 5.83 (s, 1H), 3.83 (s, 3H), 3.78 (s, 3H), 2.14 (s, 3H).
285	^1H NMR (300 MHz, DMSO- d_6) δ 12.24 (s, 1H), 9.26 (s, 1H), 7.63 (d, $J = 8.4$ Hz, 2H), 7.56 (s, 1H), 7.15 (t, $J = 9.2$ Hz, 3H), 6.99 (d, $J = 8.4$ Hz, 2H), 6.14 (s, 1H), 3.78 (s, 3H), 3.62 (s, 3H), 2.20 (s, 3H).
286	^1H NMR (300 MHz, DMSO- d_6) δ 12.22 (s, 1H), 8.92 (s, 1H), 7.62 (d, $J = 8.6$ Hz, 3H), 7.51 - 7.32 (m, 5H), 7.19 (s, 1H), 7.07 (d, $J = 8.4$ Hz, 2H), 6.69 (d, $J = 9.9$ Hz, 1H), 6.12 (s, 1H), 5.15 (s, 2H), 2.14 (s, 3H).
287	^1H NMR (300 MHz, DMSO- d_6) δ 12.24 (s, 1H), 9.26 (s, 1H), 7.63 (d, $J = 8.4$ Hz, 2H), 7.56 (s, 1H), 7.15 (t, $J = 9.2$ Hz, 3H), 6.99 (d, $J = 8.4$ Hz, 2H), 6.14 (s, 1H), 3.78 (s, 3H), 3.62 (s, 3H), 2.20 (s, 3H).
288	^1H NMR (300 MHz, DMSO- d_6) δ 12.32 (s, 1H), 9.31 (s, 1H), 7.63 (d, $J = 8.4$ Hz, 2H), 7.56 (s, 1H), 7.15 (t, $J = 9.2$ Hz, 3H), 6.99 (d, $J = 8.4$ Hz, 2H), 6.40 (s, 1H), 3.63 (s, 3H), 3.24 (s, 3H), 2.20 (s, 3H).
289	^1H NMR (300 MHz, DMSO- d_6) δ 12.33 (s, 1H), 9.32 (s, 1H), 7.63 (d, $J = 8.4$ Hz, 2H), 7.56 (s, 1H), 7.15 (t, $J = 9.2$ Hz, 3H), 6.99 (d, $J = 8.4$ Hz, 2H), 6.40 (s, 1H), 3.64 (s, 3H), 3.24 (s, 3H), 2.21 (s, 3H).
290	^1H NMR (500 MHz, Acetone- d_6) δ 8.41 (s, 1H), 7.73 (dd, $J = 8.6, 2.0$ Hz, 2H), 7.59 (dt, $J = 8.6, 2.0$ Hz, 2H), 7.52 (d, $J = 8.6$ Hz, 1H), 7.36 (s, 1H), 7.31 (d, $J = 8.7$ Hz, 1H), 6.81 (d, $J = 5.3$ Hz, 1H), 6.31 (d, $J = 1.1$ Hz, 1H), 3.69 (d, $J = 1.1$ Hz, 3H), 2.29 (s, 3H).
291	^1H NMR (500 MHz, Acetone- d_6) δ 8.43 (s, 1H), 7.75 - 7.70 (m, 2H), 7.58 (ddd, $J = 8.4, 3.8, 2.2$ Hz, 2H), 7.52 (d, $J = 8.6$ Hz, 1H), 7.36 (s, 1H), 7.32 (d, $J = 8.7$ Hz, 1H), 6.84 (s, 1H), 6.32 (s, 1H), 3.70 (d, $J = 1.1$ Hz, 3H), 2.28 (s, 3H).
292	^1H NMR (300 MHz, Acetone- d_6) δ 8.38 (s, 1H), 7.80 - 7.69 (m, 2H), 7.59 (d, $J = 8.5$ Hz, 2H), 7.51 (d, $J = 8.6$ Hz, 1H), 7.40 - 7.27 (m, 2H), 6.31 (s, 1H), 3.69 (s, 3H), 2.29 (s, 3H).
293	^1H NMR (300 MHz, Acetone- d_6) δ 8.39 (s, 1H), 7.78 - 7.69 (m, 2H), 7.63 - 7.55 (m, 2H), 7.50 (d, $J = 8.6$ Hz, 1H), 7.41 - 7.27 (m, 2H), 6.31 (s, 1H), 3.69

	(s, 3H), 2.28 (s, 3H).
294	¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ 12.38 (s, 1H), 9.28 (s, 1H), 8.18 (s, 1H), 7.91 (s, 1H), 7.65 (d, <i>J</i> = 29.7 Hz, 5H), 7.20 (d, <i>J</i> = 27.7 Hz, 3H), 6.25 (s, 1H), 3.88 (s, 3H), 3.64 (s, 3H), 2.21 (s, 3H).
295	¹ H NMR (500 MHz, Chloroform- <i>d</i>) δ 7.73 (d, <i>J</i> = 8.0 Hz, 2H), 7.63 (d, <i>J</i> = 8.2 Hz, 2H), 7.15 (d, <i>J</i> = 8.5 Hz, 1H), 6.71 (s, 1H), 6.67 (d, <i>J</i> = 9.3 Hz, 1H), 6.07 (s, 1H), 2.21 (s, 4H).
296	¹ H NMR (500 MHz, Chloroform- <i>d</i>) δ 7.66 (d, <i>J</i> = 8.0 Hz, 2H), 7.51 (d, <i>J</i> = 8.0 Hz, 2H), 7.19 (d, <i>J</i> = 8.5 Hz, 1H), 7.06 (t, <i>J</i> = 32.9 Hz, 3H), 6.18 (s, 1H), 3.76 (s, 3H), 2.06 (d, <i>J</i> = 1.6 Hz, 5H).
297	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.22 (s, 1H), 10.05 (s, 1H), 9.26 (s, 1H), 7.50 (dd, <i>J</i> = 12.6, 2.1 Hz, 2H), 7.34 (d, <i>J</i> = 8.3 Hz, 1H), 7.15 (t, <i>J</i> = 10.6 Hz, 3H), 6.97 (t, <i>J</i> = 8.8 Hz, 1H), 6.14 (s, 1H), 3.62 (s, 3H), 2.49 (s, 8H), 2.19 (s, 3H).
298	¹ H NMR (300 MHz, DMSO- <i>d</i> 6) δ 12.45 (s, 1H), 9.28 (s, 1H), 7.74 (d, <i>J</i> = 8.2 Hz, 2H), 7.48 (d, <i>J</i> = 8.2 Hz, 3H), 7.28 (s, 1H), 7.15 (d, <i>J</i> = 11.2 Hz, 2H), 6.26 (s, 1H), 3.63 (s, 3H), 2.20 (s, 3H).
299	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 12.45 (s, 1H), 9.28 (s, 1H), 7.74 (d, <i>J</i> = 8.2 Hz, 2H), 7.48 (d, <i>J</i> = 8.2 Hz, 3H), 7.28 (s, 1H), 7.15 (d, <i>J</i> = 11.2 Hz, 2H), 6.26 (s, 1H), 3.63 (s, 3H), 2.20 (s, 3H).

[618]

[619] **Evaluation of Compounds**[620] 1. Efficacy of inhibiting TNIK activity, *in vitro* TNIK kinase assay using qPCR

[621] Kinase-tagged T7 phage strains were grown in parallel in an *E. coli* host derived from the BL21 strain. *E. coli* were grown to log-phase and infected with T7 phage from a frozen stock (multiplicity of infection = 0.4) and incubated with shaking at 32°C until (90-150minutes). The lysates were centrifuged (6,000 x g) and filtered (0.2 μm) to remove cell debris. The remaining kinases were produced in HEK-293 cells and subsequently tagged with DNA for qPCR detection. Streptavidin-coated magnetic beads were treated with biotinylated small molecule ligands for 30 minutes at room temperature to generate affinity resins for kinase assays. The liganded beads were blocked with excess biotin and washed with blocking buffer (SeaBlock (Pierce), 1 % BSA, 0.05 % Tween 20, 1 mM DTT) to remove unbound ligand and to reduce non-specific phage binding. Binding reactions were assembled by combining kinases, liganded affinity beads, and test compounds in 1x binding buffer (20 % SeaBlock, 0.17x PBS, 0.05 % Tween 20, 6 mM DTT). Test compounds were prepared as 40x stocks in 100% DMSO

and directly diluted into the assay. All reactions were performed in polypropylene 384-well plates in a final volume of 0.04 ml.

[622] The assay plates were incubated at room temperature with shaking for 1 hour and the affinity beads were washed with wash buffer (1x PBS, 0.05 % Tween 20). The beads were then re-suspended in elution buffer (1x PBS, 0.05 % Tween 20, 0.5 μ M non-biotinylated affinity ligand) and incubated at room temperature with shaking for 30 minutes. The kinase concentration in the eluates was measured by qPCR.

[623] Results are shown in Tables 4 and 5 below.

[624]

[Table 4]

Compound No.	TNIK activity at 1 μ M (%)	Compound No.	TNIK activity at 1 μ M (%)
2	2	61	5
3	2	62	1
5	47	67	21
6	36	68	36
9	47	70	14
12	1	71	34
13	15	73	0
14	1	74	1
15	36	75	4
16	8	76	3
17	34	77	3
18	19	79	4
19	9	80	1
20	36	81	0
21	22	82	2
22	2	83	3
23	5	85	0
24	0	86	1
25	5	87	4
27	8	88	0
30	33	89	0
31	6	90	7
32	1	91	20
33	0	92	0
34	1	93	37
35	14	94	1
36	6	96	16
37	5	98	4

38	1	99	12
39	0	100	8
40	0	101	3
41	13	102	2
42	8	103	5
43	1	108	5
44	5	109	0
45	4	110	7
46	2	111	0
47	44	112	20
48	28	113	6
49	0	114	4
50	2	115	4
51	1	116	7
52	7	117	3
53	0	118	2
54	1	119	2
55	1	121	36
56	2	122	9
57	17	123	34
58	0		
59	0		
60	17		

[625]

[Table 5]

Compound No.	Kd (nM)
12	29
24	14
29	25
34	95
39	6
40	9
49	19
58	11
59	16
62	8
73	14
81	3
88	16
89	21
94	12
111	20

[626] As shown in Tables 4 and 5 above, the compounds of the present disclosure were very effective in inhibiting TNIK activity.

[627]

[628] 2. Efficacy of inhibiting TNIK activity, *in vitro* TNIK kinase assay using ADP-Glo™ kinase assay system

[629] The inhibitory properties of compounds were evaluated with TNIK kinase enzyme system and luminescent ADP-Glo™ Kinase Assay from Promega Corporation according to the manufacturer's protocol. The tested compounds were incubated with TNIK kinases. Reactions were performed in 10 μ l kinase buffer supplemented with reaction mixture containing 2 μ l ATP, 2 μ l MBP protein and 2 μ l TNIK at 37°C for 30 min. The reactions' conditions are provided in Table 5. After kinase reaction, 5 μ l reaction mixtures were transferred to 384 well assay plate (Greiner, solid white low-binding plates). Next, 5 μ l of ADP-Glo Reagent was added to each well and incubated at 37°C for 30min. After 45 min, 10 μ l Kinase Detection Reagent was added to each well and luminescence signal was detected with the SpectraMax M5e Microplate

Reader (Molecular Devices, Menlo Park, CA) after 30 min at 37 °C incubation. The IC₅₀ values were calculated using GraphPad Prism software (GraphPad Software Inc., La Jolla, CA, USA) to determine kit performance. To the cut-off assay, the inhibitory activity of the compounds against TNIK kinase was expressed as the percentage of the kinase inhibitory activity for an indicated concentration of inhibitor. The IC₅₀ of the compounds against TNIK kinase values were calculated a log-concentration-response curve fitted with a four-parameter logistic equation and expressed dose-response curve for % of inhibitory activity versus log of the compound concentration.

[630] The reactions' conditions to be used *in vitro* TNIK kinase inhibitory assay are shown in Table 6.

[631] [Table 6]

TNIK kinase(ng/reaction)	MBP substrate(μg/reaction)	ATP(μM)	Other reagents
13	1	100	Kinase buffer (40mM Tris (pH7.5), 20mM MgCl ₂ , 0.1mg/ml BSA)

[632] Results of *in vitro* TNIK kinase assay (HTRF) are shown in Tables 7 and 8 below.

[633]

[Table 7]

Compound No.	Inhibition at 1 μ M(%)	Compound No.	Inhibition at 1 μ M(%)
33	97	201	88
36	85	202	90
58	98	203	87
74	88	204	76
77	87	205	98
80	89	206	95
81	100	207	100
85	92	208	100
86	93	209	99
96	69	210	90
97	66	212	76
109	97	213	74
110	95	214	66
114	100	215	97
115	98	216	100
116	96	217	98
117	94	218	97
119	100	219	56
126	94	220	78
127	100	221	91
129	61	222	60
132	64	223	88
133	63	224	64
135	72	226	97
136	79	227	91
138	100	228	90
139	89	229	97
140	94	231	74

141	92	234	71
142	98	237	99
143	73	240	96
144	89	241	78
146	100	242	97
147	89	243	96
148	61	244	92
149	59	245	90
150	87	246	80
151	92	265	93
152	94	272	84
153	50	273	92
154	85	274	87
155	100	275	93
156	72	276	99
157	98	277	99
158	99	278	99
160	99	279	99
161	95	280	96
164	50	281	98
165	63	282	99
169	82	283	91
170	78	284	86
171	81	285	88
172	83	286	91
173	83	287	54
174	89	288	73
175	91	289	72
176	92	290	91
177	80	291	93
178	97	292	90

180	99	293	88
181	96	294	92
182	98	295	88
183	93	296	53
185	73	297	91
188	89	298	88
189	89	299	95
190	100		
191	82		
192	90		
193	100		
194	94		
195	88		
196	78		
197	96		
198	96		
199	99		
200	89		

[634]

[Table 8]

Compound No.	IC ₅₀ (nM)	Compound No.	IC ₅₀ (nM)
33	68	201	89
58	3	202	83
74	97	203	71
77	94	204	432
80	95	205	6
81	12	206	39
85	23	207	8
86	12	208	7
109	1	209	9
110	8	210	252
114	72	212	471
115	107	213	645
117	10	214	646
119	1	215	24
126	17	216	41
133	43	217	113
138	23	218	127
139	22	219	789
140	6	220	534
141	148	221	341
142	77	222	749
144	103	223	431
146	1	224	703
147	47	226	250
151	45	227	431
152	115	228	397
154	493	229	179
155	73	231	578
156	80	234	431

157	116	237	389
158	23	240	443
160	26	241	684
161	182	242	230
162	40	243	103
170	15	244	106
171	16	245	82
172	33	246	236
173	32	265	4
174	46	272	117
175	15	273	34
176	8	274	20
178	89	275	19
180	1	276	8
181	2	277	11
182	7	278	9
183	10	279	37
185	408	280	66
187	3	281	25
188	177	282	41
189	125	283	48
190	38	284	113
191	314	285	161
192	271	286	97
193	140	287	919
194	195	288	506
195	236	289	623
196	516	290	64
197	105	291	41
198	125	292	108
199	112	293	157

200	91	294	143
		295	264
		296	745
		297	243
		298	128
		299	54

[635]

[636] 3. Efficacy of inhibiting cancer cells using colon cancer cell lines

[637] 1) Cell Culture

[638] The human colon cancer cell line SW480 (Catalog No. CCL-228) and SW620 (Catalog No. CCL-227) were obtained from the American Type Culture Collection (ATCC), and maintained in DMEM (Thermo Fisher Scientific Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS) (Thermo Fisher Scientific Inc., Waltham, MA), 100 U/ml penicillin, and 100 µg/ml streptomycin (Gibco, Gaithersburg, MD). Cells were grown at 37 °C in 5% CO₂ for confluence.

[639] 2) Cell Viability Assay

[640] Cell viability was measured using the Cell Counting Kit-8 (Dojindo Molecular Technologies). SW480 and SW620 cells (2.5X10⁴ cells/well) were seeded in 96-well plates and incubated for 24 h. After incubation, the cells were exposed to serially diluted compounds (0.1, 0.3, 1, 3, 10 and 30 µM) in 100 µl phenol free DMEM medium containing 0.1% FBS. After 48 h, CCK-8 reagent was added at 10 µl per well and incubated for 1 h at 37 °C. Optical density was measured at 450 nm using a microplate reader (Bio-Rad Laboratories, Hercules, CA). The IC₅₀ values were calculated using GraphPad Prism software (GraphPad Software Inc., La Jolla, CA, USA) to determine kit performance. Experiments were performed in triplicate.

[641] Results of cell viability assay using SW620 cell are shown in Table 9 below.

[642]

[Table 9]

Compound No	SW620, IC ₅₀ (μM)	Compound No	SW620, IC ₅₀ (μM)
2	9.1	202	3.0
6	1.8	205	11.5
19	1.0	207	22.1
24	11.4	208	16.8
32	1.2	212	18.0
40	12.5	215	12.6
45	0.4	216	8.2
46	0.2	226	19.9
56	0.4	229	28.0
58	47.5	244	11.2
80	0.5	245	11.7
81	1.7	246	11.0
85	1.0	265	6.7
94	0.8	272	17.9
110	15.9	273	1.8
114	0.1	275	16.9
115	0.2	276	8.6
116	0.3	277	3.1
117	0.4	278	9.0
119	1.0	279	12.5
138	0.7	280	19.6
140	0.9	282	27.1
146	0.4	285	8.1
154	25.5	286	7.2
160	1.0	288	15.3
190	11.2	290	9.4
193	1.8	291	10.7
194	1.0	292	4.0
199	14.9	293	7.8

200	14.9	294	18.9
		295	10.1
		296	14.0
		297	12.1
		298	22.5
		299	6.3

[643] Results of cell viability assay using SW480 cell are shown in Table 10 below.

[644]

[Table 10]

Compound No.	SW480, IC ₅₀ (μM)	Compound No.	SW480, IC ₅₀ (μM)
2	2.6	202	5.4
6	2.9	205	13.4
18	18.7	206	13.4
19	1.0	207	11.3
24	8.2	208	11.0
32	5.4	212	8.9
40	16.6	215	4.0
45	0.9	216	10.9
46	0.6	229	19.6
56	18.9	244	5.9
80	1.5	245	11.4
81	5.7	246	9.5
85	8.2	265	37.3
94	13.5	272	18.6
110	1.5	273	6.8
114	1.3	276	10.1
115	0.7	277	6.4
116	9.1	278	7.4
117	8.2	279	19.9
119	4.4	280	21.9
127	3.1	282	11.5
138	2.8	283	28.1
140	5.5	285	13.1
146	1.3	286	6.4
160	10.0	290	8.3
190	8.1	291	9.1
193	11.5	292	2.3
194	10.6	293	4.6
199	12.9	294	4.2

		295	5.2
		296	8.7
		297	7.6
		298	7.7
		299	3.0

[645]

[646] 4. Efficacy of inhibiting Wnt / β -catenin signaling

[647] 1) Cell culture

[648] CHO-K1 (KCLB no. 10061) cells were cultured (37 °C, 5% CO₂) in RPMI medium (Gibco) containing 10% FBS (Gibco) and 1% penicillin/streptomycin (Gibco) of a T75 flask. The culture was washed with DPBS (Dulbecco's Phosphate-Beffered Saline, Gibco) and 2 ml of 0.05% trypsin-EDTA (trypsin-EDTA, Gibco) was added and cultured for one minute. Next, cell suspension was centrifuged at 1,500 rpm for 2 minutes, and cell pellets were obtained for next steps.

[649] 2) Reporter gene transfection

[650] Cultured CHO-K1 cells were diluted with the cell culture medium to become 4 X 10⁴ cells/cm² and then seeded into a culture plate. The plate was cultured overnight at 37 in a 5% CO₂ condition. Next day, the transfection of reporter gene was performed as follows: 1.35 μ g of reporter DNA (M50 Super 8XTOPFlash, no. 12456, Addgene) and lipofectamine 2000 (Lipofectamine 2000 Reagent, no. 11668, Invitrogen) were diluted in an Opti-MEM medium (Gibco) to make a solution for transfection according to the protocol of those reagents. The cell medium of the CHO-K1 cells attached by overnight culture was replaced with RPMI medium not containing FBS and penicillin/streptomycin. Then the cells were transfected with the solution and cultured (37 °C, 5% CO₂) for 5 hours. After that, the transfected cells were cultured overnight in RPMI medium containing 1% FBS and 1% penicillin/streptomycin to stabilize the transfected cells. The stabilized CHO-K1 cells were diluted with RPMI medium containing 1% FBS and seeded to be 2.5X10⁴ cells in a 96 well plate. The plate was cultured overnight to attach cells. The attached cells were used for evaluating reporter activity and cell toxicity next day.

[651] 3) Addition of test compounds and recombinant Wnt3a protein

[652] DMSO solutions of test compounds were diluted with the culture medium containing 1% FBS to make the solutions having 10 times concentration of the tested concentration. 0.1 volumes of those solutions were added to the transfected CHO-K1 cells, and cultured overnight at 37°C, 5% CO₂. Next day, 40 μ g/ml of recombinant Wnt3a protein (Mouse Recombinant Wnt3a, no.1324-WN, R&D systems) was diluted to 200

ng/ml with RPMI medium containing 1% FBS. 0.1 volumes of the diluted Wnt3a protein solution were added to CHO-K1 cells containing test compounds. Next, the CHO-K1 cells were further cultivated (37°C, 5% CO₂) for 7 hours.

[653] 4) Determination of luciferase activity

[654] Luciferase assay system (no. E1960, Promega) was used. The activity of luciferase in the cells was determined by the microplate reader (SpectraMax M5e Multi-Mod Microplate Reader, Molecular Devices). The luminescence values were normalized with respect to cell viability, and IC₅₀ values of test compounds were calculated based on both the luminescence intensity of cells stimulated by Wnt3a without test compounds (100%) and the luminescence intensity of cells without test compounds and Wnt3a stimulus (0%).

[655] 5) Cell toxicity evaluation of test compounds

[656] DMSO solutions of test compounds were diluted with the culture medium containing 1% FBS to make the solutions having 10 times concentration of the tested concentration. 0.1 volumes of those solutions were added to the transfected CHO-K1 cells, and cultured overnight at 37°C, 5% CO₂. Next day, 40 µg/ml of recombinant Wnt3a protein (Mouse Recombinant Wnt3a, no.1324-WN, R&D systems) was diluted to 200 ng/ml with RPMI medium containing 1% FBS. 0.1 volumes of the diluted Wnt3a protein solution were added to CHO-K1 cells containing test compounds. Next, the CHO-K1 cells were further cultivated (37°C, 5% CO₂) for 7 hours. After that, 10 µl of CCK-8 solution (Cell Counting Kit-8, Dojindo Molecular Technologies) was added and cultured (37°C, 5% CO₂) for 1 hour. Cell viability in wells then was determined at 450 nm by the microplate reader (SpectraMax M5e Multi-Mod Microplate Reader, Molecular Devices). The determined cell viability was used for normalization of luciferase activity.

[657] Results are shown in Tables 11 and 12 below.

[658]

[Table 11]

Compound No.	Inhibition at 10 μ M (%)
114	92
115	92
119	82
138	83
140	68
146	89
193	30
194	21
216	82
222	33

[659] [Table 12]

Compound No.	IC ₅₀ (μ M)
81	3.1
114	1.5
115	1.9
119	2.8
138	2.1
140	7.0
146	1.0
216	6.1

[660]

[661] 5. Mouse xenograft model test

[662] Test 5-1

[663] Male nude mice grafted with SW620 cell which is colorectal carcinoma cell derived from human were used. Test compounds were administered several times and inhibitory effect of tumor by test compounds was evaluated.

[664] The following groups were set up: 0 mg/kg dose of negative control group, 50 mg/kg dose of Compound **81** group, 75 mg/kg dose of Compound **58** group, 50 mg/kg dose of Compound **80** group, 75 mg/kg dose of Compound **110** group, 40 mg/kg dose of irinotecan group (positive control). The number of each group was six. Compound **81**

was administered two times a day for 28 days, and other compounds were administered once a day for 28 days. The results are shown in Figure 1.

- [665] The average tumor volume of 50 mg/kg dose of Compound **81** group showed significant difference in comparison to negative control at day 15, 18, 22, 25 and 29 after administration. The inhibitory rate of tumor growth was 72.3 %, which means that the compound has a determinate effect of inhibiting tumor growth.
- [666] The average tumor volume of 75 mg/kg dose of Compound **58** group showed significant difference in comparison to negative control at day 15, 18, 22 and 25 after administration. The inhibitory rate of tumor growth was 58.5 %, which means that the compound has a determinate effect of inhibiting tumor growth.
- [667] The average tumor volume of 50 mg/kg dose of Compound **80** group showed significant difference in comparison to negative control at day 15, 18, and 25 after administration. The inhibitory rate of tumor growth was 48.8 %, which means that the compound has an effect of inhibiting tumor growth.
- [668] The average tumor volume of 75 mg/kg dose of Compound **110** group showed significant difference in comparison to negative control at day 15, 18, 22, 25 and 29 after administration. The inhibitory rate of tumor growth was 58.0 %, which means that the compound has a determinate effect of inhibiting tumor growth.
- [669] The average tumor volume of 40 mg/kg dose of positive control (Irinotecan) group showed significant difference in comparison to negative control at day 15, 18, 22, 25 and 29 after administration. The inhibitory rate of tumor growth was 81.8 %, which means that the compound has a determinate effect of inhibiting tumor growth.
- [670] In anti-cancer effect tests against colorectal carcinoma cell derived from human, SW620, 50 mg/kg dose of Compound **81**, 75 mg/kg dose of Compound **58** and Compound **110** showed an effect of inhibiting growth of the tumor.

[671]

[672] Test 5-2

[673] The effect of inhibiting the growth of tumor after administering the compounds of the present invention was evaluated. Test compounds were repeatedly administered to male nude mice grafted with colorectal carcinoma cell, SW620, derived from human.

[674] The following groups were set up: 0 mg/kg dose of negative control group, 100 mg/kg dose of Compound **81** group (oral administration), 100 mg/kg dose of Compound **81** group (intraperitoneal injection), 50, 100 and 160 mg/kg dose of Compound **138** group, 40 mg/kg dose of irinotecan group (positive control), a combined administration group of 100 mg/kg dose of Compound **81** and 40 mg/kg dose of irinotecan, and a combined administration group of 100 mg/kg dose of Compound **138** and 40 mg/kg dose of Irinotecan. The number of each group was six. Compounds **81** and **138** were administered two times a day for 15 days, and irinotecan was administered once a

week for 15 days. The results are shown in Figures 2 and 3.

[675] The average tumor volume of 100 mg/kg of oral administration of Compound **81** group showed significant difference in comparison to negative control. The inhibitory rate of tumor growth was 49.3 %.

[676] The average tumor volume of 100 mg/kg of intraperitoneal injection of Compound **81** group showed significant difference in comparison to negative control. The inhibitory rate of tumor growth was 53.2 %.

[677] The average tumor volumes of 50, 100 and 160 mg/kg dose of Compound **138** group showed significant difference in comparison to negative control. The inhibitory rates of tumor growth were 54.5, 64.7 and 57.8%.

[678] The average tumor volume of 40 mg/kg of irinotecan group showed significant difference in comparison to negative control. The inhibitory rate of tumor growth was 69.0%.

[679] The average tumor volume of the combined administration group of 100 mg/kg dose of Compound **81** and 40 mg/kg dose of irinotecan showed significant difference in comparison to negative control. The inhibitory rate of tumor growth was 78.8%.

[680] The average tumor volume of the combined administration group of 100 mg/kg dose of Compound **138** and 40 mg/kg dose of irinotecan showed significant difference in comparison to negative control. The inhibitory rate of tumor growth was 78.0%.

[681] In conclusion, in anti-cancer effect tests against colorectal carcinoma cell derived from human, SW620, Compound **81**, Compound **138** and irinotecan clearly showed an effect of inhibiting growth of the tumor. And, the combined administration of irinotecan and Compound **81** or **138** showed a synergic inhibitory effect.

[682]


[683] All mentioned documents are incorporated by reference as if herein written. When introducing elements of the present invention or the exemplary embodiment(s) thereof, the articles "a," "an," "the" and "said" are intended to mean that there are one or more of the elements. The terms "comprising," "including" and "having" are intended to be inclusive and mean that there may be additional elements other than the listed elements. Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

Claims

[Claim 1]

A compound of Chemical Formula 1:

[Chemical Formula 1]



or a pharmaceutically acceptable salt thereof,

in the Chemical Formula 1

Y is N or CH,

Z is N or C-V,

A is H, halogen, -OH, -CO₂-C₁₋₆ alkyl, -CO₂H, -CN, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -OR¹, -NH₂, -NHR², -substituted or unsubstituted piperazine, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,

B is H, -C₁₋₆ haloalkyl, C₁₋₆ alkyl, halogen, or C₁₋₆ alkoxy,

V is H, -CH₂OH, halogen, -CO₂H, -CO₂-C₁₋₆ alkyl, -OH, -NH₂, phenoxy, or -NHCO-C₁₋₆ alkyl,

X is H or F,

W is substituted or unsubstituted, aromatic ring, heteroaryl, or fused heteroaryl,

wherein,

R¹ is C₁₋₆ alkyl, benzyl, C₁₋₆ haloalkyl, or phenyl,

R² is C₁₋₆ alkyl, C₁₋₆ haloalkyl, -CH₂CH₂-morpholin, or phenyl,

R³ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, or substituted or unsubstituted phenyl,

R⁴ is C₁₋₆ alkyl, C₁₋₆ haloalkyl, -CH₂CH₂Cl, -CH₂CH₂NMe₂, -CH₂NMe₂, or -CH₂CH₂-morpholin.

[Claim 2]

The compound of Claim 1, in the Chemical Formula 1,

Y is N or CH,

Z is N or C-V,

A is H, halogen, -OH, -CO₂-C₁₋₃ alkyl, -CO₂H, -CN, -C₁₋₃ alkyl, -OR¹, -NH₂, -NHR², -substituted or unsubstituted piperazine, -NHSO₂R³, -NHCO₂-C₁₋₆ alkyl, -NHCON-C₁₋₆ alkyl, or -NHCOR⁴,

B is H, -C₁₋₃ haloalkyl, C₁₋₃ alkyl, halogen, or C₁₋₃ alkoxy,

V is H, $-\text{CH}_2\text{OH}$, halogen, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{C}_{1-3}$ alkyl, $-\text{OH}$, $-\text{NH}_2$, phenoxy, or $-\text{NHCO}-\text{C}_{1-3}$ alkyl,

X is H or F,

W is substituted or unsubstituted, phenyl, pyridyl, thiophene, thiazole, pyrrole, benzothiophene, indole, oxazole, pyrazole, imidazole, pyrimidine, benzopyrazole, benzothiazole, benzoxazole, benzimidazole, or benzothiophene,

wherein,

R^1 is benzyl, C_{1-3} haloalkyl, or phenyl,

R^2 is CF_3 , C_{1-3} alkyl, $-\text{CH}_2\text{CH}_2$ -morpholin, or phenyl,

R^3 is C_{1-3} alkyl, or substituted or unsubstituted phenyl,

R^4 is C_{1-3} alkyl, CF_3 , $-\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}_2\text{NMe}_2$, $-\text{CH}_2\text{NMe}_2$, or $-\text{CH}_2\text{CH}_2$ -morpholin.

[Claim 3]

The compound of Claim 1, in the Chemical Formula 1,

Y is N or CH,

Z is C-V,

A is $-\text{OH}$, $-\text{CO}_2\text{C}_{1-2}$ alkyl, $-\text{methyl}$, $-\text{OR}^1$, $-\text{NH}_2$, $-\text{NHR}^2$, $-\text{substituted}$ or $-\text{unsubstituted}$ piperazine, $-\text{NHSO}_2\text{R}^3$, $-\text{NHCO}_2\text{C}_{1-6}$ alkyl, $-\text{NHCON}-\text{C}_{1-6}$ alkyl, or $-\text{NHCOR}^4$,

B is H, $-\text{C}_{1-3}$ haloalkyl, C_{1-3} alkyl, halogen, or C_{1-3} alkoxy,

V is H, $-\text{CH}_2\text{OH}$, F, $-\text{CO}_2\text{H}$, $-\text{CO}_2\text{C}_{1-2}$ alkyl, $-\text{OH}$, $-\text{NH}_2$, phenoxy, or $-\text{NHCOCH}_3$,

X is H or F,

W is substituted or unsubstituted, phenyl, pyridyl, thiophene, thiazole, pyrrole, benzothiophene, or indole,

wherein,

R^1 is benzyl, CF_3 , or phenyl,

R^2 is CF_3 , C_{1-3} alkyl, $-\text{CH}_2\text{CH}_2$ -morpholin, or phenyl,

R^3 is C_{1-3} alkyl, or substituted or unsubstituted phenyl,

R^4 is C_{1-3} alkyl, CF_3 , $-\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}_2\text{NMe}_2$, $-\text{CH}_2\text{NMe}_2$, or $-\text{CH}_2\text{CH}_2$ -morpholin.

[Claim 4]

The compound of Claim 3, in the Chemical Formula 1,

Y is CH,

Z is C-V,


A is $-\text{OH}$, $-\text{NHR}^2$, $-\text{NHSO}_2\text{R}^3$, $-\text{NHCO}_2\text{C}_{1-6}$ alkyl, $-\text{NHCON}-\text{C}_{1-6}$ alkyl, or $-\text{NHCOR}^4$,

B is H, $-\text{C}_{1-3}$ haloalkyl, C_{1-3} alkyl, halogen, or C_{1-3} alkoxy,

V is H, $-\text{CH}_2\text{OH}$, F, $-\text{OH}$, or $-\text{NHCOCH}_3$,

X is H or F,

W is



wherein,

R² is CF₃, C₁₋₃ alkyl, -CH₂CH₂-morpholin, or phenyl,

R^3 is C_{1-3} alkyl, or substituted or unsubstituted phenyl,

R^4 is C_{1-3} alkyl, CF_3 , $-CH_2CH_2Cl$, $-CH_2CH_2NMe_2$, $-CH_2NMe_2$, or $-CH_2CH_2$ -morpholin.

[Claim 5]

The compound of Claim 4, in the Chemical Formula 1,

Y is CH,

Z is C-V,


A is $-OH$, $-NHR^2$, $-NH SO_2 R^3$, $-NH CO_2 C_{1-6}$ alkyl, $-NH CON-C_{1-6}$ alkyl, or $-NHCOR^4$,


B is H, $-C_{1-3}$ haloalkyl, C_{1-3} alkyl, halogen, or C_{1-3} alkoxy,

V is H, F, $-OH$, or $-NH COCH_3$,

X is H or F,

W is





wherein,

R^2 is CF_3 , or C_{1-3} alkyl,

R^3 is C_{1-3} alkyl,

R^4 is C_{1-3} alkyl, CF_3 , $-CH_2CH_2Cl$, $-CH_2CH_2NMe_2$, $-CH_2NMe_2$, or $-CH_2CH_2$ -morpholin.

[Claim 6]

The compound of Claim 5, in the Chemical Formula 1,

Y is CH,

Z is C-V,

A is $-OH$, $-NHCF_3$, $-NHSO_2R^3$, $-NHCO_2C_{1-6}$ alkyl, $-NHCON-C_{1-6}$ alkyl, or $-NHCOR^4$,

B is C_{1-3} alkyl, or halogen,

V is H, or F,

X is H or F,

W is

4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,

4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol,
 N-(4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol,
 4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)-2-methylphenol,
 N-(4-((5-(4-hydroxy-3-methylphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 2-fluoro-4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol,
 4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol,
 2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol,
 N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 ,
 4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 4-((5-(benzo[b]thiophen-3-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 4-((5-(5-chlorothiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol,
 4-((5-(5-bromopyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 N-(3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 N-(3-chloro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide,

3-ethyl-4-((5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)phenol,
 N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)butyramide,
 2,2,2-trifluoro-N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 4-(3-((4-amino-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenol,
 N-(3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 N-(2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenyl)methanesulfonamide,
 4-((5-(6-chloropyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 ethyl
 (4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 4-(3-((2-ethyl-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)-2-fluorophenol,
 1-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,
 2-fluoro-4-(3-((2-fluoro-4-hydroxyphenyl)amino)-1H-pyrazol-5-yl)phenol,
 N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide,
 4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol,
 4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol,
 N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenyl)acetamide,
 4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol,
 4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenol,
 N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenyl)acetamide,

N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-chloropropanamide,
 N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide,
 2-(dimethylamino)-N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-morpholinopropanamide,
 3-(dimethylamino)-N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)propanamide,
 N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-(trifluoromethyl)phenol,
 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-fluorophenol,
 N-(4-((5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluorophenol,
 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-chlorophenol,
 3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenol,
 N-(3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 N-(3-methyl-4-((5-phenyl-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide,
 N-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(4-cyanophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,

3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenol,
 N-(3-methyl-4-((5-(4-morpholinophenyl)-1H-pyrazol-3-yl)amino)phenyl)methanesulfonamide,
 N-(4-((5-(4-fluorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)pyrrolidin-2-one,
 1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)piperidin-2-one,
 1-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,
 methyl
 (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 ethyl
 (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 N-(3-methyl-4-((5-(4-(2-oxopyrrolidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 N-(3-methyl-4-((5-(4-(2-oxopiperidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 N-(4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(6-methoxypyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 1-(4-(3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)azetidin-2-one,
 4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 N-(4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 N-(4-((5-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(3-chloro-4-((5-(4-(2-oxopyrrolidin-1-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 4-((5-(4-(dimethylamino)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-

methylphenol,
 N-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 1-(4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,
 ethyl
 (4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 methyl
 (4-((5-(benzo[b]thiophen-2-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(3-chloro-4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,
 4-((5-(3-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 N-(4-((5-(3-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)isobutyramide,
 1-(4-(4-fluoro-3-((4-hydroxy-2-methylphenyl)amino)-1H-pyrazol-5-yl)phenyl)pyrrolidin-2-one,
 N-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)acetamide,
 methyl
 (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 ethyl
 (4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 4-((4-fluoro-5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 1-(4-((5-(4-(1H-pyrazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,
 3-ethyl-4-((4-fluoro-5-(4-iodophenyl)-1H-pyrazol-3-yl)amino)phenol,
 3-methyl-4-((5-(3-(pyridin-3-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
 ,

3-methyl-4-((5-(4-(pyridin-3-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol
 ,
 4-((5-(4-(4-fluoro-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 N-(4-((5-(4-(4-fluoro-1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide,
 4-((4-fluoro-5-(1H-indol-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 1,
 3-ethyl-4-((5-(3-iodophenyl)-1H-pyrazol-3-yl)amino)phenol,
 4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-3-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 3-methyl-4-((5-(3-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol,
 4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol,
 2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenol,
 3-methyl-4-((5-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol,
 4-((5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol,
 3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenol,
 4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,
 1-(4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,
 methyl
 (4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,
 4-((5-(4-(1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)-2-fluoro-5-methylphenol,
 1-(4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,

methyl

(4-((5-(2'-fluoro-5'-methoxy-[1,1'-biphenyl]-4-yl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,

1-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,

(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,

4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-2,5-difluorophenol,

1-methyl-3-(3-methyl-4-((5-(4-(methylsulfonyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)urea,

methyl

(3-methyl-4-((5-(4-(methylsulfonyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate,

1-methyl-3-(3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)urea,

methyl

(3-methyl-4-((5-(4-(3-(trifluoromethyl)-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate,

1-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,

methyl

(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,

methyl

(3-methyl-4-((5-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate,

3-methyl-4-((5-(4-(trifluoromethyl)phenyl)-1H-pyrazol-3-yl)amino)phenol,

methyl

(3-methyl-4-((5-(4-(trifluoromethyl)phenyl)-1H-pyrazol-3-yl)amino)phenyl)carbamate,

methyl

(4-((5-(3-fluoro-4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,

methyl

(4-((5-(4-chlorophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate, or

methyl

(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate.

[Claim 8]

The compound of Claim 1, wherein the compound is

3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenol

4-((5-(4-(dimethylamino)phenyl)-1H-pyrazol-3-yl)amino)-3-methylphenol

N-(4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide

N-(4-((5-(4-(1H-imidazol-1-yl)phenyl)-4-fluoro-1H-pyrazol-3-yl)amino)-3-methylphenyl)methanesulfonamide

4-((5-(4-(1H-pyrazol-1-yl)phenyl)-1H-pyrazol-3-yl)amino)-3-ethylphenol,

N-(3-ethyl-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)phenyl)acetamide,

2-fluoro-4-((5-(4-hydroxyphenyl)-1H-pyrazol-3-yl)amino)-5-methylphenol,

1-(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea,

(4-((5-(4-methoxyphenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate,

1-(4-((5-(4-bromophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)-3-methylurea, or

methyl

(4-((5-(4-aminophenyl)-1H-pyrazol-3-yl)amino)-3-methylphenyl)carbamate.

[Claim 9]

A composition comprising a compound of any one of Claims 1 to 8 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[Claim 10]

A composition for treating or preventing cancer, comprising as an effective agent a compound of any one of Claims 1 to 8 or a pharmaceutically acceptable salt thereof.

[Claim 11]


The composition of Claim 11, wherein the composition further comprises irinotecan or pharmaceutically acceptable salt thereof as other pharmaceutically active compound.

[Claim 12]


The composition of Claim 10 or 11, wherein the cancer is colorectal cancer, breast cancer, brain tumor, gastric cancer, liver cancer, ovarian cancer, lung cancer, gastrointestinal cancer, leukemia, or melanoma.

- [Claim 13] The composition of Claim 12, wherein the cancer is colorectal cancer.
- [Claim 14] A method for treating or preventing cancer comprising administering to a subject in need thereof a therapeutically effective amount of a compound of any one of Claims 1 to 8 or pharmaceutically acceptable salt thereof.
- [Claim 15] The method of Claim 14, wherein the method further comprises administering to a subject in need thereof a therapeutically effective amount of irinotecan or pharmaceutically acceptable salt thereof.
- [Claim 16] The method of Claim 14 or 15, wherein the cancer is colorectal cancer, breast cancer, brain tumor, gastric cancer, liver cancer, ovarian cancer, lung cancer, gastrointestinal cancer, leukemia, or melanoma.
- [Claim 17] The method of Claim 16, wherein the cancer is colorectal cancer.

[Fig. 1]



[Fig. 2]



[Fig. 3]

