



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2662084 A1 2008/03/13

(21) **2 662 084**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2007/09/04
(87) Date publication PCT/PCT Publication Date: 2008/03/13
(85) Entrée phase nationale/National Entry: 2009/02/27
(86) N° demande PCT/PCT Application No.: EP 2007/059253
(87) N° publication PCT/PCT Publication No.: 2008/028914
(30) Priorité/Priority: 2006/09/07 (EP06120305.5)

(51) Cl.Int./Int.Cl. *A61K 31/473* (2006.01),
A61P 3/10 (2006.01)

(71) Demandeur/Applicant:
NYCOMED GMBH, DE

(72) Inventeurs/Inventors:
KLEIN, THOMAS, DE;
BLASER, ANJA, DE;
RUDOLPH, BETTINA, DE;
KAUTZ, ULRICH, DE;
SELIGE, JENS, DE;
KROMER, WOLFGANG, DE

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : POLYTHERAPIE CONTRE LE DIABETE SUCRE
(54) Title: COMBINATION TREATMENT FOR DIABETES MELLITUS

(57) **Abrégé/Abstract:**

The invention relates to combinations of (2R,4aR,10bR)-6-(2,6-Dimethoxy-pyridin-3-yl)-9-ethoxy-8-methoxy-1,2,3,4,4a,10b-hexahydrophenanthridin-2-ol with other active compounds for the treatment of diabetes mellitus type 2 and/or type 1.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 March 2008 (13.03.2008)

PCT

(10) International Publication Number
WO 2008/028914 A1

(51) International Patent Classification:

A61K 31/473 (2006.01) A61P 3/10 (2006.01)

(21) International Application Number:

PCT/EP2007/059253

(22) International Filing Date:

4 September 2007 (04.09.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

06120305.5 7 September 2006 (07.09.2006) EP

(71) Applicant (for all designated States except US): **NY-COMED GMBH** [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KLEIN, Thomas** [DE/DE]; Döbelestr. 16, 78462 Konstanz (DE). **BLASER, Anja** [DE/DE]; Fliesshornstr. 10, 78465 Konstanz (DE). **RUDOLPH, Bettina** [DE/DE]; Brauneggerstr. 19, 78462 Konstanz (DE). **KAUTZ, Ulrich** [DE/DE]; Prof.-Schmider-Str. 12, 78476 Allensbach (DE). **SELIGE, Jens** [DE/DE]; Gartenstr. 28, 78462, Konstanz. **KROMER, Wolfgang** [DE/DE]; Hinterhauserstr. 5, 78464 Konstanz (DE).

(74) Agents: **WILD, Robert** et al.; Nycomed GmbH, P.O. Box 100310, 78403 Konstanz (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(54) Title: COMBINATION TREATMENT FOR DIABETES MELLITUS

(57) Abstract: The invention relates to combinations of (2R,4aR,10bR)-6-(2,6-Dimethoxy-pyridin-3-yl)-9-ethoxy-8-methoxy-1,2,3,4,4a,10b-hexahydrophenanthridin-2-ol with other active compounds for the treatment of diabetes mellitus type 2 and/or type 1.

WO 2008/028914 A1

Combination Treatment for Diabetes Mellitus

Technical field

5

The invention relates to combinations of a known PDE4 inhibitor with one or two other active compounds which are used in the treatment of diabetes mellitus type 2 and/or diabetes mellitus type 1; the invention also relates to pharmaceutical compositions, combination products and kits containing these combinations as well as uses of such combinations in the treatment of diabetes mellitus type 2 and/or

10

diabetes mellitus type 1.

Background of the invention

15

Diabetes mellitus is on the rise worldwide and is considered to be at an epidemic level by the World Health Organization. The clinical manifestation and progression of diabetes often vary considerably between countries and commonly between ethnic groups in the same country. Currently diabetes affects 151 million people worldwide and an estimate 300 million people in 2025. There are two main forms of diabetes. Type 1 (insulin-dependent diabetes mellitus, IDDM) is due primarily to autoimmune-mediated destruction of pancreatic β -cells, resulting in absolute insulin deficiency. It is the second most common chronic disease of children. By contrast, type 2 diabetes (non-insulin-dependent diabetes mellitus, NIDDM) is characterized by insulin-resistance and inadequate insulin secretion. A significant fraction of individuals originally diagnosed with type 2 diabetes evolve with time to a type 1 state, which is defined as exhibiting anti- β -cell autoimmunity.

20

25

Because genetic factors contribute to the development of diabetes, the disease displays a strong familial aggregation. Although there are monogenic syndromes of insulin resistance, in which a definite gene has been identified as the cause of insulin resistance, these are relative rare. The more common presentation of diabetes appears to be polygenic. Additionally, behavioural- and lifestyle-related risk factors exist. Type 2 diabetes is increasingly common primarily because of increases in the prevalence of a sedentary lifestyle and obesity. One of the major arguments for the role of behavioural factors in the etiology of diabetes has been the rapid increase in the prevalence and incidence of the disease in populations undergoing rapid westernization. The westernization transition is usually accompanied by increases in obesity, decreases in physical activity and alterations in dietary intake toward more calories, fat and non-complex carbohydrates.

30

35

Plasma glucose concentrations are normally maintained within a fairly narrow range despite wide fluctuations in the body's supply (e.g. meals) and demand (e.g. exercise) for nutrients. After an overnight fast, insulin-independent tissues, the brain (50%) and splanchnic organs (25%), account for most of the total body glucose disposal. Insulin-dependent tissues, adipose tissue and primarily skeletal mus-

40

cles, are responsible for the remaining 25% of glucose utilization. This basal glucose uptake is precisely matched by the release of glucose from the liver. In response to hyperglycemia after a meal, pancreatic insulin secretion is stimulated and the combination of hyperinsulinemia plus hyperglycemia promotes glucose uptake (by splanchnic and peripheral, primarily muscle, tissues) and suppresses hepatic glucose production. It follows, therefore, that defects at the level of the β -cell, muscle and liver can lead to the development of glucose intolerance and diabetes mellitus. All the abnormalities in diabetes basically result from an imbalance between insulin sensitivity and insulin secretion. The initial stage of diabetes is characterised by impaired glucose tolerance and postprandial hyperglycemia. As the disease progresses, fasting hyperglycemia is observed.

The earliest detectable abnormality in NIDDM is an impairment in the body's ability to respond to insulin. Because the pancreas is able to appropriately augment its secretion of insulin to offset the insulin resistance, glucose tolerance remains normal. With time, however, the beta-cell fails to maintain its high rate of insulin secretion and the insulin resistance leads to the development of impaired glucose tolerance and eventually overt diabetes mellitus. The cause of pancreatic "exhaustion" remains unknown, however lipo- and glucotoxicity with an increased level of oxygen radical stress are discussed more recently. Insulin resistance in NIDDM involves both hepatic and peripheral tissues. In response to both endogenously secreted or exogenously administered insulin, hepatic glucose production fails to suppress normally and muscle glucose uptake is diminished. The accelerated rate of hepatic glucose output is due mainly to augmented gluconeogenesis. In muscle many cellular defects in insulin action have been described including impaired insulin-receptor tyrosine kinase activity, diminished glucose transport, and reduced glycogen synthase and pyruvate dehydrogenase activities. The abnormalities account for disturbances in the two major intracellular pathways of glucose disposal, glycogen synthesis and glucose oxidation. In the earliest stages of NIDDM, the major defect involves the inability of insulin to promote glucose uptake and storage as glycogen. Other potential mechanisms that have been put forward to explain the glucose intolerance include increased levels of free fatty acids, chronic low-grade activation of the immune system (increased levels of $\text{TNF}\alpha$ and IL6), altered skeletal muscle blood flow, increased conversion of amylin to its insoluble amyloid form and glucose toxicity.

Diabetes is associated with a variety of physiologic disorders such as hypertension and dyslipidemia. Diabetes also increases the risk of macrovascular (coronary artery disease, stroke, amputation) and microvascular (blindness, renal failure, neuropathies) diseases. Myocardial infarction, stroke or renal failure are the cause of death for more than 70% of diabetes patients. The huge mortality and debilitating neuropathies associated with diabetes underline the importance of active medical intervention.

There are several ways to counteract diabetes. The first is lifestyle adjustments aimed at improving endogenous insulin sensitivity. This can be achieved by increased physical activity and bodyweight reduction with diet and behavioural modification. Unfortunately, most people with non-insulin-

dependent diabetes mellitus never receive sufficient nutritional education or are not capable of complying with a strict diet regimen.

Another therapeutic way involves increasing insulin availability by the administration of exogenous insulin, insulin analogues and insulin secretagogues such as sulphonylureas. The primary mode of action of sulphonylureas is through the depolarisation of the pancreatic β -cells by blocking the ATP-dependent potassium channels and causing an influx of calcium ions, which stimulate insulin secretion.

Biguanides, of which metformin is the most commonly used, also have proven to be effective anti-hyperglycemic agents. Metformin reduces hepatic gluconeogenesis and basal hepatic glucose output. Oral antidiabetics such as sulphonylureas and metformin as monotherapy or in combination have been shown to decrease fasting plasma glucose levels, but postprandial hyperglycemia persists in more than 60% of patients and probably accounts for sustained increases of hemoglobin A_{1C} levels.

α -Glucosidase inhibitors, e.g. acarbose and miglitol, primarily target postprandial hyperglycemia. The therapy of diabetes mellitus with α -glucosidase inhibitors is based on a delayed intestinal degradation of starch and sucrose. These carbohydrates must be hydrolysed by α -glucosidases to monosaccharides before they can be transported through the mucosa of the small intestine. The reversible inhibition of the brush border glucosidases results in redistribution of carbohydrate absorption from the upper portion of the gut to a more extended surface area covering the whole length of the small intestine. This is accompanied by a delayed absorption of monosaccharides and a decrease in the postprandial elevation of blood glucose.

Another class of antidiabetic drugs are thiazolidinediones, such as rosiglitazone and pioglitazone, which are insulin sensitizers and act through activation of peroxisome proliferator-activated receptor γ (PPAR γ). PPAR γ is mainly expressed in adipose tissues, plays an important role in adipogenesis and modifies fatty acid synthesis and storage. Binding of rosiglitazone to PPAR γ results in reduced endogenous glucose production and increased blood glucose uptake. It increases the sensitivity of skeletal muscle, liver and adipose tissues to insulin. Improvements in glucose metabolism with rosiglitazone treatment are closely correlated with decreased plasma free fatty acid metabolism. The stimulation by rosiglitazone of PPAR γ in adipose tissue and subsequent adipocyte differentiation results in the generation of more, but smaller, adipocytes which are more insulin sensitive and produce less free fatty acid, TNF α and leptin.

In the International Patent Application WO9914239 compositions for treating diabetes mellitus and obesity are disclosed. The compositions contain at least two of the active agents A, B and C, wherein A is at least one hormone which stimulates the production of cAMP, B is at least one substance which inhibits the breakdown of a cyclic nucleotide, and C is at least one hormone which stimulates the pro-

duction of cGMP. In the International Patent Application WO0135979 the combined use of a PDE3 and a PDE4 inhibitor for the treatment of obesity is disclosed. In the International Patent Application WO0213798 the use of a selective cGMP PDE5 inhibitor for the treatment of Insulin Resistance Syndrome is disclosed, wherein the Insulin Resistance Syndrome is defined as the concomitant existence of two or more disease states selected from dyslipidemia, hypertension, type 2 diabetes mellitus, impaired glucose tolerance, a family history of diabetes, hyperuricaemia and/or gout, a pro-coagulant state, atherosclerosis and truncal obesity. In the unexamined German application DE 10150517 tetrahydropyridazin-3-one derivatives are described which may be useful inter alia for the treatment of diabetes mellitus. In the International Patent Application WO2005023253 several PDE4 inhibitors are disclosed to be useful in the treatment of diabetes mellitus. In Diabetes 47, pp. 570-575, 1998 is disclosed that pentoxifylline and rolipram may be effective in the treatment of autoimmune diabetes or other conditions characterized by excessive production of inflammatory cytokines.

It is an object of the present invention to provide combinations and methods of treatment that take advantage of the different therapeutic pathways of a PDE4 inhibitor on the one hand side and other known anti-diabetic compounds on the other hand side to more effectively treat diabetes mellitus type 2 and/or type 1.

Description of the invention

The invention is based on the expectation that the use of the selective PDE4 inhibitor (2R,4aR,10bR)-6-(2,6-Dimethoxy-pyridin-3-yl)-9-ethoxy-8-methoxy-1,2,3,4,4a,10b-hexahydrophenanthridin-2-ol (hereinafter referred to as "Compound A") or a pharmaceutical acceptable salt thereof in combination with one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 leads to beneficial effects in the treatment of diabetes mellitus type 2 and/or type 1 in comparison to the treatment with either the selective PDE4 inhibitor (2R,4aR,10bR)-6-(2,6-Dimethoxy-pyridin-3-yl)-9-ethoxy-8-methoxy-1,2,3,4,4a,10b-hexahydrophenanthridin-2-ol or the above-mentioned active compound(s) alone.

Therefore, according to a first aspect of the present invention there is provided a pharmaceutical composition comprising a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

As an embodiment of the first aspect of the present invention there is provided a pharmaceutical composition comprising a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1,

wherein the first amount and the second amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

According to a second aspect of the present invention there is provided a pharmaceutical composition comprising a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

As an embodiment of the second aspect of the present invention there is provided a pharmaceutical composition comprising a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, amounts of two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first, second and third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

The above-mentioned pharmaceutical compositions provide for the administration of Compound A or a pharmaceutically acceptable salt thereof with one or two other active compound(s) or pharmaceutically acceptable salts thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 and is thus presented as a single formulation.

Alternatively, Compound A or a pharmaceutically acceptable salt thereof and the one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 may be presented as separate formulations, wherein at least one of those formulations comprises Compound A or a pharmaceutically acceptable salt thereof and at least one comprises one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1.

Thus, there is further provided:

A combination product comprising the components: (A) Compound A or a pharmaceutically acceptable salt thereof; and (B) one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein each of the components (A) and (B) is formulated in admixture with at least one pharmaceutically acceptable auxiliary.

As an embodiment of the above-mentioned combination product there is provided a combination product comprising the components: (A) an amount of Compound A or a pharmaceutically acceptable salt thereof; and (B) an amount of one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first and the second amount together comprise an effective amount for the treatment of diabetes mellitus type

2 and/or type 1 and wherein each of the components (A) and (B) is formulated in admixture with at least one pharmaceutically acceptable auxiliary.

A combination product comprising the components: (A) Compound A or a pharmaceutically acceptable salt thereof; (B) one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1; and (C) still another active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein each of the components (A), (B) and (C) is formulated in admixture with at least one pharmaceutically acceptable auxiliary.

As an embodiment of the above-mentioned combination product there is provided a combination product comprising the components: (A) an amount of Compound A or a pharmaceutically acceptable salt thereof; (B) an amount of one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1; and (C) an amount of still another active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first, second and third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1 and wherein each of the components (A), (B) and (C) is formulated in admixture with at least one pharmaceutically acceptable auxiliary.

A kit comprising the components: (A) a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, in admixture with at least one pharmaceutically acceptable auxiliary; and (B) a pharmaceutical formulation including one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary.

As an embodiment of the above-mentioned kit there is provided a kit comprising the components: (A) a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, in admixture with at least one pharmaceutically acceptable auxiliary; and (B) a pharmaceutical formulation including an amount of one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary, wherein the first and the second amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1.

A kit comprising the components: (A) a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, in admixture with at least one pharmaceutically acceptable auxiliary; (B) a pharmaceutical formulation including an amount of one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary; and (C) a pharmaceutical formulation including still another active compound or a pharmaceutically acceptable salt thereof which

is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary.

As an embodiment to the above-mentioned kit there is provided a kit comprising the components: (A) a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, in admixture with at least one pharmaceutically acceptable auxiliary; (B) a pharmaceutical formulation including an amount of one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary; and (C) a pharmaceutical formulation including an amount of still another active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary, wherein the first, second and third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1.

In addition, to the components A and B (or A, B and C) the kits according to the invention may include information for the sequential or separate administration of the components to the patient in need of diabetes mellitus type 2 and/or type 1 therapy.

The combinations according to the present invention are used for the treatment of diabetes mellitus type 2 and/or type 1, preferably for the treatment of diabetes mellitus type 2.

Therefore, further aspects of the invention are:

The use of Compound A or a pharmaceutically acceptable salt thereof and one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1 for the manufacture of a medicament, in particular a pharmaceutical composition according to the invention, for the treatment of diabetes mellitus type 2 and/or type 1.

The use of Compound A or a pharmaceutically acceptable salt thereof and two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1 for the manufacture of a medicament, in particular a pharmaceutical composition according to the invention, for the treatment of diabetes mellitus type 2 and/or type 1.

The use of Compound A or a pharmaceutically acceptable salt thereof and one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1 for the manufacture of a sequentially or separately co-administrable medicament, in particular the combination product or kit according to the invention, for the treatment of diabetes mellitus type 2 and/or type 1.

The use of Compound A or a pharmaceutically acceptable salt thereof and two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1 for the manufacture of a sequentially or separately co-administrable medication, in particular a combination product or kit according to the invention, for the treatment of diabetes mellitus type 2 and/or type 1.

Compound A or a pharmaceutically acceptable salt thereof and one other active compound or pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1 for the treatment of diabetes mellitus type 2 and/or type 1.

Compound A or a pharmaceutically acceptable salt thereof and two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1 for the treatment of diabetes mellitus type 2 and/or type 1.

A pharmaceutical composition, combination product or kit according to the invention for the treatment of diabetes mellitus type 2 and/or type 1.

Another aspect of the present invention is a method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a pharmaceutical composition comprising a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first amount and the second amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

Still another aspect of the present invention is a method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a pharmaceutical composition comprising a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, amounts of two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first, second and third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

A further aspect of the present invention is a method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a combination product comprising the components:

(A) an amount of Compound A or a pharmaceutically acceptable salt thereof; and

(B) an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1,

wherein the first and second amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1;

wherein each of the components (A) and (B) is formulated in admixture with at least one pharmaceutically acceptable auxiliary;

5 and wherein the components (A) and (B) are administered sequentially or separately.

Still a further aspect of the present invention is a method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a combination product comprising the components:

10 (A) an amount of Compound A or a pharmaceutically acceptable salt thereof;

(B) an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1; and

(C) an amount of still another active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1,

15 wherein the first, second and third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1;

wherein each of the components (A), (B) and (C) is formulated in admixture with at least one pharmaceutically acceptable auxiliary;

and wherein the components (A), (B) and (C) are administered sequentially or separately.

20

The pharmaceutical compositions according to the invention may be prepared by mixing the first active agent with the second (and optionally third) active agent.

In the above-mentioned mixing process the first active agent and the second (and optionally third)

25 active agent can

a) in a first step be mixed as such, afterwards be processed with at least one pharmaceutically acceptable auxiliary and finally, for example, be pressed to tablets or caplets

or

b) in a first step separately be processed with at least one pharmaceutically acceptable auxiliary to

30 give granules or pellets containing each only one of the two (or three) active agents; the pellets or granules for their part then can be mixed in an appropriate ratio and either pressed – optionally with further pharmaceutically acceptable auxiliaries - to give, for example tablets or caplets, or can be filled in loose form in capsules.

35 Therefore, in a still further aspect of the present invention there is provided a process for the preparation of a pharmaceutical composition which comprises mixing a first active agent, which is Compound A or a pharmaceutically acceptable salt thereof with a second (and optionally a third) active agent, which is (are) one or two other active compounds which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1.

40

Simultaneous administration of Compound A or a pharmaceutically acceptable salt thereof and one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 can be preferably accomplished, by administering to the patient in need of diabetes mellitus type 2 and/or type 1 therapy the pharmaceutical

5 composition according to the invention in one dosage form, such as for example, in a single capsule, tablet or injection.

Components (A), (B) and optionally (C) of the combination product as well as of the kit may be administered sequentially or separately over the course of the treatment of diabetes mellitus type 2 and/or

10 type 1.

Sequential or separate administration of Compound A or a pharmaceutically acceptable salt thereof and one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 can be preferably accomplished, by

15 administering to the patient in need of diabetes mellitus type 2 and/or type 1 therapy components (A), (B) and optionally (C) of the combination product or the kit according to the invention in (multiple) separate dosage forms, such as for example, in separate capsules, tablets or injections.

In an alternative, one (or two) of the components (A), (B) and optionally (C) may be formulated as

20 tablet or capsule and the remaining component(s) may be formulated for administration, for example, by injection or inhalation.

Sequential administration encompasses a short time period between the administration of components (A), (B) and optionally (C) of the combination product or the kit according to the invention (for example, the time that is needed to swallow one tablet after the other).

25

Separate administration encompasses both short and long time periods between the administration of components (A), (B) and optionally (C) of the combination product or the kit according to the invention. However, for the purposes of the present invention at least one of the components is administered while the other component(s) is (are) still having an effect on the patient being treated. In a preferred embodiment of the invention the effect on the patient being treated is a synergistic effect.

30

The combined administration of Compound A or a pharmaceutically acceptable salt thereof and one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1, either in form of the pharmaceutical composition, combination product or kit according to the invention, lead to an effective treatment of diabetes type 2 and/or type 1, and in a preferred embodiment is superior to the use of either active agent alone. Moreover, in a particularly preferred embodiment, the combined administration of Compound A or a pharmaceutically acceptable thereof and one or two other active compound(s) or pharmaceutically

35

acceptable salt(s) thereof which is (are) used in the treatment of diabetes type 2 and/or type 1 shows a synergistic efficacy for treating diabetes mellitus type 2 and/or type 1.

As used herein, the term "synergistic" refers to the combination of Compound A or a pharmaceutically acceptable salt thereof with one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 either in form of the pharmaceutical composition, combination product or kit according to the invention having an efficacy for the treatment of diabetes mellitus type 2 and/or type 1 that is greater than would be expected from the sum of their individuals effects. The synergistic effects of the embodiments of the present invention encompass additional unexpected advantages for the treatment of diabetes mellitus type 2 and/or type 1. Such additional advantages may include, but are not limited to, lowering the required dose of one or more of the active agents of the combination, reducing the side effects of one or more of the active agents of the combination or rendering one or more of the active agents more tolerable to the patient in need of a diabetes type 2 and/or type 1 therapy. The combined administration of Compound A or a pharmaceutically acceptable salt thereof and one or two other active compound(s) or pharmaceutically acceptable salts thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 may also be useful for decreasing the required number of separate dosages, thus, potentially improving compliance of the patient in need of diabetes mellitus type 2 and/or type 1 therapy.

The term "active agent" as used herein refers either to Compound A or a pharmaceutically acceptable salt thereof or to the active compound(s) or pharmaceutically acceptable salt(s) thereof, which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1.

The term "active compound" as used herein refers to a compound useful in the treatment of a disease, here in particular in the treatment of diabetes mellitus type 2 and/or type 1.

The term "effective amount" as used herein refers to a therapeutically effective amount for treating diabetes mellitus type 2 and/or type 1. In case of a combination therapy the term "effective amount" refers to the sum of the amounts of the combination partners, which is therapeutically effective for the treatment of diabetes mellitus type 2 and/or type 1.

In case of the pharmaceutical compositions, combination products and kits according to the invention the term "amount" is mentioned in connection with the pharmaceutical formulations which form part of the pharmaceutical compositions, or in connection with the components which form part of the combination products or kits. The term "effective amount" is mentioned in phrases, such as "wherein the first and second amount (or the first, second and third amount) together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1". In case, a once daily administration of the pharmaceutical formulation of the pharmaceutical composition, or a once daily administration of the components of the combination product is intended, the respective amount(s) correspond to the daily

dosage necessary for the treatment of diabetes mellitus type 2 and/or type 1. In case, a twice a day administration of the pharmaceutical formulation of the pharmaceutical composition, or a twice a day administration of the components (or one of the components) of the combination product or kit is intended, the respective amount(s) correspond to the daily dosage necessary for the treatment of diabetes mellitus type 2 and/or type 1 multiplied by the factor 0.5. In case, a three times a day administration of the pharmaceutical formulation of the pharmaceutical composition, or a three times a day administration of the components (or one of the components) of the combination product or kit is intended, the respective amount(s) correspond to the daily dosage necessary for the treatment of diabetes mellitus type 2 and/or type 1 multiplied by the factor 0.33 and so on.

The term "patient" includes both humans and other mammals. In a preferred embodiment of the invention the term "patient" stands for humans.

The term "one other active compound" in connection with the term "still another active compound" means that the two mentioned active compounds are not identical. As well, the term "two other active compounds" stands for two active compounds that are not identical.

The term "Compound A" as used herein stands for (2R,4aR,10bR)-6-(2,6-Dimethoxy-pyridin-3-yl)-9-ethoxy-8-methoxy-1,2,3,4,4a,10b-hexahydrophenanthridin-2-ol.

In one embodiment of the present invention a preferred pharmaceutically acceptable salt of Compound A is selected from the tosylate, esylate, hydrobromide and hydrochloride salt of Compound A.

In another embodiment of the present invention a preferred pharmaceutically acceptable salt of Compound A is the hydrochloride salt of Compound A.

Additional information with regard to the preparation and suitable dosage forms of Compound A and the pharmaceutically acceptable salts thereof can be found in the following patents/patent applications: WO2005/085225 and PCT/EP2006/060377 (WO2006092422).

Non-limiting examples of other active compounds which are used in the treatment of diabetes mellitus type 2 and/or type 1 are provided in the following list:

- Insulin and insulin analogues
- Glucagon-Like-Peptide-1 (GLP-1) receptor agonists
- Sulfonylurea agents
- Biguanide agents
- Alpha-glucosidase inhibitors
- PPAR-Agonists
- Meglitinide agents

- 13 -

- Dipeptidyl-peptidase (DPP) IV inhibitors
 - PDE1, PDE5, PDE9, PDE10 or PDE11 inhibitors
 - Amylin agonists
 - Cinnamon
 - 5 - Glucagon receptor antagonists
 - Glycogen-Phosphorylase inhibitors
 - Fructose-1,6-Bisphosphate inhibitors
 - Cannabinoid (CB1) receptor antagonists
 - Anti-obesity drugs such as appetite suppressors, satiety increasing substances, and
 - 10 energy expenditure increasing drugs
- and pharmaceutically acceptable salts thereof.

In one embodiment of the invention, the one or two other active compound(s) which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 is (are) selected from the group consisting of

- 15 - Insulin and insulin analogues
 - Glucagon-Like-Peptide-1 (GLP-1) receptor agonists
 - Sulfonylurea agents
 - Biguanide agents
 - Alpha-glucosidase inhibitors
 - 20 - PPAR-Agonists
 - Meglitinide agents
 - Dipeptidyl-peptidase (DPP) IV inhibitors
 - PDE1, PDE5, PDE9, PDE10 or PDE11 inhibitors
 - Amylin agonists
 - 25 - Cinnamon
 - Glucagon receptor antagonists
 - Glycogen-Phosphorylase inhibitors
 - Fructose-1,6-Bisphosphate inhibitors
 - Cannabinoid (CB1) receptor antagonists
 - 30 - Anti-obesity drugs such as appetite suppressors, satiety increasing substances, and
 - energy expenditure increasing drugs
- and pharmaceutically acceptable salts thereof.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin. Specific examples of insulin include, but are not limited to Humulin® [human insulin, (rDNA origin)], Novolin® [human insulin, (rDNA origin)], Velosulin® BR [human buffered regular insulin, (rDNA origin)] and Exubera® [human insulin, inhaled].

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an insulin analogue or a pharmaceutically acceptable

salt thereof. Specific examples of insulin analogues include, but are not limited to, novarapid, insulin detemir, insulin lispro, insulin glargine, insulin zinc suspension and Lys-Pro insulin.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a Glucagon-Like-Peptide-1 receptor agonist or a pharmaceutically acceptable salt thereof. Specific examples of Glucagon-Like-Peptide-1 receptor agonists include, but are not limited to BIM-51077 (CAS-No. 275371-94-3), EXENATIDE (CAS-No. 141758-74-9), LIRAGLUTIDE (CAS-No. 20656-20-2), ALBIGLUTIDE (CAS-No. 782500-75-8) and ZP-10 (CAS-No. 320367-13-3). A preferred Glucagon-Like-Peptide-1 receptor agonist is EXENATIDE.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof. Specific examples of sulfonylurea agents include, but are not limited to, TOLBUTAMIDE (CAS-No. 000064-77-7), TOLAZAMIDE (CAS-No. 001156-19-0), GLIPIZIDE (CAS-No. 029094-61-9), CARBUTAMIDE (CAS-No. 000339-43-5), GLISOXEPIDE (CAS-No. 025046-79-1), GLISENTIDE (CAS-No. 032797-92-5), GLIBORNURIDE (CAS-No. 026944-48-9), GLIBENCLAMIDE (CAS-NO. 010238-21-8), GLIQUIDONE (CAS-No. 033342-05-1), GLIMEPIRIDE (CAS-No. 093479-97-1) and GLICLAZIDE (CAS-No. 021187-98-4).

In another embodiment of the present invention the pharmaceutically acceptable salt of TOLBUTAMIDE is the sodium salt of TOLBUTAMIDE. In another embodiment of the present invention the pharmaceutically acceptable salt of GLIQUIDONE is the sodium salt of GLIQUIDONE.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof. A specific example of a biguanide agent includes, but is not limited to METFORMIN (CAS-No. 000657-24-9).

In another embodiment of the present invention the pharmaceutically acceptable salt of METFORMIN is the hydrochloride salt of METFORMIN.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an alpha-glucosidase-inhibitor or a pharmaceutically acceptable salt thereof. Specific examples of alpha-glucosidase-inhibitors include, but are not limited to ACARBOSE (Cas-No. 056180-94-0), MIGLITOL (CAS-No. 072432-03-2) and VOGLIBOSE (CAS-No. 083480-29-9).

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR-agonist or a pharmaceutically acceptable salt thereof. Specific examples of PPAR-agonists include, but are not limited to MURAGLITAZAR

(CAS-No. 331741-94-7), ROSIGLITAZONE (CAS-NO. 122320-73-4), PIOGLITAZONE (CAS-No. 111025-46-8), FARGLITAZAR (CAS-No. 196808-45-4), NAVEGLITAZAR (CAS-No. 476436-68-7), NETOGLITAZONE (CAS-NO. 161600-01-7), RIVOGLITAZONE (CAS-No. 185428-18-6), K-111 (CAS-No. 221564-97-2), SODELGLITAZAR (= GW-677954; CAS-No. 622402-24-8) and (-)-Halofenate
5 (CAS-No. 024136-23-0). Preferred PPAR-agonists are ROSGLITAZONE and PIOGLITAZONE.

In another embodiment of the present invention the pharmaceutically acceptable salt of ROSIGLITAZONE is the maleate salt of ROSIGLITAZONE. In another embodiment of the present invention the pharmaceutically acceptable salt of RIVOGLITAZONE is the hydrochloride salt of RIVOGLITAZONE.
10 In another embodiment of the present invention the pharmaceutically acceptable salt of K-111 is the sodium salt of K-111. In another embodiment of the present invention the pharmaceutically acceptable salt of PIOGLITAZONE is the dihydrochloride salt of PIOGLITAZONE.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a meglitinide agent or a pharmaceutically acceptable salt thereof. Specific examples of meglitinide agents include, but are not limited to REPAGLINIDE
15 (CAS-No. 135062-02-1), NATEGLINIDE (CAS-No. 105816-04-4) and MITIGLINIDE (CAS-No. 145375-43-5).

20 In another embodiment of the present invention the pharmaceutically acceptable salts of MITIGLINIDE are the monopotassium or the calcium salt of MITIGLINIDE.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a DPP-IV inhibitor or a pharmaceutically acceptable salt thereof. Specific examples of DPP IV inhibitors include, but are not limited to SITAGLIPTIN (CAS-
25 No. 486460-32-6), SAXAGLIPTIN (CAS-No. 361442-04-8), VILDAGLIPTIN (CAS-No. 274901-16-5), DENAGLIPTIN (CAS-No. 483369-58-0), ALOGLIPTIN (CAS-No. 850649-61-5) and P32/98 (CAS-No. 251572-70-0).

30 In another embodiment of the present invention the pharmaceutically acceptable salt of SITAGLIPTIN is the phosphate salt of SITAGLIPTIN. In another embodiment of the present invention the pharmaceutically acceptable salt of ALOGLIPTIN is the benzoate salt of ALOGLIPTIN. In another embodiment of the present invention the pharmaceutically acceptable salts of P32/98 are the fumarate or hydrochloride salt of P32/98.

35 In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PDE5 inhibitor or a pharmaceutically acceptable salt thereof. Specific examples of PDE5 inhibitors include, but are not limited to SILDENAFIL (CAS-No. 139755-83-2), VARDENAFIL (CAS-No. 224785-90-4), TADALAFIL (CAS-No. 171596-29-5), UDE-
40 NAFIL (CAS-No. 268203-93-6) and AVANAFIL (CAS-No. 330784-47-9).

In another embodiment of the present invention the pharmaceutically acceptable salts of SILDENAFIL are the hemi-citrate, the citrate or the mesilate salt of SILDENAFIL; particularly preferred is the citrate salt of SILDENAFIL. In another embodiment of the present invention the pharmaceutically acceptable salts of VARDENAFIL are the mono-hydrochloride salt of VARDENAFIL or the dihydrochloride salt of VARDENAFIL. In another embodiment of the present invention the pharmaceutically acceptable salt of AVANAFIL is the besilate salt of AVANAFIL.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PDE1, PDE9, PDE10 or PDE11 inhibitor or a pharmaceutically acceptable salt thereof. PDE1, PDE9, PDE10 or PDE11 inhibitors which may be useful employed according to the present invention, can be found, for example, in US20020160939, WO03037432, US2004220186, WO2005003129, WO2005012485, WO2005120514 and WO03077949.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an amylin agonist or a pharmaceutically acceptable salt thereof. A specific example of an amylin agonist includes, but is not limited to PRAMLINITIDE (CAS-No. 151126-32-8).

In another embodiment of the present invention the pharmaceutically acceptable salt of PRAMLINITIDE is the acetate salt of PRAMLINITIDE.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is cinnamon.

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glucagon receptor antagonist or a pharmaceutically acceptable salt thereof. A specific example of a glucagons receptor antagonist includes, but is not limited to BAY-27-9955 (CAS-No. 202855-56-9).

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glycogen-phosphorylase inhibitor or a pharmaceutically acceptable salt thereof. An example of a glycogen-phosphorylase inhibitor includes, but is not limited to INGLIFORIB (CAS-No. 186392-65-4).

In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a fructose-1,6-bisphosphate inhibitor or a pharmaceutically acceptable salt thereof. An example of a fructose-1,6-bisphosphate inhibitor includes, but is not

limited to MANAGLINAT DIALANETIL (= MB-06322; CAS-No. 280782-97-0) and MB-05032 (Cas-No. 261365-11-1).

5 In another embodiment of the present invention the pharmaceutically acceptable salt of MB-05032 is the hydrobromide salt of MB-05032.

10 In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a cannabinoid (CB1) receptor antagonist or a pharmaceutically acceptable salt thereof. Specific examples of cannabinoid (CB1) receptor antagonists include, but are not limited to AVE-1625 (CAS-No. 261922-46-7), RIMONABANT (CAS-No. 168273-06-1) and SURINABANT (CAS-No. 288104-79-0).


15 In another embodiment of the present invention the pharmaceutically acceptable salt of RIMONABANT is the hydrochloride salt of RIMONABANT.








In another embodiment of the present invention the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an anti-obesity drug or a pharmaceutically acceptable salt thereof. Specific examples of anti-obesity drugs include, but are not limited to HMR-1426 (CAS-No. 262376-75-0), CETILISTAT (CAS-No. 282526-98-1) and SIBUTRAMINE (CAS-No. 106650-56-0).








20 In another embodiment of the present invention the pharmaceutically acceptable salt of HMR-1426 is the hydrochloride salt of HMR-1426. In another embodiment of the present invention the pharmaceutically acceptable salt of SIBUTRAMINE is the hydrochloride salt of SIBUTRAMINE.







25 More details with respect to preferred combination partners for Compound A are listed in Table 1:








Table 1:






INN or Research Code	Structure/Chemical Name
BIM-51077	L-histidyl-2-methylalanyl-L-glutamyl-glycyl-L-threonyl-L-phenylalanyl-L-threonyl-L-seryl-L-aspartyl-L-valyl-L-seryl-L-seryl-L-tyrosyl-L-leucyl-L-glutamyl-glycyl-L-glutamyl-L-alanyl-L-alanyl-L-lysyl-L-glutamyl-L-phenylalanyl-L-isoleucyl-L-alanyl-L-tryptophyl-L-leucyl-L-valyl-L-lysyl-2-methylalanyl-L-argininamide
EXENATIDE	L-histidylglycyl-L-glutamylglycyl-L-threonyl-L-phenylalanyl-L-threonyl-L-seryl-L-aspartyl-L-leucyl-L-seryl-L-lysyl-glutamyl-L-methionyl-L-glutamyl-L-glutamyl-L-glutamyl-L-alanyl-L-valyl-L-arginyl-L-leucyl-L-phenylalanyl-L-isoleucyl-L-glutamyl-L-tryptophyl-L-leucyl-L-lysyl-L-asparaginylglycylglycyl-L-prolyl-L-seryl-L-serylglycyl-L-alanyl-L-prolyl-L-prolyl-L-prolyl-L-serinamide
LIRAGLUTIDE	L-histidyl-L-alanyl-L-glutamyl-glycyl-L-threonyl-L-phenylalanyl-L-threonyl-L-seryl-L-aspartyl-L-valyl-L-seryl-L-seryl-L-tyrosyl-L-leucyl-L-glutamyl-glycyl-L-glutamyl-L-alanyl-L-alanyl-Nepsilon-(Nalpha-hexadecanoyl-gamma-L-glutamyl)-L-lysyl-L-glutamyl-L-phenylalanyl-L-isoleucyl-L-alanyl-L-tryptophyl-L-leucyl-L-valyl-L-arginyl-glycyl-L-arginyl-glycine
ALBIGLUTIDE	HGEGTFTSDV SSYLEGQAAK EFIAWLVKGR HGEGTFTSDV SSYLEQAAK EFIAWLVKGR DAHKSEVAHR FKDLGEENFK ALVLIAFAQYLQQCPFEDHV KLVNEVTEFA KTCVADESAE NCDKSLHTLF GDKLCTVATLRETYGEMADC CAKQEPERNE CFLQHKDDNP NLPRLVRPEV DVMCTAFHDN EETFLKKLY EIARRHPYFY APELLFFAKR YKAAFTECCQ AADKAACLLP KLDELDEGK ASSAKQRLKC ASLQKFGERA FKAWAVARLS QRFPKAEFAEVSKLVTDLTk VHTECCHGDL LECADDRADL AKYICENQDS ISSKLKECCE KPLLEKSCI AEVENDEMPA DLPSLAADFV ESKDVCKNYA EAKDVFLGMF LYEARRHPD YSVVLLRLA KTYETTLEKC CAAADPHECY AKVFDEFKPL VEEPQLIKQ NCELFEQLGE YKFQNALLVR YTKKVPQVST PTLVEVSRNLGKVGSKCCKH PEAKRMPCAE DYLSVVLNQL CVLHEKTPVS DRVTKCCTESLVNRRPCFSA LEVDETYVPK EFNAETFTFH ADICTLSEKE RQIKKQTALV ELVKHKKAT KEQLKAVMDD FAAFVEKCK ADDKETCF AE EGKKLV AASQ AALGL
ZP-10	H-His-Gly-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Ser-Lys-Lys-Lys-Lys-Lys-Lys-NH ₂
TOLBUTAMIDE	







INN or Research Code	Structure/Chemical Name
	<i>N</i> -[(butylamino)carbonyl]-4-methylbenzenesulfonamide
TOLAZAMIDE	 <i>N</i> -[(azepan-1-ylamino)carbonyl]-4-methylbenzenesulfonamide
GLIPIZIDE	 <i>N</i> -{2-[4-({[(cyclohexylamino)carbonyl]amino}sulfonyl)phenyl]ethyl}-5-methylpyrazine-2-carboxamide
CARBUTAMIDE	 4-amino- <i>N</i> -[(butylamino)carbonyl]benzenesulfonamide
GLISOXEPIDE	 <i>N</i> -{2-[4-({[(azepan-1-ylamino)carbonyl]amino}sulfonyl)phenyl]ethyl}-5-methylisoxazole-3-carboxamide
GLISENTIDE	 <i>N</i> -{2-[4-({[(cyclopentylamino)carbonyl]amino}sulfonyl)phenyl]ethyl}-2-methoxybenzamide
GLIBORNURIDE	 <i>N</i> -{[(3-hydroxy-4,7,7-trimethylbicyclo[2.2.1]hept-2-yl)amino]carbonyl}-4-methylbenzenesulfonamide
GLIBENCLAMIDE	 5-chloro- <i>N</i> -{2-[4-





INN or Research Code	Structure/Chemical Name
	(([(cyclohexylamino)carbonyl]amino)sulfonyl)phenyl]ethyl)-2-methoxybenzamide
GLIQUIDONE	 <i>N</i>-[(cyclohexylamino)carbonyl]-4-[2-(7-methoxy-4,4-dimethyl-1,3-dioxo-3,4-dihydroisoquinolin-2(1<i>H</i>)-yl)ethyl]benzenesulfonamide
GLIMEPIRIDE	 3-ethyl-4-methyl-<i>N</i>-(2-{4-[(<i>trans</i>-4-methylcyclohexyl)amino]carbonyl}amino)sulfonyl]phenyl)ethyl)-2-oxo-2,5-dihydro-1<i>H</i>-pyrrole-1-carboxamide
GLICLAZIDE	 <i>N</i>-[(hexahydrocyclopenta[<i>c</i>]pyrrol-2(1<i>H</i>)-ylamino)carbonyl]-4-methylbenzenesulfonamide
METFORMIN	 <i>N,N</i>-dimethylimidodicarbonimidic diamide
ACARBOSE	 4,6-dideoxy-4-[[4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}hexopyranosyl-(1→4)hexopyranosyl-(1→4)hexopyranose
MIGLITOL	 1-(2-hydroxyethyl)-2-(hydroxymethyl)piperidine-3,4,5-triol
VOGLIBOSE	







INN or Research Code	Structure/Chemical Name
	(1 <i>R</i> ,2 <i>R</i> ,3 <i>S</i> ,4 <i>R</i> ,5 <i>R</i>)-5-{[2-hydroxy-1-(hydroxymethyl)ethyl]amino}-1-(hydroxymethyl)cyclohexane-1,2,3,4-tetrol
MURAGLITAZAR	 <i>N</i>-[(4-methoxyphenoxy)carbonyl]-<i>N</i>-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]benzyl}glycine
ROSIGLITAZONE	 (5<i>RS</i>)-5-(4-{2-[methyl(pyridin-2-yl)amino]ethoxy}benzyl)-1,3-thiazolidine-2,4-dione
PIOGLITAZONE	 5-{4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl}-1,3-thiazolidine-2,4-dione
FARGLITAZAR	 <i>N</i>-(2-benzoylphenyl)-<i>O</i>-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethyl]-<i>L</i>-tyrosine
NAVEGLITAZAR	 (2<i>S</i>)-2-methoxy-3-{4-[3-(4-phenoxyphenoxy)propoxy]phenyl}propanoic acid
NETOGLITAZONE	 5-({6-[(2-fluorobenzyl)oxy]-2-naphthyl}methyl)-1,3-thiazolidine-2,4-dione

INN or Research Code	Structure/Chemical Name
RIVOGLITAZONE	 (5<i>R</i>)-5-{4-[(6-methoxy-1-methyl-1<i>H</i>-benzimidazol-2-yl)methoxy]benzyl}-1,3-thiazolidine-2,4-dione
K-111	 2,2-dichloro-12-(4-chlorophenyl)dodecanoic acid
SODELGLITAZAR =GW-677954	 2-ethyl-2-(4-[[4-[[4-(4-methoxyphenyl)piperazin-1-yl]methyl]-2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-5-yl)methyl]thio}phenoxy)butanoic acid
(-)-Halofenate	 (-)-2-Acetamidoethyl 4-chlorophenyl(3-trifluoromethylphenoxy)acetate
REPAGLINIDE	 2-ethoxy-4-(2-[[[(1<i>S</i>)-3-methyl-1-(2-piperidin-1-ylphenyl)butyl]amino]-2-oxoethyl]benzoic acid
NATEGLINIDE	 <i>N</i>-[(<i>trans</i>-4-isopropylcyclohexyl)carbonyl]-D-phenylalanine
MITIGLINIDE	 (2<i>S</i>)-2-benzyl-4-[(3<i>aR</i>,7<i>aS</i>)-octahydro-2<i>H</i>-isoindol-2-yl]-4-oxobutanoic acid

INN or Research Code	Structure/Chemical Name
	acid
SITAGLIPTIN	 (2<i>R</i>)-4-oxo-4-[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-<i>a</i>]pyrazin-7(8<i>H</i>)-yl]-1-(2,4,5-trifluorophenyl)butan-2-amine
SAXAGLIPTIN	 (1<i>S</i>,3<i>S</i>,5<i>S</i>)-2-[(2<i>S</i>)-2-amino-2-(3-hydroxy-1-adamantyl)acetyl]-2-azabicyclo[3.1.0]hexane-3-carbonitrile
VILDAGLIPTIN	 (2<i>S</i>)-1-[<i>N</i>-(3-hydroxy-1-adamantyl)glycyl]pyrrolidine-2-carbonitrile
DENAGLIPTIN	 (2<i>S</i>,4<i>S</i>)-4-fluoro-1-[4-fluoro-β-(4-fluorophenyl)-<i>L</i>-phenylalanyl]pyrrolidine-2-carbonitrile
ALOGLIPTIN	 2-({6-[(3<i>R</i>)-3-aminopiperidin-1-yl]-3-methyl-2,4-dioxo-3,4-dihydropyrimidin-1(2<i>H</i>)-yl}methyl)benzonitrile

INN or Research Code	Structure/Chemical Name
P32/98	 (2S,3S)-3-methyl-1-oxo-1-(1,3-thiazolidin-3-yl)pentan-2-amine
SILDENAFIL	 5-{2-ethoxy-5-[(4-methylpiperazin-1-yl)sulfonyl]phenyl}-1-methyl-3-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one
VARDENAFIL	 2-{2-ethoxy-5-[(4-ethylpiperazin-1-yl)sulfonyl]phenyl}-5-methyl-7-propylimidazo[5,1-f][1,2,4]triazin-4(3H)-one
TADALAFIL	 (6R,12aR)-6-(1,3-benzodioxol-5-yl)-2-methyl-2,3,6,7,12,12a-hexahydropyrazino[1',2':1,6]pyrido[3,4-b]indole-1,4-dione
UDENAFIL	 3-(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-N-[2-(1-methylpyrrolidin-2-yl)ethyl]-4-propoxybenzenesulfonamide
AVANAFIL	 4-[(3-chloro-4-methoxybenzyl)amino]-2-[(2S)-2-

INN or Research Code	Structure/Chemical Name
	(hydroxymethyl)pyrrolidin-1-yl]- <i>N</i> -(pyrimidin-2-ylmethyl)pyrimidine-5-carboxamide
PRAMLINTIDE	L-lysyl-L-cysteinyl-L-asparaginyl-L-threonyl-L-alanyl-L-threonyl-L-cysteinyl-L-alanyl-L-threonyl-L-glutaminy-L-arginyl-L-leucyl-L-alanyl-L-asparaginyl-L-phenylalanyl-L-leucyl-L-valyl-L-histidyl-L-seryl-L-seryl-L-asparaginyl-L-asparaginyl-L-phenylalanylglycyl-L-prolyl-L-isoleucyl-L-leucyl-L-prolyl-L-prolyl-L-threonyl-L-asparaginyl-L- valylglycyl-L-seryl-L-asparaginyl-L-threonyl-L-tyrosinamide, cyclic (2->7)disulfide
BAY-27-9955	 2-(4'-fluoro-3,5-diisopropyl-6-propylbiphenyl-2-yl)ethanol
INGLIFORIB	 <i>N</i>-{(1<i>S</i>,2<i>R</i>)-1-benzyl-3-[(3<i>R</i>,4<i>S</i>)-3,4-dihydroxypyrrolidin-1-yl]-2-hydroxy-3-oxopropyl}-5-chloro-1<i>H</i>-indole-2-carboxamide
MANAGLINAT DIALANTENIL = MB-06322	 ethyl 4-[5-(2-amino-5-isobutyl-1,3-thiazol-4-yl)-2-furyl]-2,6-dimethyl-7-oxo-8-oxa-3,5-diaza-4-phosphadecan-1-oate 4-oxide
MB-05032	 [5-(2-amino-5-isobutyl-1,3-thiazol-4-yl)-2-furyl]phosphonic acid

INN or Research Code	Structure/Chemical Name
AVE-1625	 1-[bis(4-chlorophenyl)methyl]-3-[(3,5-difluorophenyl)(methylsulfonyl)methylene]azetidine
RIMONABANT	 5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-N-piperidin-1-yl-1H-pyrazole-3-carboxamide
SURINABANT	 5-(4-bromophenyl)-1-(2,4-dichlorophenyl)-4-ethyl-N-piperidin-1-yl-1H-pyrazole-3-carboxamide
HMR-1426	 6-chloro-2-phenyl-8,8a-dihydro-3aH-indeno[1,2-d][1,3]thiazol-3a-ol
CETILISTAT	 2-(hexadecyloxy)-6-methyl-4H-3,1-benzoxazin-4-one
SIBUTRAMINE	 1-[1-(4-chlorophenyl)cyclobutyl]-3-methylbutan-1-amine

Additional information with regard to the preparation, suitable dosage forms and dose ranges of the glucagon-like-peptide-1 receptor agonists listed in Table 1 can be found in the following patents/patent applications: WO0334331, EP0981611, WO9808871, WO0104156 and WO03059934.

5 The sulfonylurea agents TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXE-
PIDE; GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE, GLIMEPIRIDE and GLI-
CLAZIDE listed in Table 1 are commercially available. The person skilled in the art is familiar with
suitable formulations and dose ranges of these compounds.

10 The biguanide agent METFORMIN listed in Table 1 is commercially available. The person skilled in
the art is familiar with suitable formulations and dose ranges of this compound.

The alpha-glucosidase inhibitors ACARBOSE, MIGLITOL and VOGLIBOSE listed in Table 1 are com-
mercially available. The person skilled in the art is familiar with suitable formulations and dose ranges
15 of this compound.

Additional information with regard to the preparation, suitable dosage forms and dose ranges of the
PPAR-agonists listed in Table 1 can be found in the following patents/patent applications:
WO0121602, EP03306228, EP0658161, EP0193256, WO9731907, WO0140169, WO02100813,
20 EP0604983, EP0745600, WO9615784, WO0259098 and EP1183020.

The metiglinide agents REPAGLINIDE, NATEGLINIDE and MITIGLINIDE listed in Table 1 are com-
mercially available. The person skilled in the art is familiar with suitable formulations and dose ranges
of this compound.

25 Additional information with regard to the preparation, suitable dosage forms and dose ranges of the
DPP IV inhibitors listed in Table 1 can be found in the following patents/patent applications:
WO03004498, WO0168603, WO0034241, WO0302531, WO9961431 and WO2005095381.

30 Additional information with regard to the preparation, suitable dosage forms and dose ranges of the
PDE5 inhibitors listed in Table 1 can be found in the following patents/patent applications:
WO0213798, WO0260422, WO2004082667, WO0027848 and EP1219609.

35 Additional information with regard to the preparation, suitable dosage forms and dose ranges of the
amylin analogue PRAMLINTIDE listed in Table 1 can be found in EP0567626.

Additional information with regard to the preparation, suitable dosage forms and dose ranges of the
glucagon receptor antagonist listed in Table 1 can be found in WO9804528.

Additional information with regard to the preparation, suitable dosage forms and dose ranges of the glycogen-phosphorylase inhibitor listed in Table 1 can be found in WO9639385.

5 Additional information with regard to the preparation, suitable dosage forms and dose ranges of the fructose-1,6-bisphosphate inhibitors can be found in WO0001495 and WO0147935.

Additional information with regard to the preparation, suitable dosage forms and dose ranges of the cannabinoid (CB1) receptor antagonist listed in Table 1 can be found in EP1112251, EP0576357 and WO0046209.

10

Additional information with regard to the preparation, suitable dosage forms and dose ranges of HMR-1426, CETILISTAT and SIBUTRAMINE listed in Table 1 can be found in the following patents/patent applications: WO0018749, EP1144395 and EP0397831.

15 "Pharmaceutically acceptable salts" of Compound A or the other active compound(s) which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 are not limited to the specific examples given above. The term refers to non-toxic salts of these compounds. These pharmaceutically acceptable salts are generally prepared by reacting a free base with a suitable organic or inorganic acid or by reacting an acid with a suitable organic or inorganic base. Particular mention may be made of the
20 pharmaceutically acceptable inorganic and organic acids customarily used in pharmacy. Those suitable are in particular water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)-benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, em-
25 bonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 1-hydroxy-2-naphthoic acid. As examples of pharmaceutically acceptable salts with bases may be mentioned the lithium, sodium, potassium, calcium, aluminium, magnesium, titanium, ammonium, meglumine or guanidinium salts.

It is understood that Compound A and the other active compound(s) which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 and their pharmaceutically acceptable salts can also be
30 present in the form of their pharmaceutically acceptable solvates, and in particular in the form of their hydrates.

The combinations according to the invention may be administered by any suitable route, for example,
35 by the oral, sublingual, buccal, intravenous, intraarterial, intramuscular, subcutaneous, intracutaneous, topical, transdermal, intranasal, intraperitoneal, rectal or vaginal route, by inhalation or by insufflation.

Tablets, coated tablets (dragees), pills, cachets, capsules (caplets), granules, solutions, emulsions and suspensions are e.g. suitable for oral administration. In particular, said formulations can be adapted so
40 as to represent, for example, an enteric form, an immediate release form, a delayed release form, a

repeated dose release form, a prolonged release form or a sustained release form. Said forms can be obtained, for example, by coating tablets, by dividing tablets into several compartments separated by layers disintegrating under different conditions (e.g. pH conditions) or by coupling the active compound to a biodegradable polymer.

5

Administration by inhalation is preferably made by using an aerosol. The aerosol is a liquid-gaseous dispersion, a solid-gaseous dispersion or a mixed liquid/solid-gaseous dispersion.

The aerosol may be generated by means of aerosol-producing devices such as dry powder inhalers (DPIs), pressurized metered dose inhalers (PMDIs) and nebulizers. Depending on the kind of the active compound to be administered, the aerosol-producing device can contain the active compound in form of a powder, a solution or a dispersion. The powder may contain, for example, one or more of the following auxiliaries: carriers, stabilizers and fillers. The solution may contain in addition to the solvent, for example, one or more of the following auxiliaries: propellants, solubilizers (co-solvents), surfactants, stabilizers, buffers, tonicity adjusting agents, preservatives and flavorings. The dispersion may contain in addition to the dispersant, for example, one or more of the following auxiliaries: propellants, surfactants, stabilizers, buffers, preservatives and flavorings. Examples of carriers include, but are not limited to, saccharides, e.g. lactose and glucose. Examples of propellants include, but are not limited to, fluorohydrocarbons, e.g. 1,1,1,2-tetrafluoroethane and 1,1,1,2,3,3,3-heptafluoropropane.

The particle size of the aerosol particles (solid, liquid or solid/liquid particles) is preferably less than 100 µm, more preferably it is in the range of from 0.5 to 10 µm, in particular in the range of from 2 to 6 µm (D50 value, measured by laser diffraction).

25

For parenteral modes of administration such as, for example, intravenous, intraarterial, intramuscular, subcutaneous, intracutaneous and intraperitoneal administration, preferably solutions (e.g. sterile solutions, isotonic solutions) are used. They are preferably administered by injection or infusion techniques.

30

The pharmaceutical compositions (formulations) comprising Compound A or a pharmaceutically acceptable salt thereof and/or one or two other active compound(s) or pharmaceutically acceptable salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 and at least one pharmaceutically acceptable auxiliary can be manufactured in a manner known to a person skilled in the art, e.g. by dissolving, mixing, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

35

As pharmaceutically acceptable auxiliaries, any auxiliaries known to be suitable for preparing pharmaceutical compositions (formulations) can be used. Examples thereof include, but are not limited to, solvents, excipients, dispersants, emulsifiers, solubilizers, gel formers, ointment bases, antioxidants,

40

preservatives, stabilizers, carriers, fillers, binders, thickeners, complexing agents, disintegrating agents, buffers, permeation promoters, polymers, lubricants, coating agents, propellants, tonicity adjusting agents, surfactants, colorants, flavorings, sweeteners and dyes. In particular, auxiliaries of a type appropriate to the desired formulation and the desired mode of administration are used.

5

The preferred mode of administration of the combinations according to the invention depend on the specific combination partners.

As mentioned above Compound A or a pharmaceutically acceptable salt of either may be administered in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes or suppositories. The preferred form depends on the intended mode of administration and the combination partner.

15 The most preferred mode of administration of Compound A or a pharmaceutically acceptable salt of either is oral. In another preferred embodiment Compound A or a pharmaceutically acceptable salt of either is administered by intravenous infusion or injection. In a further embodiment Compound A or a pharmaceutically acceptable salt of either is administered by intramuscular or subcutaneous injection. Other routes of administration are also contemplated, including for example intranasal and transdermal routes, and by inhalation.

20 The preferred mode of administration of the other active compound(s) which is (are) used in combination with Compound A or a pharmaceutically acceptable salt of either depends on the specific agent.

25 EXENATIDE, BIM-51077, ALBIGLUTIDE, ZP-10 or PRAMLINTIDE, for example, are preferably administered via subcutaneous injection. The preferred mode of administration of compounds like TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE, GLIMEPIRIDE, GLICLAZIDE, METFORMIN, ACARBOSE, MIGLITOL, VOGLIBOSE, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-HALOFENATE, 30 REPAGLINIDE, NATEGLINIDE, MITIGLINIDE, SITAGLIPTIN, SAXAGLIPTIN, VILDAGLIPTIN, DENAGLIPTIN, ALOGLIPTIN, P32/98, SILDENAFIL; VARDENAFIL, TADALAFIL, UDENAFIL, AVANAFIL, BAY-27-9955, INFLIGORIB, MB-06322, MB-05022, AVE-1625, RIMONABANT, SURINABANT, HMR-1426, CETILISTAT, SIBUTRAMINE and cinnamon is oral. Further information with regard to the preferred mode of administration of the other active agent(s) which is (are) used in combination with Compound A or a pharmaceutically acceptable salt of either is summarized in Table 2 below.

40 As part of the combination therapy according to the invention Compound A or a pharmaceutically acceptable salt thereof and the one or two other active compound(s) or pharmaceutically acceptable

salt(s) thereof which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 are dosed in an order of magnitude customary for the mono-therapy, it more likely being possible, on account of the individual actions, which are mutually positively influencing and reinforcing, to reduce the respective doses on the combined administration of Compound A or a pharmaceutically acceptable salt thereof and the one or two other active compound(s) which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 with the norm.

In the case of oral administration of Compound A the daily dose for an adult patient for the mono-therapy is in the range from 0.2 to 30 mg per day, preferably in the range of 0.2 to 10 mg per day, more preferably in the range of 0.5 to 5 mg per day.

Further information with regard to the preferred routes of administration and typical dosages (for mono-therapy) of the other active compound(s) which is (are) used in combination with Compound A is summarized in Table 2.

The proportions in which Compound A or a pharmaceutically acceptable salt thereof and the other active compound(s) or a pharmaceutically acceptable salt thereof may be used in the pharmaceutical compositions, combinations products or kits according to the invention are variable. Depending on the choice of the Compound A or a pharmaceutically acceptable salt thereof and the choice of the other active compound(s) or a pharmaceutically acceptable salt thereof, the weight ratios which may be used within the scope of the present invention vary on the basis of the different molecular weights of the various compounds and their different potencies.

As a general rule, without intended to be limiting, the pharmaceutical compositions, combination products or kits according to the invention may contain

Compound A and BIM-51077 in ratios by weight ranging from 50:1 to 1:2, preferably from 20:1 to 1:1; Compound A and EXENATIDE in ratios by weight ranging from 500:1 to 25:1, preferably from 200:1 to 50:1;

Compound A and LIRAGLUTIDE in ratios by weight ranging from 10:1 to 1:1, preferably from 5:1 to 2:1;

Compound A and TOLBUTAMIDE in ratios by weight ranging from 1:4000 to 1:100, preferably from 1:1000 to 1:250;

Compound A and TOLAZAMIDE in ratios by weight ranging from 1:300 to 1:20, preferably from 1:150 to 1:50;

Compound A and GLIPIZIDE in ratios by weight ranging from 1:80 to 1:1, preferably from 1:40 to 1:2;

Compound A and GLISOXEPIDE in ratios by weight ranging from 1:32 to 2:1, preferably from 1:10 to 1:1;

Compound A and GLIBORNURIDE in ratios by weight ranging from 1:150 to 1:3; preferably from 1:50 to 1:6;

Compound A and GLIBENCLAMIDE in ratios by weight ranging from 1:20 to 3:1, preferably from 1:5 to 1:1;

Compound A and GLIQUIDONE in ratios by weight ranging from 1:250 to 1:3, preferably from 1:60 to 1:6;

- 5 Compound A and GLIMEPIRIDE in ratios by weight ranging from 1:12 to 5:1, preferably from 1:4 to 2:1;

Compound A and GLICLAZIDE in ratios by weight ranging from 1:250 to 1:6, preferably from 1:60 to 1:15;

- 10 Compound A and METFORMIN in ratios by weight ranging from 1:8000 to 1:200, preferably from 1:2000 to 1:500;

Compound A and ACARBOSE in ratios by weight ranging from 1:1200 to 1:30, preferably from 1:300 to 1:75;

Compound A and MIGLITOL in ratios by weight ranging from 1:600 to 1:30, preferably from 1:150 to 1:75;

- 15 Compound A and VOGLIBOSE in ratios by weight ranging from 5:1 to 1:2, preferably from 2:1 to 1:1;
Compound A and MURAGLITAZAR in ratios by weight ranging from 1:10 to 2:1, preferably from 1:5 to 1:1;

Compound A and ROSIGLITAZONE in ratios by weight ranging from 1:16 to 1:1, preferably from 1:8 to 1:2;

- 20 Compound A and PIOGLITAZONE in ratios by weight ranging from 1:90 to 1:3, preferably from 1:45 to 1:8;

Compound A and FARGLITAZAR in ratios by weight ranging from 1:20 to 10:1, preferably from 1:10 to 4:1;

- 25 Compound A and NAVEGLITAZAR in ratios by weight ranging from 50:1 to 1:2, preferably from 20:1 to 1:1;

Compound A and NETOGLITAZONE in ratios by weight ranging from 1:100 to 5:1, preferably from 1:50 to 2:1;

Compound A and RIVOGLITAZONE in ratios by weight ranging from 1:20 to 5:1, preferably from 1:10 to 2:1;

- 30 Compound A and K-111 in ratios by weight ranging from 1:40 to 1:2, preferably from 1:20 to 1:5;

Compound A and GW-677954 in ratios by weight ranging from 1:20 to 2:1, preferably from 1:10 to 1:1;

Compound A and (-)-HALOFENATE in ratios by weight ranging from 1:2000 to 1:200, preferably from 1:1000 to 1:500;

- 35 Compound A and REPAGLINIDE in ratios by weight ranging from 1:32 to 10:1, preferably from 1:16 to 4:1;

Compound A and NATEGLINIDE in ratios by weight ranging from 1:1000 to 1:40, preferably from 1:500 to 1:100;

- 40 Compound A and MITIGLINIDE in ratios by weight ranging from 1:250 to 1:8, preferably from 1:120 to 1:20;

Compound A and SITAGLIPTIN in ratios by weight ranging from 1:200 to 1:20, preferably from 1:100 to 1:50;

Compound A and SAXAGLIPTIN in ratios by weight ranging from 1:20 to 1:2; preferably from 1:10 to 1:2;

- 5 Compound A and VILDAGLIPTIN in ratios by weight ranging from 1:200 to 1:5, preferably from 1:100 to 1:10;

Compound A and DENAGLIPTIN in ratios by weight ranging from 1:100 to 1:1, preferably from 1:50 to 1:2;

- 10 Compound A and ALOGLIPTIN in ratios by weight ranging from 1:100 to 1:2, preferably from 1:50 to 1:5;

Compound A and P32/98 in ratios by weight ranging from 1:200 to 1:6, preferably from 1:100 to 1:15;

Compound A and SILDENAFIL in ratios by weight ranging from 1:200 to 1:10, preferably from 1:100 to 1:25;

- 15 Compound A and VARDENAFIL in ratios by weight ranging from 1:40 to 2:1, preferably from 1:20 to 1:1;

Compound A and TADALAFIL in ratios by weight ranging from 1:40 to 1:2, preferably from 1:20 to 1:5;

Compound A and UDENAFIL in ratios by weight ranging from 1:400 to 1:20, preferably 1:200 to 1:50;

Compound A and AVANAFIL in ratios by weight ranging from 1:600 to 1:10, preferably from 1:300 to 1:25;

- 20 Compound A and PRIMLINTIDE in ratios by weight ranging from 250:1 to 5:1, preferably from 100:1 to 10:1;

Compound A and cinnamon in ratios by weight ranging from 1:12000 to 1:200, preferably from 1:6000 to 1:500;

- 25 Compound A and BAY-27-9955 in ratios by weight ranging from 1:400 to 1:10, preferably from 1:2000 to 1:25;

Compound A and INGLIFORIB in ratios by weight ranging from 1:100 to 1:4, preferably from 1:50 to 1:10;

Compound A and MB-06322 in ratios by weight ranging from 1:1600 to 1:40, preferably from 1:800 to 1:100;

- 30 Compound A and AVE-1625 in ratios by weight ranging from 1:20 to 1:2, preferably from 1:10 to 1:5;
Compound A and RIMONABANT in ratios by weight ranging from 1:40 to 1:4, preferably from 1:20 to 1:10;

Compound A and SURINABANT in ratios by weight ranging from 1:20 to 2:1, preferably from 1:10 to 1:1;

- 35 Compound A and CETILISTAT in ratios by weight ranging from 1:1800 to 1:25, preferably from 1:900 to 1:60;

Compound A and SIBUTRAMINE in ratios by weight ranging from 1:30 to 1:2, preferably from 1:15 to 1:5;

Corresponding ratios likely may be used in the triple combinations according to the invention.

Table 2: Preferred routes of administration and dosages:

INN or Research Code	Preferred Treatment of	Preferred route of Administration	Typical Daily dose (dose ranges) used for mono-therapy
Insulin/Insulin analogs	Diabetes mellitus type 1 Diabetes mellitus type 2	subcutaneous injection	On demand
EXUBERA	Diabetes mellitus type 1 Diabetes mellitus type 2	Inhalation	On demand
BIM-51077	Diabetes mellitus type 2	subcutaneous injection	100 - 800 µg
EXENATIDE	Diabetes mellitus type 2	subcutaneous injection	10-20 µg
LIRAGLUTIDE	Diabetes mellitus type 2	subcutaneous injection	0.5 – 2 mg
ALBIGLUTIDE	Diabetes mellitus type 2	subcutaneous injection	2-20 µg
ZP-10	Diabetes mellitus type 2	subcutaneous injection	
TOLBUTAMIDE	Diabetes mellitus type 2	oral	0.5 to 2.0 g
TOLAZAMIDE	Diabetes mellitus type 2	oral	100 to 150 mg
GLIPIZIDE	Diabetes mellitus type 2	oral	5 to 40 mg, preferably 5 to 20 mg
CARBUTAMIDE	Diabetes mellitus type 2	oral	up to 17 mg/kg
GLISOXEPIDE	Diabetes mellitus type 2	oral	2 to 16 mg
GLISENTIDE	Diabetes mellitus type 2	oral	
GLIBORNURIDE	Diabetes mellitus type 2	oral	12.5 to 75 mg
GLIBENCLAMIDE	Diabetes mellitus type 2	oral	1.75 to 10.5 mg
GLIQUIDONE	Diabetes mellitus type 2	oral	15 to 120 mg
GLIMEPIRIDE	Diabetes mellitus type 2	oral	1 to 6 mg
GLICLAZIDE	Diabetes mellitus type 2	oral	30 to 120 mg
METFORMIN	Diabetes mellitus type 2	oral	1000 to 3800 mg
ACARBOSE	Diabetes mellitus type 2	oral	150 to 600 mg, preferably 150 to 300 mg
MIGLITOL	Diabetes mellitus type 2	oral	150 to 300 mg
VOGLIBOSE	Diabetes mellitus type 2	oral	0.6 to 0.9 mg
MURAGLITAZAR	Diabetes mellitus type 2	oral or by injection	2.5 to 5 mg
ROSIGLITAZONE	Diabetes mellitus type 2	oral	4 to 8 mg

INN or Research Code	Preferred Treatment of	Preferred route of Administration	Typical Daily dose (dose ranges) used for mono-therapy
PIOGLITAZONE	Diabetes mellitus type 2	oral	15 to 45 mg
FARGLITAZAR	Diabetes mellitus type 2	oral	0.5 to 10 mg
NAVEGLITAZAR	Diabetes mellitus type 2	oral	0.004 to 1.2 mg
NETOGLITAZONE	Diabetes mellitus type 2	oral	1-50 mg
RIVOGLITAZONE	Diabetes mellitus type 2	oral	1-10 mg
K-111	Diabetes mellitus type 2	oral	10 to 20 mg
GW-677954	Diabetes mellitus type 2	oral	2.5 to 20 mg
(-)-Halofenate	Diabetes mellitus type 2	oral	≅ 1000 mg
REPAGLINIDE	Diabetes mellitus type 2	oral	0.5 to 16 mg
NATEGLINIDE	Diabetes mellitus type 2	oral	180 to 540 mg
MITIGLINIDE	Diabetes mellitus type 2	oral	40 mg/meal
SITAGLIPTIN	Diabetes mellitus type 2	oral	≅ 100 mg
SAXAGLIPTIN	Diabetes mellitus type 2	oral	≅ 10 mg
VILDAGLIPTIN	Diabetes mellitus type 2	oral	25-100 mg
DENAGLIPTIN	Diabetes mellitus type 2	oral	5-50 mg
ALOGLIPTIN	Diabetes mellitus type 2	oral	10-50 mg
P32/98	Diabetes mellitus type 2	oral	30 – 100 mg
SILDENAFIL	Diabetes mellitus type 2 Diabetes mellitus type 1	oral	50 to 100 mg
VARDENAFIL	Diabetes mellitus type 2 Diabetes mellitus type 1	oral	2.5 to 20 mg
TADALAFIL	Diabetes mellitus type 2 Diabetes mellitus type 1	oral	10 to 20 mg
UDENAFIL	Diabetes mellitus type 2 Diabetes mellitus type 1	oral	100-200 mg
AVANAFIL	Diabetes mellitus type 2 Diabetes mellitus type 1	oral	50-300 mg
PRAMLINTIDE	Diabetes mellitus type 2 Diabetes mellitus type 1	subcutaneous in- jection	20 to 120 µg
cinnamon	Diabetes mellitus type 2	oral	1 to 6 g
BAY-27-9955	Diabetes mellitus type 2	oral	50 to 200 mg
INGLIFORIB	Diabetes mellitus type 2	oral	20 to 50 mg
MB-06322	Diabetes mellitus type 2	oral	200 to 800 mg
MB-05032	Diabetes mellitus type 2	oral	
AVE-1625	Diabetes mellitus type 2	oral	≅ 10 mg
RIMONABANT	Diabetes mellitus type 2	oral	20 mg

INN or Research Code	Preferred Treatment of	Preferred route of Administration	Typical Daily dose (dose ranges) used for mono-therapy
SURINABANT	Diabetes mellitus type 2	oral	3 to 10 mg
HMR-1426	Diabetes mellitus type 2	oral	
CETILISTAT	Diabetes mellitus type 2	oral	120 to 920 mg
SIBUTRAMINE	Diabetes mellitus type 2	oral	10 to 15 mg

Examples

Table 3: Preferred combinations

5

Example Number	Combination	
1	Compound A*	human insulin
2	Compound A*	Insulin analogue
3	Compound A*	BIM-51077
4	Compound A*	EXENATIDE
5	Compound A*	LIRAGLUTIDE
6	Compound A*	ALBIGLUTIDE
7	Compound A*	ZP-10
8	Compound A*	TOLBUTAMIDE
9	Compound A*	TOLBUTAMIDE sodium
10	Compound A*	TOLAZAMIDE
11	Compound A*	GLIPIZIDE
12	Compound A*	CARBUTAMIDE
13	Compound A*	GLISOXEPIDE
14	Compound A*	GLISENTIDE
15	Compound A*	GLIBORNURIDE
16	Compound A*	GLIBENCLAMIDE
17	Compound A*	GLIQUIDONE
18	Compound A*	GLIQUIDONE sodium
19	Compound A*	GLIMEPIRIDE
20	Compound A*	GLICLAZIDE
21	Compound A*	METFORMIN
22	Compound A*	METFORMIN hydrochloride

Example Number	Combination	
23	Compound A*	ACARBOSE
24	Compound A*	MIGLITOL
25	Compound A*	VOGLIBOSE
26	Compound A*	MURAGLITAZAR
27	Compound A*	ROSIGLITAZONE
28	Compound A*	ROSIGLITAZONE maleate
29	Compound A*	PIOGLITAZONE
30	Compound A*	PIOGLITAZONE dihydrochloride
31	Compound A*	FARGLITAZAR
32	Compound A*	NAVEGLITAZAR
33	Compound A*	NETOGLITAZONE
34	Compound A*	RIVOGLITAZONE
35	Compound A*	RIVOGLITAZONE hydrochloride
36	Compound A*	K-111
37	Compound A*	K-111 sodium
38	Compound A*	GW-677954
39	Compound A*	(-)-Halofenate
40	Compound A*	REPAGLINIDE
41	Compound A*	NATEGLINIDE
42	Compound A*	MITIGLINIDE
42	Compound A*	MITIGLINIDE potassium
44	Compound A*	MITIGLINIDE calcium
45	Compound A*	SITAGLIPTIN
46	Compound A*	SITAGLIPTIN phosphate
47	Compound A*	SAXAGLIPTIN
48	Compound A*	VILDAGLIPTIN
49	Compound A*	DENAGLIPTIN
50	Compound A*	ALOGLIPTIN
51	Compound A*	ALOGLIPTIN benzoate
52	Compound A*	P32/98
53	Compound A*	P32/98 fumarate
54	Compound A*	P32/98 hydrochloride
55	Compound A*	SILDENAFIL
56	Compound A*	SILDENAFIL citrate
57	Compound A*	SILDENAFIL hemi-citrate

Example Number	Combination	
58	Compound A*	SILDENAFIL mesilate
59	Compound A*	VARDENAFIL
60	Compound A*	VARDENAFIL hydrochloride
61	Compound A*	VARDENAFIL dihydrochloride
62	Compound A*	TADALAFIL
63	Compound A*	UDENAFIL
64	Compound A*	AVANAFIL
65	Compound A*	AVANAFIL besilate
66	Compound A*	PRAMLINTIDE
67	Compound A*	PRAMLINTIDE acetate
68	Compound A*	cinnamon
69	Compound A*	BAY-27-9955
70	Compound A*	INGLIFORIB
71	Compound A*	MB-06322
72	Compound A*	MB-05032
73	Compound A*	MB-05032 hydrobromide
74	Compound A*	AVE-1625
75	Compound A*	RIMONABANT
76	Compound A*	RIMONABANT hydrochloride
77	Compound A*	SURINABANT
78	Compound A*	HMR-1426
79	Compound A*	CETILISTAT
80	Compound A*	SIBUTRAMINE
81	Compound A*	SIBUTRAMINE hydrochloride

Compound A* represents Compound A or a pharmaceutically acceptable salt of Compound A, in particular the hydrochloride salt of Compound A.

Table 4: Preferred triple combinations:

Example Number	Triple Combination		
82	Compound A*	METFORMIN	Human Insulin
83	Compound A*	METFORMIN hydrochloride	Human Insulin
84	Compound A*	ROSIGLITAZONE	Human Insulin
85	Compound A*	ROSIGLITAZONE maleate	Human Insulin
86	Compound A*	ROSIGLITAZONE	METFORMIN
87	Compound A*	ROSIGLITAZONE maleate	METFORMIN
88	Compound A*	ROSIGLITAZONE maleate	METFORMIN hydrochloride
89	Compound A*	PIOGLITAZONE	Insulin
90	Compound A*	PIOGLITAZONE dihydrochloride	Insulin
91	Compound A*	PIOGLITAZONE	METFORMIN
92	Compound A*	PIOGLITAZONE	METFORMIN hydrochloride
93	Compound A*	PIOGLITAZONE dihydrochloride	METFORMIN
94	Compound A*	PIOGLITAZONE dihydrochloride	METFORMIN hydrochloride
95	Compound A*	GLIMEPIRIDE	Insulin
96	Compound A*	GLIMEPIRIDE	METFORMIN
97	Compound A*	GLIMEPIRIDE	METFORMIN hydrochloride
98	Compound A*	GLIMEPIRIDE	ROSIGLITAZONE
99	Compound A*	GLIMEPIRIDE	ROSIGLITAZONE maleate
100	Compound A*	GLIMEPIRIDE	PIOGLITAZONE
101	Compound A*	GLIMEPIRIDE	PIOGLITAZONE dihydrochloride

Compound A* represents Compound A or a pharmaceutically acceptable salt of Compound A, in particular the hydrochloride salt of Compound A.

Pharmacology

Aim of the study was to investigate the influence of a combination of RIMONABANT Hydrochloride and Compound A on HbA_{1c} and glucose tolerance.

5

Animals

Female obese db/db-mice (BKS.Cg/BomTac-*m* +/+ *Lepr^{db}*), supplied by Taconic Europe (Denmark) were used at the age of 10-11 weeks. Animals were kept under controlled conditions (22°C, 12h light / 12h dark cycle) and received standard laboratory chow and water ad libitum. For each group a number of 15 animals were used.

10

Experimental Protocol

Animals were treated with Compound A [3mg/kg orally, suspended in methocel (hypromellose, 4% aqueous solution)], RIMONABANT Hydrochloride [10mg/kg orally, suspended in methocel (hypromellose, 4% aqueous solution), Sequoia Research Products, Pangbourne, UK] or a combination of both once daily for 4 weeks. Vehicle treated animals served as control. HbA_{1c} levels were analyzed using the Micromat II Hemoglobin A1c Test (Bio-Rad Laboratories GmbH, 80939 München, Germany). An intraperitoneal glucose tolerance test was performed in 24h fasted animals. A glucose bolus (1g/kg body weight) was administered intraperitoneally. Blood glucose was monitored for the following time points: t = -120min, 0min (before glucose administration), 30min, 60min, 90min, 120min (after glucose administration). Blood was taken by tail-tip-bleeding (5µl) and was determined from hemolysate using an analyzing machine (Biosen S_line, EKF Diagnostic, Germany). Data are given as mean +/- SEM.

20

25

Results

Treatment with Compound A or RIMONABANT Hydrochloride alone had no effect on HbA_{1c} (Fig. 1), the relevant diagnostic parameter for type 2 diabetes mellitus. In contrast, the combination of both compounds resulted in a decrease of HbA_{1c} (Fig. 1). Treatment with Compound A or RIMONABANT Hydrochloride alone had no effect on glucose tolerance (Fig. 2). In contrast, the combination of both compounds resulted in an improvement of glucose tolerance (Fig. 2).

30

Conclusion

It is shown that a combination of RIMONABANT Hydrochloride and Compound A demonstrate over additive (synergistic) effects in the db/db mice model and therefore Compound A may be a ideal drug for combinational therapy to treat type 2 diabetes mellitus.

35

Description of the Figures:

5 **Fig 1: Effect of RIMONABANT Hydrochloride, Compound A and a combination of RIMONABANT Hydrochloride and Compound A on HbA_{1c} in db/db-mice**

Fig 2: Effect of RIMONABANT Hydrochloride, Compound A and a combination of RIMONABANT Hydrochloride and Compound A on glucose tolerance in db/db-mice

Patent claims

1. A pharmaceutical composition comprising a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.
2. A pharmaceutical composition comprising a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.
3. A combination product comprising the components: (A) Compound A or a pharmaceutically acceptable salt thereof; and (B) one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein each of the components (A) and (B) is formulated in admixture with at least one pharmaceutically acceptable auxiliary.
4. A combination product comprising the components: (A) Compound A or a pharmaceutically acceptable salt thereof; (B) one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1; and (C) still another active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein each of the components (A), (B) and (C) is formulated in admixture with at least one pharmaceutically acceptable auxiliary.
5. A kit comprising the components: (A) a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, in admixture with at least one pharmaceutically acceptable auxiliary; and (B) a pharmaceutical formulation including one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary.
6. A kit comprising the components: (A) a pharmaceutical formulation including Compound A or a pharmaceutically acceptable salt thereof, in admixture with at least one pharmaceutically acceptable auxiliary; (B) a pharmaceutical formulation including one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary; and (C) a pharmaceutical formulation including still another active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, in admixture with at least one pharmaceutically acceptable auxiliary.

7. Pharmaceutical composition, combination product or kit according to any of claims 1 to 6 wherein the other active compound(s) which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 is (are independently) selected from the group consisting of insulin, insulin analogues, glucagon-like-peptide-1 receptor agonists, sulfonylurea agents, biguanide agents, alpha-glucosidase inhibitors, PPAR-agonists, meglitinide agents, dipeptidyl-peptidase IV inhibitors, PDE1 inhibitors, PDE5 inhibitors, PDE9 inhibitors, PDE10 inhibitors, PDE11 inhibitors, amylin analogues, cinnamon, glucagon receptor antagonists, glycogen phosphorylase inhibitors, fructose-1,6-bisphosphate inhibitors, cannabinoid (CB1) receptor antagonists, anti-obesity drugs and the pharmaceutically acceptable salts of these compounds.

8. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin.

9. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an insulin analogue.

10. Pharmaceutical composition, combination product or kit according to any one of claims 7 and 9 wherein the insulin analogue is selected from the group consisting of novarapid, insulin detemir, insulin lispro, insulin glargine, insulin zinc suspension and LYS-Pro insulin.

11. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glucagon-like-peptide-1 receptor agonist or a pharmaceutically acceptable salt thereof.

12. Pharmaceutical composition, combination product or kit according to any of claims 7 and 11 wherein the glucagon-like-peptide-1 receptor agonist is selected from the group consisting of BIM-51077, EXENATIDE, LIRAGLUTIDE, ALBIGLUTIDE, ZP-10 and the pharmaceutically acceptable salts of these compounds.

13. Pharmaceutical composition, combination product or kit according to any of claims 7 and 11 wherein the glucagon-like-peptide-1 receptor agonist is selected from the group consisting of BIM-51077, EXENATIDE, LIRAGLUTIDE, ZP-10 and the pharmaceutically acceptable salts of these compounds.

14. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof.

15. Pharmaceutical composition, combination product or kit according to any of claims 7 and 14 wherein the sulfonylurea agent is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE GLIMEPIRIDE, GLICLAZIDE and the pharmaceutically acceptable salts of these compounds.

16. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

17. Pharmaceutical composition, combination product or kit according any of claims 7 and 16 wherein the biguanide agent is METFORMIN or a pharmaceutically acceptable salt of this compound.

18. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a alpha-glucosidase inhibitor or a pharmaceutically acceptable salt thereof.

19. Pharmaceutical composition, combination product or kit according to any of claims 7 and 18 wherein the alpha-glucosidase inhibitor is selected from the group consisting of ACARBOSE, MIGLITOL, VOGLIBOSE and the pharmaceutically acceptable salts of these compounds.

20. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR-agonist or a pharmaceutically acceptable salt thereof.

21. Pharmaceutical composition, combination product or kit according to any of claims 7 and 20 wherein the PPAR-agonist is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically acceptable salts of these compounds.

22. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a meglitinide agent or a pharmaceutically acceptable salt thereof.

23. Pharmaceutical composition, combination product or kit according to any of claims 7 and 22 wherein the meglitinide agent is selected from the group consisting of REPAGLINIDE, NATEGLINIDE, MITIGLINIDE and the pharmaceutically acceptable salts of these compounds.

5 **24.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a dipeptidyl-peptidase IV inhibitor or a pharmaceutically acceptable salt thereof.

10 **25.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 24 wherein the dipeptidyl-peptidase IV inhibitor is selected from the group consisting of SITAGLIPTIN, SAXAGLIPTIN, VILDAGLIPTIN, DENAGLIPTIN, ALOGLIPTIN, P32/98 and the pharmaceutically acceptable salts of these compounds.

15 **26.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 24 wherein the dipeptidyl-peptidase IV inhibitor is selected from the group consisting of SITAGLIPTIN, SAXAGLIPTIN, VILDAGLIPTIN, DENAGLIPTIN, P32/98 and the pharmaceutically acceptable salts of these compounds.

20 **27.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PDE5 inhibitor or a pharmaceutically acceptable salt thereof.

25 **28.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 27 wherein the PDE5 inhibitor is selected from the group consisting of SILDENAFIL, VARDENFIL, TADALAFIL, UDENAFIL, AVANAFIL and the pharmaceutically acceptable salts of these compounds.

30 **29.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 27 wherein the PDE5 inhibitor is selected from the group consisting of SILDENAFIL, VARDENFIL, TADALAFIL and the pharmaceutically acceptable salts of these compounds.

35 **30.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of a PDE1, PDE9, PDE10, PDE11 inhibitor and the pharmaceutically acceptable salts thereof.

31. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a amylin analogue or a pharmaceutically acceptable salt thereof.

32. Pharmaceutical composition, combination product or kit according to any of claims 7 and 31 wherein the amylin analogue inhibitor is PRAMLINTIDE or a pharmaceutically acceptable salt of this compound.

5 **33.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is cinnamon.

10 **34.** Pharmaceutical composition, combination product or kit according to claim 7 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is cinnamon.

35. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glucagon receptor antagonist or a pharmaceutically acceptable salt thereof.

15 **36.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 35 wherein the glucagon receptor antagonist is BAY-27-9955 or a pharmaceutically acceptable salt thereof.

20 **37.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glycogen phosphorylase inhibitor or a pharmaceutically acceptable salt thereof.

25 **38.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 37, wherein the glycogen phosphorylase inhibitor is INGLIFORIB or a pharmaceutically acceptable salt thereof.

39. Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a fructose-1,6-bisphosphate inhibitor or a pharmaceutically acceptable salt thereof.

30 **40.** Pharmaceutical composition, combination product or kit according to any of claims 7 and 39, wherein the fructose-1,6-bisphosphate inhibitor is selected from the group consisting of MB05032, MB-06322 and pharmaceutically acceptable salt of these compounds.

35 **41.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a cannabinoid (CB1) receptor antagonist or a pharmaceutically acceptable salt thereof.

42. Pharmaceutical composition, combination product or kit according to any of claims 7 and 41 wherein the cannabinoid (CB1) receptor antagonist is selected from the group consisting of AVE-1625, RIMONABANT, SURINABANT and pharmaceutically acceptable salts of these compounds.

5 **43.** Pharmaceutical composition, combination product or kit according to any of claims 1, 3 and 5 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an anti-obesity drug or a pharmaceutically acceptable salt thereof.

44. Pharmaceutical composition, combination product or kit according to any of claims 7 and 43
10 wherein the anti-obesity drug is selected from the group consisting of HMR-1426, CETILISTAT, SIBUTRAMINE and pharmaceutically acceptable salt of these compounds.

45. Pharmaceutical composition, combination product or kit according to any of claims 2, 4 and 6
15 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

46. Pharmaceutical composition, combination product or kit according to claim 45 wherein the first
20 other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is METFORMIN or a pharmaceutically acceptable salt thereof.

47. Pharmaceutical composition, combination product or kit according to any of claims 2, 4 and 6
25 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR agonist or a pharmaceutically acceptable salt thereof.

48. Pharmaceutical composition, combination product or kit according to claim 47 wherein the first
30 other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically acceptable salts of these compounds.

35 **49.** Pharmaceutical composition, combination product or kit according to any of claims 2, 4 and 6 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof.

50. Pharmaceutical composition, combination product or kit according to claim 49 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE GLIMEPIRIDE, GLICLAZIDE and the pharmaceutically acceptable salts of these compounds.

51. Pharmaceutical composition, combination product or kit according to any of claims 2, 4 and 6 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR agonist or a pharmaceutically acceptable salt thereof and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

52. Pharmaceutical composition, combination product or kit according to claim 51 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically acceptable salts of these compounds and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is METFORMIN or a pharmaceutically acceptable salt thereof.

53. Pharmaceutical composition, combination product or kit according to any of claims 2, 4 and 6 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

54. Pharmaceutical composition, combination product or kit according to claim 53 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE GLIMEPIRIDE, GLICLAZIDE and the pharmaceutically acceptable salts of these compounds and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is METFORMIN or a pharmaceutically acceptable salt thereof.

55. Pharmaceutical composition, combination product or kit according to any of claims 2, 4 and 6 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR agonist or a pharmaceutically acceptable salt thereof.

56. Pharmaceutical composition, combination product or kit according to claim 55 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE GLIMEPIRIDE, GLICLAZIDE and the pharmaceutically acceptable salts of these compounds and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically acceptable salts of these compounds.

57. Use of Compound A or a pharmaceutically acceptable salt thereof and one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes type 2 and/or type 1, for the manufacture of a medicament, in particular the pharmaceutical composition according to the invention, for the treatment of diabetes mellitus type 2 and/or type 1.

58. Use of Compound A or a pharmaceutically acceptable salt thereof and two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1, for the manufacture of a medicament, in particular the pharmaceutical composition according to the invention, for the treatment of diabetes mellitus type 2 and/or type 1.

59. Use of Compound A or a pharmaceutically acceptable salt thereof and one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1 for the manufacture of a sequentially or separately co-administrable medicament, in particular the combination product or kit according to the invention, for the treatment of diabetes type 2 and/or type 1.

60. Use of Compound A or a pharmaceutically acceptable salt thereof two other active compounds or pharmaceutically acceptable salts thereof which are used in the treatment of diabetes mellitus type 2 and/or type 1 for the manufacture of a sequentially or separately co-administrable medicament, in particular the combination product or kit according to the invention, for the treatment of diabetes type 2 and/or type 1.

61. A method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a pharmaceutical composition comprising a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first amount and the second amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

62. A method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a pharmaceutical composition comprising a pharmaceutical formulation including an amount of Compound A or a pharmaceutically acceptable salt thereof, amounts of two other active compounds or a pharmaceutically acceptable salts thereof which is are used in the treatment of diabetes mellitus type 2 and/or type 1, wherein the first, second third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1, and at least one pharmaceutically acceptable auxiliary.

63. A method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a combination product comprising the components:
(A) an amount of Compound A or a pharmaceutically acceptable salt thereof; and
(B) an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1,
wherein the first and second amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1;
wherein each of the components (A) and (B) is formulated in admixture with at least one pharmaceutically acceptable auxiliary;
and wherein the components (A) and (B) are administered sequentially or separately.

64. A method for treating diabetes mellitus type 2 and/or type 1 comprising administering to a patient in need thereof a combination product comprising the components:
(A) an amount of Compound A or a pharmaceutically acceptable salt thereof;
(B) an amount of one other active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1; and
(C) an amount of still another active compound or a pharmaceutically acceptable salt thereof which is used in the treatment of diabetes mellitus type 2 and/or type 1,
wherein the first, second and third amount together comprise an effective amount for the treatment of diabetes mellitus type 2 and/or type 1;
wherein each of the components (A), (B) and (C) is formulated in admixture with at least one pharmaceutically acceptable auxiliary;
and wherein the components (A), (B) and (C) are administered sequentially or separately.

65. Use or method according to any of claims 57 to 64 wherein the other active compound(s) which is (are) used in the treatment of diabetes mellitus type 2 and/or type 1 is (are independently) selected from the group consisting of insulin, insulin analogs, glucagon-like-peptide-1 receptor agonists, sulfonylurea agents, biguanide agents, alpha-glucosidase inhibitors, PPAR-agonists, meglitinide agents, dipeptidyl-peptidase IV inhibitors, PDE1 inhibitors, PDE5 inhibitors, PDE9 inhibitors, PDE10 inhibitors, PDE11 inhibitors, amylin analogues, cinnamon, glucagon receptor antagonists, cannabinoid (CB1)

receptor antagonists, anti-obesity drugs and the pharmaceutically acceptable salts of these compounds.

5 **66.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin.

67. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an insulin analogue.

10 **68.** Use or method according to any of claims 65 and 67 wherein the insulin analogue is selected from the group consisting of novarapid, insulin detemir, insulin lispro, insulin glargine, insulin zinc suspension and Lys-Pro insulin.

15 **69.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glucagon-like-peptide-1 receptor agonist or a pharmaceutically acceptable salt thereof.

20 **70.** Use or method according to any of claims 65 and 69 wherein the glucagon-like-peptide-1 receptor agonist is selected from the group consisting of BIM-51077, EXENATIDE, LIRAGLUTIDE, ALBIGLUTIDE, ZP-10 and the pharmaceutically acceptable salts of these compounds.

25 **71.** Use or method according to any of claims 65 and 69 wherein the glucagon-like-peptide-1 receptor agonist is selected from the group consisting of BIM-51077, EXENATIDE, LIRAGLUTIDE, ZP-10 and the pharmaceutically acceptable salts of these compounds.

72. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof.

30 **73.** Use or method according to any of claims 65 and 72 wherein the sulfonylurea agent is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE, GLIMEPIRIDE, GLICLAZIDE and the pharmaceutically acceptable salts of these compounds.

35 **74.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

40 **75.** Use or method according to any of claims 65 and 74 wherein the biguanide agent is METFORMIN or a pharmaceutically acceptable salt of this compound.

76. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a alpha-glucosidase inhibitor or a pharmaceutically acceptable salt thereof.

5

77. Use or method according to any of claims 65 and 76 wherein the alpha-glucosidase inhibitor is selected from the group consisting of ACARBOSE, MIGLITOL, VOGLIBOSE and the pharmaceutically acceptable salts of these compounds.

10

78. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR-agonist or a pharmaceutically acceptable salt thereof.

15

79. Use or method according to any of claims 65 and 78 wherein the PPAR-agonist is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, REGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halo-fenate and the pharmaceutically acceptable salts of these compounds.

20

80. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a meglitinide agent or a pharmaceutically acceptable salt thereof.

25

81. Use or method according to any of claims 65 and 80 wherein the meglitinide agent is selected from the group consisting of REPAGLINIDE, NATEGLINIDE, MITIGLINIDE and the pharmaceutically acceptable salts of these compounds.

30

82. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a dipeptidyl-peptidase IV (DPPIV) inhibitor or a pharmaceutically acceptable salt thereof.

35

83. Use or method according to any of claims 65 and 82 wherein the dipeptidyl-peptidase IV (DPPIV) inhibitor is selected from the group consisting of SITAGLIPTIN, SAXAGLIPTIN, VILDAGLIPTIN, DENAGLIPTIN, ALOGLIPTIN, P32/98 and the pharmaceutically acceptable salts of these compounds.

35

84. Use or method according to any of claims 65 and 82 wherein the dipeptidyl-peptidase IV (DPPIV) inhibitor is selected from the group consisting of SITAGLIPTIN, SAXAGLIPTIN, VILDAGLIPTIN, DENAGLIPTIN, P32/98 and the pharmaceutically acceptable salts of these compounds.

85. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PDE5 inhibitor or a pharmaceutically acceptable salt thereof.

5 **86.** Use or method according to any of claims 65 and 85 wherein the PDE5 inhibitor is selected from the group consisting of SILDENAFIL, VARDENFIL, TADALAFIL, UDENAFIL, AVANAFIL and the pharmaceutically acceptable salts of these compounds.

10 **87.** Use or method according to any of claims 65 and 85 wherein the PDE5 inhibitor is selected from the group consisting of SILDENAFIL, VARDENFIL, TADALAFIL and the pharmaceutically acceptable salts of these compounds.

15 **88.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used for the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of a PDE1, PDE9, PDE10, PDE11 inhibitor and the pharmaceutically acceptable salts thereof.

20 **89.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an amylin analogue or a pharmaceutically acceptable salt thereof.

90. Use or method according to any of claims 65 and 89 wherein the amylin analogue is PRAMLINTIDE or a pharmaceutically acceptable salt of this compound.

25 **91.** Use or method according to any of claims 57, 59, 61 and 63 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is cinnamon.

92. Use or method according to claim 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is cinnamon.

30

93. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glucagon receptor antagonist or a pharmaceutically acceptable salt thereof.

35 **94.** Use or method according to any of claims 65 and 93 wherein the glucagon receptor antagonist is BAY-27-9955 or a pharmaceutically acceptable salt of this compound.

40 **95.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a glycogen phosphorylase inhibitor or a pharmaceutically acceptable salt thereof.

96. Use or method according to any of claims 65 and 95 wherein the glycogen phosphorylase inhibitor is INGLIFORIB or a pharmaceutically acceptable salt of this compound.

5 **97.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a fructose-1,6-bisphosphate inhibitor or a pharmaceutically acceptable salt thereof.

10 **98.** Use or method according to any of claims 65 and 97 wherein the fructose-1,6-bisphosphate inhibitor is selected from the group consisting of MB-06322, MB-05032 and the pharmaceutically acceptable salts of these compounds.

15 **99.** Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a cannabinoid (CB1) receptor antagonist or a pharmaceutically acceptable salt thereof.

100. Use or method according to any of claims 65 and 99 wherein the cannabinoid (CB1) receptor antagonist is selected from the group consisting of AVE-1625, RIMONABANT, SURINABANT and the pharmaceutically acceptable salts of these compounds.

20

101. Use or method according to any of claims 57, 59, 61, 63 and 65 wherein the other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is an anti-obesity drug or a pharmaceutically acceptable salt thereof.

25 **102.** Use or method according to any of claims 65 and 101 wherein the anti-obesity drug is selected from the group consisting of HMR-1426, CETILISTAT, SIBUTRAMINE and the pharmaceutically acceptable salts of these compounds.

30 **103.** Use or method according to any of claims 58, 60, 62 and 64 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

35 **104.** Use or method according to claim 103 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is METFORMIN or a pharmaceutically acceptable salt thereof.

40 **105.** Use or method according to any of claims 58, 60, 62 and 64 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the sec-

ond other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR agonist or a pharmaceutically acceptable salt thereof.

106. Use or method according to claim 105 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically acceptable salts of these compounds.

107. Use or method according to any of claims 58, 60, 62 and 64 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof.

108. Use or method according to claim 107 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is insulin and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLIBORNURIDE, GLIBENCLAMIDE, GLIQUIDONE GLIMEPIRIDE, GLICLAZIDE and the pharmaceutically acceptable salts of these compounds.

109. Use or method according to any of claims 58, 60, 62 and 64 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR agonist or a pharmaceutically acceptable salt thereof and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.

110. Use or method according to claim 109 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR, NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically acceptable salts of these compounds and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is METFORMIN or a pharmaceutically acceptable salt thereof.

111. Use or method according to any of claims 58, 60, 62 and 64 wherein wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent or a pharmaceutically acceptable salt thereof and the second other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a biguanide agent or a pharmaceutically acceptable salt thereof.


112. Use or method according to claim 111 wherein the first other active compound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLI-
5 BORNURIDE, GLIBENCLAMIDE, GLIQUIDONE, GLIMEPIRIDE, GLICLAZIDE and the pharmaceuti-
cally acceptable salts of these compounds and the second other active compound which is used in the
treatment of diabetes mellitus type 2 and/or type 1 is METFORMIN or a pharmaceutically acceptable
salt thereof.

10 **113.** Use or method according to any of claims 58, 60, 62 and 64 wherein the first other active com-
pound which is used in the treatment of diabetes mellitus type 2 and/or type 1 is a sulfonylurea agent
or a pharmaceutically acceptable salt thereof and the second other active compound which is used in
the treatment of diabetes mellitus type 2 and/or type 1 is a PPAR agonist or a pharmaceutically ac-
ceptable salt thereof.

15 **114.** Use or method according to claim 113 wherein the first other active compound which is used in
the treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of
TOLBUTAMIDE, TOLAZAMIDE, GLIPIZIDE, CARBUTAMIDE, GLISOXEPIDE, GLISENTIDE, GLI-
BORNURIDE, GLIBENCLAMIDE, GLIQUIDONE GLIMEPIRIDE, GLICLAZIDE and the pharmaceuti-
20 cally acceptable salts of these compounds and the second other active compound which is used in the
treatment of diabetes mellitus type 2 and/or type 1 is selected from the group consisting of
MURAGLITAZAR, ROSIGLITAZONE, PIOGLITAZONE, FARGLITAZAR, NAVEGLITAZAR,
NETOGLITAZONE, RIVOGLITAZONE, K-111, GW-677954, (-)-Halofenate and the pharmaceutically
acceptable salts of these compounds.

25 **115.** Pharmaceutical composition, combination product or kit according to any one of claims 1 to 53
for the treatment of diabetes mellitus type 2 and/or type 1.

Fig. 1: Effect of RIMONABANT Hydrochloride, Compound A and a combination of RIMONABANT Hydrochloride and Compound A on HbA_{1c} in db/db-mice.



2 / 2

Fig. 2: Effect of RIMONABANT Hydrochloride, Compound A and a combination of RIMONABANT Hydrochloride and Compound A on glucose tolerance in db/db-mice

