



(86) Date de dépôt PCT/PCT Filing Date: 2011/07/05
(87) Date publication PCT/PCT Publication Date: 2012/01/12
(85) Entrée phase nationale/National Entry: 2013/01/04
(86) N° demande PCT/PCT Application No.: CN 2011/001104
(87) N° publication PCT/PCT Publication No.: 2012/003712
(30) Priorités/Priorities: 2010/07/06 (CN PCT/CN2010/001007);
2010/07/06 (CN PCT/CN2010/001008)

(51) Cl.Int./Int.Cl. *C07D 255/02* (2006.01),
C07D 403/06 (2006.01)

(71) Demandeur/Applicant:
CATEXEL LIMITED, GB

(72) Inventeurs/Inventors:
HAGE, RONALD, NL;
KOEK, JEAN HYPOLITES, NL;
RUSSELL, STEPHEN WILLIAM, NL;
WANG, XIAOHONG, CN;
WOLF, LODEWIJK VAN DER, NL;
ZHANG, JIANRONG, CN;
ZHAO, WEI, CN

(74) Agent: MARKS & CLERK

(54) Titre : 1,2-DI-(4,7-DIMETHYL-1,4,7-TRIAZACYCLONON-1-YL)-ETHANE ET INTERMEDIAIRE CORRESPONDANT
(54) Title: 1,2-BIS-(4,7-DIMETHYL-1,4,7-TRIAZACYCLONON-1-YL)-ETHANE AND INTERMEDIATE THEREOF

(57) **Abrégé/Abstract:**

The present invention provides improved processes for the synthesis of 1,4-ditosyl- 1,4,7-triazacyclonone and protected analogues thereof, and to synthesise bis(triazacyclononane) ligands.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
12 January 2012 (12.01.2012)



(10) International Publication Number
WO 2012/003712 A1

(51) International Patent Classification:

C07D 255/02 (2006.01) **C07D 403/06** (2006.01)

(21) International Application Number:

PCT/CN2011/001104

(22) International Filing Date:

5 July 2011 (05.07.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/CN2010/001008 6 July 2010 (06.07.2010) CN

PCT/CN2010/001007 6 July 2010 (06.07.2010) CN

(71) **Applicant** (for AE, AG, AU, BB, BH, BW, BZ, CA, CY, EG, GB, GD, GH, GM, IE, IL, KE, KN, LC, LK, LS, MT, MW, MY, NA, NG, NZ, OM, PG, SC, SD, SG, SL, SZ, TT, TZ, UG, VC, ZA, ZM, ZW only): **UNILEVER PLC** [GB/GB]; Unilever House, 100 Victoria Embankment, London EC4Y 0DY (GB).

(71) **Applicant** (for all designated States except AE, AG, AU, BB, BE, BF, BH, BJ, BW, BZ, CA, CF, CG, CI, CM, CY, EG, FR, GA, GB, GD, GH, GM, GN, GQ, GR, GW, IE, IL, IN, IT, KE, KN, LC, LK, LS, LV, MC, ML, MR, MT, MW, MY, NA, NE, NG, NL, NZ, OM, PG, SC, SD, SG, SI, SL, SN, SZ, TD, TG, TT, TZ, UG, US, VC, ZA, ZM, ZW): **UNILEVER N.V.** [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

(71) **Applicant** (for IN only): **HINDUSTAN LEVER LIMITED** [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Maharashtra, Mumbai 400 020 (IN).

(72) Inventors; and

(75) **Inventors/Applicants** (for US only): **HAGE, Ronald** [NL/NL]; Rahu Catalytics B.V., BioPartner Center Leiden, Wassenaarseweg 72, NL-2333 AL Leiden (NL). **KOEK, Jean Hypolites** [NL/NL]; Unilever R & D Vlaardingen B.V., Olivier van Noortlaan 120, 3133 AT Vlaardingen (NL). **RUSSELL, Stephen William**

[NL/NL]; Singel 2, 3161 BK Rhon (NL). **WANG, Xiaohong** [CN/CN]; Unilever (China) Investing Company, 66 Linxin Road, Linkong Economic Development Zone, Shanghai 200335 (CN). **WOLF, Lodewijk van der** [NL/NL]; Amalia van Solmslaan 83, 3137 DE Vlaardingen (NL). **ZHANG, Jianrong** [CN/CN]; Unilever (China) Investing Company, 66 Linxin Road, Linkong Economic Development Zone, Shanghai 200335 (CN). **ZHAO, Wei** [CN/CN]; Unilever (China) Investing Company, 66 Linxin Road, Linkong Economic Development Zone, Shanghai 200335 (CN).

(74) **Agent:** **CHINA PATENT AGENT (H. K.) LTD.**; 22/F, Great Eagle Centre, 23 Harbour Road, Wanchai, Hong Kong Special Administrative Region (CN).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** 1,2-BIS-(4,7-DIMETHYL-1,4,7-TRIAZACYCLONON-1-YL)-ETHANE AND INTERMEDIATE THEREOF

(57) **Abstract:** The present invention provides improved processes for the synthesis of 1,4-ditosyl- 1,4,7-triazacyclonone and protected analogues thereof, and to synthesise bis(triazacyclononane) ligands.

WO 2012/003712 A1

- 1 -

1,2-BIS-(4,7-DIMETHYL-1,4,7-TRIAZACYCLONON-1-YL)-ETHANE
AND INTERMEDIATE THEREOF

FIELD

5 The present invention concerns the synthesis of an intermediate useful for the synthesis of 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane (Me₄-DTNE); and the synthesis of binucleating macrocyclic ligands that may be used to form complexes that have utility as bleach and/or oxidation catalysts.

10 BACKGROUND

Manganese complexes containing the ligands Me₃-TACN (1,4,7-trimethyl-1,4,7-triazacyclononane) and Me₄-DTNE (1,2-bis-(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)-ethane) are of interest for different bleaching of cellulosic and other substrates.

15 Different methods have been disclosed to synthesise 1,4-ditosyl-1,4,7-triazacyclononane (Ts₂-TACN) from 1,4,7-tritosyl-1,4,7-triazacyclononane (Ts₃-TACN) as described below.

20 Ts₃-TACN has been treated with a mixture of bromic acid and acetic acid for 20 h at 100 °C and subsequently refluxed for 30 h to yield fully detosylated 1,4,7-triazacyclononane (H₃-TACN) as HBr salt, i.e. H₃-TACN.HBr; subsequent reaction with 2 equivalents of tosyl chloride afforded Ts₂-TACN in 60 % yield as disclosed in *Inorg. Chem.*, 1985, 24, 1230.

25 Ts₃-TACN has been treated with a mixture of bromic acid, acetic acid and phenol for 36 h at 90 °C, to furnish monotosylated Ts-TACN. Further reaction with 1 equivalent of tosyl chloride to afford Ts₂-TACN in a higher yield than using method 1 - 76% as disclosed in *Inorg. Chem.*, 1990, 29, 4143.

 Ts₃-TACN has been heated with a mixture of hydrobromic acid and acetic acid under reflux for 3 h to yield a mixture of Ts-TACN.HBr (68%) and Ts₂-TACN.HBr(30%) as disclosed in *Synthetic Communications*, 2001, 31(20), 3141.

30 Isolation of protonated Ts₂-TACN salt with bromide as counter ion has been described in *Synthetic Communications* 31(20), 3141-3144, 2001; and US 2005/112066 A1.

 The reaction of Ts₂-TACN with 2 equivalents of ditosyl-ethyleneglycol in DMF to yield 1,2-bis(4,7-ditosyl-1,4,7-triazacyclonon-1-yl)-ethane (Ts₄-DTNE) is also disclosed in *Inorg. Chem.* 1985, 24, 1230; *Inorg. Chem.* 1996, 35, 1974-1979; *Inorg. Chem.* 1998, 37(5), 3705-3713; *Inorg. Chem.* 2005, 44 (2), 401-409; and *J. Chem. Soc., Dalton Trans.* 1994, 457-464.

35

- 2 -

Ts₄-DTNE has also been obtained using O,O',N,N'-tetratosyl-N,N'-bis(2-hydroxyethyl)ethylenediamine and ethylenediamine (*Synthesis* 2001, 2381-2383; *Inorg. Chem.* 2007, 46(1), 238-250; *Green Chem.* 2007, 9, 996-1007).

Synthesis of 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane (DTNE) from methine-
 5 1,4,7-triazacyclononane and dibromoethane or diiodoethane has been disclosed in *J. Chem. Soc., Chem. Commun.*, 1987, 886; *J. Am. Chem. Soc.*, 1998, 120, 13104-13120; *Inorg. Chem.* 1993, 32, 4300-4305; *Inorg. Chem.* 1997, 36, 3125-3132; *Chem. Lett.* 2000, 416-417; *J. Chem. Soc., Dalton Trans.*, 2000, 3034-3040.

Synthesis of Me₄-DTNE from DTNE using formaldehyde and formic acid can be
 10 found in *J. Am. Chem. Soc.*, 1998, 120, 13104-13120; *Inorg. Chem.* 1993, 32(20), 4300-4305; *Chem. Lett.*, 2000, 416-7.


Using the known methods, the binucleating triazacyclononane ligand can be obtained in a reasonable yield. However, as the purity level is insufficient to obtain the dinuclear manganese complex ([Mn^{IV}Mn^{III}(μ-O)₂(μ-OAc)(Me₄-DTNE)]²⁺) in high yield, an
 15 additional purification step, such as vacuum distillation is needed. Although this gives then a high purity material, the yield loss is quite substantial. Therefore there is still a need to be able to synthesise manganese complexes using Me₄-DTNE that has been obtained in a more simple synthetic procedure, with preferably without needing to distil Me₄-DTNE prior complexation.

20

SUMMARY

We have found that partial detosylation of Ts₃-TACN in a one-pot process leads to the formation of Ts₂-TACN as its protonated salt. This is an improvement over the two-step process of complete detosylation of Ts₃-TACN followed by ditosylation of the
 25 TACN adduct. Furthermore, less tosyl chloride can be used to make Ts₂-TACN and less tosylate waste compared to the above route has been obtained.


In a first aspect the present invention provides a method of producing a compound of formula (A):



30 the method comprising the following step:


(a) reacting a compound of formula (B):

- 3 -




in an acidic medium comprising sulfuric acid, the molar ratio of B to sulfuric acid in the range from 1:0.1 to 1:10, preferably 1:0.5 to 1:10, more preferably 1:0.5 to 1:5, even more preferably from 1:1 to 1:4, wherein P is an arylsulfonate protecting group and the compound of formula (A) is isolated as a protonated salt in amorphous or crystalline form.

In a second aspect the present invention provides a method of producing a compound of formula (A):



the method comprising the following step:

(a) reacting a compound of formula (B):



in an acidic medium, wherein P is an arylsulfonate protecting group, wherein the acidic medium is worked-up when the conversion of B to A is at least 50 mol % yielding compound (A).


As disclosed in the background of invention the Ts₂-TACN may be used to form Ts₄-DTNE which can be detosylated and secondary amines of the product methylated in a similar fashion described in US 5,284,944 for Ts₃-TACN. In a similar manner the same applies to the arylsulfonates as a class of protecting groups. Such reactions relate to further aspects of the invention.

Reaction of 1,4-di(arylsulfonate)-1,4,7-triazacyclonane ((ArSO₂)₂-TACN) with dihaloethane in a solvent, optionally, in the presence of water, and a base yields 1,2-


- 4 -

bis-(4,7-diarylsulfonate-1,4,7-triazacyclonon-1-yl)-ethane in high yield. Removing the arylsulfonate protecting groups and then reacting further with formaldehyde and formic acid in one-pot reaction yields Me₄-DTNE. Surprisingly, when acetonitrile/water is employed as a solvent in the step to form Ts₄-DTNE, the purity level of Me₄-DTNE is high enough to allow complexation to form the manganese complex, ([Mn^{IV}Mn^{III}(μ-O)₂(μ-OAc)(Me₄-DTNE)]²⁺), without the need to distil the Me₄-DTNE ligand prior the complexation step.

In a third aspect the present invention provides a method of producing a compound of formula (C), the method comprising the following step:



(a) reacting a compound of formula (A):




with a bridging element of the form ZCH₂CH₂Z, wherein P is an arylsulfonate protecting group and Z is a halogen selected from: Cl; Br; and, I.

In chemistry one-pot synthesis/reaction is a strategy to improve the efficiency of a chemical reaction whereby a reactant is subjected to successive chemical reactions in just one reactor. This is much desired by chemists because avoiding a lengthy separation process and purification of the intermediate chemical compound would save time and resources while increasing chemical yield.

In a fourth aspect the present invention provides a one-pot method for the preparation of Me₄-DTNE, the method comprising deprotecting a compound of formula (C):

- 5 -



with an acidic medium and to form DTNE and subsequently adding formaldehyde and formic acid to the reaction medium, wherein P is an arylsulfonate.


5

DETAILED DESCRIPTION

The starting material 1,4,7-tri(arylsulfonate)-1,4,7-triazacyclonane ((ArSO₂)₃-TACN) is reacted in acid to yield ((ArSO₂)₂-TACN).

A preferred synthetic scheme for obtaining an (ArSO₂)₂-TACN (Ts₂-TACN) is outlined below.

10



The preferred temperature range for monodearylsulfonation of the triarylsulfonate is from 100 to 160 °C, with most preferred from 130 and 150 °C.

15

The preferred time for the method is from 1 h to 24 h, the most preferred time from 2 to 6 h.

Preferably the method is conducted as a one-pot reaction.

The preferred acid for monodearylsulfonation of the tri arylsulfonate is sulfuric acid. Other acids, such as methanesulfonic acid and sulfonic acid resins may function to provide the monodetosylation. Preferably, the acidic medium does not contain any hydrogen halides and in this regard, the acidic medium preferably has less than 1 mol% hydrogen halides with respect to B. We have surprisingly found that use of such acidic media provides advantages in relation to the use of hydrogen halides. In particular, whereas the use of a mixture of acidic acid and hydrobromic acid has been reported to provide a mixture of mono- and ditosylated (predominantly monotosylated) TACN from Ts₃-TACN (*Synthetic Communications*, 2001, 31(20), 3141), the present

20

25

- 6 -

invention advantageously, and surprisingly, permits the provision of a significantly higher proportion of the desired ditosylated (monodetosylated) product.

Additionally auxiliary anhydrides are preferably present, such as acetic acid anhydride or propionic acid anhydride when excess water is present in the reaction mixture. The amount of acid anhydride required to facilitate the reaction depends upon the amount of water initially present in the reaction.

The use of an auxiliary anhydride in a method of producing a compound of formula (A) as hereinbefore defined represents a fifth aspect of the present invention. Viewed from the suspect, the invention provides a method of producing a compound of formula (A), as hereinbefore defined, comprising reacting a compound of formula (B) in an acidic medium comprising an acid anhydride. The acidic medium may be as described herein and the compound of formula (A) is typically isolated as a protonated salt, such as an aryl sulfonic acid salt (e.g. the toluene sulfonic acid or benzene sulfonic acid salt), for example in amorphous or crystalline form.

The acid anhydride serve to maintain the molar ratio of $(\text{ArSO}_2)_3\text{-TACN}$:water at a level that aids the ideal molar ratio for the reaction, namely 1:1.

The optimum amount of acid anhydride to be added to the reaction mixture is dependent on the amount of $(\text{ArSO}_2)_3\text{-TACN}$ and the amount of water in the system (originating from the water present in $(\text{ArSO}_2)_3\text{-TACN}$ and sulfuric acid added). If the molar amount of water present in $(\text{ArSO}_2)_3\text{-TACN}$ and sulfuric acid is much larger than the molar amount of $(\text{ArSO}_2)_3\text{-TACN}$, the reaction may become less efficient, i.e. more mono(arylsulfonate)TACN or $\text{H}_3\text{-TACN}$ will be formed. It should be noted that one mol of acid anhydride will react with one mol of water to form two moles of acid.

Therefore, the following relation exists (all on molar basis):

$\text{H}_2\text{O}(\text{Ts}_3\text{-TACN}) + \text{H}_2\text{O}(\text{sulfuric acid}) - \text{acid anhydride} =$
amount of water available to react with $\text{Ts}_3\text{-TACN}$.

Therefore:

$\text{H}_2\text{O}(\text{Ts}_3\text{-TACN}) + \text{H}_2\text{O}(\text{sulfuric acid}) - \text{Ts}_3\text{-TACN} = \text{acid anhydride},$
which is equal to:

$[\text{H}_2\text{O}(\text{Ts}_3\text{-TACN}) + \text{H}_2\text{O}(\text{sulfuric acid}) - \text{Ts}_3\text{-TACN}] : \text{acid anhydride} = 1.$

Allowing variables in process conditions, this ratio should be varying between 0.1 and 10, more preferably between 0.3 and 5 and most preferably between 0.8 and 2.

It is preferred that a tosyl group is used as protecting group for the secondary amines of the TACN moiety. The tosyl group (abbreviated Ts or Tos) is $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_2$. This group is usually derived from the compound 4-toluene sulfonyl chloride,

- 7 -

CH₃C₆H₄SO₂Cl, which forms esters and amides of toluene sulfonic acid. The para orientation illustrated (p-toluenesulfonyl) is most common, and by convention tosyl refers to the p-toluenesulfonyl group. Tosylate refers to the anion of p-toluenesulfonic acid (CH₃C₆H₄SO₃⁻). Whilst the tosyl group is the preferred protecting group, other
5 arylsulfonyl groups (ArSO₂) will function to provide the advantages of the present invention. Preferably the arylsulfonyl employed is a benzenesulfonate. The skilled person will understand that, where compounds of formula (A) are prepared from compounds of formula (B) and isolated as a protonated salt, the protonated salt will typically be of the same arylsulfonic acid (e.g. p-toluene sulfonic acid) of which
10 protecting group P in compounds of formulae (A) and (B) is the aryl sulfonate.

Compared to the known procedures to make Ts₂-TACN, as outlined in the background of the invention, there will be one-step less needed to obtain this material in a high yield and purity. Furthermore, less tosylchloride (arylsulfonate) starting materials are needed to form (ArSO₂)₂-TACN (3 instead of 5 molar equivalents) and as
15 a consequence also less tosylate (arylsulfonate) waste will be generated.

In a sixth aspect of the invention the (ArSO₂)₂-TACN can be obtained and isolated as a protonated (HX) salt in which HX is selected from: toluenesulfonic acid; benzenesulfonic acid; sulfuric acid; acetic acid; formic acid; and, propionic acid, most preferably from toluenesulfonic acid, benzenesulfonic acid and sulfuric acid. One
20 skilled in the art will appreciate that some acids will support more than one protonated (ArSO₂)₂-TACN, for example sulfuric acid. Alternatively, sulfuric acid may support one protonated (ArSO₂)₂-TACN, as the HSO₄⁻ counterion. According to particular embodiments of this and other aspects of the invention, the protonated salt of the (ArSO₂)₂-TACN is Ts₂-TACN.TsOH (wherein TsOH is toluene sulfonic acid), or the
25 benzene sulfonic acid salt of 1,4-di(benzenesulfonate)-1,4,7-triazacyclonane.

From the disclosure it will be evident that conditions and some reagents may be varied to provide the desired (ArSO₂)₂-TACN. With this in mind, one skilled in the art can monitor the progress of the reaction, for example by thin layer chromatography, and determine the extent to which (ArSO₂)₂-TACN. When the conversion of B
30 ((ArSO₂)₃-TACN) to A ((ArSO₂)₂-TACN) is at least 50 mol % yielding compound (A) the reaction is worked-up. Preferably, the reaction is worked up when the conversion of B to A is at least is at least 50 mol % yielding compound (A).

The term worked-up is known in the art. In chemistry work-up refers to the series of manipulations required to isolate and purify the product(s) of a chemical
35 reaction. Typically, these manipulations include:


- quenching a reaction to deactivate any unreacted reagents

- 8 -

- changing the pH to prevent further reaction
- cooling the reaction mixture or adding an antisolvent to induce precipitation, and collecting or removing the solids by filtration, decantation, or centrifuging
- removal of solvents by evaporation
- separating the reaction mixture into organic and aqueous layers by liquid-liquid extraction
- purification by chromatography, distillation or recrystallisation.

A method for obtaining Me₄-DTNE is also provided.

A preferred synthetic scheme for obtaining Me₄-DTNE is outlined below.



The (ArSO₂)₂-TACN (e.g. Ts₂-TACN) in particular embodiments of this invention is prepared according to the first or second aspects of the invention and/or be a protonated salt in accordance with the sixth aspect of the invention, for example Ts₂-TACN.TsOH. In the discussion of the invention herein, focus is primarily upon embodiments of the present invention for obtaining Me₄-DTNE using Ts₂-TACN.TsOH. However, the invention is not limited to these embodiments, since the skilled person is aware of other ways of making (ArSO₂)₂-TACN, and protonated salts thereof, including Ts₂-TACN and protonated salts thereof, for example in accordance with the documents referred to in the Background section.

(ArSO₂)₂-TACN reacts with 1,2-dihaloethane in a solvent and a base, wherein the water level in the solvent is between 0 and 90%. The 1,2-dihaloethane is preferably selected from 1,2-dibromoethane, 1,2-diiodoethane and 1,2-dichloroethane, with 1,2-dibromoethane being most preferred. Different solvents can be employed, such as acetonitrile, dimethylformamide (DMF), xylene, toluene, dioxane, 1-butanol, 2-butanol, t-butanol, 1-propanol, and 2-propanol. The solvent may contain additional water. The water content of the solvent may be between 0 and 90%.

The base used for the coupling of (ArSO₂)₂-TACN with dihaloethane should not be too strong; the base used for the coupling reaction is preferably sodium carbonate.

It is preferred that a tosyl group is used as protecting group for the secondary amines of the TACN moiety.

- 9 -

Preferred solvents are acetonitrile, 1-butanol, 2-butanol, t-butanol, and dimethylformamide (DMF). These solvent are preferably used with additional water, preferably between 10 and 90%. Most preferably, acetonitrile/H₂O is used, as 1,2-bis(4,7-arylsulfonate-1,4,7-triazacyclonon-1-yl)-ethane obtained is of higher purity than
5 using other solvents. This allows the formation of the Me₄-DTNE ligand of higher purity and therefore the ligand does not need to be distilled prior using for the complexation step with manganese.

The protecting groups of 1,2-bis(4,7-arylsulfonate-1,4,7-triazacyclonon-1-yl)-ethane are removed by treatment with an acid to yield DTNE. The preferred acid used
10 for deprotection is concentrated sulfuric acid. After deprotection the solution containing the deprotected ligand is neutralised to pH 5 to 9, preferably pH 6 to 8.

The DTNE is preferably methylated by reaction with formaldehyde and subsequent reduction. In this regard, reaction with formaldehyde and formic acid (Eschweiler-Clarke methylation) are the preferred reagents to effect methylation. This
15 reductive amination step will not produce quaternary ammonium salts, but instead will stop at the tertiary amine stage. For the aforementioned reason the Eschweiler-Clarke methylation is preferred over other methylation procedures.

Whilst the Eschweiler-Clarke methylation step is preferred other methylation reactions may be used. Methylation of secondary amines is well known in the art.
20 Some examples of references are Ber. 1905, 38, 880; J. Am. Chem. Soc., 1933, 55, 4571; J. Org. Chem. 1972, 37(10), 1673-1674; J. Chem. Soc., Perkin Trans 1, 1994, (1), 1-2; Synth. Commun., 2002, 32(3), 457-465; Synth. Commun., 1989, 19(20), 3561-3571; Synth. Commun., 2006, 36(23), 3609-3615; EP0553954A2; US5105013; J. of the Indian Chemical Society 1967, 44(5), 430-435; J. of the Indian Chemical Society
25 1970, 8(8), 725-727.

Reductive methylation in general applying formaldehyde and a reducing agent like cyanoborohydride, formic acid, molecular hydrogen and a catalyst (Nickel, Palladium on coal, etc.) can be employed. Also direct methylation with methyl-X (X = Cl, Br, I).

30 Catalytic conversions for preparing tertiary amines from secondary and primary amines using hydrogen gas and formaldehyde can be for example found in US 4,757,144.


After the methylation is reaction is complete, increasing the pH to preferably higher than 12, more preferably higher than 13, the Me₄-DTNE ligand can be extracted
35 using a C5-C8 hydrocarbon as solvent. The C5-C8 is preferably selected from pentane, hexane, heptane, octane, cyclopentane, cyclohexane, cycloheptane, cyclooctane,

- 10 -

toluene, xylene and combinations thereof. Most preferred solvents are hexane or heptane. When not using acetonitrile to synthesise 1,2-bis-(4,7-arylsulfonate-1,4,7-triazacyclonon-1-yl)-ethane, the ligand obtained is best vacuum distilled before further complexing with manganese salts. Alternatively, the ligand may be purified by precipitating as HCl salt, after which the free Me₄-DTNE ligand was obtained by addition of concentrated NaOH solution, as exemplified in J.Am.Chem.Soc. 1998, 120, 13104-13120.


The invention may be further understood with respect to the following non-limiting clauses:

1. A method of producing a compound of formula (A):




the method comprising the following step:

- (a) reacting a compound of formula (B):



in an acidic medium comprising sulfuric acid, the molar ratio of B to sulfuric acid in the range from 1:0.5 to 1:10, wherein P is an arylsulfonate protecting group and the compound of formula (A) is isolated as a protonated salt in amorphous or crystalline form.


2. A method of producing a compound of formula (A):



- 11 -

the method comprising the following step:


(a) reacting a compound of formula (B):




in an acidic medium, wherein P is an arylsulfonate protecting group, wherein the acidic medium is worked-up when the conversion of B to A is at least 50 mol % yielding compound (A).

3. The method of clause 1 or clause 2, wherein an acid anhydride is present in the acidic medium.

4. A method of producing a compound of formula (A):



the method comprising reacting a compound of formula (B):




in an acidic medium comprising an acid anhydride, wherein P is an arylsulfonate protecting group.

5. The method of any one of clauses 2 to 4, wherein the compound of formula (A) is isolated as a protonated salt of the same arylsulfonic acid of which P is the arylsulfonate.

- 12 -

6. The method of clause 5, wherein the protonated salt of the compound of formula (A) is isolated in amorphous or crystalline form.
7. The method of any one of clauses 2 to 6, wherein the acidic medium comprises sulfuric acid.
8. The method of any one of clauses 1 or 3 to 7, wherein the protonated salt is a salt of an aryl sulfonic acid.
9. The method of clause 8, wherein the protonated salt is:
- (1,4-ditosyl-1,4,7-triazacyclonone) tosylate (which is the toluene sulfonic acid salt of 1,4-ditosyl-1,4,7-triazacyclononane); or
 - (1,4-dibenzenesulfonyl-1,4,7-triazacyclonone) benzenesulfonate (which is the benzene sulfonic acid salt of 1,4-dibenzene sulfonyl-1,4,7-triazacyclononane).
10. The method of any one preceding clause, wherein the method is conducted at a reaction temperature from 100 to 160 °C.
11. The method of any one preceding clause, further comprising reacting a compound of formula (A) with a bridging element of the form ZCH_2CH_2Z , wherein Z is a halogen selected from: Cl; Br; and, I, whereby to produce a compound of formula (C):



(C),


- wherein groups P in the compound of formula (C) are the same as groups P in the compound of formula (A).
12. The method of clause 11, wherein the compound of formula (A) is reacted with the bridging element in acetonitrile, for example aqueous acetonitrile, such as

- 13 -

aqueous acetonitrile comprising from 10 to 90 wt/wt % water, from 10 to 50 wt/wt % water or from 10 to 35 wt/wt % water.

13. The method of clause 11 or clause 12 further comprising, optionally in a one-pot method, deprotecting the compound of formula (C) with an acidic medium to form 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane and subsequently adding formaldehyde and formic acid to the reaction medium whereby to produce 1,2-bis-(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)-ethane.

14. A protonated salt of formula (A), having a counter ion HX, the protonated salt in amorphous or crystalline form:



wherein P is a tosylate or benzene sulfonate and HX is selected from: toluenesulfonic acid; benzenesulfonic acid; sulfuric acid; acetic acid; formic acid; and, propionic acid.


15. The protonated salt of clause 14, which is (1,4-ditosyl-1,4,7-triazacyclonone) tosylate, (1,4-dibenzenesulfonyl-1,4,7-triazacyclonone) benzenesulfonate or (1,4-ditosyl-1,4,7-triazacyclonone) benzenesulfonate.

16. The protonated salt of clause 14, which is (1,4-ditosyl-1,4,7-triazacyclonone) tosylate.

17. The protonated salt of clause 14, which is (1,4-dibenzenesulfonyl-1,4,7-triazacyclonone) benzenesulfonate.


- 14 -

18. A method of producing a compound of formula (C):



the method comprising the following step:


- (a) reacting a compound of formula (A):



with a bridging element of the form ZCH_2CH_2Z , wherein P is an arylsulfonate protecting group and Z is a halogen selected from: Cl; Br; and, I.

19. The method of clause 18, wherein the compound of formula (A) is reacted with the bridging element in a solvent selected from: acetonitrile; 1-butanol; 2-butanol; and, t-butanol.
20. The method of clause 19, wherein the compound of formula (A) is reacted with the bridging element in acetonitrile as solvent, for example with aqueous acetonitrile as solvent.
21. The method of clause 20, wherein the solvent comprises from 10 to 90 wt/wt % water, from 10 to 50 wt/wt % water, or from 10 to 35 wt/wt % water.
22. The method of any one of clauses 18 to 21, wherein the method further comprises, optionally in a one-pot method, deprotecting the compound of formula (C) with an acidic medium to form 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane and subsequently adding formaldehyde and formic acid to the reaction medium whereby to produce 1,2-bis-(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)-ethane.

23. A one-pot method for the preparation of 1,2-bis-(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)-ethane (Me₄-DTNE) the method comprising deprotecting a compound of formula (C):



with an acidic medium and to form 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane and subsequently adding formaldehyde and formic acid to the reaction medium, wherein P is an arylsulfonate.

24. The one-pot method of clause 23, wherein P is tosylate.

The following examples illustrate the invention more fully in which the amounts and ratios as given herein apply to the start of the method and will change during the reaction; and Ts₂-TACN·TsOH used in Examples 2, 3a & 3b, 5a-5c and 7-10 was prepared according to Example 1.

EXPERIMENTAL

1 Preparation of Ts₂-TACN·TsOH

Ts₃-TACN was synthesised as disclosed in WO9400439. Ts₃-TACN (128.3 g, 96.6% containing 3.4% water, 209.5 mmol of Ts₃-TACN, 242 mmol H₂O) and propionic acid (113 mL) were placed in a 500mL three-necked-flask with thermometer and condenser. While stirring magnetically and warming (bath 160-170 °C) most of the Ts₃-TACN dissolved. Propionic anhydride (12 g, 92 mmol) and sulfuric acid (29.5 mL, 96%, 530 mmol, containing 120 mmol H₂O) were then added. (Caution: at the beginning period of adding H₂SO₄, exothermic reaction occurred violently). Stirring was continued (reaction mixture = 142-143 °C) until the TLC showed the conversion to be complete (about 3hrs). After partial cooling, the warm (70 ~ 80°C) contents of the flask were poured into 1.5 L ice-water while stirring vigorously. The product was left at room temperature overnight, then filtered over a large frit (φ10cm) and washed with water

- 16 -

(6×300 mL) until pH=7, the obtained white solid was dried under vacuum at 60°C with P₂O₅ until the weight is constant (at least 2 days). Yield of Ts₂-TACN.TsOH: 93 g (74%) with purity: 91.5%. The filtrate was neutralized with aqueous NaOH to pH14, white solid which proved to be Ts₂-TACN appeared, filtered and washed with water, dried under vacuum to a constant weight. Another 4% product could be obtained with 90% purity. The total yield is about 78%.

¹H NMR (400 MHz, CDCl₃): δ 2.36 (s, [ArCH₃ (TsOH), 3H]), 2.44 (s, (ArCH₃ (N-Ts), 6H), 3.41 (br.s, [N-CH₂, 4H]), 3.54 (br. S, [N-CH₂, 4H]), 3.75 (br.s, [N-CH₂, 4H]), 7.20 (d, J = 7.4 Hz, [ArH, 2H]), 7.32 (d, J = 7.4 Hz, [ArH, 4H]), 7.66 (d, J = 7.4 Hz, [ArH, 4H]), 7.90 (d, J = 7.4 Hz, [ArH, 2H]).

ESI-MS(ES⁺): m/z 438 (Ts₂-TACN + H)⁺

2 Preparation of Ts₄-DTNE using acetonitrile as aprotic solvent

(Ts₄-DTNE = 1,2-bis(4,7-ditosyl-1,4,7-triazacyclonon-1-yl)-ethane)

The mixture of the protonated tosylate salt of 1,4-ditosyl-1,4,7-triazacyclonone (Ts₂-TACN.TsOH -3.0 g, 5 mmol) and Na₂CO₃ (2.12 g, 20 mmol) in 20 mL acetonitrile was stirred under reflux for 5 min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed overnight (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). Then the solvent was evaporated and to the residue 50 mL water was added and the resulting mixture was filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product 1.84 g (84%) with 84% purity.

¹H NMR (400 MHz, CDCl₃):

1.42 (s, [ArCH₃, 12H]), 2.73 (s, [bridging N-CH₂, 4H]), 2.93

(s, 8H), 3.19 (s, 8H), 3.46 (s, 8H), 7.30 (d, J = 7.4 Hz, 8H), 7.65 (d, J = 7.4 Hz, 8H).

ESI-MS (ES⁺): m/z 901(M + H)⁺

3a Preparation of Ts₄-DTNE in acetonitrile/water

The mixture of Ts₂-TACN.TsOH (3.0 g, 5 mmol) in 25 mL acetonitrile and Na₂CO₃ (2.12 g, 20 mmol) in 10 mL water was stirred at 100 °C for 5 min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed overnight (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). After being cooled to room temperature, the mixture was poured into 50 mL water and was

filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product 1.6 g (72%) with 93.3% purity.

3b Preparation of Ts₄-DTNE in acetonitrile/water (larger scale)

5 The mixture of Ts₂-TACN·TsOH (60 g, 100 mmol) in 500 mL acetonitrile and Na₂CO₃ (42.5 g, 400 mmol) in 200 mL water was stirred at 100 °C for 5 min. Then 1,2-dibromoethane (8.75 mL, 100 mmol) was added and the resulting mixture was refluxed overnight (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). After being cooled to room temperature, the mixture was poured into ca. 1000 mL water and
10 was filtered. The solid was washed with water (4×1000 mL), dried under vacuum to afford the product 33 g (74.6%) with 93.3% purity.

4a Preparation of Me₄-DTNE using Ts₄-DTNE prepared in acetonitrile/H₂O

15 Ts₄-DTNE (93.3% purity) (25 g, 26 mmol) and 96% sulfuric acid (59.2 mL, composed of 56.8 mL concentrated H₂SO₄ (98%) plus 2.4 mL water) were stirred at 110 °C (oil bath) in a 1 L 3-necked flask overnight. The reactants were cooled to 50 °C, then water (71 mL) and NaOH solution (108 g NaOH in 200 mL water) was added dropwise under ice-bath with stirring until pH = 6~7, then formaldehyde (25.3 g (37%)) and formic acid (99%) (28.7 g) were added successively with stirring, the mixture was
20 stirred at 90~100 °C (110 °C oil bath) overnight, then cooled to room temperature, the contents were made strongly alkaline by adding NaOH (32 g in 60 mL water) until pH 14 while maintaining the temperature at 30 °C, the brown slurry was stirred efficiently with hexane (200 mL) then filtered over celite. After separating the phase, the filter cake was washed with hexane (4×200 mL) which was subsequently used to extract the
25 aqueous, then the aqueous was extracted with hexane (4×500 mL), the combined hexane layer was evaporated to get the crude product 7.4 g (84%) as yellow oil with purity 85%. Similar results were obtained when heptane was used as the extraction solvent.

¹H NMR (400 MHz, CDCl₃): 2.3 (s, 12H, -CH₃), 2.6 (m, 28H, -N-CH₂). ESI-MS (ES⁺):
30 m/z 341 (M + H)⁺.

4b Preparation of Me₄-DTNE using Ts₄-DTNE prepared in acetonitrile/H₂O

35 Ts₄-DTNE obtained in step 2 (93.3% purity) (21.5 g, 22.4 mmol) and 96% sulfuric acid (54.7 mL, composed of 52.5 mL concentrated H₂SO₄ (98%) plus 2.2 mL water) were stirred at 110 °C (oil bath) in a 1 L 3-necked flask overnight. The reactants were cooled to 50 °C, then water (59 mL) and NaOH solution (90 g NaOH in 150 mL

- 18 -

water) was added dropwise under ice-bath with stirring until pH = 6~7, then formaldehyde (27 mL(37%)) and formic acid (99%) (20 mL) were added successively with stirring, the mixture was stirred at 90~100 °C (110 °C oil bath) overnight, then cooled to room temperature, 59 mL water was added and the contents were made
5 strongly alkaline by adding NaOH (27 g in 50 mL water) until pH 14 while maintaining the temperature at 30 °C, the brown slurry was stirred efficiently with hexane (200 mL) then filtered over celite. After separating the phase, the filter cake was washed with hexane (4×200 mL) which was subsequently used to extract the aqueous layer (4 × 400 mL), the combined hexane layer was evaporated to get the crude product 6.7 g
10 (88.4%) as yellow oil with purity 84% and 5.2% Me₃-TACN and 10.8% unknown impurities included.

5a Preparation of Ts₄-DTNE in butanol/water

The mixture of Ts₂-TACN·TsOH (3.0 g, 5 mmol) and Na₂CO₃ (2.12 g, 20 mmol)
15 in 10 mL water and 1.7 mL butanol was stirred at 115 °C for 5min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed for 3 hrs (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)).After being cooled to room temperature, the mixture was poured into 50 mL water and was filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product
20 2.06 g (94%) with 75% purity.

5b Preparation of Ts₄-DTNE in butanol/water

The mixture of Ts₂-TACN·TsOH (157.5 g, 262.5 mmol) and Na₂CO₃ (106 g, 1 mol) in 360 mL water and 61 mL butanol was stirred at 115 °C for 5min. Then 1,2-
25 dibromoethane(20.8 mL, 242 mmol) was added and the resulting mixture was refluxed for 3 hrs (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)).After being cooled to room temperature, the mixture was poured into 1500 mL water and was filtered. The solid was washed with water (4×1500 mL) to pH 7, dried under vacuum to afford the product 115.9 g (98%) with 75.6% purity.

30

5c Preparation of Ts₄-DTNE in butanol/water

A similar procedure to Experiment 5a was followed to make another batch of Ts₄-DTNE by reacting Ts₂-TACN·TsOH (91.5 g) in butanol/water to afford 68.7 g (100%) Ts₄-DTNE having a purity of 77%.

35

6a Preparation of Me₄-DTNE using Ts₄-DTNE prepared in butanol/water.

- 19 -

Ts₄-DTNE (52 g (77% purity), 44.5 mmol) and 96% sulfuric acid (130.6 mL, composed of 125.6 mL concentrated H₂SO₄ (98%) plus 5ml water) were stirred at 110°C (bath) in a 1L 3-necked flask until TLC showed the detosylation to be completed (about 22hrs). The reactants were cooled to 50°C, then water (121 mL) and NaOH solution (4.63 mol, 185 g NaOH in 230 mL water) was added dropwise under ice-bath with stirring until pH=6~7, then formaldehyde (0.693 mol, 56.6 g(37%)) and formic acid (1.626 mol, 64 g) were added successively with stirring, the mixture was stirred at 90 °C (110 °C oil bath) overnight, then cooled to room temperature, the contents were made strongly alkaline by adding NaOH (56 g in 68 mL water) while maintaining the temperature at 30 °C, the brown slurry was stirred efficiently with hexane (300 mL) then filtered over celite. After separating the phase, the filter cake was washed with hexane (6×200 mL) which was subsequently used to extract the aqueous solution, then the aqueous solution was extracted with hexane (4×500 mL), the combined hexane was evaporated to get the crude product 13.53g as an yellow oil which was redistilled under reduced pressure to afford the product 8.3 g (55%) at 136~138 °C/1mbar as a pale yellow liquid with purity 93%.

6b Preparation of non-distilled Me₄-DTNE using Ts₄-DTNE prepared in butanol/water.

Ts₄-DTNE obtained in step 5b (60 g (75.6% purity), 51.2 mmol) and 96% sulfuric acid (143.5 mL, composed of 138 mL concentrated H₂SO₄ (98%) plus 5.5 mL water) were stirred at 110 °C (bath) in a 1L 3-necked flask until TLC showed the detosylation to be completed (about 22hrs). The reactants were cooled to 50 °C, then water (120 mL) and NaOH solution (198 g NaOH in 300 mL water) was added dropwise under ice-bath with stirring until pH=6~7, then formaldehyde (74 mL(37%)) and formic acid (56 mL) were added successively with stirring, the mixture was stirred at 90 °C (110 °C oil bath) overnight, then cooled to room temperature, the contents were made strongly alkaline by adding NaOH (61.5 g in 78 mL water) while maintaining the temperature at 30 °C, the brown slurry was stirred efficiently with hexane (300 mL) then filtered over celite. After separating the phase, the filter cake was washed with hexane (6×200 mL) which was subsequently used to extract the aqueous solution (4×500 mL), the combined hexane was evaporated and the residue was dried under vacuum to afford the crude product 8.2 g (48%) as a yellow oil with purity 70.5%. The product contains 22.5% Me₃-TACN.

- 20 -

6c Preparation of distilled Me₄-DTNE using Ts₄-DTNE prepared in butanol/water.

Ts₄-DTNE obtained in step 5c (54 g (77% purity), 46.2 mmol) and 96% sulfuric acid (132 mL, composed of 127 mL concentrated H₂SO₄ (98%) plus 5ml water) were stirred at 110 °C (bath) in a 1L 3-necked flask until TLC showed the detosylation to be completed (about 22 hrs). The reactants were cooled to 50 °C, then water (121 mL) and NaOH solution (4.63 mol, 185 g NaOH in 230 mL water) was added dropwise under ice-bath with stirring until pH=6~7, then formaldehyde (69 mL/37%) and formic acid (52 mL) were added successively with stirring, the mixture was stirred at 90 °C (110 °C oil bath) overnight, then cooled to room temperature, the contents were made strongly alkaline by adding NaOH (55.5 g in 70 mL water) while maintaining the temperature at 30 °C, the brown slurry was stirred efficiently with hexane (300 mL) then filtered over celite. After separating the phase, the filter cake was washed with hexane (6×200 mL) which was subsequently used to extract the aqueous solution, then the aqueous solution was extracted with hexane (3×500 mL), the combined hexane was evaporated to get the crude product 16 g as an yellow oil which was redistilled under reduced pressure to afford the product 9.31 g (59.4%) at 140~142 °C/1mbar as a pale yellow liquid with purity 89.3 %. The product contains 1.3% Me₃-TACN.

7 Preparation of Ts₄-DTNE in DMF

The mixture of Ts₂-TACN·TsOH (3.0 g, 5 mmol) and Na₂CO₃ (2.12 g, 20 mmol) in 12 mL DMF was stirred at 110 °C for 5min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed for 3 hrs (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). After being cooled to room temperature, the mixture was poured into 200 mL water and was filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product 1.8 g (82%) with 66% purity.

8 Preparation of Ts₄-DTNE in DMF/H₂O

The mixture of Ts₂-TACN·TsOH (3.0 g, 5 mmol) in 24 mL DMF and Na₂CO₃ (2.12 g, 20 mmol) in 10 mL water was stirred at 110 °C for 5min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed for 4 hrs (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). After being cooled to room temperature, the mixture was poured into 200 mL water and was filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product 1.76 g (80%) with 42% purity.

9 Preparation of Ts₄-DTNE in toluene

The mixture of Ts₂-TACN-TsOH (3.0 g, 5 mmol) and Na₂CO₃ (2.12 g, 20 mmol) in 20 mL toluene was stirred at 125 °C for 5min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed for 6 hrs (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). Then the solvent was evaporated and to the residue 50 mL water was added and the resulting mixture was filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product 1.9 g (83%) with 56% purity.

10 Preparation of Ts₄-DTNE in acetone

The mixture of Ts₂-TACN-TsOH (3.0 g, 5 mmol) and Na₂CO₃ (2.12 g, 20 mmol) in 20 mL acetone was stirred at 85 °C for 5min. Then 1,2-dibromoethane(0.43 mL, 5 mmol) was added and the resulting mixture was refluxed for 3 hrs (TLC showed the completion of the reaction, CH₂Cl₂/methanol (97:3)). Then the solvent was evaporated and to the residue 50 mL water was added and the resulting mixture was filtered. The solid was washed with water (4×50 mL), dried under vacuum to afford the product 1.95 g (89%) with 66% purity.

11. General procedure for the preparation of [Mn₂(μ-O)₂(μ-CH₃COO)(Me₄-DTNE)]Cl₂

Under N₂, to Me₄-DTNE in EtOH/H₂O(2:1,v/v), solid mixture of MnCl₂•4H₂O and sodium acetate were added. The mixture was stirred for 30 min at 58 °C. After another stirring for 10 min cooled in an ice/water bath, the freshly prepared mixture of 1 M of H₂O₂ in water and 1.5 M of NaOH was added dropwise over 5 min. The mixture turned immediately dark green-brown. The mixture was stirred for 20 min in an ice water bath and then for 20 min at room temperature. 1 M of acetic acid was added. After stirring for another 20 min, the mixture was filtered to remove the brown solid and the filtering bed was washed with ethanol. Then the green filtrate was evaporated (the water bath temperature < 45 °C). The residual dark green oil was co-evaporated with ethanol and ethyl acetate to facilitate the removal of most of the remaining water. Dark green oils were taken up in ethanol, and the insoluble white salts separated by filtration were washed with ethanol. After removing all ethanol, the dark green oil was obtained again. The small amount of ethanol was added and stirred for 2 min. Then the large amount of ethyl acetate was added. The green solid was precipitated immediately. After 3 hours at -20 °C, the suspension was filtered off, with obtaining a green solid,

- 22 -

which was washed with ethyl acetate, n-hexane, and dried under vacuum at 45 °C for 5 hrs to afford dark green powder as $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$.

11.1 Preparation of $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-CH}_3\text{COO})(\text{Me}_4\text{-DTNE}))\text{Cl}_2]$ from the distilled $\text{Me}_4\text{-DTNE}$ in the BuOH/H₂O route

Distilled $\text{Me}_4\text{-DTNE}$ obtained according to Example 6a (Example 11.1a) or 6c (Example 11.1b) (89.3% purity with 1.3% $\text{Me}_3\text{-TACN}$) (765 mg, 2 mmol); EtOH/H₂O (2:1, v/v): 20 mL; $\text{MnCl}_2\cdot 4\text{H}_2\text{O}$ (840 mg, 4.2 mmol); NaAc (82 mg, 1 mmol); 1 M of H_2O_2 in water (5 mL, 5 mmol); 1.5 M of NaOH (2.5 mL, 3.75 mmol); 1 M of HAc (1.25 mL, 1.25 mmol). 1.2 g of green powder as $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$.

UV-Vis purity of 91.1%, the yield of 86.8% (The yield (%) = the weight of the compound (g) × the purity of the compound (%) / the calcd. weight of the compound (g)). UV-Vis spectrum of a purified sample: (ϵ : $\text{mol}^{-1}\cdot\text{L}\cdot\text{cm}^{-1}$, in water, Mw: 630): 271 nm (13332), 554 nm (317), 639 nm (327).

UPLC analysis confirmed 1.53 % of the free $[\text{H}_2(\text{Me}_4\text{-DTNE})]\text{Cl}_2$, 0.7 % of the free $[\text{H}(\text{Me}_3\text{-TACN})]\text{Cl}$, and 0.08% of $[(\text{Mn}_2(\mu\text{-O})_3(\text{Me}_3\text{-TACN}))\text{Cl}_2]$.

Total chloride amount was 11.17 %.

Water analysis (Karl-Fischer method): Anal. calcd. for $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$: 2.86%; Found: 1.14%.

11.2 Preparation of $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-CH}_3\text{COO})(\text{Me}_4\text{-DTNE}))\text{Cl}_2]$ from the undistilled $\text{Me}_4\text{-DTNE}$ in the BuOH/H₂O route

Undistilled $\text{Me}_4\text{-DTNE}$ obtained according to Example 6a (Example 11.2a) or 6b (Example 11.2b) (70.5% purity with 22.8% $\text{Me}_3\text{-TACN}$) (1.93 g, 4 mmol); EtOH/H₂O (2:1, v/v): 40 mL; $\text{MnCl}_2\cdot 4\text{H}_2\text{O}$ (2.22 g, 11.2 mmol); NaAc (166 mg, 2 mmol); 1 M of H_2O_2 in water (15 mL, 15 mmol); 1.5 M of NaOH (7.5 mL, 11.25 mmol); 1 M of HAc (2.5 mL, 2.5 mmol). 2.93 g of green powder as $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$.

UV-Vis purity of 84.6%, the yield of 75.5%. (The yield (%) = the weight of the compound (g) × the purity of the compound (%) / (the calcd. weight of the compound (g) + the calcd. weight of $[(\text{Mn}_2(\mu\text{-O})_3(\text{Me}_3\text{-TACN}))\text{Cl}_2]$ (g))).

UPLC analysis confirmed 6.96 % of the free $[\text{H}_2(\text{Me}_4\text{-DTNE})]\text{Cl}_2$, 3.2 % of the free $[\text{H}(\text{Me}_3\text{-TACN})]\text{Cl}$, and 4.3% of $[(\text{Mn}_2(\mu\text{-O})_3(\text{Me}_3\text{-TACN}))\text{Cl}_2]$.

Total chloride amount was 10.35 %.

- 23 -

Water analysis (Karl-Fischer method): Anal. calcd. for $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$: 2.86%; Found: 1.07%.

11.3 Preparation of $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-CH}_3\text{COO})(\text{Me}_4\text{-DTNE}))\text{Cl}_2]$ from the undistilled $\text{Me}_4\text{-DTNE}$ in the $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ route

Undistilled $\text{Me}_4\text{-DTNE}$ obtained according to Example 4a (Example 11.3a) or 4b (Example 11.3b)(84% purity with 5.2% $\text{Me}_3\text{-TACN}$): (1.62 g, 4 mmol); $\text{EtOH}/\text{H}_2\text{O}$ (2:1, v/v): 40 mL; $\text{MnCl}_2\cdot 4\text{H}_2\text{O}$ (1.78 g, 9 mmol); NaAc (166 mg, 2 mmol); 1 M of H_2O_2 in water (9 mL, 9 mmol); 1.5 M of NaOH (4.5 mL, 6.75 mmol); 1 M of HAc (2.5 mL, 2.5 mmol). 2.6 g of green powder as $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$.

UV-Vis purity of 84.8%, the yield of 88.7% (The yield (%) = the weight the compound (g) $\times$ the purity of the compound (%) / the calcd. weight of the compound (g)).

UPLC analysis confirmed 7.2 % of the free $[\text{H}_2(\text{Me}_4\text{-DTNE})]\text{Cl}_2$, 2.56 % of the free $[\text{H}(\text{Me}_3\text{-TACN})]\text{Cl}$, and 0.14 % of $[(\text{Mn}_2(\mu\text{-O})_3(\text{Me}_3\text{-TACN}))\text{Cl}_2]$.

Total chloride amount was 10.91 %.


Water analysis (Karl-Fischer method): Anal. calcd. for $[(\text{Mn}_2(\mu\text{-O})_2(\mu\text{-OAc})(\text{Me}_4\text{-DTNE}))\text{Cl}_2\cdot\text{H}_2\text{O}]$: 2.86%; Found: 1.35%.

Using acetonitrile/ H_2O as solvent for the formation of $\text{Ts}_4\text{-DTNE}$ has advantages as the purity $\text{Me}_4\text{-DTNE}$ product is much higher than when using other solvents. This leads to formation of $\text{Me}_4\text{-DTNE}$ ligand that does not need to be further purified to make the dinuclear manganese complex. The method using butanol/water leads to a need to be distilled the $\text{Me}_4\text{-DTNE}$ ligand to obtain high-purity material, leading to significant losses in yields. A yield improvement of the ligand of about 20% can be thus achieved (Experiment 4a vs experiment 6a or Experiment 4b vs Experiments 6b & 6c).

- 24 -


CLAIMS

1. A method of producing a compound of formula (A):



- 5 the method comprising the following step:

- (a) reacting a compound of formula (B):



- 10 in an acidic medium comprising sulfuric acid, the molar ratio of B to sulfuric acid in the range from 1:0.5 to 1:10, wherein P is an arylsulfonate protecting group and the compound of formula (A) is isolated as a protonated salt in amorphous or crystalline form.

2. The method of claim 1, wherein an acid anhydride is present in the acidic medium.

15

3. The method of claim 1 or 2, wherein the counterion of the salt is selected from: toluenesulfonic acid; benzenesulfonic acid; sulfuric acid; acetic acid; formic acid; and, propionic acid.

20

4. The method of claim 3, wherein the protonated salt is a salt of an aryl sulfonic acid.


5. The method of claim 3, wherein the protonated salt is the toluene sulfonic acid salt of 1,4-ditosyl-1,4,7-triazacyclononane.

25

6. The method of claim 1 or 2, wherein the method is conducted at a reaction temperature from 100 to 160 °C.


- 25 -

7. A method of producing a compound of formula (A):



the method comprising the following step:

- 5 (a) reacting a compound of formula (B):




in an acidic medium, wherein P is an arylsulfonate protecting group, wherein the acidic medium is worked-up when the conversion of B to A is at least 50 mol % yielding compound (A).

10

8. The method of claim 7, wherein the acidic medium comprises sulfuric acid.
9. The method of claim 8, wherein the acidic medium further comprises an acid anhydride.
10. The method of any one of claims 7 to 9, wherein the method is conducted at a reaction temperature from 100 to 160 °C.

15


11. A method of producing a compound of formula (A):



20

the method comprising reacting a compound of formula (B):

- 26 -



in an acidic medium comprising an acid anhydride, wherein P is an arylsulfonate protecting group.

5


12. The method of claim 11, wherein the compound of formula A is isolated as a protonated salt of the same arylsulfonic acid of which P is the arylsulfonate.

10

13. The method of claim 11 of claim 12, wherein the protonated salt is the toluene sulfonic acid salt of 1,4-ditosyl-1,4,7-triazacyclononane.

15

14. The method of claim 1, claim 7 or claim 11, further comprising reacting a compound of formula (A) with a bridging element of the form ZCH_2CH_2Z , wherein Z is a halogen selected from: Cl; Br; and, I, whereby to produce a compound of formula (C):



wherein groups P in the compound of formula (C) are the same as groups P in the compound of formula (A).


20

15. The method of claim 14 further comprising deprotecting the compound of formula (C) with an acidic medium and to form 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane and subsequently adding formaldehyde and formic acid to the reaction medium whereby to produce 1,2-bis-(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)-ethane.

25

- 27 -

16. A protonated salt of formula (A), having a counter ion HX, the protonated salt in amorphous or crystalline form:




wherein P is a tosylate or benzene sulfonate and HX is selected from:
toluenesulfonic acid and benzenesulfonic acid.

17. The protonated salt of claim 16, which is (1,4-ditosyl-1,4,7-triazacyclonone) tosylate or (1,4-dibenzenesulfonyl-1,4,7-triazacyclonone) benzenesulfonate.


18. The protonated salt of claim 16, which is (1,4-ditosyl-1,4,7-triazacyclonone) tosylate.

19. A method of producing a compound of formula (C):



the method comprising the following step:

- (a) reacting a compound of formula (A):



with a bridging element of the form ZCH_2CH_2Z , wherein P is an arylsulfonate protecting group and Z is a halogen selected from: Cl; Br; and, I.

- 28 -

20. The method of claim 19, wherein the compound of formula (A) is reacted with the bridging element in a solvent selected from: acetonitrile; 1-butanol; 2-butanol; and, t-butanol.

5 21. The method of claim 20, wherein the compound of formula (A) is reacted with the bridging element in acetonitrile as solvent.

22. The method of claim 21, wherein the solvent comprises from 10 to 90 wt/wt % water.


10

23. The method of claim 22, wherein the acetonitrile comprises from 10 to 50 wt/wt % water.

15

24. The method of claim 23, wherein the acetonitrile comprises from 10 to 35 wt/wt % water.

25. A one-pot method for the preparation of 1,2-bis-(4,7-dimethyl-1,4,7-triazacyclonon-1-yl)-ethane (Me₄-DTNE) the method comprising deprotecting a compound of formula (C):



20

with an acidic medium and to form 1,2-bis(1,4,7-triazacyclonon-1-yl)-ethane and subsequently adding formaldehyde and formic acid to the reaction medium, wherein P is an arylsulfonate.

25 26. The one-pot method of claim 25, wherein P is tosylate.