



US00PP31965P2

(12) **United States Plant Patent**
Trees

(10) **Patent No.:** **US PP31,965 P2**

(45) **Date of Patent:** **Jul. 14, 2020**

(54) **LANTANA PLANT NAME ‘BALOOMANG’**

(50) Latin Name: *Lantana camara*
Varietal Denomination: **Baloomang**

(71) Applicant: **Ball Horticultural Company**, West
Chicago, IL (US)

(72) Inventor: **Scott C. Trees**, Arroyo Grande, CA
(US)

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/602,134**

(22) Filed: **Aug. 12, 2019**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/86 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./227**

(58) **Field of Classification Search**
USPC **Plt./227**
See application file for complete search history.

Primary Examiner — Susan McCormick Ewoldt

(74) *Attorney, Agent, or Firm* — Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Lantana* plant named
‘Baloomang’, characterized by its yellow, orange, and red-
dish-purple multicolored inflorescences, dark green-colored
foliage, and moderately vigorous, upright, mounded to
rounded growth habit, is disclosed.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Lan-
tana camara*.
Variety denomination: ‘Baloomang’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Lantana* plant botanically known as *Lantana camara* and
hereinafter referred to by the cultivar name ‘Baloomang’.

The new cultivar originated in a controlled breeding
program in Guadalupe, Calif. during September 2013. The
objective of the breeding program was the development of
Lantana cultivars with sterile flowers for continuous flow-
ering and a moderately vigorous, upright, mounded to
rounded growth habit.

The new *Lantana* cultivar is the result of cross-pollina-
tion. The female (seed) parent of the new cultivar is the
proprietary *Lantana camara* breeding selection coded 9087,
not patented, characterized by its dark red-colored inflores-
cences, medium green-colored foliage, low growth vigor
and compact-mounded growth habit. The male (pollen)
parent of the new cultivar is BANDANA Peach
‘LANZ0002’, U.S. Plant Pat. No. 23,317, characterized by
its yellow, orange, and light red multicolored flowers,
medium green-colored foliage, and moderately vigorous,
relatively compact and mounding growth habit. The new
cultivar was discovered and selected as a single flowering
plant within the progeny of the above stated cross-pollina-
tion during September 2014 in a controlled environment in
Guadalupe, Calif.

Asexual reproduction of the new cultivar by terminal stem
cuttings since September 2014 in Guadalupe, Calif. and
Arroyo Grande, Calif. has demonstrated that the new culti-
var reproduces true to type with all of the characteristics, as
herein described, firmly fixed and retained through succes-
sive generations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have
been repeatedly observed and can be used to distinguish
‘Baloomang’ as a new and distinct cultivar of *Lantana* plant:

2

1. Yellow, orange, and reddish-purple multicolored inflo-
rescences;
2. Dark green-colored foliage; and
3. Moderately vigorous, upright, mounded to rounded
growth habit.

Plants of the new cultivar differ from plants of the female
parent primarily in inflorescence color, darker green foliage
color, and a more vigorous growth habit. Plants of the new
cultivar differ from plants of the male parent primarily in
having more reddish-purple color in the flower margins of
mature inflorescences.

Of the many commercially available *Lantana* cultivars,
the most similar in comparison to the new cultivar is
Bloomify Rose ‘UF-1011-2’, U.S. Plant Pat. No. 29,267.
However, in side-by-side comparisons, plants of the new
cultivar differ from plants of ‘UF-1011-2’ in at least the
following characteristics:

1. Plants of the new cultivar have darker orange-colored
inflorescences than plants of ‘UF-1011-2’; and
2. Plants of the new cultivar have darker green-colored
leaves than plants of ‘UF-1011-2’.

The new cultivar can also be compared to Lucky Peach
‘Balucepa20’, U.S. Plant patent application Ser. No. 16/602,
137. However, in side-by-side comparison, plants of the new
cultivar differ from plants of ‘Balucepa20’ in at least the
following characteristics:

1. Plants of the new cultivar are taller and wider than
plants of ‘Balucepa20’;
2. Plants of the new cultivar have flower buds with more
pink-coloration than plants of ‘Balucepa20’; and
3. Plants of the new cultivar have larger leaves than plants
of ‘Balucepa20’.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it
is reasonably possible to make the same in color illustrations
of this type, typical flower and foliage characteristics of the
new cultivar. Colors in the photographs may differ slightly

from the color values cited in the detailed description, which accurately describes the colors of 'Baloomang'. The plants were approximately 4 months old and grown in 4.5-inch pots for approximately 12 weeks in a greenhouse in West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and flowering habit of 'Baloomang'.

FIG. 2 illustrates a close-up view of an individual inflorescence of 'Baloomang'.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2015 edition, except where general color terms of ordinary significance are used. The color values were determined in April 2019 under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe approximately 4-month old plants produced from cuttings from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown in West Chicago, Ill. in 4.5-inch pots for approximately 12 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 67° F. to 72° F. (19° C. to 22° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Supplemental lighting was used. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Lantana camara* 'Baloomang'.

Parentage:

Female parent.—Proprietary *Lantana camara* breeding selection coded 9087, not patented.

Male parent.—BANDANA Peach, U.S. Plant Pat. No. 23,317.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 7 to 11 days.

Time to produce a rooted cutting.—Approximately 24 to 35 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 7 to 8 weeks from a rooted cutting to finish in a 10 cm pot.

Growth habit and general appearance.—Moderately vigorous, upright, mounded to rounded.

Size.—Height from soil level to top of plant plane: Approximately 19.0 cm. Width: Approximately 36.0 cm.

Branching habit.—Freely branching, pinching enhances branching. Quantity of lateral branches per plant: Approximately 5.

Branch.—Shape: Square in cross section. Strength: Strong, becomes woody with age. Length: Approximately 15.0 cm. Diameter: Approximately 3.0 mm. Length of central internode: Approximately 2.0 cm. Texture: Densely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Colorless,

transparent. Color of young stem: 144A. Color of mature stem: 146B becoming woody N199A with age.

Foliage description:

General description.—Quantity of leaves per lateral branch: Approximately 12. Fragrance: Strong, spicy. Form: Simple. Arrangement: Opposite.

Leaves.—Aspect: Perpendicular to obtuse angle to stem. Shape: Ovate. Margin: Serrate. Apex: Acute. Base: Obtuse. Venation pattern: Pinnate. Length of mature leaf: Approximately 7.5 cm. Width of mature leaf: Approximately 4.5 cm. Texture of upper surface: Scabrous. Texture of lower surface: Scabrous and glandular pubescent along venation. Gland color: Colorless, transparent. Color of upper surface of young foliage: 137A with venation of 146B to indistinguishable. Color of lower surface of young and mature foliage: Closest to 138B with venation of 146C. Color of upper surface of mature foliage: Closest to NN137A with 139A, venation of 146B to indistinguishable.

Petiole.—Length: Approximately 1.7 cm. Diameter: Approximately 2.0 mm. Texture: Scabrous and glandular pubescent. Gland color: Colorless, transparent. Color: 146B.

Flowering description:

Flowering habit.—'Baloomang' is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual inflorescence on the plant.—Approximately 2 to 3 weeks.

Inflorescence description:

General description.—Type: Hemispherical head, axillary or terminal. Quantity per plant: Approximately 12. Fragrance: Strong, spicy. Aspect: Facing upward or outward. Height: Approximately 2.0 cm. Width: Approximately 4.5 cm. Quantity of fully open flowers per inflorescence: Approximately 34.

Peduncle.—Strength: Strong. Shape: Square in cross section. Aspect: Acute angle to stem. Length: Approximately 4.0 cm. Diameter: Approximately 2.0 mm. Texture: Scabrous and glandular pubescent. Gland color: Colorless, transparent. Color: 146B.

Flower description:

General description.—Type: Salverform.

Bud.—Rate of opening: Generally takes 1 to 2 days for bud to progress from first color to fully open flower. Buds open in progression from the margin to the center of the inflorescence. Quantity of unopened inflorescences per plant: Approximately 14.

Bud just before opening.—Shape: Elongated, rectangular at apex. Length: Approximately 1.0 cm. Diameter: Approximately 3.0 mm. Color: 37B.

Corolla.—Depth: Approximately 2.0 cm. Diameter: Approximately 1.2 cm.

Petals.—Quantity: 4, non-imbricate, non-symmetrical petals. Petals are fused at base forming a corolla tube. Shape: Obovate. Appearance: Dull. Aspect: Cupped. Margin: Entire to erose, ruffled. Apex: Obtuse. Length of upper petal from throat: Approximately 6.0 mm. Width of upper petal: Approximately 9.0 mm. Length of lateral petals from throat: Approximately 5.0 mm. Width of lateral petals: Approximately 6.0 mm. Length of lower petal from

throat: Approximately 7.0 mm. Width of lower petal: Approximately 9.0 mm. Texture of upper surface: Glabrous. Texture of lower surface: Densely pubescent. Color of upper surface when first open: 12A with distal half of lower petal of 12D, transitioning to 15B with distal half of lower petal of 27B. Color of lower surface when first open: 4D tinted with 53C, tint becomes stronger with age. Color of upper surface when fully open: N25C to N25A with distal half of lower petal transitioning to N77C. Color of lower surface when fully open: 23D tinted with 53D.

Corolla tube.—Length: Approximately 1.3 cm. Diameter at tube opening: Approximately 1.0 mm. Diameter at base: Approximately 1.0 mm. Texture of inner surface: Densely pubescent. Texture of outer surface: Densely pubescent with a mixture of glandular and nonglandular hairs at tube opening transitioning to glabrous at base. Pubescence color: Colorless and 53A. Color of inner surface: 12B darkening to N25D. Color of outer surface: 145C with an overlay of 53D, overlay becomes heavier with age.

Calyx.—Shape: Tubular with two broadly acute tips. Length: Approximately 3.0 mm. Diameter at tip: Approximately 1.5 mm. Diameter at base: Approximately 1.5 mm. Texture of inner surface: Glabrous. Texture of outer surface: Densely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Colorless, transparent. Color of inner and outer surfaces: 145D.

Bracts.—Quantity per flower: 1 per flower. Shape: Lanceolate. Length: Approximately 7.0 mm. Width: Approximately 2.0 mm. Texture of upper surface: Sparsely pubescent. Texture of lower surface: Densely pubescent. Color of upper surface: 137A with 146B at base. Color of lower surface: 137B with 146B at base.

Reproductive organs.—Androecium: Stamen quantity: 4, adnate to corolla tube. Stamen length: Approximately 2.0 mm. Anther shape: Bilobed, ovoid. Anther length: Approximately 1 mm. Anther color: 13B. Pollen amount: None observed. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 3.0 mm. Stigma shape: Misshapen oval. Stigma length: Less than 1 mm. Stigma color: 144A. Style length: Approximately 2.0 mm. Style color: 145D, translucent. Ovary diameter: Approximately 1.0 mm. Ovary color: 144C.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Lantana* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Lantana* plant named 'Baloomang', substantially as herein illustrated and described.

* * * * *




FIG. 1



FIG. 2