



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2542854 C 2012/04/17

(11)(21) **2 542 854**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2004/11/12
(87) Date publication PCT/PCT Publication Date: 2005/06/02
(45) Date de délivrance/Issue Date: 2012/04/17
(85) Entrée phase nationale/National Entry: 2006/04/18
(86) N° demande PCT/PCT Application No.: US 2004/037975
(87) N° publication PCT/PCT Publication No.: 2005/049142
(30) Priorité/Priority: 2003/11/14 (US60/520,182)

(51) Cl.Int./Int.Cl. *A61K 31/565* (2006.01),
A61K 31/567 (2006.01), *A61K 45/06* (2006.01),
A61P 15/18 (2006.01)
(72) Inventeur/Inventor:
BOISSONNEAULT, ROGER M., US
(73) Propriétaire/Owner:
WARNER CHILCOTT COMPANY, LLC, US
(74) Agent: SIM & MCBURNEY

(54) Titre : CONTRACEPTIF D'OESTROGENE PROGRESSIF
(54) Title: GRADUATED ESTROGEN CONTRACEPTIVE

(57) **Abrégé/Abstract:**

A multiphasic estrogenic/progestogenic contraceptive regimen that provides for the reduction or elimination of estrogen in the initial phase is disclosed. Also described is a contraceptive kit that may be used to practice the method of the invention.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 June 2005 (02.06.2005)

PCT

(10) International Publication Number
WO 2005/049142 A3

(51) International Patent Classification⁷: **A61K 31/565**,
31/567, 45/06, A61P 15/18

(21) International Application Number:
PCT/US2004/037975

(22) International Filing Date:
12 November 2004 (12.11.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/520,182 14 November 2003 (14.11.2003) US

(71) Applicant (for all designated States except US):
WARNER CHILCOTT COMPANY, INC. [US/US];
Rockaway 80 Corporate Center, 100 Enterprise Drive,
Suite 280, Rockaway, NJ 07866 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **BOISSONNEAULT,**
Roger, M. [US/US]; 5 N. Bridge Drive, Long Valley, NJ
07853 (US).

(74) Agents: **MANDRA, Raymond, R.** et al.; Fitzpatrick,
Cella, Harper & Scinto, 30 Rockefeller Plaza, New York,
NY 10112-3801 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE,
SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
29 September 2005

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: GRADUATED ESTROGEN CONTRACEPTIVE

(57) Abstract: A multiphasic estrogenic/progestogenic contraceptive regimen that provides for the reduction or elimination of es-
trogen in the initial phase is disclosed. Also described is a contraceptive kit that may be used to practice the method of the invention.



WO 2005/049142 A3

- 1 -

TITLE

GRADUATED ESTROGEN CONTRACEPTIVE

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention is directed to a method of contraception that provides for the reduction or elimination of estrogen in the initial phase of a multiphasic estrogenic/progestogenic contraceptive regimen without compromising contraceptive efficacy or cycle control. The invention is also directed to a multiphase contraceptive kit that may be used to practice the method of the invention.

Related Background Art

[0002] Contraceptive compositions containing both estrogenic and progestogenic compounds are well known. The progestogenic component of the composition is primarily responsible for the contraceptive efficacy of the composition, while the estrogenic component is employed to reduce undesired side effects, such as breakthrough bleeding or spotting.

- 2 -

[0003] The earliest of these estrogenic/progestogenic contraceptive compositions contained a relatively high level of estrogenic component. A constant goal, however, has been to reduce the estrogenic potency of such compositions without reducing contraceptive efficacy and increasing undesired side effects. As described in U.S. Patent No. 5,888,543, in an attempt to achieve this goal, numerous regimens have been developed in which the progestin/estrogen combination is administered in a monophasic regimen (fixed dose) or as biphasic or triphasic regimens (varied dose).

[0004] A particularly advantageous technique for reducing total estrogenic administration is described in U.S. Patent No. 4,962,098. This describes a triphasic method of contraception using a progestogen/estrogen combination in which the amount of estrogen is increased stepwise over the three phases. The first phase is 4-7 days, the second phase is 5-8 days and the third phase is 7-12 days. Preferably, the administration of the contraceptive compositions for the three phases will be 21 days followed by a 7 day placebo period. For all three phases the progesten is 0.5 to 1.5 mg of norethindrone acetate, while about 10 to 30 mcg of ethinyl estradiol is used in the first phase, about 20 to 40 mcg of ethinyl estradiol is used in the second phase and 30 to 50 mcg of ethinyl estradiol is employed in the third phase.

[0005] There is a continuing desire, however, to further reduce the amount of estrogenic component in an estrogenic/progestogenic composition with continued contraceptive efficacy while avoiding undesired side effects. Heretofore it was believed that at least 10 mcg of ethinyl estradiol or its estrogenic equivalent was needed in an estrogenic/progestogenic composition to assure contraceptive efficacy. It has now been surprisingly discovered that the amount of estrogenic component in the first phase of a triphasic regimen can be significantly reduced or eliminated without compromising efficacy or cycle control.

- 3 -

SUMMARY OF THE INVENTION

[0006] This invention is directed to a multiphasic method of contraception that provides for the reduction or elimination of administered ethinyl estradiol in the first phase without a reduction in contraceptive efficacy or an increase in undesired side effects. The method of this invention includes administering, in sequential steps, to a female of child bearing age the following compositions: (a) composition I for about 5 to about 9 days; (b) composition II for about 5 to about 9 days; and (c) composition III for about 8 to about 12 days. Compositions I, II and III all contain a progestogen in an amount equivalent to about 0.3 to about 1.5 mg, preferably about 0.5 to about 1.5 mg of norethindrone acetate. Composition I contains an estrogen in an amount equivalent to about 0 to less than about 10 mcg of ethinyl estradiol and both compositions II and III contain an estrogen in an amount equivalent to about 10 to about 50 mcg of ethinyl estradiol.

[0007] Significantly, the sequential administration of compositions I, II and III is repeated after a period of about 1 to about 4 days has elapsed after completion of the administration of composition III. Without being bound by theory, it is believed that this relatively short interim period between the sequential administration of the estrogenic/progestogenic components allows for the advantageous reduction or elimination of estrogen from the first phase of the above-described triphasic regimen without compromising efficacy or cycle control. It is also preferable that the amount of estrogen be increased by at least an amount equivalent to 5 mcg of ethinyl estradiol between composition II and composition III. In a preferred embodiment of this invention, the estrogen is ethinyl estradiol and the progestogen is norethindrone acetate.

[0008] Yet another embodiment of this invention is directed to a multiphase combination and contraceptive kit comprising a package containing daily dosages of: (a) a Phase I composition containing a progestogen in an amount equivalent to about 0.3 to about 1.5 mg, preferably about 0.5 to about 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to about 0 to about 10 mcg of ethinyl estradiol; (b) a Phase II composition containing a progestogen in an amount

equivalent to about 0.3 to about 1.5 mg, preferably about 0.5 to about 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to about 10 to about 50 mcg of ethinyl estradiol; and (c) a Phase III composition containing a progestogen in an amount equivalent to about 0.3 to about 1.5 mg, preferably about 0.5 to about 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to about 10 to about 50 mcg of ethinyl estradiol; wherein the amount of estrogen in the Phase III composition is at least an amount equivalent to 5 mcg of ethinyl estradiol greater than the amount of estrogen in the Phase II composition. Preferably, the estrogen used in the kit is ethinyl estradiol and the progestogen is norethindrone acetate.

In accordance with an aspect of the present invention, there is provide a multiphase combination and contraceptive kit comprising a package containing daily dosages of: a) a Phase I composition containing a progestogen in an amount equivalent to 0.3 to 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to 0 to less than 10 mcg of ethinyl estradiol; b) a Phase II composition containing a progestogen in an amount equivalent to 0.3 to 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to 10 to 50 mcg of ethinyl estradiol; and c) a Phase III composition containing a progestogen in an amount equivalent to 0.3 to 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to 10 to 50 mcg of ethinyl estradiol, wherein the amount of estrogen in the Phase III composition is greater than the amount of estrogen in the Phase II composition by at least an amount equivalent to 5 mcg ethinyl estradiol.

DETAILED DESCRIPTION OF THE INVENTION

[0009] The method of this invention is practiced by administration of the compositions in a numeric sequence with the Phase I composition being used first, the Phase II composition being used second, etc. If packaging and/or other requirements dictate, the method and kit described herein can be employed as part of a larger scheme for contraception or treatment of gynecological disorders. While the sequence in which Applicant's combinations are administered is important to their operation, it should be kept in mind that variations in timing and dosage can be tolerated when medical considerations so dictate.

[0010] Significantly, the method of this invention provides that the sequential administration of compositions I, II and III is repeated after a period of about 1 to about 4 days has elapsed after the completion of the administration of composition III. More preferably, the number of days between the completion of the administration of composition III and beginning the repeated

sequential administration of compositions I, II and m is from about 2 to about 4 days. During this interim period an iron supplement and/or a placebo may be preferably administered on a daily basis, although there is no requirement for the administration of anything during this interim period, i. e., the period between the completion of the

- 5 -

prior sequential administration of compositions I-III and the start of the next sequential administration of compositions I-III.

[0011] Estrogens which may be used in the present invention include, for example, ethinyl estradiol, 17 β -estradiol, 17 β -estradiol-3-acetate, mestranol, conjugated estrogens, USP and estrone or salts thereof. The amount of estrogen used is described herein as that which is "equivalent" in estrogenic potency to an amount of ethinyl estradiol. The equivalent estrogenic potency of an estrogen to ethinyl estradiol may be readily determined by one of ordinary skill in the art. It is contemplated that each Phase could employ one or more different estrogens that deliver a potency equivalent to the recited amount of ethinyl estradiol. It is also contemplated that the estrogen used in one Phase may be different than that used in another Phase. In a most preferred embodiment of this invention, however, the estrogen for each Phase, if present, is ethinyl estradiol.

[0012] Progestogens which may be used in the present invention include, for example, progesterone and its derivatives such as 17-hydroxy progesterone esters and 19-nor-17-hydroxy progesterone esters, 17-alpha-ethinyl testosterone, 17-alpha-ethinyl-19-nortestosterone (norethindrone) and derivatives thereof, norethindrone acetate, norgestrel, norgestamate, desogestrel and D-17-beta-acetoxy-17-beta-ethyl-17-alpha-ethinyl-gon-4-en-3-one oxime. Other exemplary progestogens include demegestone, drospirenone, dydrogesterone, gestodene, medrogestone, medroxy progesterone and esters thereof. The amount of progestogen used is described herein as that which is "equivalent" in progestogenic potency to an amount of norethindrone acetate. The equivalent progestogenic potency of a progestogen to norethindrone acetate may be readily determined by one of ordinary skill in the art. It is contemplated that each Phase could employ one or more different progestogens that deliver a potency equivalent to the recited amount of norethindrone acetate. It is also contemplated that the progestogen used in one Phase may be different than that used in another Phase. In a most preferred embodiment of this invention, however, the progestogen for each Phase is norethindrone acetate.

[0013] Accordingly, in a preferred embodiment of this invention the compositions employed in accordance with the invention will contain in Phase I about 0.3-1.5 mg,

- 6 -

preferably about 0.5-1.5 mg norethindrone acetate and about 0 to less than about 10 mcg ethinyl estradiol, preferably about 0 to about 5 mcg ethinyl estradiol, in Phase II about 0.3-1.5 mg, preferably about 0.5-1.5 mg norethindrone acetate and about 10-50 mcg ethinyl estradiol, preferably about 20-40 mcg ethinyl estradiol, and in Phase III about 0.3-1.5 mg, preferably about 0.5-1.5 mg norethindrone acetate and about 10-50 mcg ethinyl estradiol, preferably about 25-50 mcg ethinyl estradiol, wherein the amount of ethinyl estradiol is increased by at least 5 mcg from Phase II to Phase III

[0014] A significant aspect of the method and kit of this invention is that the Phase I composition has a significantly lower concentration of estrogen equivalent to ethinyl estradiol than previously considered possible, while maintaining contraceptive efficacy and avoiding or minimizing unwanted side effects such as break through bleeding. In one particularly preferred embodiment the amount of estrogen equivalent to ethinyl estradiol in the Phase I composition is about 5 mcg. In another particularly preferred embodiment the Phase I composition is substantially free of estrogen, and most preferably is substantially free of ethinyl estradiol. As used herein "substantially free" means that estrogen is not detectable or only pharmacologically insignificant minor levels are present.

[0015] An optional Phase IV composition, which contains an iron supplement, e.g., ferrous fumarate, and/or one or more placebos, can be used in conjunction with the other three.

[0016] The particularly preferred compositions employed in accordance with the invention in Phases I through IV will more preferably have the administration times and drug contents set forth in the following tables when a four-phase system is used. Each table sets forth relevant values for one of Applicant's preferred embodiments, or configurations, for administration of the system to females.

- 7 -

Table 1

Phase	Days	mg Norethindrone acetate	mcg EE	mg Fumarate
I	7	1.0	5	0
II	7	1.0	30	0
III	10	1.0	35	0
IV	4	--	--	75

Table 2

Phase	Days	mg Norethindrone acetate	mcg EE	mg Fumarate
I	7	1.0	0	0
II	7	1.0	30	0
III	10	1.0	35	0
IV	4	--	--	75

[0017] The norethindrone acetate (NA) and ethinyl estradiol (EE) are well known and readily available. Clearly, the amount of NA and EE may be varied in accordance with the disclosure of this invention. For example, the amount of NA set forth in Tables 1 and 2 could readily be adjusted from 1 mg to 0.5 mg or 0.4 mg.

[0018] The designation "mcg" refers to micrograms and "mg" to milligrams.

[0019] It should be noted that these tables are presented for illustrative purposes only. The substitution of functionally equivalent amounts and kinds of reagent(s) in these schemes is contemplated. For example, the use of sugar or other placebo in place of all or part of the ferrous fumarate is envisioned.

[0020] The compositions used in this invention are administered using a suitable daily dosage form. Tablets, pills, capsules and caplets are exemplary dosage forms.

[0021] In addition, the use of other conventional additives, e.g., fillers, colorants, polymeric binders, and the like is also contemplated. In general any pharmaceutically-acceptable additive which does not interfere with the function of the active components can be used in one or more of the compositions.

- 8 -

[0022] Suitable carriers with which the compositions can be administered include lactose, starch, cellulose derivatives and the like used in suitable amounts. Lactose is a preferred carrier. Mixtures of carriers, e.g. lactose, microcrystalline cellulose and starch, are operable.

[0023] While the norethindrone acetate is preferred, as previously noted it may be replaced by a different progestogen. Similarly, while the ethinyl estradiol component is preferred it may be completely or partially replaced with one or more conventional estrogenic substances, e.g., mestranol.

[0024] While the invention is discussed as potentially one employing four phases, it clearly may employ only three. Phase IV is not essential to the operation of the other three distinct phases. Thus a method or kit which does not contain the Phase IV component is operable and, in fact, will be preferred when suitable factors, e.g., economy, dictate the non-use of the Phase IV component. As previously noted, whether a Phase IV component is used or not, it is preferably that the period between the completion of the Phase III composition and the start of the Phase I composition in the subsequent sequence not exceed about 4 days.

[0025] The terms "method" and "kit" are used herein to encompass any drug delivery systems via the use of which the 3- or 4-phase scheme outlined above can be effectively administered to human females. Combinations of various dosage forms are operable.

[0026] A unique dosage pattern, i.e., a unique sequence of administration of a novel estrogen/progestogen combination has been discovered which minimizes the administration of estrogen in the first phase of a multiphase regimen, while also minimizing certain side effects, notably breakthrough bleeding, commonly associated with conventional low dosage pills.

[0027] Reasonable variations, such as those which would occur to a skilled artisan, can be made herein without departing from the scope of the invention.

WHAT IS CLAIMED:

1. A multiphase combination and contraceptive kit comprising a package containing daily dosages of:
 - a) a Phase I composition containing a progestogen in an amount equivalent to 0.3 to 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to 0 to less than 10 mcg of ethinyl estradiol;
 - b) a Phase II composition containing a progestogen in an amount equivalent to 0.3 to 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to 10 to 50 mcg of ethinyl estradiol; and
 - c) a Phase III composition containing a progestogen in an amount equivalent to 0.3 to 1.5 mg of norethindrone acetate and an estrogen in an amount equivalent to 10 to 50 mcg of ethinyl estradiol, wherein the amount of estrogen in the Phase III composition is greater than the amount of estrogen in the Phase II composition by at least an amount equivalent to 5 mcg ethinyl estradiol.
2. The kit according to claim 1, wherein the kit contains 5 to 9 dosages of the Phase I composition; 5 to 9 dosages of the Phase II composition; and 8 to 12 dosages of the Phase III composition.
3. The kit according to claim 2, wherein the kit further comprises 1 to 4 dosages of a Phase IV composition which contains ferrous fumarate or is a placebo.
4. The kit according to claim 3, wherein the progestogen in each Phase is norethindrone acetate.
5. The kit according to claim 4, wherein the estrogen in each Phase is ethinyl estradiol.
6. The kit according to claim 5, wherein the Phase II composition contains 20 to 40 mcg of ethinyl estradiol and the Phase III composition contains 25 to 50 mcg of ethinyl estradiol.
7. The kit according to any one of claims 1 to 3, wherein each of the Phase I, II and III compositions contains a progestogen in an amount equivalent to 0.5 to 1.5 mg of norethindrone acetate.

8. The kit according to any one of claims 4 to 6, wherein each of the Phase I, II and III compositions contains norethindrone acetate in an amount from 0.5 to 1.5 mg.
9. The kit according to claim 6, wherein the Phase I composition contains 1 mg of norethindrone acetate, the Phase II composition contains 1 mg of norethindrone acetate and 30 mcg of ethinyl estradiol, the Phase III composition contains 1 mg of norethindrone acetate and 35 mcg of ethinyl estradiol, and the Phase IV composition contains 75 mg of ferrous fumarate.
10. The kit according to claim 9, wherein the kit contains 7 dosages of the Phase I composition, 7 dosages of the Phase II composition, 10 dosages of the Phase III composition and 4 dosages of the Phase IV composition.
11. The kit according to claim 9, wherein the Phase I composition contains 5 mcg of ethinyl estradiol.
12. The kit according to claim 9, wherein the Phase I composition is free of ethinyl estradiol.
13. The kit according to claim 10, wherein the Phase I composition contains 5 mcg of ethinyl estradiol.
14. The kit according to claim 10, wherein the Phase I composition is free of ethinyl estradiol.
15. The kit according to claim 2, wherein the sequential administration of the Phase I, II and III compositions is repeated after a period of 1 to 4 days has elapsed after the completion of the administration of the Phase III composition and wherein the kit is for use in a method of contraception in a female of child bearing age.
16. The kit according to claim 15, wherein the Phase I composition contains an estrogen in an amount equivalent to 0 to 5 mcg of ethinyl estradiol.
17. The kit according to claim 15, wherein the package further comprises a Phase IV composition containing ferrous fumarate administered on a daily basis between the completion of the administration of the Phase III composition and beginning the repeated sequential administration of the Phase I, II and III compositions.

18. The kit according to claim 15, wherein the package further comprises a Phase IV composition which is a placebo administered on a daily basis between the completion of the administration of the Phase III composition and beginning the repeated sequential administration of the Phase I, II and III compositions.
19. The kit according to claim 15, wherein the Phase I composition is free of estrogen.
20. The kit according to claim 15, wherein the amount of estrogen in the Phase III composition is greater than the amount of estrogen in the Phase II composition by at least an amount equivalent to 5 mcg of ethinyl estradiol.
21. The kit according to claim 15, wherein the progestogen in each Phase composition is norethindrone acetate.
22. The kit according to claim 15, wherein the estrogen in each Phase composition is ethinyl estradiol.
23. The kit according to claim 22, wherein the Phase I composition contains an amount of ethinyl estradiol from 0.5 to 5 mcg.
24. The kit according to claim 23, wherein the Phase I composition contains 5 mcg of ethinyl estradiol.
25. The kit according to claim 22, wherein the Phase I composition is free of ethinyl estradiol.
26. The kit according to claim 22, wherein the amount of ethinyl estradiol in the Phase III composition is at least 5 mcg greater than the amount of ethinyl estradiol in the Phase II composition.
27. The kit according to claim 22, wherein the Phase I composition contains 0 to 5 mcg of ethinyl estradiol and the amount of ethinyl estradiol in the Phase III composition is at least 5 mcg greater than the amount of ethinyl estradiol in the Phase II composition.
28. The kit according to claim 27, further comprising administering on a daily basis a Phase IV composition which contains ferrous fumarate or is a placebo for 1 to 4 days.

29. The kit according to claim 28, wherein the sequential administration of the Phase I, II, III and IV compositions is repeated upon completion of the administration of the Phase IV composition.
30. The kit according to claim 28, wherein the daily administration of the Phase I, II, III and IV compositions is for a 19 to 34 day period.
31. The kit according to claim 30, wherein the daily administration of the Phase I, II, III and IV compositions is for a 28 day period.
32. The kit according to any one of claims 15 to 20, wherein each of the Phase I, II and III compositions contain a progestogen in an amount equivalent to 0.5 to 1.5 mg of norethindrone acetate.
33. The kit according to any one of claims 21 to 31, wherein each of the Phase I, II and III compositions contain norethindrone acetate in an amount from 0.5 to 1.5 mg.
34. The kit according to claim 31, wherein the Phase I composition contains 1.0 mg of norethindrone acetate and is administered for 7 days, the Phase II composition contains 1.0 mg of norethindrone acetate and is administered for 7 days, the Phase III composition contains 1.0 mg of norethindrone acetate and is administered for 10 days, and the Phase IV composition contains 75 mg of ferrous fumarate and is administered for 4 days.
35. The kit according to claim 34, wherein the Phase I composition contains 5 mcg of ethinyl estradiol, the Phase II composition contains 30 mcg of ethinyl estradiol and the Phase III composition contains 35 mcg of ethinyl estradiol.
36. The kit according to claim 34, wherein the Phase I composition is free of ethinyl estradiol, the Phase II composition contains 30 mcg of ethinyl estradiol and the Phase III composition contains 35 mcg of ethinyl estradiol.
37. The kit according to claim 15, wherein the package contains instructions for the sequential administration of:
- (a) a Phase I composition containing 0.3 to 1.5 mg norethindrone acetate and 0 to 5 mcg of ethinyl estradiol for 5 to 9 days;
 - (b) a Phase II composition containing 0.3 to 1.5 mg norethindrone acetate and 20-40 mcg ethinyl estradiol for 5 to 9 days;
 - (c) a Phase III composition containing 0.3 to 1.5 mg norethindrone acetate and 25-50 mcg ethinyl estradiol for 8 to 12 days; and

(d) optionally a phase IV composition that contains ferrous fumarate or is a placebo for 1 to 4 days,

wherein the sequential administration of the Phase I, II and III compositions is repeated and the number of days between completion of the administration of the Phase III composition and beginning the repeated sequential administration of the Phase I, II and III compositions is from 1 to 4 days.

38. The kit according to claim 37, wherein the Phase III composition contains at least 5 mcg of ethinyl estradiol more than the Phase II composition.

39. The kit according to claim 38, wherein the Phase IV composition is administered on a daily basis between the completion of the administration of the Phase III composition and beginning the repeated sequential administration of the Phase I, II and III compositions.

40. The kit according to any one of claims 37 to 39, wherein each of the Phase I, II and III compositions contain 0.5 to 1.5 mg of norethindrone acetate.